

Dictionary of

Alkaloids

SECOND EDITION

with CD-ROM



EDITED BY

John Buckingham
Keith H. Baggaley
Andrew D. Roberts
László F. Szabó

Dictionary of Alkaloids

SECOND EDITION

with CD-ROM

Dictionary of Alkaloids

SECOND EDITION

with CD-ROM



EDITED BY

John Buckingham

Keith H. Baggaley

Andrew D. Roberts

László F. Szabó



CRC Press

Taylor & Francis Group

Boca Raton London New York

CRC Press is an imprint of the
Taylor & Francis Group, an **informa** business

CRC Press
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

© 2010 by Taylor and Francis Group, LLC
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works

Printed in the United States of America on acid-free paper
10 9 8 7 6 5 4 3 2 1

International Standard Book Number: 978-1-4200-7769-8 (Hardback)

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Library of Congress Cataloging-in-Publication Data

Dictionary of alkaloids / editors, John Buckingham ... [et al.]. -- 2nd ed. with CD-ROM
p. cm.
Includes index.
ISBN 978-1-4200-7769-8 (hardcover : alk. paper)
1. Alkaloids--Dictionaries. I. Buckingham, J. II. Title.

RS43L.A53D53 2010
615'.32103--dc22

2009031445

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>

and the CRC Press Web site at
<http://www.crcpress.com>

Table of Contents

Foreword	vii	M.	1177
The Editors	ix	N.	1343
Introduction	xi	O	1401
Classification of Structural Types	xxi	P	1445
Dictionary Entries		Q	1631
A	1	R	1647
B	261	S	1687
C	341	T	1815
D	523	U	1965
E	699	V	1977
F	761	W.	2015
G	805	X	2021
H	849	Y	2029
I	1007	Z	2035
J	1073	Name Index	2043
K	1085	Compound Index Contents.	2243
L	1107	Type of Compound Index.	2245

Foreword

Plants containing alkaloids have served humankind since ancient times. Indeed, some of the oldest surviving written records of commercial transactions involving medicinal plants are of alkaloid-containing plants. For thousands of years, indigenous groups around the world discovered, through self-experimentation with locally available plant extracts, that they could provide materials for hunting prey, culinary enhancement, amelioration from disease, relief of pain, and healing. These plants contributed significantly to the survival of indigenous groups throughout history. This same process of low-throughput clinical screening also identified many plants which were to be avoided because of their toxicity, again due to alkaloids. It is the oral and written records of these investigations which survive today and form the basis of both clinical care in much of the world and aspects of contemporary drug discovery.

A transition occurred at the beginning of the 19th century when investigations on the most significant medicinal and toxic plants of Europe commenced, these included studies on opium, colchicum, nux vomica, ipecac, belladonna, and pepper. As it transpired, all of the isolated, biologically active compounds were alkaloids, and it was these investigations which initiated the diverse studies of alkaloids as a class of natural products which continue today.

Since those early times, there have been six important phases to the study of alkaloids. The first was the tremendous increase in the number of structurally unique and complex alkaloids isolated from plants, structures which defined the new field of heterocyclic chemistry. The second was the dramatic reduction in time and the amount of material required to determine the precise stereochemical and conformational structure of an individual alkaloid as a result of the advances in spectroscopic and chromatographic techniques which permitted ever smaller samples to be structurally characterized. Third, there evolved the understanding that groups of structurally similar alkaloids could, sometimes, be related to each other based on one or two biosynthetic precursors from which chemically rational metabolic schemes could be used to generate the plethora of alkaloid skeleta and structures. Fourth, there was the recognition that this structural diversity, skeletal complexity, and stereo-uniqueness were at the same time a benefit and a bane. While the activity could be exquisite, alkaloids may also have significantly modulated biological activity based on a transformation at a single functional group or stereochemical center. Fifth, as more complex alkaloids were structurally elucidated they became challenging targets for synthetic organic chemists, and continue to be the basis for the stimulation of numerous brilliant synthetic strategies and innovations in organic chemistry. Finally, as knowledge of the biology associated with alkaloids grew in both breadth and depth, initially at the *in vivo* level and eventually at the molecular level, some alkaloids were established to be archetypal entity for a particular biological response.

Consequently, in this 200-year period, many alkaloids became critical components of the global pharmaceutical armamentarium, and tremendous healing has resulted from their clinical application. As our food and beverage practices have evolved, alkaloids have become a core component of every day life for almost everyone on the planet on a moment to moment basis, and caffeine, capsaicin, nicotine, theobromine, adrenaline, dopamine, serotonin, etc. are representative examples. Many of the colors of flowers and vegetables are due to alkaloids, and at the same time, alkaloids represent enormous and potent biological power. Sometimes this power can be ameliorated by dose reduction to produce health beneficent agents (for example, the ergot alkaloids and aconitine); in other instances they must simply be avoided by both animals and humans, such as those in many toxic plants, animals, and fungi. Alkaloids also have led medicinal chemists down productive pathways for drug development, although it is seldom appreciated that the molecular phylogeny of many synthetic drugs can be traced to classical alkaloids such as morphine and atropine. At the same time, there is a “dark side” to alkaloids, because at least three - morphine, cocaine, and nicotine - have powerful social implications causing tremendous human suffering through addiction, disease development, and wars which continue unabated today. It is this good, bad, and downright ugly aspect to alkaloids which has fascinated so many scientists for so long. As a global society we do indeed have this love-hate relationship with alkaloids unlike that of any other group of natural products.

As mentioned, with the rise of spectroscopy beginning in the 1950s, interest developed in examining alkaloids from wider sources than higher plants. Indeed, it appears now that almost every class of organism, insects, reptiles, birds, fungi, bacteria, the diversity of marine organisms, and mammals, including humans, all produce alkaloids. The last twenty years have witnessed an explosion in the numbers and the structural diversity of alkaloids from these sources, and these developments are dramatically reflected in this updated compilation.

From a chemical perspective, it transpires that there is almost no overlap in the structural diversity of plant, fungal, and marine alkaloids. Each is a chemically brilliant world unto itself. Thus, one would not expect to find a monoterpene indole alkaloid in a marine sponge, nor a manzamine alkaloid in a tropical liana. This unimaginable structural diversity places the alkaloid sciences on the cusp of a deeper understanding of their biosynthetic formation. While significant and successful efforts are underway to elucidate pathways and isolate genes which are related to both plant and fungal alkaloids, for marine organisms this aspect is completely unexplored, although the most spectacular new skeleta and potent biological activities are derived from these marine sources.

The past twenty years have witnessed the evolution of three new areas of alkaloid chemistry and biology. The first is that of the integrity of alkaloid biosynthesis. Alkaloids such as the ansa macrolides from *Maytenus* and from *Colubrina*, as well as the ergot alkaloids of the Convolvulaceae are now established to be derived biosynthetically from parasitic or symbiotic fungal organisms, not through the metabolic pathways of the host plant. Second, structural conclusions regarding individual alkaloids can change; an α -hydroxy group of yesterday can become a β -hydroxy today based on refined chemical, spectroscopic, and crystallographic conclusions. As one looks through the *Dictionary of Alkaloids*, it is important to be aware that there are many examples where the synthesis of a proposed structure has failed to support that structure, or where new spectroscopic evidence does not support a proposed structure. While in some cases this has led to refinement of the structure, in others a dangling question remains as to the true identity of the natural alkaloid. Third, there is the connection of alkaloids to the biotechnology revolution, and the marvelous progress that has been made for certain alkaloid pathways in elucidating the enzymes and genes of alkaloid biosynthesis.

What, then, is the “fit” for the *Dictionary of Alkaloids*, and why is it such an important resource for researchers? Burgeoning efforts at structure elucidation and biological experimentation make it extremely difficult to quickly find key literature citations to original sources of information on individual compounds. It was recognized some years ago that there was a core global need for information systems on the isolation, spectral properties, and biological properties of alkaloids. The *Dictionary of Alkaloids* is a critical, absolutely essential component in filling that need. It is the only accumulation in the world of the taxonomy, chemistry, and biology of alkaloids derived from all sources. It provides extensive information to references for the isolation, spectroscopy, and biology of an individual alkaloid, and to its other attributes and uses. Importantly, the alkaloids are frequently collected as a group based on a parent structure. This simplifies the presentation, and accumulates alkaloids having quite different trivial names. The volume is easy to use, and the diverse indexes place the most significant information at one’s fingertips. It is a unique resource, and one which is indispensable to any alkaloid chemist and biologist. The inclusion of a fully substructure searchable CD-ROM in this second edition makes the resource even more valuable.

In spite of the vast range of studies of alkaloids over the years, much remains to be done. There are many plant families which are known to contain alkaloids where the alkaloids remain to be identified; there are untold marine organisms and tens of thousands of microorganisms with unknown alkaloids. There is the genetics of alkaloid biosynthesis in more diverse organisms, and the tremendous potential for being able to produce alkaloids through isolated and stabilized systems in a sustainable manner. The key justification for alkaloid research, though, must be relevance to the enhancement of public health. For many years, justification for alkaloid research was based on their novelty; there was no relationship to biology. Indeed, it was shown several years ago that about 75% of all plant alkaloids had never been evaluated in a single bioassay. There remains a critical need for the wider biological study of known and new alkaloids in high throughput screening systems. Alkaloids will continue to have a critical role to play in the health of the Earth and of its inhabitants for the foreseeable future. They represent a significant aspect in the evolving concept of sustainable drugs and biological agents. However, only through the extensive collaboration of taxonomists, chemists, biologists, pharmacologists, formulation chemists, and clinicians can this potential be realized.

In the 12th century, according to Matthew Fox in “*Original Blessing*,” it was Saint Hildegard of Bingen who, in discussing the relationship of humankind and Nature in philosophical and practical terms, suggested that “*All nature is at the disposal of humankind. We are to work with it. For without it, we cannot survive.*” Rather like humankind, alkaloids, in all of their stunningly beautiful complexity and diversity, are an integral and essential aspect of nature with which we must work to survive.

Geoffrey A. Cordell, Ph.D.
Professor Emeritus
Evanston, IL USA

The Editors

John Buckingham is a former lecturer in organic chemistry at the University of London. He has been involved with the Chapman & Hall (now CRC) chemical database since its inception in 1980, initially as a Chapman & Hall employee, more recently as Editorial consultant. From the database has been produced various editions of the *Dictionary of Organic Compounds* and the *Dictionary of Natural Products* (both of which have been for some years solely electronic). In addition, he compiled (with I.W. Southon) the first edition of the *Dictionary of Alkaloids* (print only) in 1989. He is also the author of the popular science books *Chasing the Molecule* and *Bitter Nemesis; The Intimate History of Strychnine*

Keith H. Baggaley obtained his BSc (Hons) and PhD degrees from the University College of Wales, Aberystwyth followed by a post-doctoral fellowship in chemistry at the Royal Institute of Technology, Stockholm, Sweden, with Professor Holger Erdtman and Professor Torbjorn Norin. He then took up a Junior Research Fellowship with Professor David Ollis at the University of Sheffield before joining Beecham Research Laboratories in Surrey as a research chemist. At Beecham he worked on discovery projects concerned with cholesterol lowering drugs, platelet aggregation inhibitors and semisynthetic antibiotics as well as research on clavulanic acid biosynthesis. He has been an editor of the Chapman & Hall dictionaries since 1996.

Andrew D. Roberts took his BSc (Hons) degree at Trent Polytechnic, Nottingham (now Nottingham Trent University), UK 1973–1977 and completed studies in drug discovery projects at Pfizer (Sandwich) UK. He contributed to the *Dictionary of Organic Compounds* (Chapman & Hall) published in 1982 and has since been a Contributing Editor for other projects including the *Dictionary of Antibiotic Substances* (1988), *Dictionary of Drugs* (1990) and has since been an Editor on the *Dictionary of Natural Products* which was first published in 1993.

László Ferenc Szabó was born in Szentgotthárd (Hungary) on April 15, 1931 and studied in Pharmacy at University of Medicine, Budapest, 1949–54. Member of the Department of Medicinal Chemistry, later of the Department of Organic Chemistry of the Semmelweis University Budapest (scientific assistant, associate professor, ordinary professor), director of the same department 1977–1996, Ph.D. in organic chemistry, Doctor habilitatus. Research visitor at Université Libre de Bruxelles, 1972/73, several times in Merseburg (Germany). Presently professor emeritus of the Semmelweis University. Fields of activity: chemistry of natural products, mainly indole alkaloids (strychnine, strictosidine) and terpenoids (iridoids, secologanin), biogenetic formation of natural products, heterocyclic chemistry, mainly quinolizine compounds, polymer supported synthesis of small molecules, theoretical problems of organic chemistry (PMO theory, graph analysis).

IN HOUSE EDITORIAL TEAM

Prof. R.A. Hill, University of Glasgow

Dr A.M. Hodgson, Consultant

Dr F.M. Macdonald, CRC Press, Boca Raton

Dr S.P. Walford, CRC Press, London

Additional data collected by contributors to the *CRC Chemical Database*

Introduction

1. COVERAGE

The *Dictionary of Alkaloids* is a comprehensive database containing over 40,000 compounds. It is a subset of the *Dictionary of Natural Products* (DNP) database. DNP is an ongoing project based on a 25-year review of the natural product literature. For the present project, the subset of DNP entries referring to alkaloids were carefully checked and reviewed and enhanced with a considerable amount of additional information relating to their natural occurrence. The intention is that the Dictionary should contain data on every known alkaloid, and various checks carried out during the compilation process indicate that it comes very close to achieving this aim. The Editors would be grateful to learn of any accidental omissions.

The compounds present in the Dictionary have been classified under the **Structural Types** section below. This section also includes biosynthetic information on these compound classes.

In compiling the printed version and the CD-ROM, the primary literature has been reviewed up to mid-2008.

2. ORGANISATION OF ENTRIES

The Dictionary is arranged alphabetically by entry name. Every entry is numbered to assist ready location. Many compounds are included as derivatives of main entry compounds but important derivatives have their own individual cross-referenced entries. Use of the CD-ROM indexes enables the rapid location of all compounds in the Dictionary by name or compound type, regardless of their location.

A representative dictionary entry is shown in **Figure 1**.


Entry name	→	Irnigaine	I-177	←	Entry number
		2-Methyl-6-(9-phenylnonyl)-3-piperidinol. 3-Hydroxy-2-methyl-6-(9-phenylnonyl)piperidine			
CAS registry number	→	[180779-56-0]			
Structural formula and stereochemical descriptor	→	 <p style="text-align: center;">Absolute Configuration</p>			
Molecular formula	→	C ₂₁ H ₃₅ NO	317.514	←	Molecular weight
		Revised abs. config.. Alkaloid from tubers of <i>Arisarum vulgare</i> (Araceae). Toxic to brine shrimp larvae. Oil. [α] _D ²⁵ -14 (c, 0.3 in CHCl ₃).			
Derivative heading	→	N-Me: N-Methylirnigaine			
		[180779-57-1]			Derivative synonym
		C ₂₂ H ₃₇ NO	331.54		
		From tubers of <i>Arisarum vulgare</i> . Toxic to brine shrimp larvae. Oil. [α] _D ²⁵ -8 (c, 5 in CHCl ₃).			
		Hydrochloride: [α] _D ²⁰ +2.8 (c, 0.9 in CHCl ₃).			
Bibliographic references	→	Melhaoui, A. <i>et al.</i> , <i>Nat. Prod. Lett.</i> , 1995, 7, 101 (<i>isol, ir, pmr, cmr, struct</i>)			
		Pahl, A. <i>et al.</i> , <i>Tet. Lett.</i> , 1998, 39, 2095-2096 (<i>synth, ir, pmr, cmr, ms, cryst struct</i>)			
		Ma, N. <i>et al.</i> , <i>Chin. J. Chem.</i> , 2003, 21, 1356-1359 (<i>synth</i>)			
					Reference tags

FIGURE 1

2.1 CHEMICAL NAMES AND SYNONYMS

The Dictionary contains a wide range of synonyms which may be (a) those found in the primary literature, (b) *Chemical Abstracts* names, or (c) names added editorially to achieve as much consistency as possible with other closely related substances. Names corresponding to those used by CAS during the 9th and subsequent

Collective Index periods (1973-) are labelled 9CI Names. All important derivatives embedded within entries are named (but see comment on CAS nomenclature below). If a compound cannot be located immediately in the main body of the entries, it is important to use the indexes. The CD-ROM version of the Dictionary is much more highly indexed than the printed version.

The most authoritative current statement of good practice on natural product nomenclature is the document *IUPAC Recommendations 1999 (Pure Appl. Chem., 1999, 71, 587–643)* the full text of which can be read on the *Dictionary of Alkaloids on CD-ROM* (access from the Windows® Start menu item or via the Help menu in the main program).



Some alkaloids have been given trivial names which duplicate those already in the literature, or which are simultaneously published for two or more non-identical substances. Where such a duplication is noticed this is indicated by the dagger symbol (‡) immediately following the name. Borderline cases, especially as between presence or absence of terminal e or suffix A or 1 are considered to constitute duplicates for this purpose.

2.1.1 Nomenclature

It is expected that this Dictionary will be used by a wide variety of scientists, not all of whom will be specialist organic chemists. Therefore compounds have been named so as to facilitate access to their factual data by keeping the nomenclature as simple as possible, whilst still adhering to good practice as determined by IUPAC (the International Union of Pure and Applied Chemistry). A great deal of care has been taken to achieve this aim as nearly as possible.

A much fuller description of the detailed nomenclature of individual classes of alkaloids is given in the **Structural Types** section. The following notes are of general applicability throughout the Dictionary.

1. There are many examples in the primary literature of compounds being named in ways which are violations of good IUPAC practice, e.g. where the substituents are ordered non-alphabetically. These are not reported in their incorrect form but have been corrected.
2. The number of trivial names used for acylating substituents has been kept to a minimum but the following are used throughout the Dictionary.



3. Many other trivial appellations have from time to time appeared in the literature for other acyl groups (e.g. Seneciroyl = 3-methyl-2-butenoyl, Feruloyl = 3-(4-hydroxy-3-methoxyphenyl)-2-propenoyl or 4-hydroxy-3-methoxycinnamoyl).

The term **prenyl** for the common 3-methyl-2-butenyl substituent, $(\text{H}_3\text{C})_2\text{C}=\text{CHCH}_2-$, is used throughout the Dictionary. Several other names for this substituent have from time to time been used and appear in the primary literature including the following:

3-Methyl-2-butenyl (systematic)
 γ,γ -Dimethylallyl
 3,3-Dimethylallyl
 3,3-Dimethylpropenyl
 Dimethallyl
 Isoprenyl
 Isopentenyl
 β,β -Dimethylacrylyl

2.2 CAS REGISTRY NUMBERS

CAS numbers are identifying numbers allocated to each distinctly definable chemical substance indexed by CAS since 1965 (plus some retrospective allocation of numbers by CAS to compounds from earlier index periods). The numbers have no chemical significance but they provide a label for each substance independent of any system of nomenclature. They are extensively used for exchanging information between individuals and databases. The numbers take the form NNNNNN-NN-R, where the total number of digits is five or more and R is a check digit.

For practical purposes, CAS numbers have certain shortcomings arising from their free allocation, resulting in one substance having more than one potential number. Duplication may arise for one of several reasons to do with the detailed chemistry of the substance, for example tautomerism, solvent formation, partially unspecified stereochemistry. There are also replaced numbers. For this reason, Dictionary entries will often contain one or more *Additional CAS numbers* which may help the user to obtain further information about the substance, especially by online searching.

Clearly, the additional CAS numbers given in this Dictionary have to be used with care. Their inclusion in the entry is the result of an editorial decision by the Dictionary contributor that they refer to what is essentially the same substance, but this decision may be a subjective one. Care has been taken to ensure that the main CAS number given in this Dictionary for each substance is the correct one.

Further information on CAS number allocation policy can be obtained from CAS indexes or *The Organic Chemist's Desk Reference*.

2.3 STRUCTURAL FORMULAE

Every attempt has been made to present the structures of chemical substances as accurately as possible according to best current practice and recommendations of IUPAC. As much consistency as possible has been aimed at between closely-related structures, for example, all sugars are shown as Haworth formulae, and whenever possible in complex structures the rings are oriented in the standard Haworth convention so that structural comparisons can be quickly made.

2.4 MOLECULAR FORMULA AND MOLECULAR WEIGHT

The elements in the molecular formula are given according to the Hill convention (C, H, then other elements in alphabetical order). The molecular weights given are formula weights (or more strictly, molar masses in daltons) and are rounded to three places in decimals. In the case of some high molecular mass substances, such as proteins, the value quoted may be that taken from an original literature source and may be an aggregate molar mass.

2.5 PHYSICAL DATA

The Dictionary gives the following physical characteristics of substances, when available; appearance, melting point, boiling point, optical rotation, density, refractive index, solubility, p*K*_a. All of these fields are searchable by numerical value (including range searching) in the CD-ROM version of the Dictionary.

2.5.1 Appearance

Organic compounds are considered to be colourless unless otherwise stated. Where the compound contains a chromophore which would be expected to lead to visible colour, but no colour is mentioned in the literature, the Dictionary entry will mention this fact if it has been noticed by the contributor. An indication of crystal form and recrystallisation solvent is often given but these are imprecise items of data; most compounds can be crystallised from several solvent systems and the crystal form often varies. In the case of the small number of compounds where crystal behaviour has been intensively studied (e.g. pharmaceuticals), it is found that polymorphism is a very common phenomenon and there is no reason to believe that it is not widespread among organic compounds generally.

2.5.2 Melting points and boiling points

The policy followed in the case of conflicting data is as follows:

- Where the literature melting points are closely similar, only one figure (the highest or most probable) is quoted
- Where two or more melting points are recorded and differ by several degrees (the most likely explanation being that one sample was impure) the lower figure is given in parentheses, thus Mp 139° (134–135°)

- c. Where quoted figures differ widely and some other explanation such as polymorphism or incorrect identity seems the most likely explanation, both figures are quoted without parentheses, thus Mp 142°, Mp 205–206°
- d. Known cases of polymorphism or double melting points are noted

Boiling point determination is less precise than that of melting points and conflicting boiling point data is not usually reported except when there appears to be a serious discrepancy between the different authors.

2.5.3 Optical rotations

These are given wherever possible, and normally refer to what the Dictionary contributor believes to be the best characterised sample of highest chemical and optical purity. Where available an indication of the optical purity (op) or enantiomeric excess (ee) of the sample measured follows the specific rotation value.*

Specific rotations are dimensionless numbers and the degree sign which was formerly universal in the literature has been discontinued.

2.6 SPECTROSCOPIC DATA

Many Dictionary entries include ultraviolet spectra which are presented in the format:

[neutral] λ_{\max} 198 (log ϵ 1.55); 224 (sh) (log ϵ 0.61); 241 (sh) (log ϵ 0.55) (H₂O) (Berdy)

where ϵ is the absorption coefficient for a given UV maxima value (λ_{\max}). A description of the solvent conditions used, if reported in the literature, is listed at the beginning and end of the UV data in parentheses. All peak absorptions cited are maxima unless otherwise described, e.g. shoulder/inflection (sh) and end absorption (end). In addition, UV data may be followed by the term 'Berdy' or 'DEREP' indicating from which database the data originated. The absence of these terms implies that the data was abstracted from the primary literature.

On the CD-ROM, all the λ_{\max} values are indexed in the UV Maxima field and can be searched for numerically including range searching. Similarly, the solvent data associated with the UV data are indexed in the UV Solvent field.

2.7 HAZARD AND TOXICITY INFORMATION

2.7.1 General

Toxicity and hazard information is highlighted by the symbol ► and has been selected to assist in risk assessments for experimental, manufacturing and manipulative procedures with chemicals.

The Publishers cannot be held responsible for any inaccuracies in the reported information, neither does the omission of hazard data in the Dictionary imply an absence of this data from the literature. Widely recognised hazards are included, however, and where possible key toxicity reviews are identified in the references. Further advice on the storage, handling and disposal of chemicals is given in *The Organic Chemist's Desk Reference*.

2.7.2 RTECS® Accession Numbers

Many entries in this Dictionary contain one or more RTECS® Accession Numbers.† Possession of these numbers allows users to locate toxicity information on relevant substances from the NIOSH *Registry of Toxic Effects of Chemical Substances*, which is a compendium of toxicity data extracted from the scientific literature.

2.8 BIBLIOGRAPHIC REFERENCES

The selection of references is made with the aim of facilitating entry into the literature for the user who wishes to locate more detailed information about a particular compound. The contents of most references are indicated by reference tags (suffixes) indicating their content and in particular the stereoisomers and derivatives of the parent

*For a recent discussion of the validity and applicability of these terms, see Gawley, R.E., *J. Org. Chem.*, 2006, **71**, 2411–2416.

† RTECS® Accession Numbers are compiled and distributed by the National Institute for Occupational Safety and Health Service of the U.S. Department of Health and Human Services of The United States of America. All rights reserved (1996).

compound which they document. The number of references cited does not indicate the relative importance of a compound; one key recent citation may supersede a number of older ones.

Journal abbreviations generally follow the practice of the Chemical Abstracts Service Source Index (CASSI), except for a short list of very well known journals where the Dictionary gives shorter abbreviations to save space (e.g. *J.A.C.S.* instead of *J. Am. Chem. Soc.*)

2.8.1 Further References

Further useful information on a variety of topics concerned with the structure, description, stereochemistry and nomenclature of organic compounds can be found in the *Organic Chemist's Desk Reference* (Chapman & Hall, 1995; new edition forthcoming).

3. INDEXES

There are two printed indexes:

1. A **Name Index** which lists every compound name and synonym in the Dictionary, and
2. A **Type of Compound** index listing all compounds given in the Dictionary.

The indexes refer to the entry number.

Searches on all text and numerical indexes, as well as structure and substructure searching can be carried out on the CD-ROM version of the Dictionary. For information on the Type of Compound codes used in the Type of Compound Index, please see the **Structural Types** section.

4. ABBREVIATIONS

The following is a selection of the most common Database abbreviations used:

Abbreviation	Name
[α]	specific rotation
abs config	absolute configuration
Ac	acetyl
acc	according
AcOH	acetic acid
Ac ₂ O	acetic anhydride
alk	alkaline
amorph	amorphous
amt	amount
anal	analytical applications, analysis of detection
anhyd	anhydrous
aq	aqueous
BAN	British Approved Name
bibl	bibliography
biosynth	biosynthesis
Bp	boiling point
c	concentration
ca	(<i>circa</i>) about
cd	circular dichroism
CAS	Chemical Abstracts Service
chromatog	chromatography
cmr	carbon (¹³ C) nuclear magnetic resonance
CNS	central nervous system
col	colour, coloration
coml	commercial(ly)
compd	compound
conc	concentrated
config	configuration

conformn	conformation
constit	constituent
cryst struct	X-ray crystal structure determination
d	density
dec	decomposes, decomposition
degradn	degradation
deg	degree
deriv(s)	derivative(s)
detn	detection, determination
dil	dilute, dilution
dimorph	dimorphic
diss	dissolves, dissolved
dist(n)	distil, distillation
DMF	dimethylformamide
DMSO	dimethyl sulfoxide
ee	enantiomeric excess
epr	electron paramagnetic (spin) resonance
equilib	equilibrium
esp	especially
Et	ethyl
EtOAc	ethyl acetate
EtOH	ethanol
EtOH aq	aqueous ethanol
exp	experimental
FEMA	Flavor and Extract Manufacturers' Association
fl p	flash point
fluor	fluoresces, fluorescence
formn	formation
Fp	freezing point
g	gram
glc	gas liquid chromatography
Glc	β -D-glucopyranosyl
GRAS	Generally Recognised As Safe
ham	hamster
haz	hazard
hplc	high performance liquid chromatography
hydrol	hydrolyses, hydrolysed, hydrolysis
ihl	inhalation
ims	intramuscular
INN	International Nonproprietary Name
intermed	intermediate
ipr	intraperitoneal
ir	infra-red spectrum
isol(n)	isolation, isolated
isom	isomerism, isomers, isomerises
ivn	intravenous
JAN	Japanese Accepted Name
JMAF	Japanese Ministry for Agriculture, Forestry and Fisheries
LC	lethal concentration
LD	lethal dose: LD ₅₀ , a dose which is lethal to 50% of the animals tested
M	molecular weight (formula weight)
manuf	manufacturer, manufactured
max	maximum
Me	methyl
MeOH	methanol

Me ₂ CO	acetone
MEL	maximum exposure limit
metab	metabolite, metabolism
misc	miscible
mixt	mixture
mod	moderately
Mp	melting point
ms	mass spectrum
mus	mouse
<i>n</i>	index of refraction, e.g. n_D^{20} for 20° and sodium light
nmr	nuclear magnetic resonance spectrum (general)
obt	obtained
oc	open cup
occup	occupational
OES	Occupational Exposure Standard
op	optical purity
ord	optical rotatory dispersion
orl	oral
Ph	phenyl (C ₆ H ₅)
pharmacol	pharmacology
pmr	proton (¹ H) nuclear magnetic resonance
polarog	polarography
polym	polymerises, polymer
ppd	precipitated
ppm	parts per million
props	properties
purifn	purification
Py	pyridine
rbt	rabbit
ref	reference
resoln	resolution
rev	review
rt	room temperature
scu	subcutaneous
sepn	separation
skn	skin
sl	slightly
sol	soluble
soln	solution
solv	solvent
sp	species (singular)
spar	sparingly
spp	species (plural)
ssp	subspecies
subl	sublimation, sublimes
synth	synthesis
tautom	tautomerism
THF	tetrahydrofuran
tlc	thin layer chromatography
TLV	Threshold Limit Value
tox	toxicity
unsatd	unsaturated
USAN	United States Adopted Name
uv	ultraviolet spectrum
v	very

var	variety
vis	visible
vol	volume

4.1 REFERENCE TAGS

The following is a selection of the most common Reference Tag abbreviations used:

Abbreviation	Name
abs config	absolute configuration
anal	analysis
bibl	bibliography
biosynth	biosynthesis
cd	circular dichroism
chromatog	chromatography
cmr	¹³ C nuclear magnetic resonance spectrum
config	configuration
conformn	conformation
cryst struct	X-ray crystal structure determination
deriv(s)	derivative(s)
detn	determination, detection
dta	differential thermal analysis
glc	gas-liquid chromatography
hplc	high performance liquid chromatography
ir	infrared spectrum
isol	isolation
isom	isomerism
manuf	manufacture
metab	metabolism
ms	mass spectrum
nmr	nuclear magnetic resonance spectrum
occur	occurrence
ord	optical rotatory dispersion
pharmacol	pharmacology
pmr	proton (¹ H) nuclear magnetic resonance spectrum
props	properties (chemical or physical)
Raman	Raman spectrum
resoln	resolution
rev	review
sepn	separation
struct	structure
synth	synthesis
tautom	tautomerism
tlc	thin layer chromatography
tox	toxicity
trans	transition(s)
uv	ultraviolet spectrum
uv-vis	ultraviolet visible spectrum

5. THE DICTIONARY OF ALKALOIDS ON CD-ROM

The *Dictionary of Alkaloids* is published together with a fully searchable CD-ROM. Space considerations have precluded the inclusion of indexes other than the Name Index and the Type of Compound index in the hard-copy version. By contrast, the CD-ROM contains searchable indexes on the following 35 fields:

Accurate Mass	Chemical Name	Molecular Weight	Supplier
All Entries	Density	Optical Rotation	Type of Compound
All Text	Development Status	Partition Coefficient (Calc.)	Type of Compound Words
Biological Source	Dissociation Constant	Percent Composition	Type of Organism
Biological Use/Importance	Hazard & Toxicity	References	Type of Organism Words
Boiling Point	Hazard Flag	Refractive Index	Use/Importance
Boiling Point Pressure	Ion Charge	Rotation Conditions	UV Maxima
CAS Registry Number	Melting Point	RTECS Accession No.	UV Solvent
CRC Number	Molecular Formula	Source/Synthesis	

In addition to searching the above text fields, it is possible to search on structure and substructure.

Once installed, a Help file providing additional information on data content and guide to searching is available from the CRC Press folder in the Start Menu and from the Help menu on the CD-ROM.

When accessing the *Dictionary of Alkaloids on CD-ROM* the first screen that is obtained is the Search Form window (**Figure 2**).




FIGURE 2


The Search Form window is split into three panes:

1. Structure Search pane - allowing structure and substructure searching
2. Search Terms pane - search from one or more of the 35 available data/text fields
3. Index pane - displays the indexed terms within a selected field

From the Search Form window design your search profile using text, structure or text/structure searching. Once your search has been performed the resultant hits are listed alphabetically by chemical name in the Hit List screen. Clicking on any one of the hits in the Hit List screen will result in that entry being displayed in the Entry Display screen (**Figure 3**).

<input checked="" type="checkbox"/>	Name	CAS Registry Number	Molecular Formula
<input type="checkbox"/>	Amauromine	88360-87-6	C ₃₂ H ₃₈ N ₄ O ₂
<input type="checkbox"/>	Amauromine; 2,3-Diepimer	143168-22-3	C ₃₂ H ₃₈ N ₄ O ₂
<input type="checkbox"/>	Amauromine; 2,3-Diepimer, <i>N</i> ¹ -Me	143006-28-6	C ₃₃ H ₃₉ N ₄ O ₂
<input type="checkbox"/>	Amauromine; Stereoisomer	97859-13-7	C ₃₂ H ₃₈ N ₄ O ₂
<input type="checkbox"/>	Antibiotic Sch 52900	174285-71-3	C ₃₁ H ₃₈ N ₄ O ₇ S ₄
<input type="checkbox"/>	Antibiotic Sch 52900; 1 ¹ -Deoxy	174285-72-4	C ₃₁ H ₃₈ N ₄ O ₆ S ₄
<input type="checkbox"/>	Antibiotic T 9888		C ₃₂ H ₃₈ N ₄ O ₇ S ₂
<input type="checkbox"/>	Antibiotic WIN 64745	150881-28-0	C ₃₁ H ₃₈ N ₄ O ₄
<input type="checkbox"/>	Antibiotic WIN 64821	150881-27-9	C ₄₀ H ₃₈ N ₄ O ₄
<input type="checkbox"/>	Asperazine	198953-76-3	C ₄₁ H ₃₈ N ₄ O ₄
<input type="checkbox"/>	Brevianamide E	23454-27-5	C ₂₁ H ₂₈ N ₂ O ₃
<input type="checkbox"/>	Brevicompanine A	215121-46-3	C ₂₂ H ₂₇ N ₂ O ₂
<input type="checkbox"/>	Brevicompanine C		C ₂₁ H ₂₇ N ₂ O ₂
<input type="checkbox"/>	Chaetocin	28097-03-2	C ₃₃ H ₃₈ N ₄ O ₇ S ₄
<input type="checkbox"/>	Chaetocin; 6 <i>S</i> -Hydroxy	12794-84-2	C ₃₃ H ₃₈ N ₄ O ₇ S ₄
<input type="checkbox"/>	Chaetocin; 19-Deoxy, 6 or 6' <i>S</i> -hydroxy	12794-85-3	C ₃₃ H ₃₈ N ₄ O ₆ S ₄
<input type="checkbox"/>	Chaetocin; 6 <i>S</i> ,6' <i>S</i> -Dihydroxy	37934-52-4	C ₃₃ H ₃₈ N ₄ O ₈ S ₄
<input type="checkbox"/>	Chaetocin B	118101-82-9	C ₃₃ H ₃₈ N ₄ O ₆ S ₃
<input type="checkbox"/>	Chaetocin C	118111-08-3	C ₃₃ H ₃₈ N ₄ O ₆ S ₃
<input type="checkbox"/>	Chaetocochin A	912551-37-2	C ₃₈ H ₄₂ N ₄ O ₆ S ₄
<input type="checkbox"/>	Chaetocochin A; 3,11 <i>β</i> -Di- <i>S</i> -de-Me, 3,11 <i>α</i> -disulfide	912551-38-3	C ₃₄ H ₃₆ N ₄ O ₆ S ₄
<input type="checkbox"/>	Chaetocochin C	912551-39-4	C ₃₃ H ₃₈ N ₄ O ₆ S ₄
<input type="checkbox"/>	Chaetomin	1403-36-7	C ₃₁ H ₃₈ N ₄ O ₆ S ₄
<input type="checkbox"/>	Chaetomin; Ring D trithia homologue		C ₃₁ H ₃₈ N ₄ O ₆ S ₃
<input type="checkbox"/>	Chaetomin; Ring G trithia homologue		C ₃₁ H ₃₈ N ₄ O ₆ S ₃
<input type="checkbox"/>	Chetrach A	99615-92-6	C ₃₃ H ₃₈ N ₄ O ₆ S ₃
<input type="checkbox"/>	Dethiotetrakis(methylthio)	84260-71-9	C ₃₈ H ₄₂ N ₄ O ₆ S ₄

Entry Name: Brevianamide E



Absolute Configuration

CRC Number: CHF45-G
 CAS Registry Number: 23454-27-5
 Type of Compound Code(s): ZG3000 VX4110 WV0150 AH0150

Molecular Formula: C₂₁H₂₈N₂O₃
 Molecular Weight: 367.447
 Accurate Mass: 367.189592
 Percentage Composition: C 68.64%; H 6.86%; N 11.44%; O 13.06%
 General Statement: Diketopiperazine
 Biological Source: Metab. from *Penicillium brevi-compactum* and *Penicillium vindicatum*
 Biological Use/Importance: Hepatotoxin
 Physical Description: Noncryst.
 Optical Rotation: [α]_D²⁰ -157 (c, 0.093 in EtOH)
 UV: [neutral] λ_{max} 239 (ε 7500); 296 (ε 2050) (EtOH)

References:
 Birch, A.J. *et al.*, *Tetrahedron*, 1970, **26**, 2329-2344 (*isol, uv, ir, pmr, biosynth*)
 Ritche, R. *et al.*, *Chem. Comm.*, 1975, 611-612 (*config*)
 Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1981, 959-963 (*synth*)
 Ritche, R. *et al.*, *Tetrahedron*, 1981, **37**, 4295-4303 (*synth*)
 Schkeryantz, J.M. *et al.*, *J.A.C.S.*, 1999, **121**, 11964-11975 (*synth*)
 Williams, R.M., *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (*rev, synth, biosynth*)

[00016584-9] V0 E0

FIGURE 13

Any comments and suggestions for inclusion may be sent to:

The Editors, Dictionary of Alkaloids
 CRC Press/Taylor & Francis Group
 Albert House, 4th Floor
 1-4 Singer Street
 London EC2A 4BQ

Fax: +44 (0)20 7017 6747
 Email: steve.walford@informa.com

Classification of Structural Types

1. DEFINITION OF AN ALKALOID

Alkaloids are a large group of nitrogen-containing secondary metabolites produced by almost every variety of organisms including humans.

The term was originally ascribed to pharmacologically active bases of plant origin, but the definition has subsequently been broadened so that it is now generally considered to include the majority of nitrogen-containing natural products with the exception of the simple amino acids and proteins, and nitrogen-containing glycosidic substances such as the aminoglycoside antibiotics. Alkaloids may be of plant, animal, insect or microbial origin. Basic properties may be weak or absent as in the various types of amide alkaloids. Not all higher plants produce alkaloids, but they are fairly widely distributed. An estimated 40% of plant families contain at least one species that is known to produce alkaloids. Traditional screening methods, e.g. using Dragendorff's reagent, may fail to detect weakly basic alkaloids.

A narrower definition has also been used, according to which the nitrogen atom(s) must be in a negative oxidation state, which excludes for example nitro and nitroso compounds. Other terms which have been used to narrow the definition include 'Imperfect alkaloids' (*Alkaloida imperfecta*), 'Pseudoalkaloid' (compounds in which the skeleton is not amino acid-derived) and 'Protoalkaloid' (bases that are amino acid-derived but acyclic, or contain rings not incorporating the amino acid-derived nitrogen). However, these terms are no longer in common use. Compounds of all of these types are in general included in the Dictionary, as well as most types of depsipeptide in which one or more amino acid residues are modified to a nitrogen heterocycle.

At the other extreme, many scientists in the life sciences/nutrition area tend to define an alkaloid as any nitrogenous toxin, including proteinaceous substances such as ptomaines. These are not included in this Dictionary.

In addition to the exclusions mentioned above, other types of nitrogenous natural product which are not covered by this Dictionary include the Sphingamines (aminolipids which are components of cerebrosides and other important biolipids) and the numerically limited but important polypyrroles, including haem, chlorophyll, etc., which can be considered as pyrrole alkaloids according to some definitions. Full entries for these are given in the parent work, *Dictionary of Natural Products*.

2. BIOSYNTHETIC CONSIDERATIONS

A great deal is now known about the biochemistry and enzymology involved in the synthesis of alkaloids by higher plants and microorganisms; because of experimental difficulties and the complexity of the ecosystems, much less is known about marine alkaloids. Purely chemical biosynthetic studies involving tracer elements are currently somewhat out of fashion, but important work in the 1970s and 1980s by several research groups delineated the framework of the biosynthetic pathways for most of the major groups of alkaloids.

In recent years, biosynthetic studies have extended into genomics, and for some important alkaloids, the whole process from nucleic acids to final assembly is now known in significant detail. The details of these pathways lie outside the scope of this Dictionary. An indication of the amount of information that is potentially available is given by the fact that, for example, eight different enzymes are involved in the construction of the relatively straightforward alkaloid Berberine, including one (berberine bridge enzyme) which is solely responsible for catalysing the insertion of the final methylene bridge into a C₁₉ precursor. However, this section gives a general account of the biosynthetic and biogenetic routes to the different classes based on experimental evidence and theoretical ideas of structural interrelationships. Much more information can be obtained by consulting the biosynthesis references given in many of the individual dictionary entries.

3. NOMENCLATURE

As with all types of natural product, at least four different types of name can be distinguished. They are not given equal prominence in this Dictionary. Changes occur with time, notably associated with migrations in *Chemical Abstracts* nomenclature policy.

1. *Trivial names.* Example; Corynoxine. These convey no structural information, and if the structure of the alkaloid has not been determined, this will be the only name available. The oldest date from the first alkaloid isolations in the early nineteenth century and they are very frequently, though not always, derived from the Latin binomial or other name for the originating species, e.g. Strychnine from *Strychnos nuxvomica*. They are given prominence in this Dictionary, and every effort has been made to include every trivial name that has been used in the past. The names have the advantage that since they convey no structural information, they are unchanged if there is a structure revision of the alkaloid, a common event.

There are a number of duplications of trivial names in the literature, and these are indicated by the sign ‡.

Trivial names for at least plant-based alkaloids have traditionally ended in *-ine*.

2. *Systematic names.* Example: Methyl 6'-ethyl 1,2,2',3,6',7',8',8'*a*-octahydro- α -(methoxymethylene)-2-oxospiro[3*H*-indole-3,1'(5'*H*-indolizine]-7'-acetate, ($\alpha E,1'S,6'S,7'S,8'aS$)- is the current CAS name for Corynoxine. The names are unambiguous and put natural products on the same footing as other organic compounds, but are of course cumbersome and typically not interpretable in terms of a recognizable structure. They are also subject to change if CAS nomenclature policy changes. These names are given in the Dictionary for relatively small molecules, and the name used to head the entry may be a systematic one if it is short and convenient. For larger molecules, such as Corynoxine, where this Dictionary does not give the systematic (CAS) name, it can be readily accessed using the CAS registry numbers that are given throughout the Dictionary. Systematic names, where they are shown in an entry, are usually presented without their stereodescriptors. These can be seen by examining the structure diagram or by consulting CAS.

There are many cases in which CAS numbering does not correspond with the biogenetic schemes used by most natural products scientists, and which are given prominence in this Dictionary. Workers in the field are aware of such discrepancies which are inevitable. Another common complication is that structural changes which to the alkaloid specialist are minor (e.g. introduction of a methylenedioxy group) cause profound changes to the CAS name, in this case because the methylenedioxy being is recognised as an additional heterocyclic ring fusion. For further details, see the compound type descriptions given below and the individual dictionary entries.

3. *Semisystematic names.* Examples are the skeletons represented by Corynoxan and Ergoline. IUPAC gives lists of these and directions for introducing new ones. These semisystematic parents can be modified by operators such as *nor-*, *abeo-*, and *seco-*, etc. CAS still uses this type of name for some types of alkaloid such as ergolines (e.g. Ergometrinine = 9,10-Didehydro-*N*-(2-hydroxy-1-methylethyl)-6-methylergoline-8-carboxamide), but has abandoned it for many other alkaloid classes, e.g. Corynoxans, in favour of systematic nomenclature. In general, the construction of semisystematic names of this type has not found universal support in view of the discovery of ever more structurally complex types of natural product, and also of the potential complexities caused by different authors deriving new parent skeletons in different ways. These names are not in general given in this Dictionary.

4. *'Semitrivial' names.* These are names which are derived by appending a systematically-derived operator to a trivial parent. Examples are 8-(1,1-Dimethylallyl)confusameline and *N*-Cyano-*sec*-pseudostrychnine. Such names are widespread, but are best avoided because of the possibilities for confusion, especially when there is a structure revision, or where there is more than one numbering scheme in use for the parent skeleton. Trivial names should be preferred. These semitrivial names are however, reported throughout the Dictionary where they are found in the literature. Where a semitrivial name is no longer accurate because there has been a structure revision, a note is given in the dictionary entry, and reference is often given to the possibility of confusion caused by the use of different numbering schemes.

4. PRINCIPAL LITERATURE SOURCES

4.1 REVIEW SERIES

Progress in the Chemistry of Organic Natural Products, formerly *Fortschritte der Chemie Organischer Naturstoffe*, (various editors), Springer Verlag, 1938- (**'Prog. Chem. Org. Nat. Prod.'**).

The leading annual review series for all types of natural product, although recent volumes contain fewer contributions on alkaloids.

Alkaloids, Chemistry and Biology (ed. Manske, R.H.F., then Rodrigo, R., then Brossi, A. then Cordell, G.A.), Academic Press, then Wiley, then Elsevier, 1949- (**'Alkaloids'**).

The leading review series devoted to alkaloids. Referred to in CAS as *'Alkaloids, N.Y.'*, or *'Alkaloids (Academic Press)'*, (but no longer abstracted)

Alkaloids, Chemical and Biological Perspectives (ed. Pelletier, S.W.), Pergamon/Elsevier, 1983–2000 (*Alkaloids Chem. Biol. Perspect.*).

Another good review series, although no longer published.

Natural Product Reports, Royal Society of Chemistry, 1984–.

Timely, succinct and authoritative reviews; coverage depends on the availability of authors for particular topics.

Boit, H.-G., *Ergebnisse der Alkaloid-Chemie bis 1960*, Akademie Verlag, 1961.

Comprehensive compilation of all alkaloid isolations to 1960.

4.2 RECENT BOOKS

Raffauf, R.F., *Plant Alkaloids: A Guide to their Discovery and Distribution*, Food Products Press, 1996

Alkaloids: Biochemistry, Ecology and Medical Applications (eds. Roberts, M.A. *et al.*), Plenum Press, 1998

Compr. Nat. Prod. Chem. (ed. Kelly, J.W.), Elsevier, 1999, 4

Hesse, M., *Alkaloids: Nature's Curse or Blessing?*, Wiley-VCH, 2002

Aniszewski, T., *Alkaloids – Secrets of Life*, Elsevier, 2007

Modern Alkaloids: Structure, Isolation, Synthesis and Biology (eds. Fattorusso, E. *et al.*), Wiley-VCH, 2007

5. TYPE OF COMPOUND CLASSIFICATION SCHEME

All alkaloids in the Dictionary carry one or more Type of Compound codes in the format VX1234, where VX is an invariable code denoting alkaloids (other codes starting with V refer to other types of natural product that can be found in the *Dictionary of Natural Products*. Some alkaloids carry additional non-alkaloid codes).

These codes are shown against the headings below and can be searched in the CD-ROM version of the Dictionary or in the parent *Dictionary of Natural Products* (DVD or web version). The VX code numbers assigned to an alkaloid type have no intrinsic significance. They are allocated so as to make the groups and subgroups appear in a reasonable biochemical sequence, and to provide gaps in the numerical order so that newly discovered types of alkaloid can be inserted.

Use of these codes in searching is complementary to searching by substructure. In some classes of alkaloid (such as the various types of steroidal alkaloid), the structures are fairly homogeneous, but in other classes (for example, the *Daphniphyllum* alkaloids), a wide variety of carbon skeleton may be produced by variation of an underlying biogenetic scheme. In such cases, substructure searching cannot locate all of the possibilities and the Type of Compound coding is of great value. Some alkaloids carry more than one code where they are of an intermediate type, or contain structural fragments that can be assigned to two different categories.

Biogenetically and structurally the alkaloids are diverse, and it is usual to discuss them in terms of biogenetic origin rather than purely on the basis of structural features. The classical plant alkaloids were from an early date recognised as being mostly derived from amino acids, notably ornithine (→ pyrrolidines, tropanes, etc.), lysine (→ piperidines), phenylalanine, and tryptophan (→ many alkaloids including the numerous and structurally diverse indole alkaloids). The order of Type of Compound codes in the first edition of this Dictionary followed this general scheme, which was also followed in most treatises on alkaloids. However, the subsequent discovery of many more diverse types of alkaloid, especially those of microbial and marine origin, has blurred these traditional biogenetic categories. New categories have now been inserted where it seemed appropriate on structural grounds, but these may not necessarily be biogenetically related to the established alkaloid types. To take one example, the Ericamycins (VX3900) have been placed at the end of the isoquinoline alkaloids because they contain a condensed isoquinoline nucleus, but they are of polyketide origin and have little in common biogenetically with the 'traditional' isoquinoline alkaloids. Even some well-established plant alkaloids were found on investigation to be polyketide in origin. The best-known example is Coniine, a long-known piperidine alkaloid which, however, is not lysine-derived. Other examples include Pinidine and (probably) Carpaine.

Much less is known about the biogenesis of marine alkaloids than their terrestrial counterparts because of experimental difficulties and the complexity of the ecosystems. Where structural similarities exist between certain marine alkaloids and their terrestrial counterparts, this is frequently indicated in the following sections, but this should not be taken to imply that their biosynthesis takes place by identical routes. There may be cases of biochemical parallelism arising from convergent evolution. In most instances the relevant information is simply not yet available for the marine alkaloids.

Details are now being uncovered about formerly unsuspected enzyme systems, which superimpose new biosynthetic possibilities on top of the traditional routes, thus blurring the traditional alkaloid subtypes.

Foremost among these are the 'Diels-Alderases' (not confined to alkaloids) which are capable of catalysing the formation of new carbon skeletons from known types. The Type of Compound categories in the database are therefore subject to future revision in the light of new research results.

Brief descriptions of the alkaloid types follow. These are not meant to be definitive, as the amount of information available for most groups of alkaloid is too great to summarise in such a brief introduction. However, the Dictionary as a whole will point the user to the key literature sources for every group; either via the general reviews given below under each category, or by going to the individual dictionary entries for the alkaloids in each group and from there to the references cited in the entry.

Herbert, R.B., *Nat. Prod. Rep.*, 2003, **20**, 494–508 (rev, *biosynth*)

Oikawa, H. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 321–352 (rev, *Diels-Alder enzymes*)

Moore, B.S., *Nat. Prod. Rep.*, 2005, **22**, 580–593; 2006, **23**, 615–629 (rev, *biosynth*)

SIMPLE ACYCLIC AMINE ALKALOIDS WITH ONE N (VX0100)

These include simple amines of mostly widespread biological occurrence such as **Choline** and **Ethylamine**. Various miscellaneous polyamines are categorised under VX9000, miscellaneous acyclic alkaloids.

SIMPLE GUANIDINES (VX0150)

This heading covers low molecular weight guanidinoid bases, many of marine origin. Their high polarity demands special isolation techniques. The guanidine or modified guanidine group is common as a structural component of higher molecular weight alkaloids, including hybrid polyketide alkaloids and peptide-alkaloids which are classified elsewhere below.

Berlinck, R.G.S., *Prog. Chem. Org. Nat. Prod.*, 1995, **66**, 119–26 (rev)

Nagasawa, K. *et al.*, *Chem. Rec.*, 2003, **3**, 201–211 (rev, *synth*)

Berlinck, R.G.S. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 516–520; 2008, **25**, 919–954 (revs)

NITRILES, ISONITRILES AND RELATED COMPOUNDS (VX0200)

Compounds containing the -NC, -NCS and -NHCHO groups frequently occur together and are biogenetically related. Compounds having the same carbon skeleton are usually included in the same entry. It now appears that the thiocyanates are central in their biosynthesis.

Edenborough, M.S. *et al.*, *Nat. Prod. Rep.*, 1988, **5**, 229–246 (rev, *isonitriles*)

Fleming, F.F., *Nat. Prod. Rep.*, 1999, **16**, 597–606 (rev, *nitriles*)

Garson, M.J. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 164–179 (rev)

SIMPLE ISOBUTYLAMIDE ALKALOIDS (VX0220); OTHER SIMPLE AMIDE ALKALOIDS (VX0250)

Simple isobutylamide amides are common plant constituents, VX0250 includes a variety of other simple amides such as the marine **Variceramides** and a wide range of other miscellaneous amides. The most intensively studied are the capsaicinoids, such as **Capsaicin**, which are responsible for the pungent flavours of capsicum peppers (Solanaceae) and related plants.

Suzuki, T. *et al.*, *Alkaloids*, 1984, **23**, 228–300 (rev, *capsicum alkaloids*)


Cordell, G.A. *et al.*, *Ann. Pharmacother.*, 1993, **27**, 330–336 (rev)

AZO AND AZOXY ALKALOIDS (VX0280)

These are not very numerous and their classification as alkaloids is borderline. They include toxic (carcinogenic) metabolites such as **Cycasin**.

SIMPLE PYRROLIDINE ALKALOIDS (VX0300, VX0380)

Several simple pyrrolidine alkaloids are known from plants. These include **Hygrine** and **Stachydrine**. In *Nicandra physaloides*, Hygrine is biosynthesised from ornithine. Condensation of two ornithine units with acetoacetate gives **Cuscohygrine**. Other alkaloids containing a pyrrolidine ring include **Ficine** (in which the pyrrolidine ring is attached to a flavone nucleus), and **Brevicolline** (in which it is attached to a β -carboline unit). Clearly the biogenesis of these molecules requires other precursors.



Hygrine


Other simple pyrrolidine alkaloids may arise by different biosynthetic pathways. The kainoids, e.g. **α -Kainic acid**; **Domoic acid**, are a group of non-proteinogenic pyrrolidinedicarboxylic acids. α -Kainic acid was first isolated from a rhodophyte alga but has since been found in terrestrial organisms. The kainoids are biosynthesised from glutamate and geranyl pyrophosphate, while many others are presumably proline-derived. Simple pyrrolidines based on a longer-chain skeleton such as the **Broussonetines** are biosynthesised analogously to sphingosines. Many other compounds indexed under miscellaneous pyrrolidine alkaloids (VX0380) contain a pyrrolidine function as part of a larger structure.

Massiot, G. *et al.*, *Alkaloids*, 1986, **27**, 270–332 (rev)

Parsons, A.F., *Tetrahedron*, 1996, **52**, 4149–4174 (kainoids, rev)

CHROMONE AND FLAVONOID ALKALOIDS (VX0340, VX0350)

A structure consisting of a pyrrolidine, piperidine or pyridine ring linked to the A ring of chromone is referred to as a chromone alkaloid. This group of compounds can be sub-divided into two types, namely those in which the chromone nucleus exists as Noreugenin (5,7-Dihydroxy-2-methylchromone) — indexed as chromone alkaloids VX0340 — and those which bear a phenyl substituent at C-2 (indexed as flavonoid alkaloids, VX0350). The former group is typified by **Rohitukine** and **Schumannificine**. Typical flavonoid alkaloids include **Ficine** and **Vochysine**. Compared with the noreugenin-related alkaloids, which have only been isolated from the plant families Meliaceae and Rubiaceae, the flavonoid alkaloids are more widely distributed throughout the higher plants.




Ficine

Houghton, P.J. *et al.*, *Alkaloids*, 1987, **31**, 67–100 (chromone alkaloids; rev)

NICOTINE-LIKE ALKALOIDS (VX0360)


This is a small group of well-known pharmacological and toxicological importance, **Nicotine** being the major alkaloid of tobacco. The pyridine ring in nicotine is derived from nicotinic acid and the pyrrolidine ring from *N*-methylpyrrolinium derived from arginine or ornithine via putrescine. The genomics and enzymology of nicotine biosynthesis are well studied. **Nornicotine** is a deleterious tobacco constituent which is a major cause of carcinogenesis through *N*-nitrosation.



Nicotine

TETRAMIC ACIDS (VX0390)

These are longer-chain pyrrolidines exemplified by the halogenated **Aurantosides** and **Rubrosides** from sponges (microbial products). A range of nonhalogenated tetramic acids has been isolated from terrestrial microorganisms, such as **Erythrokyrin** and the **Lipomycins**. They are polyketide in origin.




Erythroskyrin

Schobert, R. *et al.*, *Bioorg. Med. Chem.*, 2008, **16**, 4203–4221 (rev)

TROPANE ALKALOIDS (VX0400)

These are characteristic constituents of the Solanaceae. Almost all of them are esters of mono-, di-, and trihydroxytropans, with a wide variety of hydroxylation patterns (see for example **Calystegines**). In view of their pharmacological importance (**Cocaine**, **Atropine** and relatives) their synthesis and biosynthesis has been much studied since the nineteenth century. They are derived from ornithine and acetoacetate. One or two condensed structures occur, e.g. **Bellendine**, **Darlingine**.



Cocaine

Hemschiedt, T., *Top. Curr. Chem.*, 2001, **209**, 175–206 (rev, biosynth)


Gossauer, A., *Prog. Chem. Org. Nat. Prod.*, 2003, **86**, 1–188 (rev)

Biastoff, S., *et al.*, *Alkaloids*, 2007, **64**, 49–102 (*Calystegines*)

PYRROLIZIDINE ALKALOIDS (VX0440, VX0500, VX0520)

These occur in species of the *Senecio* genus, and elsewhere in the Asteraceae and Fabaceae. They have been shown to be responsible for the toxic effects, particularly liver damage, in livestock grazing on pastures infested by these species, and are of public health interest in human liver disease. In view of this they have been extensively reviewed. Many, and perhaps the majority, of pyrrolizidine alkaloids occur in the plant as *N*-oxides, the *N*-oxide function being lost during isolation. Toxicity appears to be the result of oxidation *in vivo* to pyrrole derivatives. Pyrrolizidine alkaloids enter the food chain and become antifeedants in animal species such as butterflies.

The majority of pyrrolizidine alkaloids are either relatively simple esters formed from a pyrrolizidine base, the necine, exemplified by **Retronecine**, and a necic acid, such as senecioic (VX0440), or more complex cyclic esters formed between a necine and a necic acid (VX0500), an example being **Monocrotaline**. The necic acids are probably derived from an amino acid (e.g. isoleucine), rather than acetate or mevalonate. **Callosine** represents an unusual variation.



Retronecine

Monocrotaline

The category of miscellaneous pyrrolizidine alkaloids includes some unusual pyrrolizidines from totally different sources, e.g. **Antibiotic PF 1018** from a fungus, **Nitropolyzonamine** from a millipede, and the cyclic polysulfide **Cassipourine** from a higher plant.

Robins, D.J., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 115–204 (rev)

Wróbel, J.T., *Alkaloids*, 1985, **26**, 327–384 (rev)

Powell, R.G. *al.*, *Alkaloids Chem. Biol. Perspect.*, 1990, **8**, 320–338 (rev, *Loline group*)

Naturally Occurring Pyrrolizidine Alkaloids, (ed. Rizk, A.-M.), CRC Press, 1991

Hartmann, Y. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1994, **9**, 155–233 (rev)

Robins, D.J., *Alkaloids*, 1995, **46**, 1–61 (rev, biosynth)

Hartmann, T. *et al.*, *Top. Curr. Chem.*, 2001, 209, 207–244 (*biosynth and metabolism in plants and insects*)

Liddell, J.R., *Nat. Prod. Rep.*, 2002, **19**, 773–781 (*rev*)


Jiang, Y. *et al.*, *Asian J. Pharmacodyn. Pharmacokinet.*, 2006, **6**, 187–192 (*tox*)

SIMPLE PIPERIDINE ALKALOIDS; MISCELLANEOUS PIPERIDINE ALKALOIDS (VX0680, VX0700)

Like the pyrrolidines, these alkaloids may arise by different biosynthetic routes, so the class is structural rather than biosynthetic. Some may be derived from lysine, acetate, acetoacetate, etc., in analogous fashion to the simple pyrrolidine alkaloids. The majority of the alkaloids in this group are plant-derived. **Piperidine** itself is a lysine-derived alkaloid.

All these structural types have their analogues among the pyrrolidine alkaloids, and while it is tempting to assume biosynthesis from lysine it may not in all cases be true; **Coniine**, for example, is acetate-derived. The longer-chain **Batzellasides**, present in sponges, are iminoglycosides structurally related to the intensively-studied terrestrial iminosugars such as **Nojirimycin**, evidently with the incorporation of a fatty-acid derived fragment.

Dioscorine, from *Dioscorea hispida*, provides a fascinating example of the unexpected in alkaloid biosynthesis. At first sight it seems plausible to postulate that it may be formed from lysine and a polyketide fragment. However, lysine is not a precursor, and it would appear that Dioscorine is formed from nicotinic acid and, probably, a polyacetate unit.



Dioscorine

Strunz, G.M. *et al.*, *Alkaloids*, 1985, **26**, 89–184 (*rev*)


Fodor, G.B. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1985, **3**, 1–90 (*rev*)

Schneider, M.J., *Alkaloids Chem. Biol. Perspect.*, 1993, **10**, 155–300 (*rev*)

Andersen, R.J. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1996, **10**, 301 (*rev, 3-alkylpiperidines*)

ANABASINE ALKALOIDS (VX0620)

These are piperidine homologues of the nicotine group (VX0360 above). They also occur in *Nicotiana* spp., in *Anabasis* spp. and others. The biosynthesis is, at least in its major features, analogous to that of the nicotine group, with lysine → cadaverine → *N*-methyl- Δ^1 -piperideine replacing ornithine → putrescine → *N*-methyl- Δ^1 -pyrrolideine. **Anabasine** has been isolated from marine animals and insects, as well as plants.



Anabasine

However, the piperidine ring in **Anatabine**, from *Nicotiana glutinosa*, appears not to be derived from lysine or from a polyacetate precursor; instead, both rings are derived from nicotinic acid.

LOBELIA-TYPE ALKALOIDS (VX0660)

These have no analogy among the pyrrolidine alkaloids. An example is **Lobelanine**.

Fel'pin, F.X. *et al.*, *Tetrahedron*, 2004, **60**, 10127–10153 (*rev*)


XESTOSPONGINS (VX0690)

These are a group of exclusively marine-elaborated piperidines. Their biosynthesis would appear to imply the involvement of a C₁₀ dialdehyde equivalent as hypothesised for the Manzamine group (VX2250) below, and they appear closely related to intermediates postulated in the Baldwin scheme.

Baldwin, J.E. *et al.*, *Tet. Lett.*, 1992, **33**, 2059–2062 (*biosynth*)

LYTHRACEAE ALKALOIDS (VX0760)

These are characterised by several unusual structural features. **Lythrancine** contains a quinolizidine ring system attached to a diphenyl residue, one of the benzene rings in which is derived from cinnamic acid. Other alkaloids in this group contain a diphenyl ether grouping, e.g. **Decaline**, and others a piperidine ring instead of a quinolizidine ring, e.g. **Lythranidine**. These alkaloids have not been studied recently, and their biosynthesis has not been fully worked out. It involves lysine as source of the quinolizidine or piperidine ring, and phenylalanine as precursor of one of the aromatic rings.



The numbering system adopted here for the lactonic Lythraceae alkaloids (e.g. Decaline) is the one generally accepted, and corresponds closely to that for piperidine and quinolizidine metacyclophane alkaloids (e.g. Lythranidine, Lythrancine). The carbon atoms that correspond in biogenetic origin to the three alkaloid types thus maintain corresponding numbers. CAS originally employed stereoparent names for these alkaloids, but now uses systematic nomenclature; in each case the numbering is different.

Golebiewski, W.M. *et al.*, *Alkaloids*, 1981, **18**, 263–273 (rev)




Fuji, K., *Alkaloids*, 1989, **35**, 155–176 (rev)

QUINOLIZIDINE ALKALOIDS (VX0900, VX0920, VX0940, VX0960, VX0980)

Quinolizidine alkaloids are sometimes reviewed alongside pyrrolizidine and indolizidine alkaloids under the joint heading 'Izidines'. The well-known alkaloids which have been long known were isolated from higher plants, particularly the Fabaceae, but the quinolizidine nucleus also occurs in some marine alkaloids and amphibian toxins.

Two-ring quinolizidine alkaloids (VX0900) come from various sources and range from simple bases such as **Epiquinamide**, from a frog toxin, to more elaborated examples such as the marine **Pictamine** and the Fabaceae alkaloid **Epilupinine**, to others which are clearly seco-derivatives of larger alkaloids, e.g. **Cadiamine**. A few relatively simple marine quinolizidine alkaloids are known such as **Pictamine** and the **Clavepictines**. The **Saraines** are more complex sponge products classified under miscellaneous quinolizidines (VX0980), together with **Halichlorine**, a popular synthetic target, and the **Petrosines**, further examples of dimeric alkaloids in which the units are linked in a medium-sized aliphatic ring.

Of the three-ring alkaloids (VX0920) the most characteristic is **Cytisine**; other alkaloids such as **Angustifoline** having a carbon substituent are clearly intermediate between Cytisine and the four-ring alkaloids. The **Tsukushinamines** (which actually have four rings) represent an unusual variation.



The four-ring alkaloids (VX0940) are the most studied group. Fusion of an additional ring onto the cytisine skeleton produces the **Sparteine** type with potentially increased (C_2) symmetry. Both enantiomers of Sparteine and many of its relatives occur naturally, the enantiomers (and sometimes the racemate) often being characterised in the older literature under different names. The symmetry properties mean that the nomenclature and numbering of Sparteine derivatives is confusing, especially since in the past many alkaloids have sometimes been poorly named. Every effort has been made to regularise the nomenclature and numbering, and to show the absolute configurations correctly.

Another type of four-ring alkaloid is represented by **Matrine** and its stereoisomers (**Isomatrine**, **Allomatrine**, **Sophoridine**, **Darvasamine**). These groups contain some dimers, e.g. **Dithermamine**.

The miscellaneous group (VX0980) includes some alkaloids with larger structures, for example **Jamine** and **Panamine** (each six rings).

Howard, A.S., *Alkaloids*, 1986, **28**, 183–308 (rev)

Aslanov, Kh.A., *Alkaloids*, 1987, **31**, 118–193 (rev)

Robins, D.J. *et al.*, *Alkaloids*, 1995, **46**, 1–61 (rev, synth)


Ohmiya, S. *et al.*, *Alkaloids*, 1995, **47**, 2–115 (rev, lupine alkaloids)

Michael, J.P., *Alkaloids*, 2001, **55**, 91–258 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2008, **25**, 139–165 (rev)

CYLINDRICINE ALKALOIDS (VX0960)

The Cylandricine/Fasicularine/Lepadiformine alkaloids (VX0960) are a small cluster of marine alkaloids containing the perhydrobenzo[*i*]quinolizine nucleus and ring-contracted homologues, plus some terrestrial alkaloids which are loosely related structurally.



Fasicularine

Biosynthetic studies do not yet appear to have been carried out. Benzologous structures have been proposed for a small group of terrestrial alkaloids (Menispermaceae) related to **Cohirsine**.

Weinreb, S.M., *Chem. Rev.*, 2006, **106**, 2531–2549 (rev, synth)

AZEPINE ALKALOIDS (VX1000)

These are mostly microbial (or marine, presumably microbial) products; the exceptions are fungal pigments such as **Chalciporone** and its congeners. Marine alkaloids include the **Bengamides**, which are thought to be symbiotic products deriving from bacterial fatty acids, cyclised lysine and a 4-carbon diketide.

NICOTINIC ACID DERIVED ALKALOIDS (VX1020); OTHER PYRIDINE ALKALOIDS (VX1040)

The number of small alkaloids derived by the transformation of nicotinic acid is limited. There are, however, a large number of terpenoid ester alkaloids in which nicotinic acid itself esterifies a variety of terpenoid alcohols, for example dihydro- β -agarofurans (e.g. the **Cangorins** and many others).

Arecoline, from betel nuts, and **Ricinine**, from the castor oil plant, are clearly derivable from nicotinic acid; in the case of Ricinine this has been established.

Strunz, G.M. *et al.*, *Alkaloids*, 1985, **26**, 89–184 (rev)

Fodor, G. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1985, **3**, 1–90 (rev)

Schneider, M.J., *Alkaloids Chem. Biol. Perspect.*, 1993, **10**, 155–300 (rev)

Almeida, A.M.P. *et al.*, *Quim. Nova*, 1997, **20**, 170–185 (rev, alkyipyridines from sponges)


Sepeć, K., *J. Toxicol., Toxin Rev.*, 2000, **19**, 139–160 (rev, alkyipyridinium compounds from sponges)

Liao, L.M., *Alkaloids*, 2003, **60**, 287–343 (rev, sesquiterpene pyridine alkaloids)

9B-AZAPHENALENE ALKALOIDS (VX1120)

(Systematically pyrido[2,1,6-*de*]quinolizines). This small piperidine-based group is exemplified by **Coccinelline**, the defensive agent of the common ladybird, *Coccinella septempunctata*, and **Porantherine**, from the shrub


Poranthera corymbosa. The insect alkaloids appear to be genuine insect metabolites which are biosynthesised from a lipid chain forming the outer perimeter, plus glutamine as the preferred nitrogen source.



Coccinelline

NAPHTHALENE-ISOQUINOLINE ALKALOIDS (VX1140)

This group is isolated mainly from the plant family Ancistrocladaceae, with some isolations from the Dioncophyllaceae. Several skeletal types are known and are based on the point of linkage between the two ring systems, e.g. 5,1'-coupled alkaloids (**Ancistrocladine**, **Dioncophylline C**), 5,8'-coupled (**Korupensamine A**), 7,1'-coupled (**Ancistrocladisine**, **Dioncophylline A**), 7,3'-coupled (**Ancistrocladidine**), 7,6'-coupled (**Dioncophylline B**), etc. These alkaloids are chiral due to diastereoisomerism at the methyl groups and in the biaryl linkage due to restricted rotation. They are biosynthesised via construction of a naphthalenoid portion onto monomeric dihydroisoquinolines.




Ancistrocladine

Bringmann, G. *et al.*, *Alkaloids*, 1986, **29**, 141–184; 1995, **46**, 128–272 (*revs*)

Bringmann, G. *et al.*, *Tetrahedron*, 2007, **63**, 1755–1761 (*biosynth*)

ELAEOCARPUS ALKALOIDS (VX1160)

These are a series of mostly indolizidine-based alkaloids from Far-eastern *Elaeocarpus* spp. A proposed biosynthetic scheme which was suggested in 1969, and which apparently lacks experimental evidence, is through the condensation of a twelve-carbon polyketide chain with a 1-pyrroline derived from ornithine. Alternatively, the whole skeleton may be polyacetate derived.




Elaeocarpiline

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 793–800 (*biosynth*)

Johns, S.R. *et al.*, *Alkaloids*, 1973, **14**, 325–346 (*rev*)

GALBULIMIMA ALKALOIDS (VX1240)

These are a numerically limited class of alkaloids e.g. **Himbacine**, based on the elaboration of a piperidine theme, and which are found in *Galbulimima* (*Himantandra*) spp. (Himantandraceae). A biosynthetic scheme for these alkaloids was suggested in 1967 which proposed their construction from a common polyacetate precursor. Synthetic studies have been carried out to provide a biomimetic route along these lines, however, direct experimental proof is still lacking.




Himbacine

Shah, U. *et al.*, *Total Synthesis of Galbulimima Alkaloids*, VDM Verlag Dr. Muller, Aktiengesellschaft, 2008 (*rev, synth*)


STEMONA ALKALOIDS (VX1260)

These represent a unique type of alkaloid isolated only from the family Stemonaceae (monocots). Three main types are recognised, the **Croomine**, **Stichoneurine** and **Protostemonine** types, although a further subdivision into 11 subtypes can be made. Biosynthetic studies of Stemon alkaloids have not been reported, but a scheme based on a terpenoid origin of the C and D rings and a spermidine origin of the A ring has been proposed by Seger, and an alternative scheme involving a polyketide-geranate condensation by Pyne and co-workers.



Croomine

Stichoneurine A



Protostemonine

Pilli, R.A. *et al.*, *Nat. Prod. Rep.*, 2000, **17**, 117–128 (*rev*)

Seeger, C. *et al.*, *Chem. Biodiversity*, 2004, **1**, 265–279 (*biosynth*)

Pilli, R.A. *et al.*, *Alkaloids*, 2005, **62**, 77–173 (*rev*)


Greger, H., *Planta Med.*, 2006, **72**, 99–113 (*rev*)

Pyne, S.G. *et al.*, *Maejo Int. J. Sci. Technol.*, 2007, **1**, 157–165 (*rev, biosynth*)

LYCOPodium ALKALOIDS (VX1280)

These are constituents of the club mosses (Lycopodiaceae). Whereas the earliest proposal concerning their biosynthesis implicated two C₈ units derived from acetate, it was later established that two lysine units are involved. Numerous skeletal variants are known, all of which can be related to the Lycopodine skeleton; older

examples are **Fawcettidine** and **Serratinine**. More recently some intriguing new skeletons such as those in the **Lycionadines**, **Lyconesidine A**, the **Lycopladines** and **Neohuperzidine** have been found.



Fawcettidine

Maclean, D.B., *Alkaloids*, 1985, **26**, 241–298 (rev)

Blumenkopf, T.A. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1985, **3**, 185–240 (rev)

Ayer, W.A., *Nat. Prod. Rep.*, 1991, **8**, 455–464 (rev)


Ayer, W.A., *Alkaloids*, 1994, **45**, 233–274 (rev)

Ma, X. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 752–772 (rev)

Peng, J. *et al.*, *Alkaloids*, 2005, **61**, 1–57 (rev)

CYTOCHALASAN ALKALOIDS (VX1300)

Cytochalasins are metabolites of several different and unrelated fungi, including some marine actinomycetes. They are characterised structurally by the presence of a perhydroisoindolone system fused to a macrocyclic ring of 11, 13 or 14 atoms. The macrocycle may be a carbocycle, a lactone or a carbonate. In addition, the isoindole ring carries either a phenyl or an indolyl substituent at position 10; the latter group includes the **Chaetoglobosins**. Cytochalasins possess a range of distinctive biological properties; these include inhibition of cytoplasmic cleavage leading to polynucleate cells, nuclear extrusion and the inhibition of cell mobility. Biosynthetically, cytochalasins arise from phenylalanine or tryptophan and a polyketide derived from acetate and methionine. They have not been reviewed recently.




Chaetoglobosin A

Pendse, G.S., *Recent Advances in Cytochalasins*, Chapman & Hall, 1987

INDOLIZIDINE ALKALOIDS (VX1360)

These are a significant class of plant alkaloids, e.g. **Castanospermine**, an important biologically active enzyme inhibitor and a major synthetic target, and **Swainsonine**, biosynthesised from lysine.



Castanospermine

A large number of indolizidine alkaloids known are amphibian skin toxins, of which the greatest number are 5,8-disubstituted, e.g. **Indolizidine 203A**. These are now all known to be of dietary origin, possibly entering the frogs through arachnid and insect prey. Many of these are available only in traces and have currently been characterised only by mass spectroscopic methods, with a number of structures uncertain or incomplete. They are

homologous with the ring-expanded quinolizidine series (see VX0900 above), and other related skeletons by ring enlargement. There are no conclusive studies on the biosynthesis of these related series of amphibian and related toxins, but a polyketide origin is plausible.

A few simple indolizidines (e.g. the **Stellettamides**) have been isolated from sponges.

Howard, A.S., *Alkaloids*, 1986, **28**, 183–308 (rev)

Elbein, A.E. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1987, **5**, 1–54 (rev)

Takahata, H. *et al.*, *Alkaloids*, 1993, **44**, 189–256 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2004, **21**, 625–649; 2005, **22**, 603–626; 2008, **25**, 139–165 (revs)

Daly, J.W. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1556–1575 (*amphibian toxins*)


Amat, M. *et al.*, *Angew. Chem. Internat. Ed. Engl.*, 2008, **47**, 3348–3351 (*amphibian toxins*)

SIMPLE ANTHRANILIC ACID ALKALOIDS (VX1460)

A number of diverse structural groups of alkaloid are derived biogenetically from anthranilic acid, the major ones being the quinolone, acridine and quinazoline groups categorised under their respective headings. The number of unelaborated anthranilic acid alkaloids is small and includes **Damascenine** from *Nigella damascena*. Biosynthetically, the alkaloid is derived from anthranilic acid (chorismate derived), which is then hydroxylated and methylated.

MAYTANSINOIDS (VX1470)

The maytansinoids are a group of ansa macrocyclic lactams originally isolated from three plant families (Euphorbiaceae, Celastraceae, Rhamnaceae), and subsequently isolated from a soil bacterium *Actinosynnema pretiosum*, other microorganisms and mosses. They are homologous with the **Ansamitocins** from microorganisms, and may be derived from plant-associated fungi. They show strong antitumour activity and are biosynthesised from the simple aromatic precursor 3-amino-5-hydroxybenzoic acid, arising from the chorismate pathway, which is substituted by a polyketide chain and then amide ring formation.



Cassady, J.M. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1–26 (rev)

SIMPLE QUINOLINES (VX1480); FURANOQUINOLINES (VX1520); PYRANOQUINOLINES (VX1540); MISCELLANEOUS QUINOLINE ALKALOIDS (VX1580)

The majority of simple quinoline alkaloids are from higher plants, but a variety of simple heteroaromatic quinolines (VX1480) have been isolated from various marine sources, including **2-Heptyl-4-hydroxyquinoline** from a marine pseudomonad, and **4,8-Quinolinediol** from cephalopod ink. Plant alkaloids include **Echinopsine** and the phenethyl-quinoline, **Cusparine**. The plant-derived furanoquinoline (VX1520) (e.g. **Platydesmine**), and pyranoquinoline (VX1540) alkaloids (e.g. **Flindersine**) are straightforwardly derived from simple alkylquinolines (especially prenyl-substituted) by cyclisation. There are also more elaborated hydrogenated quinolines, especially the **Lepadins** from tunicates and flatworms.

Bryozoans contain a number of simple quinoline-quinone pigments and there are also the **Trididemnic acids** from ascidians. Simple hydrogenated quinolines also occur in amphibian toxin mixtures, for example the numerous **Dendrobates Alkaloids**. This extensive series of alkaloids occurs in amphibian skin extracts and in tunicates, flatworms and insects (myrmicine ants), in a mixture which includes hydrogenated quinolines, and structural variants such as indolizidines (VX1360, see for further information).

Macrorine, from *Macrorungia longistrobus*, is obviously derived from anthranilic acid and histidine.


Grundon, M.F., *Alkaloids Chem. Biol. Perspect.*, 1988, **6**, 339–522 (rev)

Fernandes da Silva, M.F. *et al.*, *Alkaloids*, 2007, **64**, 139–214 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2008, **25**, 166–189 (rev)

QUINAZOLINE ALKALOIDS (VX1600)

These include **Vasicine**, first isolated in 1888 from *Adhatoda vasica*, and **Rutaecarpine**, from *Euodia rutaecarpa*; this latter base is clearly derived from tryptophan and anthranilic acid. Quinazoline alkaloids have been isolated from several plant families as well as from microorganisms, fungi and animal spp. **Febrifugine** is a pharmacologically important (antimalarial) member of the group. A number of simple quinazolines also occur, as well as **Bouchardatine**, which is a quinazoline-indole dimer. They have been well reviewed; see also Tryptoquivalines (VX6030) below.



Vasicine

Johne, S., *Prog. Chem. Org. Nat. Prod.*, 1984, **46**, 159–218 (rev)


Johne, S., *Alkaloids*, 1986, **29**, 99–140 (rev)

Mhaske, S.B., *Tetrahedron*, 2006, **62**, 9787–9826 (rev)

Michael, J. *et al.*, *Nat. Prod. Rep.*, 2008, **25**, 166–187 (rev)

ACRIDONE ALKALOIDS (VX1620-VX1690)

These are higher plant alkaloids characteristic of the Rutaceae, and are biosynthesised from anthranilic acid by elaboration of an additional malonate-derived aromatic ring. The simple acridones (VX1620) may be exemplified by **Xanthevodine** and by a large number of alkaloids in which the nucleus is prenylated followed frequently by formation of a furano- or pyranoacridine ring system, e.g. **Acronycine**. Biosynthesis of the nucleus is from anthranilic acid and a malonate fragment. The **Acrimarines** are acridone-coumarin dimers (VX1690).



Acronycine

Gerzon, K. *et al.*, *Alkaloids*, 1983, **21**, 1–28 (rev)

Grundon, M.F., *Alkaloids Chem. Biol. Perspect.*, 1988, **6**, 339–522 (rev)


Tillequin, F. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1998, **12**, 1–102 (*Acronycines*, rev)

Skaltsounis, A. *et al.*, *Alkaloids*, 2000, **54**, 259–377 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2008, **25**, 166–187 (rev)

PYRIDO[2,3,4-*KL*]ACRIDINES (VX1700)

This is a group of exclusively marine alkaloids which includes the **Kuanoniamines**, the **Ascididemins**, the **Cystodytins**, the **Styelsamines**, the **Varamines** and others. The isolation of some of them from a variety of organisms argues for a microbial origin, but it could alternatively be a case of convergent evolution. They are biosynthesised from common amino acids.




Styelsamine A

Molinski, T.F., *Chem. Rev.*, 1993, **93**, 1825–1838 (rev)
Ding, Q. *et al.*, *Curr. Med. Chem.*, 1999, **6**, 1–27 (rev)
Salomon, C.E. *et al.*, *Mar. Biol. (Berlin)*, 2001, **139**, 313–319 (biosynth)
Moore, B.S., *Nat. Prod. Rep.*, 2006, **23**, 615–629 (rev)


1,4-BENZOXAZIN-3-ONE ALKALOIDS (VX1720)

2,4-Dihydroxy-7-methoxy-2H-1,4-benzoxazin-3(4H)-one (known as DIMBOA) is the main representative of this group of alkaloids which have plant hormone significance. DIMBOA itself protects the maize plant against corn borer attack. They usually occur in the plant as glucosides, but cell injury releases a glucosidase which catalyses hydrolysis to the 2-hydroxy derivatives. All four oxygen atoms in DIMBOA arise by incorporation of molecular O₂.




BENZODIAZEPINE ALKALOIDS (VX1760)

This is a growing class of microbial metabolites typified by **Anthramycin** and others. One or two, e.g. **Aplysepine**, are of marine origin; apparently no plant alkaloids with this skeleton are known to date. They are biosynthesised from amino acids via kynurenine and its derivatives.



CRYPTOLEPINE ALKALOIDS (VX1800)


Cryptolepine (5-Methyl-5H-quindoline) is the most prominent representative of this small group isolated from *Cryptolepis* spp. The parent alkaloid, 10H-Indolo[3,2-b]quinoline (**Quindoline**), has been isolated from *Cryptolepis sanguinolenta*. Two isomeric skeletons (**Isocryptolepine**, **Neocryptolepine**) were also obtained. A further variant represented by **Cryptoheptine** is doubtful. The marine **Communesins** (currently classified under miscellaneous alkaloids VX9400) contain the Neocryptolepine skeleton embedded in a polycyclic ring system. No biosynthetic studies of these skeletons have been carried out, but a derivation from indole and anthranilic acid can be envisaged.



SIMPLE PHENETHYLAMINE ALKALOIDS (VX2000, VX2005, VX2008, VX2015)

Alkaloids derived from phenylalanine or tyrosine represent an extremely large and varied major subdivision of the alkaloids, ranging from simple β -phenylethylamine derivatives to the much more complex structures exemplified by the alkaloids of the Amaryllidaceae and the very numerous isoquinoline alkaloids.

The simplest derivatives of phenylalanine or tyrosine are the β -phenylethylamines, which are presumably obtained by decarboxylation and obvious oxidative/alkylation stages, and which are widespread among plant alkaloids (e.g. **Mescaline**) and elsewhere. Compounds showing side-chain oxygenation (VX2005) include alkaloids from higher plants, such as **Macromerine**, as well as animal hormones such as **Adrenaline** and **Octopamine**, which is an endometabolic neurotransmitter of molluscs. Brominated compounds (VX2008) such as the **Purpuramines** and the **Bastadins** are common in marine organisms. These are biosynthesised from the simple brominated tyrosines.



Lundström, J., *Alkaloids*, 1989, **35**, 77–154 (rev)

Peng, J. *et al.*, *Alkaloids*, 2005, **61**, 59–262 (rev, marine bromotyrosine alkaloids)

Bentley, K.W., *Nat. Prod. Rep.*, 2006, **23**, 444–463 (rev)

EPHEDRA ALKALOIDS (VX2010)

This is a small, and important, group of low molecular weight alkaloids centred on **Ephedrine** and its relatives, isolated from *Ephedra* spp. (Ma Huang) and showing strong physiological activity. They carry a side-chain amino or substituted amino substituent. This structural feature is common in higher molecular weight alkaloids.

Ma Huang and the Ephedra Alkaloids, (ed. Karch, S.B.), Springer/Humana Press, 2007 (book)


CINNAMIC ACID AMIDES (VX2020)

cis- and *trans*-**Fagaramide**, **Herclavine** and **Subaphylline** are simple members of this series, which are mostly isolated from terrestrial plants. These amides are typically derived from a *trans*-cinnamic acid; **Astrophylline** is an example of an *N-cis*-cinnamoyl derivative. Simple cinnamate residues are scarcer among marine alkaloids. Examples are found in the **Tunichromes**, for which phenylalanine can function as a biosynthetic precursor.

Taylor, S.W. *et al.*, *Chem. Rev.*, 1997, **97**, 333–346 (rev, tunichromes)

SECURINEGA ALKALOIDS (VX2100)


This small group of polycyclic alkaloids occurs in the Euphorbiaceae. The biosynthesis of these alkaloids, which is not readily apparent, has been shown to involve tyrosine and lysine in the case of **Securinine**. Alkaloids of both enantiomeric series occur naturally, sometimes in closely related plants.



Snieckus, V., *Alkaloids*, 1973, **14**, 425–506 (rev)

BETALAIN ALKALOIDS (VX2140)

This is a group of zwitterionic water-soluble alkaloidal pigments whose distribution is limited to the order Caryophyllales of the dicots, (more or less coterminous with the older term Centrospermae), for which they have been used as a taxonomic marker, plus some basidiomycete fungi. **Betanin** (the red pigment of beetroot) is the best-known example, other examples include various flower pigments instead of anthocyanins. Both halves of the molecule are biosynthesised from DOPA/Dopamine via Dopaxanthin; the formation of the upper unit, Betalamic acid, involves an enzymatic aromatic ring fission process.



Stintzing, F. *et al.*, *Trends Food Sci. Technol.*, 2007, **18**, 514–525 (rev)

ISOQUINOLINE ALKALOIDS (VX2200-VX2400)

This is an extremely large group of mainly higher plant alkaloids, although one or two groups of isoquinolinoid marine alkaloids, notably the Manzamine group (VX2250) are also interpolated into the listing.

Shamma, M., *The Isoquinoline Alkaloids*, Academic Press, New York, 1972

Shamma, M. *et al.*, *Isoquinoline Alkaloid Research, 1972–1978*, Plenum Press, New York, 1978

Krane, B.D., *J. Nat. Prod.*, 1982, **45**, 377–384 (rev)

Lundström, J., *Alkaloids*, 1983, **21**, 255–328 (rev)

Phillipson, J.D. *et al.*, *The Chemistry and Biology of Isoquinoline Alkaloids*, (eds. Phillipson, J.D. *et al.*), Springer, 1985

Menachery, M.D., *J. Nat. Prod.*, 1986, **49**, 745–778 (rev)

Baker, B.J. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1996, **10**, 357–407 (rev, marine isoquinolines)


Bentley, K.W., *Nat. Prod. Rep.*, 2006, **23**, 444–463 (rev, revs cited therein)

SIMPLE ISOQUINOLINE ALKALOIDS (VX2200)

This group includes simple isoquinolines with (e.g. **Salsoline**) or without (e.g. **Anhalamine**) a C-1 substituent, and various quinones (**Mimosamycin**, **Renierone**). Marine examples are restricted to some bases related to **Renierol** and the **Cribrostatins**.

MANZAMINE ALKALOIDS (VX2250)

This fairly extensive group (a typical representative is **Manzamine A**) is exclusively of marine origin and represents the most structurally complex and characteristic type of marine alkaloid for which a biosynthetic scheme has been proposed. Their biosynthesis has been postulated to involve the condensation of tryptamine with one or more C₃ units and one or more straight-chain C₁₀ dialdehyde equivalents, followed by an enzyme-assisted Diels-Alder cyclisation. The origin of the C₁₀ and C₃ units is uncertain, but it has been suggested that they could arise by oxidative fission of a hydroxylated fatty acid. The full biosynthetic scheme is shown in the reviews below and more elaborately in the companion *Dictionary of Marine Natural Products*.



Manzamine A

- Baldwin, J.E. *et al.*, *Tet. Lett.*, 1992, **33**, 2059–2062 (*biosynth*)
Tsuda, M. *et al.*, *Heterocycles*, 1997, **46**, 765–794 (*rev*)
Baldwin, J.E. *et al.*, *Chem. Eur. J.*, 1999, **5**, 3154–3161 (*biosynth*)
Hu, J.F. *et al.*, *Alkaloids*, 2003, **60**, 207–285 (*rev*)
Thomas, R., *Nat. Prod. Rep.*, 2004, **21**, 224–248 (*rev, biosynth*)

BENZYLISOQUINOLINE ALKALOIDS (VX2320)

The simple benzylisoquinoline skeleton is derived from two molecules of tyrosine, via DOPA, and is the precursor of a wide variety of alkaloids in categories listed below, belonging to numerous different ring systems. The alkaloids with the unmodified benzylisoquinoline skeleton consist mainly of 1,2,3,4-tetrahydrobenzylisoquinolines, e.g. **Reticuline**, of central importance in the elaboration of other alkaloids, and their fully aromatic analogues, e.g. **Papaverine**. Alkaloids with a carbon substituent at C-2', such as **Canadaline**, may be regarded as ring-opened berberines. There are also some benzylisoquinolines in which a pyrrolidine ring is attached to C-4, e.g. **Macrostromine**. The isolation of one or two marine benzylisoquinoline alkaloids (**Imbricatine**; **Theoneberine**) is noteworthy.

- Deulofeu, V. *et al.*, *Alkaloids*, 1968, **10**, 401–461 (*rev*)

PSEUDOBENZYLISOQUINOLINE ALKALOIDS (VX2330)

The term pseudobenzylisoquinoline alkaloid is used to describe a benzylisoquinoline skeleton in which the pendant aromatic ring is oxygenated at C-2', C-3' and C-4'. These alkaloids are derived biogenetically from protoberberinium salts by C8-C8a bond scission. **Polycarpine**, **Taxilamine** and **Ledecorine** are typical examples.

BISBENZYLISOQUINOLINE ALKALOIDS (VX2340–VX2400)

This large group of alkaloids is composed of two benzylisoquinoline units attached to each other by one, two, or three bonds. In most cases the units are joined via ether linkages, but carbon-carbon bonds between the benzyl groups are also known. The monomeric units involved are mainly hydroxylated or methoxylated benzylisoquinolines, although aporphine units occur in many alkaloids, and a few contain a proaporphine component. The alkaloids may be subdivided into the five following major groups (the classification, proposed by Shamma, contains at least 28 subgroups). All alkaloids classified here are composed of unmodified benzylisoquinoline units. Dimeric alkaloids containing aporphine, proaporphine, or otherwise modified benzylisoquinoline components are discussed under separate headings).


CAS used to employ a number of stereoparent skeletons, such as Berbaman, Thaliceran, Tubocuraran, in naming the cyclic bisbenzylisoquinoline alkaloids, but these have now been replaced by systematic nomenclature. Trivial nomenclature is given prominence in this Dictionary, but the main semisystematic parent skeletons are

shown below. Care is needed in numbering the two benzyloquinoline residues. In a few cases, CAS has in the past applied the primes to the opposite residue to that proposed in the Shamma-Moniot convention. The latter is used in this Dictionary.

The Shamma-Moniot convention is as follows;

1. The numbering system for a bisbenzyloquinoline half of the dimer is as shown below. The lower aromatic ring, namely ring C, is always numbered so as to assign the smallest numbers to the substituents on that ring
2. Each bisbenzyloquinoline half of the dimer is described in terms of its oxygenation pattern. The more highly oxygenated bisbenzyloquinoline half constitutes the left hand side of the dimer. The right hand side of the dimer is given prime numbering.

For an example see the structure of Hernandezine below.



Hernandezine with Shamma-Moniot convention numbering

Shamma, M. *et al.*, *Heterocycles*, 1976, **4**, 1817–1824 (*Shamma-Moniot numbering convention*)

Buck, K.T., *et al.*, *Alkaloids*, 1987, **30**, 1–222 (*rev*)

Schiff, P.L., *J. Nat. Prod.*, 1997, **60**, 934–953 (*rev, revs cited therein*)

BISBENZYLISOQUINOLINE ALKALOIDS CONTAINING ARYL LINKS ONLY (VX2340)

The bark of *Popowia pisocarpa* has yielded a group of alkaloids which contain a single aromatic linkage between C-11 and C-11'. See the entry for **Pisopowetine**. **Isopythaldine** and **Isopyruthaldine** represent another type in which the two benzyloquinoline groups are joined by methylene bridges.

BISBENZYLISOQUINOLINE ALKALOIDS CONTAINING ONE ETHER LINK (VX2360)

The ether linkage is in most cases between C-11 and C-12', as in **Dauricine**. However, other attachments are known, e.g. between C-10 and C-11', as in **Vanuatine**, between C-10 and C-7', as in **Malekulatine** and **Ambrimine**, and between C-11 and C-7', as in **Neferine**.

BISBENZYLISOQUINOLINE ALKALOIDS CONTAINING ONE AROMATIC LINK AND ONE OR TWO ETHER LINKS (VX2370, VX2390)

These alkaloids are mainly based on the **Rodiasine** and **Tiliacorine** skeletons.

BISBENZYLISOQUINOLINE ALKALOIDS CONTAINING TWO ETHER LINKS (VX2380)

These are the most numerous. The largest subgroup containing two ether linkages possesses the berbaman skeleton, exemplified by **Berbamine**, which contains ether linkages between C-8 and C-7', and between C-11 and C-12'. Almost as large as the berbaman group is the oxyacanthan group, e.g. **Oxyacanthine**, which contains ether linkages between C-7 and C-8', and between C-11 and C-12'. Smaller groups include the **Thalicerine** (C-8 to C-6')


and C-11 to C-12'), **Thalidasine** (C-8 to C-5' and C-11 to C-12'), and **Thalmine** (C-7 to C-5' and C-11 to C-12') types. All of these types contain ether linkages between the benzyl rings and between the aromatic rings of the tetrahydroisoquinoline component. The **Tubocurarine** sub-group contains ether linkages between the benzyl ring of one unit and the aromatic ring of the isoquinoline component of the other. Other linkages between benzyl and isoquinoline rings are also known.

BISBENZYLISOQUINOLINE ALKALOIDS WITH THREE ETHER LINKS (VX2400)

These include alkaloids with 6',7'-epoxyoxyacanthan (e.g. **Trilobine**), 7,8'-epoxyoxyacanthan, and 8,12'-epoxytubocuraran skeletons.

BENZYLISOQUINOLINE-APORPHINE DIMERS (VX2700)

Some alkaloids consist of a benzyloisoquinoline unit attached to an aporphine unit, through a single ether linkage. Of these, **Thalicarpine** is typical.




Miscellaneous bisbenzyloisoquinoline-related alkaloids (listed under their component monomeric units) include those containing a dienone ring in one of the isoquinoline moieties (e.g. **Repanduline**), some with an aporphine unit attached to a pavine component (e.g. **Pennsylvavine**), those with degraded benzyloisoquinoline units (e.g. **Baluchistanamine**) or with a proaporphine unit (e.g. **Epiberbivaldine**), and **Cancertrine**, which is really a combination of cularine and morphinan components.

Buck, K.T. *et al.*, *Alkaloids*, 1987, **30**, 1–222 (*rev*)

Schiff, P.L., *Alkaloids Chem. Biol. Perspect.*, 1987, **5**, 271–638; 2000, **14**, 1–234 (*revs*)

SECOBISBENZYLISOQUINOLINE ALKALOIDS (VX2430)

The secobisbenzyloisoquinolines are degraded alkaloids in which one of the benzyloisoquinoline units is cleaved between the C-1 and the α -carbon atom. A typical example is **Baluchistanamine**, the apparent biogenetic precursor of which is Oxyacanthine. Different oxidation states are found; aldehyde lactams (e.g. **Baluchistanamine**, **Punjabine**), lactam esters (e.g. **Punjabine**, **Curacautine**) or aldehyde amines (**Jhelumine**, **Chenabine**).




Like the bisbenzyloisoquinoline precursors, these alkaloids differ in the number and position of the diphenyl ether linkages. Karakoramine has only one such bond, Baluchistanamine has two, and Punjabine has three. **Secantioquine** and **Secolucidine** are examples of a biphenyl system.

CULARINE ALKALOIDS (VX2440)

The cularines are tetracyclic isoquinoline alkaloids which contain a dihydrooxepine or oxepine ring between C-8 and C-2'. They are formed by intramolecular oxidative coupling of 7,8,3',4'- or 7,8,4',5'-tetraoxygenated

tetrahydrobenzylisoquinolines (classical cularines and isocularines, respectively), although the biogenesis of **Gouregine** probably proceeds via oxidation of an aporphine precursor. Note the parallelism between the cularine and aporphine alkaloids.



A small number of **Cacentrine**-type alkaloids are known which are cularine-morphinan dimers.

Castedo, L. *et al.*, *Alkaloids*, 1986, **29**, 287–324 (*rev*)

SECOCULARINE ALKALOIDS (VX2450)


The secocularines can be divided into two sub-groups, namely B- and C-ring seco compounds. B-Ring secocularines, exemplified by **Secocularidine**, are structurally related to phenanthrene alkaloids derived from aporphines and are probably formed *in vivo* by Hofmann degradation of cularines. C-Ring cularines, represented by **Noyaine**, constitute a type of alkaloid without counterpart among aporphinoids.

INDENOBENZAZEPINE ALKALOIDS (VX2500)

This is a small group, e.g. **Lahorine**, which may be derived from the spirobenzylisoquinolines (VX3220 below).

QUETTAMINE ALKALOIDS (VX2470)

Whereas the classical-type cularine alkaloids of the Fumariaceae are biogenetically derived from intramolecular oxidative coupling of tetraoxygenated tetrahydroisoquinoline precursors, the quettamines are obtained from *in vivo* intramolecular oxidation of a trioxygenated tetrahydroisoquinoline. So far, few naturally occurring quettamines are known. These alkaloids, all found in *Berberis baluchistanica*, incorporate either a benzofuran or a dihydrobenzofuran ring within the molecular framework, and the seco bases possess an *N,N*-dimethylaminoethyl substituent.




DIBENZOPYRROCOLINE ALKALOIDS (VX2480)

These alkaloids, only a few of which are known, are clearly derived by oxidation of a benzylisoquinoline precursor; indeed, the ring system was prepared *in vitro* by this route before it was encountered in nature. **Oubatchensine** is a 7,13-seco alkaloid presumably derived through 13-hydroxylation and tautomerism to the open-chain form.

Elliot, I.W., *Alkaloids*, 1987, **31**, 101–117 (*rev*)

PAVINE AND ISOPAVINE ALKALOIDS (VX2520, VX2540)

These alkaloids are formed by alternative modes of oxidative cyclisation of benzylisoquinoline precursors. In addition there are bisbenzylisoquinoline alkaloids composed of pavine and aporphine units, e.g. **Pennsylvavine**.




Because of the two-fold rotational symmetry of the pavane skeleton, there are numerous variations in numbering schemes in the literature. Numbering in this Dictionary has been regularised according to IUPAC principles.

Some mixed pavinoind-benzylisoquinoline and pavinoind-aporphine dimers (**Hervelines**, **Fauripavine**) have been isolated relatively recently.

Gözler, B. *et al.*, *Alkaloids*, 1987, **31**, 317–390 (*rev*)

PROAPORPHINE ALKALOIDS (VX2600)

This group of alkaloids, e.g. **Orientalinone**, represents an intermediate stage in the conversion of the benzylisoquinoline alkaloids by phenol oxidative coupling into the aporphines. The spirocyclohexenone ring may occur in various oxidation levels from cyclohexadienone to cyclohexanol. There are various numbering schemes in the literature. That shown has been adopted here as standard.



There are also proaporphine-tryptamine dimers centred on **Roehybridine**. These heptacyclic alkaloids, found in *Roemeria hybrid* (Papaveraceae) and *Phoebe grandis* (Lauraceae), are probably derived biogenetically by a Mannich-type condensation of a ketonic tetrahydroproaporphine with a tryptamine analogue.

PROAPORPHINE-BENZYLISOQUINOLINE ALKALOID DIMERS (VX2620)

This group is self-explanatory, e.g. **Pakistanamine**.

APORPHINE ALKALOIDS (VX2640, VX2700)


This large group of alkaloids contains the tetracyclic ring system formed by the phenol oxidative coupling of a benzylisoquinoline precursor. The structural variations among the numerous simple aporphines (VX2640) are mostly 6a,7-dehydrogenation and 7-oxidation, accompanied by isoquinoline-ring aromatisation in the so-called oxoaporphines, e.g. **Liriodenine** (VX2640). To this listing must be added a small group of bisaporphines. The majority of these alkaloids possess a carbon-to-carbon linkage between C-7 and C-7' (e.g. **Urabaine**), although examples of C-C coupled bisaporphines and oxygen-bonded dimers (e.g. **Dehatriphine**) are also known. There is a considerable number of aporphine-benzylisoquinoline dimers listed separately (VX2700).

Various alkaloid groups which follow (VX2750-VX2840) may be considered as closely related to the true aporphines and are often reviewed alongside them.

Guinaudeau, H. *et al.*, *Lloydia*, 1975, **38**, 275–338; *J. Nat. Prod.*, 1979, **42**, 133–149, 325–360; 1983, **46**, 761–835; 1984, **47**, 565–580; 1988, **51**, 389–474, 1025–1053; 1994, **57**, 1033–1135 (*tabulations*)
Shamma, M., *Nat. Prod. Rep.*, 1986, **3**, 345–352 (*rev*)
Gözler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 675–685 (*aporphine dimers*)

OXOISOAPORPHINES (VX2750)


This is a small group of alkaloids, mostly from *Menispermum* spp. It has been suggested that they arise from aporphines via a spirodienone rearrangement, however, biosynthetic studies have not been reported. One of its representatives is **Menisporphine**. **Lakshminine**, from *Sciadoteria* sp., in the Menispermaceae, contains an additional nitrogen substituent.



Menisporphine

ARISTOLOCHIC ACIDS AND ARISTOLACTAMS (VX2780, VX2800)

The aristolochic acids represent a class of phenanthrenoid alkaloid derived from an aporphinoid precursor with ring-B fission and the loss of C-5 (aporphine numbering). **Aristolochic acid VIa** is a typical alkaloid. Some of the alkaloids are esters of these acids with, for example, terpenoids, e.g. **Aristolin**. They are higher plant products characteristic of the Aristolochiaceae. Since they have a seco-aporphinoid skeleton, they have been numbered in the literature in different ways, either as aporphines or as phenanthrenes, and for many of them (such as those with a methylenedioxy function), the CAS systematic numbering is different again. In this Dictionary, the simpler members are named and numbered as substituted phenanthrenes, and different alkaloid groups are collected into entries under the parent phenanthrene.



Aristolochic acid VIa


The biosynthetically-related Aristolactams, mostly nonbasic, have a five-membered lactam ring e.g. **Cepharanone A**.

These classes of alkaloid also include some members in which nitrogen has been oxidised to a nitro group, e.g. **Aristolochic acid A**.

Bick, I.R.C. *et al.*, *Alkaloids*, 1985, **24**, 113–152 (*rev*)
Chen, Z., *Alkaloids*, 1987, **31**, 29–66 (*rev*)
Kumar, V. *et al.*, *Nat. Prod. Rep.*, 2003, **20**, 565–583 (*rev*)

PHENANTHRENE ALKALOIDS (VX2820)

These alkaloids, e.g. **Thalicpureine**, are related to aporphines by fission of the *N*-6a bond. The group includes a few alkaloids showing oxidative changes elsewhere in the molecule (**Andesine**; **Chiloenine**; **Santiagoamine**). **Eupolauramine** is related by a further ring-contraction to a lactam.




Thalicipureine

Taylor, W.C., *Aust. J. Chem.*, 1984, 37, 1095–1104 (*biosynth*)

Castedo, L. *et al.*, *Alkaloids*, 1990, 39, 99–138 (*rev*)

AZAAPORPHINE ALKALOIDS (VX2830)


This fairly small group, e.g. **Sampangine**, is thought to arise by A-ring fission of an aporphinoid precursor to an azaanthraquinone intermediate, followed by ring closure.



Sampangine

AZAFLUORANTHENE ALKALOIDS (VX2840)

This is another small group of alkaloids found in the Menispermaceae. Two biosynthetic mechanisms have been proposed; cyclisation of a 1-arylisquinoline (which is considered unlikely since 1-arylisquinolines are rare) or a more persuasive route involving the decarbonylation of an oxoaporphine. **Imerubrine** and its relatives are then related by an aryl→tropolone expansion. **Eupolauridine** could arise by a combination of this route and the aza-replacement shown above under VX2830.




Imerubrine

Buck, K.T., *Alkaloids*, 1984, 23, 301–327 (*rev*)

Taylor, W.C., *Aust. J. Chem.*, 1984, 37, 1095–1104 (*biosynth*)

MORPHINE ALKALOIDS (VX2900)


This extremely important group is formed by phenol oxidative coupling of a hydroxylated benzyloquinoline precursor such as Norlaudanoline, itself originating from two molecules of tyrosine. The group may be subdivided into bases of the **Salutaridine** type, those related to **Morphine**, which have a 4,5-oxide bridge, and those related to **Sinoacutine**, which are enantiomeric with the salutaridine group. **Acutumine** and its congeners represent an *abeo* shift to a spiro skeleton.



Szantay, C. *et al.*, *Alkaloids*, 1994, **45**, 128–232 (rev)

ERYTHRINA ALKALOIDS (VX2940)

This group of alkaloids is derived from two tyrosine units by oxidative coupling and intramolecular rearrangement. Alkaloids which contain the erythrinan skeleton (e.g. **Erythraline**) constitute the majority.



Major structural variations consist of oxidative modification of the aromatic ring to a lactone, e.g. **β -Erythroidine**, and a small group containing an additional nitrogen atom in ring D. These 16-azaerythrinanes, e.g. **Erymelanthine**, are possibly derived biogenetically by *in vivo* oxidation of the aromatic ring, followed by nitrogen uptake and recyclicalisation. There are also a few so-called dimeric alkaloids that incorporate a tryptophan moiety, e.g. **Eryspinophorine**.

Dyke, S.F. *et al.*, *Alkaloids*, 1981, **18**, 1


Chawla, A.S. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1991, **9**, 85–154 (rev)

Amer, M.E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 329–363 (rev)

Tsuda, Y. *et al.*, *Alkaloids*, 1996, **48**, 249–337 (rev)

DIBENZAZECINE AND HASUBANAN ALKALOIDS (VX2980, VX3000)

These two groups may appear at first sight to belong to quite different structural groups, but there is little doubt that biosynthetically they are related. Both groups are derived from two tyrosine units, but their biosynthesis is not simple. A pivotal intermediate appears to be a hydroxylated benzyloquinoline which can cyclise, by alternative phenol oxidative coupling processes, to a hydroxysalutaridine, or its isomer. This biosynthetic route is exceptional since it would appear that two hydroxy-groups need to be present in one of the aromatic rings. In all other known cases of oxidative coupling, only one hydroxy-group seems to be essential. The biosynthesis has affinities with that of the morphine alkaloids, although there is clearly an important divergence in the later stages.




Inubushi, Y. *et al.*, *Alkaloids*, 1977, **16**, 393–340 (*Hasubanans*, rev)

Matsui, M., *Alkaloids*, 1988, **33**, 307–347 (*Hasubanans*, rev)

PROTOBERBERINE ALKALOIDS (VX3100, VX3240)

These tetracyclic alkaloids are derived from benzyloquinolines by condensation with a one-carbon unit (the berberine bridge), the additional carbon atom coming from the methionine-derived C₁ pool. The alkaloids most closely related to benzyloquinolines are the simple tetrahydropROTOBERBERINES, e.g. **Tetrahydropalmatine**, which are dehydrogenated to Protoberberines, such as **Berberine**. There are also alkaloids carrying extra carbon substituents at C-12 and C-13 such as **Orientalidine** and **Corydaline**, respectively, and ring-opened protoberberines (secoberberines) which can be regarded as benzyloquinolines with a carbon substituent at C-2'. The latter may occur in different oxidation states: as an aldehyde (e.g. **Aobamine**), an alcohol (e.g. **Macrantaline**) or as a carboxylic acid (e.g. **Macrantoridine**).



Berberine

Structure determination in this series, i.e. the assignment of correct regiochemistry to the substituents on the protoberberine ring system, has been a matter of some difficulty in the past and it is possible that some of the current assignments will prove to be incorrect.

Jeffs, P.W., *Alkaloids*, 1967, **9**, 41–115 (rev)

Santavy, F., *Alkaloids*, 1979, **17**, 439–461 (rev)

Bhakuni, D.S. *et al.*, *Alkaloids*, 1986, **28**, 95–181 (rev)

Beecher, C.W.W. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1988, **6**, 297–337 (rev)

Leitao da Cunha, E.V. *et al.*, *Alkaloids*, 2005, **62**, 1–75 (rev)


Grycova, L. *et al.*, *Phytochemistry*, 2007, **68**, 150–175 (rev)

ISOINDOLOBENZAZEPINE ALKALOIDS (VX3130)

This is a small group, one example is **Lennoxamine**, which is a popular synthetic target.

NARCEINE ALKALOIDS (VX3140)

These alkaloids constitute further examples of oxidation products of protoberberines, in which the nitrogen to C-8 bond has been cleaved. The narceine group contain an ethanamine chain and, as well as relatives of **Narceine**, include bases which contain an enol lactone function, and those with a higher (**Bicucullinine**) or lower (**Peshawarine**) oxidation level than **Narceine**.




Narceine

Blaskó, G. *et al.*, *J. Nat. Prod.*, 1982, **45**, 105–122 (rev)

PROTOPINE ALKALOIDS (VX3160)

These tricyclic bases are simply formed by oxidative ring fission of protoberberine N-metho salts. Two of these bases (**Corycavamine**, **Corycavidine**) have an additional methyl group at C-13.




Corycavamine

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1982, **45**, 237–246 (*rev*)

Onda, M. *et al.*, *Alkaloids*, 1988, **34**, 181–209 (*rev*)

RHOEADINE ALKALOIDS (VX3180)

This group of alkaloids has been encountered only in *Papaver* and related genera. Their biogenesis appears to be from two tyrosine units, via tetrahydroberberine and protopine intermediates. Oxidative fission of the nitrogen to C-8 bond followed by oxidative cyclisation of nitrogen on to C-15 and lactol formation results in a ring system in which C-8 becomes the lactol carbon atom.




Rhoeadine

CAS formerly employed the semisystematic skeleton Rheadan for naming these alkaloids, but this has been abandoned in favour of systematic nomenclature (Rheadan = [2]Benzopyrano[3,4-*a*][3]benzazepine).

Rönsch, H., *Alkaloids*, 1986, **28**, 1–94 (*rev*)

PHTHALIDEISOQUINOLINE ALKALOIDS (VX3200)

These are closely related to the rhoeadine group, and like them are formed in nature by oxidation of tetrahydroprotoberberines; **Hydrastine** and **Narcotine** are specifically derived from Reticuline via Scoulerine.




Hydrastine

MacLean, D.B., *Alkaloids*, 1985, **24**, 253–286 (*rev*)

SPIROBENZYLISOQUINOLINE ALKALOIDS (VX3220)

These alkaloids are derived from protoberberines by a 1,2-shift of C-8 from nitrogen to C-14. Several mechanisms are possible, but since several of the alkaloids contain oxygen at C-13 a route through an enolate ylid is attractive. **Ochotensine** and its relatives contain an additional carbon atom, again derived from methionine. Closely related to this group is **Lahorine**, an indenobenzazepine derivative, which may be derived biogenetically from the spirobenzylisoquinolines. **Hyperectine** represents a unique type of alkaloid substituted with an imidazole group.




Ochotensine

Presiner, R.M. *et al.*, *J. Nat. Prod.*, 1980, **43**, 305–318 (*rev*)



BENZO[C]PHENANTHRIDINE ALKALOIDS (VX3300)

These alkaloids are derived from tetrahydroprotoberberine precursors by oxidative cleavage of the C-6 to nitrogen bond, followed by cyclisation of C-6 on to position 13. Various oxidation stages exist, e.g. partially reduced benzophenanthridines, as in **Chelidonine**, and fully aromatic systems, as in **Sanguinarine**. There are also secoalkaloids (**Arnottianamide**), dimers, e.g. **Sanguidimerine**, and a group formed by the addition of a carbon substituent to C-8 e.g. **Corynoline**, which in some alkaloids has undergone further elaboration (**Buesgeniine**, **Chelelactam**, **Nitrotyrasanguinarine**).

Several numbering systems have been used for the benzophenanthridine alkaloids, but the one shown below, based on biogenetic considerations, has been adopted in this Dictionary (this is not the numbering scheme for Benzo[*c*]phenanthridine itself).



Tetrahydrocoptisine, R = H
Tetrahydrocorysamine, R = Me



Chelidonine, R = H
Corynoline, R = Me

Simanek, V., *Alkaloids*, 1985, **26**, 185–240 (*rev*)

Iwasa, K. *et al.*, *J.A.C.S.*, 1989, **111**, 7925–7931 (*biosynth*)

PHENETHYLISOQUINOLINE ALKALOIDS (VX3360)

This group (e.g. **Autumnaline**) arises from a phenethylisoquinoline precursor, which is itself generated by condensation of tyrosine with a C₆-C₃ unit derived from phenylalanine, probably via cinnamic acid.

In addition to the small group of simple phenethylisoquinolines (including dimers; **Jolantine**; **Melanthiodine**), several other members of the following groups are related to them by further elaboration.


Kametani, T. *et al.*, *Alkaloids*, 1973, **14**, 265–323; 1989, **36**, 172–223 (*revs*)

HOMOMORPHINANDIEONE ALKALOIDS (VX3370)

This is a small group, e.g. **Androcymbine**, which are biosynthetic intermediates between the phenethylisoquinolines and the colchicines (VX3400). Androcymbine arises by phenol oxidative coupling, probably of Autumnaline, by a process analogous to that involved in the biosynthesis of the morphine alkaloids.

HOMOAPORPHINE AND HOMOPROAPORPHINE ALKALOIDS (VX3380, VX3390)

The sequence from tyrosine and phenylalanine through a phenethylisoquinoline to the homoproaporphines and homoaporphines appears superficially to be exactly analogous to the course of biosynthesis of the aporphine alkaloids. However, although Autumnaline (VX3360 above) is an efficient precursor for both **Kreysiginone** and **Kreysigine** in *Kreysigia multiflora*, dienone intermediates such as Kreysiginone are not involved in the biosynthesis of the homoaporphines, such as Kreysigine. **Bitlisine** represents an anomalous type.




(-)-Kreysigine

Tojo, E., *J. Nat. Prod.*, 1989, **52**, 909–921 (*listing*)

COLCHICINE ALKALOIDS (VX3400)

Autumnaline (VX3360) and *O*-Methylandrocymbine (VX3370) (but not Androcymbine itself) are efficient precursors for **Colchicine**, an important pharmaceutical and research tool, and the details of the biosynthesis have been studied in detail.



Colchicine

Lumicolchicines are the product of u.v. irradiation of Colchicine, and while they have been reported to occur naturally, they could be regarded as artifacts. The rearrangement product **Demecolcinone** has also been isolated more recently.

Capraro, H.-G. *et al.*, *Alkaloids*, 1984, **23**, 1–72 (*rev*)

Boyè, O. *et al.*, *Alkaloids*, 1992, **41**, 125–176 (*rev*)


Battersby, A.R. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1998, 2989–2994; 2995–3001; 3002–3009 (*biosynth*)

Le Hello, C., *Alkaloids*, 2000, **53**, 287–352 (*pharmacol*)

DIBENZOCYCLOHEPTYLAMINE ALKALOIDS (VX3410)


Dibenzocycloheptylamine alkaloids are found in plants of the genera *Colchicum* and *Androcymbium*. To date only a few are known; these include **Jerusalemine** and **Salimine**. Jerusalemine may be formed through decarbonylation

of the tropolone ring of 2-Demethyldemecolcine by a peroxidase system present in the plant, with accompanying oxidation. Salimine, on the other hand, may arise from Colchicine by enzymatic peroxidation of ring C followed by hydroxylation and methylation.



CEPHALOTAXUS ALKALOIDS (VX3420)

A group of alkaloids from *Cephalotaxus* spp. (Cephalotaxaceae) in which the hydroaromatic component of the Erythrina group (VX2940 above) has undergone skeletal rearrangement. A typical alkaloid of this group is **Cephalotaxine**. Alkaloids occur in both enantiomeric series and some are potent antitumour agents.




Huang, L., *Alkaloids*, 1984, **23**, 157–227 (rev)

Miah, M.A.J., *Alkaloids*, 1998, **51**, 199–269 (rev)

HOMOERYTHRINA ALKALOIDS (VX3440)

E.g. **Schelhammeridine**. These would appear to be formed by a route analogous to that adopted in the Erythrina group (VX2940). **Selaginoidine** and **Phellibiline** represent unusual types involving oxidative degradation of the aryl ring.



Bick, I.R.C. al, *Alkaloids Chem. Biol. Perspect.*, 1989, **7**, 1–42 (rev)


AMARYLLIDACEAE (NARCISSUS) ALKALOIDS (VX3510-VX3599)

This large group of alkaloids is also derived from two tyrosine units which combine, with loss of one carbon atom, to give a benzylphenylethylamine precursor unit, e.g. **Norbelladine**. Various oxidative cyclisation processes, prominent among which are phenol oxidative coupling reactions, can give rise to the major skeletal groups. The unprimed aromatic ring derives from a simple aromatic precursor such as vanillin.

Oxidation of a norbelladine-type precursor at C-7, followed by cyclisation at C-2', gives the simple 1-aryltetrahydroisoquinolines (VX3510) exemplified by **Cryptostylin** **I** and **II**. Some representatives are also found in the Orchidaceae; and both enantiomers of Cryptostyline and of some other Amaryllidaceae alkaloids have been isolated from different plants. Oxidation at the alternative benzylic position (C-9) and cyclisation at C-6 gives the 4-aryltetrahydroisoquinolines related to **Cherylline** (VX3520). Some 4-aryltetrahydroquinolines (e.g. **Viridicatin**) have been isolated from microorganisms. Oxidative coupling of positions 2 and 2' gives the apogalanthamine skeleton of **Nivalidine** (VX3530), although this may be an artifact derived from Galanthamine.

Oxidative coupling of C-2 with C-1' in Norbelladine gives the **Galanthamine** (VX3540) skeleton which, by a straightforward cyclisation process (nitrogen to position 1), can give rise to the **Haemanthidine** (VX3550) ring system. The more recently discovered **Graciline** subgroup (VX3555) is related to Haemanthidine by N(5)-C(6) fission and pyran ring formation.

Opening of the carbinolamine function in Haemanthidine, followed by a redox reaction and cyclisation of the oxygen at C-6 on to position 11 (Haemanthidine numbering), then affords the **Tazettine** (VX3560) skeleton.



A further possibility is the migration of C-18 in the haemanthidine skeleton to position 11, which gives rise to the ring systems present in **Pancracine** (VX3570) type alkaloids.

A double cyclisation of C-2 to C-3' and nitrogen to C-2' provides the tetracyclic skeleton characteristic of **Lycorine** and its analogues (VX3580). Further modification of this ring system by oxidative fission of the nitrogen to C-7 bond followed by attachment of oxygen at C-7 to position 1 (galanthan numbering) then gives rise to the **Lycorenine** group (VX3585). A small number of Lycorenine alkaloids exemplified by **Galanthindole** show aromatisation of the indole portion, leading to a type of non-tryptophan-derived, biogenetically distinct type of indole alkaloid.

The small **Narciclasine** group (VX3590) also stems from Norbelladine, and appears to be formed via Crinine (but not 3-Epicrinine) by loss of the two-carbon bridge and appropriate oxidations.

Category VX3599, Miscellaneous Amaryllidaceae alkaloids, includes **Ismine**, an aromatised biaryl derived from a Haemanthidine precursor, and some novel two-nitrogen alkaloids such as **Plicamine**, and **Augustamine**, which cannot be assigned to known subtypes. There also remain a considerable number of alkaloids from earlier work to which structures were not assigned. **Varadine**, isolated in 1979, deserves reinvestigation.

Martin, S.F., *Alkaloids*, 1987, **30**, 252–369 (rev)

Hoshino, O., *Alkaloids*, 1998, **51**, 323–424 (rev)

Prabhakar, S. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 2001, **15**, 433–572 (rev, synth)

Bastida, J. *et al.*, *Alkaloids*, 2006, **63**, 87–179 (rev)


Jin, Z., *Nat. Prod. Rep.*, 2007, **24**, 886–905 (rev)

Unver, N., *Phytochem. Rev.*, 2007, **6**, 125–135 (rev, new types of Amaryllidaceae alkaloid)

MESEMBRENOID ALKALOIDS (VX3600)

Derived from two phenylalanine units with loss of one of the ethanamine sidechains, this smallish group of alkaloids is typified by **Mesembrine**; a second sub-group contains alkaloids in which the pyrrolidine ring has not

been formed, as in **Joubertiamine**. A third variant contains bases in which a pyridine ring has been fused on, as in **Sceletium alkaloid A4** and **Tortuosamine**. Mesembrenoid alkaloids occur in *Sceletium* spp. (Aizoaceae).




Jeffs, P.W. *et al.*, *Alkaloids*, 1981, **19**, 1–80 (rev)

Jin, Z., *Nat. Prod. Rep.*, 2007, **24**, 886–905 (rev)

EMETINE (IPECAC) ALKALOIDS (VX3690)

The emetine alkaloids are unique among the isoquinoline group in that they are derived from a monoterpenoid unit via Secologanin in an analogous manner to the monoterpenoid indole alkaloids. They occur in the Alangiaceae, Rubiaceae and Icacinaceae. Incorporation of one phenylalanine/tyrosine unit gives the alkaloids exemplified by **Ipecoside** and **Protoemetine**; the latter, by combination with a second amino acid unit, gives rise to the **Emetine** subgroup. Alternatively, combination with a tryptamine unit gives the typical alkaloids of *Alangium lamarckii*, e.g. **Tubulosine**.





Fujii, T.M. *et al.*, *Alkaloids*, 1983, **22**, 1–50; 1998, **51**, 271–321 (revs)

Wiegrebe, W. *et al.*, *J. Nat. Prod.*, 1984, **47**, 397–408 (rev)

PHENANTHROINDOLIZIDINE AND PHENANTHROQUINOLIZIDINE ALKALOIDS (VX3700, VX3760)

These alkaloids are derived from two molecules of phenylalanine or tyrosine, together with, presumably, ornithine (phenanthroindolizidines, e.g. **Tylophorine**) or lysine (phenanthroquinolizidines, e.g. **Cryptopleurine**).




Bick, I.R.C. *et al.*, *Alkaloids*, 1981, **19**, 193–220 (rev)

Gellert, E. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1987, **5**, 55–132 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2004, **21**, 625–649; 2005, **22**, 603–626; 2008, **25**, 139–165 (revs)

ERICAMYCINS (VX3900)

This is a relatively small group of polycyclic substances containing a fused isoquinoline system which has been little reviewed. The name Ericamycins is used here; although they are also referred to as the Albofungin group. They are of polyketide origin, with some unusual features to their biosynthesis. Various numbering systems have been used, and the numbering in CAS changes for trivial reasons (presence or absence of additional alicyclic rings, etc.)



Ericamycin

Bockholt, H. *et al.*, *J. Org. Chem.*, 1994, **59**, 2064–2069 (*biosynth*)

INDOLE ALKALOIDS

The group of alkaloids derived from tryptophan constitutes the largest, most varied and most fascinating of all alkaloid groups. The alkaloids include simple tryptamine derivatives, carbazoles (in which the ethanamine chain has been lost), a variety of alkaloids in which tryptamine has combined with one or more prenyl residues, and others in which regular monoterpene or diterpene units have been incorporated. However, the largest group, and the most extensively studied, is the thousands of alkaloids derived from tryptophan and a monoterpene unit based on Secologanin. The classical researches on indole alkaloids were carried out on terrestrial plant alkaloids, and then moved to studies of fungal-derived alkaloids. Indole alkaloids are also numerous in marine organisms, and the range of structural types differs from those found in terrestrial organisms. Many of the 'traditional' indole alkaloid types produced by terrestrial plants have not been found; polyhalogenation is a common feature of these alkaloids.

SIMPLE INDOLE ALKALOIDS (VX4000)

Under this heading appear various unhalogenated (plant) and halogenated (marine) indoles, together with higher molecular weight alkaloids containing unsubstituted indole ring systems, such as **Aplysinopsin** and its relatives (which are also imidazoles). Also included are a variety of simple indoles carrying prenyl, etc. substituents. The simplest plant indole alkaloid is **Gramine**.

Husson, H.-P., *Alkaloids*, 1985, **26**, 1–51 (*rev*)

Higuchi, K. *et al.*, *Nat. Prod. Rep.*, 2007, **24**, 843–865 (*rev*)

BISINDOLES (VX4020)

This category includes **Indirubin**, **6,6'-Dibromoindigotin** and other bisindoles; many from marine sources are polybrominated.

Yang, C.G. *et al.*, *Curr. Org. Chem.*, 2004, **8**, 1691–1720 (*rev, bisindole alkaloids from sponges*)

Kam, T.-S. *et al.*, *Alkaloids*, 2006, **63**, 181–337 (*rev*)

SIMPLE TRYPTAMINE ALKALOIDS (VX4040)

Tryptamine itself and some simple relatives have been isolated from marine tissues, and **Bufotenine**, long known as an amphibian product, has more recently been detected in a gorgonian. More highly elaborated marine alkaloids containing a (dehydro)tryptamine residue include the **Kottamides**. There are also some simple tryptamine dimers and oligomers such as the **Gelliusines**.


Somei, M. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 278–311 (*rev*)

Kawasaki, T. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 761–793 (*rev*)

Kam, T.-S. *et al.*, *Alkaloids*, 2006, **63**, 181–337 (*rev*)

CYCLOTRYPTAMINE ALKALOIDS (VX4100)

This group is formally derived from tryptophan by cyclisation and decarboxylation. (The presence of the cyclised form, cyclotryptophan, can be demonstrated in tryptophan solutions). **Physostigmine**, the prototype of this group of alkaloids, was first isolated from *Physostigma venenosum* and has also been produced by *Streptomyces* spp. The alkaloid is characterised by a urethane group which is readily hydrolysed with aqueous base to afford Eseroline. In addition to plant alkaloids with this skeleton, other representatives of this class have been isolated from marine organisms (e.g. the **Flustramines**) and from skin extracts of the Australian frog *Pseudophryne coriacea* (e.g. **Pseudophrynamine A**, where there is evidence for *de novo* biosynthesis by the amphibian rather than a dietary source). A number of interesting oligomers of this class occur in plants, e.g. **Chimonanthine**, a dimer, **Hodgkinsine** (a trimeric species), the **Quadrigenines** (tetramers) and higher.




Physostigmine

Anthoni, U. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1999, **13**, 163–236 (rev)

Kam, T.-S. *et al.*, *Alkaloids*, 2006, **63**, 181–337 (rev)

CHAETOCIN ALKALOIDS (VX4110)

Also called Epipolythiodioxopiperazines, although some members of the series lack the polysulfur bridge. These are microbial products, but a considerable number, including the extensive series of **Leptosins**, have been isolated from marine sources. The prototype is **Echinulin**, from *Aspergillus echinulatus*. They are biosynthesised from diketopiperazines, which are cyclic amino acid anhydrides.



Chaetocin

Williams, R.M. *et al.*, *Top. Curr. Chem.*, 2000, **209**, 97–173 (*prenylated tryptophans, biosynth*)


Somei, M. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 278–311 (rev)

Gardiner, D.M. *et al.*, *Microbiology (Reading, U.K.)*, 2005, **151**, 1021–1032 (*biosynth*)

Kawasaki, T. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 761–793 (rev)

EUODIA (QUINAZOLINOCARBAZOLE) ALKALOIDS (VX4120)

These alkaloids are characteristic of the Rutaceae. The most studied is **Rutaecarpine**, which has useful hypotensive activity. They arise by incorporation of a C₁ unit into a β -carboline precursor.




Rutaecarpine

Bergmann, J., *Alkaloids*, 1983, **21**, 29–55 (rev)

INDOLACTAM ALKALOIDS (VX4200)

The indolactam group contains an unusual nine-membered lactam ring derived from tryptophan and valine, via *N*-Methyl-L-isoleucyl-L-tryptophanol. The grouping occurs in the tumour-promoting **Teleocidins** and **Lyngbyatoxins**, which are microbial products. These contain a monoterpene portion which is derived from glucose, not mevalonate.




Lyngbyatoxin A

Irie, K *et al.*, *Tet. Lett.*, 1990, **31**, 7337–7340; 1998, **39**, 7929–7930 (*biosynth*)

β -CARBOLINE ALKALOIDS (VX4240)

A large number of relatively simple β -carboline (pyrido[3,4-*b*]indole) derivatives occur naturally. These include β -carbolines unsubstituted at *C*-1, e.g. **1,2,3,4-Tetrahydro-6-methoxy-2-methyl- β -carboline**, those containing a methyl group at *C*-1, i.e. the **Harman** group, and several which contain a substituent at *C*-1 and/or *C*-3. The substituents at position 1 may be acyl, carboxyl, or they may be more complex, as in **Perlolyrine**



Harman

Long known as terrestrial alkaloids, these are also widespread in marine organisms. β -Carboline (**Norharman**) itself has been found in an ascidian and in dinoflagellates, and examples of simple brominated analogues include the **Eudistomins** and some of the **Arborescidines**. Another group of β -carboline derivatives, presumably derived from tryptophan and cysteine, have been isolated from *Eudistoma olivaceum*, a Caribbean tunicate. The β -Carboline nucleus is also found in some of the **Manzamines** and other more complex alkaloids.

Husson, H.-P., *Alkaloids*, 1985, **26**, 1–51 (*rev*)

Baker, B.J. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1997, **10**, 357–407 (*rev, marine β -carbolines*)


Somei, M. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 278–311 (*rev*)

Fresneda, P.M. *et al.*, *Synlett*, 2004, 1–17 (*rev, synth*)

Kawasaki, T. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 761–793 (*rev*)

CARBAZOLE ALKALOIDS (VX4300)

Nearly all carbazole alkaloids have a substituent at *C*-3, although **Carbazole** itself and one or two simple marine halogenated derivatives have also been characterised. The main source of carbazole alkaloids are the plants of the Rutaceae subfamily Aurantioideae, especially the genera *Clausenia*, *Murraya* and *Glycosmis*. They are also found in some other higher plants, as well as streptomycetes and cyanobacteria. **3-Methylcarbazole** appears to be the key biosynthetic intermediate, although there is little direct experimental evidence. Elaboration into prenylcarbazoles (e.g. **Ekeberginine**) and dimerisation occurs. They have been well-reviewed.




Ekeberginine

Husson, H.-P., *Alkaloids*, 1985, **26**, 1–51 (rev)
 Chakraborty, D.P., *Alkaloids*, 1993, **44**, 257–364 (rev)
 Knölker, H.-J. *et al.*, *Chem. Rev.*, 2002, **102**, 4303–4428 (rev)
 Knölker, H.-J., *Curr. Org. Synth.*, 2004, **1**, 309–331 (rev, synth)
 Somei, M. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 278–311 (rev)
 Kawasaki, T. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 761–793 (rev)
 Knölker, H.-J. *et al.*, *Alkaloids*, 2008, **65**, 3–158; 159–180; 189–194; 195–384 (occurrence, biosynth, biochem, chemistry)

INDOLO[2,3-*A*]CARBAZOLE ALKALOIDS (VX4350)

These are a class of alkaloids exemplified by **Staurosporine**, which was first isolated in 1977 from a terrestrial microorganism, and which has since been found to be widely distributed. They are essentially microbial metabolites from a variety of organisms (field-collected or cultured), and show a wide range of biological activities. Others are the **Tjipanazoles** (from the blue-green alga *Tolypothrix tjipanasensis*) and several metabolites from slime moulds of the genus *Arcyria* (e.g. the **Arcyriarubins** and **Arcyriaflavins**). The isolation of **4'-*N,O*-Didemethylstaurosporine** from a flatworm *Pseudoceros* sp. is noteworthy. Their biosynthesis has been intensively studied; they are produced from bisindolyl precursors resembling **Violacein**; the indolyl residues are derived from tryptophan via 7-chlorotryptophan followed by an oxidative ring closure.




Staurosporine

Gribble, G.W. *et al.*, *Studies in Natural Product Chemistry*, (ed. Atta-ur-Rahman), Elsevier, 1993, **12**, 365 (rev)
 Walsh, C.T. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 517–531 (rev, biosynth)
 Sanchez, C. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 1007–1045 (rev, biosynth)

INDOLONAPHTHYRIDINE ALKALOIDS (VX4400)

These are the canthinones, a numerous and relatively straightforward subgroup of the β -carboline, e.g. **Canthin-6-one**, in which an additional C₃ unit is attached between C-1 and the indole nitrogen to form an additional ring. The C-3 unit appears to be derived from ketoglutarate. The group includes a few dimeric examples, e.g. **Haplophytine**. They occur especially in the Simaroubaceae and Rutaceae, and some representatives have been isolated from basidiomycete fungi.




Canthin-6-one

Aragozzini, F. *et al.*, *Plant Cell Rep.*, 1988, **7**, 213–215 (biosynth)
 Ohmoto, T. *et al.*, *Alkaloids* 1989, **36**, 135–170 (rev)

ERGOT ALKALOIDS (VX4460)

These alkaloids, which classically occur in the fungus *Claviceps purpurea* which in the past was responsible for ergotism, are derived from 4-prenyltryptophan by cyclisation to give a tricyclic base related to **Chanoclavine I**, a representative of the simplest subgroup. Further elaboration gives the tetracyclic ergoline skeleton, as in

Elymoclavine, which is present in the majority of alkaloids in this group. The most important alkaloids, many of which have useful medical applications, are more complex peptide alkaloids formed from lysergic acid, in which C-17 in the elymoclavine-type precursor has been oxidised to a carboxyl group, by attachment to one or more amino acids. **Ergocristine**, which is based on the ergotaman nucleus, is typical.



Chanoclavine I

They have been considered exclusively the metabolites of terrestrial fungi and microorganisms, but the recent isolation of the **Pibocines**, simple brominated analogues of Festuclavine, from *Eudistoma ascidians*, have provided a first marine occurrence. The presence of the bromine substituent in the Pibocines argues for their being genuine ascidian products and this is presumably a case of biochemical parallelism.

A few other metabolites, which may also be included in this group, are the result of skeletal rearrangement. **Clavicipitic acid** is one such compound; another is **α -Cyclopiazonic acid**. However, whereas the former is definitely a product of prenyl-tryptophan metabolism the latter, from *Penicillium cyclopium* is not, since it appears to arise from reaction of a tryptophan-acetoacetate unit with prenyl pyrophosphate.

Floss, H.G., *Tetrahedron*, 1976, **32**, 873–912 (rev, biosynth)

Horwell, D.C., *Tetrahedron*, 1980, **36**, 3123–3149 (rev, synth)

Ergot Alkaloids: Chemistry, Biological Effects, Biotechnology, (eds. Řehaček, Z. et al.), Elsevier, 1990 (rev)

Groger, D. et al., *Alkaloids*, 1997, **50**, 171–218 (rev, biochem)


Somei, M. et al., *Alkaloids*, 2000, **54**, 191–257 (rev, synth)

Mukherjee, J. et al., *Biochem. Eng. Biotechnol.*, 2000, **68**, 1–20 (rev)

Schardl, C.L. et al., *Alkaloids*, 2006, **63**, 45–86 (rev, biochem)

INDOLOTERPENOID ALKALOIDS (VX4620)

This group consists of a variety of alkaloids in which a tryptamine unit is attached to a terpenoid unit. It includes the *Aristotelia* alkaloids, of which **Aristoteline** is typical, and some other terrestrial alkaloids isolated from *Borreria* spp. There are numerous variations in the structure of the terpenoid part of the skeleton. Also included here are the **Hapalindoles** and related alkaloids mostly from freshwater cyanophytes, which are mostly isocyano/ isothiocyano substituted.



Aristoteline

Alkaloids included in this classification code are not 'monoterpenoid indole alkaloids' in the strict sense as defined below for subsequent categories derived from secologanin. They are also distinguished in the classification code from the Penitrem group, VX6470.

Bick, I.R.C., *Alkaloids*, 1985, **24**, 113–151 (rev)

Borschberg, H.J., *Chimia*, 1991, **45**, 325–341 (rev)

Borschberg, H.J., *Alkaloids*, 1996, **48**, 190 (rev)

MONOTERPENOID-DERIVED INDOLE ALKALOIDS (VX4640-VX5980)


The extremely important and varied group of monoterpenoid indole alkaloids originate from the condensation of tryptophan with secologanin to give **Strictosidine**, which is further elaborated to give the corynanthe alkaloids, together with an impressive array of structural variants which are divided here into the numerous sub-groups

tabulated in Table 1, and which are also described in the category descriptions which follow. Not all of the smaller groups of indole alkaloids are described in detail in the paragraphs which follow. If there is no description, their generic structure can be seen in Table 1, and any important references and reviews concerning them can be accessed from the individual dictionary entries.


Various schemes have been used to divide the indole alkaloids into major groups based on the major biogenetic branch points. The most widely used recognises classes 1-5, where class 1 contains those skeletons most closely related to secologanin. These schemes are not further referred to in the discussion which follows. Certain biogenetic generalisations can be made, e.g. class 3 alkaloids are found only in the Apocynaceae.

Biogenetic numbering is used throughout this introduction and the individual entries. (Le Men, J., *et al.*, *Experientia*, 1965, **21**, 508–510; Anon., *Alkaloids*, 2001, **56**, xxiii–xxiv). This almost invariably differs from systematic (e.g. CAS) numbering. However, it allows the biogenetic origin of the individual carbons in the different types of indole alkaloid to be traced in the numerous skeletons. It does have the slight disadvantage that the numbering becomes discontinuous whenever there are new bond-making and bond-breaking processes, as illustrated in the chart. CAS numbering may be encountered for some groups, such as the Iboga alkaloids, but the Le Men biogenetic scheme is preferred by the majority of specialists in the field.


Table 1. Monoterpenoid indole alkaloid skeletons




Ajmalicine skeleton
VX4860




Akuammiline skeleton
VX5200




Camptothecin skeleton
VX4700




Ajmaline skeleton
VX5120




Aspidofractine skeleton
VX5540




Chippiine skeleton
VX5940




Akageran skeleton
VX4840




Aspidosperma skeleton
VX5400




Cinchona skeleton
VX5240




Akuammicine skeleton
VX5260




Cadamban skeleton
VX4940




Cleavamine skeleton
VX5820




Condylorcarpan skeleton
VX5320




Iboga skeleton
VX5700




Melonine skeleton
VX5460




Correantane skeleton
VX4840




Kopsane skeleton
VX5560




Pandoline skeleton
VX5800




Corynanthean skeleton
VX4800




Koumine skeleton
VX5150




Pleiocarpaman skeleton
VX5220




Eburna skeleton
VX5900




Macroline skeleton
VX4900




Pyridocarbazole skeleton
VX5840




Ervatamia skeleton
VX5180




Malindan skeleton
VX4920




Quebrachamine skeleton
VX5500




Gelsemium skeleton
VX5000




Melodinus skeleton
VX5580




Rhazinilam skeleton
VX5450




Sarpagine skeleton
VX5100




Strychnidine skeleton
VX5280




Vallesiachotaman skeleton
VX4960




Schizozygine skeleton
VX5930




Tacaman skeleton
VX5920




Vindoline skeleton
VX5550




Secodine skeleton
VX5360




Ulean skeleton
VX5860




Vobasine skeleton
VX5140



Strictosidine skeleton
VX4640



Vallesaman skeleton
VX5880



Yohimbine skeleton
VX5040

In the main types of indole alkaloids having the secologanin subunit in unrearranged form, stereoisomerism at positions 3, 19, and 20 is common, but C-15 is invariant based on the corresponding carbon in secologanin. C-22 is frequently absent (for example, in all known actual alkaloids of the Pleiocarpamine group VX5220), but is shown in the skeletons.

Some indole alkaloids in various groups are found as variants resulting from indole to indoxole or pseudoindoxyl rearrangement. These are often treated as a separate category, oxindole alkaloids, but here they have been listed under the corresponding parent indole structure type (change from the first edition), as new types continue to be found. For example, **Rhynchophylline**, based on the corynoxan skeleton, is here categorised as a corynanthe alkaloid, VX4800. Another type of rearrangement found in some of the groups (Camptothecins, Cinchona alkaloids) is conversion of the indole to a quinoline nucleus. This means that the resulting structures do not strictly speaking contain an indole nucleus. However, they are considered to be indole alkaloids on biogenetic grounds.

Indole alkaloids are most common in the Apocynaceae, Loganiaceae and Rubiaceae, and are also found in the Alangiaceae, Icacinaceae, Nyssaceae, and possibly other families.

The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids, (ed, Saxton, J.E.), Wiley, 1983, **25**, part 4; supplement, 1994 (*rev. bibl*)
Saxton, J.E., *Nat. Prod. Rep.*, 1997, **14**, 559–590 and earlier (*revs*)
Verpoorte, R. *et al.*, *Alkaloids*, 1997, **49**, 221–299 (*rev*)
Leonard, J., *Nat. Prod. Rep.*, 1999, **16**, 301–318 (*rev*)
O'Connor, S.E. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 532–537 (*rev, biosynth*)
Szabó, L.F., *J. Phys. Org. Chem.*, 2006, **19**, 579–591 (*rev, classification*)
Szabó, L.F., *Molecules*, 2008, **13**, 1875–1896 (*rev, classification*)

STRICTOSIDINE ALKALOIDS (VX4640)

This category covers alkaloids related to the biogenetic parent alkaloid **Strictosidine** without further carbon-skeleton modification.

CAMPTOTHECIN ALKALOIDS (VX4700)

These alkaloids, which contain the quinoline ring system, are probably derived from Strictosidine via an intermediate related, possibly, to **Rubescine**. In this case, conversion of the indole group into the quinoline ring involves ring enlargement of ring B at the expense of ring C; otherwise the changes in the formation of **Camptothecin** from Strictosidine are straightforward.

Soepenber, O. *et al.*, *Alkaloids*, 2003, **60**, 1–50 (*rev, biochem*)

CORYNANTHEAN ALKALOIDS (VX4800)

This large group, formed by simple cyclisation of the precursor, is exemplified by **Geissoschizine** and **Sitsirikine**.

In corynanthe alkaloids and other groups closely related to them (e.g. yohimbanes, below), stereoisomeric variation is common except at C-15, which is invariably 15 α H-.

Szántay, C. *et al.*, *Alkaloids*, 1986, **27**, 131–269 (*rev*)

SECOLOGANIN-DERIVED SESQUIMER ALKALOIDS (VX4820)

These are derived from two biogenic amine subunits and one monoterpenoid subunit (normal sesquimers), or one biogenic amine subunit and two secologanin units (reversed sesquimers). The greatest number are formed by condensation of a (usually) Corynanthe portion with an additional tryptamine or canthine subunit, as in the **Ochrolifuanines** and **Usambarensines**. The dictionary entries carry the code numbers for the constituent fragments also.

Brown, R.T., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 63–146 (*rev*)

Kam, T.-S. *et al.*, *Alkaloids*, 2006, **63**, 181–337 (*rev*)

AKAGERAN AND RELATED ALKALOIDS (VX4840)

This relatively small group of alkaloids typified by **Akagerine** appears to be formed by bond-breaking of a corynanthe precursor followed by formation of a 7-membered ring. Included here are a small number related to **Adifoline**.

AJMALICINE-LIKE ALKALOIDS (VX4860)

These are based on a 17,19-secoyohimban skeleton which is invariably present as an ether (oxayohimban). **Ajmalicine** was formerly considered a key intermediate in indole alkaloid biosynthesis, but this has been disproved.

Lewis, S.E., *Tetrahedron*, 2006, **62**, 8655–8681 (*rev*)

MACROLINE ALKALOIDS (VX4900)

This group is related to the Akuammilines (VX5200 below) by scission of the C-21 to N-4 bond. The macroline skeleton mostly occurs in bisindole alkaloids; Macroline itself has not yet been found as a free alkaloid and only a small number of monomeric analogues are known.

MALINDAN ALKALOIDS; CADAMBAN ALKALOIDS; VALLESIAHOTAMAN ALKALOIDS (VX4920, VX4940, VX4960)

See chart. These are small groups, e.g. **Malindine**; **Cadambine**, **Vallesiachotamine**, some alkaloids having an elaborate additional pyridine ring.

GELSEMIUM ALKALOIDS (VX5000)

These alkaloids contain an oxindole function and a cage-like, hydroaromatic residue which can be imagined, in a formal sense, to arise from an intermediate related to anhydrovobasinediol by formation of a 6,20 bond and rearrangement to an oxindole. The major alkaloids in this group are related to **Gelsemine**; however, a smaller group, characterised by **Gelsedine**, lack the 6,20 bond, and have also lost C-21.

Takayama, H. *et al.*, *Alkaloids*, 1997, **49**, 1–78 (*rev*)

YOHIMBINOID ALKALOIDS (VX5040)

The Yohimbinoïd alkaloids contain a carbocyclic ring E arising through C-17 to C-18 bond formation in a corynantheine precursor. As in the corynantheine and ajmalicine groups, stereoisomerism at all chiral centres, except C-15, is known. Structural variations include the presence of methoxy-groups in the aromatic ring, hydroxy- or acyloxy- groups at C-18, as in **Reserpine**, and various degrees of unsaturation in rings C-E, as in **Alstoniline**.

Brown, R.T., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part4, 147–199 (*rev*)

Szántay, C. *et al.*, *Alkaloids*, 1986, **27**, 131–269 (*rev*)

Baxter, E.W. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1990, **8**, 197–319 (*rev, synth*)

SARPAGINE ALKALOIDS (VX5100)

The Sarpagine (Akuammidine) group, based on the sarpagan nucleus, arises from bond formation between C-16 and C-5 of the corynantheine precursor and consists of simple **Akuammidine** derivatives and oxindoles obtained following the migration of C-3 from C-2 to C-7, e.g. **Gardneramine**. The closely related Vobasine (VX5140) and Ervatamia (VX5180) groups are now listed separately.

Joule, J.A., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 201–264 (*rev*)

Hamaker, L.K. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1994, **9**, 23–84 (*rev, synth*)

Lounasmaa, M. *et al.*, *Alkaloids*, 1999, **52**, 103–195 (*rev*)

Lewis, S.E., *Tetrahedron*, 2006, **62**, 8655–8681 (*rev*)

AJMALINE ALKALOIDS (VX5120)

The ajmaline group contains both 5,16 and 7,17 bonds. **Ajmaline** itself is the best known example. Almost all the bases in this group contain the same skeleton, but **Perakine** and its congeners afford a rare structural variation in which the 21,N bond has been replaced by a 19,N bond.

Lewis, S.E., *Tetrahedron*, 2006, **62**, 8655–8681 (*rev*)

Lounasmaa, M. *et al.*, *Alkaloids*, 2001, **55**, 1–87 (*rev*)

VOBASINE ALKALOIDS (VX5140)

These are related to the sarpagine group by C-3 to N-4 bond scission.

KOUMINE ALKALOIDS (VX5150)

These are 7,20-cyclovobasine alkaloids.

ERVATAMIA ALKALOIDS (VX5180)

These alkaloids (e.g. **Ervatamine**, **Methuenine**) can be successfully produced *in vitro* from vobasine-type alkaloids (VX5140) and this has also been achieved by enzymic preparations. The process involves 5,6- bond cleavage and 6,16- bond formation.

Thai, C. *et al.*, *J.A.C.S.*, 1981, **103**, 4956–4957 (*biosynth*)

AKUAMMILINE ALKALOIDS (VX5200)

The ring system in this group is formed from a precursor of the corynantheine type by bond formation between C-16 and C-7. In addition to close relatives of **Akuammiline**, variations in this subgroup include alkaloids derived by C-3 to N-4 bond fission and C-2 to N-4 bonding, e.g. **Echitamine**, and alkaloids in which the N-4 to C-5 bond in the Akuammiline skeleton has been severed, e.g. **Aspidodasycarpine**. **Nareline** is a unique alkaloid with an additional bond between C-21 and C-6.

Lewis, S.E., *Tetrahedron*, 2006, **62**, 8655–8681 (*rev*)

PLEIOCARPAMAN ALKALOIDS (VX5220)

In this group, a corynantheine precursor has cyclised via C-16 on to N-1, as in **Pleiocarpamine**.

Gilbert, B., *Alkaloids*, 1968, **11**, 205–306 (*rev*)

CINCHONA ALKALOIDS (VX5240)

This important and well-known group, which includes the valuable antimalarial quinine, consists of two subgroups: the **Cinchonamine** group, derived from a corynantheine-type precursor by fission of the N-4 to C-5 bond plus attachment of N-4 to C-17, and the **Quinine** group, which contain a quinoline ring system generated from a precursor of the cinchonamine type by 2,7-bond fission followed by bonding of N-1 to C-5. **Corialstonine** represents an abeo-variant.

Grethe, G. *et al.*, *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 729–752 (*rev*)

Hoffmann, H.M.R. *et al.*, *Eur. J. Org. Chem.*, 2004, 4293–4312 (*rev*)

AKUAMMICINE (CURAN) ALKALOIDS (VX5260)

The biogenesis of these alkaloids presumably involves migration of C-3 in a corynanthe precursor from C-2 to C-7, followed by formation of the 2,16-bond. An early alkaloid in the group is therefore **Preakuammicine**, which loses formaldehyde to give **Akuammicine**. Included in this group is **Stemmadenine**, a 3,7-secoalkaloid which appears to be the intermediate on the pathway to the Ulean group below (VX5860).

STRYCHNIDINE ALKALOIDS (VX5280)

Formation of the skeleton of these typical *Strychnos* alkaloids involves the addition of two carbon atoms to a curan precursor, presumably, from an acetate unit

Husson, H.-P., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton J.E.), Wiley, 1983, **25**, part 4, 293–330 (*rev*)

Bosch, J. *et al.*, *Alkaloids*, 1996, **48**, 75–189 (*rev*)

Ohshima, T., *Chem. Pharm. Bull.*, 2004, **52**, 1031–1052 (*rev, synth*)

Shibasaki, M. *et al.*, *Alkaloids*, 2007, **64**, 103–138 (*rev, synth*)

CONDYLOCARPAN ALKALOIDS (VX5320)

These alkaloids contain a ring system similar to that of the curan group, but are formed by cyclisation of C-21 on to C-7 in a Corynanthe precursor, rather than the formation of a 3,7 bond; **Condylocarpine** is representative. Note that loss of the ethanamine carbons 5 and 6 gives the ring system of uleans (VX5860 below), which thus suggests an alternative biogenetic route to the one given below. Extensive modification of this skeleton appears to have occurred in the formation of **Goniomine**, which can be postulated to be formed by ring-opening and epoxidation of an indolenine related to Condylocarpine followed by N-1 to C-19 bonding.

Lounasmaa, M., *Prog. Chem. Org. Nat. Prod.*, 1986, **50**, 27–56 (*rev*)

SECODINE ALKALOIDS (VX5360, VX4740)

This group of tricyclic alkaloids is formed by ring-opening of a precursor of the Preakuammicine type. The alkaloids occur in various stages of reduction, and in monomeric and dimeric forms. **Andranginine**, the product of an unusual cyclisation of a dehydrosecodine, may also be included here.

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767 (rev, synth)

ASPIDOSPERMA ALKALOIDS (VX5400)

The skeleton of the aspidospermidine alkaloids is formed by cyclisation of a dehydrosecodine, itself obtained from a precursor related to Preakuammicine. The alkaloids in this group are based on several structural variants. A methoxycarbonyl group at C-16 may be present (**Tabersonine**) or absent (**Aspidospermine**); in each case, C-18/C-19 may be an ethyl group, or may be functionalised, e.g. into a lactone system. **Obscurinervidine** contains a dihydro-1,4-oxazine ring between N-1 and C-12.

Saxton, J.E., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 331–437 (rev)

Overman, L.E. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1985, **3**, 275–307 (rev)

Cordell, G.A., *Alkaloids*, 1997, **50**, 327–384 (rev)

Saxton, J.E., *Alkaloids*, 1998, **51**, 1–197 (rev)

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767 (rev, synth)

RHAZINILAM ALKALOIDS (VX5450)

These are alkaloidal artifacts related to the aspidosperma group by C2-C7 fission, the indole functionality thus being lost.

Hájíček, J., *Collect. Czech Chem. Commun.*, 2007, **72**, 821–898 (rev, synth)

MELONINE ALKALOIDS (VX5460)

This small group has undergone a (6 → 2) abeo shift from an Aspidosperma-like precursor.

QUEBRACHAMINE ALKALOIDS (VX5500)

Quebrachamine alkaloids are derived from the Aspidosperma group (type III skeleton) by fission of the 7,21-bond. They may have lost the C-16 methoxycarbonyl group (e.g. **Quebrachamine**) or it may have been retained, as in **Vincadine**.

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767; 2007, **72**, 821–898 (rev, synth)

ASPIDOFRACTINE ALKALOIDS (VX5540)

These are related to the Aspidosperma group by formation of a C-2 to C-18 bond.

Hájíček, J., *Collect. Czech Chem. Commun.*, 2007, **72**, 821–898 (rev, synth)

VINDOLININE ALKALOIDS (VX5550)

These are related to the Aspidosperma group by formation of a C-2 to C-19 bond.

KOPSANE ALKALOIDS (VX5560)

The skeleton of the kopsane group of alkaloids is simply formed by attachment of C-22 (the methoxycarbonyl carbon) of Venalstonine to C-6, as in **Kopsine**. Skeletal variations include the alternative acyloin structure, as in **Fruticosine**, in which C-22 is attached to C-17.

Hájíček, J., *Collect. Czech Chem. Commun.*, 2007, **72**, 821–898 (rev, synth)

Kam, T. S. *et al.*, *Alkaloids*, 2008, **66**, 1–111 (rev)

MELODINUS ALKALOIDS (VX5580)

These have been isolated only from *Melodinus* and *Kopsia* spp. (e.g. **Meloscine**, **Scandine**). They are plausibly biosynthesised from alkaloids of the Tabersonine type (VX5400). There are two known dimer alkaloids.

Hugel, G. *et al.*, *J.O.C.*, 1986, **51**, 1594–1595 (*biosynth*)

IBOGA ALKALOIDS (VX5700)

A further mode of cyclisation of a secodine-type precursor involves formation of a 16,21 bond, which gives rise to the bridged ring system (type II indole alkaloid) found in **Ibogamine**, **Catharanthine**, and numerous related alkaloids. They are often components of heterodimer indole alkaloids.

This group of alkaloids exists in both enantiomeric series which may be defined by the chirality of C-14. The best-known example of the 14*S* series is Catharanthine. Most other alkaloids of this group belong to the 14*R* series. Many alkaloids retain the methoxycarbonyl group, whereas others (e.g. Ibogamine) have lost it. Other variations include oxidation at C-7 to give the related hydroxyindolenines, e.g. **Iboxygaine hydroxyindolenine**; oxidation followed by rearrangement to the related indoxyl, e.g. **Desmethoxyiboluteine**, oxidative rearrangement to the corresponding oxindole, as in **Tabernoxidine**; and oxidation at C-19, C-3, C-5, or C-6 with, occasionally, ether formation between oxidised positions.

Also **16-Hydroxyalloibogamine**, the only alkaloid having the analogous but isomeric bridged ring system (type III skeleton) is also ranked into this class of alkaloids.

Cordell, G.A., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 467–537 (*rev*)

Popik, P. *et al.*, *Alkaloids*, 1998, **52**, 197–231 (*rev, pharmacol*)

Sundberg, R.J. *et al.*, *Alkaloids*, 2002, **59**, 281–366 (*rev*)

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767 (*rev, synth*)

PANDOLINE ALKALOIDS (VX5800)

The Pandoline (pseudoaspidospermidine) (type II indole alkaloid) nucleus can be imagined to be formed by cyclisation of a secodine derivative isomeric with that postulated as a precursor for Aspidospermidine. **Ibophyllidine** represents an obvious ring contraction, **Iboxyphylline** a ring extension.

CLEAVAMINE ALKALOIDS (VX5820)

The small **Cleavamine** group is more often encountered as degradation products of other alkaloids; these may arise by fission of the 3,7-bond. Alternatively, and perhaps more likely, this ring system can be generated by fission of the 16,21-bond in an Iboga skeleton (see below).

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767; 2007, **72**, 821–898 (*rev, synth*)

PYRIDOCARBAZOLE ALKALOIDS (VX5840)

This small and pharmacologically important group is based on the 6*H*-pyrido[4,3-*b*]carbazole ring system, and is exemplified by **Ellipticine** and **Olivacine**. Although these aromatic bases may superficially seem to be unrelated to the mainstream indole monoterpenoid alkaloids a possible biogenesis from Stemmadenine can be postulated. However, the details of the biosynthesis do not appear to have been investigated experimentally since the 1970s, when only low incorporation of tryptophan was found.

There are two subtypes in that Ellipticine is derived biogenetically from the Ajmalicine type (VX4860) and Olivacine from the Malindan type (VX4920). Because of this biogenetic diversity and their separation structurally from the mainstream indole alkaloids, systematic (CAS) numbering is now usually employed in the literature for the ellipticine nucleus.

Gribble, G.W. *et al.*, *Heterocycles*, 1985, **23**, 1277–1315 (*rev, synth*)

Kansai, V.K. *et al.*, *Tetrahedron*, 1986, **42**, 2389–2408 (*rev*)

Alvarez, M. *et al.*, *Alkaloids*, 2001, **57**, 235–272 (*rev*)

ULEAN AND VALLESAMAN ALKALOIDS (VX5860, VX5880)

These alkaloids may well arise, like the Ellipticine group, from an oxidative fission of a Stemmadenine-like intermediate. The genesis of the three types, e.g. **Vallesamine**, **Uleine**, and **Apparicine** can thus readily be explained.

Joule, J.A., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 265–292 (rev)
Alvarez, M. *et al.*, *Alkaloids*, 2001, **57**, 235–271 (rev)

EBURNA AND SCHIZOZYGINE ALKALOIDS (VX5900; VX5930)

The skeleton of the Eburna alkaloids is generated by rearrangement of the aspidospermidine ring system, involving migration of C-21 from C-7 to C-2, fission of the 2,16-bond, and attachment of C-16 to N-1. This rearrangement has been very successfully imitated *in vitro*. **Vincamine** and its derivatives retain the methoxycarbonyl group; **Eburnamine** and **Eburnamenine** have lost it. There are some alkaloids in which C-18 or C-19 is oxidised, e.g. **Cuanzine**.

The **Schizozygine** group (VX5930) contain an additional bond between C-2 and C-18. Included are **Andrangine** and **Vallesamidine**, which can be considered as secoschizozygines or as Eburna alkaloids in which C-21 has simply migrated to C-2.

Saxton, J.E., *The Chemistry of Heterocyclic Compounds: Indoles-The Monoterpenoid Indole Alkaloids*, (ed. Saxton, J.E.), Wiley, 1983, **25**, part 4, 439–465 (rev)
Lounasmaa, M. *et al.*, *Alkaloids*, 1992, **42**, 1–116 (rev)

TACAMAN ALKALOIDS (VX5920)

A small group is apparently obtained by processes analogous to the formation of the Eburna group, but from a precursor to a pseudoaspidospermidine (type II skeleton) derivative, by the formation of new bonds at C-2 → C-3 and C-14 → C-17. Again, C-22 may be present (**Tacamine**) or absent (**Tacamoline**).

Danieli, B. *et al.*, *Alkaloids*, 1986, **27**, 95–103 (rev)

CHIPPIINE ALKALOIDS (VX5940)

These represent a numerically limited group. Their structures can be derived from an Iboga-type precursor by cleavage of the C-3 to N-4 bond and formation of an N-1 to C-3 bond.

BISINDOLE ALKALOIDS (VX5980)

This large group of complex alkaloids consists of a wide variety of structures, depending on the identity of the monomeric alkaloid components. The individual alkaloids are normally coded in addition with the Type of Compound codes relevant to the component halves of the molecule. (e.g. **Conodurine** = a Vobasine (VX5140)-Iboga (VX5700) dimer). The most practically important indole alkaloid dimers are those of the Cleavamine-Aspidosperma group. **Vinblastine** and **Vincristine** are the best known examples.

Whilst in these dimers and many others the monomeric units are linked by single covalent bonds, in others the assembly is more complex. The **Vobtusine** group, for example, is composed of two aspidospermidine-type units linked by a spirocyclic system involving C-14 (two bonds) of one unit with C-22' of another unit, together with an additional carbon atom attached to N-1'. A small group contains three inter-unit bonds as in **Ervafoline**.

Other types of dimeric alkaloids from *Strychnos* Spp. form the major constituents of calabash curare. These are composed of two units linked via N-1 and C-17', and N-1' and C-17; **C-Toxiferine** is representative. In some alkaloids additional bonds are present; for example, **C-Curarine** has an ether bridge between C-16 and C-16', and **C-Calebassine** has an additional carbon-carbon bond between C-17 and C-17'.

Cordell, G.A., *Alkaloids*, 1981, **20**, 3-295; 1983, **25**, 539–728 (revs)

Hájíček, J., *Collect. Czech Chem. Commun.*, 2004, **69**, 1681–1767; 2007, **72**, 821–898 (rev, synth)


Kam, T.-S. *et al.*, *Alkaloids*, 2006, **63**, 181–337 (rev)

ISOINDOLES (VX6000)

This is a heterogeneous group including terpenoids which are also coded as such. Small isoindoles include microbial products, e.g. **Cichorine**.

TRYPTOQUIVALINES (VX6030)

The **Tryptoquivalines**, which are toxic metabolites from *Aspergillus clavatus*, are derived from anthranilic acid and tryptophan precursors, and half of the molecule is related to the Quinazoline alkaloids above (VX1600). The reviews of quinazoline alkaloids cited there normally cover the tryptoquivalines.




Tryptoquivaline A

There has been some uncertainty concerning their absolute configurations. It seems now established that they all have the (2*S*,3*S*,12*R*)- stereochemistry, but in the case of a few of the tryptoquivalines corrected stereochemistries have not been published.

The group also includes some more recently isolated analogues such as the **Fumiquinazolines**.

MITOMYCINS (VX6050)

This is a small group of purple microbial pigments (their typical representative is **Mitomycin A**) showing antitumour properties. They are usually quinones containing an aziridinyl function. They are biosynthesised from 3-amino-5-hydroxybenzoic acid, the aziridine nitrogen coming from glucosamine. The **Albomitomycins**, rearrangement products of mitomycins, contain a diazatricyclic ring system unique among natural products.




Mitomycin A

Remers, W.A. *et Alkaloids Chem. Biol. Perspect.*, 1987, **5**, 1–74 (rev)

PYRROLO[4,3,2-DE]QUINOLINE ALKALOIDS (VX6070)

This skeleton is characteristic of and almost exclusive to marine (sponge) alkaloids. Major subgroups include the simple **Batzellines**, **Makaluvamines** and **Damirones**, and those in which the pyrroloquinoline nucleus forms part of a larger skeleton such as in the **Discorhabdins** and the closely related **Epinardines**. They are thought to be biosynthesised from tyrosine via **3,5-Dibromoverongiaquinol**.



Batzelline A

Ding, Q. *et al.*, *Curr. Med. Chem.*, 1999, **6**, 1–27 (rev)

MISCELLANEOUS INDOLE ALKALOIDS (VX6100)

Under this heading are listed indole alkaloids for which a classification remains premature, either because they currently belong to a very limited subtype, and/or because there has been no work on their biosynthesis. Many of them probably represent minor branches on the biosynthetic pathways to the larger groups; for example the **Rhopaladins** are probably biosynthetically close to the indolo[2,3-*α*]carbazole alkaloids (VX4350), but this is not yet established experimentally. **Granulatimide** may represent a lower benzologue of these alkaloids incorporating a histamine residue in place of one tryptamine. Plant alkaloids representing variations in skeleton of which only one or two examples are known include **Pandine**, **Iboxyphylline** and **Iboxyphyllidine** (all skeletal variations apparently related to the Pandoline type). Others are degraded indole alkaloids of unknown parental type.


Somei, M. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 278–311 (rev)
Kawasaki, T. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 761–793 (rev)

TERPENOID ALKALOIDS (VX6240–VX6500)

Further information on terpenoid structures and biosynthesis generally can be found in the *Dictionary of Natural Products*.

MONOTERPENOID ALKALOIDS (VX6240)

This is a diverse group. The carbon skeleton is mostly C₁₀, but in many alkaloids it is C₉ and in some it is C₁₁. There are two major groups; those derived from iridodial-like precursors, e.g. **α-Skytanthine**, and those derived from Secologanin, typified by **Gentianine**, **Bakankoside**, and **Gentioflavine**. However, this heading also covers a variety of miscellaneous amines and amides based on small terpenoid skeletons, e.g. **Alfileramine** (menthane-type) and **Fontaineine** (simple acyclic monoterpene).



Gentianine


Cordell, G.A., *Alkaloids*, 1998, **52**, 260–327 (rev)

SESQUITERPENE ALKALOIDS (VX6300)


A variety of sesquiterpene alkaloids have been isolated from plant (e.g. **Guaipyridine**; **Muscicapines**) and marine sources and are nearly all relatively simple amine or amide derivatives of well-known sesquiterpenoid (or meroterpenoid) skeletons. They may be acyclic (e.g. **Axinyssimides**), monocyclic (**Smenospongine**) or bicyclic (**Nakijiquinones**).

MACROCYCLIC SESQUITERPENE ALKALOIDS (VX6320)

This group of plant alkaloids contains the ring system of dihydroagarofuran, a sesquiterpene of the eudesmane group, esterified with nicotinic acid, or with any of several dicarboxylic acids. Most of the alkaloids, which occur in *Euonymus* and *Maytenus* species among others, contain a medium ring dilactone involving one of the dicarboxylic acids; **Evonine** is typical. Some alkaloids with two dilactone medium rings have also been isolated.



Dihydroagarofuran




Per(deacyl)euojaponin

In the dictionary entries, series of these alkaloids which are esters of the same basic skeleton are usually shown as derivatives of the parent polyol, such as **Per(deacyl)euojaponin** (parent alkaloidal polyol of the **Euojaponins** and of various other alkaloids, e.g. **Cathedulins**). The names chosen for these parent structures (often uncharacterised as such) in the Dictionary are chemically reasonable, but may not be found elsewhere in the literature.

DENDROBIUM ALKALOIDS (VX6340)

These alkaloids fall biogenetically into two quite distinct groups; a group of sesquiterpene alkaloids typified by **Dendrobine**, from *Dendrobium nobile*, with variants involving oxygenation at C-2 or C-6, and fission of the


nitrogen to C-2 bond, and a group of indolizidine bases exemplified by **Crepidamine** from *Dendrobium crepidatum*. These are probably not terpenoid in origin, and may be derived from shikimic acid, acetate, and ornithine.



Dendrobine

NUPHAR ALKALOIDS (VX6360)

The *Nuphar* (water-lily) alkaloids contain a normal sesquiterpene carbon skeleton, and can be divided into two main sub-groups: furylpiperidines, e.g. **Nuphamine**; and furylquinolizidines, e.g. **Deoxynupharidine**. The latter group includes a number of sulfur-containing dimers such as **Neothiobinupharidine**, plus a few miscellaneous bases. A number of these alkaloids have also been isolated from the scent glands of beavers, *Castor* spp., acquired presumably by a dietary route.



Nuphamine


Wróbel, J.T., *Alkaloids*, 1977, **16**, 181–214 (rev)

Cybulski, J. *et al.*, *Alkaloids*, 1989, **35**, 215–257 (rev)

Michael, J.P., *Nat. Prod. Rep.*, 2004, **21**, 625–649; 2005, **22**, 603–626; 2008, **25**, 139–165 (revs)

C₁₉ AND C₂₀ DITERPENOID ALKALOIDS AND 4-NOR ANALOGUES (VX6400, VX6420)

These alkaloids are very numerous and include some highly toxic bases with strong physiological properties, such as **Aconitine**. Although they are obviously diterpenoid in origin, few biogenetic studies have been reported, apart from relatively early reports of the incorporation of acetate and mevalonate into **Browniine** and **Lycoctonine**, and of mevalonate and glycine into **Delcosine**. A recent hypothesis suggests their formation directly from the complete diterpenoids by a process involving Mannich reactions.



Aconitine

The first group (VX6400) is based on the C₁₉ aconitane ring system, characteristic of the Ranunculaceae. Alkaloids of this group are fairly homogeneous structurally and differ only in the pattern of substitution by hydroxy, methoxy, acetyloxy, benzoyloxy and, frequently, anthranilate-based groups (e.g. **Delsemines**), although recently other structural variations such as unsaturation and epoxidation have been found.

The simpler alkaloids are collected in the Dictionary under the parent aconitane skeleton. Note that in the CAS parent aconitane, C-18 is missing, and following CAS nomenclature, the parent skeleton is 4-methylnaconitane. The extra carbon atom is numbered C-18.

A few alkaloids belong to the heteratisane group, formed from the aconitane framework by oxidative fission of the 13,14-bond.

The second major group are the C₂₀ alkaloids (VX6420), based on atidane and more widely distributed among dicot families. Few alkaloids, as it happens, are based on the parent ring system, since many skeletal variations

are known. These include the hetisans (e.g. **Davisine**), in which additional rings are introduced into the atidane ring system by formation of 14,20 and *N*,6 bonds. The unmodified hetisans are presented in the Dictionary under the systematic hetisan name. There is a small group of atidane 7,20 cyclic ethers (**Ajaconine**) and various other modifications involving ring fission, contraction, migration etc. of the hetisan skeleton (**Denudatine**, **Delnudine**, **Cardionidine**, etc.)

Another major modification leads to the veatchine skeleton, as in **Cuauchichicine**. There are also 7,20-Cycloveatchine bases, e.g. **Lucidusculine**, 14,20-Cycloveatchine bases, e.g. **Anopterine** and miscellaneous bases.

Pelletier, S.W. *et al.*, *Alkaloids*, 1979, **17**, 2–104; 1981, **18**, 100–216 (*revs*)

Benn, M.H. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1983, **1**, 153–210 (*toxicity, pharmacology*)


Amiya, T. *et al.*, *Alkaloids*, 1988, **34**, 95–179 (*rev*)

Joshi, B.S. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1999, **13**, 289–370; 2001, **15**, 1–414 (*revs, pmr, cmr*)

Wang, F. P. *et al.*, *Alkaloids*, 2002, **59**, 1–280 (*rev*)

ERYTHROPHLEUM ALKALOIDS (VX6460)

The alkaloids of *Erythrophleum* spp. (Fabaceae) are based on the diterpene skeleton related to Cassaic acid. The oxidation pattern is relatively simple, involving only *C*-3, *C*-6, *C*-7 and *C*-19. In most alkaloids, *C*-19 is at the carboxylic acid oxidation level. All the alkaloids are esters or amides of a *C*-16 carboxylic acid with *N,N*-dimethylethanolamine or *N*-methylethanolamine. **Erythroplamine** is a typical example. There have been some confusing structure revisions in this series; they have been little studied recently.




Erythroplamine

Morin, R.B., *Alkaloids*, 1967, **10**, 287–303 (*rev*)

PENITREMS (VX6470)

This heading covers indoloterpenoid metabolites from microorganisms including the **Penitrems**, **Emindoles**, **Nodulisporic acids** and others. They are mostly toxic, many with tremorigenic activity. The intermediary of **Emindole DA** and **Paxilline** in their biosynthesis from geranylindole has been recently demonstrated. **Shearinine I** is an interesting ring-cleaved compound which has lost the indolo functionality.



Steyn, P.S., *Prog. Chem. Org. Nat. Prod.*, 1985, **48**, 1–80 (rev)
Sings, H. *et al.*, *Alkaloids*, 2003, **60**, 51–163 (rev)
Fueki, S. *et al.*, *Org. Lett.*, 2004, **6**, 2697–2700 (biosynth)

MISCELLANEOUS DITERPENOID ALKALOIDS (VX6480)

This category includes diterpenes linked by an ester function to a non-terpenoid nitrogen-containing unit. Examples are **Ryanodine**, **Concavine** and the **Decaturins**. The **Agelasines**, **Agelasimines** and related alkaloids are a series of guanidinoid- or purinoid- substituted simple acyclic, mono- or bi-cyclic terpenoids, and there are one or two simple amino-substituted cembranoids.

Appendino, G. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1996, **11**, 237–268 (rev, *Taxus alkaloids*)


SESTERTERPENE ALKALOIDS (VX6490)

A few nitrogen-containing sesterterpenoids are present in marine organisms, however, the number so far characterised is relatively low in comparison with the number of nitrogen-free tetracyclic sesterterpenes, which are common in marine organisms. The **Spongidines** and the **Molliorins**, in which a ring-E pyridine or pyrrole ring, respectively, is fused to one of the common tetracyclic skeletons, are the main types. There are also linear pyrrolic compounds including the **Sarcotrines** and **Sarcotragins** which are considered to have a common biosynthetic origin, the latter being degraded norsesesterterpenoids.

Liu, Y. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 630–651 (rev)

DAPHNIPHYLLINE ALKALOIDS (VX6500)

The alkaloids of *Daphniphyllum* species constitute a unique group of complex bases derived from squalene. They are structurally diverse and a number of different numbering schemes have been used in the literature. The structures given in this Dictionary and their numbering have been revised to follow the scheme given by Kobayashi and Morita (2003), which is based on current knowledge about their biosynthesis. This system is based on various cyclisations of squalene-like intermediates, which have been modelled with success *in vitro*. The origin of the nitrogen atom is not certain; it may come from pyridoxamine or from an amino acid.



Daphniphylline

The first Edition of this Dictionary distinguished six structural subtypes of *Daphniphyllum* alkaloid. However, since new structural variations continue to be discovered regularly (e.g. **Bukittinggine**, **Calydaphninone**, containing a unique 4-membered ring), it is best to browse the individual entries or to consult the reviews listed below.

Yamamura, S., *Alkaloids*, 1986, **29**, 265–286 (rev)
Kobayashi, J. *et al.*, *Alkaloids*, 2003, **60**, 165–205 (rev, biosynth)
Jin, Z., *Nat. Prod. Rep.*, 2003, **20**, 608–614 (rev)

STEROIDAL ALKALOIDS (VX6640-VX6790)

The large number of steroidal alkaloids may be divided into several subgroups, of which the simplest are various types of aminopregnanes. Steroidal alkaloids are typically from higher plants, and are most numerous in the Apocynaceae, Liliaceae, Buxaceae and Solanaceae, however, some are from other sources, including amphibians.

A few naturally occurring steroidal aminoglycosides have been found (e.g. Holacurtinol), but these are not included in the Dictionary.




STEROIDAL ALKALOIDS (SALAMANDRA TYPE) (VX6640)

The small class of Salamandra-type alkaloids were isolated from amphibians, and are ring-A modified, mostly into aza-homosteroids, except for **Cycloneosamandione**.

Habermehl, G., *Alkaloids*, 1967, **9**, 427–440 (rev)

STEROIDAL ALKALOIDS (JERVERATRUM, SPIROSOLANE AND SOLANIDINE TYPES) (VX6660, VX6720, VX6740)

These are alkaloids in which the side-chain of a C_{27} steroid has been converted into a piperidine ring. In the jerveratrum (VX6660) (e.g. **Jervine**) and spirosolane (VX6720) (e.g. **Soladulcidine**) types, the piperidine ring is part of a spiro system, in the latter type forming an aza-analogue of the numerous and well-known spirostane steroids. Like the spirostanes, they show variation in the stereochemistry of the two new centres of chirality formed in the side-chain. In the solanidine type (VX6740), e.g. **Rubijervine**, an additional ring is formed by C-16 to N bond formation. They are present in the Solanaceae, and are of significance as potential trace toxins present in potatoes, tomatoes and other food crops.



Tomko, J. *et al.*, *Alkaloids*, 1973, **14**, 1–82 (rev)


Ripperger, H., *Alkaloids*, 1981, **19**, 81–192 (rev, *Solanum alkaloids*)

Ripperger, H. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1982, **12**, 103–186 (rev, *Solanum alkaloids*)

Keeler, R.F. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1986, **4**, 389–425 (rev, *teratogenic steroidal alkaloids*)

STEROIDAL ALKALOIDS (CERVERATRUM (CEVANE) TYPE) (VX6680)

In this group of alkaloids, from Monocot families such as the Orchidaceae, there has been an abeo shift from a Procevine-type precursor, in which ring D has been enlarged at the expense of ring C. They show extensive stereochemical variation within the basic cevane skeleton, and the stereochemistry of the various alkaloids is still being studied. See for example the numerous stereoisomers of **3,6-Cevanediol**.



Cevane (IUPAC)

The semisystematic nomenclature of cevanes, especially unsaturated ones, has shown some annoying minor variations over the years, e.g. Cev-5-enine, Cev-5-enene, Δ^5 -Cevenine, 4,5-Didehydrocevine, etc. In this Dictionary the shortest form Cev-5-ene is used.

Ussuriedine represents an interesting tricyclic variation.

Tomko, J. *et al.*, *Alkaloids*, 1973, **14**, 1–82 (rev)

STEROIDAL ALKALOIDS (CONANINE TYPE) (VX6700)

Alkaloids containing the conanine skeleton occur in the Apocynaceae. Nearly all of these bases contain an amino or an oxygen function at C-3. There are some skeletal variations, such as C-4 methylation (**Kurchollessine**) and 9,10-seco-A ring aromatisation (**Didymeline**).

Atta-ur-Rahman *et al.*, *Alkaloids*, 1988, **32**, 79–237 (rev)

STEROIDAL ALKALOIDS (BUXUS TYPE) (VX6760)

The *Buxus* alkaloids are a large group of bases, the majority of which are pentacyclic 4,4,14-trimethyl-9,19-cyclopregnanes. A smaller number contain a tetracyclic system in which 9,10 bond fission has occurred to give a seven-membered ring B. In some, one or both of the carbon atoms attached to C-4 have been lost. All the alkaloids have a nitrogen function at C-3 and/or C-20, which may be unmethylated, partially methylated, or fully methylated. Further minor structural variations continue to be isolated from various *Buxus* spp. The suffix letters used in the nomenclature of this group indicate the degree of methylation of the nitrogen atom(s):

In this Dictionary, the entries for these alkaloids are organised under the (usually unknown) unsubstituted parents of the I, O, or P type.

The recent characterisation of the **Cortistatin** alkaloids from a *Corticium* sponge, represent a new development. Structurally their 9(10→19)-abeosteroid moiety resembles that of one subgroup of the terrestrial *Buxus* alkaloids, but the presence of an isoquinoline ring system at the other end of the molecule is without precedent. They are coded as Buxus alkaloids on structural grounds, although presumably this is a case of evolutionary parallelism.

Černý, V. *et al.*, *Alkaloids*, 1967, **9**, 305–426 (rev)

Tomko, J. *et al.*, *Alkaloids*, 1973, **14**, 1–82 (rev)

Atta-ur-Rahman *et al.*, *Alkaloids*, 1988, **32**, 79–237 (rev)

Ata, A. *et al.*, *Alkaloids*, 2008, **66**, 191–213 (rev)


STEROIDAL ALKALOIDS (PREGNANE TYPE) (VX6780)

These are fairly numerous and occur in the Apocynaceae, Buxaceae and Didymelaceae. The majority contain amino or related functions at C-3 and/or C-20, and often at other positions.

STEROIDAL ALKALOIDS (CEPHALOSTATINS/RITTERAZINES) (VX6785)

An important new group of marine steroidal metabolites isolated since the 1980s are the **Cephalostatins/Ritterazines**, which are dimeric steroidal pyrazines (also coded as pyrazine alkaloids). One of their representatives is **Ritterazine A**. They show significant biological activities, but are isolable only in small amounts and are the subject of intense synthetic efforts.

Flessner, T. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 2004, **87**, 1–80 (rev)



Ritterazine A

MISCELLANEOUS STEROIDAL ALKALOIDS (VX6790)


This includes non-nitrogenous steroids linked by an ester or acetal bond to a nitrogen containing unit, as in **Bufotoxin**, the **Batrachotoxins** (elaboration of a basic pregnane theme), and a small number of marine steroidal alkaloids of novel type such as the **Plakinamines**, as well as biologically active D-ring azahomosteroids based on **15-Azasterol**. **Petisidine**, **Tomatillidine** and their congeners are the only group of steroidal alkaloids containing a pyrrolidinyl side-chain. A 7-membered azepinyl ring structure was proposed for certain alkaloids such as **Edpetilidinine**, but most of these have now been reassigned as piperidines.

AZAANTHRACENE AND AZAFLUORENE ALKALOIDS (VX6820, VX6840)

There have been a number of structural reassignments in these groups, and some structures are still uncertain. It now appears that they are related to each other, and to the azaaporphine alkaloids (VX2830) above, via the azaanthraquinone intermediate as shown in the biosynthesis of Sampangine. They are therefore currently misplaced in the Type of Compound classification scheme.

PYRAZOLE ALKALOIDS (VX6900)


This is a small scattered group containing a few miscellaneous azapurine bases and simple hydrogenated pyrazoles. The most interesting alkaloid is **Withasomnine**. Little is known of its biosynthesis. Some early tracer studies showed incorporation of phenylalanine and ornithine, but at very low levels, and the significance of this is unclear. There is one marine example, **1-Methyl-1H-pyrazole-5-carboxylic acid**, from a gorgonian.



Withasomnine

IMIDAZOLE ALKALOIDS (VX6920)

This group contains many structurally diverse examples, especially among marine and microbial alkaloids. There are various bases obviously containing a histamine moiety, e.g. **Casimiroedine**, and some more complex plant alkaloids. The bases isolated from marine sources include **Aplysinopsin** and related alkaloids and the **Rhopaladins**. They show a range of biological activities and considerable pharmaceutical potential. The most pharmacologically significant is **Pilocarpine**, obtainable only from *Pilocarpus* spp. (Rutaceae). Two possible biosynthetic pathways were suggested; the most probable involves condensation of a polyketide precursor of the lactone ring with a late intermediate in histidine biosynthesis as the source of the imidazole. This work of the 1970s does not appear to have been followed up more recently. **Anosmine** is biosynthesised from lysine.



Casimiroedine


Hill, R.K., *Alkaloids Chem. Biol. Perspect.*, Pelletier, 1982, **2**, 49–104 (rev)

Maat, L. *et al.*, *Alkaloids*, 1983, **22**, 282–334 (rev, biosynth)

Jin, Z. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 464–496 (rev)

PYRROLE-IMIDAZOLE ALKALOIDS (VX6922)

These form a varied class of marine alkaloid exemplified by **Oroidin** which with various relatives (e.g. the dimeric **Sceptrine**) appears to be a genuine sponge product. Their significance appears to be as key intermediates in the biogenesis of many of the more complex marine alkaloids. A recent biogenetically-based classification (see Hofmann) recognises a class of nearly 100 pyrrole-imidazole alkaloids arising in various ways from this key intermediate and containing 0–4 further rings, additional to the pyrrole and imidazole rings already present. Future organisation of this database may recognise this category; at present, many of these are placed in the miscellaneous categories VX9000–VX9400.




Oroidin

Hofmann, H. *et al.*, *Synthesis*, 2003, 1753–1783 (rev)

Walsh, C.T. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 577–531 (rev, biosynth)

CYCLOHEPTADIIMIDAZOLES (ZOANTHOXANTHINS) (VX6925)


These are a group of marine alkaloids which are condensed imidazoles closely related in structure, and which in this Dictionary are grouped under just two entries; those for **Pseudozoanthoxanthin A** and **Parazoanthoxanthin A**. Their biosynthesis does not appear to have been studied in detail, however, they are presumably derived from histidine.



Pseudozoanthoxanthin A

OXAZOLE AND BENZOXAZOLE ALKALOIDS (VX6930)

Numerous naturally occurring oxazoles are known. Structurally they are diverse and may be open-chain (e.g. **Bengazoles**) or macrocyclic (e.g. **Patellamides**). Like the imidazoles, they show considerable drug potential. They have been isolated from various sources—plants of the Poaceae (e.g. **Annuloline**) and Rutaceae (e.g. **Halfordinol**), nudibranch egg masses (**Ulapualides**), algae and microorganisms. The latter have furnished the majority of the alkaloids, ranging from the simple indolyl alkaloids (**Pimprinine**) to complex peptide antibiotics such as the **Virginiamycin** family. The marine and bacterial oxazoles appear to have been formed from peptides of aliphatic amino acids, while the oxazoles of the Poaceae and Rutaceae arise from the chorismic acid-phenylalanine pathway.



Halfordinol

Jacobs, H.M. *et al.*, *Alkaloids*, 1989, **35**, 259–310 (rev)

Pattenden, G., *J. Het. Chem.*, 1992, **29**, 607–618 (rev, synth)

Yeh, V.S.C., *Tetrahedron*, 2004, **60**, 11995–12042 (rev, synth)

Jin, Z. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 464–496 (rev)


ISOXAZOLE ALKALOIDS (VX6932)

This is a fairly small group, most of which are modified amino acids (e.g. **Ibotenic acid**).

Rahboek, L. *et al.*, *Alkaloids*, 2001, **57**, 185–233 (*rev*)

SPIROBENZOXAZOLINE ALKALOIDS (VX6934)

These exclusively marine alkaloids, such as the **Psammaphysins**, are derived by cyclisation of open-chain halogenated tyrosinoid oxime precursors (see VX2008 above). This is probably through the intermediacy of an arene oxide, although attempts to replicate this chemically have given only low yields.



Psammaphysin A

Okamoto, K.T. *et al.*, *Tet. Lett.*, 1987, **28**, 4969–4972 (*biosynth*)

THIAZOLE ALKALOIDS (VX6935, VX6936, VX6937)

A large number of naturally occurring alkaloids that incorporate the thiazole nucleus have now been isolated. Some simple thiazoles (e.g. **2,4-Dialkylthiazoles**; **Benzothiazole**) have been obtained from plant and animal sources, but typically they are microbial products, including from marine sources. These alkaloids are a heterogeneous group ranging in complexity from **Aeruginic acid** and the simple peptide **Herbamide A** to antineoplastic cyclopeptides such as **Ulicyclamide**, **Ulithiacyclamide**, **Patellamides** and **Dolastatins**. The **Latrunculins** (VX6936) are a subclass of mostly macrocyclic tetrahydrothiazoles isolated from sponges. Thiazoles show a range of interesting drug activities.

The biosynthesis of the thiazole ring in the highly bioactive **Curacin A** has been shown to be from cysteine, with the remainder of the carbon chain, including the cyclopropane functionality, being acetate-derived.


Pattenden, G., *J. Het. Chem.*, 1992, **29**, 607–618 (*rev, synth*)

Jin, Z. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 196–229; 2006, **23**, 464–496 (*revs*)

Mulzer, J., *Prog. Chem. Org. Nat. Prod.*, 2009, **90**, 1–237 (*Epothilones, rev, synth, biosynth, pharmacol*)

PYRAZINE AND QUINOXALINE ALKALOIDS (VX6940)

Pyrazines have been isolated from widely differing sources: from microorganisms, plants, mushrooms, animals, insects (especially ants, where they are considered to function as alarm pheromones) and various marine organisms, where they are the actual light emitters in bioluminescence processes. Bioluminescence emitters are found scattered throughout different types of organism, which implies multiple independent origins during the course of evolution. The imidazopyrazine, **Coelenterazine**, and closely related alkaloids though account for the great majority of observed bioluminescence. There is evidence for Coelenterazine *de novo* biosynthesis in shrimps. **Cypridina Luciferin** is biosynthesised from arginine and tryptophan. A series of tetrahydroquinoxalines has been isolated from the scent gland of the Canadian beaver, *Castor fiber*.




Coelenterazine

Pyrazines also contribute to the aroma of various foodstuffs, including coffee, cocoa, tea and cooked meats. However, from these sources they are generated by pyrolytic processes and are not true alkaloids. The cephalostatin group of alkaloids described under VX6785 are steroidal dimers linked by pyrazine formation.

Rees, J.-F. *et al.*, *J. Exp. Biol.*, 1998, **201**, 1211–1221 (*rev, bioluminescence*)

BENZO[1,2-*B*:3,4-*B'*]DIPYRROLES (VX6945)

This is a numerically limited group of alkaloid antibiotics all isolated from Streptomyces. **Duocarmycin SA** appears to be a key biosynthetic intermediate, however, detailed biosynthetic studies have not been reported. They are DNA alkylating agents. Some members, e.g. **Rachelmycin**, are cyclopropa-compounds, however, this feature does not appear to be necessary for activity.



Duocarmycin SA

PYRROLO[1,2-*A*]PYRAZINES (VX6950)

These are mostly cyclic anhydrides of proline plus another amino acid residue. Their inclusion in this Dictionary is a marginal decision. Other cyclodipeptides not based on proline are not included; they are classified under the amino acid code VV0150 (diketopiperazine) and are found in the *Dictionary of Natural Products*.

MORPHOLINE ALKALOIDS (VX6955)


These are relatively few in number, e.g. **Agropine**, **Chelonins A and C** and the **Oxazinines**.

PYRIMIDINE ALKALOIDS (VX6960)

Simple thymine-derived bases such as **Thyminol** are found in various organisms, together with some more highly elaborated substances such as the **Meridianins**. The category also includes miscellaneous hydrogenated pyrimidines, e.g. **Aberiamine**.

PTILOCAULINS (VX6970)

This group is based on the (partially hydrogenated) cyclopenta[*de*]quinazoline skeleton, which contains a pyrimidine nucleus. It comprises **Ptilocaulin** and relatives, the **Mirabilins** and a few other related alkaloids, mostly isolated from sponges, but which may be symbiont metabolites. It has been suggested that these compounds arise by a late-stage addition of guanidine to a polyketide precursor.




Ptilocaulin

Snider, B.B. *et al.*, *J. Org. Chem.*, 1993, **58**, 3828–3839 (*biosynth*)

TRIAZAACENAPHTHYLENE ALKALOIDS (VX6980)

This is another group based on a tricyclic pyrimidine-related nucleus, and includes the **Batzelladines** and the **Crambescidins**. They are isolated from sponges and echinoderms. Their biosynthesis does not appear to have been


studied in detail. However, they are formally composed of a guanidine unit, with which the biological activity is associated, embedded in a linear unbranched perimeter, presumably derived from a lipid precursor.



Batzelladine A

TETRODOTOXINS (VX6990)

Tetrodotoxin and its homologues are well-known and important causes of human poisoning and are known principally from certain tissues of the Japanese puffer fish, and also found in other organisms; they are now thought to be *Pseudomonas* products. They have been extensively reviewed. The carbocycle ring in tetrodotoxins may be of isoprenoid origin.



Tetrodotoxin

Miyazawa, K. *et al.*, *J. Toxicol., Toxin Rev.*, 2001, **20**, 11–25 (*rev*)

Yotsu-Yamashita, M., *J. Toxicol., Toxin Rev.*, 2001, **20**, 51–66 (*rev*)

Daly, J.W., *J. Nat. Prod.*, 2004, **67**, 1211–1215 (*rev, occur*)

PHENAZINE AND PHENOXAZINE ALKALOIDS (VX7000, VX7005)

An increasing number of simple phenazines (e.g. the **Pelagiomicins**) and phenoxazines have been reported from microorganisms, including marine bacteria from sponges. There are also some more complex diphenazines such as **Saphenamycin** and the **Esmeraldines**. These, as well as the simpler phenazines, are biosynthesised by the shikimic acid pathway.

Van't Land, C.W. *et al.*, *J. Org. Chem.*, 1993, **58**, 6576–6582 (*biosynth*)

SIMPLE PYRROLE ALKALOIDS; MISCELLANEOUS PYRROLE ALKALOIDS (VX7010, VX7014)

The category VX7010 includes very simple brominated pyrroles (e.g. **2,3-Dibromo-1H-pyrrole**, which come from a variety of sources. Marine examples include the dimeric **Tambjamines**. Pyrrole-imidazole alkaloids are now listed above under VX6922, while the metabolically important tetrapyrroles, (chlorophyll, haem and their relatives) are not included in this Dictionary; consult the parent *Dictionary of Natural Products*. The pyrrole nucleus is a component of the indole alkaloids, but whereas these are tryptophan-derived, the polypyrroles are biosynthesised from 5-Amino-4-oxopentanoic acid (5-aminolaevulinic acid, ALA). Lower molecular weight alkaloids may arise from either route, or from proline/ornithine pathways via pyrrolidines (see above under VX0300); in the **Prodigosins**, each of the three pyrrole residues is biosynthesised differently. The process of biohalogenation differs between Cl and Br. Chlorination takes place oxidatively by a combination of chloride ion, dioxygen and reduced FAD producing a Cl⁺ equivalent, whilst bromination is mediated by vanadium bromoperoxidases.

- LeQuesne, P.W. *et al.*, *Alkaloids Chem. Biol. Perspect.*, 1999, **13**, 210–258, 237–288 (*revs*)
 Gossauer, A., *Prog. Chem. Org. Nat. Prod.*, 2003, **86**, 1–188 (*rev*)
 Berlinck, R.G.S., *Prog. Chem. Org. Nat. Prod.*, 2004, **87**, 2–188 (*rev*)
 Walsh, C.T. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 517–531 (*rev, biosynth*)

POLYAMINE ALKALOIDS (VX7018, VX7020, VX7025, VX7050, VX7060, VX7080, VX7085, VX7090)

These categories cover the various types of alkaloid containing one or more $-N(CH_2)_nN-$ groups, with n usually 3 or 4, less often 2 or 5. A useful subclassification scheme uses PA numbers defining the constituent diamine units, e.g. **Spermine** $H_2N(CH_2)_3NH(CH_2)_4NH(CH_2)_3NH_2$ is the parent substance of the PA343 subclass. They show markedly different properties (reactivity, basicity, complexing ability, mass spectrometry) from monobasic amines. Alkaloids based on the ornithine-derived **Putrescine** (PA4 alkaloids) (VX7020) are common in terrestrial plants where they are mostly simple cinnamoylputrescines, e.g. **Subaphylline**. Some have been characterised from marine organisms, such as the **Monodontamides** from gastropods. Numerous other alkaloids, again, many terrestrial and some marine, are derived from **Spermine** (PA343) (VX7080) or **Spermidine**, (PA34) (VX7050) themselves derived (in plants, certainly) from ornithine via putrescine. See for example the **Penaramides** and the unusual medium-ring azaaliphatic **Motuporamines**. Classes PA33, PA35, PA44 and PA33433 have also been found in nature, but less frequently. The largest number are the arachnid toxins, such as the extensive series of **Agel toxins** from the spider *Agelenopsis*. PA structures also occur in more complex alkaloids such as **Crambescidin 816** (PA34 substructure) and as steroidal conjugates, in **Squalamine** and related compounds from fish livers (PA34 and PA343 substructures). In terrestrial plants it is well established that these alkaloids derive from the decarboxylation and coupling of amino acids.


The spermidine group includes a variety of medium ring cyclic examples, e.g. **Codonocarpine**, **Lunaridine**, **Cannabisativine**. The PA44 class, the so-called homospermidine alkaloids, includes derivatives of the PA44 amine **Solamine** isolated from the Solanaceae.

- Guggisberg, A. *et al.*, *Alkaloids*, 1983, **22**, 85–188; 1997, **50**, 219–256 (*revs*)
 Schafer, A. *et al.*, *Alkaloids*, 1994, **45**, 1–125 (*arthropod toxins*)
 Mueller, A.L. *et al.*, *Alkaloids*, 1995, **46**, 63–94 (*toxins, pharmacol*)
 Bienz, S. *et al.*, *Alkaloids*, 2002, **58**, 83–338 (*rev*)
 Bienz, S., *Nat. Prod. Rep.*, 2005, **22**, 647–658 (*rev*)
 Estrada, G., *Nat. Prod. Rep.*, 2007, **24**, 145–161 (*rev, spider toxins*)

ANSA-PEPTIDE ALKALOIDS (VX7100)

Peptides consisting only of unmodified proteinaceous amino acid chains, whether linear or cyclic, lie outside the scope of this Dictionary. Among the marine natural products there are a considerable number of peptide-like alkaloids containing highly modified residues, e.g. the Lissoclinamides. These are classified elsewhere (in the case of the Lissoclinamides, under VX6737, macrocyclic thiazole alkaloids).

A large class of peptide derivatives occurring exclusively in higher plants, particularly in the Rhamnaceae, are the ansa-peptide alkaloids, in which a medium-sized ring is formed by meta- or para-substitution of an aromatic peptide residue, as in **Frangulanine**. Many of these alkaloids incorporate D-amino acid residues, but the stereochemical information concerning them is sometimes incomplete. They have however been fully reviewed, with extensive tabulations.




Frangulanine

- Schmidt, U. *et al.*, *Alkaloids*, 1985, **26**, 299–326 (*rev*)
 Itokawa, H. *et al.*, *Alkaloids*, 1997, **49**, 301–387 (*rev*)
 Gournelis, D. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1998, **75**, 1–180 (*rev*)

Tan, N.H. *et al.*, *Chem. Rev.*, 2006, **106**, 840–895 (rev)
El-Seedi, H., *et al.*, *Phytochem. Rev.*, 2007, **6**, 143–165 (rev)

AMANITA ALKALOIDS (VX7120)

The toxins of the European death cap mushroom *Amanita phalloides* and other *Amanita* spp. constitute a complex group of macrocyclic peptides, mostly containing sulfur. These include the amatoxins (e.g. **α -Amanitin**, **β -Amanitin**), the phallotoxins (e.g. **Phalloidin**) and the virotoxins (e.g. **Viroidin**). They have been little studied recently apart from a new synthesis of Phalloidin.




OTHER CYCLOPEPTIDE ALKALOIDS (VX7150)

This is a heterogeneous group.

PYOVERDINS (VX7170)

These are a fairly extensive and increasing series of peptide-alkaloidal siderophore pigments secreted by plant-growth promoting rhizobacteria *Pseudomonas* spp. under conditions of restricted iron availability. Their chelating properties protect the plant against attack by fungi and other pathogens by depriving them of iron. Although they were described in the nineteenth century, the first structure was not elucidated until 1981; by 1999 about 50 were known, some still with only partially determined structures.



The chromophore is a pyrimido[1,2- α]quinoline system, sometimes modified as in the **Azotobactins**. The chromophoric properties of this heterocycle are associated with the predominance of a zwitterionic form having a positively charged quaternary nitrogen with delocalisation of charge over the two *N*s, but for clarity they are all shown in the Dictionary in the uncharged form. The chromophore is linked to a short nitrogenous substituent – NHR (R = usually a dioic acid, sometimes glycine) and an oligopeptide chain X, giving a typical total molecular weight of about 1000-1500. The peptide chain may be cyclic or acyclic, and may contain one or more unusual modified amino acid residues, such as 3-Amino-1-hydroxy-2-piperidinone (*N*-hydroxycycloornithine).


Caution; Compilation of Dictionary entries for this group has been complicated by inconsistencies in the literature concerning their structures and comparison with the molecular formulae, which are often not reported. Some inconsistencies remain in the Dictionary entries; checking continues.

Budzikiewicz, H., *Prog. Chem. Org. Nat. Prod.*, 2004, **87**, 81–237 (rev)

PYRROLO[2,3-D]PYRIMIDINE ALKALOIDS (VX7200)

This is a numerically limited group consisting of the nucleosides **Tubercidin**, the **Mycalisines** and related bases and some simple halogenated pyrrolopyrimidines, as well as the **Rigidins**. They have been isolated from marine

organisms, and appear to be microbial products. It has been shown that the pyrimidine ring is derived intact from a purine base precursor which undergoes ring opening and reformation with the incorporation of a C₂ fragment to form the pyrrole ring.




PURINE ALKALOIDS (VX7300)

Purines are involved along with pyrimidines as bases in DNA and RNA. The ubiquitous, well-known oxypurines, exemplified by **Caffeine**, are usually considered alkaloidal, alongside adenine derivatives such as the plant hormone, **Zeatin**; and various miscellaneous bases. The well-known purines, **Xanthine**, **Inosine**, **Theobromine** and **Theophylline**, long known as plant alkaloids, have now been found in marine organisms along with substituted purines (e.g. **Phidolopin**) and a range of terpenoid-purine alkaloids, such as the **Agelines** and others.

Rosemeyer, H., *Chem. Biodiversity*, 2004, **1**, 361–401 (rev)

PTERIDINE ALKALOIDS (VX7350)

Pteridines are a widely distributed class of naturally occurring alkaloids. They owe their exceptional position in the field of heterocyclic chemistry mainly to their unusual chemical properties, their conspicuous fluorescence and their importance in metabolism, and partly to their discovery as pigments in butterfly wings. Three of the most common butterfly pigments are **Leucopterin**, **Xanthopterin** and **Isoxanthopterin**. The red pigments in the eye of the fruitfly *Drosophila melanogaster*, e.g. **Drosopterin**, **Isodrosopterin** and **Neodrosopterin** are more complex pteridines. **Folic acid**, a water-soluble growth factor in bacteria and an anti-pernicious anaemia factor in animals is also a pterin derivative with a *p*-aminobenzoylglutamic acid sidechain at the 6-position. It occurs naturally as the dihydro derivative. Marine pteridines are represented by **Leucettidine**, the **Urochordamines** and **Isoxanthopterin**.



The biosynthesis of pteridine pigments has been well studied in terrestrial organisms. As in the case of the pyrrolopyrimidines above, the imidazole ring is derived unchanged from a purine precursor, in this case guanosine triphosphate, and the pyrazine ring is built by ring fission and incorporation of two carbons derived from a ribose fragment.


Pfleiderer, W., *J. Het. Chem.*, 1992, **29**, 583–605 (rev)

Chemistry and Biology of Pteridines and Folates, (eds. Milstien, S. *et al.*), Kluwer, 2002

SAXITOXINS (VX7400)

The saxitoxins are a group of highly important modified purinoid toxins known as Paralytic Shellfish Toxins or PSTs. Originally characterised as a cause of dangerous shellfish poisoning, they are now known to be biosynthesised only by dinoflagellates (and some freshwater cyanobacteria), although it is possible that the red

alga *Jania* also produces them. Their presence in shellfish tissues probably results by diffusion according to a yet unknown mechanism. Their exact biosynthesis is not yet known, but it appears from radiolabelling experiments that the two guanidine groups derive from arginine and the carbamoyl side-chain from methionine.



Saxitoxin

Shimizu, Y., *Prog. Chem. Org. Nat. Prod.*, 1984, **45**, 235–264 (rev)

Garson, M.J., *Chem. Rev.*, 1993, **93**, 1699–1733 (rev, biosynth)

Llewellyn, L.E., *Nat. Prod. Rep.*, 2006, **23**, 200–222 (rev)

ALKALOIDAL METAL COMPLEXES (VX7450)


Alkaloidal chelates are additionally indexed here for ease of reference. The metals which have been identified so far in alkaloidal complexes are iron, copper, molybdenum and nickel.

NAPHTHYRIDINOMYCINS (VX7700)

This is a small group of microbial products, and labelling experiments have demonstrated the incorporation of tyrosine, ornithine and methionine.

SAFRAMYCINS (VX7800)

These are representative of the group known as the tetrahydroisoquinoline antitumour antibiotics, including some having slightly different skeletons, such as **Lemonomycin**. They are microbial products produced by *Streptomyces* spp. Close analogues such as the **Renieramycins** and the **Ecteinascidins** were isolated from marine organisms (sponges or tunicates) and are all presumably microbial products. The skeleton (studied for **Saframycin A**) is derived from two tyrosine molecules and the methyl groups from methionine.



Saframycin A

Arai, T. *et al.*, *Alkaloids*, 1983, **21**, 56–100 (rev)

Murkami, Y. *et al.*, *J. Biol. Chem.*, 1985, **260**, 344–348 (biosynth)

MISCELLANEOUS ALKALOIDS WITH 0-4+ RINGS (VX9000, VX9100, VX9200, VX9300, VX9400)

These categories are self-explanatory. They include alkaloids whose relationships with the main groups has not been established, and individual alkaloids or small groups of alkaloids which are clearly related to one of the main groups, but which contain a modified carbon skeleton and are not numerous enough to merit special treatment.

ALKALOIDS OF UNKNOWN STRUCTURE (VX9999)

The Dictionary documents over 1000 alkaloids from the older literature where the structure has remained completely or partially undetermined. In cases where the Authors were able to propose a structural class for the

unknown isolate, the dictionary entry also carries the classification code relevant to that category, e.g. *Buxus* alkaloid.

Many of these isolates will remain forever unidentified, because the original sample is no longer available and the properties reported in the literature are not precise enough for a definite identification. Others, especially from the early literature, were doubtless ill-defined mixtures. However, the number of reinvestigations of these old isolations on a year-by-year basis is now surprisingly small, and renewed investigation of certain groups, for example the Amaryllidaceae alkaloids, using modern methods would probably yield interesting results.

6. DATABASE UPDATES

The database from which this Dictionary is produced, and which is represented by the CD-ROM version, is a specially enhanced subset of the *Dictionary of Natural Products*. Updates to the alkaloid information are being continuously incorporated into the DNP database and released to DNP subscribers. New editions of the *Dictionary of Alkaloids* (probably electronic only) will be published at future intervals.


Subscription information for DNP is available from e-reference@taylorandfrancis.com and www.chemnetbase.com.

Any comments and suggestions for inclusion to DNP may be sent to:

The Editors, DNP
CRC Press
4th Floor, Albert House
1-4 Singer Street
London
EC2A 4BQ

AAL Toxin

A-1



Prod. by *Alternaria alternata* f. sp. *lycopersici*. Host-specific phytotoxin complex. Similar to Fumonisin A₁.

AAL Toxin TA₁ [79367-52-5]

C₂₅H₄₇NO₁₀ 521.647
Sol. H₂O.

5-Deoxy: AAL Toxin TB₁
[149849-90-1]

C₂₅H₄₇NO₉ 505.648

From *Alternaria alternata* f. sp. *lycopersici*. Phytotoxin.

5-Deoxy, N-Ac: AAL Toxin TD₁

[176590-35-5]

C₂₇H₄₉NO₁₀ 547.685

4,5-Dideoxy: AAL Toxin TC₁

[176590-33-3]

C₂₅H₄₇NO₈ 489.648

4,5-Dideoxy, N-Ac: AAL Toxin TE₁

[176590-37-7]

C₂₇H₄₉NO₉ 531.685

AAL Toxin TA₂ [79367-51-4]

C₂₅H₄₇NO₁₀ 521.647
Sol. H₂O.

5-Deoxy: AAL Toxin TB₂

[149849-91-2]

C₂₅H₄₇NO₉ 505.648

From *Alternaria alternata* f. sp. *lycopersici*. Phytotoxin.

5-Deoxy, N-Ac: AAL Toxin TD₂

[176590-36-6]

C₂₇H₄₉NO₁₀ 547.685

4,5-Dideoxy: AAL Toxin TC₂

[176590-34-4]

C₂₅H₄₇NO₈ 489.648

4,5-Dideoxy, N-Ac: AAL Toxin TE₂

[176590-38-8]

C₂₇H₄₉NO₉ 531.685

Bottini, A.T. et al., *Tet. Lett.*, 1981, **22**, 2723 (isol)

Caldas, E.D. et al., *J. Agric. Food Chem.*, 1994, **42**, 327 (struct)

Boyle, C.D. et al., *J.A.C.S.*, 1994, **116**, 4995 (struct)

Oikawa, H. et al., *Tet. Lett.*, 1994, **35**, 1223 (synth)

Oikawa, H. et al., *Tetrahedron*, 1994, **50**, 13347 (abs config)


Shier, W.T. et al., *Tet. Lett.*, 1995, **36**, 1571 (abs config)

Oikawa, H. et al., *Tet. Lett.*, 1999, **40**, 6621-6625 (synth, TA₁)

Aaptamine

A-2

8,9-Dimethoxy-1H-benzo[de][1,6]-naphthyridine, 9CI
[85547-22-4]



C₁₃H₁₂N₂O₂ 228.25

Alkaloid from the marine sponge *Aaptos aaptos*. α-Adrenoreceptor blocker. Antineoplastic agent. Brilliant green cryst; bright yellow cryst. (MeOH/Me₂CO) (as hydrochloride). Sol. MeOH, CHCl₃. Mp 110-113° (107°) (hydrochloride). λ_{max} 215 (€ 13700); 236 (€ 14700); 255 (€ 17900); 309 (€ 3640); 350 (€ 3750); 380 (€ 5000); 394 (€ 4570) (H₂O) (Derep). λ_{max} 220 (€ 8900); 239 (€ 11750); 257 (€ 13200); 274 (€ 9700); 312 (€ 3160); 354 (€ 3300); 384 (€ 5620) (MeOH) (Berdy).

▶DI2410500

N⁴-Me: N⁴-Methylaaptamine

C₁₄H₁₄N₂O₂ 242.277

Alkaloid from *Aaptos aaptos*. Antiviral agent. Pale yellow oil. λ_{max} 216 (€ 1318); 239 (€ 13356); 258 (€ 11025); 269 (sh) (€ 9522); 277 (sh) (€ 8654); 314 (€ 2420); 360 (€ 2939); 394 (sh) (€ 2780) (MeOH).

O⁹-De-Me: 9-De-O-methylaaptamine

[88839-98-9]

C₁₂H₁₀N₂O₂ 214.223

Alkaloid from *Aaptos aaptos*. Cytotoxic and antimicrobial agent. Greenish-yellow powder + 1.5H₂O (as hydrochloride). Mp 248-251° (dec., sealed tube) (hydrochloride). λ_{max} 241 (€ 18300); 313 (€ 3890); 370 (€ 4470); 400 (€ 4130) (H₂O) (Derep).

O⁹-De-Me, N¹-Me: 9-De-O-methyl-N¹-methylaaptamine. Isoaaptamine

[117173-75-8]

C₁₃H₁₂N₂O₂ 228.25

Alkaloid from the sponges *Aaptos aaptos*, *Hymeniacidon* sp. and *Suberites* sp. Antineoplastic agent and β-glucanase inhibitor. Amorph. yellow powder. Mp 200-205° dec.

Di-O-de-Me: Dide-O-methylaaptamine

C₁₁H₈N₂O₂ 200.196

Isol. from *Aaptos* sp. Yellow solid (as TFA salt). λ_{max} 241 (log € 4.43); 267 (log € 4.19); 313 (log € 3.63); 363 (log € 3.67); 402 (log € 3.58) (no solvent reported).

Di-O-de-Me, 9-sulfate: Dide-O-methylaaptamine 9-sulfate

C₁₁H₈N₂O₅S 280.261

Isol. from *Aaptos* sp. Pale yellow solid (as TFA salt). λ_{max} 216 (log € 4.14); 236 (log € 4.12); 257 (log € 4.18); 310 (log € 3.49); 373 (log € 3.63) (no solvent reported).

Nakamura, H. et al., *Tet. Lett.*, 1982, 5555 (isol, uv, pmr, cmr, struct)

Pelletier, J.C. et al., *Tet. Lett.*, 1985, **26**, 1259 (synth, uv, pmr)

Kelly, T.R. et al., *Tetrahedron*, 1985, **41**, 3033 (synth)

Sakamoto, T. et al., *Chem. Pharm. Bull.*, 1986, **34**, 2760 (synth)

Bassoli, A. et al., *Chem. Comm.*, 1987, 150 (synth)

Nakamura, H. et al., *J.C.S. Perkin 1*, 1987, 173 (deriv)

Pelletier, J.C. et al., *J.O.C.*, 1987, **52**, 616 (synth, ir, pmr)

Andrew, R.G. et al., *Tetrahedron*, 1987, **43**, 4803 (synth, uv, ir, pmr, ms)

Hibino, S. et al., *J.C.S. Perkin 1*, 1988, 2429 (synth, pmr)

Fedoreev, S.A. et al., *Khim.-Farm. Zh.*, 1988, **22**, 943-946 (isol)

Sova, V.V. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 420-422 (*Demethyl-N-methylaaptamine*)

Balczewski, P. et al., *J.C.S. Perkin 1*, 1990, **31**, 3193 (synth)

Molina, P. et al., *Synthesis*, 1996, 1199 (synth)

Sugino, E. et al., *Heterocycles*, 1999, **50**, 543-559 (rev, synth)

Walz, A.J. et al., *J.O.C.*, 2000, **65**, 8001-8010 (*Isoaaptamine*, synth)

Coutinho, A.F. et al., *Heterocycles*, 2002, **57**, 1265-1272 (*N⁴-Methylaaptamine*)

Calcul, L. et al., *Tetrahedron*, 2003, **59**, 6539-6544 (isol, pmr, cmr)

Pettit, G.R. et al., *J. Nat. Prod.*, 2004, **67**, 506-509 (*Isoaaptamine*, pmr, cmr, cryst struct)

Pettit, G.R. et al., *J.O.C.*, 2004, **69**, 2251-2256 (*Isoaaptamine*, synth)


Herlt, A. et al., *Tetrahedron*, 2004, **60**, 6101-6104 (*Didemethylaaptamines*)

Larghi, E.L. et al., *Tetrahedron*, 2008, **64**, 5236-5245 (synth)

Aaptosamine

A-3

[219642-41-8]



C₁₅H₁₄N₂O₃ 270.287


Alkaloid from the Red Sea sponge *Aaptos aaptos*. Orange gum. λ_{max} 240 (log € 4.01); 336 (log € 3.93) (MeOH).

Tinto, W.F. et al., *Heterocycles*, 1998, **48**, 2089-2093 (isol, uv, ir, pmr, cmr, ms)

Aaptosine

A-4

1,9-Dimethoxy-8H-5,8-diazabenz[cd]azulene, 9CI
[151041-63-3]



C₁₃H₁₂N₂O₂ 228.25


Trace alkaloid from the Red Sea sponge *Aaptos aaptos*. Cytotoxic agent. Yellow oil. λ_{max} 377 (MeOH/HCl) (Derep). λ_{max} 225 (€ 9700); 346 (€ 3800) (MeOH) (Derep).

Rudi, A. et al., *Tet. Lett.*, 1993, **34**, 4683 (isol, uv, ir, pmr, cmr, ms, struct)

Abbeymycin

A-5

1,2,3,10,11,11a-Hexahydro-2-hydroxy-11-methoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI. Antibiotic SA4-3.
SA4-3
[108073-64-9]
[115793-07-2]




C₁₃H₁₆N₂O₃ 248.281

Anthramycin-type antibiotic from *Streptomyces* sp. Has weak activity against a limited number of anaerobic bacteria. Mp 142-144° dec. [α]_D²⁵ +303 (c, 0.741 in H₂O). Stereochem. of SA4-3 not confirmed. λ_{max} 216 (ε 37200); 236 (ε 18100); 316 (ε 3600) (H₂O) (Derep).

► AA3000000

Hochlowski, J.E. *et al.*, *J. Antibiot.*, 1987, **40**, 145 (isol, uv, pmr, ms, struct)
Japan. Pat., 1987, 87 185 087; *CA*, **109**, 92661m (SA4-3)

2(3→20)-Abeo-2,5,7,10,13-pentahydroxy-4(20),11-taxadien-9-one A-6



C₂₀H₃₀O₆ 366.453

(2α,5α,7β,10β,13α)-form

5-(3-Dimethylamino-3-phenylpropanoyl), 2,7,13-tri-Ac: [240805-52-1]
C₃₇H₄₉NO₁₀ 667.795
Constit. of *Taxus cuspidata*. Gum. [α]_D²⁴ -31 (c, 0.01 in CHCl₃).

5-(3S-Dimethylamino-2S-hydroxy-3-phenylpropanoyl), 13-Ac: **2-Deacetyl-taxine A**. *Taxine C*
[160561-50-2]
C₃₃H₄₅NO₉ 599.72
Isol. from the leaves of *Taxus baccata* (Taxaceae). Amorph. or cryst. [α]_D²⁰ -106 (c, 0.47 in CHCl₃).

5-(3S-Dimethylamino-2S-hydroxy-3-phenylpropanoyl), 2,13-di-Ac: **Taxine A**
[1361-49-5]
C₃₅H₄₇NO₁₀ 641.757
Major constit. of Taxine, the alkaloid mixt. from *Taxus baccata* (yew) (Taxaceae). Taxine is a common causal agent of cattle poisoning by *T.* spp. Cryst. (Me₂CO). Mp 204-206°. [α]_D -140 (CHCl₃).

5-(3S-Dimethylamino-2S-hydroxy-3-phenylpropanoyl), 2,7,13-tri-Ac: **7-Acetyl-taxine A**
[161956-28-1]
C₃₇H₄₉NO₁₁ 683.794
Constit. of *Taxus baccata*. Cryst. Mp 178-180°. [α]_D²⁵ -96 (c, 1 in CHCl₃).

5-(3S-Dimethylamino-2S-hydroxy-3-phenylpropanoyl), 2,7,10,13-tetra-Ac: **7,10-Diacetyl-taxine A**
[244167-03-1]

C₃₉H₅₁NO₁₂ 725.831

Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -75 (c, 0.01 in CHCl₃).


Graf, E. *et al.*, *Annalen*, 1982, 376-381 (*Taxine A*, *cryst struct*)

Poupat, C. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1468-1469 (*2-Deacetyl-taxine A*)

Barboni, L. *et al.*, *Annalen*, 1995, 345-349 (*7-Acetyl-taxine A*, *2-Deacetyl-taxine A*)

Shi, Q.-W. *et al.*, *Tetrahedron*, 1999, **55**, 8365-8376 (*dimethylaminophenylpropanoyl tri-Ac*)

11(15→1)-Abeo-4(20),11-taxadiene-5,9,10,13,15-pentol A-7



C₂₀H₃₂O₅ 352.47

(5α,9α,10β,13α)-form

9-Ac: [153229-37-9]

C₂₂H₃₄O₆ 394.507
Constit. of *Taxus wallichiana*. Needles (MeOAc/hexane). Mp 160-162°. [α]_D²⁰ -24 (c, 1.1 in MeOH). Struct. revised in 1993.

9,13-Di-Ac: [240805-50-9]

C₂₄H₃₆O₇ 436.544
Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -17 (c, 0.01 in CHCl₃).

5,9,10,13-Tetra-Ac: [497879-22-8]

C₂₈H₄₀O₉ 520.619
Constit. of *Taxus wallichiana*. Oil. [α]_D²⁰ +85 (MeOH).

5-Cinnamoyl, 9-Ac: [268728-61-6]

C₃₁H₄₀O₇ 524.653
Constit. of *Taxus yunnanensis*. Gum.

5E-Cinnamoyl, 13-Ac:

C₃₁H₄₀O₇ 524.653
Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -17 (c, 0.01 in CHCl₃). λ_{max} 202 (ε 11700); 277 (ε 8400) (MeOH).

5-Cinnamoyl, 9,13-di-Ac: **Chinentaximine**

[260402-68-4]
[327024-58-8]
C₃₃H₄₂O₈ 566.69
Constit. of *Taxus chinensis*. Amorph. solid. [α]_D²⁸ +43 (c, 0.49 in CHCl₃). [α]_D²⁴ -23 (c, 0.01 in CHCl₃). λ_{max} 222; 277 (MeOH).

5-Cinnamoyl, 9,10,13-tri-Ac: **7-Deacetyl-taxuspine J**

C₃₅H₄₄O₉ 608.727
Constit. of *Taxus cuspidata*. Amorph. solid. [α]_D²⁰ -0.25 (c, 0.1 in MeOH). λ_{max} 202 (ε 11700); 277 (ε 8400) (MeOH).

5E-Cinnamoyl, 10-benzoyl, 9,13-di-Ac:

C₄₀H₄₆O₉ 670.798
Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -11 (c, 0.01 in CHCl₃).

5-(3-Dimethylamino-3-phenylpropanoyl), 13-Ac: [260361-94-2]

C₃₃H₄₇NO₇ 569.737
Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -27 (c, 0.01 in CHCl₃).

5-(3-Dimethylamino-3-phenylpropanoyl), 10-benzoyl, 9,13-di-Ac: [240805-51-0]
C₄₂H₅₃NO₉ 715.882

Constit. of *Taxus yunnanensis*. Gum. [α]_D²⁴ -13 (c, 0.01 in CHCl₃).

Appendino, G. *et al.*, *Chem. Comm.*, 1993, 1587-1589 (*struct*)

Barboni, L. *et al.*, *Phytochemistry*, 1993, **33**, 145-150 (*9-Ac*)

Murakami, R. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1660-1663

Shi, Q.-W. *et al.*, *J. Asian Nat. Prod. Res.*, 1999, **2**, 71-79; *CA*, **132**, 205456u (*deriv*)

Shen, Y.-C. *et al.*, *Phytochemistry*, 1999, **52**, 1565-1569 (*Chinentaximine*)

Shi, Q.-W. *et al.*, *Tetrahedron*, 1999, **55**, 8365-8376 (*isol, pmr, cmr*)

Shi, Q.-W. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 869-872 (*5-cinnamoyl-10-benzoyl-9,13-di-Ac*)

Shi, Q.-W. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 235-238 (*5-cinnamoyl-5-Ac*)


Shi, Q.-W. *et al.*, *J. Asian Nat. Prod. Res.*, 2000, **2**, 311-319; *CA*, **134**, 190713r (*5-cinnamoyl-9,13-di-Ac*)

Kiyota, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 2001, **65**, 35-40 (*5-cinnamoyl-13-Ac*)

Choudhary, M.I. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1488-1490 (*5,9,10,13-tetra-Ac*)

Abereamine 1 A-8

[104386-96-1]



R¹ = R² = OH, R³ = R⁴ = H

C₂₅H₃₄N₂O₆ 458.553

Related to the Corymine/Isocorymine groups but with an additional isopropyl group the origin of which is not yet clear. Alkaloid from the seeds of *Hunteria umbellata* (Apocynaceae). Yellow solid. Sol. H₂O. Mp 199° dec.

Adegoke, E.A. *et al.*, *Phytochemistry*, 1986, **25**, 1461-1468 (*isol, ir, pmr, ms, struct*)

Abereamine 2 A-9

[104406-77-1]

As Abereamine 1, A-8 with

R¹ = R³ = H, R² = R⁴ = OH

C₂₅H₃₄N₂O₆ 458.553

See note under Abereamine 1, A-8.

Alkaloid from the seeds of *Hunteria umbellata* (Apocynaceae). Red solid. Sol. H₂O. Mp 230° dec.

Adegoke, E.A. *et al.*, *Phytochemistry*, 1986, **25**, 1461-1468 (*isol, ir, pmr, ms, struct*)

Abereamine 3 A-10

[104386-97-2]

As Abereamine 1, A-8 with

R¹ = R² = H, R³ = R⁴ = OH

C₂₅H₃₄N₂O₆ 458.553

See note under Abereamine 1, A-8.

Alkaloid from the seeds of *Hunteria umbellata* (Apocynaceae). Brown solid. Sol. H₂O. Mp 280° dec.

Adegoke, E.A. *et al.*, *Phytochemistry*, 1986, **25**, 1461-1468 (*isol, ir, pmr, cmr, ms, struct*)

Abereamine 4 A-11

[104386-98-3]

As Abereamine 1, A-8 with R¹ = R⁴ = H, R² = R³ = OH


C₂₅H₃₄N₂O₆ 458.553

See note under Abereamine 1, A-8. Alkaloid from the seeds of *Hunteria umbellata* (Apocynaceae). Red solid. Sol. H₂O. Mp 260-262°.

Adegoke, E.A. *et al.*, *Phytochemistry*, 1986, **25**, 1461-1468 (*isol, ir, pmr, ms, struct*)

Aberiamide A-12

[283167-82-8]




C₂₈H₅₅N₃O₂ 465.761

Alkaloid from the leaves of *Aberia caffra*. Pale yellow solid. [α]_D²⁵ +20 (c, 1 in MeOH). *Aberia* is not a recognised genus (in the Kew system a synonym for *Dovyalis*).

Sayed, H.M. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 215-219 (*isol, pmr, cmr, ms*)

Aberiamine A-13

N-Hexylhexahydro-1,3-dimethyl-*N*-pentyl-2-pyrimidinedecanamine, 9CI [283167-81-7]




C₂₇H₅₇N₃ 423.767

Alkaloid from the leaves of *Aberia caffra*. Pale yellow solid. [α]_D²⁵ +3 (c, 1 in MeOH). *Aberia* is not a generally recognised genus (in the Kew system a synonym for *Dovyalis*).

Sayed, H.M. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 215-219 (*isol, pmr, cmr, ms*)

Abikoviromycin A-14

7-Ethylidene-1a,2,3,7-tetrahydrocyclopent[*b*]oxireno[*c*]pyridine, 9CI, 8CI. 5-Ethylidene-4,4a-epoxy-2,3,4,4a-tetrahydro-1-pyridine. *Virocidin*. *SF* 973A. *Antibiotic SF* 973A. *Thomsomycin*. *Latumcidin* [31774-33-1]



Absolute Configuration

C₁₀H₁₁NO 161.203

Pyridine numbering shown, for uniformity with other related compds. Prod. by *Streptomyces abikoensis*, *Streptomyces*

rubescens and *Streptomyces latumcidicus*. Polyketide synthase inhibitor. Antifungal and antiviral agent. Highly unstable, polymerising on isol. from broth cultures. Can be handled in dilute soln. [α]_D²¹ +148.9 (c, 1 in 0.1N NaOH). λ_{max} 238 (ε 9980); 338 (ε 11600) (0.1N HCl) (Derep). λ_{max} 245 (ε 8690); 290 (ε 7730) (0.1N NaOH) (Derep). λ_{max} 243 (ε 9000); 289 (sh) (ε 8710); 345 (sh) (ε 4000) (EtOH or pH 7) (Derep). λ_{max} 244 (ε 13900) (MeOH) (Derep).

Sulfate: [43043-59-0]

Cryst. Mp 140-141° dec. [α]_D²⁵ +24 (c, 1 in H₂O).

►GY8431000

Picrate:

Yellow needles (EtOAc/MeOH). Mp 136-137°.

1,7a-Dihydro: **Dihydroabikoviromycin**.

Dihydrolatumcidin. *Antibiotic SF* 973B. *SF* 973B

[20421-29-8]

C₁₀H₁₃NO 163.219

Prod. by *Streptomyces olivaceus*, *Streptomyces viridochromogenes*, *Streptomyces reticuli* and *Streptomyces anulatus*. Mutagenic and genotoxic agent. Mp 60-61° (57-59°). [α]_D²² +276 (c, 1 in MeOH). pK_a 7.7 (50% EtOH aq.). pK_a 8.07. λ_{max} 244 (E1%/1cm 850) (MeOH) (Berdy).

1,7a-Dihydro; hydrochloride: Mp 130-132°.

1,7a-Dihydro, 7-hydroxy: **Dihydro-N-hydroxyabikoviromycin**. *N*-Hydroxydihydroabikoviromycin

[109292-19-5]

C₁₀H₁₃NO₂ 179.218

From *Streptomyces* sp. SANK 65986. Exhibits antimicrobial activity. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{max} 210; 242 (MeOH) (Berdy). λ_{max} 243 (MeOH-HCl) (Berdy). λ_{max} 210; 242 (MeOH-NaOH) (Berdy).

►LD₅₀ (mus, ivn) 100 - 200 mg/kg.

Umezawa, H. *et al.*, *Jpn. Med. J.*, 1951, **4**, 331; *CA*, **46**, 7167c (*isol*)

Sakagami, Y. *et al.*, *J. Antibiot., Ser. A*, 1958, **11**, 6; 231; *CA*, **54**, 2483h; 2484a (*isol, uv, ir, struct*)

Gurevich, A.I. *et al.*, *Tet. Lett.*, 1968, 2209 (*struct, pmr, ir, uv*)

Kono, Y. *et al.*, *J. Antibiot.*, 1970, **23**, 572 (*pmr, cryst struct*)

Kono, Y. *et al.*, *Acta Cryst. B*, 1971, **27**, 2341 (*cryst struct*)

Ogawa, Y. *et al.*, *J. Antibiot.*, 1973, **26**, 186; 609 (*Dihydroabikoviromycin*)

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 2916 (*isol, nmr, ms*)

Takahashi, S. *et al.*, *Sankyo Kenkyusho Nenpo*, 1986, **38**, 105; *CA*, **107**, 55392n (*N*-Hydroxydihydroabikoviromycin)

Takahashi, S. *et al.*, *CA*, 1987, **107**, 55392 (*deriv*)

Maruyama, H. *et al.*, *J. Antibiot.*, 2003, **56**, 801-804 (*isol, pmr, cmr*)

Abrasine A-15

C₁₈H₂₁N₃O₃ 327.382

Struct. unknown. Alkaloid from roots of *Abrus precatorius* (Fabaceae). Mp 218-


220° dec.

Hydrochloride: Mp 226-227°.

Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1966, **3**, 203-207; *CA*, **67**, 90987n (*isol, ir*)

Aburatubolactam A A-16

[170894-24-3]



C₃₀H₄₀N₂O₅ 508.656


Tetramic acid deriv. Closely related to Alteramide A, A-661. Prod. by *Streptomyces* sp. SCRC-A20 from a marine mollusc. Cytotoxic agent, apoptosis inducer. Inhibitor of superoxide anion generation.

Japan. Pat., 1995, 95 228 583; *CA*, **123**, 337551v (*isol*)

Bae, M.-A. *et al.*, *Heterocycl. Commun.*, 1996, **2**, 315-318 (*isol, pmr, struct*)

Aburatubolactam B A-17

[170894-25-4]



C₃₀H₄₀N₂O₆ 524.656

Prod. by *Streptomyces* sp. SCRC A-20. Antiinflammatory and cytotoxic agent. *Cryst.* [α]_D +197 (Py).

25-Deoxy: **Aburatubolactam C**

[170894-26-5]

C₃₀H₄₀N₂O₅ 508.656

Prod. by *Streptomyces* sp. SCRC A-20. Apoptosis-inducing substance. Antiinflammatory and cytotoxic agent. Powder. [α]_D +136 (Py).


Japan. Pat., 1995, 95 228 583; *CA*, **123**, 337551v

Bae, M.-A. *et al.*, *J. Microb. Biotechnol.*, 1998, **8**, 455-460

Abyssenine B A-18

Abyssinine B

[54519-15-2]



C₂₅H₃₈N₄O₄ 458.6

The alkaloids included in the entry need not necessarily have the same stereochem. Alkaloid from the bark of *Zizyphus abyssinica* and *Zizyphus mucronata* and from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Needles (MeOH/petrol), Mp 229-230°. [α]_D²⁰ +151 (c, 0.16 in CHCl₃).

N-De-Me: Abyssinine C. Abyssinine C
[55857-03-9]

C₂₄H₃₆N₄O₄ 444.573

Alkaloid from the bark of *Zizyphus abyssinica* and from bark and leaves of *Zizyphus mucronata* (Rhamnaceae). Shows antibacterial and antifungal activity. Amorph. [α]_D²⁰ +144 (c, 0.12 in CHCl₃). [α]_D²⁰ -15 (c, 0.13 in MeOH).

17-Methoxy, N-de-Me: Mucronine G

[55856-94-5]

C₂₅H₃₈N₄O₅ 474.599

Alkaloid from the bark and leaves of *Zizyphus mucronata* (Rhamnaceae). Amorph. [α]_D²⁰ -50 (c, 0.084 in MeOH).

Z-Isomer, 1'-hydroxy: Zizyphine D

[55323-68-7]

C₂₅H₃₈N₄O₅ 474.599

Alkaloid from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Needles (MeOH aq.). Mp 195°. [α]_D²⁰ +236 (c, 0.10 in CHCl₃). [α]_D²⁰ -121 (c, 0.10 in MeOH).

Z-Isomer, 1'-hydroxy, N-de-Me: Zizyphine E

[55323-71-2]

C₂₄H₃₆N₄O₅ 460.572

Alkaloid from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Amorph. [α]_D²⁰ +150 (c, 0.10 in CHCl₃). [α]_D²⁰ -111 (c, 0.10 in MeOH).

Tschesche, R. *et al.*, *Annalen*, 1974, 1915-1928 (*Abyssinines B,C, Mucronine G*)

Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 2328 (*occur*)

Cassels, B.K. *et al.*, *Tetrahedron*, 1974, **30**, 2461-2466 (*Abyssinine B, Zizyphines D,E*)


Hindenlang, D.M. *et al.*, *Annalen*, 1980, 447 (*Zizyphine D, emr*)

Wang, J. *et al.*, *Tet. Lett.*, 2007, **48**, 6717-6721 (*synth, struct*)

Abysomicin B

A-19

[725254-08-0]



Absolute Configuration

C₁₉H₂₃NO₇ 377.393

Prod. by *Verrucosporina* sp. AB 18-032. Powder.


Bister, B. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 2574-2576 (*isol, struct*)

Riedlinger, J. *et al.*, *J. Antibiot.*, 2004, **57**, 271-279 (*isol, activity*)

Acacipetalin

A-20

2-(β -D-Glucopyranosyloxy)-3-methyl-2-butenenitrile, 9CI
[644-68-8]



C₁₁H₁₇NO₆ 259.258

Some confusion occurred in the literature around 1975. Isol. from *Acacia sieberiana* and *Acacia hebeclada*. Mp 177-179°. [α]_D²⁶ -36.6 (H₂O).

Tetra-Ac.: [66890-86-6]

Mp 101-103°.

2,3-Dihydro: **Dihydroacacipetalin**

C₁₁H₁₉NO₆ 261.274

Isol. from *Acacia sieberiana*.

Rimington, C. *et al.*, *S. Afr. J. Sci.*, 1935, **32**, 154 (*isol*)


Ettlinger, M.G. *et al.*, *Chem. Comm.*, 1977, 952 (*struct*)

Jaroszewski, J.W. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 555 (*pmr, cmr*)

Acalyphamide

A-21

N-[2-(4-Hydroxyphenyl)ethyl]dotriacontanamide
[81119-33-7]



C₄₀H₇₃NO₂ 600.022

Isol. from *Acalypha indica*. Mp 135°.

Ac.:


Amorph. solid (CHCl₃/MeOH). Mp 126°.

Talapatra, B. *et al.*, *Indian J. Chem., Sect. B*, 1981, **20**, 974

Acalyphin

A-22

3-(β -D-Glucopyranosyloxy)-1,2,3,6-tetrahydro-2-hydroxy-4-methoxy-1-methyl-6-oxo-3-pyridinecarbonitrile, 9CI
[81861-72-5]



C₁₄H₂₀N₂O₉ 360.32

Alkaloid from the aerial parts of *Acalypha indica* (Euphorbiaceae). Also reported from Te Xian Cai (*Acalypha australis*) and *Acalypha farnesiana*.

Hygroscopic powder. Mp 185-186° (182-184°). λ _{max} 223 (ε 6310); 255 (sh) (ε 2450) (EtOH) (Derep).


Nährstedt, A. *et al.*, *Phytochemistry*, 1982, **21**, 101 (*uv, ir, pmr, cmr, struct*)

Li, T.S.C. *et al.*, *Chinese and Related North American Herbs*, CRC Press, 2002, 4 (*occur*)

Acanthamolide

A-23

[64852-96-6]



C₁₉H₂₅NO₅ 347.41


Constit. of *Acanthospermum glabratum*. Cryst. (C₆H₆/MeOH). Mp 249-251°. λ _{max} 225 (ε 10000) (MeOH) (Derep).

Saleh, A.A. *et al.*, *J.C.S. Perkin 1*, 1980, 1090

Acanthonitrile 3

A-24

[112766-99-1]



C₁₆H₂₅N 231.38

Metab. of marine sponge *Acanthella acuta*. Oil. [α]_D -31.5 (c, 1.2 in CHCl₃).

Isothiocyanate: Acanthiothiocyanate 3

[112767-00-7]

C₁₆H₂₅NS 263.446


Metab. of *Acanthella acuta*. Oil. [α]_D -12.9 (c, 0.3 in CHCl₃). Has -NCS replacing -NC.

Mayol, L. *et al.*, *Tetrahedron*, 1987, **43**, 5381-5388

Acanthoine

A-25

N,N''-[1-(3,4-Dimethoxystyryl)-2-butenylene]diformamidine, 8CI
[6793-31-3]



C₁₆H₂₂N₄O₂ 302.375

Alkaloid from *Carduus acanthoides* (Asteraceae). Mp 221° (192-193°) (as dihydrochloride). [α]_D²⁹ +7.1 (c, 0.4 in H₂O).

Tetrahydro: Acanthoidine. Ruscopine

[94347-15-6]

[4192-85-2]

C₁₆H₂₆N₄O₂ 306.407


Alkaloid from *Carduus acanthoides* (Asteraceae). Mp 250-251° (as hydrochloride). [α]_D²² +6.8 (c, 0.4 in H₂O). Saturated at both olefinic double bonds. The duplication of

trivial names is unexpected since both clearly refer to the same isol.

Frydman, B. *et al.*, *Tetrahedron*, 1962, **18**, 1063-1072 (*Acanthoine*, *Acanthoidine*)
U.S. Pat., 1964, 3 130 231; *CA*, **62**, 9188 (*synth*)
U.S. Pat., 1964, 3 130 230; *CA*, **61**, 3159 (*Ruscopine*)

3,5-Acarnidine A-26

*C*_{12:0}-Acarnidine
 [67534-25-2]



*C*₂₆H₅₁N₅O₂ 465.721

Isol. from the sponge *Acarnus erithacus* as part of an inseparable mixt. Possesses broad antimicrobial activity and modest antiviral activity.

5,6*Z*-Didehydro: *C*_{12:1}-Acarnidine

[67534-26-3]
*C*₂₆H₄₉N₅O₂ 463.705

Component of *Acarnus erithacus* acarnidine mixt. λ_{\max} 218 (ϵ 17000) (no solvent reported) (Derep).

N-Dedodecanoyl, *N*-(5*Z*,8*Z*,11*Z*-tetradecatrienoyl): *C*_{14:3}-Acarnidine

[67534-27-4]
*C*₂₈H₄₉N₅O₂ 487.727


Isol. from *Acarnus erithacus*.

Carter, G.T. *et al.*, *J.A.C.S.*, 1978, **100**, 4302-4304 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Yorke, S.C. *et al.*, *Aust. J. Chem.*, 1986, **39**, 447-455 (*synth*)

Accedine A-27

1-Methylsarpagan-3,17-diol, 9*CI*
 [58262-63-8]



Absolute Configuration

*C*₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Tabernaemontana acedens* (Apocynaceae). Needles (MeOH). Mp 148-149°. $[\alpha]_{\text{D}}^{20}$ +72 (c, 0.096 in CHCl₃).

N-*De*-*Me*: Amerovolfine. *N*-Demethylaccedine

[121312-84-3]
*C*₁₉H₂₂N₂O₂ 310.395

Alkaloid from the stem bark of *Rauwolfia tetraphylla* and *Rauwolfia cubana* (Apocynaceae). Needles (MeOH). Mp 183-184°. $[\alpha]_{\text{D}}^{20}$ +67 (c, 0.95 in MeOH).

N-*De*-*Me*: hydrochloride: Mp 280-282°.

16-*Epimer*, *N*-*de*-*Me*: *N*-Demethyl-16-epiaccedine. Sarpagan-3,7-diol, 9*CI*
 [59190-62-4]

*C*₁₉H₂₂N₂O₂ 310.395

Alkaloid from *Tabernaemontana acedens* (Apocynaceae). Mp 170-172°. $[\alpha]_{\text{D}}^{20}$ +50 (c, 0.06 in CHCl₃).

10-Hydroxy, *N*-*de*-*Me*: 3-Hydroxysarpagine

*C*₁₉H₂₂N₂O₃ 326.394

Alkaloid from the roots of *Rauwolfia serpentina*. Powder. $[\alpha]_{\text{D}}^{21}$ +44 (c, 1 in MeOH). λ_{\max} 205 (log ϵ 4.36); 224 (log ϵ 4.23); 278 (log ϵ 3.75) (MeOH).

Achenbach, H. *et al.*, *Chem. Ber.*, 1975, **108**, 3842-3854 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Achenbach, H. *et al.*, *Tet. Lett.*, 1976, 351-352 (*N*-Demethyl-16-epiaccedine)


Martinez, J.A. *et al.*, *Phytochemistry*, 1989, **28**, 961-962 (*Amerovolfine*)

Martinez, J.A. *et al.*, *Planta Med.*, 1989, **55**, 283-285 (*Amerovolfine*)

Itoh, A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 848-852 (3-Hydroxysarpagine)

Accedinisine A-28

Methyl 3-(17-hydroxy-1-methylsarpagan-10-yl)vobasan-17-oate, 9*CI*
 [61551-76-6]



*C*₄₁H₄₈N₄O₃ 644.855

Alkaloid from *Tabernaemontana acedens* (Apocynaceae). Needles (Me₂CO). Mp 235° dec. $[\alpha]_{\text{D}}^{20}$ -60 (c, 0.4 in CHCl₃).

*N*⁴-*De*-*Me*: Demethylaccedinisine

[160427-83-8]
*C*₄₀H₄₆N₄O₃ 630.828

Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae). Also a constit. of *Peschiera van heurkii*. $[\alpha]_{\text{D}}^{20}$ -30 (c, 0.3 in MeOH).

3'-Hydroxy: Accedinine

[61551-77-7]
*C*₄₁H₄₈N₄O₄ 660.855


Alkaloid from *Tabernaemontana acedens* (Apocynaceae). Oil. $[\alpha]_{\text{D}}^{20}$ -81 (c, 0.2 in CHCl₃).

Achenbach, H. *et al.*, *Chem. Ber.*, 1976, **109**, 3527 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)

Azoug, M. *et al.*, *Phytochemistry*, 1995, **39**, 1223 (*Demethylaccedinisine*)

Aceropterin A-29

[176227-10-4]



Relative configuration

*C*₂₃H₂₉NO₇ 431.485

Alkaloid from the Caribbean sea plume *Pseudopterogorgia acerosa*. $[\alpha]_{\text{D}}$ +0.44 (c, 2.0 in CHCl₃). Closely related to Tobagolide and Pseudopterolide. λ_{\max} 258

(ϵ 3750) (MeOH) (Berdy).

Rodríguez, A.D. *et al.*, *Tet. Lett.*, 1996, **37**, 2687 (*isol*, *pmr*, *cmr*, *ms*, *struct*)

Acetamide, 9*CI* A-30

Ethanamide. Methanecarboxamide

[60-35-5]

H₃CCONH₂

*C*₂H₅NO 59.068

Isol. from some plants e.g. *Chimonanthus fragrans*. Produced industrially by distillation of ammonium acetate or by hydration of Acetonitrile. Solubiliser, plasticiser, stabiliser, used industrially as solv. in molten form. Dissolves virtually all classes of organic and inorganic compds. Deliquescent, hexagonal cryst. Odourless when pure but usually has characteristic "mouse" odour. V. sol. H₂O, EtOH; sol. CHCl₃; prac. insol. Et₂O. Mp 82-83°. Bp 222° Bp₅ 92°. p*K*_{a1} -1.4 (25°). Triboluminescent.

▶ Fl. p. >104°. Possible human carcinogen (IARC 2B). Irritant. Exp. carcinogen (v. large dose). Exp. reprod. and teratogenic effects (large doses). AB4025000

N-(13-Methyltetradecyl): Capsiamide

[64317-66-4]

*C*₁₇H₃₅NO 269.47

Constit. of fruit of hot pepper varieties of *Capsicum annuum*.

Takahashi, M. *et al.*, *Yakugaku Zasshi*, 1977, **97**, 758 (*Capsiamide*)

Speakman, J.C. *et al.*, *J. Chem. Res., Synop.*, 1979, 277 (*cryst struct*)

ul Hassan, M. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 447 (*cmr*)


Gonzalez, G. *et al.*, *J.C.S. Faraday 2*, 1981, 2231 (*pmr*)

Kerridge, D.H. *et al.*, *Chem. Soc. Rev.*, 1988, **17**, 181 (*rev*)

Liu, L.K. *et al.*, *Acta Cryst. C*, 1994, **50**, 1333 (*cryst struct*, *hydrochloride*)

2-Acetamidobenzoic acid A-31

2-(Acetylamino)benzoic acid, 9*CI*. *N*-Acetylanthranilic acid. Lappaconitic acid
 [89-52-1]



*C*₉H₉NO₃ 179.175

Needles (AcOH). Mp 185°. p*K*_a 5.64.

▶ LD₅₀ (mus, orl) 1114 mg/kg. CB2455000

Me ester: [2719-08-6]

*C*₁₀H₁₁NO₃ 193.202

Needles (EtOH). Mp 101°.

Et ester: [20628-20-0]

*C*₁₁H₁₃NO₃ 207.229

Needles (EtOH). Mp 64-65°.

Amide: 2-Acetamidobenzamide. 2-(Acetylamino)benzamide. Antibiotic NP 101A. NP 101A
 [33809-77-7]

*C*₉H₁₀N₂O₂ 178.19

Prod. by *Streptomyces aurantiogriseus* and *Cytophaga marinoflava* sp. AM13.1. Antifungal agent. Needles

(EtOH). Sol. MeOH, EtOH, EtOAc, C₆H₆, DMSO, Py; poorly sol. H₂O. Mp 179-180° (α -form) Mp 189-190° (β -form). Exists in two cryst. forms, accounting for unreliable melting point determinations. λ_{\max} 219 (log ϵ 3.56); 252 (log ϵ 3.31); 299 (log ϵ 2.72) (MeOH). λ_{\max} 219 (ϵ 3630); 252 (ϵ 2042); 299 (ϵ 550) (MeOH) (Berdy).

► CU8702020

N-Ac: Diacetylanthranilic acid

C₁₁H₁₁NO₄ 221.212

Prisms (EtOH). Mp 180°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 356D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 1362B (nmr)

Matsuo, M. et al., Chem. Pharm. Bull., 1972, 20, 990 (nmr)

Erikson, J. et al., J. Chem. Educ., 1972, 49, 688 (synth)

Matsuda, V. et al., Bull. Chem. Soc. Jpn., 1973, 46, 430 (synth)

Ger. Pat., 1976, 2 556 590; CA, 85, 160115 (synth)

Errede, L.A. et al., J.C.S. Perkin 2, 1981, 233-238 (Acetamidobenzamide)


Phay, N. et al., J. Antibiot., 1996, 49, 703-705 (Acetamidobenzamide)

Shaaban, M. et al., Dissertation, Univ. of Göttingen, 2004, (Cytophaga marinoflava isol)

Kelleher, J.M. et al., ARKIVOC, 2007, xvi, 209-226 (Acetamidobenzamide, powder struct, polymorphism)

1-(2-Acetamidophenyl)-2-hydroxy-1-propanone A-32

N-[2-(2-Hydroxy-1-oxopropyl)phenyl]acetamide, 9CI. 2-(2-Hydroxypropanoyl)acetanilide [150641-66-0]



C₁₁H₁₃NO₃ 207.229

Prod. by Nocardia sp. DSM 43130. [α]_D -8 (CHCl₃). λ_{\max} 217 ; 253 ; 339 (MeOH). λ_{\max} 217 ; 253 ; 339 (MeOH) (Berdy).

Abraham, W.R. et al., Phytochemistry, 1993, 33, 929-930

2-[(2-Acetamidopropanoyl)amino]benzamide A-33

N²-[2-(Acetylamino)propanoyl]anthranilamide. N²-Acetylalanyl anthranilamide C₁₂H₁₅N₃O₃ 249.269

(S)-form


Antibiotic NI 15501A. NI 15501A [208345-44-2]

Prod. by the marine fungus Penicillium sp. NI15501. Solid. Sol. MeOH, butanol; poorly sol. H₂O. [α]_D²⁰ -62.8 (c, 0.15 in MeOH). λ_{\max} 216 (ϵ 12000); 253 (ϵ 6100); 290 (ϵ 1800) (MeOH).

Onuki, H. et al., J. Antibiot., 1998, 51, 442-444 (isol, uv, ir, pmr, cmr, ms)

8-Acetylndihydrochelerythrine A-34

12,13-Dihydro-1,2-dimethoxy-12-methyl-13-(2-oxopropyl)-1,3-benzodioxolo[5,6-c]phenanthridine, 9CI. 13-Acetylndihydrochelerythrine, 8CI. 11-Acetylndihydrochelerythrine. Alkaloid ZT₁, 6-Acetylndihydrochelerythrine [22864-92-2]



C₂₄H₂₃NO₅ 405.449

Probably an artifact. Numbering systems vary. Alkaloid from the root bark of Toddalia aculeata and Xylocarpus granatum, and from the bark of Zanthoxylum conspersipunctatum and Zanthoxylum tshanimposa (Rutaceae, Meliaceae). Also obt. by the base-catalysed addn. of acetone to Chelerythrine. Plates (Me₂CO). Mp 199° (190-192°, 194-195°). λ_{\max} 228 (log ϵ 4.52); 283 (log ϵ 4.61); 320 (log ϵ 4.18) (solvent not reported).

O^o-De-Me: O-Desmethyldihydrochelerythrinyl-8-acetone. O-Desmethyldihydrochelerythrinyl-11-acetone. Alkaloid ZT₃ [51888-34-7]

C₂₃H₂₁NO₅ 391.423

Isol. from the bark of Zanthoxylum tshanimposa (Rutaceae). Cryst. (MeOH). Mp 201-205°. Opt. inactive, prob. an artifact. λ_{\max} 229 (log ϵ 4.55); 284 (log ϵ 4.62); 320 (log ϵ 4.17) (EtOH). λ_{\max} 234 (log ϵ 4.48); 297 (log ϵ 4.52); 338 (log ϵ 4.33) (EtOH + NaOH).

Desai, P.D. et al., Indian J. Chem., 1967, 5, 41-42 (uv, ir, pmr, ms, struct)

MacLean, D.B. et al., Can. J. Chem., 1969, 47, 1951-1956 (synth, ir, pmr, ms)

Corrie, J.E.T. et al., Aust. J. Chem., 1970, 23, 133-145 (synth, uv, ir, ms)

Krajniak, E.R. et al., Aust. J. Chem., 1973, 26, 687-689 (isol)


Decadain, N. et al., Phytochemistry, 1974, 13, 505-511 (O^o-de-Me)

Chou, F.Y. et al., Heterocycles, 1977, 7, 969-975 (occur)

Sharma, P.N. et al., Phytochemistry, 1981, 20, 2781-2783 (isol)

8-Acetylndihydronitidine A-35

[80330-39-8]



Relative Configuration

C₂₄H₂₃NO₅ 405.449

Alkaloid from the stem bark of Zanthoxylum tetraspermum. Antibacterial agent. Cryst. Mp 165-167°. [α]_D 0 (CHCl₃). λ_{\max} 316 (log ϵ 3.87) (EtOH).

10,11-Di-O-de-Me, 10,11-methylene ether: 8-Acetylndihydroavicine [348098-59-9]

C₂₃H₁₉NO₅ 389.407

Alkaloid from the stem bark of Zanthoxylum tetraspermum. Antibacterial agent. Cryst. Mp 184-185°. [α]_D²⁵ -6.6 (CHCl₃). λ_{\max} 321 (log ϵ 4.2) (EtOH).


Nissanka, A.P.K. et al., Phytochemistry, 2001, 56, 857-861 (isol, pmr, cmr, 8-Acetylndihydroavicine)

Morel, A.F. et al., Acta Cryst. C, 2002, 58, 606-607 (Acetylndihydroavicine, crystal struct)

Wang, X.-L. et al., Acta Cryst. E, 2006, 62, 2247-2248 (crystal struct)

8-Acetylndihydrosanguinarine A-36

6-Acetylndihydrosanguinarine. 6-Acetylndihydrosanguinarine [37687-34-6]



C₂₃H₁₉NO₅ 389.407

(S)-form

Alkaloid from the roots of Corydalis flabellata. Needles (CHCl₃/EtOH). Mp 209-210°. [α]_D²⁰ +25.3 (c, 0.02 in CHCl₃). λ_{\max} 232 (log ϵ 4.4); 282 (log ϵ 4.6); 323 (log ϵ 4.8) (CHCl₃).

(E)-form

Alkaloid from Argemone mexicana, the callus tissue of Papaver somniferum (opium poppy) and the roots of Glaucium flavum var. vestitum (Papaveraceae). Also obt. by the base-catalysed condensation of sanguinarine with acetone. Pale yellow needles (MeOH/CHCl₃). Mp 194-195.5°.

Furuya, T. et al., Phytochemistry, 1972, 11, 3041-3044 (isol, uv, ir, pmr, ms, struct, synth)


Döpke, W. et al., Z. Chem., 1976, 16, 54-55 (occur, uv, ir, pmr, ms, struct)

Castedo, L. et al., Heterocycles, 1981, 16, 533-536 (occur)

Koul, S. et al., Planta Med., 2002, 68, 262-265 (isol, pmr, cmr, ms)


Acetonylevodiamine A-37

[848601-89-8]



C₂₂H₂₁N₃O₂ 359.427Alkaloid from the fruit of *Evodia rutae-carpa*. Mp 163-164°.Zuo, G.Y. *et al.*, *Yunnan Zhiwu Yanjiu*, 2003, **25**, 103-106; *CA*, **142**, 332872h (*isol*)**Acetopine**


A-38

N²-(Carboxymethyl)arginine, 9CI. DemethyloctopineC₈H₁₆N₄O₄ 232.239**(S)-form***L*-form


[18416-86-9]

Isol. from cotton (*Gossypium hirsutum*) and soybean (*Glycine max*) callus. Cryst. (EtOH aq.). Mp 281-282°. [α]_D²⁴ +24 (c, 2.5 in H₂O).Herbst, R.M. *et al.*, *J.O.C.*, 1946, **11**, 368(*synth*)Christou, P. *et al.*, *Plant Physiol.*, 1986, **82**, 218 (*isol*)**1-(Acetoxymethyl)-2-formyl-1,2-dihydro-5-hydroxy-7-methoxyisoquinoline**


A-39

C₁₄H₁₅NO₅ 277.276**(+)-form**Alkaloid from the sponge *Petrosia similis*. Amorph. solid. [α]_D²⁵ +208 (c, 0.72 in CHCl₃). λ_{max} 224 (ε 6660); 228 (ε 6680); 231 (ε 6650); 299 (ε 4790) (MeOH).Ramesh, P. *et al.*, *J. Nat. Prod.*, 1999, **62**, 780-781 (*isol, uv, pmr, cmr, ms*)**6-Acetyl-2-amino-7,8-dihydro-4(3H)-pteridinone, 9CI**


A-40

Sepiapterin C
[42310-08-7]C₈H₉N₅O₂ 207.191Pigment from *Drosophila melanogaster*. Yellow cryst. (H₂O).Sugiura, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 3312 (*isol, struct*)**2-Acetyl-3-amino-5-hydroxy-1,4-naphthoquinone**


A-41

2-Acetyl-3-amino-5-hydroxy-1,4-naphthalenedione, 9CIC₁₂H₉NO₄ 231.207Constit. of the stem bark of *Goniothalamus marcanii*. Yellow powder. λ_{max} 260 (log ε 4.31); 296 (log ε 4.08) (MeOH).Soonthornchareonnon, N. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1390-1394**4-Acetyl-2(3H)-benzoxazolone, 9CI**


A-42

4-Acetylbenzoxazolone-2-one
[70735-79-4]C₉H₇NO₃ 177.159Found in kernels of *Zea mays* (sweet corn) (Poaceae). Needles (Me₂CO aq.). Mp 217-218°. λ_{max} 250 (ε 9777); 320 (ε 5128) (MeOH) (Berdy). λ_{max} 268 (ε 8912); 351 (ε 7413) (MeOH/NaOH) (Berdy).Fielder, D.A. *et al.*, *Tet. Lett.*, 1994, **35**, 521-524 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)Fielder, D.A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 456-458 (*synth*)Escobar, C.A. *et al.*, *J. Het. Chem.*, 1997, **34**, 1407-1414 (*synth*)Kluge, M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 821-822 (*synth*)**3-Acetyl-5-benzyl-2,4-pyrrolidinedione**

A-43

C₁₃H₁₃NO₃ 231.251Enolised β-tricarboxyl compd. with several tautomers possible. Isol. from *Leocarpus fragilis*. No phys. props. reported. Steglich, W. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 281-288**1-Acetyl-β-carboline**


A-44

1-(9H-Pyrido[3,4-b]indol-1-yl)ethanone, 9CI
[50892-83-6]C₁₃H₁₀N₂O 210.235The compd. lycii Alkaloid I, isol. from several plant spp. and erroneously assigned the struct. N⁹-Formylharman, has now been shown by Bracher *et al* to be identical with 1-Acetyl-β-carboline. Alkaloid from the bark of *Ailanthus malabarica* (Simaroubaceae). Also isol. from the sponge *Tedania ignis*. Light yellow solid. Mp 203-205°.N²-Oxide:C₁₃H₁₀N₂O₂ 226.234

Mp 180°.

Joshi, B.S. *et al.*, *Heterocycles*, 1977, **7**, 193 (*isol, uv, ir, pmr, ms, struct*)Dillman, R.L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1056-1061; 1141 (*isol, sponge, ir, pmr, cmr, ms*)Bracher, F. *et al.*, *Annalen*, 1993, 837; 1335 (*synth, pmr, cmr, lycii Alkaloid I*)Bracher, F. *et al.*, *Synth. Commun.*, 1995, **25**, 1557 (*synth*)Zhou, T.-S. *et al.*, *Phytochemistry*, 1998, **49**, 1807-1809 (*isol, uv, ir, pmr, cmr*)Kast, O. *et al.*, *Synth. Commun.*, 2003, **33**, 3843-3850 (N²-oxide)**1-Acetyl-β-carboline-3-carboxylic acid**

A-45

1-Acetyl-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI. 1-Acetyl-3-carboxy-β-carboline
[73818-29-8]C₁₄H₁₀N₂O₃ 254.245Alkaloid from aerial parts of *Vestia lycioides* (Solanaeae). Mp 347-350°.*Me ester: 1-Acetyl-3-methoxycarbonyl-β-carboline*

[66154-37-8]

C₁₅H₁₂N₂O₃ 268.271Alkaloid from *Vestia lycioides* (Solanaeae). Yellow needles (CHCl₃/MeOH). Mp 234-236°.*Amide: 1-Acetyl-9H-pyrido[3,4-b]indole-3-carboxamide, 9CI. Stellarine A†*
[157341-07-6]C₁₄H₁₁N₃O₂ 253.26Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*.

[2Z-(Methoxycarbonyl)vinyl]amide:

Stellarine B†

[180681-40-7]

C₁₈H₁₅N₃O₄ 337.334Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*.[2-(Methoxycarbonyl)ethyl]amide: *Dichotomide I*

[755036-54-5]


C₁₈H₁₇N₃O₄ 339.35Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. λ_{max} 220 (log ε 3.43); 286 (log ε 3.49) (MeOH).*1'S-Alcohol: 1-(1-Hydroxyethyl)-β-carboline-3-carboxylic acid. Dichotomine A*

[755036-38-5]
 $C_{14}H_{12}N_2O_3$ 256.26
 Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. $[\alpha]_D^{27}$ -9.7 (c, 0.85 in MeOH). λ_{max} 236 (log ϵ 4.14); 269 (log ϵ 4.24) (MeOH).

Faini, F. et al., *Phytochemistry*, 1978, **17**, 338 (isol, uv, ir, pmr, ms, struct, synth, deriv)
 Razmilić, I. et al., *J. Het. Chem.*, 1980, **17**, 595 (synth, uv, ir, pmr, ms)
 Faini, F. et al., *Planta Med.*, 1980, **38**, 128 (isol)
 Cui, Z.-H. et al., *Nat. Prod. Lett.*, 1995, **7**, 59-64 (Stellarines)
 Sun, B. et al., *J. Nat. Prod.*, 2004, **67**, 1464-1469 (Dichotomide I, Dichotomine A)
 Omura, K. et al., *Chem. Pharm. Bull.*, 2008, **56**, 237-238 (Dichotomine A)

5-Acetyl-2-chlorotetrahydro-4-methyl-2H-1,3-oxazine

1-(2-Chlorotetrahydro-4-methyl-2H-1,3-oxazin-5-yl)ethanone



(2R*,4S*,5R*)-form

$C_7H_{12}ClNO_2$ 177.63

(2R*,4S*,5R*)-form [1010693-65-8]
 Prod. by *Geotrichum* sp. AL4.
 Amorph. solid. $[\alpha]_D^{25}$ +18 (c, 0.09 in $CHCl_3$).

(2R*,4S*,5S*)-form [1010693-64-7]
 Prod. by *Geotrichum* sp. AL4.
 Amorph. solid. $[\alpha]_D^{25}$ +25 (c, 0.1 in $CHCl_3$).

Li, G.-H. et al., *Chem. Biodiversity*, 2007, **4**, 1520-1524 (isol, pmr, cmr)

Acetylcholine(1+) A-47

2-Acetyloxy-N,N,N-trimethylethanaminium, 9CI. Choline acetate

[51-84-3]
 $Me_3N^{\oplus}CH_2CH_2OAc$

$C_7H_{16}NO_2^{\oplus}$ 146.209

Occurs in blood, spleen, ergot and plants and in brain tissue in complexed form e.g. in ergot and *Capsella bursa-pastoris* (shepherd's purse) (Brassicaceae). Neurotransmitter, cholinergic and miotic agent, cardiac depressant, peripheral vasodilator. Easily hyd. by alkalis, and *in vivo* by cholinesterases.

► FZ9700000

Chloride: *Acetylcholine chloride*, BAN, INN, USAN. Miochol. Ovisot [60-31-1]

$C_7H_{16}ClNO_2$ 181.662

V. deliquescent cryst. powder. Sol. H_2O . Mp 149-152°.

► LD₅₀ (rat, orl) 2500 mg/kg. LD₅₀ (rat, ivn) 22 mg/kg. FZ9800000

Bromide: [66-23-9]

$C_7H_{16}BrNO_2$ 226.113

Deliquescent prisms (EtOH). Mp 143°.

► FZ9680000

Iodide: [2260-50-6]

$C_7H_{16}INO_2$ 273.113

Mp 161°.

► KH3300000

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 678C; 678D; 679A (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **1**, 1078A (nmr)

Dudley, H.W. et al., *Biochem. J.*, 1929, **23**, 1064 (synth, bibl)

Marquardt, P. et al., *Arzneim.-Forsch.*, 1956, **6**, 168; 309 (biosynth)

Culvenor, C.C.J. et al., *Chem. Comm.*, 1966, 537 (pmr)

Beveridge, D.L. et al., *J.A.C.S.*, 1971, **93**, 3759 (struct)

Micelson, M.J. et al., *Acetylcholine; Approach of Mol. Mech. of Action*, Pergamon, 1974,

Hanin, I. et al., *Choline Acetylcholine: Handb. Chem. Assay Methods*, Raven Press, NY, 1974, (anal)

Svinning, T. et al., *Acta Cryst. B*, 1975, **31**, 1581 (cryst struct)

Sax, M. et al., *Acta Cryst. B*, 1976, **32**, 1953 (conformn)

Takasuka, M. et al., *J.C.S. Perkin 2*, 1982, 585 (ir)


Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1112

Al-Badr, A.A. et al., *Anal. Profiles Drug Subst.*, 2004, **31**, 3-115 (chloride, rev)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ABO000; CMF250

3-Acetyl-2,5-diamino-1,4-benzoquinone A-48

[16791-08-5]



$C_8H_8N_2O_3$ 180.163

Constit. of *Cynanchum wilfordii*.

Amorph. red powder. Mp > 300°. λ_{max} 304 (log ϵ 4.21); 325 (log ϵ 4.34); 477 (log ϵ 2.87) (DMSO).

N^2, N^5 -Di-Me: [16791-09-6]

$C_{10}H_{12}N_2O_3$ 208.216

Mp 197°.

N^5, N^5 -Di-Me: [16789-18-7]

$C_{10}H_{12}N_2O_3$ 208.216

Mp 164°.

N^2, N^2, N^5, N^5 -Tetra-Me: [16735-85-6]

$C_{12}H_{16}N_2O_3$ 236.27


Mp 180°.

Schaefer, W. et al., *Tet. Lett.*, 1967, 4307 (synth)

Yeo, H. et al., *Phytochemistry*, 1997, **46**, 1103-1105 (isol, uv, ir, pmr, cmr, ms)

5-Acetyl-3,4-dihydro-2H-pyrrole A-49

1-(3,4-Dihydro-2H-pyrrol-5-yl)ethanone, 9CI. 2-Acetyl-1-pyrroline [85213-22-5]



C_6H_9NO 111.143

Isol. from the flowers of *Vallis glabra*. Key aroma component of rice. Liq. with popcorn odour. Bp₁₅ 26-28°. Unstable. Odour threshold of 0.02ng/L in air.

Buttery, R.G. et al., *Chem. Ind. (London)*, 1982, 958; 1983, 478 (isol, synth)

Favino, T.F. et al., *J.O.C.*, 1996, **61**, 8975-8979 (synth, pmr, bibl)

Hofman, T. et al., *J. Agric. Food Chem.*,

1998, **46**, 616-619; 2270-2277 (synth, ms, bibl)

Wongpornchai, S. et al., *J. Agric. Food Chem.*, 2003, **51**, 457-462 (isol)


Harrison, T.J. et al., *J.O.C.*, 2005, **70**, 10872-10874 (synth)

Adams, A. et al., *Chem. Rev.*, 2006, **106**, 2299-2319 (rev)

Snowdon, E.M. et al., *J. Agric. Food Chem.*, 2006, **54**, 6465-6474 (rev)

Fuganti, C. et al., *Tetrahedron*, 2007, **63**, 4762-4767 (synth)

2-Acetyl-4,6-dimethoxy-furo[2,3-b]quinoline 2-Acetylpteleine A-50



$C_{15}H_{13}NO_4$ 271.272

Alkaloid from the root bark of *Melicope semecarpifolia*. Yellow needles (MeOH). Mp 123-126°. λ_{max} 236 (log ϵ 4.01); 303 (log ϵ 3.7); 330 (log ϵ 3.76) (MeOH).

6-Demethoxy, 7-methoxy: 2-Acetyl-4,7-dimethoxyfuro[2,3-b]quinoline. 2-Acetyllevolitrine


$C_{15}H_{13}NO_4$ 271.272

Alkaloid from the root bark of *Melicope semecarpifolia*. Yellow needles (MeOH). Mp 121-124°. λ_{max} 235 (log ϵ 4.41); 260 (log ϵ 4.4); 351 (log ϵ 4.43) (MeOH).

Chen, I.S. et al., *J. Nat. Prod.*, 2001, **64**, 1143-1147

6-Acetyl-1,3-dimethyl-2,4(1H,3H)-pteridinedione, 9CI A-51

6-Acetyl-1,3-dimethylmazine [94591-18-1]




$C_{10}H_{10}N_4O_3$ 234.214

Isol. from the marine polychaete worm *Odontosyllis undecimdonta*. Needles (MeOH). Mp 177-178° (173°).

Kakoi, H. et al., *Heterocycles*, 1995, **41**, 789-797 (isol, synth, pmr, cmr)

6-Acetylhomopterin A-52

6-Acetyl-2-amino-1,7,8,9-tetrahydro-4H-pyrimido[4,5-b][1,4]diazepin-4-one, 9CI. Quench spot [80003-63-0]




C₉H₁₁N₅O₂ 221.218
Naturally occurring diazepine from *Drosophila melanogaster*. Intermed. involved in the biosynth. of Drosopterin, D-941. Greenish-yellow cryst.

Wiederrecht, G.J. *et al.*, *J. Biol. Chem.*, 1981, **256**, 10399 (*isol, struct, ms, pmr*)
Jacobson, K.B. *et al.*, *Biochemistry*, 1982, **21**, 5700 (*isol, struct, pmr, cmr, uv*)
Boyle, P.H. *et al.*, *Tet. Lett.*, 1987, **28**, 5331 (*synth*)

1-Acetyl-4-hydroxy-β-carboline A-53

1-(4-Hydroxy-9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI



C₁₃H₁₀N₂O₂ 226.234

Me ether: 1-Acetyl-4-methoxy-β-carboline

[65236-63-7]
C₁₄H₁₂N₂O₂ 240.261
Alkaloid from the bark and roots of *Ailanthus malabarica* and from the root bark of *Ailanthus altissima* (Simaroubaceae). Prisms. Mp 208° 204-205° Mp 212.5-214°. λ_{max} 238 (log ε 4.15); 270 (log ε 4.26); 288 (log ε 4.23); 370 (log ε 3.82) (EtOH).

Joshi, B.S. *et al.*, *Heterocycles*, 1977, **7**, 193-200 (*isol, uv, ir, pmr, ms, struct*)
Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 390-395 (*isol, uv, ir, pmr, cmr, ms*)
Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 471-473 (*cmr*)
Suzuki, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 2170-2172 (*synth*)
Suzuki, H. *et al.*, *Tetrahedron*, 1997, **53**, 1593-1606 (*synth*)

1-Acetyl-6-hydroxy-β-carboline A-54

1-(6-Hydroxy-9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI

C₁₃H₁₀N₂O₂ 226.234

O-(4-O-Methyl-β-D-glucopyranoside):

BB 4
C₂₀H₂₂N₂O₇ 402.403
Isol. from the oriental crude drug *bombyx batryticatus* (dried larvae of *Bombyx mori*).

Kikuchi, H. *et al.*, *Tet. Lett.*, 2004, **45**, 367-370 (**BB4**)

1-Acetyl-7-hydroxy-β-carboline A-55

1-(7-Hydroxy-9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI. **Arenarine D** [123520-96-7]

C₁₃H₁₀N₂O₂ 226.234
Alkaloid from *Arenaria kansuensis* (Caryophyllaceae). Pale yellow needles (EtOAc). Mp 176-177° dec.

O-(4-O-Methyl-β-D-glucopyranoside):

BB 3
C₂₀H₂₂N₂O₇ 402.403
Isol. from the oriental crude drug, *bombyx batryticatus* (dried silkworms, *Bombyx* spp.).

Me ether: 1-Acetyl-7-methoxy-β-carboline. Arenarine C

[62230-10-8]
C₁₄H₁₂N₂O₂ 240.261
Alkaloid from *Banisteriopsis caapi* (Malpighiaceae) and from the whole plant of *Arenaria kansuensis* (Caryophyllaceae). Light yellow needles (CHCl₃ or hexane/EtOAc). Mp 224-225° dec. (190° dec.). λ_{max} 260 (log ε 4.18); 289 (log ε 4.37); 334 (log ε 4.02) (CHCl₃).

Hashimoto, Y. *et al.*, *Phytochemistry*, 1976, **15**, 1559-1560 (*Me ether, isol, synth, pmr*)
Wu, F. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1808 (*isol, uv, ir, pmr, cmr, ms*)
Kikuchi, H. *et al.*, *Tet. Lett.*, 2004, **45**, 367-370 (**BB 3**)

1-Acetyl-8-hydroxy-β-carboline A-56


1-(8-Hydroxy-9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI

C₁₃H₁₀N₂O₂ 226.234
Alkaloid from *Hypodematum squamuloso-pilosum*. Pale yellow powder (MeOH aq.). Mp 242-243°. λ_{max} 256; 277; 310; 390 (MeOH).

Zhou, T.-S. *et al.*, *Phytochemistry*, 1998, **49**, 1807-1809 (*isol, uv, ir, pmr, cmr, ms*)

3-Acetyl-4-(hydroxymethyl)pyrrole A-57

1-[4-(Hydroxymethyl)-1H-pyrrol-3-yl]ethanone, 9CI. **Verrucarin E** [24445-13-4]



C₇H₉NO₂ 139.154


Pyrrole antibiotic. Metab. of *Myrothecium verrucaria*. Shows cytostatic activity. Cryst. (Et₂O/Me₂CO). Sol. EtOAc; poorly sol. H₂O, acids, bases. Mp 90.5-91°. λ_{max} 260 (ε 17800) (MeOH or EtOH) (Derep). λ_{max} 249 (ε 10000) (EtOH) (Berdy).

Ac:
C₉H₁₁NO₃ 181.191
Cryst. (Et₂O/Me₂CO). Mp 102°.
Härrri, E. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 839 (*isol, props*)
Pfäffli, P. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 1911-1920; 1921-1929 (*struct, biosynth*)

Gossauer, A. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 1698-1704 (*synth*)
Sheldrick, W.S. *et al.*, *Acta Cryst. B*, 1978, **34**, 1248-1253 (*cryst struct*)
Chexal, K.K. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 761 (*biosynth*)
Muchowski, J.M. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 1168 (*synth*)
Adinolfi, A. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 1598-1603 (*isol, cmr, ms*)

2-Acetyl-3-hydroxy-1H-pyrrole A-58

1-(3-Hydroxy-1H-pyrrol-2-yl)ethanone [95232-59-0]



C₆H₇NO₂ 125.127

O-[β-D-Fucopyranosyl-(1→3)-β-D-fucopyranosyl-(1→4)-[6-deoxy-β-D-glucopyranosyl-(1→2)]-6-deoxy-β-D-glucopyranoside]:

C₃₀H₄₇NO₁₈ 709.697
Alkaloid from the starfish *Asterina pectinifera*. Amorph. powder. Mp 240-245°. [α]_D¹⁸ +0.2 (c, 0.3 in MeOH).

Me ether: 2-Acetyl-3-methoxy-1H-pyrrole [70718-05-7]


C₇H₉NO₂ 139.154
Needles (H₂O). Mp 115-116°.

Fisher, B.E. *et al.*, *J.O.C.*, 1964, **29**, 776-781 (*Me ether*)

Zhang, L.-X. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 229-233 (*isol, pmr, cmr, ms*)

3-Acetylindole A-59

1-(1H-Indol-3-yl)ethanone, 9CI. 3-Indolyl methyl ketone [703-80-0]



C₁₀H₉NO 159.187

Constit. of *Strychnos cathayensis*. Cryst. (cyclohexane/EtOH). Spar. sol. H₂O. Mp 194°. pK_a 12.99 (25°, NH).

▶ LD₅₀ (mus, ipr) 300 mg/kg. OB4520000

Picrate:
Yellow cryst. (petrol). Mp 183°.

Oxime: [87052-79-7]
[40747-12-4 (E-form), 40747-13-5 (Z-form)]
C₁₀H₁₀N₂O 174.202
Mp 144-147°. Oximation produces the E-form.


N-Ac: [17537-64-3]
C₁₂H₁₁NO₂ 201.224
Sol. hot H₂O. Mp 151°. Sublimes.

N-Me: 3-Acetyl-1-methylindole [19012-02-3]
C₁₁H₁₁NO 173.214
Cryst. (EtOH). Mp 104-105°. Bp_{0.01} 130-140°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 666A (*ir*)


Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 135C (nmr)
 Baker, J.W. *et al.*, *J.C.S.*, 1946, 461-463 (*synth*, *N-Ac*)
 Pyang, H.-S. *et al.*, *Chem. Comm.*, 1972, 77 (*synth*)
 Rosenberg, E. *et al.*, *Org. Magn. Reson.*, 1976, **8**, 117 (nmr)
 Pindur, U. *et al.*, *Annalen*, 1986, 1621 (*synth*, *ir*, *pmr*)
 Kasahara, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 927 (*synth*)
 Majchrzak, M.W. *et al.*, *Synthesis*, 1986, 956-958 (*N-Me*, *synth*, *pmr*)
 Shner, V.F. *et al.*, *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1989, 272-274 (*N-Ac*, *synth*)
 Bergman, J. *et al.*, *Tetrahedron*, 1990, **46**, 6061 (*synth*)
 Pindur, U. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 727-738 (*E-form*, *oxime*, *synth*, *pmr*, *ms*)
 Yang, C.X. *et al.*, *Synth. Commun.*, 1997, **27**, 2125-2132 (*synth*, *pmr*, *cmr*, *ms*)
 Cheng, M.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 235-239 (*isol*)
 Taira, S. *et al.*, *Tet. Lett.*, 2002, **43**, 8893-8896 (*N-Ac*, *synth*, *pmr*)
 Wynne, J.H. *et al.*, *Synthesis*, 2004, 2277-2282 (*synth*, *ir*, *pmr*, *cmr*)

4-[2-(Acetylmethylamino)ethyl]-4-(4-hydroxy-3-methoxyphenyl)-2,4-cyclohexadien-1-one **A-60**
N-Methyl-4'-O-demethyl-N,7a-secomesebradionone
 [82545-13-9]



$C_{18}H_{21}NO_4$ 315.368
 Achiral molecule. Alkaloid from *Sceletium namaquense* (Aizoaceae). Mp 107.5°.
 2,3,5,6-Tetrahydro, *Me ether*: 4-[2-(Acetylmethylamino)ethyl]-4-(3,4-dimethoxyphenyl)cyclohexanone. *N-Acetyl-N-methyl-N,7a-secomesebrine*
 [82545-07-1]
 $C_{19}H_{27}NO_4$ 333.427
 Alkaloid from *Sceletium namaquense* (Aizoaceae). Oil.
 Jeffs, P.W. *et al.*, *J.O.C.*, 1982, **47**, 3611 (*isol*, *ir*, *pmr*, *ms*, *struct*, *synth*)


2-Acetyl-3-methylindole **A-61**
1-(3-Methyl-1H-indol-2-yl)ethanone, 9CI. *Salvadoricine*
 [16244-23-8]



$C_{11}H_{11}NO$ 173.214
 The first naturally occurring 2-acetylindole alkaloid. Alkaloid from the leaves


of *Salvadora persica* (Salvadoraceae). Needles (hexane). Mp 146-147° (143-144°).
 Ishizumi, K. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 863 (*synth*)
 Jackson, A.H. *et al.*, *Chem. Comm.*, 1978, 779 (*synth*)
 Malik, S. *et al.*, *Tet. Lett.*, 1987, **28**, 163 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)
 Pindur, U. *et al.*, *J. Het. Chem.*, 1992, **29**, 145 (*synth*, *pmr*, *cmr*, *ms*)

2-Acetyl-3-methylpyrazine **A-62**
1-(3-Methylpyrazinyl)ethanone, 9CI.
Methyl (3-methylpyrazinyl) ketone, 8CI
 [23787-80-6]



$C_7H_8N_2O$ 136.153
 Present in spiny lobster *Panulirus argus*. Liq. with burnt, popcorn-like odour. d 1.11. Bp_{0.5} 56°. n_D^{20} 1.5216. Odour threshold 2×10^{-2} ppm in H₂O.
 Mookherjee, B.D. *et al.*, *J.O.C.*, 1972, **37**, 511-513 (*synth*, *ir*, *pmr*, *ms*)
 Wolt, J. *et al.*, *J.O.C.*, 1975, **40**, 1178-1179 (*synth*, *pmr*, *ms*)
 Mihara, S. *et al.*, *J. Agric. Food Chem.*, 1988, **36**, 1242-1247 (*props*)
 Nakamura, S. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 1891-1899 (*glc*, *occur*)
 Cadwallader, K.R. *et al.*, *J. Agric. Food Chem.*, 1995, **43**, 2432-2437 (*detn*, *occur*)

2-Acetyl-4-methylpyridine **A-63**
1-(4-Methyl-2-pyridinyl)ethanone, 9CI.
Methyl 4-methyl-2-pyridyl ketone, 8CI. 2-Acetyl-4-picoline
 [59576-26-0]



C_8H_9NO 135.165
 Component of tobacco smoke and fig leaf absolute (*Ficus carica*). Cryst. (petrol). Sol. EtOH. Mp 33-34°. Bp₁₅ 95-97°.
Oxime (E-): [23089-35-2]
 $C_8H_{10}N_2O$ 150.18
 Used as a 0.01M soln. in EtOH for photometric detn. of Co (λ_{max} 325 nm, ϵ 11800, EtOH aq.), Cu(I) (λ_{max} 409 nm, ϵ 5300, EtOH aq.), extraction-photometric detn. of Fe(II) (λ_{max} 541 nm, ϵ 14000, isopentanol). Cryst. (C₆H₆/petrol). Mp 98-99° Mp 125-128°.

Semicarbazone:
 Solid. Mp 195-198°.
 [18103-84-9]
 Case, F.H. *et al.*, *J.A.C.S.*, 1956, **78**, 5842 (*synth*)
 Nishimoto, N. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 88; *CA*, **55**, 13421b (*synth*)
 Case, F.H. *et al.*, *J. Het. Chem.*, 1968, **5**, 161 (*synth*, *oxime*)
 Schilt, A. *et al.*, *Talanta*, 1969, **16**, 448 (*detn*, *Co*, *Cu*, *Fe*)

2-Acetyl-6-methylpyridine **A-64**
1-(6-Methyl-2-pyridinyl)ethanone, 9CI.
Methyl 6-methyl-2-pyridyl ketone, 8CI. 6-Acetyl-2-picoline
 [6940-57-4]

C_8H_9NO 135.165
 Minor component of rum and tobacco smoke. Flavour modifying agent for coffee. Liq. Bp₁ 46°.
Oxime (E-): [23089-39-6]
 $C_8H_{10}N_2O$ 150.18
 Used as a 0.01M soln. in EtOH for photometric detn. of Cu(I) (λ_{max} 422 nm, ϵ 8400, EtOH aq.). Cryst. (petrol). Sol. EtOH. Mp 49-50°.

Dimethylhydrazone: [33785-80-7]
 $C_{10}H_{15}N_3O$ 193.248
 Liq. Bp_{0.1} 53-54°. n_D^{20} 1.5370.
 [18103-88-3]
 Case, F.H. *et al.*, *J. Het. Chem.*, 1968, **5**, 161 (*synth*, *oxime*)
 Schilt, A. *et al.*, *Talanta*, 1969, **16**, 448 (*detn*, *Cu*)
 Newkome, G.R. *et al.*, *J.O.C.*, 1972, **37**, 1329 (*deriv*)
 Zanger, M. *et al.*, *Anal. Chem.*, 1974, **46**, 2042 (*pmr*)
 Amin, H.B. *et al.*, *J.C.S. Perkin 2*, 1979, 624 (*synth*)

3-Acetyl-2-methylpyridine **A-65**
1-(2-Methyl-3-pyridinyl)ethanone, 9CI.
Methyl 2-methyl-3-pyridyl ketone, 8CI. 3-Acetyl-2-picoline
 [1721-12-6]

C_8H_9NO 135.165
 Pheromone present in ferret (*Mustela putorius*) urine. Prisms. Mp 30-31°. Bp₁₅ 99-100°.
Dipicrate:
 Needles (EtOH). Mp 174-176°.
 Baumgarten, P. *et al.*, *Ber.*, 1939, **72**, 563 (*synth*)
 Sanders, E.B. *et al.*, *J.O.C.*, 1978, **43**, 324 (*synth*, *pmr*, *ir*)

3-Acetyl-4-methylpyridine **A-66**
1-(4-Methyl-3-pyridinyl)ethanone, 9CI.
Methyl 4-methyl-3-pyridyl ketone, 8CI. 3-Acetyl-4-picoline
 [51227-30-6]

C_8H_9NO 135.165
 Pheromone from ferret (*Mustela putorius*) urine. Volatile liq. Bp_{1.5} 57-58° (lit. gives a pressure range).
Picrate:
 Solid (EtOH). Mp 146-147°.
 Webb, J.L. *et al.*, *J.A.C.S.*, 1944, **66**, 1456 (*synth*)
 Deuchert, K. *et al.*, *Chem. Ber.*, 1979, **112**, 2045 (*synth*, *pmr*)
 Rey, A.W. *et al.*, *Can. J. Chem.*, 1992, **70**, 2922 (*synth*, *ir*, *pmr*, *cmr*, *ms*)

4-Acetyl-2-methylpyridine **A-67**
1-(2-Methyl-4-pyridinyl)ethanone, 9CI.
Methyl 2-methyl-4-pyridyl ketone, 8CI. 4-Acetyl-2-picoline
 [2732-28-7]

C_8H_9NO 135.165
 Component of roasting coffee aroma. Pheromone present in ferret (*Mustela putorius*) urine. Liq. Bp₁₀ 120-124° Bp₄ 130°.

Picrate: [31931-61-0]
Solid (EtOH). Mp 174-175°.

[80882-67-3]

Suzuki, Y. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 795; *CA*, **55**, 24742d (*synth*)
Eilhauer, H.D. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1965, **298**, 131 (*synth*)
Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1966, **4**, 398 (*synth*)

4-Acetyl-3-methylpyridine A-68

1-(3-Methyl-4-pyridinyl)ethanone, 9CI.
Methyl 3-methyl-4-pyridyl ketone, 8CI. 4-Acetyl-3-picoline
[82352-00-9]

C₈H₉NO 135.165

Pheromone from ferret (*Mustela putorius*) urine. Bp₁ 93-95°.

Hibino, S. *et al.*, *J. Het. Chem.*, 1990, **27**, 1751 (*synth, pmr, ms*)

5-Acetyl-2-methylpyridine A-69

1-(6-Methyl-3-pyridinyl)ethanone, 9CI.
Methyl 6-methyl-3-pyridyl ketone, 8CI. 5-Acetyl-2-picoline
[36357-38-7]

C₈H₉NO 135.165

Minor component of palmarosa oil and tobacco smoke. Used in perfume compositions. Liq. Bp₁₅ 105-106°.


2,4-Dinitrophenylhydrazones: [40624-50-8]
Solid. Mp 224-225°.

[65305-38-6]

Muruoka, M. *et al.*, *Nippon Kagaku Zasshi*, 1961, **82**, 1279; *CA*, **59**, 563b (*synth*)
Hawkins, E.G.E. *et al.*, *J.C.S. Perkin I*, 1972, 2882 (*synth*)
Kozerski, L. *et al.*, *Org. Magn. Reson.*, 1977, **9**, 395 (*cmr*)
Markova, N.K. *et al.*, *Zh. Org. Khim.*, 1984, **20**, 962 (*synth*)

4-Acetyl-6-methyl-2(1H)-pyridinone A-70

4-Acetyl-2-hydroxy-6-methylpyridine.
Streptokordin




C₈H₉NO₂ 151.165

Prod. by a marine-derived *Streptomyces* sp. KORDI-3238. Cytotoxic. Amorph. powder. λ_{max} 217 (ε 10500) (MeOH).

Jeong, S.-Y. *et al.*, *J. Antibiot.*, 2006, **59**, 234-240 (*isol, pmr, cmr*)

3-Acetyl-4-methylpyrrole A-71

1-(4-Methyl-1H-pyrrol-3-yl)ethanone, 9CI. *Deoxyverrucarin E*
[18818-30-9]



C₇H₉NO 123.154

Pyrrole antibiotic. Metab. of *Eupenicillium hirayamae*. Possesses antitumour activity. Cryst. (C₆H₆/hexane). Sol. MeOH, Et₂O; poorly sol. H₂O, hexane. Mp 115°. λ_{max} 247 (ε 11070); 277 (ε 2204) (EtOH) (Berdy).

Semicarbazone:

Pale-yellow needles (MeOH). Mp 195°.

V. Leusen, A.M. *et al.*, *Tet. Lett.*, 1972, 5337 (*pmr*)


Arndt, R.R. *et al.*, *Phytochemistry*, 1974, **13**, 1865-1870 (*isol, struct*)

Cheng, D.O. *et al.*, *Tet. Lett.*, 1977, 1469 (*synth*)

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 925 (*synth, uv, ir, pmr*)

3-Acetyl-2,7-naphthyridine A-72

1-(2,7-Naphthyridin-3-yl)ethanone, 9CI
[73607-00-8]



C₁₀H₈N₂O 172.186


Alkaloid from valerian (*Valeriana officinalis*).

Janot, M.M. *et al.*, *Ann. Pharm. Fr.*, 1979, **37**, 413-420 (*isol, cryst struct*)

Barbu, E. *et al.*, *Heterocycl. Commun.*, 2000, **6**, 25-28 (*synth*)

2-Acetylpyridine A-73

1-(2-Pyridinyl)ethanone, 9CI. *Methyl 2-pyridyl ketone*. 2-Acetopyridine. FEMA 3251
[1122-62-9]



C₇H₇NO 121.138

Present in various cooked and treated food. Present in coriander seed and black tea. Organoleptic, flavouring agent. Liq. with roasted odour. Bp 192°. n_D²⁰ 1.5203. pK_a 2.64 (25°). Turns yellow in air.

Hydrochloride: Mp 183-185° dec.

Methiodide: Mp 161°.

Ethiodide: Mp 205°.

Picrate:

Yellow cryst. (EtOH). Mp 131°.

Oxime: Acepox

[1758-54-9]

C₇H₈N₂O 136.153

Plant protectant. Used for photometric detn. of Re (λ_{max} 490 nm, ε 3800). Prisms (EtOH). Sol. H₂O. Mp 121°.

Hydrazones: [59742-91-5]

C₇H₉N₃ 135.168

Used as 0.5mM aq. soln. for fluorimetric detn. of Tc(VII) (λ_{max} 430 nm, 0.01-10μg/ml, =1.4M HCl). Cryst. (EtOH). Sol. H₂O, EtOH.

Phenylhydrazones: [7734-05-6]

Yellow cryst. (EtOH). Mp 155°.

Thiosemicarbazones: [6839-90-3]

C₈H₁₀N₄S 194.26

Used as EtOH soln. or 1% Me₂CO aq. soln. for photometric detn. of Mo (λ_{max} 470 nm, ε 17000), V (λ_{max} 400 nm, ε 56000, pH 3.5). Cryst. (EtOH). Sol. EtOH, DMF, Me₂CO.

Ethylene ketal:

C₉H₁₁NO₂ 165.191

Bp_{0.8} 91-95°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 779A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 310A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1542B (*ir*)

Thompson, R.J. *et al.*, *Anal. Chim. Acta*, 1964, **31**, 590 (*oxime, detn, Re*)

Miyajama, G. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 429 (*cmr*)

Lesman, T. *et al.*, *Org. Mass Spectrom.*, 1973, **7**, 1321 (*ms*)

Seth-Paul, W.A. *et al.*, *Spectrochim. Acta A*, 1974, **30**, 1817 (*ir*)

Reimann, E. *et al.*, *Annalen*, 1976, 1351 (*synth*)

Klayman, D.L. *et al.*, *J. Med. Chem.*, 1979, **22**, 855 (*synth, thiosemicarbazone*)

Grases, F. *et al.*, *Anal. Chim. Acta*, 1984, **166**, 71 (*synth, detn, Te*)

Thimmaiah, K.N. *et al.*, *Microchem. J.*, 1985, **32**, 8 (*detn, Mo*)

Norman, M.H. *et al.*, *J.O.C.*, 1987, **52**, 226 (*deriv, synth, ir, pmr, cmr*)


Cook, I.B. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1493 (*cmr*)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 35 (*occur*)

Girardot, M. *et al.*, *J.O.C.*, 1998, **63**, 10063-10068 (*synth*)

2-Acetylpyrrole A-74

1-(1H-Pyrrol-2-yl)ethanone, 9CI. *Methyl 2-pyrrolyl ketone*. 2-Acetopyrrole. *Pseudoacetylpyrrole*. FEMA 3202
[1072-83-9]



C₆H₇NO 109.127

Found in *Valeriana officinalis* (Valerianaceae), *Camellia sinensis* (Theaceae), *Paeonia moutan* (Chinese drug Botani) (Peoniaceae), *Lycium chinense* (Solanaceae). Prod. by

Streptomyces sp. A-5071. Organoleptic which contributes to many aromas, including tobacco smoke. Flavouring agent. Possesses hepatoprotective props. Cryst. (petrol) with bread-like odour. Sol. H₂O. Mp 90°. Bp 220°. λ_{max} 247; 287 (MeOH) (Berdy).

Oxime: [63547-59-1]

C₆H₈N₂O 124.142

Mp 145-146°.

Phenylhydrazones: Mp 146°.

2,4-Dinitrophenylhydrazones:

Reddish-brown cryst. Mp 297° dec.

Semicarbazones: Mp 190°.

N-(4-Methylbenzenesulfonyl): [129666-99-5]

C₁₃H₁₃NO₃S 263.317

Solid (CHCl₃/hexane). Mp 110-111°.

N-Me: 2-Acetyl-1-methylpyrrole. FEMA 3184

[932-16-1]
 C₇H₉NO 123.154
 Bp 200-202°. n_D²⁰ 1.5420.

N-Et: 2-Acetyl-1-ethylpyrrole. FEMA 3147

[39741-41-8]
 C₈H₁₁NO 137.181
 Bp₁₂ 82°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 567D; 568A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 5B; 5C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1447B; 1447C (ir)

Cionga, E. et al., C. R. Hebd. Seances Acad. Sci., 1935, 200, 780 (isol)

Sugisawa, H. et al., Chem. Ind. (London), 1958, 887 (synth)

Budzikiewicz, H. et al., J.C.S., 1964, 1949 (ms)

Abraham, R.J. et al., J.C.S. Perkin 2, 1974, 1004 (cmr)

Sannai, A. et al., Agric. Biol. Chem., 1983, 47, 2397 (isol)

Miyayawa, M. et al., Agric. Biol. Chem., 1983, 47, 2925; 1984, 48, 2847 (isol)

Kakushima, M. et al., J.O.C., 1983, 48, 3214 (synth)

Garrido, D.O.A. et al., J.O.C., 1984, 49, 2619 (synth)

Anderson, H.J. et al., Can. J. Chem., 1985, 63, 896 (synth)

Eyley, S.C. et al., Tet. Lett., 1985, 26, 4649 (synth)

Lewis, R.J. et al., Food Additives Handbook, Van Nostrand Reinhold International, New York, 1989, ADA 375

Pindur, H. et al., J. Het. Chem., 1989, 26, 1563 (ir)

Kuroda, Y. et al., Tet. Lett., 1989, 30, 2411 (synth)

Ito, M. et al., Agric. Biol. Chem., 1991, 55, 2117 (isol, props)

Goldberg, Yu. et al., Synth. Commun., 1991, 21, 557 (N-alkyl derivs, synth)

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 2, 215; 468; 560

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 944; 1709; 1853-1854 (occur, props, use)


Hou, S. et al., Macromolecules, 2003, 36, 3826-3832 (synth, N-tosyl)

Camarillo, C.A. et al., Acta Cryst. E, 2007, 63, o2593-o2594 (cryst struct)

2-Acetyl-4(3H)-quinazolinone

A-75

[17244-28-9]



C₁₀H₈N₂O₂ 188.185

Fungal metab. prod. from a toxic strain of *Fusarium culmorum*. Also isol. from cultures of *Fusarium sambucinum*. Prisms (EtOAc or Me₂CO). Mp 197-200° dec Mp 205°. Has also been descr. as a transformation prod. of 2-(1-Hydroxyethyl)-4(3H)-quinazolinone, H-499, and of 2-Pyruvoylaminobenzamide, from *P. chrysogenum*, *P. notatum* and *Colletotrichum lagenarium*. Although isol. from the *P. chrysogenum* fermentation, it was considered in that instance to be an

artifact. λ_{max} 302 (ε 11600) (MeOH) (Berdy). λ_{max} 232 (H₂O) (Berdy).

Suter, P.J. et al., J.C.S. (C), 1967, 2240 (synth)


Blight, M.M. et al., J.C.S. Perkin 1, 1974, 1691 (isol, uv, ir, pmr)

Niederer, D. et al., Tet. Lett., 1992, 33, 3997 (isol)

6-Acetyl-1,2,3,4-tetrahydro-pyridine

A-76

Methyl 3,4,5,6-tetrahydro-2-pyridyl ketone, 8CI. 2-Acetyl-3,4,5,6-tetrahydropyridine. 2-Acetyl-Δ¹-piperidine. 2-Acetyl-Δ²-piperideine. 1-(1,4,5,6-Tetrahydro-2-pyridinyl)ethanone, 9CI [27300-27-2]



C₇H₁₁NO 125.17

Tautomeric, 1H-form predominates ca. 2:1. Constit. of the leaves of *Semnostachya menglaensis* (preferred genus name *Strobilanthes*). Key aroma component of bread. Responsible for mousy taint in wines. Unstable liq. Bp₃ 65-67°. Odour threshold 0.06 ng/L in air.

Hydrochloride: Mp 112-119°.

Hunter, I.R. et al., Cereal Chem., 1969, 46, 189 (isol)

Büchi, G. et al., J.O.C., 1971, 36, 609 (ir, uv, pmr, ms, synth)

Strauss, C.R. et al., Chem. Ind. (London), 1984, 109 (isol)

De Kimpe, N. et al., J.O.C., 1993, 58, 2904 (synth, pmr, cmr, ms)

De Kimpe, N. et al., Tetrahedron, 1995, 51, 2387 (synth)

Hofman, T. et al., J. Agric. Food Chem., 1998, 46, 616-619; 2270-2277 (synth, ms, bibl)

Naef, R. et al., J. Agric. Food Chem., 2005, 53, 9161-9164 (isol, pmr, ms)

Harrison, T.J. et al., J.O.C., 2005, 70, 10872-10874 (synth)

Adams, A. et al., Chem. Rev., 2006, 106, 2299-2319 (rev)


Snowdon, E.M. et al., J. Agric. Food Chem., 2006, 54, 6465-6474 (rev)

Fuganti, C. et al., Tetrahedron, 2007, 63, 4762-4767 (synth)

Acidissiminol

A-77

[126006-00-6]



C₂₅H₃₁NO₃ 393.525

Alkaloid from the fruit of *Limonia acidissima* (wood apple) (Rutaceae). Cryst. (EtOAc/hexane). Mp 85-87°.

O-Ac: N-Benzoyl-4-(4-acetoxyneryloxy)-phenethylamine

[33055-26-4]

C₂₇H₃₃NO₄ 435.562

Alkaloid from fruit of *Swinglea glutinosa* (Rutaceae). Cryst. (EtOAc/hexane). V. sol. MeOH. Mp 64-65°. Opt. inactive. Related to Severine.

O-Octadecanoyl: Acidissiminin

[126005-91-2]

C₄₃H₆₅NO₄ 659.991

Alkaloid from the fruits of *Limonia acidissima* (wood apple) (Rutaceae). Amorph. solid (EtOAc/hexane). Mp 65°. [α]_D²⁰ 0 (CHCl₃). λ_{max} 225; 277 (sh); 286 (MeOH) (Derrep).

6',7'-Epoxide: Acidissiminol epoxide.

Severine†

[139165-01-8]

[33055-25-3]

C₂₅H₃₁NO₄ 409.524

Alkaloid from fruits of *Limonia acidissima* (wood apple) (Rutaceae). Cryst. (MeOH). Mp 146-149° (141-143°). Severine originally considered to be a diol with formula C₂₅H₃₃NO₄. Identity of Acidissiminol epoxide and Severine not confirmed; stereochem. not established.

6',7'-Epoxide, octadecanoyl: Acidissiminin epoxide. Severine palmitate

[139083-09-3]

C₄₃H₆₅NO₅ 675.99

Alkaloid from the fruit of *Limonia acidissima* (wood apple), leaves of *Severinia buxifolia* and from *Pamburus missionis*, *Atalantia monophylla* and *Hesperethusa crenulata* (preferred genus name *Naringi*) (Rutaceae). Cryst. (C₆H₆/hexane). Mp 105-106° Mp 113-114°. Identity of Acidissiminin epoxide and Severin palmitate not definitely establ. Mps are similar.

6',7'-Dihydro, 6',7'-dihydroxy: Dihydroxy-acidissiminol

[160387-10-0]

C₂₅H₃₃NO₅ 427.539

Alkaloid from fruits of *Limonia acidissima* (wood apple) (Rutaceae). Amorph. semi-solid.

Dreyer, D.L. et al., Tetrahedron, 1967, 23, 4613; 1970, 26, 5745; 1980, 36, 827 (Severine, Severine palmitate, N-Benzoyl-4-(4-acetoxyneryloxy)phenethylamine)

Ghosh, P. et al., J. Nat. Prod., 1989, 52, 1323;

1991, 54, 1389 (Acidissiminol, Acidissiminin)

Ghosh, P. et al., Phytochemistry, 1994, 37, 757 (Dihydroxyacidissiminol, Acidissiminol epoxide)

Acidomycin

A-78

4-Oxo-2-thiazolidinehexanoic acid, 9CI.


2-(5-Carboxypentyl)-4-thiazolidone, 8CI.

Actithiazic acid. Cinnamonin. Mycobactin.

Thiazolidomycin. PA 95. Antibiotic

6604-4. Antibiotic PA 95

[539-35-5]



C₉H₁₅NO₃S 217.288

Amino acid derived antibiotic. Abs. config. not known. Log P 0.86 (calc).


▶ LD₅₀ (mus, ipr) 2500 mg/kg. XJ6000000**(-)-form** [28223-69-0]

Prod. by *Streptomyces* spp. Active against mycobacteria *in vitro*. Needles (H₂O, MeOH or Me₂CO). Sol. MeOH, CHCl₃, bases; fairly sol. CH₂Cl₂; poorly sol. H₂O, acids, hexane, C₆H₆. Mp 139-140°. [α]_D²³ -54 (MeOH). [α]_D²⁵ -60 (c, 1 in EtOH). Blue fluor. in uv light. Racemised by dil. alkalis. λ_{max} 245 (MeOH) (Berdy). ▶ LD₅₀ (mus, ivn) 1500 mg/kg, LD₅₀ (mus, ipr) 2500 mg/kg, LD₅₀ (mus, scu) 20000 mg/kg.

*Me ester:*Needles. Mp 53-54°. [α]_D²⁵ -50.9 (MeOH).**(±)-form** [106448-99-1]Needles (H₂O or CHCl₃). Mp 122-123° (116-117°).McLamore, W.M. *et al.*, *J.A.C.S.*, 1952, **74**, 2946-2947 (*struct*)Sobin, B.A. *et al.*, *J.A.C.S.*, 1952, **74**, 2947-2948 (*isol*)Miyake, A. *et al.*, *Pharm. Bull.*, 1953, **1**, 84-88 (*isol, struct*)Japan. Pat., 1954, ((Takeda))54 448; *CA*, **48**, 14130d (*synth*)Caltrider, P.G. *et al.*, *Antibiotics (N.Y.)*, 1967, **1**, 666-668 (*rev*)Lokshin, G.B. *et al.*, *Antibiotiki (Moscow)*, 1970, **15**, 14-18; *CA*, **72**, 109788a (*isol*)Eisenberg, M.A. *et al.*, *Antimicrob. Agents Chemother.*, 1982, **21**, 5-10 (*activity*)Nadkarni, R.S. *et al.*, *J. Antibiot.*, 1983, **36**, 1567-1568 (*activity*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CMP885**Acinetobactin**

A-79

[160472-93-5]


C₁₆H₁₈N₄O₅ 346.342

Closely related to Anguibactin, A-1009. Isol. from low-iron cultures of *Acinetobacter baumannii* ATCC 19606 and *Acinetobacter haemolyticus*. Siderophore. Sol. H₂O. λ_{max} 250; 317 (MeOH) (Berdy). Yamamoto, S. *et al.*, *Arch. Microbiol.*, 1994, **162**, 249 (*isol, pmr, cmr, ms*)

Acinetoferrin

A-80

[156737-07-4]


C₂₈H₄₈N₄O₉ 584.709Isol. from *Acinetobacter haemolyticus*.

Siderophore.

Okujo, N. *et al.*, *BioMetals*, 1994, **7**, 170-176 (*isol, pmr, cmr, ms*)Wang, Q.X. *et al.*, *J.O.C.*, 1998, **63**, 1491-1495 (*synth, pmr*)Luo, M. *et al.*, *J.A.C.S.*, 2005, **127**, 1726-1736 (*propos*)**Acivicin, INN, USAN**

A-81

α-Amino-3-chloro-4,5-dihydro-5-isoxazoleacetic acid, 9CI. AT 125. NSC 163501. U 42126. Antibiotic AT 125. Antibiotic U 42126 [52583-41-2]

C₅H₇ClN₂O₃ 178.575

Amino acid antibiotic.

(α,S,5S)-form [42228-92-2]

Prod. by *Streptomyces sviveus*. Antineoplastic agent. Cryst. (MeOH aq.). Sol. H₂O; fairly sol. MeOH; poorly sol. butanol, hexane. Mp 228-230°. [α]_D²⁰ +135 (c, 0.159 in H₂O). Related to Ibotenic acid, I-8.

▶ NY2103000


4*S*-Hydroxy: Antibiotic U 43795. U 43795 [54549-02-9]C₅H₇ClN₂O₄ 194.574

From *Streptomyces sviveus*. Antineoplastic, antimetabolite. Sol. H₂O; poorly sol. Me₂CO, hexane.

Hanka, L.J. *et al.*, *Antimicrob. Agents Chemother.*, 1973, **3**, 425; 1975, **7**, 807 (*isol, propos*)Martin, D.G. *et al.*, *Tet. Lett.*, 1973, 2549 (*isol, abs config*)Martin, D.G. *et al.*, *J. Antibiot.*, 1975, **28**, 91 (*deriv*)Kelly, R.C. *et al.*, *J.A.C.S.*, 1979, **101**, 1054 (*synth*)Silverman, R.B. *et al.*, *J.A.C.S.*, 1981, **103**, 7357 (*synth*)Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 12317O'Dwyer, P.J. *et al.*, *J. Clin. Oncol.*, 1984, **2**, 1064 (*rev, pharmacol*)Earhart, W.D. *et al.*, *Adv. Enzyme Regul.*, 1985, **24**, 179 (*rev, pharmacol*)Baldwin, J.E. *et al.*, *Tetrahedron*, 1985, **41**, 5241 (*synth, bibl*)Mzengeza, S. *et al.*, *J.O.C.*, 1988, **53**, 4074 (*synth, bibl*)Cintas, P. *et al.*, *Tetrahedron*, 1991, **47**, 6079 (*synth*)**Aclidinomycin B**

A-82

[342814-66-8]

C₂₁H₂₅N₃O₇ 431.444


Related to Naphthyridinomycin A, N-32. Prod. by *Streptomyces halstedii* KB012 and a marine-derived *Streptomyces* sp. [α]_D¹⁸ +188 (c, 0.16 in MeOH). λ_{max} 286 (ε 21500); 398 (ε 5000) (MeOH).

N-De-Me: Aclidinomycin DC₂₀H₂₃N₃O₇ 417.418Prod. by a marine-derived *Streptomyces* sp.**9-Deoxy: Aclidinomycin A**

[342814-65-7]


C₂₁H₂₅N₃O₆ 415.445Prod. by *Streptomyces halstedii* KB012.[α]_D¹⁸ 0 (MeOH). λ_{max} 282 (ε 19000); 403 (ε 3500) (MeOH).Thorwest, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2001, (*marine, isol*)Cang, S. *et al.*, *J. Antibiot.*, 2001, **54**, 304-307 (*isol, pmr, cmr*)**Aclidinomycin C**

A-83

C₁₉H₂₁N₃O₆ 387.391Prod. by a marine-derived *Streptomyces* sp.Thorwest, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2001,**Aconcaguine**

A-84


[96624-39-4]

C₂₀H₂₃NO₆ 373.405

Alkaloid from the twigs of *Berberis actinacantha* (Berberidaceae). Yellow amorph. solid. Positive opt. rotn.

Weiss, I. *et al.*, *Chem. Comm.*, 1985, 3-4 (*uv, ir, pmr, cmr, ms, struct*)**Aconitane-1,4,6,7,8,14,16-heptol**

A-85

C₁₈H₂₇NO₇ 369.414

(1 α ,5 β ,6 β ,14 α ,16 β)-form O^6, O^{16} -Di-Me, N-Et: **Delbine**[†]


[1356-54-3]

C₂₂H₃₅NO₇ 425.521Alkaloid from the roots of *Delphinium bonvalotii* (Ranunculaceae). Mp 116-118°. [α]_D²⁷ +53.3 (c, 0.1 in MeOH). O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **Sinomontanine H**C₃₂H₄₄N₂O₉ 600.708Alkaloid from *Aconitum sinomontanum*. [α]_D¹⁷ -28.8 (c, 0.5 in CHCl₃). O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 8-Ac: **Hispaconitine**

[128581-45-3]

C₂₆H₄₁NO₈ 495.612Alkaloid from the roots of *Aconitum barbatum* var. *hispidum* (Ranunculaceae). Mp 185-187°. [α]_D +43.4 (c, 1.19 in CHCl₃).7-Hydroperoxide, O^1, O^6, O^{14}, O^{16} -tetra-Me, N-Et: **Lineariline**C₂₄H₃₉NO₈ 469.574Alkaloid from the roots of *Delphinium linearilobum*. Amorph. solid. [α]_D²⁰ +18.4 (c, 0.5 in CHCl₃). λ_{\max} 201 (log ϵ 4); 226 (sh) (log ϵ 3.1); 257 (log ϵ 2.5) (MeOH).Jiang, Q.P. et al., *Heterocycles*, 1985, **23**, 11 (Delbine)Lao, A. et al., *Heterocycles*, 1990, **31**, 27 (Hispaconitine)Peng, C.-S. et al., *Youji Huaxue*, 2005, **25**, 1235-1239; *CA*, **144**, 103949n (Sinomontanine H)Kolak, U. et al., *Phytochemistry*, 2006, **69**, 2170-2175 (Lineariline)**Aconitane-1,4,7,8,9,14,16-heptol**

A-86

**(1 α ,14 α ,16 β)-form**C₁₈H₂₇NO₇ 369.414**(1 α ,14 α ,16 β)-form** O^{14}, O^{16} -Di-Me, N-Et: **Sinomontanine E**C₂₂H₃₅NO₇ 425.521Alkaloid from the roots of *Aconitum sinomontanum*. [α]_D¹⁷ +7.8 (c, 0.5 in CHCl₃). O^1, O^{14}, O^{16} -Tri-Me, 4-O-(2-acetamidobenzoyl): **Sinomontanine F**

[872977-64-5]

C₃₀H₄₀N₂O₉ 572.654Alkaloid from the roots of *Aconitum sinomontanum*. [α]_D¹⁷ +44.4 (c, 0.75 in CHCl₃). O^1, O^{14}, O^{16} -Tri-Me, N-Et: **Ranaconine. Hydroxylappaconine**

[70474-32-7]

C₂₃H₃₇NO₇ 439.548Alkaloid detected by glc-ms in *Aconitum leucostomum*. Alkaline hydrol.

product of Ranaconitine (Ranunculaceae). Mp 107-109°.

 O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-aminobenzoyl): **N-Deacetylraconitine**

[82872-80-8]

C₃₀H₄₂N₂O₈ 558.67Alkaloid from the roots of *Aconitum finetianum* (Ranunculaceae). Shows good analgesic activity. Mp 125-127°. O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **Ranaconitine**


[1360-76-5]

C₃₂H₄₄N₂O₉ 600.708Alkaloid from *Aconitum ranunculaeifolium*, *Aconitum sinomontanum*, *Aconitum finetianum* and *Aconitum barbatum* var. *puberulum* (Ranunculaceae). Mp 132-134°. [α]_D²⁷ +33.2 (c, 0.5 in CHCl₃). O^1, O^{14}, O^{16} -Tri-Me, 4-O-(2-acetamidobenzoyl), N-Ac: **Sinaconitine A**C₃₂H₄₂N₂O₁₀ 614.691Alkaloid from the roots of *Aconitum sinomontanum*. Prisms (petrol/Et₂O). Mp 150-151.5°. [α]_D²⁰ +35 (c, 1 in CHCl₃).**(1 β ,14 α ,16 β)-form** O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **Puberanine**

[83685-20-5]

C₃₂H₄₄N₂O₉ 600.708Alkaloid from the roots of *Aconitum barbatum* var. *puberulum* (Ranunculaceae). Amorph. [α]_D²⁰ +16.6 (c, 0.6 in CHCl₃).Mollov, N. et al., *Dokl. Bulg. Akad. Nauk*, 1964, **17**, 251; *CA*, **61**, 12324h (Ranaconitine, *isol.*, *ir*)Pelletier, S.W. et al., *Tet. Lett.*, 1978, 5045 (Ranaconitine, *pmr*, *cmr*, *struct*)Pelletier, S.W. et al., *Can. J. Chem.*, 1979, **57**, 1652 (*cmr*)Plugar, V.N. et al., *Khim. Prir. Soedin.*, 1982, **18**, 80; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 75 (Ranaconine)Jiang, S. et al., *Yaoxue Xuebao*, 1982, **17**, 283; 1983, **18**, 440; *CA*, **97**, 20736a; **100**, 20505e (Deacetylraconitine)Yu, D. et al., *Planta Med.*, 1983, **49**, 85 (Puberanine)Peng, C.-S. et al., *Yaoxue Xuebao*, 2000, **35**, 201-203 (Sinomontanine E)Peng, C.-S. et al., *Youji Huaxue*, 2005, **25**, 1235-1239; *CA*, **144**, 103949n (Sinomontanine F)Tan, J.-J. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 535-539 (Sinaconitine A)**Aconitane-1,4,8,9,10,14,16-heptol**

A-87

C₁₈H₂₇NO₇ 369.414**(1 α ,5 β ,14 α ,16 β)-form** O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-aminobenzoyl): **Sepaconitine**


[114622-05-8]

C₃₀H₄₂N₂O₈ 558.67Alkaloid from the aerial parts of *Aconitum septentrionale* (Ranunculaceae). Mp 250-253°. [α]_D²⁰ +25 (c, 0.60 in CHCl₃). O^1, O^{14}, O^{16} -Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **N-Acetylsepaconitine**

[118201-55-1]

C₃₂H₄₄N₂O₉ 600.708Alkaloid from aerial parts of *Aconitum leucostomum* (Ranunculaceae). Amorph.Usmanova, S.K. et al., *Khim. Prir. Soedin.*, 1987, **23**, 879; *Chem. Nat. Compd. (Engl. Transl.)*, 734 (Sepaconitine)Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1988, **24**, 556; *Chem. Nat. Compd. (Engl. Transl.)*, 472 (N-Acetylsepaconitine)**Aconitane-1,6,7,8,10,14,16-heptol**

A-88

C₁₈H₂₇NO₇ 369.414**(1 α ,5 β ,6 β ,14 α ,16 β)-form** O^1, O^{14}, O^{16} -Tri-Me, N-Et: **Acoseptrine**

[146028-66-2]

C₂₃H₃₇NO₇ 439.548Alkaloid from roots of *Aconitum septentrionale*. Cryst. (Me₂CO/hexane). Mp 105-107°. [α]_D +19.6 (c, 0.245 in CHCl₃). O^7, O^8 -Methylene, O^1, O^{16} -di-Me, N-Et, 6-Ac: **Anthriscifolcine C**

[948308-30-3]

C₂₅H₃₇NO₈ 479.569Alkaloid from *Delphinium anthriscifolium* var. *savatieri*. Needles. Mp 222-224°. [α]_D²⁰ -11.4 (c, 0.5 in CHCl₃). O^7, O^8 -Methylene, O^1, O^{14}, O^{16} -tri-Me, N-Et: **Anthriscifolcine E**

[948308-32-5]


C₂₄H₃₇NO₇ 451.559Alkaloid from *Delphinium anthriscifolium* var. *savatieri*. Amorph. powder. Mp 150-152°. [α]_D²⁰ -34.8 (c, 0.5 in CHCl₃). O^7, O^8 -Methylene, O^1, O^{14}, O^{16} -tri-Me, N-Et, 6-Ac: **Anthriscifolcine D**

[948308-31-4]

C₂₆H₃₉NO₈ 493.596Alkaloid from *Delphinium anthriscifolium* var. *savatieri*. Amorph. powder. Mp 185-187°. [α]_D²⁰ -41.3 (c, 0.5 in CHCl₃).Sayed, H.M. et al., *J. Nat. Prod.*, 1992, **55**, 1595-1606 (Acoseptrine)Song, L. et al., *Chem. Pharm. Bull.*, 2007, **55**, 918-921 (Anthriscifolcines C-E)

Aconitane-1,4,6,8,14,16-hexol

A-89

(1 α ,5 β ,6 α ,14 α ,16 β)-formC₁₈H₂₇NO₆ 353.414(1 α ,5 β ,6 α ,14 α ,16 β)-formO¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, N-Et, 4-Ac:**Kirimine**

[161068-74-2]

C₂₆H₄₁NO₇ 479.612Alkaloid from *Aconitum kirinense*.(1 α ,5 β ,6 β ,14 α ,16 β)-formO⁶,O¹⁴,O¹⁶-Tri-Me, N-Et: **Akiramidine**

[287724-50-9]

C₂₃H₃₇NO₆ 423.548Alkaloid from the epigeal parts of *Aconitum kirinense*. Amorph. powder.O⁶,O¹⁴,O¹⁶-Tri-Me, N-Et, 4-Ac: **Akiramine**

[270073-83-1]

C₂₅H₃₉NO₇ 465.586Alkaloid from the epigeal part of *Aconitum kirinense*. Cryst. (Me₂CO). Mp 162-164°.O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, N-Et: **Akiranine**

[213478-70-7]


C₂₄H₃₉NO₆ 437.575Alkaloid from *Aconitum kirinense*. Amorph. Mp 102-104°.O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, N-Et, 4-Ac: **Akirane**

[171119-09-8]

C₂₆H₄₁NO₇ 479.612Alkaloid from aerial parts of *Aconitum kirinense*. Cryst. (Me₂CO). Mp 214-217°.Sultankhodzhaev, M.N. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1994, **30**, 603; 1997, **33**, 700-701 (*Akirane*, *Akiranine*)Feng, F. et al., *CA*, 1995, **122**, 128605f (*Kirimine*)Teshbaeva, U.T. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1999, **35**, 445-447; 659-660 (*Akiramidine*, *Akiramine*)

Aconitane-1,4,7,8,14,16-hexol

A-90


C₁₈H₂₇NO₆ 353.414(1 α ,5 β ,14 α ,16 β)-formO¹,O¹⁴,O¹⁶-Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **Isolappaconitine**

[114216-94-3]

C₃₂H₄₄N₂O₈ 584.708Alkaloid from the roots of *Aconitum finetianum* (Ranunculaceae) and *Aconitum rubicundum*. Cryst (Et₂O). Mp 186-188°.Jiang, S. et al., *Huaxue Xuebao*, 1988, **46**, 26; *CA*, **108**, 183673n (*Isolappaconitine*)Nishanov, A.A. et al., *Khim. Prir. Soedin. (Engl. Transl.)*, 352, **1991**, 27-349 (*Isolappaconitine*, *isol*, *pmr*, *ms*)

Aconitane-1,4,8,9,14,16-hexol

A-91

C₁₈H₂₇NO₆ 353.414(1 α ,5 β ,14 α ,16 β)-formO¹⁴,O¹⁶-Di-Me, N-Et: **Lappaconidine***Lappaconidine*

[31000-13-2]

C₂₂H₃₅NO₆ 409.522Alkaloid from the roots of *Aconitum leucostomum* (*Aconitum excelsum*) (Ranunculaceae). Mp 206-207°. [α]_D +12.9 (CHCl₃). Forms benzene solvate, Mp 118-120°.O¹⁴,O¹⁶-Di-Me, N-Et, tetra-Ac: Mp 195-197°.O¹⁴,O¹⁶-Di-Me, N-Et, 4-O-(2-aminobenzoyl): **4-Anthranoyllappaconidine**

[139441-64-8]

C₂₉H₄₀N₂O₇ 528.644Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Amorph. [α]_D +48.8 (c, 0.16 in CHCl₃).O¹⁴,O¹⁶-Di-Me, N-Et, 4-O-(2-acetamidobenzoyl): **1-De-O-methylappaconitine***Sinomontanine B*

[139085-68-0]

C₃₁H₄₂N₂O₈ 570.681Alkaloid from *Aconitum orientale* and *Aconitum sinomontanum*. Powder. [α]_D +52 (c, 0.1 in CHCl₃).O¹,O¹⁴,O¹⁶-Tri-Me, 4-O-(2-acetamidobenzoyl): **Sinomontanine A**C₃₀H₄₀N₂O₈ 556.655Alkaloid from the roots of *Aconitum sinomontanum*. Amorph. powder. [α]_D +31.2 (c, 0.5 in CHCl₃).O¹,O¹⁴,O¹⁶-Tri-Me, N-Ac, 4-O-(2-acetamidobenzoyl): **Sinaconitine B**C₃₂H₄₂N₂O₉ 598.692Alkaloid from the roots of *Aconitum sinomontanum*. Amorph. powder. [α]_D +17 (c, 0.35 in CHCl₃).O¹,O¹⁴,O¹⁶-Tri-Me, N-Et: **Lappaconine**

[23943-93-3]

C₂₃H₃₇NO₆ 423.548Detected in *Aconitum leucostomum* (Ranunculaceae). Cryst. + 1½ H₂O (H₂O). Mp 96°. [α]_D²⁵ +27 (c, 3.17 in CHCl₃).

▶AR5569540

O¹,O¹⁴,O¹⁶-Tri-Me, N-Et, 4-O-(2-aminobenzoyl): **Puberanidine**. N-Deacetyl-lappaconitine

[11033-64-0]

C₃₀H₄₂N₂O₇ 542.671Alkaloid from *Aconitum ranunculaefolium* and from the roots of *Aconitum barbatum* var. *puberulum*, *Aconitum septentrionale*, *Aconitum finetianum* (Ranunculaceae) and *Delphinium cashmirianum* (Ranunculaceae). Prisms (MeOH), needles (Me₂CO/hexane), or amorph. Mp 213-214.5° (120-121°, 209-214°). [α]_D²⁰ +23.2 (c, 0.9 in CHCl₃) (for amorph. product). [α]_D²⁷ +42 (c, 0.34 in EtOH).

▶OE5784120

O¹,O¹⁴,O¹⁶-Tri-Me, N-Et, 4-O-(2-acetamidobenzoyl): **Lappaconitine**

[32854-75-4]


C₃₂H₄₄N₂O₈ 584.708Alkaloid from *Aconitum septentrionale*, *Aconitum orientale*, *Aconitum excelsum*, *Aconitum sinomontanum*, *Aconitum finetianum*, *Aconitum ranunculaefolium*, *Aconitum barbatum* var. *puberulum* and *Delphinium cashmirianum* (Ranunculaceae). Cryst. (Et₂O). Mp 227° (219-221°). [α]_D²⁵ +28.2 (c, 2.07 in CHCl₃).

▶OE5784100

O¹,O¹⁴,O¹⁶-Tri-Me, N-Et, 4-O-(2-methylaminobenzoyl): **Septefine**. *Septephine*C₃₁H₄₄N₂O₇ 556.698Alkaloid from *Aconitum septentrionale* (Ranunculaceae). Cryst. (MeOH). Mp 194-195°.Mollov, N. et al., *Dokl. Bulg. Akad. Nauk*, 1964, **17**, 251; *CA*, **61**, 12324h (*Puberanidine*, *isol*, *struct*)Khamova, M. et al., *Tet. Lett.*, 1964, 2711 (*Lappaconidine*, *Lappaconitine*, *ir*, *pmr*, *struct*)Marion, L. et al., *Can. J. Chem.*, 1967, **45**, 969 (*Lappaconidine*, *Lappaconitine*, *isol*, *ir*, *pmr*, *struct*)Mollov, N. et al., *Tet. Lett.*, 1969, 2189 (*Lappaconidine*, *pmr*, *struct*)Birnbau, G.I. et al., *Acta Cryst. B*, 1970, **26**, 755 (*Lappaconine*, *cryst struct*)Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1970, **6**, 583; 1971, **7**, 622; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 598; 1971, **7**, 601 (*Lappaconidine*, *Lappaconitine*)Pelletier, S.W. et al., *Can. J. Chem.*, 1979, **57**, 1652 (*cmr*)Shamma, M. et al., *J. Nat. Prod.*, 1979, **42**, 615 (*Puberanidine*, *isol*, *ir*, *pmr*, *struct*)Plugar, V.N. et al., *Khim. Prir. Soedin.*, 1982, **18**, 80; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 75 (*Lappaconidine*, *glc*, *ms*)Yu, D. et al., *Planta Med.*, 1983, **49**, 85 (*Puberanidine*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)Sayed, H.M. et al., *J. Nat. Prod.*, 1992, **55**, 1595 (*4-Anthranoyllappaconidine*)Usmanova, S.K. et al., *Khim. Prir. Soedin. (Engl. Transl.)*, 1996, **32**, 225-228; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 198-200 (*Septefine*)

- Ulubelen, A. *et al.*, *Phytochemistry*, 1996, **41**, 957 (*1*-Demethylappaconitine)
 Wang, F.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 15-22 (*Sinomontanines*)
 Tan, J.J. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 535-539 (*Sinaconitine B*)
 Wang, Y.-P. *et al.*, *Acta Cryst. E*, 2007, **63**, o1645-o1647 (*Lappaconitine, crystal*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LBD000

Aconitane-1,6,7,8,14,16-hexol A-92




$C_{18}H_{27}NO_6$ 353.414

(1 α ,5 β ,6 β ,14 α ,16 β)-form

- O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*: *Umbrophine*.
Umbrophine. Acosepticine
 [144049-71-8]
 $C_{23}H_{37}NO_6$ 423.548
 Alkaloid from roots of *Aconitum septentrionale* and *Aconitum umbrosum* (Ranunculaceae). Amorph. solid. $[\alpha]_D^{25} +23.4$ (c, 0.385 in $CHCl_3$). Identity with Umbrophine (cryst., Mp. 110-112°, no opt. rotn. reported) is tentative.
 O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*, 6-*Ac*: **6-O-Acetylacosepticine**. 6-*Acetylu*mbrophine
 [144074-85-1]
 $C_{25}H_{39}NO_7$ 465.586
 Alkaloid from roots of *Aconitum septentrionale* and *Aconitum umbrosum* (Ranunculaceae). Cryst. (Et_2O). Mp 168.5-170.5°. $[\alpha]_D -1.2$ (c, 0.2 in $CHCl_3$).
 O^1, O^6, O^{14}, O^{16} -*Tetra-Me*, N-*Et*: **Exceconidine**
 $C_{24}H_{39}NO_6$ 437.575
 Alkaloid from *Aconitum excelsum*. Powder.
 O^7, O^8 -*Methylene*, O^1, O^{14}, O^{16} -*tri-Me*, N-*Et*: **Anthriscifolcine B**
 [948308-29-0]
 $C_{24}H_{37}NO_6$ 435.559
 Alkaloid from *Delphinium anthriscifolium* var. *savatteri*. Amorph. powder. Mp 75-77°. $[\alpha]_D^{20} -27$ (c, 0.5 in $CHCl_3$).
 O^7, O^8 -*Methylene*, O^1, O^{14}, O^{16} -*tri-Me*, N-*Et*, 6-*Ac*: **Anthriscifolcine A**
 [948308-28-9]
 $C_{26}H_{39}NO_7$ 477.597
 Alkaloid from *Delphinium anthriscifolium* var. *savatteri*. Amorph. powder. Mp 135-137°. $[\alpha]_D^{20} -12.2$ (c, 0.5 in $CHCl_3$).
 6-*Ketone*, O^1, O^{14}, O^{16} -*tri-Me*, N-*Et*: **Leucostine**
 [192386-08-6]
 $C_{23}H_{35}NO_6$ 421.533

- Alkaloid from *Aconitum leucostomum*.
 $[\alpha]_D^{25} +19.4$ (c, 0.05 in $EtOH$). Not to be confused with Leucostine A in M-309.
 Sayed, H.M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1595 (*Acosepticine*)
 Ross, S.A. *et al.*, *Tetrahedron*, 1992, **48**, 1183 (*6-O-Acetylacosepticine*)
 Tel'nov, V.A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 73-77; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 60-63 (*Umbrophine, 6-Acetylu*mbrophine)
 Wei, X. *et al.*, *Zhivui Xuebao (Acta Bot. Sin.)*, 1996, **38**, 995-997; *CA*, **127**, 147062w (*Leucostine*)
 Zhang, S.X. *et al.*, *Chin. Chem. Lett.*, 1999, **10**, 133-134 (*Exceconidine*)
 Zinurova, E.G. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 387-389 (*Leucostine*)
 Song, L. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 918-921 (*Anthriscifolcines A,B*)

Aconitane-1,3,4,7,8,9,14,16-octol A-93




$C_{18}H_{27}NO_8$ 385.413

(1 α ,3 α ,14 α ,16 β)-form

- O^{14}, O^{16} -*Di-Me*, N-*Et*: **Sinomontanine D**
 $C_{22}H_{35}NO_8$ 441.52
 Alkaloid from the roots of *Aconitum sinomontanum*. $[\alpha]_D^{17} +26.6$ (c, 0.5 in $CHCl_3$).
 Peng, C. *et al.*, *Yaouxue Xuebao*, 2000, **35**, 201-203

Aconitane-1,4,7,8,9,10,14,16-octol A-94




$C_{18}H_{27}NO_8$ 385.413

(1 α ,5 β ,14 α ,16 β)-form

- O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*, 4-*O*-(2-aminobenzoyl): **N-Deacetylfinaconitine**
 [82872-81-9]
 $C_{30}H_{42}N_2O_9$ 574.67
 Alkaloid from the roots of *Aconitum finetianum* (Ranunculaceae). Mp 121-123°. O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*, 4-*O*-(2-acetamidobenzoyl): **Finaconitine**. 10-*Hydroxy*-ranaconitine
 [81161-27-5]

- $C_{32}H_{44}N_2O_{10}$ 616.707
 Alkaloid from the roots of *Aconitum finetianum* (Ranunculaceae). Mp 220-221°.
 Jiang, S. *et al.*, *Yaouxue Xuebao*, 1982, **17**, 283; 1983, **18**, 440; *CA*, **97**, 20736a; **100**, 20505e (*Finaconitine, Deacetylfinaconitine*)

Aconitane-1,4,8,14,16-pentol A-95




$C_{18}H_{27}NO_5$ 337.415

(1 α ,5 β ,14 α ,16 β)-form

- O^{14}, O^{16} -*Di-Me*, N-*Et*: **Dihydromonticamine**
 [81037-23-2]
 $C_{22}H_{35}NO_5$ 393.522
 Alkaloid from the aerial parts of *Aconitum monticola* (Ranunculaceae). Cryst. (Et_2O/Me_2CO). Mp 156-157°. O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*, 4-*O*-(2-aminobenzoyl): **Neofinaconitine**. *Delphicrispuline*
 [114244-02-9]
 $C_{30}H_{42}N_2O_6$ 526.672
 Alkaloid from the roots of *Aconitum finetianum* and *Delphinium crispulum* (Ranunculaceae). $[\alpha]_D^{20} +23.8$ (c, 0.8 in $CHCl_3$) (*Delphicrispuline*). O^1, O^{14}, O^{16} -*Tri-Me*, N-*Et*, 4-*O*-(2-acetamidobenzoyl): **9-Deoxylappaconitine**
 [114216-95-4]
 $C_{32}H_{44}N_2O_7$ 568.709
 Alkaloid from roots of *Aconitum finetianum* (Ranunculaceae).
 Ametova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1982, **504**; *Chem. Nat. Compd. (Engl. Transl.)*, **472** (*Dihydromonticamine*)
 Jiang, S. *et al.*, *Huaxue Xuebao*, 1988, **46**, 26; *CA*, **108**, 183673n (*Neofinaconitine, 9-Deoxylappaconitine*)
 Ulubelen, A. *et al.*, *Phytochemistry*, 1999, **50**, 513-516 (*Delphicrispuline*)

Aconitane-1,6,7,14,16-pentol A-96



$C_{18}H_{27}NO_5$ 337.415


(1 α ,5 β ,14 α ,16 β)-form

- 6-*Ketone*, O^1, O^{14}, O^{16} -*tri-Me*, N-*Et*: **Leucconine**
 [192386-06-4]
 $C_{23}H_{35}NO_5$ 405.533

Alkaloid from *Aconitum leucostomum*. Cryst. (Me₂CO/hexane). Mp 195-197°.

Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1992, **28**, 538-540; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 470-471 (isol. ms, cmr)
Wei, X. et al., *Zhivuo Xuebao (Acta Bot. Sin.)*, 1996, **38**, 995-997; *CA*, **127**, 147062w (isol)

Aconitane-1,8,9,14,16-pentol A-97



C₁₈H₂₇NO₅ 337.415

(1α,5β,14α,16β)-form

O¹,O¹⁶-Di-Me, N-Et: **Episcopalisinine** [83692-39-1]

C₂₂H₃₅NO₅ 393.522

Alkaloid from the roots of *Aconitum episcopale* (Ranunculaceae).


O¹,O¹⁶-Di-Me, N-Et, 14-benzoyl: **Episcopalisine. 14-Benzoylepiscopalisine** [83685-22-7]

C₂₉H₃₉NO₆ 497.63

Alkaloid from the roots of *Aconitum episcopale* (Ranunculaceae).

Wang, F. et al., *Yaoxue Xuebao*, 1983, **18**, 514-521; *CA*, **100**, 64981m (*Episcopalisinine*, *Episcopalisine*)

Aconitane-1,8,13,14,16-pentol A-98



C₁₈H₂₇NO₅ 337.415

(1α,5β,14α,16β)-form

O¹,O¹⁶-Di-Me, N-Et, 14-Ac: **Delavaconitine F** [1022894-28-5]

C₂₄H₃₇NO₆ 435.559

Alkaloid from the roots of *Aconitum delavayi*. Amorph. powder.

O¹,O¹⁶-Di-Me, N-Et, 14-benzoyl: **Delavaconitine** [1356-52-1]

C₂₉H₃₉NO₆ 497.63

Alkaloid from the roots of the Chinese drug Tzu-Tsao-Wu (*Aconitum delavayi*). Anaesthetic. Mp 241° (as perchlorate). Log P 0.24 (uncertain value) (calc).

▶ AR5566000

O¹,O¹⁶-Di-Me, N-Et, 14-benzoyl, 13-Ac: **Delavaconitine E**

[126262-74-6]

C₃₁H₄₁NO₇ 539.667

Alkaloid from *Aconitum delavayi* (Ranunculaceae).

O¹,O¹⁶-Di-Me, N-Et, 14-O-(4-methoxybenzoyl): **Contortumine**

[131653-96-8]

C₃₀H₄₁NO₇ 527.656

Alkaloid from the roots of *Aconitum contortum* (Ranunculaceae). Amorph. [α]_D²⁰ -4.8 (c, 0.97 in EtOH).

Chu, J.-H. et al., *Huaxue Xuebao*, 1959, **25**, 321; *CA*, **54**, 17440i (*Delavaconitine*)


Zhu, Y. et al., *Heterocycles*, 1982, **17**, 607-644 (*Delavaconitine*)

Jiang, S. et al., *Huaxue Xuebao*, 1989, **47**, 1178; *CA*, **112**, 175625r (*Delvaconitine E*)

Niitsu, K. et al., *Heterocycles*, 1990, **31**, 1517-1524 (*Contortumine*)

Jiang, S.H. et al., *Chin. Chem. Lett.*, 2007, **18**, 409-411 (*Delavaconitines E,F*)

Aconitane-1,8,14,16-tetrol A-99



C₁₈H₂₇NO₄ 321.416

(1α,5β,14α,16β)-form

O¹⁶-Me, N-Et: **Scopaline**

[80787-49-1]

C₂₁H₃₃NO₄ 363.496

Alkaloid from *Aconitum episcopale* (Ranunculaceae).

O¹,O¹⁶-Di-Me, N-Et: **Aconosine**

[38839-95-1]

C₂₂H₃₅NO₄ 377.523

Alkaloid from *Aconitum nasutum* (all parts of the plant), the aerial parts and roots of *Aconitum fischeri*, the foliage and roots of *Aconitum arcuatum*, and from the roots of *Aconitum forrestii*, *Aconitum napellus* and *Aconitum stapfianum* var. *pubipes* (Ranunculaceae). Cryst. (MeOH aq.). Mp 150° (148°). [α]_D²⁰ -25 (c, 0.57 in MeOH). [α]_D²⁰ -21 (c, 1 in MeOH).

▶ AR5540000

O¹,O¹⁶-Di-Me, N-Et, 14-Ac: **Episcopalitine. Dolaconine**

[82841-75-6]

C₂₄H₃₇NO₅ 419.56

Alkaloid from the roots of *Aconitum episcopale* and *Aconitum stapfianum* var. *pubipes* (Ranunculaceae). Mp 44-46°.

O¹,O¹⁶-Di-Me, N-Et, 8,14-di-Ac: **8-Acetyldolaconine. 8,14-Diacetylaconosine**

[132160-38-4]

C₂₆H₃₉NO₆ 461.597

Alkaloid from the roots of *Aconitum campylorrhynchum* (Ranunculaceae).

O¹,O¹⁶-Di-Me, N-Et, 14-benzoyl: **Delavaconitine C**

[126234-20-6]

C₂₉H₃₉NO₅ 481.631

Alkaloid from *Aconitum delavayi* (Ra-

nunculaceae).

O¹,O¹⁶-Di-Me, N-Et, 14-benzoyl, 8-Ac:

Delavaconitine D

[126234-21-7]

C₃₁H₄₁NO₆ 523.668

Alkaloid from *Aconitum delavayi* (Ranunculaceae).

Murav'eva, D.A. et al., *Khim. Prir. Soedin.*, 1972, **8**, 128-129; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 132-133 (*Aconosine*, *isol*, *ir*, *pmr*, *ms*, *struct*)

Yang, C. et al., *Huaxue Xuebao*, 1981, **39**, 445-452; *CA*, **96**, 82689a (*Scopaline*)


Luo, S. et al., *Huaxue Xuebao*, 1981, **39**, 808-810; *CA*, **97**, 107079n (*Aconosine*, *Dolaconine*)

Edwards, O.E. et al., *Can. J. Chem.*, 1983, **61**, 1194-1196 (*synth*, *ir*, *pmr*, *cmr*, *ms*)

Wang, F. et al., *Yaoxue Xuebao*, 1983, **18**, 514-521; *CA*, **100**, 64981m (*Episcopalitine*)

Ding, L.S. et al., *Yaoxue Xuebao*, 1990, **25**, 441-444; *CA*, **114**, 98113 (8-Acetyldelaconine)

Aconitane-1,14,16-triol A-100



C₁₈H₂₇NO₃ 305.416

(1α,5β,14α,16β)-form

8-O-[2-(4-Hydroxyphenyl)ethoxy],

O¹,O¹⁶-di-Me, N-Et: **Piepunendine B**

[947321-59-7]

C₃₀H₄₃NO₅ 497.673

Alkaloid from the roots of *Aconitum piepunense*. Mp 85-87°. [α]_D²⁰ -10.6 (c, 0.5 in CHCl₃).

14-Ketone, O¹,O¹⁶-di-Me, N-Et: **8-Deoxy-14-dehydroaconosine**


C₂₂H₃₃NO₃ 359.508

Alkaloid from *Aconitum stapfianum* (Ranunculaceae).

Liang, X.-T. et al., *Pure Appl. Chem.*, 1986, **58**, 711-718 (*rev*)

Cai, L. et al., *Nat. Prod. Commun.*, 2006, **1**, 191-194 (*Piepunendine B*)

Aconit-19(N)-ene-1,6,7,8,14,16-hexol A-101



C₁₈H₂₅NO₆ 351.399

(1α,5β,14α,16β)-form

O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me: **Lamarckinine**†

[150375-24-9]


C₂₂H₃₃NO₆ 407.506

Alkaloid from aerial parts of *Aconitum lamarekii* (Ranunculaceae). Resin. $[\alpha]_D^{25} +76.6$ (c, 1.1 in CHCl_3).

de la Fuente, G. *et al.*, *Heterocycles*, 1993, **36**, 1455 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Aconit-19(N)-ene-1,8,13,14,16-pentol

A-102



$\text{C}_{18}\text{H}_{25}\text{NO}_5$ 335.399

(1 α ,5 β ,14 α ,16 β)-form

*O*¹,*O*¹⁶-Di-Me, 14-benzoyl, 8,13-di-Ac:

Delavaconitine G

[1022894-29-6]

$\text{C}_{31}\text{H}_{37}\text{NO}_8$ 551.635


Alkaloid from the roots of *Aconitum delavayi*. Cryst. Mp 232-233°.

Jiang, S.H. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 409-411 (*isol*, *pmr*, *cmr*)

Aconit-19(N)-ene-1,8,14,16-tetrol

A-103

19,20-Didehydroaconitane-1,8,14,16-tetrol



$\text{C}_{18}\text{H}_{25}\text{NO}_4$ 319.4

(1 α ,5 β ,14 α ,16 β)-form

*O*¹,*O*¹⁶-Di-Me: **Liconosine A**

[126621-43-0]

$\text{C}_{20}\text{H}_{29}\text{NO}_4$ 347.453


Alkaloid from *Aconitum forrestii* (Ranunculaceae).

Chen, S. *et al.*, *Yunnan Zhiwu Yanjiu*, 1989, **11**, 267; *CA*, **112**, 175578

Aconitine

A-104

[302-27-2]



Absolute Configuration

$\text{C}_{34}\text{H}_{47}\text{NO}_{11}$ 645.745

Alkaloid from *Aconitum napellus*, *Aconi-*

tum fauriei, *Aconitum grossedentatum*, *Aconitum hakusanense*, *Aconitum mokchangense*, *Aconitum zuccarini*, *Aconitum bullatifolium* var. *homotrichum*, *Aconitum carmichaeli*, *Aconitum karakolicum* and many others (Ranunculaceae). Toxic agent implicated in poisoning by *Aconitum* spp., esp. *Aconitum chasmanthum* in India, exp. arrhythmia effects. Gastric anaesthetic. Homeopathic remedy against neuralgia and febrile diseases. Cryst. V. spar. sol. H_2O . Mp 204°. $[\alpha]_D^{25} +19$ (CHCl_3). pK_a 8.35 (15°). Hydrol. to 4-Methylaconitane-1,3,6,8,13,14,15,16,18-nonol, M-331.

► Very toxic if swallowed or by skin absorption. Human systemic effects by ingestion. LD_{50} (mus, orl) 1 mg/kg, LD_{50} (mus, ivn) 0.175 mg/kg, exp. lethal doses by subcutaneous route reported; LD_{50} (mus, ipr) 0.328 mg/kg. AR5960000

Hydrochloride: Mp 170-172° dec. $[\alpha]_D^{25} -31$ (H_2O).

3-Ac: Flaconitine. 3-Acetylaconitine

[77181-26-1]

$\text{C}_{36}\text{H}_{49}\text{NO}_{12}$ 687.783

Alkaloid from the roots of *Aconitum flavum* and *Aconitum pendulum* (Ranunculaceae). Mp 196-197°. $[\alpha]_D^{24} +18.6$ (c, 1 in CHCl_3).

► AR5562000

Przybylska, M. *et al.*, *Can. J. Chem.*, 1959, **37**, 1116-1118; 1843-1845 (*struct*)

Wiesner, K. *et al.*, *Tet. Lett.*, 1959, **2**, 15-24 (*struct*)

Bachelor, F.W. *et al.*, *Tet. Lett.*, 1960, **31**, 1-9 (*struct*)

Birnbaum, K.B. *et al.*, *Tet. Lett.*, 1971, 867-870 (*config*, *struct*)

Chang, X. *et al.*, *Yaoxue Xuebao*, 1981, **16**, 474-476; *CA*, **97**, 3590f (*Flaconitine*, *isol*)

Wang, H. *et al.*, *Heterocycles*, 1985, **23**, 803-807 (*Flaconitine*, *cmr*)

Seeger, R. *et al.*, *Dtsch. Apoth. -Ztg.*, 1994, **134**, 43-50; 53-54; *CA*, **121**, 308035 (*pharmacol*, *rev*)

Chan, T.Y. *et al.*, *Vet. Hum. Toxicol.*, 1994, **36**, 326-328 (*pharmacol*, *rev*)

Ho, K. *et al.*, *Planta Med.*, 1997, **63**, 75-79 (*ms*)

Ohta, H. *et al.*, *J. Chromatogr. B*, 1998, **714**, 215-221 (*hplc*, *ms*)

Chen, Y. *et al.*, *J. Nat. Prod.*, 1999, **62**, 701-704 (*isol*, *ms*)

Wang, Z. *et al.*, *Fenxi Huaxue*, 2001, **29**, 391-395; *CA*, **135**, 127303 (*ms*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, ADH750

Acophine

A-105

Acofine

[169626-13-5]



$\text{C}_{25}\text{H}_{38}\text{ClNO}_3$ 436.033


Alkaloid from roots of *Aconitum karakolicum*. Cryst. (Me_2CO). Mp 159-160°. Poss. artifact.

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 267; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 222

Acoseptine

A-106

[257860-39-2]



$\text{C}_{32}\text{H}_{44}\text{N}_2\text{O}_7$ 568.709

Unusual (7 → 8)abeoaconitane skeleton.

Alkaloid from the roots of *Aconitum septentrionale*. Cryst. (MeOH). Mp 127-128°.

N-Succinimide: Anhydrolycaconitine

[321554-91-0]

$\text{C}_{36}\text{H}_{46}\text{N}_2\text{O}_9$ 650.767

Alkaloid from the roots of *Aconitum septentrionale*. Cryst. (Me_2CO). Mp 215-216°. $[\alpha]_D^{20} +21$ (c, 0.1 in CHCl_3). Misleading name.

Usmanova, S.K. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1999, **35**, 91-93 (*Acoseptine*)


Yunusov, M.S. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2000, **49**, 1629-1633 (*Anhydrolycaconitine*)

Khairitdinova, E.D. *et al.*, *Russ. J. Org. Chem. (Engl. Transl.)*, 2008, **44**, 536-541 (*Anhydrolycaconitine*)

Acosmine

A-107

[250343-47-6]



Absolute Configuration

$\text{C}_{21}\text{H}_{33}\text{N}_3\text{O}_2$ 359.511

Alkaloid from *Acosmium panamense* and *Bowdichia virgilioides*. Amorph. yellow solid. $[\alpha]_D^{25} -10.8$ (c, 1.3 in CHCl_3). λ_{max} 242 ; 290 (sh) (MeOH).

O-Ac: Acosmine acetate. O-Acetylacosmine

[250279-26-6]

$\text{C}_{23}\text{H}_{35}\text{N}_3\text{O}_3$ 401.548

Alkaloid from the seeds of *Acosmium panamense*. Amorph. yellow solid. $[\alpha]_D^{25} -15.7$ (c, 0.86 in CHCl_3). λ_{max} 242 ; 290 (sh) (MeOH).

O-(3,4,5-Trimethoxybenzoyl): Bowdichine

[730992-48-0]

$\text{C}_{31}\text{H}_{43}\text{N}_3\text{O}_6$ 553.697

Alkaloid from the stem bark of *Bowdichia virgilioides*. Viscous oil. λ_{\max} 225 ; 250 ; 298 (MeOH).

Nuzillard, J.-M. *et al.*, *Tetrahedron*, 1999, **55**,


11511-11518 (*Acosmine*)

Barbosa-Filho, J.M. *et al.*, *J. Asian Nat.*

Prod. Res., 2004, **6**, 11-17 (*Bowdichine*, *pmr*, *cmr*)

Acozerine**A-108**

[155210-48-3]




$C_{31}H_{42}N_2O_3$ 490.684

Alkaloid from *Aconitum zeravshanicum* (Ranunculaceae).

Vaisov, Z.M. *et al.*, *Mendelevy Commun.*, 1993, 237; *CA*, **120**, 299052m (*isol*, *ir*, *pmr*, *ms*)

Acremoauxin A**A-109**

2-(3-Indolyl)propanoyl-D-arabinitol
[125537-93-1]



$C_{16}H_{21}NO_6$ 323.345

Prod. by *Acronium roseum*. Plant growth regulator. Needles (EtOH/EtOAc). Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 142-144° (135-136°). $[\alpha]_D^{19} +53.6$ (c, 0.35 in MeOH). λ_{\max} 219 (€ 25000); 225 (€ 9600); 278 (€ 7000); 286 (€ 5700); 288 (€ 6000) (MeOH) (Berdy).

Sassa, T. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 3105 (*isol*, *struct*)

Yoshida, N. *et al.*, *Agric. Biol. Chem.*, 1990, **54**, 2681 (*synth*)

Acremoauxin B**A-110**

3-Indolylacetyl-D-arabinitol, 9CI
[138111-07-6]

As Acremoauxin A, A-109 with R = H


$C_{15}H_{19}NO_6$ 309.318

Prod. by *Acronium roseum*. Plant growth regulator. Cryst.

Japan. Pat., 1991, 91 83 968; *CA*, **116**, 39794a

Acremolactone B**A-111**

[850723-60-3]



Absolute Configuration


$C_{26}H_{31}NO_5$ 437.535

Prod. by *Acronium roseum* 14267. Phytotoxin. Needles (Me₂CO), Mp 255-256°. $[\alpha]_D^{24} +37$ (c, 0.08 in Me₂CO). λ_{\max} 212 (€ 26700); 240 (sh) (€ 8600); 283 (€ 15400); 330 (€ 1000) (MeOH).

Sassa, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2004, **68**, 2633-2636 (*isol*, *pmr*, *cmr*)

Acridone**A-112**

9(10H)-Acridinone, 9CI. 9-Hydroxyacridine. 9-Acridinol. Acridol
[578-95-0]



$C_{13}H_9NO$ 195.22

NH-form predominates. Constit. of the roots and rhizomes of *Thamnosma montana* (Rutaceae). Antineoplastic agent. Yellow leaflets (EtOH). Insol. Et₂O, C₆H₆, CHCl₃. Mp 356-358° Mp 362-365° dec. pK_{a1} -0.32 (20°, H₂SO₄ aq.).

▶AR6976000

NH-form

N-Benzoyl: 10-Benzoyl-9(10H)-acridinone, 9CI
[57479-66-0]

$C_{20}H_{13}NO_2$ 299.328

Yellow plates (hexane or by subl.). Mp 198-200°.

N-Me: see 10-Methylacridone, M-382

N-Et: 10-Ethylacridone

[2207-41-2]

$C_{15}H_{13}NO$ 223.274

Mp 159°.

OH-form [643-62-9]

Me ether: 9-Methoxyacridine

[10228-90-7]

$C_{14}H_{11}NO$ 209.247

Yellow needles (petrol). Mp 65°. pK_a 7 (20°). Forms monohydrate with Mp 103°.

Et ether: 9-Ethoxyacridine

$C_{15}H_{13}NO$ 223.274

Anal. reagent for 1-Hydrazinophthalazine and 4-Amino-3-isoxazolidinone, A-832. Mp ca. 83° Mp 92°.

Ph ether: 9-Phenoxyacridine

[2148-14-3]

$C_{19}H_{13}NO$ 271.318

Cryst. (C₆H₆). Mp 127-128°.

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **2**, 922B (*nmr*)

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 904B (*ir*)

Kliegl, A. *et al.*, *Ber.*, 1914, **47**, 1629-1640 (*synth*)

Lehmstedt, K. *et al.*, *Ber.*, 1935, **68**, 1455-1464 (*synth*, *Me ether*)

Org. Synth., Coll. Vol., **2**, 1943, 15 (*synth*)

Dupré, D.J. *et al.*, *J.C.S.*, 1945, 549-551 (*Ph ether*, *synth*)

Kitani, K. *et al.*, *Nippon Kagaku Kaishi*, 1954, **75**, 396-398; *CA*, **49**, 10297 (*Me ether*, *Et ether*, *synth*)

Albert, A. *et al.*, *The Acridines*, 2nd edn., E. Arnold, London, 1966, (*rev*)

Altiparmakian, R.H. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 85 (*pmr*)

Gagan, J.M.F. *et al.*, *Acridines*, (Ed. Acheson, R.M.), 2nd edn., Interscience, N.Y., 1973, 141 (*rev*)

Chen, G.J. *et al.*, *J.C.S. Perkin 1*, 1975, 1138 (*synth*, *N-benzoyl*)

Chang, P.T.O. *et al.*, *J. Nat. Prod.*, 1976, **39**, 134 (*isol*, *uv*, *ir*, *pmr*, *ms*)

Beak, P. *et al.*, *J.A.C.S.*, 1976, **98**, 171 (*uv*)

Adams, J.H. *et al.*, *J.C.S. Perkin 1*, 1977, 2173 (*synth*)

Faure, R. *et al.*, *Spectrosc. Lett.*, 1981, **14**, 223 (*cmr*)

Stewart, J.T. *et al.*, *Int. J. Pharm.*, 1983, **17**, 161-166 (*Me ether*, *synth*, *use*)

Stewart, J.T. *et al.*, *J. Pharm. Sci.*, 1988, **77**, 452-454 (*Me ether*, *synth*, *use*)

Potts, G.D. *et al.*, *Acta Cryst. C*, 1995, **51**, 267 (*cryst struct*)


Phanstiel, O. *et al.*, *J.O.C.*, 2000, **65**, 5590-5599 (*Ph ether*, *synth*, *pmr*)

Dadabhoy, A. *et al.*, *J.C.S. Perkin 2*, 2002, 348-357 (*N-Et*, *synth*, *ir*, *uv*, *pmr*, *cmr*)

Ebead, Y. *et al.*, *Acta Cryst. C*, 2005, **61**, o85-o87 (*Ph ether*, *cryst struct*)

Acriginine A**A-113**

[149155-25-9]




$C_{30}H_{29}NO_9$ 547.56

Acridone-lignan hybrid. Alkaloid from roots of *Citrus grandis* f. *Hirado* and *Citrus yuko* (Rutaceae). Yellow prisms. Mp 177-179°. Racemic.

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 406 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cryst struct*)

Acrimarine A**A-114**

[119152-47-5]



$C_{31}H_{29}NO_8$ 543.572

Alkaloid from the roots of *Citrus funadoko* (Rutaceae). Pale yellow oil. $[\alpha]_D -9.76$ (c, 0.082 in CHCl₃).

3-Me ether: *Acrimarine N*

$C_{32}H_{31}NO_8$ 557.599

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) x Dancy tangerine (*Citrus tangerina*)].

Yellow oil. Opt. inactive.

3-Me ether, N-de-Me: Acrimarine B

[119152-48-6]

C₃₁H₂₉NO₈ 543.572

Alkaloid from the roots of *Citrus funadoko* (Rutaceae). Yellow prisms (Me₂CO). Mp 288-290°. [α]_D²⁵ -7.14 (c, 0.056 in CHCl₃).

O⁶-De-Me: Acrimarine K

[147395-91-3]

C₃₀H₂₇NO₈ 529.545

Alkaloid from roots of *Citrus funadoko* (Rutaceae). Yellow oil. [α]_D²⁵ 0 (c, 0.0695 in CHCl₃).

O⁶-De-Me, 3-Me ether: Acrimarine F

[129722-89-0]

C₃₁H₂₉NO₈ 543.572

Alkaloid from the roots of *Citrus funadoko* (Rutaceae). Pale yellow powder.

O⁶-De-Me, 3-Me ether, N⁶-de-Me: Acrimarine E

[129722-88-9]

C₃₀H₂₇NO₈ 529.545

Alkaloid from the roots of *Citrus funadoko* (Rutaceae). Pale yellow prisms (Me₂CO). Mp 274-276°. [α]_D²⁵ +20.1 (c, 0.05 in Me₂CO).

6-De-methoxy, O⁵-de-Me: Acrimarine G

[129722-97-0]

C₂₉H₂₅NO₇ 499.519

Alkaloid from the roots of *Citrus funadoko* (Rutaceae). Yellow oil. [α]_D²⁵ +8 (c, 0.075 in CHCl₃).

6-De-methoxy, O⁵-de-Me, 3-Me ether: Acrimarine H

[132185-44-5]

C₃₀H₂₇NO₇ 513.546

Alkaloid from roots of *Citrus* spp. Yellow oil. [α]_D²⁵ 0 (Me₂CO).

Bis(demethoxy): Acrimarine M

[147513-68-6]

C₂₉H₂₅NO₆ 483.52

Alkaloid from roots of *Citrus funadoko* (Rutaceae). Yellow oil. [α]_D²⁵ 0 (c, 0.052 in CHCl₃).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 2586 (*Acrimarine H*)

Furukawa, H. *et al.*, *J.C.S. Perkin 1*, 1990, 1593 (*isol, pmr, cmr, struct*)


Takemura, Y. *et al.*, *Heterocycles*, 1992, **34**, 2363 (*Acrimarine K, Acrimarine M*)

Takemura, Y. *et al.*, *Heterocycles*, 1994, **38**, 1937 (*Acrimarine N*)

Acrimarine C

A-115

[119206-24-5]



C₃₀H₂₇NO₈ 529.545

(-)-form

Alkaloid from the roots of *Citrus funa-*

doko (Rutaceae). Yellow oil. [α]_D²⁵ -6.17 (c, 0.081 in CHCl₃).

6-Me ether: Acrimarine D

[129722-87-8]

C₃₁H₂₉NO₈ 543.572

Alkaloid from the root of *Citrus funadoko* (Rutaceae). Yellow oil. [α]_D²⁵ -3 (c, 0.103 in CHCl₃).

(±)-form

Acrimarine L

[147512-44-5]

From roots of *Citrus funadoko* (Rutaceae). Yellow oil.


Furukawa, H. *et al.*, *J.C.S. Perkin 1*, 1990, 1593 (*Acrimarine C, Acrimarine D*)

Takemura, Y. *et al.*, *Heterocycles*, 1992, **34**, 2363 (*Acrimarine L*)

Acrimarine J

A-116

[147395-90-2]



C₃₅H₃₃NO₈ 595.648

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) × Dancy tangerine (*Citrus tangerina*)] (Rutaceae). Yellow oil. [α]_D²⁸ 0 (c, 0.106 in CHCl₃).

6-Deoxy, O⁵-de-Me: Acrimarine I

[147395-89-9]

C₃₄H₃₁NO₇ 565.621

Alkaloid from roots of Yalaha (Rutaceae). Yellow oil. [α]_D²⁶ +27.8 (c, 0.036 in CHCl₃).


Takemura, Y. *et al.*, *Heterocycles*, 1992, **34**, 2363 (*isol, uv, ir, pmr, cmr, ms, struct*)

Acrodontiolamide

A-117

α-(Dichloromethoxymethyl)-*β*-hydroxy-4-nitrobenzenepropanamide

[152053-09-3]



C₁₁H₁₂Cl₂N₂O₅ 323.132

Metab. of the fungus *Acrodontium salmoneum*. Antifungal agent. Plates (MeOH). Mp 145-147°. λ_{max} 273 (ε 386000) (MeOH) (Berdy).


Buarque de Gusmao, N. *et al.*, *Spectrosc. Lett.*, 1993, **26**, 1373 (*isol, uv, ir, ms, pmr, cmr*)

Steiman, R. *et al.*, *Pharmazie*, 1995, **50**, 693 (*isol, activity*)

Acromelic acid A

A-118

5-[5-Carboxy-4-(carboxymethyl)-3-pyrrolidiny]-1,6-dihydro-6-oxo-2-pyridine-carboxylic acid, 9CI. Acromelinic acid A [86630-09-3]



C₁₃H₁₄N₂O₇ 310.263

Trace alkaloid from fruiting bodies of the poisonous mushroom *Clitocybe acromelalga*. Shows very potent neuroexcitatory action. Cryst. Sol. H₂O; poorly sol. Me₂CO, hexane. Mp 310°. [α]_D²⁵ +27.8 (c, 0.35 in H₂O). Struct. established by synth. Phys. props. are of synthetic material. λ_{max} 240 (ε 4370); 315 (ε 8910) (pH 2) (Derep). λ_{max} 241 (ε 5130); 312 (ε 9120) (pH 12) (Derep). λ_{max} 243 (ε 5500); 315 (ε 9550) (H₂O pH 7) (Derep). λ_{max} 242; 317 (pH 7 buffer) (Berdy). λ_{max} 240; 313 (pH 2 buffer) (Berdy). λ_{max} 241; 312 (pH 12 buffer) (Berdy). λ_{max} 240 (ε 5130); 313 (ε 9550) (H₂O) (Berdy).

6-Deoxo, 5,6-didehydro: Acromelic acid D

[102329-71-5]

C₁₃H₁₄N₂O₆ 294.263

Minor constit. of the toxic mushroom *Clitocybe acromelalga*. Sol. H₂O. [α]_D²⁵ +17.6 (c, 0.07 in H₂O). λ_{max} 221; 269 (H₂O) (Berdy).

6-Deoxo, 5,6-didehydro, 4-decarboxy, 6-carboxy: Acromelic acid E

[145237-01-0]

C₁₃H₁₄N₂O₆ 294.263

Minor constit. of the toxic mushroom *Clitocybe acromelalga*. Sol. H₂O. λ_{max} 266 (H₂O) (Berdy).

Takano, S. *et al.*, *J.A.C.S.*, 1987, **109**, 5523 (*synth*)

Konno, K. *et al.*, *J.A.C.S.*, 1988, **110**, 4807; 6926 (*isol, synth, ir, uv, pmr*)

Takano, S. *et al.*, *Heterocycles*, 1989, **29**, 1473 (*synth, bibl*)

Fushiya, S. *et al.*, *Heterocycles*, 1992, **34**, 1277 (*isol, pmr, cd, struct*)


Barco, A. *et al.*, *Gazz. Chim. Ital.*, 1993, **123**, 185 (*synth*)

Baldwin, J.E. *et al.*, *Tetrahedron*, 1998, **54**, 7465-7484 (*synth*)

Acromelic acid B

A-119

3-[5-Carboxy-4-(carboxymethyl)-3-pyrrolidiny]-1,6-dihydro-6-oxo-2-pyridine-carboxylic acid, 9CI. Acromelinic acid B [86630-10-6]



C₁₃H₁₄N₂O₇ 310.263

Trace alkaloid from the poisonous mushroom *Clitocybe acromelalga*. Shows very potent neuroexcitatory action. Amorph. powder. Sol. H₂O; poorly sol. Me₂CO, hexane. [α]_D²⁰ +50.1 (c, 0.45 in H₂O). Struct. established by synth. Phys. props. are of synthetic material. λ_{max} 231; 308 (pH 2) (Derep). λ_{max} 241; 311 (pH 12) (Derep). λ_{max} 227; 300 (pH 7) (Derep). λ_{max} 236 (ε 6320); 302 (ε 2920) (MeOH) (Berdy).

2'-Decarboxy: **Acromelic acid C**

[133740-47-3]

C₁₂H₁₄N₂O₅ 266.253

From *Clitocybe acromelalga*. Powder. [α]_D²⁰ +31.9 (c, 0.23 in H₂O). λ_{max} 230 (ε 11000); 302 (ε 6150) (MeOH) (Derep). λ_{max} 230 (ε 10592); 301 (ε 6150) (H₂O) (Berdy).

► LD₅₀ (mus, ipr) 10 mg/kg.

Hashimoto, K. *et al.*, *Chem. Lett.*, 1986, 1399

(*synth, uv, ir, cd, abs config*)

Konno, K. *et al.*, *J.A.C.S.*, 1988, **110**, 4807;

6926 (*isol, uv, pmr, synth, struct*)

Takano, S. *et al.*, *Heterocycles*, 1989, **29**, 1473

(*synth, bibl*)


Fushiya, S. *et al.*, *Tet. Lett.*, 1990, **31**, 3901

(*Acromelic acid C*)

Acronyline

A-120

3,12-Dihydro-6-methoxy-3,3,12-trimethyl-7H-pyrano[2,3-c]acridin-7-one, 9CI, 8CI. **Acronine**, INN, USAN. NSC 403169. Compound 42339 [7008-42-6]



C₂₀H₁₉NO₃ 321.375

Alkaloid from *Acronychia baueri* (*Bauerella baueri*), *Melicope leptococca* and *Sarcomelicope argyrophylla*. Antineoplastic agent. Yellow needles (EtOH). Mp 175-176°. Log P 3.7 (calc). λ_{max} 280 (ε 39800); 291 (ε 34700); 304 (ε 19100); 392 (ε 6920) (EtOH) (Derep). λ_{max} 224 (ε 24500); 281 (ε 48000); 293 (ε 43500); 308 (ε 40700); 380 (ε 10250) (MeOH) (Berdy).

► LD₅₀ (mus, orl) 522 mg/kg. Exp. carcinogenic data. UQ0330000

Hydrochloride: [70509-95-4]

Red needles (H₂O). Mp 125-130°.

Heating above melting point causes O-demethylation.

N-De-Me: De-N-methylacronyline. Des-N-methylacronyline

[13255-08-8]

C₁₉H₁₇NO₃ 307.348

Alkaloid from *Glycosmis pentaphylla* and *Murraya paniculata* (Rutaceae). Yellow needles (Me₂CO). Mp 268-270° dec. (with sintering at 260°). λ_{max} 280 (ε 39800); 291 (ε 34700); 304 (ε 19100); 392 (ε 6920) (EtOH) (Derep).

N-De-Me, hydrochloride:

Orange-red needles (MeOH/Et₂O). Mp 137-141°.

N-De-Me, picrate:

Orange needles (EtOH). Mp 222-224° dec.

O-De-Me: Noracronyline. Noracronine

[13161-79-0]

C₁₉H₁₇NO₃ 307.348

Alkaloid from *Glycosmis pentaphylla*, *Murraya paniculata* and *Boenninghausenia albiflora*. Bright yellow needles (C₆H₆ or EtOAc). Mp 203-205°.

O-De-Me, Ac:

Yellow rods (EtOAc). Mp 201-203°.

O,N-Di-de-Me: De-N-methylnoracronyline. Des-N-methylnoracronyline

[13396-93-5]

C₁₈H₁₅NO₃ 293.321

Alkaloid from *Glycosmis pentaphylla* (Rutaceae). Has antitumour activity. Bright yellow needles (C₆H₆/EtOAc). Mp 245-246°.

► **Carcinogenic.**

O,N-Di-de-Me, O-Ac: [36385-14-5]

Fluffy needles (CHCl₃/EtOAc). Mp 235-236.5°.

1,2-Epoxyde: Acronyline epoxide

[118964-07-1]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the bark of *Sarcomelicope simplicifolia* ssp. *neoscotica*. Pale yellow foam. [α]_D²⁰ +2 (c, 0.25 in CH₂Cl₂).

1,2-Dihydro, 1-oxo, 2ξ-hydroxy, O,N-di-de-Me: 1,2-Dihydro-2-hydroxy-N-de-methyl-1-oxonoracronyline

C₁₈H₁₅NO₅ 325.32

Alkaloid from the bark of *Medicosma subsessilis*. Amorph. yellow solid. λ_{max} 221 (log ε 4.98); 255 (log ε 4.99); 383 (log ε 4.04) (EtOH).

11-Methoxy, O-de-Me: 5-Methoxynoracronyline. Baiyumine A

[51179-68-1]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the bark of *Citrus junos* (yuzu) and *Citrus grandis* f. *hakunikuyu* (Rutaceae). Light yellow prisms, or orange plates (Et₂O). Mp 146-148° Mp 160-161° (155-157°).

Hughes, G.K. *et al.*, *Nature (London)*, 1948,

162, 223 (*isol*)

Drummond, L.J. *et al.*, *Aust. J. Sci. Res., Ser. A*,

1949, **2**, 630 (*isol, struct*)

Brown, R.D. *et al.*, *Aust. J. Sci. Res., Ser. A*,

1950, **3**, 593 (*struct*)

MacDonald, P.L. *et al.*, *Aust. J. Chem.*, 1966,

19, 275 (*struct*)

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1966, **29**,

206 (*rev*)

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1966, **55**,

758 (*pharmacol*)

Govindachari, T.R. *et al.*, *Tetrahedron*, 1966,

22, 3245 (*Noracronyline, De-N-methylacronyline*)

Beck, J.R. *et al.*, *J.A.C.S.*, 1968, **90**, 4706 (*synth*)

Gougoutas, J.Z. *et al.*, *Acta Cryst. B*, 1970, **26**,

853 (*cryst struct*)

Hlubucek, J. *et al.*, *Aust. J. Chem.*, 1970, **23**,

1881 (*synth*)

Sullivan, H.R. *et al.*, *J. Med. Chem.*, 1970, **13**,

904 (*metab*)

Gout, P.W. *et al.*, *J. Cell Physiol.*, 1972, **78**, 127

(*biochem*)

Schneider, J. *et al.*, *J. Med. Chem.*, 1972, **15**,

266 (*synth, tox, derivs*)

Lahey, F.M. *et al.*, *Aust. J. Chem.*, 1973, **26**,

2311 (*synth*)

Tan, P. *et al.*, *Cancer Res.*, 1973, **33**, 2320

(*pharmacol*)

Saxton, J.E. *et al.*, *Chem. Heterocycl. Compd.*,

1973, **9**, 379 (*rev*)

Betts, R.E. *et al.*, *J. Med. Chem.*, 1974, **17**,

599

Brannon, D.R. *et al.*, *J. Med. Chem.*, 1974, **17**,

653

Bandranayake, W. *et al.*, *J.C.S. Perkin I*, 1974,

998 (*synth*)

Bert, M. *et al.*, *Phytochemistry*, 1974, **13**, 301

(*isol*)

Blechert, S. *et al.*, *Chem. Ber.*, 1978, **111**, 439

(*synth*)

Rózsa, Zs. *et al.*, *Phytochemistry*, 1978, **17**, 169

(*isol, uv, ir, derivs*)

Adams, J.H. *et al.*, *Tetrahedron*, 1981, **37**, 209

(*synth*)

Gerzon, K. *et al.*, *Alkaloids (Academic Press)*,

1983, **21**, 1 (*rev*)

Funayama, S. *et al.*, *J. Nat. Prod.*, 1983, **46**,

391 (*cmr*)

Skaltsounis, A.L. *et al.*, *J. Nat. Prod.*, 1983, **46**,

732 (*isol*)

Reisch, J. *et al.*, *Annalen*, 1984, 31 (*synth*)

Watanabe, M. *et al.*, *Chem. Pharm. Bull.*, 1984,

32, 1264 (*synth, bibl*)

Funayama, S. *et al.*, *Planta Med.*, 1984, **50**,

121 (*pmr*)

Brum-Bousquet, M. *et al.*, *Planta Med.*, 1985,

51, 536; 1988, **54**, 470 (*isol, uv, ir, pmr, ms, struct, epoxide*)

Ju-ichi, M. *et al.*, *Heterocycles*, 1986, **24**, 1595

(*5-Methoxynoracronyline*)

Wu, T.S. *et al.*, *Phytochemistry*, 1987, **26**, 871

(*Baiyumine A*)

Anand, R.C. *et al.*, *Heterocycles*, 1990, **31**,

1733 (*synth*)

Loughhead, D.G. *et al.*, *J.O.C.*, 1990, **55**, 2245

(*synth, Acronyline, Des-N-methylacronyline*)

Horne, S. *et al.*, *Chem. Comm.*, 1991, 1046

(*synth*)

Reisch, J. *et al.*, *Annalen*, 1994, 317 (*synth, Acronyline epoxide*)

Anand, R.C. *et al.*, *Chem. Comm.*, 1996, 199

(*synth, Acronyline, Des-N-methylacronyline*)

Tillequin, F. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1998, **12**, 1-102 (*rev*)

Minh, N.T. *et al.*, *Z. Naturforsch., B*, 2003, **58**,

1234-1236

(*Dihydrohydroxydemethylxonoracronyline*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, ADR750

Acronyline

A-121

5,11-Dimethoxy-3,3-dimethyl-3H-furano[2,3-b]pyrano[3,2-f]quinoline, 9CI [518-68-3]



C₁₈H₁₇NO₄ 311.337

Alkaloid from *Acronychia baueri*, *Melicope leptococca* and *Sarcomelicope leio-carpa* (Rutaceae). Prisms (MeOH). Mp 151-153°.

Picrate:

Yellow needles (propanol or EtOH). Mp 216-218° dec.

1,2-Dihydro, 1R,2R-dihydroxy: cis-1,2-Dihydro-1,2-dihydroxyacronyridine cis-1,2-Dihydroxy-1,2-dihydroacronyridine [126661-84-5]
C₁₈H₁₉NO₆ 345.351
Alkaloid from bark of *Sarcomelicope dogniensis*. Noncryst. [α]_D +2 (c, 0.25 in MeOH). λ_{max} 225 (sh) (log ε 4.17); 254 (log ε 4.43); 303 (sh) (log ε 3.88); 317 (log ε 4.05); 329 (log ε 4.04); 344 (sh) (log ε 3.83) (MeOH).

1,2-Dihydro, 1R,2S-dihydroxy: trans-1,2-Dihydro-1,2-dihydroxyacronyridine trans-1,2-Dihydroxy-1,2-dihydroacronyridine [126661-85-6]
C₁₈H₁₉NO₆ 345.351
Alkaloid from bark of *Sarcomelicope dogniensis*. Mp 178-180°. [α]_D -8 (c, 0.2 in MeOH). λ_{max} 225 (sh) ; 254 (log ε 4.43); 303 (sh) (log ε 3.88); 317 (log ε 4.05); 329 (log ε 4.04); 344 (sh) (log ε 3.83) (MeOH).

5-Demethoxy: 11-Methoxy-3,3-dimethyl-3H-furano[2,3-b]pyrano[3,2-f]quinoline. Medicosmine [35306-86-6]
C₁₇H₁₅NO₃ 281.31

Alkaloid from *Boronella koniambiensis* and *Medicosma cunninghamii*. Pale cream needles (EtOH or ETOAc/petrol). Mp 138.5-139.5°.

5-Demethoxy, picrate:
Fine yellow needles (EtOH). Mp 190-191°.

5-Demethoxy, 1,2-dihydro, 1R*,2*R-dihydroxy: cis-1,2-Dihydro-1,2-dihydroxy-medicosmine cis-1,2-Dihydroxy-1,2-dihydromedicosmine
C₁₇H₁₇NO₅ 315.325
Alkaloid from the aerial parts of *Boronella koniambiensis*. Amorph. solid. [α]_D -5 (c, 0.1 in CHCl₃). λ_{max} 245 ; 300 ; 312 ; 342 ; 357 (MeOH).

Lamberton, J.A. et al., *Aust. J. Chem.*, 1953, **6**, 66; 173 (*Acronyridine, Medicosmine, isol*)

Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1975, (*pmr, struct*)

Govaindachari, T.R. et al., *Indian J. Biochem.*, 1971, **9**, 1031 (*Medicosmine, synth, uv, pmr*)
Skaltsounis, A.L. et al., *J. Nat. Prod.*, 1983, **46**, 732

Baudouin, G. et al., *J. Nat. Prod.*, 1985, **48**, 260

Mitaku, S. et al., *Ann. Pharm. Fr.*, 1989, **47**, 149-156 (*1,2-Dihydro-1,2-dihydroxyacronyridine*)

Grougnet, R. et al., *J. Nat. Prod.*, 2005, **68**, 1083-1086 (*1,2-Dihydro-1,2-dihydroxy-medicosmine, Medicosmine*)

Ac sine

A-122

[1353-77-1]

C₂₁H₂₉NO₅ 375.464

Struct. unknown. Alkaloid from the roots of *Aconitum excelsum* (Ranunculaceae). Mp 182-185° (slow heating) Mp 192-195° (rapid heating). Contains OAc and 2 OH groups, plus an ether linkage.


Sulfate: Mp 220° dec.

Platonova, T.F. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1958, **28**, 259 (*isol, ir*)

Aconine

A-123

[367452-76-4]



C₃₁H₄₁NO₈ 555.667


Unusual 7,17-epoxyaconitane struct. Alkaloid from the roots of *Aconitum kusnezoffii*. Amorph. powder.

Zinurova, E.G. et al., *Russ. Chem. Bull. (Engl. Transl.)*, 2001, **50**, 311-312 (*isol, pmr, cmr*)

Actiketel

A-124

4-(2,3-Dihydro-2-hydroxy-5,7-dimethyl-3-oxo-2-benzofuranyl)-2,6-piperidinedione, 9CI. RK 4415. Antibiotic RK 4415 [133658-47-6]



C₁₅H₁₅NO₅ 289.287

Glutarimide-type antibiotic. Prod. by *Streptomyces pulveraceus* ssp. *epiderstagenes*. Antitumour agent. Powder. Mp 96-100°. [α]_D 0. λ_{max} 222 (ε 9710); 264 (ε 7630); 350 (ε 2490) (MeOH) (Derep).


Sonoda, T. et al., *J. Antibiot.*, 1991, **44**, 160 (*isol, pmr, cmr*)

Japan. Pat., 1991, 91 255 082; *CA*, **116**, 126985g (*isol*)

Kiyota, H. et al., *Tet. Lett.*, 2000, **41**, 5887-5890 (*synth*)

Actinidine

A-125



(S)-form

C₁₀H₁₃N 147.219

(R)-form [15524-81-9]

Synthetic. Liq. Bp₁₀ 92-94°. [α]_D²⁰ +10.8 (c, 0.36 in CHCl₃).

Picrate: Mp 146-147°.

(S)-form [524-03-8]

Alkaloid from *Actinidia arguta* (taravine), *Actinidia polygama*, *Tecoma fulva*, *Tecoma radicans* and *Valeriana officinalis*

(valerian). Constit. of defensive secretions of various insects incl. *Megacrania alpheus*, *Ontholestes murinus* and *Tapi-noma erraticum*). Powerful attractant for Felidae (cats). Oil. Bp₉ 100-103°. [α]_D¹¹ -7.2 (CHCl₃).

Picrate: Mp 143°.

N-[2-(4-Hydroxyphenyl)ethyl]: N-(p-Hydroxyphenethyl)actinidine

[15794-92-0]
C₁₈H₂₂NO⁺ 268.378

Quaternary alkaloid from the roots of *Valeriana officinalis* (valerian). Mp 201-203° dec. (as chloride). [α]_D²² +50.5 (MeOH).

10-Acetoxy: 10-Acetoxyactinidine

C₁₂H₁₅NO₂ 205.256

Alkaloid from the roots of *Argyria radiata*. Amorph. solid. Stereochem. not certain.

(±)-form [79254-93-6]

Synthetic. Oil. Mp 146-146.5° (as picrate). Bp₉ 100-103°.

Sakan, T. et al., *Bull. Chem. Soc. Jpn.*, 1959, **32**, 315; 1155; 1960, **33**, 712 (*uv, ir, isol, struct, synth*)

Djerassi, C. et al., *Chem. Ind. (London)*, 1961, 210 (*pmr*)

Torrsell, K. et al., *Acta Chem. Scand.*, 1967, **21**, 53 (*N-4-Hydroxyphenethylactinidine*)

Cavill, G.W.K. et al., *Aust. J. Chem.*, 1967, **20**, 349 (*synth, uv, pmr*)

Johnson, R.D. et al., *Phytochemistry*, 1971, **10**, 3334 (*isol, ms*)

Wuest, J.D. et al., *J.O.C.*, 1977, **42**, 2111 (*synth*)

Nitta, M. et al., *Chem. Lett.*, 1981, 933 (*synth*)
Davies, L.B. et al., *J.C.S. Perkin I*, 1981, 1909 (*synth, uv, ir, pmr, ms*)

Tomalski, M.D. et al., *J. Chem. Ecol.*, 1987, **13**, 253-263 (*isol, ms*)

Cossy, J. et al., *Tet. Lett.*, 1988, **29**, 6113 (*synth*)

Ranarivelo, Y. et al., *Heterocycles*, 1990, **31**, 1727-1731 (*(S)-form, synth*)

Huth, A. et al., *J. Chem. Ecol.*, 1990, **16**, 2691-2711 (*isol*)

Cossy, J. et al., *J.O.C.*, 1993, **58**, 2351 (*synth*)

Shiao, M.J. et al., *J.O.C.*, 1993, **58**, 3162 (*synth*)

Stepanov, A.V. et al., *Russ. Chem. Bull. (Engl. Transl.)*, 1998, **47**, 2286-2291 (*synth*)

Jones, K. et al., *Tetrahedron*, 1998, **54**, 2275-2280 (*synth*)

Bianco, A. et al., *Nat. Prod. Lett.*, 2002, **16**, 77-80 (*10-Acetoxyactinidine*)

Actinine

A-126

3-Carboxy-N,N,N-trimethyl-1-propanaminium hydroxide inner salt, 9CI. Tri-N-methyl-γ-butyrobetaine. 4-Aminobutanoic acid betaine. γ-Butyrobetaine. Butyrobetaine. 4-Aminobutanoic acid trimethylbetaine

[407-64-7]

Me₃N⁺CH₂CH₂CH₂COO⁻

C₇H₁₅NO₂ 145.201

Inner salt. Occurs in the sea-rose *Actinia equina*, in muscles of various snakes and in urine in cases of pernicious anaemia. Isol. from brain tissue. Also from *Polyporus sulphureus*. Biosynthetic intermed. in synth. of Carnitine. Plates + 3H₂O (EtOH/Et₂O aq.). Sol. EtOH. Dec. at 220°.

▶ BP3930000

Hydrochloride: [6249-56-5]

Mp 203° Mp 214-219°.

▶ BP3850000

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, 1, 681A (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, 1, 878B (nmr)

Engelard, R. et al., Ber., 1921, 54, 2212

(derivs)

Linneweh, W. et al., Hoppe-Seyler's Z. Physiol.

Chem., 1928, 175, 95; 176, 220; 1929, 181,

48; 182, 9 (isol)

List, P.H. et al., Arch. Pharm. (Weinheim,

Ger.), 1959, 292, 260 (isol)

Hosain, E.A. et al., Nature (London), 1960,

187, 321 (isol)

Cox, R.A. et al., Biochem. J., 1973, 136, 1083

(biosynth)


Anderson, L. et al., Synthesis, 1981, 468

(synth)

Actinophyllic acid

A-127

[845959-90-2]

Absolute
ConfigurationC₁₉H₂₀N₂O₄ 340.378

Indole alkaloid of unknown biogenetic relationships. Alkaloid from the leaves of *Alstonia actinophylla*. Inhibitor of Carboxypeptidase U/Hippuricase. Brown gum. $[\alpha]_D^{25}$ -29 (c, 0.001 in MeOH). λ_{\max} 217 (ε 24410); 280 (ε 17000) (MeOH).


Carroll, A.R. et al., J.O.C., 2005, 70, 1096-1099 (isol, biosynth, pmr, cmr)

Martin, C.L. et al., J.A.C.S., 2008, 130, 7568-7569 (synth)

Actinoplanone D

A-128

3,4,8a,13-Tetrahydro-3,15,16-trihydroxy-1,4-dimethoxy-12-methyl-1H-xantheno[4',3',2':4,5][1,3]benzodioxino[7,6-g]isoquinoline-14,17(2H,9H)-dione, 9CI [116200-81-8]

Absolute
ConfigurationC₂₈H₂₅NO₁₀ 535.506

CAS numbering shown. Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 222-224° dec. $[\alpha]_D^{25}$ -624.1 (c, 0.7 in CHCl₃). λ_{\max} 221 (ε 20900); 249 (ε 24500); 298 (ε 12000); 312 (ε 11220); 328 (ε 10960); 358 (ε 16200); 372 (ε 17400) (EtOH) (Derep).

N-Amino: Actinoplanone C

[116200-80-7]

C₂₈H₂₆N₂O₁₀ 550.521

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 278-280° dec. $[\alpha]_D^{25}$ -646.1 (c, 0.35 in CHCl₃). λ_{\max} 228 (ε 19100); 253 (ε 20400); 302 (ε 14100); 336 (sh); 360 (ε 14100); 375 (ε 15500) (EtOH) (Derep).

N-(3-Oxo-2-butenylideneamino): Actinoplanone G

[116200-82-9]

C₃₂H₃₀N₂O₁₁ 618.596

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Yellowish solid. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 173-176°. $[\alpha]_D^{25}$ -582.3 (c, 0.44 in CHCl₃). λ_{\max} 227 (ε 30900); 253 (ε 31600); 304 (ε 14500); 340 (ε 12600); 380 (ε 16600) (EtOH) (Derep).

11-Chloro: Actinoplanone B

[115655-87-3]

C₂₈H₂₄ClNO₁₀ 569.951

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Yellow amorph. solid. Sol. MeOH, EtOAc, bases, CHCl₃; poorly sol. H₂O. Mp 240-243°. $[\alpha]_D^{25}$ -649.4 (c, 1.54 in CHCl₃). λ_{\max} 228 (ε 30900); 253 (ε 33100); 305 (ε 13500); 336 (sh); 365 (ε 22900); 380 (ε 25100) (EtOH) (Derep). λ_{\max} 226 (ε 25100); 252 (ε 28950); 303 (ε 13180); 363 (ε 17790); 377 (ε 19500) (EtOH) (Berdy).

11-Chloro, N-amino: Actinoplanone A

[115655-86-2]

C₂₈H₂₅ClN₂O₁₀ 584.966

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Yellow needles (Me₂CO). Sol. CHCl₃, bases, MeOH, EtOAc; poorly sol. H₂O. Mp 276-278°. $[\alpha]_D^{25}$ -619.8 (c, 0.29 in CHCl₃). λ_{\max} 228 (ε 30900); 253 (ε 33100); 305 (ε 13500); 336 (sh); 365 (ε 22900); 380 (ε 25100) (EtOH) (Derep).

11-Chloro, N-(isopropylideneamino): Actinoplanone E

[116229-70-0]

C₃₁H₂₉ClN₂O₁₀ 625.03

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Yellowish solid. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 190-193°. $[\alpha]_D^{25}$ -516.7 (c, 0.21 in CHCl₃). λ_{\max} 232 (ε 20900); 253 (ε 26300); 308 (ε 10700); 344 (sh); 370 (ε 17400); 386 (ε 18200) (EtOH) (Derep).

11-Chloro, N-(3-oxo-2-butenylideneamino): Actinoplanone F

[116200-79-4]

C₃₂H₂₉ClN₂O₁₁ 653.041

Isol. from *Actinoplanes* sp. R-304. Cytotoxic agent. Yellowish solid. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 281-282° dec. λ_{\max} 232 (ε 20900); 253 (ε 26300); 308 (ε 10700); 344 (sh); 370 (ε 17400); 386 (ε 18200) (EtOH) (Derep). λ_{\max} 231 (ε 31500); 254 (ε 33700); 308 (ε 13800); 386 (ε 18200) (EtOH) (Berdy).


Kobayashi, K. et al., J. Antibiot., 1988, 41, 502-511 (Actinoplanones A,B)

Kobayashi, K. et al., J. Antibiot., 1988, 41, 741-750 (Actinoplanones C-G)

Actiphenol

A-129

4-[2-(2-Hydroxy-3,5-dimethylphenyl)-2-oxoethyl]-2,6-piperidinedione, 9CI. 3-(2-Hydroxy-3,5-dimethylphenacyl)glutaramide, 8CI. C 73. Antibiotic C 73 [526-02-3]

C₁₅H₁₇NO₄ 275.304

Prod. by *Streptomyces noursei*, *Streptomyces albulus*, *Streptomyces griseus* and *Streptomyces pulveraceus*. Fungicide and shows antitumor propp. Needles (MeOH/CH₂Cl₂). Mp 201-203°.

3-Hydroxy (3,4-trans-): Antibiotic C

73X. C 73X

[20988-33-4]

C₁₅H₁₇NO₅ 291.303

Prod. by *Streptomyces griseus*. Cryst. λ_{\max} 262 (ε 11700); 347 (ε 4300) (MeOH).

▶ LD₅₀ (mus, ipr) 100 - 200 mg/kg.

1'-Hydroxy: Nong-kang 101G

[75314-78-2]

C₁₅H₁₇NO₅ 291.303

Isol. from *Streptomyces aureus*. λ_{\max} 267 (ε 11620); 356 (ε 3820) (EtOH) (Berdy).

1'''-Hydroxy: Antibiotic AH 135Y. AH 135Y

[145066-21-3]

C₁₅H₁₇NO₅ 291.303

Prod. by *Streptomyces albobinaceus*. Antiherpetic agent. Powder. Sol. MeOH, Me₂CO, DMSO, EtOH, EtOAc; fairly sol. CHCl₃, C₆H₆; poorly sol. H₂O, Et₂O, hexane. Mp 220°. λ_{\max} 219 (sh) (ε 14800); 260 (ε 7070); 342 (ε 3260) (MeOH/HCl) (Derep). λ_{\max} 219 (sh) (ε 18500); 260 (ε 9070); 343 (ε 4030) (MeOH) (Derep). λ_{\max} 219 (ε 15472); 260 (ε 9066); 343 (ε 4026) (MeOH) (Berdy). λ_{\max} 219 (ε 14921); 260 (ε 7066); 342 (ε 3257) (MeOH-HCl) (Berdy).

Highet, R.J. et al., Helv. Chim. Acta, 1959, 42, 1523 (isol, struct)

Rao, K.V. et al., J.O.C., 1960, 25, 661 (isol)

Johnson, F. et al., J.O.C., 1962, 27, 3658

(synth)

Kharatyan, S. et al., Chem. Ind. (London),

1963, 1038 (biosynth)

Vondracek, M. et al., Chem. Ind. (London),

1964, 1686 (isol, uv)

Spizek, J. et al., Folia Microbiol. (Prague),

1965, 10, 263 (synth)

Siegel, M.R. et al., Biochem. Pharmacol., 1966,

15, 1213 (pharmacol)

Aszalos, A. et al., J. Med. Chem., 1967, 10,

281-284 (C 73X)

Hua, J.C. et al., CA, 1980, 93, 219277 (Non-

kang 101G)

Zhang, H. et al., CA, 1982, 96, 118576 (isol)

Onan, K.D. et al., Acta Cryst. C, 1985, 41, 428

(cryst struct)

Uyeda, M. et al., J. Antibiot., 1992, 45, 1370

(Antibiotic AH 135Y)

Aculeacin B

A-130

[58814-87-2]

Struct. unknown. Isol. from *Aspergillus aculeatus*. Antibiotic. Amorph. powder. Sol. MeOH, EtOH; fairly sol. EtOAc, H₂O; poorly sol. Me₂CO, hexane, CHCl₃. Mp 148-151°. [α]_D²⁴ -45 (c, 1 in MeOH). λ_{max} 279 (E1%/1cm 18.5) (MeOH) (Berdy). λ_{max} 247 (E1%/1cm 104); 297 (E1%/1cm 21.6) (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 300 - 900 mg/kg. AU3030000


Japan. Pat., 1976, 76 98 387; CA, 86, 15206w (isol)

Satoi, S. et al., J. Antibiot., 1977, 30, 303-307 (isol, props)

Canadian Pat., 1978, 1 041 446; CA, 90, 101942x (manuf)

Acuminaminoside

A-131



C₂₂H₂₃NO₈ 429.426


Constit. of the leaves of *Glochidion acuminatum*. Needles (MeOH). Mp 212-215°. [α]_D²⁸ -33.3 (c, 0.75 in MeOH). λ_{max} 214 (log ε 4.26); 236 (log ε 4); 256 (log ε 4.04); 296 (log ε 3.99); 310 (log ε 4.01) (MeOH).

Otsuka, H. et al., Chem. Pharm. Bull., 2004, 52, 591-596 (isol, pmr, cmr, cryst struct)

Acutiaporberine

A-132

[335311-66-5]




C₄₂H₄₈N₂O₉ 724.849

Aporphine-berberine dimer. Alkaloid from the roots of *Thalictrum acutifolium*. Apoptosis-inducing agent. Light yellow flakes. Mp 183.5-184°. [α]_D²⁴ -14.2 (c, 3.14 in MeOH).

Chen, Q. et al., Planta Med., 2002, 68, 550-553 (isol, activity)

Acutodaurine

A-133



C₂₀H₂₇NO₆ 377.436


Stereochem. not determined; but likely to be the same as the related Acutumine, A-134. Alkaloid from the roots of *Menispermum dauricum*. Oil. [α]_D²⁰ -102 (c, 0.01 in EtOH). λ_{max} 255 (log ε 4.34) (EtOH).

Furumoto, T. et al., Planta Med., 2001, 67, 194-195

Acutumine

A-134

[17088-50-5]



Absolute configuration

C₁₉H₂₄ClNO₆ 397.854

Alkaloid from *Sinomenium acutum* and *Menispermum dauricum* (Menispermaceae). Needles. Mp 238-240° dec. [α]_D¹⁵ -206 (c, 0.69 in Py). pK_a 5.3 (50% EtOH). λ_{max} 245 (ε 18970); 270 (ε 9750) (EtOH).

Ac:

Needles (Et₂O/hexane). Mp 162-164°. [α]_D²⁰ -94 (c, 0.38 in CHCl₃).

N-De-Me: **Acutumidine**

[18145-26-1]

C₁₈H₂₂ClNO₆ 383.828

Alkaloid from *Sinomenium acutum* and *Menispermum dauricum* (Menispermaceae). Needles. Mp 239-241° dec. [α]_D¹⁹ -212.5 (c, 0.16 in Py). pK_a 6.6 (50% EtOH). λ_{max} 244 (ε 13100); 269 (ε 7370) (EtOH).

11-Deoxy: **Acutuminine**

[23512-32-5]

C₁₉H₂₄ClNO₅ 381.855

Alkaloid from the leaves of *Menispermum dauricum* (Menispermaceae). Mp 175-177°. [α]_D -110 (CHCl₃). λ_{max} 243 (ε 18900); 270 (ε 9200) (EtOH).

11-Deoxy, N-de-Me, N-[3-(4-hydroxyphenyl)-2ξ-nitropropyl]: **Nitrotyrasacutuminine**

C₂₇H₃₁ClN₂O₈ 547.003

Alkaloid from the roots of *Menispermum dauricum*. Powder. [α]_D²⁵ -98 (c, 0.2 in MeOH). λ_{max} 251 (log ε 3.42); 275 (log ε 3.63) (MeOH).

Dechloro: **Dechloroacutumine**

[219794-33-9]

C₁₉H₂₅NO₆ 363.41

Alkaloid from *Menispermum dauricum*. Cryst. (EtOAc/hexane). Mp 178-178.5°. [α]_D²⁵ -54 (c, 0.1 in MeOH). λ_{max} 241 (log ε 4.27); 268 (log ε 3.99) (no solvent reported).

11-Epimer: **Dauricumine**

[345641-00-1]

C₁₉H₂₄ClNO₆ 397.854

Alkaloid from *Menispermum dauricum*. Cryst. (MeOH). Mp 205° dec. [α]_D²⁵ -42 (c, 0.1 in Py). λ_{max} 244 (log ε 4.16) (MeOH).

11-Epimer, N-de-Me: **Dauricumidine**

[345640-99-5]

C₁₈H₂₂ClNO₆ 383.828

Alkaloid from *Menispermum dauricum*. Powder. Mp 180° dec. [α]_D²³ +43 (c, 0.1 in Py). λ_{max} 244 (log ε 4.11) (MeOH).

11-Epimer, dechloro: **Dechlorodauricumine**

C₁₉H₂₅NO₆ 363.41

Alkaloid from the roots of *Menispermum dauricum*. Amorph. powder. [α]_D²⁵ +20.7 (c, 0.1 in MeOH). λ_{max} 242 (log ε 4.16) (MeOH).

Nishikawa, M. et al., J.C.S. (B), 1968, 652-658 (cryst struct, abs config)

Barton, D.H.R. et al., J.C.S. (C), 1968, 929-936 (biosynth)

Okamoto, Y. et al., Tet. Lett., 1969, 1933-1935 (Acutumine)

Tomita, M. et al., Chem. Pharm. Bull., 1971, 19, 770-791 (Acutumidine, Acutumine, Dechloroacutumine)

Sugimoto, Y. et al., Phytochemistry, 1998, 49, 1293-1297 (Dechloroacutumine, cryst struct)

Babiker, H.A.A. et al., Biosci., Biotechnol., Biochem., 1999, 63, 515-518 (biosynth, cmr)

Sugimoto, Y. et al., J.O.C., 2001, 66, 3299-3302 (Dauricumidine, Dauricumine)


Yu, B.-W. et al., Nat. Prod. Lett., 2002, 16, 155-159 (Nitrotyrasacutuminine)

Sugimoto, Y. et al., Phytochemistry, 2005, 66, 2627-2631 (Dechlorodauricumine)

Acutopyrrocoline

A-135

[163047-26-5]



Absolute Configuration

C₂₀H₂₄NO₄ 342.414

Alkaloid from the rhizomes of *Sinomenium acutum*.


Moriyasu, M. et al., Nat. Med. (Tokyo), 1994, 48, 287-290

Adalinine

A-136

6-(2-Oxopropyl)-6-pentyl-2-piperidinone, 9CI

[200940-76-7]



C₁₃H₂₃NO₂ 225.33

(R)-form [175669-30-4]


Alkaloid from the European ladybird beetles *Adalia bipunctata* and *Adalia decempunctata*. Oil. [α]_D²⁰ -26 (c, 0.13 in CH₂Cl₂) (natural). [α]_D²⁰ -28.3 (c, 1.6 in CH₂Cl₂) (synthetic).

Lognay, G. et al., J. Nat. Prod., 1996, 59, 510-511 (isol, ir, pmr, cmr, ms, struct)

- Broeders, F. *et al.*, *Bull. Soc. Chim. Belg.*, 1997, **106**, 377-382 (*synth*)
 Yamazaki, N. *et al.*, *Tet. Lett.*, 1999, **40**, 739-742 (*synth, abs config*)
 Honda, T. *et al.*, *Org. Lett.*, 2000, **2**, 3925-3927 (*synth*)
 Laurent, P. *et al.*, *Tetrahedron*, 2001, **57**, 3403-3412 (*biosynth*)

Adenine, JAN, USAN **A-137**

1H-Purin-6-amine, 9CI. 6-Aminopurine. *Vitamin B₄*. *Angustmycin B* [73-24-5]



C₅H₅N₅ 135.128

9H-form is favoured in free base. CAS refers mainly to 1H struct. which is not in reality a favoured tautomer. Widespread throughout animal and plant tissue, purine components of DNA, RNA, and coenzymes and biosynthetic intermediates. Antiviral agent, granulocyte antagonist. Vitamin. Pharmaceutical aid used to extend storage life of whole blood. Needles + 3H₂O (H₂O). Mp 360-365° (anhyd.) dec. pK_{a1} 4.12; pK_{a2} 9.83 (25°).

- Exp. reprod. and teratogenic effects. LD₅₀ (rat, orl) 745 mg/kg. AU6125000

Hydrochloride: [2922-28-3]

Cryst. + 0.5H₂O. Mp 285°.

6N-Me: 6-(Methylamino)purine

[443-72-1]

C₆H₇N₅ 149.155

Minor constit. of bacterial and viral DNA. Mp 312-314° dec. pK_{a1} 4.18; pK_{a2} 9.99 (H₂O).

6N-Benzyl: N-Benzyladenine, 8CI. 6-Benzylaminopurine. **Cytokinin B. BAP** [1214-39-7]

C₁₂H₁₁N₅ 225.252

Widespread in plants. Plant growth regulator. Cryst. (C₆H₆). Mp 229°.

- AU6252200

6N-(3-Hydroxybenzyl): 6N-(3-Hydroxybenzyl)adenine. **meta-Topolin**

C₁₂H₁₁N₅O 241.252

Isol. from *Populus x canadensis* cv. *robusta*. Cytokinin. No phys. props. reported.

1H-form

1-(3-Methyl-2-butenyl): 1-(3-Methyl-2-butenyl)adenine. **1-Prenyladenine**

C₁₀H₁₃N₅ 203.246

Constit. of the leaves of *Bridelia balansae*. Prisms. Mp >300°. λ_{max} 209 (log ε 4.13); 274 (log ε 4.04) (MeOH).

3H-form

3-(3-Methyl-2-butenyl): see Triacanthine, T-451

9H-form

6N-Me, 9-O-β-D-glucopyranosyl:

[253179-56-5]

C₁₂H₁₇N₅O₅ 311.297

Constit. of *Maerua crassifolia*.

6N-(3-Hydroxybenzyl), 9-β-D-glucopyranosyl: **meta-Topolin 9-glucoside**

C₁₈H₂₁N₅O₆ 403.394

Isol. from *Populus x canadensis* cv. *robusta*. Cytokinin. No phys. props. reported.

N⁹-(3-Methyl-2-butenyl): 9-(3-Methyl-2-butenyl)adenine. 9-Prenyladenine

C₁₀H₁₃N₅ 203.246

Constit. of the leaves of *Bridelia balansae*. Prisms. Mp 188-189°. λ_{max} 212 (log ε 4.1); 274 (log ε 2.8) (MeOH).

N⁹-[1-(3,4-Dihydroxyphenyl)ethyl]: 6-Amino-9-[1-(3,4-dihydroxyphenyl)ethyl]purine. 9-[1-(3,4-Dihydroxyphenyl)ethyl]adenine

C₁₃H₁₃N₅O₂ 271.278

Isol. from *Artemisia capillaris*.

Amorph. powder.

[6055-72-7, 2312-73-4]

Skinner, C.G. *et al.*, *J.A.C.S.*, 1955, **77**, 6692-6693 (*Cytokinin B*)

Dunn, D.B. *et al.*, *Biochem. J.*, 1958, **68**, 627-636 (*isol, 6-Methylaminopurine*)

Kistenmacher, T.J. *et al.*, *Acta Cryst. B*, 1974, **30**, 166-168; 1528-1533; 1977, **33**, 253-256 (*cryst struct, 9-Me*)

Lawley, P.D. *et al.*, *Biochem. J.*, 1975, **145**, 73-84 (*ms, derivs*)

Chenon, M.T. *et al.*, *J.A.C.S.*, 1975, **97**, 4636-4642 (*cmr, tautom*)

Kos, N.J. *et al.*, *J.O.C.*, 1979, **44**, 3140-3143 (*synth, nmr*)

Lin, J. *et al.*, *J.A.C.S.*, 1980, **102**, 4627-4631 (*pe, tautom*)

Mathlouth, M. *et al.*, *Carbohydr. Res.*, 1984, **131**, 1-15 (*ir, Raman*)

Remaud, G. *et al.*, *Tetrahedron*, 1986, **42**, 5073-5080 (*N-15 nmr, derivs*)

Strnad, M. *et al.*, *Phytochemistry*, 1997, **45**, 213-218 (*meta-Topolin*)

Ramadan, M.A. *et al.*, *CA*, 2000, **132**, 61604r (*6-N-Me-9-glucosyl*)

Tsai, Y.H. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2452-2457 (*Bridelia prenyladenines*)


Ma, H.-Y. *et al.*, *Molecules*, 2008, **13**, 267-271 (*9-N-Dihydroxyphenylethyl*)

Adenocarpine **A-138**

1,2,3,4-Tetrahydro-1-(1-oxo-3-phenyl-2-propenyl)-5-(2-piperidinyl)pyridine, 9CI.

1'-Cinnamoyl-1,1',2,3,4,4',5,5',6,6'-decahydro-2,3'-bipyridine. N-Cinnamoyl-Δ²-tetrahydroanabasine. **Orensine. Teidine**

[6793-63-1]



C₁₉H₂₄N₂O 296.411

(R)-form [28052-98-4]

Alkaloid from *Adenocarpus commutatus* and *Adenocarpus grandiflorus* (Fabaceae). [α]_D¹⁸ -30.9 (c, 2 in EtOH).

Hydrochloride: Mp 64-65°.

Picrate: Mp 206° dec.

(S)-form [28976-53-6]

Alkaloid from *Adenocarpus complicatus*, *Adenocarpus foliosus* and *Adenocarpus viscosus* (Fabaceae). Resin. [α]_D¹⁸ +29.3 (c, 2 in EtOH).

Hydrochloride: Mp 65° (hydrate) Mp 133-147° dec. (anhyd.).

Picrate: Mp 213°.

(±)-form [494-06-4]

Alkaloid of *Adenocarpus commutatus* and *Adenocarpus grandiflorus* (Fabaceae). Resin or pale-yellow oil.

Hydrochloride: Mp 208-210° Mp 82-83° (dihydrate).

Picrate: Mp 210-211° dec.

(Z)-Cinnamoyl isomer: **Isoorensine**

[28168-92-5]

C₁₉H₂₄N₂O 296.411

Alkaloid from *Adenocarpus complicatus* leaves (Fabaceae).

Ribas, I. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1950, **46**, 489; 1951, **47**, 533; 1953, **49**, 707; 1955, **51**, 55; 1958, **54**, 157; 215 (*isol, struct, resoln, Isoorensine*)

González, A.G. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1951, **47**, 67; 1953, **49**, 783

Ribas, I. *et al.*, *Ann. Pharm. Fr.*, 1952, **19**, 54 (*isol*)

Schöpf, C. *et al.*, *Annalen*, 1964, **674**, 87 (*synth*)


Schöpf, C. *et al.*, *Naturwissenschaften*, 1966, **53**, 274 (*Isoorensine*)

Fitch, W.L. *et al.*, *J.A.C.S.*, 1974, **96**, 4917 (*ms*)

Nehme, M. *et al.*, *An. Quim.*, 1977, **73**, 307 (*isol*)

Adiantifoline**A-139**

[20823-96-5]



C₄₂H₅₀N₂O₉ 726.865

Alkaloid from the roots and tops of *Thalictrum minus* var. *adiantifolium*, the underground parts of *Thalictrum minus* var. *microphyllum* and the roots of *Thalictrum minus* race B (Ranunculaceae). Pale-yellow needles (EtOH). Mp 143.5-144°. [α]_D²⁸ +90 (c, 0.11 in MeOH).

2'-N-De-Me: 2'-Noradiantifoline

[83348-50-9]

C₄₁H₄₈N₂O₉ 712.838

Alkaloid from the roots and rhizomes of *Thalictrum minus* (Ranunculaceae). Amorph. [α]_D²⁵ +39 (c, 0.082 in MeOH).

O³-De-Me: **Thalilitine**

[66408-21-7]

C₄₁H₄₈N₂O₉ 712.838

Minor alkaloid from *Thalictrum revolutum* roots (Ranunculaceae). Amorph. [α]_D²⁰ +92 (c, 0.175 in MeOH).

O⁶-De-Me: **O⁶-Demethyladiantifoline**

C₄₁H₄₈N₂O₉ 712.838Alkaloid from the roots of *Thalictrum minus* (Ranunculaceae). Mp 125-126°. [α]_D²² +18 (c, 0.4 in CHCl₃).**O⁷-De-Me: Thaliadanine**


[31199-54-9]

C₄₁H₄₈N₂O₉ 712.838Alkaloid from *Thalictrum minus* roots and from *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Shows antibacterial activity against *Mycobacterium smegmatis*. Amorph. [α]_D²⁶ +81 (c, 0.41 in MeOH). λ_{\max} 281; 302; 312 (MeOH) (Berdy).**O⁷, O¹³-Di-de-Me: Bursanine**

[82958-13-2]

C₄₀H₄₆N₂O₉ 698.811Alkaloid from the roots and rhizomes of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). [α]_D²⁵ +117 (c, 0.17 in MeOH).Doskotch, R.W. *et al.*, *Tet. Lett.*, 1968, 4999; *J. Nat. Prod.*, 1969, **32**, 29 (*isol, uv, cd, pmr, ms, struct*)Mollov, N.M. *et al.*, *CA*, 1971, **74**, 61584t (*isol, ir, pmr, O-Demethyladiantifoline*)Doskotch, R.W. *et al.*, *J.O.C.*, 1971, **36**, 2409 (*synth*)Geiselman, C.W. *et al.*, *J. Nat. Prod.*, 1972, **35**, 296 (*isol*)Shamma, M. *et al.*, *Tet. Lett.*, 1973, 1859 (*pmr*)Wu, W.-N. *et al.*, *Tetrahedron*, 1977, **33**, 2919 (*uv, cd, pmr, ir, ms, struct, Thalilutine*)Liao, W.T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 271 (*isol, uv, cd, ir, pmr, ms, struct, O-Demethyladiantifoline, Th*)Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1982, **45**, 505 (*uv, cd, pmr, ms, struct, Noradiantifoline*)Guinaudeau, H. *et al.*, *Tet. Lett.*, 1982, **23**, 2523 (*uv, pmr, ms, cd, struct, Bursanine*)**Adifoline****A-140**


[20072-28-0]

C₂₂H₂₀N₂O₇ 424.409Recent work has queried the struct. It is suggested, but not proved, that the prev. isolated Adifoline is identical with Neoadifoline, N-98. Alkaloid from the heartwood of *Adina cordifolia* (Rubiaceae). Yellow needles. Mp 300°.**10-Deoxy: Deoxyadifoline**

[21451-49-0]

C₂₂H₂₀N₂O₆ 408.41Alkaloid from *Adina cordifolia* (Rubiaceae). Mp 253-255° (as Ac, Me ester). [α]_D²⁰ +560 (c, 0.105 in CHCl₃) (Ac, Me ester).Cross, A.D. *et al.*, *J.C.S.*, 1961, 2714 (*isol, uv, ir*)Brown, R.T. *et al.*, *Chem. Comm.*, 1968, 350 (*uv, ir, pmr, ms, struct*)Merlini, L. *et al.*, *Gazz. Chim. Ital.*, 1968, **98**, 974 (*Deoxyadifoline*)Balázs, B. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 751-753**Adirubine****A-141**

[38474-14-5]



Absolute Configuration

C₂₂H₂₈N₂O₅ 400.474Alkaloid from *Adina rubescens* (Rubiaceae).*Me ester*: [α]_D²⁵ -34 (CHCl₃).*O-Ac*: Mp 151-154°. [α]_D²⁵ -19 (CHCl₃).**19E,20-Didehydro, O-Ac: 19,20-Dehydroadirubine acetate**


[77485-21-3]

C₂₄H₂₈N₂O₆ 440.495Trace alkaloid from the leaves of *Rauwolfia oreogiton* (Apocynaceae). Off-white amorph. powder. λ_{\max} 227 (log ϵ 4.4); 285 (log ϵ 3.8); 291 (log ϵ 3.82) (MeOH).**17-Deoxy, 16,17-didehydro: Anhydroadirubine**

[58514-04-8]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from *Adina rubescens* (Rubiaceae). Amorph. powder (as Me ester). [α]_D²⁵ -28 (CHCl₃) (Me ester). λ_{\max} 225; 273; 282; 289 (MeOH) (Me ester).Brown, R.T. *et al.*, *Chem. Comm.*, 1972, 1007-1008; 1976, 530-531 (*ir, pmr, ms, struct, config*)Brown, R.T. *et al.*, *Phytochemistry*, 1975, **14**, 2527-2529 (*Anhydroadirubine*)Van Tamelen, E.E. *et al.*, *Bioorg. Chem.*, 1976, **5**, 283-308 (*ester, synth*)Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 2741-2745 (*19,20-Dehydroadirubine acetate*)**Adlumidiceine enol lactone****A-142**


6-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]methylene]furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI [75832-77-8]



R = R' = Me

C₂₂H₂₃NO₆ 397.427Alkaloid from *Fumaria schrammii*.Popova, M. *et al.*, *Planta Med.*, 1980, **40**, 156**Adlumidiceine****A-143**

5-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]acetyl]-1,3-benzodioxole-4-carboxylic acid, 9CI [51059-65-5]

C₂₁H₂₁NO₇ 399.399Alkaloid from *Corydalis sempervirens* (*Corydalis glauca*), *Corydalis cava* (*Corydalis tuberosa*), *Corydalis lutea*, *Fumaria schrammii* and *Papaver rhoeas* (corn poppy) (Papaveraceae, Papaveraceae). Cryst. (MeOH). Mp 244-246°.**1-Hydroxy: Narceimicine**

[116368-95-7]

C₂₁H₂₁NO₈ 415.399Alkaloid from the seeds of *Fumaria indica* (Papaveraceae). Pale-yellow granules (MeOH aq.). Mp 242-246° dec. Shown in the paper as the enediol tautomer.**1-Hydroxy, Me ester: Paprafumine**

[169626-17-9]

C₂₂H₂₃NO₈ 429.426Alkaloid from aerial parts of *Fumaria indica* (Papaveraceae). Amorph. solid. Shown in the paper as the enediol tautomer.**1-Oxo: Narceimine. Bicucullinine. Alkaloid F45**

[59443-00-4]

C₂₁H₁₉NO₈ 413.383Alkaloid from *Fumaria indica* and *Corydalis ochroleuca* (Papaveraceae). Mp 268° (259-260°).**1-Oxo, Me ester: Mp 188°.****1-Oxo, N-Me: N-Methylnarceimine. N-Methylnarceimicine (incorr.)**

[777007-71-3]

C₂₂H₂₂NO₈⁺ 428.418Quaternary alkaloid from *Corydalis saxicola*. Yellow cryst. (MeOH). Mp 194-196°. Counterion not specified. λ_{\max} 207 (log ϵ 4.48); 331 (log ϵ 4.02) (MeOH).Preininger, V. *et al.*, *Phytochemistry*, 1973, **12**, 2513 (*ir, uv, pmr, ms, struct*)Preininger, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1975, **40**, 699 (*ir, uv, pmr, ms, struct*)Rodrigo, R.G.A. *et al.*, *Can. J. Chem.*, 1976, **54**, 471 (*isol, ir, pmr, cmr, ms, struct, Narceimine*)Seth, K.K. *et al.*, *Chem. Ind. (London)*, 1979, 744 (*ir, uv, pmr, ms, struct, Narceimine*)Popova, M.E. *et al.*, *Planta Med.*, 1982, **45**, 120 (*isol*)Tripathi, Y.C. *et al.*, *Phytochemistry*, 1988, **27**, 1918 (*Narceimine, isol, ir, uv, pmr, ms, struct*)Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1995, **40**, 593 (*Parafumine*)Wu, Y.-R. *et al.*, *Planta Med.*, 2007, **73**, 787-791 (*N-Methylnarceimine*)

Adlumidicine enol lactone A-144

6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI

As Adlumiceine enol lactone, A-142 with $R, R' = -CH_2-$

$C_{21}H_{19}NO_6$ 381.384

(E)-form [51059-67-7]

Alkaloid from *Fumaria schrammii*, *Corydalis sempervirens*, *Corydalis ochotensis* and *Corydalis lutea* (Papaveraceae).

Cryst. (Et₂O). Mp 200-203°.

(Z)-form**Aobamidine**

[59614-38-9]

Alkaloid from *Corydalis ochotensis* and *Corydalis lutea* (Papaveraceae). Yellowish powder (Et₂O). Mp 195-197°.

Preininger, V. et al., *Phytochemistry*, 1973, **12**, 2513 (uv, ir, pmr, struct)


Kametani, T. et al., *J.C.S. Perkin 1*, 1977, 390 (isol, uv, ir, pmr, ms, struct)

Preininger, V. et al., *Planta Med.*, 1978, **33**, 396 (isol)

Adlumidine

A-145

6-(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI

**(+)-form**

$C_{20}H_{17}NO_6$ 367.357

Diastereoisomer of Bicuculline, B-119.

(+)-form [550-49-2]

Alkaloid from *Corydalis decumbens*, *Corydalis gigantea*, *Corydalis incisa*, *Corydalis ochotensis*, *Corydalis ochroleuca*, *Corydalis remota*, *Corydalis sibirica*, *Corydalis thalictrifolia*, *Fumaria indica*, *Fumaria parviflora* and *Adlumia fungosa* (*Adlumia cirrhosa*) (Papaveraceae, Papaveraceae). Cryst. (Me₂CO). Mp 239-240° (237°). $[\alpha]_D^{20} +116.2$ (c, 2 in CHCl₃).

▶ AV3085000

(-)-form**Capnoidine**

[485-50-7]

Alkaloid from *Corydalis cava*, *Corydalis crystallina*, *Corydalis gigantea*, *Corydalis gortschakovii*, *Corydalis marshalliana*, *Corydalis pseudo-adunca*, *Corydalis sempervirens* and *Fumaria vaillantii* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 239° (235°). $[\alpha]_D^{25} -113.2$ (c, 0.8 in CHCl₃) (-100).

▶ EX8585000

(±)-form [64397-08-6]

Alkaloid from *Corydalis rosea*. Cryst. (CHCl₃/MeOH, MeOH or C₆H₆/Me₂CO). Mp 205° (184-186°, 198-199°).

Manske, R.H.F. et al., *J.A.C.S.*, 1950, **32**, 3207 (isol, struct)

Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1964, **29**, 2328 (config)

Snatzke, G. et al., *Tetrahedron*, 1969, **25**, 5059 (ord, cd)

Margvelashvili, N.N. et al., *Khim. Prir. Soedin.*, 1972, **8**, 127; 1978, **14**, 592; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 131; 1978, **14**, 509 (isol, ir, uv)


Lu, S.-T. et al., *J.C.S. Perkin 1*, 1976, 63 (isol)

Blaskó, G. et al., *J. Nat. Prod.*, 1982, **45**, 121 (occur)

Ribár, B. et al., *Acta Cryst. C*, 1991, **47**, 2191 (cryst struct)

Adlumine

A-146

**(+)-form**

$C_{21}H_{21}NO_6$ 383.4

Diastereoisomer of Corlumine, C-653.

(+)-form [524-46-9]

Alkaloid from *Adlumia fungosa* (*Adlumia cirrhosa*) and *Fumaria rostellata* (Papaveraceae, Papaveraceae). Mp 180°. $[\alpha]_D +42$ (CHCl₃).

(-)-form [21414-43-7]

Alkaloid from *Corydalis scouleri*, *Corydalis ophiocarpa*, *Corydalis rosea*, *Corydalis gortschakovii*, *Corydalis gigantea*, *Corydalis sempervirens*, *Fumaria kralikii*, *Fumaria parviflora* and *Fumaria vaillantii* (Papaveraceae). Cryst. (CHCl₃ or CHCl₃/MeOH). Mp 178-180°. $[\alpha]_D^{20} -42$ (c, 0.38 in CHCl₃). $[\alpha]_D^{20} -51$.

N-Me: N-Methyladlumine

[80550-26-1]

$C_{22}H_{24}NO_6$ 398.435

Quaternary alkaloid from *Fumaria vaillantii*. Cryst. (MeOH) (as iodide). Mp 198-199° (iodide). $[\alpha]_D -45$ (c, 0.5 in MeOH) (iodide).

O⁶-De-Me: Corledine

$C_{20}H_{19}NO_6$ 369.373

Alkaloid from *Corydalis ledebouriana* (Papaveraceae). Cryst. (MeOH). Mp 210-212°. $[\alpha]_D -100$ (c, 0.2 in MeOH). Methylation gives Adlumine.

O⁷-De-Me: Severtzine. 1-Adlumine. Severtcine

[59272-73-0]

$C_{20}H_{19}NO_6$ 369.373

Alkaloid from *Corydalis severtzovii* (Papaveraceae). Cryst. (MeOH). Mp 94-95° (foams). $[\alpha]_D -52$ (c, 0.91 in CHCl₃).

(±)-form [38184-69-9]

Alkaloid from *Corydalis rosea* (Papaveraceae). Mp 191° (175°, 184-186°).

Safe, S. et al., *Can. J. Chem.*, 1964, **42**, 160 (pmr, config)

Snatzke, G. et al., *Tetrahedron*, 1969, **25**, 5059 (ord)

Margvelashvili, N.N. et al., *Khim. Prir.*

Soedin., 1972, **8**, 127; 1976, **12**, 832; 1978, **14**, 592; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 131; 1976, **12**, 509; 1978, **14**, 754 (isol, ir, uv)

Shamma, M. et al., *Tet. Lett.*, 1974, 2339 (synth, pmr)

Preininger, V. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 699 (isol)

Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 811; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 826 (Severtzine)

Hughes, D.W. et al., *Can. J. Chem.*, 1976, **54**, 2252 (cmr)

Nalliah, B.C. et al., *Can. J. Chem.*, 1977, **55**, 922; 1979, **57**, 1546 (synth)

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1978, **14**, 103; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 82 (cd)

Alimova, M. et al., *Khim. Prir. Soedin.*, 1981, **17**, 602-604; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 437-438 (N-Methyladlumine)


Blaskó, G. et al., *J. Nat. Prod.*, 1982, **45**, 105 (occur)

Seeger, C. et al., *Magn. Reson. Chem.*, 2004, **42**, 882-886 (pmr, cmr)

Adociaquinone B

A-147

[113831-00-8]



Absolute Configuration

$C_{22}H_{17}NO_6S$ 423.445

Isol. from the sponges *Adocia* and *Xestospongia* spp. Mildly cytotoxic. Topoisomerase inhibitor. Yellow solid. $[\alpha]_D +22$. Slowly dec. >300°. λ_{max} 288 (ε 36000); 340 (ε 12000) (EtOH) (Derep). λ_{max} 294 (ε 14400) (MeOH) (Berdy).

3-Oxo: 3-Oxoadociaquinone B. 3-Ketoadociaquinone B

[848242-73-9]

$C_{22}H_{15}NO_7S$ 437.429

Isol. from a *Xestospongia* sp. Yellowish powder. $[\alpha]_D^{23} +13$ (c, 0.12 in MeOH). λ_{max} 273 (log ε 3.6) (MeOH).

Schmitz, F.J. et al., *J.O.C.*, 1988, **53**, 3922-3925 (isol, pmr, cmr)

Concepcion, G.P. et al., *J. Med. Chem.*, 1995, **38**, 4503-4507 (isol)


Harada, N. et al., *Tetrahedron: Asymmetry*, 1995, **6**, 375-376 (synth, abs config)

Cao, S. et al., *Bioorg. Med. Chem.*, 2005, **13**, 999-1003 (3-Oxoadociaquinone B)

Adonifoline

A-148

[115712-88-4]



$C_{18}H_{23}NO_7$ 365.382

Revised struct. (1992). Two previous


isolates shown to be identical with the alkaloid now named Adonifoline. Alkaloid from *Senecio adonidifolius* and *Senecio dolichodoryius* (Asteraceae). Cryst. (Me₂CO). Mp 200°. [α]_D²⁰ +84.74 (c, 0.78 in MeOH).

Witte, L. *et al.*, *Phytochemistry*, 1992, **31**, 1027-1028 (*isol*, *pmr*, *cmr*, *ms*)

Adouetine X

Ceanothamine B
[19542-37-1]

A-149



Absolute
Configuration

C₂₈H₄₄N₄O₄ 500.68

Alkaloid from *Waltheria americana*, the root bark of *Ceanothus americanus* (New Jersey tea) and *Zizyphus jujuba* var. *inermis*, and the leaves of *Alphitonia macrocarpa* (Sterculiaceae, Rhamnaceae). Needles (MeOH or CH₂Cl₂/EtOAc). Mp 279-280.5°. [α]_D²⁵ -370 (c, 0.205 in CHCl₃).

N-De-Me: Discarine F. N-Demethyladouetine X

[96562-84-4]
C₂₇H₄₂N₄O₄ 486.653

Alkaloid from the bark of *Discaria febrifuga* (Rhamnaceae). Mp 264°. [α]_D²⁰ -191 (CHCl₃).

Païs, M. *et al.*, *Ann. Pharm. Fr.*, 1963, **21**, 139-146; *CA*, **59**, 5215c (*isol*, *ir*, *pmr*)

Warnhoff, E.W. *et al.*, *Can. J. Chem.*, 1965, **43**, 2594-2602 (*isol*, *uv*, *ms*, *pmr*)

Païs, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 1145-1148 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Servis, R.E. *et al.*, *J.A.C.S.*, 1969, **91**, 5619-5624 (*isol*, *ms*)

Branch, G.B. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2209-2216 (*isol*)


Otsuka, H. *et al.*, *Phytochemistry*, 1974, **13**, 2016 (*isol*, *ir*, *pmr*, *ms*)

Morel, A. *et al.*, *Z. Naturforsch., B*, 1984, **39**, 1825 (*Discarine F*)

Adouetine Y

[19542-39-3]

A-150



Absolute
Configuration

C₃₁H₄₂N₄O₄ 534.697

Alkaloid from *Waltheria americana*, *Waltheria douradhina*, *Myrianthus arboreus*, *Melochia corchorifolia*, *Discaria longispina*, *Discaria febrifuga*, *Discaria americana* and *Ceanothus sanguineus*. Cryst. (CHCl₃/Et₂O). Mp 295-297°. [α]_D²⁰ -390 (c, 1 in CHCl₃).

N-De-Me: N-Demethyladouetine Y

[73045-49-5]
C₃₀H₄₀N₄O₄ 520.67

Alkaloid from the root bark of *Ceanothus sanguineus* (Rhamnaceae). Mp 229°. Opt. rotn. not recorded.

10,11-Dihydro, 11-hydroxy: Discarine G

[94901-64-1]
C₃₁H₄₄N₄O₅ 552.712

Alkaloid from the root bark of *Discaria febrifuga* (Rhamnaceae). Mp 257°. [α]_D²⁰ -366 (c, 0.1 in MeOH). Stereochem. not determined.

Stereoisomer (1): Lotusanine A

[164455-28-1]
C₃₁H₄₂N₄O₄ 534.697

Alkaloid from *Zizyphus lotus*. Amorph. solid. Has (3*R*,4*R*)-config. with the other centres unassigned. Stated to be racemic. λ_{max} 208 (log ε 4.2) (MeOH).

Stereoisomer (2): Myrianthine B. Alkaloid AM 1

[24532-77-2]
C₃₁H₄₂N₄O₄ 534.697

Alkaloid from *Myrianthus arboreus* and leaves of *Antidesma montana*. Needles (MeOH). Mp 302° dec. (292°). [α]_D²⁰ -360 (c, 0.14 in CHCl₃) (-294). Has (4*S*,7*S*)-config. with the other centres undetermined.

Marchand, J. *et al.*, *Ann. Pharm. Fr.*, 1968, **26**, 771-778; *CA*, **71**, 42203q (*Myrianthine B*)

Païs, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 1145-1148 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Tschesche, R. *et al.*, *Tet. Lett.*, 1968, 3817-3818 (*isol*, *ms*)

Merkuza, V.M. *et al.*, *Phytochemistry*, 1974, **13**, 1279-1282 (*ir*, *ms*)

Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 663-668 (*N-Demethyladouetine Y*)

Herzog, R. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1984, **365**, 1351-1354 (*Discarine G*)

Arbain, D. *et al.*, *Phytochemistry*, 1993, **33**, 1263-1266 (*AM-1*)

Abu-Zarga, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 504-511 (*Lotusanine A*)


Da Silva, U.F. *et al.*, *Phytochem. Anal.*, 1996, **7**, 20-23 (*config*)

Tan, N.H. *et al.*, *Chem. Rev.*, 2006, **106**, 840-895 (*bibl*)

Adouetine Z

A-151

N,N-Dimethylphenylalanyl-N-[5,8-dioxo-3-phenyl-7-(phenylmethyl)-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]prolinamide, 9CI. N-Methylferetine. Adouetine
[19542-40-6]



Absolute
Configuration

C₄₂H₄₅N₅O₅ 699.848

Alkaloid from *Waltheria americana*, and

from the leaves of *Feretia apodanthera* and *Melochia pyramidata* (Sterculiaceae, Rubiaceae). Cryst. (cyclohexane). Mp 135-140° Mp 140-145°. [α]_D²⁰ -184 (c, 1 in CHCl₃).

▶ TW3577000

N-De-Me: Feretine. N-Demethyladouetine Z

[56031-09-5]
C₄₁H₄₃N₅O₅ 685.821

Alkaloid from the leaves of *Feretia apodanthera* (Rubiaceae). Cryst. (cyclohexane). Mp 123°. [α]_D²⁰ -139 (c, 1 in CHCl₃).

10,11-Dihydro:

Cryst. (Me₂CO/hexane). Mp 221°. [α]_D²⁰ -87 (CHCl₃).

17-Hydroxy: Oxyphylline A

[959855-77-7]
C₄₂H₄₅N₅O₆ 715.847

Alkaloid from the stem bark of *Zizyphus oxyphylla*. Cryst. Mp 204-206°. λ_{max} 233 (MeOH).

Païs, M. *et al.*, *Ann. Pharm. Fr.*, 1963, **21**, 139-146; *CA*, **59**, 5215c (*isol*, *ir*, *pmr*)

Païs, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 1145-1148 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Bailleul, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **279**, 949-951 (*Feretine*)


Medina, E. *et al.*, *Annalen*, 1981, 538-545 (*isol*, *ir*, *cmr*, *ms*)

Inayat-ur-Rahman, *et al.*, *Nat. Prod. Res.*, 2007, **21**, 243-253 (*Oxyphylline A*)

Adrenaline, BAN

A-152

4-[1-Hydroxy-2-(methylamino)ethyl]-1,2-benzenediol, 9CI. 3,4-Dihydroxy-α-(methylamino)methylbenzyl alcohol, 8CI. 1-(3,4-Dihydroxyphenyl)-2-methylaminoethanol. Epinephrine, INN, USAN. Bronkaid. Epifrin. Glaucon. Bosmin. Surprenin. NSC 62786
[6912-68-1]



C₉H₁₃NO₃ 183.207

Used in photometric detn. of Ti. Active constit. of the adrenal gland. α,β-Adrenoceptor agonist, vasoconstrictor, inotropic agent. Mydriatic. Used in management of anaphylaxis, and in cardiopulmonary resuscitation. Orally inactive. Used in combination with fluorouracil in the treatment of genital warts (AccuSite). Log P -0.68 (calc). First hormone to be isol. (in 1856) and synthesised (1904).

▶ May cause contact dermatitis.

(R)-form

L-form (absol.)

[51-43-4]

Powder. Sol. alkalis, acids; spar. sol. H₂O, EtOH; insol. Et₂O. Mp 216° dec. [α]_D²⁰ -53 (c, 4 in 1 M HCl). Pharmacol. more active isomer.

▶ LD₅₀ (rat, skn) 62 mg/kg. LD₅₀ (rat, ivn)

0.15 mg/kg. DO2625000

Hydrochloride: [55-31-2]

Mp 157°.

► DO3150000

Tartrate: **Epinephrine bitartrate, USAN.**

Epitrate

[51-42-3]

[24351-82-4] Ophthalmic adrenergic agent. Cryst. Mp 147-154°.

► DO3500000

Cyclic borate: **Epinephryl borate, USAN.**

Epinalf. Epinephrine borate

[5579-16-8] Ophthalmic adrenergic agent.

N-Me: 4-[2-(Dimethylamino)-1-hydroxyethyl]-1,2-benzenediol, 9CI. N-

Methyladrenaline

[554-99-4]

C₁₀H₁₅NO₃ 197.233

Alkaloid from tubers of *Aconitum nasutum* (Ranunculaceae).

► DO5425000

N-De-Me: see Norepinephrine, N-298

O³,O⁴,N-Tri-Me: see Macromerine, M-18

(S)-form

D-form (obsol.)

[150-05-0]

Light brown cryst. Mp 211-212°. About 12 times less hormonally active than the (R)-enantiomer.

► DO2800000

(±)-form

Racpinephrine, INN. Racpinephrine, USAN.

Vaponephrin

[329-65-7]

Microscopic cryst. Sl. sol. H₂O. Mp 230° dec.

► DO2975000

Hydrochloride: **Racpinephrine hydrochloride, USAN**

[329-63-5]

Cryst. (EtOH). Mp 157°.

► DO3330000

Di-Ac:

C₁₃H₁₇NO₅ 267.281

Mp 168°.

Di-Ac; hydrochloride: Mp 155-156°.

(ξ)-form

Isol. from callus tissue of *Portulaca grandiflora*.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 1296A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 625B (nmr)

Vulpian, M. et al., *C. R. Hebd. Seances Acad. Sci.*, 1856, **43**, 663-665

(isol)

Stolz, F. et al., *Ber.*, 1904, **37**, 4149-4154

(synth)

Ciocca, B. et al., *Boll. Chim. Farm.*, 1934, **73**,

241-245; *CA*, **28**, 4398 (synth)

Pratesi, P. et al., *J.C.S.*, 1959, 4062-4065 (abs config)

Payne, K.R. et al., *Ind. Chem.*, 1961, **37**, 523-527 (manuf, bibl)

Thies, H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 194-196 (uv)

Jerman, L. et al., *Anal. Chim. Acta*, 1966, **36**, 240-241 (use)

Carlström, D. et al., *Acta Cryst. B*, 1973, **29**, 161-167 (cryst struct)

Hawkins, C.J. et al., *Aust. J. Chem.*, 1973, **26**, 2553-2554 (resoln)

Samokish, I.I. et al., *CA*, 1975, **84**, 102283 (N-Me, isol)

Nagatsu, T. et al., *Method. Chim.*, 1977, **11**, 194-199 (rev)

Szuczewski, D. et al., *Anal. Profiles Drug Subst.*, 1978, **7**, 193-229 (rev)

Weisser, U. et al., *Med. Welt*, 1980, **31**, 40-44 (synth)

Lai, A. et al., *J.C.S. Faraday 2*, 1981, **77**, 227-233 (cmr)

Endress, R. et al., *J. Plant Physiol.*, 1984, **115**, 291-295; *CA*, **101**, 127016u (occur)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms, 6th edn., Akademie-Verlag*, 1987, 1288 (synonyms)

Kirk-Othmer Encycl. Chem. Technol., 4th edn., Wiley, 1991, **9**, 715-730 (rev)


Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 813

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 10th edn., J. Wiley*, 2000, AES000; VG000; MJV000; AES250; AES500; EBB500; AES650

Aegeline

A-153

N-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]-3-phenyl-2-propenamide, 9CI. N-β-Hydroxy-β-p-methoxyphenethylcinnamamide. N-Cinnamoyl-2-hydroxy-2-(4-methoxyphenyl)ethylamine. Cinnamic acid 2-hydroxy-2-(p-methoxyphenyl)ethylamide. Egeline
[456-12-2]



C₁₈H₁₉NO₃ 297.353

Originally assigned the formula

C₁₈H₁₈O₄.

(+)-form [15298-36-9]

Synthetic. Cryst. (EtOH). Mp 196-197°. [α]_D²² +36 (c, 0.4 in CHCl₃). [α]_D²¹ -48.1 (c, 0.5 in EtOH).

(-)-form [15298-37-0]

Synthetic. Cryst. (EtOH). Mp 196-197°. [α]_D -35.1 (CHCl₃). [α]_D +47.5 (c, 0.5 in EtOH).

(±)-form [37791-13-2]

Alkaloid from the leaves of *Aegle marmelos* (bael), *Zanthoxylum coriaceum* and *Zanthoxylum ocumarense* (Rutaceae). Antihyperglycaemic and antidiyslipidaemic agent. Cryst. (EtOH/EtOAc). Mp 176° (173-175°).

Ac:

C₂₀H₂₁NO₄ 339.39

Plates (EtOAc). Mp 124°.

Me ether: N-[2-Methoxy-2-(4-methoxyphenyl)ethyl]cinnamide. 7-O-Methyl-aegeline

[70546-93-9]

C₁₉H₂₁NO₃ 311.38

Isol. from *Aegle marmelos* (bael).

Cryst. (C₆H₆/hexane). Mp 135°. Artifact.

Et ether: N-[2-Ethoxy-2-(4-methoxyphenyl)ethyl]cinnamide

[70546-94-0]

C₂₀H₂₃NO₃ 325.407

Isol. from *Aegle marmelos* (bael).

Cryst. (C₆H₆/hexane). Mp 99-100°.

Artifact.

3-Methoxy: **3-Methoxyaegeline**

C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Zanthoxylum syncarpum*. Cryst. (EtOAc/MeOH). Mp 138-139°. λ_{max} 218 (log ε 4.37); 225 (log ε 4.57); 278 (log ε 4.65) (MeOH).

3-Methoxy, Ac: **7-O-Acetyl-3-methoxyaegeline**

C₂₁H₂₃NO₅ 369.416

Alkaloid from the leaves of *Zanthoxylum syncarpum*. Pale yellow powder. λ_{max} 242 (log ε 4.82); 252 (log ε 4.67); 282 (log ε 4.98) (MeOH).

Chatterjee, A. et al., *J.O.C.*, 1959, **24**, 687 (isol, uv, ir, struct, synth)

Albónico, S.M. et al., *J.C.S. (C)*, 1967, 1327 (synth)

Della Casa de Marcano, D.

et al., *Phytochemistry*, 1972, **11**, 1531 (isol)

Manandhar, M.D. et al., *Phytochemistry*, 1978, **17**, 1814 (derivs)

Patra, A. et al., *Indian J. Chem., Sect. B*, 1979, **17**, 385 (isol, uv, ir)

Swinehart, J. et al., *Phytochemistry*, 1980, **19**, 1219 (isol)

Patra, A. et al., *Org. Magn. Reson.*, 1981, **16**, 65 (cmr)

Sharma, B.R. et al., *Phytochemistry*, 1981, **20**, 2606 (isol)

Somanathan, R. et al., *Synth. Commun.*, 1983, **13**, 273 (synth, ir, ms, pmr, cmr)

Brown, R.F.C. et al., *Tetrahedron: Asymmetry*, 1993, **4**, 205 (synth)


Ross, S.A. et al., *J. Nat. Prod.*, 2005, **68**, 1297-1299 (3-Methoxyaegeline, 7-Acetyl-3-methoxyaegeline)

Narender, T. et al., *Bioorg. Med. Chem. Lett.*, 2007, **17**, 1808-1811 (isol, activity)

Aerophobin 1

A-154

[87075-24-9]



n = 2, X = H

C₁₅H₁₆Br₂N₄O₄ 476.124

Isol. from the sponges *Verongia aerophoba* and *Verongula rigida*. Also from *Tylodina perversa*. Mp 164-167° (Ac). [α]_D +187 (c, 2.0 in MeOH).

Cimino, G. et al., *Tet. Lett.*, 1983, **24**, 3029 (isol, pmr, cmr, ms, struct)

Nishiyama, S. et al., *Bull. Chem. Soc. Jpn.*, 1985, **58**, 3453 (synth)

Gunasekera, M. et al., *J. Nat. Prod.*, 1989, **52**, 753 (isol, uv, ir, pmr, cmr, ms)

Teeyapant, R. et al., *Z. Naturforsch., C*, 1993, **48**, 640-644 (isol)

Aerophobin 2

A-155

[87075-23-8]

As Aerophobin 1, A-154 with $n = 3$, $X = \text{NH}_2$ $\text{C}_{16}\text{H}_{19}\text{Br}_2\text{N}_5\text{O}_4$ 505.165Isol. from the sponges *Verongia aerophoba* and *Aiolochoira crassa*. $[\alpha]_{\text{D}}^{20} +139$ (c, 1.9 in MeOH).N-Me ($X = \text{NHMe}$): **N-Methylaerophobin 2**

[209472-96-8]

 $\text{C}_{17}\text{H}_{21}\text{Br}_2\text{N}_5\text{O}_4$ 519.192Alkaloid from the sponge *Aiolochoira crassa*.**13,14-Dihydro, 14-oxo: 14-Oxo-aerophobin 2**

[232259-26-6]


 $\text{C}_{16}\text{H}_{19}\text{Br}_2\text{N}_5\text{O}_5$ 521.165Isol. from *Aplysina insularis*.Cimino, G. et al., *Tet. Lett.*, 1983, **24**, 3029

(Aerophobin 2)


Assmann, M. et al., *Z. Naturforsch., C*, 1998, **53**, 398-401 (N-Methylaerophobin 2)Fendert, T. et al., *Z. Naturforsch., C*, 1999, **54**, 246-251 (14-Oxo-aerophobin 2)**Aerophysinillin**

A-156

[1056030-28-4]

 $\text{C}_{23}\text{H}_{24}\text{Br}_4\text{N}_4\text{O}_8$ 804.081Closely related to Aerothionin, A-158. Isol. from an *Aplysinella* sp. Powder. $[\alpha]_{\text{D}}^{20} +38.2$ (c, 0.2 in MeOH). λ_{max} 223 (log ϵ 5.5); 256 (log ϵ 5.48) (MeOH).Ankudey, F.J. et al., *Planta Med.*, 2008, **74**, 555-559 (isol, cd, pmr, cmr)**Aerophysinin 1**

A-157

3,5-Dibromo-1,6-dihydroxy-4-methoxy-2,4-cyclohexadiene-1-acetonitrile, 9CI. 3,5-Dibromo-1-cyanomethyl-4-methoxy-3,5-cyclohexadiene-1,2-diol. Zamamistatin $\text{C}_9\text{H}_9\text{Br}_2\text{NO}_3$ 338.983

Structure of Zamamistatin revised twice; finally shown to be the same as Aerophysinin 1 in 2008.


(+)-form [28656-91-9]Constit. of the sponges *Aplysina aerophoba* (*Verongia aerophoba*), *Aplysina archeri*, *Aiolochoira crassa*, *Verongula rigida*, *Psammaphysilla purpurea*, *Ianthella* sp. and others. Cytotoxic and antibacterial agent. Protein-tyrosine kinase inhibitor. Algicide, molluscicide. Sol.MeOH, Et₂O, Me₂CO; poorly sol. hexane. Mp 120-121° (112-113°). $[\alpha]_{\text{D}}^{20} +193$ (c, 0.63 in Me₂CO). $[\alpha]_{\text{D}}^{20} +185$ (c, 0.17 in MeOH). λ_{max} 284 (ε 4910) (MeOH) (Derep). λ_{max} 231 (ε 3220); 284 (ε 4910) (MeOH) (Berdy).**►GU4735000****Di-Ac:**Cryst. (C₆H₆/petrol). Mp 114°. $[\alpha]_{\text{D}}^{20} +218$ (CHCl₃).**(-)-form**Constit. of the sponges *Ianthella ardis*, *Verongula gigantea*, *Suberea creba*, *Psammaphysilla arabica* and *Pseudoceratina crassa*. Mp 116-117°. $[\alpha]_{\text{D}}^{20} -176$ (c, 0.125 in MeOH). $[\alpha]_{\text{D}}^{20} -198$ (c, 0.5 in Me₂CO). λ_{max} 284 (ε 4910) (MeOH) (Derep).**(±)-form**Constit. of *Verongula gigantea* and *Aiolochoira crassa*. Mp 153-154°. λ_{max} 284 (ε 4910) (MeOH) (Derep).

[55057-73-3, 55057-74-4, 66141-25-1, 30951-40-7]

Fulmor, W. et al., *Tet. Lett.*, 1970, 4551 (ir, pmr, cd, abs config)Cosulich, D.B. et al., *Chem. Comm.*, 1971, 397 (cryst struct, abs config)Mazzarella, L. et al., *Gazz. Chim. Ital.*, 1972, **102**, 391 (cryst struct, abs config)Fattorusso, E. et al., *J.C.S. Perkin 1*, 1972, 16 (isol, uv, ir, pmr, ms, struct)Andersen, R.J. et al., *J.A.C.S.*, 1975, **97**, 936 (synth)Chang, C.W.J. et al., *Tet. Lett.*, 1977, 4005 (isol)Makariev, T.N. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1981, **68**, 481 (isol, uv, ir, ms)Koulman, A. et al., *J. Nat. Prod.*, 1996, **59**, 591 (activity)Kita, M. et al., *Tet. Lett.*, 2008, **49**, 5383-5384 (Zamamistatin)**Aerothionin**

A-158

[28714-26-3]

 $\text{C}_{24}\text{H}_{26}\text{Br}_4\text{N}_4\text{O}_8$ 818.107Numbering systems vary. Isol. from *Aplysina aerophoba* (*Verongia aerophoba*), *Aplysina fistularis* and *Aplysina thiona*. Also from *Pseudoceratina durissima*, *Psammaphysilla purpurea* and the crinoid *Himerometra magnipinna*. Plates (Me₂CO/C₆H₆). Mp 134-137° dec. $[\alpha]_{\text{D}}^{20} +252$ (Me₂CO). λ_{max} 234 (ε 14450); 284 (ε 13490) (EtOH) (Derep). λ_{max} 284 (ε 12660) (MeOH) (Berdy).**Di-Ac:**Needles (Me₂CO). Mp 206-208°. $[\alpha]_{\text{D}}^{20} +236$ (CHCl₃).**2''-Hydroxy: 11-Hydroxyaerothionin**

[73622-27-2]

 $\text{C}_{24}\text{H}_{26}\text{Br}_4\text{N}_4\text{O}_9$ 834.107Constit. of *Aplysina caissara*, *Aplysina**lacunosa* and *Pseudoceratina durissima*. Glass. $[\alpha]_{\text{D}}^{20} +189$ (c, 0.15 in MeOH). λ_{max} 233 (ε 19780); 284 (ε 11100); 295 (ε 18900) (MeOH) (Berdy).**2'',3''-Dihydroxy: Dihydroxyaerothionin**

[122759-72-2]

 $\text{C}_{24}\text{H}_{26}\text{Br}_4\text{N}_4\text{O}_{10}$ 850.106Metab. of *Verongula rigida*. Powder. Mp 162-164°. $[\alpha]_{\text{D}}^{25} -64.2$ (c, 0.1 in MeOH).**2''-Oxo: 11-Oxo-aerothionin**

[73622-23-8]

 $\text{C}_{24}\text{H}_{24}\text{Br}_4\text{N}_4\text{O}_9$ 832.091Metab. from the Caribbean sponge *Aplysina lacunosa* and from *Verongia cavernicola*. Exhibits pronounced and selective antitumour activity against human colon (HCT116) cell line. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 174.6-176.6° dec. $[\alpha]_{\text{D}}^{25} +181.15$ (c, 2.17 in DMSO). λ_{max} 262 (ε 11600); 284 (ε 11500) (MeOH) (Berdy).**2''-Oxo, 3''R-hydroxy: 12R-Hydroxy-11-oxo-aerothionin**

[157544-68-8]

 $\text{C}_{24}\text{H}_{24}\text{Br}_4\text{N}_4\text{O}_{10}$ 848.09Metab. from the Caribbean sponge *Aplysina fistularis* forma *fulva*. $[\alpha]_{\text{D}}^{25} +160.7$.**2''-Oxo, 3''S-hydroxy: 12S-Hydroxy-11-oxo-aerothionin**

[157497-57-9]

 $\text{C}_{24}\text{H}_{24}\text{Br}_4\text{N}_4\text{O}_{10}$ 848.09Metab. from the sponge *Aplysina fistularis* forma *fulva*. $[\alpha]_{\text{D}}^{25} +152.5$.**Homologue: Homo-aerothionin**

[34232-66-1]

[35036-48-7]

 $\text{C}_{25}\text{H}_{28}\text{Br}_4\text{N}_4\text{O}_8$ 832.134Constit. of the sponges *Aplysina aerophoba*, *Verongia thiona* and *Verongia cavernicola*. Amorph. solid. Mp 166-167° (as di-Ac). $[\alpha]_{\text{D}}^{20} +191.5$ (CHCl₃) (di-Ac). Has a C₅ bridging chain instead of C₄. λ_{max} 289 (ε 13348) (MeOH) (Berdy).**Homologue, 2''ξ-hydroxy: 11-Hydroxy-homo-aerothionin. Caissarin C** $\text{C}_{25}\text{H}_{28}\text{Br}_4\text{N}_4\text{O}_9$ 848.134Constit. of *Aplysina caissara*. Amorph. solid. $[\alpha]_{\text{D}}^{28} +175$ (c, 0.002 in MeOH).**Homologue, 3''-hydroxy: 12-Hydroxy-homo-aerothionin**

[938464-19-8]


 $\text{C}_{25}\text{H}_{28}\text{Br}_4\text{N}_4\text{O}_9$ 848.134Isol. from the crinoid *Himerometra magnipinna*. Yellow powder. $[\alpha]_{\text{D}}^{20} +106$ (c, 0.39 in MeOH). λ_{max} 231 (log ϵ 4.33); 281 (log ϵ 3.99) (MeOH).**Homologue, 2''-oxo: 11-Oxohomo-aerothionin**

[191112-17-1]

 $\text{C}_{25}\text{H}_{26}\text{Br}_4\text{N}_4\text{O}_9$ 846.118Constit. of *Aplysina cavernicola*. λ_{max} 232 (ε 19000); 284 (10500) (MeOH).Fattorusso, E. et al., *Chem. Comm.*, 1970, 752-753 (uv, ir, pmr, struct)Fattorusso, E. et al., *Gazz. Chim. Ital.*, 1971, **101**, 61-63 (Homo-aerothionin)Moody, K. et al., *J.C.S. Perkin 1*, 1972, 18-24 (isol, uv, ir, pmr, struct, Homo-aerothionin)Forrester, A.R. et al., *Annalen*, 1978, 66-73 (synth)


- McMillan, J.A. *et al.*, *Tet. Lett.*, 1981, **22**, 39-42 (*cryst struct, uv, pmr, cd, abs config*)
- Nishiyama, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1985, **58**, 3453-3456 (*synth*)
- Okamoto, K.T. *et al.*, *Tet. Lett.*, 1987, **28**, 4969-4972 (*biosynth*)
- Gunasekera, M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 753-756 (*Dihydroxyaerothionin*)
- Kernan, M.R. *et al.*, *J. Nat. Prod.*, 1990, **53**, 615-622 (*11-Hydroxyaerothionin*)
- Acosta, A.L. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1007-1012 (*11-Oxo-aerothionin*)
- Ciminiello, P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 705-712 (*12-Hydroxy-11-oxo-aerothionin*)
- Ciminiello, P. *et al.*, *Tetrahedron*, 1997, **53**, 6565-6572 (*11-Oxohomoaerothionin*)
- Wasserman, H.H. *et al.*, *J.O.C.*, 1998, **63**, 5581-5586 (*synth*)
- De Lira, T.O. *et al.*, *J. Braz. Chem. Soc.*, 2006, **17**, 1233-1240 (*Caissarine C*)
- Ogamino, T. *et al.*, *Tet. Lett.*, 2006, **47**, 727-731 (*synth, abs config*)
- Shao, N. *et al.*, *J. Nat. Prod.*, 2007, **70**, 869-871 (*12-Hydroxyhomoaerothionin*)

Aeruginosamide **A-159**
[241483-64-7]



- $C_{30}H_{48}N_4O_4S$ 560.8
Modified peptide antibiotic. Isol. from *Microcystis aeruginosa*. Moderate cytotoxic agent. Pale green oil. $[\alpha]_D -71.4$ (c, 0.01 in $CHCl_3$).
- Lawton, L.A. *et al.*, *J.O.C.*, 1999, **64**, 5329-5332 (*isol, pmr, cmr, ir, ms*)
- Chen, Z. *et al.*, *New J. Chem.*, 2006, **30**, 518-520 (*synth*)


Aeruginosin 98A **A-160**
[167172-80-7]



- $C_{29}H_{45}ClN_6O_9S$ 689.228
Isol. from the blue-green alga *Microcystis aeruginosa* NIES98. Inhibitor of trypsin, thrombin and serine proteases. Amorph. powder. Sol. MeOH, H_2O . $[\alpha]_D -7.6$ (c, 0.2 in H_2O). λ_{max} 279 (ε 1472) (H_2O) (Berdy).
- 5'-Chloro: Aeruginosin 101**
 $C_{29}H_{44}Cl_2N_6O_9S$ 723.673
Isol. from *Microcystis aeruginosa*. Protease inhibitor. $[\alpha]_D -11$ (c, 0.5 in MeOH aq.). λ_{max} 290 (ε 1810) (MeOH).
- 3'-Dechloro: Aeruginosin 98B**
[167228-01-5]
 $C_{29}H_{46}N_6O_9S$ 654.783


- Isol. from *Microcystis aeruginosa* NIES98. Inhibitor of trypsin, thrombin, plasmin and serine proteases. Amorph. powder. Sol. MeOH, H_2O . $[\alpha]_D -5.2$ (c, 0.2 in H_2O). λ_{max} 276 (ε 1177) (H_2O) (Berdy).
- 3'-Dechloro, 3'-bromo: Aeruginosin 98C**
[167172-73-8]
 $C_{29}H_{45}BrN_6O_9S$ 733.679
Isol. from *Microcystis aeruginosa* NIES98. Serine protease inhibitor. Powder. Sol. MeOH, H_2O . $[\alpha]_D -13$ (c, 0.25 in H_2O). λ_{max} 281 (ε 860) (H_2O).
- Murakami, M. *et al.*, *Tet. Lett.*, 1995, **36**, 2785-2788 (*isol, pmr, cmr*)
- Ishida, K. *et al.*, *Tetrahedron*, 1999, **55**, 10971-10988 (*isol, abs config*)
- Ersmark, K. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1202-1223 (*rev*)

Aeruginosin 205A **A-161**
[184015-81-4]



- $C_{34}H_{53}ClN_6O_{12}S$ 805.344
Glycopeptide antibiotic. Structure probably requires revision. Isol. from freshwater *Oscillatoria agardhii* NIES 205. Serine protease inhibitor. Microcryst. Sol. MeOH, butanol; fairly sol. H_2O ; poorly sol. Et_2O , hexane. $[\alpha]_D^{20} +17.7$ (c, 0.1 in MeOH). λ_{max} 258 (ε 520) (MeOH).
- 2,2',3-Triepimer: Aeruginosin 205B**
[187521-09-1]
 $C_{34}H_{53}ClN_6O_{12}S$ 805.344
Isol. from freshwater *Oscillatoria agardhii* NIES 205. Serine protease inhibitor. Sol. MeOH, butanol; fairly sol. H_2O ; poorly sol. Et_2O , hexane. $[\alpha]_D^{20} +40.3$ (c, 0.1 in MeOH). λ_{max} 258 (ε 570) (MeOH).
- Shin, H.J. *et al.*, *J.O.C.*, 1997, **62**, 1810-1813 (*isol, uv, pmr, cmr, ms*)
- Valls, N. *et al.*, *Tet. Lett.*, 2006, **47**, 3701-3705 (*struct*)
- Ersmark, K. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1202-1223 (*rev*)

Aeruginosin B **A-162**
7-Amino-1-carboxy-5-methyl-3-sulfophenazinium hydroxide inner salt, 9CI. 7-Amino-1-carboxy-5-methyl-3-sulfophenazinium betaine. Eruginosin B
[6508-65-2]




- $C_{14}H_{11}N_3O_5S$ 333.324
Pigment from *Pseudomonas aeruginosa*. Red cryst. (aq. acid). Sol. H_2O . Dec. on heating without melting.
- Herbert, R.B. *et al.*, *J.C.S.(C)*, 1969, 2517 (*struct, ir, uv*)
- Bentley, R.K. *et al.*, *J.C.S.(C)*, 1970, 2447 (*synth*)

Aestivine† **A-163**

- Estivine*
[1356-93-0]
Struct. unknown
 $C_{17}H_{21}NO_4$ 303.357
Amaryllidaceae alkaloid. Isol. from the bulbs of *Leucojum aestivum* (Amaryllidaceae). Cryst. + $1H_2O$ (EtOH). Mp 194-195°. $[\alpha]_D -151$ (c, 0.1 in EtOH).
- Hydrochloride:*
Cryst. + $1/2 H_2O$. Mp 205-207°.
- Hydrobromide:* Mp 153-154°.
- Proskurnina, N.F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1963, **33**, 1643-1644 (*isol*)

Aestivophoenin A **A-164**
[171864-91-8]



- $C_{31}H_{32}N_2O_7$ 544.603
Phenazine antibiotic. Prod. by *Streptomyces purpeofuscus*. Neuronal cell protecting agent, neuroprotectant, lipid peroxidation inhibitor. Orange powder. Sol. MeOH, EtOAc, Me_2CO . Mp 59-61°. $[\alpha]_D^{20} -106.3$ (c, 0.005 in MeOH). λ_{max} 229 (ε 28900); 245 (ε 27400); 296 (ε 26000); 367 (ε 4000); 490 (ε 11700) (MeOH) (Berdy).
- Shin-ya, K. *et al.*, *J. Antibiot.*, 1995, **48**, 1378 (*isol, pmr, cmr, uv, ir*)

Aestivophoenin B **A-165**

- [171864-92-9]
As Aestivophoenin A, A-164 with
R = $-CH_2CH=C(CH_3)_2$
 $C_{36}H_{40}N_2O_7$ 612.721
Phenazine antibiotic. Prod. by *Streptomyces purpeofuscus*. Neuronal cell protecting agent, neuroprotectant, lipid peroxidation inhibitor. Orange powder. Sol. MeOH, Me_2CO , EtOAc. Mp 63-64°. $[\alpha]_D -139.8$ (c, 0.006 in MeOH). λ_{max} 232 (ε 26900); 245 (ε 25800); 298 (ε 19000); 368 (ε 3300); 495 (ε 9700) (MeOH) (Berdy).
- Debenzoyl: Aestivophoenin C*
 $C_{29}H_{36}N_2O_6$ 508.613
Prod. by *Streptomyces purpeofuscus*. Neuronal cell protecting agent, anti-oxidant. Yellow powder. Mp 61-62°.

λ_{\max} 247 (ϵ 24600); 328 (ϵ 2000); 453 (ϵ 4100) (MeOH).

Shin-ya, K. *et al.*, *J. Antibiot.*, 1995, **48**, 1378


(*isol*, *pmr*, *cmr*, *uv*, *ir*)

Kunigami, T. *et al.*, *J. Antibiot.*, 1998, **51**, 880-882 (*isol*, *uv*, *pmr*, *cmr*)

Affinine†

A-166

17-Hydroxyvobasan-3-one, 9CI
[2134-82-9]



Absolute
Configuration

$C_{20}H_{24}N_2O_2$ 324.422

Alkaloid from *Peschiera affinis*, *Peschiera laeta* and *Tabernaemontana psychotrifolia* (Apocynaceae). Cryst. (MeOH/EtOAc, CH_2Cl_2 /heptane, or EtOH). Mp 273-275° (264-265°) dec. $[\alpha]_D^{24}$ -180 (c, 0.78 in EtOH). λ_{\max} 236 (sh) (ϵ 13200); 320 (ϵ 13400) (EtOH).

Hydrochloride:

Cryst. (MeOH/EtOAc). Mp 267-269° dec. $[\alpha]_D^{25}$ -105.4 (c, 0.5 in MeOH).

O-Ac:

Cryst. as solvate (Me₂CO). Mp 95-115°.

10-Methoxy: **Pelirine**

[30435-26-8]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from the roots of *Rauwolfia perakensis* (Apocynaceae). Pale yellow plates (MeOH aq.). Mp 130-131°. $[\alpha]_D^{22}$ -121 (c, 1 in EtOH). The 16-config. was uncertain until 1987.

10-Methoxy, sulfate (1:2):

Cryst. + 1/2 H₂O. Mp 223-224° dec.

10-Methoxy, picrate:

Yellow needles (EtOH). Mp 159-159.5°.

16-Epimer: **16-Epiaffinine**

[38990-09-9]

$C_{20}H_{24}N_2O_2$ 324.422

Alkaloid from the stem bark of *Pleio-carpa talbotii* (Apocynaceae). Cryst. (Et₂O/pentane). Mp 152-154°. $[\alpha]_D^{23}$ -190 (c, 0.95 in CHCl₃). pK_a 6.9 (methylcellosolve).

16-Epimer, picrate:

Cryst. (MeOH). Mp 187-189°.

16-Epimer, N-Me: **16-Epi-N-methylaffi-**

nine. N-Methyl-16-epiaffinine

[58262-66-1]

$C_{21}H_{26}N_2O_2$ 338.449

Alkaloid from *Tabernaemontana acedens* (Apocynaceae). Plates (MeOH). Mp 208-210° dec. $[\alpha]_D^{20}$ -243 (c, 0.05 in CHCl₃).

16-Epimer, Me ether: **Hystrixnine**. 16-Epi-17-O-methylaffinine

$C_{21}H_{26}N_2O_2$ 338.449

Alkaloid from the root bark of *Tabernaemontana hystrix*. Amorph. solid. $[\alpha]_D^{25}$ -100 (c, 0.66 in MeOH). λ_{\max} 223 (ϵ 42570); 282 (ϵ 6290) (MeOH).

16-Epimer, 10-hydroxy: **16-Epi-10-hydroxyaffinine**. 10-Hydroxy-16-epiaffinine

[82513-70-0]

$C_{20}H_{24}N_2O_3$ 340.421

Alkaloid from *Hunteria zeylanica* (Apocynaceae). Amorph. $[\alpha]_D$ -122 (c, 0.46 in CHCl₃). λ_{\max} 212 (log ϵ 4.46); 228 (sh) (log ϵ 4.29); 277 (sh); 328 (log ϵ 4.17) (MeOH).

Kiang, A.K. *et al.*, *J.C.S.*, 1960, 1394-1398 (*Pelirine*, *isol*, *uv*)

Weisbach, J.A. *et al.*, *J. Pharm. Sci.*, 1963, **52**, 350-353 (*isol*, *ir*)

Cava, M.P. *et al.*, *Chem. Ind. (London)*, 1964, 1193-1194 (*struct*, *pmr*)

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1971, **49**, 307-316 (*isol*, *uv*, *ir*, *pmr*, *struct*)

Naranjo, J. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 752-771 (*16-Epiaffinine*)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 929-933 (*Pelirine*, *cd*)

Achenbach, H. *et al.*, *Chem. Ber.*, 1975, **108**, 3842-3854 (*16-Epi-N-methylaffinine*)

Voticky, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 1403-1406 (*isol*, *uv*, *ir*, *ms*)

Lavaud, C. *et al.*, *Phytochemistry*, 1982, **21**, 445-447 (*16-Epi-10-hydroxyaffinine*)


Wan, A.S.C. *et al.*, *Heterocycles*, 1987, **26**, 1211-1214 (*Pelirine*, *cryst struct*, *abs config*)

Monnerat, C.S. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 1331-1335 (*Hystrixnine*)

Affinisine oxindole

A-167

[701950-31-4]



Absolute
Configuration

$C_{20}H_{24}N_2O_2$ 324.422


Alkaloid from the leaves of *Alstonia angustifolia* var. *latifolia*. Light yellow oil. $[\alpha]_D$ -70 (c, 0.06 in CHCl₃). λ_{\max} 211 (log ϵ 4.31); 233 (log ϵ 3.82); 255 (log ϵ 3.87); 290 (log ϵ 3.35) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 2004, **65**, 603-608 (*isol*, *pmr*, *cmr*)

Aflastatin A

A-168

[179729-59-0]



Absolute
Configuration

$C_{62}H_{115}NO_{24}$ 1258.583

Abs. config. revised in 2007. Prod. by *Streptomyces* sp. MRI 142. Inhibits production of Aflatoxin by aflatoxigenic fungi. Active against *Candida*, *Trichophyton* and *Staphylococcus* infections. Powder. Sol. DMSO; fairly sol. MeOH; poorly sol. H₂O, CHCl₃, hexane, EtOAc, Me₂CO. $[\alpha]_D^{19}$ -2.6 (c, 0.5 in DMSO). λ_{\max} 247 (ϵ 11000); 299 (ϵ 6200) (MeOH aq.). λ_{\max} 237 (ϵ 7900); 314 (ϵ 7300) (MeOH/HCl). λ_{\max} 247 (ϵ 11000); 299 (ϵ 6200)

(MeOH/NaOH).

▶ LD₅₀ (mus, ipr) 6.17 mg/kg, LD₅₀ (mus, orl) 2000 mg/kg.

N-De-Me: **Aflastatin B**

$C_{61}H_{113}NO_{24}$ 1244.556

Prod. by *Streptomyces* MRI142. $[\alpha]_D^{23}$ -0.6 (c, 0.16 in DMSO). λ_{\max} 246 (ϵ 10600); 299 (ϵ 6300) (MeOH aq.). λ_{\max} 237 (ϵ 8000); 314 (ϵ 7800) (MeOH/HCl). λ_{\max} 299 (ϵ 6300) (MeOH/NaOH).

Sakuda, S. *et al.*, *J.A.C.S.*, 1996, **118**, 7855-7856 (*biosynth*, *struct*)

Ono, M. *et al.*, *J. Antibiot.*, 1997, **50**, 111-118; 1998, **51**, 1019-1029 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *props*, *ms*)


Ikedo, H. *et al.*, *J.O.C.*, 2000, **65**, 438-444 (*abs config*)

Sakuda, S. *et al.*, *Tet. Lett.*, 2007, **48**, 2527-2531 (*abs config*)

Aflatrem

A-169

α, α -Dimethylallylpaspalinine
[70553-75-2]



$C_{32}H_{39}NO_4$ 501.664

Isol. from *Aspergillus flavus* and *Claviceps paspali*. Tremorgenic mycotoxin. Needles (Me₂CO/Et₂O). Sol. MeOH, CHCl₃. Mp 222-224°. λ_{\max} 231 (ϵ 27700); 250 (ϵ 10000); 282 (ϵ 9000) (MeOH) (Berdy).

▶ Toxic.

Wilson, B.J. *et al.*, *Science (Washington, D.C.)*, 1964, **144**, 177 (*isol*)

Gallagher, R.T. *et al.*, *Tet. Lett.*, 1980, 239 (*struct*, *uv*, *ir*, *pmr*, *cmr*)


Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 293 (*isol*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 410

Aflatrem B

A-170

β -Aflatrem
[144446-23-1]




$C_{32}H_{39}NO_4$ 501.664

Prod. by *Aspergillus flavus*, *Aspergillus parasiticus* and *Aspergillus subulivaceus*. Mycotoxin, exhibits insecticidal activity. Yellow cryst. Mp 188-190°. $[\alpha]_D$ +77.9 (c, 0.011 in CHCl₃). λ_{\max} 233 (ϵ 18500); 264 (ϵ 6300); 296 (ϵ 2700) (MeOH) (Berdy).

TePaske, M.R. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1080 (*isol*, *uv*, *pmr*, *cmr*, *ms*, *struct*)

Aflavazole

[133401-09-9]


C₂₈H₃₅NO₂ 417.59

Closely related to Aflavinine, A-172. Metab. of *Aspergillus flavus* and *Aspergillus parasiticus*. Exhibits pronounced insect antifeedant activity. Light yellow cryst. Mp 156-160° dec. [α]_D +2.8 (c, 0.35 in MeOH). λ_{max} 219 (ε 15300); 243 (ε 17300); 263 (ε 7500); 297 (ε 7600); 327 (ε 1300); 341 (ε 1600) (MeOH) (Derep).

TePaske, M.R. *et al.*, *J.O.C.*, 1990, **55**, 5299 (isol, pmr, cmr)

Aflavinine

[74328-59-9]

C₂₈H₃₉NO 405.622

Indolic diterpenoid antibiotic. Related to biosynthetic intermeds. in the penitrem series. Different numbering systems in use. Isol. from *Aspergillus flavus*, *Eupenicillium crustaceum* and *Eupenicillium molle*. Tremorgenic mycotoxin, shows insecticidal activity. Needles + 1EtOAc (EtOAc). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 102-104°. λ_{max} 226 (ε 27000); 284 (ε 5300); 292 (ε 4500) (MeOH) (Berdy).

10β,11α-Dihydro, 24,25-didehydro-10,11-dihydroaflavinine [124693-66-9]

C₂₈H₃₉NO 405.622

Metab. from the sclerotia of *Aspergillus tubingensis*. Also shown by hplc to be present in the sclerotia of *Aspergillus flavus* and *Aspergillus parasiticus*. Mp 192-194°. [α]_D -1.2 (c, 0.5 in CHCl₃).

20β-Hydroxy: 20-Hydroxyaflavinine

[116865-08-8]

C₂₈H₃₉NO₂ 421.622

From *Aspergillus flavus*. Tremorgenic mycotoxin, shows antifeedant and insecticidal props. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 174-176° dec. (161-162°). [α]_D +64 (c, 0.08 in MeOH). [α]_D +23.8 (c, 0.56 in MeOH).

A-171

λ_{max} 224 (ε 14900); 283 (ε 2760); 290 (ε 2520) (MeOH) (Derep).

20α-Hydroxy, 10β,11α-dihydro, 24,25-didehydro: 24,25-Dehydro-10,11-dihydro-20-epi-20-hydroxyaflavinine

C₂₈H₃₉NO₂ 421.622

Metab. from the sclerotia of *Aspergillus tubingensis*. Also shown by hplc to be present in the sclerotia of *Aspergillus flavus* and *Aspergillus parasiticus*. Mp 79-82°. [α]_D -5.6 (c, 1.06 in CHCl₃). λ_{max} 226 (ε 16400); 284 (ε 2290); 292 (ε 2100) (MeOH) (Berdy).

20β-Hydroxy, 10β,11α-dihydro, 11,12-didehydro: 11,12-Didehydro-10,11-dihydro-20-hydroxyaflavinine

[116865-10-2]

C₂₈H₃₉NO₂ 421.622

From *Aspergillus flavus*. Mycotoxin showing insecticidal and antifeedant props. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 276-278°. [α]_D +1.7 (c, 0.1 in MeOH). λ_{max} 224 (ε 20030); 281 (ε 3600); 285 (ε 3380) (MeOH) (Berdy).

20β-Hydroxy, 10β,11α-dihydro, 24,25-didehydro: 24,25-Dehydro-10,11-dihydro-20-hydroxyaflavinine. Monohydroxyisoaflavinine

C₂₈H₃₉NO₂ 421.622

Metab. from *Aspergillus flavus*. Insecticide, tremorgenic agent. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 146-148° Mp 259-262°. [α]_D +0.9 (c, 0.34 in MeOH). [α]_D²⁴ +56 (c, 0.10 in MeOH). Identical structs. assigned to Dehydrodihydro-20-hydroxyaflavinine and Monohydroxyisoaflavinine (Nozawa *et al*) but very different props. reported (lower Mp and higher opt. rotn. refer to Monohydroxyisoaflavinine). λ_{max} 226; 227 (ε 26260); 284 (ε 6060); 291 (ε 5220) (MeOH) (Berdy). λ_{max} 226 (ε 21380); 283 (ε 4266); 291 (ε 3990) (EtOH) (Berdy).

24-Hydroxy, 10,11-dihydro: 10,11-Dihydro-24-hydroxyaflavinine

[171569-81-6]

C₂₈H₄₁NO₂ 423.637

Prod. by *Eupenicillium crustaceum*. Shows insecticidal props. Powder. [α]_D -22 (CHCl₃).

20β,25-Dihydroxy: 20,25-Dihydroxyaflavinine

[76410-56-5]

C₂₈H₃₉NO₃ 437.621

From *Aspergillus flavus*. Tremorgenic mycotoxin, shows insecticidal and antifeedant props. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 254-256°. [α]_D +22.9 (c, 0.5 in MeOH). λ_{max} 224 (ε 20190); 283 (ε 3850); 291 (ε 3670) (MeOH) (Berdy).

13-Oxo, 10β,11α-dihydro, 24,25-didehydro: 24,25-Dehydro-10,11-dihydro-13-oxoaflavinine

[124693-67-0]

C₂₈H₃₇NO₂ 419.606

From *Aspergillus tubingensis*. Shows insecticidal props. Mp 138-141°. [α]_D -13.5 (c, 0.72 in CHCl₃). λ_{max} 224 (ε 18200); 267 (ε 1200); 273 (ε 1400); 283 (ε 2100); 291 (ε 1750) (MeOH) (Berdy).

Gallagher, R.T. *et al.*, *Tet. Lett.*, 1980, 243 (isol, pmr, cmr, cryst struct)

Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 293 (deriv)

Danishesky, S. *et al.*, *J.A.C.S.*, 1985, **107**, 2474 (synth)

Gloer, J.B. *et al.*, *J.O.C.*, 1988, **53**, 5457 (isol, derivs)


Nozawa, K. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 626 (Monohydroxyisoaflavinine, isol, uv, pmr, cmr, ms, cryst struct)

TePaske, M.R. *et al.*, *Tetrahedron*, 1989, **45**, 4961 (derivs)

Wang, H.J. *et al.*, *Appl. Environ. Microbiol.*, 1995, **61**, 4429-4435 (10,13-Dihydro-24-hydroxyaflavinine)

Africanine

[107019-97-6]


C₂₀H₁₇NO₆ 367.357

Alkaloid from *Rupicapnos africana* (Papaveraceae). Cryst. (MeOH). Mp 237°. [α]_D²⁵ +22 (c, 0.322 in CHCl₃).

Castedo, L. *et al.*, *Heterocycles*, 1986, **24**, 2781 (uv, ir, pmr, ms, struct)

Afrocuarine

[55599-19-4]



Absolute Configuration


C₃₉H₄₂N₄O₂⁺ 582.787

Alkaloid from *Strychnos usambarensis* (Loganiaceae). Weak neuromuscular blocking agent (activity ca. 0.05 × Tubocuarine).

Caprasse, M. *et al.*, *Planta Med.*, 1984, **50**, 131-133 (isol, uv, ir, pmr, ms, cd, struct)

Agaricone

[99280-73-6]

C₁₀H₁₁N₃O 189.216

Isol. from the toadstool *Agaricus xanthoderma*. Red cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 115° (dec.).

A-173

A-172

A-174

A-175

Formed on injury of the fungus by air or enzyme oxidn. of Leucoagaricone, L-154. λ_{\max} 242 (log ϵ 3.74); 427 (log ϵ 4.19) (MeOH).

Hilbig, S. *et al.*, *Angew. Chem., Int. Ed.*, 1985, **24**, 1063-1065 (*isol, synth, uv, pmr, ms*)

Agel 452


A-176

N-(20-Amino-4-hydroxy-4,8,12,17-tetrazaeicos-1-yl)-4-hydroxybenzamide, 9CI.

AG 452

[128549-98-4]

[128550-23-2 (trifluoroacetate salt)]



$C_{23}H_{44}N_6O_3$ 452.639

See also *Agelenopsis aperta* Hydroxybenzoylpentamine toxins, H-413 for related toxins. Alkaloid from the venom of the spider *Agelenopsis aperta*. λ_{\max} 251 (log ϵ 4.12) (H_2O) (trifluoroacetate).

4-Deoxy, 2,5-dihydroxy: **Agel 468**. AG 468

[128549-99-5]

 $C_{23}H_{44}N_6O_4$ 468.638

Alkaloid from the venom of the spider *Agelenopsis aperta*.

N,4-Dideoxy, 2,5-dihydroxy: **Agel 452a**.

AG 452a. HO 452

[133805-35-3]

 $C_{23}H_{44}N_6O_3$ 452.639

Alkaloid from the venom of the spiders *Agelenopsis aperta* and *Hololea curta*.

N¹⁷-De(aminopropyl), N¹⁷-(aminoimino-methyl), N-deoxy: **PB 421**

[333401-98-2]

 $C_{21}H_{39}N_7O_2$ 421.585

Alkaloid from the venom of the spider *Paracoelotes birulai*.

[128550-24-3]


Jasys, V.J. *et al.*, *J.A.C.S.*, 1990, **112**, 6696-6704 (*isol, synth, ms, struct*)

Chesnov, S. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 3295-3305; 2001, **84**, 2178-2197 (*PB 421, Agel 452a*)

Ageladine A

A-177

[643020-13-7]



$C_{10}H_7Br_2N_5$ 357.007

Isol. from *Agelas nakamura*. Antiangiogenic matrixmetalloproteinase inhibitor. Yellow powder (as di-TFA salt). λ_{\max} 227 (ϵ 15400); 250 (ϵ 9960); 381 (ϵ 12200)

(MeOH) (di-TFA salt).

Fujita, M. *et al.*, *J.A.C.S.*, 2003, **125**, 15700-15701 (*Ageladine A*)

Shengule, S.R. *et al.*, *Org. Lett.*, 2006, **8**, 4083-4084 (*synth*)


Meketa, M.L. *et al.*, *J.O.C.*, 2007, **72**, 4892-4899 (*synth*)

Meketa, M.L. *et al.*, *Tetrahedron*, 2007, **63**, 9112-9119 (*synth*)

Agelasidine A

A-178

2-[[[1-(1-Ethenyl-1,5,9-trimethyl-4,8-decadienyl)sulfonyl]ethyl]guanidine, 9CI



$C_{18}H_{33}N_3O_2S$ 355.544

(R)-form [915159-59-0]

Isol. from *Agelas clathrodes*. Oil. $[\alpha]_D^{20}$ -14.5 (c, 1.5 in MeOH).

(S)-form

Isol. from the Okinawan sea sponge *Agelas nakamura* and from a Pacific sponge *Agelas* sp. Shows antispasmodic and antibacterial activity. Unstable yellow oil; cryst. + $\frac{1}{3} H_2O$ (EtOAc) (as hydrochloride). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 108-108.5° (hydrochloride). $[\alpha]_D^{25}$ +19.1 (c, 1 in MeOH) (hydrochloride). λ_{\max} 227 (ϵ 5400); 265 (ϵ 1800) (EtOH) (Berdy). λ_{\max} 226 (ϵ 4000); 270 (ϵ 2000) (EtOH/HCl) (Berdy).

Capon, R.J. *et al.*, *J.A.C.S.*, 1984, **106**, 1819-1822 (*isol, uv, ir, pmr, cmr, ms*)

Nakamura, H. *et al.*, *J.O.C.*, 1985, **50**, 2494-2497 (*isol, uv, ir, pmr, cmr*)


Ichikawa, Y. *et al.*, *J.C.S. Perkin 1*, 1992, 1497-1500 (*synth*)

Medeiros, M.A. *et al.*, *Z. Naturforsch., C*, 2006, **61**, 472-476 (*R-form, isol, pmr, cmr*)

Agelasidine B

A-179

[96617-50-4]




$C_{23}H_{41}N_3O_2S$ 423.662

Isol. from the Okinawan sea sponge *Agelas nakamura*. Shows antispasmodic and antibacterial activity. Syrup (as hydrochloride). Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{25}$ -2.5 (c, 0.43 in MeOH).

Nakamura, H. *et al.*, *J.O.C.*, 1985, **50**, 2494 (*isol, uv, ir, pmr, cmr, struct*)

Agelasidine C

A-180



(+)-form

$C_{23}H_{41}N_3O_2S$ 423.662

(+)-form [96617-52-6]

Isol. from the sponges *Agelas nakamura* and *Agelas dispar*. Shows antispasmodic and antibacterial activity. Syrup (as hydrochloride). Sol. MeOH, $CHCl_3$. $[\alpha]_D^{25}$ +8.5 (c, 2.0 in MeOH) (hydrochloride).

(-)-form [139757-42-9]

Isol. from *Agelas clathrodes*. Oil. Sol. MeOH, $CHCl_3$. $[\alpha]_D^{29}$ -5.6 (c, 7.2 in MeOH).

20-Hydroxy: **Agelasidine D**

[139680-61-8]

 $C_{23}H_{41}N_3O_3S$ 439.661

Isol. from *Agelas clathrodes*. Oil. Sol. MeOH, $CHCl_3$. $[\alpha]_D^{29}$ -3.6 (c, 2.75 in MeOH).

[96617-52-6]

Nakamura, H. *et al.*, *J.O.C.*, 1985, **50**, 2494-2498 (*isol, uv, ir, pmr, cmr, struct*)


Asao, K. *et al.*, *Chem. Lett.*, 1989, 1813-1814 (*synth*)

Morales, J.J. *et al.*, *J. Nat. Prod.*, 1992, **55**, 389-394 (*isol*)

Agelasimine A

A-181

[114216-85-2]



$C_{27}H_{43}N_5O$ 453.67

Metab. from the marine sponge *Agelas mauritiana*. Exhibits cytotoxicity, inhibition of adenosine transfer into rabbit erythrocytes, Ca^{2+} -channel antagonistic action and α_1 adrenergic blockade. Yellowish oil. Sol. MeOH, EtOH, $CHCl_3$. $[\alpha]_D^{25}$ +2.3 (MeOH). λ_{\max} 223 (ϵ 9700); 287 (ϵ 14700) (MeOH) (Berdy).

2',3'-Dihydro, 3'N-Me, N^{4'}-de-Me: **Agelasimine B**

[114216-87-4]

 $C_{27}H_{45}N_5O$ 455.685

Metab. from the marine sponge *Agelas mauritiana*. Exhibits cytotoxicity, inhibition of adenosine transfer into rabbit erythrocytes, Ca^{2+} -channel antagonistic action and α_1 adrenergic blockade. Viscous yellowish oil. Sol. MeOH, $CHCl_3$, EtOH. $[\alpha]_D^{25}$ +2.46 (MeOH). λ_{\max} 245 (ϵ 7100); 327 (ϵ 5400) (MeOH) (Berdy).

Fathi-Afshar, R. *et al.*, *Can. J. Chem.*, 1988,

66, 45 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ohba, M. *et al.*, *Tetrahedron*, 1997, **53**, 16977-16986 (*synth, abs config*)

Agelazine†

A-182

Struct. unknown

 $C_{26}H_{40}N_5^{\oplus}$ 422.635

Quaternary 9-methyladenine deriv. of an unidentified bicyclic diterpene. Isol. from the sponge *Agelas dispar*. Cryst. (butanone/THF) (as chloride). Mp 197-200°


dec. (chloride). Mp of chloride does not appear to correspond with any of the Agelasines of known struct. λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

Cullen, E. *et al.*, *Can. J. Chem.*, 1975, **53**, 1690-1691 (*isol, uv, ir, pmr, ms*)

Agelasine I

A-183

[205382-10-1]



$C_{26}H_{40}N_5O^{\oplus}$ 438.635


Isol. from a sponge *Agelas* sp. Amorph. solid (as chloride). $[\alpha]_D^{25}$ -2.5 (c, 0.2 in MeOH). λ_{\max} 272 (ϵ 8230) (MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 548-550 (*isol, uv, ir, pmr, cmr*)

Agelasine A[†]

A-184

[56271-74-0]



$C_{26}H_{40}N_5^{\oplus}$ 422.635

Constit. of the Okinawan sponge *Agelas nakamura*. ATPase inhibitor, spasmolytic agent, smooth muscle contractant. Cryst. (as chloride). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 173-174° (chloride). $[\alpha]_D^{25}$ -31.3 (c, 0.59 in MeOH). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

18-Hydroxy: Agelasine H

[205382-09-8]

$C_{26}H_{40}N_5O^{\oplus}$ 438.635

Constit. of an *Agelas* sp. Amorph. solid (as chloride). $[\alpha]_D^{25}$ -63.9 (c, 0.36 in MeOH). λ_{\max} 272 (ϵ 9015) (MeOH).

18-Hydroxy, 18-O-(2-pyrrolicarbonyl):**Ageline B**

[88874-28-6]

[88855-18-9, 88855-19-0]

$C_{31}H_{43}N_6O_2^{\oplus}$ 531.72

Constit. of sponge *Agelas* sp. Active against gram-positive bacteria. ATPase inhibitor. Smooth muscle contractor. Phytotoxin. Weak ichthyotoxin. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . λ_{\max}

210 (ϵ); 274 (ϵ) (EtOH) (Derep). λ_{\max} 268 (ϵ 8320) (MeOH) (Berdy). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10111) (EtOH) (Berdy). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Berdy).

18-Hydroxy, 18-O-(4-bromo-2-pyrrolicarbonyl): Agelasine G

[142808-37-5]

$C_{31}H_{42}BrN_6O_2^{\oplus}$ 610.616

Isol. from the Okinawan marine sponge *Agelas* sp. Exhibits cytotoxicity against murine lymphoma L1210 cells *in vitro*. $[\alpha]_D^{27}$ -85 (c, 0.02 in $CHCl_3$) (as chloride). λ_{\max} 212 (ϵ 24000); 272 (ϵ 19000) (MeOH) (Berdy).

8,9,10-Triepimer: Agelasine B

[92664-76-1]

$C_{26}H_{40}N_5^{\oplus}$ 422.635

Constit. of *Agelas nakamura*. ATPase inhibitor, spasmolytic agent. Cryst. (as chloride). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 167-170° (chloride). $[\alpha]_D^{25}$ -21.5 (c, 1.00 in MeOH). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

Cullen, E. *et al.*, *Can. J. Chem.*, 1975, **53**, 1690-1691 (*isol*)

Capon, R.J. *et al.*, *J.A.C.S.*, 1984, **106**, 1819-1822 (*Ageline B*)

Nakamura, H. *et al.*, *Tet. Lett.*, 1984, **25**, 2989-2992 (*uv, pmr, cmr, struct*)

Hideo, I. *et al.*, *Chem. Comm.*, 1985, 774-775 (*Ageline B, synth*)

Wu, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2495-2504 (*isol, pmr, cmr, uv, Agelasine B*)

Ishida, K. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 766-767 (*Agelasine G*)

Piers, E. *et al.*, *J.C.S. Perkin 1*, 1995, 963-966 (*synth*)


Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 548-550 (*Agelasine H*)

Agelasine C

A-185

[92664-78-3]

[92664-79-4 (chloride)]



$C_{26}H_{40}N_5^{\oplus}$ 422.635

Stereochem. revised in 2005. Isol. from the sponge *Agelas nakamura*. Shows antimicrobial activity. ATPase inhibitor. Spasmolytic agent. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 176-179° (as chloride). $[\alpha]_D^{25}$ -55.1 (c, 2.04 in MeOH) (chloride). $[\alpha]_D^{25}$ -212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

5,9-Diepimer: Epiagelasine C

[187825-31-6]

$C_{26}H_{40}N_5^{\oplus}$ 422.636

Constit. of sponge *Agelas mauritiana*.

Antifouling substance against macroalgae. Powder (as chloride). $[\alpha]_D^{25}$ +33.9 (c, 0.056 in MeOH) (chloride). Stereochem. revised in 2005. λ_{\max} 273 (ϵ 8140) (MeOH).

Nakamura, H. *et al.*, *Tet. Lett.*, 1984, **25**, 2989-2992 (*isol, uv, pmr, cmr*)

Wu, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2495-2504 (*isol, pmr, cmr, uv*)

Hattori, T. *et al.*, *J. Nat. Prod.*, 1997, **60**, 411-413 (*5-epimer*)


Marcos, I. *et al.*, *Tetrahedron*, 2005, **61**, 11672-11678 (*synth, abs config*)

Agelasine D

A-186

[92664-80-7]

[92664-81-8 (chloride)]



$C_{26}H_{40}N_5^{\oplus}$ 422.635

Isol. from the Okinawan sea sponge *Agelas nakamura*. Shows antimicrobial activity. Spasmolytic agent. Na/K ATPase inhibitor. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 175-176° (as chloride). $[\alpha]_D^{25}$ +10.4 (c, 1.1 in MeOH) (chloride). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

Nakamura, H. *et al.*, *Tet. Lett.*, 1984, **25**, 2989-2992 (*uv, pmr, cmr, struct*)

Wu, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2495-2504 (*isol, uv, pmr, cmr*)

Utenova, B.T. *et al.*, *Tet. Lett.*, 2004, **45**, 4233-4235 (*synth*)


Vik, A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 381-386 (*synth, activity*)

Agelasine E

A-187

[92599-00-3]

[92599-01-4 (chloride)]



$C_{26}H_{40}N_5^{\oplus}$ 422.635


Isol. from the Okinawan sea sponge *Agelas nakamura*. ATPase inhibitor, spasmolytic agent, shows antimicrobial props. Powder (as chloride). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 180-182° (chloride). $[\alpha]_D^{23}$ -17.1 (c, 1.88 in MeOH) (chloride). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ϵ 18100); 272 (ϵ 10000) (EtOH) (Derep).

Wu, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2495-2504 (*uv, pmr, cmr, ms, struct*)

Bakkestuen, A.K. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 1025-1033 (*synth*)

Agelaspongin

[122893-39-4]

Relative
ConfigurationC₁₁H₁₃N₅O₂ 247.256

CA numbering shown. Synthetic. Cryst. (MeOH). Mp 220° dec.

Hydrochloride:

Cryst. (EtOH). Mp 250° dec.

10,11-Dibromo: Dibromoagelaspongin


[122893-38-3]

C₁₁H₁₁Br₂N₅O₂ 405.048Isol. from an *Agelas* sp. Glucanase inhibitor. Yellow-green cryst. (MeOH/Me₂CO) (as hydrochloride). Mp 233-235° dec. (hydrochloride). λ_{max} 235 (ε 11250); 290 (ε 7420) (MeOH) (Berdy).**10,11-Dibromo, Me ether: Dibromoagelaspongin methyl ether**

[934842-17-8]

C₁₂H₁₃Br₂N₅O₂ 419.075Isol. from *Agelas dispar*.Fedoreyev, S.A. *et al.*, *Tetrahedron*, 1989, **45**, 3487-3492 (isol, pmr, cmr, ms, cryst struct)Pina, I.C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 613-617 (dibromo Me ether)Feldman, K.S. *et al.*, *J.A.C.S.*, 2008, **130**, 14964-14965 (dibromo, synth)**Agelastatin A**

[152406-28-5]

Absolute
ConfigurationC₁₂H₁₃BrN₄O₃ 341.164Alkaloid from the sponge *Agelas dendromorpha* and *Cymbastela* sp. Exhibits marked cytotoxicity; insecticide. Anti-neoplastic agent. Prevents metastatic cancer spread. [α]_D²⁰ -84.3 (c, 0.3 in EtOH) (as *N,N,O*-tri-Me). Not obt. completely pure. λ_{max} 203 (ε 12400); 232 (ε 8400); 279 (ε 11900) (EtOH) (Derep).***N-De-Me: Agelastatin D***

[201338-45-6]

C₁₁H₁₁BrN₄O₃ 327.137Alkaloid from the sponge *Cymbastela* sp. Insecticide. Solid. λ_{max} 227 (ε 7900); 277 (ε 10800) (MeOH).**2-Bromo: Agelastatin B**

[159903-65-8]

C₁₂H₁₂Br₂N₄O₃ 420.06Minor alkaloid from *Agelas dendromorpha*. [α]_D -84.4 (c, 0.49 in EtOH) (as *N,N,O*-tri-Me). Obt. as insep. mixt. with Agelastatin A but characterised as the tri-Me deriv.


A-188

6b-Hydroxy: Agelastatin C

[201338-44-5]

C₁₂H₁₃BrN₄O₄ 357.163Alkaloid from the sponge *Cymbastela* sp. Cytotoxic; insecticide. Solid. [α]_D -5 (c, 0.06 in MeOH). λ_{max} 228 (ε 7900); 278 (ε 11000) (MeOH).D'Ambrosio, M. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1895; 1996, **79**, 727 (*Agelastatins A,B*)Hong, T.W. *et al.*, *J. Nat. Prod.*, 1998, **61**, 158-161 (*Agelastatins C,D*)Stien, D. *et al.*, *J.A.C.S.*, 1999, **121**, 9574-9579 (synth)Feldman, K.S. *et al.*, *J.O.C.*, 2002, **67**, 7096-7109 (synth)Hale, K.J. *et al.*, *Org. Lett.*, 2003, **5**, 2927-2930 (synth)Domostoj, M.M. *et al.*, *Org. Lett.*, 2004, **6**, 2615-2618 (synth)Davis, F.A. *et al.*, *Org. Lett.*, 2005, **7**, 621-623 (synth)Trost, B.M. *et al.*, *J.A.C.S.*, 2006, **128**, 6054-6055 (synth)Ichikawa, Y. *et al.*, *Org. Lett.*, 2007, **9**, 2989-2992 (synth)Mason, C.K. *et al.*, *Mol. Cancer Ther.*, 2008, **7**, 548-558 (*agelastatin A, pharmacol*)**Ageliferin**

[117417-64-8]



A-190

C₂₂H₂₄Br₂N₁₀O₂ 620.305Alkaloid from the marine sponges *Agelas confifera*, *Agelas novaecaledoniae*, *Agelas cf. mauritiana* and *Xestospongia* sp.

Potent actomyosin ATPase activator.

Antifouling agent. Somatostatin antagonist. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D³³ +15.5 (c, 0.11 in MeOH). λ_{max} 204 (ε 25500); 220 (ε 27100); 267 (ε 25900); 393 (ε 300) (MeOH) (Derep).***1'-N-Me: 1'-N-Methylageliferin***

[175861-97-9]

C₂₃H₂₆Br₂N₁₀O₂ 634.332Alkaloid from the sponge *Astrosclera willeyana*. Pale yellow glass (as bis(trifluoroacetate)). [α]_D +40 (c, 0.01 in MeOH).***1,1'-Di-N-Me: N¹,N^{1'}-Dimethylageliferin***

[175861-98-0]

C₂₄H₂₈Br₂N₁₀O₂ 648.359Alkaloid from the sponge *Astrosclera willeyana*. Pale buff glass (as dihydrochloride). [α]_D +40 (c, 0.01 in MeOH).**2-Bromo: 2-Bromoageliferin**

[117417-67-1]

C₂₂H₂₃Br₃N₁₀O₂ 699.201Alkaloid from the sponges *Agelas confifera*, *Agelas cf. mauritiana* and *Astrosclera willeyana*. Potent actomyosin ATPase activator. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D³³ +8.8 (c, 0.08 in MeOH). λ_{max} 203 (ε 30600); 220 (ε 26300); 270 (ε 24500);396 (ε 700) (MeOH) (Derep). λ_{max} 204 (ε 15000); 222 (ε 12200); 274 (ε 12500); 396 (ε 700) (EtOH) (Berdy).**2-Bromo, 1'-N-Me: 1'-N-Methyl-2-bromoageliferin**

[175862-01-8]

C₂₃H₂₅Br₃N₁₀O₂ 713.228Alkaloid from the sponge *Astrosclera willeyana*. Off-white glass (as bis(trifluoroacetate)). [α]_D +50 (c, 0.01 in MeOH).**2'-Bromo, 1'-N-Me: 1'-N-Methyl-2'-bromoageliferin**

[175862-02-9]

C₂₃H₂₅Br₃N₁₀O₂ 713.228Alkaloid from the sponge *Astrosclera willeyana*. Off-white glass (as bis(trifluoroacetate)). [α]_D +40 (c, 0.01 in MeOH).**2,2'-Dibromo: 2,2'-Dibromoageliferin**

[117417-69-3]

C₂₂H₂₂Br₄N₁₀O₂ 778.098Alkaloid from the sponges *Agelas confifera*, *Agelas cf. mauritiana* and *Astrosclera willeyana*. Potent actomyosin ATPase activator. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D³³ +3 (c, 0.1 in MeOH). λ_{max} 204 (ε 12200); 220 (ε 9100); 278 (ε 8200); 394 (ε 300) (no solvent reported) (Derep). λ_{max} 202 (ε 24300); 214 (ε 24500); 274 (ε 25000) (MeOH) (Derep). λ_{max} 204 (ε 12200); 220 (ε 9100); 278 (ε 8200); 394 (ε 300) (EtOH) (Berdy).**2,2'-Dibromo, 1'-N-Me: 1'-N-Methyl-2,2'-dibromoageliferin**

[175862-03-0]

C₂₃H₂₄Br₄N₁₀O₂ 792.124Alkaloid from the sponge *Astrosclera willeyana*. Off-white glass (as bis(trifluoroacetate)). [α]_D +30 (c, 0.01 in MeOH).**3'-Debromo, 2'-bromo, 1'-N-Me: 1'-N-Methylisoageliferin**

[175861-99-1]

C₂₃H₂₆Br₂N₁₀O₂ 634.332Alkaloid from the sponge *Astrosclera willeyana*. Pale yellow glass (as bis(trifluoroacetate)). [α]_D +40 (c, 0.01 in MeOH).**3'-Debromo, 2'-bromo, 1,1'-di-N-Me: N¹,N^{1'}-Dimethylisoageliferin**

[175862-00-7]

C₂₄H₂₈Br₂N₁₀O₂ 648.359Alkaloid from the sponge *Astrosclera willeyana*. Off-white glass (as dihydrochloride). [α]_D +40 (c, 0.01 in MeOH).**10-Epimer: Nagelamide E**

[690627-58-8]

C₂₂H₂₄Br₂N₁₀O₂ 620.305Alkaloid from an *Agelas* sp. Amorph. solid. [α]_D¹⁷ -11.3 (c, 1 in MeOH). λ_{max} 202 (ε 32000); 215 (sh) (ε 18000); 272 (ε 17200) (MeOH).**10-Epimer, 2-bromo: Nagelamide F**

[690627-59-9]

C₂₂H₂₃Br₃N₁₀O₂ 699.202Alkaloid from an *Agelas* sp. Amorph. solid. [α]_D¹⁷ -14.1 (c, 1 in CHCl₃). λ_{max} 202 (ε 24000); 216 (sh) (ε 20100); 272 (ε 17700) (MeOH).


10-Epimer, 2,2'-dibromo: Nagelamide G
[690627-60-2]
C₂₂H₂₂Br₄N₁₀O₂ 778.098
Alkaloid from an *Agelas* sp. Inhibitor of protein phosphatase 2A. Amorph. solid. $[\alpha]_D^{17} +6.7$ (c, 1 in MeOH). λ_{\max} 202 (ε 32000); 215 (sh) (ε 28900); 277 (ε 27400) (MeOH).

Kobayashi, J. *et al.*, *Tetrahedron*, 1990, **46**, 5579-5586 (isol, pmr, cmr, struct)
Keifer, P.A. *et al.*, *J.O.C.*, 1991, **56**, 2965-2975; 6728 (isol)
Vassas, A. *et al.*, *Planta Med.*, 1996, **62**, 28-30 (isol, activity)
Williams, D.H. *et al.*, *Tetrahedron*, 1996, **52**, 5381-5390 (*N*-Methylageliferins)
Kawasaki, I. *et al.*, *Tet. Lett.*, 2002, **43**, 4377-4380 (deriv, synth)
Hoffmann, H. *et al.*, *Synthesis*, 2003, 1753-1783 (rev)
Endo, T. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1262-1267 (Nagelamides)
O'Malley, D.P. *et al.*, *J.A.C.S.*, 2007, **129**, 4762-4775; 7702 (synth)

Ageline A

A-191

Agelazine F
[88929-28-6]
[8885-51-6]




C₂₆H₄₀N₅⁺ 422.635
Constit. of sponge *Agelas* sp. Active against gram-positive bacteria and fungi. ATPase inhibitor, weak ichthyotoxin. Active against *Mycobacterium tuberculosis*. Plates (MeCN) or powder (as chloride). Mp 178-180° (chloride). $[\alpha]_D^{25} -5.5$ (c, 2.5 in MeOH). $[\alpha]_D -8.4$ (c, 3 in CHCl₃). λ_{\max} 212 (ε 18100); 272 (ε 10000) (EtOH/HCl) (Derep). λ_{\max} 212 (ε 18100); 272 (ε 10000) (EtOH) (Derep).

Capon, R.J. *et al.*, *J.A.C.S.*, 1984, **106**, 1819-1822 (isol, uv, pmr, cmr)
Wu, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2495-2504 (isol, pmr, cmr, uv, struct)
Mangalindan, G.C. *et al.*, *Planta Med.*, 2000, **66**, 364-365 (activity)

Agelocaisarine A₁

A-192

[927409-64-1]



C₂₂H₂₂Br₄N₄O₉ 806.053
Related to Aerothionin, A-158. Constit. of *Aplysina caissara*. Obt. as an inseparable mixt. with Agelocaisarine A₂.
3,3'-Diepimer: Agelocaisarine A₂
[927409-65-2]
C₂₂H₂₂Br₄N₄O₉ 806.053
Constit. of *Aplysina caissara*.

Homologue: Agelocaisarine B₁
[927409-66-3]
C₂₃H₂₄Br₄N₄O₉ 820.08
Constit. of *Aplysina caissara*. Has a C₅ bridging chain (-CH₂CH(OH)CH₂CH₂CH₂-) instead of C₄. Obt. as an inseparable mixt. with Agelocaisarine B₂.

Homologue, 2''-deoxy: Aplysionone B
[943837-23-8]
C₂₃H₂₄Br₄N₄O₈ 804.081
Constit. of *Aplysina gerardogreeni*. Amorph. solid. $[\alpha]_D^{25} -4$ (c, 0.1 in Me₂CO). Has a C₅ bridging chain instead of C₄.


Homologue, 3,3'-diepimer: Agelocaisarine B₂
[927409-67-4]
C₂₃H₂₄Br₄N₄O₉ 820.08
Constit. of *Aplysina caissara*. Has a C₅ bridging chain (-CH₂CH(OH)CH₂CH₂CH₂-) instead of C₄.

De Lira, T.O. *et al.*, *J. Braz. Chem. Soc.*, 2006, **17**, 1233-1240 (Agelocaisarines A₁, A₂, B₁, B₂)
Hernández-Guerrero, C.J. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 5275-5282 (Aplysionone B)

Agelongine

A-193

[163564-84-9]



C₁₃H₁₁BrN₂O₄ 339.145
Alkaloid from *Agelas longissima* and *Axinella damicornis*. Exhibits antiserotonergic activity. Amorph. solid. Sol. MeOH, butanol. λ_{\max} 262 (log ε 4.2) (MeOH).


Debromo: Daminine
[862288-76-4]
C₁₃H₁₂N₂O₄ 260.249
Alkaloid from *Axinella damicornis*. Amorph. solid. Mp 193° dec.

Cafieri, F. *et al.*, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 799-804 (isol, uv, pmr, cmr)
Aiello, A. *et al.*, *Tetrahedron*, 2005, **61**, 7266-7270 (Daminine)

Agelolin A

A-194

[149998-47-0]



C₂₉H₂₆Br₆N₄O₁₁ 1085.969
Hydrol. prod. of Fistularin 3, F-70 on standing. Metab. from the tropical marine sponge *Agelas oroides*. Shows antibacterial activity. Amorph. off-white powder. $[\alpha]_D^{25} -17.1$ (c, 1.26 in Me₂CO). λ_{\max} 220 (ε 12600); 250 (ε 7740) (EtOH) (Berdy).

11,17-Dideoxy: 11,17-Dideoxyagelolin A
C₂₉H₂₆Br₆N₄O₉ 1053.97
Isol. from the sponge *Suberea* aff. *praetensa*.

2,2'-Diepimer: Agelolin B
[150133-30-5]
C₂₉H₂₆Br₆N₄O₁₁ 1085.969
Isol. from *Agelas oroides*. Shows antibacterial activity. Amorph. powder. $[\alpha]_D^{25} +50$ (c, 0.27 in Me₂CO). λ_{\max} 215 (ε 12570); 250 (ε 7940) (EtOH) (Berdy).


2,2'-Diepimer, 11,17-dideoxy: 11,17-Dideoxyagelolin B
C₂₉H₂₆Br₆N₄O₉ 1053.97
Isol. from the sponge *Suberea* aff. *praetensa*.

König, G.M. *et al.*, *Heterocycles*, 1993, **36**, 1351 (isol, uv, ir, pmr, cmr, ms)
Kijjoo, A. *et al.*, *Z. Naturforsch., C*, 2001, **56**, 1116-1119 (11,17-Dideoxyagelolins)
Rogers, E.W. *et al.*, *J. Nat. Prod.*, 2005, **68**, 891-896 (stereochem)
Bardhan, S. *et al.*, *Org. Lett.*, 2006, **8**, 927-930 (stereochem)

Agesamide A

A-195

[911296-34-9]



Relative Configuration

C₁₁H₁₀Br₂N₄O₃ 406.033
Related to Cyclooroidin, C-887. Isol. from an *Agelas* sp. Amorph. solid. $[\alpha]_D^{20} +3.2$ (c, 0.5 in MeOH). λ_{\max} 204 (ε 12000); 282 (ε 2800) (MeOH).


4'-Epimer: Agesamide B
[911296-35-0]
C₁₁H₁₀Br₂N₄O₃ 406.033
Isol. from an *Agelas* sp. Amorph. solid. $[\alpha]_D^{20} +2.8$ (c, 0.5 in MeOH). λ_{\max} 204 (ε 12300); 282 (ε 3050) (MeOH).

Tsuda, M. *et al.*, *Org. Lett.*, 2006, **8**, 4235-4238 (isol, pmr, cmr)
Trost, B.M. *et al.*, *Org. Lett.*, 2007, **9**, 2357-2359 (synth)

Aglaforbesine A

A-196

[177262-33-8]



C₃₆H₄₂N₂O₈ 630.736
Alkaloid from *Aglaia forbesii*. Amorph. solid. [α]_D²⁰ -18 (c, 1 in MeOH). λ_{max} 271 (log ε 3.25) (MeOH).


10-Epimer: Aglaforbesine B

[177468-86-9]
C₃₆H₄₂N₂O₈ 630.736
Alkaloid from *Aglaia forbesii*. Amorph. solid. [α]_D²⁰ +1 (c, 1 in MeOH). λ_{max} 270 (log ε 3.25) (MeOH).

Dumontet, V. et al., *Tetrahedron*, 1996, **52**, 6931-6942 (isol, uv, ir, pmr, cmr)

Aglaiastratin

Aglaroxin D
[176785-78-7]



Absolute Configuration

C₃₁H₃₀N₂O₆ 526.588
Isol. from leaves of *Aglaia odorata* and *Aglaia roxburghiana*. Protein synthesis inhibitor. K-Ras transformed fibroblast morphology inducer. Powder + 1½H₂O. Mp 157-160°. [α]_D²³ +59.7 (c, 0.3 in MeOH). λ_{max} 281 (ε 4163); 304 (ε 4480) (MeOH).

5a''', N^{6''''}-Didehydro: **Dehydroaglaiastratin**. *Didehydroaglaiastratin* [155595-93-0]
C₃₁H₂₈N₂O₆ 524.572
Isol. from roots of *Aglaia odorata* and twigs of *Aglaia dupperreana* (Meliaceae). Protein synthesis inhibitor. Plates (CH₂Cl₂/MeOH). Mp 256-257°. [α]_D²⁰ -50.1 (c, 0.41 in CHCl₃). λ_{max} 215 (ε 50000); 234 (ε 19000); 272 (ε 6000) (EtOH) (Derep).

5a''', N^{6''''}-Didehydro, 7-O-de-Me: **Mari-karin** [349642-85-9]
C₃₀H₂₆N₂O₆ 510.545
Alkaloid from *Aglaia gracilis*. Oil. [α]_D²⁰ -45 (c, 0.3 in CHCl₃). λ_{max} 273 (MeOH).

3'-Hydroxy, 5a''', N^{6''''}-didehydro: **Didehydro-3'-hydroxyaglaiastratin** [259143-58-3]
C₃₁H₂₈N₂O₇ 540.571
Isol. from *Aglaia dupperreana*.

3'-Hydroxy, 5a''', N^{6''''}-didehydro, 7-O-de-Me: **3'-Hydroxymari-karin** [349484-01-1]
C₃₀H₂₆N₂O₇ 526.545
Alkaloid from *Aglaia gracilis*. Oil. [α]_D²⁰ -53 (c, 0.1 in CHCl₃). λ_{max} 279 (MeOH).

3'-Methoxy, 5a''', N^{6''''}-didehydro: **Aglaiformosanin**. *Didehydro-3'-methoxyaglaiastratin* [371163-97-2]
C₃₂H₃₀N₂O₇ 554.598

Isol. from *Aglaia formosana*. Cytotoxic. Powder. Mp 260-262°. [α]_D²² -78.9 (c, 0.01 in CHCl₃). λ_{max} 213 (log ε 4.39); 256 (log ε 3.78); 274 (log ε 3.68) (MeOH).

Stereoisomer: [189343-27-9]

C₃₁H₃₀N₂O₆ 526.588
Isol. from twigs of *Aglaia dupperreana*. [α]_D²⁰ +45.7 (c, 0.19 in CHCl₃).

Kokpol, U. et al., *Chem. Comm.*, 1994, 773-774 (isol, ir, pmr, cmr, *cryst struct*)
Ohse, T. et al., *J. Nat. Prod.*, 1996, **59**, 650-652 (isol, uv, ir, pmr, cmr, *cryst struct*)
Pat. Coop. Treaty (WIPO), 1996, 96 04 284; *CA*, **124**, 335673h (*Aglaroxin D*)
Nugroho, B.W. et al., *Phytochemistry*, 1997, **44**, 1455-1461 (isol, pmr, cmr, *cd, ms*)
Watanabe, T. et al., *Chem. Comm.*, 1998, 1097-1098 (*synth*)
Chaidir, et al., *Phytochemistry*, 1999, **52**, 837-842 (*Didehydro-3'-hydroxyaglaiastratin*)
Watanabe, T. et al., *Heterocycles*, 2000, **53**, 1051-1064 (*synth*)
Greger, H. et al., *Phytochemistry*, 2001, **57**, 57-64 (*Mari-karin*, 3'-Hydroxymari-karin)
Wang, S.-K. et al., *Planta Med.*, 2001, **67**, 555-557 (*Aglaiformosanin*)

Aglaidithioduline

A-198

N,N'-Bis[3-(methylthio)-2-propenyl]putrescine. *Hemileptagline*
MeSCH=CH
CONH(CH₂)₄NHCOCH=CHSMe
C₁₂H₂₀N₂O₂S 256.368
Struct. of *Hemileptagline* revised in 2002.

(*E,E*)-form [261158-70-7]


Alkaloid from *Aglaia leptantha* and *Aglaia edulis*. Pale orange needles (MeOH). Mp 164-165°. λ_{max} 271 (log ε 4.6) (MeOH).

Saifah, E. et al., *Phytochemistry*, 1999, **52**, 1085-1088 (*Aglaidithioduline*)
Greger, H. et al., *J. Nat. Prod.*, 2000, **63**, 616-620 (*Hemileptagline*)
Detterback, R. et al., *Tetrahedron*, 2002, **58**, 6887-6893 (*synth*)

Aglaine B

A-199

Aglaroxin K
[177262-32-7]
[251982-74-8 (*Aglaroxin K*)]



C₃₆H₄₂N₂O₈ 630.736
Alkaloid from *Aglaia argentea* and *Aglaia roxburghiana*. Amorph. powder. [α]_D²⁰ -10 (c, 1 in MeOH). λ_{max} 272 (log ε 3.41) (MeOH).

5-Ac: 5-Acetylglaine B. *Aglaroxin J*
[251982-73-7]

C₃₈H₄₄N₂O₉ 672.774
Alkaloid from the stem bark of *Aglaia roxburghiana*.

10-Ac: **10-Acetylglaine B**

[177468-88-1]
C₃₈H₄₄N₂O₉ 672.774
Alkaloid from the leaves of *Aglaia elliptica*. Amorph. powder. [α]_D²⁰ +20.4 (c, 0.83 in MeOH).

3'-Hydroxy: **3'-Hydroxyaglaïne B**

[243136-41-6]
C₃₆H₄₂N₂O₉ 646.736
Alkaloid from *Aglaia odorata*. [α]_D²⁰ -11.4 (c, 0.25 in CHCl₃).

10-Epimer: **Deacetylglaine A**

C₃₆H₄₂N₂O₈ 630.736
Alkaloid from *Aglaia gracilis*. Oil. [α]_D²⁰ -14 (c, 0.1 in CHCl₃). Possible artifact. C-19 config. not determined. λ_{max} 271 ; 277 (sh) (MeOH).

10-Epimer, 10-Ac: **Aglaïne A**

[177262-31-6]
C₃₈H₄₄N₂O₉ 672.774
Alkaloid from *Aglaia argentea* and *Aglaia forbesii*. Amorph. powder. [α]_D²⁰ -8 (c, 1 in MeOH). λ_{max} 272 (log ε 3.6) (MeOH).

10-Epimer, 19,20-didehydro(E-): **Elliptifoline**

[319917-12-9]
C₃₆H₄₀N₂O₈ 628.721
Alkaloid from *Aglaia elliptifolia*. Cytotoxic agent. Powder. Mp 184-185°. [α]_D²² -88.9 (c, 0.6 in CHCl₃). λ_{max} 273 (log ε 3.36) (MeOH).

3,4-Diepimer: **Aglaïne C**

[177468-85-8]
C₃₆H₄₂N₂O₈ 630.736
Alkaloid from *Aglaia argentea*. Cryst. (MeOH). Mp 180°. [α]_D²⁰ -105 (c, 1 in CHCl₃). λ_{max} 271 (log ε 3.27) (MeOH).

3,4-Diepimer, 3'-hydroxy: **3'-Hydroxyaglaïne C**

[243136-38-1]
C₃₆H₄₂N₂O₉ 646.736
Alkaloid from *Aglaia odorata*. [α]_D²⁰ -103.4 (c, 0.43 in CHCl₃).

3,4-Diepimer, 19ξ-hydroxy: **Aglaroxin L**

[251982-75-9]
C₃₆H₄₂N₂O₉ 646.736
Alkaloid from the stem bark of *Aglaia roxburghiana*. C-19 config. not determined.

3,4-Diepimer, 3',19-dihydroxy: **3',19-Dihydroxyaglaïne C**

[243136-39-2]
C₃₆H₄₂N₂O₁₀ 662.735
Alkaloid from *Aglaia odorata*. [α]_D²⁰ -86 (c, 0.18 in CHCl₃). C-19 config. not determined.

3,4-Diepimer, 3'-methoxy, 19-hydroxy: **19-Hydroxy-3'-methoxyaglaïne C**

[243136-40-5]
C₃₇H₄₄N₂O₁₀ 676.762
Alkaloid from *Aglaia odorata*. [α]_D²⁰ -111 (c, 0.18 in CHCl₃). C-19 config. not determined.

4,10-Diepimer, 10-Ac: **4-Epiaglaïne A**

[372119-00-1]
C₃₈H₄₄N₂O₉ 672.774


Alkaloid from the leaves of *Aglaia elliptica*. Amorph. powder. $[\alpha]_D^{25}$ -1 (c, 1 in CHCl_3).

- Dumontet, V. *et al.*, *Tetrahedron*, 1996, **52**, 6931-6942 (*isol, uv, ir, pmr, cmr*)
 Molleyres, L.-P. *et al.*, *Pestic. Sci.*, 1999, **55**, 486-503 (*Aglaroxins J,K,L*)
 Nugroho, B.W. *et al.*, *Phytochemistry*, 1999, **51**, 367-376 (*isol, derivs*)
 Inada, A. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1226-1228 (*4-Epiaglaïne A, 10-Acetylglaine B*)
 Wang, S.-K. *et al.*, *J. Nat. Prod.*, 2001, **64**, 92-94 (*Elliptifoline*)
 Greger, H. *et al.*, *Phytochemistry*, 2001, **57**, 57-64 (*Deacetylglaine A*)

Aglaroxin I

A-200

[251982-72-6]



Absolute Configuration

 $\text{C}_{31}\text{H}_{28}\text{N}_2\text{O}_7$ 540.571Alkaloid from the stem bark of *Aglaia roxburghiana*.**5a,6-Didehydro: Aglaroxin C**

[176785-77-6]


 $\text{C}_{31}\text{H}_{26}\text{N}_2\text{O}_7$ 538.556Alkaloid from *Aglaia roxburghiana*.Insecticide. λ_{max} 274 (ϵ 9600) (MeCN).**3'-Methoxy: Aglaroxin H**

[251982-71-5]

 $\text{C}_{32}\text{H}_{30}\text{N}_2\text{O}_8$ 570.598Alkaloid from the stem bark of *Aglaia roxburghiana*.**3'-Methoxy, 5a''',6-didehydro: Aglaroxin G** $\text{C}_{32}\text{H}_{28}\text{N}_2\text{O}_8$ 568.582Alkaloid from *Aglaia roxburghiana*.Pat. Coop. Treaty (WIPO), 1996, 96 04 284; CA, **124**, 335673h (*Aglaroxin C*)Molleyres, L.-P. *et al.*, *Pestic. Sci.*, 1999, **55**, 494-497 (*Aglaroxins G-I*)**Aglaxiflorin C**

A-201

[269739-77-7]




R = OAc

 $\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_{10}$ 688.773Flavagline compd. Alkaloid from *Aglaia laxiflora*. Cryst. (CHCl_3). Mp 147-149°. $[\alpha]_D^{25}$ +34.9 (c, 0.05 in CHCl_3). λ_{max} 222 (log ϵ 4.42); 272 (log ϵ 3.29) (MeOH).Xu, Y.-J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 473-476**Aglaxiflorin D**

A-202

[269739-78-8]

 $\text{C}_{36}\text{H}_{42}\text{N}_2\text{O}_9$ 646.736Flavagline compd. Alkaloid from *Aglaia laxiflora*. Cryst. (CHCl_3). Mp 131-133°. $[\alpha]_D^{25}$ -102.1 (c, 0.04 in CHCl_3). λ_{max} 210 (log ϵ 5.62); 214 (log ϵ 5.66); 268 (log ϵ 4.66) (MeOH).**12-Epimer, 12-Ac: Aglaxiflorin B**

[269739-75-5]


 $\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_{10}$ 688.773Alkaloid from *Aglaia laxiflora*. Cryst. (CHCl_3). Mp 158-160°. $[\alpha]_D^{25}$ -105.5 (c, 0.13 in CHCl_3). λ_{max} 222 (log ϵ 4.47); 274 (log ϵ 3.66) (MeOH).**10,12-Diepimer, 12-Ac: Aglaxiflorin A**

[269739-76-6]

 $\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_{10}$ 688.773Alkaloid from *Aglaia laxiflora*. Cryst. (CHCl_3). Mp 217-219°. $[\alpha]_D^{25}$ +8.1 (c, 0.12 in CHCl_3). λ_{max} 214 (log ϵ 4.64); 268 (log ϵ 3.73) (MeOH).Xu, Y.-J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 473-476**Agrobactin**

A-203

[70393-50-9]

 $\text{C}_{32}\text{H}_{36}\text{N}_4\text{O}_{10}$ 636.657Isol. from *Agrobacterium tumefaciens*. Siderophore. Cryst. (EtOAc/hexane). Sol. H_2O ; fairly sol. MeOH. Dec. at 108-112°. λ_{max} 252 ; 316 (H_2O) (Berdy).**3'-Deoxy: Parabactin. Parabactin A**

[74149-70-5]

 $\text{C}_{32}\text{H}_{36}\text{N}_4\text{O}_9$ 620.658From *Paracoccus denitrificans*. Siderophore with antitumour and antiviral activity. Iron chelator. Cryst. (EtOAc/hexane). Sol. MeOH, Et_2O , bases; poorly sol. H_2O . Mp 114-117°. $[\alpha]_D^{21}$ +99.6 (c, 1.8 in MeOH). Formerly given a struct. with the oxazole ring in open-form under the name Parabactin A. λ_{max} 250 (ϵ 26600); 310 (ϵ 11000) (EtOH) (Berdy). λ_{max} 208 (ϵ 50000); 310 (ϵ 7700) (pH 7 buffer) (Berdy). λ_{max} 210 (ϵ 56800); 253 (ϵ 27300); 319 (ϵ 8200) (pH 1.5 buffer) (Berdy).

[54135-85-2]


Tait, G.H. *et al.*, *Biochem. J.*, 1975, **146**, 191

(Parabactin)

Ong, S.A. *et al.*, *J. Biol. Chem.*, 1979, **254**, 1860 (*isol, props*)Sciortino, C.V. *et al.*, *J. Lab. Clin. Med.*, 1980, **96**, 1081 (*props*)Peterson, T. *et al.*, *J.A.C.S.*, 1980, **102**, 7715 (*synth, struct*)Eng-Wilmot, D.L. *et al.*, *J.A.C.S.*, 1980, **102**, 7719 (*cryst struct*)Neilands, J.B. *et al.*, *Methods Enzymol.*, 1983, **94** (*rev*)Bergeron, R.J. *et al.*, *Biochem. Biophys. Res. Commun.*, 1984, **121**, 848 (*activity*)Nagao, Y. *et al.*, *J.C.S. Perkin 1*, 1984, 183 (*Parabactin*)Bergeron, R.J. *et al.*, *J.O.C.*, 1985, **50**, 2780 (*synth*)Bergeron, R.J. *et al.*, *Cancer Res.*, 1987, **47**, 6010 (*Parabactin, pharmacol*)Bergeron, R. *et al.*, *Blood*, 1993, **82**, 2552 (*Parabactin, pharmacol*)**Agrochelin**

A-204


[247115-74-8]



Relative Configuration

 $\text{C}_{23}\text{H}_{34}\text{N}_2\text{O}_4\text{S}_2$ 466.665Alkaloid from a marine *Agrobacterium* sp. Cytotoxic agent. Siderophore. Pale yellow oil. $[\alpha]_D^{25}$ -20.5 (c, 0.2 in CHCl_3). Chelates Zn^{2+} ion. λ_{max} 210 (log ϵ 5.18); 264 (log ϵ 4.8) (MeOH).Acebal, C. *et al.*, *J. Antibiot.*, 1999, **52**, 983-987 (*isol*)Canedo, L.M. *et al.*, *Tet. Lett.*, 1999, **40**, 6841-6844 (*isol, uv, cd, pmr, cmr*)**Agroclavine**

A-205

8,9-Didehydro-6,8-dimethylergoline, 9CI

(-)-form

 $\text{C}_{16}\text{H}_{18}\text{N}_2$ 238.332**(-)-form** [548-42-5]Alkaloid from ergot and from several spp. in the genera *Argyriaea*, *Rivea*, *Cuscuta*, *Ipomoea* (Convolvulaceae). Prod. by *Penicillium*, *Aspergillus* and *Rhizopus* spp. CNS excitator acting by stimulation of sympathetic centres. Active against gram-positive bacteria. Mycotoxin. Sol. MeOH, CHCl_3 , Py; fairly sol. C_6H_6 ; poorly sol. H_2O . Mp 208-209° dec. $[\alpha]_D$ -151 (CHCl_3). $[\alpha]_D$ -182 (Py). λ_{max} 225 (ϵ 29512); 284 (ϵ 7586); 293 (ϵ 6456) (MeOH) (Berdy).▶ LD₅₀ (mus, ipr) 25 mg/kg. KE6325000

Dihydro: see Festuclavine, F-49

18-Hydroxy: Elymoclavine

[548-43-6]

C₁₆H₁₈N₂O 254.331

Alkaloid from ergot and several other *Claviceps* spp. and from *Stictocardia tiliaefolia*, *Stictocardia campanulata*, *Argyria* spp. and several other spp. in the Convolvulaceae. Prod. by *Penicillium aurantiovirens*. 5HT receptor antagonist. CNS excitatory agent acting by stimulation of sympathetic centres. Mycotoxin. Sol. MeOH, CHCl₃; fairly sol. C₆H₆; poorly sol. H₂O. Mp 250-252° dec. [α]_D²⁰ -152 (c, 1 in Py). λ_{max} 227 (ε 20417); 283 (ε 6918); 293 (ε 5754) (MeOH) (Berdy).

▶ KE6340000

18-Hydroxy, O-β-D-fructofuranoside: Elymoclavine O-β-D-fructofuranoside
[28986-92-7]
[121844-72-2]

C₂₂H₂₈N₂O₆ 416.473

Isol. from the *Claviceps* strain SD 58 grown in a culture medium containing saccharose. Amorph. Prob. artifact formed in a secondary process from Elymoclavine and fructose from saccharose in the culture medium.

18-Hydroxy, O-[β-D-fructofuranosyl-(2→1)-β-D-fructofuranoside]: Elymo-clavine O-β-D-fructofuranosyl-(2→1)-O-β-D-fructofuranoside
[121864-94-6]

C₂₈H₃₈N₂O₁₁ 578.615

Isol. from saprophytic cultures of *Claviceps* strain SD 58 and *Claviceps purpurea* strain 88 EP on sucrose medium.

18-Hydroxy, O-4-nitrobenzoyl: Mp 176-177° dec. [α]_D²⁰ -127 (c, 0.4 in CHCl₃).

18-Hydroxy, 5,10-didehydro: 4,6,7,8-Tetrahydro-7-methylindolo[4,3-fg]quinoline-9-methanol. Dehydroelymo-clavine
[141869-64-9]

C₁₆H₁₆N₂O 252.315

Alkaloid from roots of *Securidaca longipedunculata*.

10-Epimer: Agroclavine I

[82597-81-7]

C₁₆H₁₈N₂ 238.332

Metab. of *Penicillium kapuscinskii* strain UCM F-2156D. λ_{max} 211 (sh) (ε 14600); 225 (ε 25500); 276 (sh) (ε); 282 (ε 5470); 292 (ε 4560) (MeOH) (Derep).

10-Epimer, N¹-methoxy: 1-Methoxyagro-clavine I

[207349-44-8]

C₁₇H₂₀N₂O 268.358

Prod. by *Penicillium* sp. WC75209. Inhibitor of Lck tyrosine kinase. [α]_D²⁰ -63 (c, 0.08 in MeOH). λ_{max} 228; 282 (MeOH).

10-Epimer, 8β,9β-epoxide: Epoxyagro-clavine I

[82564-34-9]

C₁₆H₁₈N₂O 254.331

Metab. from *Penicillium kapuscinskii* and *Penicillium sizovae* VKM.F1073. Mycotoxin. λ_{max} 211 (sh) (ε 14600);

225 (ε 25500); 276 (sh) (ε); 282 (ε 5470); 292 (ε 4560) (MeOH) (Derep).

(±)-form

Synthetic. Mp 189-191° dec.

18-Hydroxy: Synthetic. Cryst. (CHCl₃/MeOH). Mp 207-210° dec.

10-Epimer: Synthetic.

Cryst. (Et₂O). Mp 157-158° (152-154°).

Stoll, A. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1815-1825 (*Elymoclavine, isol, uv*)

Schreier, E. *et al.*, *Helv. Chim. Acta*, 1958, **41**, 1984-1997 (*Elymoclavine, config*)

Floss, H.G. *et al.*, *Z. Naturforsch., B*, 1967, **22**, 399-402 (*Elymo-clavin fructoside*)

Inoue, T. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 409-411 (*ms*)

Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435-2440 (*occur*)

Bach, N.J. *et al.*, *J.O.C.*, 1974, **39**, 1272-1276 (*pmr, cmr*)

Maier, W. *et al.*, *Biochem. Physiol. Pflanz.*, 1978, **172**, 531-536 (*biosynth*)

Wurst, M. *et al.*, *J. Chromatogr.*, 1978, **150**, 477; 1979, **174**, 401-407 (*hplc*)

Mrtek, R.G. *et al.*, *Phytochemistry*, 1978, **7**, 1535 (*Elymo-clavine, pmr*)

Saini, M. *et al.*, *Phytochemistry*, 1978, **17**, 799 (*biosynth*)

Kiguchi, T. *et al.*, *Heterocycles*, 1984, **22**, 43-45; 1985, **23**, 2891-2893 (*Agroclavine I, synth*)

Sakharovsky, V.G. *et al.*, *Tet. Lett.*, 1984, **25**, 109-112 (*Agroclavine I, Epoxyagroclavine I*)

Eich, E. *et al.*, *Arzneim.-Forsch.*, 1985, **35**, 1760-1762 (*activity*)

Kozikowski, A.P. *et al.*, *J.A.C.S.*, 1985, **107**, 2569-2571 (*Agroclavine I, synth*)

Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 948-950 (*synth*)

Wheeler, W.J. *et al.*, *Tet. Lett.*, 1986, **27**, 3469-3470 (*synth*)

Somei, M. *et al.*, *Heterocycles*, 1987, **26**, 895-898; 1997, **45**, 1263-1266 (*Agroclavine I, synth*)

Flieger, M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 506-510; 1991, **54**, 390-395 (*cmr, Elymo-clavine fructosides*)

Ninomiya, I. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 23-30 (*Agroclavine I, synth*)

Costa, C. *et al.*, *J. Het. Chem.*, 1992, **29**, 1641-1647 (*Dehydroelymo-clavine*)

Padmanabha, R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 569-574 (*1-Methoxyagroclavine*)


Yamada, F. *et al.*, *Heterocycles*, 2007, **72**, 599-620 (*Agroclavine I, synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AEY375; EAJ000 (*Agroclavine, Elymo-clavine*)

Agrocybenine

A-206

1,2,3,5-Tetrahydro-2,2,5,5,7-pentamethyl-6H-pyrrolo[2,3-b]pyridin-6-one, 9CI
[178764-92-6]



C₁₂H₁₈N₂O 206.287

Alkaloid from the edible Korean mushroom yangimatusutake (*Agrocybe*


cylindracea). Prolyl endopeptidase inhibitor. Mycotoxin. Yellow powder. λ_{max} 223 (ε 16800); 333 (ε 9200) (MeOH).

Koshino, H. *et al.*, *Tet. Lett.*, 1996, **37**, 4549 (*isol, uv, ir, pmr, cmr, N-15 nmr, struct*)

Agropine

A-207

2-Oxo-6-(1,2,3,4-tetrahydroxybutyl)-3-morpholinepropanamide, 9CI
[70699-77-3]



C₁₁H₂₀N₂O₇ 292.288

Struct. revised in 1982. Obt. from crown gall tumours.

Coxon, D.T. *et al.*, *Tet. Lett.*, 1980, 495 (*synth, spectra*)


Tate, M.E. *et al.*, *Carbohydr. Res.*, 1982, **104**, 105 (*ms, struct*)

Chilton, W.S. *et al.*, *Phytochemistry*, 1995, **40**, 619 (*occur*)

Ailanindole

A-208

9-Aminopyrido[1,2-a]indole-6,10-dione, 9CI
[159903-51-2]



C₁₂H₈N₂O₂ 212.207


Alkaloid from the wood of *Ailanthus malabarica*. Orange needles (MeOH). Mp 300°. Intense green fluorescence in org. solvs. λ_{max} 238 (log ε 4.16); 252 (sh) (log ε 3.97); 296 (sh) (log ε 3.33); 452 (log ε 3.83); 474 (log ε 3.82) (MeOH).

Aono, H. *et al.*, *Phytochemistry*, 1994, **37**, 579-584 (*isol, uv, ir, pmr, cmr, ms*)

Ailanthoidine

A-209

[86003-22-7]




C₂₉H₂₅N₃O₄ 479.534

Alkaloid from the bark of *Zanthoxylum ailanthoides* (Rutaceae). Mp 260-263°. [α]_D 0.

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1340 (*isol, pmr, cmr, cryst struct*)

Ajabicine

[148225-24-5]

C₂₂H₃₃NO₂ 343.508

Unique methyleneaconitane struct. Alkaloid from leaves of *Delphinium ajacis* (Ranunculaceae). Amorph.

1-Ketone: Actaline

[122279-81-6]

C₂₂H₃₁NO₂ 341.492


Alkaloid from the epigeal part of *Aconitum talassicum* (Ranunculaceae). Cryst. (hexane). Mp 125-127°.

Nishanov, A.A. et al., *Khim. Prir. Soedin.*, 1989, **25**, 39; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 32 (isol, ir, pmr, ms, cryst struct)

Joshi, B.S. et al., *Tet. Lett.*, 1993, **34**, 1441 (isol, pmr, cmr, struct)

Ajaconine, 9CI

[545-61-9]



Absolute configuration

C₂₂H₃₃NO₃ 359.508

Alkaloid from *Delphinium virescens* and *Delphinium carolinianum* (whole plants), the roots of *Delphinium tatsienense*, and the seeds of *Delphinium consolida*, *Delphinium ajacis* and *Consolida ambigua* (Ranunculaceae). Also isol. from *Consolida oliveriana*. Mp 167°. [α]_D -122 (c, 1.75 in EtOH). pK_{a1} 11.8 (50% MeOH aq.). pK_{a1} 11.3 (80% EtOH aq.).

Sulfate (2:1):

Cryst. + 7H₂O (Me₂CO aq.). Mp 113° (hydrate) Mp 231° (anhyd.).

15-Epimer: 15-Epiajaconine. Deacetyl-spiramine FC₂₂H₃₃NO₃ 359.508

Alkaloid from *Spiraea japonica* var. *ovalifolia*. Plates (Me₂CO). Mp 149-151°.

15-Epimer, 15-Ac: Spiramine F

[142750-37-6]

C₂₄H₃₅NO₄ 401.545

Alkaloid from roots of *Spiraea japonica* var. *acuminata* (Rosaceae). Amorph. [α]_D¹⁸ -101 (c, 2.5 in CHCl₃).

15-Epimer, di-Ac: Spiramine E

[142750-36-5]

C₂₆H₃₇NO₅ 443.582

Alkaloid from roots of *Spiraea japonica* var. *acuminata* (Rosaceae). Amorph. [α]_D²⁰ -97 (c, 1.9 in CHCl₃).

15-Epimer, 19-oxo, 15-Ac: Spiramine R

[162858-00-6]

C₂₄H₃₃NO₅ 415.528

Alkaloid from roots of *Spiraea*

A-210

japonica var. *incisa* (Rosaceae). Needles (EtOAc). Mp 190-192°. [α]_D -180 (c, 0.39 in CHCl₃).

Keller, O. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1913, **251**, 207; *CA*, **8**, 674 (isol)

Dvornik, D. et al., *Tetrahedron*, 1961, **14**, 54 (isol, ir, struct)

Scott, A.I. et al., *Tetrahedron*, 1971, **27**, 4787 (cd)

Sastry, S.D. et al., *Chem. Ind. (London)*, 1972, 381 (ms)

Pelletier, S.W. et al., *Heterocycles*, 1979, **12**, 779; 1981, **16**, 747; 1983, **20**, 1347 (occur)

Pelletier, S.W. et al., *J. Nat. Prod.*, 1980, **43**, 395 (isol, ir, pmr)

Hao, X.-J. et al., *Heterocycles*, 1993, **36**, 825 (Spiramine E, Spiramine F)

Hao, X.-J. et al., *Phytochemistry*, 1995, **38**, 545 (Spiramine R)


Ulubelen, A. et al., *J. Nat. Prod.*, 1996, **59**, 907 (cmr)

Zuo, G.Y. et al., *Heterocycles*, 2001, **55**, 487-493 (15-Epiajaconine)

González-Coloma, A. et al., *Chem. Biodiversity*, 2004, **1**, 1327-1335 (isol, pmr)

Ajmalicine**A-212**

Methyl 16,17-didehydro-19-methyloxayohimban-16-carboxylate, 9CI. Tetrahydroserpentine. Raubasine. δ-Yohimbine. Vincine. Vincaine†. Py-tetrahydroserpentine. Lamuran



Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Log P 2.76 (calc).

(-)-form [483-04-5]

Alkaloid from *Rauwolfia serpentina*, many other *Rauwolfia* spp., *Catharanthus roseus* and many other spp. in the Apocynaceae. Antidiuretic agent, CNS depressant. Vasodilator. Has been used to treat peripheral and cerebral vascular disorders. Active against fungi. Prisms (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 259° dec. [α]_D -66 (CHCl₃). λ_{max} 227 (ε 40740); 246 (sh) (ε 11750); 280 (ε 6920); 292 (ε 6160) (MeOH) (Berdy).

▶ LD₅₀ (mus, orl) 400 mg/kg. Human systemic effects by ingestion. AX7875000

Hydrochloride:

Leaflets (EtOH). Mp 289°. [α]_D²⁰ -17 (c, 0.5 in MeOH).

3-Epimer: 3-Isoajmalicine

[483-03-4]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Mitragyna* and *Uncaria* spp. (Rubiaceae). Needles (Et₂O). Mp 193-194°. λ_{max} 225 (log ε 4.61); 250 (log ε 4.14); 282 (log ε 3.93); 291 (log ε 3.87) (EtOH).

19-Epimer: Mayumbine. 19-Epiajmalicine [25532-45-0]C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Pseudocinchona mayumbensis* (syn. *Corynanthe mayumbensis*), *Uncaria elliptica* and

Rauwolfia yunnanensis (Rubiaceae, Rubiaceae, Apocynaceae). Mp 216°.

[α]_D -68 (Py). λ_{max} 227 (log ε 4.6); 280 (log ε 3.84); 292 (log ε 3.8) (MeOH).

19-Epimer, hydrochloride: Mp 293-294°.

[α]_D²⁰ -96 (MeOH).

19-Epimer, 20,21-didehydro: see Cathenamamine, C-207

20-Epimer: Tetrahydroalstonine

[6474-90-4]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Alstonia*, *Amsonia*, *Cabucala*, *Catharanthus*, *Mitragyna*, *Rauwolfia*, *Uncaria*, *Rhazya* and *Voa-canga* spp. Active against gram-positive bacteria. Lustrous plates (EtOH aq.). Mp 230-231°. [α]_D²⁷ -108 (c, 0.5 in CHCl₃). λ_{max} 226 (log ε 4.64); 248 (sh) (log ε 4.07); 272 (log ε 3.78); 280 (sh) (log ε 3.85); 291 (log ε 3.83) (MeOH) (Berdy).

20-Epimer, N^b-oxide(R-): Tetrahydroalstonine (4R)-N-oxide

[41590-28-7]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Uncaria* spp. (Rubiaceae). Cryst. (Me₂CO/Et₂O). Mp 172-174° dec. [α]_D²⁰ +33.8 (c, 0.11 in MeOH). λ_{max} 224 (ε 43400); 245 (sh) (ε 11800); 273 (sh) (ε 7100); 282 (ε 7450); 290 (ε 6400) (EtOH).

20-Epimer, N^b-Me: Melinonine A. N^b-Methyltetrahydroalstonine

[6801-41-8]

C₂₂H₂₇N₂O₅[⊕] 367.467

Quaternary alkaloid from *Strychnos melinoniana* and *Rauwolfia volkensii* (Loganiaceae, Apocynaceae). Hygroscopic needles (EtOH/Me₂CO) (as chloride). Mp 260-261° (chloride). [α]_D²⁰ -120 (H₂O) (chloride). λ_{max} 225 (log ε 4.8); 279 (log ε 4); 290 (log ε 3.8) (MeOH).

3,19-Diepimer: 19-Epi-3-isoajmalicine

[25532-46-1]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Uncaria elliptica*, *Uncaria attenuata* and *Catharanthus roseus* (Rubiaceae, Apocynaceae). λ_{max} 225; 273 (sh); 283; 290 (MeOH).

3,20-Diepimer: Akuammigine

[642-17-1]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Picralima nitida* (*Picralima klaineana*) and some *Uncaria* spp. (Apocynaceae, Rubiaceae). Mp 125° (113°). [α]_D²⁰ -44.4 (EtOH). [α]_D^{18.5} -42 (c, 1.6 in EtOH).

3,20-Diepimer, hydrochloride: Mp 287°.[α]_D²⁰ -38 (MeOH).**3,20-Diepimer, (4R)-N-oxide: Akuammigine (4R)-N-oxide**

[41590-26-5]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from an *Uncaria* sp. Cryst. (Me₂CO/Et₂O). Mp 214-215°. [α]_D²⁰ -20 (c, 0.11 in MeOH). [α]_D²⁰ -46 (c, 0.41 in Py). λ_{max} 223 (ε 43000); 245 (sh) (ε 13150); 281 (ε 8300); 291 (ε 6600) (EtOH).

3,20-Diepimer, (4S)-N-oxide: Akuammigine (4S)-N-oxide

[41590-27-6]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from an *Uncaria* sp. (Rubiaceae). Cryst. (Me₂CO/Et₂O). Mp 195-197° dec. [α]_D²⁰ -112 (c, 0.12 in MeOH). λ_{max} 224 (ε 43400); 246 (sh) (ε 10200); 273 (sh) (ε 7900); 282 (ε 8350); 290 (ε 7150) (EtOH).

19,20-Diepimer: **Rauniticine**. *Ervine*[†] [5299-09-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Rauwolfia nitida*, *Uncaria elliptica*, *Uncaria attenuata*, *Vinca libanotica*, *Vinca major* and *Veratrum album lobelianum* (Apocynaceae, Rubiaceae, Liliaceae). Cryst. (MeOH). Mp 233-235°. [α]_D -6.6 (Py). [α]_D -38.4 (CHCl₃). λ_{max} 228 (log ε 4.65); 246 (sh) (log ε 4.09); 282 (log ε 3.91); 288 (log ε 3.79) (EtOH).

19,20-Diepimer, 14α-hydroxy: **14α-Hydroxyrauniticine** [77101-58-7]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the leaves of *Uncaria attenuata* and *Uncaria elliptica* (Rubiaceae). Plates (MeOH). Mp 174-175° dec. [α]_D -17.5 (c, 0.1 in MeOH). Originally proposed as 14β-Hydroxy-3-isorauniticine.

3,19,20-Triepimer: **3-Isorauniticine** [6870-40-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Corynanthe mayumbensis* and the leaves of *Uncaria elliptica* (Apocynaceae, Rubiaceae). Cryst. + 1/2 H₂O (as hydrochloride). Mp 277° (hydrochloride). [α]_D +25 (c, 1 in MeOH) (as hydrochloride).

(±)-**form** [24196-16-5]

Synthetic. Mp 228-231°.

Hydrochloride: Mp 285-286° dec.

3,20-Diepimer: [20361-85-7]

Synthetic. Mp 143° (125-126°).

Henry, T.A. *et al.*, *J.C.S.*, 1932, 2759-2768 (*Akuammigine, isol*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 29-45 (*Melinonine A*)

Klohs, M.W. *et al.*, *J.A.C.S.*, 1954, **76**, 1332-1334 (*isol, uv, ir*)

Robinson, R. *et al.*, *J.C.S.*, 1954, 3479-3482 (*Akuammigine, isol, uv, ir, struct*)

Raymond-Hamet, *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1961, **253**, 2776-2778 (*Akuammigine, pharmacol*)

Salkin, R. *et al.*, *J. Pharm. Sci.*, 1961, **50**, 1038-1041 (*Rauniticine, isol, uv, ir, struct*)

Antonaccio, L.D. *et al.*, *J.A.C.S.*, 1962, **84**, 2161-2169 (*ms*)

Shamma, M. *et al.*, *J.A.C.S.*, 1963, **85**, 2507-2512 (*ir, pmr, config*)

Hesse, M. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 674-689 (*Melinonine A, ms*)

Finch, N. *et al.*, *Tetrahedron*, 1966, **22**, 1327-1333 (*Rauniticine, uv, ord, stereochem*)

Laubie, M. *et al.*, *Arzneim.-Forsch.*, 1969, **19**, 1820-1826 (*pharmacol*)

Winterfeld, E. *et al.*, *Chem. Ber.*, 1969, **102**, 3558-3572 (*epimer, synth*)

Van Tamelen, E.E. *et al.*, *J.A.C.S.*, 1969, **91**, 7359-7371 (*synth*)

Gutzwiller, J. *et al.*, *J.A.C.S.*, 1971, **93**, 5907-5908 (*synth, ir, ms, pmr, Akuammigine*)

Merlini, L. *et al.*, *Tetrahedron*, 1972, **28**, 5971-5975 (*Akuammigine oxides, Tetrahydroalstonine oxide, isol, synth*)

Levin, R.H. *et al.*, *J.O.C.*, 1973, **38**, 1983-1992 (*cmr*)

Lohr, J.P. *et al.*, *Arzneim.-Forsch.*, 1975, **25**, 870-873 (*metab*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1976, **98**, 3645-3655 (*Akuammigine, Tetrahydroalstonine, synth, cmr*)

Brown, R.T. *et al.*, *J.C.S. Perkin 1*, 1976, 160-162 (*Mayumbine, Tetrahydroalstonine, synth*)

Brown, R.T. *et al.*, *Chem. Comm.*, 1977, 636-638 (*synth, Akuammigine, Tetrahydroalstonine*)

Melchio, J. *et al.*, *Tet. Lett.*, 1977, 315-316 (*Mayumbine, 3-Isorauniticine, bibl, config*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 2596-2598 (*Akuammigine, 3-Isorauniticine, synth*)

Stöckigt, J. *et al.*, *Phytochemistry*, 1979, **18**, 965-971 (*biosynth*)

Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 307-311 (*Melinonine A*)

Ponglux, D. *et al.*, *Phytochemistry*, 1980, **19**, 2013-2016 (*Rauniticine, isol, uv, ir, ms*)

Gutzwiller, J. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 1663-1671 (*synth*)

Wenkert, E. *et al.*, *Acta Chem. Scand., Ser. B*, 1982, **36**, 607-611 (*stereochem, config*)

Kohl, W. *et al.*, *Z. Naturforsch., B*, 1982, **37**, 1346-1351 (*3,19-diepimer, isol, uv, pmr, ms*)

Seguin, E. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 2059-2068 (*cd*)

Phillipson, J.D. *et al.*, *Phytochemistry*, 1983, **22**, 1809-1813 (*3-Isorauniticine, Rauniticine, 3,19-diepimer, Tetrahydroalstonine N-oxide, isol*)

Massiot, G. *et al.*, *Chem. Comm.*, 1984, 715-716 (*synth*)

Lala, P.K. *et al.*, *J. Inst. Chem. (India)*, 1984, **56**, 42-47 (*ms*)

Palmisano, G. *et al.*, *J.C.S. Perkin 1*, 1985, 923-926 (*19-Epiajmalicine, Mayumbine, synth, uv, ir, pmr, ms*)

Hatakeyama, S. *et al.*, *Tet. Lett.*, 1985, **26**, 865-868 (*synth*)

Yamanaka, E. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 3713-3721 (*14α-Hydroxyrauniticine, synth, pmr*)

Naito, T. *et al.*, *Heterocycles*, 1986, **24**, 2117-2120 (*synth*)

Mekki, A.G. *et al.*, *Anal. Profiles Drug Subst.*, 1987, **16**, 731 (*rev*)

Morales-Rios, M.S. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 377-395 (*cmr*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 6133

Takano, S. *et al.*, *Chem. Comm.*, 1988, 59-60 (*synth, Tetrahydroalstonine*)

Hirai, Y. *et al.*, *J.A.C.S.*, 1988, **110**, 958-960 (*synth, Tetrahydroalstonine*)

Martin, S.F. *et al.*, *J.A.C.S.*, 1988, **110**, 5925-5927 (*Tetrahydroalstonine, synth, bibl*)

Baggiolini, E.G. *et al.*, *Tetrahedron*, 1988, **44**, 3203-3208 (*19-Epiajmalicine, Mayumbine, synth, ir, pmr, ms*)

Oppolzer, W. *et al.*, *J.A.C.S.*, 1991, **113**, 9660-9661 (*3-Isorauniticine, synth*)

Hanessian, S. *et al.*, *J.O.C.*, 1991, **56**, 2947-2949 (*19-Epiajmalicine, Mayumbine, synth*)

Honda, T. *et al.*, *J.C.S. Perkin 1*, 1993, 539-540 (*19-Epiajmalicine, Mayumbine, synth*)

Lögers, M. *et al.*, *J.A.C.S.*, 1995, **117**, 9139-9150 (*synth*)

Arbain, D. *et al.*, *Aust. J. Chem.*, 1998, **51**, 961-964 (*Rauniticine, 14-Hydroxyrauniticine, isol, pmr, cryst struct*)


Laus, G. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 831-837 (*Akuammigine, cryst struct*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AFG750

Ajmalicine hydroxyindolenine

A-213

[5802-03-9]

C₂₁H₂₄N₂O₄ 368.432


Alkaloid from *Catharanthus roseus* tissue culture (Apocynaceae).

Guéritte, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 144-148 (*isol*)

Ajmalicinial

A-214

[81737-77-1]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from *Strychnos johnsonii*. No phys. props. reported.

N¹-Methoxycarbonyl: **Ajmalicidine**

[109031-00-7]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the roots of *Rauwolfia serpentina* (Apocynaceae). Plates (MeOH/EtOAc). Mp 235-236°. [α]_D²⁰ +190 (CHCl₃). The struct. assigned is prob. incorrect. Proposed by Lounasmaa *et al* to be possibly identical with Ajmalicinine, A-215. λ_{max} 208 ; 225 ; 285 (no solvent reported).

N¹-Methoxycarbonyl, *Ac*: [109031-01-8] Needles. Mp 251-252°.

20-Epimer: **Tetrahydroalstonial**

[85717-32-4]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from *Strychnos johnsonii*. No phys. props. reported.

Stereoisomer (?): **16-Descarbomethoxy-16,17-dihydro-17-hydroxy-19-epiajmalicine** [78184-81-3]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from the bark of *Hunteria zeylanica* and the root bark of *Aspidosperma margravianum* (Apocynaceae). Mp 185°. [α]_D -86 (Py). Stereochem. unclear. Shown in both papers as having same 19-config. as Ajmalicine, but both papers refer to it as an 19-epiajmalicine. May be identical with Ajmalicinial.


Arambewela, L.S.R. *et al.*, *Phytochemistry*, 1981, **20**, 349-350 (*Descarbomethoxydihydroxyepiajmalicine*)

- Chatterjee, A. *et al.*, *J.O.C.*, 1982, **47**, 3113-3117 (*synth, pmr*)
 Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694-707 (*Descarboxymethoxydihydroxy-droxyepiajmalicine*)
 Palmisano, G. *et al.*, *J.C.S. Perkin 1*, 1985, 923-926 (*synth, pmr, ms*)
 Siddiqui, S. *et al.*, *Phytochemistry*, 1987, **26**, 875-877 (*Ajmalicine*)
 Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*Ajmalicinial, Tetrahydroalstonial*)
 Lounasmaa, M. *et al.*, *Planta Med.*, 1994, **60**, 480-481 (*Ajmalicine*)

Ajmalicine

A-215

[55529-55-0]



Absolute Configuration

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the root bark of *Cabucala striolata* and the stem bark of *Rauwolfia cumminsii* and *Rauwolfia caffra* (Apocynaceae).

11-Methoxy: Tetraphyllinine

[23924-91-6]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Rauwolfia discolor* and from the leaves of *Neisosperma glomerata* (Apocynaceae). Plates (MeOH). Mp 231-234°. [α]_D²⁰ -35 (c, 0.42 in Py).

10,11-Dimethoxy: 10,11-Dimethoxyajmalicine

[55511-11-0]

C₂₃H₃₀N₂O₆ 430.5

Alkaloid from the root bark of *Cabucala striolata* and the trunk bark of *Ochrosia moorei* (Apocynaceae).

Stereoisomer, 20,21-didehydro: 16,17-Dihydro-17-hydroxy-19-epicathenamine

[69284-02-2]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Guettarda eximia* (Rubiaceae). Isol. as an inseparable mixt. of two epimers at C-16 and/or C-17. Not named in the original papers.

Winternitz, F. *et al.*, *Phytochemistry*, 1969, **8**, 1797-1802 (*Tetraphyllinine, synth, pmr, abs config*)

Bombardelli, E. *et al.*, *Fitoterapia*, 1974, **45**, 183-187; *CA*, **83**, 25051u (*Ajmalicine, Dimethoxyajmalicine, isol, struct*)
 Kan-Fan, C. *et al.*, *Chem. Comm.*, 1978, 618-619 (*16,17-Dihydro-17-hydroxy-19-epicathenamine*)

Iwu, M.M. *et al.*, *Phytochemistry*, 1978, **17**, 1651-1654 (*isol*)

Brown, R.T. *et al.*, *Chem. Comm.*, 1979, 877-879 (*16,17-Dihydro-17-hydroxy-19-epicathenamine, synth*)

Ahond, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 193-199 (*Dimethoxyajmalicine*)

Seguin, E. *et al.*, *J. Nat. Prod.*, 1982, **45**, 738-744 (*Tetraphyllinine, isol*)


Seguin, E. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 2059-2068 (*cd*)

Nasser, A.M.A.G. *et al.*, *J. Ethnopharmacol.*, 1984, **11**, 99-117; *CA*, **102**, 3224m (*isol*)

Ajmaline, BAN, JAN

A-216

Ajmalan-17,21-diol, 9CI. Ajmaline. Gilyrytmal. Merabitol. Raugalline. Rauwolfine. Rytmalin. Tachmalin. Many other names
 [4360-12-7]



Absolute Configuration

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Rauwolfia serpentina* and most *Rauwolfia* spp., *Melodinus balansae* and *Tonduzia longifolia* (Apocynaceae). Antiarrhythmic (Class Ia) drug which functions by inhibition of glucose uptake by heart tissue mitochondria. Platelet activating factor antagonist. Antiinflammatory agent. Mp 158-160° (MeOH solvate) Mp 205-207° (anhyd.). [α]_D +144 (CHCl₃). Log P 1.26 (uncertain value) (calc).

► High doses may cause cardiac arrhythmias, coma and death. Adverse neurological, hepatic, and haematological effects also recorded. LD₅₀ (rat, orl) 360 mg/kg. LD₅₀ (rat, ipr) 94 mg/kg. AX8050000

Hydrochloride: [4410-48-4]

Amber prisms + 2H₂O. Mp 133-134° Mp 253-255° (anhyd.).

► AX8100000

Hydrochloride (1:2):

Plates. Mp 305-306° dec.

Picrate:

Plates (EtOH). Mp 126-127° Mp 223° (anhyd.).

21-O-β-D-Glucopyranoside: Rauglucine

[119365-83-2]

C₂₆H₃₆N₂O₇ 488.58

Alkaloid from *Rauwolfia serpentina* (Apocynaceae).

17-Ac: 17-O-Acetyljmaline

[19918-92-4]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from cell suspension cultures of *Rauwolfia serpentina*. Rods (Et₂O), cryst. (EtOH). Mp 150° (rods) Mp 214-215° (cryst.).

17-Ac, 21-O-β-D-glucopyranoside: Acetyljauglucine

[119329-01-0]

C₂₈H₃₈N₂O₈ 530.617

Alkaloid from *Rauwolfia serpentina* (Apocynaceae). [α]_D²⁰ +17 (c, 0.36 in EtOH).

21-Ac:

Needles (EtOAc). Mp 190-192°.

Di-Ac: [19775-56-5]

Needles (metastable) or rods. Mp 132° Mp 187-189° (double Mp).

17-O-(3,4,5-Trimethoxybenzoyl): Ajmalimine. Willcourttine

[59846-31-0]

[110941-51-0]

C₃₀H₃₆N₂O₆ 520.624

Alkaloid from *Rauwolfia obscura*, *Rauwolfia serpentina* and *Rauwolfia*

vomitaria (Apocynaceae). Needles (MeOH/EtOAc). Mp 188-189°. [α]_D²⁰ +105. Struct. of Ajmalimine revised in 2001.

17-O-Chloroacetyl: Lorajmine, INN

[47562-08-3]

C₂₂H₂₇ClN₂O₃ 402.92

Cardiac depressant, antiarrhythmic agent. Cryst. Mp 232-235°. [α]_D +27.5 (CHCl₃). Log P 2.32 (uncertain value) (calc).

17-O-Chloroacetyl; hydrochloride: Lorajmine hydrochloride, USAN. Nevergor.*Ritmos. Ritmosel. Viaductor. Win*

11831

[40819-93-0]

Cryst. Mp 230-235°. [α]_D +40 (CHCl₃).

► LD₅₀ (rat, orl) 480 mg/kg. LD₅₀ (rat, ivn) 17.3 mg/kg. AX7710000

N^d-Me: N^b-MethylajmalineC₂₁H₂₉N₂O₂[⊕] 341.472

Alkaloid from the roots of *Rauwolfia serpentina*. Amorph. powder (as chloride). [α]_D²⁸ +113 (c, 1.1 in MeOH) (chloride). λ_{max} 205 (log ε 4.4); 244 (log ε 3.86); 288 (log ε 3.4) (MeOH) (chloride).

N^d-Propyl, hydrogen tartrate: Prajmalium bitartrate, BAN, INN. Neorythmin. GT

1012. NPAB

[2589-47-1]

C₂₇H₃₈N₂O₈ 518.606

Cardiac antiarrhythmic agent. Cryst. (EtOH/Et₂O). Mp 149-152°.

► LD₅₀ (rat, orl) 54 mg/kg. LD₅₀ (rat, ivn) 3.4 mg/kg. Hepatotoxic. AX7750000

N-De-Me: Norajmaline

[23944-24-3]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from *Rauwolfia macrophylla*, *Rauwolfia obscura* and *Rauwolfia suaveolens* (Apocynaceae). Yellow-grey amorph. powder. [α]_D +36 (c, 0.67 in CHCl₃).

N-De-Me, 17-Ac: 17-O-Acetylnorajmaline

[80202-83-1]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from cell suspension cultures of *Rauwolfia serpentina*.

N-De-Me, 17-Ac, 21-O-β-D-glucopyranoside: Acetylnorrauglucine

[119365-66-1]

C₂₇H₃₆N₂O₈ 516.59

Alkaloid from *Rauwolfia serpentina* (Apocynaceae).

17-Ketone: Ajmalidine

[639-30-5]

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Rauwolfia sellowii*, *Rauwolfia maiuensis* and *Rauwolfia vomitoria* (Apocynaceae). Prisms (MeOH). Mp 241-242°. [α]_D²⁵ -80 (c, 1 in AcOH). pK_{a1} 6.3 (80% DMF). pK_{a1} 6.6 (60% DMF).

17-Ketone, O-Ac:

Needles (EtOH). Mp 218-220°. [α]_D²⁵ +242 (c, 1 in Py).

11-Hydroxy: Ajmalinol

[73012-74-5]

C₂₀H₂₆N₂O₃ 342.437

Alkaloid from *Rauwolfia vomitoria*


- (Apocynaceae). Plates (C₆H₆/MeOH). Mp 209-210°. [α]_D³⁰ +132 (CHCl₃). λ_{\max} 254 ; 294 (no solvent reported).
- 12-Hydroxy: 12-Hydroxyajmaline**
[152014-38-5]
C₂₀H₂₆N₂O₃ 342.437
Alkaloid from hairy roots of *Rauwolfia serpentina* (Apocynaceae). λ_{\max} 250 ; 290 (MeOH).
- 12-Methoxy, 17-ketone: Vomalidine**
[639-28-1]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from the roots of *Rauwolfia vomitoria*, and root bark and stems of *Rauwolfia obscura* (Apocynaceae). Prisms (Me₂CO). Mp 242-243°. [α]_D²⁰ +210 (c, 0.5 in Py). [α]_D +318 (c, 1 in CHCl₃).
- 12-Methoxy, 17-ketone, hydrochloride:**
Cryst. (EtOH). Mp 239-242°.
- 12-Methoxy, 17-ketone, O-Ac:**
Cryst. (Et₂O). Mp 252-254°. [α]_D²⁰ +195 (c, 1 in Py).
- 2-Epimer: [51019-46-6]**
Noncryst. [α]_D +55 (c, 1 in CHCl₃).
- 17-Epimer: Sandwicine. Epiajmaline**
[509-37-5]
C₂₀H₂₆N₂O₂ 326.438
Alkaloid from *Rauwolfia sandwicensis*, *Rauwolfia maiuensis* and *Rauwolfia vomitoria* (Apocynaceae). Antiarhythmic agent. Amorph. (MeOH aq. or Me₂CO/hexane). [α]_D²⁰ +174 (c, 1 in MeOH).
- 17-Epimer, hydrochloride (1:2):**
Cryst. (Me₂CO). Mp 210-213°. [α]_D +129 (MeOH).
- 17-Epimer, di-Ac:**
Cryst. (petrol). Mp 105-108°. [α]_D²⁰ +104 (c, 1 in CHCl₃).
- 20,21-Diepimer: Isoajmaline**
[6989-79-3]
C₂₀H₂₆N₂O₂ 326.438
Alkaloid from the roots of *Rauwolfia serpentina* (Apocynaceae). Prisms (MeOH aq.), plates (Et₂O). Mp 265° dec. [α]_D¹⁸ +72 (c, 0.7 in CHCl₃). p*K*_a 8.05.
- 20,21-Diepimer, hydrochloride (1:2):**
Plates. Mp 310° dec.
- 20,21-Diepimer, di-Ac:**
Plates (Et₂O). Mp 224-225°.
- 20,21-Diepimer, N⁴-Me: N^b-Methylisoajmaline**
C₂₁H₂₉N₂O₂⁺ 341.472
Alkaloid from the roots of *Rauwolfia serpentina*. Powder (as chloride). [α]_D²¹ +88 (c, 1.2 in MeOH) (chloride). λ_{\max} 204 (log ϵ 4.41); 244 (log ϵ 3.88); 288 (log ϵ 3.42) (MeOH) (chloride).
- 17,20,21-Triepimer: Isosandwicine**
[6835-90-1]
C₂₀H₂₆N₂O₂ 326.438
Alkaloid from the roots of *Rauwolfia vomitoria* (Apocynaceae). Cryst. (MeOH aq.). Mp 250° (sinters at 160°). [α]_D²⁰ +130 (c, 1.18 in CHCl₃).
- 17,20,21-Triepimer, hydrochloride:** Mp 244-245°. [α]_D²⁰ +133 (c, 1 in MeOH).
- 17,20,21-Triepimer, di-Ac:**
Cryst. (EtOH aq.). Mp 160° (sinters at

- 110°). [α]_D²⁰ +101 (c, 1 in CHCl₃).
- [31081-68-2 ; 35080-11-6 ; 110906-81-5]
- Anet, F.A.L. *et al.*, *J.C.S.*, 1954, 1242-1260 (*isol, uv, struct, synth*)
- Pakrashi, S.C. *et al.*, *J.A.C.S.*, 1955, **77**, 6687-6689 (*Ajmalidine, isol, uv*)
- Woodward, R.B. *et al.*, *Angew. Chem.*, 1956, **68**, 13-20 (*rev, struct*)
- Djerassi, C. *et al.*, *J.A.C.S.*, 1956, **78**, 1259-1260 (*Ajmalidine, struct*)
- Hofmann, A. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 1866-1874 (*Vomalidine*)
- Gorman, M. *et al.*, *Tetrahedron*, 1957, **1**, 328-337 (*Sandwicine*)
- Bartlett, M.F. *et al.*, *J.A.C.S.*, 1962, **84**, 622-630 (*abs config*)
- Bonati, A. *et al.*, *Farmaco, Ed. Sci.*, 1963, **18**, 851-863 (*Prajmalium, synth*)
- Biemann, K. *et al.*, *Tet. Lett.*, 1963, 1969-1973 (*Ajmalidine, ms*)
- Keck, J. *et al.*, *Z. Naturforsch., B*, 1963, **18**, 177-179 (*Prajmalium*)
- Biemann, K. *et al.*, *J.A.C.S.*, 1964, **86**, 4624-4628 (*Ajmaline, Vomalidine, ms*)
- Albright, J.D. *et al.*, *J.A.C.S.*, 1967, **89**, 2416-2423 (*Ajmalidine, synth, ir*)
- Masamune, S. *et al.*, *J.A.C.S.*, 1967, **89**, 2506-2509 (*synth*)
- Mashimo, K. *et al.*, *Tetrahedron*, 1970, **26**, 803-812 (*Isoajmaline, synth*)
- Ronchetti, F. *et al.*, *Phytochemistry*, 1971, **10**, 1385-1388 (*Sandwicine, Isosandwicine*)
- Koch, M. *et al.*, *Arzneim.-Forsch.*, 1972, **22**, 2079; 2085; 1973, **23**, 642 (*Prajmalium*)
- Majumdar, S.P. *et al.*, *Phytochemistry*, 1973, **12**, 1167-1169 (*Norajmaline*)
- Petter, A. *et al.*, *Arzneim.-Forsch.*, 1974, **24**, 873; 874; 876 (*rev, pharmacol, props*)
- Timmins, P. *et al.*, *Phytochemistry*, 1974, **13**, 281-282; 1997; 1976, **15**, 733-735 (*Norajmaline, Vomalidine*)
- Hubert-Brierre, Y. *et al.*, *Tetrahedron*, 1975, **31**, 3049-3054 (*synth, epimer*)
- Kuhnert-Brandstaetter, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1976, **309**, 699-706 (*polymorphs*)
- Van Tamelen, E.E. *et al.*, *Bioorg. Chem.*, 1976, **5**, 309-326 (*synth*)
- Timmins, P. *et al.*, *Planta Med.*, 1976, **29**, 283-288 (*Ajmalimine*)
- Prewo, R. *et al.*, *Acta Cryst. B*, 1978, **34**, 454-460 (*cryst struct*)
- Chatterjee, A. *et al.*, *Tet. Lett.*, 1978, 3879-3882 (*cmr*)
- Siddiqui, S. *et al.*, *J. Chem. Soc. Pak.*, 1979, **1**, 1-4; *CA*, **92**, 111204p (*Ajmalinol, isol*)
- Capra, C. *et al.*, *Farmaco, Ed. Prat.*, 1980, **35**, 49-70 (*Lorajmine, pharmacol*)
- Libot, F. *et al.*, *Phytochemistry*, 1980, **19**, 989-991 (*Isoajmaline, cmr*)
- Iwu, M.M. *et al.*, *Planta Med.*, Suppl, 1980, 13 (*Ajmalimine*)
- Stöckigt, J. *et al.*, *Plant Cell Rep.*, 1981, **1**, 36-39 (*17-O-Acetylajmaline, 17-O-Acetylnorajmaline*)
- Danieli, B. *et al.*, *Tetrahedron*, 1984, **40**, 5255-5261 (*cmr*)
- Johnston, M.D. *et al.*, *J. Het. Chem.*, 1988, **25**, 1803-1807 (*cmr*)
- Ruyter, C.M. *et al.*, *Z. Naturforsch., C*, 1988, **43**, 479-484 (*Rauhlucine, Acetylrauhglucine, Acetylnorrauhglucine*)
- Alvarez, J.L. *et al.*, *J. Cardiovasc. Pharmacol.*, 1992, **20**, 43-49 (*Prajmalium, pharmacol*)
- Köppel, C. *et al.*, *J. Chromatogr.*, 1992, **575**, 87-91 (*Ajmaline, hplc*)
- Falkenhagen, H. *et al.*, *Can. J. Chem.*, 1993, **71**, 2201-2203 (*12-Hydroxyajmaline*)
- Saeed, S.A. *et al.*, *J. Pharm. Pharmacol.*, 1993, **45**, 715-719 (*Ajmaline, pharmacol*)
- Bailey, P.D. *et al.*, *J.C.S. Perkin I*, 1993, 441-449 (*synth*)

- Martindale. The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 58; 67
- Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 7912 (*synonyms*)
- Jokela, R. *et al.*, *Planta Med.*, 1996, **62**, 577-579 (*Isoajmaline, Isosandwicine, pmr, cmr*)
- Li, J. *et al.*, *J.A.C.S.*, 1999, **121**, 6998-7010 (*synth*)
- Hanhinen, P. *et al.*, *J. Nat. Prod.*, 2001, **64**, 686-687 (*Ajmalimine*)
- Wang, T. *et al.*, *Org. Lett.*, 2001, **3**, 345-348 (*synth*)
- Itoh, A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 848-852 (*N^b-Methylajmaline, N^b-Methylisoajmaline*)
- Lewis, S.E. *et al.*, *Tetrahedron*, 2006, **62**, 8655-8681 (*rev*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AFH250; DNB000; AFH280; PNC875

Ajmalinimine

A-217

10-Acetyl-17-O-acetylajmaline

Absolute Configuration

C₂₄H₃₀N₂O₄ 410.512

Shown to be an artifact formed from 17-O-Acetylajmaline in A-216. Alkaloid from the roots of *Rauwolfia serpentina*. Elongated rods (moist MeOH). Mp 198-199°. [α]_D²⁰ +205 (c, 0.3 in CHCl₃).

Ac:

Needles. Mp 205-206°.

- Siddiqui, S. *et al.*, *Heterocycles*, 1987, **26**, 463 (*isol, uv, ir, pmr, cmr, ms, struct*)
- Lounasmaa, M. *et al.*, *Heterocycles*, 2007, **72**, 647-648

Ajmalinine

A-218

[73144-99-7]

C₂₂H₂₆N₂O₃ 366.459

Struct. unknown. Alkaloid from *Rauwolfia vomitoria* and the roots of *Rauwolfia serpentina* and *Rauwolfia sellowii* (Apocynaceae). Antihypertensive agent. Weak sympatholytic agent. Hexagonal prisms + 1H₂O (EtOAc). Mp 180-181°. [α]_D²⁵ -97. Conts. OH and OMe but no NMe. Prob conts. a 5-Methoxyindole chromophore. Props. very close to Isorauhimbine in Y-13 isol. from the same sources (Bader *et al.*, *Experientia*, 1954). λ_{\max} 273 ; 300 (no solvent reported).

Hydrochloride:

Powder (EtOH/Et₂O). Mp 240-245° dec. [α]_D⁴⁰ -44 (c, 1 in H₂O).

Methodide:

Rods (EtOH). Mp 233-234° dec.

Picrate:


Yellow powder. Mp 200-205°.

- Siddiqui, S. *et al.*, *J. Indian Chem. Soc.*, 1931, **8**, 669-680; 1932, **9**, 539-544; 1935, **12**, 37-47 (*isol*)
- Raymond-Hamet, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1953, **237**, 1435-1438 (*uv*)
- Bader, F.E. *et al.*, *Experientia*, 1954, **10**, 298 (*bibl*)

- Seba, R.A. *et al.*, *CA*, 1955, **49**, 14270e (*isol*)
 Bader, F.E. *et al.*, *J.A.C.S.*, 1955, **77**, 3547-3550 (*bibl*)
 Malik, A. *et al.*, *Pak. J. Sci. Ind. Res.*, 1979, **22**, 121-123 (*isol*)

Ajudazol B

A-219



Relative Configuration

$C_{34}H_{44}N_2O_7$ 592.731
 Prod. by *Chondromyces crocatus*. Inhibitor of mitochondrial electron transport. Amorph. solid. $[\alpha]_D^{21} +6.1$ (c, 1.34 in MeOH). λ_{max} 214 (log ϵ 4.58); 237 (log ϵ 4.54); 306 (sh); 319 (log ϵ 3.66) (MeOH).

15,29-Didehydro: Ajudazol A

$C_{34}H_{42}N_2O_7$ 590.715
 Prod. by *Chondromyces crocatus*. Inhibitor of mitochondrial electron transport. Amorph. powder. $[\alpha]_D^{21} -44.3$ (c, 1 in MeOH). λ_{max} 213 (log ϵ 4.67); 235 (log ϵ 4.59); 320 (log ϵ 3.69) (MeOH).

Jansen, R. *et al.*, *Eur. J. Org. Chem.*, 2002, 917-921 (*isol*, *pmr*, *cmr*)


Kunze, B. *et al.*, *J. Antibiot.*, 2004, **57**, 151-155 (*activity*)

Buntin, K. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 4595-4599 (*biosynth*)

AK toxin I

A-220

N-Acetyl- β -methyl-L-phenylalanine 7-carboxy-1-(2-methyloxiranyl)-2,4,6-heptatrienyl ester, 9CI
 [85146-09-4]



$C_{23}H_{27}NO_6$ 413.469
 Prod. by *Alternaria alternata* Japanese pear pathotype and *Alternaria kikuchiana*. Phytotoxin. Cryst. (MeOH). Mp 168° dec. $[\alpha]_D^{25} +164$ (c, 0.128 in MeOH). λ_{max} 285 (ϵ 27600) (MeOH) (Derep).

Nakashima, T. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 807 (*isol*, *struct*, *props*)

Uemura, I. *et al.*, *Tetrahedron*, 2002, **58**, 2351 (*synth*)

AK toxin II

A-221

[85146-10-7]
 As AK toxin I, A-220 with

R = H

$C_{22}H_{25}NO_6$ 399.443
 Prod. by *Alternaria alternata*. Phytotoxin. Cryst. (MeOH). Mp 163° dec. $[\alpha]_D^{23} +125$ (c, 0.132 in MeOH). λ_{max} 285 (ϵ 27600) (MeOH) (Derep).

Nakashima, T. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 807 (*isol*, *struct*, *props*)

Ando, K. *et al.*, *Heterocycles*, 1989, **29**, 1023 (*synth*)


Irie, H. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1451 (*synth*)

Uemura, I. *et al.*, *Tetrahedron*, 2002, **58**, 2351 (*synth*)

Akagerine

A-222

α -Ethylidene-1,2,3,3a,4,5,6,7-octahydro-7-hydroxy-3-methyl-3,7a-diazacyclohepta[jk]fluorene-5-acetaldehyde, 9CI
 [56519-07-4]



Absolute configuration

$C_{20}H_{24}N_2O_2$ 324.422
 Alkaloid from *Strychnos usambarensis*, *Strychnos gardneri*, *Strychnos jobertiana*, *Strychnos parvifolia*, *Strychnos camptoneura*, *Strychnos spinosa*, *Strychnos nigriflora* and *Strychnos decussata*. Convulsant. Platelets (hexane). Sol. MeOH, CHCl₃. Mp 188° dec. $[\alpha]_D^{20} -16.6$ (c, 1 in MeOH). λ_{max} 227 (ϵ 32360); 276 (ϵ 6600); 283 (ϵ 6610); 293 (ϵ 5370) (MeOH) (Berdy).

Me ether: 17-O-Methylakagerine

[69241-17-4]
 $C_{21}H_{26}N_2O_2$ 338.449

Alkaloid from *Strychnos dale*, *Strychnos decussata* and *Strychnos elaeocarpa* (Loganiaceae). Convulsant. Cryst. (EtOH). Mp 187-189°.

▶HM1068000

Et ether: 17-O-Ethylakagerine

$C_{22}H_{28}N_2O_2$ 352.475
 Alkaloid from the root bark of *Strychnos johnsonii* (Loganiaceae). $[\alpha]_D -13$ (c, 0.6 in MeOH). Poss. an artifact.

10-Hydroxy: 10-Hydroxyakagerine

[74765-89-2]
 $C_{20}H_{24}N_2O_3$ 340.421
 Alkaloid from leaves of *Strychnos spinosa* (Loganiaceae). Cryst. (CHCl₃/EtOAc). Mp 149-150°. $[\alpha]_D^{20} -4.6$ (c, 1 in MeOH).

10-Hydroxy, 17-Me ether: 10-Hydroxy-17-O-methylakagerine

[73360-05-1]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from *Strychnos decussata* stem bark (Loganiaceae).

▶HM1060000

Angenot, L. *et al.*, *Tet. Lett.*, 1975, 1357 (*isol*, *ir*, *uv*, *ms*, *pmr*, *cryst struct*)

Rolfesen, W. *et al.*, *Planta Med.*, 1978, **34**, 264-273 (*deriv*)

Oguakwa, J. *et al.*, *Gazz. Chim. Ital.*, 1980, **110**, 97-100 (*10-Hydroxyakagerine*)

Rolfesen, W. *et al.*, *J. Nat. Prod.*, 1980, **43**, 97-102 (*10-Hydroxyakagerine*, *10-Hydroxy-17-O-methylakagerine*)

Marini-Bettolo, G.B. *et al.*, *J. Nat. Prod.*, 1980, **43**, 717 (*isol*)

Benson, W. *et al.*, *Heterocycles*, 1981, **15**, 935 (*synth*)

Verpoorte, R. *et al.*, *Planta Med.*, 1981, **42**, 32-36 (*isol*, *uv*, *pmr*)

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*O-Ethylakagerine*)

Danieli, B. *et al.*, *J.O.C.*, 1995, **60**, 2506 (*synth*)


Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur*, *Strychnos*)

Bennasar, M.-L. *et al.*, *J.O.C.*, 1999, **64**, 9605-9612 (*synth*)

Akagerine lactone

A-223

[75667-85-5]



Absolute configuration

$C_{20}H_{22}N_2O_3$ 338.405
 Alkaloid from *Strychnos decussata* and from *Strychnos johnsonii* (Loganiaceae). Mp 184-186°.

Et ether: O-Ethylakagerine lactone


[113141-70-1]
 $C_{22}H_{26}N_2O_3$ 366.459
 Alkaloid from root bark of *Strychnos johnsonii* (Loganiaceae). $[\alpha]_D -12$ (c, 0.4 in CHCl₃).

Olaniyi, A.A. *et al.*, *J. Nat. Prod.*, 1980, **43**, 595-597 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*O-Ethylakagerine lactone*)

Akanthomycin

A-224




$C_{16}H_{25}NO_4$ 295.378
 Isol. from the entomopathogenic fungus *Akanthomyces gracilis*. Antibacterial agent. Cryst. Isol. as a mixt. of two atropisomers.

Wagenaar, M.M. *et al.*, *Org. Lett.*, 2002, **4**, 671-673 (*isol*, *pmr*, *cmr*)

Akashin C

A-225

[428517-19-5]




C₂₆H₂₅Cl₂N₃O₆ 546.405

N-Glycoside of 6,6'-Dichloroindigotin, D-353. Prod. by *Streptomyces* sp. GW 48/1497. $[\alpha]_D^{25} +3100$ (c, 0.0058 in MeOH). λ_{\max} 241 (log ϵ 4.58); 290 (log ϵ 4.43); 619 (log ϵ 4.2) (MeOH).

Maskey, R.P. et al., *Angew. Chem., Int. Ed.*, 2002, **41**, 597-599 (isol, pmr, cmr, ms)
Maskey, R.P. et al., *Nat. Prod. Res.*, 2005, **19**, 137-142 (struct)

Akirine**A-226**

[402821-48-1]

C₂₂H₃₁NO₆ 405.49

Alkaloid from *Aconitum kirinense* (Ranunculaceae). Cryst. (Me₂CO). Mp 214-217°.

Nishanov, A.A. et al., *Khim. Prir. Soedin.*, 1992, **28**, 534-537; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 466-469 (isol, pmr, ms, cryst struct)

Akuammenine**A-227**

Struct. unknown


C₂₀H₂₂N₂O₄ 354.405

Minor alkaloid from *Picralima klaineana* (Apocynaceae). Scarlet flakes (MeO-H)(as picrate). Mp 225° (picrate).

Henry, T.A. et al., *J.C.S.*, 1932, 2759-2768

Akuammicine**A-228**

Methyl 2,16,19,20-tetrahydrocuran-17-oate, 9CI

C₂₀H₂₂N₂O₂ 322.406**(-)-form** [639-43-0]

Alkaloid from *Catharanthus roseus*, *Picralima nitida*, *Picralima klaineana*, *Amsonia brevifolia*, *Alstonia angustifolia*, *Alstonia scholaris* and other *Alstonia* spp., several *Vinca* spp., *Strychnos angolensis*, *Rauwolfia volkensii*, seeds of *Hunteria congolana*, *Catharanthus microphyllus*, *Cabucala erythrocarpa* leaves, and *Pandaca ochrascens*. Mp 177.5°. $[\alpha]_D^{19} -738$ (CHCl₃). pK_a 7.45. λ_{\max} 228 (ϵ 12300); 298 (ϵ 11800); 328 (ϵ 17400) (MeOH) (Berdy).

Hydrochloride: Mp 144° (hydrate) Mp 171° (anhyd.).

Picrate: Mp 169°.

N^b-Oxide: Akuammicine N^b-oxide [60048-86-4]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia scholaris* root, *Alstonia angustifolia* stem bark, *Rauwolfia* spp. and *Tabernaemontana* spp. (Apocynaceae). Amorph. powder. Mp 200.7-200.8°. $[\alpha]_D^{22} -624.5$ (c, 0.55 in CHCl₃).

N^b-Me: Akuammicine N^b-methosalt

[100680-12-4]

C₂₁H₂₅N₂O₂[⊕] 337.441

Alkaloid from *Alstonia scholaris*, root and stem bark of *Hunteria eburnea* and leaves of *Vinca minor* (Apocynaceae). Mp 270° dec. (as chloride). $[\alpha]_D^{20} -120$ (c, 0.1 in EtOH) (chloride). CAS no. refers to chloride.

19,20S-Dihydro: 19,20-Dihydroakuammicine

[2912-08-5]

C₂₀H₂₄N₂O₂ 324.422

Minor alkaloid from leaves of *Pleiocarpa pycnantha* var. *tubicina* and *Rauwolfia caffra* (Apocynaceae). Mp 169-171°. $[\alpha]_D^{27} -636.6$ (c, 0.74 in MeOH).

19 α ,20 α -Epoxide: 19,20-Epoxyakuammicine

[173658-92-9]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from leaves of *Rauwolfia sellowii* (Apocynaceae). No phys. props. reported.

10-Hydroxy: Sewarine. 10-Hydroxyakuammicine

[27160-72-1]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from leaves of *Rauwolfia stricta* and leaves and roots of *Rhazya stricta* (Apocynaceae). Shows cytotoxic activity. Cryst. or off-white powder. Mp 245° dec. $[\alpha]_D^{20} -720$ (c, 0.1 in EtOH). λ_{\max} 220 (ϵ 10800); 311 (ϵ 14400); 340 (ϵ 11900) (MeOH).

10-Hydroxy; hydrochloride: Mp 210° dec. $[\alpha]_D^{25} -724$ (EtOH).

12-Hydroxy: Vinervine. 12-Hydroxyakuammicine

[1963-86-6]

C₂₀H₂₂N₂O₃ 338.405

Originally considered to be 11-Hydroxyakuammicine. Alkaloid from *Vinca erecta* upper parts, also obt. from cell suspension cultures of *Stemmadenia tomentosa* (Apocynaceae). Mp 154-155° dec. λ_{\max} 234 (log ϵ 4.22); 290 (log ϵ 3.86); 336 (log ϵ 4.22) (EtOH).

19,20-Dihydro, 19-hydroxy: see Echitamidine, E-30

19,20-Dihydro, hydroxy: Hydroxy-19,20-dihydroakuammicine

[59981-20-3]

C₂₀H₂₄N₂O₃ 340.421

Minor alkaloid from *Alstonia scholaris* root bark (Apocynaceae). Amorph. powder. Appears to be distinct from the 18- or 19-hydroxy compd. above. Could be identical with Echitamidine, E-30 or a stereo- or regioisomer of it.

19,20-Dihydro, 18- or 19-hydroxy:

18(19)-Hydroxy-19,20-dihydroakuammicineC₂₀H₂₄N₂O₃ 340.421

Minor alkaloid from *Alstonia scholaris* root bark (Apocynaceae). Amorph. powder. Not compared with Echitamidine, could be identical or a stereo- or regioisomer of it.

11-Methoxy: 11-Methoxyakuammicine

[54484-54-7]

C₂₁H₂₄N₂O₃ 352.432

Minor alkaloid from *Alstonia muelleriana* bark, *Alstonia lanceolifera* stem bark, *Alstonia odontophora* and other *Alstonia* spp. (Apocynaceae). $[\alpha]_D -505$ (EtOAc). Originally considered to be identical with Vinervine, but when the struct. of Vinervine was revised to 12-methoxyakuammicine it was reexamined and found to be different. λ_{\max} 232 (ϵ 11500); 305 (ϵ 5620); 324 (ϵ 6525) (MeOH).

11-Methoxy, N^b-oxide: 11-Methoxyakuammicine N^b-oxide

[94444-35-6]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. $[\alpha]_D^{25} -419.1$ (c, 2.65 in MeOH).

11-Methoxy, 19 α ,20 α -epoxide: Alstolaguamine. 11-Methoxy-19,20 α -epoxyakuammicine

[154849-51-1]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. $[\alpha]_D^{22} -73.4$ (c, 1.30 in MeOH).

12-Methoxy: Vinervinine. 12-Methoxyakuammicine

[17366-53-9]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the epigeal parts of *Vinca major* and *Vinca erecta* (Apocynaceae). Mp 190-191°. $[\alpha]_D^{20} -564$ (c, 1.877 in CHCl₃).

12-Methoxy, N^b-Me: 12-Methoxy-N^b-methylakuammicine

[123064-76-6]

C₂₂H₂₇N₂O₃[⊕] 367.467

Quaternary alkaloid from the root bark of *Alstonia congensis* (Apocynaceae). $[\alpha]_D +70$ (c, 0.15 in CHCl₃) (as 3-pyridinecarboxylate). λ_{\max} 225; 282; 289; 333 (MeOH) (3-pyridinecarboxylate).

12-Methoxy, 19 α ,20 α -epoxide: 12-Methoxy-19 α ,20 α -epoxyakuammicine

[104696-13-1]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the leaves of *Amsonia lenormandii* var. *lenormandii* (Apocynaceae). Cryst. (Et₂O). Mp 188°. $[\alpha]_D -590$ (c, 1 in CHCl₃).

19,20 α -Dihydro, 19-oxo, 20-hydroxy: LagumicineC₂₀H₂₂N₂O₄ 354.405

Alkaloid from the leaves of *Alstonia angustifolia* var. *latifolia*. Light yellow oil. $[\alpha]_D -552$ (c, 0.07 in CHCl₃). λ_{\max} 234 (log ϵ 3.79); 294 (log ϵ 3.71); 328 (log ϵ 3.85) (EtOH).

 Δ^{18} -Isomer: Angustimicine

[54086-20-3]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Amsonia angustifolia* (Apocynaceae). Struct. appears dubious. The obvious intercorrelation with Akuammicine has not been carried out.

(±)-form

Pseudakuammicine. *Pseudoakuammicine* [7344-80-1]

Alkaloid from seeds of *Picalima nitida* (Apocynaceae). Mp 187.5°.

Hydrochloride:

Minute needles (EtOH or H₂O). Mp 216°.

Henry, T.A. *et al.*, *J.C.S.*, 1927, 1950-1959; 1932, 2759-2768 (*isol*)

Aghoramurthy, K. *et al.*, *Tetrahedron*, 1957, **1**, 172 (*isol, struct*)

Lévy, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1960, 979-981 (*Akuammicine*)

Edwards, P.N. *et al.*, *J.C.S.*, 1961, 152-165 (*abs config*)

Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 1445-1449 (*isol, synth, methosalt*)

Budzikiewicz, H. *et al.*, *Tetrahedron*, 1963, **19**, 1265-1276 (*ms*)

Kump, W.G. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1497-1503 (*19,20-Dihydroakuammicine*)

Kuchenkova, M.A. *et al.*, *Izv. Akad. Nauk SSSR. Ser. Khim.*, 1965, 2152; *CA*, **64**, 11269c (*Vinervine, isol*)

Yuldashev, P.Kh. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 34-42; 1967, **3**, 310-315; 1974, **10**, 260-261; 1983, **19**, 210-212; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 25-30; 1967, **3**, 263-266; 1974, **10**, 276-277; 1983, **19**, 199-201 (*Vinervine, Vinervinine, isol, cmr, struct, bibl*)

Ahmad, Y. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1581-1583 (*Sewarine, isol, uv, ms*)

Karle, J.M. *et al.*, *Chem. Comm.*, 1972, 416-417 (*Sewarine, cryst struct*)

Burke, D.E. *et al.*, *Phytochemistry*, 1973, **12**, 1467-1474 (*11-Methoxyakuammicine*)

Böjthe-Korvath, K. *et al.*, *CA*, 1974, **81**, 136347 (*Angustimicine*)

Kutney, J.P. *et al.*, *Heterocycles*, 1975, **3**, 197-204 (*synth*)

Burke, D.E. *et al.*, *J.O.C.*, 1975, **40**, 1367-1368 (*11-Methoxyakuammicine*)

Boonchuay, W. *et al.*, *Phytochemistry*, 1976, **15**, 821 (*Alstonia scholaris alkaloids*)

Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 307-311 (*Sewarine, isol, uv, ms, ir*)

Mukopadhyay, S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 696-700 (*Sewarine, isol, pharmacol*)

Ravao, T. *et al.*, *Phytochemistry*, 1982, **21**, 2160-2161 (*11-Methoxyakuammicine, Angustimicine*)

Stöckigt, J. *et al.*, *Z. Naturforsch., C*, 1982, **37**, 857-860 (*Vinervine, isol*)

Legseir, B. *et al.*, *Phytochemistry*, 1986, **25**, 1735-1738 (*12-Methoxy-19,20-epoxyakuammicine*)

Caron, C. *et al.*, *Phytochemistry*, 1989, **28**, 1241-1244 (*12-Methoxy-N^b-methylakuammicine*)

Proksa, B. *et al.*, *Planta Med.*, 1989, **55**, 188-190 (*Akuammicine N-methosalt*)

Hu, W.-L. *et al.*, *Planta Med.*, 1989, **55**, 463-466 (*Akuammicine N-oxide*)

Amat, M. *et al.*, *J.O.C.*, 1990, **55**, 6299-6312 (*19,20-Dihydroakuammicine, synth*)

Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 249-252; 253-257 (*11-Methoxyakuammicine N^b-oxide, 11-Methoxy-19,20-epoxyakuammicine*)

Solé, D. *et al.*, *J.O.C.*, 1996, **61**, 4194-4195 (*synth*)

Batista, C.V.F. *et al.*, *Phytochemistry*, 1996, **41**, 969-973 (*epoxide*)

Bonjoch, J. *et al.*, *J.A.C.S.*, 1997, **119**, 7230-7240 (*synth*)

Ito, M. *et al.*, *J.A.C.S.*, 2001, **123**, 8003-8010 (*synth*)

Salim, A.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1591-1594 (*oxide*)


Kam, T.-S. *et al.*, *Phytochemistry*, 2004, **65**, 603-608 (*Lagumicine*)

Akuammidine

A-229

Methyl 17-hydroxysarpagan-16-carboxylate, 9CI†. Rhazine

[639-36-1]



Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Pmr and cmr spectral data revised in 1996. Alkaloid from *Picalima nitida*, *Amsonia brevifolia*, *Rauwolfia vomitoria*, *Picalima nitida*, *Rhazya stricta*, *Strychnos angolensis* and *Strychnos potatorum*. Mp 234-236° Mp 265°.

Picrate:

Yellow spheroids (EtOH). Mp 215°.

N^d-Me: Macusine C

[6801-17-8]

C₂₂H₂₇N₂O₃⁺ 367.467

Quaternary alkaloid from the bark of *Strychnos toxifera* (Loganiaceae).

Cryst. (EtOH/Et₂O) (as chloride). Mp 264-265° (260-261°) dec. (chloride).

[α]_D²⁴ -60.8 (c, 2.13 in H₂O) (chloride).

O-Ac:

Prisms (CHCl₃/EtOH). Mp 272°.

Z-Isomer (Z)-Akuammidine

[113973-31-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from roots of *Gelsemium elegans*. Needles (Me₂CO). Mp 240-242°. [α]_D¹⁶ +9 (c, 0.16 in MeOH).

10-Methoxy, N¹-Me: N^d-Methyl-10-methoxyakuammidine

[56440-63-2]

C₂₃H₂₈N₂O₄ 396.485

Alkaloid from the aerial parts of *Alstonia lanceolifera* and *Alstonia bou-lindaensis* (Apocynaceae). Mp 234-236°. [α]_D +52 (c, 1 in CHCl₃).

16-Epimer: see Polyneuridine, P-562

Chatterjee, A. *et al.*, *Chem. Ind. (London)*, 1961, 1034-1035 (*isol, uv, ir*)

Chatterjee, A. *et al.*, *J. Sci. Ind. Res., Sect. B*, 1962, **21**, 147 (*pmr*)

Silvers, S. *et al.*, *Tet. Lett.*, 1962, 339-343 (*cryst struct*)

Ohashi, M. *et al.*, *Tetrahedron*, 1963, **19**, 2241-2246 (*ms*)

Battersby, A.R. *et al.*, *J.C.S.*, 1964, 4419-4427 (*Macusine C*)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 3168-3176 (*uv, cd, ord*)

Lewin, G. *et al.*, *Phytochemistry*, 1975, **14**, 2067-2071 (*N^d-Methyl-10-methoxyakuammidine*)

Jewers, K. *et al.*, *Planta Med.*, 1980, **38**, 359-362 (*cmr*)

Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075-5094 (*Z-isomer*)


Lin, L. *et al.*, *Phytochem. Anal.*, 1990, **1**, 26 (*pmr, cmr, Z-isomer*)

Jokela, R. *et al.*, *Heterocycles*, 1996, **43**, 1015-1020 (*pmr, cmr*)

Akuammiginone

A-230

[761445-62-9]



Relative Configuration

C₂₁H₂₂N₂O₄ 366.416


Alkaloid from the bark of *Alstonia scholaris*. Amorph. light yellow powder. [α]_D²⁵ +23.4 (c, 0.5 in MeOH). Zwitterionic. λ_{max} 221 (log ε 3.7); 232 (log ε 3.69); 286 (log ε 3.36) (MeOH).

Salim, A.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1591-1594 (*isol, pmr, cmr*)

Akuammiline

A-231

Methyl 17-(acetyloxy)akuammilan-16-carboxylate, 10CI. Methyl 16-[(acetyloxy)methyl]akuammilan-17-oate, 9CI [1897-26-3]



Absolute Stereochemistry

C₂₃H₂₆N₂O₄ 394.469

Alkaloid from *Picalima nitida*, *Cono-pharyngia durissima*, *Vinca minor*, *Rauwolfia oreogiton* and *Rauwolfia vomitoria* (Apocynaceae). Cryst. (Et₂O). Mp 157-161°. [α]_D²⁴ +83 (c, 0.5 in CHCl₃).

Hydrochloride:

Cryst. + 1H₂O (H₂O or EtOH). Mp 196°. [α]_D²⁰ -29.6 (c, 3.84 in H₂O).

Methiodide:

Needles (H₂O). Mp 233°. [α]_D²⁰ -83 (c, 1.36 in EtOH).

N^d-Oxide: Akuammiline N^d-oxide

C₂₃H₂₆N₂O₅ 410.469

Alkaloid from *Kopsia griffithii*. [α]_D -144 (c, 0.13 in CHCl₃). λ_{max} 222 (log ε 4.35); 270 (log ε 3.77) (EtOH).

O-De-Ac: Rhazimol. Deacetylakuammiline. Ercinaminine

[1897-30-9]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Vinca minor*, *Picalima nitida*, *Rauwolfia vomitoria*, *Rauwolfia oreogiton* and *Rhazya stricta* (Apocynaceae). Pale yellow amorph. powder. [α]_D²¹ +19.7 (MeOH).

O-De-Ac, O-(3,4,5-trimethoxybenzoyl):

Alstolenine


[85769-33-1]

- $C_{31}H_{34}N_2O_7$ 546.619
Minor alkaloid from the leaves of *Alstonia venenata*. Amorph. $[\alpha]_D^{20}$ -70.9 (EtOH).
- 1,2 β -Dihydro-1,2 β -Dihydroakuammiline**
[77485-26-8]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from the leaves of *Rauwolfia oreogiton* (Apocynaceae). Off-white amorph. powder. $[\alpha]_D^{21}$ -9.4 (MeOH).
- 1,2 β -Dihydro, O-de-Ac: Deacetyl-1,2 β -dihydroakuammiline**
[77485-27-9]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from the leaves of *Rauwolfia oreogiton* (Apocynaceae). Mp 228-230°. $[\alpha]_D^{25}$ +99 (MeOH).
- 5-Oxo, 1,2 ζ -dihydro, O-de-Ac: Deacetyl-1,2-dihydro-5-oxoakuammiline**
[870995-65-6]
 $C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from the leaves of *Alstonia scholaris*. Powder. Mp 290-292°. $[\alpha]_D^{13}$ -147.5 (c, 0.35 in $CHCl_3$).
- 10-Hydroxy, O-de-Ac: Ercinamine. 10-Hydroxydeacetylakuammiline**
[85783-98-8]
 $C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from *Catharanthus roseus* and *Vinca erecta*. Mp 238-240°. $[\alpha]_D$ +53.
- 10-Methoxy: Raufloricine**
[38734-62-2]
 $C_{24}H_{28}N_2O_5$ 424.496
Alkaloid from *Vinca minor* and from the root bark of *Rauwolfia confertiflora* (Apocynaceae). Cryst. (Me₂CO/hexane or EtOAc). Mp 190-193°. $[\alpha]_D^{20}$ +129 (c, 1.5 in $CHCl_3$).
- 10-Methoxy, O-de-Ac: Nervobscurine. 10-Methoxydeacetylakuammiline**
[38734-63-3]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Vinca minor*. Amorph. $[\alpha]_D^{25}$ +83 (c, 0.72 in EtOH).
- 5 α ,10-Dimethoxy, O-de-Ac, O-benzoyl: Deacetyl-O-benzoyl-5,10-dimethoxyakuammiline**
 $C_{30}H_{32}N_2O_6$ 516.593
Alkaloid from *Alstonia villosa*. Solid. $[\alpha]_D^{20}$ -169.7 (c, 0.38 in MeOH).
- 5 α ,10-Dimethoxy, O-de-Ac, O-(3,4,5-trimethoxybenzoyl): 5,10-Dimethoxyalstolenine**
 $C_{33}H_{38}N_2O_9$ 606.671
Alkaloid from *Alstonia villosa*. Prisms. Mp 190-192°. $[\alpha]_D^{25}$ -213 (c, 0.82 in $CHCl_3$). λ_{max} 218 (log ϵ 4.71); 279 (log ϵ 4.22); 300 (sh) (log ϵ 4.13) (MeOH).
- 5 β ,11-Dimethoxy, O-de-Ac: Deacetyl-5 β ,11-dimethoxyakuammiline**
 $C_{23}H_{28}N_2O_5$ 412.485
Alkaloid from the roots of *Rauwolfia yunnanensis*. Amorph. yellow solid. $[\alpha]_D^{24}$ -124 (c, 0.46 in Py). λ_{max} 208 (log ϵ 4.24); 224 (log ϵ 4.2); 230 (log ϵ 4.21); 289 (log ϵ 3.53) (MeOH).
- 5 β -Hydroxymethyl: 5-Hydroxymethylakuammiline**
[77485-24-6]
 $C_{24}H_{28}N_2O_5$ 424.496
Alkaloid from the leaves of *Rauwolfia oreogiton* (Apocynaceae). Off-white

- amorph. powder. $[\alpha]_D^{21}$ -141 (MeOH).
- 16-Epimer: 16-Epiakuammiline**
[958247-21-7]
 $C_{23}H_{26}N_2O_4$ 394.469
Alkaloid from *Kopsia singapurensis*. Oil. $[\alpha]_D^{25}$ +116 (c, 0.23 in $CHCl_3$). λ_{max} 210 (log ϵ 3.99); 220 (log ϵ 4.05); 265 (log ϵ 3.35) (EtOH).
- 16-Epimer, O-de-Ac: 16-Epideacetylakuammiline**
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from the stem bark of *Kopsia deverrei*. $[\alpha]_D^{20}$ +232 (c, 0.53 in $CHCl_3$). λ_{max} 220 (log ϵ 4.22); 265 (log ϵ 3.18) (EtOH).
- 16-Epimer, O-de-Ac, N⁴-oxide: 16-Epideacetylakuammiline N⁴-oxide**
 $C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from *Kopsia griffithii*. $[\alpha]_D$ -66 (c, 0.15 in $CHCl_3$). λ_{max} 222 (log ϵ 4.12); 272 (log ϵ 3.57) (EtOH).
- Henry, T.A. *et al.*, *J.C.S.*, 1932, 2759 (*isol*)
Dugan, J.J. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 701 (*isol*, uv, ir, pmr, ms)
Savařkan, S. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 2861 (*Akuammiline*, Raufloricine, Nervobscurine)
De Maindreville, M.D. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1975, **280**, 131 (*config*)
Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 2741 (*Dihydroakuammiline*, Rhazimol, Deacetyldihydroakuammiline)
Majumder, P. *et al.*, *Phytochemistry*, 1982, **21**, 2389 (*Alstolenine*)
Gueritte, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 144 (*Ercinamine*)
Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 483; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 454 (*Ercinamine*, Ercinaminine)
Kan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 275 (*16-Epideacetylakuammiline*)
Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1235-1238 (5,10-Dimethoxydeacetylakuammilines)
Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **50**, 75-79 (*N⁴-oxides*)
Zhou, H.-U. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 2508-2512 (*Deacetyldihydrooxoakuammiline*)
Hu, X.-J. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1344-1350 (*Deacetyl-5,11-dimethoxyakuammiline*)
Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1783-1789 (*16-Epiakuammiline*)

Akuammine

Vincamajoridine
[3512-87-6]



Absolute configuration

- $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Picalima nitida* (*Picalima klaineana*), *Cabucala erythrocarpa* and *Vinca herbacea* (Apocynaceae). Needles (EtOH). Mp 255° Mp 278-280° (295°). $[\alpha]_D^{20}$ -73.4 (c, 0.87 in $CHCl_3$).
- Hydrochloride:**
Cryst. + 1H₂O. Mp 227°. $[\alpha]_D^{22}$ -26.6 (c, 0.88 in H₂O).


Picrate:

- Yellow needles (EtOH). Mp 199°.
- N⁴-Oxide: Akuammine N-oxide**
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Vinca major*. Cryst. (EtOH). Mp 178-180°. λ_{max} 243 (log ϵ 2.91); 312 (log ϵ 3.21) (EtOH).
- Ac:**
Prisms (EtOH). Mp 226°. $[\alpha]_D^{20}$ -52.1 (c, 0.64 in EtOH).
- Me ether: O-Methylakuammine**
[36101-52-7]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Rauwolfia oreogiton*, *Vinca erecta* and *Cabucala fasciculata* (Apocynaceae). Needles (Me₂CO). Mp 242-243°.
- 10-Deoxy: Pseudoakuammigine. ψ -Akuammigine**
[2447-70-3]
 $C_{22}H_{26}N_2O_3$ 366.459
Alkaloid from *Alstonia scholaris*, *Picalima klaineana* and *Picalima nitida* (Apocynaceae). Prisms (EtOH aq.), plates (C₆H₆/cyclohexane). Mp 165° (158°). $[\alpha]_D^{20}$ -53.8 (c, 3.42 in EtOH).
- 10-Deoxy; hydrochloride:**
Prisms (H₂O). Mp 183° (solvate) Mp 218° (anhyd.). $[\alpha]_D^{20}$ -15.4 (c, 1.065 in EtOH).
- 10-Deoxy, picrate:**
Yellow needles (Me₂CO). Mp 223°.
- 10-Deoxy, N⁴-oxide: Pseudoakuammigine N-oxide**
[125205-49-4]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from the stem bark of *Alstonia angustifolia* (Apocynaceae). Mp 204.3-204.4°. $[\alpha]_D^{22}$ -11.5 (c, 0.17 in $CHCl_3$).
- 10-Deoxy, 17S-hydroxy: 17-Hydroxy- ψ -akuammigine. 10-Deoxy-17-hydroxyakuammigine. 17-Hydroxypseudoakuammigine**
[79839-00-2]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Hunteria congolana* (Apocynaceae). Mp 131°. $[\alpha]_D$ -8.6 (c, 0.5 in $CHCl_3$).
- 10-Deoxy, 17S-methoxy: 17-Methoxy- ψ -akuammigine**
[84959-48-8]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Hunteria congolana* (Apocynaceae). Cryst. (Et₂O). $[\alpha]_D$ -8 (c, 0.5 in Et₂O).
- Henry, T.A. *et al.*, *J.C.S.*, 1927, 1950 (*isol*)
Joule, J.A. *et al.*, *J.C.S.*, 1962, 312 (*struct. derivs*)
Olivier, L. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 868 (*struct. abs config*)
Rakhimov, D.A. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 677; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 663 (*isol*, uv, ir, pmr, ms, *struct.*, O-Methylakuammine)
Morita, Y. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1419 (*isol*, Pseudoakuammigine)
Banerji, A. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 390 (*isol*, uv, ir, pmr, ms, Pseudoakuammigine)
Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 2741 (*isol*, O-Methylakuammine)
Vercauteren, J. *et al.*, *Tet. Lett.*, 1981, **22**, 2871 (*17-Hydroxy- ψ -akuammigine*, Pseudoakuammigine)

- Vercauteren, J. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1982, 291 (*derivs*)
 Zhukovich, E.N. *et al.*, *Khim. Prir. Soedin.*, 1989, **25**, 434-435; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 378 (*N-oxide*)
 Hu, W.-L. *et al.*, *Planta Med.*, 1989, **55**, 463 (*Pseudoakumigmine N-oxide*)


Alalagine A-233

22-Ethyl-19-hydroxy-15,16-dimethoxy-4,5-secoobscurinerivan-21-oic acid γ -lactone, 9CI
 [68346-07-6]



C₂₅H₃₂N₂O₅ 440.538
 Alkaloid from *Aspidosperma album* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 195°. [α]_D -38 (CHCl₃).

Urrea, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1978, **287**, 63 (*ir, uv, ms, pmr, struct*)

Alamaridine A-234

C₁₈H₂₀N₂O₂ 296.368

Natural-form

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Light-yellow needles (MeOH). Mp 196°. Opt. rotn. not determined due to paucity of material. λ_{\max} 226 (sh) (log ϵ 3.85); 258 (log ϵ 3.35); 286 (log ϵ 3.31) (EtOH).


(±)-form [107588-71-6]

Synthetic. Pale yellow granules (MeOH). Mp 242-244°.

Bhattacharjya, A. *et al.*, *Tetrahedron*, 1988, **44**, 3477-3488 (*isol, uv, ir, pmr, ms, synth, struct*)
 Reimann, E. *et al.*, *Monatsh. Chem.*, 1994, **125**, 1397-1406 (*synth*)

Alangimaridine A-235

12-Ethenyl-5,6,13,13a-tetrahydro-2-hydroxy-3-methoxy-8H-isoquino[2,1-b][2,7]naphthyridin-8-one, 9CI
 [77173-59-2]



C₁₉H₁₈N₂O₃ 322.363

(R)-form [178121-46-5]

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Cryst. (MeOH/CHCl₃). Mp 278°. [α]_D +429 (c, 0.35 in CHCl₃).

Ac:

Cryst. (MeOH). Mp 232°.

13,13a-Didehydro: Alangimarine

[77156-16-2]

C₁₉H₁₆N₂O₃ 320.347

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Mp 247°.

Achiral.

13,13a-Didehydro, Ac: Mp 202°.**13,13a-Didehydro, O-de-Me, O²-Me: Isoalangimarine**

[96422-49-0]

C₁₉H₁₆N₂O₃ 320.347

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Yellow needles (CHCl₃/MeOH). Mp 249°.

14-Oxo, 13,13a-didehydro, 14,15-dihydro: Alangimarionone

[96422-50-3]

C₁₉H₁₆N₂O₄ 336.346

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Yellow needles. Mp 282°.

14 ξ -Hydroxy, 14,15-dihydro: Dihydroalamarine

[97168-61-1]

C₁₉H₂₀N₂O₄ 340.378

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Pale-yellow needles (CHCl₃/MeOH). Mp 252-253°. [α]_D +347 (c, 0.35 in Py).

14 ξ -Hydroxy, 14,15-dihydro, O-de-Me, O²-Me: Dihydroisalamarine

[96422-51-4]

C₁₉H₂₀N₂O₄ 340.378

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Yellow needles (CHCl₃/MeOH). Mp 268-270°. [α]_D +140 (c, 0.07 in Py).

14 ξ -Hydroxy, 13,13a-didehydro, 14,15-dihydro: Alamarine

[77156-18-4]

C₁₉H₁₈N₂O₄ 338.362

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Yellow needles (CHCl₃/MeOH). Mp 288°. [α]_D 0 (CHCl₃).

14 ξ -Hydroxy, 13,13a-didehydro, 14,15-dihydro, O-de-Me, O²-Me: Isoalamarine

[77156-20-8]

C₁₉H₁₈N₂O₄ 338.362

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Yellow needles (MeOH/CHCl₃). Mp 301-303°.

(±)-form

Synthetic. Cryst. (MeOH/CHCl₃). Mp 242-245°.

Pakrashi, S.C. *et al.*, *Indian J. Chem., Sect. B*, 1985, **24**, 19-28 (*Alangimarine, Alangimaridine, Isoalangimarine, Alangimarionone, Alamarine, Dihydroalamarine, Dihydroisalamarine, Isoalamarine, isol, uv, ir, pmr, ms, cd, struct*)
 Jahangir, *et al.*, *Can. J. Chem.*, 1987, **65**, 2362-2368 (*Alangimarine, Alangimaridine, synth, ir, pmr, cmr, ms*)

Chowdhury, U.S. *et al.*, *Tetrahedron*, 1990, **46**, 7893-7900 (*Isoalamarine, synth*)

Itoh, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 535-538 (*Alangimaridine, Alangimarine, synth, cd, abs config*)

Reimann, E. *et al.*, *Monatsh. Chem.*, 2005, **136**, 193-209 (*Alangimarine, synth*)

Alangine A

A-236

[6793-39-1]

C₁₅H₂₃N₂O₂ 249.352

Struct. unknown. Alkaloid from *Alangium lamarckii* (Alangiaceae). Cryst. (Et₂O/hexane). Mp 85°. [α]_D -41.3 (c, 0.9 in CHCl₃). Was assigned the struct. 3-(4-methoxyphenyl)-2-(1-piperidyl)-1-propanol, but synthetic material differed from the alkaloid.

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 250°.

Methiodide:

Cryst. (Me₂CO/Et₂O). Mp 184°.

Picrate:

Cryst. (EtOH). Mp 138°.

Bhakuni, D.S. *et al.*, *J. Sci. Ind. Res., Sect. B*, 1960, **19**, 8-10; *CA*, **54**, 21168h

Kapil, R.S. *et al.*, *J. Sci. Ind. Res.*, 1961, **20**, 136; *CA*, **55**, 18792g

Alanginine

A-237


Struct. unknown. Alkaloid from the root bark of *Alangium lamarckii* (Alangiaceae). Cream-coloured amorph. solid. Mp 245-247°.

Singh, M.P. *et al.*, *Proc. Natl. Acad. Sci., India, Sect. A*, 1948, **17**, 1-6; *CA*, **45**, 10489h

Alangiobussine

A-238

[171090-88-3]



C₂₂H₂₀N₄O 356.426

Alkaloid from leaves of *Alangium bussonianum*. Cryst. (MeOH). Mp 152-154°. λ_{\max} 220 (log ϵ 4.93); 250 (sh) (log ϵ 4.49); 290 (log ϵ 4.18); 334 (log ϵ 4.31) (MeOH).

3,4-Didehydro: Alangiobussinine

[171090-89-4]

C₂₂H₁₈N₄O 354.41


Trace alkaloid from leaves of *Alangium bussonianum* (Alangiaceae). Cryst. (MeOH). Mp 188-190°.

Diallo, A.O. *et al.*, *Phytochemistry*, 1995, **40**, 975-977 (*isol, uv, pmr, cmr, ms*)

Alangiside

A-239

[34482-51-4]




- $C_{25}H_{31}NO_{10}$ 505.521
Alkaloid from the roots, leaves and fruit of *Alangium lamarckii* (best source unripe fruit) (Alangiaceae). Amorph. powder + 2.5 H₂O. Mp 187° dec. (shrinks at 164°). $[\alpha]_D^{26}$ -105 (c, 1.0 in MeOH).
- Tetra-Ac:**
Amorph. powder. $[\alpha]_D^{21}$ -50.4 (c, 1.81 in CHCl₃).
- 2'-(4-Hydroxy-3-methoxy-E-cinnamoyl): 2'-O-trans-Feruloylalangiside**
[173180-42-2]
 $C_{35}H_{39}NO_{13}$ 681.692
Alkaloid from fruits of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{24}$ -153 (c, 1.0 in MeOH).
- 2'-(4-Hydroxy-3,5-dimethoxy-E-cinnamoyl): 2'-O-trans-Sinapoylalangiside**
[173180-45-5]
 $C_{36}H_{41}NO_{14}$ 711.718
From fruits of *Alangium lamarckii* (Alangiaceae). Cryst. (H₂O). Mp 179-181°. $[\alpha]_D^{23}$ -157 (c, 1.0 in MeOH).
- 2'-O-[4-(1,3-Dihydroxypropoxy)-3-methoxy-E-cinnamoyl]: 2'-O-trans-[4-(1,3-Dihydroxypropoxy)-3-methoxycinnamoyl]alangiside**
[173180-47-7]
 $C_{38}H_{45}NO_{15}$ 755.771
From fruits of *Alangium lamarckii* (Alangiaceae). Cryst. (H₂O). Mp 166-169°. $[\alpha]_D^{25}$ -153 (c, 0.51 in MeOH).
- Me ether:**
Needles + 1.5 H₂O (EtOAc). Mp 236°.
- O-De-Me: Demethylalangiside**
[47763-23-5]
 $C_{24}H_{29}NO_{10}$ 491.494
Alkaloid from the roots of *Cephaelis ipecacuanha* (Rubiaceae) and leaves of *Alangium platanifolium* var. *trilobum* (Alangiaceae). Needles (MeOH aq.). Mp 180-182°. $[\alpha]_D^{24}$ -73 (c, 0.2 in MeOH).
- O-De-Me, 2-O-β-D-glucopyranoside:**
[408363-51-9]
 $C_{30}H_{39}NO_{15}$ 653.636
Alkaloid from the roots of *Cephaelis acuminata*. Amorph. powder. $[\alpha]_D^{27}$ -24 (c, 0.34 in MeOH). λ_{max} 206 (log ε 4.4); 231 (log ε 4.25); 242 (sh) (log ε 4.18); 280 (log ε 3.66) (MeOH).
- O-De-Me, 6'-(4-hydroxy-3-methoxycinnamoyl): 6'-O-Feruloyldemethylalangiside**
[141361-35-5, 141434-40-4]
 $C_{34}H_{37}NO_{13}$ 667.665
Alkaloid from leaves of *Alangium platanifolium* var. *trilobum* (Alangiaceae). Powder. Isol. as a mixt. of *cis*- and *trans*-forms.
- O-De-Me, 2'-(4-hydroxy-3-methoxy-E-cinnamoyl): 2'-O-trans-Feruloyldemethylalangiside**
[173180-41-1]
 $C_{34}H_{37}NO_{13}$ 667.665
From fruits of *Alangium lamarckii* (Alangiaceae). Cryst. (H₂O). Mp 175-177°. $[\alpha]_D^{27}$ -162 (c, 0.44 in MeOH).
- O-De-Me, 2'-(4-hydroxy-3,5-dimethoxy-E-cinnamoyl): 2'-O-trans-Sinapoyldemethylalangiside**
[173180-44-4]
 $C_{35}H_{39}NO_{14}$ 697.691
From fruits of *Alangium lamarckii* (Alangiaceae). Cryst. (H₂O). Mp 183.5-186°. $[\alpha]_D^{26}$ -155 (c, 0.43 in MeOH).
- O-De-Me, 6'-(4-hydroxy-3,5-dimethoxycinnamoyl): 6'-Sinapoyldemethylalangiside**
[141435-01-0, 141361-36-6]
 $C_{35}H_{39}NO_{14}$ 697.691
Alkaloid from leaves of *Alangium platanifolium* var. *trilobum* (Alangiaceae). Powder. Isol. as mixt. of *cis*- and *trans*-isomers.
- O-De-Me, O²-Me: 3-O-Demethyl-2-O-methylalangiside**
[57071-85-9]
 $C_{25}H_{31}NO_{10}$ 505.521
Alkaloid from fruits of *Alangium lamarckii* (Alangiaceae) and roots of *Cephaelis ipecacuanha* (Rubiaceae). Needles (MeOH aq.). Mp 275-277°. $[\alpha]_D^{26}$ -48 (c, 1.0 in MeOH).
- O-De-Me, O²-Me, 2'-(4-hydroxy-3-methoxy-E-cinnamoyl): 2'-O-trans-Feruloyl-3-O-demethyl-2-O-methylalangiside**
[173180-43-3]
 $C_{35}H_{39}NO_{13}$ 681.692
From fruits of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{28}$ -115 (c, 0.43 in MeOH).
- O-De-Me, O²-Me, 2'-(4-hydroxy-3,5-dimethoxy-E-cinnamoyl): 2'-O-trans-Sinapoyl-3-O-demethyl-2-O-methylalangiside**
[173180-46-6]
 $C_{36}H_{41}NO_{14}$ 711.718
From fruits of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{28}$ -126 (c, 0.39 in MeOH).
- 3-Demethoxy, 1-hydroxy: Demethylneolangiside**
[169304-90-9]
 $C_{24}H_{29}NO_{10}$ 491.494
Alkaloid from the fruits of *Alangium lamarckii*. Cryst. (MeOH aq.). Mp 193-195°. $[\alpha]_D^{28}$ -9.8 (c, 1 in MeOH). λ_{max} 231 (sh) (log ε 4.24); 239 (log ε 4.25); 287 (sh) (log ε 3.37) (MeOH).
- 3-Demethoxy, 1-hydroxy, O²-Me: Neoalangiside**
[169391-62-2]
 $C_{25}H_{31}NO_{10}$ 505.521
Alkaloid from the fruits of *Alangium lamarckii*. Amorph. powder. $[\alpha]_D^{23}$ +7.7 (c, 0.25 in MeOH). λ_{max} 230 (sh) (log ε 4.27); 234 (log ε 4.27); 286 (sh) (log ε 3.38) (MeOH).
- A^{8a,12a}-Isomer, 9R-methoxy, 13,13a-didehydro: [206875-77-6]**
 $C_{26}H_{31}NO_{11}$ 533.531
Alkaloid from the fruit of *Alangium lamarckii*. Pale yellow powder. $[\alpha]_D^{27}$ +8.5 (c, 1 in MeOH). λ_{max} 221 (log ε 4.32); 226 (log ε 4.32); 243 (sh) (log ε 4.02); 258 (sh) (log ε 3.79); 267 (log ε 3.84); 351 (log ε 4.33); 366 (sh) (log ε 4.24) (MeOH).
- A^{8a,12a}-Isomer, 9S-methoxy, 13,13a-didehydro: [206875-80-1]**
 $C_{26}H_{31}NO_{11}$ 533.531
Alkaloid from the fruit of *Alangium lamarckii*. Pale yellow needles (MeOH). Mp 219-220°. $[\alpha]_D^{20}$ -105 (c, 1 in MeOH). λ_{max} 226 (log ε 4.38); 244 (sh) (log ε 4.1); 260 (sh) (log ε 3.89); 266 (log ε 3.91); 334 (sh) (log ε 4.21); 351 (log ε 4.37); 367 (sh) (log ε 4.28) (MeOH).
- 6'-O-α-D-Xylopyranosyl: 6'-O-α-D-Xylopyranosylalangiside**
[201555-41-1]
 $C_{30}H_{39}NO_{14}$ 637.636
Alkaloid from the fruit of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{29}$ -38 (c, 0.3 in MeOH). λ_{max} 233 (log ε 4.32); 284 (log ε 3.68) (MeOH).
- 3'-O-β-D-Glucopyranosyl: 3'-O-β-D-Glucopyranosylalangiside**
[201555-37-5]
 $C_{31}H_{41}NO_{15}$ 667.663
Alkaloid from the fruit of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{26}$ -107 (c, 0.5 in MeOH). λ_{max} 233 (log ε 4.34); 284 (log ε 3.71) (MeOH).
- 6'-O-α-D-Glucopyranosyl: 6'-O-α-D-Glucopyranosylalangiside**
[201555-38-6]
 $C_{31}H_{41}NO_{15}$ 667.663
Alkaloid from the fruit of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{26}$ -33 (c, 1 in MeOH). λ_{max} 234 (log ε 4.34); 284 (log ε 3.69) (MeOH).
- 6'-O-β-D-Glucopyranosyl: 6'-O-β-D-Glucopyranosylalangiside**
[201555-36-4]
 $C_{31}H_{41}NO_{15}$ 667.663
Alkaloid from the fruit of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{26}$ -81 (c, 0.7 in MeOH). λ_{max} 233 (log ε 4.32); 284 (log ε 3.67) (MeOH).
- 6'-O-α-D-Glucopyranosyl, O-de-Me, O²-Me: 6'-O-α-D-Glucopyranosyl-3-O-demethyl-2-O-methylalangiside**
[201555-39-7]
 $C_{31}H_{41}NO_{15}$ 667.663
Alkaloid from the fruit of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{30}$ +2.4 (c, 0.7 in MeOH). λ_{max} 234 (log ε 4.32); 283 (log ε 3.65); 292 (sh) (log ε 3.55) (MeOH).
- 13a-Epimer: Isoalangiside**
[57128-91-3]
 $C_{25}H_{31}NO_{10}$ 505.521
Alkaloid from fruits of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{25}$ -118 (c, 0.86 in MeOH).
- 13a-Epimer, Me ether: Methylisoalangiside**
 $C_{26}H_{33}NO_{10}$ 519.547
Alkaloid from fruits of *Alangium lamarckii* (Alangiaceae). Amorph. powder. $[\alpha]_D^{25}$ -141 (c, 0.28 in MeOH).
- 13a-Epimer, O-de-Me: Demethylisoalangiside**
[169531-70-8]
 $C_{24}H_{29}NO_{10}$ 491.494
Alkaloid from the roots of *Cephaelis acuminata*. Amorph. powder. $[\alpha]_D^{27}$ -143 (c, 0.24 in MeOH). λ_{max} 234 (log ε 4.24); 287 (log ε 3.64) (MeOH).

13a-Epimer, O-de-Me, O²-Me: 3-O-Demethyl-2-O-methylisoalangiside

[57128-93-5]

C₂₅H₃₁N₃O₁₀ 505.521Alkaloid from fruits of *Alangium lamarekii* (Alangiaceae). Amorph. powder. [α]_D²⁸ -169 (c, 0.45 in MeOH).Shoeb, A. *et al.*, *J.C.S. Perkin 1*, 1975, 1245-1248 (isol, uv, ir, pmr, ms, struct, abs config)
Nagakura, N. *et al.*, *Chem. Comm.*, 1978, 896-897 (cmr)Höfle, G. *et al.*, *Chem. Ber.*, 1980, **113**, 566-576 (synth, uv, ir, pmr, cmr, ms)Itoh, A. *et al.*, *Phytochemistry*, 1991, **30**, 3117-3123; 1992, **31**, 1037-1040; 1994, **36**, 383-387 (Demethylalangiside, 6'-O-Feruloyldemethylalangiside, 6'-O-Sinapoyldemethylalangiside, 3-O-Demethyl-2-O-methylalangiside)Itoh, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1228-1239 (Isoalangiside, Methylisoalangiside, 3-O-Demethyl-2-O-methylisoalangiside, Neoalangiside, Demethylneoalangiside)Itoh, A. *et al.*, *Phytochemistry*, 1996, **41**, 651-656; 1997, **46**, 1225-1229 (esters, glycosyl derivs)Itoh, A. *et al.*, *Heterocycles*, 1998, **48**, 499-505 (9-methoxy derivs)Itoh, A. *et al.*, *Phytochemistry*, 2002, **59**, 91-97 (Demethylisoalangiside, O-de-Me glucoside)**Alantrypinone****A-240**

PF 1198A. Antibiotic PF 1198A

**(+)-form**C₂₁H₁₆N₄O₃ 372.382**(+)-form** [212911-06-3]Prod. by *Aspergillus terreus* PF1198 and *Penicillium thymicola*. Mycotoxin. Insecticidal agent. Amorph. solid. [α]_D²² +37 (c, 2.1 in EtOH), λ_{\max} 210 (log ϵ 3.78); 257 (log ϵ 3.15); 266 (log ϵ 3.17); 277 (log ϵ 3.08); 292 (log ϵ 2.73); 303 (log ϵ 2.77); 316 (log ϵ 2.62) (EtOH).**16-Hydroxy: (-)-Serantrypinone**

[956006-97-6]


C₂₁H₁₆N₄O₄ 388.382Prod. by *Penicillium thymicola*. Mp > 240°. [α]_D²² -12 (c, 0.12 in EtOH), λ_{\max} 218 (log ϵ 4.55); 267 (log ϵ 3.54); 278 (log ϵ 3.51); 290 (log ϵ 3.36); 305 (log ϵ 3.16); 318 (log ϵ 3.05) (EtOH).**(-)-form****16-Hydroxy: (+)-Serantrypinone. Antibiotic PF 1198B. PF 1198B**

[366785-28-6]

C₂₁H₁₆N₄O₄ 388.382Prod. by *Aspergillus terreus* PF 1198. Mp > 250°. [α]_D +21.7 (c, 0.1 in MeOH), λ_{\max} 208; 258; 266; 278; 303; 315 (MeOH).Larsen, T.O. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1154-1157 (isol, uv, pmr, cmr, cryst struct, abs config)Ariza, M.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1590-1592 (Serantrypinone)Hart, D.J. *et al.*, *J.A.C.S.*, 2001, **123**, 5892-5899 (synth, abs config)Kuriyama, T. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 3884-3887 (isol, pmr, cmr)Chen, Z. *et al.*, *J.O.C.*, 2004, **69**, 79-85 (synth)Hart, D.J. *et al.*, *Tet. Lett.*, 2007, **48**, 7069-7071 (Serantrypinone, synth, abs config)**N- β -Alanyldopamine****A-241****3-Amino-N-[2-(3,4-dihydroxyphenyl)ethyl]propanamide, 9CI**

[54653-62-2]


[58077-93-3]

C₁₁H₁₆N₂O₃ 224.259

Occurs in insects. Precursor to sclerotized insect cuticle; functions as cross-linking agent between peptide chains. No phys. props. reported.

3-O- β -D-Glucopyranoside: [96735-97-6]C₁₇H₂₆N₂O₈ 386.401

Occurs in insects. Precursor in insect cuticle sclerotization.

4-O- β -D-Glucopyranoside: [96848-78-1]C₁₇H₂₆N₂O₈ 386.401Isol. from *Manduca sexta* insects. No phys. props. reported.**N-Ac, 3-O- β -D-glucopyranoside: [88874-85-5]**C₁₉H₂₈N₂O₉ 428.438Isol. from praying mantis *Hierodula patellifera*, *Manduca sexta* and *Mantis religiosa*. No phys. props. reported.Hopkins, T.L. *et al.*, *Science (Washington, D.C.)*, 1982, **217**, 364-366 (isol, synth, biochem)Yago, M. *et al.*, *Insect Biochem.*, 1984, **14**, 7-9; 487-489 (N-Ac glucoside)Mueller, D.D. *et al.*, *Bioconjug. Chem.*, 1993, **4**, 47-53 (cmr, pmr)Hopkins, T.L. *et al.*, *Insect Biochem. Mol. Biol.*, 1995, **25**, 29-37 (3-glucoside)Merritt, M.E. *et al.*, *J.A.C.S.*, 1996, **118**, 11278-11282 (props)**Alatamide****A-242****N-[2-(4-Methoxyphenyl)ethyl]benzamide, 9CI. N-(p-Methoxystyryl)benzamide**C₁₆H₁₅NO₂ 253.3**(E)-form** [54797-23-8]Obt. from leaves of *Pleiospermium alatum* (Rutaceae) and aerial parts of *Piper guayranum* (Piperaceae). Cryst. (C₆H₆/CHCl₃), granular cryst. (CH₂Cl₂/petrol), flakes (MeOH/CHCl₃). Mp 169° Mp


178-180° Mp 190-193°. Although thought to be principally the (E)-form, Mp variations may be the result of config. inhomogeneity.

Galat, A. *et al.*, *J.A.C.S.*, 1950, **72**, 4436

(synth)

Chatterjee, A. *et al.*, *Aust. J. Chem.*, 1975, **28**, 457 (isol, ir, uv, ms)Ghosh, S. *et al.*, *Synth. Commun.*, 1987, **17**, 299 (synth, ir)Maxwell, A. *et al.*, *J. Nat. Prod.*, 1989, **52**, 411 (isol, uv, ir, pmr, ms)**Albine****A-243****Dehydroalbine**

[53915-26-7]




Absolute configuration

C₁₄H₂₀N₂O 232.325Alkaloid from *Lupinus albus* and *Lupinus termis* seeds (Fabaceae). Mp 50°. [α]_D²⁵ -103. The N-Methyl deriv. was formerly reported erroneously (see Alkaloid LC2, A-554).Perchlorate: Mp 253°. [α]_D²⁵ -76 (H₂O).**N-Formyl: N-Formylalbine**C₁₅H₂₀N₂O₂ 260.335Trace alkaloid in seeds and leaves of *Lupinus albus* (Fabaceae). Provisional identification.**5,6-Didehydro: Δ^5 -Dehydroalbine**C₁₄H₁₈N₂O 230.309Alkaloid from seeds of *Lupinus termis* (Fabaceae). Oil. [α]_D²⁵ -44.6 (c, 0.02 in MeOH).Wiewiórowski, M. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1964, **12**, 213; 217 (isol, ir)Chekhlov, A.N. *et al.*, *J. Struct. Chem. (Engl. Transl.)*, 1974, **15**, 848 (cryst struct)Wolńska-Moczydłarz, J. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1976, **24**, 613 (ir, struct)Mohamed, M.H. *et al.*, *Phytochemistry*, 1991, **30**, 3111 (Δ^5 -Dehydroalbine)Planchuelo-Ravelo, A.M. *et al.*, *Z. Naturforsch., C*, 1993, **48**, 414 (N-Formylalbine)**Albofungin****A-244****Kanchanomycin. BA 180265A. P 42-1.**

Antibiotic P 42-1

[37895-35-5]



Absolute Configuration

C₂₇H₂₄N₂O₉ 520.495Stereochem. revised in 1992. Metab. of *Actinomyces albus* var. *fungatus* and *Actinomyces tumemacerans*. Antifungal agent with antineoplastic props. Yellow cryst. (MeNO₂). Mp 304-307°. [α]_D²⁰ -670

(DMF). Log P 0.34 (uncertain value) (calc). λ_{\max} 250 (ϵ 9000); 320 (ϵ 5000); 388 (ϵ 8000) (MeOH/NaOH) (Derep). λ_{\max} 230 (ϵ 36400); 253 (ϵ 36500); 300 (ϵ 14600); 364 (sh) (ϵ 22400); 373 (ϵ 23700) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 2 mg/kg. ZD5706000

Penta-Ac: Mp 230-233°.

4-Chloro: **Chloroalbofungin**. *Albofungin B* [37891-66-0]

C₂₇H₂₃ClN₂O₉ 554.939

From *Actinomyces albus* var. *fungatus* and from *Streptomyces* spp. Possesses antifungal and antineoplastic activities. Cryst. (MeNO₂). Sol. Me₂CO, DMF, DMSO, Py, CHCl₃, dioxan, AcOH; fairly sol. MeOH, EtOH, CHCl₃, C₆H₆, butanol; poorly sol. H₂O, Et₂O, hexane. Mp 327-330°. $[\alpha]_{\text{D}}^{20}$ -560 (DMF). Log P 1.27 (uncertain value) (calc). λ_{\max} 230 (E1%/1cm 530); 257 (E1%/1cm 540); 311 (E1%/1cm 330); 365 (E1%/1cm 320) (MeOH) (Berdy). λ_{\max} 233 (ϵ 26300); 254 (ϵ 31600); 305 (ϵ 12300); 384 (ϵ 20900) (EtOH) (Berdy).

Gurevich, A.I. et al., *Tet. Lett.*, 1972, **13**, 1751-1754; 1974, **15**, 2801 (*Albofungin*, *Chloroalbofungin*, *isol*, *uv*, *pmr*)

Fukushima, K. et al., *J. Antibiot.*, 1973, **26**, 65-69 (*P42-1*)

Kogan, G.A. et al., *J. Struct. Chem. (Engl. Transl.)*, 1975, **15**, 793-799 (*cd*)


Onoprienko, V.V. et al., *Bioorg. Khim.*, 1978, **4**, 1418-1422 (*struct*)

Cooper, R. et al., *J. Antibiot.*, 1992, **45**, 444-453 (*pmr*, *cmr*, *stereochem*, *abs config*)

Alboinone

A-245

4-(Dimethylamino)-6-(1H-indol-3-yl)-2H-1,3,5-oxadiazin-2-one, 9CI [188547-40-2]




C₁₃H₁₂N₄O₂ 256.263

First naturally occurring oxadiazinone. Alkaloid from the ascidian *Dendrodoa grossularia*. λ_{\max} 210 ; 252 ; 270 ; 334 (MeOH).

Bergmann, T. et al., *Tetrahedron*, 1997, **53**, 2055-2060 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *synth*)

Albomitomycin A

Antibiotic AX 1 [110934-24-2]



Absolute Configuration

C₁₆H₁₉N₃O₆ 349.343

Quinone antibiotic. Metab. of *Streptomyces caespitosus*. Prod. by intramolecular rearrangement of Mitomycin A, M-653. Plates (CHCl₃). $[\alpha]_{\text{D}}^{23}$ -2.7 (c, 0.5 in CHCl₃). Dec. at >130°. λ_{\max} 215 (ϵ 3300); 288 (ϵ 10000) (MeOH) (Berdy).

7-*Demethoxy*, 7-*amino*: **Albomitomycin C** [111750-67-5]

C₁₅H₁₈N₄O₅ 334.331

From *Streptomyces* sp. Prisms. Mp 120° dec. $[\alpha]_{\text{D}}^{20}$ +34.7 (c, 0.3 in CHCl₃). Incorrect MF given in paper. λ_{\max} 242 (ϵ 5012); 344 (ϵ 6310) (MeOH) (Berdy).

Fukuyama, T. et al., *Tet. Lett.*, 1986, **27**, 6299 (*synth*)

Kono, M. et al., *J.A.C.S.*, 1987, **109**, 7224 (*isol*, *struct*)


Hirayama, N. et al., *Acta Cryst. C*, 1991, **47**, 409 (*cryst struct*)

Kono, M. et al., *J. Antibiot.*, 1995, **48**, 179 (*Albomitomycin C*, *isol*, *pmr*, *cmr*, *uv*)

Alborine

A-247

Pangrenine. Alkaloid PO5. Alkaloid R-K [23943-91-1]



C₂₂H₂₂NO₆⁺ 396.419

Alkaloid from *Papaver pseudocanescens*, *Papaver oreophilum*, *Papaver nudicaule* var. *leiocarpum*, *Papaver pyrenaicum*, *Papaver alborosum* and *Papaver angrenicum*. Yellow needles (MeOH) (as iodide or perchlorate). Mp 255-258° dec. (sinters from 228°) (as perchlorate) Mp 281-283° (as nitrate).

Preininger, V. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 124; 1440 (*uv*, *pmr*, *struct*, *bibl*)

Pfeifer, S. et al., *Pharmazie*, 1972, **27**, 48


Novák, V. et al., *Coll. Czech. Chem. Comm.*, 1974, **39**, 883 (*isol*)

Siddikov, D. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 442-445 (*isol*, *cryst struct*)

Alboviolaconitine D

A-248

[138842-68-9]



C₃₄H₄₂N₂O₁₀ 638.713


Alkaloid from roots of *Aconitum alboviolaceum* (Ranunculaceae).

Chen, D. et al., *Huaxue Xuebao*, 1992, **50**, 1211; *CA*, **118**, 230146m

Albovionitine

A-249

[138935-81-6]



C₂₃H₃₅NO₄ 389.534


Alkaloid from *Aconitum alboviolaceum*. Cryst. (CHCl₃). Mp 150-152°.

Zhigang, H. et al., *Phytochemistry*, 1991, **30**, 3494-3496 (*isol*, *pmr*, *cmr*, *ms*)

Alcaligin

A-250

1,8,11,18-Tetrahydroxy-1,6,11,16-tetraazacycloeicosane-2,5,12,15-tetrone, 9CI [117959-43-0]



C₁₆H₂₈N₄O₈ 404.419

Isol. from *Alcaligenes denitrificans* ssp. *xyloxydans*, *Bordetella pertussis* and *Bordetella bronchiseptica*. Siderophore. Cryst. + 2H₂O (H₂O). $[\alpha]_{\text{D}}^{27}$ +40.4 (c, 0.54 in MeOH). λ_{\max} 206 ; 426 (H₂O) (Berdy).

8,18-Dideoxy: 1,11-Dihydroxy-1,6,11,16-tetraazacycloeicosane-2,5,12,15-tetrone, 9CI. **Putrebactin** [188944-66-3]

C₁₆H₂₈N₄O₆ 372.42

Prod. by *Shewanella putrefaciens*. Siderophore.

Nishio, T. et al., *J.A.C.S.*, 1988, **110**, 8733 (*isol*, *struct*)

Nishio, T. et al., *Agric. Biol. Chem.*, 1990, **54**, 1837 (*isol*)

Bergeron, R.J. et al., *J.O.C.*, 1991, **56**, 5560 (*synth*)


Moore, C.H. et al., *J. Bacteriol.*, 1995, **177**, 1116 (*isol*)

Ledyard, K.M. et al., *J. Biol. Inorg. Chem.*, 1997, **2**, 93-97 (*Putrebactin*)

Alchorneine

A-251

1,2,3,7-Tetrahydro-1-methoxy-7,7-dimethyl-2-(1-methylethenyl)imidazo[1,2-a]pyrimidine, 9CI



Absolute configuration

C₁₂H₁₉N₃O 221.302

(R)-form [28340-21-8]

Alkaloid from *Alchornea floribunda* and *Alchornea hirtella* (Euphorbiaceae). Effective as a spasmolytic agent in dogs. Ganglioplegic parasympathomimetic agent. Strong vagolytic agent and inhibitor of intestinal peristalsis. Mp 43°. [α]_D -105 (c, 1 in CHCl₃).

Methiodide:

Cryst. (Me₂CO/MeOH). Mp 159°. [α]_D -28 (EtOH).

Cesario, M. *et al.*, *Acta Cryst. B*, 1972, **28**, 151 (cryst struct)

Fr. Pat., 1972, 2 087 982; *CA*, **77**, 88757a (isol, use)


Khuong-Huu, F. *et al.*, *Tetrahedron*, 1972, **28**, 5207 (isol, ir, uv, pmr, ms, struct, abs config)

Büchi, G. *et al.*, *J.O.C.*, 1989, **54**, 4494 (synth)

Alchorneinone

A-252

[41758-41-2]



C₁₂H₂₀N₂O₃ 240.302


Alkaloid from the leaves of *Alchornea floribunda* (Euphorbiaceae). Liq. [α]_D -66 (c, 1 in CHCl₃).

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1972, **28**, 5207 (isol, ir, pmr, ms, struct)

Alchornidine

A-253

[25801-15-4]



Preferred structure

C₁₆H₂₃N₃O₂ 289.377


Spectroscopic data do not allow a firm choice to be made between two alternative structs. Minor alkaloid from the leaves of *Alchornea javanensis* (Euphorbiaceae). Glistening needles by subl. Mp 96-97°. [α]_D -18 (c, 1.1 in CHCl₃).

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1679-1693 (isol, ir, pmr, ms, struct)

Alchornine

A-254

[25819-91-4]



C₁₁H₁₇N₃O 207.275

A major constit. of both the leaves and

bark of *Alchornea javanensis* (Euphorbiaceae). Needles (petrol/C₆H₆). Mp 134-135°. [α]_D +74 (c, 1.5 in CHCl₃).

Picrate:


Yellow prisms (EtOH). Mp 275-278°.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1679 (isol, ir, pmr, ms, struct)

Aldisine

A-255

6,7-Dihydropyrrolo[2,3-c]azepine-4,8(1H,5H)-dione, 9CI [72908-87-3]



C₈H₈N₂O₂ 164.163

Isol. from the marine sponges *Hymeniacidon aldus*, *Axinella carteri* and *Pseudaxinyssa cantharella*. Mp 275-277° (269° dec.). [α]_D -6 (c, 0.12 in MeOH). λ_{\max} 216 (ε 19100); 245 (ε 8370); 295 (ε 6770) (MeOH) (Derep).

2-Bromo- 2-Bromoaldisine

[96562-96-8]

C₈H₇BrN₂O₂ 243.059

Isol. from *Hymeniacidon aldus*, an unidentified sponge from Fiji, a *Lisodendoryx* sp. of sponge from Sri Lanka and from *Pseudaxinyssa cantharella*. Cryst. (MeOH). Mp 265° (243°). [α]_D +5 (c, 0.47 in MeOH). λ_{\max} 220 (ε 29600); 244 (sh) (ε 12100); 303 (ε 8130) (MeOH) (Derep).

Sharma, G.M. *et al.*, *Chem. Comm.*, 1980, 435 (synth, uv, pmr, cmr)

Schmitz, F.J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 47 (isol, uv, ir, pmr, cmr, ms, struct, deriv)

Utkina, N.K. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 578; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 547 (isol)


De Nanteuil, G. *et al.*, *Tetrahedron*, 1985, **41**, 6019 (isol, uv, ir, pmr, cmr, ms, cd, deriv)

Xu, X.-H. *et al.*, *J. Struct. Chem. (Engl. Transl.)*, 2001, **20**, 173-175 (cryst struct)

α-Aldotripiperidine

A-256

[642-10-4]



C₁₅H₂₇N₃ 249.398

Alkaloid from *Coelidium fourcadei* dried branches and leaves (Fabaceae). Cryst. (hexane). Mp 121° (115-118°). Prob. an artifact of thermal dec.


Schöpf, C. *et al.*, *Angew. Chem.*, 1950, **62**, 452 (synth)

Arndt, R.R. *et al.*, *J. S. Afr. Chem. Inst.*, 1968, **21**, 54; *CA*, **70**, 4342f (isol)

Aleicide B

A-257

2,3-Dimethoxy-6-(10-methoxy-3,7,11-trimethyl-2,4,7,11-tridecatetraenyl)-5-methyl-4-pyridinol, 9CI [156204-42-1]



C₂₅H₃₇NO₄ 415.572

Prod. by *Streptomyces aurantiacus*. Insecticide.

Shopotova, L.P. *et al.*, *Zh. Prikl. Khim. (Leningrad)*, 1993, **66**, 1111-1117 (isol, pmr, cmr, ms)

Aleicide C

A-258

C₂₅H₃₇NO₅ 431.571


Prod. by *Streptomyces aurantiacus*. Insecticide.

Navashin, S.M. *et al.*, *Antibiotiki (Moscow)*, 1994, **39**, 3-12

Aleurodisconitrile

A-259

[1000180-33-5]



C₁₃H₁₆N₂O₆ 296.279

Isol. from the fruiting bodies of *Aleurodiscus amorphus*. Solid. λ_{\max} 212 (log ε 4.07); 246 (log ε 3.31); 274 (log ε 2.79) (MeOH).

Decyano, 1'-oxo- Aleurodiscoester

[1000180-34-6]

C₁₂H₁₅NO₇ 285.253


Isol. from fruiting bodies of *Aleurodiscus amorphus*. Solid. [α]_D²⁵ +70 (c, 0.13 in MeOH). λ_{\max} 217 (log ε 3.57); 277 (log ε 3.18) (MeOH).

Kindler, B.L.J. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 8076-8078 (isol, cd, pmr, cmr)

Alfileramine

A-260

[73326-84-8]



C₃₀H₄₂N₂O₂ 462.674

A bishordeninylterpene alkaloid related structurally to the cannabinoids. Exists in soln. as an equilib. mixt. of 2 rotamers. Alkaloid from the leaves of *Zanthoxylum punctatum*, *Zanthoxylum chiriquinum* and *Zanthoxylum coriaceum* (Rutaceae). Mp 185-187°. Opt. inactive.

Methiodide (1:2): Mp 253-255°.

N^a-De-Me: N^a-Demethylalfileramine

[129743-97-1]

C₂₉H₄₀N₂O₂ 448.647

Alkaloid from the leaves of *Zanthoxylum coriaceum* (Rutaceae). Opt. inactive.

N^b-De-Me: N^b-Demethylalfileramine

[129743-98-2]

C₂₉H₄₀N₂O₂ 448.647

Alkaloid from the leaves of *Zanthoxylum coriaceum* (Rutaceae). Opt. inactive.

N^a,N^b-Di-de-Me: N^a,N^b-Didemethylalfileramine

[129743-95-9]

C₂₈H₃₈N₂O₂ 434.62

Alkaloid from the leaves of *Zanthoxylum coriaceum* (Rutaceae). Amorph. Opt. inactive.

Me ether: O-Methylalfileramine

[128700-82-3]

C₃₁H₄₄N₂O₂ 476.701

Alkaloid from the leaves of *Zanthoxylum chiriquinum* (Rutaceae). Oil (solidifies upon standing). Opt. inactive.

Caolo, M.A. *et al.*, *Tetrahedron*, 1979, **35**, 1487-1492 (*isol, uv, ir, pmr, ms, cryst struct*)

Swinehart, J.A. *et al.*, *Phytochemistry*, 1980, **19**, 1219-1223 (*isol*)

Marcos, M. *et al.*, *Tetrahedron*, 1989, **45**, 7477-7484 (*uv, ir, pmr, cmr, ms*)

Marcos, M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 459-461 (*O-Methylalfileramine*)

Marcos, M. *et al.*, *Phytochemistry*, 1990, **29**, 2315-2319 (*Demethylalfileramines*)

Alginidine

A-261

[1353-94-2]

Struct. unknown

C₂₇H₄₃NO₄ 445.641

Steroidal alkaloid. Alkaloid from *Fritillaria severzowii* (Liliaceae). Mp 164°. [α]_D -24.6 (MeOH).

Hydrochloride: Mp 230-232°.

Methiodide: Mp 288-289°.

Nuriddinov, R.N. *et al.*, *CA*, 1962, **57**, 15165

Alginine†

A-262

C₂₃H₃₉NO₃ 377.566

Steroidal alkaloid of unknown struct. contg. 3 OH groups. Alkaloid from *Fritillaria severzowii* (Liliaceae). CNS stimulant, adrenergic blocking agent, local anaesthetic. Mp 271-272°. [α]_D +108.5.

Hydrochloride: Mp 323-325°.


Methiodide: Mp 310-311°.

Yunusov, S. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1939, **9**, 1911; *CA*, **34**, 4070

Alihirsutine A

A-263

[147526-80-5]




C₂₅H₂₉NO₄ 407.508

Alkaloid from aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Cryst. (CHCl₃/MeOH 1:1). Mp 296-298°.

Ali, M. *et al.*, *Fitoterapia*, 1992, **63**, 243 (*isol, uv, ir, pmr, ms, struct*)

Alioline

A-264



Absolute Configuration

C₃₀H₃₆N₂O₃ 472.626

Unusual iboga-type alkaloid containing an additional C₉ terpenoid unit, the origin of which is unclear. Alkaloid from *Catharanthus roseus*. [α]_D²⁴ +105.5 (MeOH). λ_{\max} 227 (log ϵ 4.13); 283 (log ϵ 3.69); 292 (log ϵ 3.61) (MeOH).

Habib-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 2005, **60**, 870-874 (*isol, pmr, cmr*)

Aconitum nemorosum C₂₀ Alkaloid

A-265

C₂₀H₂₅NO₃ 327.422

Struct. unknown. Alkaloid from *Aconitum nemorosum*. Cryst. (Me₂CO). Mp 244-246°. Could be identical with Hetisinone (see under Hetisan-2,11,13-triol, H-202).

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 113; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 88

Aconitum nemorosum C₂₃ Alkaloid

A-266

C₂₃H₂₉NO₆ 415.485

Struct. unknown. Alkaloid from *Aconitum nemorosum*. Mp 270-272° (as hydrochloride). MF erroneously given as C₂₃H₂₉NO₃ in ref. Anal. figures given correspond to O₆.

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*,

1965, **1**, 113; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 88

Erythrophleum C₂₄ Alkaloid

A-267

C₂₄H₃₇NO₇ 451.559

Struct. unknown. Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Glass.

Hydrochloride:

Needles (EtOH/EtOAc). Mp 191° (solvate).

3-Ac: C₂₄ Amine 3 β -acetate

C₂₆H₃₉NO₈ 493.596

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae).

C₂₄ amide (rearrangement product):

Cryst. (EtOAc). Mp 186°. Obt. by treating the C₂₄ amine with 5M NaOH.

Falkiner, M.J. *et al.*, *Aust. J. Chem.*, 1975, **28**, 645-650 (*isol, pmr*)

Aconitum kirinense Alkaloid

A-268

C₂₆H₄₁NO₈ 495.612

Struct. unknown. Isol. from *Aconitum kirinense*. Mp 134.5° (as nitrate). Conts. 4 OH, 3 OMe and 1 NMe groups.

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 113-116; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 88

Aconitum koreanum Alkaloid

A-269

C₃₅H₄₁NO₁₀ 635.71

Struct. unknown. Isol. from *Aconitum koreanum*. Mp 210° (as hydrobromide).

[α]_D²⁷ -18.7 (c. 4.27 in EtOH). Species name given as coreanum in the English-language translation of the paper. No further reports to 2007. Later investigations of this species do not refer.

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 113-116; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 88-90

Kao, H.C. *et al.*, *Yaoxue Xuebao*, 1966, **13**, 186-194; *CA*, **65**, 3922g (*Aconitum koreanum constits*)

Alstonia constricta Alkaloid

A-270

Struct. unknown

C₄₃H₅₀N₄O₇ 734.891

Bisindole alkaloid. Alkaloid from the root bark of *Alstonia constricta* (Apocynaceae). Needles (Me₂CO). Mp 196-197°. [α]_D +9 (c. 3.3 in CHCl₃).

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2489-2501 (*isol, uv, ir, pmr*)

Ammodendron karelinii Alkaloid

A-271

Struct. unknown

C₁₅H₂₄N₂O 248.367


Quinolizidine alkaloid, prob. of Spar-teine group. Alkaloid from *Ammodendron karelinii* (Fabaceae). Oil.

Kushmuradov, Yu.K. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 717-718; *Chem. Nat. Compd. (Engl. Transl.)*, 604-605 (*isol*)

Aplysina archeri Alkaloid

A-272

[179523-37-6]

C₂₂H₂₀Br₄N₄O₉ 804.037

Probable abs. config. illus. Metab. from the Caribbean sponge *Aplysina archeri*. Shows antifungal activity. λ_{\max} 252 (ε 7850) (MeOH).

Ciminiello, P. *et al.*, *Tetrahedron*, 1996, **52**, 9863 (*isol, uv, ir, pmr, cmr, cd, struct*)

Artemisia rutifolia Alkaloid

A-273

C₂₂H₃₁NO₈ 437.489

Struct. unknown. Isol. from epigeal parts of *Artemisia rutifolia*. Amorph. solid. Mp 189-190°. Prob. artifact (NH₃ used in extraction). *Artemisia* is not an alkaloid-bearing genus.

Brutko, L.I. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 292; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 247

Astragalus berterianus Alkaloid

A-274

Struct. unknown. Alkaloid from *Astragalus berterianus* (Fabaceae). Mp 210°.

Gutierrez, C.P. *et al.*, *An. Fac. Quim. Farm. (Univ. Chile)*, 1960, **12**, 120; *CA*, **56**, 5121h

Buxus Alkaloid 1

A-275

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 199°. [α]_D +31 (CHCl₃). Props. similar to Buxaminol E in B-456 and *Buxus* Alkaloid B398, A-398.

Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*isol*)

Buxus Alkaloid 2

A-276

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 195°. [α]_D +92 (CHCl₃).

Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*isol*)

Buxus Alkaloid 3

A-277

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 281°. [α]_D +68 (CHCl₃).

Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*isol*)

Buxus Alkaloid 4

A-278

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 177°. [α]_D +147 (CHCl₃). Props. corresp. to Buxtauine M, B-482.

Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964,

47, 968-981 (*isol*)**Buxus Alkaloid 5**

A-279

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 156°. [α]_D +40 (CHCl₃).

Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*isol*)

C-Alkaloid I


A-280

Minimum formula. Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Prisms (Me₂CO aq.) (as picrate). Mp 194° (picrate). Blue-violet col. with Ce(SO₄)₂, becoming carmine on standing.

Kebrle, J. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 102-121; 1384-1386 (*isol*)

Callyspongia fibrosa Alkaloid

A-281



High molecular weight oligomer or polymer. Alkaloid from the Micronesian sponge *Callyspongia fibrosa*. Inhibitor of the epidermal growth factor receptor.

Davies-Coleman, M.T. *et al.*, *J.O.C.*, 1993, **58**, 5925

Chamaecytisus Alkaloid

A-282

C₁₂H₂₀N₂O 208.303

Struct. unknown. Alkaloid from *Chamaecytisus austriacus stefanoffi*, *Chamaecytisus ciliatus* and *Chamaecytisus polytrichus polytrichus* (Fabaceae). Mp 168-170°. Ir 1675 cm⁻¹. Prob. a tertiary amide.

Deili, A. *et al.*, *Farmatsiya (Sofia)*, 1977, **27**, 15-18; *CA*, **89**, 56427z (*isol*)

Colchicum luteum Alkaloid

A-283

C₁₉H₂₃NO₄ 329.395

Struct. unknown. Alkaloid from *Colchicum luteum* (Liliaceae). Mp 237-238°.

Chommadov, B. *et al.*, *Khim. Prir. Soedin.*, 1985, **810**; *Chem. Nat. Compd. (Engl. Transl.)*, 770-773 (*isol, ir, pmr, ms*)

Crotalaria retusa Alkaloid

A-284

Struct. unknown. Isol. from *Crotalaria retusa*. Cryst. (Me₂CO). Mp 130-132°.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1957, **10**, 464-473 (*isol*)

Cytisus laburnum Alkaloid

A-285

C₁₂H₂₂N₂O 210.319

Struct. unknown. Pyrrolizidine alkaloid. Alkaloid from *Cytisus laburnum* (*Laburnum laburnum*) (Fabaceae). Mp 128-129°. [α]_D¹⁹ +18.6 (EtOH). Prob. conts. one NH (?lactam) group, no NMe group.

Galinovsky, F. *et al.*, *Sci. Pharm.*, 1953, **21**, 256; *CA*, **49**, 6977

Cytisus ruthenicus Alkaloid

A-286

C₁₅H₁₆N₂O₅ 304.302

Prob. a quinolizidine (Sparteine-group) alkaloid. Struct. unknown. Alkaloid from *Cytisus ruthenicus* (Fabaceae). Mp 171.5-173°. [α]_D¹⁸ -15.26. Conts. *N*-Me group. Co-occurs with Sparteine.

Hydroiodide: Mp 231-233°.

Perchlorate: Mp 198-200°.

Dipicrate: Mp 199-201°.

Alekseev, V.S. *et al.*, *Farm. Zh. (Kiev)*, 1967, **22**, 59-60; *CA*, **67**, 54325m (*isol*)

Dendrobates Alkaloid 185

A-287

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula not known; m/e 185 (1), 170 (100). Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 197

A-288

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula not known; m/e 197 (1), 180 (100), 126 (35). Cannot be hydrogenated.


Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 203

A-289

Decahydro-2-(2-penten-4-ynyl)quinoline



R = CH₂CH=CHC≡CH, R' = H

C₁₄H₂₁N 203.327

Tentative struct. A major alkaloid in skin extracts of *Dendrobates fulguritus*, trace constit. in *Dendrobates auratus*, *Dendrobates histrionicus*, *Dendrobates minutus* and *Dendrobates pumilio* (Dendrobatidae). m/e 203(1), 202(2), 138(100). H₆-deriv., m/e 209, 138. Does not appear to form an *N*-Ac deriv.

Dihydro (?): **Dendrobates Alkaloid 205**

C₁₄H₂₃N 205.342

Trace alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Tentative mol. formula. m/e 205(1), 204(2), 138(100).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)
Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 217 A-290C₁₅H₂₃N 217.353

Struct. unknown. A major alkaloid from skin extracts of one population of *Dendrobates bombetes* and *Dendrobates minutus*; minor constit. in *Dendrobates tricolor* (Dendrobatidae). m/e 217 (2), 216 (3), 152 (C₁₀H₁₈N, 100). H₆-deriv, m/e 223, 152. Did not form an N-Ac deriv.

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 225 A-291C₁₄H₂₇NO 225.373

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates tinctorius* (Dendrobatidae). Mol. formula tentative; m/e 225 (3), 224 (6), 208 (2), 168 (100), 152 (25). Possibly a hydroxypumiliotoxin C. Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 233 A-292C₁₆H₂₇N 233.396

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates* sp. (Colombia) (Dendrobatidae). Mol. formula tentative; m/e 233 (2), 232 (2), 166 (100). Probably higher homologue of *Dendrobates* Alkaloid 223B, A-404 with 2 double bonds in the R side chain. H₄-deriv, m/e 237, 166.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 295 A-293C₁₉H₃₇NO 295.507

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula tentative; m/e 295 (3), 278 (4), 138 (100).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 301 A-294C₂₁H₃₅N 301.514

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates azureus* (Dendrobatidae). Mol. formula tentative; m/e 301 (<1), 260 (100). Possibly a member of Pumiliotoxin-C class with an allyl R-substituent and a C₉H₁₅ R'-substituent, see *Dendrobates* Alkaloid 167A, A-354.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 325 A-295C₁₉H₃₅NO₃ 325.49

Struct. unknown. A major alkaloid from skin extracts of *Dendrobates fulguritus* and *Dendrobates minutus* (Dendrobatidae). Mol. formula tentative; m/e 325 (3), 308 (12), 280 (3), 210 (2), 182 (56), 114 (16), 112 (9), 70 (100). Forms H₂-deriv, m/e similar to H₄-deriv of Pumiliotoxin B.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 351 A-296C₂₁H₃₇NO₃ 351.528

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula tentative; m/e 351 (6), 350 (2), 336 (4), 152 (38), 138 (65), 70 (100). Probably a member of the Pumiliotoxin A class, but the large peak at 138 is atypical.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Fritillaria Alkaloid 5 A-297

Fritillaria Base 5

Struct. unknown. Alkaloid from *Fritillaria raddeana* (Liliaceae), occurring with Raddeamine, R-7. Mp 245-247°.


Aslanov, Kh.A. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 579; *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 623

Fritillaria Alkaloid 6 A-298

Fritillaria Base 6

Struct. unknown. Alkaloid from *Fritillaria raddeana* (Liliaceae). Mp 197-199° (as hydrochloride).

Aslanov, Kh.A. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 579; *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 623

Fromia monilis Alkaloid A-299C₂₃H₄₉N₃O₃ 415.658**(S)-form** [162340-03-6]

Alkaloid from the New Caledonian starfish *Fromia monilis*. Cytotoxic agent. [α]_D +3.5.

Palagiano, E. *et al.*, *Tetrahedron*, 1995, **51**, 3675 (*isol, pmr, cmr, struct*)

Galanthus caucasicus Alkaloid A-300C₁₇H₁₉NO₄ 301.341

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Galanthus caucasicus*

(Amaryllidaceae). Cryst. (H₂O). Mp 214°. [α]_D²⁰ +94.3 (c, 0.71 in MeOH).

Tsakadze, D.M. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 331; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 281 (*isol*)

Gentiana Alkaloid I A-301C₁₃H₁₆N₂O₃ 248.281

Struct. unknown. Monoterpene alkaloid. Alkaloid from *Gentiana asclepiadea* and *Gentiana punctata* (Gentianaceae). Mp 183-187°.

Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1965, **18**, 947-949; *CA*, **64**, 10084h (*isol, uv, ir*)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (*rev*)

Gentiana Alkaloid V A-302C₁₀H₉NO₂ 175.187

Struct. unknown. Monoterpene alkaloid. Alkaloid from *Gentiana asclepiadea*, *Gentiana bulgarica*, *Gentiana cruciata*, *Gentiana lutea* (yellow gentian) and *Gentiana punctata* (Gentianaceae). Mp 240°.


Marekov, N. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1965, **18**, 999-1002; *CA*, **64**, 11270a (*isol, ir*)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (*rev*)

Gynura segetum Alkaloid A-303C₁₈H₂₅NO₅ 335.399

Pyrrrolizidine alkaloid. Struct. unknown. Possibly related to Seneciphylline, S-240. Alkaloid from *Gynura segetum*. Antimalarial agent. Cryst. Mp 223-224°.

Tang, S.-R. *et al.*, *Zhongcaoyao*, 1980, **11**, 193-195; *CA*, **94**, 36187w

Heliotropium europaeum Alkaloid A-304C₂₄H₃₇N₂O₆[⊕] 449.566


Dimeric pyrrolizidine alkaloid, originally thought to be a C₁₂ monomer. Alkaloid from *Heliotropium europaeum* (Boraginaceae). Prisms (Me₂CO/MeOH) (as chloride). Mp 158-159° dec. (as chloride). Results from alkylation of Heliotrine, H-95 by dehydroheliotrine in the plant. Does not appear to be an artifact.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1954, **7**, 287 (*isol*)


Culvenor, C.C.J. *et al.*, *Tet. Lett.*, 1969, 3603 (*pmr, struct*)

Helleborus Alkaloid V A-305C₂₅H₄₃NO₆ 453.618Struct. unknown. Alkaloid from *Helleborus viridis*. Mp 267-268° (browns from 210°). V. weak base.Keller, O. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1928, **266**, 545-572**Hovea longipes Alkaloid** A-306Struct. unknown. Alkaloid from the whole plant of *Hovea longipes* (Fabaceae). Cryst. (EtOH/HCl)(as hydrochloride). Mp 238-239° (hydrochloride).Fitzgerald, J.S. *et al.*, *An. Quim.*, 1972, **68**, 737 (isol)**Jacmaia incana Alkaloid** A-307

[78869-20-2]

C₂₃H₂₇NO₇ 429.469Alkaloid from the aerial parts of *Jacmaia incana* (Asteraceae). Gum. [α]_D²⁴ -12 (c, 0.9 in CHCl₃). Closely related to the Senampelines (see Senampelines, S-233).Bohlmann, F. *et al.*, *Phytochemistry*, 1981, **20**, 831-832 (ir, pmr, ms, struct)**Leuconotis Alkaloid 376** A-308

[131242-30-3]

C₂₃H₂₄N₂O₃ 376.454Alkaloid from *Leuconotis griffithii* and *Leuconotis eugenifolia*.Goh, S.H. *et al.*, *Tetrahedron*, 1989, **45**, 7899 (isol, uv, ir, pmr, cmr, ms, struct)**Leuconotis griffithii Alkaloid 308** A-309Struct. and MF unknown. Alkaloid from *Leuconotis griffithii*.Goh, S. *et al.*, *Tetrahedron*, 1989, **45**, 7899-7820 (isol, uv, pmr, cmr, ms)**Lycopodium Alkaloid V** A-310

Lycopodium Base V

C₁₇H₂₇NO₂ 277.406Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).Methiodide: Mp 304°. [α]_D²⁰ -66.

Methochloride: Mp 258°.

Methoperchlorate: Mp 321°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (isol, uv)**Lycopodium Alkaloid X** A-311

Lycopodium Base X

C₁₇H₂₅NO₃ 291.389Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).Methiodide: Mp 315°. [α]_D¹⁸ -30.9.

Methochloride: Mp 269°.

Methoperchlorate: Mp 335°.

Methopicate: Mp 151°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (isol, uv)**Lycopodium Alkaloid 168** A-312

Lycopodium Base 168

C₁₆H₂₃NO₃ 277.363Struct. unknown. Trace alkaloid from *Lycopodium clavatum* (Lycopodiaceae). Mp 168°.Rodewald, W.J. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1977, **51**, 1271 (isol, ir, ms)**Lycopodium Alkaloid 258** A-313

Lycopodium Base 258

Molecular formula not known; MW = 263. Struct. unknown. Trace alkaloid from *Lycopodium clavatum* (Lycopodiaceae). Mp 258°.Rodewald, W.J. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1977, **51**, 1271 (isol)**Lycopodium annotinum Alkaloid†** A-314Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).

Perchlorate:

Cryst. (H₂O). Mp 234° dec.

Picrate: Mp 241-241.5° dec.

Methiodide:

Cryst. (EtOH). Mp 261° dec.

Bertho, A. *et al.*, *Chem. Ber.*, 1952, **85**, 663 (isol)**Lycopodium annotinum Alkaloid†** A-315Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).

Picrate:

Yellow rosettes (EtOH). Mp 121° dec. (sint. at 110°).

Methiodide:

Cryst. (EtOH). Mp 290°.

Bertho, A. *et al.*, *Chem. Ber.*, 1952, **85**, 663 (isol)**Lycopodium clavatum Alkaloid** A-316Struct. unknown. Alkaloid from *Lycopodium clavatum* var. *megastachyon* (Lycopodiaceae). Mp 261-263°.Ayer, W.A. *et al.*, *Can. J. Chem.*, 1962, **40**, 2088 (isol)**Lycopodium thyoides Alkaloid** A-317C₁₈H₂₇NO₃ 305.416Struct. unknown. Alkaloid from *Lycopodium thyoides* and *Lycopodium contiguum* (Lycopodiaceae).

Perchlorate:

Cryst. (Me₂CO). Mp 210-212°.Ayer, W.A. *et al.*, *Phytochemistry*, 1974, **13**, 653 (isol, ir, ms)**Mantella Alkaloid 161** A-318C₉H₁₁N₃ 161.206Struct. unknown. Trace alkaloid from skin extracts of the Madagascan frog *Mantella laevigata*. Minor alkaloid in *Mantella* sp. cf. *madagascariensis*.Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ir, ms)**Mantella Alkaloid 191** A-319Struct. and MF unknown. Possibly related to Precoccinelline. Trace alkaloid from skin extracts of the Madagascan frog *Mantella betsileo*.Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ms)**Mantella Alkaloid 317** A-320

Homopumiliotoxin 317

Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascan frog *Mantella betsileo*.Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ms)**Mucuna pruriens Alkaloid X** A-321

Mucuna pruriens Base X

C₁₁H₂₅NO₃ 219.323Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Mp 94-95°.

Picrate: Mp 130-132°.

Rakhit, S. *et al.*, *Indian J. Pharm.*, 1956, **18**, 285; *CA*, **52**, 5748e (isol)**Pachysandra Alkaloid V** A-322

Pachysandra Base V

Steroidal alkaloid. Struct. unknown. Isol. from *Pachysandra terminalis* (Buxaceae). Mp 218-221°.Kikuchi, T. *et al.*, *Tet. Lett.*, 1964, 1817Kikuchi, T. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 215; *CA*, **67**, 32888v**Pancreatium Alkaloid 6** A-323

Pancreatium Base 6

C₁₆H₁₇NO₄ 287.315Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Pancreatium maritimum* (Amaryllidaceae). Needles (MeOH). Mp 267-270°. [α]_D²¹ -184 (c, 1.04 in MeOH).Sandberg, F. *et al.*, *J. Nat. Prod.*, 1963, **26**, 78-90 (isol, ir)**Pancreatium Alkaloid 31** A-324

Pancreatium Base 31

C₁₈H₂₁NO₅ 331.368Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Pancreatium maritimum* (Amaryllidaceae). Prisms (Me₂CO). Mp 213-216° dec. [α]_D²¹ -61 (c, 1.23 in CHCl₃).

Sandberg, F. *et al.*, *J. Nat. Prod.*, 1963, **26**, 78-90 (*isol, ir*)

Pancratium sickenbergi Alkaloid A-325

C₉H₁₃NO₃ 183.207

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Pancratium sickenbergi* (Amaryllidaceae). Mp 161° dec.

Fahmy, I.R. *et al.*, *J. Pharm. Sci. U.A.R.*, 1960, **1**, 133-147; *CA*, **56**, 10280i (*isol*)

Papaver rhoeas Alkaloid A-326

C₁₉H₁₉NO₆ 357.362

Struct. unknown. Alkaloid from *Papaver rhoeas*. Cryst. (CHCl₃/EtOH). Mp 208-209°. [α]_D²³ +235 (CHCl₃).

Slavik, J. *et al.*, *Chem. Listy*, 1958, **52**, 1957-1964; *CA*, **53**, 1640h (*isol*)

Pseudaxinyssa Alkaloid A-327

C₉H₁₄BrN₅O 288.146

Struct. unknown. Isol. from the marine sponge *Pseudaxinyssa cantharella*. Amorph. [α]_D +15.

De Nanteuil, G. *et al.*, *Tetrahedron*, 1985, **41**, 6019-6033

Rhynchosia pyramidalis Alkaloid A-328

Struct. unknown. Alkaloid from *Rhynchosia pyramidalis* (Fabaceae). Mp 132° (as picrate).

Ristic, S. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 510; *CA*, **57**, 14180e

Rosmarinus officinalis Alkaloid 2 A-329

C₂₀H₂₇NO₄ 345.438

Struct. unknown. Isol. from rosemary (*Rosmarinus officinalis*) using NH₃ during extraction. Cryst. (toluene). Mp 197-198° dec. [α]_D +36 (c, 1.48 in dioxan). Artifact.

Hydrochloride:

Cryst. (H₂O). Mp 214-216° dec.


N,O,O-Tri-Ac:

Cryst. (MeOH). Mp 291-292° dec.

Yakhontova, L.D. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 140; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 118

Sceletium namaquense Alkaloid A-330

[82545-08-2]



C₂₀H₂₆N₂O₃ 342.437

Compd. not named in reference. Alkaloid

from *Sceletium namaquense* (Aizoaceae). Oil. Related to *Sceletium* Alkaloid A₄, A-346.

Jeffs, P.W. *et al.*, *J.O.C.*, 1982, **47**, 3611 (*isol, ir, cd, pmr, ms, struct*)

Senecio borysthenicus Alkaloid A-331

C₁₈H₂₃NO₅ 333.383

Struct. unknown. Pyrrolizidine alkaloid. Alkaloid from *Senecio borysthenicus* (Asteraceae). Mp 195° dec. Conts. an OH group.

Red'ko, A.L. *et al.*, *CA*, 1959, **53**, 20695

Senecio renardi Alkaloid 4 A-332


Senecio renardi Base 4

Struct. unknown. Alkaloid from *Senecio renardi* (Asteraceae). Cryst. (Me₂CO). Mp 176-178°.

Danilova, A.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1950, **20**, 1921; *CA*, **45**, 2960

Sophora Alkaloid 9 A-333

Sophora Base 9



C₁₆H₂₈N₂O₃ 296.409

Struct. of acyl group not known. Alkaloid from *Sophora alopecuroides* (Fabaceae). Related to Sophorine in O-59.

Kuchkarov, S. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 413-414; *Chem. Nat. Compd. (Engl. Transl.)*, 364-365

Strepeliopsis strepelioides Alkaloid A-334

Strepeliopsis Bisindole base

C₃₈H₄₆N₄ 558.808

Struct. unknown. Alkaloid from the roots of *Strepeliopsis strepelioides* (Apocynaceae). Mp 158-171°. [α]_D²⁰ +210 (CHCl₃).

Laguna, A. *et al.*, *Planta Med.*, 1984, **50**, 285-288 (*isol, uv, ir, ms*)

Thermopsis lanceolata Alkaloid A-335

Struct. unknown. Alkaloid from *Thermopsis lanceolata* epigeal parts (Fabaceae). Cryst. (Me₂CO). Mp 235°. [α]_D +121.8 (c, 1 in CHCl₃).

Vinogradova, V.I. *et al.*, *Khim. Prir. Soedin.*, 1971, 463-466; *Chem. Nat. Compd. (Engl. Transl.)*, 440 (*isol*)

Ulex europaeus Alkaloid A-336

C₁₅H₂₀N₂O₅ 308.333

Struct. unknown. Alkaloid from shoots of *Ulex europaeus* (Fabaceae). Lustrous plates (petrol). Mp 170°.

Clemo, G.R. *et al.*, *J.C.S.*, 1935, 10 (*isol*)

Uncaria callophylla Alkaloid A-337

Bisindole alkaloid; probably derived

from Gambirine, G-13 and Pseudoyohimbine. Struct. unknown. Minor alkaloid from the leaves of *Uncaria callophylla* (Rubiaceae). Yellow powder. Mp 330°. Mol. formula not reported.

Goh, S.H. *et al.*, *Phytochemistry*, 1985, **24**, 880-881 (*isol, uv, ir, pmr*)

Ungernia severtzovii Alkaloid A-338

C₁₇H₁₉NO₃ 285.342

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the leaves of *Ungernia severtzovii* (Amaryllidaceae). Mp 186-187°. Opt. inactive.

Nitrate: Mp 245-246°.

Picrate: Mp 241-243°.

Smirnova, L.S. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1964, **154**, 171-173; *CA*, **60**, 9324b (*isol*)

Veratrum Alkaloid X A-339

Steroidal alkaloid. Struct. unknown.

Isol. from *Veratrum lobelianum*

(Liliaceae). Needles (MeOH).

Mp 215-217°. ir shows C=C, OH and NH₂ groups.

Shinkarenko, A.L. *et al.*, *Khim. Prir. Soedin.*, 1966, **2**, 293; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **2**, 239

Voacanga thouarsii Alkaloid I A-340

[50924-03-3]

C₄₃H₄₈N₄O₆ 716.875

Bisindole alkaloid. Struct. unknown.

Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Amorph. Intense blue col. with HNO₃. λ_{max} 223 ; 264 ; 300 ; 327 (EtOH).

Rolland, Y. *et al.*, *Phytochemistry*, 1973, **12**, 2039-2042 (*isol, uv, ir, pmr*)

Zephyranthes Alkaloid A-341

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Zephyranthes robusta* and *Zephyranthes sulphurea* (Amaryllidaceae). Stout prisms (MeOH). Mp 252-254°.

Rao, R.V.K. *et al.*, *Indian J. Pharm.*, 1969, **31**, 62-63; 86-87; *CA*, **71**, 109780w; 128623p (*isol*)

Alkaloid 1 (Calabash curare) A-342

C₂₀H₂₁N₂[⊕] 289.399

Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 239-243° (as chloride). [α]_D²⁰ -134 (c, 8.95 in H₂O) (chloride). Deep-blue col. with Ce(SO₄)₂.

Wieland, T. *et al.*, *Chem. Ber.*, 1952, **85**, 731-743

Alkaloid 2 (Calabash curare) A-343

Struct. unknown. Quaternary alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). [α]_D²⁰ -659 (as chloride). Red-violet col. with Ce(SO₄)₂.

▶ Toxic.

Wieland, T. *et al.*, *Chem. Ber.*, 1952, **85**, 731

Lycopodium Alkaloid IV A-344


Lycopodium Base IV

C₁₆H₂₃NO 245.364Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).Methiodide: Mp 265°. [α]_D²⁰ -150.1.

Methochloride: Mp 210°.

Methoperchlorate: Mp 270°.

Methopicrate: Mp 230-232°.

Achmatowicz, O. et al., *Pol. J. Chem. (Roc. Chem.)*, 1958, **32**, 485-498 (*isol, uv*)**Daphniphyllum Alkaloid A₁** A-345C₂₃H₃₃NO₃ 371.519Struct. unknown. Alkaloid from the bark and leaves of *Daphniphyllum macropodum* (Daphniphyllaceae). Mp 225-226° (as methiodide).Toda, M. et al., *Tetrahedron*, 1972, **28**, 1477 (*isol, ir, pmr, ms*)**Scelletium Alkaloid A₄** A-3463a-(3,4-Dimethoxyphenyl)-2,3,3a,4,5,9b-hexahydro-1-methyl-1H-pyrrolo[2,3-f]quinoline, 9CI
[35135-35-4]


Absolute Configuration

C₂₀H₂₄N₂O₂ 324.422Alkaloid from *Scelletium namaquense* and *Scelletium tortuosum* (Aizoaceae). Prisms (EtOAc). Mp 153.3-154.5°. [α]_D +131 (MeOH).Luhan, P.A. et al., *J.C.S. Perkin 2*, 1972, 2006 (*cryst struct*)Jeffs, P.W. et al., *J.O.C.*, 1974, **39**, 2703 (*isol, uv, ir, pmr, ms, cd, struct*)Martin, N.H. et al., *Org. Mass Spectrom.*, 1976, **11**, 1 (*ms*)Stevens, R.V. et al., *Acc. Chem. Res.*, 1977, **10**, 193 (*synth*)Kamikubo, T. et al., *Chem. Comm.*, 1998, 783-784 (*synth, abs config*)Yamada, O. et al., *Tet. Lett.*, 1998, **39**, 7747-7750 (*synth*)Hayashi, M. et al., *Tet. Lett.*, 2002, **43**, 1461-1464 (*synth*)**Abrus precatorius Alkaloid A** A-347Struct. unknown. Alkaloid from seeds of *Abrus precatorius*. Mp 143° dec.Tung, Y.-C. et al., *C.A.*, 1960, **55**, 17770g; **61**, 9777c; *Biol. Abstr.*, **36**, 65720**Buxus Alkaloid A1** A-348Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (CH₂Cl₂). Mp 195-197°. Contains NMe₂ group at C-3.Huong, L. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (*isol, ms*)**Buxus Alkaloid A2** A-349Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO/MeOH). Mp 138-142°.Huong, L. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (*isol, ms*)**Buxus Alkaloid A3** A-350Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO/MeOH). Mp 272-274°. [α]_D²² +20 (c, 0.25 in EtOH).Huong, L. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (*isol, ms*)**Buxus Alkaloid A4** A-351Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (CH₂Cl₂). Mp 252-256°. [α]_D²² -109 (c, 0.51 in MeOH).Huong, L. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (*isol, ms*)**Buxus Alkaloid A5** A-352C₄₀H₆₀N₂O₇ 680.923Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (CH₂Cl₂). Mp 125-127°. [α]_D²² -10 (c, 0.3 in CH₂Cl₂).Huong, L. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (*isol, ms*)**Chondrodendron limaciifolium Alkaloid A** A-353C₃₆H₃₈N₂O₆ 594.706Struct. unknown. Bisbenzylisoquinoline alkaloid. Alkaloid from the wood of *Chondrodendron limaciifolium* (Menispermaceae). Thin plates or flattened needles (Py, CHCl₃/MeOH or MeOH). Mp 270-300° dec.


Sulfate:

Rhombic plates +10H₂O. Mp 289° (efferv.).Bartrop, J.A. et al., *J.C.S.*, 1954, 159 (*isol*)**Dendrobates Alkaloid 167A** A-354

2-Ethyldecahydroquinoline

R = CH₂CH₃, R' = HC₁₁H₂₁N 167.294Tentative struct. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula tentative; m/e 167(1), 166(1), 138(100). Cannot be hydrogenated.Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)Spande, T.F. et al., *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)**Dendrobates Alkaloid 181A** A-355

2-Ethyldecahydro-5-methylquinoline

As *Dendrobates* Alkaloid 167A, A-354 withR = CH₂CH₃, R' = CH₃C₁₂H₂₃N 181.32Tentative struct. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). Mol. formula tentative; m/e 181(2), 180(1), 152(100). Cannot be hydrogenated.Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)Spande, T.F. et al., *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)**Dendrobates Alkaloid 209A** A-356C₁₃H₂₅NO 211.347Struct. unknown. Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mol. formula tentative; m/e 209 (5), 168 (100). H₂-deriv, m/e 211, 168.Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)**Dendrobates Alkaloid 219A** A-357Decahydro-2,5-bis(2-propenyl)quinoline.
2,5-DiallyldecahydroquinolineR = CH₂CH=CH₂C₁₅H₂₅N 219.369A major constit. in skin extracts of *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates granuliferus*, *Dendrobates histrionicus*, *Dendrobates parvulus* and *Dendrobates truncatus* (Dendrobatidae). [α]_D²⁴ +9.7 (c, 2.0 in MeOH).Tetrahydro: Decahydro-2,5-dipropylquinoline. **Dendrobates Alkaloid 223A**†. *Pumiliotoxin C_{II}*

[63983-61-9]

[112791-03-4, 112790-00-8, 112789-97-6, 67920-43-8, 112789-98-7]


C₁₅H₂₉N 223.401Minor or trace alkaloids in skin extracts of *Dendrobates auratus*, *Dendrobates histrionicus* (various populations), *Dendrobates lehmanni*, *Dendrobates minutus*, *Dendrobates occultator*, *Dendrobates truncatus* and *Dendrobates* sp. of Colombia (Dendrobatidae).5-Epimer: **Dendrobates Alkaloid 219A'**C₁₅H₂₅N 219.369Present in skin extracts from one population of *Dendrobates histrionicus* and *Dendrobates auratus* (Dendrobatidae). [α]_D +5.8 (c, 0.31 in CHCl₃).

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)
 Tokuyama, T. *et al.*, *Tetrahedron*, 1986, **42**, 3453; 1991, **47**, 5401 (*isol. pmr, cmr, ms. cryst. struct. abs config.*)
 McCloskey, P.J. *et al.*, *J.O.C.*, 1988, **53**, 1380 (*tetrahydro, synth.*)
 Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur.*)
 Toyooka, N. *et al.*, *J.O.C.*, 2002, **67**, 6078-6081 (*synth.*)

Dendrobates Alkaloid 231A A-358

5-Ethyldecahydro-2-(2-penten-4-ynyl)quinoline

As *Dendrobates* Alkaloid 167A, A-354 with



C₁₆H₂₅N 231.38


Tentative struct. Minor alkaloid from skin extracts of *Dendrobates histrionicus* and *Dendrobates minutus* (Dendrobatidae). Mol. formula tentative; m/e 231(2), 230(1), 166(100). Forms H₆-deriv., m/e 237, 166. Does not appear to form an *N*-Ac deriv.

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)
 Spande, T.F. *et al.*, *Phytochemistry*, 1999, **62**, 5-21 (*pmr, cmr, occur.*)

Dendrobates Alkaloid 237A A-359

Octahydro-8-methyl-6-(2-methylpentylidene)-8-indolizolinol, 9CI. 8-Hydroxy-8-methyl-6-(2-methylpentylidene)indolizidine

[73376-36-0]



C₁₅H₂₇NO 237.384

Proposed struct. A major alkaloid from skin extracts of *Dendrobates abditus*; trace constit. in *Dendrobates histrionicus* (Dendrobatidae).

7-Hydroxy: **Dendrobates Alkaloid 253A** [73376-37-1]

C₁₅H₂₇NO₂ 253.384

A major alkaloid of *Dendrobates abditus*; minor constit. in *Dendrobates lehmanni* and *Dendrobates auratus* (Dendrobatidae). Stereochem. undefined.

7-Hydroxy, didehydro (?): **Dendrobates Alkaloid 251E**

C₁₅H₂₅NO₂ 251.368

Minor alkaloid from skin extracts of *Dendrobates minutus* (Dendrobatidae). Mol. formula and structure tentative;

m/e 251(3), 250(1), 234(2), 168(30), 84(18), 70(100).

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *J.A.C.S.*, 1980, **102**, 830 (*struct.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)
 Thanh, G.V. *et al.*, *Tet. Lett.*, 1999, **40**, 3713-3716 (*synth.*)

Dendrobates Alkaloid 239A A-360

C₁₅H₂₉NO 239.4

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates occultator* (Dendrobatidae). Mol. formula tentative; m/e 239 (2), 238 (3), 182 (100). Cannot be hydrogenated.

- Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 241A A-361

Dendrobates Alkaloid 241

Struct. unknown


Trace alkaloid from skin extracts of *Dendrobates occultator* (Dendrobatidae). Mol. formula not known; m/e 241 (2), 240 (3), 166 (100), 126 (48). Does not separate from major alkaloids 239 A, B, C, D.

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 243A A-362

Decahydro-5-(2-penten-4-ynyl)-2-(2-propenyl)quinoline, 2-Allyl-5-(2-penten-4-ynyl)decahydroquinoline

As *Dendrobates* Alkaloid 219A, A-357 with



C₁₇H₂₅N 243.391

A major alkaloid in *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates granuliferus*, *Dendrobates histrionicus*; minor or trace constit. in *Dendrobates pictus*, *Dendrobates tinctorius*, *Dendrobates trivittatus* and *Dendrobates truncatus* (Dendrobatidae). $[\alpha]_D^{16}$ -15.2 (c, 1.37 in MeOH). $[\alpha]_D^{16}$ -30.7 (c, 1.37 in CHCl₃).

Stereoisomer: *Dendrobates Alkaloid 243A'*

C₁₇H₂₅N 243.391

Present in some skin extracts of *Dendrobates* spp. (Dendrobatidae). $[\alpha]_D^{16}$ -0.96 (c, 0.73 in MeOH). $[\alpha]_D^{16}$ -18.6 (c, 0.73 in CHCl₃). Of unknown config.

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)
 Tokuyama, T. *et al.*, *Tetrahedron*, 1986, **42**, 3453 (*isol. pmr, cmr, struct.*)
 Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur.*)

Dendrobates Alkaloid 251A A-363

C₁₇H₃₃N 251.454

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates histrionicus*

(Dendrobatidae). Mol. formula tentative; m/e 251 (2), 208 (6), 152 (100). Possibly in Pumiliotoxin-C class with R = C₇H₁₅ (see *Dendrobates* Alkaloid 223C, A-438 and *Dendrobates* Alkaloid 167A, A-354).

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 257A A-364

C₁₈H₂₅N 255.402

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates trivittatus* (Dendrobatidae). Mol. formula tentative; m/e 257 (1), 256 (2), 216 (100). Cannot be acetylated, but consumes 4 molar equivs. of H₂. Possibly a perhydro-pyrroloquinoline of the gephyrotoxin class.

- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 265A A-365

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates abditus*, *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates histrionicus*, *Dendrobates lehmanni*, *Dendrobates occultator*, *Dendrobates parvulus* and *Dendrobates tinctorius* (Dendrobatidae). Mol. formula not known; m/e 265 (50), 264 (100), 222 (58), 180 (72). Very atypical mass spectrum for a dendrobatid alkaloid. Possibly a degradation artifact.


- Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 269A A-366

Decahydro-2-(3,4-pentadienyl)-5-(2-penten-4-ynyl)quinoline, 9CI. *Dendrobates histrionicus* Alkaloid III

[63983-62-0]

As *Dendrobates* Alkaloid 167A, A-354 with




C₁₉H₂₇N 269.429

Tentative struct. Minor or trace alkaloid in skin extracts of *Dendrobates granuliferus*, *Dendrobates histrionicus* (several populations), *Dendrobates occultator*, *Dendrobates trivittatus* and *Dendrobates truncatus* (Dendrobatidae), also in skin of the Madagascan frog *Mantella madagascariensis* (Ranidae, subfamily Mantellinae). m/e 269(4), 268(12), 204(100). H₁₀-deriv., m/e 279, 208.

- Daly, J.W. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1128 (*isol. ms.*)
 Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163; 1984, **22**, 905 (*isol. ms. rev.*)
 Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)
 Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur.*)

Dendrobates Alkaloid 275A A-367

Octahydro-5-methyl-10-(8-nonyl)-1H-pyrrolo[1,2-a]azepine, 9CI. 5-Methyl-10-(8-nonyl)lehmizidine



C₁₉H₃₃N 275.476

Major alkaloid from the skin extracts of the frog *Dendrobates lehmanni*.

8,9-Dihydro: Octahydro-5-methyl-10-(8-nonyl)-1H-pyrrolo[1,2-a]azepine.

Dendrobates Alkaloid 277A. 5-Methyl-10-(8-nonyl)lehmizidine

C₁₉H₃₅N 277.492

Alkaloid from *Dendrobates lehmanni*.

Oxo: Octahydro-5-methyl-10-(oxo-8-nonyl)-1H-pyrrolo[1,2-a]azepine. **Dendrobates Alkaloid 289A**

C₁₉H₃₁NO 289.46

Alkaloid from *Dendrobates lehmanni*.

Posn. of oxo group not known.

6,7'-Didehydro, 8,9-dihydro: Octahydro-5-methyl-10-(6,8-nonadienyl)-1H-pyrrolo[1,2-a]azepine. **Dendrobates Alkaloid 275G.** 5-Methyl-10-(6,8-nonadienyl)lehmizidine

C₁₉H₃₃N 275.476

Alkaloid from *Dendrobates lehmanni*.

Tentative struct. assigned.

3,5-Diepimer: **Dendrobates Alkaloid 275A'**

C₁₉H₃₃N 275.476

Alkaloid from *Dendrobates lehmanni*.

[120037-67-4]

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 421-427

Dendrobates Alkaloid 281A A-368

C₁₇H₃₁NO₂ 281.437

Struct. unknown. A major alkaloid from skin extracts of *Dendrobates minutus*; trace constit. in *Dendrobates abditus*.

Mol. formula tentative; m/e 281(4), 280(2), 264(2), 194(12), 166(72), 70(100). H₂-deriv., m/e 283(1), 282(2), 266(4), 208(40), 138(10), 110(10), 84(100), 70(85).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol. ms. rev.*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 297A A-369

C₁₇H₃₁NO₃ 297.437

Struct. unknown. A major alkaloid from skin extracts of *Dendrobates minutus* (Dendrobatidae). m/e 297 (3), 296 (4), 280 (9), 236 (2), 210 (3), 194 (4), 193 (3), 182 (21), 114 (27), 112 (16), 70 (100). Lowest molecular-weight member of the Pumiliotoxin-A class to contain 3 oxygens. H₂-deriv., m/e 299 (4), 282 (12), 256 (6), 224 (4), 110 (50), 84 (100), 70 (75).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms. rev.*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Dendrobates Alkaloid 309A A-370

Dihydropumiliotoxin A

C₁₉H₃₅NO₂ 309.491

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates minutus* and *Dendrobates viridis* (Dendrobatidae). Mol. formula tentative; m/e 309 (9), 308 (3), 292 (2), 280 (4), 206 (4), 194 (15), 176 (5), 166(100), 110 (10), 84 (20), 70 (51). Forms H₂-deriv.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol. ms. rev.*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms. rev.*)

Erythrophleum Alkaloid A A-371

C₂₅H₃₇NO₆ 447.57

Struct. unknown. Dimethylaminoethanol esterified with an α,β-unsaturated diterpene acid carrying a methoxycarbonyl group and 3 Me groups. UV shows no evidence for a 6,7-dione as observed in other *E.* alkaloids. Alkaloid from the bark of *Erythrophleum africanum* (Fabaceae). Cryst. (Et₂O/hexane). Mp 105°. Poss. identical with an alkaloid from *E. guineense* with the same MF. No further data to 2007.

Friedrich-Fichtl, J. *et al.*, *Chem. Ber.*, 1971, **104**, 3535-3548 (*Erythrophleum guineense alkaloid*)

Jansson, S. *et al.*, *Acta Pharm. Suec.*, 1976, **13**, 51-54; *CA*, **84**, 147726j (*isol. uv. ir. pmr. ms*)

Euonymus Alkaloid A A-372

C₂₉H₃₇NO₁₃ 607.61

Struct. unknown. Alkaloid from the seeds of *Euonymus europaea*. Mp 258-260°. [α]_D +21 (CHCl₃).

De-Ac: **Euonymus Alkaloid B**

C₂₇H₃₅NO₁₂ 565.573

Alkaloid from the seeds of *Euonymus europaea*. Mp 288-290°. [α]_D +14 (CHCl₃).

Ac: **Euonymus Alkaloid C**

C₃₁H₃₉NO₁₄ 649.647


Alkaloid from the seeds of *Euonymus europaea*. Mp 164-168°. [α]_D +8 (CHCl₃).

Doebel, K. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 592-597 (*isol.*)

Fritillaria Alkaloid A A-373

Struct. unknown. Alkaloid from *Fritillaria meleagris* (Liliaceae). Mp 156-158°. Cooccurs with Imperialine in C-298.

Bauer, S. *et al.*, *Chem. Zvesti.*, 1958, **12**, 584

Mantella Alkaloid 384A A-374


C₂₆H₄₄N₂ 384.647

Tentative struct. shown. Obt. as mixt. with *Mantella* Alkaloid 384B. Major constit. of skin extracts of the Madagascan frog *Mantella betsileo*. Minor constit. in *Mantella* sp. cf. *madagascariensis* and *Mantella laevigata*. No CAS no. found to 15CI.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol. ir. ms*)

Securidaca longipedunculata Alkaloid A A-375

[146442-66-2]



C₂₀H₁₆N₂O₄ 348.357

Alkaloid from roots of *Securidaca longipedunculata* (Polygalaceae).

Costa, C. *et al.*, *J. Het. Chem.*, 1992, **29**, 1641-1647

Stephania glabra Alkaloid A A-376

Stephania glabra Base A

C₁₉H₂₁NO₄ 327.379

Struct. unknown, conts. 2 OMe groups. Authors suggest the struct. to be *N*-methylhydroxystepharine. Isol. from herbage of *Stephania glabra*. Mp 205-206°. [α]_D²¹ -70. λ_{max} 242; 282 (no solvent reported).

Shchelchkova, I.I. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 271-275; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 210-212 (*isol.*)

Stephania glabra Alkaloid A A-377

C₁₉H₁₈NO₂ 292.357

Struct. unknown. The MF assigned is impossible. Isol. from tubers of *Stephania glabra*. Cryst. (Et₂O). Mp 79-81°. [α]_D -88 (c, 1 in EtOH). Prob. an aporphine alkaloid. Contains an NMe group. No further reports, although later investigations of the species have been reported.

Hydrochloride: Mp 252-253°.

N-Ac:

Cryst. (Me₂CO). Mp 146°.

Kin, F.K. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 392-394; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 308-309 (*isol.*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2785-2789 (*Stephania glabra* constits)

Stephania japonica Alkaloid A A-378

Stephania Base A

C₂₁H₂₆ClNO₇ 439.892

Struct. unknown. Alkaloid from the stems and rhizomes of *Stephania japonica* var. *australis* (Menispermaceae). Prisms (MeOH). Mp 248°. Ir shows OH

and γ -lactam groups, pmr shows 4 OMe groups, 1 NMe group, and 2 aromatic protons. Ms m/e 257.1068, indicating a hasubanan alkaloid.

Matsui, M. *et al.*, *Phytochemistry*, 1979, **18**, 1087 (isol, ir, pmr, ms)

Taxus baccata Alkaloid A A-379

C₃₃H₄₃N₂O₁₄ 691.708

Struct. unknown. The MF given is impossible. Isol. from leaves of *Taxus baccata*. Cryst. (petrol). Mp 112-113°. [α]_D²⁵ +10.7 (c, 0.835 in CHCl₃). Co-occurs with Taxol, T-52.

Mirzoev, Kh.M. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 777-778; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 797

Thermopsis lanceolata Alkaloid A A-380

[78040-76-3]

Struct. unknown. Constit. of *Thermopsis lanceolata*.

Chao, K.-S. *et al.*, *CA*, 1981, **95**, 35461q (isol, props)

Alangium Alkaloid AL64 A-381

C₂₉H₃₇N₃O₃ 475.63

Struct. unknown. Poss. identical with Alangimarckine in T-675. Alkaloid from the root bark of *Alangium lamareckii* (Alangiaceae). Glistening needles (CH₂Cl₂ and/or MeOH). Mp 272°. [α]_D²⁵ -64 (CHCl₃). λ_{\max} 226 (log ϵ 4.28); 283 (log ϵ 4.18) (EtOH).


Mono-Ac: Mp 184°.

Di-Ac: Mp 150°.

Pakrashi, S. *et al.*, *Indian J. Chem.*, 1964, **2**, 468 (isol, uv)

Alkaloid AM 3 A-382

N-Deacetyl-N,N-dimethyl- γ -lumicornigerine
[99499-80-6]



C₂₁H₂₃NO₅ 369.416

Alkaloid from the aerial parts of *Androcymbium melanthioides* var. *stricta* (Liliaceae). Needles. Mp 245-247°. [α]_D²³ -391 (c, 1.0 in CHCl₃).

Potěšilová, H. *et al.*, *Planta Med.*, 1985, **344** (isol, uv, ir, pmr, cmr, ms, cd, struct)

Alangium Alkaloid B1 A-383

C₂₀H₃₁NO₅ 365.469

Struct. unknown. Alkaloid from the bark of *Alangium lamareckii*. Mp 197-198° Mp 228-229° (as hydrochloride).

Basu, N.K. *et al.*, *J. Indian Chem. Soc.*, 1957, **34**, 629-639; *CA*, **52**, 7337g

Alangium Alkaloid B2 A-384

C₂₇H₄₃NO₆ 477.64

Struct. unknown. Alkaloid from the bark of *Alangium lamareckii*. Mp 119-120° Mp 188° (as hydrochloride).

Basu, N.K. *et al.*, *J. Indian Chem. Soc.*, 1957, **34**, 629-639; *CA*, **52**, 7337g

Alangium Alkaloid B3 A-385

Struct. unknown. Alkaloid from the bark of *Alangium lamareckii*. Mp 160-161° Mp 181° (as hydrochloride).

Basu, N.K. *et al.*, *J. Indian Chem. Soc.*, 1957, **34**, 629-639; *CA*, **52**, 7337g

Alangium Alkaloid B4 A-386

C₁₉H₂₇NO₇ 381.425

Struct. unknown. Alkaloid from the bark of *Alangium lamareckii*. Mp 149-151° Mp 199° (as hydrochloride).

Basu, N.K. *et al.*, *J. Indian Chem. Soc.*, 1957, **34**, 629-639; *CA*, **52**, 7337g

Alangium Alkaloid B5 A-387

C₂₁H₃₁NO₈ 425.478

Struct. unknown. Alkaloid from the bark of *Alangium lamareckii*. Mp 177-179° Mp 208° (as hydrochloride).

Basu, N.K. *et al.*, *J. Indian Chem. Soc.*, 1957, **34**, 629-639; *CA*, **52**, 7337g

Asteracantha Alkaloid B1 A-388

Struct. unknown. Alkaloid from *Asteracantha longifolia* (preferred genus name *Hygrophila*). Mp 294°.

Basu, N.K. *et al.*, *Indian J. Pharm.*, 1952, **14**, 212-213; *CA*, **47**, 4044g

Asteracantha Alkaloid B2 A-389

C₃H₁₃NO₃ 135.163

Struct. unknown. Alkaloid from *Asteracantha longifolia* (preferred genus name *Hygrophila*). Mp 194°.

Basu, N.K. *et al.*, *Indian J. Pharm.*, 1952, **14**, 212-213; *CA*, **47**, 4044g

Buxus Alkaloid B A-390

C₂₄H₄₂N₂O 374.609

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 205-207°.

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 2209-2226 (isol)

Buxus Alkaloid B1 A-391

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO/MeOH). Mp 280° dec. [α]_D²² +11.8 (c, 0.55 in MeOH). Contains MeNH group. λ_{\max} 243 (log ϵ 4.19) (MeOH).

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, uv, ms)

Buxus Alkaloid B4 A-392

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (CHCl₃). Mp 150-152°.

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, ms)

Buxus Alkaloid B5 A-393

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (CHCl₃). Mp 179-182°. [α]_D²² +9.6 (c, 0.5 in CHCl₃).

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, ms)

Buxus Alkaloid B6 A-394

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (MeOH). Mp 214-216°. [α]_D²² +6.6 (c, 0.15 in MeOH). Contains OH group and C₆-C₇ double bond.

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, ir, ms)

Buxus Alkaloid B7 A-395

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO). Mp 245-247°.

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, ms)

Buxus Alkaloid B8 A-396

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO). Mp 265°. [α]_D²² +34.6 (c, 0.54 in CHCl₃). Mass spec. indicated a MeNH group at C-20 and a dimethyl group at C-3.

Huong, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 1425-1432 (isol, ms)

Buxus Alkaloid B387 A-397

C₂₅H₄₁NO₂ 387.604

Steroidal alkaloid. Struct. unknown. Alkaloid from *Buxus microphylla* var. *sinica* (Buxaceae). Mp 188-189°. [α]_D²² +6.7 (CHCl₃). ORD spectrum resembles that of 4,4a-dihydrobuxpiene.

Bauerová, O. *et al.*, *Pharmazie*, 1973, **28**, 212-214; *CA*, **78**, 156665s (isol, ir, uv, ms, ord)

Buxus Alkaloid B398 A-398

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *argentea* (Buxaceae). Cryst. (C₆H₆/MeOH). Mp 197-199°.

Kuchkova, K.I. *et al.*, *Chem. Zvesti*, 1976, **30**, 174-178 (isol)

Buxus Alkaloid B426 A-399

Struct. unknown. Alkaloid from *Buxus sempervirens* var. *argentea* (Buxaceae). Cryst. (MeOH/CH₂Cl₂). Mp 208-210°.

Kuchkova, K.I. *et al.*, *Chem. Zvesti*, 1976, **30**, 174-178; *CA*, **87**, 2362q (isol)

C-Alkaloid B A-400

C₂₀H₂₃N₂O[⊕] 307.414

Minimum formula. Struct. unknown. Alkaloid from calabash curare and *Strychnos mitscherlichii* (Loganiaceae).

Mp 270° (as picrate). Red-violet col. with Ce(SO₄)₂, becoming brown on standing.

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 1864-1879; 1953, **36**, 102-121

Dendrobates Alkaloid 181B A-401

Decahydro-2-propylquinoline

As *Dendrobates* Alkaloid 167A, A-354 with

R = C₃H₇, R' = H

C₁₂H₂₃N 181.32

Tentative struct. Trace alkaloid from skin extracts of *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates fulguritus* and *Dendrobates trivittatus* (Dendrobatidae). Mol. formula tentative; m/e 181(2), 180(2), 138(100). Cannot be hydrogenated.


Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 205B A-402

Alkaloid 205B



Absolute Configuration

C₁₄H₂₃N 205.342

Proposed gross struct. confirmed and stereochem. revised in 2003. Trace alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates punilio* (Dendrobatidae). [α]_D -8.5 (c, 0.59 in CHCl₃).

Tokuyama, T. *et al.*, *Tetrahedron*, 1987, **43**, 643 (*isol, pmr, cmr, ms, struct*)

Toyooka, N. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3808-3810 (*synth, abs config*)

Smith, A.B. *et al.*, *J.O.C.*, 2006, **71**, 2547-2557 (*synth*)

Dendrobates Alkaloid 219B A-403

Decahydro-5-methyl-2-(2,4-pentadienyl)-quinoline

As *Dendrobates* Alkaloid 167A, A-354 with

R = -CH₂CH=CHCH=CH₂,

R' = CH₃

C₁₅H₂₅N 219.369

Tentative struct. Minor alkaloid from skin extracts of *Dendrobates histrionicus* and *Dendrobates* sp. (Colombia), trace constit. in *Dendrobates azureus* (Dendrobatidae). Mol. formula tentative, m/e 219(1), 218(2), 152(100). H₄-deriv., m/e 223, 152.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 223B A-404

As *Dendrobates* Alkaloid 167A, A-354 with

R = C₄H₉, R' = CH₂CH₃

C₁₅H₂₉N 223.401

Tentative struct. Minor or trace alkaloid in skin extracts of *Dendrobates auratus*, *Dendrobates histrionicus*, *Dendrobates minutus*, *Dendrobates truncatus* and *Dendrobates* sp. of Colombia (Dendrobatidae). m/e 223(1), 222(2), 166(100). Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 225B A-405

[103881-40-9]

C₁₅H₃₁N 225.417

Struct. unknown. May be a 2,6-dipentylpiperidine. Trace constit. of *Dendrobates histrionicus* and a minor constit. in an undescribed *Dendrobates* sp. from Panama (Dendrobatidae).

Daly, J.W. *et al.*, *J. Nat. Prod.*, 1986, **49**, 265-280

Dendrobates Alkaloid 231B A-406

As *Dendrobates* Alkaloid 167A, A-354 with

R = C₆H₇, R' = CH₃

C₁₆H₂₅N 231.38

Tentative struct. Minor or trace alkaloid from skin extracts of *Dendrobates azureus*, *Dendrobates histrionicus*, *Dendrobates lehmanni*, *Dendrobates occaltator*, *Dendrobates tinctorius* and *Dendrobates* sp. (Colombia) (Dendrobatidae). Mol. formula tentative; m/e 231(2), 230(1), 152(100). Forms H₆-deriv. m/e 237, 152.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol. ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 237B A-407

C₁₄H₂₃NO₂ 237.341

Struct. unknown. Trace constit. in skin extracts of *Dendrobates abditus* and *Dendrobates fulguritus* (Dendrobatidae). Mol. formula tentative; m/e 237 (1), 236 (2), 182 (60), 114 (30), 112 (25), 70 (100).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 239B A-408

C₁₅H₂₉NO 239.4

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates bombetes*; trace constit. in *Dendrobates occaltator* (Dendrobatidae). Mol. formula tentative;

m/e 239 (2), 238 (3), 180 (100). Cannot be hydrogenated.

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 253B A-409

C₁₆H₃₃NO 255.443

Struct. unknown. Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mol. formula tentative; m/e 253 (1), 138 (100). Cannot be hydrogenated.

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 257B A-410

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates lehmanni* (Dendrobatidae). Mol. formula not known; m/e 257 (60), 256 (100), 152 (20). Very atypical mass spectrum for a dendrobatid alkaloid. Possibly a degradation artifact.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 267B A-411

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* and *Dendrobates minutus* (Dendrobatidae). Mol. formula not known; m/e 267 (7), 266 (4), 250 (1), 170 (100), 152 (4), 112 (13). The base peak is atypical for a dendrobatid alkaloid.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 269B A-412

As *Dendrobates* Alkaloid 167A, A-354 with

R = CH₂CH=CHC≡CH,

R' = CH₂CH=CHCH=CH₂

C₁₉H₂₇N 269.429

Tentative struct. Minor or trace alkaloid in skin extracts of *Dendrobates granuliferus*, *Dendrobates histrionicus* (several populations), *Dendrobates occaltator*, *Dendrobates trivittatus* and *Dendrobates truncatus* (Dendrobatidae). m/e 269(4), 286(12), 202(100). H₁₀-deriv., m/e 279, 208.

Daly, J.W. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1128 (*isol, ms*)

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 281B A-413

C₁₈H₃₅NO 281.481

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates granuliferus* (Dendrobatidae). Mol. formula tentative; m/e 281 (4), 264 (12), 208 (25), 206 (20), 150 (65), 98 (5), 96 (20), 70 (100). Atypical spectrum for a dendrobatid alkaloid. Probably in

Pumiliotoxin-A class. Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol*, *ms*, *rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Dendrobates Alkaloid 283B A-414

C₁₇H₃₃NO₂ 283.453

Struct. unknown. A major alkaloid in skin extracts of *Dendrobates histrionicus* and *Dendrobates occultator* (Dendrobatidae). m/e 283 (<1), 282 (1), 254 (2), 212 (C₁₂H₂₂NO₂, 40), 152 (C₁₀H₁₈N, 23), 140 (C₉H₁₈N, 100). Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol*, *ms*, *rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Dendrobates Alkaloid 297B A-415

C₁₈H₃₅NO₂ 297.48

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). m/e 297 (10), 166 (92), 70 (100). Forms H₄-deriv.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol*, *ms*, *rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Dendrobates Alkaloid 307B A-416

Isopumiliotoxin A
[67255-98-5]

C₁₉H₃₃NO₂ 307.475

Struct. unknown. Isomer of Pumiliotoxin A, P-791 without a large fragment ion for loss of OH. Minor or trace alkaloid in skin extracts of *Dendrobates auratus*, *Dendrobates granuliferus*, *Dendrobates lehmanni*, *Dendrobates minutus* and *Dendrobates pumilio* (Dendrobatidae). m/e 307 (12), 306 (4), 290 (2), 194 (24), 193(45), 166 (100), 70 (56). H₂-deriv, m/e 309(1), 280 (3), 208 (10), 138 (25), 110 (60), 84(100), 70 (45). H₄-deriv, m/e 311 (1), 110(25), 84 (100), 70 (30).

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (*isol*, *ms*, *rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Dendrobates Alkaloid 309B A-417

C₂₀H₃₉NO 309.534

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates auratus* and *Dendrobates granuliferus* (Dendrobatidae). Mol. formula tentative; m/e 309 (1), 152 (100). Possibly member of Hydroxy pumiliotoxin-C class with OH group in the side chain.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol*, *ms*, *rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Gentiana macrophylla Alkaloid B A-418

C₉H₉NO₂ 163.176


Struct. unknown. Monoterpene alkaloid. Alkaloid from the roots of *Gentiana*

macrophylla (Gentianaceae). Cryst. (C₆H₆/petrol). Mp 128-130°.

Fu, F.-Y. *et al.*, *Yaouxue Xuebao*, 1958, **6**, 198-203; *CA*, **53**, 8310b (*isol*, *uv*)

Mantella Alkaloid 253B A-419

(Octahydro-8-methyl-5-indolizinyloxy)-sec-heptanol, 9CI. Indolizidine 253B
[151871-22-6]



C₁₆H₃₁NO 253.427

Provisional struct. Secondary alcohol. Minor alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (*isol*, *ir*, *ms*)

Mantella Alkaloid 271B A-420

Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascar frog *Mantella betsileo*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*, *ir*, *ms*)

Mantella Alkaloid 293B A-421

Struct. and MF unknown. Minor alkaloid from skin extracts of the Madagascar frog *Mantella* sp. cf. *madagascariensis*. Minor or trace constit. in *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*, *ir*, *ms*)

Mantella Alkaloid 295B A-422

Struct. and MF unknown. Tentatively classified as an indolizidine. Forms a di-O-Ac deriv. Minor alkaloid from skin extracts of the Madagascar frog *Mantella crocea*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*, *ir*, *ms*)

Pseudophryne Alkaloid 252B A-423

C₁₄H₂₄N₂O₂ 252.356

Struct. unknown. Possibly analogous to Spiropyrolizidine 252, S-437 but with the hydroxy group in another part of the struct. Minor alkaloid from skin extracts of one population of the Australian frog *Pseudophryne coriacea* (trace constit. in a different population of the same frog).

Daly, J.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 407-421 (*isol*, *ms*)

Skytanthus acutus Alkaloid B A-424

C₁₁H₂₁N 167.294

Struct. unknown. Monoterpene alkaloid. Alkaloid from *Skytanthus acutus* (Apocynaceae). Bp₅ 190°.

Appel, H.H. *et al.*, *Scientia (Valparaiso)*, 1961, **28**, 5-9; *CA*, **57**, 2332g

Sophora griffithii Alkaloid B A-425

Struct. unknown. Constit. of *Sophora griffithii*.

Primukhamedov, I. *et al.*, *Rastit. Resur.*, 1969, **5**, 572-575; *CA*, **72**, 118549

Strychnos camptoneura Alkaloid B A-426

[58572-16-0]

Struct. unknown. Bisindole alkaloid. Alkaloid from *Strychnos camptoneura* (Loganiaceae). Mol. formula not reported (MW 622).

Verpoorte, R. *et al.*, *Acta Pharm. Suec.*, 1975, **12**, 455-460 (*isol*, *uv*, *ir*, *ms*)

Thermopsis lanceolata Alkaloid B A-427

[78040-77-4]

Struct. unknown. Constit. of *Thermopsis lanceolata*.

▶AZ7675000

Chao, K.-S. *et al.*, *CA*, 1981, **95**, 35461q (*isol*, *props*)

Alkaloid 241B A-428

[97380-36-4]

C₁₆H₃₅N 241.459

Struct. unknown. Major alkaloid from skin extracts of the Madagascar frog *Mantella madagascariensis* (family Ranidae, subfamily Mantellinae). m/e 241 (15), 135 (45), 58 (100). Cannot be hydrogenated. No exchangeable hydrogen. Apparently lacks rings. Has not been detected in dendrobatid frogs.

Daly, J.W. *et al.*, *Toxicol.*, 1984, **22**, 905-919 (*isol*, *ms*)

C-Alkaloid BL A-429

C₄₀H₅₀N₄O₂[⊕] 618.861

An oxidn. prod. of C-Alkaloid D, A-476. Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae).

▶Toxic (0.02 x Toxiciferine).

Dichloride:

Cryst. + 2H₂O. Dec. without melting, turning yellow, red then black. Deep-blue col. with Ce(SO₄)₂, becoming blue-green on standing.

Dipicrate: Mp 172-178°.

Schmidt, I. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 1218 (*isol*, *ir*, *uv*)

Gorman, A.A. *et al.*, *Alkaloids (London)*, 1971, **1**, 200-338

Buxus Alkaloid BX6 A-430

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 207-212°.

Döpke, W. *et al.*, *Pharmazie*, 1969, **24**, 649; *CA*, **72**, 43929p (*isol*)

Buxus Alkaloid BX10 A-431

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 221-224°.

Döpke, W. *et al.*, *Pharmazie*, 1969, **24**, 649; *CA*, **72**, 43929p (*isol*)

Chamaecytisus Alkaloid C₁ A-432C₁₅H₂₄N₂O₂ 264.367

Struct. unknown. Prob. a hydroxylupanine. Alkaloid from *Chamaecytisus friwaldskyanus*, *Chamaecytisus albus* and *Chamaecytisus calcareus* (Fabaceae). Mp 207-209°. $[\alpha]_D^{25} +59.2$ (c, 0.38 in EtOH). Daily, A. et al., *Planta Med.*, 1977, **32**, 380-383

Chamaecytisus Alkaloid C₂ A-433C₁₂H₂₀N₂O 208.303

Struct. unknown. Alkaloid from *Chamaecytisus austriacus*, *Chamaecytisus ciliatus* and *Chamaecytisus polytrichus* (Fabaceae). Mp 168-170°. $[\alpha]_D^{25} +38$ (c, 0.4 in EtOH).

Daily, A. et al., *Planta Med.*, 1977, **32**, 380-383,

Rauwolfia obscura Alkaloid A-434C₂C₃₀H₃₄N₄O 466.625

Bisindole alkaloid. Struct. unknown. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Mp 142-144°.

Timmins, P. et al., *Planta Med.*, 1975, **27**, 105-111 (isol, uv, ir, pmr, ms)

Buxus Alkaloid C A-435C₂₄H₄₂N₂O 374.609

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 212-214°. $[\alpha]_D +60$ (CHCl₃). Props. similar to Cyclobuxamine H in C-846.


Schlittler, E. et al., *Helv. Chim. Acta*, 1949, **32**, 2209-2226 (isol)

C-Alkaloid C A-436

Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Characterised as picrate, dec. >270° without melting. Red-violet col. with Ce(SO₄)₂, becoming brown on standing. Schmid, H. et al., *Helv. Chim. Acta*, 1952, **35**, 1864-1879; 1953, **36**, 102-121

Crotalaria goreensis A-437**Alkaloid C**

Crotalaria goreensis Base C



Possible structure

C₈H₁₃NO 139.197

Alkaloid from *Crotalaria goreensis* whole plant (Fabaceae). Bp₃ 112-114°. $[\alpha]_D +9.3$ (EtOH), pK_a 8.8. Conts. one C-Me group. Gives a dihydro deriv.

Culvenor, C.C.J. et al., *Aust. J. Chem.*, 1961, **14**, 284

Dendrobates Alkaloid 223C A-438

[67217-78-1]

As *Dendrobates* Alkaloid 167A, A-354 with

R = C₅H₁₁, R' = CH₃C₁₅H₂₉N 223.401

Tentative struct. Minor or trace alkaloid in skin extracts of *Dendrobates auratus*, *Dendrobates histrionicus* (various populations), *Dendrobates truncatus* and *Dendrobates* sp. of Colombia (Dendrobatidae). m/e 223(1), 222(2), 152(100). Cannot be hydrogenated.

Hydroxy (?) (1): Dendrobates Alkaloid 239EC₁₅H₂₉NO 239.4

Trace alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Mol. formula tentative. m/e 239(2), 238(3), 210(40), 152(100). Poss. a side-chain substd. deriv. of 223C. Cannot be hydrogenated.

Hydroxy (?) (2): Dendrobates Alkaloid 239FC₁₅H₂₉NO 239.4

Trace alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Mol. formula tentative. m/e 239(1), 168(100). Cannot be hydrogenated, forms O-Ac deriv. Poss. ring-subst. deriv. of 223C.

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Spande, T.F. et al., *J. Nat. Prod.*, 1999, **62**, 5-21 (pmr, cmr, occur)

Dendrobates Alkaloid 225C A-439C₁₅H₃₁N 225.417

Struct. unknown. Tentatively proposed to be a 2-*n*-butyl-5-*n*-heptylpyrrolidine. Trace constit. in *Dendrobates histrionicus* (Dendrobatidae).

Daly, J.W. et al., *J. Nat. Prod.*, 1986, **49**, 265-280

Dendrobates Alkaloid 237C A-440C₁₆H₃₁N 237.428

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Mol. formula tentative; m/e 237 (2), 236 (1), 180 (100). Probably higher homologue of 223B. Cannot be hydrogenated.

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 239C A-441C₁₅H₂₉NO 239.4

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates occulator* (Dendrobatidae). Mol. formula tentative; m/e 239 (2), 238 (3), 196 (100). Cannot be hydrogenated.

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 251C A-442C₁₆H₂₉NO 251.411

Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates minutus* (Dendrobatidae). Mol. formula tentative;

m/e 251 (2), 234 (4), 154 (100). H₂-deriv, m/e 253, 154.

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 283C A-443C₁₇H₃₃NO₂ 283.453

Struct. unknown. A major alkaloid from skin extracts of *Dendrobates histrionicus* and *Dendrobates occulator* (Dendrobatidae). m/e 283 (<1), 282 (1), 240 (5), 226 (C₁₃H₂₄NO₂, 28), 224 (C₁₅H₃₀N, 10), 166 (C₁₁H₂₀N, 60), 126 (C₈H₁₆N, 100). Cannot be hydrogenated.

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 307C A-444C₁₉H₃₃NO₂ 307.475

Struct. unknown. Minor alkaloid from skin extracts of *Dendrobates lehmanni* and *Dendrobates pumilio* (Dendrobatidae). m/e 307 (9), 290 (11), 182 (62), 70 (100). H₄-deriv, m/e 311 (3), 294 (3), 268 (5), 236(2), 110 (30), 84 (100), 70 (35).

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 309C A-445C₁₉H₃₅NO₂ 309.491

Struct. unknown. Minor alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mol. formula tentative; m/e 309 (3), 308 (2), 292 (1), 280 (4), 266 (4), 194 (15), 166 (100), 70(90). H₂-deriv, m/e 311 (3), 110 (25), 84(100), 70 (30).

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Erythrophleum Alkaloid C A-446

Struct. unknown. Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Cryst. (petrol). Mp 116°. Readily dec. Possibly an artifact.

Cronlund, A. et al., *Acta Pharm. Suec.*, 1975, **12**, 467-478 (isol, uv, ir, pmr, ms)

Gentiana macrophylla Alkaloid C A-447

Struct. unknown. Mol. formula not recorded. Monoterpene alkaloid. Alkaloid from the roots of *Gentiana macrophylla* (Gentianaceae). Cryst. (Me₂CO). Mp 206-208°.

Fu, F.-Y. et al., *Yaoxue Xuebao*, 1958, **6**, 198-203; *CA*, **53**, 8310b (isol, uv)

Leucojum Alkaloid C A-448

Leucojum Base C

C₁₆H₁₇NO₅ 303.314

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Leucojum* sp.

(Amaryllidaceae). Mp 185-186°. $[\alpha]_D^{25}$ -29 (CHCl₃).

Raffauf, R.F. *et al.*, *Handb. Alkaloids Alkaloid-Containing Plants*, Wiley, 1970, 284

Mantella Alkaloid 205C A-449

Struct. and MF unknown. Trace alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.


Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (isol, ms)

Mantella Alkaloid 211C A-450

Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascar frog *Mantella laevigata*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ir, ms)

Mantella Alkaloid 235C A-451



C₁₅H₂₅NO 235.369

Structure revised in 2005. Alkaloid from skin extracts of the Madagascar frogs *Mantella aurantiaca*, *Mantella madagascariensis* and *Mantella crocea*.

Ketone: Mantella Alkaloid 233F

C₁₅H₂₃NO 233.353

Trace alkaloid from skin extracts of *Mantella madagascariensis*, *Mantella aurantiaca* and *Mantella crocea*.

12 ξ -Hydroxy: Mantella Alkaloid 251G


C₁₅H₂₅NO₂ 251.368

Alkaloid from a *Mantella* sp. Tentative struct. assigned.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ir, ms)

Andriamaharavo, N.R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1743-1748 (synth, ir, pmr, ms)

Mantella Alkaloid 261C A-452



C₁₈H₃₁N 261.45

Major alkaloid from skin extracts of the frog *Mantella betsileo*.

Dihydro(?): Mantella Alkaloid 263G

C₁₈H₃₃N 263.465

Alkaloid from skin extracts of *Mantella betsileo*.

Didehydro(?): Mantella Alkaloid 259D

C₁₈H₂₉N 259.434

Alkaloid from the skin extracts of *Mantella betsileo*.

Hydroxy(?): Mantella Alkaloid 277G

C₁₈H₃₁NO 277.449

Alkaloid from the skin extracts of *Mantella betsileo*.

Hydroxy(?): Mantella Alkaloid 277H

C₁₈H₃₁NO 277.449

Alkaloid from the skin extracts of *Mantella betsileo*.

Kaneko, T. *et al.*, *Heterocycles*, 2003, **59**, 745-757 (isol, pmr, cmr, ms)

Toyooka, N. *et al.*, *Synlett*, 2005, 3109-3110 (synth)

Mantella Alkaloid 321C A-453

Allopumiliotoxin 321C

Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascar frog *Mantella betsileo*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (isol, ms)

Salicornia herbacea A-454

Alkaloid C

Struct. unknown. Alkaloid from *Salicornia herbacea* (Chenopodiaceae). Mp 236-238°.

Perchlorate: Mp 210-212°.

Picrate: Mp 117°.

Borkowski, B. *et al.*, *Pharmazie*, 1965, **20**, 390-393; *C.A.*, **63**, 15223c

Stephania glabra Alkaloid C A-455

C₂₁H₂₅NO₄ 355.433

Struct. unknown. Not reported in further investigations of this species. Isol. from tubers of *Stephania glabra*. Cryst. (Me₂CO). Mp 182-183°. $[\alpha]_D^{25}$ +259 (c, 1 in EtOH). Conts. 3 OMe and 1 NMe group. λ_{max} 267 ; 304 (no solvent reported).

Hydrochloride: Mp 230-232°.

Kin, F.K. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 392-394; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 308-309

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2785-2789 (*Stephania glabra* constits)

Voacanga thouarsii A-456

Alkaloid C

[50924-00-0]

C₄₂H₅₄N₄O₁₀ 774.909

Bisindole alkaloid. Struct. unknown. Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Cryst. (MeOH). Mp 290-295° dec. $[\alpha]_D^{25}$ -282. Intense blue col. with HNO₃. λ_{max} 223 (log ϵ 4.57); 263 (log ϵ 4.1); 299 (log ϵ 4.17); 328 (log ϵ 4.21) (EtOH).

Rolland, Y. *et al.*, *Phytochemistry*, 1973, **12**, 2039-2042 (isol, uv, ir, pmr)

Alkaloid C2 A-457


C₃₀H₃₄N₄O 466.625

Prob. of Usambarine type. Struct. unknown. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Cryst. Mp 142-144°.

Timmins, P. *et al.*, *Planta Med.*, 1975, **27**, 105-111 (isol, uv, ir, pmr, ms)

Alkaloid CB2 A-458

α -D-Glucopyranose 1,3,6-tri-3-pyridine-carboxylate, 9CI. 1,3,6-Tri-O-nicotinoyl- α -D-glucopyranose [76129-64-1]




C₂₄H₂₁N₃O₉ 495.445

Alkaloid from stems of *Cryptolepis buchanani* (Asclepiadaceae). Mp 131-133°.

Dutta, S.K. *et al.*, *Phytochemistry*, 1980, **19**, 1278 (isol, uv, ir, pmr, ms, struct)

Alkaloid CC 2 A-459

[35320-76-4]



Absolute Configuration

C₂₁H₂₇NO₅ 373.448

Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Cryst. (EtOAc/Et₂O). Mp 172-174° (168-170°). $[\alpha]_D^{25}$ +40 (c, 0.48 in CHCl₃).

Ac: Mp 135-137°.

5,6-Didehydro, 7-ketone: Alkaloid CC 20 [35320-75-3]

C₂₁H₂₃NO₅ 369.416

Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Cryst. (EtOAc/Et₂O). Mp 210-212°. $[\alpha]_D^{25}$ +324 (c, 0.26 in CHCl₃).

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413 (isol)

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540 (isol, uv, ir)

Battersby, A.R. *et al.*, *J.C.S.(C)*, 1971, 3514 (uv, ir, pmr, cryst struct)

Alkaloid CC 3 A-460

C₂₀H₂₅NO₅ 359.421

Struct. unknown. Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Fine needles (EtOAc/Et₂O). Mp 197-199°. $[\alpha]_D^{25}$ +155 (c, 0.76 in CHCl₃).

El-Hamidi, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 2111-2118 (isol, uv, ir)

Alkaloid CC 11 A-461

Mol. formula not reported. Struct. unknown. Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Mp 251-253°.

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413-3423 (isol)

Alkaloid CC 13 A-462

Mol. formula not reported. Struct. unknown. Alkaloid from the seeds of *Colchicum cornigerum* (Liliaceae). Cryst. (MeOH). Mp 115-117°.

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413-3423 (*isol*)

Alkaloid CC 14 A-463

Struct. unknown. Mol. formula not reported. Alkaloid from the stems of *Colchicum cornigerum* (Liliaceae). Mp 182-185°.

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413-3423 (*isol*)


Alkaloid CC 23 A-464

Struct. unknown. MF = C₂₁₋₂₂H₂₃₋₂₇NO₅₋₆. Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Mp 240-243°. [α]_D²² -22 (c, 0.2 in CHCl₃). [α]_D²² -4 (c, 0.3 in MeOH).

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540-3552 (*isol, uv, ir*)

Alkaloid CC 3b A-465

[35320-77-5]



C₂₁H₂₉NO₅ 375.464

Probable struct. May be the O⁴-de-Me, O²-Me isomer. Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Cryst. (EtOAc). Mp 210-212°. [α]_D²² +25 (c, 0.76 in CHCl₃).

Di-Ac: Mp 205-206°.

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540 (*isol, uv, ir*)

Battersby, A.R. *et al.*, *J.C.S. (C)*, 1971, 3514 (*uv, ir, pmr, struct*)

Papaver Alkaloid A-466

C₂₁H₂₁NO₆

C₂₁H₂₁NO₆ 383.4

Struct. unknown. Alkaloid from the flowers of *Papaver rhoeas*. Yellow prisms. Mp 172-174°.

Awe, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1957, **290**, 367-376 (*isol*)

Hortia Alkaloid A-467

C₂₂H₁₉N₃O₃

C₂₂H₁₉N₃O₃ 373.41

Struct. unknown. Alkaloid from the bark of *Hortia arborea*. Cryst. (Et₂O). Mp 136-139° Mp 191-194° (as hydrochloride).

Antonaccio, L.D. *et al.*, *Ann. Acad. Bras. Cinc.*, 1956, **28**, 183-188; *CA*, **51**, 7385d (*isol*)

Alkaloid CN1 A-468

[126161-02-2]

C₂₃H₂₅NO₅ 395.454

Struct. unknown. Minor alkaloid from the aerial parts of *Corydalis nobilis* (Papaveraceae). Long thin needles (MeOH). Mp 210-211°.

Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1989, **54**, 2009-2020 (*isol, ms*)

Alkaloid CN2 A-469

[126161-13-5]

Struct. unknown. Minor alkaloid from the aerial parts of *Corydalis nobilis* (Papaveraceae). Light yellow prismatic needles (MeOH). Mp 260-261°. Ir 1660 cm⁻¹ (CO), OH band absent.

Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1989, **54**, 2009-2020 (*isol, uv, ir*)

Alkaloid CR1 A-470

C₂₂H₃₄NO₂⁺ 344.516

Struct. unknown. Alkaloid from whole plants of *Consolida regalis*. Mp 288° (as iodide). [α]_D²⁴ +29 (c, 0.2 in MeOH) (iodide). Contains a CO group (1760 cm⁻¹) and an OH group (3320 cm⁻¹). Prob. a quaternary C₂₀-diterpenoid alkaloid.

Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1987, **52**, 804-811 (*isol, uv, pmr, ms*)

Alkaloid CRB 11 A-471

C₂₁H₂₆N₂O₃ 354.448

Struct. unknown. Trace alkaloid from root of *Rauwolfia cumminsii* (Apocynaceae). Off-white amorph. powder. Prob. an indolenine.

Iwu, M.M. *et al.*, *Planta Med.*, 1978, **34**, 390-396 (*isol, uv, ir, ms*)

Alkaloid CSA 3 A-472

CSA 3

Struct. unknown. Mol. formula not reported. Co-occurs with Cryptolepine and Indolol[3,2-*b*]quinoline, I-109. Could possibly be identical with Isocryptolepine in I-110. Alkaloid from roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Mp 300°.

Dwuma-Badu, D. *et al.*, *J. Pharm. Sci.*, 1978, **67**, 433-434 (*isol*)

Rauwolfia obscura Alkaloid D₁ A-473

C₃₀H₃₄N₄O 466.625

Struct. unknown. Bisindole alkaloid. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Mp 190-192°.

Timmins, P. *et al.*, *Planta Med.*, 1975, **27**, 105-111 (*isol, uv, ir, ms*)

Rauwolfia obscura Alkaloid D₂ A-474

C₃₁H₃₆N₄O 480.652

Struct. unknown. Bisindole alkaloid. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Amorph. yellow powder.

Timmins, P. *et al.*, *Planta Med.*, 1975, **27**, 105-111 (*isol, uv, ms*)

Buxus Alkaloid D A-475

Struct. unknown


C₂₉H₅₀N₂O 442.727

Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 182-183°. [α]_D²⁰ +50 (CHCl₃).

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 2209-2226 (*isol*)

C-Alkaloid D A-476

[6902-18-7]



C₄₀H₄₈N₄O₂⁺ 616.845

Alkaloid from calabash curare (*Strychnos* spp.) and from *Strychnos castelneana* (Loganiaceae), also obt. by acid treatment of C-dihydrotoxiciferine I. Shows low curarising activity. [α]_D²⁵ -51 (50% Me₂CO aq.) (as dichloride). Characterised as dichloride and dipicrate, dec. >270° without melting. Red-violet col. with Ce(SO₄)₂ turning yellowish on standing.

▶ EY 5200000

Di-N-de-Me: Bisnor-C-alkaloid D

[62510-51-4]

C₃₈H₄₂N₄O₂ 586.775

Alkaloid from stem bark of *Strychnos dolichothyrsa* and from *Strychnos matopensis*. Poss. artifact. Tentative identification.

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 1864; 1953, **36**, 102; 1954, **37**, 1993; 1961, **44**, 34 (*isol, uv, props*)

Battersby, A.R. *et al.*, *Proc. Chem. Soc., London*, 1961, 413 (*struct*)

Delle Monache, F. *et al.*, *J. Nat. Prod.*, 1970, **33**, 279 (*isol*)


Verpoorte, R. *et al.*, *J. Nat. Prod.*, 1976, **39**, 357-362 (*Bisnor-C-alkaloid D*)

Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304 (*isol, cmr*)

Cassia Alkaloid D A-477

16-(5-Hydroxy-6-methyl-2-piperidinyl)-2,13-hexadecanediol, 9CI. 6-(4,15-Dihydroxyhexadecyl)-3-hydroxy-2-methylpiperidine

[38839-06-4]



C₂₂H₄₅NO₃ 371.602

Minor alkaloid from *Cassia carnavall* whole plant (Fabaceae). Oil. Cryst. salts could not be obt.

Lythgoe, D. *et al.*, *An. Asoc. Quim. Argent.*, 1972, **60**, 317; *CA*, **77**, 164901 (*isol, uv, ir, ms*)

Coelidium Alkaloid D A-478

[11051-60-8]

C₁₆H₂₇N₃ 261.409

Struct. unknown. Alkaloid from *Coelidium fourcadei* dried leaves and branches (Fabaceae). Cryst. (hexane). Mp 138-139°. Interconverts with *Coelidium* Alkaloid E, A-497 in dil. acid. Cooccurs with Isotripiperidine, I-340 and α -Al-dotripiperidine.

Arndt, R.R. *et al.*, *J. S. Afr. Chem. Inst.*, 1968, **21**, 54-57 (*isol, ir, uv, pmr*)

Dendrobates Alkaloid 223D A-479

As *Dendrobates* Alkaloid 167A, A-354 with

R = C₆H₁₃, R' = HC₁₅H₂₉N 223.401

Tentative struct. Minor or trace alkaloid in skin extracts of *Dendrobates histrionicus* (various populations), *Dendrobates minutus*, *Dendrobates pumilio* and *Dendrobates truncatus* (Dendrobatidae). m/e 223(1), 222(2), 138(100). Cannot be hydrogenated.

Hydroxy (?): **Dendrobates Alkaloid 239G**C₁₅H₂₉NO 239.4

Trace alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Mol. formula tentative. m/e 239(1), 238(3), 138(100). Poss. side-chain hydroxy deriv. of 223D. Cannot be hydrogenated.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)


Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 251D A-480

Octahydro-8-methyl-6-(2-methylhexylidene)-8-indolizolinol, 9CI. *Pumiliotoxin 251D*

[73376-35-9]

C₁₆H₂₉NO 251.411

A major alkaloid in skin extracts of the neotropical poison-frogs *Dendrobates tricolor*, *Dendrobates silverstonei*, *Dendrobates bomboetes*, *Dendrobates* sp. (Colombia); minor or trace constit. of *Dendrobates auratus*, *Dendrobates histrionicus*, *Dendrobates lehmanni* and *Dendrobates minutus* (Dendrobatidae).

[α]_D²⁰ -8.5 (c, 1 in CHCl₃) (synthetic).**Hydrochloride**:

Cryst. (hexane/EtOAc). Mp 206-206.5° (evac. capillary). [α]_D²⁵ +32 (c, 0.5 in MeOH) (synthetic).

7 β -Hydroxy: **Allopumiliotoxin 267A**

[73376-38-2]

C₁₆H₂₉NO₂ 267.411

A major alkaloid in *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates fulguritus*, *Dendrobates granuliferus*, *Dendrobates lehmanni*, *Dendrobates leucomelas*, *Dendrobates minutus* (all populations), *Dendrobates tinctorius*; minor or trace constit. in *Dendrobates auratus* (different population), *Dendrobates histrionicus* and *Dendrobates pumilio* (Dendrobatidae). [α]_D²⁵ +24.7 (c, 0.17 in MeOH).

5' ξ -Hydroxy: [94596-77-7]C₁₆H₂₉NO₂ 267.411

Major alkaloid from skin of the Brazilian toad *Melanophryniscus moreirae*. [α]_D²⁵ +7.2 (c, 0.8 in MeOH). Has not been detected in dendrobatid frogs.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms*)

Daly, J.W. *et al.*, *J.A.C.S.*, 1980, **102**, 830 (*isol, pmr, cmr, cryst struct, abs config*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1984, **106**, 4192 (*synth, pmr, cmr, ms*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1984, **40**, 1183 (*deriv*)

Daly, J.W. *et al.*, *Toxicol.*, 1984, **22**, 905 (*Pumiliotoxin 267C*)

Fox, D.N.A. *et al.*, *J.A.C.S.*, 1991, **113**, 2652 (*synth*)

Honda, T. *et al.*, *Heterocycles*, 1992, **34**, 1515 (*synth*)

Goldstein, S.W. *et al.*, *J.O.C.*, 1992, **57**, 1179 (*Allopumiliotoxin 267A, synth*)

Aoyagi, S. *et al.*, *J.A.C.S.*, 1993, **115**, 11393 (*Allopumiliotoxin 267A, synth*)

Honda, T. *et al.*, *J.C.S. Perkin 1*, 1994, 2091 (*synth*)

Cossy, J. *et al.*, *Synlett*, 1996, 909 (*synth*)

Martin, S.F. *et al.*, *Tetrahedron*, 1999, **55**, 8905-8914 (*synth*)

Tang, X.-Q. *et al.*, *J.A.C.S.*, 2000, **122**, 6950-6954 (*Allopumiliotoxin 267A, synth*)

Ni, Y. *et al.*, *J.C.S. Perkin 1*, 2000, 3264-3266 (*synth*)

Comins, D.L. *et al.*, *Org. Lett.*, 2001, **3**, 469-471 (*Allopumiliotoxin 267A, synth*)

Sudau, A. *et al.*, *Eur. J. Org. Chem.*, 2002, 3315-3325 (*synth, pmr, cmr, ms*)

Woodin, K.S. *et al.*, *J.O.C.*, 2007, **72**, 7451-7454 (*synth*)

Dendrobates Alkaloid 253D A-481

2-Allyl-5-(2,3-dihydroxypropyl)decahydroquinoline

As *Dendrobates* Alkaloid 219A, A-357 with

R = -CH₂CH(OH)CH₂OHC₁₅H₂₇NO₂ 253.384

Alkaloid from skin extracts of the Colombian poison-frog *Dendrobates histrionicus* (Dendrobatidae).

Tokuyama, T. *et al.*, *Tetrahedron*, 1986, **42**, 3453 (*isol, pmr, cmr, ms, struct*)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (*pmr, cmr, occur*)

Dendrobates Alkaloid 285D A-482

Struct. unknown. Trace alkaloid from

skin extracts of an undescribed *Dendrobates* sp. (Colombia)(Dendrobatidae). Mol. formula not known; m/e 285 (3), 270 (2), 256 (2), 180 (35), 140 (100). Atypical spectrum for a dendrobatid alkaloid.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-180 (*isol, ms, rev*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Dendrobates Alkaloid 307D A-483C₁₈H₃₁NO₃ 309.448

Struct. unknown. Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mol. formula tentative; m/e 307 (8), 306 (5), 292 (3), 290 (5), 264 (4), 262 (11), 206 (11), 194(18), 166 (100), 70 (85). H₄-deriv. m/e 311, 110, 84, 70.

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Leucojum Alkaloid D A-484

Leucojum Base D

C₁₆H₁₉NO₃ 273.331

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Leucojum* sp. (Amaryllidaceae).

Raffauf, R.F. *et al.*, *Handb. Alkaloids Alkaloid-Containing Plants*, Wiley, 1970, 284


Mantella Alkaloid 211D A-485

Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascan frogs *Mantella* sp. cf. *madagascariensis*, *Mantella betsileo* and *Mantella laevigata*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)

Mantella Alkaloid 243D A-486

8-Ethyl-5-(1,4-heptadien-6-ynyl)octahydroindolizine, 9CI. *Indolizidine 243D* [151834-95-6]

C₁₇H₂₅N 243.391

Provisional struct. Minor or trace alkaloid from skin extracts of 2 populations of the Madagascan frog *Mantella madagascariensis*.


4',5'-Dihydro: 8-Ethyl-5-(1-hepten-6-ynyl)octahydroindolizine, 9CI. **Mantella Alkaloid 245C**. *Indolizidine 245C* [151805-29-7]

C₁₇H₂₇N 245.407

Minor or trace alkaloid in 2 populations of *Mantella madagascariensis*. Provisional struct.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (*isol, ir, ms*)

Mantella Alkaloid 279D A-487
(Octahydro-8-methyl-5-indolizinyI)-sec-nonenol, 9CI. Indolizidine 279D
[151871-23-7]



C₁₈H₃₃NO 279.465

Provisional struct. Secondary alcohol contg. a Z-double bond. Minor or trace alkaloid from skin extracts of 2 populations of the Madagascar frog *Mantella madagascariensis*.

Garraffo, H.M. et al., *J. Nat. Prod.*, 1993, **56**, 1016 (isol, ir, ms)

Melodinus aeneus Alkaloid D A-488
Struct. unknown. Bisindole alkaloid. Alkaloid from *Melodinus aeneus* (Apocynaceae). No phys. props. recorded. M.W. = 534.

Baassou, S. et al., *Phytochemistry*, 1978, **17**, 1449-1450 (isol)

Narcissus Alkaloid D A-489
Narcissus Base D
C₁₇H₁₉NO₃ 285.342
Amaryllidaceae alkaloid. Approx. MF given (may be H₂₁). Struct. unknown. Isol. from *Narcissus* sp. (Amaryllidaceae). Prisms (Me₂CO). Mp 228-229° dec. [α]_D²⁵ -175 (c, 0.2 in CHCl₃).

Boit, H.-G. et al., *Chem. Ber.*, 1957, **90**, 2197-2202 (isol)

Pandaca caducifolia Alkaloid D A-490
[C₄₀H₄₄₋₄₆N₄O₃₋₄]
Struct. unknown. Bisindole alkaloid. Alkaloid from *Pandaca caducifolia* (Apocynaceae). Noncryst. [α]_D +236 (MeOH). Alternative mol. formula C₄₀H₄₆N₄O₄.
Zeches, M. et al., *Phytochemistry*, 1975, **14**, 1122-1124 (isol, uv, ir, pmr, ms)

Salicornia herbacea Alkaloid D A-491
Struct. unknown. Alkaloid from *Salicornia herbacea* (Chenopodiaceae). Mp 178-180°.
Borkowski, B. et al., *Pharmazie*, 1965, **20**, 390-393; *CA*, **63**, 15223c

Stephania glabra Alkaloid D A-492
C₂₀H₂₅NO₄ 343.422
Struct. unknown. Isol. from tubers of *Stephania glabra*. Cryst. (C₆H₆). Mp 153-154°. Conts. 2 OMe and 1 NMe groups. Not reported in further investigations of this species. λ_{max} 265 (no solvent reported).
Hydrochloride: Mp 227-228°.
Hydrobromide: Mp 229-230°.
Kin, F.K. et al., *Khim. Prir. Soedin.*, 1965, **1**,

392-394; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 308-309 (isol)
Cava, M.P. et al., *J.O.C.*, 1968, **33**, 2785-2789 (*Stephania glabra constits*)

Alkaloid D1† A-493
Prob. of Usambarine type. Struct. unknown. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Off-white cryst. Mp 190-192°.
Timmins, P. et al., *Planta Med.*, 1975, **27**, 105-111 (isol, uv, ir, pmr, ms)


Alkaloid D2 A-494
Prob. of Usambarine type. Struct. unknown. Alkaloid from the leaves of *Rauwolfia obscura* (Apocynaceae). Amorph. yellow powder.
Timmins, P. et al., *Planta Med.*, 1975, **27**, 105-111

Alkaloid 267D A-495
[97380-35-3]
C₁₆H₂₉NO₂ 267.411
Struct. unknown. Alkaloid from skin extracts of the Australian frog *Pseudophryne semimarmorata* (Myobatrachidae) and the Madagascar frog *Mantella aurantiaca* (family Ranidae, subfamily Mantellinae). m/e 267 (13), 250 (10), 194 (16), 166 (100), 70(80). Forms H₂-deriv. Two exchangeable hydrogens. Has not been detected in dendrobatid frogs.
Daly, J.W. et al., *Toxicol.*, 1984, **22**, 905-919 (isol, ms)

Buxus Alkaloid E A-496
C₂₇H₅₀N₂O₃ 450.704
Struct. unknown. Alkaloid from *Buxus balearica* (Buxaceae). Mp 287-289°. [α]_D²⁰ +12 (c, 0.6 in CHCl₃/MeOH).
Kurakina, I.O. et al., *Khim. Prir. Soedin.*, 1969, **5**, 26-28; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 20-21 (isol)

Coelidium Alkaloid E A-497
[11051-61-9]
C₁₆H₂₇N₃ 261.409
Struct. unknown. Alkaloid from *Coelidium fourcadei* dried leaves and branches (Fabaceae). Cryst. (hexane). Mp 148-149°. Interconverts with *Coelidium* Alkaloid D, A-478 in dil. acid. Cooccurs with α-aldotripiperidine and Isotripiperidine, I-340.
Arndt, R.R. et al., *J. S. Afr. Chem. Inst.*, 1968, **21**, 54-57 (isol, ir, uv, pmr)

Croton linearis Alkaloid E A-498
[58206-97-6]



C₁₈H₂₃NO₃ 301.385
Belongs to diastereomeric series to alkaloids covered by Amuroline, A-957 and Cryprochine, C-781. Alkaloid from *Croton linearis* (Euphorbiaceae).
Di-Ac: Mp 190-193°. [α]_D²⁶ -18 (c, 0.5 in MeOH).

4-Ketone, N-de-Me: Jacularine
[20361-70-0]
C₁₇H₁₉NO₃ 285.342
Alkaloid from *Croton linearis* (Euphorbiaceae). Enantiomer of Isocrotosparinine in A-957.
Haynes, L.J. et al., *J.C.S. (C)*, 1966, 1680-1685 (*Alkaloid E*)
Stuart, K.L. et al., *Tet. Lett.*, 1968, 4473-4474 (*Jacularine*)
Casagrande, C. et al., *J.C.S. Perkin I*, 1975, 1659-1663 (stereochem)

Dendrobates Alkaloid 223E A-499
C₁₄H₂₅NO 223.358
Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates lehmanni*, *Dendrobates occulator* and *Dendrobates tinctorius* (Dendrobatidae). Mol. formula tentative; m/e 223 (2), 222 (3), 168 (100). Possibly an analogue of *Dendrobates* Alkaloid 225, A-291 with double bond in side chain. H₂-deriv, m/e 223, 168.
Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)
Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Dendrobates Alkaloid 237E A-500
C₁₅H₂₇NO 237.384
Struct. unknown. Trace alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobatidae). Mol. formula tentative; m/e 237 (1), 236 (3), 208 (70), 152 (100). Possibly a hydroxypumiliotoxin C. H₂-deriv, m/e 239, 152.
Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)
Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Lycopodium Alkaloid E A-501
Lycopodium Base E
C₁₇H₂₅NO₂ 275.39
Struct. unknown. Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Mp 267-269° dec. (as perchlorate).
Burnell, R.H. et al., *J.C.S.*, 1959, 3091 (isol, ir)

Muntafara sessilifolia Alkaloid E A-502
Struct. unknown. Bisindole alkaloid. Alkaloid from *Muntafara sessilifolia* (preferred genus name *Tabernaemontana*) (Apocynaceae). Other dimeric alkaloids isol. from *M. sessilifolia* designated F, I, J, N, O, P, Q and R. No phys. props. recorded.
Panas, J.M. et al., *Phytochemistry*, 1975, **14**, 1120-1122 (isol)

Pandaca caducifolia Alkaloid E A-503
C₄₀H₄₆N₄O₄ 646.828

Struct. unknown. Bisindole alkaloid.

Alkaloid from *Pandaca caducifolia* (Apocynaceae). Noncryst. $[\alpha]_D^{25}$ -51 (MeOH).

Zeches, M. *et al.*, *Phytochemistry*, 1975, **14**, 1122-1124 (*isol*, *uv*, *ir*, *pmr*)

Skytanthus acutus Alkaloid E A-504

$C_{10}H_{19}N$ 153.267

Struct. unknown. Monoterpene alkaloid. Alkaloid from *Skytanthus acutus* (Apocynaceae). $Bp_{1.5}$ 120-140°.

Appel, H.H. *et al.*, *Scientia (Valparaiso)*, 1961, **28**, 5-9; *CA*, **57**, 2332g

Sophora microphylla Alkaloid E A-505

Sophora Base E

Struct. unknown. Alkaloid from *Sophora microphylla* (Fabaceae). Mp 168-171°.

Briggs, L.H. *et al.*, *J.C.S.*, 1937, 1795 (*isol*)

Voacanga thouarsii Alkaloid E A-506

[50924-06-6]

$C_{43}H_{50}N_4O_7$ 734.891

Bisindole alkaloid. Struct. unknown. Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Cryst. (MeOH). Mp 235-238°. $[\alpha]_D^{25}$ -140. Intense blue col. with HNO_3 . λ_{max} 225 (log ϵ 4.66); 265 (log ϵ 4.08); 283 (sh) (log ϵ 3.96); 291 (sh) (log ϵ 3.9) (EtOH).

Rolland, Y. *et al.*, *Phytochemistry*, 1973, **12**, 2039-2042 (*isol*, *uv*, *ir*, *pmr*)

Buxus Alkaloid F A-507

Struct. unknown. Alkaloid from

Buxus wallichiana (Buxaceae). Mp 190°. $[\alpha]_D^{25}$ +26.6 (c, 0.76 in EtOH). MW 442.

Vassová, A. *et al.*, *Pharmazie*, 1970, **25**, 363; *CA*, **73**, 127748n

Dendrobates Alkaloid 223F A-508


$C_{14}H_{25}NO$ 223.358

Struct. unknown. Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mol. formula tentative; m/e 223 (1), 138 (100). H_2 -deriv, m/e 225, 182.

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms*, *rev*)

Dendrobates Alkaloid 251F A-509

Dodecahydro-3,7,10-trimethylcyclopenta[b]quinolizine-2-methanol, 9CI. Indolizidine 251F [142543-01-9]



R = H

Absolute Configuration

$C_{16}H_{29}NO$ 251.411

Major alkaloid from the skin extracts of the Colombian dendrobatid frog *Minyobates bombetes*. Pale yellow oil. $[\alpha]_D^{25}$ -11.1 (c, 0.96 in MeOH).

Aldehyde: Dodecahydro-3,7,10-trimethylcyclopenta[b]quinolizine-2-carboxaldehyde, 9CI. Dendrobates Alkaloid 249B. Indolizidine 249B [142543-05-3]

$C_{16}H_{27}NO$ 249.395

Alkaloid from *Minyobates bombetes*. Tentative struct.

2,3,6,7-Tetrahydro: Dendrobates Alkaloid 247 [142543-06-4]

$C_{16}H_{25}NO$ 247.38

Alkaloid from *Minyobates bombetes*. Tentative struct.

2,3,6,7-Tetrahydro, aldehyde: Dendrobates Alkaloid 245 [142543-07-5]

$C_{16}H_{23}NO$ 245.364

Alkaloid from *Minyobates bombetes*. Tentative struct.

Deoxy: Dodecahydro-2,3,7,10-tetramethylcyclopenta[b]quinolizine, 9CI. Dendrobates Alkaloid 235H. Indolizidine 235H [142543-02-0]

$C_{16}H_{29}N$ 235.412

Alkaloid from *Minyobates bombetes*. Tentative struct.

Deoxy, 2 ζ -hydroxy: Dodecahydro-2,3,7,10-tetramethylcyclopenta[b]quinolizine-2-ol, 9CI. Dendrobates Alkaloid 251J. Indolizidine 251J [142543-08-6]

$C_{16}H_{29}NO$ 251.411

Alkaloid from *Minyobates bombetes*. Tentative struct.

2-Epimer: Dendrobates Alkaloid 251F'. Indolizidine 251F' [142631-46-7]

$C_{16}H_{29}NO$ 251.411

Alkaloid from *Minyobates bombetes*. Tentative struct.

Homologue (R = CH_3): Dodecahydro-3,4,7,10-tetramethylcyclopenta[b]quinolizine-2-methanol, 9CI. Dendrobates Alkaloid 265B. Indolizidine 265B [142543-03-1]

$C_{17}H_{31}NO$ 265.438

Alkaloid from *Minyobates bombetes*. Tentative struct.

Homologue (R = CH_2CH_3): 4-Ethyl-dodecahydro-3,7,10-trimethylcyclopenta[b]quinolizine-2-methanol, 9CI. Dendrobates Alkaloid 279B. Indolizidine 279B [142543-04-2]

$C_{18}H_{33}NO$ 279.465

Alkaloid from *Minyobates bombetes*. Tentative struct.

Homologue (R = CH_2CH_3), deoxy, 2 ζ -hydroxy: 4-Ethyl-dodecahydro-2,3,7,10-tetramethylcyclopenta[b]quinolizine-2-ol, 9CI. Dendrobates Alkaloid 279C. Indolizidine 279C [142543-09-7]

$C_{18}H_{33}NO$ 279.465

Alkaloid from *Minyobates bombetes*.


Tentative struct.

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1992, **55**, 707-722 (*isol*, *pmr*, *ms*)

Taber, D.F. *et al.*, *J.A.C.S.*, 1995, **117**, 5757-5762 (*synth*, *pmr*, *cmr*)

Wroblewski, A. *et al.*, *J.A.C.S.*, 2004, **126**, 5475-5481 (*synth*)

Mantella Alkaloid 221F A-510



$C_{14}H_{23}NO$ 221.342


Structure tentatively reassigned in 2005. Trace alkaloid from skin extracts of the Madagascan frog *Mantella aurantiaca*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*, *ms*)

Adriamaharavo, N.R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1743-1748 (*struct*)

Mantella Alkaloid 241F A-511

8-(3-Butynyl)octahydro-5-(2-penten-4-ynyl)indolizine, 9CI. Indolizidine 241F [151805-25-3]



$C_{17}H_{23}N$ 241.375

Provisional struct. Minor alkaloid from skin extracts of 2 populations of the Madagascan frog *Mantella madagascariensis*.

2',3'-Dihydro: 8-(3-Butynyl)octahydro-5-(4-pentenyl)indolizine, 9CI. Mantella Alkaloid 243B. Indolizidine 243B [151805-26-4]

$C_{17}H_{25}N$ 243.391

Trace alkaloid in one population of *Mantella madagascariensis*. Provisional struct.

3'',4''-Dihydro: 8-(3-Butenyl)octahydro-5-(2-penten-4-ynyl)indolizine, 9CI. Mantella Alkaloid 243C. Indolizidine 243C [151805-27-5]

$C_{17}H_{25}N$ 243.391

Minor or trace alkaloid in 2 populations of *Mantella madagascariensis*. Provisional struct.

2',3',3'',4''-Tetrahydro: 8-(3-Butenyl)octahydro-5-(4-pentenyl)indolizine, 9CI. Mantella Alkaloid 245B. Indolizidine 245B [151805-28-6]

$C_{17}H_{27}N$ 245.407

- Trace alkaloid in 1 population of *Mantella madagascariensis*. Provisional struct.
Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (*isol, ir, ms*)
- Mantella Alkaloid 249F** A-512
Homopumiliotoxin 249F
Struct. and MF unknown. Trace alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.
Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)
- Mantella Alkaloid 265F** A-513
 $C_{16}H_{27}NO_2$ 265.395
Struct. unknown. May be related to alkaloids of the 235C class. Trace alkaloid from skin extracts of the Madagascar frog *Mantella aurantiaca*.
Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ms*)
- Mantella Alkaloid 281F** A-514
Struct. unknown. Probably belongs to the pumiliotoxin A class of alkaloid (see Pumiliotoxin A, P-791). Alkaloid from skin extracts of two populations of the Madagascar frog *Mantella madagascariensis*.
Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)
- Pandaca caducifolia Alkaloid F** A-515
 $C_{40}H_{46}N_4O_4$ 646.828
Struct. unknown. Bisindole alkaloid. Alkaloid from *Pandaca caducifolia* (Apocynaceae). Noncryst. $[\alpha]_D +56$ (MeOH).
Zeches, M. *et al.*, *Phytochemistry*, 1975, **14**, 1122-1124 (*isol, uv, ir, pmr*)
- Lycopodium Alkaloid G** A-516
Lycopodium Base G
 $C_{18}H_{27}NO_3$ 305.416
Struct. unknown. Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Mp 198-200° (as perchlorate).
Burnell, R.H. *et al.*, *J.C.S.*, 1959, 3091 (*isol, ir*)
- Mantella Alkaloid 207G** A-517
Homopumiliotoxin 207G
Struct. and MF unknown. Trace alkaloid from skin extracts of the Madagascar frog *Mantella viridis*.
Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ms*)
- Pandaca caducifolia Alkaloid G** A-518
 $C_{40}H_{46}N_2O_4$ 618.814
Struct. unknown. Bisindole alkaloid. Alkaloid from *Pandaca caducifolia* (Apocynaceae). Noncryst. $[\alpha]_D +35$ (MeOH).
Zeches, M. *et al.*, *Phytochemistry*, 1975, **14**, 1122-1124 (*isol, uv, ir, pmr*)
- Voacanga thouarsii Alkaloid G** A-519
[50923-99-4]
 $C_{42}H_{50}N_4O_5$ 690.881
Bisindole alkaloid. Struct. unknown. Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Cryst. (Et₂O). Mp 280-285° dec. $[\alpha]_D +46$. Pink col. with HNO₃. λ_{max} 221 (log ϵ 4.63); 260 (log ϵ 4.29); 298 (log ϵ 3.7) (EtOH).
Rolland, Y. *et al.*, *Phytochemistry*, 1973, **12**, 2039-2042 (*isol, uv, ir, pmr*)
- Alkaloid 251G** A-520
[97380-38-6]
 $C_{15}H_{25}NO_2$ 251.368
Struct. unknown. Alkaloid from skin extracts of the Madagascar frog *Mantella aurantiaca* (family Ranidae, subfamily Mantellinae). m/e 251 (26), 250 (45), 162 (100). Forms H₂-deriv. Two exchangeable hydrogens. Hydroxy congener of Alkaloid 235C. Has not been detected in dendrobatid frogs.
Daly, J.W. *et al.*, *Toxicon*, 1984, **22**, 905-919 (*isol, ms*)
- Alkaloid GB14** A-521
 $C_{24}H_{33}NO_5$ 415.528
Struct. unknown. Alkaloid from the bark of *Himantandra belgraveana* (*Galbulimima belgraveana*) (Himantandraceae). Prisms + 1H₂O (Me₂CO/heptane). Mp 106°.
Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)
- Alkaloid GB15** A-522
 $C_{22}H_{35}NO_3$ 361.523
Struct. unknown. Alkaloid from the bark of *Himantandra* (*Galbulimima*) *belgraveana* (Himantandraceae). Needles (C₆H₆). Mp 230°. $[\alpha]_D +58$ (c, ca. 1 in CHCl₃).
Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)
- Alkaloid GB16** A-523
 $C_{20}H_{27}NO_2$ 313.439
Struct. unknown. Alkaloid from the bark of *Himantandra* (*Galbulimima*) *belgraveana* (Himantandraceae). Prisms (EtOAc). Mp 203°. $[\alpha]_D +550$ (c, ca. 1 in CHCl₃).
Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)
- Alkaloid GB17** A-524
 $C_{21}H_{31}NO_3$ 345.481
Struct. unknown. Alkaloid from the bark of *Himantandra* (*Galbulimima*) *baccata* (Himantandraceae). Needles + 1H₂O (MeOH aq.). Mp 115°. $[\alpha]_D -21$ (c, ca. 1 in CHCl₃).
Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)
- Alkaloid GB18** A-525
 $C_{22}H_{33}NO_3$ 359.508
Struct. unknown. Alkaloid from the bark of *Himantandra baccata* (preferred genus name *Galbulimima*) (Himantandraceae). Prisms (heptane). Mp 120°. $[\alpha]_D +37$ (c, ca. 1 in CHCl₃).
Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)
- Gentiana Alkaloid II** A-526
Struct. unknown. Monoterpene alkaloid. Alkaloid from *Gentiana asclepiadea*, *Gentiana bulgarica*, *Gentiana cruciata*, *Gentiana lutea* and *Gentiana punctata* (Gentianaceae).
Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1965, **18**, 947; *CA*, **64**, 10084h (*isol, ir*)
Cordell, G.A. *et al.*, *Alkaloids* (*Academic Press*), 1977, **16**, 431 (*rev*)
- Sternbergia Alkaloid III** A-527
Sternbergia Base III
Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Sternbergia fischeriana* (Amaryllidaceae). Mp 212-213°. Opt. inactive, mol. formula not recorded.
Hydrobromide: Mp 174-175°. *Picrate*: Mp 217-219°.
Proskurnina, N.F. *et al.*, *Chem. Zentralbl.*, 1954, **125**, 8353 (*isol*)
- Solanum Alkaloid IV** A-528
Struct. unknown. A glycoalkaloid cont. NH, OH and conjugated C=O. Isol. from epigeal parts of *Solanum nigrum* (Solanaceae). Cryst. (MeOH). Mp 320-325°. Cooccurs with Solasonine, Solamargine and β -Solamargine (see Spirosol-5-en-3-ol, S-456).
Aslanov, S.M. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 674; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 658
- Ungernia Alkaloid IV** A-529
Ungernia Base IV
Amaryllidaceae alkaloid. Struct. unknown. Isol. from the leaves of *Ungernia minor* (Amaryllidaceae). Cryst. (MeOH). Mp 193-194°.
Normatov, M. *et al.*, *Uzb. Khim. Zh.*, 1965, **9**, 25-30; *CA*, **63**, 7061f (*isol*)
- Gentiana cruciata Alkaloid IVb** A-530
 $C_{10}H_{16}N_2O_7$ 276.246
Struct. unknown. Monoterpene alkaloid. Alkaloid from *Gentiana cruciata* (Gentianaceae). Mp 138-140°.
Marekov, N. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1965, **18**, 999-1002; *CA*, **64**, 11270a (*isol, ir*)
Cordell, G.A. *et al.*, *Alkaloids* (*Academic Press*), 1977, **16**, 431-510 (*rev*)
- Lycopodium annotinum Alkaloid IX** A-531
Lycopodium Base IX
 $C_{17}H_{25}NO_2$ 275.39
Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae).

Methiodide: Mp 324°. $[\alpha]_D^{20}$ -49.8.

Methochloride: Mp 274°.

Methoperchlorate: Mp 316°.

Methopicrate: Mp 78-80°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol, uv*)

C-Alkaloid J A-532

C₁₉H₂₁N₂ 277.388

Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 260° dec. (as picrate). Red-orange col. with Ce(SO₄)₂.

Kebrle, J. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 102-121 (*isol, uv*)

Mantella Alkaloid 207J A-533

Struct. and MF unknown. Possibly a homologue of Precocinelline in D-886. Trace alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)

Mantella Alkaloid 235J A-534

Homopumiliotoxin 235J

Struct. and MF unknown. Trace alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)

Mantella Alkaloid 235K A-535

Struct. and MF unknown. Considered to be a tricyclic alkaloid. Trace alkaloid from skin extracts of the Madagascar frog *Mantella betsileo*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ms*)

Alkaloid K2 A-536

C₂₁H₂₇NO₆ 389.447


Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Mp 226-228°.

Methiodide: Mp 247-249°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K4 A-537

2-Demethylalcolcolchicine



C₂₁H₂₃NO₆ 385.416

(S)-form

Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Mp 238-240° (natural) Mp 259-260° (semisynthetic). $[\alpha]_D^{20}$ -152 (c, 0.92 in MeOH) (semisynthetic).

Ac:

Cryst. (Et₂O/CHCl₃). Mp 230-232°.

$[\alpha]_D^{22}$ -150 (c, 1.0 in MeOH).

Me ether: Allocolchicine. Suhailamine.

Alkaloid K3

[641-28-1]

C₂₂H₂₅NO₆ 399.443

Alkaloid from the flowers of *Colchicum autumnale*, from *Colchicum decaisnei* and above ground parts of *Colchicum kesselringii* (Liliaceae). Cryst. (Et₂O/CHCl₃) or amorph. solid. Mp 257-258°. $[\alpha]_D^{22}$ -150 (c, 1.2 in MeOH) (semisynthetic). First descr. as Alkaloid K3 without abs. config. or opt. rotn. Suhailamine (1991) assigned (S)-config. However, spectroscopic and phys. props. of authentic Allocolchicine differ from those of Suhailamine and therefore the true struct. of the latter remains unclear.

►HO7884000

Yusupov, M.K. *et al.*, *Zh. Obshch. Khim.*, 1964, **34**, 1677-1680; *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1686-1688 (*isol struct*)

Yusupov, M.K. *et al.*, *Khim. Prir. Soedin. (Engl. Transl.)*, 1973, **9**, 188-189

Mackay, M.F. *et al.*, *Acta Cryst. C*, 1989, **45**, 795-799 (*Allocolchicine, cryst struct*)

Abu Zarga, M.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 936-940 (*Suhailamine*)

Banwell, M.G. *et al.*, *Chem. Comm.*, 1994, 2647-2649 (*Suhailamine*)

Leblanc, M. *et al.*, *Org. Lett.*, 2005, **7**, 2849-2852 (*synth*)

Alkaloid K5 A-538

C₁₉H₂₅NO₄ 331.411

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Mp 224-226°.

Hydrochloride: Mp 239-241°.

Methiodide: Mp 278-280°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K6 A-539

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae).

Methiodide: Mp 253-254°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K7 A-540

C₂₁H₂₉NO₄ 359.464

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Mp 144-146°.

Hydrochloride: Mp 232-234°.

Methiodide: Mp 247-248°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K9 A-541

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae).

Methiodide: Mp 245-247°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K10 A-542

C₁₈H₂₃NO₄ 317.384

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Cryst. (Me₂CO/MeOH). Mp 232-234°.

Hydrochloride: Mp 261-263°.

Methiodide: Mp 249-251°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

Alkaloid K11 A-543

Struct. unknown. Alkaloid from the above-ground parts of *Colchicum kesselringii* (Liliaceae). Mp 274-276°.

Yusupov, M.K. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 1682 (*isol*)

C-Alkaloid L A-544

Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Prisms (Me₂CO aq.) (as picrate). Mp 171° (picrate). Red col. with Ce(SO₄)₂.

Kebrle, J. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 102-121 (*isol, uv*)

Colubrina Alkaloid L A-545

C₃₇H₄₀N₂O₆ 608.733

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the stem bark of *Colubrina faralaotra* ssp. *faralaotra* (Rhamnaceae).

Guinaudeau, H. *et al.*, *Planta Med.*, 1975, **27**, 304-318 (*isol, uv, pmr, ms*)

Mantella Alkaloid 251L A-546

Homopumiliotoxin 251L

Struct. and MF unknown. Occurs mainly as the O-Ac deriv. Trace alkaloid from skin extracts of one population of the Madagascar frog *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol, ir, ms*)

Alkaloid L10 A-547


C₁₆H₂₇NO 249.395

Struct. unknown. Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Mp 223° (as perchlorate).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 92-96 (*isol*)

Alkaloid LA 4 A-548

[116097-93-9]



R = H

C₁₉H₂₂N₂O₄ 342.394

Alkaloid artifact generated by NH₃ and secoiridoid glucoside from ripe fruits of the common privet *Ligustrum vulgare* (Oleaceae). Semi-solid. $[\alpha]_D^{20}$ -0.23 (c, 1.0 in EtOH).

Willems, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 229 (*isol, uv, ir, pmr, cmr, ms, struct*)

Alkaloid LA 5

A-549

[115531-75-4]

As Alkaloid LA 4, A-548 with

R = CH₃C₂₀H₂₄N₂O₄ 356.421


Alkaloid artifact generated by NH₃ and secoiridoid glucosides from ripe fruit of the common privet *Ligustrum vulgare* (Oleaceae). Semi-solid. $[\alpha]_D^{20}$ -0.39 (c, 1.0 in EtOH).

Willems, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 229 (*isol, uv, ir, pmr, cmr, ms, struct*)

Alkaloid LA 6

A-550

[113145-61-2]

C₁₉H₂₁NO₅ 343.379


Alkaloid artifact generated by NH₃ and secoiridoid glucosides from ripe fruits of the common privet *Ligustrum vulgare* (Oleaceae). Needles (EtOAc/MeOH). Mp 115-116°.

Willems, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1987, **320**, 1245 (*isol, uv, ir, pmr, cmr, ms, struct*)

Alkaloid LA 9

A-551

[116057-84-2]

C₂₅H₃₁NO₁₀ 505.521

Alkaloid artifact generated by NH₃ and secoiridoid glucosides from ripe fruits of the common privet *Ligustrum vulgare* (Oleaceae). Cryst. (EtOAc/MeOH). Mp 64-66°. $[\alpha]_D^{20}$ -0.1 (c, 1.0 in EtOH).

Willems, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 357 (*isol, uv, ir, pmr, cmr, struct*)

Alkaloid LB1

A-552

Prob. a spermidine related alkaloid.


Struct. unknown. Minor alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Mp 220° dec.

Hairs, E. *et al.*, *Bull. Acad. R. Med. Belg.*, 1909, 1042-1048; *CA*, 1892, **4**, 1042 (*isol*)

Alkaloid LBZ

A-553

[38143-11-2]

C₂₆H₃₃N₃O₄ 451.564

CAS numbering shown. Minor alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Amorph.

22-Ketone: **Alkaloid LBX**

[38143-10-1]

C₂₆H₃₁N₃O₄ 449.549

Minor alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Cryst. (MeOH/Me₂CO or MeOH/Et₂O). Mp 250° dec. $[\alpha]_D^{20}$ +201 (c, 0.6 in CHCl₃).


Doskotch, R.W. *et al.*, *Experientia*, 1972, **28**, 382 (*struct*)

Poupat, C. *et al.*, *Tetrahedron*, 1972, **28**, 3087 (*uv, ir, pmr, struct*)

Alkaloid LC2

A-554

3-(3-Butenyl)-1,2,3,4,5,6,11,11a-octahydro-1,5-methano-10H-pyrido[1,2-a][1,5]diazocin-10-one, 9CI. 1,13-Didehydro-10,11-secomultiflorine. 11,12-Seco-12-dehydromultiflorine. N-Methylalbine (*incorr.*) [71635-26-2]

C₁₅H₂₂N₂O 246.352

Struct. revised in 1988. Some isolates were prev. reported erroneously as N-methylalbine (see Albine, A-243). Alkaloid from the leaves and seeds of *Lupinus cosentinii*, *Lupinus albus* and *Lupinus formosus* (Fabaceae). Cryst. (cyclohexane). Mp 60°. $[\alpha]_D^{25}$ -520 (c, 1.0 in MeOH).

[97906-66-6, 6822-63-5]

Beck, A.B. *et al.*, *J. Nat. Prod.*, 1979, **42**, 385 (*isol, uv, ir, pmr, ms, struct*)

Wysocka, W. *et al.*, *Planta Med.*, 1988, **54**, 522 (*cmr, ms, struct*)

Brukwicki, T. *et al.*, *J. Mol. Struct.*, 1989, **196**, 343 (*cmr, conform, bibl*)


Wyrzykiewicz, E. *et al.*, *Org. Mass Spectrom.*, 1990, **25**, 453 (*ms*)

Dutkiewicz, G. *et al.*, *Acta Cryst. C*, 1995, **51**, 1182 (*cryst struct*)

Alkaloid LC7

A-555

Alkaloid LV3†



Probable structure

C₄₀H₅₄N₂O₈ 690.875

Truxillate-type dimer of Epilupinyl transferulate in E-95. Alkaloid from leaves of

Lupinus cosentinii (Fabaceae). Cryst.

(MeOH/CHCl₃). Mp 228°. $[\alpha]_D^{25}$ +23 (c, 1 in CHCl₃/MeOH 1:1).

Beck, A.B. *et al.*, *J. Nat. Prod.*, 1979, **42**, 385 (*isol, struct, uv, ir, pmr, ms*)

Alkaloid LE1

A-556

C₂₈H₄₇N₃O 441.699

Struct. unknown. Alkaloid from *Lycopodium erythraeum* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LO1

A-557

C₃₁H₃₉N₃O₂ 485.668

Struct. unknown. Alkaloid from *Lycopodium obtusifolium* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LO2

A-558

C₃₀H₄₉N₃O 467.737

Struct. unknown. Alkaloid from *Lycopodium obtusifolium* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LO3

A-559

C₂₉H₄₉N₃ 439.726

Struct. unknown. Alkaloid from *Lycopodium obtusifolium* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LS14

A-560

C₃₀H₄₉N₃O 467.737

Struct. unknown. Alkaloid from *Lycopodium saururus* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LV1†

A-561

C₁₅H₂₂N₂O 246.352

Not the same as Alkaloid LV1, A-562.

Struct. unknown. Alkaloid from *Lupinus digitatus* (Fabaceae). Needles (Me₂CO/petrol). Mp 108-109°. $[\alpha]_D$ -314 (MeOH). Deep-red col. with FeCl₃.

Perchlorate:

Cream needles + ½H₂O (EtOH). Mp 160-162° dec. Cryst. with difficulty.

Methiodide:

Cream nodules (EtOH). Mp 246-248° dec.

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1955, **8**, 136-139; 1957, **10**, 177-181

Gladstones, J.S. *et al.*, *J. R. Soc. West. Aust.*, 1958, **41**, 29

Alkaloid LV1†

A-562

C₂₉H₄₅N₃ 435.695

Not the same as Alkaloid LV1, A-561. Struct. unknown. Alkaloid from *Lycopodium verticillatum* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Alkaloid LV2

A-563

C₁₅H₂₄N₂O₂ 264.367

Struct. unknown. Tentative mol. formula. Trace alkaloid from seeds of *Lupinus varius* (Fabaceae).

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1955, **8**, 136-139

Alkaloid LV3† A-564

C₂₀H₂₇N₄ 345.438

Struct. unknown. Alkaloid from *Lupinus digitatus* (Fabaceae). Needles (MeOH). Mp 232-233° dec. Conts. one OMe group. λ_{max} 235 nm. Possibly the monomer of Alkaloid LC7, A-555.

Picrate:

Yellow needles (Me₂CO aq.). Mp 186°. Dec. sharply at 231°.

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1957, **10**, 177-181

Gladstones, J.S. *et al.*, *J. R. Soc. West. Aust.*, 1958, **41**, 29

Alkaloid LV4 A-565

C₁₇H₂₃N₅ 321.372

Provisional mol. formula. Struct. unknown. Alkaloid from *Lupinus digitatus* (Fabaceae). Needles (CHCl₃/petrol). Mp 235-237° dec.

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1957, **10**, 177-181

Gladstones, J.S. *et al.*, *J. R. Soc. West. Aust.*, 1958, **41**, 29

Buxus Alkaloid M† A-566

See also *Buxus* Alkaloid M, A-567.

Struct. unknown. Alkaloid from *Buxus wallichiana* (Buxaceae). Mp 261°. [α]_D²³ +27 (c, 0.33 in CHCl₃).

Vassová, A. *et al.*, *Pharmazie*, 1970, **25**, 363-365; *CA*, **73**, 127748n

Buxus Alkaloid M† A-567

C₂₇H₄₆N₂O 414.673

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 203-205°. [α]_D -80 (CHCl₃).

Friedrich, W. *et al.*, *Helv. Chim. Acta*, 1950, **33**, 873-878 (*isol*)

C-Alkaloid M A-568

C₂₀H₂₃N₂O₂⁺ 323.414

Struct. unknown. Minimum formula. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Slowly developing yellow col. with Ce(SO₄)₂.

Asmies, H. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1968-1973 (*isol, uv*)

Alkaloid MB 10 A-569

C₂₃H₃₀N₂O₅ 414.5

Struct. unknown. Alkaloid from roots of *Rauwolfia mombasiana* (Apocynaceae). Off-white amorph. powder. Conts. 2 OMe groups.

Iwu, M.M. *et al.*, *Planta Med.*, 1980, **38**, 260-263 (*isol, uv, ir, ms*)

Buxus Alkaloid N A-570

C₂₂H₃₅N₂O 345.524


Struct. unknown. Poss. identical with Buxtauine M, B-482. Alkaloid from

Buxus sempervirens (Buxaceae). Mp 178-179°. [α]_D +150 (CHCl₃). Props. corresp. to Buxtauine M, B-482.

Friedrich, W. *et al.*, *Helv. Chim. Acta*, 1950, **33**, 873-878 (*isol*)

Alkaloid ND 370 A-571

[37304-95-3]



C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Cryst. by subl. Mp 209-211°.


MacLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**, 1496; 1976, **54**, 1262 (*uv, ir, pmr, ms, struct*)

C-Alkaloid O A-572

19,20-Didehydro-9-methoxy-4,17-cyclo-corynanium, 9CI

[86138-68-3]

[86138-69-4]



Absolute Configuration

C₂₀H₂₅N₂O⁺ 309.43

Quaternary alkaloid from calabash curare and *Strychnos guianensis* (Loganiaceae). Mp 288-289° dec. (as chloride). [α]_D²⁵ -150 (c, 0.283 in 5%Py aq.) (as chloride). λ_{max} 227 (log ε 4.1); 266 (log ε 3.88); 282 (log ε 3.77); 291 (log ε 3.73) (MeOH).

Giesbrecht, E. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1974-1982 (*isol*)

Borris, R.P. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 405 (*struct, spectra*)

Penelle, J. *et al.*, *Phytochemistry*, 2000, **53**, 1057-1066; 2001, **58**, 619-626 (*isol, uv, ir, cd, pmr, cmr, activity*)

Alkaloid Or₂ A-573

Struct. unknown

Alkaloid from *Papaver orientale* and *Papaver pseudo-orientale* (Papaveraceae). Cryst. (petrol). Mp 124-127° (120-121°). Unstable, not obt. completely pure. m/e 333.

Délenk-Heydenreich, K. *et al.*, *Pharmazie*, 1969, **24**, 635-645 (*isol, uv, ms*)

Shafiee, A. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 1570-1572 (*isol, uv, pmr*)

Baptisia Alkaloid P₂ A-574

C₁₁H₁₈N₂O 194.276

Struct. unknown. Isol. from *Baptisia*

australis and *Baptisia perfoliata*. Cryst. (MeOH/Et₂O). Mp 300°.

Perchlorate:

Needles (MeOH). Mp 198°.

Marion, L. *et al.*, *J.A.C.S.*, 1948, **70**, 691-692; 3253-3254 (*isol*)

Oung-Boran, *et al.*, *Planta Med.*, 1969, **17**, 301-318 (*isol*)

Baptisia Alkaloid P₄ A-575

Struct. unknown. Isol. from *Baptisia minor*. Oil. Bp_{0.05} 175-200°.

Perchlorate:

Needles. Mp 286°.

Marion, L. *et al.*, *J.A.C.S.*, 1948, **70**, 3472-3474 (*isol*)

C-Alkaloid P A-576

C₂₀H₂₃N₂O⁺ 307.414

Struct. unknown. Quaternary alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 224-232° (as picrate). Blue col. with Ce(SO₄)₂.

Giesbrecht, E. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1974-1982 (*isol, uv*)

Lupinus Alkaloid P1 A-577

C₁₅H₂₂N₂O 246.352

Tentative mol. form. Struct. unknown. Alkaloid from *Lupinus macoumii* (Fabaceae). Small plates (EtOH/petrol). Mp 126° (sinters at 123°). Salts could not be obt. crystalline. Cooccurs with Anagyryne, A-970 and Lupanine, L-302.

Marion, L. *et al.*, *J.A.C.S.*, 1946, **68**, 759-760

Lupinus Alkaloid P2 A-578

C₁₁H₁₈N₂O 194.276

Tentative mol. formula. Lupine alkaloid. Struct. unknown. Alkaloid from *Baptisia australis* and *Baptisia perfoliata* (Fabaceae). Cryst. (MeOH/Et₂O). Mp 300°.

Perchlorate:

Needles (MeOH). Mp 198°.

Picrate(?): Mp 241° (sinters at 185°).

Marion, L. *et al.*, *J.A.C.S.*, 1948, **70**, 691-692; 3253-3254 (*isol*)

Mucuna pruriens Alkaloid P A-579

Mucuna pruriens Base P

C₁₇H₂₆N₂O₆ 340.395

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Bp 118-119°.

Picrate: Mp 100-101°.

Rakhit, S. *et al.*, *Indian J. Pharm.*, 1956, **18**, 285; *CA*, **52**, 5748e (*isol*)

Alkaloid PAR 1 A-580

Struct. unknown. Mol. formula not recorded. Minor alkaloid from *Papaver argemone* (Papaveraceae). Prisms (MeOH). Mp 193-195°.

Táborská, E. *et al.*, *Coll. Czech. Chem. Comm.*, 1988, **53**, 1845 (*isol, uv*)

Alkaloid PAR 2 A-581

Struct. unknown. Mol. formula not

recorded. Minor alkaloid from *Papaver argemone* (Papaveraceae). Prisms (MeOH). Mp 252-256°.

Táborská, E. et al., *Coll. Czech. Chem. Comm.*, 1988, **53**, 1845 (*isol, uv*)

Alkaloid PAR 3 A-582


Struct. unknown. Quaternary alkaloid. Mol. formula and counterion not specified. Minor alkaloid from *Papaver argemone* (Papaveraceae). Amorph.

Táborská, E. et al., *Coll. Czech. Chem. Comm.*, 1988, **53**, 1845 (*isol, uv*)

Alkaloid PO3 A-583

1-Hydroxy-2,11-dimethoxy-6-methyl-7H-dibenzo[de,g]quinolinium hydroxide inner salt, 9CI

[27699-47-4]
[101064-66-8]



C₁₉H₁₅NO₄ 321.332

Alkaloid from *Papaver orientale* (Papaveraceae). Amorph.

Perchlorate: Mp 253-255°.

Preininger, V. et al., *CA*, 1967, **67**, 54290w (*isol*)


Preininger, V. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1969, **302**, 808 (*uv, ir, pmr*)

Saá, C. et al., *Tet. Lett.*, 1985, **26**, 4559 (*synth*)

Atanes, N. et al., *J.O.C.*, 1991, **56**, 2984 (*synth*)

Alkaloid PO4 A-584

Dehydroorientalidine
[23943-92-2]



C₂₂H₂₀NO₆[⊕] 394.403

Alkaloid from *Papaver orientale*, *Papaver oreophilum*, *Papaver nudicaule* var. *leio-carpum*, *Papaver pyrenaicum* ssp. *rhoeatiticum* and *Papaver alboroseum* (Papaveraceae).

Preininger, V. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 124 (*uv, pmr, struct, bibl*)

Alkaloid PP1† A-585

C₁₉H₁₅NO₅ 337.331

Struct. unknown. Quaternary phenolic protoberberine. Minor alkaloid from

Papaver pavoninum (Papaveraceae). Orange needles (MeOH) (as iodide). Mp 250° dec. (iodide). λ_{max} 222 ; 239 ; 251 ; 272 ; 352 (sh) ; 360 ; 468 (MeOH).

Táborská, E. et al., *Coll. Czech. Chem. Comm.*, 1988, **53**, 1845 (*isol, uv, ir, ms*)

Alkaloid PP1†, 9CI A-586

[158827-53-3]

C₁₇H₁₉NO₃ 285.342

Struct. unknown, prob. a normorphinane. Alkaloid from *Papaver pinnatifidum* (Papaveraceae). Mp 179°.

Laevorotatory.

Slavik, J. et al., *Coll. Czech. Chem. Comm.*, 1994, **59**, 1879-1883 (*isol*)

C-Alkaloid Q A-587

C₂₂H₂₇N₃O₃ 381.474

Minimum formula. Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 276-283° dec. No col. with Ce(SO₄)₂.

Meyer, H. et al., *Helv. Chim. Acta*, 1956, **39**, 1208-1213 (*isol, uv*)

Mucuna pruriens Alkaloid Q A-588

Mucuna pruriens Base Q

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Bp 220-221°.

Picrate: Mp 152-154°.

Rakhit, S. et al., *Indian J. Pharm.*, 1956, **18**, 285; *CA*, **52**, 5748e (*isol*)

Veratrum Alkaloid Q A-589

C₃₂H₅₃NO₄ 515.775

Steroidal alkaloid. Struct. unknown. Alkaloid from *Veratrum californicum* (Liliaceae). Mp 209-210°. [α]_D²⁵ -95 (CHCl₃).

Keeler, R.F. et al., *Phytochemistry*, 1968, **303** (*isol*)

C-Alkaloid R A-590

C₂₁H₂₇N₂O₂[⊕] 339.456

Minimum formula. Struct. unknown. Quaternary alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Violet col. with Ce(SO₄)₂. UV suggests indoline chromophore resembling C-Alkaloid B, A-400, C-Alkaloid C, A-436 and C-Alkaloid D, A-476.

Chloride: Mp 312° dec.

Perchlorate: Mp 317° dec.

Meyer, H. et al., *Helv. Chim. Acta*, 1956, **39**, 1208-1213 (*isol, uv*)

Fritillaria imperialis A-591

Alkaloid R

Fritillaria Base R

Prob. a steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria imperialis* var. *lutea* (Liliaceae). Mp 134-136° Mp 213-215° (double Mp). Co-occurs with Imperonine, I-53.

Hydroiodide: Mp 205-207°.

Perchlorate: Mp 193-196°.


Methiodide: Mp 269° dec.

Paul, L. et al., *Chem. Ber.*, 1958, **91**, 1968

Lycopodium fawcettii A-592

Alkaloid R

Lycopodium Base R
[53938-16-2]



Absolute Configuration

C₁₆H₂₄N₂O₂ 276.378

Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Mp 129-130°. [α]_D +104.

Perchlorate: Mp 198-199°.

Burnell, R.H. et al., *Can. J. Chem.*, 1963, **41**, 3091 (*isol, ir*)

Burnell, R.H. et al., *Chem. Comm.*, 1974, **391** (*cryst struct*)

Mucuna pruriens Alkaloid R A-593

Mucuna pruriens Base R

C₂₃H₃₅NO₄ 389.534

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Bp 320°.

Oxalate: Mp 130-131° dec.


Picrate: Mp 172-173°.

Methiodide: Mp 95-96°.

Rakhit, S. et al., *Indian J. Pharm.*, 1956, **18**, 285; *CA*, **52**, 5748e (*isol*)

Alkaloid RMB 10 A-594

[71635-30-8]



C₂₀H₂₆N₂O₂ 326.438

Alkaloid from the stem bark of *Rauwolfia mombasiana* (Apocynaceae). Yellow amorph. powder.

Iwu, M.M. et al., *Planta Med.*, 1979, **36**, 208 (*isol, uv, ir, ms, struct*)

Voacanga schweinfurthii A-595

Alkaloid S₂

Struct. unknown. Bisindole alkaloid (*Aspidosperma-Aspidosperma* type).

Alkaloid from the leaves of *Voacanga schweinfurthii* (Apocynaceae). Mol. formula not reported.

Newcombe, F. et al., *Planta Med.*, 1969, **17**, 276-280 (*isol, uv*)

Cordell, G.A. et al., *Alkaloids (Academic Press)*, 1981, **20**, 1; 266 (*rev*)

Voacanga schweinfurthii A-596

Alkaloid S₃

Struct. unknown. Bisindole alkaloid (*Aspidosperma-Aspidosperma* type).

Alkaloid from the leaves of *Voacanga schweinfurthii* (Apocynaceae). Mol. formula not reported.

Newcombe, F. *et al.*, *Planta Med.*, 1969, **17**, 276-280 (*isol*, *uv*)
Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 266 (*rev*)

C-Alkaloid S A-597C₁₉H₂₂N₂ 278.396

Approx. minimum formula. Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 250° (as picrate). Red-violet col. with Ce(SO₄)₂ becoming colourless on standing. Tertiary base.

Meyer, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1208-1213 (*isol*, *uv*)

Mucuna pruriens Alkaloid S A-598

Mucuna pruriens Base S

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Bp 320°.


Picrate: Mp 165-166°.

Rakhit, S. *et al.*, *Indian J. Pharm.*, 1956, **18**, 285; *CA*, **52**, 5748e (*isol*)

Alkaloid S-F A-599C₁₅H₂₁NO₃ 263.336

Pyrrolizidine alkaloid. Struct. unknown. Alkaloid from *Senecio viscosus* dried plants (Asteraceae). Mp 228-230° dec. [α]_D²² -148 (c, 0.573 in CHCl₃).

Šantavý, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 1666-1671

Alkaloid TC-D A-600R = C₂H₆NO₂C₂₃H₃₁N₃O₄ 413.516

Similar to Pagisulfine, P-11. Minor alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae).

van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400-423 (*isol*, *uv*, *pmr*, *cmr*, *ms*)

Saxton, J.E. *et al.*, *Nat. Prod. Rep.*, 1987, **4**, 591 (*rev*, *struct*)

Veratrum Alkaloid U A-601C₃₆H₅₅NO₁₃ 709.829


Steroidal alkaloid, poss. an ester of Protoverine. Struct. unknown. Alkaloid from *Veratrum lobelianum* (Liliaceae). Mp 181-183°. [α]_D +7.6 (CHCl₃).

Bondarenko, N.V. *et al.*, *Zh. Obshch. Khim.*, 1967, **37**, 332; *J. Gen. Chem. USSR (Engl. Transl.)*, 1967, **37**, 310 (*isol*)

C-Alkaloid UB A-602C₁₉H₂₃N₂O₃⁺ 327.402

Minimum formula. Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 238-240° dec. (as picrate). Carmine col. with Ce(SO₄)₂, becoming brown on standing.

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1947, **30**, 2081-2091; 1952, **35**, 1846-1879 (*isol*, *uv*)

Alkaloid Us 7 A-603Us 7
[188790-08-1]C₂₂H₂₆N₂O₆ 414.457

Alkaloid from *Uncaria attenuata*. Pale yellow oil. [α]_D²⁵ -12.9 (c, 1.10 in MeOH). λ_{max} 208 ; 244 (MeOH).

3-Epimer: Alkaloid Us 8. Us 8

[188790-09-2]

C₂₂H₂₆N₂O₆ 414.457

From *Uncaria attenuata*. Yellow oil. [α]_D²⁵ -9.6 (c, 1.25 in MeOH). λ_{max} 208 ; 244 (MeOH).

Aimi, N. *et al.*, *J.C.S. Perkin 1*, 1997, 187 (*isol*, *uv*, *pmr*, *cmr*, *struct*)

Gentiana Alkaloid VI A-604C₁₆H₁₈N₂O₃ 286.33

Struct. unknown. Monoterpene alkaloid. Alkaloid from *Gentiana asclepiadea*, *Gentiana bulgarica* and *Gentiana cruciata* (Gentianaceae). Mp 248-252°.

Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1965, **18**, 947-949; *CA*, **64**, 10084h (*isol*, *ir*)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (*rev*)

Lycodium annotinum A-605

Alkaloid VI

Lycodium Base VI

C₁₆H₂₃NO₂ 261.363

Struct. unknown. Alkaloid from *Lycodium annotinum* (Lycopodiaceae).

Methodide: Mp 294°. [α]_D²⁰ -13.9.

Methochloride: Mp 263°.

Methoperchlorate: Mp 287.5°.

Methopicrate: Mp 163°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol*, *uv*)

Lycodium annotinum A-606

Alkaloid VII

Lycodium Base VII

C₂₀H₂₉NO₄ 347.453

Struct. unknown. Alkaloid from *Lycodium annotinum* (Lycopodiaceae).

Methodide: Mp 292°. [α]_D²⁰ -11.1.

Methochloride: Mp 261°.

Methoperchlorate: Mp 295°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol*, *uv*)

Pachysandra terminalis A-607

Alkaloid VI

Pachysandra Base VI

Steroidal alkaloid. Struct. unknown. Isol. from *Pachysandra terminalis* (Buxaceae). Mp 290-295°.

Kikuchi, T. *et al.*, *Tet. Lett.*, 1964, 1817

Kikuchi, T. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 215; *CA*, **67**, 32888v

Rhodophiala bifida A-608

Alkaloid VI

Rhodophiala Base VI

C₁₆H₁₇NO₄ 287.315

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Rhodophiala bifida* (Amaryllidaceae). Mp 272-273°. [α]_D²⁵ -74 (MeOH).

Di-O-Ac: Mp 163-165°.

Dihydro: Mp 271°.

Wildman, W.C. *et al.*, *Pharmazie*, 1967, **22**, 725; *CA*, **69**, 19337m (*isol*)

Lycodium annotinum A-609

Alkaloid VIII

Lycodium Base VIII

C₁₆H₂₁NO₃ 275.347

Struct. unknown. Alkaloid from *Lycodium annotinum* (Lycopodiaceae).

Methodide: Mp 216-217°.

Methochloride: Mp 255°.

Methoperchlorate: Mp 234-236°.

Methopicrate: Mp 134-136°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol*, *uv*)

Lycodium annotinum A-610

Alkaloid XI

Lycodium Base XI

C₁₈H₂₅NO₃ 303.4

Struct. unknown. Alkaloid from *Lycodium annotinum* (Lycopodiaceae).

Methodide: Mp 272°. [α]_D²⁰ -95.8.

Methochloride: Mp 250°.

Methoperchlorate: Mp 267-268°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol*, *uv*)

Lycodium annotinum A-611

Alkaloid XII

Lycodium Base XII

C₁₈H₂₅NO₄ 319.4

Struct. unknown. Alkaloid from *Lycodium annotinum* (Lycopodiaceae).

Methodide: Mp 283°. [α]_D²⁰ -168.8.

Methochloride: Mp 270°.

Methyl perchlorate: Mp 259-260°.


Methopicrate: Mp 219°.

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1958, **32**, 485 (*isol*, *uv*)

Pachysandra terminalis A-612

Alkaloid XI

20-(Dimethylamino)-2',3,3',4-tetrahydro-3'-methylpregn-3-eno[3,4-d]oxazole, 8CI [15027-86-8]



C₂₅H₄₄N₂O 388.635Alkaloid from *Pachysandra terminalis* (Buxaceae) after hydrol. Mp 201-202°. [α]_D²⁰ -60 (CHCl₃). Could be an artifact.Kikuchi, T. *et al.*, *Tet. Lett.*, 1964, 1817-1823 (isol)**Rhodophiala bifida Alkaloid XIII** A-613

Rhodophiala Base XIII

C₁₆H₁₇NO₄ 287.315Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Rhodophiala bifida* (Amaryllidaceae). Mp 248-250°. [α]_D²¹ +12 (MeOH).Wildman, W.C. *et al.*, *Pharmazie*, 1967, 22, 725; *CA*, 69, 19337m (isol)**Pachysandra terminalis Alkaloid XVI** A-614


Pachysandra Base XVI

Steroidal alkaloid. Struct. unknown. Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 272-276°.Kikuchi, T. *et al.*, *Tet. Lett.*, 1964, 1817
Kikuchi, T. *et al.*, *Yakugaku Zasshi*, 1967, 87, 215; *CA*, 67, 32888v**Rhodophiala bifida Alkaloid XXV** A-615

Rhodophiala Base XXV




C₂₀H₂₃NO₅ 357.405Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Rhodophiala bifida* (Amaryllidaceae). Mp 204-206°. [α]_D²¹ -65 (MeOH).Wildman, W.C. *et al.*, *Pharmazie*, 1967, 22, 725; *CA*, 69, 19337m (isol)**Pachysandra terminalis Alkaloid XXVII** A-616

Pachysandra Base XXVII

Steroidal alkaloid. Struct. unknown. Alkaloid from *Pachysandra terminalis* (Buxaceae).Kikuchi, T. *et al.*, *Yakugaku Zasshi*, 1967, 87, 215; *CA*, 67, 32888v**C-Alkaloid Y** A-617*C-Proflurocurine*, 2,7-Dihydroxymavacurine [6822-67-9]C₂₀H₂₇N₂O₃⁺ 343.445Quaternary alkaloid from *Strychnos guianensis*, *Strychnos toxifera* and calabash curare (*Strychnos* spp.) (Loganiaceae). Cryst. (as chloride). Red-violet colour with Ce(IV) becoming olive-green on standing.

17-Carboxylic acid, N-de-Me, Me ester:

2,7-Dihydro-2,7-dihydroxypleiocarpamine. 2,7-Dihydroxypleiocarpamine [87614-66-2]

C₂₀H₂₄N₂O₄ 356.421Alkaloid from *Alstonia plumosa* (Apocynaceae). Cryst. (Me₂CO). Mp 152-158°. [α]_D²⁰ +66 (c, 0.5 in MeOH). Called 2,7-Dihydroxypleiocarpamine in the lit., which is not strictly correct. No stereochem. assigned. CAS no. not found 8-14 CI.Asmis, H. *et al.*, *Helv. Chim. Acta*, 1954, 37, 1968-1973 (isol, uv)Fritz, H. *et al.*, *Annalen*, 1958, 611, 268-276 (uv)Hesse, M. *et al.*, *Helv. Chim. Acta*, 1964, 47, 878-911 (struct)Jacquier, M.J. *et al.*, *Phytochemistry*, 1982, 21, 2973 (2,7-Dihydroxypleiocarpamine)Penelle, J. *et al.*, *Phytochemistry*, 2000, 53, 1057-1066 (isol, ir)**Alkaloid YC2** A-618Struct. unknown. Alkaloid from *Peganum harmala* (Zygophyllaceae). Characterised mainly by colour reactions and chem. props.McKenzie, E. *et al.*, *Phytochemistry*, 1975, 14, 273-275 (isol, uv)**2-(3-Alkyl-5-chloro-2,6-dihydroxybenzoyl)-4,5-dichloro-1H-pyrroles** A-619**2-(3-Chloro-5-hexyl-2,6-dihydroxybenzoyl)-4,5-dichloro-1H-pyrrole** [188988-62-7]C₁₇H₁₈Cl₃NO₃ 390.692Prod. by *Streptomyces* sp. PA48424. GABA receptor antagonist, antibacterial agent. Yellow cryst. Sol. CHCl₃, EtOAc, MeOH; poorly sol. H₂O. Mp 145-147°. λ_{\max} 225; 310 (€ 10000); 358 (€ 10700) (MeOH) (Berdy). λ_{\max} 260 (€ 5200); 310 (€ 15000); 340 (MeOH/HCl) (Berdy). λ_{\max} 233; 280 (€ 6400); 369 (€ 21300) (MeOH/NaOH) (Berdy).**2-[3-Chloro-2,6-dihydroxy-5-(4-methylpentyl)benzoyl]-4,5-dichloro-1H-pyrrole** [188988-63-8]C₁₇H₁₈Cl₃NO₃ 390.692Prod. by *Streptomyces* sp. PA48424. GABA receptor antagonist, antibacterial agent. Yellow cryst. Sol. CHCl₃, EtOAc, MeOH; poorly sol. H₂O. λ_{\max} 255; 310 (€ 10400); 358 (€ 10300) (MeOH) (Berdy). λ_{\max} 260 (€ 5300); 310 (€ 14900); 340 (MeOH/HCl) (Berdy).**2-[3-Chloro-2,6-dihydroxy-3-(4-methylhexyl)benzoyl]-4,5-dichloro-1H-pyrrole** [189008-36-4]C₁₈H₂₀Cl₃NO₃ 404.719Prod. by *Streptomyces* sp. PA48424. GABA receptor antagonist, antibacterial agent. Yellow cryst. Sol. CHCl₃, MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 225; 310 (€9300); 358 (€ 10000) (MeOH) (Berdy). λ_{\max} 260 (€ 4800); 310 (€ 14100) (MeOH/HCl) (Berdy). λ_{\max} 279 (€ 6100); 370 (€ 16000) (MeOH/NaOH) (Berdy).**2-[3-Chloro-2,6-dihydroxy-5-(5-methylhexyl)benzoyl]-4,5-dichloro-1H-pyrrole** [188988-64-9]C₁₈H₂₀Cl₃NO₃ 404.719Prod. by *Streptomyces* sp. PA48424. GABA receptor antagonist, antibacterial agent. Yellow cryst. Sol. CHCl₃, MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 225; 310 (€ 9800); 357 (€ 4800) (MeOH) (Berdy). λ_{\max} 258 (€ 4900); 310 (€ 14200) (MeOH/HCl) (Berdy). λ_{\max} 233; 281 (€ 6100); 371 (€ 19700) (MeOH/NaOH) (Berdy).Japan. Pat., 1997, 97 59 249; *CA*, 126, 276425v**2-Alkyl-3,4-dihydroxy-5-hydroxymethylpyridines** A-620
2-Alkyl-5-hydroxymethyl-3,4-pyridinediols**3,4-Dihydroxy-5-hydroxymethyl-2-propylpyridine**
5-Hydroxymethyl-2-propyl-3,4-pyridinediolC₉H₁₃NO₃ 183.207Prod. by *Streptomyces* sp. GW23/1540.**3,4-Dihydroxy-5-hydroxymethyl-2-isopropylpyridine**
5-Hydroxymethyl-2-(1-methylethyl)-3,4-pyridinediolC₉H₁₃NO₃ 183.207Prod. by *Streptomyces* sp. GW23/1540.**3,4-Dihydroxy-5-hydroxymethyl-2-(1-methylpropyl)pyridine**
5-Hydroxymethyl-2-(1-methylpropyl)-3,4-pyridinediol. 2-sec-Butyl-3,4-dihydroxy-5-hydroxymethylpyridineC₁₀H₁₅NO₃ 197.233Prod. by *Streptomyces* sp. GW63/1571.**3,4-Dihydroxy-5-hydroxymethyl-2-(2-methylpropyl)pyridine**
5-Hydroxymethyl-2-(2-methylpropyl)-3,4-pyridinediol. 3,4-Dihydroxy-5-hydroxymethyl-2-isobutylpyridineC₁₀H₁₅NO₃ 197.233Prod. by *Streptomyces* sp. GW23/1540.Maskey, R.P. *et al.*, *Z. Naturforsch., B*, 2005, 60, 63-66 (isol, pmr, cmr, ms)**2-Alkyl-3,4-dihydroxyquinolines** A-621
2-Alkyl-3-hydroxy-4(1H)-quinolinones. 2-Alkyl-3,4-quinolinediols

Metabs. of *Pseudomonas aeruginosa*. Signalling compds. involved in cell-to-cell signalling.

3,4-Dihydroxy-2-pentylquinoline

2-Pentyl-3,4-quinolinediol, 9CI. 3-Hydroxy-2-pentyl-4(1H)-quinolinone
[521313-34-8]
C₁₄H₁₇N₂O₂ 231.294

2-Hexyl-3,4-dihydroxyquinoline

2-Hexyl-3,4-quinolinediol, 2-Hexyl-3-hydroxy-4(1H)-quinolinone
[678172-89-9]
C₁₅H₁₉N₂O₂ 245.321

2-Heptyl-3,4-dihydroxyquinoline

2-Heptyl-3,4-quinolinediol, 9CI. 2-Heptyl-3-hydroxy-4(1H)-quinolinone. PQS
[521313-35-9]
C₁₆H₂₁N₂O₂ 259.347

3,4-Dihydroxy-2-octylquinoline

2-Octyl-3,4-quinolinediol. 3-Hydroxy-2-octyl-4(1H)-quinolinone
[678172-90-2]
C₁₇H₂₃N₂O₂ 273.374

3,4-Dihydroxy-2-nonylquinoline

2-Nonyl-3,4-quinolinediol, 9CI. 3-Hydroxy-2-nonyl-4(1H)-quinolinone
[521313-36-0]
C₁₈H₂₅N₂O₂ 287.401

3,4-Dihydroxy-2-undecylquinoline

2-Undecyl-3,4-quinolinediol, 9CI. 3-Hydroxy-2-undecyl-4(1H)-quinolinone
[678172-91-3]
C₂₀H₂₉N₂O₂ 315.455

Takeda, R. *et al.*, *CA*, 1960, **54**, 16533i (*isol*)
Pesci, E.C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1999, **96**, 11229-11234 (*heptyl, isol, synth, pmr, cmr*)

Lepine, F. *et al.*, *Biochim. Biophys. Acta*, 2003, **1622**, 36-41 (*detm*)


Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*occur, biosynth, ms*)

Bredenbruch, F. *et al.*, *J. Bacteriol.*, 2005, **187**, 3630-3635 (*biosynth*)

3-(2-Alkyl-5-oxazolyl)-1H-indoles

A-622

2-Alkyl-5-(1H-indol-3-yl)oxazole



Revised structs. (1999) for several members of this series which were formerly descr. as pyrazoloisoquinolines.

3-(2-Methyl-5-oxazolyl)-1H-indole, 9CI

5-(3-Indolyl)-2-methyloxazole. *Pimprinine*. WS 30581C. Antibiotic WS 30581C. APHE 3
[13640-26-1]
[179728-23-5]
C₁₂H₁₀N₂O 198.224

Metab. of *Streptomyces pimprina*, *Streptovorticillium griseocarneum* and *Streptovorticillium olivoreticuli*. Monoamine oxidase inhibitor. Shows antiepileptic effects. Prisms (C₆H₆), needles. Mp 204-205°. λ_{max} 224 (ε 24600); 266 (ε 15500); 282 (sh) (ε 12100); 298 (sh) (ε 11000) (EtOH) (Derep).

3-(2-Ethyl-5-oxazolyl)-1H-indole, 9CI

2-Ethyl-5-(3-indolyl)oxazole. *Pimprinethine*. APHE 1
[73053-81-3]
[146426-35-9]
C₁₃H₁₂N₂O 212.251
Metab. of *Streptovorticillium olivoreticuli*, *Streptovorticillium griseocarneum* and *Streptomyces cinnamomeus*. Cryst. (C₆H₆ or CHCl₃). Mp 163-165° (161-163°).

3-(2-Propyl-5-oxazolyl)-1H-indole

5-(3-Indolyl)-2-propyloxazole. *Antibiotic WS 30581A*. APHE 2. WS 30581A
[93773-64-9]
[146426-36-0]
C₁₄H₁₄N₂O 226.277
Prod. by *Streptovorticillium waksmanii* and *Streptovorticillium griseocarneum*. Antiplatelet, cytotoxic and antithrombotic agent. Needles. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 128-130°. λ_{max} 224 (ε 24000); 266 (ε 15400); 282 (sh) (ε 13000); 298 (sh) (ε 11200) (EtOH) (Derep).
▶ LD₅₀ (mus, ipr) 250-750 mg/kg.

3-(2-Butyl-5-oxazolyl)-1H-indole, 9CI

2-Butyl-5-(3-indolyl)oxazole. *Antibiotic WS 30581B*. APHE 4. WS 30581B
[93773-63-8]
[179728-24-6]
C₁₅H₁₆N₂O 240.304
Prod. by *Streptovorticillium waksmanii* and *Streptovorticillium griseocarneum*. Platelet aggregation inhibitor. An antithrombotic agent. Needles (C₆H₆/hexane). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 123-125°. λ_{max} 224 (ε 24000); 266 (ε 15400); 282 (sh) (ε 13000); 298 (sh) (ε 11200) (EtOH) (Derep).

3-[2-(2-Methylpropyl)-5-oxazolyl]-1H-indole

5-(3-Indolyl)-2-isobutyloxazole. *Labradorin 1*
[477951-20-5]
C₁₅H₁₆N₂O 240.304
Prod. by *Pseudomonas syringae* pv. *coronafaciens*. Cytotoxic. Prisms (Me₂CO). Mp 147-148°.

3-(2-Pentyl-5-oxazolyl)-1H-indole

5-(3-Indolyl)-2-pentyloxazole. *Labradorin 2*
[326494-56-8]
C₁₆H₁₈N₂O 254.331
Prod. by *Pseudomonas syringae* pv. *coronafaciens*. Cytotoxic. Small plates (MeOH). Mp 130-132°.

3-(2-Benzyl-5-oxazolyl)-1H-indole, 9CI

3-[2-(Phenylmethyl)-5-oxazolyl]-1H-indole. 2-Benzyl-5-(3-indolyl)oxazole. *Pimprinaphine*
[73053-85-7]

C₁₈H₁₄N₂O 274.321

Metab. of *Streptovorticillium olivoreticuli*. Cryst. (C₆H₆). Mp 200-201° (198-200°).

Bhate, D.S. *et al.*, *Experientia*, 1960, **16**, 504 (*Pimprinine, isol*)

Joshi, B.S. *et al.*, *Tetrahedron*, 1963, **19**, 1437-1439 (*Pimprinine, uv, ir, struct, synth*)

Oikawa, Y. *et al.*, *Heterocycles*, 1979, **12**, 1457-1462 (*synth*)

Koyama, Y. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 1285-1287 (*Pimprinine, Pimprinethine, Pimprinaphine, isol, uv, pmr, synth*)

Yoshioka, T. *et al.*, *J. Chem. Res., Synop.*, 1981, 194-195; *J. Chem. Res., Miniprint*, 2252-2281 (*synth, uv, ir, pmr, ms*)

Noltemeyer, M. *et al.*, *J. Antibiot.*, 1982, **35**, 549-555 (*Pimprinethine, isol, uv, ir, pmr, cmr, ms, cryst struct*)

Umehara, K. *et al.*, *J. Antibiot.*, 1984, **37**, 1153-1160 (*WS 30581*)

Somei, M. *et al.*, *Heterocycles*, 1985, **23**, 1101-1106 (*synth*)

Doyle, K.J. *et al.*, *Synthesis*, 1994, 1021-1022 (*synth, ir, pmr, cmr*)

Kelly, T.R. *et al.*, *Tet. Lett.*, 1999, **40**, 1857-1860 (*APHE 1-4*)


Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1793-1797 (*Labradorins, cryst struct*)

Kumar, D. *et al.*, *Tet. Lett.*, 2008, **49**, 867-869 (*synth*)

3-Alkyl-1H-pyrrole-2-carboxaldehydes

A-623

3-Alkyl-2-formylpyrroles



Isol. as a mixt. of 3.3% n = 18, 12.5% n = 19, 49% n = 20, 25% n = 21, and 10.2% n = 22. The substitution pattern of these constds. has been questioned (Stierle *et al.*). Metabs. of the marine sponge *Oscarella lobularis*.

3-Nonadecyl-1H-pyrrole-2-carboxaldehyde

[57992-51-5]
C₂₄H₄₃NO 361.61
n = 18.

3-Eicosyl-1H-pyrrole-2-carboxaldehyde

3-Icosyl-1H-pyrrole-2-carboxaldehyde
[57992-52-6]
C₂₅H₄₅NO 375.637
n = 19.

3-Heneicosyl-1H-pyrrole-2-carboxaldehyde

[57992-53-7]
C₂₆H₄₇NO 389.663
n = 20.

12',13'-Didehydro(E)-: 13-(12-Heneicosenyl)-1H-pyrrole-2-carboxaldehyde, 9CI
[57992-57-1]
C₂₆H₄₅NO 387.648
Isol. from the marine sponge *Oscarella lobularis*.


3-Docosyl-1H-pyrrole-2-carboxaldehyde

[57992-54-8]
C₂₇H₄₉NO 403.69
n = 21.

3-Tricosyl-1*H*-pyrrole-2-carboxaldehyde
[57992-55-9]C₂₈H₅₁NO 417.717
n = 22.Cimino, G. et al., *Experientia*, 1975, **31**, 1387-1389 (isol, uv, ir, pmr, ms, struct)
Stierle, D.B. et al., *J.O.C.*, 1980, **45**, 4980-4982 (struct)**5-Alkyl-1*H*-pyrrole-2-carboxaldehydes**

A-624

2-Alkyl-5-formylpyrroles

**5-Nonyl-1*H*-pyrrole-2-carboxaldehyde**
[89631-84-5]C₁₄H₂₃NO 221.342Isol. from a soft coral, *Telesto* sp., and its associated, but unidentified, demospunge. Yellow oil. Not possible to identify the exact source of the natural product since it perfused both organisms.**5-Pentadecyl-1*H*-pyrrole-2-carboxaldehyde**
[75233-97-5]C₂₀H₃₅NO 305.503Metab. from a marine sponge *Laxosuberites* sp. Isol. in admixture (46%) with the hexadecyl (12%), heptadecyl (23%) and nonadecyl (19%) homologues. λ_{max} 297 (ε 16000) (MeCN) (Derep).**6',7'E-Didehydro: 5-(6-Pentadecenyl)-1*H*-pyrrole-2-carboxaldehyde**C₂₀H₃₃NO 303.487Metab. of the sponge *Mycale tenuispiculata*. Pale yellow solid. Mp 42-44°. λ_{max} 297 (ε 15400) (MeCN).**5-(13-Methyltetradecyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-63-3]C₂₀H₃₅NO 305.503Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-Hexadecyl-1*H*-pyrrole-2-carboxaldehyde**
[75233-98-6]C₂₁H₃₇NO 319.529Metab. from *Laxosuberites* sp. See note under pentadecyl above.**5-(14-Methylpentadecyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-64-4]C₂₁H₃₇NO 319.529Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-Heptadecyl-1*H*-pyrrole-2-carboxaldehyde**
[75233-99-7]C₂₂H₃₉NO 333.556Metab. from *Laxosuberites* sp. See note under pentadecyl above.**9,10Z-Didehydro: 5-(9Z-Heptadecenyl)-1*H*-pyrrole-2-carboxaldehyde**

[233744-73-5]

C₂₂H₃₇NO 331.54Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-Octadecyl-1*H*-pyrrole-2-carboxaldehyde**
[262351-87-1]C₂₃H₄₁NO 347.583Isol. from *Mycale mytilorum*. Pale yellow flakes. Mp 56-58°. λ_{max} 302 (ε 16000) (EtOH).**5-(16-Methylheptadecyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 6*

[705973-02-0]

C₂₃H₄₁NO 347.583Isol. from the sponge *Mycale cecilia*. Amorph. powder. λ_{max} 203 (ε 6260); 249 (ε 2080); 300 (ε 9790) (MeOH).**5-Nonadecyl-1*H*-pyrrole-2-carboxaldehyde**
[75234-00-3]C₂₄H₄₃NO 361.61Metab. from *Laxosuberites* sp. See note under pentadecyl above.**9,10Z-Didehydro: 5-(9Z-Nonadecenyl)-1*H*-pyrrole-2-carboxaldehyde**

[233744-74-6]

C₂₄H₄₁NO 359.594Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**12,13Z-Didehydro: 5-(12Z-Nonadecenyl)-1*H*-pyrrole-2-carboxaldehyde.***Mycalazal 4*

[705972-97-0]

C₂₄H₄₁NO 359.594Isol. from the sponge *Mycale cecilia*. Amorph. powder. λ_{max} 202 (ε 7585); 249 (ε 2340); 300 (ε 11760) (MeOH).**5-Eicosyl-1*H*-pyrrole-2-carboxaldehyde**
[233744-67-7]C₂₅H₄₅NO 375.637Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-(9Z-Heneicosenyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-65-5]C₂₆H₄₅NO 387.648Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-(14Z-Heneicosenyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 8*

[705973-25-7]

Isol. from the sponge *Mycale cecilia*.Amorph. powder. λ_{max} 202 (ε 7800); 248 (ε 2320); 300 (ε 11290) (MeOH).**5-(12Z,15Z-Heneicosadienyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 5*

[705972-98-1]

C₂₆H₄₃NO 385.632Isol. from the sponge *Mycale cecilia*. Oil. λ_{max} 204 (ε 9250); 248 (ε 2390); 300 (ε 11150) (MeOH).**5-(12Z,15Z,18Z-Heneicosatrienyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 3*

[705972-95-8]

C₂₆H₄₁NO 383.616Isol. from the sponge *Mycale cecilia*. Oil. λ_{max} 204 (ε 14690); 248 (ε 3440); 300 (ε 15400) (MeOH).**5-(12Z-Docosenyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-66-6]C₂₇H₄₇NO 401.674Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*. Indexed as the Δ¹¹-isomer in CAS.**5-Tricosyl-1*H*-pyrrole-2-carboxaldehyde**
[233744-70-2]C₂₈H₅₁NO 417.717Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**9,10Z-Didehydro: 5-(9Z-Tricosenyl)-1*H*-pyrrole-2-carboxaldehyde**

[233744-68-8]

C₂₈H₄₉NO 415.701Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**14,15Z-Didehydro: 5-(14Z-Tricosenyl)-1*H*-pyrrole-2-carboxaldehyde**

[233744-69-9]

C₂₈H₄₉NO 415.701Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**14,15Z,17,18Z-Tetradecydro: 5-(14Z,17Z-Tricosadienyl)-1*H*-pyrrole-2-carboxaldehyde.***Mycalazal 9*

[705973-27-9]

C₂₈H₄₇NO 413.685Isol. from the sponge *Mycale cecilia*. Oil. λ_{max} 201 (ε 14170); 248 (ε 4060); 300 (ε 21160) (MeOH).**14,15Z,17,18Z,20,21Z-Hexadecydro: 5-(14Z,17Z,20Z-Tricosatrienyl)-1*H*-pyrrole-2-carboxaldehyde.***Mycalazal 7*

[705973-15-5]

C₂₈H₄₅NO 411.67Isol. from the sponge *Mycale cecilia*. Oil. λ_{max} 204 (ε 9250); 248 (ε 2390); 300 (ε 11150) (MeOH).**5-(11Z-Pentacosenyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-71-3]C₃₀H₅₃NO 443.755Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-(15Z-Pentacosenyl)-1*H*-pyrrole-2-carboxaldehyde** [233744-72-4]C₃₀H₅₃NO 443.755Isol. from the sponges *Desmapsamma anchorata* and *Mycale microsigmatosa*.**5-(16Z-Pentacosenyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 12*

[705973-36-0]

C₃₀H₅₃NO 443.755Isol. from the sponge *Mycale cecilia*.**5-(18Z-Pentacosenyl)-1*H*-pyrrole-2-carboxaldehyde***Mycalazal 13*

[705973-38-2]

C₃₀H₅₃NO 443.755Isol. from the sponge *Mycale cecilia*.

5-(16Z,19Z-Pentacosadienyl)-1H-pyrrole-2-carboxaldehyde**Mycalazal 11**

[705973-35-9]

Isol. from the sponge *Mycale cecilia*. Oil. λ_{\max} 202 (ϵ 6040); 248 (ϵ 1800); 301 (ϵ 7630) (MeOH).

5-(16Z,19Z,22Z-Pentacosatrienyl)-1H-pyrrole-2-carboxaldehyde**Mycalazal 10**

[705973-28-0]

Isol. from the sponge *Mycale cecilia*. Oil. λ_{\max} 204 (ϵ 11920); 248 (ϵ 3690); 300 (ϵ 16900) (MeOH).

5-(10Z,13Z,16Z,19Z,22Z-Pentacosapentaenyl)-1H-pyrrole-2-carboxaldehyde, 9CI**Mycalazal 2**

[185389-83-7]

C₃₀H₄₅NO 435.692

Metab. from *Mycale micracanthoxea*. Cytotoxic agent. Oil.

5-(7Z,10Z,13Z,16Z,19Z,22Z-Pentacosahexaenyl)-1H-pyrrole-2-carboxaldehyde**Mycalazal 1**

[185389-71-3]

Metab. of the sponge *Mycale micracanthoxea*. Oil.

Stierle, D.B. *et al.*, *J.O.C.*, 1980, **45**, 4980 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Bowden, B.F. *et al.*, *Aust. J. Chem.*, 1984, **37**, 227 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *synth*)

Ortega, M.J. *et al.*, *Tetrahedron*, 1997, **53**, 331-340; 2004, **60**, 2517-2524 (*Mycalazals 1-13*)

Compagnone, R.S. *et al.*, *Nat. Prod. Lett.*, 1999, **13**, 203-211 (*alkenyls*, *branched-chain alkyls*)


Reddy, G.B.S. *et al.*, *Bioorg. Med. Chem.*, 2000, **8**, 27-36 (*octadecyl*)

Venkatesham, U. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1318-1320 (*6-pentadecenyl*)

Hansen, T.V. *et al.*, *Tet. Lett.*, 2004, **45**, 2809-2811 (*Mycalazal 2*, *synth*)

Allantoin, BAN, USAN A-625

(2,5-Dioxo-4-imidazolidinyl)urea, 9CI. 5-Ureido-2,4-imidazolidinedione. Glyoxylic diureide. 5-Ureidohydantoin. Dermalex. Cordianin. Actinac. Alphosyl



(R)-form

C₄H₆N₄O₃ 158.116**(±)-form [5377-33-3]**

Occurs in allantoinic fluid. A product of purine metabolism. V. widely distributed in biol. systems, *isol.* from numerous plants. *Isol.* from *Bambusa oldhamii* (green tomato), *Allophylus edulis* var *edulis*, *Allophylus edulis* var *gracilis*, *Strychnos cathayensis*, roots of *Pyrostegia venusta*, marine sponges *Cymbastela cantharella* and *Tedania digitata*, and cows milk. Formed in animals, except primates, by enzymic oxidn. of Uric acid, U-45. Antiinflammatory, anti-psoriatic (disputed), topical vulnerary

agent. Promotes wound healing. Monoclinic prisms (H₂O). V. spar. sol. H₂O; spar. sol. org. solvs. Mp 238-240° dec. Log P -1.54 (calc). Component of numerous preps. Actinac, Alphosyl, Dermalex.

▶YT1600000

[97-59-6 ((±)-form), 61504-48-1]

Org. Synth., Coll. Vol., 2, 1943, 21-23 (*synth*, *bibl*)

Karrer, W. *et al.*, *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 2569 (*occur*)

Mecca, S.B. *et al.*, *Soap. Perfum. Cosmet.*, 1976, **49**, 434; 481; 483-485 (*rev*)

Coxon, B. *et al.*, *J.O.C.*, 1977, **42**, 3132-3140 (*cmr*)

Mecca, S.B. *et al.*, *Cosmet. Toiletries*, 1978, **93**, 39-41 (*rev*)

Cook, A.F. *et al.*, *J.O.C.*, 1980, **45**, 4020-4025 (*isol*)

Chen, K.J. *et al.*, *Taiwan Kexue*, 1989, **42**, 21-24; *CA*, **113**, 55816 (*(±)-form*, *isol*)

Hoffmann-Bohm, K. *et al.*, *Planta Med.*, 1992, **58**, 544-548 (*(±)-form*, *isol*)

Mourabit, A.A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 290-291 (*(±)-form*, *isol*)


Shingfield, K.J. *et al.*, *J. Chromatogr., B*, 1998, **706**, 342-346 (*hplc*)

Ferreira, D.T. *et al.*, *Quim. Nova*, 2000, **23**, 42-46 (*isol*)

Alline**A-626**

2,3,8,8a-Tetrahydro-1-methylpyrrolo[2,3-b]indol-3a(1H)-ol, 9CI

[101053-34-3]



Relative Configuration

C₁₁H₁₄N₂O 190.244

Alkaloid from Chinese chives (*Allium odorum*) and several other *Allium* spp. Mp 91-92°. [α]_D +136.3 (c, 1.218 in HCl). λ_{\max} 245 (log ϵ 3.68); 303 (log ϵ 3.14) (no solvent reported).

Hydrochloride: Mp 196-197°.

Me ether, N⁸-Me: 2,3,8,8a-Tetrahydro-3a-methoxy-1,8-dimethylpyrrolo[2,3-b]indole. **Alkaloid CPC I**. **CPC 1**

C₁₃H₁₈N₂O 218.298

Alkaloid from *Chimonanthus praecox* f. *concolor*. Amorph. solid. [α]_D²⁵ -88 (c, 0.1 in MeOH). (3aR,8aR)-config. determined. λ_{\max} 205 (log ϵ 4.15); 251 (log ϵ 3.72); 301 (log ϵ 3.14) (MeOH).

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 687-691; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 645-649 (*isol*, *cryst struct*, *uv*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 383; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 362 (*occur*)


Antsupova, T.P. *et al.*, *Rastit. Resur.*, 1987, **23**, 436; *CA*, **107**, 214830x (*occur*)

Kitajima, M. *et al.*, *Tet. Lett.*, 2006, **47**, 3199-3202 (**CPCI**)

Allithiamine**A-627**

N-[(4-Amino-2-methyl-5-pyrimidinyl)-methyl]-N-[4-hydroxy-1-methyl-2-(2-propenylidithio)-1-butenyl]formamide, 9CI

[554-44-9]

C₁₅H₂₂N₄O₂S₂ 354.496


Naturally-occurring open-chain thiamine analogue. Obt. from garlic (*Allium sativum*). Imparts meaty flavour to foods. Has vitamin B₁ activity. Investigated as a dietary supplement to enhance muscle performance and as treatment for thiamine deficiency. Cryst. (C₆H₆). Mp 132-133° dec.

Matsukawa, T. *et al.*, *Science (Washington, D.C.)*, 1953, **118**, 325 (*isol*, *struct*)

Matsukawa, T. *et al.*, *CA*, 1955, **49**, 7572g (*synth*)

Baker, H. *et al.*, *J. Nutr. Sci. Vitaminol., Suppl.*, 1976, **22**, 63-68 (*pharmacol*)

Doyle, M.R. *et al.*, *Int. J. Sport Nutr.*, 1997, **7**, 39-47 (*use*)

Allomatrine**A-628**

(+) -form

C₁₅H₂₄N₂O 248.367

Stereoisomer of Matrine, M-121, Isomatrine, I-244, Sophoridine, S-380 and Darvasamine, D-79.

(+) -form [641-39-4]

Obt. by isomerisation of Matrine, M-121. Mp 103-105°. [α]_D²⁵ +77.9. *Incorr.* stated by Ibragimov *et al* (1979) to have been *isol.* by Ueno *et al* from *Sophora flavescens*. This was Isomatrine.

Methodide: Dec. at 315°.

Picrate: Mp 177°.

(-) -form**Leontine†. Isoleontine**

[6783-60-4]

Alkaloid from *Leontice eversmannii* and *Leontice darvasica* (Leonticeaceae). Mp 107-108°. Bp₃ 175-190°. [α]_D -78.2 (EtOH). The original isolate of Leontine from *L. eversmannii* (Yunusov, 1949) was opt. inactive. It may have been the (±)-form, or a different alkaloid. Platonova *et al* (1953) isolated the (-)-form, which they called Isoleontine, but later workers used the name Leontine for the (-)-form.

Perchlorate: Mp 252°.

Methodide: Mp 297°.

13-Oxo: LeontalbamineC₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Leontice alberti*. Mp 195°. [α]_D²⁰ -93. Struct. dubious. Given in the ref. as the unlikely 12,13-didehydro, 13-hydroxy enol tautomer. However struct. and MF do not agree in ref. CAS no. not found 8-14 CI. λ_{max} 220 (log ε 3.4) (no solvent reported).

(±)-form

Mp 85-87°.

Hydrochloride: Mp 215-217°.

Platonova, T.F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1953, **23**, 921-926 (*Leontine*)
Bohlmann, F. *et al.*, *Chem. Ber.*, 1958, **91**, 2176-2189 (*ir*)

Rulko, F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1961, **31**, 282-285 (*Leontine*)
Mandell, L. *et al.*, *J.A.C.S.*, 1965, **87**, 5234-5236 (*synth*)

Vul'fson, N.S. *et al.*, *Khim. Geterotsikl. Soedin.*, 1974, 251-260; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1974, **10**, 221-229 (*ir*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1975, **108**, 1043-1051 (*cmr*)


Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 416-417; *Chem. Nat. Compd. (Engl. Transl.)*, 368-369 (*cryst struct*)

Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 71-75; *Chem. Nat. Compd. (Engl. Transl.)*, 66-69 (*rev. stereochem*)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 615-681; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 596-675 (*Leontalbamine*)

Allopumiliotoxin 309D**A-629**

6-(2,5-Dimethyloctylidene)octahydro-8-methyl-7,8-indolizinediol, 9CI
[141643-33-6]


C₁₉H₃₅NO₂ 309.491

Alkaloid from skin extracts of the Panamanian poison frog *Dendrobates pumilio*.

Tokuyama, T. *et al.*, *Tetrahedron*, 1991, **47**, 5415 (*isol, cmr, ms, struct*)

Allosecurinine**A-630**

Phyllochryisine. 2-*Episecurinine*. *Viroallosecurinine*

**(-)-form**C₁₃H₁₅NO₂ 217.267

Viroallosecurinine is the (+)-enantiomer.

(-)-form [884-68-4]

Alkaloid from the leaves and roots of *Securinega suffruticosa*. Minor alkaloid from the roots of *Phyllanthus discoides*. Also isol. from bark and roots of *Margaritaria indica* (Euphorbiaceae). Yellow needles (Et₂O). Mp 136-138° (128°). [α]_D²⁶ -1082 (EtOH). [α]_D -1140 (CHCl₃).

Hydrochloride:

Cubes (EtOH). Mp 248° dec. [α]_D³⁰ -335 (EtOH).

Picrate:

Yellow needles (EtOH). Mp 231-232° dec.

14,15-Dihydro:

Needles (petrol). Mp 91°.

14,15-Dihydro, 15β-hydroxy: 14,15-Dihydroallosecurinine-15-ol

[119817-92-4]

C₁₃H₁₇NO₃ 235.282

Alkaloid from the leaves of *Phyllanthus discoides* (Euphorbiaceae). Orange-yellow cryst. (1-propanol). Mp 157-158°. [α]_D²⁵ -96 (c, 0.48 in EtOH).

14,15-Dihydro, 15α-methoxy: 14,15-Dihydro-15-methoxyphyllochryisine

15-Methoxy-14,15-dihydrophyllochryisine
[128855-48-1]

C₁₄H₁₉NO₃ 249.309

Alkaloid from bark and roots of *Margaritaria indica* (Euphorbiaceae). Needles (petrol). Mp 141°. [α]_D +71 (c, 0.6 in MeOH).

14,15-Dihydro, 15α-methoxy, hydrobromide:

Plates (MeOH/Me₂CO). Mp 182-183°.

(+)-form [1857-30-3]

Alkaloid from the leaves of *Securinega virosa* (Euphorbiaceae). Yellow prisms (Me₂CO). Mp 136-138°. [α]_D²⁰ +1084.6 (c, 0.05 in EtOH).

14,15-Dihydro, 15β-ethoxy: 15β-Ethoxy-14,15-dihydroviroallosecurinine

[936951-59-6]

C₁₅H₂₁NO₃ 263.336

Alkaloid from *Securinega suffruticosa*.

Parello, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 898-910 (*uv, ir, pmr*)

Nakano, T. *et al.*, *J.O.C.*, 1963, **28**, 2619-2621; 1964, **29**, 3441-3443 (*uv, ord, cd, conformn, abs config*)

Saito, S. *et al.*, *Chem. Ind. (London)*, 1964, 1263-1264 (*struct, Viroallosecurinine*)

Bevan, C.W.L. *et al.*, *Chem. Ind. (London)*, 1964, 2054 (*isol*)

Chatterjee, A. *et al.*, *J. Indian Chem. Soc.*, 1964, **41**, 163-172 (*isol, ir, pmr, struct*)
Saito, S. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1126; *CA*, **62**, 5498d (*isol, Viroallosecurinine*)

Mukherjee, R. *et al.*, *Indian J. Chem.*, 1966, **4**, 459 (*isol, uv*)

Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 2009-2012 (*synth*)

Sankawa, U. *et al.*, *Tet. Lett.*, 1974, 1867-1868 (*biosynth*)

Beutler, J.A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 677-681 (*cmr*)

Mensah, J.L. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1113-1115 (*14,15-Dihydroallosecurinine-15β-ol*)

Arbain, D. *et al.*, *Aust. J. Chem.*, 1990, **43**, 439-445 (*15α-Methoxy-14,15-dihydrophyllochryisine, cryst struct, abs config*)

Honda, T. *et al.*, *Tet. Lett.*, 2004, **45**, 5211-5213 (*synth*)

Wang, Y. *et al.*, *Zhongguo Tianran Yaowu*, 2006, **4**, 260-263; *CA*, **146**, 518050b (*Ethoxydihydroviroallosecurinine*)

Yuan, W. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 234-242 (*isol, pmr, cmr*)

Bardaji, G.G. *et al.*, *J.O.C.*, 2008, **73**, 7657-7662 (*synth*)

Allothiobinupharidine**A-631**

Struct. unknown


C₃₀H₄₂N₂O₂S 494.74

Nuphar alkaloid. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Plates (as diperchlorate). Mp 320-325° (diperchlorate).

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1962, **36**, 1815-1825 (*isol*)

Alloxazine**A-632**

Benzo[g]pteridine-2,4(1H,3H)-dione, 9CI. 6,7-Benzolumazine
[490-59-5]

C₁₀H₆N₄O₂ 214.183Parent compd. exists in 1*H*,3*H*-form.

In the older lit., posns. 5,6,7,8,9,10 are often numbered 10,5,6,7,8,9 respectively. Greyish-green powder. V. spar. sol. EtOH; insol. H₂O, Et₂O. Mp 300° dec. The original synthesis of alloxazine (Kühling, 1891) prob. gave a material containing little of the desired product.

(3*H*,10*H*)-form

*Benzo[g]pteridine-2,4(3*H*,10*H*)-dione*. *Isoalloxazine*. *Flavin*
Parent residue present in Riboflavin. Many subst. derivs. synthesised. Same CAS no. as Alloxazine.

3,10-Di-Me: [4074-59-3]C₁₂H₁₀N₄O₂ 242.237

Orange-yellow needles (AcOH). Mp 320-325° dec. Bitter taste.

10-Ph: [6851-14-5]C₁₆H₁₀N₄O₂ 290.281

Orange-yellow prisms (AcOH). V. spar. sol. H₂O. Tasteless.

10-Ph, 3-Me: [35804-39-8]C₁₇H₁₂N₄O₂ 304.307

Orange-yellow cryst. (AcOH). Stable on heating to at least 360°. Tasteless.

10-Benzyl: [21708-43-0]C₁₇H₁₂N₄O₂ 304.307


Mp 273-276°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 894C (*ir*)

- Kuhn, R. *et al.*, *Ber.*, 1934, **67**, 1459; 1935, **68**, 1282 (*Isoalloxazine derivs*)
 Tishler, M. *et al.*, *J.A.C.S.*, 1945, **67**, 2165 (*synth*)
 Lambooy, J.P. *et al.*, *Heterocycl. Compd.*, (ed. Elderfield, R.C.) Wiley, New York, 1967, **9**, 118 (*rev*)
 Szymusiak, H. *et al.*, *J.C.S. Perkin 2*, 1990, 229 (*struct, props*)
 Villemain, D. *et al.*, *Synth. Commun.*, 1995, **25**, 2315 (*synth*)
 Kaupp, G. *et al.*, *Eur. J. Org. Chem.*, 2002, 1368-1373 (*synth, ir, pmr, cmr*)
 Chattopadhyay, P. *et al.*, *Synth. Commun.*, 2006, **36**, 1857-1861 (*10-benzyl*)

Almazole A

A-633

C₂₁H₂₁N₃O₃ 363.415**(S)-form** [157382-33-7]

Alkaloid from the red alga *Haraldio-phyllum* sp. Semisolid. $[\alpha]_D^{20} +103$ (c, 0.155 in MeOH). λ_{max} 204 (ϵ 36700); 237 (ϵ 26400); 278 (ϵ 18200); 320 (ϵ 6000) (MeOH) (Derep).

Deformyl: Almazole B [157382-34-8]

C₂₀H₂₁N₃O₂ 335.405
 Alkaloid from *Haraldio-phyllum* sp. Yellow semisolid. $[\alpha]_D^{20} +92$ (c, 0.05 in MeOH). λ_{max} 203 (ϵ 40300); 234 (ϵ 22600); 260 (ϵ 16500); 396 (ϵ 6300) (MeOH) (Derep).


N'Diaye, I. *et al.*, *Tet. Lett.*, 1994, **35**, 4827-4830 (*isol, uv, cd, pmr, cmr, ms*)

Khalafy, J. *et al.*, *Aust. J. Chem.*, 1999, **52**, 31-36 (*synth*)

Almazole C

A-634

[161068-69-5]

C₂₁H₂₁N₃O 331.416


Alkaloid from the red alga *Haraldio-phyllum* sp. CNS active. $[\alpha]_D^{20} +168$ (c, 1.08 in MeOH). λ_{max} 222 (ϵ 26000); 271 (ϵ 15100); 282 (ϵ 14500); 300 (ϵ 12900) (MeOH) (Berdy).

Guella, G. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1999-2006 (*isol, synth, cd, uv, pmr, cmr, ms*)

Almazole D

A-635

[176739-66-5]

C₂₂H₂₁N₃O₃ 375.426

Alkaloid from the red alga *Haraldio-phyllum* sp. Shows antifungal activity. Powder. $[\alpha]_D^{20} +20$ (c, 0.07 in MeOH).

Me ether

Powder. $[\alpha]_D^{20} +70$ (c, 0.1 in MeOH).


N'Diaye, I. *et al.*, *Tet. Lett.*, 1996, **37**, 3049-3050 (*isol, uv, pmr, cmr, ms*)

Almazolone

A-636

[878143-16-9]

[878143-15-8 (Z-form), 878143-13-6 (E-form)]

C₂₀H₁₆N₂O₂ 316.359


Isol. from the red alga *Haraldio-phyllum* sp. Yellow powder. Mp 245-247°. Isol. as an 80:20 mixt. of (Z/E)-isomers to which data refers.

Guella, G. *et al.*, *Tetrahedron*, 2006, **62**, 1165-1170 (*isol, synth, pmr, cmr, ms*)

Aloperine

A-637

16,17-Didehydro-9-de-2-piperidinylormosanine, 9CI. Allopterin [56293-29-9]



Absolute Configuration

C₁₅H₂₄N₂ 232.368

Alkaloid from seeds and green parts of *Sophora alopecuroides* (Fabaceae). Large prisms (petrol). Mp 73-75° (70-71°). $[\alpha]_D^{20} +82.1$ (c, 0.85 in EtOH). Incorr. stereochem. assigned in CA (1997).

►RM2979500

Hydrochloride: Mp 208°.

Hydrochloride (1:2): Mp 265°.

N-Benzoyl:

Cryst. (Et₂O). Mp 161-162°.

N-Me: N-Methylaloperine

[63128-33-6]

C₁₆H₂₆N₂ 246.395

Alkaloid from *Sophora alopecuroides* (Fabaceae). Mp 94-95°. $[\alpha]_D +140$ (EtOH).

N-(2-Propenyl): N-Allylaloperine

[56595-96-1]

C₁₈H₂₈N₂ 272.433

Alkaloid from *Sophora alopecuroides* (Fabaceae). Oil. Mp 235-237° (as hydrobromide)(synthetic).

11,12-Didehydro: 11,12-Didehydroaloperine

[142808-31-9]

C₁₅H₂₂N₂ 230.352

Alkaloid from leaves and stalks of *Sophora alopecuroides* (Fabaceae).

5,6,11,12,13,14,15,16-Octadehydro:**5,6,11,12,13,14,15,16-Octadehydroaloperine**C₁₅H₁₆N₂ 224.305

Alkaloid from *Sophora alopecuroides*. Oil. Unpublished data (1999).

Stereoisomer(1)(?): Sophora alopecuroides Alkaloid A₁C₁₅H₂₄N₂ 232.368

Alkaloid from *Sophora alopecuroides* (Fabaceae). Cryst. (Me₂CO). Mp 92°. $[\alpha]_D +195.5$ (MeOH).

Stereoisomer(2)(?): Sophora alopecuroides Alkaloid A₂C₁₅H₂₄N₂ 232.368

Alkaloid from *Sophora alopecuroides* (Fabaceae). Cryst. (petrol). Mp 206-208°. $[\alpha]_D +25.5$ (CHCl₃). Possibly the 11-epimer of Aloperine, since it gave the same salts but a different N-Me deriv.

Orechoff, A. *et al.*, *Ber.*, 1935, **68**, 431 (*isol*)
 Aslanov, Kh.A. *et al.*, *CA*, 1965, **63**, 2050h (*stereoisomers*)

Tolkachev, O.N. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 30; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 29 (*ms, pmr, struct*)

Kuchkarov, S. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 413; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 364 (*isol*)

Wang, Z. *et al.*, *Zhivyu Xuebao (Acta Bot. Sin.)*, 1991, **33**, 727; *CA*, **117**, 86697b (*11,12-Didehydroaloperine*)

Brosius, A.D. *et al.*, *Acta Cryst. C*, 1997, **53**, 1510-1512 (*cryst struct, abs config*)

Brosius, A.D. *et al.*, *J.O.C.*, 1997, **62**, 440 (*abs config, synth*)


Brosius, A.D. *et al.*, *J.A.C.S.*, 1999, **121**, 700-709 (*synth*)

Passarella, D. *et al.*, *Org. Lett.*, 2002, **4**, 2925-2928 (*synth*)

Alpinigenine

A-638

2,3,10,11-Tetramethoxy-16-methylrheadan-8-ol, 9CI. Alkaloid E†

**(+)-form**C₂₂H₂₇NO₆ 401.458

Biogenetic numbering shown. Other systems also used.

(+)-form [14028-91-2]

Alkaloid from *Papaver alpinum* various ssp., *Papaver bracteatum*, *Papaver fugax* (= *Papaver acausicum*), *Papaver orientale*

and *Papaver pseudo-orientale* (Papavera-
ceae). Prismatic cryst. (C₆H₆/MeOH or
EtOAc). Mp 193-195° (186.5-187.5°).
[α]_D²² +286 (c, 0.63 in MeOH). [α]_D²² +306
(c, 1.08 in MeOH).

Hydrochloride: Mp 161-163° dec. [α]_D¹⁸
+210 (c, 0.796 in MeOH).

Me ether: Epialpinine. O-Methylalpi-
nigine
[15210-99-8]
C₂₃H₂₉NO₆ 415.485

Alkaloid from *Papaver alpinum*, var-
ious ssp. (Papaveraceae). Cryst.
(MeOH aq.). Mp 105-106° Mp 122-
123°. [α]_D²⁵ +302 (c, 0.511 in MeOH).

Et ether: O-Ethylalpinigine

C₂₄H₃₁NO₆ 429.512
Isol. from *Papaver fugax* (Papavera-
ceae). Prob. an artifact.

8-Epimer, Me ether: Alpinine†. N-
Methylpapaverrubine G
[14028-90-1]

C₂₃H₂₉NO₆ 415.485
Alkaloid from *Papaver alpinum*, various
ssp., *Papaver bracteatum* and *Papaver
pseudo-orientale* (Papaveraceae). Non-
cryst. [α]_D²² +288 (c, 0.82 in CHCl₃).

*8-Epimer, Me ether, N-de-Me: Papaver-
rubine G*. N-Demethylalpinine
[17597-20-5]

C₂₂H₂₇NO₆ 401.458
Alkaloid from *Papaver alpinum*, var-
ious ssp., and *Papaver anomalum* (*Pa-
paver nudicaule* ssp. *amurense*)
(Papaveraceae). [α]_D +397 (c, 0.12 in
MeOH). Indefinite Mp.

(±)-*form* [40170-39-6]

Synthetic. Needles (EtOAc). Mp 172-
173° (168-169°).

Maturová, M. *et al.*, *Coll. Czech. Chem.
Comm.*, 1967, **32**, 419 (*ir, pmr, ms, struct*)
Pfeifer, S. *et al.*, *Pharmazie*, 1967, **22**, 343;
1968, **23**, 585; 1972, **27**, 48 (*Papaverrubine G*)

Dolejš, L. *et al.*, *Tetrahedron*, 1967, **23**, 2997
(*ms*)

Irie, H. *et al.*, *J.C.S. Perkin I*, 1972, 2986
(*synth*)

Rönsch, H. *et al.*, *Helv. Chim. Acta*, 1977, **60**,
2402 (*Epialpinine*)

Meshulam, H. *et al.*, *Phytochemistry*, 1980, **19**,
2633 (*isol*)

Lavie, D. *et al.*, *J.C.S. Perkin I*, 1981, 1019
(*isol, cmr, cryst struct, abs config*)

Prabhaker, S. *et al.*, *J.C.S. Perkin I*, 1981, 1273
(*synth*)


Phillipson, J.D. *et al.*, *Planta Med.*, 1981, **43**,
105; 261 (*O-Ethylalpinigine*)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1985,
33, 2273 (*synth*)

Alpkinidine

A-639

[486992-29-4]



C₁₉H₁₃N₅O₃ 331.33


Alkaloid from *Xestospongia cf. carbo-
naria*. Purple solid.

Thale, Z. *et al.*, *J.O.C.*, 2002, **67**, 9384-9391
(*isol, pmr, cmr, cryst struct*)

Alschomine

A-640

[123045-72-7]



Absolute
Configuration

C₂₁H₂₄N₂O₅ 384.431

Alkaloid from the leaves of *Alstonia
scholaris* (Apocynaceae). Prisms
(MeOH). Mp 233-240° dec. [α]_D²⁷ +116.1
(c, 0.25 in MeOH).

5-Epimer: Isoalschomine

[123062-73-7]

C₂₁H₂₄N₂O₅ 384.431


Alkaloid from leaves of *Alstonia scho-
laris* (Apocynaceae). Prisms (MeOH).
Mp 220-230° dec. [α]_D²⁷ +112.3 (c, 0.14
in MeOH).

Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**,
887 (*isol, uv, pmr, cmr, cryst struct*)

Alstocraline

A-641

[120416-01-5]



Absolute Configuration

C₄₃H₅₂N₄O₅ 704.908

Alkaloid from the leaves of *Alstonia
angustifolia* (Apocynaceae). [α]_D +3 (c,
0.7 in MeOH). λ_{max} 227 (log ε 4.02); 282
(log ε 4.45); 295 (sh) (log ε 4.44) (MeOH).

1'-N-De-Me, 1',2'-didehydro: Foliacraline
[120374-18-7]

C₄₂H₄₈N₄O₅ 688.865


Alkaloid from the leaves of *Alstonia
angustifolia* (Apocynaceae). [α]_D +55
(c, 0.3 in CHCl₃). λ_{max} 220 ; 228 ; 289 ;
302 (sh) (MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1988, **27**,
3955-3962 (*isol, uv, ir, pmr, cmr, ms, struct*)

Alstolactone

A-642

[701950-29-0]



Absolute
Configuration

C₂₀H₂₂N₂O₂ 322.406


Alkaloid from the leaves of *Alstonia
angustifolia* var. *latifolia*. Light yellow oil.
[α]_D -10 (c, 0.05 in CHCl₃). λ_{max} 216 (log
ε 4.23); 228 (log ε 4.3); 284 (log ε 3.76);
293 (log ε 3.71) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 2004, **65**,
603-608 (*isol, pmr, cmr*)

Alstomacroine

A-643

[160866-97-7]



C₂₁H₂₈N₂O₂ 340.464


Alkaloid from the leaves of *Alstonia
macrophylla*. [α]_D +2.6 (c, 0.02 in
CHCl₃). λ_{max} 230 (log ε 4.6); 285 (log ε
4.5) (MeOH).

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1994,
5, 201-209 (*isol, pmr, cmr, ms*)

Alstomacroline

A-644

[199800-14-1]



C₄₂H₅₀N₄O₅ 690.881


Alkaloid from the root bark of *Alstonia
macrophylla*. Amorph. powder. [α]_D²⁰ +54.6
(c, 0.4 in CHCl₃). λ_{max} 232 (log ε 4.44); 251
(log ε 3.88); 293 (log ε 3.78) (MeOH).

Keawpradub, N. *et al.*, *Phytochemistry*, 1997,
46, 757-762 (*isol, uv, ir, pmr, cmr, ms*)

Alstomacrophylline

A-645

[199800-12-9]




C₄₄H₅₄N₄O₅ 718.934

Alkaloid from the root bark of *Alstonia macrophylla* (Apocynaceae). Amorph. powder. $[\alpha]_D^{20}$ +42.8 (c, 0.09 in MeOH). λ_{\max} 232 (log ϵ 4.27); 260 (log ϵ 3.67); 292 (log ϵ 3.51) (MeOH).

Keawpradub, N. *et al.*, *Phytochemistry*, 1997, **46**, 757-762 (*isol, uv, ir, pmr, cmr, ms*)

Alstomaline A-646

[685508-21-8]


C₂₀H₂₂N₂O₃ 338.405

Represents a new structural subtype related to the akuammiline alkaloids, which has however been found previously in a small number of dimers, e.g. Flexicorine, F-88. Alkaloid from the leaves of *Alstonia macrophylla*. Yellowish oil. $[\alpha]_D$ -244 (c, 0.04 in CHCl₃). λ_{\max} 202 (log ϵ 3.77); 273 (log ϵ 3.85); 317 (log ϵ 3.12) (EtOH).

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*isol, pmr, cmr*)

Alstomicine A-647

[5594-43-4]


C₂₀H₂₆N₂O₂ 326.438

Alkaloid from the leaves of *Alstonia macrophylla*. Light yellowish oil. $[\alpha]_D$ +74 (c, 0.14 in CHCl₃). λ_{\max} 230 (log ϵ 3.8); 289 (log ϵ 3.11) (EtOH).

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*isol, pmr, cmr, ms*)

Alstonal A-648

20-Deacetyl-20-formyl-17-methylalstonisine, 9CI
[172924-28-6]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from bark of *Alstonia macrophylla*. Needles (petrol/Me₂CO). Mp 198-199°. $[\alpha]_D$ +167 (c, 0.062 in CHCl₃).

N-De-Me: N-Demethylalstonal

[301323-29-5]

C₁₉H₂₀N₂O₃ 324.379

Alkaloid from *Alstonia macrophylla*. Not obt. pure.

16-Hydroxy: 16-Hydroxyalstonal

[683239-99-8]

C₂₀H₂₂N₂O₄ 354.405

Alkaloid from the leaves of *Alstonia macrophylla*. Amorph. powder. $[\alpha]_D$ +153 (c, 0.26 in CHCl₃). λ_{\max} 207 (log ϵ 4.61); 263 (log ϵ 4.35) (EtOH).

11-Methoxy: N⁴-Demethylalstophyllal**oxindole. 11-Methoxyalstonal**

[172927-64-9]

C₂₁H₂₄N₂O₄ 368.432

From bark of *Alstonia macrophylla*. Needles (petrol/Me₂CO). Mp 190-191°. $[\alpha]_D$ +174 (c, 0.063 in CHCl₃).

11-Methoxy, 16-hydroxy: 16-Hydroxy-11-methoxyalstonal

[683240-00-8]

C₂₁H₂₄N₂O₅ 384.431

Alkaloid from the leaves of *Alstonia macrophylla*. Amorph. powder. $[\alpha]_D$ +203 (c, 0.11 in CHCl₃). λ_{\max} 220 (log ϵ 4.54); 267 (log ϵ 4.2) (EtOH).


Wong, W.-H. *et al.*, *Phytochemistry*, 1996, **41**, 313 (*isol, pmr, cmr, ms, struct*)

Kam, T.-S. *et al.*, *Tetrahedron*, 2000, **56**, 6143-6150 (*N-Demethylalstonal*)

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*16-Hydroxyalstonal, 16-Hydroxy-11-methoxyalstonal*)

Alstonerine A-649

Alstophyllan-19-one, 9CI. Alkaloid D[†]
[25921-22-6]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from bark of *Alstonia muelleriana* (Apocynaceae). Cryst. (Et₂O). Mp 172-173°. $[\alpha]_D^{25}$ -195 (c, 0.6 in EtOH).

N⁴-De-Me: N⁴-Demethylalstonerine

[701304-79-2]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from the leaves of *Alstonia angustifolia* var. *latifolia*.

11-Methoxy: Alstophylline

[3382-41-0]

C₂₂H₂₆N₂O₃ 366.459

Alkaloid from *Alstonia macrophylla* and *Alstonia glabriflora* (Apocynaceae). Cryst. (CH₂Cl₂/hexane). Mp 155-158°. $[\alpha]_D^{26}$ -151 (c, 0.326 in MeOH).

11-Methoxy, N¹-de-Me: N¹-DemethylalstophyllineC₂₁H₂₄N₂O₃ 352.432

Alkaloid from the bark of *Alstonia macrophylla*.

11-Methoxy, 6-oxo: 6-Oxoalstophylline

[683240-01-9]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from the leaves of *Alstonia macrophylla*.

Kishi, T. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 1349; 1966, **49**, 946 (*Alstonerine, Alstophylline, isol, uv, ir, pmr, struct, synth*)

Cook, J.M. *et al.*, *Chem. Comm.*, 1969, 1306 (*uv, ir, pmr, ms, struct*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2739 (*Alstophylline, isol*)

Elderfield, R.C. *et al.*, *Phytochemistry*, 1972, **11**, 339 (*isol, ir, uv*)

Garnick, R.L. *et al.*, *J.A.C.S.*, 1978, **100**, 4213 (*synth, ms*)

Zhang, L.H. *et al.*, *J.A.C.S.*, 1990, **112**, 4088 (*synth*)

Takayama, H. *et al.*, *Tetrahedron*, 1991, **47**, 1383 (*synth*)

Bi, Y. *et al.*, *J.A.C.S.*, 1994, **116**, 9027 (*synth*)

Yu, P. *et al.*, *J.O.C.*, 2000, **65**, 3173-3191 (*synth*)

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*6-Oxoalstophylline*)

Kam, T.-S. *et al.*, *Phytochemistry*, 2004, **65**, 603-608 (*N⁴-Demethylalstonerine*)

Kam, T.-S. *et al.*, *Tetrahedron*, 2004, **60**, 3957-3966 (*N¹-Demethylalstophylline*)


Tran, Y.S. *et al.*, *Org. Lett.*, 2005, **7**, 4289-4291 (*synth*)

Liao, X. *et al.*, *J.O.C.*, 2006, **71**, 8884-8890 (*synth*)

Miller, K.A. *et al.*, *Org. Lett.*, 2007, **9**, 1113-1116 (*synth*)

Alstonidine A-650

[25394-75-6]



Relative Configuration

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from *Alstonia constricta* (Apocynaceae). Mp 188-190°. $[\alpha]_D$ -99 (c, 1.0 in CHCl₃).

Ac:

Needles + 3H₂O (Me₂CO aq.). Mp 92-96°.

14-Oxo: 14-KetoalstonidineC₂₂H₂₂N₂O₅ 394.426

Alkaloid from stem bark of *Alstonia constricta* (Apocynaceae). Noncryst. Labile.


Boaz, H. *et al.*, *J. Am. Pharm. Assoc.*, 1957, **46**, 510 (*isol, uv, ir, struct*)

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2489 (*isol, pmr, ms, config*)

Allam, K. *et al.*, *J. Nat. Prod.*, 1987, **50**, 623 (*14-Ketoalstonidine*)

Alstonilidine A-651

[31185-76-9]


C₂₃H₁₈N₂O₆ 418.405

A secoyohimbane. Alkaloid from the

root bark of *Alstonia constricta* (Apocynaceae). Needles (EtOH). Mp 244-245°. Crow, W.D. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2489 (*isol, uv, ir, pmr, ms, struct*)

Alstonine**A-652**

[47485-83-6]



Absolute Configuration

C₂₁H₂₀N₂O₃ 348.401

The early history of alkaloids called Alstonine is confused. Alkaloid from *Alstonia constricta*, *Rauwolfia obscura*, *Rauwolfia volkensii*, *Rauwolfia nitida*, *Rauwolfia serpentina*, *Catharanthus roseus*, *Strychnos camptoneura* and *Strychnos gossweileri* (Apocynaceae, Loganiaceae). Shows neoplasm-inhibiting props. Mp 205-210° dec. λ_{max} 252; 289; 309; 336; 369 (MeOH) (Berdy).

Hydrochloride: Mp 278-279° dec.*Perchlorate*: Mp 239-240°. [α]_D²⁰ +154 (Me₂CO).

10,11-Dimethoxy: Bleekerine. 10,11-Dimethoxyalstonine
[41758-43-4]

C₂₃H₂₄N₂O₅ 408.453

Alkaloid from the stem bark of *Bleekeria vitiensis* (preferred genus name *Ochrosia*) (Apocynaceae). Pale yellow prisms (EtOH). Mp 276-277°. [α]_D^{22.5} +612 (MeOH).

10,11-Dimethoxy, 5,6-dihydro: Ochroposine

[38146-72-4]

C₂₃H₂₆N₂O₅ 410.469

Alkaloid from the bark of *Ochrosia oppositifolia* and *Ochrosia moorei* (Apocynaceae). Mp 312-314° dec. (as perchlorate). λ_{max} 222 (log ε 4.44); 401 (log ε 4.43) (no solvent reported) (perchlorate).

19-Epimer: 19-Epialstonine

C₂₁H₂₀N₂O₃ 348.401

Alkaloid from the roots of *Amphicome emodi* (preferred genus name *Incarvillea*). Cryst. Mp 245° dec.

20-Epimer: Serpentine

[18786-24-8]

C₂₁H₂₀N₂O₃ 348.401

Alkaloid from *Rauwolfia serpentina* and many other *Rauwolfia* spp., *Strychnos camptoneura*, *Catharanthus roseus*, *Catharanthus ovalis* and *Vinca major* (Apocynaceae). Shows antitumour activity. Mp 156-157° dec. (170°). [α]_D +292 (MeOH). λ_{max} 251 (log ε 4.5); 307 (log ε 4.31); 368 (log ε 3.55) (MeOH).

20-Epimer, hydrochloride: Mp 246-248°. [α]_D²⁵ +178 (H₂O).

19,20-Diepimer: 19-Episerpentine

[64753-57-7]

C₂₁H₂₀N₂O₃ 348.401

Alkaloid from the roots of *Rauwolfia cumminsi* (Apocynaceae). Yellow

cryst. powder. Mp 240° dec.

19,20-Diepimer, 11-methoxy: Serpentine

[81149-04-4]

C₂₂H₂₂N₂O₄ 378.427

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae). Yellow plates + 5H₂O (MeOH/C₆H₆). Mp 275-276°. λ_{max} 265 (log ε 4.4); 310 (log ε 4.2); 365 (log ε 3.6) (no solvent reported).


19,20-Diepimer, 11-methoxy, hydrochloride: Mp 289-290°.

19,20-Diepimer, 11-methoxy, picrate: Mp 196-198°.

[642-18-2]

Sharp, T.M. *et al.*, *J.C.S.*, 1934, 287-291 (*isol*)Raymond-Hamet, *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1948, **227**, 344-345 (*uv*)Elderfield, R.C. *et al.*, *J.O.C.*, 1951, **16**, 506-523 (*struct*)Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 271-276; 1954, **37**, 1912-1920 (*isol*, *Serpentine, uv, ir, struct*)Bader, F.E. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 215-226 (*struct*)Wenkert, E. *et al.*, *J.A.C.S.*, 1957, **79**, 1519-1520; 1958, **80**, 1613-1619 (*config, synth*)Fritz, H. *et al.*, *Annalen*, 1962, **655**, 148-167 (*Serpentine, config, ir, uv*)Saxton, J.E. *et al.*, *Alkaloids (Academic Press)*, 1965, **8**, 159-202 (*bibl*)Battersby, A.R. *et al.*, *Chem. Comm.*, 1966, 888-890 (*biosynth*)Peube-Locou, N. *et al.*, *Phytochemistry*, 1972, **11**, 2109-2111 (*Ochroposine*)Sainsbury, M. *et al.*, *Phytochemistry*, 1972, **11**, 2337-2339 (*Bleekerine*)Timmins, P. *et al.*, *Phytochemistry*, 1976, **15**, 733-735 (*isol*)Iwu, M.M. *et al.*, *Planta Med.*, 1977, **32**, 158-161 (*19-Episerpentine*)Malik, A. *et al.*, *Heterocycles*, 1981, **16**, 1727-1733 (*Serpentine*)Ahond, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 193-199 (*Ochroposine, isol*)Dinda, B. *et al.*, *Indian J. Chem., Sect. B*, 2002, **41**, 2698-2700 (*19-Epialstonine*)Wachsmuth, O. *et al.*, *Phytochemistry*, 2002, **61**, 705-709 (*Serpentine, isol, pmr, cmr, ms*)**Alstonisidine****A-653**

[36474-13-2]



Absolute Configuration

C₄₂H₄₈N₄O₄ 672.866

Alkaloid from the bark of *Alstonia muelleriana* (Apocynaceae). Rods (MeOH). Mp 325° dec. [α]_D -234 (c, 1.0 in EtOH). [α]_D -133 (c, 0.208 in CHCl₃) (natural). [α]_D -144 (c, 0.335 in CHCl₃) (synthetic).


Hydrochloride (1:2):

Cryst. (MeOH). Mp 268° (dec. *in vacuo*). [α]_D²⁵ +137 (c, 0.6 in H₂O).

Cook, J.M. *et al.*, *J.O.C.*, 1971, **36**, 582 (*uv, ir,*
pmr, ms)Elderfield, R.C. *et al.*, *Phytochemistry*, 1972, **11**, 339 (*isol, uv, ir*)Burke, D.E. *et al.*, *J.A.C.S.*, 1973, **95**, 546 (*synth, struct*)Hoard, L.D. *et al.*, *Diss. Abstr. Int., B*, 1977, **38**, 2690 (*cryst struct*)**Alstonisine, 9CI****A-654**

Alkaloid C†

[35804-91-2]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the bark of *Alstonia muelleriana* and *Alstonia macrophylla* (Apocynaceae). Cryst. (MeOH aq.). Mp 168-169°. [α]_D²⁵ +200 (c, 1.0 in EtOH).

Hydrochloride: Mp 250-260° dec. [α]_D²⁵ +155 (c, 1.0 in H₂O).

N⁴-Formyl: N⁴-Formylalstonisine. Alstofoline
[301521-97-1]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from *Alstonia macrophylla*. [α]_D +39 (c, 0.21 in CHCl₃). λ_{max} 214 (log ε 4.02); 246 (log ε 4.01); 254 (log ε 4.05); 290 (log ε 3.13) (EtOH).

N-De-Me: N-Demethylalstonisine

[301323-28-4]

C₁₉H₂₀N₂O₃ 324.379Alkaloid from *Alstonia macrophylla*.

16-Hydroxy: 16-Hydroxyalstonisine

[683239-98-7]

C₂₀H₂₂N₂O₄ 354.405

Alkaloid from the leaves of *Alstonia macrophylla*. Amorph. powder. [α]_D +170 (c, 0.15 in CHCl₃). λ_{max} 209 (log ε 4.17); 256 (log ε 3.95) (EtOH).

11-Methoxy: N^b-Demethylalstophylline oxindole

[108195-73-9]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from leaves and bark of *Alstonia macrophylla*. Prisms (petrol/Me₂CO). Mp 205-206°. [α]_D +126 (c, 0.068 in CHCl₃).

11-Methoxy, 16-hydroxy: 16-Hydroxy-N^b-demethylalstophylline oxindole. 14-Hydroxy-11-methoxyalstonisine, 9CI

[114466-39-6]

C₂₁H₂₄N₂O₅ 384.431

Alkaloid from leaves of *Alstonia macrophylla*. Mp 215°. [α]_D +104.47 (c, 0.067 in CHCl₃).

7-Epimer: Isoalstonisine

[301521-98-2]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia macrophylla*. [α]_D +207 (c, 0.07 in CHCl₃). λ_{max} 210 (log ε 3.93); 255 (log ε 3.8) (EtOH).


Nordman, C.E. *et al.*, *J.A.C.S.*, 1963, **85**, 353 (*cryst struct*)Elderfield, R.C. *et al.*, *Phytochemistry*, 1972, **11**, 339 (*isol, uv, ir*)

- Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1987, **26**, 865 (*N^b*-Demethylalstophylline oxindole)
 Atta-ur-Rahman, *et al.*, *Heterocycles*, 1988, **27**, 725 (*16-Hydroxy-N^b*-demethylalstophylline oxindole)
 Wong, W.-H. *et al.*, *Phytochemistry*, 1996, **41**, 313 (*N^b*-Demethylalstophylline oxindole)
 Kam, T.-S. *et al.*, *Tetrahedron*, 2000, **56**, 6143-6150 (*Isaalstonisine*, *N*-Demethylalstonisine, *Alstofoline*)
 Wearing, X.Z. *et al.*, *Org. Lett.*, 2002, **4**, 4237-4240 (*synth*)
 Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*16-Hydroxyalstonisine*)

Alstonoxine B

A-655

[301521-96-0]



$C_{19}H_{26}N_2O_3$ 330.426
 Alkaloid from *Alstonia macrophylla*. $[\alpha]_D$ -12 (c, 0.41 in $CHCl_3$). λ_{max} 213 (log ϵ 4.15); 255 (log ϵ 3.65) (EtOH).

19-Ketone: Alstonoxine A

[301521-95-9]

 $C_{19}H_{24}N_2O_3$ 328.41


Alkaloid from *Alstonia macrophylla*. $[\alpha]_D$ -34 (c, 0.19 in $CHCl_3$). λ_{max} 216 (log ϵ 3.89); 255 (log ϵ 3.7); 285 (log ϵ 3.18) (EtOH).

Kam, T.-S. *et al.*, *Tetrahedron*, 2000, **56**, 6143-6150

Alstophyllal

A-656

[685508-23-0]



$C_{22}H_{26}N_2O_3$ 366.459
 Alkaloid from the leaves of *Alstonia macrophylla*.

6-Oxo: 6-Oxoalstophyllal

[683240-02-0]


 $C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the leaves of *Alstonia macrophylla*.

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 547-552 (*isol*, *pmr*, *cmr*)

Alstoscholarine

A-657



Absolute
 Configuration

(E)-form

 $C_{22}H_{20}N_2O_3$ 360.412**(E)-form** [937245-47-1]

Alkaloid from the leaves of *Alstonia scholaris*. Green prisms. Mp 168-170°. $[\alpha]_D^{20}$ -53 (c, 0.065 in $CHCl_3$). λ_{max} 223; 291; 300 ($CHCl_3$).


(Z)-form [937245-48-2]

Alkaloid from the leaves of *Alstonia scholaris*. Powder. $[\alpha]_D^{25}$ -140 (c, 0.05 in $CHCl_3$). λ_{max} 220; 286; 293 ($CHCl_3$).
 Cai, X.-H. *et al.*, *Org. Lett.*, 2007, **9**, 1817-1820 (*isol*, *pmr*, *cmr*, *cryst struct*)

Alstoumerine

A-658

[135824-73-6]



Absolute
 Configuration


$C_{20}H_{24}N_2O_2$ 324.422
 Alkaloid from *Alstonia macrophylla* (Apocynaceae). Pale yellow cryst. Mp 170°. $[\alpha]_D$ -5.5 (c, 0.0034 in $CHCl_3$).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1991, **54**, 750-754 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Altamycin A

A-659

[60202-22-4]




$C_{30}H_{43}NO_9$ 561.671
 Polyene-type antibiotic. Sol. MeOH, EtOH, Me_2CO , DMF, bases, AcOH, Py, butanol; poorly sol. dioxan, MeCN, $CHCl_3$, acids, H_2O , hexane. Mp 140-142°. $[\alpha]_D$ -72 (c, 0.5 in EtOH). No biol. source referred to in abstract. λ_{max} 262 (E1%/1cm 105); 343 (E1%/1cm 400); 355 (E1%/1cm 450); 375 (E1%/1cm 320) (MeOH) (Berdy).

Shenin, Yu.D. *et al.*, *CA*, 1987, **106**, 66971f (*struct*)

Altemicidin

A-660

Artemicidine
 [125399-82-8]



$C_{13}H_{20}N_4O_7S$ 376.39
 Prod. by *Streptomyces sioyaensis*. Possesses acaricidal, mitocidal and antitumour activities. Powder. Sol. H_2O ; fairly sol. MeOH, EtOH; poorly sol. Me_2CO , hexane. Mp 195-199° dec. $[\alpha]_D^{27}$ -7.6 (c, 1 in H_2O). λ_{max} 305 (ϵ 11300) (0.1M HCl)

(Derep). λ_{max} 300 (ϵ 20800) (0.1M NaOH) (Derep). λ_{max} 300 (ϵ 20300) (pH 6.8 phosphate buffer) (Derep).

▶ LD₅₀ (mus, ivn) 0.3 mg/kg. UW8797000

5'-N-(2S-Amino-3S-methylpentanoyl):

Antibiotic SB 203207 $C_{19}H_{31}N_5O_8S$ 489.549

Prod. by *Streptomyces* sp. NCIMB 40513. Isoleucyl tRNA synthetase inhibitor. Powder. λ_{max} 299 (H_2O).

5'-N-(2S-Amino-3S-methylpentanoyl), 6-

O-(2S-amino-3S-phenylbutanoyl):

Antibiotic SB 203208 $C_{29}H_{42}N_6O_9S$ 650.752

Prod. by *Streptomyces* sp. NCIMB 40513. Isoleucyl tRNA synthetase inhibitor. Powder. λ_{max} 297 (H_2O).

Takahashi, A. *et al.*, *J. Antibiot.*, 1989, **42**,

1556-1561; 1562-1566 (*isol*, *struct*)

Takahashi, A. *et al.*, *Tetrahedron*, 1991, **47**,

3621-3632 (*pmr*, *cmr*)

Kende, A.S. *et al.*, *J.A.C.S.*, 1995, **117**, 10597-

10598 (*synth*)

Stefanska, A.L. *et al.*, *J. Antibiot.*, 2000, **53**,


357-363 (*SB 203207/8*)

Houge-Frydrych, C.S.V. *et al.*, *J. Antibiot.*,

2000, **53**, 364-372 (*SB 203207/8*)**Alteramide A**

A-661

[142131-06-4]

 $C_{29}H_{38}N_2O_6$ 510.629

Macrocyclic lactam antibiotic. Isol. from an *Alteromonas* sp. associated with the sponge *Halichondria okadai*. Exhibits cytotoxicity against murine leukaemia P388 cells, murine lymphoma L1210 cells and the human epidermoid carcinoma KB cells *in vitro*. Yellow powder. Sol. MeOH, EtOAc, $CHCl_3$; poorly sol. H_2O . Mp 200° (dec.). $[\alpha]_D^{22}$ +36.2 (c, 0.1 in MeOH). λ_{max} 268 (ϵ 30300); 347 (ϵ 11000) (MeOH) (Derep).

25-Deoxy: Alteramide B $C_{29}H_{38}N_2O_5$ 494.63

From an *Alteromonas* sp. Sol. MeOH, EtOAc, $CHCl_3$; poorly sol. H_2O . λ_{max} 268; 347 (MeOH) (Berdy).


[142131-07-5]

Shigemori, H. *et al.*, *J.O.C.*, 1992, **57**, 4317-4320 (*isol*, *ir*, *uv*, *pmr*, *cmr*, *struct*)

Alternatamide B

A-662

[191212-58-5]



C₁₇H₂₂Br₃N₃O 524.029

Alkaloid from the marine bryozoan *Amathia alternata*. Moderate antibacterial agent. Amorph. powder.

N¹-Me: **Alternatamide A**

[191212-57-4]

C₁₈H₂₄Br₃N₃O 538.119

Alkaloid from *Amathia alternata*. Amorph. powder.

2-Debromo: **Alternatamide D**

[191212-60-9]

C₁₇H₂₃Br₂N₃O 445.196

Alkaloid from *Amathia alternata*. Amorph. powder.

6-Debromo: **Alternatamide C**

[191212-59-6]


C₁₇H₂₃Br₂N₃O 445.196

Alkaloid from *Amathia alternata*. Amorph. powder.

Lee, N.-K. *et al.*, *J. Nat. Prod.*, 1997, **60**, 697-699 (*isol, uv, cd, ir, pmr, cmr, ms*)

Altersetin

A-663



Absolute Configuration

C₂₄H₃₃NO₄ 399.529


Tetramic acid antibiotic. Enolised triketone. Related to Equisetin, E-142 and Phomasetin, P-373. Prod. by *Alternaria* spp. P 0506 and P 0535. Active against gram-positive and -negative bacteria and yeasts. Solid. λ_{\max} 234 (log ϵ 4.37); 286 (log ϵ 3.84) (MeOH).

Hellwig, V. *et al.*, *J. Antibiot.*, 2002, **55**, 881-892 (*isol, cd, pmr, cmr, ms*)

Althiomycin, 8CI

A-664

Matamycin. Antibiotic 116A [12656-40-5]

C₁₆H₁₇N₅O₆S₂ 439.472

Peptide antibiotic. The nat. prod. has the thiazole-ring S-config. as illus. but is prob. a mixt. of side-chain diastereoisomers. The oxime config. also appears uncertain although the Z-isomer has been prepd. synthetically. Isol. from *Cystobacter fuscus*, *Myxococcus virescens*, *Myxococcus xanthus*, *Streptomyces althioticus*, *Streptomyces chartreusis* and *Streptomyces matensis*. Active against gram-positive organisms. Also shows anticoccidial and antihelminthic activity. Cryst. (CH₂Cl₂/EtOH). Mp 181-184°

(dec.). $[\alpha]_D^{25} +37.8$ (c, 2 in EtOH/CH₂Cl₂). λ_{\max} 235 (ϵ 22800); 312 (ϵ 11500) (0.1N NaOH) (Derep). λ_{\max} 222 (ϵ 36600); 238 (ϵ 28300); 285 (ϵ 8200) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 720 mg/kg.

Yamaguchi, H. *et al.*, *J. Antibiot., Ser. A*, 1957, **10**, 195 (*isol*)

Cram, D.J. *et al.*, *J.A.C.S.*, 1963, **85**, 1430 (*uv, ir, pmr*)

Sakakibara, H. *et al.*, *J. Antibiot.*, 1974, **27**, 897 (*pmr, cmr, cryst struct*)

Pestka, S. *et al.*, *Antibiotics (N.Y.)*, 1975, **3**, 323 (*rev*)

Bycroft, B.W. *et al.*, *Chem. Comm.*, 1975, 121 (*struct, ms, ir, uv, pmr, cmr*)

Kirst, H.A. *et al.*, *J. Antibiot.*, 1975, **28**, 286 (*struct, nmr, ir, uv, ms*)

Kunze, B. *et al.*, *J. Antibiot.*, 1982, **35**, 635 (*isol, bibl*)

Shiba, T. *et al.*, *Tet. Lett.*, 1984, **25**, 2009 (*synth*)

Inami, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 2185 (*activity, synth, derivs*)

Toogood, P.L. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 1543 (*synth*)

Alvanidine

A-665

C₂₀H₃₃NO₂ 319.486

Prob. a steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria raddeana* (Liliaceae) occurring with Alvanine, A-666 and Raddeamine, R-7. Cryst. Mp 235-236°.

Hydrochloride: Mp 174-175°.

Aslanov, Kh.A. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 579; *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 623

Alvanine

A-666

C₂₆H₄₃NO₃ 417.631

Prob. a steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria raddeana* (Liliaceae). Cryst. Mp 185-187°. Co-occurs with Alvanidine, A-665 and Raddeamine, R-7.


Hydrochloride: Mp 163-165°.

Aslanov, Kh.A. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 579-583; *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 623

Alvimine

A-667

1-Acetyl-17-(acetyloxy)-20,21-didehydro-19-hydroxy-10,11-dimethoxy-4-methyl-3,4-seco-curran-3-one, 9CI [82083-63-4]



Absolute configuration

C₂₆H₃₄N₂O₇ 486.564


Alkaloid from bark of *Strychnos alvimiana* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 124-126°. $[\alpha]_D^{20} +282$ (c, 0.5 in EtOH).

Marini-Bettolo, G.B. *et al.*, *An. Asoc. Quim. Argent.*, 1982, **70**, 263 (*isol, ir, uv, pmr, ms, struct*)

Alviminine

A-668

[82083-62-3]



Probable Absolute Configuration

C₂₄H₂₈N₂O₅ 424.496


Tentative struct. Alkaloid from the bark of *Strychnos alvimiana* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 103-106°.

Marini-Bettolo, G.B. *et al.*, *An. Asoc. Quim. Argent.*, 1982, **70**, 263 (*isol, uv, pmr, cmr, ms, struct*)

Amabiline†

A-669

[17958-43-9]

C₁₅H₂₅NO₄ 283.367

Amabiline and its stereoisomers are esters of Supinidine, S-644 with stereoisomers of 2,3-Dihydroxy-2-isopropylbutanoic acid. Alkaloid from *Cynoglossum amabile* dried whole plants and from whole plants of *Neatostema apulum* (Boraginaceae). Noncryst. $[\alpha]_D^{20} -7.1$ (c, 2.02 in EtOH). Cryst. derivs. could not be obt.

N-Oxide: **Amabiline N-oxide**

[145265-23-2]

C₁₅H₂₅NO₅ 299.366

Alkaloid from whole plants of *Neatostema apulum* (Boraginaceae). Noncryst. $[\alpha]_D -12$ (c, 1.4 in EtOH).

8-Epimer: **Cynaustine**

[17958-39-3]

C₁₅H₂₅NO₄ 283.367

Minor alkaloid from *Cynoglossum australe* dried plants (Boraginaceae). Pale-yellow gum. $[\alpha]_D^{20} +13.2$ (c, 1.59 in EtOH). Ester of (+)-supinidine with (-)-viridifloric acid.

▶ EK7792000

8-Epimer, **picrate**:

Yellow plates (EtOH). Mp 135-136°.

3'-Epimer: **Supinine**

[551-58-6]

C₁₅H₂₅NO₄ 283.367

Alkaloid from *Heliotropium supinum* and *Heliotropium europaeum* (Boraginaceae). Shows antineoplastic props. Needles (Me₂CO). Mp 148-149° (146-147.5°). $[\alpha]_D -23.8$ (EtOH). $[\alpha]_D -12.1$ (c, 1.98 in EtOH). Log P 0.2 (calc).

▶ Hepatotoxin. WU2050000

3'-Epimer, O³-Me, N-oxide: **Helewine N-oxide**

[132886-10-3]

C₁₆H₂₇NO₅ 313.393

Alkaloid from aerial parts of *Heliotropium hirsutissimum* (Boraginaceae).

Gummy solid. $[\alpha]_D^{20} +19.5$ (c, 0.55 in CHCl_3).

3'-Epimer, O³-Me: **Heleurine**. Alkaloid C†

[488-00-6]
C₁₆H₂₇NO₄ 297.394

Alkaloid from *Heliotropium indicum* and *Heliotropium europaeum* (Boraginaceae). Large transparent prisms. Sol. H₂O. Mp 67-68°. $[\alpha]_D^{15} -12$ (c, 5.15 in EtOH). Discolours rapidly unless v. pure.

▶EK7879200

2'-Epimer: **Heliovinine**

[84471-36-3]
C₁₅H₂₅NO₄ 283.367

Minor alkaloid from *Heliotropium curassavicum* (Boraginaceae). Gum. Probable struct. Obt. only as a mixt. with Coromandalinine and Curassavinine.

2',3'-Diepimer: **Coromandalinine**

[85798-06-7]
C₁₅H₂₅NO₄ 283.367

Probable struct. Obt. only as a mixt. with Heliovinine and Curassavinine. Minor alkaloid from *Heliotropium curassavicum* (Boraginaceae). Gum.

Men'shikov, G.P. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1949, **19**, 1382; *CA*, **44**, 3486 (isol, *Supinine*)

Culvenor, C.C.J. et al., *Aust. J. Chem.*, 1954, **7**, 287; 1967, **20**, 2499 (*Supinine*, *Amabiline*, *Cynaustine*, *Heleurine*)

Culvenor, C.C.J. et al., *Chem. Ind. (London)*, 1959, 20 (*synth*, *Supinine*)

Mattocks, A.R. et al., *Nature (London)*, 1968, **217**, 723 (*tox*, *Supinine*)

Šimánek, V. et al., *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*, *Supinine*)

Culvenor, C.C.J. et al., *J.C.S. (C)*, 1971, 3653 (*cd*)

Mohanraj, S. et al., *Phytochemistry*, 1982, **21**, 1775 (*Coromandalinine*, *Heliovinine*)

Mackay, M.F. et al., *Acta Cryst. C*, 1985, **41**, 722-725 (*cryst struct*, *Supinine*, *Heleurine*)

Roeder, E. et al., *Phytochemistry*, 1992, **31**, 3613 (*Amabiline N-oxide*)

Constantinidis, T. et al., *Phytochemistry*, 1993, **32**, 1335 (*Heleurine N-oxide*)


Logie, C.G. et al., *Phytochemistry*, 1994, **37**, 43-109 (*rev*, *pmr*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SOW500

Amabiline†

A-670

Crinan-1,2-diol, 9CI
[151204-56-7]



Absolute Configuration

C₁₆H₁₉NO₄ 289.33

Alkaloid from bulbs of *Crinum amabile* and *Crinum kirkii* (Amaryllidaceae). Mp 210° dec. $[\alpha]_D^{20} -32$ (c, 0.3 in EtOH). λ_{max} 209 (log ϵ 4.13); 234 (log ϵ 3.49); 292 (log ϵ 3.61) (EtOH).

2-Epimer, O²-Ac: **Josephine**

[157469-84-6]
C₁₈H₂₁NO₅ 331.368

Alkaloid from bulbs of *Brunsvigia josephinae* (Amaryllidaceae). Mp 230-232°. $[\alpha]_D^{22} -30.9$ (c, 0.5 in EtOH).

1,2-Diepimer: **4a-Deoxycrinamabine**. 4a-Dehydroxycrinamabine

[211234-46-7]
C₁₆H₁₉NO₄ 289.33

Alkaloid from the bulbs of *Crinum amabile*. Mp 208-210° dec. $[\alpha]_D^{29} +28$ (c, 0.1 in MeOH). λ_{max} 205 (log ϵ 4.07); 237 (sh) (log ϵ 3.4); 289 (log ϵ 3.24) (MeOH).

1,2-Diepimer, 1,2-di-Ac: **1-Epidemethoxybowdensine**

[184900-14-9]
C₂₀H₂₃NO₆ 373.405

Alkaloid from bulbs of *Brunsvigia orientalis*. Mp 96-98°. $[\alpha]_D^{22} +20$ (c, 0.78 in CHCl_3).

Likhitwitayawuid, K. et al., *J. Nat. Prod.*, 1993, **56**, 1331-1338 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Viladomat, F. et al., *Phytochemistry*, 1994, **35**, 809-812; 1996, **43**, 1379-1384 (*Josephine*, *1-Epidemethoxybowdensine*)

Pearson, W.H. et al., *J.O.C.*, 1998, **63**, 3607-3617 (*synth*)


Pham, L.H. et al., *Phytochemistry*, 1998, **48**, 371-376 (*4a-Deoxycrinamabine*)

Machocho, A.K. et al., *Phytochemistry*, 2004, **65**, 3143-3149 (*isol*)

Amaminol A

A-671

[261622-18-8]



Absolute Configuration

C₁₈H₃₁NO 277.449

Isol. from an unidentified tunicate of the family Polyclinidae. Cytotoxic agent. Pale yellow oil. $[\alpha]_D^{24} -170.8$ (c, 0.2 in MeOH).

2-Epimer: **Amaminol B**

[261622-22-4]
C₁₈H₃₁NO 277.449

Isol. from an unidentified tunicate of the family Polyclinidae. Cytotoxic agent. Pale yellow oil. $[\alpha]_D^{24} -112.4$ (c, 0.2 in MeOH).


Sata, N.U. et al., *Tet. Lett.*, 2000, **41**, 489-492 (*isol*, *pmr*, *cmr*)

Kumpulainen, E.T.T. et al., *Org. Lett.*, 2007, **9**, 5043-5045 (*synth*, *abs config*)

Amamistatin A

A-672

[225505-59-9]



Absolute Configuration

C₃₇H₅₅N₅O₁₁ 745.868

Prod. by *Nocardia asteroides* SCRC-A 2359. Antitumour and antiproliferative agent. Mast cell growth inhibitor. $[\alpha]_D^{26} -9.8$ (c, 0.61 in MeOH). Related to Formobactin, F-130 and Nocobactin NA, N-267. λ_{max} 272 (c 22000); 335 (c 11000) (MeOH).

Demethoxy: **Amamistatin B**

[225505-77-1]
C₃₆H₅₃N₅O₁₀ 715.842

Prod. by *Nocardia asteroides* SCRC-A 2359. Antitumour and antiproliferative agent. Mast cell growth inhibitor. $[\alpha]_D^{25} -8.2$ (c, 0.47 in MeOH). λ_{max} 266 (c 22000); 307 (c 12000) (MeOH).

Yokokawa, F. et al., *Tetrahedron*, 2000, **56**, 3027-3034 (*synth*)


Kokubo, S. et al., *Tetrahedron*, 2000, **56**, 6435-6440 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *abs config*)

Fennell, K.A. et al., *J.O.C.*, 2008, **73**, 1018-1024 (*synth*)

Amanitin†

A-673

Amanitin. *Amatoxin*



α -Amanitin R = NH₂, R' = R'' = OH
 β -Amanitin R' = R'' = OH
 γ -Amanitin R = NH₂, R' = OH, R'' = H
 ϵ -Amanitin R = R' = OH, R'' = H
 Amanullin R = NH₂, R' = R'' = H
 Amanullin acid R = OH, R' = R'' = H

α -Amanitin [23109-05-9]

C₃₉H₅₄N₁₀O₁₄S 918.98

Toxic constit. of *Amanita phalloides*. Also from *Amanita verna*, *Amanita bisporigera*, *Galerina autumnalis*, *Galerina marginata* and *Galerina venenata*. Needles. Sol. MeOH, H₂O. Mp 254-255° dec. $[\alpha]_D^{20} +191$ (H₂O). λ_{max} 302 (MeOH) (Berdy).

▶ Systemic effects in humans by ingestion. Can cause hepatic necrosis, liver failure and death if untreated. Nephrotoxic. LD₅₀ (mus, ipr) 0.1 mg/kg. Some human and exp. toxic effects are delayed (2-8 d after administration) cf. phallotoxins. Gastrointestinal effects occur 2-4 h after ingestion. BD6195000

Deoxy: **Amaninamide**

[58311-65-2]
C₃₉H₅₄N₁₀O₁₃S 902.981

Isol. from *Amanita virosa*. Lacks the OH group on the tryptophan ring.

β -Amanitin

1-L-Aspartic acid- α -amanitin, 9CI
[21150-22-1]

C₃₉H₅₃N₉O₁₅S 919.965

Toxic constit. of several spp. of *Amanita*, notably *Amanita phalloides* and also

found in *Galerina autumnalis*, *Galerina venenata*, *Galerina marginata* and *Amanita bisporigera*. Cryst. (EtOH). Sol. H₂O, EtOH. Mp 300°. λ_{\max} 302 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 0.4 mg/kg. NJ8324000

Deoxy: Amanin. 1-L-Aspartic acid-4-(2-mercapto-L-tryptophan)- α -amanitin, 9CI

[21150-21-0]

C₃₉H₅₃N₉O₁₄S 903.966

Constit. of *Amanita phalloides* toxin.

Lacks the OH-group on the tryptophan ring.

► LD₅₀ (mus, ipr) 0.5 mg/kg. NJ8326000

γ -Amanitin

3-(4-Hydroxy-L-isoleucine)- α -amanitin, 9CI

[21150-23-2]

C₃₉H₅₄N₁₀O₁₃S 902.981

Constit. of *Amanita phalloides*.

► LD₅₀ (mus, ipr) 0.2 mg/kg. NJ8323000

ϵ -Amanitin

1-L-Aspartic acid-3-(S)-4-hydroxy-L-isoleucine- α -amanitin, 9CI

[21705-02-2]

C₃₉H₅₃N₉O₁₄S 903.966

► LD₅₀ (mus, ipr) 0.3 mg/kg. NJ8325000

Amanullinic acid

1-L-Aspartic acid-3-isoleucine- α -amanitin, 9CI

[54532-45-5]

C₃₉H₅₃N₉O₁₃S 887.966

Constit. of *Amanita phalloides*.

► LD₅₀ (mus, ipr) >20 mg/kg. BD6093000

Amanullin [21803-57-6]

C₃₉H₅₄N₁₀O₁₂S 886.981

Constit. of *Amanita phalloides*.

► LD₅₀ (mus, ipr) >20 mg/kg. BD6200000

Deoxy: Proamanullin. 2-L-Proline-3-isoleucine- α -amanitin

[54532-46-6]

C₃₉H₅₄N₁₀O₁₁S 870.982

Constit. of *Amanita phalloides*. Lacks the 4-hydroxypropyl OH group.

► LD₅₀ >20mg/Kg (mus, ipr). BD6300000

[11030-71-0]

Wieland, T. et al., *Prog. Chem. Org. Nat. Prod.*, 1967, **25**, 214 (rev)

Wieland, T. et al., *Annalen*, 1974, 1561; 1570; 1580; 1587 (bibl, abs config)

Yocum, R.R. et al., *Biochemistry*, 1978, **17**, 3786; 3790 (isol, purifn, cryst struct)

Wieland, T. et al., *Crit. Rev. Biochem.*, 1978, **5**, 185 (rev)

Buku, A. et al., *Experientia*, 1980, **36**, 33-34 (Amaninamide)

Wieland, T. et al., *Peptides of Poisonous Amanita Mushrooms*, Springer-Verlag, 1986, (book)

Michelot, D. et al., *Drug Metab. Drug Interact.*, 1988, **6**, 265 (biol props)

Shoham, G. et al., *J.A.C.S.*, 1989, **111**, 4791 (cryst struct)

Pehk, T. et al., *Magn. Reson. Chem.*, 1989, **27**, 173 (pmr, conformn)

Antkowiak, R. et al., *Alkaloids (Academic Press)*, 1991, **40**, 189 (rev)


Tagliaro, F. et al., *J. Chromatogr.*, 1991, **563**, 299 (hplc)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1390

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AH1625

Amarbellisine

[767356-23-0]




C₁₇H₁₉NO₄ 301.341

Alkaloid from the bulbs of *Amaryllis belladonna*. Needles (MeOH). Mp > 300°. $[\alpha]_D^{25}$ -39.2 (c, 0.7 in CHCl₃). λ_{\max} 244 (log ϵ 2.9); 293 (log ϵ 2.9) (MeOH).

Evidente, A. et al., *Phytochemistry*, 2004, **65**, 2113-2118 (isol, cd, pmr, cmr, ms)

Amaryllisine

[6874-70-0]



C₁₈H₂₃NO₄ 317.384

Alkaloid from *Amaryllis belladonna* (*Brunsvigia rosea*) (Amaryllidaceae). Prisms (MeOH). Mp 255-258° dec. $[\alpha]_D^{24}$ +2.4. $[\alpha]_{436}^{24}$ -6.6 (c, 0.27 in CHCl₃).

Ac: Mp 181-182°.


Me ether:

Fine prisms (cyclohexane). Mp 99-100°. $[\alpha]_D^{24}$ -16.4 (c, 1.6 in CHCl₃).

Burlingame, A.L. et al., *J.A.C.S.*, 1964, **86**, 4976 (isol, uv, ir, pmr, ms, struct)

Amataine

Grandifoline†. *Subsessiline*†
[31148-60-4]



C₄₃H₄₈N₄O₆ 716.875

Alkaloid from the root bark of *Hedranthera* (*Callichilia*) *barteri* and *Voacanga cholutiana*, the stems of

Callichilia subsessilis and the leaves of *Voacanga grandifolia* (Apocynaceae). Shows antimicrobial activity. Cryst. (hexane). Mp 216-221°. $[\alpha]_D^{26}$ -262 (c, 0.727 in CHCl₃).

18-Oxo: Subsessiline lactone. 18-Oxoamataine. Alkaloid H†
[59796-72-4]

C₄₃H₄₆N₄O₇ 730.859

Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Cryst.

(Me₂CO). Mp 237-239° dec. $[\alpha]_D$ -292.

Agwada, V. et al., *Helv. Chim. Acta*, 1970, **53**, 1567; 1977, **60**, 2830 (isol, uv, ir, pmr, ms, ord, struct)

Rolland, Y. et al., *Phytochemistry*, 1973, **12**, 2039 (deriv)

Gabetta, B. et al., *Fitoterapia*, 1974, **45**, 32; *CA*, **81**, 166291t (isol)


Rolland, Y. et al., *J.O.C.*, 1976, **41**, 3270 (cmr, struct, deriv)

Rolland, Y. et al., *Helv. Chim. Acta*, 1977, **60**, 2854 (cd, ms, deriv)

Villar, A. et al., *Planta Med.*, 1986, **52**, 556-557 (activity)

Amathamide A

[99615-75-5]



C₁₅H₁₈Br₂N₂O₂ 418.127

Alkaloid from the marine bryozoan *Amathia wilsoni*. Mp 189-190.5°.

(Z)-Isomer: *Amathamide B*


[99615-76-6]

Alkaloid from *Amathia wilsoni*. Oil.

Blackman, A.J. et al., *Heterocycles*, 1985, **23**, 2829-2833 (isol, ir, pmr, cmr, ms, cd, struct, abs config)

Amathamide E

[112515-28-3]



C₁₅H₁₇Br₃N₂O₂ 497.023

Alkaloid from the bryozoan *Amathia wilsoni*. Light yellow oil.

N-Me: *Amathamide C*

[112515-26-1]

C₁₆H₁₉Br₃N₂O₂ 511.05

Alkaloid from *Amathia wilsoni* and *Amathia pinnata*. Pale yellow oil.

Dihydro: *Amathamide D*

[112515-27-2]

C₁₅H₁₉Br₃N₂O₂ 499.039

Alkaloid from *Amathia wilsoni*. Light green oil.

(Z)-Isomer: *Amathamide F*

[112515-29-4]


C₁₅H₁₇Br₃N₂O₂ 497.023Alkaloid from *Amathia wilsoni*. Oil.**N-Me, 15-methoxy: Amathamide G**

[149355-72-6]

C₁₇H₂₁Br₃N₂O₃ 541.076Alkaloid from *Amathia convoluta*. Oil.Blackman, A.J. *et al.*, *Aust. J. Chem.*, 1987, **40**, 1655-1662 (*isol, ir, pmr, cmr, ms, struct*)Blackman, A.J. *et al.*, *Aust. J. Chem.*, 1993, **46**, 401-405 (*Amathamide G*)**Amathaspiramide C**

A-679

[226993-86-8]

C₁₅H₁₈Br₂N₂O₃ 434.127Alkaloid from the marine bryozoan *Amathia wilsoni*. Amorph. solid. [α]_D²⁵ -2 (c, 0.005 in MeOH). λ_{max} 215 (log ε 4.16); 289 (log ε 3.15); 294 (log ε 3.15) (MeOH).**N¹-Me: Amathaspiramide A**

[226993-84-6]

C₁₆H₂₀Br₂N₂O₃ 448.154Alkaloid from *Amathia wilsoni*. Amorph. solid. [α]_D²⁵ -3 (c, 0.004 in MeOH). λ_{max} 216 (log ε 4.2); 289 (log ε 3.34); 295 (log ε 3.36) (MeOH).**1,2-Didehydro: Amathaspiramide E**

[226993-88-0]

C₁₅H₁₆Br₂N₂O₃ 432.111Alkaloid from *Amathia wilsoni*. Amorph. solid. [α]_D²⁵ -21 (c, 0.002 in MeOH). λ_{max} 220 (log ε 4.13); 289 (log ε 3.17); 295 (log ε 3.17) (MeOH).**2-Oxo: Amathaspiramide D**

[226993-87-9]

C₁₅H₁₆Br₂N₂O₄ 448.11Alkaloid from *Amathia wilsoni*. Amorph. solid. [α]_D²⁵ -44 (c, 0.002 in MeOH). λ_{max} 215 (log ε 4.03); 289 (log ε 3.22); 295 (log ε 3.22) (MeOH).**2-Oxo, N¹-Me: Amathaspiramide B**

[226993-85-7]

C₁₆H₁₈Br₂N₂O₄ 462.137Alkaloid from *Amathia wilsoni*. Amorph. solid. [α]_D²⁵ -13 (c, 0.002 in MeOH). λ_{max} 208 (log ε 4.5); 310 (log ε 4.38); 436 (log ε 4.6) (MeOH).**8-Epimer: Amathaspiramide F**

[226993-89-1]


C₁₅H₁₈Br₂N₂O₃ 434.127Alkaloid from *Amathia wilsoni*. Amorph. solid. [α]_D -10 (c, 0.002 in MeOH). λ_{max} 214 (log ε 4.12); 290 (log ε 3.08); 296 (log ε 3.08) (MeOH).Morris, B.D. *et al.*, *J. Nat. Prod.*, 1999, **62**, 688-693 (*isol, uv, ir, pmr, cmr, ms*)Hughes, C.C. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 4556-4559 (*Amathaspiramide F, synth*)Sakaguchi, K. *et al.*, *Org. Lett.*, 2008, **10**, 5449-5452 (*Amathaspiramide F, synth*)**Amauromine**

A-680

FR 900220. WF 6237. Antibiotic FR

900220. Antibiotic WF 6237

[88360-87-6]



Absolute Configuration

C₃₂H₃₆N₄O₂ 508.662Diketopiperazine antibiotic. Isol. from *Amauroascus* sp. ATCC 20595 and *Penicillium nigricans* IMI 228669. Hypotensive vasodilator; gastric juice secretion inhibitor, calcium channel blocker.Prisms (EtOH). Sol. MeOH, Me₂CO, Py, CHCl₃; poorly sol. H₂O, hexane. Mp 156-158°. [α]_D²³ -583 (c, 1 in CHCl₃). λ_{max} 245 (ε 11000); 300 (ε 4200) (EtOH) (Derep). ▶ LD₅₀ (mus, ipr) 200 mg/kg. UQ3987000**2,3-Diepimer: Epiaumuromine**

[143168-22-3]

C₃₂H₃₆N₄O₂ 508.662Metab. from sclerotia of *Aspergillus ochraceus* NRRL 3519. Insect antifeedant, insecticide. Solid. Mp 134°. [α]_D -50 (c, 0.18 in CHCl₃). λ_{max} 215 (ε 100000); 243 (ε 100000); 300 (ε 40000) (MeOH) (Berdy).**2,3-Diepimer, N¹-Me: N-Methylepiaumuromine**

[143086-28-6]

C₃₃H₃₈N₄O₂ 522.689Metab. from the sclerotia of *Aspergillus ochraceus* NRRL 3519. Insect antifeedant, insecticide. [α]_D -29.1 (c, 0.46 in CHCl₃). λ_{max} 214 (ε 25000); 244 (ε 12500); 303 (ε 6400) (MeOH) (Berdy).**Stereoisomer: Nigrifortine**

[97859-13-7]

C₃₂H₃₆N₄O₂ 508.662Metab. of *Penicillium nigricans*. Stereochem. not determined. May be identical with Amauromine or Epiaumuromine.Takase, S. *et al.*, *J. Antibiot.*, 1984, **37**, 1320-1323 (*isol, props*)Laws, I. *et al.*, *Phytochemistry*, 1985, **24**, 1395-1397 (*Nigrifortine*)Takase, S. *et al.*, *Tetrahedron*, 1985, **41**, 3037-3048; 1986, **42**, 5887-5894 (*cryst struct, synth*)De Guzman, F.S. *et al.*, *J. Nat. Prod.*, 1992, **55**, 931-939 (*Epiaumuromine, N-Methylepiaumuromine*)Marsden, S.P. *et al.*, *J.A.C.S.*, 1994, **116**, 11143-11144 (*synth*)Depew, K.M. *et al.*, *J.A.C.S.*, 1999, **121**, 11953-11963 (*synth*)**Ambaline**

A-681


C₃₈H₄₂N₂O₁₀ 686.757Struct. unknown. Alkaloid from *Pycnarhena manillensis*. Mp 123° Mp 265° (as dihydrochloride). [α]_D +143 (CHCl₃). Nonphenolic.Quibilan, G.Q. *et al.*, *CA*, 1934, **28**, 4735² (*isol*)**Ambalinine**

A-682

C₁₈H₂₁NO₃ 299.369Struct. unknown. Nonphenolic, conts. 1 N-Me and 1 OMe group. Alkaloid from *Pycnarhena manillensis*. Mp 203-204° Mp 238° (as picrate).Villanos, M.I. *et al.*, *CA*, 1936, **30**, 4171⁷ (*isol*)**Ambelline**

A-683

[3660-62-6]



Absolute Configuration

C₁₈H₂₁NO₅ 331.368C-12 config. incorr. given as 12S- in CAS. Alkaloid of *Brunsvigia rosea* (*Amaryllis belladonna*) and many other spp. in the Amaryllidaceae. Cytotoxic to P-388 sarcoma cells *in vitro*, shows no activity *in vivo*. Weak transient hypotensive agent and cardiac stimulant. Epinephrine potentiator, respiratory paralytic. Sol. MeOH, CHCl₃, Mp 260-261° (254-256°). [α]_D²² +81 (c, 0.56 in CHCl₃). [α]_D²⁵ -13.6 (c, 0.11 in MeOH). λ_{max} 213 (log ε 4.6); 244 (sh) (log ε 3.64); 286 (log ε 3.24) (MeOH).▶ Toxic, LD₅₀ (mus, scu) 5 mg/kg.*Hydrochloride*: Mp 227-230° dec.*Perchlorate*: Mp 200° dec.**Ac: 11-O-Acetylabelline**

[98983-14-3]

C₂₀H₂₃NO₆ 373.405Alkaloid from bulbs of *Brunsvigia josephinae* (Amaryllidaceae). Mp 80-82°. [α]_D²² -23.5 (c, 0.59 in CHCl₃).**11-O-(3-Pyridinecarbonyl): Filifoline†.****11-O-Nicotinoylabelline**C₂₄H₂₄N₂O₆ 436.463Alkaloid from the bulbs of *Nerine filifolia*. Powder. Mp 191-193°. [α]_D²⁰ +12 (c, 0.85 in MeOH). λ_{max} 280 (log ε 5.93) (EtOH).**1β,2β-Epoxyde: 1,2-Epoxyambelline**

[93771-88-1]

C₁₈H₂₁NO₆ 347.367Alkaloid from *Crinum latifolium* (Amaryllidaceae). Prisms (Me₂CO). Mp 245-246°. [α]_D²² +22.4 (c, 0.32 in CHCl₃).**1β,2β-Epoxyde, Ac: 11-O-Acetyl-1,2-epoxyambelline**

[93771-89-2]

C₂₀H₂₃NO₇ 389.404Alkaloid from *Ammocharis tinneana*. Needles (Me₂CO). Mp 201-203° (195-197°).**Dihydro: Dihydroambelline**C₁₈H₂₃NO₅ 333.383Isol. from *Nerine crispa* (Amaryllidaceae). Mp 198-199°. [α]_D²⁴ -12.6 (c, 0.5 in CHCl₃).**6α-Hydroxy: Crinafoline**

[106534-56-9]

C₁₈H₂₁NO₆ 347.367

Alkaloid from the mature fruits of *Crinum latifolium* (Amaryllidaceae). Has antitumour props. Microcryst. (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 234-237°. [α]_D²⁸ +44.7 (c, 0.35 in MeOH). 12-Config. incorrectly given as 12*S*- in CAS.

6 α -Hydroxy, di-Ac:

Prisms (MeOH/Me₂CO). Mp 214-215°.

6 α -Hydroxy, N-Me:

C₁₉H₂₄NO₅⁺ 346.402
Straw-coloured solid (as iodide). Mp 262-265° dec.(iodide).

3-Epimer: Brunsbelline

[171119-08-7]
C₁₈H₂₁NO₅ 331.368
Alkaloid from the bulbs of *Brunsvigia josephinae*. Mp 226-228°. [α]_D²² -97.8 (c, 0.1 in EtOH).

Stereoisomer, 1,2-epoxide: Cavinine


[58189-38-1]
C₁₈H₂₁NO₆ 347.367
Alkaloid from *Hippeastrum punicum* (Amaryllidaceae). Identified by ms only and assigned the same plane struct. as 1,2-Epoxyambelline without assignment of stereochem. May be identical with 1,2-Epoxyambelline.

- Mason, L.H. *et al.*, *J.A.C.S.*, 1955, **77**, 1253-1256 (*isol, uv, ir*)
Döpke, W. *et al.*, *Naturwissenschaften*, 1962, **49**, 469 (*Dihydroambelline*)
Naegeli, P. *et al.*, *J.O.C.*, 1963, **28**, 206-210 (*ir, pmr, struct*)
Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902-4912 (*ms*)
Samuel, E.H.S. *et al.*, *Org. Mass Spectrom.*, 1975, **10**, 427-431 (*Cavinine*)
Roques, R. *et al.*, *Acta Cryst. B*, 1976, **32**, 1394-1398 (*cryst struct*)
Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1984, 232-233; 1986, 312-313 (*Epoxyambelline, Crinafoline*)
Viladomat, F. *et al.*, *Phytochemistry*, 1994, **35**, 809-812 (*isol, cd, pmr, cmr, ms*)
Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 961-965 (*11-O-Acetylabelline, Brunsbelline*)
Pham, L.H. *et al.*, *Phytochemistry*, 1998, **48**, 371-376 (*isol, cd, uv, pmr, cmr, ms*)
Machocho, A. *et al.*, *Phytochemistry*, 1999, **51**, 1185-1191 (*Epoxyambelline, O-Acetylepoxymbelline*)
Nair, J.J. *et al.*, *Phytochemistry*, 2005, **66**, 373-382 (*Filifoline*)

Ambiguine B isonitrile

A-684

A 89271C. Antibiotic A 89271C [138630-60-1]



Relative Configuration

C₂₆H₃₁ClN₂O 422.996

Alkaloid from the terrestrial blue-green alga *Fischerella ambigua*, also from *Hapalosiphon hibernicus* and *Westrellopsis prolifica*. Amorph. solid. [α]_D -44.3 (c, 0.1 in MeOH). λ_{\max} 223 (€ 40700); 281 (€ 9400); 291 (€ 7700) (MeOH).

Deoxy: Ambiguine A isonitrile. A 89271B.

Antibiotic A 89271B [138630-59-8]

C₂₆H₃₁ClN₂ 406.997
From the terrestrial blue-green algae *Fischerella ambigua* and *Hapalosiphon hibernicus*, also *Westrellopsis prolifica*. Needles (hexane/CH₂Cl₂). Mp > 300° dec. [α]_D -37 (c, 0.1 in MeOH). λ_{\max} 225 (€ 32400); 282 (€ 7600); 295 (sh) (€ 6500) (MeOH).

Dechloro: Ambiguine C isonitrile. A

89271E. Antibiotic A 89271E [138630-61-2]

C₂₆H₃₂N₂O 388.552
From *Fischerella ambigua*. Amorph. solid. [α]_D -9.5 (c, 0.1 in MeOH). λ_{\max} 223 (€ 36000); 281 (€ 8400); 291 (€ 6800) (MeOH).

Dechloro, deoxy: Ambiguine H isonitrile


C₂₆H₃₂N₂ 372.552
From a *Fischerella* sp. Amorph. solid. [α]_D²⁵ -65 (c, 0.51 in MeOH). λ_{\max} 222 (log € 4.38); 282 (log € 3.72); 291 (log € 3.62) (MeOH).

Smitka, T.A. *et al.*, *J.O.C.*, 1992, **57**, 857-861 (*isol, cd, uv, ir, pmr, cmr*)
Raveh, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 196-201 (*isol, pmr, cmr*)

Ambiguine D isonitrile

A-685

A 89271A. Antibiotic A 89271A [138666-15-6]



Relative Configuration

C₂₆H₂₉ClN₂O₃ 452.979

Alkaloid from the terrestrial blue-green algae *Fischerella ambigua* and *Westrellopsis prolifica*. Cryst. (MeOH aq.). Mp > 300° dec. [α]_D -30.3 (c, 0.1 in MeOH). λ_{\max} 227 (€ 13900); 301 (€ 2170) (MeOH).

Dechloro: Ambiguine J isonitrile


C₂₆H₃₀N₂O₃ 418.535
From a *Fischerella* sp. Amorph. solid. [α]_D²⁵ -32 (c, 0.1 in MeOH). λ_{\max} 222 (log € 3.68); 270 (log € 2.97); 308 (log € 2.78) (MeOH).

Smitka, T.A. *et al.*, *J.O.C.*, 1992, **57**, 857-861 (*isol, cd, uv, ir, pmr, cmr, cryst struct*)
Raveh, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 196-201 (*isol, pmr, cmr*)

Ambiguine E isonitrile

A-686

A 89271D. Antibiotic A 89271D [138666-16-7]



Relative Configuration

C₂₆H₂₉ClN₂O₂ 436.98

Alkaloid from the terrestrial blue-green algae *Fischerella ambigua*, *Hapalosiphon hibernicus* and *Westrellopsis prolifica*. Needles (hexane/CH₂Cl₂). Mp > 300° dec. [α]_D -59.7 (c, 0.1 in MeOH). λ_{\max} 223 (€ 36000); 272 (€ 7100); 279 (€ 6800) (MeOH).

Dechloro: Ambiguine I isonitrile


C₂₆H₃₀N₂O₂ 402.535
From a *Fischerella* sp. Amorph. solid. [α]_D²⁵ -39 (c, 0.29 in MeOH). λ_{\max} 222 (log € 4.04); 282 (log € 3.69); 286 (log € 3.67) (MeOH).

Smitka, T.A. *et al.*, *J.O.C.*, 1992, **57**, 857-861 (*isol, cd, uv, ir, pmr, cmr*)
Raveh, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 196-201 (*isol, pmr, cmr*)

Ambiguine F isonitrile

A-687

A 89271F. Antibiotic A 89271F [138630-62-3]



Relative Configuration

C₂₆H₃₁ClN₂O₃ 454.995


Alkaloid from the terrestrial blue-green alga *Fischerella ambigua*. Amorph. solid. [α]_D -18.2 (c, 0.1 in MeOH). λ_{\max} 224 (€ 36000); 280 (€ 7200); 290 (€ 5500) (MeOH).

Smitka, T.A. *et al.*, *J.O.C.*, 1992, **57**, 857 (*isol, uv, ir, pmr, cmr, cd, struct*)

Ambiguine G nitrile

A-688

[213911-41-2]




C₂₆H₂₇CIN₂ 402.965
Isol. from the blue-green alga *Hapalosiphon delicatulus*. [α]_D²⁵ +138.3 (c, 0.6 in CHCl₃). λ_{max} 230 (ε 32400); 274 (ε 13900); 390 (ε 6750) (MeOH).

Huber, U et al., *J. Nat. Prod.*, 1998, **61**, 1304-1306 (isol, uv, ir, pmr, cmr, ms)

Ambinine

A-689

[96935-26-1]



Absolute Configuration


C₂₃H₂₇NO₆ 413.469
Minor alkaloid from the stem tuber of *Corydalis ambigua* (Papaveraceae). Prisms (EtOH). Mp 218-219°. [α]_D²² -106 (c, 1.3 in CHCl₃).

Hydrobromide: [132296-33-4]
Fine needles (EtOH). Mp 199-202°.
Cui, Z. et al., *Yaoxue Xuebao*, 1984, **19**, 904; *CA*, **103**, 19853x (isol, uv, ir, pmr, cmr, ms, struct)
Cui, Z. et al., *J. Nat. Prod.*, 1990, **53**, 1182 (abs config, cd, cryst struct)
Hanaoka, M. et al., *Chem. Pharm. Bull.*, 1991, **39**, 242 (synth)

Americine

A-690

N-[7-(1*H*-Indol-3-ylmethyl)-3-(1-methyl-ethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-3-methyl-2-(methylamino)butanamide, 9*CI*
[18867-84-0]



Absolute Configuration

C₃₁H₃₉N₅O₄ 545.68
Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) (Rhamnaceae). Mp 135.5-137°. [α]_D²⁰ -198 (c, 0.51 in MeOH).

N-Me: *N-Methylamericine*. 5-β-Indolylmethyl-8*N*-(*N,N*-dimethylvalyl)-9-iso-propylphenylcyclopeptide
[70403-89-3]

C₃₂H₄₁N₅O₄ 559.707
Alkaloid from *Ceanothus integerrimus* and *Ceanothus sanguineus*. Mp 233° (229°).

Klein, F.K. et al., *J.A.C.S.*, 1968, **90**, 2398-2404 (isol, uv, ir, pmr, ms, struct)
Lagarias, J.C. et al., *J. Nat. Prod.*, 1979, **42**, 220-227; 663-668 (*N-Me*)

Amianthine

A-691

C₂₇H₄₁NO₂ 411.626
Steroidal alkaloid. Struct. unknown. Alkaloid from *Amianthium muscaetoxicum* (*Zygadenus muscaetoxicum*) (Liliaceae). Long prisms (Me₂CO). Mp 251-253° dec. [α]_D²⁰ -87 (c, 0.173 in CHCl₃). p*K*_a 9.7.


Tertiary base contg. conjugated CO and acylable OH gps.

Ac: Mp 206-207° dec.
Neuss, N. et al., *J.A.C.S.*, 1953, **75**, 2772-2773 (isol)

Amidinomycin

A-692

N-(2-*Amidinoethyl*)-3-aminocyclopentanecarboxamide. *Myxoviomycin*
[3572-60-9]



C₉H₁₈N₄O 198.267
Prod. by *Streptomyces flavochromogenes*. Antifungal antiviral agent. Antineoplastic antibiotic. Sol. H₂O, EtOH, MeOH; poorly sol. butanol, hexane. Log P -2.26 (uncertain value) (calc). λ_{max} 205 (ε 4460) (0.1*N* HCl) (Derep). λ_{max} 219 (ε 1580) (0.1*N* NaOH) (Derep). λ_{max} 212 (ε 1170) (pH 3 H₂O) (Derep). λ_{max} 212 (E1%/1cm 59) (H₂O) (Berdy). λ_{max} 205 (E1%/1cm 225) (HCl) (Berdy). λ_{max} 219 (E1%/1cm 79) (NaOH) (Berdy).

► LD₅₀ (mus, orl) 140 mg/kg, LD₅₀ (mus, ivn) 18 mg/kg, LD₅₀ (mus, ipr) 200 mg/kg. GY2555000

Sulfate:
Plates or needles (MeOH aq.). Mp 285-288° dec. [α]_D²¹ -3.9 (c, 3 in H₂O).
[19946-52-2]

Nakamura, S. et al., *Chem. Pharm. Bull.*, 1961, **9**, 641 (synth, struct)
Nakamura, S. et al., *J. Antibiot., Ser. A*, 1961, **14**, 103 (isol, struct)
Katsube, J. et al., *Chem. Pharm. Bull.*, 1968, **16**, 232 (synth, ir, pmr)
Allan, R.D. et al., *Aust. J. Chem.*, 1979, **32**, 2517 (synth, struct)
Kameda, M. et al., *J. Antibiot.*, 1980, **33**, 778 (cryst struct, abs config)
Sapse, A.M. et al., *J. Biomol. Struct. Dyn.*, 1993, **10**, 709 (conformn)
Chenevert, R. et al., *Chem. Lett.*, 1994, 93 (synth, abs config)

2-Amino-12 methyl-1,3,12-tetradecanetriol

A-693

H₃CCH₂C(OH)(CH₃)(CH₂)₈-CH(OH)CH(NH₂)CH₂OH
C₁₅H₃₃NO₃ 275.431

N-Hydroxy, N-nitroso, 2-(Hydroxynitrosoamino)-12-methyl-1,3,12-tetradecanetriol, 9CI, Nitrosostromelin
[182064-61-5]

C₁₅H₃₂N₂O₅ 320.428
Prod. by a *Streptomyces* sp. Stromelysin inhibitor.
U.K. Pat., 1996, 2 297 324; *CA*, **125**, 245813v (isol)

2-Aminoacetophenone, 8CI

A-694

2-Amino-1-phenylethanone, 9*CI*. Phenacylamine. Phenomydrol. Benzoylmethylamine. (Aminoacetyl)benzene
[613-89-8]
PhCOCH₂NH₂

C₈H₉NO 135.165
Component of tortilla aroma and of other corn flour products. Constit. of *Castanopsis cuspidata* and *Vitis* sp. Aroma threshold 0.2 ppb in H₂O.

► Exp. reprod. effects. AM5775000
Hydrochloride: [5468-37-1]
Needles (Me₂CO/EtOH). Mp 188°.
► AM5940000

Oxime:
C₈H₁₀N₂O 150.18
Mp 140°.

2,4-Dinitrophenylhydrazone: Mp 221°.

Di-Et ketal:
C₁₂H₁₉NO₂ 209.288
Liq. Bp_{0.5} 68°.

N-Formyl: [73286-37-0]
C₉H₉NO₂ 163.176
Cryst. (Me₂CO/hexane). Mp 80-82°.

N-Ac: [1846-33-9]
C₁₀H₁₁NO₂ 177.202
Mp 85°.

N-Benzoyl: 2-(Benzoylamino)acetophenone. *N-Phenacylbenzamide, Muricatinone*
[4190-14-1]
C₁₅H₁₃NO₂ 239.273
Alkaloid from *Oxytropis muricata* and *Oxytropis puberula*. Cryst. (EtOH). Mp 122-124°.

N-Me: α-Methylaminoacetophenone
[35534-19-1]
C₉H₁₁NO 149.192
Yellow oil.

N-Me, hydrochloride: [23826-47-3]
Plates. Mp 219° dec.

N-Me, N-Ac:
C₁₁H₁₃NO₂ 191.229
Mp 156°.

N,N-Di-Me: 2-(*N,N*-Dimethylamino)acetophenone
[3319-03-7]
C₁₀H₁₃NO 163.219
Isol. from *Desmodium gangeticum*. Liq. Bp₂₀ 130-132°.

[25384-14-9]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 46D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 867A (nmr)
Tiffeneau, M. et al., *Bull. Soc. Chim. Fr.*, 1931, **49**, 1761 (rev)

Stalham, F.S. et al., *J.C.S.*, 1951, 213 (synth)
Chapman, N.B. et al., *J.C.S.*, 1963, 1385 (synth, deriv)

Tanaka, H. et al., *CA*, 1971, **75**, 147849 (struct)
Ghosal, S. et al., *Planta Med.*, 1972, **22**, 434 (isol, deriv)

Bellamy, L.J. et al., *Spectrochim. Acta A*, 1972, **28**, 1869 (ir)
Yamaguchi, K. et al., *J. Agric. Food Chem.*, 1979, **27**, 847 (isol)


Muchowski, J.M. et al., *J.O.C.*, 1986, **51**, 3374 (*N-formyl*)
Abdalla, G.M. et al., *J. Het. Chem.*, 1987, **24**, 297 (synth, ir, pmr)

Yamagama, C. et al., *Aust. J. Chem.*, 1989, **42**, 463 (*N-15 nmr*)
Baumstark, A.L. et al., *Chem. Comm.*, 1989, 767 (*O-17 nmr*)

Buttery, R.G. et al., *J. Agric. Food Chem.*, 1994, **42**, 1-2 (occur, anal)

Shure, K.B. *et al.*, *J. Agric. Food Chem.*, 1994, **42**, 350 (*isol*)
 Demeuov, N.B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1998, **34**, 484-485 (*Muricatisine*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AHR250

α -Amino-5-[[[(2-aminocarbonyl-1-carboxylethyl)amino]carbonyl]-2-pyrrolopropanoic acid, 9CI A-695




$C_{12}H_{16}N_4O_6$ 312.282
 Minor metab. of *Pantoea agglomerans* (formerly *Erwinia herbicola*). λ_{max} 200 (ϵ 6440); 220 (sh) (ϵ 2994); 275 (ϵ 10890) (H_2O).

Jin, M. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2898-2901; 2902-2905 (*isol, pmr, cmr, ms*)

2-Amino-6-(2-aminoethyl)-3-(3,4-dihydroxyphenyl)-1,4-benzodioxan A-696

2-Amino-3-(3,4-dihydroxyphenyl)-2,3-dihydro-1,4-benzodioxin-6-ethanamine



$C_{16}H_{18}N_2O_4$ 302.329
 Dimer of Dopamine, D-920.


(2S,3R)-form

N,N' -Ac:
 $C_{20}H_{22}N_2O_6$ 386.404
 Isol. from the cast-off shells of the cicada *Cryptotympana* sp. Component of Zentai. Yellow powder. Mp 115-120°. $[\alpha]_D^{25} +28.2$ (c, 1 in MeOH).

1',2'-Didehydro(E-), N,N' -di-Ac:
 $C_{20}H_{20}N_2O_6$ 384.388
 Isol. from the cast-off shells of *Cryptotympana pustulata*. Yellow powder. $[\alpha]_D^{25} +33.8$ (c, 0.1 in MeOH). λ_{max} 282 (log ϵ 4.32) (MeOH).

Noda, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1749-1752 (*isol, pmr, cmr*)
 Xu, M.-Z. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 7826-7834 (*deriv*)

3-Amino-1-(2-amino-3-hydroxyphenyl)-1-propanone A-697
 2',3-Diamino-3'-hydroxypropiophenone. 2-Amino-3-(3-aminopropanoyl)phenol. 3-Hydroxykynuramine




$C_9H_{12}N_2O_2$ 180.206
 N^3 -Ac: **Erebisinone**. *Eribusinone* [325477-44-9] [194615-13-9]

$C_{11}H_{14}N_2O_3$ 222.243
 Isol. from the sponge *Isodictya erinacea*. Yellow solid. λ_{max} 238 (ϵ 1240); 372 (ϵ 180) (MeOH).

Moon, B. *et al.*, *Tetrahedron*, 2000, **56**, 9057-9062 (*Erebisinone*)

2-Amino-3-(3-amino-1H-inden-2-yl)propanoic acid A-698

$\alpha,3$ -Diamino-1H-indene-2-propanoic acid, 9CI. 3-(3-Amino-1H-inden-2-yl)alanine. **Cycasindene** [192801-71-1]




$C_{12}H_{14}N_2O_2$ 218.255
 Constit. of the seeds of *Cycas revoluta* (Cycadaceae). Microcryst.

Pan, M. *et al.*, *Phytochemistry*, 1997, **45**, 517-519 (*isol, pmr, cmr*)

2-Amino-5-(aminomethyl)-pyrrolo[2,3-d]pyrimidin-4(3H)-one, 9CI A-699

7-Aminomethyl-7-deazaguanine. Pre-Q1 base [69251-45-2]




$C_7H_9N_5O$ 179.181
 Isol. from *Escherichia coli* tRNA. Biosynthetic precursor of Queuine, Q-9. Solid; cryst. + 2H₂O (as hydrochloride). Mp 220-225° dec. (hydrochloride).

Ohgi, T. *et al.*, *Chem. Lett.*, 1979, 1283 (*synth, ir, ms, pmr, uv, derivs*)
 Okada, N. *et al.*, *J. Biol. Chem.*, 1979, **254**, 3067 (*isol, synth*)
 Farkas, W.R. *et al.*, *Biochim. Biophys. Acta*, 1984, **781**, 64 (*metab*)
 Akimoto, H. *et al.*, *J.C.S. Perkin 1*, 1988, 1637 (*synth, ir, pmr, bibl*)
 Klepper, F. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 2610-2616 (*synth, ir, pmr, cryst struct*)

3-Amino-1-(2-aminophenyl)-1-propanone, 9CI A-700

2',3-Diaminopropiophenone. **Kynuramine** [363-36-0]



$C_9H_{12}N_2O$ 164.207
 Found in rat brains, and derived from


tryptophan. Liq.
 Hydrochloride: Mp 185-187° dec.
 Picrate: Mp 176-177° dec.

N^3 -Ac: N^3 -Acetylkynuramine
 $C_{11}H_{14}N_2O_2$ 206.244
 Prod. by the marine-derived *Janibacter limosus* Hel 1. Light yellow oil. λ_{max} 202 (log ϵ 4.21); 227 (log ϵ 4.31); 362 (log ϵ 3.61) (MeOH). λ_{max} 208 (log ϵ 4.3); 227 (log ϵ 4.31); 255 (log ϵ 3.77); 360 (log ϵ 3.68) (MeOH/NaOH).

Butenandt, A. *et al.*, *Z. Naturforsch., B*, 1953, **86**, 454 (*synth, pharmacol*)
 Seiler, N. *et al.*, *CA*, 1971, **74**, 39005 (*ms*)
 Gal, E.M. *et al.*, *J. Neurochem.*, 1978, **30**, 607 (*occur*)
 Asolkar, R.N. *et al.*, *J. Antibiot.*, 2004, **57**, 17-23 (N^3 -Acetylkynuramine)

2-Amino-4-(3-amino-1-propenyl)-1H-imidazole A-701

4-(3-Amino-1-propenyl)-1H-imidazol-2-amine, 9CI. 3-(2-Amino-1H-imidazol-4-yl)-2-propen-1-amine. 3-Amino-1-(2-aminoimidazolyl)-1-propene




$C_6H_{10}N_4$ 138.172

(E)-form [140381-65-3]

Alkaloid from *Agelas oroides*, *Ptilocaulis walpersi* and *Teichaxinella morchella*. Light brown oil. λ_{max} 276 (ϵ 8790) (MeOH) (Derep).

Wright, A.E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1684-1686 (*isol, uv, pmr, cmr*)
 Tasdemir, D. *et al.*, *Bioorg. Med. Chem.*, 2008, **15**, 6834-6845 (*isol, pmr, cmr*)

3-Aminoandrost-4-en-17-ol A-702



$C_{19}H_{31}NO$ 289.46

(3 α ,17 β)-form

N-Carbamoyl: 3-Ureidoandrost-4-en-17-ol
 $C_{20}H_{32}N_2O_2$ 332.485
 Alkaloid from the musk of *Moschus moschiferus*. Amorph. powder. Mp 210-212°. $[\alpha]_D^{25} +250$ (c, 0.02 in MeOH). Dec. at 215°.


17-Ketone, N-carbamoyl: 3-Ureidoandrost-4-en-17-one
 $C_{20}H_{30}N_2O_2$ 330.469
 Alkaloid from the musk of *Moschus moschiferus*. Amorph. powder. Mp

220-223°. $[\alpha]_D^{25} +115$ (c, 0.002 in MeOH).

Oh, S.-R. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 663-664 (*isol.*, *pmr.*, *cmr*)

1-Amino-1(10)-aristolene-2,8,9-trione

A-703



$C_{15}H_{19}NO_3$ 261.32

ent-form

Lepidamine

[609332-21-0]


Prod. by *Russula lepida*. Pale yellow oil. $[\alpha]_D +125$ (c, 0.36 in $CHCl_3$). λ_{max} 242; 409 (Et₂O).

Tan, J. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 307-309 (*isol.*, *pmr.*, *cmr.*, *ms*)

1-Amino-8-azabicyclo[3.2.1]octane-2,3,4-triol

A-704

1-Amino-2,3,4-trihydroxynortropane



$C_7H_{14}N_2O_3$ 174.199

(1S,2R,3R,4S)-form

Calystegine N₁

[177794-03-5]

Alkaloid from *Hyoscyamus niger* (Solanaceae). Shows glycosidase inhibitory activity. $[\alpha]_D +59.4$ (c, 0.2 in H₂O).

Asano, N. *et al.*, *Carbohydr. Res.*, 1996, **284**, 169-178 (*isol.*, *pmr.*, *cmr.*, *activity*)

Bekkouche, K. *et al.*, *Phytochemistry*, 2001, **58**, 455-462 (*ms.*, *occur*)


Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

4-Amino-1,3-benzenediol, 9CI

A-705

4-Aminoresorcinol, 8CI. 2,4-Dihydroxylaniline

[13066-95-0]



$C_6H_7NO_2$ 125.127

Used in hair dyes. pK_{a1} 7.91; pK_{a2} 9.16; pK_{a3} 11.24 (20°). Unstable.

3-Me ether: 4-Amino-3-methoxyphenol [61638-01-5]

$C_7H_9NO_2$ 139.154

Reddish-brown needles (toluene). Mp 175-180°. Blackens at 160°. Aq. solns. discolour in air.

3-Me ether, N-Ac: 4-Hydroxy-2-methoxyacetanilide [5307-06-2]

$C_9H_{11}NO_3$ 181.191

Isol. from a fungal endophyte of spruce needles and from a soil fungus. Also prod. by a marine-derived *Penicillium* sp. Red needles (toluene). Mp 169-171°.

Di-Me ether: 2,4-Dimethoxyaniline

[2735-04-8]

$C_8H_{11}NO_2$ 153.18

Alarm pheromone isol. from *Neamura muscorum*. Plates (petrol) or oil. Mp 32.5-33.5°. Bp_{0.006} 75-80°.

▶ LD₅₀ (rat, orl) 464 mg/kg. BX4200000

Di-Me ether; hydrochloride: [54150-69-5] Mp 230° dec.

[37966-57-7]

Collins, R.F. *et al.*, *J.C.S.(C)*, 1966, 366

(*synth*)

Messer, C. *et al.*, *CA*, 1999, **131**, 71415y (2,4-Dimethoxyaniline, *isol*)


2-Aminobenzoic acid, 9CI

A-706

Anthranilic acid, 8CI. Vitamin L₁

[118-92-3]

[1321-11-5]



$C_7H_7NO_2$ 137.138

Prod. by bacteria and fungi. An important intermed. in the synth. of many compds., corrosion inhibitor for metals. Used as 3% aq. soln. of Na salt for gravimetric detn. of Co, Ni, Cu(II), Hg(II), Mn, Pb, Cd, U(VI), Zn. Inducer of fruiting body formation. Transient receptor potential (TRP) channel blocker. Leaflets. Sol. hot H₂O, EtOH, Et₂O. Mp 144-148°. pK_{a1} 1.97; pK_{a2} 4.79 (25°). pK_{a1} 2.17; pK_{a2} 4.85 (25°, 0.1M KCl). Sublimes, triboluminescent. λ_{max} 218 (ε 20200); 248 (ε 7000); 335 (ε 4700) (MeOH) (Berdy).

▶ LD₅₀ (mus, orl) 1400 mg/kg. Exp. reprod. effects. CB2450000

Hydrochloride: [2099-63-0]

Mp 193-194°.

β-D-Glucopyranosyl ester: β-D-Glucopyranosyl anthranilate

[55798-72-6]

$C_{13}H_{17}NO_7$ 299.28

Constit. of the fruit of piñuela (*Bromelia plumieri*, Bromeliaceae) and of *Arabidopsis thaliana*.

α-L-Rhamnopyranosyl ester: α-L-Rhamnopyranosyl anthranilate

$C_{13}H_{17}NO_6$ 283.28

Prod. by *Actinomyces* sp. Stamm K 17/2. Amorph. solid. $[\alpha]_D^{20} -76$ (c, 0.1 in MeOH). λ_{max} 219 (log ε 4.25); 249 (log ε 3.72); 341 (log ε 3.55) (MeOH).

6-Deoxy-α-L-talopyranosyl ester: 2-Aminobenzoyl 6-deoxy-α-L-talopyranoside

$C_{13}H_{17}NO_6$ 283.28

Prod. by *Streptomyces* sp. GÖM1.

Amorph. solid. λ_{max} 219 (log ε 3.93); 248 (log ε 3.43); 340 (log ε 3.19) (MeOH).

Me ester: Methyl 2-aminobenzoate.

FEMA 2682

[134-20-3]

$C_8H_9NO_2$ 151.165

Found in essential oils, including bergamot, orange peel, lemon peel, jasmine, ylang-ylang and neroli. Also present in concord grape, strawberry, star fruit, black tea and rice bran.

Found in egg masses of the muricid mollusc *Trunculariopsis trunculus*.

Used in perfumery as artificial orange blossom fragrance, also employed extensively in the form of its Schiff's bases (e.g. with Hydroxycitronellal). Flavouring agent. Cryst. Mp 24-25°. Bp₁₅ 133.5°. pK_a 2.32 (25°, 1% EtOH aq.). Steam-volatile.

▶ Skin irritant. LD₅₀ (rat, orl) 2910 mg/kg. Exp. reprod. effects. CB3325000

Et ester: Ethyl 2-aminobenzoate. FEMA 2421

[87-25-2]

$C_9H_{11}NO_2$ 165.191

Present in orange and grape. Flavouring ingredient. Liq. with faint orange-flower odour and flavour. d_4^{20} 1.12. Mp 13°. Bp 266-268° Bp₁₅ 145-147°.

▶ DG2448000

Amide: 2-Aminobenzamide. Anthranilamide

[88-68-6]

$C_7H_8N_2O$ 136.153

Prod. by a marine *Cytophaga marino-flava* sp. AM13.1. Acetaldehyde scavenger for polyethylene beverage bottles. Fluorescent label for polysaccharides etc. Leaflets. Mp 109-111.5° dec.

▶ CU8993000

Amide, N²-(2S-hydroxypropanoyl): 2-[(2-Hydroxypropanoyl)amino]benzamide

[18326-27-7]

$C_{10}H_{12}N_2O_3$ 208.216

Prod. by *Penicillium chrysogenum*. Oil.

N-Eicosanoyl: N-Eicosanoylanthranilic acid

N-Arachidylanthranilic acid

[165393-50-0]

$C_{27}H_{45}NO_3$ 431.657

Constit. of *Inula japonica* and *Ononis natrix*. Cryst. (MeOH). Mp 86-87°.

N-Docosanoyl: N-Docosanoylanthranilic acid

[85563-88-8]

$C_{29}H_{49}NO_3$ 459.711

Constit. of *Inula oculus-christi* and *Inula japonica*. Cryst. (Me₂CO/petrol). Mp 91-93°.

N-Docosanoyl, Et ester: Ethyl N-docosanoylanthranilate

[209523-04-6]

$C_{31}H_{53}NO_3$ 487.765

Constit. of the roots of *Gentiana tibetica*. Antifungal agent.

N-(13*Z*-Docosenoyl): N-(13-*Docosenoyl*)anthranilic acid
[129277-41-4]
C₂₉H₄₇NO₃ 457.695
Constit. of *Ononis natrix* ssp. *hispanica*. Oil.

N-Tricosanoyl: N-Tricosanoylanthranilic acid
C₃₀H₅₁NO₃ 473.738
Constit. of *Inula japonica*.

N-Tetracosanoyl: N-Tetracosanoylanthranilic acid
C₃₁H₅₃NO₃ 487.765
Constit. of *Inula japonica*.

N-(Carboxyacetyl): 2-[(Carboxyacetyl)amino]benzoic acid, 9CI. 2-(Malonylamino)benzoic acid
[53947-84-5]
C₁₀H₉NO₅ 223.185
Isol. from the leaves of the peanut (*Arachis hypogaea*). Cryst. (Me₂CO/hexane). Mp 190-192°.

N-(3ξ-Hydroxyeicosanoyl): N-(3-Hydroxyeicosanoyl)anthranilic acid
[856686-30-1]
C₂₇H₄₅NO₄ 447.657
Constit. of the aerial parts of *Ononis natrix*.

N-Succinoyl: N-Succinoylanthranilic acid
[5694-37-1]
[120572-43-2 (Et ester)]
C₁₁H₁₁NO₅ 237.212
Alkaloid from *Aconitum septentrionale*. Cryst. (Me₂CO) (as di-Et ester). Mp 57-58° (di-Et ester). Prob. artifact.

N-Succinoyl, Me ester: Methyl N-succinoylanthranilate
[108540-96-1]
C₁₂H₁₃NO₅ 251.238
Constit. of *Aconitum laeve*. Amorph. powder.

N-[3-(Methoxycarbonyl)propanoyl], Me ester: [59868-50-7]
C₁₃H₁₅NO₅ 265.265
Constit. of the alga *Jolyana laminarioides*. Chymotrypsin inhibitor. Amorph. powder. Mp 50°. λ_{max} 222; 251; 307 (MeOH).

N-(4-Methylbenzyl): 2-[(4-Methylbenzyl)amino]benzoic acid. **Onosmin A**
C₁₅H₁₅NO₂ 241.289
Constit. of *Onosma hispidum*. Lipoxygenase inhibitor. Amorph. solid. Mp 185-187°. λ_{max} 284; 343 (MeOH).


N-(4-Methylbenzyl), Me ester: **Onosmin B**
C₁₆H₁₇NO₂ 255.316
Constit. of *Onosma hispidum*. Lipoxygenase inhibitor. Amorph. solid. Mp 137-140°. λ_{max} 275; 353 (MeOH).

[99196-75-5; 552-37-4; 99197-00-9; 117808-63-6; 99196-74-4]

Kimura, Y. et al., *Agric. Biol. Chem.*, 1974, **38**, 1507-1510 (2-Malonylamino)benzoic acid)
Malakov, P.Y. et al., *Phytochemistry*, 1982, **21**, 2589-2590 (*Docosanoylanthranilic acid*)
Barrero, A.F. et al., *Phytochemistry*, 1990, **29**, 1967-1969 (*Docosanoylanthranilic acid*)
Al-Khalil, S. et al., *J. Nat. Prod.*, 1995, **58**, 760-763 (*Eicosanoylanthranilic acid*)

Usmanova, S.K. et al., *Khim. Prir. Soedin.*, 1996, **32**, 77-81; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 62-65 (*N-succinoyl*)
Tan, R.X. et al., *Phytochemistry*, 1998, **47**, 1223-1226 (*Ethyl docosanoylanthranilate*)
Nawasreh, M. et al., *Alexandria J. Pharm. Sci.*, 2004, **18**, 165-170; *CA*, **143**, 93919r (*N-3-Hydroxyeicosanoylanthranilic acid*)
Shaaban, M. et al., *Dissertation*, Univ. of Göttingen, 2004, (*marine, isol*)
Ahmad, I. et al., *Chem. Pharm. Bull.*, 2005, **53**, 907-910 (*Onosmins A,B*)
Shaheen, F. et al., *Phytochemistry*, 2005, **66**, 935-940 (*Methyl N-succinoylanthranilate*)
Wang, J. et al., *Plant J.*, 2005, **44**, 606-619 (*biosynth*)
Bitzer, J. et al., *Eur. J. Org. Chem.*, 2006, 3661-3666 (*6-deoxytalopyranosyl ester*)
Qin, J.J. et al., *Chin. Chem. Lett.*, 2008, **19**, 556-558 (*Inula japonica N-acyl derivs*)

2-Amino-8-benzoyl-6-hydroxy-3*H*-phenoxazin-3-one A-707




C₁₉H₁₂N₂O₄ 332.315
Prod. by marine bacterium *Halomonas* sp. GWS-BW-H8hM. Moderately cytotoxic. Red solid. λ_{max} 257 (log ε 3.81); 309 (log ε 3.59); 432 (log ε 3.46) (MeOH). λ_{max} 254 (log ε 3.73); 305 (log ε 3.53); 454 (log ε 3.57) (MeOH/HCl). λ_{max} 243 (log ε 3.89); 438 (log ε 3.67) (MeOH/NaOH).

4'-Hydroxy: 2-Amino-6-hydroxy-8-(4-hydroxybenzoyl)-3*H*-phenoxazin-3-one
C₁₉H₁₂N₂O₅ 348.314
Prod. by *Halomonas* sp. GWS-BW-H8hM. Red solid. λ_{max} 266 (log ε 3.54); 300 (log ε 3.52); 428 (log ε 3.44); 434 (log ε 3.45) (MeOH). λ_{max} 249 (log ε 3.62); 304 (log ε 3.58); 465 (log ε 3.35) (MeOH/HCl). λ_{max} 239 (log ε 3.61); 346 (log ε 3.56); 445 (log ε 3.49) (MeOH/NaOH).

Bitzer, J. et al., *J. Antibiot.*, 2006, **59**, 86-92 (*isol, uv, pmr, cmr, ms*)

2-(2-Aminobenzylamino)benzyl alcohol A-708


2-[[(2-Aminophenyl)methyl]amino]benzenemethanol, 9CI
[83326-78-7]



C₁₄H₁₆N₂O 228.293
Isol. from the Indian medicinal plant *Justicia gendarussa* (Acanthaceae). Fine needles (CHCl₃/petrol). Mp 131°.
Me ether: 2-Amino-N-[2-(methoxymethyl)phenyl]benzenemethanamine, 9CI
[83326-77-6]

C₁₅H₁₈N₂O 242.32
From *Justicia gendarussa* (Acanthaceae). Viscous oil.
Chakravarty, A.K. et al., *Tetrahedron*, 1982, **38**, 1797 (*isol*)
Kamat, V.P. et al., *Org. Prep. Proced. Int.*, 1994, **26**, 494 (*synth, pmr*)
Bhattacharyya, S. et al., *Synth. Commun.*, 1995, **25**, 3597 (*synth*)

3-Amino-8,10-bisaboladiene A-709



C₁₅H₂₇N 221.385

(3α,6β,7ξ,8E)-form


N-Formyl: 3-Formamido-8,10-bisaboladiene
[134781-22-9]
C₁₆H₂₇NO 249.395
Isol. from the sponge *Halichondria cf. lendenfeldi*. Dec. on isol.

Isocyanide: 3-Isocyano-8,10-bisaboladiene
[134781-21-8]
C₁₆H₂₅N 231.38
Isol. from the sponge *Halichondria cf. lendenfeldi* and the mollusc *Phyllidia pustulosa*. Oil. [α]_D -5.6 (c, 0.2 in CHCl₃). Has -NC replacing NH₂. λ_{max} 230 (sh) (ε 14000); 237 (ε 15300); 246 (sh) (ε 11100) (no solvent reported).

Kassühlke, K.E. et al., *J.O.C.*, 1991, **56**, 3747-3750 (*isol, pmr, cmr*)

7-Amino-2,10-bisaboladiene A-710

7-Amino-7,8-dihydro-α-bisabolene. 7-Aminobisabolene



(6*R*,7*R*)-form

C₁₅H₂₇N 221.385

(6*R*,7*R*)-form [105281-34-3]

Constit. of *Ciocalypta* sp. Oil. [α]_D -15 (c, 0.4 in MeOH).

Hydrochloride: [105281-33-2]
Yellow oil. [α]_D²³ -8.3 (c, 0.5 in MeOH).

(6*R*,7*S*)-form [105281-43-4]

Constit. of a *Halichondria* sp. and a sponge *Theonella* sp. Oil. [α]_D +39 (MeOH). [α]_D²⁰ +59.9 (c, 3 in CHCl₃).

Δ⁹-Isomer, 11-hydroxy: 7-Amino-2,9-bisaboladien-11-ol. **Aminobisabolanol**
[108384-61-8]

C₁₅H₂₇NO 237.384
Constit. of *Theonella* sp. Oil. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O. [α]_D +29 (MeOH).

*Δ*¹¹-Isomer, 10*R*-hydroxy: 7-Amino-2,11-bisaboladien-10*R*-ol. **Isoaminobisabolanol b**

[108384-62-9]
C₁₅H₂₇NO 237.384

Constit. of *Theonella* sp. Oil. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O. [α]_D +40 (MeOH).

*Δ*¹¹-Isomer, 10*S*-hydroxy: 7-Amino-2,11-bisaboladien-10*S*-ol. **Isoaminobisabolanol a**

[108449-15-6]
C₁₅H₂₇NO 237.384

Constit. of *Theonella* sp. Oil. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O. [α]_D +34 (MeOH).


Sullivan, B.W. *et al.*, *J.O.C.*, 1986, **51**, 5134-5136 (*Halichondria constits*)

Gulavita, N.K. *et al.*, *J.O.C.*, 1986, **51**, 5136-5139 (*Ciocalypia constit*)

Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 928 (*Theonella constits, cryst struct*)

7-Amino-2,8-bis(2,4-dihydroxyphenyl)-3H-phenoxazin-3-one, 9CI A-711

Resorcein
[33869-21-5]



C₂₄H₁₆N₂O₆ 428.4

Tentative isolation. Constit. of tobacco leaves. Used in hair preparations and as an elastin stain. Violet needles (C₆H₆/Me₂CO). Dec. at 300°. λ_{max} 587 (MeOH).


Musso, H. *et al.*, *Chem. Ber.*, 1963, **96**, 1579 (*synth, uv*)

Lillie, R.D. *et al.*, *Acta Histochem.*, Suppl., 1971, **No. 9**, 625 (*synth, use*)

Avunddzhyan, E.S. *et al.*, *CA*, 1974, **81**, 166553e (*isol*)

2-Amino-3-(3-bromo-5-chloro-4-hydroxyphenyl)propanoic acid A-712

3-Bromo-5-chlorotyrosine, 9CI. 3-Chloro-5-bromotyrosine (incorr.)



C₉H₉BrClNO₃ 294.532

(S)-form

L-form


Isol. from the operculum of *Buccinum undatum* and cuticle of *Limulus polyphemus*.

Hunt, S. *et al.*, *Biochim. Biophys. Acta*, 1971, **252**, 401-404 (*isol*)

Welinder, B.S. *et al.*, *Biochim. Biophys. Acta*, 1972, **279**, 491-497 (*isol, synth, ms*)

4-Amino-5-bromo-pyrrolo[2,3-d]pyrimidine A-713

5-Bromo-1*H*-pyrrolo[2,3-d]pyrimidin-4-amine, 9CI
[22276-99-9]



C₆H₅BrN₄ 213.036


Isol. from a sponge *Echinodictyum* sp. Bronchodilator and hypotensive agent. Adenosine kinase inhibitor. Needles (EtOH aq.). Mp 240-241° (238-239°) dec. λ_{max} 232 (ε 16600); 282 (ε 8710) (pH 1) (Derep). λ_{max} 225 (ε 14800); 279 (ε 8910) (pH 11) (Derep).

Gerster, J.F. *et al.*, *J. Het. Chem.*, 1969, **6**, 207-213 (*synth, uv*)

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1983, **36**, 165-170 (*isol, uv, pmr, cmr, ms*)

4-Amino-1,3-butanediol A-714

[130239-34-8]



C₄H₁₁NO₂ 105.136

(S)-form [637343-51-2]
Oil.

(±)-form
Oil.

N-E-Cinnamoyl: **Piperlotine K**
[958296-18-9]

C₁₃H₁₇NO₃ 235.282
Alkaloid from the leaves of *Piper lolot*. Syrup. Racemate. λ_{max} 211; 218; 224; 280 (MeOH).

N-E-Cinnamoyl (E): **Piperlotine J**
[958296-17-8]

C₁₅H₁₉NO₄ 277.319
Alkaloid from the leaves of *Piper lolot*. Syrup. Racemate. λ_{max} 212; 218; 223; 280 (MeOH).

(ξ)-form

N-E-Cinnamoyl, 1,3-di-Ac: **Piperlotine L**
[958296-19-0]

C₁₇H₂₁NO₅ 319.357
Alkaloid from the leaves of *Piper lolot*. Syrup. [α]_D +42.2 (c, 0.1 in MeOH). λ_{max} 210; 216; 222; 274 (MeOH).

Kuriyama, N. *et al.*, *Synthesis*, 1990, 735-738 (*synth*)

Pat. Coop. Treaty (WIPO), 2003, 03 106 413; *CA*, **140**, 59633 (*S*-form, *synth*)

Li, C.-Y. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 9436-9442 (*Piperlotines J,K,L*)

4-Aminobutanoic acid, 9CI A-715

γ-Aminobutyric acid. *Piperidinic acid*.

Piperidic acid. *Aminalon*. *Gamalon*.

GABA

[56-12-2]
[28805-76-7]

H₂NCH₂CH₂CH₂COOH

C₄H₉NO₂ 103.121

Widely distributed in higher plants.

Prod. by *Monascus pilosus* (Ifo4520).

Antihypertensive agent. The natural inhibitory transmitter at synaptic junctions in some regions of the mammalian brain and spinal cord. Prisms (EtOH). V. sol. H₂O; sl. sol. EtOH, Me₂CO; insol. Et₂O. Mp 203° dec. Log P -3.33 (calc). Homopolymers (polyamides) are usually derived from the lactam 2-Pyrrolidinone, P-956.

▶ LD₅₀ (mus, orl) 12680 mg/kg. ES6300000
Hydrochloride: Mp 135-136°.

Hydrobromide: [36551-16-3]

Orange solid. Mp 215°.

Me ester: [13031-60-2]

C₅H₁₁NO₂ 117.147

Solid (EtOH/Et₂O) (as hydrochloride). Mp 119° (114-115°) (hydrochloride).

▶ ES7065000

Et ester: [5959-36-4]

C₆H₁₃NO₂ 131.174

Bp₁₂ 75-77°.

Et ester: hydrochloride: Mp 65-72°.

Hygroscopic.

▶ LD₅₀ (mus, ivn) 425 mg/kg. Exp. reprod. effects.

N-Ac: [3025-96-5]

C₆H₁₁NO₃ 145.158

Mp 129-131°.

▶ ES5610000

N-Ac; hydrochloride: Mp 133°.

N-(2,2-Dimethylpropanoyl)-4-[(2,2-Dimethyl-1-oxopropyl)amino]butanoic acid, 9CI. Pivagabine, INN

[69542-93-4]

C₉H₁₇NO₃ 187.238

Anticonvulsant.

N-(5-Hydroxy-3-methyl-2*Z*-pentenoyl):

Fuscoatramide

C₁₀H₁₇NO₄ 215.249

Prod. by *Humicola fuscoatra* NRRL 22980. Oil. λ_{max} 233 (ε 3900) (MeOH).

N-tert-Butyloxycarbonyl: [57294-38-9]

C₉H₁₇NO₄ 203.238

Cryst. (EtOAc/hexane). Mp 53-54°. Bp_{0.2} 78-82°.

N-tert-Butyloxycarbonyl, anhydride:

[89231-63-0]

C₁₈H₃₂N₂O₇ 388.46

Cryst. (EtOAc/petrol). Mp 106°.

N-Benzoyloxycarbonyl: [5105-78-2]

C₁₂H₁₅NO₄ 237.255

Needles (EtOAc/hexane). Mp 65-67°.

N-Benzoyloxycarbonyl, amide: [35821-20-6]

C₁₂H₁₆N₂O₃ 236.27

Cryst. Mp 125-128°.

N-Me: [1119-48-8]

C₅H₁₁NO₂ 117.147

Cryst. Mp 151.5-153°.

N-Me; hydrochloride: [6976-17-6]

Mp 124-126°.

N,N-Di-Me: 4-(Dimethylamino)butanoic acid

[693-11-8]

C₆H₁₃NO₂ 131.174

Mp 102-104°. Bp₁₀ 135-137°.

- N,N-Di-Me; hydrochloride: Mp 145-147°.
- N,N-Di-Me, Me ester: [22041-22-1] C₇H₁₅NO₂ 145.201 Sol. H₂O. Mp 171.5-173°.
- ES9850000
- N-(2-Nitrobenzenesulfonyl): [80977-11-3] [104809-33-8] C₁₀H₁₂N₂O₄S 256.282 Solid. Mp 102° (92.5-94°).
- Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 577C; 577D (ir)
- Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 877C (nmr)
- Org. Synth., Coll. Vol., 2, 1943, 25 (synth)
- Syngé, R.L.M. et al., Biochem. J., 1951, 48, 429 (isol)
- Thompson, J.F. et al., Arch. Biochem. Biophys., 1953, 46, 248 (synth)
- Reppe, W. et al., Annalen, 1955, 596, 158 (derivs)
- Goerdler, J. et al., Angew. Chem., 1959, 71, 775 (N-(2-nitrophenylsulfonyl), synth)
- Evans, R.L. et al., J.O.C., 1959, 24, 863-864 (benzyloxycarbonyl)
- Taddei, F. et al., J.C.S., 1964, 1553 (pmr)
- Voellmin, J. et al., Microchem. J., 1966, 11, 73 (ms)
- Karrer, W. et al., Konstitution und Vorkommen der Organischen Pflanzenstoffe, 2nd edn., Birkhäuser Verlag, 1972, no. 2427 (occur)
- Steward, E.G. et al., Acta Cryst. B, 1973, 29, 2038; 2825 (cryst struct)
- Tomita, K. et al., Bull. Chem. Soc. Jpn., 1973, 46, 2199 (cryst struct)
- Galzigna, L. et al., Arch. Int. Pharmacodyn. Ther., 1978, 235, 73 (Pivagabine)
- Terano, S. et al., Phytochemistry, 1978, 17, 550 (biosynth)
- Krogsgaard-Larsen, P. et al., Gaba-Neuro-Transm.: Pharmacochem., Biochem., Pharmacol. Aspects, 1979, Academic Press, N.Y., 1979,
- Rauls, D.O. et al., J. Med. Chem., 1979, 22, 81-86 (benzyloxycarbonyl amide)
- Stewart, F.H.C. et al., Aust. J. Chem., 1981, 34, 2439-2447 (N-(2-nitrophenylsulfonyl), synth)
- Meldrum, B. et al., Clin. Neuropharmacol., 1982, 5, 293
- Battersby, A.R. et al., J.C.S. Perkin 1, 1982, 455 (biosynth)
- Weber, H.P. et al., Acta Cryst. B, 1983, 39, 360 (cryst struct)
- Shashua, V.E. et al., J. Med. Chem., 1984, 27, 659-664 (tert-butyloxycarbonyl anhydride)
- Blankespoor, R.L. et al., J.O.C., 1984, 49, 4441-4446 (benzyloxycarbonyl, synth, ir, pmr)
- Galakatos, N.G. et al., J.O.C., 1985, 50, 1302 (deriv, synth, pmr)
- De Feudis, F.V. et al., Drugs of Today (Barcelona), 1986, 22, 563 (rev)
- Roberts, E. et al., Recept. Biochem. Methodol., 1986, 5, 1 (rev)
- Negwer, M. et al., Organic-Chemical Drugs and their Synonyms, 6th edn., Akademie-Verlag, 1987, 223 (synonyms)
- Kirk, D.N. et al., J.C.S. Perkin 1, 1988, 2979-2982 (tert-butyloxycarbonyl)
- Houssin, R. et al., Synthesis, 1988, 259-261 (tert-butyloxycarbonyl)
- Nisticò, G. et al., GABA: Basic Research and Clinical Applications, Pythagora Press, 1989, (book)
- Ramek, M. et al., Int. J. Quantum Chem., 1990, 17, 45 (conformn)
- Erdoe, S.L. et al., J. Neurochem., 1990, 54, 363 (rev)
- Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1336
- Bon, E. et al., J.O.C., 1994, 59, 1904 (synth, ir, pmr, cmr)
- Joshi, B.K. et al., J. Nat. Prod., 2002, 65, 1734-1737 (Fuscoatramide)
- Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold, 1992, AAJ350; PIM500

4-Amino-1-butanol **A-716**

[13325-10-5]

H₂NCH₂CH₂CH₂CH₂OHC₄H₁₁NO 89.137Oil. Sol. EtOH, insol. Et₂O. Bp 206° Bp₃₄ 125°. Forms a hydrate in water. Absorbs CO₂ and H₂O from air.► LD₅₀ (rat, orl) 1150 mg/kg. EK9700000

Picrate: Mp 86.6-88.6°.

N-Benzyloxycarbonyl: [17996-13-3]

C₁₂H₁₇NO₃ 223.271

Cryst. (EtOAc/hexane). Mp 78-79°.

N-[3-(Methylthio)-2E-propenoyl]: N-(4-Hydroxybutyl)-3-(methylthio)-2-propenamide. **Aglatenol**C₈H₁₅NO₂S 189.278Alkaloid from the leaves of *Aglaia tenuicaulis*. λ_{max} 224 (log ε 3.78); 274 (log ε 4.32) (MeOH aq.).

N-O-Bis[3-(methylthio)-2E-propenoyl]:

AglateninC₁₂H₁₉NO₃S₂ 289.419Alkaloid from the leaves of *Aglaia tenuiformis*. Cryst. Mp 81-83°. λ_{max} 224 (log ε 3.75); 274 (log ε 4.49) (MeOH aq.).N-E-Cinnamoyl, O-[3-(methylthio)-2E-propenoyl]: **Tenuicaulin A**C₁₇H₂₁NO₃S 319.424Alkaloid from the leaves of *Aglaia tenuicaulis*. Cryst. Mp 75-77°. λ_{max} 216 (log ε 4.18); 221 (sh) (log ε 4.12); 276 (log ε 4.52) (MeOH aq.).N-E-Cinnamoyl, O-[3-(methylthio)-2Z-propenoyl]: **Isotenuicaulin A**C₁₇H₂₁NO₃S 319.424Alkaloid from the leaves of *Aglaia tenuicaulis*. Isol. as a mixt. with Tenuicaulin B. λ_{max} 216 (log ε 4.17); 221 (sh) (log ε 4.12); 282 (log ε 4.48) (MeOH aq.).N-E-Cinnamoyl, O-benzoyl: **Caulitenin**C₂₀H₂₁NO₃ 323.391Alkaloid from the stem bark of *Aglaia tenuicaulis*. λ_{max} 218 (sh) (log ε 4.32); 222 (log ε 4.32); 231 (sh) (log ε 4.18); 274 (log ε 4.35) (MeOH aq.).N,O-Di-E-cinnamoyl: **Tenaglin**C₂₂H₂₃NO₃ 349.429Alkaloid from the stem bark of *Aglaia tenuicaulis*. λ_{max} 216 (log ε 4.4); 221 (sh) (log ε 4.35); 274 (log ε 4.54) (MeOH aq.).N-Z-Cinnamoyl, O-[3-(methylthio)-2E-propenoyl]: **Tenucaulin B**C₁₇H₂₁NO₃S 319.424Alkaloid from the leaves of *Aglaia tenuicaulis*. Isol. as a mixt. with Isotenucaulin A. λ_{max} 274 (log ε 4.16) (MeOH aq.).

N-Me: [42042-68-2]

C₅H₁₃NO 103.164Liq. Bp_{0.36} 71°.

N,N-Di-Me: 4-(Dimethylamino)-1-butanol, 9CI

[13330-96-6]

C₆H₁₅NO 117.191Oil. Bp 186-189° Bp₁₄ 80-81°.

Me ether: 4-Methoxy-1-butylamine. 1-Amino-4-methoxybutane

[34039-36-6]

C₅H₁₃NO 103.164Sol. H₂O, EtOH, Et₂O. Bp 142-145°.

N,N-Dibutyl: [94473-23-1]

C₁₂H₂₇NO 201.351Bp₄₀ 172-175°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 338D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 538A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 429A (ir)

Henry, L. et al., Ber., 1900, 33, 3170 (synth)

Lunsford, C.D. et al., J.O.C., 1957, 22, 1225 (N,N-dibutyl)

Kaluszyner, A. et al., J.O.C., 1961, 26, 3536 (deriv)

Hoffman, R.V. et al., J.C.S. Perkin 1, 1989, 1375 (deriv)

Zoppetti, G. et al., Biochim. Biophys. Acta, 1992, 1156, 92 (N,N-dibutyl)

Wilk, A. et al., J.O.C., 1999, 64, 7515-7522 (N-Me, synth, pmr, cmr)

Wakayama, T. et al., Bull. Chem. Soc. Jpn., 2004, 77, 331-340 (N-benzyloxycarbonyl)

Greger, H. et al., Phytochemistry, 2008, 69, 928-938 (*Aglaia tenuicaulis* alkaloids)**4-Amino-2-butenic acid**, **A-717**
9CI

4-Aminocrotonic acid

[25747-40-4]

H₂NCH₂CH=CHCOOHC₄H₇NO₂ 101.105**(E)-form** [38090-53-8] Neurochemical transmission inhibitor. Prisms (EtOH). Sol. H₂O. Mp 164° dec.**(E)-form**

N-Ac: 4-(Acetylamino)-2-butenic acid,

9CI

[64120-63-4]

C₆H₉NO₃ 143.142Metab. of *Fusarium graminearum*.Cryst. (MeOH/CHCl₃). Mp 140°.

Balenovic, K. et al., J.O.C., 1954, 19, 1589 (synth)

Jones, G.P. et al., J.C.S. Perkin 2, 1975, 1059 (cryst struct)

Vesonder, R.F. et al., Phytochemistry, 1977, 16, 1296 (deriv, isol)

Allan, R.D. et al., Aust. J. Chem., 1978, 31, 2283 (synth)

N-(4-Aminobutyl)-N-(3-ami- **A-718**
nopropyl)-1,4-butanediamine, 9CI

5-(3-Aminopropyl)-1,9-diamino-5-azan-

nane. N³-(3-Aminopropyl)homospermi-

dine

[143085-75-0]

(H₂NCH₂CH₂CH₂CH₂)₂NCH₂CH₂C-H₂NH₂C₁₁H₂₈N₄ 216.369

Isol. from thermophilic bacteria.

Hamana, K. et al., Biochem. J., 1992, 284, 741-

747

(4-Aminobutyl)guanidine, 9CI A-719

4-Guanidinobutylamine. 1-Amino-4-guanidinobutane. **Agmatine**

[306-60-5]



$\text{C}_5\text{H}_{14}\text{N}_4$ 130.192

Present in ergot, pollen of *Ambrosia artemisiifolia* (Asteraceae), the sea anemone *Anthopleura japonica* and herring semen (*Clupea* sp.). Occurs in *Stichopus japonicus*, *Halocynthia roretzi*, *Crassostrea gigas* and *Tapes philippinarum*. Metab. intermed. for polyamines. Synth. in mammalian brain. Endogenous neurotransmitter. Inhibits hyperalgesia and tolerance to morphine. NMDA receptor antagonist and nitric oxide synthase inhibitor. Mp 101.5-103°.

Sulfate: [2482-00-0]

Cryst. Mp 229° (224-225°).

***N*¹-Ac: *N*¹-Acetylglutamine**

[3031-89-8]

$\text{C}_7\text{H}_{16}\text{N}_4\text{O}$ 172.23

Isol. from *Actinia equina* and *Actinia fragacea*.

N*¹-(3*S*-Methyldecanoyl): *Aplysillamide B

[164301-83-1]

$\text{C}_{16}\text{H}_{34}\text{N}_4\text{O}$ 298.471

Alkaloid from *Psammaphysilla purea*. Exhibits modest antimicrobial activity against some fungi and bacteria. Not cytotoxic. Oil. Sol. MeOH. $[\alpha]_{\text{D}}^{21}$ -2.4 (c, 0.1 in MeOH) (natural). $[\alpha]_{\text{D}}^{25}$ -5.1 (c, 1.6 in MeOH) (synthetic).

N*¹-(3-Methyl-2*Z*-decenoyl): *Aplysillamide A

[164301-82-0]

$\text{C}_{16}\text{H}_{32}\text{N}_4\text{O}$ 296.455

Alkaloid from the Okinawan marine sponge *Psammaphysilla purea*. Exhibits modest antimicrobial activity against some fungi and bacteria. Cytotoxic against murine lymphoma L1210 and human epidermoid carcinoma KB cells. Oil. Sol. MeOH. λ_{max} 225 (ε 10000) (MeOH) (Berdy).

***N*¹-(4-Hydroxy-E-cinnamoyl): *N*¹-trans-p-Coumaroylagmatine**

[47096-24-2]

[7295-86-5]

$\text{C}_{14}\text{H}_{20}\text{N}_4\text{O}_2$ 276.338

Alkaloid from barley seedlings, *Hordeum bulbosum*, *Hordeum distichon*, *Hordeum jubatum*, *Hordeum murinum* and *Hordeum spontaneum* (Poaceae). Exhibits weak antifungal activity. Sol. H_2O . Mp 215-217° (as picrate). λ_{max} 229 (ε 33100); 300 (ε 26300) (H_2O) (Berdy).

***N*¹-(4-Hydroxy-Z-cinnamoyl): *N*¹-cis-p-Coumaroylagmatine**

[191330-81-1]

$\text{C}_{14}\text{H}_{20}\text{N}_4\text{O}_2$ 276.338

Constit. of *Albizzia julibrissin*. Leaf-opening factor. Syrup or powder. λ_{max} 278 (ε 2000) (H_2O).

***N*¹-(4-Hydroxy-3-methoxy-E-cinnamoyl): *N*¹-trans-Feruloylagmatine**

[188305-06-8]

$\text{C}_{15}\text{H}_{22}\text{N}_4\text{O}_3$ 306.364

Isol. from *Corydalis saxicola* and from *Triticum aestivum* (wheat) exposed to low temps. Antifungal agent.

***N*¹-(3,4-Dimethoxycinnamoyl): *N*¹-(3,4-Dimethoxycinnamoyl)agmatine**

[128009-18-7]

[146072-40-4]

$\text{C}_{16}\text{H}_{24}\text{N}_4\text{O}_3$ 320.391

Alkaloid from *Verbesina caracasana*.

***N*⁶-Me: N-(4-Aminobutyl)-*N*⁶-methylguanidine. *N*⁶-Methylglutamine**

[77414-15-4]

$\text{C}_6\text{H}_{16}\text{N}_4$ 144.219

Constit. of *Glycine max* and *Medicago sativa*.

***N*⁶-(3-Methyl-2-butenyl): N-(4-Aminobutyl)-*N*⁶-prenylguanidine. *N*⁶-(3-Methyl-2-butenyl)agmatine. *N*⁶-Prenylglutamine**

[128009-17-6]

$\text{C}_{10}\text{H}_{22}\text{N}_4$ 198.311

Alkaloid from *Verbesina caracasana*.

Antihypertensive.

***N*⁶-(3-Methyl-2-butenyl), *N*¹-(3,4-dimethoxycinnamoyl):** see Caracasamide, C-107

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 821B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1330A (nmr)

Kossel, A. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1910, **66**, 257-261; **68**, 170-171 (isol. struct. synth)

Heyl, F.W. et al., *J.A.C.S.*, 1919, **41**, 670-682 (isol)

Stoessel, A. et al., *Phytochemistry*, 1965, **4**, 973-976 (isol, uv, struct, synth, *N*-coumaroyl)

Boldt, A. et al., *Phytochemistry*, 1971, **10**, 731-738 (biosynth)

Smith, T.A. et al., *Phytochemistry*, 1978, **17**, 1093-1098 (occur, *N*-coumaroyl)

Kowabata, T. et al., *CA*, 1979, **89**, 178349 (isol)

Robin, Y. et al., *Oceanis*, 1980, **5**, 575-580; *CA*, **94**, 118125p (*N*¹-Acetylglutamine)

Bird, C.R. et al., *Phytochemistry*, 1981, **20**, 2345-2346 (biosynth, *N*-coumaroyl)

Chandrasekhar, K. et al., *Acta Cryst. B*, 1982, **38**, 2538-2540 (cryst struct)

Cho, Y.B. et al., *Anal. Biochem.*, 1987, **160**, 429-433 (*N*⁶-Methylglutamine)

Matsuzaki, S. et al., *Phytochemistry*, 1990, **29**, 1313-1315 (*N*⁶-Methylglutamine)

Hamana, K. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur, hplc)

Mitchinson, A. et al., *Chem. Comm.*, 1994, 2613-2614 (synth)

Honma, K. et al., *Tetrahedron*, 1995, **51**, 3745 (*Aplysillamides*)

Ueda, M. et al., *Biosci., Biotechnol., Biochem.*, 1998, **62**, 2133-2137 (*Z*-coumaroyl)

Reis, D.J. et al., *Ann. N.Y. Acad. Sci.*, 1999, **881**, 65-80 (rev, activity)

Monache, G.D. et al., *Bioorg. Med. Chem. Lett.*, 1999, **9**, 3249-3254 (*Prenylglutamine*, *Dimethoxycinnamoylagmatine*)

Jin, S. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 1614-1617 (*Feruloylagmatine*)

Fairbanks, C.A. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 10584-10589 (pharmacol)

Reis, D.J. et al., *Trends Pharmacol. Sci.*, 2000, **21**, 187-193 (pharmacol, rev)

Demady, D.R. et al., *Mol. Pharmacol.*, 2001, **59**, 24-29 (pharmacol)

Cheng, X. et al., *Chem. Biodiversity*, 2008, **5**, 1335-1344 (*Feruloylagmatine*)

(4-Aminobutyl)urea, 8CI A-720

***N*-Carbamoylputrescine**

[6851-51-0]



$\text{C}_5\text{H}_{13}\text{N}_3\text{O}$ 131.177

Alkaloid from *Hordeum vulgare* (barley) and *Sesamum indicum* (sesame) (K-deficient plants)(Poaceae, Pedaliaceae).

Hydrochloride: Mp 185-186°.

Dipicrate: Mp 164-165°.


Linneweh, F. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1932, **205**, 126; *CA*, **26**, 3276 (synth)

Smith, T.A. et al., *Phytochemistry*, 1964, **3**, 23 (occur, synth)

Crocomo, O.J. et al., *Phytochemistry*, 1970, **9**, 1487 (occur)

4-(2-Amino-2-carboxyethyl)-1*H*-pyrrole-2-carboxylic acid A-721

3-(2-Carboxy-4-pyrrolyl)alanine. 2-Amino-3-(2-carboxy-4-pyrrolyl)propanoic acid



$\text{C}_8\text{H}_{10}\text{N}_2\text{O}_4$ 198.178

(*S*)-form

L-form

[137309-97-8]

Isol. from the toxic mushroom *Clitocybe acromelalga*. Mp 200-202° dec.


(±)-form

Mp 200-202° dec.

Yamano, K. et al., *Tetrahedron*, 1992, **48**, 1457 (isol, synth, struct)

 α -Amino-6-carboxy-2-oxo-3-pyridinepropanoic acid A-722

3-(6-Carboxy-2-oxo-4-pyridinyl)alanine



$\text{C}_9\text{H}_{10}\text{N}_2\text{O}_5$ 226.188


Sol. H_2O , MeOH; poorly sol. EtOAc, hexane. λ_{max} 241 (ε 12000); 315 (ε 13500) (H_2O) (Berdy).

(*S*)-form [148001-20-1]

Isol. from the toxic mushroom *Clitocybe acromelalga*. $[\alpha]_{\text{D}}^{23}$ -5.8 (c, 0.13 in H_2O).

Yamano, K. et al., *Heterocycles*, 1993, **35**, 125 (isol, struct)

Yamano, K. et al., *Tetrahedron*, 1993, **49**, 2427 (isol)

17-Amino-3,7,11-cembratrien-16,2-olid-19-oic acid A-723C₂₀H₂₉NO₄ 347.453**(1S,2S,3E,7E,11E,15R)-form**

N,N-Di-Me: 17-Dimethylaminolobohedrolide (incorr.)


[267007-12-5]

C₂₂H₃₃NO₄ 375.507Constit. of a *Lobophytum* sp.

Shows moderate anti-HIV activity.

Gum. [α]_D²⁰ +13.1 (c, 0.25 in CHCl₃).λ_{max} 218 (log ε 3.6); 222 (log ε 3.56) (EtOH).Rashid, M.A. et al., *J. Nat. Prod.*, 2000, **63**, 531-533 (isol, pmr, cmr)**2-Amino-3-(3-chloro-4-hydroxyphenyl)propanoic acid** A-724

3-Chlorotyrosine. Monochlorotyrosine [7298-90-0]

C₉H₁₀ClNO₃ 215.636**(R)-form**

D-form

[162599-96-4]

[905309-56-0 (hydrochloride)]

Mp 255-256° dec. [α]_D²⁰ +26 (c, 0.9 in H₂O). [α]_D²⁴ -12 (c, 3.05 in MeOH).

N-(3-Methoxycarbonyl-2E-propenyl):


Xylariamide A

[853259-88-8]


C₁₄H₁₄ClNO₆ 327.72Isol. from *Xylaria* sp. (FRR 5657).Pale yellow gum. [α]_D²⁴ -22 (c, 0.06 in MeOH). λ_{max} 202 (log ε 4.05); 217 (sh) (log ε 3.74); 278 (log ε 3.12) (MeOH).**(S)-form**

L-form

[7423-93-0]

Isol. from hydrolysates of mollusc (*Buccinum undatum*) scleroproteins and insect cuticle (*Schistocerca gregaria* and *Limulus polyphemus*). [α]_D²⁰ -26.2.Dibbo, A. et al., *J.C.S.*, 1961, 2645-2651 (R-form, synth)Welinder, B.S. et al., *Biochim. Biophys. Acta*, 1972, **279**, 491-497 (isol, ms)Hunt, S. et al., *FEBS Lett.*, 1972, **24**, 109-112 (isol)Patel, V.F. et al., *J. Med. Chem.*, 1999, **42**, 2588-2603 (synth, pmr, cmr)Davis, R.A. et al., *J. Nat. Prod.*, 2005, **68**, 769-772 (*Xylariamide A*)Davis, R.A. et al., *Tet. Lett.*, 2005, **46**, 5199-5201 (*Xylariamide A*, synth)**3-(2-Amino-3-chlorophenyl)-4-chloro-1H-pyrrole, 8CI** A-7252-Chloro-6-(4-chloro-1H-pyrrol-3-yl)benzenamine, 9CI. **Aminopyrrolnitrin**. WB 2838. Antibiotic WB 2838 [16386-65-5]C₁₀H₈Cl₂N₂ 227.092Prod. by *Pseudomonas aureofaciens* and *Pseudomonas cepacia*. Androgen-receptor antagonist and antifungal agent.Pale yellow needles (MeOH aq.). Mp 91° (synthetic). Similar to Pyrrolnitrin, P-961. λ_{max} 216 (ε 21400); 270 (ε); 299 (ε 3890) (solvent not reported) (Derep).

2-Chloro-4-(2-Amino-3-chlorophenyl)-2,3-dichloro-1H-pyrrole. Aminochloropyrrolnitrin [124201-43-0]

C₁₀H₇Cl₃N₂ 261.537From *Pseudomonas cepacia*. λ_{max} 212 (ε 34500); 302 (ε 4080) (MeOH) (Berdy).Salcher, O. et al., *Tet. Lett.*, 1978, 3097 (isol, pmr)Roitman, J.N. et al., *Appl. Microbiol. Biotechnol.*, 1990, **34**, 381 (isol)Roitman, J.N. et al., *J. Agric. Food Chem.*, 1990, **38**, 538 (isol)Hori, Y. et al., *J. Antibiot.*, 1993, **46**, 1327 (isol, props)**5-Amino-5-deoxygalactose** A-7266-Hydroxymethyl-2,3,4,5-piperidinetetrol, 9CI. **Galactostatin**. galacto-*Nojirimycin*. 5-Amino-5-deoxygalactopyranose

α-D-Pyranose-form

C₆H₁₃NO₅ 179.172

Log P -3.55 (calc).

D-form [107537-94-0]Isol. from *Streptomyces lydicus*β-Galactosidase inhibitor. Amorph. solid + ½ H₂O or syrup. Mp 94-98°. [α]_D²³ +85.6 (c, 1 in H₂O). Stable up to 5d at 4°.**Bisulfite adduct**: Mp 149-150°. [α]_D +19 (c, 0.8 in H₂O).**D-Pyranose-form****Pentakis(trimethylsilyl)**: [40222-80-8]C₂₁H₅₃NO₅Si₅ 540.081

Liq.

α-D-Furanose-form

1,2-Isopropylidene-5-Amino-5-deoxy-1,2-O-isopropylidene-α-D-galactofuranose

C₉H₁₇NO₅ 219.237Hygroscopic solid. [α]_D²² -25.3 (c, 1.0 in MeOH).


6-Trityl, 1,2-isopropylidene-5-Amino-5-deoxy-1,2-O-isopropylidene-6-O-trityl-α-D-galactofuranose

C₂₈H₃₁NO₅ 461.557Mp 135°. [α]_D²³ -11.5 (c, 1.0 in CHCl₃).

[40222-81-9, 108147-56-4, 109718-63-0]

Inouye, S. et al., *CA*, 1973, **78**, 47847g (silyl deriv)Legler, G. et al., *Carbohydr. Res.*, 1986, **155**, 119 (α-D-fur deriv)Miyake, Y. et al., *J. Antibiot.*, 1987, **40**, 122-123 (isol)Miyake, Y. et al., *Agric. Biol. Chem.*, 1988, **52**, 153-158; 661-666 (isol, pmr, cmr, struct, props)Aoyagi, S. et al., *J.O.C.*, 1991, **56**, 815-819 (synth)Chida, N. et al., *Chem. Comm.*, 1994, 1247-1248 (synth)Kirihata, M. et al., *Heterocycles*, 1995, **41**, 2271 (synth)Dondoni, A. et al., *J.O.C.*, 1995, **60**, 4749 (synth, pmr)Pérez, P.D. et al., *Eur. J. Org. Chem.*, 2005, 2903-2913 (α-D-fur 1,2-isopropylidene)**5-Amino-5-deoxyglucose, 9CI, 8CI** A-727

6-(Hydroxymethyl)-2,3,4,5-piperidinetetrol



D-Pyranose-form

C₆H₁₃NO₅ 179.172**D-Pyranose-form****Nojirimycin**

[15218-38-9]

[19130-94-0 (α-D-pyr-form)]

Amino sugar antibiotic. Produced by several *Streptomyces* spp. Primarily active against gram-positive bacteria. Sol. H₂O; fairly sol. MeOH, EtOH; poorly sol. butanol, hexane. Mp 126-130° dec. [α]_D²⁴ +100 (3 min.) → +73.5 (20 hr.).▶ LD₅₀ (mus, ipr) 1600 mg/kg; LD₅₀ (mus, ivn) 1250 mg/kg. LZ5655000

1-Deoxy: see 2-(Hydroxymethyl)-3,4,5-piperidinetriol, H-616

L-Pyranose-formMp 122-124°. [α]_D²⁰ -72.1 (c, 0.3 in H₂O).**α-D-Furanose-form**

1,2-O-Isopropylidene-5-Amino-5-deoxy-1,2-O-isopropylidene-α-D-glucofuranose [16958-27-3]

C₉H₁₇NO₅ 219.237Cryst. (EtOAc). Mp 125-126° Mp 175°. [α]_D²⁵ -13.75 (c, 4 in MeOH).


Higher Mp product (2004) may be the β-L-ido-analogue.

1,2:3,6-Diisopropylidene-5-Amino-5-deoxy-1,2:3,6-di-O-isopropylidene-α-D-glucofuranose

C₁₂H₂₁NO₅ 259.302

Needles (EtOAc/heptane). Mp 68-69°.


1,2-O-Isopropylidene, N-3,6-tri-Ac: 5-Acetamido-3,5-di-O-acetyl-5-deoxy-1,2-O-isopropylidene- α -D-glucofuranose [16958-28-4]

C₁₅H₂₃NO₈ 345.349Needles (Et₂O). Mp 145-146°. [α]_D²⁵ +1.3 (c, 6 in CHCl₃).Inouye, S. *et al.*, *J. Antibiot.*, 1966, **19**, 288 (struct, nmr)Saeki, H. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 962 (synth)Nayak, U.G. *et al.*, *J.O.C.*, 1968, **33**, 3582-3585 (α -D-fur isopropylidene, α -D-fur isopropylidene tri-Ac)Inouye, S. *et al.*, *Tetrahedron*, 1968, **24**, 2125 (ir, ms, nmr, struct, synth)Ger. Pat., 1978, 2 658 561; *CA*, **89**, 152713 (use)Kinast, G. *et al.*, *Angew. Chem., Int. Ed.*, 1981, **20**, 805 (bibl)Austrian Pat., 1982, 366 032; *CA*, **97**, 90401 (synth)Vasella, A. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 1134 (synth)Ezure, Y. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 1119 (isol, bibl)Iida, H. *et al.*, *J.O.C.*, 1987, **52**, 3337 (synth)Tsuda, Y. *et al.*, *Heterocycles*, 1988, **27**, 63 (synth)Kayakiri, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1397-1401 (α -D-fur isopropylidene)Chida, N. *et al.*, *Carbohydr. Res.*, 1992, **237**, 185 (synth)Dondoni, A. *et al.*, *Tetrahedron*, 1993, **49**, 2939 (synth, L-form)Dondoni, A. *et al.*, *Chem. Eur. J.*, 1995, **1**, 505-520 (α -D-fur isopropylidene)Moutel, S. *et al.*, *J.C.S. Perkin 1*, 1999, 1403-1406 (synth)Gottschaldt, M. *et al.*, *Carbohydr. Res.*, 2004, **339**, 1941-1952 (α -D-fur isopropylidene, α -D-fur diisopropylidene)Morwenna, M.S.M. *et al.*, *Eur. J. Org. Chem.*, 2005, 2159-2191 (rev, synth)**5-Amino-5-deoxymannose** A-728**Nojirimycin B.** Mannonojirimycin. Mannojirimycin α -D-Pyranose-formC₆H₁₃NO₅ 179.172

Aminoglycoside antibiotic.

D-Pyranose-form [62362-40-7]

Prod. by *Streptomyces lavendulae* SF-425. Weakly active against *Xanthomonas oryzae*. Glucosidase and mannosidase inhibitor. Pale yellow powder. Sol. H₂O; poorly sol. Me₂CO, hexane. Unstable.

Bisulfite adduct:Needles (H₂O). Mp 163-165° dec. [α]_D²⁰ +4.6 (c, 0.5 in H₂O).**L-Pyranose-form**Mp 162-164° (as bisulfite adduct). [α]_D²⁰ -4.5 (c, 0.31 in H₂O).Japan. Pat., 1976, 76 151 393; *CA*, **86**, 153954 (isol)Niwa, T. *et al.*, *J. Antibiot.*, 1984, **37**, 1579 (isol, cryst struct, props, nmr, ir)Dondoni, A. *et al.*, *Tetrahedron*, 1993, **49**, 2939 (L-form)**2-Amino-4,6-dichloropyridine** A-729
4,6-Dichloro-2-pyridinamineC₅H₄Cl₂N₂ 163.005Reported from Ku Shu (*Picrasma quasiosoides*). Cryst. (petrol). Mp 112.5°.

N-Ac: [63763-91-7]

C₇H₆Cl₂N₂O 205.043

Cryst. (EtOH). Mp 218-219°.

2-N,N-Di-Me: 4,6-Dichloro-N,N-dimethyl-2-pyridinamine, 9CI. 2,4-Dichloro-6-(dimethylamino)pyridine [849937-99-1]

C₇H₈Cl₂N₂ 191.059

Platelets (hexane). Mp 45-46°.

Den Hertog, H.J. *et al.*, *Rec. Trav. Chim.*(J. R. Neith. Chem. Soc.), 1950, **69**, 673-699 (synth)Li, T.S.C. *et al.*, *Chinese and Related North American Herbs*, CRC Press, 2002, 113 (occur)Schlosser, M. *et al.*, *J.O.C.*, 2005, **70**, 2494-2502 (di-Me)**4-Amino-1,13-diguanidino-5-tridecanone** A-730HN=C(NH₂)NH(CH₂)₈CO⁴CH(NH₂)CH₂CH₂CH₂NHC(NH₂)=NHC₁₅H₃₃N₇O 327.472N⁴-Ac: [167103-52-8]C₁₇H₃₅N₇O₂ 369.509

Isol. from the fermentation broth of an actinomycete (SCC 2268) prob. belonging to the genus *Streptomyces*. Muscarinic receptor antagonist. Sol. H₂O, MeOH; poorly sol. EtOAc, hexane. Mp 250°.

N⁴-(9-Guanidinononanoyl):C₂₅H₅₂N₁₀O₂ 524.752


Isol. from SCC 2268 broth. Muscarinic receptor antagonist. Sol. H₂O, MeOH; poorly sol. EtOAc, hexane. Mp 250°. [α]_D²⁶ -30 (c, 0.3 in H₂O).

Hegde, V.R. *et al.*, *J. Nat. Prod.*, 1995, **58**, 843 (isol, ir, pmr, cmr, struct)**2-Amino-4,5-dihydro-1H-imidazole-1-acetic acid, 9CI** A-731

2-Imino-1-imidazolidineacetic acid. 1-(Carboxymethyl)-2-iminoimidazolidine.

Cyclocreatine

[35404-50-3]

C₅H₉N₃O₂ 143.145Substrate for creatine kinase *in vitro*.

3-Phosphate: 1-(Carboxymethyl)-2-imi-

no-3-phosphonoimidazolidine


[61839-19-8]

C₅H₁₀N₃O₃P 223.125


Source of high energy phosphate. Stored in muscle, heart and brain.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1329B (nmr)Struve, G.E. *et al.*, *J.O.C.*, 1977, **42**, 4035 (synth, cmr, struct)Woznicki, D.T. *et al.*, *J. Neurochem.*, 1979, **33**, 75 (formn)Phillips, G.N. *et al.*, *J.A.C.S.*, 1979, **101**, 7120 (bibl, cryst struct, phosphate)Roberts, J.J. *et al.*, *Arch. Biochem. Biophys.*, 1983, **220**, 563 (activity)**2-Amino-4,5-dihydro-1H-imidazole-4-acetic acid, 9CI** A-732

2-[2-Amino-2-imidazolin-4-yl]acetic acid [69098-41-5]

C₅H₉N₃O₂ 143.145**(R)-form** [65388-06-9]Isol. from the seeds of *Lonchocarpus* spp. Cryst. (EtOH/Et₂O).Fellows, L.E. *et al.*, *Phytochemistry*, 1977, **16**, 1957 (isol, ir, pmr, cmr, ms, struct)Evans, S.V. *et al.*, *Biochem. Syst. Ecol.*, 1985, **13**, 271 (isol) **α -Amino-2,5-dihydro-5-oxo-4-isoxazolepropanoic acid, 9CI** A-733

α -Amino-5-oxo-3-isoxazoline-4-propionic acid, 8CI. 3-(Isoxazolin-5-on-4-yl)alanine. 4-Alanyl-3-isovalin-5-one. TAN 950A. Antibiotic TAN 950A

C₆H₈N₂O₄ 172.14

Exists in tautomeric equilibrium with Antibiotic TAN 950B. λ_{\max} 253 (ϵ 8060) (H₂O) (Derep).

(S)-form

L-form

[127607-88-9]

Prod. by *Streptomyces platensis*. Antifungal agent. Powder + 1H₂O (as Na salt). Sol. H₂O, DMSO, DMF; poorly sol. Me₂CO, CHCl₃, EtOAc. [α]_D²³ -69.5 (c, 0.52 in H₂O) (as Na salt). λ_{\max} 253 (E1%/1cm 380) (H₂O) (Berdy). λ_{\max} 259 (E1%/1cm 421) (HCl) (Berdy).

N²- β -D-Glucosyl: [29790-46-3]C₁₂H₁₈N₂O₉ 334.282


Isol. from *Pisum sativum* and *Lathyrus odoratus*. Cryst. (MeOH/propanol). Mp 182° dec.

[130620-29-0, 130621-41-9, 130621-42-0, 130621-61-3]

- Lambein, F. *et al.*, *Biochem. Biophys. Res. Commun.*, 1970, **40**, 557; 1974, **61**, 155 (*isol. struct.*, *uv. pmr.*, *ir*)
 Murakoshi, I. *et al.*, *Phytochemistry*, 1975, **14**, 1515 (*biosynth*)
 Iwama, T. *et al.*, *Eur. J. Pharmacol.*, 1991, **197**, 187 (*props*)
 Tsubotani, S. *et al.*, *Tetrahedron*, 1991, **47**, 8079-8090 (*synth. cryst. struct.*)
 Hakoda, S. *et al.*, *J. Antibiot.*, 1992, **45**, 854 (*isol*)

2-Amino-4-(2,5-dihydro-5-oxo-2-isoxazolyl)butanoic acid A-734

2-Amino-4-(isoxazolin-5-on-2-yl)butanoic acid
 [59476-62-9]



C₇H₁₀N₂O₄ 186.167

Incorrect (3-isoxazolyl) *struct.* in published work. *Isol.* from sweet pea extracts. *Cryst.* L-Config. not stated but assumed on biogenetic grounds.

- Lambein, F. *et al.*, *Biochem. Biophys. Res. Commun.*, 1974, **61**, 155 (*isol. struct.*)
 Kuo, Y.H. *et al.*, *Arch. Int. Physiol. Biochim.*, 1976, **84**, 169 (*isol*)
 Tamura, N. *et al.*, *Chem. Pharm. Bull.*, (Footnote to), 1992, **40**, 381 (*struct.*)


5-Amino-1,6-dihydro-7H-1,2,3-triazolo[4,5-d]pyrimidin-7-one, 9CI A-735

8-Azaguanine. Guanazole†. Pathocidin.

NSC 749. SF 337. Antibiotic SF 337.

Azan

[134-58-7]



C₄H₄N₆O 152.115

The name 8-Azaguanine comes from the alternative purine numbering. Systematic (CAS) numbering shown. *Isol.* from *Streptomyces albus* and *Streptomyces morookaensis*. Purine antagonist. Tumour inhibitor. Active against fungi. *Cryst.* (H₂O). *Mp* 305° dec. λ_{max} 250 (ε 12500); 270 (sh) (ε 9880) (0.1N HCl) (*Derep.*). λ_{max} 247 (ε 4860); 278 (ε 6540) (0.1N NaOH) (*Derep.*). λ_{max} 247 (ε 9580); 273 (ε 6300) (H₂O) (*Derep.*)

- *Exp. reprod.* and teratogenic effects. LD₅₀ (mus, orl) 1500 mg/kg. XZ6157000

3N-β-D-Ribofuranosyl: 8-Azaguanosine [2133-80-4]
 C₉H₁₂N₆O₅ 284.231
 Purine antagonist. *Mp* 250-252° dec.

λ_{max} 255 (ε 13600); 269 (ε 10300) (0.01N HCl). λ_{max} 221 (ε 23000); 279 (ε 11600) (0.03N NaOH).

► XZ6157200

3N-β-D-Ribofuranosyl, 5'-phosphate: 8-Azaguanosine 5'-phosphate

[1165-66-8]

C₉H₁₃N₆O₈P 364.211

Appears to be the key intermed. in the conversion of the anticancer drug, 8-Azaguanine, into its active form. λ_{max} 256 (ε 12900) (pH 2).

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **2**, 716C (*ir*)

Roblin, R.K. *et al.*, *J.A.C.S.*, 1945, **67**, 290-294 (*synth*)

Friedkin, M. *et al.*, *J. Biol. Chem.*, 1954, **209**, 295-301 (8-Azaguanosine)

Davoll, J. *et al.*, *J.C.S.*, 1958, 1593-1599 (8-Azaguanosine)

Blank, H.U. *et al.*, *J.O.C.*, 1970, **35**, 1131-1138 (*synth*)

Crit. Rev. Toxicol., 1973, **2**, 159-209 (*tox. rev*)

Kozlowski, D.L. *et al.*, *Acta Cryst. B*, 1975, **31**, 1751-1753 (*cryst. struct.*)

Elliott, R.D. *et al.*, *J. Med. Chem.*, 1976, **19**, 1186-1191 (8-Azaguanosine)

Luedemann, H.D. *et al.*, *Z. Naturforsch., C*, 1976, **31**, 135-140; *CA*, **84**, 146335u (8-Azaguanosine, *conform.*, *pmr*)

Bose, S.N. *et al.*, *Biomed. Mass Spectrom.*, 1977, **4**, 305-309 (*ms*)

Purnell, L.G. *et al.*, *Org. Magn. Reson.*, 1977, **10**, 1-4 (*cmr*)

Hirasawa, K. *et al.*, *J. Antibiot.*, 1978, **31**, 628-629 (*isol. props*)

Grunberger, D. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 110-123 (*rev*)


Contreras, J.G. *et al.*, *Bioorg. Chem.*, 1998, **26**, 345-355 (*tautom.*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, AJO500

2-Amino-3',4'-dihydroxyacetophenone, 8CI A-736

2-Amino-1-(3,4-dihydroxyphenyl)ethanone. Noradrenalone. Arterenone. 2-Oxodopamine

[499-61-6]



C₈H₉NO₃ 167.164

Isol. from hydrolysates of sclerotized insect cuticle.

► AM5900000

N-Ac: 2-Acetamido-3',4'-dihydroxyacetophenone. N-Acetyl-2-oxodopamine
 C₁₀H₁₁NO₄ 209.201

Isol. from the cast-off shells of the cicada, *Cryptotympana* sp. Yellow *cryst.* *Mp* 168-175°.

N-Me: 1-(3,4-Dihydroxyphenyl)-2-(methylamino)ethanone. 3',4'-Dihydroxy-2-methylaminoacetophenone. Adrenolone, INN; USAN. Chemosan. Damazor. Haemodan. Kephriane. Ketogaze. Remestyp†. Stryphnon. Stypnon. Adrenone

[99-45-6]

C₉H₁₁NO₃ 181.191

Shows similar physiological activity to Adrenaline, A-152. Sympathomimetic, vasoconstrictor and haemostatic agent. Needleless. *Mp* 235-236° dec. *Log P* 0.32 (*calc.*)

- LD₅₀ (mus, ivn) 275 mg/kg. AM8050000

N-Me; hydrochloride: [62-13-5]

Mp 242°.

- AM8225000

Stolz, F. *et al.*, *Ber.*, 1904, **37**, 4152

Friedmann, E. *et al.*, *Chem. Zentralbl.*, 1906, **1**, 1620 (*synth*)

Lands, A.M. *et al.*, *J. Pharmacol. Exp. Ther.*, 1948, **92**, 369 (*pharmacol*)

Ravina, A. *et al.*, *Therapie*, 1953, **8**, 224 (*props. use*)

Remizov, A.L. *et al.*, *Zh. Obshch. Khim.*, 1958, **28**, 2530 (*synth*)

Binięcki, S. *et al.*, *Acta Pol. Pharm.*, 1963, **20**, 245 (*synth*)

Anderson, S.O. *et al.*, *J. Insect Physiol.*, 1970, **16**, 1951 (*isol*)

Bergin, R. *et al.*, *Acta Cryst. B*, 1971, **27**, 2139 (*N-Me, cryst. struct.*)

Hussain, A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 1510 (*deriv*)

Martindale, *The Extra Pharmacopoeia, 30th edn.*, Pharmaceutical Press, 1993, 1240


Noda, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1749-1752 (*isol, N-Ac*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MGC350

2-Amino-4,5-dihydroxybenzoic acid A-737

6-Aminoprotocatechuic acid. 4,5-Dihydroxyanthranilic acid

[114874-99-6]



C₇H₇NO₄ 169.137

Prod. by *Botrytis* sp. AM13791. Inhibitor of protein-tyrosine kinase.

N-Et, amide: 2-(Ethylamino)-4,5-dihydroxybenzamide, 9CI
 [127793-87-7]

C₉H₁₂N₂O₃ 196.205

Alkaloid from *Piper nigrum* (pepper).

4-Me ether: 2-Amino-5-hydroxy-4-methoxybenzoic acid. 2-Amino-5-hydroxy-p-anisic acid, 8CI
 [31839-21-1]

C₈H₉NO₄ 183.163

Mp 214-215° (194-195°).


4-Me ether, N-(4-hydroxy-3-methoxy-E-cinnamoyl): Avenanthramide 2
 [154992-25-3]

C₁₈H₁₇NO₇ 359.335

Constit. of *Avena sativa* (oat).

Bandyopadhyay, C. *et al.*, *J. Agric. Food Chem.*, 1990, **38**, 1696-1699 (*N-Et amide*)

Dimberg, L.H. *et al.*, *Cereal Chem.*, 1993, **70**, 637-641 (*Avenanthramide 2*)

3-Amino-6,7-dihydroxy-2H-1-benzopyran-2-one, 9CI A-7383-Amino-6,7-dihydroxycoumarin
[22065-08-3]C₉H₇NO₄ 193.159

Shows hypotensive props. Mp 258-260°.

N-Formyl: 3-(Formylamino)-6,7-dihydroxycoumarin. **Pseudoverdin**
[150624-46-7]
C₁₀H₇NO₅ 221.169
Prod. by *Pseudomonas aeruginosa*.
Chromophore.

Di-Me ether: 3-Amino-6,7-dimethoxy-2H-1-benzopyran-2-one. 3-Amino-6,7-dimethoxycoumarin
[150358-93-3]
C₁₁H₁₁NO₄ 221.212
Cryst. (EtOH aq.). Mp 170°.

Di-Me ether. *N*-formyl: [150358-91-1]
C₁₂H₁₁NO₅ 249.223
Mp 246°.


Di-Me ether, *N*-Ac: 3-(Acetylamino)-6,7-dimethoxycoumarin
[150358-92-2]
C₁₃H₁₃NO₅ 263.249
Cryst. (EtOH aq.). Mp 248°.

Fr. Pat., 1968, 1 523 317; *CA*, **72**, 31615v (synth)

Longerich, I. et al., *Z. Naturforsch.*, **C**, 1993, **48**, 425 (isol, synth, derivs)

4-Amino-2,5-dihydroxybenzyl alcohol A-739

2-Amino-5-(hydroxymethyl)-1,4-benzenediol

C₇H₉NO₃ 155.153

N-Ac: 2,5-Dihydroxy-4-(hydroxymethyl)acetanilide

C₉H₁₁NO₄ 197.19

Prod. by *Actinomyces* sp. Lu 9419
isol. from *Cetonia aureata*. Solid.
Mp 176°. λ_{max} 209 (ε 12300); 245
(ε 4840); 302 (ε 3400)
(MeOH).


l'-Me ether, *N*-Ac: 2,5-Dihydroxy-4-

(methoxymethyl)acetanilide

C₁₀H₁₃NO₄ 211.217

Prod. by *Actinomyces* sp. Lu 9419
isol. from *Cetonia aureata*. Solid.

Schlörke, O. et al., *J. Antibiot.*, 2002, **55**,
635-642 (*N*-Ac, isol, pmr, cmr, ms)

26-Amino-3,16-dihydroxy-cholestan-22-one A-740C₂₇H₄₇NO₃ 433.673**(3β,5α,16α,25ξ)-form**

N-Ac, 3-O-[β-D-xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Abutiloside B**
[177262-59-8]

C₄₆H₇₇NO₁₇ 916.111

Constit. of *Solanum abutiloides*. Solid.
[α]_D²⁵ -41.9 (c, 0.49 in MeOH).

N-Butanoyl, 3-O-[α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Abutiloside K**
[449178-14-7]

C₄₃H₇₃NO₁₃ 812.049

Constit. of *Solanum abutiloides*. Powder.
[α]_D²⁵ -50.4 (c, 0.25 in MeOH).

N-Butanoyl, 3-O-[β-D-xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Abutiloside J**
[449176-35-6]

C₄₈H₈₁NO₁₇ 944.164

Constit. of *Solanum abutiloides*. Powder.
[α]_D²⁵ -54.1 (c, 0.95 in MeOH).

N-(3-Methylbutanoyl), 3-O-[α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Abutiloside I**
[449176-33-4]

C₄₄H₇₅NO₁₃ 826.075

Constit. of *Solanum abutiloides*.
Powder. [α]_D²⁵ -38.7 (c, 0.15 in
MeOH).

N-(3-Methylbutanoyl), 3-O-[β-D-xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]:
Abutiloside A

[171864-79-2]


C₄₉H₈₃NO₁₇ 958.191

Constit. of *Solanum abutiloides*.
Amorph. powder. [α]_D²² -49.2 (MeOH).

Tian, R.-H. et al., *Chem. Pharm. Bull.*, 1996,
44, 1119-1121 (isol, pmr, cmr)

Tian, R.-H. et al., *Phytochemistry*, 1997, **44**,
723-726 (*Abutiloside A*)


Yoshimitsu, H. et al., *Chem. Pharm. Bull.*,
2002, **50**, 284-286 (*Abutilosides I, J, K*)

26-Amino-3,16-dihydroxy-cholest-5-en-22-one A-741C₂₇H₄₅NO₃ 431.657**(3β,16α,26ξ)-form**

N-Ac, 3-O-[β-D-xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Abutiloside H**
[449176-31-2]

Constit. of *Solanum abutiloides*. Powder.
[α]_D²⁵ -107 (c, 0.2 in MeOH).

Yoshimitsu, H. et al., *Chem. Pharm. Bull.*,
2002, **50**, 284-286

4-Amino-2-(2,3-dihydroxy-3-methylbutyl)benzoic acid A-742C₁₂H₁₇NO₄ 239.271**(-)-form**

Me ester: [1009318-74-4]

C₁₃H₁₉NO₄ 253.297

Isol. from *Xylaria* sp. BCC 9653. Gum.
[α]_D²⁷ -19.2 (c, 0.9 in CHCl₃).

Pongcharoen, W. et al., *Chem. Pharm. Bull.*,
2007, **55**, 1647-1648 (isol, pmr, cmr, ms)

2-Amino-1,7-dihydroxy-phenanthrene A-743

2-Amino-1,7-phenanthrenediol, 9CI

C₁₄H₁₁NO₂ 225.246


Di-Me ether, *N*-Ac: *N*-(1,7-Dimethoxy-2-phenanthrenyl)acetamide. 2-Acetamido-1,7-dimethoxyphenanthrene
C₁₈H₁₇NO₃ 295.337

Alkaloid from *Sinomenium acutum*.
Yellow-green needles. λ_{max} 256 ; 282 ;
310 (no solvent reported).

Cheng, W.M. et al., *Chin. Chem. Lett.*, 2005, **16**,
1481-1483 (*di*-Me ether *N*-Ac, isol, pmr, cmr)

3-Amino-2-(3,4-dihydroxyphenyl)-2,3-dihydro-1,4-benzodioxin-6-ethanamine A-744

2-Amino-7-(2-aminoethyl)-3-(3,4-dihydroxyphenyl)-1,4-benzodioxan

C₁₆H₁₈N₂O₄ 302.329

Dimer of Dopamine, D-920.

(2R,3S)-form*N,N'*-Di-Ac:


C₂₀H₂₂N₂O₆ 386.404

Isol. from the cast-off shells of the cicada *Cryptotympana* sp. Component of Zentai. Yellow powder. Mp 118-126°. [α]_D +51.7 (c, 1 in MeOH).

Noda, N. et al., *Chem. Pharm. Bull.*, 2000, 48, 1749-1752 (isol, pmr, cmr)

2-Amino-3-(3,4-dihydroxyphenyl)propanoic acid A-745

3-Hydroxytyrosine, 9CI. 3,4-Dihydroxyphenylalanine. DOPA [587-45-1]

C₉H₁₁NO₄ 197.19

Used in treatment of the Parkinsonian syndrome. The immediate precursor of the neurotransmitter dopamine. Usually administered with a peripheral DOPA decarboxylase inhibitor. Intermed. in biosynth. of various marine alkaloids, e.g. the Lamellarins. Log P -2.82 (uncertain value) (calc).

(R)-form

D-form

[5796-17-8]

Prisms (H₂SO₃ aq.). Mp 282°. [α]_D +11.9.

(S)-form

L-form. *Levodopa*, **BAN**, **INN**, **JAN**, **USAN**. *Bendopa*. *Brocadopa*. *Dopar*. *Larodopa*. *Levopa*. *Veldopa*. Many other names

[59-92-7]

Occurs in seedlings and pods of *Vicia faba*, in *Mucuna pruriens*, *Sarothamnus scoparius*, *Stizolobium deeringianum*, *Stizolobium hassjoo*, *Aristolochia clematidis* and other plants. Also prod. by *Bacillus* spp. Prisms, needles (H₂O + SO₂) or plates (EtOH aq.). Mp 285.5° dec. [α]_D²⁰ -12.15 (c, 4 in 1N HCl). Pharmacol. active isomer. Rapidly turns green in air, aq. solns. darken in air. Green col. with FeCl₃. Reduces AgNO₃, NH₃ and acid KMnO₄. Component of Sinemet. High-mol.-wt. polymers have been synth.

► Systemic (e.g. CNS) and adverse effects when used therapeutically. LD₅₀ (rat, orl) 1780 mg/kg. Exp. reprod. and teratogenic effects. AY5600000

Hydrochloride: [5796-14-5]

Prisms. Mp 209°.

3'-O-Sulfate: 3-(Sulfooxy)tyrosine, 9CI [96253-55-3]

C₉H₁₁NO₇S 277.254

Constit. of the brown alga *Ascophyllum nodosum*.

3'-O-β-D-Glucopyranoside: [2275-95-8]

C₁₅H₂₁NO₉ 359.332

Isol. from *Pisum sativum* (peas) and *Vicia faba*. Resin or syrup.

Me ester: *Melevodopa*, **INN**. *Levomet*

[7101-51-1]

[1421-65-4]

C₁₀H₁₃NO₄ 211.217

Antiparkinsonian agent. Dopamine receptor agonist. Mp 170.5-171.5° (as hydrochloride). [α]_D²² +14.7 (c, 12.5 in MeOH) (hydrochloride). CAS no. refers to the hydrochloride. Prodrug of levodopa.

Et ester: *Etilevodopa*, **USAN**

[37178-37-3]

C₁₁H₁₅NO₄ 225.244

Sol. prodrug of levodopa.

Et ester, hydrochloride: [39740-30-2]

Cryst. (Et₂O/EtOH). Mp 126-129°.[α]_D²² -8.95 (c, 1.05 in H₂O).

N-Ac: N-Acetyl-3,4-dihydroxyphenylalanine

[19641-90-8]

C₁₁H₁₃NO₅ 239.227

Prod. by *Streptomyces akiyoshiensis* ATCC13480. Sol. H₂O, MeOH. [α]_D²⁰ +68.2 (c, 1 in MeOH). λ_{\max} 198 (ε 34200); 280 (ε 2210) (H₂O) (Berdy).

O,O,N-Tribenzoyl: [97857-68-6]

C₃₀H₂₃NO₇ 509.514

Needles (AcOH). Mp 170°.

N-(4-Hydroxy-E-cinnamoyl): N-trans-p-Coumaroyl-DOPA

[77201-64-0]

C₁₈H₁₇NO₆ 343.335

Constit. of the bark of *Dalbergia melanoxylon* and of roasted cocoa powder.

N-(4-Hydroxy-Z-cinnamoyl): N-cis-p-Coumaroyl-DOPA

[77201-63-9]

C₁₈H₁₇NO₆ 343.335

Constit. of the bark of *Dalbergia melanoxylon*.

N-(4-Methylbenzenesulfonyl): [37169-34-9]

C₁₆H₁₇NO₆S 351.379

Cryst. (EtOAc/petrol). Mp 188°. [α]_D²⁵ -5.4 (c, 1 in MeOH).

N-tert-Butyloxycarbonyl: [30033-24-0]

C₁₄H₁₉NO₆ 297.307

Solid (EtOAc/cyclohexane). Mp 148° (142-144°). [α]_D²⁵ +16.4 (c, 1 in MeOH).

N-tert-Butyloxycarbonyl, Me ester:

[37169-36-1]

C₁₅H₂₁NO₆ 311.334

Solid (MeOH aq.). Mp 140-141° (133-135°). [α]_D²⁵ +12.3 (c, 1 in MeOH) (+7.6).

N-tert-Butyloxycarbonyl, benzyl ester:

[37169-37-2]

C₂₁H₂₅NO₆ 387.432

Brown glass. [α]_D²⁵ -4.2 (c, 1 in MeOH).

N-(9-Fluorenylmethyloxycarbonyl):

Fmoc-L-DOPA

[137018-93-0]

C₂₄H₂₁NO₆ 419.433

Solid.

N,N,N-Tri-Me, betaine: α-Carboxy-3,4-

dihydroxy-N,N,N-trimethylbenzene-

neethanaminium(1+). DOPA betaine

[81130-72-5]

C₁₂H₁₈NO₄⁺ 240.279

Constit. of *Lobaria laetevirens*.

N,N,N-Tri-Me, Me ester: 3,4-Dihydroxy-

α-(methoxycarbonyl)-N,N,N-tri-

methylbenzeneethanaminium(1+), 9CI.

Sticticin

[77035-53-1]

C₁₃H₂₀NO₄⁺ 254.305

Isol. from the thallus of *Lobaria laetevirens* and present in other lichens of the Stictaceae. V. hygroscopic solid (as chloride).

3'-Me ether: 4-Hydroxy-3-methoxyphenylalanine. 3-Methoxytyrosine

[300-48-1]

C₁₀H₁₃NO₄ 211.217

Isol. from *Cortinarius brunneus*, *Pachymatisma johnstoni* and the blood of Parkinsonian patients. Cryst. (H₂O). [α]_D -34 (c, 0.39 in H₂O). [α]_D -15.4 (c, 0.36 in 0.1M HCl).

5,5'-Dimer: α,α'-Diamino-5,5',6,6'-tetrahydroxy-3,3'-biphenyldipropanoic acid, 8CI

[17785-52-3]

C₁₈H₂₀N₂O₈ 392.365

Powder (H₂O containing SO₂). Mp 300°.

N-Jasmonoyl: N-Jasmonoyl DOPA

[866421-54-7]

C₂₁H₂₇NO₆ 389.447

Constit. of the flowers of *Vicia faba* (broad bean).

(±)-form [63-84-3]

Cryst. (H₂O or NaHSO₃ aq.). Mp 271-272° dec.

► AY5250000

Hydrochloride: [63302-79-4]

Plates (MeOH). Mp 246° dec.

Hydrobromide:

Plates (MeOH). Mp 212° dec.

Me ester: [41439-83-2]

Mp 126°.

Me ester, hydrochloride: [40611-00-5]

Mp 180-181°.

Et ester: [23234-42-6]

Cryst. (EtOH aq.). Mp 129°.

O,O,N-Tri-Ac: [47302-76-1]

C₁₅H₁₇NO₇ 323.302

Prisms (H₂O). Mp 171-172°.

N-Benzoyl: [38250-04-3]

C₁₆H₁₅NO₅ 301.298

Cryst. Mp 190-195° (anhyd.).

N-tert-Butyloxycarbonyl: [59686-55-4]

Solid (EtOAc/C₆H₆). Mp 140-142°.

N-tert-Butyloxycarbonyl, Me ester:

[59686-54-3]

Solid (MeOH aq.). Mp 186-188°.

N-tert-Butyloxycarbonyl, di-Ac: [59686-

56-5]

C₁₈H₂₃NO₈ 381.382

Solid. Mp 132-134°.

3'-Me ether: [4214-13-5]

Prisms + 2H₂O (H₂O). Mp 255-256° dec.

3'-Me ether, N-benzoyl: [2901-78-2]

C₁₇H₁₇NO₅ 315.325

Plates (H₂O or EtOH). Mp 164°.

3',4'-Di-Me ether: [33522-62-2]

C₁₁H₁₅NO₄ 225.244

Needles (EtOH aq.). Mp 252-255°.

3',4'-Di-Me ether, N-benzoyl: [34996-90-

2]

C₁₈H₁₉NO₅ 329.352

Mp 173-174°.

[57308-51-7, 127441-81-0]


Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 257A; 257B (ir)

Aldrich Library of 13C and 1H

FT NMR Spectra, 1992, 2, 1190A; 1190B (nmr)

O'Neill, J.J. et al., *J.O.C.*, 1956, 21, 363-364 (Me ester)Losse, G. et al., *Chem. Ber.*, 1961, 94, 2271-2277 (Et ester, synth)Andrews, R.S. et al., *Nature (London)*, 1965, 205, 1213 (glucoside)Kaiser, A. et al., *Helv. Chim. Acta*, 1970, 53, 1708-1712 (N-Ac, Boc-L-DOPA, synth, pmr)Chapman, R.F. et al., *J.C.S.(C)*, 1970, 865 (dimer)Vorbruggen, H. et al., *Chem. Ber.*, 1972, 105, 1168 (synth)


Ger. Pat., 1972, 2 153 811; CA, 77, 62311a (Boc derivs)

Griffith, T. et al., *Phytochemistry*, 1973, 12, 1651 (biosynth)Mostad, A. et al., *Acta Chem. Scand., Ser. B*, 1974, 28, 1161 (cryst struct)Renth, E.-O. et al., *Angew. Chem., Int. Ed.*, 1975, 14, 361 (synth)Fellman, J.H. et al., *Biochim. Biophys. Acta*, 1975, 381, 9 (isol, deriv)Bartholini, G. et al., *Pharmacol. Ther., Part B*, 1975, 1, 407 (rev, pharmacol)Gomez, R. et al., *Anal. Profiles Drug Subst.*, 1976, 5, 189 (rev)Bannerjee, S.N. et al., *J.O.C.*, 1976, 41, 3056-3058 (Boc derivs)Boder, N. et al., *J. Med. Chem.*, 1977, 20, 1435-1445 (S-form Me ester, synth, pmr)Dardenne, G. et al., *Phytochemistry*, 1977, 16, 1822 (deriv)Yamamoto, H. et al., *Polymer*, 1977, 18, 979; 1978, 19, 1115 (polym)Fuller, W.D. et al., *Biopolymers*, 1978, 17, 2939 (polym)Rudd, E.A. et al., *J. Med. Chem.*, 1979, 22, 233-237 (Boc-L-DOPA Me ester)Bernard, T. et al., *Phytochemistry*, 1980, 19, 1967 (Sticticin)Van Heerden, F.R. et al., *Phytochemistry*, 1980, 19, 2125 (N-p-coumaroyl)Bernard, T. et al., *Phytochemistry*, 1981, 20, 2325 (betaine)Danishevsky, S. et al., *Tetrahedron*, 1981, 37, 4081 (synth)Nutt, J.G. et al., *Clin. Neuropharmacol.*, 1984, 7, 35 (rev, metab)Laycock, M.V. et al., *J. Nat. Prod.*, 1984, 47, 1033 (isol, sulfate)Lee, M. et al., *Chem. Pharm. Bull.*, 1987, 35, 235 (hplc, bibl, anal)Cooper, D.R. et al., *J. Pharm. Pharmacol.*, 1987, 39, 627-635; 809-818 (Et ester)Behrman, E.J. et al., *Org. Prep. Proced. Int.*, 1989, 21, 351 (synth, sulfate)Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 1471 (synonyms)Smith, K.C. et al., *J. Nat. Prod.*, 1995, 58, 1274 (isol, N-Ac)Brunner-Guenat, M. et al., *J. Pharm. Pharmacol.*, 1995, 47, 861 (Me ester)Djalalietti, R. et al., *Ann. Neurol.*, 1996, 39, 400-404 (Et ester)Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1137Chen, F.-Y. et al., *J.O.C.*, 2001, 66, 3650-3652 (R-form, ir, pmr, cmr)Sever, M.J. et al., *Tetrahedron*, 2001, 57, 6139-6146 (Fmoc/TBDMS derivs)Kramell, R. et al., *J. Nat. Prod.*, 2005, 68, 1345-1349 (N-jasmonoyl)Stark, T. et al., *J. Agric. Food Chem.*, 2006, 54, 2859-2867 (N-4-hydroxycinnamoyl, isol)Aubry, S. et al., *Eur. J. Org. Chem.*, 2007, 5212-5225 (S-form, Me ester)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DNA200**6-Amino-4,5-dihydroxy-3-piperidinecarboxylic acid** A-746C₆H₁₂N₂O₄ 176.172**(3S,4S,5R,6S)-form**N-Ac: **Siastatin B**. A 72363B. Antibiotic A 72363B


[54795-58-3]

C₈H₁₄N₂O₅ 218.209Isol. from *Streptomyces verticillus* var. *quintum* and *Streptomyces nobilis*. Potent neuraminidase inhibitor. Needles. Mp 137° dec. [α]_D²⁵ +57.2 (c, 1 in H₂O).

▶ TM6125400

(3R*,4R*,5R*,6R*)-formN-Ac: **Antibiotic A 72363A1**. A 72363A-1C₈H₁₄N₂O₅ 218.209Prod. by *Streptomyces nobilis*. Heparanase inhibitor. Sol. H₂O; poorly sol. Me₂CO, hexane. [α]_D²³ +29 (c, 1 in H₂O).**(3R*,4S*,5R*,6R*)-form**N-Ac: **Antibiotic A 72363A2**. A 72363A-2C₈H₁₄N₂O₅ 218.209Prod. by *Streptomyces nobilis*. Heparanase inhibitor. Sol. H₂O; poorly sol. Me₂CO, hexane. [α]_D²³ -31 (c, 1 in H₂O).**(3R*,4R*,5S*,6S*)-form**N-Ac: **Antibiotic A 72363C**. A 72363CC₈H₁₄N₂O₅ 218.209Prod. by *Streptomyces nobilis*. Heparanase inhibitor. Sol. H₂O; poorly sol. Me₂CO, hexane. [α]_D²⁷ -59 (c, 0.7 in H₂O).Umezawa, H. et al., *J. Antibiot.*, 1974, 27, 963 (isol, ir, props)Aoyagi, T. et al., *Experientia*, 1975, 31, 896 (struct)Nishimura, Y. et al., *J.A.C.S.*, 1988, 110, 7249 (synth, abs config)Nishimura, Y. et al., *Bull. Chem. Soc. Jpn.*, 1992, 65, 978 (synth)Nishimura, Y. et al., *Stud. Nat. Prod. Chem.*, 1995, 16, 75 (rev, synth)Takatsu, T. et al., *J. Antibiot.*, 1996, 49, 54; 61 (A 73263, pmr, cmr, activity)Knapp, S. et al., *Org. Lett.*, 2000, 2, 4037-4040 (*Siastatin B*, synth)**3-Amino-11,12-dihydroxy-pregnan-20-one** A-747C₂₁H₃₅NO₃ 349.512**(3β,5α,11α,12β)-form**N,N-Di-Me, 11-O-(3,4-dimethyl-3-pentenoyl), 12-Ac: **Pachysanonin**


[873799-27-0]

C₃₂H₅₁NO₅ 529.759Constit. of *Pachysandra axillaris*.Cryst. (Me₂CO). Mp 153.5-154.5°.Qiu, M.-H. et al., *Chem. Biodiversity*, 2005, 2, 866-871 (*Pachysanonin*)**2-Amino-7-(1,2-dihydroxypropyl)-1,5-dihydro-4,6-pteridine-dione, 9CI** A-748**6-Oxoprimapterin**. 6-Oxo-7-biopterin [115991-76-9]C₉H₁₁N₅O₄ 253.217

Occurs in the urine of humans with hyperphenylalaninemia.

Curtuis, H.C. et al., *J. Biol. Chem.*, 1990, 265, 3923-3930 (isol, ms, struct)**2-Amino-6-(1,2-dihydroxypropyl)-4(1H)-pteridinone** A-749

2-Amino-4-hydroxy-6-(1,2-dihydroxypropyl)pterin

C₉H₁₁N₅O₃ 237.218**(1'R,2'R)-form***D*-threo-form. **Dictyopterin**

[13019-52-8]

Isol. from *Dictyostelium discoideum*. Mp 300°. pK_{a1} 2.2; pK_{a2} 7.92.**(1'R,2'S)-form***L*-erythro-form. **Biopterin**. *Ranachrome 1* [22150-76-1]

Widely distributed in microorganisms, insects, algae, amphibia and mammals.

Found in urine. Growth factor. Pale yellow cryst. (AcOH aq.). Mp 250-280° dec. [α]_D²⁰ -66 (c, 0.2 in 0.1M HCl). pK_{a1} 2.23; pK_{a2} 7.89.

▶ UO3506000

1'-O-*D*-Glucopyranoside: *Biopterin glucose*

[32838-67-8]

C₁₅H₂₁N₅O₈ 399.36

Constit. of a marine plankton.

2'-O-*α*-*D*-Glucopyranoside: [235416-12-3]C₁₅H₂₁N₅O₈ 399.36Isol. from *Spirulina (Arthrospira) platensis*.2'-O-(2-Acetamido-2-deoxy-β-*D*-glucopyranoside): **Limipterin**

[164803-20-7]
C₁₇H₂₄N₆O₈ 440.412

Prod. by *Chlorobium limicola* f. *thio-sulfatophilum*.

3-Me: 2-Amino-6-(1,2-dihydroxypropyl)-3-methylpterin-4-one

[111317-37-4]
C₁₀H₁₃N₅O₃ 251.244

Isol. from the marine anthozoan *Asroides calycularis*. Pale yellow cryst. powder (MeOH). Sol. MeOH, H₂O. Mp 229-231°. [α]_D -60 (c, 0.3 in 0.1M HCl). λ_{max} 230 (ε 28200); 319 (ε 11700) (pH 1 H₂O) (Derep). λ_{max} 242 (ε 28200); 276 (ε 24000); 356 (ε 9550) (H₂O pH 7) (Derep).

7,8-Dihydro: 7,8-Dihydrobiopterin

[6779-87-9]
[7644-44-2]
C₉H₁₃N₅O₃ 239.233

Hygroscopic needles (H₂O or MeOH aq.). [α]_D²² +35 (c, 0.4 in 0.1M NaOH).

5,6,7,8-Tetrahydro: see 5,6,7,8-Tetrahydrobiopterin, T-142

(1'S,2'R)-form

D-erythro-form
[13039-62-8]
Mp 300°. pK_{a1} 2.23; pK_{a2} 7.9.

(1'S,2'S)-form

L-threo-form. **Orinapterin. Ciliapterin**
[13039-82-2]
Isol. from human urine. Prod. by *Aphanizomenon flos-aquae*. Pale yellow cryst. (AcOH aq.). Mp 300°. [α]_D +72 (0.1M HCl). pK_{a1} 2.24; pK_{a2} 7.87.

2'-O-(2-Acetamido-2-deoxy-β-D-glucopyranoside): Tepidopterin

[188778-28-1]
C₁₇H₂₄N₆O₈ 440.412
Prod. by *Chlorobium tepidum*.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 480B (nmr)

Patterson, E.L. *et al.*, *J.A.C.S.*, 1955, **77**, 3167
Kidder, G.W. *et al.*, *J. Biol. Chem.*, 1968, **243**, 826 (*Ciliapterin*)

Kidder, G.W. *et al.*, *Methods Enzymol.*, 1971, **18B**, 739 (*Ciliapterin*)

Rembold, H. *et al.*, *Angew. Chem., Int. Ed.*, 1972, **11**, 1061 (*biochem*)

Blair, J.A. *et al.*, *Tet. Lett.*, 1973, 203-204 (*synth, uv, dihydro*)

Sugimoto, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 3767 (*synth*)

Taylor, E.C. *et al.*, *J.A.C.S.*, 1976, **98**, 2301 (*synth*)

Schircks, B. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 211; 1978, **61**, 2731-2738; 1985, **68**, 1639 (*synth, pmr, cmr, derivs*)

Fukushima, T. *et al.*, *Methods Enzymol.*, 1980, **66**, 508-511 (*synth, uv, dihydro*)

Armarego, W.L.F. *et al.*, *Aust. J. Chem.*, 1982, **35**, 785 (*synth*)

Kappel, M. *et al.*, *Annalen*, 1984, 1815 (*synth*)
Aiello, A. *et al.*, *Experientia*, 1987, **43**, 950 (*deriv*)

Mori, K. *et al.*, *Annalen*, 1989, 958; 1212 (*synth, bibl*)

Klein, R. *et al.*, *Eur. J. Biochem.*, 1990, **187**, 665 (*Dictyopterine*)

Ogiwara, S. *et al.*, *Biol. Chem. Hoppe-Seyler*, 1992, **373**, 1061 (*Orinapterin*)

Cha, K.W. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 600-614 (*Limipterin*)

Ikawa, M. *et al.*, *Phytochemistry*, 1995, **38**, 1229 (*isol, pmr, ms*)

Fernandez, A.M. *et al.*, *J.O.C.*, 1996, **61**, 8698 (*synth*)

Cho, S.-H. *et al.*, *Biochim. Biophys. Acta*, 1998, **1379**, 53-60 (*Tepidopterine*)


Noguchi, Y. *et al.*, *Mar. Biotechnol.*, 1999, **1**, 207-210 (*2'-glucoside*)

Hanaya, T. *et al.*, *Tetrahedron*, 2008, **64**, 2090-2100 (*Limipterin, Tepidopterine, synth*)

2-Amino-4,6-dihydroxypteridine A-750

2-Amino-1,5-dihydro-4,6-pteridinedione, 9CI. Xanthopterine. Uropterine

[119-44-8]
[5979-01-1]



C₆H₅N₅O₂ 179.138

Important naturally occurring pteridine. Normal constit. of human urine. Yellow wing pigment of butterflies e.g. *Gonepteryx rhamni*, *Appias nero*, *Colias edusa*. Inhibits cell proliferation. Sol. acids, alkalis. Mp 410° (carbonises above 360°). pK_{a1} 6.25; pK_{a2} 9.23 (20°).

6-O-Sulfate: Xanthopterinsulfonic acid

C₆H₅N₅O₅S 259.202
Prod. by *Azotomonas insolita* and *Escherichia coli*. Needles (H₂O) (as Na salt). Mp 300° (Na salt).

Schöpf, C. *et al.*, *Annalen*, 1933, **507**, 266
Goto, M. *et al.*, *Arch. Biochem. Biophys.*, 1965, **111**, 8 (*Xanthopterinsulfonic acid*)

Taylor, E.C. *et al.*, *J.A.C.S.*, 1973, **95**, 4455 (*synth*)

Bieri, J.H. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2374 (*cryst struct*)


Pfleiderer, W. *et al.*, *J. Het. Chem.*, 1992, **29**, 583 (*rev*)

Rubenstein, M. *et al.*, *Int. J. Biochem.*, 1993, **25**, 1873

2-Amino-4,7-dihydroxypteridine A-751

2-Amino-4,7(1H,8H)-pteridinedione, 9CI. Isoxanthopterine. Ranachrome 4

[529-69-1]



C₆H₅N₅O₂ 179.138

Widespread insect pigment found in amphibian and fish skin; normal constit. of urine. Mp 300°. pK_{a1} 7.34; pK_{a2} 10.06 (20°, H₂O).

► UO3425000

1H,3H-form

Me ether, 1,3-di-Me: N¹,N³,O⁷-Tri-methylisoxanthopterine

C₉H₁₁N₅O₂ 221.218

Isol. from a *Eudistoma* sp. Solid. λ_{max} 208 (sh) (log ε 4); 236 (sh) (log ε 3.7); 268 (log ε 3.4); 316 (log ε 3.6); 328 (sh) (log ε 3.6); 350 (sh) (log ε 3) (MeOH).

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 769A (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 895D (*ir*)

Pfleiderer, W. *et al.*, *Chem. Ber.*, 1961, **94**, 1 (*struct, bibl, uv*)

Konrad, G. *et al.*, *Chem. Ber.*, 1970, **103**, 735 (*uv, props*)

Taylor, E.C. *et al.*, *J.O.C.*, 1975, **40**, 2341 (*synth, bibl*)


Pfleiderer, W. *et al.*, *J. Het. Chem.*, 1992, **29**, 583 (*rev*)

Van Wagoner, R.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1100-1101 (*N,N,O-trimethyl*)

2-Amino-4,7-dihydroxy-6-pteridinedicarboxylic acid A-752

2-Amino-1,4,7,8-tetrahydro-4,7-dioxo-6-pteridinedicarboxylic acid, 9CI. Iso-xanthopterincarboxylic acid, 8CI. Cypri-no-Pourpre B

[3254-85-1]



C₇H₅N₅O₄ 223.148

Isol. from the scales and skin of various fish. Cryst. Mp > 360°. λ_{max} 224 (log ε 4.57); 259 (log ε 4); 282 (sh) (log ε 3.53); 347 (log ε 4.17) (0.1M NaOH).

Purrmann, R. *et al.*, *Annalen*, 1941, **548**, 284-292 (*synth*)

Matsuura, S. *et al.*, *J. Biochem. (Tokyo)*, 1955, **42**, 419-422 (*isol*)


Taylor, E.C. *et al.*, *J.A.C.S.*, 1959, **81**, 2474-2479 (*synth*)

Taylor, E.C. *et al.*, *Tet. Lett.*, 1973, 2093-2095 (*synth*)

2-Amino-6,8-dihydroxypurine A-753

2-Amino-7,9-dihydro-1H-purine-6,8-dione, 9CI. 8-Oxoguanine. 7,8-Dihydro-8-oxo-guanine. 2-Aminopurine-6,8-diol. 8-Hydroxy-guanine

[5614-64-2]



C₅H₅N₅O₂ 167.127

Other tautomers possible. 8-Oxo form predominates. Prod. *in vivo* by free radical hydroxylations of guanine radicals in DNA. Involved in base pair mismatches in mutagenesis. Crystalline powder. Mp 300°.

► Potent mutagen.

7-Me: 2-Amino-6,8-dihydroxy-7-methyl-purine. 2-Amino-7-methyl-6,8-purine-diol. 8-Hydroxy-7-methylguanine

[1688-85-3]
C₆H₇N₅O₂ 181.154
Platelets.

9-Me: 2-Amino-6,8-dihydroxy-9-methyl-purine. 2-Amino-9-methyl-6,8-purine-diol. 8-Hydroxy-9-methylguanine. 9-Methylguanin-8(7H)-one

[21823-84-7]
C₆H₇N₅O₂ 181.154
Cryst. + H₂O. Mp 400°.

1,7-Di-Me: 1,7-Dimethyl-8-oxo-9H-guanine

[65879-11-0]

C₇H₉N₅O₂ 195.18

Cryst. (H₂O). Gradual dec. >240°.

7,0⁶-Di-Me: 2-Amino-8-hydroxy-6-methoxy-7-methylpurine. 2-Amino-7,9-dihydro-6-methoxy-7-methyl-8H-purin-8-one. 6-Methoxy-7-methyl-8-oxoguanine

[247165-80-6]

C₇H₉N₅O₂ 195.18

Isol. from the ascidian *Symplegma rubra*. Solid. λ_{max} 210 (log ε 4.2); 246 (log ε 3.6); 284 (log ε 3.9) (MeOH).

9-Et: 2-Amino-9-ethyl-6,8-dihydroxypurine. 2-Amino-9-ethyl-6,8-(1H,9H)-purinedione

[21823-85-8]

C₇H₉N₅O₂ 195.18

Cryst. (KOH aq./AcOH). Mp 359-360°.

[112009-08-2]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 716D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 217C (nmr)

Fischer, E. *et al.*, *Ber.*, 1897, 30, 570 (synth)

Karrer, P. *et al.*, *Helv. Chim. Acta*, 1948, 31, 1214 (isol)

Cavaliere, L.F. *et al.*, *J.A.C.S.*, 1950, 72, 2587 (synth, uv)

Borowitz, I.J. *et al.*, *Biochemistry*, 1965, 4, 650 (7-Me)

Perini, F. *et al.*, *J.O.C.*, 1970, 35, 812 (synth, uv, derivs)

Brown, R. *et al.*, *J.C.S. Perkin 1*, 1977, 1003 (9-Me)

Parham, J.C. *et al.*, *J.O.C.*, 1978, 43, 2325 (di-Me)

Dizdaroglu, M. *et al.*, *Biochemistry*, 1985, 24, 4476 (formn)

Kohda, K. *et al.*, *Biochem. Biophys. Res. Commun.*, 1987, 149, 1141 (formn)

Kasai, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1987, 60, 3799 (cryst struct, 9-Et)

Floyd, R.A. *et al.*, *Carcinogenesis (London)*, 1990, 11, 1447 (rev)

Doi, M. *et al.*, *J.C.S. Perkin 1*, 1991, 55 (cryst struct, 9-Et)


Brown, T. *et al.*, *Chem. Ber.*, 1993, 484 (rev)

Lindsay, B.S. *et al.*, *J. Nat. Prod.*, 1999, 62, 1573-1575 (2-Amino-8-hydroxy-6-methoxy-7-methylpurine)

6-Amino-2,8-dihydroxypurine A-754

6-Amino-1H-purine-2,8(3H,7H)-dione, 9CI. 6-Aminopurine-2,8-diol. 2,8-Dihydroxyadenine

[30377-37-8]



C₅H₅N₅O₂ 167.127

Other tautomers possible. Prod. by *Alcaligenes aquamarinus*. Found in urine of humans with 2,8-dihydroxyadenine urolithiasis associated with adenine phosphoribosyltransferase deficiency. Xanthine oxidase inhibitor. Sol. bases; fairly sol. H₂O; poorly sol. MeOH, DMSO, hexane. Mp 335-340° dec. λ_{max}

305 (ε 19950) (HCl) (Berdy). λ_{max} 300 (ε 14800) (NaOH) (Berdy).

(1H,3H)-form

1,3-Di-Me: 6-Amino-1,3-dimethyl-1H-purine-2,8(3H,8H)-dione. 1,3-Dimethyl-8-oxoisoguanine

[683228-71-9]

C₇H₉N₅O₂ 195.18

Isol. from the ascidian *Pseudodistoma cereum* and from *Phestilla melanobantanchia* and a *Tubastrea* sp. Amorph. solid. λ_{max} 203 (log ε 3.89); 308 (log ε 3.57) (MeOH).

Bendich, A. *et al.*, *J. Biol. Chem.*, 1950, 183, 267-277 (isol)

Cavaliere, L.F. *et al.*, *J.A.C.S.*, 1950, 72, 2587-2594 (synth)

Klenow, H. *et al.*, *Biochem. J.*, 1952, 50, 404-407 (isol)

Stevens, M.A. *et al.*, *J.A.C.S.*, 1960, 82, 1148-1152 (synth)

Simmonds, H.A. *et al.*, *Biochem. J.*, 1976, 157, 485-487 (isol)

Sunahara, N. *et al.*, *Agric. Biol. Chem.*, 1977, 41, 1103-1109 (isol)

Simmonds, H.A. *et al.*, *Clin. Chim. Acta*, 1986, 160, 103-108 (biochem, rev)

Kamatani, N. *et al.*, *Pharma Med.*, 1988, 6, 39-42 (rev)

Kojima, T. *et al.*, *Biomed. Chromatogr.*, 1991, 5, 57-61 (hplc)

Simmonds, H.A. *et al.*, *Lancet*, 1992, ii, 1295-1296 (biochem)

Sumi, S. *et al.*, *J. Chromatogr. B*, 1995, 672, 233-239 (hplc)


Sevcik, J. *et al.*, *Clin. Chim. Acta*, 1996, 245, 85-92 (occur)

Engle, S.J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1996, 93, 5307-5312 (biochem)

Appleton, D.R. *et al.*, *Nat. Prod. Res.*, 2004, 18, 39-42 (1,3-di-Me)

4-Amino-3,6-dihydroxy-1H-pyrazolo[3,4-d]pyrimidine A-755

4-Amino-1H-pyrazolo[3,4-d]pyrimidine-3,6(2H,5H)-dione, 9CI. *Hydroxyakalone* [182056-35-5]




C₅H₅N₅O₂ 167.127

Prod. by the marine *Agrobacterium aurantiacum*. Xanthine oxidase inhibitor. Shows antigout activity. Powder. Sol. bases, DMSO; poorly sol. MeOH, hexane, CHCl₃. λ_{max} 298 (H₂O). λ_{max} 298 (pH12 buffer) (Berdy).

Izumida, H. *et al.*, *J. Antibiot.*, 1997, 50, 916-918 (isol, pmr, cmr)

25-Amino-3,23-dihydroxys-tigmast-5-ene-7,22-dione A-756



C₂₉H₄₇NO₄ 473.695

(3β,23R,24S)-form

N-Me, 3-O-β-D-glucopyranoside: *Triumfettoside A*

[551929-44-3]

C₃₆H₅₉NO₉ 649.863

Alkaloid from the aerial parts of *Triumfetta flavesces*. Amorph. powder. Mp > 300°.


Ahmed, B. *et al.*, *Pharmazie*, 2002, 57, 709-713 (isol, pmr, cmr, ms)

7-Amino-1,6-dimethyl-5,8-isoquinolinedione, 9CI

A-757

Cribrostatin 1

[144279-35-6]



C₁₁H₁₀N₂O₂ 202.212

Alkaloid from the blue marine sponge *Cribrorchalina* sp. Exhibits cytotoxicity against the P388 lymphocytic leukaemia cell line. Red-orange cryst. (CH₂Cl₂/MeOH). Mp 220-235° dec. λ_{max} 207 (ε 8730); 232 (ε 4700); 256 (ε 4170); 272 (ε 4160); 304 (ε 537) (MeOH) (Berdy).

Petit, G.R. *et al.*, *Can. J. Chem.*, 1992, 70, 1170 (isol, uv, ir, pmr, cmr, cryst struct)

2-Amino-14,16-dimethyl-3-octadecanol A-758

Paecilaminol. Antibiotic FKI 0550. FKI 0550

H₃CCH₂CH(CH₃)CH₂CH(CH₃)(C-H₂)₁₀CH(OH)CH(NH₂)CH₃

C₂₀H₄₃NO 313.566

Prod. by *Fusarium avenaceum* and *Paecilomyces* sp. FKI-0550. NADH-fumarate reductase inhibitor. Cytotoxic. Mycotoxin. Pale yellow oil. [α]_D²⁵ +26.7 (c, 0.24 in MeOH). λ_{max} 203 (ε 8250); 220 (sh) (ε 4400); 246 (ε 2040); 265 (sh) (ε 1570) (MeOH).

Uhlig, S. *et al.*, *Toxicol.*, 2005, 46, 513-522 (isol, pmr, cmr, ms)


Ui, H. *et al.*, *J. Antibiot.*, 2006, 59, 591-596 (isol, pmr, cmr, activity)

6-Amino-1,2-dithiolo[4,3-*b*]pyrrol-5(4H)-one, 9CI

A-759

Holothin

[488-03-9]



C₅H₄N₂OS₂ 172.231

Hydrol. prod. of *Holomycin*. Greenish-black cryst. (as hydrochloride). Mp 300° (hydrochloride).


▶ AB7450000

N⁶-Ac: *Holomycin*. N-Demethylthiolutin.


- Antibiotic MM 21801. MM 21801*
[488-04-0]
C₇H₆N₂O₂S₂ 214.269
Prod. by *Streptomyces* sp. P6621 and *Streptomyces clavuligerus*. Shows antibiotic props. Orange-yellow flakes (MeOH/EtOAc). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 268-270° dec. λ_{max} 246 (ε 6460); 302 (ε 3090); 388 (ε 11200) (MeOH or EtOH) (Derep). λ_{max} 246 (ε 4665); 301 (ε 2354); 388 (ε 7918) (MeOH) (Berdy). λ_{max} 245 (ε 6310); 302 (ε 3310); 390 (ε 11750) (EtOH) (Berdy).
- ▶ AB7450000
N⁶-Propanoyl: **N-Propionylholothin**. N-(4,5-Dihydro-5-oxo-1,2-dithiolo[4,3-b]pyrrol-6-yl)propanamide, 9CI [4708-23-0]
C₈H₈N₂O₂S₂ 228.295
Isol. from *Streptomyces* P6621. Orange-yellow prisms. Sol. MeOH. Mp 255-264° dec. λ_{max} 246 (ε 6460); 302 (ε 3090); 388 (ε 11200) (MeOH or EtOH) (Derep). λ_{max} 246 (ε 7800); 302 (ε 4250); 388 (ε 14400) (MeOH) (Berdy).
- N⁶-Hexanoyl: **Xenorhabdin 1**. N-(4,5-Dihydro-5-oxo-1,2-dithiolo[4,3-b]pyrrol-6-yl)hexanamide, 9CI [92680-94-9]
C₁₁H₁₄N₂O₂S₂ 270.376
Prod. by *Xenorhabdus* spp. Possesses antimicrobial and insecticidal props. Sol. MeOH, EtOAc, DMSO; fairly sol. hexane; poorly sol. H₂O. Mp 192-193°. Related to Thiolutin and Aureothricin. λ_{max} 243 (sh) (ε 6000); 300 (ε 3100); 387 (ε 11000) (MeOH) (Derep). λ_{max} 250 ; 310 ; 390 (MeOH) (Berdy).
- N⁶-Octanoyl: **Xenorhabdin 3**
[92680-91-6]
C₁₃H₁₈N₂O₂S₂ 298.429
Prod. by *Xenorhabdus* spp. Possesses antimicrobial and insecticidal props. Sol. MeOH, DMSO, EtOAc; fairly sol. hexane; poorly sol. H₂O. Mp 360°. λ_{max} 243 (sh) (ε 6000); 300 (ε 3100); 387 (ε 11000) (MeOH) (Derep). λ_{max} 250 ; 310 ; 390 (MeOH) (Berdy). λ_{max} 300 ; 387 (MeOH) (Berdy).
- N⁶-Tetradecanoyl: **N-Tetradecanoylholothin**
C₁₉H₃₀N₂O₂S₂ 382.59
Prod. by a marine-derived *Alteromonas* sp.
- N⁶-(5-Methylhexanoyl): **Xenorhabdin 2**
[92680-90-5]
C₁₂H₁₆N₂O₂S₂ 284.403
Prod. by *Xenorhabdus* spp. Possesses antimicrobial and insecticidal props. Sol. MeOH, EtOAc, DMSO; fairly sol. hexane; poorly sol. H₂O. Mp 210-213°. λ_{max} 243 (sh) (ε 6000); 300 (ε 3100); 387 (ε 11000) (MeOH) (Derep). λ_{max} 244 ; 300 ; 387 (MeOH) (Berdy).
- 4-Me: 6-Amino-4-methyl-1,2-dithiolo[4,3-b]pyrrol-5(4H)-one. **Pyrrhothine** [642-77-3]
C₆H₆N₂O₂S₂ 186.258
Isol. from a *Streptomyces* sp. Active against gram-positive and -negative bacteria. Amorph. yellow solid. Mp 191-194° dec.
- 4-Me: **hydrochloride**:
Yellow prisms + 1H₂O. Sinters at ca. 200°.
- 4-Me, N⁶-Ac: **Thiolut.** N-Acetylpyrrhothine. **Farcinicin**†. **Acetopyrrhothine** [87-11-6]
C₈H₈N₂O₂S₂ 228.295
Prod. by *Streptomyces albus*, *Streptomyces pimprina*, *Streptomyces thioleutus*, other *Streptomyces* spp. and *Saccharothrix* sp. SA 233. Antibiotic which inhibits microbiological growth in beer. Phytotoxic in high concentrations. Shows platelet aggregation props. Brilliant yellow needles (1-butanol). Sol. MeOH, CHCl₃; poorly sol. H₂O, C₆H₆, Et₂O, hexane. Mp 273-276° dec. Bp_{0.1} 200° subl. λ_{max} 248 (ε 6300); 311 (ε 5700); 388 (ε 10800) (MeOH) (Derep). λ_{max} 250 (ε 6030); 311 (ε 5600); 388 (ε 10960) (MeOH) (Berdy).
- ▶ LD₅₀ (mus, scu) 25 mg/kg. JPI355000
- 4-Me, N⁶-propanoyl: **Aureothricin**. **Propionylpyrrhothine**. **Farcinicin**† [574-95-8]
C₉H₁₀N₂O₂S₂ 242.322
Isol. from *Streptomyces celluloflavus* 39a, *Streptomyces cyanoflavus*, *Streptomyces kasugaensis* and *Streptomyces farcinicus*. Antibiotic and antifungal agent. Platelet aggregation inhibitor. Golden-yellow cryst. (EtOAc). Sol. MeOH, CHCl₃; fairly sol. C₆H₆; poorly sol. Et₂O, H₂O, hexane. Mp 260-270° dec. λ_{max} 246 (ε 6500); 313 (ε 4000); 390 (ε 11100) (EtOH) (Berdy).
- ▶ LD₅₀ (mus, scu) 10 mg/kg.
- 4-Me, N⁶-butanoyl: **N-(4,5-Dihydro-4-methyl-5-oxo-1,2-dithiolo[4,3-b]pyrrol-6-yl)butanamide**. **N-Butanoylpyrrhothine**. **Antibiotic PSC₂**. **PSC₂** [112843-01-3]
C₁₀H₁₂N₂O₂S₂ 256.349
Prod. by *Xenorhabdus bovienii* A2 and *Saccharothrix* sp. SA 233. Yellow powder. λ_{max} 308 (log ε 3.7); 389 (log ε 3.92) (MeOH).
- 4-Me, N⁶-(2-methylpropanoyl): **N-(2-Methylpropanoyl)pyrrhothine**. **Iso-butyropyrrhothine**†. **N-Isobutyrylpyrrhothine**. **Antibiotic PSC₁**. **PSC₁** [39859-18-2]
C₁₀H₁₂N₂O₂S₂ 256.349
Prod. by *Streptomyces pimprina* and *Saccharothrix* sp. SA 233. Shows platelet aggregation props. Orange-red plates (C₆H₆). Sol. CHCl₃, EtOAc, Me₂CO; fairly sol. C₆H₆; poorly sol. H₂O, hexane. Mp 228-229°. λ_{max} 245 ; 290 ; 310 ; 388 (MeOH) (Berdy). λ_{max} 246 (ε 5400); 302 (ε 3000); 386 (ε 11000) (MeOH) (Derep).
- 4-Me, N⁶-hexanoyl: **Xenorhabdin 4**. **N-Pentanoylpyrrhothine**
[92680-92-7]
C₁₂H₁₆N₂O₂S₂ 284.403
From *Xenorhabdus* spp. Possesses antimicrobial and insecticidal props. Sol. MeOH, DMSO, EtOAc; fairly sol. hexane; poorly sol. H₂O. Mp 165°. λ_{max} 246 (sh) (ε 6300); 308 (ε 5700); 388 (ε 11000) (MeOH) (Derep). λ_{max} 245 ; 310 (ε 1000); 312 ; 390 (ε 10000) (MeOH) (Berdy).
- 4-Me, N⁶-(3-methylbutanoyl): **N-(4,5-Dihydro-4-methyl-5-oxo-1,2-dithiolo[4,3-b]pyrrol-6-yl)-3-methylbutanamide**. **N-(3-Methylbutanoyl)pyrrhothine** [167559-98-0]
C₁₁H₁₄N₂O₂S₂ 270.376
Isol. from *Xenorhabdus bovienii* A2. Solid.
- 4-Me, N⁶-(5-methylhexanoyl): **N-(5-Methylhexanoyl)pyrrhothine**. **Xenorhabdin 5**
[92680-93-8]
C₁₃H₁₈N₂O₂S₂ 298.429
From *Xenorhabdus* spp. Possesses antimicrobial and insecticidal props. Sol. MeOH, EtOAc, DMSO; fairly sol. hexane; poorly sol. H₂O. λ_{max} 246 (sh) (ε 6300); 308 (ε 5700); 388 (ε 11000) (MeOH) (Derep). λ_{max} 246 ; 250 ; 308 ; 310 (ε 1000); 388 ; 390 (ε 10000) (MeOH) (Berdy).
- 4-Me, N⁶-(3-methyl-2-butanoyl): **N-Sec-necioylpyrrhothine**. **PSA**. **Antibiotic PSA**
C₁₁H₁₂N₂O₂S₂ 268.36
Prod. by *Saccharothrix* sp. SA 233. Yellow-orange powder. λ_{max} 302 (log ε 3.87); 402 (log ε 3.97) (MeOH).
- 4-Me, N⁶-tigloyl: **N-Tigloylpyrrhothine**. **PSB**. **Antibiotic PSB**
C₁₁H₁₂N₂O₂S₂ 268.36
Prod. by *Saccharothrix* sp. SA 233. Yellow-orange powder. λ_{max} 302 (log ε 3.85); 402 (log ε 3.96) (MeOH).
- Celmer, W.D. et al., *J.A.C.S.*, 1952, **74**, 6304; 1955, **77**, 2861 (*Pyrrhothine*, *Aureothricin*, *Thiolutin*, *isol*, *ir*, *uv*, *struct*)
Ettlinger, L. et al., *Helv. Chim. Acta*, 1959, **42**, 563 (*isol*)
Bhate, D.S. et al., *Experientia*, 1960, **16**, 504-505 (*Isobutyropyrrhothine*)
Schmidt, U. et al., *Chem. Ber.*, 1964, **97**, 1511 (*synth*, *uv*, *ir*)
Büchi, G. et al., *J.A.C.S.*, 1964, **86**, 5654 (*synth*)
Yamagishi, S. et al., *Yakugaku Zasshi*, 1971, **91**, 351 (*Aureothricin*, *isol*, *struct*, *ir*, *pmr*, *ms*)
Hagio, K. et al., *Bull. Chem. Soc. Jpn.*, 1974, **47**, 1484 (*Holothin*, *Thiolutin*, *Aureothricin*, *synth*, *ir*, *uv*, *pmr*)
Okamura, K. et al., *J. Antibiot.*, 1977, **30**, 334-336 (*isol*, *ir*, *ms*)
Ellis, J.E. et al., *J.O.C.*, 1977, **42**, 2891-2893 (*synth*, *ir*, *uv*, *ms*, *nmr*)
Kenig, M. et al., *J. Antibiot.*, 1979, **32**, 549-554 (*MM 21801*)
Ninomiya, Y.T. et al., *Chem. Pharm. Bull.*, 1980, **28**, 3157-3162 (*Pyrrhothine*, *Thiolutin*, *props*, *ms*)
Deb, P.R. et al., *Curr. Sci. (India)*, 1984, **53**, 659 (*Thiolutin*, *props*)
Pat. Coop. Treaty (WIPO), 1984, 01 775; *CA*, **101**, 189727 (*Xenorhabdins*)
McInerney, B.V. et al., *J. Nat. Prod.*, 1991, **54**, 774 (*Xenorhabdins*)
Dell, I. et al., *ACS Symp. Ser.*, 1992, **504**, 384 (*synth*)
Sato, A. et al., *Annu. Rep. Sankyo Res. Lab.*, 1995, **47**, 1-58 (*N-Tetradecanoylholothin*)
Li, J. et al., *J. Nat. Prod.*, 1995, **58**, 1081-1086 (*N-Butanoylpyrrhothine*, *N-3-Methylbutanoylpyrrhothine*)
Lamari, L. et al., *J. Antibiot.*, 2002, **55**, 696-701; 702-706 (*Saccharothrix pyrrhothines*)
Hjelmgaard, T. et al., *Org. Biomol. Chem.*, 2007, **5**, 344-348 (*synth*)

2-(12-Amino-3,7,9-dodecatrienyl)pyridine A-760

12-(2-Pyridinyl)-3,5,9-dodecatrien-1-amine

C₁₇H₂₄N₂ 256.39**(all-E)-form****Naloamine**


[179118-58-2]

Alkaloid from the mollusc *Smaragdinella calyculata*. Amorph. solid. λ_{\max} 206 (ϵ 12000); 230 (ϵ 15000); 262 (ϵ 4300); 268 (ϵ 4200); 280 (ϵ 2700) (MeOH).Szabo, C.M. *et al.*, *Tetrahedron*, 1996, **52**, 9681 (isol, uv, pmr, cmr, ms, struct)**2-Amino-11-dodecen-3-ol** A-761C₁₂H₂₅NO 199.336**(2S,3R)-form****Halaminol B**


[389125-59-1]

Isol. from a marine sponge *Haliclona* n. sp. Oil. $[\alpha]_D^{25}$ +2.1 (c, 0.06 in CH₂Cl₂).Clark, R.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1568-1571 (*Halaminol B*, isol, pmr, cmr)**4-Amino-2-dodecyltetrahydro-3-furanol** A-762

4-Amino-2-dodecyl-3-hydroxytetrahydrofuran

C₁₆H₃₃NO₂ 271.442**(2R*,3S*,4S*)-form**N-(2R-Hydroxyheptadecanoyl): **Suillamide**

[1067910-08-0]

C₃₃H₆₅NO₄ 539.881Isol. from *Suillus luteus*. Amorph. solid. $[\alpha]_D^{25}$ +16.3 (c, 0.1 in CHCl₃).León, F. *et al.*, *Chem. Biodiversity*, 2008, **5**, 120-125 (isol, pmr, cmr, ms)**3-Aminoestra-1,3,5(10)-tri-en-17-ol** A-763C₁₈H₂₅NO 271.402**17β-form****Hedonegamini**

[10427-24-4]

Constit. of the female of fossil plant *Palaeogelos doleros*. Cryst. (EtOAc). Mp 144-146° (63-65°).Dannenberg, H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1967, **348**, 775-782 (synth)Roedder, W. *et al.*, *Pharm. Ztg.*, 1976, **121**, 479-484 (isol)Smith, W.B. *et al.*, *Org. Prep. Proced. Int.*, 1990, **22**, 501-506 (synth, cmr)**2-Aminoethanesulfinic acid, 9CI** A-764**Hypotaaurine**. *Cystaminesulfinic acid*

[300-84-5]

H₂NCH₂CH₂SO₂HC₂H₇NO₂S 109.149Constit. of rat brain. Found in the marine vestimentiferan worm *Riftia pachyptila* and other marine invertebrates. Mp 186-188°. pK_{a1} 2.16; pK_{a2} 9.56.Chatagner, F. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1951, **232**, 448 (isol)*Biochem. Prep.*, 1963, **10**, 72-75 (synth)Hope, D.B. *et al.*, *J.C.S. (C)*, 1970, 270 (synth)Fellman, J.H. *et al.*, *J. Labelled Compd. Radiopharm.*, 1981, **18**, 765 (synth)Stipanuk, M.H. *et al.*, *Methods Enzymol.*, 1987, **143**, 155 (hplc)Fugler-Domenico, L. *et al.*, *Biol. Amines*, 1989, **6**, 289 (conform)Chaimbault, P. *et al.*, *Anal. Biochem.*, 2004, **332**, 215-225 (marine, occur)**2-Aminoethanesulfonic acid, 9CI** A-765**Taurine**, **8CI**, **INN**. *Aminoethylsulfonic acid*. *Ethylaminesulfonic acid*. **FEMA** 3813

[107-35-7]

H₂NCH₂CH₂SO₃HC₂H₇NO₃S 125.148Occurs free in animal tissues, bacteria, sponges, red algae, e.g. isol. from *Macrocallista nimbosa*, *Turbo stenogyrus*, *Calyx nicaeensis*, *Godia gigas*, *Mytilus edulis*. Also from green algae, e.g. *Caulerpa okamurai*, *Caulerpa racemosa*, *Chlorodesmis comosa*, *Codium adherens*, *Codium fragile* and *Enteromorpha linza*. Isol. from marine vestimentarian worm *Riftia pachyptila* and from higher plants, e.g. leguminous seedlings. A non-essential amino acid. Intermed. in metab. of cysteine. Used as an adjunct in treatment of hypercholesterolaemia. Metabolic regulator. Monoclinic prismatic rods with sharp taste. Sol. H₂O, insol. EtOH. Mp 328° (320-325° dec.). pK_{a1} 1.5; pK_{a2} 8.74. Dec. at 300°.▶ LD₅₀ (mus, scu) 6000 mg/kg. WX0175000*Amide*: [4378-70-5]C₂H₈N₂O₂S 124.163

Plates (EtOH) (as hydrochloride). Mp 133° (hydrochloride).

N-Me: [107-68-6]C₃H₉NO₃S 139.175Widely distributed in marine algae including the red alga *Ptilota pectinata* and green alga *Chlorodesmis comosa*. Also found in sponges *Calyx nerets* and *Halichondria* sp. Prisms. V. sol.H₂O; insol. EtOH, Et₂O. Mp 241-242°.*N,N-Di-Me*: [637-95-6]C₄H₁₁NO₃S 153.202Obt. from *Corallina officinalis*. Present in *Furcellaria fastigiata* and other red algae. Prisms (MeOH). V. sol. H₂O, AcOH, insol. EtOH, Et₂O. Mp 315-316° dec.*N,N-Bis(2-chloroethyl)*: **Taumustine** [98277-87-3]C₆H₁₃Cl₂NO₃S 250.145Cryst. (H₂O). Mp 178-180°.*N-(2,3-Dihydroxypropyl)*: *N-Glyceryl-taurine*

[65222-42-6]

C₅H₁₃NO₅S 199.227Isol. from the red alga *Gigartina leptorhynchus*. Cryst. (EtOH aq.). Mp 163-164°. $[\alpha]_D^{25}$ -21 (H₂O).*N-(15-Methyl-9Z-hexadecenyl)*: **N-(15-Methyl-9-hexadecenyl)taurine**

[679834-30-1]

C₁₉H₃₇NO₄S 375.572Isol. from the sponge *Erylus nobilis*.*N-(5Z,8Z,11Z,14Z-Eicosatetraenyl)*:*N-(5,8,11,14-Eicosatetraenyl)taurine*

[679834-28-7]

C₂₂H₃₇NO₄S 411.605Isol. from the starfish *Certonaroda semiregularis*.

[7347-25-3, 4316-74-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 890A (ir)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 1433C (nmr)*Org. Synth., Coll. Vol.*, 2, 1943, 563; 564 (synth)Ishidate, M. *et al.*, *Chem. Pharm. Bull.*, 1954, **2**, 275-279 (synth)Lindberg, B. *et al.*, *Acta Chem. Scand.*, 1955, **9**, 1093-1096; 1323-1326 (isol, *N-Me*, *N,N-di-Me*)Wickberg, B. *et al.*, *Acta Chem. Scand.*, 1956, **10**, 1097-1099 (*Glyceryltaurine*)Sutherland, H.H. *et al.*, *Acta Cryst.*, 1963, **16**, 897-901 (cryst struct)Huxtable, R. *et al.*, *Taurine*, [Int. Symp.], 1st, 1975 (1976), (Eds.), Raven Press, New York, 1975, (book)Utkina, N.K. *et al.*, *Khim. Prir. Soedin.*, 1984, 124-125; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 126-127 (*N-Me*, occur)Hashem, K.M.E. *et al.*, *Bull. Soc. Chim. Belg.*, 1985, **94**, 735-754 (cmr)*Taurine: Nutritional Value and Mechanisms of Action*, (eds. Lombardini, J.B. *et al.*), Plenum Press, 1992, (book)*Martindale, The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1419*Fenaroli's Handbook of Flavor Ingredients*, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 1665-1666 (use, occur)*Merck Index*, 13th edn., 2001, No. 9163 (bibl) Hibbs, D.E. *et al.*, *Chem. Eur. J.*, 2003, **9**, 1075-1084 (cryst struct)Wang, W. *et al.*, *Nat. Prod. Sci.*, 2003, **9**, 241-244; *CA*, **140**, 335912 (*N-Methylhexadecenyl*, *N-Eicosatetraenyl*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TAG750**2-Aminoethanesulfonylthioic acid, 9CI** A-766**S-(2-Aminoethyl) thiosulfate**, **8CI**. **Thio-taurine**

[2937-54-4]

[31999-89-0 (Na salt)]

H₂NCH₂CH₂SO₂SH

C₂H₇NO₂S₂ 141.215

Isol. from various marine invertebrates incl. *Bathymodiolus* spp., *Calyptogena magnifica* and *Riftia pachyptila*. Biomarker of sulfur-based symbiosis in deep-sea bivalves. Cryst. (EtOH aq.). Mp 213° dec.

Sörbö, B.H. et al., *Acta Chem. Scand.*, 1958, **12**, 146 (synth)

Biochem. Prep., 1963, **10**, 72-75 (synth)

Pruski, A.M. et al., *J. Exp. Biol.*, 2003, **206**, 2923-2930 (biosynth)

Chaimbault, P. et al., *Anal. Biochem.*, 2004, **332**, 215-225 (occur)

2-Aminoethanol, 9CI A-767

2-Hydroxyethylamine. Ethanolamine.

Monoethanolamine. Colamine. Olamine

[141-43-5]

H₂NCH₂CH₂OH

C₂H₇NO 61.083

Manuf. from ethylene oxide and NH₃. Widely distributed in biol. tissues. Component of lecithin. Constit. of defence secretions of *Chrysolina* and related beetles. Chemical intermed., org. base. Removes CO₂ and H₂S from gases. Reagent for fluorimetric anal. of carbohydrates by hplc. Selectively cleaves the glycosyl ester bond of peracylated aldoses. Curing agent for epoxy resins. Viscous, hygroscopic liq. Misc. H₂O, MeOH, Me₂CO; spar. sol. C₆H₆, Et₂O, d₄²⁵ 1.01. Mp 10.5°. Bp 171°. n_D²⁰ 1.4539. pK_a 9.47. Vp 0.48 mmHg (20°).

- Fl. p. 85°, autoignition temp. 410°. Corrosive and irritating to skin, eyes and mucous membranes. LD₅₀ (rat, orl) 1720 mg/kg. Exp. teratogen. OES: long-term 3 ppm; short-term 6 ppm. KJ5775000

O-Ac: O-Acetyethanolamine. 2-Acetoxyethylamine

[1854-30-4]

C₄H₉NO₂ 103.121

Isol. from seeds of *Lens culinaris* (lentil), also present in other Fabaceae. Shows antiinflammatory props., stimulates cockroach heart. Bp_{0.3} 134-135°.

N-(9Z,12Z-Octadecadienyl): N-(2-Hydroxyethyl)-9,12-octadecadienamide, 9CI. N-(2-Hydroxyethyl)linoleamide

[68171-52-8]

[10015-67-5]

C₂₀H₃₇NO₂ 323.518

Alkaloid from *Vaccaria segetalis*.

[96-80-0, 20739-39-3]

Hayman, A.R. et al., *Phytochemistry*, 1987, **26**, 839 (O-Acetyethanolamine)

Sang, S. et al., *CA*, 1999, **130**, 349718n (N-Hydroxyethyl)linoleamide, isol)

2-Aminoethyl dihydrogen phosphate A-768

Ethanolamine O-phosphate. O-Phosphoethanolamine

[1071-23-4]

H₂NCH₂CH₂OP(O)(OH)₂

C₂H₈NO₄P 141.063

Isol. from eye lens tissue and gram-negative bacteria. Intermed. in phosphoglyceride biosynth. in mammals. Cryst. (EtOH aq.). Mp 242° (231.5-234°).

Mono-Et ester: Ethyl hydrogen 2-aminoethyl phosphate

C₄H₁₂NO₄P 169.117

Cryst. (EtOH/Me₂CO). Mp 238°.

Diisopropyl ester: Diisopropyl 2-aminoethyl phosphate

[14646-04-9]

C₈H₂₀NO₄P 225.224

Liq. Bp_{0.6} 75°.

Mono-Ph ester: Phenyl hydrogen 2-aminoethyl phosphate

C₈H₁₂NO₄P 217.161

Solid. V. sol. H₂O; spar. sol. EtOH; insol. C₆H₆, CHCl₃, Et₂O.

N-Ac: 2-Acetamidoethyl phosphate

[89603-45-2]

C₄H₁₀NO₅P 183.101

Occurs in O-antigen polysaccharide of *Proteus mirabilis* 038. No phys. props. reported.

[87111-65-7, 3724-89-8, 18672-70-3, 19077-76-0, 131703-98-5, 61020-38-0, 31512-55-7, 32607-10-6, 37785-54-9, 63441-18-9]

Plapinger, R.E. et al., *J.A.C.S.*, 1953, **75**, 5757

Cherbuliez, E. et al., *Helv. Chim. Acta*, 1956, **39**, 1455-1461 (N-Ac, synth)

Tsizin, Y.S. et al., *Zh. Obshch. Khim.*, 1963, **33**, 2873; *J. Gen. Chem. USSR (Engl. Transl.)*, 1963, **33**, 2800 (synth, ir)

Greenhalgh, R. et al., *Can. J. Chem.*, 1967, **45**, 495 (diisopropyl ester, synth, ir, pnr, Pnmr)

Greiner, J.V. et al., *Exp. Eye Res.*, 1982, **34**, 545 (P-31 nmr)

Glonek, T. et al., *J. Neurochem.*, 1982, **39**, 1210 (P-31 nmr)

Trigalo, F. et al., *J.C.S. Perkin 1*, 1982, 1733 (monoethyl ester)

Weber, H.P. et al., *Acta Cryst. B*, 1984, **40**, 506 (cryst struct)

Cates, L.A. et al., *J. Med. Chem.*, 1984, **27**, 654 (synth)

Hayase, Y. et al., *Chem. Pharm. Bull.*, 1986, **34**, 2251 (isol, ir, pnr, cmr)

Bissinger, P. et al., *Chem. Ber.*, 1991, **124**, 509 (synth, salts, cryst struct)

Ger. Pat., 1993, 4 141 842; *CA*, **120**, 57142 (N-Ac, synth)


Kandakova, A.N. et al., *Carbohydr. Res.*, 2003, **338**, 2387-2392 (N-Ac, occur)

4-(2-Aminoethyl)-2-bromo-phenol, 9CI A-769

2-(3-Bromo-4-hydroxyphenyl)ethylamine.

3-Bromotyramine

[104616-74-2]



C₈H₁₀BrNO 216.077

Isol. from the ascidian *Cnemidocarpa bicornuta*. Cryst. Mp 162-164°. λ_{max} 203 (log ε 4.4); 283 (log ε 3.3) (MeOH).

Hydrochloride:

Cryst. (MeOH/EtOAc). Mp 205.5-207°.

N,N,N-Tri-Me: 3-Bromo-4-hydroxy-N,N,N-trimethylbenzeneethanaminium, 9CI. 3-Bromo-N,N,N-trimethyltyramine

[259533-84-1]

C₁₁H₁₇BrNO⁺ 259.165

Isol. from the sponge *Verongula gigantea*. Isol. as trifluoroacetate salt. λ_{max} 281 (ε 1485) (MeOH).

Tolkachev, O.N. et al., *Zh. Obshch. Khim.*, 1958, **28**, 3320-3323; *J. Gen. Chem.*


USSR (Engl. Transl.), 1958, **28**, 3345-3348 (synth)

Lindsay, B.S. et al., *J. Nat. Prod.*, 1998, **61**, 857-858 (isol, uv, ir, pnr, cmr, ms)

Ciminiello, P. et al., *J. Nat. Prod.*, 2000, **63**, 263-266 (N-tri-Me deriv)

6-(2-Aminoethyl)-2,2-dimethyl-2H-1-benzopyran A-770

(2,2-Dimethyl-2H-1-benzopyran-6-yl)ethanamine



C₁₃H₁₇NO 203.283

N-(2-Methylpropanoyl): [124704-80-9]

C₁₇H₂₃NO₂ 273.374

Alkaloid from roots of *Amyris sylvatica* (Rutaceae).

N-(3-Methylbutanoyl): [124704-79-6]

C₁₈H₂₅NO₂ 287.401

Alkaloid from roots of *Amyris sylvatica* (Rutaceae). Mp 104-105°.

N-Benzoyl: N-[2-(2,2-Dimethyl-2H-1-benzopyran-6-yl)ethyl]benzamide, 9CI.

6-(2-Benzamidoethyl)-2,2-dimethyl-2H-1-benzopyran

[89764-18-1]

C₂₀H₂₁NO₂ 307.391

Alkaloid from the aerial portions of *Amyris plumieri* (Rutaceae). Mp 133.5-134.5°.

N-(3-Pyridinecarbonyl): N-[2-(2,2-Dimethyl-2H-1-benzopyran-6-yl)ethyl]-3-pyridinecarboxamide, 9CI

[69142-83-2]

C₁₉H₂₀N₂O₂ 308.379

Alkaloid from *Amyris plumieri* (Rutaceae). Mp 99-100°.

N-(3-Pyridinecarbonyl), N-Me: N-[2-(2,2-Dimethyl-2H-1-benzopyran-6-yl)ethyl]-N-methyl-3-pyridinecarboxamide

[137132-29-7]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Amyris texana* (Rutaceae). Oil.

Burke, B.A. et al., *Tet. Lett.*, 1978, **19**, 2723-2726 (N-3-pyridinecarbonyl)


Philip, S. et al., *Heterocycles*, 1984, **22**, 9-12 (N-benzoyl)

Hasbun, C. et al., *J. Nat. Prod.*, 1989, **52**, 868-870 (*Amyris sylvatica* constit)

De la Fuente, G. et al., *Phytochemistry*, 1991, **30**, 2677-2684 (*Amyris texana* constit)


3-(2-Aminoethyl)-2-nitroindole A-771

2-Nitro-1H-indole-3-ethanamine. 2-Nitro-tryptamine

C₁₀H₁₁N₃O₂ 205.216N^b-Ac: N^b-Acetyl-2-nitrotryptamineC₁₂H₁₃N₃O₃ 247.253Isol. from the marine-derived *Flavobacterium* sp. T436.Schuhmann, I. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*Ac, isol*)**(2-Aminoethyl)phosphonic acid, 9CI** A-772

(2-Aminoethyl)phosphonous acid. 2-(Hydroxyphosphinyl)ethylamine. Antibiotic MP 105. MP 105

[85618-16-2]

C₂H₈NO₂P 109.064Prod. by *Streptomyces hygrosopicus* SF-1293. Amorph. powder. Mp 208-210° dec. (natural) Mp 255° (synthetic).Seto, H. *et al.*, *Biochem. Biophys. Res. Commun.*, 1983, **111**, 1008-1014 (*isol*)Dingwall, J.G. *et al.*, *Tetrahedron*, 1989, **45**, 3787-3808 (*synth, pmr, P-31 nmr*)Berlicki, L. *et al.*, *J.O.C.*, 2005, **70**, 6340-6349 (*synth*)**(2-Aminoethyl)phosphonic acid, 9CI** A-773*Ciliatine. AEP*

[2041-14-7]

C₂H₈NO₂P 125.064Constit. as glyceryl esters of rumen protozoan, *Tetrahymina pyriformis*, sea anemones, bovine brain, mycobacteria, and a fungus *Pythium prolatum*. Rhombic cryst. or needles (EtOH aq.). Mp 295-297° dec. pK_{a1} 2.45; pK_{a2} 7; pK_{a3} 10.8. pK_{a1} 1.3; pK_{a2} 6.51; pK_{a3} 11.5 (25°). The metastable rhombic form is difficult to prepare. First known natural compd. contg. the C-P bond.*Di-Et ester*: [41468-36-4]C₆H₁₆NO₃P 181.171Liq. Bp_{0.025} 54-56°.*N-Benzoyl*: 2-(Benzoylamino)ethylphosphonic acidC₉H₁₂NO₄P 229.172


Solid. Mp 191-192°.

N-Me: N-Methylciliatine. 2-(Methylamino)ethylphosphonic acid

[14596-55-5]

C₃H₁₀NO₃P 139.091Isol. from the sea anemone *Anthopleura xanthogrammica*. Needles (MeOH aq.). Mp 291° dec.*N,N-Di-Me*: 2-Dimethylaminoethylphosphonic acid. N,N-Dimethylciliatine

[14596-56-6]

C₄H₁₂NO₃P 153.117Isol. from *Anthopleura xanthogrammica*. Pale yellow cryst. (MeOH aq.). Mp 249.5° dec.*N,N-Di-Me, Et ester*: [41948-36-1]C₆H₁₆NO₃P 181.171Bp₁ 100°.*N-Phthalimide, di-Et ester*: Diethyl (2-phthalimidoethyl)phosphonateC₁₄H₁₈NO₅P 311.274Cryst. (Et₂O/petrol). Mp 58°.*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 1477A (*nmr*)Kosolapoff, G.M. *et al.*, *J.A.C.S.*, 1947, **69**, 2112 (*synth*)*Top. Phosphorus Chem.*, Interscience, 1966, **4**, 23; **11**; 297 (*rev*)Barycki, J. *et al.*, *Tet. Lett.*, 1970, 3147 (*synth*)Isbell, A.F. *et al.*, *J.O.C.*, 1972, **37**, 4399 (*synth, bibl*)Harvey, D.J. *et al.*, *J. Chromatogr.*, 1973, **79**, 65 (*ms*)Lagrange, C.G. *et al.*, *Can. J. Chem.*, 1978, **56**, 663 (*ir, Raman*)Brigot, D. *et al.*, *Tetrahedron*, 1979, **35**, 1345 (*synth, pmr, derivs*)Fabre, G. *et al.*, *Can. J. Chem.*, 1981, **59**, 2864 (*synth, di-Et ester, ir, pmr*)Varlet, J.M. *et al.*, *Tetrahedron*, 1981, **37**, 3713 (*derivs*)Al-Rawi, J.M.A. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 75 (*di-Et ester, cmr*)Appleton, T.G. *et al.*, *Aust. J. Chem.*, 1984, **37**, 1833 (*props, cmr, P-31 nmr*)Yamauchi, K. *et al.*, *J.O.C.*, 1984, **49**, 1158 (*N-phthalimide deriv*)Gubnitskaya, E.S. *et al.*, *Zh. Obshch. Khim.*, 1986, **56**, 2017; *J. Gen. Chem. USSR*(*Engl. Transl.*), 1986, **56**, 1779 (*esters, synth, ir*)Osapay, G. *et al.*, *Tetrahedron*, 1987, **43**, 2977 (*ir, pmr*)Lazukina, L.A. *et al.*, *Zh. Obshch. Khim.*, 1988, **58**, 939; *J. Gen. Chem. USSR*(*Engl. Transl.*), 1988, **58**, 833 (*synth, P-31 nmr*)Thayer, J.S. *et al.*, *Appl. Organomet. Chem.*, 1989, **3**, 203 (*rev*)Warielowski, C. *et al.*, *J. Prakt. Chem.*, 1989, **331**, 507 (*synth, ir*)Harris, R.K. *et al.*, *Magn. Reson. Chem.*, 1989, **27**, 470 (*cmr, N-15 nmr, P-31 nmr*)Glowacki, Z. *et al.*, *Magn. Reson. Chem.*, 1989, **27**, 897 (*cmr*)Warielowski, C. *et al.*, *Synthesis*, 1989, **52** (*synth, ir, pmr*)Ohashi, K. *et al.*, *Tetrahedron*, 1989, **45**, 2557 (*N-Me, synth, ir*)Hayashi, A. *et al.*, *Chem. Phys. Lipids*, 1990, **52**, 57 (*fab-ms*)Neidlein, R. *et al.*, *Helv. Chim. Acta*, 1992, **75**, 2545 (*N-phthalimide deriv*)Ohno, K. *et al.*, *J. Mol. Struct.*, 1993, **298**, 1 (*ir, Raman, conformm*)Fields, S.C. *et al.*, *Tetrahedron*, 1999, **55**, 12237-12273 (*rev, synth*)**5-(1-Aminoethyl)-3-pyridine-carboxylic acid, 9CI** A-774

(+/-)-form

C₈H₁₀N₂O₂ 166.179

(±)-form


Me ester:

Syrup.

p-Nitrobenzoyl:

Needles. Mp 154-157°.

(±)-form

Me ester: Methyl 5-(1-aminoethyl)-3-pyridinecarboxylate, 9CI. 3-(1-Aminoethyl)-5-carbomethoxy-pyridine [38940-68-0]C₉H₁₂N₂O₂ 180.206Alkaloid from *Nauclea diderrichii* (Rubiaceae). Oil. [α]_D²⁵ +27 (c, 0.34 in MeOH).McLean, S. *et al.*, *Can. J. Chem.*, 1971, **50**, 1478 (*isol, ms, pmr, uv, synth*)**4-Amino-11-eudesmene** A-775C₁₅H₂₇N 221.385**(4α,7α)-form** [442851-12-9]


[442851-11-8]

Constit. of *Axinyssa ambrosia*. Oil; cryst. (as hydrochloride). Mp 210-220° (hydrochloride). [α]_D -17.5 (c, 0.9 in CHCl₃).*N-Formyl*: 4-Formamido-11-eudesmene

[442851-14-1]

C₁₆H₂₇NO 249.395Constit. of *Axinyssa ambrosia*. Cryst. Mp 104-107°. [α]_D -10 (c, 0.22 in CHCl₃).*4-Isocyanide*: 4-Isocyanato-11-eudesmene

[442851-13-0]

C₁₆H₂₅N 231.38Constit. of *Axinyssa ambrosia*. Oil. [α]_D -15 (c, 0.2 in CHCl₃). Has -NC in place of NH₂.Petrichtcheva, N.V. *et al.*, *J. Nat. Prod.*, 2002, **65**, 851-855 (*isol, pmr, cmr*)**6-Amino-4(15)-eudesmene** A-776C₁₅H₂₇N 221.385**6α-form***Halichonadin D*

[847605-79-2]


Constit. of a *Halichondria* sp. Yellow oil. [α]_D²⁷ +18 (c, 0.2 in CHCl₃).*N-Methoxycarbonyl*: Halichonadin B

[847605-77-0]

C₁₇H₂₉NO₂ 279.422Constit. of a *Halichondria* sp. Amorph. solid. [α]_D²⁴ -10 (c, 1 in CHCl₃).

Ishiyama, H. *et al.*, *Tetrahedron*, 2005, **61**, 1101-1105 (isol, pmr, cmr)

4-Amino-11-gorgonene A-777



C₁₅H₂₇N 221.385

(4α,6α)-form

N-Formyl: 4-Formamido-11-gorgonene [134781-19-4]

C₁₆H₂₇NO 249.395

Isol. from the molluscs *Phyllidia pustulosa* and *Phyllidia varicosa*. Cryst. [α]_D²⁴ -61.8 (c, 0.11 in CHCl₃). Mp >300° dec.

4-Isocyanide: 4-Isocyano-11-gorgonene

[134781-18-3]

C₁₆H₂₅N 231.38

Isol. from the molluscs *Phyllidia pustulosa* and *Phyllidia varicosa*. Oil. [α]_D²⁴ -66.9 (c, 1.6 in CHCl₃).

4-Isothiocyante: 4-Isothiocyanto-11-gorgonene


[134781-20-7]

C₁₆H₂₅NS 263.446

Isol. from *Phyllidia pustulosa*. Oil. [α]_D²⁴ -101.6 (c, 0.5 in CHCl₃).

Kassühlke, K.E. *et al.*, *J.O.C.*, 1991, **56**, 3747-3750 (isol, pmr, cmr, ms)

14-Amino-1(5),7,10(14)-guaiatricene-6,9-dione A-778



C₁₅H₁₉NO₂ 245.321

(4ξ,10Z)-form

Muscicapine A

[873684-18-5]

Alkaloid from the roots of *Croton muscicapa*. Amorph. yellow solid. [α]_D²⁰ -44 (c, 0.05 in CHCl₃).

N-(2-Methylbutyl): **Muscicapine C**

[873684-20-9]

C₂₀H₂₉NO₂ 315.455

Alkaloid from the roots of *Croton muscicapa*. Yellow oil. [α]_D²⁰ -9 (c, 0.05 in CHCl₃).

N-(3-Methylbutyl): **Muscicapine B**

[873684-19-6]


C₂₀H₂₉NO₂ 315.455

Alkaloid from the roots of *Croton muscicapa*. Yellow oil. [α]_D²⁰ -6 (c, 0.02 in CHCl₃).

Araújo-Júnior, V.T. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 553-557 (isol, pmr, cmr, ms)

3-Amino-5-guanidinopentanoic acid A-779

3-Amino-5-[(aminoiminomethyl)amino]pentanoic acid, 9CI. β-Arginine



C₆H₁₄N₄O₂ 174.202

N³-Me: 5-[(Aminoiminomethyl)amino]-3-(methylamino)pentanoic acid, 9CI. 5-Guanidino-3-methylaminopentanoic acid

[67396-07-0]

C₇H₁₆N₄O₂ 188.229

Component of Antibiotic LL-BM

547β.

McGahren, W.J. *et al.*, *J.O.C.*, 1977, **42**, 1282

Nomoto, S. *et al.*, *Chem. Lett.*, 1978, 589

(synth)

1-Amino-20-guanidino-4,8,12,16-tetrazaicosane A-780

22-Amino-2,7,11,15,19-pentaazadocosanimidamide

[864812-20-4]



C₁₇H₄₂N₈ 358.572

Isol. from the venom of the spider

Paracoelotes birulai.

N^ω-Ac: [864812-18-0]

C₁₉H₄₄N₈O 400.61


Isol. from the venom of *Paracoelotes birulai*.

Tzouros, M. *et al.*, *Toxicol.*, 2005, **46**, 350-354

1-Amino-16-guanidino-4,8,12-triazahexadecane A-781

18-Amino-2,7,11,15-tetraazaoctadecanimidamide, 9CI

[864812-13-5]



C₁₄H₃₅N₇ 301.477

Isol. from the venom of the spider

Paracoelotes birulai.

N¹-Ac: [864812-15-7]

C₁₆H₃₇N₇O 343.515

Isol. from the venom of the spider

Paracoelotes birulai.

4-N-Hydroxy: [864812-14-6]

C₁₄H₃₅N₇O 317.477


Isol. from the venom of the spider

Paracoelotes birulai.

Tzouros, M. *et al.*, *Toxicol.*, 2005, **46**, 350-354

(isol, 4-N-hydroxy, N¹-Ac)

2-Amino-7,15-hexadecadien-3-ol A-782



C₁₆H₃₁NO 253.427

(2S,3R,7Z)-form

Obscuraminol B


[350485-82-4]

Isol. from the tunicate *Pseudodistoma obscurum*. Oil (as di-Ac). [α]_D²⁴ -21.1 (c,

0.65 in MeOH) (di-Ac).

Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-4588 (isol, pmr, cmr, abs config)

2-Amino-11,15-hexadecadien-3-ol A-783



C₁₆H₃₁NO 253.427

(2S,3R,11E)-form

Obscuraminol C

[350484-95-6]

Isol. from the tunicate *Pseudodistoma obscurum*. Amorph. solid (as di-Ac). [α]_D²⁴ -24.5 (c, 0.83 in MeOH) (di-Ac).

11,12-Dihydro: 2-Amino-15-hexadecen-3-ol.

Obscuraminol F

[350485-01-7]

C₁₆H₃₃NO 255.443

Isol. from *Pseudodistoma obscurum*. Amorph. solid (as di-Ac). [α]_D²⁴ -19.2 (c, 0.71 in MeOH) (di-Ac).

15,16-Dihydro: 2-Amino-11-hexadecen-3-ol.

Obscuraminol E

[350485-00-6]


C₁₆H₃₃NO 255.443

Isol. from *Pseudodistoma obscurum*.

Oil (as di-Ac).

Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-4588 (*Obscuraminols C,E,F*, isol, pmr, cmr, abs config)

2-Amino-9-hexadecen-3-ol A-784



C₁₆H₃₃NO 255.443

(2S,3R,9Z)-form

Obscuraminol D

[350484-99-0]

Isol. from the tunicate *Pseudodistoma obscurum*. Oil (as di-Ac). [α]_D²⁴ -21.6 (c,

0.44 in MeOH) (di-Ac).


Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-

4588 (isol, ir, pmr, cmr, abs config)

3-Aminohexahydro-2H-azepin-2-one, 9CI A-785

3-Amino-ε-caprolactam

[671-42-1]



(R)-form

C₆H₁₂N₂O 128.174

(R)-form [28957-33-7]


Hydrochloride: [26081-03-8]

Solid (MeOH). [α]_D²⁶ +26.4 (c, 4 in 1M HCl) (>99% ee).

- Hydrobromide*: [16473-63-5]
[α]_D²⁰ +20.7 (c, 5 in H₂O).
- (S)-form** [21568-87-6]
Mp 71-72°. [α]_D²⁵ -34 (c, 4 in 1M HCl) (100% ee).
- Hydrochloride*: [26081-07-2]
Mp 270°. [α]_D²⁵ -27 (c, 2.98 in 1M HCl).
- Hydrobromide*: [16473-62-4]
Mp 285-288° dec. [α]_D²⁰ -20.7 (c, 5 in H₂O).
- N³-(6-Methyloctanoyl): Caprolactin B**
[151379-42-9]
C₁₅H₂₈N₂O₂ 268.398
Isol. from a marine bacterium. Cytotoxic agent. [α]_D²² +5.4 (c, 1.03 in CH₂Cl₂). Isol. as a mixt. with Caprolactin A. Data given is for the mixt. λ _{max} 219 ; 274 (sh) (MeOH).
- N³-(7-Methyloctanoyl): Caprolactin A**
[151379-41-8]
C₁₅H₂₈N₂O₂ 268.398
Isol. from a marine bacterium. Cytotoxic agent. Isol. as a mixt. with Caprolactin B. λ _{max} 219 ; 274 (MeOH) (Berdy).
- (±)-form** [17929-90-7]
Cryst. (EtOAc). Mp 68-71° Mp 147-149°. Bp_{0.1} 105°.
- Hydrochloride*: [29426-64-0]
Mp 294-296°.
- Picrate*:
Needles (H₂O). Mp 233° dec.
- N-Ac**:
C₈H₁₄N₂O₂ 170.211
Cryst. (EtOAc). Mp 160-162°.
- N-Benzoyl**:
C₁₃H₁₆N₂O₂ 232.282
Cryst. (EtOH). Mp 203-211°.
- Pellegata, R. *et al.*, *Synthesis*, 1978, 614 (*synth, S-form*)
Boyle, W.J. *et al.*, *J.O.C.*, 1979, **44**, 4841 (*synth*)
Davidson, B.S. *et al.*, *Tetrahedron*, 1993, **49**, 6569-6574 (*Caprolactins*)
Uchikawa, J. *et al.*, *J. Het. Chem.*, 1994, **31**, 877 (*N-Ac, N-Benzoyl*)
Sakai, K. *et al.*, *Tetrahedron: Asymmetry*, 2003, **14**, 3713-3718 (*resoln*)
Urbanczyk-Lipkowska, Z. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 1254-1256 (*resoln, S-form*)

2-Amino-4'-hydroxyacetophenone A-786

2-Amino-1-(4-hydroxyphenyl)ethanone,
9CI. p-Hydroxyphenacylamine
[77369-38-1]



C₈H₉NO₂ 151.165
Plates (EtOH). Mp 190-193° dec.

Hydrochloride: [19745-72-3]
Needles (HCl aq.), prisms (EtOH). Mp 241-245° Mp 249-251° dec.


- N-(4-hydroxy-3-methoxy-E-cinnamoyl): Terrestriamide**
[157536-49-7]
C₁₈H₁₇NO₅ 327.336
Alkaloid from the fruit of *Tribulus terrestris*. Light yellow cryst. Mp 218-220°.
- N-Di-Me**:
C₁₀H₁₃NO₂ 179.218
Prisms (Et₂O/petrol). Mp 142°.
- Me ether: 2-Amino-4'-methoxyacetophenone**
[3883-94-1]
C₉H₁₁NO₂ 165.191 HCl
Cryst. (EtOH) (as hydrochloride). Mp 200-201° (hydrochloride). CAS no. refers to hydrochloride.

[40513-43-7]

- Tutin, F. *et al.*, *J.C.S.*, 1910, **97**, 2520
Mannich, C. *et al.*, *Ber.*, 1911, **44**, 1547
Voswinckel, H. *et al.*, *Ber.*, 1912, **45**, 1005
Mannich, C. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1915, **253**, 193
Corrigan, J.R. *et al.*, *J.A.C.S.*, 1945, **67**, 1894
Yinglin, H. *et al.*, *Synthesis*, 1990, 615 (*Me ether, synth, ir, pmr*)
Ren, Y.J. *et al.*, *Yaoxue Xuebao*, 1994, **29**, 204 (*Terrestriamide*)

2-Amino-3-hydroxybenzoic acid A-787

3-Hydroxyanthranilic acid
[548-93-6]



- C₇H₇NO₃ 153.137
Metab. of Tryptophan, T-640 in humans, rats and *Claviceps* spp. Isol. from cultures of *Klebsiella pneumoniae*. Also isol. from *Brassica oleracea* (cauliflower). Leaflets (H₂O). Mp 164°. pK_{a2} 5.19; pK_{a3} 10.12 (20°).

▶ Exp. carcinogen. DG2625000

Hydrochloride: [4920-81-4]
Cryst. (HCl). Mp 198-200°.

- N-Ac, amide: 2-Acetamido-3-hydroxybenzamide**
C₉H₁₀N₂O₃ 194.19
Prod. by *Streptomyces* sp. GW37/3236. Solid. λ _{max} 240 (sh) ; 295 (log ϵ 3.68) (MeOH).

Me ether: 2-Amino-3-methoxybenzoic acid. 3-Methoxyanthranilic acid
[3177-80-8]
C₈H₉NO₃ 167.164
Leaflets (AcOH). Mp 170-171°.

Trifluoromethyl ether:
C₈H₆F₃NO₃ 221.136
Needles (H₂O). Mp 99-101°.

O,N-Di-Me: 3-Methoxy-2-methylamino-benzoic acid. Damascenic acid. Damascenic acid
[485-27-8]
C₉H₁₁NO₃ 181.191
Mp 144°. Forms a trihydrate, Mp 80°.

O,N-Di-Me; hydrochloride: Mp 210-211°.

O,N-Di-Me, Me ester: **Damascenine**
[483-64-7]

C₁₀H₁₃NO₃ 195.218
Alkaloid from seeds of *Nigella damascena* (Ranunculaceae). Mp 24-26°. Bp 270° dec. Bp₁₀ 147-148°.

O,N-Di-Me, Me ester; hydrochloride:
Cryst. + 1H₂O. Mp 122° Mp 156° (anhyd.).

O,N-Di-Me, Me ester, picrate: Mp 158-159°.

Me ether, nitrile: 2-Amino-3-methoxybenzonitrile. 2-Cyano-6-methoxyaniline
[148932-68-7]
C₈H₈N₂O 148.164
Powder. Mp 64-67°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 206C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1098C; 1123C (*nmr*)
CRC Atlas of Spectral Data and Physical Constants, b842 (*uv, ir*)

Ewins, A.J. *et al.*, *J.C.S.*, 1912, **101**, 544-552 (*Damascenine*)

Froelicher, V. *et al.*, *J.C.S.*, 1921, **119**, 1425-1432 (*Me ether, synth*)

Muelle, F. *et al.*, *Planta*, 1961, **57**, 403-477 (*isol*)
Munsche, D. *et al.*, *Phytochemistry*, 1965, **4**, 705-712 (*biosynth*)

Kuhn, R. *et al.*, *Chem. Ber.*, 1968, **101**, 3597-3603 (*Damascenine*)

Kuznezova, L.E. *et al.*, *Nature (London)*, 1969, **222**, 484-485 (*props*)

Thoinet, M. *et al.*, *Ann. Pharm. Fr.*, 1978, **36**, 337-339 (*synth, Damascenine*)

Mohr, N. *et al.*, *Annalen*, 1981, 1515-1518 (*isol*)

Okabe, N. *et al.*, *Acta Cryst. C*, 1996, **52**, 2345-2347 (*cryst struct*)

Goto, K. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 547-551 (*Me ether nitrile*)

Nagasaka, T. *et al.*, *J.O.C.*, 1998, **63**, 6797-6801 (*Me ether, synth, ir, pmr*)

Leroux, F. *et al.*, *J.O.C.*, 2003, **68**, 4693-4699 (*trifluoromethyl ether, synth, pmr, cmr*)

Fotso, S. *et al.*, *Z. Naturforsch., B*, 2003, **58**, 1242-1246 (*2-Acetamido-3-hydroxybenzamide*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, AKE750

2-Amino-4-hydroxybenzoic acid A-788

4-Hydroxyanthranilic acid
[38160-63-3]

C₇H₇NO₃ 153.137
Needles (H₂O). Mp 148° (dec.).

Hydrochloride:
Needles.

Me ester:
C₈H₉NO₃ 167.164
Needles. Mp 132-133°.

Me ether: 2-Amino-4-methoxybenzoic acid. 2-Aminoanisic acid. 4-Methoxyanthranilic acid
[4294-95-5]
C₈H₉NO₃ 167.164
Plates (EtOH). Mp 166° (dec.) Mp 190°. pK_{a1} 2.06; pK_{a2} 4.88 (25°, 0.1M KCl).

N-(4-Hydroxycinnamoyl) (E)-: 4-Hydroxy-N-(4-hydroxycinnamoyl)anthranilic acid. Avenanthramide G
C₁₆H₁₃NO₅ 299.282

Stress metab. in oats (*Avena sativa*) induced by Victorin M. Cryst. + 1/4 MeOH (MeOH/1% AcOH aq.). Mp

290-291°.

CRC Atlas of Spectral Data and Physical Constants, b843 (uv)Drain, D.G. *et al.*, *J.C.S.*, 1949, 1489 (synth)
van der Stelt, C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1953, **72**, 195 (synth)Jilek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1985, **50**, 519 (synth, uv, ir, pmr, deriv)Paulidis, V.H. *et al.*, *Synth. Commun.*, 1994, **24**, 533 (Me ether, synth, ir, pmr, cmr)Miyagawa, H. *et al.*, *Phytochemistry*, 1996, **41**, 1473 (Avenanthramide G)Boojamra, C.G. *et al.*, *J.O.C.*, 1997, **62**, 1240-1256 (Me ether)**2-Amino-5-hydroxybenzoic acid** A-789*5-Hydroxyanthranilic acid*
[394-31-0]C₇H₇NO₃ 153.137Metab. of Tryptophan in rats. Oral hypoglycaemic agent. Violet cryst. (H₂O). Mp 252° dec. pK_{a1} 2.72; pK_{a2} 5.37 (25°, 0.1M KCl). Log P 0.66 (calc).*Me ester*: [1882-72-0]C₈H₉NO₃ 167.164

Yellow needles. Mp 158°.

Et ester:C₉H₁₁NO₃ 181.191

Cryst. (EtOH). Mp 140°.

8-Methylonyl ester: Isodecyl 5-hydroxyanthranilate. Antibiotic BU 4601A. BU 4601A
[148915-76-8]C₁₇H₂₇NO₃ 293.405Prod. by a *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. MeOH, CCl₄, EtOH, DMSO, EtOAc; fairly sol. hexane; poorly sol. H₂O. λ_{max} 212 (ε 29200); 234 (ε 9100); 300 (ε 3800) (MeOH/HCl) (Derep). λ_{max} 208 (ε 33800); 225 (ε 24600); 372 (ε 4600) (MeOH/NaOH) (Derep). λ_{max} 219 (ε 24600); 246 (sh) (ε); 360 (ε 5200) (MeOH) (Derep).*9-Methyldecyl ester: Isondecyl 5-hydroxyanthranilate. Antibiotic BU 4601B. BU 4601B*
[148915-77-9]C₁₈H₂₉NO₃ 307.432Prod. by a *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. MeOH, EtOAc, EtOH, DMSO, CCl₄; fairly sol. hexane; poorly sol. H₂O. λ_{max} 212 (ε 29200); 234 (ε 9100); 300 (ε 3800) (MeOH/HCl) (Derep). λ_{max} 208 (ε 33800); 225 (ε 24600); 372 (ε 4600) (MeOH/NaOH) (Derep). λ_{max} 219 (ε 24600); 246 (sh) (ε); 360 (ε 5200) (MeOH) (Derep).*10-Methylundecyl ester: Isolauryl 5-hydroxyanthranilate. Antibiotic BU 4601C. BU 4601C*
[148915-78-0]C₁₉H₃₁NO₃ 321.459Prod. by a *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. MeOH, CCl₄, EtOH, DMSO, EtOAc; fairly sol. hexane; poorly sol. H₂O. λ_{max} 212 (ε 29200); 234 (ε 9100); 300 (ε 3800) (MeOH/HCl) (Derep). λ_{max} 208 (ε 33800); 225 (ε 24600); 372 (ε 4600) (MeOH/NaOH) (Derep). λ_{max} 219 (ε 24600); 246 (sh) (ε); 360 (ε 5200)(MeOH) (Derep). λ_{max} 208 (ε 33900); 223 (ε 4700); 372 (ε 4100) (MeOH-NaOH) (Berdy).*N-(4-Hydroxybenzoyl): 5-Hydroxy-N-(4-hydroxybenzoyl)anthranilic acid.***Melandrin**

[110846-17-8]

C₁₄H₁₁NO₅ 273.245Constit. of *Melandrin firmum*. Prisms (MeOH). λ_{max} 237; 290; 336 (MeOH).*N-(4-Hydroxycinnamoyl) (E-): 5-Hydroxy-N-(4-hydroxycinnamoyl)anthranilic acid. Avenanthramide A. Avenalumin I*
[108605-70-5]C₁₆H₁₃NO₅ 299.282Isol. from the grains of oats (*Avena sativa*). Phytoalexin. Pale yellow needles (Me₂CO aq.). Mp 277°. Darkens at 235°. Originally descr. as Avenalumin I, given the wrong struct. λ_{max} 320 (ε 15400); 344 (ε 13400) (MeOH/HCl) (Derep). λ_{max} 366 (ε 16600) (MeOH/NaOH) (Derep). λ_{max} 317 (ε 15200); 336 (ε 15900) (MeOH) (Derep).*N-(4-Hydroxycinnamoyl) (Z-):* [116764-20-6]C₁₆H₁₃NO₅ 299.282Isol. from oats (*Avena sativa*).*N-(3,4-Dihydroxycinnamoyl) (E-): N-(3,4-Dihydroxycinnamoyl)-5-hydroxyanthranilic acid. Avenanthramide C*
C₁₆H₁₃NO₆ 315.282Isol. from the grains of *Avena sativa*.*N-(3,4-Dihydroxycinnamoyl) (Z-):*
[116764-22-8]C₁₆H₁₃NO₆ 315.282Isol. from oats (*Avena sativa*).*N-(4-Hydroxy-3-methoxycinnamoyl) (E-): N-Feruloyl-5-hydroxyanthranilic acid. Avenanthramide B. Avenanthramide 1*
[108605-69-2]C₁₇H₁₅NO₆ 329.309Isol. from the grains of *Avena sativa*. Long pale yellow needles (Me₂CO aq.). Sol. EtOAc, Et₂O, H₂O-Me₂CO; poorly sol. CHCl₃, C₆H₆, H₂O. Mp 246°. λ_{max} 228; 315; 361; 392 (MeOH-NaOH) (Berdy).*N-(4-Hydroxy-3-methoxycinnamoyl) (Z-):* [116764-21-7]C₁₇H₁₅NO₆ 329.309Isol. from oats (*Avena sativa*).*Me ether: 2-Amino-5-methoxybenzoic acid. 5-Methoxyanthranilic acid*
[6705-03-9]C₈H₉NO₃ 167.164Needles (H₂O). Mp 179-180° (155-156° dec.). pK_{a1} 2.37; pK_{a2} 5.57 (25°, 0.1M KCl).*Me ether, nitrile: 2-Amino-5-methoxybenzonitrile, 9CI. 2-Cyano-4-methoxyaniline*
[23842-82-2]C₈H₈N₂O 148.164

Solid. Mp 46°.

Trifluoromethyl ether: 2-Amino-5-(trifluoromethoxy)benzoic acid, 9CI
[83265-56-9]C₈H₆F₃NO₃ 221.136Needles (H₂O). Mp 137-138°.*Et ether: 2-Amino-5-ethoxybenzoic acid*C₉H₁₁NO₃ 181.191

Yellowish solid. Mp 179-180°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1122B (nmr)*CRC Atlas of Spectral Data and Physical Constants*, 6844 (uv)Puxeddu, E. *et al.*, *Gazz. Chim. Ital.*, 1929, **59**, 14; 492van der Stelt, C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1953, **72**, 195Zeitler, H.J. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1965, **340**, 73 (synth, props)Mitchell, S.C. *et al.*, *J. Chromatogr.*, 1978, **151**, 249Woo, W.S. *et al.*, *Phytochemistry*, 1987, **26**, 2099 (Melandrin)Collins, F.W. *et al.*, *J. Agric. Food Chem.*, 1989, **37**, 60 (Avenanthramides)Crombie, L. *et al.*, *Tet. Lett.*, 1990, **31**, 2647 (synth, Avenanthramides)Dimberg, L.H. *et al.*, *Cereal Chem.*, 1993, **70**, 637-641 (Avenanthramides)Ohkuma, H. *et al.*, *J. Antibiot.*, 1993, **46**, 705 (BU 4601)Paulidis, V.H. *et al.*, *Synth. Commun.*, 1994, **24**, 533 (Me ether)Zhang, P. *et al.*, *J. Med. Chem.*, 1995, **38**, 1679 (Et ether, synth, ir, pmr)Ishihara, A. *et al.*, *Phytochemistry*, 1999, **50**, 237-242 (Avenanthramides, biosynth)Leroux, F. *et al.*, *J.O.C.*, 2003, **68**, 4693-4699 (trifluoromethyl ether, synth, pmr, cmr)Manetsch, R. *et al.*, *Chem. Eur. J.*, 2004, **10**, 2487-2506 (Me ether nitrile)**3-Amino-5-hydroxybenzoic acid** A-790

[76045-71-1]

C₇H₇NO₃ 153.137

Key biogenetic precursor of aromatic nuclei in ansamycin, mitomycin and maytansinoid antibiotics.

Hydrochloride: [14206-69-0]

Mp 200-230°.

Me ester: [67973-80-2]C₈H₉NO₃ 167.164Cryst. (MeOAc/CHCl₃). Mp 125-127°.*N-Ac*: [93561-88-7]C₉H₉NO₄ 195.174


Amorph. (MeOH aq.). Mp 245-253°.

Me ether, nitrile: 3-Amino-5-methoxybenzonitrile, 9CI. 3-Cyano-5-methoxyaniline
[269411-71-4]C₈H₈N₂O 148.164

Yellow solid. Mp 81-85°.

Bickel, H. *et al.*, *Tetrahedron, Suppl.*, No. 8, 1966, 171-179 (synth)Herlt, A.J. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1319 (synth, ms, pmr)Kibby, J.J. *et al.*, *J. Antibiot.*, 1981, **34**, 605 (isol)Becker, A.M. *et al.*, *Tetrahedron*, 1983, **20**, 4189 (synth, pmr, ms)Becker, A.M. *et al.*, *Aust. J. Chem.*, 1984, **37**, 2103-2109 (use, N-Ac)Kim, C.G. *et al.*, *J.A.C.S.*, 1992, **114**, 4941; 1996, **118**, 7486 (biosynth)Watson, T.J.N. *et al.*, *Synth. Commun.*, 1999, **29**, 1379-1382 (synth, pmr, cmr)Reich, S.H. *et al.*, *J. Med. Chem.*, 2000, **43**, 1670-1683 (Me ether nitrile)Clark, J.D. *et al.*, *Org. Process Res. Dev.*, 2004, **8**, 571-575 (synth)Wang, Z. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 2242-2252 (synth, pmr, cmr)

2-Amino-4-(4-hydroxybenzyl)-1-methyl-1H-imidazole A-791
4-[(2-Amino-1-methyl-1H-imidazol-4-yl)methyl]phenol. **Dorimidazole A** [138935-50-9]




C₁₁H₁₃N₃O 203.243

Alkaloid from the nudibranch *Notodoris gardineri* and *Notodoris citrina*. Anthelmintic, antiparasitic. Yellow powder; cryst. (MeCN) (as hydrobromide). Mp 175-176° (hydrobromide). λ_{max} 228; 269; 297; 300 (MeOH) (Derep). λ_{max} 228; 269; 297; 360 (MeOH) (Berdy).

Alvi, K.A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1509-1515 (isol, uv, pmr, cmr, ms, struct, synth)
Molina, P. *et al.*, *J.O.C.*, 1999, **64**, 2540-2544 (synth, pmr, cmr)

3-Amino-4-hydroxybutanoic acid, 9CI A-792

γ-Hydroxy-β-aminobutyric acid. **GOBAB** [589-44-6]



C₄H₉NO₃ 119.12

Antiinflammatory, antifungal and antiseptic agent.

(R)-form [16504-56-6]

Hydrochloride: [196950-52-4]
Mp 177-178°.

Lactone: 4-Aminodihydro-2(3H)-furanone. β-Amino-γ-butyrolactone
C₄H₇NO₂ 101.105
Mp 175-177° (as hydrochloride). [α]_D²⁰ +56.7 (c, 1 in H₂O).

N-Benzyl, lactone: [152783-42-1]

C₁₁H₁₃NO₂ 191.229
Oil. [α]_D²⁵ +16.5 (c, 1.0 in CHCl₃).

(S)-form [16504-57-7] Pharmacol. active isomer.

Lactone: Mp 198-201° (as hydrobromide). [α]_D²⁰ -42.6 (c, 1.08 in H₂O).

N-Benzoyl, lactone:

C₁₁H₁₁NO₃ 205.213
Mp 123.5-126°. [α]_D²⁰ -97 (c, 1.41 in CHCl₃).

N-tert-Butyloxycarbonyl, lactone:

C₉H₁₅NO₄ 201.222
Solid. Mp 113-114° (106-108°). [α]_D²⁵ -54.3 (c, 1.15 in CHCl₃).

N-Benzoyloxycarbonyl, lactone:

C₁₂H₁₃NO₄ 235.239
Cryst. Mp 103-105°. [α]_D²⁰ -54.9 (c, 2.27 in CHCl₃).

N-Trifluoroacetyl, benzyl ester: [673470-50-3]

C₁₃H₁₄F₃NO₄ 305.253

Cryst. solid. Mp 70-71°. [α]_D²⁵ +9.1 (c, 0.99 in CHCl₃).

(±)-form [16504-55-5]

Needles. Mp 184°.

Lactone: [16504-58-8]

Mp 185-186° (as hydrochloride).

Lactone, N-benzyl: [157037-11-1]

C₁₁H₁₃NO₂ 191.229

Oil.

(ξ)-form

O-(4-Hydroxybenzoyl), N,N,N-tri-Me, betaine: 1-Carboxy-3-[(4-hydroxybenzoyl)oxy]-N,N,N-trimethyl-2-propanaminium inner salt, 9CI.

Zoaxanthellabetaïne A

[208256-89-7]

C₁₄H₁₉NO₅ 281.308

Isol. from the cultures of a symbiotic dinoflagellate *Symbiodinium* sp. Oil. [α]_D²⁰ -13 (c, 0.0035 in MeOH).

Jolles, P. *et al.*, *Bull. Soc. Chim. Fr.*, 1951, 862-863 (synth)

Chibnall, A.C. *et al.*, *Biochem. J.*, 1958, **68**, 122-128 (synth)

Nagai, K. *et al.*, *Arzneim.-Forsch.*, 1967, **17**, 1575-1577 (synth, isomers, pharmacol)

Kim, Y.B. *et al.*, *Arch. Pharmacol. Res.*, 1985, **8**, 1-6; *C.A.*, **103**, 98858d (cryst struct)

McGarvey, G.J. *et al.*, *J.A.C.S.*, 1986, **108**, 4943-4952 (lactone N-benzyloxycarbonyl, synth, ir, pmr)

Hvidt, T. *et al.*, *Can. J. Chem.*, 1988, **66**, 779-782 (synth, pmr, cmr)

Hanessian, S. *et al.*, *Can. J. Chem.*, 1993, **71**, 1407-1411 (lactone, N-tert-butyloxycarbonyl, synth, ir, pmr)

Maude, A.B. *et al.*, *J.C.S. Perkin 1*, 1997, 2513-2526 (synth, ir, pmr)

Calvisi, G. *et al.*, *Synlett*, 1997, 71-74 (lactone, synth, pmr)

Nakamura, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1998, **71**, 781-787 (Zooanthellabetaïne A)

El Hadri, A. *et al.*, *J. Med. Chem.*, 2002, **45**, 2824-2831 (N-benzyl lactone, synth)

Jackson, R.F.W. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 110-113 (N-trifluoroacetyl benzyl ester, synth, pmr, cmr)

1-Amino-3-hydroxy-8-decen-5-one A-793

H₃CCH=CHCH₂CH₂COCH₂-

CH(OH)CH₂CH₂NH₂

C₁₀H₁₉NO₂ 185.266

(+)-(E)-form

N-Ac: **Streptenol E**

C₁₂H₂₁NO₃ 227.303


Prod. by *Streptomyces* sp. strain A1.

Oil. [α]_D²² +33 (c, 0.34 in CHCl₃).

Puder, C. *et al.*, *J. Nat. Prod.*, 2001, **64**, 42-45

(2-Amino-1-hydroxyethyl)-phosphonic acid, 9CI A-794

[41744-58-5]



C₂H₈NO₄P 141.063

(R)-form [115511-00-7]

Isol. from the plasma membrane of *Acanthamoeba castellanii*. Cryst. (EtOH aq.). Mp 244-246° dec. [α]_D²⁰ -31.4 (c, 0.52 in H₂O).

(S)-form [121421-25-8]

Cryst. (EtOH aq.). Mp 245-247° dec. [α]_D²² +32.5 (c, 1 in H₂O).

(±)-form [114157-09-4]

Cryst. (EtOH aq.). Mp 260-262° dec.


Korn, E.D. *et al.*, *J. Biol. Chem.*, 1973, **248**, 2257 (isol)

Hammerschmidt, F. *et al.*, *Annalen*, 1988, 537-542; 955-960; 961-964; 1989, 577-583 (synth, abs config)

Heisler, A. *et al.*, *Tetrahedron: Asymmetry*, 1993, **4**, 959-960 (synth)

4-Amino-2-hydroxy-5-(hydroxymethyl)pyrimidine A-795

4-Amino-2-hydroxy-5-pyrimidinemethanol. 4-Amino-5-hydroxymethyl-2(1H)-pyrimidinone. 5-Hydroxymethylcytosine [1123-95-1]



C₅H₇N₃O₂ 141.129

DNA constit. of various plants and viruses. Cryst. + 0.5H₂O (H₂O). Mp 300° dec. pK_{a1} 4.32; pK_{a2} 13.

OH-form

2-Me ether: 4-Amino-5-(hydroxymethyl)-2-methoxypyrimidine. 4-Amino-2-methoxy-5-pyrimidinemethanol. **Bacimethrin** C₆H₉N₃O₂ 155.156

Prod. by *Bacillus megaterium* and *Streptomyces albus*. Active against gram-positive and -negative bacteria. Thiamine antagonist. Needles (MeOH or EtOH/Et₂O). Sol. H₂O, MeOH, AcOH; fairly sol. EtOH, Py; poorly sol. butanol, hexane. Mp 175°. λ_{max} 230 (ε 8540); 263 (ε 9590) (0.1M HCl) (Derep). λ_{max} 228 (ε 8760); 271 (ε 8170) (0.1M NaOH) (Derep). λ_{max} 228 (ε 8310); 270 (ε 7630) (H₂O) (Derep). λ_{max} 229 (ε 6600); 273 (ε 6200) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) 300 mg/kg.

2-Me ether, O-Ac:

C₈H₁₁N₃O₃ 197.193

Mp 131°.

2-Me ether, O,N-Di-Ac:

C₁₀H₁₃N₃O₄ 239.23

Mp 142°.

Wyatt, G.R. *et al.*, *Biochem. J.*, 1953, **55**, 774 (isol)

Miller, C.S. *et al.*, *J.A.C.S.*, 1955, **77**, 752 (synth, uv)

Tanaka, F. *et al.*, *J. Antibiot., Ser. A*, 1961, **14**, 161 (Me ether, isol, struct)

Koppel, H.C. *et al.*, *J.O.C.*, 1962, **27**, 1492; 3614 (Me ether, synth, uv)


Drautz, M. *et al.*, *J. Antibiot.*, 1987, **40**, 1431 (Me ether, isol, synth, props)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (rev)

Hudson, R.H.E. *et al.*, *Can. J. Chem.*, 2007, 302-312 (*synth, pmr, cmr*)

2-Amino-4-hydroxy-4-(5-hydroxy-2-pyridinyl)-3-methylbutanoic acid A-796

α -Amino- γ ,5-dihydroxy- β -methyl-2-pyridinebutanoic acid, 9CI



C₁₀H₁₄N₂O₄ 226.232

(2S,3R,4R)-form [76249-78-0]

Amino acid residue from Nikkomycin Z, N-218.

(2S,3S,4S)-form

Nikkomycin D

[67676-66-8]

The terminal amino acid of the nikkomycins and neopolyoxins. Fine needles (H₂O). Mp 230° dec. Mp 207° dec. [α]_D²⁰ -13.4 (c, 0.59 in H₂O).

[76612-52-7, 83572-53-6, 83572-54-7, 83572-56-9, 83572-55-8]


Koenig, W.A. *et al.*, *Annalen*, 1980, 1728; 1982, 1615 (*isol, synth, cryst struct*)

Teintze, M. *et al.*, *Biochemistry*, 1981, 20, 6446 (*isol*)

Uramoto, M. *et al.*, *Tetrahedron*, 1982, 38, 1599 (*isol*)

2-Amino-3-hydroxy-3-(4-imidazolyl)propanoic acid A-797

β -Hydroxyhistidine, 9CI



(2R,3R)-form

C₆H₉N₃O₃ 171.155

(2R,3R)-form

D-threo-form

[35849-15-1 (hydrochloride)]

Pale yellow crust. (as hydrochloride).

[α]_D²⁵ +15 (c, 1.2 in H₂O) (hydrochloride).

N⁺-Ac: [α]_D²⁵ -17 (c, 1.25 in H₂O).

(2S,3R)-form

L-erythro-form

[41215-80-9]

[51077-31-7 (hydrochloride)]

Component of antibiotics of the Bleomycin A₂ group. Cryst. Mp 205° dec. [α]_D²⁸ +40 (c, 1 in H₂O) (hydrochloride).

(2S,3S)-form

L-threo-form

[148219-09-4]

Constit. of a pyoverdin isol. from *Pseudomonas fluorescens* 244. Pale yellow solid. [α]_D -9 (c, 1 in H₂O).

(2RS,3RS)-form

(\pm)-threo-form

[35849-17-3]

Solid (as hydrochloride). CAS no. refers to hydrochloride.

(2RS,3SR)-form

(\pm)-erythro-form

[71389-30-5]

Cryst. Mp 205° dec.

Hydrochloride: [35849-19-5]

Needles (2-propanol/EtOH aq.). Mp 228°.

Takita, T. *et al.*, *J. Antibiot.*, 1971, 24, 795 (*isol, nmr, struct, synth*)

Ohashi, Y. *et al.*, *Agric. Biol. Chem.*, 1973, 37, 2283 (*isol*)

Koyama, G. *et al.*, *J. Antibiot.*, 1973, 26, 109 (*cryst struct*)

Naganawa, H. *et al.*, *J. Antibiot.*, 1977, 30, 388 (*cmr*)

Hecht, S.M. *et al.*, *J.A.C.S.*, 1979, 101, 3982 (*synth, pmr*)

Owa, T. *et al.*, *Chem. Lett.*, 1988, 83; 1873


(*synth*)

Hancock, D.K. *et al.*, *Chem. Comm.*, 1993, 468 (*isol, pmr, cmr*)

Dong, L. *et al.*, *J.O.C.*, 2002, 67, 4759-4770

(*S,S*-form, *synth, pmr, cmr*)

3-Amino-4-hydroxymethyl-14-methyl-9,19-cyclopregn-4-en-20-one A-798



C₂₃H₃₅NO₂ 357.535

3 β -form

N-Me: **30-Hydroxycyclomicrobuxene**


[220705-48-6]

C₂₄H₃₇NO₂ 371.562

Alkaloid from the roots of *Buxus sempervirens*. Amorph. solid. [α]_D²⁰ +84 (c, 0.69 in CHCl₃). Misleading synonym. λ _{max} 202 (MeOH).

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, 12, 299-306 (*isol, pmr, cmr*)

7-Amino-1-hydroxymethyl-6-methyl-5,8-isoquinolinedione, 9CI A-799



C₁₁H₁₀N₂O₃ 218.212

O-Angeloyl: **Cribrostatins 3**

[276682-84-9]

C₁₆H₁₆N₂O₄ 300.313

Alkaloid from *Cribrorchalina* sp. Cytotoxic agent. Orange-red needles (CH₂Cl₂/MeOH). Mp 190-192°.

N⁷-Me, *O*-angeloyl: **Cribrostatins 5**

[276682-90-7]

C₁₇H₁₈N₂O₄ 314.34

Alkaloid from *Cribrorchalina* sp. Cytotoxic agent. Red-brown plates (MeOH/CH₂Cl₂).

N⁷-(2-Sulfoethyl), *O*-angeloyl: **Cribrostatins 7**

[683276-94-0]

C₁₈H₂₀N₂O₇S 408.431

Alkaloid from a *Petrosia* sp. Cytotoxic agent. Dark red film. λ _{max} 210 (€ 14500); 282 (€ 6500); 338 (€ 2000); 474 (€ 1700) (MeOH).


Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2000, 63, 793-798 (*Cribrostatins 3, Cribrostatins 5*)

Sandoval, I.T. *et al.*, *Nat. Prod. Res.*, 2004, 18, 89-93 (*Cribrostatins 7*)

2-Amino-4-hydroxy-3-methylpentanoic acid A-800

2-Amino-2,3,5-trideoxy-3-methylpentonic acid. 4-Hydroxyisoleucine

[50764-07-3]



(2R,3R,4R)-form

C₆H₁₃NO₃ 147.174

(2R,3R,4R)-form

D-xylo-form. 2-Amino-2,3,5-trideoxy-3-methyl-*D*-xyloic acid

[60010-78-8]

Cryst. (EtOH aq.). Mp ca.° 230. [α]_D²⁰ -35.6 (c, 1.2 in D₂O).

1,4-Lactone: 3-Aminodihydro-4,5-dimethyl-2(3H)-furanone

C₆H₁₁NO₂ 129.158

Cryst. (EtOH/petrol) (as hydrochloride). Mp 222° (hydrochloride). [α]_D²⁵ +88.7 (c, 1 in MeOH).

1,4-Lactone, *N*-Ac:

Needles (EtOH). Mp 140°. [α]_D²⁵ +88 (c, 1 in DMSO).

(2R,3R,4S)-form

L-arabino-form. 2-Amino-2,3,5-trideoxy-3-methyl-*L*-arabinoic acid

[60010-73-3]

Minor amino acid constit. of *Trigonella foenum-graecum* (fenugreek) seeds. [α]_D²⁰ +1 (c, 1 in H₂O). Originally thought to be the 2R,3R,4R-form.

1,4-Lactone:

Cryst. (as hydrochloride). Mp 170-175° (hydrochloride).

1,4-Lactone, *N*-Ac: 3-Acetamidodihydro-4,5-dimethyl-2(3H)-furanone. **Desmodilactone**

[60010-74-4]

C₈H₁₃NO₃ 171.196

Constit. of *Desmodium styracifolium*. Component of Guang Jin Qian Cao.

Cryst. Mp 84-85°. [α]_D¹⁸ -16.4 (c, 0.11 in MeOH).

(2S,3R,4R)-form

D-lyxo-form. 2-Amino-2,3,5-trideoxy-3-methyl-*D*-lyxoic acid

[55399-92-3]

Cryst. (EtOH aq.). Mp ca.° 220. [α]_D²⁰ -

24.9 (c, 1.17 in D₂O).

1,4-Lactone:

Needles (EtOH/petrol) (as hydrochloride). Mp 255° (hydrochloride). $[\alpha]_D^{25} +24.3$ (c, 1 in MeOH).

1,4-Lactone, N-Ac:

Needles (EtOH). Mp 141°.

(2S,3R,4S)-form

L-ribo-form. 2-Amino-2,3,5-trideoxy-3-methyl-L-ribonic acid

[55399-93-4]

Major constit. of *Trigonella foenum-graecum* (fenugreek). Cryst. (EtOH aq.). Mp 224-225°. $[\alpha]_D^{20} +31$ (c, 1 in H₂O). Stereochem. revised in 1989.

1,4-Lactone:

Needles (EtOH/petrol) (as hydrochloride). Mp 230° (hydrochloride). $[\alpha]_D^{25} -15.5$ (c, 1 in MeOH).

1,4-Lactone, N-Ac:

Needles (EtOH). Mp 94°.

(2S,3S,4R)-form

D-arabino-form. 2-Amino-2,3,5-trideoxy-3-methyl-D-arabinonic acid

[21704-86-9]

Constit. of the flowers of *Quararibea funebris*. Mp 205-207°. $[\alpha]_D^{27} +2.9$ (c, 0.1 in H₂O). λ_{\max} 197 (ε 250) (H₂O).

1,4-Lactone: [71392-28-4]

[90693-46-2]

Constit. of the flowers of *Quararibea funebris*. Powder (as hydrochloride). Mp 212-215° (hydrochloride). $[\alpha]_D^{25} -14.8$ (c, 0.7 in MeOH).

Fowden, L. *et al.*, *Phytochemistry*, 1973, **12**, 1707-1711 (*Trigonella foenum-graecum* constits)

Gieren, A. *et al.*, *Annalen*, 1974, 1561-1569 (*synth*)

Hasan, M. *et al.*, *Annalen*, 1976, 781-787 (*lactone, synth, pmr*)

Raffauf, R.F. *et al.*, *J.O.C.*, 1984, **49**, 2714-2718 (*Quararibea funebris* constits)

Alcock, N.W. *et al.*, *Phytochemistry*, 1989, **28**, 1835-1841 (*struct*)

Inghardt, T. *et al.*, *Tetrahedron*, 1991, **47**, 6469-6482 (*synth, abs config, cryst struct*)


Yang, J. *et al.*, *Yaoxue Xuebao*, 1993, **28**, 197-201; *CA*, **119**, 156209m (*Desmodilactone*)

Wang, Q. *et al.*, *Eur. J. Org. Chem.*, 2002, 834-839 (*synth, pmr, cmr*)

Rolland-Fulcrand, V. *et al.*, *Eur. J. Org. Chem.*, 2004, 873-877 (*synth, pmr, cmr*)

Aouadi, K. *et al.*, *Synthesis*, 2007, 3399-3405 (*2S,3R,4R*-form, *synth, pmr, cmr, ms*)

2-Amino-8-(hydroxymethyl)-3H-phenoxazin-3-one A-801



C₁₃H₁₀N₂O₃ 242.234

N-Ac: 2-Acetamido-8-(hydroxymethyl)-3H-phenoxazin-3-one. **Exfoliazone**. *BT* 38. *Antibiotic BT* 38 [132627-73-7]

C₁₅H₁₂N₂O₄ 284.271

Prod. by *Streptomyces exfoliatus*. Active against *Valsa ceratosperma*. Or-


ange needles (CHCl₃). Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O, butanol. Mp 294-296°. λ_{\max} 238 (ε 75300); 400 (ε 42300) (solvent not reported) (Derep). λ_{\max} 238 (ε 75300); 400 (ε 42300) (MeOH) (Berdy).

Imai, S. *et al.*, *J. Antibiot.*, 1990, **43**, 1606

2-Amino-6-(hydroxymethyl)-4(1H)-pteridinone, 9CI A-802

6-Hydroxymethylpterine. **Ranachrome 3**. 2-Amino-4-hydroxy-6-(hydroxymethyl)pteridine

[712-29-8]



C₇H₇N₅O₂ 193.165

Occurs widely in nature. The 7,8-dihydro deriv. is the biological precursor of folic acid. Cryst. (H₂O). Mp 320° dec.

► UO3450000

O-[4-O-Methyl-α-D-glucuronopyranosyl-(1→6)-β-D-galactopyranoside]: **Cyanopterin**

[223532-84-1]

C₂₀H₂₇N₅O₁₃ 545.459

Isol. from *Synechocystis* sp. PCC 6803. λ_{\max} 192 ; 236 ; 276 ; 345 (H₂O).

O-Ac: [73978-45-7]

C₉H₉N₅O₃ 235.202

Cryst. (H₂O). Mp 300° dec.

N²,O-Di-Ac: [32363-58-9]

C₁₁H₁₁N₅O₄ 277.239

Cryst. (Me₂CO). Mp 239-242° dec. (227-229°).

[1501-52-6]

Blair, J.A. *et al.*, *Org. Mass Spectrom.*, 1969, **2**, 923 (*ms*)

Thijssen, H.H.W. *et al.*, *Anal. Biochem.*, 1973, **54**, 609 (*synth*)

Boyle, P.H. *et al.*, *Chem. Ber.*, 1980, **113**, 1514 (*synth, uv, pmr*)

Waring, P. *et al.*, *Aust. J. Chem.*, 1985, **38**, 629 (*synth, bibl*)

Zeitler, H.J. *et al.*, *Methods Enzymol.*, 1986, **122**, 273 (*rev, hplc*)


Burgmayer, S.J.N. *et al.*, *Inorg. Chem.*, 1988, **27**, 4059 (*ir*)

Lee, H.W. *et al.*, *Biochim. Biophys. Acta*, 1999, **1410**, 61-70 (*Cyanopterin*)

4-Amino-2-hydroxy-5-methylpyrimidine A-803

4-Amino-5-methyl-2(1H)-pyrimidinone, 9CI. 4-Amino-5-methyl-2-pyrimidinol. 5-Methylcytosine

[554-01-8]



C₅H₇N₃O 125.13

Constit. of DNA of plants and animals. Cryst. (H₂O). Mp 287-292°.

Hydrochloride:

Plates or prisms. Mp 288-290°.

1-β-D-(2,3-Dideoxyribofuranosyl): 2',3'-Dideoxy-5-methylcytidine, 9CI

[107036-56-6]

C₁₀H₁₅N₃O₃ 225.247

Lyophilised solid.

Fox, J.J. *et al.*, *J.A.C.S.*, 1951, **73**, 3256 (*synth*)

Kream, J. *et al.*, *J.A.C.S.*, 1952, **74**, 5157 (*isol*)

Thomas, A.J. *et al.*, *Biochem. J.*, 1956, **62**, 1 (*isol*)

Kokko, J. *et al.*, *J.A.C.S.*, 1962, **84**, 1042 (*pmr*)

Rice, J.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4569 (*ms*)

Kloetzer, W. *et al.*, *Monatsh. Chem.*, 1965, **96**, 1721 (*synth*)


Padmaja, N. *et al.*, *Acta Cryst. C*, 1987, **43**, 2157 (*cryst struct*)

Kim, C.H. *et al.*, *J. Med. Chem.*, 1987, **30**, 862 (β-D-2,3-dideoxyribofuranosyl, *synth, pmr, ms*)

Buchini, S. *et al.*, *Eur. J. Org. Chem.*, 2006, 3152-3168 (*synth, pmr, cmr, ms*)

3-Amino-5-hydroxy-7-oxabicyclo[4.1.0]hept-3-en-2-one, 9CI A-804

2-Amino-5,6-epoxy-4-hydroxy-2-cyclohexen-1-one



(1R,5R,6R)-form

C₆H₇NO₃ 141.126

(1R,5R,6R)-form

N-Ac: *Antibiotic MT* 35214. *MT* 35214 [89020-32-6]

C₈H₉NO₄ 183.163

Semisynthetic. Weakly active against gram-positive and -negative bacteria and *Candida* spp. Prisms (Me₂CO/Et₂O). Mp 149-151° dec. $[\alpha]_D^{20} +104$ (c, 1 in MeOH). λ_{\max} 265 (ε 5910); 321 (ε 12700) (0.1N NaOH) (Derep). λ_{\max} 212 (ε 11000); 276 (ε 3510) (H₂O) (Derep).

N-Ac, 5-ketone: 3-(Acetylamino)-7-oxabicyclo[4.1.0]hept-3-ene-2,5-dione. 2-(Acetylamino)-5,6-epoxy-2-cyclohexene-1,4-dione. *Antibiotic MT* 36531. *MT* 36531

[89020-31-5]

C₈H₇NO₄ 181.148

Semisynthetic. Shows weak antibacterial activity. Needles (EtOAc/Et₂O). Mp 144-145°. $[\alpha]_D^{20} -99$ (c, 0.5 in MeOH).

(1S,5S,6S)-form

Antibiotic MM 14201. *MM* 14201

[89020-30-4]

Isol. from *Streptomyces* spp. Sol. MeOH, H₂O; poorly sol. hexane. $[\alpha]_D^{20} -202$ (c, 0.334 in MeOH). λ_{\max} 304 (H₂O) (Derep). λ_{\max} 304 (EtOH) (Berdy).

N-Ac: N-(5-Hydroxy-2-oxo-7-oxabicyclo[4.1.0]hept-3-en-3-yl)acetamide.

Antibiotic LL C10037α. *LL* C10037α


[93752-54-6]
 $C_8H_9NO_4$ 183.163
 Isol. from *Streptomyces* spp. Shows antitumour props. Weakly active against bacteria. Fluffy powder. Sol. H_2O , MeOH, EtOH; poorly sol. Me_2CO , hexane. Mp 153°. $[\alpha]_D^{25}$ -155 (c, 0.1 in H_2O). $[\alpha]_D^{20}$ -202 (c, 0.3 in MeOH). Revised struct. λ_{max} 265 (ε 5910); 321 (ε 12700) (0.1N NaOH) (Derep). λ_{max} 212 (ε 11000); 276 (ε 3510) (H_2O) (Derep). λ_{max} 212 (ε 10970); 275 (ε 3510) (H_2O) (Berdy). λ_{max} 265 (ε 5200); 321 (ε 15200) (NaOH) (Berdy).

(1RS,5RS,6RS)-form


N-Ac: [157183-80-7]
 Cryst. (Me_2CO/Et_2O). Mp 167° dec.
 N-Ac, 5-ketone:
 Pale yellow solid ($EtOAc$ /petrol). Mp 169-170° dec.

(1RS,5SR,6RS)-form


N-Ac: (±)-Epi-LL-C10037α
 [157183-81-8]
 Mp 154° dec.
 [121055-94-5, 121055-92-3]
 Box, S.J. et al., *J. Antibiot.*, 1983, **36**, 1631 (isol, props)
 Lee, M.D. et al., *J. Antibiot.*, 1984, **37**, 1149 (isol, struct)
 Whittle, Y.G. et al., *J.A.C.S.*, 1987, **109**, 5043 (biosynth, struct)
 Gould, S.J. et al., *J.A.C.S.*, 1989, **111**, 7932 (biosynth)
 Shen, B. et al., *J.O.C.*, 1990, **55**, 4422 (pmr, cryst struct)
 Wipf, P. et al., *J.O.C.*, 1994, **59**, 3520 (synth, LL C10037)
 Wipf, P. et al., *Synthesis*, 1995, 1549 (synth)
 Kapfer, I. et al., *Tet. Lett.*, 1996, **37**, 2101 (synth, LL C10037α)
 Taylor, R.J.K. et al., *Synthesis*, 1998, 775-790 (synth, LL-C10037α, 5-ketone, ir, pmr, cmr, ms)
 Macdonald, G. et al., *Tet. Lett.*, 1998, **39**, 5433-5436 (synth)
 Murphy, S.T. et al., *Org. Lett.*, 1999, **1**, 1483-1485 (synth, LL C10037α)
 Block, O. et al., *J.O.C.*, 2000, **65**, 716-721 (synth, LL C10037α)

8-Amino-1-hydroxyphenazine A-805
8-Amino-1-phenazolinol, 9CI

$C_{12}H_9N_3O$ 211.223
 Prod. by an unidentified bacterium. λ_{max} 275; 376; 480 (MeOH) (Berdy).
 Me ether: 8-Amino-1-methoxyphenazine
 [23448-77-3]
 $C_{13}H_{11}N_3O$ 225.249
 Bright orange solid. Mp 180° Mp 240-243° (dec.) (double Mp). λ_{max} 275; 376; 480 (EtOH).
 Gerber, N.N. et al., *J. Het. Chem.*, 1969, **6**, 297-300 (isol)

2-Amino-5-[[[3-(4-hydroxyphenyl)-2-methoxy-2-propenoyl]amino](imino)methyl]amino]pentanoic acid A-806

$C_{16}H_{22}N_4O_5$ 350.374
 Isol. from the ascidian *Atrium robustum*. Amorph. solid. $[\alpha]_D^{25}$ -3.2 (c, 0.15 in H_2O). λ_{max} 295 (ε 34000) (H_2O).
 Kehraus, S. et al., *J. Med. Chem.*, 2004, **47**, 2243-2255 (isol)

1-(3-Amino-4-hydroxyphenyl)-1,2-propanediol A-807


$C_9H_{13}NO_3$ 183.207

(1R,2R)-form

4'-Me ether, N-Ac: *Agaridiol*
 [243864-56-4]
 $C_{12}H_{17}NO_4$ 239.271
 Prod. by an *Agaricus* sp. Antibacterial agent.
 Berg, A. et al., *J. Basic Microbiol.*, 1999, **39**, 213-215

3-Amino-2-hydroxy-3-phenylpropanoic acid A-808

β-Amino-α-hydroxybenzenepropanoic acid, 9CI. 3-Amino-3-phenyllactic acid. 3-Phenylisoserine
 [6049-55-4]



$C_9H_{11}NO_3$ 181.191


(2R,3S)-form

(-)-threo-form
 [136561-53-0] Residue present in, and intermed. in synth. of, the sidechain of Taxol, T-52.
 Solid. Mp 224-226° Mp 238° dec. (as hydrochloride). $[\alpha]_D$ -14.9 (c, 0.55 in 6M HCl). $[\alpha]_D^{25}$ -14.5 (c, 0.37 in MeOH) (as hydrochloride).
 N-Benzoyl: [132201-33-3]
 $C_{16}H_{15}NO_4$ 285.299
 Isol. from *Taxus baccata*. Sidechain of Taxol, T-52. Cryst. ($EtOAc$). Mp 177-178° (166-168°). $[\alpha]_D^{25}$ -36.3 (c, 0.73 in

EtOH). $[\alpha]_D^{25}$ -31.7 (c, 1.0 in EtOH).
 Harada, K. et al., *J.O.C.*, 1966, **31**, 1407 (abs config)
 Kamandi, E. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1974, **307**, 871-878 (synth, amide)
 Harada, K. et al., *Bull. Chem. Soc. Jpn.*, 1974, **47**, 2911 (synth, abs config)
 Bruncko, M. et al., *Angew. Chem., Int. Ed.*, 1997, **36**, 1483-1486 (synth, 2R,3S-form)
 Zhou, Z. et al., *Synth. Commun.*, 2001, **31**, 3609-3615; 2003, **33**, 723-728 (synth, ir, pmr, amide, N-benzoyl)


2-Amino-6-[3-hydroxy-4-(phosphonoxy)-1-butenyl]-4(1H)-pteridinone, 9CI A-809

2-Amino-4-hydroxy-6-(3-hydroxy-4-phosphonoxy-1-butenyl)pteridine
 [136338-57-3]



$C_{10}H_{12}N_5O_6P$ 329.208
 Isol. from *Rhodospseudomonas sphaeroides*.

Seo, N. et al., *Biochim. Biophys. Acta*, 1991, **1074**, 439 (isol, synth)

20-Amino-18-hydroxypregna-1,4-dien-3-one A-810


$C_{21}H_{31}NO_2$ 329.481

(20S)-form

N-Me: 18-Hydroxy-20-(methylamino)-pregna-1,4-dien-3-one
 [113846-36-9]
 $C_{22}H_{33}NO_2$ 343.508
 Alkaloid from the leaves of *Didymeles madagascariensis* (Didymelaceae). $[\alpha]_D$ +31 (c, 1.71 in EtOH).
 N,N-Di-Me: 20-(Dimethylamino)-18-hydroxypregna-1,4-dien-3-one
 [91147-28-3]
 $C_{23}H_{35}NO_2$ 357.535
 Alkaloid from the stem bark of *Didymeles madagascariensis* (Didymelaceae). λ_{max} 241 (log ε 4.39); 322 (log ε 4.07) (EtOH).
 18-Aldehyde, N,N-di-Me: 20-(Dimethylamino)-3-oxopregna-1,4-dien-18-al
 [91147-29-4]
 $C_{23}H_{33}NO_2$ 355.519
 Alkaloid from the stem bark of *Didymeles madagascariensis* (Didymelaceae). Cryst. (Et_2O). Mp 157°. $[\alpha]_D$ +69 (c, 0.01 in $CHCl_3$).
 Sánchez, V. et al., *Bull. Soc. Chim. Fr., Part II*, 1984, **71**; 1987, 877 (isol, uv, cd, pmr, cmr, ms, struct)

3-Amino-12-hydroxy-pregnan-20-one

A-811

C₂₁H₃₃NO₂ 331.497C₂₁H₃₅NO₂ 333.513**(3β,5α,12β)-form****Bokitamine**

[5224-04-4]

Alkaloid from a *Holarrhena* sp. (poss. *Holarrhena wulfsbergii*) (Apocynaceae). Cryst. (EtOAc). Mp 191°.

N-Ac:

C₂₃H₃₇NO₃ 375.55Mp 247-248°. [α]_D +22.5 (CHCl₃).

N,O-Di-Ac:

C₂₅H₃₉NO₄ 417.587


Mp 202-204°.

15,16-Didehydro-3-Amino-12-hydroxy-pregn-15-en-20-one. Kisantamine


[69375-36-6]

C₂₁H₃₃NO₂ 331.497Alkaloid from *Holarrhena congolensis* (Apocynaceae). [α]_D²⁰ -16 (c, 0.1 in CHCl₃) (as N-Ac).Nellé, S. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1970, **271**, 153 (isol, struct, synth)Dadoun, H. et al., *Plant. Med. Phytother.*, 1978, **12**, 225; *CA*, **90**, 100106r (deriv)**3-Amino-14-hydroxypregnan-20-one**

A-812

C₂₁H₃₅NO₂ 333.513**(3α,5α,14β)-form** [215595-77-0]Alkaloid from *Holarrhena curtisii*. Cytotoxic and leishmanicidal agent. Amorph. [α]_D +23.3 (c, 0.26 in CHCl₃).Kam, T.-S. et al., *J. Nat. Prod.*, 1998, **61**, 1332-1336 (isol, pmr, cmr, ms)**3-Amino-2-hydroxypregn-5-en-16-one**


A-813

**(2α,3β)-form**N-Me: 2-Hydroxy-3-(methylamino)-pregn-5-en-16-one. **Kurchiphyllamine** [1433-87-0]C₂₂H₃₅NO₂ 345.524Alkaloid from the leaves of *Holarrhena antidysenterica* (Apocynaceae). Mp 161°. [α]_D -211 (c, 1.3 in CHCl₃).N,N-Di-Me: 3-(Dimethylamino)-2-hydroxypregn-5-en-16-one. **Kurchiphylline** [1433-88-1]C₂₃H₃₇NO₂ 359.551Alkaloid from the leaves of *Holarrhena antidysenterica* (Apocynaceae). Mp 184°. [α]_D -173 (c, 1 in CHCl₃).

N,N-Di-Me, O-Ac: Mp 125°.

Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1966, 1212-1216 (*Kuchiphyllamine*, *Kurchiphylline*)**3-Amino-2-hydroxypregn-5-en-20-one**

A-814



Absolute Configuration

C₂₁H₃₃NO₂ 331.497**(2α,3β)-form**N,N-Di-Me: 3-(Dimethylamino)-2-hydroxypregn-5-en-20-one. **Kurchiline** C₂₃H₃₇NO₂ 359.551Alkaloid from the leaves of *Holarrhena antidysenterica* (Apocynaceae). Mp 219°. [α]_D +46 (c, 1 in CHCl₃).


N,N-Di-Me, O-Ac: Mp 139°.

Ghosh, S. et al., *J. Indian Chem. Soc.*, 1928, **5**, 477 (isol)Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1964, 2158 (isol, ir, pmr, ms, struct)**2-Amino-3-hydroxy-1-propanesulfonic acid**

A-815

Cysteinolic acid

[3687-17-0]

**(R)-form**C₃H₉NO₄S 155.174

Stereochem. and registry nos. are confused. Wickberg states that the L-form was isol., but his exptl. work shows clearly that it was D (i.e. S as it is related to D-Cysteine). CAS uses the same registry no. for L- (in 8CI) and (S-) (in 10CI).

(R)-form

L-form

Isol. from red alga *Polysiphonia fastigiata*. Mp 279-281° dec. [α]_D -6 (c, 2.0 in H₂O).**(S)-form**


D-form

Isol. from brown (e.g. *Hijikia fusiforme*) and green (*Ulva pertusa*, *Enteromorpha linza*) algae. Also from diatoms, e.g. *Navicula pelliculosa* and from the starfish *Asterina pectinifera*. Mp 279-282° dec. [α]_D +7 (H₂O).

[15509-62-3, 16421-58-2, 56942-41-7]

Wickberg, B. et al., *Acta Chem. Scand.*, 1957, **11**, 506 (isol)Yoneda, T. et al., *CA*, 1967, **66**, 113388p (isol) Higashiura, K. et al., *J.O.C.*, 1992, **57**, 764 (synth)**2-Amino-4-hydroxypteridine**

A-816

2-Amino-4(1H)-pteridinone, 9CI. 2-Amino-4-pteridinol. **Pterin**. **Pteridoxamine** [2236-60-4]C₆H₅N₅O 163.138

4-One-form predominates. Isol. from numerous natural sources. Yellow cryst. (HCOOH aq.). Fluoresces: violet in neutral/alkaline soln.; yellow-green in acid soln.

▶ UO3505000

8-Oxide: [42346-89-4]

C₆H₅N₅O₂ 179.138

Bright-yellow cryst. Mp 360°.

2-N-Ac:

C₈H₇N₅O₂ 205.176

Cryst. (propanol). Mp 350°. Browns at 270°.

2-N-Me: 2-(Methylamino)-4(1H)-pteridinone, 9CI

[13005-84-0]

C₇H₇N₅O 177.165

Pale-yellow cryst. Mp 350°.

N¹-Me: 2-Amino-1-methyl-4(1H)-pteridinone, 9CI

[13005-86-2]

C₇H₇N₅O 177.165


Mp 335-337° dec.

[938-42-1]

Dick, G.P.G. et al., *J.C.S.*, 1955, 1379 (synth) Pfeleiderer, W. et al., *Chem. Ber.*, 1960, **93**, 2015 (synth, struct)Brown, D.J. et al., *J.C.S.*, 1961, 4413 (tautom) Viscontini, M. et al., *Helv. Chim. Acta*, 1963, **46**, 1181 (pmr)McCormack, J.J. et al., *J.O.C.*, 1964, **29**, 3370 (synth)Müller, G. et al., *Helv. Chim. Acta*, 1973, **56**, 2680 (cmr)Williams, V.P. et al., *J. Het. Chem.*, 1973, **10**, 827 (ms)Albert, A. et al., *J.C.S. Perkin 2*, 1973, 1101 (uv)Taylor, E.C. et al., *J.O.C.*, 1975, **40**, 2341 (oxide)Tobias, S. et al., *Chem. Ber.*, 1985, **118**, 354 (cmr)

2-Amino-4-hydroxy-6-pteridinecarboxylic acid A-817

2-Amino-1,4-dihydro-4-oxo-6-pteridinecarboxylic acid. 2-Amino-4(3H)-pteridinone-6-carboxylic acid. Pterin-6-carboxylic acid. Ranachrome 5 [948-60-7]



C₇H₅N₅O₃ 207.148

Isol. from various biol. sources incl. human blood, sheep pineals, various amphibians, fish and soybeans. Also from the Chinese drug Jin Yu. Metab. of Pteroylglutamic acid, P-772. Cream cryst. Mp 360°.

Me ester: [31010-61-4]

C₈H₇N₅O₃ 221.175

Cryst. (DMF). Mp 285° dec.

2-N-Ac: [31010-65-8]

C₉H₇N₅O₄ 249.185

Cryst. (MeOH). Mp 210°.

2-N-Ac, Me ester: [31010-67-0]

C₁₀H₉N₅O₄ 263.212

Mp 283°.

7,8-Dihydro: 7,8-Dihydropterin-6-carboxylic acid

C₇H₈N₅O₃ 210.172

Light emitter of the millipede *Luminesmus sequoiae*.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 896C (ir)

Baugh, C.M. et al., *J.O.C.*, 1964, 29, 3610 (synth)

Pfleiderer, W. et al., *Annalen*, 1970, 741, 64 (synth, derivs)

Iwanami, Y. et al., *Tet. Lett.*, 1972, 3219 (ms)

Nair, M.G. et al., *J.O.C.*, 1973, 38, 2185 (synth)

Mengel, R. et al., *Chem. Ber.*, 1978, 111, 3790 (synth)

Kohashi, M. et al., *J. Biochem. (Tokyo)*, 1980, 87, 1581 (isol)

Zeitler, M. et al., *Methods Enzymol.*, 1986, 122, 273 (isol)


Sato, N. et al., *J. Het. Chem.*, 1988, 25, 1737 (synth, derivs)

Suga, T. et al., *J. Nat. Prod.*, 1988, 51, 713 (isol)

Kuse, M. et al., *Bioorg. Med. Chem. Lett.*, 2001, 11, 1037-1040 (7,8-dihydro)

6-Amino-2-hydroxypurine A-818

6-Amino-1,3-dihydro-2H-purin-2-one, 9CI. Isoguanine. Guanopterin [3373-53-3]



C₅H₅N₅O 151.127

Several tautomers possible. Aglycone from *Croton tiglium*. Isol. from the wings of the butterfly *Prionis thestylis*. Amorph. powder. Mp 360°. pK_{a1} 4.47; pK_{a2} 9.03 (20°).

9-(β-D-Arabinofuranosyl): [38819-11-3]

C₁₀H₁₃N₅O₅ 283.243

Light yellow powder. Mp 269-272° dec. [α]_D²⁴ +29.5 (c, 0.5 in H₂O).

9-β-D-Ribofuranosyl: see Isoguanosine

1-Me: 6-Amino-1,3-dihydro-1-methyl-2H-purin-2-one, 9CI

[73691-67-5]

C₆H₇N₅O 165.154

Powder. Mp 300°.

1-Me, 9-β-D-arabinofuranosyl: Ara-doridose

[77856-33-8]

C₁₁H₁₅N₅O₅ 297.27

Shows antiviral props. Plates (H₂O).

Mp 223-225° dec. [α]_D²⁴ +25.5 (c, 0.5 in H₂O).

1-Me, 6-N-Ac: [98933-71-2]

C₈H₉N₅O₂ 207.191

Cryst. (MeOH). Mp 222-223° Mp 274° dec.

1-Me, 6,9-di-Ac: [98933-69-8]

C₁₀H₁₁N₅O₃ 249.229

Cryst. Mp 222-223° dec. Exists as 6-imino tautomer.

6-N-Me: 6-(Methylamino)-1,3-dihydro-2H-purin-2-one, 9CI. 2-Hydroxy-6-methylamino-9H-purine

[24391-35-3]

C₆H₇N₅O 165.154

Isol. from the blue coral NIO-156 and from the bacterium *Pseudomonas syringae*. Cytokinin. Sol. H₂O. λ_{max} 265; 282 (MeOH) (Berdy).

1,3-Di-Me: 1,3-Dimethylisoguanine. 1,3-Dimethylisoguaninium

[191614-40-1]

C₇H₉N₅O 179.181

Isol. from the sponges *Amphimedon viridis*, *Amphimedon paraviridis* and *Xestospongia exigua*. Also from *Cnemidocarpa bicornuta*. Powder; cryst. (as salt). Isol. by different groups both as the free base and as salts protonated on the imino N. λ_{max} 207 (log ε 4.5); 295 (log ε 3.9) (MeOH).

3,7-Di-Me: 3,7-Dimethylisoguanine

[18904-09-1]

C₇H₉N₅O 179.181

Isol. from the Caribbean sponge *Agelas longissima* and *Zyzzya fuliginosa*. Sol. MeOH, butanol. λ_{max} 245 (ε 8500); 285 (ε 9900) (MeOH) (Berdy).

N³,O-Di-Me: Mucronatine†

C₇H₉N₅O 179.181

Isol. from the sponge *Stryphnus mucronatus*. Solid. Mp 200-202°. Tautomeric.

7,8-Dihydro, 6-N-(3-methylbutyl): 7,8-Dihydro-2-hydroxy-N⁶-(3-methylbutyl)adenine. 7,8-Dihydro-2-hydroxy-6-(3-methylbutylamino)purine

[188542-24-7]

C₁₀H₁₇N₅O 223.277

Constit. of *Artemisia annua*.

Spies, J.R. et al., *J.A.C.S.*, 1939, 61, 350-351 (*Isoguanine, isol*)

Purrmann, R. et al., *Annalen*, 1940, 544, 182-190 (*Isoguanine, isol*)

Taylor, E.C. et al., *J.A.C.S.*, 1959, 81, 2442-2448 (*Isoguanine, synth, uv*)

Veillard, A. et al., *J. Chim. Phys.*, 1962, 59, 1056-1066 (pmr)

Pal, B.C. et al., *J.C.S.*, 1964, 400-405 (ir)

Pettit, G.R. et al., *J. Nat. Prod.*, 1976, 39, 363-367 (*Isoguanine, isol*)

Yamazaki, A. et al., *Nucleic Acids Res.*, 1976, 3, 251-259 (*Isoguanine, synth*)

Nachman, R.J. et al., *J.C.S. Perkin 1*, 1985, 1315-1321 (*1-Me, 1-Me-6,9-di-Ac, synth, cmr, pmr, ms, uv, ir, cryst struct*)

Farooqi, A.H.A. et al., *Phytochemistry*, 1990, 29, 2061-2063 (*6-N-Me, isol*)

Cafieri, F. et al., *Tet. Lett.*, 1995, 36, 7893-7896 (*3,7-Dimethylisoguanine*)

Shukla, A. et al., *J. Indian Chem. Soc.*, 1997, 74, 59 (*Artemisia annua constit*)

Mitchell, S.S. et al., *J. Nat. Prod.*, 1997, 60, 727-728 (*1,3-Dimethylisoguanine*)

Chehade, C.C. et al., *J. Nat. Prod.*, 1997, 60, 729-731 (*1,3-Dimethylisoguanine*)

Do Prado, G.M.T. et al., *Acta Cryst. C*, 1999, 55, 1585-1587 (*trihydrate, 1,3-di-Me, cryst struct*)

Bourguet-Kondracki, M.-L. et al., *Tet. Lett.*, 2001, 42, 7257-7259 (*Mucronatine*)

Jeong, S.-J. et al., *Chem. Pharm. Bull.*, 2003, 51, 731-733 (*1,3-Dimethylisoguaninium*)


Blas, J.R. et al., *J.A.C.S.*, 2004, 126, 154-164 (*tautom*)

Panthong, K. et al., *Acta Cryst. C*, 2006, 62, o193-o195 (*1,3-Dimethylisoguanine, isol, cryst struct*)

Bats, J.W. et al., *Acta Cryst. E*, 2006, 62, 2040-2042 (*cryst struct*)

6-Amino-8-hydroxypurine A-819

6-Amino-1,7-dihydro-8H-purin-8-one, 9CI. 6-Amino-8-purinol, 8CI. 7,8-Dihydro-8-oxoadenine. 8-Hydroxyadenine. 8-Oxoadenine [21149-26-8]



C₅H₅N₅O 151.127

8-Oxo-form predominates. Other tautomers are possible. Results *in vivo* from hydroxylation of DNA by OH radicals. Found in the urine of humans suffering from leukaemia. Isol. from the ascidian *Sympyegma rubra* and a Thorectidae sponge. Potential cause of DNA mismatch in mutagenesis. Needles + ½H₂SO₄. Mp 300° (H₂SO₄). Unlike 2-Amino-6,8-dihydroxypurine, A-753, is not particularly mutagenic.

6-N-Me: [89073-90-5]

C₆H₇N₅O 165.154

Needles (MeOH aq.). Mp >300°. λ_{max} 273 (ε 16900) (H₂O).

9-Me: 6-Amino-7,9-dihydro-9-methyl-8H-purin-8-one, 9CI. 9-Methyl-8-oxoadenine

[119138-92-0]

C₆H₇N₅O 165.154

Isol. from a Thorectidae sponge.

Prisms (H₂O). Mp 300°.

6-N,9-Di-Me: 7,9-Dihydro-9-methyl-6-(methylamino)-8H-purin-8-one, 9CI. N⁶,9-Dimethyl-8-oxoadenine

[119138-91-9]

C₇H₉N₅O 179.181

Needles (EtOH). Mp 300°.

N⁶-(3,5-Dibromo-4-hydroxybenzyl):

Aplidiamine

[188348-59-6]

C₁₂H₉Br₂N₅O₂ 415.043

Isol. from the marine ascidian *Aplidiopsis* sp. Sol. H₂O, MeOH. Mp 241-243°. λ_{max} 221 (ε 17300); 273 (ε 15800) (MeOH/CH₂Cl₂ 1:1). λ_{max} 223 (ε 19000); 290 (ε 15200) (MeOH/CH₂Cl₂/HCl). λ_{max} 223 (ε 52500); 250 (sh); 283 (ε 30700); 313 (sh); 377 (sh) (MeOH/CH₂Cl₂/NaOH).

1H-form*l*-Me: [186391-69-5]C₆H₇N₅O 165.154

Needles +1H₂O (MeOH aq.). Mp > 300°. Various tautomers possible. λ_{max} 216 (ε 22000) (H₂O).

l,9-*Di*-Me: see 1,6,7,9-Tetrahydro-6-imino-1,9-dimethyl-8H-purin-8-one, T-194

3H-form

3-Me: 6-Amino-3,7-dihydro-3-methyl-8H-purin-8-one. 8-Hydroxy-3-methyladenine

[185201-03-0]

C₆H₇N₅O 165.154

Needles. Mp > 300°. pK_a 4.1 (30°H₂O). λ_{max} 232 (ε 18800) (95% EtOH).

3,9-*Di*-Me: 3,6,7,9-Tetrahydro-6-imino-3,9-dimethyl-8H-purin-8-one, 9CI

[198758-37-1]

[198023-18-6]

C₇H₉N₅O 179.181

Plates (HCl aq.) (as hydrochloride). Mp 274-276° dec. (hydrochloride).

Robins, R.K. *et al.*, *J.A.C.S.*, 1958, **80**, 6671-6679 (*synth*, *uv*)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 2146-2150 (*Me derivs*, *synth*, *ir*, *uv*, *pmr*)

Brown, T. *et al.*, *Chem. Br.*, 1993, **29**, 484-486 (*rev*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2025-2032; 2318-2321; 1997, **45**, 1582-1588; 1867-1869 (*synth*, *ir*, *uv*, *pmr*, *bibl*, *Me derivs*)

Kang, H. *et al.*, *Tet. Lett.*, 1997, **38**, 941-944 (*Aplidiamine*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1297-1300 (*Aplidiamine*, *synth*, *struct*)

Lindsay, B.S. *et al.*, *J. Nat. Prod.*, 1999, **62**,


1573-1575 (*Symplegma rubra constii*)

Madyastha, K.M. *et al.*, *J.C.S. Perkin 1*, 1999, 677-680 (*isol*)

Makariev, T.N. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 711-714 (*9-Methyl-8-oxoadenine*, *isol*)

4-Amino-3-hydroxy-1H-pyrazolo[3,4-*d*]pyrimidine A-820

4-Amino-1,2-dihydro-3H-pyrazolo[3,4-*d*]pyrimidin-3-one, 9CI. *Akalone* [128850-54-4]

C₅H₅N₅O 151.127

Prod. by the marine bacterium *Agrobacterium aurantiacum* and *Flavobacterium* sp. N-81106. Xanthine oxidase inhibitor.

Powder or light green needles. Sol. bases, DMSO; poorly sol. MeOH, hexane, CHCl₃. Mp > 350°. λ_{max} 269 (ε 1690) (H₂O).


Anderson, J.D. *et al.*, *J. Het. Chem.*, 1990, **27**, 439-453 (*synth*)

Japan. Pat., 1993, 93 317 070; *CA*, **120**, 189879d (*isol*)

Izumida, H. *et al.*, *J. Mar. Biotechnol.*, 1995, **2**, 115-118 (*isol*, *ir*, *cmr*, *pmr*)

2-Amino-6-hydroxy-3-pyridine-necarboxaldehyde A-821

2-Amino-3-formyl-6-hydroxypyridine

C₆H₆N₂O₂ 138.126

Me ether: 2-Amino-6-methoxy-3-pyridine-carboxaldehyde. 2-Amino-3-formyl-6-methoxypyridine


C₇H₈N₂O₂ 152.152

Alkaloid from the basidiomycete *Cortinarius umidicola*. Needles (MeOH). Mp 186-187°. λ_{max} 205 (log ε 4.32); 252 (log ε 4.38) (MeOH).

Hu, L. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 659-662 (*isol*, *pmr*, *cmr*, *ms*)

α-Amino-5-hydroxy-2-pyridine-nepropanoic acid, 9CI A-822

β-(5-Hydroxy-2-pyridyl)alanine. *Azatyrosine*. Antibiotic SF 1346. SF 1346

C₈H₁₀N₂O₃ 182.179

Amino acid antibiotic.

(S)-form [58525-82-9]

Isol. from *Streptomyces chibaensis*. Antiviral, antitumour, antibacterial and anti-neoplastic agent. Also shows antiinflammatory activity. Cryst. (H₂O). Sol. H₂O, MeOH, butanol-H₂O, bases; fairly sol. EtOH; poorly sol. Me₂CO, hexane. Mp 262-263° dec. [α]_D²⁵ -33 (c, 1.0 in H₂O). [α]_D²⁵ +55 (c, 1.1 in 1M HCl). λ_{max} 219 (E1%/1cm 330); 280 (E1%/1cm 140); 320 (E1%/1cm 15) (H₂O) (Berdy). λ_{max} 228 (E1%/1cm 175); 289 (E1%/1cm 235) (HCl) (Berdy). λ_{max} 241 (E1%/1cm 445); 302 (E1%/1cm 165) (NaOH) (Berdy).

Hydrochloride (1:3):

Hygroscopic foam + 2½H₂O. [α]_D²⁵ +28 (c, 0.06 in 1M HCl).

Me ester: [58485-16-8]

Cryst. (as hydrochloride). Mp 224-225° dec. (hydrochloride). [α]_D²⁰ +38 (c, 1.0 in H₂O).

(±)-form [943-82-8]

Prisms (H₂O). Mp 272-273° dec.

Inouye, S. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2669 (*isol*)

Harris, R.L.N. *et al.*, *Aust. J. Chem.*, 1977, **30**, 649 (*isol*)

Ger. Pat., 1978, 2 823 346; *CA*, **90**, 133025s (*synth*, *props*)

Fukuyasu, H. *et al.*, *CA*, 1984, **101**, 48223 (*props*)

Izawa, M. *et al.*, *Cancer Res.*, 1992, **52**, 1628-1630 (*pharmacol*)

Schow, S.R. *et al.*, *J.O.C.*, 1994, **59**, 6850 (*synth*)

Ye, B. *et al.*, *J.O.C.*, 1995, **60**, 2640 (*synth*, *ir*, *pmr*)


Coper, M.S. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 2613 (*synth*)

Myers, A.G. *et al.*, *J.O.C.*, 1996, **61**, 813 (*synth*, *ir*, *pmr*, *bibl*)

Monden, Y. *et al.*, *Ann. N.Y. Acad. Sci.*, 1999, **886**, 109-121 (*rev*, *activity*)

2-Amino-4-hydroxypyrimidine A-823

2-Amino-4(1H)-pyrimidinone, 9CI. 2-Amino-4-pyrimidinol. *Isocytosine* [108-53-2]

C₄H₅N₃O 111.103

NH-forms predominate. Prisms (H₂O). Mp 280° dec. pK_{a1} 3.91; pK_{a2} 9.54 (25°).

1H-form*l*-β-D-Ribopyranosyl: *Isocytidine*

[489-59-8]


C₉H₁₃N₃O₅ 243.219

Isol. from a strain of *Penicillium brevicompactum*. Glass. λ_{max} 256 nm, λ_{min} 240 nm (H₂O, pH1).

Lis, A.W. *et al.*, *Biochim. Biophys. Acta*, 1962, **61**, 250 (*Isocytidine*)

3-Amino-5-hydroxy-5-vinyl-2-cyclopenten-1-one A-824

3-Amino-5-ethenyl-5-hydroxy-2-cyclopenten-1-one, 9CI

C₇H₉NO₂ 139.154

Incorrect struct. given in CAS.

(R)-form*Myrothenone B*

[878633-73-9]

[858126-05-3]

Prod. by *Streptomyces* sp. GT-20026114 obt. from the mangrove plant *Aegiceras coniculatum* and from a marine-derived fungus *Myrothecium* sp. Yellowish oil. [α]_D²⁰ +127.1 (c, 1.54 in MeOH). [α]_D²⁰ +35 (c, 0.6 in MeOH). λ_{max} 270 (MeOH). λ_{max} 203 (log ε 3.7); 268 (log ε 4) (MeOH).

N-Formyl: Myrothenone AC₈H₉NO₃ 167.164

Prod. by *Myrothecium* sp. Tyrosinase inhibitor. Oil. [α]_D²⁰ +61 (c, 0.6 in MeOH). λ_{max} 203 (log ε 3.7); 272 (log ε 4.2) (MeOH).

N-Methoxycarbonyl: [878633-74-0]
C₉H₁₁NO₄ 197.19
Prod. by *Streptomyces* sp. GT-20026114. Yellowish oil. [α]_D²⁰ +67.6 (c, 1.16 in MeOH). λ_{max} 261 (MeOH).

N,N-Di-Me: 3-(Dimethylamino)-5-hydroxy-5-vinyl-2-cyclopenten-1-one. 3-(Dimethylamino)-5-ethenyl-5-hydroxy-2-cyclopenten-1-one
[179422-20-9]

Isol. from *Trichoderma koningii*. Amorph. powder. Sol. H₂O, MeOH, CHCl₃, DMSO; fairly sol. Me₂CO, MeCN, EtOAc; poorly sol. hexane. Mp 137-139°. [α]_D²⁰ +69.3 (c, 0.2 in H₂O). Biol. inactive. λ_{max} 280 (ε 30000) (MeOH).

N-[2-(4-Hydroxyphenyl)ethyl]: [878633-75-1]

C₁₅H₁₇NO₃ 259.304
Prod. by *Streptomyces* sp. GT-20026114. Yellowish oil. [α]_D²⁰ +59.3 (c, 1.73 in MeOH). λ_{max} 232 (MeOH).

2-Bromo-3-Amino-2-bromo-5-hydroxy-5-vinyl-2-cyclopenten-1-one. 2-Bromomyrothenone **B**

[929276-07-3]
C₇H₈BrNO₂ 218.05
Prod. by a marine-derived *Botrytis* sp. Red oil. [α]_D²⁰ +61 (c, 0.9 in MeOH). λ_{max} 210 (log ε 3.7); 277 (log ε 4.2) (MeOH).

Mukhopadhyay, T. *et al.*, *J. Antibiot.*, 1996, **49**, 210-211 (*N,N*-di-Me)


Li, X. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 453-455 (*Myrothenones A,B*)

Lin, W. *et al.*, *J. Antibiot.*, 2005, **58**, 594-598 (*isol, cd, pmr, cmr, ms*)

Li, X. *et al.*, *J. Nat. Prod.*, 2007, **70**, 307-309 (*2-Bromomyrothenone B*)

2-Aminoimidazole, 8CI A-825

1*H*-Imidazol-2-amine, 9CI. 2-Amino-glyoxaline
[7720-39-0]



C₃H₅N₃ 83.093

Isol. from the sponges *Halichondria* sp. and *Reniera cratera*. Also occurs in the seeds of the legume *Mundulea sericea*, and has been shown to be a precursor of the antibiotic 2-Nitroimidazole, N-250 in *Streptomyces eurocidicus*.

Hydrochloride: [57575-96-9]

Mp 152°. Hygroscopic.

Nitrate: Mp 135-136°.

Sulfate (2:1): [1450-93-7]
[42383-61-9]

Mp 270° dec.

Picrate: Mp 236°.

2-*N*-Ac: 2-Acetamidoimidazole

[52737-49-2]
C₅H₇N₃O 125.13
Mp 285° (270-275°) dec.

1-*Me*: [6646-51-1]

C₄H₇N₃ 97.119

Prismatic needles (EtOH/EtOAc) (as

hydrochloride). Mp 84° (hydrochloride).

[1450-94-8]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 614D (*ir*)

Lawson, A. *et al.*, *J.C.S.*, 1956, 307 (*synth*)
Seki, Y. *et al.*, *J. Biochem. (Tokyo)*, 1970, **67**, 389 (*isol, ir, pmr*)

Cimino, G. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1974, **47**, 895 (*isol, uv, pmr, ms*)

Fellows, L.E. *et al.*, *Phytochemistry*, 1977, **16**, 1399 (*isol, pmr, ms*)

Kirk, K.L. *et al.*, *J.O.C.*, 1978, **43**, 4381-4383 (*synth, pmr*)

Utkina, N.K. *et al.*, *Khim. Prir. Soedin.*, 1984, 124-125; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 126-127 (*isol, Halichondria*)

Weinmann, H. *et al.*, *Tet. Lett.*, 2002, **43**, 593-595 (*synth, ir, pmr*)

Yano, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **12**, 3443-3450 (*1-Me, synth, pmr*)

4(5)-Aminoimidazole A-826

1*H*-Imidazol-4-amine. 4-Aminoglyoxaline
[4919-03-3]

C₃H₅N₃ 83.093

Present in biotin-restricted yeast. Mp 184° (as hydrochloride).

*N*⁴-Ac: [51741-72-1]

C₅H₇N₃O 125.13

Cryst. (dioxan). Sol. H₂O. Mp 226°.

*N*⁴-Ac, picrate: Mp 208°.

1*H*-form

1-Benzyl: 4-Amino-1-benzyl-1*H*-imidazole
[53594-77-7]

C₁₀H₁₁N₃ 173.217

Cryst. (MeCN) (as hydrochloride). Mp 179-182° (hydrochloride). CAS no. refers to hydrochloride.

1-Benzyl, *N*⁴-Ac: [7524-03-0]

C₁₂H₁₃N₃O 215.254

Cryst. (EtOAc). Mp 168-168.8°.

3*H*-form

3-*Me*: 5-Amino-1-methyl-1*H*-imidazole

[66787-75-5]

C₄H₇N₃ 97.119

Purple solid. Mp 107-109°.

Hunter, G. *et al.*, *Can. J. Res., Sect. B*, 1941, **19**, 296; *CA*, **36**, 1321 (*synth, N-Ac*)

Whitaker, J.M. *et al.*, *Arch. Biochem. Biophys.*, 1962, **96**, 541 (*isol*)

Coburn, M.D. *et al.*, *J. Het. Chem.*, 1970, **7**, 1391-1393 (*synth*)


Buchman, R. *et al.*, *J. Med. Chem.*, 1974, **17**, 1168-1171 (*N-benzyl derivs*)

Al-Shaar, A.H.M. *et al.*, *J.C.S. Perkin 1*, 1992, 2779 (*3-Me*)

5-Aminoimidazole riboside A-827

1-β-*D*-Ribofuranosyl-1*H*-imidazol-5-amine, 9CI

[30597-39-8]



C₈H₁₃N₃O₄ 215.208

Prod. by *Escherichia coli* strain

K-12. Hygroscopic pale grey solid.

Mp 92-94°. λ_{max} 300 (no solvent reported).

5'-Phosphate: 5-Aminoimidazole ribonucleotide. AIR

[25635-88-5]

C₈H₁₄N₃O₇P 295.188

Intermed. in biosynth. of purine ribonucleotides. Precursor of thiamine in *Escherichia coli*. Solid.

Love, S.H. *et al.*, *Biochim. Biophys. Acta*, 1959, **35**, 367 (*isol*)


Schrimsher, J.L. *et al.*, *Biochemistry*, 1986, **25**, 4356-4365 (*synth, pmr, cmr*)

Bhat, B. *et al.*, *J.A.C.S.*, 1990, **112**, 4891-4897 (*synth, pmr*)

Firestone, S.M. *et al.*, *Biochemistry*, 1994, **33**, 11917-11926 (*synth, ir, pmr, cmr*)

Humphries, M.J. *et al.*, *Synthesis*, 1999, 985-992 (*synth, pmr*)

N-[2-(2-Amino-1*H*-imidazol-4-yl)ethyl]-3-(3,5-dibromo-4-methoxyphenyl)-2-propenamide, 9CI A-828



C₁₅H₁₆Br₂N₄O₂ 444.125


(*E*)-form [160491-72-5]

Metab. from the Caribbean sponge *Verongula* sp. Amorph. solid (as trifluoroacetate). CAS no. refers to trifluoroacetate.

Ciminiello, P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1564 (*isol, uv, ir, pmr, cmr, struct*)

2-Amino-1*H*-imidazo[4,5-*b*]pyrazine A-829

1*H*-Imidazo[4,5-*b*]pyrazin-2-amine. Zarzissine



C₅H₅N₅ 135.128

Struct. of Zarzissine revised in 2001.

Alkaloid from the Mediterranean

sponge *Anchinoe paupertas*. Cytotoxic.

Active against *Candida*, bacteria and

some tumour lines. Cryst. (H₂O or

MeOH). Mp > 300° dec. λ_{max} 217 (sh);

244 (log ε 3.3); 318 (log ε 4.03)

(MeOH). λ_{max} 206 (log ε 4.31); 230 (sh)

; 282 (sh); 326 (log ε 5.39) (MeOH/


NaOH).

Bouaicha, N. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1455 (*isol*)

Wan, Z.K. *et al.*, *Tetrahedron*, 2001, **57**, 5497-5507 (*synth, uv, pmr, cmr*)

2-Amino-3-(3-indolyl)-1-propanol A-830

β-Amino-1H-indole-3-propanol, 9CI.
Tryptophanol
[526-53-4]



C₁₁H₁₄N₂O 190.244

▶ NM1320000

(R)-form

D-form

[52485-52-6]

Cryst. (EtOAc). Mp 80-81°. [α]_D²⁵ +20.9 (c, 1 in MeOH).

Oxalate salt (1:1): [58889-66-0]

Mp 202-203°. [α]_D²⁰ +22.1 (c, 1.02 in H₂O).

(S)-form

L-form

[2899-29-8]

Cryst. (EtOAc). Mp 84-85°. [α]_D²⁵ -20.5 (c, 1 in MeOH).

Oxalate salt (1:1): [2899-30-1]

Mp 203-205°. [α]_D²⁰ -22.4 (c, 1.03 in H₂O).

N-Ac: *N*-Acetyltryptophan

C₁₃H₁₆N₂O₂ 232.282

Prod. by *Streptomyces pleomorphus* (strain YIM33176). Amorph. powder. [α]_D²⁵ +27 (c, 1.2 in CHCl₃). λ_{max} 221 (log ε 4.49); 281 (log ε 3.77) (no solvent reported).

(±)-form [154-09-6]

Cryst. (C₆H₆). Mp 85-86°.

Hydrochloride: [25825-57-4]

Cryst. (EtOH/Et₂O). Mp 178-179°.

Hellmann, H. *et al.*, *Chem. Ber.*, 1958, **91**, 2290 (*synth*)

Ito, A. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 3106 (*synth*)


Repke, D.B. *et al.*, *J. Het. Chem.*, 1976, **13**, 775 (*synth*, *pmr*)

Hvidt, T. *et al.*, *Can. J. Chem.*, 1988, **66**, 779 (*synth*, *pmr*, *cmr*)

Li, Y.Q. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 613-615 (*N*-Acetyltryptophanol)

7-Amino-5,8-isoquinoline-dione A-831

7-Amino-5,8-isoquinolinequinone



C₉H₆N₂O₂ 174.159

*N*⁷-*Me*: 7-(Methylamino)-5,8-isoquinolinedione, 9CI. *Caulibugulone A* [662167-15-9]

C₁₀H₈N₂O₂ 188.185

Alkaloid from the marine bryozoan *Caulibugula intermis*. Cytotoxic. Dark

red solid. λ_{max} 237 (log ε 4.24); 268 (log ε 4.1); 451 (log ε 3.61) (MeOH).

*N*⁷-(2-Hydroxyethyl): 7-[(2-Hydroxyethyl)amino]-5,8-isoquinolinedione.

Caulibugulone D

[662167-18-2]

C₁₁H₁₀N₂O₃ 218.212

Alkaloid from *Caulibugula intermis*.

Cytotoxic. Dark orange solid. λ_{max} 273 (log ε 3.95); 450 (log ε 3.35) (MeOH).

*N*⁷-*Me*, 5-imine: *Caulibugulone E*

[662167-19-3]

C₁₀H₉N₃O 187.201

Alkaloid from *Caulibugula intermis*.

Cytotoxic. Dark orange solid (as TFA salt). λ_{max} 213 (log ε 3.71); 245 (log ε 3.75); 441 (log ε 3.09) (MeOH) (TFA salt).

*N*⁷-*Me*, 5-(2-hydroxyethyl)imide: *Caulibugulone F*

[662167-20-6]

C₁₂H₁₃N₃O₂ 231.254

Alkaloid from *Caulibugula intermis*.

Cytotoxic. Dark orange solid (as TFA salt). λ_{max} 215 (log ε 3.79); 245 (log ε 3.75); 440 (log ε 3.16) (MeOH) (TFA salt).

6-Chloro, *N*⁷-*Me*: 6-Chloro-7-(methylamino)-5,8-isoquinolinedione. *Caulibugulone B*

[662167-17-1]

C₁₀H₇ClN₂O₂ 222.63

Alkaloid from *Caulibugula intermis*.

Cytotoxic. Dark red solid. λ_{max} 251 (log ε 4.03); 274 (log ε 3.97); 475 (log ε 3.36) (MeOH).

6-Bromo, *N*⁷-*Me*: 6-Bromo-7-(methylamino)-5,8-isoquinolinedione. *Caulibugulone C*

[662167-16-0]

C₁₀H₇BrN₂O₂ 267.082

Alkaloid from *Caulibugula intermis*.

Cytotoxic. Dark red solid. λ_{max} 254 (log ε 4); 272 (log ε 3.97); 474 (log ε 3.49) (MeOH).

Milanowski, D.J. *et al.*, *J. Nat. Prod.*, 2004, **67**, 70-73 (*Caulibugulones*, *isol*, *pmr*, *cmr*)


Wipf, P. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 2173-2174 (*synth*, *activity*)

Alagille, D. *et al.*, *Tet. Lett.*, 2004, **45**, 6179-6181 (*synth*)

4-Amino-3-isoxazolidinone, A-832

9CI, 8CI

Cycloserine, *BAN*, *INN*, *USAN*. *Cyclo-mycin*. *Closina*. *Micoserina*. *Farmiserina*. *Orientomycin*. *Oxymycin*. *Antibiotic 106-7*. *Antibiotic 5915*. *Antibiotic 8217*. *Antibiotic 17452*. *Antibiotic E 733A*. *Antibiotic I 1431*. *Antibiotic K 300*. *Antibiotic NJ 21*. *Antibiotic PA 94*. *Antibiotic Ro 1-9213*. *Oxamycin*. *Seromycin*. *Orientmycin*. *Many other names*
[4834-58-6]



C₃H₆N₂O₂ 102.093

λ_{max} 226 (ε 4140) (H₂O) (Derep).

(R)-form [68-41-7]

Prod. by *Streptomyces garyphalus*, *Streptomyces orchidaceus*, *Streptomyces lavendulae* and *Streptomyces nagasakiensis*. Shows antibiotic activity primarily against mycobacteria. Tuberculostatic. Cryst. Sol. H₂O, alkalis. Mp 155-156° dec. [α]_D²³ +116 (c, 1.17 in H₂O). Of limited clinical use due to toxicity.

▶ CNS adverse effects reported when used therapeutically. LD₅₀ (mus, orl) 5290 mg/kg. NY2975000

N-Ac:

C₅H₈N₂O₃ 144.13

Needles. Mp 179-180°.

N,N'-*Di*-Ac:

C₇H₁₀N₂O₄ 186.167

Chunky needles (Me₂CO). Mp 121-122°.

(S)-form

Levycloserine, *INN*, *USAN*

[339-72-0]

Enzyme inhibitor used in the treatment of Gaucher's disease. Mp 146° dec. [α]_D²⁰ -104 (c, 1 in H₂O).

▶ NY2976000

(±)-form [68-39-3]

Mp 137-140°.

▶ NY2974900

[339-72-0]

Aldrich Library of NMR Spectra, 2nd edn., 1983, **1**, 678B; 678C; 678D (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 810A; 810D; 811A (*ir*)

Harned, R.L. *et al.*, *Antibiot. Chemother. (Washington, D.C.)*, 1955, **5**, 204 (*struct*)

Stammer, C.H. *et al.*, *J.A.C.S.*, 1955, **77**, 2344; 2345; 2346; 1957, **79**, 3236 (*struct*, *synth*, *isol*, *resoln*)

Neuhaus, F.C. *et al.*, *Antibiotics (N.Y.)*, 1967, **1**, 40 (*rev*)

Milne, G.W.A. *et al.*, *Tetrahedron*, 1967, **23**, 65 (*nmr*, *ms*)

Lamb, J.W. *et al.*, *Anal. Profiles Drug Subst.*, 1972, **1**, 53 (*rev*, *synth*, *anal*)

O'Brien, P. *et al.*, *Met. Ions Biol. Syst.*, 1985, **19**, 295 (*rev*, *pharmacol*)


Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., *Akademie-Verlag*, 1987, 107 (*synonyms*)

El-Obeid, H.A. *et al.*, *Anal. Profiles Drug Subst.*, 1989, **18**, 567 (*rev*)

Martindale, The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 156

Hofmann, S.G. *et al.*, *CNS Drug Rev.*, 2006, **12**, 208-217 (*rev*, *pharmacol*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., *Van Nostrand Reinhold*, 1992, CQH000

8-Amino-4-methylaconitane-1,3,6,13,14,16,18-heptol A-833

C₁₉H₃₀N₂O₇ 398.455

(1 α ,3 α ,5 β ,6 α ,14 α ,16 β)-form


O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*: **Hemsleyatine**
[618456-73-8]
C₂₅H₄₂N₂O₇ 482.616
Alkaloid from the roots of *Aconitum hemsleyanum*. Amorph. powder. Mp 89-90°. [α]_D +36.5 (c, 0.55 in CHCl₃).
Zhou, X.L. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 592-594 (*Hemsleyatine*)

8-Amino-4-methylaconitane-1,3,6,13,14,15,16,18-octol A-834

C₁₉H₃₀N₂O₈ 414.455

(1 α ,3 α ,5 β ,6 α ,14 α ,15 α ,16 β)-form


O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*: **Lasianine**
[799782-01-7]
C₂₅H₄₂N₂O₈ 498.615
Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrum*. Needles. Mp 134-136°. [α]_D +12.9 (c, 0.4 in MeOH).
Ji, H. *et al.*, *Heterocycles*, 2004, **63**, 2363-2370 (*isol*, *pmr*, *cmr*)

1-Amino-3-methyl-2,3-butanediol A-835

C₅H₁₃NO₂ 119.163

(S)-form

N-*Ac*: N-(2,3-Dihydroxy-3-methylbutyl)acetamide, **9CI**
[81892-89-9]
C₇H₁₅NO₃ 161.2
Isol. from *Verbesina encelioides*. [α]_D²⁵ -20.7 (c, 0.15 in CHCl₃).
Eichholzer, J.V. *et al.*, *Phytochemistry*, 1982, **21**, 97 (*isol*, *struct*, *synth*)
Eichholzer, J.V. *et al.*, *Aust. J. Chem.*, 1986, **39**, 1907 (*synth*, *abs config*, *pmr*, *cmr*)


3-(Aminomethylene)-1,3-dihydro-2H-indole-2-thione A-836

C₉H₈N₂S 176.242
Intermed. in the detoxification of the phytoalexins Cyclobrassinin, C-842 and Brassinin, B-284 by the fungus *Leptosphaeria maculans*/Phoma lingam. Mp 197-200°.

Pedras, M.S.C. *et al.*, *Chem. Comm.*, 1998, 1565-1566 (*synth*, *pmr*, *cmr*)
Pedras, M.S.C. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 1196-1202 (*isol*, *synth*, *pmr*, *cmr*)
Pedras, M.S.C. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 2002-2007 (*isol*)

5-(Aminomethyl)-3(2H)-isoxazolone, 9CI A-837

Pantherine. Agarin. Muscimol. Pyroibotenic acid. β -Toxin
[2763-96-4]




C₄H₆N₂O₂ 114.104

Exists as betaine in solid phase. Constit. of fly agaric (*Amanita muscaria*). GA-BA_A agonist. Used in neurochemical research. Insecticide against flies. Cryst. (EtOH). Sol. H₂O. Mp 174-176° dec. λ _{max} 212 (ε 9700) (H₂O) (Berdy). λ _{max} 220 (ε 7500) (pH 12 buffer) (Berdy). λ _{max} 212 (ε 8700) (pH 2 buffer) (Berdy).
► LD₅₀ (rat, orl) 45 mg/kg. NY3325000
Nakamura, N. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 46 (*synth*)
Brehm, C. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 1298 (*cryst struct*)
Krogsgaard-Larsen, P. *et al.*, *Acta Chem. Scand., Ser. B*, 1976, **30**, 281; 1981, **35**, 311 (*synth*, *pharmacol*)
McCarry, B.E. *et al.*, *Tet. Lett.*, 1981, 5153 (*synth*)
Konda, Y. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 1083 (*ir*, *pmr*, *cmr*, *struct*, *synth*, *bibl*)
Chiarrino, D. *et al.*, *Tet. Lett.*, 1986, **27**, 3181 (*synth*)
Pevarello, P. *et al.*, *Synth. Commun.*, 1992, **22**, 1939 (*synth*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AKT750

2-Amino-4-methyl-1-pentanol, 9CI A-838

2-Amino-1-hydroxy-4-methylpentane.
Leucinol
[502-32-9]



(R)-form

C₆H₁₅NO 117.191

(R)-form

D-form
[53448-09-2]
Bp₁₁ 98-99°.
Oxalate: Mp 216°. [α]_D²⁰ -7.

(S)-form
L-form

[7533-40-6]
[14438-11-0] Bp₁₁ 98-99°. [α]_D¹⁷ +4.15 (EtOH).

N-*tert*-Butyloxycarbonyl: [82010-31-9]
C₁₁H₂₃NO₃ 217.308
Oil. Bp₅ 141°. [α]_D²⁰ -28.36 (c, 0.95 in CHCl₃).

N-*Benzyl*: [10249-88-4]
C₁₃H₂₁NO 207.315
Cryst. Mp 70°. [α]_D²⁰ +31 (c, 1.0 in CHCl₃).

(±)-**form** [16369-17-8] Sol. H₂O, EtOH, spar. sol. Et₂O. Bp 198-200°.
Hydrochloride: Mp 161-162°.

(ξ)-form

N-*Ac*: N-[1-(Hydroxymethyl)-3-methylbutyl]acetamide, **9CI**. N-[2-(1-Hydroxy-4-methylpentyl)]acetamide
[150206-13-6]
C₈H₁₇NO₂ 159.228
Prod. by *Streptomyces globisporus* and *Streptomyces griseus*.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 337A (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 541C (*nmr*)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 426D; 427A (*ir*)
Karrer, W. *et al.*, *Helv. Chim. Acta*, 1921, **4**, 90; 1948, **31**, 1617 (*synth*, *abs config*)
Enz, W. *et al.*, *Helv. Chim. Acta*, 1946, **29**, 1048 (*synth*, *use*)
Adkins, H. *et al.*, *J.A.C.S.*, 1947, **69**, 3039 (*synth*)
Dillon, J. *et al.*, *J.A.C.S.*, 1975, **97**, 5409; 5825 (*config*)
Jizba, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1993, **58**, 1452-1456 (N-*Ac*, *isol*, *pmr*, *cmr*, *ms*)
Ibuka, T. *et al.*, *J.O.C.*, 1997, **62**, 999-1015 (N-*tert*-butoxycarbonyl, *synth*, *ir*, *pmr*)
Braüner-Osborne, H. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 3926-3936 (N-*benzyl*)

4-Amino-4-methyl-2-pentanone, 9CI A-839

Diacetonamine
[625-04-7]
(H₃C)₂C(NH₂)CH₂COCH₃

C₆H₁₃NO 115.175
Isol. from *Genista hystrix* subspecies. Also isol. as a salt from *Tennopleurus hardwickii*. Mod. sol. H₂O. Bp 178-181° Bp_{0.2} 25°. G. isolates stated to be prob. as artifacts.

Oxime: [69151-99-1]
C₆H₁₄N₂O 130.189
Needles. Mp 58°. Bp₁₄ 130°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 460A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 659B (*nmr*)
Haeseler, P.R. *et al.*, *J.A.C.S.*, 1925, **47**, 1195 (*synth*)

Org. Synth., Coll. Vol., **1**, 1932, 191 (*synth*)
Smith, M.E. *et al.*, *J.A.C.S.*, 1938, **60**, 408 (*synth*)

U.S. Pat., 1950, 2 497 548; *CA*, **44**, 4494 (*synth*)

Steinegger, E. *et al.*, *Pharm. Acta Helv.*, 1976, **51**, 172; 203 (*isol*, *Genista*)
Chen, L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1787-1789 (*isol*, *Tennopleurus*)

1-Amino-2-methyl-2-propanol, 9CI A-840

Amino-tert-butyl alcohol. 2-Hydroxyisobutylamine. Aminotrimethylcarbinol
[2854-16-2]
(H₃C)₂C(OH)CH₂NH₂
C₄H₁₁NO 89.137
Component of phospholipids of *Neurospora crassa*. d₂₀²⁰ 0.93. Bp 151° Bp₂₀ 81°. n_D¹⁸ 1.4460.

▶ UA6125000

Hydrochloride: [30533-50-7]
Hygroscopic cryst. Mp 62-65°.

N-Benzoyl: [33561-46-5]
C₁₁H₁₅NO₂ 193.245
Mp 108°.

N,N-Di-Me: 1-(Dimethylamino)-2-methyl-2-propanol
[14123-48-9]
C₆H₁₅NO 117.191
Aggregation pheromone of cockroach *Blattella germanica*. Liq. Bp₄₈ 60°.

N-(2,4-Dinitrophenyl):
Yellow-orange cryst. (C₆H₆/petrol).
Mp 107-108°.

[119330-21-1]

Krassusky, K. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1908, **146**, 238 (*synth*)

Cairn, T.L. *et al.*, *J.A.C.S.*, 1941, **63**, 1034 (*synth*)

Vanderwerf, C.A. *et al.*, *J.A.C.S.*, 1954, **76**, 1231 (*synth*)

Ellman, G.L. *et al.*, *J.A.C.S.*, 1954, **76**, 4028 (*isol*)

Pfleiderer, W. *et al.*, *Chem. Ber.*, 1966, **99**, 3008 (*synth*)

Magnet, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1978, 539; 550 (*synth*)

Sakuma, M. *et al.*, *Appl. Entomol. Zool.*, 1990, **25**, 355 (*N-di-Me, isol*)

Rai, B.L. *et al.*, *J. Med. Chem.*, 1998, **41**, 3347-3359 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ALB250

2-(Aminomethyl)-2-propenoic acid, 9CI A-841

3-Amino-2-methylenepropanoic acid. α-Methylene-β-alanine
[4452-16-8]

H₂C=C(COOH)CH₂NH₂C₄H₇NO₂ 101.105

Toxic amino acid from the sponge *Fasciospongia cavernosa* and other sponges. Also prod. by a *Streptomyces* sp. Shows herbicidal props. Sol. H₂O.

Hydrochloride: [75509-28-3]
Cryst. (MeOH/Me₂CO). Mp 153-154°.

Me ester: [87375-90-4]
C₅H₉NO₂ 115.132
From *Fasciospongia cavernosa*. Mp 107-108° (as hydrochloride).

N-Tetradecanoyl: [70290-27-6]
C₁₈H₃₃NO₃ 311.464
Isol. from *Spongia* cf. *zimocca*.

N-Pentadecanoyl: [70290-28-7]
C₁₉H₃₅NO₃ 325.49
Isol. from *Spongia* cf. *zimocca*.

N-Hexadecanoyl: [70290-29-8]
C₂₀H₃₇NO₃ 339.517
Isol. from *Spongia* cf. *zimocca*.

N-Tetradecanoyl, Me ester: [70290-25-4]
C₁₉H₃₅NO₃ 325.49
Isol. from *Fasciospongia cavernosa* and *Spongia* cf. *zimocca*.

N-Pentadecanoyl, Me ester: [70290-26-5]
C₂₀H₃₇NO₃ 339.517
Isol. from *Fasciospongia cavernosa* and *Spongia* cf. *zimocca*.

N-Hexadecanoyl, Me ester: [52634-29-4]
C₂₁H₃₉NO₃ 353.544
Isol. from *Fasciospongia cavernosa*, a *Hippospongia* sp. and *Spongia* cf. *zimocca*.

N-(14-Methylpentadecanoyl), Me ester: Hurghamide A
C₂₁H₃₉NO₃ 353.544
Isol. from a *Hippospongia* sp. Wax.

N-(15-Methylhexadecanoyl), Me ester: Hurghamide B
C₂₂H₄₁NO₃ 367.571
Isol. from a *Hippospongia* sp. Wax.

N-(Methylhexadecanoyl), Me ester: Hurghamide C
C₂₂H₄₁NO₃ 367.571
Isol. from a *Hippospongia* sp. Wax.

Posn. of methyl on *N*-acyl group not determined.

N-(13-Octadecanoyl), Me ester: Hurghamide D
C₂₃H₄₁NO₃ 379.582
Isol. from a *Hippospongia* sp.

N-Octadecanoyl, Me ester: [52634-31-8]
C₂₃H₄₃NO₃ 381.598
Isol. from *Fasciospongia cavernosa*.

N-Nonadecanoyl, Me ester: [52634-32-9]
C₂₄H₄₅NO₃ 395.624
Isol. from *Fasciospongia cavernosa*.

N-Eicosanoyl, Me ester: [52634-33-0]
C₂₅H₄₇NO₃ 409.651
Isol. from *Fasciospongia cavernosa*.

N-(2-Oxotetradecanoyl), Me ester: [70290-19-6]
C₁₉H₃₃NO₄ 339.474
Isol. from *Spongia* cf. *zimocca* and *Simularia dissecta*.

N-(2-Oxopentadecanoyl), Me ester: [70290-20-9]
C₂₀H₃₅NO₄ 353.501
Isol. from *Spongia* cf. *zimocca* and *Simularia dissecta*.

N-(2-Oxohexadecanoyl), Me ester: [70290-21-0]
C₂₁H₃₇NO₄ 367.528
Isol. from *Spongia* cf. *zimocca*.

N-(2-Hydroxytetradecanoyl), Me ester: [70290-22-1]
C₁₉H₃₅NO₄ 341.49
Isol. from *Spongia* cf. *zimocca*.

N-(2-Hydroxypentadecanoyl), Me ester: [70290-23-2]
C₂₀H₃₇NO₄ 355.517
Isol. from *Spongia* cf. *zimocca*.

N-(2-Hydroxyhexadecanoyl), Me ester: [70290-24-3]
C₂₁H₃₉NO₄ 369.543
Isol. from *Spongia* cf. *zimocca*.

N-(2-Acetoxy-2-tetradecenoyl), Me ester: C₂₁H₃₅NO₅ 381.511
Isol. from *Fasciospongia cavernosa*.

N-(2-Acetoxy-2-pentadecenoyl), Me ester: C₂₂H₃₇NO₅ 395.538
Isol. from *Fasciospongia cavernosa*.

N-(15-Methyl-8-hexadecenoyl), Me ester: Hurghamide E
[330203-64-0]
C₂₂H₃₉NO₃ 365.555
Isol. from a *Hippospongia* sp. Wax.

λ_{max} 205 (ε 14700) (MeOH).

N-(2-Alkylcyclopropyl)alkanoyl, Me ester (1): Hurghamide F
[330203-65-1]
C₂₂H₃₉NO₃ 365.555
Isol. from a *Hippospongia* sp. Wax.

Posn. of cyclopropyl ring in side-chain not yet determined.

N-(2-Alkylcyclopropylalkanoyl), Me ester (2): Hurghamide G
[330203-66-2]
C₂₄H₄₃NO₃ 393.609
Isol. from a *Hippospongia* sp. Wax.

Posn. of cyclopropyl not determined.

Kashman, Y. *et al.*, *Tetrahedron*, 1973, **29**, 3655 (*isol*)

Yunker, M.B. *et al.*, *Tet. Lett.*, 1978, 4651 (*isol*)

Holm, A. *et al.*, *Tet. Lett.*, 1980, 1125 (*synth*)

Neeman, J. *et al.*, *Arch. Toxicol., Suppl.*, 1983, **6**, 258 (*isol*)

Jackson, W.R. *et al.*, *Tet. Lett.*, 1988, **29**, 1983 (*synth*)

Isaac, B.G. *et al.*, *J. Antibiot.*, 1991, **44**, 795 (*isol*)

Guo, Y. *et al.*, *Nat. Prod. Lett.*, 1997, **9**, 281-288 (*Hurghamides A-D*)


Venkateswarlu, Y. *et al.*, *Indian J. Chem., Sect. B*, 1998, **37**, 832-834 (*N-2-acetoxy-2-alkenoyl derivis*)

Ramesh, P. *et al.*, *Biochem. Syst. Ecol.*, 1999, **27**, 661-662 (*2-oxoalkyl derivis, isol, Simularia*)

Guo, Y.-W. *et al.*, *J. Asian Nat. Prod. Res.*, 2000, **2**, 251-256 (*Hurghamides E-G*)

6-Amino-1-methylpurine A-842

1-Methyl-1H-purin-6-amine, 9CI. 1-Methyladenine. 1-Methyl-6-iminopurine. Spongopurine
[5142-22-3]

C₆H₇N₅ 149.155

Exists in the imino form in nonpolar solvs. and the amino form in H₂O. *Isol.* from starfishes *Asterias amurensis*, *Asterias rubens* and *Marthasterias glacialis*, and sponges *Geodia gigas* and *Hymeniacidon sanguinea*. Spawning-inducing factor in *Asterias*. Cryst. (H₂O). Mp 296-299° dec. pK_{a1} 7.2; pK_{a2} 11.

Picrate: Mp 255-257°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 714A (*ir*)


Ackermann, D. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1958, **312**, 210-213; 1961, **323**, 192-193 (*isol, ir, struct*)

Brookes, P. *et al.*, *J.C.S.*, 1960, 539-545 (*synth*)

Karatani, H. *et al.*, *Nature (London)*, 1969, **221**, 273-274 (*isol*)
 Lichtenberg, D. *et al.*, *J. Magn. Reson.*, 1972, **6**, 600-604 (*nmr*)
 Garrett, E.R. *et al.*, *J.A.C.S.*, 1972, **94**, 8532-8541 (*synth, uv*)
 Doree, M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1976, **73**, 1669-1673 (*biochem, bibl*)
 Dreyfus, M. *et al.*, *J.A.C.S.*, 1977, **99**, 7027-7037 (*tautom, bibl*)
 Cimino, G. *et al.*, *J. Nat. Prod.*, 1985, **48**, 523-528 (*isol, pmr*)

6-Amino-2-methylpurine A-843

2-Methyl-1H-purin-6-amine, 9CI. 2-Methyladenine
 [1445-08-5]




C₆H₇N₅ 149.155
 Obt. by hydrol. of RNAs. Cryst. (EtOH).
 Mp 350°.

▶ AU6519000

Robins, R.K. *et al.*, *J.A.C.S.*, 1953, **75**, 263 (*synth*)
 Littlefield, J.W. *et al.*, *Biochem. J.*, 1958, **70**, 642 (*isol*)
 Starr, J.L. *et al.*, *J. Biol. Chem.*, 1964, **239**, 3457 (*isol*)
 Huynh-Dinh, T. *et al.*, *J. Het. Chem.*, 1975, **12**, 111 (*synth, ir, pmr, uv, ms*)

6-Amino-3-methylpurine A-844

3-Methyl-3H-purin-6-amine, 9CI. 3-Methyladenine, 8CI
 [5142-23-4]



C₆H₇N₅ 149.155
 Isol. from the marine sponges *Plakortis aff. simplex* and *Topsentia genitrix*. Cryst. (H₂O). Mp 300° (291-292° dec.). pK_a 6.1. λ_{max} 273 (ε 14000) (MeOH) (Derep). λ_{max} 273 (ε 14010) (MeOH) (Berdy).

▶ AU6520000

9-β-D-Ribofuranosyl: 3-Methyladenosine [72055-63-1]
 C₁₁H₁₅N₅O₄ 281.271
 Plates (MeOH) (as tosylate salt). Mp 150° dec. (tosylate salt). [α]_D²⁰ -28.2 (c, 1.00 in H₂O) (tosylate salt). Imino at C-6.

9-(2-Deoxy-β-D-ribofuranosyl): 2'-Deoxy-3-methyladenosine [76227-26-4]
 C₁₁H₁₅N₅O₃ 265.271
 Powder (as tosylate salt). Mp 120° dec. (tosylate salt). Imino at C-6.


7-Oxide:

C₆H₇N₅O 165.154
 Sl. dark prisms. Mp 250-260° (dec.).
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 217A (*nmr*)

Brookes, P. *et al.*, *J.C.S.*, 1960, 539 (*synth*)
 Denayer, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 1358 (*synth, uv*)
 Jones, J.W. *et al.*, *J.A.C.S.*, 1962, **84**, 1914 (*synth, uv*)
 Elion, G.B. *et al.*, *J.O.C.*, 1962, **27**, 2478 (*synth, uv*)
 Pal, B.C. *et al.*, *J.C.S.*, 1964, 400 (*ir, cryst struct*)
 Lichtenberg, D. *et al.*, *J. Magn. Reson.*, 1972, **6**, 600 (*nmr*)
 Yamauchi, K. *et al.*, *J.O.C.*, 1975, **40**, 385 (*synth*)
 Stoller, C. *et al.*, *J. Nat. Prod.*, 1988, **51**, 383 (*isol, uv, pmr, cmr, ms*)
 Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2601 (*nucleosides*)
 Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2025 (7-oxide)
 Rudi, A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 682-685 (*isol, pmr, cmr*)

2-Amino-4-methylpyridine A-845

2-Amino-γ-picoline. 4-Methyl-2-pyridinamine, 9CI. W 45
 [695-34-1]



C₆H₈N₂ 108.143
 Alkaloid from *Acacia rigidula*. Analgesic, cardiac stimulant. Leaflets (petrol). Mp 98°. Bp 200-250°. pK_a 7.38. Sublimes. Component of Ascensil.

▶ LD₅₀ (rat, orl) 200 mg/kg. TJ5150000

Hydrochloride: [2403-84-1]
 Prisms (EtOH). Mp 176-177°.
Camphorsulfonate: *Piricardio. Varunax* [12261-97-1]
N-Ac: [5327-32-2]
 C₈H₁₀N₂O 150.18
 Prisms (C₆H₆). Mp 102-103° Mp 75-76°.

N-Benzoyl: [33120-20-6]
 C₁₃H₁₂N₂O 212.251
 Cryst. (petrol). Mp 114°.

N-Me: [45699-12-5]
 C₇H₁₀N₂ 122.169
 Mp 90-92°.

N-Benzyl: [13021-71-1]
 C₁₃H₁₄N₂ 198.267
 Cryst. (EtOH aq.). Mp 98-100°.

N-Nitro: [33245-30-6]
 C₆H₇N₃O₂ 153.14
 Solid. Mp 220°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 771B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 298C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1539B (*ir*)

Seide, O. *et al.*, *Ber.*, 1924, **57**, 791-794 (*synth, derivs*)

Sprinzak, Y. *et al.*, *J.A.C.S.*, 1956, **78**, 3207-3208 (*N-benzyl*)

Marchetti, G. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1963, **143**, 385-400 (*pharmacol*)


Clement, B.A. *et al.*, *Phytochemistry*, 1998, **49**, 1377-1380 (*isol*)

Bhattacharya, A. *et al.*, *Org. Process Res. Dev.*, 2007, **11**, 885-888 (*N-nitro*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, ALC250

3-Amino-N^b-methylseco-vocarpine A-846


[169133-06-6]



Absolute Configuration

C₂₂H₂₉N₃O₃ 383.489
 Alkaloid from leaves of *Hunteria zeylanica* (Apocynaceae). Mp 154-155°.

Subhadhirasakul, S. *et al.*, *Heterocycles*, 1995, **41**, 2049-2056 (*isol, uv, pmr, cmr, ms, cd, struct*)

2-Amino-5,9,12,15,17-octadecapentaen-3-ol A-847

C₁₈H₂₉NO 275.433

(2R,3S,5E,9Z,12Z,15Z)-form

Crucigasterin 275
 [150151-84-1]
 Isol. from *Pseudodistoma crucigaster*. Light yellow oil. λ_{max} 204 (ε 6200); 227 (ε 8700) (MeOH).

N,O-Di-Ac: [149849-81-0]
 Oil. [α]_D²⁰ +36 (c, 0.26 in MeOH).


17,18-Dihydro: 2-Amino-5,9,12,15-octadecatetraen-3-ol, 9CI. *Crucigasterin* 277
 [150151-83-0]

C₁₈H₃₁NO 277.449
 Isol. from *Pseudodistoma crucigaster*. Oil. λ_{max} 204 (ε 7900) (MeOH).

17,18-Dihydro, *N,O-di-Ac*: [149849-80-9]
 Oil. [α]_D²³ +36 (c, 0.53 in MeOH).

Jares-Erijman, E.A. *et al.*, *J.O.C.*, 1993, **58**, 5732-5737 (*isol*)

Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-4588 (*abs config*)

2-Amino-6,9,12,15-octadecatetraen-3-ol A-848

C₁₈H₃₁NO 277.449

(2S,3R,6Z,9Z,12Z,15Z)-form

Obscuraminol A. Halaminol D
 [350484-80-9]


Isol. from the tunicate *Pseudodistoma obscurum*. Oil. [α]_D²⁰ +5 (c, 0.14 in MeOH). [α]_D -23 (c, 0.78 in MeOH) (as di-Ac).

Clark, R.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1568-1571 (*Halaminol D*)

Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-4588 (*Obscuraminol A*)

 α -Amino-5-oxo-2(5H)-isoxazolebutanoic acid, 9CI A-849

α -Amino- γ -(isoxazolin-5-on-2-yl)butyric acid
[60102-46-7]



C₇H₁₀N₂O₄ 186.167

Isol. from sweet pea root exudates. Cryst. L-Config. not stated but presumed on biogenetic grounds.

Van Rompuy, L. *et al.*, *J.C.S. Perkin I*, 1973, 2503 (*pmr, uv*)


Lambein, F. *et al.*, *Biochem. Biophys. Res.*

Commun., 1974, **61**, 155 (*isol, struct, uv, pmr*)

Kuo, Y.H. *et al.*, *Arch. Int. Physiol. Biochim.*, 1976, **84**, 169 (*isol*)

 α -Amino-5-oxo-2(5H)-isoxazolepropanoic acid, 9CI A-850

α -Amino-5-oxo-3-oxazoline-2-propionic acid, 8CI. β -(Isioxazolin-5-on-2-yl)alanine. 2-Alanyl-3-isoxazolin-5-one



C₆H₈N₂O₄ 172.14

(S)-form

L-form

[59476-61-8]

Amino acid from the roots of pea seedlings (*Pisum* spp.) and *Lathyrus* spp. Fairly sol. H₂O. Mp 203-205° dec. [α]_D²⁰ -62 (c, 1.4 in H₂O). Base-labile. λ_{\max} 265 (€ 12300) (pH 2 buffer) (Berdy). λ_{\max} 267 (€ 13190) (pH 10 buffer) (Berdy).

N-(4-Amino-4-carboxybutanoyl): γ -Glutamyl- β -(isoxazolin-5-on-2-yl)alanine [142115-24-0]

C₁₁H₁₅N₃O₇ 301.255

Constit. of *Lens culinaris* (lentil) and some *Lathyrus* spp. Cryst. (Me₂CO aq.).

Lambein, F. *et al.*, *Biochem. Biophys. Res. Commun.*, 1969, **37**, 375 (*isol*)

Lambein, F. *et al.*, *Heterocycles*, 1976, **4**, 567 (*isol*)

Baldwin, J.E. *et al.*, *Tet. Lett.*, 1985, **26**, 5931 (*synth, pmr, ir*)

Lambein, F. *et al.*, *Phytochemistry*, 1992, **31**, 887 (*glutamyl deriv*)

Kuo, Y.-H. *et al.*, *Phytochemistry*, 1998, **49**, 43-48 (*occur, metab, biosynth*)

8-Amino-7-oxononanoic acid, A-851
9CI

[4707-58-8]

H₃CCH(NH₂)CO(CH₂)₅COOH

C₉H₁₇NO₃ 187.238


Isol. from *Penicillium chrysogenum* as a biotin intermediate. Cryst. (EtOH/Et₂O). Mp 133-134°.

Suyama, T. *et al.*, *CA*, 1964, **60**, 4013 (*synth*)
Iwahara, S. *et al.*, *Agric. Biol. Chem.*, 1966, **30**, 304

Eisenburg, M.A. *et al.*, *Biochemistry*, 1970, **9**, 108

2-Amino-3-oxo-3H-phenoxazine-1-carboxylic acid A-852

2-Amino-1-carboxy-3H-phenoxazin-3-one
[14994-68-4]



C₁₃H₈N₂O₄ 256.217


Prod. by a *Nocardia* strain. Dark orange needles. Sol. bases, MeOH, CHCl₃; poorly sol. H₂O, acids. Mp 310-320°. λ_{\max} 233 (€ 28220); 425 (€ 10620); 442 (€ 10380) (EtOH) (Berdy). λ_{\max} 230; 441 (HCl) (Berdy).

Gerber, N.N. *et al.*, *Biochemistry*, 1966, **5**, 3824 (*isol, synth, uv, ir*)

Gerber, N.N. *et al.*, *Can. J. Chem.*, 1968, **46**, 790 (*synth, ir*)

2-Amino-3-oxo-3H-phenoxazine-8-carboxylic acid A-853

Umycin C. SM 76. Antibiotic SM 76
[149837-40-1]



C₁₃H₈N₂O₄ 256.217

Prod. by *Streptomyces* sp. DSM 3813. Anthelmintic. Red powder. λ_{\max} 247 (€ 21900); 257 (€ 23400); 426 (€ 15800); 436 (€ 16200); 485 (sh) (€ 5010) (MeOH) (Derep).

N²-Ac: **Umycin A. SM 35. Antibiotic SM 35**
[116511-07-0]

C₁₅H₁₀N₂O₅ 298.254

Prod. by *Streptomyces* sp. DSM 3813. Anthelmintic. Red powder.

N²-Me: 2-(Methylamino)-3-oxo-3H-phenoxazine-8-carboxylic acid. **Texazone**
[87081-53-6]

C₁₄H₁₀N₂O₄ 270.244

Prod. by an *Actinomyces* strain WRAT-210. Dark red needles (Me₂CO). Sol. DMSO, Py; fairly sol. EtOAc, EtOH, Et₂O, CHCl₃; poorly sol. H₂O. Dec. without melting at 240°. λ_{\max} 247 (€ 21900); 257 (€ 23400); 426 (€ 15800); 436 (€ 16200); 485 (sh) (€ 5010) (MeOH). λ_{\max} 247 (€ 22000); 257 (€ 23440); 426 (€

15850); 436 (€ 16218); 480 (€ 5110) (MeOH) (Berdy).

N²-Me, Et ester: [87081-50-3]

Reddish-orange. Mp 256-257°.

N²-Me, amide: **Antibiotic BE 39907B. BE 39907B**

[181469-60-3]


C₁₄H₁₁N₃O₃ 269.259

Prod. by *Saccharopolyspora* sp. A39907. Cytotoxic agent.

Gerber, N.N. *et al.*, *J. Antibiot.*, 1983, **36**, 688 (*Texazone*)

Eur. Pat., 1988, 260 486; *CA*, **109**, 142586p (*Umycins*)

Japan. Pat., 1996, 96 198 861; *CA*, **125**, 219739f (*BE 39907B*)

2-Amino-3-oxo-3H-phenoxazine-1,8-dicarboxylic acid A-854

C₁₄H₈N₂O₆ 300.227

1-Amide: 8-Amino-9-(aminocarbonyl)-7-oxo-7H-phenoxazine-2-carboxylic acid.

Elloxazinone B

[943756-13-6]

C₁₄H₉N₃O₅ 299.242

Prod. by *Streptomyces griseus* Acta 2871. Antitumour agent. Red solid.

λ_{\max} 240 (log € 2.12); 256 (log € 1.93); 412 (log € 1.42); 431 (log € 1.5) (MeOH).

1-Amide, 8-Me ester: **Elloxazinone A**

[943756-12-5]


C₁₅H₁₁N₃O₅ 313.269

Prod. by *Streptomyces griseus* Acta 2871. Red solid. λ_{\max} 239 (log € 2.73); 259 (log € 2.57); 411 (log € 2.07); 428 (log € 2.09) (MeOH).

Graf, E. *et al.*, *J. Antibiot.*, 2007, **60**, 277-284 (*isol, uv, pmr, cmr, activity*)

1-(2-Amino-1-oxopropyl)-2,3,3a,13a,14,15,16,18a-octahydro-8-methoxy-5,9-metheno-9H-dipyrrrolo[3,2-b:1',2'-e][1,5,8]oxadiazacyclopentadecine-13,18(1H,12H)-dione, 9CI A-855

[147471-65-6]




C₂₂H₂₈N₄O₅ 428.487

Related to the Zizyphines. Alkaloid from flowers of *Sphaeranthus indicus* (Asteraceae). Mp 72°.

Chughtai, M.I.D. *et al.*, *Sci. Int. (Lahore)*, 1992, **4**, 151; *CA*, **118**, 230141f (*isol, struct*)

2-Amino-6-(1,2,3,4,5-pentahydroxypentyl)-4(1H)-pteridinone A-856




$C_{11}H_{15}N_5O_6$ 313.269
Cryst. (EtOH aq.). $[\alpha]_D^{25} +3.7$ (c, 0.1 in NH_4OH aq.).

2'-O-β-D-Glucuronopyranoside:

$C_{17}H_{23}N_5O_{12}$ 489.395
Isol. from cultures of *Mycobacterium phlei* and *Mycobacterium smegmatis*. Mp 250-280° (dec.). $[\alpha]_D^{25} +7.2$ (c, 0.26 in H_2O).

Goto, M. *et al.*, *Annalen*, 1965, **689**, 221 (*isol, uv, ir, ms*)

2-Amino-3-pentanol, 9CI A-857
[116836-16-9]



(2*S*,3*R*)-form

$C_5H_{13}NO$ 103.164

(2*S*,3*R*)-form [111061-02-0]
Oil. $[\alpha]_D^{23} +14$ (c, 1 in $CHCl_3$).

N-Ac: [138145-16-1]
 $C_7H_{15}NO_2$ 145.201
Isol. from the fungus *Scolecotrichum graminis*. $[\alpha]_D^{22} -25$ (c, 1 in EtOH).

N,N-Dibenzyl: [111060-70-9]
 $C_{19}H_{25}NO$ 283.413
Oil. $[\alpha]_D^{23} +48.2$ (c, 1 in $CHCl_3$).

(2*S*,3*S*)-form [180187-00-2]
Solid (hexane). Mp 67-68°. $[\alpha]_D^{23} -15.9$ (c, 0.34 in MeOH).

N-Ac: [138145-17-2]
Isol. from *Scolecotrichum graminis*. $[\alpha]_D^{22} -188$ (c, 0.5 in EtOH).

N,N-Dibenzyl: [111060-69-6]
Oil. $[\alpha]_D^{23} +70.5$ (c, 1 in $CHCl_3$).

(2*R*,3*R*)-form
(±)-threo-form
[147975-87-9]
Mp 104-105° (as oxalate).

(2*R*,3*S*)-form
(±)-erythro-form
[147975-86-8]
Mp 124-126° (as oxalate).

[141824-38-6, 599207-02-0]


Reetz, M.T. *et al.*, *Angew. Chem., Int. Ed.*, 1987, **26**, 1141-1143 (*synth*)

Tabuchi, H. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 2669-2671 (*N-Ac, isol*)

Andrés, J.M. *et al.*, *J.O.C.*, 1996, **61**, 4210-4213 (*synth, ir, pmr, cmr*)

Ueda, S. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 4101-4116 (*synth, pmr*)

2-Amino-3-pentanone A-858



$C_5H_{11}NO$ 101.148

(*S*)-form [138145-15-0]

N-Ac: 2-Acetamido-3-pentanone

$C_7H_{13}NO_2$ 143.185
Isol. from the fungus *Scolecotrichum graminis*. $[\alpha]_D^{25} -69.8$ (c, 1 in EtOH).

(±)-form [93222-93-6]

Off-white cryst. (as hydrochloride). Mp 127-129° (hydrochloride). CAS no. refers to hydrochloride.

N,N-Di-Me: [71504-23-9]

$C_7H_{15}NO$ 129.202

Bp₄₀ 63-64°.

[79851-68-6]

Coffen, D.L. *et al.*, *J.O.C.*, 1984, **49**, 5109

(*synth*)

Tabuchi, H. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 2669 (*isol, deriv, synth*)

Collina, S. *et al.*, *Tetrahedron: Asymmetry*, **10**, 2387-2397 (*N,N-di-Me, synth*)

(5-Aminopentyl)guanidine, A-859
9CI

1-Amino-5-guanidinopentane. Homoagmatine

[18431-52-2]
 $HN=C(NH_2)NH(CH_2)_4CH_2NH_2$

$C_6H_{16}N_4$ 144.219
Constit. of *Lathyrus sativus* (chickling pea). Major intermed. in homoarginine metab. pathway. Mp 135-137° (as hydrochloride).

N⁵-Ac: Acetylhomoagmatine

[180251-16-5]


$C_8H_{18}N_4O$ 186.256
Constit. of *Cliona celata*. Yellow amorph. solid. λ_{max} 206 (log ϵ 3.23) (MeOH).

Ramakrishna, S. *et al.*, *Phytochemistry*, 1973, **12**, 2691-2695 (*isol, synth, ir*)

Lenis, L.A. *et al.*, *Nat. Prod. Lett.*, 1996, **8**, 15-23 (*Acetylhomoagmatine*)

Castellanos, L. *et al.*, *Mar. Drugs*, 2006, **4**, 286-289 (*Acetylhomoagmatine, synth*)

7-Amino-1-phenazine-carboxylic acid, 8CI A-860
[24827-33-6]



$C_{13}H_9N_3O_2$ 239.233

Red needles (PhNO₂). Mp 340° dec.

Me ester: [24535-36-2]

$C_{14}H_{11}N_3O_2$ 253.26
Red needles (toluene). Mp 202-204°.

N⁵-Me, betaine: 7-Amino-1-carboxy-5-methylphenazinium hydroxide inner salt, 9CI. Aeruginosin A. Eruginosin A [21668-67-7]

$C_{14}H_{11}N_3O_2$ 253.26

Pigment from *Pseudomonas aeruginosa*. Insol. nonpolar. solvs., sol. H_2O , acids, bases, poorly sol. butanol, hexane. Red-purple fluor. in uv light. Dec. on heating. λ_{max} 235 (ϵ 31620); 280 (ϵ 44670); 396 (ϵ 11750); 515 (ϵ 14120) (pH 7 buffer) (Berdy). λ_{max} 235 ; 285 ; 380 ; 538 (HCl) (Berdy). λ_{max} 235 ; 275 ; 375 ; 525 (NaOH aq.) (Berdy).


Holliman, F.G. *et al.*, *Tetrahedron*, 1963, **19**, 1841-1848 (*synth, Me ester*)

Hansford, G.S. *et al.*, *J.C.S. Perkin 1*, 1972, 103 (*betaine, synth, biosynth*)

2-Aminophenol, 9CI A-861

o-Hydroxyaniline. 1-Amino-2-hydroxybenzene. Quetiomyacin B

[95-55-6]
[27598-85-2]



C_6H_7NO 109.127

Manuf. by redn. of 2-nitrophenol. Isol. from *Streptomyces* sp. and *Penicillium notatum*. Also isol. from a purple bacterium from a sponge of *Adocia* sp. Electrochem. polym. gives electroactive polymer films. Dye intermediate. Tuberculostatic. Possesses antibacterial props. Cryst. (H_2O). Sol. MeOH, H_2O , Et₂O, EtOAc; fairly sol. CCl_4 ; poorly sol. hexane. Mp 174°. pK_{a1} 4.78; pK_{a2} 9.97 (20°). Free base readily oxidised. λ_{max} 235 (ϵ 28400); 467 (ϵ 1700) (0.25*N* HCl) (Derep). λ_{max} 335 (ϵ 1230) (0.1*N* KOH) (Derep). λ_{max} 240 (ϵ 26300); 435 (ϵ 2330) (pH 7.4) (Derep). λ_{max} 229 ; 283 (H_2O) (Berdy). λ_{max} 268 (HCl) (Berdy). λ_{max} 300 (NaOH) (Berdy).

▶ Eye and respiratory tract irritant. LD₅₀ (rat, orl) 1300 mg/kg. LD₅₀ (mus, ipr) 350 mg/kg. Exp. reprod. and teratogenic effect. SJ4950000

Hydrochloride: [51-19-4]

Mp 207°.

▶ SJ6069000

N-Formyl: N-(2-Hydroxyphenyl)formamide, 9CI. 2-Hydroxyformanilide. N-Formylquetiomyacin B

[2843-27-8]

$C_7H_7NO_2$ 137.138

Prod. by *Penicillium notatum*. Needles (H_2O). Mp 129-129.5°. λ_{max} 218 (ϵ 10280); 245 (ϵ 11910); 286 (ϵ 6295) (MeOH) (Berdy).

N-Ac: 2-Acetamidophenol. 2-Hydroxyacetanilide. N-(2-Hydroxyphenyl)acetamide

[614-80-2]


$C_8H_9NO_2$ 151.165

Prod. by *Streptomyces* sp. Plates (EtOH). Sol. hot H₂O. Mp 209°.

► Eye irritant. AE4025000

- Anzai, K. *et al.*, *J. Antibiot.*, Ser. A, 1960, **13**, 125 (isol)
 Pfeifer, S. *et al.*, *Pharmazie*, 1972, **27**, 536-542 (isol)
 Oclarit, J.M. *et al.*, *Fish. Sci.*, 1994, **60**, 559 (isol)
 Pusecker, K. *et al.*, *J. Antibiot.*, 1997, **50**, 479 (N-Ac, isol, ir, pmr, ms)
 Rahaim, R.J. *et al.*, *Synthesis*, 2006, 3316-3340 (synth, pmr, cmr, N-Ac)

2-Amino-3H-phenoxazin-3-one, 9CI, 8CI
Questioniomycin A. AV toxin C
 [1916-59-2]



C₁₂H₈N₂O₂ 212.207

Isol. from *Acrospermum viticola*, *Brevibacterium iodinum*, *Calocybe gambosa* (St George's mushroom), *Microbispora aerata*, *Streptomyces thioluteus*, *Pycnoporus* sp., *Waksmania* sp., and mushroom *Lepiota americana*. Used as a 1mM soln. in EtOH as redox indicator for titanometric detn. of Fe(III), Au(III), Cr₂O₇²⁻, VO₄³⁻ and some organic compds.; redox indicator for stannometry. Aromatase and sulfatase inhibitor. Active against gram-positive bacteria, mycobacteria, *Candida albicans* and shows antitumour activity. Phytotoxin. Dark brown or red cryst. (EtOH). Sol. EtOH. Subl. 255-257. Sometimes occurs in amorph. form with Mp 296-297°. E° + 0.372V (pH 0.92, 23°). λ_{max} 235 (ε 28400); 467 (ε 1700) (0.25N HCl) (Derep). λ_{max} 335 (ε 1230) (0.1N KOH) (Derep). λ_{max} 240 (ε 26300); 435 (ε 2330) (pH 7.4) (Derep).

► LD₅₀ (mus, ipr) 200 mg/kg. SP7695000

N-β-D-Glucopyranosyl: N-β-D-Glucopyranosylquestioniomycin A

C₁₈H₁₈N₂O₇ 374.349

Prod. by *Microbispora* sp. TP-A0184. Antibacterial and cytotoxic agent. Red powder. Sol. DMSO, Py; fairly sol. MeOH; poorly sol. CHCl₃, EtOAc. Mp > 195° dec. [α]_D²⁵ +16.5 (c, 0.1 in Py). λ_{max} 240 (log ε 4.33); 423 (log ε 4.23) (MeOH).

N-Ac: N-(3-Oxo-3H-phenoxazin-2-yl)acetamide, 9CI. 2-Acetamido-3H-phenoxazin-3-one. N-Acetylquestioniomycin A
 [1916-55-8]

C₁₄H₁₀N₂O₃ 254.245

From *Brevibacterium iodinum*, *Microbispora aerata*, *Streptomyces thioluteus* and *Waksmania aerata*. Used as a 1mM soln. in EtOH as redox indicator for titanometric detn. of Fe(III), Au(III), Ce(IV), Cr₂O₇²⁻, VO₄³⁻ and some organic compounds. Active against *Sarcina lutea* and *Trichophyton* sp. Orange cryst. Sol. EtOH, Et₂O, C₆H₆, MeOH; poorly sol. H₂O. Subl. 165°. E° + 0.375V (pH 0.84, 23°). λ_{max}

240 (E1%/1cm 1400); 405 (E1%/1cm 1100) (EtOH) (Berdy).

N-Hydroxyacetyl: N-Hydroxyacetylquestioniomycin A. Chandrananimycin B

C₁₄H₁₀N₂O₄ 270.244

Prod. by a marine *Actinomadura* sp. M045. Orange solid. λ_{max} 238 (log ε 3.85); 399 (log ε 3.66) (MeOH).

6-Hydroxy: 2-Amino-6-hydroxy-3H-phenoxazin-3-one. 6-Hydroxyquestioniomycin A

C₁₂H₈N₂O₃ 228.207

Prod. by the marine *Halomonas* sp. GWS-BW-H8hM. Red solid. λ_{max} 236 (log ε 3.93); 277 (log ε 3.75); 430 (log ε 3.72) (MeOH). λ_{max} 230 (log ε 3.91); 284 (log ε 3.61); 464 (log ε 3.5) (MeOH/HCl).

9-Hydroxy, N-Ac: N-Acetyl-9-hydroxyquestioniomycin A. Chandrananimycin A

C₁₄H₁₀N₂O₄ 270.244

Prod. by a marine *Actinomadura* sp. M048. Orange solid. λ_{max} 226 (log ε 4.52); 270 (log ε 4.38); 423 (log ε 4.45) (MeOH).

7,8-Dimethoxy: 2-Amino-7,8-dimethoxy-3H-phenoxazin-3-one. 7,8-Dimethoxyquestioniomycin A. Peristrophine

C₁₄H₁₂N₂O₄ 272.26

Alkaloid from *Peristrophe roxburghiana*. Mp 250° dec. λ_{max} 235; 280 (sh); 484 (MeOH). λ_{max} 239; 273 (sh); 320 (sh); 544 (sh); 583 (MeOH/HCl).

Fischer, O. *et al.*, *Ber.*, 1961, **27**, 2784 (synth)
 Gerber, N.N. *et al.*, *Biochemistry*, 1964, **3**, 598; 1966, **5**, 3824 (isol, uv, ir)

Gerber, N.N. *et al.*, *J.O.C.*, 1967, **32**, 4055 (isol, uv, ir, bibl)

Ruzička, E. *et al.*, *Mikrochim. Acta*, 1967, 277 (use)

Ruzička, E. *et al.*, *CA*, 1968, **69**, 40930j (stannometry)

Ikekawa, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 1705 (synth, ir)

Sullivan, G. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1097 (isol)

Baer, H. *et al.*, *Pharmazie*, 1971, **26**, 108; 314 (isol)

Hishida, T. *et al.*, *Chem. Lett.*, 1974, 293 (synth)

Schlunegger, U.V. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 1383 (isol)

Motohashi, N. *et al.*, *Yakugaku Zasshi*, 1983, **103**, 364 (synth, praps)

Bolognese, A. *et al.*, *J. Het. Chem.*, 1986, **23**, 1003 (synth, pmr, uv, deriv)

Kinjo, J. *et al.*, *Tet. Lett.*, 1987, **28**, 3697 (isol)

Simandi, L.I. *et al.*, *Tet. Lett.*, 1993, **34**, 717 (synth)

Igarashi, Y. *et al.*, *J. Antibiot.*, 1998, **51**, 915-920 (*Glucosylquestioniomycin A*)

Qin, J.P. *et al.*, *Yaoxue Xuebao*, 1999, **34**, 599-603 (*Peristrophine*)

Kim, D.S. *et al.*, *Planta Med.*, 2000, **66**, 78 (*Questioniomycin A*, isol, activity)

Bolognese, A. *et al.*, *J. Med. Chem.*, 2002, **45**, 5205-5216 (synth, N-Ac, uv, pmr)

Maskey, R.P. *et al.*, *J. Antibiot.*, 2003, **56**, 622-629 (*Chandrananimycins*)

Bitzer, J. *et al.*, *J. Antibiot.*, 2006, **59**, 86-92 (*6-Hydroxyquestioniomycin A*)


Giurg, M. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 2006, **80**, 297-306 (synth)

Giurg, M. *et al.*, *Synth. Commun.*, 2007, **37**, 1779-1789 (synth, ir, uv, pmr)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, QC1275

2-[(4-Aminophenyl)amino]-pyrimidine A-863

N-(2-Pyrimidinyl)-1,4-benzenediamine
 [743449-54-9]



C₁₀H₁₀N₄ 186.216


Alkaloid from the bulbs of *Scilla maderensis*. α-Adrenoceptor antagonist. Needles (MeOH). Mp 270-271°. λ_{max} 213; 270 (MeOH).

Dias, C. *et al.*, *Planta Med.*, 2003, **69**, 1060-1062 (isol, pmr, cmr, ms)

2-Amino-1-phenylethanol A-864

α-Aminomethylbenzenemethanol, 9CI. α-(Aminomethyl)benzyl alcohol, 8CI. β-Amino-α-hydroxyethylbenzene. (Aminomethyl)phenylcarbinol. 2-Hydroxy-2-phenylethylamine. 1-Phenylethanolamine. **Resedine**†. Apophedrin. Norpheadrin. **Bisnorephedrine**

[7568-93-6]



C₈H₁₁NO 137.181

Log P 0.27 (calc).

► LD₅₀ (mus, ipr) 250 mg/kg. DN5500000

(R)-form [2549-14-6]

Mp 54-58°. [α]_D²⁰ -42.2 (c, 1 in EtOH).

O-Benzoyl: [111025-00-4]

C₁₅H₁₅NO₂ 241.289

Isol. from *Oxytropis pseudoglandulosa*. Needles (pentane). Mp 38-40°. [α]_D²⁵ +181.9 (c, 2.275 in MeOH).

N-Benzoyl: N-Benzoyl-2-hydroxy-2-phenylethylamine. 2-(Benzoylamino)-1-phenylethanol
 [111059-46-2]

C₁₅H₁₅NO₂ 241.289

Constit. of *Oxytropis pseudoglandulosa* (Fabaceae). Plates (MeOH/Et₂O). Mp 151-152° (143-144°). [α]_D²⁴ -16.2 (c, 0.81 in MeOH). [α]_D -21 (c, 0.7 in MeOH).

N-(3-Pyridinecarbonyl): N-Nicotinoyl-2-hydroxy-2-phenylethylamine. 2-(Nicotinoylamino)-1-phenylethanol
 [168780-03-8]

C₁₄H₁₄N₂O₂ 242.277


Isol. from aerial parts of *Oxytropis puberula*. Mp 157-158°. [α]_D -25.3 (c, 0.79 in MeOH).

N-Me: Halostachine. α-[(Methylamino)-methyl]benzenemethanol, 9CI. α-[(Methylamino)methyl]benzyl alcohol, 8CI. 2-(Methylamino)-1-phenethanol. β-Hydroxy-N-methylphenethylamine
 [495-42-1]

Alkaloid from *Halostachys caspica* (Chenopodiaceae). Hypertensive agent showing ephedrine-like activity. Mp 43-45°. [α]_D -47.

- N*-Me, hydrochloride: Mp 113-114°. $[\alpha]_D^{25}$ -52.4.
- N,N*-Di-Me: 2-Dimethylamino-1-phenylethanol. α -[(Dimethylamino)methyl]-benzenemethanol, 9CI. α -[(Dimethylamino)methyl]benzyl alcohol, 8CI. *N,N*-Dimethyl-1-phenylethanolamine. **Ubine** [34469-09-5]
Alkaloid from *Dolichothele uberiformis* (Cactaceae), also detected by ms in *Dolichothele longimamma*. Bp_{0.15} 58-60°. $[\alpha]_D^{25}$ -62.5 (c, 1.0 in MeOH). n_D^{25} 1.5164.
- Me ether*: 2-Methoxy-2-phenylethylamine. β -Methoxybenzeneethanamine, 9CI [55163-76-3]
C₉H₁₃NO 151.208
 d_4^{20} 1.02. Bp₂₀ 110-113°. $[\alpha]_D^{22}$ -118.5 (neat).
- Me ether, hydrochloride*: [62064-68-0]
Mp 158-159°.
- SH8090000
- (S)-form** [56613-81-1]
Mp 61-62° (55-57°). $[\alpha]_D$ +47.9 (c, 2.4 in EtOH).
- Hydrochloride*: [71025-82-6]
Mp 209-210°. $[\alpha]_D^{15}$ +48.8 (c, 5.08 in H₂O).
- N*-Ac: [155073-71-5]
C₁₀H₁₃NO₂ 179.218
 $[\alpha]_D$ +73 (c, 0.2 in CHCl₃) (>90% ee; room temp.).
- N*-Benzoyl: Constit. of *Oxytropis trichophysa* and *Oxytropis muricata*. Cryst. (Me₂CO), Mp 154-155°. $[\alpha]_D^{30}$ +35.2 (c, 0.68 in MeOH).
- N*-Benzoyl, *O*-Ac: 2-Acetyloxy-*N*-benzoyl-2-phenylethylamine. **Muricatide** [111025-01-5]
[128820-22-4, 109393-37-5]
C₁₇H₁₇NO₃ 283.326
Alkaloid from *Oxytropis muricata*. Needles (hexane/petrol). Mp 114-115° (110-112°). $[\alpha]_D^{20}$ +40.7 (c, 0.43 in CHCl₃). λ_{max} 208 (sh); 226 (EtOH).
- N*-tert-Butyloxycarbonyl:
C₁₃H₁₉NO₃ 237.298
Solid. Mp 66-68°. $[\alpha]_D^{20}$ +3.5 (c, 1 in EtOH) (99% ee).
- N*-Me: 2-(Methylamino)-1-phenylethanol [65058-52-8]
C₉H₁₃NO 151.208
Solid. Mp 45-47° (39-40°). Bp_{0.1} 99-102°. $[\alpha]_D^{20}$ +40.41 (c, 1.89 in EtOH).
- N,N*-Di-Me: [2202-69-9]
[939-45-7]
Mp 113.5° (as hydrochloride). $[\alpha]_D^{23}$ +77.6 (c, 0.1 in EtOH).
- N*-Benzyl:
C₁₃H₁₇NO 227.305
Cryst. (EtOAc/petrol). Mp 105-107°. $[\alpha]_D^{20}$ +33.8 (c, 2 in EtOH) (99% ee).
- (±)-form** [1936-63-6]
Pale yellow cryst. Mp 56-57°. Bp₁₇ 160°. p*K*_{a1} 8.79; p*K*_{a2} 11.9 (25°, 0.1M KCl).
- DA4727250
- Sulfate*: [50835-37-5]
Cryst. Mp 275-276°.
- DA4729900
N-Ac: [3306-05-6]
C₁₀H₁₃NO₂ 179.218
Solid. Mp 125-126°.
- N*-Benzoyl: [96328-15-3]
Leaflets (EtOH). Mp 148-149.5°.
- N*-tert-Butyloxycarbonyl:
Solid. Mp 120-121°.
- N*-Me: [68579-60-2]
Cryst. (CH₂Cl₂/hexane). Mp 72-73°.
- DA4976000
N,N-Di-Me: [2202-68-8]
[1797-76-8]
Isol. from leaves of *Aptenia cordifolia*, as partial racemate of low opt. rotn. Mp 144-145° (as hydrochloride)(synthetic). $[\alpha]_D^{25}$ +1.1 (c, 0.08 in CHCl₃) (nat.).
- N*-Benzyl:
Solid (EtOAc/petrol). Mp 93-96°.
- Me ether*: [3490-79-7]
Bp₁₂ 103-105°.
- (±)-form**
Alkaloid from *Reseda luteola* (Reseda-ceae), also obt. by hydrol. of 5-Phenyl-2-oxazolidinone, P-347. Vasoconstrictor. Mp 112-114°. Prob. artifact. The identification is dubious in view of the Mp which is higher than that of either the chiral or racemic form of authentic substance.
- O*-Benzoyl: **Trichophydine**. *Trichophidine* [67031-54-3]
C₁₅H₁₅NO₂ 241.289
Isol. from epigeal parts of *Oxytropis trichophysa*. Cryst. (hexane/petrol). Mp 200-203° (as hydrochloride). It is claimed that a previous alkaloid, Mp 38-40°, isolated from *Oxytropis pseudoglandulosa* given this struct. was erroneous and that it is an oxazoline of undetermined struct.
- N,O*-Dibenzoyl: **Trichophysine**
C₂₂H₁₉NO₃ 345.397
Isol. from epigeal parts of *Oxytropis trichophysa*. Cryst. (hexane/Et₂O). Mp 100-103°.
- [36297-03-7, 61217-79-6, 6027-95-8, 6589-55-5, 6853-14-1 (*N*-Me), 18867-43-1, 80997-83-7]
- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, 1, 1272B (*ir*)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 1277A (*N*-Me, *ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 585B (*N*-Me, *nmr*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 575C (*nmr*)
Rosenmund, K.W. *et al.*, *Ber.*, 1913, 46, 1034-1050 (*Me ether*)
Mannich, C. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1927, 2, 10 (*Me ether*)
Men'shikov, G.P. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1943, 13, 801; 1947, 17, 1569; *CA*, 39, 1172; 42, 2245 (*N*-Me, *isol. struct., synth*)
Rebstock, M.C. *et al.*, *J.A.C.S.*, 1951, 73, 3666-3670 (*synth*)
Dornow, A. *et al.*, *Chem. Ber.*, 1955, 88, 1267-1275 (*synth*)
Schöpf, C. *et al.*, *Annalen*, 1959, 626, 150-154 (*abs config*)
Poos, G.I. *et al.*, *J. Med. Chem.*, 1963, 6, 266-272 (*synth*)
- Lukes, R. *et al.*, *CA*, 1964, 61, 693g (*N*-Me, *abs config., synth*)
Reisch, J. *et al.*, *Fresenius' Z. Anal. Chem.*, 1968, 238, 29-35 (*nmr, struct*)
Cocolas, G.H. *et al.*, *J. Pharm. Sci.*, 1971, 60, 1749 (*N*-di-Me, *synth*)
Sabelli, H.C. *et al.*, *Adv. Behav. Biol.*, 1974, 10, 331 (*rev, props*)
Kirmse, W. *et al.*, *Chem. Ber.*, 1975, 108, 79-87 (*Me ether*)
Tadzhibaev, M.M. *et al.*, *Khim. Priir. Soedin.*, 1976, 12, 270-271; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, 12, 245-246 (*isol*)
Angeloni, A.S. *et al.*, *Gazz. Chim. Ital.*, 1977, 107, 421 (*N*-di-Me, *abs config*)
Ranieri, R.L. *et al.*, *J. Nat. Prod.*, 1977, 40, 173 (*N*-di-Me, *isol*)
Kruger, T.L. *et al.*, *J.O.C.*, 1977, 42, 4161 (*N*-di-Me, *ms, occur*)
Frank, H. *et al.*, *Angew. Chem.*, 1978, 90, 396-398 (*glc, resoln*)
Knizeo, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, 43, 1917-1923 (*synth*)
Wehrli, H. *et al.*, *Helv. Chim. Acta*, 1980, 63, 1915 (*N*-Me, *synth, ir, pmr*)
Meyers, A.I. *et al.*, *J.O.C.*, 1980, 45, 2785-2791 (*synth*)
Shannon, H.E. *et al.*, *J. Pharmacol. Exp. Ther.*, 1981, 217, 379-385 (*pharmacol, rev*)
Midland, M.M. *et al.*, *J.O.C.*, 1985, 50, 3237-3239 (*synth*)
Coote, S.J. *et al.*, *J.C.S. Perkin 1*, 1989, 2223-2228 (*S*-form, *N*-Me, *synth, ir, pmr, cmr*)
Ziegler, T. *et al.*, *Synthesis*, 1990, 575-578 (*R*-form, *synth, pmr*)
Brussee, J. *et al.*, *Tetrahedron*, 1990, 46, 1653-1658 (*R*-form, *synth*)
Izumi, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1993, 66, 1216-1221 (*Muricatide, synth*)
Akhmedzhanova, V.I. *et al.*, *Khim. Priir. Soedin.*, 1993, 29, 90-91; 1994, 30, 414-416; 1996, 32, 212-216; 1997, 33, 424-427; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, 29, 76; 1994, 30, 379-380; 1996, 32, 187-189; 1997, 33, 326-328 (*N*-nicotinoyl, *Trichophydine, Trichophysine, Muricatide*)
Uccello-Barretta, G. *et al.*, *J.O.C.*, 1997, 62, 827-835 (*R and S forms, synth, pmr*)
Laib, T. *et al.*, *Tetrahedron: Asymmetry*, 1998, 9, 168-178 (*S*-form, *N*-Ac, *synth*)
Kawamoto, A.M. *et al.*, *J.C.S. Perkin 1*, 2001, 1916-1928 (*N*-benzyl, *N*-Boc, *synth, ir, pmr, cmr*)
Becker, C.W. *et al.*, *Synthesis*, 2005, 2549-2561 (*S*-form, *N*-Me, *resoln*)
Veum, L. *et al.*, *Eur. J. Org. Chem.*, 2006, 1664-1671 (*N*-Ac)
- Tanis, S. *et al.*, *Tetrahedron*, 2006, 17, 2154-2182 (*R*-form, *S*-form, *N*-Me)
- Green, M.D. *et al.*, *Chem. Biodiversity*, 2007, 4, 118-128 (*N*-di-Me, *isol, pmr, cmr*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HNF000

1-Amino-4-phenyl-2,7-naphthyridine A-865
4-Phenyl-2,7-naphthyridin-1-amine, 9CI.
Lophocladine B




C₁₄H₁₁N₃ 221.261
Alkaloid from *Lophocladia* sp. Cytotoxic. Yellowish gum. λ_{max} 220 (log ϵ

4.01); 254 (log ϵ 4.02); 336 (log ϵ 3.89) (MeOH).

Gross, H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 640-644 (*isol, pmr, cmr, ms*)

5-(4-Aminophenyl)-2,4-pentadienoic acid A-866



$C_{11}H_{11}NO_2$ 189.213

Prod. by *Streptomyces* sp. Tu 3946. Orange-brown solid. Mp 178°. *Isol.* as a mixt. of (2E,4E)- and (2E,4Z)-isomers.

Amide: 5-(4-Aminophenyl)-2,4-pentadienamamide

$C_{11}H_{12}N_2O$ 188.229


Prod. by *Streptomyces* sp. Tu 3946.

Yellow solid. Mp 205°. Mixt. of (2E,4E)- and (2E,4Z)-isomers.

[154824-88-1, 154824-87-0, 154824-83-6, 154824-82-5]

Potterat, O. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 569 (*isol, synth*)

4-Amino-1-phenyl-1-penten-3-ol, 9CI A-867



$C_{11}H_{15}NO$ 177.246

(1E,3R,4S)-form

Merucathine

[107673-74-5]

Alkaloid from *Catha edulis* (Celastraceae). Needles. Mp 129-130°. $[\alpha]_D^{20} +29.6$ (c, 0.6 in MeOH).

Hydrochloride: [107673-75-6]

Mp 142-144°. $[\alpha]_D^{20} -5.2$ (c, 0.4 in H_2O).

(1E,3S,4S)-form

Pseudomerucathine

[96861-89-1]

Alkaloid from *Catha edulis* (Celastraceae). Cryst. Mp 103-104°. $[\alpha]_D^{20} -15$ (mixed solvent).

Hydrochloride: [107673-76-7]

Powder. Mp 194-196°. $[\alpha]_D^{20} -17.4$ (c, 0.4 in H_2O).


Brenneisen, R. *et al.*, *Planta Med.*, 1984, **531** (*isol, struct*)

Wolf, J.-P. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 918 (*struct, pmr, uv, cd, synth, abs config*)

Adam, W. *et al.*, *Synthesis*, 1995, 1066 (*synth*)

4-Amino-1-phenyl-1-penten-3-one, 9CI A-868

Merucathinone



$C_{11}H_{13}NO$ 175.23


(1E,4S)-form [107638-80-2]

Alkaloid detected in *Catha edulis* (Celastraceae). Mp 144° dec. (as oxalate salt). $[\alpha]_D +7.5$ (c, 0.85 in 2M HCl).

Wolf, J.-P. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 1498 (*synth, abs config, ir, uv, cd, pmr*)

2-Amino-1-phenyl-1-propanol A-869

α -(1-Aminoethyl)benzenemethanol, 9CI [48115-38-4]



$C_9H_{13}NO$ 151.208

See also refs. under 2-(Methylamino)-1-phenyl-1-propanol, M-386. Anorectic agent, resolving agent. Log P 0.58 (calc).

(1R,2R)-form

D-threo-form. *Nor-ψ-ephedrine*. *Norpseudoephedrine*. *Norisoeephedrine*. *Cathine*, *INN*

[37577-07-4]

Found in "Ma Huang" and *Catha edulis* (Celastraceae) (Khat), used as a stimulant drink in Arab countries. Plates (MeOH). Mp 77°. $[\alpha]_D^{20} +33.14$ (EtOH). Pharmacol. active isomer.

Hydrochloride: [53643-20-2]

Mp 180-181°. $[\alpha]_D^{20} +42.5$ (H_2O).

▶ RC9350000

N-Me: see 2-(Methylamino)-1-phenyl-1-propanol, M-386

(1S,2S)-form

(-)-*Nor-ψ-ephedrine*. *Katine*

[492-39-7]

Synthetic. Mp 77.5-78°. $[\alpha]_D^{20} -32.64$ (EtOH).

▶ LD₅₀ (rat, orl) 1538 mg/kg. RC9100000

Hydrochloride: Mp 180-181°. $[\alpha]_D^{20} -42.7$ (H_2O).

N-Me: see 2-(Methylamino)-1-phenyl-1-propanol, M-386

N,N-Di-Me: see 2-Dimethylamino-1-phenyl-1-propanol, D-738

(1S,2R)-form

(+)-*Norephedrine*

[37577-28-9]

Synthetic. Mp 52°. $[\alpha]_D^{27} +14.76$ (EtOH).

Hydrochloride: [40626-29-7]

Mp 171-172°. $[\alpha]_D^{27} +33.4$ (EtOH).

▶ RC3325000

N-Me: see 2-(Methylamino)-1-phenyl-1-propanol, M-386

N,N-Dibutyl: [114389-70-7]

$C_{17}H_{29}NO$ 263.422

Chiral catalyst, e.g. for enantioselective addn. of R_2Zn to aldehydes. Oil. Bp₂ 170°. $[\alpha]_D^{22} -24.4$ (c, 2 in hexane).

(1R,2S)-form

Norephedrine

[492-41-1]

Alkaloid from "Ma Huang", *Ephedra vulgaris* and *Catha edulis* (Ephedraceae, Celastraceae). Mp 51°. $[\alpha]_D -15$ (EtOH).

▶ RC2275000

Hydrochloride: [3198-15-0]

Mp 171-172°. $[\alpha]_D^{20} -33.27$ (H_2O).

▶ RC3240000

N-Formyl: *N-Formylnorephedrine*

$C_{10}H_{13}NO_2$ 179.218

Alkaloid from leaves of *Catha edulis* (Celastraceae). Needles (Et₂O/hexane). Mp 72-73°. $[\alpha]_D^{25} -45$ (c, 1.1 in $CHCl_3$).

(1RS, 2RS)-form

(±)-*Nor-ψ-ephedrine*

[54680-46-5]

Plates (petrol). Mp 77°.

Hydrochloride: [1485-15-0]

Mp 172-173°.

N-Me: see 2-(Methylamino)-1-phenyl-1-propanol, M-386

(1RS,2SR)-form

(±)-*Norephedrine*. **Phenylpropanolamine**, **BAN**, **INN**. *Propadrine*.

Mydratine

[14838-15-4]

Adrenergic, vasoconstrictor. Nasal decongestant. Plates (H_2O). Mp 104-105° (101-101.5°). Log P 0.58 (calc).

▶ LD₅₀ (rat, scu) 850 mg/kg. RC2625000

Hydrochloride: **Phenylpropanolamine hydrochloride**, **USAN**

[154-41-6] Nasal decongestant. Sol. H_2O . Mp 194°. Component of Allerest, Demazin, Dimetapp, Sinarest.

▶ Hypertensive and CNS effects reported when used therapeutically. LD₅₀ (rat, orl) 1490 mg/kg. Exp. teratogen. DN4200000

Complex with sulfonated styrenedivinylbenzene copolymer: **Phenylpropanolamine polistirex**, **USAN**

Adrenergic, vasoconstrictor.

N-Me: see 2-(Methylamino)-1-phenyl-1-propanol, M-386

[36393-56-3, 2153-98-2, 54705-32-7, 53631-70-2, 700-65-2, 4345-16-8, 115651-77-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1273A; 1273B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 577A; 577B; 577C; 578A (*nmr*)

Wolfes, O. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1930, **268**, 81 (*isol*)

Leithe, W. *et al.*, *Ber.*, 1932, **65**, 660 (*abs config*)


Hoover, F.W. *et al.*, *J.O.C.*, 1947, **12**, 506 (*synth*)

Heacock, R.A. *et al.*, *Can. J. Pharm. Sci.*, 1974, **9**, 64 (*pharmacol*)

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*occur*)
 Baudet, M. *et al.*, *Anal. Lett.*, 1979, **12**, 641 (*cmr*)
 Kaufer, I. *et al.*, *Anal. Profiles Drug Subst.*, 1983, **12**, 357
 Al-Meshal, I.A. *et al.*, *Phytochemistry*, 1986, **25**, 2241 (*N-Formylnorephedrine*)
 Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 1277
 Lasagna, L. *et al.*, *Phenylpropanolamine: A Review*, Wiley, New York, 1988, (*book*)
 Jackson, W.R. *et al.*, *Aust. J. Chem.*, 1990, **43**, 2045 (*synth*)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1990, **15**, 219 (*use*)
 Soai, K. *et al.*, *J.O.C.*, 1991, **56**, 4264 (*synth, ir, pmr, use, deriv*)
Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1222; 1252
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NNM000; NNN000; NNO000; NNM500; NNW500; PMJ500; NNV500; NNN500

2-Amino-1-phenyl-1-propanone, 9CI A-870

α -Aminopropiophenone. α -Benzoylethylamine. *Cathinone*, INN. *Norephedrone* [5265-18-9]



C₉H₁₁NO 149.192
 Analgesic, anorectic, antinociceptive agent. Possesses psychostimulant props. Log P 0.84 (calc).

(R)-form [80096-54-4]

► UC0418990

Hydrochloride: [76333-53-4]
 Cryst. (2-propanol/THF). Mp 189-190° dec. (175-177°). [α]_D²⁵ +47.3 (c, 1 in H₂O).

(S)-form [71031-15-7]

Isol. from leaves of *Catha edulis* (Khat) (Celastraceae) which are chewed for narcotic effect. [α]_D²⁶ -26.5 (c, 0.24 in CH₂Cl₂). Pharmacol. active enantiomer. Unstable except in dilute non-polar non-hydroxylic soln. Cathinone is the true alkaloid but is transformed in unfresh leaves to Ephedrine and Pseudoephedrine.

► UC0419000

Hydrochloride: [72739-14-1]
 Cryst. (2-propanol/Et₂O). Mp 188-190° (176-178°). [α]_D¹ -46.9 (c, 1 in H₂O).

Oxalate salt: [81626-17-7]
 Fine cryst. (EtOH). Mp 173-175°. [α]_D²⁵ -40.5 (c, 0.3 in MeOH).

(±)-form [75925-46-1]

Unstable solid. Mp 112-114°.

Hydrochloride: [42787-61-1]
 Needles (EtOH/Et₂O). Mp 187°.

N-Formyl: [102831-14-1]

C₁₀H₁₁NO₂ 177.202
 Cryst. (Me₂CO/hexane). Mp 93-96°.

N-Ac:

C₁₁H₁₃NO₂ 191.229
 Cryst. (C₆H₆). Mp 90-91°.

N-Me: 2-Methylamino-1-phenyl-1-propanone. *Ephedrone*

[5650-44-2]
 C₁₀H₁₃NO 163.219
 Yellow oil. Bp₁₁ 120-121°.

N-Et: [51553-17-4]

C₁₁H₁₅NO 177.246
 Shows anorexigenic activity in rats.
 Mp 196-198° (as hydrochloride). CAS no. refers to hydrochloride.

Gabriel, S. *et al.*, *Ber.*, 1908, **41**, 1127 (*synth*)
 Eberhard, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1915, **253**, 62

Kirchner, G. *et al.*, *Annalen*, 1959, **625**, 104

(*synth*)
 Schorno, X. *et al.*, *Experientia*, 1979, **35**, 572 (*isol*)

Szendrei, K. *et al.*, *Bull. Narc.*, 1980, **32**, 5 (*rev*)
 Glennon, R.A. *et al.*, *J. Med. Chem.*, 1980, **23**, 294 (*pharmacol*)

Berrang, B.D. *et al.*, *J.O.C.*, 1982, **47**, 2643 (*resoln*)

Martindale, The Extra Pharmacopoeia, 28th/29th edn., Pharmaceutical Press, 1982, 12537

Kalix, P. *et al.*, *Alcohol Alcohol.*, 1984, **19**, 319


(*pharmacol, use*)
 Wolf, J.-P. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 918; 1498 (*synth, bibl*)

Muchowski, J.M. *et al.*, *J.O.C.*, 1986, **51**, 3374 (*deriv*)

Farouz-Grant, F. *et al.*, *J. Med. Chem.*, 1997, **40**, 1977-1981 (*synth, pmr*)

4-Amino-2-piperidine-carboxylic acid, 9CI A-871

4-Aminopipercolic acid, 8CI



C₆H₁₂N₂O₂ 144.173

(2S,4R)-form

Trifluoroacetate salt: [756486-08-5]
 Solid. Mp 314° dec. [α]_D²⁵ -5.1 (c, 0.75 in H₂O).

(2S,4S)-form

L-trans-form

Trifluoroacetate salt: [756486-10-9]
 Solid. Mp 290° dec. [α]_D²⁵ -10.4 (c, 0.64 in H₂O).

N-Ac: 4-Acetamidopipercolic acid. 2-Carboxy-4-acetylaminopiperidine
 [72015-68-0]

C₈H₁₄N₂O₃ 186.21
 Alkaloid from leaves of *Calliandra haematocephala* (Fabaceae). Cryst. + ½ H₂O (Me₂CO aq.). [α]_D²⁰ -6 (c, 0.20 in H₂O). [α]_D²⁰ +0.5 (c, 0.196 in 2M HCl).
 Not correlated with the parent amine above due to scarcity of material.
 Incorr. descr. as (2S,4R) in the lit.

(-)-form

Present in *Strophanthus scandens* (Apocynaceae). Powder. Hygroscopic. Config. unknown.


Schenk, W. *et al.*, *Naturwissenschaften*, 1961, **48**, 223 (*isol*)

Marlier, M. *et al.*, *Phytochemistry*, 1979, **18**, 479 (*deriv*)

Machetti, F. *et al.*, *Eur. J. Org. Chem.*, 2004, 2928-2935 (2*S,4R*-form, 2*S,4S*-form, *synth, pmr, cmr*)

3-Amino-2-piperidinone, 9CI A-872

Ornithine lactam. Cycloornithine [1892-22-4]



C₅H₁₀N₂O 114.147

(R)-form [88763-76-2]

Residue occurring in pyoverdins. Cryst. (EtOAc). Mp 85.5-87.5°.

N-tert-Butyloxycarbonyl: [221874-51-7]

C₁₀H₁₈N₂O₃ 214.264
 Oil. [α]_D²⁴ -57.4 (c, 1.07 in CHCl₃).

N-Hydroxy; hydrochloride: [37524-32-6]

Cryst. (MeOH/EtOH). Mp 208-215° dec. [α]_D²⁴ -0.8 (c, 0.83 in H₂O).

(S)-form [34294-79-6]

Cryst. (EtOAc). Mp 42° Mp 86-87°. [α]_D²⁵ -12.4 (c, 3.44 in CHCl₃). [α]_D²⁵ -19.9 (dioxan).

Hydrochloride: [42538-31-8]

Mp 218-220° dec.

N³-Benzoyl: [148824-28-6]

[3328-28-7 (*L*-form)]
 C₁₂H₁₄N₂O₂ 218.255
 Mp 170-172°.

N-Hydroxy: [40716-77-6]

C₅H₁₀N₂O₂ 130.146
 Isol. from cultures of *Rhodotorula pilimanae*. Needles (MeOH/EtOH) (as hydrochloride). Mp 211-212° dec. (*hydrochloride*).

N¹-4-Methylbenzenesulfonyl: [20803-07-0]

Cryst. (MeOH). Mp 185-186°. [α]_D²⁵ +60.9 (c, 2.04 in DMF).

(±)-form [4077-38-7]

Mp 42°. Bp_{2.5} 125-135° lit. gives a pressure range.

N³-Ac: [98336-92-6]

C₇H₁₂N₂O₂ 156.184
 Needles (CHCl₃/Et₂O). Mp 187-188°.

N-Hydroxy; hydrochloride:

[7438-77-9 (free base)]
 Cryst. (EtOH). Mp 211-212°.

N-Hydroxy, N³-Ac: [37524-31-5]

C₇H₁₂N₂O₃ 172.183
 Cryst. (MeOH). Mp 203-205°.

Bergmann, M. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1926, **159**, 179-189 (*N³-Ac, synth*)


Emery, T.F. *et al.*, *Biochemistry*, 1966, **5**, 3694-3701 (*S*-form, *hydroxy, synth*)

Isowa, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 1464-1466 (*R*-form, *N-hydroxy, (±)-form, N-hydroxy N³-Ac, synth*)

Akers, H.A. *et al.*, *Biochemistry*, 1973, **12**, 1006 (*isol, synth, S*-form, *N-hydroxy*)

Terashima, S. *et al.*, *Tetrahedron*, 1973, **29**, 1487 (*synth*)
 Merguro, H. *et al.*, *Tet. Lett.*, 1975, 1309 (*cd*)
 Pellegata, R. *et al.*, *Synthesis*, 1978, 614-616 (*S-form, synth*)
 Blade-Font, A. *et al.*, *Tet. Lett.*, 1980, **21**, 2443 (*synth*)
 Winter, W. *et al.*, *Annalen*, 1983, 2021 (*synth*)
 Miyamura, K. *et al.*, *J.C.S. Dalton*, 1987, 1127 (*synth*)
 Toshima, H. *et al.*, *Tetrahedron*, 1999, **55**, 5793-5808 (*R-form, N-tert-butylloxycarbonyl*)
 Abe, M. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 785-797 (*S-form, synth, pmr, cmr*)
 Kubota, D. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 4158-4171 (*S-form, synth, pmr*)
 Urbanczyk-Lipkowska, Z. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 1254-1256 (*resoln, S-form*)

3-Aminopregna-5,17(20)-diene-2,16-diol A-873




C₂₁H₃₃NO₂ 331.497

(2α,3β,16α,17Z)-form

N,N-Di-Me: 3-(Dimethylamino)pregna-5,17(20)-diene-2,16-diol. **Kurchaline** [6885-72-9]
 C₂₃H₃₇NO₂ 359.551
 Alkaloid from the leaves of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (Me₂CO). Mp 185°. [α]_D -37 (c, 1 in CHCl₃).
 Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 1212-1216 (*struct, ir, pmr, ms*)

20-Aminopregna-5,16-diene-2,3-diol A-874




C₂₁H₃₃NO₂ 331.497

(2β,3β,20S)-form

3-Me ether, N²⁰,N²⁰-di-Me: 3-Methoxy-20-(dimethylamino)pregna-5,16-dien-2-ol. **Qasmine A**. 2-Hydroxy-N-methylsalignamine [657411-72-8]
 C₂₄H₃₉NO₂ 373.578
 Alkaloid from *Sarcococca saligna*. Acetylcholinesterase inhibitor. Yellow gum. [α]_D²⁰ -26 (c, 0.02 in CHCl₃). Incorrect synonym assigned in ref. λ_{max} 206 (log ε 3.8) (MeOH).

Zaheer-ul-Haq, *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4375-4380 (*isol*)
 Atta-ur-Rahman, *et al.*, *Helv. Chim. Acta*, 2004, **87**, 439-448 (*isol, pmr, cmr, ms*)

3-Aminopregna-5,16-dien-20-ol A-875




C₂₁H₃₃NO 315.498

(3α,20S)-form

N-Me: **Holadysamine** [1433-91-6]
 C₂₂H₃₅NO 329.525
 Minor alkaloid from the leaves of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (hexane). Mp 173°. [α]_D -78 (c, 1 in CHCl₃).
 N-Me, O,N-di-Ac: Mp 136°. [α]_D -74 (c, 1.3 in CHCl₃).
 Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 2158-2166; 1966, 1212-1216 (*isol, struct, ir, pmr, ms*)

20-Aminopregna-5,16-dien-3-ol A-876



C₂₁H₃₃NO 315.498

(3β,20S)-form

Me ether, N-Me: 3-Methoxy-20-(methylamino)pregna-5,16-diene. **Salignamine**. **Qasmine B** [657411-73-9]
 C₂₃H₃₇NO 343.551
 Alkaloid from *Sarcococca saligna*. Cholinesterase and acetylcholinesterase inhibitor. Yellow gum. [α]_D²⁰ -23 (c, 0.12 in CHCl₃). λ_{max} 202 (log ε 2.9) (MeOH).


Me ether, N-Me, N-formyl: N²⁰-Demethyl-N²⁰-formylsalonine **B** [657403-05-9]
 C₂₄H₃₇NO₂ 371.562
 Alkaloid from *Sarcococca saligna*. Cholinesterase inhibitor. Yellow gum. [α]_D²⁰ -21 (c, 0.07 in MeOH). λ_{max} 237 (log ε 2.86) (MeOH).

Me ether, N,N-di-Me: 20-(Dimethylamino)-3-methoxypregna-5,16-diene. **Salonine B** [618852-70-3]

C₂₄H₃₉NO 357.578
 Alkaloid from *Sarcococca saligna*. Cholinesterase inhibitor. Powder. [α]_D²⁰ -116 (c, 0.04 in MeOH). λ_{max} 206 (MeOH).

Zaheer-ul-Haq, *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4375-4380 (N²⁰-Demethyl-N²⁰-formylsalonine **B**)
 Atta-ur-Rahman, *et al.*, *Nat. Prod. Res.*, 2003, **17**, 235-241 (*Salonine B*)
 Atta-ur-Rahman, *et al.*, *Helv. Chim. Acta*, 2004, **87**, 439-448 (*Salignamine*)

3-Aminopregna-5,17(20)-dien-16-one A-877



C₂₁H₃₁NO 313.482

(3β,17E)-form


N-Me: 3-Methylaminopregna-5,17(20)-dien-16-one [113846-33-6]
 C₂₂H₃₃NO 327.509
 Alkaloid from leaves of *Didymeles perrieri* (Didymelaceae). [α]_D -37 (c, 0.60 in EtOH).

(3β,17Z)-form

Alkaloid from leaves of *Didymeles perrieri* (Didymelaceae). [α]_D -63 (c, 0.41 in CHCl₃).
 N-Me: [113846-32-5]
 Alkaloid from leaves of *Didymeles perrieri* (Didymelaceae). [α]_D -16 (c, 0.68 in CHCl₃).
 Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1987, 877

20-Aminopregna-1,4-dien-3-one A-878


[77546-81-7]



C₂₁H₃₁NO 313.482

(20S)-form [78548-54-6]

N,N-Di-Me: 20-(Dimethylamino)pregna-1,4-dien-3-one [91147-27-2]
 C₂₃H₃₅NO 341.536
 Alkaloid from the stem bark of *Didymeles cf. madagascariensis* (Didymelaceae). [α]_D +48 (c, 0.1 in CHCl₃).
 Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1984, 71; 1987, 877 (*isol, uv, cd, pmr, cmr, ms, struct*)
 De, D. *et al.*, *Steroids*, 1991, **56**, 189 (*synth*)

20-Aminopregnane-3,4-diol A-879C₂₁H₃₇NO₂ 335.529**(3β,4α,5α,20S)-form**N,N-Di-Me: *Terminaline*

[15112-49-9]

C₂₃H₄₁NO₂ 363.582

Alkaloid from *Pachysandra terminalis* and *Sarcococca hookeriana*. Cryst. Mp 243-244° (216-218°). [α]_D²⁵ +115 (c, 0.04 in MeOH). λ_{max} 242 (ε 1000) (MeOH).


N,N-Di-Me, Di-O-Ac: Mp 202-204°. [α]_D +40 (CHCl₃).

Kikuchi, T. *et al.*, *Tet. Lett.*, 1965, **24**, 1993-1999 (*isol*, *pmr*, *ms*)

Choudhary, M.I. *et al.*, *Steroids*, 2005, **70**, 295-303 (*isol*, *pmr*, *cmr*)

3-Aminopregnane-20-ol A-880

3-Amino-20-hydroxypregnane



(3α,5α,20R)-form

C₂₁H₃₇NO 319.529**(3α,5α,20R)-form***Isofuntumidine*, *Epifuntumidine*

[516-52-9]

Cryst. (MeOH). Mp 172-173°. [α]_D -6.2 (c, 2.7 in CHCl₃).

(3α,5α,20S)-form*Funtumidine*

[474-44-2]

Alkaloid from *Funtumia latifolia* (Apocynaceae). Hypocholesterolaemic agent. Shows antigonadotrophic, corticotrophic antiinflammatory and hypotensive activity. Respiratory stimulant. Mp 182°. [α]_D +10 (c, 1.5 in CHCl₃).

▶ TU4200000

(3α,5β,20R)-form

Cryst. (DMF). Mp 160-163°. [α]_D^{23.5} +14.4 (c, 1 in CHCl₃).

(3β,5α,20R)-form

Cryst. Mp 172-174°. [α]_D^{21.5} +5.4 (c, 1 in CHCl₃).

(3β,5β,20R)-form


Cryst. (EtOAc). Mp 150-152°. [α]_D²⁰ 0 (c, 1 in CHCl₃).

Janot, M.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1958, **246**, 3076; 1959, **248**, 982 (*ir*, *struct*, *Funtumidine*)

U.S. Pat., 1963, 3 098 082; *CA*, **59**, 11632d (*synth*)

Schmitt, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 761 (*synth*)

Dodoun, H. *et al.*, *Phytochemistry*, 1973, **12**, 923 (*Isofuntumidine*)

20-Aminopregnane-3-ol A-881C₂₁H₃₇NO 319.529**(3β,5α,20S)-form***Funtuphyllamine A*

[667-71-0]

Alkaloid from *Funtumia africana* (Apocynaceae). Mp 173°. [α]_D +13 (c, 1.3 in CHCl₃).

N-Ac: *Holacetine*

[60882-55-5]

C₂₃H₃₉NO₂ 361.567

Alkaloid from *Holarrhena antidysenterica* root bark (Apocynaceae). Mp 258°. [α]_D²⁵ +6.9 (EtOH).

N-Me: 20-Methylaminopregnane-3-ol.

Funtuphyllamine B

[474-81-7]

C₂₂H₃₉NO 333.556

Alkaloid from *Funtumia africana* and *Malouetia bequaertiana* (Apocynaceae). Mp 214°. [α]_D +24 (CHCl₃).

N,N-Di-Me: 20-(Dimethylamino)pregnane-3-ol. *Funtuphyllamine C*, *Sarcococinine D*

[474-51-1]

C₂₃H₄₁NO 347.583

Alkaloid from *Funtumia africana* (Apocynaceae). Also from *Sarcococca ruscifolia* (Buxaceae). Mp 172°. [α]_D +24 (CHCl₃).

N,N-Di-Me, O-β-D-glucopyranoside: *Pachyaxioside A*

[128255-18-5]

C₂₉H₅₁NO₆ 509.725

Alkaloid from *Pachysandra axillaris*.

N,N-Di-Me, O-(3-methyl-2-butenyl):

*O-β,β-Dimethylacrylylfuntuphyllamine C*C₂₈H₄₇NO₂ 429.685

Alkaloid from *Sarcococca pruniformis* (Buxaceae). Mp 274-276°. [α]_D²⁶ +10 (CHCl₃).

Černý, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1957, **22**, 76 (*synth*)

Janot, M.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1960, **250**, 2445 (*isol*, *ir*, *struct*)

Chatterjee, A. *et al.*, *Chem. Ind. (London)*, 1966, 769 (*Dimethylacrylylfuntuphyllamine C*)


Demailly, G. *et al.*, *Tet. Lett.*, 1975, 2471 (*synth*)

Rej, R.N. *et al.*, *Phytochemistry*, 1976, **15**, 1173 (*Holacetine*)

Biesemans, M. *et al.*, *Bull. Soc. Chim. Belg.*, 1985, **94**, 59 (*pmr*, *cmr*, *conformn*)

Qiu, M. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 535; *CA*, **113**, ; 74724n (*Sarcococinine D*)

Qiu, M. *et al.*, *CA*, 1990, **113**, 55838m (*N,N-Di-Me glucoside*)

3-Aminopregnane-20-one A-882

(3α,5α)-form

C₂₁H₃₅NO 317.514**(3α,5α)-form***Funtumine*

[474-45-3]

Alkaloid from *Funtumia latifolia*, *Holarrhena febrifuga* and *Holarrhena congolensis* (Apocynaceae).

Hypocholesterolaemic agent, shows antigonadotropic, corticotropic, antiinflammatory and hypotensive props. Respiratory stimulant. Mp 126°. [α]_D +95 (c, 1.7 in CHCl₃).

▶ LD₅₀ (mus, ivn) 30 mg/kg. TU4375000

Hydrochloride: Mp 280°. [α]_D +60 (c, 0.67 in MeOH).

N-Ac: [4465-66-1]

Mp 240°.

N-Me: 3-(Methylamino)pregnane-20-one.

N-Methylfuntumine

[40855-51-4]

C₂₂H₃₇NO 331.54

Alkaloid from *Holarrhena febrifuga* and *Sarcococca coriacea*. Yellow cryst. (CHCl₃). Mp 129-130°. [α]_D²⁰ +92 (c, 0.9 in CHCl₃). λ_{max} 202 (log ε 4.8) (MeOH).

N,N-Di-Me: 3-(Dimethylamino)pregnane-20-one. *N,N-Dimethylfuntumine*

[3022-47-7]

C₂₃H₃₉NO 345.567

Alkaloid from *Holarrhena febrifuga* (Apocynaceae). Mp 128-130°. [α]_D²⁰ +92 (c, 1.8 in CHCl₃).

(3β,5α)-form*Dihydroholaphyllamine*

[2136-36-9]

[24202-19-5 (hydrochloride)]

Alkaloid from *Holarrhena floribunda*

(Apocynaceae). Noncryst.

Janot, M.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1958, **246**, 3076; 1959, **248**, 982 (*isol*, *ir*, *struct*)

Leboeuf, M. *et al.*, *Ann. Pharm. Fr.*, 1969, **27**, 217-218 (*Dihydroholaphyllamine*)

Boutigue, M.H. *et al.*, *Bull. Soc. Chim. Fr.*, 1973, 750-753 (*synth*, *N-Methylfuntumine*, *N,N-Dimethylfuntumine*)


Dadoun, H. *et al.*, *Phytochemistry*, 1973, **12**, 923-928 (*Funtumine*, *N-Methylfuntumine*, *N,N-Dimethylfuntumine*)

Kapnang, H. *et al.*, *Tet. Lett.*, 1977, 3469-3472 (*N,N-Dimethylfuntumine*, *synth*)

Kalauni, S.K. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1423-1426 (*N-Methylfuntumine*, *pmr*, *cmr*)

20-Aminopregnan-3-one

A-883

C₂₁H₃₅NO 317.514(5 α ,20S)-form**Wrightiamine B**

[640266-39-3]

Alkaloid from the leaves of *Wrightia japonica*. Amorph. solid. $[\alpha]_D^{25} +5$ (c, 0.04 in MeOH).

N-Me: 20-(Methylamino)pregnan-3-one.

Funtumafrine B

[474-82-8]

C₂₂H₃₇NO 331.54

Alkaloid from the leaves of *Funtumia africana*. Mp 160°. $[\alpha]_D +43$ (c, 1.2 in CHCl₃).

N,N-Di-Me: 20-(Dimethylamino)pregnan-3-one.

Funtumafrine C

[474-52-2]

C₂₃H₃₉NO 345.567

Alkaloid from *Funtumia africana*, *Chonemorpha macrophylla* and *Sarcococca coriacea*. Cryst. (CHCl₃). Mp 176° (174°). $[\alpha]_D^{25} +50$ (c, 0.6 in CHCl₃). λ_{max} 202 (log ϵ 4.7) (MeOH).

Janot, M.M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1960, **250**, 2445-2447 (*isol. struct. synth*)


Banerji, J. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 346 (*isol*)

Kalauni, S.K. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1423-1426 (*Funtumafrine C*, *pmr, cmr*)

Kawamoto, S. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 737-739 (*Wrightiamine B*)

20-Aminopregn-5-ene-3,16-diol

A-884

C₂₁H₃₅NO₂ 333.513(3 β ,16 α ,20S)-form

N-Me: 20-(Methylamino)pregn-5-ene-3,16-diol

[16378-59-9]

C₂₂H₃₇NO₂ 347.54Mp 265°. $[\alpha]_D^{24} -69.9$ (c, 0.5 in propanol).

N-Me, N,3,16-tri-Ac: Mp 214°. $[\alpha]_D^{30.5} -142$ (c, 1 in CHCl₃).

(3 β ,16 β ,20S)-form

3-Me ether, N,N-di-Me: 20-(Dimethylamino)-3-methoxypregn-5-en-16-ol.

Pachyaximine B

[128286-24-8]

C₂₄H₄₁NO₂ 375.593

Alkaloid from *Pachysandra axillaris* (Buxaceae).

Di-Me ether, N,N-di-Me: 3,16-Dimethoxy-N,N-dimethylpregn-5-en-20-amine, 9CI. 20-(Dimethylamino)-3,16-dimethoxypregn-5-ene. **Sarcosaligmine**.

[873804-22-9]

C₂₅H₄₃NO₂ 389.62

Alkaloid from *Sarcococca saligna*. Solid. Mp 212-217°. $[\alpha]_D^{27} +77$ (c, 0.44 in CHCl₃).


Schmitt, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2240-2248 (*N-Me, synth*)

Qui, M. *et al.*, *Zhiviu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 535-539; *CA*, **113**, 74724n (*Pachyaximine B*)

Naeem, I. *et al.*, *CA*, 2005, **144**, 146530m (*Sarcosaligmine*)

3-Aminopregn-5-en-20-ol

A-885

C₂₁H₃₅NO 317.514(3 α ,20R)-form**Holaminol**

[28375-11-3]

Minor alkaloid from the leaves of *Holarrhena febrifuga* (Apocynaceae). Mp 205°. $[\alpha]_D -78.6$ (CHCl₃). Phys. const. for synthetic sample.

O,N-Di-Ac: Mp 226°. $[\alpha]_D^{20} -23.8$ (c, 0.9 in CHCl₃).

(3 β ,20R)-formN-Me: **Holaphyllinol**

[5750-08-3]

C₂₂H₃₇NO 331.54

Alkaloid from *Holarrhena floribunda* (Apocynaceae). Mp 226-227°. $[\alpha]_D -71$ (c, 1.3 in CHCl₃).

(3 β ,20S)-formN-Me: **Holaphyllidine**

[3837-07-8]

C₂₂H₃₇NO 331.54

Alkaloid from *Holarrhena africana* (*Holarrhena floribunda*) (Apocynaceae).

Goutarel, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 646 (*synth*)

Leboeuf, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1964, **259**, 3401 (*Holaphyllidine*)


Leboeuf, M. *et al.*, *Ann. Pharm. Fr.*, 1969, **27**, 217 (*Holaphyllinol*)

Dadoun, H. *et al.*, *Ann. Pharm. Fr.*, 1973, **31**, 237 (*isol, pmr, ms*)

20-Aminopregn-5-en-3-ol

A-886

20-Amino-3-hydroxypregn-5-ene

C₂₁H₃₅NO 317.514(3 β ,20S)-form(3 β ,20S)-form**Holafefrine**

[5035-10-9]

Alkaloid from *Holarrhena febrifuga*, *Holarrhena mitis*, *Holarrhena crassifolia* and *Kibatalia arborea* (Apocynaceae). Mp 177°. $[\alpha]_D -61.5$ (c, 0.85 in CHCl₃). $[\alpha]_D^{25} -66.6$ (90% Py aq.).

O- β -D-Glucopyranoside: **Conopharyngine**[†]

[7067-29-0]

C₂₇H₄₅NO₆ 479.656

Alkaloid from *Conopharyngia pachysiphon* (Apocynaceae). Hypotensive agent. Mp 285-288°.

O- β -D-Glucopyranoside; hydrochloride:

Mp 259-260°. $[\alpha]_D^{25} -66.6$ (90% Py aq.).

O,N-Di-Ac: Mp 250°. $[\alpha]_D -56.6$ (c, 0.7 in CHCl₃).

N-Me: 20-Methylaminopregn-5-en-3-ol.

Irehamine

[7083-27-4]

C₂₂H₃₇NO 331.54

Alkaloid from *Funtumia elastica*, *Funtumia latifolia* and *Dictyophleba lucida* (Apocynaceae). Mp 230°. $[\alpha]_D -33$ (c, 0.9 in CHCl₃).

N,N-Di-Me: 20-Dimethylaminopregn-5-en-3-ol.

Irephine. Buxomegine

[2309-39-9]

C₂₃H₃₉NO 345.567

Alkaloid from *Funtumia elastica* and *Buxus sempervirens*. Mp 174°. $[\alpha]_D^{25} +40$ (c, 1.3 in CHCl₃). λ_{max} 204 (MeOH).

N,N-Di-Me, O- β -D-glucopyranoside:

Pachyaxioside B

[128255-19-6]

C₂₉H₄₉NO₆ 507.709

Alkaloid from *Pachysandra axillaris*.

Me ether, N,N-di-Me: 3-Methoxy-20-dimethylaminopregn-5-ene. **Pachyaximine A**

[128255-08-3]

[32164-75-3]

C₂₄H₄₁NO 359.594

Alkaloid from *Pachysandra axillaris* and *Sarcococca prunifoliosa*. Needles. Mp 155° (152-153°). $[\alpha]_D^{30} -32$ (CHCl₃). Alkaloid C from *S. prunifoliosa* originally assigned 3 α -config.

Lucas, R.A. *et al.*, *J.A.C.S.*, 1960, **82**, 5688 (*glucoside*)

Janot, M.-M. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 285 (*isol. struct*)

Vetter, W. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 1324 (*ms*)

Truong-Ho, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2332 (*derivs*)

Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 348 (*derivs*)

Robinson, C.H. *et al.*, *Chem. Ind. (London)*, 1966, 377 (*pmr*)

Kohli, K.M. *et al.*, *Phytochemistry*, 1971, **10**, 442 (*Alkaloid C*)


Einhorn, J. *et al.*, *Phytochemistry*, 1972, **11**, 769 (*isol*)

Dadoun, H. *et al.*, *Ann. Pharm. Fr.*, 1973, **31**, 237 (*isol*)

Qiu, M. *et al.*, *Zhiviu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 535; *CA*, **113**, 74724 (*Pachyaximine A*)

Qiu, M. *et al.*, *CA*, 1990, **113**, 55838m (*glucoside*)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Res.*, 2003, **17**, 235-241 (*Alkaloid C*)
 Babar, Z.U. *et al.*, *Steroids*, 2006, **71**, 1045-1051 (*Irehine*)

3-Aminopregn-5-en-16-one A-887

$C_{21}H_{33}NO$ 315.498

3 α -form

N-Me: 3-Methylaminopregn-5-en-16-one.

Holadysine


$C_{22}H_{35}NO$ 329.525

Alkaloid from the leaves of *Holarrhena antidyenterica* (Apocynaceae). Cryst. (hexane). Mp 120°. $[\alpha]_D$ -199 (c, 1.2 in $CHCl_3$).

N-Me, *N-Ac*:

Cryst. by subl. Mp 175°. $[\alpha]_D$ -163 (c, 1.3 in $CHCl_3$).

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 2158; 1966, 1212 (*isol, struct, ir, pmr, ms*)

3-Aminopregn-5-en-20-one A-8883 α -form

$C_{21}H_{33}NO$ 315.498

Log P 4.13 (uncertain value) (calc).

3 α -form**Holamine**

[28840-94-0]

Alkaloid from *Holarrhena curtisii* and *Holarrhena floribunda* and leaves of *Holarrhena febrifuga* (Apocynaceae). Antigonadotropic, corticotropic and antiinflammatory agent. Strong local anaesthetic. Respiratory depressant. Diuretic causing sodium retention. Cytotoxic and leishmanicidal agent. Cryst. (EtOAc). Mp 135-136°. $[\alpha]_D$ +23 (c, 0.88 in $CHCl_3$).

N-Me: 3-Methylaminopregn-5-en-20-one.

Monomethylholamine

[41567-48-0]

$C_{22}H_{35}NO$ 329.525

Alkaloid from leaves of *Holarrhena febrifuga* (Apocynaceae). Cryst. (Me_2CO). $[\alpha]_D^{20}$ +25 (c, 1 in $CHCl_3$). Constants refer to a synthetic sample.

N,N-Di-Me: 3-Dimethylaminopregn-5-en-20-one. **Dimethylholamine**

[41515-65-5]

$C_{23}H_{37}NO$ 343.551

Alkaloid from leaves of *Holarrhena febrifuga* (Apocynaceae). Cryst. (Me_2CO). Mp 135°. $[\alpha]_D$ +18 (c, 1 in

$CHCl_3$). Constants are for a synthetic sample.

15 α -Hydroxy: 3-Amino-15-hydroxypregn-5-en-20-one. **15-Hydroxyholamine**

[215595-78-1]

$C_{21}H_{33}NO_2$ 331.497

Alkaloid from *Holarrhena curtisii*. Cytotoxic and leishmanicidal agent. Amorph. $[\alpha]_D$ +47.2 (c, 0.23 in $CHCl_3$).

3 β -form**Holaphyllamine**

[7068-92-0 (hydrochloride)]

Alkaloid from *Holarrhena floribunda* and *Holarrhena crassifolia* (Apocynaceae). Antiinflammatory agent. Noncryst. Mp 300° dec. (260°) (as hydrochloride). $[\alpha]_D$ +33 (c, 0.68 in MeOH) (hydrochloride). Log P 4.13 (uncertain value) (calc).

N-Me: **Holaphylline**

[562-02-7]

$C_{22}H_{35}NO$ 329.525

Alkaloid from *Holarrhena floribunda*, *Holarrhena febrifuga*, *Holarrhena congoensis* and *Holarrhena crassifolia* (Apocynaceae). Antigonadotropic, corticotropic and antiinflammatory agent. Respiratory depressant, diuretic. Mp 128°. $[\alpha]_D$ +23.7 ($CHCl_3$).

N-Me, *N-Ac*: [2842-66-2]

Mp 205-206°. $[\alpha]_D$ +16.7 (c, 1.2 in $CHCl_3$).

N,N-Di-Me: **Dimethylholaphyllamine.**

Methylholaphylline

[6405-70-5]

$C_{23}H_{37}NO$ 343.551

Alkaloid from *Holarrhena floribunda* (Apocynaceae). Cryst. (MeOH). Mp 121°. $[\alpha]_D$ +30.5 (c, 0.86 in $CHCl_3$).

Janot, M.-M. *et al.*, *Bull. Soc. Chim. Fr.*, 1959, 896-900 (*Holamine*, *Monomethylholamine*, *Dimethylholamine*)


Janot, M.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1960, **251**, 559-561 (*Holaphyllamine*, *Holamine*)

Goutarel, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 4575-4582 (*synth, pmr*)

Einhorn, J. *et al.*, *Phytochemistry*, 1972, **11**, 769-777 (*Holaphyllamine*, *Holaphylline*)

Dadoun, H. *et al.*, *Phytochemistry*, 1973, **12**, 923-928 (*Holamine*, *Methylholamine*, *Dimethylholamine, isol, synth*)

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1332-1336 (*Holamine*, *15 α -Hydroxyholamine, pmr, cmr*)

3-Aminopregn-16-en-20-one A-889

$C_{21}H_{33}NO$ 315.498


(3 α ,5 α)-form

N-Di-Me: 3-Dimethylaminopregn-16-en-20-one. **Sarcorucinine B**
 [143086-42-4]

$C_{23}H_{37}NO$ 343.551

Alkaloid from aerial parts of *Sarcococca ruscifolia* (Buxaceae).

Qiu, M. *et al.*, *Yunnan Zhiwu Yanjiu*, 1991, **13**, 445; *CA*, **117**, 108079f (*isol, ir, pmr, ms*)

3-Aminopregn-17(20)-en-16-one A-890

$C_{21}H_{33}NO$ 315.498

(3 β ,5 α ,17E)-form

N-Me: **3 β -(Methylamino)pregn-17(20)-en-16-one**

[113846-34-7]

$C_{22}H_{35}NO$ 329.525

Alkaloid from the leaves of *Didymelea perrieri* (Didymelaceae). $[\alpha]_D$ -64 (c, 0.52 in $CHCl_3$).

N,N-Di-Me: 3-(Dimethylamino)pregn-17(20)-en-16-one. **E-Salignone**

[204450-32-8]

$C_{23}H_{37}NO$ 343.551

Constit. of *Sarcococca saligna*. Oil. $[\alpha]_D^{25}$ -26 (c, 0.076 in $CHCl_3$). λ_{max} 242 (log ϵ 3.8) (MeOH).

(3 β ,5 α ,17E)-form

N,N-Di-Me: **Z-Salignone**


[204450-33-9]

$C_{23}H_{37}NO$ 343.551

Constit. of *Sarcococca saligna*. Oil. $[\alpha]_D^{25}$ -46 (c, 0.086 in $CHCl_3$). λ_{max} 242 (log ϵ 3.7) (MeOH).

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1987, 877 (*isol, uv, ir, pmr, ms, cd, struct*)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, **11**, 81-91 (*Salignones*)

20-Aminopregn-4-en-3-one A-891

$C_{21}H_{33}NO$ 315.498

(20S)-form

N-Me: 20-Methylaminopregn-4-en-3-one.

4-Dehydrofuntumafrine B

[16316-42-0]

$C_{22}H_{35}NO$ 329.525

Alkaloid from the leaves of *Didymelea madagascariensis*. $[\alpha]_D$ +91 (c, 0.72 in EtOH).

N,N-Di-Me: 20-Dimethylaminopregn-4-en-3-one. **4-Dihydrofuntumafrine C.**

Buxaprogestine

[113762-72-4]

C₂₃H₃₇NO 343.551

Alkaloid from the leaves of *Didymelus perrieri* and from leaves of *Buxus papilosa* (Buxaceae). $[\alpha]_D^{25} +49$ (c, 1.8 in EtOH). $[\alpha]_D^{25} +26$ (c, 1.78 in CHCl₃). Identity of the two samples not establ.

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1987, 877 (isol, uv, ir, pmr, ms, cd, struct)
Choudhary, M.I. *et al.*, *Phytochemistry*, 1988, 27, 271 (*Buxaprogestine*)

3-Aminopropanal, 9CI A-892

3-Aminopropionaldehyde, 8CI
[352-92-1]

H₂NCH₂CH₂CHOC₃H₇NO 73.094Liq. Bp₁₂ 45-46°.

2,4-Dinitrophenylhydrazone:

Cryst. (EtOH). Mp 178°.

Di-Et acetal: 3,3-Diethoxy-1-propylamine. APEA

[41365-75-7]

C₇H₁₇NO₂ 147.217Liq. Bp₂₀ 68-70°.

N-(3-Methylbutanoyl): 3-Methyl-N-(3-oxopropyl)butanamide, 9CI. Diplocardia *Luciferin*

[58947-91-4]

C₈H₁₅NO₂ 157.212

Bioluminescent constit. of the earthworm *Diplocardia longa*. Oil.

N-(3-Methylbutanoyl), 2,4-dinitrophenylhydrazone:

Yellow cryst. Mp 174°.

N-tert-Butyloxycarbonyl: [58885-60-2]

C₈H₁₅NO₃ 173.211

Oil.

N-Benzoyloxycarbonyl: [65564-05-8]

C₁₁H₁₃NO₃ 207.229

Mp 57-58°.

Albers, H. *et al.*, *Ber.*, 1946, 79, 623Birkhofer, L. *et al.*, *Chem. Ber.*, 1958, 91, 2383Ohtsuka, H. *et al.*, *Biochemistry*, 1976, 15, 1001 (*Luciferin*)

Geall, A.J. *et al.*, *Tetrahedron*, 2000, 56, 2449-2460 (*N-benzoyloxycarbonyl*, synth, pmr, cmr)

Delfourne, E. *et al.*, *J. Med. Chem.*, 2003, 46, 3536-3545 (*N-tert-butyloxycarbonyl*, synth, pmr, cmr)

2-Amino-1,3-propanediol, 9CI, 8CI A-893

Serinol

[534-03-2]

(HOCH₂)₂CHNH₂C₃H₉NO₂ 91.11Mp 128°. Bp 264-265° Bp_{0.06} 115-120°.

Hydrochloride: [73708-65-3]

Mp 104°.

Oxalate: [24070-20-0]

Hygroscopic cryst. Mp 202-203°.

N-Ac: [2655-79-0]

C₅H₁₁NO₃ 133.147

Mp 89-90°.

N-(tert-Butyloxycarbonyl): [125414-41-7]

C₈H₁₇NO₄ 191.227

Needles (hexane/EtOAc). Mp 84-85°.

N-(2-Hydroxybenzoyl): N-Salicyloyl-2-amino-1,3-propanediol

C₁₀H₁₃NO₄ 211.217

Prod. by *Streptomyces hygroscopicus*. Antibacterial agent. Amorph. solid. λ_{\max} 225 (log ϵ 4.32); 283 (sh) (log ϵ 3.54); 302 (sh) (log ϵ 3.5) (MeOH).

N-(2-Hydroxybenzoyl), 1-Ac:

C₁₂H₁₅NO₅ 253.254

Prod. by *Streptomyces hygroscopicus*. Antibacterial agent. Yellow amorph. solid. $[\alpha]_D^{25} -29$ (c, 0.14 in MeOH). λ_{\max} 228 (log ϵ 4.56); 282 (sh) (log ϵ 4.17); 301 (sh) (log ϵ 4.13) (MeOH).

Dibenzyl ether:

C₁₇H₂₁NO₂ 271.358

Oil.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 551B (nmr)

Den Otter, H.P. *et al.*, *Rec. Trav. Chim.*

(*J. R. Neth. Chem. Soc.*), 1938, 57, 13 (synth)

Karrer, P. *et al.*, *Helv. Chim. Acta*, 1948, 31, 1617 (*oxalate*, synth)

Adkins, H. *et al.*, *J.A.C.S.*, 1948, 70, 3121 (synth)

Langenbeck, W. *et al.*, *Naturwissenschaften*, 1955, 42, 389 (synth)

Foster, A.B. *et al.*, *J.C.S.*, 1958, 1890 (synth)

Szammer, J. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1969, 61, 417 (synth)

Japan. Pat., 1976, 76 67 788; *CA*, 85, 157975

(biosynth)

Rai, B.L. *et al.*, *J. Med. Chem.*, 1998, 41, 3347-3359 (*dibenzyl ether*)

Neri, C. *et al.*, *Adv. Synth. Catal.*, 2003, 345, 835-848 (*tert-butyloxycarbonyl*)

Hossain, M.S. *et al.*, *Phytochemistry*, 2004, 65, 2147-2151 (*N-salicyloyl amides*)

Gu, K. *et al.*, *Bioorg. Med. Chem.*, 2006, 14, 1339-1347 (synth, pmr, cmr)

2-Amino-1-propanol, 9CI A-894

Alaninol. 2-Hydroxyisopropylamine
[78-91-1]

C₃H₉NO 75.11

(S)-form [2749-11-3]

Oil. Bp₁₁ 72-73°. $[\alpha]_D^{20} +15.8$.

N-(2-Hydroxybenzoyl): N-Salicyloyl-2-amino-1-propanol

C₁₀H₁₃NO₃ 195.218Prod. by *Streptomyces hygroscopicus*.

Antibacterial agent. Amorph. solid.

 $[\alpha]_D^{25} -2.5$ (c, 0.08 in MeOH).

[27646-78-2, 92596-51-5, 40916-69-6, 40916-61-8, 40916-73-2, 57707-57-0, 54490-98-1, 15521-18-3]

Hossain, M.S. *et al.*, *Phytochemistry*, 2004, 65, 2147-2151 (*N-salicyloyl amide*)

(3-Aminopropoxy)guanidine, 9CI A-895 γ -Guanidinoxypropylamine

[97091-01-5]

HN=C(NH₂)NHOCH₂CH₂CH₂NH₂C₄H₁₂N₄O 132.165


Constit. of *Canavalia gladiata* (sword-bean) and *Wisteria floribunda*.

Hamana, K. *et al.*, *Biochem. Biophys. Res. Commun.*, 1985, 129, 46 (*isol*)

Matsuzaki, S. *et al.*, *Phytochemistry*, 1990, 29, 1311 (*isol*)

4-(3-Aminopropyl)-1,2-benzenediol, 9CI A-896

3-(3,4-Dihydroxyphenyl)-1-propylamine. Homodopamine
[52336-45-5]

C₉H₁₃NO₂ 167.207

Hydrobromide: [52336-30-8]

Cryst. (2-propanol). Mp 156-158°.

Di-Me ether: 3,4-Dimethoxybenzenepropylamine, 9CI. 3-(3,4-Dimethoxyphenyl)propylamine, 8CI
[14773-42-3]

C₁₁H₁₇NO₂ 195.261

Constit. of *Piper arboricola*. Bp₁₅ 158-161°.

Di-Me ether, hydrochloride: [59734-58-6]

Cryst. (MeOH/EtOAc). Mp 166-167°.

Di-Me ether, picrate:

Cryst. (EtOH). Mp 145-146°.

Benington, F. *et al.*, *J.O.C.*, 1956, 21, 1545-1546 (synth, di-Me ether)

Winn, M. *et al.*, *J. Med. Chem.*, 1975, 18, 434-437 (synth)

Ho, C.F. *et al.*, *CA*, 1981, 95, 138470w (*isol*, deriv)

N-(3-Aminopropyl)-1,5-pentanediamine, 9CI A-897

N-(3-Aminopropyl)cadaverine. 1,9-Diamino-4-azanonane. 1,5,11-Triazaundecane. as-Homospermidine
[56-19-9]

H₂NCH₂(CH₂)₃CH₂NHCH₂CH₂-CH₂NH₂

C₈H₂₁N₃ 159.274


Isol. from *Hydrogenobacter halophilus* and *Paracoccus* sp. Cryst. (EtOH) (as trihydrochloride). Mp 258-259° (trihydrochloride). Bp 94-95°.

Mervyn, I. *et al.*, *J. Med. Chem.*, 1964, 7, 710-716 (synth)

Hamana, K. *et al.*, *Biochem. J.*, 1992, 284, 741-747 (*isol*)

4-(2-Aminopropyl)phenol, 9CI, 8CI A-898

1-(4-Hydroxyphenyl)-2-propylamine. p- β -Aminopropylphenol. Hydroxyamfetamine, IN. Hydroxyamphetamine, BAN. Methyltyramine. Norpholedrine. Norveritol. Oxamphetamine. Paredrinex
[103-86-6]

C₉H₁₃NO 151.208

Sympathomimetic and mydriatic drug. Amphetamine metab. Log P 0.92 (calc).

▶ SJ6475010

(R)-form [1518-89-4]

Mp 109-111.5°. [M]_D²⁰ -56° (c, 0.02 in EtOH).

Me ether: [50505-80-1]

C₁₀H₁₅NO 165.235

Needles (EtOH/Et₂O) (as hydrochloride). Mp 251-253° dec. (hydrochloride). [α]_D²⁵ -23.2 (c, 2.03 in H₂O). CAS no. refers to hydrochloride.

(S)-form [1693-66-9]

Mp 110-111.5°. [M]_D²⁰ + 59° (c, 0.02 in EtOH).

(±)-form [1518-86-1]

Alkaloid from *Acacia rigidula*. Cryst. (C₆H₆). Mp 125°.

Hydrobromide: **Hydroxyamphetamine hydrobromide, USAN**. *Mycadrine*. *Paredrine*. *Pedrolon*

[140-36-3]

Cryst. (EtOH/Et₂O). Mp 171-172°.

▶ SJ6650000

Me ether: **4-Methoxy-α-methylbenzeneethanamine. 1-(4-Methoxyphenyl)-2-propylamine. 4-Methoxyamphetamine** [23239-32-9]

[64-13-1]

C₁₀H₁₅NO 165.235

Alkaloid from *Acacia rigidula*.

Liq. Mp 210° (as hydrochloride). Bp₁₄ 141°.

Mannich, C. *et al.*, *Ber.*, 1910, **43**, 189

(*synth*)

Hoover, F.W. *et al.*, *J.O.C.*, 1947, **12**, 501

(*synth*)

Dirkx, I.P. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1964, **83**, 535 (*ord*)

v. Dijk, J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1965, **84**, 521 (*resoln*)

Sever, P.S. *et al.*, *Biochem. Soc. Trans.*, 1973, **1**, 1158 (*metab*)

Hahn, W.E. *et al.*, *Acta Pol. Pharm.*, 1979, **36**, 259 (*synth*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., *Pharmaceutical Press*, 1993, 1246


Clement, B.A. *et al.*, *Phytochemistry*, 1998, **49**, 1377-1380 (*Me ether, isol*)

Kohno, H. *et al.*, *Synth. Commun.*, 1998, **28**, 1935-1945 (*Me ether*)

Aminopteroic acid

A-899

4-[[[(2,4-Diamino-6-pteridiny)]methyl]amino]benzoic acid, 9CI. p-[N-2,4-Diamino-6-pteridyl]methylaminobenzoic acid [36093-85-3]



C₁₄H₁₃N₇O₂ 311.302

Parent acid present in Aminopterin.

N^{acr}-Me: **4-[[[(2,4-Diamino-6-pteridiny)]methyl]methylamino]benzoic acid** [19741-14-1]

C₁₅H₁₅N₇O₂ 325.329

Orange powder. Mp > 300°.

Sato, H. *et al.*, *Nippon Kagaku Kaishi*, 1951, **72**, 866; *CA*, **47**, 5946

U.S. Pat., 1951, 2 568 597; *CA*, **46**, 5094

Venditti, J.M. *et al.*, *Cancer Res.*, 1960, **20**, 691


Elliott, R.D. *et al.*, *J.O.C.*, 1970, **35**, 1676-1680 (*N-Me*)

Piper, J.R. *et al.*, *J.O.C.*, 1977, **42**, 208

6-Amino-9H-purine-9-propa-noic acid, 9CI

A-900

3-(9-Adeniny)]propionic acid [4244-47-7]



C₈H₉N₅O₂ 207.191

Isol. from *Lentinus edodes* (shiitake).

Exhibits anticholesteraeamic activity. Mp 277-278° dec. λ_{max} 262 (ε 14400) (0.1N NaOH). λ_{max} 259 (ε 13800) (0.1N HCl).

Me ester: [70259-15-3]

C₉H₁₁N₅O₂ 221.218

Cryst. (Et₂O/MeOH). Mp 182-183°.

Et ester: [7083-40-1]

C₁₀H₁₃N₅O₂ 235.245

Solid. Mp 167-168°.

Nitrile: **6-Amino-9-(2-cyanoethyl)purine** [4244-45-5]

C₈H₈N₆ 188.191

Solid (H₂O). Mp 258-261° (245-247°).

Lira, E.P. *et al.*, *J.O.C.*, 1966, **31**, 2188-2191

(*acid, nitrile, synth*)

Chakraborti, S.K. *et al.*, *Indian J. Chem.*, 1969, **7**, 426 (*synth*)

Saito, Y. *et al.*, *Tet. Lett.*, 1970, 4863 (*isol, struct, synth*)


Holy, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 3444-3465 (*synth*)

Poriterra, S.E. *et al.*, *Khim. Geterotsikl. Soedin.*, 1982, 539-541; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1982, 415-417 (*nitrile*)

4-Amino-2,3-pyridinedicar-boxylic acid, 9CI

A-901

[122475-55-2]



C₇H₆N₂O₄ 182.135


Isol. from fruit bodies of the poisonous mushroom *Clitocybe acromelalga*. Cryst. (H₂O). Sol. H₂O, MeOH; poorly sol. Me₂CO, hexane. λ_{max} 262 (ε 9550) (0.1N HCl) (Derep). λ_{max} 245 (sh) (ε 8710); 290 (ε 2090) (H₂O at pH 11) (Derep). λ_{max} 265 (ε 9120) (H₂O pH 7) (Derep). λ_{max} 252 (ε 10000) (MeOH) (Berdy). λ_{max} 265 (ε 9800); 290 (ε 2080) (H₂O) (Berdy).

Hirayama, F. *et al.*, *Phytochemistry*, 1989, **28**, 1133 (*isol, uv, ir, pmr, cmr, ms, synth, struct*)

2-Amino-3-(2-pyridinyl)pro-panoic acid

A-902

α-Amino-2-pyridinepropanoic acid, 9CI. **3-(2-Pyridinyl)alanine**. Antibiotic SF 2538. SF 2538 [17407-39-5] [37535-52-7]



(*S*)-form

C₈H₁₀N₂O₂ 166.179

(S)-form [37535-51-6]

[120191-52-8]

Prod. by *Streptomyces albus*. Cryst. Sol. H₂O; poorly sol. Me₂CO, hexane. [α]_D -33.6 (H₂O). Incorrectly indexed in CA. λ_{max} 254 (ε 3720); 260 (ε 4170); 266 (ε 3100) (H₂O). λ_{max} 254 (ε 3720); 260 (ε 4170); 266 (ε 3100) (H₂O) (Berdy). λ_{max} 261 (ε 7600) (HCl) (Berdy).

(±)-form [17407-44-2]

Cryst. (EtOH aq.). Mp 209-212° (209-210°).

[177164-50-0, 169547-60-8, 33715-92-3, 172927-00-3, 163513-22-2, 33560-87-1]

Slater, G. *et al.*, *Tetrahedron*, 1967, **23**, 2823-2828 (*synth*)

Watanabe, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1968, **41**, 1634-1638 (*synth*)

Veselova, L.N. *et al.*, *Zh. Obshch. Khim.*, 1972, **42**, 1123-1125; 1973, **43**, 1637-1640 (*synth, resoln*)


Iwata, M. *et al.*, *CA*, 1989, **110**, 169903x (*isol*)

Zhao, J.S. *et al.*, *Yaoxue Xuebao*, 1994, **29**, 558-560; *CA*, **122**, 10513w (*synth*)

2-Amino-4-pyrimidinecar-boxylic acid

A-903

[2164-65-0]



C₅H₅N₃O₂ 139.113

Constit. of *Lathyrus tingitanus* seeds.

Precursor of the pyrimidine moiety of Lathyrine. Fine needles (H₂O). Mp 265° dec. Mp 285° dec.

Nitrile: **2-Amino-4-cyanopyrimidine**

[36314-98-4]

C₅H₄N₄ 120.113

Mp 222° dec.

Matsukawa, T. *et al.*, *Yakugaku Zasshi*, 1952, **72**, 909; *CA*, **47**, 6425d (*synth*)


Whitlock, B.J. *et al.*, *J.O.C.*, 1965, **30**, 115 (*synth, uv*)

Brown, E.C. *et al.*, *Phytochemistry*, 1996, **42**, 61 (*occur*)

Doláková, P. *et al.*, *Heterocycles*, 2007, **71**, 1107-1115 (*nitrile*)

α -Amino-1H-pyrrole-1-hexanoic acid, 9CI A-904

N^ε-Pyrrolylnorleucine. Pyrrole-1-norleucine. 1-(5'-Amino-5'-carboxypentyl)pyrrole



C₁₀H₁₆N₂O₂ 196.249

(S)-form

L-form

[156539-32-1]

Secondary lipid peroxidation prod. derived from *L*-lysine and 4,5-epoxy-2-alkenols. Widely distributed in fresh foodstuffs, e.g. meats, fish, vegetables and nuts. Characterised spectroscopically.

Chiang, G.H. *et al.*, *J. Agric. Food Chem.*, 1988, **36**, 506-509 (*hplc, occur*)

Zamora, R. *et al.*, *Lipids*, 1994, **29**, 243-249;

1995, **30**, 477-483 (*synth, detn, pmr, cmr*)

Zamora, R. *et al.*, *Biochim. Biophys. Acta*, 1995, **1258**, 319-327 (*formn*)


Hidalgo, F.J. *et al.*, *J. Agric. Food Chem.*, 1995, **43**, 1023-1028 (*occur, formn*)

Zamora, R. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 1942-1947 (*occur*)

4-Amino-2-pyrrolidinecarboxylic acid A-905

4-Aminoproline, 9CI

[3285-76-5]



(2*R*,4*S*)-form

C₅H₁₀N₂O₂ 130.146

(2*R*,4*S*)-form

(+)-*trans*-form

Prod. by *Ascochyta caulina*. Phytotoxin. [α]_D²¹ +47.2 (c, 0.4 in H₂O).

(2*S*,4*R*)-form

(-)-*trans*-form

[16257-88-8]

Prisms. Mp 229-230° dec. [α]_D²¹ -57.8 (H₂O).

Hydrochloride (1:2): [16257-89-9]

Mp 257-258°.

N-tert-*Butyloxycarbonyl*: [132622-69-6]

C₁₀H₁₈N₂O₄ 230.263

Solid. Mp 240-241° dec. [α]_D -35.9 (c, 1.0 in H₂O).

*N*¹-(4-*Methylbenzenesulfonyl*): [16257-87-7]

C₁₂H₁₆N₂O₄S 284.335

Prisms (EtOH aq.). Mp 275-276° dec.

[α]_D²⁰ -62.7 (EtOH aq.).

(2*S*,4*S*)-form

(-)-*cis*-form

[16257-83-3]

Prisms (EtOH aq.). Mp 191-193° dec. [α]_D²⁰ -57.4 (H₂O). Unstable.

Hydrochloride (1:2): [16257-84-4]

Mp 247°.

N-tert-*Butyloxycarbonyl*: [132622-66-3]

Solid. Mp 225-228° dec. [α]_D²⁰ +21 (c, 0.24 in H₂O).

*N*¹-(4-*Methylbenzenesulfonyl*): [16257-82-2]

Needles (EtOH aq.). Mp 284-285° dec. [α]_D²⁰ -72 (EtOH aq.).

Andreata, R.H. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1493; 2701 (*synth, pmr*)

Webb, T.R. *et al.*, *J.O.C.*, 1991, **56**, 3009-3016 (*N-Boc*)


Evidente, A. *et al.*, *Phytochemistry*, 2000, **53**, 231-237 (*isol, pmr, cmr*)

Fisher, A. *et al.*, *J. Med. Chem.*, 2006, **49**, 307-317 (*N-Boc*)

1-Aminopyrrolizidine A-906

Hexahydro-1H-pyrrolizin-1-amine

[170442-12-3]



C₇H₁₄N₂ 126.201

(1*S*,7*aR*)-form

(-)-*cis*-form

Waxy solid. [α]_D²⁰ -24.1 (c, 5.5 in MeOH).

*N*¹-(2-*Methylbutanoyl*): *Laburnamine*

[2072-58-4]

C₁₂H₂₂N₂O 210.319

Alkaloid from the seeds of *Cytisus laburnum*. Mp 125-127°.

*N*¹-(4-*Methoxy-E-cinnamoyl*): *Absouline*

[112513-33-4]

C₁₇H₂₂N₂O₂ 286.373

Alkaloid from *Hugonia oreogena* and *Hugonia penicillanthemum*. Cryst. (Me₂CO). Mp 186°. [α]_D +56 (c, 1 in EtOH).

*N*¹-(4-*Methoxy-E-cinnamoyl*), *N*⁴-oxide:

Absouline N-oxide

[112494-54-9]

C₁₇H₂₂N₂O₃ 302.372

From *Hugonia oreogena* and *Hugonia penicillanthemum*. Amorph. [α]_D +13 (c, 0.9 in EtOH).

*N*¹-(4-*Methoxy-Z-cinnamoyl*): *Isoabsouline*

[112513-34-5]

C₁₇H₂₂N₂O₂ 286.373

From *Hugonia oreogena* and *Hugonia penicillanthemum*. Amorph.

*N*¹-(4-*Methoxy-Z-cinnamoyl*), *N*⁴-oxide:

Isoabsouline N-oxide

[112494-55-0]

C₁₇H₂₂N₂O₃ 302.372

From *Hugonia oreogena* and *Hugonia penicillanthemum*. Amorph.

(1*R*,*S*,7*aRS*)-form

(±)-*trans*-form

[141197-03-7]

Amorph. solid.

(1*R*,*S*,7*aSR*)-form

(±)-*cis*-form

[145511-58-6]

Characterised spectroscopically.

[66393-07-5, 66393-06-4]

Neuner-Jehle, N. *et al.*, *Monatsh. Chem.*, 1965, **96**, 321-338 (*Laburnamine*)

Ikhiri, K. *et al.*, *J. Nat. Prod.*, 1987, **50**, 626-

630 (*Absouline, Absouline N-oxide, Isoabsouline, Isoabsouline N-oxide*)

Christine, C. *et al.*, *Tetrahedron*, 2000, **56**,

1837-1850 (*SR-form, RS,RS-form, RS,SR-form, synth, pmr, cmr*)

Giri, N. *et al.*, *J.O.C.*, 2004, **69**, 7303-7308 ((-)-*cis*-form)


Vargas-Sanchez, M. *et al.*, *Org. Lett.*, 2005, **7**, 5861-5864 (*synth*)

Matos, M.N. *et al.*, *Tetrahedron*, 2005, **61**, 1221-1244 (*Laburnamine, synth*)

1-Aminoquinolizidine A-907

Octahydro-2H-quinolizin-1-amine, 9CI

[80220-52-6]



C₉H₁₈N₂ 154.255

(1*S*,9*aS*)-form

*N*¹-*Ac*: 1-Acetamidoquinolizidine. *Epiquinamide*. Alkaloid 196. *Quinolizidine 196* [616235-95-1]

C₁₁H₂₀N₂O 196.292

Alkaloid from the skin extracts of the frog *Epipedobates tricolor*. Nicotinic agonist. [α]_D²⁰ +19 (c, 0.9 in CHCl₃).

Fitch, R.W. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1345-1350 (*isol, pmr, cmr*)

Kanakubo, A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 4648-4651 (*synth*)

Huang, P.-Q. *et al.*, *Org. Lett.*, 2006, **8**, 1435-1438 (*synth*)

Suyama, T.L. *et al.*, *Org. Lett.*, 2006, **8**, 4541-4543 (*synth*)


Tong, S.T. *et al.*, *Tet. Lett.*, 2006, **47**, 5017-5020 (*synth*)

Voituriez, A. *et al.*, *Org. Lett.*, 2007, **9**, 4705-4708 (*synth*)

Wijdeven, M.A. *et al.*, *Org. Lett.*, 2008, **10**, 4001-4003 (*synth, abs config*)

3-Aminosolanidan-23-ol, 9CI A-908

3-Amino-23-hydroxysolanidane



(3 β ,5 α ,22*R*,23*S*,25*R*)-form

C₂₇H₄₆N₂O 414.673

(3 β ,5 α ,22*R*,23*S*,25*R*)-form

Solanogantine

[63785-19-3]

Alkaloid from leaves of *Solanum giganteum* (Solanaceae). Glass. Unusual 2 β H-config., first natural 3-aminosolanidane.

N,*O*-*Di-Ac*: Mp 219-221° dec. [α]_D +27.2 (CHCl₃).

N,N-Di-Me: Mp 111-112°. [α]_D +41.3 (CHCl₃).

(3 β ,5 α ,22S,23S,25R)-form**Solanogantamine**

[72075-33-3]

Alkaloid from *Solanum giganteum* and *Solanum vegum* (major alkaloid) (Solanaceae). Plates (MeOH/MeCN). Mp 180°. [α]_D +35 (CHCl₃).

N,O-Di-Ac:

Needles (Me₂CO aq.). Mp 229-231°. [α]_D +11 (CHCl₃).

N,N-Di-Me: Mp 207-210°. [α]_D +50 (CHCl₃).

(3 α ,5 α ,22S,23S,25R)-form**Isosolanogantamine**

[72075-53-7]

Alkaloid from *Solanum giganteum* (Solanaceae). Flakes (CHCl₃/petrol). Mp 252-254°. [α]_D +31 (CHCl₃).

N,O-Di-Ac:

Cryst. (Me₂CO). Mp 257-259°. [α]_D +35.5 (CHCl₃).

N,N-Di-Me:

Flakes (MeOH). Mp 200-203°. [α]_D +41 (CHCl₃).

(3 β ,5 α ,22S,23S,25S)-form**Solanopubamine**

[97858-53-2]

Alkaloid from the aerial parts of *Solanum pubescens* (Solanaceae). Needles (Me₂CO/MeOH). Mp 263°. [α]_D +30.5 (MeOH).

N-Formyl: **Solanopubamide A**

[105705-84-8]

C₂₈H₄₆N₂O₂ 442.684

Alkaloid from the aerial parts of *Solanum pubescens* (Solanaceae). Needles (MeOH). Mp 209-210°. [α]_D +26.1 (CHCl₃).

N-Ac: **Solanopubamide B**

[105814-55-9]

C₂₉H₄₈N₂O₂ 456.71

Alkaloid from the aerial parts of *Solanum pubescens* (Solanaceae). Needles (MeOH). Mp 255-256°. [α]_D +40 (CHCl₃).

N,O-Di-Ac:

Needles (MeOH). Mp 232°.

Pakrashi, S.C. *et al.*, *Tet. Lett.*, 1977, 645 (*isol, ms, pmr, struct, synth*)

Pakrashi, S.C. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 1109 (*isol, ir, pmr*)


Chakravarty, A.K. *et al.*, *J.C.S. Perkin 1*, 1984, 467 (*config*)

Krishna, G.N. *et al.*, *Phytochemistry*, 1985, **24**, 1369 (*Solanopubamine*)

Kumari, G.N.K. *et al.*, *Phytochemistry*, 1986, **25**, 2003 (*Solanopubamides*)

3-Aminospirostan-6-ol, 9CI

A-909

C₂₇H₄₅NO₃ 431.657**(3 β ,5 α ,6 α ,25R)-form****Isojuripidine**

[23656-00-0]

Alkaloid from *Solanum asterophorum* and *Solanum paniculatum*. Cryst. (MeOH). Mp 204-205° (200°). [α]_D -47 (c, 0.013 in CHCl₃).

O,N-Di-Ac: Mp 153-154°.

Glycoside: **Isojuripine**C₃₃H₅₅NO₈ 593.799

From *Solanum paniculatum* (Solanaceae). Mp 162-171°. [α]_D -20 (Py). β -Glycoside of undetd. struct.

(3 β ,5 α ,6 α ,25S)-form**Juripidine**

[89945-92-6]

Alkaloid from roots of *Solanum hispidum*.

N-Ac:

Cryst. (MeCN). Mp 260-262°. [α]_D -78.12 (c, 0.32 in CHCl₃).

N,O-Di-Ac: Mp 264-266°. [α]_D -63.9 (c, 0.36 in CHCl₃).

Gandolfi, C. *et al.*, *Tet. Lett.*, 1970, **11**, 1677-1680 (*synth*)

Cambiaghi, S. *et al.*, *Ann. Chim. (Rome)*, 1971, **61**, 99; *CA*, **75**, 36397x


(*Isojuripidine*)

Chakravarty, A.K. *et al.*, *Phytochemistry*, 1983, **22**, 2843-2845 (*Juripidine*)

Silva, T.M.S. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 1467-1471 (*Isojuripidine*)

3-Aminospirostan-9-ol, 9CI

A-910

C₂₇H₄₅NO₃ 431.657**(3 β ,5 α ,25R)-form****Isopaniculidine**

[16577-35-8]

Alkaloid from *Solanum paniculatum* (Solanaceae). Mp 202-204°. [α]_D -70 (c, 0.01 in CHCl₃).

N-Ac: Mp 282-284°.

(3 β ,5 α ,25S)-form**Paniculidine**

[16577-34-7]

Alkaloid from *Solanum paniculatum* (Solanaceae). Needles (MeOH). Mp 228-244°. [α]_D -82.2 (c, 1.447 in CHCl₃).

Cambiaghi, S. *et al.*, *Ann. Chim. (Rome)*, 1971, **61**, 99; *CA*, **75**, 36397x (*ir, pmr, isol, struct*)

14-Aminotetradecanoic acid

A-911

14-Aminomyristic acid

[17437-20-6]

H₂NCH₂(CH₂)₁₂COOHC₁₄H₂₉NO₂ 243.389


Constit. of the roots of *Decalepis hamiltonii*. Antioxidant. Mp 178-179°.

Hünig, S. *et al.*, *Chem. Ber.*, 1967, **100**, 3039-3044 (*synth*)

Srivastava, A. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 790-795 (*isol, pmr, cmr*)

2-Amino-5-tetradecen-3-ol

A-912

C₁₄H₂₉NO 227.389**(2S,3R,5Z)-form****Halaminol C**

[389125-68-2]

Isol. from a marine sponge

Haliclona n. sp. Oil. [α]_D +1.9 (c, 0.025 in CH₂Cl₂).

(2S,3S,5E)-form [181523-19-3]

N,O-Di-Ac: [181523-11-5]

C₁₈H₃₃NO₃ 311.464

Isol. from a marine ascidian (*Pseudodistoma* sp.).


Hooper, G.J. *et al.*, *Nat. Prod. Lett.*, 1995, **6**, 31-35 (*isol*)

Clark, R.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1568-1571 (*isol, pmr, cmr*)

Garrido, L. *et al.*, *Tetrahedron*, 2001, **57**, 4579-4588 (*abs config*)

2-Amino-13-tetradecen-3-ol

A-913

C₁₄H₂₉NO 227.389**(2S,3R)-form****Halaminol A**

[389125-56-8]

Isol. from the marine sponge *Haliclona* n. sp. Oil. [α]_D +1.7 (c, 0.04 in CH₂Cl₂). [α]_D -30 (c, 0.003 in MeOH) (as N,O-di-Ac).


Clark, R.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1568-1571 (*Halaminol A, isol, pmr, cmr*)

7-Amino-3,4,4a,10b-tetrahydro-

A-914

dro-2-(hydroxymethyl)-4-methylbenzo[*f*]quinolin-6(5H)-one


[134677-39-7]

C₁₅H₁₈N₂O₂ 258.319

Isol. from *Claviceps purpurea*. Cryst. (Et₂O). Prob. a degradn. prod. of ergot alkaloids.

Deoxy: **7-Amino-3,4,4a,10b-tetrahydro-2,4-dimethylbenzo[*f*]quinolin-6(5H)-one**

[134677-38-6]

C₁₅H₁₈N₂O 242.32Isol. from *Claviceps purpurea*. Cryst. (Me₂CO). Prob. a degradn. prod. of ergot alkaloids.Flieger, M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 390 (isol, pmr, cmr, ms, struct)**2-Amino-6-(1,2,3,4-tetrahydroxybutyl)-4(1H)-pteridinone, 9CI** A-915
2-Amino-4-hydroxy-6-(1,2,3,4-tetrahydroxybutyl)pteridine. *Rs I* [13392-23-9]C₁₀H₁₃N₅O₅ 283.243**(1'R,2'S,3'R)-form***D*-arabino-form

[735-67-1]

[α]_D -97.3 (c, 0.3 in 0.1M NaOH). Dec. slowly above 220°.**(1'R,2'S,3'S)-form***L*-xylo-form


[153829-74-4]

[α]_D -69 (c, 0.4 in 0.1M NaOH).**(1'S,2'S,3'R)-form***D*-ribo-form

[153829-70-0]

Isol. from *Rhodospseudomonas spheroides*.

[735-67-1, 95840-18-9, 153829-73-3, 153829-71-1, 153829-72-2, 95191-28-9]

Petering, H.G. *et al.*, *J.A.C.S.*, 1949, **71**, 3977 (synth)Henseke, G. *et al.*, *Chem. Ber.*, 1956, **89**, 2904 (synth)Taylor, E.C. *et al.*, *J.A.C.S.*, 1976, **98**, 2301 (synth)Seo, N. *et al.*, *Biochim. Biophys. Acta*, 1991, **1074**, 439 (isol)Hariu, T. *et al.*, *Pteridines*, 1993, **4**, 63 (abs config)**16-Amino-2,6,10,14-tetramethyl-5,10,14-hexadecatrien-4-one** A-916

(5E,10E,14E)-form

C₂₀H₃₅NO 305.503**(5E,10E,14E)-form**N-Formyl: *Malonganenone H*


[945458-51-5]

C₂₁H₃₅NO₂ 333.513Isol. from *Euplexaura nuttingi*. Oil.**(5Z,10E,14E)-form**N-Formyl: *Malonganenone C*

[882403-71-6]


C₂₁H₃₅NO₂ 333.513Isol. from *Leptogorgia gilchristi*. Yellow solid. λ_{max} 230 (ε 8800); 276 (ε 1400) (MeOH).Keyzers, R.A. *et al.*, *Tetrahedron*, 2006, **62**, 2200-2206 (*Malonganenone C*)Sorek, H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1104-1109 (*Malonganenone H*)**1-Amino-4,12-tridecadien-2-ol, 9CI** A-917C₁₃H₂₅NO 211.347**(±)-(E)-form** [181523-21-7]

N,O-Di-Ac: [181523-13-7]

C₁₇H₂₉NO₃ 295.421Isol. from a marine ascidian (*Pseudodistoma* sp.).Hooper, G.J. *et al.*, *Nat. Prod. Lett.*, 1995, **6**, 31-35 (isol)**1-Amino-3-tridecen-2-ol** A-918C₁₃H₂₇NO 213.362**(2R,3E)-form** [150922-61-5]Isol. from the ascidian *Didemnum* sp.Searle, P.A. *et al.*, *J.O.C.*, 1993, **58**, 7578-7580 (isol)**1-Amino-5-tridecen-2-ol** A-919C₁₃H₂₇NO 213.362**(2R,5E)-form** [150922-60-4]Isol. from the ascidian *Didemnum* sp.Antifungal agent. [α]_D +1.9 (c, 0.4 in MeOH) (as trifluoroacetate salt).Searle, P.A. *et al.*, *J.O.C.*, 1993, **58**, 7578-7580 (isol, ir, pmr, abs config)**5-(12-Aminotridecyl)hexahydro-2,1-b[oxazole]** A-920


Hexahydro-α-methylpyrrolo[2,1-b]oxazole-5-dodecanamine, 9CI

[192213-01-7]

C₁₉H₃₈N₂O 310.522Alkaloid from the ladybirds *Epilachna boraelis* and *Epilachna varivestis*.Radford, P. *et al.*, *J. Nat. Prod.*, 1997, **60**, 755-759 (isol, pmr, cmr, ms)**2-(12-Aminotridecyl)pyrrolidine** A-921

α-Methyl-2-pyrrolidinedodecanamine,


9CI. 13-(2-Pyrrolidinyl)-2-tridecylamine

C₁₇H₃₆N₂ 268.485**(2S,12'R)-form** [151751-64-3]Alkaloid from whole body extracts of the Mexican bean beetle *Epilachna varivestis*.[α]_D²² +9.3 (c, 0.15 in CDCl₃).N¹-(2-Hydroxyethyl)-2-(12-Aminotridecyl)-1-pyrrolidineethanol, 9CI. 1-(2-Hydroxyethyl)-2-(12-aminotridecyl)-pyrrolidine

[151751-65-4]

C₁₉H₄₀N₂O 312.538Alkaloid present in eggs, larvae, pupae and adults of *Epilachna varivestis*. [α]_D²² +38.8 (c, 0.18 in CDCl₃).Attygalle, A.B. *et al.*, *Tetrahedron*, 1993, **49**, 9333 (isol, ir, pmr, cmr, ms, struct)Shi, X. *et al.*, *Tetrahedron*, 1996, **52**, 6859 (isol, pmr, cmr, ms, synth, abs config)Shi, X. *et al.*, *Tet. Lett.*, 1997, **38**, 6479 (synth, abs config)**2-Amino-3,4,5-trihydroxybenzoic acid, 9CI** A-922

3,4,5-Trihydroxyanthranilic acid, 9CI

C₇H₇NO₅ 185.136

5-Me ether: 2-Amino-3,4-dihydroxy-5-methoxybenzoic acid

[1035536-86-7]

C₈H₉NO₅ 199.163Alkaloid from the leaves of *Cyclocaryapallivus*. Needles. Mp 186-188°. λ_{max} 290 (ε 3450) (CHCl₃).

Tri-Me ether: 2-Amino-3,4,5-trimethoxybenzoic acid

[61948-85-4]


C₁₀H₁₃NO₅ 227.216

Mp 138-140°.

Tri-Me ether, Me ester: [5035-82-5]

C₁₁H₁₅NO₅ 241.243Mp 44-45°. Bp_{0.1} 127-140°.Hammen, P.D. *et al.*, *J. Het. Chem.*, 1987, **24**, 1701-1704Li, J. *et al.*, *Planta Med.*, 2008, **74**, 287-289 (5-Me ether, isol)**2-Amino-6-(1,2,3-trihydroxypropyl)-4(1H)-pteridinone, 9CI** A-923

[670-65-5]

C₉H₁₁N₅O₄ 253.217**(1'R,2'R)-form***D*-threo-form. *D*-Monapterin. *Umanapterin*

[10162-32-0]

Isol. from *Tetrahymena pyriformis* and from the urine of cancer patients. [α]_D²⁵ -92 (c, 0.3 in 0.1M HCl).5,6R,7,8-Tetrahydro: 5,6,7,8-Tetrahydro-*D*-monapterin

[99630-31-6]

C₉H₁₅N₅O₄ 257.249Isol. from *Tetrahymena pyriformis*.**(1'S,2'S)-form***L*-threo-form. **L-Monapterin**

[2277-42-1]

Constit. of human blood and urine.

Yellow cryst. [α]_D²⁵ +97 (c, 0.3 in 0.1M HCl).

2-N,N-Di-Me: 2-Dimethylamino-6-

(1,2,3-trihydroxypropyl)-4(3H)-pteridinone. **Euglenapterin**

[73789-39-6]

[73789-45-4]

C₁₁H₁₅N₅O₄ 281.271Found in the alga *Euglena gracilis*.Yellow cryst. (H₂O). Mp 200°. Yellow fluor. in soln.

2-N,N-Di-Me, 3'-phosphate:

C₁₁H₁₆N₅O₇P 361.251Found in the alga *Euglena gracilis*.

Yellow fluor. in soln.

2-N,N-Di-Me, 2',3'-cyclic phosphate:

C₁₁H₁₄N₅O₆P 343.235Found in the alga *Euglena gracilis*.

Yellow fluor. in soln.

5,6R,7,8-Tetrahydro: 5,6,7,8-Tetrahydro-**L-monapterin**

[348622-74-2]

C₉H₁₅N₅O₄ 257.249Isol. from *Escherichia coli*. Needles (CHCl₃) (as hexa-Ac). Mp 139.5-141.5° (hexa-Ac).**(1'S,2'R)-form***D*-erythro-form. **Neopterin**

[2009-64-5]

Found in human blood and urine. First isol. from the pupae of bees. Precursor in biosynth. of 2-Amino-6-(1,2-dihydroxypropyl)-4(1H)-pteridinone, A-749. [α]_D²⁵ +45 (c, 0.3 in 0.1M HCl).3'-O- β -D-Glucopyranoside: **Solfapterin**.**Neopterin 3-O- β -D-glucoside**

[114312-02-6]

C₁₅H₂₁N₅O₉ 415.359Isol. from *Sulfolobus solfataricus*.**(1'R,2'S)-form***L*-erythro-form. **Bufochrome**

[2277-43-2]

Found in human blood and urine. Also isol. from toad skins. Growth factor for the protozoan *Crithidia fasciculata*. [α]_D²⁵ -44 (c, 0.3 in 0.1M HCl). λ_{\max} 275 (ε 14600); 348 (ε 6180) (H₂O). λ_{\max} 255 (ε 24000); 365 (ε 7050) (0.1M NaOH). λ_{\max} 248 (ε 11600); 321 (ε 8020) (0.1M HCl). [39923-31-4, 87174-05-8, 81873-20-3]*Aldrich Library of 13C and 1H FT NMR**Spectra*, 1992, **3**, 480C (nmr)Rembold, H. et al., *Chem. Ber.*, 1963, **96**, 1406

(struct, synth)

Andrews, K.J.M. et al., *Chem. Comm.*, 1968,

120

v. Viscontini, M. et al., *Helv. Chim. Acta*, 1968,**51**, 1495 (synth, pmr)Fukushima, T. et al., *J. Biol. Chem.*, 1972, **247**,

4549 (isol)

Elstner, E. et al., *Arch. Biochem. Biophys.*,1976, **173**, 614 (*Euglenapterin*)Schircks, B. et al., *Helv. Chim. Acta*, 1976, **59**,

248 (nmr)

Sugimoto, T. et al., *Bull. Chem. Soc. Jpn.*,1980, **53**, 2344 (synth)Fukushima, T. et al., *Methods Enzymol.*, 1980, **66**, 508 (rev, isol)Jacobi, P.A. et al., *J.O.C.*, 1981, **46**, 5416(*Euglenapterin*, synth)Sun, J.H. et al., *Diss. Abstr. Int.*, **B**, 1983, **43**,

3990 (synth)

Böhme, M. et al., *Annalen*, 1986, 1705(*Euglenapterin*, isol, cryst struct, uv, cd, pmr,


synth)

Zeitler, H.J. et al., *Methods Enzymol.*, 1986,**122**, 273 (rev, detn)Huber, C. et al., *Dtsch. Med. Wochenschr.*,1987, **112**, 107 (rev)Lin, X. et al., *J. Bacteriol.*, 1988, **170**, 1396(*Solfapterin*)Klein, R. et al., *Biochimie*, 1991, **73**, 1281(*D-Monapterin*)Ogiwara, S. et al., *Biol. Chem. Hoppe-Seyler*,1992, **373**, 1061 (isol)Ogiwara, S. et al., *Tet. Lett.*, 1992, **33**, 1341(*Umanapterin*)Sugimoto, T. et al., *Helv. Chim. Acta*, 2001, **84**,918-927 (*Tetrahydro-D-monapterin*)Ikemoto, K. et al., *Biol. Chem.*, 2002, **383**,325-330 (*Tetrahydro-L-monapterin*)Chen, N. et al., *Heterocycles*, 2005, **65**, 2917-

2924 (acetates, cd, abs config)

2-Amino-7-(1,2,3-trihydroxypropyl)-4(1H)-pteridinone A-924

7-(1,2,3-Trihydroxypropyl)pterin

C₉H₁₁N₅O₄ 253.217

Abnormal pterin isol. from urine of patients with atypical phenylketonuria (abs. config. of nat. metab. not yet known).

(1'R,2'S)-form*L*-erythro-form[α]_D -13.3'-Deoxy: **Primapterin**. 7-Biopterin

[2636-52-4]

C₉H₁₁N₅O₃ 237.218Occurs in urine of patients with atypical phenylketonuria. Also in patients with hyperphenylalaninemia. [α]_D²⁵ -37.4 (c, 0.1 in 0.1M HCl).**(1'S,2'R)-form***D*-erythro-form. **Anapterin**. 7-Neopterin

[126779-19-9]

Occurs in urine of patients with hyperphenylalaninemia. Microcryst.

(H₂O). [α]_D²² +19.5 (c, 0.4 in

1M HCl). CA erroneously names

Anapterin as

7-(12-dihydroxypropyl)pterin.

Rembold, H. et al., *Chem. Ber.*, 1963, **96**, 1395-

1405 (7-Biopterin, synth, uv)

Curtis, H.-C. et al., *Biochem. Biophys. Res.**Commun.*, 1988, **153**, 715-721 (*Anapterin*,*Primapterin*, isol)Viscontini, M. et al., *Helv. Chim. Acta*, 1990,**73**, 337-345 (*Primapterin*, *Anapterin*, synth,


cd, pmr, bibl)

Curtius, H.-C. et al., *J. Biol. Chem.*, 1990,**265**, 3923-3930 (7-Neopterin, 7-Biopterin,

isol)

2-Amino-4,6,7-trihydroxypteridine A-9252-Amino-5,8-dihydro-4,6,7(1H)-pteridinetrione, 9CI. **Leucopterin**

[492-11-5]

C₆H₅N₅O₃ 195.137Found in the wings of many butterflies, e.g. *Pieris brassicae*. Cryst. Blue fluor. in alkaline soln. Forms yellow salts.Wieland, Th. et al., *Annalen*, 1940, **544**, 172

(synth)

Pfleiderer, W. et al., *Chem. Ber.*, 1957, **90**,2631; 1961, **94**, 118 (synth, struct)Mueller, G. et al., *Helv. Chim. Acta*, 1973, **56**,

2680 (nmr)

Pfleiderer, W. et al., *J. Het. Chem.*, 1992, **29**,


583 (rev)

4-Amino-N,N,N-tris(3-aminopropyl)-1-butanaminium, 9CI A-926N⁺,N⁺-Bis(3-aminopropyl)spermidine

[143085-76-1]


(H₂NCH₂CH₂CH₂)₃N⁺(CH₂)₃CH₂NH₂C₁₃H₃₄N₅[⊕] 260.445Isol. from the thermophilic bacteria *Bacillus schlegelii*, *Thermoleophilum album*, *Thermoleophilum minutum*, *Hydrogenobacter thermophilus* and *Hydrogenobacter halophilus*. Counterion not specified.Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747 (isol)**5-(10-Aminoundecyl)hexahydroppyrrrolo[2,1-b]oxazole** A-927Hexahydro- α -methylpyrrrolo[2,1-b]oxazole-5-decanamine, 9CI

[192213-02-8]

C₁₇H₃₄N₂O 282.468Alkaloid from the ladybird *Epilachna borealis*.Radford, P. et al., *J. Nat. Prod.*, 1997, **60**, 755-759 (isol, ms)**Aminozooanemonin** A-928

2,3-Dihydro-2-imino-1,3-dimethyl-1H-imidazole-4-acetic acid, 9CI

[213749-79-2]



C₇H₁₁N₃O₂ 169.183

Prototropic guanidine system showing zwitterion formation. Shown here and in the CAS name as the neutral species. Alkaloid from the sponge *Agelas dispar*. Amorph. solid. λ_{\max} 226 (ϵ 3500); 262 (ϵ 1500) (MeOH).

Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1171-1173 (*isol, uv, ir, pmr, cmr*)

Ammocalline**A-929**

[1354-30-9]


C₁₉H₂₂N₂ 278.396

Struct. unknown. Alkaloid from roots of *Catharanthus roseus* (Apocynaceae). Needles (Me₂CO). Mp 335° dec. pK_a 7.3 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (*isol, ir, uv*)

Ammodendrine**A-930**

1-Acetyl-1,2,3,4-tetrahydro-5-(2-piperidinyl)pyridine, 9CI. *1'-Acetyl-1,1',2,3,4,4',5,5',6,6'-decahydro-2,3'-bipyridine*. *N-Acetyl-Δ²-tetrahydroanabasine*. *Sphaerocarpine* [27542-15-0]

C₁₂H₂₀N₂O 208.303

► Exp. teratogen.

(R)-form*Isoammodendrine*

[494-15-5]

Alkaloid from *Genista sphaerocarpa*, *Ammodendron conollyi*, *Lupinus formosus* and other plant spp. Cryst. or oil. Mp 43-46°. $[\alpha]_D^{24}$ +5.4 (c, 1.3 in MeOH). Many isolates with higher opt rotations were probably contaminated.

► UR9150000

Perchlorate:Cryst. (H₂O or Me₂CO). Mp 210-211°.*N-Formyl: N'-Formylammodendrine*

[53508-17-1]

C₁₃H₂₀N₂O₂ 236.313

Alkaloid from *Dichilus strictus* and *Spartidium saharae*.

N-Me: N'-Methylammodendrine

[52196-10-8]

C₁₃H₂₂N₂O 222.33

Alkaloid in *Lupinus formosus* (Fabaceae). Oil (natural), cryst. (petrol)-synthetic. Mp 65-66°. $[\alpha]_D^{23}$ +62.4 (c, 0.5 in MeOH).

(±)-form [20824-32-2]

Alkaloid from *Ammodendron conollyi*, other *Ammodendron* spp. *Sophora franchetiana* and *Coelidium fourcadei* (Fabaceae). Mp 50-60°.

Monohydrate: Mp 73-74°.*Hydroiodide*:

Cryst. (EtOH). Mp 221-222°.

N-Me:

Cryst. (petrol). Mp 65-66°.

N-Me; hydroiodide: Mp 184-185°.**(ξ)-form**

Alkaloid from *Ammodendron karelinii*, *Ammodendron longiracemosum*, *Ammodendron lehmanni*, *Genista hystrix*, *Lupinus mutabilis*, *Castilleja miniata*, *Ammopiptanthus mongolicus*, *Thermopsis lupinoides* and other plants (opt. rotn. of these isolates mostly unreported) (Fabaceae).

Orekhov, A.P. *et al.*, *Ber.*, 1935, **68**, 1807 (*isol*)

Orekhov, A.P. *et al.*, *Bull. Soc. Chim. Fr.*, 1938,

5, 29 (*struct*)

Dominguez, J. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1956, **43**; *C.A.*, **51**, 1212h

(struct, resoln)

Ribas-Marques, I. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1962, **57**, 781; *C.A.*, **58**, 5738c

(abs config)

Schöpf, C. *et al.*, *Annalen*, 1964, **674**, 87

(synth)

Arndt, R.R. *et al.*, *J. S. Afr. Chem. Inst.*, 1968,

21, 54; *C.A.*, **70**, 4342f (*isol, pmr*)

Fitch, W.L. *et al.*, *J.A.C.S.*, 1974, **96**, 4917 (*ms*)

Fitch, W.L. *et al.*, *J.O.C.*, 1974, **39**, 2974 (*isol, deriv*)

Kushmuradov, Yu.K. *et al.*, *Khim. Prir. Soedin.*, 1977, 717; 1979, 871; *Chem. Nat. Compd. (Engl. Transl.)*, 604; 780

(isol)

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1982, 661; *Chem. Nat. Compd. (Engl. Transl.)*, 631 (*cryst struct*)

Van Wyk, B.E. *et al.*, *S. Afr. J. Bot.*, 1989, **55**,

523 (*N'-Formylammodendrine*)

Lee, S.T. *et al.*, *J. Nat. Prod.*, 2005, **68**, 681-685

*(isol, resoln, tox)***Ammorosine****A-931**


[1354-31-0]

Struct. unknown. Alkaloid from roots of *Catharanthus roseus* (Apocynaceae). Blades (MeOH). Mp 221-225°. pK_a 7.3 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**,

141-153 (*isol, ir, uv*)**Amphisterin C₁****A-932**

[330680-37-0]

R = -(CH₂)₃CH=CH(CH₂)₉CH₃(E-)

Relative Configuration

C₂₇H₄₅NO₆ 479.656

Isol. from *Plakortis quasiamphiaster*. Oil. $[\alpha]_D^{20}$ -19.3 (c, 0.14 in CHCl₃).

4'',5''-Dihydro: Amphisterin C₃

[330680-39-2]

C₂₇H₄₇NO₆ 481.671Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -8 (c, 0.4 in CHCl₃).*3-Epimer: Amphisterin D₁*

[330680-41-6]

C₂₇H₄₅NO₆ 479.656Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -16 (c, 0.06 in CHCl₃).*3,4,5-Triepimer: Amphisterin E₁*

[330680-44-9]

C₂₇H₄₅NO₆ 479.656Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -3.8 (c, 0.11 in CHCl₃).

Zampella, A. *et al.*, *Tetrahedron*, 2001, **57**, 257-263 (*Amphisterins C₁, C₃, D₁, E₁*)

Amphisterin C₂**A-933**

[330680-38-1]

As Amphisterin C₁, A-932 withR = -(CH₂)₅CH=CH(CH₂)₉CH₃(E-)C₂₉H₄₉NO₆ 507.709Isol. from *Plakortis quasiamphiaster*. Oil. $[\alpha]_D^{20}$ -12.3 (c, 0.65 in CHCl₃).*6'',7''-Dihydro: Amphisterin C₄*

[330680-40-5]

C₂₉H₅₁NO₆ 509.725Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -11.5 (c, 0.13 in CHCl₃).*3-Epimer: Amphisterin D₂*

[330680-42-7]

C₂₉H₄₉NO₆ 507.709Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -2.9 (c, 0.24 in CHCl₃).*3-Epimer, 6'',7''-dihydro: Amphisterin D₃*

[330680-43-8]


C₂₉H₅₁NO₆ 509.725Isol. from *Plakortis quasiamphiaster*.Oil. $[\alpha]_D^{20}$ -1.5 (c, 0.1 in CHCl₃).

Zampella, A. *et al.*, *Tetrahedron*, 2001, **57**, 257-263 (*Amphisterins C₂, C₄, D₂, D₃*)

Amphibine I**A-934**

L-Valyl-N-[1-(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)ethyl]glycinamide, 9CI

[53098-67-2]

C₂₁H₃₄N₄O₄ 406.524

Consists of mixt. of diastereoisomer shown with 1,9-diepimer. Alkaloid from *Zizyphus amphibia* (Rhamnaceae). Pale yellow foam. $[\alpha]_D^{20}$ -50 (c, 0.6 in C₆H₆).

Hydrochloride (1:2):Cryst. (CHCl₃/petrol). Mp 175° dec. $[\alpha]_D^{20}$ -3.4 (c, 0.24 in MeOH).*N-Ac*:


Needles (CH₂Cl₂). Mp 188-190°. $[\alpha]_D^{20}$ -20 (c, 0.1 in EtOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 1329; 1975, **108**, 2247 (*ir, uv, ms, pmr, isol, struct, synth, config, cd*)

Koyama, J. *et al.*, *Heterocycles*, 1978, **9**, 443
(*synth, cd*)

Amphibine A
[36535-97-4]

A-935



Absolute
Configuration

C₃₃H₄₃N₅O₄ 573.734

Prob. identical with Discarine A and possibly also with Waltherine B. Alkaloid from the bark of *Zizyphus amphibia* and *Zizyphus spinachristi* and from the root bark of *Zizyphus nummularia* (Rhamnaceae). Needles (CH₂Cl₂/MeOH). Mp 237-239° dec. [α]_D²⁰ -310 (c, 0.021 in MeOH).

Stereoisomer (1) (?) : Waltherine B

[243464-60-0]

C₃₃H₄₃N₅O₄ 573.734

Alkaloid from the root bark of *Waltheria douradinha*. Needles (CHCl₃/MeOH). Mp 242-243°. [α]_D²⁰ -201.8 (c, 0.21 in MeOH).

Stereoisomer (2) (?) : Discarine A

[36211-12-8]

C₃₃H₄₃N₅O₄ 573.734

Alkaloid from the roots of *Discaria longispina* (Rhamnaceae). Cryst. (CH₂Cl₂/Et₂O). Mp 229-231°. [α]_D -282 (c, ca. 0.05 in CHCl₃).

González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915-916 (*Discarine A, pmr, stereochem*)

Mascaretti, O.A. *et al.*, *Phytochemistry*, 1972, **11**, 1133 (*Discarine A, isol, struct*)

Tschesche, R. *et al.*, *Tet. Lett.*, 1972, 865-868 (*Amphibine A, ms, struct*)

Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 1633 (*Amphibine A, isol*)


Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (*Amphibine A, isol*)

Pais, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869-1872 (*Discarine A, cmr*)

Morel, A.F. *et al.*, *Phytochemistry*, 1999, **51**, 473-477 (*Waltherine B, isol*)

Amphibine B
[38541-74-1]

A-936



R¹ = R³ = CH₂Ph
R² = CH(CH₃)CH₂CH₃

C₃₉H₄₇N₅O₅ 665.831

Stereochem. not yet determined. Alkaloid from the bark of *Zizyphus amphibia* and *Zizyphus rugosa*, and from *Zizyphus oenoplia*, *Zizyphus mauritiana* (Rhamnaceae). Amorph. [α]_D²⁰ -181 (c, 0.08 in MeOH). λ_{max} 210 (end) ; 260 (sh)

(MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1972, **105**, 3094-3105 (*Zizyphus amphibia constit, struct*)

Tschesche, R. *et al.*, *Annalen*, 1974, 1694-1701 (*Zizyphus mauritiana constit*)

Singh, A. *et al.*, *J. Indian Chem. Soc.*, 2008, **85**, 658-659 (*Zizyphus rugosa constit*)

Amphibine C

A-937

[38496-01-4]

As Amphibine B, A-936 with

R¹ = CH₂CH(CH₃)₂, R² = CH(CH₃)CH₂CH₃, R³ = CH₂Ph

C₃₆H₄₉N₅O₅ 631.814

Alkaloid from the bark of *Zizyphus amphibia* (Rhamnaceae). Amorph. [α]_D²⁰ -224 (c, 0.075 in MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1972, **105**, 3094 (*isol, uv, cd, ir, pmr, ms, struct*)

Amphibine D

A-938

[38496-02-5]

As Amphibine B, A-936 with

R¹ = CH₂Ph, R² = R³ = CH(CH₃)CH₂CH₃

C₃₆H₄₉N₅O₅ 631.814

Alkaloid from the bark of *Zizyphus amphibia*, *Zizyphus mauritiana*, *Zizyphus rugosa* and *Zizyphus juazeiro* (Rhamnaceae). Amorph. [α]_D²⁰ -203 (c, 0.09 in MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1972, **105**, 3094-3105 (*isol, uv, cd, ir, pmr, ms, struct*)


Hindenlang, D.M. *et al.*, *Annalen*, 1980, 447-450 (*cmr*)

Tschesche, R. *et al.*, *Pharmazie*, 1981, **36**, 511 (*isol, uv, ir, pmr, ms*)

Amphibine E

A-939

[38541-75-2]



C₃₈H₅₀N₆O₅ 670.85

Alkaloid from the bark of *Zizyphus amphibia*, *Zizyphus mauritiana* and *Zizyphus spinachristi* (Rhamnaceae). Amorph. [α]_D²⁰ -175 (c, 0.14 in MeOH).

N-De-Me: Mauritine J

[177715-71-8]

C₃₇H₄₈N₆O₅ 656.823

Alkaloid from root bark of *Zizyphus mauritiana*. Amorph. [α]_D -175.9 (c, 1.0 in MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1972, **105**, 3094-3105 (*isol, uv, cd, ir, pmr, ms, struct*)

Tschesche, R. *et al.*, *Annalen*, 1974, 1694-1702 (*isol*)

Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 1633 (*isol*)


Hindenlang, D.M. *et al.*, *Annalen*, 1980, 447-450 (*cmr*)

Jossang, A. *et al.*, *Phytochemistry*, 1996, **42**, 565-567 (*Mauritine J*)

Amphibine F

A-940

[52617-26-2]



R¹ = H, R² = -CH₂Ph
R³ = -CH(CH₃)CH₂CH₃

C₂₉H₃₆N₄O₄ 504.628

Alkaloid from the bark of *Zizyphus amphibia*, *Zizyphus mauritiana* and *Zizyphus spinachristi* (Rhamnaceae). Amorph. [α]_D²⁰ -171 (c, 0.26 in CHCl₃).

Dihydro, N-Me:

Needles (Me₂CO/cyclohexane). Mp 152-156°. [α]_D²⁰ -119 (c, 0.05 in CHCl₃).

Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 686 (*isol, uv, ir, pmr, ms, struct*)

Tschesche, R. *et al.*, *Annalen*, 1974, 1694 (*isol*)
Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 1633 (*isol*)

Amphibine G

A-941

[52617-27-3]

As Amphibine F, A-940 with

R¹ = Me, R² = -CH₂CH(CH₃)₂, R³ = 3-Indolyl-CH₂-

C₃₂H₃₉N₅O₄ 557.691

Alkaloid from the bark of *Zizyphus amphibia* (Rhamnaceae). Needles (CCl₄). Mp 182-185°. [α]_D²⁰ -218 (c, 0.24 in CHCl₃).

Dihydro:


Amorph. [α]_D²⁰ +119 (c, 0.2 in CHCl₃).

Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 686 (*isol, uv, cd, ir, pmr, ms, struct*)

Amphibine H

A-942

[52659-55-9]



C₃₃H₄₃N₅O₆ 605.733

Alkaloid from the bark or root bark of *Zizyphus amphibia*, *Zizyphus nummularia*, *Zizyphus jujuba*, *Zizyphus spina-christi* and *Zizyphus xylopyra*. Shows antibacterial activity. Cryst. (CHCl₃/petrol). Mp 205°. [α]_D²⁰ -570 (c, 0.12 in MeOH).

N-De-Me: Nummularine B. N-Demethylamphibine H. *Daechuine S27* [53947-96-9]

C₃₂H₄₁N₃O₆ 591.706

Alkaloid from the root bark or stem bark of *Zizyphus nummularia*, *Zizyphus jujuba*, *Zizyphus sativa* and *Zizyphus xylopyra* (Rhamnaceae). Shows antibacterial activity. Needles (MeOH). Mp 230-231°. [α]_D²⁰ -390 (c, 0.2 in CHCl₃).

N-De-Me, N-formyl: Nummularine T [99694-95-8]

C₃₃H₄₁N₃O₇ 619.716

Alkaloid from bark of *Zizyphus nummularia* (Rhamnaceae). Granules (MeOH). Mp 188-190°.

Dihydro: Shows antibacterial activity. Needles (CH₂Cl₂/Me₂CO). Mp 281°. [α]_D²⁰ -381 (c, 0.15 in MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 686; 3180 (*isol, uv, ir, pmr, ms, struct*)

Tschesche, R. *et al.*, *Phytochemistry*, 1976, **15**, 541; 1979, **18**, 702 (*isol*)


Pandey, V.B. *et al.*, *Phytochemistry*, 1984, **23**, 2118 (*isol*)

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443 (*isol, Nummularine B*)

Pandey, V.B. *et al.*, *Planta Med.*, 1990, **56**, 649 (*activity*)

Singh, B. *et al.*, *Phytochemistry*, 1995, **38**, 271 (*Nummularine T*)

Amphikuemin **A-943**
[105870-54-0]




C₁₆H₂₆N₃O⁺ 308.4

Isol. from the sea anemone *Radianthus kuekenhali*. Induces symbiosis between sea anemone and anemone fish.

Murata, M. *et al.*, *Science (Washington, D.C.)*, 1986, **234**, 585-587 (*isol, uv, pmr, struct*)

Konno, K. *et al.*, *Heterocycles*, 1990, **30**, 247-251 (*synth*)

Amphimedine **A-944**
10-Methyl-8H-benzo[b]pyrido[4,3,2-de][1,8]phenanthroline-8,11(10H)-dione, 9CI
[86047-14-5]



C₁₉H₁₁N₃O₂ 313.315

Alkaloid from an *Amphimedon* sp. of Pacific sponge and *Xestospongia carbo-naria*. Cytotoxic; topoisomerase II inhibitor. Yellow solid. Fairly sol. MeOH; poorly sol. butanol, hexane. Mp 360°. λ_{\max} 210 (ε 19700); 233 (ε 39400); 281 (ε 9100); 341 (ε 6060) (EtOH) (Derep). λ_{\max} 245 (ε 38000); 281 (ε 10000); 340 (ε 7000) (MeOH) (Berdy). λ_{\max} 235 (ε 38000); 281 (ε 10000); 340 (ε 7000) (EtOH) (Berdy).

Schmitz, F.J. *et al.*, *J.A.C.S.*, 1983, **105**, 4835 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kubo, A. *et al.*, *Heterocycles*, 1988, **27**, 2095 (*synth, uv, ir, pmr, ms*)


Echavarren, A.M. *et al.*, *J.A.C.S.*, 1988, **110**, 4051 (*synth*)

Prager, R.H. *et al.*, *Aust. J. Chem.*, 1991, **44**, 277 (*synth, uv, ir*)

Guillier, F. *et al.*, *J.O.C.*, 1995, **60**, 292 (*synth*)

Bracher, F. *et al.*, *Annalen*, 1996, 115 (*synth*)
Nakahara, S. *et al.*, *Heterocycles*, 1996, **43**, 2113 (*synth*)

Amphimedoside A **A-945**
[915721-96-9]



m = 4, n = 10

C₂₈H₄₆N₂O₆ 506.681

Similar to Hachijodines, H-3. Alkaloid from *Amphimedon* sp. Cytotoxic.

Takekawa, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1503-1505 (*isol, pmr, cmr, ms*)

Amphimedoside B **A-946**
[915722-03-1]

As Amphimedoside A, A-945 with m = 4, n = 8

C₂₆H₄₂N₂O₆ 478.628

Alkaloid from *Amphimedon* sp. Cytotoxic.

Takekawa, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1503-1505 (*isol, pmr, cmr, ms*)

Amphimedoside C **A-947**
[915722-04-2]


As Amphimedoside A, A-945 with m = 2, n = 10

C₂₆H₄₂N₂O₆ 478.628

Alkaloid from *Amphimedon* sp. Cytotoxic.

Takekawa, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1503-1505 (*isol, pmr, cmr, ms*)

Amphimedoside E **A-948**
[915722-06-4]




C₂₆H₄₄N₂O₆ 480.643

Alkaloid from *Amphimedon* sp. Cytotoxic.

Takekawa, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1503-1505 (*isol, pmr, cmr, ms*)

Amphistin **A-949**
[199783-52-3]




C₁₃H₂₁N₃O₆ 343.339

Prod. by an actinomycete strain KP-3052. Melanogenesis inhibitor. Powder. Mp 260° (dec.). [α]_D²⁵ -9 (c, 0.6 in H₂O). λ_{\max} 220 (ε 7500) (H₂O).

Arai, N. *et al.*, *J. Antibiot.*, 1997, **50**, 808-814 (*isol, uv, ir, pmr, cmr, N-15 nmr*)

Amphitoxin **A-950**
[165724-12-9]




X = or (ratio about 1:1)

n = 5 (mean value)

Polymeric alkaloid from the Caribbean sponge *Amphimedon compressa*. Exhibits antifeedant activity. λ_{\max} 266 (H₂O).

Albrizio, S. *et al.*, *J. Nat. Prod.*, 1995, **58**, 647 (*isol, uv, ir, pmr, cmr*)

Ampullicin **A-951**



C₁₅H₁₇NO₃ 259.304

Metab. of an *Ampulliferina*-like fungal sp. Needles + 1/2 H₂O (Me₂CO/C₆H₆/hexane). Mp 197-198°. [α]_D²⁰ -32 (c, 0.5 in MeOH). λ_{\max} 281 (ε 14700) (EtOH) (Derep). λ_{\max} 281 (ε 14700) (EtOH) (Berdy).

5,6-Dihydro: Dihydroampullicin

[149182-93-4]

C₁₅H₁₉NO₃ 261.32

Metab. of an *Ampulliferina*-like fungus. Plant growth regulator. Needles. Sol. MeOH, Me₂CO, EtOAc. Mp 180-181°. [α]_D²⁰ +36 (c, 0.25 in MeOH). λ_{\max} 244 (ε 2490); 248 (ε 3000); 254 (ε 3350); 260 (ε 2400) (EtOH). λ_{\max} 244 (ε 2940); 248 (ε 3000); 254 (ε 3350); 260 (ε 2400) (EtOH) (Berdy).

5Z-Isomer: Isoampullicin

[127943-13-9]

C₁₅H₁₇NO₃ 259.304

From an *Ampulliferina*-like fungal sp. Mp 202-203°. [α]_D²⁰ +110 (c, 0.5 in MeOH). λ_{\max} 281 (ε 14700) (EtOH) (Derep). λ_{\max} 284 (ε 20500) (EtOH) (Berdy).

- Kimura, Y. *et al.*, *Agric. Biol. Chem.*, 1990, **54**, 813 (*isol*, *pmr*, *cmr*)
 Kimura, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 687-688 (*Dihydroampullicin*)
 Rico, R. *et al.*, *Tet. Lett.*, 1995, **36**, 7889-7892; 1996, **37**, 5809-5812 (*synth*)
 Marcos, I. *et al.*, *Tet. Lett.*, 2000, **41**, 8451-8455 (*synth*, *Dihydroampullicin*)
 Bermejo, F.A. *et al.*, *J.O.C.*, 2001, **66**, 8287-8292 (*synth*, *abs config*)

Amsoniaefoline A-952


Struct. unknown

$C_{25}H_{32}N_2O_5$ 440.538
 Alkaloid from *Rauwolfia amsoniaefolia* (Apocynaceae). Mp 220-223°.

Gomez, R. *et al.*, *J. Philipp. Pharm. Assoc.*, 1957, **44**, 101; 127

Amosinine A-953

Methyl 19-hydroxy-19-methylxayohimban-16-carboxylate, 9CI. 19-Hydroxy-16,17-dihydromayumbine [136092-57-4]



$C_{21}H_{26}N_2O_4$ 370.447
 Alkaloid from *Amsonia sinensis* (Apocynaceae). Cryst. Mp 221-222°. $[\alpha]_D^{25}$ -51.6 (c, 0.05 in Me_2CO).

16,19-Diepimer: Diangoutengjian I


[344927-27-1]
 $C_{21}H_{26}N_2O_4$ 370.447
 Alkaloid from *Uncaria yunnanensis*. Cryst. (MeOH). Mp 190-191°.

Liu, H.-M. *et al.*, *Planta Med.*, 1991, **57**, 566 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Tao, Z.-Y. *et al.*, *Yaoxue Xuebao*, 2001, **36**, 120-122 (*Diangoutengjian I*)

Amurensine† A-954

10,11-Dihydro-7-methoxy-13-methyl-11,5-(iminomethano)-5H-benzo[4,5]cyclohepta[1,2-f]-1,3-benzodioxol-8-ol, 9CI. *Xanthopetaline* [10481-92-2]



Absolute configuration

$C_{19}H_{19}NO_4$ 325.363
 Alkaloid from *Papaver alpinum*, *Papaver alpinum* var. *tatricum*, var. *burseri*, var. *kernerii*, *Papaver pseudocanescens*, *Papaver nudicaule* var. *amurense*, var. *xanthopetalum*, var. *leiocarpum*, var. *rubroaurantiacum*, *Papaver pyrenaicum*, *Papaver tatricum*, *Papaver suaveolens* and *Papaver anomalum* (Papaveraceae). Cryst. (Me_2CO or $MeOH/Et_2O$ /hexane). Mp 221-223° (206-8°, 213°, 216-217°). $[\alpha]_D^{22}$ -178 (c, 0.819 in MeOH). $[\alpha]_D^{22}$ -194 (c, 0.25 in $CHCl_3$).

Hydroiodide: Mp 255° dec.

Picrate: Mp 132°.

Me ether: Amurensinine


[10470-47-0]

 $C_{20}H_{21}NO_4$ 339.39

Alkaloid from *Papaver tauricola*, *Papaver anomalum*, *Papaver pyrenaicum*, *Papaver pseudocanescens*, *Papaver alpinum*, *Papaver alpinum* var. *rhoeticum*, var. *burseri*, var. *kernerii*, var. *tatricum*, *Papaver nudicaule* var. *xanthopetalum*, var. *leiocarpum*, var. *rubroaurantiacum*, *Papaver radicum* and *Papaver suaveolens* (Papaveraceae). Cryst. (Et_2O or $MeOH$). Mp 136-138° Mp 144-146° Mp 162-164°. $[\alpha]_D^{22}$ -162 (c, 0.80 in $CHCl_3$).

Me ether: hydroiodide:Cryst. ($MeOH/Et_2O$). Mp 236-245° dec.**Me ether, N-oxide(-): Amurensinine N-oxide B** $C_{20}H_{21}NO_5$ 355.39Alkaloid from *Meconopsis horridulavar. racemosa*. $[\alpha]_D^{27}$ -89 (c, 0.38 in $CHCl_3$).**Me ether, N-oxide(+): Amurensinine N-oxide A** $C_{20}H_{21}NO_5$ 355.39Alkaloid from *Meconopsis horridulavar. racemosa*. $[\alpha]_D^{27}$ -101 (c, 0.3 in $CHCl_3$).Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 180 (*isol*)Maturová, M. *et al.*, *Planta Med.*, 1962, **10**, 345; 1966, **14**, 22 (*isol*)Šantavý, F. *et al.*, *Chem. Comm.*, 1966, 36; 144 Šantavý, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1966, **31**, 4286 (*uv*, *ir*, *pmr*, *struct*)Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 600 (*ms*)Pfeifer, S. *et al.*, *Pharmazie*, 1968, **23**, 585 (*isol*, *uv*)Shamma, M. *et al.*, *Tet. Lett.*, 1971, 3425 (*uv*, *cd*) Pfeifer, S. *et al.*, *Pharmazie*, 1972, **27**, 48 (*isol*, *uv*, *pmr*, *ms*)Novák, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 883 (*isol*, *uv*)Böhm, V.H. *et al.*, *Planta Med.*, 1975, **28**, 210 (*isol*)Dyke, S.F. *et al.*, *Tetrahedron*, 1978, **34**, 241 (*cd*) Sariyar, G. *et al.*, *Phytochemistry*, 1980, **19**, 2189 (*isol*, *ms*)Carrillo, L. *et al.*, *J.O.C.*, 1997, **62**, 6716-6721 (*Amurensinine*, *synth*)Xie, H. *et al.*, *Fitoterapia*, 2001, **72**, 120-123 (*Amurensinine N-oxides*)Tambar, U.K. *et al.*, *J.A.C.S.*, 2006, **128**, 11752-11753 (*Amurensinine*, *synth*)**Amurine** A-955

5,6,8,14-Tetradehydro-6-methoxy-17-methyl-2,3-[methylenebis(oxy)]morphinan-7-one, 9CI

 $C_{19}H_{19}NO_4$ 325.363**(+)-form** [4984-99-0]**(+)-form**

Alkaloid from *Papaver* spp., esp. *Papaver nudicaule* (Papaveraceae). Cryst. (Me_2CO). Mp 213-215°. $[\alpha]_D^{25}$ +10 (c, 1.0 in $CHCl_3$). λ_{max} 238 (log ϵ 3.67); 290 (log ϵ 3.7) (EtOH).

N-De-Me: Synthetic. Oil. $[\alpha]_D^{25}$ +29.6 (c, 0.3 in $CHCl_3$).

7 α -Alcohol: Nudaurine. Amurinol I

[4850-04-8]

 $C_{19}H_{21}NO_4$ 327.379

Alkaloid from *Papaver nudicaule* var. *aurantiacum* and *Papaver croceum* (Papaveraceae). Cryst. (Me_2CO). Mp 201-202°. $[\alpha]_D^{22}$ -52 (c, 1 in $CHCl_3$). Abs. config. revised in 1993. λ_{max} 244 (log ϵ 4); 292 (log ϵ 3.78) (EtOH).

7 α -Alcohol, hydroiodide: Mp 198° dec.**7 α -Alcohol, picrate:** Mp 150° dec.**8,14 β -Dihydro, 7 α -alcohol: Dihydro-nudaurine** $C_{19}H_{23}NO_4$ 329.395

Alkaloid from aerial parts of *Papaver pilosum*. Noncryst. $[\alpha]_D^{20}$ +100 (c, 0.32 in $CHCl_3$). Incorrectly indexed in CA. λ_{max} 233 (log ϵ 3.97); 292 (log ϵ 3.85) (MeOH).

(-)-form [77449-68-4]

Alkaloid from *Roemeria refracta* and *Stephania aculeata*. Amorph. $[\alpha]_D$ -14 (c, 0.11 in MeOH). $[\alpha]_D$ -10 (c, 0.15 in $CHCl_3$). λ_{max} 242 (log ϵ 4.18); 291 (log ϵ 3.87) (MeOH).

N-De-Me: Noramurine

[132209-25-7]

 $C_{18}H_{17}NO_4$ 311.337

Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D$ -8 (c, 0.1 in $CHCl_3$). $[\alpha]_D^{25}$ -33.3 (c, 0.3 in $CHCl_3$) (*synthetic*). λ_{max} 240 (log ϵ 4.08); 290 (log ϵ 3.84) (MeOH).

(±)-form [21124-18-5]

Yellowish oil.

N-Me:

Cryst. (MeOH) (as iodide). Mp 230-233° (222-224°, 202-206° dec.) (iodide).

7 α -Alcohol: (±)-Nudaurine. Alkaloid PNI

PNI

Alkaloid from *Papaver nudicaule* ssp. *xanthopetalum* and *Papaver nudicaule* ssp. *album* (Papaveraceae). Needles (MeOH). Mp 116-117°. Suggested struct. λ_{max} 215 (log ϵ 4.23); 245 (sh) (log ϵ 3.95); 294 (log ϵ 3.91) (MeOH).

8,14 ξ -Dihydro: 8,14-Dihydroamurine

[1012311-30-6]


 $C_{19}H_{21}NO_4$ 327.379

Alkaloid from *Papaver nudicaule*. Amorph. solid. Racemic. λ_{max} 222 (log ϵ 3.79); 254 (log ϵ 3.66); 268 (log ϵ 3.5); 280 (log ϵ 3.49) (MeOH).

Snatzke, G. *et al.*, *J.C.S.(C)*, 1966, 1681-1685 (*cd*)Döpke, W. *et al.*, *Tetrahedron*, 1968, **24**, 4459-4476 (*struct*, *pmr*)Kametani, T. *et al.*, *J.C.S.(C)*, 1969, 801; 1971, 2446-2448; 2712-2714 (*synth*)Kotani, E. *et al.*, *Tet. Lett.*, 1973, 4759-4762 (*synth*)Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 583-586 (*synth*)Roblot, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1984, 139-141 (*cmr*)

- Vezník, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1985, **50**, 1745-1752; 1987, **52**, 1634-1640 (*Amurine*, Alkaloid P_{NI}, *isol*)
 Gozler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 986-988 (*isol*, *pmr*, *struct*, *cd*, *abs config*)
 Gollwitzer, J. *et al.*, *Nat. Prod. Lett.*, 1993, **2**, 197-201 (*Nudaurine*, *cryst struct*, *abs config*)
 Hara, H. *et al.*, *Tetrahedron: Asymmetry*, 1995, **6**, 1683-1692 (*Amurine*, *Noramurine*, *synth*)
 Blanchfield, J.T. *et al.*, *Phytochemistry*, 2003, **63**, 711-720 ((-)-*Amurine*, *isol*, *pmr*, *cmr*, *cryst struct*)
 Philipov, S. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 852-856 (8,14-Dihydroamurine)

Amurinine **A-956**
 [87245-86-1]




C₁₉H₂₁NO₄ 327.379
 Alkaloid from the aerial parts of *Papaver pilosum* and *Papaver apokrinomenon* (Papaveraceae). [α]_D²⁰ +42 (c, 0.18 in CHCl₃). [α]_D²⁰ +42 (c, 0.18 in MeOH).

6-Epimer: Epiamurinine

[87220-76-6]
 C₁₉H₂₁NO₄ 327.379
 Alkaloid from the aerial parts of *Papaver pilosum* and *Papaver apokrinomenon* (Papaveraceae). [α]_D²⁰ -68 (c, 0.58 in CHCl₃).

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 342 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)
 Oztekin, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 560 (*isol*)

Amuroline **A-957**



(+)-form

C₁₉H₂₅NO₃ 315.411
 Diastereomeric with Cryprochine, C-781.

(+)-form [10214-57-0]

Alkaloid from *Papaver nudicaule* var. *amurense* (Papaveraceae). Prisms (Me₂CO). Mp 165-166°. [α]_D²⁶ +106 (c, 0.3 in CHCl₃).

Perchlorate: Mp 151-153°.

Ac: Mp 152-154°. [α]_D²⁴ +147 (c, 0.85 in CHCl₃).

4-Ketone: Amurinine

[10214-56-9]
 C₁₉H₂₃NO₃ 313.396
 Alkaloid from *Papaver nudicaule* var. *amurense* (Papaveraceae). Mp 119-120°
 Mp 132-133° (double Mp). [α]_D²⁶ +140

(c, 0.3 in CHCl₃).

4-Ketone, O'-de-Me: N-Methylisocrot-sparinine

C₁₈H₂₁NO₃ 299.369
 Alkaloid from *Croton sparsiflorus* (Euphorbiaceae). Mp 197-199°. [α]_D +111.5 (c, 0.34 in MeOH).

4-Ketone, O'-de-Me, N-de-Me: Isocrot-sparinine

[58166-01-1]
 C₁₇H₁₉NO₃ 285.342
 Alkaloid from the aerial parts of *Croton sparsiflorus* (Euphorbiaceae). Cryst. (EtOAc). Mp 198-199°. [α]_D +112.5 (c, 0.23 in CHCl₃).

4-Ketone, O²-de-Me: Linearisine

[10214-55-8]
 C₁₈H₂₁NO₃ 299.369
 Alkaloid from *Croton linearis* (Euphorbiaceae). Mp 219-222°. [α]_D²⁸ +116 (c, 0.83 in MeOH).

4-Ketone, O²-de-Me, hydrochloride: Mp 300° dec.

10-Epimer, O-de-Me (?): Discolorine

C₁₈H₂₃NO₃ 301.385
 Alkaloid from *Croton discolor* (Euphorbiaceae). Mp 206-208°. [α]_D +99 (c, 0.5 in EtOH). Struct. assigned here on the basis of later work on the stereochem. of these alkaloids. The Me ether was not the same as Amurinine, but MnO₂ oxidn. of the Me ether gave Amurinine (incl. opt. rotn.).

(±)-form

Synthetic. Noncryst.

Hydrochloride:

Cryst. (MeOH/Et₂O). Mp 235° dec. (preheated to 230°).

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1959, **46**, 514 (*isol*)

Haynes, L.J. *et al.*, *J.C.S.*, 1963, 1784-1788; 1789-1793 (*Linearisine*, *Amurinine*, *isol*, *uv*, *ir*, *pmr*)

Haynes, L.J. *et al.*, *J.C.S.(C)*, 1966, 1676-1679 (*Linearisine*, *Amurinine*, *struct*)

Snatzke, G. *et al.*, *J.C.S.(C)*, 1966, 1681-1685 (*cd*, *abs config*)

Baldwin, M. *et al.*, *J.C.S.(C)*, 1967, 154-161 (*ms*)

Bernauer, K. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1119 (*synth*)

Döpke, W. *et al.*, *Tetrahedron*, 1968, **24**, 2297 (*pmr*, *cd*, *struct*, *stereochem*)


Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 1558 (*ord*)

Stuart, K.L. *et al.*, *J.C.S.(C)*, 1970, 1228-1230 (*Discolorine*)

Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 571 (*ms*)

Casagrande, C. *et al.*, *J.C.S. Perkin I*, 1975, 1659-1663 (*Isocrot-sparinine*, *N-Methylisocrot-sparinine*)

Amycomycin **A-958**
 [344362-08-9]




C₆₅H₁₁₅NO₁₈ 1198.619

Tetramic acid antibiotic. Prod. by *Amycolatopsis* sp. ST101170. Active against gram-positive bacteria. Amorph. solid. λ_{max} 229 (log ε 3.29); 281 (log ε 3.16) (no solvent reported).

Eur. Pat., 2001, 1 106 604; *CA*, **135**, 32809t (*isol*)

Amythiamicin A **A-959**
 MI 481-42F4A. Antibiotic MI 481-42F4A [152741-89-4]




C₅₀H₅₁N₁₅O₈S₆ 1182.445

CycliC thiazole peptide. Prod. by *Amycolatopsis* sp. MI 481-42F4. Active against gram-positive bacteria. Sol. MeOH, DMSO; poorly sol. H₂O, hexane. [α]_D²⁸ +133 (c, 0.7 in DMSO). λ_{max} 203 (ε 83200); 221 (ε 77600); 250 (ε 53700); 310 (ε 35500); 345 (ε 13500) (MeOH) (Derep). λ_{max} 203 (ε 85113); 223 (ε 79432); 303 (ε 33884) (MeOH/HCl). λ_{max} 204 (ε 288400); 309 (ε 33880) (MeOH/NaOH).

Shimamaka, K. *et al.*, *J. Antibiot.*, 1994, **47**, 668-674; 1145-1152; 1153-1159 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *props*)
 Nicolaou, K.C. *et al.*, *Chem. Comm.*, 2008, 2632-2634 (*synth*)

Amythiamicin B **A-960**
 MI 481-42F4B. Antibiotic MI 481-42F4B [156620-48-3]




C₅₀H₅₃N₁₅O₈S₆ 1200.46

Cyclic polythiazole peptide. Prod. by *Amycolatopsis* sp. MI 481-42F4. Active against gram-positive bacteria. Sol. MeOH, DMSO; poorly sol. H₂O, hexane. $[\alpha]_D^{23} +155$ (c, 0.25 in MeOH). Nontoxic. $\lambda_{\max} 203$ (ε 83200); 221 (ε 77600); 250 (sh) (ε 52500); 308 (ε 35500); 345 (sh) (ε 13500) (MeOH) (Derep). $\lambda_{\max} 204$ (ε 58884); 222 (ε 56234); 308 (ε 23988) (MeOH/HCl).

Shimanaka, K. *et al.*, *J. Antibiot.*, 1994, **47**, 668-674; 1145-1152; 1153-1159 (*isol, uv, ir, pmr, cmr, props*)

Nicolaou, K.C. *et al.*, *Chem. Comm.*, 2008, 2632-2634 (*synth*)

Amythiamicin C **A-961**
MI 481-42F4C. Antibiotic MI 481-42F4C
[156620-47-2]



C₅₀H₅₀N₁₄O₉S₆ 1183.429

Cyclic polythiazole peptide. Prod. by *Amycolatopsis* sp. MI 481-42F4. Active against gram-positive bacteria. Poorly sol. H₂O, hexane. $[\alpha]_D^{24} +112$ (c, 0.25 in MeOH). $\lambda_{\max} 203$ (ε 83200); 221 (ε 77600); 250 (sh) (ε 52500); 308 (ε 35500); 345 (sh) (ε 13500) (MeOH) (Derep). $\lambda_{\max} 203$ (ε 60256); 221 (ε 57543); 306 (ε 25700) (MeOH/HCl). $\lambda_{\max} 203$ (ε 27542); 306 (ε 24547) (MeOH/NaOH).

Shimanaka, K. *et al.*, *J. Antibiot.*, 1994, **47**, 668-674; 1145-1152; 1153-1159 (*isol, uv, ir, pmr, cmr, props*)

Nicolaou, K.C. *et al.*, *Chem. Comm.*, 2008, 2632-2634 (*synth*)

Amythiamicin D **A-962**
MI 481-42F4D. Antibiotic MI 481-42F4D
[156620-46-1]

As Amythiamicin A, A-959 with
R = -COOMe


C₄₃H₄₂N₁₂O₇S₆ 1031.277

Cyclic polythiazole peptide. Prod. by *Amycolatopsis* sp. MI 481-42F4. Active against gram-positive bacteria. Sol. MeOH, DMSO; poorly sol. H₂O, hexane. $[\alpha]_D^{25} +179$ (c, 0.5 in MeOH). $\lambda_{\max} 204$ (ε 52500); 224 (ε 61700); 250 (sh) (ε 36300); 307 (ε 30900); 345 (sh) (ε 11000) (MeOH) (Derep). $\lambda_{\max} 205$ (ε 52480); 224 (ε 63095); 307 (ε 31622) (MeOH/HCl). $\lambda_{\max} 204$ (ε 60260); 224 (ε 61660); 308 (ε 30900) (MeOH/NaOH).

Shimanaka, K. *et al.*, *J. Antibiot.*, 1994, **47**, 668-674; 1145-1152; 1153-1159 (*isol, uv, ir, pmr, cmr, props*)

Hughes, R.A. *et al.*, *J.A.C.S.*, 2005, **127**, 15644-15651 (*synth*)

Anabasamine **A-963**
5-(1-Methyl-2-piperidinyl)-2,3'-bipyridine, 9CI
[20410-87-1]



C₁₆H₁₉N₃ 253.346

Alkaloid from the seeds of *Anabasis aphylla* (Chenopodiaceae). Mp 65-66°. $[\alpha]_D +107$ (c, 3.65 in EtOH).

►BV4299600

N-De-Me: 1'',2'',3'',4'',5'',6''-Hexahydro-2,3':5,2''-terpyridine. 5-(2-Piperidinyl)-2,3'-bipyridine. **Noranabasamine**

[85365-52-2]

C₁₅H₁₇N₃ 239.319


Minor alkaloid from skin extracts of the Colombian poison-dart frog *Phylllobates terribilis*. Also identified by gc-ms in skin extracts of *Phylllobates aurotaenia* and *Phylllobates bicolor*. $[\alpha]_D^{25} -14.4$ (MeOH).

Sadykov, A.S. *et al.*, *Dokl. Akad. Nauk SSSR*, 1967, **24**, 34; *CA*, **68**, 78473e (*isol, uv, ir, pmr, ms, struct*)

Mukhamedzhanov, S.Z. *et al.*, *Khim. Priir. Soedin.*, 1968, **4**, 158; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 136 (*uv, ir, pmr, ms, struct*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1983, **39**, 41 (*Noranabasamine*)

Anabilsine **A-964**
1-(1-Amino-1-carboxypentyl)-3-[1-(1-amino-1-carboxypentyl)-2-piperidinyl]-pyridinium(1+)
[66517-63-3]




C₂₂H₃₇N₄O₄⁺ 421.559

Isol. from acid hydrolysates of glutaraldehyde-treated ovalbumin. CAS no. refers to chloride. $\lambda_{\max} 263$ (ε 400) (NH₄OH).

Hardy, P.M. *et al.*, *J.C.S. Perkin 1*, 1979, 2282 (*isol, struct, synth, pmr, cmr*)

Anachelin H **A-965**
[326490-92-0]



Absolute Configuration


C₃₅H₅₁N₆O₁₄⁺ 779.82

Isol. from *Anabaena cylindrica* CCAP 1403/2A. Siderophore.

Beiderbeck, H. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 681-687 (*isol*)

Gademann, K. *et al.*, *J.O.C.*, 2007, **72**, 8361-8370 (*synth*)

Anacine **A-966**
[154613-22-6]



Absolute Configuration

C₁₈H₂₂N₄O₃ 342.397

Struct. revised in 1999. Related to Fumiquinazoline A, F-185. Metab. of marine-derived and terrestrial *Penicillium aurantiogriseum*. Also from *Penicillium polonicum*. Mycotoxin. Amorph. solid. Sol. H₂O; poorly sol. Me₂CO, hexane. $[\alpha]_D^{24} +233.3$ (c, 0.21 in MeOH). $\lambda_{\max} 226$ (log ε 4.23); 270 (log ε 3.6) (MeOH).

1,12-Didehydro(Z): **Aurantiomide C**
[915190-86-2]

C₁₈H₂₀N₄O₃ 340.381

Prod. by the marine-derived *Penicillium aurantiogriseum* SP0-16. Amorph. powder. $[\alpha]_D^{24} +25.8$ (c, 0.1 in CHCl₃). $\lambda_{\max} 314$ (log ε 4.48) (MeOH).

1-Hydroxy: **Aurantiomide B**

[915190-84-0]

C₁₈H₂₂N₄O₄ 358.396

Prod. by the marine-derived *Penicillium aurantiogriseum* SP0-19. Amorph. powder. $[\alpha]_D^{24} +96.5$ (c, 0.09 in CHCl₃). $[\alpha]_D +172$ (c, 1.8 in MeOH). $\lambda_{\max} 280$ (log ε 4.6); 311 (log ε 4.28) (MeOH).

1-Methoxy: **Aurantiomide A**

[915190-85-1]

C₁₉H₂₄N₄O₄ 372.423


Prod. by the marine-derived *Penicillium aurantiogriseum* SP0-19. Oil. $[\alpha]_D^{24} +16$ (c, 0.05 in CHCl₃). $\lambda_{\max} 282$ (log ε 3.9); 311 (log ε 3.71) (MeOH).

- Boyes-Korkis, J.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1707-1717 (*isol, uv, ir, pmr, cmr, ms*)
 Larsen, T.O. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1578-1580 (*struct*)
 Wang, H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1497-1501 (*synth, pmr, cmr*)
 Xin, Z.H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 853-855 (*Aurantionides A-C*)

Anacrotine

A-967

6,12-Dihydroxysenecionan-11,16-dione,
 9CI. Crotalaburnine
 [5096-49-1]



Absolute
 configuration

$C_{18}H_{25}NO_6$ 351.399

Cyclic diester of Crotanecine, C-775 with senecic acid. Alkaloid from *Crotalaria laburnifolia* (Fabaceae). Mp 191-192°. $[\alpha]_D^{20} +30$ (EtOH).

▶ Hepato- and pneumotoxin. VT5707000

O⁶-Ac: Acetylanacrotine

$C_{20}H_{27}NO_7$ 393.436

Alkaloid from *Crotalaria agatiflora* (Fabaceae). Mp 106-107°. $[\alpha]_D^{20} +74$ (c, 0.4 in EtOH).

6-Epimer: Uspallatine

[98264-41-6]

Alkaloid from the roots of *Senecio uspallatensis* (Asteraceae). Cryst. (MeOH/CHCl₃). Mp 205-207°. $[\alpha]_D^{20} +4.11$ (c, 0.0764 in MeOH).

(E)-Isomer: trans-Anacrotine

$C_{18}H_{25}NO_6$ 351.399

Minor alkaloid from seeds of *Crotalaria capensis* (Fabaceae). $[\alpha]_D^{22} +11$ (c, 1.7 in CHCl₃).

(E)-Isomer, O⁶-Ac: Acetyl-trans-anacrotine

$C_{20}H_{27}NO_7$ 393.436

Alkaloid from *Crotalaria agatiflora* (Fabaceae). Prisms (EtOH). Mp 96-97°. $[\alpha]_D^{20} +65$ (c, 1.37 in EtOH).

(E)-Isomer, O⁶-angeloyl, N-oxide: Ange-loyl-trans-anacrotine N-oxide

$C_{23}H_{31}NO_8$ 449.5

Alkaloid from *Crotalaria agatiflora* (Fabaceae). Mp 175-176°.

Atal, C.K. *et al.*, *Tet. Lett.*, 1966, 537 (*isol, struct, pmr*)

Culvenor, C.C.J. *et al.*, *An. Quim.*, 1972, **68**, 883 (*derivs*)

Crout, D.H.G. *et al.*, *J.C.S. Perkin I*, 1972, 1602 (*isol*)

Culvenor, C.C.J. *et al.*, *Chem. Biol. Interact.*, 1976, **12**, 299 (*tox*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Mackay, M.F. *et al.*, *Acta Cryst. C*, 1984, **40**, 1073 (*cryst struct*)

Pestchanker, M.J. *et al.*, *Phytochemistry*, 1985, **24**, 1622 (*Uspallatine*)

Verdoorn, G.H. *et al.*, *Phytochemistry*, 1992, **31**, 369 (*trans-Anacrotine*)

Anacycline

A-968

N-(2-Methylpropyl)-2,4-tetradecadiene-8,10-diynamide, 9CI. 2,4-Tetradecadiene-8,10-diynoic acid isobutylamide
 [502-57-8]



$C_{18}H_{25}NO$ 271.402

λ_{max} 230 (ϵ 10000); 259 (ϵ 33500); 265 (sh) (ϵ 29800) (EtOH) (Derep).

(E,E)-form [94413-18-0]

Alkaloid from the roots of *Anacyclus pyrethrum* (Asteraceae) and *Achillea* spp. Cryst. (Et₂O/petrol). Mp 122°.

Maleic anhydride adduct:

Cryst. (Et₂O/petrol). Mp 196°.

N-Me: N-Methyl-N-(2-methylpropyl)-2,4-tetradecadiene-8,10-diynamide. N-Methylanacycline
 [38340-83-9]

$C_{19}H_{27}NO$ 285.428

Constit. of the roots of *Anacyclus pyrethrum* (Asteraceae). Oil. λ_{max} 251 nm.

Tetrahydro:

Waxy needles. Mp 40°.

12,13-Didehydro: N-Isobutyl-2,4,12-tetradecatriene-8,10-diynamide. Didehydroanacycline. 12,13-Didehydroanacycline
 [29428-83-9]

$C_{18}H_{23}NO$ 269.386

Alkaloid from *Anacyclus fuscata* and *Achillea* spp. (Asteraceae). Cryst. (petrol). Mp 150.5°. Becomes rose-pink on prolonged exposure to sunlight.

(E,Z)-form [37064-16-7]

Synthetic. Cryst. (Et₂O/petrol). Mp 85.5°.

[112711-13-4, 94481-83-1]

Crombie, L. *et al.*, *J.C.S.*, 1955, 999 (*isol, uv, ir, struct*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1956, **89**, 1276; 1970, **103**, 2856; 1974, **107**, 2120 (*synth, biosynth, uv, ir, Didehydroanacycline*)

Crombie, L. *et al.*, *Chem. Ind. (London)*, 1956, 409 (*synth, uv*)

Crombie, L. *et al.*, *J.C.S.*, 1957, 2767 (*synth*)

Jente, R. *et al.*, *Chem. Ber.*, 1972, **105**, 1694 (*isol, synth, ir, pmr, uv*)

Kuropka, G. *et al.*, *Planta Med.*, 1987, **53**, 440 (*isol*)


Crombie, L. *et al.*, *Tet. Lett.*, 1987, **28**, 4875 (*synth*)

Greger, H. *et al.*, *Phytochemistry*, 1989, **28**, 2363 (*isol*)

Anaferine

A-969

1,3-Di-2-piperidinyl-2-propanone, 9CI



(R,R)-form

$C_{13}H_{24}N_2O$ 224.345

The isolated alkaloid is the *meso*-form, but Anaferine readily racemises and the (±)-form epimerises to the *meso*-form in aq. soln., so it is probable that the nat.

alkaloid is opt. active.

(R,R)-form [19519-55-2]

[19519-54-1 dihydrobromide]

Mp 242.5-243.5° (as dihydrochloride).

$[\alpha]_D^{22} -49.8$ (c, 0.529 in MeOH) (dihydrochloride).

(S,S)-form [19519-53-0]

[19519-52-9 dihydrochloride]

Mp 242.5-244° (as dihydrochloride). $[\alpha]_D^{22} +50.7$ (c, 0.736 in MeOH) (dihydrochloride).

(RS,RS)-form

(±)-form

[28666-51-5]

Dihydrobromide: Mp 247-249°.

Dipicrate: Mp 177-179°.

(RS,SR)-form

meso-form

[28699-15-2]

Constit. of root of *Withania somnifera* (Solanaceae). Mp 16-17°. Bp_{0.01} 55°.

Dihydrochloride: Mp 226-228°.

Dihydrobromide: Mp 227-229°.

Dipicrate: Mp 180-182° Mp 193-195° (dimorph.).

Rother, A. *et al.*, *Chem. Ind. (London)*, 1962, 654 (*isol, ir, ms, pmr, struct*)

Schöpf, C. *et al.*, *Annalen*, 1970, **737**, 1 (*synth, ir, abs config*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1971, **90**, 1326 (*abs config*)

Cymerman Craig, J. *et al.*, *J.O.C.*, 1978, **48**, 347 (*ord, cd*)

Blechert, S. *et al.*, *Eur. J. Org. Chem.*, 2002, 2855-2858 (*synth, abs config*)


Anagryrine

A-970

7,7a,8,9,10,11,13,14-Octahydro-7,14-methano-4H,6H-dipyrido[1,2-a;1',2'-e][1,5]diazocin-4-one, 9CI. Rhombinine. Monolupine. Alkaloid III

[486-89-5]

[34389-11-2 ((±)-form)]



Absolute
 Configuration

$C_{15}H_{20}N_2O$ 244.336

Alkaloid from *Anagryris foetida*, several *Cytisus*, *Genista*, *Lupinus*, *Sophora* and *Ammodendron* spp, *Ulex europaeus*, *Ulex nanus*, *Templetonia retusa* and *Thermopsis chinensis*.

Antiarrhythmic, diuretic, purgative agent. Cardiotonic agent inducing tachycardia. Major toxic alkaloid responsible for "crooked calf disease" caused by leguminous plants. Noncryst. $[\alpha]_D -165$ (EtOH). Log P 0.82 (calc). A reported crystal structure determination (1991) of Anagryrine was erroneous; it was Thermopsine, T-363.

▶ Exp. teratogen. BV5620000

Hydrochloride: Mp 295-297°.



Methiodide: Mp 264° dec.

Hydroxy: Hydroxyanagyrine

[67984-05-8]

C₁₅H₂₀N₂O₂ 260.335Alkaloid from *Anagryis foetida* (Fabaceae). No phys. props. reported.

Subst. in the D-ring (ring remote from the carbonyl function); posn. of subn. unknown.

11-Epimer: see Thermopsine, T-363Couch, J.F. *et al.*, *J.A.C.S.*, 1939, **61**, 3327 (*isol*)Ribas, I. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1952, **48**, 161; 167 (*isol*)van Tamelen, E.E. *et al.*, *J.A.C.S.*, 1958, **80**, 4659 (*synth*)Fales, H.M. *et al.*, *J.A.C.S.*, 1970, **92**, 1590 (*ms*)Goldberg, S.I. *et al.*, *J.O.C.*, 1972, **37**, 1823 (*synth*)Bohlmann, F. *et al.*, *Chem. Ber.*, 1975, **108**, 1043 (*cmr*)Keeler, R.F. *et al.*, *CA*, 1976, **85**, 138290f (*tox*)
Lobo, J.M. *et al.*, *An. Quim.*, 1977, **73**, 1366 (*Hydroxyanagyrine*)Rycroft, D.S. *et al.*, *Magn. Reson. Chem.*, 1991, **29**, 936 (*pmr, cmr, conformn*)Robins, D.J. *et al.*, *Magn. Reson. Chem.*, 1992, **30**, 1125 (*config*)Rode, J.E. *et al.*, *J. Mol. Struct.*, 2005, **749**, 1 (*cryst struct*)Gray, D. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 2419-2423 (*synth*)**Anahygrine****A-971**C₁₃H₂₄N₂O 224.345Alkaloid from roots of *Withania somnifera* (Solanaceae). Pale yellow oil. Bp_{0.2} 106°.**Dihydrochloride:** Mp 216.5-217.5°.**Dipicrate:** Mp 173-174.5°.Schwartz, A.E. *et al.*, *J. Nat. Prod.*, 1963, **26**, 258 (*isol*)Leary, J.D. *et al.*, *Chem. Ind. (London)*, 1964, 283 (*isol*)El-Olemy, M.M. *et al.*, *Experientia*, 1965, **21**, 249 (*synth*)**Anantine†****A-972**C₁₅H₁₅N₃O 253.303λ_{max} 218 (ε 15800); 277 (ε 25000) (prob. MeOH) (Derep).**(R)-form** [50656-82-1]Alkaloid from the leaves of *Cynometra ananta* and from the root bark, trunk bark and leaves of *Cynometra lujae* (Fabaceae). Mp 204°. [α]_D -549.**N-Ac:** Mp 99°. [α]_D -365.**N-De-Me:** **Noranantine**


[85651-90-7]

C₁₄H₁₃N₃O 239.276Alkaloid from the root bark and trunk bark of *Cynometra lujae* (Fabaceae).Cryst. (Me₂CO/MeOH). Mp 205°.[α]_D²⁰ -431 (c, 1 in CHCl₃).**3'-Hydroxy: Hydroxyanantine**

[85644-21-9]

C₁₅H₁₅N₃O₂ 269.302Alkaloid from the root bark and trunk bark of *Cynometra lujae* (Fabaceae). Cryst. (Me₂CO/MeOH). Mp 170°. [α]_D²⁰ -433 (c, 1 in CHCl₃). Hydroxylated in the phenyl ring.**(±)-form** [68069-26-1]


Synthetic. Mp 179°.

Khuong-Huu, F. *et al.*, *Tet. Lett.*, 1973, 1757 (*ir, uv, pmr, cmr, struct*)Tchissambou, L. *et al.*, *Tet. Lett.*, 1978, 1801 (*synth*)Tchissambou, L. *et al.*, *Tetrahedron*, 1982, **38**, 2687 (*isol, cmr, struct, derivs*)Naito, T. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1932-1939 (*synth*)**Anatalline****A-973****3,3'-(2,4-Dipiperidinediyl)bispyridine**,
9CI. 2,4-Di-3-pyridylpiperidine
[18793-19-6]

Relative configuration

C₁₅H₁₇N₃ 239.319Alkaloid from the leaves and roots of *Nicotiana tabacum*. Also found in *Duboisia hopwoodii*. Amorph. solid. Bp₃ 225°. [α]_D 0. *Isol.* as a mixt. with its *trans*-isomer. Dehydrogenation gives 3,2':4',3''-Terpyridine, T-98.**Perchlorate (1:3):** [18793-18-5]Cryst. (MeOH/Et₂O). Mp 244-252°.**Picrate:** [18793-20-9]

Mp 258.5°.

Kisaki, T. *et al.*, *Phytochemistry*, 1968, **7**, 323-327 (*isol, ir, pmr, ms*)Rao, H.S.P. *et al.*, *Indian J. Chem., Sect. B*, 1997, **36**, 557-561 (*synth, pmr, cmr, config*)Häkkinen, S.T. *et al.*, *Planta Med.*, 2004, **70**, 936-941 (*isol, pmr, cmr, ms*)**Anatoxin a****A-974****1-(9-Azabicyclo[4.2.1]non-2-en-2-yl)ethanone**, **9CI. 2-Acetyl-9-azabicyclo[4.2.1]non-2-ene**. *Very fast death factor*. **VFDF. AnTx****(+)-form**C₁₀H₁₅NO 165.235**(+)(HCl)-form** [64285-06-9]


[64314-16-5 (hydrochloride)]

Alkaloid from *Anabaena flos-aquae* NRC-44h (Euphorbiaceae). Also *isol.* from *Oscillatoria* spp., *Aphanizomenon flos-aquae* and *Cylindrospermum* sp.Depolarising agent. Potent agonist for the nicotinic acetylcholine receptor. Oil. Sol. H₂O, MeOH. [α]_D²⁴ +43.2 (c, 0.676 in EtOH). λ_{max} 226 (ε 8300) (MeOH) (Berdy). λ_{max} 227 (ε 10000) (EtOH) (Berdy).► *Exp. reprod. effects.* Very toxic by intraperitoneal route; LD₅₀ (mus, ipr) 0.3 mg/kg, LD₅₀ (mus, ipr) 0.2 mg/kg. KM5527000**N-Ac:** Mp 117-118°. [α]_D -127 (c, 4.8 in EtOH).**(-)(HCl)-form** [92142-32-0]Synthetic. [α]_D²³ -45 (c, 0.3 in EtOH) (hydrochloride).Devlin, J.P. *et al.*, *Can. J. Chem.*, 1977, **55**, 1367-1371 (*isol, pmr, cmr, ms*)Campbell, H.F. *et al.*, *Can. J. Chem.*, 1977, **55**, 1372 (*synth*)Bates, H.A. *et al.*, *J.A.C.S.*, 1979, **101**, 1259-1265 (*synth*)Spivak, C.E. *et al.*, *Mol. Pharmacol.*, 1980, **18**, 384-394; 1983, **23**, 337-343; 1986, **29**, 250-257 (*pharmacol*)Petersen, J.S. *et al.*, *J.A.C.S.*, 1984, **106**, 4539-4547 (*synth*)Koskinen, A.M.P. *et al.*, *J. Med. Chem.*, 1985, **28**, 1301-1309 (*synth, cryst struct*)Danheiser, R.L. *et al.*, *J.A.C.S.*, 1985, **107**, 8066 (*synth*)Tufariello, J.J. *et al.*, *Tetrahedron*, 1985, **41**, 3447 (*synth*)Wiseman, J.R. *et al.*, *J.O.C.*, 1986, **51**, 2485 (*synth, bibl*)Vernon, P. *et al.*, *Chem. Comm.*, 1987, 245 (*synth, bibl*)Stjernloef, P. *et al.*, *Acta Chem. Scand.*, 1989, **43**, 917 (*synth*)Matsunaga, S. *et al.*, *J.A.C.S.*, 1989, **111**, 8021 (*isol*)Sardina, F.J. *et al.*, *J.O.C.*, 1989, **54**, 4654; 1990, **55**, 5025 (*synth, bibl*)Gallon, J.R. *et al.*, *Phytochemistry*, 1990, **29**, 1107; 1994, **35**, 1195 (*biosynth*)Somfai, P. *et al.*, *Tet. Lett.*, 1992, **33**, 3791 (*synth*)Skrinjar, M. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 1263 (*synth*)Harada, K. *et al.*, *Tetrahedron*, 1993, **49**, 9251 (*isol*)Newcombe, N.J. *et al.*, *Chem. Comm.*, 1995, 831 (*synth*)Hemscheidt, T. *et al.*, *Chem. Comm.*, 1995, 1361-1362 (*biosynth*)Molloy, L. *et al.*, *Eur. J. Pharmacol.*, 1995, **289**, 447-453 (*pharmacol*)Mansell, H.L. *et al.*, *Tetrahedron*, 1996, **52**, 6025 (*rev, synth*)Parsons, P.J. *et al.*, *Tetrahedron*, 1996, **52**, 11637 (*synth*)Oh, C.-Y. *et al.*, *Tet. Lett.*, 1998, **39**, 2133-2136 (*synth*)*Food Sci. Technol., Seafood and Freshwater Toxins*, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (revs)Parsons, P.J. *et al.*, *Tetrahedron*, 2000, **56**, 309-315 (*synth*)Mori, M. *et al.*, *Tet. Lett.*, 2004, **45**, 4397-4399 (*synth*)Brenneman, J.B. *et al.*, *Tetrahedron*, 2004, **60**, 7301-7314 (*synth*)Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 134-137 (*synth*)

- Wonnacott, S. et al., *Mar. Drugs*, 2006, **4**, 228-254 (rev. activity)
 Tomita, T. et al., *Tetrahedron*, 2006, **62**, 10518-10527 (synth)
 Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AOO120

Anatoxin a(s) **A-975**

2-Amino-4,5-dihydro-1-[(hydroxymethoxyphosphinyl)oxy]-N,N-dimethyl-1H-imidazole-5-methanamine, 9CI
 [103170-78-1]



C₇H₁₇N₄O₄P 252.209


Isol. from the blue-green alga, *Anabaena flos-aquae* NRC525.17. Potent neurotoxin with anticholinesterase activity. Dec. slowly at -20°. λ_{max} 200 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 0.05 mg/kg.
 BV6593100

Matsunaga, S. et al., *J.A.C.S.*, 1989, **111**, 8021-8023 (isol, pmr, cmr)

Anchinopeptolide A **A-976**

[152369-61-4]



C₃₄H₄₆N₁₀O₈ 722.8

Stereochem. at C-5, C-4' and C-5' is relative. Isol. from the marine sponge *Anchinoe tenacior*. [α]_D -103.6 (c, 4 in MeOH). λ_{max} 217 (ε 13600); 285 (ε 20000) (MeOH) (Derep).

7-Demethyl: Anchinopeptolide C

[160041-36-1]

C₃₃H₄₄N₁₀O₈ 708.773

From *Anchinoe tenacior*. [α]_D -6.3 (c, 1 in MeOH). λ_{max} 284 (ε 14490); 320 (MeOH) (Berdy).

7-Demethyl: Anchinopeptolide B

[160072-37-7]

C₃₃H₄₄N₁₀O₈ 708.773

From *Anchinoe tenacior*. [α]_D -12.4 (c, 0.7 in MeOH). λ_{max} 284 (ε 12070); 318 (MeOH) (Berdy).

7,7'-Bis(demethyl): Anchinopeptolide D

[160072-38-8]

C₃₂H₄₂N₁₀O₈ 694.746

From *Anchinoe tenacior*. [α]_D +11.4 (c, 1 in MeOH). λ_{max} 284 (ε 16940); 318 (MeOH) (Berdy).


Casapullo, A. et al., *Tet. Lett.*, 1993, **34**, 6297 (isol, uv, ir, pmr, cmr, struct)

Casapullo, A. et al., *J. Nat. Prod.*, 1994, **57**, 1227 (*Anchinopeptolides B-D*)

Snider, B.B. et al., *J.O.C.*, 2000, **65**, 793-800 (*Anchinopeptolide D*, synth)

Ancustrigosine

[701974-94-9]




C₂₄H₄₁NO₈ 471.59

Alkaloid from the roots of *Ancusa strigosa*. Red oil. [α]_D²⁵ -4 (c, 0.1 in MeOH). λ_{max} 239 (sh); 286 (MeOH).

Braca, A. et al., *Planta Med.*, 2003, **69**, 835-841 (isol, pmr, cmr)

Ancisheynine

[603933-04-6]



C₂₆H₂₈NO₄⁺ 418.512


Quaternary alkaloid from the aerial parts of *Ancistrocladus heyneanus*. Yellow solid. Counterion not specified. Racemate. λ_{max} 233 (log ε 4.47); 264 (log ε 4.29); 307 (log ε 3.89); 358 (log ε 3.71) (MeOH).

Yang, L.-K. et al., *Tet. Lett.*, 2003, **44**, 5827-5829

Bringman, G. et al., *Org. Lett.*, 2006, **8**, 1037-1040 (synth)

Ancistrobrevine A

[169276-08-8]



C₂₇H₃₃NO₄ 435.562

Alkaloid from *Ancistrocladus abbreviatus* (Ancistrocladaceae).

1-Epimer, 5',8-di-O-de-Me, N-de-Me:**Ancistrogriffine C**

[478796-16-6]

C₂₄H₂₇NO₄ 393.482

Alkaloid from *Ancistrocladus griffithii*. Light yellow powder (MeOH). Mp 165°. [α]_D²⁵ +13.8 (c, 0.25 in MeOH).

Atropisomer, 1-epimer, 4',6-di-O-de-Me:**Ancistrogriffine A**

[478796-14-4]

C₂₅H₂₉NO₄ 407.508


Alkaloid from *Ancistrocladus griffithii*. Cryst. (MeOH). Mp 148°. [α]_D²⁵ +26 (c, 0.4 in MeOH).

Bringmann, G. et al., *Magn. Reson. Chem.*, 1997, **35**, 297-301 (pmr, config)

Bringmann, G. et al., *Phytochemistry*, 2002, **61**, 195-204 (*Ancistrogriffines*)

A-977**Ancistrobrevine B**

[146471-74-1]



Absolute Configuration

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus abbreviatus* and *Ophiopogon japonicus*. Amorph. powder (MeOH/Me₂CO). Mp 122-124°. [α]_D -68 (c, 0.81 in CHCl₃).

6-Me ether: Ancistroretoriline A

[302543-63-1]

C₂₆H₃₁NO₄ 421.535

Alkaloid from *Ancistrocladus tectorius*. Amorph. solid. Mp 103-105°. [α]_D²⁵ +1.3 (c, 0.75 in CHCl₃).

6-Me ether, N-Me: 5-Epi-6-O-methylancistrobertsonine A. N-Methylancistroretoriline A

C₂₇H₃₃NO₄ 435.562

Alkaloid from a Congolese *Ancistrocladus* sp. Amorph. solid. [α]_D²⁵ +1.2 (c, 0.09 in MeOH). λ_{max} 237 (log ε 1.29); 307 (log ε 0.4); 321 (log ε 0.33); 337 (log ε 0.26) (CH₂Cl₂).

4',8-Di-O-de-Me, 6-Me ether, N-Me:

[222409-76-9]

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus likoko*.

1,2-Didehydro, 6-Me ether: Ancistrotanzanine B

[604767-80-8]

C₂₆H₂₉NO₄ 419.519

Alkaloid from *Ancistrocladus likoko* and *Ancistrocladus tanzaniensis*. Yellow oil. [α]_D²⁵ +44 (c, 0.01 in MeOH). λ_{max} 230 (log ε 1.89); 305 (log ε 1.42) (MeOH).

1-Epimer, 4'-O-de-Me, 6-Me ether, N-Me: 5-Epi-4'-O-demethylancistrobertsonine C

C₂₆H₃₁NO₄ 421.535

Alkaloid from a Congolese *Ancistrocladus* sp. Cryst. (MeOH). Mp 92° (synthetic). [α]_D²⁵ +15.7 (c, 0.12 in MeOH) (natural). [α]_D²⁰ +51.8 (c, 1 in MeOH) (synthetic). λ_{max} 230 (log ε 1.05); 307 (log ε 0.42) (CH₂Cl₂).

Epimer, 6-Me ether, N-Me: [222409-80-5]

C₂₇H₃₃NO₄ 435.562

Alkaloid from *Ancistrocladus likoko*. Possesses *cis*-config.

Atropisomer, N-Me: Ancistrobertsonine A

C₂₆H₃₁NO₄ 421.535

Alkaloid from *Ancistrocladus robertsoniorum* (Ancistrocladaceae). Cryst. (MeOH aq.). Mp 220-222°. [α]_D²⁵ +29 (c, 0.5 in CHCl₃).

Atropisomer, 5'-O-de-Me: Ancistroguineine A

[202343-74-6]

C₂₄H₂₇NO₄ 393.482

Alkaloid from *Ancistrocladus guineen-*

sis. Needles (CHCl₃). Mp 202-204°. $[\alpha]_D^{25} +191.4$ (c, 0.5 in CHCl₃).

Atropisomer, 5'-O-de-Me, 6-Me ether:

Ancistroalaine B

[303752-19-4]

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus ealaensis*.

Yellow powder (MeOH). Mp 248°.

$[\alpha]_D^{20} -16.7$ (c, 0.71 in EtOH).

Atropisomer, 1,2-didehydro, 6-O-De-

methylancistroalaine A

C₂₅H₂₇NO₄ 405.493

Alkaloid from a Congolese *Ancistrocladus* sp. Yellow cryst. Mp 155-160°. $[\alpha]_D^{25} -63.2$ (c, 0.1 in MeOH). λ_{max} 232 (log ϵ 1.06); 310 (log ϵ 0.34); 316 (log ϵ 0.34); 321 (log ϵ 0.35); 334 (log ϵ 0.33) (CH₂Cl₂).

Atropisomer, 1,2-didehydro, 6-Me ether:

Ancistroalaine A

[303752-18-3]

C₂₆H₂₉NO₄ 419.519

Alkaloid from *Ancistrocladus ealaensis*.

Light yellow powder (MeOH). Mp 94-96°. $[\alpha]_D^{25} -34.3$ (c, 0.55 in EtOH).

Atropisomer, 1,2-didehydro, 4'-O-de-Me:

Ancistrolikokine D

C₂₄H₂₅NO₄ 391.466

Alkaloid from *Ancistrocladus likoko*. Amorph. solid. Mp 122-124°. $[\alpha]_D^{25} +191.6$ (c, 0.15 in CHCl₃).

Atropisomer, 1,2-didehydro, 5'-O-de-Me:

5',6-Di-O-demethylancistroalaine A

C₂₄H₂₅NO₄ 391.466

Alkaloid from a Congolese *Ancistrocladus* sp. Amorph. yellow solid. $[\alpha]_D^{25} -68.6$ (c, 0.1 in MeOH). λ_{max} 235 (log ϵ 1.15); 313 (log ϵ 0.47); 316 (log ϵ 0.47); 321 (log ϵ 0.48); 325 (log ϵ 0.47); 335 (log ϵ 0.45) (CH₂Cl₂).

Atropisomer, 1-epimer, 6-Me ether, N-Me:

Ancistrobertsonine C

[250130-68-8]

C₂₇H₃₃NO₄ 435.562

Alkaloid from *Ancistrocladus robertsoniorum*. Mp 148-149°. $[\alpha]_D^{25} -3$ (c, 0.09 in CHCl₃).

Atropisomer, 3-epimer, 5'-O-de-Me:

Ancistroguineine B

[202420-24-4]

C₂₄H₂₇NO₄ 393.482

Alkaloid from *Ancistrocladus guineensis*. Amorph. solid. $[\alpha]_D^{25} -141.2$ (c, 0.04 in CHCl₃).

Bringmann, G. et al., *Phytochemistry*, 1992, **31**, 4011; 1998, **47**, 31-35; 37-43; 1999, **52**, 321-332 (*isol, ir, cd, pmr, ms, struct, derivs*)

Bringmann, G. et al., *Phytochemistry*, 1998, **49**, 1667 (*Ancistrobertsonine A*)

Hoye, T.R. et al., *J.O.C.*, 1999, **64**, 7184-7201 (*synth*)

Bringmann, G. et al., *Magn. Reson. Chem.*, 1999, **37**, 98-102 (*Ancistrocladus likoko derivs*)

Tang, C.-P. et al., *J. Nat. Prod.*, 2000, **63**, 1384-1387 (*Ancistrocladus A*)

Bringmann, G. et al., *J. Nat. Prod.*, 2000, **63**, 1465-1470; 2003, **66**, 1159-1165 (*Ancistroalaines A-B, Ancistrotananine B, Ancistrocladus A*)

Bringmann, G. et al., *Phytochemistry*, 2003, **62**, 631-636; 2008, **69**, 1065-1075 (*Ancistrolikokine D, 6-Demethylancistroalaine, 5',6-*

Didemethylancistroalaine, 5-Epiancistrobertsonines)


Bringmann, G. et al., *Tetrahedron*, 2004, **60**, 4349-4360 (*Ancistroalaine A, Ancistrotananine B, synth*)

Bringmann, G. et al., *Tetrahedron*, 2008, **64**, 5563-5568 (*5-Epidemethylancistrobertsonine C, synth*)

Ancistrobreve C

A-981

7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-3,4-dihydro-6-methoxy-1,3-dimethyl-8-isoquinolinol
[151870-96-1]



C₂₅H₂₇NO₄ 405.493

Alkaloid from stem bark and roots of *Ancistrocladus abbreviatus* (Ancistrocladaceae). Amorph. yellow solid. Mp 180-183°. $[\alpha]_D^{25} +13$ (c, 0.69 in CHCl₃).

1,2-Dihydro, stereoisomer: Ancistine

[58738-33-3]

C₂₅H₂₉NO₄ 407.508

Alkaloid from the roots of *Ancistrocladus ealaensis* (Ancistrocladaceae). Cryst. (Me₂CO). Mp 275-276°. $[\alpha]_D^{20} -34$ (c, 1 in CHCl₃/MeOH 1:1). Stereochem. not determined.


Foucher, J.P. et al., *Phytochemistry*, 1975, **14**, 2699 (*Ancistine*)

Bringmann, G. et al., *Phytochemistry*, 1993, **33**, 1511 (*Ancistrobreve C*)

Ancistrocladine

A-982

5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-8-methoxy-1,3-dimethyl-6-isoquinolinol, 9CI
[32221-59-3]



Absolute Configuration

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus heyneanus*, *Ancistrocladus congolensis*, *Ancistrocladus hamatus* and *Ancistrocladus tectorius*. Shows antimalarial activity. Cryst. (MeOH). Mp 265-267° dec.

Hydrochloride:

Needles (MeOH/Me₂CO). Mp 220-224° dec. $[\alpha]_D^{25} -25.5$ (c, 2.3 in MeOH).

N-Ac:

Cryst. (Et₂O). Mp 277-279°.

Me ether: O-Methylancistrocladine

[32215-20-6]

C₂₆H₃₁NO₄ 421.535

Alkaloid from the root and stem bark

of *Ancistrocladus congolensis* (Ancistrocladaceae). Fluffy needles (MeOH). Mp 200-202°.

Me ether, hydrochloride:

Needles (Me₂CO). Mp 315-317° dec.

$[\alpha]_D^{25} -56.1$ (c, 1.9 in CHCl₃).

O,N-Di-Me: O,N-Dimethylancistrocladine

[32215-13-7]

Alkaloid from the leaves of *Ancistrocladus tanzaniensis*. Prisms (Et₂O). Mp 183-185° (163° dec.). $[\alpha]_D^{25} -21$ (c, 2.55 in CHCl₃).

4'-O-De-Me: 4'-O-Demethylancistrocladine

[165816-69-3]

C₂₄H₂₇NO₄ 393.482

Alkaloid from bark of *Ancistrocladus tectorius* (Ancistrocladaceae). Gum.

$[\alpha]_D +2.5$ (c, 1 in CHCl₃).

4'-O-De-Me, 6-Me ether: 4'-O-Demethyl-6-O-methylancistrocladine

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus tectorius*. Light yellow powder. $[\alpha]_D^{25} -3.1$ (c, 0.38 in MeOH).

1,2-Didehydro: Ancistrocladinine

[36209-87-7]

C₂₅H₂₇NO₄ 405.493

Minor alkaloid from the roots of *Ancistrocladus heyneanus* (Ancistrocladaceae). Mp 235-238° dec. $[\alpha]_D^{25} -321.8$ (c, 1.06 in Py).

1,2-Didehydro, Me ether: 6-O-Methylancistrocladinine

[134984-06-8]

C₂₆H₂₉NO₄ 419.519

Alkaloid from the leaves of *Ancistrocladus tanzaniensis*. Amorph. solid. Mp 102°. $[\alpha]_D^{25} -31$ (c, 0.4 in MeOH). λ_{max} 231 (log ϵ 0.22); 303 (log ϵ 0.04) (MeOH).

1,2,3,4-Tetrahydro: Ancistrocladeine

[58738-34-4]

C₂₅H₂₅NO₄ 403.477

Alkaloid from roots of *Ancistrocladus ealaensis* and *Ancistrocladus tectorius* (Ancistrocladaceae). Cryst. (Me₂CO). Mp 275-277°.

1-Epimer: Isoancistrocladine. 1-Epiancistrocladine

[36209-88-8]

C₂₅H₂₉NO₄ 407.508

Alkaloid from roots of *Ancistrocladus heyneanus* (Ancistrocladaceae). Cryst. (MeOH). Mp 230-232°. $[\alpha]_D +59.5$ (c, 0.6 in CHCl₃).

1-Epimer, N-Me: Ancistrocline

[82189-88-6]

C₂₆H₃₁NO₄ 421.535

Alkaloid from *Ancistrocladus tectorius* (Ancistrocladaceae). Mp 223-224°. $[\alpha]_D^{25} +59.1$ (c, 0.23 in CHCl₃).

1-Epimer, 5'-O-de-Me, N-Me: 5'-O-De-methylancistrocline

[828935-65-5]

C₂₅H₂₉NO₄ 407.508

Alkaloid from the stem bark of *Ancistrocladus benomensis*. Amorph. brown solid (MeOH). Mp 213°. $[\alpha]_D^{25} +61.7$ (c, 0.01 in MeOH). λ_{max} 203 (log ϵ 0.35); 227 (log ϵ 0.42); 307 (log ϵ 0.07); 319

(log ϵ 0.06); 335 (log ϵ 0.05) (MeOH).

1- Andlor 3-epimer, Me ether: Ancistroealaensine

[54382-93-3]
C₂₆H₃₁NO₄ 421.535

Alkaloid from the roots of *Ancistrocladus ealaensis* (Ancistrocladaceae). Mp 84° (softens). $[\alpha]_D^{20}$ -26 (c, 1 in MeOH). Config. unknown.

1- Andlor 3-epimer, Me ether, perchlorate: Cryst. (MeOH). Mp 132-135°. $[\alpha]_D^{20}$ -84 (c, 1 in MeOH).

Atropisomer: Hamatine

[56688-90-5]
C₂₅H₂₉NO₄ 407.508

Alkaloid from the roots of *Ancistrocladus hamatus* (Ancistrocladaceae). Mp 250-252°. $[\alpha]_D$ +77.44 (CHCl₃) (+68). Has opposite chirality of the biaryl system.

Atropisomer, Me ether:

Cryst. (hexane). Mp 160-162°. $[\alpha]_D$ +27.2 (c, 1 in CHCl₃).

Atropisomer, O,N-di-Me:

Needles (Et₂O). Mp 170-172°.

Atropisomer, 4'-O-de-Me, 6-Me ether: 4'-O-Demethyl-6-O-methylhamatine

C₂₅H₂₉NO₄ 407.508

Alkaloid from *Ancistrocladus tectorius*. Light yellow powder. $[\alpha]_D^{25}$ +12.4 (c, 0.36 in MeOH).

Atropisomer, 5'-O-de-Me: 5'-O-Demethylhamatine

C₂₄H₂₇NO₄ 393.482

Alkaloid from a Congolese *Ancistrocladus* sp. Amorph. solid. Mp 174-180°. $[\alpha]_D^{25}$ +26 (c, 0.1 in MeOH). λ_{\max} 237 (log ϵ 1.42); 309 (log ϵ 0.41); 322 (log ϵ 0.37); 337 (log ϵ 0.34) (CH₂Cl₂).

Atropisomer, 1,2-didehydro: Hamatinine

[194811-19-3]
C₂₅H₂₇NO₄ 405.493

Alkaloid from leaves of *Ancistrocladus cochinchinensis*. Isol. as a ca. 1:1 inseparable mixt. with Ancistrocladinine.

Atropisomer, 1,2-didehydro, Me ether: 6-O-Methylhamatinine

[194242-88-1]
C₂₆H₂₉NO₄ 419.519

Alkaloid from leaves of *Ancistrocladus cochinchinensis*. Amorph. $[\alpha]_D^{22}$ +34.1 (c, 1.0 in CHCl₃).

Atropisomer, 1,2-didehydro, 5'-O-de-Me: 5'-O-Demethylhamatinine

C₂₄H₂₅NO₄ 391.466

Alkaloid from a Congolese *Ancistrocladus* sp. Amorph. yellow solid. $[\alpha]_D^{25}$ +25 (c, 0.1 in MeOH). λ_{\max} 235 (log ϵ 1.05); 311 (log ϵ 0.34); 321 (log ϵ 0.33); 335 (log ϵ 0.3) (CH₂Cl₂).

Atropisomer, 1,2,3,4-tetrahydro, 6-Me ether: 6-O-Methylhamatine

C₂₆H₂₇NO₄ 417.504

Alkaloid from the leaves of *Ancistrocladus cochinchinensis*. Cryst. (CHCl₃/hexane). Mp 233-236°. $[\alpha]_D^{22}$ -41.4 (c, 0.46 in CHCl₃).

Atropisomer, 10-hydroxy, 1,2,3,4-tetrahydro, 4'-O-de-Me: 6-O-Demethylancistrobenomine A

[828935-64-4]

C₂₄H₂₃NO₅ 405.449

Alkaloid from the stem bark of *Ancistrocladus benomensis*. Pale yellow solid (MeOH). Mp 169°. $[\alpha]_D^{25}$ -34.4 (c, 0.05 in MeOH). λ_{\max} 231 (log ϵ 0.44); 259 (log ϵ 0.28); 307 (log ϵ 0.07); 335 (log ϵ 0.08); 363 (log ϵ 0.05) (MeOH).

Atropisomer, 10-hydroxy, 1,2,3,4-tetrahydro, 4'-O-de-Me, 6-Me ether: Ancistrobenomine A

[828935-63-3]

C₂₅H₂₅NO₅ 419.476

Alkaloid from the stem bark of *Ancistrocladus benomensis*. Pale yellow powder (MeOH). Mp 270° dec. $[\alpha]_D^{25}$ -21 (c, 0.1 in MeOH). λ_{\max} 219 (log ϵ 1.79); 235 (log ϵ 2.09); 259 (log ϵ 1.95); 307 (log ϵ 0.85); 323 (log ϵ 0.81); 335 (log ϵ 0.85); 363 (log ϵ 0.58) (MeOH).

Atropisomer, 1-epimer, Me ether: Ancistrobertsonine B

C₂₆H₃₁NO₄ 421.535

Alkaloid from *Ancistrocladus robertsoniorum*. Mp 192-193°. $[\alpha]_D^{25}$ +4 (c, 0.09 in CHCl₃).

Stereoisomer, O,N-di-Me: 5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,2,3-trimethylisoquinoline. Ancistrocladinine

[54382-91-1]

C₂₇H₃₃NO₄ 435.562

Alkaloid from the roots of *Ancistrocladus ealaensis* (Ancistrocladaceae). Needles (Et₂O). Mp 82-83°. $[\alpha]_D^{20}$ +20 (c, 1 in MeOH). Stereochem. unknown.

Stereoisomer, O,N-di-Me, perchlorate: Cryst. (MeOH). Mp 256-258°. $[\alpha]_D^{20}$ +71 (c, 1 in MeOH).

Govindachari, T.R. et al., *Tetrahedron*, 1971, **27**, 1013 (isol, uv, ir, pmr, struct)

Govindachari, T.R. et al., *J.C.S. Perkin 1*,

1974, 1413 (config, cryst struct, pmr)

Foucher, J.P. et al., *Phytochemistry*, 1974, **13**,

1253; 1975, **14**, 2699 (Ancistrocladine,

Ancistroealaensine, Ancistrocladinine)

Govindachari, T.R. et al., *Indian J. Chem.,*

Sect. B, 1977, **15**, 871 (Hamatine)

Bringmann, G. et al., *Angew. Chem., Int. Ed.*,

1982, **21**, 200; 1986, **25**, 913 (synth, biosynth)

Bringmann, G. et al., *Heterocycles*, 1989, **28**,

137 (synth)

Rizzacasa, M.A. et al., *J.C.S. Perkin 1*, 1991,

2773 (synth, Ancistrocladinine)

Bringmann, G. et al., *Phytochemistry*, 1992, **31**,

3297; 1994, **35**, 259; 1999, **52**, 321-332

(Ancistrocline, Isoancistrocladine,

Ancistorobertsonine B)

Bringmann, G. et al., *Tetrahedron*, 1993, **49**,

3305 (cd)

Fleischhauer, J. et al., *Z. Naturforsch., B*, 1993,

48, 140 (cd, Ancistrocladine)

Montagnac, A. et al., *Phytochemistry*, 1995,

39, 701 (4'-O-Demethylancistrocladine)

Anh, N.H. et al., *Phytochemistry*, 1997, **45**,

1287 (Hamatine, O-Methylhamatinine)

Fleischhauer, J. et al., *Z. Naturforsch., A*, 1998,

53, 993-996 (Isoancistrocladine, abs config)

Tang, C.-P. et al., *J. Nat. Prod.*, 2000, **63**, 1384-

1387 (4'-Demethyl-6-methylancistrocladine,

4'-Demethyl-6-methylhamatine)

Bringmann, G. et al., *J. Nat. Prod.*, 2004, **67**,

743-748; 2058-2062 (O,N-Dimethylancistrocladine, 6-O-

Methylancistrocladinine, Ancistrobenomine


A, 5'-O-Demethylancistrocline)

Bringmann, G. et al., *Phytochemistry*, 2008, **69**, 1065-1075 (5'-Demethylhamatine, 5'-Demethylhamatinine)

Ancistrocladinium A

A-983

[918875-18-0]



Absolute Configuration

C₂₆H₃₀NO₄⁺ 420.527

Alkaloid from the leaves of *Ancistrocladus* sp. Pale yellow cryst. Mp > 230° dec. $[\alpha]_D^{20}$ -6 (c, 0.05 in MeOH). λ_{\max} 214 (log ϵ 2.73); 225 (log ϵ 2.69); 335 (log ϵ 1.65) (MeOH).


Bringmann, G. et al., *J.O.C.*, 2006, **71**, 9348-9356 (isol, cd, pmr, cmr)

Ancistrocladinium B

A-984

[918875-19-1]

[918875-20-4]



(R)-form

C₂₅H₂₈NO₄⁺ 406.501

Alkaloid from the leaves of *Ancistrocladus* sp. Pale yellow cryst. Mp > 230° dec. $[\alpha]_D^{20}$ -8 (c, 0.06 in MeOH). Occurs as a mixt. of the two atropisomers, to which the data refers. λ_{\max} 227 (log ϵ 0.94); 346 (log ϵ 0.34) (MeOH).

(R)-form

$[\alpha]_D$ -4.5 (c, 0.04 in MeOH).

(S)-form

$[\alpha]_D^{20}$ -11.3 (c, 0.04 in MeOH).


Bringmann, G. et al., *J.O.C.*, 2006, **71**, 9348-9356 (isol, cd, pmr, cmr)

Ancistrocladisine

A-985

7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-3,4-dihydro-6,8-dimethoxy-1,3-dimethylisoquinoline, 9CI

[41787-65-9]



Absolute configuration

C₂₆H₂₉NO₄ 419.519

Alkaloid from the roots of *Ancistrocladus heyneanus* and *Ancistrocladus hamatus* (Ancistrocladaceae). Pale-brown needles (Et₂O or petrol). Mp 178-180°. $[\alpha]_D^{25}$ +7.8 (c, 0.53 in CHCl₃) (synthetic).

Hydrochloride:

Needles (MeOH/Me₂CO). Mp 220-222° dec. $[\alpha]_D$ -16.13 (c, 1.29 in CHCl₃).

Methodide:


Pale-yellow needles. Mp 220-224° dec.

O⁶-De-Me: 1,2-Didehydroancistrobertsonine D. 6-O-Demethylancistrocladisine

[250131-08-9]

C₂₅H₂₇NO₄ 405.493Alkaloid from *Ancistrocladus robertsoniorum*. Cryst. (MeOH aq.). Mp 185-187°. [α]_D²⁵ -30 (c, 0.04 in CHCl₃).Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1972, **10**, 1117 (*uv, pmr, isol, struct*)Govindachari, T.R. *et al.*, *J.C.S. Perkin 1*, 1975, 2134 (*uv, abs config*)Parthasarathy, P.C. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 590 (*cryst struct*)Bringmann, G. *et al.*, *Angew. Chem., Int. Ed.*, 1989, **28**, 1672 (*synth*)Bringmann, G. *et al.*, *Phytochemistry*, 1999, **52**, 321-332 (*1,2-Didehydroancistrobertsonine D*)**Ancistrocongine A-986***1,2-Dihydro-5-(4-methoxy-2-methyl-1-naphthalenyl)-3-methyl-6,8-isoquinoline-diol, 9CI*

[56973-82-1]


C₂₂H₂₁NO₃ 347.413Alkaloid from the roots of *Ancistrocladus congolensis* (Ancistrocladaceae). Cryst. (Me₂CO). Mp 298-299°. [α]_D 0 (MeOH).Foucher, J.P. *et al.*, *Plant. Med. Phytother.*, 1975, **9**, 87 (*isol, spectra, struct*)**Ancistrocongolensine A-987***5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2-dihydro-8-methoxy-3-methyl-6-isoquinolinol, 9CI*

[56973-81-0]

As Ancistrocongine, A-986 with R = R' = OMe

C₂₄H₂₅NO₄ 391.466Alkaloid from the roots of *Ancistrocladus congolensis* (Ancistrocladaceae). Cryst. (Me₂CO). Mp 258°. [α]_D 0 (MeOH).Foucher, J.P. *et al.*, *Plant. Med. Phytother.*, 1975, **9**, 87 (*isol, spectra, struct*)**Ancistrogriffithine A A-988**


[346702-11-2]



Absolute Configuration

C₄₈H₅₂N₂O₈ 784.947Alkaloid from *Ancistrocladus griffithii*. Light yellow powder (MeOH). Mp 230° dec. [α]_D²⁵ +73.5 (c, 0.3 in MeOH).Bringmann, G. *et al.*, *Phytochemistry*, 2002, **61**, 195-204 (*isol, pmr, cmr, ms, cd*)**Ancistroheynine A A-989**


[185417-34-9]



Absolute Configuration

C₂₄H₂₅NO₄ 391.466Alkaloid from *Ancistrocladus heynanus*. Antimalarial agent. Yellow amorph. powder. [α]_D²⁵ +68 (c, 0.3 in CHCl₃).Bringmann, G. *et al.*, *Phytochemistry*, 1996, **43**, 1405 (*isol, ir, pmr, cms, ms, cd, struct*)**Ancistrobertsonine D A-990**

[250131-22-7]



Absolute Configuration

C₂₅H₂₉NO₄ 407.508Alkaloid from *Ancistrocladus robertsoniorum*. Amorph. yellow powder. Mp 133-134°. [α]_D²⁵ +82 (c, 0.51 in CHCl₃).N-Me: **6-O-Demethyl-8-O-methyl-7-epiancistrobrevine D**. N-Methylancistrobertsonine D

[169168-91-6]

C₂₆H₃₁NO₄ 421.535Alkaloid from the leaves of *Ancistrocladus cochinchinensis*.8-O-De-Me, N-Me: **6-O-Demethyl-7-epiancistrobrevine D**

[194141-89-4]

C₂₅H₂₉NO₄ 407.508Alkaloid from the leaves of *Ancistrocladus cochinchinensis*. Cryst. (CHCl₃/hexane). Mp 229-232°. [α]_D²² +51.8 (c, 0.49 in CHCl₃).8-O-De-Me, O⁶,N-di-Me: **7-Epiancistrobrevine D**

[194242-89-2]

C₂₆H₃₁NO₄ 421.535Alkaloid from the leaves of *Ancistrocladus cochinchinensis*. Cryst. (CHCl₃/hexane). Mp 177-178°. [α]_D²² +46.1 (c, 1 in CHCl₃).1-Epimer, 4'-O-de-Me, 6-Me ether: **Ancistrotectoriline B**

[302543-64-2]

C₂₅H₂₉NO₄ 407.508Alkaloid from *Ancistrocladus tectorius*. Light yellow powder. [α]_D²⁵ +79.1 (c, 0.46 in CHCl₃).1-Epimer, 8-O-de-Me: **Ancistrogriffine B**

[478796-15-5]

C₂₄H₂₇NO₄ 393.482Alkaloid from *Ancistrocladus griffithii*.3-Epimer: **Ancistrocongoline D**

[455255-22-8]

C₂₅H₂₉NO₄ 407.508Alkaloid from the root bark of *Ancistrocladus congolensis*. Brownish powder. Mp 232°. [α]_D²⁴ -6.7 (c, 0.52 in CHCl₃).Stereoisomer: 7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-8-methoxy-1,3-dimethyl-6-isoquinolinol. **Ancistrine**


[58738-35-5]

C₂₅H₂₉NO₄ 407.508Alkaloid from the roots of *Ancistrocladus ealaensis*. Cryst. (Me₂CO). Mp 230-231°. [α]_D²⁰ -35 (c, 1 in MeOH). Stereochem. not determined-identity uncertain.Atropisomer, 8-O-de-Me, O⁶,N-di-Me: **Ancistrobrevine D**

[167503-63-1]

C₂₆H₃₁NO₄ 421.535Alkaloid from *Ancistrocladus abbreviatus*. Mp 172°. [α]_D²⁵ +24.9 (c, 0.37 in CHCl₃).Foucher, J.P. *et al.*, *Phytochemistry*, 1975, **14**, 2699-2702 (*Ancistrine*)Bringmann, G. *et al.*, *Planta Med.*, Suppl.1, 1992, **58**, 703-704 (*Ancistrobrevine D*)Anh, N.H. *et al.*, *Phytochemistry*, 1997, **45**, 1287-1291 (*7-Epiancistrobrevine D derivis*)Bringmann, G. *et al.*, *Phytochemistry*, 1999, **52**, 321-332 (*isol, cd, ir, pmr, cmr, ms, abs config*)Tang, C.-P. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1384-1387 (*Ancistrotectoriline B*)Bringmann, G. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1096-1101 (*Ancistrocongoline D*)Bringmann, G. *et al.*, *Phytochemistry*, 2002, **61**, 195-204 (*Ancistrogriffine B*)**Ancistrotanzanine A A-991**

[613245-54-8]



Absolute Configuration


C₂₅H₂₇NO₄ 405.493Alkaloid from *Ancistrocladus tanzanien-*

sis. Pale yellow powder. $[\alpha]_D^{25} +69.5$ (c, 0.1 in EtOH). λ_{\max} 231 (log ϵ 1.87); 307 (log ϵ 1.43) (MeOH).

Bringmann, G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1159-1165 (*isol, cd, pmr, cmr, ms*)

Ancistrotoctorine A-992

8-Methoxy-3-methyl-2-(1,2,3,4-tetrahydro-6,8-dimethoxy-1,2,3-trimethyl-7-isoquinolinyl)-1-naphthalenol. 1,2,3,4-Tetrahydro-7-(1-hydroxy-8-methoxy-3-methyl-2-naphthalenyl)-6,8-dimethoxy-1,2,3-trimethylisoquinoline [98985-59-2]



Absolute Configuration

$C_{26}H_{31}NO_4$ 421.535

Alkaloid from the leaves of *Ancistrocladus guineensis* and *Ancistrocladus tectorius*. Shows antimalarial props. Pale yellow needles (Me₂CO). Mp 134-140° Mp 150-151°. $[\alpha]_D^{25} -3.6$ (c, 0.2 in EtOH).

O⁶-De-Me: Ancistrotanzanine C

[692755-31-0]

$C_{25}H_{29}NO_4$ 407.508

Alkaloid from the leaves of *Ancistrocladus tanzaniensis*. Yellow oil. $[\alpha]_D -75.5$ (c, 0.01 in CHCl₃). λ_{\max} 231 (log ϵ 0.74); 307 (log ϵ 0.07) (MeOH).

1,2-Didehydro, N-de-Me: Ancistrocladidine

[52659-52-6]

$C_{25}H_{27}NO_4$ 405.493

Alkaloid from *Ancistrocladus heyneanus* and *Ancistrocladus tectorius*. Pale yellow needles (Me₂CO). Mp 255-258° dec. $[\alpha]_D^{25} -149.7$ (c, 1.13 in CHCl₃). λ_{\max} 238 (log ϵ 4.22); 255 (sh) (log ϵ 3.89); 285 (log ϵ 3.63); 306 (sh) (log ϵ 3.41); 320 (log ϵ 3.34); 335 (log ϵ 3.42) (MeOH).

1,2-Didehydro, O⁶-de-Me, N-de-Me: Ancistroheynine B

[813458-56-9]

$C_{24}H_{25}NO_4$ 391.466

Alkaloid from the leaves of *Ancistrocladus heyneanus*. Yellow oil. $[\alpha]_D^{25} -194.9$ (c, 0.02 in MeOH). λ_{\max} 231; 319; 335 (EtOH).

Govindachari, T.R. *et al.*, *J.C.S. Perkin 1*, 1975, 2134-2136 (*Ancistrocladidine*)

Ruangrungsri, N. *et al.*, *J. Nat. Prod.*, 1985, **48**, 529-535 (*isol, uv, ir, pmr, ms, cd, cryst struct, abs config*)

Meksuriyen, D. *et al.*, *Phytochemistry*, 1990, **29**, 2750-2752 (*Ancistrocladidine, pmr, cmr*)

Bringmann, G. *et al.*, *Phytochemistry*, 1998, **47**, 37-43 (*isol, cd, cmr, abs config*)


Bringmann, G. *et al.*, *J. Nat. Prod.*, 2004, **67**, 743-748 (*Ancistrotanzanine C*)

Bringmann, G. *et al.*, *Phytochemistry*, 2004, **65**, 2903-2907 (*Ancistroheynine B*)

Bungard, C.J. *et al.*, *J.O.C.*, 2006, **71**, 7354-7363 (*Ancistrocladidine, synth*)

Ancorinazole A-993

[473740-10-2]




$C_{18}H_{12}N_2O_8S_2$ 448.433

Alkaloid from the sponge *Ancorina* sp. Powder (as di-Na salt). Registry number refers to di-Na salt. λ_{\max} 242 (log ϵ 4.28); 280 (log ϵ 4.23); 296 (sh) (log ϵ 3.97); 308 (log ϵ 4); 340 (log ϵ 3.78); 356 (log ϵ 3.95) (MeOH) (di-Na salt).

Meragelman, K.M. *et al.*, *J.O.C.*, 2002, **67**, 6671-6677 (*isol, pmr, cmr, ms*)

Ancorinoside A A-994

[194426-98-7]



$C_{41}H_{69}NO_{17}$ 847.993

Tetramic acid deriv. Isol. from the sponges *Ancorina* sp. and *Penares sollasi*. Specifically inhibits blastulation of starfish embryos. Inhibits MT1-matrix metalloproteinase. Oil. $[\alpha]_D^{25} -5.5$ (c, 0.09 in MeOH). λ_{\max} 240 (ϵ 5600); 283 (ϵ 7700) (MeOH).

Mg salt:

$C_{41}H_{67}MgNO_{17}$ 870.282

Isol. from *Ancorina* sp. Powder. Mp 200-208° dec. $[\alpha]_D^{25} -7.7$ (c, 0.18 in H₂O). λ_{\max} 243 (ϵ 8900); 285 (ϵ 9300) (H₂O).


Ohta, S. *et al.*, *J.O.C.*, 1997, **62**, 6452-6453 (*isol, uv, ir, pmr, cmr*)

Fujita, M. *et al.*, *Tetrahedron*, 2001, **57**, 1229-1234 (*isol, activity*)

Ohta, E. *et al.*, *Tetrahedron*, 2001, **57**, 4699-4703 (*Mg salt, isol, cd, pmr, cmr, ms*)

Ancorinoside B A-995

[334519-25-4]



$C_{41}H_{69}NO_{17}$ 847.993

Enolised β -triketone. Isol. from the sponge *Penares sollasi*. Powder. $[\alpha]_D^{24} +1.5$ (c, 0.1 in MeOH). λ_{\max} 282 (ϵ 8540) (MeOH).

Fujita, M. *et al.*, *Tetrahedron*, 2001, **57**, 1229-1234 (*isol, pmr, cmr, uv*)

Ancorinoside C A-996

[334519-26-5]

As Ancorinoside B, A-995 with R = CH₃


$C_{42}H_{71}NO_{17}$ 862.019

Enolised β -triketone. Isol. from the sponge *Penares sollasi*. Powder. $[\alpha]_D^{24} +2.8$ (c, 0.1 in MeOH). λ_{\max} 282 (ϵ 8430) (MeOH).

Fujita, M. *et al.*, *Tetrahedron*, 2001, **57**, 1229-1234 (*isol, pmr, cmr, uv*)

Ancorinoside D A-997

[334519-27-6]




$C_{43}H_{71}NO_{17}$ 874.03

Enolised β -triketone. Isol. from the sponge *Penares sollasi*. Powder. $[\alpha]_D^{24} -5.2$ (c, 0.1 in MeOH). λ_{\max} 282 (ϵ 9630) (MeOH).

Fujita, M. *et al.*, *Tetrahedron*, 2001, **57**, 1229-1234 (*isol, pmr, cmr, uv*)

Andersonine A-998

[119419-69-1]




$C_{39}H_{54}N_2O_{12}$ 742.862

Mixt. of regioisomers. Alkaloid from the aerial parts of *Delphinium andersonii* (Ranunculaceae). Amorph. $[\alpha]_D +34$ (c, 1.0 in CHCl₃). Possibly an artifact formed from Nudicauline in G-91 during the isol. process.

Pelletier, S.W. *et al.*, *Heterocycles*, 1988, **27**, 2387 (*isol, ir, pmr, cmr, ms, struct*)

Andesine† A-999

[96158-02-0]



$C_{20}H_{21}NO_6$ 371.389


Alkaloid from the stems of *Berberis*

darwinii and the stems and twigs of *Berberis actinacantha* (Berberidaceae). Amorph. red solid.

Weiss, I. *et al.*, *Heterocycles*, 1985, **23**, 301 (isol, uv, ir, pmr, ms, struct)

Andrachnidine A-1000

2-(2-Hydroxypentyl)-6-(2-oxopropyl)pyperidine
[274925-36-9]



Absolute Configuration

$C_{13}H_{25}NO_2$ 227.346

Alkaloid from *Andrachne aspera*. Oil. $[\alpha]_D^{22}$ -20 (c, 1.6 in $CHCl_3$).

Mill, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 762-764 (isol)

Shu, C. *et al.*, *J.A.C.S.*, 2003, **125**, 2878-2879 (synth)

Jung, H.H. *et al.*, *J.O.C.*, 2007, **72**, 7359-7366 (synth)

Krishna, P.R. *et al.*, *Tet. Lett.*, 2007, **48**, 7279-7282 (synth)

Andrachnine A-1001

[11023-51-1]


$C_{11}H_{17}NO_2$ 195.261

Struct. unknown. Alkaloid from subterranean parts of *Andrachne rotundifolia* (Euphorbiaceae). Cryst. (C_6H_6). Mp 97-99°.

Vil'yams, V.V. *et al.*, *Khim. Prir. Soedin.*, 1966, **2**, 257-260; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **2**, 207

Andrangine A-1002

[52659-54-8]



Absolute Configuration

$C_{19}H_{24}N_2O$ 296.411


Abs. config. enantiomeric with Vallesamidine, V-8. Alkaloid from *Craspidospermum verticillatum* and *Crioceris dipladeniiflorus* (Apocynaceae). Cryst. (hexane). Mp 132°. $[\alpha]_D^{20}$ -42 (c, 1 in $CHCl_3$). λ_{max} 248 (log ϵ 4.12); 299 (log ϵ 3.71) (EtOH).

Cavé, A. *et al.*, *Tet. Lett.*, 1973, 5081-5084 (cmr)

Kan-Fan, C. *et al.*, *Bull. Soc. Chim. Fr.*, 1974, 2839-2841 (isol, uv, ms, pmr, struct)

Andranginine A-1003

[53912-96-2]



$C_{21}H_{22}N_2O_2$ 334.417

Unusual type of iboga alkaloid showing C18 → C17 in place of C 16 → C21 cyclisation. Alkaloid from *Craspidospermum verticillatum* var. *petiolarie* (Apocynaceae). Mp 240°. Racemic.

Kan-Fan, C. *et al.*, *Chem. Comm.*, 1974, 164 (uv, ir, pmr, cmr, ms, struct, synth)


Massiot, G. *et al.*, *J.A.C.S.*, 1975, **97**, 3277 (pmr)

Andriamialisoa, R.Z. *et al.*, *Tetrahedron*, 1975, **31**, 2347 (synth)

Riche, P.G. *et al.*, *Acta Cryst. B*, 1979, **35**, 666 (cryst struct)

Andrimide A-1004

[108868-95-7]



$C_{27}H_{33}N_3O_5$ 479.575

Peptide antibiotic. Prod. by a bacterial symbiont from *Nilaparvata lugens*, a marine *Pseudomonas fluorescens* and an *Enterobacter* sp. Active against *Xanthomonas* sp. Mp 172-173.5°. λ_{max} 297 (ϵ 38000) (MeOH/AcOH) (Derep). λ_{max} 297 (ϵ 51100) (MeOH/NaOH) (Derep). λ_{max} 297 (ϵ 41200) (MeOH) (Derep).

3-Epimer, 3-hydroxy: *Moiramide C*

[155233-32-2]

$C_{27}H_{33}N_3O_6$ 495.574

Metab. of a marine *Pseudomonas fluorescens*. Amorph. solid. Sol. MeOH, EtOAc. λ_{max} 289 (ϵ 29711) (MeOH) (Berdy).

Fredenhagen, A. *et al.*, *J.A.C.S.*, 1987, **109**, 4409

Kenny, P.T. *et al.*, *Pestic. Sci.*, 1987, **27**, 117 (isol)


McWhorter, W. *et al.*, *Chem. Comm.*, 1989, 299 (synth)

Rama Rao, A.V. *et al.*, *Tet. Lett.*, 1991, **32**, 4393 (synth)

Needham, J. *et al.*, *J.O.C.*, 1994, **59**, 2058-2063 (*Andrimid*, *Moiramide C*)

Davies, S.G. *et al.*, *J.C.S. Perkin 1*, 1998, 2635-2643 (synth)

Androcymbine A-1005



(-)-form

$C_{21}H_{25}NO_5$ 371.432

(-)-form [2115-98-2]

Alkaloid from *Androcymbium melanthioides* var. *stricta* (Liliaceae). Mp

199-201°. $[\alpha]_D^{22}$ -260 ($CHCl_3$). λ_{max} 211 (log ϵ 4.6); 241 (log ϵ 4.28); 280 (log ϵ 3.7) (EtOH).

Me ether: **O-Methylandrocymbine**

[6890-90-0]

[38739-42-3, 31735-12-3]

$C_{22}H_{27}NO_5$ 385.459

Alkaloid from *Colchicum autumnale* and *Colchicum szovitsii* (Liliaceae).

Cryst. (Et₂O/hexane). Mp 176-178°.

$[\alpha]_D^{24}$ -295 (c, 0.13 in $CHCl_3$). $[\alpha]_D$ -140 ($CHCl_3$).

Me ether, O²-de-Me: **Isoandrocymbine**

[13205-97-5]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from *Colchicum stevenii*.

Me ether, O⁴-de-Me: **Collutine**

[20055-75-8]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from *Colchicum luteum* (Liliaceae). Cryst. (Me₂CO). Mp 192-194°. $[\alpha]_D$ -182 (c, 1.7 in $CHCl_3$). λ_{max} 238 (log ϵ 4.3); 275 (sh) (log ϵ 2.81) (MeOH).

Me ether, O⁴-de-Me, N-oxide: **Collutine N-oxide**

[102719-92-6]

$C_{21}H_{25}NO_6$ 387.432

Alkaloid from *Colchicum luteum* (Liliaceae). Cryst. (Me₂CO). Mp 217-219°. $[\alpha]_D$ -204 (c, 0.36 in $CHCl_3$).

(+)-form

Me ether, O²-de-Me: **Alkaloid CC 10**

[35320-74-2]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Prisms (EtOAc). Mp 209-212° (200-204°). $[\alpha]_D^{22}$ +308 (c, 0.87 in $CHCl_3$). $[\alpha]_D^{22}$ +336 (c, 0.10 in EtOH). Enantiomer of Isoandrocymbine.

(±)-form [31730-26-4]

Noncryst. λ_{max} 240 (log ϵ 4.18); 279 (log ϵ 3.62) (MeOH).

(ξ)-form

Me ether, 7ξ-alcohol: **Szovitsidine**

[57912-36-4]

$C_{22}H_{29}NO_5$ 387.475

Alkaloid from *Colchicum szovitsii* (Liliaceae). No phys. props. reported.

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 255-267 (isol, ir, uv)

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413 (*Alkaloid CC 10*, isol, uv)

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540 (*Alkaloid CC 10*, isol, uv, ir)

Battersby, A.R. *et al.*, *J.C.S.(C)*, 1971, 3514 (*Alkaloid CC 10*, uv, ir, struct)

Kametani, T. *et al.*, *J.O.C.*, 1971, **36**, 3729-3733; 3733-3736 (synth)

Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1972, 1736-1740 (ir, pmr, ms, struct)

Mukhamed'yarova, N.L. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 781-783 (*Collutine*, isol, uv, pmr, struct)


Yusupov, M.K. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 271; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 289 (*Szovitsidine*)

Chommadov, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 770-773 (*Collutine N-oxide*)

Nasreen, A. *et al.*, *Tet. Lett.*, 1996, **37**, 8161-8164 (*Isoandrocybine*)
 Barker, A.C. *et al.*, *J.C.S. Perkin 1*, 1998, 2989-2994; 3003-3009 (*isol. synth. Me ether*)
 Al-Mahmoud, M.S. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 153-160 (*Isoandrocybine*)

Angchibangkinge A-1006

[187109-61-1]




$C_{35}H_{34}N_2O_5$ 562.664
 Alkaloid from stem bark of *Pachygone dasycarpa*. Exhibits antiplasmodial activity. $[\alpha]_D^{20} +450$ (c, 0.1 in $CHCl_3$). λ_{max} 234 (sh) (log ϵ 4.67); 290 (log ϵ 3.71) (MeOH).

Me ether: [187086-18-6] Shows antiplasmodial activity. $[\alpha]_D^{20} +392$ (c, 0.05 in $CHCl_3$).

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1997, **60**, 258 (*isol, uv, cd, pmr, ms, struct*)

Angelicastigmine A-1007

[312584-11-5]




$C_{21}H_{22}N_2O_4$ 366.416
 Alkaloid from the roots of *Angelica polymorpha*. Brownish amorph. solid. $[\alpha]_D^{20} -92$ (c, 0.79 in MeOH). λ_{max} 212 (ϵ 18740); 268 (ϵ 10750); 292 (ϵ 6750) (MeOH).

Pachaly, P. *et al.*, *Pharmazie*, 2000, **55**, 777-778 (*isol, pmr, cmr, ms*)

6-Angeloyloxyditropan-3-yl itaconate A-1008

[182015-05-0]




$C_{26}H_{38}N_2O_6$ 474.596
 Alkaloid from the leaves of *Schizanthus litoralis*. Oil. $[\alpha]_D^{20} -10$ (c, 0.09 in EtOH).

Muñoz, O. *et al.*, *Phytochemistry*, 1996, **43**, 709-713 (*isol, pmr, cmr, ms*)

Anguibactin A-1009

2-(2,3-Dihydroxyphenyl)-4,5-dihydro-N-hydroxy-N-[2-(1H-imidazol-4-yl)ethyl]-4-thiazolecarboxamide, 9CI
 [117308-63-1]




$C_{15}H_{16}N_4O_4S$ 348.382
 Isol. from the iron-deficient cultures of a marine *Vibrio anguillarum*. Siderophore. λ_{max} 216 ; 266 ; 327 (MeOH) (Berdy). λ_{max} 214 ; 260 (ϵ 11000); 315 ; 410 (pH 7 buffer) (Berdy). λ_{max} 214 ; 258 ; 335 ; 415 (pH 10 buffer) (Berdy).

Actis, L.A. *et al.*, *J. Bacteriol.*, 1986, **167**, 57 (*isol*)

Jalal, M.A.F. *et al.*, *J.A.C.S.*, 1989, **111**, 292-296 (*Anguibactin*)

Angusteine A-1010

[54163-65-4]




$C_{21}H_{24}N_2O_3$ 352.432
 Stereochem. undefined, position of double bond in side chain at C-20 unknown. Alkaloid from *Amsonia angustifolia* (Apocynaceae).

Kocsis, A. *et al.*, *Acta Pharm. Hung.*, 1974, **44**, 70-73; *CA*, **81**, 120843w (*isol, uv, ir, ms, struct*)

Angusticaline A-1011

[120374-19-8]




$C_{44}H_{56}N_4O_6$ 736.95
 Alkaloid from the leaves of *Alstonia angustifolia* (Apocynaceae). $[\alpha]_D^{20} +6$ (c, 1.1 in MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1988, **27**, 3955 (*isol, uv, ir, pmr, cmr, ms, struct*)

Angustidine A-1012

8,13-Dihydro-2-methylindolo[2',3':3,4]pyrido[1,2-b][2,7]naphthyridin-5(7H)-one, 9CI
 [40217-50-3]



$C_{19}H_{15}N_3O$ 301.347
 Alkaloid from the leaves of *Strychnos angustiflora* and from several other *Strychnos*, *Mitragyna*, *Nauclea* and *Uncaria* spp. (Loganiaceae, Rubiaceae). Yellow needles ($CHCl_3$). Mp 309-310°.

Au, T.Y. *et al.*, *J.C.S. Perkin 1*, 1973, 13 (*isol, uv, ir, pmr, struct*)

Phillipson, J.D. *et al.*, *Phytochemistry*, 1974, **13**, 973 (*isol*)

Ninomiya, I. *et al.*, *J.C.S. Perkin 1*, 1976, 1865 (*synth, ir, pmr*)


Repke, D.B. *et al.*, *Tetrahedron*, 1989, **45**, 2541 (*synth*)

Delaupe, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, Strychnos*)

Abreu, P. *et al.*, *Heterocycles*, 1998, **48**, 885-891 (*isol, uv, ir, pmr, cmr, ms*)

Angustifoline† A-1013

Decahydro-4-(2-propenyl)-1,5-methano-8H-pyrido[1,2-a][1,5]diazocin-8-one, 9CI. Jamaicensine
 [550-43-6]



$C_{14}H_{22}N_2O$ 234.341

(-)-form

Alkaloid from *Lupinus angustifolius*, *Lupinus polyphyllus*, *Lupinus albus*, *Ormosia jamaicensis* and *Ormosia panamensis* (Fabaceae). Active against gram-positive bacteria. Mp 80.5-81°. $[\alpha]_D^{20} -7.5$ (+5.2) (EtOH).

Hydrochloride: Mp 96-97°.

Picrate: Mp 182-185°.

N-Formyl: N-Formylangustifoline

[4697-81-8]

$C_{15}H_{22}N_2O_2$ 262.351

Alkaloid from combined leaf and hypocotyl extracts of *Lupinus polyphyllus* (Fabaceae).

N-Ac: Mp 151-152°.

N-Methoxycarbonyl: Angustifoline N-carboxymethyl ester

[145889-87-8]

$C_{16}H_{24}N_2O_3$ 292.377

Alkaloid from combined leaf and

hypocotyl extracts of *Lupinus polyphyllus* (Fabaceae).

N-Ethoxycarbonyl: Angustifoline N-carboxyethyl ester

[145889-88-9]
C₁₇H₂₆N₂O₃ 306.404

Alkaloid from combined leaf and hypocotyl extracts of *Lupinus polyphyllus* (Fabaceae).

N-Me: N-Methylangustifoline

[4697-79-4]
C₁₅H₂₄N₂O 248.367

A principal alkaloid of the oil of *Lupinus mutabilis* (Fabaceae). Mp 86° (synthetic). [α]_D²⁰ +7.24 (c, 2.35 in EtOH) (synthetic). Isoln. is poorly documented.

Deoxo: Deoxoangustifoline. Deoxyangustifoline

[72710-32-8]
C₁₄H₂₄N₂ 220.357

Alkaloid from *Thermopsis mongolica* (Fabaceae).

8,9-Didehydro: Dehydroangustifoline. Alkaloid W102

[85530-32-1]
C₁₄H₂₀N₂O 232.325

Alkaloid from *Lupinus albus* and *Lupinus angustifolius* (Fabaceae). Mp 105°.

10-Epimer: Isoangustifoline

[82189-28-4]
C₁₄H₂₂N₂O 234.341

Alkaloid from phloem exudate of *Lupinus angustifolius*, and from *Lupinus polyphyllus*. Identified only by ms.

(+)-form [150283-69-5]

Alkaloid from seeds of *Lupinus termis*. Minor constit. in seeds of *Lupinus albus* cv. BAC. Oil. Mp 77°. [α]_D²⁰ +4.9 (sol. unspecified). [α]_D²⁵ +5.2 (c, 0.1 in MeOH).

Bohlmann, F. *et al.*, *Chem. Ber.*, 1960, **93**, 1956-1960; 1962, **95**, 944-948 (*isol, ir, struct, synth*)

Lloyd, H.A. *et al.*, *J.O.C.*, 1960, **25**, 1959-1962 (*isol*)

Marion, L. *et al.*, *Tet. Lett.*, 1960, **No. 19**, 1-5 (*struct, Angustifoline, Dehydroangustifoline*)

Fales, H.M. *et al.*, *J.A.C.S.*, 1970, **92**, 1590-1597 (*ms*)

Lloyd, H.A. *et al.*, *Phytochemistry*, 1975, **14**, 155-159 (*biosynth*)

Bratek-Wiewiorowska, M.D. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1979, **55**, 69-87 (*ir, pmr, conformn*)

Hatzold, T. *et al.*, *Fette, Seifen, Anstrichm.*, 1982, **84**, 59-64 (*isol, N-Methylangustifoline*)

Wink, M. *et al.*, *Planta Med.*, 1982, **44**, 15-20 (*Isoangustifoline*)

Fraser, A.M. *et al.*, *Chem. Comm.*, 1984, 1477-1479 (*biosynth*)

Tyski, S. *et al.*, *J. Plant Physiol.*, 1988, **133**, 240-242 (*activity*)

Mohamed, M.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1578-1580 (*(+)-form, isol*)

Christov, V. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1413-1415 (*Deoxoangustifoline*)

Veen, G. *et al.*, *Phytochemistry*, 1992, **31**, 4343-4345 (*Lupinus polyphyllus* constits)


Wysocka, W. *et al.*, *Planta Med.*, 1993, **59**, 289 (*(+)-form, isol, cmr*)

Bäumel, P. *et al.*, *Z. Naturforsch., C*, 1993, **48**, 436-443 (*Isoangustifoline*)

Wysocka, W. *et al.*, *Monatsh. Chem.*, 1994, **125**, 1267-1272 (*pmr, cmr, conformn*)

Angustilobine A

[112464-20-7]



C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia angustiloba*, *Alstonia congensis* and *Alstonia pneumatophora* (Apocynaceae). [α]_D +84 (c, 0.45 in CHCl₃).

15-Hydroxy: 15-Hydroxyangustilobine A

[112464-21-8]
C₂₀H₂₂N₂O₄ 354.405


Alkaloid from *Alstonia angustiloba* and *Alstonia pneumatophora* (Apocynaceae). [α]_D -160 (c, 0.5 in CHCl₃).

Zeche, M. *et al.*, *J. Nat. Prod.*, 1987, **50**, 714 (*isol*)

Caron, C. *et al.*, *Phytochemistry*, 1989, **28**, 1241 (*isol*)

Angustilobine B

[112448-50-7]



C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia angustifolia*, *Alstonia congensis* and *Alstonia pneumatophora* (Apocynaceae). [α]_D -136 (c, 0.5 in CHCl₃).

N^b-Oxide: Angustilobine B N^b-oxide

[123085-03-0]
C₂₀H₂₂N₂O₄ 354.405

Alkaloid from *Alstonia congensis* and *Alstonia scholaris*. [α]_D +97 (c, 0.2 in CHCl₃).

N^b-Me: N^b-Methylangustilobine B

C₂₁H₂₅N₂O₃⁺ 353.44

Alkaloid from *Alstonia scholaris*. Counterion not specified.

Parent acid: Angustilobine B acid

[126594-74-9]
C₁₉H₂₀N₂O₃ 324.379

Alkaloid from leaves of *Alstonia scholaris* (Apocynaceae). Fine prisms. Mp 230-235° dec. [α]_D²⁴ -117.4 (c, 0.66 in Py).

15-Epimer(?): Alstonamine

[1394-99-6]
C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Amorph. solid. Mp 126°. [α]_D²⁰ +46 (c, 2 in CHCl₃). Same gross struct. assigned to Alstonamine as to Angustilobine B but with undetd. 15-conf. Clearly not the same as Angustilobine B.

A-1014

Zeche, M. *et al.*, *J. Nat. Prod.*, 1987, **50**, 714 (*Angustilobine B*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1987, **26**, 2139 (*Alstonamine*)

Caron, C. *et al.*, *Phytochemistry*, 1989, **28**, 1241 (*oxide*)


Yamauchi, T. *et al.*, *Phytochemistry*, 1990, **29**, 3321; 3547 (*isol, pmr, cmr*)

Macabeo, A.P.G. *et al.*, *Phytochemistry*, 2005, **66**, 1158-1162 (*N-Methylangustilobine B*)

Angustilodine

A-1016

[740811-97-6]



C₂₀H₂₄N₂O₄ 356.421


Alkaloid from the leaves of *Alstonia angustiloba*. Oil. [α]_D -622 (c, 0.16 in CHCl₃). λ_{max} 224 (log ε 4.23); 281 (log ε 3.6) (EtOH).

Kam, T.-S. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 366-369 (*isol, pmr, cmr*)

Angustimalal

A-1017

[714252-05-8]



C₁₃H₁₉NO₃ 237.298


Alkaloid from the bark of *Alstonia macrophylla*. Oil. [α]_D +78 (c, 0.06 in CHCl₃). Prob. derived from a macroline-type precursor. λ_{max} 211 (log ε 3.09); 256 (log ε 2.42) (EtOH).

Kam, T.-S. *et al.*, *Tetrahedron*, 2004, **60**, 3957-3966 (*isol, pmr, cmr, ms*)

Angustimaline

A-1018

[186772-63-4]



C₁₃H₁₉NO₃ 237.298


Alkaloid from the stem bark of *Alstonia angustifolia*. Amorph. [α]_D +115 (c, 0.1 in CHCl₃). Prob. derived from a macroline-type precursor. λ_{max} 257 (log ε 4.11) (No solvent reported).

Kam, T.-S. *et al.*, *Tet. Lett.*, 1997, **38**, 477 (*isol, uv, ir, pmr, cmr, ms, struct*)

Angustine

A-1019

1-Ethenyl-8,13-dihydroindolo[2',3':3,4]pyrido[1,2-b][2,7]naphthyridin-5(7H)-one, 9CI
[40041-96-1]



$C_{20}H_{15}N_3O$ 313.358

Alkaloid from *Strychnos angustiflora* and other *Strychnos* spp., *Mitragyna*, *Nauclea* and *Uncaria* spp. (Loganiaceae, Rubiaceae). Shows anticancer props. Yellow plates. Mp 283-284° dec., (>340°). λ_{max} 220 (log ϵ 4.54); 255 (log ϵ 4.32); 291 (log ϵ 4.1); 304 (log ϵ 4.03); 380 (log ϵ 4.58); 400 (log ϵ 4.59) (EtOH).

3,14-Dihydro: 3,14-Dihydroangustine.

13b,14-Dihydroangustine

[116965-55-0]

$C_{20}H_{17}N_3O$ 315.374

Alkaloid from the root bark of *Nauclea pobeguini* (Rubiaceae). Shows anticancer props. The synthetic racemate had Mp 305-306°.

18,19-Dihydro: 18,19-Dihydroangustine

[40042-52-2]

$C_{20}H_{17}N_3O$ 315.374

Alkaloid from leaves of *Nauclea orientalis* (Rubiaceae). Exhibits *in vitro* antiproliferative activity. Orange cryst. solid. λ_{max} 221 (log ϵ 4.44); 252 (log ϵ 4.15); 290 (log ϵ 3.9); 300 (log ϵ 3.81); 375 (log ϵ 4.46); 395 (log ϵ 4.48) (EtOH).

3,14,18,19-Tetrahydro: 3,14,18,19-Tetrahydroangustine

[52410-21-6]

$C_{20}H_{19}N_3O$ 317.39

Alkaloid from leaves of *Nauclea orientalis* (Rubiaceae). Exhibits *in vitro* antiproliferative activity. Shows anticancer props. Yellowish amorph. solid.

10-Hydroxy: 10-Hydroxyangustine

[142115-23-9]

$C_{20}H_{15}N_3O_2$ 329.357

Alkaloid from leaves of *Nauclea orientalis* and from *Nauclea latifolia* (Rubiaceae). Exhibits *in vitro* antiproliferative activity. Orange cryst. solid. λ_{max} 205 (ϵ 20000); 253 (ϵ 6300); 303 (ϵ 4000); 310 (ϵ 4000); 384 (ϵ 12500); 404 (ϵ 12500) (EtOH) (Berdy).

Au, T.Y. et al., *J.C.S. Perkin I*, 1973, 13-16

(isol, uv, ir, pmr, struct)

Phillipson, J.D. et al., *Phytochemistry*, 1974,

13, 973-978 (isol, uv, ir, pmr)

Kametani, T. et al., *J.O.C.*, 1976, 41, 2542-

2545 (synth, uv, ir, pmr)

Zeches, M. et al., *J. Nat. Prod.*, 1985, 48, 42-46

(3,14-Dihydroangustine)

Repke, D.B. et al., *Tetrahedron*, 1989, 45, 2541-

2550 (synth, Angustine, 3,14-

Dihydroangustine)

Erdelmeier, C.A.J. et al., *Planta Med.*, 1990,

56, 507 (activity)

Erdelmeier, C.A.J. et al., *Planta Med.*, 1992,

58, 43-48 (10-Hydroxyangustine, 18,19-

Dihydroangustine, 3,14,18,19-

Tetrahydroangustine)

Abreu, P. et al., *Heterocycles*, 1998, 48, 885-


891 (isol, uv, ir, pmr, cmr, ms)

Lavilla, R. et al., *Eur. J. Org. Chem.*, 1999, 373-

378 (synth)

Angustoline

A-1020



$C_{20}H_{17}N_3O_2$ 331.373

(R)-form [40041-95-0]

Alkaloid from *Strychnos angustiflora* and several other *Strychnos* spp., and from *Uncaria* spp. (Loganiaceae, Rubiaceae). Yellow needles (CHCl₃/MeOH). Mp 310-314°. $[\alpha]_D^{20}$ -34 (CHCl₃).

Me ether: 19-O-Methylangustoline

[132074-99-8]

$C_{21}H_{19}N_3O_2$ 345.4

Alkaloid from the seeds of *Campotheca acuminata* (Nyssaceae). Cytotoxic agent. Yellow cryst. (Me₂CO). Mp 222-225°. $[\alpha]_D^{20}$ -6.3 (c, 0.008 in MeOH). λ_{max} 204 (ϵ 56234); 370 (ϵ 17380); 395 (ϵ 14125) (MeOH) (Berdy).

3,14-Dihydro: 3,14-Dihydroangustoline

[142115-22-8]

[142186-12-7]

$C_{20}H_{19}N_3O_2$ 333.389

Alkaloid from leaves of *Nauclea orientalis* (Rubiaceae). Yellowish amorph. solid. $[\alpha]_D^{20}$ -172.5 (c, 0.20 in EtOH). $[\alpha]_D^{20}$ -301.4 (c, 0.33 in EtOH). Two diastereoisomers isol. λ_{max} 204; 265; 373; 396 (EtOH) (Berdy).

Ketone: Nauclefine

[54698-29-2]

$C_{20}H_{15}N_3O_2$ 329.357

Alkaloid from the root bark of *Nauclea latifolia* (Rubiaceae). Orange cryst. (MeOH). Mp 310°.

10-Hydroxy: Naucleofficine E

$C_{20}H_{17}N_3O_3$ 347.373

Alkaloid from the stems of *Nauclea officinalis*. Orange-yellow powder. Mp 279-281°. $[\alpha]_D^{20}$ -29.3 (c, 0.21 in MeOH). Probable abs. config. λ_{max} 220 (log ϵ 4.31); 251 (log ϵ 4.02); 307 (log ϵ 3.9); 400 (log ϵ 4.67) (MeOH).

(±)-form

Synthetic. Pale-yellow cryst. (MeOH/CHCl₃). Mp 292-294°.

(ξ)-form**O-Ac: O-Acetylanguustoline**

[40042-53-3]

$C_{22}H_{19}N_3O_3$ 373.41

Alkaloid from the root bark of *Nauclea*

pobeguini (Rubiaceae). Yellow needles + 1.5H₂O (CHCl₃). Mp 148-152° (synthetic). Nat. alkaloid obt. only in small amt., no info. on opt. rotn. or abs. config.

Et ether: 19-O-Ethylanguustoline

[210895-10-6]

$C_{22}H_{21}N_3O_2$ 359.427

Alkaloid from the stem bark of *Sarcocephalus latifolius*. Mp 230-240°. $[\alpha]_D^{20}$ -0.01 (CHCl₃). Appears to be a virtual racemate.

Au, T.Y. et al., *J.C.S. Perkin I*, 1973, 13-16

(isol, uv, ir, pmr, ms, struct)

Phillipson, J.D. et al., *Phytochemistry*, 1974,

13, 973-978 (occur)

Hotellier, F. et al., *Phytochemistry*, 1975, 14,

1407-1409 (Nauclefine, isol, uv, ir, pmr, ms, struct)

Ninomiya, I. et al., *J.C.S. Perkin I*, 1976,

1865-1868 (synth, ir, pmr)

Zeches, M. et al., *J. Nat. Prod.*, 1985, 48, 42-46

(O-Acetylanguustoline)

Lin, L.-Z. et al., *Phytochemistry*, 1990, 29,

2744-2746 (19-O-Methylanguustoline)

Erdelmeier, C.A.J. et al., *Planta Med.*, 1992,

58, 43-48 (3,14-Dihydroanguustoline)

Abreu, P. et al., *Heterocycles*, 1998, 48, 885-

891 (19-O-Ethylanguustoline, isol, uv, ir, ord,

pmr, cmr, abs config)

Lavilla, R. et al., *Eur. J. Org. Chem.*, 1999, 373-

378 (synth)

Sun, J. et al., *Phytochemistry*, 2008, 69, 1405-

1410 (Naucleofficine E)

Anhalonine


A-1021

6,7,8,9-Tetrahydro-4-methoxy-9-methyl-

1,3-dioxolo[4,5-h]isoquinoline, 9CI.

1,2,3,4-Tetrahydro-6-methoxy-1-methyl-

7,8-methylenedioxyisoquinoline



$C_{12}H_{15}NO_3$ 221.255

(R)-form

Obt. by resoln. Mp 84.5-85.5°. $[\alpha]_D^{25}$ +56.7 (CHCl₃).

(S)-form [519-04-0]

Alkaloid from *Anhalonium lewinii* (*Lophophora williamsii*), *Gymnocalycium lecanum* and *Trichocereus terscheckii* (Cactaceae). Also obt. by resoln. and by synth. Cryst. (Et₂O/petrol). Mp 85-86°. $[\alpha]_D^{25}$ -56.3 (CHCl₃). $[\alpha]_D^{25}$ -62.9 (c, 1 in MeOH). $[\alpha]_D^{25}$ -51 (c, 1 in 1M HCl).

Hydrochloride: Mp 258-259°. $[\alpha]_D^{25}$ -40 (c, 1 in 50% EtOH).

Hydrobromide:

Cryst. (H₂O). Mp 270-271°. $[\alpha]_D^{25}$ -30 (c, 1 in H₂O).

N-Me: Lophophorine. 6,7,8,9-Tetrahydro-4-methoxy-8,9-dimethyl-1,3-dioxolo[4,5-h]isoquinoline, 9CI. 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl-7,8-methylenedioxyisoquinoline

[17627-78-0]

$C_{13}H_{17}NO_3$ 235.282

Alkaloid from *Gymnocalycium gibbosum*, *Gymnocalycium lecanum*, *Lopho-*

phora diffusa and *Lophophora williamsii* (Cactaceae). Respiratory stimulant, convulsive agent. Oil. Bp_{0.05} 140-145°. [α]_D²⁵ -47.3 (CHCl₃).

► Toxic, LD₅₀ (rbt, ivn) 15-20 mg/kg. JI4800000

N-Me; hydrochloride:

Cryst. (EtOH). Mp 236-237°. [α]_D²⁵ -15.6 (c, 1 in H₂O).

N-Me, picrate: Mp 162-163°.

N-Formyl: N-Formylanhalonine

C₁₃H₁₅NO₄ 249.266

Isol. from *Anhalonium lewinii* (Cactaceae). Abs. config. of this isolate unknown; identified by glc/ms comparison with an authentic prepn.

N-Ac: N-Acetylanhalonine

C₁₄H₁₇NO₄ 263.293

Constit. of *Anhalonium lewinii* (Cactaceae). Identified by glc/ms comparison.

N-Et: Peyophorine. N-Ethylanhalonine [22030-12-2]

C₁₄H₁₉NO₃ 249.309

Alkaloid from *Anhalonium lewinii* (Cactaceae). Abs. config. not determined.

N-Et, picrate: Mp 155-156°.

N,N-Di-Me: 6,7,8,9-Tetrahydro-4-methoxy-8,8,9-trimethyl-1,3-dioxolo[4,5-h]isoquinolinium(1+), 9CI. Lophotine [19267-94-8]

[19445-63-7]

C₁₄H₂₀NO₃⁺ 250.317

Alkaloid from *Anhalonium lewinii*.

Cryst. (EtOH/EtOAc) (as iodide).

Mp 223° (softens) Mp 240-242°

(iodide). Abs. config. not certain.

Double Mp attributed to thermal racemisation.

(±)-form

Synthetic. Cryst. (MeOH) (as hydrochloride). Mp 262-264° (hydrochloride).

N-Me; hydrobromide:

Cryst. (EtOH). Mp 221-222°.

N,N-Di-Me, iodide: Mp 242-243°.

Heffter, A. et al., *Ber.*, 1896, **29**, 216 (*isol*)

Späth, E. et al., *Monatsh. Chem.*, 1923, **44**, 103

Späth, E. et al., *Ber.*, 1935, **68**, 501; 1663

(*struct, synth, resoln*)

Battersby, A.R. et al., *J.C.S.*, 1960, 1214 (*abs config*)

Battersby, A.R. et al., *Tet. Lett.*, 1967, 563 (*biosynth*)

Kapadia, G.J. et al., *Chem. Comm.*, 1968, 1688 (*N-Formylanhalonine, N-Acetylanhalonine*)

Kapadia, G.J. et al., *J. Pharm. Sci.*, 1968, **57**, 254; 2017 (*isol, Lophotine, Peyophorine*)

Brossi, A. et al., *J.A.C.S.*, 1971, **93**, 6248

(*synth, ord, cd, ms, uv, pmr, abs config, cryst struct*)

Cymerman Craig, J. et al., *J.A.C.S.*, 1977, **99**, 7996 (*uv, cd*)

Mata, R. et al., *Phytochemistry*, 1983, **22**, 1263 (*cmr, Lophophorine*)

Anhweiaconitine

A-1022

[1354-38-7]

Struct. unknown

C₃₀H₄₁NO₉ 559.655

Probably a C₁₉-diterpene alkaloid. Alka-

loid from the roots of the Chinese drug Tzu-Tsao-Wu (*Aconitum* spp.) (Ranunculaceae). Mp 199°. [α]_D¹¹ +25.2 (c, 1 in CHCl₃). Conts. N-Me, 4 OH, 3 OMe and PhCOO groups.

Nitrate: Mp 188-189°.

Perchlorate: Mp 226°.

Methiodide: Mp 210-213°.


Tetra-Ac: Mp 181-183°.

Chu, J.-H. et al., *Huaxue Xuebao*, 1959, **25**, 214-217; *CA*, **54**, 4642e

Anhweidelphinine

A-1023

[102503-60-6]



C₃₅H₄₄N₂O₁₀ 652.74

Alkaloid from the roots of *Delphinium anhweinsae* and whole plants of *Delphinium nuttallianum*. Defective diag. in CAS.

O¹⁴-De-Me, 14-Ac: 14-O-Demethyl-14-acetylanhweidelphinine

C₃₆H₄₄N₂O₁₁ 680.75

Alkaloid from *Delphinium pentagynum*. Amorph. solid. [α]_D²⁵ +67.1 (c, 0.16 in CHCl₃).

O¹⁴-De-Me, 14-O-(2-methylpropanoyl): 14-O-Demethyl-14-isobutyrylanhweidelphinine

C₃₈H₄₈N₂O₁₁ 708.804

Alkaloid from *Delphinium pentagynum*. Amorph. solid. [α]_D²⁵ +65.2 (c, 0.14 in CHCl₃).

Jin, J. et al., *Zhongcaoyao*, 1986, **17**, 1; *CA*,

105, 3523v

Bai, Y. et al., *Heterocycles*, 1989, **29**, 1017 (*isol*)


Joshi, B.S. et al., *J. Nat. Prod.*, 1990, **53**, 1028 (*config*)

Diaz, J.G. et al., *Phytochemistry*, 2004,

65, 2123-2127 (*14-demethyl-14-acyl derivs*)

Anhydroberberillic acid

A-1024



C₂₀H₁₇NO₈ 399.356

Alkaloid from the roots of *Phellodendron amurense* var. *wilsonii*. Powder (MeOH). Mp 252-253°. λ_{\max} 224 ; 254 ; 299 ; 335 (MeOH).

Me ester:

C₂₁H₁₉NO₈ 413.383


Alkaloid from the roots of *Phellodendron amurense* var. *wilsonii*. Powder (MeOH). Mp 183-184°. λ_{\max} 224 ; 260 ; 316 (MeOH).

Chiang, Y.-L. et al., *Heterocycles*, 2006, **68**, 339-345 (*isol, pmr, cmr, ms*)

Anhydrodehydrotylophorinidine

A-1025

[52985-89-4]



C₂₂H₂₀NO₃⁺ 346.405

Trace quaternary alkaloid from the leaves and stems of *Tylophora asthmatica* (Asclepiadaceae). No phys. props. reported. Prob. an artifact.

Me ether: Anhydrodehydrotylophorinine.

Tylophoridine D

[52936-83-1]

[415898-72-5]

C₂₃H₂₂NO₃⁺ 360.432

Trace quaternary alkaloid from leaves and stems of *Tylophora asthmatica* and from *Tylophora atrofoliculata*. Yellow amorph. powder (as chloride). Mp 295-310° (chloride). [α]_D¹⁶ +0.5 (c, 1 in MeOH) (chloride). Prob. an artifact. λ_{\max} 257 (log ϵ 4.68); 284 (log ϵ 4.56); 309 (sh) (log ϵ 4.06); 340 (log ϵ 3.46) (MeOH).

2-Methoxy, Me ether: Dehydrotylophorine [52936-55-7]

C₂₄H₂₄NO₄⁺ 390.458

Quaternary alkaloid from leaves and stems of *Tylophora asthmatica* (Asclepiadaceae), also obt. by air oxidn. of Tylophorine in diffused daylight. Yellow needles + 2H₂O (MeOH) (as chloride). Mp 245-250° dec. (chloride). Prob. an artifact.

7-DEMethoxy, 2-methoxy, Me ether: Dehydroantofine

[126262-29-1]

C₂₃H₂₂NO₃⁺ 360.432

Quaternary alkaloid from the leaves, twigs, roots and fruit of *Cryptocarya phyllostemon* (Lauraceae). Yellow cryst. (MeOH) (as perchlorate). Mp 230° dec. (perchlorate). Poss. artifact. CAS no. refers to perchlorate.

Wiegreb, W. et al., *Annalen*, 1969, **721**, 154

(*Dehydroantofine, synth*)

Govindachari, T.R. et al., *Indian J. Chem.*, 1973, **11**, 1215

(*Anhydrodehydrotylophorinidine,*

Anhydrodehydrotylophorinine,

Dehydrotylophorine)


Cavé, A. et al., *Aust. J. Chem.*, 1989, **42**, 2243

(*Dehydroantofine, isol*)

Huang, X.S. et al., *Planta Med.*, 2004, **70**, 441-445 (*Tylophoridine D*)

Anhydrohapaloxindole A A-1026

[109217-16-5]

C₂₁H₂₁ClN₂O 352.862

Minor alkaloid from the blue-green alga *Hapalosiphon fontinalis*. Mp 123° dec. [α]_D²⁵ +150 (c, 0.4 in EtOH). Appears to be a ¹O₂ oxidn. prod. of Hapalindole A, H-48, the major alkaloid in this cyanophyte.

Isothiocyanate: Anhydrohapaloxindole B [123519-73-3]

C₂₁H₂₁ClN₂OS 384.928

Minor alkaloid from the blue-green alga *Hapalosiphon fontinalis*. Has -NCS replacing -NC.

Dechloro, isothiocyanate: Anhydrohapaloxindole M

[123498-03-3]

C₂₁H₂₂N₂OS 350.484


Minor alkaloid from *Hapalosiphon fontinalis*.

Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 3773

(Anhydrohapaloxindole A)

Moore, R.E. *et al.*, *Phytochemistry*, 1989, **28**, 1565 (Anhydrohapaloxindoles B,M)**Anhydroharringtonine** A-1027


[142735-74-8]

C₂₈H₃₅NO₈ 513.586

Alkaloid from *Cephalotaxus fortunei*. Shows significant antileukaemic activity.

Wang, D.Z. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 173; *CA*, **117**, 86698c**Anhydroholantogenin** A-1028

14,20:18,20-Diepoxypregnan-3-ol [28719-41-7]

C₂₁H₃₂O₃ 332.482

Aglycone from Holantossine B. Cryst. (Me₂CO). Mp 179°. [α]_D²⁰ -38 (CHCl₃).

4-Amino-2,4,6-trideoxy-3-O-methyl-β-D-ribo-hexopyranoside: **Holantossine B**

[28719-39-3]

C₂₈H₄₅NO₅ 475.667

Constit. of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (Me₂O) (as *N*-Ac). Mp 0° (-290° as *N*-Ac). [α]_D²⁰ -29 (CHCl₃) (as *N*-Ac).

4-Amino-2,4,6-trideoxy-3-O-methyl-α-L-arabino-hexopyranoside: **Holantossine D**

[33662-04-3]

C₂₈H₄₅NO₅ 475.667

Constit. of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (Me₂CO)(as Ac). Mp 262-263° (Ac). [α]_D²⁰ -67 (c, 6 in CHCl₃).

4-Amino-2,4,6-trideoxy-3-O-methyl-β-D-xylo-hexopyranoside: **Holantossine F**


[40738-41-8]

C₂₈H₄₅NO₅ 475.667

Isol. from *Holarrhena antidysenterica* (Apocynaceae). Mp 134° (as *N*-Ac).

[α]_D²⁰ -43 (c, 1 in CHCl₃) (as *N*-Ac).Schaffner, B.P. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 2266 (synth)Janot, M.-M. *et al.*, *Tetrahedron*, 1970, **26**, 1695 (isol)Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1971, 864 (isol)Goutarel, R. *et al.*, *Carbohydr. Res.*, 1972, **24**, 297 (Holantossine F)**Anhydrolycodoline** A-1029

[18834-79-2]



Absolute Configuration

C₁₆H₂₃NO 245.364

Alkaloid from *Lycopodium alopecuroides*, *Lycopodium carolinum*, *Lycopodium clavatum* var. *borbonicum*, *Lycopodium gni-dioides*, *Lycopodium inundatum*, *Lycopodium phlegmaria* and *Lycopodium saururus* (Lycopodiaceae). Oil; off-white cryst. (as hydrochloride). Mp 274-276° dec. (as hydrochloride). Darkens and dec. on exp. to air.

***N*-Me: Lycopodatine B**

[872610-91-8]

C₁₇H₂₆NO[⊕] 260.399

Alkaloid from *Lycopodium inundatum*. Solid. [α]_D³³ -36 (c, 0.3 in MeOH). Counterion not specified.

7-Hydroxy: Serratidine

[20084-82-6]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum* (Lycopodiaceae). Mp 143-144°. First *Lycopodium* alkaloid with a bridge-head OH group.

10β-Hydroxy: Lycoplegmine

[82841-97-2]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Mp 121-123°.

4,7-Dihydroxy: 4-Hydroxyserratidine

[485817-34-3]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*.

Prisms (Me₂CO/MeOH). Mp 230-232° dec. [α]_D²⁵ -0.7 (c, 0.18 in MeOH).

6α,7-Dihydroxy: 6α-Hydroxyserratidine

[485817-33-2]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*.

Prisms (Me₂CO/MeOH). Mp 223-225°. [α]_D²⁵ -0.5 (c, 1.2 in MeOH).

4,6α,7-Trihydroxy: 4,6α-Dihydroxyserratidine

[485817-35-4]

C₁₆H₂₃NO₄ 293.362

Alkaloid from *Huperzia serrata*. Yellow amorph. solid. [α]_D²⁵ -0.2 (c, 0.17 in MeOH).

4-Epimer: Lycopodatine C

[872610-92-9]

C₁₆H₂₃NO 245.364


Alkaloid from *Lycopodium inundatum*. Solid. [α]_D³³ -118 (c, 1 in MeOH).

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1964, **42**, 2514-2522; 1968, **46**, 14-20 (isol, ir, pmr, ms, synth)Inubushi, Y. *et al.*, *Chem. Comm.*, 1968, 1138-1139 (Serratidine)Kim, S.-W. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 3150-3153 (synth, ir, pmr, ms)Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 434-439; *CA*, **97**, 107030q

(Lycoplegmine)

Tan, C.-H. *et al.*, *Nat. Prod. Lett.*, 2002, **16**, 149-153 (Serratidine, Hydroxyserratidines)Morita, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1809-1812 (Lycopodatines B,C)**Anhydrolycorinium** A-1030


[61221-42-9 (chloride)]

C₁₆H₁₂NO₂[⊕] 250.276

Quaternary alkaloid from the bulbs of *Amaryllis belladonna* (Amaryllidaceae). Cryst. + 1.33 H₂O (EtOH)(as chloride). Sol. MeOH, CHCl₃, H₂O, poorly sol. hexane. Mp 280-285° (264-270° dec.) (chloride). λ_{max} 258 (ε 32300); 267 (ε 29500); 279 (ε 28800); 341 (ε 12100) (MeOH) (chloride).

Humber, L.G. *et al.*, *J.C.S.*, 1954, 4622 (synth, uv)Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 796 (isol, uv, ir, pmr, cmr)Boger, D.L. *et al.*, *J.O.C.*, 2000, **65**, 9120-9124 (synth)**4,6'-Anhydrooxysporidinone** A-1031

[933044-87-2]



C₂₈H₄₁NO₅ 471.636


Anhydro deriv. of Oxysporidinone, O-226. Prod. by *Fusarium oxysporum* EPH2R_{AA}. Cryst. (MeOH). Mp 160-161°. [α]_D²⁵ -78.6 (c, 0.3 in MeOH). λ _{max} 290 (log ϵ 3.7) (MeOH).

Zhan, J. et al., *J. Nat. Prod.*, 2007, **70**, 227-232 (isol, pmr, cmr, ms)

Anhydroperforine

A-1032

[22952-50-7]



Relative configuration

C₁₈H₂₃NO₄ 317.384

Alkaloid from the seeds of *Haplophyllum perforatum* (Rutaceae). Mp 143-144°. [α]_D³⁰ -35.1 (c, 2.19 in MeOH).


Bessonova, I.A. et al., *Khim. Prir. Soedin.*, 1974, **10**, 682; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 710 (pmr, struct)

Karimov, Z. et al., *Khim. Prir. Soedin.*, 1979, **15**, 805; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 712 (cryst struct)

Anhydroperipentamine

A-1033

N-[3-[[3-(3,4-Dihydro-5-methyl-4-oxo-2H-1-benzopyran-2-yl)propyl]amino]propyl]hexanamide, 9CI [100595-82-2]

C₂₂H₃₄N₂O₃ 374.522


Minor alkaloid from the bark of *Peripentadenia mearsii* (Elaeocarpaceae). Also obt. by dehydration of Peripentamine under mild acid conditions. Needles + 2½ H₂O (Me₂CO aq.). Mp 96°.

Bick, I.R.C. et al., *Tetrahedron*, 1985, **41**, 5627 (isol, uv, ir, pmr, cmr, ms, struct, synth)

Anhydrotetracycline

A-1034

4-(Dimethylamino)-1,4,4a,5,12,12a-hexahydro-3,10,11,12a-tetrahydroxy-6-methyl-1,12-dioxo-2-naphthacene-carboxamide, 9CI. Deschloroanhydroaureomycin [1665-56-1]

C₂₂H₂₂N₂O₇ 426.425

Toxic degradn. and photolysis prod. derived from Tetracycline. Intermed. in tetracycline biosynth. Cryst. (C₆H₆/hexane). Mp 225-226° (215-220° dec.). [α]_D²⁵ +25 (c, 1 in 2-ethoxyethanol).

[7518-17-4]

Waller, C.W. et al., *J.A.C.S.*, 1952, **74**, 4981-4982 (synth)

Stephens, C.R. et al., *J.A.C.S.*, 1954, **76**, 3568-3575 (synth)

Simmons, D.L. et al., *J. Pharm. Sci.*, 1966, **55**, 219-220 (synth)

Hasan, T. et al., *J.O.C.*, 1985, **50**, 1755-1757


Dos Santos, H.F. et al., *J. Pharm. Sci.*, 1998, **87**, 190-195 (conformm)

Zhang, W. et al., *J.A.C.S.*, 2008, **130**, 6068-6069 (biosynth)

4,9-Anhydrotetrodotoxin

A-1035

[13072-89-4]

C₁₁H₁₅N₃O₇ 301.255

Isol. from newts, other marine animals incl. *Cynops ensicauda*, *Fugu* spp. and *Octopus maculosus*, prob. as endobacterial prod. Toxic sodium channel blocker; phycotoxin. Amorph.

▶ Paralytic poison.

6-Epimer: 4,9-Anhydro-6-epitetrodotoxin [113565-40-5]

C₁₁H₁₅N₃O₇ 301.255

Isol. from the newt *Cynops ensicauda*. Amorph. [α]_D²⁵ +10 (c, 0.08 in AcOH aq.).

11-Deoxy: 4,9-Anhydro-11-deoxytetrodotoxin [113507-22-5]

C₁₁H₁₅N₃O₆ 285.256

Isol. from the newt *Cynops ensicauda*. Amorph.

Tsuda, K. et al., *Chem. Pharm. Bull.*, 1964, **12**, 634-642; 642-645; 1357-1374

Tamura, C. et al., *Acta Cryst. C*, 1966, **21**, 226-236 (cryst struct)

Nakamura, M. et al., *Toxicol.*, 1985, **23**, 271-276 (isol)

Yasumoto, T. et al., *Agric. Biol. Chem.*, 1986, **50**, 793-795 (isol)


Yasumoto, T. et al., *J.A.C.S.*, 1988, **110**, 2344-2345 (isol, pmr, cmr)

Dyankonov, A.L. et al., *Khim. Prir. Soedin.*, 1997, **33**, 297-351; *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 221-267 (rev)

Anibamine

A-1036

[725265-00-9]

C₃₀H₅₀N⁺ 424.732

Quaternary alkaloid from the stems of an *Aniba* sp. Amorph. solid (as TFA salt). λ _{max} 222 (ϵ 36270); 284 (ϵ 17140)


(MeOH) (TFA salt).

Jayasuriya, H. et al., *J. Nat. Prod.*, 2004, **67**, 1036-1038 (isol, pmr, cmr)

Anibine

A-1037

4-Methoxy-6-(3-pyridinyl)-2H-pyran-2-one, 9CI [643-91-4]

C₁₁H₉NO₃ 203.197

Alkaloid from *Aniba rosaeodora* and other *Aniba* spp. (Lauraceae). Mp 179-180°.

Picrate: Mp 199-201°.

Methiodide: Mp 233-236° dec.


Mors, W.B. et al., *J.A.C.S.*, 1957, **79**, 4507 (ir, uv, isol, struct)

Ziegler, E. et al., *Monatsh. Chem.*, 1958, **89**, 391; 716 (synth)

Anicanine

A-1038

1,2,3,4-Tetrahydro-1-(3-hydroxybenzyl)-6,7-dimethoxy-2-methylisoquinoline



(R)-form

C₁₉H₂₃NO₃ 313.396

(R)-form [151841-47-3]

Trace alkaloid from the stem bark of *Aniba canelilla* (Lauraceae). Isol. only as a 1:2 mixt. with Armepevine, A-1444.

N-De-Me: Noranicanine

[38974-83-3]

C₁₈H₂₁NO₃ 299.369

Alkaloid from the stem bark of *Aniba canelilla* (Lauraceae). Amorph. solid or needles (synthetic). Mp 182° synthetic. [α]_D +36 (c, 1 in CHCl₃).

(S)-form

N-De-Me: Synthetic. Flakes (EtOAc). Mp 188-189.5°. [α]_D²⁵ -25.6 (c, 0.5 in CHCl₃).

N-De-Me, hydrochloride:

Off-white solid (Me₂CO/Et₂O). Mp 140° dec.

O⁶-De-Me: Canelilline

[151757-04-9]

C₁₈H₂₁NO₃ 299.369

Alkaloid from the stem bark of *Aniba canelilla* (Lauraceae). [α]_D +43 (c, 0.5 in MeOH).

O⁶,N-Di-de-Me: Norcanelilline

[151757-03-8]


C₁₇H₁₉NO₃ 285.342

Alkaloid from the stem bark of *Aniba canelilla* (Lauraceae). [α]_D -18 (c, 1 in

CHCl₃).Oger, J.M. *et al.*, *Heterocycles*, 1992, **34**, 17 (Noranicanine)Oger, J.M. *et al.*, *Can. J. Chem.*, 1993, **71**, 1128 (Anicanine, Canelilline, Norcanelilline)Meyers, A.I. *et al.*, *Heterocycles*, 1994, **39**, 513 (synth, Noranicanine)Komori, K. *et al.*, *Heterocycles*, 1996, **43**, 1681 (synth, Noranicanine)**Aniflorine**

A-1039

3-[2-(Dimethylamino)phenyl]-2,3-dihydro-3-hydroxy-5-methoxypyrrrolo[2,1-b]quinazolin-9(1H)-one, 9CI [29048-81-5]

C₂₀H₂₁N₃O₃ 351.404Minor alkaloid from the leaves and branches of *Anisotes sessiliflorus* (Acanthaceae). Cryst. (MeOH). Mp 195-197°. Opt. inactive.**Deoxy: Deoxyaniflorine**


[16688-22-5]

C₂₀H₂₁N₃O₂ 335.405Minor alkaloid from leaves and branches of *Anisotes sessiliflorus* (Acanthaceae). Cryst. (MeOH). Mp 168-172°. Opt. inactive.**Demethoxy: Demethoxyaniflorine**

[178631-30-6]

C₁₉H₁₉N₃O₂ 321.378Alkaloid from leaves of *Adhatoda vasica*. Mp 128-129°.Arndt, R.R. *et al.*, *Tetrahedron*, 1967, **23**, 3521 (isol, uv, ir, pmr, ms, struct, synth)Thappa, R.K. *et al.*, *Phytochemistry*, 1996, **42**, 1485 (Demethoxyaniflorine)**2-Anilino-1,4-naphthoquinone, 8CI**

A-1040


2-(Phenylamino)-1,4-naphthalenedione, 9CI. 4-Hydroxy-2-(phenylimino)-1(2H)-naphthalenone, 9CI. **Lutine** [66855-47-8]C₁₆H₁₁NO₂ 249.268Alkaloid from *Reseda luteola* (Resedaceae). Dye for wool and silk. Cherry-red cryst. (CHCl₃ or EtOH aq.). Mp 190-191°. λ_{max} 220 (log ε 4.2); 271 (log ε 4.53); 325 (log ε 3.68); 465 (log ε 3.77) (EtOH).

[6628-97-3]

Blackhall, A. *et al.*, *J.C.S.*, 1953, 1138-1143 (synth)Inoue, A. *et al.*, *CA*, 1961, **55**, 13853i (synth)Jerzy, M. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1974, **48**, 147-153 (uv)Lutfullin, K.L. *et al.*, *Khim. Prir. Soedin.*, 1977, 826-830; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **13**, 696-699 (isol, ir, ms, uv)**Anisessine**

A-1041

Methyl 2-[(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-b]quinazolin-3-yl)amino]benzoate, 9CI. Methyl N-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-b]quinazolin-3-yl)anthranilate, 8CI [16688-20-3]


C₂₀H₁₉N₃O₃ 349.388Minor alkaloid from the leaves and branches of *Anisotes sessiliflorus* (Acanthaceae). Cryst. (MeOH). Mp 170-171°. Opt. inactive.**Me ester analogue: Vasnetine**

[158734-26-0]

C₁₉H₁₇N₃O₃ 335.362Alkaloid from leaves of *Adhatoda vasica* (Acanthaceae). Mp 185-187°.Arndt, R.R. *et al.*, *Tetrahedron*, 1967, **23**, 3521 (isol, uv, ir, pmr, ms, struct)Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4387 (synth) Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 953 (Vasnetine)**Anisocycline**

A-1042

5,6-Dihydro-2,3,4,9,10-pentamethoxydibenzo[a,g]quinolininium(1+), 9CI [125085-14-5]

C₂₂H₂₄NO₅[⊕] 382.435Quaternary alkaloid from the roots of *Anisocycla cymosa* (Menispermaceae). Yellow needles (MeOH). Mp 195°. Counterion not specified.**O⁴-De-Me: Lincangine**


[142741-22-8]

C₂₁H₂₂NO₅[⊕] 368.409Alkaloid from the roots of *Stephania lincangensis* (Menispermaceae).**O⁴,O¹⁰-Di-de-Me: Fissisaine**

[211235-37-9]

C₂₀H₂₀NO₅[⊕] 354.382Alkaloid from *Fissistigma balansae*. Yellow needles (as chloride). Mp >300° (chloride). λ_{max} 206; 230; 282; 337 (EtOH).Kanyinda, B. *et al.*, *Planta Med.*, 1989, **55**, 394 (isol, uv, ir, pmr, ms, struct)Chen, Y. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1991, **33**, 552; *CA*, **117**, 86637g (Lincangine)Chia, Y.C. *et al.*, *Phytochemistry*, 1998, **48**, 367-369 (Fissisaine)**Anisodine**


A-1043

9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl α-hydroxy-α-(hydroxymethyl)-benzeneacetate, 9CI. 3α-(2,3-Dihydroxy-2-phenylpropionyl)-6β,7β-epoxytropolane. 6,7-Epoxy-3-tropanyl 2,3-dihydroxy-2-phenylpropionate. α-Hydroxyscopolamine. Daturamine [52646-92-1]C₁₇H₂₁NO₅ 319.357Ester of Scopine, S-158 with 2,3-Dihydroxy-2-phenylpropanoic acid. Antispasmodic, anticholinergic agent. Mp 182-183° (as hydrochloride). [α]_D -18.5 (c,3 in H₂O). Log P -0.51 (calc).**(S)-form**Minor alkaloid from *Datura sanguinea* (Solanaceae), impurity in coml. Scopolamine. Present in many Chinese plants. Shows strong depressant effect in rabbits. Some claimed syntheses are dubious.**(±)-form**

Mp 217-219° (as hydrochloride).

Moorhoff, C.F. *et al.*, *Planta Med.*, 1975, **28**, 106-108 (isol)Minina, S.A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 712-713; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 598 (isol)Bode, J. *et al.*, *Acta Cryst. B*, 1982, **38**, 333-335 (cryst struct, abs config)Bian, C. *et al.*, *CA*, 1982, **96**, 46211e (pharmacol)Xie, J. *et al.*, *CA*, 1982, **97**, 216529c; **98**, 215843m; 1984, **100**, 139413 (S-form, synth, abs config)Fodor, G. *et al.*, *Nat. Prod. Rep.*, 1984, **1**, 232-234 (rev)He, L. *et al.*, *CA*, 1991, **114**, 129215 (occur)Wu, Y.F. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 1287-1289 (synth)**Anisomycin**

A-1044

2-(4-Methoxyphenylmethyl)-3,4-pyrrolidinediol 3-acetate, 9CI. Flagecidin. PA 106. SA 3097C₁. Antibiotic SA 3097C₁. Antibiotic PA 106. Anhydroscopecin A. NSC 76712

(-)-form

C₁₄H₁₉NO₄ 265.308Log P 1.13 (calc). λ_{max} 223 (ε 10900); 276 (ε 1790); 283 (ε 1550) (MeOH) (Derep).**►BZ9800000****(-)-form [22862-76-6]**Isol. from *Streptomyces griseolus*, *Streptomyces anhydroscopecinus* and *Streptomyces roseochromogenes*. Antibiotic and

antifungal agent. Protein synthesis inhibitor. Has been used to induce amnesia in animal models. Long needles (EtOAc, H₂O or toluene). Mp 141.6-142.2°. [α]_D²³ -30 (c. 1 in MeOH). [α]_D -44.6 (CHCl₃). pK_a 7.9. Pharmacol. active isomer. ▶ LD₅₀ (rat, orl) 72 mg/kg. LD₅₀ (rat, ipr) 345 mg/kg.

Hydrochloride:

Cryst. (EtOAc/EtOH). Mp 187-188°.

O-De-Ac: Deacetylanisomycin. Antibiotic SA 3097D₁. SA 3097D₁

[27958-06-1]

C₁₂H₁₇NO₃ 223.271

Prod. by *Streptomyces* sp. SA3097.

Solid. Sol. MeOH, H₂O. [α]_D²⁵ -22.5 (c. 1 in MeOH). λ_{\max} 223 (ε 11000); 276 (ε 1100); 282 (sh) (ε 1000) (MeOH) (Derrep). λ_{\max} 225; 277 (MeOH) (Berdy).

O-De-Ac, 3-propanoyl: Antibiotic SA 3097B₁. SA 3097B₁

[139755-62-7]

C₁₅H₂₁NO₄ 279.335

Prod. by *Streptomyces* sp. SA3097.

Powder. Sol. MeOH, H₂O. λ_{\max} 223 (ε 10900); 276 (ε 1790); 283 (ε 1550) (MeOH) (Derrep). λ_{\max} 224; 275; 282 (MeOH) (Berdy). λ_{\max} 223; 276; 282 (EtOH) (Berdy).

O-De-Ac, 3-butanoyl: Anisomycin C [96740-19-1]

C₁₆H₂₃NO₄ 293.362

Prod. by *Streptomyces* sp. C4826.

Cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 225 (ε 12300); 277 (ε 1380); 283 (ε 1090) (EtOH).

O-De-Ac, 3-(3-methylbutanoyl): Anisomycin D

[96751-33-6]

C₁₇H₂₅NO₄ 307.389

Prod. by *Streptomyces* sp. C4826.

Cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D -43 (MeOH). λ_{\max} 225 (ε 14500); 276 (ε 2190); 283 (ε 1820) (EtOH).

O-De-Me: De-O-methylanisomycin. Antibiotic SA 3097C₂. SA 3097C₂

[139755-64-9]

C₁₃H₁₇NO₄ 251.282

Prod. by *Streptomyces* sp. SA3097. Oil.

Sol. MeOH, H₂O. λ_{\max} 223 (ε 10900); 276 (ε 1790); 283 (ε 1550) (MeOH) (Derrep). λ_{\max} 225; 275; 280 (MeOH) (Berdy). λ_{\max} 225; 277 (EtOH) (Berdy).

O-De-Me, O-de-Ac: Deacetylde-O-methylanisomycin. Antibiotic SA 3097D₂. SA 3097D₂

[152338-66-4]

C₁₁H₁₅NO₃ 209.244

Prod. by *Streptomyces* sp. SA3097. Sol.

MeOH, H₂O. λ_{\max} 223 (ε 11000); 276 (ε 1100); 282 (sh) (ε 1000) (MeOH) (Derrep). λ_{\max} 225; 277 (MeOH) (Berdy).

O-De-Me, 3-propanoyl, O-de-Ac: Antibiotic SA 3097B₂. SA 3097B₂

[139755-63-8]

C₁₄H₁₉NO₄ 265.308

Prod. by *Streptomyces* sp. SA3097. Oil.

Sol. MeOH, H₂O. λ_{\max} 223 (ε 10900); 276 (ε 1790); 283 (ε 1550) (MeOH) (Derrep). λ_{\max} 225; 275; 280 (MeOH)

(Berdy). λ_{\max} 225; 277 (EtOH) (Berdy).

(±)-form

Cryst. (EtOAc). Mp 126-127°.

Sobin, B.A. *et al.*, *J.A.C.S.*, 1954, **76**, 4053 (*isol*)
Schaefer, J.P. *et al.*, *J.O.C.*, 1968, **33**, 166 (*cryst struct*)

Butler, K. *et al.*, *J.O.C.*, 1968, **33**, 2136 (*struct, pmr, conformn*)

Wong, C.M. *et al.*, *Can. J. Chem.*, 1969, **47**, 2421 (*synth, ir*)

Oida, S. *et al.*, *Chem. Pharm. Bull.*, 1969, **17**, 1405

Felner, I. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 754 (*synth, ir, ms*)

Verheyden, J.P. *et al.*, *Pure Appl. Chem.*, 1978, **50**, 1363 (*rev*)

Jimenez, A. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 1 (*rev*)

Martinez, J.L. *et al.*, *Prog. Neurobiology*, 1981, **16**, 155 (*pharmacol, rev*)

Schumacher, D.P. *et al.*, *J.A.C.S.*, 1982, **104**, 6076 (*synth*)

Buchanan, J.G. *et al.*, *Chem. Comm.*, 1983, 486 (*synth*)

Shono, T. *et al.*, *Chem. Lett.*, 1987, 697 (*synth, bibl*)

Baer, H.H. *et al.*, *J.O.C.*, 1988, **53**, 4786 (*synth, bibl*)

Takano, S. *et al.*, *Heterocycles*, 1989, **29**, 1861 (*synth*)

Ballini, R. *et al.*, *J.O.C.*, 1992, **57**, 1316 (*synth*)

Hosoya, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 1300 (*sar*)

LeFevre, J.W. *et al.*, *Magn. Reson. Chem.*, 1993, **31**, 318 (*cmr*)

Cano, E. *et al.*, *Mol. Cell. Biochem.*, 1994, **14**, 7352 (*pharmacol*)

Yoda, H. *et al.*, *Heterocycles*, 1995, **41**, 2423 (*synth*)

Huang, P.Q. *et al.*, *Nat. Prod. Lett.*, 1998, **11**, 101-106 (*synth*)

Reddy, G.V. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 133-136 (*synth, de-Ac*)

Delair, P. *et al.*, *J.O.C.*, 1999, **64**, 1383-1386 (*synth*)

Hulme, A.N. *et al.*, *Org. Lett.*, 2002, **4**, 265-267 (*synth*)

Chandrasekhar, S. *et al.*, *Synthesis*, 2002, 1867-1870 (*de-Ac*)

Merino, P. *et al.*, *Eur. J. Org. Chem.*, 2003, 2877-2881 (*de-Ac, synth*)

Chang, M.-Y. *et al.*, *Heterocycles*, 2003, **60**, 1203-1209 (*synth*)


Ono, M. *et al.*, *Tetrahedron*, 2004, **60**, 10187-10195 (*synth*)

Joo, J.-E. *et al.*, *Org. Lett.*, 2007, **9**, 3627-3630 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AOY000

Anisotine**A-1045**

Methyl 2-(methylamino)-5-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-b]quinazolin-3-yl)benzoate, 9CI. Methyl N-methyl-5-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-b]quinazolin-3-yl)anthranilate, 8CI [16688-19-0]



C₂₀H₁₉N₃O₃ 349.388

Minor alkaloid from the leaves and branches of *Anisotes sessiliflorus*. Also isol. from *Adhatoda vasica* (Acanthaceae). Cryst. (Me₂CO/hexane). Mp 189-190°. Opt. inactive.

Deoxo: Adhatodine

[33903-14-9]

C₂₀H₂₁N₃O₂ 335.405

Alkaloid from *Adhatoda vasica* (Acanthaceae). Cryst. (Me₂CO/hexane or CHCl₃/heptane). Mp 183°.

3-Hydroxy: 3-Hydroxyanisotine

[16688-24-7]

C₂₀H₁₉N₃O₄ 365.388

Alkaloid from *Adhatoda vasica*. Mp 185-187° (182-184°). λ_{\max} 226 (log ε 4.74); 263 (log ε 4.46); 302 (log ε 3.98); 314 (log ε 3.87); 353 (log ε 3.69) (EtOH).

Arndt, R.R. *et al.*, *Tetrahedron*, 1967, **23**, 3521-3532 (*isol, uv, ir, pmr, ms, struct*)

Johne, S. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 826 (*isol, uv, ir, pmr, ms, struct, Anisotine, Adhatodine*)


Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 953-962 (*isol, pmr, cmr*)

Ankoline**A-1046**

C₁₇H₃₆N₂O₄ 332.482

Struct. unknown. Alkaloid from *Alangium lamarckii*. Brownish-yellow cryst. Mp 110° Mp 153° (as picrate).

Basu, N.K. *et al.*, *Indian J. Pharm.*, 1950, **12**, 98-99 (*isol*)

Ankorine**A-1047**

(-)-form

C₁₉H₂₉NO₄ 335.442

The 8-hydroxy congener of Protoemetinol in P-664.

(-)-form [13849-54-2]

Alkaloid from the leaves of *Alangium lamarckii* and the branches of *Alangium salvifolium* (Alangiaceae). Prisms (Me₂CO). Mp 176-177° (174-176°). [α]_D¹⁶ -58 (c. 0.23 in CHCl₃). λ_{\max} 272 (log ε 2.96) (EtOH). λ_{\max} 287 (log ε 3.43) (0.1M NaOH). λ_{\max} 230 (sh) (log ε 3.97); 273 (log ε 3.01); 280 (sh) (log ε 2.94) (MeOH). λ_{\max} 273 (log ε 3.02); 279 (sh) (log ε 3.01) (0.1M NaCl).

Hydrochloride:

Cryst. + 1H₂O (EtOH/Et₂O). Mp 233-234°.

Hydrobromide: Mp 220-222° dec.

Carboxylic acid: Alancine

[92631-63-5]

C₁₉H₂₇NO₅ 349.426

Alkaloid from the stem bark of *Alangium lamarckii* (Alangiaceae). Mp 229-

230°. $[\alpha]_D^{25}$ -40 (c, 0.1 in MeOH). Forms hemihydrate, Mp 216-220.5° dec.

Carboxylic acid, hydrochloride:

Minute prisms (EtOH). Mp 247.5-248.5°. $[\alpha]_D^{22}$ -28 (c, 0.101 in MeOH).

(±)-form [56816-22-9]

Synthetic. Mp 200-202° (192.5°).

Hydrochloride: Mp 132-133°.

Carboxylic acid: Synthetic. Prisms + 1/2 H₂O (EtOH aq.). Mp 217-218° dec.

Carboxylic acid, hydrochloride:

Minute needles (EtOH aq.). Mp 241.5-245° dec.

Dasgupta, B. *et al.*, *J. Pharm. Sci.*, 1965, **54**, 481-483 (*isol, uv, ir, pmr*)

Battersby, A.R. *et al.*, *Tet. Lett.*, 1966, 4965-4971 (*uv, ir, pmr, ms*)

Yoshifujii, S. *et al.*, *Tet. Lett.*, 1975, 1965-1968 (*abs config*)

Szántay, C. *et al.*, *Chem. Ber.*, 1976, **109**, 2420-2429 (*struct, synth, ir, pmr, ms*)

Fujii, T. *et al.*, *J.O.C.*, 1980, **45**, 1889-1893 (*uv, ir, pmr, ms, abs config*)

Fujii, T. *et al.*, *Tetrahedron*, 1980, **36**, 965-968 (*synth*)

Chen, M.-J. *et al.*, *Zhivuo Xuebao (Acta Bot. Sin.)*, 1980, **22**, 257; *CA*, **94**, 12818b (*isol*)

Chattopadhyay, S.K. *et al.*, *Heterocycles*, 1984, **22**, 1965-1968 (*Alcane, isol, uv, ir, pmr, cmr, ms, struct, synth*)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 669-676; 1987, **35**, 3470-3474; 1988, **36**, 2665-2668 (*Alcane, Ankorine, synth, uv, ir, pmr, cd*)

Annapawine**A-1048**

C₁₆H₁₇NO₃ 271.315

Amaryllidaceae alkaloid. Struct unknown. *Isol.* from the *Hippeastrum* hybrid "Anna Pavlovna" (Amaryllidaceae). Cryst. (Me₂CO). Mp 219°. $[\alpha]_D^{25}$ +74 (c, 0.2 in CHCl₃).


Hydroiodide: Mp 221° dec.

Picrate: Mp 249° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 470-471 (*isol*)

Annocherine A**A-1049**

Annocherine C



C₁₇H₁₅NO₄ 297.31

Annocherine A was the *S*-enantiomer and Annocherine C the *R*-form.

(R)-form [431982-10-4]

Alkaloid from the leaves of *Annona cherimola* (cherimoya). Yellow amorph. powder. Mp 123-125°. $[\alpha]_D^{25}$ -125 (c, 0.3 in CHCl₃). λ_{max} 260 ; 300 ; 331 (MeOH).

(S)-form

Alkaloid from *Annona cherimola* (cherimoya). Yellow needles. Mp 156-158°. $[\alpha]_D^{24}$ +135 (c, 0.1 in CHCl₃). λ_{max} 260

(log ϵ 3.41); 300 (log ϵ 3.1); 330 (log ϵ 3.52) (MeOH).

 α -Me ether: Annocherine B

C₁₈H₁₇NO₄ 311.337


Alkaloid from *Annona cherimola* (cherimoya). Amorph. yellow powder. Mp 196-198°. $[\alpha]_D^{24}$ +115 (c, 0.1 in CHCl₃). λ_{max} 260 (log ϵ 3.32); 300 (log ϵ 3.11); 331 (log ϵ 3.6) (MeOH).

Chen, C.-Y. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 1203-1206 (*Annocherine C*)

Chen, C.-Y. *et al.*, *Phytochemistry*, 2001, **56**, 753-757 (*Annocherine A,B*)

Annotatine**A-1050**

[149998-52-7]




C₁₉H₁₅NO₅ 337.331

Alkaloid from leaves of *Annona montana* (Annonaceae). Green amorph. powder.

Wu, Y.-C. *et al.*, *Phytochemistry*, 1993, **33**, 497 (*isol, uv, ir, pmr, ms, struct*)

Annomontine**A-1051**

4-(9H-Pyrido[3,4-b]indol-1-yl)-2-pyridinamine, 9CI. 1-(2-Amino-4-pyrimidinyl)- β -carboline. Annomurine [82504-00-5]



C₁₅H₁₁N₅ 261.285

Alkaloid from the bark of *Annona foetida* and *Annona montana*. Yellow cryst. (CH₂Cl₂). Mp 249-250°. λ_{max} 246 ; 312 ; 398 ; 428 (CHCl₃).

N⁹-Hydroxy: N⁹-Hydroxyannonomontine

C₁₅H₁₁N₅O 277.285

Alkaloid from the bark of *Annona foetida*. Red needles (CHCl₃). Mp 249-250°. Hydroxylated at the indolic N. λ_{max} 244 (log ϵ 4.3); 314 (log ϵ 3.97); 434 (log ϵ 3.63) (CHCl₃).

6-Methoxy: 4-(6-Methoxy-9H-pyrido[3,4-b]indol-1-yl)-2-pyrimidinamine, 9CI. 1-(2-Amino-4-pyrimidinyl)-6-methoxy- β -carboline. **Methoxyannonomontine** [82504-01-6]

C₁₆H₁₃N₅O 291.312


Alkaloid from the root bark of *Annona montana* (Annonaceae). Orange cryst. (EtOH). Mp 229-230° (223°).

Yokomori, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1982, **55**, 2236 (*isol, cryst struct*)

Leboeuf, M. *et al.*, *J.C.S. Perkin 1*, 1982, 1205 (*uv, ir, pmr, cmr, ms, struct, deriv*)

Bracher, F. *et al.*, *Annalen*, 1993, 837 (*synth*)

Costa, E.V. *et al.*, *J. Nat. Prod.*, 2005, **69**, 292-294 (*N-Hydroxyannonomontine*)

Annonelliptine**A-1052**

C₁₉H₂₃NO₄ 329.395

(R)-form [96400-46-3]

Alkaloid from the leaves and stems of *Annona elliptica* (Annonaceae). Needles (MeOH). Mp 198-200°. $[\alpha]_D^{22}$ -101 (c, 0.38 in CHCl₃/MeOH 1:1). λ_{max} 225 (sh) (log ϵ 4.22); 282 (log ϵ 3.51) (MeOH).

N-De-Me: Anomoline

[142287-93-2]

C₁₈H₂₁NO₄ 315.368

Alkaloid from stems of *Annona cherimola* (Annonaceae). Mp 193-194.5°. $[\alpha]_D^{25}$ +27.82 (MeOH).

(±)-form

Synthetic. Mp 182.5-184°. CAS No. not found 10-14CI.

Hydrochloride: [173789-57-6]

Cryst. (EtOH). Mp 230-231°.

N-De-Me, hydrochloride: [173789-58-7]

Cryst. (EtOH). Mp 238.5-240°.


Sandoval, D. *et al.*, *Phytochemistry*, 1985, **24**, 375-376 (*isol, uv, ir, pmr, cmr, ms, ord, struct*)

Yang, T.H. *et al.*, *Chin. Pharm. J. (Taipei)*, 1991, **43**, 457-463; *CA*, **117**, 44523m (*Anomoline*)

Chen, C.-M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1767-1771 (*synth, Annonelliptine, Anomoline*)

Annonidine A**A-1053**

[99102-30-4]



C₂₆H₂₈N₂ 368.521

Alkaloid from the stem bark of *Anonidium mannii* (Annonaceae) and roots of *Esenbeckia leiocarpa* (Rutaceae). Needles (petrol/EtOAc). Mp 106-108°. $[\alpha]_D^{20}$ 0 (c, 1.0 in CHCl₃).

Achenbach, H. *et al.*, *Heterocycles*, 1985, **23**, 2075 (*isol, uv, pmr, cmr, ms, struct*)

Somei, M. *et al.*, *Heterocycles*, 1987, **26**, 1783 (synth)
 Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (isol)

Annonidine B A-1054


[99102-31-5]
 As Annonidine A, A-1053 with
 R = 3-indolyl

C₂₆H₂₈N₂ 368.521
 Alkaloid from the stem bark of *Anonidium mannii* (Annonaceae).

Achenbach, H. *et al.*, *Heterocycles*, 1985, **23**, 2075 (isol, uv, pmr, cmr, ms, struct)

Annonidine C A-1055

[99102-32-6]



C₂₆H₂₈N₂ 368.521
 Alkaloid from the stem bark of *Anonidium mannii* (Annonaceae) and roots of *Esenbeckia leiocarpa* (Rutaceae). Oil.

2',3'-Epoxide: Annonidine E

[99102-36-0]
 C₂₆H₂₈N₂O 384.52

Alkaloid from the stem bark of *Anonidium mannii*.

Achenbach, H. *et al.*, *Heterocycles*, 1985, **23**, 2075 (isol, uv, pmr, cmr, ms, struct)
 Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (isol)

Annonidine D A-1056


[99102-34-8]
 As Annonidine C, A-1055 with
 R = 6-indolyl

C₂₆H₂₈N₂ 368.521
 Alkaloid from the stem bark of *Anonidium mannii* (Annonaceae). Oil.

Achenbach, H. *et al.*, *Heterocycles*, 1985, **23**, 2075 (isol, uv, pmr, cmr, ms, struct)

Annonidine F A-1057

[820965-40-0]




C₂₆H₂₈N₂ 368.521
 Alkaloid from *Monodora angolensis*. Yellow oil.

Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (isol, pmr, cmr)

Annopholine A-1058

5,10-Dimethoxy-4-methylbenzo[g]quinoline, 9CI
 [112494-58-3]




C₁₆H₁₅NO₂ 253.3
 Alkaloid from the stem bark of *Annona haysii* (Annonaceae). Amorph.

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1987, **50**, 759 (isol, uv, pmr, ms, struct)
 Vallejos, G. *et al.*, *Synth. Commun.*, 1999, **29**, 809-814 (synth, ir, pmr, cmr)

Annopodine A-1059

[20390-90-3]



C₁₇H₂₅NO₃ 291.389
 Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Mp 211-212°.


Perchlorate: Mp 210-212°.

Methiodide: Mp 240-242°.

Ayer, W.A. *et al.*, *Tet. Lett.*, 1968, 4597 (ir, uv, pmr, cryst struct)

Annorettine A-1060

[149998-37-8]




C₁₉H₁₉NO₂ 293.365
 Alkaloid from leaves of *Annona montana* (Annonaceae). Exhibits significant cytotoxicity. Purple cryst. Mp 160-163°. λ_{max} 205 (ε 12020); 231 (ε 9770); 255 (ε 22387); 302 (ε 12020); 343 (ε 3300) (MeOH) (Berdy). λ_{max} 205 (ε 15488); 258 (ε 12882); 312 (ε 5128); 345 (ε 4900) (MeOH-NaOH) (Berdy).

Wu, Y.-C. *et al.*, *Phytochemistry*, 1993, **33**, 497 (isol, uv, ir, pmr, cmr, ms, struct)

Pampin, M.C. *et al.*, *Tet. Lett.*, 2001, **42**, 2307-2308 (synth, cryst struct)

Annosqualine A-1061

[770737-96-7]



C₁₉H₁₉NO₅ 341.363
 Alkaloid from the stems of *Annona squamosa* (sugar apple). Syrup. [α]_D²⁷ +56 (c 0.19 in MeOH).


Yang, Y.-L. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1392-1399 (isol, pmr, cmr, ms)

Shigehisa, H. *et al.*, *Tet. Lett.*, 2006, **47**, 7301-7306 (synth)

Annotine A-1062

Alkaloid L11

[5096-59-3]



C₁₆H₂₁NO₃ 275.347
 Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Prisms (EtOH). Mp 174.5-176°. [α]_D²⁰ -114 (c 1 in CHCl₃).

Methiodide: Mp 236-237° dec.

Dihydro: Mp 203-204°.

1:1 Complex with Acrifoline: Annotoxine

C₃₂H₄₄N₂O₅ 536.71

Isol. from *Lycopodium annotinum* (Lycopodiaceae). Cryst. (Me₂CO or MeOH). Mp 196-197°. [α]_D²⁰ -187.1 (c 1 in CHCl₃).

Bertho, A. *et al.*, *Chem. Ber.*, 1952, **85**, 663 (isol, uv)

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1955, **29**, 509 (Annotoxine)

Perry, G.S. *et al.*, *Can. J. Chem.*, 1956, **34**, 1189 (Annotoxine)

Szarek, W.A. *et al.*, *Can. J. Chem.*, 1964, **42**, 2584 (ir, pmr, struct)


MacLean, D.B. *et al.*, *Can. J. Chem.*, 1966, **44**, 611 (ms)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1989, **67**, 1765 (pmr, cmr, struct)

Annotinine A-1063

Alkaloid L7

[559-49-9]



C₁₆H₂₁NO₃ 275.347
 Alkaloid from *Lycopodium annotinum*, *Lycopodium annotinum* var. *acrifolium*, *Lycopodium flabelliforme* and *Lycopodium obscurum* (Lycopodiaceae). Mp 232°.

▶ CA1190000

Hydrochloride:

Cryst. + ½ H₂O. Mp 210-211° dec.


Perchlorate: Mp 267°.

- Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 92 (*isol*)
 Przybylska, M. *et al.*, *Acta Cryst.*, 1958, **11**, 718 (*cryst struct*)
 Wiesner, K. *et al.*, *Tet. Lett.*, 1961, 187 (*abs config*)
 MacLean, D.B. *et al.*, *Can. J. Chem.*, 1963, **41**, 2654 (*ms*)
 Wiesner, K. *et al.*, *Can. J. Chem.*, 1969, **47**, 433 (*synth*)
 Ho, T.-L. *et al.*, *Tet. Lett.*, 1969, 1307 (*abs config*)
 Hughes, D.W. *et al.*, *Can. J. Chem.*, 1989, **67**, 1765 (*pmr, cmr*)
 Ayer, W.A. *et al.*, *Can. J. Chem.*, 1999, **77**, 1514-1520 (*pmr, conformn*)
 Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (*synth, rev*)

Annuloline

A-1064

[3988-51-0]




- $C_{20}H_{19}NO_4$ 337.374
 Alkaloid from *Lolium multiflorum* (annual rye grass) (Poaceae). Mp 114° (105-106°).
Hydrochloride: Mp 174-177°.
Picrate: Mp 216-218°.
 Axelrod, B. *et al.*, *J.O.C.*, 1958, **23**, 919 (*isol, uv*)
 Karimoto, R.S. *et al.*, *Tet. Lett.*, 1962, 83 (*pmr, synth*)
 O'Donovan, D.G. *et al.*, *J.C.S.(C)*, 1971, 331 (*biosynth*)
 Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*occur*)
 Molina, P. *et al.*, *Heterocycles*, 1993, **36**, 2255 (*synth*)

Anodendrine

A-1065

1-Carboxyhexahydro-4-(3-methyl-2-butenyl)-1H-pyrrolizinium, 9CI. N-Isopentenylburninic acid
 [28942-07-6]




- $C_{13}H_{21}NO_2$ 223.314
 Alkaloid from *Anodendron affine* (Apocynaceae). $[\alpha]_D^{25} +9.5$ (EtOH). pK_a 3.7 (MeOH).
Picrate:
 Yellow needles. Mp 123-124°.
 1-Epimer: *Alloanodendrine*
 [29391-56-8]
 $C_{13}H_{21}NO_2$ 223.314
 From *Anodendron affine* (Apocynaceae). Oil. $[\alpha]_D^{25} +18$ (EtOH). pK_a 3.2 (H₂O).
 Sasaki, K. *et al.*, *Tetrahedron*, 1970, **26**, 2119

Anolobine

A-1066

9-Hydroxy-1,2-methylenedioxyynoraporphine. *Anolobine*



(R)-form

 $C_{17}H_{15}NO_3$ 281.31

(R)-form [641-17-8]

Alkaloid from a variety of genera in the Annonaceae (*Annona*, *Asimina*, *Duguetia*, *Guatteria*, *Monodora*, *Polyalthia*, *Schefferomitra*) and Magnoliaceae (*Magnolia*). Shows antimicrobial activity. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 242-244° (>310°). $[\alpha]_D^{30} -19$ (c, 0.42 in CHCl₃/MeOH).

N-Formyl: N-Formylanolobine

$C_{18}H_{15}NO_4$ 309.321
 Alkaloid from *Sciadotenia toxifera*. Purple-grey needles (CH₂Cl₂). Mp 284° dec. λ_{max} 246 (log ϵ 2.8); 282 (log ϵ 2.8); 294 (log ϵ 2.81); 310 (log ϵ 2.79) (MeOH).

N-Ac: N-Acetylanolobine

[220464-95-9]
 $C_{19}H_{17}NO_4$ 323.348
 Alkaloid from *Magnolia coco*. Cryst. (MeOH). Mp 295-297°. λ_{max} 201 (log ϵ 3.56); 282 (log ϵ 3.04) (MeOH).

N-Me: 9-Hydroxy-1,2-methylenedioxyynoraporphine. **Roemeroline**. **Remeroline**. N-Methylanolobine
 [21059-86-9]
 $C_{18}H_{17}NO_3$ 295.337

Alkaloid from *Roemeria refracta*, *Stephania pierrei*, *Stephania sasakii* and *Duguetia obovata*. Shows antimalarial props. Needles + 1MeOH (MeOH). Mp 218-220° (sint. at 133-135°). $[\alpha]_D -32.1$ (c, 0.437 in CHCl₃). λ_{max} 281 (ϵ 8912) (MeOH) (Berdy).

N-Me; hydrochloride:

Cryst. (HCl aq.). Mp 220-225°.

Me ether: 9-Methoxy-1,2-methylenedioxyynoraporphine. **Xylopine**. O-Methylanolobine. **Artabotrinine**

[517-71-5]
 $C_{18}H_{17}NO_3$ 295.337
 Alkaloid from *Guatteria amplifolia*, *Xylophia discreta* and *Xylophia papuana*, poss. also from the bark of *Artabotrys suaveolens* (Annonaceae). CNS depressant, shows antimicrobial activity. Mp 78-102°. $[\alpha]_D^{22} -28.2$ (c, 0.001 in MeOH). Struct. of Artabotrinine not definitely establ., identity with Xylopine not certain.

Me ether, hydrochloride: Mp 250°.*Me ether, N-formyl*: N-Formylxylopine

[89368-31-0]
 $C_{19}H_{17}NO_4$ 323.348
 Trace alkaloid from the leaves of *Duguetia obovata* (Annonaceae).

Cryst. (MeOH). Mp 173°. $[\alpha]_D -248$ (c, 1.70 in CHCl₃).

Me ether, N-methoxycarbonyl: 9-Methoxy-N-methoxycarbonyl-1,2-methylenedioxyynoraporphine
 $C_{20}H_{19}NO_5$ 353.374

Alkaloid from the stems of *Fissistigma bracteolatum*. Powder. Mp 174-176°. $[\alpha]_D^{20} -281.7$ (c, 0.01 in CHCl₃). MF incorr. in ref. λ_{max} 215 (log ϵ 4.38); 239 (log ϵ 4.03); 282 (log ϵ 4.24); 325 (log ϵ 3.41) (no solvent reported).

Me ether, N-Me: 9-Methoxy-1,2-methylenedioxyynoraporphine. (-)-**Isolaureline**.

N-Methylxylopine

[475-84-3]

 $C_{19}H_{19}NO_3$ 309.364

Trace alkaloid from the leaves of *Duguetia obovata* and from *Stephania pierrei*. Antimalarial. Needles (Me₂CO/Et₂O). Mp 108-110°. $[\alpha]_D^{26} -36.7$ (EtOH).

Me ether, N-Me, hydrochloride:

Prisms (MeOH/2-propanol). Mp 243-246° dec.

Me ether, N-Me, N-oxide: **Isolaureline N-oxide**

[104385-28-6]

 $C_{19}H_{19}NO_4$ 325.363Alkaloid from *Magnolia obovata*.

Me ether, N,N-di-Me: N,N-Dimethylxylopine

 $C_{20}H_{22}NO_3^{\oplus}$ 324.399

Quaternary alkaloid from *Monodora grandidieri* and *Monodora junodii*.

Prisms (MeOH) (as iodide). Mp 229-232° dec. (iodide).

Me ether, N-nitroso: N-Nitrosoxylopine $C_{18}H_{16}N_2O_4$ 324.335

Alkaloid from *Duguetia furfuracea*. Amorph. solid (MeOH/CHCl₃). $[\alpha]_D^{25} -264$ (c, 0.5 in CHCl₃).

(S)-form

Me ether, N-Me: (+)-**Isolaureline** $C_{19}H_{19}NO_3$ 309.364

Alkaloid from *Liriodendron tulipifera* (Magnoliaceae). Needles (Et₂O). Mp 108-109°. $[\alpha]_D +40.3$ (EtOH).

(±)-form

Synthetic. Pale brown cryst. (CHCl₃/MeOH). Mp 242-244°.

N-Me: Synthetic. Cryst. (MeOH). Mp 218-220°.

Me ether, N-Me:Cryst. (C₆H₆). Mp 109-110°.

Me ether, N-Me, hydrochloride: Mp 238-240°.

Manske, R. *et al.*, *Can. J. Res.*, 1938, **16**, 76 (*isol*)

Barger, G. *et al.*, *J.C.S.*, 1939, 991 (*Artabotrinine*)

Faltis, F. *et al.*, *Ber.*, 1944, **77**, 686 (*synth*)

Marion, L. *et al.*, *J.A.C.S.*, 1944, **66**, 1125 (*synth*)

Schmutz, J. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 335 (*Xylopine, Isolaureline, isol, uv, synth*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 4066 (*Roemeroline*)


Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **21**, 1383 (*Xylopine, isol, pmr*)

Gellert, E. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2477 (*isol, pmr, ir, uv, ms*)

- Ziyaev, R. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 685; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 714 (*Isolaureline*)
- Pai, B.R. *et al.*, *Heterocycles*, 1977, **6**, 1993 (*Roemeroline, synth*)
- Suguna, H. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 416 (*synth, uv, pmr*)
- Ricca, G.S. *et al.*, *Gazz. Chim. Ital.*, 1979, **109**, 1 (*pmr, cmr, Xylopine*)
- Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 2251 (*Roemeroline*)
- Roblot, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 862 (*pmr, cmr, N-Formylxylopine*)
- Villar, A. *et al.*, *Farm. Tijdschr. Belg.*, 1984, **61**, 300 (*activity*)
- Simeon, S. *et al.*, *Pharmazie*, 1990, **45**, 442-443 (*activity*)
- Sturua, M. *et al.*, *CA*, 1998, **128**, 268208c (*Isolaureline N-oxide*)
- Yu, H.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1998, **45**, 773-778 (*N-Acetylanolobine*)
- Freyer, A.J. *et al.*, *Heterocycles*, 1999, **51**, 2221-2226 (*N-Formylanolobine*)
- Deng, Y. *et al.*, *Chin. Chem. Lett.*, 2002, **13**, 862-864 (*Me ether, N-methoxycarbonyl*)
- Nishiyama, Y. *et al.*, *Nat. Med. (Tokyo)*, 2004, **58**, 303-306; *CA*, **143**, 363319h (*N,N-Dimethylxylopine*)
- Carollo, C.A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1222-1224 (*N-Nitrosoxylopine*)

Anomaline† **A-1067**

1-Acetyl-16-methoxyaspidospermidine-6,17-diol, 9CI
[88660-02-0]



Relative
Configuration

$C_{22}H_{30}N_2O_4$ 386.49
Alkaloid from the bark and leaves of *Microplumeria anomala* (Apocynaceae). Needles (EtOAc). Mp 216-218°. λ_{max} 226 (log ϵ 4.27); 259 (log ϵ 3.85) (MeOH).


O¹²-Me: 12-O-Methylanomaline. *1-Acetyl-16,17-dimethoxyaspidospermidin-6-ol*, 9CI
[88660-03-1]
 $C_{23}H_{32}N_2O_4$ 400.517
Minor alkaloid from leaves of *Microplumeria anomala* (Apocynaceae).

Demethoxy: Demethoxyanomaline. *1-Acetylaspidospermidine-6,17-diol*, 9CI
[88669-30-1]
 $C_{21}H_{28}N_2O_3$ 356.464
Minor alkaloid from leaves of *Microplumeria anomala* (Apocynaceae). Needles (EtOAc). Mp 263-265°. λ_{max} 221 (log ϵ 4.13); 260 (log ϵ 3.72); 288 (log ϵ 3.47) (MeOH).

Luz, A.I.R. *et al.*, *Phytochemistry*, 1983, **22**, 2301-2304 (*isol, uv, ir, pmr, cmr, ms, struct*)

Anomoian A **A-1068**

[129602-20-6]




$C_{24}H_{31}Br_4N_3O_3$ 729.143
Metab. of the verongid sponge *Anom-*

oianthella popeae. Powder (as hydrochloride). Mp 200° dec. (hydrochloride). $[\alpha]_D^{25} +5.1$ (c, 0.013 in MeOH). λ_{max} 245 (ϵ 12300); 276 (ϵ 9400); 283 (ϵ 7800) (MeOH).

Kernan, M.R. *et al.*, *J. Nat. Prod.*, 1990, **53**, 720-723 (*isol, uv, ir, pmr, cmr, ms*)

Anomuricine **A-1069**

1,2,3,4-Tetrahydro-6,7-dimethoxy-1-[(4-methoxyphenyl)methyl]-5-isoquinolinol, 9CI. *1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-1-(4-methoxybenzyl)isoquinoline*
[78416-89-4]



$C_{19}H_{23}NO_4$ 329.395

(R)-form

Me ether, N,N-di-Me: N,N-Dimethylanomorurine
Quaternary alkaloid from the roots of *Xylopi parviflora*. Amorph. powder (as perchlorate). $[\alpha]_D^{22} -26.3$ (c, 0.32 in MeOH) (perchlorate). λ_{max} 277 (log ϵ 3.49); 283 (log ϵ 3.47) (MeOH) (perchlorate).

(ξ)-form


Minor alkaloid from the root and stem barks of *Annona muricata* (soursop) (Annonaceae). Noncryst.

Me ether 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1-[(4-methoxyphenyl)methyl]isoquinoline, 9CI. *Anomurine*
[78478-27-0]
 $C_{20}H_{25}NO_4$ 343.422
Minor alkaloid from the root and stem bark of *Annona muricata* (soursop) (Annonaceae). Noncryst.

Leboeuf, M. *et al.*, *Planta Med.*, 1981, **42**, 37 (*Anomurine, Anomuricine*)
Nishiyama, Y. *et al.*, *Phytochemistry*, 2004, **65**, 939-944 (*N,N-Dimethylanomorurine*)

Anopterimine **A-1070**

19,21-Didehydro-4-methyl-16-methylene-14,20-cycloveatchane-11,12-diol 12-(2-methyl-2-butenolate), 9CI
[60718-14-1]



$C_{25}H_{33}NO_3$ 395.541

Minor alkaloid from the leaves of *Anopterus macleayanus* (Escalloniaceae). Prisms (Me₂CO). Mp 235-238°. $[\alpha]_D^{25} +106$ (c, 0.6 in CHCl₃).

N-Oxide: Anopterimine N-oxide

[60706-77-6]


$C_{25}H_{33}NO_4$ 411.54

Minor alkaloid from the leaves of *Anopterus macleayanus* (Escalloniaceae). Needles (Me₂CO). Mp 233-235°. $[\alpha]_D^{25} +95$ (c, 1.1 in CHCl₃).

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1976, **29**, 1295; 1319 (*isol, ir, pmr, cmr, ms, struct*)

Anopteryl alcohol**A-1071**

[38826-63-0]



Absolute
Configuration

$C_{21}H_{31}NO_5$ 377.48

12-Tigloyl: Anopteryl 12-tiglate

$C_{26}H_{37}NO_6$ 459.581

Minor alkaloid from the root bark of *Anopterus macleayanus* (Escalloniaceae). Cryst. (Me₂CO). Mp 172-173°. $[\alpha]_D^{25} +15$ (c, 0.21 in CHCl₃). Sometimes resolidifies and remelts at 205° when slowly heated.

11,12-Ditigloyl: Anopterine. Anopteryl 11,12-ditiglate

[38826-62-9]

$C_{31}H_{43}NO_7$ 541.683

Alkaloid from leaves and bark of *Anopterus macleayanus* and *Anopterus glandulosus* (Escalloniaceae). Antineoplastic agent. Prisms (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 222-223°. $[\alpha]_D^{25} -12$ (c, 1.47 in CHCl₃). Log P 0.88 (uncertain value) (calc).

▶ LD₅₀ (mus, ipr) 100 mg/kg.

11-O-(4-Hydroxybenzoyl), 12-tigloyl: Anopteryl 11-4'-hydroxybenzoate 12-tiglate

$C_{33}H_{41}NO_8$ 579.689

Minor alkaloid from *Anopterus macleayanus* (Escalloniaceae). Prisms (Me₂CO). Mp 273-276°. $[\alpha]_D^{25} -28$ (c, 1.0 in CHCl₃/MeOH, 1:1).

7β-Hydroxy, 11,12-ditigloyl: Hydroxy-anopterine

$C_{31}H_{43}NO_8$ 557.683

Alkaloid from leaves and bark of *Anopterus macleayanus* and *Anopterus glandulosus* (Escalloniaceae). Needles (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 247-249°. $[\alpha]_D^{25} -14$ (c, 1.1 in MeOH).

7β-Hydroxy, 11-O-(4-hydroxy-3-methyl-2E-butenoyl), 12-tigloyl: Dihydroxy-anopterine

$C_{31}H_{43}NO_9$ 573.682

Minor alkaloid from *Anopterus macleayanus* and *Anopterus glandulosus*

(Escalloniaceae). Cryst. (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 242–244°. [α]_D²⁰ -9 (c, 1.2 in MeOH).

7β-Hydroxy, 11-benzoyl, 12-tigloyl: 7-Hydroxyanopteryl 11-benzoate 12-tiglate

C₃₃H₄₁NO₈ 579.689

Minor alkaloid from *Anopterus macleayanus* (Escalloniaceae). Amorph.

7β-Hydroxy, 11-O-(4-hydroxybenzoyl), 12-tigloyl: 7-Hydroxyanopteryl 11-4'-hydroxybenzoate 12-tiglate

C₃₃H₄₁NO₉ 595.688

Minor alkaloid from *Anopterus macleayanus* (Escalloniaceae). Noncryst.

Denne, W.A. *et al.*, *Tet. Lett.*, 1972, 2727 (*Anopterine, cryst struct*)


Hart, N.K. *et al.*, *Aust. J. Chem.*, 1976, **29**, 1295 (*isol, ir, pmr, cmr, ms, struct*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1985, **38**, 1091 (*Hydroxyanopterine, Dihydroxyanopterine*)

Anorldianine

A-1072

[123690-71-1]



C₂₇H₄₀N₄O₄ 484.637

Alkaloid from the stem bark of *Canthium anorldianum* (Rubiaceae). Pinkish cryst. (Et₂O). Mp 160°.

N²⁷-Oxide: Anorldianine N²⁷-oxide

C₂₇H₄₀N₄O₅ 500.637

Alkaloid from *Heisteria nitida*.


Dongo, E. *et al.*, *J. Nat. Prod.*, 1989, **52**, 840 (*isol, uv, ir, pmr, cmr, ms, struct*)

El-Seedi, H.R. *et al.*, *Phytochemistry*, 1999, **52**, 1739-1744 (*N²⁷-oxide*)

Anrakorinine

A-1073

20-(1,5-Dimethyl-2-piperidinyl)-pregn-5-ene-3,16,18-triol, 9CI
[78285-94-6]



C₂₈H₄₇NO₃ 445.684

Alkaloid from the aerial parts of *Fritillaria camtschatcensis* (Liliaceae). Needles (Me₂CO). Mp 248–251°. [α]_D²⁰ -50.4 (c, 0.79 in CHCl₃).


Tri-Ac:

Amorph.

Kaneko, K. *et al.*, *Tet. Lett.*, 1978, 2099-2102 (*isol, struct, synth, pmr*)

Antazirine

A-1074



C₁₇H₂₅Br₂NO₂ 435.198

(2*S*,4*E*)-form [170385-24-7]

Isol. from the sponge *Dysidea fragilis*. Oil. [α]_D²⁴ +16.7 (c, 0.58 in hexane) (30% e.e.). λ_{max} 213 ; 220 (sh) (MeOH).

(2*S*,4*Z*)-form [170554-75-3]


Isol. from *Dysidea fragilis*. Oil. [α]_D²⁴ +98.9 (c, 3.33 in hexane) (78% e.e.). λ_{max} 215 ; 220 (sh) (MeOH).

Salomon, C.E. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1463-1466 (*isol, uv, ir, pmr, cmr, ms, struct*)
Skepper, C.K. *et al.*, *J.O.C.*, 2008, **73**, 2592-2597 (*isol, pmr, cmr*)

Antholorin A

A-1075

[358334-50-6]



C₃₃H₄₇NO₃ 505.739

Prod. by *Aspergillus versicolor* OUPS-N136 isol. from the sea urchin *Anthocidaris crassispina*.

3-Hydroxy: Antholorin G

[358334-56-2]

C₃₃H₄₇NO₄ 521.739

Prod. by *Aspergillus versicolor* OUPS-N136 from the sea urchin *Anthocidaris crassispina*.

3-Epimer: Antholorin B

[358334-51-7]

C₃₃H₄₇NO₃ 505.739

Prod. by *Aspergillus versicolor* OUPS-N136 from the sea urchin *Anthocidaris crassispina*.

3-Epimer, 3-hydroxy: Antholorin H

[358334-57-3]

C₃₃H₄₇NO₄ 521.739

Prod. by *Aspergillus versicolor* OUPS-N136 from the sea urchin *Anthocidaris crassispina*.

13'-Epimer: Antholorin C

[358334-52-8]

C₃₃H₄₇NO₃ 505.739

Prod. by *Aspergillus versicolor* OUPS-N136 from the sea urchin *Anthocidaris crassispina*.

3,13'-Diepimer: Antholorin D

[358334-53-9]

C₃₃H₄₇NO₃ 505.739

Prod. by *Aspergillus versicolor* OUPS-


N136 from the sea urchin *Anthocidaris crassispina*.

Yamada, K. *et al.*, *CA*, 2001, **135**, 204898 (*isol*)

Antholorin E

A-1076

[358334-54-0]



C₃₃H₄₉NO₄ 523.754

Prod. by *Aspergillus versicolor* OUPS-N136 isol. from the sea urchin *Anthocidaris crassispina*.

3-Epimer: Antholorin F

[358334-55-1]

C₃₃H₄₉NO₄ 523.754


Prod. by *Aspergillus versicolor* OUPS-N136.

Yamada, K. *et al.*, *CA*, 2001, **135**, 204898 (*isol*)

Anthelvencin A

A-1077

[58616-25-4]



C₁₉H₂₅N₉O₃ 427.465

Oligopeptide antibiotic. Similar to Kikumycin A, K-43. Isol. from *Streptomyces venezuelae* ATCC14583.

Anthelmintic, antiparasitic and nematocidal agent. Sol. MeOH, acids; fairly sol. H₂O; poorly sol. Me₂CO, hexane. Mp 170-175° dec. [α]_D²² +9.7 (H₂O). λ_{max} 235 (ε 18700); 300 (ε 19300) (HCl salt/H₂O) (Derep). λ_{max} 235 (E1%/1cm 437); 300 (E1%/1cm 451) (H₂O) (Berdy).

▶ LD₅₀ (mus, ipr) 177 mg/kg. CA7808000

[113598-34-8, 113598-28-0]


Probst, G.W. *et al.*, *Antimicrob. Agents Chemother.*, 1965, 789

Lee, M. *et al.*, *J.O.C.*, 1988, **53**, 1855 (*synth*)

Anthocerdiazonin

A-1078

[159903-66-9]




C₁₉H₁₈N₂O₅ 354.362

Alkaloid from *in vitro* cultures of *Anthoceros agrestis*. Amorph. solid. $[\alpha]_D^{20} +73$.

Trennheuser, F. *et al.*, *Phytochemistry*, 1994, **37**, 899 (*isol, uv, ir, pmr, cmr, struct*)

Anthosamine A **A-1079**
3,4,5,6-Tetrahydro-3-[(4-hydroxyphenyl)-methylene]-2-pyridinecarboxylic acid, 9CI [165074-97-5]



$C_{13}H_{13}NO_3$ 231.251

Shows reversible equilib. between lactone and zwitterion form; the latter predominates in protic solv. the former in aprotic solvs. Metab. from the marine sponge *Anthosigmella* aff. *raramicroscclera*. Induces larval metamorphosis in ascidians. Yellow solid. λ_{max} 229 (ϵ 7300); 249 (ϵ 6400); 283 (ϵ 4200); 368 (ϵ 14500) (MeOH) (Berdy). λ_{max} 287 (ϵ 7000) (DMSO) (Berdy).


N-Me: **Anthosamine B**. 1,3,4,5-Tetrahydro-5-(4-hydroxyphenyl)-1-methylfuro[3,4-b]pyridin-7(2H)-one, 9CI [165074-96-4]

$C_{14}H_{15}NO_3$ 245.277

Metab. of *Anthosigmella* aff. *raramicroscclera*. Induces larval metamorphosis in ascidians. Yellow solid. $[\alpha]_D^{24} 0$ (c, 0.15 in MeOH). λ_{max} 249 (ϵ 7100); 369 (ϵ 18400); 453 (ϵ 1900) (MeOH) (Berdy). λ_{max} 285 (ϵ 6000); 357 (ϵ 2400); 486 (ϵ 3600) (DMSO) (Berdy).

Tsakamoto, S. *et al.*, *Tetrahedron*, 1995, **51**, 6687 (*isol, uv, ir, pmr, cmr, struct*)

Anthracyclinone blue B **A-1080**
Methyl 2-ethyl-5,7,12-trihydroxy-4-[(2-hydroxyethyl)amino]-6,11-dioxo-1-naphthacene-1-carboxylate [140900-60-3]



$C_{24}H_{21}NO_8$ 451.432

Prod. by a blocked mutant of *Streptomyces griseus* sp. ZIMET 43707. Deep blue solid.


12-Deoxy: **Anthracyclinone blue A** [140900-59-0]

$C_{24}H_{21}NO_7$ 435.432

Prod. by a blocked mutant of *Streptomyces galilaeus* S 383. Blue solid.

Eckardt, K. *et al.*, *J. Basic Microbiol.*, 1991, **31**, 371-376

Anthramycin, USAN **A-1081**
3-(5,10,11,11a-Tetrahydro-9,11-dihydroxy-8-methyl-5-oxo-1H-pyrrolo[2,1-c][1,4]benzodiazepin-2-yl)-2-propenamide, 9CI. **Anthramycin, INN**. *Refuin.* Ro 5-9000. *Antibiotic Ro* 5-9000 [4803-27-4]



$C_{16}H_{17}N_3O_4$ 315.328

Isol. from *Streptomyces refuineus* var. *thermotolerans* and *Streptomyces spadicogriseus*. Antibiotic with antineoplastic activity. DNA complexing agent. Small yellow prisms (Me₂CO aq.). Sol. DMF, DMSO, MeOH, butanol; poorly sol. hexane. Mp 188-194°. $[\alpha]_D^{25} +930$ (c, 1 in DMF). Log P -1.1 (uncertain value) (calc). Cryst. from hot aq. MeOH gives the Me ether. Epimerises in soln. λ_{max} 235 (ϵ 18200); 333 (ϵ 31800) (MeCN) (Derep). λ_{max} 220 (ϵ 34000); 235 (sh) (ϵ 30000); 335 (ϵ 39400) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 0.65 mg/kg. UY8429000
11-Me ether:

Pale-yellow needles (H₂O). Mp 120° dec. $[\alpha]_D^{25} +1002$ (c, 1 in DMF). λ_{max} 218 (ϵ 30000); 231 (ϵ 22100); 232 (sh); 334 (ϵ 40800); 360 (sh) (MeCN) (Derep).

Stereoisomer(?): **Antibiotic PBA**

[16397-13-0]

$C_{16}H_{17}N_3O_4$ 315.328

Isol. from *Streptomyces spadicogriseus*. Antitumour agent. Cryst. (Me₂CO/EtOAc). Stereochem. not determined, may be identical with Anthramycin.

Leimgruber, W. *et al.*, *J.A.C.S.*, 1965, **87**, 5791; 1968, **90**, 5641 (*isol, ir, uv, ms, nmr, struct, synth*)

Kohn, K.W. *et al.*, *Antibiotics (N.Y.)*, 1975, **3**, 3 (*rev*)

Horwitz, S.B. *et al.*, *Handb. Exp. Pharmacol.*, 1975, **38**, 642 (*rev. pharmacol*)

Hurley, L.H. *et al.*, *J. Antibiot.*, 1977, **30**, 349-370 (*rev, biosynth*)

U.S. Pat., 1977, 4 011 140; CA, **86**, 153957 (*Antibiotic PBA*)

Mostad, A. *et al.*, *Acta Chem. Scand., Ser. B*, 1978, **32**, 639 (*cryst struct*)

Komatsu, N. *et al.*, *J. Antibiot.*, 1980, **33**, 54 (*isol*)

Hurley, L.H. *et al.*, *Antibiotics (N.Y.)*, 1981, **4**, 262 (*rev*)

Ishikura, M. *et al.*, *Chem. Comm.*, 1982, 741 (*synth*)

Peña, M.R. *et al.*, *J.A.C.S.*, 1989, **111**, 5417 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, API000

Antibiotic 68-1147 **A-1082**
[66771-52-6]

Thiazole-peptide antibiotic complex. Struct. unknown. Prod. by *Micromonospora arborensis*. Antibacterial agent.

Antibiotic 68-1147-I
Sch 18640A. *Antibiotic Sch* 18640A [66771-53-7]

Mp 229-231° dec. $[\alpha]_D^{26} -87.5$ (c, 0.4 in

dioxan). λ_{max} 202 ; 247 ; 308 (EtOH).

Antibiotic 68-1147-II
68-1147-II. *Sch* 18640B. *Antibiotic Sch* 18640B

[66771-54-8] Sol. CHCl₃, EtOAc; poorly sol. H₂O, hexane, Et₂O. Mp 300° dec. $[\alpha]_D^{26} -65.5$ (c, 0.4 in dioxan). λ_{max} 240 ; 300 (EtOH). λ_{max} 240 (E1%/1cm 188); 300 (E1%/1cm 54) (EtOH) (Berdy).

U.S. Pat., 1978, 4 078 056; CA, **89**, 105873c

Antibiotic 69 **A-1083**


Struct. unknown. Related to Lyngbyatoxin A, L-367. Prod. by *Streptomyces* sp. No. 69. Antiparasitic and anthelmintic agent. Brown powder. Sol. H₂O, MeOH, AcOH, Me₂CO aq., Py; poorly sol. EtOAc, hexane.

Mori, R. *et al.*, *J. Antibiot.*, 1961, **14**, 280-285; 286-288 (*isol*)

Antibiotic 2158 **A-1084**

Antibiotic 1270

[182320-34-9]



$C_{33}H_{45}NO_4$ 519.723

Tetramic acid deriv. Similar to Oteromycin, O-134. Prod. by *Sclerophoma pythiophila* and *Sclerophoma* sp. MO1270. PAF synthesis inhibitor. Powder.

U.S. Pat., 1996, 5 550 148; CA, **125**, 257170w

Antibiotic 5102-2 **A-1085**

[83137-91-1]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces hygrosopicus-yingchengensis*. Powder. λ_{max} 224 ; 275 (1-propanol).

Zhang, S. *et al.*, CA, 1982, **97**, 143076t

Antibiotic 9408 **A-1086**

Tunicamycin-type antibiotic. Struct. unknown. Prod. by *Streptomyces* sp. 9408. Powder. $[\alpha]_D +68.6$. CAS no. not found to 2008.

► LD₅₀ (mus, ivn) 3.6 mg/kg, LD₅₀ (mus, ipr) 16.5 mg/kg.


Jpn. J. Antibiot., 1956, **9**, 201

Antibiotic X 14885A **A-1087**

X 14885A

[83917-57-1]

[83874-20-8 (Na salt)]



C₂₇H₃₂N₂O₇ 496.559

Polyether antibiotic related to Calcimycin, C-26. Isol. from *Streptomyces* sp. X-14885. Active against gram-positive bacteria and *Treponema hyodysenteriae*.

Cryst. + 1H₂O (as Na salt). Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O. Mp 264-266° (Na salt). [α]_D +177 (c, 1 in CHCl₃) (Na salt). λ_{max} 204 (ε 30700); 257 (ε 15100); 306 (ε 21800) (EtOH) (Derep).


Liu, C.-M. *et al.*, *J. Antibiot.*, 1983, **36**, 1118; 1275 (isol, uv, ir, struct)

U.S. Pat., 1983, 4 547 523; CA, **104**, 49844 (deriv)

Albrecht-Gary, A.M. *et al.*, *J.A.C.S.*, 1989, **111**, 8598 (pmr, props)

Antibiotic 0231A A-1088

8,8,9-Trimethoxy-5-methylbenz[cd]isoindolo[2,1-a]indol-1(8H)-one
[357924-11-9]



C₂₂H₁₉NO₄ 361.396

Prod. by *Streptomyces* sp. HKI 0231. Inhibitor of 3α-hydroxysteroid dehydrogenase. Yellow solid. Mp 125°. Fluorescent. λ_{max} 240 ; 285 ; 365 ; 380 ; 425 ; 440 ; 462 (MeOH).

5-Demethoxy: **Antibiotic 0231B**. 0231B
[357924-12-0]

C₂₁H₁₇NO₃ 331.37

Prod. by *Streptomyces* sp. HKI 0231. Inhibitor of 3α-hydroxysteroid dehydrogenase. Yellow solid. Mp 135°. Racemic. λ_{max} 240 ; 285 ; 365 ; 380 ; 420 ; 440 ; 460 (MeOH).


Kleinwachter, P. *et al.*, *J. Antibiot.*, 2001, **54**, 510-512 (isol)

Komoda, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2003, **67**, 659-662 (0231B, synth)

Kraus, G.A. *et al.*, *Tetrahedron*, 2005, **61**, 9502-9505 (0231B, synth)

Antibiotic 593A A-1089

3,6-Bis(5-chloro-2-piperidinyl)-2,5-piperazine-dione, 9CI
[41871-93-6]



Absolute configuration

C₁₄H₂₂Cl₂N₄O₂ 349.259

Diketopiperazine antibiotic. Isol. from *Streptomyces griseoluteus*. Shows antitumour activity. Sol. H₂O, MeOH, CHCl₃; poorly sol. hexane, Et₂O. Mp 235-240° dec. (as sulfate).

▶ LD₅₀ (mus, ipr) 5 mg/kg.

Gitterman, C.O. *et al.*, *J. Antibiot.*, 1970, **23**, 305 (isol)

Arison, B.H. *et al.*, *Tetrahedron*, 1973, **29**, 2743 (ir, uv, ms, nmr, struct)

Pettit, G.R. *et al.*, *J.A.C.S.*, 1976, **98**, 6742 (cryst struct, abs config)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1980, **102**, 2122 (synth)

Von Dreele, R.B. *et al.*, *Acta Cryst. B*, 1981, **37**, 93 (cryst struct)

Frank, R.K. *et al.*, *Diss. Abstr. Int., B*, 1983, **44**, 495 (synth)

Antibiotic 859A A-1090

859A

[97955-48-1]

Nucleoside-type antibiotic complex with 2 components. Struct. unknown. Prod. by *Streptomyces griseoruber*. Active against gram-positive and -negative bacteria. Powder.

U.S. Pat., 1985, 4 521 408; CA, **103**, 103458e

Antibiotic 875A A-1091

875A


[89491-83-8]

Nucleoside-type antibiotic complex. Prod. by *Streptomyces hygrosopicus* ATCC39665. Active against gram-positive and -negative bacteria. λ_{max} 262 (H₂O).

U.S. Pat., 1984, 4 427 655; CA, **100**, 155206d

Antibiotic 927A A-1092

2-Amino-5-(1H-indol-3-ylmethyl)-4(5H)-oxazolone, 9CI. C,N-Didemethyl-lindolmycin
[65954-89-4]




C₁₂H₁₁N₃O₂ 229.238

Isol. from *Streptomyces griseus*. Shows activity against gram-positive bacteria. Prisms (EtOAc). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 219-222°. λ_{max} 217 (ε 44550); 273 (ε 5080); 280 (ε 5310); 288 (ε 4580) (MeOH) (Berdy).

Hirota, A. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 147 (isol)

Antibiotic 2096A A-1093



C₁₁H₁₄N₈O₄ 322.283

Prod. by *Streptomyces* sp. IM 2096. Unstable.

Stereoisomer: **Antibiotic 2096B**

C₁₁H₁₄N₈O₄ 322.283

Prod. by *Streptomyces* sp. IM 2096.

Unstable.

Wang, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1641-1646

Antibiotic 2197A A-1094

SF 2197A

[98565-52-7]

Related to Matlystatin A₁. Struct. unknown. Prod. by *Microbispora* sp. SF-2197. Antibacterial agent. Cryst. (EtOAc/hexane). Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O, hexane. [α]_D -132 (MeOH). λ_{max} 200 (H₂O).

▶ LD₅₀ (mus, ipr) 100-300 mg/kg.

Japan. Pat., 1985, 85 83 585; CA, **103**, 159107g (isol)

Antibiotic A 6984 A-1095

A 6984

Thiazole-peptide antibiotic complex.

Struct. unknown. Prod. by *Actinoplanes utahensis* A6984. Antibacterial agent. No description found of component A.

Antibiotic A 6984B

A 6984B

Yellow powder. Sol. Py, DMSO; poorly sol. H₂O, C₆H₆, hexane. [α]_D +33 (Py). λ_{max} 218 ; 325 ; 416 ; 440 (MeOH). λ_{max} 218 (E1%/1cm 794); 231 (E1%/1cm 246); 351 (E1%/1cm 237) (MeOH/HCl) (Berdy). λ_{max} 215 (E1%/1cm 288); 414 (E1%/1cm 136) (MeOH/NaOH) (Berdy).

Antibiotic A 6984C

A 6984C

Yellow cryst. Sol. Py, DMSO; poorly sol. H₂O, C₆H₆, hexane. [α]_D +75 (Py). λ_{max} 218 (E1%/1cm 689); 324 (E1%/1cm 222); 412 (E1%/1cm 93) (MeOH) (Berdy). λ_{max} 218 (E1%/1cm 715); 300 (E1%/1cm 190); 333 (E1%/1cm 215); 351 (E1%/1cm 211) (MeOH/HCl) (Berdy). λ_{max} 232 (E1%/1cm 553); 295 (E1%/1cm 248); 408 (E1%/1cm 151) (MeOH/NaOH) (Berdy).

Antibiotic A 6984D

A 6984D

Yellow cryst. Sol. Py, DMSO; poorly sol. H₂O, hexane, C₆H₆. [α]_D +53 (Py). λ_{max} 219 (E1%/1cm 641); 327 (E1%/1cm 267); 414 (E1%/1cm 141) (MeOH) (Berdy). λ_{max} 219 (E1%/1cm 715); 300 (E1%/1cm 215); 333 (E1%/1cm 247); 352 (E1%/1cm 242) (MeOH/HCl) (Berdy). λ_{max} 232 (E1%/1cm 577); 298 (E1%/1cm 268); 410 (E1%/1cm 177) (MeOH/NaOH) (Berdy). Japan. Pat., 1979, 79 32 401; 79 32 402; CA, **91**, 18344; 18345

Antibiotic A 7413 A-1096

A 7413

Thiazole-peptide antibiotic complex.

Struct. unknown. Prod. by *Actinoplanes* sp. NRRL 8122.

Antibiotic A 7413A

A 7413A

[64296-61-3]

Light yellow cryst. (EtOH). Sol. C₂H₄Cl₂, MeOH, CHCl₃, DMSO, DMF; fairly sol. EtOH; poorly sol.

Me₂CO, C₆H₆, CCl₄, CH₂Cl₂, EtOAc, H₂O, Et₂O. Mp 205-212° dec. [α]_D²⁵ +54.5 (c, 2 in CHCl₃). λ_{max} 215 ; 260 ; 300 ; 358 (EtOH). λ_{max} 217 (E1%/1cm 440); 265 (E1%/1cm 227); 293 (E1%/1cm 210); 358 (E1%/1cm 95) (EtOH/HCl) (Berdy). λ_{max} 278 (E1%/1cm 255); 408 (E1%/1cm 80) (EtOH/NaOH) (Berdy).

► LD₅₀ (mus, ipr) 400 - 1600 mg/kg.

Antibiotic A 7413B

A 7413B

[64295-99-4]

Light yellow powder. Sol. MeOH, DMF, CHCl₃, DMSO, C₂H₄Cl₂; fairly sol. EtOH; poorly sol. Me₂CO, CCl₄, C₆H₆, H₂O, Et₂O, hexane. [α]_D²⁵ -26.2 (c, 7.5 in DMSO). λ_{max} 268 (E1%/1cm 104); 357 (E1%/1cm 30) (EtOH) (Berdy). λ_{max} 268 (E1%/1cm 108); 357 (E1%/1cm 35) (EtOH/HCl) (Berdy). λ_{max} 268 (E1%/1cm 179) (EtOH/NaOH) (Berdy).

Antibiotic A 7413C

A 7413C

[64296-00-0]

Light yellow powder. Sol. MeOH, CHCl₃, DMF, C₂H₄Cl₂, DMSO; fairly sol. EtOH; poorly sol. Me₂CO, H₂O, hexane, Et₂O, CCl₄, C₆H₆. Mp > 250°. λ_{max} 205 (E1%/1cm 356); 235 (E1%/1cm 180); 260 (E1%/1cm 127); 290 (E1%/1cm 104) (EtOH) (Berdy). λ_{max} 205 (E1%/1cm 356); 235 (E1%/1cm 180); 260 (E1%/1cm 127); 290 (E1%/1cm 103); 355 (E1%/1cm 40) (EtOH/HCl) (Berdy). λ_{max} 260 (E1%/1cm 268); 325 (E1%/1cm 189) (EtOH/NaOH) (Berdy).

U.S. Pat., 1979, 4 174 390; CA, 92, 56811g

Antibiotic A 10947

A-1097

A 10947

[72509-93-4]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Actinoplanes* sp. A10947. Antibacterial agent. Cryst. Sol. AcOH, Py, DMSO, CHCl₃; fairly sol. MeOH, EtOAc, EtOH, butanol; poorly sol. C₆H₆, hexane, H₂O. Mp 205-220° dec. [α]_D²⁵ +64.1 (c, 0.66 in CHCl₃). λ_{max} 217 ; 261 ; 302 ; 372 ; 409 (MeOH).

► LD₅₀ (mus, ipr) 500 - 1500 mg/kg.

Yaginuma, S. et al., *J. Antibiot.*, 1979, 32, 967-969


Japan. Pat., 1979, 79 157 502; CA, 92, 126912s

Antibiotic A 33853

A-1098

2-[3-Hydroxy-2-[(3-hydroxy-2-pyridinyl)carbonyl]amino]-4-benzoxazolecarboxylic acid, 9CI. A 33853

[80148-45-4]



C₂₀H₁₃N₃O₆ 391.339

Isol. from *Streptomyces* sp., NRRL 12068. Active against gram-positive bacteria, viruses, *Eimeria tenella*, trichomonads and mosquito larvae. Cryst. (CHCl₃/MeOH). Sol. DMSO, bases, CHCl₃, Py; fairly sol. Me₂CO, EtOAc, H₂O, MeOH; poorly sol. hexane. Mp 310-315° dec. Possible identity with Antibiotic U 60394. λ_{max} 255 (€ 39000); 310 (€ 17000); 360 (€ 11000); 400 (€ 11000) (dioxan/KOH) (Derep). λ_{max} 260 (sh) (€ 18000); 317 (€ 28300) (dioxan) (Derep). λ_{max} 322 (€ 28000) (dioxan-HC) (Berdy). λ_{max} 254 (€ 39000); 310 (€ 17000); 360 (€ 11000); 400 (€ 10500) (dioxan-NA) (Berdy).

► LD₅₀ (mus, ipr) 300 - 700 mg/kg. DM4675350

Tetra-Ac: [80161-47-3] Less active than parent compd. Less active than parent compd. Cryst. (CHCl₃/EtOH). Mp 184-189°.

U.S. Pat., 1980, 4 293 649; CA, 96, 33349 (isol) Michel, K.H. et al., *J. Antibiot.*, 1984, 37, 441 (isol, uv, ir, cryst struct)

Antibiotic A 65636

A-1099

A 65636

C₁₉H₂₃N₃O₄ 357.408


Benzodiazepine antibiotic. Struct. may be known, but lit. is inaccessible. Prod. by an actinomycete. Antibacterial agent. Yellow powder. λ_{max} 236 (€ 12630); 336 (€ 20870) (MeOH). λ_{max} 236 (€ 12632); 336 (€ 20870) (MeOH) (Berdy). λ_{max} 208 (€ 15500); 336 (€ 20000) (MeOH-HCl) (Berdy).

Intersci. Conf. Antimicrob. Agents Chemoth., 28th, 1987, 27

Antibiotic A 75943

A-1100

A 75943




C₁₅H₂₃NO₄ 281.351

Related to Cycloheximide in D-765. Prod. by *Streptomyces* sp. SANK 61296. Bone resorption inhibitor. Amorph. powder. [α]_D²⁵ -2.7 (c, 0.5 in CHCl₃). Morishita, T. et al., *J. Antibiot.*, 1998, 51, 531-538 (isol, synth, ir, pmr, cmr)

Antibiotic A 58365A

A-1101

3-Carboxy-1,2,3,5-tetrahydro-8-hydroxy-5-oxo-6-indolizinepropanoic acid, 9CI. A 58365A



C₁₂H₁₃NO₆ 267.238

λ_{max} 243 (€ 7200); 355 (€ 7400) (MeOH/NaOH) (Derep). λ_{max} 232 (€ 6000); 327 (€ 7600) (MeOH) (Derep).

(S)-form [87896-52-4]

Isol. from *Streptomyces chromofuscus*. Angiotensin converting enzyme inhibitor. Amorph. powder. Sol. H₂O, DMSO, EtOH, MeOH; fairly sol. Me₂CO; poorly sol. C₆H₆, hexane. [α]_D²⁵ -199.5 (c, 1 in H₂O).


(±)-form

Pale yellow plates (H₂O). Mp 163-165°. Nakatsukasa, W.M. et al., *J. Antibiot.*, 1985, 38, 997 (isol) Samlaska, S.K. et al., *J. Antibiot.*, 1985, 38, 1003 (isol, uv, ir, pmr, struct) Hunt, A.H. et al., *J. Antibiot.*, 1988, 41, 771 (pmr, cmr, uv, cryst struct) Fang, F.G. et al., *Tet. Lett.*, 1989, 30, 3621 (synth) Clive, D.L.J. et al., *Tet. Lett.*, 1998, 39, 2519-2522 (synth) Straub, C.S. et al., *Org. Lett.*, 1999, 1, 83-85 (synth) Reichelt, A. et al., *Tetrahedron*, 2002, 58, 6323-6328 (synth)

Antibiotic A 500359A

A-1102

A 500359A



C₂₄H₃₃N₅O₁₂ 583.551

Homologue of Capuramycin, C-103. Prod. by *Streptomyces griseus* SANK 60196. Powder. [α]_D²⁰ +94.7 (c, 1 in MeOH). λ_{max} 257 (€ 10300) (H₂O).

2'-O-Carbamoyl: *Antibiotic A 503083A*. A 503083A

C₂₅H₃₄N₆O₁₃ 626.576
Prod. by *Streptomyces* sp. SANK 62799. Powder. [α]_D²⁰ +84 (c, 0.7 in H₂O). Stereochem. not confirmed. λ_{max} 257 (€ 11000) (H₂O).

O-De-Me: *Antibiotic A 500359C*. A 500359C

C₂₃H₃₁N₅O₁₂ 569.524
Prod. by *Streptomyces griseus* SANK 60196. Powder. [α]_D²⁰ +89 (c, 0.44 in H₂O). λ_{max} 257 (€ 10700) (H₂O).

3''-Deoxy: *Antibiotic A 500359D*. A 500359D

C₂₄H₃₃N₅O₁₁ 567.552
Prod. by *Streptomyces griseus* SANK 60196. Powder. [α]_D²⁰ +68 (c, 0.69 in H₂O). λ_{max} 244 (€ 10000) (H₂O).



Muramatsu, Y. et al., *J. Antibiot.*, 2003, 56, 243-252; 253-258; 259-267; 268-279 (isol, pmr, cmr, biosynth)

Muramatsu, Y. et al., *J. Antibiot.*, 2004, 57, 639-646 (A 503083A)

Muramatsu, Y. et al., *J. Antibiot.*, 2006, 59, 601-606 (prodn)

Antibiotic A 10255B
Thioplabin B. A 10255B
[119166-38-0]


A-1103

Clive, D.L.J. *et al.*, *J.O.C.*, 1999, **64**, 1447-1454 (synth)C₅₃H₄₈N₁₆O₁₅S₃ 1245.258

Cyclic thiopentapeptide antibiotic. Prod. by *Streptomyces gardneri* and *Streptomyces* sp. R1401. Feed additive. Noncryst. Sol. DMSO, THF-H₂O, CHCl₃-MeOH, Py, DMF, MeOH, CHCl₃, Me₂CO; poorly sol. H₂O. λ_{max} 247 (ε 78700) (EtOH) (Derep). λ_{max} 245 (ε 66000) (MeOH) (Berdy). λ_{max} 247 (ε 78700) (MeOH/HCl) (Berdy). λ_{max} 245 (ε 66000) (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 300 - 700 mg/kg.Boeck, L.D. *et al.*, *J. Antibiot.*, 1992, **45**, 1222; 1278 (isol, biosynth)Favret, M.E. *et al.*, *J. Antibiot.*, 1992, **45**, 1809-1811 (biosynth)Debono, M. *et al.*, *J.O.C.*, 1992, **57**, 5200-5208 (isol, pmr, cmr, struct)Ohyama, S. *et al.*, *J. Antibiot.*, 2002, **55**, 83-91 (Thioplabin)

Antibiotic A 58365B A-1104
4-Carboxy-1,3,4,6-tetrahydro-9-hydroxy-6-oxo-2H-quinolizine-7-propanoic acid, 9CI. A 58365B. Alkaloid A 58365B
[87896-53-5]

C₁₃H₁₅NO₆ 281.265

Isol. from *Streptomyces chromofuscus*. Angiotensin converting enzyme inhibitor. Amorph. powder. Sol. H₂O, MeOH, DMSO, EtOH; fairly sol. Me₂CO; poorly sol. C₆H₆, hexane. [α]_D²⁵ -141.2 (c, 0.16 in H₂O). λ_{max} 243 (ε 7200); 355 (ε 7400) (MeOH/NaOH) (Derep). λ_{max} 232 (ε 6000); 327 (ε 7600) (MeOH) (Derep). λ_{max} 233 (ε 3800); 332 (ε 4500); 333 (ε 4500) (MeOH) (Berdy). λ_{max} 244; 360 (MeOH-NaOH) (Berdy).

Nakatsukasa, W.M. *et al.*, *J. Antibiot.*, 1985, **38**, 997 (isol)Samlaska, S.K. *et al.*, *J. Antibiot.*, 1985, **38**, 1003 (isol, uv, ir, pmr, struct)Hunt, A.H. *et al.*, *J. Antibiot.*, 1988, **41**, 771 (pmr, cmr, uv, cryst struct)

Antibiotic A 10255C A-1105
A 10255C
[119166-39-1]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces gardneri*. Feed additive. Yellow powder. Sol. DMSO, DMF, THF-H₂O, MeOH, CHCl₃-MeOH, CHCl₃, Me₂CO, Py; poorly sol. H₂O. Mp 221-224°. [α]_D +66.6 (EtOH). λ_{max} 243 (ε 63000) (MeOH).

▶ LD₅₀ (mus, ipr) 300 - 700 mg/kg.U.S. Pat., 1993, 5 229 362; CA, **119**, 269171m

Antibiotic A 58365C A-1106
A 58365C
[87913-25-5]

C₁₄H₁₇NO₆ 295.291

Quinolizine antibiotic. Related to Antibiotic A 58365A, A-1101. Prod. by *Streptomyces chromofuscus* NRRL 15098. ACE inhibitor. Hypotensive agent. Powder. Sol. H₂O, DMSO, MeOH, EtOH; fairly sol. Me₂CO; poorly sol. C₆H₆, hexane. λ_{max} 233 (ε 3800); 332 (ε 4500) (MeOH). λ_{max} 244; 360 (MeOH/NaOH) (Berdy).

Eur. Pat., 1982, 103 403; CA, **100**, 207884v

Antibiotic A 7413D A-1107
A 7413D

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Actinoplanes* sp. Antibacterial agent.

Ger. Pat., 1977, 2 703 938; CA, **87**, 165997q

Antibiotic A 10255E A-1108
Thioplabin C. A 10255E
[145427-74-3]

As Antibiotic A 10255B, A-1103 with R¹ = CONHC(=CH₂)CONHC(=CH₂)CONHC(=CH₂)COOH, R² = CH(CH₃)₂

C₅₄H₅₀N₁₆O₁₅S₃ 1259.285

Thiazole peptide antibiotic. Prod. by *Streptomyces gardneri* and *Streptomyces* sp. R1401. Yellow powder.

Favret, M.E. *et al.*, *J. Antibiot.*, 1992, **45**, 1499-1511; 1809-1811 (isol, pmr, cmr)U.S. Pat., 1993, 5 229 362; CA, **119**, 269171m (isol)Ohyama, S. *et al.*, *J. Antibiot.*, 2002, **55**, 83-91 (Thioplabin)

Antibiotic A 10255F A-1109
A 10255F
[119166-41-5]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces gardneri*. Feed additive. Yellow powder. Sol. DMSO, CHCl₃, Me₂CO, DMF, THF-H₂O, MeOH, CHCl₃-MeOH, Py; poorly sol. H₂O. λ_{max} 245 (ε 71500) (MeOH).

▶ LD₅₀ (mus, ipr) 300 - 700 mg/kg.U.S. Pat., 1993, 5 229 362; CA, **119**, 269171m

Antibiotic A 10255H A-1110
A 10255H
[119166-43-7]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces gardneri*. Feed additive. Yellow powder. Sol.

DMSO, Me₂CO, CHCl₃, MeOH, DMF, Py, CHCl₃-MeOH, THF-H₂O; poorly sol. H₂O. λ_{max} 244 (ε 74000) (MeOH). λ_{max} 245 (ε 74500) (MeOH/HCl) (Berdy). λ_{max} 211 (ε 23800) (MeOH/NaOH) (Berdy).


▶ LD₅₀ (mus, ipr) 300 - 700 mg/kg.U.S. Pat., 1993, 5 229 362; CA, **119**, 269171m

Antibiotic AB 97 A-1111
AB 97
[68652-25-5]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Actinomadura helvata* var. *antibiotica*. Active against gram-positive bacteria. Yellow powder. Sol. EtOAc, bases, DMSO, MeOH; poorly sol. acids, hexane. [α]_D +764 (DMSO). λ_{max} 245; 290; 350; 410 (MeOH). λ_{max} 245; 360 (MeOH/HCl) (Berdy). λ_{max} 295; 415 (MeOH/NaOH) (Berdy).

Japan. Pat., 1978, 78 101 301; CA, **90**, 37630q

Antibiotic AB 3217A A-1112
AB 3217A
[139158-99-9]

C₁₇H₂₃NO₇ 353.371

Prod. by *Streptomyces platensis*. Active against mites. Needles (MeOH). Sol. H₂O, MeOH; fairly sol. EtOAc, CHCl₃; poorly sol. hexane. Mp 241°. [α]_D²⁴ -52.5 (c, 1 in H₂O). λ_{max} 226 (ε 11400); 272 (ε 1100); 278 (ε 900) (H₂O) (Derep). λ_{max} 226 (ε 10900); 272 (ε 1100); 279 (ε 900) (NaOH) (Berdy). λ_{max} 226 (ε 11400); 272 (ε 1100); 278 (ε 900) (H₂O) (Berdy). λ_{max} 227 (ε 11300); 272 (ε 1100); 279 (ε 900) (HCl) (Berdy).

13-O-(6-Hydroxy-6-methylheptanoyl):

Antibiotic AB 3217B. AB 3217B

[139159-00-5]

C₂₅H₃₇NO₉ 495.569

Prod. by *Streptomyces platensis*. Active against mites. Sol. H₂O, MeOH; fairly sol. EtOAc, CHCl₃; poorly sol. hexane. Mp 116° (monohydrochloride). [α]_D²⁵ -68.3 (c, 1 in H₂O). λ_{max} 226 (ε 11500); 272 (ε 990); 278 (ε 842) (H₂O) (Derep). λ_{max} 226 (E1%/1cm 233); 272 (ε 20); 278 (E1%/1cm 17) (H₂O) (Berdy).

13-O-(4-Methylpentanoyl): **Antibiotic****AB 3217C. AB 3217C**

[139159-01-6]

C₂₃H₃₃NO₈ 451.516


Prod. by *Streptomyces platensis*. Active against mites. Sol. H₂O, MeOH; fairly sol. EtOAc, CHCl₃; poorly sol. hexane. Mp 117° (monohydrochloride). [α]_D²⁵ -76.8 (c, 1 in H₂O). λ_{max} 226 (ε 13000);

272 (ϵ 1260); 278 (ϵ 1130) (H₂O) (Derep). λ_{\max} 226 (E1%/1cm 288); 272 (E1%/1cm 28); 278 (E1%/1cm 25) (H₂O) (Berdy).

Kanbe, K. *et al.*, *J. Antibiot.*, 1992, **45**, 458; 568 (*isol, pmr, cmr, props*)
Nakata, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 3057 (*synth*)

Antibiotic AB 4015B

AB 4015B
[162857-79-6]




C₂₆H₃₉NO₃ 413.599

Tetracycline deriv. Prod. by *Phoma* sp. AB4015. Active against bacteria and fungi. Orange oil. Sol. CHCl₃, MeOH. λ_{\max} 228 (ϵ 4450); 290 (ϵ 6740) (MeOH). λ_{\max} 228 (ϵ 4450); 290 (ϵ 8580) (MeOH-HCl) (Berdy). λ_{\max} 248 (ϵ 7390); 290 (ϵ 9370) (MeOH-NaOH) (Berdy).

Japan. Pat., 1995, 95 48 348; *CA*, **122**, 289047r

Antibiotic AB 4063B

AB 4063B
[160041-33-8]



C₂₆H₃₉NO₃ 413.599

Tetracycline deriv. Prod. by *Phoma* sp. AB4063. Active against phytopathogenic fungi. λ_{\max} 228 (ϵ 6200); 290 (ϵ 12800) (MeOH) (Berdy). λ_{\max} 204 (ϵ 12600); 226 (ϵ 6600); 290 (ϵ 11100) (MeOH-HCl) (Berdy). λ_{\max} 209 (ϵ 9100); 248 (ϵ 10300); 290 (ϵ 12700) (MeOH-NaOH) (Berdy).

1'',2''-Epoxide: **Antibiotic AB 4063A**. AB 4063A

[160041-32-7]

C₂₆H₃₉NO₄ 429.598


Prod. by *Phoma* sp. AB4063. Active against phytopathogenic fungi. λ_{\max} 228 (ϵ 6500); 292 (MeOH) (Berdy). λ_{\max} 227 (ϵ 6700); 292 (ϵ 13400) (MeOH-HCl) (Berdy). λ_{\max} 246 (ϵ 11200); 290 (ϵ 14300) (MeOH-NaOH) (Berdy).

Japan. Pat., 1994, 94 277 084; *CA*, **122**, 54155k

Antibiotic AF 140

A-1115

9,10,13,14,15,16,20,20a-Octahydro-15-hydroxy-19-methoxy-2,6-dimethyl-9-phenylcyclopent[1]azacyclonadecine-7,17(8H,19H)-dione, 9CI. AF 140 [157375-70-7]



C₃₀H₃₇NO₅ 491.626


Prod. by *Amycolatopsis* sp. TA 0234. Antifungal and cytotoxic agent. Cryst. (MeOH). Sol. MeOH.

Japan. Pat., 1994, 94 178 693; *CA*, **122**, 131134g (*isol*)

Antibiotic AJI 9561

A-1116

AJI 9561



C₂₂H₁₄N₂O₅ 386.363

Similar to Antibiotic UK 1, A-1302. Prod. by *Streptomyces* sp. AJ9561. Cytotoxic. Powder. Mp 251-252°. λ_{\max} 250 (ϵ 27700); 272 (ϵ 23800); 281 (ϵ 24900); 311 (ϵ 37500); 319 (ϵ 38100); 328 (ϵ 37200); 347 (ϵ 27000); 361 (ϵ 21600) (MeOH).

Sato, S. *et al.*, *J. Antibiot.*, 2001, **54**, 102-104

Antibiotic ASK 753

A-1117

ASK 753

[54650-03-2]

Related to Ferrimycin A₁, F-37. Prod. by *Streptomyces* sp. AS-K-753. Siderophore. Antibacterial agent. Off-white cryst. Sol. MeOH, Et₂O, CHCl₃; fairly sol. H₂O, EtOAc; poorly sol. C₆H₆, hexane. Mp 120° dec. λ_{\max} 270 (EtOH). λ_{\max} 270 (E1%/1cm 120) (EtOH) (Berdy). λ_{\max} 270 (E1%/1cm 120) (EtOH-HCl) (Berdy). λ_{\max} 287 (E1%/1cm 135) (EtOH-NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 58 mg/kg.

Shimi, I.R. *et al.*, *J. Antibiot.*, 1969, **22**, 106-111

Antibiotic AS-N-7A

A-1118

AS-N-7A

[60730-59-8]

Related to Ferrimycin A₁, F-37. Prod. by *Streptomyces erythrochromogenes*. Siderophore. Antibacterial agent.


▶ LD₅₀ (mus, ims) 1.1 mg/kg.

Abdallah, N.M. *et al.*, *CA*, 1976, **85**, 171695q

Antibiotic AT 2433

A-1119

AT 2433



Antibiotic AT 2433A₁ R¹ = Cl, R² = CH₃
AT 2433A₂ R¹ = Cl, R² = H
AT 2433B₁ R¹ = H, R² = CH₃
AT 2433B₂ R¹ = R² = H

Nucleoside-type antibiotic complex. Stereochem. of sugar residues revised in 1999. Closely related to Rebeccamycin, R-32. Prod. by *Actinomadura mellioura* and *Actinomadura fulva*. Shows antibacterial and antineoplastic props.

Antibiotic AT 2433A₁

AT 2433A₁

[102644-20-2]

C₃₄H₃₅ClN₄O₉ 679.125

Sol. MeOH, EtOAc, CHCl₃-MeOH, DMSO; fairly sol. THF, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 200 (ϵ 30800); 235 (ϵ 39900); 284 (ϵ 34300); 315 (ϵ 45500); 395 (ϵ 3860) (MeOH) (Derep). λ_{\max} 234 (ϵ 38645); 286 (ϵ 33598); 314 (ϵ 43890); 394 (ϵ 3453) (MeOH) (Berdy).

Antibiotic AT 2433A₂

AT 2433A₂

[102644-19-9]

C₃₃H₃₃ClN₄O₉ 665.098

Sol. MeOH, CHCl₃-MeOH, DMSO, EtOAc; fairly sol. THF, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 200 (ϵ 30800); 235 (ϵ 39900); 284 (ϵ 34300); 315 (ϵ 45500); 395 (ϵ 3860) (MeOH) (Derep). λ_{\max} 234 (ϵ 38645); 286 (ϵ 33598); 314 (ϵ 43890); 394 (ϵ 3453) (MeOH) (Berdy).

Antibiotic AT 2433B₁

AT 2433B₁

[102622-96-8]

C₃₄H₃₆N₄O₉ 644.68

Sol. MeOH, CHCl₃-MeOH, EtOAc, DMSO; fairly sol. THF, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 202 (ϵ 29000); 234 (ϵ 41300); 283 (ϵ 33600); 316 (ϵ 46900); 400 (ϵ 4060) (MeOH) (Derep). λ_{\max} 234 (ϵ 41280); 284 (ϵ 33617); 316 (ϵ 46948); 400 (ϵ 4057) (MeOH) (Berdy).

Antibiotic AT 2433B₂

AT 2433B₂

[102622-95-7]

C₃₃H₃₄N₄O₉ 630.653

Sol. DMSO, CHCl₃-MeOH; fairly sol. THF, EtOAc; poorly sol. cyclohexane. λ_{\max} 202 (ϵ 29000); 234 (ϵ 41300); 283 (ϵ 33600); 316 (ϵ 46900); 400 (ϵ 4060) (MeOH) (Derep). λ_{\max} 233 (ϵ 41076); 282

(ϵ 33390); 315 (ϵ 46935); 400 (ϵ 3969) (MeOH) (Berdy).

Eur. Pat., 1986, 175 309; *CA*, **105**, 5184 (*isol*)
Matson, J.A. *et al.*, *J. Antibiot.*, 1989, **42**, 1547;
1784 (*isol*, *pmr*, *cmr*, *struct*)
Chisholm, J.D. *et al.*, *J.A.C.S.*, 1999, **121**,
3801-3802 (*synth*, *config*)
Chisholm, J.D. *et al.*, *J.O.C.*, 2000, **65**, 7541-
7543; 8406 (*synth*)

Antibiotic 1401B A-1120
1401B

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces corchorusii*. Antibacterial agent. Yellow powder. Sol. DMF, CHCl_3 , dioxan, AcOH; fairly sol. MeOH, EtOH; poorly sol. acids, bases. $[\alpha]_D^{25}$ -93.8 (dioxan).

Gauze, G.F. *et al.*, *Antibiotiki (Moscow)*, 1983, **28**, 643-647

Antibiotic 4181B A-1121
4181B

[115088-45-4]

$\text{C}_{28}\text{H}_{19}\text{NO}_9$ 513.459

Quinone antibiotic. Struct. was under investigation; prob. identical with De-O-methylcevinomycin A_2 in C-287, but no further reports to 2007. *Isol.* from *Streptomyces griseus*. Exhibits antibacterial, antifungal and antitumour props. Red-orange powder. Sol. MeOH, EtOAc; fairly sol. CHCl_3 , EtOAc, DMSO, MeOH; poorly sol. H_2O , hexane, Et_2O . Mp 260° dec. λ_{max} 258 (ϵ 22700); 327 (ϵ 27000); 377 (sh) (ϵ 5220); 418 (ϵ 3770) (MeOH) (Derep). λ_{max} 280 ; 330 ; 410 (MeOH) (Berdy). λ_{max} 258 (ϵ 22700); 327 (ϵ 18000); 418 (ϵ 3770) (CHCl_3) (Berdy).


► LD_{50} (mus, ipr) 40 - 60 mg/kg.
CB9133810

Otani, T. *et al.*, *J. Antibiot.*, 1988, **41**, 275-281 (*isol*, *props*)

Antibiotic B 4317 A-1122

B 4317

[173075-46-2]



$\text{C}_{28}\text{H}_{32}\text{N}_4\text{O}_9$ 568.582

Prod. by *Ateromonas rava*. Leucotriene antagonist. Antibacterial and anti-inflammatory agent. Siderophore. Powder. $[\alpha]_D^{25}$ -50.2 (c, 1.3 in CHCl_3). λ_{max} 207 (ϵ 59600); 240 (ϵ 19800); 303 (ϵ 1050) (MeOH).


Japan. Pat., 1995, 95 291 992; *CA*, **124**, 115565j

Antibiotic B 90063 A-1123

5,5'-Dithiobis[1,4-dihydro-4-oxo-6-(2-pentyl-4-oxazolyl)-2-pyridinecarboxaldehyde], 9CI. *B 90063*. *5,5'-Dithiobis[4-hydroxy-6-(2-pentyl-4-oxazolyl)-2-pyridinecarboxaldehyde]*

[215377-24-5]

[181634-31-1]



$\text{C}_{28}\text{H}_{30}\text{N}_4\text{O}_6\text{S}_2$ 582.7

Prod. by the marine bacterium *Blastobacter* sp. SANK 71894. Endothelin-converting enzyme inhibitor. Yellow cryst. Mp 73-74° Mp 116-118° (+ 2MeOH). The alternative CAS no. refers to the bis(4-hydroxypyridine) tautomeric struct. under which it was originally descr. λ_{max} 214 (ϵ 47400); 261 (ϵ 37200); 340 (sh) (ϵ 8100); 360 (sh) (ϵ 4100) (MeCN).

Japan. Pat., 1996, 96 208 646; *CA*, **125**, 219743c
Takaishi, S. *et al.*, *J. Antibiot.*, 1998, **51**, 805-
815 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Antibiotic B 1625FA 2β -1 A-1124

B 1625FA 2β -1

[111420-30-5]



$\text{C}_{37}\text{H}_{49}\text{N}_7\text{O}_{12}$ 783.834

Actinomycin-related antibiotic. *Isol.* from *Streptomyces antibioticus*. Orange-red powder. Sol. MeOH, CHCl_3 ; poorly sol. H_2O . Mp 185°. $[\alpha]_D^{25}$ +3 (CHCl_3). λ_{max} 427 (ϵ 32100); 443 (ϵ 32300) (MeOH- CHCl_3) (Berdy).

Fujimoto, Y. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 803 (*isol*, *struct*, *props*)

Antibiotic BA 181314A A-1125

BA 181314A

$\text{C}_{28}\text{H}_{26}\text{N}_2\text{O}_{11}$ 566.52


Struct. unknown. Related to Albofungin, A-244. Prod. by *Streptomyces aspergilloides* ATCC14808. Pale yellow plates (MeOH). Sol. Py, DMF, DMSO; fairly sol. H_2O , Me_2CO , CHCl_3 , EtOAc, EtOH, MeOH; poorly sol. C_6H_6 , hexane. Dec. at 315-320°. λ_{max} 250 (E1%/1cm 700); 361 (E1%/1cm 400); 375 (E1%/1cm 450) (MeOH) (Berdy).

U.S. Pat., 1967, 3 328 248; *CA*, **67**, 89788s

Antibiotic BE 10988 A-1126

2-(5-Amino-4,7-dihydro-1-methyl-4,7-dioxo-1H-indol-3-yl)-4-thiazolecarboxamide, 9CI. *BE 10988*

[135261-89-1]



$\text{C}_{13}\text{H}_{10}\text{N}_4\text{O}_3\text{S}$ 302.313

Prod. by *Streptomyces* sp. BA10988.

Cytotoxic. Topoisomerase II inhibitor. Dark red cryst. Sol. DMSO; fairly sol. MeOH; poorly sol. H_2O . Mp 300°. λ_{max} 213 (ϵ 43200); 235 (sh) (ϵ 24200); 278 (ϵ 22300); 305 (sh) (ϵ 13300); 318 (sh) (ϵ 12100); 385 (ϵ 7300); 525 (ϵ 2400) (MeOH).

Oka, H. *et al.*, *J. Antibiot.*, 1991, **44**, 486-491 (*isol*, *uv*, *ir*, *pmr*)


Suda, H. *et al.*, *Tet. Lett.*, 1991, **32**, 2791-2792 (*pmr*, *cmr*)

Rauhut, T. *et al.*, *J.O.C.*, 2008, **73**, 5279-5286 (*biosynth*, *cmr*, *ms*)

Antibiotic BE 14106 A-1127

BE 14106. *GT 32A*. *Antibiotic GT 32A*

[140212-86-8]



$\text{C}_{27}\text{H}_{37}\text{NO}_3$ 423.594

Macrocyclic lactam antibiotic. Prod. by *Streptomyces spheroides* A14106 and *Streptomyces* sp. GT 32. Cytotoxic agent. Amorph. Sol. DMSO; fairly sol. MeOH; poorly sol. H_2O . Mp 270°. λ_{max} 230 (sh) (ϵ 11800); 278 (ϵ 91800); 288 (ϵ 110800); 310 (sh) (ϵ 22000); 325 (sh) (ϵ 16100) (MeOH) (Derep).

► CL6400000

8-Deoxy: Antibiotic GT 32B. *GT 32B*.

Immunosuppressant GT 32B

[178560-35-5]

$\text{C}_{27}\text{H}_{37}\text{NO}_2$ 407.595

Prod. by *Streptomyces* sp. GT 32.

Immunosuppressant. Powder. Sol. MeOH, DMSO, butanol. λ_{max} 280 (ϵ 74000); 289 (ϵ 92000); 310 (sh) (ϵ 20000); 325 (sh) (ϵ 15000) (MeOH).

Kojiri, K. *et al.*, *J. Antibiot.*, 1992, **45**, 868-874 (*isol*, *ir*, *pmr*, *cmr*)


Takahashi, I. *et al.*, *J. Antibiot.*, 1997, **50**, 186-188 (*GT 32*)

Mitchell, S.S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1400-1402 (*pmr*, *cmr*)

Antibiotic BE 22179 A-1128

BE 22179

[147816-33-9]



Absolute Configuration


$\text{C}_{46}\text{H}_{48}\text{N}_{10}\text{O}_{12}\text{S}_4$ 1061.209

Depsipeptide antibiotic. Prod. by *Streptomyces* sp. A22179. Topoisomerase II inhibitor. Antitumour agent. Active against gram-positive bacteria. Pale yellow cryst. $[\alpha]_D^{20}$ -96.6 (c, 0.47 in CHCl_3). λ_{max} 218 (ϵ 87500); 226 (ϵ 84600); 290 (ϵ 12800); 360 (ϵ 12000) (MeOH).

Okada, H. *et al.*, *J. Antibiot.*, 1994, **47**, 129-135 (*isol*, *uv*, *ir*, *pmr*, *activity*)


Boger, D.L. *et al.*, *J.A.C.S.*, 2001, **123**, 561-568
(*synth, abs config*)

Antibiotic BE 48021 A-1129
BE 48021
[182230-51-9]




$C_{22}H_{27}NO_5$ 385.459
Prod. by *Streptomyces* sp. A48021. Cytotoxic agent. Sol. MeOH. λ_{max} 227 ; 249 ; 269 ; 291 ; 332 ; 344 (MeOH) (Berdy).
Japan. Pat., 1996, 96 198 874; *CA*, **125**, 273720f

Antibiotic BE 52211 A-1130
BE 52211
[205449-00-9]




$C_{24}H_{35}NO_3$ 385.545
Prod. by *Streptomyces* sp. A52211. Cytotoxic agent. λ_{max} 212 ; 240 (MeOH).
Japan. Pat., 1998, 98 67 725; *CA*, **128**, 269581f (*isol*)

Antibiotic BE 54017 A-1131
BE 54017
[274694-01-8]




$C_{23}H_{18}ClN_3O_5$ 451.865
Prod. by *Streptomyces* sp. A54017. Antileukaemic agent.
Japan. Pat., 2000, 178 274; *CA*, **133**, 42244c

Antibiotic BE 54476 A-1132
BE 54476
[261360-41-2]




$C_{22}H_{33}NO_5$ 391.506
Prod. by a *Streptomyces* sp. Antibacterial agent.
Japan. Pat., 2000, 86 627; *CA*, **132**, 221452v

Antibiotic BE 55051 A-1133
3,5,6,10-Tetrachloro-1-(5-chloro-2-hydroxybenzoyl)-2H,8H-dipyrrolo[3,4-b:1',2'-d][1,4]benzoxazocin-8-one, 9Cl. BE 55051
[222548-15-4]




$C_{22}H_9Cl_5N_2O_4$ 542.588
Prod. by *Streptomyces* sp. A55051. Antitumour agent.
Japan. Pat., 1999, 99 71 379; *CA*, **130**, 280924g

Antibiotic BE 67251 A-1134
BE 67251
[261625-75-6]



$C_{28}H_{39}NO_4$ 453.62
Prod. by *Streptomyces* sp. A67251. Antitumour agent.
Japan. Pat., 2000, 86 664; *CA*, **132**, 221455y

Antibiotic BE 19412A A-1135
BE 19412A
[173774-90-8]




$C_{28}H_{22}ClNO_8$ 535.936
Prod. by *Streptomyces* sp. A19412. Cy-

tototoxic agent. Greyish-green powder.
 λ_{max} 243 ; 287 ; 412 ; 431 (MeOH).

▶ LD_{50} (mus, ipr) 200 mg/kg.
10,12-Di-Me ether: Antibiotic BE 19412B. BE 19412B
[173774-91-9]
 $C_{30}H_{26}ClNO_8$ 563.99
Semisynthetic. Cytotoxic agent. Dark yellow powder. λ_{max} 234 ; 290 ; 338 ; 404 (MeOH) (Berdy).
Tsukamoto, M. *et al.*, *J. Antibiot.*, 1998, **51**, 908-914 (*isol, uv, ir, pmr, cmr*)

Antibiotic BE 54238A A-1136
BE 54238A
[205433-26-7]




Absolute Configuration

$C_{22}H_{23}NO_6$ 397.427
Prod. by *Streptomyces* sp. A54238. Cytotoxic agent. Orange-yellow powder. λ_{max} 276 (ϵ 16700); 411 (ϵ 27700) (MeOH).

1 α -Hydroxy, 2' \rightarrow 1-lactone: Antibiotic BE 54238B. BE 54238B
[205433-27-8]
 $C_{22}H_{21}NO_6$ 395.411
Prod. by *Streptomyces* sp. A54238. Cytotoxic agent. Orange-yellow powder. Mp 209° dec. (hydrochloride). $[\alpha]_D^{25}$ -521 (c, 0.29 in DMSO) (hydrochloride). λ_{max} 278 (ϵ 21500); 420 (ϵ 27600) (MeOH).

Tsukamoto, M. *et al.*, *J. Antibiot.*, 2000, **53**, 26-32 (*isol, pmr, N-15 nmr*)
Tatsuta, K. *et al.*, *J. Antibiot.*, 2004, **57**, 291-297 (*synth, pmr*)

Antibiotic BE 52211C A-1137
BE 52211C



Absolute Configuration

$C_{23}H_{33}NO_2$ 355.519
Analogue of Antibiotic BE 52211, A-1130. Prod. by *Streptomyces* sp. Cytotoxic. Pale yellow oil. $[\alpha]_D^{25}$ -38 (c, 0.008 in MeOH). Obt. as a mixt. with BE 52211B to which the data refers. λ_{max} 224 (log ϵ 4.41); 238 (sh) (log ϵ 4.37) (MeOH).

Δ^{17} -Isomer: Antibiotic BE 52211B. BE 52211B
 $C_{23}H_{33}NO_2$ 355.519
Prod. by a *Streptomyces* sp. Obt. as a mixt. with BE52211C.

Kubota, N.K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 85-87 (*isol, pmr, cmr, ms*)

Antibiotic BMY 42428

A-1138

BMY 42428

[140879-94-3]

C₅₂H₆₉NO₂₃ 1076.11

Related to Albofungin, A-244. Struct. unknown. Prod. by *Actinoadura mardurae* ATCC53806. Cytotoxic agent active against gram-positive bacteria. Greenish-orange cryst. (MeOH/CHCl₃/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 226 (E1%/1cm 239); 254 (E1%/1cm 262); 324 (E1%/1cm 157); 396 (E1%/1cm 76) (MeOH) (Berdy). λ_{max} 224 (E1%/1cm 292); 254 (E1%/1cm 317); 324 (E1%/1cm 187); 396 (E1%/1cm 85) (MeOH/HCl) (Berdy). λ_{max} 222 (E1%/1cm 302); 246 (E1%/1cm 315); 344 (E1%/1cm 120); 444 (E1%/1cm 95) (MeOH/NaOH) (Berdy).


U.S. Pat., 1992, 5 087 567; CA, 116, 172425x

Antibiotic BR 040

A-1139

BR 040

[167503-59-5]

C₂₆H₃₇NO₈ 491.58

Glutarimide-type antibiotic. Prod. by *Streptomyces hygroscopicus* var. *ossamyceticus*. Osteophatic and bone absorption inhibitor. Powder. [α]_D -160 (MeOH). λ_{max} 204 (ε 17000) (MeOH). λ_{max} 204 (ε 17000) (MeOH) (Berdy).

8-Me ether: **Antibiotic BR 042**. BR 042 [167503-60-8]

C₂₇H₃₉NO₈ 505.607

Prod. by *Streptomyces hygroscopicus* var. *ossamyceticus*. Osteophatic and bone absorption inhibitor. Powder. [α]_D -64 (MeOH). λ_{max} 205 (ε 14000) (MeOH). λ_{max} 205 (ε 14000) (MeOH) (Berdy).


Japan. Pat., 1995, 95 138 257; CA, 123, 167722g (isol)

Antibiotic BU 4514N

A-1140

BU 4514N

[151013-39-7]

C₂₇H₄₂N₂O₅ 474.639

Prod. by a *Microtetraspora* sp. Neurotoxic agent. Also active against gram-positive bacteria. Fine needles + ³H₂O (MeOH aq.). Sol. acids, DMSO, MeOH; poorly sol. H₂O, CHCl₃, EtOAc. Mp 185-186°. [α]_D²⁵ -140 (c, 0.5 in 0.1M HCl). Related to Lydicamycin, L-353. λ_{max} 204 (ε 6540); 254 (sh) (ε 4320); 286 (ε 9000) (MeOH/HCl) (Derep). λ_{max} 205 (ε 8440); 248 (ε 9200); 287 (ε 8200) (MeOH/NaOH) (Derep). λ_{max} 204 (ε 6300); 251 (ε 4360); 286 (ε 8530) (MeOH) (Derep). λ_{max} 204 (ε 6540); 286 (ε 9000) (MeOH-HCl) (Berdy).


Toda, S. et al., J. Antibiot., 1993, 46, 875 (isol, pmr, cmr, struct, props)

Antibiotic CJ 12662

A-1141

CJ 12662

[169062-89-9]



Absolute Configuration

C₂₉H₃₇ClN₂O₇ 561.073

Related to Paeciloxazine, P-9. Prod. by *Aspergillus fischeri* var. *thermomutans* ATCC 18618. Antiparasitic agent. Sol. MeOH, Me₂CO, EtOAc, CHCl₃; fairly sol. H₂O. λ_{max} 217; 248; 287 (MeOH).

Dechloro: **Antibiotic CJ 12663**. CJ 12663 [169062-90-2]C₂₉H₃₈N₂O₇ 526.628

Prod. by *Aspergillus fischeri* var. *thermomutans* ATCC 18618. Antiparasitic agent. Sol. MeOH, CHCl₃, EtOAc, Me₂CO; fairly sol. H₂O. λ_{max} 217; 248; 287 (MeOH).

8'a-Deoxy, deacetoxy: **Antibiotic UK**

88051. UK 88051

[138399-19-6]

C₂₇H₃₅ClN₂O₄ 487.037

Prod. by *Chryso sporium* sp. N845. Antiparasitic agent. Powder. Stereochem. not confirmed. λ_{max} 217 (ε 17530); 248 (ε 4300); 287 (ε 1390) (MeOH).

U.K. Pat., 1991, 2 240 100; CA, 116, 39777x (UK 88051)

Pat. Coop. Treaty (WIPO), 1995, 95 19 363;


CA, 123, 254718d (CJ 12662, CJ 12663)

Didier, C. et al., J.O.C., 2004, 69, 7875-7879 (synth, abs config)

Antibiotic CJ 16170

A-1142

2,3-Dihydro-4-hydroxy-3-methyl-2-(1-methyl-1-propenyl)-7-phenylfuro[3,2-c]pyridine-3-methanol. CJ 16170

C₁₉H₂₁NO₃ 311.38

Prod. by *Cladobotryum varium* CL12284. Glass. [α]_D²⁴ +15.4 (c, 0.26 in MeOH). λ_{max} 206 (ε 22000); 246 (ε 21000) (MeOH).

Sakemi, S. et al., J. Antibiot., 2002, 55, 6-18 (isol, pmr, cmr)


Clive, D.L.J. et al., Tetrahedron, 2002, 58, 10243-10250 (epimer, synth)

Snider, B.B. et al., Org. Lett., 2004, 6, 2877-2880 (synth)

Antibiotic CJ 16171

A-1143

CJ 16171

C₁₉H₁₉NO₃ 309.364


Prod. by *Cladobotryum varium* CL12284. Needles (MeOH). Mp 243-244°. [α]_D²⁴ -22.7 (c, 0.33 in MeOH). λ_{max} 207 (ε 28000); 247 (ε 27000) (MeOH).

Sakemi, S. et al., J. Antibiot., 2002, 55, 6-18 (isol, pmr, cmr, cryst struct)

Antibiotic CJ 16173

A-1144

CJ 16173

C₁₉H₁₉NO₄ 325.363


Prod. by *Cladobotryum varium* CL12284. Needles (MeOH). Mp 234-235°. [α]_D²⁴ +205 (c, 0.12 in MeOH). λ_{max} 208 (ε 28000); 235 (ε 25000) (MeOH).

Sakemi, S. et al., J. Antibiot., 2002, 55, 6-18 (isol, pmr, cmr, cryst struct)

Antibiotic CJ 16174

A-1145

2,3-Dihydro-4-hydroxy-2-(1-methyl-1-propenyl)-5-phenylfuro[2,3-b]pyridine-3,3-dimethanol. CJ 16174

C₁₉H₂₁NO₄ 327.379

Prod. by *Cladobotryum varium* CL12284. Needles (MeOH). Mp 214-216°. [α]_D²⁴ +40 (c, 0.02 in MeOH). λ_{max} 207 (ε 15000); 234 (ε 13000) (MeOH).

6'-Deoxy: **Antibiotic CJ 16169**. CJ 16169

C₁₉H₂₁NO₃ 311.38
Prod. by *Cladobotryum varium*
CL12284. Needles (MeOH). Mp 207-208°. [α]_D²⁴ +52.3 (c, 0.09 in MeOH). λ_{\max} 208 (ε 24000); 233 (ε 19000) (MeOH).

6'-Deoxy, 5'-aldehyde: Cladobotryal.

Antibiotic CJ 15696. CJ 15696

[189010-01-3]

C₁₉H₁₉NO₃ 309.364

Prod. by *Cladobotryum varium*.

Needles (MeOH) (as hemiacetal).

Mp 198-199° (hemiacetal). [α]_D²⁴ +21.2

(c, 0.52 in MeOH) (CJ 15696). [α]_D²⁴

+65 (c, 0.8 in CHCl₃) (Cladobotryal).

λ_{\max} 209 (ε 24000); 234 (ε 19000)

(MeOH).

Brinholt, J. et al., *Acta Chem. Scand.*, 1998,

52, 631-634 (Cladobotryal)

Sakemi, S. et al., *J. Antibiot.*, 2002, 55, 6-18


(isol, pmr, cmr, cryst struct)

Snider, B.B. et al., *Org. Lett.*, 2004, 6, 2877-2880 (synth)

Antibiotic CJ 16196

A-1146

2,3-Dihydro-3-(3-hydroxy-2-methyl-1-butenyl)-3-methyl-7-phenylfuro[3,2-c]pyridine-2,4-diol. CJ 16196



C₁₉H₂₁NO₄ 327.379

Prod. by *Cladobotryum varium*

CL12284. Glass. [α]_D²⁴ -7.4 (c, 0.11 in

MeOH). λ_{\max} 206 (ε 23000); 245 (ε

23000) (MeOH).

3'-Epimer: Antibiotic CJ 16197. CJ 16197

C₁₉H₂₁NO₄ 327.379

Prod. by *Cladobotryum varium*

CL12284. Glass. [α]_D²⁴ +2.2 (c, 0.14 in

MeOH). λ_{\max} 207 (ε 22000); 245 (ε

22000) (MeOH).


Sakemi, S. et al., *J. Antibiot.*, 2002, 55, 6-18

(isol, pmr, cmr)

Antibiotic CJ 16264

A-1147

CJ 16264



Relative Configuration

C₂₃H₃₁NO₅ 401.502

Related to Antibiotic UCS 1025B, A-

1301. Prod. by the fungal strain

CL39457. Antibacterial agent. Powder.

[α]_D²³ +27.3 (c, 0.11 in MeOH). λ_{\max} 208

(ε 12000); 225 (sh) (ε 6200) (MeOH).


Sugie, Y. et al., *J. Antibiot.*, 2001, 54, 917-925

(isol, uv, pmr, cmr, activity)

Antibiotic CJ 16367

A-1148

CJ 16367



C₂₄H₃₃NO₅ 415.528

Prod. by the fungal strain CL39457.

Antibacterial agent. Powder. [α]_D²⁴ +17 (c,

0.11 in MeOH). λ_{\max} 207 (ε 24000); 226

(sh) (ε 10000) (MeOH).


Sugie, Y. et al., *J. Antibiot.*, 2001, 54, 917-925

(isol, uv, pmr, cmr, activity)

Antibiotic CJ 17572

A-1149

CJ 17572



C₂₁H₃₁NO₄ 361.48

Tetramic acid antibiotic. Prod. by the

fungus *Pezicula* sp. CL11877. Antibac-

terial incl. MDR bacteria. Amorph.

powder. [α]_D²⁵ +110.5 (c, 0.42 in MeOH).

λ_{\max} 255 (ε 12700); 292 (ε 15700)

(MeOH).


Sugie, Y. et al., *J. Antibiot.*, 2002, 55, 19-24

(isol, uv, pmr, cmr, activity)

Antibiotic CJ 21058

A-1150

CJ 21058



C₂₃H₃₃NO₄ 387.518

Tetramic acid antibiotic. Methyl analo-

gue of Equisetin, E-142. Prod. by an

unidentified fungus CL47745. SecA in-

hibitor. Antibacterial agent. Cytotoxic.

Powder. [α]_D²³ +318 (c, 0.5 in MeOH).

λ_{\max} 205 (ε 13500); 252 (ε 6100); 293 (ε

7900) (MeOH).

Sugie, Y. et al., *J. Antibiot.*, 2002, 55, 25-29

(isol, pmr, cmr, activity)

Antibiotic CP 35763

A-1151

Antibiotic 35763. CP 35763

[64735-13-3]

C₂₆H₃₇N₃O₇ 503.594

Cyclic peptide antibiotic (Mikamycin A

group). Probably identical with Madu-

mycin II, M-35. Prod. by *Actinoplanes*

auranticolor and *Actinoplanes azureus*.

Active against gram-positive bacteria.

Animal growth promoter. Powder. Sol.

MeOH, CHCl₃, CH₂Cl₂; poorly sol.

Et₂O, hexane. Mp 100° dec. [α]_D²⁵ -114 (c,

1 in EtOH). Inseparable from A2315A.

λ_{\max} 218 (EtOH). λ_{\max} 218 (E1%/1cm

669) (EtOH) (Berdy).

Ger. Pat., 1975, 2 516 020; CA, 84, 87960

U.S. Pat., 1977, 4 038 383; CA, 88, 20525v

Antibiotic CP 41043

A-1152

CP 41043. Antibiotic 41043

[63799-02-0]

Thiazole-peptide antibiotic. Struct. un-

known. Prod. by *Pseudonocardia fasti-*

diosa. Antibacterial agent. Amorph.

solid. Sol. Me₂CO, DMSO, DMF,

EtOH, CHCl₃, EtOAc; poorly sol.

Et₂O, H₂O, hexane. [α]_D +77 (c, 1 in

Me₂CO). λ_{\max} 225 (E1%/1cm 444); 270

(E1%/1cm 217); 350 (E1%/1cm 91)

(EtOH) (Berdy).

U.S. Pat., 1977, 4 031 206; CA, 87, 100678s

Antibiotic CP 41494

A-1153

CP 41494. Antibiotic 41494

[63799-03-1]

Thiazole-peptide antibiotic. Struct. un-

known. Prod. by *Pseudonocardia fasti-*

diosa. Antibacterial agent. Amorph.

solid. Sol. Me₂CO, EtOH, DMF, DMSO,

EtOAc, CHCl₃; poorly sol. Et₂O, hexane,

H₂O. [α]_D +29 (c, 0.5 in Me₂CO). λ_{\max}

270 ; 295 ; 350 (EtOH).


U.S. Pat., 1977, 4 031 206; CA, 87, 100678s

Antibiotic CV 1

A-1154

CV 1

[108351-34-4]



C₇H₁₄N₂O₆ 222.197

Aminoglycoside-related antibiotic. Isol.

from *Streptomyces* sp. CO 1. Lipopoly-

saccharide synthesis inhibitor active

against gram-negative bacteria and *E.*

coli. Synergistic with Spiramycin. Pow-

der. Sol. H₂O, DMSO; poorly sol.

MeOH, hexane. Mp 85-90°. [α]_D²⁸ +66 (c,

0.9 in H₂O).


Ichimura, M. et al., *J. Antibiot.*, 1987, 40, 723;

727 (isol, struct, props)

Antibiotic 0089D

A-1155

5-Methyl-2-[(1-methyl-1H-indol-3-yl)-methylene]-3,5-hexadienenitrile, 9CI
[215876-60-1]



$C_{17}H_{16}N_2$ 248.327


Prod. by *Nocardia brasiliensis*. Antitumour and antibacterial agent.

Japan. Pat., 1998, 98 287 647; CA, 130, 3119j

Antibiotic DC 81

A-1156

1,2,3,11a-Tetrahydro-8-hydroxy-7-methoxy-5H-pyrrolo[2,1-c]benzodiazepin-5-one. DC 81
[89824-22-6]



$C_{13}H_{14}N_2O_3$ 246.265


Anthramycin-type antibiotic. Prod. by *Streptomyces roseiscleroticus*. Weakly active against gram-positive and -negative bacteria and tumours. Powder. Sol. MeOH, EtOAc, Py; poorly sol. H₂O, hexane. Mp 98-105°. $[\alpha]_D^{22} +135$ (c, 0.2 in MeOH). λ_{max} 224 ; 236 ; 316 (MeOH) (Berdy).

- ▶ LD₅₀ (mus, ipr) 42 mg/kg. UY8546700
Japan. Pat., 1983, 83 180 487; CA, 100, 173150 (isol, struct, nmr)
- Weidner-Wells, M.A. et al., J.O.C., 1989, 54, 5746 (synth)
- Wang, T. et al., Org. Lett., 1999, 1, 1835-1837 (synth)
- Hu, W.-P. et al., J.O.C., 2001, 66, 2881-2883 (synth)

Antibiotic DC 102

A-1157

DC 102
[115722-50-4]



Absolute Configuration

$C_{24}H_{35}N_3O_6$ 461.557

Isol. from a *Streptomyces* sp. Weakly active against gram-positive bacteria. Active against murine leukaemia. Powder. Sol. MeOH, Me₂CO; fairly sol. H₂O; poorly sol. CHCl₃, hexane, EtOAc. Mp 120° dec. λ_{max} 210 (€ 14000); 244 (sh) (€ 11000); 310 (€ 6500) (MeOH) (Derep).

- ▶ LD₅₀ (mus, ipr) 1-5 mg/kg. UY8534200

Demethoxy, 12,13-didehydro: **Sibanomicin**. SF 2364. Antibiotic SF 2364

[117782-84-0]

[119180-42-6]

$C_{23}H_{31}N_3O_5$ 429.515

Anthramycin-type antibiotic. Prod. by *Micromonospora* sp. SF2364. Antitumour agent. Amorph. solid (as hydrochloride). Sol. H₂O, butanol, MeOH; poorly sol. EtOAc, hexane. Mp 185-195° dec. (hydrochloride). $[\alpha]_D^{22} +59$ (c, 1 in H₂O). $[\alpha]_D^{22} +371$ (c, 0.2 in DMSO). λ_{max} 206 (€ 37700); 240 (sh) (€ 15400); 310 (€ 2140) (0.1N HCl) (Derep). λ_{max} 217 (€ 21400); 240 (sh) (€ 14900); 314 (€ 4330) (0.1N NaOH) (Derep). λ_{max} 214 (€ 35400); 240 (sh) (€ 13500); 314 (€ 3630) (H₂O) (Derep).

- ▶ LD₅₀ (mus, ipr) 1.3 mg/kg, LD₅₀ (mus, ivn) 2.1 mg/kg.


Hara, M. et al., J. Antibiot., 1988, 41, 702 (DC 102)

Itoh, J. et al., J. Antibiot., 1988, 41, 1281 (Sibanomicin)

Antibiotic DC 105

A-1158

DC 105
[123731-93-1]



$C_{24}H_{31}N_3O_6$ 457.525

Benzodiazepine antibiotic. Similar to Sibiromycin. Prod. by *Streptomyces* sp. DO-105. Cytotoxic agent. Yellow cryst. Sol. CHCl₃, MeOH, DMSO; fairly sol. butanol; poorly sol. H₂O. $[\alpha]_D^{25} +567.5$ (CHCl₃). λ_{max} 268 (MeOH). λ_{max} 268 (H₂O) (Berdy). λ_{max} 270 ; 302 ; 440 (HCl) (Berdy). λ_{max} 280 ; 360 ; 410 ; 425 (NaOH) (Berdy).


- ▶ LD₅₀ (mus, ipr) .147 mg/kg.

Japan. Pat., 1989, 89 121 296; CA, 111, 230672k

Antibiotic DC 8118A

A-1159

DC 8118A
[150045-16-2]



$C_{13}H_{20}N_2O_5$ 284.311

Prod. by *Streptomyces* sp. (DO-118; BP3595). Antibacterial and antitumour agent. Sol. MeOH, Me₂CO; poorly sol. H₂O. $[\alpha]_D^{25} -176$. λ_{max} 212 (€ 4600); 288 (€ 2600) (MeOH) (Berdy).

8'-Ketone: Antibiotic DC 8118B. DC 8118B

[150045-17-3]

$C_{13}H_{18}N_2O_5$ 282.296


Prod. by *Streptomyces* sp. (DO-118; BP3595). Antibacterial and antitumour agent. Sol. MeOH, Me₂CO; poorly sol. H₂O. $[\alpha]_D^{25} -245$. λ_{max} 286 (€ 3800) (MeOH) (Berdy).

Pat. Coop. Treaty (WIPO), 1993, 09 110; CA, 119, 158353 (isol, pmr, cmr, uv)

Antibiotic DOB 41

A-1160

6-[1-(3-Hydroxy-2-methoxy-1-oxopropoxy)ethyl]-1-phenazinecarboxylic acid, 9CI. DOB 41
[115666-98-3]



Relative configuration

$C_{19}H_{18}N_2O_6$ 370.361


Phenazine antibiotic. Isol. from a *Pseudomonas* sp. Active against gram-positive bacteria and a murine leukaemia P388. Yellow needles. Sol. MeOH, DMSO, CHCl₃; poorly sol. Et₂O, hexane, H₂O. Mp 201°. λ_{max} 255 (€ 76700); 370 (€ 15700) (CHCl₃) (Derep).

Shoji, T. et al., J. Antibiot., 1988, 41, 589 (isol, pmr, cmr, struct)

Antibiotic ECO 02301

A-1161

ECO 02301



$C_{70}H_{108}N_2O_{20}$ 1297.625


Prod. by *Streptomyces aizunensis* NRRL B-11277. Antifungal agent. Yellow solid. λ_{max} 319 (€ 110000); 334 (€ 146000); 351 (€ 128000) (MeOH).

McAlpine, J.B. et al., J. Nat. Prod., 2005, 68, 493-496 (isol, pmr, cmr)

Antibiotic EGS 1300-1

A-1162

EGS 1300-1
[218937-20-3]




$C_{62}H_{95}ClNO_{12}P$ 1112.859

Prod. by *Penicillium* sp. G-1300. Sialyltransferase inhibitor. Red glassy solid. Related to Isochromophilone VI, I-205.

Japan. Pat., 1998, 98 330 364; CA, 130, 80425h

Antibiotic F 1778 A-1163

F 1778
[204992-07-4]




C₂₈H₃₃NO₆ 479.572

Tetramic acid deriv. Prod. by the discomycete *Tapesia* sp. SANK 18896. Antifungal agent.

Tanaka, A. et al., *Annu. Rep. Sankyo Res. Lab.*, 1997, 49, 135-141; CA, 128, 241597s

Antibiotic F 1839A A-1164

F 1839A
[159096-43-2]



Probable Absolute Configuration

C₂₃H₃₁NO₅ 401.502

Struct. of Stachybotrylactam revised in 2003. The structs. of other members of the group must currently be regarded as probable only. Prod. by *Stachybotrys* sp. F-1839. Inhibitor of pancreatic cholesterol esterase. Amorph. powder. Sol. MeOH, Py, DMSO; fairly sol. EtOAc, 2-butanone; poorly sol. H₂O, hexane. [α]_D²⁵ -29.6 (c, 0.1 in MeOH). λ_{max} 218 (ε 37400); 262 (ε 6550); 301 (ε 3180) (MeOH) (Derep).

2,3-Di-Ac: [163391-78-4]

C₂₇H₃₅NO₇ 485.576

Prod. by *Stachybotrys chartarum* and *Stachybotrys complementi*. Immunosuppressive. Powder. [α]_D -29 (MeOH). λ_{max} 216 ; 258 ; 299 (MeCN).

N-(2-Hydroxyethyl): **Antibiotic F 1839E**, F 1839E

[159096-46-5]

C₂₅H₃₅NO₆ 445.555

From *Stachybotrys* sp. F-1839. Inhibitor of cholesterol esterase, anticholesterolemic. Amorph. powder. Sol. MeOH, Py, DMSO; fairly sol. EtOAc, 2-butanone; poorly sol. H₂O, hexane. λ_{max} 217 (ε 31500); 262 (ε 6410); 302 (ε 2310) (MeOH) (Derep).

N-(3-Carboxypropyl): [214132-31-7]

C₂₇H₃₇NO₇ 487.592

Metab. of *Stachybotrys chartarum*.

N-(3-Methoxycarbonylpropyl): **Antibiotic F 1839F**, F 1839F

[159096-48-7]

C₂₈H₃₉NO₇ 501.619

From *Stachybotrys* sp. F-1839. Inhibitor of cholesterol esterase, anticholesterolemic. Amorph. powder. Sol. MeOH, Py, DMSO; fairly sol. EtOAc, 2-butanone; poorly sol. H₂O, hexane. λ_{max} 217 (ε 33100); 267 (ε 5290); 300 (ε 2530) (MeOH) (Derep).

N-(4-Carboxybutyl): **Antibiotic Q 11270B**, Q 11270B

[150459-98-6]

C₂₈H₃₉NO₇ 501.619

Prod. by *Stachybotrys chartarum*.

Powder. [α]_D²⁵ +25.5 (c,0.19 in MeOH). C-2 config. not certain.

N-(1-Carboxy-2-methylpropyl): [685141-56-4]

C₂₈H₃₉NO₇ 501.619

Metab. of *Stachybotrys chartarum*.

Amorph. solid.

N-(1-Carboxy-2-methylbutyl): [685141-58-6]

C₂₉H₄₁NO₇ 515.645

Constit. of *Stachybotrys chartarum*.

N-(1-Carboxy-3-methylbutyl): [685141-57-5]

C₂₉H₄₁NO₇ 515.645

Constit. of *Stachybotrys chartarum*.

N-(1-Carboxy-2-phenylethyl): [685141-59-7]

C₃₂H₃₉NO₇ 549.663

Constit. of *Stachybotrys chartarum*.

Pale yellow solid.

2-Deoxy: **Stachybotrylactam**, Spirodihydrobenzofuranlactam I

[163391-76-2]

[173220-01-4]

C₂₃H₃₁NO₄ 385.502

Prod. by *Stachybotrys chartarum* and *Stachybotrys complementii*.

Immunosuppressant. Endothelin

antagonist, inhibitor of HIV-1 protease. Powder. λ_{max} 216 ; 257 ; 298 (MeCN).

2-Deoxy, 3-Ac: [163391-77-3]

C₂₅H₃₃NO₅ 427.539

Prod. by *Stachybotrys chartarum* and *Stachybotrys complementi*

ATCC20511. Immunosuppressant.

Powder. λ_{max} 216 ; 257 ; 298

(MeCN).

2-Deoxy, N-(2-hydroxyethyl): **Stachybotramide**, Spirodihydrobenzofuranlactam II, Stachybotrin†

[149598-71-0]

[149691-32-7 ; 207453-12-1 ; 173220-02-5]

C₂₅H₃₅NO₅ 429.555

Prod. by *Stachybotrys alternans*,

Stachybotrys chartarum and

Stachybotrys cylindrospora. Cryst.

(MeOH). Mp 202°. [α]_D²⁴ -12.5 (c, 0.8 in CHCl₃/MeOH) (tri-Ac).

2-Deoxy, N-(1-carboxy-2-hydroxypropyl): [685141-55-3]

C₂₇H₃₇NO₇ 487.592

Metab. of *Stachybotrys chartarum*.

Amorph. solid.

2-Deoxy, N-(1,3-dicarboxypropyl): **Spirodihydrobenzofuranlactam IV**

[173220-03-6]

[685141-54-2]

C₂₈H₃₇NO₈ 515.602

Prod. by *Stachybotrys chartarum*.

Enzyme inhibitor, HIV-1 protease; tyrosine kinase inhibitor. Amorph. solid.

2-Deoxy, N-(4-carboxybutyl): **Spirodihydrobenzofuranlactam III**, Q 11270A.

Antibiotic Q 11270A

[150459-97-5]

C₂₈H₃₉NO₆ 485.619

Prod. by *Stachybotrys chartarum* and

another *Stachybotrys* sp. [α]_D²⁵ -167.5 (c,0.04 in MeOH).

2-Deoxy, N-(3-methoxycarbonylbutyl): **Antibiotic F 1839J**, F 1839J

[159096-47-6]

C₂₉H₄₁NO₆ 499.646

From *Stachybotrys* sp. F-1839. Cho-

lesterol esterase inhibitor, anticholesterolemic. Amorph. powder. Sol.

MeOH, Py, DMSO; fairly sol. EtOAc,

2-butanone; poorly sol. H₂O, hexane.

λ_{max} 217 (ε 33100); 267 (ε 5290); 300 (ε

2530) (MeOH) (Derep). λ_{max} 218 (ε

32041); 261 (ε 9412); 301 (ε 2323)

(MeOH) (Berdy).

2-Deoxy, 4'-Me ether, N-(2-hydroxyethyl): **Stachybotrin A†**

[256520-70-4]

C₂₆H₃₇NO₅ 443.582

Prod. by *Stachybotrys alternans*.

8'-Oxo, 2-deoxy: **Spirodihydrobenzofuranlactam V**

[173220-04-7]

C₂₃H₂₉NO₅ 399.486

Prod. by *Stachybotrys chartarum* and

Stachybotrys atra.

7'-Deoxy, 8'-oxo: **Antibiotic F 1839D**, F 1839D

[159096-45-4]

C₂₃H₃₁NO₅ 401.502

From *Stachybotrys* sp. F-1839. Inhibi-

tor of cholesterol esterase, antichole-

lesterolemic. Amorph. powder. Sol.

MeOH, Py, DMSO; fairly sol.

EtOAc, 2-butanone; poorly sol. H₂O,

hexane. λ_{max} 225 (ε 31600); 255 (ε

10400); 298 (ε 4920) (MeOH) (De-

rep).

8'-Methoxy: **Antibiotic F 1839B**, F 1839B

[159096-44-3]

C₂₄H₃₃NO₆ 431.528

From *Stachybotrys* sp. F-1839. Inhi-

bitor of cholesterol esterase, anti-

cholesterolaemic. Amorph. powder.

Sol. MeOH, DMSO, Py; fairly sol.

EtOAc, 2-butanone; poorly sol. H₂O,

hexane. λ_{max} 218 (ε 41300); 271 (ε

6230); 308 (ε 2910) (base in MeOH)

(Derep).

8'-Epimer, 8'-methoxy: **Antibiotic F 1839C**, F 1839C

[159170-51-1]

C₂₄H₃₃NO₆ 431.528

From *Stachybotrys* sp. F-1839. Inhibi-

tor of cholesterol esterase, antichole-

sterolemic. Amorph. powder. Sol.

MeOH, Py, DMSO; fairly sol. EtOAc,

2-butanone; poorly sol. H₂O, hexane.

λ_{max} 217 (ε 33100); 267 (ε 5290); 302 (ε

2530) (MeOH) (Derep). λ_{max} 218 (ε

41300); 271 (ε 6230); 308 (ε 2910)

(MeOH) (Derep). λ_{\max} 217 (ϵ 33069); 267 (ϵ 5293); 302 (ϵ 2525) (MeOH) (Berdy).

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1993, **71**, 487-493 (*Stachybotramide*)
Sakai, K. *et al.*, *J. Antibiot.*, 1995, **48**, 447-456 (*isol*, *pmr*, *cmr*)
Jarvis, B.B. *et al.*, *Nat. Toxins*, 1995, **3**, 10-16 (*Stachybotrylactam*)
Kamalov, L.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 599-607; 1998, **34**, 666-669; 1999, **35**, 103-107 (*Stachybotrin*, *Stachybotrin A*)
Deng, W.P. *et al.*, *J.O.C.*, 2003, **68**, 7422-7427 (*synth*, *struct*, *cryst struct*)
Vázquez, M.J. *et al.*, *Tetrahedron*, 2004, **60**, 2379-2385 (*Stachybotrys chartarum* *metabs*)

Antibiotic F 10C A-1165


F 10C

Pyrrhothine-related antibiotic. Struct. unknown. Prod. by *Streptomyces luteoreticuli*. Antibacterial and antifungal agent. Orange cryst. Sol. EtOAc. λ_{\max} 245; 310; 340 (MeOH).

Fukui Kenkyo Nempo, 1957, **10**, 52

Antibiotic FA 4283 A-1166

2-(3-Heptenyl)-2,6-dihydro-3-methyl-5H-pyrano[3,2-c]pyridin-5-one. FA 4283 [115501-93-4]



$C_{16}H_{21}NO_2$ 259.347

Prod. by *Penicillium funiculosum*. Inhibitor of fatty acid synth. Antiobesity agent. Yellow cryst. (C_6H_6 /hexane). Sol. MeOH, DMSO, DMF, $CHCl_3$; fairly sol. EtOAc, Et_2O ; poorly sol. H_2O . $[\alpha]_D^{+38.3}$ (EtOH). λ_{\max} 235 ($E1\%/1cm$ 810); 330 ($E1\%/1cm$ 350) (MeOH) (Berdy).


► LD_{50} (mus, ipr) 400-1200 mg/kg.

Eur. Pat., 1988, 259 811; *CA*, **109**, 53213x

Antibiotic FD 838 A-1167

FD 838

[110341-78-1]



$C_{22}H_{21}NO_7$ 411.41


Tetramic acid deriv. Related to Antibiotic FD 839, A-1168. Prod. by *Streptomyces fumigatus fresenius*. Antitumour agent. Pale green powder. Sol. MeOH, $CHCl_3$, C_6H_6 ; fairly sol. Et_2O , hexane; poorly sol. H_2O . Mp 98-99°. λ_{\max} 252 (ϵ 7500); 283 (ϵ 4110); 336 (ϵ 10010) (MeOH).

Eur. Pat., 1987, 216 607; *CA*, **107**, 132627x
Orellana, A. *et al.*, *Chem. Comm.*, 2008, (*synth*)

Antibiotic FD 839 A-1168

FD 839

[135862-94-1]



$C_{22}H_{23}NO_8$ 429.426


Prod. by *Aspergillus fumigatus fresenius*. Antitumour agent. Mp 105-107°. λ_{\max} 249 (ϵ 11100); 283 (ϵ 3170) (MeOH).

Japan. Pat., 1991, 91 10 689; *CA*, **115**, 112816v (*isol*)

Antibiotic FL 120B' A-1169

FL 120B'

[156429-12-8]



Absolute Configuration

$C_{20}H_{14}N_2O_7$ 394.34

Prod. by *Streptomyces chattanoogensis* ssp. *taitungensis*. Active against gram-positive bacteria. Orange cryst. ($EtOAc/CHCl_3$). Sol. MeOH, $EtOAc$; poorly sol. H_2O . Mp 155-160° dec. Similar to Kinamycin F. λ_{\max} 207 (ϵ 14000); 247 (ϵ 15300); 275 (ϵ 10300); 295 (sh) (ϵ 6500); 309 (sh) (ϵ 5300); 394 (ϵ 5700); 445 (ϵ 4200) (MeOH). λ_{\max} 218 (ϵ 11600); 240 (ϵ 15400); 274 (ϵ 13900); 309 (ϵ 7600); 392 (ϵ 5300); 545 (ϵ 5000) (MeOH/NaOH) (Berdy).

O-De-Ac, 4-O-(2-methylpropanoyl):

Antibiotic FL 120B. FL 120B

[156429-11-7]

$C_{22}H_{18}N_2O_7$ 422.393

Prod. by *Streptomyces chattanoogensis* ssp. *taitungensis*. Active against gram-positive bacteria. Orange needles ($CHCl_3/EtOAc$). Sol. MeOH, $EtOAc$; poorly sol. H_2O . Mp 185-190° dec. λ_{\max} 206 (ϵ 12500); 243 (ϵ 10600); 274 (ϵ 7600); 294 (sh) (ϵ 4600); 308 (sh) (ϵ 3200); 394 (ϵ 3400); 445 (ϵ 3700) (MeOH). λ_{\max} 218 (ϵ 14300); 238 (ϵ 14800); 275 (ϵ 14100); 308 (ϵ 6400); 389 (ϵ 3900); 539 (ϵ 4000) (MeOH/NaOH) (Berdy).

Chang, L.-R. *et al.*, *J. Antibiot.*, 1994, **47**, 675-680; 681-687 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *props*)

Antibiotic FR 3383 A-1170

FR 3383

[60351-50-0]

Nucleoside-type antibiotic. Struct. unknown. Prod. by *Streptomyces odainensis*. Antibacterial agent. Yellow cryst. (butanol). Sol. H_2O ; poorly sol. MeOH, Me_2CO , $EtOAc$, $CHCl_3$. λ_{\max} 215 ($E1\%/1cm$ 340); 258 ($E1\%/1cm$ 210) (MeOH)

(Berdy). λ_{\max} 215 ($E1\%/1cm$ 340); 258 ($E1\%/1cm$ 210) (H_2O) (Berdy). λ_{\max} 212 ($E1\%/1cm$ 360); 240 ($E1\%/1cm$ 300) (HCl) (Berdy).


► LD_{50} (mus, ipr) 1000 - 2000 mg/kg.

Japan. Pat., 1976, 76 54 988; *CA*, **85**, 121791x
Japan. Pat., 1977, 77 93 701; *CA*, **87**, 199159h

Antibiotic FR 66979 A-1171

FR 66979

[102409-60-9]



$C_{14}H_{17}N_3O_6$ 323.305

Exists as tautomeric mixt. Prod. by *Streptomyces sandaensis*. Shows weak activity against bacteria, fungi and yeasts. Powder. Sol. H_2O , MeOH; poorly sol. Me_2CO , $CHCl_3$. Mp 165-170° dec. $[\alpha]_D^{23} +12.5$. Related to Antibiotic FR 900482, A-1177. λ_{\max} 214; 240 (sh); 280 (H_2O/HCl) (Derep). λ_{\max} 223; 250 (sh); 298 ($H_2O/NaOH$) (Derep). λ_{\max} 215 (ϵ 20000); 240 (sh) (ϵ 5000); 280 (ϵ 1600) (H_2O) (Derep).

► LD_{50} (mus, ivn) 40 mg/kg. KC5010000

Terano, H. *et al.*, *J. Antibiot.*, 1989, **42**, 145


(*isol*, *pmr*, *cmr*, *struct*)

Ducray, R. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 4688-4691 (*synth*)

Judd, T.C. *et al.*, *J.O.C.*, 2004, **69**, 2825-2830 (*synth*)

Antibiotic FR 225656 A-1172

FR 225656



R = CH_2CH_3

$C_{37}H_{44}ClN_7O_8$ 750.249

Prod. by *Helicomyces* sp. No. 19353. Gluconeogenesis inhibitor. Powder. Mp 230° dec. $[\alpha]_D^{23} +37$ (c, 0.5 in DMSO). λ_{\max} 232 (ϵ 63000) (MeOH).

5-Me ether:

$C_{38}H_{46}ClN_7O_8$ 764.276

Prod. by *Helicomyces* sp. No. 19353. Gluconeogenesis inhibitor. Powder. Mp 210-215° dec. $[\alpha]_D^{23} +31$ (c, 0.3 in DMSO). λ_{\max} 232 (ϵ 49000) (MeOH).

Ohtsu, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 682-688; 689-693; 694-699 (*isol*, *ir*, *pmr*, *cmr*, *activity*)

Antibiotic FR 225659 A-1173

FR 225659

As Antibiotic FR 225656, A-1172 with R = CH_3

C₃₆H₄₂CIN₇O₈ 736.223
 Prod. by *Helicomyces* sp. No. 19353.
 Gluconeogenesis inhibitor. Potential anti-diabetic agent. Powder. Mp 225-230° dec. $[\alpha]_D^{23} +40$ (c, 0.5 in DMSO). λ_{\max} 233 (ε 54000) (MeOH).

5-Me ether:

C₃₇H₄₄CIN₇O₈ 750.249
 Prod. by *Helicomyces* sp. No. 19353.
 Gluconeogenesis inhibitor. Powder. Mp 230-235° dec. $[\alpha]_D^{23} +35$ (c, 0.1 in DMSO). λ_{\max} 228 (ε 43000) (MeOH).

3''-Deoxy:


C₃₆H₄₂CIN₇O₇ 720.223
 Prod. by *Helicomyces* sp. No. 19353.
 Gluconeogenesis inhibitor. Powder. Mp 225-230° dec. $[\alpha]_D^{23} +22$ (c, 0.06 in DMSO). λ_{\max} 232 (ε 47000) (MeOH).

Ohtsu, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 682-688; 689-693; 694-699 (isol, ir, pmr, cmr, crystal, activity)

Antibiotic FR 252921

FR 252921

A-1174



C₂₉H₄₄N₂O₅ 500.677
 Prod. by *Pseudomonas fluorescens* No. 408813. Immunosuppressant. Powder. Mp 140-144°. $[\alpha]_D^{23} -222$ (c, 0.2 in DMSO). λ_{\max} 239 (ε 32200); 304 (ε 39700) (MeCN).

Fujine, K. *et al.*, *J. Antibiot.*, 2003, **56**, 55-61; 62-67; 68-71 (isol, activity)

Yu, S. *et al.*, *Tet. Lett.*, 2006, **47**, 9155-9157 (synth)

Falck, J.R. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 4527-4529 (synth, abs config)

Antibiotic FR 252922

FR 252922

A-1175

As Antibiotic FR 252921, A-1174 with R = -(CH₂)₆CH₃

C₃₁H₄₈N₂O₅ 528.731
 Prod. by *Pseudomonas fluorescens* No. 408813. Immunosuppressant. Powder. Mp 142-146°. $[\alpha]_D^{23} -235$ (c, 0.2 in DMSO). λ_{\max} 239 (ε 29300); 304 (ε 36700) (MeCN).

5',6'-Didehydro: Antibiotic FR 256523.

FR 256523
 C₃₁H₄₆N₂O₅ 526.715
 Prod. by *Pseudomonas fluorescens* No. 408813. Immunosuppressant. Powder. Mp 88-93°. $[\alpha]_D^{23} -72$ (c, 0.3 in DMSO). λ_{\max} 266 (sh); 278 (ε 60700); 291 (ε 59400) (MeCN).


Fujine, K. *et al.*, *J. Antibiot.*, 2003, **56**, 55-61; 62-67; 68-71 (isol, activity)

Antibiotic FR 900148

FR 900148

A-1176

[74242-74-3]
 [73706-57-7]



C₁₀H₁₅CIN₂O₅ 278.692

Peptide antibiotic. Struct. revised in 1991. Isol. from *Streptomyces xanthocidicus*. Active against gram-positive and -negative bacteria. Inhibits cell wall synth. by spheroplast formn. Amphoteric powder + 2H₂O (as Na salt). Sol. H₂O, MeOH; fairly sol. EtOH; poorly sol. Me₂CO, HCl. Mp 143-147° dec. (Na salt). $[\alpha]_D^{20} +77.1$ (c, 0.75 in H₂O). pK_{a1} 3.25; pK_{a2} 7.9.

▶ LD₅₀ (mus, ivn) 3500-6500 mg/kg.


Kuroda, Y. *et al.*, *J. Antibiot.*, 1980, **33**, 259; 267 (isol)

Yasuda, N. *et al.*, *J. Antibiot.*, 1991, **44**, 801-802 (struct)

Antibiotic FR 900482

A-1177

FR 900482
 [102363-08-6]



C₁₄H₁₅N₃O₆ 321.289

Exhibits epimerisation at anomeric centre. Prod. by *Streptomyces sandaensis* isol. from soil. Active against gram-positive and -negative bacteria and tumours. Powder. Sol. MeOH, H₂O; poorly sol. Me₂CO, CHCl₃. Mp 175° dec. $[\alpha]_D^{23} +8$ (c, 1 in H₂O). Synthetic derivs. showed antileukaemic activity. λ_{\max} 214 (ε 31000); 240 (sh) (ε 7400); 282 (ε 2500); 331 (ε 600) (MeOH/HCl) (Derep). λ_{\max} 238 (ε 15700); 302 (ε 2600); 374 (ε 2200) (MeOH/NaOH) (Derep). λ_{\max} 236 (ε 19200); 281 (ε 6100); 330 (ε 2200) (MeOH) (Derep). λ_{\max} 218 (E1%/1cm 630); 236 (E1%/1cm 600); 281 (E1%/1cm 190); 330 (E1%/1cm 70) (MeOH) (Berdy). λ_{\max} 216; 282; 331 (MeOH/HCl) (Berdy). ▶ LD₅₀ (mus, ipr) 32 mg/kg. KC5000000

1,7,9-Tri-Ac: FK973

[114580-45-9]

C₂₀H₂₁N₃O₉ 447.401

Isol. from *Streptomyces sandaensis*. Antitumour antibiotic.

7-Me ether, 1,9-di-Ac: FK317

[102409-92-7]

C₁₉H₂₁N₃O₈ 419.39

Antineoplastic antibiotic. Used in the treatment of solid tumours. Metab. of FK973.

Eur. Pat., 1986, ((Fujisawa))166 389; *CA*, **104**, 223624y (FK 317)

Kiyoto, S. *et al.*, *J. Antibiot.*, 1987, **40**, 589-593; 594-599; 600-606; 607-611 (isol, struct, props)

Uchida, I. *et al.*, *J.A.C.S.*, 1987, **109**, 4108-4109 (struct)

Fujita, T. *et al.*, *J. Antibiot.*, 1988, **41**, 392-394 (biosynth)

Masuda, K. *et al.*, *Jpn. J. Pharmacol.*, 1989, **51**, 219-226 (FK 973)

Yasuda, N. *et al.*, *Tet. Lett.*, 1989, **30**, 3397-3400 (synth)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1992, **114**, 383-385 (synth)

Schkeryantz, J.M. *et al.*, *J.A.C.S.*, 1995, **117**, 4722-4723 (synth)

Katoh, T. *et al.*, *Tetrahedron*, 1997, **53**, 10229-10238; 10239-10252; 10253-10270 (synth)

Naoue, Y. *et al.*, *Cancer Chemother. Pharmacol.*, 1998, **42**, 31-36 (FK 317)

Hoshi, A. *et al.*, *Drugs of the Future*, 1998, **23**, 483-484 (FK 317)

Inami, M. *et al.*, *Cancer Lett. (Shannon, Irel.)*, 2002, **181**, 39-45 (FK 317, pharmacol)

Paleo, M.R. *et al.*, *J.O.C.*, 2003, **68**, 130-138 (synth)

Judd, T.C. *et al.*, *J.O.C.*, 2004, **69**, 2825-2830 (synth)


Suzuki, M. *et al.*, *J.O.C.*, 2004, **69**, 2831-2843 (synth)

Trost, B.M. *et al.*, *Org. Lett.*, 2008, **10**, 1369-1372 (epimer, synth)

Antibiotic FR 901483

A-1178

FR 901483. WF 11231A. Antibiotic WF 11231A
 [150151-86-3]



C₂₀H₃₁N₂O₆P 426.449

Prod. by *Cladobotryum* sp. No. 11231. Immunosuppressant. Lymphocyte proliferation inhibitor. Purine nucleotide biosynth. inhibitor. Mycotoxin. Needles. Sol. MeOH, H₂O; poorly sol. Me₂CO, EtOAc. Mp 210-213°. $[\alpha]_D^{25} -11$ (c, 0.7 in CHCl₃). λ_{\max} 225; 275; 285 (sh) (MeOH).

[150284-39-2]

Sakamoto, K. *et al.*, *J. Antibiot.*, 1996, **49**, 37-44 (isol, uv, ir, pmr, cmr, props)

Ousmer, M. *et al.*, *J.A.C.S.*, 2001, **123**, 7534-7538 (synth)

Ousmer, M. *et al.*, *Org. Lett.*, 2001, **3**, 765-769 (synth)

Maeng, J.-H. *et al.*, *Org. Lett.*, 2001, **3**, 1125-1128 (synth)


Kan, T. *et al.*, *Org. Lett.*, 2004, **6**, 2729-2731 (synth)

Brummond, K.M. *et al.*, *J.O.C.*, 2005, **70**, 907-916 (synth)

Antibiotic G 1499-2

A-1179

3-Methyl-2-(pentylidenecyclopropyl)-4(1H)-quinolinone, 9CI. G 1499-2
 [68978-12-1]



C₁₈H₂₁NO 267.37

Prod. by *Cytophaga johnsonii*. Possesses limited antibiotic activity against a few bacteria. Rosettes (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 192-


193°. $[\alpha]_D^{25} +172$ (c, 0.5 in CHCl_3). λ_{\max} 236 (ϵ 49100); 310 (ϵ 10900) (MeOH/HCl) (Derep). λ_{\max} 244 (ϵ 33600); 322 (ϵ 11200); 336 (ϵ 11900) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 500 - 1500 mg/kg.
VC8298500

Evans, J.R. *et al.*, *J. Antibiot.*, 1978, **31**, 952
(isol, uv, ir, pmr, cmr, ms)

Antibiotic GE 2270
GE 2270

A-1180



- A R¹ = CH₂OMe, R² = CH₃
B₂ R¹ = R² = CH₃
C₁ R¹ = H, R² = CH₃
C_{2a} R¹ = CH₂OMe, R² = CH₂OH
C_{2b} R¹ = CH₂OMe, R² = H
D₂ R¹ = CH₂OH, R² = CH₃

Peptide antibiotic complex. Structs. revised in 1994. Prod. by *Planobispora rosea* ATCC 53773. Active against gram-positive bacteria and anaerobes. Sol. MeOH, EtOAc; poorly sol. hexane.

Antibiotic GE 2270A

GE 2270A. MDL 62879. Antibiotic MDL 62879

C₅₆H₅₅N₁₅O₁₀S₆ 1290.541
Amorph. powder. $[\alpha]_D +140.8$ (EtOH). λ_{\max} 240 (sh) (ϵ 54100); 309 (ϵ 34400); 340 (sh) (ϵ 13800) (MeOH/HCl) (Derep). λ_{\max} 240 (sh) (ϵ 58800); 310 (ϵ 34200); 340 (sh) (ϵ 15500) (MeOH/NaOH) (Derep). λ_{\max} 240 (sh) (ϵ 58800); 310 (ϵ 34200); 340 (sh) (ϵ 15500) (MeOH) (Derep).

N-De-Me: Antibiotic GE 2270B₁. GE 2270B₁

[139351-31-8]
C₅₅H₅₃N₁₅O₁₀S₆ 1276.514
Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 310 (MeOH) (Berdy).

8,9-Didehydro: Antibiotic GE 2270T. GE 2270T

[138967-24-5]
C₅₆H₅₃N₁₅O₁₀S₆ 1288.525
Powder. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 250; 280; 345 (MeOH) (Berdy). λ_{\max} 311 (MeOH/NaOH) (Berdy). λ_{\max} 250 (HCl) (Berdy).

Antibiotic GE 2270B₂
GE 2270B₂

[139351-32-9]

C₅₅H₅₃N₁₅O₉S₆ 1260.515
Sol. MeOH, EtOAc; poorly sol. H₂O, hexane.

Antibiotic GE 2270C₁

GE 2270C₁

[139351-33-0]
C₅₄H₅₁N₁₅O₉S₆ 1246.488
Sol. MeOH, EtOAc; poorly sol. H₂O, hexane.

N-De-Me: Antibiotic GE 2270D₁. GE 2270D₁

[138948-59-1]
C₅₃H₄₉N₁₅O₉S₆ 1232.461
Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 310 (MeOH) (Berdy).

Antibiotic GE 2270C_{2a}

GE 2270C_{2a}

[145118-96-3]
C₅₆H₅₅N₁₅O₁₁S₆ 1306.54

Antibiotic GE 2270C_{2b}

GE 2270C_{2b}

[139351-34-1]
C₅₅H₅₃N₁₅O₁₀S₆ 1276.514

Antibiotic GE 2270D₂

GE 2270D₂

[138948-60-4]
C₅₅H₅₃N₁₅O₁₀S₆ 1276.514
Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 310 (MeOH) (Berdy).


N-De-Me: Antibiotic GE 2270E. GE 2270E

C₅₄H₅₁N₁₅O₁₀S₆ 1262.487
Selva, E. *et al.*, *J. Antibiot.*, 1991, **44**, 693-701; 702-715; 1995, **48**, 1039-1042 (isol, pmr, cmr)
Colombo, L. *et al.*, *Org. Mass Spectrom.*, 1992, **27**, 219-225 (ms)
Tavecchia, P. *et al.*, *J. Antibiot.*, 1994, **47**, 1564-1567 (ms, struct)
Colombo, L. *et al.*, *Rapid Commun. Mass Spectrom.*, 1995, **9**, 717-722 (ms)
De Pietro, M.T. *et al.*, *J. Antibiot.*, 2001, **54**, 1066-1071 (biosynth)
Müller, H.M. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 4771-4774 (synth)
Nicolaou, K.C. *et al.*, *Chem. Asian J.*, 2008, **3**, 413-429 (GE 2270A, GE 2270T, synth)

Antibiotic GE 37468A

A-1181

GE 37468A
[160262-65-7]



C₅₉H₅₂N₁₄O₁₂S₅ 1309.476
Prod. by *Streptomyces* sp. ATCC 55365. Bacterial protein synthesis inhibitor. Active against gram-positive bacteria. Solid.

Mp 180-189°. $[\alpha]_D^{20} +105$ (c, 0.04 in MeOH/DMSO). λ_{\max} 250 (sh) (ϵ 43000); 305 (ϵ 34400); 340 (sh) (ϵ 18300) (MeOH/MeCN aq.).

3-N-De(carboxyalkyl): Antibiotic GE

37468B. GE 37468B

[160262-64-6]
C₅₆H₅₀N₁₄O₁₀S₅ 1239.429
Prod. by *Streptomyces* sp. ATCC 55365. Yellow powder. λ_{\max} 302; 317; 334 (NaOH) (Berdy).

6-N-De(alkylaminoalkyl): Antibiotic GE

37468C. GE 37468C

[160262-66-8]
C₅₃H₄₇N₁₃O₉S₅ 1170.366
Prod. by *Streptomyces* sp. ATCC 55365. Yellow powder. λ_{\max} 302; 317; 334 (NaOH) (Berdy).


Pat. Coop. Treaty (WIPO), 1994, 94 14 838;

CA, **122**, 128491r (derivs)
Stella, S. *et al.*, *J. Antibiot.*, 1995, **48**, 780-786; 1304-1311 (isol, uv, pmr, ms, props)

Antibiotic GKK 1032A₂

A-1182

GKK 1032A₂
[358375-09-4]



Relative Configuration

C₃₂H₄₁NO₄ 503.68

Similar to Pyrrocidine B, P-940. Prod. by *Penicillium* sp. GKK1032. Antitumour agent. Mp 147-148°. $[\alpha]_D^{19} +104$ (c, 0.2 in MeOH). λ_{\max} 203 (ϵ 17400); 226 (sh) (ϵ 6200) (MeOH).

2'-Me ether: Antibiotic GKK 1032A₁.

GKK 1032A₁
[358375-08-3]
C₃₃H₄₃NO₄ 517.707
Prod. by *Penicillium* sp. GKK1032. Antitumour agent. Mp 238-239°. $[\alpha]_D^{19} +87$ (c, 0.4 in MeOH). λ_{\max} 209 (ϵ 19200); 225 (sh) (ϵ 12400) (MeOH).


Japan. Pat., 2001, 01 247 574; CA, **135**, 209979b (isol)

Oikawa, H. *et al.*, *J.O.C.*, 2003, **68**, 3552-3557 (pmr, cmr, biosynth)

Antibiotic GKK 1032B

A-1183

GKK 1032B
[358375-11-8]



Relative Configuration

C₃₂H₃₉NO₄ 501.664

Similar to Pyrrocidine B, P-940. Prod. by *Penicillium* sp. GKK1032. Antitumour agent. Mp 159-161°. [α]_D²⁰ +172 (c, 0.2 in MeOH). λ_{\max} 202 (ε 23000); 230 (sh) (ε 4600) (MeOH).


Japan. Pat., 2001, 01 247 574; *CA*, **135**, 209979b

Oikawa, H. *et al.*, *J.O.C.*, 2003, **68**, 3552-3557 (biosynth)

Antibiotic HA 23

A-1184

HA 23

C₃₄H₅₂N₂O₇ 600.794

Prod. by *Fusarium* sp. CANU-HA 23. Oil. [α]_D²⁰ -46 (c, 0.001 in MeOH). λ_{\max} 232 (log ε 3.23); 278 (log ε 2.8) (MeOH).

Feng, Y. *et al.*, *Org. Lett.*, 2002, **4**, 2095-2096 (isol, pmr, cmr)

Antibiotic IC 202A

A-1185

IC 202A

[221456-27-5]

H₂N(CH₂)₅N(OH)COCH₂CH₂CONH-(CH₂)₅N(OH)COCH₂CH₂CONH-(CH₂)₅N(O)=CHCH₂CH₂CH₃

C₂₇H₅₂N₆O₇ 572.744


Closely related to Deferoxamine, D-142. Prod. by *Streptoalloteichus* sp. 1454-19. Siderophore. Immunosuppressant. Powder.

Iijima, M. *et al.*, *J. Antibiot.*, 1999, **52**, 20-24; 25-28

Antibiotic IC 202B

A-1186

IC 202B

C₂₃H₄₄N₆O₈ 532.636

Prod. by *Streptoalloteichus* sp. 1454-19. Siderophore. Immunosuppressant. Pale

yellow powder. Mp 117-119°.

Iijima, M. *et al.*, *J. Antibiot.*, 1999, **52**, 775-780

Antibiotic IC 202C

A-1187

IC 202C

As Antibiotic IC 202B, A-1186 with R = -CH=NOH

C₂₃H₄₄N₆O₇ 516.637


Prod. by *Streptoalloteichus* sp. 1454-19. Siderophore. Immunosuppressant. Powder. Mp 140-141°. Isol. as a mixt. of *E*- and *Z*- isomers.

Iijima, M. *et al.*, *J. Antibiot.*, 1999, **52**, 775-780; 2000, **53**, 1411-1415

Antibiotic ICM 0201

A-1188

ICM 0201

C₁₂H₁₃NO₆ 267.238

Prod. by *Cunninghamella* sp. F-1490. Inhibitor of osteoclastogenesis. Pale yellow powder. Mp 125-131° dec. [α]_D²² -57.5 (c, 0.4 in MeOH).


Inoue, H. *et al.*, *J. Antibiot.*, 2003, **56**, 209-213; 214-218 (isol, synth, pmr, cmr)

Antibiotic K 252b

A-1189

K 252b. SF 2370. Antibiotic SF 2370

[99570-78-2]



Absolute Configuration

C₂₆H₁₉N₃O₅ 453.453

No stereochem. indicated for SF 2370 but it appears to be identical with K 252b. Isol. from *Nocardiopsis* sp. and *Actinomadura* sp. Protein kinase C inhibitor. Histamine inhibitor. Neurotrophin-3 potentiator. Calmodulin inhibitor. Shows activity against bacteria and fungi. Protects against rice blast disease. Pale yellow powder or cryst. Sol. Me₂CO, MeCN, dioxan, DMSO, THF, Py, CHCl₃; fairly sol. MeOH, EtOAc, EtOH, butanol; poorly sol. H₂O, Et₂O, acids. Mp 262-266° dec. [α]_D²⁰ +97 (c, 0.6 in DMF). [α]_D²⁵ +57 (c, 0.1 in MeOH). λ_{\max} 227 (ε 42000); 249 (ε 41000); 264 (sh) (ε 44000); 280 (sh) (ε 68000); 290 (ε 102000); 320 (sh) (ε 19000); 334 (ε 26000); 350 (ε 19000); 367 (ε 21000) (MeOH) (Derep). λ_{\max} 371 (ε 13000)

(MeOH) (Berdy).

Me ester: Antibiotic K 252a. K 252a [99533-80-9]

C₂₇H₂₁N₃O₅ 467.48

Isol. from *Nocardiopsis* sp. and *Streptomyces longisporoflavus* R19. Protein kinase C and phosphorylase kinase inhibitor. Calmodulin inhibitor. Anti-allergic agent, vasorelaxant. Pale yellow cryst. (Me₂CO/MeOH). Sol. Me₂CO, dioxan, Py, THF, DMSO, MeCN, CHCl₃; fairly sol. MeOH, butanol, EtOAc, EtOH; poorly sol. H₂O, acids, Et₂O. Mp 262-273° dec. [α]_D²⁰ -23 (c, 0.5 in CHCl₃). [α]_D²⁰ +52 (c, 0.1 in MeOH). λ_{\max} 227 (ε 42000); 249 (ε 41000); 264 (sh) (ε 44000); 280 (sh) (ε 68000); 290 (ε 102000); 320 (sh) (ε 19000); 334 (ε 26000); 350 (ε 19000); 367 (ε 21000) (MeOH) (Derep).

Hexyl ester: KT 5720

[108068-98-0]

[161510-88-9]

C₃₂H₃₁N₃O₅ 537.614

Protein kinase inhibitor. Reverses multidrug resistance. Mp 145-147°.

3'-Deoxy, 3'-(methylamino), Me ester: [183145-61-1]

C₂₈H₂₄N₄O₄ 480.522

Prod. by *Streptomyces longisporoflavus*. Cryst. (CH₂Cl₂). Mp 152-155°.

Koyama, M. *et al.*, *J. Antibiot.*, 1985, **38**, 1437-1439 (SF 2370)

Kase, H. *et al.*, *J. Antibiot.*, 1986, **39**, 1059-1065 (isol, props)

Nakanishi, S. *et al.*, *J. Antibiot.*, 1986, **39**, 1066-1071 (isol, props)

Yasuzawa, T. *et al.*, *J. Antibiot.*, 1986, **39**, 1072-1078 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Eur. Pat., 1994, ((Kyowa Hakko Kogyo))630 898; *CA*, **122**, 10020h (KT 5720, synth, pharmacol)

Galski, H. *et al.*, *Eur. J. Cancer, Part A*, 1995, **31**, 380-388 (KT 5720, pharmacol)

Lowinger, T.B. *et al.*, *Tet. Lett.*, 1995, **36**, 8383-8386 (synth)

Cai, Y. *et al.*, *J. Antibiot.*, 1996, **49**, 1060-1062 (*3'-deoxy-3'-methylamino Me ester*)

Wood, J.L. *et al.*, *J.A.C.S.*, 1996, **118**, 10656-10657 (synth)

Kobayashi, Y. *et al.*, *J.A.C.S.*, 1999, **121**, 6501-6502 (synth)

Antibiotic K 252c


A-1190

6,7,12,13-Tetrahydro-5H-indolo[2,3-

a]pyrrolo[3,4-c]carbazol-5-one, 9CI.

Staurosporine aglycone. Staurosporinone. K 252c

[85753-43-1]

C₂₀H₁₃N₃O 311.342

Isol. from *Nocardiopsis* sp. K290e and from the marine ascidian *Eudistoma* sp., also from *Streptomyces longisporoflavus*

R19, *Streptomyces mediocidicus* ATCC 13279 and the fruiting bodies of the myxomycete *Lycogala epidendrum*. Protein kinase C inhibitor, calmodulin inhibitor, serotonin release inhibitor and cell adhesion inhibitor. Pale yellow needles. Sol. DMSO; fairly sol. MeOH, CHCl₃; poorly sol. H₂O, acids, bases. Mp > 300°. λ_{max} 230 (ε 37000); 238 (sh) (ε 34000); 246 (sh) (ε 28000); 257 (sh) (ε 29000); 287 (ε 86000); 320 (sh) (ε 16000); 331 (ε 20000); 341 (ε 16000); 358 (ε 11000) (MeOH) (Derep).

N¹³-(α-L-Rhamnopyranosyl): **Antibiotic K 252d**. K 252d [105114-22-5] C₂₆H₂₃N₃O₅ 457.485
Isol. from *Nocardiosis* sp. K290e. Protein kinase C inhibitor, calmodulin inhibitor and serotonin release inhibitor. Pale yellow needles. Sol. MeOH, CHCl₃, DMSO; poorly sol. H₂O, bases, acids. Mp 240-245° dec. [α]_D²⁰ +35 (c, 0.4 in MeOH). λ_{max} 223 (ε 40000); 233 (ε 39700); 248 (sh) (ε 31000); 260 (sh) (ε 34000); 268 (sh) (ε 38000); 280 (sh) (ε 61000); 290 (ε 98800); 320 (sh) (ε 17000); 335 (ε 20300); 347 (ε 13200); 364 (ε 11500) (MeOH) (Derep).

N¹³-(2,6-Dideoxy-α-D-ribo-hexopyranosyl): **Antibiotic RK 286D**. RK 286D [140429-37-4] C₂₆H₂₃N₃O₄ 441.485
Prod. by *Streptomyces* sp. RK-286/P10634. Inhibitor of protein kinase C, also antiinflammatory, platelet agglutination inhibitor, bleb formation inhibitor and antioncotic promoter. Pale yellow powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 225-230°. [α]_D²⁰ -60 (c, 0.13 in MeOH). λ_{max} 223 (ε 40000); 233 (ε 39700); 248 (sh) (ε 31000); 260 (sh) (ε 34000); 268 (sh) (ε 38000); 280 (sh) (ε 61000); 290 (ε 98800); 320 (sh) (ε 17000); 335 (ε 20300); 347 (ε 13200); 364 (ε 11500) (MeOH) (Derep).

N¹³-(3-Amino-2,3,6-trideoxy-α-L-ribo-hexopyranosyl): **Holyrine A** [249512-77-4] C₂₆H₂₄N₄O₃ 440.501
Prod. by a marine actinomycete.
N¹³-(3-Amino-2,3,6-trideoxy-5-hydroxy-α-L-erythro-hexopyranosyl): **Holyrine B** [249512-78-5] C₂₆H₂₄N₄O₄ 456.5
Prod. by a marine actinomycete.

N⁶-(1-Methylethoxy)methyl: [178276-03-4] C₂₄H₂₁N₃O₂ 383.449
Prod. by *Streptomyces longisporoflavus*. Yellowish solid. Mp 45-50°.

7-Oxo: see Arcyriaflavin A, A-1393

2-Hydroxy: **2-Hydroxystaurosporinone** C₂₀H₁₃N₃O₂ 327.342
Isol. from *Lycogala epidendrum*. Protein tyrosine kinase inhibitor. Cytotoxic. Light brown solid. λ_{max} 226 (log ε 10.2); 294 (log ε 10.9); 342 (log ε 9.8) (MeOH).

7-Methoxy: [137888-73-4] C₂₁H₁₅N₃O₂ 341.368
Prod. by *Saccharothrix aerocolonigenes* ssp. *copiosa* ATCC 53856. Yellow powder. λ_{max} 206 (ε 24000); 234 (ε 23000); 296 (ε 45600); 332 (ε 8000); 344 (ε 6500); 361 (ε 4700) (MeOH) (Berdy).

Sarstedt, B. et al., *Heterocycles*, 1983, **20**, 469-476 (synth)
Nakanishi, S. et al., *J. Antibiot.*, 1986, **39**, 1066-1071 (isol, props)
Yasuzawa, T. et al., *J. Antibiot.*, 1986, **39**, 1072-1078 (isol, uv, ir, pmr, cmr, ms, struct)
Hughes, I. et al., *J.C.S. Perkin 1*, 1990, 2475-2480 (synth, uv, pmr)
Pat. Coop. Treaty (WIPO), 1991, 91 9 034; CA, **116**, 39768v (7-methoxy)
Osada, H. et al., *J. Antibiot.*, 1992, **45**, 278-279 (Antibiotic RK286D)
Moody, C.J. et al., *J.O.C.*, 1992, **57**, 2105-2114 (synth)
Horton, P.A. et al., *Experientia*, 1994, **50**, 843-845 (isol)
Xie, G. et al., *Tet. Lett.*, 1994, **35**, 5555-5558 (synth)
Cai, Y. et al., *J. Antibiot.*, 1996, **49**, 519-526 (N⁶-(1-methylethoxy)methyl, ir, pmr, cmr, cd)
Mahboobi, S. et al., *J.O.C.*, 1999, **64**, 4697-4704 (synth)
Williams, D.E. et al., *Tet. Lett.*, 1999, **40**, 7171-7174 (Holyrines)
Hosoya, T. et al., *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2776-2780 (2-Hydroxystaurosporinone)


Antibiotic KG 431A A-1191
KG 431A [149146-28-1]
Indole antibiotic. Struct. unknown. Prod. by *Clostridium* sp. KH-431. Antibacterial and antifungal agent. Powder. Sol. MeOH, Me₂CO, EtOH; poorly sol. H₂O, C₆H₆, hexane, CHCl₃.
Hong, S.H. et al., *CA*, 1993, **109**, 90899c; 113035j

Antibiotic L 11-1 A-1192
L 11-1 [105054-65-7] C₃₇H₅₃N₃O₉ 683.84
Possibly identical with Tredimycin B in O-155. Prod. by *Streptomyces* sp. L11-1. Active against *Streptococcus* sp. Powder. Sol. MeOH, CHCl₃; poorly sol. cyclohexane. [α]_D²⁰ +43.4 (MeOH). λ_{max} 233 (E1%/1cm 496); 247 (E1%/1cm 510); 267 (E1%/1cm 419); 297 (E1%/1cm 389) (MeOH) (Berdy).
Japan. Pat., 1986, 86 111 696; *CA*, **105**, 189459g

Antibiotic L 13365 A-1193
L 13365 [66105-52-0]
Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Actinoplanes sarveparensis*. Antibacterial agent. Light yellow cryst. (MeOH/EtOAc). Sol. DMSO, CHCl₃-MeOH, DMF; fairly sol. CHCl₃-MeOH, AcOH; poorly sol. EtOH, hexane, H₂O. Mp 210°. [α]_D²⁰ +125 (MeOH/CHCl₃). λ_{max} 237; 290 (sh); 378 (H₂O). λ_{max} 237 (E1%/1cm 326); 378 (E1%/1cm 67) (pH 7.5 buffer) (Berdy). λ_{max} 238 (E1%/1cm 336); 408 (E1%/1cm 99) (pH 9


buffer) (Berdy). λ_{max} 237 (E1%/1cm 326); 360 (E1%/1cm 103) (HCl) (Berdy).
▶ LD₅₀ (mus, ipr) 1000 - 2000 mg/kg.
Ger. Pat., 1978, 2 724 090; *CA*, **88**, 168482p

Antibiotic L 654040 A-1194
L 654040 [127964-76-5]




C₂₀H₂₆N₄O₇ 434.448
Struct. in abstract is misleading. Prod. by *Streptoverticillium syroense*. Sol. MeOH, H₂O. λ_{max} 255 (E1%/1cm 338); 318 (E1%/1cm 79) (MeOH) (Berdy). λ_{max} 255 (E1%/1cm 303); 315 (E1%/1cm 680) (H₂O) (Berdy). λ_{max} 267 (E1%/1cm 390); 340 (E1%/1cm 65) (HCl) (Berdy).
Eur. Pat., 1989, 332 248; *CA*, **113**, 38891 (isol)

Antibiotic L 681176 A-1195
L 681176. Antibiotic 176 [91386-17-3]



C₁₂H₂₃N₅O₇ 349.343
Amino acid antibiotic. Isol. from *Streptomyces* sp. MA5143a. Inhibits angiotensin converting enzyme and the pressor response to angiotensin I in rat. Needles (H₂O). Sol. H₂O, MeOH.
Huang, L. et al., *J. Antibiot.*, 1984, **37**, 462 (isol, props)
Hensens, O.D. et al., *J. Antibiot.*, 1984, **37**, 466 (pmr, ms, struct)

Antibiotic L 697318 A-1196
L 697318 [142494-53-9]



C₃₀H₃₉NO₄ 477.642
Metab. of *Hypoxylon fragiforme*. Amorph. Sol. DMSO, CH₂Cl₂, MeOH; poorly sol. H₂O, hexane.

Deoxy: **Antibiotic L 696475**. L 696475 [142494-54-0] C₃₀H₃₉NO₃ 461.643
Metab. of *Hypoxylon fragiforme*. Amorph. Sol. DMSO, MeOH, CH₂Cl₂; poorly sol. H₂O, hexane.
18β-Hydroxy: **Cytochalasin U**†

[145040-61-5]
C₃₀H₃₉NO₅ 493.642
Isol. from a *Pestalotia* sp. Immuno-suppressant. λ_{\max} 286 (ϵ 330) (MeOH) (Derep).

Deoxy, 18 β -hydroxy: **Antibiotic RKS**

1778. RKS 1778

[191475-89-5]

C₃₀H₃₉NO₄ 477.642

Metab. of *Phoma* sp. SNF-1778.

Mammalian cell cycle inhibitor. Powder. Mp 117-120°. $[\alpha]_D$ -20 (c, 0.05 in MeOH).

Ondeyka, J. et al., *J. Antibiot.*, 1992, **45**, 679-685; 686-691 (isol, pmr, cmr, struct)


Burres, N.S. et al., *J. Antibiot.*, 1992, **45**, 1367-1369 (Cytochalasin U)

Kakeya, H. et al., *J. Nat. Prod.*, 1997, **60**, 669-672 (RK 1778)

Antibiotic L 755807

A-1197

4-Hydroxy-4-(1-methylethyl)-1-(8,10,12-trimethyl-1-oxo-2,4,6,8-tetradecatetraenyl)-6-oxa-3-azabicyclo[3.1.0]hexan-2-one, 9CI. L 755807
[173293-97-5]



C₂₄H₃₅NO₄ 401.545

Tetramic acid deriv. Isol. from an endophytic *Microsphaeropsis* sp. Bradykinin binding inhibitor. Yellow gum. Sol. MeOH, CHCl₃. $[\alpha]_D^{21}$ -87.3 (c, 0.65 in MeOH). λ_{\max} 256 (log ϵ 3.86); 361 (log ϵ 4.44); 381 (sh) (log ϵ 4.33) (MeOH). λ_{\max} 256 (ϵ 7245); 361 (ϵ 27542); 381 (MeOH) (Berdy).


Stahl, M. et al., *J.O.C.*, 1996, **61**, 8083-8088 (config)

Lam, Y.K.T. et al., *Tetrahedron*, 1996, **52**, 1481-1486 (isol, struct)

Antibiotic L 970843

A-1198

L 970843



Relative Configuration

C₂₃H₂₈N₂O₄ 396.485

Related to Xanthocillin X, X-7. Prod. by fungal species HIL Y-903146. Antifungal agent. Pale yellow solid. Mp 118-119°. $[\alpha]_D^{20}$ -20 (c, 0.02 in MeOH). λ_{\max} 224; 288; 300 (MeOH).

Stereoisomer: **Antibiotic L 970844**. L 970844

C₂₃H₂₈N₂O₄ 396.485


Prod. by fungal species HIL Y-903146. Antifungal agent. Pale yellow shiny flakes. Mp 116-117°. $[\alpha]_D$ +20 (c, 0.02 in MeOH).

Vijayakumar, E.K.S. et al., *J. Antibiot.*, 2001, **54**, 973-976 (isol, pmr, cmr)

Antibiotic LL-BH872 α

A-1199

[3-(1-Hexenyl)-ONN-azoxy]-1-hydroxy-2-butanone, 9CI. LL-BH872 α
[24397-77-1]



Absolute Configuration

C₁₀H₁₈N₂O₃ 214.264

Prod. by *Streptomyces hinnulinus*. Potent antifungal agent. Light yellow oil. $[\alpha]_D^{25}$ +157 (c, 1.7 in MeOH). Dec. v. readily. λ_{\max} 238 (ϵ 9000) (MeOH) (Berdy).

Ac: [36846-66-9]

Semisynthetic. Antifungal agent. More stable than parent compd.

2 ξ -Alcohol: [3-(1-Hexenyl)-ONN-azoxy]-1,2-butanediol

[131205-70-4]

[637035-36-0]

C₁₀H₂₀N₂O₃ 216.28

Prod. by an *Actinomadura* sp. isol. from the roots of *Prunus armeniaca* (apricot). Also prod. by *Streptomyces misionensis* FERM 10459.

Antihepatotoxic agent. Yellow oil. $[\alpha]_D^{25}$ -30 (c, 0.2 in MeOH). Stereochemical information incomplete. The two isolates may be stereoisomeric. λ_{\max} 228 (ϵ 9900) (EtOH).

4' ξ -Hydroxy, 2 ξ -alcohol: 3-[4-Hydroxy-1-hexenyl]-ONN-azoxy]-1,2-butanediol. **Antibiotic MH 072**. MH 072

[131205-71-5]

C₁₀H₂₀N₂O₄ 232.279

Isol. from *Streptomyces misionensis*. Antihepatotoxic.

McGahren, W.J. et al., *J.A.C.S.*, 1969, **91**, 2808-2810; 1970, **92**, 1587-1590 (isol, cd)

McGahren, W.J. et al., *J.O.C.*, 1972, **37**, 902-906 (cd, struct)

U.S. Pat., 1972, 3 647 776; CA, 77, 3778 (isol, synth)

Japan. Pat., 1990, 90 202 867; CA, 114, 23409b (*Streptomyces misionensis* isolates)

Bianchi, G. et al., *Planta Med.*, 2003, **69**, 574-576 (2-alcohol, isol, pmr, cmr)

Antibiotic LL-BO 2964

A-1200

LL-BO 2964

[96119-50-5]

Nucleoside-type antibiotic complex.

Struct. unknown. Prod. by *Streptomyces coeruleorubidus* ssp. *rubidus*. Active against gram-positive and -negative bacteria and mycobacteria.

Antibiotic LL-BO 2694 α

LL-BO 2694 α

[96119-51-6] Cell wall biosynth. inhibitor. Powder. Sol. MeOH, EtOH,

H₂O; poorly sol. hexane, CHCl₃, toluene. $[\alpha]_D$ -10 (H₂O). λ_{\max} 221; 254 (H₂O). λ_{\max} 240 (NaOH aq.) (Berdy). λ_{\max} 221; 254 (HCl) (Berdy).

Antibiotic LL-BO 2964 β

LL-BO 2964 β

[96119-52-7] Cell wall biosynthesis inhibitor. Powder. Sol. MeOH, EtOH, H₂O; poorly sol. CHCl₃, toluene, hexane. $[\alpha]_D$ -19 (H₂O). λ_{\max} 220; 256

(HCl) (Berdy). λ_{\max} 238 (NaOH aq.) (Berdy). λ_{\max} 220; 256 (H₂O) (Berdy).

Antibiotic LL-BO 2964 γ

LL-BO 2964 γ

[96119-53-8] Cell Wall Biosynthesis Inhibitor. Powder. Sol. MeOH, EtOH, H₂O; poorly sol. hexane, CHCl₃, toluene. $[\alpha]_D$ -20 (H₂O). λ_{\max} 225; 259 (H₂O). λ_{\max} 245 (NaOH aq.) (Berdy). λ_{\max} 220; 262 (HCl) (Berdy).


U.S. Pat., 1985, 4 499 075; CA, 102, 183743m

Antibiotic LL-49F233 α

A-1201

LL-49F233 α

[220756-92-3]



C₂₅H₃₅NO₄ 413.556

Tetramic acid deriv. Similar to Equisetin, E-142. Prod. by an unknown fungus, LL-49F233.

Singh, M.P. et al., *J. Antibiot.*, 1998, **51**, 1109-1112

Antibiotic M 259

A-1202

Antibiotic A 6413. A 6413. M 259. NSC 51954

Benzodiazepine-type antibiotic. Struct. unknown. Prod. by *Streptomyces nigellus*. Antiinflammatory and cytotoxic agent. Yellow prisms (Me₂CO/EtOAc). Sol. CHCl₃, DMSO, DMF, MeOH, EtOH; fairly sol. butanol, Me₂CO, EtOAc, hexane, C₆H₆; poorly sol. H₂O. Mp 155-157° (dec.). $[\alpha]_D^{25}$ +0.9 (EtOH). λ_{\max} 235; 335 (EtOH). λ_{\max} 235 (E1%/1cm 480); 335 (E1%/1cm 960) (EtOH) (Berdy).

► LD₅₀ (mus, ipr) 1.47 mg/kg, LD₅₀ (mus, ivn) 2.21 mg/kg.


U.S. Pat., 1963, 3 094 461; CA, 59, 5733d

Antibiotic MBH 001

A-1203

2-(1-Hydroxyethyl)-2-methyl-4-(2-methylpropyl)-5(2H)-oxazolone, 9CI. MBH 001

[147368-20-5]




C₁₀H₁₇NO₃ 199.249

Prod. by an insect symbiotic microorganism N1-Ishi-YC50807-1-7. Shows herbicidal props. Sol. MeOH, butanol, EtOAc, DMSO, hexane, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{25}$ 0.

Japan. Pat., 1992, 92 342 576; CA, 118, 211447 (isol, uv, ir, pmr, cmr)

Antibiotic MBP 039-06 A-1204

2,5-Dihydro-3-hydroxy-5-oxo-4-(1-oxo-2,4,6-octatrienyl)-1H-pyrrole-2-acetic acid, 9CI. MBP 039-06 [152158-05-9]




C₁₄H₁₅NO₅ 277.276
Prod. by *Phaeosphaeria* sp. L288. Proline hydroxylase inhibitor. Mp 212-214°. λ_{max} 254 (ε 8450); 386 (ε 19500) (MeOH) (Berdy). λ_{max} 257 (ε 6770); 347 (ε 15300) (MeOH-NaOH) (Berdy).

Japan. Pat., 1993, 93 239 023; CA, 120, 52821 (isol, pmr, cmr, ir, uv)

Antibiotic MBP 049-13 A-1205

MBP 049-13 [143208-09-7]




C₂₅H₃₇NO₃ 399.572
Prod. by *Ophiobolus rubellus*. Proline hydroxylase inhibitor. [α]_D +58.9. λ_{max} 248 (ε 5023); 292 (ε 8069) (MeOH) (Berdy). λ_{max} 224 (ε 8405); 291 (ε 6028) (MeOH/HCl) (Berdy).

Japan. Pat., 1992, 92 74 163; CA, 117, 169564g (isol, pmr, cmr, ir, uv)

Antibiotic 610MF A-1206

610MF




C₂₅H₃₁N₃O₃ 421.538
Pyrrole antibiotic. Struct. not certain, may have oxo and OH functions interchanged. Prod. by *Actinomadura madurae*. λ_{max} 547 (CHCl₃) (Berdy). λ_{max} 538 (EtOH/HCl) (Berdy).

Gerber, N.N. et al., J. Het. Chem., 1973, 10, 925-929

Antibiotic MPC 1001F A-1207

MPC 1001F




C₂₉H₂₆N₂O₁₁S 610.597
Related to Emestrin. Prod. by *Cladorrhinum* sp. KY4922. Powder. Mp 138-139°. [α]_D²⁸ +19.7 (c, 0.12 in CHCl₃). λ_{max} 262 (ε 21600) (MeOH).

Onodera, H. et al., Org. Lett., 2004, 6, 4101-4104 (isol, ir, pmr, cmr, ms)

Antibiotic MS 681 A-1208

MS 681



Antibiotic MS 681A X = Aib, Y = Trp
MS 681B X = Aib, Y = Phe
MS 681C X = Iva, Y = Trp
MS 681D X = Iva, Y = Phe

Peptide antibiotic complex. Prod. by *Myrothecium* sp. (KY6568; FERM-BP-4634). Inhibitor of myosin light chain kinase and cyclic nucleotide phosphodiesterase.

Antibiotic MS 681A

MS 681A

[173485-72-8]

C₆₁H₉₁N₁₃O₉ 1150.472
Pale yellow solid. Sol. MeOH, Me₂CO; fairly sol. EtOAc, CHCl₃; poorly sol. hexane, H₂O. [α]_D²⁶ +20.4 (c, 0.2 in MeOH). λ_{max} 217 (ε 26700); 281 (ε 3700); 289 (ε 3100) (MeOH).

Antibiotic MS 681B

MS 681B

[173485-73-9]

C₅₉H₉₀N₁₂O₉ 1111.435
Pale yellow solid. Sol. MeOH, CHCl₃, Me₂CO; fairly sol. EtOAc; poorly sol. hexane, H₂O. [α]_D²⁶ +28.2 (c, 0.2 in MeOH). λ_{max} 260 (ε 1200) (MeOH).

Antibiotic MS 681C

MS 681C

[173485-74-0]

C₆₂H₉₃N₁₃O₉ 1164.498
Pale yellow solid. Sol. MeOH, Me₂CO; fairly sol. EtOAc, CHCl₃; poorly sol. hexane, H₂O. [α]_D²⁶ +24.7 (c, 0.2 in MeOH). λ_{max} 218 (ε 31100); 281 (ε 4100); 289 (ε 3600) (MeOH).

Antibiotic MS 681D

MS 681D


[173485-75-1]

C₆₀H₉₂N₁₂O₉ 1125.462
Pale yellow solid. Sol. MeOH, CHCl₃, Me₂CO; fairly sol. EtOAc; poorly sol. hexane, H₂O. [α]_D²⁶ +24.4 (c, 0.2 in MeOH). λ_{max} 260 (ε 1500) (MeOH).

Japan. Pat., 1995, 7 285 993 (isol)
Yano, H. et al., J. Antibiot., 1997, 50, 992-997; 998-1006 (isol, uv, ir, pmr, cmr, ms)

Antibiotic MY 336a A-1209

1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-6-methyl-1,3-isoquinolinedimethanol, 9CI. MY 336a [95727-08-5]



C₁₃H₁₉NO₄ 253.297
Isol. from *Streptomyces gabonae* ATCC15282. Effective in blocking sympathetic nerve receptors. β-Adrenergic antagonist. Needles. Sol. acids, MeOH; fairly sol. Me₂CO, H₂O, Et₂O; poorly sol. hexane. Mp 177-178°. λ_{max} 293 (ε 3500) (EtOH/NaOH) (Derep). λ_{max} 285 (ε 2800) (EtOH) (Derep).

Eur. Pat., 1985, 130 554; CA, 102, 147548 (isol)

Kase, H. et al., J. Antibiot., 1986, 39, 354 (isol)

Hirayama, N. et al., Acta Cryst. C, 1990, 46, 86 (cryst struct)

Kaufman, T.S. et al., J.C.S. Perkin 1, 1996, 2497 (synth)

Antibiotic MYC 8005 A-1210

MYC 8005

[37372-39-7]

Tunicamycin-type antibiotic. Struct. unknown. Prod. by *Streptomyces exfoliatus* var. *echinosporus*. Acaricide and insecticide. Cryst. (1-butanol aq.). Sol. Py, MeOH, bases, Me₂CO-H₂O; poorly sol. Me₂CO, hexane. [α]_D +60 (Py). λ_{max} 219; 259 (MeOH). λ_{max} 219 (E1%/1cm 170); 259 (E1%/1cm 118) (MeOH) (Berdy).


▶ LD₅₀ (mus, ipr) 1.5 mg/kg, LD₅₀ (mus, orl) 37 mg/kg.

Meltzer, J. et al., Eur. J. Plant Pathol. (Neth. J. Plant Pathol.), 1972, 78, 1-14; 77-78

Antibiotic NA 22598A₁ A-1211


NA 22598A₁

[188674-15-9]





C₂₀H₃₈N₈O₇ 502.57
Prod. by *Streptomyces* sp. NA22598. Cytotoxic agent. Powder. Mp 190-195°.

Kuwahara, A. et al., J. Antibiot., 1997, 50, 712-713 (isol, pmr)


Antibiotic NA 30851ANA 30851A
[221209-96-7]C₃₄H₄₂N₂O₁₂ 670.712Prod. by *Streptomyces* sp. NA 30851A.

Japan. Pat., 1999, 99 49 768; CA, 130, 337075n

Antibiotic NFAT 68Methyl 7-(3-acetylamino-2,5-dihydroxyphenyl)-7-hydroxy-4,6-dimethyl-2,4-heptadienoate. NFAT 68
[165133-84-6]C₁₈H₂₃NO₆ 349.383Prod. by a *Streptomyces* sp. Blocks NFAT-dependent transcription and shows immunosuppressant props. Oil.Burren, N.S. et al., *J. Antibiot.*, 1995, 48, 380 (isol, pmr, cmr, props)**Antibiotic NG 311**2,6,6a,7,8,9,10,10a-Octahydro-6,8,10-trimethyl-1H-[2]benzopyrano[4,3-c]pyridin-1-one, 9CI. NG 311
[172318-70-6]C₁₅H₂₁NO₂ 247.336

Related to Fusaricidin, F-225. Prod. by a fungal sp. strain TF0407. Shows nerve growth factor-like activity. Neuroprotective and antidementia agent. Powder.

Japan. Pat., 1995, 95 233 168; CA, 124, 86990m

Antibiotic NG 391NG 391
[185850-14-0]C₂₂H₂₇NO₇ 417.458Tetramic acid deriv. Closely related to Fusarin C, F-227. Prod. by *Fusarium* sp.


A-1212

TF-0452, *Fusarium* sp. RK 97-94 and *Metarhizium anisopliae* (KOB1-3). Neuroprotectant. Mutagenic. Yellow powder. Sol. MeOH. [α]_D²⁰ +39.3 (MeOH). λ_{max} 283 (ε 9100); 366 (ε 35500) (MeOH).(3Z)-Isomer: Antibiotic NG 392. NG 392
[185850-15-1]C₂₂H₂₇NO₇ 417.458Prod. by *Fusarium* sp. TF-0452. Neuroprotectant. Yellow powder. Sol. MeOH. [α]_D²⁰ -13.4 (MeOH). λ_{max} 290 (ε 7600); 343 (ε 11900) (MeOH).(7Z)-Isomer: Antibiotic NG 393. NG 393
[185850-16-2]C₂₂H₂₇NO₇ 417.458Prod. by *Fusarium* sp. TF-0452 and *Metarhizium anisopliae* (KOB1-3). Neuroprotectant. Mutagenic. Yellow powder. Sol. MeOH. [α]_D²⁰ +32 (MeOH). λ_{max} 273 (ε 8000); 367 (ε 61200) (MeOH).


Japan. Pat., 1996, 96 319 289; CA, 126, 103149e

Kakeya, H. et al., *J. Antibiot.*, 2001, 54, 850-853 (isol)Hayashi, Y. et al., *Tetrahedron*, 2002, 58, 9839-9846 (synth, abs config)Krasnoff, S.B. et al., *J. Agric. Food Chem.*, 2006, 54, 7083-7088 (isol, activity)**Antibiotic NK 30424A**

NK 30424A

C₃₀H₄₆N₂O₁₀S 626.767Glutarimide-macrolide antibiotic. Prod. by *Streptomyces* sp. NA30424. Antitumour agent. Powder. Mp 129-131° dec. [α]_D²⁰ +10.8 (c, 0.5 in H₂O). λ_{max} 206 (log ε 4.28); 286 (log ε 2.53) (no solvent reported).

Stereoisomer: Antibiotic NK 30424B. NK 30424B

C₃₀H₄₆N₂O₁₀S 626.767Prod. by *Streptomyces* sp. NA30424. Antitumour agent. Powder. Mp 129-133° dec. [α]_D²⁰ +11.2 (c, 0.5 in H₂O). λ_{max} 206 (log ε 4.28); 283 (log ε 2.72) (no solvent reported).Takayasu, Y. et al., *J. Antibiot.*, 2001, 54, 1111-1115 (isol, pmr, cmr)**Antibiotic NK 170204B**4-Hydroxy-6-(10-hydroxy-3,7,9,12,13-pentamethyl-2,5,7,11,13-pentadecapent-2-enyl)-3-methoxy-5-methyl-2(1H)-pyridinone, 9CI. NK 170204B
[161161-46-2]

A-1216

C₂₇H₃₉NO₄ 441.609Closely related to Piericidin, P-421. Prod. by *Streptomyces* sp. NK170204.

Japan. Pat., 1994, 94 293 736; CA, 122, 158767c

Antibiotic NP 522

NP 522

C₁₄H₁₅N₃O₂ 257.291Struct. unknown. Prod. by *Streptomyces hygroscopicus*. Antibacterial agent. Prisms (MeOH/EtOAc). Sol. MeOH, Me₂CO, EtOH, butyl acetate, CHCl₃, EtOAc; fairly sol. H₂O, C₆H₆, Et₂O; poorly sol. hexane. Mp 204-205°. [α]_D²⁰ -190 (c, 2 in MeOH). λ_{max} 220 (E1%/1cm 2741); 274 (E1%/1cm 250); 281 (E1%/1cm 266); 290 (E1%/1cm 229) (MeOH) (Berdy).▶ LD₅₀ (mus, ims) 800 mg/kg; LD₅₀ (mus, scu) 800 mg/kg.

Japan. Pat., 1964, 64 10 247; CA, 61, 12593d

Antibiotic NRCS 15


NRCS 15

[52229-10-4]

Related to Ferrimycin A₁, F-37. Prod. by *Streptomyces* sp. NRCS15. Siderophore. Antibacterial agent. Powder. Sol. H₂O, MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. hexane. λ_{max} 210 (EtOH). λ_{max} 210 (E1%/1cm 122.5) (EtOH) (Berdy). λ_{max} 208 (E1%/1cm 112) (EtOH-HCl) (Berdy). λ_{max} 222 (E1%/1cm 12.2) (EtOH-NaOH) (Berdy).▶ LD₅₀ (mus, scu) 50 mg/kg.Haroun, B.M. et al., *J. Antibiot.*, 1974, 27, 14-19**Antibiotic OPC 15161**

OPC 15161

[121071-92-9]

C₁₈H₂₁N₃O₃ 327.382Artifact. Degradn. prod. of Antibiotic OPC 15160 (isol. from *Thielavia major*). Inhibitor of superoxide anion generation. More potent than OPC 15160. Pale yellow prisms (MeOH). Mp 223.5-225.5°. [α]_D²⁰ 0. λ_{max} 220 (ε 49400); 279 (ε 10320); 354 (ε 7370) (EtOH) (Derep).Nakano, Y. et al., *J. Antibiot.*, 1991, 44, 52 (isol, pmr, cmr, struct)Jing, H. et al., *Heterocycles*, 1992, 34, 1847 (synth, bibl)Shinham, K. et al., *Tetrahedron*, 2000, 56, 7427-7431 (synth)


A-1218

A-1219

A-1220

Antibiotic PD 116152 A-1221

Methyl 6-formyl-4,7,9-trihydroxy-8-methyl-1-phenazinecarboxylate, 9CI. PD 116152 [101708-64-9]



$C_{16}H_{12}N_2O_6$ 328.281


Phenazine antibiotic. Similar to Lomofungin, L-241. Prod. by *Streptomyces lomondensis* subsp. *galanosa*. Active against *Streptococcus pneumoniae* and murine P388 leukemia. Fairly sol. MeOH, EtOAc; poorly sol. $CHCl_3$, hexane. λ_{max} 262 (€ 21000); 294 (€ 23800); 351 (€ 19800); 415 (€ 5100); 440 (€ 4700) (MeOH/KOH) (Derep). λ_{max} 226 (€ 15900); 271 (€ 26540); 385 (€ 6070) (MeOH) (Derep).

Tunac, J.B. *et al.*, *J. Antibiot.*, 1986, **39**, 192 (isol, props)

Smitka, T.A. *et al.*, *J. Antibiot.*, 1986, **39**, 800 (isol, cryst struct)

Antibiotic PDE I A-1222

PDE I [62497-62-5]



R = -CONH₂

$C_{13}H_{13}N_3O_5$ 291.263

Isol. from culture closely related to *Streptomyces griseoflavus*. Shows antitumour activity. Inhibitor of cyclic adenosine-3',5'-monophosphate phosphodiesterase. Cryst. (MeOH). Sol. DMSO, Py; fairly sol. MeOH; poorly sol. EtOH, C_6H_6 . Mp 235° dec. λ_{max} 253 (€ 41000); 310 (€ 17000) (MeOH) (Derep). λ_{max} 256 (€ 39000); 320 (€ 19000) (HCl) (Berdy). λ_{max} 234 (€ 33000); 253 (€ 38000); 338 (€ 22000) (NaOH) (Berdy).

▶ DF4936665

Enomoto, Y. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 1331; 1337; 1979, **43**, 559 (isol, uv, ir, pmr, cmr, cryst struct, synth)

Carter, P. *et al.*, *J.A.C.S.*, 1987, **109**, 2711 (synth, bibl)

Boger, D.L. *et al.*, *J.A.C.S.*, 1987, **109**, 2717 (synth)

Bolton, R.E. *et al.*, *J.C.S. Perkin 1*, 1987, 931 (synth)

Antibiotic PDE II A-1223

PDE II [62874-94-6]

As Antibiotic PDE I, A-1222 with R = Ac

$C_{14}H_{14}N_2O_5$ 290.275

Isol. from culture closely related to *Streptomyces griseoflavus*. Shows antitumour props. Inhibitor of cyclic adenosine-3',5'-monophosphate phosphodiesterase. Sol. DMSO, Py; fairly sol. MeOH; poorly sol. EtOH, C_6H_6 . Mp 253° dec. λ_{max} 266 (€ 35600); 323 (€ 14600) (pH 2) (Derep). λ_{max} 266 (€ 26000); 344 (€ 17900) (pH 12) (Derep). λ_{max} 263 (€ 38200); 275 (€ 17000); 310 (€ 11700) (MeOH) (Derep).

▶ DF4936663

Enomoto, Y. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 1331; 1337; 1979, **43**, 559 (isol, struct, synth)


Bolton, R.E. *et al.*, *J.C.S. Perkin 1*, 1987, 931 (synth)

Carter, P. *et al.*, *J.A.C.S.*, 1987, **109**, 2711 (synth, bibl)

Boger, D.L. *et al.*, *J.A.C.S.*, 1987, **109**, 2717 (synth)

Antibiotic PF 1018 A-1224

PF 1018 [131256-42-3]



$C_{28}H_{35}NO_3$ 433.589


Prod. by *Humicola* sp. Insecticidal agent. Pale yellow cryst. (MeOH). Sol. $CHCl_3$, Me_2CO , EtOAc; poorly sol. H_2O . Mp 182-184°. $[\alpha]_D^{24}$ -185 (c, 1 in $CHCl_3$). λ_{max} 251 (E1%/1cm 294); 320 (E1%/1cm 355) (MeOH) (Berdy). λ_{max} 235 (E1%/1cm 229); 332 (E1%/1cm 471) (MeOH/HCl) (Berdy). λ_{max} 254 (E1%/1cm 326); 316 (E1%/1cm 340) (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 1.25-2.5 mg/kg, LD₅₀ (mus, orl) 37 mg/kg.

Gomi, S. *et al.*, *J. Antibiot.*, 1994, **47**, 571

Antibiotic PF 1052 A-1225

PF 1052 [147317-15-5]



$C_{26}H_{39}NO_4$ 429.598

Tetramic acid deriv. Similar to Vermisporin, V-77. Prod. by *Phoma* sp. PF1052 and *Phoma* sp. FK1-1840. Antibacterial agent. Oil. $[\alpha]_D$ +52.9 ($CHCl_3$). λ_{max} 229 (€ 6100); 291 (€ 12100) (MeOH).


Japan. Pat., 1992, 92 316 578; *CA*, **118**, 211424j

Kock, I. *et al.*, *Dissertation*, Univ. of Paderborn, Germany, 2005, (isol, pmr, cmr, ms)

Koyama, N. *et al.*, *J. Antibiot.*, 2005, **58**, 338-345 (isol, pmr, cmr)

Antibiotic PF 1140 A-1226

2,6,6a,7,8,9,10,10a-Octahydro-2-hydroxy-6,6a,8,10-tetramethyl-1H-[2]benzopyrano[4,3-c]pyridin-1-one, 9CI. Fungicide PF 1140. PF 1140 [173560-58-2]



Absolute Configuration

$C_{16}H_{23}NO_3$ 277.363

Related to Cordyipyridone D, C-647 and Fusaricide, F-225. Prod. by *Eupenicillium* sp. PF1140. Antifungal agent. Cryst. Sol. MeOH. $[\alpha]_D$ -148 (MeOH). λ_{max} 286 (MeOH).


Japan. Pat., 1995, 95 267 962; *CA*, **124**, 143744k (isol, props)

Fujita, Y. *et al.*, *J. Antibiot.*, 2005, **58**, 425-427 (abs config)

Fujita, Y. *et al.*, *Tet. Lett.*, 2005, **46**, 5885-5888 (biosynth)

Antibiotic PF 1163A A-1227

PF 1163A



Absolute Configuration

$C_{27}H_{43}NO_6$ 477.64

Prod. by *Penicillium* sp. PF1163. Antifungal agent. Oil. $[\alpha]_D^{25}$ -91.8 (c, 1 in MeOH). λ_{max} 224 (€ 11500); 276 (€ 1700); 282 (€ 1400) (MeOH).

11-Deoxy: Antibiotic PF 1163B. PF 1163B

$C_{27}H_{43}NO_5$ 461.64

Prod. by *Penicillium* sp. PF 1163. Antifungal agent. Oil. $[\alpha]_D^{25}$ -111.6 (c, 1.2 in MeOH). λ_{max} 224 (€ 11000); 276 (€ 1500); 282 (€ 1300) (MeOH).

Tatsuta, K. *et al.*, *J. Antibiot.*, 1999, **52**, 1146-1151 (synth, abs config)

Nose, H. *et al.*, *J. Antibiot.*, 2000, **53**, 33-37; 38-44; 2002, **55**, 969-974 (isol, pmr, cmr, activity)


Bouazza, F. *et al.*, *Org. Lett.*, 2003, **5**, 4049-4052 (synth)

Antibiotic PI 091 A-1228

PI 091

[129051-63-4]

[169869-85-6, 169735-66-4]



$C_{17}H_{29}NO_4$ 311.42


Prod. by *Paecilomyces* sp. F-3430. Platelet aggregation inhibitor. Oil. $[\alpha]_D^{29}$ -26.3 (c, 0.8 in CHCl_3). Obt. as a diastereoisomeric 1:1 mixture. Probable artifact.

[169735-66-4]

Shiraki, R. *et al.*, *J.O.C.*, 1996, **61**, 2845-2852 (synth, ir, pmr, cmr)
Iwasama, N. *et al.*, *J.O.C.*, 1997, **62**, 1918-1919 (synth)

Antibiotic PM 94128 A-1229

PM 94128
[182292-49-5]



$\text{C}_{22}\text{H}_{34}\text{N}_2\text{O}_6$ 422.52


Prod. by a marine *Bacillus* sp. Antitumour agent. Mp 172-173°. $[\alpha]_D^{25}$ -88.9 (c, 2 in CHCl_3). Related to Amicoumacin B. λ_{max} 208 (ε 27000); 246 (ε 6400); 314 (ε 4380) (no solvent reported).

Canedo, L.M. *et al.*, *J. Antibiot.*, 1997, **50**, 175-176 (isol, uv, ir, pmr, cmr)

Patel, S.K. *et al.*, *Org. Lett.*, 2003, **5**, 4081-4084 (synth)

Antibiotic R 39MF A-1230

R 39MF
[52288-96-7]




$\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}_2$ 377.485

Pyrrole antibiotic. Prod. by *Actinomadura madurae*. λ_{max} 554 (CHCl_3) (Berdy). λ_{max} 547 (EtOH/HCl) (Berdy).

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1973, **10**, 925-929

Antibiotic R 39SF A-1231

R 39SF
[52288-97-8]



$\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}_2$ 379.501


Pyrrole antibiotic. Prod. by *Actinomadura madurae*. λ_{max} 547 (CHCl_3) (Berdy).

λ_{max} 538 (EtOH/HCl) (Berdy).

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1973, **10**, 925-929

Antibiotic RK 1441A A-1232

1,2,3,10,11,11a-Hexahydro-8,11-dihydroxy-3,7-dimethoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI. RK 1441A
[132035-29-1]



$\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}_5$ 294.307

Anthramycin-type antibiotic. Prod. by *Streptomyces* sp. RK-1441. Possesses antibacteriophage activity. Yellow powder. Mp 115-120° dec. $[\alpha]_D^{21}$ +692 (c, 0.52 in MeOH). Related to Neothramycin. λ_{max} 235 (ε 31000); 264 (ε 11900); 318 (ε 5900) (50% EtOH) (Derep).

11-Ketone, 3-O-de-Me: Antibiotic RK

1441B. RK 1441B

[132138-94-4]

$\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_5$ 278.264

Prod. by *Streptomyces* sp. RK-1441. Possesses antibacteriophage activity. Pale yellow powder. Mp 224-225°. $[\alpha]_D^{21}$ +219 (c, 0.54 in EtOH). λ_{max} 235 (ε 55000); 267 (sh) (ε 15400); 300 (ε 8600); 310 (sh) (ε 6600) (50% EtOH) (Derep).

[132035-30-4]

Osada, H. *et al.*, *Agric. Biol. Chem.*, 1990, **54**, 2875; 2883 (isol, pmr, cmr, uv, props)

Antibiotic RP 6798† A-1233

RP 6798†

$\text{C}_{42}\text{H}_{54}\text{N}_{10}\text{O}_{13}$ 906.948

Nucleoside-peptide type antibiotic. Struct. unknown. Prod. by *Streptomyces peruviansis*. Powder. Sol. H_2O , DMF; fairly sol. butanol- H_2O ; poorly sol. Me_2CO , hexane. Mp 240-245°. $[\alpha]_D$ -21 (NaOH aq.). λ_{max} 222; 257 (MeOH). λ_{max} 222 (E1%/1cm 519); 257 (E1%/1cm 235) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) .1 mg/kg, LD₅₀ (mus, scu) .5 mg/kg.

U.K. Pat., 1960, 846 801; CA, **55**, 6778h

Antibiotic RP 13252 A-1234

RP 13252

Benzodiazepine-type antibiotic. Struct. unknown. Prod. by *Streptomyces croceus*. Cytotoxic agent. Powder. Sol. MeOH, CHCl_3 ; fairly sol. Et_2O , H_2O ; poorly sol. hexane. λ_{max} 222; 238; 313 (MeOH). λ_{max} 222 (E1%/1cm 760); 238 (E1%/1cm 690); 313 (E1%/1cm 126) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) 2.5 mg/kg, LD₅₀ (mus, ipr) .75 mg/kg, LD₅₀ (mus, scu) 2.5 mg/kg, LD₅₀ (mus, orl) 6 mg/kg.

Fr. Pat., 1968, 1 516 743; CA, **71**, 2128

Antibiotic S 433 A-1235

S 433

[82642-91-9]

Related to Cervinomycin A₁, C-287. Struct. unknown. Prod. by *Streptomyces* sp. ATCC31668. Antibacterial and antitumour agent. Orange-red cryst. (CHCl_3). Sol. DMF, DMSO, AcOH; fairly sol. CHCl_3 , CH_2Cl_2 ; poorly sol. MeOH, H_2O , hexane, Et_2O . λ_{max} 266; 321 (H_2O). λ_{max} 265 (E1%/1cm 414); 315 (E1%/1cm 335) (HCl) (Berdy). λ_{max} 266 (E1%/1cm 414); 321 (E1%/1cm 326) (pH 7.4 buffer) (Berdy). λ_{max} 266 (E1%/1cm 403); 322 (E1%/1cm 326) (pH 9.3 buffer) (Berdy).

Eur. Pat., 1982, 50 749; CA, **97**, 90410p

Antibiotic S 53210A A-1236

S 53210A

Thiazole-peptide antibiotic complex.

Struct. unknown. Prod. by *Microclavosporia brunnea* sp. S53210A.

Antibiotic S 53210A-I

S 53210A-I

[69670-88-8]

Berdy 6614. Yellow powder. Sol. DMF, MeCN, dioxan, DMSO; fairly sol. CHCl_3 , MeOH, EtOH; poorly sol. H_2O , hexane. $[\alpha]_D$ +104 (CHCl_3). λ_{max} 215; 305; 355 (MeCN) (Berdy).

Antibiotic S 53210A-II

S 53210A-II

[69670-89-9]

Yellow powder. Sol. CHCl_3 , DMF, MeCN, dioxan, DMSO; fairly sol. MeOH, EtOH; poorly sol. H_2O , hexane. $[\alpha]_D$ +107 (CHCl_3). λ_{max} 215; 305; 355 (MeCN) (Berdy).

Antibiotic S 53210A-III

S 53210A-III


[69670-90-2]

Yellow powder. Sol. CHCl_3 , DMSO, MeCN, DMF, dioxan; fairly sol. MeOH, EtOH; poorly sol. H_2O , hexane. $[\alpha]_D$ +67.2 (CHCl_3). λ_{max} 215; 300; 350 (MeCN) (Berdy).

U.S. Pat., 1980, 4 216 206; CA, **93**, 184280r

Antibiotic S 54832A A-1237

S 54832A



Depsideptide antibiotics. Struct. shown is S 54832A-I. Isol. from *Micromonospora globosa*. Active against gram-positive bacteria, mycoplasmas and *Neisseriae* sp.

Antibiotic S 54832A-I

S 54832A-I

[73666-05-4]

C₅₉H₅₅N₁₃O₁₉S₅ 1410.489Yellow powder. Mp 310° dec. [α]_D²⁰
+118.7 (c, 1.140 in Py). pK_{a1} 7.65; pK_{a2}
10.7 (84% 2-methoxyethanol).Dihydro: **Antibiotic S 54832A-IV. S****S 54832A-IV**

[73666-08-7]

C₅₉H₅₇N₁₃O₁₉S₅ 1412.505Pale yellow powder. Mp 310° dec. [α]_D²⁰
+153.2 (c, 0.752 in Py).**Antibiotic S 54832A-II****S 54832A-II**

[73666-06-5]

Pale yellow powder. Mp 310° dec.
Struct. unknown.**Antibiotic S 54832A-III****S 54832A-III**

[73666-07-6]


Pale yellow powder. Mp 310° dec.
Struct. unknown.

[94588-49-5, 94588-48-4]

Ger. Pat., 1979, 2 921 148; CA, **93**, 6124 (*isol*,
props, *nmr*)U.S. Pat., 1984, 4 478 831; CA, **102**, 77266
(*isol*)**Antibiotic SAB 711****SAB 711**


[89342-57-4]

A-1238


C₂₇H₅₁N₇O₅ 553.744Oligopeptide antibiotic. Isol. from
Streptomyces kaizukaensis. Protease in-
hibitor. Amorph. powder + 4H₂O. Sol.
H₂O. Mp 176-182° dec. [α]_D²³ -35.1 (c,
0.115 in MeOH). Similar to Leupeptin.Japan. Pat., 1983, 83 198 453; CA, **100**, 137364
(*isol*, *props*)**Antibiotic SB 212305****SB 212305**

[170591-56-7]


A-1239

C₂₀H₁₇N₃O₈S 459.436Phenazine antibiotic. Prod. by a *Streptomyces* sp. Inhibitor of metalloenzymes such as metallolactamases. Sol. MeOH, bases, butanol; poorly sol. H₂O. λ_{\max} 212 (€ 25422); 249 (€ 18754); 305 (€ 33757); 380 (€ 5000); 530 (€ 2084) (MeOH) (Berdy). λ_{\max} 212 (€ 22288); 251 (€ 22921); 278 (€ 23755); 378 (€ 7084); 440 (€ 2917) (MeOH/HCl) (Berdy). λ_{\max} 249 (€ 18754); 305 (€ 36674); 370 (€ 2917); 515 (€ 2500) (MeOH/NaOH) (Berdy).Gilpin, M.L. et al., *J. Antibiot.*, 1995, **48**, 1081
(*isol*, *pmr*, *cmr*, *uv*, *props*)**Antibiotic SB 217452****SB 217452**


A-1240

C₁₆H₂₄N₆O₉S 476.466Prod. by *Streptomyces* sp. ATCC 700974.
Enzymic cleavage prod. of Albomycin.
Seryl tRNA synthetase inhibitor. [α]_D -
14.9 (c, 0.22 in H₂O). λ_{\max} 215; 229; 306
(1M HCl).Stefanska, A.L. et al., *J. Antibiot.*, 2000, **53**,
1346-1353**Antibiotic SB 219383****SB 219383**


A-1241

C₁₇H₂₃N₃O₉ 413.383Prod. by *Micromonospora* sp. NCIMB
40684. Tyrosyl tRNA synthetase inhi-
bitor. Powder. [α]_D²² +25.3 (EtOH aq./4M
HCl) (as Et ester). λ_{\max} 225; 275 (H₂O).Stefanska, A.L. et al., *J. Antibiot.*, 2000, **53**,
345-350 (*isol*)Houge-Frydrych, C.S.V. et al., *J. Antibiot.*,
2000, **53**, 351-356 (*struct*)**Antibiotic SB 291071****SB 291071**

A-1242


C₂₆H₄₂N₂O₉ 526.626Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of lipoprotein
associated phospholipase A₂
(LpPLA₂).*Homologue* (R = -C₁₁H₂₁): **Antibiotic SB**
291072. SB 291072C₂₈H₄₄N₂O₉ 552.664Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of LpPLA₂.
Posn. of double bond in R group not
determined.*Homologue* (R = -(CH₂)₁₀CH₃): **Anti-**
biotic SB 253514. SB 253514C₂₈H₄₆N₂O₉ 554.679Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of LpPLA₂.
Powder. λ_{\max} 252 (€ 14900)
(MeOH).*Homologue* (R = -C₁₃H₂₅): **Antibiotic SB**
253517. SB 253517C₃₀H₄₈N₂O₉ 580.717Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of LpPLA₂.
Posn. of double bond in R group not
determined.*Homologue* (R = -(CH₂)₁₂CH₃): **Anti-**
biotic SB 253518. SB 253518C₃₀H₅₀N₂O₉ 582.733Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of LpPLA₂.*Homologue* (R = -C₁₅H₂₉): **Antibiotic SB**
291073. SB 291073C₃₂H₅₂N₂O₉ 608.771Prod. by *Pseudomonas fluorescens*
DSM 11579. Inhibitor of LpPLA₂.
Posn. of double bond in R group not
determined.Thirkettle, J. et al., *J. Antibiot.*, 2000, **53**, 664-
669; 733-735 (*isol*, *activity*)Busby, D.J. et al., *J. Antibiot.*, 2000, **53**,
670-676 (*pmr*, *cmr*, *ms*, *cryst struct*)**Antibiotic SB 315021****SB 315021**

A-1243

C₂₈H₄₆N₂O₉ 554.679Major component of a series of
unstable factors. Related to
Cyclocarbamide A, C-852. Prod. by
Pseudomonas fluorescens DSM 11579.
Inhibitor of lipoprotein-associated
phospholipase A₂ (LpPLA₂). Off-white
powder. λ_{\max} 280 (€ 10400)
(EtOH).Thirkettle, J. et al., *J. Antibiot.*, 2000, **53**, 664-
669 (*isol*, *activity*)Busby, D.J. et al., *J. Antibiot.*, 2000, **53**, 670-
676 (*pmr*, *cmr*, *ms*, *cryst struct*)**Antibiotic Sch 40832****Sch 40832**

[205925-96-8]

A-1244



C₈₄H₁₀₄N₁₈O₂₆S₅ 1942.181

Prod. by *Micromonospora carbonaceae* var. *africana*. Solid. $[\alpha]_D^{26}$ -163.1 (c, 0.9 in CHCl₃). λ_{\max} 234 (ε 68320); 295 (ε 16960); 307 (ε 11630) (MeOH).


Puar, M.S. *et al.*, *J. Antibiot.*, 1998, **51**, 221-224 (pmr, cmr, ms, struct)

Antibiotic Sch 40873

Sch 40873

[120201-09-4]

[119948-40-2 (hydrochloride)]

C₂₄H₅₁N₉O₂ 497.726

Prod. by an *Actinomadura* sp. Antifungal agent. Solid (as hydrochloride). Sol. MeOH, H₂O. Mp 270° (hydrochloride). $[\alpha]_D^{26}$ -1.5 (c, 0.5 in MeOH).


► LD₅₀ (mus, ivn) 1.5 mg/kg.

Hegde, V.R. *et al.*, *J.O.C.*, 1989, **54**, 2402 (isol, struct)

Antibiotic Sch 42137

Sch 42137

[142383-44-6]



Absolute Configuration

C₂₉H₂₇NO₁₀ 549.533

Prod. by an *Actinoplanes* sp. ATCC53878. Antifungal agent. Yellow solid. $[\alpha]_D^{26}$ +681 (c, 0.5 in DMF). λ_{\max} 253 (ε 35500); 320 (ε 12900); 379 (sh) (ε 22900); 395 (ε 25100) (MeOH) (Derep). λ_{\max} 214 (ε 1.66); 225; 255 (ε 1.817); 260; 310; 319 (ε 0.66); 370; 381 (ε 1156); 385; 396 (ε 1.297) (MeOH) (Berdy).

11-Ac: [144050-51-1]C₃₁H₂₉NO₁₁ 591.57

Prod. by *Actinoplanes* sp. ATCC53878. λ_{\max} 225; 260; 310; 370; 385 (MeOH) (Berdy).

14-Ac: [144074-97-5]C₃₁H₂₉NO₁₁ 591.57

Prod. by *Actinoplanes* sp. ATCC53878. λ_{\max} 225; 260; 310; 370; 385 (MeOH) (Berdy).


Cooper, R. *et al.*, *J. Antibiot.*, 1992, **45**, 444-453; C3 (isol, pmr, cmr, struct)

U.S. Pat., 1992, 5126350; CA, **117**, 198525a (isol, derivs)

Antibiotic Sch 52900

Sch 52900

[174285-71-3]



Relative Configuration

C₃₁H₃₀N₆O₇S₄ 726.878

Epithiodioxopiperazine antibiotic. Closely related to Verticillin A, V-89. Prod. by *Gliocladium* sp. SCF1168 and *Gliocladium roseum* 1A. Signal transduction pathway inhibitor. Oncogene induction inhibitor. Powder. Sol. MeOH, DMSO, THF, CH₂Cl₂; poorly sol. hexane, H₂O. Mp 202-205° dec. $[\alpha]_D^{22}$ +769 (c, 0.1 in CHCl₃). λ_{\max} 242; 301 (MeOH) (Berdy).

1''-Deoxy: **Antibiotic Sch 52901**. Sch 52901

[174285-72-4]

C₃₁H₃₀N₆O₆S₄ 710.879

Prod. by *Gliocladium* sp. SCF1168 and *Gliocladium roseum* 1A. Signal transduction pathway inhibitor. Oncogene induction inhibitor. Powder. Sol. MeOH, CH₂Cl₂, THF, DMSO; poorly sol. hexane, H₂O. Mp 199-200° dec. $[\alpha]_D^{22}$ +688 (c, 0.1 in CHCl₃). λ_{\max} 241; 305 (MeOH) (Berdy).


Chu, M. *et al.*, *J. Antibiot.*, 1995, **48**, 1440-1445 (isol, uv, ir, pmr, cmr)

Dong, J.-Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1510-1513 (isol, pmr, cmr)

Antibiotic Sch 54445

Sch 54445

[175448-19-8]



Relative Configuration

C₃₀H₂₉ClN₂O₉ 597.02

Prod. by *Actinoplanes* sp. SCC2314. Antifungal agent. Brownish-yellow needles. Mp 201-203° dec. $[\alpha]_D^{23}$ -558 (c, 0.2 in CHCl₃). λ_{\max} 215; 251; 270 (sh); 326; 393 (MeOH).

Chu, M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 525-528 (isol, pmr, cmr)


Antibiotic Sch 56036

Sch 56036

[215596-10-4]

[175448-20-1]

A-1249



Relative Configuration

C₃₀H₃₁NO₈ 533.577

Related to Albofungin, A-244. Prod. by an *Actinoplanes* sp. SCC2314. Antifungal agent. Amorph. yellow powder. Sol. CHCl₃, MeOH, CH₂Cl₂, DMSO; poorly sol. H₂O, hexane. Mp 195°. $[\alpha]_D^{25}$ +212.5 (c, 0.2 in CHCl₃). λ_{\max} 260; 290 (sh); 315; 354 (sh); 365 (MeOH).


► LD₅₀ (mus, ipr) 2.5 mg/kg.

U.S. Pat., 1994, 5 494 913; CA, **124**, 258681e
Chu, M. *et al.*, *Tet. Lett.*, 1998, **39**, 7649-7652 (isol, uv, ir, pmr, cmr)

Antibiotic Sch 210971

Sch 210971

A-1250



Relative Configuration

C₂₅H₃₅NO₆ 445.555

Enolised β-triketone. Prod. by *Chaetomium globosum* (Mer-0229). Mp 126-128°. $[\alpha]_D^{24}$ +127.5 (c, 0.1 in Me₂CO). λ_{\max} 220; 290 (MeOH).

6'-Epimer: **Antibiotic Sch 210972**. Sch 210972C₂₅H₃₅NO₆ 445.555

Prod. by *Chaetomium globosum* (Mer-0229). Chemokine receptor CCR-5 inhibitor. Mp 149-150°. $[\alpha]_D^{24}$ +40 (c, 0.1 in Me₂CO). λ_{\max} 220; 295 (MeOH).

6'-Epimer, Me ester: **Antibiotic Sch 213766**. Sch 213766

[364779-57-7]

C₂₆H₃₇NO₆ 459.581

Prod. by *Chaetomium globosum* (Mer-0229). Chemokine receptor CCR-5 inhibitor. Gum. λ_{\max} 220; 295 (MeOH).


Yang, S.-W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1025-1028 (Sch 210971, Sch 210972, *cryst struct*)

Yang, S.-W. *et al.*, *J. Antibiot.*, 2007, **60**, 524-528 (Sch 213766)

Antibiotic Sch 538415

A-1251

1,4,6,9-Tetramethylpyrido[3,2-g]quinoline-2,5,8,10-(1H,9H)-tetrone, 9CI. Sch 538415


C₁₆H₁₄N₂O₄ 298.298

Related to Diazaquinomycin B, D-295. Isol. from an unidentified bacterium and a *Streptomyces* sp. Acyl carrier protein synthase inhibitor. Antineoplastic agent. Orange-red powder. λ_{\max} 254; 268; 282; 302; 313; 348; 435 (no solvent reported).

Chu, M. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 3827-3829 (*isol*, *pmr*, *cmr*)

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2006, **69**, 804-806 (*isol*, *pmr*, *cmr*)

Antibiotic Sch 725424 A-1252
Sch 725424



$C_{18}H_{23}NO_4$ 317.384

Enolised β -diketone. Prod. by *Kitasatospora* sp. (SPRI-0408). Antibacterial agent.

Yang, S.-W. *et al.*, *J. Antibiot.*, 2005, **58**, 192-195 (*isol*, *pmr*, *cmr*)

Antibiotic SF 1508 A-1253
SF 1508

[56378-68-8]

Nucleoside-type antibiotic. Struct. unknown. Prod. by *Streptomyces hygrosopicus* SF1508. Antifungal agent. Needles. Sol. acids, AcOH; fairly sol. H_2O ; poorly sol. EtOH, hexane. $[\alpha]_D^{20} +21.2$ (H_2O). Dec. at 192-195°. λ_{\max} 286 (E1%/1cm 230) (H_2O) (Berdy). λ_{\max} 295 (E1%/1cm 240) (HCl) (Berdy). λ_{\max} 242 (E1%/1cm 280); 243 (E1%/1cm 282); 312 (E1%/1cm 180); 318 (E1%/1cm 180) (NaOH aq.) (Berdy).

► LD₅₀ (mus, ipr) 25 mg/kg.

Japan. Pat., 1975, 75 19 992; *CA*, **83**, 56712q

Takashi, O. *et al.*, *CA*, 1977, **86**, 53844k

Antibiotic SF 1739 A-1254
SF 1739

[61991-01-3]

$C_{22}H_{29}N_3O_6$ 431.488

Struct. unknown. See also Antibiotic SF 1739HP in N-32. Isol. from *Streptomyces griseoplanus*. Antibiotic active against gram-positive and -negative bacteria. Orange powder (as hydrochloride). Sol. H_2O , MeOH; poorly sol. C_6H_6 , hexane, Et₂O. Mp 120-135° (hydrochloride). $[\alpha]_D^{20} +163$ (hydrochloride). λ_{\max} 268 (E1%/1cm 190); 455 (E1%/1cm 30) (H_2O) (Berdy). λ_{\max} 262 (E1%/1cm 166); 410 (E1%/1cm 25) (HCl) (Berdy). λ_{\max} 272 (E1%/1cm 144); 510 (E1%/1cm 43) (NaOH aq.) (Berdy).


► LD₅₀ (mus, ivn) .54 mg/kg. VU6306000

Japan. Pat., 1976, 76 125 798; *CA*, **86**, 119253a (*isol*)

Antibiotic SF 2140 A-1255

SF 2140

[91284-30-9]



$C_{18}H_{20}N_2O_6$ 360.366

Nucleoside antibiotic. Prod. by *Actinomadura albolutea*. Antiviral agent. Shows weak broad spectrum of antibacterial activity. Cryst. ($CHCl_3$ /MeOH). Sol. MeOH, EtOAc, EtOH, Me₂CO, $CHCl_3$; fairly sol. C_6H_6 ; poorly sol. H_2O , hexane. Mp 174-176°. $[\alpha]_D^{20} +59$ (c, 1.0 in MeOH). Similar to Neosidomycin, N-138. λ_{\max} 222 (ϵ 34600); 258 (sh) (ϵ 7630); 265 (ϵ 8210); 284 (ϵ 6260); 294 (ϵ 6910) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 2000-4000 mg/kg. MQ3160000

Di-Ac:

Cryst. Mp 114°.

[83652-17-9, 93207-27-3]

Japan. Pat., 1982, 82 85 397; *CA*, **97**, 214253 (*isol*)


Ito, T. *et al.*, *J. Antibiot.*, 1984, **37**, 931 (*isol*, *cryst struct*)

Tohyama, H. *et al.*, *J. Antibiot.*, 1984, **37**, 1145 (*isol*)

Buchanan, J.G. *et al.*, *J.C.S. Perkin I*, 1994, 1417 (*synth*)

Antibiotic SF 2312 A-1256

(1,5-Dihydroxy-2-oxo-3-pyrrolidinyloxy)phosphonic acid, 9CI. SF 2312 [107729-45-3]



$C_4H_8NO_6P$ 197.084

Prod. by *Micromonospora* sp. Active against *Pseudomonas* and *Proteus* spp. Sol. H_2O ; poorly sol. EtOH, MeOH, C_6H_6 . Isol. as Na salt.


Mono-Na salt:

Cryst. + 1/2 H_2O . Mp 160-164° dec.

Japan. Pat., 1985, 85 224 493; *CA*, **104**, 107918
Watabe, H. *et al.*, *CA*, 1987, **106**, 134903; 152686 (*isol*, *struct*, *props*)

Antibiotic SF 2369 A-1257

2,5-Bis[(3-isocyanobutanoyl)amino]-1-pentanol. SF 2369



$C_{15}H_{24}N_4O_3$ 308.38

Isol. from *Actinomadura* sp. SF-2369.


Japan. Pat., 1986, 86 280 286; *CA*, **107**, 152846w (*isol*)

Sasaki, T. *et al.*, *CA*, 1988, **108**, 201434s (*isol*)

Antibiotic SF 2425 A-1258

SF 2425

[120692-47-9]



$C_{25}H_{54}N_{10}O_2$ 526.767

Prod. by *Streptomyces* sp. SF 2425.

Powder + 2 H_2O (as trihydrochloride).

Sol. H_2O , MeOH; poorly sol. Me₂CO, hexane. Mp 124-127° (trihydrochloride). $[\alpha]_D^{22} +14.8$ (c, 1 in H_2O).

Japan. Pat., 1989, 89 63 587; *CA*, **111**, 193068v (*isol*)

Antibiotic SF 2695 A-1259

SF 2695

[130151-71-2]

$C_{29}H_{42}N_8O_{18}S$ 822.76

Related to Actinoplanone A in A-128.

Structs. unknown. Prod. by *Actinomadura* sp. SF2695. Antitumour agents.

Active against gram-positive bacteria and mycoplasma.

Antibiotic SF 2695A

SF 2695A

[138634-42-1]

$C_{36}H_{35}NO_{13}$ 689.671

Yellow powder. Sol. $CHCl_3$, Me₂CO, EtOAc; fairly sol. MeOH, hexane; poorly sol. H_2O . $[\alpha]_D -94.9$ ($CHCl_3$). λ_{\max} 211 (E1%/1cm 899); 237 (E1%/1cm 585); 263 (E1%/1cm 508); 399 (E1%/1cm 115) (MeOH) (Berdy). λ_{\max} 210 (E1%/1cm 880); 239 (E1%/1cm 574); 262 (E1%/1cm 547); 293 (E1%/1cm 255); 298 (E1%/1cm 117) (MeOH/HCl) (Berdy). λ_{\max} 213 (E1%/1cm 1231); 270 (E1%/1cm 376); 347 (E1%/1cm 155); 414 (E1%/1cm 169) (MeOH/NaOH) (Berdy).

Antibiotic SF 2695B

SF 2695B

[138634-41-0]

$C_{35}H_{33}NO_{13}$ 675.645


Yellow powder. Sol. $CHCl_3$, EtOAc, Me₂CO; fairly sol. MeOH, hexane; poorly sol. H_2O . $[\alpha]_D -106.7$ ($CHCl_3$). λ_{\max} 211 (E1%/1cm 916); 238 (E1%/1cm 598); 263 (E1%/1cm 511); 296 (E1%/1cm 245); 399 (E1%/1cm 117) (MeOH) (Berdy). λ_{\max} 211 (E1%/1cm 891); 238 (E1%/1cm 595); 262 (E1%/1cm 560); 295 (E1%/1cm 255); 397 (E1%/1cm 120) (MeOH/HCl) (Berdy). λ_{\max} 215 (E1%/1cm 1301); 270 (E1%/1cm 367); 346 (E1%/1cm 170); 414 (E1%/1cm 172) (MeOH/NaOH) (Berdy).

Japan. Pat., 1991, 91 198 783; *CA*, **116**, 57529x

Antibiotic SF 2768

A-1260

3-Cyano-N-[6-[[[(3-cyano-1-oxobutyl)amino]methyl]tetrahydro-2-hydroxy-2H-pyran-3-yl]butanamide, 9CI. SF 2768 [159392-35-5]



$C_{16}H_{24}N_4O_4$ 336.39

Aminoglycoside antibiotic. Prod. by *Streptomyces* sp. SF2768. Active against a wide spectrum of bacteria and fungi. Melanin formation inhibitor. Oil. Sol. $CHCl_3$, Me_2CO , MeOH, EtOAc; poorly sol. H_2O , hexane. $[\alpha]_D^{25} +25.6$ (MeOH). λ_{max} 200 (MeOH). λ_{max} 210 (MeOH) (Berdy).


Japan. Pat., 1994, 94 234 784; CA, 122, 8152j (isol)

Tabata, Y. et al., CA, 1996, 124, 255435s (isol)

Antibiotic SF 2771

A-1261

Hexahydro-6-methyl-3-methylenepyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. SF 2771 [168074-99-5]



$C_9H_{12}N_2O_2$ 180.206


Prod. by *Streptomyces* sp. SF2771. Antibacterial and cytotoxic agent. Cryst. Sol. MeOH, Me_2CO . $[\alpha]_D^{25} -109.1$ (MeOH). λ_{max} 225 (MeOH).

Japan. Pat., 1995, 95 165 761; CA, 123, 254695u (isol)

Antibiotic SF 2809

A-1262

SF 2809



SF 2809-I $R^1 = H$, $R^2 = CH_2CH_2NHAc$

SF 2809-II $R^1 = 4-Hydroxyphenyl$, $R^2 = CH_2CH_2NHAc$

SF 2809-III $R^1 = H$, $R^2 = CH_2CH_2OH$

SF 2809-IV $R^1 = 4-Hydroxyphenyl$, $R^2 = CH_2CH_2OH$

SF 2809-V $R^1 = Ph$, $R^2 = CH_2CH_2NHAc$

SF 2809-VI $R^1 = Ph$, $R^2 = CH_2CH_2OH$

Prod. by *Dactylosporangium* sp. SF2809. Chymase inhibitors.

Antibiotic SF 2809-I

SF 2809-I

[271580-72-4]

$C_{23}H_{23}N_3O_3$ 389.453

Pale yellow powder. λ_{max} 227 (€ 45900); 277 (€ 10900); 284 (€ 10900); 319 (€ 6420);

332 (€ 5250) (MeOH/1M HCl). λ_{max} 206 (€ 54900); 222 (€ 54500); 257 (sh) (€ 14000); 292 (€ 12100); 310 (€ 11700) (MeOH/1M NaOH).

Antibiotic SF 2809-II

SF 2809-II

[271580-73-5]

$C_{29}H_{27}N_3O_4$ 481.55

Pale red powder. λ_{max} 227 (€ 68600); 278 (€ 15500); 286 (€ 15500); 322 (€ 10600); 335 (€ 7730) (MeOH/1M HCl). λ_{max} 207 (€ 93200); 223 (€ 81600); 260 (sh) (€ 29000); 283 (€ 23200); 292 (€ 22700); 310 (sh) (€ 18800) (MeOH/1M NaOH).

Antibiotic SF 2809-III

SF 2809-III

[271580-74-6]

$C_{21}H_{20}N_2O_3$ 348.401

Pale yellow powder. λ_{max} 227 (€ 47000); 276 (€ 10800); 284 (€ 11000); 319 (€ 6450); 332 (€ 5050) (MeOH/1M HCl). λ_{max} 205 (€ 44300); 223 (€ 46300); 257 (sh) (€ 12200); 292 (€ 10500); 310 (€ 10100) (MeOH/1M NaOH).

Antibiotic SF 2809-IV

SF 2809-IV

[271580-75-7]

$C_{27}H_{24}N_2O_4$ 440.498

Pale red powder. λ_{max} 228 (€ 57400); 280 (€ 14500); 323 (€ 8020); 334 (€ 6170) (MeOH/1M HCl). λ_{max} 205 (€ 75700); 223 (€ 58000); 260 (sh) (€ 16700); 286 (€ 16000); 293 (€ 16000); 310 (€ 13600) (MeOH/1M NaOH).

Antibiotic SF 2809-V

SF 2809-V

[271580-76-8]

$C_{29}H_{27}N_3O_3$ 465.551

Powder. λ_{max} 227 (€ 62800); 278 (€ 14400); 284 (€ 14000); 323 (€ 9120); 335 (€ 6980) (MeOH/1M HCl). λ_{max} 207 (€ 83700); 222 (€ 75300); 257 (sh) (€ 23700); 284 (€ 17700); 292 (€ 18100); 310 (€ 15800) (MeOH/1M NaOH).

Antibiotic SF 2809-VI

SF 2809-VI

[271580-77-9]

$C_{27}H_{24}N_2O_3$ 424.498

Powder. λ_{max} 227 (€ 50200); 278 (€ 11900); 322 (€ 7490); 334 (€ 5960) (MeOH/1M HCl). λ_{max} 205 (€ 51700); 223 (€ 51100); 258 (sh) (€ 13200); 293 (€ 10500); 310 (€ 9790) (MeOH/1M NaOH).

Tani, M. et al., J. Antibiot., 2004, 57, 83-88; 89-96 (isol, uv, pmr, cmr, activity)

Antibiotic SF 2197B

SF 2197B

[105634-06-8]


Related to Matlystatin A₁. Struct. unknown. Prod. by *Microbispora* sp. SF-2197. Antibacterial agent. Powder. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{25} -20.2$ (MeOH). λ_{max} 200 (MeOH).

Japan. Pat., 1986, 86 141 889; CA, 106, 3851w (isol)

Antibiotic SMTP 7

A-1264

SMTP 7



$C_{51}H_{68}N_2O_{10}$ 869.106

Related to Stachybotrin A in S-490. Abs. config. only known at C-27. Rel. configs. determined for other centres. Prod. by *Stachybotrys microspora*. Light brown oil. $[\alpha]_D^{25} -35.9$ (c, 0.85 in MeOH). λ_{max} 213 (€ 82500); 257 (€ 18800); 302 (€ 5820) (MeOH).

27-Epimer: SMTP 7D. Antibiotic SMTP 7D

Prod. by *Stachybotrys microspora* fed with D-Orn. Light brown oil. $[\alpha]_D^{20} -2.4$ (c, 0.8 in MeOH).

Hu, W. et al., J. Antibiot., 2000, 53, 241-247; 2003, 56, 832-837 (isol, pmr, cmr)

Antibiotic SMTP 8

A-1265

SMTP 8

As Antibiotic SMTP 7, A-1264 with n = 4

$C_{52}H_{70}N_2O_{10}$ 883.132

Prod. by *Stachybotrys microspora*. Light brown oil. $[\alpha]_D^{25} -30$ (c, 1 in MeOH). λ_{max} 213 (€ 79500); 257 (€ 18200); 302 (€ 5600) (MeOH).

27-Epimer: SMTP 8D. Antibiotic SMTP 8D


Prod. by *Stachybotrys microspora* fed with D-Lys. Light brown oil. $[\alpha]_D^{20} -5.2$ (c, 0.8 in MeOH).

Hu, W. et al., J. Antibiot., 2000, 53, 241-247; 2003, 56, 832-837 (isol, pmr, cmr)

Antibiotic SPF 32629A

A-1266

SPF 32629A



R = H

$C_{17}H_{19}NO_4$ 301.341

Prod. by *Penicillium* sp. SPF-32629. Human chymase inhibitor. Pale yellow powder. $[\alpha]_D^{25} +26.4$ (c, 0.58 in $CHCl_3$). λ_{max} 206 (€ 22000); 286 (€ 4300) (MeOH).

Shimatani, T. et al., J. Antibiot., 2006, 59, 29-34 (isol, pmr, cmr)

Vegi, S.R. et al., Tet. Lett., 2008, 49, 6297-6299 (synth)

Antibiotic SPF 32629B

A-1267

SPF 32629B

As Antibiotic SPF 32629A, A-1266 with R = -COOH

$C_{18}H_{19}NO_6$ 345.351


Prod. by *Penicillium* sp. SPF-32629.

Specific inhibitor of human chymase.

Pale yellow powder. $[\alpha]_D^{25} +24.5$ (c. 0.4 in CHCl_3). λ_{max} 203 (ϵ 16500); 215 (sh) (ϵ 13900); 303 (ϵ 6300) (MeOH).

Shimatani, T. *et al.*, *J. Antibiot.*, 2006, **59**, 29-34 (*isol, pmr, cmr*)

Antibiotic SQ 02-S-L1 A-1268
SQ 02-S-L1




$\text{C}_{56}\text{H}_{74}\text{N}_2\text{O}_{12}$ 967.207

Related to Stachybotrin A, S-489. Prod. by *Stachybotrys* sp. RF-7260. Powder. $[\alpha]_D^{24} +131$ (c, 0.63 in MeOH). λ_{max} 218 (ϵ 64100); 261 (ϵ 14000); 301 (ϵ 4300) (MeOH).

Minagawa, K. *et al.*, *J. Antibiot.*, 2002, **55**, 239-248 (*isol, pmr, cmr*)

Antibiotic SQ 02-S-V2 A-1269
SQ 02-S-V2




$\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_4$ 426.555

Related to Stachybotrin and Stachyflin, S-492. Prod. by *Stachybotrys* sp. RF-7260. Antiviral agent. Powder (as hydrochloride). $[\alpha]_D^{24} +113.6$ (c, 0.47 in MeOH) (hydrochloride). λ_{max} 223 (ϵ 27700); 273 (ϵ 6200); 314 (ϵ 2600) (MeOH) (hydrochloride).

Minagawa, K. *et al.*, *J. Antibiot.*, 2002, **55**, 239-248 (*isol, pmr, cmr*)

Antibiotic SS 228B A-1270
SS 228B




$\text{C}_{29}\text{H}_{19}\text{NO}_7$ 493.472

Angucycline antibiotic. Prod. by a *Chaetia* sp. Blue solid.

Rohr, J. *et al.*, *Nat. Prod. Rep.*, 1992, **9**, 103-137

Antibiotic SW 163C A-1271
SW 163C




$\text{C}_{52}\text{H}_{60}\text{N}_{10}\text{O}_{14}\text{S}_2$ 1113.237

Depsipeptide antibiotic. Related to Quinomycin A, Q-58. Prod. by *Streptomyces* sp. SNA15896. Antitumour agent. Pale yellow powder. Mp 240-243°. $[\alpha]_D^{28} -78.6$ (c, 0.2 in CHCl_3). λ_{max} 214 (ϵ 78300); 230 (ϵ 77400); 299 (ϵ 9800); 359 (ϵ 9800) (MeOH).

Takahashi, K. *et al.*, *J. Antibiot.*, 2001, **54**, 615-621; 622-627 (*isol, struct*)

Nakaya, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 2969-2976 (*isol*)

Antibiotic T 988B A-1272
T 988B




$\text{C}_{25}\text{H}_{26}\text{N}_4\text{O}_4\text{S}_2$ 510.637

Prod. by *Tilachlidium* sp. (CANU-T988). Cytotoxic. Amorph. solid. $[\alpha]_D^{20} +330$ (c, 0.001 in MeOH). λ_{max} 210 (log ϵ 4.14); 232 (log ϵ 4.2); 258 (log ϵ 4.22); 292 (log ϵ 4.22) (MeOH).

Feng, Y. *et al.*, *J. Nat. Prod.*, 2004, **67**, 2090-2092 (*isol, pmr, cmr*)

Antibiotic TA 3037A A-1273
3-Benzylidene-3,4-dihydro-2-oxo-2H-1,4-benzoxazine-5-carboxylic acid. TA 3037A
[143651-44-9]



$\text{C}_{16}\text{H}_{11}\text{NO}_4$ 281.267

Prod. by *Streptomyces* sp. TA-3037. Inhibitor of glutathione S-transferase. Antiallergic agent. Yellow needles (MeOH/ CHCl_3). Sol. THF, DMF, dioxan; poorly sol. H_2O , MeOH, hexane, EtOAc. Mp 248-250°. λ_{max} 205 (ϵ 21900); 241 (ϵ 18200); 282 (ϵ 9120); 350 (sh) (ϵ 8710); 382 (ϵ 14500) (EtOH/HCl) (Derrep). λ_{max} 324 (ϵ 9120) (EtOH/NaOH) (Derrep). λ_{max} 204 (ϵ 26900); 242 (ϵ 15800); 282 (ϵ 9330); 383 (ϵ 14500) (EtOH) (Derrep).

Komagata, D. *et al.*, *J. Antibiot.*, 1992, **45**,

1117-1121; 1122-1124 (*isol, pmr, cmr, uv, ir, props*)
Kelly, T.R. *et al.*, *Heterocycles*, 1997, **45**, 87-93 (*synth*)

Antibiotic TAN 883 A-1274
TAN 883

Tetramic acid derivs. Structs. not known except probably for 883B in X-6. Prod. by *Lysobacter gummosus*. Angiogenic and antiulcer agent.

Antibiotic TAN 883A
TAN 883A

[138636-81-4]

Antibiotic TAN 883C
TAN 883C

[138636-82-5]

Antibiotic TAN 883D
TAN 883D

[138634-39-6]

$\text{C}_{29}\text{H}_{38}\text{N}_2\text{O}_5$ 494.63

Yellow cryst. $[\alpha]_D +724$ (DMF). λ_{max} 259; 340 (MeOH).

Japan. Pat., 1991, 91 178 988; CA, **116**, 57524s

Antibiotic TAN 1263 A-1275
TAN 1263

Tetramic acid derivs. Prod. by *Corollospora gracilis* IFO32111. Antifungal agent.

Antibiotic TAN 1263A

TAN 1263A

[148092-20-0]

$\text{C}_{22}\text{H}_{23}\text{NO}_6$ 397.427

Powder. Sol. MeOH, EtOAc. Mp 139-140°. $[\alpha]_D^{27} -55.1$ (c, 0.5 in MeOH). λ_{max} 254; 279 (MeOH).

Antibiotic TAN 1263B

TAN 1263B

[148092-21-1]

$\text{C}_{22}\text{H}_{25}\text{NO}_7$ 415.442

Powder. Sol. MeOH, EtOAc. $[\alpha]_D^{27} +22.4$ (c, 0.53 in MeOH). λ_{max} 253; 283 (MeOH).

Antibiotic TAN 1263C

TAN 1263C

[148092-22-2]

$\text{C}_{22}\text{H}_{25}\text{NO}_7$ 415.442

Powder. Sol. MeOH, EtOAc. Mp 93.5-95.5°. $[\alpha]_D^{22} +44.8$ (c, 0.28 in MeOH). λ_{max} 252; 282 (MeOH).

Antibiotic TAN 1263D

TAN 1263D

[148092-23-3]

$\text{C}_{22}\text{H}_{25}\text{NO}_7$ 415.442

Powder. Sol. MeOH, EtOAc. $[\alpha]_D^{22} -62.9$ (c, 0.25 in MeOH). λ_{max} 252; 282 (MeOH).

Japan. Pat., 1992, 92 330 091; CA, **119**, 15319k

Antibiotic TAN 1280 A-1276
TAN 1280

[138264-01-4]


$\text{C}_7\text{H}_5\text{N}_3\text{O}_3$ 179.135

Struct. unknown. Prod. by *Streptomyces* sp. TAN-1280. Antifungal agent. Brown

cryst. Sol. H₂O. λ_{\max} 205 (E1%/1cm 883); 239 (E1%/1cm 621); 272 (E1%/1cm 467); 312 (E1%/1cm 360); 451 (E1%/1cm 193) (pH 7 buffer) (Berdy).

Japan. Pat., 1991, 91 161 494; CA, 116, 39773t (isol)

Antibiotic TAN 1307 A-1277
 α -Amino-4-(2-chloroethenyl)-1H-pyrrole-3-propanoic acid, 9CI. TAN 1307 [137400-60-3]




C₉H₁₁ClN₂O₂ 214.651

Prod. by a *Streptomyces* sp. Antifungal agent. Powder. Sol. H₂O, MeOH; poorly sol. CHCl₃, EtOAc. [α]_D +1.3 (H₂O). λ_{\max} 210 (H₂O).

Japan. Pat., 1991, 91 157 366; CA, 115, 255901e (isol)

Antibiotic TAN 1813 A-1278
 TAN 1813 [172924-31-1]




C₂₆H₃₅NO₆ 457.566

Prod. by *Phoma* sp. FL4150. Ras-Farnesyltransferase inhibitor. Antitumour agent. Yellow powder. Sol. MeOH, DMSO, CHCl₃, EtOAc; poorly sol. H₂O, hexane. [α]_D -48.9 (MeOH). λ_{\max} 240 (€ 16500) (MeOH).

Eur. Pat., 1995, 677 513; CA, 124, 106637g
 Ishii, T. et al., *J. Antibiot.*, 2000, 53, 765-778

Antibiotic TAN 2020 A-1279
 1-(1-Carboxyethyl)-2-methylpropyl 3-[[5-[(aminoiminomethyl)amino]-1-oxopentyl]oxy]-2,4-dimethylpentanoate. TAN 2020 [177191-46-7]




C₂₀H₃₇N₃O₆ 415.529

Isol. from *Actinomadura* sp. Cell adhesion inhibitor.

Japan. Pat., 1996, 08 67 664; CA, 125, 8686y (isol)

Antibiotic TAN 876A A-1280
 2,3,8-Trichloro-6-ethyl-5-hydroxy[1]benzopyrano[2,3-b]pyrrol-4(1H)-one, 9CI. TAN 876A [119059-71-1]




C₁₃H₈Cl₃NO₃ 332.569

Pyrrrole antibiotic. Isol. from the culture broth of *Streptomyces* sp. C-70899. Shows antimicrobial activity with immunomodulating props. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 247 (E1%/1cm 1134); 357 (E1%/1cm 332) (MeOH/HCl) (Berdy). λ_{\max} 242 (E1%/1cm 830); 266 (E1%/1cm 648); 300 (E1%/1cm 540); 370 (E1%/1cm 356) (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, scu) 400 - 800 mg/kg.

Funabashi, Y. et al., *Takeda Kenkyushoho*, 1992, 51, 73; CA, 118, 55722x (isol)

Antibiotic TAN 1030A A-1281
 TAN 1030A [126221-76-9]



Absolute Configuration

C₂₇H₂₂N₄O₄ 466.495

Isol. from culture broths of *Streptomyces longisporoflavus* R19 and *Streptomyces* sp. C-71799. Possesses macrophage-activating props. Cryst. + 1H₂O. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O, hexane. Mp 290-295° dec. [α]_D²⁵ 0 (c, 0.5 in DMF). Related to Staurosporine, S-498. λ_{\max} 233 (€ 29400); 244 (sh) (€ 28000); 263 (sh) (€ 31300); 275 (sh) (€ 42000); 289 (€ 71000); 319 (sh) (€ 13400); 333 (€ 17700); 352 (€ 12100); 369 (€ 13400) (MeOH) (Derep). λ_{\max} 233 (€ 29400); 289 (€ 71000); 333 (€ 17700); 352 (€ 12100); 369 (€ 13400) (MeOH) (Berdy).

N⁶-(Methoxymethyl): [178276-02-3]

C₂₉H₂₆N₄O₅ 510.548

Prod. by *Streptomyces longisporoflavus* R19. Cryst. (CH₂Cl₂/2-propanol). Mp >300° dec.

N⁶-(Isopropoxymethyl): [178276-04-5]

C₃₁H₃₀N₄O₅ 538.602

Prod. by *Streptomyces longisporoflavus* R19. Cryst. (CH₂Cl₂/2-propanol). Mp 266-272° (dec.).

7ξ-Hydroxy: [178276-00-1]

C₂₇H₂₂N₄O₅ 482.495

Prod. by *Streptomyces longisporoflavus*. Cryst. (MeCN aq.). Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. Mp 240° (dec.).

7-Oxo: [178276-01-2]

C₂₇H₂₀N₄O₅ 480.479

Prod. by *Streptomyces longisporoflavus*. Yellow cryst. (CH₂Cl₂/2-propanol). Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. Mp 265-270° (dec.).

4'-De(hydroxyimino), 4'-oxo: [178276-05-6]

C₂₇H₂₁N₃O₄ 451.481

Prod. by *Streptomyces longisporoflavus*. Cryst. (CH₂Cl₂). Mp 236-243°.

3'-Epimer, 4'-de(hydroxyimino), 4'-oxo: [178455-10-2]

C₂₇H₂₁N₃O₄ 451.481

Prod. by *Streptomyces longisporoflavus* R19. Cryst. (CH₂Cl₂). Mp 171-176°.


Tanida, S. et al., *J. Antibiot.*, 1989, 42, 1619-1630 (isol, struct, props)

Tsubotani, S. et al., *Tetrahedron*, 1991, 47, 3565-3574 (pmr, cmr)

Cai, Y. et al., *J. Antibiot.*, 1996, 49, 519-526 (*Streptomyces longisporoflavus* isolates, abs config)

Wood, J.L. et al., *Tet. Lett.*, 1996, 37, 7335-7338 (synth)

Antibiotic TAN 1057A A-1282
 TAN 1057A [128126-44-3]



C₁₃H₂₅N₉O₃ 355.399

Prod. by *Flexibacter* spp. Active against gram-positive and -negative bacteria. Powder (as dihydrochloride). [α]_D²² -39.1 (c, 0.53 in H₂O) (dihydrochloride). λ_{\max} 215 (€ 23200); 245 (€ 11800) (H₂O) (Derep).

5-Epimer: Antibiotic TAN 1057B. TAN 1057B [128126-45-4]

C₁₃H₂₅N₉O₃ 355.399

Prod. by *Flexibacter* spp. Active against gram-positive and -negative bacteria. Powder (as dihydrochloride). [α]_D²² +72.6 (c, 0.52 in H₂O) (dihydrochloride). λ_{\max} 215 (€ 23200); 245 (€ 11800) (H₂O) (Derep).

▶ MN7880200

Katayama, N. et al., *J. Antibiot.*, 1993, 46, 606 (isol, props)

Funabashi, Y. et al., *Tetrahedron*, 1993, 49, 13 (pmr, cmr, cd)


Yuan, C. et al., *J.A.C.S.*, 1997, 119, 11777-11784 (synth)

Sokolov, V.V. et al., *Eur. J. Org. Chem.*, 1998, 777-783 (synth)

Belov, V.N. et al., *Tetrahedron*, 2004, 60, 7579-7589 (synth)

Antibiotic TAN 876B A-1283

(3-Chloro-5-ethyl-2,6-dihydroxyphenyl)(4,5-dichloro-1H-pyrrol-3-yl)methanone. 2,3-Dichloro-4-(3-chloro-5-ethyl-2,6-dihydroxybenzoyl)pyrrole. TAN 876B [119059-72-2]



C₁₃H₁₀Cl₃NO₃ 334.585


Pyrrole antibiotic. Prod. by *Streptomyces* sp. C-70899. Shows antimicrobial activity with immunomodulating props. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{max} 308 (E1%/1cm 417) (MeOH) (Berdy). λ_{max} 220 ; 308 (E1%/1cm 417) (MeOH/HCl) (Berdy). λ_{max} 321 (E1%/1cm 721) (MeOH/NaOH) (Berdy).

► LD₅₀ (mus, scu) 400 - 800 mg/kg, LD₅₀ (mus, orl) 50 mg/kg.

Funabashi, Y. et al., *Takeda Kenkyushoho*, 1992, **51**, 73; *CA*, **118**, 55722x (isol, struct)

Antibiotic TAN 1251B A-1284

TAN 1251B
[138214-46-7]



C₂₄H₃₂N₂O₃ 396.528

Incorrect name in CA. Prod. by *Penicillium thomii* RA-89. Muscarinic receptor antagonist. Hydriatic, antispasmodic and antiulcer agent. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. λ_{max} 265 (ε 22700); 304 (ε 1700) (MeOH) (Berdy).

3'-Deoxy: **Antibiotic TAN 1251A**. TAN 1251A

[138214-45-6]
[211506-06-8]

C₂₄H₃₂N₂O₂ 380.529

Prod. by *Penicillium thomii* RA-89. Muscarinic receptor antagonist.

Hydriatic, antispasmodic and antiulcer agent. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. λ_{max} 265 (ε 23800); 304 (ε 1600) (MeOH) (Berdy).

3'-Deoxy, 3,9-dihydro: **Antibiotic TAN 1251D**. TAN 1251D

[138214-48-9]

C₂₄H₃₄N₂O₂ 382.545

Prod. by *Penicillium thomii* RA-89.

3'-Deoxy, Δ^{3,4}-isomer: **Antibiotic TAN 1251C**. TAN 1251C

[138214-47-8]

C₂₄H₃₂N₂O₂ 380.529

Prod. by *Penicillium thomii* RA-89.

Pat. Coop. Treaty (WIPO), 1991, 91 13887; *CA*, **116**, 39780t (isol)

Snider, B.B. et al., *Org. Lett.*, 2000, **2**, 643-646 (synth)


Ousmer, M. et al., *J.A.C.S.*, 2001, **123**, 7534-7538 (TAN 1251C, synth)

Nagumo, S. et al., *Tetrahedron*, 2002, **58**, 4917-4924; 9871-9877 (TAN 1251A, synth)

Mizutani, H. et al., *Heterocycles*, 2004, **62**, 343-355 (TAN 1251A, synth)

Antibiotic TAN 1057C A-1285

TAN 1057C
[128126-46-5]



C₁₃H₂₅N₉O₃ 355.399

Prod. by *Flexibacter* spp. Active against gram-positive and -negative bacteria.

2-Epimer: **Antibiotic TAN 1057D**. TAN 1057D

[128126-47-6]

C₁₃H₂₅N₉O₃ 355.399


Prod. by *Flexibacter* spp. Active against gram-positive and -negative bacteria.

Katayama, N. et al., *J. Antibiot.*, 1993, **46**, 606

Yuan, C. et al., *J.A.C.S.*, 1997, **119**, 11777-11784 (synth)

Antibiotic TM 64 A-1286

2-(1-Aminoethyl)-N-[2-(1H-indol-3-yl)ethyl]-4-thiazolecarboxamide, 9CI. TM 64



(S)-form

C₁₆H₁₈N₄OS 314.41

(S)-form [58887-22-2]

Prod. by *Thermoactinomyces* sp. TM-64. Sol. acids, MeOH, CHCl₃; fairly sol. C₆H₆; poorly sol. H₂O. Mp 120° (145-147°). [α]_D²⁰ -6 (c, 1 in MeOH). [α]_D -8.3 (c, 1 in EtOH). λ_{max} 223 (ε 51750); 275 (ε 10500); 282 (ε 10500); 291 (ε 8350) (MeOH).

1''-N-Ac: **Microbiaeratin**

[58887-23-3]

[1020001-70-0]

C₁₈H₂₀N₄O₂S 356.448

Prod. by the marine-derived *Microbispora aerata* IMBAS-11A and from *Thermoactinomyces* sp. TA66-2. Powder. λ_{max} 224 ; 282 ; 290 (MeOH).

(ξ)-form

1''-N-Propanoyl: [1020001-66-4]

C₁₉H₂₂N₄O₂S 370.474

Prod. by *Thermoactinomyces* sp. TA66-2. Powder.

Omura, S. et al., *J. Antibiot.*, 1975, **28**, 609-610 (isol)

Konda, Y. et al., *Chem. Pharm. Bull.*, 1976, **24**, 92-96 (uv, ir, pmr, ms, struct)


Onda, M. et al., *Chem. Pharm. Bull.*, 1978, **26**, 2167-2169 (synth)

Ivanova, V. et al., *Prep. Biochem. Biotechnol.*, 2007, **37**, 161-168 (*Microbiaeratin*)

Korkmaz, C.A. et al., *Magn. Reson. Chem.*, 2008, **46**, 80-83 (*N-propanoyl*)

Antibiotic TMC 52 A-1287

TMC 52



TMC 52A, m = 3, n = 4

TMC 52B, m = 4, n = 3

Similar to Cathestatin B, C-208. Prod. by a *Gliocladium* sp. (fungal strain F-2665). Cysteine proteinase inhibitors.

Antibiotic TMC 52A

TMC 52A

[213417-23-3]

C₂₀H₃₀N₄O₆ 422.48

Sol. H₂O, MeOH, DMSO, EtOH; poorly sol. CHCl₃, EtOAc. [α]_D²⁴ +22 (c, 0.4 in H₂O). λ_{max} 275 (ε 1200); 281 (sh) (ε 1000) (H₂O). λ_{max} 275 (ε 1200); 281 (ε 1000) (H₂O) (Berdy).

4'-Deoxy: **Antibiotic TMC 52C**. TMC 52C

[213417-24-4]

C₂₀H₃₀N₄O₅ 406.481

Sol. H₂O, DMSO, MeOH, EtOH; poorly sol. CHCl₃, EtOAc. [α]_D²⁴ +17 (c, 0.5 in H₂O). λ_{max} 251 (ε 180); 257 (ε 210); 263 (ε 170) (H₂O). λ_{max} 251 (ε 180); 257 (ε 210); 263 (ε 170) (H₂O) (Berdy).

Antibiotic TMC 52B

TMC 52B

[213330-57-5]

C₂₀H₃₀N₄O₆ 422.48

Sol. H₂O, MeOH, EtOH, DMSO; poorly sol. CHCl₃, EtOAc. [α]_D²⁴ +22 (c, 0.4 in H₂O). λ_{max} 275 (ε 1200); 281 (sh) (ε 1000) (H₂O). λ_{max} 275 (ε 1200); 281 (ε 1000) (H₂O) (Berdy).

4'-Deoxy: **Antibiotic TMC 52D**. TMC 52D

[213417-25-5]

C₂₀H₃₀N₄O₅ 406.481


Sol. H₂O, DMSO, EtOH, MeOH; poorly sol. CHCl₃, EtOAc. [α]_D²⁴ +11 (c, 0.5 in H₂O). λ_{max} 251 (ε 180); 257 (ε 210); 263 (ε 170) (H₂O). λ_{max} 251 (ε 180); 257 (ε 210); 263 (ε 170) (H₂O) (Berdy).

Isshiki, K. et al., *J. Antibiot.*, 1998, **51**, 629-634 (isol, uv, ir, pmr, cmr)

Antibiotic TMC 66 A-1288

TMC 66

[245660-81-5]



Absolute Configuration

C₂₉H₂₁NO₉ 527.486


Related to Pradimicin A. Prod. by *Streptomyces* sp. A5008. Endothelin converting enzyme inhibitor. Reddish-brown powder. Mp 215-220° (dec.). $[\alpha]_D^{24}$ -327 (c, 0.01 in CHCl₃). λ_{\max} 215 (ε 27400); 245 (ε 28600); 310 (sh) (ε 11700); 365 (ε 4400); 465 (ε 10500) (MeOH).

Asai, Y. *et al.*, *J. Antibiot.*, 1999, **52**, 607-612 (isol, uv, ir, pmr, cmr)

Hosokawa, S. *et al.*, *Tet. Lett.*, 2007, **48**, 7305-7308 (synth, abs config)

Antibiotic TMC 69
TMC 69

A-1289

C₂₆H₃₁NO₄ 421.535


Prod. by *Chrysosporium* sp. TC 1068. Antitumour agent. Amorph. powder. Related to Funiculosin, F-197 and Sambutoxin, S-48. λ_{\max} 245 (sh) (log ε 4.54); 262 (log ε 4.69); 275 (log ε 4.76); 287 (log ε 4.63) (MeOH).

Hirano, N. *et al.*, *J. Antibiot.*, 2001, **54**, 421-427 (isol, activity)

Kohno, J. *et al.*, *Tetrahedron*, 2001, **57**, 1731-1735 (isol, pmr, cmr)

Antibiotic TMC 260
TMC 260

A-1290

C₁₇H₂₇NO₅ 325.404


Prod. by *Acremonium kiliense* Grütz TC 1703. IL-4 signal transduction inhibitor. Powder (EtOAc/hexane). Mp 140-144°. $[\alpha]_D^{20}$ -29 (c, 0.096 in MeOH). λ_{\max} 245 (ε 14050) (MeOH).

Sakurai, M. *et al.*, *J. Antibiot.*, 2003, **56**, 787-791 (isol, pmr, cmr)

Antibiotic TMC 1A

A-1291

TMC 1A
[177079-60-6]

C₂₈H₃₆N₂O₇ 512.602

Prod. by *Streptomyces* sp. A-230. Cytotoxic agent. Yellow powder. Sol. MeOH, Me₂CO, butanol, EtOAc, DMSO, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{24}$ -55

(c, 0.1 in CHCl₃). Mp >95° dec. Related to Manumycin A, M-87. λ_{\max} 262 (ε 38800); 304 (ε 43400) (MeOH) (Berdy).


Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1212-1220 (isol, uv, ir, pmr, cmr)

Cronje Grove, J.J. *et al.*, *Chem. Comm.*, 1999, 421-422 (synth)

Antibiotic TMC 2A

A-1292

TMC 2A
[195976-77-3]

C₂₈H₃₄N₄O₉ 570.598

Prod. by *Aspergillus oryzae*. Dipeptidyl peptidase IV inhibitor. Powder. Mp 166-168° (dec.). $[\alpha]_D^{24}$ +2.4 (c, 0.2 in H₂O). λ_{\max} 212 (ε 9300); 272 (sh) (ε 1550); 280 (ε 1800); 288 (ε 1350) (MeOH).

5-Deoxy: **Antibiotic TMC 2B**. TMC 2B
[196212-08-5]

C₂₈H₃₄N₄O₈ 554.599
Prod. by *Aspergillus oryzae*. Dipeptidyl peptidase IV inhibitor. Powder. Mp 168-169° (dec.). $[\alpha]_D$ +11.4 (c, 0.1 in H₂O). λ_{\max} 212 (ε 9200); 272 (sh) (ε 1500); 280 (ε 1650); 288 (ε 1200) (MeOH).

4-Epimer, 5-deoxy: **Antibiotic TMC 2C**. TMC 2C

C₂₈H₃₄N₄O₈ 554.599
Prod. by *Aspergillus oryzae*. Dipeptidyl peptidase IV inhibitor. Powder. Mp 175-181° (dec.). $[\alpha]_D^{24}$ -17.5 (c, 0.1 in MeOH). λ_{\max} 212 (ε 10400); 272 (sh) (ε 1650); 280 (ε 1750); 288 (ε 1300) (MeOH).


Nonaka, N. *et al.*, *J. Antibiot.*, 1997, **50**, 646-652; 653-658 (isol, uv, ir, pmr, cmr, cryst struct)

Ohnuki, T. *et al.*, *Drugs of the Future*, 1999, **24**, 665-670 (rev)

Antibiotic TMC 95A

A-1293

TMC 95A
[220666-21-7]

C₃₃H₃₈N₆O₁₀ 678.697

Cyclic peptide antibiotic. Prod. by *Apiospora montagnei* Sacc. TC 1093. Proteasome inhibitor. Cytotoxic agent. Powder. $[\alpha]_D^{23}$ +102 (c, 0.54 in MeOH). λ_{\max} 206 (log ε 4.59); 222 (log ε 4.63); 294 (log ε 3.74) (MeOH).

7-Epimer: **Antibiotic TMC 95C**. TMC 95C

[262854-07-9]
C₃₃H₃₈N₆O₁₀ 678.697

Prod. by *Apiospora montagnei*. Proteasome inhibitor. Powder. $[\alpha]_D^{23}$ -18 (c, 0.23 in MeOH). λ_{\max} 207 (log ε 4.55); 222 (log ε 4.6); 295 (log ε 3.72) (MeOH).

36-Epimer: **Antibiotic TMC 95B**. TMC 95B

[262854-05-7]
C₃₃H₃₈N₆O₁₀ 678.697

Prod. by *Apiospora montagnei*. Proteasome inhibitor. Powder. $[\alpha]_D^{23}$ +74 (c, 0.47 in MeOH). λ_{\max} 206 (log ε 4.6); 222 (log ε 4.62); 295 (log ε 3.73) (MeOH).

7,36-Diepimer: **Antibiotic TMC 95D**. TMC 95D

[262854-09-1]
C₃₃H₃₈N₆O₁₀ 678.697

Prod. by *Apiospora montagnei*. Proteasome inhibitor.

Koguchi, Y. *et al.*, *J. Antibiot.*, 2000, **53**, 105-109 (isol, activity)

Kohno, J. *et al.*, *J.O.C.*, 2000, **65**, 990-995 (uv, ir, pmr, cmr, struct)

Ma, D. *et al.*, *Tet. Lett.*, 2000, **41**, 9089-9093 (synth)

Inoue, M. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2654-2657 (synth)

Inoue, M. *et al.*, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 663-665 (TMC-95A, pharmacol)

Kaiser, M. *et al.*, *ChemBioChem*, 2004, **5**, 1256-1266 (TMC-95A, pharmacol)


Lin, S. *et al.*, *J.A.C.S.*, 2004, **126**, 6347-6355 (synth)

Albrecht, B.K. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 11949-11954 (synth)

Antibiotic TMC 120A

A-1294

2,3-Dihydro-7-methyl-2-(1-methylethyl)-furo[3,2-h]isoquinolin-3-one, 9CI. TMC 120A
[250231-80-2]

C₁₅H₁₅NO₂ 241.289

Prod. by *Aspergillus ustus*. Pale yellow solid. Mp 115-116°. $[\alpha]_D^{23}$ -18 (c, 0.5 in MeOH). λ_{\max} 210 (log ε 4.5); 250 (log ε 4.53); 323 (sh) (log ε 3.85); 340 (log ε 4.05); 354 (log ε 4.11) (MeOH).

2,11-Didehydro: **Antibiotic TMC 120B**. TMC 120B

[250231-81-3]
C₁₅H₁₃NO₂ 239.273

Prod. by *Aspergillus ustus* and the

marine *Aspergillus pseudodeflectus*. Moderate inhibitor of eosinophil survival. Pale yellow needles. Mp 176–177°. λ_{\max} 212 (log ϵ 4.54); 233 (log ϵ 4.11); 239 (log ϵ 4.11); 271 (log ϵ 4.37); 295 (sh) (log ϵ 4.33); 302 (log ϵ 4.33); 365 (log ϵ 3.89) (MeOH).

2-Hydroxy: Antibiotic TMC 120C. TMC 120C

[250231-82-4]

$C_{15}H_{15}NO_3$ 257.288

Prod. by *Aspergillus ustus*. Pale yellow needles. Mp 176–177° dec. Possibly racemic. λ_{\max} 211 (log ϵ 4.51); 253 (log ϵ 4.5); 354 (log ϵ 4.01); 413 (sh) (log ϵ 3.05) (MeOH).

Kohno, J. *et al.*, *J. Antibiot.*, 1999, **52**, 913-916 (*isol, activity*)

Kohno, J. *et al.*, *Tetrahedron*, 1999, **55**, 11247-11252 (*uv, ir, pmr, cmr, TMC 120B, TMC 120C, cryst struct*)


Ogawa, A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 3539-3543 (*TMC 120B, isol, pmr, cmr*)

Kumemura, T. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 393-397 (*TMC 120B, synth*)

Antibiotic TMC 135A

A-1295

TMC 135A



$C_{39}H_{49}N_3O_8S$ 719.897


Ansamycin-type antibiotic. Related to Thiazinotrienomycin A. Prod. by *Streptomyces* sp. TC 1190. Cytotoxic agent. Pale yellow powder. Mp 166–168°. λ_{\max} 216 (log ϵ 4.42); 259 (log ϵ 4.5); 270 (log ϵ 4.46); 282 (log ϵ 4.34); 310 (log ϵ 3.33) (MeOH).

Nishio, M. *et al.*, *J. Antibiot.*, 2000, **53**, 724-727

Antibiotic TMC 135B

A-1296

TMC 135B



$C_{39}H_{49}N_3O_8S$ 719.897


Ansamycin antibiotic. Related to Thiazinotrienomycin C. Prod. by *Streptomyces* sp. TC 1190. Cytotoxic agent. Pale yellow powder. Mp 190–192°. λ_{\max} 217 (log ϵ 4.46); 259 (log ϵ 4.63); 270 (log ϵ 4.58); 281 (log ϵ 4.43); 310 (log ϵ 3.56) (MeOH).

Nishio, M. *et al.*, *J. Antibiot.*, 2000, **53**, 724-727

Antibiotic TMC 1C

A-1297

TMC 1C
[177079-62-8]



$C_{30}H_{38}N_2O_7$ 538.639

Prod. by *Streptomyces* sp. A-230. Cytotoxic agent. Yellow powder. Sol. MeOH, Me₂CO, butanol, EtOAc, DMSO, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{24} +116$ (c, 0.1 in CHCl₃). Mp >106° dec. Related to Manumycin A, M-87. λ_{\max} 265 (sh) (ϵ 49500); 285 (ϵ 52700); 306 (sh) (ϵ 47800) (MeOH).

4',5'-Dihydro: Antibiotic TMC 1D. TMC 1D

[177079-63-9]

$C_{30}H_{40}N_2O_7$ 540.655

Prod. by *Streptomyces* sp. A-230.

Cytotoxic agent. Yellow powder. Sol. MeOH, DMSO, Me₂CO, EtOAc, CHCl₃, butanol; poorly sol. H₂O, hexane. $[\alpha]_D^{24} +16$ (c, 0.1 in CHCl₃). Mp >89° dec. λ_{\max} 263 (ϵ 40700); 278 (ϵ 40200); 304 (ϵ 42700) (MeOH).


Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1212-1220 (*isol, uv, ir, pmr, cmr*)

Antibiotic U 62162

A-1298

2-Hydroxy-5-oxo-4-[(4,6,8-trimethyl-1-oxo-2,7-nonadienyl)amino]-7-oxabicyclo[4.1.0]hept-3-ene-2-pentanoic acid. U 62162

[82516-67-4]



$C_{23}H_{33}NO_6$ 419.517

Isol. from *Streptomyces verdensis*. Active *in vitro* against gram-positive bacteria esp. *Staphylococcus aureus*. Needles (EtOAc/petrol). Sol. MeOH, toluene, DMF, DMSO, C₆H₆; poorly sol. H₂O, hexane. Mp 96–98°. $[\alpha]_D^{25} +74$ (c, 0.97 in Me₂CO). λ_{\max} 235 (ϵ 14000); 276 (sh) (ϵ 8150) (no solvent reported) (Derep). λ_{\max} 235 (ϵ 14000); 276 (ϵ 8150) (MeOH) (Berdy). λ_{\max} 232 (ϵ 14750); 276 (ϵ 8100) (MeOH-HCl) (Berdy).


Slechtla, L. *et al.*, *J. Antibiot.*, 1982, **35**, 556
Alcaraz, L. *et al.*, *Chem. Comm.*, 1998, 1157-1158 (*synth*)

Antibiotic U 106305

A-1299

U 106305

[170591-54-5]



$C_{28}H_{41}NO$ 407.638

Prod. by *Streptomyces* sp. UC 11136.

Potent inhibitor of the cholesteryl ester transfer protein. $[\alpha]_D^{20} -270$ (c, 0.3 in CHCl₃). Similar to Jawsamycin. λ_{\max} 215; 240 (MeOH) (Berdy).

Kuo, M.S. *et al.*, *J.A.C.S.*, 1995, **117**, 10629-10634 (*isol, pmr, cmr, struct*)

Barrett, A.G.M. *et al.*, *J.A.C.S.*, 1996, **118**, 7863-7864 (*synth, config*)

Charette, A.B. *et al.*, *J.A.C.S.*, 1996, **118**, 10327-10328 (*synth*)


Pietruszka, J. *et al.*, *Chem. Rev.*, 2003, **103**, 1051-1070 (*rev*)

Antibiotic UCH 15A

A-1300

UCH 15A

[189080-82-8]



$C_{23}H_{29}N_{11}O_5$ 539.553

Oligopeptide antibiotic. Related to Anthelvincin A, A-1077. Prod. by *Streptomyces* sp. UCH15. Antibacterial and antitumour agent. Powder. Sol. H₂O, DMSO, MeOH; poorly sol. CHCl₃, λ_{\max} 233 (ϵ 26300); 297 (ϵ 29600) (MeOH).

N^d-Hydroxy: Antibiotic UCH 15B. UCH 15B

[189080-83-9]

$C_{23}H_{29}N_{11}O_6$ 555.552

Prod. by *Streptomyces* sp. UCH15.

Antibacterial and antitumour agent. Powder. Sol. H₂O, DMSO, MeOH; poorly sol. CHCl₃. λ_{\max} 232 (ϵ 27000); 297 (ϵ 32500) (MeOH).


Pat. Coop. Treaty (WIPO), 1997, 97 10 208; CA, **126**, 292503z (*isol*)

Antibiotic UCS 1025B

A-1301

UCS 1025B

[209126-30-7]



Absolute Configuration

$C_{20}H_{25}NO_6$ 375.421

Prod. by *Acremonium* sp. KY4917. Needles (hexane/EtOAc). Mp 221–223°. $[\alpha]_D^{28} -31.8$ (c, 0.1 in MeOH).

2-Deoxy: Antibiotic UCS 1025A. UCS 1025A

[209126-31-8]

$C_{20}H_{25}NO_5$ 359.421

Prod. by *Acremonium* sp. KY4917.

Telomerase inhibitor. Antibacterial and antitumour agent. Needles (hexane/EtOAc). Mp 135-137°. $[\alpha]_D^{28} +30.1$ (c, 0.1 in MeOH). λ_{\max} 260 (ε 7100) (MeOH).


Nakai, R. *et al.*, *J. Antibiot.*, 2000, **53**, 294-296 (isol, pmr, cmr)

Agatsuma, T. *et al.*, *Org. Lett.*, 2002, **4**, 4387-4390 (cryst struct, abs config)

Lambert, T.H. *et al.*, *J.A.C.S.*, 2006, **128**, 426-427 (UCS 1025A, synth)

Hoye, T.R. *et al.*, *J.A.C.S.*, 2006, **128**, 2550-2551 (UCS 1025A, synth)

Antibiotic UK 1 A-1302
UK 1
[151271-53-3]



$C_{22}H_{14}N_2O_5$ 386.363

Prod. by a close relative of *Streptomyces morookaense*. Mg ion-dependent DNA binding agent and inhibitor of human topoisomerase II. Cytotoxic. Needles (MeOH). Sol. $CHCl_3$, C_6H_6 , Py, CH_2Cl_2 ; fairly sol. MeOH, EtOAc, hexane, DMF; poorly sol. H_2O . Mp 217-219°. λ_{\max} 249 (ε 21000); 263 (ε 20200); 271 (ε 19800); 314 (ε 29000); 325 (ε 32500); 348 (ε 27800); 363 (ε 18700); 420 (ε 1480) (MeOH) (Derep).

Parent acid: Demethyl Antibiotic UK 1.

Demethyl UK 1

$C_{21}H_{12}N_2O_5$ 372.336


Prod. by *Actinomyces* sp. Stamm K 17/9. Yellow solid. λ_{\max} 233 (log ε 4.09); 264 (sh); 270 (log ε 4); 308 (sh); 324 (log ε 4.15); 340 (log ε 4.19); 415 (log ε 3.92) (MeOH).

Ueki, M. *et al.*, *J. Antibiot.*, 1993, **46**, 1089-1094; 1095-1100 (isol, uv, pmr, cmr)

Reynolds, M.B. *et al.*, *Bioorg. Chem.*, 1999, **27**, 326-337 (activity)

Strösch, K. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (Demethyl UK1)

Antibiotic UK 2 A-1303
UK 2



UK 2A R = -CH(CH₃)₂

UK 2B R = -C(CH₃)=CHCH₃(E-)

UK 2C R = -CH₂CH(CH₃)₂

UK 2D R = -CH(CH₃)CH₂CH₃

Prod. by *Streptomyces morookaense*.

Antifungal agent.

Antibiotic UK 2A

UK 2A

[167173-85-5]

$C_{26}H_{30}N_2O_9$ 514.531

Needles. Mp 207-209°. $[\alpha]_D^{23} +89.1$ (c, 0.8 in $CHCl_3$).

Demethoxy: **Antibiotic UK 3A**. UK 3A

[194931-82-3]

$C_{25}H_{28}N_2O_8$ 484.505

Prod. by *Streptomyces* sp. 517-02.

Antifungal agent. Needles. Stereochem. not confirmed.

Antibiotic UK 2B

UK 2B

[167173-86-6]

$C_{27}H_{30}N_2O_9$ 526.542

$[\alpha]_D^{17} +87.5$ (c, 0.3 in $CHCl_3$).

Antibiotic UK 2C

UK 2C

[167173-87-7]

$C_{27}H_{32}N_2O_9$ 528.558

Inseparable mixture with UK 2D.

Antibiotic UK 2D

UK 2D

[167173-88-8]

$C_{27}H_{32}N_2O_9$ 528.558

Inseparable mixture with UK 2C.

Ueki, M. *et al.*, *J. Antibiot.*, 1996, **49**, 639; 1997, **50**, 1052 (isol, activity)


Hanafi, M. *et al.*, *J. Antibiot.*, 1996, **49**, 1226 (pmr, cmr, struct)

Ueki, M. *et al.*, *J. Antibiot.*, 1997, **50**, 551 (UK 3A)

Shibata, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1113-1116 (abs config, conformm)

Shimano, M. *et al.*, *Tetrahedron*, 1998, **54**, 12745-12774 (synth, abs config)

Antibiotic UK 63052 A-1304



UK 63052 R = -CH(CH₃)CH₂CH₃

UK 63598 R = -CH₃

UK 65662 R = -CH(CH₃)₂

UK 163E R = -CH₂CH₃

Depsipeptide antibiotics. Related to Quinomycin A, Q-58. Isol. from *Streptomyces braegensis* ssp. *japonicus*. Several uncharacterised components are also prod.

Antibiotic UK 63052

UK 63052. L 52-71A. SW 163G. Anti-

biotic L 52-71A. Antibiotic SW 163G

[120763-23-7]

$C_{56}H_{68}N_{10}O_{14}S_2$ 1169.344

Also from *Streptomyces* sp. SNA15896. Off-white solid + 1H₂O. Sol. Py, DMSO, CH_2Cl_2 , $CHCl_3$; fairly sol. MeOH, EtOAc; poorly sol. H_2O , hexane. Mp 252-255°. $[\alpha]_D -166.2$ (c, 0.1 in $CHCl_3$).

λ_{\max} 223 (ε 65600); 247 (ε 82000); 399 (ε 11200) (MeOH/NaOH) (Derep). λ_{\max}

219 (ε 77300); 231 (ε 78300); 300 (ε 9400); 306 (ε 9200); 359 (ε 10200) (MeOH) (Derep).

Antibiotic UK 63598

UK 63598. SW 163D. Antibiotic SW 163D

[120832-02-2]

$C_{53}H_{62}N_{10}O_{14}S_2$ 1127.263

Also from *Streptomyces* sp. SNA15896. Off-white solid + 1H₂O. Sol. Py, CH_2Cl_2 , $CHCl_3$, DMSO; fairly sol. MeOH, EtOAc; poorly sol. H_2O , hexane. Mp 235-241°. $[\alpha]_D -126.8$ (c, 0.1 in $CHCl_3$). λ_{\max} 223 (ε 67800); 247 (ε 80600); 399 (ε 11500) (MeOH/NaOH) (Derep). λ_{\max} 219 (ε 75100); 231 (ε 76700); 300 (ε 9750); 308 (ε 9600); 359 (ε 10500) (MeOH) (Derep).

Antibiotic UK 65662

UK 65662. L 52-71B. SW 163F. Antibiotic L 52-71B. Antibiotic SW 163F

[120796-23-8]

$C_{55}H_{66}N_{10}O_{14}S_2$ 1155.317

Also from *Streptomyces* sp. SNA15896. Sol. Py, $CHCl_3$, DMSO, CH_2Cl_2 ; fairly sol. MeOH, EtOAc; poorly sol. H_2O , hexane. Mp 240-244°. $[\alpha]_D -168.1$ (c, 0.1 in $CHCl_3$). λ_{\max} 223 (ε 65600); 247 (ε 82000); 399 (ε 11200) (MeOH/NaOH) (Derep). λ_{\max} 219 (ε 77300); 231 (ε 78300); 300 (ε 9400); 306 (ε 9200); 359 (ε 10200) (MeOH) (Derep).

Antibiotic UK 163E

SW 163E. Antibiotic SW 163E. UK 163E

$C_{54}H_{64}N_{10}O_{14}S_2$ 1141.29

Prod. by *Streptomyces* sp. SNA15896.

Antitumour agent. Pale yellow needles.

Mp 234-237°. $[\alpha]_D^{28} -157.1$ (c, 0.2 in $CHCl_3$). λ_{\max} 214 (ε 70500); 230 (ε 66200); 299 (ε 9800); 359 (ε 9200) (MeOH).

Range, M.J. *et al.*, *J. Antibiot.*, 1989, **42**, 206-217 (isol, struct, props)


Takahashi, K. *et al.*, *J. Antibiot.*, 2001, **54**, 615-621; 622-627 (SW 163E)

Nakaya, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 2969-2976 (SW 163)

Antibiotic USF 19A A-1305

USF 19A

[165337-87-1]



$C_{22}H_{28}N_2O_9$ 464.471

Prod. by *Streptomyces* sp. USF-19. Li-


poxxygenase inhibitor. Amorph. powder.

Mp 180°. Similar to Antimycin A. λ_{\max} 240 (ε 1850); 302 (ε 1510) (EtOH). λ_{\max} 240 (ε 1853); 302 (ε 1510) (EtOH) (Berdy).

Komoda, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 924-926 (isol, pmr, cmr, uv, ir)

Antibiotic VM 55595 A-1306

VM 55595
[138797-22-5]




C₂₇H₃₃N₅O₃ 447.576
Prod. by a *Penicillium* sp. IMI 332995.
Mycotoxin, antiparasitic agent. Related to a Paraherquamide A, P-90. λ_{max} 245 (MeOH) (Berdy).

Blanchflower, S.E. *et al.*, *J. Antibiot.*, 1993, **46**, 1355 (isol)

Antibiotic VM 55599 A-1307

VM 55599
[148717-88-8]



Absolute Configuration

C₂₂H₂₇N₃O 349.475
Shunt metab. of *Penicillium* sp. IMI33995 and *Penicillium fellutanum*. Related to Paraherquamide A, P-90. λ_{max} 226 (ε 29700); 280 (ε 5600) (MeOH) (Derep).

Blanchflower, S.E. *et al.*, *J. Antibiot.*, 1993, **46**, 1355-1363 (isol, cryst struct)


Sanz-Cervera, J.F. *et al.*, *J.A.C.S.*, 2002, **124**, 2556-2559 (synth, abs config)

Domingo, L.R. *et al.*, *J.O.C.*, 2003, **68**, 2895-2902 (biosynth)

Ding, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1574-1578 (isol, biosynth, ms)

Antibiotic WF 14865 A-1308

WF 14865



WF 14865A R¹ = CH₃, R² = CH₂CH₃
WF 14865B R¹ = H, R² = CH(CH₃)₂

Prod. by *Aphanoascus fulvescens* and *Anixiopsis stercoraria*.

Antibiotic WF 14865A

WF 14865A
[178439-23-1]

C₁₆H₂₅N₅O₅ 367.404
Cathepsin B inhibitor, shows antiarthritic props. Powder. Sol. MeOH, H₂O; fairly sol. Me₂CO; poorly sol. CHCl₃. [α]_D +30 (MeOH). λ_{max} 200 (MeOH).

Antibiotic WF 14865B

WF 14865B
[178439-24-2]


C₁₆H₂₅N₅O₅ 367.404
Powder. Sol. MeOH, H₂O; fairly sol. Me₂CO; poorly sol. CHCl₃. [α]_D +35 (MeOH). λ_{max} 200 (MeOH).

Japan. Pat., 1996, 96 119 983; *CA*, **125**, 86398v (isol)

Otsuka, T. *et al.*, *J. Antibiot.*, 2000, **53**, 449-458 (isol, activity)

Antibiotic WF 14861A A-1309

WF 14861A
[176777-37-0]



Absolute Configuration

C₂₀H₃₀N₄O₆ 422.48

Prod. by *Colletotrichum* sp. WF 14861A. Inhibitor of Cathepsins B and L. Amorph. powder. Sol. H₂O, MeOH; poorly sol. CHCl₃, EtOAc. [α]_D +43 (c, 0.5 in MeOH).

4'-Deoxy: **Antibiotic 460B**. 460B

[207442-33-9]

C₂₀H₃₀N₄O₅ 406.481

Prod. by *Aspergillus* sp. Cathepsin L inhibitor.

Japan. Pat., 1996, 96 34 792; *CA*, **124**, 341065c

Japan. Pat., 1998, 98 101 662; *CA*, **129**, 3915v (Antibiotic 460B)


Otsuka, T. *et al.*, *J. Antibiot.*, 1999, **52**, 536-541; 542-547 (isol, pmr, cmr, ms, activity)

Detterbeck, R. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 222-232 (synth, abs config)

Antibiotic WIN 64745 A-1310

WIN 64745. Q 20547C. Antibiotic Q

[150881-28-0]



Absolute Configuration

R = -CH₂CH(CH₃)₂

C₃₇H₃₈N₆O₄ 630.745

Isol. from *Aspergillus* sp. ATCC74177. Neurokinin antagonist. Solid. Mp 194-196°. [α]_D +280 (c, 0.012 in MeOH). Related to Chaetocin, C-320. λ_{max} 242 (ε 10000); 301 (ε 4500) (MeOH).

Barrow, C.J. *et al.*, *J.O.C.*, 1993, **58**, 6016-6021 (isol, uv, ir, pmr, cmr, cd, struct)

Sedlock, D.M.J. *et al.*, *J. Antibiot.*, 1994, **47**, 391-398; 399-410; 411-419 (isol, biochem, biosynth)

Japan. Pat., 1995, 258 266; *CA*, **124**, 28114t (Q 20547C)

Pérez-Balado, C. *et al.*, *Org. Lett.*, 2008, **10**, 3701-3704 (synth)

Antibiotic WIN 64821 A-1311

WIN 64821. Q 20547A. Antibiotic Q

20547A

[150881-27-9]

As Antibiotic WIN 64745, A-1310 with

R = CH₂Ph

C₄₀H₃₆N₆O₄ 664.762

Isol. from *Aspergillus* sp. SC 319. Competitive antagonist to substance P at the human NK1 receptor. Solid. Mp 203-205°. [α]_D +200 (c, 0.15 in MeOH). Related to Chaetocin, C-320. λ_{max} 241 (ε 12100); 301 (ε 5300) (MeOH).

Barrow, C.J. *et al.*, *J.O.C.*, 1993, **58**, 6016-6021 (isol, uv, ir, pmr, cmr, cd, struct)

Sedlock, D.M. *et al.*, *J. Antibiot.*, 1994, **47**, 391-398; 399-410; 411-419 (R = CH₂Ph, isol, biochem, biosynthetic analogues, pharmacol)

Barrow, C.J. *et al.*, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 377-380 (sar)

Japan. Pat., 1995, 258 266; *CA*, **124**, 28114t (Q 20547A)

Overman, L.E. *et al.*, *J.A.C.S.*, 2001, **123**, 9465-9467 (synth)


Movassaghi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1485-1487 (synth)

Pérez-Balado, C. *et al.*, *Org. Lett.*, 2008, **10**, 3701-3704 (synth)

Antibiotic WJ 35 A-1312

WJ 35

[894085-53-1]



C₁₉H₁₇NO₈ 387.345

Prod. by engineered *Streptomyces coelicolor* strain CH999.


Zhang, W. *et al.*, *Appl. Environ. Microbiol.*, 2006, **72**, 2573-2580 (isol, biosynth, pmr, cmr)

Antibiotic WJ 85 A-1313

4-Hydroxy-6-(1,7,9,10-tetrahydroxybenz[*g*]isoquinolin-3-yl)-2H-pyran-2-one.

WJ 85

[917571-64-3]



C₁₉H₁₃NO₇ 367.314

Prod. by *Streptomyces coelicolor* strain CH999. Pale yellow solid. λ_{max} 262 (sh) (log ε 4.05); 293 (log ε 4.22); 376 (log ε 3.45) (MeCN).

5-Hydroxy, 5,10-quinone: **Antibiotic WJ 85B**. WJ 85B

[917571-67-6]

C₁₉H₁₁NO₈ 381.298

Prod. by *Streptomyces coelicolor* CH999. Yellow solid. λ_{max} 248 (log ε 3.36); 279 (sh) (log ε 3.35); 322 (log ε 3.46); 431 (log ε 2.99) (MeCN).

Zhang, W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1633-1636 (*isol, pmr, cmr*)



Antibiotic WS 1921 A-1314
WS 1921
[12704-00-6]

C₈H₁₀N₂O₅S 214.312

Pyrothine-related antibiotic. Struct. unknown. Prod. by *Streptomyces griseolus-sulfoantibioticus*. Antibacterial agent. Orange-yellow cryst. Sol. MeOH, CHCl₃; fairly sol. C₆H₆; poorly sol. hexane, H₂O. Mp 191-193° (dec.). λ_{max} 230 (E1%/1cm 450); 370 (E1%/1cm 1150) (MeOH) (Berdy).

Japan. Pat., 1972, 72 00 679; *CA*, **76**, 125429x (*isol*)

Antibiotic WS 75624A A-1315
WS 75624A
[170663-44-2]



C₁₈H₂₄N₂O₅S 380.464

Prod. by *Saccharothrix* sp. 75624. Endothelin converting enzyme inhibitor; Collagenase:Nep II processing protease inhibitor. Powder. Sol. MeOH, CHCl₃, Me₂CO; fairly sol. EtOAc; poorly sol. H₂O. Mp 53-54°. [α]_D²³ 0 (c, 0.5 in MeOH). λ_{max} 215; 245; 298 (MeOH). λ_{max} 215; 242; 313 (MeOH/HCl) (Berdy). λ_{max} 215; 238; 290; 303 (MeOH/NaOH).

3-Hydroxy, amide: **Karnamicin D₁**, BU 3557D₁. Antibiotic BU 3557D₁
[122535-59-5]

C₁₈H₂₅N₃O₅S 395.479

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, C₆H₆, EtOH, CHCl₃, EtOAc, DMSO; poorly sol. hexane, H₂O. Mp 146.5-148°. [α]_D²⁵ 0 (c, 1 in CHCl₃). λ_{max} 219 (ε 26300); 242 (ε 23500); 317 (ε 6800) (MeOH). λ_{max} 220 (sh) (ε 23300); 243 (ε 26400); 317 (ε 8000) (MeOH/HCl). λ_{max} 221 (ε 22900); 252 (ε 20800); 292 (ε 16400); 342 (ε 8800) (MeOH/NaOH).

1'',3-Dihydroxy, amide: **Karnamicin A₃**, BU 3557A₃. Antibiotic BU 3557A₃
[122535-50-6]

C₁₈H₂₅N₃O₆S 411.478

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder + ½H₂O. Sol. MeOH, EtOH, CHCl₃, EtOAc, DMSO, C₆H₆; poorly sol. hexane, H₂O. Mp 69-72°. [α]_D²⁵ -19 (c, 0.25 in CHCl₃). λ_{max} 218 (ε 26400); 242 (ε 23700); 317 (ε 6800) (MeOH). λ_{max} 220 (ε 25000); 243 (ε 25000); 317 (ε 7300) (MeOH/HCl). λ_{max} 220 (ε 24200); 251 (ε 22200); 282 (ε 17400);

342 (ε 9200) (MeOH/NaOH).

[157242-76-7, 157242-77-8]

Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (*Karnamicins*)

Tsurumi, Y. *et al.*, *J. Antibiot.*, 1995, **48**, 1066-1072; 1073-1075 (*WS 75624A*)

Antibiotic WS 75624B A-1316
WS 75624B

[157242-75-6]

As Antibiotic WS 75624A, A-1315 with R = -(CH₂)₃CH(OH)CH₃

C₁₈H₂₄N₂O₅S 380.464

Prod. by *Saccharothrix* sp. 75624. Endothelin converting enzyme inhibitor, collagenase; NEP II inhibitor. Powder. Sol. MeOH, Me₂CO, CHCl₃; fairly sol. EtOAc; poorly sol. H₂O. Mp 139-140°. [α]_D²³ +3 (c, 1 in MeOH). λ_{max} 215; 245; 298 (MeOH). λ_{max} 215; 242; 313 (MeOH/HCl). λ_{max} 215; 238; 285; 298 (MeOH/NaOH).

[170129-40-5]

Tsurumi, Y. *et al.*, *J. Antibiot.*, 1995, **48**, 1066-1072; 1073-1075 (*isol, pmr, cmr, synth, props*)

Patt, W.C. *et al.*, *Tet. Lett.*, 1997, **38**, 1297 (*synth*)

Huang, S.-T. *et al.*, *Tet. Lett.*, 1998, **39**, 9335-9338 (*synth*)

Antibiotic XK 211 A-1317
XK 211

[76930-06-8]

C₁₃H₂₀N₄O₇S 376.39


Possibly identical with Altemicidin, A-660. Prod. by *Streptomyces nigrescens* XK 211. Cryst. (H₂O). Sol. bases, CHCl₃; fairly sol. acids, MeOH; poorly sol. EtOAc, hexane. Mp 210°. [α]_D²⁰ -27 (c, 0.1 in H₂O). λ_{max} 302 (H₂O). λ_{max} 308 (HCl) (Berdy). λ_{max} 305 (NaOH aq.) (Berdy).

Japan. Pat., 1980, 80 122 795; *CA*, **94**, 137795y

Antibiotic Y 03559J-A A-1318

3-Cyano-μ-hydroxy-3-methyloxiranetridentricanoic acid, 9CI. Y 03559J-A

[162559-36-6]



C₁₇H₂₉NO₄ 311.42

Prod. by *Micromonospora* sp. Y-03559J.


Active against gram-positive bacteria incl. MRSA. Oil.

Japan. Pat., 1995, 95 02 821; *CA*, **122**, 263680s (*isol*)

Antibiotic Y 05460MA A-1319

Y 05460MA

[126262-07-5]




C₂₁H₃₂N₂O₆ 408.494

Prod. by a *Bacillus* sp. Y-05460M. Active against gram-positive and -negative bacteria. Shows antitumour and antiulcer activities. Powder. Mp 104-106°. Related to Amicoumacin B. λ_{max} 248 (sh); 347 (0.1M NaOH) (Derep). λ_{max} 208 (ε 27300); 248 (ε 6000); 314 (ε 4000) (H₂O) (Derep). λ_{max} 208 (ε 27300); 246 (ε 6400); 314 (ε 4380) (MeOH) (Berdy). λ_{max} 245; 313 (EtOH) (Berdy).

Sato, T. *et al.*, *J. Antibiot.*, 1992, **45**, 1949

Antibiotic YCM 1008A A-1320
YCM 1008A

[944271-69-6]



Absolute Configuration

C₂₁H₂₉NO₄ 359.464


Similar to Antibiotic FA 4283, A-1166. Prod. by *Fusarium* sp. Calcium signalling inhibitor. Pale yellow powder. [α]_D²⁵ -58.1 (c, 0.03 in MeOH).

Koizumi, F. *et al.*, *J. Antibiot.*, 2007, **60**, 455-458 (*isol, pmr, cmr*)

Tatsuta, K. *et al.*, *Tet. Lett.*, 2007, **48**, 4187-4190 (*synth, abs config*)

Antibiotic YM 193221 A-1321

2-(Dimethylamino)-1-(4-hydroxyphenyl)-8,10-dimethyl-6-dodecen-3-one. YM 193221



C₂₂H₃₅NO₂ 345.524


Prod. by *Pseudallescheria ellipsoidea* CBS 128.78. Antifungal agent. Yellow oil. λ_{max} 226; 280 (MeOH).

Kamigiri, K. *et al.*, *J. Antibiot.*, 2004, **57**, 569-572 (*isol, pmr, cmr*)

Antibiotic YM 216391 A-1322

YM 216391

[230645-58-6]



Absolute Configuration

C₃₄H₅₂N₈O₇S 696.742


Cyclic peptide antibiotic. Prod. by *Streptomyces nobilis* JCM 4274. Cytotoxic. Powder. Mp 238° dec. [α]_D²⁵ +48 (c, 0.1 in MeCN). λ_{max} 261 (ε 31000); 273 (ε 31200); 287 (ε 32000); 322 (sh) (ε 8700) (MeCN).

Sohda, K. *et al.*, *J. Antibiot.*, 2005, **58**, 27-31; 32-36 (*isol, pmr, cmr, ms, activity*)

Antibiotic YN 0165JA A-1323

4-[[(Methyl-aci-nitro)acetyl]amino]butanamide, 9CI. YN 0165JA [102692-05-7]
 $\text{MeON(O)}=\text{CHCONH}(\text{CH}_2)_3\text{CONH}_2$
 $\text{C}_7\text{H}_{13}\text{N}_3\text{O}_4$ 203.197
 Prod. by *Streptomyces* sp. Active against gram-positive bacteria. Cryst. (EtOAc). Sol. MeOH, DMSO. Mp 120-121° dec. Related to Enteromycin. λ_{max} 252 (ε 14800) (MeOH) (Derep). λ_{max} 252 (ε 14837) (MeOH) (Berdy).

Imai, H. et al., *J. Antibiot.*, 1986, **39**, 601 (isol. struct, nmr)

Antibiotic ZHD 0501 A-1324

Absolute Configuration


$\text{C}_{28}\text{H}_{22}\text{N}_4\text{O}_4$ 478.506

Related to Staurosporine, S-498. Prod. by a marine-derived *Actinomadura* sp. 007. Cytotoxic. Pale yellow cryst. ($\text{CHCl}_3/\text{MeOH}$). Mp 283.4-285.5°. $[\alpha]_{\text{D}}^{20}$ +83.2 (c, 0.1 in MeOH). λ_{max} 232 (ε 19610); 243 (sh) (ε 19020); 275 (sh) (ε 19720); 290 (ε 33210); 318 (ε 9940); 332 (ε 9640); 351 (ε 6750); 369 (ε 6920) (MeOH).

Han, X.-X. et al., *Tet. Lett.*, 2005, **46**, 6137-6140 (isol. uv, pmr, cmr)

Antidesmone A-1325

6,7-Dihydro-3-methoxy-2-methyl-5-octyl-4,8(1H,5H)-quinolinedione. Hyeronine A [222629-77-8] [221037-18-9]



n = 7

Absolute Configuration

$\text{C}_{19}\text{H}_{29}\text{NO}_3$ 319.443

Struct. revised in 2000. Alkaloid from *Antidesma membranaceum*, many other *Antidesma* spp., *Hieronima oblonga* and *Melochia chamaedrys*. Shows strong fungitoxic activity. Pale yellow oil. $[\alpha]_{\text{D}}^{25}$ +24 (c, 0.2 in CHCl_3). $[\alpha]_{\text{D}}^{25}$ +48.6 (c, 1 in CHCl_3). Genus name given as Hyeronima. The higher opt. rotn. was reported for Hyeronine A from *H. oblonga*. λ_{max} 247 (log ε 4.21); 275 (log ε 3.44); 327 (log ε 3.52) (MeOH).

8α-Alcohol: 5,6,7,8-Tetrahydro-8-hydro-

xy-3-methoxy-2-methyl-5-octyl-4(1H)-quinolinone. **Hyeronimone**

[138690-48-9]
 $\text{C}_{19}\text{H}_{31}\text{NO}_3$ 321.459
 Alkaloid from roots of *Hieronima alchorneoides* (Euphorbiaceae). Mp 85-86°. $[\alpha]_{\text{D}}$ +115 (c, 0.06 in CHCl_3). Abs. config. not determined.

8α-Alcohol, O-Ac: **8-O-Acetylhyeronimone**

[138690-49-0]
 $\text{C}_{21}\text{H}_{33}\text{NO}_4$ 363.496
 From roots of *Hieronima alchorneoides* (Euphorbiaceae). Mp 75-77°. $[\alpha]_{\text{D}}$ +178 (c, 0.06 in CHCl_3).

7'-(β-D-Glucopyranosyloxy):

$\text{C}_{25}\text{H}_{39}\text{NO}_9$ 497.584
 Alkaloid from *Antidesma membranaceum*. $[\alpha]_{\text{D}}^{25}$ +0.2 (c, 0.16 in MeOH). λ_{max} 248 (log ε 4.1); 276 (log ε 3.31); 328 (log ε 3.49) (MeOH).

8-Deoxo: 6,7,8,9-Tetrahydro-3-methoxy-2-methyl-5-octyl-4(1H)-quinolinone. **8-Deoxoantidesmone**

$\text{C}_{19}\text{H}_{31}\text{NO}_2$ 305.459
 Alkaloid from *Antidesma* spp.

8-Deoxo, 17-hydroxy: 6,7,8,9-Tetrahydro-2-(hydroxymethyl)-3-methoxy-5-octyl-4(1H)-quinolinone. **Vanessine**

$\text{C}_{19}\text{H}_{31}\text{NO}_3$ 321.459
 Alkaloid from the stems of *Waltheria douradinha*. Yellow viscous oil. $[\alpha]_{\text{D}}^{25}$ +18.5 (c, 0.12 in CHCl_3).

8-Deoxo, 7'-(β-D-glucopyranosyloxy):

$\text{C}_{25}\text{H}_{41}\text{NO}_8$ 483.601
 Alkaloid from *Antidesma membranaceum*. $[\alpha]_{\text{D}}^{25}$ -49.1 (c, 0.13 in MeOH). λ_{max} 203 (log ε 4.22); 219 (log ε 4.12); 231 (log ε 4.11); 275 (log ε 3.9) (MeOH).

Lower homologue (n = 5): 5-Hexyl-6,7-dihydro-3-methoxy-2-methyl-4,8(1H,5H)-quinolinedione. 17,18-Bisnorantidesmone. **Hyeronine B** [221037-19-0]

$\text{C}_{17}\text{H}_{25}\text{NO}_3$ 291.389
 Alkaloid from *Antidesma velutinosum*, *Hieronima alchorneoides* and *Hieronima oblonga*. Amorph. yellow powder. $[\alpha]_{\text{D}}^{25}$ +16.4 (c, 1 in CHCl_3). λ_{max} 210 (sh); 248 (no solvent reported).

Lower homologue (n = 6): 5-Heptyl-6,7-dihydro-3-methoxy-2-methyl-4,8(1H,5H)-quinolinedione. **18-Norantidesmone**

$\text{C}_{18}\text{H}_{27}\text{NO}_3$ 305.416
 Alkaloid from *Antidesma* spp. and *Hieronima alchorneoides*.

Tinto, W.F. et al., *J. Nat. Prod.*, 1991, **54**, 1309-1313 (Hyeronimone, Acetylhyeronimone)

Alves, T.M.A. et al., *Tet. Lett.*, 1999, **40**, 205-208 (Hyeronimines)

Bringmann, G. et al., *J.A.C.S.*, 2000, **122**, 9905-9910 (biosynth)

Bringmann, G. et al., *Tetrahedron*, 2000, **56**, 3691-3695 (Antidesmone)

Buske, A. et al., *Eur. J. Org. Chem.*, 2001, 3537-3543 (glucosyloxy derivs)


Buske, A. et al., *Phytochemistry*, 2002, **60**, 489-496 (isol, occur, ms, struct, activity)

Dias, G.O.C. et al., *Planta Med.*, 2007, **73**, 289-292 (isol, pmr, cmr)

Gressler, V. et al., *Phytochemistry*, 2008, **69**, 994-999 (Vanessine)

Antifungin A-1326

Phenazine antibiotic. Struct unknown. Prod. by *Pseudomonas mycophaga*. Antifungal agent. Cryst. Sol. MeOH, Et₂O. ▶ LD₅₀ (mus, scu) 300 - 700 mg/kg. Khudyakov, Y.P. et al., *Prikl. Biokhim. Mikrobiol.*, 1965, **1**, 186-190; *CA*, **63**, 7619f

Antiostatin B A-1327

Antiostatin B₂ R = (CH₂)₃CH₃
 B₃ R = (CH₂)₄CH(CH₃)₂
 B₄ R = (CH₂)₆CH₃
 B₅ R = (CH₂)₅CH(CH₃)₂

Carbazole antibiotic complex. Isol. from *Streptomyces cyaneus*. Antioxidative agent.

Antiostatin B₂ [131602-18-1]

$\text{C}_{25}\text{H}_{34}\text{N}_4\text{O}_3$ 438.569
 Sol. MeOH, EtOAc, CHCl_3 , Me₂CO; poorly sol. H₂O, hexane. Mp 119-120°. Grad. starts to subl. at 90°. λ_{max} 218 (ε 44300); 238 (ε 35300); 301 (ε 20900); 338 (ε 6160); 352 (ε 6290) (MeOH) (Derep).

Antiostatin B₃ [131602-19-2]

$\text{C}_{26}\text{H}_{36}\text{N}_4\text{O}_3$ 452.595
 Sol. MeOH, EtOAc, CHCl_3 , Me₂CO, EtOAc; poorly sol. H₂O, hexane. Mp 117-118°. Grad. starts to subl. at 90°. λ_{max} 218 (ε 44300); 238 (ε 35300); 301 (ε 20900); 338 (ε 6160); 352 (ε 6290) (MeOH) (Derep).

Antiostatin B₄ [131643-62-4]

$\text{C}_{26}\text{H}_{36}\text{N}_4\text{O}_3$ 452.595
 Sol. MeOH, Me₂CO, CHCl_3 , EtOAc; poorly sol. H₂O, hexane. Mp 118-120°. Grad. starts to subl. at 90°. λ_{max} 218 (ε 44300); 238 (ε 35300); 301 (ε 20900); 338 (ε 6160); 352 (ε 6290) (MeOH) (Derep).


Antiostatin B₅ [131602-20-5]

$\text{C}_{27}\text{H}_{38}\text{N}_4\text{O}_3$ 466.622
 Sol. MeOH, EtOAc, CHCl_3 , Me₂CO; poorly sol. H₂O, hexane. Mp 92-94°. Grad. starts to subl. at 90°. λ_{max} 218 (ε 44300); 238 (ε 35300); 301 (ε 20900); 338 (ε 6160); 352 (ε 6290) (MeOH) (Derep).

Mo, C.-J. et al., *J. Antibiot.*, 1990, **43**, 1337 (isol, cmr)

Antirhine A-1328

Rhazimine. Anthirine [16049-28-8]



Absolute Configuration

$\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}$ 296.411

Alkaloid from *Antirhea putaminosa*, *Rhazya stricta*, *Catharanthus longifolius* and some other spp. in the Apocynaceae. Mp 115-116°. $[\alpha]_D^{25}$ -2 (CHCl₃) ((+4)). λ_{\max} 225 ; 275 (sh) ; 282 ; 290 (MeOH).

Hydrochloride: Mp 220°. $[\alpha]_D^{25}$ +2 (EtOH).

N⁴- α -Me: Antirhine α -N-methosalt
[50315-65-6]
[50315-63-4] Synthetic. Cryst. (Me₂CO aq.) (as chloride). Mp 325-328° dec. (chloride). $[\alpha]_D$ -17.9 (c, 0.27 in MeOH aq.) (chloride). CAS no. refers to chloride.

N⁴- β -Me: Antirhine β -N-methosalt. N⁴-Methylantirhine
[58580-53-3]
C₂₀H₂₇N₂O⁺ 311.446
Quaternary alkaloid from *Hunteria eburnea*, *Amsonia elliptica* and *Strychnos usambarensis* and *Strychnos camp-toneura* (Apocynaceae, Loganiaceae). Cryst. (EtOH) (as chloride). Mp 306° dec. (chloride). $[\alpha]_D$ +75 (c, 0.04 in EtOH) (chloride).

18,19-Dihydro: Dihydroantirhine
[16049-29-9]

C₁₉H₂₆N₂O 298.427
Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Mp 106-108° (120-125°). $[\alpha]_D$ +23 (c, 0.12 in CHCl₃).

18,19-Dihydro, N⁴- α -Me: Dihydroantirhine α -methochloride
[53729-45-6]
Synthetic. Cryst. (EtOH aq.) (as chloride). Mp 310-312° dec. (chloride). $[\alpha]_D$ -8 (c, 0.13 in EtOH aq.) (chloride). CAS no. refers to chloride.

18,19-Dihydro, N⁴- β -Me: Dihydroantirhine β -N-methosalt. H-Alkaloid J
[53643-59-7]
C₂₀H₂₉N₂O⁺ 313.462
Quaternary alkaloid from the root and stem bark of *Hunteria eburnea* (Apocynaceae). Cryst. (EtOH) (as chloride). Mp 305° dec. (chloride). $[\alpha]_D$ +71.4 (c, 0.13 in H₂O) (chloride). CAS no. refers to chloride.

15-Epimer, A¹⁹-isomer (Z-): Isoantirhine†. Isoantirhine
[113122-68-2]
C₁₉H₂₄N₂O 296.411
Isol. from stem bark of *Strychnos johnsonii* and *Aspidosperma marcgravianum*. Yellow cryst. $[\alpha]_D$ -15 (c, 0.1 in MeOH). λ_{\max} 228 ; 275 ; 282 ; 290 (MeOH).

10-Hydroxy: Hunterburnine. 10-Hydroxyantirhine
[6870-62-8]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from leaves of *Ochrosia alyxioides* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ +15 (c, 0.5 in MeOH). λ_{\max} 230 (log ϵ 3.76); 276 (log ϵ 3.62); 291 (log ϵ 3.52); 310 (log ϵ 3.33) (MeOH). λ_{\max} 280 (log ϵ 3.66); 300 (sh) (log ϵ 3.56); 312 (sh) (log ϵ 3.5); 330 (sh) (log ϵ 3.23) (MeOH/NaOH).

10-Hydroxy, N⁴- α -Me: Hunterburnine α -N-metho salt
C₂₀H₂₇N₂O₂⁺ 327.445
Quaternary alkaloid from *Hunteria eburnea*, *Ochrosia sandwicensis* and *Pleiocarpa mutica* (Apocynaceae). Cryst. (H₂O) (as chloride). Mp 335° (322-324°) (chloride). No CAS no. to 14CI. λ_{\max} 268 (sh) (log ϵ 4.05); 274 (log ϵ 4.08); 302 (log ϵ 3.78); 311 (sh) (log ϵ 3.72) (MeOH) (chloride).

10-Hydroxy, N⁴- β -Me: Hunterburnine β -N-metho salt
C₂₀H₂₇N₂O₂⁺ 327.445
Quaternary alkaloid from *Hunteria eburnea* and *Pleiocarpa mutica* (Apocynaceae). Cryst. (Me₂CO aq.) (as chloride). Mp 307-308° (chloride). $[\alpha]_D$ +105 (72.5% MeOH aq.) (chloride). No CAS no. to 14CI. λ_{\max} 276 (log ϵ 4); 300 (sh) (log ϵ 3.66) (95% EtOH) (chloride).

10-Methoxy: 10-Methoxyantirhine
[126223-76-5]
C₂₀H₂₆N₂O₂ 326.438
Alkaloid from leaves and stem bark of *Ochrosia alyxioides* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ +17 (c, 0.2 in MeOH). λ_{\max} 230 (log ϵ 3.78); 276 (log ϵ 3.64); 291 (log ϵ 3.53); 310 (log ϵ 3.33) (MeOH).

10-Methoxy, 11-hydroxy: Alangine
[262369-77-7]
C₁₈H₂₅NO₃ 303.4
Alkaloid from the fruit of *Alangium lamarekii*. Amorph. powder. $[\alpha]_D$ -0.9 (c, 0.21 in MeOH). $[\alpha]_D$ -2.5 (c, 0.2 in CHCl₃). λ_{\max} 225 (sh) (log ϵ 3.74); 285 (log ϵ 3.45) (MeOH).

ar-Methoxy: Methoxyantirhine
[88608-78-0]
C₂₀H₂₆N₂O₂ 326.438
Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Obt. in admixture with Antirhine. May be identical with 10-Methoxyantirhine.

3-Epimer: 3-Epiantirhine. 3-Isoantirhine
[34384-65-1]
C₁₉H₂₄N₂O 296.411
Alkaloid from the leaves of *Guettarda heterosepala* (Rubiaceae). Cryst. (CHCl₃). Mp 207°. $[\alpha]_D^{20}$ +57 (c, 0.42 in EtOH).

20-Epimer: 20-Epiantirhine
[155418-10-3]
C₁₉H₂₄N₂O 296.411
Alkaloid from roots of *Antirhea portoricensis* (Rubiaceae) and from *Strychnos potatorum*. Amorph. $[\alpha]_D^{25}$ +57 (c, 0.5 in CHCl₃).


20-Epimer, 18,19-dihydro: (20S)-Dihydroantirhine
[144606-86-0]
C₁₉H₂₆N₂O 298.427
Alkaloid from *Strychnos potatorum*. No phys. props. reported.
Bartlett, M.F. et al., *J.O.C.*, 1963, **28**, 1445-1449 (*Hunterburnine, isol, uv, struct*)
Khan, Z.M. et al., *Helv. Chim. Acta*, 1965, **48**, 1957-1965 (*Hunterburnine, isol, uv, ir, ms*)

Asher, J.D.M. et al., *J.C.S.*, 1965, 6355-6367 (*Hunterburnine, cryst struct*)
Jordan, W. et al., *Tetrahedron*, 1965, **21**, 3731-3740 (*Hunterburnine α -N-methosalt, isol, uv, pmr*)
Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1463-1471 (*isol, uv, ir, pmr, ms, struct*)
Sawa, Y.K. et al., *Tetrahedron*, 1969, **25**, 5319-5328 (*Hunterburnine N-methosalts, abs config*)
Sakai, S. et al., *Yakugaku Zasshi*, 1973, **93**, 483; *CA*, **79**, 63538h (*Antirhine N-methosalt*)
Burnell, R.H. et al., *Can. J. Chem.*, 1974, **52**, 2327-2330 (*Antirhine N-methosalt, Dihydroantirhine N-methosalt*)
Bisset, N.G. et al., *Phytochemistry*, 1974, **13**, 1265-1267 (*Antirhine N-methosalt*)
Ficini, J. et al., *J.A.C.S.*, 1979, **101**, 1318-1319 (*synth*)
Takano, S. et al., *Chem. Comm.*, 1981, 1155-1156 (*synth*)
Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 694-707; 708-722 (*Dihydroantirhine, 10-Methoxyantirhine*)
Suzuki, T. et al., *Chem. Pharm. Bull.*, 1986, **34**, 3135-3141 (*synth, ir, pmr*)
Kan-Fan, C. et al., *J. Nat. Prod.*, 1986, **49**, 1130-1132 (*3-Epiantirhine*)
Massiot, G. et al., *Phytochemistry*, 1987, **26**, 2839-2846 (*Isoantirhine*)
Boughandjiou, N. et al., *J. Nat. Prod.*, 1989, **52**, 1107-1112 (*10-Hydroxyantirhine, 10-Methoxyantirhine*)
Lounasmaa, M. et al., *Tetrahedron*, 1989, **45**, 7449-7458 (*18,19-Dihydroantirhine, synth*)
Brown, R.T. et al., *Chem. Comm.*, 1991, 825-826 (*synth*)
Pancrazi, A. et al., *Tet. Lett.*, 1991, **32**, 4483-4486 (*synth*)
Massiot, G. et al., *Phytochemistry*, 1992, **31**, 2873-2876 (*Dihydroantirhine, (20S)-Dihydroantirhine*)
Wei, S.Y. et al., *Tetrahedron*, 1993, **49**, 1025-1042 (*synth*)
Weniger, B. et al., *J. Nat. Prod.*, 1994, **57**, 287-290 (*20-Epiantirhine*)
Danieli, B. et al., *Tetrahedron*, 1994, **50**, 8837-8852 (*synth*)
Tietze, L.F. et al., *Liebigs Ann./Recl.*, 1997, 881-886 (*synth*)
Hanhinen, P. et al., *Heterocycles*, 1999, **51**, 1827-1842 (*synth*)
Itoh, A. et al., *J. Nat. Prod.*, 2000, **63**, 723-725 (*Alangine, isol, struct*)
Takayama, H. et al., *Chem. Pharm. Bull.*, 2002, **50**, 1141-1143 (*Alangine, synth*)

Antirhine lactone

Anthirine lactone
[113122-67-1]

A-1329



C₁₉H₂₀N₂O₂ 308.379


CAS gives the opposite abs. config., apparently incorrectly. Alkaloid from stem bark of *Strychnos johnsonii*. λ_{\max} 228 ; 282 ; 290 (MeOH).

Massiot, G. et al., *Phytochemistry*, 1987, **26**, 2839-2846 (*isol, uv, ir, pmr, cmr, ms, struct*)

Antofine

A-1330

9,11,12,13,13a,14-Hexahydro-2,3,6-trimethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinoline, 9CI. 2,3,6-Trimethoxyphenanthroindolizidine. Tylophorine B



(R)-form

C₂₃H₂₅NO₃ 363.455

Tylophorine B was the racemate.

(R)-form [32671-82-2]

Alkaloid from *Antitoxicum funebre* (preferred genus name *Vincetoxicum*), *Cynanchum vincetoxicum*, *Vincetoxicum nigrum*, *Tylophora asthmatica* and *Ficus septica* (Asclepiadaceae, Moraceae). Possesses pronounced antibacterial and antifungal props. Cryst. (EtOH or CHCl₃/EtOH). Mp 212-214°. [α]_D²² -165 (c, 1.9 in CHCl₃). λ_{max} 259 (log ε 4.71); 286 (log ε 4.46); 342 (log ε 2.99); 360 (log ε 2.61) (MeOH).

Methiodide:

Needles (EtOH). Mp 206°.

N-Oxide (α-): Antofine α-N-oxide

[142698-18-8]

C₂₃H₂₅NO₄ 379.455Alkaloid from *Cynanchum vincetoxicum* (Asclepiadaceae).**N-Oxide (β-): Antofine β-N-oxide**

[314744-19-9]

C₂₃H₂₅NO₄ 379.455Alkaloid from *Cynanchum vincetoxicum* (Asclepiadaceae). Shows cytotoxic activity. [α]_D²¹ -37.3 (c, 0.13 in MeOH).**O⁶-De-Me: 6-Demethylantofine. Alkaloid C†**

[32392-19-1]

C₂₂H₂₃NO₃ 349.429Alkaloid from aerial parts of *Cynanchum vincetoxicum* and *Cynanchum hancockianum*. Cryst. (EtOH). Mp 226-228°. [α]_D²² -125 (c, 1.05 in Py). [α]_D²⁶ -51.5 (c, 0.27 in Py). λ_{max} 260 (ε 4.74); 287 (ε 4.47); 344 (ε 3.05); 362 (ε 2.75) (MeOH).**14R-Hydroxy: 14-Hydroxyantofine**

[32684-56-3]

C₂₃H₂₅NO₄ 379.455Alkaloid from *Cynanchum komarovii* and *Cynanchum vincetoxicum* (Asclepiadaceae). [α]_D²⁵ -72.2 (c, 1 in CHCl₃).**14R-Hydroxy, N-oxide (β-): 14-Hydroxyantofine β-N-oxide**

[142330-17-4]

C₂₃H₂₅NO₅ 395.454Alkaloid from *Cynanchum komarovii* and *Cynanchum vincetoxicum*. Gum.**(S)-form** [51745-04-1]Synthetic. Mp 206-211°. [α]_D²² +66 (CHCl₃).**(±)-form** [26648-81-7]Alkaloid from *Albizzia julibrissin*. Needles (Me₂CO). Mp 213-215°.**Methiodide:**

Cryst. (EtOH). Mp 204-205°.

O⁶-De-Me: [26648-80-6]


Synthetic. Mp 225-228° (216-218).

[142698-18-8]

Wiegerebe, W. *et al.*, *Annalen*, 1969, **721**, 154-162; 1970, **733**, 125-140 (*isol, uv, pmr, ms, struct, synth, Alkaloid C*)Wiegerebe, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1970, **303**, 1009-1012; 1981, **314**, 240-250 (*14-Hydroxyantofine, isol, uv, ir, pmr, ms, struct*)Chauncy, B. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2503-2516 (*synth, uv, ir, pmr*)Herbert, R.B. *et al.*, *Phytochemistry*, 1972, **11**, 1184 (*isol*)Faber, L. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 2882-2884 (*synth*)Govindachari, T.R. *et al.*, *J.C.S. Perkin 1*, 1974, 1161-1165 (*ord, abs config*)Bhakuni, D.S. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 393-395 (*synth, Alkaloid C*)Iwao, M. *et al.*, *Tetrahedron*, 1983, **39**, 1955-1962 (*synth*)Li, X. *et al.*, *Heterocycles*, 1989, **29**, 1797-1808 (*6-O-Demethylantofine*)Capo, M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 389-390 (*isol*)Zhang, R. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1991, **33**, 870-875 (*14-Hydroxyantofine β-N-oxide*)Mi, J.F. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 197-203 (*14-Hydroxyantofine, cd, abs config*)Lavault, M. *et al.*, *Pharm. Acta Helv.*, 1994, **68**, 225; *CA*, **121**, 17809k (*N-oxides*)Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1996, **118**, 12082-12089 (*synth*)Lebrun, S. *et al.*, *Tetrahedron*, 1999, **55**, 2659-2670 (*synth*)Wang, Q. *et al.*, *Acta Cryst. C*, 2000, **56**, 197-198 (*Tylophorine B, cryst struct*)Staerk, D. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1584-1586; 2002, **65**, 1299-1302 (*isol, pmr, cmr, N-oxide, activity*)Kim, S. *et al.*, *J.O.C.*, 2004, **69**, 3144-3149 (*synth*)Fürstner, A. *et al.*, *Chem. Eur. J.*, 2006, **12**, 7398-7410 (*synth*)Camacho-Davila, A. *et al.*, *J.O.C.*, 2006, **71**, 6682-6685 (*synth*)Toribio, A. *et al.*, *Org. Lett.*, 2006, **8**, 3825-3828 (*isol, pmr, cmr, 6-O-Demethylantofine, 14-Hydroxyantofine*)Wang, K.-L. *et al.*, *Tetrahedron*, 2008, **64**, 7504-7510 (*synth*)**Apateline**

A-1331

[68779-85-1]

C₃₄H₃₂N₂O₅ 548.637Alkaloids covered by this entry are diastereomeric with those under Isotriolobine, I-339. Alkaloid from the bark of *Daphnandra apatela* (Monimiaceae). Cryst. + 1MeOH (MeOH). Mp 197-200° dec. [α]_D +270 (CHCl₃).**Dipicrate:**Cryst. + 1.5H₂O. Mp 205-210° dec.**N-Me: N-Methylapateline**

[68779-86-2]

C₃₅H₃₄N₂O₅ 562.664Alkaloid from the leaves, stems and bark of *Daphnandra johnsonii* and from the stems of *Cocculus pendulus* (Monimiaceae, Menispermaceae). Cryst. (MeOH). Mp 165-167° (sinters) Mp 198-200° dec.

O-De-Me: Mp 286-292° dec.

O-De-Me, N-Me: N-Methylnorapateline

[69088-72-8]

C₃₄H₃₂N₂O₅ 548.637Alkaloid from the leaves, stems and bark of *Daphnandra johnsonii* (Monimiaceae). Needles + 0.2CHCl₃. Mp 257-259° dec. [α]_D¹⁹ +235 (CHCl₃).**Me ether: Telobine**

[41758-42-3]


C₃₅H₃₄N₂O₅ 562.664Alkaloid from the bark of *Doryphora apatela* and *Doryphora* sp. Dt-7 (Monimiaceae). Yellow powder + 1/3 CCl₄. Mp 185-195° dec. [α]_D¹⁹ +188 (CHCl₃).**Me ether, dipicrate:**Cryst. + 1H₂O. Mp 198-204° dec.**Me ether, N-Ac:** Mp 178-182° dec. [α]_D¹⁸ +115 (CHCl₃).**Me ether, N-Me: N-Methyltelobine**C₃₆H₃₆N₂O₅ 576.691Alkaloid from tubers of *Stephania erecta* (Menispermaceae). Cryst. (Me₂CO aq.). Mp 175-180° dec. (172°). [α]_D¹⁸ +248 (+226) (CHCl₃). λ_{max} 207 (ε 44600); 233 (ε 36300); 280 (ε 44600); 305 (ε 2950) (MeOH) (Berdy).**5'-Hydroxy: 5'-Hydroxyapateline**C₃₄H₃₂N₂O₆ 564.637Alkaloid from the leaves of *Cocculus pendulus*. [α]_D²⁵ +185 (c, 0.07 in MeOH).**5'-Hydroxy, O¹²-Me: 5'-Hydroxytelobine**C₃₅H₃₄N₂O₆ 578.663Alkaloid from the leaves of *Cocculus pendulus*. [α]_D²⁵ +154 (c, 0.11 in CHCl₃).Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1978, **31**, 2077; 2539 (*Telobine, Apateline, N-Methylapateline, N-Methylnorapateline*)Bick, I.R.C. *et al.*, *Heterocycles*, 1981, **16**, 2105 (*biosynth*)Hussain, S.F. *et al.*, *Tetrahedron*, 1984, **40**,2513 (*N-Methylapateline, 1,2-**Dehydroapateline*)Atta-ur-Rahman, *et al.*, *Pure Appl. Chem.*, 1986, **58**, 663 (*5'-Hydroxyapateline, 5'-Hydroxytelobine*)Guinaudeau, H. *et al.*, *Phytochemistry*, 1987,**26**, 829 (*5'-Hydroxyapateline, 5'-**Hydroxytelobine*)Likhitwitayawuid, K. *et al.*, *J. Nat. Prod.*,1993, **56**, 30 (*2-N-Methyltelobine*)**Apc†**

A-1332

Aphonopelma chalcodes ToxinTwo toxins, Apc₆₀₀ and Apc₇₂₈ partially characterised. Appear to be alkaloidal not polypeptide, contg. spermine, 1,3-diaminopropane and a tyramine-like residue. Isol. from the venom of the tarantula *Aphonopelma chalcodes*. Paralytic toxins.Skinner, W.S. *et al.*, *Toxicol.*, 1990, **28**, 541-546 (*isol*)

Aphanorphine A-1333

2,3,4,5-Tetrahydro-1,3-dimethyl-1,4-methano-1H-benzazepin-8-ol, 9CI
[118964-06-0]



Absolute
Configuration

C₁₃H₁₇NO 203.283

Isol. from the freshwater blue-green alga *Aphanizomenon flos-aquae*. Prisms (Me₂CO aq.). Mp 223-229°. [α]_D²⁵ -43.7 (c, 0.47 in H₂O) (as hydrochloride). [α]_D²⁵ -20.2 (c, 0.15 in MeOH).

Gulavita, N. *et al.*, *Tet. Lett.*, 1988, **29**, 4381 (isol. struct, props)

Takano, S. *et al.*, *Chem. Comm.*, 1989, 1591; 1990, 290 (synth, abs config)

Honda, T. *et al.*, *J.C.S. Perkin I*, 1992, 531 (synth)

Meyers, A.I. *et al.*, *Heterocycles*, 1995, **40**, 525 (synth, *O*-Methylaphanorphine)

Hulme, A.N. *et al.*, *J.O.C.*, 1995, **60**, 1265 (synth)

Fadel, A. *et al.*, *Tetrahedron: Asymmetry*, 1995, **6**, 893; 1997, **8**, 371-374 (synth)

Shimizu, M. *et al.*, *Heterocycles*, 1997, **46**, 21-26 (synth)

Tamura, O. *et al.*, *Org. Lett.*, 2001, **3**, 2427-2429 (synth)

Tanaka, K. *et al.*, *Tet. Lett.*, 2001, **42**, 1049-1052 (synth)

Kita, Y. *et al.*, *J.O.C.*, 2003, **68**, 5917-5924 (synth)

Bower, J.F. *et al.*, *Chem. Comm.*, 2005, 5793-5795 (synth)

Grainger, R.S. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 5377-5380 (synth)

Katoh, M. *et al.*, *Heterocycles*, 2007, **72**, 497-516 (synth)

Li, M. *et al.*, *Synthesis*, 2007, 55-60 (synth, pmr, cmr)


Ma, Z. *et al.*, *Tetrahedron*, 2007, **63**, 7523-7531 (synth)

Yang, X. *et al.*, *Org. Lett.*, 2008, **10**, 2457-2460 (synth)

Yoshimitsu, T. *et al.*, *Tet. Lett.*, 2008, **49**, 4473-4475 (synth)

Aphelandrine A-1334

11-Epiephedradine A. 11-Epiorantine



(+)-form

C₂₈H₃₆N₄O₄ 492.617

(+)-form [69721-57-9]

Alkaloid from *Aphelandra squarrosa* and other *Aphelandra* spp., from *Encephalospaera lasiandra* and *Premna integrifolia* (Acanthaceae, Verbenaceae). Plates (MeOH), needles (H₂O). Mp 263° dec.

[α]_D +214 (c, 0.674 in CHCl₃/MeOH 9:1).

Perchlorate (1:2):

Cryst. (EtOH). Mp 252-253° dec.

N⁶-Hydroxy: N⁶-Hydroxyaphelandrine

[259813-07-5]

C₂₈H₃₆N₄O₅ 508.616

Alkaloid from *Aphelandra fuscopunctata*. Cryst. Mp 186-190° (dec.). [α]_D²¹ +125 (c, 0.18 in EtOH). λ_{max} 232; 280; 286 (EtOH).

N⁶-Acetoxy: N⁶-Acetoxyaphelandrine

[259813-08-6]

C₃₀H₃₈N₄O₆ 550.653

Alkaloid from *Aphelandra fuscopunctata*. Cryst. Mp 176-184° (dec.). [α]_D²² +192 (c, 0.2 in EtOH). λ_{max} 232; 280; 286 (EtOH).

17,18-Diepimer: Orantine. Ephedradine A

[71461-13-7]

C₂₈H₃₆N₄O₄ 492.617

Isol. from the crude drug "Mao-kon", made from the roots of *Ephedra* spp. (Ephedraceae). Hypotensive principle. Noncryst.; cryst. + 1H₂O (as dihydrochloride). Mp 222-225° (dihydrochloride).

17,18-Diepimer, N,N,O-tri-Ac: Mp 165-175°.

17,18-Diepimer, Me ether: O-Methylorantine

[64366-67-2]

C₂₉H₃₈N₄O₄ 506.644

Alkaloid from *Chaenorhinum minus*, *Chaenorhinum villosum* and *Aphelandra sinclairiana* (Scrophulariaceae, Acanthaceae). Cryst. (C₆H₆). Mp 167° dec. [α]_D -84 (c, 0.88 in CHCl₃/MeOH).

17,18-Diepimer, 26-methoxy: Ephedradine D

[81349-26-0]

C₂₉H₃₈N₄O₅ 522.643

Isol. from the crude drug "Mao-kon", made from the roots of *Ephedra* spp. (Ephedraceae). Hypotensive principle. Noncryst. Mp 219-221° (as hydrobromide). [α]_D -85.3 (H₂O) (hydrobromide).

17,18-Diepimer, 30-methoxy, Me ether:

Ephedradine C

[73276-37-6]

C₃₀H₄₀N₄O₅ 536.67

Isol. from the crude drug "Mao-kon", made from the roots of *Ephedra* spp. (Ephedraceae). Hypotensive principle. Noncryst.; cryst. + 1H₂O (as hydrobromide). Mp 224-225° (hydrobromide). [α]_D -100.7 (H₂O).

17,18-Diepimer, deoxy, 30-methoxy:

Ephedradine B

[71327-57-6]

C₂₉H₃₈N₄O₅ 522.643

Isol. from the crude drug "Mao-kon", made from the roots of *Ephedra* spp. (Ephedraceae). Hypotensive principle. Noncryst.; cryst. + 1H₂O (as hydrobromide). Mp 219-221° (hydrobromide). [α]_D -101.5 (H₂O).

(-)-form [133162-95-5]

Alkaloid from the whole plant of *Schweinfurthia papilionacea* (Scrophular-

iaceae). Mp 200° dec. [α]_D -13 (c, 0.3 in MeOH). This alkaloid, claimed as new and named as 11-Epiephedradine A or 11-Epiorantine, appears to be (-)-Aphelandrine of low opt. purity. However this requires confirmation.

Dätwyler, P. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 2646-2671; 1979, **62**, 2712-2723

(*Aphelandrine, Orantine, O-Methylorantine*)

Tamada, M. *et al.*, *Heterocycles*, 1979, **12**, 783-786 (*Ephedradine B*)

Tamada, M. *et al.*, *Tet. Lett.*, 1979, 873-876

(*Ephedradine A, cryst struct, abs config*)

Konno, C. *et al.*, *Heterocycles*, 1980, **14**, 295-298 (*Ephedradine C*)

Hikino, H. *et al.*, *Heterocycles*, 1982, **17**, 155-158 (*Ephedradine D*)

Wasserman, H.M. *et al.*, *J.A.C.S.*, 1985, **107**, 519-521 (synth)

Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 1012-1016 (*Aphelandrine, cryst struct, abs config*)

Ahmad, V.U. *et al.*, *J. Nat. Prod.*, 1990, **53**,

1162-1167 (*11-Epiephedradine A*)


Youhnovski, N. *et al.*, *Phytochemistry*, 1999, **52**, 1717-1723 (*N-Hydroxyaphelandrine, N-acetoxyaphelandrine*)

Nezbedova, L. *et al.*, *Tet. Lett.*, 2000, **41**, 7859-7862 (*biosynth*)

Kurosawa, W. *et al.*, *Tetrahedron*, 2004, **60**, 9615-9628 (*Orantine, synth*)

Aphyllic acid A-1335

Octahydro-3-(2-piperidinyl)-2H-quinoline-1-carboxylic acid, 9CI. *Aphyllinic acid*
[642-67-1]



Absolute
configuration

C₁₅H₂₆N₂O₂ 266.383

Artifact obt. during coml. acid treatment of *Anabasis aphylla* alkaloids (Chenopodiaceae). Powder, cryst. + 5H₂O. Mp 218-219° (anhyd.). [α]_D +14. Almost certainly derived by ring-opening of Aphylline, which it readily reforms (with some stereomutation).

Hydrochloride (1:2): Mp 262-264°.

Me ester: Methyl aphyllate

Alkaloid from seeds of *Anabasis aphylla* (Chenopodiaceae). Cryst.

(petrol). Mp 80-81°. [α]_D +8.9 (synthetic).

2',3'-Didehydro: Aphyllidinic acid

C₁₅H₂₄N₂O₂ 264.367

Alkaloid artifact from *Anabasis aphylla* acid treatment, also obt. from Aphyllidine by treatment with H₂SO₄ aq. (Chenopodiaceae). Mp 215-216° dec. [α]_D²⁰ -8 (c, 7 in MeOH).

2',3'-Didehydro, picrate: Mp 196° dec.

Labenskii, A.S. *et al.*, *Zh. Obshch. Khim.*, 1958, **28**, 547; *J. Gen. Chem. USSR (Engl. Transl.)*, 537 (isol. struct)


Sadykov, A.S. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 1733; *J. Gen. Chem. USSR (Engl. Transl.)*, 1720 (isol. struct)

Aslanov, Kh.A. *et al.*, *CA*, 1967, **67**, 73730f (isol)

- Cho, Y.D. *et al.*, *Anal. Biochem.*, 1971, **44**, 49 (glc, ms)
 Ishbaev, A.I. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 328; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 322 (ord)

Aphyllidine A-1336

3,4,7,7a,8,9,10,11,13,14-Decahydro-7,14-methano-2H,6H-dipyrido[1,2-a:1',2'-e][1,5]diazocin-6-one, 9CI
 [643-32-3]



(+)-form

C₁₅H₂₂N₂O 246.352**(+)-form** [124223-08-1]

Alkaloid from *Anabasis aphylla* (Chenopodiaceae) and *Lupinus argenteus* ssp. *rubricaulis* (Fabaceae). Mp 112-113°. [α]_D²⁶ +7.3 (c, 2.18 in EtOH). [α]_D¹⁸ +5.6 (MeOH).

Hydrochloride: Mp 235-237°. [α]_D +30 (H₂O).

2R-Hydroxy: 2R-Hydroxyaphyllidine
 [124223-06-9]

C₁₅H₂₂N₂O₂ 262.351

Constit. of *Lupinus argenteus* ssp. *rubricaulis* (Fabaceae). Solid. Mp 150-155°. [α]_D²⁶ -15 (c, 0.2 in CHCl₃).

2S-Hydroxy: 2S-Hydroxyaphyllidine
 [124223-07-0]

C₁₅H₂₂N₂O₂ 262.351

Constit. of *Lupinus argenteus* ssp. *rubricaulis* (Fabaceae).

2R,9R-Dihydroxy: 2R,9R-Dihydroxyaphyllidine

[124125-78-6]

C₁₅H₂₂N₂O₃ 278.35

Constit. of *Lupinus argenteus* ssp. *rubricaulis* (Fabaceae). Oil (as di-Ac). [α]_D²⁶ +38.5 (c, 0.62 in CHCl₃) (as di-Ac).

2S,9R-Dihydroxy: 2S,9R-Dihydroxyaphyllidine

[124125-77-5]

C₁₅H₂₂N₂O₃ 278.35

Constit. of *Lupinus argenteus* ssp. *rubricaulis* (Fabaceae). Prisms (MeOH/Et₂O). Mp 122-123°. [α]_D²⁶ +53.6 (c, 1.48 in CHCl₃).

17-Oxo: Oxoaphyllidine. Oxoaphyllidine
 C₁₅H₂₀N₂O₂ 260.335

Alkaloid from *Anabasis aphylla* (Chenopodiaceae). Also obt. by oxidn. of Aphyllidine. Mp 188-189° (182-184°). [α]_D²⁰ -29.8 (c, 0.56 in EtOH) (-21). The higher Mp and opt. rotn. are for semisynthetic material, the lower for the nat. prod. An incorr. abs. config. was prev. given. λ_{max} 244 (log ε 0.75) (EtOH).

13S-Hydroxy: 13α-Hydroxyaphyllidine

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Lupinus hartwegii* (Fabaceae). Oil. [α]_D²⁵ +44 (c, 0.1 in CHCl₃). λ_{max} 239 (log ε 3.9) (MeOH).

(-)-form

Alkaloid from *Argyrolobium megarhizum* (Fabaceae). Prisms (Et₂O/hexane). Mp 101-106°. [α]_D -9.1 (c, 2.2 in EtOH).

Hydrochloride:

Prisms (MeOH/Me₂CO/Et₂O). Mp 228-230° dec.

2ξ-Hydroxy: Argyrolobine

[6871-18-7]

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Argyrolobium megarhizum* whole plants (Fabaceae). Prisms (Me₂CO). Mp 168-169°. [α]_D +13.3 (c, 1.4 in CHCl₃). Subl. undec. Unstable in CHCl₃ soln.

2ξ-Hydroxy; hydrochloride:

Prisms (MeOH/EtOAc). Mp 170-172°.

2ξ-Acetoxy:

Prisms. Mp 61°.

Orechoff, A. *et al.*, *Ber.*, 1932, **65**, 234; 1934, **67**, 1845; 1976

Späth, E. *et al.*, *Ber.*, 1942, **75**, 805 (isol)

Tsuda, Y. *et al.*, *Can. J. Chem.*, 1964, **42**, 764

(isol, uv, struct, Aphyllidine, Argyrolobine)

Schumann, D. *et al.*, *Monatsh. Chem.*, 1968, **99**, 390 (ms)

Zakharov, V.P. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 805; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 775 (isol)

Nurimov, E. *et al.*, *CA*, 1974, **80**, 130536s

(biosynth)

Ishbaev, A.I. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 113; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 128 (Oxoaphyllidine, pmr, uv)

Arslanian, R.L. *et al.*, *J.O.C.*, 1990, **55**, 1204

(isol, derivs)


Mohamed, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 365 (13-Hydroxyaphyllidine)

Aphylline

A-1337

10-Oxosparteine

[577-37-7]

C₁₅H₂₄N₂O 248.367

Alkaloid from *Anabasis aphylla*, *Virgilia oroboides*, *Cytisus canariensis* and *Lupinus hartwegii* (Chenopodiaceae, Fabaceae). Mp 52-53°. Bp₄ 200°. [α]_D +10.3 (20% MeOH aq.). Cryst. with difficulty.

Hydrochloride:

Cryst. + 1H₂O. [α]_D²⁰ +13.6 (H₂O).

Picolonate: Mp 230-231° dec.

Methiodide: Mp 212-213° dec.

N¹⁶-Oxide: Aphylline N¹⁶-oxide

[41451-56-3]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Anabasis aphylla* leaves (Chenopodiaceae).

2α-Hydroxy: 2α-Hydroxyaphylline

[197297-29-3]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Lupinus hartwegii*.

Amorph. [α]_D²⁵ +30.4 (c, 0.1 in CHCl₃).

11-Epimer: Epiaphylline. Alkaloid A1. α-Isoaphylline

[1218-51-5]

C₁₅H₂₄N₂O 248.367

Alkaloid from *Lupinus hartwegii* whole plants (Fabaceae). Cryst. Config. incorr. descr. as 11S-.

11-Epimer, 2,3-didehydro: 2,3-Dehydroepiaphylline. 2,3-Dehydro-α-isoaphylline.

2,3-Dehydro-10-oxo-α-isoarteine

C₁₅H₂₂N₂O 246.352

Alkaloid from *Uresiphita reversalis* larvae fed on *Cytisus monspessulamus* leaves. Needles. Mp 98-103°. [α]_D²⁶ +132 (c, 0.6 in EtOH). λ_{max} 257 (log ε 3.07) (MeOH).

[10159-81-6]

Orechoff, A. *et al.*, *Ber.*, 1932, **65**, 234 (isol)

Galinovsky, F. *et al.*, *Monatsh. Chem.*, 1953, **84**, 199 (struct)

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1954, **32**, 235 (config)

Wiewiorowski, M. *et al.*, *Can. J. Chem.*, 1967, **45**, 1447 (ir, pmr, conformm)

Schumann, D. *et al.*, *Monatsh. Chem.*, 1968, **99**, 390 (ms)

Steinegger, E. *et al.*, *An. Quim.*, 1972, **68**, 893 (isol)

Klyne, W. *et al.*, *J.C.S. Perkin I*, 1974, 2565 (cd)

Anderson, J.N. *et al.*, *J.O.C.*, 1976, **41**, 3441

(isol, ms, Aphylline, Epiaphylline)

Nizamkodzaeva, A.N. *et al.*, *CA*, 1978, **88**, 121499v (isol, oxide)

Abdusalomov, B.A. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 56; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 46

(biosynth)


Mohamed, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 365-369 (2-Hydroxyaphylline)

Nihei, K. *et al.*, *Phytochemistry*, 2002, **61**, 987-990 (Dehydroepiaphylline)

Apicularen A

A-1338

[220757-04-0]



Absolute Configuration

C₂₅H₃₁NO₆ 441.523

Prod. by *Chondromyces apiculatus*, *Chondromyces robustus* and other *Chondromyces* spp. Cytotoxic agent. Vacuolar-type ATPase inhibitor. Antineoplastic agent. Mp 139-141° dec. [α]_D -36.1 (c, 1 in MeCN). λ_{max} 278 (log ε 4.43) (MeOH).

4-O-(2-Acetamido-2-deoxy-β-D-glucopyranoside): Apicularen B

[220757-06-2]

C₃₃H₄₄N₂O₁₁ 644.717

Prod. by *Chondromyces* spp. Cytotoxic agent. [α]_D -5.5 (c, 0.3 in MeOH). λ_{max} 280 (log ε 4.49) (MeOH).

Kunze, B. *et al.*, *J. Antibiot.*, 1998, **51**, 1075-1080 (isol, ir, activity)

Jansen, R. *et al.*, *Eur. J. Org. Chem.*, 2000, 913-919 (pmr, cmr, ms, abs config)


Bhattacharjee, A. *et al.*, *Tet. Lett.*, 2001, **42**, 1217-1220 (synth)

Yet, L. *et al.*, *Chem. Rev.*, 2003, **103**, 4283-4306 (rev)

- Lewis, A. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 104-116 (*synth*)
 Graetz, B.R. *et al.*, *Org. Lett.*, 2003, **5**, 3357-3360 (*synth*)
 Petri, A.F. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 5821-5823 (*synth*)
 Nicolaou, K.C. *et al.*, *Chem. Eur. J.*, 2004, **9**, 6177-6191 (*synth*)
 Su, Q. *et al.*, *J.A.C.S.*, 2004, **126**, 2425-2430 (*synth*)
 Hilli, F. *et al.*, *Org. Lett.*, 2004, **6**, 1289-1292 (*synth*)
 Huss, M. *et al.*, *BMC Biochem.*, 2005, **6**, 13 (*pharmacol*)
 Hong, J. *et al.*, *J. Pharmacol. Exp. Ther.*, 2005, **312**, 968-977 (*pharmacol*)
 Li, M. *et al.*, *Org. Lett.*, 2006, **8**, 6087-6090 (*synth*)
 Jung, Y.-H. *et al.*, *Chem. Asian J.*, 2007, **2**, 656-661 (*synth*)

Apiodionene A-1339

3-[5,6-Dihydro-5-methyl-6-(1-propenyl)-2H-pyran-2-ylidene]-2,4-pyrrolidinedione, 9CI
 [142808-38-6]




C₁₃H₁₅NO₃ 233.266

Tetramic acid deriv. Isol. from *Apiosordaria effusa*. Inhibits topoisomerase. Cytotoxic. Antiinflammatory and anticancer agent. Sol. MeOH, DMF, Me₂CO, DMSO; poorly sol. H₂O, hexane. Mp 190-192°. [α]_D²⁵ +155.4. λ_{max} 222 (ε 3700); 327 (ε 18600) (MeOH) (Berdy). λ_{max} 245 (ε 12300); 277 (ε 8700) (MeOH-HCl) (Berdy).

Japan. Pat., 1992, 92 49 289; *CA*, **117**, 88730f (*isol, ir, uv, pmr, cmr*)
 Takahashi, S. *et al.*, *CA*, 1993, **118**, 187475; **119**, 24279z (*isol, pmr, cmr, props*)

Apiosporamide A-1340

[162414-45-1]



Absolute Configuration

C₂₄H₃₁NO₆ 429.512

Prod. by the fungus *Apiospora montagnei* (terrestrial strain). Antifungal agent. Sol. MeOH; fairly sol. CHCl₃, Me₂CO; poorly sol. H₂O, C₆H₆, hexane. Mp 240-250° dec. [α]_D²⁵ -97.4 (c, 0.04 in MeOH). λ_{max} 210 (ε 13700); 231 (ε 11400); 279 (ε 4200); 330 (ε 7100) (MeOH) (Berdy).

N-Hydroxy: *N*-Hydroxyapiosporamide

[179638-09-6]

C₂₄H₃₁NO₇ 445.511

Metab. of a fungus *Cytospora* sp. associated with the ant *Pogonomyrmex badius*. Cholesteryl ester transfer protein inhibitor. Light yellow solid. Mp 148° (dec.). [α]_D²⁵ -59.7 (c, 0.5 in MeOH). λ_{max} 231 (ε 19000); 283 (ε

12000); 339 (ε 12000) (MeOH).

4''-Ketone: *Antibiotic YM 215343*. YM 215343

C₂₄H₂₉NO₆ 427.496

Prod. by *Phoma* sp. QN04621. Antifungal agent. Powder. [α]_D²⁵ -44 (c, 0.1 in MeOH). λ_{max} 231 (ε 15300); 279 (ε 5200); 330 (ε 8900) (MeOH).

Alfatafta, A.A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1696-1702 (*isol, uv, ir, pmr, cmr, ms*)

Lee, J.C. *et al.*, *J. Antibiot.*, 1996, **49**, 693-696 (*N*-Hydroxyapiosporamide)


Shibazaki, M. *et al.*, *J. Antibiot.*, 2004, **57**, 379-382 (YM 215343)

Williams, D.R. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 6715-6718 (*synth, abs config*)

Aplaminal

[1005010-20-7]

A-1341



Absolute Configuration

C₁₆H₁₉N₃O₅ 333.343

Isol. from *Aplysia kurodai*. Cytotoxic. Plates (MeOH). Mp 235-237°. [α]_D²⁰ -133 (c, 0.02 in MeOH).

Kuroda, T. *et al.*, *Org. Lett.*, 2008, **10**, 489-491 (*isol, pmr, cmr, cryst struct*)


Smith, A.B. *et al.*, *Org. Lett.*, 2008, **10**, 4363-4365 (*synth*)

Aplaminone

A-1342

12-[3-Bromo-2-[2-(dimethylamino)ethyl]-5-hydroxy-4-methoxyphenyl]-2,6,10-trimethyl-2,10-dodecadien-5-one, 9CI

[131815-20-8]



C₂₆H₄₀BrNO₃ 494.511

(*R*)-form

Isol. from the marine mollusc *Aplysia kurodai*. Cytotoxic. Oil. [α]_D²³ -2.9 (c, 1.18 in MeOH). λ_{max} 225 (ε 11000); 285 (ε 2900) (MeOH) (Berdy).

Δ⁹-Isomer(*E*-), 11'-hydroxy: *Neoaplaminone*

[131815-21-9]

C₂₆H₄₀BrNO₄ 510.51

Alkaloid from the marine mollusc *Aplysia kurodai*. Cytotoxic. Oil. [α]_D²³ -5.3 (c, 0.65 in MeOH). λ_{max} 226 (ε 11000); 284 (ε 2500) (MeOH) (Derep).

Δ⁹-Isomer(*E*-), 11'-hydroxy, 3-O-sulfate: *Neoaplaminone sulfate*

[131815-22-0]

C₂₆H₄₀BrNO₅S 590.574

Isol. from the marine mollusc *Aplysia kurodai*. Cytotoxic. [α]_D²⁷ -3 (c, 1.29 in


CHCl₃). λ_{max} 227 (ε 11000); 282 (ε 1700) (MeOH) (Derep).

Kigoshi, H. *et al.*, *Tet. Lett.*, 1990, **31**, 4911-4914; 1992, **33**, 4195-4198 (*isol, uv, ir, pmr, cmr, ms, struct, abs config*)

Aplicyanin A

[936633-59-9]

A-1343



C₁₂H₁₃BrN₄ 293.166

Alkaloid from *Aplidium cyaneum*. Pale yellow oil. [α]_D²⁵ -0.8 (c, 0.1 in CHCl₃).

N²-Ac: *Aplicyanin B*

[936633-60-2]

C₁₄H₁₅BrN₄O 335.203

Alkaloid from *Aplidium cyaneum*. Cytotoxic. Pale yellow oil. [α]_D²⁵ +8.7 (c, 0.1 in CHCl₃).

1-Methoxy: *Aplicyanin C*

[936633-61-3]

C₁₃H₁₅BrN₄O 323.192

Alkaloid from *Aplidium cyaneum*. Pale yellow oil. [α]_D²⁵ +3.1 (c, 0.1 in CHCl₃).

1-Methoxy, N²-Ac: *Aplicyanin D*

[936633-62-4]

C₁₅H₁₇BrN₄O₂ 365.229

Alkaloid from *Aplidium cyaneum*. Cytotoxic. Pale yellow oil. [α]_D²⁵ +9.5 (c, 0.1 in CHCl₃).

5-Bromo, 1-methoxy: *Aplicyanin E*

[936633-63-5]

C₁₃H₁₄Br₂N₄O 402.088

Alkaloid from *Aplidium cyaneum*. Cytotoxic. Pale yellow oil. [α]_D²⁵ +0.5 (c, 0.1 in CHCl₃).

5-Bromo, 1-methoxy, N²-Ac: *Aplicyanin F*

[936633-64-6]

C₁₅H₁₆Br₂N₄O₂ 444.125


Alkaloid from *Aplidium cyaneum*. Pale yellow oil. [α]_D²⁵ +1.9 (c, 0.1 in CHCl₃).

Reyes, F. *et al.*, *Tetrahedron*, 2008, **64**, 5119-5123 (*isol, pmr, cmr*)

Aplidinone A

[875057-52-6]

A-1344



R = -OMe

C₁₉H₂₅NO₅S 379.476

Alkaloid from the ascidian *Aplidium conicum*. Orange amorph. solid. λ_{max} 310 (ε 5700) (MeOH).

Aiello, A. *et al.*, *Eur. J. Org. Chem.*, 2005, 5024-5030 (*isol*, *pmr*, *cmr*)

Aplidinone B A-1345

[875081-15-5]

As Aplidinone A, A-1344 with

R = -NH₂

C₁₈H₂₄N₂O₄S 364.465

Alkaloid from the ascidian *Aplidium conicum*. Violet amorph. solid. λ_{max} 330 (ε 2600) (MeOH).

N⁷-(2-Sulfoethyl): **Aplidinone C**

[875082-39-6]


C₂₀H₂₈N₂O₇S₂ 472.582

Alkaloid from *Aplidium conicum*. Red-violet amorph. solid. λ_{max} 340 (ε 2900) (MeOH).

Aiello, A. *et al.*, *Eur. J. Org. Chem.*, 2005, 5024-5030 (*isol*, *pmr*, *cmr*)

Aplidite A A-1346

[165689-34-9]



C₂₇H₃₂N₂O₈ 512.558

Unique orthonitrite functionality. Alkaloid from the Australian marine tunicate *Aplidium* sp. Solid. Mp 126-128°. [α]_D -15.8 (c, 0.58 in MeOH). λ_{max} 213 (ε 14300); 284 (ε 3785) (MeOH) (Berdy).

N,O-Di-Ac: Synthetic. Yellow solid. Mp 84-86.5°.

12-Hydroxy: **Aplidite C**

[165689-35-0]

C₂₇H₃₂N₂O₉ 528.558

From *Aplidium* sp. Solid. Mp 103.5-105.5°. λ_{max} 208 (ε 11733); 280 (ε 6746) (MeOH) (Berdy).

12-Hydroxy, 15-Ac: **Aplidite E**

[165689-36-1]

C₂₉H₃₄N₂O₁₀ 570.595

From *Aplidium* sp. Solid. Mp 104-105°. [α]_D -50 (c, 0.71 in MeOH). λ_{max} 208 (ε 12892); 280 (ε 8686) (MeOH) (Berdy).

Geom. isomer: **Aplidite B**

[165883-78-3]

C₂₇H₃₂N₂O₈ 512.558

From *Aplidium* sp. Solid. Mp 124.5-126°. [α]_D -32.9 (c, 0.665 in MeOH). Possibly has opposite config. at the Δ¹ double bond. λ_{max} 213 (ε 14300); 284 (ε 3785) (MeOH) (Berdy).

Geom. isomer, 12-hydroxy: **Aplidite D**

[165883-79-4]

C₂₇H₃₂N₂O₉ 528.558

From *Aplidium* sp. Solid. Mp 107.5-109°. [α]_D -17.8 (c, 0.105 in MeOH). λ_{max} 206 (ε 15000); 282 (ε 8800) (MeOH) (Berdy).

Geom. isomer, 12-hydroxy, 15-Ac: **Aplidite F**

[165883-80-7]

C₂₉H₃₄N₂O₁₀ 570.595

From *Aplidium* sp. Solid. Mp 105-109°. [α]_D -8.6 (c, 0.245 in MeOH). λ_{max} 208 (ε 5286); 285 (ε 2830) (MeOH) (Berdy).

Geom. isomer, 12-acetoxy: **Aplidite G**

[165689-37-2]


C₂₉H₃₄N₂O₁₀ 570.595

From *Aplidium* sp. Solid. Mp 106-108°. [α]_D -26.9 (c, 0.60 in MeOH). λ_{max} 201 (ε 27400); 286 (ε 11000) (MeOH) (Berdy).

Murray, L. *et al.*, *Aust. J. Chem.*, 1995, 48, 1253-1266 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Aplysepine

[151756-71-7]




C₁₉H₂₅N₃O₃ 343.425

Alkaloid from the sea hare *Aplysia kurodai*. [α]_D¹² -3 (c, 0.47 in MeOH). λ_{max} 231 (ε 33600); 256 (sh) (ε 9400); 298 (sh) (ε 1600) (MeOH) (Derep).

Ojika, M. *et al.*, *Tet. Lett.*, 1993, 34, 5307-5308 (*isol*, *ir*, *pmr*, *cmr*, *struct*)

Aplysinafulvin

[1025766-28-2]




C₁₀H₁₃Br₂NO₄ 371.025

Isol. from *Aplysina fulva*. Glassy solid. [α]_D²⁵ +130 (c, 0.002 in MeOH). λ_{max} 230 (log ε 3.68); 286 (log ε 3.69) (MeOH).

Nuñez, C.V. *et al.*, *Biochem. Syst. Ecol.*, 2008, 36, 283-296 (*isol*, *cd*, *pmr*, *cmr*)

Aplysinamisine I

[150417-67-7]



C₁₆H₁₇Br₂N₅O₄ 503.149


Alkaloid from the Caribbean sponge *Aplysina cauliformis*. Oil. [α]_D²⁶ +121.9 (c, 5.7 in MeOH). λ_{max} 226 (ε 13700); 266 (ε 13100) (MeOH) (Berdy).

Rodríguez, A.D. *et al.*, *J. Nat. Prod.*, 1993, 56, 907 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Aplysinamisine II

A-1350

[150417-68-8]



C₁₆H₂₃Br₂N₅O₄ 509.197


Alkaloid from the Caribbean sponge *Aplysina cauliformis*. Semisolid. [α]_D⁵⁶ +47 (c, 7.9 in MeOH). λ_{max} 218 (ε 4600); 284 (ε 2500) (MeOH) (Berdy).

Rodríguez, A.D. *et al.*, *J. Nat. Prod.*, 1993, 56, 907 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Aplysinellin A

A-1351

[325155-76-8]



C₃₁H₂₉Br₃N₄O₁₃S₃ 1001.499


Isol. from the sponge *Aplysina rhax*. Amorph. yellow solid (as di-Na salt). Mp 191-194° dec. (di-Na salt). λ_{max} 204 (log ε 4.83); 291 (log ε 4.11) (MeOH) (di-Na salt).

Shin, J. *et al.*, *Tetrahedron*, 2000, 56, 9071-9077

Aplysinellin B

A-1352

[325155-78-0]




C₃₁H₂₇Br₃N₄O₁₃S₃ 999.483

Isol. from the sponge *Aplysina rhax*. Amorph. yellow solid (as mono-Na salt). Mp 236-239° dec. (Na salt). λ_{max} 205 (log ε 4.88); 280 (log ε 3.93) (MeOH) (Na salt).

Shin, J. *et al.*, *Tetrahedron*, 2000, 56, 9071-9077

Aplysintetral A

[129138-56-3]



R = (CH₂)₃CH₃

C₁₃H₁₉Br₂NO₄ 413.105

Related to 3,5-Dibromo-1-hydroxy-4,4-dimethoxy-2,5-cyclohexadiene-1-acetamide, D-318. Metab. of *Aplysina thiona*. Cryst. Mp 194-195°.

Cruz, F. et al., *J. Nat. Prod.*, 1990, **53**, 543-548 (isol, pmr, ms)

Toscano, R.A. et al., *Acta Cryst. C*, 1992, **48**, 2235-2237 (cryst struct)

Aplysintetral B

[129138-57-4]

As Aplysintetral A, A-1353 with

R = (CH₂)₄CH₃


C₁₄H₂₁Br₂NO₄ 427.132

Metab. of *Aplysina thiona*. Cryst. Mp 194-195°.

Cruz, F. et al., *J. Nat. Prod.*, 1990, **53**, 543-548 (isol, pmr, ms)

Aplysinone A

[943837-22-7]



Absolute Configuration

n = 5

C₂₄H₂₆Br₄N₄O₈ 818.107

Constit. of *Aplysina gerardogreeni*. Cytotoxic. Amorph. solid. [α]_D²⁵ +65 (c, 0.1 in Me₂CO).

3'-Debromo, 2'-deoxy, 2'α,3'α-epoxide:

Aplysinone C

[943837-24-9]

C₂₄H₂₅Br₃N₄O₈ 737.196

Constit. of *Aplysina gerardogreeni*. Amorph. solid. [α]_D²⁵ +102 (c, 0.1 in Me₂CO).

Hernández-Guerrero, C.J. et al., *Bioorg. Med. Chem.*, 2007, **15**, 5275-5282 (isol, pmr, cmr)

Aplysinone D

[943837-25-0]

As Aplysinone A, A-1355 with

n = 4

C₂₃H₂₄Br₄N₄O₈ 804.081

Constit. of *Aplysina gerardogreeni*. Amorph. solid. [α]_D²⁵ +86 (c, 0.1 in Me₂CO).


Hernández-Guerrero, C.J. et al., *Bioorg. Med. Chem.*, 2007, **15**, 5275-5282 (isol, pmr, cmr)

Aplysinopsin

2-Imino-5-(1H-indol-3-ylmethylene)-1,3-dimethyl-4-imidazolidinone, 9CI

[63153-56-0]

[138127-90-9]



C₁₄H₁₄N₄O 254.291

Metab. of the marine sponges *Verongia spengelii* and *Thorecta* sp. and from the anthozoan *Astroides calycularis*. Antibiotic with antitumour props. Fine yellow needles + 1H₂O (MeOH aq. + Et₂NH). Mp 235-237° dec. (232-233°). λ_{max} 217 (ε 27500); 280 (ε 10300); 406 (ε 24800) (EtOH aq./HCl) (Derep). λ_{max} 260 (ε 10000); 376 (ε 21200) (EtOH aq./NaOH) (Derep). λ_{max} 225 (ε 23000); 268 (ε 8800br); 383 (ε 21500) (50% EtOH) (Derep).

Di-Ac:

Fine yellow needles (EtOH). Mp 221-222° dec. (217-220°).

N¹-Propanoyl: N¹-Propionylaplysinopsin [100667-70-7]

C₁₇H₁₈N₄O₂ 310.355

Metab. from the anthozoan *Astroides calycularis*. Cryst. (MeOH). Mp 300°.

3'-N-Me: N-3'-Methylaplysinopsin

[66492-97-5]

[66492-99-7]

C₁₅H₁₆N₄O 268.318

Prod. by *Aplysinopsis reticulata*. Monoamine oxidase inhibitor, serotonergic neurotransmission potentiator. Sol. MeOH, CHCl₃; poorly sol. H₂O.

3'-N-Et: N-3'-Ethylaplysinopsin

C₁₆H₁₈N₄O 282.344

Isol. from the Jamaican sponge *Smenospongia aurea*. Yellowish gum. No CAS no. found to CA 136-139. λ_{max} 219 (log ε 4.13); 276 (log ε 3.74) (MeOH).

N²-De-Me: 2'-N-Demethylaplysinopsin

[72479-07-3]

C₁₃H₁₂N₄O 240.264

Metab. of the marine sponge *Dercitus* sp., also isol. from the nudibranch *Phestilla melanobranchia* and the corals *Tubastrea coccinea* and *Dendrophyllia* sp. Mp 235° (with charring from 180°). λ_{max} 217 (ε 27500); 280 (ε 10300); 406 (ε 24800) (EtOH aq./HCl) (Derep). λ_{max} 260 (ε 10000); 376 (ε 21200) (EtOH aq./NaOH) (Derep). λ_{max} 225 (ε 23000); 268 (ε 8800br); 383 (ε 21500) (50% EtOH) (Derep). λ_{max} 385 (ε 25000) (MeOH) (Berdy).

N²-De-Me, 3'-N-Me: 2'-N-Demethyl-N-3'-methylaplysinopsin

[126149-80-2]

C₁₄H₁₄N₄O 254.291

Isol. from *Dendrophyllia* sp. Yellow

solid. Mp > 250° dec. Obt. as an E/Z-mixt., Z-form predominates.

N⁴-De-Me, 3'-N-Me: **Isoplysin A**

[158761-04-7]

C₁₄H₁₄N₄O 254.291

Metab. from the Okinawan marine sponge *Aplysina* sp. Weakly cytotoxic. Yellow needles. Mp 310° dec. λ_{max} 220 (ε 14000); 278 (ε 5400); 405 (ε 13000) (MeOH) (Berdy).

1',8-Dihydro: 1',8-Dihydroaplysinopsin

[85079-46-5]

C₁₄H₁₆N₄O 256.307

Isol. from the coral *Tubastrea coccinea*.

3'-Deimino, 3'-oxo: [63153-57-1]

[117604-23-6, 117603-69-7]

C₁₄H₁₃N₃O₂ 255.276

Alkaloid from the coral *Tubastrea* sp. Obt. as an E/Z-mixt.

3'-Deimino, 3'-oxo, N², N⁴-di-de-Me:

[117490-33-2, 117490-34-3]

C₁₂H₉N₃O₂ 227.222

Alkaloid from the coral *Leptopsammia pruvoti*. Obt. as an E/Z-mixt.

6-Bromo: **6-Bromoaplysinopsin**

[85079-45-4]

C₁₄H₁₃BrN₄O 333.187

Isol. from the marine sponge *Smenospongia aurea*, the anthozoan *Astroides calycularis* and the coral *Tubastrea coccinea*. Cryst. (MeOH). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 283-285°. λ_{max} 230 (ε 35000); 284 (ε 15000); 364 (ε 26000); 385 (ε 24000) (MeOH) (Berdy).

6-Bromo, N¹-propanoyl: **6-Bromo-N¹-propionylaplysinopsin**

[100667-71-8]

C₁₇H₁₇BrN₄O₂ 389.251

Metab. from the anthozoan *Astroides calycularis*. Cryst. (MeOH). Mp 300°.

6-Bromo, N²-de-Me: **6-Bromo-2'-N-de-methylaplysinopsin**

[72479-08-4]

C₁₃H₁₁BrN₄O 319.116

Metab. of the marine sponge *Dercitus* sp. Also isol. from the nudibranch *Phestilla melanobranchia* and the coral *Tubastrea coccinea*. Mp 186-188° (with charring from 175°). λ_{max} 390 (ε 23000) (MeOH) (Derep).

6-Bromo, N²-de-Me, 3'-N-Me: **6-Bromo-2'-N-demethyl-N-3'-methylaplysinopsin**

[126149-84-6 (E-form), 126149-81-3 (Z-form)]

C₁₄H₁₃BrN₄O 333.187

Isol. from *Dendrophyllia* sp. Yellow solid. Mp > 280° dec. Obt. as an E/Z-mixt., Z-form predominates. λ_{max} 238 (ε 8500); 286 (ε 5200); 385 (ε 10700) (MeOH).

6-Bromo, N⁴-de-Me: **6-Bromo-4'-N-demethylaplysinopsin**

[97480-15-4]

C₁₃H₁₁BrN₄O 319.116

Metab. of the marine sponge *Smenospongia aurea*. Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 290° (as Ac). Characterised as the Ac deriv. λ_{max} 390 (ε 23000) (MeOH) (Derep). λ_{max} 230 (ε 35000); 284 (ε 15000); 364 (ε 26000)

(MeOH) (Berdy).

6-Bromo, 1',8-dihydro- 6-Bromo-1',8-dihydroaplysinopsin

[85079-47-6]
C₁₄H₁₅BrN₄O 335.203

Isol. from the coral *Tubastrea coccinea*.

6-Bromo, 1',8-dihydro, 1'ξ-hydroxy- 6-Bromo-1',8-dihydro-1'-hydroxyaplysinopsin

[868830-18-6]
C₁₄H₁₅BrN₄O₂ 351.202

Isol. from a *Thorectandra* sp. Red solid. [α]_D²⁵ +1 (c, 0.5 in MeOH).

6-Bromo, 1',8-dihydro, 1'ξ-methoxy- 6-Bromo-1',8-dihydro-1'-methoxyaplysinopsin

[868830-19-7]
C₁₅H₁₇BrN₄O₂ 365.229

Isol. from a *Thorectandra* sp. Red solid. [α]_D²⁵ +3 (c, 1.4 in MeOH).

6-Bromo, 1',8-dihydro, 1'ξ-ethoxy- 6-Bromo-1'-ethoxy-1',8-dihydroaplysinopsin

[868830-20-0]
C₁₆H₁₉BrN₄O₂ 379.256

Isol. from a *Smenospongia* sp. Red solid. [α]_D²⁵ +19.3 (c, 0.05 in MeOH).

6-Bromo, 3'-deimino, 3'-oxo- 6-Bromo-3'-deimino-3'-oxoaplysinopsin

[117490-31-0, 117490-32-1]
C₁₄H₁₂BrN₃O₂ 334.172

Alkaloid from the coral *Tubastrea* sp. Obt. as an *E-Z*-mixture.

6-Bromo, 3'-deimino, 3'-oxo, N^{2'},N^{4'}-di-de-Me- 6-Bromo-3'-deimino-2',4'-bis(-demethyl)-3'-oxoaplysinopsin

[117490-30-9, 117490-29-6]
C₁₂H₈BrN₃O₂ 306.118

Alkaloid from the coral *Leptopsammia pruvotii*. Obt. as an *E-Z*-mixture.

5,6-Dibromo, N^{2'}-de-Me- 5,6-Dibromo-2'-N-demethylaplysinopsin

[376598-41-3]
C₁₃H₁₀Br₂N₄O 398.056

Isol. from the sponge *Hyrtios erecta*. Yellow powder. λ_{max} 246 (ε 8500); 293 (ε 2700); 385 (ε 8600) (MeOH).

Z-Isomer, 5,6-dibromo, N^{2'}-de-Me-

[376598-39-9]
C₁₃H₁₀Br₂N₄O 398.056

Isol. from *Hyrtios erecta*. Yellow powder. λ_{max} 247 (ε 17300); 294 (ε 4400); 383 (ε 20700) (MeOH).

Hollenbeak, K.H. *et al.*, *J. Nat. Prod.*, 1977, **40**, 479-481 (isol, uv, ir, pmr, ms, struct)

Kazlauskas, R. *et al.*, *Tet. Lett.*, 1977, 61-64 (uv, ir, pmr, ms, struct)

Djura, P. *et al.*, *J.O.C.*, 1980, **45**, 735-737 (2'-N-Demethylaplysinopsin, 6-Bromo-2'-N-demethylaplysinopsin)

Okuda, R.K. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 1907-1914 (1',8-Dihydroaplysinopsin, 6-Bromo-1',8-dihydroaplysinopsin)

Fattorusso, E. *et al.*, *J. Nat. Prod.*, 1985, **48**, 924-927 (N¹-Propionylaplysinopsin, 6-Bromo-N¹-propionylaplysinopsin)

Tymiak, A.A. *et al.*, *Tetrahedron*, 1985, **41**, 1039-1047 (6-Bromoaplysinopsin, 6-Bromo-4'-N-demethylaplysinopsin)

Guella, G. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 773-781; 1989, **72**, 1444-1450 (*Dendrophyllia constits*, *Tubastrea constits*)

Dalkafouki, A. *et al.*, *Tet. Lett.*, 1991, **32**, 5325-5328 (synth)

Gulati, D. *et al.*, *Indian J. Chem.*, 1994, **33**, 4-9; 10-16 (synth)

Kondo, K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1008-1011 (*Isoplysin A*)

Molina, P. *et al.*, *Tetrahedron*, 1994, **50**, 2241-2254 (synth, derivs)


Aoki, S. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1372-1374 (5,6-Dibromo-N-demethylaplysinopsin)

Hu, J.-F. *et al.*, *J. Nat. Prod.*, 2002, **65**, 476-480 (N-3'-Ethylaplysinopsin, N-3'-Methylaplysinopsin)

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1484-1488 (6-Bromo-1',8-dihydro-1'-hydroxyaplysinopsin)

Aplyzanzine A

[327165-67-3]



C₂₅H₃₃Br₄N₃O₃ 743.17


Isol. from an *Aplysina* sp. Pale orange oil.

Evan, T. *et al.*, *J. Nat. Prod.*, 2001, **64**, 226-227

Apoatropine

A-1359

8-Methyl-8-azabicyclo[3.2.1]oct-3-yl α-methylenebenzeneacetate, 9CI. Tropine atropate. Atropamine. Atropyltropine [500-55-0]



C₁₇H₂₁NO₂ 271.358

Alkaloid from *Atropa belladonna*, *Datura pruinosa*, *Datura stramonium*, *Duboisia leichhardtii*, *Anisodus tanguticus* root, *Hyoscyamus orientalis*, *Mandragora* spp., *Physochlaina alaiica* and other spp. (Solanaceae). Also obt. by dehydration of atropine with acids, acetic anhydride, etc. Antispasmodic agent. Cryst. (Et₂O). Mp 60-62°. Log P 2.64 (calc).

►YM4025000

Hydrochloride:

Leaflets (H₂O). Mp 245-246° (237-239°).

Picrate:

Yellow needles. Mp 166-168°.

N-Oxide:

Cryst. (EtOH/Me₂CO). Mp 127-128°.

N-De-Me: Aponoratropine. Atropoylnor-tropan-3-ol

C₁₆H₁₉NO₂ 257.332

Minor alkaloid from leaves of *Anthro-troche myoporoides* (Solanaceae).

Ladenburg, A. *et al.*, *Annalen*, 1883, **217**, 74 (synth)

Hesse, O. *et al.*, *Annalen*, 1893, **277**, 290 (synth)

Wolffenstein, R. *et al.*, *Ber.*, 1908, **41**, 723 (synth)

Takeuchi, Y. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2603 (synth, ir, pmr)

Sharova, E.G. *et al.*, *Khim. Prir. Soedin.*, 1977, 126 (isol)


Kagei, K. *et al.*, *Yakugaku Zasshi*, 1980, **100**, 216 (isol)

Evans, W.C. *et al.*, *Phytochemistry*, 1981, **20**, 497 (isol, Aponoratropine)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AQO250

Apocinchonamine

A-1360



Absolute Configuration

C₁₉H₂₂N₂O 294.396

Parent not isol.

O-Ac: 5-O-Acetylapocinchonamine

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from the leaves of *Remijia peruviana*. Amorph. [α]_D²⁵ +4.6 (c, 0.22 in EtOH). λ_{max} 207 (log ε 4.26); 302 (log ε 3.8) (EtOH).

18,19-Dihydro: Apodihydrocinchonamine

[151271-88-4]

Alkaloid from the leaves of *Isertia haenkeana*. Amorph. solid. [α]_D²⁰ +12 (c, 2 in CHCl₃). λ_{max} 224 (log ε 4.46); 347 (log ε 4.34) (MeOH).


Bruix, M. *et al.*, *Phytochemistry*, 1993, **33**, 1257-1261 (*Apodihydrocinchonamine*)

Diaz, J.G. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1667-1671 (*5-Acetylapocinchonamine*)

Apogalanthamine

A-1361

5,6,7,8-Tetrahydro-6-methyldibenz[c,e]azocine-1,2-diol, 9CI [26955-02-2]



C₁₆H₁₇NO₂ 255.316

Mp 202-203° dec.

►CD9380000

O^δ-Me: Nivalidine

[18126-83-5]

C₁₇H₁₉NO₂ 269.343

Alkaloid from *Galanthus nivalis* var. *gracilis* (Amaryllidaceae). Antiarhythmic, hypertensive. Approved for use as a hypertensive agent ca. 1969 (USSR). Needles (C₆H₆). Mp 206-208°. Log P 3.4 (calc). Also produced by treatment of Galanthamine with HCl aq. Probably an artifact.

O^δ-Me, hydrochloride: Apochlorine.

Apokhlorin

[24620-67-5]

Creamy cryst. (EtOH). Mp 163-166° (foaming).

▶ CD9380000

O⁸-Me, hydrobromide: Mp 238-239°.

O⁸-Me, Ac:

Cryst. (Et₂O). Mp 107-109.5°.

O⁹-Me: **8-O-Demethylbuflavine**

[61972-44-9]

C₁₇H₁₉NO₂ 269.343

Alkaloid from bulbs of *Boophone flava* (Amaryllidaceae). Amorph. solid. Mp 112-114°. Genus name given as Boophane.

Di-Me ether: **Buflavine**

[65762-70-1]

C₁₈H₂₁NO₂ 283.369

Alkaloid from bulbs of *Boophone flava* (Amaryllidaceae). Amorph. solid. Mp 106-108°.

Kobayashi, S. *et al.*, *J.C.S.*, 1957, 638-645 (*Apogalanthamine*)

Bubewa-Iwanova, L. *et al.*, *Chem. Ber.*, 1962, **95**, 1348 (*Nivalidine*, synth, ir, uv)

Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 194; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 168 (*Nivalidine*, synth, use)

Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 307-311 (*Buflavine*, *Demethylbuflavine*)


Hoarau, C. *et al.*, *J.O.C.*, 2002, **67**, 5846-5849 (*Buflavine*, synth)

Sahakitpichan, P. *et al.*, *Tet. Lett.*, 2003, **44**, 5239-5241 (*Buflavine*, synth)

Kodama, S. *et al.*, *Tetrahedron*, 2004, **60**, 4901-4907 (*Buflavine*, synth)

Apogeissoschizine

[6900-80-7]



Absolute Configuration

C₂₁H₂₂N₂O₂ 334.417

Obt. by treatment of Geissospermine, G-35 with conc. HCl. Cryst. (MeOH/EtOAc/hexane) (as hydrochloride). Mp 139-142° (hydrochloride). λ_{max} 273 (ε 19300); 324 (ε 19000) (EtOH) (as hydrochloride).

2,7-Dihydro, 2α,7α-dihydroxy: **2,7-Dihydroxyapogeissoschizine**

[154850-61-0]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from root bark of *Strychnos gossweileri* (Loganiaceae). λ_{max} 211 (ε 12600); 285 (ε 10000); 316 (ε 15800) (MeOH/HCl) (Derep). λ_{max} 219 (ε 12600); 288 (ε 10000); 326 (ε 15800) (MeOH/NaOMe) (Derep). λ_{max} 211 (ε 12600); 288 (ε 10000); 323 (ε 15800) (MeOH) (Derep).

Rapoport, H. *et al.*, *J.A.C.S.*, 1958, **80**, 1601-1604; 1960, **82**, 4404-4414

(*Apogeissoschizine*)


Quetin-Leclercq, J. *et al.*, *Phytochemistry*, 1994, **35**, 533-536 (2,7-

Dihydroxyapogeissoschizine)

Apohaemanthamine

Aphoemanthamine

[6900-81-8]



C₁₆H₁₅NO₃ 269.299

Alkaloid from *Eucharis amazonica*. Amorph. solid. Mp 145-147°. [α]_D²² +198 (c, 0.63 in CHCl₃).


Campbell, W.E. *et al.*, *Phytochemistry*, 2000, **53**, 587-591 (*synth*, pmr, cmr)

Cabezas, F. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 315-317 (*isol*, cd)

Apohyoscine

A-1364

9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]-non-7-yl α-methylbenzeneacetate, 9CI. 6,7-Epoxy-3-atropolyoxytropane. *Oscine atropate*. *Aposcopolamine* [535-26-2]



C₁₇H₁₉NO₃ 285.342

Alkaloid from the aerial parts of *Datura innoxia*, *Datura meteloides*, *Hyoscyamus niger* and *Anthocercis frondosa* (stem-bark and roots) (Solanaceae). Needles (Et₂O). Mp 97°. Opt. inactive (*meso*-).

Picrate:

Needles (MeOH aq.). Mp 216° dec.

N-De-Me: **Aponorhyoscine**

[25650-56-0]

C₁₆H₁₇NO₃ 271.315

Minor alkaloid from roots of *Anthocercis genistoides* (Solanaceae). Plates (EtOH aq.) (as picrate). Mp 224-225° (picrate).

Evans, W.C. *et al.*, *J.C.S.*, 1965, 4936 (*isol*)

Sharova, E.G. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 126; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 117 (*isol*)


Evans, W.C. *et al.*, *J. Pharm. Pharmacol. Suppl.*, 1979, **31**, 9P (*isol*)

Evans, W.C. *et al.*, *Phytochemistry*, 1981, **20**, 497 (*isol*, *Aponorhyoscine*)

Apparicine

A-1365

4-Ethylidene-1,3,4,5,6,7-hexahydro-6-methylene-2,5-ethano-2H-azocino[4,3-b]indole, 9CI. *Gomezine*. *Pericalline*. *Tabernoschizine*



(R)-form

C₁₈H₂₀N₂ 264.369

(R)-form [3463-93-2]

Alkaloid from the bark of *Aspidosperma dasycarpon* and from *Aspidosperma olivaceum*, *Vallesia antillana*, *Parkia speciosa*, *Tabernaemontana dichotoma* and *Sapitum sebiferum*. Cryst. (Me₂CO). Mp 192-194°. [α]_D²⁷ +176 (c, 2.16 in CHCl₃). λ_{max} 303 (ε 46500) (EtOH) (Berdy).

(S)-form [2122-36-3]

Found in *Aspidosperma olivaceum* and several other *Aspidosperma* spp., *Catharanthus ovalis*, *Catharanthus roseus*, *Pandaca ochrascens*, *Pandaca eusepala*, *Ervatamia heyneana*, *Tabernaemontana cumminsii*, *Schizozygia coffaeoides* and other spp. in the Apocynaceae. Shows cytotoxic activity against mouse lymphocyte P-388 cells. Also shows weak antibacterial activity. Antiviral agent active against Polio virus but not Vaccinia. Shows pronounced analeptic props. Cryst. (Me₂CO or petrol/CHCl₃). Mp 196-202° (188-190°). [α]_D²⁷ -179 (c, 2.16 in CHCl₃). pK_{a1} 8.05 (33% DMF). pK_{a1} 7.26 (80% 2-methoxyethanol). λ_{max} 218 (ε 31300); 230 (ε 49000); 307 (ε 41800) (MeOH) (Berdy).

19R,20R-Epoxyde: **Epchrosine**. 19,20-Epoxyapparicine

[103763-50-4]

C₁₈H₂₀N₂O 280.369

Alkaloid from cell suspension cultures of *Ochrosia elliptica* (Apocynaceae).

10-Hydroxy: **10-Hydroxyapparicine**

[70503-70-7]

C₁₈H₂₀N₂O 280.369

Alkaloid from the leaves of *Ochrosia oppositifolia* (Apocynaceae). Noncryst. [α]_D²⁵ -90 (CHCl₃). λ_{max} 314 (log ε 4.92) (no solvent reported).

16α-Hydroxy, 16,22-dihydro: **16,22-Dihydro-16-hydroxyapparicine**

[94061-32-2]

C₁₈H₂₂N₂O 282.385

Alkaloid from the leaves of *Tabernaemontana dichotoma* (Apocynaceae). Also isol. from whole plants and callus cultures of *Tabernaemontana elegans*. [α]_D²⁰ +129 (c, 0.1 in EtOH). λ_{max} 220 (log ε 4.43); 284 (log ε 3.79); 292 (sh) (EtOH).

10-Methoxy: **10-Methoxyapparicine**

[70503-71-8]

C₁₉H₂₂N₂O 294.396

Alkaloid from the leaves of *Ochrosia oppositifolia* (Apocynaceae). Cryst. (Me₂CO). Mp 199-202°. [α]_D²⁵ -129 (CHCl₃). λ_{max} 314 (log ε 4.92) (no solvent reported).

(ξ)-form

19,20-Dihydroxy, 19,20-dihydro: **Brafouedine**

[104021-43-4]

C₁₈H₂₂N₂O₂ 298.384

Alkaloid from the stem bark of *Strychnos dinklagei*. Noncryst. [α]_D²⁰ -26 (c, 0.02 in MeOH). Stereochem. undefined. CAS assigns an abs. config. although none was determined. λ_{max} 220 (log ε 4.02); 230 (sh) (log ε 3.93);

300 (log ϵ 3.65); 313 (sh) (log ϵ 3.6) (EtOH).

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1963, **52**, 407-408 (*Pericalline*)

Joule, J.A. *et al.*, *J.C.S.*, 1965, 4773-4780 (*struct*)

Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141-1166 (*isol, ir*)

Joule, J.A. *et al.*, *Tetrahedron*, 1965, **21**, 1717-1734 (*isol, uv, pmr, struct*)

Farnsworth, N.R. *et al.*, *J. Pharm. Sci.*, 1969, **58**, 280 (*props*)

Cosson, J.P. *et al.*, *Phytochemistry*, 1970, **9**, 1353-1354 (*isol*)

Segelman, A.B. *et al.*, *J. Pharm. Sci.*, 1974, **63**, 1419-1422 (*isol, uv*)

Panas, J.M. *et al.*, *Phytochemistry*, 1974, **13**, 1969-1974 (*isol, uv, ms*)

Akhter, L. *et al.*, *Tet. Lett.*, 1978, 4137 (*10-Hydroxyapparicine, 10-Methoxyapparicine*)

Langlois, N. *et al.*, *Phytochemistry*, 1979, **18**, 467-471 (*isol*)

Laguna, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1980, **45**, 1419 (*isol, uv, ir, pmr, ms*)

Heatley, F. *et al.*, *J.C.S. Perkin 2*, 1980, 919-924 (*pmr, conformn*)

Gunasekera, S.P. *et al.*, *Phytochemistry*, 1980, **19**, 1213-1218 (*isol, uv, ir, pmr, ms*)

Massiot, G. *et al.*, *Chem. Comm.*, 1982, 768-769 (*cmr*)

Perera, P. *et al.*, *J. Nat. Prod.*, 1984, **47**, 835-838 (*16,22-Dihydro-16-hydroxyapparicine*)

Michel, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 452-455 (*Brafouedine*)


Van der Heijden, R. *et al.*, *Phytochemistry*, 1986, **25**, 843-846 (*16,22-Dihydro-16-hydroxyapparicine*)

Pawelka, K.-H. *et al.*, *Plant Cell Rep.*, 1986, **5**, 147-149 (*Epchrosine*)

Apratoxin A

A-1366

[350791-64-9]



Absolute Configuration

R = -C(CH₃)₃C₄₅H₆₉N₅O₈S 840.135

Isol. from *Lyngbya majuscula*. Potent cytotoxin. Amorph. solid. $[\alpha]_D^{25}$ -161 (c, 1.3 in MeOH).

N¹²-De-Me: Apratoxin B

[444885-29-4]

C₄₄H₆₇N₅O₈S 826.108

Isol. from *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25}$ -73 (c, 0.2 in MeOH). λ_{max} 200 (log ϵ 4.64); 230 (log ϵ 4.36) (MeOH).

Luesch, H. *et al.*, *J.A.C.S.*, 2001, **123**, 5418-5423 (*isol, pmr, cmr*)

Luesch, H. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 1973-1978 (*Apratoxin B*)

Chen, J. *et al.*, *J.A.C.S.*, 2003, **125**, 8734-8735 (*synth*)

Ma, D. *et al.*, *Chem. Eur. J.*, 2006, **12**, 7615-7626 (*synth*)

Doi, T. *et al.*, *Org. Lett.*, 2006, **8**, 531-534 (*synth*)

Apratoxin C

A-1367

[444885-30-7]

As Apratoxin A, A-1366 with

R = -CH(CH₃)₂C₄₄H₆₇N₅O₈S 826.108Isol. from *Lyngbya* sp. Cytotoxic agent.

Amorph. solid. $[\alpha]_D^{25}$ -171 (c, 0.22 in MeOH). λ_{max} 201 (log ϵ 4.64); 230 (log ϵ 4.31) (MeOH).

Luesch, H. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 1973-1978

Aquaticine

A-1368

C₁₈H₂₅NO₅ 335.399

Struct. unknown. Alkaloid from *Senecio aquaticus* (Asteraceae). Octahedral cryst. (EtOH). Mp 220° dec. $[\alpha]_D^{25}$ -83 (c, 1 in CHCl₃).


Picrate:

Yellow needles (MeOH). Mp 182°.

Evans, W.C. *et al.*, *Nature (London)*, 1949, **164**, 30-31

Aquileidine

A-1369

C₂₀H₂₀N₂O₅ 368.388Constit. of *Aquilegia ecalcarata*.

Amorph. powder (MeOH). Mp 214-215°. $[\alpha]_D^{25}$ +21 (c, 0.54 in MeOH). λ_{max} 288; 368 (sh) (MeOH).

Chen, S.-B. *et al.*, *J. Nat. Prod.*, 2001, **64**, 85-87

Aquileginine

A-1370

C₄₁H₅₆N₂O₂²⁺ 608.906


Struct. unknown. Prob. a bisbenzylisoquinoline alkaloid. Alkaloid from the roots of *Aquilegia hybrida* "Scott-Elliott" (Ranunculaceae). Plates (MeOH) (as diiodide). Mp 238-239° (diiodide). $[\alpha]_D^{22}$ -7.8 (c, 0.36 in CHCl₃) (diiodide).

Winek, C.L. *et al.*, *J. Nat. Prod.*, 1964, **27**, 111-114 (*isol, uv*)

Aragupetrosine A

A-1371

[125236-58-0]

C₃₀H₅₂N₂O₂ 472.753


Alkaloid from an Okinawan marine sponge *Xestospongia* sp. Exhibits vasodilatory activity. $[\alpha]_D$ -18.8 (CHCl₃).

Kobayashi, M. *et al.*, *Tet. Lett.*, 1989, **30**, 4149 (*isol, ir, pmr, cmr, struct, abs config*)

Araguspongine C

A-1372

[122908-10-5]



Relative configuration

C₂₈H₅₀N₂O₄ 478.714

Alkaloid from an Okinawan marine sponge, *Xestospongia* sp. and from the sponge *Haliclona exigua*. Exhibits vasodilatory activity. $[\alpha]_D$ +11.1 (CHCl₃).

N-Oxide: Araguspongine LC₂₈H₅₀N₂O₅ 494.713Alkaloid from *Xestospongia exigua*.Needles. $[\alpha]_D^{25}$ +1.7 (c, 0.36 in CHCl₃).**3 α -Methyl: 3 α -Methylaraguspongine C**

[159392-28-6]

C₂₉H₅₂N₂O₄ 492.741

Alkaloid from the sponge *Haliclona exigua*. Amorph. solid. $[\alpha]_D$ +1.2 (c, 0.15 in CHCl₃).

Kobayashi, M. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1676 (*isol, pmr, cmr, struct*)


Venkateswarlu, Y. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1283 (*3 α -Methylaraguspongine C*)

Orabi, K.Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1782-1785 (*Araguspongines C,L*)

Araguspongine F

A-1373

[122908-11-6]



Absolute configuration

C₂₉H₅₂N₂O₂ 460.742


Alkaloid from an Okinawan marine sponge, *Xestospongia* sp. Exhibits vasodilatory activity. $[\alpha]_D$ +8.6 (CHCl₃).

Work in 1998 suggested that the stereochemistries may require revision.

3-Epimer: Araguspongine G

[123000-04-4]
 $C_{29}H_{52}N_2O_2$ 460.742
 Alkaloid from *Xestospongia* sp. Shows vasodilatory activity. $[\alpha]_D -16$ (CHCl₃).
 Kobayashi, M. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1676 (*isol, pmr, cmr, struct, abs config*)
 Baldwin, J.E. *et al.*, *J.A.C.S.*, 1998, **120**, 8559-8560 (*configs*)

Aragspongine J A-1374
 [123000-05-5]




$C_{30}H_{54}N_2O_2$ 474.769
 Alkaloid from an Okinawan marine sponge, *Xestospongia* sp. Exhibits vasodilatory activity. $[\alpha]_D -23.4$ (CHCl₃).
 Work in 1998 suggested that the stereochemistries may need revision.

3'-Epimer: Aragspongine H
 [122908-12-7]
 $C_{30}H_{54}N_2O_2$ 474.769
 Alkaloid from *Xestospongia* sp. Shows vasodilatory activity. $[\alpha]_D -9$ (CHCl₃).

3,3',5,9-Tetraepimer: 3R,3'R-Dimethylxestospongine C
 [203576-22-1]
 $C_{30}H_{54}N_2O_2$ 474.769
 Isol. from a *Xestospongia* sp.

Kobayashi, M. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1676 (*isol, ir, pmr, cmr, struct, abs config*)
 Reddy, M.V.R. *et al.*, *Nat. Prod. Lett.*, 1997, **11**, 53-59 (*3,3'-Dimethylxestospongine C*)
 Baldwin, J.E. *et al.*, *J.A.C.S.*, 1998, **120**, 8559-8560 (*configs*)

Aralionine A A-1375
 N-(7-Benzoyl-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl)-2-(dimethylamino)-3-methylpentanamide, 9CI. Aralio-nine
 [21761-48-8]




$C_{34}H_{38}N_4O_5$ 582.698
 Major alkaloid from the leaves of *Araliorhamnus vaginatus* (Rhamnaceae).
 Cryst. (petrol or EtOH aq.). Mp 165-

167°. $[\alpha]_D^{20} +82$ (c, 0.2 in MeOH). λ_{max} 246 (ε 19000) (MeOH). λ_{max} 305 (ε 10600) (MeOH/KOH).

1''-Alcohol(R): Aralio-nine C
 [64309-20-2]
 $C_{34}H_{40}N_4O_5$ 584.714
 Alkaloid from the bark of *Araliorhamnus vaginatus* (Rhamnaceae).
 Needles (EtOH aq.). Mp 95-97°. $[\alpha]_D^{20} -17$ (c, 0.015 in MeOH).

1''-Deoxo: Deoxoaralio-nine A. Deoxyaralio-nine A
 [70420-57-4]
 $C_{34}H_{40}N_4O_4$ 568.714
 Alkaloid from the root bark of *Ceanothus integerrimus* var. *integerrimus* and *Ceanothus integerrimus* var. *californicus* (Rhamnaceae). Mp 350°. No stereochem. indicated.
 Tschesche, R. *et al.*, *Chem. Ber.*, 1969, **102**, 50-63 (*isol, uv, cd, ir, pmr, ms, struct*)
 González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915-916 (*stereochem*)
 Tschesche, R. *et al.*, *Phytochemistry*, 1977, **16**, 1025-1028 (*Aralio-nine C*)
 Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 220 (*Deoxoaralio-nine A*)


Araliopdimerine A A-1376
 [119089-40-6]



$C_{34}H_{38}N_2O_8$ 602.683
 Alkaloid from the bark of *Araliopsis tabouensis* (Rutaceae). Cryst. (CHCl₃/hexane). Mp 259-260°. Racemic.

17,18-Dihydro, 18-hydroxy: Araliopdi-merine B
 [119089-41-7]
 $C_{34}H_{40}N_2O_9$ 620.698
 Alkaloid from the bark of *Araliopsis tabouensis* (Rutaceae). Yellow granules (Et₂O). Mp 246-247°. Racemic.
 Authors' numbering.
 Ngadjui, B.T. *et al.*, *Phytochemistry*, 1988, **27**, 2979 (*isol, uv, ir, pmr, cmr, ms, struct*)

Araliopsine A-1377
 3,5-Dihydro-2-(1-hydroxy-1-methylethyl)-5-methylfuro[3,2-c]quinolin-4(2H)-one, 9CI. Pseudoisoplatydesmine



$C_{15}H_{17}NO_3$ 259.304
 Absolute configuration revised in 2000.


(S)-form [82226-76-4]
 [60354-62-3]
 Alkaloid from the root bark of *Araliopsis soyauxii* (Rutaceae). Beige cryst. (CHCl₃). Mp 152°. $[\alpha]_D +40$ (c, 1 in CHCl₃).

(±)-form [89708-41-8]
 Synthetic. Mp 141-144°.

(E)-form
1'-Deoxy, 1',2'-didehydro: 3,5-Dihydro-5-methyl-2-(1-methylethenyl)furo[3,2-c]quinolin-4(2H)-one. 3,5-Dihydro-2-isopropenyl-5-methylfuro[3,2-c]quinolin-4(2H)-one. Almeine
 [88839-93-4]
 $C_{15}H_{15}NO_2$ 241.289
 Alkaloid from the stem and root barks of *Almeida guyanensis* (Rutaceae). Noncryst.

Vaquette, J. *et al.*, *Phytochemistry*, 1976, **15**, 743 (*isol, ir, uv, ms, pmr, struct*)
 Coppola, G.M. *et al.*, *J. Het. Chem.*, 1983, **20**, 1589 (*synth, uv, ir, pmr, cmr*)
 Moulis, C. *et al.*, *Phytochemistry*, 1983, **22**, 2095 (*Almeine*)
 Reisch, J. *et al.*, *Monatsh. Chem.*, 1989, **120**, 363; 1993, **124**, 1225 (*Almeine, synth, uv, ir, pmr, cmr, ms*)
 Barr, S.A. *et al.*, *Chem. Comm.*, 1994, 153 (*synth*)
 Reisch, J. *et al.*, *Pharmazie*, 1994, **49**, 62 (*Almeine, synth*)
 Boyd, D.R. *et al.*, *J.C.S. Perkin 1*, 2000, 3397-3405 (*synth, abs config*)
 Bar, G. *et al.*, *Tetrahedron*, 2001, **57**, 4719-4728 (*synth*)

Araplysin I A-1378
 N-[3-[4-(2-Aminoethyl)-2,6-dibromophe-noxy]propyl]-7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxamide, 9CI. Hexa-dellin A
 [129313-98-0]



$C_{21}H_{23}Br_4N_3O_5$ 717.046
 Relative configs. only known in this series. The compds. covered by this entry have not been correlated. Identity of Araplysin I and Hexadellin A not confirmed. Isol. from the sponge *Psammaphysilla arabica*. ATPase inhibitor. Shows antimicrobial props. Amorph. Mp 140-142°. $[\alpha]_D -70$ (c, 0.7 in MeOH).

N²⁰-Sulfo: Araplysin I N²⁰-sulfamate
 [945408-71-9]
 $C_{21}H_{23}Br_4N_3O_8S$ 797.11
 Isol. from *Aplysina fulva*. Amorph.

solid (as Na salt). $[\alpha]_D^{23} +100$ (c, 0.06 in MeOH) (Na salt). λ_{\max} 206 (log ϵ 4.7); 278 (log ϵ 3.84) (MeOH) (Na salt).

N^{20} -(13-Methyltetradecanoyl): **Araplysin II**

[129277-05-0]

$C_{36}H_{51}Br_4N_3O_6$ 941.431

From the sponge *Psammaplysilla arabica*. ATPase inhibitor. Shows antimicrobial props. Amorph. Mp 40-42°.

$[\alpha]_D -38$ (c, 0.73 in $CHCl_3$). λ_{\max} 270 (ϵ 11650) (MeOH) (Berdy).

N^{20}, N^{20} -Di-Me: **Purealidin P**

[167394-81-2]

$C_{23}H_{27}Br_4N_3O_5$ 745.099

Alkaloid from the Okinawan sponge *Psammaplysilla purea*. Cytotoxic against murine lymphoma and human KB carcinoma cells. Oil (as trifluoroacetate). Sol. MeOH, EtOAc; poorly sol. H_2O . $[\alpha]_D +6.6$ (c, 0.75 in MeOH). λ_{\max} 277 (ϵ 1700); 284 (ϵ 1400) (MeOH) (Berdy).

N^{20}, N^{20}, N^{20} -Tri-Me: **Purealidin B**

[138590-55-3]

$C_{24}H_{30}Br_4N_3O_5^{\oplus}$ 760.134

Quaternary alkaloid from *Psammaplysilla purea* and *Pseudoceratina verrucosa*. Antibacterial and cytotoxic. Amorph. solid (prob. as trifluoroacetate). $[\alpha]_D^{18} -4.5$ (c, 1.3 in MeOH). λ_{\max} 220 (ϵ 10000); 284 (ϵ 1000) (MeOH) (Berdy).

14-Debromo: **14-Debromoaraplysin I**

[136685-29-5]

$C_{21}H_{24}Br_3N_3O_5$ 638.15

From the sponge *Psammaplysilla purea*. Glass. $[\alpha]_D +21$ (c, 0.47 in MeOH/ $CHCl_3$). λ_{\max} 240 (ϵ 23000); 287 (ϵ 16700) (MeOH) (Berdy).

19-Hydroxy, N^{20} -Ac: **Aplysinamisine III**

[150417-69-9]

$C_{23}H_{25}Br_4N_3O_7$ 775.082

Isol. from the Caribbean sponge *Aplysina cauliformis*. Semisol. $[\alpha]_D^{26} +69$ (c, 6.4 in MeOH). λ_{\max} 218 (ϵ 14500); 282 (ϵ 5200) (MeOH) (Berdy).

[121135-00-0]

Morris, S.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 677 (*Hexadellin A*, *isol*, *ir*, *pmr*, *cmr*, *ms*)

Longeon, A. *et al.*, *Experientia*, 1990, **46**, 548 (*Araplysin I*)

James, D.M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1137 (*14-Debromoaraplysin I*)

Kobayashi, J. *et al.*, *Tetrahedron*, 1991, **47**, 6617 (*Purealidin B*, *isol*, *pmr*, *cmr*, *uv*, *ir*, *ms*, *cd*)

Rodriguez, A.D. *et al.*, *J. Nat. Prod.*, 1993, **56**, 907 (*Aplysinamisine III*)

Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403 (*Purealidin P*)

Rogers, E.W. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1191-1194 (*sulfamate*)

Araplysin III

A-1379

[245436-90-2]

As Araplysin I, A-1378 with

R = COOH


$C_{22}H_{23}Br_4N_3O_7$ 761.056

Isol. from the sponge *Aiolochoira crassa*. Amorph. solid. $[\alpha]_D +96.3$ (c, 0.19 in MeOH). λ_{\max} 208 (ϵ 8100); 284 (ϵ 1200) (EtOH).

Gao, H. *et al.*, *Tetrahedron*, 1999, **55**, 9717-9726

Arboflorine

A-1380



$C_{19}H_{21}N_3O$ 307.394


Modified Tubotaiwine-type alkaloid. Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D +71$ (c, 0.07 in $CHCl_3$). λ_{\max} 224; 283; 291 (EtOH).

Lim, K.-H. *et al.*, *Org. Lett.*, 2006, **8**, 1733-1735 (*isol*, *pmr*, *cmr*)

Arbolosine

A-1381

[926927-38-0]



Absolute Configuration

$C_{20}H_{24}N_2O_3$ 340.421


Highly rearranged aspidosperma-type skeleton. Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D +137$ (c, 0.15 in $CHCl_3$). λ_{\max} 210 (log ϵ 3.83); 247 (log ϵ 4.14); 265 (log ϵ 3.99); 321 (log ϵ 3.38) (EtOH).

Lim, K.-H. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 31-35 (*isol*, *pmr*, *cmr*, *ms*)

Arbophylline

A-1382

[917600-41-0]



Relative Configuration

$C_{21}H_{22}N_2O_4$ 366.416


Sarpagine-related skeleton. Alkaloid from the leaves of *Kopsia arborea*. Oil. $[\alpha]_D +2$ (c, 0.16 in $CHCl_3$). λ_{\max} 210; 242; 297 (EtOH).

Lim, K.-H. *et al.*, *Tet. Lett.*, 2006, **47**, 8653-8655 (*isol*, *pmr*, *cmr*)

Arborescidine C

A-1383

[147395-94-6]



$C_{16}H_{19}BrN_2O$ 335.243

Alkaloid from the marine tunicate *Pseudodistoma arborescens*. Cryst. (MeOH). Mp 172-173°. $[\alpha]_D +3$ (c, 1 in $CHCl_3$).

17-Epimer: **Arborescidine D**

[147512-45-6]

$C_{16}H_{19}BrN_2O$ 335.243

Alkaloid from the marine tunicate *Pseudodistoma arborescens*. Exhibits moderate cytotoxicity *in vitro* against KB human buccal carcinoma.

Amorph. solid. λ_{\max} 231 (ϵ 29800); 285 (ϵ 8000) (MeOH) (Berdy).

17-Deoxy, 16,17-didehydro: **Arborescidine B**

[147395-93-5]

$C_{16}H_{17}BrN_2$ 317.228

Alkaloid from *Pseudodistoma arborescens*. Oil. $[\alpha]_D +70$ (c, 0.6 in $CHCl_2$).


Chbani, M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 99-104 (*isol*, *cd*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Burm, B.E.A. *et al.*, *Tetrahedron*, 1998, **54**, 6135-6146 (*synth*, *pmr*, *cmr*)

Santos, L.S. *et al.*, *J.O.C.*, 2004, **69**, 1283-1289 (*synth*)

Arboreumine

A-1384



$C_{25}H_{40}N_2O_5$ 448.601


Alkaloid from the leaves of *Piper arboreum*. Antifungal agent. Amorph. solid. λ_{\max} 240 (ϵ 2607) (MeOH).

Vasques da Silva, R. *et al.*, *Phytochemistry*, 2002, **59**, 521-527 (*isol*, *pmr*, *cmr*)

Arboricine†

A-1385

[929606-10-0]



$C_{19}H_{22}N_2O$ 294.396


Alkaloid from the leaves of *Kopsia arborea*. Prisms. Mp 126-128°. $[\alpha]_D -110$ (c, 0.14 in $CHCl_3$). λ_{\max} 225; 282; 290 (EtOH).

Lim, K.-H. *et al.*, *Tet. Lett.*, 2007, **48**, 1143-1145 (*isol*, *pmr*, *cmr*)

Arboricine

A-1386


[929606-12-2]



C₁₉H₂₂N₂O 294.396
Alkaloid from the leaves of *Kopsia arborea*. Oil. $[\alpha]_D^{25}$ -18 (c, 0.14 in CHCl₃).
Lim, K.-H. *et al.*, *Tet. Lett.*, 2007, **48**, 1143-1145 (*isol*, *pmr*, *cmr*)

Archaeosine A-1387


2-Amino-4,7-dihydro-4-oxo-7-β-D-ribofuranosyl-1H-pyrrolo[2,3-d]pyrimidine-5-carboximidamide. 7-Formamidino-7-deazaguanosine
[148608-52-0]



C₁₂H₁₆N₆O₅ 324.296
Constit. of archaeal transfer RNA.
Gregson, J.M. *et al.*, *J. Biol. Chem.*, 1993, **268**, 10076 (*isol*, *synth*)

Archazolid A A-1388

[149633-98-7]



C₄₂H₆₂N₂O₇S 739.027
Prod. by *Archangium gephyra* and *Cystobacter violaceus*. V-ATPase inhibitor. Antifungal and antineoplastic agent. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 228 (ε 39810) (MeOH) (Berdy).

7-O-β-D-Glucopyranoside: **Archazolid C**
C₄₈H₇₂N₂O₁₂S 901.169
Isol. from *Cystobacter violaceus* Cb vil05. Amorph. solid. $[\alpha]_D^{25}$ -33.9 (c, 6.2 in MeOH). λ_{\max} 228 (log ε 4.6) (MeOH).

26-Hydroxy, 7-O-β-D-glucopyranoside: **Archazolid D**
[945027-76-9]
C₄₈H₇₂N₂O₁₃S 917.168
Prod. by *Cystobacter violaceus* strain Cb vil05. V-ATPase inhibitor. Amorph. solid. $[\alpha]_D^{25}$ -26.8 (c, 0.22 in MeOH). λ_{\max} 231 (MeOH).

Ger. Pat., 1993, 4 142 951; *CA*, **119**, 137527 (*Archazolid A*)

Sasse, F. *et al.*, *J. Antibiot.*, 2003, **56**, 520-525 (*Archazolid A*, *activity*)


Hassfeld, J. *et al.*, *Org. Lett.*, 2006, **8**, 4751-4754 (*Archazolid A*, *abs config*)
Menche, D. *et al.*, *Eur. J. Org. Chem.*, 2007, 1196-1202 (*Archazolid C*)
Menche, D. *et al.*, *J. Antibiot.*, 2007, **60**, 328-331 (*Archazolid D*)
Menche, D. *et al.*, *J.A.C.S.*, 2007, **129**, 6100-6101 (*synth*)

Archazolid B A-1389

As Archazolid A, A-1388 with R = H

C₄₁H₆₀N₂O₇S 725
Prod. by *Archangium gephyra*. V-ATPase inhibitor. Cytotoxic.

Sasse, F. *et al.*, *J. Antibiot.*, 2003, **56**, 520-525 (*isol*, *activity*)
Hassfeld, J. *et al.*, *Org. Lett.*, 2006, **8**, 4751-4754 (*pmr*, *abs config*)
Roethle, P.A. *et al.*, *J.A.C.S.*, 2007, **129**, 8960-8961 (*synth*)


Archerine A-1390

C₃₂H₃₆Br₄N₁₀O₈ 1008.315
Isol. from *Aplysina archeri*. Amorph. brown solid. $[\alpha]_D^{25}$ +111.4 (c, 0.07 in MeOH). λ_{\max} 221 (ε 17600); 279 (ε 9500) (MeOH).

Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2001, 55-60

Arcutinidine A-1391

[441064-17-1]



C₂₀H₂₇NO₂ 313.439

15-O-(2-Methylpropanoyl): **Arcutinine**
[441064-16-0]

C₂₄H₃₃NO₃ 383.53
Alkaloid from the aerial parts of *Aconitum arcuatum*.

15-O-(2-Methylbutanoyl): **Arcutine**
[336819-46-6]


C₂₅H₃₅NO₃ 397.556
Alkaloid from the aerial parts of *Aconitum arcuatum*. Cryst. (hexane). Mp 225-226°.

Tashkhodzhaeva, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 79-83 (*Arcutine*)

Saidkhodzhaeva, S.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 466-469 (*Arcutinine*)

Arcyriacyanin A A-1392

5,9-Dihydro-1H-pyrrolo[3',4':6,7]cyclohepta[2,1-b:4,3-c'd']indole-1,3(2H)-dione, 9CI
[188548-81-4]



C₂₀H₁₁N₃O₂ 325.326
Pigment from *Arcyria nutans*. Cytotoxic agent. Protein kinase C and protein tyrosine kinase inhibitor. Green-blue solid. Mp > 300°. λ_{\max} 224 (log ε 4.06); 243 (log ε 4.15); 264 (sh) (log ε 3.88); 360 (log ε 3.72); 625 (log ε 2.49) (MeOH).


4c,8a-Dihydro (cis-): **Dihydroarcyriacyanin A**

C₂₀H₁₃N₃O₂ 327.342
Pigment from *Arcyria nutans* and *Arcyria obvelata*. Amorph. powder. λ_{\max} 344 (ε 38000); 363 (ε 32000) (MeOH).

Gill, M. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1987, **51**, 216 (*rev*, *uv*)
Brenner, M. *et al.*, *Chem. Eur. J.*, 1997, **3**, 70-74 (*synth*, *uv*, *ir*, *pmr*, *cmr*)
Murase, M. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 889-892; 2000, **48**, 81-84 (*synth*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *activity*)
Kamata, K. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1252-1254 (*Dihydroarcyriacyanin A*)
Kraus, G.A. *et al.*, *Org. Lett.*, 2008, **10**, 3061-3063 (*synth*)

Arcyriaflavin A A-1393

12,13-Dihydro-5H-indolo[2,3-a]pyrrolo[3,4-c]carbazole-5,7(6H)-dione, 9CI
[118458-54-1]



C₂₀H₁₁N₃O₂ 325.326
Minor pigment from the fruiting bodies of the slime mould *Arcyria denudata*. Also isol. from the tunicate *Eudistoma* sp. and from *Nocardioopsis* sp. Orange solid. Mp 360°. λ_{\max} 235 (log ε 4.8); 257 (log ε 4.42); 272 (log ε 4.45); 281 (log ε 4.55); 300 (sh) (log ε 4.64); 314 (log ε 4.86); 402 (log ε 3.81) (MeOH).

N¹²-β-D-Glucopyranosyl: **Antibiotic BMY 41219**. BMY 41219. Didechloro-4'-O-demethylbeccamycin
[158204-55-8]

C₂₆H₂₁N₃O₇ 487.468
Prod. by *Saccharothrix aerocoloni-genes*. Orange-yellow powder. Sol.

MeOH, DMSO, THF; poorly sol.
H₂O. λ_{\max} 236 (ϵ 29970); 284 (ϵ 17750);
316 (ϵ 37420); 404 (ϵ 3020) (MeOH).

Deoxo: see Antibiotic K 252c, A-1190

2-Hydroxy: Arcyriaflavin B

[73697-64-0]
C₂₀H₁₁N₃O₃ 341.325

Isol. from *Arcyria denudata* and *Metatrichia vesparium*. Pale yellow. Mp 350°. λ_{\max} 229 (log ϵ 4.07); 271 (log ϵ 3.7); 280 (log ϵ 3.75); 323 (log ϵ 4.11); 414 (log ϵ 3.19) (MeOH).

1,11-Dihydroxy: Antibiotic BE 13793C.

BE 13793C. Antibiotic J 104303. J 104303

[133805-03-5]

C₂₀H₁₁N₃O₄ 357.325

Prod. by *Streptomyces mobaraensis* sp. BA-13793. Antitumour agent.

Topoisomerase inhibitor. Yellow-orange powder. Sol. THF, DMSO; fairly sol. MeOH; poorly sol. H₂O. Mp 295°. λ_{\max} 245 (ϵ 63500); 298 (ϵ 38100); 307 (ϵ 37400); 325 (sh) (ϵ 21900); 430 (ϵ 5600) (MeOH) (Derep).

► NM3457000

2,3-Dihydroxy: 2,3-Dihydroxyarcyriaflavin A

C₂₀H₁₁N₃O₄ 357.325

Isol. from *Lycogala epidendrum*. Cytotoxic. Amorph. yellow solid. λ_{\max} 230 (log ϵ 10.5); 284 (log ϵ 9.8); 330 (log ϵ 10.5) (MeOH).

2,9-Dihydroxy: Arcyriaflavin D

[179077-55-5]

C₂₀H₁₁N₃O₄ 357.325

Pigment from *Dictydiaethalium plumbeum*. Yellow cryst. powder. Mp > 300°.

2,10-Dihydroxy: Arcyriaflavin C

[73697-65-1]

C₂₀H₁₁N₃O₄ 357.325

Isol. from *Arcyria denudata*, *Metatrichia vesparium* and *Streptosporangium vulgare* K254. Enzyme inhibitor, calmodulin antagonist, shows insecticidal props. Pale yellow. Mp 350°. λ_{\max} 229 (ϵ 13800); 255 (sh) (ϵ 5890); 270 (ϵ 4570); 280 (ϵ 5010); 318 (sh) (ϵ 12300); 331 (ϵ 19500); 422 (ϵ 2190) (MeOH) (Derep).

Steglich, W. et al., *Angew. Chem., Int. Ed.*, 1980, **19**, 459-460 (*Arcyriaflavins*, *isol*, *uv*, *ir*, *pmr*, *struct*)

Bergman, J. et al., *J.O.C.*, 1989, **54**, 824-828 (*synth*, *ir*, *pmr*)

Steglich, W. et al., *Pure Appl. Chem.*, 1989, **61**, 281-288 (*Arcyriaflavin D*)

U.S. Pat., 1990, 5 326 754; CA, **121**, 221985u (BMV 41219)

Kojiri, K. et al., *J. Antibiot.*, 1991, **44**, 723-728 (BE 13793C)

Horton, P.A. et al., *Experientia*, 1994, **50**, 843-845 (*isol*)

Ohkubo, M. et al., *Tetrahedron*, 1996, **52**, 8099-8112 (*synth*)

Marques, M.M.B. et al., *Tet. Lett.*, 1999, **40**, 3795-3796 (*synth*)

Ohuchi, T. et al., *J. Antibiot.*, 2000, **53**, 393-403 (BE13793C, *biosynth*)

Bergman, J. et al., *J.C.S. Perkin 1*, 2000, 2609-2614 (*synth*)

Marques, M.M.B. et al., *Tet. Lett.*, 2000, **41**, 9835-9838 (*synth*)

Hosoya, T. et al., *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2776-2780 (2,3-Dihydroxyarcyriaflavin A)


Alonso, D. et al., *Tet. Lett.*, 2005, **46**, 4839-4841 (*synth*)

Hinze, C. et al., *Synthesis*, 2007, 608-612 (*synth*)

Arcyriarubin A A-1394

3,4-Di-1H-indol-3-yl-1H-pyrrole-2,5-dione, 9CI. 2,3-Bis(1H-indol-3-yl)maleimide

[119139-23-0]



C₂₀H₁₃N₃O₂ 327.342

Red pigment from the fruiting bodies of the slime mould *Arcyria denudata*. Protein kinase C inhibitor. Mp 161° Mp 281°.

1-Me: Ro 31-6045

[113963-68-1] Negative control for protein kinase C inhibitory activity. Solid (Me₂CO). Mp 278°.

6'-Hydroxy: Arcyriarubin B

[73697-62-8]

C₂₀H₁₃N₃O₃ 343.341

From *Arcyria denudata*. Red cryst. (MeOH). Mp 177° (154-155°). λ_{\max} 281 (ϵ 8510); 392 (sh) (ϵ 3720); 465 (ϵ 5890) (MeOH) (Derep).

6'-Sulfooxy: Arcyriarubin B 6'-sulfate

C₂₀H₁₃N₃O₆S 423.405

Isol. from *Arcyria denudata*. Amorph. powder. λ_{\max} 278 (ϵ 37000); 374 (ϵ 15000) (MeOH).

6'-Hydroxy, 3,4-dihydro: Dihydroarcyriarubin B

C₂₀H₁₅N₃O₃ 345.357

From *Arcyria denudata*. Cryst. Mp 250° dec. Prob. the *trans*-isomer.

6',6''-Dihydroxy: 3,4-Bis(6-hydroxy-1H-indol-3-yl)-1H-pyrrole-2,5-dione, 9CI.

Arcyriarubin C

[73697-63-9]

C₂₀H₁₃N₃O₄ 359.34

Red pigment from *Arcyria denudata* and *Arcyria ferruginea*. Mp 205-206°. λ_{\max} 283 (ϵ 8510); 474 (ϵ 5750) (MeOH) (Derep).

6',6''-Dihydroxy, 3 α ,4 β -dihydro:

Dihydroarcyriarubin C

C₂₀H₁₅N₃O₄ 361.356

From *Arcyria ferruginea*. Cell-cycle inhibitor. Amorph. solid. Possesses *trans*-config. λ_{\max} 221 ; 274 (no solvent reported).

Gill, M. et al., *Prog. Chem. Org. Nat. Prod.*, 1987, **51**, 218 (*isol*)

Brenner, M. et al., *Tetrahedron*, 1988, **44**, 2887-2892 (*synth*, 1-Me)

Davis, P.D. et al., *J. Med. Chem.*, 1992, **35**, 177-184 (*synth*, *cryst struct*, *pharmacol*)

Faul, M.M. et al., *Synthesis*, 1995, 1511-1516 (*synth*, *pmr*, *cmr*, *ir*, *uv*, 1-Me)

Marmy-Conus, N. et al., *FEBS Lett.*, 2002, **519**, 135-140 (1-Me, *pharmacol*)


Nakatani, S. et al., *Bioorg. Med. Chem. Lett.*, 2003, **13**, 2879-2881 (*Dihydroarcyriarubin C*)

Kamata, K. et al., *J. Nat. Prod.*, 2006, **69**, 1252-1254 (*Arcyriarubin B 6'-sulfate*)

Kaniwa, K. et al., *Bioorg. Med. Chem. Lett.*, 2007, **17**, 4254-4257 (*Dihydroarcyriarubin C*, *synth*, *config*)

Arcyriaverdin C

A-1395



C₂₀H₁₁N₃O₆ 389.323


Green pigment from *Arcyria denudata*. Mp > 300°. λ_{\max} 256 (sh) (log ϵ 3.91); 336 (log ϵ 3.51); 433 (log ϵ 3.69); 634 (log ϵ 2.82) (MeOH).

Gill, M. et al., *Prog. Chem. Org. Nat. Prod.*, 1987, **51**, 216 (*rev*, *uv*)

Arcyirin A

A-1396

2,9-Dihydro-1H-indolo[3,2-j]pyrrolo[2,3,4-gh]phenanthridine-1,8-diol



C₂₀H₁₃N₃O₂ 327.342

Systematic numbering shown. Isol. from *Arcyria* sp. No phys. props. reported. Not indexed by CAS.

1-Deoxy, 5-hydroxy: Arcyirin B


C₂₀H₁₃N₃O₂ 327.342

Isol. from *Arcyria* sp. No phys. props. reported. Not indexed by CAS.

Steglich, W. et al., *Pure Appl. Chem.*, 1989, **61**, 281-288 (*occur*)

Arcyroxindole A

A-1397




C₂₀H₁₁N₃O₃ 341.325

Orange pigment from *Arcyria denudata*. Mp 240° (dec.). λ_{\max} 247 (sh) (log ϵ 4.08); 280 (sh) (log ϵ 3.89); 341 (sh) (log ϵ 3.43); 493 (log ϵ 3.45) (MeOH).

Gill, M. et al., *Prog. Chem. Org. Nat. Prod.*,

1987, **51**, 216 (rev, uv)**Arcyroxocin A** A-1398

5,10-Dihydro-1H-pyrrolo[3',4':4,5]oxocino[2,3-b:6,7,8-c'd']diindole-1,3(2H)-dione, 9CI [179632-33-8]



$C_{20}H_{11}N_3O_3$ 341.325

Red pigment from the slime mould *Arcyria denudata*. Red cryst. Mp > 300°. λ_{max} 276 (log ϵ 4.15); 285 (log ϵ 4.12); 364 (log ϵ 3.76); 465 (log ϵ 3.91) (MeOH).

4c,4d-Dihydro: Dihydroarcyroxocin A

$C_{20}H_{13}N_3O_3$ 343.341

Pigment from *Arcyria nutans*. No phys. props reported. Possesses *cis*-config.

2-Hydroxy: Arcyroxocin B

$C_{20}H_{11}N_3O_4$ 357.325

Pigment from *Arcyria denudata*. Amorph. powder. Mp 235° (dec.). λ_{max} 212 (ϵ 91000); 474 (ϵ 10000) (MeOH).

N⁶-Hydroxy: Arcyroxepin A

[73697-66-2]

$C_{20}H_{11}N_3O_4$ 357.325

Red pigment from fruiting bodies of the slime mould *Arcyria denudata*. Mp 268-270°. Originally assigned an erroneous struct. λ_{max} 226 (log ϵ 4.44); 273 (log ϵ 3.82); 283 (sh) (log ϵ 3.83); 362 (log ϵ 3.51); 471 (log ϵ 3.68) (MeOH).

N⁶,2-Dihydroxy: Arcyroxepin B. N-Hydroxyarcyroxocin B

$C_{20}H_{11}N_3O_5$ 373.324

Violet pigment from *Arcyria denudata*. Originally assigned an erroneous struct. Not indexed by CA to 2001. λ_{max} 274; 388; 484 (MeOH).

Steglich, W. *et al.*, *Angew. Chem., Int. Ed.*, 1980, **19**, 459-460 (*Arcyroxepin A*)

Gill, M. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1987, **51**, 216-297 (*isol, uv, struct, Arcyroxepin B, Arcyroxocin B*)

Steglich, W. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 281-288 (*Dihydroarcyroxocin A*)


Gribble, G.W. *et al.*, *Stud. Nat. Prod. Chem.*, 1993, **12**, 365-409 (*struct, Arcyroxepin A*)

Mayer, G. *et al.*, *Tet. Lett.*, 1996, **37**, 4483-4486 (*synth*)

Kamata, K. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1252-1254 (*Arcyroxocin B*)

Aardeemin A-1399

[148441-25-2]



Absolute Configuration

$C_{26}H_{26}N_4O_2$ 426.517

Prod. by *Aspergillus fischeri* var. *brasiliensis*. Antitumour drug resistance reversing agent. Mycotoxin. Cytotoxicity enhancer. Amorph. solid. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{25}$ -92 (c, 0.24 in MeOH). Related to Asperlicin E, A-1481. λ_{max} 210 (ϵ 12600); 224 (ϵ 7400); 268 (ϵ 2700); 274 (ϵ 2600); 302 (ϵ 1700); 314 (ϵ 2600) (MeOH) (Derep).

5-N-Ac: 5-N-Acetylardeemin

[148441-26-3]

$C_{28}H_{28}N_4O_3$ 468.554

Prod. by *Aspergillus fischeri* var. *brasiliensis*. Antitumour drug resistance reversing agent. Shows MDR modulatory props. Mycotoxin. Cryst. (MeOH). Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 226-228°. $[\alpha]_D^{25}$ -33 (c, 0.78 in MeOH). λ_{max} 210 (ϵ 10700); 224 (ϵ 10700); 264 (ϵ 4300); 274 (ϵ 4000); 302 (ϵ 1700); 314 (ϵ 1360) (MeOH) (Derep).

15b-Hydroxy, 5-N-Ac: 5-N-Acetyl-15b-hydroxyardeemin

[148717-81-1]

$C_{28}H_{28}N_4O_4$ 484.554

Prod. by *Aspergillus fischeri* var. *brasiliensis*. Antitumour drug resistance reversing agent. Mycotoxin. Amorph. solid. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{25}$ -245 (c, 0.21 in MeOH). λ_{max} 210 (ϵ 10700); 224 (ϵ 10700); 264 (ϵ 4300); 274 (ϵ 4000); 302 (ϵ 1700); 314 (ϵ 1360) (MeOH) (Derep).

Karwowski, J.P. *et al.*, *J. Antibiot.*, 1993, **46**, 374-379; 380-386 (*isol, struct, props, activity*)

Marsden, S.P. *et al.*, *J.A.C.S.*, 1994, **116**, 11143-11140 (*synth, 5-N-Acetylardeemin*)


Chou, T.-C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1998, **95**, 8369-8374 (*activity*)

Depew, K.M. *et al.*, *J.A.C.S.*, 1999, **121**, 11953-11963 (*synth*)

Arecolidine A-1400

1,2-Dihydro-3,4-dimethoxy-1-methylpyridine, 9CI

[57680-57-6]



$C_8H_{13}NO_2$ 155.196

Although arecolidine has been referred to in the recent lit., the struct. proposed for it, if it is a genuine alkaloid, is very poorly founded. Recent synthetic studies show that the stability of compds. decreases the closer the target struct. is approached. Minor alkaloid from *Areca catechu* (betel) nuts (Arecaceae). Needles (Et_2O or by subl.). Mp 110°.

Hydrochloride: Mp 250°.

Hydrobromide: Mp 268-271°.


Emde, H. *et al.*, *Chem. Zentralbl.*, 1915, 1381 (*isol*)

Marion, L. *et al.*, *Alkaloids (Academic Press)*, 1949, **1**, 175

Dehmlow, E.V. *et al.*, *Heterocycles*, 1994, **37**, 355-366

Arenaine

[35471-10-4]



Relative Configuration


$C_{11}H_{17}N_3O$ 207.275

Alkaloid from seeds of *Plantago arenaria* (sand plantain) (Plantaginaceae). Mp 208-210°. $[\alpha]_D^{22}$ +305 (c, 1.7 in $CHCl_3$).

Rabaron, A. *et al.*, *J.A.C.S.*, 1971, **93**, 6270-6271 (*uv, ir, pmr, cmr, struct*)

Arenarine B A-1402

α -(Methoxymethyl)-9H-pyrido[3,4-b]indole-1-methanol, 9CI. 1-(1-Hydroxy-2-methoxyethyl)- β -carboline [123520-95-6]



$C_{14}H_{14}N_2O_2$ 242.277

Alkaloid from the whole plant of *Arenaria kansuensis* (Caryophyllaceae). Pale-yellow needles (EtOAc). Mp 157-158°.

1'-Ketone: 2-Methoxy-1-(9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI. 1-(Methoxyacetyl)- β -carboline. **Arenarine A** [123520-94-5]

$C_{14}H_{12}N_2O_2$ 240.261


Alkaloid from *Arenaria kansuensis* (Caryophyllaceae). Yellow needles (EtOAc). Mp 182-183°.

Wu, F. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1808 (*isol, uv, ir, pmr, cmr, ms, struct*)

Bracher, F. *et al.*, *J. Het. Chem.*, 2004, **41**, 173-176 (*synth, ir, pmr, cmr, ms, 1'-ketone*)

Arenine

[68676-55-1]



$C_{23}H_{24}N_2O_4$ 392.454

(R)-form

Me ether: Synthetic. Mp 95-100°. $[\alpha]_D^{25}$ +37 (c, 0.9 in $CHCl_3$).

(S)-form

Alkaloid from *Papaver arenarium* (Papa-

veraceae). Noncryst. $[\alpha]_D^{25}$ -31 (c, 1.03 in MeOH).

Me ether: Macrostromine

[53912-94-0]

$C_{24}H_{26}N_2O_4$ 406.48

Alkaloid from *Papaver macrostomum* (Papaveraceae). Mp 107-110°. $[\alpha]_D^{25}$ -51 (c, 0.89 in $CHCl_3$).

Me ether, methiodide: Mp 220-230° dec.

Me ether, N-oxide (α): Macrostromine α -N-oxide

[90138-27-5]

$C_{24}H_{26}N_2O_5$ 422.48

Alkaloid from *Papaver arenarium*. Cryst. (Me₂CO). Mp 141-142°. $[\alpha]_D^{25}$ +68 (c, 0.8 in MeOH).

Me ether, N-oxide (β): Macrostromine β -N-oxide

$C_{24}H_{26}N_2O_5$ 422.48

Alkaloid from *Papaver arenarium*. Amorph. $[\alpha]_D^{25}$ +35 (c, 0.8 in MeOH).

Mnatsukanyan, V.A. *et al.*, *Tet. Lett.*, 1974, 851 (*Macrostromine*)

Mnatsukanyan, V.A. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 1421 (*isol, uv, pmr, ms, struct, Macrostromine*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 417; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 358 (*uv, ir, pmr, ms, struct*)

Wykypiel, W. *et al.*, *Tet. Lett.*, 1980, 1927 (*synth, Macrostromine*)

Sharma, R.B. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 141 (*synth, Macrostromine*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 76-79; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 71-73 (*Macrostromine N-oxides*)


Brunner, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 73 (*synth, pmr, Macrostromine*)

Mahboobi, S. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 275 (*synth, pmr, Macrostromine*)

Arenosclerin A

A-1404

[289041-16-3]



$C_{32}H_{54}N_2O$ 482.791

Alkaloid from the sponge *Arenosclera brasiliensis*. Glassy solid. $[\alpha]_D^{25}$ -3 (c, 0.01 in MeOH). λ_{max} 235 (€ 15100) (MeOH).

Deoxy: Haliconacyclamine E

[289041-15-2]

$C_{32}H_{54}N_2$ 466.792

Alkaloid from *Arenosclera brasiliensis*. Glassy solid. $[\alpha]_D^{25}$ +14 (c, 0.02 in MeOH). λ_{max} 236 (€ 14900) (MeOH).

Deoxy, 23,24-dihydro: Halichondramine

$C_{32}H_{56}N_2$ 468.808

Alkaloid from *Halichondria* sp. Yellow oil. $[\alpha]_D^{25}$ +3.3 (c, 0.5 in MeOH).

2,3-Diepimer: Arenosclerin C

[289041-18-5]

$C_{32}H_{54}N_2O$ 482.791

Alkaloid from *Arenosclera brasiliensis*. Glassy solid. $[\alpha]_D^{25}$ -17 (c, 0.02 in MeOH). λ_{max} 235 (€ 15150) (MeOH).

7,9-Diepimer, deoxy: Haliconacyclamine F

[934986-37-5]

$C_{32}H_{54}N_2$ 466.792

Alkaloid from *Pachychalina alcaloidifera*. Glassy solid. $[\alpha]_D^{25}$ +5.4 (c, 0.004 in MeOH). λ_{max} 226 (€ 2450) (MeOH).

2,3,7-Triepimer: Arenosclerin B

[289041-17-4]

$C_{32}H_{54}N_2O$ 482.791

Alkaloid from *Arenosclera brasiliensis*. Glassy solid. $[\alpha]_D^{25}$ +8.6 (c, 0.008 in MeOH). λ_{max} 235 (€ 15000) (MeOH).

Stereoisomer: Arenosclerin D

[934986-38-6]

$C_{32}H_{54}N_2O$ 482.791

Alkaloid from *Pachychalina alcaloidifera*. Glassy solid. $[\alpha]_D^{25}$ +6.9 (c, 0.01 in MeOH). Prob. 9-epimer or 7,9-diepimer of Arenosclerin A. λ_{max} 236 (€ 28456) (MeOH).

Torres, Y.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1098-1105 (*isol*)

Chill, L. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1738-1741 (*Halichondramine*)


Torres, Y.R. *et al.*, *Toxicol.*, 2002, **40**, 885-891 (*props*)

De Oliveira, J.H.H.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 538-543 (*Arenosclerin D, Haliconacyclamide F*)

Arenosclerin E

A-1405

[934986-39-7]



$C_{32}H_{54}N_2O$ 482.791


Alkaloid from *Pachychalina alcaloidifera*. Cytotoxic. Glassy solid. $[\alpha]_D^{25}$ +14.5 (c, 0.01 in MeOH). λ_{max} 235 (€ 23780) (MeOH).

De Oliveira, J.H.H.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 538-543 (*isol, pmr, cmr, ms*)

Arestrictin A

A-1406

[929101-59-7]



Absolute Configuration

$C_{29}H_{39}N_3O_3$ 477.645

Prod. by *Aspergillus penicilloides* and *Aspergillus restrictus*. Pale yellow cryst. (MeOH). Mp 209°. λ_{max} 228 (log € 4.62);

274 (log € 3.95); 283 (log € 3.92); 291 (log € 3.82) (MeOH).

8-Deoxy: Arestrictin B

[929101-60-0]

$C_{29}H_{39}N_3O_2$ 461.646

Prod. by *Aspergillus penicilloides* and *Aspergillus restrictus*. Cryst. (MeOH). Mp 228°. $[\alpha]_D^{25}$ -36.8 (c, 0.51 in $CHCl_3$). λ_{max} 230 (log € 4.62); 278 (log € 3.96); 283 (log € 3.96); 294 (log € 3.86) (MeOH).


Itabashi, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 1639-1641 (*isol, cd, pmr, cmr*)

Argadin

A-1407

Antibiotic FO 7314. FO 7314

[289665-92-5]



Absolute Configuration

$C_{29}H_{42}N_{10}O_9$ 674.712

Prod. by *Clonostachys* sp. FO-7314. Chitinase inhibitor. Powder. Mp 270° dec. $[\alpha]_D^{25}$ +52.1 (c, 0.1 in H_2O). λ_{max} 200 (€ 31400) (H_2O).


Arai, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1442-1446 (*isol, pmr, cmr*)

Dixon, M.J. *et al.*, *Eur. J. Org. Chem.*, 2006, 5002-5006; 5602 (*synth, stereochem*)

Argemexicaine A

A-1408

[624736-05-6]



$C_{21}H_{23}NO_5$ 369.416


Alkaloid from the whole plant of *Argemone mexicana*. Cryst. (hexane). Mp 158-160°. λ_{max} 225 (log € 4.35); 290 (log € 4.21) (EtOH).

Chang, Y.-C. *et al.*, *Planta Med.*, 2003, **69**, 148-152 (*isol, pmr, cmr, ms*)

Argemexicaine B

A-1409

[624736-06-7]



C₂₁H₂₃NO₅ 369.416


Alkaloid from the whole plant of *Argemone mexicana*. Powder. Mp 131-133°. λ_{\max} 225 (log ϵ 4.33); 290 (log ϵ 4.04) (EtOH).

Chang, Y.-C. *et al.*, *Planta Med.*, 2003, **69**, 148-152 (*isol*, *pmr*, *cmr*, *ms*)

Argemone

A-1410

5,6,11,12-Tetrahydro-2,3,8,9-tetra-methoxy-13-methyldibenzo[a,e]cycloocten-5,11-imine, 9CI. 2,3,8,9-Tetramethoxypavinane. N-Methylpavine



(-)-form

C₂₁H₂₅NO₄ 355.433**(+)-form** [16584-62-6]

Alkaloid from *Leontice smirnovii*, also obt. by resoln. (Leonticeaceae). Shows weak *in vitro* antimicrobial activity against *Mycobacterium smegmatis*. Cryst. (CHCl₃). Mp 152-153°. $[\alpha]_D^{25}$ +218 (c, 2.11 in EtOH).

(-)-form [6901-16-2]

Alkaloid from *Argemone gracilentia*, other *Argemone* spp., *Berberis buxifolia*, *Thalictrum revolutum* and *Thalictrum strictum* (Papaveraceae, Berberidaceae, Ranunculaceae). Mp 155.5-156.5° (147-148°). $[\alpha]_D^{25}$ -214.22 (c, 1.01 in EtOH). $[\alpha]_D^{20}$ -222 (c, 0.1 in CHCl₃). Forms a hydrate Mp 125-135°. λ_{\max} 230 (sh) (log ϵ 4.15); 276 (sh) (log ϵ 3.79); 282 (log ϵ 3.91); 287 (log ϵ 3.93); 292 (log ϵ 3.89) (95% EtOH).

N-Oxide: (-)-Argemone N-oxide

[18841-61-7]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the above-ground parts of *Argemone gracilentia* (Papaveraceae). Glass + 1H₂O + 0.5CHCl₃. Mp 140-160° dec. (efferv.). $[\alpha]_D^{25}$ -185 (c, 2.81 in CHCl₃).

N-Me: (-)-Argemone N-methosalt

[55056-96-7]

[18826-69-2]

C₂₂H₂₈NO₄[±] 370.467

Quaternary alkaloid from above-ground parts of *Argemone gracilentia*, the roots and aerial parts of *Argemone platyceras* and the fruits of *Thalictrum revolutum* (Papaveraceae, Ranunculaceae). Needles (MeOH) (as iodide). Mp 272-274° (iodide). $[\alpha]_D^{25}$ -200 (c, 0.14 in MeOH). CAS no. refers to iodide.

O²-De-Me: 2-Hydroxy-3,8,9-trimethoxypavinane. Norargemone

[5876-16-4]

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Argemone brevicornuta*, *Argemone hispida*, *Argemone platyceras*, *Argemone munita* var. *rotundata*, *Eschscholtzia californica*, *Eschscholtzia glauca*, *Eschscholtzia douglasii*, *Berberis buxifolia*, *Thalictrum dasycarpum* and *Cryptocarya longifolia* (Papavera-

ceae, Berberidaceae, Ranunculaceae, Lauraceae). Cryst. (CHCl₃, CHCl₃/MeOH, C₆H₆/CHCl₃ or MeOH aq.). Mp 255-256° dec. (241-242°, 244-250°). $[\alpha]_D^{25}$ -154 (c, 2.79 in CHCl₃) (-147). λ_{\max} 230 (sh) (log ϵ 4.12); 278 (sh) (log ϵ 3.81); 283 (log ϵ 3.92); 287 (log ϵ 3.93); 293 (log ϵ 3.92) (95% EtOH).

O³-De-Me: 3-Hydroxy-2,8,9-trimethoxypavinane. Isonorargemone

[18826-67-0]

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Argemone munita* var. *argentea*, the above-ground parts of *Argemone gracilentia*, and the tops of *Thalictrum revolutum* (Papaveraceae, Ranunculaceae). Cryst. + 1H₂O (MeOH aq.). Mp 219-221°. $[\alpha]_D^{25}$ -202 (c, 3.31 in CHCl₃). λ_{\max} 287 (log ϵ 3.96) (MeOH).

O²,O⁹-Di-de-Me: 2,9-Dihydroxy-3,8-dimethoxypavinane. Bisorargemone.

Rotundine†

[6808-63-5]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Argemone hispida*, *Argemone munita* var. *rotundata*, *Thalictrum dasycarpum*, *Eschscholtzia californica*, *Eschscholtzia douglasii* and *Eschscholtzia glauca*, and *Cryptocarya longifolia* (Papaveraceae, Ranunculaceae, Lauraceae). Cryst. (MeOH or EtOH). Mp 254-255° (244-246°). $[\alpha]_D^{20}$ -244 (c, 0.19 in CHCl₃). $[\alpha]_D^{25}$ -265.8 (c, 0.158 in MeOH). λ_{\max} 230 (sh) (log ϵ 4.07); 279 (sh) (log ϵ 3.83); 285 (log ϵ 3.92); 288 (log ϵ 3.93); 294 (log ϵ 3.89) (95% EtOH).

(±)-form [5531-95-3]

Synthetic. Cryst. (Et₂O/CH₂Cl₂). Mp 138-140°.

O²-De-Me: [5876-37-9]

Synthetic. Needles (EtOH or EtOH aq.). Mp 222-223° (218-220°).

O³-De-Me: [5876-50-6]

Synthetic. Plates (heptane). Mp 175-177°.

O²,O⁹-Di-de-Me: [29944-24-9]

Synthetic. Cryst. (CHCl₃ or butyronitrile). Mp 233.5-235° (226-227°).

Soine, T.O. *et al.*, *J. Am. Pharm. Assoc.*, 1944, **33**, 185-188 (*isol*)

Battersby, A.R. *et al.*, *J.C.S.*, 1955, 2888-2896 (*synth*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 1728-1737 (*Norargemone, isol, uv, ir*)

Martell, M.J. *et al.*, *J.A.C.S.*, 1963, **85**, 1022-1023 (*pmr, struct*)

Stermitz, F.R. *et al.*, *J.O.C.*, 1966, **31**, 2925-2933 (*Argemone, Norargemone, Bisorargemone, isol, pmr, ms*)

Červinka, O. *et al.*, *Tet. Lett.*, 1966, 5375-5377 (*abs config*)

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 414-417; *CA*, **65**, 5500e (*ms*)

Martell, M. *et al.*, *J. Pharm. Sci.*, 1967, **56**, 973-976 (*resoln*)

Barker, A.C. *et al.*, *J.C.S. (C)*, 1967, 1317-1323 (*abs config, ord*)

Chan, R.P.K. *et al.*, *Tetrahedron*, 1967, **23**, 4209-4214 (*uv, ord, cd*)

Lee, K.-H. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 1922-1926 (*Norargemone, synth*)

Stermitz, F.R. *et al.*, *J.O.C.*, 1969, **34**, 555-559 (*isol, ir, pmr, ms, struct, synth*,

Argemone N-methosalt, N-oxide, Isonorargemone)

Kupchan, S.M. *et al.*, *J.O.C.*, 1969, **34**, 1062-1065 (*Norargemone, Bisorargemone*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1971, **36**, 2067-2069 (*Norargemone, Bisorargemone, isol*)

Chen, C.-H. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1634-1638 (*Bisorargemone, synth*)

Kaneda, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 1263-1266 (*cryst struct*)

Wenkert, E. *et al.*, *Top. Carbon-13 NMR Spectrosc.*, Vol. 2, (Levy, G.C., Ed.), Wiley, New York, 1976, 110 (*cmr*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 593-601 (*Isonorargemone*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1980, **43**, 270-277 (*isol, uv, cd, pmr, ms, synth, Argemone N-methosalt*)

Rice, K.C. *et al.*, *J.O.C.*, 1980, **45**, 601-607 (*Bisorargemone, synth, ms*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1981, **34**, 195-207 (*Norargemone, Bisorargemone*)

Nomoto, T. *et al.*, *Chem. Comm.*, 1982, 1113-1115 (*synth, pmr*)

Stadler, R. *et al.*, *Phytochemistry*, 1989, **28**, 1083-1086 (*Bisorargemone, pmr, cmr*)

Johnson, A.P. *et al.*, *J.C.S. Perkin I*, 1996, 907-913 (*Norargemone, synth, ir, pmr*)

Munchhof, M.J. *et al.*, *J.O.C.*, 1996, **61**, 4607-4610 (*synth, pmr, cmr*)

Necas, M. *et al.*, *Acta Cryst. C*, 2001, **57**, 66-71 (*cryst struct*)

Ruchirawat, S. *et al.*, *Tet. Lett.*, 2001, **42**, 1359-1361 (*synth*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr*)

Youte, J.-I. *et al.*, *J.O.C.*, 2004, **69**, 2737-2740 (*synth*)

Argentinamine

A-1411

C₁₅H₂₀N₂O₂ 260.335

Prob. Quinolizidine alkaloid. Struct. unknown. Alkaloid from *Ammodendron argenteum* (Fabaceae). Mp 201-202°. $[\alpha]_D$ -142.3 (EtOH). Cooccurs with Sparteine, S-385, Lupanine, L-302 and Cytisine, C-938.


Kushmuradov, Yu.K. *et al.*, *Nauch. Tr. Tashk. Gos. Univ. im. V.I. Lenina*, 1968, **341**, 95-98; *CA*, **73**, 32298b (*isol*)

Argentine

A-1412

3,3'-Carbonylbis[1,2,3,4,5,6-hexahydro-1,5-methano-2H-pyrido[1,2-a][1,5]diazocin-5-one], 9CI. 3,3'-Carbonylbiscytisine

[37551-61-4]



Absolute Configuration

C₂₃H₂₆N₄O₃ 406.483

Alkaloid from *Ammodendron longiracemosum*, *Sophora griffithii* and *Thermopsis alterniflora* (Fabaceae). Mp 258-259°. $[\alpha]_D$ -305 (c, 3.28 in EtOH).

Octahydro:

Cryst. (Me₂CO/petrol). Mp 208-209°. $[\alpha]_D$ -143 (c, 0.81 in EtOH).


Primukhamedov, I. *et al.*, *CA*, 1970, **72**, 79280j (*isol*)

Aslanov, Kh.A. *et al.*, *CA*, 1972, **77**, 140388x (*ir, pmr, ms, uv, struct*)

Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1972, 218-222; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 216-218 (*isol*)
 Karakozova, S.A. *et al.*, *Khim. Prir. Soedin.*, 1975, 664-665; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 703 (*isol*)
 Ajaz, M.S. *et al.*, *Thesis*, Uni. of Karachi, 1993, (*cmr*)

Arginine, INN, USAN A-1413

2-Amino-5-[(aminoiminomethyl)amino]pentanoic acid. 2-Amino-5-guanidinovaleic acid. Arg
 [7004-12-8]



C₆H₁₄N₄O₂ 174.202

Basic genetically coded amino acid. Essential amino acid for human development. Precursor of nitric oxide and is synth. by the body from 2,5-Diaminopentanoic acid. Diuretic agent. Growth hormone releaser. Immune stimulating agent and wound healing agent. Acidifying agent. Dietary supplement. Ammonia detoxicant (hepatic failure). Diagnostic aid for pituitary function. Log P -4.79 (*calc*).

(S)-form

L-form. *L*-Arginine. FEMA 3819 [74-79-3]

Found in seeds/shoots of pine, fir, acacia, arachis etc. and in many proteins. Reagent for the fluorimetric anal. of reducing sugars by hplc. Diagnostic aid (pituitary function determination); ammonia detoxicant; dietary supplement, nutrient. Plates or prisms + 2H₂O (H₂O), plates (EtOH). Sol. H₂O (15.0g/100g at 21°). Mp 207° Mp 244° dec. (anhyd. 105°). [α]_D²⁵ +21.8 (H₂O). [α]_D²⁵ +48.1 (c, 1 in 5M HCl). pK_{a1} 2.17; pK_{a2} 9.04; pK_{a3} 12.48 (guanido). Isoelectric point 10.76. Bitter taste. *N*-Protected derivs. useful in peptide synth. have been listed alphabetically elsewhere.

► CF1934200

N-Tetradecanoyl: **N-Tetradecanoylarginine**

C₂₀H₄₀N₄O₃ 384.561

Isol. from environmental DNA expressed in *Escherichia coli*.

N-Pentadecanoyl: **N-Pentadecanoylarginine**

C₂₁H₄₂N₄O₃ 398.588

Isol. from environmental DNA expressed in *Escherichia coli*.

N-Hexadecanoyl: **N-Hexadecanoylarginine**

C₂₂H₄₄N₄O₃ 412.615

Isol. from environmental DNA expressed in *Escherichia coli*.

N-(9*Z*-Hexadecenyl): **N-(9*Z*-Hexadecenyl)arginine**

C₂₂H₄₂N₄O₃ 410.599

Isol. from environmental DNA expressed in *Escherichia coli*.

*N*²-(4-Hydroxybenzoyl): **N²-(4-Hydroxybenzoyl)arginine**

[178495-39-1]

C₁₃H₁₈N₄O₄ 294.31

Isol. from the ascidian *Leptoclinides dubius*. Sol. MeOH, butanol. [α]_D²⁰ +19.5 (c, 2 in MeOH). λ_{max} 210 (log ε 3.46); 256 (log ε 3.49) (MeOH). λ_{max} 216 ; 294 (MeOH/NaOH) (Berdy).

*N*²-(2-Carboxyethyl): **N²-(2-Carboxyethyl)arginine**

[151636-36-1]

C₉H₁₈N₄O₄ 246.266

Prod. by a *Streptomyces clavuligerus* mutant. Biosynth. intermed. of Clavulanic acid. Related to Octopine.

*N*²-(3-Carboxypropanoyl): **N²-Succinylarginine**

[2478-02-6]

C₁₀H₁₈N₄O₅ 274.276

Constit. of the shoots of pear trees.

*N*²-Oxalyl: **N²-Oxalylarginine**

[90250-86-5]

C₈H₁₄N₄O₅ 246.222

Constit. of the shoots of apple and pear trees (*Malus* spp.).

*N*²-(3-Carboxy-2-hydroxypropanoyl):

N²-(2-Hydroxysuccinoyl)arginine. **N²-(3-Carboxy-2-hydroxy-1-oxopropyl)arginine**

[87605-92-3]

C₁₀H₁₈N₄O₆ 290.275

Constit. of the seeds of *Vicia faba*, the tubers of *Smilax china* and the shoots of apple and pear trees.

*N*²-(3-Carboxy-3-hydroxypropanoyl):

N²-(3-Hydroxysuccinoyl)arginine

[90250-85-4]

C₁₀H₁₈N₄O₆ 290.275

Constit. of the shoots of apple and pear trees.

*N*²-[3-Carboxy-2-(carboxymethyl)-2-hydroxypropanoyl]:

N²-(2-Carboxymethyl-2-hydroxysuccinoyl)arginine

[87605-91-2]

C₁₂H₂₀N₄O₈ 348.312

Constit. of *Lilium maximowiczii* and the shoots of apple and pear tree.

[14975-30-5 , 50912-92-0 , 32042-43-6 , 63238-98-2 , 78851-84-0]

Karrer, W. *et al.*, *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 2392 (*occur*)

Elson, S.W. *et al.*, *Chem. Comm.*, 1993, 1211; 1212 (*N*²-(2-Carboxylethyl)arginine)


Garcia, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 782-785 (4-hydroxybenzoyl)

Brady, S.F. *et al.*, *Org. Lett.*, 2005, **7**, 3613-3616 (*N*-acyl derivs)

Argiopin

Argiotoxin 636

[105029-41-2]



C₂₉H₅₂N₁₀O₆ 636.793

Isol. from venom of the spider *Argiope lobata*. Glutamate receptor antagonist.

Adams, M.E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1987, **148**, 678-683 (*isol, struct, activity*)

Shih, T.L. *et al.*, *Tet. Lett.*, 1987, **28**, 6015 (*struct, bibl*)

Jasys, V.J. *et al.*, *Tet. Lett.*, 1988, **29**, 6223 (*synth, cmr, pmr*)

Herlitz, S. *et al.*, *Neuron*, 1993, **10**, 1131-1140 (*activity*)


Scott, R.H. *et al.*, *Neuropharmacology*, 1998, **37**, 1563-1578 (*bibl, activity*)

Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)

Argiopin II

A-1415

[117233-42-8]



C₃₅H₆₀N₁₂O₆ 744.936

*N*¹ Numbered acc. to CAS. Isol. from the venom of the *Argiope lobatum* spider.

*N*¹-Me: **Argiopin I**

[117233-41-7]

C₃₆H₆₃N₁₂O₆[⊕] 759.971

Isol. from the venom of *Argiope lobatum*.

4-Deoxy, *N*¹-de-Me: **Pseudoargiopin II**

[117233-45-1]

C₃₄H₅₈N₁₂O₅ 714.91

Isol. from the venom of *Argiope lobatum*.

4-Deoxy, *N*¹-Me: **Pseudoargiopin I**

[117255-12-6]

C₃₆H₆₃N₁₂O₅[⊕] 743.971

Isol. from the venom of *Argiope lobatum*.


Grishin, E.V. *et al.*, *Bioorg. Khim.*, 1988, **14**, 883 (*isol*)

Grishin, E.V. *et al.*, *Toxicon*, 1989, **27**, 541 (*isol*)

Argiopin IV

A-1416

[117233-43-9]



C₃₁H₅₄N₁₀O₄ 630.832

Isol. from the venom of the spider *Argiope lobatum*.

4',9'-Di-Me: **Argiopin V**

[117233-44-0]


C₃₃H₅₈N₁₀O₄ 658.886

Isol. from the venom of the spider *Argiope lobatum*.

Grishin, E.V. *et al.*, *Toxicon*, 1989, **27**, 541 (*isol*)

Argiotoxin 659Argiopinin III
[111944-83-3]

A-1417

 $C_{31}H_{53}N_{11}O_5$ 659.83


Argiopinin III had unspecified stereochem. and the two substances may not therefore be identical. Isol. from the venom of *Argiope aurantia* spider and (as Argiopinin III) from *Argiope lobatum* venom.

[117306-99-7]

Adams, M.E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1987, **148**, 678 (isol)Jasys, V.J. *et al.*, *Tet. Lett.*, 1988, **29**, 6223 (synth, cmr, pmr)Grishin, E.V. *et al.*, *Toxicol.*, 1989, **27**, 541 (Argiopinin III)**Argiotoxin 673**

[111924-44-8]


A-1418

 $C_{32}H_{55}N_{11}O_5$ 673.857

Isol. from the venom of *Argiope aurantia* spider.

Adams, M.E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1987, **148**, 678 (isol)Jasys, V.J. *et al.*, *Tet. Lett.*, 1988, **29**, 6223 (synth, cmr, pmr)**Arglecin**[3-[1,6-Dihydro-5-(2-methylpropyl)-6-oxopyrazinyl]propyl]guanidine, 9CI
[34098-41-4]

A-1419


 $C_{12}H_{21}N_5O$ 251.331

Diketopiperazine derived antibiotic. Produced by *Streptomyces toxytricini* and other *Streptomyces* sp. Enzyme inhibitor. Antiarrhythmic agent. Needles (EtOH/EtOAc)(as dihydrochloride). Sol. H_2O , DMSO, Py, MeOH; poorly sol. butanol, hexane. Mp 176-177° (hydrochloride). pK_{a1} 1.4; pK_{a2} 8.9. Log P -0.5 (calc). λ_{max} 226 (ϵ 8700); 338 (ϵ 9400) (0.1N HCl) (Derep). λ_{max} 232 (ϵ 8200); 321 (ϵ 9100) (0.1N NaOH) (Derep). λ_{max} 226 (ϵ 8000); 322 (ϵ 9600) (H_2O) (Derep). λ_{max} 258 (ϵ 8200); 321 (ϵ 9100) (NH_4OH) (Berdy).

Umezawa, S. *et al.*, *Tet. Lett.*, 1971, 259 (ir, uv, pmr, cmr, isol)Tatsuta, K. *et al.*, *J. Antibiot.*, 1972, **25**, 674 (ir, ms, nmr, synth, struct)MacDonald, J.C. *et al.*, *Tetrahedron*, 1976, **32**, 655 (pmr, cmr)MacDonald, J.C. *et al.*, *Can. J. Biochem.*, 1977, **55**, 165 (biosynth)Ohta, A. *et al.*, *Heterocycles*, 1987, **26**, 3181 (synth)Turck, A. *et al.*, *J. Het. Chem.*, 1994, **31**, 1449 (synth)**Argutane A**

[1003567-49-4]

A-1420



Relative Configuration

 $C_{20}H_{22}N_2O_2$ 322.406

Alkaloid from the roots of *Incarvillea arguta*. Yellow oil. $[\alpha]_D^{25} +25.4$ (c, 0.05 in MeOH).

10'-Epimer: **Argutane B**


[1003567-50-7]

 $C_{20}H_{22}N_2O_2$ 322.406

Alkaloid from the roots of *Incarvillea arguta*. Yellow oil. $[\alpha]_D^{25} +21$ (c, 0.02 in MeOH).

Fu, J.-J. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2151-2155 (isol, pmr, cmr, ms)**Argvalin**[3-[1,6-Dihydro-5-(1-methylethyl)-6-oxopyrazinyl]propyl]guanidine, 9CI. 6-(3-Guanidinypropyl)-3-isopropyl-2(1H)-pyrazinone
[52159-72-5]

A-1421


 $C_{11}H_{19}N_5O$ 237.304

Microbial metab. of *Streptomyces* spp. Shows antibiotic props. Needles (EtOAc/2-propanol)(as hydrochloride). Sol. H_2O , MeOH. Mp 175-176° (hydrochloride). λ_{max} 226 (ϵ 8300); 321 (ϵ 9600) (H_2O) (Berdy). λ_{max} 226 (ϵ 9100); 333 (ϵ 9600) (HCl) (Berdy).

Tatsuta, K. *et al.*, *J. Antibiot.*, 1973, **26**, 606 (isol, struct, nmr)Ohta, A. *et al.*, *Heterocycles*, 1988, **27**, 437 (synth)**Ariakemicin A**

[1032399-11-3]

A-1422

 $C_{32}H_{38}N_4O_7$ 590.675

Alkaloid from the marine bacterium *Rapidithrix* sp. strain HC35. Reddish-brown amorph. solid. Isol. as a mixt. with Ariakemicin B to which data refers. λ_{max} 228 (ϵ 35000); 278 (ϵ 44000); 302 (ϵ 43000) (MeOH).

 Δ^{11} -Isomer: **Ariakemicin B**


[1032399-13-5]

 $C_{32}H_{38}N_4O_7$ 590.675

Alkaloid from *Rapidithrix* sp. strain HC35.

Oku, N. *et al.*, *Org. Lett.*, 2008, **10**, 2481-2484 (isol, uv, ir, pmr, cmr)**Aricine***Heterophylline*†. *Cinchovatine*
[482-91-7]

A-1423

 $C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from several *Cabucala*, *Ochrosia*, *Rauwolfia* and *Aspidosperma* spp., also from *Cinchona ledgeriana* and *Cinchona pelteteriana* (Apocynaceae, Rubiaceae). Cryst. (EtOH aq.). Mp 190° dec. $[\alpha]_D^{20} -91$ (c, 1.4 in $CHCl_3$). $[\alpha]_D^{20} -63$ (c, 1.5 in Py). $[\alpha]_D^{20} -57$ (c, 1.0 in EtOH). λ_{max} 227 (log ϵ 4.54); 250 (log ϵ 4.05); 279 (log ϵ 3.98); 298 (sh) (log ϵ 3.88) (EtOH or MeOH).

Hydrochloride:

Plates (MeOH/Me₂CO). Mp 254°. $[\alpha]_D^{20} -5$ (c, 0.9 in 50% EtOH aq.).

Hydrobromide:

Needles (MeOH). Mp 262-263° dec.

20-Epimer: **Cabucine**. 10-Methoxyajmalicine

[16812-97-8]

 $C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from *Cabucala madagascariensis*, *Cabucala erythrocarpa*, *Cabucala fasciculata*, *Cabucala striolata*, *Cabucala torulosa* and *Cabucala cryptophlebia* (Apocynaceae). Mp 125°. $[\alpha]_D -60$ ($CHCl_3$).

19,20-Diepimer: **Raumitorine**

[549-74-6]


 $C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae). Needles (MeOH). Mp 138°. $[\alpha]_D^{20} +60$ (c, 0.54 in $CHCl_3$).

Goutarel, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1954, 1481-1482 (*Raumitorine*)Stoll, A. *et al.*, *Helv. Chim. Acta*, 1955, **38**, 270-283 (isol, uv, ir, struct)Shamma, M. *et al.*, *J.A.C.S.*, 1963, **85**, 2507-2512 (stereochem)Finch, N. *et al.*, *Tetrahedron*, 1966, **22**, 1327-1337 (uv, ord, stereochem)Douzoua, L.L. *et al.*, *Ann. Pharm. Fr.*, 1972, **30**, 199-204; *CA*, **77**, 72560v (*Cabucine*)Sakai, S. *et al.*, *Yakugaku Zasshi*, 1978, **98**, 850-862; *CA*, **89**, 197771u (synth)

Aristo red

[27323-39-3]

C₁₉H₁₅NO₆ 353.331

Alkaloid from *Aristolochia reticulata*, *Aristolochia serpentaria*, and from the roots of *Aristolochia bracteata* (Aristolochiaceae). Red needles (EtOH). Mp 286.5-287.5°.


Coutts, R.T. *et al.*, *J.C.S.*, 1957, 4120-4124 (*isol, uv*)

Jagannadha Rao, K.V. *et al.*, *Curr. Sci.*, 1958, 27, 168; *J. Sci. Ind. Res., Sect. B*, 1959, 18, 245; *CA*, 52, 20421f; 54, 14373a (*isol*)

Coutts, R.T. *et al.*, *J. Pharm. Pharmacol.*, 1959, 11, 607-617 (*isol*)

Aristoaristone

[155210-53-0]



Relative Configuration


C₄₀H₄₆N₄O₂ 614.829

Alkaloid from *Aristolochia australisica* (Elaeocarpaceae).

Quirion, J.C. *et al.*, *Nat. Prod. Lett.*, 1993, 2, 41-48

Aristofrucosine

[116965-59-4]



Absolute Configuration

C₂₀H₂₄N₂ 292.423

Unique 3-azatricyclo[3.3.1.0^{3,7}]nonane nucleus. Alkaloid from *Aristolochia fruticosa* (Elaeocarpaceae). Amorph. Mp 146° (synthetic). [α]_D²⁵ +50.5 (c, 0.53 in CHCl₃). [α]_D²⁵ +15.5 (c, 0.8 in CHCl₃) (synthetic).

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1988, 29, 3355-3356 (*isol, pmr, ms, struct*)

Beerli, R. *et al.*, *Helv. Chim. Acta*, 1991, 74, 110-116 (*synth, ir, pmr, cmr, ms, abs config*)


A-1424

Aristolactam I

A-1427

8-Methoxybenzo[*f*]-1,3-benzodioxolo[6,5,4-*cd*]indol-5(6H)-one, 9CI. 10-Amino-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid lactam.

Aristolactam. Aristolactam
[13395-02-3]

C₁₇H₁₁NO₄ 293.278

Alkaloid from the roots of *Aristolochia argentina* and *Aristolochia indica*, and from the tubers of *Aristolochia rotunda*, also *isol.* from *Aristolochia debilis* and the Chinese drug Fang-chi (Aristolochiaceae). Cytotoxic. Yellow cryst. (by *subl.*). Mp 315-317°.

▶DE5375000

N-β-D-Glucopyranosyl: **Aristolactam I N-glucoside**

[17413-41-1]

C₂₃H₂₁NO₉ 455.42

Alkaloid from *Aristolochia indica*, *Aristolochia longa*, *Aristolochia contorta*, *Aristolochia kankunensis* and *Aristolochia tuberosa* (Aristolochiaceae). Pale yellow needles (MeOH aq.). Mp 331-333°. [α]_D²⁵ -14 (c, 0.02 in H₂O) (c, 0.11 in DMF). λ_{max} 238 (ε 30900); 243 (ε 33200); 249 (ε 31400); 258 (ε 36500); 290 (ε 15300); 298 (ε 15200); 328 (ε 9521) (EtOH) (Derep). λ_{max} 214; 242; 260; 290; 330; 396 (MeOH) (Berdy).

N-β-D-Glucopyranosyl, tetra-O-Ac:

Pale yellow cryst. (MeOH/CHCl₃). Mp 287-289°. [α]_D²⁵ -97 (c, 0.12 in CHCl₃).

N-[4-Hydroxy-E-cinnamoyl-(→6)-β-D-glucopyranosyl]: **Aristolactam I N-(6'-trans-p-coumaroyl)-β-D-glucopyranoside**

C₃₂H₂₇NO₁₁ 601.565

Alkaloid from roots of *Aristolochia contorta* (Aristolochiaceae). Exhibits significant antibacterial activity against gram-positive bacteria. Yellow needles. Mp 260-264°. [α]_D²⁰ -20.2 (c, 0.043 in MeOH). λ_{max} 208; 235; 295; 390 (MeOH) (Berdy).

O-De-Me: **Aristolactam Ia**

[85814-30-8]

C₁₆H₉NO₄ 279.251

Alkaloid from the rhizomes of *Aristolochia argentina* and from *Aristolochia longa*, *Aristolochia indica*, *Aristolochia kankunensis* and *Stephania cepharantha*. Cryst. (AcOH). Mp 350°. λ_{max} 210; 238; 243; 250; 260; 292; 298 (MeOH) (Berdy).

O-De-Me, N-β-D-glucopyranosyl: **Aristolactam Ia N-β-D-glucoside**

[213530-43-9]

C₂₂H₁₉NO₉ 441.393

Alkaloid from *Aristolochia foveolata*. Yellow powder (MeOH). Mp >280°.

O-De-Me, O-β-D-glucopyranoside: **Aristolactam Ia O-glucoside. Manshurienine B**

C₂₂H₁₉NO₉ 441.393

Alkaloid from the stems of *Aristolochia manshuriensis*. Yellow powder. Mp 292-293°. [α]_D²⁵ -10.5 (c, 0.2 in MeOH). λ_{max} 246; 264; 275; 318 (MeOH).

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1959, 79, 973; 1470 (*isol, uv, ir*)

Carboni, S. *et al.*, *Gazz. Chim. Ital.*, 1966, 96, 641 (*isol*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1968, 33, 3735 (*isol, uv, ir, pmr, struct, glucoside*)

Priestap, H.A. *et al.*, *An. Asoc. Quim. Argent.*, 1971, 59, 245; *CA*, 76, 43943n (*isol*)

Zhu, D. *et al.*, *Zhongcaoyao*, 1981, 12, 529; *CA*, 97, 39261p (*isol, glucoside*)

Priestap, H.A. *et al.*, *Phytochemistry*, 1985, 24, 849-852 (*isol, uv, ir, pmr, ms, struct*)

Lee, H.S. *et al.*, *J. Nat. Prod.*, 1992, 55, 1165 (*glucosides*)

Leu, Y.-L. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1998, 45, 539-541 (*Aristolactam Ia N-glucoside*)


Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, 48, 1006-1009 (*activity*)

Zhang, Y.-T. *et al.*, *Helv. Chim. Acta*, 2006, 89, 2665-2670 (*Manshurienine B*)

Aristolactam C II

A-1428

[106283-30-1]

C₁₈H₁₅NO₄ 309.321

Struct. revised in 1988. CAS numbering shown. Alternative (phenanthrene) numbering frequently used, in which 1,2,9-become 4,3,6- respectively. Alkaloid from the rhizomes of *Aristolochia argentina* (Aristolochiaceae). Pale yellow needles (C₆H₆). Mp 190°.

N,O-Di-Ac: Mp 207-208°.

I'-Carboxylic acid, O²-de-Me: **Aristolactam D II**

[106283-33-4]

C₁₇H₁₁NO₅ 309.278

Alkaloid from the rhizomes of *Aristolochia argentina*. Yellow needles (DMSO/propanol). Mp 278-280°. Struct. revised in 1988.

9-Hydroxy, O²-de-Me: **Aristolukine A**

C₁₇H₁₃NO₅ 311.293

Alkaloid from *Aristolochia kaempferi*. Yellowish needles (MeOH). Mp 274-276°. λ_{max} 239; 256; 280 (sh); 345 (sh); 404 (MeOH).

9-Methoxy: **Aristolactam C III**

[106283-32-3]

C₁₉H₁₇NO₅ 339.347

Alkaloid from *Aristolochia argentina* rhizomes (Aristolochiaceae). Yellow needles (C₆H₆). Mp 230-235°.

9-Methoxy, O²-de-Me: **Aristolactam C IV**
C₁₈H₁₅NO₅ 325.32
Alkaloid from *Aristolochia heterophylla*. Yellowish needles. Mp 206-207°. λ_{max} 207 (log ε 4.54); 239 (log ε 4.56); 253 (log ε 4.47); 281 (log ε 4.33); 293 (log ε 4.34); 316 (sh) (log ε 4.09); 399 (log ε 3.63) (MeOH).

9-Methoxy, 1'-carboxylic acid, O²-de-Me: **Aristolactam D III**
[106283-35-6]
C₁₈H₁₃NO₆ 339.304
Trace alkaloid from rhizomes of *Aristolochia argentina*.

Priestap, H.A. *et al.*, *Phytochemistry*, 1985, **24**, 3035 (isol, uv, ir, pmr, cmr, ms)


Talapatra, S.K. *et al.*, *Phytochemistry*, 1988, **27**, 903 (struct)

Wu, T.S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1624-1626 (Aristolokine A)

Wu, T.S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 357-361 (Aristolactam C IV)

Aristolactam IIIa A-1429

10-Hydroxybenzo[f]-1,3-benzodioxolo[6,5,4-cd]indol-5(6H)-one, 9CI [97399-89-8]



C₁₆H₉NO₄ 279.251

Alkaloid from *Aristolochia tuberosa* and from the rhizomes of *Aristolochia argentina* (Aristolochiaceae). Cytotoxic. Cryst. (AcOH). Mp 350°.

O-β-D-Glucopyranoside: **Aristolactam IIIa O-glucoside. Manshurienine A**
C₂₂H₁₉NO₉ 441.393

Alkaloid from the stems of *Aristolochia manshuriensis*. Yellow powder. Mp 236-237°. [α]_D²⁵ -8.5 (c, 0.2 in MeOH). λ_{max} 238 ; 253 ; 280 ; 324 (MeOH).

N-β-D-Glucopyranosyl: **Aristolactam IIIa N-glucoside**
[80311-26-8]

C₂₂H₁₉NO₉ 441.393

Alkaloid from the roots of *Aristolochia indica* and from *Aristolochia tuberosa* (Aristolochiaceae). Deep brown amorph. powder. Mp 320°.

Me ether: **Aristolactam III**

[81451-90-3]

C₁₇H₁₁NO₄ 293.278

Alkaloid from the rhizomes of *Aristolochia argentina* (Aristolochiaceae). Cryst. (AcOH). Mp 297-304°.

Achari, B. *et al.*, *Phytochemistry*, 1981, **20**, 1444 (isol, ir, pmr, ms, struct, glucoside)

Zhu, D. *et al.*, *Zhongcaoyao*, 1981, **12**, 529; *CA*, **97**, 39261p (isol)

Priestap, H.A. *et al.*, *Phytochemistry*, 1985, **24**, 849 (isol, uv, ir, pmr, ms, struct)


Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **148**, 1006-1009 (activity)

Nascimento, I.R. *et al.*, *Phytochemistry*, 2003, **63**, 953-957 (glucoside)

Zhang, Y.-T. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 2665-2670 (Manshurienine A)

Aristolarine A-1430

[90930-69-1]



Absolute Configuration

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Aristolotelia peduncularis* (Elaeocarpaceae). Yellow cryst.

Mp 181-185.5°. [α]_D +264 (c, 1.12 in CHCl₃) (synthetic). λ_{max} 233 (sh) (log ε 4.24); 236 (log ε 4.24); 256 (sh) (log ε 3.8); 266 (sh) (log ε 3.63); 290 (sh) (log ε 3.07); 325 (sh) (log ε 2.89) (EtOH).

Deoxy: **Aristoloteline**

[59863-01-3]

C₂₀H₂₆N₂O 310.438

Minor alkaloid from the leaves and stems of *Aristolotelia chilensis* (Elaeocarpaceae). Also obt. from Serratoline, S-259 by boiling with dil. acid. Cryst. (MeOH). Mp 218-222°. [α]_D +41 (CHCl₃).

2-Epimer, deoxy: **2-Epiaristoloteline**

[144789-06-0]

C₂₀H₂₆N₂O 310.438

Alkaloid from *Aristolotelia chilensis*.

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1976, **15**, 574-575 (Aristoloteline)

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 804-814 (Aristolarine)


Gueller, R. *et al.*, *Helv. Chim. Acta*, 1993, **76**, 1847-1862 (Aristoloteline, synth)

Stoermer, D. *et al.*, *J.O.C.*, 1993, **58**, 564-568 (Aristoloteline, synth)

He, K. *et al.*, *Int. J. Pharmacogn.*, 1997, **35**, 215-217 (2-Epiaristoloteline)

Aristolasene A-1431

[119459-70-0]



Relative Configuration


C₂₀H₂₂N₂ 290.407

Minor alkaloid from the aerial parts of *Aristolotelia australasica* (Elaeocarpaceae). Yellow oil. [α]_D²⁰ +493 (c, 0.9 in CHCl₃). λ_{max} 215 (log ε 4.13); 262 (log ε 3.9); 330 (log ε 3.89) (EtOH).

Quirion, J.C. *et al.*, *Phytochemistry*, 1988, **27**, 3337-3339 (isol, uv, ir, pmr, ms, struct)

Aristolasicone A-1432

[117611-47-9]



Absolute Configuration

C₂₀H₂₄N₂O 308.422

New inverted indole subgroup of Aristotelia alkaloids. Revised struct. (1992). Formerly assigned as 15-Oxoaristoloteline. Alkaloid from *Aristolotelia australasica* (Elaeocarpaceae). Mp 205-210°. [α]_D -161.

15R-Alcohol: **Aristolasicol**

[117611-48-0]

C₂₀H₂₆N₂O 310.438

Alkaloid from aerial parts of *Aristolotelia australasica* (Elaeocarpaceae). Amorph. [α]_D²⁰ +20.3 (c, 0.77 in CHCl₃). Revised struct. Prev. descr. as 15-hydroxyaristoloteline. λ_{max} 226 (log ε 4.41); 282 (log ε 3.84); 289 (log ε 3.78) (EtOH).

Deoxy: **Alloaristoloteline. Deoxoaristolasicone**

[117678-03-2]

C₂₀H₂₆N₂ 294.439

Alkaloid from *Aristolotelia australasica* (Elaeocarpaceae). Cryst. (THF/cyclohexane/Et₃N). Mp 219.5° (211.5-214°). [α]_D²⁰ -32 (c, 1.4 in CHCl₃). Struct. revised in 1991. Previously descr. as 11-Epiaristoloteline. λ_{max} 226 (log ε 4.32); 282 (log ε 3.85); 290 (log ε 3.8) (EtOH).

8-Hydroxy, 15R-alcohol: **Aristolasicolone**

[117611-50-4]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from aerial parts of *Aristolotelia australasica* (Elaeocarpaceae). Amorph. [α]_D²⁰ -150 (c, 0.45 in CHCl₃). Revised struct. Previously descr. as 15-hydroxyaristolotelinone. λ_{max} 213 (log ε 4.26); 244 (log ε 4.13); 266 (log ε 2.99); 301 (log ε 3.94) (EtOH).

[134779-80-9]

Kan-Fan, C. *et al.*, *Tetrahedron*, 1988, **44**, 1651-1660 (Aristolasicol, Aristolasicolone)

Burkard, S. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 275-289; 1636-1642 (synth, struct, bibl)


Quirion, J.C. *et al.*, *J.O.C.*, 1992, **57**, 5848-5851 (Alloaristoloteline, pmr, cmr, cryst struct)

Gueller, R. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 1197-1204 (Alloaristoloteline, synth, abs config)

Stoermer, D. *et al.*, *J.O.C.*, 1993, **58**, 564-568 (synth, Alloaristoloteline)

Aristolasol A-1433

[119483-74-8]



Relative Configuration

C₂₀H₂₆N₂O₂ 326.438


Minor alkaloid from the aerial parts of *Aristolotelia australasica* (Elaeocarpaceae). Oil. $[\alpha]_D^{20} +36$ (c, 0.6 in CHCl₃). λ_{\max} 222 (log ϵ 4.33); 281 (log ϵ 3.85); 289 (log ϵ 3.83) (EtOH).

Quirion, J.C. *et al.*, *Phytochemistry*, 1988, **27**, 3337-3339 (*isol, uv, ir, pmr, cmr, ms, struct*)

Aristolin

[1398-05-6]

A-1434

C₃₇H₄₃NO₈ 629.749

Ester of 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262 with 16,17-Kauranediol. Constit. of *Aristolochia elegans* and *Aristolochia pubescens*. Yellow cryst. (CHCl₃). Mp 119-120°. $[\alpha]_D^{25} -63.9$ (c, 0.04 in CHCl₃). λ_{\max} 226 (log ϵ 4.19); 254 (log ϵ 4.14); 321 (log ϵ 3.93); 390 (log ϵ 3.8) (no solvent reported).


Wu, T.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1522-1525 (*isol, pmr, cmr*)

Nascimento, I.R. *et al.*, *Phytochemistry*, 2003, **63**, 953-957 (*isol, cmr*)

Aristoloin I

[611211-57-5]

A-1435

C₃₇H₄₃NO₈ 629.749

Ester of 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262 with 16,17-Kauranediol. Alkaloid from the tubercula of *Aristolochia pubescens*. Amorph. yellow solid. $[\alpha]_D^{25} -42.1$ (c, 0.2 in CHCl₃). λ_{\max} 223 (log ϵ 4.42); 256 (log ϵ 4.24); 289 (log ϵ 4.06); 316 (log ϵ 3.86); 389 (log ϵ 3.68) (MeOH).

Demethoxy: Aristoloin II

[611211-59-7]

C₃₆H₄₁NO₇ 599.722


Alkaloid from the tubercula of *Aristolochia pubescens*. Amorph. yellow solid. $[\alpha]_D^{25} -53.4$ (c, 0.16 in CHCl₃). λ_{\max} 219 (log ϵ 4.28); 250 (log ϵ 4.19); 314 (log ϵ 3.8); 387 (log ϵ 3.5) (MeOH).

Nascimento, I.R. *et al.*, *Phytochemistry*, 2003, **63**, 953-957 (*isol, pmr, cmr*)

Aristoloterpenate I

[184955-22-4]

A-1436

C₃₂H₃₁NO₈ 557.599

Constit. of *Aristolochia mollissima* and *Aristolochia heterophylla*. Yellow needles (Me₂CO). Mp 247-249°. $[\alpha]_D -39.5$ (c, 0.02 in CHCl₃). Ester of 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262. Abs. config. revised in 1999. λ_{\max} 226 (log ϵ 4.5); 239 (log ϵ 4.4); 250 (log ϵ 4.36); 268 (log ϵ 4.21); 287 (log ϵ 3.98); 322 (log ϵ 3.99); 393 (log ϵ 3.76) (MeOH). λ_{\max} 226; 239; 250; 268; 287; 322; 393 (MeOH) (Berdy).

2'E-Isomer: Aristoloterpenate III

[222412-32-0]

C₃₂H₃₁NO₈ 557.599

Constit. of *Aristolochia heterophylla*. Yellow needles (Me₂CO). Mp 245-247°. $[\alpha]_D -86$ (c, 0.038 in CHCl₃). λ_{\max} 225 (log ϵ 4.31); 251 (log ϵ 4.31); 267 (log ϵ 4.18); 284 (log ϵ 3.98); 321 (log ϵ 3.96); 391 (log ϵ 3.74) (MeOH).

8-Demethoxy: Aristoloterpenate II

[222403-62-5]

C₃₁H₂₉NO₇ 527.573

Constit. of *Aristolochia heterophylla*. Yellow needles (Me₂CO). Mp 241-243°. $[\alpha]_D -35.4$ (c, 0.016 in CHCl₃). Ester of 3,4-Methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-447. λ_{\max} 226 (log ϵ 4.57); 241 (log ϵ 4.46); 250 (log ϵ 4.41); 267 (log ϵ 4.27); 283 (log ϵ 4.08); 321 (log ϵ 4.05); 392 (log ϵ 3.82) (MeOH).

8-Demethoxy, 2'E-isomer: Aristoloterpenate IV

[222403-63-6]

C₃₁H₂₉NO₇ 527.573


Constit. of *Aristolochia heterophylla*. Yellow needles (Me₂CO). Mp 234-236°. $[\alpha]_D -88.3$ (c, 0.011 in CHCl₃). λ_{\max} 210 (log ϵ 4.6); 219 (log ϵ 4.62); 243 (log ϵ 4.57); 251 (log ϵ 4.6); 266 (log ϵ 4.44); 298 (log ϵ 4.17); 375 (log ϵ 3.74) (MeOH). λ_{\max} 210; 219; 243; 251; 266; 298; 375 (MeOH) (Berdy).

Wu, T.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 415-418 (*isol, uv, ir, pmr, cmr, ms*)

Aristomakine

[80787-63-9]

A-1437



Probable Absolute Configuration

C₂₀H₂₆N₂ 294.439

Not apparently related biogenetically to the *Aristolotelia* group of alkaloids. Abs. config. based on biogenetic considerations. Alkaloid from *Aristolotelia serrata* (Elaeocarpaceae). Noncryst. $[\alpha]_D^{22} -79.1$ (c, 1.5 in CHCl₃).

N-Deisopropyl: AristomakinineC₁₇H₂₀N₂ 252.358

Alkaloid from *Aristolotelia serrata* (Elaeocarpaceae). Amorph. $[\alpha]_D^{19} -72$ (CHCl₃).

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1981, **22**, 3275-3276 (*ms, pmr, cmr, struct*)


Bick, I.R.C. *et al.*, *Alkaloids (Academic Press)*, 1985, **24**, 113-151 (*rev, Aristomakinine*)

Burkard, S. *et al.*, *Helv. Chim. Acta*, 1990, **73**, 298-302 (*Aristomakine, Aristomakinine, synth*)

Aristone

[66833-17-8]

A-1438



Relative Configuration

C₂₀H₂₄N₂O 308.422

Trace alkaloid from *Aristolotelia chilensis*, also *isol.* from aerial parts of *Aristolotelia australasica* (Elaeocarpaceae). Mp 240-242°. Mp 268-270°. $[\alpha]_D^{20} -130$ (c, 0.4 in CHCl₃). λ_{\max} 213 (log ϵ 3.85); 239 (log ϵ 2.32); 293 (log ϵ 0.8) (EtOH).

Bittner, M. *et al.*, *Chem. Comm.*, 1978, 79-80 (*uv, struct*)


Zabel, V. *et al.*, *J.C.S. Perkin 1*, 1980, 2842-2844 (*cryst struct*)

Quirion, J.-C. *et al.*, *J.O.C.*, 1987, **52**, 4527-4530 (*isol, uv, ir, pmr, cmr, ms*)

Aristophyllide A

[221148-60-3]

A-1439

C₃₂H₃₁NO₈ 557.599

Alkaloid from *Aristolochia heterophylla*. Yellow needles. Mp 226-228°. $[\alpha]_D +120$ (c, 0.02 in CHCl₃). λ_{\max} 224 (log ϵ 4.34); 251 (log ϵ 4.16); 266 (log ϵ 4.07); 283 (log ϵ 3.86); 320 (log ϵ 3.82); 390 (log ϵ 3.59) (MeOH).

5',12'-Diepimer: Aristophyllide B

[221148-61-4]

C₃₂H₃₁NO₈ 557.599

Constit. of *Aristolochia heterophylla*. Yellow needles. Mp 229-231°. $[\alpha]_D -79$

(c, 0.02 in CHCl₃). λ_{\max} 224 (log ϵ 4.55); 249 (log ϵ 4.4); 268 (log ϵ 4.28); 283 (log ϵ 4.04); 318 (log ϵ 4.04); 390 (log ϵ 3.79) (MeOH).

Demethoxy: Aristophyllide C

[221148-62-5]

C₃₁H₂₉NO₇ 527.573

Alkaloid from *Aristolochia heterophylla*. Yellow needles. Mp 217-219°. [α]_D +113 (c, 0.06 in CHCl₃). λ_{\max} 218 (log ϵ 4.33); 243 (log ϵ 4.32); 252 (log ϵ 4.37); 266 (log ϵ 4.2); 298 (log ϵ 3.98); 377 (log ϵ 3.48) (MeOH).

5',12'-Diepimer, demethoxy: Aristophyllide D

[221148-63-6]

C₃₁H₂₉NO₇ 527.573


Alkaloid from *Aristolochia heterophylla*. Yellow needles. Mp 220-222°. [α]_D -100 (c, 0.01 in CHCl₃). λ_{\max} 219 (log ϵ 4.6); 242 (log ϵ 4.55); 252 (log ϵ 4.61); 266 (log ϵ 4.47); 297 (log ϵ 4.19); 355 (log ϵ 3.76); 373 (log ϵ 3.73) (MeOH).

Wu, T.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 348-351 (*isol, uv, ir, pmr, cmr, ms*)

Aristoquinoline C

A-1440

[710319-87-2]

C₂₆H₂₇NO₇ 465.502

Alkaloid from *Aristolochia elegans*. Yellowish syrup. [α]_D²⁵ -52 (c, 0.01 in MeOH). λ_{\max} 248 (log ϵ 4.24); 274 (log ϵ 3.85) (MeOH).

2''-Me ether: Aristoquinoline B

[710319-86-1]

C₂₇H₂₉NO₇ 479.529

Alkaloid from *Aristolochia elegans*. Yellowish syrup. [α]_D²⁵ -77.1 (c, 0.016 in MeOH). λ_{\max} 235 (log ϵ 4.2); 280 (log ϵ 3.75) (MeOH).

N-Epimer, 2''-Me ether: Aristoquinoline A

[710319-85-0]

C₂₇H₂₉NO₇ 479.529


Alkaloid from *Aristolochia elegans*. Yellowish syrup. [α]_D²⁵ -31.1 (c, 0.03 in MeOH). λ_{\max} 234 (log ϵ 4.26); 284 (log ϵ 3.75) (MeOH).

Shi, L.-S. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 439-446 (*isol, pmr, cmr, ms*)

Aristoserratenine

A-1441

[95727-41-6]



Absolute Configuration

C₂₀H₂₆N₂ 294.439

Alkaloid from *Aristolochia serrata* (Elaeocarpaceae). Amorph. [α]_D¹⁹ +58 (CHCl₃). λ_{\max} 226 (log ϵ 4.38); 259 (log ϵ 3.57) (MeOH).

3-Epimer: 3-Epiaristoserratenine

[117678-05-4]

C₂₀H₂₆N₂ 294.439


Alkaloid from aerial parts of *Aristolochia australasica* (Elaeocarpaceae). Amorph. [α]_D +36 (c, 0.8 in CHCl₃). λ_{\max} 226 (log ϵ 4.35); 258 (log ϵ 3.58) (EtOH).

Hai, M.A. *et al.*, *Tetrahedron*, 1984, **40**, 4359-4361 (*Aristoserratenine, isol, uv, ir, pmr, cmr, struct*)

Kan-Fan, C. *et al.*, *Tetrahedron*, 1988, **44**, 1651-1660 (*3-Epiaristoserratenine*)

Aristoteline

A-1442

C₂₀H₂₆N₂ 294.439**(+)-form [57103-59-0]**

Major alkaloid from *Aristolochia serrata*, *Aristolochia peduncularis* and from the leaves and stems of *Aristolochia chilensis* (Elaeocarpaceae). Shows hypotensive props., reduced pulse rate. Cryst. (MeOH). Mp 164°. [α]_D²⁰ +16 (MeOH). [α]_D²⁵ +23 (c, 1.84 in CHCl₃). λ_{\max} 228 (log ϵ 4.52); 283 (log ϵ 3.88); 289 (sh) (log ϵ 3.81) (EtOH).

9,10-Didehydro: 9,10-Dehydroaristoteline

[117611-51-5]

C₂₀H₂₄N₂ 292.423

Alkaloid from aerial parts of *Aristolochia australasica* (Elaeocarpaceae). Amorph. [α]_D²⁰ +272 (c, 0.8 in CHCl₃). λ_{\max} 223 (log ϵ 4.82); 275 (log ϵ 4.2); 281 (log ϵ 4.2); 290 (log ϵ 4.14) (EtOH).

17-Hydroxy: Aristocarbinoil. 17-Hydroxyaristoteline

[117611-49-1]

C₂₀H₂₆N₂O 310.438

Alkaloid from aerial parts of *Aristolochia australasica* (Elaeocarpaceae). Amorph. [α]_D²⁰ +4.3 (c, 0.7 in CHCl₃). Struct. needs reinvestigating after struct. revision of Aristolasicol. λ_{\max} 226 (log ϵ 4.3); 280 (log ϵ 3.79) (EtOH).

8-Oxo: Aristotelinone

[74260-87-0]

C₂₀H₂₄N₂O 308.422

Minor alkaloid from *Aristolochia serrata* (Elaeocarpaceae). Fine needles. Mp 320°. [α]_D¹⁹ +122.7 (MeOH/CHCl₃ 1:1). Crystal change to larger needles at ca. 255°.

8-Oxo, 9,10-didehydro: Makonine

[79592-57-7]

C₂₀H₂₂N₂O 306.407

Alkaloid from *Aristolochia serrata* (Elaeocarpaceae). Cryst. (MeOH). Mp 310-312° dec. [α]_D¹⁹ +431.1 (c, 0.93 in MeOH + CHCl₃).

13-Oxo: Aristoserratine

[76202-18-1]

C₂₀H₂₄N₂O 308.422

Minor alkaloid from *Aristolochia peduncularis* and *Aristolochia serrata* (Elaeocarpaceae). Mp 199°. [α]_D¹⁹ +22.5 (c, 1.9 in CHCl₃). λ_{\max} 228 (log ϵ 4.42); 282 (log ϵ 3.82); 290 (log ϵ 3.75) (EtOH).

13-Oxo, 15,16-didehydro: Peduncularistine

[90930-65-7]

C₂₀H₂₂N₂O 306.407

Alkaloid from *Aristolochia peduncularis* (Elaeocarpaceae). Amorph. λ_{\max} 228 (log ϵ 4.56); 278 (sh) (log ϵ 3.89); 286 (log ϵ 3.93); 294 (log ϵ 3.88) (EtOH).

(-)-form

Alkaloid from *Aristolochia chilensis* (Elaeocarpaceae). Mp 164-165°. [α]_D²⁵ -23 (c, 0.3 in CHCl₃). Not indexed by CAS.

(±)-form [79171-58-7]

Synthetic. Mp 173-175°.

13-Oxo: Synthetic. Yellow amorph. powder.

Anderson, B.F. *et al.*, *Chem. Comm.*, 1975, 511-512 (*cryst struct*)

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1976, **15**, 574-575 (*isol, uv, ir, pmr, ms*)

Hai, M.A. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 2130-2134 (*Aristoserratine*)

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1980, **21**, 545-546 (*Aristotelinone*)

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 2555-2561 (*isol, uv, ir, pmr, ms, cd*)

Bick, I.R.C. *et al.*, *Heterocycles*, 1981, **16**, 1301-1303 (*Makonine*)

Mirand, C. *et al.*, *J.O.C.*, 1982, **47**, 4169-4170 (*synth*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1037-1042 (*Aristoserratine, cryst struct*)

Stevens, R.V. *et al.*, *Chem. Comm.*, 1983, 384-386 (*synth*)

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 804-818 (*Peduncularistine*)

Gribble, G.W. *et al.*, *J.O.C.*, 1985, **50**, 5900-5902 (*synth, ir, pmr, cmr, ms*)

Kan-Fan, C. *et al.*, *Tetrahedron*, 1988, **44**, 1651-1660 (*Aristocarbinoil, 9,10-Dehydroaristoteline*)

Burkard, S. *et al.*, *Helv. Chim. Acta*, 1989, **72**, 254-263 (*Aristoserratine*)


Quirion, J.-C. *et al.*, *J.O.C.*, 1992, **57**, 5848-5851 (*pmr, cmr, struct*)

Stoermer, D. *et al.*, *J.O.C.*, 1993, **58**, 564-568 (*synth*)

Aristoyagonine

A-1443

[95377-98-3]

C₁₉H₁₇NO₅ 339.347


Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Yellow needles (MeOH). Mp 165-166°. λ_{\max} 220 ; 230 ; 250 ; 296 ; 330 (sh) (EtOH).

Campello, M.J. *et al.*, *Tet. Lett.*, 1984, **25**, 5933

(uv, ir, pmr, cmr, ms)
de Lera, A.R. *et al.*, *J. Het. Chem.*, 1987, **24**, 313 (*synth*)
Tojo, E. *et al.*, *Phytochemistry*, 1991, **30**, 1005-1010 (*isol, uv, ir, pmr, cmr, ms*)
Moreau, A. *et al.*, *J.O.C.*, 2004, **69**, 4527-4530 (*synth*)

Artemepavine **A-1444**

4-[(1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)methyl]phenol, 9CI. 1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-6,7-dimethoxyisoquinoline. *Evoeuropine*



(R)-form

C₁₉H₂₃NO₃ 313.396

(R)-form [524-20-9]

Alkaloid from *Papaver armeniacum* and many other spp. in several different families (Papaveraceae). Convulsive and irritant agent. Shows no spasmolytic, parasympatholytic or anaesthetic props. Produces cardiac irregularity. Mp 148-149° (138-148°). [α]_D¹⁸ -109 (CHCl₃).

Hydrochloride: Mp 151-152°.

Me ether: (-)-O-Methylartemepavine

[5701-00-8]
C₂₀H₂₅NO₃ 327.422

Alkaloid from *Discaria serratifolia*, *Xylopia pancheri* and *Aconitum leucostomum* (Rhamnaceae, Annonaceae, Ranunculaceae). Cryst. (petrol). Mp 65°. [α]_D²⁰ -81 (c, 1.2 in CHCl₃).

Me ether, N-oxide: N-Methylartemepavine N-oxide

C₂₀H₂₅NO₄ 343.422

Alkaloid from *Xylopia pancheri* (Annonaceae). Cryst. (Me₂CO). Mp 170-172°. [α]_D -7.9 (EtOH).

N-De-Me: Norartemepavine. N-Demethylartemepavine

C₁₈H₂₁NO₃ 299.369

Alkaloid from the wood of *Magnolia kachirachirai*. Mp 157-158°. [α]_D²² +31.5.

(S)-form

From reductive cleavage of dimeric isoquinoline alkaloids, e.g. Stepinonine. Mp 149-150°. [α]_D²⁰ +118 (CHCl₃).

Me ether: (+)-O-Methylartemepavine

[3423-02-7]
C₂₀H₂₅NO₃ 327.422

Alkaloid from *Annona squamosa* (sugar apple) and *Magnolia acuminata* (Annonaceae, Magnoliaceae). Cryst. (petrol). Mp 60-62°. [α]_D²⁰ +85.

Me ether, hydrochloride:

Cryst. (CHCl₃/petrol). Mp 198-199°.

O,N-Di-Me: Zanoxyline

[73529-17-6]
C₂₁H₂₈NO₃ 342.457

Quaternary alkaloid from the stem bark of *Zanthoxylum oxyphyllum*. Mp 128° (as iodide) (nat.) Mp 135-137° (sinters at 128°) (iodide) (synthetic). [α]_D²⁰ +66.3 (MeOH) (hydroxide) (nat.).

[α]_D²⁴ +91.7 (c, 0.193 in MeOH) (iodide) (synthetic). No abs. config. was assigned to the natural alkaloid but from the reported opt. rotation it is clearly S-.

N-De-Me: Alkaloid from leaves and stems of *Nelumbo lutea* and wood of *Machilus kusanoi*. Weak mydriatic and local anaesthetic. Cryst. (Me₂CO). Mp 152-153°. [α]_D²⁵ -23 (c, 1.33 in CHCl₃).

N-De-Me, hydrochloride:
Cryst. (MeOH). Mp 159-162°.

(±)-form

Synthetic. Mp 166°.

Me ether: [1934-93-6]
Synthetic. Prismatic needles (Et₂O). Mp 92° (62.5-63°).

O,N-Di-Me:

Small yellow prisms (MeOH) (as iodide). Mp 135-136° (iodide).

Marion, L. *et al.*, *J.O.C.*, 1950, **15**, 216 (*deriv, synth*)

Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1953, **1**, 10 (*deriv, synth*)

Ferrari, C. *et al.*, *Tetrahedron*, 1962, **18**, 419 (*deriv, synth*)

Kupchan, S.M. *et al.*, *Tetrahedron*, 1963, **19**, 227 (*isol, uv, ir, pmr, struct, Norarmepavine*)

Yang, T.H. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 22; *CA*, **59**, 3974d (*isol, uv, ir, struct, Norarmepavine*)

Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1964, **53**, 1140 (*deriv, isol, pmr, struct*)

Craig, J.C. *et al.*, *Tetrahedron*, 1965, **21**, 401 (*ord*)

Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1966, **14**, 232 (*ms*)

Albonico, S.M. *et al.*, *J.C.S.(C)*, 1966, 1340 (*config*)

Cava, M.P. *et al.*, *Tetrahedron*, 1969, **25**, 2795 (*synth*)

Bishay, D.W. *et al.*, *Phytochemistry*, 1973, **12**, 693 (*isol, uv, ir, pmr, ms*)

Niéto, M. *et al.*, *Planta Med.*, 1976, **30**, 48 (*oxide*)

Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 1636 (*synth*)

Sheth, J.P. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 595 (*deriv, synth, ms*)

Shono, T. *et al.*, *Tet. Lett.*, 1978, 4819 (*deriv, synth*)

Torres, R. *et al.*, *J. Nat. Prod.*, 1979, **42**, 430 (*deriv, isol, uv, ir, pmr, ms*)

Tiwari, K.P. *et al.*, *Phytochemistry*, 1979, **18**, 517 (*Zanoxyline*)

Bhaumik, P.K. *et al.*, *Phytochemistry*, 1979, **18**, 1584 (*deriv, isol, uv, pmr*)

Marek, R. *et al.*, *Coll. Czech. Chem. Comm.*, 1997, **62**, 1623-1630 (*isol, pmr, N-15 nmr, cryst struct*)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Baxendale, I.R. *et al.*, *Heterocycles*, 2003, **60**, 2707-2715 (*Norarmepavine, synth*)

Arnoamine A

A-1445

Pyrido[4,3,2-mn]*pyrrolo*[3,2,1-de]*acridin-4-ol*, 9CI. 4-Hydroxypyrido[4,3,2-mn]*pyrrolo*[3,2,1-de]*acridine*, 9CI [202652-44-6]



C₁₇H₁₀N₂O 258.279

Alkaloid from the ascidian *Cystodytes* sp. Yellow glass. λ_{max} 278 (log ε 4.11); 401 (log ε 3.09); 486 (log ε 3.17) (trifluoroacetic acid/MeOH). λ_{max} 258 (ε 14125); 328 (ε 7762); 431 (ε 1023) (MeOH) (Berdy).

Me ether: 4-Methoxypyrido[4,3,2-mn]*pyrrolo*[3,2,1-de]*acridine*. **Arnoamine B** [202652-47-9]

C₁₈H₁₂N₂O 272.306

Alkaloid from *Cystodytes* sp. Yellow glass. λ_{max} 280 (log ε 4.15); 398 (log ε 3.15); 467 (log ε 3.22) (trifluoroacetic acid/MeOH). λ_{max} 254 (ε 13800); 309 (ε 7582); 400 (ε 1483) (MeOH) (Berdy).

Plubrukarn, A. *et al.*, *J.O.C.*, 1998, **63**, 1657-1659 (*isol, uv, ir, pmr, cmr, ms*)


Delfourne, E. *et al.*, *J.O.C.*, 2000, **65**, 5476-5479 (*synth*)

Nakahara, S. *et al.*, *Heterocycles*, 2007, **71**, 1801-1806 (*Arnoamine B, synth*)

Arnottianamide

A-1446

N-[6-(2-Hydroxy-3,4-dimethoxyphenyl)-naphtho[2,3-d]-1,3-dioxol-5-yl]-N-methylformamide, 9CI [60394-88-9]



C₂₁H₁₉NO₆ 381.384

A secobenzo[c]phenanthridine alkaloid. Alkaloid from the bark of *Zanthoxylum* (*Zanthoxylum*, *Fagara*) *arnottianum*, *Zanthoxylum cuspidatum* and *Zanthoxylum ailanthoides*, the root wood of *Zanthoxylum integrifolium*, the roots of *Toddalia asiatica*, and *Zanthoxylum bungeanum*, and the root bar. Cryst. (EtOAc/MeOH or CHCl₃/MeOH). Mp 271-273° (264-267°, 267-269°).

Ac: Mp 237°.

O¹⁰-De-Me: Iwamide

[61774-74-1]

C₂₀H₁₇NO₆ 367.357

Minor alkaloid from the bark of *Zanthoxylum arnottianum* (Rutaceae). Prisms (CHCl₃/MeOH). Mp 271-273° (267-268.5°).

9-Demethoxy, 11-methoxy: Isoarnottianamide

[60394-91-4]

C₂₁H₁₉NO₆ 381.384

Alkaloid from the bark of *Zanthoxylum cuspidatum* (Rutaceae). Cryst. (CHCl₃). Mp 254-257° dec.

9-Demethoxy, 11-methoxy, O¹⁰-de-Me, O⁸-Me: Turraeanthine A


C₂₁H₁₉NO₆ 381.384

Alkaloid from the stem bark of *Turraeanthus africanus*. Brown crystal. (Me₂CO). Mp 237-239°. λ_{max} 237 (log ε 4.73); 291 (log ε 4); 332 (log ε 3.85)

(MeOH).

- Ishii, H. *et al.*, *Tet. Lett.*, 1976, 1203 (*Arnottianamide*)
 Ishii, H. *et al.*, *Yakugaku Zasshi*, 1976, **96**, 1458; 1977, **97**, 890 (*Iwamide*, *Isoarnottianamide*, *isol*, *struct*, *ir*, *pmr*, *synth*)
 Chou, F.Y. *et al.*, *Heterocycles*, 1977, **7**, 969 (*isol*)
 Sharma, P.N. *et al.*, *Phytochemistry*, 1982, **21**, 252 (*isol*)
 Vardamides, J.C. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 1034-1036 (*Turraeanthine A*)

Aromadendrane 1-isonitrile A-1447
1-Isocyanoaromadendrane

(1 α ,4 β ,5 α ,6 β ,7 β ,10 α)-formC₁₆H₂₅N 231.38**(1 α ,4 β ,5 α ,6 β ,7 β ,10 α)-form***Isothiocyanate*:C₁₆H₂₅NS 263.446

Constit. of *Axinyssa aphysinoides*. Oil. [α]_D +128 (c, 0.24 in CHCl₃). Has -NCS replacing -NC.

(1 β ,4 β ,5 α ,6 β ,7 β ,10 β)-form [112757-34-3]

Metab. of marine sponge *Acanthella acuta*. Ichthyotoxin, spongicide. Cryst. Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O. Mp 65-66°. [α]_D -13.7 (c, 1.4 in CHCl₃).

Isothiocyanate: Aromadendrane 1-isothiocyanate. 1-Isothiocyanatoaromadendrane

[112767-01-8]

C₁₆H₂₅NS 263.446

Metab. of *Acanthella acuta*. Oil. [α]_D -32.8 (c, 0.7 in CHCl₃).

Formamide: 1-Formamidoaromadendrane

[126609-93-6]

C₁₆H₂₇NO 249.395


Constit. of *Hexabranchus sanguineus*. Oil. [α]_D²⁰ +16.7 (c, 1 in CHCl₃). Has -NHCHO replacing -NC.

Mayol, L. *et al.*, *Tetrahedron*, 1987, **43**, 5381-5388 (*Acanthella acuta* constits)

He, H.-Y. *et al.*, *J.O.C.*, 1992, **57**, 3191-3194 (*Axinyssa aphysinoides* constiti)

Zhang, W. *et al.*, *Tetrahedron*, 2007, **63**, 4725-4729 (*formamide*)

Artabonatine A A-1448
 [247244-62-8]



Relative Configuration

C₁₈H₁₃NO₄ 307.305

Alkaloid from fresh unripe fruit of *Artabotrys uncinatus*. Amorph. powder. [α]_D²⁴ -102.7 (c, 0.4 in CHCl₃). λ_{\max} 205 (log ϵ 4.11); 275 (log ϵ 3.94); 325 (log ϵ

3.76) (EtOH).

3-Methoxy: Artabonatine E


[368422-66-6]

C₁₉H₁₅NO₅ 337.331Alkaloid from *Artabotrys uncinata*.

Amorph. powder. [α]_D²⁴ -116.7 (c, 0.8 in CHCl₃). λ_{\max} 250 (log ϵ 3.94); 280 (log ϵ 3.76); 325 (log ϵ 2.64) (EtOH).

Hsieh, T.-J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1192-1193; 2001, **64**, 1157-1161 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Artabonatine C A-1449
1,4,5-Trimethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI
 [368422-64-4]

C₁₉H₁₅NO₄ 321.332Alkaloid from *Artabotrys uncinata*.

Green amorph. powder. λ_{\max} 268 (log ϵ 3.34); 436 (log ϵ 3.42); 592 (log ϵ 4.62) (EtOH).

O⁵-De-Me: Artabonatine D


[368422-65-5]

C₁₈H₁₃NO₄ 307.305

Constit. of *Artabotrys uncinata*. Red amorph. powder. λ_{\max} 204 (log ϵ 3.04); 280 (log ϵ 3.71); 506 (log ϵ 4.52) (EtOH).

Hsieh, T.-J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1157-1161

Artacinate A-1450
 [123131-49-7]

C₁₉H₂₁NO₄ 327.379

Unusual reduced aporphine. Alkaloid from the stem parts of *Artabotrys uncinatus* (Annonaceae). Greenish prisms (CHCl₃/Me₂CO). Mp 160-162°. [α]_D²⁴ 0 (c, 0.1 in MeOH). λ_{\max} 214 (log ϵ 5.03); 261 (log ϵ 4.92); 290 (sh) (log ϵ 4.55); 344 (sh) (log ϵ 4.43); 382 (log ϵ 4.79) (no solvent reported).

4,5-Dioxo: 4,5-Dioxoartacinate

[1001858-73-6]

C₁₉H₁₇NO₆ 355.346

Alkaloid from the stems of *Artabotrys uncinatus*. Yellow needles. Mp 167-169°. [α]_D +52.2 (c, 0.03 in MeOH). λ_{\max} 230; 252; 282; 345; 355; 396 (no solvent reported).

Wu, Y.-C. *et al.*, *Phytochemistry*, 1989, **28**, 2191-2195 (*Artacinate*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cryst* *struct*)

Lan, Y.-H. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 1597-1599 (*pmr*, *cmr*, *4,5-Dioxoartacinate*)

Arthrobactin A-1451

Terregens factor


[39007-57-3]

[AcN(OH)(CH₂)₅NHCOCH₂]₂C(OH)-COOHC₂₀H₃₆N₄O₉ 476.526Growth factor from *Arthrobacter pas-**cens*. Microbial iron chelator. Powder.Sol. H₂O; poorly sol. CHCl₃, hexane. λ_{\max} 405 (ϵ 1590) (pH 7.6 buffer) (Berdy). λ_{\max} 477 (ϵ 940) (HCl) (Berdy).

Linke, W.D. *et al.*, *Arch. Microbiol.*, 1972, **85**, 44

Lee, B.H. *et al.*, *J.O.C.*, 1983, **48**, 24 (*synth*)

Arundamine A-1452
 [475977-53-8]

C₂₃H₂₈N₄O 376.5Alkaloid from the roots of *Arundo donax*.Cryst. + 1H₂O (EtOH). Mp 104-105°.**N⁻Ac: Arundacine**C₂₅H₃₀N₄O₂ 418.538

Alkaloid from the roots of *Arundo donax*. Cryst. (Me₂CO). Mp 192-193°.

N⁻Me: Arundanine

[618852-71-4]


C₂₄H₃₀N₄O 390.527Constit. of the roots of *Arundo donax*.Cryst. (CHCl₃/hexane). Mp 198-199°. λ_{\max} 223 (log ϵ 4.65); 284 (log ϵ 4.07) (EtOH).

Zhalolov, I.Z. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2002, **38**, 83-86; 276-279;

280-283 (*isol*, *pmr*, *cmr*, *cryst* *struct*)

Khuzhaev, V.U. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2003, **52**, 745-747 (*Arundanine*)

Arundarine A-1453
 [844830-22-4]

C₂₄H₂₈N₄O 388.511


Alkaloid from the roots of *Arundo donax*. Cryst. Poorly sol. CHCl_3 , MeOH, Me_2CO . Mp 250-252°.

Khuzhaev, V.U. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2004, **53**, 1765-1767 (*isol, pmr, cmr*)

Arundinine

A-1454

[246856-96-2]




Absolute Configuration

 $\text{C}_{23}\text{H}_{28}\text{N}_4\text{O}$ 376.5Alkaloid from *Arundo donax*. Mp 148-150°.

Zhalolov, I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1998, **34**, 706-710 (*isol, cryst struct*)

Asaramide

A-1455

 $\text{C}_{16}\text{H}_{25}\text{NO}_2$ 263.379


Alkaloid from *Asiasarum sieboldii*. Amorph. solid. λ_{max} 244 (ϵ 24145); 264 (ϵ 23710) (MeOH).

Opitz, M. *et al.*, *Pharmazie*, 1999, **54**, 218-223

Ascidiacyclamide

A-1456

[86701-12-4]

 $\text{C}_{36}\text{H}_{52}\text{N}_8\text{O}_6\text{S}_2$ 756.989

Cyclic peptide antibiotic. Prod. by an unidentified ascidian. Cytotoxic. Active against PV_4 cultured cells transformed with Polyoma virus. Prisms (C_6H_6). Sol. Me_2CO , CHCl_3 ; poorly sol. H_2O . Mp 245-246° (139-139.5°). $[\alpha]_{\text{D}}^{25} +164$ (c, 0.466 in CHCl_3). λ_{max} 232 (ϵ 21000) (MeOH) (Derep).

Hamamoto, Y. *et al.*, *Chem. Comm.*, 1983, 323-324 (*struct, pmr, cmr*)

Hamada, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 3223 (*synth, config*)


Ishida, T. *et al.*, *J.O.C.*, 1988, **53**, 107 (*struct, conformn*)

In, Y. *et al.*, *Acta Cryst. C*, 1994, **50**, 2015 (*cryst struct*)

Ascidiathiazone A

A-1457

[880137-41-7]

 $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_6\text{S}$ 308.271


Isol. from *Aplidium* sp. Antiinflammatory agent. Yellow powder. Mp 155° dec. λ_{max} 216 (log ϵ 4.37); 236 (log ϵ 4.46); 268 (log ϵ 4.21); 416 (log ϵ 3.4) (MeOH). λ_{max} 205 (log ϵ 4.43); 237 (log ϵ 4.55); 274 (log ϵ 4.18); 421 (log ϵ 3.56) (MeOH/trifluoroacetic acid). λ_{max} 211 (log ϵ 5.07); 236 (log ϵ 4.52); 329 (log ϵ 3.97); 467 (log ϵ 3.59) (MeOH/KOH).

Pearce, A.N. *et al.*, *J. Nat. Prod.*, 2007, **70**, 936-940 (*isol, synth, uv, pmr, cmr*)

Ascidiathiazone B

A-1458

[880137-42-8]

 $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_6\text{S}$ 306.255


Isol. from *Aplidium* sp. Pink powder. Mp 280° dec. λ_{max} 214 (log ϵ 4.5); 237 (log ϵ 4.02); 269 (log ϵ 3.91); 417 (log ϵ 3.11) (MeOH/trifluoroacetic acid). λ_{max} 207 (log ϵ 4.95); 237 (log ϵ 4.04); 279 (log ϵ 3.81); 311 (log ϵ 3.62); 472 (log ϵ 3.29) (MeOH/KOH).

Pearce, A.N. *et al.*, *J. Nat. Prod.*, 2007, **70**, 936-940 (*isol, uv, pmr, cmr*)

Ascidieminn

A-1459

9*H*-Quino[4,3,2-*de*][1,10]phenanthrolin-9-one, 9*Cl*. Leptoclinidinone. Ascidieminn [114622-04-7]

 $\text{C}_{18}\text{H}_9\text{N}_3\text{O}$ 283.289

CAS numbering shown. Other systems have been used. The name Ascidieminn found in the lit. appears to be an error. Alkaloid from the Okinawan tunicate *Didemnum* sp., *Cystodytes delle chiaiei*, *Didemnum rubeum* and the Seychelles

tunicate *Eudistoma* sp. Exhibits potent antineoplastic activity. Ca release agent. DNA intercalator and cleavage agent. Yellow solid. Sol. MeOH, EtOAc. Mp 300°. λ_{max} 220 (ϵ 49500); 248 (ϵ 48000); 273 (sh) (ϵ 27500); 298 (ϵ 17000); 308 (ϵ 15700); 340 (sh) (ϵ 11300); 377 (ϵ 13600) (MeOH) (Derep). λ_{max} 220 ; 247 ; 270 ; 308 ; 340 ; 375 (EtOH) (Berdy).

10-Hydroxy-11-Hydroxyascidieminn.

Cystodamine

[129741-41-9]

[158761-11-6]

 $\text{C}_{18}\text{H}_9\text{N}_3\text{O}_2$ 299.288

Alkaloid from the ascidian *Leptoclinides* sp. and *Cystodytes delle chiaiei*. Cytotoxic agent. Yellow amorph. solid. Mp 250°. Struct. of Cystodamine revised in 2000. λ_{max} 203 (ϵ 25000); 227 (ϵ 38000); 275 (ϵ 18000); 285 (ϵ 17000); 370 (ϵ 11000) (MeOH) (Derep).

10-Hydroxy, 12,13-dihydro-8,9-Dihydro-11-Hydroxyascidieminn

[151392-04-0]

 $\text{C}_{18}\text{H}_{11}\text{N}_3\text{O}_2$ 301.304

Alkaloid from the Okinawan marine sponge *Biemna* sp. Cytotoxic agent. Yellow amorph. powder. Mp 300°. λ_{max} 218 (ϵ 17000); 273 (ϵ 9300); 319 (ϵ 6700); 355 (ϵ 5300) (MeOH) (Berdy).

6-Bromo-2-Bromoleptoclinidinone

[109802-17-7]

 $\text{C}_{18}\text{H}_8\text{BrN}_3\text{O}$ 362.185

Alkaloid from an ascidian, prob. *Leptoclinides* sp. Toxic to lymphocytic leukaemia cells *in vitro*. Protein phosphatase inhibitor. Yellow powder ($\text{CHCl}_3/\text{MeOH}$). Sol. MeOH, CHCl_3 ; poorly sol. H_2O . Mp 300°. Revised struct. λ_{max} 227 (ϵ 152000); 247 (ϵ 27800); 254 (sh) (ϵ 27300); 278 (ϵ 32000); 298 (ϵ 30600); 335 (ϵ 18500); 371 (ϵ 21000) (EtOH) (Derep).

5-Hydroxy-Neocalliactine

[143370-24-5]

Obt. from Calliactine, C-34 by treatment with dil. acid. Cryst. ($\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$)(as Ac). Mp 280-290° (Ac).

Bloor, S.J. *et al.*, *J.A.C.S.*, 1987, **109**, 6134-6136 (2-Bromoleptoclinidinone)

Kitahara, Y. *et al.*, *Tetrahedron*, 1987, **53**, 17029-17038 (Neocalliactine)

Kobayashi, J. *et al.*, *Tet. Lett.*, 1988, **29**, 1177-1180 (*isol, uv, ir, pmr, cmr, ms, struct*)

Bracher, F. *et al.*, *Heterocycles*, 1989, **29**, 2093-2095 (*synth*)

De Guzman, F.S. *et al.*, *Tet. Lett.*, 1989, **30**, 1069-1070 (2-Bromoleptoclinidinone, *struct*)

Bracher, F. *et al.*, *Annalen*, 1990, 205; 1992, 1205-1207 (2-Bromoleptoclinidinone, Neocalliactine, *synth, ir, pmr, cmr, ms*)

Schmitz, F.J. *et al.*, *J.O.C.*, 1991, **56**, 804-808 (11-Hydroxyascidieminn)

He, H.-Y. *et al.*, *J.O.C.*, 1991, **56**, 5369-5371 (*isol*)

Moody, C.J. *et al.*, *Tetrahedron*, 1992, **48**, 3589-3602 (*synth*)

Zeng, C.-M. *et al.*, *Tetrahedron*, 1993, **49**, 8337-8342 (8,9-Dihydro-11-hydroxyascidieminn)


Gellerman, G. *et al.*, *Synthesis*, 1994, 239-241 (*synth*)

- Lindsay, B.S. *et al.*, *J. Chem. Crystallogr.*, 1998, **28**, 645-648 (2-Bromoleptoclidinone, *cryst struct*)
- Alvarez, M. *et al.*, *Eur. J. Org. Chem.*, 2000, 849-855 (*synth*)
- Delfourne, E. *et al.*, *Tet. Lett.*, 2000, **41**, 3863-3864 (Cystodamine, *struct*)
- Alvarez, M. *et al.*, *Tetrahedron*, 2000, **56**, 3703-3708 (11-Hydroxyascididemin, *synth*)
- Cuerva, J.M. *et al.*, *J.C.S. Perkin 1*, 2002, 1360-1365 (*synth*)

Ascochalasin

A-1460

[116095-60-4]



$C_{29}H_{39}NO_4$ 465.631
Isol. from *Ascochyta heteromorpha*.
Amorph. solid.

21,22-Didehydro(E-): Deoxaphomin

[51053-39-5]

 $C_{29}H_{37}NO_4$ 463.616


Prod. by *Ascochyta heteromorpha* and a *Phoma* sp. Active against phytopathogenic fungi. Amorph. powder. $[\alpha]_D^{25}$ -50.8 (c, 0.18 in $CHCl_3$). λ_{max} 217 (log ϵ 4.15); 234 (log ϵ 3.94) (MeOH). λ_{max} 215 (ϵ 10715); 237 (ϵ 7410) (MeOH) (Berdy).

- Binder, M. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 966-976 (Deoxaphomin)
- Capasso, R. *et al.*, *J. Nat. Prod.*, 1988, **51**, 567 (*isol, uv, ir, pmr, cmr, ms, struct*)
- Evidente, A. *et al.*, *Nat. Toxins*, 1998, **5**, 228-233; *CA*, **129**, 199012r (*activity*)

Ascorbigen

A-1461

2-C-(1H-Indol-3-ylmethyl)-3-hexulofuranosonic acid γ -lactone, 9CI
[8075-98-7]



(2R)-form

$C_{15}H_{15}NO_6$ 305.287
Present in plants, esp. cabbage and other crucifers. Bound form of Ascorbic acid.

(2R)-form β -L-lyxo-form. **Ascorbigen B**

[26548-49-2]

Light yellow amorph. powder. Mp 70° (sinters). $[\alpha]_D^{25}$ +12.5 (c, 1.0 in MeOH).

(2S)-form β -L-xylo-form. **Ascorbigen A**

[26676-89-1]

Amorph. powder. Mp ca. 65° (sinters). $[\alpha]_D^{25}$ +11 (c, 2.0 in EtOH).


[29400-26-8]

- Piironen, E. *et al.*, *Acta Chem. Scand.*, 1962, **16**, 1286 (*synth, bibl*)
- Kiss, G. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 989 (*synth, ir, uv, pmr, struct*)
- Korolev, A.M. *et al.*, *Bioorg. Khim.*, 1991, **17**, 981

Ascosalipyrrolidinone B

A-1462

[303052-08-6]



$C_{24}H_{35}NO_3$ 385.545
Prod. by the marine fungus *Ascochyta salicorniae*. Powder. Racemic. Related to Oteromycin, O-134. λ_{max} 207 (log ϵ 4.38); 225 (log ϵ 4.08); 273 (log ϵ 3.24) (EtOH).

O-Me, O-butyl: Ascosalipyrrolidinone A

[303052-07-5]

 $C_{27}H_{41}NO_3$ 427.626


Prod. by *Ascochyta salicorniae*. Antiplasmodial and antimicrobial agent. Inhibitor of tyrosine kinase. Amorph. powder. $[\alpha]_D^{20}$ -51.3 (c, 0.16 in EtOH). λ_{max} 211 (log ϵ 4.1); 225 (sh) (log ϵ 3.89); 267 (log ϵ 3.08) (EtOH).

Osterhage, C. *et al.*, *J.O.C.*, 2000, **65**, 6412-6417 (*Ascosalipyrrolidinones A, B*)

Asimicilone

A-1463

7-Hydroxy-3,4-dimethoxy-5-methylbenzo[h]quinolin-2(1H)-one, 9CI
[145701-06-0]




$C_{16}H_{15}NO_4$ 285.299
Alkaloid from twigs of *Asimina parviflora* (Annonaceae). Toxic to brine shrimp. Mp 191° (with dec. at 151°). λ_{max} 242 (ϵ 50100); 282 (ϵ 61700); 293 (ϵ 61700); 331 (ϵ 15500); 348 (ϵ 12900) (EtOH) (Derrep). λ_{max} 242 (ϵ 50120); 282 (ϵ 61660); 293 (ϵ 61660); 331 (ϵ 15500); 348 (ϵ 12880) (MeOH) (Berdy).

Ratnayake, S. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1462 (*isol, uv, ir, pmr, cmr, ms, struct*)

Asmarine I

A-1464

[805228-53-9]



$C_{25}H_{37}N_5O$ 423.6
Alkaloid from the sponge *Raspailia* sp. Oil. $[\alpha]_D^{25}$ +24 (c, 0.6 in MeOH).

Deoxy: Asmarine J

[805228-54-0]

 $C_{25}H_{37}N_5$ 407.601


Alkaloid from a *Raspailia* sp. Oil. $[\alpha]_D^{25}$ +20 (c, 0.6 in MeOH).

Rudi, A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1932-1935 (*isol, pmr, cmr*)

Asmarine B

A-1465

[208391-67-7]



$C_{25}H_{37}N_5O$ 423.6
Alkaloid from the sponge *Raspailia* sp. Cytotoxic agent. Oil. $[\alpha]_D^{25}$ +60 (c, 0.5 in $CHCl_3$). λ_{max} 292 (ϵ 1000) (MeOH) (Berdy).

Deoxy: Asmarine K

[805228-55-1]

 $C_{25}H_{37}N_5$ 407.601

Alkaloid from a *Raspailia* sp. Oil. $[\alpha]_D^{25}$ +28 (c, 0.5 in MeOH).

5-Epimer: Asmarine A

[208391-66-6]

 $C_{25}H_{37}N_5O$ 423.6

Alkaloid from a *Raspailia* sp. Cytotoxic agent. Cryst. (MeOH). Mp 232°. $[\alpha]_D^{25}$ +55 (c, 0.5 in $CHCl_3$). λ_{max} 292 (ϵ 10000) (MeOH) (Berdy).

5-Epimer, Me ether: Asmarine G

[662166-94-1]

 $C_{26}H_{39}N_5O$ 437.627

Alkaloid from a *Raspailia* sp. Oil. $[\alpha]_D^{25}$ +40 (c, 1.2 in CH_2Cl_2).

5-Epimer, deoxy: Asmarine H

[662166-95-2]

 $C_{25}H_{37}N_5$ 407.601

Alkaloid from a *Raspailia* sp. Oil. $[\alpha]_D^{25}$ +10 (c, 0.37 in CH_2Cl_2).


Yosief, T. *et al.*, *J. Nat. Prod.*, 2000, **63**, 299-304 (*Asmarines A, B, isol, pmr, cmr, crystal struct, activity*)

Rudi, A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 106-109; 1932-1935 (*Asmarines G,H,K*)

Asmarine C

A-1466

[208391-68-8]

C₂₆H₃₉N₅O 437.627Alkaloid from the sponge *Raspailia* sp. Oil.N^{10'}-Methoxy: *Asmarine F*

[261630-39-1]

C₂₇H₄₁N₅O₂ 467.653Constit. of a *Raspailia* sp.5-Epimer: *Asmarine D*


[261630-37-9]

C₂₆H₃₉N₅O 437.627Alkaloid from a *Raspailia* sp.5-Epimer, N^{10'}-methoxy: *Asmarine E*

[261630-38-0]


C₂₇H₄₁N₅O₂ 467.653Alkaloid from a *Raspailia* sp.Yosief, T. *et al.*, *J. Nat. Prod.*, 2000, **63**, 299-304 (*isol, pmr, cmr*)**L-Aspartyl-L-N²-hydroxyaspartyl-D-cycloserine**

A-1467

C₁₁H₁₆N₄O₉ 348.269Incorr. named by CAS. Isol. from cultures of *Corynebacterium kutscheri* grown on an Fe-limiting medium. Cryst. (EtOH aq.).McCullough, W.G. *et al.*, *J. Bacteriol.*, 1979, **137**, 243-247 (*isol, pmr, cmr*)**Asperazine**

A-1468

[198953-76-3]



Absolute Configuration


C₄₀H₃₆N₆O₄ 664.762

Alkaloid from the sponge-derived culture


of *Aspergillus niger*. Amorph. powder. [α]_D²⁵ +53 (c, 0.2 in MeOH). λ_{max} 225; 285; 300 (MeOH).Varoglu, M. *et al.*, *J.O.C.*, 1997, **62**, 7078-7079 (*isol, uv, ir, pmr, ms*)Govek, S.P. *et al.*, *Tetrahedron*, 2007, **63**, 8499-8513 (*synth*)**Asperfumoid**

A-1469

3-Hydroxy-2,5',6-trimethoxy-7'-methylspiro[2,5-cyclohexadiene-1,3'(2'H)-quinoline]-2',4,4'(1'H)-trione, 9CI [848073-02-9]

C₁₈H₁₇NO₇ 359.335Prod. by *Aspergillus fumigatus* CY018.Antifungal agent. Mp 284-286°. [α]_D²⁰ +55.6 (c, 0.45 in CHCl₃).Liu, J.Y. *et al.*, *J. Biotechnol.*, 2004, **114**, 279-287 (*isol, pmr, cmr, ms*)**Aspergamide B**

A-1470



(+) -form

C₂₆H₂₇N₃O₃ 429.518**(+) -form**Prod. by *Aspergillus ochraceus*. No phys. props. reported. Incorr. named in CAS.N¹⁶-Oxide: *Avrainvillamide*. *CJ* 17665.*Antibiotic CJ* 17665

[269741-97-1]

C₂₆H₂₇N₃O₄ 445.517Prod. by a marine *Aspergillus* sp.CNC358 and *Aspergillus ochraceus* CL41582. Cytotoxic and antibacterial agent. Powder or yellow gum. [α]_D²⁵ +12(c, 0.6 in MeOH). λ_{max} 280 (ε 6300);

370 (ε 2720) (MeOH).

16,21-Dihydro: Stephacidin A

[360765-74-8]

C₂₆H₂₉N₃O₃ 431.533Prod. by *Aspergillus ochraceus* WC76466. Cytotoxic. Amorph. solid. [α]_D+61.5 (c, 0.26 in CH₂Cl₂/MeOH). λ_{max}

211 (log ε 3.96); 242 (log ε 4.54); 309

(log ε 4.54); 335 (sh) (log ε 3.64)

(MeOH).

20β-Hydroxy, 20,21-dihydro, N¹⁶-oxide:**Aspergamide A**

[863238-39-5]


C₂₆H₂₉N₃O₅ 463.532Prod. by *Aspergillus ochraceus*. No

phys. props. reported.

(-) -form**16,21-Dihydro: (-)-Stephacidin A**C₂₆H₂₉N₃O₃ 431.533Prod. by *Aspergillus versicolor* NRRL35600. [α]_D²⁵ -32 (c, 0.05 in CHCl₃).U.S. Pat., 2000, 6 066 635; *CA*, **132**, 346709r (*Avrainvillamide*)Sugie, Y. *et al.*, *J. Antibiot.*, 2001, **54**, 911-916 (*CJ* 17665)U.S. Pat., 2001, 6 291 461; *CA*, **135**, 2364119 (*Stephacidin A*)Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (*Aspergamides A,B*)Qian-Cutrone, J. *et al.*, *J.A.C.S.*, 2002, **124**, 14556-14557 (*Stephacidin A*)Von Nussbaum, F. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3068-3071 (rev)Baran, P.S. *et al.*, *J.A.C.S.*, 2006, **128**,8678-8693 (*Stephacidin A, Avrainvillamide, synth*)Grubbs, A.W. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2257-2261; 2262-2265 (*Stephacidin A, synth*)Greshock, T.J. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 3573-3577 (*(-) -Stephacidin A*)**Aspergillamide A**

A-1471

[217489-31-1]

C₂₈H₃₄N₄O₃ 474.602Prod. by a marine *Aspergillus* sp. Cyto-toxic agent. Amorph. powder. [α]_D -26.2

(c, 3.05 in MeOH). Exists as a 1:1 mixt.

of *cis*- and *trans*-amide rotational con-formers. λ_{max} 200 (log ε 4.68); 221 (log ε

4.59); 286 (log ε 4.44) (MeOH).

(E)-Isomer: Aspergillamide B


[217489-32-2]

C₂₈H₃₄N₄O₃ 474.602Prod. by a marine *Aspergillus* sp.

Amorph. powder.

Toske, S.G. *et al.*, *Tetrahedron*, 1998, **54**,13459-13466 (*isol*)Rivas, L. *et al.*, *Tet. Lett.*, 2002, **43**, 7639-7641(*Aspergillamide B, synth*)Su, S. *et al.*, *Tetrahedron*, 2003, **59**, 8931-8946(*synth, pmr, cmr*)**Aspergillazine A**

A-1472

C₂₀H₂₀N₂O₈S 448.453

Related to Trichodermamide A, T-487.

Prod. by *Aspergillus unilateralis* (MST-F8675) and the marine-derived *Spicaria**elegans*. Cytotoxic. Amorph. yellow sol-id. [α]_D -103 (c, 0.14 in MeOH). [α]_D -366.4 (c, 0.11 in MeOH). λ_{max} 205 (ε

19270); 245 (sh); 337 (ε 11100) (MeOH).

λ_{max} 270 (log ε 3.95); 338 (log ε 4.29)


(MeOH).

Liu, R. *et al.*, *Arch. Pharmacol. Res.*, 2005, **28**,1042-1046 (*isol, pmr, cmr, ms*)

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 123-129 (*isol, pmr, cmr*)

Aspergillazine B

A-1473



$C_{20}H_{22}N_2O_8S$ 450.468

Related to Trichoderamide A, T-487. Prod. by *Aspergillus unilateralis* (MST-F8675). Amorph. yellow solid. $[\alpha]_D -172$ (c, 0.07 in MeOH). λ_{max} 203 (€ 17850); 243 (sh); 335 (€ 9400) (MeOH).

2-Epimer: Aspergillazine C

$C_{20}H_{22}N_2O_8S$ 450.468

Prod. by *Aspergillus unilateralis* (MST-F8675). Amorph. yellow solid. $[\alpha]_D +23$ (c, 0.08 in MeOH). λ_{max} 205 (€ 16400); 243 (sh); 335 (€ 7700) (MeOH).

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 123-129 (*isol, pmr, cmr*)

Aspergillazine D

A-1474

As Aspergillazine B, A-1473 with X = O

$C_{20}H_{22}N_2O_9$ 434.402

Prod. by *Aspergillus unilateralis* (MST-F8675). Amorph. yellow powder. $[\alpha]_D -79$ (c, 0.14 in MeOH). Isol. as an equilibrating mixt. with Aspergillazine E to which data refers. λ_{max} 204 (€ 17280); 228 (sh); 333 (€ 11400) (MeOH).

2-Epimer: Aspergillazine E

$C_{20}H_{22}N_2O_9$ 434.402


Prod. by *Aspergillus unilateralis* (MST-F8675).

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 123-129 (*isol, pmr, cmr*)

Aspergillilic acid, 8CI

A-1475

1-Hydroxy-6-(1-methylpropyl)-3-(2-methylpropyl)-2(1H)-pyrazinone, 9CI. Aspergillin†. Granegillin [490-02-8]



(S)-form

$C_{12}H_{20}N_2O_2$ 224.302

Identity of Granegillin with Aspergillilic acid is tentative.

► C19625000

(S)-form

Prod. by *Aspergillus flavus* and *Aspergillus sojae*. Shows antibiotic and hypotensive props. Pale yellow needles (MeOH) with odour of black walnuts. Sol. most org. solvs.; insol. H_2O . Mp 97-99°. $[\alpha]_D^{18} +13.3$ (c, 3.9 in EtOH). pK_a 5.5 (20°, H_2O). λ_{max} 238 (€ 10900); 326 (€ 8900) (EtOH) (Berdy). λ_{max} 235 (€

15900); 332 (€ 10400) (pH 7.5 buffer) (Berdy). λ_{max} 242 (€ 7300); 350 (€ 9700) (HCl) (Berdy).

► LD₅₀ (mus, ipr) 150 mg/kg, LD₅₀ (mus, orl) 250 mg/kg.

Hydrochloride: Mp 182° dec.

Fe complex: Ferriaspergillin

[77423-90-6]

$C_{36}H_{57}FeN_6O_6$ 725.73

Pigment from *Aspergillus flavus*, *Aspergillus parasiticus* and *Aspergillus subolivaceus*. Red oil.

Zn complex: Zn-Aspergillin

$C_{24}H_{38}N_4O_4Zn$ 511.979

Isol. from *Aspergillus sojae*.

Deoxy: Deoxyaspergillilic acid

[21641-71-4]

$C_{12}H_{20}N_2O$ 208.303

Metab. of *Aspergillus sojae*. Needles.

Mp 85° Mp 102°. No antibiotic activity.

(ξ)-form

1''ξ-Hydroxy: 1-Hydroxy-6-(1-hydroxy-1-methylpropyl)-3-(2-methylpropyl)-2(1H)-pyrazinone. Hydroxyaspergillilic acid

[4562-39-4]

$C_{12}H_{20}N_2O_3$ 240.302

Isol. from *Aspergillus flavus*, *Aspergillus sojae* and *Aspergillus oryzae*. Shows antibiotic props. Needles (hexane). Sol. MeOH, bases, C_6H_6 ; fairly sol. Et_2O ; poorly sol. H_2O , hexane. Mp 148-150°. $[\alpha]_D +36$ (c, 1 in EtOH). pK_a 4.9. λ_{max} 232 (€ 7500); 330 (€ 7900) (MeOH) (Berdy). λ_{max} 235 (€ 8250); 328 (€ 8800) (EtOH) (Berdy). λ_{max} 245 (E1%/1cm 440); 363 (E1%/1cm 310) (H_3PO_4) (Berdy).

► LD₅₀ (mus, ivn) 101 mg/kg, LD₅₀ (mus, ipr) 90 - 100 mg/kg.

Deoxy, 1''ξ-hydroxy: Deoxyhydroxyaspergillilic acid

$C_{12}H_{20}N_2O_2$ 224.302

Metab. of *Aspergillus sojae*. Mp 107°.

Dutcher, J.D. *et al.*, *J. Biol. Chem.*, 1947, **171**, 321-339; 341-353 (*Aspergillilic acid, Deoxyaspergillilic acid, isol, uv, struct*)

Newbold, G.T. *et al.*, *J.C.S.*, 1951, 2679 (*Deoxyaspergillilic acid*)

Gallagher, J.J. *et al.*, *J.C.S.*, 1952, 4870 (*synth*)

Cieleszky, V. *et al.*, *CA*, 1958, **52**, 15307 (*Granegillin*)

Dutcher, K. *et al.*, *J. Biol. Chem.*, 1958, **232**, 785 (*Hydroxyaspergillilic acid*)

Nakamura, T. *et al.*, *Bull. Agric. Chem. Soc. Jpn.*, 1959, **23**, 418 (*Hydroxyaspergillilic acid*)

MacDonald, J.C. *et al.*, *J. Biol. Chem.*, 1962, **237**, 1977 (*biosynth*)

Chigira, V.Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1966, **39**, 632 (*ir, uv, nmr, synth*)

Masaki, M. *et al.*, *J.O.C.*, 1966, **31**, 4143 (*synth, ir, uv, nmr*)

Bergeron, R.J. *et al.*, *J.O.C.*, 1980, **45**, 163 (*Deoxyaspergillilic acid*)

Assante, G. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 785 (*Ferriaspergillin*)

Ohta, A. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 20-24 (*Hydroxyaspergillilic acid, synth*)

Yamamoto, Y. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 185 (*rev, bibl, derivs*)


Okada, Y. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2259-2262 (*synth, abs config*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 794

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ARO000

Aspergillin PZ

A-1476



Relative Configuration


$C_{24}H_{35}NO_4$ 401.545

Cytochalasin related to Aspochalasin C, A-1502. Prod. by *Aspergillus awamori*. Square cryst. ($CHCl_3/MeOH$).

Zhang, Y. *et al.*, *J. Antibiot.*, 2002, **55**, 693-695 (*isol, pmr, cmr*)

Aspergillitine

A-1477



$C_{15}H_{13}NO_2$ 239.273


Prod. by the fungus *Aspergillus versicolor* isol. from the sponge *Xestospongia exigua*. Yellow powder (MeOH). λ_{max} 214; 245; 265; 295; 315; 370 (MeOH aq.).

Lin, W. *et al.*, *J. Nat. Prod.*, 2003, **66**, 57-61 (*isol, pmr, cmr, ms*)

Asperlicin

A-1478

[93413-04-8]



Absolute Configuration

R = H

$C_{31}H_{29}N_5O_4$ 535.601

Isol. from *Aspergillus alliaceus*. Nonpeptide cholecystokinin antagonist used for treatment of gastrointestinal and CNS disorders. Cryst. (MeOH). Sol. MeOH,

CHCl₃; poorly sol. H₂O. Mp 211-213°. [α]_D^{26.5} -185.3 (c, 1.1 in MeOH). Log P 3.08 (uncertain value) (calc). λ_{\max} 230 (sh) (ϵ 48500); 258 (sh) (ϵ 17700); 266 (sh) (ϵ 14400); 278 (sh) (ϵ 10300); 310 (ϵ 4080); 322 (sh) (ϵ 3100) (MeOH) (Derep).

N^{1'}-Hydroxy: Asperlicin B

[93413-08-2]

C₃₁H₂₉N₃O₅ 551.601

Minor metab. from *Aspergillus alliaceus*. Cholecystokinin antagonist (7 times more potent than Asperlicin). Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Log P 2.93 (uncertain value) (calc). λ_{\max} 230 (sh) (ϵ 48500); 258 (sh) (ϵ 17700); 266 (sh) (ϵ 14400); 278 (sh) (ϵ 10300); 310 (ϵ 4080); 322 (sh) (ϵ 3100) (MeOH) (Derep).

Albers-Schönberg, G. *et al.*, *CA*, 1985, **104**, 205424 (rev)

Chang, R.S.L. *et al.*, *J. Antibiot.*, 1985, **38**, 1633-1637; 1638-1641 (isol, uv, ir, pmr, cmr, struct)

Chang, R.S.L. *et al.*, *Science (Washington, D.C.)*, 1985, **230**, 177-179 (isol, props)

Goetz, M.A. *et al.*, *J. Antibiot.*, 1988, **41**, 875-877; 878-881; 882-891 (isol, pmr, uv, ms, struct, Asperlicin B, biosynth, derivs)

Wisner, J.R. *et al.*, *Pancreas*, 1988, **3**, 174-179 (pharmacol)

Monaghan, R.L. *et al.*, *J. Ind. Microbiol.*, 1989, **4**, 97-104 (isol, bibl)


Sun, H.H. *et al.*, *J. Antibiot.*, 1994, **47**, 599-601 (pmr, cmr)

He, F. *et al.*, *J.A.C.S.*, 1995, **120**, 6417-6418 (synth)

Asperlicin C

A-1479

6,7-Dihydro-7-(1H-indol-3-ylmethyl)quinazolino[3,2-a][1,4]benzodiazepine-5,13-dione, 9CI



C₂₅H₁₈N₄O₂ 406.443

(S)-form [93413-06-0]

Isol. from *Aspergillus alliaceus* ATCC 20655. Cholecystokinin antagonist. Off-white powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 222 (ϵ 56300); 268 (ϵ 15100); 289 (sh) (ϵ 8480); 310 (ϵ 4650); 321 (sh) (ϵ 3590) (MeOH) (Derep).

Bock, M.G. *et al.*, *J.O.C.*, 1987, **52**, 1644-1646 (synth)


Goetz, M.A. *et al.*, *J. Antibiot.*, 1988, **41**, 875-877; 878-881; 882-891 (isol, uv, ir, pmr, cmr, struct)

Liu, J.-F. *et al.*, *J.O.C.*, 2005, **70**, 10488-10493 (synth)

Asperlicin D

A-1480

7-(1H-Indol-3-ylmethyl)quinazolino[3,2-d][1,4]benzodiazepine-6,9(5H,7H)-dione, 9CI



C₂₅H₁₈N₄O₂ 406.443

(S)-form [93413-07-1]

Isol. from *Aspergillus alliaceus* ATCC 20655. Cholecystokinin antagonist. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 222 (ϵ 61100); 283 (sh) (ϵ 13800); 290 (ϵ 14600); 310 (ϵ 5920) (MeOH) (Derep).


Goetz, M.A. *et al.*, *J. Antibiot.*, 1988, **41**, 875; 878; 882 (isol, uv, ir, pmr, cmr)

Al-Said, N.H. *et al.*, *Tet. Lett.*, 2006, **47**, 693-694 (synth)

Asperlicin E

A-1481

13,17b,18,18a-Tetrahydro-17b-hydroxy-11H-indolo[3',2':4,5]pyrrolo[2,1-c]quinazolino[3,2-a][1,4]benzodiazepine-5,11(12aH)-dione, 9CI [93413-05-9]



Absolute Configuration

C₂₅H₁₈N₄O₃ 422.442

Isol. from *Aspergillus alliaceus* ATCC20655. Cholecystokinin antagonist. Off-white powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 213 (sh) (ϵ 37200); 227 (ϵ 38500); 268 (ϵ 8680); 277 (sh) (ϵ 7830); 310 (sh) (ϵ 4850); 324 (ϵ 3200) (MeOH) (Derep).


Bock, M.G. *et al.*, *J.O.C.*, 1987, **52**, 1644-1646 (synth)

Goetz, M.A. *et al.*, *J. Antibiot.*, 1988, **41**, 875; 878; 882 (isol, uv, ir, pmr, cmr, struct)

Liu, J.-F. *et al.*, *J.O.C.*, 2005, **70**, 10488-10493 (synth)

Aspernigerin

A-1482



C₂₆H₃₂N₄O₂ 432.564

Prod. by *Aspergillus niger* IFB-E003.


Cytotoxic. Cryst. (CHCl₃/MeOH). Mp 186-188°. λ_{\max} 215 (log ϵ 4.37); 246 (log ϵ 4.23) (MeOH).

Shen, L. *et al.*, *Chem. Eur. J.*, 2006, **12**, 4393-4396 (isol, pmr, cmr, ms, cryst struct)

Aspernigrin A

A-1483

1,4-Dihydro-4-oxo-6-(phenylmethyl)-3-pyridinecarboxamide. 2-Benzyl-5-carbamoyl-4(1H)-pyridinone [773855-60-0]



C₁₃H₁₂N₂O₂ 228.25

Struct. revised in 2005. Prod. by *Aspergillus niger* isol. from the sponge *Axinella damicornis*. Also prod. by the terrestrial *Cladosporium herbarum* IFB-E002 and *Pestalotiopsis theae*. Prisms (CHCl₃/MeOH). Mp 197-199°. λ_{\max} 217 (log ϵ 4.25); 257 (log ϵ 4.68) (MeOH).


Hiort, J. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1532-1543; 2005, **68**, 1821 (isol, pmr, cmr)

Ye, Y.H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1106-1108 (isol, pmr, cmr, ms, cryst struct)

Aspernigrin B

A-1484

[773855-63-3]



Absolute Configuration

C₂₇H₂₄N₂O₅ 456.497


Struct. revised in 2005. Prod. by *Aspergillus niger* isol. from the sponge *Axinella damicornis*. Neuroprotective agent. Oil. [α]_D²⁰ +37.8 (c, 0.5 in DMSO).

Hiort, J. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1532-1543; 2005, **68**, 1821 (isol, cd, pmr, cmr)

Aspernomine

A-1485

[138605-80-8]



Absolute Configuration

C₂₈H₃₉NO₂ 421.622


Prod. by *Aspergillus nomius* NRRL6552/NRRL18836. Exhibits cytotoxic and insecticidal activities. Mycotoxin. Fine needles. [α]_D²⁷ +225 (c, 0.12 in MeOH).

λ_{\max} 222 (ϵ 5460); 244 (ϵ 5760); 302 (ϵ 2510); 333 (ϵ 1150) (MeOH) (Derep).

Staub, G.M. *et al.*, *J.A.C.S.*, 1992, **114**, 1015-1017 (*isol*, *pmr*, *cmr*, *struct*)

Liu, Y. *et al.*, *Org. Lett.*, 2003, **3**, 333-335 (*biosynth*)

Asperparaline A A-1486
VM 55598. Antibiotic VM 55598. *Aspergillimide*
[195966-93-9]



Relative Configuration

R = CH₃

C₂₀H₂₉N₃O₃ 359.467

Related to Paraherquamide A, P-90. Prod. by *Aspergillus* sp. IMI 337664 and *Aspergillus japonicus* JV-23. Anthelmintic agent. Paralytic agent against *Bombyx mori*. Prisms (toluene). Mp 203-205°. [α]_D²⁰ -8.5 (c, 0.2 in MeOH). λ_{\max} 206 (no solvent reported).

N²-De-Me: Asperparaline B

[227598-73-4]

C₁₉H₂₇N₃O₃ 345.441

Prod. by *Aspergillus japonicus* JV-23. Prisms (EtOAc). Mp 250-252°. [α]_D²⁰ -3 (c, 0.08 in MeOH).

16-Oxo: 16-Ketoaspergillimide. 16-Ox-oaspergillimide. SB 202327. Antibiotic SB 202327

[199784-50-4]

C₂₀H₂₇N₃O₄ 373.451

Prod. by *Aspergillus* sp. IMI 337664. Anthelmintic agent.

Lower homologue (R = H): Asperparaline C

[227598-74-5]

C₁₉H₂₇N₃O₃ 345.441

Prod. by *Aspergillus japonicus* JV-23. Prisms (toluene). Mp 175-177°. [α]_D²⁰ -20 (c, 0.05 in MeOH).

Banks, R.M. *et al.*, *J. Antibiot.*, 1997, **50**, 840-846 (*isol*, *pmr*, *cmr*)

Hayashi, H. *et al.*, *Tet. Lett.*, 1997, **38**, 5655-5658 (*isol*, *ir*, *pmr*, *cmr*, *cryst struct*)

Hayashi, H. *et al.*, *CA*, 1999, **131**, 99865u (*Asperparaline B*)


Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 111-115 (*Asperparalines*)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (*rev*, *synth*, *biosynth*)

Gray, C. *et al.*, *J.A.C.S.*, 2003, **125**, 14692-14693 (*biosynth*)

Aspertine B A-1487

1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxyhexyl)pyridine
[442155-61-5]



Absolute Configuration

C₁₅H₂₉NO₂ 255.4

Alkaloid from the aerial parts of *Andrachne aspera*. Gum. [α]_D -85 (c, 0.8 in CHCl₃).

Ahmad, V.U. *et al.*, *Turk. J. Chem.*, 2002, **26**, 245-250 (*isol*, *pmr*, *cmr*)

Aspidexcelcine A-1488

[1354-51-4]

Bisindole alkaloid. Struct. unknown. Alkaloid from *Aspidosperma excelsum* (Apocynaceae). Mp 76-81°. [α]_D²⁰ +72 (c, 0.2 in CHCl₃). Mol. formula not recorded (MW 622 by ms).

Relyveld, P. *et al.*, *Pharm. Weekbl.*, 1965, **100**, 614-615; *CA*, **63**, 7254d (*isol*, *uv*, *ir*)

Aspidexcine A-1489

[1354-55-8]

C₄₂H₅₆N₄O₄ 680.929


Bisindole alkaloid. Struct. unknown. May be the same as Tetrahydrosecamine in S-174 (same source and MF, but no further information). Alkaloid from *Aspidosperma excelsum* and *Aspidosperma marcgravianum* (Apocynaceae). Sol. Me₂CO, H₂O. Mp 76-81°. [α]_D²⁰ -14 (c, 0.2 in CHCl₃). λ_{\max} 220; 283; 292 (MeOH) (Berdy).

Hydrochloride: Mp 212° dec.

Relyveld, P. *et al.*, *Pharm. Weekbl.*, 1965, **100**, 614-615; *CA*, **63**, 7254d (*isol*, *uv*, *ir*)

Aspidoalbidine A-1490

19,21-Epoxyaspidospermidine, 9CI. Fendleridine
[36459-35-5]



Absolute Configuration

C₁₉H₂₄N₂O 296.411

Alkaloid from seeds of *Aspidosperma fendleri* (Apocynaceae). Cryst. (Me₂CO). Mp 185-186°. λ_{\max} 242 (ϵ 7300); 293 (ϵ 3050) (EtOH).

N-Ac: N-Acetylaspidoalbidine. Dehydroxyhaplocidine

[2671-45-6]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Vallesia dichotoma* and bark of *Aspidosperma pyriformis* (Apocynaceae). Mp 173-174°. [α]_D +46 (CHCl₃) (+1). The low opt. rotn. of the *A. pyriformis* sample (Dehydroxyhaplocidine) suggests that it was largely racemic. λ_{\max} 212 (log ϵ 4.43); 253 (log ϵ 4.17) (EtOH).

12-Hydroxy, N-Ac: Haplocidine. 1-Acetyl-12-hydroxyaspidoalbidine

[2671-46-7]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Haplophyton cimidicum*, *Aspidosperma discolor* and *Vallesia dichotoma* (Apocynaceae). Cryst. (MeOH/Et₂O). Mp 184-189°. [α]_D²³ +214 (c, 0.35 in CHCl₃).

12-Hydroxy, N-propanoyl: Haplocine

[6879-62-5]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Haplophyton cimidicum*

(Apocynaceae). Mp 186-187°. [α]_D +196 (CHCl₃).

12-Methoxy: Fannine

[79405-65-5]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from stems of *Strempelepis strempeleioides* (Apocynaceae).

12-Methoxy, N-formyl: N-Formyl-12-methoxyaspidoalbidine. N-Formyl-17-methoxyaspidoalbidine

[2671-44-5]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Vallesia dichotoma* (Apocynaceae). Mp 208-210°. [α]_D +20 (CHCl₃).

12-Methoxy, N-propanoyl: [100979-51-9] Plates (MeOH). Mp 240-241°. [α]_D³⁰ +5 (c, 1.25 in CHCl₃).

11,12-Dihydroxy, N-Ac: Aspidoendlerine

[4913-97-7]

C₂₁H₂₆N₂O₄ 370.447

Constit. of the seeds of *Aspidosperma fendleri* (Apocynaceae). Cryst. (EtOH). Mp 278° dec. λ_{\max} 225 (ϵ 11500); 259 (ϵ 5000) (EtOH).

11-Methoxy, 12-hydroxy, N-Ac: Aspidoalbidine

[2494-58-8]

C₂₂H₂₈N₂O₄ 384.474

Constit. of *Aspidosperma limae*, *Aspidosperma fendleri* and *Aspidosperma album* (Apocynaceae). Prisms (MeOH). Mp 196-199°. [α]_D +245 (c, 1.26 in CHCl₃).

11-Methoxy, 12-hydroxy, N-propanoyl: Fendlerine. N-Propionyl-N-deacetylaspidolimidine

[4913-93-3]

C₂₃H₃₀N₂O₄ 398.501

Alkaloid from the stem bark of *Aspidosperma album*, also from fruit and bark of *Aspidosperma fendleri* (Apocynaceae). Glass or cryst. (Me₂CO). Mp 179-181°. [α]_D +226 (c, 1.08 in CHCl₃). λ_{\max} 225 (ϵ 19200); 258 (ϵ 3560) (EtOH).

10,11-Dimethoxy, 12-hydroxy, N-Ac: Kromantine. N-Acetyl-N-depropionylaspidoalbidine

[2122-26-1]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from the bark of *Aspidosperma album* and other *Aspidosperma* spp. (Apocynaceae). Mp 194-195° (176°). [α]_D +174 (c, 1.93 in CHCl₃) (+159). λ_{\max} 227 (log ϵ 4.43); 267 (log ϵ 4.11) (EtOH).

10,11-Dimethoxy, 12-hydroxy, N-propanoyl: Aspidoalbine

[2122-25-0]

C₂₄H₃₂N₂O₅ 428.527

Alkaloid from the bark of *Aspidosperma album* and other *Aspidosperma* spp. (Apocynaceae). Mp 174-177°. [α]_D²⁵ +164 (c, 2.12 in CHCl₃). λ_{\max} 227 (log ϵ 4.16); 267 (log ϵ 3.88) (EtOH).

10,11,12-Trimethoxy, N-propanoyl: Mp 128-131°. [α]_D +9.2 (c, 1.84 in MeOH).


Gilbert, B. *et al.*, *Chem. Ind. (London)*, 1962, 1949-1950 (*Aspidolimidine*, *pmr*, *ms*, *struct*)
Djerassi, C. *et al.*, *Tet. Lett.*, 1962, **3**, 1001-1009 (*Aspidoalbine*, *uv*, *ir*, *pmr*, *ms*, *struct*)

- Ferrari, C. *et al.*, *Can. J. Chem.*, 1963, **41**, 1531-1534 (*Aspidoalbine*, *N*-Acetyl-*N*-depropionylaspidoalbine)
- Relyveld, P. *et al.*, *Pharm. Weekbl.*, 1963, **98**, 47-48 (*Kromantine*)
- Brown, K.S. *et al.*, *Tet. Lett.*, 1963, **4**, 1731-1736 (*N*-Ac, *isol*, *ir*, *uv*, *ms*, *struct*)
- Ferrari, C. *et al.*, *Can. J. Chem.*, 1964, **42**, 2705-2709 (*Aspidolimidine*, *Fendlerine*)
- Cava, M.P. *et al.*, *Tetrahedron*, 1964, **20**, 581-583 (*Haplocidine*)
- Walser, A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 391 (*N*-Formyl-12-methoxyaspidoalbine, *Haplocidine*, *isol*)
- Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141-1166 (*Kromantine*)
- Burnell, R.H. *et al.*, *Can. J. Chem.*, 1966, **44**, 28-31 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *Aspidofendlerine*, *Fendlerine*)
- Dastoor, N.J. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 213-231 (*isol*, *uv*, *ir*, *ms*, *Haplocidine*)
- Ban, Y. *et al.*, *Tet. Lett.*, 1975, **16**, 727-730 (*synth*, *Aspidoalbidine*)
- Honma, Y. *et al.*, *Heterocycles*, 1976, **5**, 47-51 (*synth*, *ms*, *uv*, *Aspidoalbidine*)
- Laguna, A. *et al.*, *CA*, 1981, **95**, 165600v (*Fannine*)
- Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694-707 (*18-Oxohaplocidine*)
- Laguna, A. *et al.*, *Planta Med.*, 1984, **50**, 285-288 (*Fannine*)
- Overman, L.E. *et al.*, *J.A.C.S.*, 1991, **113**, 2598-2610 (*N*-Acetylaspidoalbine, *synth*)
- Mitaine, A.C. *et al.*, *Planta Med.*, 1996, **62**, 458 (*Dehydroxyhaplocidine*)
- Mitaine, A.-C. *et al.*, *Planta Med.*, 1998, **64**, 487 (*Fendlerine*, *pmr*, *cmr*)
- Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (*synth*, *rev*)

Aspidochibine

Alkaloid AQC2
[139955-86-5]

A-1491



Relative configuration

 $C_{19}H_{22}N_2O_3$ 326.394


New structural class of the quebrachamine series. Alkaloid from cell cultures of *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Amorph.

Aimi, N. *et al.*, *Tet. Lett.*, 1991, **32**, 4949 (*isol*, *pmr*, *cmr*, *struct*)

Aspidodasycarpine

Methyl 2,5-epoxy-1,2-dihydro-16-(hydroxymethyl)-4,5-secoakummlan-17-oate, 9CI
[2744-47-0]

A-1492



Absolute Configuration

 $C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from the bark of *Aspidosperma dasycarpon* and *Aspidosperma cuspa*

(Apocynaceae). Cryst. (Me₂CO). Mp 207-209°. [α]_D²⁵ -101 (c, 1.42 in CHCl₃).

N^d-Ac: Mp 250-253°. [α]_D -147.

N^d,O-Di-Ac:

Cryst. (Me₂CO). Mp 175°. [α]_D -174 (CHCl₃).

16-Epimer: **Lonicerine**[†]. 16-Epiaspidoasycarpine

[26988-11-4]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the root bark of *Callichilia barteri* (Apocynaceae). Plates (Me₂CO/hexane). Mp 105-107°. [α]_D²⁴ -127 (c, 0.98 in CHCl₃).

Joule, J.A. *et al.*, *Tetrahedron*, 1965, **21**, 1717 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Burnell, R.H. *et al.*, *Phytochemistry*, 1968, **7**, 2045 (*isol*, *uv*, *ir*, *pmr*, *ms*)


Naranjo, J. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 749 (*Lonicerine*)

Simões, J.C. *et al.*, *Phytochemistry*, 1976, **15**, 543 (*isol*)

Aspidodispermine

[21451-67-2]

A-1493



Absolute configuration

 $C_{19}H_{24}N_2O_3$ 328.41

Alkaloid from *Aspidosperma dispernum* (Apocynaceae). Noncryst. [α]_D +119 (MeOH).

O¹⁷-Me: Mp 189-191°. [α]_D -43 (MeOH).

O¹⁷-Me; hydrobromide: Mp 259-261°.

17-Deoxy: **Deoxyaspidodispermine**

[21446-30-0]

[54191-59-2]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from *Aspidosperma dispernum* (Apocynaceae). Amorph.; cryst. + 1H₂O (as picrate). Mp 184-186° (picrate). [α]_D -20 (MeOH).

Ikeda, M. *et al.*, *Tet. Lett.*, 1968, 5837 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Ling, N.C. *et al.*, *Tet. Lett.*, 1970, 3015 (*cryst struct*)

Ohnuma, T. *et al.*, *Chem. Comm.*, 1974, 296 (*synth*, *deriv*)

Ban, Y. *et al.*, *Lect. Heterocycl. Chem.*, 1976, **3**, 55 (*synth*)

Honma, Y. *et al.*, *Heterocycles*, 1977, **6**, 129 (*synth*)


Aspidofractinine

2,3,4,5,11,12-Hexahydro-6H,13aH-3a,5a-ethano-1H-indolizino[8,1-cd]carbazole, 9CI

[6871-25-6]

[60365-41-5 ((±)-form)]

A-1494



Absolute Configuration

 $C_{19}H_{24}N_2$ 280.412

Considerable confusion regarding the abs. configs. of this group of alkaloids. All appear to have the abs. config indicated. The key ref. is Guggisberg 1969 but this has often been overlooked in CAS and elsewhere. Note also that in *Aspidofractinine* and other alkaloids unsubstituted on the ethylene bridges, C-2 is not a chiral centre. Alkaloid from *Aspidosperma refractum*, the leaves of *Pleiocarpa tubicina*, and the stem and root barks of *Hunteria elliptica* (Apocynaceae). Prisms (pentane). Mp 101-102°. Bp_{0.001} 110°. [α]_D²³ -20 (c, 0.58 in CHCl₃). λ_{max} 241 (log ϵ 3.83); 291 (log ϵ 3.46) (EtOH).

N-Ac: Mp 127-130°. [α]_D²⁵ +27 (c, 0.25 in CHCl₃).

N^l-Me: N^l-Methylaspidofractinine

[132356-83-3]

C₂₀H₂₆N₂ 294.439

Alkaloid from the roots of *Vinca sardoa* (Apocynaceae). Amorph. solid. λ_{max} 208 (log ϵ 4.13); 257 (log ϵ 3.7); 300 (log ϵ 3.15) (MeOH).

14,15-Didehydro, N^l-formyl: N-Formyl-

14,15-didehydroaspidofractinine

C₂₀H₂₂N₂O 306.407

Alkaloid from *Vinca sardoa* (Apocynaceae). Cryst. (cyclohexane). Mp 138-140°. [α]_D²³ +55.8 (c, 0.5 in MeOH). CAS no. not found to 14CI. λ_{max} 211 (log ϵ 4.34); 254 (log ϵ 4.13); 278 (log ϵ 3.76); 287 (log ϵ 3.7) (MeOH).

14,15-Didehydro, N^l-Me: N^l-Methyl-

14,15-didehydroaspidofractinine

[131711-44-9]

C₂₀H₂₄N₂ 292.423

Alkaloid from the roots of *Vinca sardoa* (Apocynaceae). Cryst. (EtOH) (hydrobromide). Mp 285-295° (hydrobromide).

12-Hydroxy, N-Ac: **Aspidofiline**

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from the leaves of *Aspidosperma pyriformis* (Apocynaceae). Needles (Et₂O). Mp 190-191°. [α]_D²⁰ -174 (CHCl₃). Publ. details on abs. config. are fragmentary. The abs. config. given here is probable on biogenetic grounds. λ_{max} 231 (log ϵ 3.76); 308 (log ϵ 4.31) (EtOH/base). λ_{max} 258 (log ϵ 3.85); 282 (sh) (EtOH).

12-Hydroxy, N-Ac, picrate:

Cryst. (Me₂CO). Mp 146°.

12-Hydroxy, 14,15-didehydro, N^l-formyl:

N^l-Formyl-14,15-didehydro-12-hydro-

xyaspidofractinine

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Vinca sardoa* (Apocynaceae). Cryst. (cyclohexane). Mp 165° dec. [α]_D²⁰ -39 (c, 0.3 in MeOH). CAS no. not found to 14CI. λ_{max} 216 (log ϵ 4.41); 256 (log ϵ 3.94); 286 (log ϵ 3.69) (MeOH).

12-Hydroxy, 14,15-didehydro, N^l-Me:

14,15-Didehydro-12-hydroxy-N^l-methyl-

aspidofractinine. 12-Hydroxy-N^l-

methyl-14,15-didehydroaspidofractinine

C₂₀H₂₄N₂O 308.422

Alkaloid from *Vinca sardoa* (Apocyn-

naceae). Amorph. $[\alpha]_D^{23} +98.4$ (c, 0.2 in MeOH). CAS no. not found to 14CI. λ_{\max} 214 (log ϵ 4.3); 253 (log ϵ 3.76); 292 (log ϵ 3.46) (MeOH).

12-Methoxy-12-Methoxyaspidofractinine. 17-Methoxyaspidofractinine

$C_{20}H_{26}N_2O$ 310.438
Alkaloid from *Aspidosperma populifolium* (Apocynaceae). Cryst. (Me₂CO). Mp 128-130°. $[\alpha]_D^{25} +3$ (c, 0.67 in CHCl₃). CAS no. not found 8-14 CI.

12-Methoxy, N¹-formyl: N-Formyl-12-methoxyaspidofractinine

$C_{21}H_{26}N_2O_2$ 338.449
Alkaloid from *Aspidosperma populifolium* (Apocynaceae). Amorph. CAS no. not found 8-14CI. λ_{\max} 255 (log ϵ 4.04); 288 (log ϵ 3.42) (EtOH).

12-Methoxy, 14,15-didehydro, N¹-Me-14,15-didehydro-12-methoxy-N¹-methylaspidofractinine. 12-Methoxy-N¹-methyl-14,15-didehydroaspidofractinine

$C_{21}H_{26}N_2O$ 322.449
Alkaloid from *Vinca sardoa* (Apocynaceae). Cryst. (EtOAc/cyclohexane). Mp 138-140°. $[\alpha]_D^{23} +159$ (c, 0.1 in MeOH). CAS no. not found to 14CI. λ_{\max} 211 (log ϵ 4.35); 257 (log ϵ 3.96); 295 (log ϵ 3.56) (MeOH).

11,12-Dimethoxy: 11,12-Dimethoxyaspidofractinine. 16,17-Dimethoxyaspidofractinine

$C_{21}H_{28}N_2O_2$ 340.464
Alkaloid from *Aspidosperma populifolium* (Apocynaceae). Cryst. (Me₂CO). Mp 140-142°. $[\alpha]_D^{27} -141$ (c, 2.61 in CHCl₃). CAS no. not found. λ_{\max} 214 (log ϵ 4.5); 298 (log ϵ 3.39) (EtOH).

11,12-Dimethoxy, N¹-formyl: N-Formyl-11,12-dimethoxyaspidofractinine

$C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Aspidosperma populifolium* (Apocynaceae). Amorph. CAS no. not found. λ_{\max} 253 (log ϵ 3.86); 287 (log ϵ 3.42) (EtOH).

17-Oxo: see Kopsinone, K-80

Antonaccio, L.D. *et al.*, *J.O.C.*, 1960, **25**, 1262-1263 (*Aspidofiline*, *isol*, *uv*, *ir*)

Djerassi, C. *et al.*, *Experientia*, 1962, **18**, 397-398 (*Aspidofiline*, *pmr*, *ms*, *struct*)

Djerassi, C. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 742-751 (*ms*, *struct*)

Bycroft, B.W. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1147-1152 (*isol*, *uv*, *ir*, *pmr*)

Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141-1166 (*Aspidosperma populifolium* *constits*)

Klyne, W. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1168-1184 (*ord*)

Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 76-89 (*abs config*)

Ban, Y. *et al.*, *Tet. Lett.*, 1976, 1111-1114 (*synth*, *uv*, *ir*, *ms*)

Vercauteren, J. *et al.*, *Phytochemistry*, 1980, **19**, 1959-1961 (*isol*)

Craveiro, A.A. *et al.*, *Phytochemistry*, 1983, **22**, 1526-1528 (*Aspidofiline*, *isol*, *ir*, *pmr*)

Cartier, D. *et al.*, *Tet. Lett.*, 1989, **30**, 1951-1954 (*synth*)

Dufour, M. *et al.*, *Tet. Lett.*, 1989, **30**, 3429-3432 (*synth*)

Crippa, S. *et al.*, *Heterocycles*, 1990, **31**, 1663-1667 (*N-Methylaspidofractinine*)


Wenkert, E. *et al.*, *J.O.C.*, 1994, **59**, 7677-7682 (*synth*)

Nicoletti, M. *et al.*, *Phytochemistry*, 1998, **47**, 149-151 (*didehydro derivs*)

Aspidophylline A

[958247-22-8]

A-1495



$C_{21}H_{24}N_2O_4$ 368.432


Alkaloid from *Kopsia singapurensis*. Cryst. (CHCl₃). Mp 110-111°. $[\alpha]_D^{25} -86$ (c, 0.09 in CHCl₃). λ_{\max} 206 (log ϵ 4.52); 239 (log ϵ 4.04); 293 (log ϵ 3.58) (EtOH).

Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1783-1789 (*isol*, *pmr*, *cmr*, *ms*)

Aspidophytine

A-1496

3,4-Didehydro-19-hydroxy-16,17-dimethoxy-1-methylaspidospermidin-21-oic acid, γ -lactone, 9CI
[16625-21-1]



$C_{22}H_{26}N_2O_4$ 382.458

Possible biosynthetic precursor of Haplophytine, H-58. Needles (EtOH). Mp 201-203°. $[\alpha]_D^{25} -122$ (c, 0.16 in CHCl₃). λ_{\max} 222 (ϵ 28700); 256 (ϵ 5900); 304 (ϵ 2400) (EtOH).

Yates, P. *et al.*, *J.A.C.S.*, 1973, **95**, 7842-7850 (*isol*)

Sumi, S. *et al.*, *Tetrahedron*, 2003, **59**, 8571-8587 (*synth*, *pmr*, *cmr*)

Mejia-Oneto, J.M. *et al.*, *Org. Lett.*, 2006, **8**, 3275-3278 (*synth*)


Marino, J.P. *et al.*, *Tet. Lett.*, 2006, **47**, 7711-7713 (*synth*)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 2008, **130**, 14942-14943 (*synth*)

Aspidospermatidine

A-1497

14,19-Didehydrocondyfolan, 9CI
[3890-05-9]



$C_{18}H_{22}N_2$ 266.385

Alkaloid from *Aspidosperma quebrachoblanco* (quebracho) (Apocynaceae). Cryst. (Et₂O/hexane). Mp 168-170° Mp 184-186°. λ_{\max} 242 (log ϵ 3.83); 296 (log ϵ 3.48) (EtOH).

Absolute Configuration

N-Ac: N-Acetylaspidospermatidine

$C_{20}H_{24}N_2O$ 308.422

Alkaloid from *Vallesia dichotoma* (Apocynaceae). Noncryst. $[\alpha]_D -28$ (EtOH). CAS No. not found 8-14CI. λ_{\max} 210 ; 253 ; 289 (CHCl₃).

N-Me: N-Methylaspidospermatidine

$C_{19}H_{24}N_2$ 280.412

Alkaloid in *Aspidosperma quebrachoblanco* (quebracho) (Apocynaceae). Amorph. CAS No. not found 8-14CI.

11-Hydroxy, N-Ac: N-Acetyl-11-hydroxyaspidospermatidine

$C_{20}H_{24}N_2O_2$ 324.422

Minor alkaloid from *Aspidosperma compactinervium* (Apocynaceae). Noncryst. CAS No. not found 8-14CI. λ_{\max} 218 (log ϵ 4.23); 252 (log ϵ 3.89); 294 (log ϵ 3.65); 300 (log ϵ 3.65) (EtOH).

12-Hydroxy, N-Ac: Limatinine

[11035-45-3]

[19046-21-0]

$C_{20}H_{24}N_2O_2$ 324.422

Alkaloid from *Aspidosperma tomentosum*, *Aspidosperma limae* and *Aspidosperma subincanum* (Apocynaceae). Prisms (Et₂O/pentane). Mp 162-163° (158-160°). $[\alpha]_D^{22} +170$ (c, 0.24 in EtOH). $[\alpha]_D^{25} +162$ (c, 0.83 in CHCl₃). 19-Config. uncertain. λ_{\max} 220 (log ϵ 4.36); 259 (log ϵ 3.85); 290 (log ϵ 3.49) (EtOH).

12-Hydroxy, N-Ac, picrate:

Cryst. (Me₂CO/MeOH or Me₂CO aq.). Mp 250° dec.

12-Hydroxy, N-propanoyl: Limatine

$C_{21}H_{26}N_2O_2$ 338.449

Alkaloid from the stem bark of *Aspidosperma limae* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 175-176°. $[\alpha]_D^{20} +166$ (c, 0.85 in CHCl₃). 19-Config. uncertain. CAS No. not found 8-14CI. λ_{\max} 220 (log ϵ 4.66); 259 (log ϵ 4.51); 290 (log ϵ 3.83) (EtOH).

12-Hydroxy, N-propanoyl, picrate:

Cryst. (Me₂CO/EtOAc or Me₂CO aq.). Mp 265° dec.

11-Methoxy, 12-hydroxy: 11-Methoxylimatine

[11044-89-6]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from the stem bark of *Aspidosperma limae* (Apocynaceae). Prisms (MeOH/Et₂O/pentane). Mp 75-78°. $[\alpha]_D^{23} +168$ (c, 0.21 in CHCl₃). 19-Config. uncertain. λ_{\max} 227 (log ϵ 4.39); 262 (log ϵ 3.91) (EtOH).

11-Methoxy, 12-hydroxy, N-Ac: 11-Methoxylimatine

[11035-48-6]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from *Aspidosperma limae* (Apocynaceae). Prisms (Et₂O/pentane or Me₂CO/hexane). Mp 139-140°. $[\alpha]_D^{23} +181$ (c, 0.17 in CHCl₃). 19-Config. uncertain. λ_{\max} 229 (log ϵ 4.38); 261 (log ϵ 3.77) (EtOH).

11-Methoxy, 12-hydroxy, N-Ac, picrate:

Cryst. (Me₂CO/EtOH). Mp 238-244° dec.

11-Methoxy, 12-hydroxy, N-propanoyl, picrate:

Cryst. (Me₂CO/EtOH). Dec. >250° without melting.

12-Methoxy, N-Ac: Aspidospermatine
[5794-14-9]

C₂₁H₂₆N₂O₂ 338.449
Alkaloid from *Aspidosperma quebracho-blanco* (quebracho) and *Vallesia glabra*. Mp 162° (157-159°). [α]_D -73 (EtOH). 19-Config. uncertain. λ_{max} 219 (log ε 4.54); 255 (log ε 4.1); 290 (log ε 3.62) (EtOH).

12-Methoxy, 19,20-dihydro, N-Ac: Dihydroaspidospermatine

C₂₁H₂₈N₂O₂ 340.464
Alkaloid from *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Noncryst. CAS No. not found 8-14CI.

Biemann, K. et al., *J.A.C.S.*, 1963, **85**, 631-638 (*Aspidospermatidine, N-Methylaspidospermatidine*).

Aspidospermatine, Dihydroaspidospermatine
Walsler, A. et al., *Helv. Chim. Acta*, 1965, **48**, 391-404 (*N-Acetylaspidospermatidine*)

Pinar, M. et al., *Helv. Chim. Acta*, 1965, **48**, 822-825; 1967, **50**, 89-93 (*Aspidosperma limae* alkaloids)

Gilbert, B. et al., *Tetrahedron*, 1965, **21**, 1141-1166 (*N-Acetyl-11-hydroxyaspidospermatidine*)


Klyne, W. et al., *Helv. Chim. Acta*, 1966, **49**, 833-841 (*Limatine, isol, ord, abs config*)

Arndt, R.R. et al., *Phytochemistry*, 1967, **6**, 1653-1658 (*Limatine, abs config*)

Zeches, M. et al., *Planta Med.*, 1995, **61**, 89-91 (*Aspidospermatine*)

Aspidospermidine

A-1498



(+)-form

C₁₉H₂₆N₂ 282.428

(+)-form [2912-09-6]

Alkaloid from *Amsonia tabernaemontana*, *Gonioma kamassi*, *Aspidosperma quebracho-blanco* (quebracho) and several other *Aspidosperma* spp. (Apocynaceae). Mp 119.5-121°. [α]_D²³ +21 (EtOH).

N¹-Ac: N-Acetylaspidospermidine. Demethoxyaspidospermine
[2111-81-1]

C₂₁H₂₈N₂O 324.465
Alkaloid from *Aspidosperma exalatum*, *Aspidosperma discolor*, *Geissospermum argenteum* and *Vallesia dichotoma* (Apocynaceae). Cryst. (hexane or Et₂O/hexane). Mp 114° (103-105°). [α]_D²⁰ -15 (c, 1 in CHCl₃).

N¹-Ac: perchlorate:

Cryst. (MeOH/Et₂O). Mp 164-170° dec.

N-Propanoyl: Demethoxyपालosine
[17472-57-0]

C₂₂H₃₀N₂O 338.492
Alkaloid from seeds of *Aspidosperma exalatum*, bark of *Aspidosperma rhombosignatum*, *Aspidosperma disco-*

lor and *Aspidosperma limae* (Apocynaceae). Cryst. (pentane). Mp 117-120°. [α]_D -20 (CHCl₃).

N-(β-threo-Pentopyranos-4-uloside): Aspidospermiöse
[183803-80-7]

C₂₄H₃₂N₂O₅ 428.527
Alkaloid from leaves of *Rhazya stricta*. Stereochem. of sugar residue not totally determined. λ_{max} 207 (log ε 4.12); 235 (log ε 3.18); 250 (log ε 3.14); 300 (log ε 2.3) (MeOH).

N¹-(β-D-xylo-Hexos-4-uloside): Aspidospermiöse

[110325-67-2]
C₂₅H₃₄N₂O₅ 442.554
Alkaloid from leaves of *Rhazya stricta* (Apocynaceae). [α]_D²⁴ +203 (MeOH). The first alkaloidal glycoside with an oxidised sugar unit.

N¹-Me: α-Methylaspidospermidine

[2671-47-8]
C₂₀H₂₈N₂ 296.455
Alkaloid from *Vinca minor* (Apocynaceae). [α]_D²⁴ +24.4 (c, 1.25 in CHCl₃).

14,15-Didehydro, N¹-Me: 14,15-Dehydro-N-methylaspidospermidine. 6,7-Didehydro-1-methylaspidospermidine, 9CI

C₂₀H₂₆N₂ 294.439
Alkaloid from *Vallesia herbacea* (Apocynaceae). Cryst. (EtOH). Mp 110-112°. [α]_D -40 (c, 0.3 in CHCl₃).

21-Hydroxy: Fendlispermine

[52309-71-4]
C₁₉H₂₆N₂O 298.427
Alkaloid from *Aspidosperma fendleri* (Apocynaceae).

14β,15α-Dihydroxy, N¹-Me: 14,15-Dihydroxy-N-methylaspidospermidine

C₂₀H₂₈N₂O₂ 328.453
Alkaloid from *Tabernaemontana bovina*. Cryst. (Me₂CO). Mp 200-203°. [α]_D³⁰ +15.3 (c, 0.06 in MeOH).

3-Oxo: Melocelinine

[52628-61-2]
C₁₉H₂₄N₂O 296.411
Alkaloid from *Melodinus celastroides* (Apocynaceae).

5-Oxo: Meloceline

[52617-21-7]
C₁₉H₂₄N₂O 296.411
Alkaloid from *Melodinus celastroides* (Apocynaceae).

10-Methoxy: 10-Methoxyaspidospermidine

C₂₀H₂₈N₂O 312.454
Alkaloid from stem bark of *Aspidosperma pyriformis*. [α]_D +2.4 (c, 0.5 in CHCl₃). Largely racemic.

20,21-Diepimer: 20,21-Epiaspidospermidine. 5,19-Epiaspidospermidine

[52152-79-1]
C₁₉H₂₆N₂ 282.428
Alkaloid from *Melodinus celastroides* (Apocynaceae).

20,21-Diepimer, N^b-oxide: 20,21-Epiaspidospermidine N-oxide. N^b-Oxy-20,21-epiaspidospermidine

[52077-40-4]
C₁₉H₂₆N₂O 298.427
Alkaloid from *Melodinus celastroides* (Apocynaceae).

(-)-form

14,15-Didehydro, N¹-Me: ent-14,15-Didehydro-N¹-methylaspidospermidine

[16718-72-2]
C₂₀H₂₆N₂ 294.439
Alkaloid from the roots of *Vinca sardoa* (Apocynaceae). Cryst. (petrol). Mp 118-120°. Double bond omitted from struct. diag. in ref.

(±)-form [7689-02-3]

Mp 108-110°.

N¹-Ac: [56197-28-5]

Resin.

Ferreira, J.M. et al., *Experientia*, 1963, **19**, 585 (*N-Acetylaspidospermidine, Demethoxyपालosine*)

Biemann, K. et al., *J.A.C.S.*, 1963, **85**, 631

(ms)

Smith, G.F. et al., *J.C.S.*, 1963, 4002 (*synth, uv*)

Walsler, A. et al., *Helv. Chim. Acta*, 1965, **48**,

391 (*N-Acetylaspidospermidine*)

Gilbert, B. et al., *Tetrahedron*, 1965, **21**, 1141

(*N-Acetylaspidospermidine, isol*)

Mokry, J. et al., *Coll. Czech. Chem. Comm.*,

1967, **32**, 2523 (*N-Methylaspidospermidine*)

Dastoor, N.J. et al., *Helv. Chim. Acta*, 1967,

50, 213 (*N-Acetylaspidospermidine, isol*)

Klyne, W. et al., *Helv. Chim. Acta*, 1968, **51**,

1169 (*ord*)

Rabaron, A. et al., *Plant. Med. Phytother.*,

1973, **7**, 319 (*Oxoaspidospermidine, epimers*)

Medina, J.D. et al., *Rev. Latinoam. Quim.*,

1973, **4**, 73; *C.A.*, **80**, 27419g (*Fendlispermine*)

Larozze, J.-Y. et al., *Tet. Lett.*, 1974, 491

(*synth*)

Babaev, N.A. et al., *Khim. Prir. Soedin.*, 1975,

11, 267; *Chem. Nat. Compd. (Engl. Transl.)*,

1975, **11**, 283 (*14,15-Didehydro-N-methylaspidospermidine*)

Seki, K. et al., *Tet. Lett.*, 1975, 723 (*synth*)

Paccioni, J.P. et al., *Phytochemistry*, 1978, **17**,

2146

Gallagher, T. et al., *J.A.C.S.*, 1983, **105**, 4750

(*synth*)

Atta-ur-Rahman, et al., *J.C.S. Perkin 1*, 1987,

1701 (*Aspidospermiöse*)

Mandal, S.B. et al., *J.O.C.*, 1988, **53**, 4236

(*synth*)

Crippa, S. et al., *Heterocycles*, 1990, **31**, 1663

(*ent-14,15-Didehydro-N-methylaspidospermidine*)

Node, M. et al., *J.O.C.*, 1990, **55**, 517 (*synth*)

Le Ménez, P. et al., *J.O.C.*, 1991, **56**, 2915

(*synth*)

Desmaële, D. et al., *J.O.C.*, 1994, **59**, 2292

(*synth*)

Wenkert, E. et al., *J.O.C.*, 1994, **59**, 7677

(*synth*)

Habib-ur-Rehman, et al., *Fitoterapia*, 1996,

67, 145 (*Aspidospermiöse*)

Forns, P. et al., *J.O.C.*, 1996, **61**, 7882 (*synth*)

Mitaine, A.C. et al., *Planta Med.*, 1996, **62**,

458 (*10-Methoxyaspidospermidine*)

Schultz, A.G. et al., *J.O.C.*, 1997, **62**, 6855-

6861 (*synth*)

Lien, T.P. et al., *Phytochemistry*, 1998, **49**,

1457-1461 (*14,15-Dihydroxy-N-methylaspidospermidine*)

Callaghan, O. et al., *J.C.S. Perkin 1*, 1999, 995-

1001 (*synth*)

Callaghan, O. et al., *Tet. Lett.*, 1999, **40**, 161-

164; 2225 (*synth, bibl*)

Toczko, M.A. et al., *J.O.C.*, 2000, **65**, 2642-

2645 (*synth*)

Kozmin, S.A. et al., *J.A.C.S.*, 2002, **124**, 4628-

4641 (*synth*)

Marino, J.P. et al., *J.A.C.S.*, 2002, **124**, 13398-

13399 (*synth*)

Banwell, M.G. et al., *Aust. J. Chem.*, 2005, **58**,


722-737 (*synth*)

Iyengar, R. *et al.*, *J.O.C.*, 2005, **70**, 10645-10652 (*synth*)

Callier-Dublanchet, A.-C. *et al.*, *Tetrahedron*, 2008, **64**, 4803-4816 (*synth*)

Aspidospermine A-1499

1-Acetyl-17-methoxyaspidospermidine, 9CI
[466-49-9]



(-)-form

$C_{22}H_{30}N_2O_2$ 354.491

Alkaloid from *Aspidosperma quebrachoblanco* (quebracho) and several other *Aspidosperma* spp., *Vallesia glabra*, *Vallesia dichotoma* and *Strychnos angolensis*. Diuretic, respiratory stimulant. λ_{max} 218 (ε 37500); 255 (ε 11000); 285 (ε 3400) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 40 mg/kg.

N-De-Ac: **Deacetylaspido-spermine**. De-formylvallesine. 17-Methoxyaspidospermidine
[2447-50-9]

$C_{20}H_{28}N_2O$ 312.454

Alkaloid from *Strepeliopsis strempelioides* and *Vallesia antillana*. Cryst. (pentane). Mp 107-109°. $[\alpha]_D^{20}$ +8 (MeOH).

N-De-Ac, hydroiodide (1:2):

Prisms. Mp 243° (after sintering from 235°).

N-De-Ac, N-formyl: **Vallesine**

[466-46-6]

$C_{21}H_{28}N_2O_2$ 340.464

Alkaloid from *Strepeliopsis strempelioides*, *Vallesia glabra* and *Vallesia dichotoma* (Apocynaceae). Needles (Me₂CO/Et₂O). Mp 154-155°. $[\alpha]_D$ -92 (c, 0.12 in CHCl₃).

N-De-Ac, N-propanoyl: **Palosine**

[466-48-8]

$C_{23}H_{32}N_2O_2$ 368.518

Alkaloid from *Aspidosperma polyneuron* and stem bark of *Aspidosperma pyriformis* (Apocynaceae). Needles (Et₂O/petrol). Mp 149-152°. $[\alpha]_D^{24}$ -85.9 (c, 0.734 in CHCl₃).

O-De-Me: **Demethylaspido-spermine**

[2122-21-6]

$C_{21}H_{28}N_2O_2$ 340.464

Alkaloid from *Aspidosperma discolor*, *Aspidosperma neblinae*, *Aspidosperma eburneum*, *Geissospermum argenteum*, *Strepeliopsis strempelioides* and *Vallesia antillana* (Apocynaceae). Needles. Mp 112-114°. $[\alpha]_D^{20}$ +114 (CHCl₃).

O-De-Me, perchlorate:

Cryst. + 1 MeOH (MeOH). Mp 170° dec. $[\alpha]_D^{25}$ +94 (c, 0.98 in MeOH).

O-De-Me, N-de-Ac: **Aspidosine**

[466-47-7]

$C_{19}H_{26}N_2O$ 298.427

Alkaloid from the leaves of *Strepeliopsis strempelioides* (Apocynaceae).

Mp 255.5-257.5° (250-253°). $[\alpha]_D^{20}$ -12 (EtOH). λ_{max} 246 (log ε 3.91); 289 (log ε 3.42) (MeOH).

O-De-Me, N-de-Ac, N-propanoyl: **O-De-methylpalosine**

[16049-40-4]

$C_{22}H_{30}N_2O_2$ 354.491

Alkaloid from *Tabernaemontana amygdalaeifolia* roots (Apocynaceae). Mp 169°. $[\alpha]_D^{20}$ +118 (c, 0.36 in CHCl₃).

Ewins, A.J. *et al.*, *J.C.S.*, 1914, **105**, 2738

(*synth*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1948, **31**, 446 (*synth, uv*)

Witkop, B. *et al.*, *J.A.C.S.*, 1954, **76**, 5603 (*synth, uv*)

Chalmers, J.R. *et al.*, *J.C.S.*, 1957, 1115 (*uv*)

Conroy, H. *et al.*, *J.A.C.S.*, 1958, **80**, 5178 (*ir, pmr*)

Schmutz, J. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 874 (*Palosine*)

Taylor, W.I. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 2750 (*Palosine*)

Holker, J.S.E. *et al.*, *J.O.C.*, 1959, **24**, 314 (*Vallesine*)

Gilbert, B. *et al.*, *Chem. Ind. (London)*, 1962, 1949 (*Demethoxypalosine*)

Ferreira, J.M. *et al.*, *Experientia*, 1963, **19**, 585 (*uv, ir, pmr, ms, struct, deriv*)

Biemann, K. *et al.*, *J.A.C.S.*, 1963, **85**, 631 (*ms*)

Walsler, A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 391 (*deriv*)

Klyne, W. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 443; 1966, **49**, 833 (*ord, uv, abs config*)

Ban, Y. *et al.*, *Tet. Lett.*, 1965, 2261 (*synth*)

Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141 (*isol*)

Achenbach, H. *et al.*, *Tet. Lett.*, 1966, 5027 (*Demethylpalosine*)

Craven, B.M. *et al.*, *Experientia*, 1968, **24**, 770 (*cryst struct, abs config*)

Sakabe, N. *et al.*, *Tet. Lett.*, 1969, 2527 (*cryst struct*)

Stevens, R.V. *et al.*, *Chem. Comm.*, 1971, 857 (*synth*)

Paccioni, J.P. *et al.*, *Phytochemistry*, 1978, **17**, 2146 (*isol*)

Laguna, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1980, **45**, 1419-1423 (*Vallesine, Aspidosine, Demethylaspido-spermine*)

Meyers, A.I. *et al.*, *J.O.C.*, 1989, **54**, 4673 (*synth*)


Campbell, W.E. *et al.*, *Spectrosc. Lett.*, 1993, **26**, 707 (*pmr, cmr*)

Mitaine, A.C. *et al.*, *Planta Med.*, 1996, **62**, 458 (*Palosine, isol, pmr, cmr*)

Fukuda, Y.-I. *et al.*, *Org. Lett.*, 2003, **5**, 749-751 (*synth*)

Aspochalamin A

A-1500



$C_{44}H_{55}N_5O_6$ 749.948

Prod. by *Aspergillus niveus* LU 9575.

17-Deoxy: **Aspochalamin C**

$C_{44}H_{55}N_5O_5$ 733.949

Prod. by *Aspergillus niveus* LU 9575.

17,18-Dideoxy: **Aspochalamin D**

$C_{44}H_{55}N_5O_4$ 717.95

Prod. by *Aspergillus niveus* LU 9575.

Stereoisomer: **Aspochalamin B**


$C_{44}H_{55}N_5O_6$ 749.948

Prod. by *Aspergillus niveus* LU 9575.

Gebhardt, K. *et al.*, *J. Antibiot.*, 2004, **57**, 707-714; 715-720 (*isol, pmr, cmr*)

Aspochalasin I

A-1501



$C_{24}H_{35}NO_5$ 417.544

Prod. by *Aspergillus flavipes*. Solid. Mp 136-138°. $[\alpha]_D^{27}$ -166.6 (c, 0.2 in CHCl₃).

17-Deoxy: **Aspochalasin J**

$C_{24}H_{35}NO_4$ 401.545


Prod. by *Aspergillus flavipes*. Solid.

Zhou, G.-X. *et al.*, *J. Nat. Prod.*, 2004, **67**, 328-332 (*isol, pmr, cmr*)

Aspochalasin C

A-1502

[72401-79-7]



$C_{24}H_{35}NO_4$ 401.545

Metab. of *Aspergillus microcysticus*. Metab. inactive. Amorph. powder. $[\alpha]_D^{25}$ -86 (c, 1.37 in CHCl₃). λ_{max} 244 (ε 7320) (EtOH) (Derep). λ_{max} 240 (EtOH) (Berdy).

18-Ketone: **Aspochalasin B**

[72363-47-4]

$C_{24}H_{33}NO_4$ 399.529

From *Aspergillus microcysticus*. Inactive. Yellow amorph. powder. $[\alpha]_D^{25}$ -118 (c, 1.37 in CHCl₃). λ_{max} 225 (ε 9770) (EtOH) (Derep). λ_{max} 225 (ε 9900) (EtOH) (Berdy).

17,18-Diketone: **Aspochalasin A**. *Asposterol*

[72363-48-5]

$C_{24}H_{31}NO_4$ 397.513

From *Aspergillus microcysticus*. Antibiotic. Light yellow amorph. powder. $[\alpha]_D^{25}$ -20 (c, 0.27 in CHCl_3). λ_{max} 265 (ε 407); 410 (ε 12) (EtOH) (Derep).

17-Deoxy: 17-Deoxyaspochalasin C.

TMC 169. Antibiotic TMC 169

$\text{C}_{24}\text{H}_{35}\text{NO}_3$ 385.545

Prod. by *Aspergillus flavipes*. Powder. Mp 103-105° Mp 112-114°. $[\alpha]_D^{23}$ -50 (c, 0.31 in MeOH). C-18 config. not determined.

17,18-Dideoxy: Aspochalasin Z

$\text{C}_{24}\text{H}_{35}\text{NO}_2$ 369.546

Prod. by *Aspergillus niveus* LU 9575.

17,18-Dideoxy, 17,18-epoxide: Aspochalasin G

$\text{C}_{24}\text{H}_{33}\text{NO}_3$ 383.53

Prod. by *Aspergillus* sp. FO-4282. Inhibits lipid droplet formation. Pale yellow solid. Mp 95°. $[\alpha]_D^{25}$ -77.9 (c, 1 in EtOH). Stereochem. not determined. λ_{max} 205 (ε 9800); 240 (sh); 280 (sh) (MeOH).

19,20-Dihydro, 19ξ-hydroxy: Aspochalasin E

[149598-68-5]

$\text{C}_{24}\text{H}_{37}\text{NO}_5$ 419.56

Prod. by *Aspergillus flavipes* and an unidentified fungal sp. FA-2277. Powder. Sol. MeOH, CHCl_3 . Mp 129-131°. $[\alpha]_D^{24}$ -52 (c, 0.5 in CHCl_3). λ_{max} 265 (ε 407); 410 (ε 12) (EtOH) (Derep).

19,20-Dihydro, 19ξ-hydroxy, 18-Me ether: Aspochalasin K

$\text{C}_{25}\text{H}_{39}\text{NO}_5$ 433.587

Prod. by *Aspergillus flavipes*. Amorph. solid. Mp 185-186°. $[\alpha]_D^{25}$ -81.4 (c, 1 in CHCl_3). Stereochem. not confirmed.

17-Epimer: Aspochalasin D

[71968-02-0]

$\text{C}_{24}\text{H}_{35}\text{NO}_4$ 401.545

Prod. by *Aspergillus flavipes* and *Aspergillus microcysticus*. Shows selective cytotoxicity. Needles (EtOAc). Mp 148° dec. $[\alpha]_D^{25}$ -81 (c, 1.43 in EtOH). Unknown stereochem. λ_{max} 244 (ε 7320) (EtOH) (Derep). λ_{max} 248 (ε 9300) (EtOH) (Berdy).

17-Epimer, 19α, 20β-epoxide: Aspochalasin H

$\text{C}_{24}\text{H}_{35}\text{NO}_5$ 417.544

Prod. by *Aspergillus* sp. AJ117509.

Powder.

Heberle, W. *et al.*, *Arch. Microbiol.*, 1974, **100**, 73 (isol)

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 1501 (struct)

Neupert-Laves, K. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 1426 (cryst struct)

Trost, B.M. *et al.*, *J.A.C.S.*, 1989, **4**, 8284 (synth)

Naruse, N. *et al.*, *J. Antibiot.*, 1993, **46**, 679-681 (Aspochalasin E)

Fang, F. *et al.*, *J. Antibiot.*, 1997, **50**, 919-925 (Aspochalasin G)

Kohno, J. *et al.*, *J. Antibiot.*, 1999, **52**, 575-577 (TMC 169)

Namatame, I. *et al.*, *J. Antibiot.*, 2000, **53**, 19-25 (Aspochalasin G, activity)

Tomikawa, T. *et al.*, *J. Antibiot.*, 2001, **54**, 379-381; 2002, **55**, 666-668 (Aspochalasins D,H)


Gebhardt, K. *et al.*, *J. Antibiot.*, 2004, **57**, 707-714; 715-720 (Aspochalasin Z)

Zhou, G.-X. *et al.*, *J. Nat. Prod.*, 2004, **67**, 328-332 (Aspochalasin K)

Aspochalasin F

[200631-32-9]

A-1503



$\text{C}_{24}\text{H}_{33}\text{NO}_4$ 399.529


Prod. by *Aspergillus* sp. FO-4282. Inhibits lipid droplet formation. Pale yellow solid. Mp 96-97°. $[\alpha]_D^{25}$ +9.8 (c, 1 in EtOH). Stereochem. not determined. λ_{max} 203 (ε 10200) (MeOH).

Fang, F. *et al.*, *J. Antibiot.*, 1997, **50**, 919-925 (isol, uv, ir, pmr, cmr)

Namatame, I. *et al.*, *J. Antibiot.*, 2000, **53**, 19-25 (activity)

Aspoquinolone A

A-1504



$\text{C}_{27}\text{H}_{31}\text{NO}_6$ 465.545

Prod. by *Aspergillus nidulans* HKI0410. Amorph. solid. λ_{max} 201 (ε 34200); 217 (ε 27900); 279 (ε 15100); 323 (ε 14000) (MeOH).

20,21-Diepimer: Aspoquinolone B

$\text{C}_{27}\text{H}_{31}\text{NO}_6$ 465.545


Prod. by *Aspergillus nidulans*

HKI0410. Amorph. solid. λ_{max} 201 (ε 33000); 217 (ε 27100); 279 (ε 14400); 323 (ε 13200) (MeOH).

Scherlach, K. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3517-3520 (isol, pmr, cmr, ms)

Aspoquinolone C

A-1505



$\text{C}_{27}\text{H}_{33}\text{NO}_7$ 483.56

Closely related to Penigequinolone A, P-191. Prod. by *Aspergillus nidulans* HKI0410. Amorph. solid. Isol. as an inseparable mixt. with Aspoquinolone D

to which data refers. λ_{max} 201 (ε 30600); 279 (ε 11800); 323 (ε 10600) (MeOH).

Stereoisomer: Aspoquinolone D

$\text{C}_{27}\text{H}_{33}\text{NO}_7$ 483.56

Prod. by *Aspergillus nidulans*


HKI0410. Stereoisomeric on pyran ring.

Scherlach, K. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3517-3520 (isol, pmr, cmr, ms)

Aspyridone A

A-1506

[935863-26-6]



Relative Configuration

$\text{C}_{19}\text{H}_{23}\text{NO}_4$ 329.395

Prod. by *Aspergillus nidulans*. Amorph. powder. λ_{max} 203 (log ε 4.39); 247 (log ε 4.37); 344 (log ε 3.74) (MeOH).

3''-Hydroxy: Aspyridone B

[935863-27-7]

$\text{C}_{19}\text{H}_{23}\text{NO}_5$ 345.394


Prod. by *Aspergillus nidulans*. Amorph. powder. λ_{max} 204 (log ε 4.4); 214 (log ε 4.41); 251 (log ε 4.21); 344 (log ε 3.61) (MeOH).

Bergmann, S. *et al.*, *Nat. Chem. Biol.*, 2007, **3**, 213-217 (isol, biosynth, pmr, cmr, ms)

Assamicadine

A-1507

[126260-96-6]



$\text{C}_{16}\text{H}_{23}\text{NO}_5$ 309.361


Rel. config. only of lactone portion detd. Minor alkaloid from seeds of *Crotalaria assamica* (Fabaceae). Viscous oil. $[\alpha]_D^{19}$ -9.4 (c, 0.4 in MeOH).

Cheng, D. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1153 (isol, ir, pmr, cmr, ms, cd, struct)

Assoanine

A-1508

4,5-Dihydro-9,10-dimethoxy-7H-pyrrolo[3,2,1-de]phenanthridine, 9CI. Anhydromethylpseudolycorine
[107208-75-3]



$\text{C}_{17}\text{H}_{17}\text{NO}_2$ 267.327

Alkaloid from the bulbs of *Narcissus pseudonarcissus*, the aerial parts of *Narcissus assoanus* collected at flowering and from the whole plant of *Narcissus jace-*

tanus (Amaryllidaceae). Cryst. (EtOH). Air oxidn. occurs during determination of the Mp, and the dec. point finally observed is that of the 7-oxo compd.

N-Me: N-Methylasosanin

[132368-07-1]
C₁₈H₂₀NO₂⁺ 282.361

Alkaloid from *Lapiedra martinezii* (Amaryllidaceae). CAS no. refers to the chloride.

O¹⁰-De-Me: Anhydroseudolycorine

C₁₆H₁₅NO₂ 253.3

Alkaloid from *Hymenocallis guianensis*, *Hymenocallis lobata* and *Hymenocallis tubiflora*.

7(N)-Dehydro: see Vasconine, V-35

4,5-Didehydro, O¹⁰-de-Me: 4,5-Dehydroanhydroseudolycorine

C₁₆H₁₃NO₂ 251.284

Alkaloid from *Hymenocallis tubiflora* and *Hymenocallis venezuelensis*.

7-Oxo: Oxoasosanine. 7-Oxoanhydro-methylseudolycorine

[65367-74-0]

C₁₇H₁₅NO₃ 281.31

Alkaloid from the aerial parts of *Narcissus assoanus* collected at flowering and from the whole plant of *Narcissus jacetanus* (Amaryllidaceae). Prisms (CHCl₃/EtOH). Mp 272-274° (247-250°).

7-Oxo, N-oxide: Oxoasosanine N-oxide

[129225-26-9]

C₁₇H₁₅NO₄ 297.31

Alkaloid from the bulbs of *Narcissus bicolor* (Amaryllidaceae). Cryst. + 2½H₂O. Mp 196-198°.

8-Methoxy, 7-oxo: 8-Methoxyoxoasosanine

C₁₈H₁₇NO₄ 311.337

Alkaloid from *Eucharis amazonica*. Amorph. solid. Mp 245-247°.

Fales, H.M. *et al.*, *J.A.C.S.*, 1956, **78**, 4145 (isol, uv, struct, synth, deriv)

Llabrés, J.M. *et al.*, *Phytochemistry*, 1986, **25**, 2637 (isol, uv, ir, pmr, cmr, ms, deriv)

Bastida, J. *et al.*, *Planta Med.*, 1988, **54**, 362 (isol, Assoanine, Oxoasosanine)

Suau, R. *et al.*, *An. Quim.*, 1990, **86**, 672; *CA*, **114**, 98281n (*N-Methylasosanin*)

Viladomat, F. *et al.*, *Phytochemistry*, 1990, **29**, 1307 (*Oxoasosanine N-oxide*)

Meyers, A.I. *et al.*, *Tet. Lett.*, 1993, **34**, 6185 (synth, Oxoasosanine)

Iwao, M. *et al.*, *Heterocycles*, 1994, **38**, 1717

Parnes, J.S. *et al.*, *J.O.C.*, 1994, **59**, 3497 (synth, Assoanine, Oxoasosanine)

Banwell, M.G. *et al.*, *Chem. Comm.*, 1995, 2551 (synth, Oxoasosanine)

Hutchings, R.H. *et al.*, *J.O.C.*, 1996, **61**, 1004 (synth, Oxoasosanine)

Rosa, A.M. *et al.*, *Tetrahedron*, 1997, **53**, 299 (synth, Assoanine, Oxoasosanine)

Padwa, A. *et al.*, *J.O.C.*, 1998, **63**, 3986-3997 (*Oxoasosanine*, synth)

Cabezas, F. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 315-317 (*8-Methoxyoxoasosanine*)


Harayama, T. *et al.*, *Heterocycles*, 2003, **60**, 2429-2434 (synth)

Rivero, N. *et al.*, *Pharm. Biol.*, 2004, **42**, 280-285 (*Anhydroseudolycorine*, *4,5-Dehydroanhydroseudolycorine*)

Harayama, T. *et al.*, *Tetrahedron*, 2004, **60**, 1611-1616 (synth)

Ganton, M.D. *et al.*, *Org. Lett.*, 2005, **7**, 4777-4779 (synth)

Astechrome
[75310-10-0]



C₆₀H₆₆FeN₉O₉ 1113.083


Isol. from *Aspergillus terreus* grown in the presence of Fe ions. Shows weak antibiotic activity. Dark-red needles (CHCl₃/hexane). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 188-189° dec. λ_{max} 223 (ε 151360); 282 (ε 24550); 292 (ε 21900); 347 (ε 26300); 450 (ε 4790) (MeOH) (Berdy).

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1510 (isol, uv, ir, ms, struct)

Ohta, A. *et al.*, *Heterocycles*, 1992, **34**, 111 (synth)

Jing, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 277 (synth)

Asteranthine A-1510
2,9-Dihydro-2,2-dimethyl-7-(3-methylbutyl)pyrano[3,2-g]indole



C₁₈H₂₃NO 269.386

Parent not known.

2',3'-Didehydro, 2',3'-epoxide: 2',3'-Epoxyasteranthine

[187523-31-5]

C₁₈H₂₁NO₂ 283.369

Alkaloid from *Asteranthe asterias*. Yellowish oil. [α]_D²¹ +4 (c, 1 in CHCl₃). Mixt. of stereoisomers.

2',3'-Dihydroxy: 2',3'-Dihydroxyasteranthine

[187523-32-6]

C₁₈H₂₃NO₃ 301.385


Alkaloid from *Asteranthe asterias*. Cryst. Mp 138-140°. [α]_D²¹ -8 (c, 0.7 in CHCl₃). Mixt. of stereoisomers.

Nkunya, M.H.H. *et al.*, *Nat. Prod. Lett.*, 1996, **9**, 71-78

Asterin A†

[151964-51-1]

A-1511



C₂₅H₃₃N₅O₈ 531.564

Oligopeptide. Constit. of the roots of *Aster tataricus*.

Me ester: Asterin B

[151964-52-2]

C₂₆H₃₅N₅O₈ 545.591

Constit. of the roots of *Aster tataricus*.


Chen, D. *et al.*, *Chin. Chem. Lett.*, 1993, **4**, 605 (isol, struct)

Asterinin A

A-1512

2,3,4,5-Tetrahydroprolylallothreonylseryl-3-phenyl-β-alanyl-2-aminobutanoic acid, 9CI

[157536-40-8]



C₂₅H₃₃N₅O₈ 531.564

Constit. of the roots of *Aster tataricus*.

Amorph. powder (MeOH aq.). Mp 272-274°. [α]_D +38.5 (c, 0.33 in Py).

Me ester: Asterinin B

[157536-41-9]

C₂₆H₃₅N₅O₈ 545.591

Constit. of the roots of *Aster tataricus*.


Amorph. powder (MeOH aq.). Mp 235-237°. [α]_D +3.1 (c, 0.57 in Py).

Cheng, D. *et al.*, *Phytochemistry*, 1994, **36**, 945 (isol, pmr, cmr, uv, ir, ms)

Asterinin C

A-1513

[157605-23-7]



C₂₆H₃₅N₅O₈ 545.591


Constit. of the roots of *Aster tataricus*

(Asteraceae). Amorph. powder. Mp 250-252°. [α]_D -4.9 (c, 0.33 in Py). λ_{max} 209 (ε 13800); 267 (ε 18200) (MeOH).

Cheng, D. *et al.*, *Phytochemistry*, 1994, **36**, 945 (isol, uv, pmr, cmr, ms)

Asterionellin A

A-1514



R = CH₃

C₁₀H₁₄N₂O₅ 242.231

Isol. from an *Asterionella* sp. Not indexed in CAS.

Shimizu, Y. *et al.*, *Chem. Rev.*, 1993, **93**, 1685-1698

Asterionellin B A-1515

As Asterionellin A, A-1514 with R = H

$C_9H_{12}N_2O_5$ 228.204

Isol. from the microalga *Asterionella* sp. Antibiotic.

6-Hydroxy: **Asterionellin C**


$C_9H_{12}N_2O_6$ 244.204

Isol. from *Asterionella* sp.

Shimizu, Y. *et al.*, *Chem. Rev.*, 1993, **93**, 1685-1698

Asteropterin A-1516

[1041186-56-4]




$C_{12}H_{13}N_7O_2$ 287.28

Alkaloid from *Asteropus simplex*. Cathepsin B inhibitor. Yellowish powder. λ_{max} 279 (ε 1360); 416 (ε 290) (MeOH).

Murayama, S. *et al.*, *Tet. Lett.*, 2008, **49**, 4186-4188 (*isol*, *pmr*, *cmr*)

Asterredione A-1517




$C_{24}H_{18}N_2O_5$ 414.417

Prod. by *Aspergillus terreus* isol. from *Opuntia versicolor*. Cytotoxic. Yellow powder. $[\alpha]_D^{25}$ -2.3 (c, 0.5 in $CHCl_3$). Dec. at $>112^\circ$. λ_{max} 218 (log ε 5.68); 269 (log ε 5.23); 411 (log ε 4.92) (EtOH).

Wijeratne, E.M.K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1567-1573 (*isol*, *pmr*, *cmr*)

Asterridinone A-1518

[161300-75-0]




$C_{32}H_{28}N_2O_4$ 504.584

Metab. from the mycelium of *Aspergillus terreus*. Orange needles (MeOH). Mp 261-263° dec.

Kaji, A. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1682 (*isol*, *uv*, *ir*, *pmr*, *struct*)

Asterriquinol D A-1519

2,5-Bis(1*H*-indol-3-yl)-3,6-dimethoxy-1,4-benzenediol. Asterriquinone D hydroquinone



$C_{24}H_{20}N_2O_4$ 400.433

Parent compd. (currently unknown as a natural product) of a series of prenylated metabolites related to the asterriquinones; see Asterriquinone, A-1520 and Asterriquinone D, A-1521.

Di-Me ether: **Di-O-methylasterriquinol D**

$C_{26}H_{24}N_2O_4$ 428.487

Prod. by *Aspergillus solerotiorum* and *Aspergillus terreus*. Solid. Mp 187-189°.

2',3'-Dihydro, 2'-oxo, di-Me ether: **2'-Oxoasterriquinol D di-Me ether**

$C_{26}H_{24}N_2O_5$ 444.486

Prod. by *Aspergillus sclerotiorum*. Solid. Mp 145-147° (dec.). $[\alpha]_D$ -13.9 (c, 0.001 in MeOH). λ_{max} 232 (ε 5500); 276 (ε 2600) (MeOH).

1'-(1,1-Dimethyl-2-propenyl), di-Me ether: **1-(1,1-Dimethyl-2-propenyl)-3-[4-(1*H*-indol-3-yl)-2,3,5,6-tetramethoxyphenyl]-1*H*-indole, 9CI**

[78279-80-8]

$C_{31}H_{32}N_2O_4$ 496.605

Metab. of *Aspergillus terreus* var. *africanus*. Mp 202-204°.

2'-(1,1-Dimethyl-2-propenyl): **2-[2-(1,1-Dimethyl-2-propenyl)-1*H*-indol-3-yl]-5-(1*H*-indol-3-yl)-3,6-dimethoxy-1,4-benzenediol, 9CI. Asterriquinone C1 hydroquinone**

[78279-79-5]

$C_{29}H_{28}N_2O_4$ 468.551

Metab. of *Aspergillus terreus* var. *africanus*. Mp 206-208°. Oxidised gradually in air, rapidly by aq. $FeCl_3$, to Asterriquinone C1 in A-1521.

1',1''-Bis(1,1-dimethyl-2-propenyl): **2,5-Bis[1-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dimethoxy-1,4-benzenediol, 9CI. Asterriquinone A1 hydroquinone**

[78279-78-4]

$C_{34}H_{36}N_2O_4$ 536.669

Metab. of *Aspergillus terreus* var. *africanus*. Mp 242-245°. Oxidised gradually in air, rapidly by $FeCl_3$ aq., to

Asterriquinone A1 in A-1520.

1',1''-Bis(1,1-dimethyl-2-propenyl), di-Me ether: **3,3'-(2,3,5,6-Tetramethoxy-1,4-phenylene)bis[1-(1,1-dimethyl-2-propenyl)-1*H*-indole], 9CI**

[78302-49-5]

$C_{36}H_{40}N_2O_4$ 564.723

Metab. of *Aspergillus terreus* var. *africanus*. Mp 158-159°.

7'-(3-Methyl-2-butenyl), di-Me ether: **3-[4-(1*H*-indol-3-yl)-2,3,5,6-tetramethoxyphenyl]-7-(3-methyl-2-butenyl)-1*H*-indole, 9CI**

[78279-81-9]

$C_{31}H_{32}N_2O_4$ 496.605


Metab. of *Aspergillus terreus* var. *africanus*. Mp 226.5-228.5°.

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1005

Whyte, A.C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1006-1009 (2'-Oxoasterriquinol D)

Asterriquinone A-1520

2,5-Bis[1-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dihydroxy-2,5-cyclohexadiene-1,4-dione, 9CI. NSC 289487 [60696-52-8]



$C_{32}H_{30}N_2O_4$ 506.6

Benzoquinone antibiotic. Metab. of *Aspergillus terreus*. Antineoplastic agent, shows anti-HIV activity. Dark purple needles (MeOH). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 218-220° dec. Log P 5.92 (uncertain value) (calc). λ_{max} 298 (ε 32400); 508 (ε 5500) ($CHCl_3$) (Derp). λ_{max} 298 (ε 30000); 508 (ε 6600) ($CHCl_3$) (Berdy).

Mono-Ac: **Asterriquinone monoacetate**

[160786-85-6]

$C_{34}H_{32}N_2O_5$ 548.637

Metab. of *Aspergillus terreus*. Blue-purple needles (hexane/ CH_2Cl_2). Mp 172-174° dec. λ_{max} 225 (ε 46000); 292 (ε 27600); 487 (ε 6400) (EtOH) (Berdy).

Mono-Me ether: **O-Methylasterriquinone.**

Asterriquinone SU5504

$C_{33}H_{32}N_2O_4$ 520.627

Prod. by *Aspergillus candidus*. Purple powder. λ_{max} 225; 292; 489 (no solvent reported).

Di-Me ether: **Asterriquinone A1**

[60924-75-6]

$C_{34}H_{34}N_2O_4$ 534.654

Metab. of *Aspergillus terreus* var. *africanus*. Purple needles (EtOH). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 168°. λ_{max} 295; 299; 490 (EtOH)


(Berdy).

- Kiriyama, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1853 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 961; 991; 1005; 1990, **38**, 2929 (*derivs, biosynth*)
 Shimizu, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1896; 1990, **38**, 2617 (*pharmacol*)
 Ono, K. *et al.*, *Biochem. Biophys. Res. Commun.*, 1991, **174**, 56 (*anti-HIV activity*)
 Kaji, A. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1682 (*monoacetate*)
 Alvi, K.A. *et al.*, *J. Antibiot.*, 1999, **52**, 215-223 (*Asterriquinone SU5504*)

Asterriquinone D

A-1521

[78723-19-0]

 $C_{24}H_{18}N_2O_4$ 398.417

Benzoquinone antibiotic. *Isol.* from *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Purple powder (Me₂CO). Mp 178-180°. λ_{max} 222 (log ϵ 5.69); 282 (log ϵ 5.42); 480 (log ϵ 4.71) (EtOH).

N-(1,1-Dimethyl-2-propenyl): **Asterriquinone B3**

[78708-37-9]

 $C_{29}H_{26}N_2O_4$ 466.535

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Reddish-purple powder (cyclohexane). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 148-150°. λ_{max} 240; 284; 484 (EtOH) (Berdy).

Di-O-de-Me: **Didemethylasterriquinone D**

[78860-40-9]

 $C_{22}H_{14}N_2O_4$ 370.364

Prod. by *Aspergillus terreus* and *Chryso sporium merdarium*.

2'-(1,1-Dimethyl-2-propenyl): **Asterriquinone C1**

[78723-18-9]

 $C_{29}H_{26}N_2O_4$ 466.535

From *Aspergillus terreus* var. *africanus*. No biol. activity. Purple needles (C₆H₆). Mp 162-164° Mp 213-214°. λ_{max} 222 (log ϵ 5.49); 281 (log ϵ 5.12); 367 (log ϵ 3.97); 489 (log ϵ 4.19) (EtOH).

2',N^{1'}-Bis(1,1-dimethyl-2-propenyl): **Asterriquinone A3**

[78723-15-6]

 $C_{34}H_{34}N_2O_4$ 534.654

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Deep purple powder (C₆H₆). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 112-114°. λ_{max} 284; 292; 376; 516 (EtOH) (Berdy).

2',2''-Bis(1,1-dimethyl-2-propenyl): **Asterriquinone B4**

[78723-17-8]

 $C_{34}H_{34}N_2O_4$ 534.654

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Reddish-purple plates (MeOH). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 300°. λ_{max} 284; 292; 385; 502 (EtOH) (Berdy).

2'-(3-Methyl-2-butenyl), di-O-de-Me:

Asterriquinone SU5228

[78860-46-5]

 $C_{27}H_{22}N_2O_4$ 438.482

Prod. by *Aspergillus candidus*. Purple powder. Mp 150-154°. λ_{max} 260; 285; 290; 489 (no solvent reported).

5'-(3-Methyl-2-butenyl), di-O-de-Me:

Semicochliodinol A

[194224-72-1]

 $C_{27}H_{22}N_2O_4$ 438.482

Prod. by *Chryso sporium merdarium*. Mp 195-197°.

6'-(3-Methyl-2-butenyl), di-O-de-Me:

Semicochliodinol B

[194224-73-2]

 $C_{27}H_{22}N_2O_4$ 438.482

Prod. by *Chryso sporium merdarium*. Mp > 240°.

7'-(3-Methyl-2-butenyl): **Asterriquinone C2**

[78708-38-0]

 $C_{29}H_{26}N_2O_4$ 466.535

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Purple needles (MeOH). Mp 202-203°. λ_{max} 285; 290; 485 (EtOH) (Berdy).

7'-(3-Methyl-2-butenyl), 2'-(1,1-dimethyl-2-propenyl): **Asterriquinone B2**

[78708-36-8]

 $C_{34}H_{34}N_2O_4$ 534.654

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Deep purple prisms (C₆H₆). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 199-200°. λ_{max} 283; 292; 372; 518 (EtOH) (Berdy).

7'-(3-Methyl-2-butenyl), 2',2''-bis(1,1-dimethyl-2-propenyl): **Asterriquinone A4**

[78723-16-7]

 $C_{39}H_{42}N_2O_4$ 602.772

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Reddish-purple prisms (C₆H₆). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 252-253°. λ_{max} 284; 291; 390; 497 (EtOH) (Berdy).

7''-(3-Methyl-2-butenyl), 2'-(1,1-dimethyl-2-propenyl): **Asterriquinone B1**

[78708-35-7]

 $C_{34}H_{34}N_2O_4$ 534.654

From *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Deep purple prisms (C₆H₆). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 208-209°. λ_{max} 285; 292; 382; 522 (EtOH) (Berdy).

7''-(3-Methyl-2-butenyl), N^{1'}-(1,1-dimethyl-2-propenyl), di-O-de-Me: **Asterriquinone SU 5501**

[226562-16-9]

 $C_{32}H_{30}N_2O_4$ 506.6

Prod. by *Aspergillus candidus*. Purple powder. λ_{max} 260; 285; 290; 489 (no solvent reported).

2',2''-Bis(3-methyl-2-butenyl), di-O-de-Me: **Neoasterriquinone Asterriquinone CT5**

Asterriquinone E

[78860-48-7]

 $C_{32}H_{30}N_2O_4$ 506.6

Metab. of *Aspergillus terreus*. Inhibits binding of proteins to the EGF receptor tyrosine kinase. Dark purple needles or powder. Mp 300° dec., approx (192°-193° dec.). λ_{max} 295 (log ϵ 4.49); 448 (log ϵ 3.7) (no solvent reported). λ_{max} 260; 279; 290; 492 (MeOH). λ_{max} 295 (ϵ 30900); 448 (ϵ 5110) (MeOH) (Berdy).

2',2''-Bis(3-methyl-2-butenyl), di-O-de-Me, O-Ac: **Asterriquinone SU5500**

[213831-76-6]

 $C_{34}H_{32}N_2O_5$ 548.637

Prod. by *Aspergillus candidus*. Purple powder. λ_{max} 225; 285; 292; 487 (no solvent reported).

2',2''-Bis(3-methyl-2-butenyl), 7'-carboxy, di-O-de-Me: **Asterriquinone SU 5503**

[226562-14-7]

 $C_{33}H_{30}N_2O_6$ 550.61

Prod. by *Aspergillus candidus*. Purple powder. λ_{max} 260; 282; 294; 479 (no solvent reported).

6',6''-Bis(3-methyl-2-butenyl), di-O-de-Me: **Asterriquinone CT3**

[182182-93-0]

 $C_{32}H_{30}N_2O_4$ 506.6

Prod. by *Humicola grisea*. Dark purple powder. Mp ca. 300° dec. λ_{max} 297 (log ϵ 4.45); 452 (log ϵ 3.61) (no solvent reported). λ_{max} 297 (ϵ 28200); 452 (ϵ 4073) (MeOH) (Berdy).

7',7''-Bis(3-methyl-2-butenyl), di-O-de-Me: **Asterriquinone CT4**

[182182-94-1]

 $C_{32}H_{30}N_2O_4$ 506.6

Prod. by *Humicola fuscoatra*. Dark purple powder. Mp 295-300° dec. λ_{max} 303 (log ϵ 4.43); 470 (log ϵ 3.79) (no solvent reported). λ_{max} 303 (ϵ 26900); 470 (ϵ 6166) (MeOH) (Berdy).

7',7''-Bis(3-methyl-2-butenyl), 2'-(1,1-dimethyl-2-propenyl): **Asterriquinone A2**

[78723-14-5]

 $C_{39}H_{42}N_2O_4$ 602.772

Prod. by *Aspergillus terreus* var. *africanus*. Shows weak antitumour activity. Deep purple prisms (C₆H₆). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 189-192°. λ_{max} 283; 292; 374; 505 (EtOH) (Berdy).

5'-(3-Methyl-1,3-butadienyl): **Asterriquinone PM 53**

PM 53

[198992-52-8]

 $C_{29}H_{24}N_2O_4$ 464.52

From *Petromyces muricatus*. Deep blue cryst. (hexane/CH₂Cl₂). Mp 240° dec. λ_{max} 219 (log ϵ 4.42); 274 (log ϵ 4.44); 308 (log ϵ 4.21); 614 (log ϵ 3.14) (dioxane).

6''-(3-Methyl-1,3-butadienyl), 6'-(3-methyl-2-butenyl), di-O-de-Me: **Asterriquinone CT2**

[182182-92-9]

C₃₂H₂₈N₂O₄ 504.584

Prod. by *Botryotrichum* sp. Black powder. Mp ca. 350° dec. λ_{max} 303 (log ε 4.82); 470 (log ε 3.64) (no solvent reported). λ_{max} 303 (ε 66000); 470 (ε 4365) (MeOH) (Berdy).

6',6''-Bis(3-methyl-1,3-butadienyl)(E,E-), di-O-de-Me: **Asterriquinone CT1** [182182-91-8]

C₃₂H₂₆N₂O₄ 502.568

Prod. by *Humicola fuscoatra*. Black powder. Mp 350-355° dec. λ_{max} 305 (log ε 4.68); 470 (log ε 3.79) (no solvent reported). λ_{max} 305 (ε 12022); 470 (ε 6166) (MeOH) (Berdy).

7''-(3-Methyl-2-butenyl), 2''-(1,1-dimethyl-2-propenyl), di-O-de-Me: **Demethylasterriquinone B1**. LY 783281 [78860-34-1]

C₃₂H₃₀N₂O₄ 506.6

Isol. from *Pseudomassaria* fungus. Small mol. insulin mimetic. Shows antidiabetic activity. Dark purple needles (EtOAc/hexane). Mp 199-201°.

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 961-969; 991-999; 1005-1012; 1990, **38**, 2929-2932 (LY 783281, biosynth)

Mocek, U. *et al.*, *J. Antibiot.*, 1996, **49**, 854-859 (Asterriquinones CT1-CT5)

Ooike, M. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1694-1696 (PM 53)

Fredenhagen, A. *et al.*, *J. Antibiot.*, 1997, **50**, 395-401 (Semicochliodinols, Didemethylasterriquinone)

Alvi, K.A. *et al.*, *J. Antibiot.*, 1999, **52**, 215-223 (Asterriquinone SU derivs, isol, activity)

Harris, G.D. *et al.*, *Org. Lett.*, 1999, **1**, 431-433 (synth, deriv)

Zhang, B. *et al.*, *Science (Washington, D.C.)*, 1999, **284**, 974-977 (LY 783281, isol, pharmacol)

Liu, K. *et al.*, *Tet. Lett.*, 1999, **40**, 5119-5122 (Asterriquinone B1, synth)

Tatsuta, K. *et al.*, *J. Antibiot.*, 2001, **54**, 105-108 (Asterriquinone B1, synth)

Wijeratne, E.M.K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1567-1573 (isol, pmr)

Pirrung, M.C. *et al.*, *J.A.C.S.*, 2005, **127**, 4609-4624 (Demethylasterriquinone B1, synth)

Asterubin

A-1522

2-[[[(Dimethylamino)iminomethyl]amino]ethanesulfonic acid, 9CI [6249-86-1]

HN=C(NMe₂)NHCH₂CH₂SO₃HC₅H₁₃N₃O₃S 195.242


Isol. from starfish. Needles (MeOH aq.). Mp 272-273° dec.

Ackermann, D. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1935, **232**, 206; **235**, 233 (isol, synth)

Astin J

A-1523

Asterinin D [163136-31-0]

C₂₅H₃₃N₅O₇ 515.565

Constit. of the roots of *Aster tataricus* (Asteraceae). Amorph. powder (EtOH aq.). Mp 280° dec. (261°-265° dec.). [α]_D -

19.8 (c, 0.2 in EtOH aq.). [α]_D +6 (c, 0.1 in Py). λ_{max} 210 (ε 12600); 269 (ε 22400) (MeOH).

Me ester: Asterinin F

[148057-47-0]

C₂₆H₃₅N₅O₇ 529.592

Constit. of the roots of *Aster tataricus* (Asteraceae). Amorph. powder (EtOH aq.). Mp 236-241° dec. [α]_D -30.9 (c, 0.2 in EtOH aq.). λ_{max} 210 (ε 30900); 267 (ε 26900) (MeOH).

4'S,4'''S-Dihydroxy, Me ester: **Asterinin E**

[172548-92-4]

C₂₆H₃₅N₅O₉ 561.591


Constit. of the roots of *Aster tataricus* (Asteraceae). Amorph. powder (EtOH aq.). Mp 277-281° dec. [α]_D +43.1 (c, 0.3 in EtOH aq.). λ_{max} 212 (ε 34700); 267 (ε 21900) (MeOH).

Morita, H. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 271 (*Astin J*)
Cheng, D.-L. *et al.*, *Phytochemistry*, 1996, **41**, 225-227 (*Asterinins*)

Astrocasin

A-1524

1,3,4,12b-Tetrahydro-3-(1-methyl-2-piperidinyl)pyrido[2,1-a][2]benzazepin-6(2H)-one, 9CI [2114-92-3]

C₂₀H₂₆N₂O 310.438

Alkaloid from *Astrocasia phyllanthoides* (Euphorbiaceae). Mp 171-172°. [α]_D²⁴ -270 (EtOH).

Perchlorate: Mp 149-151°.**Methiodide:** Mp 227-228°.


Lloyd, H.A. *et al.*, *Tet. Lett.*, 1965, 1761 (isol, uv, ir, pmr, struct)

Bright, W.M. *et al.*, *J.O.C.*, 1976, **41**, 2454 (cryst struct, abs config)

Astrophylline

A-1525

1'-(1-Oxo-3-phenyl-2-propenyl)-2,3'-bipiperidine, 9CI. 1'-Cinnamoyl-2,3'-bipiperidine [5081-53-8]



Absolute Configuration

C₁₉H₂₆N₂O 298.427

Alkaloid from *Astrocasia phyllanthoides* (Euphorbiaceae). Viscous oil. Bp_{0.001} 115°. [α]_D²⁰ +27 (c, 0.35 in EtOH).

Perchlorate: Mp 172-174°.**Picrate:** Mp 146-148° dec.

N-Me:

Oil. [α]_D +92 (EtOH).


Lloyd, H.A. *et al.*, *Tet. Lett.*, 1965, 4537-4540 (isol, uv, ir, pmr, struct, abs config)

Schaudt, M. *et al.*, *J.O.C.*, 2003, **68**, 2913-2920 (synth, pmr, cmr, ms)

Atalanine

A-1526

trans-Atalantine [50906-85-9]

C₃₄H₃₀N₂O₉ 610.619


Alkaloid from *Atalantia ceylanica* (Rutaceae). Mp 216.5-217.5°.

Fraser, A.W. *et al.*, *Chem. Comm.*, 1973, 615 (isol, struct)

Atalaphyllidine†

A-1527

3,12-Dihydro-6,11-dihydroxy-3,3-dimethyl-7H-pyrano[2,3-c]acridin-7-one, 9CI [57959-88-3]

C₁₈H₁₅NO₄ 309.321

See also 1,3,8-Trihydroxy-2,7-diprenylacridone, T-548. Alkaloid from the root bark of *Atalantia monophylla* (Rutaceae). Mp 275° dec.

N-Me: N-Methylatalaphyllidine. 5-Hydroxynoracronycine [27067-70-5]

[27067-70-5]

C₁₉H₁₇NO₄ 323.348

Alkaloid from *Atalantia ceylanica*, *Citrus depressa* and *Citrus maxima*; metab. of Acronycine, A-120 in the guinea pig. Red needles (Et₂O/petrol). Mp 252-254°.

N-Me, di-Ac:

Prisms (Et₂O/petrol). Mp 178-181°.

11-Me ether: Mp 200°.

11-Me ether, N-Me: see 11-Methoxy, O-de-Me, in Acronycine, A-120

Di-Me ether, N-Me: 5-Methoxyacronycine Yellow needles (Et₂O/petrol). Mp 97-98°.

13-Hydroxy, N-Me: 5,13-Dihydroxynoracronycine [904327-13-5]

[904327-13-5]

C₁₉H₁₇NO₅ 339.347


Alkaloid from the stem bark of *Citrus maxima* (pummelo). Yellow. Mp 162-163°. λ_{max} 266 (log ε 4.34); 283 (log ε 4.21) (MeOH).

- Sullivan, H.R. *et al.*, *J. Med. Chem.*, 1970, **13**, 904 (*isol, deriv*)
 Fraser, A.W. *et al.*, *J.C.S. Perkin 1*, 1973, 1173 (*isol, uv, ir, nmr, ms, struct, deriv*)
 Basa, S.C. *et al.*, *Experientia*, 1975, **31**, 1387 (*isol, uv, ir, pmr, ms, struct*)
 Adams, J.H. *et al.*, *J. Nat. Prod.*, 1976, **39**, 399 (*synth, deriv*)
 Wu, T.-S. *et al.*, *Heterocycles*, 1982, **19**, 273 (*isol, deriv*)
 Teng, W.-Y. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2005, **52**, 1253-1255 (*5,13-Dihydroxynoracronycine*)

Atalane

A-1528

[50906-86-0]




Relative Configuration
Probable structure

C₃₈H₃₄N₂O₉ 662.695Alkaloid from *Atalantia ceylanica* (Rutaceae). Mp 209-210°.Fraser, A.W. *et al.*, *Chem. Comm.*, 1973, 615 (*isol, struct*)**Atalane**

A-1529

4-Methoxy-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CI. 2-Hydroxy-4-methoxy-3-prenylquinoline [7282-19-1]

C₁₅H₁₇NO₂ 243.305

Alkaloid from the heartwood of *Fagara xanthoxyloides* and the leaves of *Ravenia spectabilis* (Rutaceae). Also present in *Evodia rutaecarpa*. Possesses anthelmintic and antiparasitic activity. Needles (Me₂CO). Mp 133°.

N-Me: N-Methylatanine

[22826-43-3]

C₁₆H₁₉NO₂ 257.332

Alkaloid from the root bark and stem bark of *Almeidea guyanensis* (Rutaceae). Needles (MeOH). Mp 130°.

O-De-Me, N-Me: 4-Demethyl-N-methylatanine

[15954-07-1]

C₁₅H₁₇NO₂ 243.305

Alkaloid from the root bark of *Almeidea guyanensis* (Rutaceae). Needles (Et₂O). Mp 164°.

O-De-Me, O-(3-methyl-2-butenyl): 3-(3-Methyl-2-butenyl)-4-[(3-methyl-2-butenyl)oxy]-2(1H)-quinolinone. 3-(3,3-Dimethylallyl)-4-(3,3-dimethylallyloxy)-2-quinolone. 2-Hydroxy-3-prenyl-4-prenyloxyquinoline. 3-Prenyl-4-pre-

nyloxy-2(1H)-quinolinone

[18118-29-1]

C₁₉H₂₃NO₂ 297.396

Alkaloid from the leaves and stems of *Haplophyllum tuberculatum* (Rutaceae). Also *isol.* from *Evodia roxburghiana*. Exhibits modest anti-HIV activity against HIV-1 in cultured human lymphoblastoid CEM-SS cells. Cryst. (Me₂CO). Mp 114-115°.

2',3'-Epoxy, O-de-Me, N-Me: 3-(2,3-Epoxy-3-methylbutyl)-4-hydroxy-N-methyl-2-quinolone. 2',3'-Epoxy-4-O-demethyl-N-methylatanine

C₁₅H₁₇NO₃ 259.304

Alkaloid from the bark of *Galipea officinalis*.

Paul, B.D. *et al.*, *J. Indian Chem. Soc.*, 1968, **45**, 552 (*occur*)

Eshiett, I.T. *et al.*, *J.C.S. (C)*, 1968, 481 (*isol, uv, ir, pmr, struct*)

Lavie, D. *et al.*, *Tetrahedron*, 1968, **24**, 3011-3018 (*Prenylprenyloxyquinolinone*)

Moulis, C. *et al.*, *Phytochemistry*, 1983, **22**, 2095 (*N-Methylatanine, 4-Demethyl-N-methylatanine*)

Perrett, S. *et al.*, *Planta Med.*, 1995, **61**, 276 (*isol, activity*)

McCormick, J.L. *et al.*, *J. Nat. Prod.*, 1996, **59**, 469 (*Prenylprenyloxyquinolinone, isol*)

Shin, H.-K. *et al.*, *Planta Med.*, 1998, **64**, 764-765 (*isol, uv, pmr, cmr, ms*)


Jacquemond-Collet, I. *et al.*, *Fitoterapia*, 2000, **71**, 605-606 (*Epoxydemethyl-N-methylatanine*)

Jones, K. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 4380-4383 (*synth, pmr, cmr*)

Atanisatine

A-1530

2-Methoxy-8-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-2-methoxy-8-prenylcarbazole [58523-26-5]

C₁₉H₁₉NO₂ 293.365


Alkaloid from the stems of *Clausena anisata* (Rutaceae). Isomeric with Indizoline, I-75.

Okorie, D.A. *et al.*, *Phytochemistry*, 1975, **14**, 2720 (*occur*)

Atherospermidine

A-1531

4-Methoxy-8H-benzo[g]benzodioxolo[6,5,4-de]quinolin-8-one. 3-Methoxy-1,2-methylenedioxyxoaporphine. Psilopine [3912-57-0]

C₁₈H₁₁NO₄ 305.289

Alkaloid from the bark of *Atherosperma moschatum*, the leaves and stems of *Guatteria psilopus* and the leaves of *Enantia polycarpa* (Monimiaceae, Annonaceae). Yellow needles (CHCl₃ or Py). Mp 276-278° dec Mp 283-285°. λ_{max} 247 (ε 23000); 280 (ε 36000); 315 (ε 8900) (MeOH) (Berdy). λ_{max} 247 (ε 17000); 281 (ε 34000); 312 (ε 9600) (EtOH) (Berdy).

Hydrochloride:

Scarlet needles (5% HCl). Mp 256-258° dec.

Oxime:

Cryst. (1-butanol). Mp 247-250° dec.

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 111; 1965, **18**, 1997 (*isol, uv, struct, synth*)

Harris, W.M. *et al.*, *J.O.C.*, 1965, **30**, 432 (*isol, uv*)

Pai, B.R. *et al.*, *Tetrahedron*, 1965, **21**, 2579 (*synth*)


Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1403 (*ms*)

Jossang, A. *et al.*, *Planta Med.*, 1977, **32**, 249 (*isol*)

Atherosperminine

A-1532

3,4-Dimethoxy-N,N-dimethyl-1-phenanthreneethanamine, 9CI. 1-Dimethylaminoethyl-3,4-dimethoxyphenanthrene [5531-98-6] [98900-05-1]

C₂₀H₂₃NO₂ 309.407

Alkaloid from the bark of *Atherosperma moschatum* and *Cryptocarya angulata*, the leaves of *Enantia chlorantha*, the stem bark of *Duguetia calycina* and *Annona muricata* (sour sop), the leaves and stem bark of *Annona montana*. Needles (Me₂CO/petrol). Mp 199-200°.

Hydrochloride: [54749-97-2] Shows anti-platelet and vasorelaxing activity. Mp 234-235° dec.

Picrate: [5531-99-7]

Yellow needles (Me₂CO/MeOH). Mp 189-190°.

N-Oxide: Atherosperminine N-oxide

[91174-15-1]

C₂₀H₂₃NO₃ 325.407

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Shows antiplatelet and vasorelaxing activity. Noncryst.

N-Me: N-Methylatherosperminium

[98900-03-9]

[98900-04-0, 5532-00-3]

C₂₁H₂₆NO₂⁺ 324.442

Alkaloid from the stem bark and stem wood of *Fissistigma glaucescens* (Annonaceae). Needles (EtOH)(as iodide). Mp 282-284° (iodide).

N-De-Me: Noratherosperminine

[74606-53-4]

C₁₉H₂₁NO₂ 295.38

Alkaloid from *Duguetia* spp. (Annonaceae). Mp 180°.

2-O-De-Me: 1-[2-(Dimethylamino)ethyl]-4-methoxy-3-phenanthrenol. 1-(2-Dimethylaminoethyl)-3-hydroxy-4-methoxyphenanthrene. **Argentinine** [16625-57-3]

C₁₉H₂₁NO₂ 295.38

Alkaloid from the roots of *Aristolochia argentina*, the leaves of *Enantia chlorantha*, the stem bark and leaves of *Annona montana* and from *Monodora angolensis* (Aristolochiaceae, Annonaceae). Oil.

2-O-De-Me, N-oxide: Argentinine N-oxide

[138690-45-6]

C₁₉H₂₁NO₃ 311.38

Alkaloid from *Monocyclanthus vignei* (Annonaceae). Oil. Possible artifact.

Cooke, R.G. et al., *Aust. J. Chem.*, 1954, **7**, 99 (isol, uv)

Bick, I.R.C. et al., *Aust. J. Chem.*, 1956, **9**, 111; 1965, **18**, 1997 (isol, pmr, uv)

Priestap, H.A. et al., *Chem. Comm.*, 1967, 754 (*Argentinine, Atherosperminine*)

Aguilar-Santos, G. et al., *Philipp. J. Sci.*, 1967, **96**, 399; *CA*, **74**, 1066r (isol)

Priestap, H.A. et al., *An. Asoc. Quim. Argent.*, 1972, **60**, 309; *CA*, **77**, 164898q (*Argentinine, Atherosperminine*)

Roblot, F. et al., *Plant. Med. Phytother.*, 1978, **12**, 259

Guinaudeau, H. et al., *J. Nat. Prod.*, 1979, **42**, 325 (*Noratherosperminine*)

Leboeuf, M. et al., *Planta Med.*, 1980, **38**, 33; 1981, **42**, 37

Castedo, L. et al., *Heterocycles*, 1982, **19**, 209 (*Noratherosperminine, synth*)

Hocquemiller, R. et al., *J. Nat. Prod.*, 1984, **47**, 353 (*Atherosperminine N-oxide*)

Lu, S.T. et al., *Phytochemistry*, 1985, **24**, 1829 (*N-Methylatherosperminium*)

Achenbach, H. et al., *J. Nat. Prod.*, 1991, **54**, 1331 (*Argentinine N-oxide*)


Chen, K.-S. et al., *J. Nat. Prod.*, 1996, **59**, 531-534 (activity)

Lopez-Martin, J. et al., *Chem. Pharm. Bull.*, 2002, **50**, 1613-1615 (*Argentinine, isol, pmr, cmr*)

Atidine

A-1533

[467-91-4]



C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Prisms (C₆H₆ or Et₂O). Mp 182.5-183.5°. [α]_D²⁵ -47 (CHCl₃). pK_a 7.53 (50% MeOH).

Hydrochloride:

Rosettes (MeOH/Me₂CO). Mp 204-215° dec.

7α-Alcohol: Dihydroajaconine

[1466-07-5]

C₂₂H₃₅NO₃ 361.523

Minor alkaloid from the seeds of *Consolida ambigua* (*Delphinium ajacis*),

also obt. by mild redn. of Ajaconine (Ranunculaceae). Needles (EtOH aq.). Mp 99-100°. [α]_D²⁴ -35 (c, 1.0 in EtOH).

7α-Alcohol, 2'-aldehyde: Consorientaline

C₂₂H₃₃NO₃ 359.508

Alkaloid from *Consolida orientalis*.

Amorph. [α]_D²⁵ -22.7 (c, 0.33 in CHCl₃).

1',2'-Didehydro, 7α-alcohol: Uncinatine

C₂₂H₃₃NO₃ 359.508

Alkaloid from *Delphinium uncinatum*. Amorph.

7-Deoxo: Dihydroatisine

[17934-91-7]

C₂₂H₃₅NO₂ 345.524

Alkaloid from roots of *Aconitum heterophyllum* (Ranunculaceae). Cryst. (Me₂CO). Mp 159-161°. [α]_D²⁵ -44.5 (EtOH). [α]_D²⁷ -43.9 (c, 1.41 in CHCl₃).

7-Deoxo, 2'-aldehyde: Chellespontine

[149998-46-9]

C₂₂H₃₃NO₂ 343.508

Alkaloid from aerial parts of *Consolida hellespontica* (Ranunculaceae). Plates. Mp 227-230°. [α]_D²⁵ +14.6 (c, 0.56 in MeOH).

7-Deoxo, 19-oxo: 19-Oxodihydroatisine

C₂₂H₃₃NO₃ 359.508

Alkaloid from *Delphinium staphisagria*. Cryst. (CHCl₃). Mp 221-223°. [α]_D²⁵ -31.2 (c, 0.57 in CHCl₃).

7-Deoxo, 19-oxo, 2'-Ac: 22-O-Acetyl-19-oxodihydroatisine

C₂₄H₃₅NO₄ 401.545

Alkaloid from *Delphinium staphisagria*. Gum. [α]_D²⁵ -35.2 (c, 0.54 in CHCl₃).

14-Oxo, 15-deoxy, 7α-alcohol: Spiramine G

[142750-44-5]

C₂₂H₃₃NO₃ 359.508

Alkaloid from roots of *Spiraea japonica* var. *acuminata* (Rosaceae). Needles (Et₂O). Mp 160-162°. [α]_D²⁰ -16 (c, 0.95 in CHCl₃).

15-Epimer, 7α-alcohol, 2'-aldehyde: Spiramine A

C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Spiraea japonica* var. *acuta*. Amorph. powder. [α]_D²⁴ -6.2 (c, 1 in MeOH).

15-Epimer, 7-deoxo, 14-oxo: Spiramine H

[142750-45-6]

C₂₂H₃₃NO₃ 359.508

Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae) and *Aconitum kusnezoffii*. Cryst. (Me₂CO/cyclohexane). Mp 150-151°. [α]_D¹⁷ -39.7 (c, 0.295 in EtOH).

15-Epimer, 7-deoxo, 14-oxo, 15-Ac: Spiramine I

[142761-31-7]

C₂₄H₃₅NO₄ 401.545

Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae).

15-Epimer, 7-deoxo, 14-oxo, 1α-hydroxy: Beiwusine A

[224790-29-8]

C₂₂H₃₃NO₄ 375.507

Alkaloid from the roots of *Aconitum kusnezoffii*. Amorph. powder. [α]_D¹⁷ -34.1 (c, 0.41 in EtOH).

15-Epimer, 7-deoxo, 14-oxo, 1β-hydroxy: Beiwusine B

[224790-30-1]

C₂₂H₃₃NO₄ 375.507

Alkaloid from the roots of *Aconitum kusnezoffii*. Amorph. powder. [α]_D¹⁷ -42 (c, 0.41 in EtOH).

Dvornik, D. et al., *Tetrahedron*, 1961, **14**, 54 (*Atidine, synth*)

Pelletier, S.W. et al., *J.A.C.S.*, 1965, **87**, 777; 799 (*Atidine, ir, pmr, struct, synth*)

Pelletier, S.W. et al., *Phytochemistry*, 1968, **7**, 625 (*Atidine, isol*)

Pelletier, S.W. et al., *Tetrahedron*, 1968, **24**, 2019 (*Atidine, pmr*)

Pelletier, S.W. et al., *J.A.C.S.*, 1978, **100**, 7976 (*Dihydroatisine, cryst struct, abs config*)

Finer-Moore, J. et al., *Cryst. Struct. Commun.*, 1979, **8**, 649 (*Atidine, cryst struct*)

Pelletier, S.W. et al., *J. Nat. Prod.*, 1980, **43**, 395 (*Dihydroajaconine*)

Hao, X.-J. et al., *Yunnan Zhiwu Yanjiu*, 1991, **13**, 452; 1994, **16**, 301; *CA*, **117**, 108080z; **122**, 101603s (*Spiramines H,I*)

Hao, X.-J. et al., *Heterocycles*, 1993, **36**, 825 (*Spiramine G, isol, ir, pmr, cmr, cryst struct*)

Desai, H.K. et al., *Heterocycles*, 1993, **36**, 1081 (*Chellespontine*)

Li, Z.-B. et al., *J. Asian Nat. Prod. Res.*, 1998, **1**, 87-92 (*Beiwusines A,B*)

Ulubelen, A. et al., *Phytochemistry*, 1998, **47**, 1141-1144 (*Uncinatine*)

Diaz, J.G. et al., *J. Nat. Prod.*, 2000, **63**, 1136-1139 (*19-Oxodihydroatisine, 22-Acetyl-19-oxodihydroatisine*)

He, H.-P. et al., *J. Nat. Prod.*, 2001, **64**, 379-380 (*Spiratine A*)


Merikli, F. et al., *J. Nat. Prod.*, 2001, **64**, 787-789 (*Consortientaline*)

Atisine, 9CI

A-1534

Anthorine

[466-43-3]



Absolute Configuration

C₂₂H₃₃NO₂ 343.508

Exists as a mixt. of C-20 epimers in soln. Alkaloid from *Aconitum heterophyllum*, *Aconitum gigas*, *Aconitum anthara* and *Aconitum heterophyllodes* (Ranunculaceae). Resin. [α]_D²¹ -21 (c, 2.4 in EtOH). [α]_D²⁰ -30.7 (EtOH). pK_a 12.8 (50% MeOH). Undergoes a ready isomerisation of the oxazolidine ring to Isoatisine, I-192 on treatment with methanolic alkali or even by boiling MeOH.

Hydrochloride: Atisinium chloride. Guanfu base H

[4758-99-0]

Mp 311-312° (303°) dec. [α]_D²⁵ +28 (c, 1.10 in H₂O). Exists as a quaternary iminium salt rather than a protonated NH⁺ salt.

7β,19R-Dihydroxy: Delphatisine A [1003600-46-1]

C₂₂H₃₃NO₄ 375.507

Alkaloid from *Delphinium chrysotrichum*. Amorph. powder. [α]_D²⁷ +16 (c, 1 in CHCl₃).

6β,7α-Diacetoxy, 15-deoxy, 19-oxo: Spiramide†

[301541-30-0]
 $C_{26}H_{35}NO_6$ 457.566
 Alkaloid from the roots of *Spiraea japonica* var. *acuta*. Amorph. powder. $[\alpha]_D^{26}$ -63.4 (c, 4.2 in $CHCl_3$).

15-Epimer, 19-oxo, 7 α -hydroxy: **Deacetylsiramine S**
 [340814-10-0]
 $C_{22}H_{31}NO_4$ 373.491
 Alkaloid from *Spiraea japonica* var. *ovalifolia*. Prisms ($CHCl_3/MeOH$). Mp 113-115°. $[\alpha]_D^{21}$ -74.3 (c, 0.27 in MeOH).


15-Epimer, 19-oxo, 7 α -hydroxy, 7-Ac: **Siramine V**
 [501327-99-7]
 $C_{24}H_{33}NO_5$ 415.528
 Alkaloid from *Spiraea japonica* var. *ovalifolia*.

15-Epimer, 19-oxo, 7 α -hydroxy, 15-Ac: **Siramine S**
 $C_{24}H_{33}NO_5$ 415.528
 Alkaloid from *Spiraea japonica*. Incorrect struct. given in one ref.

Lawson, A. et al., *J.C.S.*, 1937, 1640-1643 (isol)
 Jacobs, W.A. et al., *J. Biol. Chem.*, 1942, **143**, 589-603 (isol)
 Edwards, O.E. et al., *Can. J. Chem.*, 1954, **32**, 465-473; 1955, **33**, 448-451 (isol, uv, ir, struct)
 Nagata, W. et al., *J.A.C.S.*, 1963, **85**, 2342-2343; 1967, **89**, 1483-1499 (synth)
 Dvornik, D. et al., *Can. J. Chem.*, 1964, **42**, 137-149 (struct)
 Masamune, S. et al., *J.A.C.S.*, 1964, **86**, 291-292 (synth)
 Pelletier, S.W. et al., *Phytochemistry*, 1968, **7**, 625-635 (isol, isom)
 Pelletier, S.W. et al., *Tetrahedron*, 1968, **24**, 2019-2038 (pmr)
 Pradhan, S.K. et al., *Chem. Comm.*, 1970, 644-645 (struct)
 Scott, A.I. et al., *Tetrahedron*, 1971, **27**, 4787-4819 (cd)
 Pelletier, S.W. et al., *J.A.C.S.*, 1977, **99**, 284-286; 1978, **100**, 7976-7987 (cmr, cryst struct, abs config)
 Mody, N.V. et al., *Tetrahedron*, 1978, **34**, 2421-2431 (cmr)
 Shishido, K. et al., *Chem. Comm.*, 1987, 1360-1361 (synth)
 Ihara, M. et al., *J.A.C.S.*, 1990, **112**, 1164-1171 (synth)
 Banerjee, A.K. et al., *Bull. Soc. Chim. Belg.*, 1994, **103**, 67-82 (rev, synth)
 Hao, X.Y. et al., *Guizhou Kexue*, 1995, **13**, 25-28 (Siramine S)
 Zuo, G.Y. et al., *Heterocycles*, 2001, **55**, 487-493 (Deacetylsiramine S)
 He, H.P. et al., *J. Nat. Prod.*, 2001, **64**, 379-380 (Spiramide, Siramine S)
 Li, L. et al., *Eur. J. Pharmacol.*, 2002, **449**, 23-28 (Siramine V)
 He, Y.Q. et al., *Chin. Chem. Lett.*, 2007, **18**, 545-547 (Delphatisine A)

Atpenin

A-1535



Atpenin A_4 $R^1 = Cl$, $R^2 = H$
 A_5 $R^1 = R^2 = Cl$
 B $R^1 = R^2 = H$

Antibiotic complex of which 3 components have been isol. Prod. by *Penicillium* sp. FO-125. Antifungal agent.

Atpenin A_4 [119509-25-0]
 $C_{15}H_{22}ClNO_5$ 331.795
 Powder. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 98°. $[\alpha]_D^{22}$ -8.6 (c, 1 in EtOH). λ_{max} 240 (€ 11500); 270 (€ 7500); 318 (€ 7000) (EtOH/HCl) (Derep). λ_{max} 242 (€ 9500); 290 (€ 9600); 355 (€ 4300) (EtOH/NaOH) (Derep). λ_{max} 235 (€ 12200); 267 (€ 8400); 320 (€ 5800) (EtOH) (Derep).
 ▶ LD₅₀ (mus, ipr) 50 - 150 mg/kg.
 CJ8794000

Atpenin A_5
 WF 16775A₁. Antibiotic WF 16775A₁ [119509-24-9]
 $C_{15}H_{21}Cl_2NO_5$ 366.24
 Also prod. by *Chaetabolisia erysiphoides*. Angiogenesis inhibitor. Needles. Mp 86°. $[\alpha]_D^{22}$ -0.8 (c, 1 in EtOH). λ_{max} 240 (€ 11500); 270 (€ 7000); 318 (€ 7000) (EtOH/HCl) (Derep). λ_{max} 242 (€ 9500); 290 (€ 9500); 355 (€ 4300) (EtOH/NaOH) (Derep). λ_{max} 236 (€ 16800); 270 (€ 11300); 319 (€ 9500) (EtOH) (Derep).
 ▶ CJ8800000


5'-Chloro: Antibiotic WF 16775A₂. WF 16775A₂
 $C_{15}H_{20}Cl_3NO_5$ 400.685
 Prod. by *Chaetabolisia erysiphoides*. Angiogenesis and neovascularisation inhibitor. Needles. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 119.5-121°. $[\alpha]_D^{20}$ -52 (c, 0.5 in $CHCl_3$). λ_{max} 240 (€ 11500); 270 (€ 7500); 318 (€ 7000) (MeOH/HCl) (Derep). λ_{max} 242 (€ 9500); 290 (€ 9600); 355 (€ 4300) (MeOH/NaOH) (Derep). λ_{max} 236 (€ 19200); 269 (€ 12400); 322 (€ 11600) (MeOH) (Derep). λ_{max} 235 (€ 14600); 270 (€ 9600); 322 (€ 8400) (MeOH-HCl) (Berdy). λ_{max} 245 (€ 11400); 292 (€ 11200); 350 (€ 4000) (MeOH-NAOH) (Berdy).

Atpenin B [119509-26-1]
 $C_{15}H_{23}NO_5$ 297.35
 Prod. by *Chaetabolisia erysiphoides*. Powder. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 78°. $[\alpha]_D^{22}$ -27 (c, 1 in EtOH). λ_{max} 240 (€ 11500); 270 (€ 7500); 318 (€ 7000) (EtOH/HCl) (Derep). λ_{max} 242 (€ 9500); 290 (€ 9600); 355 (€ 4300) (EtOH/NaOH) (Derep). λ_{max} 237 (€ 5600); 270 (€ 3600); 318 (€ 3100) (EtOH) (Derep).
 ▶ LD₅₀ (mus, ipr) 50 - 150 mg/kg.
 CJ8810000
 Omura, S. et al., *J. Antibiot.*, 1988, **41**, 1769 (isol)
 Kumagai, H. et al., *J. Antibiot.*, 1990, **43**, 1553 (pmr, cmr, struct)
 Otsuka, T. et al., *J. Antibiot.*, 1992, **45**, 1970
 Trecourt, F. et al., *J.O.C.*, 1994, **59**, 6173 (synth, Atpenin B)

Atractylenolactam

A-1536

[193757-67-4]




$C_{15}H_{19}NO$ 229.321
 Constit. of *Atractyloides macrocephala*. Needles. Mp 225-227°.
 Chen, Z.-L. et al., *Phytochemistry*, 1997, **45**, 765-767 (isol, pmr, cmr)

Aucherine

A-1538

[123715-12-8]




$C_{17}H_{25}NO_6$ 339.388
 Alkaloid from aerial parts of *Senecio integrifolius* subsp. *aucheri* (Asteraceae).

Sener, B. et al., *Gazi Univ. Eczacilik Fak. Derg.*, 1988, **5**, 157; *CA*, **111**, 211907z

Augustamine

A-1539

[86734-72-7]



Absolute Configuration

$C_{17}H_{19}NO_4$ 301.341
 Alkaloid from the leaves, bulbs and roots of *Crinum augustum*. Prisms (Me_2CO). Mp 173-175°. $[\alpha]_D^{20}$ -83 (c, 1.4 in $CHCl_3$). λ_{max} 214 (€ 83440); 241 (€ 45510); 292 (€ 46300) (MeOH).

N-De-Me: **Noraugustamine**

$C_{16}H_{17}NO_4$ 287.315
 Alkaloid from the bulbs of *Crinum kirkii*. Mp 149-151°. $[\alpha]_D^{20}$ -50 (c, 0.9 in MeOH).

N,4a-Didehydro, N-de-Me: **4a,N-Didehydronoraugustamine**


$C_{16}H_{15}NO_4$ 285.299
 Alkaloid from the bulbs of *Crinum kirkii*. Mp 127-130°. $[\alpha]_D^{20}$ -242.6 (c, 0.27 in MeOH).

Ali, A.A. et al., *Phytochemistry*, 1981, **20**, 1121-1123; 1983, **22**, 283-287 (isol, uv, ir, pmr, cmr, ms)
 Pearson, W.H. et al., *J.O.C.*, 1998, **63**, 3607-3617 (synth)

Joshi, B.S *et al.*, *J. Chem. Crystallogr.*, 2000, **30**, 135-138 (*cryst struct*)
 Machocho, A.K. *et al.*, *Phytochemistry*, 2004, **65**, 3143-3149 (*Noraugustamines*)

Aulosirazole**A-1540**

5-Hydroxy-3-methoxynaphth[2,3-d]isothiazole-4,9-dione, 9CI
 [159194-92-0]




C₁₂H₇NO₄S 261.258

Isol. from the blue-green alga *Aulosira fertilissima*. Cytotoxic agent. Deep yellow cryst. (MeOH aq.). λ_{max} 216 (ε 20500); 248 (ε 11200); 272 (ε 8500); 424 (ε 6300) (MeOH). λ_{max} 226 ; 280 ; 312 ; 536 (MeOH/NaOH) (Berdy).

Stratmann, K. *et al.*, *J.O.C.*, 1994, **59**, 6279-6281

Aurachin A**A-1541**

α-(4,8-Dimethyl-3,7-nonadienyl)-1,2-dihydro-α,4-dimethylfuro[2,3-c]quinoline-2-methanol 5-oxide, 9CI
 [108354-15-0]




C₂₅H₃₃NO₃ 395.541

Prod. by *Stigmatella aurantiaca* Sg a15. Active against gram-positive bacteria and a few yeasts and molds. Cryst. (Et₂O). Sol. MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 111-112°. [α]_D -49.2 (c, 0.4 in MeOH). λ_{max} 219 (ε 22100); 239 (ε 24400); 249 (ε 26000); 304 (ε 3600); 317 (ε 4200); 354 (ε 5200); 369 (ε 6100) (MeOH) (Derep). λ_{max} 237 (ε 32000); 300 (ε 1700); 315 (ε 2400); 354 (ε 5100); 365 (ε 6100) (MeOH/HCl aq.).

Kunze, B. *et al.*, *J. Antibiot.*, 1987, **40**, 258-265 (*isol, props*)
 Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*pmr, cmr, biosynth*)

Aurachin B**A-1542**

[108354-12-7]




C₂₅H₃₃NO₂ 379.541

Prod. by *Stigmatella aurantiaca* Sg a15. Active against gram-positive bacteria and a few yeasts and molds. Cryst. (Et₂O). Mp 93-94°. λ_{max} 242 (ε 39600); 288 (sh) (ε 2900); 302 (ε 5500); 315 (ε 6900); 342 (ε 7100); 352 (ε 7600) (pH 2.5 MeOH) (Derep).

Kunze, B. *et al.*, *J. Antibiot.*, 1987, **40**, 258-265 (*isol, props*)
 Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*pmr, cmr, biosynth*)

Aurachin D**A-1543**

[108354-13-8]



C₂₅H₃₃NO 363.542

Prod. by *Stigmatella aurantiaca* Sg a15. Active against gram-positive bacteria and a few yeasts and molds. Cryst. (Et₂O). Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 165-168°. λ_{max} 240 (ε 33000); 245 (ε 33500); 280 (sh) (ε 3300); 291 (sh) (ε 4300); 308 (sh) (ε 7400); 321 (ε 11400); 334 (ε 11400) (MeOH) (Derep).

N-Hydroxy: **Aurachin C**

[108354-14-9]


C₂₅H₃₃NO₂ 379.541

From *Stigmatella aurantiaca* Sg a15. Active against gram-positive bacteria and a few yeasts and molds. Cryst. (Et₂O). Mp 124-125°. λ_{max} 212 (ε 34000); 249 (ε 30600); 334 (ε 10400) (MeOH).

Kunze, B. *et al.*, *J. Antibiot.*, 1987, **40**, 258-265 (*isol, props*)
 Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*biosynth, pmr, cmr*)

Aurachin E**A-1544**

4-Methyl-5-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-6H-imidazo[4,5,1-ij]quinoline-2,6(1H)-dione, 9CI
 [113366-10-2]



C₂₆H₃₂N₂O₂ 404.551


Isol. from *Stigmatella aurantiaca* Sg a15. Needles (Et₂O). Mp 168-169°. λ_{max} 241 (log ε 4.43); 249 (log ε 4.49); 300 (sh) (log ε 3.68); 339 (log ε 4.12) (MeOH).

Ger. Pat., 1986, 3 520 229; CA, **110**, 73858k
 Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*biosynth, pmr, cmr, ms*)

Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1967-1969 (*synth*)

Aurachin F**A-1545**

3-(4,8-Dimethyl-3,7-nonadienyl)-3,5-dimethyl-3H-pyrano[2,3-c]quinoline 6-oxide, 9CI
 [113366-11-3]



C₂₅H₃₁NO₂ 377.525


Alkaloid from *Stigmatella aurantiaca* Sg a15. Amorph. solid. λ_{max} 241 (sh) (log ε 4.27); 253 (log ε 4.43); 267 (sh) (log ε 4.1); 279 (sh) (log ε 3.89); 305 (log ε 3.49); 320 (log ε 3.62); 336 (log ε 3.75); 373 (log ε 3.64); 390 (log ε 3.69) (MeOH).

Ger. Pat., 1986, 3 520 229; CA, **110**, 73858k

Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*pmr, cmr, biosynth*)

Aurachin G**A-1546**

α-(4,8-Dimethyl-1,3,7-nonatrienyl)-2,3-dihydro-α,4-dimethylfuro[3,2-c]quinoline-2-methanol 5-oxide, 9CI
 [113366-12-4]



C₂₅H₃₁NO₃ 393.525

Isol. from *Stigmatella aurantiaca* Sg a15. Amorph. solid. λ_{max} 210 (log ε 4.47); 238 (log ε 4.52); 247 (log ε 4.55); 350 (sh) (log ε 4.02) (MeOH).

1',2'-Dihydro: **Aurachin H**

[113366-13-5]

C₂₅H₃₃NO₃ 395.541

From *Stigmatella aurantiaca* Sg a15. Amorph. solid. λ_{max} 216 (log ε 4.62); 249 (log ε 4.57); 339 (log ε 4.07) (MeOH).

3ξ-Methoxy, 1',2'-dihydro: **Aurachin I**

[113366-14-6]

C₂₆H₃₅NO₄ 425.567


From *Stigmatella aurantiaca* Sg a15. Amorph. solid. λ_{max} 216 (log ε 4.62); 249 (log ε 4.56); 350 (log ε 3.95) (MeOH).

Ger. Pat., 1986, 3 520 229; CA, **110**, 73858k

Höfle, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1843-1849 (*pmr, cmr, biosynth*)


Auranthine

[107290-05-1]

C₁₉H₁₄N₄O₂ 330.345

Metab. from sporing cultures of *Penicillium aurantiogriseum*. Mycotoxin. Ne-phrotoxic agent. Amorph. solid. Sol. MeOH, C₆H₆; poorly sol. H₂O. $[\alpha]_D^{25}$ -164 (c, 1 in EtOH). λ_{\max} 228 (ε 38400); 268 (ε 8790); 280 (ε 7470); 310 (ε 3760); 312 (ε 3850) (MeOH) (Berdy).

Yeulet, S.E. *et al.*, *J.C.S. Perkin 1*, 1986, 1891 (isol, uv, ir, pmr, cmr, ms, struct)


AurantiamineAlkaloid 302
[143085-86-3]Absolute
ConfigurationC₁₆H₂₂N₄O₂ 302.375

Metab. from *Penicillium aurantiogriseum* var. *aurantiogriseum* and *Penicillium aurantiogriseum* var. *neoechinulatum*. Mycotoxin. Cryst. Mp 238-239°. $[\alpha]_D^{23}$ -116 (c, 0.5 in MeOH). λ_{\max} 230; 320 (MeCN aq.) (Berdy).

Larsen, T.O. *et al.*, *Phytochemistry*, 1992, 31, 1613-1615 (isol, uv, pmr, cmr, ms)

Aurantioclavine

3,4,5,6-Tetrahydro-6-(2-methyl-1-propenyl)-1H-azepino[5,4,3-cd]indole, 9CI

C₁₅H₁₈N₂ 226.321**(-)-form** [80152-02-9]

Isol. from *Penicillium aurantio-virens*. No opt. rotn. reported.

N⁵-Et: N⁵-Ethylaurantioclavine

[174792-01-9]

C₁₇H₂₂N₂ 254.374

A-1547

Prod. by *Penicillium janczewskii*.**(±)-form** [99211-67-3]

Off-white powder. Mp 194-195°.

Kozlovskii, A.G. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1981, 260, 230-233; *CA*, 96, 3403b (isol, struct)

Sakharovskii, V.G. *et al.*, *Khim. Prir. Soedin.*, 1983, 656-657; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, 19, 626-627 (pmr, conform)


Yamada, F. *et al.*, *Chem. Pharm. Bull.*, 1985, 33, 2162-2163 (synth, ir, pmr)

Hegedus, L.S. *et al.*, *J.O.C.*, 1987, 52, 3319-3322 (synth, bibl)

Kozlovskii, A.G. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 1997, 33, 70-74 (N-Ethylaurantioclavine)

Aurantioside A

[137895-70-6]

C₃₆H₄₆Cl₂N₂O₁₅ 817.669

Tetramic acid glycoside. Geom. isom. of terminal double bond revised in 1999.

Isol. from the sponge *Theonella* sp. Cytotoxic agent. Orange solid. $[\alpha]_D^{23}$ -568 (c, 0.1 in MeOH). λ_{\max} 242 (ε 12100); 414 (ε 46700) (H₂O).

O-De-Me: **Aurantioside B**

[137895-71-7]

C₃₅H₄₄Cl₂N₂O₁₅ 803.642


Isol. from *Theonella* sp. Cytotoxic agent. Orange solid. $[\alpha]_D^{23}$ -492 (c, 0.1 in MeOH). λ_{\max} 242 (ε 10900); 414 (ε 49100) (H₂O).

Matsunaga, S. *et al.*, *J.A.C.S.*, 1991, 113, 9690-9692 (isol, uv, ir, pmr, cmr, ms)

Sata, N.U. *et al.*, *J. Nat. Prod.*, 1999, 62, 969-971 (struct)

Aurantioside C

[219297-37-7]

C₃₇H₄₆Cl₂N₂O₁₅ 829.68

Tetramic acid deriv. Isol. from the sponge

A-1550

Homophymia conferta. Amorph. red powder. $[\alpha]_D^{20}$ -480 (c, 0.2 in CH₂Cl₂/MeOH). λ_{\max} 252 (ε 9700); 426 (ε 41600) (H₂O).

Wolf, D. *et al.*, *J. Nat. Prod.*, 1999, 62, 170-172 (isol, uv, ir, pmr, cmr, ms)

Aurantioside E

A-1552

[233608-98-5]

As Aurantoside A, A-1550 with

R = -CH=CHCH=CCl(CH₃) (E,Z-)C₃₈H₄₈Cl₂N₂O₁₅ 843.707

Tetramic acid glycoside. Isol. from the sponge *Siliquariaspongia japonica*. Antifungal agent. Red solid. $[\alpha]_D^{24}$ -1038 (c, 0.001 in MeOH). λ_{\max} 250 (log ε 4.12); 423 (log ε 4.99) (MeOH).

O-De-Me: **Aurantioside D**

[233608-94-1]

C₃₇H₄₆Cl₂N₂O₁₅ 829.68

Isol. from *Siliquariaspongia japonica*. Antifungal agent. Red solid. $[\alpha]_D^{24}$ -536 (c, 0.001 in MeOH). λ_{\max} 244 (log ε 4.3); 433 (log ε 4.83) (MeOH).

Sata, N.U. *et al.*, *J. Nat. Prod.*, 1999, 62, 969-971 (isol, uv, pmr, cmr)

Aurantioside F

A-1553

[233608-99-6]

As Aurantoside A, A-1550 with


R = -CH=CHCH=CHCH=CClCH₃ (Z-)C₄₀H₅₀Cl₂N₂O₁₅ 869.745

Tetramic acid glycoside. Isol. from the sponge *Siliquariaspongia japonica*. Antifungal agent. Red solid. $[\alpha]_D^{24}$ -1012 (c, 0.001 in MeOH). λ_{\max} 440 (log ε 4.79); 465 (log ε 4.85) (MeOH).

Sata, N.U. *et al.*, *J. Nat. Prod.*, 1999, 62, 969-971 (isol, uv, pmr, cmr)

Aurantioside G

A-1554

C₂₃H₂₇ClN₂O₈ 494.928

Tetramic acid glycoside. Isol. from the sponge *Theonella swinhoei*. Orange solid. $[\alpha]_D^{24}$ -116 (c, 0.1 in MeOH). λ_{\max} 240 (log ε 3.72); 280 (log ε 3.69); 376 (log ε 3.57) (MeOH).

2'-O-β-D-Arabinopyranosyl: **Aurantioside H**C₂₈H₃₅ClN₂O₁₂ 627.043

Isol. from *Theonella swinhoei*. Orange solid. $[\alpha]_D^{24}$ -295 (c, 0.1 in MeOH). λ_{\max} 240 (log ε 3.79); 280 (log ε 3.79); 434 (log ε 3.95) (MeOH).

2'-O-[5-Deoxy-2-O-methyl-β-D-arabinofuranosyl-(1→4)-β-D-arabinopyranosyl]: **Aurantioside I**


C₃₄H₄₅ClN₂O₁₅ 757.187

Isol. from *Theonella swinhoei*. Orange solid. $[\alpha]_D^{24}$ -210 (c, 0.1 in MeOH). λ_{\max} 242 (log ϵ 3.68); 282 (log ϵ 3.68); 430 (log ϵ 3.72) (MeOH).

Ratnayake, A.S. *et al.*, *J. Nat. Prod.*, 2005, **68**, 104-107 (*isol*, *pmr*, *cmr*)

Aureoverticillactam A-1555

22-(2-Hexenyl)-9,10,12-trihydroxy-17-methylazaacyclodocosa-3,5,7,13,15,17,19-heptaen-2-one, 9CI
[682774-28-3]

C₂₈H₃₉NO₄ 453.62


Closely related to Antibiotic BE 14106, A-1127. Prod. by the marine *Streptomyces aureoverticillatus* (NPS001583). Shows moderate cytotoxic activity. Amorph. solid. λ_{\max} 283 (sh); 290; 310 (sh); 330 (sh) (MeCN aq.).

Mitchell, S.S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1400-1402 (*isol*, *pmr*, *cmr*)

Austamide

[34427-31-1]

A-1556



Absolute Configuration

C₂₁H₂₁N₃O₃ 363.415

Metab. of *Aspergillus ustus*. Tremorgenic toxin. Yellow amorph. solid. Sol. MeOH, CHCl₃. $[\alpha]_D^{20}$ +152 (c, 1 in EtOH). λ_{\max} 234 (ϵ 26300); 256 (ϵ 2344); 282 (ϵ 8700); 392 (ϵ 2700) (EtOH) (Berdy).

12*S*,13-Dihydro: 12,13-Dihydroaustamide
[34506-79-1]

C₂₁H₂₃N₃O₃ 365.431

Minor metab. of *Aspergillus ustus*. Cryst. (Me₂CO). Mp 235-238°. $[\alpha]_D^{22}$ +55 (c, 1.1 in CHCl₃). Fluorescent.

12*ξ*-Hydroxy, 12,13-dihydro: 12,13-Dihydro-12-hydroxyaustamide
[59476-59-4]

C₂₁H₂₃N₃O₄ 381.43

Minor metab. from *Aspergillus ustus*. Cryst. (MeOH). Mp 164-165°.

Coetzer, J. *et al.*, *Acta Cryst. B*, 1973, **29**, 685 (*cryst struct*)

Steyn, P.S. *et al.*, *Tetrahedron*, 1973, **29**, 107-120 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *deriv*)

Steyn, P.S. *et al.*, *Phytochemistry*, 1976, **15**, 355-356 (*dihydro*)

Harrison, D.M. *et al.*, *Tet. Lett.*, 1981, **22**, 2501-2504 (*synth*)


Stocking, E.M. *et al.*, *J.A.C.S.*, 2000, **122**, 9089-9098 (*biosynth*)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (*rev*, *synth*, *biosynth*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 484

Austrodimerine

[206440-45-1]



Relative Configuration

C₂₁H₂₇N₃O₅ 401.461


Alkaloid from *Osmanthus austrocaledonica*. $[\alpha]_D^{20}$ +75 (C, 1 in CHCl₃). λ_{\max} 207 (log ϵ 4.27); 232 (log ϵ 3.79); 290 (log ϵ 4.17) (MeOH).

Benkrief, R. *et al.*, *Phytochemistry*, 1998, **47**, 825-832 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Autumnaline

A-1558

1,2,3,4-Tetrahydro-7-hydroxy-1-(3-hydroxy-4,5-dimethoxyphenethyl)-6-methoxy-2-methylisoquinoline. Alkaloid CC22



(R)-form

C₂₁H₂₇NO₅ 373.448

(R)-form [23068-65-7]

Alkaloid from *Colchicum cornigerum*, *Colchicum visianii*, *Colchicum latifolium* and *Colchicum ritchii* (Liliaceae). Mp 166-168°. $[\alpha]_D^{22}$ -5 (c, 0.2 in CHCl₃). $[\alpha]_D$ -8 (c, 0.29 in MeOH). Incorr. given (S)-config. in CA.

O⁶-De-Me, O⁷-Me: Isoautumnaline
[111509-11-6]

C₂₁H₂₇NO₅ 373.448

Alkaloid from *Colchicum ritchii* (Liliaceae). Amorph. $[\alpha]_D$ -6 (c, 0.4 in CHCl₃). $[\alpha]_D$ -1 (c, 0.4 in MeOH).

(±)-form

Mp 170-172°.

Hydrochloride: Mp 235°.

O⁶-De-Me, O⁷-Me:Prisms (CHCl₃/hexane). Mp 190-191°.

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540 (*isol*, *uv*, *ir*)

Battersby, A.R. *et al.*, *J.C.S. (C)*, 1971, 3514 (*uv*, *ir*, *ms*, *pmr*, *struct*)


Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1972, 1741 (*synth*, *biosynth*)

Freyer, A.J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 684 (*Autumnaline*, *Isoautumnaline*, *isol*, *cd*, *pmr*, *abs config*)

Nimgirawath, S. *et al.*, *Aust. J. Chem.*, 1994, **47**, 957 (*Isoautumnaline*, *synth*)

Auxarconjugatin A

A-1559



(all-E)-form

C₁₉H₁₈ClNO₃ 343.809

Related to Rumbrin, R-160.

(all-E)-form

Alkaloid from *Auxarthron conjugatum* and *Gymnoascus reessii*. Cytotoxic. Red powder (MeOH aq.). Mp >300°. λ_{\max} 205 (log ϵ 4.19); 222 (log ϵ 4.14); 267 (log ϵ 4.34); 327 (sh) (log ϵ 4.25); 340 (log ϵ 4.31); 442 (log ϵ 4.91) (MeOH).

Dechloro: Auxarconjugatin C

C₁₉H₁₉NO₃ 309.364

Alkaloid from *Auxarthron conjugatum*. Red prisms (MeOH aq.). Mp >300°. λ_{\max} 204 (log ϵ 4.27); 224 (log ϵ 4.17); 266 (log ϵ 4.38); 328 (sh) (log ϵ 4.16); 340 (log ϵ 4.21); 448 (log ϵ 4.91) (MeOH).

Lower homologue (R = H): Auxarconjugatin B

C₁₈H₁₆ClNO₃ 329.782

Alkaloid from *Auxarthron conjugatum*. Red prisms (MeCN aq.). Mp 203-205°. λ_{\max} 220 (log ϵ 3.6); 261 (log ϵ 3.75); 321 (sh) (log ϵ 3.6); 331 (log ϵ 3.66); 435 (log ϵ 4.33) (MeOH).

(1'E,3'Z,5'E,7'E)-form

Alkaloid from *Auxarthron conjugatum*. Red cryst. (MeOH aq.). Mp 123-125°. λ_{\max} 206 (log ϵ 4.36); 268 (log ϵ 4.36); 329 (sh) (log ϵ 4.56); 341 (log ϵ 4.61); 440 (log ϵ 4.83) (MeOH).


Hosoe, T. *et al.*, *Phytochemistry*, 1999, **52**, 459-463 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Clark, B.R. *et al.*, *Org. Lett.*, 2006, **8**, 701-704 (*isol*)

Avarol

A-1560

[55303-98-5]



Absolute Configuration

C₂₁H₃₀O₂ 314.467Constit. of *Dysidea avara*.

Immunostimulant. Shows antiinflammatory, analgesic, ichthyotoxic, antimutagenic and anti-HIV activity. Cryst. (CHCl₃). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 148-150°. $[\alpha]_D$ +6.1. Log P 6.98 (calc). λ_{\max} 298 (ϵ 3900) (no solvent reported) (Derep).

▶ LD₅₀ (mus, ipr) 269 mg/kg. Antimutagenic activity with inhibiting effect on

benzo[*a*]pyrene monooxygenase.
CZ9026700

2',5'-Quinone, 3'-amino: 3'-Aminoavarone
C₂₁H₂₉NO₂ 327.466
Isol. from a *Dysidea* sp. Red oil. [α]_D²²
+45 (c, 0.1 in CH₂Cl₂).

2',5'-Quinone, 3'-(methylamino): 3'-Methylaminoavarone
[83995-06-6]
C₂₂H₃₁NO₂ 341.492
Found in the sponge *Dysidea avara*.
Inhibitor of cell division in sea urchin eggs. Red cryst. Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 153-155°. Poss. artifact. Called 2-methylamino in the lit. λ_{\max} 288 (ϵ 50029); 486 (ϵ 2088) (MeOH) (Berdy).

2',5'-Quinone, 4'-(methylamino): 4'-Methylaminoavarone
[83995-05-5]
C₂₂H₃₁NO₂ 341.492
Isol. from the sponge *Dysidea avara*.
Inhibitor of cell division in sea urchin eggs. Red cryst. Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 160-163°. Poss. artifact. Called 3-methylamino in the lit. λ_{\max} 288 (ϵ 5030); 486 (ϵ 2090) (MeOH) (Derep). λ_{\max} 289 (ϵ 4688); 485 (ϵ 1065) (MeOH) (Berdy).

2',5'-Quinone, 3'-(2-phenylethylamino): 3'-(2-Phenylethylamino)avarone
C₂₉H₃₇NO₂ 431.617
Isol. from a *Dysidea* sp. Reddish oil. [α]_D²² -89 (c, 0.1 in CH₂Cl₂).


Cimino, G. *et al.*, *Experientia*, 1982, **38**, 896 (*Methylaminoavarones*)

Puliti, R. *et al.*, *J. Mol. Struct.*, 2000, **516**, 31-41 (*3'-Methylaminoavarone, cryst struct*)

Diaz-Marrero, A.R. *et al.*, *Org. Lett.*, 2006, **8**, 3749-3752 (*3'-Aminoavarone, 3'-Phenylethylaminoavarone*)

Avenalumin II A-1561

2-[2-(3-Methoxyphenyl)ethenyl]-4H-3,1-benzoxazin-4-one, 9CI. 2-(3-Methoxycinnamoyl)-4H-3,1-benzoxazin-4-one
[78214-14-9]



C₁₇H₁₃NO₃ 279.295

The proposed struct. illus. has been questioned (see Crombie *et al.*). Phytoalexin from oat (*Avena sativa*) infected with rust fungus *Puccinia coronata*. Poorly sol. hexane. λ_{\max} 317; 336 (MeOH) (Berdy).


Mayama, S. *et al.*, *Physiol. Plant Pathol.*, 1981, **19**, 217 (*isol*)

Collins, F.W. *et al.*, *Spectroscopy (Ottawa)*, 1985, **4**, 171 (*synth, ms*)

Crombie, L. *et al.*, *Tet. Lett.*, 1990, **31**, 2647 (*synth*)

Avenalumin III A-1562

2-[4-(4-Hydroxyphenyl)-1,3-butadienyl]-4H-3,1-benzoxazin-4-one, 9CI
[78214-15-0]



C₁₈H₁₃NO₃ 291.306


The proposed struct. illus. has been questioned (see Crombie *et al.*). Phytoalexin isol. from oat (*Avena sativa*) infected with rust fungus (*Puccinia coronata*). Poorly sol. hexane.

Mayama, S. *et al.*, *Physiol. Plant Pathol.*, 1981, **19**, 217 (*isol*)

Crombie, L. *et al.*, *Tet. Lett.*, 1990, **31**, 2647 (*synth*)

Avenanthramide D A-1563

N-p-Hydroxycinnamoylanthranilic acid. Dianthramide P
[115610-36-1]



C₁₆H₁₃NO₄ 283.283

Sol. EtOAc, Et₂O, H₂O-Me₂CO; poorly sol. CHCl₃, H₂O, C₆H₆. λ_{\max} 218 (ϵ 19950); 329 (ϵ 29500) (MeOH) (Berdy). λ_{\max} 213 (ϵ 26900); 314 (ϵ 12020); 371 (ϵ 32360) (MeOH/NaOH) (Berdy).

(E)-form

Isol. from the oat *Avena sativa* and *Dianthus caryophyllus*. Rods (Me₂CO aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 219°. λ_{\max} 300; 329 (MeOH) (Berdy).

3'-Methoxy: Avenanthramide E. N-Feruloylanthranilic acid
[93755-77-2]

C₁₇H₁₅NO₅ 313.309

Isol. from the oat *Avena sativa*. Pale yellow needles (Me₂CO aq.). Sol. EtOAc, Et₂O, H₂O-Me₂CO; poorly sol. CHCl₃, H₂O, C₆H₆. Mp 235°. λ_{\max} 211; 338 (MeOH) (Berdy). λ_{\max} 219; 385 (MeOH/NaOH) (Berdy).

(Z)-form

Isol. from *Avena sativa* (oat) and *Dianthus caryophyllus*.

3'-Methoxy: Z-Avenanthramide E

[116764-24-0]

C₁₇H₁₅NO₅ 313.309

Isol. from oat (*Avena sativa*).

Collins, F.W. *et al.*, *J. Chromatogr.*, 1988, **445**, 363 (*hplc*)

Ponchet, M. *et al.*, *Phytochemistry*, 1988, **27**, 725


Collins, F.W. *et al.*, *J. Agric. Food Chem.*, 1989, **37**, 60 (*isol, pmr, cmr, uv, ms*)

Crombie, L. *et al.*, *Tet. Lett.*, 1990, **31**, 2647 (*synth*)

Ishihara, A. *et al.*, *Phytochemistry*, 1999, **50**, 237-242 (*biosynth*)

Avenanthramide L A-1564

[172549-38-1]




C₁₈H₁₅NO₅ 325.32

Isol. from oat, *Avena sativa*, inoculated with *Puccinia coronata* f.sp. *avenae*. Phytoalexin. Yellow cryst. Mp 267-269°. λ_{\max} 355 (log ϵ 4.8) (MeOH).

Miyagawa, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 2305-2306 (*isol, uv, pmr*)

Ishihara, A. *et al.*, *Phytochemistry*, 1999, **50**, 237-242 (*biosynth*)

Avicennamine A-1565




C₃₂H₄₈N₂O₃ 508.743

Alkaloid from *Zanthoxylum avicennae*. Amorph. [α]_D²⁸ -52.9 (c, 1 in CHCl₃).

Thuy, T.T. *et al.*, *Phytochemistry*, 1999, **50**, 903-907 (*isol, cd, pmr, cmr, ms*)

Avicine A-1566

5-Methyl-1,3-benzodioxolo[5,6-c][1,3]dioxolo[4,5-j]phenanthridinium(1+), 9CI
[24939-31-9]



C₂₀H₁₄NO₄⁺ 332.335

Alkaloid from the root bark of *Zanthoxylum avicennae* and the bark of *Zanthoxylum inerme* (= *Fagara boninensis*) (Rutaceae). Rather unstable; disproportionates into its dihydro- and oxy-derivs.

Acetate:

Bright yellow needles + 2H₂O (EtOH). Mp 160° dec.

N-De-Me: Des-N-methylavicine. Noravicine

[217-52-7]

C₁₉H₁₁NO₄ 317.3

Alkaloid from the bark of
Zanthoxylum cuspidatum (Rutaceae).
Needles (Py). Mp 325° dec. (290-
295°).


8-Methoxy, N-de-Me: Rhoifoline AC₂₀H₁₃NO₅ 347.326

Alkaloid from the stem bark of
Zanthoxylum rhoifolium. Solid. Mp
168-169°.

Arthur, H.R. *et al.*, *J.C.S.*, 1959, 4007 (*isol*)Gopinath, K.W. *et al.*, *Tetrahedron*, 1961, **14**,
322 (*synth, uv, deriv*)Ishii, H. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 118;
1976, **96**, 1458; *CA*, **77**, 16530y; **86**, 136297k
(*isol, pmr, deriv*)Kessar, S.V. *et al.*, *Tet. Lett.*, 1974, 2269
(*synth*)Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**,
4139 (*synth*)Green, G.R. *et al.*, *J.C.S. Perkin I*, 1996, 1647
(*synth*)Gonzaga, W. de A. *et al.*, *Planta Med.*, 2003,
69, 371-374 (*Rhoifoline A*)**Avinosol**

A-1567

[909576-71-2]


C₃₁H₄₀N₄O₆ 564.68

Conjugate of Avarol, A-1560 and 2'-
Deoxyinosine. Isol. from a *Dysidea* sp.
Unstable solid. [α]_D²² -26.9 (c, 0.35 in
MeOH).

Diaz-Marrero, A.R. *et al.*, *Org. Lett.*, 2006, **8**,
3749-3752 (*isol, pmr, cmr*)**Awaine**

A-1568

**3-Phenylpropanoic acid 2,3-didehydro-4-
hydroxypiperidide**
[593254-86-5]


C₁₄H₁₇NO₂ 231.294

Alkaloid from the aerial parts of *Piper
methysticum*. Oil.

Dragull, K. *et al.*, *Phytochemistry*, 2003, **63**,
193-198 (*isol, pmr, cmr*)**Awajanomycin**

A-1569

[913260-20-5]


C₁₇H₂₇NO₅ 325.404

Prod. by a marine-derived *Acremonium*
sp. AWA16-1. Cytotoxic. Gum. [α]_D²⁵ +78
(c, 0.1 in MeOH). λ_{max} 274 (log ε 3.73)
(MeOH).

Jang, J.-H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1358-
1360 (*isol, pmr, cmr*)**Axamide 1**

A-1570

[56012-88-5]

C₁₆H₂₇NO 249.395

Constit. of *Axinella cannabina*. Oil. [α]_D
+10 (c, 1.2 in CHCl₃).

7,11-Didehydro: Axamide 4

[62078-09-5]

C₁₆H₂₅NO 247.38

Constit. of *Axinella cannabina*. Cryst.
Mp 81-84°. [α]_D +63.3.

Isothiocyanate: Axisothiocyante 1

[53822-97-2]

C₁₆H₂₅NS 263.446

Isol. from *Axinella cannabina*. Oil. [α]_D
+5.9 (c, 2.5 in CHCl₃). Has -NCS
replacing -NHCHO.

Isocyanide: Axisonitrile 1

[53822-96-1]

C₁₆H₂₅N 231.38

From *Axinella cannabina* and *Phyllidia
pultzeri*. Ichthyotoxic agent. Cryst.
Mp 43-45°. [α]_D +22.6 (c, 1 in CHCl₃).
Has -NC replacing -NHCHO.

7,11-Didehydro, isocyanide: Axisonitrile 4

[62078-10-8]

C₁₆H₂₃N 229.364

Isol. from *Axinella cannabina*. Cryst.
Mp 56-58°. [α]_D +51.4 (c, 1 in CHCl₃).

**7,11-Didehydro, isothiocyante: Axi-
sothiocyante 4**


[62078-11-9]

C₁₆H₂₃NS 261.43

Isol. from *Axinella cannabina*. Oil. [α]_D
-35.9 (c, 1.2 in CHCl₃).

Cafieri, F. *et al.*, *Tetrahedron*, 1973, **29**, 4259
(*isol*)Fattorusso, E. *et al.*, *Tetrahedron*, 1975, **31**,
269 (*isol*)Iengo, A. *et al.*, *Experientia*, 1977, **33**, 11;
1979, **35**, 10 (*isol, struct*)Adinolfi, M. *et al.*, *Tet. Lett.*, 1977, 2815
(*struct*)Piers, E. *et al.*, *Can. J. Chem.*, 1986, **64**, 2475
(*synth*)Piers, E. *et al.*, *Tetrahedron*, 1987, **43**, 5521
(*synth*)Chenera, B. *et al.*, *J.O.C.*, 1992, **57**, 2018
(*synth*)Ohkubo, T. *et al.*, *Tet. Lett.*, 1995, **36**, 3365
(*synth*)Guevel, A.-C. *et al.*, *J.O.C.*, 1996, **61**, 473
(*synth*)**Axamide 2**

A-1571

C₁₆H₂₇NO 249.395**(+)-form [56012-89-6]**

Constit. of *Axinella cannabina*. Oil. [α]_D
+37.5 (c, 0.9 in CHCl₃).

Isocyanide: Axisonitrile 2

[55907-33-0]

C₁₆H₂₅N 231.38

From *Axinella cannabina*, *Phyllidia
pustulosa* and *Cadlina luteomarginata*.
Oil. [α]_D +29 (c, 0.5 in CHCl₃). Has -
NC relacing -NHCHO.

Isothiocyanate: Axisothiocyante 2

[56012-90-9]

C₁₆H₂₅NS 263.446

Isol. from *Axinella cannabina*. Oil. [α]_D
+12.8 (c, 1.5 in CHCl₃). Has -NCS
replacing -NHCHO.

**1-Epimer: 10α-Formamidoallovomaden-
drane**

[108739-41-9]

C₁₆H₂₇NO 249.395

Constit. of *Axinella cannabina*. Oil.

**1-Epimer, isocyanide: 10α-Isocyanooal-
loaromadendrane**

[108739-39-5]

C₁₆H₂₅N 231.38

Isol. from *Axinella cannabina*. Oil. [α]_D
-17.21 (c, 0.7 in CHCl₃).

**1-Epimer, isothiocyante: 10α-Isothiocya-
natoallovomadenrane**

[108739-40-8]

C₁₆H₂₅NS 263.446

From *Axinella cannabina*. Oil. [α]_D -
6.95 (c, 0.9 in CHCl₃).


(-)-form**1-Epimer, isothiocyante:**C₁₆H₂₅NS 263.446

Constit. of *Acanthella cavernosa*. Oil.
[α]_D +8 (c, 0.1 in CHCl₃). Not indexed
by CAS.

Fattorusso, E. *et al.*, *Tetrahedron*, 1974, **30**,
3911-3913; 1975, **31**, 269-270 (*Axisonitrile 2*,
Axisothiocyante 2, *Axamide 2*)Ciminiello, P. *et al.*, *Can. J. Chem.*, 1987, **65**,
518-522 (*Axinella cannabina* constits)Fusetani, N. *et al.*, *Tet. Lett.*, 1991, **32**, 7291-
7294 (*Axisonitrile 2*)Da Silva, C.C. *et al.*, *J.O.C.*, 1994, **59**, 2880-
2881 (*synth*)Hirota, H. *et al.*, *Tetrahedron*, 1996, **52**, 2359
(*isol, pmr, cmr*)

Axillarine†

[19637-66-2]

C₁₈H₂₇NO₇ 369.414

Cyclic ester of retronecine. Alkaloid from the seed of *Crotalaria axillaris*. Major alkaloid from *Crotalaria scassellatii* (Fabaceae). Cryst. (MeOH/CHCl₃). Mp 205° dec. [α]_D²⁰ +65.1 (c, 0.82 in Py). The config. of the isopropyl group is not clear from the lit. and may be the opposite of that shown.

Hydrochloride:

Needles (MeOH). Mp 228°.

Picrate: Mp 214-216° dec.**1'-Deoxy: Axillaridine**

[23506-96-9]

C₁₈H₂₇NO₆ 353.414

From *Crotalaria axillaris* and *Crotalaria scassellatii* (Fabaceae). Cryst. (H₂O). Mp 148-152° dec. [α]₄₀₀²¹ +241 (c, 0.12 in MeOH).

1'-Deoxy, picrate:

Silky yellow needles. Mp 235° dec.

12-Deoxy: 12-Deoxyaxillarine. 12-Desoxyaxillarine

[96400-47-4]

C₁₈H₂₇NO₆ 353.414

Minor alkaloid from seeds of *Crotalaria scassellatii* (Fabaceae). [α]_D²⁰ -5 (CHCl₃).

Stereoisomer, 1'-deoxy: Othonnine

[119565-25-2]

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio othonnae* (Asteraceae). Partial stereochem. known.

Crout, D.H.G. *et al.*, *J.C.S.(C)*, 1969, 1379

(isol, ms, struct)

Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653

(cd)

Stoekli-Evans, H. *et al.*, *Helv. Chim. Acta*,1976, **59**, 2168 (cryst struct)Weidenfeld, H. *et al.*, *Phytochemistry*, 1985, **24**,

376 (12-Desoxyaxillarine)

Sener, B. *et al.*, *CA*, 1988, **110**, 132140v


(Othonnine)

Axinellamide

A-1573

1,5-Dihydro-5-hydroxy-5-(6-methyl-2,4-octadienyl)-2H-pyrrol-2-one, 9CI. 5-Hydroxy-5-(6-methyl-2,4-octadienyl)-3-pyrrolin-2-one

[167425-78-7]

C₁₃H₁₉NO₂ 221.299

Alkaloid from the marine sponge *Axinella* sp. Pale yellow gum. [α]_D -26.2 (c,

0.08 in CHCl₃). λ_{max} 277 (ε 8700)


(MeOH) (Berdy).

Miller, S.L. *et al.*, *Tet. Lett.*, 1995, **36**, 5851

(isol, uv, ir, pmr, cmr, struct)

Axinellamine A†

[221225-27-0]



A-1574

C₂₂H₂₃Br₄ClN₁₀O₄ 846.557

CAS nos. refer to bis(trifluoroacetates).

Alkaloid from the Australian sponge *Axinella* sp. Powder (as bistrifluoroacetate salt). [α]_D²⁰ -18 (c, 0.16 in MeOH). λ_{max} 277 (ε 11400) (MeOH).

1-Me ether: Axinellamine C

[221226-05-7]

C₂₃H₂₅Br₄ClN₁₀O₄ 860.584

Alkaloid from an *Axinella* sp. Pale yellow oil. [α]_D²⁰ -9 (c, 1.1 in MeOH). λ_{max} 277 (ε 6500) (MeOH).

5,9-Diepimer: Axinellamine B†

[221226-09-1]

C₂₂H₂₃Br₄ClN₁₀O₄ 846.557

Alkaloid from an *Axinella* sp. Pale yellow oil. [α]_D²⁰ -7 (c, 0.21 in MeOH). λ_{max} 277 (ε 8000) (MeOH).

5,9-Diepimer, 1-Me ether: Axinellamine D


[221226-07-9]

C₂₃H₂₅Br₄ClN₁₀O₄ 860.584

Alkaloid from an *Axinella* sp. Pale yellow oil. [α]_D²⁰ -6 (c, 0.5 in MeOH). λ_{max} 276 (ε 8000) (MeOH).

Urban, S. *et al.*, *J.O.C.*, 1999, **64**, 731-735 (isol, uv, ir, pmr, cmr, ms)Dransfield, P.J. *et al.*, *Tetrahedron*, 2006, **62**, 5223-5247 (synth)O'Malley, D.P. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 3581-3583 (synth)**Axinellamine B†**

[213269-07-9]



Probable Configuration

C₂₉H₄₂N₂ 418.664


Alkaloid from the marine sponge *Axinella* sp. Pale yellow gum. [α]_D -8.9 (c, 0.2 in CHCl₃). λ_{max} 228 (ε 6500) (MeOH).

Bascombe, K.C. *et al.*, *Heterocycles*, 1998, **48**, 1461-1464 (isol, uv, ir, pmr, cmr, ms)Seki, M. *et al.*, *Eur. J. Org. Chem.*, 2001, 503-506 (config)**Axinohydantoin**

A-1576

5-(2-Bromo-5,6,7,8-tetrahydro-8-oxopyrrolo[2,3-c]azepin-4(1H)-ylidene)-2,4-imidazolidinedione, 9CI

[132160-42-0]

C₁₁H₉BrN₄O₃ 325.121

Alkaloid from the sponges *Axinella* sp. and *Monanchora* sp. Yellow prisms (MeOH). Mp 350°. λ_{max} 264 ; 345 (ε 14400) (MeOH) (Berdy).

Debromo: DebromoaxinohydantoinC₁₁H₁₀N₄O₃ 246.225

Isol. from the Papua New Guinea sponge *Monanchora* sp. Mp 300° dec. λ_{max} 208 (ε 16000); 225 (ε 11200); 255 (ε 15500); 339 (ε 27500) (MeOH) (Berdy).

(Z)-Isomer: Spongiacidin D. (Z)-Axinohydantoin. Fuscin†

[193411-88-0]

C₁₁H₉BrN₄O₃ 325.121

Isol. from the sponges *Phakellia fusca*, *Stylorella aurantium* and *Hymeniacidon* sp. Inhibitor of protein kinase C. Amorph. solid. Mp 221-227° (dec.). There is confusion in CAS w.r.t. the config. of the *Phakellia* isolate. λ_{max} 272 (ε 10000); 310 (ε 4000) (MeOH).

Debromo, (Z)-isomer: Spongiacidin C.**(Z)-Debromoaxinohydantoin**

[193202-57-2]

C₁₁H₁₀N₄O₃ 246.225

Isol. from the sponge *Stylorella aurantium* and *Hymeniacidon* sp. Inhibitor of protein kinase C. Amorph. solid. Mp 213-218° (dec.). λ_{max} 273 (ε 9700); 312 (ε 3800) (MeOH).

[132160-43-1]

Pettit, G.R. *et al.*, *Can. J. Chem.*, 1990, **68**, 1621 (isol, uv, ir, pmr, cryst struct)Fu, X. *et al.*, *Chem. Res. Chin. Univ.*, 1991, **7**, 178; *CA*, **118**, 36148y (Fuscine)Groszek, G. *et al.*, *Annalen*, 1995, **715**

(Debromoaxinohydantoin)

Patil, A.D. *et al.*, *Nat. Prod. Lett.*, 1997, **9**,

201-207 (Z-isomers)

Inaba, K. *et al.*, *J. Nat. Prod.*, 1998, **61**, 693-

695 (Spongiacidins C and D)


Sosa, A.C.B. *et al.*, *J.O.C.*, 2002, **67**, 4498-4500

(synth)

Axinyssimide A

A-1577


[184173-50-0]

C₁₆H₂₄Cl₃NO 352.73

Constit. of an *Axinyssa* sponge and *Phyllidia pustulosa*. Antifouling agent. $[\alpha]_D +1.7$ (c, 0.06 in CHCl_3).

Hirota, H. *et al.*, *Tetrahedron*, 1998, **54**, 13971-13980 (*isol*, *pmr*, *cmr*)

Axinyssimide B A-1578
[184173-51-1]



$\text{C}_{16}\text{H}_{26}\text{Cl}_3\text{NO}_2$ 370.745
Constit. of an *Axinyssa* sponge. Antifouling agent. $[\alpha]_D -22.5$ (c, 0.02 in CHCl_3).


10-Epimer: Axinyssimide C

[217481-01-1]
 $\text{C}_{16}\text{H}_{26}\text{Cl}_3\text{NO}_2$ 370.745

Constit. of an *Axinyssa* sponge. Antifouling agent. $[\alpha]_D +7.3$ (c, 0.015 in CHCl_3).

Hirota, H. *et al.*, *Tetrahedron*, 1998, **54**, 13971-13980 (*isol*, *pmr*, *cmr*)

Azaacridone A A-1579
Azaacridone A
[150050-14-9]




$\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_3$ 308.336

First naturally occurring azaacridone alkaloid. Alkaloid from roots of *Citrus paradisi* (grapefruit) (Rutaceae). Light brown oil.

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 789 (*isol*, *uv*, *ir*, *pmr*, *struct*)

Scopton, A. *et al.*, *Org. Lett.*, 2004, **6**, 3869-3871 (*synth*)

9-Azabicyclo[3.3.1]nonan-3-one, 9CI A-1580
Granatonine. *Norgranatan-3-one*. *Norpseudopelletierine*
[4390-39-0]



$\text{C}_8\text{H}_{13}\text{NO}$ 139.197

Found in pomegranate (*Punica granatum*) bark (Punicaceae). Mp 123°.

► **CL5593500**

Picrate:

Cryst. (MeOH). Mp 216°.

N-(Trifluoroacetyl): [180406-55-7]

$\text{C}_{10}\text{H}_{12}\text{F}_3\text{NO}_2$ 235.206
Cryst. (Et₂O/hexane). Mp 70-73°.

N-Benzoyl: [36146-90-4]

$\text{C}_{15}\text{H}_{17}\text{NO}_2$ 243.305

Oil.

N-(4-Methylbenzenesulfonyl): [180406-57-9]

$\text{C}_{15}\text{H}_{19}\text{NO}_3\text{S}$ 293.386

Cryst. (EtOH). Mp 149-151°.

N-Me: 9-Methyl-9-azabicyclo[3.3.1]nonan-3-one, 9CI. *Granatan-3-one*. *N*-Methylgranatonine. *Pseudopelletierine*. *Ψ*-Pelletierine

[552-70-5]

$\text{C}_9\text{H}_{15}\text{NO}$ 153.224

Found in bark of pomegranate (*Punica granatum*). Prismatic plates (petrol). V. sol. H_2O . Mp 62-64°. Bp 246°. Strong base.

N-Me, *oxime*: [6164-67-6]

$\text{C}_9\text{H}_{16}\text{N}_2\text{O}$ 168.238

Plates (Et₂O). Mp 128-129°.

N-Me, *N*-oxide:

$\text{C}_9\text{H}_{15}\text{NO}_2$ 169.223

Mp 160-162°.

N-Me, *N*-oxide; hydrochloride: Mp 224°.

N-Benzyl:

$\text{C}_{15}\text{H}_{19}\text{NO}$ 229.321

Mp 72°. Bp_{0.005} 115-120°.

Org. Synth., *Coll. Vol.*, 4, 1963, 816 (*synth*, *bibl*)
Chen, C.-Y. *et al.*, *J.C.S.(B)*, 1966, 539 (*conform*)

Guthrie, R.D. *et al.*, *J.C.S.(C)*, 1966, 1207 (*ms*, *N*-Me)

Wiseman, J.R. *et al.*, *J.O.C.*, 1977, **42**, 629 (*synth*, *N*-Me)

Dupeyre, R.M. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1978, 612-620 (*synth*, *pmr*, *N*-benzyl)


Hill, R.K. *et al.*, *Tetrahedron*, 1982, **38**, 1959 (*synth*, *N*-Me)

Neuhofer, H. *et al.*, *Pharmazie*, 1993, **48**, 389-391 (*isol*)

Momose, T. *et al.*, *J.C.S. Perkin I*, 1997, 1307-1313 (*N*-benzyl)

Kirihara, M. *et al.*, *Tetrahedron*, 1999, **55**, 2911-2926 (*derivs*)

8-Azabicyclo[3.2.1]octane-2,3-diol A-1581
2,3-Nortropanediol. *2,3-Dihydroxynortropane*



$\text{C}_7\text{H}_{13}\text{NO}_2$ 143.185

(2R,3R)-form

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D -33.9$ (c, 0.32 in H_2O).

(2S,3R)-form

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D -34$ (c, 0.61 in H_2O).

(2S,3S)-form

Alkaloid from *Calystegia soldanella*. $[\alpha]_D +48.4$ (c, 0.82 in H_2O). Abs. config. not detd. but clearly enantiomeric with the (2R,3R)-form above.

2-Benzoyl: 2-Benzoyloxy-3-hydroxynortropane

[34622-25-8]

$\text{C}_{14}\text{H}_{17}\text{NO}_3$ 247.293

Alkaloid from *Peripentadenia mearsii* (Elaeocarpaceae). Needles (Me₂CO). Mp 187-188°. $[\alpha]_D +68$ (c, 0.27 in CHCl_3).

N-Me: 2,3-Tropanediol

$\text{C}_8\text{H}_{15}\text{NO}_2$ 157.212

Synthetic. Mp 133-135°.

(2R*,3S*)-form

N-Me, 2-O-(1-methyl-2-pyrrolocarbonyl): *Vaccinine A*

[864962-36-7]

$\text{C}_{14}\text{H}_{20}\text{N}_2\text{O}_3$ 264.324

Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. $[\alpha]_D^{20} +37.1$ (c, 0.51 in EtOH). λ_{max} 268 (log ϵ 4.19) (EtOH).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1971, **24**, 2399 (*isol*, *ir*, *ms*, *pmr*, *struct*)


Asano, N. *et al.*, *Phytochemistry*, 2001, **57**, 721-726 (*isol*, *pmr*, *cmr*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192 (*isol*, *pmr*, *cmr*, *abs config*)

Zanolari, B. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1153-1158 (*Vaccinine A*)

8-Azabicyclo[3.2.1]octane-2,6-diol A-1582

2,6-Nortropanediol. *2,6-Dihydroxynortropane*



(2R*,6R*)-form

$\text{C}_7\text{H}_{13}\text{NO}_2$ 143.185

(2R*,6R*)-form

(2β,6β)-form. *Baogongteng C*

[107259-50-7]

Alkaloid from *Erycibe elliptilimba*, *Erycibe hainanensis* and *Erycibe obtusifolia* (Convolvulaceae). Oil.

6-Ac: Baogongteng A

[74239-84-2]

$\text{C}_9\text{H}_{15}\text{NO}_3$ 185.222

Alkaloid from *Erycibe obtusifolia* and *Erycibe hainanensis* (Convolvulaceae). Myotic agent. $[\alpha]_D^{28} -7.21$ (c, 0.97 in H_2O).

(2RS,6RS)-form

6-Ac: [126640-93-5]

2-Benzoyl, 6-Ac: [126640-96-8]

Needles (Me₂CO). Mp 158-159°.

(2RS,6SR)-form

6-Ac: [153222-79-8]

Needles. Mp 125-128°.

(2ξ,6ξ)-form

N-Me: 8-Methyl-8-azabicyclo[3.2.1]octane-2,6-diol. *2,6-Tropanediol*. *2,6-Dihydroxytropone*

[65356-02-7]

$\text{C}_8\text{H}_{15}\text{NO}_2$ 157.212


2,6-Tropanediol of undefined config. was isol. from epigeal parts and roots of *Datura stramonium* (Solanaceae).

[107633-96-5]

- Sharova, E.G. *et al.*, *Khim. Prir. Soedin.*, 1977, 126; *Chem. Nat. Compd. (Engl. Transl.)*, 117 (*isol. N-Me*)
 Yao, T. *et al.*, *Yaouxue Xuebao*, 1981, **16**, 582; *CA*, **96**, 48972c
 Chen, Z. *et al.*, *Zhongcaoyao*, 1986, **17**, 386; *CA*, **106**, 153016s (*2,6-Dihydroxynortropane*)
 Wang, P. *et al.*, *Huaxue Xuebao*, 1989, **47**, 1002 (*isol*)
 Jung, M.E. *et al.*, *J.O.C.*, 1992, **57**, 3528 (*synth. pmr, Baogongteng A*)
 Pei, X. *et al.*, *Heterocycles*, 1993, **36**, 2549 (*synth. ir, pmr*)
 Pham, V.C. *et al.*, *J.O.C.*, 1995, **60**, 8051 (*Baogongteng A, synth*)
 Rezel, S. *et al.*, *Heterocycles*, 1999, **51**, 989-1002 (*Baogongteng A, synth*)
 Zhang, Y. *et al.*, *J.A.C.S.*, 2006, **128**, 465-472 (*Baogongteng A, synth*)

8-Azabicyclo[3.2.1]octane-2,7-diol A-1583

2,7-Dihydroxynortropane. 2,7-Nortropenediol



(2*R**,7*R**)-form

C₇H₁₃NO₂ 143.185

(2*R,7*R**)-form** [357426-27-8]

Alkaloid from *Calystegia* spp. and *Ipomoea* spp. [α]_D²⁵ +13.1 (c, 2.3 in H₂O).

N-Me, 2-O-(1-methyl-2-pyrrolicarboxyl): **Vaccinine B**

[864962-37-8]
 C₁₄H₂₀N₂O₃ 264.324

Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D²⁰ -2 (c, 0.47 in EtOH). λ_{max} 267 (log ε 4.1) (EtOH).

(2*R,7*S**)-form**

Erycibelline

[107633-95-4]

Alkaloid from the stems of *Erycibe elliptimba*.

Lu, Y. *et al.*, *Yaouxue Xuebao*, 1986, **21**, 829-835; *CA*, **106**, 153028x (*Erycibelline*)

Asano, N. *et al.*, *Phytochemistry*, 2001, **57**, 721-726 (*isol, pmr, cmr*)

Zanolari, B. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1153-1158 (*Vaccinine B*)

8-Azabicyclo[3.2.1]octane-3,6-diol A-1584

3,6-Dihydroxynortropane. 3,6-Nortropenediol



(3*S*,6*R*)-form

C₇H₁₃NO₂ 143.185

(3*S*,6*R*)-form

Alkaloid from the fruit of *Morus alba* (white mulberry). Powder. [α]_D²⁵ -1.3 (c, 0.6 in H₂O).

(3*R,6*R**)-form**

N-Me: see 8-Methyl-8-azabicyclo[3.2.1]octane-3,6-diol, M-391

3-O-Tigloyl: 3-Tigloyloxynortropan-6-ol

[65636-83-1]
 [65636-84-2]

C₁₂H₁₉NO₃ 225.287

Alkaloid from *Datura sanguinea* (Solanaceae). Plates (EtOH) (as hydrochloride). Mp 289-290° (hydrochloride). Opt. rotn. not measured.

3-O-Benzoyl: 3-Benzoyloxynortropan-6-ol

C₁₄H₁₇NO₃ 247.293

Alkaloid from the leaves of *Erythroxylum macrocarpum* and *Erythroxylum sideroxyloides* (Erythroxylaceae). Mp 225° (as picrate).

6-O-Benzoyl, 3-(2-methylpropanoyl): 6-Benzoyloxy-3-isobutanoyloxynortropane

C₁₈H₂₃NO₄ 317.384

Alkaloid from the leaves of *Erythroxylum moonii*. Light brown powder. [α]_D²⁵ -21.4 (c, 0.07 in MeOH). λ_{max} 232 (log ε 4.07); 275 (log ε 2.87) (MeOH).

3-O-(3,4,5-Trimethoxybenzoyl): 3-(3,4,5-Trimethoxybenzoyloxy)nortropan-6-ol

C₁₇H₂₃NO₆ 337.372

Alkaloid from *Erythroxylum zambeziacum* (Erythroxylaceae). Prisms (EtOH aq.) (as picrate). Mp 201° (picrate).

6-O-E-Cinnamoyl, 3-O-propanoyl: 6-trans-Cinnamoyloxy-3-propanoyloxynortropane

C₁₉H₂₃NO₄ 329.395

Alkaloid from the leaves of *Erythroxylum moonii*. Amorph. solid. [α]_D²⁵ -23.1 (c, 0.31 in MeOH). λ_{max} 280 (log ε 4.37) (MeOH).

6-O-(3,4,5-Trimethoxy-Z-cinnamoyl), 3-O-(2-methylpropanoyl): 3-Isobutanoyloxy-6-(3,4,5-trimethoxy-cis-cinnamoyloxy)nortropane

C₂₃H₃₁NO₇ 433.5

Alkaloid from the leaves of *Erythroxylum moonii*. Semisolid. [α]_D²⁵ -25.2 (c, 0.07 in CHCl₃). λ_{max} 326 (log ε 4.43) (MeOH).

(3*R*,6*S*)-form

3β,6β-Dihydroxynortropane

Alkaloid from the fruit of *Morus alba* (white mulberry). [α]_D²⁵ -8.2 (c, 0.34 in H₂O).

Di-O-tigloyl: 3,6-Ditigloyloxynortropane

[359723-70-9]

C₁₇H₂₅NO₄ 307.389

Alkaloid from the leaves of *Erythroxylum argentinum*.

Evans, W.C. *et al.*, *Phytochemistry*, 1978, **17**, 171 (*3-tiglate*)

Al-Said, M.S. *et al.*, *Phytochemistry*, 1986, **25**, 851-853 (*3-benzoate*)

El-Iman, Y.M.A. *et al.*, *Phytochemistry*, 1987, **26**, 2385-2389 (*3-trimethoxybenzoate*)

Zuanazzi, J.A.S. *et al.*, *Biochem. Syst. Ecol.*, 2001, **29**, 819-825 (*ditiglate*)


Asano, N. *et al.*, *J. Agric. Food Chem.*, 2001, **49**, 4208-4213 (*3β,6β-form*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192 (*Morus alba isolate*)

Khattak, K.F. *et al.*, *Heterocycles*, 2003, **60**, 917-924 (*Erythroxylum moonii esters*)

8-Azabicyclo[3.2.1]octane-3,7-diol A-1585

3,7-Dihydroxynortropane. 3,7-Nortropenediol



C₇H₁₃NO₂ 143.185

(3*R*,7*R*)-form

Alkaloid from *Duboisia leichardtii*.

3-(2-Methylpropanoyl): 3-Isobutyryloxynortropan-7-ol

C₁₁H₁₉NO₃ 213.276

Alkaloid from the leaves of *Erythroxylum moonii*. Amorph. powder. [α]_D²⁵ -13.4 (c, 0.04 in MeOH). λ_{max} 208 (log ε 3.85); 264 (log ε 2.37) (MeOH).

7-Phenylacetyl: 7-Phenylacetoxynortropan-3-ol

C₁₅H₁₉NO₃ 261.32


Alkaloid from the leaves of *Erythroxylum moonii*. Amorph. powder. [α]_D²⁵ +7.2 (c, 0.25 in MeOH). λ_{max} 208 (log ε 3.85); 264 (log ε 2.37) (MeOH).

Asano, N. *et al.*, *Phytochemistry*, 2001, **57**, 721-726 (*isol, pmr, cmr*)

Khattak, K.F. *et al.*, *J. Nat. Prod.*, 2002, **65**, 929-931 (*3-isobutyryl-, 7-phenylacetyl derivs*)

8-Azabicyclo[3.2.1]octane-1,2,3,4,6-pentol A-1586

1,2,3,4,6-Pentahydroxynortropane. 1,2,3,4,6-Nortropanepentol



(1*R**,2*R**,3*R**,4*S**,6*R**)-form

C₇H₁₃NO₅ 191.183

(1*R*,2*R*,3*R*,4*S*,6*R*)-form

Calystegine C₂

[190957-44-9]

Alkaloid from *Duboisia leichardtii* and *Lycium chinense* (Chinese boxthorn). [α]_D²⁵ -40.6 (c, 0.32 in H₂O).

(1*R*,2*S*,3*R*,4*S*,6*R*)-form

Calystegine C₁

[156705-04-3]

Alkaloid from *Morus alba* (white mulberry) and *Lycium chinense* (Chinese boxthorn). Powder. [α]_D²⁵ +23.1 (c, 0.8 in H₂O).

N-Me: N-Methylcalystegine C₁

[197449-07-3]

C₈H₁₅NO₅ 205.21


Alkaloid from the roots of *Lycium chinense* (Chinese boxthorn). [α]_D²⁵ +27.6 (c, 0.3 in H₂O).

Asano, N. *et al.*, *Eur. J. Biochem.*, 1995, **229**, 369-376; 1997, **248**, 296-303 (*isol, pmr, cmr*)

- Kato, A. *et al.*, *Phytochemistry*, 1997, **45**, 425-429 (*Calystegine C₂*)
 Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (*occur*)
 Watson, A.A. *et al.*, *Phytochemistry*, 2001, **56**, 265-295 (*rev*)
 Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (*rev*)
 Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

8-Azabicyclo[3.2.1]octane-1,2,3,4-tetrol A-1587

1,2,3,4-Tetrahydroxynortropane. 1,2,3,4-Nortropanetetrol



(1R,2S,3R,4S)-form

C₇H₁₃NO₄ 175.184

(1R,2S,3R,4S)-form

Calystegine B₂. Nortropanoline
 [127414-85-1]

Alkaloid from *Atropa belladonna*, leaves of *Solanum tuberosum*, *Solanum dulcamara*, *Solanum melongena*, *Solanum dimidiatum*, *Solanum kwebense* and *Datura wrightii* (Solanaceae), *Calystegia arvensis* and *Calystegia sepium* (Convolvulaceae). [α]_D +2.9 (c, 0.2 in H₂O). [α]_D +17.1 (c, 0.41 in H₂O) (synthetic).

N-Me: N-Methylcalystegine B₂

[184045-65-6]

C₈H₁₅NO₄ 189.211

Alkaloid from the roots of *Lycium chinense* (Chinese boxthorn). [α]_D +22.2 (c, 0.22 in H₂O).

(1S,2R,3S,4R)-form [146275-92-5]

Synthetic. [α]_D -17.5 (c, 0.37 in H₂O).

4-O-α-D-Galactopyranoside: 4-O-α-D-Galactopyranosylcalystegine B₂

C₁₃H₂₃NO₉ 337.326

Alkaloid from the fruit of *Morus alba* (white mulberry). [α]_D +114.5 (c, 0.48 in H₂O).

(1R,2R,3R,4S)-form

Calystegine B₃

[178231-95-3]

Alkaloid from *Hyoscyamus niger* and *Physalis alkekengi* var. *francheti* (Solanaceae). Powder. [α]_D +82.8 (c, 0.5 in H₂O).

(1R,2S,3R,4R)-form

Calystegine B₄

[184046-85-3]

Alkaloid from *Scopolia japonica* (Solanaceae). Trehalase inhibitor. Powder. [α]_D -63 (c, 0.65 in H₂O).

Goldmann, A. *et al.*, *Phytochemistry*, 1990, **29**, 2125 (*isol*)

Ducrot, P.-H. *et al.*, *Tet. Lett.*, 1990, **31**, 3879; 3883 (*pmr, cmr, struct, synth*)

Boyer, F.-D. *et al.*, *Synlett*, 1992, 969 (*synth*)

Duclos, O. *et al.*, *Tet. Lett.*, 1992, **33**, 8061 (*synth*)

Nash, R.J. *et al.*, *Phytochemistry*, 1993, **34**, 1281 (*occur*)

Boyer, F.-D. *et al.*, *Tetrahedron*, 1994, **50**, 10443 (*synth*)

Asano, N. *et al.*, *Eur. J. Biochem.*, 1995, **229**, 369 (*Calystegine B₃*)

Asano, N. *et al.*, *Carbohydr. Res.*, 1996, **284**, 169; **293**, 195 (*Calystegine B₃*, *Calystegine B₄*)

Goldmann, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1137

Souli, J. *et al.*, *Tetrahedron*, 1996, **52**, 15137 (*synth*)

Asano, N. *et al.*, *Eur. J. Biochem.*, 1997, **248**, 296-303 (*N-Methylcalystegine B₂*)

Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (*occur*)

Asano, N. *et al.*, *J. Agric. Food Chem.*, 2001, **49**, 4208-4213 (*4-Galactosylcalystegine B₂*)

Watson, A.A. *et al.*, *Phytochemistry*, 2001, **56**, 265-295 (*rev*)

Bekkouche, K. *et al.*, *Phytochemistry*, 2001, **58**, 455-462 (*occur*)

Boyer, F.-D. *et al.*, *Tet. Lett.*, 2001, **42**, 1275-1277 (*Calystegine B₂*, *synth*)

Marco-Contelles, J. *et al.*, *J.O.C.*, 2002, **67**,

3705-3717 (*Calystegine B₂*, *synth*)

Skaanderup, P.R. *et al.*, *J.O.C.*, 2003, **68**, 2115-2122 (*Calystegine B₂*, *B₃*, *B₄*, *synth*)

Scholl, Y. *et al.*, *Phytochemistry*, 2003, **62**, 325-332 (*biosynth*)


Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (*rev*)

Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

Chen, Y.-L. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 3330-3339 (*Calystegine A₃*, *synth*)

8-Azabicyclo[3.2.1]octane-1,2,3,6-tetrol A-1588

1,2,3,6-Tetrahydroxynortropane. 1,2,3,6-Nortropanetetrol



C₇H₁₃NO₄ 175.184

(1S,2R,3S,6S)-form

Calystegine B₁

[127414-86-2]

Alkaloid from *Atropa belladonna* and *Nicandra physalodes* (Solanaceae), *Calystegia arvensis* and *Calystegia sepium* (Convolvulaceae). Powder. [α]_D -13 (c, 0.6 in H₂O).

3-O-β-D-Glucopyranoside: [176678-49-2]

C₁₃H₂₃NO₉ 337.326

Alkaloid from fruits of *Nicandra physalodes*.

Goldmann, A. *et al.*, *Phytochemistry*, 1990, **29**, 2125 (*isol*)

Ducrot, P.-H. *et al.*, *Tet. Lett.*, 1990, **31**, 3879; 3883 (*pmr, cmr, struct, synth*)

Asano, N. *et al.*, *Eur. J. Biochem.*, 1995, **229**, 369 (*isol, pmr, cmr*)

Griffiths, R.C. *et al.*, *Tet. Lett.*, 1996, **37**, 3207 (*glucoside*)

Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (*occur*)

Watson, A.A. *et al.*, *Phytochemistry*, 2001, **56**, 265-295 (*rev*)


Bekkouche, K. *et al.*, *Phytochemistry*, 2001, **58**, 455-462 (*occur*)

Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (*rev*)

Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

8-Azabicyclo[3.2.1]octane-1,2,4,7-tetrol A-1589

1,2,4,7-Tetrahydroxynortropane. 1,2,4,7-Nortropanetetrol



C₇H₁₃NO₄ 175.184

(1R*,2R*,4S*,7S*)-form

Calystegine B₅

[197565-91-6]

Alkaloid from the roots of *Lycium chinense* (Chinese boxthorn). [α]_D +9.6 (c, 0.31 in H₂O).


Asano, N. *et al.*, *Eur. J. Biochem.*, 1997, **248**, 296-303 (*Calystegine B₅*)

Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (*rev*)

Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

8-Azabicyclo[3.2.1]octane-1,2,3-triol A-1590

1,2,3-Trihydroxynortropane. 1,2,3-Nortropanetriol



C₇H₁₃NO₃ 159.185

(1R,2S,3R)-form

Calystegine A₃

[131580-36-4]

Alkaloid from *Atropa belladonna* (Solanaceae), *Calystegia arvensis* and *Calystegia sepium* (Convolvulaceae) and *Solanum tuberosum* (potato) (Solanaceae). Powder. [α]_D -17.3 (c, 0.5 in H₂O).

Goldmann, A. *et al.*, *Phytochemistry*, 1990, **29**, 2125 (*isol*)

Ducrot, P.-H. *et al.*, *Tet. Lett.*, 1990, **31**, 3879; 3883 (*pmr, cmr, struct, synth*)

Boyer, F.-D. *et al.*, *Synlett*, 1992, 357 (*synth*)

Nash, R.J. *et al.*, *Phytochemistry*, 1993, **34**, 1281 (*occur*)

Asano, N. *et al.*, *Eur. J. Biochem.*, 1995, **229**, 369 (*isol, pmr, cmr*)

Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (*occur*)


Bekkouche, K. *et al.*, *Phytochemistry*, 2001, **58**, 455-462 (*occur*)

Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (*rev*)

Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (*rev*)

8-Azabicyclo[3.2.1]octane-1,2,4-triol


1,2,4-Trihydroxynortropane. 1,2,4-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1R*,2S*,4S*)-form****Calystegine A₇**

[197565-90-5]

Alkaloid from the roots of *Lycium chinense* (Chinese boxthorn). [α]_D -10.8 (c, 0.27 in H₂O).Asano, N. *et al.*, *Eur. J. Biochem.*, 1997, **248**, 296-303 (isol)Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (rev)Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (rev)Csuk, R. *et al.*, *Tetrahedron*, 2008, **64**, 9417-9422 (synth)**8-Azabicyclo[3.2.1]octane-1,2,7-triol**


1,2,7-Trihydroxynortropane. 1,2,7-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1R*,2R*,7R*)-form****Calystegine A₆**

[177794-04-6]

Alkaloid from *Hyoscyamus niger* and *Lycium chinense*. [α]_D -27.6 (c, 0.4 in H₂O).Asano, N. *et al.*, *Carbohydr. Res.*, 1996, **284**, 169-178 (isol, pmr, cmr)Asano, N. *et al.*, *Eur. J. Biochem.*, 1997, **248**, 296-303 (isol)Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (occur)Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (rev)Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (rev)**8-Azabicyclo[3.2.1]octane-1,3,4-triol**


1,3,4-Trihydroxynortropane. 1,3,4-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1R*,3S*,4S*)-form****Calystegine A₅**


[165905-26-0]

Alkaloid from *Hyoscyamus niger* and *Physalis alkekengi* var. *francheti* (Solanaceae). Powder. [α]_D +68.7 (c, 0.6 in H₂O).Asano, N. *et al.*, *Eur. J. Biochem.*, 1995, **229**, 369-376 (isol, pmr, cmr)Asano, N. *et al.*, *Carbohydr. Res.*, 1996, **284**, 169-175 (isol)Schimming, T. *et al.*, *Phytochemistry*, 1998, **49**, 1989-1995 (occur)Dräger, B. *et al.*, *Nat. Prod. Rep.*, 2004, **21**, 211-223 (rev)Biastoff, S. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 49-102 (rev)**8-Azabicyclo[3.2.1]octane-2,3,4-triol**


2,3,4-Trihydroxynortropane. 2,3,4-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1RS,2SR,3RS,4RS,5SR)-form**Alkaloid from the fruit of *Morus alba*. Powder. meso-Stereoisomer.Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192**8-Azabicyclo[3.2.1]octane-2,3,6-triol**

2,3,6-Trihydroxynortropane. 2,3,6-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1S,2R,3R,5S,6R)-form**Alkaloid from the fruit of *Morus alba*. Powder. [α]_D -27.3 (c, 0.55 in H₂O).Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192**8-Azabicyclo[3.2.1]octane-3,6,7-triol**

3,6,7-Trihydroxynortropane. 3,6,7-Nortropanetriol

C₇H₁₃NO₃ 159.185**(1R*,3S*,6S*,7R*)-form**

(3α,6β,7β)-form

3-O-Benzoyl, 6-O-(2ξ-hydroxy-3-phenylpropanoyl): 3-Benzoyloxy-6-hydroxy-7-(2-hydroxy-3-phenylpropanoyloxy)nortropane

C₂₃H₂₅NO₆ 411.454Alkaloid from the leaves of *Erythroxylum moonii*. Semisolid. [α]_D²⁵ +17.6 (c, 0.13 in CHCl₃). Chiral by asymmetric substitution but the abs. config. is not known. λ_{max} 209 (log ε 3.93); 230 (log ε 3.99); 266 (log ε 2.41); 276 (log ε 2.91) (MeOH).

N-Me: see 8-Methyl-8-azabicyclo[3.2.1]octane-3,6,7-triol, M-392

Khattak, K.F. *et al.*, *Heterocycles*, 2003, **60**, 917-924 (isol, pmr, cmr, ms)**8-Azabicyclo[3.2.1]octan-3-ol, 9CI**

3-Hydroxynortropane. 3-Nortropanol



(1RS,3RS,5SR)-form

C₇H₁₃NO 127.186**(1RS,3SR,5SR)-form**

exo-form. Nor-ψ-tropine. Norpseudotropine

[501-33-7]

Alkaloid from the fruit of *Morus alba* (white mulberry). Cryst. (C₆H₆/pentane). Mp 134.5-135° (129-130°).▶ LD₅₀ (mus, orl) 227 mg/kg. RD0875000

Hydrochloride: [14383-51-8]

Cryst. Mp 280-282° dec.

▶ RD0600000

N-Benzoyl: Nortropacocaine

[18470-33-2]

C₁₄H₁₇NO₂ 231.294Alkaloid in *Erythroxylum mamacoca*

(Erythroxylaceae). Feathery cryst.

(synthetic, as picrate). Mp 234° (picrate).

N-Me: see 8-Methyl-8-azabicyclo[3.2.1]octan-3-ol, M-393

(1RS,3RS,5SR)-form

endo-form. Nortropeine. Tropigenin. Nortropenol. Nortropine

[538-09-0]

▶ LD₅₀ (mus, orl) 1420 mg/kg. RD0700000

Hydrochloride: [17366-48-2]

Cryst. Mp 200°.

O-(2-Methylbutanoyl): Isoporoidine

[537-28-0]

C₁₂H₂₁NO₂ 211.303Alkaloid from *Duboisia myoporoides* and *Erythroxylum deckindtii* (Solanaceae, Erythroxylaceae).

O-(3-Methylbutanoyl): Poroidine

[500-58-3]

C₁₂H₂₁NO₂ 211.303Alkaloid from *Duboisia myoporoides* and root-bark of *Erythroxylum deckindtii* (Solanaceae, Erythroxylaceae).

Co-isol. with Isoporoidine, difficult to separate.

O-Benzoyl: 3 α -Benzoyloxynortropane

C₁₄H₁₇NO₂ 231.294

Alkaloid from leaves, root bark and stem bark of *Erythroxylum macrocarpum* and *Erythroxylum sideroxyloides* (Erythroxylaceae). Mp 232° (as picrate).

O-(4-Methoxybenzoyl): Merresectine A

C₁₅H₁₉NO₃ 261.32

Alkaloid from *Merremia dissecta* and *Merremia quinquefolia*. Yellow solid.

O-Phenylacetyl: Nortropan-3-yl phenylacetate

C₁₅H₁₉NO₂ 245.321

Minor alkaloid from the root bark of *Erythroxylum hypericifolium* (Erythroxylaceae).

O-Cinnamoyl (E-): Nortropinyl cinnamate

[126394-79-4]

C₁₆H₁₉NO₂ 257.332

Alkaloid from *Pellacalyx axillaris* (Rhizophoraceae). Yellow gum.

O-Cinnamoyl, N-Ac: [136945-63-6]

Needles (CH₂Cl₂/petrol). Mp 143°.

N-Me: see 8-Methyl-8-azabicyclo[3.2.1]octan-3-ol, M-393

Barger, G. et al., *J.C.S.*, 1937, 1820 (*Poroidine, Isoporoidine*)

Werner, G. et al., *Annalen*, 1967, **708**, 210 (*synth*)

Kraiss, G. et al., *Tet. Lett.*, 1971, 57 (*synth*)
Chappell, G.S. et al., *J. Pharm. Sci.*, 1973, **62**, 414 (*nmr*)

Tufariello, J.J. et al., *J.A.C.S.*, 1979, **101**, 2435 (*synth*)

Al-Yahya, M.A.I. et al., *J.C.S. Perkin 1*, 1979, 2130 (*isol, Poroidine, Isoporoidine*)

El-Imam, Y.M.A. et al., *Phytochemistry*, 1985, **24**, 2285-2289 (*Nortropacocaine*)

Al-Said, M.S. et al., *J.C.S. Perkin 1*, 1986, 957 (*Phenylacetate*)

Al-Said, M.S. et al., *Phytochemistry*, 1986, **25**, 851-853 (*isol, ir, pmr, ms, struct, deriv*)


Arbain, D. et al., *Aust. J. Chem.*, 1991, **44**, 1013 (*Nortropinyl cinnamate*)

Kusano, G. et al., *Chem. Pharm. Bull.*, 2002, **50**, 185-192 (*isol, pmr, cmr*)

Jenett-Siems, K. et al., *Phytochemistry*, 2005, **66**, 1448-1464 (*Merresectine A*)

Azacyclo-6-undecene, 9CI

A-1598



C₁₀H₁₉N 153.267

(Z)-form Keramaphidin C

[126412-12-2]

Isol. from the Okinawan marine sponge *Amphimedon* sp. Plausible biogenetic precursor of Manzamine C, M-93. Amorph. solid. Mp 106-109°.

Torisawa, Y. et al., *Tetrahedron*, 1991, **47**, 8067 (*synth*)


Tsuda, M. et al., *Tet. Lett.*, 1994, **35**, 4387 (*isol, ir, pmr, cmr*)

Tsuda, M. et al., *Heterocycles*, 1997, **46**, 765-794 (*rev*)

Azamerone

A-1599

[897385-39-6]



Absolute Configuration

C₂₅H₃₂Cl₂N₂O₅ 511.444


Isol. from the marine-derived *Streptomyces* sp. CNQ766. Cryst. Mp 210-212°. [α]_D -8.8 (c, 0.002 in MeOH). λ_{max} 217 (sh) (log ϵ 3.9); 250 (sh) (log ϵ 3.4); 323 (log ϵ 3.3) (MeOH).

Cho, J.Y. et al., *Org. Lett.*, 2006, **8**, 2471-2474 (*isol, pmr, cmr, cryst struct*)

Azaspiracid 1

A-1600

Killarytoxin 3
[214899-21-5]



Absolute Configuration

C₄₇H₇₁NO₁₂ 842.077

Struct. finally confirmed in 2004. There are various errors in the assignment of CAS numbers to the minor congeners. Alkaloid from *Mytilus edulis* (blue mussel). Shellfish toxin. Amorph. solid. [α]_D²⁰ -21 (c, 0.1 in MeOH).

3R-Hydroxy: Azaspiracid 7

[629656-50-4]

C₄₇H₇₁NO₁₃ 858.077

Alkaloid from *Mytilus edulis* (blue mussel).

23R-Hydroxy: Azaspiracid 8

[629656-53-7]

C₄₇H₇₁NO₁₃ 858.077

Alkaloid from *Mytilus edulis* (blue mussel).

22-Demethyl: Azaspiracid 3

[265996-93-8]

C₄₆H₆₉NO₁₂ 828.051

Alkaloid from *Mytilus edulis* (blue mussel).

22-Demethyl, 3R-hydroxy: Azaspiracid 4

[344422-49-7]

C₄₆H₆₉NO₁₃ 844.05

Alkaloid from *Mytilus edulis* (blue mussel).

22-Demethyl, 23S-hydroxy: Azaspiracid 5

[344422-51-1]

C₄₆H₆₉NO₁₃ 844.05

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl: Azaspiracid 2

[265996-92-7]

C₄₈H₇₃NO₁₂ 856.104

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl, 3R-hydroxy: Azaspiracid 11

[629656-62-8]

C₄₈H₇₃NO₁₃ 872.104

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl, 23R-hydroxy: Azaspiracid 12

[629656-65-1]

C₄₈H₇₃NO₁₃ 872.104

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl, 22-demethyl: Azaspiracid 6

[629656-47-9]

C₄₇H₇₁NO₁₂ 842.077

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl, 22-demethyl, 3R-hydroxy: Azaspiracid 9

[629656-56-0]

C₄₇H₇₁NO₁₃ 858.077

Alkaloid from *Mytilus edulis* (blue mussel).

8-Methyl, 22-demethyl, 23R-hydroxy: Azaspiracid 10

[629656-59-3]

C₄₇H₇₁NO₁₃ 858.077

Alkaloid from *Mytilus edulis* (blue mussel). Cahn-Ingold-Prelog priorities change at C-23.

Satake, M. et al., J.A.C.S., 1998, **120, 9967-9968 (*Azaspiracid 1*)**

Ofuji, K. et al., *Nat. Toxins*, 1999, **7**, 99-102 (*Azaspiracids 2,3*)

Food Sci. Technol., Seafood and Freshwater Toxins, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**,

Ofuji, K. et al., *Biosci., Biotechnol., Biochem.*, 2001, **65**, 740-742 (*Azaspiracids 4,5*)

James, K.J. et al., *Toxicol.*, 2003, **41**, 277-283 (*isol, ms*)

Lehane, M. et al., *J. Chromatogr., A*, 2004, **1024**, 63-70 (*hplc-ms*)

Nicolaou, K.C. et al., *Chem. Asian J.*, 2006, **1**, 245-263 (*synth*)

Nicolaou, K.C. et al., *J.A.C.S.*, 2006, **128**, 2244-2257; 2258-2267; 2859-2872 (*synth, struct*)

Evans, D.A. et al., *J.A.C.S.*, 2008, **130**, 16295-16309 (*synth*)


Twiner, M.J. et al., *Mar. Drugs*, 2008, **6**, 39-72 (*rev*)

Rehmann, N. et al., *Rapid Commun. Mass Spectrom.*, 2008, **22**, 549-558 (*isol, ms*)

Azaspirene

A-1601

[461644-34-8]



Absolute Configuration

C₂₁H₂₃NO₅ 369.416

Related to Pseurotins, P-738 and Synerazol, S-660. Prod. by the fungus *Neosartorya* sp. Angiogenesis inhibitor. Pale

yellow powder. Mp 166-167°. $[\alpha]_D^{25}$ -204.4 (c, 0.16 in MeOH). λ_{\max} 231 (ε 5900); 341 (ε 27860) (MeOH).


Hayashi, Y. *et al.*, *J.A.C.S.*, 2002, **124**, 12078-12079 (*synth, abs config*)

Asami, Y. *et al.*, *Org. Lett.*, 2002, **4**, 2845-2848 (*isol, pmr, cmr*)

Aoki, S. *et al.*, *Heterocycles*, 2004, **62**, 161-166 (*synth, pmr, cmr*)

15-Azasterol**A-1602**

15-Aza-D-homoergosta-8,14,24(28)-trien-3-ol, 9CI. A 25822B. Antibiotic A 25822B. *Geotrichum alkaloid* A 25822B. UCA 1064A. Antibiotic UCA 1064A [50686-98-1]



$C_{28}H_{45}NO$ 411.67

Main alkaloid from cultures of *Geotrichum flavo-brunneum*. Also prod. by *Wallemia sebi*. Antifungal agent, active against *Candida* and *Trichophyton* spp. Sol. MeOH, $CHCl_3$; fairly sol. C_6H_6 ; poorly sol. H_2O , hexane. Mp 115-118°. $[\alpha]_D^{25}$ -20 (c, 0.78 in MeOH). λ_{\max} 277 (ε 13400) (EtOH/HCl) (Derep). λ_{\max} 240 (ε 12300); 270 (sh) (EtOH) (Derep). λ_{\max} 241; 270 (MeOH) (Berdy). λ_{\max} 238 (ε 12300) (EtOH) (Berdy). λ_{\max} 277 (ε 13400) (EtOH-HCl) (Berdy). λ_{\max} 279 (MeOH-HCl) (Berdy). λ_{\max} 241 (MeOH-NaOH) (Berdy).

Ac: *Geotrichum alkaloid* A 25822M. A 25822M. Antibiotic A 25822M [50886-40-3]

$C_{30}H_{47}NO_2$ 453.707

Prod. by *Geotrichum flavo-brunneum*. Yellowish amorph. powder. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ -15 (c, 0.021 in MeOH). pK_a 7.9 (66% DMF aq.). λ_{\max} 278 (ε 14200) (EtOH/HCl) (Derep). λ_{\max} 239 (ε 13900) (EtOH/NaOH) (Derep). λ_{\max} 239 (ε 13900) (EtOH) (Berdy). λ_{\max} 278 (ε 14200) (EtOH-HCl) (Berdy).

3-Ketone: *Geotrichum alkaloid* A 25822N. A 25822N. Antibiotic A 25822N [50687-00-8]

$C_{28}H_{43}NO$ 409.654

Prod. by *Geotrichum flavo-brunneum*. Microcryst. solid (Me_2CO). Sol. MeOH, Et_2O ; fairly sol. C_6H_6 , hexane; poorly sol. H_2O . Mp 165°. $[\alpha]_D^{25}$ -14 (c, 0.05 in MeOH). pK_a 7.45 (66% in DMF aq.). λ_{\max} 277 (ε 13400) (EtOH/HCl) (Derep). λ_{\max} 240 (ε 12300); 270 (sh) (EtOH) (Derep). λ_{\max} 239 (ε 13900) (EtOH) (Berdy). λ_{\max} 278 (ε 14100) (EtOH-HCl) (Berdy).

24,28-Dihydro: 24,28-Dihydro-15-azasterol. UCA 1064B. Antibiotic UCA 1064B $C_{28}H_{47}NO$ 413.685

Prod. by *Wallemia sebi*. Antitumour agent. Powder. $[\alpha]_D^{24}$ -28.5 (c, 0.5 in MeOH). λ_{\max} 241; 270 (MeOH) (Berdy). λ_{\max} 279 (MeOH-HCl) (Berdy). λ_{\max} 241 (MeOH-NaOH) (Berdy).

16ξ-Hydroxy: *Geotrichum alkaloid* A 25822D. Antibiotic A 25822D. A 25822D

[50687-01-9]

$C_{28}H_{45}NO_2$ 427.669

Minor metab. of *Geotrichum flavo-brunneum*. Shows antifungal props. Amorph. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ +39 (c, 0.722 in MeOH). pK_a 7.7 (66% DMF aq.). Indefinite Mp. λ_{\max} 277 (ε 13400) (EtOH/HCl) (Derep). λ_{\max} 240 (ε 12300); 270 (sh) (EtOH) (Derep). λ_{\max} 235 (ε 10400) (EtOH) (Berdy). λ_{\max} 270 (ε 8600) (EtOH-HCl) (Berdy).

16-Oxo: *Geotrichum alkaloid* A 25822L.

Antibiotic A 25822L. A 25822L

[50687-02-0]

$C_{28}H_{43}NO_2$ 425.653

Minor metab. of *Geotrichum flavo-brunneum*. Shows antifungal props. Amorph. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ +75 (c, 0.072 in MeOH). pK_a 4.75 (66% DMF aq.). λ_{\max} 278 (ε 10800) (EtOH/HCl) (Derep). λ_{\max} 262 (ε 10200) (EtOH/NaOH) (Derep). λ_{\max} 262 (ε 10200) (EtOH) (Derep). λ_{\max} 262 (ε 10200) (EtOH) (Berdy). λ_{\max} 278 (ε 10800) (EtOH-HCl) (Berdy).

[51517-53-6, 74608-58-5]

Chamberlin, J.W. *et al.*, *J. Antibiot.*, 1974, **27**, 992; 1975, **28**, 102 (*isol, uv, ir, pmr, cryst struct*)

Boeck, L.D. *et al.*, *J. Antibiot.*, 1975, **28**, 95 (*deriv*)

Michel, K.H. *et al.*, *J. Antibiot.*, 1975, **28**, 102 (*isol, struct, nmr, ir*)

Rodriguez, R.J. *et al.*, *Antimicrob. Agents Chemother.*, 1980, **18**, 822 (*isol, props*)

Parks, L.W. *et al.*, *Biochem. Soc. Trans.*, 1983, **11**, 656 (*rev*)


Chrisp, P. *et al.*, *Z. Naturforsch., C*, 1990, **45**, 179 (*isol*)

Takahashi, I. *et al.*, *J. Antibiot.*, 1993, **46**, 1312 (UCA 1064)

Back, T.G. *et al.*, *Tetrahedron*, 1993, **49**, 337 (*synth*)

1-Aza-2,5,8-trioxacyclononane**A-1603**

Hexahydro-1,4,7,2-trioxazonine, 9CI



$C_5H_{11}NO_3$ 133.147

N-Butyl: *Trigoxazonane*

[957466-26-1]


$C_9H_{19}NO_3$ 189.254

Alkaloid from the roots of *Trigonella foenum-graecum* (fenugreek). Allelopathic agent. Oil.

Evidente, A. *et al.*, *Phytochemistry*, 2007, **68**, 2487-2492 (*isol, pmr, cmr, ms*)

Azcarpine, 9CI**A-1604**

[17420-53-0]




$C_{26}H_{46}N_2O_4$ 450.66

Unsymmetrical homologue intermediate between Carpaine, C-168 and Azimine, A-1608. Alkaloid from dried leaves and stems of *Azima tetracantha* (Salvadoraceae). Plates (petrol). Mp 78-80°. $[\alpha]_D^{220}$ -343 (c, 0.11 in MeOH).

Smalberger, T.M. *et al.*, *Tetrahedron*, 1968, **24**, 6417 (*isol, struct, abs config*)

Azepinomycin**A-1605**

4,5,6,7-Tetrahydro-6-hydroxyimidazo[4,5-e][1,4]diazepin-8(1H)-one, 9CI [89354-15-4]



$C_6H_8N_4O_2$ 168.155

Isol. from *Streptomyces* MF718-03. Possesses antiganase and antitumour activities. Sol. MeOH, EtOH, butanol; fairly sol. H_2O ; poorly sol. C_6H_6 , hexane. λ_{\max} 203 (ε 12600); 279 (ε 8250) (H_2O) (Derep). λ_{\max} 290 (MeOH) (Berdy). λ_{\max} 290 (ε 3600) (H_2O) (Berdy).

Japan. Pat., 1983, 58 159 494; *CA*, **100**, 137362 (*isol*)

Isshiki, K. *et al.*, *J. Antibiot.*, 1987, **40**, 1461 (*synth*)


Fujii, T. *et al.*, *Heterocycles*, 1988, **27**, 1163 (*synth*)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (*rev*)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1231 (*synth*)

3-Azetidinone**A-1606**

[54044-11-0]



C_3H_5NO 71.079

Claimed syntheses prior to 1988 were prob. erroneous.

Covalent hydrate: 3,3-Azetidinediol. 3,3-Dihydroxyazetidone. Antibiotic BM 1. BM 1

[244145-62-8]

$C_3H_7NO_2$ 89.094

Prod. by *Bacillus mesentericus*. Promotes growth of *Bifidobacterium* sp.

Hydrochloride: [17557-84-5]

Solid. Sol. H_2O . Mp 110-140°. Con-

tains ca. 20% MeOH.

N-Ac: [179894-05-4]
C₅H₇NO₂ 113.116
Oil.

N-Ac, oxime: [179894-09-8]
C₅H₈N₂O₂ 128.13
Microcryst. solid (Me₂CO). Mp 220-221° dec.


N-Benzoyl: [25566-02-3]
C₁₀H₉NO₂ 175.187
Cryst. Mp 85-86°.

N-(4-Methylbenzenesulfonyl): N-Tosyl-3-azetidione
[76543-27-6]
C₁₀H₁₁NO₃S 225.268
Mp 149°.

N-Nitro: [157952-92-6]
C₃H₄N₂O₃ 116.076
Microcryst. solid. Mp 61-62°.

Baumann, H. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 1035 (*synth, ir, pmr*)
Katritzky, A.R. *et al.*, *J. Het. Chem.*, 1994, **31**, 271 (*tosyl*)
Marchand, A.P. *et al.*, *J.O.C.*, 1994, **59**, 5499 (*N-nitro*)
Marchand, A.P. *et al.*, *J. Het. Chem.*, 1996, **33**, 837 (*N-benzoyl*)
Dave, P.R. *et al.*, *J.O.C.*, 1996, **61**, 5453-5455 (*N-Ac, N-nitro, synth, pmr, cmr*)
Eur. Pat., 1999, 943 309; *CA*, **131**, 242089b (*Antibiotic BMI*)
Marchand, A.P. *et al.*, *Synth. Commun.*, 1999, **29**, 885-890 (*N-Ac, N-benzoyl*)
Seo, G. *et al.*, *Microbios*, 2000, **101**, 105-114 (*Antibiotic BMI*)
Dejaegher, Y. *et al.*, *Chem. Rev.*, 2002, **102**, 29-60 (*derivs, rev*)

6-Azidotetrazolo[5,1-a]phthalazine, 9CI A-1607
[51627-67-9]




C₈H₄N₈ 212.173

Isol. from the toxic flagellate *Gymnodinium breve*. Ichthyotoxin. Mp 152° (*synthetic*).

- LD₅₀ 0.4 µg/ml (fish). Potentially explosive.
Reynolds, G.A. *et al.*, *J.O.C.*, 1959, **24**, 1205 (*synth, struct*)
Waller, A.R. *et al.*, *J. Chromatogr.*, 1979, **179**, 392 (*ir*)
Hossain, M.B. *et al.*, *Acta Cryst. C*, 1985, **41**, 1199 (*cryst struct*)

Azimine, 9CI A-1608
[17398-27-5]



C₂₄H₄₂N₂O₄ 422.607


Lower homologue of Carpaine, C-168. Alkaloid from dried leaves and stems of *Azima tetracantha* (Salvadoraceae). Cubes (Et₂O). Mp 112-113°. [α]_D²⁵ -899 (c, 0.11 in MeOH).

Hydrochloride: Mp 284-287°.

Di-N-Me: Mp 92-93°.

Smalberger, T.M. *et al.*, *Tetrahedron*, 1968, **24**, 6417 (*isol, struct, abs config*)
Brown, E. *et al.*, *J.C.S. Perkin 1*, 1976, 2190 (*synth*)
Hanessian, S. *et al.*, *Tet. Lett.*, 1979, 3391 (*synth*)
Natsume, M. *et al.*, *Heterocycles*, 1980, **14**, 169 (*synth*)
Sato, T. *et al.*, *Org. Lett.*, 2003, **5**, 3839-3842 (*synth*)

Azitrine A-1609
Atisine azomethine
[18041-82-2]



C₂₀H₂₉NO 299.455

Alkaloid from *Aconitum zeravschanicum* and *Consolida hellespontica*. Cryst. (hexane/Me₂CO). Mp 177-179° Mp 180-182°.

13-(2-Methylbutanoyloxy): *13-(2-Methylbutanoyloxy)azitrine*

C₂₅H₃₇NO₃ 399.572
Alkaloid from the roots of *Delphinium scabriflorum*. Amorph. [α]_D²⁰ -7.6 (c, 0.2 in CHCl₃). λ_{\max} 214 (log ϵ 3.43) (EtOH).

$\Delta^{19(N)}$ -Isomer: *Isoazitrine*

[290818-20-1]
C₂₀H₂₉NO 299.455
Alkaloid from *Delphinium staphisagria*. Gum. [α]_D²⁵ -6.7 (c, 0.63 in CHCl₃).

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1992, **70**, 1397-1405 (*synth, pmr, cmr*)
Salimov, B.T. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 70 (*isol*)
Desai, H.K. *et al.*, *Heterocycles*, 1993, **36**, 1081 (*isol, pmr, cmr, struct*)
Diaz, J.G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1136-1139 (*Isoazitrine*)
Shrestha, P.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1574-1576 (*13-(2-Methylbutanoyloxy)azitrine*)

4-Azoniaspiro[3.3]heptane-2,6-diol A-1610
Charamin
[106500-22-5]




C₆H₁₂NO₂⁺ 130.166

Isol. from the green alga *Chara globularis*. Possesses antibacterial props. No phys. props. reported.

Anthoni, U. *et al.*, *J.O.C.*, 1987, **52**, 694 (*isol, synth*)

Azotobactin D A-1611
[109336-19-8]




C₅₅H₇₉N₁₆O₂₈⁺ 1412.32

Struct. revised in 1988. Positive charge delocalised over the two ring Ns. Prod. by *Azotobacter vinelandii*. Siderophore. λ_{\max} 336 (sh) (ϵ 19600); 380 (ϵ 23500) (pH 5 buffer).

Demange, P. *et al.*, *Biochemistry*, 1988, **27**, 2745-2752 (*isol, pmr, cmr, uv, struct*)

Azotobactin G173 A-1612
[535994-94-6]




C₄₆H₆₄N₁₃O₁₉⁺ 1103.087

Prod. by *Pseudomonas fluorescens* G173. Siderophore.

Fernandez, D.U. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 1-10 (*isol, pmr, cmr, ms*)

Azotobactin P19 A-1613
[280142-34-9]




C₄₃H₅₉N₁₂O₂₂⁺ 1096.006

Positive charge delocalised over the two ring N atoms. Prod. by *Pseudomonas fluorescens* P19. Siderophore.

Vossen, W. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 153-164

Azoverdin A

[176450-81-0]

R = $-\text{COCH}_2\text{CH}_2\text{COOH}$ $\text{C}_{46}\text{H}_{67}\text{N}_{12}\text{O}_{19}^{\oplus}$ 1092.104

Isol. from *Azomonas macrocytogenes* ATCC 12334. Siderophore. Struct. revised in 1996. λ_{max} 400 (€ 19500) (MeOH) (Berdy). λ_{max} 366 (€ 11220); 375 (€ 10964) (pH 3 buffer) (Berdy). λ_{max} 399 (€ 17350) (H_2O) (Berdy).

4'-Amide: Azoverdin

[131688-65-8]

 $\text{C}_{46}\text{H}_{68}\text{N}_{13}\text{O}_{18}^{\oplus}$ 1091.12

Isol. from *Azomonas macrocytogenes* ATCC 12334. Siderophore. Sol. H_2O . Azoverdin and Azoverdin A are different compounds. λ_{max} 380 ; 450 (pH 5 buffer) (Berdy).

Linget, C. *et al.*, *Tet. Lett.*, 1992, **33**, 1889-1892 (*isol, ms, pmr, cmr*)

A-1614

Bernardini, J.-J. *et al.*, *BioMetals*, 1996, **9**, 107-120 (*isol, pmr, cmr, N-15 nmr*)

Michalke, R. *et al.*, *Z. Naturforsch., C*, 1996, **51**, 772-780 (*isol, uv, pmr, cmr*)

Azoverdin G

[187271-50-7]

As Azoverdin A, A-1614 with

R = $-\text{COCH}_2\text{CH}_2\text{CH}(\text{NH}_2)\text{COOH}$ $\text{C}_{47}\text{H}_{70}\text{N}_{13}\text{O}_{19}$ 1121.146Isol. from *Azomonas macrocytogenes*.

Siderophore. λ_{max} 366 (€ 10920); 375 (€ 11200) (pH 3 buffer) (Berdy). λ_{max} 400 (€ 11220) (H_2O) (Berdy).

5 β ,6-Dihydro: 5,6-Dihydroazoverdin G

[203264-21-5]


 $\text{C}_{47}\text{H}_{72}\text{N}_{13}\text{O}_{19}$ 1123.162Isol. from *Azomonas macrocytogenes*.

Siderophore. λ_{max} 301 (MeOH) (Berdy). λ_{max} 250 ; 318 ; 535 (pH 3 buffer) (Berdy). λ_{max} 250 ; 328 ; 517 (H_2O) (Berdy).

Michalke, R. *et al.*, *Z. Naturforsch., C*, 1996, **51**, 772-780; 1997, **52**, 855-857 (*isol, uv, pmr, cmr, Dihydroazoverdin G*)

Azoxybacilin**A-1616**

2-Amino-4-(methyl-NNO-azoxy)butanoic acid, 9CI

 $\text{C}_5\text{H}_{11}\text{N}_3\text{O}_3$ 161.16**(Z,S)-form** [157998-96-4]Prod. by *Bacillus cereus* NR2991.

Methionine antagonist. Antifungal agent. Needles (MeOH aq.). Sol. H_2O , MeOH, DMSO; poorly sol. Me_2CO , hexane. Mp 203-205°. $[\alpha]_{\text{D}}^{24}$ +9.4 (c, 1 in H_2O). λ_{max} 215 (€ 8400) (MeOH) (Berdy).

Ohwada, J. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1703 (*synth*)

Fujiu, M. *et al.*, *J. Antibiot.*, 1994, **47**, 833 (*isol, struct*)

Aoki, Y. *et al.*, *J. Antibiot.*, 1994, **47**, 909 (*props*)

Aztequine**A-1617**

[57608-18-1]

 $\text{C}_{30}\text{H}_{40}\text{N}_2\text{O}_7$ 540.655


Bisbenzylisoquinoline alkaloid. Struct. unknown. A struct. was originally proposed, but is v. improbable. Alkaloid from the powdered leaves of *Yoloxochitl* (*Embriophyta* sp.; not a recognised genus, prob. *Magnolia* sp.) (Magnoliaceae). Light-brown. Mp 176°.

Pallares, E.S. *et al.*, *Arch. Biochem.*, 1948, **16**, 275-282

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 949-952

Bacillamide A

Microbiaeratimin
[637344-98-0]



C₁₆H₁₅N₃O₂S 313.379

Prod. by the marine *Bacillus* sp. SY-1, *Bacillus endophyticus* SP31, *Microbispora aerata* IMBAS-11A and *Thermoactinomyces* sp. TA66-2. Algicide active against *Cochlodinium polykrikoides*. Amorph. powder. λ_{max} 279 (log ε 3.72) (no solvent reported).

15R-Alcohol: Bacillamide B

[959853-21-5]

C₁₆H₁₇N₃O₂S 315.395

Prod. by *Bacillus endophyticus* SP31. Light brown solid. [α]_D²⁰ +7.4 (c, 0.09 in MeOH).

15-Deoxo, 15R-acetamido: Bacillamide C

[959853-22-6]

C₁₈H₂₀N₄O₂S 356.448

Prod. by *Bacillus endophyticus* SP31. Light brown solid. [α]_D²⁴ -15.2 (c, 0.08 in MeOH). λ_{max} 223 (log ε 4.05); 280 (log ε 3.5); 288 (log ε 3.41); 340 (log ε 2.71) (MeOH).


Jeong, S.-Y. *et al.*, *Tet. Lett.*, 2003, **44**, 8005-8007 (*isol, pmr, cmr*)

Socha, A.M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1793-1795 (*Bacillamides B, C*)

Ivanova, V. *et al.*, *Prep. Biochem. Biotechnol.*, 2007, **37**, 161-168 (*isol, pmr, cmr*)

Bacillibactin

Corynebactin
[95536-41-7]



C₃₉H₄₂N₆O₁₈ 882.79

Related to Enterobactin, E-86. Prod. by *Bacillus subtilis* ATCC 21332 and *Corynebacterium glutamicum* ATCC 14067. Siderophore.

Budzkievicz, H. *et al.*, *Z. Naturforsch., C.*, 1997, **52**, 551-554 (*Corynebactin*)

May, J.J. *et al.*, *J. Biol. Chem.*, 2001, **276**, 7209-7217 (*isol*)


May, J.J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2002, **99**, 12120-12125 (*synth*)

Dertz, E.A. *et al.*, *J.A.C.S.*, 2006, **128**, 22-23 (*activity*)

B-1

Bagougeramine A

[104840-35-9]



C₁₇H₂₈N₁₀O₇ 484.471

Nucleoside antibiotic. Prod. by *Bacillus circulans*. Shows broad spectrum antimicrobial activity and active against two-spotted spider-mite. Sol. H₂O; poorly sol. MeOH, Me₂CO. λ_{max} 274 (ε 13060) (0.1N HCl) (Derep). λ_{max} 235 (sh); 265 (ε 9680) (0.1N NaOH) (Derep). λ_{max} 232 (ε 9220); 265 (ε 9310) (H₂O) (Derep).

► LZ6560000

Sulfate (1:1.5):

Hygroscopic powder + 3H₂O. Mp 230° dec. [α]_D²⁷ +34.1 (c, 0.5 in H₂O).

Takahashi, A. *et al.*, *J. Antibiot.*, 1986, **39**, 1033; 1041 (*isol, struct, props*)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (*rev*)

Bagougeramine B

[104840-34-8]

As Bagougeramine A, B-3 with

R = -(CH₂)₃NH(CH₂)₄NH₂

C₂₄H₄₄N₁₂O₇ 612.688

Nucleoside antibiotic. Prod. by *Bacillus circulans*. Miticide, acaricide. Shows broad spectrum antimicrobial activity and active against two-spotted spider mite. Hygroscopic powder + 6H₂O (as sulfate). Sol. H₂O; poorly sol. MeOH, Me₂CO. Mp 237° dec. (sulfate). [α]_D²⁷ +22.1 (c, 0.5 in H₂O) (sulfate). λ_{max} 274 (ε 13060) (0.1M HCl) (Derep). λ_{max} 235 (sh); 265 (ε 9680) (0.1M NaOH) (Derep). λ_{max} 232 (ε 9220); 265 (ε 9310) (H₂O) (Derep).


► LZ6550000

Takahashi, A. *et al.*, *J. Antibiot.*, 1986, **39**, 1033; 1041 (*isol, struct, props*)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (*rev*)

Bahamamide

2-(3-Decenyl)-1,6-diazacyclododec-8-ene-7,12-dione, 9CI
[190782-45-7]



B-3


C₂₀H₃₄N₂O₂ 334.501

Prod. by an unidentified marine bacterium. Oil. [α]_D +27.8 (c, 0.5 in CHCl₃). λ_{max} 227 (ε 1700) (THF).

Boehler, M. *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 75-78 (*isol, uv, ir, pmr, cmr*)

Bahienoside A

[552333-59-2]



C₄₄H₅₈N₂O₁₈ 902.945

Alkaloid from the aerial parts of *Psychotria bahiensis*. Creamy amorph. solid (EtOAc). Mp 156-158°. [α]_D²⁵ -128 (c, 0.003 in MeOH). λ_{max} 228 (log ε 4.63); 284 (sh) (MeOH).

3-Epimer: Bahienoside B

[552333-60-5]


C₄₄H₅₈N₂O₁₈ 902.945

Alkaloid from the aerial parts of *Psychotria bahiensis*. Pale yellowish amorph. solid (EtOAc). Mp 164-166°. [α]_D²⁵ +65 (c, 0.003 in MeOH). λ_{max} 226 (log ε 4.5); 286 (log ε 3.79) (MeOH).

Paul, J.H.A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 752-754 (*isol, pmr, cmr, ms*)

Baicaline

5,6,6a,7-Tetrahydro-9,10,12-trimethoxy-4H-benzo[g]-1,3-benzodioxolo[4,5,6-de]-quinoline, 9CI. 1,9,10-Trimethoxy-2,3-methylenedioxyranoraphrine



C₂₀H₂₁NO₅ 355.39

(S)-form [83008-38-2]

Alkaloid from *Thalictrum baicalense* (Ranunculaceae). Mp 169-172°. [α]_D +48 (MeOH).

N-Me: Baicalidine

[83008-39-3]

[109010-09-5]

C₂₁H₂₃NO₅ 369.416

Alkaloid from *Thalictrum baicalense* (Ranunculaceae). Cryst. (Me₂CO). Mp 146-147°. [α]_D +55 (MeOH). λ_{max} 217; 242; 291; 306; 318 (MeOH).

Maekh, S.K. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 227; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 208 (*isol, uv, pmr, ms, struct*)

B-2

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4

B-4


B-4

Maekh, S.K. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 791; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 761 (*Baicalidine*)

Bakankoside

B-8

Bakankosine
[1398-17-0]



Absolute
configuration

$C_{16}H_{23}NO_8$ 357.36

Alkaloid from the seeds of *Strychnos vacacoua* and from *Strychnos madagascariensis* (Loganiaceae). Cryst. + H_2O (H_2O). Mp 162° (solidifies and remelts at 211-212°). $[\alpha]_D^{18}$ -195.

Tetra-O-Ac: Mp 212-214°. $[\alpha]_D^{18}$ -158 (c, 1.6 in EtOH).

Balenović, K. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 2519 (*uv, ir, struct*)

Büchi, G. *et al.*, *Tet. Lett.*, 1960, No. **26**, 5 (*struct*)

Inouye, H. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1406 (*synth, struct*)

Inouye, H. *et al.*, *J. Chromatogr.*, 1976, **118**, 201 (*ms*)

Tietze, L.F. *et al.*, *Tet. Lett.*, 1976, 2535-2538 (*isol, struct*)


Bailleul, F. *et al.*, *Phytochemistry*, 1977, **16**, 723 (*cmr*)

Tietze, L.F. *et al.*, *Tetrahedron*, 1989, **45**, 681 (*synth, uv, ir, pmr*)

Balanol

B-9

Azepinostatin. Ophiocordin
[63590-19-2]



$C_{28}H_{26}N_2O_{10}$ 550.521

Prod. by *Cordyceps ophioglossoides*, *Fusarium merismoides* and *Verticillium balanoides*. Protein kinase C inhibitor. Antifungal agent. Pale yellow amorph. powder. Mp 175° dec. $[\alpha]_D$ -127.8 (0.3 in MeOH). λ_{max} 217 (ε 19600); 292 (ε 13500) (MeOH/NaOH) (Derep). λ_{max} 213 (ε 18600); 258 (ε 9600); 274 (ε 9020); 285 (sh); 360 (ε 1380) (MeOH) (Derep).

[158931-29-4, 147397-03-3]

Kneifel, H. *et al.*, *Arch. Microbiol.*, 1977, **113**, 121 (*isol, ms, ir, nmr*)

Koenig, W.A. *et al.*, *Chem. Ber.*, 1980, **113**, 2221 (*ms, nmr, struct*)

Kulanthaivel, P. *et al.*, *J.A.C.S.*, 1993, **115**, 6452 (*isol*)

Ohshima, S. *et al.*, *J. Antibiot.*, 1994, **47**, 639 (*isol*)

Boros, C. *et al.*, *J. Antibiot.*, 1994, **47**, 1010 (*isol, pmr*)

Miyabe, H. *et al.*, *J.O.C.*, 1998, **63**, 4397-4407 (*synth, bibl*)

Phansavath, P. *et al.*, *Eur. J. Org. Chem.*, 2000, 3903-3907 (*synth*)

Laursen, B. *et al.*, *Tetrahedron*, 2002, **58**, 2231-2238 (*synth, bibl*)

Patil, M.L. *et al.*, *Tetrahedron*, 2004, **60**, 1869-1873 (*synth*)


Sullivan, B. *et al.*, *J. Nat. Prod.*, 2008, **71**, 346-350 (*synth*)

Sullivan, B. *et al.*, *Tet. Lett.*, 2008, **49**, 5211-5213 (*synth, bibl*)

Balasubramide

B-10

[190905-98-7]



$C_{20}H_{20}N_2O_2$ 320.39


Alkaloid from leaves of *Clausena indica*. Oil. $[\alpha]_D$ +7 (c, 0.5 in $CHCl_3$). λ_{max} 224; 274 (sh); 283; 292 (Et_2O).

Riemer, B. *et al.*, *Phytochemistry*, 1997, **45**, 337-341 (*isol, uv, ir, pmr, cmr, ms, struct*)

Yang, L. *et al.*, *Org. Lett.*, 2007, **9**, 1387-1390 (*synth*)

Baleabuxaline I

B-11



$C_{24}H_{42}N_2O_3$ 406.607

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Baleabuxaline I, is unknown.

N^{20}, N^{20} -*Di-Me*, N^3 -(2-methylpropanoyl): ***N*-Isobutyrylbaleabuxaline F** [14155-71-6]

$C_{30}H_{52}N_2O_4$ 504.752

Alkaloid from *Buxus balearica* (Buxaceae). Mp 275°. $[\alpha]_D$ -32 ($CHCl_3$).


N^{20}, N^{20} -*Di-Me*, N^3 -(2-methylpropanoyl), O,O' -*di-Ac*: Mp 261°. $[\alpha]_D$ -10 ($CHCl_3$).

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*isol, ir, pmr*)

Baleabuxidine I

B-12

3,20-Diamino-16-hydroxy-4-hydroxy-methyl-4,14-dimethyl-9,19-cyclopregnan-11-one. Cycloabuxidine I



$C_{24}H_{40}N_2O_3$ 404.592

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Baleabuxidine I, is unknown.

N^{20} -*Me*, N^3 -(2-methylpropanoyl): ***3-N*-Isobutyrylcycloabuxidine H** [19330-62-2]

$C_{29}H_{48}N_2O_4$ 488.709

Alkaloid from *Buxus balearica* (Buxaceae). Cryst. (EtOAc). Mp 285°. $[\alpha]_D$ +75 ($CHCl_3$).

N^{20}, N^{20} -*Di-Me*: ***Baleabuxidine F***. *Cycloabuxidine F*. *Buxidine F*† [4947-53-9]

$C_{26}H_{44}N_2O_3$ 432.645

Alkaloid from *Buxus balearica* (Buxaceae). Mp 242° (227-230°). $[\alpha]_D$ +127 (+114) ($CHCl_3$). Reacn. with HCHO gives Cyclobuxoxazine A.

N^{20}, N^{20} -*Di-Me*, N^3 -(2-methylpropanoyl): ***Isobutyrylbaleabuxidine F***. *Baleabuxidine*. N^3 -*Isobutyrylcycloabuxidine F*

$C_{30}H_{50}N_2O_4$ 502.736

Alkaloid from *Buxus balearica* (Buxaceae). Mp 257° (236-238°). $[\alpha]_D$ +71 ($CHCl_3$).

N^{20}, N^{20} -*Di-Me*, N^3 -benzoyl: ***Benzoylbaleabuxidine F***. N^3 -*Benzoylcycloabuxidine F*

[15524-62-6]

$C_{33}H_{48}N_2O_4$ 536.753

Alkaloid from *Buxus balearica* and *Buxus sempervirens* (Buxaceae). Mp 277° (274-276°). $[\alpha]_D$ +57 ($CHCl_3$).

N^{20}, N^{20} -*Di-Me*, N^3 -benzoyl, O^{16} -*Ac*: ***N*-Benzoyl-16-acetylcycloabuxidine** [108567-60-8]

$C_{35}H_{50}N_2O_5$ 578.79

Alkaloid from the leaves of *Buxus papilosa* (Buxaceae). Amorph. needles. $[\alpha]_D$ +42. Shown in the reference with the incorrect C_4 -config. which has been reversed for these alkaloids.

Herlem-Gaulier, D. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 3478-3486; 1968, 763-773 (*Buxus balearica constitis*)


Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*Benzoylbaleabuxidine F*, *Isobutyrylbaleabuxidine F*)

Kurakina, I.O. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 26-28; 1970, **6**, 231-235; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 20-21; 1970, **6**, 225-227 (N^3 -*Isobutyrylcycloabuxidine F*)

- Kupchan, S.M. *et al.*, *Tetrahedron*, 1973, **23**, 4563-4586 (*Benzoylbaleabuxidine F*)
 Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*cmr, struct*)
 Guilhem, J. *et al.*, *Tet. Lett.*, 1975, **16**, 2937-2938 (*cryst struct, Baleabuxidine*)
 Atta-ur-Rahman, *et al.*, *Planta Med.*, 1987, **53**, 75 (*N-Benzoyl-16-acetylcyclohexobuxidine*)

Balfourodine**B-13**

3,9-Dihydro-2-(1-hydroxy-1-methylethyl)-8-methoxy-9-methylfuro[2,3-b]quinolin-4(2H)-one, 9CI. *Hydroxy-lunacrine*



$C_{16}H_{19}NO_4$ 289.33

(R)-form [484-61-7]

Alkaloid from *Balfourodendron riedelianum* (Rutaceae). Cryst. ($CHCl_3/C_6H_6$). Mp 188-189°. $[\alpha]_D^{25} +49$ (c, 1 in EtOH). *Perchlorate*: Mp 212-213°.

l'-Deoxy: **Lunacrine**

[82-40-6]

$C_{16}H_{19}NO_3$ 273.331

Alkaloid from leaves of *Lunasia amara* and other *Lunasia* spp. (Rutaceae).

Hypotensive agent. Needles (EtOAc). Mp 117-119°. $[\alpha]_D^{25} -50$ (c, 0.8 in EtOH). Forms monohydrate, Mp 96°.

▶ Toxic, LD₅₀ (mus, ivn) 80 mg/kg.**(S)-form** [17958-35-9]

Alkaloid from *Lunasia amara* (Rutaceae). Mp 201-203°.

Perchlorate: Mp 216-218°. $[\alpha]_D^{24} -14.6$ (EtOH).

(±)-form

Prisms (EtOAc). Mp 189-191°.

Perchlorate: Mp 212-213°.

l'-Deoxy:

$C_{16}H_{19}NO_3$ 273.331

Alkaloid from bark of *Lunasia amara* (Rutaceae). Mp 145-147°.

(ξ)-form

l'-Deoxy, *l',2'*-didehydro: **Neoacutifolin**

[145237-09-8]

$C_{16}H_{17}NO_3$ 271.315

Alkaloid from leaves of *Zanthoxylum petiolare* (Rutaceae). Viscous oil. $[\alpha]_D +124$ (c, 0.0004 in $CHCl_3$).

Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 1908; 3065; 6209 (*Lunacrine, Balfourodine*)

Rapoport, H. *et al.*, *J.A.C.S.*, 1959, **81**, 3738; 1960, **82**, 4395 (*Balfourodine*)

Clarke, E.A. *et al.*, *J.C.S.*, 1964, 438; 4196 (*synth*)

Bowman, R.M. *et al.*, *J.C.S. Perkin 1*, 1973, 1051 (*config*)

Hammerum, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1977, **31**, 31 (*ms*)

Rideau, M. *et al.*, *Phytochemistry*, 1979, **18**, 155 (*isol, pmr, ms*)

Ramesh, M. *et al.*, *Tetrahedron*, 1984, **40**, 3431 (*synth, pmr*)


Arruda, M.S.P. *et al.*, *Phytochemistry*, 1992, **31**, 3617; 1994, **36**, 1303 (*Neoacutifolin*)

Anand, R.C. *et al.*, *J. Chem. Res., Synop.*, 1998, 6-7 (*synth, abs config*)

Sekar, M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 294-296 (*synth, pmr*)

Balfourodinium**B-14**

2,3-Dihydro-2-(1-hydroxy-1-methylethyl)-4,8-dimethoxy-9-methylfuro[2,3-b]quinolinium, 9CI. *O⁺-Methylbalfourodinium*

*(R)-form*

$C_{17}H_{22}NO_4^{\oplus}$ 304.365

(R)-form [38487-24-0]

Alkaloid from the bark of *Balfourodendron riedelianum* (Rutaceae). Cryst. (H_2O or EtOH/Et₂O) (as perchlorate). Mp 124-125° (perchlorate). $[\alpha]_D^{25} +9$ (EtOH).

O⁸-De-Me: 2,3-Dihydro-8-hydroxy-2-(1-hydroxy-1-methylethyl)-4-methoxy-9-methylfuro[2,3-b]quinolinium. Pteleatine. Pteleatinium

[34443-73-7]

$C_{16}H_{20}NO_5^{\oplus}$ 290.338

Alkaloid from *Ptelea trifoliata* (Rutaceae). Shows antimicrobial activity. Cryst. (MeOH) (as chloride). Mp 267-270° (chloride). λ_{max} 213 ; 233 ; 257 ; 277 (MeOH).

l'-Deoxy: *2,3-Dihydro-4,8-dimethoxy-9-methyl-2-(1-methylethyl)furo[2,3-b]quinolinium. 2,3-Dihydro-2-isopropyl-4,8-dimethoxy-9-methylfuro[2,3-b]quinolinium. Lunasine*

[6901-22-0]

$C_{17}H_{22}NO_3^{\oplus}$ 288.366

Alkaloid from the bark and leaves of *Lunasia quercifolia* (Rutaceae). Microcryst. powder (MeOH/Et₂O) (as perchlorate). Mp 195-196° (perchlorate). $[\alpha]_D^{20} -29.3$ (c, 0.96 in MeOH).

(S)-form [70494-63-2]

Alkaloid from the leaves, roots and stems of *Choisya ternata* (Rutaceae). Mp 184-185° (as chloride). $[\alpha]_D^{20} -15.3$ (c, 0.02 in MeOH).

l'-Deoxy: Synthetic. Tan cryst. (MeOH) (as perchlorate). Mp 195-196° (perchlorate). $[\alpha]_D^{24} +33.9$ (c, 0.193 in $CHCl_3$).

(±)-form

Synthetic. Prisms (MeOH/Et₂O) (as perchlorate). Mp 203-204° (perchlorate).

N-De-Me: 8-Methoxyplatydesmine

$C_{16}H_{19}NO_4$ 289.33

Alkaloid from the leaves of *Melicope semecarpifolia*. Prisms ($CHCl_3/MeOH$). Mp 166-168°. λ_{max} 217 (log ε 4.44); 248 (log ε 4.59); 280 (log ε 3.82); 314 (log ε 3.45); 327 (log ε 3.44) (EtOH).

l'-Deoxy: Synthetic. Mp 130-132° dec. (as iodide).

(ξ)-form

l'-Deoxy, *l',2'*-didehydro: **Ptelecuninium**

[111509-10-5]

$C_{17}H_{20}NO_3^{\oplus}$ 286.35

Quaternary alkaloid from roots and callus strains of *Ptelea trifoliata* (Rutaceae). Mp 210-212° dec. (as chloride).

Price, J.R. *et al.*, *Aust. J. Chem.*, 1959, **12**, 458 (*Lunacrine, isol, ir*)

Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 1908 (*Lunacrine, synth, uv, ir*)

Rapoport, H. *et al.*, *J.A.C.S.*, 1959, **81**, 3738 (*isol, uv, struct, synth*)

Clarke, E.A. *et al.*, *J.C.S.*, 1964, 4196 (*synth, iv*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1966, **19**, 2185 (*Lunacrine, isol*)

Bowman, R.M. *et al.*, *J.C.S. Perkin 1*, 1973, 1051 (*Lunacrine, config*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1975, **38**, 109-116 (*Pteleatine, activity*)

Rideau, M. *et al.*, *Phytochemistry*, 1979, **18**, 155 (*isol, pmr, ms, struct*)

Ramesh, M. *et al.*, *Tetrahedron*, 1984, **40**, 3431 (*synth*)


Petit-Paly, G. *et al.*, *Plant Cell Rep.*, 1987, **6**, 309 (*Ptelecuninium*)

Petit-Paly, G. *et al.*, *Planta Med.*, 1989, **55**, 209 (*Ptelecuninium*)

Chen, J.-J. *et al.*, *Planta Med.*, 2003, **69**, 542-546 (*8-Methoxyplatydesmine*)

Bannucine**B-15**

[104380-51-0]



Absolute Configuration


$C_{29}H_{37}N_3O_7$ 539.627

Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Mp 152-154°. $[\alpha]_D -33$ (c, 0.26 in $CHCl_3$).

Atta-ur-Rahman, *et al.*, *J.C.S. Perkin 1*, 1986, 923 (*isol, uv, ir, pmr, cmr, ms, struct*)

Banyascyclamide B**B-16**

[501904-00-3]

R = -CH₂CH(CH₃)₂

$C_{22}H_{30}N_6O_5S_2$ 522.648

Isol. from *Nostoc* sp. TAU strain IL-235.

Ploutno, A. *et al.*, *Tetrahedron*, 2002, **58**, 9949-9957 (*isol, pmr, cmr, ms*)

Banyascyclamide C**B-17**

[501904-01-4]

As Banyascyclamide B, B-16 with

R = -CH₂Ph


$C_{25}H_{28}N_6O_5S_2$ 556.665

Isol. from *Nostoc* sp. TAU strain IL-235. $[\alpha]_D^{25} +50.1$ (c, 0.1 in MeOH). This is an isomer of Didmolamide B, D-382. λ_{max} 222 (ε 15100); 238 (ε 16300) (MeOH).

Ploutno, A. *et al.*, *Tetrahedron*, 2002, **58**, 9949-9957 (*isol, pmr, cmr, ms*)

Banyaside A

B-18



$C_{40}H_{64}N_8O_{19}S$ 993.054

Isol. from a *Nostoc* sp. Trypsin and thrombin inhibitor. Glassy solid. $[\alpha]_D^{20} +28.7$ (c, 1 in MeOH). λ_{max} 218 (ϵ 4000); 271 (ϵ 2400) (MeOH).

Pluotno, A. *et al.*, *Tetrahedron*, 2005, **61**, 575-583 (isol, pmr, cmr)

Banyaside B

B-19

As Banyaside A, B-18 with $R^1 = R^2 = H, R^3 = -CO(CH_2)_4CH_3$

$C_{39}H_{63}N_7O_{18}S$ 950.029


Isol. from a *Nostoc* sp. Trypsin and thrombin inhibitor. Glassy solid. $[\alpha]_D^{20} +17.3$ (c, 0.3 in MeOH). λ_{max} 234 (ϵ 3600); 268 (ϵ 2300) (MeOH).

Pluotno, A. *et al.*, *Tetrahedron*, 2005, **61**, 575-583 (isol, pmr, cmr)

Baptifoline

B-20

13-Hydroxyanagyridine. Alkaloid P3† [732-50-3]



Absolute Configuration

$C_{15}H_{20}N_2O_2$ 260.335

Alkaloid from *Baptisia perfoliata*, *Baptisia minor*, *Sophora alopecuroides*, *Sophora flavescens* (seeds) and *Caulophyllum thalictroides* (blue cohosh) (Fabaceae, Leonticaceae). Hexagonal or triangular plates (MeOH), needles (Me₂CO). Mp 210°. $[\alpha]_D -147.7$ (-140) (EtOH). $[\alpha]_D^{16} -89.05$ (c, 1.415 in H₂O).

Perchlorate:

Plates (MeOH). Mp 286-287° (279°).

Picrate:

Yellow needles (MeOH). Mp 256° (sinters at 145°).

Ac: 13-Acetoxyanagyridine. O-Acetylbaptifoline

$C_{17}H_{22}N_2O_3$ 302.372

Alkaloid in *Baptisia australis* and *Thermopsis chinensis* (Fabaceae). No phys. props. detd. for nat. alkaloid.

13-Epimer: 13-Epibaptifoline. 13-Epilydroxyanagyridine

$C_{15}H_{20}N_2O_2$ 260.335

Alkaloid from *Genista sphaerocarpa* leaves and from *Genista monosperma* (Fabaceae). Mp 215°. $[\alpha]_D^{21} -138.9$.

13-Epimer; hydrochloride:

Cryst. + 1H₂O. Mp 270° dec. $[\alpha]_D -165.3$ (H₂O).

13-Epimer; picrate: Mp 250°.

11-Epimer: **Argentamine**. 13-Hydroxythermopsine

[27773-56-4]

$C_{15}H_{20}N_2O_2$ 260.335

Alkaloid from *Ammodendron argenteum* and *Thermopsis licentiana* (Fabaceae). Cryst. (C₆H₆). Mp 203°. $[\alpha]_D -142.3$ (c, 2.04 in EtOH).

11-Epimer; hydrochloride: Mp 298°.

11-Epimer; picrate:

Cryst. (EtOH). Mp 230°.

Marion, L. *et al.*, *J.A.C.S.*, 1948, **70**, 3253; 3472 (isol)

Martin-Smith, M. *et al.*, *Can. J. Chem.*, 1957, **35**, 37 (struct)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1958, **91**, 2189; 2194; 1962, **95**, 944 (isol, struct, synth)

Vazquez, M.D. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1966, **62**, 837; *CA*, **66**, 76216h (epimer)

Flom, M.S. *et al.*, *J. Pharm. Sci.*, 1967, **56**, 1515 (isol)

Pham Hoang Ngok, *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 111; *Chem. Nat. Compd. (Engl. Transl.)*, 103 (Argentamine)

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 59; *Chem. Nat. Compd. (Engl. Transl.)*, 52 (isol)

Ohmiya, S. *et al.*, *Phytochemistry*, 1974, **13**, 1016 (isol)


Wink, M. *et al.*, *J. Nat. Prod.*, 1981, **44**, 14 (isol, ms, acetate)

Jia, Z. *et al.*, *Gaodeng Xuexiao Huaxue Xuebao*, 1990, **11**, 1014; *CA*, **115**, 110591 (Argentamine, O-Acetylbaptifoline)

Kennelly, E.J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1385-1389 (isol, pmr, cmr)

Barbaleucamide A

B-21



$C_{16}H_{20}Cl_6N_2O_2S$ 517.128

Isol. from a *Dysidea* sp. Pale yellow oil.

N-Me: **Barbaleucamide B**

$C_{17}H_{22}Cl_6N_2O_2S$ 531.155


Isol. from a *Dysidea* sp. Pale yellow oil.

Harrigan, G.G. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1133-1138

Barbamide

B-22

[174630-03-6]



Absolute Configuration

$C_{20}H_{23}Cl_3N_2O_2S$ 461.838

Metab. from the Caribbean cyanobacterium *Lyngbya majuscula*. Possesses molluscicidal activity. Pale yellow oil. Sol. MeOH, CHCl₃, EtOAc; poorly sol.

hexane, H₂O. $[\alpha]_D^{26} -89$ (c, 1.9 in MeOH). The first marine nat. prod. for which the biosynthetic gene cluster has been sequenced. λ_{max} 240 (ϵ 16000) (MeOH) (Berdy).

Dechloro: **Dechlorobarbamide**

[327613-98-9]

$C_{20}H_{24}Cl_2N_2O_2S$ 427.393

Metab. of *Lyngbya majuscula*. Pale yellow oil. $[\alpha]_D^{25} -67$ (c, 0.05 in MeOH). λ_{max} 238 (ϵ 16000) (MeOH).

Orjala, J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 427-430 (isol, uv, ir, pmr, cmr, struct)

Hartung, J. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 1209-1211 (biosynth)

Sitachitta, N. *et al.*, *Tetrahedron*, 2000, **56**, 9103-9113 (isol, biosynth, abs config, Dechlorobarbamide)


Nguyen, V.-A. *et al.*, *Chem. Comm.*, 2001, 1934-1935 (synth)

Flatt, P.M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 938-944 (biosynth)

Barbital, INN

B-23

5,5-Diethyl-2,4,6-(1H,3H,5H)-pyrimidine-trione, 9CI. 5,5-Diethylbarbituric acid. Diethylmalonylurea. Barbitone, BAN. Veronal. Many other names [57-44-3]



$C_8H_{12}N_2O_3$ 184.194

Exists in several tautomeric forms. Isol. from tissues of the Indian puffer fish *Sphaeroides oblongus*. Powerful hypnotic and sedative of long duration. Faintly bitter cryst. (H₂O); polymorphic forms; trigonal cryst., monoclinic prisms, monoclinic needles, triclinic cryst. Mp 190° Mp 183° (181°, 176°). pK_{a1} 8.02 (25°). pK_a 7.43. Log P 0.66 (calc). λ_{max} 224 (EtOH) (Berdy). λ_{max} 224; 240 (EtOH/NaOH) (Berdy). λ_{max} 243 (NaOH) (Berdy).


▶ LD₅₀ (mus, orl) 600 mg/kg. CQ3500000

Mitra, S.K. *et al.*, *Chem. Comm.*, 1989, 16 (isol, uv, ir, pmr, cmr)

Baretin

B-24

[104311-70-8]



$C_{17}H_{19}BrN_6O_2$ 419.28

Struct. and MF revised in 2002. Isol. from the sponge *Geodia baretii*. Selective 5-HT ligand. Yellow solid. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 207-210°. $[\alpha]_D -25$ (c, 3 in MeOH). λ_{max}

235 (ε 13000); 294 (sh) (ε 5200); 340 (ε 9700) (MeOH) (Derep).

8,9-Dihydro: 8,9-Dihydrobarettin

$C_{17}H_{21}BrN_6O_2$ 421.296

Isol. from *Geodia barretti*. Brownish solid. $[\alpha]_D^{25}$ -24 (c, 0.096 in MeOH). λ_{max} 225 (log ε 1.27); 286 (log ε 0.16) (MeOH).

Sölter, S. *et al.*, *Tet. Lett.*, 2002, **43**, 3385-3386 (isol, pmr, cmr, struct)


Sjögren, M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 368-372 (isol, activity)

Johnson, A.-L. *et al.*, *Tetrahedron*, 2004, **60**, 961-965 (synth)

Hedner, E. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1421-1424 (activity)

Barettin bromobenzisoxazolone **B-25**

Bromobenzisoxazolone Barettin
[1008125-63-0]




$C_{24}H_{23}Br_2N_7O_5$ 649.298

Related to Barettin, B-24. Alkaloid from *Geodia barretti*. Antifouling agent. Brownish-yellow solid. Mp 150-151°. $[\alpha]_D^{25}$ +0.01 (c, 0.01 in MeOH). λ_{max} 198; 229; 265; 348 (MeCN).

Hedner, E. *et al.*, *J. Nat. Prod.*, 2008, **71**, 330-333 (isol, pmr, cmr, activity)

Bargustanine **B-26**
[169626-12-4]




$C_{29}H_{34}N_2O_7$ 522.597

Alkaloid from roots of *Berberis vulgaris* (barberry). Cryst. (MeOH). Mp 193-194°. $[\alpha]_D^{20}$ +114.2 (c, 0.3 in MeOH). λ_{max} 218 (sh) (log ε 4.85); 286 (log ε 3.98) (EtOH).

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 44-47; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 35-38

Barrenazine A **B-27**



$C_{24}H_{42}N_4$ 386.623

Isol. from an unidentified tunicate. Cytotoxic. Oil. $[\alpha]_D^{21}$ -55 (4) (c, 0.48 in MeOH). λ_{max} 288 (ε 6950); 313 (sh)

(MeOH).

6',6'',7',7''-Tetrahydro: Barrenazine B

$C_{24}H_{38}N_4$ 382.591

Isol. from an unidentified tunicate. Oil. $[\alpha]_D^{21}$ -21 (c, 0.26 in MeOH). λ_{max} 288 (ε 6950); 313 (sh) (MeOH).


Chill, L. *et al.*, *Org. Lett.*, 2003, **5**, 2433-2435 (isol, pmr, cmr)

Focken, T. *et al.*, *Org. Lett.*, 2006, **8**, 2985-2988 (synth)

Martinez, M.M. *et al.*, *Tet. Lett.*, 2007, **48**, 8536-8539 (synth)

Bassianin **B-28**

3-(6,8-Dimethyl-1-oxo-2,4,6-decatrienyl)-1,4-dihydroxy-5-(4-hydroxyphenyl)-2(1H)-pyridinone, 9CI
[54278-73-8]



$C_{23}H_{25}NO_5$ 395.454

Metab. of *Beauveria bassiana*. Similar to Tenellin.


Brewer, D. *et al.*, *Can. J. Bot.*, 1968, **46**, 441 (isol)

McInnes, A.G. *et al.*, *Chem. Comm.*, 1974, 281 (struct)

McInnes, A.G. *et al.*, *Can. J. Chem.*, 1977, **55**, 4090 (struct)

Bassiatin **B-29**

4-Methyl-6-(1-methylethyl)-3-(phenylmethyl)-2,5-morpholinedione, 9CI. 3-Benzyl-6-isopropyl-4-methyl-2,5-morpholinedione. PI 290. Antibiotic PI 290
[173008-70-3]



$C_{15}H_{19}NO_3$ 261.32

Identity with PI 290 not confirmed. Prod. by *Beauveria bassiana*. Platelet aggregation inhibitor. Cryst. (EtOAc/hexane). Mp 143-148°. $[\alpha]_D^{25}$ +181 (c, 0.02 in $CHCl_3$). Lateritin isol. from *Gibberella lateritium* is given the same gross struct. but data do not correspond. It was concluded that the struct. of Lateritin was incorrectly assigned.

3-Epimer: 3-Epibassiatin

$C_{15}H_{19}NO_3$ 261.32

Prod. by the fruiting bodies of *Isaria japonica*. Apoptosis-inducing agent. Powder. $[\alpha]_D^{23}$ -48.8 (c, 0.43 in $CHCl_3$). λ_{max} 209 (ε 5056) (MeOH).

[164107-91-9, 172721-96-9, 65454-13-9, 172721-98-1, 172721-97-0]

Hasumi, K. *et al.*, *J. Antibiot.*, 1993, **46**, 1782 (Lateritin)


Kagamizono, T. *et al.*, *J. Antibiot.*, 1995, **48**, 1407 (isol, pmr, cmr, ir, synth)

Japan. Pat., 1995, 95 97 389; *CA*, **123**, 31396w (PI 290)

Oh, H. *et al.*, *Planta Med.*, 2002, **68**, 345-348 (3-Epibassiatin)

Hughes, A.B. *et al.*, *J.O.C.*, 2005, **70**, 3079-3088 (synth)

Bastadin 1 **B-30**
[75513-48-3]



$C_{34}H_{30}Br_4N_4O_8$ 942.249

Isol. from the sponge *Ianthella basta*. Active against gram-positive bacteria. Foam. λ_{max} 220 (ε 63100); 283 (ε 10700); 288 (sh) (ε 8510) (MeOH) (Derep). λ_{max} 220 (ε 63000); 282 (ε 11400); 288 (ε 11000) (MeOH) (Berdy).

6-Bromo: Bastadin 2

[75513-47-2]

$C_{34}H_{29}Br_5N_4O_8$ 1021.145


From *Ianthella basta*. Active against gram-positive bacteria. Foam. λ_{max} 220 (sh) (ε 50100); 280 (ε 8510); 293 (sh) (ε 7410) (MeOH) (Derep).

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1981, **34**, 765 (isol, uv, pmr, cmr, ms, struct)

Nishiyama, S. *et al.*, *Tet. Lett.*, 1982, **23**, 1281 (synth, ir)

Guo, Z. *et al.*, *J.O.C.*, 1998, **63**, 4269-4276 (Bastadin 2, synth)

Bastadin 3 **B-31**
[79067-77-9]



$C_{34}H_{30}Br_4N_4O_8$ 942.249

Isol. from the sponge *Ianthella basta*. Active against gram-positive bacteria. Pale yellow foam. Poorly sol. hexane. λ_{max} 220 (ε 63100); 283 (ε 10700); 288 (sh) (ε 8510) (MeOH) (Derep). λ_{max} 220 (ε 57200); 285 (ε 8700) (MeOH) (Berdy).

10-O-Sulfate: 10-O-Sulfatobastadin 3

[184478-47-5]

$C_{34}H_{30}Br_4N_4O_{11}S$ 1022.313

Isol. from *Ianthella basta*. Calcium ion release inducer. Calcium ion channel modulator. Powder (as Na salt). λ_{max} 209 (ε 84400); 279 (ε 5550) (MeOH) (as Na salt).

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1981, **34**, 765 (isol, uv, pmr, cmr, ms, struct)


Nishimaya, S. *et al.*, *Tet. Lett.*, 1982, **23**, 1281 (synth, ir)

- Franklin, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1121-1127 (10-sulfate)
 Guo, Z. *et al.*, *J.O.C.*, 1998, **63**, 4269-4276 (synth)

Bastadin 4

B-32

Cyclobastadin 1
 [79067-76-8]



$C_{34}H_{25}Br_5N_4O_8$ 1017.113

Some Bastadins were renumbered in 1993. Isol. from the sponge *Ianthella basta*. Antiinflammatory agent. Active against gram-positive bacteria. Yellow needles (DMF aq.). Poorly sol. hexane. Mp 250° dec. λ_{max} 209 (€ 126000); 289 (€ 20000); 300 (sh) (€ 15800); 310 (€ 15800) (MeOH) (Derep). λ_{max} 210 (€ 68400); 228 (€ 10000); 315 (€ 10000) (MeOH) (Berdy).

5,6-Dihydro: **Bastadin 5**. *Cyclobastadin 2*
 [79067-75-7]

$C_{34}H_{27}Br_5N_4O_8$ 1019.129

From *Ianthella basta*. Cryst. (MeCN) (as tetra-Me ether). Poorly sol. hexane. Mp 262-264° (tetra-Me ether). λ_{max} 208 (€ 126000); 280 (€ 7940) (MeOH) (Derep).

5,6-Dihydro, 6S-hydroxy: **Bastadin 8**

[127709-45-9]

$C_{34}H_{27}Br_5N_4O_9$ 1035.129

From *Ianthella basta*. Film. Fortuitously given the same trivial name by 2 independent groups. λ_{max} 208 (€ 126000); 280 (€ 7940) (MeOH) (Derep).

11-Bromo, 5,6-dihydro: **Bastadin 6**. *Cyclobastadin 3*

[79067-74-6]

$C_{34}H_{26}Br_6N_4O_8$ 1098.025

From *Ianthella basta* and *Psammaphysilla purpurea*. Antiangiogenic agent. Powder. Poorly sol. hexane. λ_{max} 220 (sh) (€ 50000); 281 (€ 5500) (MeOH) (Derep).

11-Bromo, 20,21-didehydro, 5,6-dihydro: **Bastadin 22**

$C_{34}H_{24}Br_6N_4O_8$ 1096.01

Isol. from *Dendrilla cactos*. Amorph. solid. Mp 198-202°.

11-Bromo, 25 ξ -hydroxy, 5,6-dihydro: **Bastadin 24**

$C_{34}H_{26}Br_6N_4O_9$ 1114.025

Isol. from *Ianthella quadrangulata*. Cytotoxic. Amorph. solid. $[\alpha]_D^{25}$ -36 (c, 0.88 in MeOH). λ_{max} 207 (€ 109225); 280 (€ 4873) (MeOH).

11-Bromo, 28-debromo, 20,21-didehydro, 5,6-dihydro: **Bastadin 23**

$C_{34}H_{25}Br_5N_4O_8$ 1017.114

Isol. from *Dendrilla cactos*. Amorph. solid. Mp 205-208°.

16-Debromo: **Bastadin 11**

[127687-09-6]

$C_{34}H_{26}Br_4N_4O_8$ 938.217

From *Ianthella basta*. Antiinflammatory agent. Film. λ_{max} 208 (€ 20000); 285 (€ 2510); 331 (€ 3160) (MeOH) (Derep).

16-Debromo, 5,6-dihydro: **Bastadin 9**†

[127687-07-4]

$C_{34}H_{28}Br_4N_4O_8$ 940.233

From *Ianthella basta*. Antiinflammatory agent. Powder (MeOH aq.). λ_{max} 208 (€ 126000); 279 (€ 10000) (MeOH) (Derep).

16-Debromo, 5,6-dihydro, 34-O-sulfate: **34-O-Sulfobastadin 9**

$C_{34}H_{28}Br_4N_4O_{11}S$ 1020.297

Isol. from *Ianthella basta*. Amorph. solid (as Na salt).

28-Debromo: **Bastadin 7**. *Cyclobastadin 4*

[79067-73-5]

$C_{34}H_{26}Br_4N_4O_8$ 938.217

From *Ianthella basta*. Foam. Poorly sol. hexane. λ_{max} 290 (€ 15100); 315 (€ 15500) (MeOH) (Derep).

28-Debromo, 15,34-disulfate: **15,34-Di-O-sulfatobastadin 7**

[184478-46-4]

$C_{34}H_{26}Br_4N_4O_{14}S_2$ 1098.346

From *Ianthella basta*. Calcium ion release inducer. Calcium ion channel modulator. Yellow solid (as di-Na salt). λ_{max} 207 (€ 82300); 321 (€ 12400) (MeOH) (di-Na salt).

28-Debromo, 6S-hydroxy, 5,6-dihydro: **Bastadin 10**

[127687-08-5]

$C_{34}H_{28}Br_4N_4O_9$ 956.233

From *Ianthella basta*. Antiinflammatory agent. Inosine 5'-phosphate dehydrogenase inhibitor. Oil. λ_{max} 208 (€ 126000); 279 (€ 10000) (MeOH) (Derep). λ_{max} 210 (€ 50000); 277 (€ 5000) (MeOH) (Berdy).

28-Debromo, 11-bromo: **Bastadin 14**. *Iso-bastadin 4*

[146345-73-5]

$C_{34}H_{25}Br_5N_4O_8$ 1017.113

Isol. from sponge *Psammaphysilla purpurea*. Dehydrofolate reductase inhibitor. λ_{max} 284 (€ 20000); 292 (€ 20000); 314 (€ 25000) (MeOH) (Berdy).

28-Debromo, 11-bromo, 5,6-dihydro: **Bastadin 15**

[149420-79-1]

$C_{34}H_{27}Br_5N_4O_8$ 1019.129

Metab. of a temperate marine sponge *Ianthella* sp. Small -ve opt. rotn.

28-Debromo, 11-bromo, 5,6-dihydro, 6S-hydroxy: **Bastadin 12**†. *Bastadin 9*

(*obsol.*)†

[150204-46-9]

$C_{34}H_{27}Br_5N_4O_9$ 1035.129

From *Ianthella basta*. Amorph. Renumbered in 1993. Name changed from Bastadin 9 to Bastadin 12 as the name Bastadin 9 was duplicated.

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1981, **34**, 765 (*isol. uv, ir, pmr, ms, cryst struct, derivs*)

Nishiyama, S. *et al.*, *Tet. Lett.*, 1982, **23**, 3699 (*synth*)

Miao, S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1441 (*isol, pmr, cmr, struct*)

Pordesimo, E.O. *et al.*, *J.O.C.*, 1990, **55**, 4704 (*isol, pmr, cmr, struct*)

Carney, J.R. *et al.*, *J. Nat. Prod.*, 1993, **56**, 153-157 (*nomencl*)

Dexter, A.F. *et al.*, *J. Nat. Prod.*, 1993, **56**, 782 (*Bastadin 15*)

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1995, **58**, 680-688 (*abs config*)

Franklin, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1121-1127 (*15,34-Disulfatobastadin 7*)

Guo, Z. *et al.*, *J.O.C.*, 1998, **63**, 4269-4276 (*Bastadin 6, synth*)

Masuno, M.N. *et al.*, *Mar. Drugs*, 2004, **2**, 176-184 (*34-Sulfobastadin 9*)

Kotoku, N. *et al.*, *Tetrahedron*, 2005, **61**, 7211-7218 (*Bastadin 6, synth*)

Reddy, A.V. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 4452-4457 (*Bastadins 6, 12, 15, 22, 23*)


Greve, H. *et al.*, *J. Nat. Prod.*, 2008, **71**, 309-312 (*Bastadin 24*)

Bastadin 13

B-33

Bastadin 12 (*obsol.*)†

[134981-78-5]



$C_{34}H_{28}Br_4N_4O_8$ 940.233

Originally isol. as Bastadin 12. Renumbered in 1993. Constit. of the sponge *Ianthella basta*. Powder. Mp 177-179°. λ_{max} 204 (€ 141000); 280 (€ 11200) (MeOH).

34-Sulfate: **34-Sulfobastadin 13**

[152213-67-7]

$C_{34}H_{28}Br_4N_4O_{11}S$ 1020.297

Constit. of an *Ianthella* sp. Endothelin A binding inhibitor; inhibitor of ATP citrate lyase. CAS no. refers to the Na salt. λ_{max} 204 (€ 74000); 280 (€ 4800) (MeOH).

9-Bromo: **Bastadin 19**

[158982-26-4]

$C_{34}H_{27}Br_5N_4O_8$ 1019.129

Isol. from *Ianthella basta*. Calcium channel modulator.

28-Debromo: **Bastadin 21**

$C_{34}H_{29}Br_3N_4O_8$ 861.337

Isol. from *Ianthella quadrangulata*. Amorph. solid. λ_{max} 195 (log € 4.56); 279 (log € 3.4); 384 (log € 2.99) (MeOH).

28-Debromo, 9-bromo: **Bastadin 20**

[184679-29-6]

$C_{34}H_{28}Br_4N_4O_8$ 940.233


Isol. from *Ianthella basta*. Calcium

channel modulator. Solid. λ_{\max} 209 (ε 84400); 279 (ε 5550) (MeOH).

- Butler, M.S. *et al.*, *Aust. J. Chem.*, 1991, **44**, 287-296 (*isol*, *pmr*, *cmr*)
 Carney, J.R. *et al.*, *J. Nat. Prod.*, 1993, **56**, 153-157 (*nomencl*)
 Gulavita, N.K. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1613-1617 (*34-sulfate*)
 Franklin, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1121-1127 (*Bastadin 19*, *Bastadin 20*)
 Coll, J.C. *et al.*, *J. Nat. Prod.*, 2002, **65**, 753-756 (*Bastadin 21*)

Bastadin 16

[156312-12-8]



$C_{34}H_{27}Br_5N_4O_8$ 1019.129
 Constit. of the sponge *Ianthella basta* and *Psammaphysilla purpurea*. λ_{\max} 280 (ε 3980) (MeOH) (Berdy).

6-Hydroxy, 11-debromo: Bastadin 17

[156312-13-9]

 $C_{34}H_{28}Br_4N_4O_9$ 956.233

Constit. of *Ianthella basta*. λ_{\max} 278 (ε 12600) (MeOH) (Berdy).

11,30-Bisdebro, 16-bromo: Bastadin 18

[157536-51-1]

 $C_{34}H_{28}Br_4N_4O_8$ 940.233


Constit. of *Ianthella basta*. Inosine 5'-phosphate dehydrogenase inhibitor. Amorph. solid.

Park, S.K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 407 (*isol*, *pmr*, *cmr*, *struct*)

Jaspars, M. *et al.*, *Tetrahedron*, 1994, **50**, 7367 (*Bastadin 18*)

Batrachamine**B-35**

3-Amino-1,2,5,6-tetrahydro-5-imino-2,6-pyrazinedibutanamine, 9CI. 2,6-Bis(4-aminobutyl)-3,5-diiminopiperazine [255865-64-6]


 $C_{12}H_{26}N_6$ 254.378

Drawn in the ref. as the diimino tautomer. Tautomer shown corresponds to the CAS name. *Isol.* from J774 cells. Cell regulation modulator. Inhibitor of sphingosine burst.

Warden, L.A. *et al.*, *J. Biol. Chem.*, 1999, **274**, 33875-33880 (*isol*, *pmr*, *cmr*)

Batrachotoxinin A

[19457-37-5]



Absolute Configuration

 $C_{24}H_{35}NO_5$ 417.544

Toxic principle from skin extracts of Colombian poison-dart frogs *Phylllobates aurotaenia*, *Phylllobates bicolor* and *Phylllobates terribilis*. Constit. of the skin and feathers of the toxic birds *Pitohui dichrous*, *Pitohui kirhocephalus* and *Ifrita kowaldi*, also melyrid beetles (*Choresine* sp.). Depolarizing agent for neurones and muscle cells *via* specific interaction with sodium channels. Noncryst. Mp 222-224° (as carbonate salt).

► Very toxic. LD₅₀ (mus, scu) 1 mg/kg. Avoid skin contact. CR4025000

 O^{20} -(4-Bromobenzoyl): Mp 213°.

O^{20} -(2E-Butenoyl): *Batrachotoxinin A trans-crotonate* [832099-84-0]

 $C_{28}H_{39}NO_6$ 485.619

Constit. of melyrid beetles (*Choresine* sp.), food of the toxic birds *Pitohui* spp. and *Ifrita kowaldi*.

O^{20} -(2Z-Butenoyl): *Batrachotoxinin A cis-crotonate* [320591-33-1]

 $C_{28}H_{39}NO_6$ 485.619

Constit. of feathers and skin of the toxic birds *Pitohui dichrous*, *Pitohui kirhocephalus* and *Ifrita kowaldi*, also melyrid beetles, a food of these birds (*Choresine* sp.).

O^{20} -Pentenoyl: *Batrachotoxinin A homo-crotonate*

 $C_{29}H_{39}NO_6$ 497.63

Constit. of melyrid beetles (*Choresine* sp.), a food of the toxic birds *Pitohui* spp. and *Ifrita kowaldi*. Full structure not assigned. 2004.

O^{20} -(3-Hydroxypentanoyl): [320600-82-6]

 $C_{29}H_{43}NO_7$ 517.661

Constit. of the feathers and skin of the toxic birds *Pitohui dichrous* and *Pitohui kirhocephalus*.

O^{20} -(2,4-Dimethyl-3-pyrrolicarboxylate): *Batrachotoxin* [23509-16-2]

 $C_{31}H_{42}N_2O_6$ 538.683

Alkaloid from poisonous dart frogs *Phylllobates aurotaenia*, *Phylllobates terribilis* and feathers and skin of the toxic birds *Pitohui dichrous*, *Pitohui kirhocephalus* and *Ifrita kowaldi*. Also constit. of melyrid beetles (*Choresine* sp.), food of *Pitohui* and *Ifrita* spp. Depolarising agent for neurones and muscle cells *via* specific interaction with sodium channels in plasma membranes. Noncryst.

► V. toxic, avoid skin contact. LD₅₀ (mus, ipr) 0.002 mg/kg. LD₅₀ (mus, scu) 0.002

B-36

mg/kg. Very toxic by intraperitoneal and subcutaneous routes. CR3990000

O^{20} -(2,4-Dimethyl-3-pyrrolicarboxylate), *N-de-Me*:

 $C_{30}H_{40}N_2O_6$ 524.656

Constit. of melyrid beetles (*Choresine* sp.) - a food of the toxic birds *Pitohui* spp. and *Ifrita kowaldi*. Full structure not confirmed, 2004. May be an artifact.

O^{20} -(2-Ethyl-4-methyl-3-pyrrolicarboxylate): *Homobatrachotoxin* [23509-17-3]

 $C_{32}H_{44}N_2O_6$ 552.709

Alkaloid from *Phylllobates aurotaenia* and *Phylllobates terribilis*. Also *isol.* from the feathers and skin of toxic birds *Pitohui dichrous*, *Pitohui kirhocephalus* and *Ifrita kowaldi*. Constit. of melyrid beetles (*Choresine* sp.), which are food of these birds. Noncryst.

► CR5075000

O^3 -Me, O^{20} -(2,4-dimethyl-3-pyrrolicarboxylate): *3-O-Methylbatrachotoxin*

 $C_{32}H_{44}N_2O_6$ 552.709

Alkaloid from *Phylllobates terribilis*. Probably an artifact. CAS no. not found 8-11CI.

O^3 -Me, O^{20} -(2-ethyl-4-methyl-3-pyrrolicarboxylate): *3-O-Methylhomobatrachotoxin*

 $C_{33}H_{46}N_2O_6$ 566.736

Alkaloid from *Phylllobates terribilis*. Probably an artifact.

O^3 -Et, O^{20} -(2-butenoyl):

 $C_{30}H_{43}NO_6$ 513.673

Constit. of melyrid beetles (*Choresine* sp.), a food of the toxic birds *Pitohui* spp. and *Ifrita kowaldi*. Full structure not confirmed, 2004. May be an artifact.

16,17-Dihydro, O^{20} -Ac:

 $C_{26}H_{39}NO_6$ 461.597

Constit. of feathers and skin of the toxic birds *Pitohui dichrous*, *Pitohui kirhocephalus* and *Ifrita kowaldi*. Also constit. of melyrid beetles (*Choresine* sp.), a food of the *Pitohui* spp. and *Ifrita kowaldi*. Incorrectly indexed in CA 134.

4β-Hydroxy, O^{20} -(2,4-dimethyl-3-pyrrolicarboxylate): *4-Hydroxybatrachotoxin*

[85439-39-0]

 $C_{31}H_{42}N_2O_7$ 554.682

Alkaloid from *Phylllobates terribilis*.

4β-Hydroxy, O^{20} -(2-ethyl-4-methyl-3-pyrrolicarboxylate): *4-Hydroxyhomobatrachotoxin*

[85439-40-3]

 $C_{32}H_{44}N_2O_7$ 568.709

Alkaloid from *Phylllobates terribilis*.

20-Deoxy, 20-chloro:

 $C_{24}H_{34}ClNO_4$ 435.99

Constit. of melyrid beetles (*Choresine* sp.), a food of the toxic birds *Pitohui* spp. and *Ifrita kowaldi*. Full structure not confirmed, 2004. May be an artifact.

Epimer: $C_{24}H_{35}NO_5$ 417.544


Constit. of feathers and skin of the

toxic bird *Pitohui dichrous*. Structure not elucidated. Unique CAS no not assigned by CA.

- Daly, J.W. *et al.*, *J.A.C.S.*, 1965, **87**, 124-126 (*isol, ms, ir, pmr*)
 Tokuyama, T. *et al.*, *J.A.C.S.*, 1969, **91**, 3931-3938 (*isol, ir, uv, pmr, ms, cryst struct*)
 Imhof, R. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 139-162 (*synth*)
 Witkop, B. *et al.*, *Alkaloids (Academic Press)*, 1983, **21**, 219-223 (*pharmacol*)
 Tokuyama, T. *et al.*, *Tetrahedron*, 1983, **39**, 41-47 (*Phyllobates terribilis constits, cmr*)
 Daly, J.W. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1986, **4**, 23 (*rev, pharmacol*)
 Dumbacher, J.P. *et al.*, *Science (Washington, D.C.)*, 1992, **258**, 799-801 (*Homobatrachotoxin, occur*)
 Deligeorges, S. *et al.*, *Recherches*, 1996, **284**, 50-51 (*Homobatrachotoxin, isol*)
 Kurosu, M. *et al.*, *J.A.C.S.*, 1998, **120**, 6627-6628 (*synth*)
 Weldon, P.J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **21**, 12948-12949 (*rev*)
 Dumbacher, J.P. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 12970-12975; 2004, **101**, 15857-15860 (*occur, birds, insects*)
 Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (*synth, rev*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, BAR750

Batzelladine A**B-37**

[147664-18-4]

C₄₂H₇₃N₉O₄ 768.097

Major component of complex where in addn. *n* = 9 and 10. Alkaloids from the Caribbean sponge *Batzella* sp. Inhibits the binding of HIVgp-120 to CD4. Amorph. powder. [α]_D²⁵ +8.9 (c, 2.3 in MeOH). λ_{max} 205 (ε 20100); 288 (ε 6820) (MeOH) (Derep).

[162047-80-5, 162047-79-2]

- Patil, A.D. *et al.*, *J.O.C.*, 1995, **60**, 1182 (*isol, uv, ir, pmr, cmr, struct*)
 Snider, B.B. *et al.*, *J.O.C.*, 1999, **62**, 1707-1711 (*config*)
 Franklin, A.S. *et al.*, *J.O.C.*, 1999, **64**, 1512-1519 (*config*)
 Shimokawa, J. *et al.*, *Chem. Eur. J.*, 2005, **11**, 6878-6888 (*synth*)
 Arnold, M.A. *et al.*, *J.A.C.S.*, 2006, **128**, 13255-13260 (*synth*)

Batzelladine B**B-38**

[161503-23-7]

As Batzelladine A, B-37 with *n* = 6, Δ^{24,32}, 25-epimer

C₄₀H₆₇N₉O₄ 738.027

Major component of complex where in addn. *n* = 7 and 8. Alkaloids from the Caribbean sponge *Batzella* sp. Inhibitor


of HIVgp-120-human CD4 binding. Interleukin binding inhibitor. Amorph. powder. Sol. H₂O; fairly sol. MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁵ +44.3 (c, 3.7 in MeOH). λ_{max} 206 (ε 16600); 289 (ε 9300); 340 (ε 4380) (MeOH) (Derep).

[162047-82-7, 162047-81-6]

- Patil, A.D. *et al.*, *J.O.C.*, 1995, **60**, 1182 (*isol, uv, ir, pmr, cmr, struct*)
 Franklin, A.S. *et al.*, *J.O.C.*, 1999, **64**, 1512-1519 (*synth, config*)

Batzelladine C**B-39**

[161503-24-8]

C₂₇H₄₈N₆O₂ 488.715


Major component of complex where in addn. *n* = 7 and 8. Alkaloids from the Caribbean sponge *Batzella* sp. Amorph. powder. [α]_D²⁵ -3.7 (c, 2.4 in MeOH). λ_{max} 211 (ε 8080); 230 (ε 7830); 298 (ε 6940) (MeOH) (Derep).

[162047-84-9, 162047-83-8]

- Patil, A.D. *et al.*, *J.O.C.*, 1995, **60**, 1182 (*isol, uv, ir, pmr, cmr, struct*)

Batzelladine D**B-40**

[161596-65-2]

C₂₅H₄₆N₆O₂ 462.677

Major component of complex where in addn. *n* = 9 and 10. Alkaloids from the Caribbean sponge *Batzella* sp. Amorph. powder. [α]_D²⁵ -1.2 (c, 0.9 in MeOH). λ_{max} 205 (ε 9170); 298 (ε 2380) (MeOH) (Derep).

30-Epimer: 30-Epibatzelladine D

[147664-20-8]

C₂₅H₄₆N₆O₂ 462.677


Alkaloid from a *Batzella* sp. Cytotoxic agent. Powder. Sol. MeOH, H₂O.

[162047-86-1, 162047-85-0]

- Pat. Coop. Treaty (WIPO)*, 1993, 93 1 193; *CA*, **118**, 225662x (*30-epimer*)
 Patil, A.D. *et al.*, *J.O.C.*, 1995, **60**, 1182 (*isol, uv, ir, pmr, cmr, struct*)
 Franklin, A.S. *et al.*, *J.O.C.*, 1999, **64**, 1512-1519 (*config*)
 Cohen, F. *et al.*, *Org. Lett.*, 1999, **1**, 2169-2172 (*synth*)
 Ishiwata, T. *et al.*, *Org. Lett.*, 2002, **4**, 2921-2924 (*synth*)
 Shimokawa, J. *et al.*, *Chem. Eur. J.*, 2005, **11**, 6878-6888 (*synth*)
 Arnold, M.A. *et al.*, *J.A.C.S.*, 2006, **128**, 13255-13260 (*synth*)
 Evans, P.A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 7417-7419 (*synth*)

Batzelladine E**B-41**

[161503-25-9]


C₂₇H₄₆N₆O₂ 486.699

Alkaloid from the Caribbean sponge *Batzella* sp. Protein kinase C and interleukin binding inhibitor. Gum. Sol. H₂O; fairly sol. MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁵ +87.1 (c, 1.9 in MeOH). λ_{max} 207 (ε 7954); 227 (ε 7012); 291 (ε 3311); 345 (ε 2876) (MeOH) (Berdy).

- Patil, A.D. *et al.*, *J.O.C.*, 1995, **60**, 1182 (*isol, uv, ir, pmr, cmr, struct*)
 Snider, B.B. *et al.*, *Tet. Lett.*, 1998, **39**, 5697-5700 (*synth, config*)

Batzelladine F**B-42**

[188112-82-5]

C₃₇H₆₄N₆O₂ 624.952

Struct. finalised in 2001. Alkaloid from the sponge *Monanchora arbuscula* (formerly *Batzella* sp.). Gum. [α]_D +19.4 (c, 0.87 in MeOH). λ_{max} 236; 254; 296 (MeOH).

21,22-Didehydro: Batzelladine NC₃₇H₆₂N₆O₂ 622.936

Alkaloid from *Monanchora unguifera*. Gum. [α]_D +8 (c, 0.1 in MeOH). C-18 and C-29 configs. not confirmed. λ_{max} 206 (ε 13340); 260 (ε 4320); 296 (ε 4184) (MeOH).

- Patil, A.D. *et al.*, *J.O.C.*, 1997, **62**, 1814-1819 (*isol, uv, ir, pmr, cmr, struct*)
 Snider, B.B. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1707-1711 (*config*)
 Black, G.P. *et al.*, *Tetrahedron*, 1999, **55**, 6547-6554 (*synth, pmr, cmr, config*)
 Cohen, F. *et al.*, *J.A.C.S.*, 2006, **128**, 2594-2603; 2604-2608 (*synth, struct*)
 Hua, H.-M. *et al.*, *Tetrahedron*, 2007, **63**, 11179-11188 (*Batzelladine N*)

Batzelladine G**B-43**

[188112-84-7]

As Batzelladine F, B-42 with

R = OH, m = 8, n = 7


C₃₉H₆₈N₆O₃ 669.005

Alkaloid from the sponge *Batzella* sp. Immunosuppressant. Powder. Sol. H₂O; fairly sol. MeOH; poorly sol. Me₂CO, hexane. [α]_D +14.7 (c, 0.79 in MeOH). λ_{max} 233; 258; 295 (MeOH).

- Patil, A.D. *et al.*, *J.O.C.*, 1997, **62**, 1814-1819 (*isol, uv, ir, pmr, cmr, struct*)

Batzelladine H

[188112-85-8]

 $C_{35}H_{57}N_6O_3^{\oplus}$ 609.874

Isol. as a 43:57 inseparable mixt. with Batzelladine I. Alkaloid from the sponge *Batzella* sp. Oil (as formate). Sol. H_2O ; fairly sol. MeOH; poorly sol. Me_2CO , hexane. $[\alpha]_D^{25} +33.7$ (c, 0.56 in MeOH) (formate). λ_{max} 217; 258 (MeOH).

Isomer: Batzelladine I

[188112-86-9]


 $C_{35}H_{57}N_6O_3^{\oplus}$ 609.874

From *Batzella* sp. Isomer of Batzelladine H in which the N-hydroxy group is located on the N between C-9 and C-10.

Patil, A.D. *et al.*, *J.O.C.*, 1997, **62**, 1814-1819 (isol, uv, ir, pmr, cmr, struct)

Batzelladine J

B-45

 $C_{41}H_{67}N_9O_4$ 750.038


Alkaloid from the sponge *Monanchora unguifera*. $[\alpha]_D^{25} -11.8$ (c, 0.13 in MeOH). λ_{max} 228 (ϵ 8800); 295 (ϵ 7200) (MeOH).

Gallimore, W.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1420-1423 (isol, pmr, cmr, ms)

Batzelladine K

B-46

[956260-43-8]

 $C_{15}H_{27}N_3$ 249.398

Alkaloid from *Monanchora unguifera*. Gum. $[\alpha]_D^{25} +6.4$ (c, 0.14 in MeOH). λ_{max} 208 (ϵ 2880) (MeOH).

Hua, H.-M. *et al.*, *Tetrahedron*, 2007, **63**, 11179-11188 (isol, cd, pmr, cmr)

Batzelladine L

B-47

[956260-45-0]

As Batzelladine F, B-42 with

R = H, m = 8, n = 7


 $C_{39}H_{68}N_6O_2$ 653.005

Alkaloid from *Monanchora unguifera*. Gum. $[\alpha]_D^{25} +5$ (c, 0.1 in MeOH). λ_{max} 212 (ϵ 18440) (MeOH).

Hua, H.-M. *et al.*, *Tetrahedron*, 2007, **63**, 11179-11188 (isol, cd, pmr, cmr)

Batzelladine M

[956260-47-2]

 $C_{35}H_{58}N_6O_2$ 594.882


Alkaloid from *Monanchora unguifera*. Gum. $[\alpha]_D^{25} +41.1$ (c, 0.09 in MeOH). λ_{max} 210 (ϵ 14600); 258 (ϵ 10203); 298 (ϵ 3530) (MeOH).

Hua, H.-M. *et al.*, *Tetrahedron*, 2007, **63**, 11179-11188 (isol, cd, pmr, cmr)

Batzellaside A

B-49

6-(2-Hydroxydodecyl)-2-(hydroxymethyl)-3,4-piperidinediol

 $C_{18}H_{37}NO_4$ 331.495

Alkaloid from the sponge *Batzella* sp. Oil. $[\alpha]_D^{25} +8.1$ (c, 0.44 in MeOH).

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**, 118-121 (isol, pmr, cmr)

Batzellaside B

B-50

2-(Hydroxymethyl)-6-(2-hydroxyundecyl)-3,4-piperidinediol

As Batzellaside A, B-49 with

n = 8

 $C_{17}H_{35}NO_4$ 317.468

Alkaloid from the sponge *Batzella* sp. Oil. $[\alpha]_D^{25} +10$ (c, 0.5 in MeOH).

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**, 118-121 (isol, pmr, cmr)

Batzellaside C

B-51

2-(Hydroxymethyl)-6-(2-hydroxytridecyl)-3,4-piperidinediol

As Batzellaside A, B-49 with

n = 10

 $C_{19}H_{39}NO_4$ 345.521

Alkaloid from the sponge *Batzella* sp. Oil. $[\alpha]_D^{25} +12$ (c, 0.4 in MeOH).


Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**, 118-121 (isol, pmr, cmr)

Batzelline A

B-52

6-Chloro-1,3,4,5-tetrahydro-1-methyl-2-(methylthio)pyrrolo[4,3,2-de]quinoline-7,8-dione, 9CI

[123064-89-1]

 $C_{12}H_{11}ClN_2O_2S$ 282.75

Alkaloid from the deep water Bahamas sponge *Batzella* sp. Black prisms ($CHCl_3/MeOH$). Mp 205°. λ_{max} 214 (ϵ 8900); 269 (ϵ 22100); 356 (ϵ 7900); 376 (ϵ 8000); 548 (ϵ 600) (MeOH) (Derep).

N-Me: Mp 202°.

N-De-Me: Batzelline B

[123064-90-4]

 $C_{11}H_9ClN_2O_2S$ 268.723

Isol. from *Batzella* sp. Herbicide. Dark brown solid. No definite Mp (dec.). λ_{max} 211 (ϵ 58300); 266 (ϵ 120800); 330 (sh); 387 (ϵ 51800); 550 (ϵ 4000) (MeOH) (Derep).

De(methylthio): Batzelline C

[123064-91-5]

 $C_{11}H_9ClN_2O_2$ 236.657

Isol. from *Batzella* sp. and *Zyzzya massalis*. Herbicide. Dark brown solid. No definite Mp (dec.). λ_{max} 211 (ϵ 7900); 250 (ϵ 17300); 331 (ϵ 9100); 380 (sh); 540 (ϵ 900) (MeOH) (Derep).

De(methylthio), N-de-Me: Batzelline D

[437980-22-8]

 $C_{10}H_7ClN_2O_2$ 222.63

Isol. from *Zyzzya fuliginosa* and *Zyzzya massalis*. Purple solid. λ_{max} 248 (log ϵ 4); 328 (log ϵ 3.73) (MeOH).

Sakemi, S. *et al.*, *Tet. Lett.*, 1989, **30**, 2517 (isol, uv, ir, pmr, cmr, ms, struct)

Tao, X.L. *et al.*, *Tetrahedron*, 1994, **50**, 2017 (Batzelline C, synth)

Yamada, F. *et al.*, *Heterocycles*, 1995, **41**, 1905 (Batzelline C, synth)

Roberts, D. *et al.*, *J.O.C.*, 1997, **62**, 568 (Batzelline C, synth)

Alvarez, M. *et al.*, *Eur. J. Org. Chem.*, 1999, 1173-1183 (synth)


Chang, L.C. *et al.*, *J. Nat. Prod.*, 2002, **65**, 776-778 (Batzelline D)

Dijou, M.-G. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 6035-6044 (Batzelline C, D)

Beccapoline

B-53

[85643-65-8]

 $C_{37}H_{24}N_2O_7$ 608.606

Alkaloid from *Polyalthia beccarii* and *Polyalthia cauliflora* var. *beccarii* (Annonaceae). Amorph. Mp 280° dec. λ_{max} 218 (sh) (log ϵ 4.52); 231 (log ϵ 4.63); 250 (sh) (log ϵ 4.53); 279 (log ϵ 4.4); 310 (log ϵ 4.01); 366 (sh) (log ϵ 4.07); 428 (log ϵ 4.22); 440 (log ϵ 4.21) (MeOH).

N⁶-Me: Beccapolinium

[85643-68-1]

 $C_{38}H_{27}N_2O_7^{\oplus}$ 623.64

Alkaloid from *Polyalthia beccarii* and *Polyalthia cauliflora* var. *beccarii* (Annonaceae). Mp 250° dec. (as hydroxide). CAS no. refers to hydroxide.

4',5'-Dihydro, 4',5'-dioxo: Beccapolydione
[91794-12-6]
C₃₇H₂₂N₂O₉ 638.589
Alkaloid from *Polyalthia cauliflora* var. *beccarii* (Annonaceae). Mp 280° dec. λ_{max} 218 (sh) (log ε 4.53); 224 (log ε 4.7); 250 (log ε 4.67); 276 (log ε 4.47); 314 (log ε 4.19); 325 (sh) (log ε 4.19); 370 (sh) (log ε 4.02); 438 (log ε 4.29) (MeOH).

8-Demethoxy: Polybeccarine
[91794-11-5]
C₃₆H₂₂N₂O₆ 578.58
Alkaloid from *Polyalthia cauliflora* var. *beccarii* (Annonaceae). Mp 280°. λ_{max} 216 (log ε 4.66); 234 (log ε 4.7); 248 (sh) (log ε 4.68); 273 (log ε 4.55); 308 (log ε 4.17); 344 (log ε 4.05); 406 (log ε 4.24); 425 (log ε 4.23) (MeOH).


Jossang, A. *et al.*, *Tet. Lett.*, 1982, **23**, 5147-5150 (*Beccapoline, struct*)
Jossang, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 504-513 (*Beccapolinium, Beccapolydione, Polybeccarine*)

Beilupeimine B-54

C₂₇H₄₃NO₃ 429.642
Steroidal alkaloid. Struct. unknown. Isol. from the Chinese drug Bei-lu-pei-mu. Mp 155-157°. [α]_D -52.23 (EtOH). λ_{max} 289 (no solvent reported).
Chu, T.T. *et al.*, *CA*, 1951, **51**, 445

Belemine B-55

[96400-75-8]




C₁₉H₁₇NO₃ 307.348
Alkaloid from the stem bark of *Gutteria schomburgkiana* (Annonaceae). Shows antimicrobial activity. Noncryst.

Cortes, D. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C, Ser. 2*, 1984, **299**, 311 (*uv, pmr, ms, struct*)
Villar, A. *et al.*, *Farm. Tijdschr. Belg.*, 1984, **61**, 300 (*activity*)

Belladonnine B-56

1,2,3,4-Tetrahydro-1-phenyl-1,4-naphthalenedicarboxylic acid bis(8-methyl-8-azabicyclo[3.2.1]oct-3-yl) ester, 9CI. Isatropyliditropein. 1,4-Ditropine isatropate
[510-25-8]



(1*RS*,4*RS*)-form

C₃₄H₄₂N₂O₄ 542.717
Alkaloid from *Mandragora officinarum* and *Orobanche aegyptiaca* (stereo undifferentiated). Also a thermal dimerisation prod. of Apoptropine, A-1359 and alkaloid occurrence may be as an artifact (Solanaceae, Orobanchaceae). Local anaesthetic potentiator.

(1*RS*,4*RS*)-form

cis-form. **β-Belladonnine. 1,4-Ditropine β-isatropate**
[6696-63-5]
Alkaloid from *Hyoscyamus niger* (Solanaceae). Mp 58°. *Hydrochloride* (1:2): Mp 195-196° dec. *Dipicrate*: Mp 230-232° dec. (anhyd.).


(1*RS*,4*SR*)-form

trans-form. **α-Belladonnine. 1,4-Ditropine α-isatropate**
[5878-33-1]
Alkaloid from *Hyoscyamus niger* (Solanaceae). Mp 129°. *Perchlorate* (1:2): Cryst. + 1H₂O. Mp 180° dec. *Dipicrate*: Mp 150° dec.

Voigtländer, H.W. *et al.*, *Annalen*, 1959, **625**, 196-205 (*struct, synth*)
Voigtländer, H.W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1959, **292**, 632 (*synth*)
Staub, H. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 2297-2305 (*isol*)
Wartman-Hafner, F. *et al.*, *Pharm. Acta Helv.*, 1966, **41**, 406-427 (*tlc*)
Jackson, B.P. *et al.*, *Phytochemistry*, 1973, **12**, 1165-1166 (*isol*)
Sharova, E.G. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 126-127; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 117-118 (*isol*)

Bellendine B-57

6,7,8,9-Tetrahydro-3,10-dimethylcyclohepta[b]pyran-5,8-imin-4(5H)-one, 9CI



Absolute Configuration

C₁₂H₁₅NO₂ 205.256

(+)-form [32152-73-1]

Alkaloid from the upper leaves, flowers and flowering stems of *Bellendena montana* (Proteaceae). Fine needles (Et₂O/petrol). Mp 162-163°. [α]_D¹⁹ +168 (CHCl₃).

2,3-Dihydro: 2,3-Dihydrobellendine

[72362-47-1]
C₁₂H₁₇NO₂ 207.272
Isol. from *Bellendena montana* (Proteaceae). Noncryst. Mp 188-190° (as picrate). [α]_D¹⁹ +59 (CHCl₃).

2,3-Dihydro, 3-epimer: 2,3-Epidihydrobellendine

[72401-75-3]
C₁₂H₁₇NO₂ 207.272
Isol. from *Bellendena montana* (Proteaceae). Noncryst. Mp 172-175° (as picrate). [α]_D¹⁹ +47 (CHCl₃).


(±)-form [52438-01-4]
Synthetic. Mp 128.5°.

Motherwell, W.D.S. *et al.*, *Chem. Comm.*, 1971, 133-134 (*cryst struct*)
Bick, I.R.C. *et al.*, *Phytochemistry*, 1971, **10**, 475-477 (*isol*)
Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 1827-1840 (*isol, pmr, cmr, ms, struct, synth, derivs*)

Bengacarboline

B-58

[192947-81-2]




C₂₉H₂₇N₅ 445.566
Isol. from the ascidian *Didemnum* sp. Cytotoxic agent. Brown solid. λ_{max} 226 (ε 6719); 284 (ε 1877) (MeOH).
Foderaro, T.A. *et al.*, *J.O.C.*, 1997, **62**, 6064-6065 (*isol, uv, ir, cd, pmr, cmr*)

Bengamide K

B-59

[193894-98-3]




C₂₂H₄₀N₂O₄ 396.569
Isol. from the sponge *Jaspis carteri*. [α]_D²⁵ +78.7 (c, 0.08 in MeOH).

D'Auria, M.V. *et al.*, *J. Nat. Prod.*, 1997, **60**, 814-816 (*isol, pmr, cmr, ms*)

Bengamide Z

B-60

[118477-10-4]



Absolute Configuration

C₁₈H₃₂N₂O₇ 388.46
Isol. from the sponge *Jaspis* cf. *coriacea* and from *Pachastrissa* sp. Cytotoxic agent; anthelmintic; nematocide; shows antiproliferative action. [α]_D²⁰ +45 (c, 0.11 in MeOH).

6-O-(3*R*,4*S*,5*R*-Trihydroxy-2*R*-methoxy-8-methyl-6-nonenoyl): Bengamide D

[118477-02-4]
C₂₉H₅₀N₂O₁₂ 618.72
From a Choriistid sponge (*Jaspidae*). Anthelmintic, nematocide. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ +19.8 (c, 0.086 in MeOH). Unstable in CDCl₃ soln.

6-O-Tridecanoyl: Bengamide H


[193894-95-0]
C₃₁H₅₆N₂O₈ 584.792
Isol. from *Jaspis carteri* and *Jaspis* cf. *coriacea*. [α]_D²⁵ +9.2 (c, 0.1 in MeOH).

- 6-O-(12-Methyltridecanoyl): **Bengamide O**
 $C_{32}H_{58}N_2O_8$ 598.819
 Isol. from *Jaspis cf. coriacea*. Oil. $[\alpha]_D^{20}$ +35.8 (c, 23.9 in MeOH).
- 6-O-Tetradecanoyl: **Bengamide B. NSC 646846**
 [104947-69-5]
 $C_{32}H_{58}N_2O_8$ 598.819
 Isol. from *Jaspis cf. coriacea* and *Pachastrissa sp.* Anthelmintic and cytotoxic agent. Exhibits antiparasitic and antimicrobial activity. Shows antiproliferative action. Viscous oil. $[\alpha]_D^{20}$ +34.6 (c, 0.075 in MeOH).
- 6-O-(13-Methyltetradecanoyl): **Bengamide M**
 $C_{33}H_{60}N_2O_8$ 612.846
 Isol. from *Jaspis cf. coriacea*. Oil. $[\alpha]_D^{20}$ +2.1 (c, 61.9 in MeOH).
- 6-O-Pentadecanoyl: **Bengamide J**
 [193894-97-2]
 $C_{33}H_{60}N_2O_8$ 612.846
 Isol. from *Jaspis carteri*. $[\alpha]_D^{25}$ +33 (c, 0.1 in MeOH).
- N-De-Me: **Bengamide Y**
 [118477-09-1]
 $C_{17}H_{30}N_2O_7$ 374.433
 Isol. from *Jaspis cf. coriacea*. Cytotoxic agent. Shows antiproliferative action. $[\alpha]_D^{20}$ +14 (c, 0.11 in MeOH).
- N-De-Me, 6S-(3R,4S,5R-trihydroxy-2R-methoxy-8-methyl-6-nonenoyl): **Bengamide C**
 [104975-72-6]
 $C_{28}H_{48}N_2O_{12}$ 604.693
 Constit. of a Choristid sponge (*Jaspidae*). Anthelmintic, nematocide. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Unstable in $CDCl_3$ soln.
- N-De-Me, 6-O-tridecanoyl: **Bengamide G**
 [193894-94-9]
 $C_{30}H_{54}N_2O_8$ 570.765
 Isol. from *Jaspis carteri* and *Jaspis cf. coriacea*. Shows antiproliferative action. $[\alpha]_D^{25}$ +14 (c, 0.1 in MeOH).
- N-De-Me, 6-O-(12-methyltridecanoyl): **Bengamide N**
 $C_{31}H_{56}N_2O_8$ 584.792
 Isol. from *Jaspis cf. coriacea*. Oil. $[\alpha]_D^{20}$ +20.7 (c, 38 in MeOH).
- N-De-Me, 6-O-tetradecanoyl: **Bengamide A. NSC 613012**
 [104947-68-4]
 $C_{31}H_{56}N_2O_8$ 584.792
 Isol. from *Jaspis cf. coriacea* and *Pachastrissa sp.* Anthelmintic, antiparasitic, antimicrobial and cytotoxic agent. Shows antiproliferative action. Cryst. (MeOH). Mp 114-115°. $[\alpha]_D^{20}$ +30.3 (c, 0.081 in MeOH).
- N-De-Me, 6-O-(13-methyltetradecanoyl): **Bengamide L**
 [226922-85-6]
 $C_{32}H_{58}N_2O_8$ 598.819
 Isol. from the sponges *Pachastrissa sp.* and *Jaspis cf. coriacea*. $[\alpha]_D^{20}$ +18.5 (c, 1.4 in MeOH).
- N-De-Me, 6-O-pentadecanoyl: **Bengamide I**
 [193894-96-1]
 $C_{32}H_{58}N_2O_8$ 598.819

- Isol. from *Jaspis carteri* and *Jaspis cf. coriacea*. $[\alpha]_D^{25}$ +32.4 (c, 0.7 in MeOH).
- 6-Deoxy: **Bengamide F**
 [118477-04-6]
 $C_{18}H_{32}N_2O_6$ 372.461
 Isol. from *Jaspis cf. coriacea*. Cytotoxic agent. Shows antiproliferative action. $[\alpha]_D^{20}$ +27.9 (c, 0.039 in MeOH).
- 6-Deoxy, 5'-O-tetradecanoyl: **Bengamide Q**
 $C_{32}H_{58}N_2O_7$ 582.819
 Isol. from *Jaspis cf. coriacea*. Oil. $[\alpha]_D^{20}$ +14.1 (c, 10.7 in MeOH).
- 6-Deoxy, N-de-Me: **Bengamide E**
 [118477-03-5]
 $C_{17}H_{30}N_2O_6$ 358.434
 Isol. from *Jaspis cf. coriacea* and *Pachastrissa sp.* Cytotoxic agent; shows antiproliferative action; anthelmintic; nematocide. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{20}$ +36.9 (c, 0.043 in MeOH).
- 6-Deoxy, N-de-Me, 5'-O-tetradecanoyl: **Bengamide P**
 $C_{31}H_{56}N_2O_7$ 568.793
 Isol. from *Jaspis cf. coriacea*. Oil. $[\alpha]_D^{20}$ +47.2 (c, 3.6 in MeOH).
- 6-Deoxy, N-de-Me, 5'-O-hexadecanoyl: **Bengamide R**
 $C_{33}H_{60}N_2O_7$ 596.846
 Isol. from *Jaspis cf. coriacea*. Oil (impure).
- Quiñó, E. et al., *J.O.C.*, 1986, **51**, 4494-4497 (isol, ir, pmr, cmr, ms, struct)
- Adamczeski, M. et al., *J.A.C.S.*, 1989, **111**, 647-654 (isol, ir, pmr, cmr, ms, struct)
- Adamczeski, M. et al., *J.O.C.*, 1990, **55**, 240-242 (abs config)
- Chida, N. et al., *Chem. Comm.*, 1992, 1064 (synth, abs config)
- Marshall, J.A. et al., *J.O.C.*, 1993, **58**, 6229-6234 (synth, bibl)
- D'Auria, M.V. et al., *J. Nat. Prod.*, 1997, **60**, 814-816 (*Bengamides G-J*)
- Fernandez, R. et al., *J. Nat. Prod.*, 1999, **62**, 678-680 (*Bengamide L*)
- Groweiss, A. et al., *J. Nat. Prod.*, 1999, **62**, 1691-1693 (*Bengamides Y,Z*)
- Thale, Z. et al., *J.O.C.*, 2001, **66**, 1733-1741 (*Bengamides M-Q*, isol, activity)
- Kinder, F.R. et al., *J.O.C.*, 2001, **66**, 2118-2122 (*Bengamides B,E*, synth)
- Banwell, M.G. et al., *J.O.C.*, 2001, **66**, 6768-6774 (*Bengamide E*, synth)
- Boeckman Jr., R.K. et al., *Org. Lett.*, 2002, **4**, 2109-2112 (synth)
- Liu, W. et al., *Tet. Lett.*, 2002, **43**, 1373-1375 (*Bengamide E*, synth)
- Sarabia, F. et al., *Tet. Lett.*, 2005, **46**, 1131-1135 (synth)

Bengazole Z**B-61**

[153549-05-4]




Absolute Configuration

- $C_{13}H_{18}N_2O_6$ 298.295
 Isol. from a *Jaspis sp.* sponge. Oil. $[\alpha]_D^{20}$ -7.5 (c, 0.17 in MeOH).
- O²-Tetradecanoyl: **Bengazole C₂**
 [153516-65-5]

- $C_{27}H_{44}N_2O_7$ 508.654
 Alkaloid from the Papua New Guinean marine sponge *Jaspis cf. coriacea*. Isol. as an inseparable mixture with 6 benzazoles C_3 , C_4 , C_6 , D_2 - D_4 .
- O²-(13-Methyltetradecanoyl): **Bengazole D₂**
 [153516-66-6]
 $C_{28}H_{46}N_2O_7$ 522.681
 From *Jaspis cf. coriacea*.
- O³-Tetradecanoyl: **Bengazole C₃**
 [153590-03-5]
 $C_{27}H_{44}N_2O_7$ 508.654
 From *Jaspis cf. coriacea*.
- O³-(13-Methyltetradecanoyl): **Bengazole D₃**
 [153590-04-6]
 $C_{28}H_{46}N_2O_7$ 522.681
 From *Jaspis cf. coriacea*.
- O⁴-Tetradecanoyl: **Bengazole C₄**
 [153590-05-7]
 $C_{27}H_{44}N_2O_7$ 508.654
 From *Jaspis cf. coriacea*.
- O⁴-(13-Methyltetradecanoyl): **Bengazole D₄**
 [153590-06-8]
 $C_{28}H_{46}N_2O_7$ 522.681
 From *Jaspis cf. coriacea*.
- O⁶-Tetradecanoyl: **Bengazole C₆**
 [153516-67-7]
 $C_{27}H_{44}N_2O_7$ 508.654
 From *Jaspis cf. coriacea*.
- O²-Pentadecanoyl: [226922-32-3]
 $C_{28}H_{46}N_2O_7$ 522.681
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -10.5 (c, 0.6 in MeOH). λ_{max} 209 (MeOH).
- O²-(14-Methylpentadecanoyl): [226922-28-7]
 $C_{29}H_{48}N_2O_7$ 536.707
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -12.9 (c, 0.2 in MeOH). λ_{max} 209 (MeOH).
- O⁶-Pentadecanoyl: [226922-37-8]
 $C_{28}H_{46}N_2O_7$ 522.681
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -9.3 (c, 1.8 in MeOH). λ_{max} 209 (MeOH).
- O⁶-(14-Methylpentadecanoyl): [226922-30-1]
 $C_{29}H_{48}N_2O_7$ 536.707
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -8.5 (c, 0.9 in MeOH). λ_{max} 209 (MeOH).
- O²-Hexadecanoyl: [226922-26-5]
 $C_{29}H_{48}N_2O_7$ 536.707
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -9.5 (c, 0.3 in MeOH). λ_{max} 209 (MeOH).
- O⁶-Hexadecanoyl: [226922-27-6]
 $C_{29}H_{48}N_2O_7$ 536.707
 Isol. from *Pachastrissa sp.* $[\alpha]_D^{20}$ -11.4 (c, 1.1 in MeOH). λ_{max} 209 (MeOH).
- O⁶-Heneicosanoyl: **Digonazole**
 [157536-06-6]
 $C_{34}H_{58}N_2O_7$ 606.841
 Metab. from the marine sponge *Jaspis digonoxea*. Viscous oil. $[\alpha]_D^{20}$ -8.9 (c, 2.2 in $CHCl_3$).
- Rodríguez, J. et al., *J. Nat. Prod.*, 1993, **56**, 2034-2040 (isol, pmr, cmr, struct)
- Rudi, A. et al., *J. Nat. Prod.*, 1994, **57**, 829-836 (*Digonazole*)
- Fernandez, R. et al., *J. Nat. Prod.*, 1999, **62**, 678-680 (*Pachastrissa Bengazoles*)
- Groweiss, A. et al., *J. Nat. Prod.*, 1999, **62**, 1691-1693 (isol)

Benhamycin

[1011287-15-2]


C₂₃H₁₈N₂O₄ 386.406

Prod. by *Streptomyces* sp. NR12. Anti-bacterial agent. Yellow solid. λ_{\max} 215 (log ϵ 3.52); 245 (log ϵ 3.61); 348 (log ϵ 3.33); 452 (log ϵ 3.21) (MeOH).

Shaaban, M. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1205-1211 (*isol*, *pmr*, *cmr*)

Beninine

6,21-Epoxy-17-methoxyaspidospermidine, 9CI [25480-48-2]



Absolute Configuration

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Hedranthera barteri* (Apocynaceae). Cryst. (EtOAc). Mp 225-227°.

N-Ac: Mp 220°. $[\alpha]_D^{25} +11$ (c, 0.047 in CHCl₃).

1,2-Didehydro: **1,2-Dehydrobeninine**. 1,2-Didehydrobeninine

[31146-53-9]

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Hedranthera barteri* (Apocynaceae).


Gorman, A.A. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 2072 (*ir*, *uv*, *pmr*, *ms*, *struct*)

Agwada, V. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1567 (*Dehydrobeninine*)

Kunesch, N. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 2854 (*cd*)

Benthocyanin A**B-64**

5-(3,7-Dimethyl-2,6-octadienyl)-2,5-dihydro-2-oxo-3-phenylfuro[2,3-b]phenazine-9-carboxylic acid, 9CI [133410-66-9]

C₃₁H₂₈N₂O₄ 492.573

Phenazine antibiotic. Prod. by *Strepto-*

B-62

myces prunicolor. Free radical scavenger. Antioxidant. Dark blue diamond shaped cryst. Mp 195-196°. λ_{\max} 246 (ϵ 30600); 340 (ϵ 18800); 635 (ϵ 16200); 690 (sh) (ϵ 10000) (MeOH/HCl) (Derep). λ_{\max} 248 (ϵ 31400); 341 (ϵ 17400); 410 (ϵ 8000); 429 (ϵ 8000); 616 (ϵ 18400); 660 (sh) (ϵ 7000) (MeOH/NaOH) (Derep). λ_{\max} 248 (ϵ 29000); 343 (ϵ 18600); 415 (ϵ 6100); 427 (ϵ 7400); 619 (ϵ 17200); 675 (sh) (ϵ 7000) (MeOH) (Derep).

11-Carboxy isomer: 5-(3,7-Dimethyl-2,6-octadienyl)-2,5-dihydro-2-oxo-3-phenylfuro[2,3-b]phenazine-11-carboxylic acid, 9CI. **Benthocyanin B**. BK 97B1. Antibiotic BK 97B1 [133683-28-0]

C₃₁H₂₈N₂O₄ 492.573

From *Streptomyces prunicolor*.

Antioxidant, free radical scavenger.

Dark blue plates. Mp 235-236°. λ_{\max} 246 (ϵ 30600); 340 (ϵ 18800); 635 (ϵ 16200); 690 (sh) (ϵ 10000) (MeOH/HCl) (Derep). λ_{\max} 248 (ϵ 31400); 341 (ϵ 17400); 410 (ϵ 8000); 429 (ϵ 8000); 616 (ϵ 18400); 660 (sh) (ϵ 7000) (MeOH/NaOH) (Derep). λ_{\max} 248 (ϵ 29000); 343 (ϵ 18600); 415 (ϵ 6100); 427 (ϵ 7400); 619 (ϵ 17200); 675 (sh) (ϵ 7000) (MeOH) (Derep). λ_{\max} 205 (ϵ 21900); 249 (ϵ 24400); 344 (ϵ 15900); 415 (ϵ 6100); 430 (ϵ 6100); 615 (ϵ 15700) (MeOH) (Berdy). λ_{\max} 208 (ϵ 34900); 250 (ϵ 26400); 343 (ϵ 17400); 415 (ϵ 6500); 430 (ϵ 6500); 615 (ϵ 17300) (MeOH/NaOH) (Berdy). λ_{\max} 209 (ϵ 34900); 250 (ϵ 26400); 343 (ϵ 17400); 415 (ϵ 6500); 430 (ϵ 6500); 615 (ϵ 17300) (MeOH/HCl) (Berdy).


Hayakawa, Y. *et al.*, *CA*, 1991, **114**, 225267 (*isol*)

Shin-ya, K. *et al.*, *Tet. Lett.*, 1991, **32**, 943 (*cryst struct*)

Shin-ya, K. *et al.*, *J.O.C.*, 1993, **58**, 4170 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Benthocyanin C**B-65**

3-(Cyanophenylmethylene)-5-(3,7-dimethyl-2,6-octadienyl)-2,3,5,10-tetrahydro-2-oxo-1-phenazinecarboxylic acid, 9CI. BK 97B2. Antibiotic BK 97B2 [133683-29-1]

C₃₁H₂₉N₃O₃ 491.588

Phenazine antibiotic. Prod. by *Streptomyces prunicolor*. Antioxidant. Free radical scavenger. Violet powder. Mp 157-158°. λ_{\max} 247 (ϵ 21100); 280 (ϵ 7800); 325 (ϵ 12100); 410 (ϵ 5300); 430 (ϵ 6400);


665 (ϵ 13100) (MeOH/HCl) (Derep). λ_{\max} 251 (ϵ 16300); 335 (ϵ 8000); 410 (ϵ 3600); 440 (ϵ 4400); 535 (ϵ 11600) (MeOH/NaOH) (Derep). λ_{\max} 246 (ϵ 18100); 283 (ϵ 11300); 445 (ϵ 5700); 470 (ϵ 5300); 570 (ϵ 11700) (MeOH) (Derep). λ_{\max} 208 (ϵ 19700); 246 (ϵ 18100); 283 (ϵ 11300); 445 (ϵ 5700); 470 (ϵ 5300); 570 (ϵ 11700) (MeOH) (Berdy). λ_{\max} 209 (ϵ 29500); 251 (ϵ 16300); 335 (ϵ 8000); 410 (ϵ 3600); 440 (ϵ 4400); 535 (ϵ 11600) (MeOH/HCl) (Berdy). λ_{\max} 247 (ϵ 21200); 280 (ϵ 7800); 325 (ϵ 12100); 410 (ϵ 5300); 430 (ϵ 6400); 665 (ϵ 13100) (MeOH/NaOH) (Berdy).

Hayakawa, Y. *et al.*, *CA*, 1991, **114**, 225267 (*isol*)

Shin-ya, K. *et al.*, *J.O.C.*, 1993, **58**, 4170 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Benthophoenin**B-66**

[152406-25-2]


C₃₇H₃₄N₂O₄ 570.687

Isol. from the mycelium of *Streptomyces prunicolor*. Free radical scavenger, haemolysis inhibitor. Red powder. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O. Mp 189-190°. λ_{\max} 234 (ϵ 28000); 309 (ϵ 36500); 388 (ϵ 7100); 520 (ϵ 12400) (MeOH) (Berdy).

Shin-ya, K. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1255 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

2-Benzamidobenzoic acid**B-67**

2-(Benzoylamino)benzoic acid, 9CI. N-Benzoylanthranilic acid. **Dianthramide B'** [579-93-1]

C₁₄H₁₁NO₃ 241.246

The naming system for dianthramides was changed in 1988. Isol. from *Dianthus caryophyllus* (carnation). Phytoalexin. Cryst. (EtOH aq.). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 183-184°. λ_{\max} 207; 245; 314 (MeOH) (Berdy). λ_{\max} 239; 281; 319 (MeOH/NaOH) (Berdy).

Me ester: [7510-49-8]

C₁₅H₁₃NO₃ 255.273

Cryst. (petrol). Mp 101°.


2'-Hydroxy: N-(2-Hydroxybenzoyl)anthranilic acid. (2-Hydroxybenzamido)benzoic acid. N-Salicylanthranilic

- acid. Dianthramide S*
[13316-98-8]
C₁₄H₁₁NO₄ 257.245
Constit. of *Dianthus caryophyllus*.
Phytoalexin. Cryst. (Me₂CO). Mp 219-220°.
- 2'-Hydroxy, Me ester: Dianthramide S methyl ester*
C₁₅H₁₃NO₄ 271.272
Constit. of *Delphinium staphisagria*.
Amorph.
- 4-Hydroxy: 2-Benzamido-4-hydroxybenzoic acid. N-Benzoyl-4-hydroxyanthranilic acid. 4-Hydroxydianthramide B*
[85915-70-4]
C₁₄H₁₁NO₄ 257.245
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 4-Hydroxy, Me ester: 4-Hydroxydianthramide B methyl ester*
C₁₅H₁₃NO₄ 271.272
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 2,4-Dihydroxy: 4-Hydroxy-N-(2-hydroxybenzoyl)anthranilic acid. 4-Hydroxy-N-salicylanthranilic acid. 4-Hydroxydianthramide S*
[115610-40-7]
C₁₄H₁₁NO₅ 273.245
Constit. of *Dianthus caryophyllus*.
Phytoalexin. Cryst. (EtOH). Mp 262-264°.
- 2,4-Dihydroxy, Me ester: Methyl 2-[(2-hydroxybenzoyl)amino]-4-hydroxybenzoate, 9CI. Methyl N-salicyl-4-hydroxyanthranilate. Hydroxydianthramide S methyl ester. Dianthramide B (obsol.)†*
[93289-91-9]
C₁₅H₁₃NO₅ 287.271
Constit. of *Dianthus caryophyllus*.
Phytoalexin. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{max} 206 ; 252 ; 277 ; 313 (MeOH) (Berdy). λ_{max} 219 ; 239 ; 269 ; 302 (MeOH/NaOH) (Berdy).
- 2',4'-Dihydroxy: N-(2,4-Dihydroxybenzoyl)anthranilic acid. N-Resorcy lanthranilic acid. Dianthramide R*
[115610-37-2]
C₁₄H₁₁NO₅ 273.245
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 2',4,4'-Trihydroxy: N-(2,4-Dihydroxybenzoyl)-4-hydroxyanthranilic acid. 4-Hydroxy-N-resorcy lanthranilic acid. 4-Hydroxydianthramide R*
[115610-38-3]
C₁₄H₁₁NO₆ 289.244
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 4-Methoxy: 2-Benzamido-4-methoxybenzoic acid. N-Benzoyl-4-methoxyanthranilic acid. 4-Methoxydianthramide B*
[109437-82-3]
C₁₅H₁₃NO₄ 271.272
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 4-Methoxy, Me ester: 4-Methoxydianthramide B methyl ester*
C₁₆H₁₅NO₄ 285.299
Constit. of *Dianthus superbus*. Cytoxin. Powder. λ_{max} 214 (log ε 4.21);

- 251 (log ε 4.44); 278 (log ε 4.22); 311 (sh) (log ε 4.01) (MeOH).
- 4-Methoxy, 2'-hydroxy: N-(2-Hydroxybenzoyl)-4-methoxyanthranilic acid. 4-Methoxy-N-salicylanthranilic acid. 4-Methoxydianthramide S. Dianthramide A (obsol.)†*
[93289-90-8]
C₁₅H₁₃NO₅ 287.271
Constit. of *Dianthus caryophyllus*.
Phytoalexin. Cryst. (Me₂CO). Mp 216-217°.
- 4-Methoxy, 2',4'-dihydroxy: N-(2,4-Dihydroxybenzoyl)-4-methoxyanthranilic acid. 4-Methoxy-N-resorcy lanthranilic acid. 4-Methoxydianthramide R*
[115610-39-4]
C₁₅H₁₃NO₆ 303.271
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- 4,4'-Dimethoxy, 2'-hydroxy: N-(2-Hydroxy-4-methoxybenzoyl)-4-methoxyanthranilic acid. 4-Methoxydianthramide M*
[115610-35-0]
C₁₆H₁₅NO₆ 317.298
Constit. of *Dianthus caryophyllus*.
Phytoalexin.
- Zentmeyer, D.T. *et al.*, *J.O.C.*, 1949, **14**, 967-981 (*synth*)
Ponchet, M. *et al.*, *Phytochemistry*, 1984, **23**, 1901-1903; 1988, **27**, 725-730 (*isol, uv, ir, pmr, ms*)
Hauteville, M. *et al.*, *J. Het. Chem.*, 1988, **25**, 715 (*synth, ir, pmr*)
Lee, C.K. *et al.*, *J.O.C.*, 1989, **54**, 3744-3747 (*synth, uv, ir, pmr*)
Hsieh, P.-W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1522-1527 (*4-Methoxydianthramide B Me ester*)
Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 733-739 (*Dianthramide S Me ester*)

Benzastatin A B-68

4-Amino-3-[3-(methoxymethyl)-6,7-dimethyl-2,6-octadienyl]benzamide, 9CI
[173429-75-9]



- C₁₉H₂₈N₂O₂ 316.442
Prod. by *Streptomyces nitrosporeus*. Free radical scavenger. Powder. Sol. MeOH, Me₂CO, CHCl₃, EtOAc, DMSO; poorly sol. H₂O. Mp 74°. λ_{max} 206 (ε 33880); 284 (ε 12880) (MeOH).
- Demethoxy: 4-Amino-3-(3,6,7-trimethyl-2,6-octadienyl)benzamide. Benzastatin B*
[173429-76-0]
C₁₈H₂₆N₂O 286.416
Prod. by *Streptomyces nitrosporeus*. Free radical scavenger. Powder. Sol. MeOH, CHCl₃, DMSO, Me₂CO, EtOAc; poorly sol. H₂O. Mp 98°. λ_{max} 207 (ε 32360); 282 (ε 13500) (MeOH).
- Demethoxy, 10'-hydroxy: Benzastatin H*

C₁₈H₂₆N₂O₂ 302.416
Prod. by *Streptomyces nitrosporeus*. Neuronal cell protectant. Powder. λ_{max} 206 (log ε 4.27); 286 (log ε 4.11) (no solvent reported).


Demethoxy, 11'-hydroxy: Benzastatin I
C₁₈H₂₆N₂O₂ 302.416
Prod. by *Streptomyces nitrosporeus*. Neuronal cell protectant. Powder. λ_{max} 206 (log ε 4.32); 286 (log ε 4.14) (no solvent reported).

Kim, W.-G. *et al.*, *J. Antibiot.*, 1996, **49**, 20-25; 26-30 (*Benzastatins A,B*)

Kim, W.-G. *et al.*, *J. Antibiot.*, 2001, **54**, 513-516 (*Benzastatins H,I*)

Benzastatin F B-69

[189309-08-8]



C₁₈H₂₆N₂O₂ 302.416
Prod. by *Streptomyces nitrosporeus*. Neuronal cell protecting agent. Powder. [α]_D¹⁸ +18 (c, 0.1 in MeOH). λ_{max} 206 (ε 19500); 230 (sh) ; 303 (ε 11000) (MeOH). λ_{max} 207 (ε 21300); 230 ; 303 (ε 11000) (MeOH) (Berdy).

16-Methoxy: Benzastatin E

[189309-07-7]
C₁₉H₂₈N₂O₃ 332.442
Prod. by *Streptomyces nitrosporeus*. Neuronal cell protecting agent. Powder. [α]_D¹⁸ +17 (c, 0.1 in MeOH). λ_{max} 207 (ε 21300); 230 (sh) ; 303 (ε 11000) (MeOH).

Kim, W.-G. *et al.*, *Tetrahedron*, 1997, **53**, 4309-4316 (*isol, uv, ir, pmr, cmr, ms*)

Toda, N. *et al.*, *Org. Lett.*, 2003, **5**, 269-271 (*synth*)

Benzastatin G B-70

[189309-09-9]

As Benzastatin F, B-69 with


R = H

C₁₇H₂₄N₂O₂ 288.389
Prod. by *Streptomyces nitrosporeus*. Neuronal cell protecting agent. Powder. [α]_D¹⁸ +23 (c, 0.1 in MeOH). λ_{max} 206 (ε 18000); 230 (sh) ; 301 (ε 8900) (MeOH).

Kim, W.-G. *et al.*, *Tetrahedron*, 1997, **53**, 4309-4316 (*isol, uv, ir, pmr, cmr, ms*)

Benz[glisoquinoline-5,10-dione B-71

Biquidone. 2-Azaanthraquinone. 3,4-Phthaloylpyridine. β-Anthrapyridinequinone
[46492-08-4]



C₁₃H₇NO₂ 209.204

Alkaloid from *Mitracarpus scaber* and *Psychotria camponutans*. Shows antimicrobial activity. Yellow needles (C₆H₆ or CCl₄). Mp 181–183° (179°). Insect teratogen. λ_{max} 209 ; 249 (MeOH) (Berdy).

N-Oxide: [21628-57-9]

C₁₃H₇NO₃ 225.203

Yellow cryst. (EtOH). Mp 263°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 462A (nmr)

Philips, A. et al., Ber., 1894, 27, 1923 (synth)

Étienne, A. et al., C. R. Hebd. Seances Acad. Sci. Ser. C, 1978, 267, 1826 (deriv)

Walton, B.T. et al., Science (Washington, D.C.), 1983, 222, 422 (tox)

Khanapure, S.S.P. et al., Heterocycles, 1988, 27, 2643 (synth)

Ohgaki, E. et al., J.C.S. Perkin I, 1990, 3109 (synth, ir, pmr)

Solis, P.N. et al., Planta Med., 1995, 61, 62 (isol, uv, ir, pmr, cmr, ms, struct)

Okunade, A.L. et al., Planta Med., 1999, 65, 447-448 (isol, cmr, activity)


Rebstock, A.-S. et al., Org. Biomol. Chem., 2004, 2, 291-295 (synth)

1,2-Benzisoxazole-3,6-diol **B-72**

6-Hydroxy-1,2-benzisoxazol-3(2H)-one, 9CI. 3,6-Dihydroxy-1,2-benzisoxazole.

Bactropone. Y-T 0678H. Antibiotic Y-T 0678H

[86004-57-1]

C₇H₅NO₃ 151.121

Exists in NH-form. Prod. by *Chromobacterium violaceum*. Active against gram-negative bacteria. Cryst. (EtOAc). Mp 248–250° dec. pK_{a1} 6.2; pK_{a2} 10.7 (2-methoxyethanol). λ_{max} 206 (€ 21100); 252 (€ 8230); 260 (sh); 280 (sh); 283 (€ 6340); 290 (€ 6340) (MeOH/HCl) (Derep). λ_{max} 220 (sh); 237 (€ 18100); 264 (€ 4380); 273 (sh); 296 (€ 5590) (MeOH/NaOH) (Derep). λ_{max} 220 (€ 18000); 251 (€ 6800); 259 (sh) (€ 6160); 273 (sh) (€ 4380); 279 (sh) (€ 5160); 283 (€ 6000); 289 (€ 5800) (MeOH) (Derep).


▶ DE4936000

[75661-62-0]

Japan. Pat., 1980, 80 102 597; CA, 93, 236929 (isol)

Hamada, M. et al., J. Antibiot., 1983, 36, 445

Imai, H. et al., J. Antibiot., 1983, 36, 911

Benzo[c]cinnoline-1,10-diol, 9CI **B-73**C₁₂H₈N₂O₂ 212.207


Di-Me ether: 1,10-Dimethoxybenzo[c]cinnoline. Antibiotic 4849F. 4849F

C₁₄H₁₂N₂O₂ 240.261Prod. by *Streptomyces* sp. 4849. Inhibitor of IL-4 receptor. Yellow needles. Mp 233–235°. λ_{max} 270 (MeOH).

Wang, K. et al., J. Antibiot., 2007, 60, 325-327 (di-Me ether, isol, pmr, cmr)

Benzomalvin B **B-74**

6,7-Dihydro-6-methyl-7-(phenylmethylene)quinazolino[3,2-a][1,4]benzodiazepine-5,13-dione, 9CI [157047-97-7]

C₂₄H₁₇N₃O₂ 379.417

Chiral molecule. Prod. by a *Penicillium* sp. Substance P inhibitor. Neurokinin receptor antagonist. Solid. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 260°. [α]_D +158 (c, 0.59 in MeOH). Related to Asperlicin C, A-1479. λ_{max} 234 (€ 19600); 269 (€ 7400); 280 (€ 6000); 312 (€ 3100); 323 (€ 2300) (MeOH) (Derep). λ_{max} 229 (€ 30000); 236 (€ 30100); 265 (€ 13800); 284 (€ 12700); 312 (€ 6500); 326 (€ 4700) (MeOH) (Berdy).

19α,20α-Epoxyde: **Benzomalvin C**

[157047-98-8]

C₂₄H₁₇N₃O₃ 395.417

Prod. by a *Penicillium* sp. Substance P inhibitor. Needles (MeOH aq.). Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D +69.7 (c, 0.38 in MeOH). λ_{max} 234 (€ 19600); 269 (€ 7400); 280 (€ 6000); 312 (€ 3100); 323 (€ 2300) (MeOH) (Derep). λ_{max} 229 (€ 27600); 273 (€ 6700); 283 (€ 5900); 312 (€ 2800); 323 (€ 2200) (MeOH) (Berdy).

19α,20-Dihydro: **Benzomalvin A**

[157047-96-6]

C₂₄H₁₉N₃O₂ 381.433

Prod. by a *Penicillium* sp. Substance P inhibitor. Solid. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 105–115°. [α]_D -106 (c, 1 in MeOH). λ_{max} 234 (€ 19600); 269 (€ 7400); 280 (€ 6000); 312 (€ 3100); 323 (€ 2300) (MeOH) (Derep).

19,20-Dihydro, atropisomer: **Benzomalvin D**

[173007-03-9]

C₂₄H₁₉N₃O₂ 381.433

Prod. by a *Penicillium* sp. Solid. Sol. MeOH, EtOAc. [α]_D +48. Benzomalvins A and D interconvert in soln. to form a 4:1 mixt. at r.t. and are separable by hplc. λ_{max} 234 ; 269 ; 280 ; 312 ; 323 (MeOH) (Berdy).

Sun, H.H. et al., J. Antibiot., 1994, 47, 515 (isol, pmr, cmr, uv, ir)

Sun, H.H. et al., J. Nat. Prod., 1995, 58, 1575 (Benzomalvin D, isol, pmr, cmr)

Sugimori, T. et al., Tetrahedron, 1998, 54, 7997-8008 (synth, Benzomalvin A)

Liu, J.-F. et al., J.O.C., 2005, 70, 10488-10493 (Benzomalvin A, synth)


Benzo[c][2,7]naphthyridin-4-ol **B-75**

Benzo[c][2,7]naphthyridin-4(3H)-one, 9CI. 4-Hydroxybenzo[c][2,7]naphthyridine. 2,9-Diazaphenanthren-1(2H)-one.

Perlolidine. Samoquasine A

[7344-61-8]

[62574-41-8 (Samoquasine A)]

C₁₂H₈N₂O 196.208

Samoquasine A finally shown to be identical with Perlolidine in 2008. Alkaloid from *Lolium perenne* and *Annona squamosa* (sugar apple). Mp 338–345° (332–334°). pK_{a1} 4.01; pK_{a2} 11.39 (18°). λ_{max} 239 (€ 50300); 243 (€ 49300); 252 (€ 45000); 322 (€ 11600); 337 (€ 13800) (EtOH aq.). λ_{max} 240 (€ 19400); 249 (€ 23500); 256 (€ 28000); 272 (€ 10700); 283 (€ 8800); 377 (€ 12200) (0.01M HCl). λ_{max} 237 ; 284 ; 325 ; 355 (0.01M NaOH).

OH-form

Me ether: 4-Methoxybenzo[c][2,7]-naphthyridine

[156780-62-0]

C₁₃H₁₀N₂O 210.235

Cryst. Mp 88° Mp 139–141°.

Et ether: 4-Ethoxybenzo[c][2,7]-naphthyridine

[399039-97-5]

C₁₄H₁₂N₂O 224.262

Beige cryst. Mp 104°.

Grimmett, R.E.R. et al., N.Z. J. Sci. Technol., Sect. B, 1943, 24, 151 (isol)

White, E.P. et al., N.Z. J. Sci. Technol., Sect. B, 1946, 27, 242

Powers, J.C. et al., J.A.C.S., 1968, 90, 7102 (synth, uv, ir, ms)

Lalezari, I. et al., J. Het. Chem., 1980, 17, 1761 (synth, ms)

Bracher, F. et al., Arch. Pharm. (Weinheim, Ger.), 1989, 322, 511 (synth, pmr, ms)

Rocca, P. et al., J.O.C., 1993, 58, 7832 (synth, pmr, ir)

Björk, P. et al., Heterocycles, 1997, 44, 237-253 (synth, Me ether)


Duvey, G. et al., J. Het. Chem., 2001, 38, 1039-1044 (Me ether, Et ether)

Yang, Y.-L. et al., Tet. Lett., 2003, 44, 319-322 (isol, pmr, cmr)

Timmons, C. et al., J.O.C., 2008, 73, 9168-9170 (Samoquasine A)

Benzosimuline **B-76**

[198336-58-2]




C₂₀H₁₉NO₂ 305.376

Alkaloid from the bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil. λ_{\max} 210 (log ϵ 4.19); 233 (log ϵ 4.42); 250 (sh) (log ϵ 4.05); 268 (log ϵ 3.95); 310 (sh) (log ϵ 3.74); 320 (sh) (log ϵ 3.79); 331 (log ϵ 3.86); 346 (log ϵ 3.94); 365 (log ϵ 3.81) (EtOH).

Chen, I.-S. *et al.*, *Phytochemistry*, 1997, **46**, 525-529 (*isol.*, *uv.*, *ir.*, *pmr.*, *cmr*)

Benzothiazole, 9CI

[95-16-9]

C₇H₅NS 135.189

Isol. from tail glands of the red deer *Cervus elaphus* and cranberries. Used in synth. of α -hydroxy carbonyl compounds. Sol. EtOH, CS₂, v. spar. sol. H₂O. Bp 223-225° Bp₂₅ 119-120°. pK_a 1.2 (H₂O). Steam-volatile.

- Fl. p. 112°. LD₅₀ (rat, orl) 466 mg/kg; LD₅₀ (mus, orl) 900 mg/kg. DL0875000 *Picrate*: Mp 176°.

N-Oxide: [27655-38-5]

C₇H₅NOS 151.189

Prisms + 1H₂O (MeOAc or Et₂O). Mp 45-48°.

Methiodide: [2786-31-4]

Needles. Mp 210°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 699C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 200A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1511A (*ir*)

Reid, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 143 (*synth*)

Ellis, B. *et al.*, *Spectrochim. Acta*, 1965, **21**, 1881 (*uv*)

Anjou, K. *et al.*, *Acta Chem. Scand.*, 1967, **21**, 2076-2082 (*isol*)

Takahashi, S. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1176 (*oxide*)

Witanowski, M. *et al.*, *Tetrahedron*, 1972, **28**, 637 (*nmr*)

Selva, A. *et al.*, *Org. Mass Spectrom.*, 1974, **9**, 1161 (*ms*)

Diehl, P. *et al.*, *Org. Magn. Reson.*, 1976, **8**, 155 (*pmr*)

Faure, R. *et al.*, *Can. J. Chem.*, 1978, **56**, 46 (*cmr*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1980, **8**, 274; 1990, **15**, 20 (*use*)

Angelilli, J.M. *et al.*, *Spectrosc. Lett.*, 1980, **13**, 741 (*ir*)

Bakke, J.M. *et al.*, *J. Chem. Ecol.*, 1983, **9**, 513


Bieräugel, H. *et al.*, *Tetrahedron*, 1983, **39**, 3971 (*synth*)

Bordwell, F.G. *et al.*, *J.A.C.S.*, 1991, **113**, 985 (*deriv.*, *synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BDE500

2,6-Benzothiazoleliol

2,6-Dihydroxybenzothiazole. 6-Hydroxy-2-(3H)-benzothiazolone
[80567-65-3]

C₇H₅NO₂S 167.188

Mp 233-235°.

6-Me ether: 6-Methoxy-2-(3H)-benzothiazolone
[40925-65-3]

C₇H₅NO₂S 167.188

Mp 163-165°.

NH-form

N-Me: 6-Hydroxy-3-methyl-2-(3H)-benzothiazolone
[96489-25-7]

C₈H₇NO₂S 181.215

Prod. by a *Micrococcus* sp. found in *Tedania ignis*. Solid (EtOH). Mp 182-183°.

Jones, G.H. *et al.*, *J. Med. Chem.*, 1987, **30**, 295 (*synth*)

Stierle, A.A. *et al.*, *Tet. Lett.*, 1991, **32**, 4847 (*isol.*, *deriv*)

Delhomel, J.F. *et al.*, *J. Het. Chem.*, 2001, **38**, 633-639 (*N-Me*)

2(3H)-Benzothiazolethione,**9CI**

2-Benzothiazolethiol. 2-Mercaptoben-zothiazole. 2-Benzothiazolinethione. Accelerator M. Accel M. Captax. Kaptax. Pneumax MBT. Thiotax. Vulkacit M. Wobezit M
[149-30-4]

C₇H₅NS₂ 167.255

NH-form predominates. *Isol.* from a *Micrococcus* sp. in *Tedania ignis*. Also present in cranberries. Vulcanisation accelerator. Used as 1% CHCl₃ soln. for separation pptn. of Au, I[⊖]; photometric detn. of Pd, Os, Rh, Se, Ni (λ_{\max} 750 nm). Needles (MeOH aq.). Spar. sol. EtOH, Et₂O, AcOH; insol. H₂O; sol. alkalis. Mp 177-179°. Readily oxidised to the disulfide.

- LD₅₀ (rat, orl) 100 mg/kg. Exp. carcinogen. Exp. reprod. and teratogenic effects. DL6475000

[2492-26-4 ; 37437-20-0 ; 155-04-4]

Chesick, J.P. *et al.*, *Acta Cryst. B*, 1971, **27**, 1441 (*cryst struct*)

DeJongh, D.C. *et al.*, *Adv. Mass Spectrom.*, 1974, **6**, 99 (*ms*)

Faure, R. *et al.*, *Org. Magn. Reson.*, 1978, **11**, 617 (*cmr*, *tautom*)


Cristiani, F. *et al.*, *Phosphorus Sulfur Relat. Elem.*, 1984, **20**, 231 (*ir*)

Radha, A. *et al.*, *Z. Kristallogr., Kristallgeom., Kristallphys., Kristallchem.*, 1985, **171**, 225 (*cryst struct*)

Stierle, A.A. *et al.*, *Tet. Lett.*, 1991, **32**, 4847-4848 (*isol*)

2-Benzothiazolol


2-(3H)-Benzothiazolone, 9CI. 2-Benzothiazolinone. 2-Hydroxybenzothiazole

C₇H₅NOS 151.189

NH-form predominates. Prod. by a *Micrococcus* sp. found in *Tedania ignis*. Gives colour reaction with Pd (λ_{\max} 368 nm). Needles (EtOH). Mp 138-139°. Faure, R. *et al.*, *Org. Magn. Reson.*, 1978, **11**, 617 (*tautom*, *cmr*)
Stierle, A.A. *et al.*, *Tet. Lett.*, 1991, **32**, 4847 (*isol*)

2,4-Benzoxazoleliol

4-Hydroxy-2-(3H)-benzoxazolone, 9CI. 4-Hydroxy-2-benzoxazolinone. 2,4-Dihydroxybenzoxazole
[28955-70-6]

C₇H₅NO₃ 151.121

Exists in NH-form. Needles (H₂O). Mp 281-283° dec.

4-O- β -D-Glucopyranoside:C₁₃H₁₅NO₈ 313.263


Constit. of *Acanthus ilicifolius*. Pale yellow solid. Mp 131-133°. λ_{\max} 228 ; 265 (MeOH).

Zinner, H. *et al.*, *Chem. Ber.*, 1960, **93**, 1331-1339 (*synth*)

Huo, C. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 343-345 (*glucoside*)

2,6-Benzoxazoleliol

6-Hydroxy-2-(3H)-benzoxazolone, 9CI. 6-Hydroxy-2-benzoxazolinone. 2,6-Dihydroxybenzoxazole
[78213-03-3]

C₇H₅NO₃ 151.121

Exists in NH-form. Alkaloid from *Acanthus arboreus*. Needles (H₂O) or pale yellow cryst. Mp 288-292° (260°). λ_{\max} 268 ; 302 ; 305 ; 340 (MeOH).

6-Me ether: 2-Hydroxy-6-methoxybenzoxazole. 6-Methoxy-2-(3H)-benzoxazolone. 6-Methoxy-2-benzoxazolinone. 6-Methoxy-2-benzoxazolol. MBOA. Coixol
[532-91-2]

C₈H₇NO₃ 165.148

Isol. from roots of *Coix lacryma jobi* (Job's tears) (Poaceae) and from seedlings of several cereal plants, e.g. wheat (*Triticum aestivum*), corn (*Zea mays*) and rye (*Secale cereale*). Also present in all plant parts of *Scoparia dulcis* (Scrophulariaceae) and *isol* from a

marine sponge *Oceanapia* sp. Disease and insect attack inhibitor in plants. Toxic to brine shrimp. Pink cryst. (MeOH aq.). Sol. bases, EtOH, Et₂O, C₆H₆; fairly sol. H₂O; poorly sol. hexane. Mp 160–161° (154–155°). λ_{max} 229 ; 286 (H₂O) (Berdy).

▶ DM5275200

NH-form

N-Me: [359434-29-0]

C₈H₇NO₃ 165.148

Solid (EtOH). Mp 203–204°.

6-Me ether, N-Ac: [62655-08-7]

C₁₀H₉NO₄ 207.185

Needles. Mp 147.5°.

6-Me ether, N-benzoyl:

C₁₅H₁₁NO₄ 269.256

Needles. Mp 162–162.5°.

Koyama, T. *et al.*, *Yakugaku Zasshi*, 1955, **75**, 699–701; *CA*, **50**, 34029 (6-Me ether)

Beck, S.D. *et al.*, *J. Agric. Food Chem.*, 1957, **5**, 933–935 (6-Me ether, detn)

Smissman, E.E. *et al.*, *J.A.C.S.*, 1957, **79**, 4697–4698 (6-Me ether, isol, synth)

Hietala, P.K. *et al.*, *Acta Chem. Scand.*, 1958, **12**, 119–123 (detn)

Wahlroos, O. *et al.*, *Acta Chem. Scand.*, 1958, **12**, 124–128 (activity, 6-Me ether)

List, P.H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1959, **292**, 452–456 (6-Me ether)

Zinner, H. *et al.*, *Chem. Ber.*, 1960, **93**, 1331–1339 (synth)

Allen, E.H. *et al.*, *J.O.C.*, 1971, **36**, 2004–2005 (6-Me ether)

Richey, J.D. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 2413–2416 (6-Me ether, synth)

Kubo, I. *et al.*, *Experientia*, 1983, **39**, 355 (6-Me ether, synth)

Cole, E.R. *et al.*, *Aust. J. Chem.*, 1986, **39**, 295–301 (synth, ir, pmr)

Sicher, D. *et al.*, *Synthesis*, 1989, 875–876 (6-Me ether)

Maleski, R.J. *et al.*, *J. Het. Chem.*, 1991, **28**, 1937–1939 (synth)

Bjostad, L.B. *et al.*, *J. Chem. Ecol.*, 1992, **18**, 931–944 (6-Me ether, isol)

Hayashi, T. *et al.*, *Phytochemistry*, 1994, **37**, 1611–1614 (6-Me ether, occur)

Venkateswarlu, Y. *et al.*, *Biochem. Syst. Ecol.*, 1999, **27**, 519–520 (6-Me ether, isol, activity)

Delhomel, J.F. *et al.*, *J. Het. Chem.*, 2001, **38**, 633–639 (N-Me)

Amer, M.E. *et al.*, *J. Braz. Chem. Soc.*, 2004, **15**, 262–266 (isol)


2-Benzoxazolol

B-83

2-(3H)-Benzoxazolone, 9CI. 2-Benzoxazolinone, 8CI. 2-Hydroxybenzoxazole.

BOA

[59-49-4]



C₇H₅NO₂ 135.122

Tautomeric. NH-form predominates.

Found in rye seedlings. Sol. bases, Et₂O; fairly sol. H₂O; poorly sol. hexane. Mp 141–142°. pK_{a1} 8.9 (23°). λ_{max} 270 (H₂O) (Berdy). λ_{max} 275 (EtOH) (Berdy).

▶ LD₅₀ (rat, orl) 700 mg/kg. LD₅₀ (mus, orl) 554 mg/kg. DM4905000

Thompson, M.L. *et al.*, *Can. J. Chem.*, 1973, **51**, 3313 (ms)

Linares, J. *et al.*, *Can. J. Chem.*, 1979, **57**, 937 (cmr, tautom)

Barnes, J.P. *et al.*, *Phytochemistry*, 1987, **26**, 1385–1390 (isol)

Maleski, R.J. *et al.*, *J. Het. Chem.*, 1991, **28**, 1937 (synth)

El-Faham, A. *et al.*, *J. Het. Chem.*, 2006, **43**, 599–606 (synth, ir, pmr, cmr, ms)

Benzylamine, 8CI

B-84

Benzenemethanamine, 9CI. α-Aminotoluene. Moringine

[100-46-9]

PhCH₂NH₂

C₇H₉N 107.155

Alkaloid from *Moringa oleifera* (horseradish tree) (Moringaceae). Reagent for gc anal. of alkyl isocyanates. d₄¹⁹ 0.98. Bp 185° Bp₁₂ 90°. n_D²⁰ 1.5401. pK_a 8.82 (20°, 60% dioxan aq.). Absorbs CO₂ from air.

▶ Fl. p. 63°. Reacts violently with N-haloimides. Corrosive and irritating to all tissues. DP1488500

Hydrochloride: [3287-99-8]

Mp 255–257°.

▶ DP5425000

Hydrobromide: [37488-40-7]

Mp 204°.

Hydroiodide: [45579-91-7]

Leaflets. Mp 162°.

Picrate: Mp 194°.

Sulfate: [2208-34-6]

Needles. Mp 93°.

N-Ac: N-(Phenylmethyl)acetamide, 9CI

[588-46-5]

C₉H₁₁NO 149.192

Leaflets (petrol). Mp 61°.

▶ AB4546300

N-Benzoyl: N-Benzylbenzamide

[1485-70-7]

C₁₄H₁₃NO 211.263

Constit. of the stems of *Salvadora persica*. Leaflets. Mp 105–106°.

N-tert-Butyloxycarbonyl:

C₁₂H₁₇NO₂ 207.272

Needles (petrol). Mp 53.5–55°.

N-(4-Methylbenzenesulfonyl):

C₁₄H₁₅NO₂S 261.344

Mp 185°.

N-Me: [103-67-3]

C₈H₁₁N 121.182

Minor constit. of *Ma Huang* (from *Ephedra* sp.). Bp 180–181° Bp₁₄ 78°.

N-Me; hydrochloride: [13426-94-3]

Needles (EtOH/Et₂O). Mp 178°.

▶ DP5950000

N-Me, N-nitroso: Benzylmethylnitrosamine

[937-40-6]

C₈H₁₀N₂O 150.18

Liq. Bp₁₈ 145° Bp_{0.3} 80°. n_D^{31.5} 1.5388.

N,N-Di-Me: [103-83-3]

C₉H₁₃N 135.208

Liq. Bp 181° Bp₁₅ 66–67°. Steam-volatile.

▶ Corrosive and irritating to all tissues. LD₅₀ (rat, orl) 265 mg/kg. LD₅₀ (rbt, skp) 1660 mg/kg. Flammable, fl. p. 54°. DP4500000

N,N-Di-Me; hydrochloride: [1875-92-9]

Mp 175°.

▶ DP4700000

N,N-Di-Et: [772-54-3]

C₁₁H₁₇N 163.262

Bp 211–212° Bp₁₅ 94°.

▶ Fl. p. 77° (oc).

N-Propyl: [2032-33-9]

C₁₀H₁₅N 149.235

Oil. Bp 214–216° Bp₁₀₀ 129° Bp_{0.2} 40–42°.

N-Isopropyl: N-Benzylisopropylamine

[102-97-6]

C₁₀H₁₅N 149.235

Bp 200°.

N-Butyl: [2403-22-7]

C₁₁H₁₇N 163.262

Bp₃ 87–89°. n_D²⁰ 1.5010.

N-tert-Butyl: [3378-72-1]

C₁₁H₁₇N 163.262

Bp₃ 80°.

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 1266B; 1266C; 1274A; 1274C;

1274D; 1275A; 1279B; 2, 3, 666B (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **2**, 565B; 565C; 581B; 582A;

582B; 582C; 589A; 1385A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase,

1989, **3**, 1163A; 1165D; 1166B; 1166C;

1167D (ir)

Chen, A.L. *et al.*, *J. Am. Pharm. Assoc.*, 1931,

20, 339 (isol, derivs)

Wegler, R. *et al.*, *Ber.*, 1936, **69**, 2074 (N,N-di-

Et)

Chakravarti, R.N. *et al.*, *CA*, 1955, **50**, 16891f

(isol)

Looney, C.E. *et al.*, *J.A.C.S.*, 1957, **79**, 6136

(N-Me N-nitroso, synth, pmr, ir)

Leyson, R. *et al.*, *Spectrochim. Acta*, 1963, **19**,

243 (ir)

Fieser and Fieser's Reagents for Organic

Synthesis, Wiley, 1967, **1**, 51; 1975, **5**, 479;

1980, **8**, 70; 1982, **10**, 26 (use)

Witanowski, M. *et al.*, *Can. J. Chem.*, 1969, **47**,

1321 (N-15 nmr)

Egli, R.A. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 47

(synth)

Aihara, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1972,

45, 1942 (pmr)

Lichter, R.L. *et al.*, *J.A.C.S.*, 1972, **94**, 2495

(N-15 nmr)

Shapiro, M.J. *et al.*, *J.O.C.*, 1976, **41**, 3197

(cmr)

Kostyanovsky, R.G. *et al.*, *Org. Mass*

Spectrom., 1976, **11**, 237 (ms)

Knights, R.J. *et al.*, *Anal. Biochem.*, 1977, **77**,

106 (use)

Bastaert, G. *et al.*, *C. R. Hebd. Seances Acad.*

Sci. Ser. C, 1980, **290**, 193 (uv)

Bradamante, S. *et al.*, *J.O.C.*, 1980, **45**, 105

(cmr)

Garrido, D.O.A. *et al.*, *J.O.C.*, 1984, **49**, 2021

(N-Alkyl derivs, synth, cmr)

Boyer, J.H. *et al.*, *Synthesis*, 1985, 677–679 (N-

Me N-nitroso)

Elmore, D.T. *et al.*, *Acta Cryst. C*, 1992, **48**,

1331 (cryst struct)

Ding, C.Z. *et al.*, *J. Med. Chem.*, 1993, **36**,

1711 (N,N-di-Me)

Bhattacharyya, S. *et al.*, *Synth. Commun.*,

1995, **25**, 2061 (N,N-di-Me)

Kanie, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1998,

71, 1973–1991 (N-propyl)

Collins, C.J. *et al.*, *J.O.C.*, 1999, **64**, 2574–2576

(N,N-di-Me, synth, pmr, cmr)

Degani, I. *et al.*, *Synthesis*, 1999, 1200–1208

(N-benzoyl)

Gibson, S.E. *et al.*, *Org. Biomol. Chem.*, 2003,

1, 676–683 (N-boc, synth, ir, pmr)

Khalil, A.T. *et al.*, *Arch. Pharmacol. Res.*, 2006,

29, 952–956 (N-Benzylbenzamide, isol, pmr,

cmr)


Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 2631

Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 122

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BDY000; DQQ000; DQP800

5-Benzyl-2,5-dihydro-4-hydroxy-3-(2,4,6-octatrienyl)-1H-pyrrol-2-one **B-85**

2,5-Dihydro-4-hydroxy-3-(1-oxo-2,4,6-octatrienyl)-5-(phenylmethyl)-1H-pyrrol-2-one



C₁₉H₁₉NO₃ 309.364

Enolised β-tricarbonyl compd. Isol. from *Leocarpus fragilis*. No phys. props. reported.


N-Me: 2,5-Dihydro-4-hydroxy-1-methyl-3-(1-oxo-2,4,6-octatrienyl)-5-(phenylmethyl)-1H-pyrrol-2-one

C₂₀H₂₁NO₃ 323.391

Isol. from *Leocarpus fragilis*.

Steglich, W. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 281-288

4-Benzyl-1,4-dihydro-3-oxo-3H-pyrrolo[2,1-c][1,4]oxazine-6-carboxaldehyde **B-86**




C₁₅H₁₃NO₃ 255.273

Alkaloid from the fruit of *Celastrus orbiculatus*. Cryst. (CHCl₃). Mp 112-114°. λ_{max} 291 (MeOH).

Guo, Y. *et al.*, *Fitoterapia*, 2005, **76**, 273-275 (isol, pmr, cmr)

5-Benzyl-3-(2,4,6,8,10-dodecapentaenyl)-2,5-dihydro-4-hydroxy-1H-pyrrol-2-one **B-87**

2,5-Dihydro-4-hydroxy-3-(1-oxo-2,4,6,8,10-dodecapentaenyl)-5-(phenylmethyl)-1H-pyrrol-2-one




C₂₃H₂₃NO₃ 361.44

Enolised β-tricarbonyl compd. Isol. from *Leocarpus fragilis*. No phys. props. reported.

Steglich, W. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 281-288

5-Benzyl-3-hydroxy-2-isopropylpyrazine **B-88**

3-(1-Methylethyl)-6-(phenylmethyl)-2(1H)-pyrazinone. 6-Benzyl-3-isopropyl-2(1H)-pyrazinone. 6-Benzyl-3-isopropyl-2-pyrazinol. **Phevalin** [170713-71-0]




C₁₄H₁₆N₂O 228.293

Prod. by *Streptomyces* sp. SC433. Calpain inhibitor. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{max} 205 (ε 2860); 325 (ε 1296) (MeOH).

Alvarez, M.E. *et al.*, *J. Antibiot.*, 1995, **48**, 1165 (isol, uv, ir, pmr, cmr, ms)

2-Benzyl-3-hydroxy-1-methyl-5-nonylpyrrolidine **B-89**

1-Methyl-5-nonyl-2-(phenylmethyl)-3-pyrrolidinol, 9CI [119463-16-0]



C₂₁H₃₅NO 317.514

(2S,3S,5R)-form

Preussin. L 657398. Antibiotic L 657398 [125356-66-3]

Isol. from a *Preussia* sp. and *Aspergillus ochraceus*. Broad-spectrum antifungal agent. Yellow oil. [α]_D²⁵ +22 (c, 1 in CHCl₃) (natural). [α]_D²⁵ +31.1 (c, 1 in CHCl₃) (synthetic). λ_{max} 203 (ε 12900) (MeOH) (Derep).

(2R,3R,5S)-form

Synthetic. Yellowish wax-like solid. [α]_D²⁶ -21.6 (c, 1.0 in CHCl₃).

Schwartz, R.E. *et al.*, *J. Antibiot.*, 1988, **41**, 1774 (L 657398)

Johnson, J.H. *et al.*, *J. Antibiot.*, 1989, **42**, 1184 (*Preussin*)

Pak, C.S. *et al.*, *J.O.C.*, 1991, **56**, 1128 (synth) Shimazaki, M. *et al.*, *Heterocycles*, 1993, **36**, 1823 (synth)

Deng, W. *et al.*, *J.A.C.S.*, 1994, **116**, 11241 (synth, bibl)

Overhand, M. *et al.*, *J.O.C.*, 1994, **59**, 4721 (synth)

Verma, R. *et al.*, *Chem. Comm.*, 1997, 1601 (synth)

Kadota, I. *et al.*, *Heterocycles*, 1997, **46**, 335-348 (synth)

Beier, C. *et al.*, *Synthesis*, 1997, 1296-1300 (synth)

Kanazawa, A. *et al.*, *J.O.C.*, 1998, **63**, 4660-4663 (synth)

De Armas, P. *et al.*, *Tet. Lett.*, 1998, **39**, 131-134 (synth, bibl)

Veerasa, G. *et al.*, *Tetrahedron*, 1998, **54**, 15673-15678 (synth)

Verma, R. *et al.*, *J.C.S. Perkin 1*, 1999, 265-270 (synth, pmr, cmr)

Lee, K.-Y. *et al.*, *Org. Lett.*, 2000, **2**, 4041-4042 (synth)

Krasinski, A. *et al.*, *Heterocycles*, 2001, **54**, 581-584 (synth)

Okue, M. *et al.*, *Tetrahedron*, 2001, **57**, 4107-4110 (synth)

Okue, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 1093-1096 (synth, bibl)

Davis, F.A. *et al.*, *Tetrahedron*, 2004, **60**, 5111-5115 (synth)


Gogoi, N. *et al.*, *Eur. J. Org. Chem.*, 2006, 1722-1725 (synth)

Bertrand, M.B. *et al.*, *Org. Lett.*, 2006, **8**, 2353-2356 (synth)

Caldwell, J.J. *et al.*, *ARKIVOC*, 2007, xii, 67-90 (synth)

2-Benzyl-3-hydroxypyrazine **B-90**

3-(Phenylmethyl)-2(1H)-pyrazinone, 9CI. 3-Benzyl-2(1H)-pyrazinone. 3-Benzylpyrazinol



C₁₁H₁₀N₂O 186.213

OH-form


Me ether: 2-Benzyl-3-methoxypyrazine [57674-19-8]

C₁₂H₁₂N₂O 200.24

Prod. by *Chondromyces crocatus*.

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (isol)

3-Benzylidene-6-(hydroxymethyl)-1,4-dimethyl-6-(methylthio)-2,5-piperazinedione **B-91**



C₁₅H₁₈N₂O₃S 306.385


(R,Z)-form [1028330-17-7]

Prod. by a marine-derived fungus of the Pleosporales (ascomycetes). Powder. [α]_D²⁵ +416 (c, 0.47 in CHCl₃).

Prachyawarakorn, V. *et al.*, *Planta Med.*, 2008, **74**, 69-72 (isol, pmr, cmr)

6-Benzyl-4-oxo-4H-pyran-3-carboxylic acid **B-92**

4-Oxo-6-(phenylmethyl)-4H-pyran-3-carboxylic acid



C₁₃H₁₀O₄ 230.22

Amide: 6-Benzyl-4-oxo-4H-pyran-3-carboxamide. **Carbonarone A** [937182-85-9]


C₁₃H₁₁NO₃ 229.235

Prod. by the marine-derived fungus *Aspergillus carbonarius* WZ-4-11 and from *Pestalotiopsis theae*. Cytotoxic. Amorph. yellow powder. λ_{max} 279 (log ε 4.2); 317 (log ε 3.66); 401 (log ε 3.6) (MeOH).

Zhang, Y. et al., *J. Antibiot.*, 2007, **60**, 153-157 (isol, pmr, cmr)

2-Benzyl-4(3H)-quinazolinone**B-93**

2-(Phenylmethyl)-4(3H)-quinazolinone, 9CI. **Glycosminine**. Glycophymine. 2-Benzyl-4-hydroxyquinazolinone. 2-(Phenylmethyl)-4-quinazolinol [4765-56-4]

C₁₅H₁₂N₂O 236.273

Minor alkaloid from the leaves of *Glycosmis arborea*. Also isol. from the flowers of *Glycosmis pentaphylla* (arbor-ea). Cryst. (MeOH, EtOH, CHCl₃ or C₆H₆). Mp 254-256° (248-249°).

OH-form

Me ether: 4-Methoxy-2-(phenylmethyl)-quinazolinone. 2-Benzyl-4-methoxyquinazolinone. **Glycophymoline** [72361-61-6]

C₁₆H₁₄N₂O 250.299

Minor alkaloid from the flower heads of *Glycosmis pentaphylla* (Rutaceae). Cryst. (C₆H₆/MeOH). Mp 165°.

1H-form

N-Me: 1-Methyl-2-(phenylmethyl)-4(1H)-quinazolinone. 2-Benzyl-1-methyl-4(1H)-quinazolinone. **Arborine**. **Glycosine** [6873-15-0]

C₁₆H₁₄N₂O 250.299

Major alkaloid from leaves of *Glycosmis arborea*. Also present in *Ruta graveolens* (rue) and in the fruits of *Zanthoxylum budrunga* (Rutaceae). Acetylcholine inhibitor, central hypotensive agent. *G. arborea* is a folk medicine against fever, anaemia and jaundice. Cryst. (MeOH/CHCl₃). Mp 161-162° (155-156°).

N-Me, hydrochloride:

Cryst. + 2H₂O (H₂O). Mp 215° dec. (partial melting at 106-108°).

N-Me, picrate:

Yellow needles (EtOH). Mp 172-173°.

Chakravarti, D. et al., *Tetrahedron*, 1961, **16**,224-250 (*Arborine*, uv, ir, pmr, struct)Pakrashi, S.C. et al., *Tetrahedron*, 1963, **19**,


1011 (uv, ir, pmr, ms, struct, synth)

Pakrashi, S.C. et al., *Indian J. Chem.*, 1968, **6**,

472 (synth)

O'Donovan, D.G. et al., *J.C.S.(C)*, 1970,2466-2470 (*biosynth*)Patel, V.S. et al., *J. Indian Chem. Soc.*, 1972,**49**, 59 (uv)Kametani, T. et al., *J.A.C.S.*, 1977, **99**, 2306-2309 (*synth*, ir, pmr)Rhee, R.P. et al., *J.O.C.*, 1977, **42**, 3650 (*synth*, ir, pmr)Sarkar, M. et al., *Phytochemistry*, 1977, **16**, 2007 (*isol*, uv, ir, ms, synth)Kametani, T. et al., *Heterocycles*, 1978, **9**, 1585 (*synth*)Bhattacharyya, J. et al., *Heterocycles*, 1979, **12**, 929; 1980, **14**, 1469 (uv, ir, pmr, cmr)Naik, N.R. et al., *J. Indian Chem. Soc.*, 1979, **56**, 708-711 (*synth*, uv, ir)Sarkar, M. et al., *Phytochemistry*, 1979, **18**, 694 (*Glycophymoline*)Yamato, M. et al., *Chem. Pharm. Bull.*, 1981, **29**, 3124 (*synth*)Chakraborty, D.P. et al., *Synthesis*, 1981, 977-979 (*synth*)Teshima, T. et al., *J. Biol. Chem.*, 1982, **257**, 5085 (*pharmacol*)Ganjan, I. et al., *Synth. Commun.*, 1984, **14**, 33 (*synth*, pmr)Johns, S. et al., *Alkaloids (Academic Press)*, 1986, **29**, 129-140 (*Arborine*, rev. *pharmacol*)Uozumi, Y. et al., *J.A.C.S.*, 1989, **111**, 3725 (*synth*, ir, pmr, ms)Muthukrishnan, J. et al., *Phytochemistry*, 1999, **50**, 249-254 (*Arborine*, *isol*, pmr, cmr)**2-Benzyl-3,6-tropanediol****B-94**

8-Methyl-2-(phenylmethyl)-8-azabicyclo[3.2.1]octane-3,6-diol, 9CI. 2-Benzyl-3,6-dihydroxytropane

C₁₅H₂₁NO₂ 247.336

(2α,3α,6β)-form Abs config. not known - can only be deduced.

O⁶-Benzoyl: **Alkaloid C†**


[50656-88-7]

C₂₂H₂₅NO₃ 351.444

Alkaloid from the leaves of *Knightsia deplanchei* (Proteaceae). Cryst. (C₆H₆). Mp 168-170°.

Kan-Fan, C. et al., *Acta Chem. Scand.*, 1973, **27**, 1039-1052 (*isol*, ms, uv, pmr)Lounasmaa, M. et al., *J.O.C.*, 1975, **40**, 3694-3697 (*cmr*, *struct*, *config*)Lounasmaa, M. et al., *Planta Med.*, 1978, **34**, 68-72 (*pmr*, *struct*)**4-Benzyl-3,6-tropanediol****B-95**

8-Methyl-4-(phenylmethyl)-8-azabicyclo[3.2.1]octane-3,6-diol, 9CI. 2-Benzyl-3,7-dihydroxytropane

C₁₅H₂₁NO₂ 247.336

(3α,4α,6β)-form

3-Ac: **Knightsoline**

[77053-08-8]

C₁₇H₂₃NO₃ 289.374

Alkaloid from the leaves of *Knightsia strobilina* (Proteaceae). Cryst.

(Me₂CO). Mp 191-192°. [α]_D²⁰ +10(CHCl₃). Abs. config. not known.3-(E-Cinnamoyl): **Knightsia Alkaloid D**


[56761-54-7]

C₂₄H₂₇NO₃ 377.482

Alkaloid from the leaves of *Knightsia deplanchei* (Proteaceae). Cryst. (C₆H₆/hexane). Mp 174-175°. Racemate.

Kan-Fan, C. et al., *Acta Chem. Scand.*, 1973, **27**, 1039-1052 (*isol*, ms, pmr, uv)Lounasmaa, M. et al., *J.O.C.*, 1975, **40**, 3694-3697 (*cmr*, *struct*, *config*)Lounasmaa, M. et al., *Planta Med.*, 1978, **34**, 66-72 (*pmr*, *struct*)Lounasmaa, M. et al., *Phytochemistry*, 1980, **19**, 949-955 (*isol*, ir, pmr, ms, *struct*)**2-Benzyl-3-tropanol****B-96**

8-Methyl-2-(phenylmethyl)-8-azabicyclo[3.2.1]octan-3-ol, 9CI. 2-Benzyl-8-methyl-8-azabicyclo[3.2.1]octan-3-ol. 2-Benzyl-3-hydroxytropane

C₁₅H₂₁NO 231.337(1*RS*,2*SR*,3*RS*)-form

(±)-(2α,3α)-form

Cryst. (Me₂CO). Mp 123-124°.Ac: **Alkaloid KDI**. 2-Benzyl-O-acetyltropanol. **Knightsia Alkaloid B**

[50656-87-6]

C₁₇H₂₃NO₂ 273.374

Alkaloid from the leaves of *Knightsia deplanchei* (Proteaceae). Viscous oil.

[α]_D 0.Benzoyl: **Knightsia Alkaloid A**. 2-Benzyl-3-O-benzoyltropanol

[50656-86-5]

C₂₂H₂₅NO₂ 335.445

Alkaloid from the leaves of *Knightsia deplanchei* (Proteaceae). Viscous oil.

[α]_D 0.

1'-Hydroxy: see 2-(α-Hydroxybenzyl)-3-tropanol, H-429

Kan-Fan, C. et al., *Acta Chem. Scand.*, 1973, **27**, 1039-1052 (*isol*, ms, pmr, uv)Lounasmaa, M. et al., *Tet. Lett.*, 1974, 2509-2512 (*synth*)Lounasmaa, M. et al., *J.O.C.*, 1975, **40**, 3694-3697 (*cmr*, *config*)Lounasmaa, M. et al., *Planta Med.*, 1978, **34**, 66-72 (*pmr*, *struct*)Majewski, M. et al., *J.O.C.*, 1995, **60**, 5825-5830 (*synth*, O²-Ac, pmr, cmr)**Benzylurea, 9CI****B-97****Benzylcarbamide**

[538-32-9]

PhCH₂NHCONH₂C₈H₁₀N₂O 150.18


Constit. of the stems of *Salvadora persica*. Mod. sol. hot H₂O. Mp 147-148°.

▶ LD₅₀ (rat, orl) 2700 mg/kg. YS1780000Boivin, J.L. et al., *Can. J. Chem.*, 1951, **29**, 478 (*synth*)Argabright, P.A. et al., *J.O.C.*, 1967, **32**, 3261 (*synth*)Patonay, T. et al., *Synth. Commun.*, 1996, **26**, 4253 (*synth*)Khalil, A.T. et al., *Arch. Pharmacol. Res.*, 2006, **29**, 952-956 (*isol*, pmr, cmr)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BFN125

Berberamunine**B-98**

[485-18-7]

C₃₆H₄₀N₂O₆ 596.722

Alkaloids covered by this entry are enantiomeric with those covered by Grisabine, G-177 (1*S*,1'*R*)-config. and diastereomeric with those covered by Dauricine, D-88 (*R,R*-config.) and Thallibrine, T-309 (*S,S*-config.). Alkaloid from *Berberis amurensis*, *Berberis integrifolia* and *Berberis oblonga*. Cryst. (Et₂O/Me₂CO). Mp 190-191°. [α]_D +87.

N²-De-Me: 2-NorberbamunineC₃₅H₃₈N₂O₆ 582.695

Alkaloid from callus cultures of *Berberis stolonifera*, *Berberis beaniana*, *Berberis giraldii*, *Berberis julianae*, *Berberis papillifera*, *Berberis taliensis*, *Berberis turcomanica* and *Berberis wilsonae* (Berberidaceae).

O⁷-Me: Temuconine

[18210-69-0]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from *Berberis valdiviana* (Berberidaceae). Shows selective anti-plasmodial activity. [α]_D²⁵ +68 (c, 0.24 in MeOH). Struct. of this alkaloid recently revised from that publ. (Shamma, M. *et al.*, unpubl. results, personal communication).

O¹²-Me: Thaligrisine

[93780-78-0]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from the roots and rhizomes of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Needles (MeOH). [α]_D²⁵ +57 (c, 0.13 in MeOH).

O⁶-De-Me, O⁷-Me: Espinine

[26137-40-6]

C₃₆H₄₀N₂O₆ 596.722

Alkaloid from the root bark of *Berberis laurina* (Berberidaceae). Mp 123-125°. [α]_D +25 (CHCl₃).

O⁶-De-Me, O⁷, O¹²-di-Me: Espinidine

[26137-41-7]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from root bark of *Berberis laurina* (Berberidaceae). Amorph. [α]_D +31 (CHCl₃).

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1955, **75**, 753; 1957, **77**, 1075; 1079 (*isol, struct*)

Kametani, T. *et al.*, *Yakugaku Zasshi*, 1968, **88**, 1163 (*synth*)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1969, **17**, 2120 (*cd, ord, pmr*)

Falco, M.R. *et al.*, *Experientia*, 1969, **25**, 1236 (*Espinine*)

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 80; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 68 (*isol*)

Guinaudeau, H. *et al.*, *Heterocycles*, 1982, **19**, 1009 (*Temuconine*)


Guinaudeau, H. *et al.*, *Tetrahedron*, 1984, **40**, 1975 (*Thaligrisine*)

Cassels, B.K. *et al.*, *Phytochemistry*, 1987, **26**, 1005 (*2-Norberbamunine*)

Angerhoffer, C.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 59-66 (*Espinine, activity*)

Berberal†**B-99**

6-(1,3-Dihydro-6,7-dimethoxy-3-oxo-1-isobenzofuran-1-yl)-7,8-dihydro-1,3-dioxolo[4,5-g]isoquinolin-5(6H)-one, 9CI [66408-44-4] [64939-64-6]

C₂₀H₁₇NO₇ 383.357

Not to be confused with Berberal in B-101. Alkaloid from leaves and young shoots of *Berberis heterobotrys*. Cryst. (Et₂O). Mp 151-153°. No evidence for abs.config. of nat. alkaloid but prob. racemic. λ_{max} 226 (log ε 4.92); 255 (log ε 4.6); 305 (log ε 4.41) (EtOH).


Perkin, W.H. *et al.*, *J.C.S.*, 1890, **57**, 1079-1080 (*synth*)

Kondo, Y. *et al.*, *J.C.S. Perkin I*, 1980, 911-918 (*synth, pmr*)

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, 869-873; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 774-777 (*isol, pmr, cmr, ms*)

Berberilycine**B-100**

[186358-50-9]


C₂₀H₁₇NO₄ 335.359

Alkaloid from the roots of *Berberis lycium* (Berberidaceae). Cryst. (MeOH). Mp 180-181°. λ_{max} 235 (log ε 7.5); 253 (log ε 6.5); 344 (log ε 6.2) (MeOH).

Ali, M. *et al.*, *Indian J. Heterocycl. Chem.*, 1996, **6**, 127-130 (*isol, uv, ir, pmr, ms*)

Berberine**B-101**

5,6-Dihydro-9,10-dimethoxybenzo[g]-1,3-benzodioxolo[5,6-a]quinolinizinium(1+), 9CI. Umbellatine†. Berbericine. Natural yellow 18. C.I. 75160. Majorine [2086-83-1]

C₂₀H₁₈NO₄⁺ 336.366

Quaternary alkaloid from many *Berberis* and *Mahonia* spp. (Berberidaceae) and very many other spp. in several different families. Fluorescent stain for heparin. Pharmacol. extensively studied. Antiarhythmic agent. α₂-Adrenoceptor agonist. Cholinesterase, tyrosine decarboxylase and tryptophanase inhibitor. Shows a wide variety of pharmacol. effects including respiratory stimulation, transient hypotension. Antianaemic, anticonvulsant, antiinflammatory agent. Shows antibacterial and antifungal props. and some cytotoxic and antineoplastic activity. Possesses immunostimulant, sedative and antitrypanosomal activity. Of clinical use in treatment of gastrointestinal disorders, has been used for cholera and infantile diarrhoea. Log P -0.4 (uncertain value) (calc).

▶ Adverse human effects including reduced body temperature. Can cause death by central paralysis. LD₅₀ (mus, orl) 329 mg/kg. DR9870000

Chloride: Dirin. Kyoberin. Phelloberin. T-Up. Wakamoto. Pool Eyes [633-65-8] λ_{max} 230 (log ε 4.42); 266 (log ε 4.39); 352 (log ε 4.35); 432 (log ε 3.7) (96% EtOH).

▶ DR9866400

Iodide: [4263-84-7]

Mp 250° dec.

Hydroxide: [117-74-8] Traditional dye-stuff. Mp 145° (CHCl₃ solvate). Forms hydrate, Mp variable, dec. ca. 160°.

▶ DR9866500

Sulfate: Berisol. Berberal†. Stogerin-Amp. Stopnin. NSC 5355

[633-66-9]

[6190-33-6]

Mp 274°. Not to be confused with Berberal, B-99.

▶ DR9867000

O⁹-De-Me: Berberrubine. Chileninone

[17388-19-1]

C₁₉H₁₆NO₄⁺ 322.34

Alkaloid from the twigs of *Berberis actinacantha* and from the stems of *Berberis darwinii* and *Berberis valdiviana*. Also from *Berberis vulgaris* (barberry), *Thalictrum polygamum* and *Fibraurea chloroleuca* (Berberidaceae, Ranunculaceae, Menispermaceae). Active against gram-positive bacteria. Deep red amorph. solid (as chloride).

O¹⁰-De-Me: Thalifendine

[18207-71-1]

C₁₉H₁₆NO₄⁺ 322.34

Quaternary alkaloid from *Thalictrum fendleri* (Ranunculaceae). Shows anti-plasmodial and antiamebic activities. Yellow cryst. (as chloride). Mp 230° (sinters) (chloride).

7,8-Dihydro, N-Me: see Lambertine, L-23

[79236-58-1]

Späth, E. *et al.*, *Monatsh. Chem.*, 1929, **52**, 117-128 (*Berberubine, isol*)

Shamma, M. *et al.*, *Tet. Lett.*, 1965, 3595-3598 (*Thalifendine*)


Kametani, T. *et al.*, *J.C.S.(C)*, 1969, 2036-2038 (*synth, ir*)

Preiminger, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 124-135 (*isol, uv*)

- Jewers, K. *et al.*, *J.C.S. Perkin 2*, 1972, 1393-1396 (*pmr*)
- Gharbo, S.A. *et al.*, *J. Nat. Prod.*, 1973, **36**, 349-351 (*Berberrubine*, activity)
- Holland, H.L. *et al.*, *Can. J. Chem.*, 1979, **57**, 1588-1597 (*biosynth*)
- Dimlich, R.V.W. *et al.*, *Stain Technol.*, 1980, **55**, 217-223 (*use*)
- Siwon, J. *et al.*, *Planta Med.*, 1981, **41**, 65-68 (*Berberrubine*, occur)
- Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957-3962 (*Berberrubine*, isol, uv, ir, pmr, ms)
- Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 191-193 (*antitumour props*)
- Shamma, M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 398-405 (*Berberrubine*, struct)
- Schiff, P.L. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1987, **5**, 363 (*rev. pharmacol*)
- Blaskó, G. *et al.*, *Heterocycles*, 1988, **27**, 911-916 (*cmr*)
- Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1989, **28**, 2833-2839 (*pmr, cmr*)
- Hui, K.K. *et al.*, *Life Sci.*, 1991, **49**, 315-324 (*pharmacol*)
- Ishii, H. *et al.*, *Heterocycles*, 1994, **37**, 897-902 (*synth*)
- Wang, Y.-X. *et al.*, *J. Cardiovasc. Pharmacol.*, 1994, **23**, 716-722 (*pharmacol*)
- Chen, C.-M. *et al.*, *J. Chromatogr. B*, 1995, **665**, 117-123 (*hplc*)
- Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)
- Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1552
- Wright, C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1638-1640 (*Thalifendine*, activity)
- Yadav, R.C. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 165-174 (*props*)
- Doggrell, S.A. *et al.*, *Expert Opin. Invest. Drugs*, 2005, **14**, 683-685 (*rev*)
- Tripathi, A.N. *et al.*, *Magn. Reson. Chem.*, 2007, **45**, 647-655 (*pmr, cmr, soln struct*)
- Grycová, L. *et al.*, *Phytochemistry*, 2007, **68**, 150-175 (*rev*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BFN500

Berbidine B-102

3,4-Dihydro-6-methoxy-2-methyl-7-[(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-8-isoquinolinyl)oxy]-1-(2H)-isoquinolinone, 9CI
[35611-57-5]




C₂₃H₂₈N₂O₅ 412.485
Alkaloid from aerial parts of *Berberis brandisiana* (Berberidaceae). Amorph.

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 317 (*isol, uv, ir, pmr, ms, struct*)

Berbitine B-103

6-(1,3-Dioxolo[4,5-g]isoquinolin-5-yl-methyl)-2,3-dimethoxyphenol, 9CI. 1-(2-Hydroxy-3,4-dimethoxybenzyl)-6,7-methylenedioxyisoquinoline
[105798-91-2]




C₁₉H₁₇NO₅ 339.347
Alkaloid from the roots of *Berberis actinacantha* (Berberidaceae). Needles (EtOH). Mp 159-160°.

α -Oxo: see Rugosinone, R-156

- Rahimzadeh, M. *et al.*, *Planta Med.*, 1986, **339** (*isol, uv, pmr, ms, struct*)
- Chantimakorn, V. *et al.*, *Aust. J. Chem.*, 1989, **42**, 209 (*synth, uv, ir, pmr, ms*)

Berninamycin A, 9CI B-104

U 27810. Antibiotic U 27810
[58798-97-3]



C₅₁H₅₁N₁₅O₁₅S 1146.121

Peptide antibiotic. Isol. from *Streptomyces bernensis*. Component of the Berninamycin complex which shows activity against gram-positive bacteria and is a potent inhibitor of bacterial protein synth. Sol. MeOH, butanol, DMF; poorly sol. H₂O, Me₂CO, hexane. Prob. identical with Theiomycetin. λ_{\max} 208 (ϵ 71500); 236 (ϵ 73800); 255 (sh) (ϵ 71500) (MeOH) (Derep).

Dihydro: Mp 268-272° dec.

Hexahydro: Mp 275-280°.

1''-Deoxy: **Berninamycin B**

[58798-98-4]

C₅₁H₅₁N₁₅O₁₄S 1130.122

Isol. from *Streptomyces bernensis*. Powder. λ_{\max} 208 (ϵ 71500); 236 (ϵ 73800); 255 (sh) (ϵ 71500) (MeOH) (Derep).

14-Demethyl: **Geninthiocin**

[158792-27-9]

C₅₀H₄₉N₁₅O₁₅S 1132.095

Prod. by *Streptomyces* sp. DD84. tipA Promotor inducing agent. Amorph. powder. Mp 270-275° dec. [α]_D¹⁹ +174.3 (c, 0.1 in CHCl₃/MeOH). λ_{\max} 215 (sh) (ϵ 76000); 240 (ϵ 85000); 315 (sh) (ϵ 13000) (EtOH) (Derep). λ_{\max} 215 (ϵ 76000); 240 (ϵ 85000); 315 (ϵ 13000) (EtOH) (Berdy).

Liesch, J.M. *et al.*, *CA*, 1976, **84**, 146412s (*struct*)

Liesch, J.M. *et al.*, *J.A.C.S.*, 1976, **98**, 299; 1977, **99**, 1645 (*isol, struct*)

Pearce, C.J. *et al.*, *J.A.C.S.*, 1979, **101**, 5069 (*biosynth*)

Abe, H. *et al.*, *Tet. Lett.*, 1988, **29**, 1401 (*struct*)


Yun, B.-S. *et al.*, *J. Antibiot.*, 1994, **47**, 969 (*Geninthiocin*)

Lau, R.C.M. *et al.*, *J. Antibiot.*, 1994, **47**, 1466 (*cmr, struct*)

Lau, R.C.M. *et al.*, *J.A.C.S.*, 1995, **117**, 7606 (*biosynth*)

Bernumicine B-105

[169626-38-4]



R¹ = R² = Me

C₂₀H₂₅NO₄ 343.422

Alkaloid from leaves of *Berberis nummularia*. Cryst. (as hydrochloride). Mp 211-212° (hydrochloride). [α]_D +14 (c, 0.02 in CHCl₃).

3'-O-De-Me: **Nummularine**

[198704-40-4]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Berberis nummularia*. Cryst. Mp 134-135°. [α]_D +21 (c, 0.03 in MeOH).

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 397; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 335

Faskhutdinov, M.F. *et al.*, *Khim. Prir. Soedin.*, 1997, **33**, 91-93; *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 70-71 (*Nummularine*)

Bernumine B-106

[169134-46-7]

As Bernumicine, B-105 with

R¹R² = -CH₂-

C₁₉H₂₁NO₄ 327.379

Alkaloid from leaves of *Berberis nummularia*. Oil. [α]_D +33 (c, 0.04 in CHCl₃).

Me ether: **Bernumidine**

[169626-37-3]


C₂₀H₂₃NO₄ 341.406

From leaves of *Berberis nummularia*. Oil. Mp 180-181° (as hydrochloride). [α]_D +21 (c, 0.01 in CHCl₃).

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 394; 397; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 331; 335

Berpodine B-107

[169238-48-6]



C₄₀H₃₈N₂O₁₃ 754.746

Alkaloid from young shoots of *Berberis heteropoda*. Cryst. (MeOH). Mp 197-198°.

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 264; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 219

Bervulcine**B-108**C₁₈H₁₉NO₃ 297.353


Struct. unknown. Alkaloid from *Berberis vulgaris* (barberry) (Berberidaceae). Mp 125-126° dec. [α]_D²⁴ -185 (c, 0.2 in CHCl₃).

Hydriodide:Prisms (H₂O). Mp 236° dec.**Methiodide:**Cryst. (H₂O). Mp 269° dec.**Picrate:**

Fine needles (MeOH). Mp 188-190° dec.

Döpke, W. et al., *Naturwissenschaften*, 1963, **50**, 595**Betalamic acid****B-109**

1,2,3,4-Tetrahydro-4-(oxoethylidene)-2,6-pyridinedicarboxylic acid
[18766-66-0]




Absolute Configuration

C₉H₉NO₅ 211.174

Precursor of betalains pigments in plants of the Centrospermae. Detected in *Beta vulgaris* (beetroot), *Celosia cristata* and *Portulaca grandiflora*. Yellow soln. in H₂O at pH 9, or amorph. green residue. Sensitive to air, acids and strong alkali. Various protected derivs. known. λ_{max} 430 (no solvent reported).

Kimler, L. et al., *Chem. Comm.*, 1971, 1329 (isol, synth, struct)Döpp, H. et al., *Chem. Ber.*, 1973, **106**, 3473 (synth, uv)Büchi, G. et al., *J.O.C.*, 1977, **42**, 2192 (synth)
Hilpert, H. et al., *Helv. Chim. Acta*, 1984, **67**, 1547 (synth)Fischer, N. et al., *Helv. Chim. Acta*, 2004, **55**, 649-658 (biosynth)**Betanidin****B-110**

2-Carboxy-1-[(2,6-dicarboxy-2,3-dihydro-4(1H)-pyridinylidene)ethylidene]-2,3-dihydro-5,6-dihydroxy-1H-indolium hydroxide inner salt
[2181-76-2]



Absolute Configuration

C₁₈H₁₆N₂O₈ 388.333

Aglycone from Betanin. Cryst. (as hydrochloride). Forms cryst. K and NH₄ salts. Shows rapid E/Z isom. in soln. λ_{max} 271 (ε 8530); 295 (sh) (ε 7170); 544 (ε 51000) (no solvent reported) (hydrochloride).

5-O-β-D-Glucopyranoside: **Betanin**. *Phytolaccanin*. C.I. Natural Red 33. E 162

[7659-95-2]

C₂₄H₂₆N₂O₁₃ 550.475

Red pigment from beetroot, *Beta vulgaris* var. *rubra* and other Centrospermae. Food dye. Chemotaxonomic marker for the Centrospermae.

► US7968100

5-O-(3-O-Sulfo-β-D-glucopyranoside):

Rivianin

[58115-21-2]

C₂₄H₂₆N₂O₁₆S 630.539Pigment from *Rivina humilis* (Phytolaccaceae).

5-O-(6-O-Sulfo-β-D-glucopyranoside):

Prebetanin

[13798-16-8]

C₂₄H₂₆N₂O₁₆S 630.539Pigment from beetroot *Beta vulgaris* (Chenopodiaceae).

5-O-(4-O-Malonyl-β-D-glucopyranoside):

Phylloactin IIC₂₇H₂₈N₂O₁₆ 636.522Constit. of the fruit peel of *Hylocereus ocamponis*.

5-O-(6-O-Malonyl-β-D-glucopyranoside):

Phylloactin

[15167-85-8]

C₂₇H₂₈N₂O₁₆ 636.522Constit. of *Phyllocactus hybridus* (preferred genus name *Epiphyllum*) and *Hylocereus polyrhizus*.5-O-[3-Hydroxy-3-methylglutaryl-(→6)-β-D-glucopyranoside]: **Hylocerem**C₃₀H₃₄N₂O₁₇ 694.602Pigment from cactus *Hylocereus polyrhizus*.

5-O-[4-Hydroxy-3-methoxy-E-cinnamoyl-(→6)-β-D-glucopyranoside]:

Lampranthin II. Monoferuloylbetanin

[32074-65-0]

C₃₄H₃₄N₂O₁₆ 726.646Isol. from petals of *Lampranthus peerisii* and *Lampranthus sociorum*.

5-O-[β-D-Apiofuranosyl-(1→2)-β-D-glucopyranoside]:

C₂₉H₃₄N₂O₁₇ 682.591Constit. of the fruit peel of *Hylocereus ocamponis*.

5-O-[β-D-Apiofuranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]:

[290313-63-2]

C₃₂H₃₆N₂O₂₀ 768.638Pigment from *Schlumbergera x buckleyi*.

5-O-[4-Hydroxy-3-methoxy-E-cinnamoyl-(→5)-β-D-apiofuranosyl-(1→2)-β-D-glucopyranoside]: [290313-64-3]

[178885-00-2]

C₃₉H₄₂N₂O₂₀ 858.762Pigment from *Phytolacca americana* (pokeberry) and *Schlumbergera x buckleyi*.

5-O-[4-Hydroxy-3-methoxy-E-cinnamoyl-(→5)-β-D-apiofuranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]:

[290313-68-7]

C₄₂H₄₄N₂O₂₃ 944.809Pigment from *Schlumbergera x buckleyi*.

5-O-[4-Hydroxy-3,5-dimethoxy-E-cinnamoyl-β-D-apiofuranosyl-(1→2)-β-D-glucopyranoside]:

C₄₀H₄₄N₂O₂₁ 888.788Constit. of the fruit peel of *Hylocereus ocamponis*.5-O-[β-D-Glucopyranosyl-(1→2)-β-D-glucopyranoside]: **Bougainvillein r-I.***Betanidin 5-sophoroside*

[30513-63-4]

C₃₀H₃₆N₂O₁₈ 712.617Pigment from a horticultural variety of *Bougainvillea* "Mrs. Butt" and from *Hylocereus ocamponis*. λ_{max} 538 (H₂O).

5-O-[β-D-Glucopyranosyl-(1→2)-4-O-malonyl-β-D-glucopyranoside]:

[951659-92-0]

C₃₃H₃₈N₂O₂₁ 798.664Constit. of the fruit of *Mammillaria* spp. Tentative struct. assigned.5-O-[β-D-Glucopyranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]: **Mammillarinin**

[951676-55-4]

C₃₃H₃₈N₂O₂₁ 798.664Constit. of the fruit of *Mammillaria* spp.5-O-[β-D-Glucuronopyranosyl-(1→2)-β-D-glucopyranoside]: **Amaranthin.***Amarantin*

[15167-84-7]

[11033-33-3]

C₃₀H₃₄N₂O₁₉ 726.601Pigment from *Amaranthus caudatus* (love-lies-bleeding), *Amaranthus tricolor* (Chinese spinach), *Celosia plumosa*, *Salicornia europaea* and other spp. (Amaranthaceae, Chenopodiaceae).5-O-[4-Hydroxy-E-cinnamoyl-(→2)-β-D-glucuronopyranosyl-(1→2)-β-D-glucopyranoside]: **Celosianin I**

[114818-87-0]

C₃₉H₄₀N₂O₂₁ 872.746Pigment from *Celosia cristata* (Amaranthaceae).5-O-[4-Hydroxy-3-methoxy-E-cinnamoyl-(→2)-β-D-glucuronopyranosyl-(1→2)-β-D-glucopyranoside]: **Celosianin II. Monoferuloylamarantin**

[114847-18-6]

C₄₀H₄₂N₂O₂₂ 902.772Isol. from cell suspension cultures of *Chenopodium rubrum* and from *Celosia cristata*.5-O-[β-D-Glucuronopyranosyl-(1→2)-[3-hydroxy-3-methylglutaryl-(→6)-β-D-glucopyranoside]: **Iresinin I**

[78413-55-5]

C₃₆H₄₂N₂O₂₃ 870.727Pigment from *Iresine herbstii* (Amaranthaceae). Violet cryst. [α]_D²⁰ +160.6-O-β-D-Glucopyranoside: **Gomphrenin I**

[17008-59-2]

C₂₄H₂₆N₂O₁₃ 550.475Violet pigment from the inflorescence of *Gomphrena globosa* (Amaranthaceae). Also isol. from bracts of *Bougainvillea glabra* (Nyctaginaceae).6-O-[4-Hydroxy-E-cinnamoyl-(→6)-β-D-glucopyranoside]: **Gomphrenin II**

[143022-02-0]

C₃₃H₃₂N₂O₁₅ 696.62Pigment from *Gomphrena globosa* and

- fruits of *Basella rubra* (Malabar spinach).
- 6-O-[4-Hydroxy-3-methoxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranoside]: **Gomphrenin III**. Gomphrenin V [11048-86-5] [16955-52-5] $C_{34}H_{34}N_2O_{16}$ 726.646
Pigment from *Gomphrena globosa* (Amaranthaceae). Gomphrenin V mentioned in a 1967 paper only. Current evidence is that it was identical with Gomphrenin III. Other Gomphrenins (IV, VI, VII and VIII) were not fully characterised. λ_{max} 290 ; 319 ; 549 (no solvent reported).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Bougainvillein-v**. Betanidin 6-sophoroside [32100-96-2] $C_{30}H_{36}N_2O_{18}$ 712.617
Pigment from bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[4-Hydroxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: [160094-28-0] $C_{39}H_{42}N_2O_{20}$ 858.762
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[4-hydroxy-E-cinnamoyl-($\rightarrow 6$)]- β -D-glucopyranoside]: [160094-29-1] $C_{39}H_{42}N_2O_{20}$ 858.762
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[4-Hydroxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranosyl-($\rightarrow 2$)-[4-hydroxycinnamoyl-($\rightarrow 6$)]- β -D-glucopyranoside]: [160094-33-7] $C_{48}H_{48}N_2O_{22}$ 1004.907
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[4-hydroxycinnamoyl-($\rightarrow 6$)]- β -D-glucopyranosyl-(1 \rightarrow 2)-3,4-dihydroxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranoside]: [160094-31-5] $C_{54}H_{58}N_2O_{28}$ 1183.049
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[4-hydroxycinnamoyl-($\rightarrow 6$)]- β -D-glucopyranosyl-(1 \rightarrow 2)-4-hydroxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranoside]: [160094-32-6] $C_{54}H_{58}N_2O_{27}$ 1167.049
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[3,4-dihydroxycinnamoyl-($\rightarrow 6$)]- β -D-glucopyranoside]: [160094-27-9] $C_{39}H_{42}N_2O_{21}$ 874.762
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- 6-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 $\rightarrow 6$)]- β -D-glucopyranoside]: Betanidin 6-(2^G-glucosyl-rutinoside). Betanidin rhamnosophoroside $C_{36}H_{46}N_2O_{22}$ 858.76
Pigment from bracts of *Bougainvillea glabra*.
- 6-O-[4-Hydroxycinnamoyl-($\rightarrow 6$)- β -D-glucopyranosyl-(1 $\rightarrow 2$)-[3,4-dihydroxycinnamoyl-($\rightarrow 6$)]- β -D-glucopyranosyl-(1 $\rightarrow 2$)- β -D-glucopyranoside]: [160094-30-4] $C_{60}H_{68}N_2O_{33}$ 1345.191
From bracts of *Bougainvillea glabra* (Nyctaginaceae).
- Glycoside(1): Bougainvillein r-II**
Pigment from bracts of horticultural variety of *Bougainvillea* "Mrs Butt" (Nyctaginaceae). Complete struct. not determined. Alkaline hydrolysis gave Bougainvillein r-I, Isobougainvillein r-I, 4-Hydroxycinnamic and 3,4-Dihydroxycinnamic acids, ratio of latter two, 2:1. λ_{max} 280 ; 320 ; 541 (no solvent reported).
- Glycoside(2): Bougainvillein r-III**
Pigment from bracts of horticultural variety of *Bougainvillea* "Mrs Butt" (Nyctaginaceae). Complete struct. not determined. Alkaline hydrolysis gave 4-Hydroxycinnamic acid and a mixt. of Bougainvillein r-I and Isobougainvillein r-I. λ_{max} 312 ; 541 (no solvent reported).
- Glycoside(3): Bougainvillein r-IV**
Pigment from bracts of horticultural variety of *Bougainvillea* "Mrs Butt" (Nyctaginaceae). Complete struct. not determined. Alkaline hydrolysis gave 4-Hydroxycinnamic acid and a mixt. of Bougainvillein r-I and Isobougainvillein r-I.
- Glycoside(4): Bougainvillein r-V**
Pigment from bracts of horticultural variety of *Bougainvillea* "Mrs Butt" (Nyctaginaceae). Complete struct. not determined. Alkaline hydrolysis gave 4-Hydroxycinnamic acid and a mixt. of Bougainvillein r-I and Isobougainvillein r-I.
- Glycoside(5): Iresinin III** [11025-56-2]
Minor pigment from *Iresine herbstii*. Complete struct. not determined. Alkaline hydrolysis gave Amarantin, Isoamarantin plus (E)-4-Hydroxycinnamic, (E)-3,4-Dihydroxycinnamic, (E)-4-Hydroxy-3-methoxycinnamic and (E)-4-Hydroxy-3,5-dimethoxycinnamic acids. λ_{max} 298 ; 537 (no solvent reported).
- Glycoside(6): Iresinin IV** [11025-57-3]
Minor pigment from *Iresine herbstii*. Complete struct. not determined. Alkaline hydrolysis gave Amarantin plus, Isoamarantin, 4-Hydroxy-3-methoxycinnamic and 4-Hydroxy-3,5-dimethoxycinnamic acids (ratio 1:1:1). λ_{max} 300 ; 321 ; 545 (no solvent reported).
- Glycoside(7): Isobougainvillein r-II**
Pigment from bracts of horticultural variety of *Bougainvillea* "Mrs Butt" (Nyctaginaceae). Complete struct. not determined. Alkaline hydrolysis gave Bougainvillein r-I, Isobougainvillein r-I, 4-Hydroxycinnamic and 3,4-Dihydroxycinnamic acids (ratio of latter two, 2:1). λ_{max} 280 ; 320 ; 541 (no solvent reported).
- 14,15-Didehydro, 5-O- β -D-glucopyranoside: Neobetainin** [71199-29-6] $C_{24}H_{24}N_2O_{13}$ 548.459
Minor pigment from roots of *Bougainvillea vulgaris* ssp. *vulgaris*. Also in flowers of Barbary fig (*Opuntia ficus-indica*), *Portulaca grandiflora*, *Zygocactus truncatus* (preferred genus name *Schlumbergera*) and *Phytolacca bogotensis* (Chenopodiaceae, Portulacaceae, Cactaceae, Phytolaccaceae). Prob. artifact. λ_{max} 267 ; 306 ; 470 (no solvent reported). λ_{max} 298 ; 365 (H₂O).
- 15-Epimer: Isobetainin** [4934-32-1] $C_{18}H_{16}N_2O_8$ 388.333
Minor congener of Betanidin. Epimeric at the piperidine COOH group.
- 15-Epimer, 5-O- β -D-glucopyranoside: Isobetainin** [15121-53-6] $C_{24}H_{26}N_2O_{13}$ 550.475
Minor congener of Betanin, e.g. from *Beta vulgaris* and *Amaranthus* spp. (Chenopodiaceae, Amaranthaceae).
- 15-Epimer, 5-O-(4-O-malonyl- β -D-glucopyranoside): Isohyllactin II** $C_{27}H_{28}N_2O_{16}$ 636.522
Constit. of the fruit peel of *Hylocereus ocamponis*.
- 15-Epimer, 5-O-(6-O-malonyl- β -D-glucopyranoside): Isohyllactin** [28378-93-0] $C_{27}H_{28}N_2O_{16}$ 636.522
Constit. of *Hylocereus polyrhizus* and *Phyllocactus hybridus*.
- 15-Epimer, 5-O-[3-hydroxy-3-methylglutaroyl-($\rightarrow 6$)- β -D-glucopyranoside]: Isohylocerenin** $C_{30}H_{34}N_2O_{17}$ 694.602
Constit. of the fruit of *Hylocereus ocamponis* and *Hylocereus polyrhizus*.
- 15-Epimer, 5-O-[β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]:** $C_{29}H_{34}N_2O_{17}$ 682.591
Constit. of the fruit peel of *Hylocereus ocamponis*.
- 15-Epimer, 5-O-[β -D-apiofuranosyl-(1 \rightarrow 2)-6-O-malonyl- β -D-glucopyranoside]:** [290313-74-5] $C_{32}H_{36}N_2O_{20}$ 768.638
Pigment from *Schlumbergera x buckleyi*.
- 15-Epimer, 5-O-[4-hydroxy-3-methoxy-E-cinnamoyl-($\rightarrow 3$)- β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]:** $C_{39}H_{42}N_2O_{20}$ 858.762
Pigment from *Phytolacca americana* (pokeberry).
- 15-Epimer, 5-O-[4-hydroxy-3,5-dimethoxy-E-cinnamoyl-($\rightarrow 5$)- β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]:** $C_{40}H_{44}N_2O_{21}$ 888.788
Constit. of the fruit peel of *Hylocereus ocamponis*.

15-Epimer, 5-O- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Isobougainvillein r-I**

[28146-90-9]

C₃₀H₃₆N₂O₁₈ 712.617

Pigment from *Bougainvillea* sp. (Nyctaginaceae). λ_{\max} 538 (H₂O).

15-Epimer, 5-O- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 2)-4-O-malonyl- β -D-glucopyranoside]: [951659-95-3]

C₃₃H₃₈N₂O₂₁ 798.664

Constit. of the fruit of *Mammillaria* spp. Tentative struct. assigned.

15-Epimer, 5-O- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 2)-6-O-malonyl- β -D-glucopyranoside]: **Isomammillaridin**

C₃₃H₃₈N₂O₂₁ 798.664

Constit. of the fruit of *Mammillaria* spp.

15-Epimer, 5-O- $[\beta$ -D-glucuronopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Isoamaranthin**

[15229-42-2]

C₃₀H₃₄N₂O₁₉ 726.601

Minor congener of Amaranthin.

15-Epimer, 5-O- $[\beta$ -D-glucuronopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Isocelosianin I**

[338948-27-9]

C₃₉H₄₀N₂O₂₁ 872.746

Pigment from *Celosia cristata* (Amaranthaceae).

15-Epimer, 5-O- $[\beta$ -D-glucuronopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Isocelosianin II**

[338948-57-5]

C₄₀H₄₂N₂O₂₂ 902.772

Pigment from *Celosia cristata* (Amaranthaceae).

15-Epimer, 5-O- $[\beta$ -D-glucuronopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Isoiresinin I**

[338947-95-8]

C₃₆H₄₂N₂O₂₃ 870.727

Pigment from *Iresine herbstii* (Amaranthaceae). Noncryst.

15-Epimer, 6-O- $[\beta$ -D-glucopyranoside]: **Iso-gomphrenin I**

[17008-60-5]

C₂₄H₂₆N₂O₁₃ 550.475

Violet pigment from *Gomphrena globosa* (Amaranthaceae).

15-Epimer, 6-O- $[\beta$ -D-glucopyranoside]: **Iso-gomphrenin II**

[143062-65-1]

C₃₃H₃₂N₂O₁₅ 696.62

Pigment from *Gomphrena globosa* and fruits of *Basella rubra* (Malabar spinach).

15-Epimer, 6-O- $[\beta$ -D-glucopyranoside]: **Iso-gomphrenin III**

[16955-53-6]

C₃₄H₃₄N₂O₁₆ 726.646

Pigment from *Gomphrena globosa* (Amaranthaceae) and fruits of *Basella pulva*.

15-Epimer, 6-O- $[\beta$ -D-glucopyranosyl-

(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside]: **Isobetanidin**

6-(2^G-glucosylrutinoside). **Isobetanidin**

rhamnosophoroside

C₃₆H₄₆N₂O₂₂ 858.76

Pigment from bracts of *Bougainvillea glabra*.

Dreiding, A.S. et al., *Helv. Chim. Acta*, 1965, **48**, 252-258; 1977, **60**, 673-683 (synth, Betanin)

Piattelli, M. et al., *Ann. Chim. (Rome)*, 1966, **56**, 1060; *CA*, **66**, 65805y (Amaranthin)

Minale, L. et al., *Phytochemistry*, 1966, **5**, 1037-1052 (Iresinins, Phyllocactin)

Wyler, H. et al., *Helv. Chim. Acta*, 1967, **50**, 545-560; 1984, **67**, 1793-1800 (Prebetanin)

Minale, L. et al., *Phytochemistry*, 1967, **6**, 703-709 (Gomphrenins)

Piattelli, M. et al., *Phytochemistry*, 1970, **9**, 455-458 (*Bougainvilleins*)

Imperato, F. et al., *Phytochemistry*, 1975, **14**, 2526-2527 (Riviniainin, glucosylrutinosides)

Hilpert, H. et al., *Helv. Chim. Acta*, 1984, **67**, 1547-1561 (synth)

Alard, D. et al., *Phytochemistry*, 1985, **24**, 2383-2385 (*Neobetainin*)

Strack, D. et al., *Phytochemistry*, 1988, **27**, 3529-3531 (*Lampranthin II*, *Celosianin II*)

Heuer, S. et al., *Phytochemistry*, 1992, **31**, 1801-1807; 1994, **37**, 3529-3531 (*sphorosides*, *Bougainvillea glabra* constits)

Schliemann, W. et al., *Phytochemistry*, 1996, **42**, 1039-1046 (*Phytolacca americana* constits)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 241 (*Betanin*, use)

Kobayshi, N. et al., *Phytochemistry*, 2000, **54**, 419-426 (*Schlumbergera betacyanins*)

Cai, Y. et al., *J. Agric. Food Chem.*, 2001, **49**, 1971 (bibl, *Gomphrenin II*, *Celosianin I*, *Isocelosianins*, *Isogomphrenin II*)

Wybraniec, S. et al., *Phytochemistry*, 2001, **58**, 1209-1212; 2007, **68**, 251-259 (*Hylocerenin*, *Isolylocerenin*, *Bougainvillein r-I*, *Hylocereus ocamponis* constits)

Stintzing, F.C. et al., *Phytochemistry*, 2004, **65**, 415-422 (pmr, cmr)


Wybraniec, S. et al., *J. Agric. Food Chem.*, 2007, **55**, 8138-8143 (*Mammillaria* constits)

Bharatamine

B-111

5,8,13,13a-Tetrahydro-3-methoxy-6H-dibenzo[a,g]quinolizin-2-ol, 9CI.

5,8,13,13a-Tetrahydro-2-hydroxy-3-methoxy-6H-dibenzo[a,g]quinolizine



C₁₈H₁₉NO₂ 281.354

The first ring D unoxygenated protoberberine alkaloid. Biogenetically related to Emetine, E-72; prob. of mixed tyrosine-terpenoid derivation.

(±)-form [85769-48-8]

Alkaloid from the seeds of *Alangium lamarckii* (Alangiaceae). Granules (CHCl₃/petrol). Mp 182-183°. λ_{\max} 224 (log ϵ 3.91); 285 (log ϵ 3.62) (MeOH).

λ_{\max} 224 (log ϵ 3.93); 286 (log ϵ 3.63) (MeOH/NaOH).

Pakrashi, S.C. et al., *Tet. Lett.*, 1983, **24**, 291-294 (isol, biosynth, uv, ir, pmr, ms, struct)

Patil, S.D. et al., *Indian J. Chem., Sect. B*, 1985, **24**, 360-362 (synth, uv, ir, pmr, ms)

Mandal, S.B. et al., *J.O.C.*, 1988, **53**, 4236-4241 (synth)


Takano, S. et al., *Heterocycles*, 1990, **31**, 1151-1156 (synth)

Grajewska, A. et al., *Tetrahedron: Asymmetry*, 2007, **18**, 2910-2914 (synth)

Bharhingine

B-112

[112448-55-2]



Absolute Configuration

C₁₉H₂₀N₂O 292.38


Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). $[\alpha]_{\text{D}}^{26}$ -50 (MeOH).

Atta-ur-Rahman, et al., *Planta Med.*, 1987, **53**, 256 (isol, uv, ir, pmr, ms, struct)

Bhimamycin C

B-113

5-Hydroxy-1-(1-hydroxyethyl)-2-(2-hydroxyethyl)-3-methyl-2H-benz[b]isoidole-4,9-dione, 9CI



C₁₇H₁₇NO₅ 315.325

(ξ)-form [701915-60-8]


Prod. by a terrestrial streptomycete. Yellow powder. $[\alpha]_{\text{D}}^{20}$ -23.4 (c, 0.55 in CHCl₃). λ_{\max} 380 (log ϵ 3.39) (MeOH).

Fotso, S. et al., *J. Antibiot.*, 2003, **56**, 931-941 (isol, ir, pmr, cmr, ms)

Bhimamycin D

B-114

[701915-61-9]




C₂₂H₁₅NO₆ 389.364

Prod. by a terrestrial streptomycete. Yellow powder. $[\alpha]_{\text{D}}^{20}$ -2.5 (c, 0.79 in CHCl₃). Bhimamycin F and Bhimamycin G are synthetic analogues. λ_{\max} 248 (log ϵ 4.43); 404 (log ϵ 3.92) (MeOH).

Fotso, S. et al., *J. Antibiot.*, 2003, **56**, 931-941 (isol, ir, pmr, cmr, ms)

10,10'-Bi[N-acetyl-11,12-dihydroxyaspidospermidine] **B-115**

[24778-13-0]


C₄₂H₅₄N₄O₆ 710.912

Alkaloid from the bark of *Aspidosperma melanocalyx* (Apocynaceae). Cryst. (hexane). Mp 274-275°. [α]_D²⁵ +108 (c, 1.01 in Py).

Tetra-Ac: Mp 135-140°.

Miranda, E.C. et al., *Experientia*, 1969, **25**, 575 (uv, ir, pmr, ms, struct)**Bianfugecine**

5,9-Dimethoxy-7H-dibenzo[de,h]quinolin-7-one, 9CI
[96681-50-4]

R¹ = R³ = H, R² = OMeC₁₈H₁₃NO₃ 291.306


Alkaloid from the rhizomes of *Menispermum dauricum* (Menispermaceae). Yellow plates (MeOH). Mp 199-201° (synthetic).

Hou, C. et al., *Yaoxue Xuebao*, 1985, **20**, 112; CA, **103**, 3691gKunitomo, J. et al., *Heterocycles*, 1986, **24**, 437 (uv, ir, pmr, ms, struct)**Bianfugedine**

2-Methoxy-12H-benzo[h][1,3]benzodioxolo[6,5,4-de]quinolin-12-one, 9CI, 5,6-Methylenedioxy-9-methoxy-7H-dibenzo[de,h]quinolin-7-one
[96681-51-5]

As Bianfugecine, B-116 with R¹ = H, R²R³ = -OCH₂O-C₁₈H₁₁NO₄ 305.289Alkaloid from the rhizomes of *Menispermum dauricum* (Menispermaceae).Hou, C. et al., *Yaoxue Xuebao*, 1985, **20**, 112; CA, **103**, 3691g**Bicucine**


5-[Hydroxy-(5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)methyl]-1,3-benzodioxole-4-carboxylic acid, 9CI
[481-60-7]

C₂₀H₁₉NO₇ 385.373

The hydroxy acid corresponding to Bicuculline, B-119 with which it equilibrates. Alkaloid from *Corydalis sempervirens* and *Dicentra cucullaria* (Papaveraceae). Stout prisms (conc. aq. NH₃). Mp 217° (215°).

Manske, R.H.F. et al., *Can. J. Res.*, 1933, **8**, 1 (isol)Robinson, R. et al., *Annu. Rev. Biochem.*, 1935, **4**, 497 (struct)**Bicuculline**

6-(5,6,7,8-Tetrahydro-6-methyl-1,3-benzodioxolo[4,5-g]isoquinolin-5-yl)furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI



(+)-form

C₂₀H₁₇NO₆ 367.357

Diastereoisomeric with Adlumidine, A-145. Alkaloid from many *Corydalis* spp. and several *Fumaria* spp. (opt. rotn. of isolates not recorded in some cases). Also isol. from *Adlumia fungosa* (*Adlumia cirrhosa*) (Papaveraceae). GABA_A-receptor antagonist. Contradictory data as to activity towards heart, intestinal and uterine muscle (relaxant and tonic actions claimed). Convulsant. Much used pharmacological tool. Log P 2.66 (uncertain value) (calc).

(+)-form [485-49-4]

Alkaloid from *Corydalis gigantea*, *Corydalis gortschakovii*, *Corydalis govianiana*, *Corydalis lutea*, *Corydalis marshalliana*, *Corydalis pseudo-adunca*, *Fumaria parviflora* (*F. indica*) and *Fumaria vaillantii* (Papaveraceae). Cryst. (EtOH or MeOH). Mp 194-195° (177°) Mp 215°. [α]_D +137.2 (CHCl₃). [α]_D²⁰ +132.7 (c, 0.049 in CHCl₃).

►DT4550000

Picrate: Mp 202°.

Methiodide: [40709-69-1]

Cryst. Mp 203-207°.

(-)-form [19730-80-4]

Alkaloid from *Corydalis severtzovii* and *Fumaria vaillantii* (Papaveraceae). Cryst. (EtOH or MeOH/Me₂CO). Mp 193-195°. [α]_D³³ -128 (c, 0.27 in CHCl₃) (-110).

Methiodide: [55950-07-7]

Mp 236° dec. [α]_D -107 (c, 1 in CHCl₃).**(±)-form** [56083-00-2]


Alkaloid from *Fumaria indica* and *Fumaria schleicheri* (Papaveraceae). Mp 216°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1310A (nmr)Edwards, O.E. et al., *Can. J. Chem.*, 1961, **39**, 1801 (isol, uv, ir, pmr, bibl)Safe, S. et al., *Can. J. Chem.*, 1964, **42**, 160 (pmr, config)Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1968, **4**, 194; 1970, **6**, 588; 1974, **10**, 411; 476; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 167; 1970, **6**, 603; 1974, **10**, 418; 481 (isol)Snatzke, G. et al., *Tetrahedron*, 1969, **25**, 5059 (ord)Curtis, D.R. et al., *Nature (London)*, 1970, **226**, 1222 (pharmacol)Teitel, S. et al., *J.O.C.*, 1972, **37**, 1879 (synth)Gorinsky, C. et al., *J. Cryst. Mol. Struct.*, 1973, **3**, 299 (cryst struct)Preininger, V. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 699 (isol)Mehra, K. et al., *Indian J. Chem., Sect. B*, 1976, **14**, 844 (isol, ir, uv, pmr)Markosyan, S.S. et al., *CA*, 1977, **86**, 136375j; 1978, **88**, 133268g (isol)Nalliah, B.L. et al., *Can. J. Chem.*, 1977, **55**, 922 (synth)Kametani, T. et al., *J.C.S. Perkin 1*, 1977, 390 (isol)Hussain, S.F. et al., *J. Nat. Prod.*, 1981, **44**, 169; 475 (isol)

Benzodiazepine/GABA Recept. Chloride Channels: 1986, (Eds., Olsen, R.W. et al), A.R. Liss, 1986, (pharmacol, book)

Marek, R. et al., *Magn. Reson. Chem.*, 2002, **40**, 687-692 (N-15 nmr)**Bicucullinidine**

[78416-83-8]

C₂₂H₂₃NO₈ 429.426

Alkaloid from *Fumaria schrammii* (Papaveraceae). Cryst. (CHCl₃/EtOH). Mp 265-266°.

1-Deoxo: 5-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]acetyl]-1,3-benzodioxole-4-carboxylic acid, 9CI, **Adlumiceine**
[51059-66-6]

C₂₂H₂₅NO₇ 415.442

Alkaloid from *Corydalis sempervirens* and *Fumaria schrammii* (Papaveraceae). Isol. as an inseparable mixt. with Adlumidiceine A-143.

9-Deoxo: 5-[2-[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]-2-oxoethyl]-1,3-dioxole-4-carboxylic acid, 9CI, **Isoadlumiceine**
[223738-87-2]

C₂₂H₂₅NO₇ 415.442


Alkaloid from *Thalictrum triternatum*. Cryst. (CHCl₃/MeOH). Mp 218-220°.

Preininger, V. et al., *Phytochemistry*, 1973, **12**, 2513 (*Adlumiceine*, pmr, ms, struct)Kiryakov, H.G. et al., *Phytochemistry*, 1981, **20**, 1721 (*Bicucullinidine*)Popova, M.E. et al., *Planta Med.*, 1982, **40**, 156 (*Adlumiceine*, isol, ir, pmr, ms)Kintsurashvili, L. et al., *CA*, 1999, **130**, 322950v (*Isoadlumiceine*)

Bicyclomahanimbicine

B-121

1,2,2a,10,10c,11,11a,11b-Octahydro-2a,7,11,11-tetramethyl-3-oxa-10-azacyclobut[3,4]indeno[5,6-a]fluorene, 9CI [28613-80-1]

R = CH₃, R' = HC₂₃H₂₅NO 331.457

Alkaloid from the leaves of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 218° dec. Probably an artifact. Also obt. by shaking a sol. of Mahanimbicine, M-46 with ion-exchange resin (H⁺).

Kureel, S.P. *et al.*, *Chem. Ind. (London)*, 1970, 958 (*uv, ir, pmr, struct, synth*)

Bicyclomahanimbicine

B-122

1,2,2a,10,10c,11,11a,11b-Octahydro-2a,4,11,11-tetramethyl-3-oxa-10-azacyclobut[3,4]indeno[5,6-a]fluorene, 9CI [31077-94-8]

As Bicyclomahanimbicine, B-121 with R = H, R' = CH₃

C₂₃H₂₅NO 331.457

Alkaloid from the leaves of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 145°. [α]_D²³ -1.23 (CHCl₃). Probably an artifact. Also obt. readily from Mahanimbicine in M-47.


N-Me: Mp 156°.

Kureel, S.P. *et al.*, *Tet. Lett.*, 1969, 3857 (*uv, ir, pmr, ms*)

Bandaranayake, W.M. *et al.*, *J.C.S. Perkin 1*, 1974, 998 (*struct*)

Biemnadin

B-123

C₂₇H₁₉N₅O 429.48**(±)-form** [152053-10-6]

Alkaloid from the Okinawan marine sponge *Biemna* sp. and *Biemna fortis*. Cytotoxic. Neuronal differentiation inducer. Yellow cryst. (as hydrochloride). Mp 300° (hydrochloride). Closely related to Eudistone A, E-291. λ_{max} 229 (ε 29000); 256 (ε 17000); 370 (ε 5000) (MeOH).

Zeng, C.-M. *et al.*, *Tetrahedron*, 1993, 49, 8337-8342 (*isol, uv, ir, pmr, cmr, cryst struct*)
Aoki, S. *et al.*, *Bioorg. Med. Chem.*, 2003, 11, 1969-1973 (*isol, pmr, cmr*)

Biflorine†

B-124

C₁₇H₁₇NO₄ 299.326

Struct. unknown. Alkaloid from *Oldenlandia biflora* (Rubiaceae) whole plant. Antibiotic. Possesses fungicidal props. Cryst. (EtOH). Mp 206°. [α]_D -135.4. Oxidn. gives Biflorone, B-125. Conts. 1MeO, 3OH and tertiary N, resistant to redn.

Hydrochloride: Mp 259° (sealed tube).

Turns yellow before melting.

Nitrate:

Yellow cryst. Mp 278°.

Methiodide: Mp 124°.

Chauhan, R.N.S. *et al.*, *J. Indian Chem. Soc.*, 1952, 29, 386-390; 1954, 31, 741-745

Biflorone

B-125

C₁₇H₁₅NO₄ 297.31

Struct. unknown. Alkaloid from *Oldenlandia biflora* (Rubiaceae), also obt. by oxidn. of Biflorine, B-124. Hygroscopic brownish-yellow cryst. Mp 98°. [α]_D +193.4.

Hydrochloride:

Cryst. (EtOH). Mp 152°.

Nitrate: Mp 156°.


Methiodide: Mp 139°.

Chauhan, R.N.S. *et al.*, *J. Indian Chem. Soc.*, 1952, 29, 386-390; 1954, 31, 741-745

4,4''-Bi[N-4-hydroxycinnamoylserotonin]

B-126

4,4''-Bis(N-p-coumaroylserotonin)

C₃₈H₃₄N₄O₆ 642.71**(E,E)-form** [175702-01-9]

Isol. from safflower (*Carthamus tinctorius*) oil cake. Antioxidant. Powder. Mp 180-182°. λ_{max} 221 (ε 43000); 294 (ε 35000); 305 (ε 34700) (MeOH).

3'-Methoxy: 4-[N-(p-Coumaroyl)serotonin-4''-yl]-N-feruloylserotonin

[175702-02-0]

C₃₉H₃₆N₄O₇ 672.736

From *Carthamus tinctorius* oil cake. Antioxidant. Powder. Mp 179-181°.

λ_{max} 221 (ε 70500); 292 (ε 51000); 309 (ε 52700) (MeOH).

3',3'''-Dimethoxy: 4,4''-Bi[N-(4-hydroxy-3-methoxycinnamoyl)serotonin]. 4,4''-Bis(N-feruloyl)serotonin

[175702-03-1]

C₄₀H₃₈N₄O₈ 702.762


From *Carthamus tinctorius* oil cake.

Antioxidant. Mp 158-160°. λ_{max} 289 (ε 35400); 317 (ε 38800) (MeOH).

Zhang, H.-L. *et al.*, *Chem. Pharm. Bull.*, 1997, 45, 1910-1914 (*isol, uv, ir, pmr, cmr*)

12,12'-Bi[11-hydroxycoronaridine]

B-127

C₄₂H₅₀N₄O₆ 706.88

Alkaloid from the leaves of *Bonafousia tetrastachya* (preferred genus name *Tabernaemontana*) and *Tabernaemontana citrifolia* (Apocynaceae). Amorph. [α]_D -43 (c, 0.37 in CHCl₃).

Mono-Me ether: Obovatine

[163135-91-9]

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from leaves of *Stemmadenia obovata* (Apocynaceae). Amorph. [α]_D²⁵ +17.8 (c, 0.13 in CHCl₃).

Damak, M. *et al.*, *Tet. Lett.*, 1976, 3531 (*uv, ir, pmr, cmr, ms, struct*)

Abaul, J. *et al.*, *J. Nat. Prod.*, 1989, 52, 1279 (*isol*)


Valencia, E. *et al.*, *J. Nat. Prod.*, 1995, 58, 134 (*Obovatine*)

1,1'-Bi[2-hydroxy-3-methylcarbazole]

B-128

3,3'-Dimethyl-[1,1'-bi-9H-carbazole]-2,2'-diol, 9CI

[155519-83-8]

C₂₆H₂₀N₂O₂ 392.456


Alkaloid from roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Oil.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, 41, 2096 (*isol, uv, ir, pmr, ms, struct*)

Knölker, H.J. *et al.*, *Synlett*, 1996, 737 (*synth*)

Bi[5-hydroxynoracronycine]

B-129




C₃₈H₃₄N₂O₈ 646.695

Alkaloid from the roots of *Citrus paradisi* (grapefruit). Yellow cubes (Me₂CO). Mp 207-209°. Racemic. λ_{max} 202 ; 231 ; 271 ; 290 (sh) ; 345 (sh) (EtOH).

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 693-696 (*isol, synth, uv, ir, pmr, cmr, ms*)

3,3'-Bi-1H-indole-5,5',6,6'-tetrol B-130

5,5',6,6'-Tetrahydroxy-3,3'-bi-1H-indole [390401-91-9]

C₁₆H₁₂N₂O₄ 296.282


The synthesis in 2004 suggests that the struct. needs revision. Constit. of beet-root peel, *Beta vulgaris*.

Kujala, T. *et al.*, *Z. Naturforsch., C*, 2001, **56**, 714-718 (*isol, pmr, cmr, ms*)

Mee, S.P.H. *et al.*, *Tetrahedron*, 2004, **60**, 3695-3712 (*synth*)

[3,3'-Biindoline]-2,2'-dione, 8CI B-131

3-(1,2-Dihydro-2-oxo-3H-indol-3-ylidene)-1,3-dihydro-2H-indol-2-one, 9CI. **Isoidigotin**. Isoindigo. Indigo brown [476-34-6] [180479-95-2]

C₁₆H₁₀N₂O₂ 262.267

Minor constit. of various plant spp. incl. *Indigofera* spp., *Isatis tinctoria* and *Polygonum tinctorium*. Dark red-violet cryst. (hexane/petrol). Mp 364° (>325°). λ_{max} 396 (ε 10900); 483 (ε 5100) (tetrachloroethane).

Perkin, A.G. *et al.*, *J.C.S.*, 1907, **91**, 279-288 (*isol*)


Eller-Pandraud, H. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1959, **248**, 2581-2583 (*cryst struct*)

Papageorgiou, C. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 1079-1083 (*synth*)

Maugard, T. *et al.*, *Phytochemistry*, 2001, **58**, 897-904 (*synth, biosynth, uv*)

8',11O-Biisocorydine B-132

8',11O-Biisocorydine

C₄₀H₄₄N₂O₈ 680.796


(6aS,6a'S)-form [176901-25-0]

Alkaloid from leaves of *Dehaasia triandra*. Amorph. solid. [α]_D²⁰ +89 (c, 1.0 in MeOH).

Lee, S.-S. *et al.*, *Tetrahedron*, 1996, **52**, 6561 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

8,8'-Biisocorydine B-133

8',11O-Biisocorydine



(6aS,6a'S,R_{axial})-form

C₄₀H₄₄N₂O₈ 680.796

(6aS,6a'S,R_{axial})-form

Alkaloid from leaves of *Dehaasia triandra*. Also obt. by oxidative coupling of Isocorydine, I-208. Amorph. solid. [α]_D²⁰ +14 (c, 0.5 in MeOH).

(6aS,6a'S,S_{axial})-form


From leaves of *Dehaasia triandra*. Also obt. by oxidative coupling. Amorph. solid. [α]_D²⁰ +176 (c, 0.5 in MeOH).

Lee, S.-S. *et al.*, *Tetrahedron*, 1996, **52**, 6561 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

8,8'-Bikoenigine B-134

8,8'-Biskoenigine

[477890-82-7]


C₃₈H₃₆N₂O₆ 616.712

Alkaloid from the dried leaves of *Murraya koenigii*. Brown gum. [α]_D¹⁷ +139.6 (c, 0.01 in CHCl₃).

Wang, Y.-S. *et al.*, *J. Nat. Prod.*, 2003, **66**, 416-418 (*isol, pmr, cmr*)

Bikoenuinone A B-135

1'-Methoxy-3,3'-dimethyl-[2,4'-bi-9H-carbazole]-1,4-dione, 9CI [155519-84-9]


C₂₇H₂₀N₂O₃ 420.467

Alkaloid from roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Orange oil.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 2096 (*isol, uv, ir, pmr, ms, struct*)

Bilain A B-136

[959779-92-1]

C₂₀H₂₆N₂O₅S₂ 438.568

Related to Bis(methylthio)silvatin. Prod. by the marine-derived *Penicillium bilaii* MST-MF667. Brown oil. [α]_D²⁰ -10 (c, 0.25 in EtOH).

β-Alanine amide: **Bilain C**

[959779-94-3]

C₂₃H₃₁N₃O₆S₂ 509.646


Prod. by *Penicillium bilaii* MST-MF667. Brown oil.

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1746-1752 (*isol, pmr, cmr*)

***Cypridina* Biluciferyl** B-137

Cypridina Biluciferol

[96700-80-0]

C₄₄H₅₂N₁₄O₂ 808.987

Dimer of *Cypridina* Luciferin, L-281. Isol. from *Cypridina* sp. Bioluminescent compd.


Toya, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 239 (*uv, pmr, struct*)

Biochanin C**B-138**C₁₆H₁₃N₃O₄ 311.296

Struct. unknown. Isol. from seeds of *Cicer arietinum* (chickpea). Large prisms (EtOH or H₂O). Sol. H₂O. Mp 315°. Conts. one OMe group, unsatd. to Br₂. Siddiqui, S. *et al.*, *J. Sci. Ind. Res.*, 1945, **4**, 68-70; *CA*, **40**, 1502
 Warsi, S.A. *et al.*, *Pak. J. Sci. Res.*, 1951, **3**, 85-88; *CA*, **47**, 3305f

Biotin, INN, USAN**B-139**

Hexahydro-2-oxo-1H-thieno[3,4-d]imidazole-4-pentanoic acid, 9CI. Vitamin H. Coenzyme R. Vitamin B₇. Factor S† [58-85-5] [22879-79-4]

C₁₀H₁₆N₂O₃S 244.314

Occurs in yeast, eggs, liver. Prod. by various microorganisms and isol. from various higher plant sources, e.g. sweet corn seedlings, leaves of *Raphanus sativus*. Bacterial growth factor for egg white injury. Fine needles (H₂O). Sol. H₂O. Mp 232-233°. [α]_D²⁰ +91 (c, 1 in 0.1M NaOH). Log P -0.18 (calc).

► Shows exp. reproductive effects.

S-Oxide: Biotin sulfoxide

[3376-83-8]

C₁₀H₁₆N₂O₄S 260.313

Isol. from *Aspergillus niger*. Polymorphic plates (H₂O). Mp 238° part. dec. [α]_D²⁰ -39.5 (c, 1.01 in 0.1M NaOH).

Me ester: [608-16-2]

C₁₁H₁₈N₂O₃S 258.341

Cryst. Mp 166.5°. [α]_D¹⁵ +82 (c, 0.45 in MeOH).

Succinimidoyl ester: [35013-72-0]

C₁₄H₁₉N₃O₅S 341.387

Reagent for biotinylation of proteins and nucleic acids. Cryst. (2-propanol). Mp 196-198°.

Amide: Biotinamide

[6929-42-6]

C₁₀H₁₇N₃O₂S 243.329

Formed by the action of *Rhodotorula flava* on biotin. Rosettes (MeOH or H₂O). Mp 242-244°. [α]_D²³ +80 (c, 2.5 in EtOH).

N^t-Benzyl: N-Benzylbiotin. γ-Biotin

[76335-62-1]

C₁₇H₂₂N₂O₃S 334.438

Needles (MeOH aq.). Mp 183-184°. Originally descr. as a natural vitamin under the name γ-biotin and descr. as N-phenylbiotin. Later shown to be an artifact with no vitamin activity.

[10406-89-0]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 809C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1317C (nmr)

Harris, S.A. *et al.*, *J.A.C.S.*, 1944, **66**, 1756-1757 (synth, resohn)

Wolf, D.E. *et al.*, *J.A.C.S.*, 1951, **73**, 4142-4144 (amide, synth)

Jansen, A.B.A. *et al.*, *J.C.S.*, 1962, 4909-4914; 1964, 1530-1531 (γ-Biotin)

Trotter, J. *et al.*, *Biochemistry*, 1966, **5**, 713-714 (cryst struct, abs config)

Green, N.M. *et al.*, *J.C.S.(C)*, 1970, 1330-1333 (cd, ord)

De Titta, G.T. *et al.*, *J.A.C.S.*, 1976, **98**, 1920-1926 (cryst struct)

Marx, M. *et al.*, *J.A.C.S.*, 1977, **99**, 6754-6756 (synth, bibl)

Ohrui, H. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 865-868 (synth)

Vasilevskis, J. *et al.*, *J.A.C.S.*, 1978, **100**, 7423-7424 (synth, bibl)

Uskokovic, M.R. *et al.*, *Kirk-Othmer Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **24**, 41 (rev)

Baggiolini, E.G. *et al.*, *J.A.C.S.*, 1982, **104**, 6460-6462 (synth)

Whitney, R.A. *et al.*, *Can. J. Chem.*, 1983, **61**, 1158-1160 (synth)

Bonjour, J.-P. *et al.*, *Handbook of Vitamins*, (ed. Machlin, L.J.), M. Dekker, New York, 1984, 403 (rev)

Dakshinamurti, K. *et al.*, *Ann. N.Y. Acad. Sci.*, (Eds.), 1985, **447**, (book)

Bentley, R. *et al.*, *Trends Biochem. Sci.*, 1985, **10**, 51-56 (rev, abs config)

Al-Hakim, A.H. *et al.*, *Nucleic Acids Res.*, 1986, **14**, 9965-9976 (succinimidoyl ester)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 1715

Lee, H.L. *et al.*, *Tetrahedron*, 1987, **43**, 4887-4903 (synth)

Corey, E.J. *et al.*, *Tet. Lett.*, 1988, **29**, 57-60 (synth)

Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, BGD100

Senuma, M. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 882-887 (rev, synth)

Parameswaran, K.N. *et al.*, *Org. Prep. Proced. Int.*, 1990, **22**, 119-121 (succinimidoyl ester, synth)

Alcázar, V. *et al.*, *Tetrahedron*, 1990, **46**, 1057-1062 (synth)

Bihovsky, R. *et al.*, *Tetrahedron*, 1990, **46**, 7667-7676 (synth)

De Titta, G.T. *et al.*, *J.A.C.S.*, 1994, **116**, 6485-6493 (cryst struct, nmr)

Moolenaar, M.J. *et al.*, *Angew. Chem., Int. Ed.*, 1995, **34**, 2391-2393 (synth)

Deroose, F.D. *et al.*, *J.O.C.*, 1995, **60**, 321-330 (synth, bibl)

Nicolaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, 285 (bibl, synth)

De Clercq, P.J. *et al.*, *Chem. Rev.*, 1997, **97**, 1755-1792 (rev, synth)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1336

Chen, F.-E. *et al.*, *Synthesis*, 2000, 2004-2008; 2003, 2155-2160 (synth, ir, pmr, ms)

Mori, Y. *et al.*, *Heterocycles*, 2002, **58**, 125-127 (synth)

Seki, M. *et al.*, *J.O.C.*, 2002, **67**, 5527-5536 (synth)

Shimizu, T. *et al.*, *Yakugaku Zasshi*, 2003, **123**, 43-52 (synth)

Seki, M. *et al.*, *Chem. Eur. J.*, 2004, **10**, 6102-6110 (synth)

Kimura, M. *et al.*, *Tet. Lett.*, 2004, **45**, 1635-1637 (synth, bibl)

Chavan, S.P. *et al.*, *Tetrahedron*, 2005, **61**, 9273-9280 (synth)


Roje, S. *et al.*, *Phytochemistry*, 2007, **68**, 1904-1921 (biosynth, rev)

Huang, J. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 1436-1443 (synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, VSU100

Bioxalomycin α1**B-140**

Antibiotic 31F508α1. 31F508α1 [157207-82-4]

C₂₀H₂₅N₃O₅ 387.435

Isol. from the marine-derived *Streptomyces viridostaticus* ssp. *littoralis* (LL-31F508). Exhibits potent antimicrobial and antitumor activity. Powder. Sol. MeOH, H₂O, DMSO, Me₂CO; poorly sol. CH₂Cl₂, Me₂CO, MeCN, CHCl₃. λ_{max} 290 ; 294 (MeOH) (Berdy).

N-Me: Bioxalomycin α2. Antibiotic

31F508α2. 31F508α2

[157207-83-5]

C₂₁H₂₇N₃O₅ 401.461

From *Streptomyces viridostaticus* ssp. *littoralis*. Exhibits potent antimicrobial and antitumor activity. Powder. Sol. MeOH, Me₂CO, DMSO, H₂O; poorly sol. CH₂Cl₂, CHCl₃, MeCN, Me₂CO. [α]_D²⁵ +31 (MeOH). Unstable in soln. λ_{max} 292 (ε 2729) (MeOH) (Berdy).

10,13-Quinone: Bioxalomycin β1. Antibiotic

31F508β1. 31F508β1

[157207-84-6]

C₂₀H₂₃N₃O₅ 385.419

Isol. from *Streptomyces viridostaticus* ssp. *littoralis*. Potent antitumor and antimicrobial agent. Sol. CH₂Cl₂, DMSO, MeOH, CHCl₃; poorly sol. H₂O, hexane.

10,13-Quinone, N-Me: Bioxalomycin β2. Antibiotic

31F508β2. 31F508β2

C₂₁H₂₅N₃O₅ 399.446

From *Streptomyces viridostaticus* ssp. *littoralis*. Exhibits potent antitumor and antimicrobial activity. Sol. CH₂Cl₂, CHCl₃, DMSO, MeOH; poorly sol. H₂O, hexane. Erroneously given the same CAS no. as Naphthrydinomycin A, N-32. λ_{max} 270 (ε 10508); 370 (MeOH) (Berdy). λ_{max} 264 ; 370 (MeOH/HCl) (Berdy).


Bernan, V.S. *et al.*, *J. Antibiot.*, 1994, **47**, 1417-1424 (isol, props)




Zaccardi, J. *et al.*, *J.O.C.*, 1994, **59**, 4045-4047 (isol, ir, uv, pmr, cmr, ms)

1,1'-Bipiperidine, 9CI**B-141**

1,1'-Dipiperidine. Fabioline

[6130-94-5]



C₁₀H₂₀N₂ 168.281Alkaloid from aerial parts of *Cassia grandis*. Needles. Mp 19.5-20.5°.*Picrate*: Mp 153-154°.Mackay, D. et al., *J.C.S. (C)*, 1966, 813 (*synth*)Ogawa, K. et al., *J.C.S. Perkin I*, 1982, 3031 (*synth*)Valencia, E. et al., *Fitoterapia*, 1995, **66**, 476 (*isol, ir, pmr, cmr, ms, struct*)**Bipowine****B-142**7,7'-Bis(6a,7-didehydrowilsonirine)
[107882-21-3]C₃₈H₃₆N₂O₈ 648.711Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Cryst. (Me₂CO). Mp 249-251°.Jossang, A. et al., *J. Nat. Prod.*, 1986, **49**, 1028 (*uv, pmr, cmr, ms, struct*)Ruchirawat, S. et al., *Heterocycles*, 2001, **55**, 371-376 (*synth*)**Bipowinone****B-143**7,7'-Bispancoridine
[107882-20-2]C₃₈H₂₈N₂O₈ 640.648Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Cryst. (Me₂CO). Mp 295°.Jossang, A. et al., *J. Nat. Prod.*, 1986, **49**, 1028 (*uv, ir, pmr, cmr, ms, struct, synth*)Ruchirawat, S. et al., *Heterocycles*, 2001, **55**, 371-376 (*synth*)**2,2'-Bipyridine, 9CI****B-144**2,2'-Bipyridyl. α,α'-Bipyridyl. Bipy
[366-18-7]
[37275-48-2]C₁₀H₈N₂ 156.187Used in photometric detn. of Fe(II) (λ_{max} 522 nm, ε 8700, pH 2-9); as a masking agent for Zn, Cd, Cu. Fe(II) chelate is used as a redox indicator. Complexingagent for prepn. of inorganic complexes. Prisms (petrol). Sol. H₂O, EtOH, Et₂O, C₆H₆, CHCl₃, dil. acids. Mp 69.5°. Bp 272.5°. pK_{a1} 4.3 (25°, H₂O). Fe^{II} salts → red col.▶ LD₅₀ (rat, orl) 100 mg/kg. Exp. teratogen. DW1750000*Methodide*: [77972-47-5]Yellow-white needles (Et₂O/MeCN). Mp 143-145°.*Chlorochromate*: [76899-34-8]C₁₀H₉ClCrN₂O₃ 292.642

Used to oxidise acid labile primary and secondary alcohols to carbonyl compounds. Yellow cryst.

Mono-N-oxide: [33421-43-1]C₁₀H₈N₂O 172.186

Mp 59°.

▶ DW1840000

N,N'-Dioxide: [7275-43-6]C₁₀H₈N₂O₂ 188.185

Mp 310° dec.

▶ DW1770000

N,N'-Di-Me: 1,1'-Dimethyl-2,2'-bipyridinium(2+)C₁₂H₁₄N₂²⁺ 186.256Isol. from mollusc *Callista chione*.*Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 731C (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 239B; 483B (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 1513B (*ir*)Fresenius, W. et al., *Fresenius' Z. Anal. Chem.*, 1965, **209**, 340 (*detn, Fe*)Schilt, A.A. et al., *Analytical Applications of 1,10-Phenanthroline and Related Compounds*, Pergamon, London, 1969, (*use*)Nordén, B. et al., *Acta Chem. Scand.*, 1972, **26**, 429 (*spectra*)French, W.J. et al., *Analyst (London)*, 1972, **97**, 828 (*detn, Fe*)*Org. Synth.*, *Coll. Vol.*, **5**, 1973, 102Keats, N.G. et al., *J. Het. Chem.*, 1976, **13**, 369 (*ms*)Guziec, F.S. et al., *Synthesis*, 1980, 691 (*chlorochromate, synth, use*)Cheng, K.L. et al., *Handbook of Organic Analytical Reagents*, CRC Press, Boca Raton, 1982, 309 (*use*)Wenkert, D. et al., *J.O.C.*, 1983, **48**, 283 (*oxides*)Moran, D.B. et al., *J. Het. Chem.*, 1986, **23**, 1071 (*monoxide*)Almenningen, A. et al., *Acta Chem. Scand.*, 1989, **43**, 932 (*ed, struct, bibl*)*Encyclopaedia of Reagents for Organic Synthesis*, (ed. Paquette, L.A.), Wiley, 1995, **1**, 409-410 (*chlorochromate, use*)Norrby, T. et al., *Acta Chem. Scand.*, 1998, **52**, 77-85 (*monoxide, methodide*)Demnitz, F.W.J. et al., *Org. Prep. Proced. Int.*, 1998, **30**, 467-469 (*monoxide*)Vagias, C. et al., *Nat. Prod. Lett.*, 2000, **14**, 425-428 (*N,N'-di-Me, isol*)Bowen, R.J. et al., *Acta Cryst. C*, 2004, **60**, o113-o114 (*cryst struct*)Rajalakshmanan, E. et al., *Synth. Commun.*, 2005, **35**, 891-895 (*synth, pmr*)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, BGO500

2,3'-Bipyridine, 9CI**B-145**2,3'-Dipyridyl. *Isonicotine*
[581-50-0]
[37275-48-2]C₁₀H₈N₂ 156.187Occurs in tobacco (*Nicotiana tabacum*) (Solanaceae) and the marine hoplone-merterine *Amphiporus angulatus*. Used for extraction-photometric detn. of Cu(II), and for extraction of Sc. Liq. Insol. H₂O. d₂₀⁴ 1.14. Bp 298°. n_D²⁰ 1.6223.

▶ Crustacean convulsant, but weak activity in mice. Mutagenic props. DW1755000

Perchlorate (1:2): Mp 215-216°.*Picrate*: Mp 153-154°.*Dipicrate*: Mp 165-168°.*Methodide* (1:2): [63095-08-9]

Cryst. (EtOH aq.). Mp 210-214° dec.


I-Oxide: [39182-27-9]C₁₀H₈N₂O 172.186

Solid. Mp 135-137°. Prepared indirectly, not obt. by direct oxidation.

I'-Oxide: [39182-28-0]C₁₀H₈N₂O 172.186Cryst. + 1H₂O (Me₂CO/hexane). Mp 78°.*I,I'-Dioxide*: [30651-23-1]C₁₀H₈N₂O₂ 188.185


Cryst. (MeOH). Mp 240-243°.

[42907-60-8]

Smith, J.R. et al., *J.A.C.S.*, 1930, **52**, 397 (*synth*)Morgan, G.T. et al., *J.C.S.*, 1932, 20 (*synth*)Späth, E. et al., *Ber.*, 1936, **69**, 2448 (*isol*)Talipov, S.T. et al., *Uzb. Khim. Zh.*, 1963, **22** (*detn, Cu*)Leete, E. et al., *J.A.C.S.*, 1969, **91**, 1697 (*synth*)Alimarin, I.P. et al., *CA*, 1973, **78**, 8393m (*sepn, Sc*)Kem, W.R. et al., *Experientia*, 1976, **32**, 684-686 (*isol, ms, tox*)Leete, E. et al., *J.A.C.S.*, 1976, **98**, 6326 (*biosynth*)Moran, D.B. et al., *J. Het. Chem.*, 1986, **23**, 1071 (*oxides*)Zoltewicz, J.A. et al., *J.O.C.*, 1992, **57**, 2392 (*deriv, synth, pmr*)Zoltewicz, J.A. et al., *Tetrahedron*, 1995, **51**, 3103 (*I-oxide*)Núñez, A. et al., *Tetrahedron*, 2004, **60**, 6217-6224 (*synth, pmr, cmr*)**9,9'-Bi-9H-pyrido[3,4-b]indole, 9CI****B-146***N,N'-Bi-β-carboline*
[85753-68-0]C₂₂H₁₄N₄ 334.379Alkaloid from the ascidian *Didemnum* sp. Also obt. by photochemical dimerisation of β-Carboline, C-130. Needles (EtOAc/hexane). Mp 209-210° dec.Erra-Balsells, R. et al., *Tetrahedron*, 1983, **39**, 33-39 (*synth*)Erra-Balsells, R. et al., *Magn. Reson. Chem.*, 1988, **26**, 1109-1112 (*cmr*)Kearns, P.S. et al., *J. Nat. Prod.*, 1995, **58**, 1075-1076 (*isol, uv, ir, pmr, cmr, ms*)

2,3'-Bipyrrolidine**B-147**

[107334-13-4]

C₈H₁₆N₂ 140.228


N,N'-Di-Me: 1,1'-Dimethyl-2,3'-bipyrrolidine. 2,3'-Bi(1-methylpyrrolidine) [6602-17-1]

C₁₀H₂₀N₂ 168.281

Alkaloid from the roots of *Nicotiana tabacum* (tobacco). Liq. Bp_{0.1} 37-38°. Isol. as a mixt. of two diastereoisomers. Properties given for unspecified isomer composition.

N,N'-Di-Me, picrate (1:2):

Yellow cryst. (EtOH aq.). Mp 215-216°.

Bocz, A.K. *et al.*, *Chem. Ber.*, 1966, **99**, 1923-1931 (*N,N'*-di-Me, synth)Wei, X. *et al.*, *Life Sci.*, 2005, **78**, 495-505 (*N,N'*-di-Me, isol, synth, pmr, ms)**Birnbaum A****B-148**C₁₆H₂₀N₆O₄ 360.372

Alkaloid from *Leucocoprinus birnbaumii*. Yellow solid. λ_{max} 212 (log ε 4.42); 250 (log ε 4.03); 322 (log ε 3.96) (MeOH).

7-Hydroxy: **Birnbaum B**C₁₆H₂₀N₆O₅ 376.371

Alkaloid from *Leucocoprinus birnbaumii*. Yellow solid. λ_{max} 214 (log ε 4.74); 248 (log ε 4.34); 267 (sh) (log ε 4.19); 356 (log ε 4.16) (MeOH).

Bartsch, A. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 2957-2959 (isol, ir, pmr, cmr, ms)**Bis(4-aminobutyl)bis(3-aminopropyl)ammonium(1+)****B-149**

4-Amino-N-(4-aminobutyl)-N,N-bis(3-aminopropyl)-1-butanaminium, 9CI.

N⁵,N⁵-Bis(3-aminopropyl)homospermidine

[143085-77-2]

(H₂NCH₂CH₂CH₂CH₂)₂N[⊕](CH₂CH₂CH₂CH₂)₂C₁₄H₃₆N₅[⊕] 274.472

Isol. from thermophilic bacteria. Counterion not specified.

Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-747**N,N'-Bis(4-aminobutyl)-1,4-butanediamine, 9CI****B-150**1,6,11,16-Tetraazahexadecane. 1,14-Diamino-5,10-diazatetradecane. **Homospermine**. PA444

[45185-87-3]

H₂NCH₂(CH₂)₃¹⁰NH(CH₂)₄⁵NH(CH₂)₃C-H₂NH₂C₁₂H₃₀N₄ 230.396

Prod. by a *Rhizobium* sp. Also found in *Vicia sativa*, *Vicia villosa*, *Crotalaria spectabilis* and *Stichopus japonica*.

N¹,N¹⁴-Di-Et: N,N'-Bis[4-(ethylamino)-butyl]-1,4-butanediamine, 9CI. N¹,N¹⁴-Diethylhomospermine. DEHSPM [119422-08-1]

[156886-86-1 (hydrochloride)]

C₁₆H₃₈N₄ 286.503

Antiproliferative agent. Plates (EtOH) (as tetrahydrochloride).

N⁵-(4-Aminobutyl): N⁵-(4-Aminobutyl)-homospermine

[139035-35-1]

C₁₆H₃₉N₅ 301.518

Constit. of *Vicia sativa* and *Vicia villosa*.

N⁵,N¹⁰-Bis(4-aminobutyl): N⁵,N¹⁰-Bis(4-aminobutyl)homospermine


[139035-38-4]

C₂₀H₄₈N₆ 372.639

Constit. of *Vicia sativa*.

Samejima, K. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3428-3435 (synth)Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (synth, cmr)Fujihara, S. *et al.*, *Biochem. Biophys. Res. Commun.*, 1989, **165**, 659-666 (occur)Casero, R.A. *et al.*, *Cancer Res.*, 1989, **49**, 639-643; 2959-2964 (DEHSPM, pharmacol)Hamana, K. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur)Hamana, K. *et al.*, *Phytochemistry*, 1991, **30**, 3319-3322 (occur)Bergeron, R.J. *et al.*, *J. Med. Chem.*, 1994, **37**, 3464-3476; 1996, **39**, 2461-2471 (DEHSPM, synth, pharmacol)Hamana, K. *et al.*, *Can. J. Bot.*, 1996, **74**, 1766-1772 (occur)Bergeron, R.J. *et al.*, *Drug Metab. Dispos.*, 1996, **24**, 334-343 (DEHSPM, metab, pharmacokinetic)**1,3-Bis(4-aminomethyl)phenyl]-2-octadecanoylglycerol****B-151**

[790693-27-5]

C₃₅H₅₆N₂O₄ 568.838

Alkaloid from the twigs of *Carapa guianensis*. Powder. Mp 115-116°. λ_{max} 209 ; 229 ; 256 ; 277 ; 287 (MeOH).

Qi, S.-H. *et al.*, *Pharmazie*, 2004, **59**, 488-490 (*Carapa guianensis* alkaloid)**N,N'-Bis(3-aminopropyl)-1,4-butanediamine****B-152**N⁴-(3-Aminopropyl)spermidine

[128837-67-2]

(H₂NCH₂CH₂CH₂)₂N(CH₂)₃CH₂NH₂C₁₀H₂₆N₄ 202.342

Isol. from the thermophilic bacteria *Bacillus schlegelii*, *Thermoleophilum album*, *Thermoleophilum minutum* and

Hydrogenobacter thermophilus TK-6.Hamana, K. *et al.*, *FEMS Microbiol. Lett.*,1990, **66**, 35-38 (*Thermoleophilum constits*)Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-747 (*Bacillus schlegelii* constit,*Hydrogenobacter thermophilus* constit)**N,N'-Bis(3-aminopropyl)-1,2-ethanediamine****B-153**

N',N''-1,2-Ethanediybis-1,3-propanediamine, 9CI. 1,5,8,12-Tetraazadodecane.

1,10-Diamino-4,7-diazadecane

[10563-26-5]

H₂NCH₂CH₂CH₂NHCH₂CH₂NHCH₂CH₂CH₂NH₂C₈H₂₂N₄ 174.289

Alkaloid from *Nuphar japonicum*. Bp₅ 150-160°. n_D²⁰ 1.4910.

Hydrochloride (1:4): [13493-17-9]

Solid (EtOH). Mp 275-276°.

Aldrich Library of NMR Spectra, 2nd edn., 1983, **1**, 267DAldrich Library of FT-IR Spectra, 1st edn., 1985, **3**, 398CBarefield, E.K. *et al.*, *Inorg. Synth.*, 1975, **16**, 220 (synth)Hamana, K. *et al.*, *Can. J. Bot.*, 1998, **76**, 130-133 (isol)Rubio, M. *et al.*, *Synth. Commun.*, 2002, **32**, 2441-2452 (synth, pmr, cmr)**N,N'-Bis(3-aminopropyl)-1,3-propanediamine, 9CI****B-154**

1,11-Diamino-4,8-diazaundecane.

1,5,9,13-Tetraazatridecane, 8CI. **Norspermine**. **Thermine**

[4605-14-5]


[16632-23-8]

H₂NCH₂CH₂CH₂NHCH₂CH₂CH₂NHCH₂CH₂CH₂NH₂C₉H₂₄N₄ 188.315

Metab. of various bacteria and plant spp. Mp 260° (as tetrahydrochloride).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 309B (ir)Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 503B (nmr)Dagnall, S.P. *et al.*, *J.C.S. Perkin 2*, 1984, 435 (cmr)Clay, R.M. *et al.*, *Inorg. Chem.*, 1985, **24**, 3330 (synth)Hamana, K. *et al.*, *J. Biochem. (Tokyo)*, 1985, **97**, 1595; 1991, **109**, 444 (occur)Vaultier, M. *et al.*, *Synth. Commun.*, 1992, **22**, 665 (synth)Rubio, M. *et al.*, *Synth. Commun.*, 2002, **32**, 2441-2452 (synth, pmr, cmr)**Bisancorinolate B****B-155**

[471913-82-3]


C₁₆H₁₂N₂O₁₆S₄ 616.539

Alkaloid from the sponge *Ancorina* sp. Powder (as tetra-Na salt). CAS no. refers to tetra Na salt. λ_{max} 221 (log ε 4.53); 268 (log ε 4.07); 301 (log ε 3.82) (MeOH) (tetra-Na salt).

Meragelman, K.M. *et al.*, *J.O.C.*, 2002, **67**, 6671-6677 (*isol*, *pmr*, *cmr*, *ms*)

Bisaristone A

[110192-13-7]



Relative Configuration


C₄₀H₄₆N₄O₂ 614.829

Alkaloid from the aerial parts of *Aris- totelia australisica* (Elaeocarpaceae). Cryst. (MeOH). Mp 303-305°. λ_{max} 209 (log ε 4.41); 245 (log ε 3.97); 299 (log ε 3.5) (EtOH).

Quirion, J.-C. *et al.*, *J.O.C.*, 1987, **52**, 4527-4530 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Bisavenanthramide B₁

[939790-85-9]



Relative Configuration


C₃₄H₂₈N₂O₁₂ 656.601

Isol. from elicited oat leaves (*Avena sativa*). Amorph. solid. Racemic. λ_{max} 210 (log ε 4.83); 233 (sh) (log ε 4.44); 285 (log ε 4.04); 300 (sh) (log ε 3.84) (MeOH).

Okazaki, Y. *et al.*, *J.O.C.*, 2007, **72**, 3830-3839 (*isol*, *pmr*, *cmr*, *ms*)

Bisavenanthramide B₂

[939790-86-0]


C₃₄H₃₀N₂O₁₃ 674.617

Isol. from elicited oat leaves (*Avena sativa*). Yellowish solid. Racemic. λ_{max} 204 (log ε 5.04); 265 (log ε 4.23); 317 (log ε 3.07) (MeOH).

Okazaki, Y. *et al.*, *J.O.C.*, 2007, **72**, 3830-3839 (*isol*, *pmr*, *cmr*, *ms*)

Bisavenanthramide B₅

[939790-89-3]



Relative Configuration


C₃₄H₃₀N₂O₁₃ 674.617

Isol. from elicited oat leaves (*Avena sativa*). Yellow solid. λ_{max} 204 (log ε 4.79); 224 (log ε 4.51); 263 (log ε 4.19); 328 (log ε 3.75) (MeOH).

Okazaki, Y. *et al.*, *J.O.C.*, 2007, **72**, 3830-3839 (*isol*, *pmr*, *cmr*, *ms*)

Bisavenanthramide B₆

B-160

C₃₄H₂₈N₂O₁₂ 656.601

Isol. from elicited oat leaves (*Avena sativa*). Phytoalexin. Racemic. λ_{max} 204 (log ε 4.61); 265 (log ε 4.31); 329 (log ε 4.45) (MeOH).

7',8'α-Dihydro, 7'ξ-hydroxy (1): **Bisavenanthramide B₃** [939790-87-1]

C₃₄H₃₀N₂O₁₃ 674.617

Isol. from elicited oat leaves (*Avena sativa*). Yellowish solid. Racemic. C-7 epimer of B₄. λ_{max} 205 (log ε 4.76); 225 (sh) (log ε 4.42); 267 (log ε 3.98); 319 (log ε 3.49) (MeOH).

7',8'α-Dihydro, 7'ξ-hydroxy (2): **Bisavenanthramide B₄** [939790-88-2]

C₃₄H₃₀N₂O₁₃ 674.617


Isol. from elicited oat leaves (*Avena sativa*). Yellowish solid. Racemic. C-7 epimer of B₃. λ_{max} 204 (log ε 4.87); 225 (sh) (log ε 4.59); 262 (log ε 4.19); 319 (log ε 3.67) (MeOH).

Okazaki, Y. *et al.*, *Tetrahedron*, 2004, **60**, 4765-4771 (*isol*, *pmr*, *cmr*, *ms*)

Okazaki, Y. *et al.*, *J.O.C.*, 2007, **72**, 3830-3839 (*isol*, *pmr*, *cmr*, *ms*)

***N,N'*-Bis(2,10-bisaboladien-7-yl)urea**

B-161



Absolute Configuration

C₃₁H₅₂N₂O 468.765**(6*R*,6'*R*,7*S*,7'*S*)-form** [105281-45-6]


Isol. from a *Halichondria* sp. Mp 139-141°. [α]_D²⁰ +44 (c, 1.1 in CHCl₃).

Sullivan, B.W. *et al.*, *J.O.C.*, 1986, **51**, 5134-5136 (*isol*, *pmr*, *cmr*)

2,2-Bis(6-bromo-3-indolyl)ethylamine

B-162

6-Bromo-β-(6-bromo-1*H*-indol-3-yl)-1*H*-indole-3-ethanamine, 9*CI* [135077-20-2]

C₁₈H₁₅Br₂N₃ 433.145

Alkaloid from the marine tunicate *Didemnum candidum*. Also isol. from the deep-water New Caledonian sponge *Orina* sp. Pale yellow oil.


Fahy, E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 564-569 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Bifulco, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1254-1260 (*isol*)

Bis(6-bromo-2-tryptaminy) disulfide

B-163

Bis[3-(2-aminoethyl)-6-bromo-1*H*-indol-2-yl] disulfide [622011-16-9]

C₂₀H₂₀Br₂N₄S₂ 540.345

Alkaloid from the defensive mucus of the marine snail *Calliostoma canaliculatum*. Potassium channel agonist. Neurotoxin.

Kelley, W.P. *et al.*, *J. Biol. Chem.*, 2003, **278**, 34934-34942 (*isol*, *pmr*, *cmr*, *ms*, *activity*)

Wolters, A.M. *et al.*, *J. Nat. Prod.*, 2005, **68**, 162-167 (*pmr*, *anal*)

***N,N'*-Bis[3-(carboxyamino)-4-methylphenyl]urea**

B-164

[Carbonylbis[imino(6-methyl-3,1-phenylene)]]biscarbamic acid

C₁₇H₁₈N₄O₅ 358.353

Di-Me ester:C₁₉H₂₂N₄O₅ 386.407

Isol. from the Pliocene fossil wood of *Pinus armandii*. Powder. Mp 152-154°. λ_{max} 216 (log ε 4.57); 258 (log ε 4.58) (MeOH).

Dibutyl ester:C₂₅H₃₄N₄O₅ 470.567

Isol. from the Pliocene fossil wood of *Pinus armandii*. Powder. Mp 153-155°.


Bis(2-methylpropyl) ester:C₂₅H₃₄N₄O₅ 470.567

Isol. from Pliocene fossil wood of *Pinus armandii*. Powder. Mp 153-155°.

Zhao, Y.-X. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 325-329 (*isol, pmr, cmr, ms*)

Biscephalezomine A**B-165**

[769120-40-3]

C₅₇H₇₂N₂O₁₉ 1089.198

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D²⁰ -151 (c, 1 in MeOH). λ_{max} 209 (ε 34700); 291 (ε 6400) (MeOH).

ω-Deoxy, 3'S-hydroxy: Biscephalezomine D

[769120-42-5]

C₅₇H₇₂N₂O₁₉ 1089.198

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D²⁰ -86 (c, 0.3 in MeOH). λ_{max} 207 (ε 41000); 290 (ε 4800) (MeOH).

Yoshinaga, M. *et al.*, *Tetrahedron*, 2004, **60**, 7861-7868 (*isol, cd, pmr, cmr*)

Biscephalezomine B**B-166**

[769120-41-4]

As Biscephalezomine A, B-165 with m = 2, n = 3

C₅₇H₇₂N₂O₁₉ 1089.198

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D²⁰ -126 (c, 1 in MeOH). λ_{max} 208 (ε 28200); 291 (ε 5300) (MeOH).

ω'-Deoxy: Biscephalezomine E

[768392-34-3]

C₅₇H₇₂N₂O₁₈ 1073.198

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D²⁰ -146 (c, 0.4 in MeOH). λ_{max} 208 (ε 33600); 290 (ε 4300) (MeOH).

Yoshinaga, M. *et al.*, *Tetrahedron*, 2004, **60**, 7861-7868 (*isol, cd, pmr, cmr*)

Biscephalezomine C**B-167**

[768392-32-1]

As Biscephalezomine A, B-165 with m = n = 2


C₅₆H₇₀N₂O₁₉ 1075.171

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D²⁰ -118 (c, 0.5 in MeOH). λ_{max} 207 (ε 27000); 290 (ε 4000) (MeOH).

Yoshinaga, M. *et al.*, *Tetrahedron*, 2004, **60**, 7861-7868 (*isol, cd, pmr, cmr*)

Biscryptolepine**B-168**


5,5'-Dimethyl-11,11'-bi-5H-quinoline, 9CI
[175178-79-7]

C₃₂H₂₂N₄ 462.553

Alkaloid from root bark of *Cryptolepis sanguinolenta*. Active against gram-positive bacteria and mycobacteria. Green amorph. powder.


Cimanga, K. *et al.*, *Tet. Lett.*, 1996, **37**, 1703 (*isol, uv, pmr, cmr, ms, struct*)

Cimanga, K. *et al.*, *Phytomedicine*, 1998, **5**, 209-214 (*activity*)

Bisdehydroestemoninine**B-169**C₁₇H₂₁NO₃ 287.358

Alkaloid from the roots of *Stemona tuberosa*. Amorph. yellow solid. [α]_D²⁰ -132 (c, 0.06 in CHCl₃).

Lin, L.-G. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1051-1054 (*isol, pmr, cmr*)

Bisdehydrostemoninine A**B-170**

Relative Configuration

C₂₂H₂₇NO₆ 401.458

Alkaloid from the roots of *Stemona tuberosa*. Cubic cryst. (hexane/EtOAc). [α]_D²⁰ -186 (c, 0.05 in CHCl₃).


1'-Deoxo: Bisdehydrostemoninine BC₂₂H₂₆NO₅ 387.475

Alkaloid from the roots of *Stemona tuberosa*. Amorph. yellow solid. [α]_D²⁰ -122 (c, 0.14 in CHCl₃).

Lin, L.-G. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1051-1054 (*isol, pmr, cmr, ms, cryst struct*)


N,N'-Bis[2-(3,5-diiodo-4-methoxyphenyl)ethyl]urea, 9CI**B-171**

[89631-87-8]

C₁₉H₂₀I₄N₂O₃ 831.997


Minor metab. from an unidentified didemnum sp. Mp 224-225°.

Sesin, D.F. *et al.*, *Tet. Lett.*, 1984, **25**, 403-404 (*pmr, cmr, struct*)

1,5-Bis(1,3-dimethyl-6-lumazinyl)-2-methyl-1,5-pentanedione**B-172**C₂₂H₂₂N₈O₆ 494.466**(S)-form [173324-68-0]**

Metab. from the marine polychaete *Odontosyllis undecimdongta*. Solid (MeOH). Mp 114-115°. [α]_D²⁵ -61.9 (c, 0.21 in CHCl₃).


Tanino, H. *et al.*, *Heterocycles*, 1996, **42**, 125-128 (*isol, uv, ir, pmr, ms, synth, struct*)

N,N'-Bis(3,7-dimethyl-2,6-octadienyl)guanidine**B-173**C₂₁H₃₇N₃ 331.543**(Z,Z)-form**

N,N'-Dimerolylguanidine. **Nitensidine B**
[173221-01-7]

Alkaloid from leaves of *Pterogyne nitens*. Liq.

Bolzani, V. da.S. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1683 (*isol, ir, pmr, cmr, ms, struct*)

N,N'-Bis(3,7-dimethyl-2,6-octadienyl)guanidine**B-174**C₂₁H₃₇N₃ 331.543**(E,E)-form**

N,N'-Digeranylguanidine. **Nitensidine A**
[173221-00-6]


Alkaloid from leaves of *Pterogyne nitens*. Liq.

N³,N⁴-Di-Me: Nitensidine C


[173221-02-8]

C₂₃H₄₁N₃ 359.597From leaves of *Pterogyne nitens*. Liq.Bolzani, Vda.S. et al., *J. Nat. Prod.*, 1995, **58**, 1683 (*isol, ir, pmr, cmr, ms, struct*)**N,N'-Bis(2,6-dimethylphenyl)-2,2,4,4-tetramethyl-1,3-cyclobutanediimine** **B-175**


N,N'-(2,2,4,4-Tetramethyl-1,3-cyclobutanediylidene)bis[2,6-dimethylbenzamine], 9CI [52199-12-9]

C₂₄H₃₀N₂ 346.514Alkaloid from *Arundo donax*. Prisms (hexane/CHCl₃). Mp 176-178° (169-170°).Miles, D.H. et al., *Phytochemistry*, 1993, **34**, 1277-1279 (*isol, pmr, cmr*)Mochizuki, K. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 647-651 (*synth*)**1,1-Bis(8,9-epoxy-6,8-dimethylergoline)** **B-176**

8,9:8',9'-Diepoxy-6,6',8,8'-tetramethyl-1,1'-biergoline, 9CI [132158-23-7]

C₃₂H₃₄N₄O₂ 506.646Prod. by *Penicillium sizovae*. Mycotoxin. Cryst. (EtOAc/hexane). Sol. MeOH, CHCl₃, Me₂CO, EtOAc, MeCN, CCl₄; poorly sol. H₂O. λ_{max} 240 (ε 5890); 280 (ε 3710); 293 (ε 3310); 302 (ε 2950) (MeOH).Zelenkova, N.F. et al., *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1990, **313**, 1494 (*isol, pmr, cmr*)Zelenkova, N.F. et al., *Prikl. Biokhim. Mikrobiol.*, 1992, **28**, 738-741 (*isol, struct*)**Bis(gorgiacerol)amine** **B-177**

[138828-03-2]

C₄₂H₄₇NO₁₂ 757.833Constit. of *Pseudopterogorgia acerosa*. Cryst. Mp 248-249°. [α]_D +40 (c, 0.3 in CHCl₃).Tinto, W.F. et al., *Tetrahedron*, 1991, **47**, 8679-8686 (*isol, pmr, cmr*)**N,N'-Bis(4-guanidinobutyl)oxamide** **B-178**N,N'-Bis[4-[(aminoiminomethyl)amino]butyl]ethanediamide, 9CI. N,N'-Bis-(agmatine)oxamide [151368-32-0]
HN=C(NH₂)NH(CH₂)₄
NHCOCNH(CH₂)₄NHC(NH₂)=NHC₁₂H₂₆N₈O₂ 314.39Alkaloid from the spider *Plectreurys tristis*.Quistad, G.B. et al., *Toxicon*, 1993, **31**, 920-924 (*isol, struct*)**Bishaconitine** **B-179**

[11011-58-8]

C₄₁H₅₃NO₁₃ 767.869Tentatively classified as a C₁₉-diterpenoid alkaloid. Struct. unknown. Alkaloid from the Indian crude drug Bikh (or Bish or Mitha telia) (tuberous roots of *Aconitum falconeri*) (Ranunculaceae). Light yellow amorph. powder. Mp 118-119°. [α]_D²⁵ +31.25.*Reineckate*:Microprisms + 4H₂O (EtOH aq.). Mp 179-181°.*Aurichloride*:

Microcryst. powder. Mp 124-127°.

Singh, N. et al., *Indian J. Chem.*, 1966, **4**, 39 (*isol, ir*)**Bishatisine** **B-180**C₂₄H₃₇NO₃ 387.561Tentatively classified as a C₂₀-diterpenoid alkaloid. Struct. unknown. Alkaloid from the Indian crude drug Bikh (or Bish or Mitha telia) (tuberous roots of *Aconitum falconeri*) (Ranunculaceae). Cryst. + 1½H₂O. Mp 295-296°. [α]_D²² +18.6 (EtOH).*Hydrochloride*:

Needles (EtOH). Mp 274°.

Hydrobromide:

Needles (EtOH/HBr). Mp 227-228°.

Hydroiodide:

Light yellow cryst. (EtOH/HI). Mp 271-272°.

Nitrate: Mp 238-239°.*Sulfate (2:1)*:



Rosettes of cryst. Mp 98-100°.

Perchlorate:


Needles. Mp 219-220°.


Picrate:Cryst. (Me₂CO). Mp 146-147°.*Methiodide*:

Cryst. (EtOH). Mp 269-270°.

Reineckate:Cryst. + 2H₂O (EtOH). Mp 138-139.5°.*Ac*:Needles + 3H₂O (EtOH). Mp 224-225°.Singh, N. et al., *Indian J. Chem.*, 1966, **4**, 39-42 (*isol, uv, ir*)**N,N-Bis(4-hydroxybenzyl)hydroxylamine** **B-181**4,4'-[(Hydroxyimino)bis(methylene)]biphenol, 9CI. *Gastrodamine* [245428-10-8]C₁₄H₁₅NO₃ 245.277Alkaloid from the tubers of *Gastrodia elata*. λ_{max} 238 (log ε 4.4); 279 (log ε 3.5); 282 (log ε 3.4) (MeOH).Hao, X.Y. et al., *Chin. Chem. Lett.*, 1999, **10**, 467-468 (*isol*)**3,4-Bis(5-hydroxy-1H-indol-3-yl)-1H-pyrrole-2-carboxylic acid** **B-182**C₂₁H₁₅N₃O₄ 373.367*Me ester: Cinereapyrrole A*C₂₂H₁₇N₃O₄ 387.394Isol. from wild fruit bodies of *Arcyria cinerea*. Dark brown powder. λ_{max} 283 (ε 11000) (MeOH).*5'-Deoxy, Me ester: Cinereapyrrole B*C₂₂H₁₇N₃O₃ 371.395Isol. from the fruiting bodies of *Arcyria cinerea*. Dark brown powder. λ_{max} 230 (ε 36000); 286 (ε 14000) (MeOH).Kamata, K. et al., *Chem. Pharm. Bull.*, 2005, **53**, 594-597 (*isol, pmr, cmr*)**3,3'-Bis-3-[(4-hydroxy-2-methoxyphenyl)-2-propenamide]** **B-183**

3,3'-(6,6'-Dihydroxy-2,2'-dimethoxy-[1,1'-biphenyl]-3,3'-diyl)bis[2-propenamide], 9CI. 4,4'-Dihydroxy-2,2'-dimethoxy-3,3'-dicinnamide

C₂₀H₂₀N₂O₆ 384.388**(Z,Z)-form** [203524-81-6]Constit. of *Sinularia flexibilis*.Anjaneyulu, A.S.R. et al., *Nat. Prod. Lett.*, 1997, **11**, 5-11 (*isol*)

2,5-Bis(1-hydroxy-3-methylbutyl)-3,6-dimethylpyrazine B-1843,6-Dimethyl- α,α' -bis(2-methylpropyl)-2,5-pyrazinemethanol, 9CIC₁₆H₂₈N₂O₂ 280.409

Isol. as 2 diastereoisomers of unknown config.

High-melting-formIsol. from ants of the genus *Mesoponera* (*Mesoponera castanea*, *Mesoponera castaneicolor*). Mp 129.5-130°.**Low-melting-form**From *Mesoponera castanea* and *Mesoponera castaneicolor*. Mp 99.5-100.5°.**Monoketone: 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methylbutanoyl)pyrazine**

[118025-76-6]

C₁₆H₂₆N₂O₂ 278.394Isol. from ants *Mesoponera castanea* and *Mesoponera castaneicolor*.**Diketone: 2,5-Dimethyl-3,6-bis(3-methylbutanoyl)pyrazine**

[118025-79-9]

C₁₆H₂₄N₂O₂ 276.378Isol. from *Mesoponera castanea* and *Mesoponera castaneicolor*. Mp 39.5-41°.**2',3'-Didehydro, 1'-ketone: 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methyl-2-butenoyl)pyrazine**

[118025-80-2]

C₁₆H₂₄N₂O₂ 276.378Isol. from *Mesoponera castanea* and *Mesoponera castaneicolor*.**2',3'-Didehydro, diketone: 2,5-Dimethyl-3-(3-methylbutanoyl)-6-(3-methyl-2-butenoyl)pyrazine**

[118025-81-3]

C₁₆H₂₂N₂O₂ 274.362Isol. from *Mesoponera castanea* and *Mesoponera castaneicolor*.**Monodeoxy: 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methylbutyl)pyrazine**

[118025-78-8]

C₁₆H₂₈N₂O 264.41Isol. from *Mesoponera castanea* and *Mesoponera castaneicolor*.**Dideoxy: 2,5-Dimethyl-3,6-bis(3-methylbutyl)pyrazine. 2,5-Diisopentyl-3,6-dimethylpyrazine**

[10132-44-2]

C₁₆H₂₈N₂ 248.411Isol. from ants *Mesoponera castanea* and *Mesoponera castaneicolor*.**Dideoxy, 1',2'-didehydro: 2,5-Dimethyl-3-(3-methyl-1-butenyl)-6-(3-methylbutyl)pyrazine**

[118025-75-5]

C₁₆H₂₆N₂ 246.395Isol. from *Mesoponera castanea* and*Mesoponera castaneicolor*. E- and Z-isomers reported. Tentative struct.**Dideoxy, 2',3'-didehydro: 2,5-Dimethyl-3-(3-methyl-2-butenyl)-6-(3-methylbutyl)pyrazine**


[118025-82-4]

C₁₆H₂₆N₂ 246.395Isol. from *Mesoponera castanea* and *Mesoponera castaneicolor*. Tentative struct.


[118025-84-6, 118025-77-7]

Fales, H.M. et al., *Tetrahedron*, 1988, **44**, 5045 (occur, ir, pmr, cmr, ms, synth, struct)**2,6-Bis(hydroxymethyl)-3,4-piperidinediol** B-185

2,3,6-Trideoxy-2,6-iminoheptitol

C₇H₁₃NO₄ 177.2**(2R,3R,4R,6S)-form***D*-manno-form. α -1-C-(Hydroxymethyl)-fagomineConstit. of the roots of *Stemona tuberosa*. Oil. [α]_D +28.9 (c, 0.2 in H₂O).Goujon, J.-Y. et al., *Bioorg. Med. Chem.*, 2005, **13**, 2313-2324 (synth, pmr, cmr)
Asano, N. et al., *J. Nat. Prod.*, 2005, **68**, 1238-1242 (isol, pmr, cmr)**2,6-Bis(hydroxymethyl)-3,4,5-piperidinetriol, 9CI** B-186

2,6-Dideoxy-2,6-iminoheptitol

C₇H₁₃NO₅ 193.199**(2R,3R,4R,5S,6R)-form***D*-glycero-*L*-gulo-form. α -Homonojirimycin

[119557-99-2]

Alkaloid from *Aglaonema treubii*, *Commelina communis*, *Adenophora* spp. *Hyacinthus orientalis* and *Omphalea diandra*. Inhibitor of several α -glucosidases. Antidiabetic agent. Mp 206-207°. [α]_D +88.2 (c, 0.54 in H₂O).*1''*-O- β -D-Glucopyranoside: **7-O- β -D-Glucopyranosyl- α -homonojirimycin**.MDL 25637. 2,6-Dideoxy-7-O- β -D-glucopyranosyl-2,6-imino-*D*-glycero-*L*-gulo-heptitol, 9CI

[104343-33-1]

[104419-80-9]

C₁₃H₂₅NO₁₀ 355.341Alkaloid from *Commelina communis*, *Aglaonema treubii*, *Hyacinthus orientalis* and *Lobelia sessilifolia*.Potent α -glucosidase inhibitor, competitive inhibitor for intestinalsucrase. Also inhibits maltase, trehalase, glucoamylase and α -amylase. Antidiabetic agent. Amorph. powder + 1H₂O. Mp 84-86° Mp 131-134° (as hydrochloride) Mp 216-219° synthetic. [α]_D²⁰ +20 (c, 0.5 in H₂O). [α]_D +24.7 (c, 0.7 in H₂O). [α]_D +27.5 (c, 1 in H₂O).**5-O- α -D-Galactopyranoside: 5-O- α -D-Galactopyranosyl- α -homonojirimycin**

[185826-28-2]

C₁₃H₂₅NO₁₀ 355.341Alkaloid from *Aglaonema treubii*. [α]_D +168.8 (c, 0.54 in H₂O).**1,3,4,5-Tetra-O-benzoyl:** [110205-71-5] Needles (Et₂O/petrol). Mp 76-78°.**(2R,3R,4R,5R,6S)-form** β -Homomannojirimycin

[154349-07-2]

Alkaloid from *Aglaonema treubii* and *Hyacinthus orientalis*. β -Mannosidase inhibitor. Hygroscopic solid. [α]_D +12.1 (c, 0.27 in H₂O).**(2R,3R,4S,5R,6S)-form***6-Epi- α -homomannojirimycin*

[127995-31-7]

[127995-28-2]

Potent inhibitor of α -fucosidase.Hygroscopic solid; cryst. (MeOH/CHCl₃) (as hydrochloride). Mp 203-205° (hydrochloride). [α]_D²⁰ +26.4 (c, 0.5 in H₂O).**(2R,3S,5S,6R)-form***D*-glycero-*L*-galacto-form. α -Homogalactostatin. **3,4-Diepi- α -homonojirimycin**

[169872-51-9]

Alkaloid from *Aglaonema treubii*. [α]_D +39.1 (c, 0.51 in H₂O). Existence of this isomer as a natural product is in doubt. C-4 is achiral.**(2R,3R,5R,6R)-form** α -Homomannojirimycin

[127995-29-3]

Alkaloid from *Hyacinthus orientalis* and *Aglaonema treubii*. Nonselective α -glucosidase inhibitor. Hygroscopic solid. [α]_D +4.4 (c, 0.55 in H₂O) (natural). [α]_D²⁰ +7.4 (c, 0.55 in H₂O) (synthetic). C-4 is achiral.**(2RS,3RS,4RS,5SR,6SR)-form** β -Homonojirimycin

[157544-15-5]

Alkaloid from *Aglaonema treubii* and *Hyacinthus orientalis*. No phys. props. reported. Meso-stereoisomer.**(2R,3R,4S,5S,6R)-form***D*-glycero-*L*-allo-form. α -Homoallojirimycin. **4-Epi- α -homonojirimycin**

[253785-13-6]

Alkaloid from *Aglaonema treubii*. [α]_D +39.1 (c, 0.51 in H₂O). Meso-stereoisomer. Stereochem. of natural product revised in 1999.Rhinehart, B.L. et al., *J. Pharmacol. Exp.**Ther.*, 1987, **24**, 915; C.A., **107**, 109172mLiu, P.S. et al., *J.O.C.*, 1987, **52**, 4717-4721 (α -*Homonojirimycin*, pmr, ms, derivs, synth)Kite, G.C. et al., *Tet. Lett.*, 1988, **29**, 6483-

6486 (isol, cmr)


Anzeveno, P.B. et al., *J.O.C.*, 1989, **54**, 2539-

2542 (MDL 25637, synth)

- Bruce, T. et al., *Tetrahedron*, 1992, **48**, 10191-10120 (α -Homomannojirimycin, 6-Epi- α -homomannojirimycin, synth, pmr, cmr, biochem)
- Holt, K.E. et al., *J.C.S. Perkin I*, 1994, 231-234 (α -Homomannojirimycin, β -Homomannojirimycin, synth, pmr, biochem)
- Asano, N. et al., *J. Nat. Prod.*, 1997, **60**, 98-101; 1998, **61**, 625-628 (isol, pmr, cmr)
- Asano, N. et al., *J. Med. Chem.*, 1998, **41**, 2565-2571 (synth, cryst struct, pmr, cmr, activity)
- Shilvlock, J.P. et al., *Tetrahedron: Asymmetry*, 1998, **9**, 3505-3516 (α -Homomannojirimycin, β -Homomannojirimycin, synth, pmr)
- Martin, O.R. et al., *Bioorg. Med. Chem. Lett.*, 1999, **9**, 3171-3174 (stereochem)
- Kim, H.S. et al., *Planta Med.*, 1999, **65**, 437-439 (isol)
- Watson, A.A. et al., *Phytochemistry*, 2000, **56**, 265-295 (rev)
- Morwenna, M.S.M. et al., *Eur. J. Org. Chem.*, 2005, 2159-2191 (rev, synth)

2,6-Bis(2-hydroxypentyl)piperidine **B-187**

α, α' -Dipropyl-2,6-piperidinediethanol, 9CI



(2*S*,2'*S*,2''*S*,6*S*)-form

$C_{15}H_{31}NO_2$ 257.415

(2*S*,2'*S*,2''*S*,6*S*)-form

Aspertine C
[442155-62-6]

Alkaloid from the aerial parts of *Andrachne aspera*. Gum. $[\alpha]_D^{20}$ -89 (c, 0.6 in $CHCl_3$).

(2*RS*,2'*RS*,2''*SR*,6*SR*)-form

N-Me: 2,6-Bis(2-hydroxypentyl)-1-methylpiperidine
[212006-67-2]

$C_{16}H_{33}NO_2$ 271.442

Alkaloid from *Siphocampylus verticillatus*. Fluffy needles ($CHCl_3$). Mp 31-32°. meso- Diastereoisomer.

(2*RS*,2'*SR*,2''*RS*,6*SR*)-form

Andrachamine

[107019-94-3]

Alkaloid from *Andrachne aspera*. meso-Diastereoisomer. Stereochem. revised in 1996.

N-Me: 8,10-Dipropyllobelidiol. N-Methylandrachamine

[212511-78-9]

$C_{16}H_{33}NO_2$ 271.442

Alkaloid from *Dialypetalum floribundum*. meso-Diastereoisomer.

[178062-05-0, 178062-06-1, 212006-69-4]

Miguel, O.G. et al., *Acta Cryst. C*, 1996, **52**, 1223-1225 (cryst struct, N-Me)

Mill, S. et al., *Can. J. Chem.*, 1996, **74**, 2434-2443 (*Andrachamine*)


Biaavatti, M.W. et al., *Phytochemistry*, 1998, **48**, 747-749 (N-Me, isol, ir, pmr, cmr, ms)

Krebs, H.C. et al., *Phytochemistry*, 1998, **48**, 911-913 (N-Me, isol, pmr, cmr, ms)

Ahmad, V.U. et al., *Turk. J. Chem.*, 2002, **26**, 245-250 (*Aspertine C*)

3,4-Bis(4-hydroxyphenyl)-1*H*-pyrrole-2,5-dicarboxylic acid **B-188**

[917608-79-8]



$C_{18}H_{13}NO_6$ 339.304

Prod. by the marine-derived *Halomonas* sp. DSM 17966. Amorph. solid. Mp 213° dec. λ_{max} 249 (log ϵ 4.27) (MeOH). λ_{max} 252 (log ϵ 4.27) (MeOH/HCl). λ_{max} 251 (log ϵ 4.28) (MeOH/NaOH).

4'-Deoxy-3-(4-Hydroxyphenyl)-4-phenyl-1*H*-pyrrole-2,5-dicarboxylic acid
[917608-78-7]


$C_{18}H_{13}NO_5$ 323.304

Prod. by *Halomonas* sp. RK377. Amorph. solid. Mp 213° dec. λ_{max} 245 (log ϵ 4.26) (MeOH). λ_{max} 243 (log ϵ 4.26) (MeOH/HCl). λ_{max} 249 (log ϵ 4.27) (MeOH/NaOH).

Wang, L. et al., *Appl. Microbiol. Biotechnol.*, 2006, **72**, 816-822 (isol, uv, pmr, cmr, ms, cryst struct)

Bisindigotin

B-189




$C_{32}H_{18}N_4O_2$ 490.52

Alkaloid from the leaves of *Isatis indigotica*. TCDD antagonist. Deep purple amorph. powder. λ_{max} 227 (log ϵ 4.94); 261 (log ϵ 4.78); 352 (log ϵ 4.62); 568 (log ϵ 4.38) (THF).

Wei, X.Y. et al., *J. Nat. Prod.*, 2005, **68**, 427-429 (isol, pmr, cmr)

3,3'-Bis(indolylmethyl)dimethylammonium **B-190**

N-(1*H*-Indol-3-ylmethyl)-N,N-dimethyl-1*H*-indole-3-methanaminium, 9CI



Quaternary alkaloid from flowers of *Arundo donax* (Poaceae).

Hydroxide: [35302-42-2]

Straw-coloured needles (EtOAc). Mp 179-180°.


Iodide:

Tan cryst. Mp 168-171°.

Ghosal, S. et al., *Phytochemistry*, 1971, **10**, 2852 (isol, uv, ms)

2,5-Bis(1*H*-indol-3-ylmethyl)pyrazine

B-191



$C_{22}H_{18}N_4$ 338.411

Prod. by *Cytophaga marinoflava* sp. strain AM13.1.


Shaaban, M. et al., *J. Nat. Prod.*, 2002, **65**, 1660-1663 (isol, pmr, cmr, ms)

Bisomahanine

B-192

Bispyrafoline D

[832726-58-6]



$C_{46}H_{48}N_2O_4$ 692.896


Alkaloid from the roots of *Glycosmis stenocarpa*. Pale cream powder. Mp 130-140°. $[\alpha]_D^{20}$ -13.1 (c, 0.25 in $CHCl_3$). λ_{max} 209 (log ϵ 4.1); 224 (log ϵ 3.7); 245 (log ϵ 5); 298 (log ϵ 4.9); 335 (log ϵ 4.4) ($CHCl_3$).

Cuong, N.M. et al., *Chem. Pharm. Bull.*, 2004, **52**, 1175-1178 (isol, cd, pmr, cmr, ms)

Bisjatrorrhizine

B-193

[35470-49-6]




$C_{40}H_{38}N_2O_8^{2+}$ 674.749

Alkaloid from the root of *Jateorhiza palmata* (Menispermaceae). Yellow-orange cryst. (as dichloride). Mp 270° dec. (dichloride).

Carvalhas, M.L. *et al.*, *J.C.S. Perkin 1*, 1972, 327 (*isol, uv, ir, pmr, ms, struct, synth*)

Bisline

15,20-Dihydro-12,15-dihydroxysenecionan-11,16-dione, 9CI
[30258-28-7]



C₁₈H₂₇NO₆ 353.414

C-15 config. revised in 2000. Alkaloid from *Senecio othonniformis* (*Senecio ruwenzoriensis*) and *Senecio petasis* (Asteraceae). Cryst. (Me₂CO/petrol). Mp 169°.

12-Ac: **Isoline**. Ruwenine

[30000-36-3]

C₂₀H₂₉NO₇ 395.452

Alkaloid from *Senecio othonniformis* (*Senecio ruwenzoriensis*) (Asteraceae). Cryst. (Me₂CO/petrol). Mp 173°. [α]_D²⁰ -4.8 (c, 1.78 in EtOH).

▶ Hepatotoxic.

15-Deoxy: 15,20-Dihydro-12-hydroxyse-necionan-11,16-dione. **Yamataimine**.

Duciformine

[67113-69-3]

C₁₈H₂₇NO₅ 337.415

Alkaloid from *Cacalia yatabei* and *Ligularia duciformis*. Mp 181-182°.

[α]_D²⁰ +63.6 (EtOH).

15-Deoxy, hydrobromide:

Needles (EtOH). Mp 249°.

15-Deoxy, 12-Ac: **O-Acetylyamataimine**

[217793-38-9]

C₂₀H₂₉NO₆ 379.452

Alkaloid from the roots of *Ligularia tsangchanensis*. Amorph. powder (Me₂CO). Mp 128-129°. [α]_D²⁰ +8.6 (c, 10.4 in CHCl₃).

15-Deoxy, 12-Ac, N-oxide: **O-Acetylyamataimine N-oxide**

[566901-96-0]

C₂₀H₂₉NO₇ 395.452

Alkaloid from the roots of *Ligularia tsangchanensis*. Yellow gum. [α]_D²⁰ +15 (c, 4.4 in CHCl₃).

Coucourakis, E.D. *et al.*, *J.C.S. (C)*, 1970, 2312-2315 (*isol*)

Coucourakis, E.D. *et al.*, *J.C.S. Perkin 1*, 1972, 2339-2343 (*Isoline*)

Gonzalez, A. *et al.*, *An. Quim.*, 1973, **69**, 1343-1345; *CA*, **80**, 96194s (*struct*)

Hikichi, M. *et al.*, *Tet. Lett.*, 1978, **19**, 767-770

(*Yamataimine, cryst struct, ir, pmr, cmr, ms*)

Benn, M. *et al.*, *Phytochemistry*, 1992, **31**, 3295-3296 (*Ruwenine*)

Niwa, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 3094-3099 (*Yamataimine, synth*)

Lin, G. *et al.*, *Rapid Commun. Mass Spectrom.*, 1998, **12**, 1445-1456


(*Duciformine*)

Susag, L. *et al.*, *Phytochemistry*, 2000, **54**, 933-935 (*struct, bibl*)

Tan, A. *et al.*, *Heterocycles*, 2003, **60**, 1195-1198 (*Acetylyamataimines*)

Bismahanine

[155519-87-2]




C₄₆H₄₈N₂O₄ 692.896

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Oil.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 2096 (*isol, uv, ir, pmr, ms, struct*)

1,1'-[[2,4-Bis(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidine, 9CI

2,4-Bis(5-methoxy-1,3-benzodioxol-6-yl)-1,3-cyclobutanedicarboxylic acid bispyrrolidide
[173866-77-8]



C₃₀H₃₄N₂O₈ 550.607


Alkaloid from leaves and stems of *Piper peepuloides*. Also obt. by irradiation (sunlight) of Peepuloidine, P-163. Mp 258°. λ_{max} 240; 304; 330 (MeOH).

3'-Methoxy: 1,1'-[[2-(6,7-Dimethoxy-1,3-benzodioxol-5-yl)-4-(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidine, 9CI. 4-(4,5-Dimethoxy-1,3-benzodioxol-6-yl)-2-(5-methoxy-1,3-benzodioxol-6-yl)-1,3-cyclobutanedicarboxylic acid bispyrrolidide
[173866-76-7]

C₃₁H₃₆N₂O₉ 580.633

From leaves and stems of *Piper peepuloides*. Mp 265°. λ_{max} 240; 305; 333 (MeOH).

Dhar, K.L. *et al.*, *Fitoterapia*, 1995, **66**, 390 (*isol, uv, ir, pmr, cmr, ms, struct*)

N-[1,3-Bis(methoxycarbonyl)propyl]-2,3-dihydro-2-(4-hydroxy-3-methoxyphenyl)-3-hydroxy-methyl-7-methoxy-5-benzofuranpropenamide

C₂₇H₃₁NO₁₀ 529.543

(2''ξ,7R,7'E,8S)-form [185311-15-3]

B-195

Constit. of *Boreava orientalis*. Pale yellow powder. Mp 82-84.5°. λ_{max} 288 (log ε 3.3); 320 (log ε 3.32) (MeOH).

(2''ξ,7R,7'Z,8S)-form [185311-14-2]


Constit. of *Boreava orientalis*. Pale yellow powder. Mp 95-99°. λ_{max} 287 (log ε 3.07); 318 (log ε 1.83) (MeOH).

Sakushima, A. *et al.*, *Phytochemistry*, 1996, **43**, 1349-1354 (*isol, uv, ir, cd, pmr, cmr, ms*)

2,4-Bis(2-methoxy-4,5-methylenedioxyphenyl)-1,3-cyclobutanedicarboxylic acid dipiperidide

1,1'-[[2,4-Bis(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidine, 9CI

[252361-69-6]



Relative Configuration

C₃₂H₃₈N₂O₈ 578.661

Alkaloid from *Piper peepuloides*. Mp 270°. λ_{max} 240; 304; 330 (MeOH).

3'-Methoxy: [252361-68-5]


C₃₃H₄₀N₂O₉ 608.687

Alkaloid from *Piper peepuloides*. Mp 280°. λ_{max} 240; 305; 353 (MeOH).

Sharma, R.L. *et al.*, *Fitoterapia*, 1999, **70**, 144-147

N,N-Bis(3-methyl-2-butenyl)guanidine

N,N-Diisopentenylguanidine. N,N-Diprenylguanidine. **Pterogynine**
[25387-60-4]



C₁₁H₂₁N₃ 195.307

Alkaloid from *Pterogyne nitens* and *Alchornea javanensis* (Fabaceae, Euphorbiaceae).

Hydrochloride: Mp 142-143°.


Picrate: Mp 157-158°.

Corral, R.A. *et al.*, *Experientia*, 1969, **25**, 1020 (*isol, pmr, struct*)

Corral, R.A. *et al.*, *Chem. Comm.*, 1970, 556 (*synth*)

3,5-Bis(3-methyl-2-butenyl)-1H-indole, 9CI

3,5-Diisopentenylindole. 3,5-Diprenylindole. 3,5-Bis(γ,γ-dimethylallyl)indole. **3,5-Hexalobine A**
[93822-62-9]



C₁₈H₂₃N 253.386

Alkaloid from the stem bark of *Hexalobus crispiflorus* (Annonaceae). Needles. Mp 78–79°. λ_{max} 226 (log ε 4.6); 286 (log ε 3.9); 297 (log ε 3.8) (MeOH).

2'R,3'-Epoxide: 3-(2,3-Epoxy-3-methylbutyl)-5-(3-methyl-2-butenyl)-1H-indole. **3,5-Hexalobine B** [165905-10-2]

C₁₈H₂₃NO 269.386

Alkaloid from the stem bark of *Hexalobus crispiflorus*. Oil. [α]_D²¹ -5 (c, 1.2 in CHCl₃). λ_{max} 226 (log ε 4.6); 286 (log ε 3.9); 297 (log ε 3.8) (MeOH).

2''S,3'''-Epoxide, 2',3'-dihydro, 2'R,3'-dihydroxy: 3-(2,3-Dihydroxy-3-methylbutyl)-5-(2,3-epoxy-3-methylbutyl)-1H-indole

[170384-65-3]

C₁₈H₂₅NO₃ 303.4

Alkaloid from the stem bark of *Hexalobus crispiflorus*. Oil. [α]_D²¹ +56 (c, 0.2 in CHCl₃). Poss. artifact. λ_{max} 227 (log ε 4.6); 286 (log ε 3.9); 295 (sh) (MeOH).

(2'R,3':2''S,3''')-Diepoxide: 3,5-Bis[(3,3-dimethyloxiranyl)methyl]-1H-indole, **9CI. 3,5-Hexalobine C**

[165942-81-4]

[93822-64-1]

C₁₈H₂₃NO₂ 285.385

Alkaloid from the stem bark of *Hexalobus crispiflorus* and *Isolona maitlandii* (Annonaceae). Oil. [α]_D²¹ +6 (c, 2.5 in CHCl₃). λ_{max} 226 (log ε 4.6); 271 (sh) (log ε 3.9); 285 (log ε 3.9); 296 (log ε 3.8) (MeOH).

1''-Oxo, 2'R,3'-epoxide: **3,5-Hexalobine D**

[165905-12-4]

C₁₈H₂₁NO₂ 283.369

Alkaloid from stem bark of *Hexalobus crispiflorus* and from *Isolona maitlandii*. Needles. Mp 130–132°. [α]_D²¹ +6 (c, 1.1 in CHCl₃). λ_{max} 216 (log ε 4.4); 267 (log ε 4.6); 310 (log ε 4.2) (MeOH).

1''-Oxo, Δ³-isomer, 2'-hydroxy: 3-(2-Hydroxy-3-methyl-3-butenyl)-5-(3-methyl-1-oxo-2-butenyl)-1H-indole

[170128-43-5]

C₁₈H₂₁NO₂ 283.369

Alkaloid from stem bark of *Isolona maitlandii*. Oil. [α]_D²¹ -6 (c, 0.13 in CHCl₃). λ_{max} 216 (log ε 4.24); 267 (log ε 4.41); 311 (log ε 4.05) (MeOH).

1''-Oxo, 2',3'-dihydro, 2'R,3'-dihydroxy: 3-(2,3-Dihydroxy-3-methylbutyl)-5-(3-methyl-1-oxo-2-butenyl)-1H-indole

[165604-39-7]

C₁₈H₂₃NO₃ 301.385

Alkaloid from stem bark of *Hexalobus crispiflorus* and from *Isolona maitlandii*. Oil. [α]_D²¹ +62 (c, 0.3 in CHCl₃). Poss. artifact. λ_{max} 217 (log ε 4.3); 269 (log ε 4.5); 313 (log ε 4.2) (MeOH).

1''-Oxo, 2',3'-dihydro, 2'R-hydroxy, 3'-methoxy: 3-(2-Hydroxy-3-methoxy-3-methylbutyl)-5-(3-methyl-1-oxo-2-butenyl)-1H-indole

[170128-42-4]

C₁₉H₂₅NO₃ 315.411

Alkaloid from the stem bark of *Isolona*

maitlandii. Oil. [α]_D²¹ +76 (c, 1.40 in CHCl₃). Poss. artifact. λ_{max} 217 (log ε 4.22); 268 (log ε 4.44); 311 (log ε 4.06) (MeOH).

Δ^{1''}-Isomer, 3''-hydroxy, 2'R,3'-epoxide:

3,5-Hexalobine E. 3-(2,3-Epoxy-3-methylbutyl)-5-(3-hydroxy-3-methyl-1-butenyl)-1H-indole

[165905-13-5]

C₁₈H₂₃NO₂ 285.385

Alkaloid from stem bark of *Hexalobus crispiflorus* and from *Isolona maitlandii*. Oil. [α]_D²¹ -6 (c, 0.25 in CHCl₃) (-3.3). λ_{max} 251 (log ε 4.7); 278 (sh); 306 (sh) (MeOH).

Δ^{1''}-Isomer, 3'',4''-didehydro, 2'R,3'-epoxide: 3-(2,3-Epoxy-3-methylbutyl)-5-(3-methyl-1,3-butadienyl)-1H-indole

[170128-44-6]

C₁₈H₂₁NO 267.37

Alkaloid from the stem bark of *Isolona maitlandii*. Oil. [α]_D²¹ -4 (c, 0.66 in Me₂CO). λ_{max} 266 (log ε 4.49); 273 (sh) (log ε 4.45); 305 (log ε 4.28) (MeOH).

Achenbach, H. et al., *Annalen*, 1995, 1327-1337 (isol, uv, pmr, cmr, ms, struct, derivs)

Achenbach, H. et al., *Phytochemistry*, 1995, **40**, 967-973 (derivs, isol, uv, ir, pmr, cmr)

3,6-Bis(3-methyl-2-butenyl)-1H-indole, 9CI

3,6-Bis(γ,γ-dimethylallyl)indole. 3,6-Diprenylindole. 3,6-Diisopentenylindole. **3,6-Hexalobine A**

[73618-54-9]

C₁₈H₂₃N 253.386

Alkaloid from the stem bark of *Uvaria elliptica*, *Hexalobus crispiflorus*, *Hexalobus monopetalus* and *Isolona maitlandii* (Annonaceae). Mp 36–37°. λ_{max} 229 (ε 39810); 283 (ε 7940); 294 (ε 6310) (MeOH) (Berdy).

2''S,3'''-Epoxide: 3-[(3,3-Dimethyloxiranyl)methyl]-6-(3-methyl-2-butenyl)-1H-indole, **9CI. 3,6-Hexalobine B**

[165905-11-3]

[93822-63-0]

C₁₈H₂₃NO 269.386

Alkaloid from the stem bark of *Hexalobus crispiflorus*, *Hexalobus monopetalus* and *Isolona maitlandii* (Annonaceae). Needles. Mp 72–73°. [α]_D²¹ +9 (c, 1.5 in CHCl₃). λ_{max} 227 (ε 39800); 283 (ε 6310); 293 (ε 5011) (MeOH) (Berdy).

2'R,3':2''R,3'''-Bisepoxide: **ent-3,6-Hexalobine C**

[172821-83-9]

C₁₈H₂₃NO₂ 285.385

Alkaloid from stem bark and leaves of *Isolona maitlandii* (Annonaceae). Oil. [α]_D²¹ -12 (c, 1.6 in CHCl₃).

2''S,3':2''S,3'''-Bisepoxide: 3,6-Bis[(3,3-dimethyloxiranyl)methyl]-1H-indole, **9CI. 3,6-Hexalobine C**

[166022-37-3]

[93822-65-2]

C₁₈H₂₃NO₂ 285.385

Alkaloid from the stem bark of *Hexalobus crispiflorus* and *Hexalobus monopetalus* (Annonaceae). Oil. [α]_D²¹ +16 (c, 0.4 in CHCl₃). λ_{max} 226 (ε 39800); 283 (ε 6310); 293 (ε 5000)

(MeOH) (Berdy).

2',3'-Dihydro, 2''S,3'''-dihydroxy: 3-(2,3-Dihydroxy-3-methylbutyl)-6-(3-methyl-2-butenyl)-1H-indole

[165604-38-6]

[93822-66-3]

C₁₈H₂₅NO₂ 287.401

Alkaloid from *Hexalobus crispiflorus*. Oil. [α]_D²¹ -35 (c, 0.75 in CHCl₃). λ_{max} 228 (log ε 4.7); 283 (log ε 3.9); 293 (log ε 3.9) (MeOH).

Δ^{1''}-Isomer(E-), 3''-hydroxy, 2'R,3'-epoxide: **(E)-3,6-Hexalobine E**

[172670-42-7]

C₁₈H₂₃NO₂ 285.385

Alkaloid from the bark and leaves of *Isolona maitlandii*. Cryst. Mp 129–131°. [α]_D²¹ +1 (c, 0.5 in CHCl₃). λ_{max} 251 (log ε 4.46); 295 (log ε 4.14) (MeOH).

Δ^{1''}-Isomer(Z-), 3''-hydroxy, 2'R,3'-epoxide: **(Z)-3,6-Hexalobine E**

[172670-43-8]

C₁₈H₂₃NO₂ 285.385

Alkaloid from leaves of *Isolona maitlandii*. Oil. [α]_D²¹ -2 (c, 1.1 in CHCl₃). λ_{max} 231 (log ε 4.39); 242 (log ε 4.38); 292 (log ε 3.99) (MeOH).

Achenbach, H. et al., *Tet. Lett.*, 1979, 2571 (uv, ir, pmr, cmr, struct)

Achenbach, H. et al., *Annalen*, 1995, 1327-1337 (isol, uv, pmr, cmr, ms, struct)

Achenbach, H. et al., *Phytochemistry*, 1995, **40**, 967-973 (bisepoxides)

3,7-Bis(3-methyl-2-butenyl)-1H-indole, 9CI

3,7-Diprenylindole. 3,7-Bis(γ,γ-dimethylallyl)indole. 3,7-Diisopentenylindole

C₁₈H₂₃N 253.386

Constit. of *Glycosmis trichanthera*. Oil. λ_{max} 225; 275 (sh); 282; 292 (sh) (MeOH).

4'-Hydroxy: 2-Methyl-4-[7-(3-methyl-2-butenyl)-1H-indol-3-yl]-2-buten-1-ol, **9CI. 3-(3-Hydroxymethyl-2-butenyl)-7-(3-methyl-2-butenyl)-1H-indole**

C₁₈H₂₃NO 269.386

Alkaloid from *Glycosmis montana*. Oil. Possesses (E)-config. λ_{max} 226; 275 (sh); 282; 292 (sh) (MeOH).


Vajrodaya, S. et al., *Phytochemistry*, 1998, **48**, 897-902 (isol, uv, ir, pmr, cmr, ms)

Wang, J. et al., *Phytochemistry*, 2005, **66**, 697-701 (*Glycosmis montana* constit)

2,5-Bis(2-methylpropyl)pyrazine, 9CI

2,5-Diisobutylpyrazine

[65032-04-4]

C₁₂H₂₀N₂ 192.303C₁₂H₂₀N₂ 192.303


Prod. by *Chondromyces crocatus* and *Paenibacillus polymyxa*.

Beck, H.C. et al., *FEMS Microbiol. Lett.*, 2003, **220**, 67-73 (isol)

Schulz, S. et al., *Tetrahedron*, 2004, **60**, 3863-3872 (isol)

3,6-Bis(1-methylpropyl)-2(1H)-pyrazinone, 9CI B-204

3,6-Bis(1-methylpropyl)-2-pyrazinol. 3-Hydroxy-2,5-bis(1-methylpropyl)pyrazine. 3,6-Di-sec-butyl-2-hydroxypyrazine [17005-09-3]



C₁₂H₂₀N₂O 208.303

Stereochem. of isolates undefined. Metab. from *Aspergillus flavus* and *Aspergillus sojae*. Mp 129°.

1''-Hydroxy: 6-(1-Hydroxy-1-methylpropyl)-3-(1-methylpropyl)-2(1H)-pyrazinone

C₁₂H₂₀N₂O₂ 224.302

Metab. of *Aspergillus sojae*. Mp 117°. λ_{max} 233 ; 326 (MeOH) (Berdy).

NH-form

N¹-Hydroxy: 1-Hydroxy-3,6-bis(1-methylpropyl)-2(1H)-pyrazinone

C₁₂H₂₀N₂O₂ 224.302

Metab. from *Aspergillus flavus* and *Aspergillus sojae*. Oil. λ_{max} 233 ; 325 (MeOH) (Berdy).

▶ **LD₅₀** (mus, ipr) 200 mg/kg.

1'',N¹-Dihydroxy: 1-Hydroxy-6-(1-hydroxy-1-methylpropyl)-3-(1-methylpropyl)-2(1H)-pyrazinone


C₁₂H₂₀N₂O₃ 240.302

Metab. from *Aspergillus sojae*. Mp 120-121°.

Yamamoto, Y. et al., *Alkaloids (Academic Press)*, 1986, **29**, 185 (rev, bibl)

Bismorphine A

[389117-13-9]



C₃₄H₃₆N₂O₆ 568.668


Dimer of Morphine, M-704. Alkaloid from wounded capsules of *Papaver somniferum*.

Morimoto, S. et al., *J. Biol. Chem.*, 2001, **276**, 38179-38184 (isol, pmr, cmr)

Morimoto, S. et al., *J. Nat. Prod.*, 2003, **66**, 987-989 (isol)

Bismorphine B

[575487-91-1]

**B-205**


C₃₄H₃₆N₂O₆ 568.668

Dimer of Morphine, M-704. Alkaloid from wounded capsules of *Papaver somniferum*. Off-white powder + 1H₂O. [α]_D²⁸ -68.3 (c, 0.15 in MeOH). λ_{max} 284 (log ε 4.12) (MeOH).

Morimoto, S. et al., *J. Nat. Prod.*, 2003, **66**, 987-989 (isol, pmr, cmr)

Bismurrayafoline A**B-207**

1-Methoxy-9-[(1-methoxy-9H-carbazol-3-yl)methyl]-3-methyl-9H-carbazole, 9CI [89368-89-8]



C₂₈H₂₄N₂O₂ 420.51

Alkaloid from *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Needles (Et₂O). Mp 176-177°.

1'-Aldehyde: Chrestifoline D

[142750-14-9]

C₂₈H₂₂N₂O₃ 434.493

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

1'-Hydroxy: Bismurrayafolinol

[107903-16-2]

C₂₈H₂₄N₂O₃ 436.509

Alkaloid from stem bark of *Murraya euchrestifolia* (Rutaceae). Oil.


Furukawa, H. et al., *Chem. Pharm. Bull.*, 1983, **31**, 4202 (uv, pmr, ms, struct)

Ito, C. et al., *Chem. Pharm. Bull.*, 1987, **35**, 450; 1992, **40**, 230 (Bismurrayafolinol, Chrestifoline D)

Itoigawa, M. et al., *J. Nat. Prod.*, 2000, **63**, 893-897 (activity)

Bismurrayafoline B**B-208**

7,7'-Dimethoxy-3,3'-dimethyl-8,8'-bis(3-methyl-2-butenyl)[2,2'-bi-9H-carbazole]-1,1'-diol, 9CI [89368-92-3]




C₃₈H₄₀N₂O₄ 588.745

Alkaloid from *Murraya euchrestifolia* (Rutaceae). Needles (Me₂CO). Mp 260° dec.

Furukawa, H. et al., *Chem. Pharm. Bull.*, 1983, **31**, 4202 (uv, ir, pmr, cmr, ms, struct)

Bismurrayafoline C**B-209**

[139726-48-0]



C₄₆H₅₂N₂O₄ 696.928

Alkaloid from the stem bark of *Murraya euchrestifolia*. Pale yellow oil. λ_{max} 226 ; 235 ; 260 (sh) ; 285 (sh) ; 311 ; 335 (sh) (MeOH).

7,7'-Di-Me ether: Bismurrayafoline D

[139726-49-1]


C₄₈H₅₆N₂O₄ 724.981

Alkaloid from the stem bark of *Murraya euchrestifolia*. Prisms (Me₂CO). Mp 198-200°. λ_{max} 224 ; 238 ; 260 (sh) ; 285 (sh) ; 310 ; 334 (sh) (MeOH).

Ito, C. et al., *Chem. Pharm. Bull.*, 1991, **39**, 2525-2528

Bismurrayafoline E**B-210**

[252350-80-4]



C₄₈H₅₆N₂O₄ 724.981


Alkaloid from the leaves of *Murraya koenigii* (curryleaf tree). Gum.

Nutan, M.T.H. et al., *Fitoterapia*, 1999, **70**, 130-133

6,8'-Bismurrayamine A**B-211**


Bis(7-hydroxygirinimbine B)

[133632-48-1]




C₃₆H₃₂N₂O₄ 556.66Alkaloid from leaves of *Murraya euchrestifolia* (Rutaceae). Pale yellow oil. Poss. artifact.Wu, T.-S. *et al.*, *Phytochemistry*, 1991, **30**, 1048; 1052**8,8'-Bismurrayamine A**


B-212

Bis(7-hydroxygirinimbine A)
[133655-90-0]C₃₆H₃₂N₂O₄ 556.66Alkaloid from leaves of *Murraya euchrestifolia* and from *Murraya exotica* (Rutaceae). Pale yellow oil. Poss. artifact.*Di-Meether: Bis(7-methoxygirinimbine A)*
[133632-47-0]C₃₈H₃₆N₂O₄ 584.713Alkaloid from *Murraya exotica* (Rutaceae).Wu, T.S. *et al.*, *Phytochemistry*, 1991, **30**, 1048; 1052Desoky, E.K. *et al.*, *Bull. Fac. Pharm. (Cairo Univ.)*, 1992, **30**, 231; *CA*, **119**, 177557p
(*Bis(7-methoxygirinimbine A)*)**Bismurrayquinone A**

B-213

3,3'-Dimethyl-[2,2'-bi-1H-carbazole]-1,1',4,4'-(9H,9'H)-tetrone, 9CI
[155519-86-1]C₂₆H₁₆N₂O₄ 420.423Alkaloid from roots of *Murraya koenigii* (curryleaf tree). Orange powder.Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 2096 (*isol, uv, ir, pmr, ms, struct*)Bringmann, G. *et al.*, *Tetrahedron*, 1995, **51**, 9353 (*synth*)Murphy, W.S. *et al.*, *J.C.S. Perkin 1*, 1998, 4115-4119 (*synth*)**Bisnorbiotin**

B-214

Hexahydro-2-oxo-1H-thieno[3,4-d]imidazole-4-propanoic acid, 9CI

(3S,4S,5R)-form

C₈H₁₂N₂O₃S 216.26**(3S,4S,5R)-form***d-Bisnorbiotin*

[16968-98-2]

Minor metab. of Biotin, B-139. Cryst. (EtOAc) (as Me ester). Mp 165-166° (Me ester). $[\alpha]_D^{25} +55.7$ (c, 0.96 in DMSO) (Me ester).**(3S,4S,5S)-form***d-Allobisnorbiotin*

[51606-79-2]


Prod. by *Pseudomonas* sp. grown on Biotin, B-139.

[39871-27-7, 57671-76-8, 39871-28-8]

Im, W.B. *et al.*, *J. Biol. Chem.*, 1973, **248**, 7798-7805 (*isol*)Confalone, P.N. *et al.*, *J.A.C.S.*, 1977, **99**, 7020-7026 (*synth*)Im, W.B. *et al.*, *Methods Enzymol.*, 1979, **62**, 385-390 (*isol*)**Bisparthenolidine**

B-215

[112078-76-9]

C₃₀H₄₃NO₆ 513.673Constit. of *Paramichelia baillonii* (preferred genus name *Michelia*) and *Michelia rajaniana* (Magnoliaceae). Shows antitumour props. Cryst. (CHCl₃). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 100-103°. $[\alpha]_D^{20} -112$ (CHCl₃).Ruangrunsi, N. *et al.*, *J. Nat. Prod.*, 1987, **50**, 891; 1988, **51**, 1230 (*isol, ir, ms, struct, pmr, cmr*)**N,N'-Bis(2-phenylethyl)urea, 9CI**

B-216

N,N'-Diphenethylurea. AM 2498. Metabolite AM 2498


[5467-84-5]

PhCH₂CH₂NHCONHCH₂CH₂PhC₁₇H₂₀N₂O 268.358Metab. from the marine ascidian *Didemnum ternatanum*. Also *isol.* from a *Streptomyces* sp. strain No. AM-2498.Weak depressant lacking acute toxicity. Needles (MeOH). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 138-141° (135°). λ_{max} 248 (ε 308); 251 (ε 411); 253 (ε 446); 256 (ε 535); 259 (ε 549); 262 (ε 543); 265 (ε 419); 269 (ε 351) (MeOH) (Derep).


► YS2768000

Iwai, Y. *et al.*, *J. Antibiot.*, 1978, **31**, 375 (*isol, uv, ir, pmr, cmr, struct*)Ireland, C.M. *et al.*, *J. Nat. Prod.*, 1981, **44**, 360 (*isol, ir, pmr, cmr, ms, synth*)**Bispsammaplin A**

B-217


C₄₄H₄₆Br₄N₈O₁₂S₄ 1326.774*Isol.* from an association of the sponges *Jaspis wondoensis* and *Poecillastra wondoensis*. Yellow oil. $[\alpha]_D^{23} -3$ (c, 0.21 in MeOH).Park, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1495-1498 (*isol, pmr, cmr*)**Biscodehydrocyclopiazonic acid**

B-218

Dehydrodisecocyclopiazonic acid. β-Cyclopiazonic acid
[31008-70-5]C₂₀H₂₂N₂O₃ 338.405*Isol.* from *Penicillium cyclopium* and *Aspergillus versicolor*. Cryst. (MeOH). Mp 168-169°. $[\alpha]_D^{20} -186$ (c, 0.42 in Py). λ_{max} 225 (ε 39800); 253 (ε 16600); 275 (sh) (ε 19100); 284 (ε 20400); 292 (sh) (ε 17400) (MeOH) (Derep). λ_{max} 226 (ε 38500); 276 (ε 11750) (MeOH) (Berdy).Holzapfel, C.W. *et al.*, *Tetrahedron*, 1970, **26**, 5239 (*isol, uv, ir, pmr, ms*)Ohmomo, S. *et al.*, *Nippon Nogei Kagaku Kaishi*, 1973, **47**, 57 (*isol, pmr*)Nolte, M.J. *et al.*, *J.C.S. Perkin 1*, 1980, 1057 (*cmr, tautom*)Holzapfel, C.W. *et al.*, *Aust. J. Chem.*, 1992, **45**, 99 (*synth*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 504**Bistelletadine A**

B-219

[233766-63-7]

C₃₀H₅₂N₁₂O₂ 612.82*Isol.* as a salt tetraprotonated at the guanidino residues. Counterion not specified. Alkaloid from the marine sponge *Stelletta* sp. $[\alpha]_D^{21} +2$ (c, 0.72 in MeOH) (salt). λ_{max} 255 (ε 13000) (MeOH).Tsukamoto, S. *et al.*, *J.O.C.*, 1999, **64**, 3794-3795 (*isol, uv, pmr, cmr*)

Bistelletadine B **B-220**

[233766-64-8]

As Bistelletadine A, B-219 with

R = -CH₂CH=C(CH₃)₂C₃₅H₆₀N₁₂O₂ 680.938

Isol. as a salt tetraprotonated at the guanidino residues. Counterion not specified. Alkaloid from the marine sponge *Stelletta* sp. [α]_D²¹ +3.9 (c, 0.1 in MeOH) (salt). λ_{max} 255 (ε 11000) (MeOH).


Tsukamoto, S. *et al.*, *J.O.C.*, 1999, **64**, 3794-3795 (*isol, uv, pmr, cmr*)

2,5-Bis(1,2,3,4-tetrahydro-2-butyl)pyrazine **B-221**

1,1'-(2,5-Pyrazinediyl)bis-1,2,3,4-butane-tetrol, 9CI

[68538-87-4]

[220145-44-8, 220145-46-0, 220145-43-7, 220145-45-9, 220145-47-1]



(1'R,1''R,2'R,2''R,3'R,3''R)-form

C₁₂H₂₀N₂O₈ 320.299**(1'R,1''R,2'R,2''R,3'R,3''R)-form***D*-lyxo-form

[13121-64-7]

Beige powder. Mp 109°.

(1'R,1''R,2'S,2''S,3'R,3''R)-form*D*-arabino-form. *D*-Fructosazine

[13185-73-4]

Formed by the action of ammonia on glucose. Exhibits DNA strand cleavage activity. Needles (EtOH). Mp 237°.

[α]_D²⁰ -84.1 (c, 1 in H₂O).*Octa-Ac*: [13051-88-2]Mp 174°. [α]_D²⁰ -7.2 (c, 1 in CHCl₃).**(1'R,1''R,2'S,2''S,3'S,3''S)-form***L*-xylo-form. *L*-Sorbosazine

[41093-14-5]

Needles (EtOH). Mp 175-177°. [α]_D²⁰ -27 (c, 0.1 in H₂O).

(1''ξ,1''ξ,2''ξ,2''ξ,3''ξ,3''ξ)-form

2',2''-Bis-O-(2-acetamido-2-deoxy-β-D-glucopyranoside): *Vibrio alginolyticus* Pyrazine. VAPY

[165956-47-8]

C₂₈H₄₆N₄O₁₈ 726.687

Prod. by *Vibrio alginolyticus* TK-24 with chitin. Platelet aggregation inhibitor. Powder. Sol. MeOH, H₂O. λ_{max} 275 (H₂O).

[68510-02-1]

Fujii, S. *et al.*, *J.O.C.*, 1966, **31**, 2239-2241 (*synth, fructosazine*)

Tsuchida, H. *et al.*, *Agric. Biol. Chem.*, 1973, **37**, 2571-2578 (*Fructosazine, synth, uv, ir*)

Teglia, M.C. *et al.*, *Carbohydr. Res.*, 1973, **26**, 377-384 (*synth, Sorbosazine*)

Tsuchida, H. *et al.*, *Dev. Food Sci.*, 1986, **13**, 85-94 (*occur*)

Tsuchida, H. *et al.*, *CA*, 1990, **113**, 229877r (*occur*)

Sumoto, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1991, **39**, 792-794 (*Frucosazine, synth, ir, pmr, cmr*)


Yanai, T. *et al.*, *J. Ferment. Bioeng.*, 1995, **80**, 406-407 (*VAPY*)

Japan. Pat., 1995, 95 118 286; *CA*, **124**, 7180w (*VAPY*)

Pat. Coop. Treaty (WIPO), 1999, 9 903 842; *CA*, **130**, 139579s (*synth, lyxo-form*)

Bistratamide I

[501909-64-4]



X = O

C₂₅H₃₆N₆O₆S 548.662


Isol. from the ascidian *Lissoclinum bistratum*. Solid. [α]_D²⁰ -122 (c, 0.5 in MeOH). λ_{max} 243 (sh) (MeOH).

Perez, L.J. *et al.*, *J. Nat. Prod.*, 2003, **66**, 247-250 (*isol, pmr, cmr, ms*)

You, S.-L. *et al.*, *Tetrahedron*, 2005, **61**, 241-249 (*synth*)

Bistratamide A

[120853-13-6]

C₂₇H₃₄N₆O₄S₂ 570.735

Cyclic hexapeptide from the ascidian *Lissoclinum bistratum* and from *Prochloron* sp. Cytotoxic agent. λ_{max} 232 (ε 13000) (CH₂Cl₂) (Derep).

11,12-Didehydro: Bistratamide B

[120881-21-2]


C₂₇H₃₂N₆O₄S₂ 568.72

From *Lissoclinum bistratum* and from *Prochloron* sp. Cytotoxic, but less so than Bistratamide A. λ_{max} 232 (ε 13000) (CH₂Cl₂) (Derep).

Degnan, B.M. *et al.*, *J. Med. Chem.*, 1989, **32**, 1354-1359 (*isol, pmr, cmr, struct*)

Bistratamide C

[145400-56-2]

**B-224**C₂₂H₂₆N₆O₄S₂ 502.617


Cyclic hexapeptide from the ascidian *Lissoclinum bistratum*. Sol. MeOH, CHCl₃. [α]_D²⁵ -65 (c, 0.42 in CHCl₃). λ_{max} 233 (ε 17000) (CH₂Cl₂) (Derep).

Foster, M.P. *et al.*, *J.O.C.*, 1992, **57**, 6671-6675 (*isol, uv, ir, pmr, cmr, struct*)

Aguilar, E. *et al.*, *Tet. Lett.*, 1994, **35**, 2477 (*synth*)

Bistratamide D

[145427-87-8]

**B-225**C₂₅H₃₄N₆O₅S 530.647

Cyclic hexapeptide from the ascidian *Lissoclinum bistratum*. Shows sedative and depressant props. Sol. MeOH, CHCl₃. [α]_D²⁵ -31 (c, 0.33 in CHCl₃). λ_{max} 233 (ε 17000) (CH₂Cl₂) (Derep).

19α,20-Dihydro: Bistratamide F

[501909-61-1]

C₂₅H₃₆N₆O₅S 532.663Isol. from *Lissoclinum bistratum*.

Cream solid. [α]_D²⁰ +23.2 (c, 1 in MeOH). Incorrect MF in ref. λ_{max} 230 (sh) (MeOH).

2,3-Didehydro: Bistratamide G

[501909-62-2]

C₂₅H₃₂N₆O₅S 528.631

Isol. from *Lissoclinum bistratum*. Solid. [α]_D²⁰ -73.8 (c, 1 in MeOH). λ_{max} 230 (sh) (MeOH).

Foster, M.P. *et al.*, *J.O.C.*, 1992, **57**, 6671-6675 (*isol, uv, ir, pmr, cmr, struct*)

Downing, S.V. *et al.*, *J.O.C.*, 1999, **64**, 826-831 (*synth, pmr, cmr*)


Perez, L.J. *et al.*, *J. Nat. Prod.*, 2003, **66**, 247-250 (*Bistratamides F,G*)

Shin, C. *et al.*, *Chem. Lett.*, 2004, **33**, 664-665 (*synth*)

You, S.-L. *et al.*, *Tetrahedron*, 2005, **61**, 241-249 (*synth*)

Bistratamide E

[501909-60-0]

B-226C₂₅H₃₄N₆O₄S₂ 546.713

Isol. from the ascidian *Lissoclinum bistratum*. Glass. [α]_D²⁰ -31 (c, 1 in MeOH). λ_{max} 237 (log ε 3030) (MeOH).

2,3-Didehydro: Bistratamide H

[501909-63-3]


C₂₅H₃₂N₆O₄S₂ 544.698Isol. from *Lissoclinium bistratum*. Solid. [α]_D -92.9 (c, 1 in MeOH). λ_{max} 237 (sh) (MeOH).Perez, L.J. *et al.*, *J. Nat. Prod.*, 2003, **66**, 247-250 (isol, pmr, cmr, ms)You, S.-L. *et al.*, *Tetrahedron*, 2005, **61**, 241-249 (synth)Nakamura, Y. *et al.*, *Tet. Lett.*, 2006, **47**, 239-243 (Bistratamide H, synth)Bertram, A. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1541-1553 (Bistratamide H, synth)**Bistratamide J****B-227**

[501909-65-5]


As Bistratamide I, B-222 with X = S

C₂₅H₃₆N₆O₅S₂ 564.729Isol. from *Lissoclinium bistratum*. Solid. [α]_D -25 (c, 0.5 in MeOH). λ_{max} 248 (sh) (MeOH).Perez, L.J. *et al.*, *J. Nat. Prod.*, 2003, **66**, 247-250 (isol, pmr, cmr, ms)You, S.-L. *et al.*, *Chem. Eur. J.*, 2004, **10**, 71-74 (synth)**Bitlisine****B-228**


[105223-81-2]

C₂₀H₁₉NO₆ 369.373Alkaloid from *Corydalis rutifolia* (Papaveraceae).Şener, B. *et al.*, *CA*, 1986, **105**, 206252d**Bizelesin, INN, USAN****B-229**

6,6'-[Carbonylbis(imino-1H-indole-5,2-diylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-methylbenzo[1,2-b:4,3-b']dipyrrol-4-ol], 9Cl. U 77779. NSC 615291. Antibiotic U 77779 [129655-21-6]

C₄₃H₃₆Cl₂N₈O₅ 815.713Prod. by *Streptomyces zeleneis*. DNA alkylating agent. Antineoplastic agent. Solid. Analogue of Rachelmycin, R-4. [129655-29-4]Eur. Pat., 1990, ((Upjohn))359 454; *CA*, **113**, 152387w (synth, pharmacol)Lee, C.S. *et al.*, *Cancer Res.*, 1991, **51**, 6586-6591 (pharmacol)Lee, C.S. *et al.*, *Biochemistry*, 1993, **32**, 2592-2600 (pharmacol)Sun, D. *et al.*, *Chem. Res. Toxicol.*, 1993, **6**, 889-894 (pharmacol)Butryn, R.K. *et al.*, *Cancer Chemother. Pharmacol.*, 1994, **34**, 44-50; 317-322 (pharmacol, hplc)Borman, S. *et al.*, *Chem. Eng. News*, November 17, 1997, 37 (rev. activity)Schwartz, G.H. *et al.*, *Ann. Oncol.*, 2003, **14**, 775-782 (clin trial)**Blasticidin A****B-230**

[100513-53-9]

C₅₈H₁₀₇NO₂₃ 1186.476


Prod. by *Streptomyces griseochromogenes*. Shows broad antifungal and antibacterial spectrum. Exhibits aflatoxin production inhibitory activity. Pale yellow powder. Sol. MeOH, butanol, Py, DMSO, bases, AcOH, DMF; fairly sol. EtOH, H₂O; poorly sol. C₆H₆, Me₂CO, Et₂O, CCl₄, EtOAc, hexane, CHCl₃. Mp 197-201°. [α]_D +10.8 (c, 1 in DMSO). λ_{max} 235; 308 (MeOH) (Berdy). λ_{max} 246 (€ 11500); 299 (€ 5800) (MeOH aq.) (Berdy). λ_{max} 237 (€ 9000); 314 (€ 6400) (MeOH-HCl) (Berdy).

▶ LD₅₀ (mus, ipr) 26 mg/kg.

Ca chelate: Mp 165.5-166.5°.

Fukunaga, K. *et al.*, *Bull. Agric. Chem. Soc. Jpn.*, 1955, **19**, 181Kono, Y. *et al.*, *J. Antibiot., Ser. A*, 1968, **21**, 433Sakuda, S. *et al.*, *J. Antibiot.*, 2000, **53**, 1265-1271 (pmr, cmr, struct)Sakuda, S. *et al.*, *Tet. Lett.*, 2007, **48**, 2527-2531 (abs config)**Blastmycetin A****B-231**

[107783-16-4]


C₃₅H₄₆N₆O₄ 614.786

Isol. from *Streptoverticillium blastmyceticus*. Amorph. Sol. MeOH, CHCl₃. [α]_D²⁶ -220 (c, 0.46 in EtOH). Related to Teleocidin B₁, T-63 and Indolactam V, I-78. λ_{max} 234 (€ 17200); 297 (sh) (€ 5900); 310 (€ 7100) (EtOH) (Derep). λ_{max} 231 (€ 49300); 259 (€ 15900); 303 (€ 16700) (EtOH) (Berdy).

▶ Tumour promotor.

Irie, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 285-287; 1988, **52**, 641-648 (isol, struct)**Blastmycetin B****B-232**

[107745-57-3]

C₁₇H₂₃N₃O₄ 333.386

Isol. from *Streptoverticillium blastmyceticus*. Tumour promotor. Amorph. Sol. MeOH, CHCl₃. [α]_D²⁰ -323 (c, 0.48 in MeOH). λ_{max} 234 (€ 17200); 297 (sh) (€ 5900); 310 (€ 7100) (EtOH) (Derep). λ_{max} 243 (€ 14300); 325 (€ 3700) (EtOH) (Berdy).

3-Deoxy: 3-Deoxyblastmycetin B

[111004-10-5]

C₁₇H₂₃N₃O₃ 317.387

From *Streptoverticillium blastmyceticus*. Tumour promotor. [α]_D²⁴ -306.5 (c, 0.21 in MeOH).

3-Epimer: Blastmycetin C

[107797-58-0]

C₁₇H₂₃N₃O₄ 333.386

From *Streptoverticillium blastmyceticus*. Tumour promotor. Amorph. Sol. MeOH, CHCl₃. [α]_D¹⁹ +166.5 (c, 0.14 in MeOH). λ_{max} 234 (€ 17200); 297 (sh) (€ 5900); 310 (€ 7100) (EtOH) (Derep). λ_{max} 247 (€ 6300); 285 (€ 1500) (EtOH) (Berdy).

3-Epimer, 3-deoxy: 3-Deoxyblastmycetin C


[98770-63-9]

C₁₇H₂₃N₃O₃ 317.387

From *Streptoverticillium blastmyceticus*. Tumour promotor. [α]_D¹⁹ +89.6 (c, 0.17 in MeOH).

Irie, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 285-287; 1988, **52**, 641-648 (isol, struct)**Blastmycetin D****B-233**


[110064-64-7]

C₂₉H₄₅N₃O₃ 483.693

Isol. from *Streptoverticillium blastmyceticum*. Poss. precursor of the Teleocidins. Amorph. [α]_D²³ -51 (c, 0.27 in MeOH). λ_{max} 234 (€ 17200); 297 (sh) (€ 5900); 310 (€ 7100) (EtOH) (Derep). λ_{max} 229 (€ 26200); 287 (€ 3600); 298 (€ 3400) (MeOH) (Berdy).

Irie, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 1733 (*isol, struct*)
Irie, K. *et al.*, *Tetrahedron*, 1990, **46**, 2773 (*isol*)


Blastmycetin E **B-234**
[121530-46-9]



Absolute Configuration


$C_{28}H_{41}N_5O_2$ 451.651
Isol. from *Streptovorticillum blastmyceticum*. Tumour promoter. Powder. $[\alpha]_D^{22}$ -64.7 (c, 0.63 in EtOH). λ_{max} 234 (ε 17200); 297 (sh) (ε 5900); 310 (ε 7100) (EtOH) (Derep).
Irie, K. *et al.*, *Tet. Lett.*, 1989, **30**, 2113 (*isol, struct*)
Irie, K. *et al.*, *Tetrahedron*, 1990, **46**, 2773 (*isol*)

Blastmycetin F **B-235**
[156312-08-2]




$C_{22}H_{28}N_4O_3$ 396.488
Metab. of *Streptovorticillum blastmyceticum*. Tumour promoter. Red amorph. powder. $[\alpha]_D^{26}$ -147 (c, 0.21 in MeOH). λ_{max} 234 (ε 17200); 297 (sh) (ε 5900); 310 (ε 7100) (EtOH) (Derep). λ_{max} 223 (ε 27800); 273 (ε 12400); 344 (ε 11200); 432 (ε 4800) (MeOH) (Berdy).
Irie, K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 363 (*isol, pmr, cmr, ms, struct*)

Bleomycin A₆ **B-236**
 N^1 -[3-[4-[3-(Aminopropyl)amino]butyl]amino]propyl]bleomycinamide, 9CI. *Zhengguangmycin A₆* [37293-17-7]




$C_{60}H_{96}N_{20}O_{21}S_2$ 1497.672
Glycopeptide antibiotic. Isol. from *Streptomyces verticillus*. Sol. MeOH, H_2O ; fairly sol. EtOH; poorly sol. Me_2CO , hexane. λ_{max} 244 ; 295 (H_2O) (Berdy).
Fujii, A. *et al.*, *J. Antibiot.*, 1973, **26**, 398 (*isol*)

Bluensidine **B-237**
1-O-Carbamoyl-3-deoxy-3-guanidinoscylo-inositol [4939-64-4]




$C_8H_{16}N_4O_6$ 264.238
Mp 194-196° dec. (as hydrochloride). $[\alpha]_D$ +1 (H_2O).
N,N,N,O,O,O-Hexa-Ac: Mp 250-251° dec. $[\alpha]_D$ +5 (c, 0.9 in $CHCl_3$).
Bannister, B. *et al.*, *J.A.C.S.*, 1963, **85**, 119

Blumeanine **B-238**
[214702-48-4]



$C_{38}H_{48}N_4O_2^{\oplus}$ 592.823
Alkaloid from *Ophiorrhiza blumeana*.
O,O-Di-Ac: [214702-53-1]
 $C_{42}H_{52}N_4O_4^{\oplus}$ 676.897
Plates (MeOH/EtOAc). $[\alpha]_D^{32}$ -155 (c, 0.003 in MeOH). Dec. without melting at 250°. Error in the MF given in the lit. The counterion is not clear from the reference. λ_{max} 220 (ε 25400); 291 (ε 7600) (MeOH).
Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1998, 2537-2540 (*isol, uv, ir, cd, pmr, cmr, ms*)

Bocconoline **B-239**
12,13-Dihydro-1,2-dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridine-13-methanol, 9CI. 8-Hydroxymethyl-dihydrochelerythrine [32906-88-0]



$C_{22}H_{21}NO_5$ 379.412
Probably an artifact. Alkaloid from *Macleaya cordata*, *Chelidonium japonicum*, *Glaucium flavum*, and from an unnamed *Fagara* sp. (Papaveraceae, Rutaceae). Pillars (MeOH). Mp 232-233° (215-218°, 221-222°).

O⁹-De-Me: 7,8-Dihydro-8-hydroxymethylfagaridine
 $C_{21}H_{19}NO_5$ 365.385
Alkaloid from stems and twigs of *Zanthoxylum microcarpum* (Rutaceae). Solid. Poss. artifact.


Deoxy: 8-Methyldihydrochelerythrine [159465-79-9]
 $C_{22}H_{21}NO_4$ 363.412
Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Needles (MeOH). Mp 203-205°.

Tani, C. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 755; *CA*, **58**, 5747f (*isol*)
Ishii, H. *et al.*, *Tet. Lett.*, 1971, 2429 (*uv, ir, pmr, ms, struct*)
Novák, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 3352 (*isol, uv*)
Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 166 (*uv, ir, pmr, ms, struct, synth*)
Itokawa, H. *et al.*, *Phytochemistry*, 1978, **17**, 839 (*isol, uv, ir, pmr, ms*)
Castedo, L. *et al.*, *Heterocycles*, 1981, **16**, 533 (*occur*)
Boulware, R.J. *et al.*, *J. Nat. Prod.*, 1981, **44**, 200 (7,8-Dihydro-8-hydroxymethylfagaridine)
Chen, I.S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1206 (8-Methyldihydrochelerythrine)

Bodamine **B-240**

$C_{17}H_{21}NO_3$ 287.358
Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine bowdenii* (Amaryllidaceae). Mp 208-210°. $[\alpha]_D$ 0 (c, 0.15 in $CHCl_3$).
Hydroiodide: Mp 245° dec.
Methiodide: Mp 265° dec.
Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109 (*isol*)

Boehmeriasine A **B-241**
11,12,13,14,14a,15-Hexahydro-3,6,7-trimethoxy-9H-phenanthro[9,10-b]quinolinezine, 9CI



$C_{24}H_{27}NO_3$ 377.482

(ξ)-form
Alkaloid from *Boehmeria siamensis*. Cytotoxic. Needles ($CH_2Cl_2/MeOH$). Mp 216-218°. $[\alpha]_D^{20}$ -80.4 (c, 0.1 in MeOH). λ_{max} 227 (log ε 4.36); 253 (log ε 3.85); 282 (log ε 3.27) (MeOH).


O⁶-De-Me: Boehmeriasine BC₂₃H₂₅NO₃ 363.455

Alkaloid from *Boehmeria siamensis*. Powder (CH₂Cl₂/MeOH). Mp 248-250°. [α]_D²⁰ -63.7 (c, 0.2 in MeOH). λ_{max} 228 (log ε 4.39); 252 (log ε 3.9); 285 (log ε 3.34) (MeOH).

Luo, Y.G. *et al.*, *Planta Med.*, 2003, **69**, 842-845 (*isol*, *pmr*, *cmr*)

Bohemamine

[72926-12-6]



Relative Configuration

C₁₄H₁₈N₂O₃ 262.308

Prod. by *Actinosporangium* sp. strain C36145, *Streptomyces* sp. UMA-044 and the marine-derived *Streptomyces* sp. CNQ-583. Cell adhesion inhibitor. Cryst. (CH₂Cl₂/Et₂O). Mp 199-200° dec. [α]_D²⁵ +16 (c, 2 in MeOH).

Deepoxy: Deepoxybohemamine. Antibiotic NP 25302. NP 25302C₁₄H₂₀N₂O₂ 248.324

Prod. by *Streptomyces* sp. UMA-044 and the marine-derived *Streptomyces* sp. CNQ-583. Cell adhesion inhibitor. Amorph. solid (CHCl₃). Mp 229-230°. [α]_D +115.5 (c, 1.1 in MeOH). λ_{max} 250; 282; 334 (MeOH).

Deepoxy, 6α-hydroxy: Bohemamine BC₁₄H₂₀N₂O₃ 264.324

Prod. by the marine-derived *Streptomyces* sp. CNQ-583. Viscous oil. [α]_D²⁵ -6.8 (c, 1 in MeOH). λ_{max} 250 (log ε 4.3); 284 (log ε 4); 332 (log ε 3.9) (MeOH).

Deepoxy, 7α-hydroxy: Bohemamine CC₁₄H₂₀N₂O₃ 264.324

Prod. by the marine-derived *Streptomyces* sp. CNQ-583. [α]_D²⁵ -12 (c, 0.6 in MeOH). λ_{max} 250 (log ε 4.2); 281 (log ε 3.7); 325 (log ε 3.4) (MeOH).

Deepoxy, 6β-chloro, 7α-hydroxy: 6-Chloroboheamine CC₁₄H₁₉ClN₂O₃ 298.768

Prod. by the marine-derived *Streptomyces* sp. CNQ-583. [α]_D²⁵ -14.5 (c, 0.4 in MeOH). λ_{max} 250 (log ε 4.1); 281 (log ε 4); 330 (log ε 3.8) (MeOH).

Doyle, T.W. *et al.*, *J.O.C.*, 1980, **45**, 1324-1326 (*isol*, *uv*, *pmr*, *cmr*, *ms*, *cryst struct*)


Zhang, Q. *et al.*, *J. Antibiot.*, 2003, **56**, 673-681 (*isol*, *pmr*, *cmr*, *activity*)

Bugni, T.S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1626-1628 (*marine*, *isol*, *pmr*, *cmr*)

Duvall, J.R. *et al.*, *J.O.C.*, 2006, **71**, 8579-8590 (*NP 25302*, *synth*)

Boldine

5,6,6a,7-Tetrahydro-1,10-dimethoxy-6-methyl-4H-dibenzo[de,g]quinoline-2,9-diol, 9CI. 2,9-Dihydroxy-1,10-dimethoxy-yaporphine
[476-70-0]

**(R)-form**C₁₉H₂₁NO₄ 327.379

Log P 2.42 (uncertain value) (calc).

▶CE0750000

(R)-form**N-De-Me: (-)-Lauroilsine**

[142794-65-8]

C₁₈H₁₉NO₄ 313.352

Alkaloid from bark of *Dehaasia kurzii* (Lauraceae). Solid. [α]_D²⁷ -162 (c, 0.01 M in MeOH).

(S)-form

Alkaloid from a variety of genera in the Annonaceae (*Desmos*, *Polyalthia*), Lauraceae (*Actinodaphne*, *Litsea*, *Neolitsea*, *Sassafras*, *Laurus*, *Machilus*), Magnoliaceae (*Liriodendron*), Monimiaceae (*Peumus*, *Hedycarya*, *Monimia*), Rhamnaceae (*Retanilla*), and Monimiaceae (*Laurelia*). Diuretic. *Peumus boldus* and boldine have been used in treatment of gall-stones. Antioxidant, shows antitrypanosomal activity. Mp 161-163°. [α]_D¹³ +111 (c, 1.03 in EtOH). λ_{max} 220 (log ε 4.6); 283 (log ε 4.21); 304 (log ε 4.23) (EtOH).

▶LD₅₀ (mus, orl) 450 mg/kg. CE0750000**Hydrochloride:** [16625-69-7]

Cryst. (MeOH/Et₂O). Mp 218-222° dec. [α]_D²⁰ +118 (c, 0.1 in EtOH).

N-Oxide(R-): Boldine N^β-oxide

[937018-76-3]

C₁₉H₂₁NO₅ 343.379

Alkaloid from the leaves of *Neolitsea sericea* var. *aurata*.

N-De-Me: 2,9-Dihydroxy-1,10-dimethoxy-yaporphine. Lauroilsine. Norboldine
[5890-18-6]

C₁₈H₁₉NO₄ 313.352

Alkaloid from a wide variety of genera in the Annonaceae (*Xylopia*), Hernandiaceae (*Hernandia*), Lauraceae (*Actinodaphne*, *Cinnamomum*, *Lindera*, *Litsea*, *Neolitsea*, *Machilus*, *Phoebe*, *Sassafras*), Monimiaceae (*Monimia*, *Peumus*), Monimiaceae (*Laurelia*) and Rhamnaceae (*Retanilla*). Mp 138-140°. [α]_D +101 (c, 0.69 in EtOH).

▶RB5935000

N-De-Me, N-Ac: Mp 255-260°. [α]_D +370 (c, 0.11 in CHCl₃).

N-Me: N-Methylboldine

[73892-20-3]

C₂₀H₂₄NO₄[⊕] 342.414

Quaternary alkaloid from the leaves of

Cocculus laurifolius (Menispermaceae). Cryst. (MeOH/Et₂O) (as chloride). Mp 252-255° dec. (chloride).

Späth, E. *et al.*, *Ber.*, 1933, **66**, 904-914 (*isol*, *struct*)

Nakasato, T. *et al.*, *Chem. Pharm. Bull.*, 1959, **7**, 780-784 (*Lauroilsine*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1277-1281 (*isol*, *pmr*, *Lauroilsine*)

Tewari, S. *et al.*, *Phytochemistry*, 1972, **11**, 1149-1152 (*Boldine*, *Lauroilsine*, *isol*, *pmr*, *uv*, *ms*)

Kupchan, S.M. *et al.*, *Chem. Comm.*, 1976, 91 (*synth*)

Bhakuni, D.S. *et al.*, *J.C.S. Perkin I*, 1977, 706 (*biosynth*)

Bremner, J.B. *et al.*, *Aust. J. Chem.*, 1978, **31**, 313-320 (*N-Methylboldine*)

Saxena, N.K. *et al.*, *J. Indian Chem. Soc.*, 1979, **56**, 1020-1023 (*N-Methylboldine*)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (*cd*)

Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1991, **62**, 261-265 (*(-)-Lauroilsine*)

Asencio, M. *et al.*, *Acta Cryst. C*, 1996, **52**, 1581 (*cryst struct*)

Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 1554


Reveco, P.G. *et al.*, *Synth. Commun.*, 2005, **35**, 341-347 (*ir*, *uv*, *derivis*)

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 637-642 (*N-oxide*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, DNZ100

Bonabiline A

[935761-54-9]



Relative Configuration

C₁₈H₂₉NO₄ 323.431

Alkaloid from the roots of *Bonamia spectabilis*. Muscarinic M₃ receptor antagonist. Yellow oil. [α]_D²⁰ -7.3 (c, 0.3 in CHCl₃).

3-Deoxy, 2,3-didehydro: Bonabiline B

[935761-55-0]


C₁₈H₂₇NO₃ 305.416

Alkaloid from the roots of *Bonamia spectabilis*. Yellow oil.

Ott, S.C. *et al.*, *Planta Med.*, 2006, **72**, 1403-1406 (*isol*, *pmr*, *cmr*, *ms*)

Bonafousine

[60820-65-7]

**B-245**


C₃₅H₄₀N₄O₃ 564.726

Alkaloid from the leaves of *Bonafousia tetrastachya* (preferred genus name *Tabernaemontana*) (Apocynaceae). Cryst. + 2MeOH (MeOH). Mp 199-200°. [α]_D²⁰ -35 (c, 1 in CHCl₃).

Hydrochloride (1:2):

Cryst. + 3H₂O (H₂O). Mp 235° dec.Damak, M. et al., *Chem. Comm.*, 1976, 510 (uv, ir, pmr, cmr, ms, cryst struct)Damak, M. et al., *Bull. Soc. Chim. Fr.*, 1980, 490 (isol, uv, ir, pmr, cmr, ms)**Boneratamide A****B-246**


[786698-99-5]

C₂₄H₃₈N₂O₄ 418.575

Constit. of *Axinyssa aphysinoides*. Deriv. of 1-Spiroaxen-6-amine, in S-425.

Williams, D.E. et al., *J. Nat. Prod.*, 2004, 67, 1752-1754 (isol, pmr, cmr, cryst struct)**Boneratamide B****B-247**

[786699-00-1]

C₂₃H₃₆N₂O₄ 404.548


Constit. of *Axinyssa aphysinoides*. Deriv. of 1-Spiroaxen-6-amine, in S-425.

Stereoisomer: **Boneratamide C**

[786699-02-3]

C₂₃H₃₆N₂O₄ 404.548Constit. of *Axinyssa aphysinoides*.Williams, D.E. et al., *J. Nat. Prod.*, 2004, 67, 1752-1754 (isol, pmr, cmr)**Bongardine****B-248**

α ,6-Dimethyl-3-pyridinemethanamine, 5-(1-Aminoethyl)-2-methylpyridine

C₈H₁₂N₂ 136.196

Alkaloid from the tubers of *Bongardia chrysogonum*. Pale yellow powder. [α]_D²⁰ -48 (c, 0.36 in MeOH). λ _{max} 205 (log ϵ 6.1); 265 (log ϵ 5.5) (MeOH).


N-Ac: 5-[(1-Acetylamino)ethyl]-2-methylpyridine. **N-Acetylbongardine**

C₁₀H₁₄N₂O 178.233

Alkaloid from the tubers of *Bongardia chrysogonum*. Amorph. powder. [α]_D²⁵ -56 (c, 0.1 in MeOH). λ _{max} 200 (log ϵ 7.2); 280 (log ϵ 4.5) (MeOH).

Atta-ur-Rahman, et al., *Phytochemistry*, 1999, 50, 333-336 (isol, uv, pmr, cmr)**Boophiline****B-249**

[199536-52-2]

C₃₃H₅₅NO₇S 609.866

Isol. from the cattle tick *Boophilus microplus*. Powder. Mp 185-190°. [α]_D²⁰ -29 (c, 0.5 in DMSO).

Potterat, O. et al., *Helv. Chim. Acta*, 1997, 80, 2066-2072 (isol, ir, pmr, cmr)**Borbonicine****B-250**


[105053-77-8]

C₁₇H₂₄N₂O 272.389

Struct. unknown. Alkaloid from *Lycopodium clavatum* var. *borbonicum* (Lycopodiaceae).

MacLean, D.B. et al., *Alkaloids (Academic Press)*, 1985, 26, 241Gerard, R.V. et al., *Phytochemistry*, 1986, 25, 1143-1150**Borrecapine****B-251**

[3-Ethenyl-3-methyl-5-(2-methyl-1-propenyl)-2-pyrrolidinyl]-1H-indol-3-ylmethanone, 9CI
[66408-14-8]

C₂₀H₂₄N₂O 308.422

Various numbering systems in use. Alkaloid from *Borreria capitata* (Rubiaceae). Mp 203-205°. [α]_D -75 (c, 0.66 in MeOH). λ _{max} 215 (log ϵ 4.34); 245 (log ϵ 4.15); 263 (log ϵ 4.01); 304 (log ϵ 4.12) (MeOH).

12,18-Didehydro: **Dehydroborrecapine**

[80733-93-3]

C₂₀H₂₂N₂O 306.407

Alkaloid from *Borreria capitata* (Rubiaceae). Cryst. (MeOH). Mp 179-181°.

12,18-Didehydro, 19 ξ ,20-epoxide: **Borrecoxine**


[80733-92-2]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Borreria capitata* (Rubiaceae). Cryst. (MeOH). Mp 193°.

Jössang, A. et al., *Tet. Lett.*, 1977, 18, 4317-4318 (isol, uv, pmr, struct)Jössang, A. et al., *Planta Med.*, 1981, 43, 301-304 (Dehydroborrecapine, Borrecoxine)**Borreline****B-252**

[64643-94-3]


C₁₇H₁₈N₂O 266.342

Alkaloid from *Borreria* spp. (Rubiaceae). Mp 215-216°. [α]_D -235 (c, 0.7 in MeOH).

12,13-Dihydro: Mp 216°.

Jössang, A. et al., *Tet. Lett.*, 1977, 1219-1220 (uv, ir, ms, cmr, pmr, cryst struct)**Borreverine****B-253**

[51109-65-0]

C₃₂H₄₀N₄ 480.695

Alkaloid from *Borreria verticillata* and *Flindersia fourmieri* (Rubiaceae, Flindersiaceae). Strongly active against gram-positive bacteria. Mp 193°. Racemic. λ _{max} 225 (ϵ 60260); 284 (ϵ 27540); 295 (ϵ 25120) (EtOH).

N⁴-Me: **Auricularine. 4-Methylborreverine**

[73706-32-8]

C₃₃H₄₂N₄ 494.722

Isol. from *Flindersia fourmieri* and *Hedyotis auricularia*. Amorph. [α]_D +41 (c, 0.03 in CHCl₃) (0.0). λ _{max} 226 (log ϵ 4.53); 250 (sh) (log ϵ 3.96); 286 (log ϵ 4.01); 294 (log ϵ 4.01) (EtOH).

14',15'-Dihydro, 15 ξ -hydroxy: **15'-Hydroxy-14',15'-dihydroborreverine**

[73706-33-9]


C₃₂H₄₂N₄O 498.71

Isol. from *Flindersia fourmieri*. Amorph.

Pousset, J.-L. et al., *Chem. Comm.*, 1977, 261-262 (uv, ms, pmr, cryst struct)Tillequin, F. et al., *Chem. Comm.*, 1978, 826-828 (synth)Tillequin, F. et al., *J. Nat. Prod.*, 1979, 42, 92-95 (isol)Tillequin, F. et al., *Phytochemistry*, 1979, 18, 1559-1561; 2066-2067 (N⁴-Me, 14',15'-dihydro-15'-hydroxy)Maynard, G. et al., *C. R. Hebd. Seances Soc. Biol. Ses Fil.*, 1980, 174, 925-928 (activity)Purushothaman, K.K. et al., *Phytochemistry*, 1981, 20, 351-352 (Auricularine)

Boschniakine
Indicaine

B-254



(*R*)-form

C₁₀H₁₁NO 161.203

(*R*)-form [18070-40-1]

Alkaloid from *Boschniakia rossica*, *Pedicularis ludwigi*, *Pedicularis olgae*, *Plantago albicans*, *Plantago indica*, *Plantago major*, *Plantago notata*, *Plantago psyllium* (African plantain), *Plantago ramosa*, *Tecoma radicans* and *Tecoma stans*. Also found in the defence substance of *Megacranium tsudai*. Bp₃ 80-90°. [α]_D²⁰ +21 (c, 0.98 in CHCl₃).

Picrate: Mp 126-128° Mp 156° (dimorph.).

Semicarbazone: Mp 217-220° (210-224°).

N-Et. Indicainine

[32152-74-2]
C₁₂H₁₆NO[⊕] 190.264

Quaternary alkaloid from *Plantago olgae* (Scrophulariaceae). Mp 125-127° (as picrate). [α]_D³⁰ +14.2 (c, 1.9 in CHCl₃) (picrate). The correctness of the struct. has been questioned (D. Gross *et al.*).

Carboxylic acid: Boschniakic acid.

Plantagonine
[21857-97-6]
C₁₀H₁₁NO₂ 177.202

Alkaloid from *Boschniakia rossica*, *Pedicularis dolichorrhiza*, *Pedicularis ludwigi*, *Pedicularis olgae*, *Plantago albicans*, *Plantago coronopus*, *Plantago crassifolia*, *Plantago crypsoides*, *Plantago cylindrica*, *Plantago indica*, *Plantago major*, *Plantago notata*, *Plantago ovata*, *Plantago psyllium* (African plantain), *Plantago ramosa*, *Verbascum songaricum* (Orobanchaceae, Scrophulariaceae, Plantaginaceae). Mp 218-220°. [α]_D +38.6 (+ 30.8). λ_{max} 271 (log ε 3.5) (MeOH).

(*S*)-form

Carboxylic acid: [21913-34-8]
Alkaloid from *Incarvillea olgae* (Bignoniaceae) Mp 218-220°. [α]_D -30.1.]

(±)-form

Carboxylic acid: Synthetic. Mp 226-227°.

(ξ)-form

Carboxylic acid, Me ester: Deoxyrhexifoline

[94054-29-2]
C₁₁H₁₃NO₂ 191.229


Alkaloid from the seeds of *Castilleja rhexifolia* aff. *miniata* (Scrophulariaceae). [α]_D²⁰ -18 (c, 0.3 in CHCl₃) (synthetic). Abs. config. of natural product not detd. due to paucity of material. The *S*-enantiomer was synthesised as a foam.

Sakan, T. *et al.*, *Tetrahedron*, 1967, **23**, 4635-4652 (*Boschniakine, Boschniakic acid, uv, ir, synth, struct*)
Torsell, K. *et al.*, *Acta Chem. Scand.*, 1968, **22**, 2715 (*struct*)
Dickinson, E.M. *et al.*, *Tetrahedron*, 1969, **25**, 1523 (*pmr*)
Khakimdzhano, S. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 126-127; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 121 (*Indicainine*)
Gross, D. *et al.*, *Z. Chem.*, 1973, **13**, 296 (*ms*)
Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (*rev*)
Roby, M.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 846-853 (*Deoxyrhexifoline*)
Ranarivelo, Y. *et al.*, *Heterocycles*, 1990, **31**, 1727-1731 (*(-)-Deoxyrhexifoline, synth*)
Ho, H.-Y. *et al.*, *J. Chem. Ecol.*, 1993, **19**, 39-46 (*Megacranium, isol*)
Lin, L.-C. *et al.*, *Chin. Pharm. J. (Taipei)*, 2004, **56**, 77-85 (*Boschniakic acid, isol, pmr, cmr*)
Fu, J.-J. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2151-2155 (*isol, pmr, cmr*)
Robert, N. *et al.*, *Tetrahedron*, 2007, **63**, 3702-3706 (*synth*)

Botrycoidin

B-255

*6,9-Dihydroxy-7-methoxy-3-methylbenz[*g*]isoquinoline-5,10-dione*
[4589-33-7]



C₁₅H₁₁NO₅ 285.256

Pigment from *Fusarium botrycoides* and *Fusarium solani*. Active against gram-positive bacteria and yeast. Antitubercular. Dark red needles. Sol. MeOH, acids, bases, CHCl₃; fairly sol. C₆H₆, Me₂CO; poorly sol. H₂O, hexane. Mp 241-243° dec. λ_{max} 252 (ε 25100); 321 (ε 5010); 475 (sh); 499 (ε 7940); 527 (sh) (EtOH) (Derep). λ_{max} 251 (ε 15850); 320 (ε 8510); 497 (ε 9550) (EtOH) (Berdy).

▶ LD₅₀ (mus, ipr) 250 mg/kg, LD₅₀ (mus, scu) 250 mg/kg.
Ac: Mp 214°.

9-Me ether: 9-O-Methylbotrycoidin

[73590-03-1]
C₁₆H₁₃NO₅ 299.282

Major pigment of a strain of *Fusarium moniliforme* and *Fusarium oxysporum*. Active against gram-positive bacteria. Red needles (CHCl₃/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 215-216°. 8-Me ether acc. to the trivial numbering system. λ_{max} 227 (ε 20900); 262 (ε 15800); 310 (ε 7590); 510 (ε 5370) (MeOH/HCl) (Derep). λ_{max} 259 (ε 27500); 306 (ε 6610); 546 (ε 10000) (MeOH/NaOH) (Derep). λ_{max} 248 (ε 31600); 318 (ε 8320); 480 (ε 6760) (MeOH) (Derep).

9-Me ether, Ac: Mp 245-246°.

7-O-De-Me: Tolypocladine

[131815-26-4]
C₁₄H₉NO₅ 271.229

Prod. by *Tolypocladium inflatum*. Me-

tal chelating agent. Sol. DMF, DMSO, bases; fairly sol. MeOH, Me₂CO, CHCl₃; poorly sol. hexane, H₂O. λ_{max} 250 ; 320 ; 505 ; 545 (MeOH) (Berdy). λ_{max} 219 (ε 14100); 280 (ε 10200); 290 ; 420 ; 525 ; 560 (MeOH/NaOH) (Berdy). λ_{max} 252 ; 340 ; 476 ; 496 ; 530 (EtOH) (Berdy).

6-Deoxy: 6-Deoxybotrycoidin

[126262-44-0]
C₁₅H₁₁NO₄ 269.256

Metab. of the fungus, *Nectria haematococca*. Yellow needles (CH₂Cl₂/hexane). Mp 195-196°. λ_{max} 207 ; 237 ; 270 ; 414 (MeOH) (Berdy).

6-Deoxy, 9-Me ether: 7,9-Dimethoxy-3-methylbenz[*g*]isoquinoline-5,10-dione.

Scorpinone

C₁₆H₁₃NO₄ 283.283
Prod. by a *Bispora*-like fungal sp. Yellow needles (CHCl₃). Mp 195°. λ_{max} 238 ; 282 ; 321 (sh) ; 405 (MeOH).

6-Deoxy, 7-O-de-Me: 7-O-Demethyl-6-deoxybotrycoidin

[132311-85-4]
C₁₄H₉NO₄ 255.229
Metab. of *Nectria haematococca*. Yellow plates (EtOAc). Mp 300-305° dec. λ_{max} 205 (ε 25100); 239 (ε 26200); 280 (ε 13200); 421 (ε 5600) (MeOH) (Berdy).

Arsenault, G.P. *et al.*, *Tet. Lett.*, 1965, 4033 (*struct*)

Steyn, P.S. *et al.*, *Tetrahedron*, 1979, **35**, 1551 (*deriv*)

Cameron, D.W. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1439 (*synth, bibl*)

McCulloch, A.W. *et al.*, *Can. J. Chem.*, 1982, **60**, 2943 (*isol, props*)

Watanabe, M. *et al.*, *Tetrahedron*, 1987, **43**, 5281 (*synth*)

Parisot, D. *et al.*, *Z. Naturforsch., B*, 1989, **44**, 1473 (*6-Deoxybotrycoidin*)

Graefe, U. *et al.*, *Biol. Met.*, 1990, **3**, 39 (*Tolypocladine*)

Parisot, D. *et al.*, *Phytochemistry*, 1990, **29**, 3364 (*deriv*)

Kesteleyn, B. *et al.*, *J.O.C.*, 2000, **65**, 640-644 (*6-Deoxybotrycoidin, synth*)

Miljkovic, A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1251-1253 (*Scorpinone*)

Van, T.N. *et al.*, *Tetrahedron*, 2005, **61**, 2295-2300 (*synth*)


Van Wagoner, R.M. *et al.*, *J. Nat. Prod.*, 2008, **71**, 426-439 (*Scorpinone, biosynth*)

Choshi, T. *et al.*, *Tet. Lett.*, 2008, **49**, 3725-3728 (*Scorpinone, synth*)

Botryllamide A

B-256

N-[2-(3,5-Dibromo-4-methoxyphenylethynyl)-3-(4-hydroxyphenyl)-2-methoxy-2-propenamide, 9CI
[163564-64-5]



C₁₉H₁₇Br₂NO₄ 483.156

Isol. from the ascidians *Botrylloides tyreum*, *Botryllus schlosseri* and a *Bo-*

tryllus sp. from the Philippines. Needles (CHCl₃). Mp 169-171°.

4'-O-De-Me: Botryllamide G

[724434-05-3]
C₁₈H₁₅Br₂NO₄ 469.129
Isol. from *Botrylloides tyreum*. Gum.
λ_{max} 208 (ε 22230); 332 (ε 22150)
(MeOH).

Monodebromo: Botryllamide C

[163564-66-7]
C₁₉H₁₈BrNO₄ 404.259
Isol. from *Botrylloides tyreum*, *Botryllus schlosseri* and *Botryllus* sp. Needles (CHCl₃). Mp 173-175°.

Bis(debromo): Botryllamide E

[724434-03-1]
C₁₉H₁₉NO₄ 325.363
Isol. from *Botrylloides tyreum*. Gum.
λ_{max} 207 (ε 15000); 222 (ε 12540); 339
(ε 29800) (MeOH).

Bis(debromo), 4'-O-de-Me: Botryllamide F

[724434-04-2]
C₁₈H₁₇NO₄ 311.337
Isol. from *Botrylloides tyreum*. Oil.
λ_{max} 208 (ε 13780); 220 (ε 12000); 316
(ε 14260) (MeOH).

(5E)-Isomer: Botryllamide B

[163564-65-6]
C₁₉H₁₇Br₂NO₄ 483.156
Isol. from *Botryllus schlosseri* and
Botryllus sp. Gum.

(5E)-Isomer, monodebromo: Botryllamide D

[163564-67-8]
C₁₉H₁₈BrNO₄ 404.259
Isol. from *Botryllus schlosseri*. Shows
marginal cytotoxicity against the hu-
man cancer cell line HCT 116 but is
inactive *in vivo*. Gum. λ_{max} 203 (ε
11100); 326 (ε 11700) (MeOH) (Berdy).


McDonald, L.A. *et al.*, *Tetrahedron*, 1995, **51**,
5237-5244 (*isol, uv, ir, pmr, cmr, ms*,
Botryllamides B,C)

Rao, M.R. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1064-
1066 (*isol, pmr, cmr, Botryllamides E-G*)

Botryllamide H

B-257

3-(2-Cyano-3-hydroxyphenyl)-2-hydroxy-N-[2-(4-hydroxyphenyl)ethenyl]-2-propanamide, 9CI
[724434-06-4]




C₁₈H₁₄N₂O₄ 322.32
Isol. from the ascidian *Botrylloides tyreum*. Powder. λ_{max} 210 (ε 17630); 249 (ε 12570); 314 (ε 7790) (MeOH).

Rao, M.R. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1064-
1066 (*isol, pmr, cmr*)

Botryllazine A

B-258

2,3-Bis(4-hydroxybenzoyl)-5-(4-hydroxyphenyl)pyrazine
[252026-23-6]



C₂₄H₁₆N₂O₅ 412.401
Alkaloid from the red ascidian *Botryllus leachi*. Yellow powder. λ_{max} 228 (ε 18500); 310 (ε 27500) (MeOH).


Duran, R. *et al.*, *Tetrahedron*, 1999, **55**, 13225-
13232 (*isol*)

Buron, F. *et al.*, *J.O.C.*, 2005, **70**, 2616-2621
(*synth*)

Botryllazine B

B-259

2-(4-Hydroxybenzoyl)-6-(4-hydroxyphenyl)pyrazine
[252026-25-8]



C₁₇H₁₂N₂O₃ 292.293
Alkaloid from the red ascidian *Botryllus leachi*. Yellow powder. λ_{max} 228 (ε 14700); 298 (ε 22800) (MeOH).

Duran, R. *et al.*, *Tetrahedron*, 1999, **55**, 13225-
13232 (*isol*)


Mahboobi, S. *et al.*, *Monatsh. Chem.*, 2004,
135, 333-342 (*synth*)

Buron, F. *et al.*, *J.O.C.*, 2005, **70**, 2616-2621
(*synth*)

Bouchardatine

B-260

[623903-29-7]




C₁₇H₁₁N₃O₂ 289.293
Alkaloid from the aerial parts of *Bouchardatia neurococca*. Amorph. yellow powder. λ_{max} 254 (log ε 3.3); 351 (log ε 3.2); 367 (sh); 391 (sh) (no solvent reported).

Wattanapiromsakul, C. *et al.*, *Phytochemistry*,
2003, **64**, 609-615 (*isol, pmr, cmr*)

Bowdensine

B-261

[1354-75-2]



Absolute
Configuration

C₂₁H₂₅NO₇ 403.431
Alkaloid from the bulbs of *Nerine bowdenii* (Amaryllidaceae). Glass. [α]_D²⁴
+17.3. [α]₄₃₆²⁴ +48.1 (c, 1.06 in CHCl₃).

Perchlorate:

Prisms + 1Me₂CO (Me₂CO). Mp 260-
262° dec. [α]_D²⁴ +5. [α]₄₃₆²⁴ +15 (c, 0.4 in
75% EtOH).

Methodide:

Cryst. (EtOH). Mp 284-285° dec. [α]_D²⁴
+9.8. [α]₄₃₆²⁴ +25.5 (c, 1.30 in 90%
EtOH).

Di-de-Ac: Deacetylbowdensine

[3660-65-9]
C₁₇H₂₁NO₅ 319.357
Isol. from the bulbs of *Nerine bowdenii*
and *Crinum erubescens* (Amaryllida-
ceae). Mp 277-278° dec. [α]_D²⁵ -43.7.
[α]₄₃₆²⁵ -86.6 (c, 0.5 in EtOH). Probably
an artifact.

1-Epimer: 1-Epibowdensine

[184900-13-8]
C₂₁H₂₅NO₇ 403.431
Alkaloid from bulbs of *Brunsvigia
orientalis*. Mp 124-126°. [α]_D²² +4 (c, 1.1
in CHCl₃).

1-Epimer, di-de-Ac: 1-Epideacetylbowdensine. Bulbisine

[101219-55-0]
C₁₇H₂₁NO₅ 319.357
Alkaloid from bulbs of *Brunsvigia
orientalis* and *Crinum bulbispermum*.
Mp 162-164°. [α]_D²² +22 (c, 0.47 in
CHCl₃).

Lyle, R.E. *et al.*, *J.A.C.S.*, 1960, **82**, 2620-2625
(*isol*)

Chapman, O.L. *et al.*, *J.A.C.S.*, 1964, **86**, 1256-
1258 (*di-de-Ac, pmr*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902-
4912 (*di-de-Ac, ms, struct*)

Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1984,
32, 3015-3022 (*Deacetylbowdensine*)

Frahm, A.W. *et al.*, *Magn. Reson. Chem.*,
1985, **23**, 804-808 (*cmr, Bulbisine*)

Ali, A.A. *et al.*, *CA*, 1990, **112**, 84290m


(*Bulbisine*)

Viladomat, F. *et al.*, *Phytochemistry*, 1996, **43**,
1379-1384 (*epimers*)

Boxazomycin A

B-262

5-Hydroxy-2-[5-hydroxy-2-(hydroxymethyl)-4-pyrimidinyl]-7-methyl-4-benzoxazolecarboxamide, 9CI
[107021-64-7]



C₁₄H₁₂N₄O₅ 316.273

Prod. by *Pseudonocardia* sp. G495-11. Active against gram-positive bacteria and anaerobes. Pale yellow needles (MeOH). Sol. bases, DMF; fairly sol. H₂O, Me₂CO, MeOH, EtOH; poorly sol. EtOAc, C₆H₆, hexane, CHCl₃. Mp 275° dec. λ_{max} 219 (ε 28800); 230 (sh) (ε 24500); 259 (sh) (ε 13200); 403 (ε 20000) (0.1M NaOH) (Derep). λ_{max} 364 (ε 13500); 379 (sh) (ε 12600); 420 (sh) (ε 6170) (EtOH) (Derep).

1''-Deoxy: Boxazomycin B

[107021-65-8]

C₁₄H₁₂N₄O₄ 300.273

Prod. by *Pseudonocardia* sp. G495-11. Similar biol. props. as parent. Pale yellow powder (MeOH). Sol. bases, DMF; fairly sol. H₂O, Me₂CO, EtOH, MeOH; poorly sol. EtOAc, CHCl₃, hexane, C₆H₆. Mp 270° dec. λ_{max} 219 (ε 28800); 230 (sh) (ε 24500); 259 (sh) (ε 13200); 403 (ε 20000) (0.1M NaOH) (Derep). λ_{max} 364 (ε 13500); 379 (sh) (ε 12600); 420 (sh) (ε 6170) (EtOH) (Derep).

5-Deoxy: Boxazomycin C

[107021-66-9]


[107021-71-6 (di-Ac)]

C₁₄H₁₂N₄O₄ 300.273

Prod. by *Pseudonocardia* sp. G495-11. Similar biol. props. as parent. Silky pink needles (as di-Ac). Mp 230-231° (di-Ac). λ_{max} 224 (ε 15800); 252 (ε 10200); 318 (ε 16200) (MeOH) (Derep).

Kusumi, T. et al., *J.A.C.S.*, 1988, **110**, 2954-2958 (pmr, cmr, struct)Suto, M.J. et al., *Tet. Lett.*, 1995, **36**, 7213-7216 (Boxazomycin B, synth)**Brachyamide A****B-263**


1-[13-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,12-tridecatrienyl]pyrrolidine, 9CI. 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienoic acid pyrrolidide

C₂₄H₃₁NO₃ 381.514**(E,E,E)-form** [119736-85-5]

Minor alkaloid from the fruits of *Piper brachystachyum* (Piperaceae). Semisolid.

Koul, S.K. et al., *Phytochemistry*, 1988, **27**, 3523 (isol, uv, ir, pmr, cmr, ms, struct)**Brachycerine****B-264**

[349647-30-9]

C₂₇H₃₄N₂O₁₀ 546.573

Studies in 2008 suggest that the C-1' config. should be revised to that shown. Alkaloid from the leaves of *Psychotria brachyceras*. Powder. [α]_D²⁰ -24.5 (c, 0.2 in MeOH). λ_{max} 226 (log ε 4.59); 280 (log ε 3.88) (MeOH).

Deoxy, 7',8'-didehydro: Croceaine A

[800391-12-2]

C₂₇H₃₂N₂O₉ 528.558

Alkaloid from the leaves of *Palicourea crocea*. Amorph. solid. [α]_D²⁵ -11 (c, 0.003 in MeOH).

Deoxy, 1,2,3,4,7',8'-hexadehydro: Croceaine B

[800391-13-3]

C₂₇H₂₈N₂O₉ 524.526

Alkaloid from the leaves of *Palicourea crocea*. Amorph. solid. [α]_D²⁵ -40 (c, 0.002 in MeOH).

1-Epimer, 7'-deoxy, 7',8'-didehydro: Psychollatine. Umbellatine†

[850307-03-8]


[850355-58-7]

C₂₇H₃₂N₂O₉ 528.558

Alkaloid from the leaves of *Psychotria umbellata*. 5HT_{2A/C} serotonin modulator. Analgesic. Amorph. powder. [α]_D²⁰ -24 (c, 0.6 in MeOH). λ_{max} 226 ; 280 (MeOH).

Kerber, V.A. et al., *J. Nat. Prod.*, 2001, **64**, 677-679; 2003, **66**, 1038 (*Brachycerine*)Düsmann, L.T. et al., *J. Nat. Prod.*, 2004, **67**, 1886-1888 (*Croceaine A,B*)Both, F.L. et al., *J. Nat. Prod.*, 2005, **68**, 374-380; 2006, **69**, 342-345 (*activity, Psychollatine*)Kerber, V.A. et al., *J. Nat. Prod.*, 2008, **71**, 697-700 (*Psychollatine, Croceaine A*)**Brachystamide B****B-265**

15-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4,14-pentadecatrienamide, 9CI. 15-(3,4-Methylenedioxyphenyl)-2,4,14-pentadecatrienoic acid isobutylamide. N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,14-pentadecatrienamide

C₂₆H₃₇NO₃ 411.583**(E,E,E)-form** [126394-65-8]

Alkaloid from the aerial parts of *Piper brachystachyum* (Piperaceae). Amorph. solid.

14,15-Dihydro: **Brachystamide A**. 15-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4-pentadecadienamide, 9CI. 15-(3,4-Methylenedioxyphenyl)-2,4-pentadecadienoic acid isobutylamide [126394-64-7]

C₂₆H₃₉NO₃ 413.599

Alkaloid from the aerial parts of *Piper brachystachyum* (Piperaceae). Mp 101°.

14,15-Dihydro, 12,13-didehydro(E)-: 15-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4,12-pentadecatrienamide,

9CI. N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,12-pentadecatrienamide. **Ridleyamide**

[170475-09-9]

C₂₆H₃₇NO₃ 411.583

Isol. from stems and leaves of *Piper ridleyi* (Piperaceae). Cryst. Mp 95-98°.

14,15-Dihydro, 13,14-didehydro(E)-: 15-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4,13-pentadecatrienamide. N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,13-pentadecatrienamide. **Brachystamide C**


C₂₆H₃₇NO₃ 411.583

Alkaloid from the aerial parts of *Piper brachystachyum*. Amorph. solid. Mp 90-95°. λ_{max} 208 (log ε 4.15); 260 (log ε 4.41) (EtOH).

Banerji, A. et al., *Phytochemistry*, 1989, **28**, 3039 (isol, uv, ir, pmr, cmr, ms, struct)Ahmad, F. et al., *Phytochemistry*, 1995, **40**, 1163 (*Ridleyamide*)Banerji, A. et al., *Phytochemistry*, 2002, **59**, 897-901 (*Brachystamide C*)**Brachystamide D****B-266**

16-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4,15-hexadecatrienamide. 16-(3,4-Methylenedioxyphenyl)-2,4,15-hexadecatrienoic acid isobutylamide. *Pergumidiene*

[208345-31-7]

C₂₇H₃₉NO₃ 425.61

Alkaloid from the aerial parts of *Piper brachystachyum* and the fruit of *Piper longum*. Amorph. solid. Mp 88-90°.


15,16-Dihydro: 16-(3,4-Methylenedioxyphenyl)-2,4-hexadecadienoic acid isobutylamide. **Brachystamide E**

C₂₇H₄₁NO₃ 427.626

Alkaloid from the aerial parts of *Piper brachystachyum*. Isol. as a mixt. with Brachystamide D.

Das, B. et al., *Nat. Prod. Sci.*, 1998, **4**, 23-25 (*Pergumidiene*)Das, B. et al., *Tet. Lett.*, 1998, **39**, 9099-9100 (synth)Banerji, A. et al., *Phytochemistry*, 2002, **59**, 897-901 (isol, pmr, cmr, ms)**Brachystemidine A****B-267**

[436807-92-0]


C₁₅H₁₈N₂O₅ 306.318

Alkaloid from the roots of *Brachystemma calycinum*. Amorph. solid. Mp 210-211.5°. Laevorotatory but unstable.

Cheng, Y.X. et al., *J. Nat. Prod.*, 2002, **65**, 750-752 (isol, pmr, cmr, ms)

Brachystemidine E

[436807-96-4]

C₁₄H₁₆N₂O₆ 308.29Alkaloid from the roots of *Brachystemma calycinum*. Gum. [α]_D²⁸ +0.76 (c, 1.65 in MeOH).**5'-Me ether: Brachystemidine C**

[436807-94-2]

C₁₅H₁₈N₂O₆ 322.317Alkaloid from the roots of *Brachystemma calycinum*. Gum. [α]_D²¹ -21 (c, 0.25 in CHCl₃).**3'-Deoxy: Brachystemidine D**


[436807-95-3]

C₁₄H₁₆N₂O₅ 292.291Alkaloid from the roots of *Brachystemma calycinum*. Amorph. solid. Mp 147.5-149°. [α]_D²⁵ +3.5 (c, 0.43 in MeOH).**3'-Deoxy, 5'-Me ether: Brachystemidine B**



[436807-93-1]

C₁₅H₁₈N₂O₅ 306.318Alkaloid from the roots of *Brachystemma calycinum*. Amorph. solid. Mp 151-152°. [α]_D²⁷ -3.1 (c, 0.32 in MeOH).Cheng, Y.-X. *et al.*, *J. Nat. Prod.*, 2002, **65**, 750-752 (*isol*, *pmr*, *cmr*, *ms*, *cryst struct*)**Brachystemidine F**

[1050363-02-4]

C₁₅H₁₉N₃O₆ 337.332Alkaloid from the roots of *Brachystemma calycinum*. Oily solid. [α]_D²⁴ +9.3 (c, 0.24 in MeOH). λ _{max} 202 ; 266 (MeOH).Lu, Q. *et al.*, *Chem. Biodiversity*, 2008, **4**, 2948-2952 (*isol*, *pmr*, *cmr*, *ms*)**Brachystemidine G**

[1050363-03-5]

**B-268**C₁₄H₁₆N₄O₆ 336.304Alkaloid from the roots of *Brachystemma calycinum*. Potent immunosuppressant. Gum. [α]_D²⁴ +71.4 (c, 0.13 in MeOH). λ _{max} 203 ; 269 (MeOH).Lu, Q. *et al.*, *Chem. Biodiversity*, 2008, **4**, 2948-2952 (*isol*, *pmr*, *cmr*, *ms*)**Brachystine****B-271***1-(1-Oxo-2,9-nonadecadienyl)pyrrolidine, 9CI. 2,9-Nonadecadienoic acid pyrrolidide*C₂₃H₄₁NO 347.583**(E,Z)-form** [119736-86-6]Minor constit. of fruits of *Piper brachystachyum* (Piperaceae). Semi-solid.Koul, S.K. *et al.*, *Phytochemistry*, 1988, **27**, 3523 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)**Bractamine****B-272**




[93586-21-1]

C₁₁H₁₅NO₂ 193.245Struct. unknown. Alkaloid from *Papaver bracteatum* (Papaveraceae). Cryst. (EtOH). Mp 216°. Opt. inactive. Conts. 1 OMe, 1NMe and 1 OH gp.**Hydrochloride:**Cryst. (EtOH/Et₂O). Mp 143°.**Methiodide:**

Cryst. (EtOH). Mp 220-221°.


Picrate: Mp 184-185°.Kiselev, V.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1948, **18**, 142; *CA*, **42**, 5037b**Bractazonine****B-273**

[93631-24-4]

C₁₉H₂₃NO₃ 313.396Alkaloid from the capsules of *Papaver bracteatum* (Papaveraceae). Mp 101°.Theuns, H.G. *et al.*, *Phytochemistry*, 1984, **23**, 1157-1166 (*isol*, *synth*, *pmr*, *ms*)Chen, W. *et al.*, *J.O.C.*, 2003, **68**, 1929-1932 (*synth*)**Bracteatine****B-274**C₂₅H₃₅N₂O₆[⊕] 459.561Quaternary alkaloid from the aerial parts of *Ophiorrhiza bracteata* (Rubiaceae).**Chloride:** [203256-99-9]C₂₅H₃₅ClN₂O₆ 495.014Needles (MeOH/EtOAc). Mp 250° (dec.). [α]_D -123 (c, 0.8 in MeOH).Arbain, D. *et al.*, *Aust. J. Chem.*, 1997, **50**, 1111-1112 (*isol*, *pmr*, *cmr*)**Bracteine****B-275**C₁₉H₂₁NO₄ 327.379Struct. unknown. Probably identical with Orientalinone, O-112. Alkaloid from *Papaver bracteatum* (Papaveraceae). Cryst. (Et₂O). Mp 230°. [α]_D +120 (CHCl₃).**Hydrochloride:** Mp 258°.**Perchlorate:** Mp 231-232° dec.**Methiodide:** Mp 205-215° (softens at 190°).Kiselev, V.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1948, **18**, 142; *CA*, **42**, 5037bShakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 761-863; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 737-858 (*rev*)**Bracteoline****B-276***5,6,6a,7-Tetrahydro-2,9-dimethoxy-6-methyl-4H-dibenzo[de,g]quinoline-1,10-diol, 9CI. 1,10-Dihydroxy-2,9-dimethoxyaporphine***(S)-form**C₁₉H₂₁NO₄ 327.379**(S)-form** [25651-04-1]Alkaloid from *Papaver bracteatum* and *Papaver orientale* (Papaveraceae). Cryst. (Et₂O). Mp 226-229° (218-221°). [α]_D²⁰ +35 (c, 0.16 in CHCl₃).*N-De-Me: 1,10-Dihydroxy-2,9-dimethoxy-9noraporphine. Norbracteoline* [89825-97-8]C₁₈H₁₉NO₄ 313.352Alkaloid from *Glaucium corniculatum*. Amorph. [α]_D +41 (c, 0.4 in MeOH).**(±)-form** [28230-74-2]Synthetic. Pale yellow needles (Et₂O/MeOH). Mp 209-211° dec.Heydenreich, K. *et al.*, *Pharmazie*, 1967, **22**, 124-125 (*isol*, *uv*, *ir*, *struct*)Delenk-Heydenreich, K. *et al.*, *Pharmazie*, 1969, **24**, 635-645 (*isol*, *uv*, *ir*, *ms*)Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 1513-1514 (*synth*, *uv*, *pmr*, *ms*)Kametani, T. *et al.*, *J.C.S. (C)*, 1971, 2446-2448 (*synth*, *uv*, *ms*)Kerekes, P. *et al.*, *Chem. Ber.*, 1972, **105**, 609-613 (*synth*, *uv*, *ir*, *pmr*)Hara, H. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1921-1925 (*synth*, *ir*, *pmr*)Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1979, **42**, 325-360 (*rev*, *cmr*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 751-753; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 714-716 (*Norbracteoline*)


Bradyoxetin **B-277**
2,2'-(Carbonimidoyldi-4,1-phenylene)bis-3-oxetanamine, 9CI
[494225-42-2]



$C_{19}H_{21}N_5O_2$ 323.394
Prod. by *Bradyrhizobium japonicum*.
Chemical signal involved in symbiotic gene regulation. λ_{max} 271 (MeOH aq.).

Loh, J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2002, **99**, 14446-14451 (*isol, pmr, cmr, activity*)


Brasiliamide A **B-278**



$C_{24}H_{26}N_2O_6$ 438.479
Prod. by *Penicillium brasilianum* Batista JV-379. Prisms (MeOH). Mp 133-134°. λ_{max} 211 (ϵ 42000); 235 (sh) (ϵ 21000); 253 (sh) (ϵ 16000) (MeOH).

Fujita, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 1697-1705 (*isol, pmr, cmr*)

Brasiliamide D **B-279**



$C_{24}H_{28}N_2O_5$ 424.496
Prod. by *Penicillium brasilianum* Batista JV-379. Amorph. powder. Mp 61-63°. $[\alpha]_D^{20}$ +52 (c, 0.72 in MeOH). λ_{max} 209 (ϵ 52000); 242 (sh) (ϵ 4700); 277 (ϵ 1200); 285 (sh) (ϵ 1000) (MeOH).

*N*¹-De-Ac: **Brasiliamide E**

$C_{22}H_{26}N_2O_4$ 382.458
Prod. by *Penicillium brasilianum* Batista JV-379. Amorph. powder. Mp 34-36°. $[\alpha]_D^{20}$ -21 (c, 0.08 in MeOH). λ_{max} 209 (ϵ 49000); 241 (sh) (ϵ 4500); 278 (ϵ 1300); 285 (sh) (ϵ 1200) (MeOH).

2,3-Didehydro: **Brasiliamide B**

$C_{24}H_{26}N_2O_5$ 422.48
Prod. by *Penicillium brasilianum* Batista JV-379. Amorph. powder. Mp 59-61°. $[\alpha]_D^{20}$ +195 (c, 0.62 in MeOH). λ_{max} 208 (ϵ 46000); 270 (ϵ 21000) (MeOH).


1',5-Didehydro(E-): **Brasiliamide C**

$C_{24}H_{26}N_2O_5$ 422.48
Prod. by *Penicillium brasilianum* Ba-

tista JV-379. Amorph. powder. Mp 72-73°. $[\alpha]_D^{20}$ +42 (c, 0.07 in MeOH). λ_{max} 210 (ϵ 47000); 252 (ϵ 13000); 262 (sh) (ϵ 13000) (MeOH).

Fujita, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 1697-1705; 2004, **68**, 820-826 (*isol, uv, pmr, cmr, ms*)

Brasilicardin A **B-280**
[211108-46-2]



$C_{45}H_{68}N_2O_{16}$ 893.036
Constit. of *Nocardia brasiliensis*. Shows potent immunosuppressive activity. Cytotoxic agent. Amorph. solid. Mp 270-273°. $[\alpha]_D^{30}$ +15 (c, 0.5 in MeOH). λ_{max} 212 (ϵ 15000); 239 (ϵ 5200); 300 (ϵ 1900) (MeOH).

4'-Deacyl, 3'-deglycosyl: **Brasilicardin C**
[799804-52-7]
 $C_{30}H_{51}NO_9$ 569.734

Constit. of *Nocardia brasiliensis*. Amorph. solid. $[\alpha]_D^{23}$ +65 (c, 1 in MeOH).

16-Demethoxy: **Brasilicardin B**


[799804-51-6]
 $C_{44}H_{66}N_2O_{15}$ 863.01
Constit. of *Nocardia brasiliensis*. Amorph. solid. $[\alpha]_D^{23}$ +17 (c, 1 in MeOH).

16-Demethoxy, 4'-deacyl, 3'-deglycosyl:

Brasilicardin D
[799804-53-8]
 $C_{29}H_{49}NO_8$ 539.708
Constit. of *Nocardia brasiliensis*. Amorph. solid. $[\alpha]_D^{20}$ +79 (c, 0.5 in MeOH).

Shigemori, H. *et al.*, *J.O.C.*, 1998, **63**, 6900-6904 (*Brasilicardin A, pmr, cmr, cryst struct*)
Komaki, H. *et al.*, *J. Antibiot.*, 1999, **52**, 13-19; 2000, **53**, 75-77 (*Brasilicardin A, activity*)
Shigemori, H. *et al.*, *Tet. Lett.*, 1999, **40**, 4353-4354 (*biosynth*)
Komatsu, K. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 5545-5551 (*Brasilicardins B-D, cryst struct*)
Komatsu, K. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 1507-1513 (*sar*)

Brasiliidine A **B-281**
3-(2-Isocyano-5-methyl-1,3,5-hexatrienyl)-1-methyl-1H-indole, 9CI
[189827-04-1]




$C_{17}H_{16}N_2$ 248.327
Alkaloid from *Nocardia brasiliensis*. Cy-

tototoxic agent. Amorph. yellow solid. λ_{max} 204 (ϵ 1800); 220 (sh); 268 (ϵ 40000); 332 (ϵ 20000) (EtOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 719-720 (*isol, uv, ir, pmr, cmr*)


Brassicinal B **B-282**
2,3-Dihydro-3-hydroxy-3-methylthiazolo[3,2-a]indole-9-carboxaldehyde, 9CI
[126654-63-5]



$C_{12}H_{11}NO_2S$ 233.29
Exists in CD₃OD as an equilibrium mixt. of the hemiaminal (illus.) and open-chain keto form. Alkaloid from Chinese cabbage *Brassica campestris* ssp. *pekinensis* inoculated with *Pseudomonas cichorii*. Phytoalexin. Mp 169-170°. $[\alpha]_D$ 0. λ_{max} 207 (ϵ 25800); 220 (ϵ 25400); 252 (ϵ 24100); 277 (ϵ 14200); 315 (ϵ 11700) (MeOH) (Berdy).

Monde, K. *et al.*, *Chem. Lett.*, 1990, 209 (*isol, uv, ir, pmr, cmr, ms, struct*)


Brassicinal C **B-283**
Methyl 3-formyl-1H-indole-2-sulfinate, 9CI
[137761-23-0]



$C_{10}H_9NO_3S$ 223.252
Alkaloid from *Brassica oleracea* (cabbage) inoculated with *Pseudomonas cichorii*. Phytoalexin. Mp 150-151°. $[\alpha]_D$ -13.3 (c, 1.04 in MeOH) (natural). $[\alpha]_D$ -231 (c, 0.2 in MeOH) (synthetic). λ_{max} 215 (ϵ 18300); 247 (ϵ 13600); 309 (ϵ 9400) (MeOH).

Monde, K. *et al.*, *Phytochemistry*, 1991, **30**, 2915-2917 (*isol, uv, ir, pmr, cmr, ms*)
Pedras, M.S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1503-1509 (*synth*)
Taniguchi, T. *et al.*, *Org. Biomol. Chem.*, 2008, 4399-4405 (*ecd, vcd, abs config*)

Brassinin **B-284**
Methyl (1H-indol-3-ylmethyl) carbamodithioate, 9CI
[105748-59-2]



$C_{11}H_{12}N_2S_2$ 236.361

Isol. from Chinese cabbage (*Brassica campestris* ssp. *pekinensis*) (Brassicaceae) heads inoculated with *Pseudomonas cichorii*. Phytoalexin. Sol. MeOH, Et₂O, Me₂CO; fairly sol. H₂O. Mp 132–133°. First report of the isol. of sulfur-containing phytoalexins (see also Cyclobrassinin, C-842). λ_{\max} 220 (ε 30000); 280 (ε 6000); 289 (ε 5000) (MeOH) (Derep). λ_{\max} 218 (ε 38300); 236 (ε 87400); 268 (ε 16100); 287 (ε 9070) (MeOH) (Berdy).

N¹-Methoxy: Methoxybrassinin

[105748-60-5]

C₁₂H₁₄N₂O₅ 266.387

Isol. from *Brassica campestris* ssp. *pekinensis* (Brassicaceae) inoculated with *Pseudomonas cichorii*. Phytoalexin. Viscous oil. λ_{\max} 218 (ε 37300); 267 (ε 15800); 287 (sh) (ε 9470); 297 (sh) (ε 5740) (MeOH) (Derep). λ_{\max} 218 (ε 37000); 241 (ε 13000); 267 (ε 15300) (MeOH) (Berdy). λ_{\max} 218 (ε 24500); 267 (ε 16200) (EtOH) (Berdy).

4-Methoxy: 4-Methoxybrassinin

[129602-03-5]

C₁₂H₁₄N₂O₅ 266.387

Isol. from white cabbage (*Brassica oleracea* var. *capitata*) (Brassicaceae) heads inoculated with *Pseudomonas cichorii*. Phytoalexin. Amorph. λ_{\max} 219 (ε 44300); 265 (ε 18500) (MeOH) (Berdy).

Takasugi, M. *et al.*, *Chem. Comm.*, 1986, 1077 (isol, synth, struct, deriv)

Kawasaki, T. *et al.*, *Heterocycles*, 1990, **31**, 1605 (synth, Methoxybrassinin)

Monde, K. *et al.*, *Phytochemistry*, 1990, **29**, 1499 (4-Methoxybrassinin)

Somei, M. *et al.*, *Heterocycles*, 1992, **33**, 77 (synth, Methoxybrassinin)

Yamada, F. *et al.*, *Heterocycles*, 1993, **36**, 2783 (synth)

Kutschy, P. *et al.*, *Tetrahedron*, 1998, **54**, 3549-3566 (synth)


Pedras, M.S.C. *et al.*, *Phytochemistry*, 2000, **53**, 161-176 (rev)

Brassitin

B-285

S-Methyl 1H-indol-3-ylmethylcarbamothioate, 9CI

[113866-42-5]



C₁₁H₁₂N₂OS 220.295

Stress metab. from Japanese radish Daikon (*Raphanus sativus* var. *hortensis*, Brassicaceae), inoculated with *Pseudomonas cichorii*. Amorph. λ_{\max} 218 (ε 36800); 271 (ε 5910); 277 (ε 5910); 287 (ε 4830) (MeOH) (Berdy).

1-Methoxy: Methoxybrassinin

[113900-63-3]

C₁₂H₁₄N₂O₅ 250.321

Isol. from Chinese cabbage (*Brassica campestris* ssp. *pekinensis*, Brassicaceae) inoculated with the bacterium *Pseudomonas cichorii*. Phytoalexin. Cryst. (Et₂O/EtOH). Mp 94–96°.

Takasugi, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 285 (Methoxybrassinin)


Monde, K. *et al.*, *Phytochemistry*, 1995, **39**, 581 (Brassitin)

Kutschy, P. *et al.*, *Tetrahedron*, 1998, **54**, 3549-3566 (synth)

Bravomicin A

B-286

[251094-90-3]



R = -CH(CH₃)CH₂CH₃

C₃₈H₃₁N₃O₁₂ 721.676

Prod. by *Micromonospora polytropa* ATCC 202091. Antibacterial agent. Orange-red powder. Mp 208–210°. [α]_D -25.4 (c, 0.66 in MeOH/CHCl₃). λ_{\max} 250 (log ε 4.46); 281 (sh) (log ε 4.32); 326 (log ε 4.1); 468 (log ε 4.17) (MeOH).

Me ester: Bravomicin B

[251094-91-4]

C₃₉H₃₃N₃O₁₂ 735.703

Prod. by *Micromonospora polytropa* ATCC 202091. Orange-red powder. λ_{\max} 249 (log ε 4.43); 281 (sh) (log ε 4.27); 325 (log ε 4); 467 (log ε 4.14) (MeOH).

Et ester: Bravomicin C

[251094-92-5]

C₄₀H₃₅N₃O₁₂ 749.729

Prod. by *Micromonospora polytropa* ATCC 202091. Orange-red powder. λ_{\max} 250 (log ε 4.44); 281 (sh) (log ε 4.26); 325 (log ε 4.03); 467 (log ε 4.18) (MeOH).

U.S. Pat., 1999, 5 994 543; CA, **132**, 2808n

Bravomicin D

B-287

[251094-93-6]

As Bravomicin A, B-286 with

R = -CH(CH₃)CH₂CH₂CH₃

C₃₉H₃₃N₃O₁₂ 735.703

Prod. by *Micromonospora polytropa* ATCC 202091. Antibacterial agent. Orange-red powder. λ_{\max} 250 (log ε 4.49); 281 (sh) (log ε 4.31); 326 (log ε 4.02); 468 (log ε 4.17) (MeOH).

Me ester: Bravomicin E

[251094-94-7]

C₄₀H₃₅N₃O₁₂ 749.729

Prod. by *Micromonospora polytropa* ATCC 202091. Orange-red powder. λ_{\max} 250 (log ε 4.52); 281 (sh) (log ε 4.32); 325 (log ε 4.03); 469 (log ε 4.18) (MeOH).

Et ester: Bravomicin F

[251094-95-8]

C₄₁H₃₇N₃O₁₂ 763.756


Prod. by *Micromonospora polytropa* ATCC 202091. Orange-red powder. λ_{\max} 249 (log ε 4.56); 280 (sh) (log ε 4.28); 326 (log ε 4.06); 468 (log ε 4.14) (MeOH).

U.S. Pat., 1999, 5 994 543; CA, **132**, 2808n

Braznitidumine

B-288

[926307-39-3]



C₂₄H₃₂N₄O₆ 472.54


Biogenetic origin uncertain. Alkaloid from the stem bark of *Aspidosperma nitidum*. Yellow solid. Mp 272.9–273.1°. Pereira, M.M. *et al.*, *J. Braz. Chem. Soc.*, 2006, **17**, 1274-1280 (isol, pmr, cmr, N-15 nmr)

Brefelamide

B-289

N-[3-[2-Amino-3-(4-hydroxyphenoxy)-phenyl]-3-oxopropyl]-4-hydroxybenzamide, 9CI

[868529-07-1]



C₂₂H₂₀N₂O₅ 392.41


Isol. from *Dictyostelium brefeldianum* and *Dictyostelium giganteum*. Inhibitor of astrocytoma. Yellow oil.

Kikuchi, H. *et al.*, *J.O.C.*, 2005, **70**, 8854-8858 (isol, synth, pmr, cmr)

Brevianamide E

B-290

[23454-27-5]



Absolute Configuration

C₂₁H₂₅N₃O₃ 367.447

Diketopiperazine. Metab. from *Penicillium brevi-compactum* and *Penicillium viridicatum*. Hepatotoxin. Noncryst. [α]_D²⁰ -157 (c, 0.093 in EtOH). λ_{\max} 239 (ε 7500); 296 (ε 2050) (EtOH).

Birch, A.J. *et al.*, *Tetrahedron*, 1970, **26**, 2329-2344 (isol, uv, ir, pmr, biosynth)

Ritchie, R. *et al.*, *Chem. Comm.*, 1975, 611-612 (config)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1981, 959-963 (synth)


Ritchie, R. *et al.*, *Tetrahedron*, 1981, **37**, 4295-4303 (synth)

Schkeryantz, J.M. *et al.*, *J.A.C.S.*, 1999, **121**, 11964-11975 (synth)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (rev, synth, biosynth)

Brevicarine**B-291**

N,1-Dimethyl-9H-pyrido[3,4-*b*]indole-4-butanamine, 9CI. 4-(4-Methylaminobutyl)-β-carboline [25978-39-6]



$C_{17}H_{21}N_3$ 267.373

Alkaloid from *Carex brevicollis* and *Carex parva* (Cyperaceae). Mp 112° (anhyd.). Sesquihydrate:

Cryst. (Me₂CO). Mp 61°.

Hydrochloride (1:2):

Cryst. (EtOH). Mp 195-196°.

Picrate: Mp 210-212° dec.


N-Ac: Mp 154°.

Terent'eva, I.V. et al., *Khim. Prir. Soedin.*, 1969, **5**, 397; 404; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 330; 335 (isol, ms, pmr, struct, synth)

Müller, W.H. et al., *Chem. Ber.*, 1977, **110**, 2424 (synth)

Brevicolline**B-292**

1-Methyl-4-(1-methyl-2-pyrrolidinyl)-9H-pyrido[3,4-*b*]indole, 9CI. 1-Methyl-4-(1-methyl-2-pyrrolidinyl)-β-carboline



(S)-form

$C_{17}H_{19}N_3$ 265.357

(S)-form [20069-02-7]

Main alkaloid of *Carex brevicollis* (Cyperaceae). Active against gram-positive bacteria and fungi. Prisms (MeOH). Mp 232-233°. $[\alpha]_D^{20}$ -164.5 (c, 1.8 in EtOH). λ_{max} 213 (ε 4.54); 232 (ε 4.68); 279 (ε 4.19); 287 (ε 4.39); 337 (ε 4.09); 351 (ε 4.12) (EtOH).

Didehydro: **Dehydrobrevicolline**. Didehydrobrevicolline, 8CI

[31049-73-7]

$C_{17}H_{17}N_3$ 263.341

Alkaloid from *Carex brevicollis* (Cyperaceae). Mp 237-238°. Struct. not clear from abstract, ref. inaccessible.

(±)-form [64364-56-3]

Noncryst.

N-Me: Mp 162°.

Vember, P.A. et al., *Khim. Prir. Soedin.*, 1967, **3**, 249; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 208 (isol, uv, ir, struct)

Terent'eva, I.V. et al., *CA*, 1971, **74**, 50497r (Dehydrobrevicolline)

Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1971, **36**, 3448 (uv, pmr, config)

Lovkova, M.Y. et al., *CA*, 1979, **91**, 1898537y (biosynth)


Towers, G.H.N. et al., *J. Nat. Prod.*, 1983, **46**, 576-581 (activity)

Mahboobi, S. et al., *J. Nat. Prod.*, 1999, **62**, 577-579 (synth, uv, ir, pmr)

Wagner, F.F. et al., *Org. Lett.*, 2006, **8**, 3549-3552 (synth)

Brevicompanine A**B-293**

[215121-46-3]



Absolute Configuration

R = CH₂CH₃

$C_{22}H_{29}N_3O_2$ 367.49

Related to Fructigenine B, F-160. Prod. by *Penicillium brevicompactum*. Plant growth regulator. Amorph. solid. Mp 61-65°. $[\alpha]_D^{20}$ -237.5 (c, 0.7 in EtOH). λ_{max} 210; 245; 303 (EtOH). λ_{max} 210; 245; 303 (MeOH) (Berdy).

Kusano, M. et al., *J.C.S. Perkin I*, 1998, 2823-2826 (isol, uv, ir, pmr, cmr)

Matsumara, K. et al., *Heterocycles*, 2001, **54**, 727-733 (synth)

Brevicompanine C**B-294**

As Brevicompanine A, B-293 with R = CH₃


$C_{21}H_{27}N_3O_2$ 353.463

Prod. by *Penicillium brevicompactum*. Plant growth regulator. Cryst. (EtOAc). Mp 94-96°. $[\alpha]_D^{20}$ -321.7 (c, 0.6 in EtOH). λ_{max} 207 (log ε 3.97); 245 (log ε 3.36); 302 (log ε 3.08) (EtOH).

Kimura, Y. et al., *J. Nat. Prod.*, 2005, **68**, 237-239 (isol, pmr, cmr)

Brevigellin**B-295**


[79432-16-9]



$C_{31}H_{41}N_5O_7$ 595.694

Depsipeptide antibiotic. Prod. by *Penicillium brevicompactum*. Glass, powder or gel. Mp 209-212°.

McCorkindale, N.J. et al., *Tetrahedron*, 1981, **37**, 1795 (isol)

Brevinic acid**B-296**


$C_{14}H_{11}NO_4S$ 289.311

Prod. by *Brevibacterium flavum*. Antibacterial agent. Deep purple cryst. (Et₂O/C₆H₆). Mp 168-172° dec. $[\alpha]_D^{23}$ +1260 (c, 0.0014 in MeOH).

Eguchi, C. et al., *Heterocycles*, Special Issue 2, 1984, **21**, 470 (isol, struct)

Brevioxime**B-297**

[198773-28-3]



Absolute Configuration

$C_{15}H_{22}N_2O_3$ 278.35

Prod. by *Penicillium brevicompactum*. Inhibitor of juvenile hormone biosynth. Cryst. Mp 147-149° (synthetic). $[\alpha]_D^{23}$ -39 (c, 0.24 in CHCl₃) (ca. 17% ee natural). $[\alpha]_D^{22}$ -209 (c, 0.26 in CHCl₃) (96% ee synthetic). λ_{max} 254 (ε 680) (no solvent reported).

Moya, P. et al., *J.O.C.*, 1997, **62**, 8544-8545 (isol, uv, pmr, cmr, ms, cryst struct)

Castillo, M. et al., *Arch. Insect Biochem. Physiol.*, 1998, **37**, 287-294 (activity)

Clive, D.L.J. et al., *J.O.C.*, 2000, **65**, 4923-4929 (synth)


Clark, D. et al., *Tetrahedron*, 2000, **56**, 6181-6184 (synth)

Karadogan, B. et al., *Tetrahedron*, 2001, **57**, 8699-8703 (synth)

Nishimura, Y. et al., *Heterocycles*, 2003, **61**, 481-491 (synth, abs config)

Bripiodionene**B-298**

[190265-70-4]




$C_{15}H_{20}N_2O_4$ 292.334

Tetramic acid deriv. Prod. by *Streptomyces* sp. WC76599. Inhibitor of human cytomegalovirus protease. Powder. Sol. MeOH, DMSO, butanol. Mp 195-198°. $[\alpha]_D$ -260 (c, 0.22 in MeOH). λ_{max} 228 (ε 3630); 328 (ε 13180) (MeOH). λ_{max} 228 (ε 3630); 328 (ε 13182) (MeOH) (Berdy).

Shu, Y.-Z. et al., *J. Nat. Prod.*, 1997, **60**, 529-532 (isol)

6-Bromo-β-carboline**B-299**

6-Bromo-9H-pyrido[3,4-*b*]indole, 9CI. **Eudistomin N**. Bromonorharman [59444-69-8]



$C_{11}H_7BrN_2$ 247.094

Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Mod. active against gram-positive bacteria, yeasts and viruses. Enzyme inhibitor. Yellow needles (MeOH/CHCl₃). Sol. MeOH, C₆H₆, CHCl₃; poorly sol. H₂O. Mp 265-268° dec. λ_{max} 233 (ε 39600); 251 (ε 21000); 283 (ε 9200); 291 (ε 15800); 342 (ε 3800); 357 (ε 3900) (MeOH) (Derep).

λ_{\max} 333; 347; 361; 373 (MeOH) (Berdy).
 λ_{\max} 235 (ϵ 23900); 292 (ϵ 10100); 335 (ϵ 2400); 347 (ϵ 2600) (hexane) (Berdy).

N-Me:

$C_{12}H_9BrN_2$ 261.121

Cryst. (Et₂O/hexane). Mp 108-112°.

Roques, B.P. *et al.*, *Bull. Soc. Chim. Fr.*, 1975, 2403 (*pmr, struct, synth*)

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**, 3378-3387 (*isol, struct, props*)

Ponce, M.A. *et al.*, *J. Het. Chem.*, 2001, **38**, 1087-1095 (*synth, pmr, cmr, ms*)

7-Bromo- β -carboline B-300

7-Bromo-9H-pyrido[3,4-b]indole. **Eudistomin O**
 [88704-40-9]

$C_{11}H_7BrN_2$ 247.094


Alkaloid from *Eudistoma olivaceum* and from the New Zealand ascidian *Ritterella sigillinoides*. Shows mod. activity against bacteria and viruses. Enzyme inhibitor. Pale-yellow cryst. Sol. MeOH, C₆H₆, CHCl₃; poorly sol. H₂O. Mp 208-210°.
 λ_{\max} 333; 347; 361; 373 (MeOH) (Berdy). λ_{\max} 235 (ϵ 23900); 292 (ϵ 10100); 335 (ϵ 2400); 347 (ϵ 2600) (hexane) (Berdy). λ_{\max} 211 (ϵ 15900); 238 (ϵ 24900); 289 (ϵ 8700); 294 (ϵ 11500); 336 (ϵ 2800) (EtOH) (Berdy).

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**, 3378-3387 (*isol, synth*)

Lake, R.J. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1201-1206 (*isol, pmr*)

7-Bromocavernicolone B-301

[98604-31-0]



$C_8H_8BrNO_4$ 262.059

Isol. from the Mediterranean sponge *Aplysina cavernicola* (*Verongia cavernicola*). Mildly antibacterial *in vitro*. Needles (MeOH). Sol. MeOH, CHCl₃, butanol; poorly sol. H₂O. Mp 165-170° dec. No observable opt. rotn. from 589 to 365 nm (MeOH). λ_{\max} 254 (ϵ 2600) (MeOH) (Berdy).

Chloro analogue: **7-Chlorocavernicolone**

[102192-93-8]

$C_8H_8ClNO_4$ 217.608

Metab. from the sponge *Aplysina* (*Verongia*) *cavernicola*. Mp 197-198° dec. No opt. rotn. from 589 to 365 nm. A small excess of one of the enantiomers cannot be ruled out.

D'Ambrosio, M. *et al.*, *Helv. Chim. Acta.*, 1985, **68**, 1453-1460 (7-Bromocavernicolone, *cryst struct*)

D'Ambrosio, M. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1986, **83**, 309-312 (7-Chlorocavernicolone)

Bromochloroacetic acid B-302

[5589-96-8]

ClCHBrCOOH

$C_2H_2BrClO_2$ 173.393

Constit. of the red alga *Asparagopsis taxiformis*.

(\pm)-**form**

Mp 31.5°. Bp 215° dec. Bp₁₁ 103-104°.

Et ester: [22524-32-9]

$C_4H_6BrClO_2$ 201.447

Liq. Bp 174° dec. Bp₁₂ 65°.

Chloride: [38282-29-0]

C_2HBrCl_2O 191.839

Bp 138-139°.

Amide: **Bromochloroacetamide**

[62872-34-8]

$C_2H_3BrClNO$ 172.408

Constit. of *Asparagopsis taxiformis*.

Cryst. Sol. hot H₂O. Mp 126° (117°).

Nitrile: **Bromochloroacetoneitrile**. **Bromochlorocyanomethane**

[83463-62-1]

$C_2HBrClN$ 154.393

Detected in drinking water. Bp 138-140°.

► **Exp. carcinogen. Exp. reprod. effects.**

AL8010000

Compton, H. *et al.*, *J.C.S.*, 1921, 119; 1874 (*synth*)

Backer, H.J. *et al.*, *J.C.S.*, 1928, 2125 (*synth*)

McBee, E.T. *et al.*, *J.A.C.S.*, 1956, **78**, 4595 (*ester*)

Woolard, F.X. *et al.*, *Tetrahedron*, 1976, **32**, 2843-2846 (*amide, isol*)

Woolard, F.X. *et al.*, *Phytochemistry*, 1979, **18**, 617-620 (*isol*)

Oliver, B.G. *et al.*, *Environ. Sci. Technol.*, 1983, **17**, 80 (*nitrile*)

Coleman, W.E. *et al.*, *Environ. Sci. Technol.*, 1984, **18**, 674 (*ms, nitrile*)

IARC Monog., 1991, **52**, 269 (*rev, tox, nitrile*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, BMY800

6-Bromo-3-chloro-1H-indole, B-303

9Cl

[57916-08-2]



C_8H_5BrClN 230.491

Isol. from the yellow hemichordate *Ptychodera flava laysanica* and from *Ptychodera flava*. Mp 55-57° (54.5-56.5°).

Ac: Mp 158-159°.

Higa, T. *et al.*, *Naturwissenschaften*, 1975, **62**, 395-396 (*isol, struct*)


Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 231-233 (*synth, ir, pmr, ms*)

Higa, T. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1980, **65**, 525-530 (*isol*)

7-Bromo-5-(2-chlorophenyl)- B-304

1,3-dihydro-3-hydroxy-2H-1,4-benzodiazepin-2-one, 9Cl

[70030-11-4]



$C_{15}H_{10}BrClN_2O_2$ 365.613


Prod. by *Streptomyces roseochromogenes*. Improbable isolation. A benzodiazepine related to various synthetic drugs.

U.S.S.R. Pat., 1980, 787 027; *C.A.*, **94**, 101315c

9-[3-Bromo-2-(2,3-dibromo- B-305

4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]adenine

[929006-58-6]



$C_{19}H_{14}Br_3N_5O_4$ 616.063

Isol. from *Rhodomela confervoides*.


Brown gum.

Ma, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 337-341 (*isol, pmr, cmr, ms*)

7-[3-Bromo-2-(2,3-dibromo- B-306

4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]-3,7-dihydro-1H-purine-2,6-dione

[929006-56-4]



$C_{19}H_{13}Br_3N_4O_6$ 633.047


Isol. from *Rhodomela confervoides*.

Amorph. solid.

Ma, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 337-341 (*isol, pmr, cmr, ms*)

N-[3-Bromo-2-(2,3-dibromo- B-307

4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]-N'-(3-methoxycarbonylpropyl)urea
Methyl N'-(3-bromo-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl) γ -ureidobutyrate




$C_{20}H_{21}Br_3N_2O_7$ 641.107

Constit. of *Rhodomela confervoides*.

Brown powder. Mp 176-178°.


N^{2-} -(2,3-Dibromo-4,5-dihydroxybenzyl):
 $C_{27}H_{25}Br_2N_2O_9$ 921.023
 Constit. of *Rhodomela confervoides*.
 Brown gum.
 Ma, M. et al., *J. Nat. Prod.*, 2006, **69**, 206-210
 (isol, pmr, cmr, ms)

7-Bromo-2,3-dichloro-1H-indole, 9CI B-308
 [68234-19-5]



$C_8H_4BrCl_2N$ 264.936
 Alkaloid from the marine red alga
Rhodophyllis membranacea. Mp 82-83°.
 Brennan, M.R. et al., *Tet. Lett.*, 1978, **19**,
 1637-1640 (isol, pmr, struct)
 Erickson, K.L. et al., *Synth. Commun.*, 1981,
11, 253-259 (synth, ir, pmr, ms)


6-Bromo-1,2-dihydro-2,2-bis(methylthio)-3H-indol-3-one, 9CI B-309
Tyrindolinone
 [50630-71-2]



$C_{10}H_{10}BrNOS_2$ 304.231
 Alkaloid from the molluscs *Dicathais orbita*, *Nucella lapillus* and *Thais clavigera*. Yellow needles (C_6H_6 /petrol). Mp 117° dec. λ_{max} 225 (sh) (ϵ 21000); 248 (ϵ 25000); 270 (sh) (ϵ 8000); 355 (sh) (ϵ 1600); 402 (ϵ 2700) (EtOH).

Baker, J.T. et al., *Aust. J. Chem.*, 1973, **26**,
 2153-2157 (isol, uv, ir, pmr, ms)
 Cooksey, C.J. et al., *Molecules*, 2001, **6**, 736-
 769 (rev)

6-Bromo-4,5-dihydroxyindole B-310
6-Bromo-1H-indole-4,5-diol, 9CI
 [211808-67-2]



$C_8H_6BrNO_2$ 228.045
 Alkaloid from the gastropod *Drupella fragum*. Amorph. solid.


Ochi, M. et al., *J. Nat. Prod.*, 1998, **61**, 1043-
 1045 (isol)

6-Bromo-4,7-dihydroxyindole B-311
6-Bromo-1H-indole-4,7-diol, 9CI
 [211808-68-3]
 $C_8H_6BrNO_2$ 228.045

Alkaloid from the gastropod *Drupella fragum*. Amorph. solid.

Ochi, M. et al., *J. Nat. Prod.*, 1998, **61**, 1043-
 1045 (isol)


7-Bromo-2,4-dihydroxyquinazoline B-312
7-Bromo-2,4(1H,3H)-quinazolinone, 9CI. 7-Bromo-2,4-quinazolinediol
 [114703-12-7]



$C_8H_5BrN_2O_2$ 241.044
 Alkaloid from the marine tunicate *Pyura sacciformis*. Cryst. (MeOH). Mp 330°.
 λ_{max} 225 (ϵ 47000); 310 (ϵ 4600) (EtOH) (Derep).


Niwa, H. et al., *J. Nat. Prod.*, 1988, **51**, 343-
 344 (isol, uv, ir, pmr, ms, synth, struct)

6-Bromo-2,4-dihydroxyquinoline B-313
6-Bromo-4-hydroxy-2(1H)-quinolinone, 9CI. 6-Bromo-2,4-quinolinediol
 [54675-23-9]



$C_9H_6BrNO_2$ 240.056
 Isol. from the sponge *Hyrtios erecta*.
 Yellow powder. Mp >340°.
 Buckle, D.R. et al., *J. Med. Chem.*, 1975, **18**,
 726-732 (synth)
 Aoki, S. et al., *Chem. Pharm. Bull.*, 2001, **49**,
 1372-1374 (isol, pmr, cmr)


6-Bromo-3a-(3,7-dimethyl-2,6-octadienyl)-1,2,3,3a,8,8a-hexahydro-1-methylpyrrolo[2,3-b]indol-7-ol B-314
 [474779-13-0]



$C_{21}H_{29}BrN_2O$ 405.377
 Alkaloid from the marine bryozoan
Flustra foliacea. Yellow oil. $[\alpha]_D^{22}$ -22 (c,
 0.1 in $CHCl_3$). λ_{max} 251 (log ϵ 4.7); 309
 (log ϵ 4.5) (no solvent reported).

Peters, L. et al., *J. Nat. Prod.*, 2002, **65**, 1633-
 1637 (isol, pmr, cmr, ms)


6-Bromo-2-(1,1-dimethyl-2-propenyl)-1H-indole-3-carboxaldehyde B-315
6-Bromo-3-formyl-2-(1,1-dimethyl-2-propenyl)indole
 [474657-70-0]



$C_{14}H_{14}BrNO$ 292.175
 Alkaloid from the marine bryozoan
Flustra foliacea. Yellow oil. λ_{max} 222 (log
 ϵ 4.6); 274 (log ϵ 4); 300 (log ϵ 3.9)
 (MeOH).

Peters, L. et al., *J. Nat. Prod.*, 2002, **65**, 1633-
 1637 (isol, pmr, cmr, ms)

6-Bromo-2-(1,1-dimethyl-2-propenyl)tryptamine B-316
6-Bromo-2-(1,1-dimethyl-2-propenyl)-1H-indole-3-ethanamine, 9CI. 3-(2-Aminoethyl)-6-bromo-2-(1,1-dimethyl-2-propenyl)-1H-indole. 6-Bromo-2-(1,1-dimethylallyl)tryptamine



$C_{15}H_{19}BrN_2$ 307.233

N^b -Me: *Deformylflustrabromine*
 [474657-72-2]

$C_{16}H_{21}BrN_2$ 321.259
 Isol. from *Flustra foliacea*. Cytotoxic.
 Yellow oil. λ_{max} 231 (log ϵ 4.6); 289
 (log ϵ 3.9) (MeOH). λ_{max} 204 (ϵ
 30190); 230 (ϵ 22730); 282 (ϵ 4770)
 (MeOH).

N^b -Me, N^b -formyl: *Flustrabromine*
 [80693-54-5]

$C_{17}H_{21}BrN_2O$ 349.27
 Alkaloid from the marine bryozoan
Flustra foliacea. Amorph. solid. Consists
 of two rotameric forms reflecting
 hindered rotation around the forma-
 mide C-N bond. Each rotamer also
 exists in equilib. with a form in which
 the amide nitrogen atom associates
 intramolecularly with the benzene ring.


N^b -Me, N^b -(methylsulfonyl): *N-[2-[6-Bromo-2-(1,1-dimethyl-2-propenyl)-1H-indol-3-yl]ethyl]-N-methylmethanesulfonamide*
 [474657-71-1]

$C_{17}H_{23}BrN_2O_2S$ 399.351
 Isol. from *Flustra foliacea*. Yellow oil.
 λ_{max} 227 (log ϵ 4.8); 288 (log ϵ 4.3)
 (MeOH).

Wulff, P. et al., *J.C.S. Perkin 1*, 1981, 2895-
 2898 (*Flustrabromine*, isol, uv, ir, pmr, cmr,
 ms, struct)


Peters, L. et al., *J. Nat. Prod.*, 2002, **65**, 1633-
 1637 (*Deformylflustrabromine*, *N*-Me *N*-
 methylsulfonyl)

Lysek, N. et al., *Z. Naturforsch., C*, 2002, **57**,
 1056-1061 (*Deformylflustrabromine*)

7-Bromo-4-(2-ethoxyethyl)-quinoline, 9CI B-317
[82503-96-6]

$C_{13}H_{14}BrNO$ 280.164
Minor alkaloid from the marine bryozoan *Frustra foliacea*. Possibly an artifact.


Wulff, P. et al., *Comp. Biochem. Physiol. B: Comp. Biochem.*, 1982, **71**, 525-526 (isol, uv, pmr, ms, struct)

6-Bromo-1-ethyl-β-carboline B-318
6-Bromo-1-ethyl-9H-pyrido[3,4-b]indole, 9CI
[113679-47-3]

$C_{13}H_{11}BrN_2$ 275.147
Metab. from the marine hydroid *Aglaophenia pluma*. Cryst. (CH_2Cl_2). Mp 155-157°.

8-Bromo-6,8-Dibromo-1-ethyl-β-carboline
[113679-48-4]
 $C_{13}H_{10}Br_2N_2$ 354.043
Major metab. from *Aglaophenia pluma*. Cryst. (CH_2Cl_2). Mp 150-152°.

Aiello, A. et al., *Tetrahedron*, 1987, **43**, 5929-5932 (isol, uv, pmr, cmr, struct, synth, deriv)

4-Bromo-2-(guanidinomethyl)pyrrole B-319


$C_6H_9BrN_4$ 217.068
Isol. from the sponge *Agelas* sp.

α-Oxo: N-(Aminoiminomethyl)-4-bromo-1H-pyrrole-2-carboxamide

$C_6H_7BrN_4O$ 231.051
Isol. from an *Agelas* sp. Not named in the lit., not listed in CA.

Stempien, M.F. et al., *Abstr. Am. Chem. Soc. Natl. Mtg., MEDI Sect., 164th*, 1972, 21 (α-oxo, isol)

Christophersen, C. et al., *Alkaloids (Academic Press)*, 1985, **24**, 36 (occur)


6-Bromo-3-(hydroxyacetyl)-1H-indole B-320
1-(6-Bromo-1H-indol-3-yl)-2-hydroxyethanone, 9CI
[126235-45-8]

$C_{10}H_8BrNO_2$ 254.083
Alkaloid from the demosponge *Pleroma menouii*. Microcryst. powder (MeOH). Mp 194-196°. λ_{max} 217 (ε 24100); 242 (ε 13100); 265 (ε 12200); 293 (ε 9200) (no solvent reported).

Guella, G. et al., *Z. Naturforsch., C*, 1989, **44**, 914 (isol, uv, ir, pmr, cmr, ms, struct)

5-Bromo-6-hydroxy-β-carboline B-321

5-Bromo-9H-pyrido[3,4-b]indol-6-ol, 9CI. 5-Bromo-6-hydroxy-9H-pyrido[3,4-b]indole. Eudistomin D
[88704-37-4]



$C_{11}H_7BrN_2O$ 263.093
Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Mod. active against gram-positive bacteria and Herpes simplex virus. Yellow amorph. solid. Mp 280°. λ_{max} 215 (ε 14000); 233 (ε 24800); 246 (ε 19800); 286 (ε 8500); 294 (ε 15000); 365 (ε 4500) (MeOH) (Derep).

N²-Me: N²-Methyleudistomin D
[380614-17-5]
 $C_{12}H_{10}BrN_2O^{\oplus}$ 278.128
Quaternary alkaloid from *Eudistoma gilboverde*. Amorph. yellow powder. Counterion not specified. λ_{max} 214 (log ε 4.02); 236 (log ε 3.8); 268 (log ε 3.96); 303 (sh) (log ε 3.71); 314 (log ε 3.79); 416 (log ε 3.19) (EtOH).

N⁹-Me: Eudistomidin D
[125422-19-7]
 $C_{12}H_9BrN_2O$ 277.12
Alkaloid from the Okinawan tunicate *Eudistoma glaucus*. Cytotoxic. Induces Ca^{2+} release from the sarcoplasmic reticulum. Yellow solid. Mp 180° dec. λ_{max} 254 (ε 7000); 299 (ε 8600); 342 (sh); 503 (ε 700) (MeOH/KOH) (Derep). λ_{max} 211 (ε 13600); 270 (ε 9800); 315 (ε 6800); 416 (ε 1800) (MeOH) (Derep).

Kobayashi, J. et al., *J.A.C.S.*, 1984, **106**, 1526-1528 (isol, uv, pmr, struct)

Rinehart, K.L. et al., *J.A.C.S.*, 1987, **109**, 3378-3387 (isol, struct, props)

Kobayashi, J. et al., *J.O.C.*, 1990, **55**, 3666-3670 (*Eudistomidin D*)

Rocca, P. et al., *Tet. Lett.*, 1995, **36**, 7085-7088 (synth)

Rashid, M.A. et al., *J. Nat. Prod.*, 2001, **64**, 1454-1456 (*N²-Methyleudistomin D*)

7-Bromo-6-hydroxy-β-carboline B-322

7-Bromo-9H-pyrido[3,4-b]indol-6-ol, 9CI. Eudistomin J
[88704-38-5]

$C_{11}H_7BrN_2O$ 263.093
Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Shows modest antibiotic activity. Calcium release indu-

cer. λ_{max} 215 (ε 14000); 233 (ε 24800); 246 (ε 19800); 286 (ε 8500); 294 (ε 15000); 365 (ε 4500) (MeOH) (Derep).


N²-Me: N²-Methyleudistomin J
[380614-18-6]

$C_{12}H_{10}BrN_2O^{\oplus}$ 278.128

Quaternary alkaloid from *Eudistoma gilboverde*. Yellow gum. λ_{max} 212 (log ε 4.04); 236 (log ε 3.99); 271 (log ε 3.97); 307 (sh) (log ε 3.83); 321 (log ε 3.96); 412 (log ε 3.35) (EtOH).


Rinehart, K.L. et al., *J.A.C.S.*, 1987, **109**, 3378-3387 (isol)

Rashid, M.A. et al., *J. Nat. Prod.*, 2001, **64**, 1454-1456 (*N²-Methyleudistomin J*)

3-Bromo-4-hydroxy-α-(hydroxyimino)-N-[2-(1H-imidazol-4-yl)ethyl]benzenepropanamide B-323
Pseudoceratina crassa Alkaloid
[165329-94-2]

$C_{14}H_{15}BrN_4O_3$ 367.201
Alkaloid from the Caribbean sponge *Pseudoceratina crassa*.

Ciminiello, P. et al., *J. Nat. Prod.*, 1995, **58**, 689 (isol, uv, ir, pmr, cmr, struct)

5-Bromo-3-hydroxy-1H-indole B-324
5-Bromo-1H-indol-3-ol. 5-Bromo-1,2-dihydro-3H-indol-3-one. 5-Bromo-3-indolone

C_8H_6BrNO 212.045

OH-form

O-β-D-Galactopyranoside: Z-gal
[97753-82-7]

$C_{14}H_{16}BrNO_6$ 374.188

Chromogenic substrate for β-galactosidase.

O-Ac: [17357-14-1]

$C_{10}H_8BrNO_2$ 254.083

Stout needles (MeOH). Mp 187-188°.

N-Ac: [106698-07-1]

$C_{10}H_8BrNO_2$ 254.083

Cryst. (EtOH aq.). Mp 184-185°.

Di-Ac: [33588-54-4]

$C_{12}H_{10}BrNO_3$ 296.12

Needles (MeOH aq./Ac₂O). Mp 125-128°.

Oxo-form

Oxime: [76983-85-2]

$C_8H_7BrN_2O$ 227.06

Brown needles (C_6H_6). Mp 233-235°.

Su, H.C.F. et al., *J.A.C.S.*, 1960, **82**, 1187-1189 (*O-Ac, N-Ac*)

Torii, S. *et al.*, *J.O.C.*, 1978, **43**, 2882 (*di-Ac*)
 Aguzzi, A. *et al.*, *Histochemistry*, 1994, **102**,
 477-481 (*O-galactopyranosyl, use*)

6-Bromo-3-hydroxy-1H-indole B-325

6-Bromo-1H-indol-3-ol, 9CI, 8CI. 6-Bromo-1,2-dihydro-3H-indol-3-one. 6-Bromo-3-indolone. 6-Bromoindoxyl
 [114224-27-0]
 [52578-60-6 (oxo-form)]

C₈H₆BrNO 212.046

Constit. of egg masses of *Dicathais orbita*.

N-Ac: [244256-22-2]

C₁₀H₈BrNO₂ 254.083
 Solid. Mp 187°.

O-Ac: [114306-17-1]

C₁₀H₈BrNO₂ 254.083
 Solid (EtOH aq.). Mp 200°.

Di-Ac: [108996-91-4]

C₁₂H₁₀BrNO₃ 296.12
 Solid (EtOH). Mp 123°.

Holt, S.J. *et al.*, *J.C.S.*, 1958, 1217-1223 (*N-Ac*)
 Holt, S.J. *et al.*, *Proc. R. Soc. London, B*, 1958,
148, 481-494 (*O-Ac, di-Ac*)

Pat. Coop. Treaty (WIPO), 1999, 99 50 538;
CA, **131**, 243537h (*N-Ac, di-Ac, synth, pmr, cmr*)

Benkendorff, K. *et al.*, *Molecules*, 2001, **6**, 70-78 (*isol*)

6-Bromo-5-hydroxy-1H-indole B-326

6-Bromo-1H-indol-5-ol
 [211808-66-1]


C₈H₆BrNO 212.046

Alkaloid from the gastropod *Drupella fragum*. Shows antioxidant props.
 Amorph. solid.

Ochi, M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1043-1045 (*isol*)

6-Bromo-5-hydroxy-1H-indole-3-carboxaldehyde B-327

6-Bromo-3-formyl-5-hydroxyindole
 [151833-94-2]




C₉H₆BrNO₂ 240.056

Isol. from sponge *Oceanapia bartschi*.
 λ_{\max} 255 (ϵ 5800) (no solvent reported).

Cafieri, F. *et al.*, *Z. Naturforsch., B*, 1993, **48**,
 1408-1410

6-Bromo-3-hydroxy-2-(methylthio)indole B-328

6-Bromo-2-(methylthio)-1H-indol-3-ol. Tyrindoxol
 [142960-47-2]



C₉H₈BrNOS 258.138

Biosynthetic precursor of 6,6'-Dibromoindigotin, D-325 and 6,6'-Dibromoindirubin in I-73.

O-Sulfate: Tyroxindoxyl sulfate

[74626-31-6]

C₉H₈BrNO₄S₂ 338.203

Constit. of *Dicathais orbita*, *Mancinella keineri*, *Murex* spp. and other molluscs. Biosynthetic precursor of

6,6'-Dibromoindigotin, D-325 and 6,6'-Dibromoindirubin in I-73. Yellow leaflets + 1H₂O (as Ag salt). Mp 118-120° dec. (Ag salt).

S,S-Dioxide, O-sulfate:

C₉H₈BrNO₆S₂ 370.201

Isol. from *Murex trunculus*, *Murex erinaceum*, other *Murex* spp. and *Purpura haemastoma*. Probable biosynthetic precursor of Tyrian Purple.

Baker, J.T. *et al.*, *Tet. Lett.*, 1968, 43-46; 1976,
 1233-1234 (*isol, pmr, struct*)


Fouquet, H. *et al.*, *Angew. Chem., Int. Ed.*,
 1971, **10**, 816-817 (*isol*)

Prota, G. *et al.*, *Marine Natural Products*, (ed.
 Scheuer, P.J.), Academic Press, 1980, **3**, 141-178 (*biosynth, rev*)

Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-769 (*rev*)

[2-(3-Bromo-4-hydroxyphenyl)ethyl]ethanediamide, 9CI B-329

[182496-50-0]



C₁₀H₁₁BrN₂O₃ 287.113

Isol. from the Papua New Guinea marine sponge *Ianthella basta*. Amorph. solid.
 λ_{\max} 212 (log ϵ 3.89); 281 (log ϵ 3.14) (MeOH).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1996, **59**, 927-934 (*isol, uv, ir, pmr, cmr, ms, struct*)

3-Bromo-1H-indole, 9CI B-330

[1484-27-1]

C₈H₆BrN 196.046

Trace odorous component of the marine hemichordate *Ptychodera flava laysanica*. Also isol. from sponge *Oceanapia bartschi*. Cryst. (heptane or EtOH aq.). Mp 66-67° (65-66° dec.).

N-(4-Methylbenzenesulfonyl): [90481-77-9]

C₁₅H₁₂BrNO₂S 350.235
 Needles (EtOH). Mp 119-120°.

Pappalardo, G. *et al.*, *Gazz. Chim. Ital.*, 1958,
88, 1147-1169 (*uv*)

Jardine, R.V. *et al.*, *Can. J. Chem.*, 1963, **41**,
 2067-2073; 2399-2401 (*synth, pmr*)

Higa, T. *et al.*, *Naturwissenschaften*, 1975, **62**,
 395-396 (*isol*)


Brennan, M.P. *et al.*, *Heterocycles*, 1986, **24**,
 2879-2885 (*synth, ir, pmr, cmr*)

Cafieri, F. *et al.*, *Z. Naturforsch., B*, 1992, **48**,
 1408-1410 (*isol, sponge*)

Hodson, H.F. *et al.*, *Tetrahedron*, 1994, **50**,
 1899-1906 (*N-tosyl*)

5-Bromo-1H-indole-3-acetic acid B-331

[40432-84-6]



C₁₀H₈BrNO₂ 254.083

Constit. of a sponge *Smenospongia* sp.
 Mp 143-145°.

Szewczyk, H. *et al.*, *Rocz. Chem.*, 1972, **46**,
 2135

Tasdemir, D. *et al.*, *Z. Naturforsch., C*, 2002,
57, 914-922 (*isol, pmr, cmr, ms*)

6-Bromo-1H-indole-3-acetic acid B-332

C₁₀H₈BrNO₂ 254.083

Me ester: Methyl 6-bromo-1H-indole-3-acetate, 9CI

[152213-63-3]

C₁₁H₁₀BrNO₂ 268.109

Alkaloid from the marine sponge *Pseudosuberites hyalinus*. Oil. λ_{\max} 228 (log ϵ 4.49) (MeOH).

Amide: 6-Bromo-1H-indole-3-acetamide, 9CI

[152213-62-2]

C₁₀H₉BrN₂O 253.098

Alkaloid from the marine sponge *Pseudosuberites hyalinus*. Mp 148-150°.

Nitrile: 6-Bromo-1H-indole-3-acetonitrile, 9CI. 6-Bromo-3-(cyanomethyl)indole

[152213-61-1]

C₁₀H₇BrN₂ 235.083

Alkaloid from the marine sponge *Pseudosuberites hyalinus*. Mp 114-115°.

Rasmussen, T. *et al.*, *J. Nat. Prod.*, 1993, **56**,
 1553-1558 (*isol, pmr, cmr, ir, uv, ms*)

6-Bromo-1H-indole-3-carboxaldehyde, 9CI B-333

6-Bromo-3-formylindole. 6-Bromoindole-3-carbaldehyde

[17826-04-9]

C₉H₆BrNO 224.056

Isol. from the marine tunicate *Pyura sacciformis*; also from *Pseudosuberites hyalinus*, *Pleroma menoui*, *Tabastrea cocinea*, *Dendrophyllia* spp. and sponges *Oceanapia bartschi*, *Halichondria* spp. and others. From *Acinetobacter* sp. associated with the ascidian *Stomozoa murrayi*. Exhibits antibacterial and antifouling activity; plant growth regulator; algicide. Solid (MeOH). Mp 203-204°.

N-(4-Methylbenzenesulfonyl): [158991-81-2]

C₁₆H₁₂BrNO₂S 378.246

Solid or cryst. (CHCl₃). Mp 115° Mp 146-147°.

Da Settimo, A. *et al.*, *Gazz. Chim. Ital.*, 1967,
97, 1304-1316 (*synth*)

Wratten, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1977, **11**, 411-414 (*isol*)


Dellar, G. *et al.*, *J.C.S. Perkin I*, 1981, 1679-1680 (*synth, pmr*)

- Okuda, R.K. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 1907-1914 (*isol*)
 Schmidt, V. *et al.*, *Annalen*, 1985, 1882-1894 (*synth*)
 Niwa, H. *et al.*, *J. Nat. Prod.*, 1988, **51**, 343-344 (*isol*)
 Guella, G. *et al.*, *Helv. Chim. Acta*, 1989, **72**, 1444-1450 (*isol, hplc*)
 Cafieri, F. *et al.*, *Z. Naturforsch., B*, 1992, **48**, 1408-1410 (*isol, sponge*)
 Jiang, B. *et al.*, *J.O.C.*, 1994, **59**, 6823-6827 (*synth, ir, pmr, ms*)
 Olguin-Urbe, G. *et al.*, *J. Chem. Ecol.*, 1997, **23**, 2507-2521 (*isol, activity*)
 Jiang, B. *et al.*, *J.O.C.*, 2001, **66**, 4865-4869 (*tosyl, synth, ir, pmr, cmr, ms*)

6-Bromo-1H-indole-3-carboxylic acid, 9CI

B-334

[101774-27-0]

C₉H₆BrNO₂ 240.056

Early lit. mistakenly reports physical data for both acid and Et ester which actually refer to inseparable equimolar mixt. of 5- and 6-bromo isomers. *Isol.* from the marine sponge *Pseudosuberites hyalinus*.

*Me ester:*C₁₀H₈BrNO₂ 254.083

Alkaloid from *Smenospongia* sp. Yellow solid.

Et ester: [103858-55-5]C₁₁H₁₀BrNO₂ 268.109

Alkaloid from the demosponge *Pleroma menoui* and the sponge *Ietrochota birotulata*. Microcryst. powder (MeOH). Mp 147-149°.

Leggetter, B.E. *et al.*, *Can. J. Chem.*, 1960, **38**, 1467 (*synth*)

Guella, G. *et al.*, *Z. Naturforsch., C*, 1989, **44**, 914-916 (*Et ester, isol*)

Rasmussen, T. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1553-1558 (*isol, ms, pmr, cmr*)


Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1484-1488 (*Me ester*)

6-Bromo-1H-indole-2,3-dione, 9CI

B-335

6-Bromoisatin

[6326-79-0]

C₈H₄BrNO₂ 226.029

Isol. from many molluscs. Plant growth regulator. Probably a precursor of 6,6'-Dibromoindirubin. Needles, yellow plates (AcOH). Mp 272.5°. pK_a 10.35.

Inagaki, S. *et al.*, *Yakugaku Zasshi*, 1938, **58**, 961-975; *CA*, **33**, 2516 (*synth*)

Sadler, P.W. *et al.*, *J.O.C.*, 1956, **21**, 169-170 (*synth*)


Budesinsky, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 3473-3483 (*synth*)

Clark, R.J.H. *et al.*, *New J. Chem.*, 1999, **23**, 323-328 (*pmr*)

Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-769 (*rev*)

3-(6-Bromo-1H-indol-3-yl)-2-propenoic acid, 9CI

B-336

C₁₁H₈BrNO₂ 266.094*(E)-form**Penaresin*

[135250-41-8]

Alkaloid from the marine sponge *Penares* sp. Powerful Ca-inducer in sarcoplasmic reticulum. Pale yellow solid. Mp 189-192° dec. λ_{max} 228 (ε 27100); 280 (ε 18400); 307 (ε 18300) (MeOH).

Me ester: [134419-23-1]

[78520-47-5]

C₁₂H₁₀BrNO₂ 280.12

Alkaloid from the marine sponges *Hymeniacion* sp., *Ietrochota* sp. and *Mycale adhaerens*. Clusters of prisms (Et₂O/hexane). Mp 186°. λ_{max} 226 (ε 34200); 283 (ε 19900); 324 (ε 26200) (MeOH) (Berdy).

Et ester: [226888-27-3]C₁₃H₁₂BrNO₂ 294.147

Alkaloid from the sponge *Hymeniacion* sp. Cryst. (Me₂CO/petrol) (synthetic). Mp 147-149° (synthetic).

*(Z)-form**Me ester:* [419567-80-9]Alkaloid from *Hymeniacion* sp.*Et ester:* [419567-79-6]Alkaloid from *Hymeniacion* sp.

Dellar, G. *et al.*, *J.C.S. Perkin 1*, 1981, 1679-1680 (*isol, ir, pmr, ms, struct, synth, Me ester*)

Kobayashi, J. *et al.*, *Heterocycles*, 1990, **31**, 2205-2208 (*isol, uv, ir, pmr, ms, struct*)

Fusetani, N. *et al.*, *J.O.C.*, 1991, **56**, 4971-4974 (*isol, Me ester*)

Beugelmans, R. *et al.*, *Tetrahedron*, 1999, **55**, 5089-5112 (*Et ester, synth*)

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2002, **65**, 368-370 (*Me esters, Et esters, isol*)

Bromiodoacetic acid

B-337

[71815-43-5]

BrCHICOOH

C₂H₂BrIO₂ 264.845

Constit. of the red alga *Asparagopsis taxiformis*.

Et ester: [62874-49-1]C₄H₆BrIO₂ 292.899

Constit. of *Asparagopsis taxiformis* and *Falkenbergia rufolonosa*.

Amide: Bromiodoacetamide

[62872-36-0]

C₂H₃BrINO 263.86

Constit. of *Asparagopsis taxiformis*.

Woolard, F.X. *et al.*, *Tetrahedron*, 1976, **32**, 2843-2846 (*amide, isol*)

Combaut, G. *et al.*, *Phytochemistry*, 1978, **17**, 1661-1663 (*isol*)


Woolard, F.X. *et al.*, *Phytochemistry*, 1979, **18**, 617-620 (*isol, ms*)

3'-Bromo-4-methoxy-2,2'-bipyrrole-5-carboxaldehyde

B-338

3-Bromo-5'-formyl-4-methoxy-2,2'-bipyrrrole

[85849-99-6]

C₁₀H₉BrN₂O₂ 269.097

Isol. from the nembrothid nudibranchs *Roboastra tigris*, *Tambja eliora*, *Tambja abdere* and the bryozoan *Sessibugula translucens*. Mp 243-245° dec. Artifact.

Carté, B. *et al.*, *J.O.C.*, 1983, **48**, 2314 (*isol, uv, ir, pmr, struct*)

5'-Bromo-4-methoxy-2,2'-bipyrrole-5-carboxaldehyde

B-339

5'-Bromo-5-formyl-4-methoxy-2,2'-bipyrrrole

[85849-98-5]

C₁₀H₉BrN₂O₂ 269.097

Isol. from the nembrothid nudibranchs *Roboastra tigris*, *Tambja eliora*, *Tambja abdere* and the bryozoan *Sessibugula translucens*. Mp 235-238° dec. Artifact.


Carté, B. *et al.*, *J.O.C.*, 1983, **48**, 2314 (*isol, uv, ir, pmr, struct*)

6-Bromo-1-methyl-β-carboline

B-340

6-Bromo-1-methyl-9H-pyrido[3,4-b]indole

[18813-71-3]

C₁₂H₉BrN₂ 261.12

Metab. from the marine hydroid *Aglao-phenia pluma*. Cryst. (CH₂Cl₂). Mp 248-250°.

Aiello, A. *et al.*, *Tetrahedron*, 1987, **43**, 5929-5932 (*isol, uv, pmr, cmr, struct, synth*)


Ponce, M.A. *et al.*, *J. Het. Chem.*, 2001, **38**, 1087-1095 (*synth, pmr, cmr, ms*)

6-Bromo-2-(methylthio)-3H-indol-3-one, 9CI

B-341

Tyrindoleninone

[50630-70-1]



C₉H₆BrNOS 256.123

Alkaloid from the molluscs *Dicathais orbita*, *Nucella lapillus*, *Thais clavigera* and *Murex brandaris*. Also occurs in egg masses of *Agnestia tritoniformis*, *Lepsiella reticularis*, *Morula marginalba* and other spp. Biosynthetic precursor of Tyrian Purple 6,6'-Dibromoindigotin, D-325. Red needles. Mp 109.5°. λ_{\max} 217 (€ 8700); 242 (€ 25000); 247 (€ 25000); 278 (€ 11000); 318 (€ 5200); 333 (€ 5000); 355 (€ 2700); 426 (€ 1700); 447 (€ 1700) (hexane).

S-Oxide:C₉H₆BrNO₂S 272.122

Isol. from egg masses of *Dicathais orbita* and *Ceratostoma erinaceum*. CAS no. not found to 2007.

S,S-Dioxide:C₉H₆BrNO₃S 288.121

Doubtless present in molluscs. Putative precursor of Tyrian Purple 6,6'-Dibromoindigotin, D-325. CAS no. not found to 2007.

Baker, J.T. et al., *Aust. J. Chem.*, 1973, **26**, 2153-2157 (*isol, uv, ir, pmr, ms*)

Baker, J.T. et al., *Endeavour*, 1974, **33**, 11-17 (*rev*)


Prota, G. et al., *Marine Natural Products*, (ed. Scheuer, P.J.), Academic Press, 1980, **3**, 141-178 (*biosynth, rev*)

Benkendorff, K. et al., *J. Chem. Ecol.*, 2000, **26**, 1037-1050 (*isol*)

Benkendorff, K. et al., *Molecules*, 2001, **6**, 70-78 (*occur, S-oxide, S,S-dioxide*)

6-Bromo- α -oxo-1H-indole-3-acetic acid B-342

(6-Bromo-1H-indol-3-yl)glyoxylic acid

C₁₀H₆BrNO₃ 268.066**Me ester:** [220407-33-0]C₁₁H₈BrNO₃ 282.093

Alkaloid from *Spongosorites* sp. Amorph. yellow powder.

Et ester: [17826-12-9]C₁₂H₁₀BrNO₃ 296.12

Alkaloid from the sponge *Ietrochota birotulata*. Pale yellow solid. λ_{\max} 270; 320 (MeOH).

Amide: 6-Bromo- α -oxo-1H-indole-3-acetamide

[108061-75-2]

C₁₀H₇BrN₂O₂ 267.082


Alkaloid from *Spongosorites* sp. Amorph. yellow powder. λ_{\max} 212 (log € 3.21); 258 (log € 2.73); 275 (log € 2.75); 320 (log € 2.61) (MeOH).

Li, L. et al., *Pharmazie*, 2003, **58**, 680-681 (*Et ester*)

Bao, B. et al., *Mar. Drugs*, 2007, **5**, 31-39 (*Me ester, amide*)

4-Bromophakellin

Monobromophakellin

C₁₁H₁₂BrN₅O 310.153**(-)-form** [31955-05-2]

Alkaloid from the marine sponge *Phakellia flabellata*. Mp 170-180° dec. λ_{\max} 227 (€ 7140); 277 (€ 5540) (as hydrochloride) (Derep). λ_{\max} 228 (€ 7780); 275 (€ 6110) (MeOH) (Derep).

Hydrochloride: Mp 215-220°. $[\alpha]_{D}^{25}$ -123 (c, 3 in MeOH).

N⁷-Me: 4-Bromo-N⁷-methylphakellin.**N⁷-Methylmonobromophakellin**

[697284-91-6]

C₁₂H₁₄BrN₅O 324.18

Isol. from an *Agelas* sp. Powder. $[\alpha]_{D}^{28}$ -35 (c, 1 in MeOH). λ_{\max} 238 (€ 1700); 280 (€ 1280) (MeOH).

5-Bromo: (-)-Dibromophakellin

[31954-96-8]

C₁₁H₁₁Br₂N₅O 389.049

Alkaloid from *Phakellia flabellata*, *Phakellia mauritiana* and *Pseudaxinyssa cantharella*. Sol. MeOH, butanol. Mp 237-245° dec. λ_{\max} 233 (€ 8880); 281 (€ 8810) (MeOH) (Derep). λ_{\max} 211 (€ 104000); 237 (€ 10700); 285 (€ 11000) (EtOH) (Berdy).

5-Bromo; hydrochloride: Mp 220-221°. $[\alpha]_{D}^{25}$ -205 (c, 2.9 in MeOH).**5-Bromo, N⁷-Me: N⁷-Methyldibromophakellin**

[697284-90-5]

C₁₂H₁₃Br₂N₅O 403.076

Isol. from an *Agelas* sp. Powder. $[\alpha]_{D}^{28}$ -19.8 (c, 1.4 in MeOH). λ_{\max} 238 (€ 6780); 280 (€ 5430) (MeOH).

(+)-form**5-Bromo: (+)-Dibromophakellin**C₁₁H₁₁Br₂N₅O 389.049

Metab. of the New Caledonian sponge *Pseudaxinyssa cantharella*. Mp 220° dec. $[\alpha]_{D}^{20}$ +159 (c, 1.55 in MeOH). λ_{\max} 233 (€ 8880); 281 (€ 8810) (MeOH) (Derep).

Sharma, G. et al., *J.O.C.*, 1977, **42**, 4118-4124 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Foley, L.H. et al., *J.A.C.S.*, 1982, **104**, 1776-1777 (*synth*)

Utkina, N.K. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 547-548 (*Dibromophakellin, isol*)

De Nanteuil, G. et al., *Tetrahedron*, 1985, **41**, 6019-6033 (*(+)-Dibromophakellin*)

Gautschi, J.T. et al., *J. Nat. Prod.*, 2004, **67**, 1256-1261 (*N⁷-Methylphakellins*)

Atodiresei, I. et al., *Chirality*, 2007, **19**, 542-549 (*Dibromophakellin, electronic cd*)


Wang, S. et al., *Angew. Chem., Int. Ed.*, 2008, **47**, 1284-1286 (*synth*)

Meyer, S.W. et al., *J. Nat. Prod.*, 2008, **71**, 1524-1529 (*pmr, cmr, N-15 nmr*)

B-343


N-(4-Bromophenyl)-2,2-diphenylacetamide B-344

N-(4-Bromophenyl)- α -phenylbenzeneacetamide, 9CI. p-Bromo- α , α -diphenylacetanilide [70481-06-0]

C₂₀H₁₆BrNO 366.257

Alkaloid from *Arundo donax*. Needles (EtOAc/petrol). Mp 203-205°.

Stevens, C.L. et al., *J.O.C.*, 1964, **29**, 34 (*synth*)
Miles, D.H. et al., *J. Nat. Prod.*, 1993, **56**, 1590-1593 (*isol, pmr, cryst struct*)


N²-(4-Bromo-1H-pyrrole-2-carbonyl)arginine B-345C₁₁H₁₆BrN₅O₃ 346.183

Isol. from *Stylyssa caribica*. Light yellow oil. $[\alpha]_{D}^{23}$ -16 (c, 0.25 in MeOH).

Grube, A. et al., *J. Nat. Prod.*, 2006, **69**, 125-127 (*isol, synth, pmr, cmr*)

N²-(4-Bromo-1H-pyrrole-2-carbonyl)homoarginine B-346


N⁶-(Aminoiminomethyl)-N²-[(4-bromo-1H-pyrrol-2-yl)carbonyl]lysine, 9CI [242147-75-7]

C₁₂H₁₈BrN₅O₃ 360.21

Isol. from *Agelas wiedenmayeri*. λ_{\max} 271 (€ 11500) (H₂O).


Assmann, M. et al., *Org. Lett.*, 1999, **1**, 455-457 (*isol, struct*)

Lindel, T. et al., *J. Nat. Prod.*, 2000, **63**, 1566-1569 (*synth, uv, ir, pmr, cmr*)

N⁶-(4-Bromo-1H-pyrrole-2-carbonyl)lysine B-347C₁₁H₁₆BrN₃O₃ 318.17

Isol. from *Stylyssa caribica*. Light yellow oil. $[\alpha]_{D}^{23}$ +5.2 (c, 0.5 in MeOH).

Grube, A. et al., *J. Nat. Prod.*, 2006, **69**, 125-127 (*isol, synth, pmr, cmr*)

4-Bromo-1H-pyrrole-2-carboxylic acid, 9CI
[27746-02-7]

$C_5H_4BrNO_2$ 189.996
Isol. from *Axinella verrucosa*. Mp 160° dec. pK_a 4.06 (25°, H_2O).

Me ester: [934-05-4]
 $C_6H_6BrNO_2$ 204.023
Isol. from sponges *Axinella tenuidigitata* and *Lissodendoryx* sp. Mp 106-106.5° (98-99°). Struct. of nat. prod. revised in 1993.

Et ester: [433267-55-1]
 $C_7H_8BrNO_2$ 218.05
Solid. Mp 55°.

Amide: [196106-96-4]
 $C_5H_5BrN_2O$ 189.011
Isol. from the sponge *Acanthella carteri*. λ_{max} 202 (ϵ 13800); 230 (ϵ 9800); 267 (ϵ 16200) (MeOH).

(*Methoxymethyl*)*amide*: 4-Bromo-N²-methoxymethyl-1H-pyrrole-2-carboxamide

$C_7H_9BrN_2O_2$ 233.064
Alkaloid from *Axinella verrucosa*.
Anderson, H.J. *et al.*, *Can. J. Chem.*, 1965, **43**, 409-414 (*synth*)
Fringuelli, F. *et al.*, *Tetrahedron*, 1969, **25**, 5815-5818 (*synth*)
Bélanger, P. *et al.*, *Tet. Lett.*, 1979, **20**, 2505-2508 (*Me ester*)
Barrow, R.A. *et al.*, *Nat. Prod. Lett.*, 1993, **1**, 243-250 (*Me ester, struct*)
Kitamura, C. *et al.*, *J.C.S. Perkin 1*, 1997, 1443-1447 (*Me ester*)
Mancini, I. *et al.*, *Tet. Lett.*, 1997, **38**, 6271-6274 (*amide, isol, uv, pmr, cmr, ms*)
Reddy, N.S. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 1145-1147 (*Me ester, isol*)
Bergauer, M. *et al.*, *Synthesis*, 2002, 274-278 (*Et ester*)
Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 17-24 (*methoxymethylamide, isol*)

5-Bromo-1H-pyrrole-2-carboxylic acid, 9CI
[28383-57-5]

$C_5H_4BrNO_2$ 189.996
Mp 138° dec. (block). pK_a 4.17 (25°, H_2O).

Me ester Methyl 5-bromo-2-pyrrolecarboxylate
[934-07-6]
 $C_6H_6BrNO_2$ 204.023


Isol. from the sponge *Lissodendoryx* sp. Cryst. (hexane). Mp 106.5-108°.

Amide: 5-Bromo-1H-pyrrole-2-carboxamide
[17543-94-1]
 $C_5H_5BrN_2O$ 189.011
Alkaloid from the sponge *Agelas nakamurai*. Needles. Mp 137-139°. λ_{max} 213 ($\log \epsilon$ 3.47); 268 ($\log \epsilon$ 3.93) (MeOH).

(*Methoxymethyl*)*amide*: 5-Bromo-N²-


(*methoxymethyl*)-1H-pyrrole-2-carboxamide
[213619-56-8]
 $C_7H_9BrN_2O_2$ 233.064
Alkaloid from *Agelas nakamurai*.
Amorph. solid. λ_{max} 214 ($\log \epsilon$ 3.6); 271 ($\log \epsilon$ 4.05) (MeOH).

Anderson, H.J. *et al.*, *Can. J. Chem.*, 1965, **43**, 409-414 (*synth*)
Hodge, P. *et al.*, *J.C.S.*, 1965, 459-470 (*synth, uv, ir*)
Friguelli, F. *et al.*, *Tetrahedron*, 1969, **25**, 5815-5818 (*synth*)
Schmitz, F.J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 47-53 (*Me ester, isol*)
Iwagawa, T. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1310-1312 (*isol, amides*)

3-Bromo-1H-pyrrole-2,5-dione B-350
2-Bromomaleimide
[45514-47-4]


$C_4H_2BrNO_2$ 175.969
Alkaloid from the marine sponge *Axinella brevistyla*. Antifungal and cytotoxic agent. λ_{max} 201 ($\log \epsilon$ 3.8); 229 ($\log \epsilon$ 3.7); 245 (sh) ($\log \epsilon$ 3.5) (MeOH).

N-Ph: [72000-67-0]
 $C_{10}H_6BrNO_2$ 252.067
Pale yellow solid (EtOH). Mp 150-153°.
Choi, D.S. *et al.*, *J.O.C.*, 1998, **63**, 2646-2655 (*N-Ph, synth*)
Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1576-1578 (*isol, uv, ir, pmr, cmr*)

5-Bromo-7H-pyrrolo[2,3-d]pyrimidine B-351
[175791-49-8]

$C_6H_4BrN_3$ 198.022
Isol. from a *Hypnea* sp. Scales. Mp 199-201°. CAS no. refers to 1H-form.
Naqvi, S.W.A. *et al.*, *Bot. Mar.*, 1981, **24**, 51-55 (*isol*)
Sakamoto, T. *et al.*, *J.C.S. Perkin 1*, 1996, 459 (*synth, pmr, ir*)

6-Bromo-1,2,3,4-tetrahydro-7,8-dihydroxy-3-isoquinoline-carboxylic acid B-352



$C_{10}H_{10}BrNO_4$ 288.097

(*S*)-*form*
7-Me ether: 6-Bromo-1,2,3,4-tetrahydro-8-hydroxy-7-methoxy-3-isoquinoline-carboxylic acid
 $C_{11}H_{12}BrNO_4$ 302.124

7-Me ether, Me ester: [929006-62-2]
 $C_{12}H_{14}BrNO_4$ 316.151
Alkaloid from *Rhodomela confervoides*. Brown gum. $[\alpha]_D^{20}$ -29 (c, 0.07 in MeOH). λ_{max} 206 ($\log \epsilon$ 4.53); 275 ($\log \epsilon$ 3.11) (MeOH).


Ma, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 337-341 (*isol, pmr, cmr, ms*)

8-Bromo-1,2,3,4-tetrahydro-6,7-dihydroxy-3-isoquinoline-carboxylic acid B-353
 $C_{10}H_{10}BrNO_4$ 288.097

(*S*)-*form*
7-Me ether: 8-Bromo-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-3-isoquinoline-carboxylic acid
[929006-60-0]
 $C_{11}H_{12}BrNO_4$ 302.124
Alkaloid from *Rhodomela confervoides*. Powder. $[\alpha]_D^{20}$ -99.1 (c, 0.11 in MeOH). λ_{max} 207 ($\log \epsilon$ 4.49); 287 ($\log \epsilon$ 3.41) (MeOH).


7-Me ether, Me ester: [929006-61-1]
 $C_{12}H_{14}BrNO_4$ 316.151
Alkaloid from *Rhodomela confervoides*. Brown gum. $[\alpha]_D^{20}$ -49.6 (c, 0.07 in MeOH). λ_{max} 205 ($\log \epsilon$ 4.56); 282 ($\log \epsilon$ 3.53) (MeOH).

Ma, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 337-341 (*isol, pmr, cmr, ms*)

4(7)-Bromo-2,3,7(2,3,4)-trichloro-1H-indole, 9CI B-354
[68124-92-5]

$C_8H_3BrCl_3N$ 299.381
Alkaloid from the marine red alga *Rhodophyllis membranacea*.

Brennan, M.R. *et al.*, *Tet. Lett.*, 1978, 1637-1640 (*isol, pmr, struct*)

5-Bromotryptamine B-355
5-Bromo-1H-indole-3-ethanamine, 9CI. 3-(2-Aminoethyl)-5-bromoindole, 8CI
[3610-42-2]

$C_{10}H_{11}BrN_2$ 239.114
Radioprotective agent. Tryptophan, T-640 antagonist, 5-Hydroxytryptamine,

H-755 antimetabolite displaying vasoconstructive and embryotoxic activity. Amber oil. Biosynth. precursor of Eudistomin H, E-288.

Hydrochloride: [81868-12-4]

Cryst. (EtOH/Et₂O) or solid. Mp 288-289° (286-287°).

N-Ac:

C₁₂H₁₃BrN₂O 281.151
Solid. Mp 153.4-153.8°.

N^b-Hydroxy: 5-Bromo-N-hydroxy-1H-indole-3-ethanamine. 5-Bromo-N^b-hydroxytryptamine

[123475-32-1]
C₁₀H₁₁BrN₂O 255.114

Pale beige resinous material.

N^b, N^b-Di-Me: 5-Bromo-N^b, N^b-dimethyltryptamine. 5-Bromo-N,N-dimethyl-1H-indole-3-ethanamine, 9CI. 5-Bromo-3-(2-dimethylaminoethyl)indole

[17274-65-6]
C₁₂H₁₅BrN₂ 267.168

Metab. of the marine sponge *Smenospongia aurea* and from *Polyfibrospongia echina*. Major alkaloid from the New Caledonian ascidian *Eudistoma fragum*. Antimicrobial compd. Cryst. (MeOH aq.). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 98-99° (90-92°). λ_{max} 225 (ε 39000); 285 (ε 4400); 305 (ε 2700) (MeOH) (Derep). λ_{max} 227 (ε 24000); 282 (ε 5500); 290 (ε 7943); 300 (ε 5012) (EtOH) (Berdy).

Quadbeck, G. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1954, **297**, 229; *CA*, **51**, 15795i (pharmacol)

Ho, B.-T. et al., *J. Pharm. Sci.*, 1969, **58**, 563 (pharmacol)

Djura, P. et al., *J.O.C.*, 1980, **45**, 1435-1441 (5-Bromo-N,N-dimethyltryptamine, isol, uv, ir, pmr, cmr, struct)

Landau, M.A. et al., *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1981, **69**, 359 (tox)

Tymiak, A.A. et al., *Tetrahedron*, 1985, **41**, 1039-1047 (5-Bromo-N,N-dimethyltryptamine, isol, uv, ir, pmr, ms, synth)

Kennaway, D.J. et al., *Aust. J. Biol. Sci.*, 1988, **41**, 393 (synth)

Debitus, C. et al., *J. Nat. Prod.*, 1988, **51**, 799-801 (5-Bromo-N,N-dimethyltryptamine, isol, uv, pmr, cmr, ms)

Still, I.W. et al., *Can. J. Chem.*, 1990, **68**, 1408-1419 (synth, ir, pmr, cmr, ms)

Shen, G.Q. et al., *Tet. Lett.*, 1994, **35**, 4923-4926 (biochem)

Somei, M. et al., *Heterocycles*, 1995, **40**, 119-122 (synth, 5-Bromo-N,N-dimethyltryptamine)

6-Bromotryptamine B-356

6-Bromo-1H-indole-3-ethanamine, 9CI. 3-(2-Aminoethyl)-6-bromoindole

[96624-18-9]

C₁₀H₁₁BrN₂ 239.114
Alkaloid from the encrusting grey marine tunicate *Didemnum candidum*. Mp 120-120.5°. λ_{max} 225 (ε 15800); 286 (ε 3100); 294 (ε 2700) (MeOH) (Berdy).

Hydrochloride: [108061-77-4]

Mp 215-217° dec.

N^b-Me, N^b-formyl: 6-Bromo-N^b-formyl-N^b-methyltryptamine. N-[2-(6-Bromo-1H-indol-3-yl)ethyl]-N-methylforma-

mide, 9CI

[82503-95-5]

C₁₂H₁₃BrN₂O 281.151

Alkaloid from the marine bryozoan *Flustra foliacea*. Amorph. solid. Shows restricted C-N bond rotn.

N^a-(3-Methyl-2-butenyl), N^b-Me: 6-Bromo-N^b-methyl-N^a-prenyltryptamine.

Deformylflustrabromine B

C₁₆H₂₁BrN₂ 321.259

Alkaloid from *Flustra foliacea*. Yellow oil. λ_{max} 230 (log ε 4.1); 291 (log ε 3.3) (MeOH).

Wulff, P. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1982, **71**, 523-524 (N^b-Me N^b-formyl, isol, pmr, ms, struct)


Groen, C. et al., *Acta Chem. Scand., Ser. B*, 1984, **38**, 709-711 (synth, pmr, cmr, ir, uv)

Rinehart, K.L. et al., *J.A.C.S.*, 1987, **109**, 3378-3387 (synth, hplc, pmr, cmr, ms, uv, ir)

Fahy, E. et al., *J. Nat. Prod.*, 1991, **54**, 564-569 (isol, pmr, cmr)

Peters, L. et al., *Planta Med.*, 2004, **70**, 883-886 (Bromomethylprenyltryptamine)

Oceanapia Bromotyrosine B-357 alkaloid



C₁₇H₂₆Br₂N₆O₃ 522.239

Isol. from the sponge *Oceanapia* sp. Inhibitor of mycothiol S-conjugate amidase. Related to Puralidin H, P-811.

Nicholas, G.M. et al., *Org. Lett.*, 2001, **3**, 1543-1545 (isol, pmr, cmr)


Kende, A.S. et al., *Tet. Lett.*, 2004, **45**, 133-135 (synth)

Chanda, B.M. et al., *Tet. Lett.*, 2005, **46**, 6461-6463 (synth)

Broussonetine B-358

Dihydro-4,5-bis(8-hydroxy-4-quinolinyl)-2(3H)-furanone, 9CI. 3,4-Bis(8-hydroxy-4-quinolinyl)-γ-butyrolactone

[92664-84-1]



C₂₂H₁₆N₂O₄ 372.379

Alkaloid from the wood of *Broussonetia zeylanica* (Moraceae). Cryst. (CHCl₃). Mp 238-239°. [α]_D 0 (c, 0.11 in DMF).


Di-Ac:

Cryst. (C₆H₆/petrol). Mp 162-163°.

Gunatilaka, A.A. et al., *Phytochemistry*, 1984, **23**, 929 (isol, uv, ir, pmr, cmr, ms, struct)

Broussonetine I B-359

[215117-10-5]



C₂₀H₃₈N₂O₆ 402.53

Alkaloid from *Broussonetia kazinoki*. Oil. [α]_D +2.9 (c, 0.26 in MeOH).

N¹-Ac: Broussonetine J

[215117-17-2]

C₂₂H₄₀N₂O₇ 444.567

Alkaloid from *Broussonetia kazinoki*. Oil. [α]_D +2.1 (c, 0.7 in MeOH).

N-De-Ac: Broussonetine J₂

C₁₈H₃₆N₂O₅ 360.493

Alkaloid from *Broussonetia kazinoki*. Oil. [α]_D +13.8 (c, 0.42 in MeOH).

N-De-Ac, N¹-Ac: Broussonetine J₁

C₂₀H₃₈N₂O₆ 402.53


Alkaloid from *Broussonetia kazinoki*. Oil. [α]_D -17.5 (c, 0.3 in MeOH).

Shibano, M. et al., *Chem. Pharm. Bull.*, 1998, **46**, 1416-1420 (isol, ir, pmr, cmr)

Tsukamoto, D. et al., *Chem. Pharm. Bull.*, 2001, **49**, 1487-1491 (Broussonetines J₁, J₂)

Shibano, M. et al., *Heterocycles*, 2002, **57**, 1539-1553 (biosynth)

Broussonetine G B-360



Absolute Configuration


C₁₈H₃₃NO₆ 359.462

Alkaloid from *Broussonetia kazinoki*. Glycosidase inhibitor. Oil. [α]_D +17.5 (c, 0.4 in MeOH).

Shibano, M. et al., *Chem. Pharm. Bull.*, 1998, **46**, 1048-1050 (isol, ir, pmr, cmr, ms)

Trost, B.M. et al., *Chem. Eur. J.*, 2006, **12**, 6607-6620 (synth, abs config)

Broussonetine H B-361




Absolute Configuration

C₁₈H₃₃NO₆ 359.462

Alkaloid from *Broussonetia kazinoki*. Glycosidase inhibitor. Oil. [α]_D +15.5 (c, 0.5 in MeOH).

Shibano, M. et al., *Chem. Pharm. Bull.*, 1998, **46**, 1048-1050 (isol, ir, pmr, cmr, ms)

Broussonetine N B-362




Absolute Configuration

C₁₈H₃₃NO₆ 359.462

Alkaloid from *Broussonetia kazinoki*.
Glycosidase inhibitor. Oil. $[\alpha]_D +5.4$ (c, 0.17 in MeOH).

Shibano, M. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 907-908


Broussonetine R**B-363**

Absolute Configuration

$C_{18}H_{31}NO_6$ 357.446

Alkaloid from *Broussonetia kazinoki*. Oil. $[\alpha]_D +21.8$ (c, 0.27 in MeOH). λ_{max} 247 (log ϵ 3.98) (MeOH).

Tsukamoto, D. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 492-496

Broussonetine U**B-364**

Absolute Configuration


$C_{18}H_{33}NO_5$ 343.462

Alkaloid from *Broussonetia kazinoki*. Oil. $[\alpha]_D -33.3$ (c, 0.2 in MeOH).

Tsukamoto, D. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 492-496 (isol, pmr, cmr)

Broussonetine W**B-365**

[386702-83-6]



Absolute Configuration

$C_{18}H_{31}NO_4$ 325.447

Alkaloid from *Broussonetia kazinoki*. Oil. $[\alpha]_D +16$ (c, 0.07 in MeOH). λ_{max} 236 (log ϵ 3.59) (MeOH).

4-Epimer, 3-O- β -D-glucopyranoside:

Broussonetine X

[386702-84-7]


$C_{24}H_{41}NO_9$ 487.589

Alkaloid from *Broussonetia kazinoki*. Oil. $[\alpha]_D +13.7$ (c, 0.51 in MeOH). λ_{max} 236 (log ϵ 3.96) (MeOH).

Tsukamoto, D. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1487-1491

Broussonetine Z**B-366**

[647025-61-4]



Relative Configuration


$C_{18}H_{33}NO_7$ 375.461

Alkaloid from the branches of *Broussonetia kazinoki*. $[\alpha]_D^{25} +25.9$ (c, 0.57 in MeOH).

Tsukamoto, D. *et al.*, *Nat. Med. (Tokyo)*, 2003, **57**, 68-72 (isol, pmr, cmr)

Broussonetine A**B-367**

[173220-06-9]



$C_{18}H_{35}NO_5$ 345.478

Alkaloid from *Broussonetia kazinoki* (Moraceae). Inhibits glycosidases. Powder. Mp 126-128°. $[\alpha]_D +25.8$ (c, 0.6 in MeOH).

3-O- β -D-Glucopyranoside: **Broussonetine A**

[173220-07-0]

$C_{24}H_{45}NO_{10}$ 507.62

Alkaloid from *Broussonetia kazinoki* (Moraceae). Powder. Mp 164-166°. $[\alpha]_D +32.1$ (c, 0.2 in MeOH).

4,13'-Di-O- β -D-glucopyranoside: **Broussonetine Q**

$C_{30}H_{55}NO_{15}$ 669.762

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +8.8$ (c, 0.25 in MeOH).

10'-Deoxo, 9'-oxo: **Broussonetine B**

[190317-93-2]

$C_{18}H_{35}NO_5$ 345.478

Alkaloid from *Broussonetia kazinoki* (Moraceae). Inhibits glycosidases. Powder. Mp 128-130°. $[\alpha]_D +15.3$ (c, 0.3 in MeOH).

10'-Deoxo, 9'-oxo, 3-O- β -D-glucopyranoside: **Broussonetine B**

[190317-46-5]

$C_{24}H_{45}NO_{10}$ 507.62

Alkaloid from *Broussonetia kazinoki* (Moraceae). Powder. Mp 154-156°. $[\alpha]_D +29.8$ (c, 0.4 in MeOH).

10'-Deoxo, 9'-oxo, 3',4'-didehydro(E-):

Broussonetine V

$C_{18}H_{33}NO_5$ 343.462

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +10.9$ (c, 0.09 in MeOH).

4-Epimer: **Broussonetine C**

[173727-05-4]

$C_{18}H_{35}NO_5$ 345.478

Alkaloid from *Broussonetia kazinoki*. Powder. Mp 147-149°. $[\alpha]_D +32$ (c, 0.96 in MeOH) (+25).

4-Epimer, 10'S-alcohol: **Broussonetine M**

$C_{18}H_{37}NO_5$ 347.494

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +5.9$ (c, 0.3 in MeOH).

4-Epimer, 3',4'-didehydro(E-): **Broussonetine O**

$C_{18}H_{33}NO_5$ 343.462

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +22.7$ (c, 0.37 in MeOH).

4-Epimer, 10'-deoxo, 9'-oxo: **Broussonetine D**

[173220-08-1]

$C_{18}H_{35}NO_5$ 345.478

Alkaloid from *Broussonetia kazinoki*. Powder. Mp 136-138°. $[\alpha]_D +22.9$ (c, 0.31 in MeOH).

4-Epimer, 10'-deoxo, 9'-oxo, 3',4'-didehydro(E-): **Broussonetine P**

$C_{18}H_{33}NO_5$ 343.462

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +28.8$ (c, 0.96 in MeOH).

4-Epimer, 1'R-hydroxy: **Broussonetine E**

[190317-56-7]

$C_{18}H_{35}NO_6$ 361.478

Alkaloid from *Broussonetia kazinoki* (Moraceae). Powder. Mp 103-105°. $[\alpha]_D +4.9$ (c, 1 in MeOH).

4-Epimer, 1'R-hydroxy, 13'-O- β -D-glucopyranoside: **Broussonetine K**

[230285-77-5]

$C_{24}H_{45}NO_{11}$ 523.62

Alkaloid from *Broussonetia kazinoki*. β -Glucosidase inhibitor. Powder. $[\alpha]_D^{25} +20.1$ (c, 0.26 in MeOH).

4-Epimer, 1'R-hydroxy, 10'S-alcohol: **Broussonetine S**

$C_{18}H_{37}NO_6$ 363.493

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +25.1$ (c, 0.18 in MeOH).

4-Epimer, 1'R-hydroxy, 10'-deoxo, 9'-oxo: **Broussonetine F**

[190317-57-8]

$C_{18}H_{35}NO_6$ 361.478

Alkaloid from *Broussonetia kazinoki* (Moraceae). Powder. Mp 105-107°. $[\alpha]_D +13.5$ (c, 0.9 in MeOH).

4-Epimer, 1'R-hydroxy, 10'-deoxo, 9'-oxo, 13'-O- β -D-glucopyranoside: **Broussonetine L**

[230285-78-6]

$C_{24}H_{45}NO_{11}$ 523.62

Alkaloid from *Broussonetia kazinoki*. β -Glucosidase inhibitor. Powder. $[\alpha]_D^{25} +21.3$ (c, 0.56 in MeOH).

4-Epimer, 9'R-hydroxy, 10'R-alcohol: **Broussonetine Y**

[647025-60-3]

$C_{18}H_{37}NO_6$ 363.493

Alkaloid from *Broussonetia kazinoki*. $[\alpha]_D^{25} +32.5$ (c, 0.16 in MeOH).

4-Epimer, 9'R-hydroxy, 10'-deoxo: **Broussonetine M₁**

$C_{18}H_{37}NO_5$ 347.494

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D +18.3$ (c, 0.56 in MeOH).

4-Epimer, 1'R,5'S-dihydroxy: **Broussonetine T**

$C_{18}H_{35}NO_7$ 377.477

Alkaloid from *Broussonetia kazinoki*. Oil. $[\alpha]_D +11$ (c, 0.49 in MeOH).

4-Epimer, 1'R,5'ξ,7'ξ,12'ξ-tetrahydroxy, 10'-deoxo: 3,4-Dihydroxy-2-(hydroxymethyl)-5-(1,5,7,12,13-pentahydroxytridecyl)pyrrolidine

[710948-63-3]

$C_{18}H_{37}NO_8$ 395.492

Alkaloid from the bulbs of *Scilla peruviana*. Glucosidase inhibitor. Syrup. $[\alpha]_D +20.5$ (c, 0.65 in H₂O).

3,5-Diepimer, 1,2-didehydro: **Broussonetine U₁**

$C_{18}H_{33}NO_5$ 343.462

Alkaloid from *Broussonetia kazinoki*. Powder. $[\alpha]_D -30.2$ (c, 0.09 in MeOH).


Shibano, M. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 505-508; 700-705; 1998, **46**, 1048-1050; 1416-1420; 1999, **47**, 472-476; 907-908; 2000, **48**, 1281-1285 (isol, ir, pmr, cmr, abs config, activity)

Yoda, H. *et al.*, *Tet. Lett.*, 1999, **40**, 1335-1336 (*Broussonetine C*, synth)

Tsukamoto, D. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 492-496; 1487-1491 (*Broussonetines S,T,V,M₁,U₁*)

- Shibano, M. *et al.*, *Heterocycles*, 2002, **57**, 1539-1553 (*biosynth*)
 Perlmutter, P. *et al.*, *J. Carbohydr. Chem.*, 2003, **22**, 719-732 (*Broussonetine C*, *synth*, *pmr*, *cmr*)
 Asano, N. *et al.*, *J. Nat. Prod.*, 2004, **67**, 846-850 (*Scilla peruviana constit*)
 Tsukamoto, D. *et al.*, *Nat. Med. (Tokyo)*, 2004, **57**, 68-72 (*Broussonetine Y*)

Bruceolline D **B-368**
 [66278-53-3]




$C_{13}H_{13}NO$ 199.252
 Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Needles (MeOH). Mp 160-162° dec.

10-Oxo: Bruceelline E
 [159903-54-5]
 $C_{13}H_{11}NO_2$ 213.235
 Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Pale yellow needles (MeOH). Mp 289-291° dec.

Ouyang, Y. *et al.*, *Phytochemistry*, 1994, **37**, 575 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Brugine **B-369**
 8-Methyl-8-azabicyclo[3.2.1]oct-3-yl 1,2-dithiolane-3-carboxylate, 9CI
 [14912-30-2]




$C_{12}H_{19}NO_2S_2$ 273.42
 Alkaloid from *Bruguiera sexangula*, *Bruguiera cylindrica* and *Crossostylis multiflora* (Rhizophoraceae). Noncryst. $[\alpha]_D^{24}$ -24 (CHCl₃). $[\alpha]_D^{24}$ +265 (EtOH).

Loder, J.W. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1271 (*isol*, *struct*, *ms*, *pmr*, *ord*)
 Kato, A. *et al.*, *Phytochemistry*, 1975, **14**, 1458 (*isol*)
 Gnecco Medina, D.H. *et al.*, *J. Nat. Prod.*, 1983, **46**, 398 (*isol*)

Brunfelsine **B-370**
 Struct. unknown. Alkaloid from roots of *Brunfelsia hopeana* (Solanaceae).
 Costa, O. de A. *et al.*, *Bol. Assoc. Bras. Pharm.*, 1933, **14**, 295-299; *CA*, **27**, 5889

Brunonine **B-371**
 [106982-88-1]



$C_{22}H_{33}NO_3$ 359.508
 Alkaloid from the whole plant of *Delphinium brunonianum* (Ranunculaceae). Mp 208-209°. $[\alpha]_D^{21}$ +174 (c, 0.067 in EtOH).

15-Epimer: Spiramine N
 [137760-60-2]
 $C_{22}H_{33}NO_3$ 359.508
 Alkaloid from the roots of *Spiraea japonica* var. *acuminata* (Rosaceae). $[\alpha]_D^{20}$ +44.4 (c, 1 in CHCl₃). Has (*R*)-config. at C-19.

15-Epimer, O-de-Et: 19-O-Deethylspiramine N
 $C_{20}H_{29}NO_3$ 331.454
 Alkaloid from *Spiraea japonica* var. *ovalifolia*. Amorph. $[\alpha]_D^{23}$ +18 (c, 0.26 in MeOH).

15-Epimer, O-de-Et, O¹⁹-Me: Spiramine O
 [137760-61-3]
 $C_{21}H_{31}NO_3$ 345.481
 Alkaloid from roots of *Spiraea japonica* var. *acuminata* (Rosaceae).
 Deng, W. *et al.*, *Heterocycles*, 1986, **24**, 869 (*pmr*, *cmr*, *struct*)
 Hao, X. *et al.*, *Chin. Chem. Lett.*, 1992, **3**, 427-430 (*Spiramine N*)
 Hao, X.-J. *et al.*, *Yunnan Zhiwu Yanjiu*, 1994, **16**, 301; *CA*, **122**, 101603s (*Spiramine O*)
 Zuo, G.Y. *et al.*, *Heterocycles*, 2001, **55**, 487-493 (*O-Deethylspiramine N*)

Brunsdonnine **B-372**
 [1354-80-9]

$C_{18}H_{21}NO_5$ 331.368
 Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Brunsdonna tubergenii* (Amaryllidaceae). Cryst. (Me₂CO/MeOH). Mp 253°. $[\alpha]_D^{23}$ +75 (c, 0.2 in CHCl₃).

Hydroiodide: Mp 248° dec.
Perchlorate: Mp 232° dec.
Methiodide: Mp 280-281° dec.


Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 159 (*isol*)

Brunsvinine **B-373**
 [1354-82-1]

$C_{17}H_{19}NO_4$ 301.341
 Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Brunsvigia cooperi* (Amaryllidaceae). Mp 202°.

Picrate: Mp 67-69°.
 Dry, L.J. *et al.*, *J.C.S.*, 1958, 4701-4704 (*isol*)


Brussalexin A **B-374**
 S-(1H-Indol-3-ylmethyl) 2-propenylthiocarbamate. S-(1H-Indol-3-ylmethyl) allylthiocarbamate
 [940005-53-8]



$C_{13}H_{14}N_2OS$ 246.332
 Constit. of *Brassica oleracea* var. *gemmifera* (Brussels sprouts). Phytoalexin. Antifungal agent.

Pedras, M.S.C. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1167-1169 (*isol*, *synth*, *pmr*, *cmr*)

Bubbialine **B-375**
Bubbialidine
 [138230-52-1, 138230-51-0]



Relative Configuration

$C_{12}H_{15}NO_3$ 221.255
 Bubbialine was the (+)-form and Bubbialidine the (-)-form.

(+)-form [131989-89-4]
 Alkaloid from the leaves of *Zygogynum pauciflorum* (Winteraceae). Cryst. (Et₂O). Mp 144-145°. $[\alpha]_D$ +129 (c, 1 in MeOH).


2-Epimer: Epibubbialine. Prenirurine
 [182209-98-9]
 [138230-52-1]
 $C_{12}H_{15}NO_3$ 221.255
 Alkaloid from *Phyllanthus amarus*. Cryst. powder. Mp 107-108°. $[\alpha]_D$ -14.5.

7-Epimer: Isobubbialine
 [182209-97-8]
 $C_{12}H_{15}NO_3$ 221.255
 Alkaloid from *Phyllanthus amarus*. Cryst. powder. Mp 112-113°. $[\alpha]_D$ +11.7.

2,7-Diepimer: Niruroidine
 [175777-84-1]
 $C_{12}H_{15}NO_3$ 221.255
 Alkaloid from *Phyllanthus niruroides*. Amorph. solid.

(-)-form [132075-02-6]
 Alkaloid from *Zygogynum pauciflorum*. Amorph. $[\alpha]_D$ -85 (c, 1.47 in MeOH).
 Ahond, A. *et al.*, *J. Nat. Prod.*, 1990, **53**, 875-881 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cd*, *struct*)
 Magnus, P. *et al.*, *Tetrahedron*, 1993, **49**, 8059-8072 (*synth*, *pmr*, *cmr*)
 Babady-Bida, *et al.*, *Phytochemistry*, 1996, **41**, 1441-1443 (*Niruroidine*)
 Houghton, P.J. *et al.*, *Phytochemistry*, 1996, **43**, 715-717 (*Epibubbialine*, *Isobubbialine*)

Buchananine **B-376**
 6-O-Nicotinoyl- α -D-glucopyranose
 [70802-12-9]




$C_{12}H_{15}NO_7$ 285.253
 Alkaloid from the stems of *Cryptolepis buchanani* (Asclepiadaceae). *C. buchanani* has been used in Indian medicine for treatment of rickets. Cryst. (MeOH aq.). Mp 137-138°. $[\alpha]_D^{20}$ +38.4 (c, 3.08 in H₂O). λ_{max} 221 (ϵ 8510); 258 (sh) (ϵ 2690); 262 (ϵ 2820); 271 (sh) (ϵ 2340) (EtOH) (Derep).

Dutta, S.K. *et al.*, *Phytochemistry*, 1978, **17**, 2047 (*uv, ir, pmr, ms, isol, struct*)

Buchapine**B-377**

3-(1,1-Dimethyl-2-propenyl)-3-(3-methyl-2-butenyl)-2,4(1H,3H)-quinoline-dione, 9CI. 3-(1,1-Dimethylallyl)-3-(3,3-dimethylallyl)-1,2,3,4-tetrahydro-2,4-quinolinedione [84017-97-0]



$C_{19}H_{23}NO_2$ 297.396

Alkaloid from the aerial parts of *Haplophyllum bucharicum* and from *Haplophyllum tuberculatum* (Rutaceae). Also isol. from *Evodia roxburghiana*. Exhibits modest anti-HIV activity against HIV-1 in cultured human lymphoblastoid CEM-SS cells. Mp 134-135° (121-123°).

Nesmelova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 532; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 507 (*isol, struct*)


Bellino, A. *et al.*, *Heterocycles*, 1986, **24**, 1821 (*synth, pmr*)

Sheriha, G.M. *et al.*, *Phytochemistry*, 1987, **26**, 3339 (*isol, uv, ir, pmr, cmr, ms, struct*)

McCormick, J.L. *et al.*, *J. Nat. Prod.*, 1996, **59**, 469 (*isol*)

Bucharaine**B-378**

[21059-47-2]



$C_{19}H_{25}NO_4$ 331.411

Alkaloid from *Haplophyllum bucharicum* (Rutaceae). Mp 151-152°.

N-Me: Mp 142-143°.


Fayzudinova, Z.Sh. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 239; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 232

Grundon, M.F. *et al.*, *Alkaloids (Academic Press)*, 1979, **17**, 173 (*rev*)

Grundon, M.F. *et al.*, *J.C.S. Perkin I*, 1981, 633 (*synth*)

Bucharamine**B-379**

[38485-09-5]




$C_{22}H_{29}NO_4$ 371.475

Alkaloid from *Haplophyllum bucharicum* (Rutaceae). Cryst. (Me₂CO). Mp 223°. Presumably artifact arising from Me₂CO during cryst.

Ubaidullaev, K. *et al.*, *Khim. Prir. Soedin.*, 1972, 343; *Chem. Nat. Compd. (Engl. Transl.)*, 337 (*isol, uv, ms, struct*)

Bucharidine**B-380**

4-Hydroxy-3-[1-(tetrahydro-5-hydroxy-2,6,6-trimethyl-2H-pyran-2-yl)ethyl]-2(1H)-quinolinone, 9CI [25865-94-5]




$C_{19}H_{25}NO_4$ 331.411

Alkaloid from *Haplophyllum bucharicum* (Rutaceae). Mp 251-252°.

Faizudinova, Z.S. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 455; 1970, **6**, 239; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 380; 1970, **6**, 232 (*synth*)

Buchtienine**B-381**

[167875-14-1]




$C_{31}H_{34}N_4O_2$ 494.635

Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae).

Azoug, M. *et al.*, *Phytochemistry*, 1995, **39**, 1223 (*isol, uv, ir, pmr, cmr, ms, struct*)

Buddamine**B-382**

[98260-41-4]



$C_{10}H_{15}NO_3$ 197.233


Alkaloid from the aerial parts of *Buddleia davidii* (Buddlejaceae). Cryst. (Me₂CO/MeOH/petrol). Mp 112°. Prob. artifact. Prev. reported as C₁₀H₁₅NO₅ with incorr. struct.

Röder, E. *et al.*, *Planta Med.*, 1985, 164 (*isol, ir, pmr, cmr, ms*)

Stermitz, F.R. *et al.*, *Tet. Lett.*, 1985, **26**, 5251 (*struct*)

Budmunchiamine A**B-383**

1,9,13-Trimethyl-8-undecyl-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI [139750-76-8]



n = 10

Absolute Configuration

$C_{27}H_{56}N_4O$ 452.766

Alkaloid from the seeds of *Albizzia amara* (Fabaceae). Yellow oil. [α]_D -16.3 (c, 1 in CHCl₃).

N⁹-De-Me: **Budmunchiamine F**

[143051-87-0]

$C_{26}H_{54}N_4O$ 438.739

Isol. from seeds of *Albizzia amara*. Yellowish solid. [α]_D -6.2 (c, 1 in MeOH). Obt. as inseparable mixt. with Budmunchiamine G to which props. refer.

8'-Oxo, N¹,N⁹,N¹³-tri-de-Me: **Budmunchiamine L3**

[165467-48-1]

$C_{24}H_{48}N_4O_2$ 424.669

Isol. from seeds of *Albizzia lebbek*. Viscous liq.

Pezzuto, J.M. *et al.*, *Heterocycles*, 1991, **32**, 1961 (*isol*)

Mar, W. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1531 (*props*)

Pezzuto, J.M. *et al.*, *Phytochemistry*, 1992, **31**, 1795 (*Budmunchiamine F*)

Misra, L.N. *et al.*, *Phytochemistry*, 1995, **39**, 247 (*Budmunchiamine L3*)

Popaj, K. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 180-186 (*synth*)

Detterbeck, R. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1742-1748 (*synth, pmr, cmr, abs config*)

Budmunchiamine B**B-384**

1,9,13-Trimethyl-8-nonyl-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI [139750-77-9]

As Budmunchiamine A, B-383 with n = 8

$C_{25}H_{52}N_4O$ 424.712


Alkaloid from the seeds of *Albizzia amara* (Fabaceae).

Pezzuto, J.M. *et al.*, *Heterocycles*, 1991, **32**, 1961 (*isol*)

Popaj, K. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 180-186 (*synth*)

Budmunchiamine L1**B-385**

8-Hexadecyl-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI [165561-01-3]



R = $-(CH_2)_{15}CH_3$

C₂₉H₆₀N₄O 480.819

Isol. as an inseparable mixt. with Budmunchiamine L2, B-386, to which props. refer. Alkaloid from seeds of *Albizia lebbek*. Viscous liq.

16'-Hydroxy: **Budmunchiamine L4†**

[195734-28-2]

C₂₉H₆₀N₄O₂ 496.819

Alkaloid from the seeds of *Albizia lebbek*. Viscous liq. $[\alpha]_D^{27} +13.5$ (c, 0.02 in MeOH). Not to be confused with Budmunchiamine L4, B-387. The same name has been applied to two different homologues. λ_{\max} 208 (log ϵ 4.01) (MeOH).

Misra, L.N. et al., *Phytochemistry*, 1995, **39**, 247-249 (*Budmunchiamine L1, isol, ir, pmr, cmr, ms, struct*)

Dixit, A.K. et al., *J. Nat. Prod.*, 1997, **60**, 1036-1037 (*Budmunchiamine LA*)

Budmunchiamine L2 B-386

8-Tetradecyl-1,5,9,13-tetraazacycloheptadecan-6-one

[165467-47-0]

As Budmunchiamine L1, B-385 with R = ⁻¹CH₂(CH₂)₁₂CH₃

C₂₇H₅₆N₄O 452.766

Isol. as an inseparable mixt. with Budmunchiamine L1, B-385, to which props. refer. Alkaloid from seeds of *Albizia lebbek*. Viscous liq.

5',6'Z,8',9'Z,11',12'Z-Hexadehydro: 8-(5,8,11-Tetradecatrienyl)-1,5,9,13-tetraazacycloheptadecan-6-one. **Felipealbizine A**

[288094-80-4]

C₂₇H₅₀N₄O 446.718

Alkaloid from the leaves of *Albizia inopinata*.

Misra, L.N. et al., *Phytochemistry*, 1995, **39**, 247-249 (*Budmunchiamine L2, isol, ir, pmr, cmr, ms*)

Simoes de Assis, T. et al., *CA*, 2000, **133**, 168223p (*Felipealbizine A*)

Budmunchiamine L4† B-387

8-Tridecyl-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI

[335153-43-0]

As Budmunchiamine L1, B-385 with R = -(CH₂)₁₂CH₃

C₂₆H₅₄N₄O 438.739

Not to be confused with Budmunchiamine L4, B-385. The name has been applied to two different homologues. Alkaloid from *Albizia adinocephala*. Inhibitor of plasmepsin II. Oil. $[\alpha]_D -13$ (c, 0.65 in MeOH).

N¹,N¹³-Di-Me: **Budmunchiamine G**

[143051-88-1]

C₂₈H₅₈N₄O 466.792

From seeds of *Albizia amara* and *Albizia lebbek*.

N¹,N⁹,N¹³-Tri-Me: 1,9,13-Trimethyl-8-tridecyl-1,5,9,13-tetraazacycloheptadecan-6-one. **Budmunchiamine C**

[139750-78-0]

C₂₉H₆₀N₄O 480.819

Alkaloid from the seeds of *Albizia amara* and stem bark of *Albizia gummifera* (Fabaceae). Light yellow oil. $[\alpha]_D -5.1$ (c, 0.08 in CHCl₃).

6'-Hydroxy, N¹,N⁹,N¹³-tri-Me: 6'-Hydroxybudmunchiamine C

[178494-83-2]

C₂₉H₆₀N₄O₂ 496.819

Alkaloid from stem bark of *Albizia schimperana*. $[\alpha]_D -6.4$ (c, 0.08 in CHCl₃).

9'-Oxo, N¹,N¹³-di-Me: **Budmunchiamine I**

[143051-90-5]

C₂₈H₅₆N₄O₂ 480.776

From seeds of *Albizia amara*. Isol. as an inseparable mixt. with Budmunchiamine H.

9'-Oxo, N¹,N⁹,N¹³-tri-Me: **Budmunchiamine E**

[143051-86-9]

C₂₉H₅₈N₄O₂ 494.803

From seeds of *Albizia amara*. Isol. as an inseparable mixt. with Budmunchiamine D.

10'-Oxo, N¹,N¹³-di-Me: **Budmunchiamine H**

[143051-89-2]

C₂₈H₅₆N₄O₂ 480.776

From seeds of *Albizia amara*. Yellowish cryst. (Me₂CO). $[\alpha]_D -8.1$ (c, 0.2 in MeOH). Isol. as an inseparable mixt. with Budmunchiamine I, to which props. refer.

10'-Oxo, N¹,N⁹,N¹³-tri-Me: **Budmunchiamine D**

[143070-37-5]

C₂₉H₅₈N₄O₂ 494.803

Isol. from seeds of *Albizia amara*. Yellow solid (Me₂CO). $[\alpha]_D 0$ (c, 1.6 in MeOH). Isol. as an inseparable mixt. with Budmunchiamine E to which props. refer.

Pezzuto, J.M. et al., *Heterocycles*, 1991, **32**, 1961 (*Budmunchiamines A-C*)

Mar, W. et al., *J. Nat. Prod.*, 1991, **54**, 1531 (*props*)

Pezzuto, J.M. et al., *Phytochemistry*, 1992, **31**, 1795 (*Budmunchiamines D-I*)

Misra, L.N. et al., *Phytochemistry*, 1995, **39**, 247 (*Budmunchiamine G*)

Rukunga, G.M. et al., *J. Nat. Prod.*, 1996, **59**, 850-853 (*Budmunchiamine G*)

Rukunga, G.M. et al., *Phytochemistry*, 1996, **42**, 1211-1215 (6'-Hydroxybudmunchiamine C)

Popaj, K. et al., *Helv. Chim. Acta*, 2001, **84**, 180-186 (\pm -Budmunchiamine LA, \pm -Budmunchiamine C, synth)

Ovenden, S.P.B. et al., *Phytochemistry*, 2002, **60**, 175-177 (*Budmunchiamine LA*)

Budmunchiamine L5† B-388

8-(13-Heptadecenyl)-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI

[195734-29-3]

As Budmunchiamine L1, B-385 with

R = -(CH₂)₁₂CH=CHCH₂CH₂CH₃C₃₀H₆₀N₄O 492.83

Not to be confused with Budmunchiamine L5, B-389. The same name has been applied to two different homologues. Alkaloid from the seeds of *Albizia lebbek*. Viscous liq. $[\alpha]_D^{27} +13.2$ (c, 0.01 in MeOH). λ_{\max} 209 (log ϵ 4) (MeOH).

Dixit, A.K. et al., *J. Nat. Prod.*, 1997, **60**, 1036-1037

Budmunchiamine L5† B-389

8-Pentadecyl-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI

[479243-93-1]

As Budmunchiamine L1, B-385 with

R = -(CH₂)₁₄CH₃C₂₈H₅₈N₄O 466.792

Not to be confused with Budmunchiamine L5, B-388. The name has been applied to two different homologues. Alkaloid from *Albizia adinocephala*. Inhibitor of plasmepsin II. Oil. $[\alpha]_D -20$ (c, 0.21 in MeOH).

N¹,N⁹-Di-Me: 13-Normethylbudmunchiamine K

[180285-78-3]

C₃₀H₆₂N₄O 494.846

Isol. from stem bark of *Albizia gummifera* and from *Albizia schimperana*. $[\alpha]_D -1.8$ (c, 0.09 in CHCl₃).

N¹,N¹³-Di-Me: 9-Normethylbudmunchiamine K†

[178494-84-3]

C₃₀H₆₂N₄O 494.846

Isol. from stem bark of *Albizia schimperana*. $[\alpha]_D -5.1$ (c, 0.08 in CHCl₃).

N⁹,N¹³-Di-Me: 1-Normethylbudmunchiamine K

[178494-86-5]

C₃₀H₆₂N₄O 494.846

Alkaloid from stem bark of *Albizia schimperana*. $[\alpha]_D -1.8$ (c, 0.09 in CHCl₃).

N¹,N⁹,N¹³-Tri-Me: **Budmunchiamine K**

[178494-87-6]

C₃₁H₆₄N₄O 508.873

Alkaloid from stem bark of *Albizia gummifera*. Yellow oil. $[\alpha]_D -9.3$ (c, 0.09 in CHCl₃).

11',12'-Didehydro: 8-(11-Pentadecenyl)-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI. **Budmunchiamine L6**

[195734-30-6]

C₂₈H₅₆N₄O 464.777

Alkaloid from the seeds of *Albizia lebbek*. Viscous liq. $[\alpha]_D^{27} +11.2$ (c, 0.01 in MeOH). λ_{\max} 209 (log ϵ 3.98) (MeOH).

3',4'Z,6',7'Z,9',10'Z-Hexadehydro: 8-(3,6,9-Pentadecatrienyl)-1,5,9,13-tetraazacycloheptadecan-6-one, 9CI. **Felipealbizine B**

[288094-81-5]

C₂₈H₅₂N₄O 460.745

Constit. of the leaves of *Albizia inopinata*.

6'-Hydroxy, N¹,N¹³-di-Me: 6'-Hydroxy-9-normethylbudmunchiamine K

[178494-85-4]

C₃₀H₆₂N₄O₂ 510.845

Isol. from stem bark of *Albizia schimperana*. $[\alpha]_D -2.3$ (c, 0.01 in CHCl₃).

6'-Hydroxy, N¹,N⁹,N¹³-tri-Me: 6'-Hydroxybudmunchiamine K


[180285-72-7]

C₃₁H₆₄N₄O₂ 524.872

Isol. from stem bark of *Albizia gummifera*. $[\alpha]_D -6.4$ (c, 0.07 in CHCl₃).

Rukunga, G.M. et al., *J. Nat. Prod.*, 1996, **59**, 850-853 (*Albizia gummifera alkaloids*)

- Rukunga, G.M. *et al.*, *Phytochemistry*, 1996, **42**, 1211-1215 (*Albizia schimperana* alkaloids)
- Dixit, A.K. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1036-1037 (*Budmunchiamine L6*)
- Simoes de Assis, T. *et al.*, *CA*, 2000, **133**, 168223p (*Felipealazine B*)
- Ovenden, S.P.B. *et al.*, *Phytochemistry*, 2002, **60**, 175-177 (*Budmunchiamine L5*)

Buesgeniine**B-390**

Relative Configuration


C₃₄H₃₇N₈ 587.668

Alkaloid from the stem bark of *Zanthoxylum buesgenii*. Cryst. (MeOH). Mp 141-142°. λ_{max} 230 (log ε 5.1); 286 (log ε 5.15); 321 (log ε 4.57) (EtOH).

Tane, P. *et al.*, *Fitoterapia*, 2005, **76**, 656-660 (*isol*, *pmr*, *cmr*)

Bufopyramide

[220786-68-5]


B-391C₁₈H₁₉N₃O₃ 325.366

Alkaloid from the Chinese traditional drug Ch'an Su (prepared skin secretions of various *Bufo* spp.). Shows cytotoxic activity. Amorph. solid. λ_{max} 208 ; 225 ; 302 (MeOH).

Kamano, Y. *et al.*, *Heterocycles*, 1999, **50**, 499-503 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *activity*)

Bufoserotonin C

[1002722-87-3]


B-392C₁₈H₁₉N₃O₃ 325.366

Alkaloid from the venom of *Bufo* spp. (ChanSu). Amorph. brown powder (MeOH). Mp 182-185°. λ_{max} 221 ; 292 ; 302 (MeOH).

Liu, R.-H. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2427-2431 (*isol*, *pmr*, *cmr*, *ms*)

Bufotenine**B-393**

3-(2-Dimethylaminoethyl)-1H-indol-5-ol, 9CI. 3-(2-Dimethylaminoethyl)-5-hydroxyindole. N^b,N^b-Dimethylserotonin. Cino-bufotenine. Mappine. DM SHT [487-93-4]

C₁₂H₁₆N₂O 204.271

Alkaloid from poisonous secretion of toads (*Bufo* spp.), from seeds of *Piptadenia peregrina*, many genera in the Fabaceae and Poaceae, and in some mushrooms, e.g. *Amanita citrina*. Also identified spectroscopically in the gorgonian *Paramuricea chamaeleon*. Hallucinogen. Sol. MeOH, acids, EtOH, bases; fairly sol. Et₂O; poorly sol. H₂O. Mp 146-147°. Bp_{0.1} 320°. λ_{max} 225 ; 274 ; 300 (MeOH) (Berdy). λ_{max} 220 (ε 10000); 265 (ε 3000) (EtOH) (Berdy).

► Adverse systemic effects by intravenous route (psychotropic). LD₅₀ (mus, ipr) 290 mg/kg. NM2800000

O-Sulfate: **Bufoviridine**. Dihydrobufothionine

[16369-08-7]

C₁₂H₁₆N₂O₄S 284.335

Isol. from the skins of various amphibia incl. *Bufo viridis*.

O-β-D-Glucopyranoside: **Bufotenine O-glucoside**

[64656-15-1]

C₁₈H₂₆N₂O₆ 366.413

Alkaloid from the leaves of *Citrus unshiu* (satsuma mandarin). [α]_D -58 (c, 0.5 in H₂O). λ_{max} 224 ; 274 (No solvent reported).

N^b-Oxide: **Bufotenine N-oxide**

[1019-44-9]

C₁₂H₁₆N₂O₂ 220.271

Alkaloid from *Desmodium pulchellum* and *Desmodium caudatum* (Fabaceae). Also *Piptadenia macrocarpa* and *Amanita* spp. Constit. of toad venom (*Bufo* spp.). Cryst. Mp 214-215°.

N^b-Oxide, O-β-D-glucopyranoside: **Bufotenine O-glucoside N-oxide**

C₁₈H₂₆N₂O₇ 382.413

Alkaloid from *Evodnia fargesii*. Amorph. brown powder. [α]_D²³ -37.4 (c, 0.42 in H₂O). λ_{max} 221 ; 279 (H₂O).

N^b-Me, inner salt: **Bufotenidine**

[487-91-2]

C₁₃H₁₈N₂O 218.298

Alkaloid from secretions of toad spp. Mp 209° (as hydriodide).

N^b-Me, inner salt, O-sulfate: **Bufotenidine O-sulfate**

C₁₃H₁₈N₂O₄S 298.362

Alkaloid from the skin of the toad *Nyctimystes tympanocryptis*. Genus name sometimes given as *Nictimystes*.

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 530D (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 663C (*ir*)

Wieland, T. *et al.*, *Annalen*, 1953, **581**, 10-16 (*isol*, *uv*)

Stromberg, V.L. *et al.*, *J.A.C.S.*, 1954, **76**, 1707 (*isol*, *uv*)

Harley-Mason, J. *et al.*, *J.C.S.*, 1954, 1165-1171 (*synth*)

Fish, M.S. *et al.*, *J.A.C.S.*, 1955, **77**, 5892-5895 (*oxide*)

Erspamer, V. *et al.*, *Biochem. Pharmacol.*, 1959, **2**, 270 (*Bufoviridine*)

Pachter, I.J. *et al.*, *J.O.C.*, 1959, **24**, 1285-1287 (*isol*)

Ghosal, S. *et al.*, *J.O.C.*, 1966, **31**, 2284-2288 (*isol*)

Paris, R.R. *et al.*, *Ann. Pharm. Fr.*, 1967, **25**, 509 (*oxide*)

Daly, J.W. *et al.*, *J.A.C.S.*, 1967, **89**, 1032-1033 (*pmr*)

Creveling, C.R. *et al.*, *Clin. Chem. (Winston-Salem, N.C.)*, 1968, **14**, 302-309 (*ms*)

Roseghini, M. *et al.*, *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1976, **54**, 31-43; 1989, **94**, 455-460 (*Bufoviridine*)

Sanders-Bush, E. *et al.*, *Life Sci.*, 1976, **19**, 1407 (*metab*)

Schultes, R.E. *et al.*, *Planta Med.*, 1976, **29**, 330

Roseghini, M. *et al.*, *Z. Naturforsch., C*, 1976, **31**, 118-120 (*Bufoviridine*, *Bufotenidine O-sulfate*)

Shaw, G.J. *et al.*, *Biomed. Mass Spectrom.*, 1977, **4**, 348-353 (*ms*)

Cimino, G. *et al.*, *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1978, **61**, 361-362 (*occur*)

Barlow, R.B. *et al.*, *Br. J. Pharmacol.*, 1980, **69**, 597-600 (*pharmacol*)

Nishida, R. *et al.*, *Agric. Biol. Chem.*, 1990, **54**, 1853-1855 (*glucoside*)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1555

Somei, M. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 87-96 (*synth*)


Zhang, P. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 1582-1586 (*toad venom constits*)

Qu, S.-J. *et al.*, *Planta Med.*, 2006, **72**, 264-266 (*N-oxide glucoside*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DPG109

Bufotoxin**B-394***Vulgarobufotoxin*

[464-81-3]



Absolute configuration

C₄₀H₆₀N₄O₁₀ 756.935

Ester of suberic acid with 14,15-Epoxy-3,5-dihydroxybufa-20,22-dienolide, E-132 and 2-Amino-4-hexynoic acid. Isol. from the toxic secretions of the common toad *Bufo vulgaris*. Cardiotoxic agent. Mp 204-205°. [α]_D²⁴ +3.9 (MeOH).

Bufotoxin is the most investigated of a series of frog toxin components known generally as Bufotoxins in which various steroidal genins are esterified at C-3 with suberoyl amino acid residues. The minor components have not all been characterised. See also 14,15-Epoxy-3,5-dihydroxybufa-20,22-dienolide, E-132.

► LD₅₀ (mus, scu) 0.4 mg/kg. LD₅₀ (cat, ivn) 0.3 mg/kg. EI3160000

L-Glutamine analogue: see 14,15-Epoxy-3,5-dihydroxybufa-20,22-dienolide, E-132

Wieland, H. *et al.*, *Annalen*, 1936, **524**, 203; 1941, **549**, 209 (*isol, uv, struct*)


Linde-Tempel, H.O. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 2188 (*struct*)

Shimada, K. *et al.*, *Tet. Lett.*, 1979, 163

Pettit, G.R. *et al.*, *J.O.C.*, 1987, **52**, 3573 (*synth*)

Bujeine

[221011-11-6]



C₂₀H₂₃NO₆ 373.405


Alkaloid from *Narcissus bujei*. Mp 140-142°. [α]_D²⁰ +129.4 (c, 0.11 in MeOH).

Labrana, J. *et al.*, *Phytochemistry*, 1999, **50**, 183-188 (*isol, ir, cd, pmr, cmr, ms*)

Bukittingine

[128412-07-7]

[138435-38-8 ((±)-form)]



Absolute Configuration

C₂₂H₃₁NO₂ 341.492

Novel skeleton, closely related to the *Daphniphyllum* alkaloids such as Methylhomosecodaphniphyllate in H-346. Major alkaloid from the leaves and terminal branchlets of *Sapium baccatum* (Euphorbiaceae). Felted needles (hexane). Mp 135-137°. [α]_D -157 (c, 0.2 in CHCl₃).

Hydrobromide:

Plates (MeOH/Me₂CO). Dec. at =310° without melting (sealed tube).

Methiodide:

Cryst. +H₂O (MeOH/Me₂CO). Mp 187-189° dec.


Arbain, D. *et al.*, *Aust. J. Chem.*, 1990, **43**, 185 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1992, **57**, 2575 (*synth*)

Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (*synth, rev*)

Bulbocapnine

6,7,7a,8-Tetrahydro-11-methoxy-7-methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-12-ol, 9Cl. 11-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine. N-Methylaunobine



(S)-form

C₁₉H₁₉NO₄ 325.363

Shows antimicrobial activity. Log P 2.59 (uncertain value) (calc).

(S)-form [298-45-3]

Alkaloid from *Corydalis cava* (Papaveraceae). Antihypertensive, cataleptic, sedative agent. Possesses adrenolytic props. In low doses depresses spinal motor-neurone synaptic reflexes. Shows antimicrobial activity. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 201-203°. [α]_D²⁰ +232 (c, 0.92 in CHCl₃). Pharmacol. active isomer. λ_{max} 335; 397 (MeOH) (Berdy). ► LD₅₀ (mus, scu) 195 mg/kg. CE1050000

N-Oxide(β-): *Bulbocapnine β-N-oxide*

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Glaucium fimbriigerum*. Needles (Me₂CO). Mp 153-155°. [α]_D²⁰ +180 (c, 0.65 in MeOH). λ_{max} 225 (log ε 4.19); 269 (log ε 3.86); 307 (log ε 3.57) (MeOH).

N-Me: *N-Methylbulbocapnine*

[5787-06-4]

C₂₀H₂₂NO₄[⊕] 340.398

Quaternary alkaloid from the tubers of *Corydalis cava*, also present in trace amount in the aerial parts (Papaveraceae). Needles (MeOH) (as iodide). Mp 268-269° (iodide). [α]_D²² +163 (c, 0.11 in MeOH).

► DE5380000

N-De-Me: 11-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine. *Launobine*. *Norbulbocapnine*

[20497-21-6]

C₁₈H₁₇NO₄ 311.337

Alkaloid from *Cassipoupa americana* and *Laurus nobilis* (bay laurel). Also occurs in the genera *Lindera* and *Illigera* (Lauraceae, Hernandiaceae). Shows bactericidal activity against *Scenedesmus obliquus*. Mp 214-215°. [α]_D²¹ +192 (c, 0.95 in CHCl₃).

Me ether: 10,11-Dimethoxy-1,2-methylenedioxyaporphine. *O-Methylbulbocapnine*. *N,O-Dimethylnandigerine*

[2490-83-7]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Lindera oldhamii* (Lauraceae). Shows antimicrobial activity. Mp 129-130°. [α]_D +248 (c, 0.67 in CHCl₃).

Me ether, *N*-oxide(α-): *O-Methylbulbocapnine α-N-oxide*

[122331-84-4]

C₂₀H₂₁NO₅ 355.39

B-397

Alkaloid from the leaves of *Polyalthia longifolia* (Annonaceae). Yellowish oil. [α]_D +153 (CHCl₃).

Me ether, *N*-oxide(β-): *O-Methylbulbocapnine β-N-oxide*

[122331-83-3]

C₂₀H₂₁NO₅ 355.39

Alkaloid from the leaves of *Polyalthia longifolia* (Annonaceae). Greyish-white needles (MeOH). Mp 115-117°. [α]_D +158 (c, 0.1 in MeOH).

Me ether, *N*-de-Me, *N*-(*N*-methylcarbamoyl): *O-Methyl-N-(N-methylcarbamoyl)aunobine*. *O-Methyl-N-(N-methylcarbamoyl)nandigerine*

C₂₁H₂₂N₂O₅ 382.415

Alkaloid from *Hernandia nymphaeifolia*. Amorph. power. [α]_D²³ +314 (c, 0.2 in CHCl₃). λ_{max} 225 (log ε 4.57); 272 (log ε 4.16); 308 (log ε 3.82) (EtOH).

4β-Hydroxy: *4-Hydroxybulbocapnine*

[70475-64-8]

C₁₉H₁₉NO₅ 341.363

Alkaloid from the aerial parts of *Glaucium vitellinum* (Papaveraceae). Mp 231-233°. [α]_D +100 (c, 0.14 in CHCl₃).

(±)-form

Synthetic. Mp 213-214° (209-210°).

(ξ)-form

Me ether, *N*-de-Me: 10,11-Dimethoxy-1,2-methylenedioxyaporphine. *Litseedine*

[56245-65-9]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the bark of *Litsea nitida* (Lauraceae). Lustrous brownish cryst. (MeOH/CHCl₃). Mp 182-183°.

Gulland, J.M. *et al.*, *J.C.S.*, 1928, 1132 (*synth*)
Tomita, M. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 763 (*Launobine*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2443 (*Bulbocapnine, Launobine, isol*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 157-176 (*activity*)

Lu, S.-T. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 910; *CA*, **77**, 101949m (*O-Methylbulbocapnine*)

Patnaik, P.C. *et al.*, *Indian J. Chem.*, 1975, **13**, 197 (*Litsedine*)

Shafiee, A. *et al.*, *J. Nat. Prod.*, 1979, **42**, 174 (*4-Hydroxybulbocapnine*)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (*cd*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1985, **48**, 646 (*4-Hydroxybulbocapnine*)

Soicke, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 149 (*synth*)

Wu, Y.-C. *et al.*, *Heterocycles*, 1989, **29**, 463 (*O-Methylbulbocapnine N-oxides*)

Ribár, B. *et al.*, *Acta Cryst. C*, 1991, **47**, 2612 (*cryst struct*)

Abbasoglu, U. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 379-380 (*activity*)

Shafiee, A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1564-1565 (*β-N-oxide*)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Chen, J.J. *et al.*, *Planta Med.*, 2000, **66**, 251-256 (*O-Methyl-N-methylcarbamoylaunobine*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HLT000

Bulbocodine†

[33792-78-8]


C₁₉H₂₃NO₃ 313.396

Alkaloid from the whole plants of *Bulbocodium vernum* (Liliaceae). Cryst. (Me₂CO). Mp 262° (220-222°). [α]_D²² +111 (CHCl₃).

Šantavý, F. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 1084 (*isol, uv, ir, pmr, ms, cd, struct*)

Bulbodione

[96681-60-6]


C₁₉H₁₅NO₅ 337.331

Minor alkaloid from *Corydalis bulbosa* (Papaveraceae). Blue-violet needles (EtOH). Mp 248-250°.

Kiryakov, H.G. *et al.*, *Planta Med.*, 1984, **50**, 136 (*isol, uv, ir, pmr, ms, struct, synth*)

Bulgaramine

[96681-78-6]

C₂₁H₂₁NO₄ 351.401

Alkaloid from Herba *Fumaria* Officinalis (commercial medicinal mixt. of several *Fumaria* spp. growing in Bulgaria) (Papaveraceae). Cryst. (EtOH). Mp 209°.


Yakimov, G.I. *et al.*, *J. Nat. Prod.*, 1984, **47**, 1048 (*isol, uv, pmr, ms, struct*)

Blasko, G. *et al.*, *Acta Chim. Hung.*, 1991, **128**, 819 (*synth*)

Giese, M.W. *et al.*, *J.O.C.*, 2005, **70**, 6222-6229 (*synth*)

Bulgarsenine

[62018-77-3]



Absolute configuration

B-398C₁₈H₂₇NO₅ 337.415

Cyclic platynecine diester. Alkaloid from *Senecio nemorensis*, *Senecio doricum* and *Senecio abrotanifolius* (Asteraceae). Mp 115° (112-113°). [α]_D²⁴ -54 (c, 0.78 in CHCl₃). [α]_D²⁰ -74.2 (EtOH).

O-Ac: 11-O-AcetylbulgarsenineC₂₀H₂₉NO₆ 379.452

Alkaloid from *Senecio callosus*. Cryst. (EtOAc) (as dihydrochloride). Mp 123-125° (2HCl). [α]_D -46.9 (c, 0.26 in CHCl₃) (2HCl). λ_{max} 223 (log ε 3.9) (MeOH) (2HCl).

O-Ac, N-oxide: 11-O-Acetylbulgarsenine N-oxideC₂₀H₂₉NO₇ 395.452

Alkaloid from *Senecio callosus*. Cryst. (EtOAc). Mp 142-144°. [α]_D -60.5 (c, 0.2 in CHCl₃). Possible artifact. λ_{max} 224 (log ε 3.87) (MeOH).

N-(Chloromethyl): N-ChloromethylbulgarsenineC₁₉H₂₉ClNO₅ 386.895

Quaternary alkaloid from *Senecio callosus*. Cryst. (hexane/EtOAc) (as chloride). Mp 173-175° (chloride). [α]_D -48.1 (c, 0.3 in CHCl₃) (chloride). Possible artifact. λ_{max} 227 (log ε 4.19) (MeOH) (chloride).

1,2-Didehydro: Doronenine

[74217-57-5]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio doricum* (Asteraceae). Mp 124-127°. [α]_D²⁰ +123.4 (EtOH). Cyclic retronecine diester.

12,13-Diepimer: IodanthineC₁₈H₂₇NO₅ 337.415

Alkaloid from *Senecio bracteatus* and *Senecio iodanthus*. Cryst. (hexane/EtOAc) (as hydrochloride). Mp 217-220° (hydrochloride). [α]_D²⁵ -103.3 (c, 2.1 in CHCl₃) (hydrochloride).

Nghia, N.T. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 2952 (*isol, uv, ms, pmr, struct*)

Stoeckli-Evans, H. *et al.*, *Acta Cryst. B*, 1980, **36**, 3150 (*cryst struct*)

Kurfel, A. *et al.*, *Cryst. Struct. Commun.*, 1980, **9**, 353 (*cryst struct, deriv*)

Roeder, E. *et al.*, *Phytochemistry*, 1980, **19**, 1275; 1984, **23**, 1761 (*isol, ms, struct*)


Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Perez-Castorena, A.-L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1288-1291; 1999, **62**, 1039-1043 (*11-Acetylbulgarsenine, Iodanthine*)

Bulgecin A

β-Lactam activator F2

[92953-54-3]

B-402C₁₆H₂₉N₃O₁₄S₂ 551.549

Aminoglycoside antibiotic. Isol. from *Pseudomonas acidophila* and *Pseudomonas mesoacidophila*. Induces formn. of bulges in bacterial cell walls by cooperation with β-lactam antibiotics. Sol. H₂O; fairly sol. MeOH, DMSO; poorly sol. EtOAc, phenol, hexane. Major component of Bulgecin complex.

▶ LD₅₀ (mus, ivn) 1000-3000 mg/kg.

Mono-Na salt:

Needles (MeOH aq.). Mp 211-212°. [α]_D²⁰ +6.5 (c, 0.5 in 1M AcOH).

Imada, A. *et al.*, *J. Antibiot.*, 1982, **35**, 1400;

1985, **38**, 17 (*isol, ir, pmr, cmr, struct*)

Shinagawa, S. *et al.*, *Tetrahedron*, 1984, **40**, 3465 (*isol, struct*)

Cintas, P. *et al.*, *Tetrahedron*, 1991, **47**, 6079 (*synth*)

Bulgecin B**B-403**F₆. Antibiotic F₆

[95863-86-8]

As Bulgecin A, B-402 with

R = -NHCH₂CH₂COOHC₁₇H₂₉N₃O₁₃S 515.494

Aminoglycoside antibiotic. From *Pseudomonas acidophila* and *Pseudomonas mesoacidophila*. Induces the formation of bulges in bacterial cell walls by cooperation with β-lactam antibiotics. Powder + 1H₂O (MeOH aq.) (as Na salt). [α]_D²⁰ -2.6 (c, 0.5 in 1M AcOH) (Na salt). Minor component of Bulgecin complex.

Imada, A. *et al.*, *J. Antibiot.*, 1982, **35**, 1400;

1985, **38**, 17 (*isol, ir, pmr, cmr, struct*)

Shinagawa, S. *et al.*, *Tetrahedron*, 1984, **40**,

3465 (*isol, struct*)

Cintas, P. *et al.*, *Tetrahedron*, 1991, **47**, 6079

(*synth*)**Bulgecin C****B-404**

[92953-56-5]

As Bulgecin A, B-402 with

R = OH

C₁₄H₂₄N₂O₁₂S 444.416

Aminoglycoside antibiotic. From *Pseudomonas acidophila* and *Pseudomonas mesoacidophila*. Causes bulges in bacterial cell walls. Synergistic with β-lactam antibiotics in antimicrobial activity. Powder + ½ H₂O (MeOH aq.) (as Na salt). Sol. H₂O. [α]_D²⁰ +2.9 (c, 0.6 in 1M AcOH) (Na salt). Minor component of Bulgecin complex.

▶ LD₅₀ (mus, ivn) 1000-2000 mg/kg.

Imada, A. *et al.*, *J. Antibiot.*, 1982, **35**, 1400;

1985, **38**, 17 (*isol, ir, pmr, cmr, struct*)

Shinagawa, S. *et al.*, *Tetrahedron*, 1984, **40**,

3465 (*isol, struct*)

Barret, A.G.M. *et al.*, *J.O.C.*, 1991, **56**, 2787

(*synth*)

Cintas, P. *et al.*, *Tetrahedron*, 1991, **47**, 6079

(*synth*)**Bullatine A****B-405**

[1354-84-3]

C₂₁H₃₁NO₂ 329.481

Struct. unknown. Unsatd. base with an ethylimino group and 2 hydroxyls. Alkaloid from the Chinese drug Hye-shang-yi-zhi-hao (*Aconitum bullatifolium* var.

homotrichum, Ranunculaceae). Mp 251-253°. $[\alpha]_D^{27}$ -55 (MeOH).

Hydrochloride: Mp 265°.

Hydrobromide: Mp 263-264°.

Nitrate: Mp 237°.

Perchlorate: Mp 231-232°.

Picrate: Mp 221°.

Di-Ac: Mp 145-147°.

Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1964, **30**, 139-145; *CA*, **61**, 8128e

Bullatine D**B-406**

[1354-87-6]

C₂₃H₃₇NO₉ 471.547


Struct. unknown. Trace alkaloid from the Chinese drug Hye-shang-yi-zhi-hao (*Aconitum bullatifolium* var. *homotrichum*, Ranunculaceae). Mp 211°.

Hydrobromide: Mp 224°.

Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1964, **30**, 139-145; *CA*, **61**, 8128e

Bulleyanaline**B-407**

8-Acetyl-5-hydroxycorynoline
[104574-44-9]



Absolute
Configuration

C₂₄H₂₅NO₇ 439.464

Various numbering systems in use. The stereochem. of Bulleyanaline and the other alkaloids covered by this entry is uncertain. Alkaloid from *Corydalis bulleyana* whole plant (Papaveraceae). Cryst. (MeOH). Mp 199-201°. $[\alpha]_D^{18}$ +128.2 (EtOH).

5-Deoxy (?): 8-Acetylcorynoline

[82189-89-7]

C₂₄H₂₅NO₆ 423.465

Minor alkaloid from *Corydalis delavayi* and *Corydalis bulleyana* (Papaveraceae). Mp 218-220° (210-212°). $[\alpha]_D^{17}$ +123. C-8 config. uncertain. The two 8-epimeric (±) isomers have been synthesized.

5-Deoxy, Ac: **Conspersine**. 6-Acetylcorynoline

[91897-67-5]

C₂₆H₂₇NO₇ 465.502

Alkaloid from *Corydalis conspersa* whole plant (Papaveraceae). Granules (CHCl₃/MeOH). Mp 215-218°. $[\alpha]_D^{20}$ +107.54 (CHCl₃). CAS gives the 8 α (14*R*-) config., but the authors state that the 8 β - config. is more probable. However in a later paper it is shown as 8 α but said to be uncertain. λ_{max} 232 (sh) (log ϵ 3.88); 287 (log ϵ 3.8) (EtOH).

Stereoisomer, 5-deoxy, Ac: **Bulleyanine**. 8-Acetylacetyliscorynoline
[104639-71-6]

C₂₆H₂₇NO₇ 465.502

Alkaloid from *Corydalis bulleyana* whole plant (Papaveraceae). Powder. Stereochem. not certain. Prob. epimeric with Conspersine at the ring junction, C-8 config. not detd. λ_{max} 235 ; 290 (EtOH).


Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1880-1889 (8-Acetylcorynoline, synth, uv, ir, pmr, cmr, ms)

Luo, S. *et al.*, *Yunnan Zhiwu Yanjiu*, 1982, **4**, 69; *CA*, **97**, 20743a (8-Acetylcorynoline, isol, struct)

Fang, Q. *et al.*, *Planta Med.*, 1984, **50**, 25-27; 1986, **52**, 193-198 (*Conspersine*, *Bulleyanine*, *Bulleyanaline*)

Bulleyaniline B**B-408**

[128717-98-6]



R¹ = CH₃, R² = H

C₃₅H₄₇N₃O₁₀ 669.77

Alkaloid from the roots of *Delphinium bulleyanum* (Ranunculaceae).

Wei, X. *et al.*, *Yunnan Zhiwu Yanjiu*, 1989, **11**, 453; *CA*, **113**, 74750 (isol)

Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1028 (config)

Bulleyaniline C**B-409**

[128717-99-7]

As Bulleyaniline B, B-408 with

R¹ = H, R² = CH₃C₃₅H₄₇N₃O₁₀ 669.77


Alkaloid from the roots of *Delphinium bulleyanum* (Ranunculaceae).

Wei, X. *et al.*, *Yunnan Zhiwu Yanjiu*, 1989, **11**, 453; *CA*, **113**, 74750 (isol)

Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1028 (config)

Bungeanine**B-410**

[113145-59-8]



Absolute
Configuration


C₂₂H₁₉NO₆ 393.395

Alkaloid from *Corydalis bungeana* (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 141-143°. $[\alpha]_D^{25}$ +40 (c, 0.37 in CHCl₃).

Weiguang, Z. *et al.*, *Planta Med.*, 1987, **53**, 418 (isol, uv, ir, pmr, cmr, ms, struct)

Buntanbismine**B-411**

[176520-67-5]

C₃₅H₃₂N₂O₉ 624.646

Alkaloid from stem bark of *Citrus grandis* f. *buntan*. Red needles (CHCl₃). Mp >300°.

5-Ac:

Yellow needles (CHCl₃). Mp >250°.

1,5-Di-Ac:

Yellow syrup.

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **42**, 221 (isol, uv, ir, pmr, cmr, ms, struct)

Buphacetine**B-412**

[1354-91-2]


C₂₀H₂₅NO₇ 391.42

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Boophone disticha* (Amaryllidaceae). Needles (Me₂CO/Et₂O). Mp 182-183°. $[\alpha]_D^{21}$ -73 (c, 0.418 in CHCl₃). $[\alpha]_D^{21}$ -51 (c, 0.418 in EtOH).

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 491-502 (isol, uv, ir)

Buphanamine**B-413**

[6793-24-4]



Absolute
Configuration

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Boophone fischeri*, *Boophone disticha* and *Crinum bulbispermum* (Amaryllidaceae). Mp 184-186°. $[\alpha]_D^{20}$ -205 (EtOH).

Perchlorate: Mp 232-234°. $[\alpha]_D^{20}$ -154 (95% EtOH).

Renz, J. *et al.*, *Helv. Chim. Acta*, 1955, **38**, 1209 (isol, uv, ir)

Fales, F.C. *et al.*, *J.O.C.*, 1961, **26**, 881 (uv, ir, struct)

Longévialle, P. *et al.*, *Org. Mass Spectrom.*, 1973, **7**, 401 (ms)

Buphanine**B-414**

[1390-55-2]

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Boophone disticha* (Amaryllidaceae). *B. disticha* has been used as an arrow poison. Amorph.

Mol. formula not determined. Converted by KOH to Nerbowdine, N-169.


► Highly toxic.

Tutin, F. *et al.*, *J.C.S.*, 1911, **99**, 1240-1248 (isol)

Buprestin A

B-415

β-D-Glucopyranose 1,2,6-tri-1H-pyrrole-2-carboxylate
[95245-60-6]



C₂₁H₂₁N₃O₉ 459.412

Isol. from the Jewel beetle *Stigmodera macularia*. Predator repellent. Amorph. solid. Mp 135-140° (softens at 125°).

Brown, W.V. *et al.*, *Aust. J. Chem.*, 1985, **38**, 197 (isol, cmr, pmr, ms)


Moore, B.P. *et al.*, *J. Aust. Entomol. Soc.*, 1985, **24**, 81; *CA*, **103**, 193347v

Schramm, S. *et al.*, *Tet. Lett.*, 2006, **47**, 7741-7743 (synth)

Buprestin B

B-416

β-D-Glucopyranose-6-(4-hydroxybenzoate) 1,2-di-1H-pyrrole-2-carboxylate, 9CI
[95258-10-9]



C₂₃H₂₂N₂O₁₀ 486.434

Isol. from the Jewel beetle *Stigmodera macularia*. Predator repellent. λ_{max} 265 nm (MeOH).

Brown, W.V. *et al.*, *Aust. J. Chem.*, 1985, **38**, 197 (isol, pmr, cmr, ms)

Schramm, S. *et al.*, *Tet. Lett.*, 2006, **47**, 7741-7743 (synth)

Burmanneline

B-417

C₂₁H₂₃NO₄ 353.417

Struct. unknown. Alkaloid from root of *Cyclea burmanni* (Menispermaceae). Mp 165°.

Saradamma, S.P. *et al.*, *CA*, 1955, **49**, 11794a

Burmantine

B-418

C₁₈H₂₁NO₃ 299.369

Struct. unknown. Alkaloid from roots of *Cyclea burmanni* (Menispermaceae). Mp 218°.

Hydrochloride: Mp 304°. [α]_D^{28.5} +235.5 (H₂O).

Sulfate: Mp 252° dec.

Methiodide: Mp 292°.

Picrate: Mp 246° dec.


Ac: Mp 140°.

Saradamma, S.P. *et al.*, *CA*, 1955, **49**, 11794a

Burnamicine

B-419

5-Ethylidene-1,2,3,4,5,6,7,9-octahydro-6-(2-hydroxyethyl)-3-methyl-8H-azecino[5,4-b]indol-8-one, 9CI
[2134-96-5]



Absolute Configuration

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Hunteria eburnea* (Apocynaceae). Cryst. (EtOH). Mp 198-200° dec. (193-195°, synthetic). [α]_D²⁴ -281 (CHCl₃). λ_{max} 311 (ε 14600) (no solvent reported).

Bartlett, M.F. *et al.*, *J.A.C.S.*, 1963, **85**, 1203-1204 (ms, struct)


Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 2197-2199 (isol, uv)

Sakai, S. *et al.*, *Heterocycles*, 1976, **4**, 981-983 (synth, cd, abs config)

Burnamine

B-420

Deacetylpicraline
[6808-68-0]



Absolute Configuration

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the seeds of *Picralima nitida*, the stem bark of *Hunteria eburnea* and *Rauwolfia cumminsii*, and the bark of *Aspidosperma cuspa* (Apocynaceae). Cryst. (C₆H₆/Et₂O). Mp 197-198°. [α]_D -131 (CHCl₃). λ_{max} 236 (ε 7000); 289 (ε 2800) (EtOH).

Picrate:

Needles (MeOH). Mp 147-149°.

17-Ac: *Picraline*

[2671-32-1]

C₂₃H₂₆N₂O₅ 410.469

Alkaloid from *Picralima klaineana*, the leaves of *Rauwolfia oregiton* and the bark of *Aspidosperma rigidum* (Apocynaceae). Needles (C₆H₆/pentane). Mp 160-162° Mp 182°. λ_{max} 237 (ε 7400); 289 (ε 3200) (EtOH).

17-Ac, *picrate*:

Yellow prisms (MeOH). Mp 162-164°.

17-O-(3,4,5-Trimethoxybenzoyl): *Deace-*

tylpicraline 3,4,5-trimethoxybenzoate
[102358-22-5]

C₃₁H₃₄N₂O₈ 562.618

Alkaloid from roots of *Alstonia yunnanensis*. Cryst. Mp 222°. [α]_D¹⁷ -185 (c, 0.05 in CHCl₃).

N¹-Me: N^a-*Methylburnamine*. N^a-

Methyldeacetylpicraline

[126594-72-7]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Amorph. solid. [α]_D²⁸ -116.4 (c, 1.25 in MeOH).

N¹-Me, 17-Ac: *Picratidine*. N-Methylpicraline

[131989-85-0]

C₂₄H₂₈N₂O₅ 424.496

Alkaloid from the seeds of *Picralima nitida* (Apocynaceae). Cryst. Mp 185-187°. [α]_D²² -100 (c, 0.07 in EtOH). λ_{max} 233 (log ε 8620); 293 (log ε 3430) (EtOH).

17-Aldehyde: *Picralinal*

[20045-06-1]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from *Picralima klaineana*, *Alstonia scholaris*, *Rhazya stricta* and *Rhazya orientalis* (Apocynaceae). Cryst. (MeOH or C₆H₆/pentane). Mp 156-158° (154-155°) Mp 169-170° dec Mp 179-180°. λ_{max} 237 (ε 7400); 293 (ε 2690) (EtOH).

17-Aldehyde, *picrate*:

Yellow needles (MeOH). Mp 170-172°.

10-Methoxy, N¹-Me, 17-benzoyl: 10-Methoxy-N^a-methylburnamine 17-O-benzoate

[154849-48-6]

C₃₀H₃₂N₂O₆ 516.593

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Prisms (MeOH). Mp 192-193°. [α]_D²⁶ -151.1 (c, 0.17 in MeOH).

10-Methoxy, N¹-Me, 17-O-(3,4-dimethoxybenzoyl): 10-Methoxy-N^a-methylburnamine 17-O-veratrate

[154849-49-7]

C₃₂H₃₆N₂O₈ 576.645

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. [α]_D²⁴ -129.8 (c, 1.24 in MeOH).

10-Methoxy, 17-aldehyde: *Vincarinine*

[33023-09-5]

C₂₂H₂₄N₂O₅ 396.442

Alkaloid from *Vinca erecta* (Apocynaceae). Cryst. (C₆H₆). Mp 178-179°. [α]_D²² -81 (c, 0.1 in MeOH). Struct. revised in 1985. λ_{max} 238 (log ε 3.89); 312 (log ε 3.53) (EtOH).

10-Methoxy, 17-aldehyde, *oxime*:

Cryst. (MeOH). Mp 211-213° dec.

10,11-Dimethoxy, N¹-Me: 10,11-Dimethoxy-1-methyldeacetylpicraline

[79667-54-2]

C₂₄H₃₀N₂O₆ 442.511

Alkaloid from the leaves of *Alstonia lanceolifera* (Apocynaceae). Cryst. (MeOH). Mp 204°. [α]_D -90 (c, 0.7 in CHCl₃). λ_{max} 217; 248; 303 (MeOH).

10,11-Dimethoxy, N¹-Me, 17-Ac: 10,11-Dimethoxy-1-methylpicraline

[79659-66-8]

- $C_{26}H_{32}N_2O_7$ 484.548
Alkaloid from the stem bark of *Alstonia lanceolata* (Apocynaceae). Cryst. (MeOH/Et₂O). Mp 228°. $[\alpha]_D^{20}$ -204 (c, 1.9 in CHCl₃) (-124). λ_{max} 215 ; 246 (log ϵ 3.84); 305 (log ϵ 3.68) (EtOH).
- 10,11-Dimethoxy, N¹-Me, 17-benzoyl: 10,11-Dimethoxy-1-methyldeacetylpicraline benzoate* [80787-53-7]
 $C_{31}H_{34}N_2O_7$ 546.619
Alkaloid from the leaves of *Alstonia lanceolata* (Apocynaceae). Cryst. (MeOH). Mp 114°. $[\alpha]_D^{20}$ -157 (c, 0.5 in CHCl₃). λ_{max} 220 ; 237 ; 282 (sh) ; 305 (MeOH).
- 10,11-Dimethoxy, N¹-Me, 17-O-(3,4,5-trimethoxybenzoyl): 10,11-Dimethoxy-1-methyldeacetylpicraline 3,4,5-trimethoxybenzoate* [79659-67-9]
 $C_{34}H_{40}N_2O_{10}$ 636.697
Alkaloid from the leaves of *Alstonia lanceolata* (Apocynaceae). Cryst. (MeOH). Mp 194°. $[\alpha]_D^{20}$ -126 (c, 0.98 in CHCl₃). λ_{max} 212 (log ϵ 4.64); 255 (log ϵ 4.09); 302 (log ϵ 3.16) (MeOH).
- Z-Isomer, 17-O-(3,4,5-trimethoxybenzoyl): O-(3,4,5-Trimethoxybenzoyl)-(Z)-burnamine* [214263-36-2]
 $C_{31}H_{34}N_2O_8$ 562.618
Alkaloid from *Alstonia villosa*. Prisms. Mp 168-173°. $[\alpha]_D^{28}$ -199.2 (c, 1.12 in MeOH). λ_{max} 213 (log ϵ 4.59); 265 (log ϵ 4.04); 293 (log ϵ 3.83) (MeOH).
- Z-Isomer, 17-aldehyde: (Z)-Picralinal* [214263-35-1]
 $C_{21}H_{22}N_2O_4$ 366.416
Alkaloid from *Alstonia villosa*. Solid. $[\alpha]_D^{28}$ -144.3 (c, 0.85 in MeOH). λ_{max} 208 (log ϵ 4.67); 230 (log ϵ 4.4); 290 (log ϵ 4.1) (MeOH).
- Olivier, L. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 646-650 (*Picraline, Picralinal, uv, ir, pmr, ms, struct*)
Britten, A.Z. *et al.*, *J.C.S.*, 1963, 3850-3854 (*Picraline, Picralinal, uv, ir, synth, struct*)
Bartlett, M.F. *et al.*, *J.O.C.*, 1963, 28, 2197-2199 (*Burnamine, isol, uv, ir*)
Taylor, W.I. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 392 (*Burnamine, uv, struct*)
Durham, L.J. *et al.*, *Tet. Lett.*, 1965, 5-12 (*Picraline, pmr*)
Evans, D.A. *et al.*, *Phytochemistry*, 1968, 7, 1429-1431 (*Picralinal*)
Burnell, R.H. *et al.*, *Phytochemistry*, 1968, 7, 2045-2051 (*Burnamine, isol, uv, ir, pmr*)
Rastogi, R.C. *et al.*, *Experientia*, 1970, 26, 1056 (*Picralinal, isol*)
Il'yasova, Kh.T. *et al.*, *Khim. Prir. Soedin.*, 1971, 7, 164-166; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, 7, 155-156 (*Vincarinine*)
Vercauteren, J. *et al.*, *Phytochemistry*, 1981, 20, 1411-1413 (*10,11-Dimethoxy-1-methylpicraline*)
Petitfrère-Auvray, N. *et al.*, *Phytochemistry*, 1981, 20, 1987-1990 (*Alstonia lanceolata alkaloids*)
Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1985, 21, 129-130; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, 21, 131-132 (*Vincarinine, pmr, cmr, struct*)
Chen, W. *et al.*, *Yaoxue Xuebao*, 1985, 20, 906-912; *CA*, 105, 3520s (*Deacetylpicraline 3,4,5-trimethoxybenzoate*)
- Ansa-Asamoah, R. *et al.*, *J. Nat. Prod.*, 1990, 53, 975-977 (*Picratidine*)
Yamauchi, T. *et al.*, *Phytochemistry*, 1990, 29, 3547-3552 (*N-Methylburnamine*)
Abe, F. *et al.*, *Phytochemistry*, 1994, 35, 253-257 (*10-Methoxy-N¹-methylburnamine 17-O-benzoate, 10-Methoxy-N¹-methylburnamine 17-O-veratrate*)
Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, 46, 1235-1238 (*Z-Picralinal, Trimethoxybenzoyl-Z-burnamine*)
- 1,3-Butadiene-1,1,4,4-tetra-carboxylic acid** **B-421**
(HOOC)₂C=CHCH=C(COOH)₂
 $C_8H_6O_8$ 230.131
Tetra-Et ester:
 $C_{16}H_{22}O_8$ 342.345
Cryst. (EtOH). Mp 56°. Bp_{0.2} 170°.
Tetrakis (methylamide): N,N',N'',N'''-Tetramethyl-1,3-butadiene-1,1,4,4-tetracarboxamide, 9CI. **Chevalieramide** [124709-40-6]
 $C_{12}H_{18}N_4O_4$ 282.299
Alkaloid from the soft coral *Lobophytum chevalieri*.
[64496-11-3]
Baganz, H. *et al.*, *Chem. Ber.*, 1961, 94, 2132 (*synth, ester*)
Li, R. *et al.*, *CA*, 1990, 112, 52550e (*Chevalieramide, isol, pmr, cmr, ms*)
- 1,4-Butanediamine, 9CI** **B-422**
1,4-Diaminobutane. Tetramethylenediamine. Putrescine
[110-60-1]
 $H_2NCH_2CH_2CH_2CH_2NH_2$
 $C_4H_{12}N_2$ 88.152
Manuf. by amination of 1,4-dihalobutanes. Prod. of dec. of excreta, dead animal matter etc. Isol. in small amts. from plants, e.g. *Datura stramonium*, and esp. from fungi. Anal sac secretion of the red fox *Vulpes vulpes*. Occurs in *Anthocharis crassispina*, *Stichopus japonica*, *Halocynthia roretzi* and *Tapes philippinarum*. Prod. by the thermophilic eubacteria, *Thermoleophilum album*, *Thermoleophilum minutum* and *Hydrogenobacter thermophilus*. Cross-linking agent for epoxy resins. Comonomer with Hexanedioic acid, H-258 for manuf. of nylon 4.6. Cryst. with strong odour. Sol. H₂O. Mp 27-28°. Bp 158-159°. pK_a 5.29. Absorbs CO₂ from air.
- EJ6800000
Hydrochloride (1:2): [333-93-7]
Needles (H₂O). Sol. H₂O; insol. MeOH. Mp 290°.
► EJ7280000
N-Ac: [5699-41-2]
 $C_6H_{14}N_2O$ 130.189
Mp 140-142°. Bp₈ 116-118°.
N,N'-Di-Ac: N,N'-1,4-Butanediyldisacetamide, 9CI
[3073-57-2]
 $C_8H_{16}N_2O_2$ 172.227
Cryst. (Me₂CO). Mp 137°.
N-Hexadecanoyl: N-Hexadecanoylputrescine. N-Palmitoylputrescine [126617-68-3]
 $C_{20}H_{42}N_2O$ 326.565
- Alkaloid from *Escherichia coli* transformed with a cosmid (pCSLF16) containing DNA. Powder.
N,N'-Dibenzoyl: N,N'-1,4-Butanediyldisbenzamide, 9CI. **Haplamide** [31991-78-3]
 $C_{18}H_{20}N_2O_2$ 296.368
Isol. from *Haplophyllum latifolium*. Cryst. (EtOH). Mp 176-177° (172-173°).
N-(2,3-Dihydroxybenzoyl): N-(4-Aminobutyl)-2,3-dihydroxybenzamide, 9CI. N-(2,3-Dihydroxybenzoyl)putrescine. **Aminocheilin** [114191-64-9]
 $C_{11}H_{16}N_2O_3$ 224.259
Prod. by *Azotobacter vinelandii*. Siderophore. Solid. Sol. H₂O.
N,N'-Bis(phenylacetyl): N,N'-1,4-Butanediyldis[benzenacetamide], 9CI. N,N'-Bis(phenylacetyl)putrescine. **Aglaudinine. Aglanthine** [102552-55-6]
 $C_{20}H_{24}N_2O_2$ 324.422
Alkaloid from *Aglaia edulis* and *Aglaia leptantha*. Needles (MeOH). Mp 162-163° Mp 175-178°. λ_{max} 222 (log ϵ 3.53); 260 (log ϵ 2.75) (MeOH).
N,N'-Bis(benzenesulfonyl): [97249-21-3]
 $C_{16}H_{20}N_2O_4S_2$ 368.477
Leaflets (EtOH). Mp 132°.
Bis(4-methylbenzenesulfonyl): [15544-47-5]
 $C_{18}H_{24}N_2O_4S_2$ 396.531
Mp 224°.
N-Cinnamoyl: see *N-Cinnamoylputrescine, C-456*
N,N'-Dicinnamoyl: see *N,N'-Dicinnamoylputrescine, D-362*
N-Me: *N-Methyl-1,4-butanediamine, 9CI* [14475-60-6]
[89690-09-5 (dihydrochloride)]
 $C_5H_{14}N_2$ 102.179
Constit. of tobacco. Cryst. (as dihydrochloride). Mp 177-179° (dihydrochloride).
N-Me, dibenzoyl:
 $C_{19}H_{22}N_2O_2$ 310.395
Mp 115.5°.
N,N-Di-Me: N,N-Dimethyl-1,4-butanediamine, 9CI
[3529-10-0]
 $C_6H_{16}N_2$ 116.206
Mp 142-143° (as dihydrochloride). Bp 150° Bp₃₅ 73°.
N,N'-Di-Me: N,N'-Dimethyl-1,4-butanediamine, 9CI. *1,4-Bis(methylamino)-butane* [16011-97-5]
 $C_6H_{16}N_2$ 116.206
Bp 168°.
N,N'-Di-Me, dibenzoyl: N,N'-1,4-Butanediyldis[N-methylbenzamide], 9CI
[16012-03-6]
 $C_{20}H_{24}N_2O_2$ 324.422
Mp 117°.
N-Tetra-Me: see *N,N,N',N'*-Tetramethyl-1,4-butanediamine, T-281
N-Et: *N-Ethyl-1,4-butanediamine, 9CI* [89690-10-8]
 $C_6H_{16}N_2$ 116.206

Cryst. (as dihydrochloride). Mp 220-221° (dihydrochloride).

N,N-Di-Et: N,N-Diethyl-1,4-butanediamine, 9CI

[27431-62-5]
C₈H₂₀N₂ 144.259
Bp₃ 55°. *n*_D²⁰ 1.4425.

N-(3-Methyl-2-butenyl): N-(3-Methyl-2-butenyl)putrescine. **N-Prenylputrescine**
C₉H₂₀N₂ 156.27

Trace alkaloid from *Eremosparton flaccidum*. Bp₁₀ 105-106°. Prob. struct. Struct. revised here from that originally proposed, N-(3-methyl-1-butenyl)-putrescine.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 291D; 292A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 465A; 465B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 376A (ir)

Hill, A.J. *et al.*, *J.A.C.S.*, 1939, **61**, 822-825 (synth, N-Ac)

Chambret, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1947, 1023-1026 (synth)

Schultz, H.P. *et al.*, *J.A.C.S.*, 1948, **70**, 2666-2667 (synth)

Merlis, V.M. *et al.*, *CA*, 1952, **46**, 7289 (N-Prenylputrescine, isol)

Jackson, E.L. *et al.*, *J.O.C.*, 1956, **21**, 1374-1375 (N-Ac)

Rama Sastry, B.V. *et al.*, *J.O.C.*, 1958, **23**, 1577-1578 (N,N-di-Et)

Lüttringhaus, A. *et al.*, *Chem. Ber.*, 1959, **92**, 1756-1765 (bisbenzenesulfonyl)

Org. Synth., Coll. Vol., 4, 1963, 819-823 (synth)
Smith, T.A. *et al.*, *Phytochemistry*, 1964, 3, 23-26 (biosynth)

Morris, D.R. *et al.*, *J. Biol. Chem.*, 1966, **241**, 3129-3135 (biosynth)

Roberts, J.D. *et al.*, *J.A.C.S.*, 1972, **94**, 2495-2500 (N-15 nmr)

Sarneski, J.E. *et al.*, *Anal. Chem.*, 1975, **47**, 2116-2124 (cmr)

Mayerl, F. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 127-133 (ms)

Albone, E.S. *et al.*, *J. Chem. Ecol.*, 1976, **2**, 101-111; 167-175 (occur)

Gaymans, R.J. *et al.*, *J. Polym. Sci., Polym. Chem. Ed.*, 1977, **15**, 537-545 (polym)

Nesmelova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 749-752; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 637-639 (Haplamide)

Eur. Pat., 1983, 77 106; *CA*, **99**, 123162f (nylon 4,6)

Cirrincone, G. *et al.*, *J.C.S. Perkin 2*, 1984, 1089-1091 (benzoyl derivs)

Frydman, B. *et al.*, *J.O.C.*, 1984, **49**, 2021-2023 (N-alkyl derivs)

Page, W.J. *et al.*, *J. Gen. Microbiol.*, 1988, **134**, 453-460 (Aminochelin, isol)

Golding, B.T. *et al.*, *Tet. Lett.*, 1988, **29**, 6651-6654 (N,N-di-Me)

Hamana, K. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur)

Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-747 (thermophilic eubacteria constits)

Linden, A. *et al.*, *Acta Cryst. C*, 1999, **55**, IUC9900046 (bisbenzenesulfonyl, cryst struct)

Saifah, E. *et al.*, *Phytochemistry*, 1999, **52**, 1085-1088 (Aglaiduline)

Thalladi, V.R. *et al.*, *Angew. Chem., Int. Ed.*, 2000, **39**, 918-922 (cryst struct)

Greger, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 616-620 (Aglanthine)

Brady, S.F. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1283-1286 (N-Hexadecanoylputrescine)

Pospieszna-Markiewicz, I. *et al.*, *Acta Cryst. E*, 2007, **63**, o3650 (cryst struct)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, BOS000; 1992, POK325

2-Butene-1,4-diamine B-423

1,4-Diamino-2-butene

[18231-61-3]
H₂NCH₂CH=CHCH₂NH₂
C₄H₁₀N₂ 86.136

(E)-form [40930-37-8]

[119874-79-2 (dihydrochloride)]
Potent irreversible inhibitor of kidney diamine oxidase. Bp 169° Bp₄₅ 92-95°.

N,N'-Dibenzoyl: [156544-32-0]

C₁₈H₁₈N₂O₂ 294.352
Cryst. (EtOH aq.). Mp 178.5-179.5°.

N,N'-Bis(benzenesulfonyl):

C₁₆H₁₈N₂O₄S₂ 366.461
Cryst. (EtOH). Mp 155°.

N,N'-Di-E-cinnamoyl: N',N'-Di-E-cinnamoyl-2,3-dehydroputrescine

[849462-51-7]
C₂₂H₂₂N₂O₂ 346.428
Prod. by the mushroom *Pholiota spumosa*.

N,N,N',N'-Tetra-Me: N,N,N',N'-Tetramethyl-2-butene-1,4-diamine. 1,4-Bis(-dimethylamino)-2-butene

[4559-79-9]
C₈H₁₈N₂ 142.244
d 0.81. Bp₇₃₅ 171-172° Bp₃₃ 74-76°. *n*_D²⁰ 1.4416. Props. may refer to a mixt. of geom. isomers.

(Z)-form [40794-72-7]

[114118-70-6 (dihydrochloride)]
Potent irreversible inhibitor of kidney diamine oxidase. Mp 292-300° (as dihydrochloride). Bp₁₅ 78°.

[111-52-4]

Amundsen, L. *et al.*, *J.A.C.S.*, 1951, **73**, 2118-2121 (E-form)

Ger. Pat., 1968, 1 274 116; *CA*, **69**, 105875s (N-tetra-Me)

Fiengenbaum, A. *et al.*, *Bull. Soc. Chim. Fr.*, 1973, 198-202 (synth)

Fabiano, E. *et al.*, *Synthesis*, 1987, 190-192 (Z-form)


Saljoughian, M. *et al.*, *J. Labelled Compd. Radiopharm.*, 1988, **25**, 313-328 (E-form, Z-form, synth, pmr, bibl)

He, Z. *et al.*, *Biochim. Biophys. Acta*, 1995, **1253**, 117-127 (E-form, Z-form, synth, pmr, activity)

Clericuzio, M. *et al.*, *Croat. Chem. Acta*, 2004, **77**, 605-611 (dicinnamoyl)

5-(2-Butenylidene)-3-ethyl-1,2,3,4,5,7a-hexahydro-4aH-1-pyridine-4,4a-diol B-424

[154593-87-0]



C₁₄H₂₁NO₂ 235.325


Prod. by *Streptomyces* sp. SCC 2313. Pale

yellow solid. [α]_D²¹ -68.6 (c.0.04 in H₂O). Similar to Pyridincine, P-932.

Hegde, V.R. *et al.*, *J. Antibiot.*, 1994, **47**, 110

5-(3-Butenyl)-2-pyridine-carboxylic acid, 9CI B-425

Dehydrofusaric acid. 5-(3-Butenyl)picolinic acid
[3626-76-4]



C₁₀H₁₁NO₂ 177.202

Isol. from *Fusarium lycopersici* and *Gibberella fujikuroi*. Plant growth inhibitor. Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O. Mp 121°. λ_{max} 260 (ε 3901) (MeOH) (Berdy).

Me ester: [78106-26-0]

C₁₁H₁₃NO₂ 191.229
Bp_{0,1} 110-120°. λ_{max} 225 (ε 7762); 264 (ε 3162) (MeOH) (Berdy).


Amide:

C₁₀H₁₂N₂O 176.218
Mp 147-148°.

Steiner, K. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 690 (synth)

Pitel, D.W. *et al.*, *Can. J. Biochem.*, 1970, **48**, 625 (biosynth)

5(4)-Butyl-2-chloro-1H-imidazole-4(5)-carboxaldehyde 4(5)-Butyl-2-chloro-5(4)-formyl-1H-imidazole B-426




C₈H₁₁ClN₂O 186.64

Alkaloid from *Jatropha curcas*. Semi-solid.

Das, B. *et al.*, *Indian J. Chem., Sect. B*, 2005, **44**, 1119-1120 (isol, pmr, cmr)

Butylcycloheptylprodigiosine B-427



C₂₅H₃₃N₃O 391.555


Following conflicting reports the structure was finally confirmed, in 2007, as the original assignment and distinct from Streptorubin B, S-572. Prod. by *Streptomyces* sp. Y-42 and *Streptomyces abikoensis*. Dark purple solid (as hydrochloride). Mp ca.170-175° (hydrochloride).

[56247-03-1]

Gerber, N.N. *et al.*, *J. Antibiot.*, 1975, **28**, 194-199 (*isol, pmr, activity*)
 Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (*rev*)
 Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 1929-1945 (*synth, pmr, cmr, struct*)
 Reeves, J.T. *et al.*, *Org. Lett.*, 2007, **9**, 1879-1881 (*synth*)

2-Butyldecahydroquinoline, 9CI B-428


Pumiliotoxin C_{III}. *Dendrobates histrionicus* Alkaloid I [63983-60-8]



C₁₃H₂₅N 195.347
 Minor alkaloid from skin extracts of *Dendrobates histrionicus* (Dendrobati-
 dae).

Daly, J.W. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1128 (*isol, ms, struct*)

7-Butyl-1,2-dichloro-6,8-dihydroxy-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI B-429




C₁₅H₁₃Cl₂NO₄ 342.177
 Prod. by *Streptomyces rimosus*.

1-Dechloro-7-Butyl-2-chloro-6,8-dihydroxy-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI

C₁₅H₁₄ClNO₄ 307.733
 Prod. by *Streptomyces rimosus*. Powder. λ_{max} 240 ; 295 ; 340 (MeCN aq.).
Pat. Coop. Treaty (WIPO), 1998, 98 25 931; *CA*, **129**, 52003e (*isol*)
 Trew, S.J. *et al.*, *J. Antibiot.*, 2000, **53**, 1-11 (*isol*)


5-Butyl-6,8-diethyloctahydroindolizine, 9CI B-430

5-Butyl-6,8-diethylindolizidine. *Dendrobates Alkaloid 237L*. *Indolizidine 237L* [185417-26-9]



C₁₆H₃₁N 237.428
 Tentative struct. Alkaloid from skin extracts of a Panamanian population of the frog *Dendrobates pumilio*.
 Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 2 (*isol, ms*)

3-Butyl-2,5-dimethylpyrazine B-431




C₁₀H₁₆N₂ 164.25
 Isol. from the ant *Iridomyrmex purpureus* and the fly *Anastrepha fraterculus*. Prod. by various marine bacteria. Bp₁₂ 101°.

Zavylov, S.I. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1973, 707 (*synth*)
 Bus, J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1973, **92**, 123 (*ir*)
 Flament, I. *et al.*, *Bull. Soc. Chim. Belg.*, 1979, **88**, 941 (*synth, ms*)
 Cavill, G.W.K. *et al.*, *Insect Biochem.*, 1984, **14**, 505 (*isol*)
 Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*marine isol*)
 Dickschat, J.S. *et al.*, *J. Chem. Ecol.*, 2005, **31**, 925-947 (*marine isol*)

2-Butyl-6-hydroxymethyl-3,4,5-piperidinetriol B-432

2-Butyl-3,4,5-trihydroxy-6-hydroxymethylpiperidine. 1-C-(5-Amino-5-deoxygalactopyranosyl)butane




C₁₀H₂₁NO₄ 219.28

(2S,3S,4R,5S,6R)-form
1β-C-Butyl-1-deoxygalactonojirimycin
 Alkaloid from *Adenophora* spp. [α]_D -12.8 (c, 0.36 in H₂O) (natural). [α]_D -14.8 (c, 0.39 in H₂O) (synthetic).

Ikeda, K. *et al.*, *Carbohydr. Res.*, 2000, **323**, 73-80

6-tert-Butyl-1-hydroxy-3-[2-(methylthio)ethyl]-2(1H)-pyrazinone B-433

1-Hydroxy-6-(1,1-dimethylethyl)-3-[2-(methylthio)ethyl]-2(1H)-pyrazinone




C₁₁H₁₈N₂O₂S 242.341
 Metab. from *Aspergillus flavus*. Mp 74-76°.

MacDonald, J.C. *et al.*, *Can. J. Biochem.*, 1972, **50**, 543 (*isol, pmr, ms, struct*)

3-Butyloctahydro-5-methylindolizine B-434

3-Butyl-5-methylindolizidine [42607-24-9]



C₁₃H₂₅N 195.347

(3R,5R,8aR)-form [53447-41-9]
 Isol. from dendrobatic frog skin. Pale yellow oil. [α]_D²² -97.1 (c, 0.12 in MeOH).
Hydrochloride: [94535-28-1]
 [α]_D²⁰ -69.2 (c, 0.55 in MeOH).

(3R,5S,8aR)-form [92841-44-6]
 Isol. from thief ant, *Solenopsis* spp. venom. Pale yellow oil. [α]_D²⁵ -36.5 (c, 0.14 in MeOH).

(3R,5S,8aS)-form Monomorine I [53447-44-2]
 Trail pheromone of Pharaoh ant, *Monomorium pharaonis*. Pale yellow oil. Bp₂₇ 119°. [α]_D²⁵ +35.1 (c, 1.33 in hexane).

(3S,5R,8aR)-form [94535-27-0]
 [α]_D²⁰ -35.8 (c, 1.35 in hexane).

(3S,5R,8aS)-form [53447-42-0]
 [α]_D²⁶ +68.5 (c, 0.145 in MeOH).

(3S,5S,8aS)-form Dendrobates Alkaloid 195B. Indolizidine 195B

[118015-64-8]
 Isol. from skin of Colombian poison-frog *Dendrobates histrionicus*. Oil. [α]_D²⁶ +99.6 (c, 0.23 in MeOH) (+65).
Hydrochloride: [α]_D²⁴ +36 (c, 0.52 in MeOH).

[53447-43-1, 79465-32-0, 82262-91-7, 107597-44-4]

Ritter, F.J. *et al.*, *Experientia*, 1973, **29**, 530 (*isol, ms*)

Sonnet, P.E. *et al.*, *J.O.C.*, 1974, **34**, 2662
 Sonnet, P.E. *et al.*, *J. Het. Chem.*, 1975, **12**, 289; 1979, **16**, 1041 (*synth, pmr, cmr*)

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)

Spande, Th.F. *et al.*, *Experientia*, 1981, **37**, 1242 (*config*)

Stevens, R.V. *et al.*, *Chem. Comm.*, 1982, 102 (*synth*)

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1984, **10**, 1233 (*isol*)

Royer, J. *et al.*, *J.O.C.*, 1985, **50**, 670 (*synth, pmr, cmr, ms*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1986, **42**, 3453 (*isol, cmr, struct*)

Jefford, C.W. *et al.*, *Helv. Chim. Acta*, 1989, **72**, 1749 (*synth*)

Yamazaki, N. *et al.*, *J.A.C.S.*, 1989, **111**, 1396 (*synth, ir, pmr, cmr, ms*)

Watanabe, Y. *et al.*, *J.O.C.*, 1989, **54**, 4088 (*synth, pmr, cmr, ms*)


Momose, T. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 2072 (*synth*)

Nagasaka, T. *et al.*, *Heterocycles*, 1, Spec. Issue, 1990, **30**, 561 (*synth*)

- Vavrecka, M. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 438 (*synth*)
 Ito, M. *et al.*, *J.A.C.S.*, 1991, **113**, 3513 (*synth*)
 Saliou, C. *et al.*, *Tet. Lett.*, 1991, **32**, 3365 (*synth*)
 Takahata, H. *et al.*, *Heterocycles*, 1993, **36**, 2777 (*synth, ir, pmr, cmr*)
 Higashiyama, K. *et al.*, *J.C.S. Perkin 1*, 1994, 351-353 (*synth, pmr, Indolizidine 195B, Monomarine I*)
 Somfai, P. *et al.*, *Acta Chem. Scand.*, 1997, **51**, 1024-1029 (*synth*)
 Berry, M.B. *et al.*, *Chem. Comm.*, 1997, 2141-2142 (*synth*)
 Muraoka, O. *et al.*, *J.C.S. Perkin 1*, 1997, 113 (*synth*)
 Momose, T. *et al.*, *J.C.S. Perkin 1*, 1997, 1315 (*synth*)
 Mori, M. *et al.*, *J.O.C.*, 1998, **63**, 4832-4833 (*synth*)
 Celimene, C. *et al.*, *Tetrahedron*, 1998, **54**, 10457-10468 (*synth*)
 Riesinger, S.W. *et al.*, *Eur. J. Org. Chem.*, 1999, 3277-3280; 2000, 867 (*synth*)
 Yuguchi, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 2004, **77**, 1031-1032 (*synth*)
 Toyooka, N. *et al.*, *J.O.C.*, 2008, **73**, 4575-4577 (*synth*)
 Kuhakarn, C. *et al.*, *Tetrahedron*, 2008, **64**, 1663-1670 (*synth*)

3-Butyloctahydro-5-(4-pentenyl)indolizine B-435

3-Butyl-5-(4-pentenyl)indolizidine



C₁₇H₃₁N 249.439


(3S,5R,8aS)-form [128855-21-0]

Alkaloid from venom of the ant *Monomorium smithii*.

Jones, T.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 375 (*isol, cmr, synth, struct*)

3-Butyloctahydro-5-propa- B-436

noylindolizine
 1-(3-Butyloctahydro-5-indoliziny)-1-propanone, 9CI. 3-Butyl-5-propanoylindolizidine



(3R,5R,8aR)-form

C₁₅H₂₇NO 237.384

(3R,5R,8aR)-form Myrmicarin 237B [165877-93-0]

Alkaloid from the poison gland secretion of the African ant *Myrmecaria eumenoides*. Oil. [α]_D¹⁹ +14.1 (c, 0.81 in heptane).

(3R,5R,8aS)-form

Synthetic. [α]_D¹⁹ +12.6 (c, 2.39 in heptane).

(3R,5S,8aR)-form Myrmicarin 237A [165877-92-9]


From *Myrmecaria eumenoides*. [α]_D²⁰ -123 (c, 1.46 in heptane).

(3R,5S,8aS)-form

Synthetic. [α]_D²⁰ -5.6 (c, 1.64 in heptane). Francke, W. *et al.*, *Annalen*, 1995, 965 (*isol, ir, pmr, cmr, ms, synth, struct*)

8-Butyloctahydro-5-propyl- B-437

1H-indolizine, 9CI
 8-Butyl-5-propylindolizidine. *Indolizidine 223V*. *Dendrobates Alkaloid 223V* [194344-43-9] [194344-44-0]




C₁₅H₂₉N 223.401

Alkaloid from the poison frog *Dendrobates pumilio*.

Toyooka, N. *et al.*, *Tetrahedron*, 2005, **61**, 1187-1198; 5139 (*synth, ms*)

3-Butyloctahydro-5-propyl-8- B-438

indolizine
 3-Butyloctahydro-8-hydroxy-5-propylindolizine. 3-Butyl-8-hydroxy-5-propylindolizidine



C₁₅H₂₉NO 239.4


(3R*,5S*,8R,8aR*)-form [929631-68-5]

Alkaloid from the venom of *Myrmecaria melanogaster*.

Jones, T.H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 160-168 (*isol, synth, ir, ms*)

3-Butyloctahydro-1H-pyrro- B-439

lo[1,2-a]azepine
 3-Butyllehmizidine



C₁₃H₂₅N 195.347


(3R,9aR)-form [929631-67-4]

Alkaloid from the venom of *Myrmecaria melanogaster*.

Jones, T.H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 160-168 (*isol, synth, ir, ms*)

4-tert-Butyl-2-oxazolidinone B-440

4-(1,1-Dimethylethyl)-2-oxazolidinone, 9CI [134668-36-3] [142758-90-5]



C₇H₁₃NO₂ 143.185

(R)-form [142618-93-7]

Cryst. (toluene). Mp 119-120°. [α]_D²⁰ +18.8 (c, 1.0 in EtOH) (>99% ee).

(S)-form [54705-42-9]

Cryst. (Et₂O/hexane) or needles (EtOAc/petrol). Mp 120° (110-111°). [α]_D²⁰ -18 (c, 0.50 in EtOH).

(±)-form

Alkaloid from *Strychnos cathayensis*.

Hassner, A. *et al.*, *Tetrahedron*, 1974, **30**, 2613-2621 (*S-form, synth, pmr*)

Ishizuka, T. *et al.*, *Chem. Lett.*, 1992, 991-994 (*synth, resoln*)

Drauz, K. *et al.*, *Chem. Eur. J.*, 1995, **1**, 538-540 (*R-form, synth, pmr*)


Tietze, L.F. *et al.*, *Chem. Eur. J.*, 1996, **2**, 139-148 (*S-form, synth, pmr*)

Takacs, J.M. *et al.*, *J.O.C.*, 1998, **63**, 2742-2748 (*R-form, synth*)

Cheng, M.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 235-239 (*isol*)

N-[2-(4-Butylphenyl)ethyl]-3- B-441

(4-hydroxy-3-methoxyphenyl)-2-propanamide
 4-Butyl-N-feruloylphenethylamine [402499-33-6]




C₂₂H₂₇NO₃ 353.46

Alkaloid from *Cuscuta reflexa*. α-Glucosidase inhibitor. Solid. λ_{max} 220 (log ε 4.1); 292 (log ε 3.95); 325 (log ε 4.42) (MeOH).

Anis, E. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 112-114 (*isol, pmr, cmr, ms*)

2-Butylpiperidine B-442

[72939-22-1]



C₉H₁₉N 141.256

(+)-form

[α]_D¹⁵ +15.7.

Hexachloroplatinate: Mp 131-132°.

(-)-form[α]_D -18.7.**Tartrate salt:**


Needles. Mp 41-42°. Forms a dihydrate.

(±)-form [68144-45-6]Liq. Bp 191-193° Bp₁₄ 75°.**Hydrochloride:** Mp 191-193° (181-182°).**Hydrobromide:** Mp 196-197°.**N-Me:**Liq. Mp 88° (as picrate). Bp₁₅ 75-80°.**(ξ)-form****N-Me: 2-Butyl-1-methylpiperidine**C₁₀H₂₁N 155.283Trace alkaloid from oil of *Pimpinella acuminata* (Apiaceae). Opt. activity not reported.


[68474-12-4]

Löffler, K. *et al.*, *Ber.*, 1907, **40**, 1310 (*synth, resoln*)Diels, O. *et al.*, *Annalen*, 1937, **530**, 68 (*synth*)Astier, A. *et al.*, *Tet. Lett.*, 1978, 2051 (*synth*)Ashraf, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1979, **22**, 79 (*isol, deriv*)Scully, F.E. *et al.*, *J.O.C.*, 1980, **45**, 1515 (*synth, pmr, ir*)Pirkle, W.H. *et al.*, *J.O.C.*, 1984, **49**, 2504 (*resoln*)**5-Butyl-8-propylindolizidine B-443****Octahydro-5-butyl-8-propyl-1H-indolizine, 9CI. Dendrobates Alkaloid 223J. Indolizidine 223J**

[194344-81-5]

C₁₅H₂₉N 223.401Tentative struct. assigned. Minor alkaloid from a frog *Dendrobates* sp. Oil. Mp 192-195° (as hydrochloride) (synthetic). [α]_D²⁶ -131.9 (c, 0.5 in CHCl₃).Toyooka, N. *et al.*, *Tetrahedron*, 1997, **53**, 9553-9574 (*synth, ir, pmr, cmr, ms*)Michel, P. *et al.*, *J.O.C.*, 2000, **65**, 8908-8918 (*synth*)**3-Butylpyridine, 9CI B-444****Fusarinin†. Fusarin†**

[539-32-2]

C₉H₁₃N 135.208Isol. from the fungus *Fusarium heterosporum*. Insol. H₂O. d₄²⁰ 0.92. Bp 205-208° Bp₇ 82°.


▶ US4056000

Hydrochloride:Cryst. (H₂O). Mp 126°.**Picrate:**

Yellow needles (EtOH). Mp 89-90°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 245A (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 1517B (*ir*)Maas, E. *et al.*, *Ber.*, 1914, **47**, 1164-1173 (*synth*)Yabuta, Y. *et al.*, *CA*, 1940, **34**, 3314 (*isol*)Frank, R.L. *et al.*, *J.A.C.S.*, 1948, **70**, 3482-3483 (*synth*)Tamari, K. *et al.*, *Nippon Nogei Kagaku Kaishi*, 1948, **22**, 16-17; *CA*, **46**, 5143 (*synth*)Plattner, P.A. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1379-1392 (*synth, ir, uv*)Kyte, C.T. *et al.*, *J.C.S.*, 1960, 4454-4472 (*synth, uv*)Chelucci, G. *et al.*, *J. Het. Chem.*, 1988, **25**, 1761-1765 (*synth, pmr*)**5-Butyl-2-pyridinecarboxylic acid, 9CI B-445****5-Butylpicolinic acid. Fusaric acid**

[536-69-6]

C₁₀H₁₃NO₂ 179.218Isol. from *Fusarium lycopersici*, *Fusarium oxysporum*, *Fusarium vasinfectum* and *Gibberella fujikuroi*. Plant growth inhibitor. Shows antihypertensive and insecticidal activity. Dopamine β-hydroxylase inhibitor. Plates (petrol). Sol. MeOH, butanol; poorly sol. H₂O. Mp 108-109°. Log P 2.88 (calc). λ_{max} 260 (MeOH) (Berdy).▶ LD₅₀ (mus, ivn) 100 mg/kg, LD₅₀ (mus, ipr) 80 mg/kg, LD₅₀ (mus, orl) 230 mg/kg. US5625000**Picrate:** Mp 83-89°.**Me ester:** [17072-92-3]C₁₁H₁₅NO₂ 193.245Bp 98-99°. λ_{max} 225 (ε 5888); 264 (ε 2455) (MeOH) (Berdy).**Amide: Bupicomide, INN, USAN. Sch 10595**

[22632-06-0]

C₁₀H₁₄N₂O 178.233


Dopamine β-hydroxylase inhibitor.

Antihypertensive agent. Mp 127-128°.

Log P 1.88 (calc).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 786D (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 319C (*nmr*)Grove, F. *et al.*, *J.C.S.*, 1958, 1236 (*isol*)Chumakov, Y.I. *et al.*, *Tet. Lett.*, 1965, 129 (*synth, ir*)Vogt, H. *et al.*, *Tet. Lett.*, 1966, 5887 (*synth*)U.S. Pat., 1970, 3 519 717; *CA*, **73**, 59319w (*bupicomide*)Korduba, C.A. *et al.*, *J. Pharmacol. Exp. Ther.*, 1973, **184**, 671 (*bupicomide*)Symchowicz, S. *et al.*, *J. Pharmacol. Exp. Ther.*, 1974, **191**, 324 (*bupicomide*)Tschesche, R. *et al.*, *Chem. Ber.*, 1978, **111**, 3502 (*synth, spectra*)Medvedeva, T.M. *et al.*, *Khim.-Farm. Zh.*, 1978, **12**, 66 (*synth, pharmacol*)Piesche, L. *et al.*, *Pharmazie*, 1979, **34**, 332 (*pharmacol*)Langhals, E. *et al.*, *Annalen*, 1982, 930 (*synth*)Schroetter, E. *et al.*, *Pharmazie*, 1984, **39**, 155 (*synth*)Sagi, M. *et al.*, *Heterocycles*, 1989, **29**, 2249 (*synth*)Waldner, A. *et al.*, *Synth. Commun.*, 1989, **19**, 2371 (*synth, pmr*)*Martindale, The Extra Pharmacopoeia, 30th edn.*, Pharmaceutical Press, 1993, 841 (*Fusaric acid*)Koyama, J. *et al.*, *Heterocycles*, 1994, **38**, 1595 (*synth*)Renslo, A.R. *et al.*, *J.O.C.*, 1998, **63**, 7840-7850 (*synth, Me ester, ir, pmr, cmr*)Song, J.J. *et al.*, *J.O.C.*, 2001, **66**, 605-608 (*synth*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, BSI000**2-Butyl-3,4,5,6-tetrahydropyridine B-446****2-Butyl-1-azacyclohexene**

[1462-94-8]

C₉H₁₇N 139.24Isol. from the mushroom *Tylophilus* sp. Toxin. Oil.Hoffmann, R.V. *et al.*, *J.O.C.*, 1988, **53**, 3316-3321 (*synth*)Watanabe, R. *et al.*, *Tet. Lett.*, 2002, **43**, 6501-6504 (*isol*)**2-Butynedioic acid, 9CI B-447****Acetylenedicarboxylic acid, 8CI. Ethynedicarboxylic acid**

[142-45-0]

HOCC≡CCOOH

C₄H₂O₄ 114.057


Monomer for acetylenic polyesters.

Cryst. + 2H₂O. Mp 179°. pK_{a1} 1.23; pK_{a2} 2.53 (28°).**Diamide: Butynediamide. Aquamycin.****Cellocidin. Lenamycin**

[543-21-5]


C₄H₄N₂O₂ 112.088Isol. from *Streptomyces chibaensis* and *Streptomyces reticuli*. Active against gram-positive and -negative bacteria and HeLa cells. Cryst. (MeOH aq.). Mp 216-218° (dec.). Unstable in alkaline solns., evolving NH₃. λ_{max} 299 (ε 3248) (dil. NaOH) (Derrep). λ_{max} 216 (ε 9150) (H₂O) (Derrep).▶ LD₅₀ (mus, orl) 89.2 mg/kg. LD₅₀ (mus, skn) 667 mg/kg. AO9900000Suzuki, S. *et al.*, *J. Antibiot., Ser. A*, 1958, **11**, 81 (*isol, amide*)**Buxafuranamide****B-448**

[123871-27-2]



C₃₅H₄₈N₂O₅ 576.775Alkaloid from *Buxus papilosa* (Buxaceae). [α]_D²⁴ +190 (CHCl₃).Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1990, **53**, 319-324 (isol, pmr, cmr, struct)**O¹⁰-Buxafuranamine****B-449**



[142674-57-5]

C₃₅H₄₈N₂O₅ 576.775Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Gum. [α]_D²⁰ -126.2 (c, 1.3 in CHCl₃).*Deoxy: 6-Dehydroxy-O¹⁰-buxafuranamine*

[142674-58-6]

C₃₅H₄₈N₂O₄ 560.775Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Gum. [α]_D²⁰ -333 (c, 1.5 in CHCl₃).Atta-ur-Rahman, et al., *Heterocycles*, 1992, **34**, 157-171 (isol, uv, ir, pmr, ms, struct)**O⁶-Buxafuranamine****B-450**

[142674-55-3]

C₃₅H₄₈N₂O₅ 576.775Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Amorph. solid. [α]_D²⁰ +20 (c, 0.7 in CHCl₃).Atta-ur-Rahman, et al., *Heterocycles*, 1992, **34**, 157-171 (isol, uv, ir, pmr, ms, struct)**Buxahyrcanine****B-451**C₂₆H₄₆N₂O 402.662

Parent not known.

N³-(2-Methylpropanoyl): N-Isobutyryl-buxahyrcanine

[552291-03-9]

C₃₀H₅₂N₂O₂ 472.753Alkaloid from the leaves of *Buxus hyrcana*. Amorph. powder (CHCl₃).Mp 234-235°. [α]_D²⁹ +14 (c, 0.12 in CHCl₃). λ_{max} 206 (log ε 4.33) (MeOH).*N³-Tigloyl: N-Tigloylbuxahyrcanine*

[552291-02-8]

C₃₁H₅₂N₂O₂ 484.764Alkaloid from the leaves of *Buxus hyrcana*. Acetylcholinesterase inhibitor. Amorph. powder (MeOH/C₆H₆). Mp 245.5°. [α]_D²⁹ +62 (c, 0.1 in CHCl₃). λ_{max} 206 (log ε 4.33) (MeOH).*N³-Benzoyl: N-Benzoylbuxahyrcanine*


[552291-00-6]

C₃₃H₅₀N₂O₂ 506.77Alkaloid from the leaves of *Buxus hyrcana*. Amorph. powder (MeOH). Mp 239.7°. [α]_D²⁹ +15 (c, 0.14 in CHCl₃). λ_{max} 223 (log ε 4.2) (MeOH).Choudhary, M.I. et al., *J. Nat. Prod.*, 2003, **66**, 739-742 (isol, pmr, cmr, cryst struct)**Buxalfine****B-452***Buxalphine*



[1354-96-7]

C₃₃H₄₆N₂O₃ 518.738Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). [α]_D +56 (c, 0.87 in CHCl₃). [α]_D +61 (EtOH).*Dihydro: Buxalfol*Mp 244-249°. [α]_D +81 (c, 0.26 in CHCl₃).*Octahydro: Mp 152-155°. [α]_D +49 (c, 0.72 in CHCl₃).*Tomko, J. et al., *Chem. Zvesti*, 1964, **18**, 721-731; *CA*, **62**, 4321d (isol)Votický, Z. et al., *CA*, 1967, **67**, 11668h (isol)**Buxalongifolamidine****B-453**

[149472-47-9]

C₃₅H₅₀N₂O₅ 578.79Alkaloid from leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. [α]_D²² +23 (c, 0.085 in CHCl₃).Atta-ur-Rahman, et al., *Phytochemistry*, 1993, **32**, 1059-1063 (isol, uv, ir, pmr, cmr, ms, struct)**Buxaltine H****B-454***Buxaltine*

[25650-66-2]

C₃₅H₄₈N₂O₃ 544.776Struct. not well established. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 188-191°.Döpke, W. et al., *Pharmazie*, 1969, **24**, 649; *CA*, **72**, 43929p (isol, ir, ms, struct)**Buxamine I****B-455***4,4,14-Trimethyl-B-(9a)-homo-19-norpregna-9(11),9a-diene-3,20-diamine, 9CI*C₂₄H₄₀N₂ 356.593The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Buxamine I, is unknown.*N³-Me: Buxamine G. Buxenine G. Norbuxamine*

[6246-56-6]

C₂₅H₄₂N₂ 370.62Alkaloid from *Buxus sempervirens* (Buxaceae). Noncryst. Mp 298-300° dec. (as dihydroiodide). [α]_D²⁰ +20 (c, 0.51 in CHCl₃).*N³-Me, N²⁰-Ac: N²⁰-Acetylbuxamine G*

[307317-34-6]

C₂₇H₄₄N₂O 412.657Alkaloid from *Buxus sempervirens*. Gum. [α]_D +20 (c, 0.41 in MeOH). λ_{max} 238 ; 245 ; 254 (sh) (no solvent reported).*N³,N³-Di-Me: Buxamine E. Buxamine*

[14317-17-0]

C₂₆H₄₄N₂ 384.647Alkaloid from *Buxus sempervirens*, *Buxus balearica*, *Buxus microphylla* and *Buxus harlandi* (Buxaceae). HIV reverse transcriptase (HIV-rt) inhibitor. [α]_D²⁰ +32 (c, 0.57 in CHCl₃).*N³,N³-Di-Me, N²⁰-Ac: N²⁰-Acetylbuxamine E*

[76199-87-6]

C₂₈H₄₆N₂O 426.684Alkaloid from *Buxus sempervirens*. Gum. [α]_D +8 (c, 0.53 in MeOH). λ_{max} 238 ; 245 ; 254 (sh) (no solvent reported).*N³,N²⁰-Di-Me: Papilamine*

[96910-87-1]

C₂₆H₄₄N₂ 384.647Alkaloid from the leaves of *Buxus papilosa* (Buxaceae). Cryst. Mp 240° dec. [α]_D +23.3. λ_{max} 210 ; 237 ; 245 ; 254 (No solvent reported).*N²⁰,N²⁰-Di-Me: Buxamine F*

[231628-56-1]

C₂₆H₄₄N₂ 384.647Alkaloid from *Buxus sempervirens*. Amorph. solid. [α]_D²⁰ +49 (c, 0.4 in CHCl₃). λ_{max} 231 (sh) (log ε 3.2); 238 (log ε 3.4); 245 (log ε 2.9); 252 (sh) (log ε 3.1) (MeOH).*N²⁰,N²⁰-Di-Me, N³-benzoyl: Buxabenza-*

- midienine**
[107259-38-1]
C₃₃H₄₈N₂O 488.755
Alkaloid from leaves of *Buxus papillosa* (Buxaceae). Amorph. powder. [α]_D +6 (c, 1.59 in CHCl₃).
- N³,N³,N²⁰-Tri-Me: Buxamine B. Papilicine**
[76186-32-8]
C₂₇H₄₆N₂O 398.674
Alkaloid from the leaves of *Buxus harlandi* and from *Buxus papillosa* (Buxaceae). Mp 248°. [α]_D²³ +61.4 (c, 0.57 in CHCl₃). [α]_D +47.6 (CHCl₃). Papilicine, which was descr. as a gum, is of unclear stereochem. It could be the C-20 epimer of Buxamine B, which is unlikely, but was probably impure Buxamine B.
- N³,N³,N²⁰-Tri-Me, N²⁰-formyl: N-Formylpapilicine. N-Formylbuxamine B**
[115713-19-4]
C₂₈H₄₆N₂O 426.684
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. [α]_D +36 (c, 0.35 in CHCl₃).
- N³,N³,N²⁰-Tri-Me, N²⁰-Ac:**
Cryst. (Me₂CO). Mp 186-190°.
- N³,N²⁰,N²⁰-Tri-Me: Buxamine C**
[51059-62-2]
C₂₇H₄₆N₂O 398.674
Alkaloid from *Buxus papillosa* (Buxaceae). Cryst. (Me₂CO). Mp 153-155°. [α]_D +24 (c, 1.12 in CHCl₃).
- N-Tetra-Me: Buxamine A**
[36127-40-9]
C₂₈H₄₈N₂O 412.701
Alkaloid from *Buxus madagascariensis* (Buxaceae). Mp 134°. [α]_D +40 (CHCl₃).
- 9a,10-Dihydro, N²⁰,N²⁰-di-Me, N³-benzoyl: Buxabenzamidine**
[109305-86-4]
C₃₃H₅₀N₂O 490.771
Alkaloid from leaves of *Buxus papillosa* (Buxaceae). Amorph. [α]_D +47 (c, 0.73 in CHCl₃).
- 1,2-Didehydro, N²⁰,N²⁰-di-Me, N³-benzoyl: Hyrcatrienine**
C₃₃H₄₆N₂O 486.739
Alkaloid from the leaves of *Buxus hyrcana*. Amorph. solid. [α]_D²⁴ +47.9 (c, 0.2 in CHCl₃). λ_{max} 232 ; 289 (CHCl₃).
- 30-Nor, N-tetra-Me: 30-Norbuxamine A**
[129225-36-1]
C₂₇H₄₆N₂O 398.674
Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Amorph. [α]_D -13 (CHCl₃).
- [89164-27-2 , 11037-97-1]
Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*Buxamine, Norbuxamine, isol, uv, ir, pmr*)
Kupchan, S.M. *et al.*, *Tet. Lett.*, 1964, 3145-3150 (*Buxenine G, uv, ir, pmr, struct*)
Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*Buxus balearic constis*)
Khuong-Huu-Laine, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 558-560 (*Buxamine A*)
Puckett, R.T. *et al.*, *J.C.S. (B)*, 1971, 935-941 (*Buxenine G, cryst struct*)

- Shamma, M. *et al.*, *Phytochemistry*, 1973, **12**, 2051-2054 (*Buxamine C*)
Vassová, A. *et al.*, *Chem. Zvesti*, 1980, **34**, 706-711 (*Buxamine B*)
Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 1984, **39**, 127-128; 1985, **40**, 567-568 (*Papilicine, Papilamine*)
Choudhary, M.I. *et al.*, *Tetrahedron*, 1986, **42**, 5747-5752 (*Buxabenzamidienine*)
Choudhary, M.I. *et al.*, *J. Nat. Prod.*, 1987, **50**, 84-88 (*Buxabenzamidine*)
Choudhary, M.I. *et al.*, *Phytochemistry*, 1988, **27**, 1561-1562 (*N-Formylpapilicine*)
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1990, **29**, 1293-1296 (*30-Norbuxamine A*)
Sun, H.D. *et al.*, *J. Nat. Prod.*, 1996, **59**, 525-528 (*activity*)
Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1999, **62**, 665-669 (*Buxamine F*)
Loru, F. *et al.*, *Phytochemistry*, 2000, **54**, 951-957 (*N²⁰-Acetyl-buxamines*)
Choudhary, M.I. *et al.*, *Chem. Biodiversity*, 2006, **3**, 1039-1052 (*Hyrcatrienine*)

Buxaminol I

B-456

- Buxodienine I**
As Buxamine I, B-455 with R = OH
C₂₄H₄₀N₂O 372.593
The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compl., Buxaminol I, is unknown.
- N²⁰-Me, N³-benzoyl, 16-Ac: Nor-16-acetoxybuxabenzamidienine**
[126025-06-7]
C₃₄H₄₈N₂O₃ 532.765
Alkaloid from leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. [α]_D²⁰ +17.5 (c, 0.4 in CHCl₃).
- N³,N³-Di-Me: Buxaminol E. Buxaminol. Buxodienine E**
[14155-76-1]
[50861-49-9 (Buxodienine E)]
C₂₆H₄₄N₂O 400.646
Alkaloid from *Buxus balearica* and *Buxus sempervirens*; main alkaloid of *Buxus sempervirens* var. *bullata* (Buxaceae). Cryst. (MeOH) or amorph. Mp 198°. [α]_D +40 (CHCl₃).
- N³,N³-Di-Me, N²⁰-formyl: N²⁰-Formyl-buxaminol E**
[307317-32-4]
C₂₇H₄₄N₂O₂ 428.657
Alkaloid from *Buxus sempervirens*. Gum. [α]_D +13 (c, 0.15 in MeOH). λ_{max} 238 ; 245 ; 254 (sh) (no solvent reported).
- N³,N³-Di-Me, N²⁰-benzoyl: N-Benzoyl-buxodienine E**
[49776-67-2]
C₃₃H₄₈N₂O₂ 504.754
Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 235-237°. [α]_D²⁴ +34.6 (CHCl₃).
- N³,N³-Di-Me, N²⁰-benzoyl, O-Ac: N-Benzoyl-O-acetyl-buxodienine E**
[49776-68-3]
C₃₅H₅₀N₂O₃ 546.792
Alkaloid from *Buxus sempervirens* (Buxaceae). Amorph. [α]_D²⁴ -6 (CHCl₃).
- N³,N³-Di-Me, O-(4-hydroxy-3,5-dimethoxybenzoyl): 16-O-Syringoylbuxaminol E**
[307317-33-5]


- C₃₅H₅₂N₂O₃ 580.806
Alkaloid from *Buxus sempervirens*. Gum. [α]_D +83 (c, 0.53 in EtOH). λ_{max} 221 ; 236 ; 245 ; 254 (sh) ; 278 (no solvent reported).
- N³,N²⁰-Di-Me, N³-Ac: [204449-46-7]**
C₂₈H₄₆N₂O₂ 442.684
Alkaloid from *Buxus papillosa*. Gum. [α]_D +124 (c, 0.95 in CHCl₃). Synonym assigned in ref. is incorrect. λ_{max} 236 (sh) ; 238 ; 245 ; 251 (sh) (MeOH).
- N²⁰,N²⁰-Di-Me, N³-benzoyl, O-Ac: 16-Acetoxybuxabenzamidienine**
[107259-39-2]
C₃₅H₅₀N₂O₃ 546.792
Alkaloid from leaves of *Buxus papillosa* and *Buxus longifolia*. Amorph. [α]_D +6 (c, 1.02 in CHCl₃). λ_{max} 225 (log ε 4.28); 230 (log ε 4.3); 237 (log ε 4.32); 245 (log ε 4.35); 254 (log ε 4.4) (MeOH).
- N³,N³,N²⁰-Tri-Me: Buxaminol B**
[58672-77-8]
C₂₇H₄₆N₂O 414.673
Alkaloid from *Buxus sempervirens* var. *bullata* and *Buxus papillosa* (Buxaceae). Cryst. (Me₂CO). Mp 225°. [α]_D²⁵ +20 (c, 0.59 in MeOH). λ_{max} 236 (log ε 4.45); 246 (log ε 4.49); 255 (log ε 4.28); 273 (log ε 2.68) (EtOH).
- N³,N²⁰,N²⁰-Tri-Me: Buxaminol C**
[191288-32-1]
C₂₇H₄₆N₂O 414.673
Alkaloid from the roots of *Buxus sempervirens* (Buxaceae). Gum. [α]_D²⁰ +88 (c, 0.6 in CHCl₃). λ_{max} 230 (sh) ; 235 ; 247 ; 251 (sh) (MeOH).
- N-Tetra-Me: Buxaminol A**
[58672-76-7]
Semisynthetic. Mp 146-147°. [α]_D²² +34 (c, 1.0 on CHCl₃).
- [11037-98-2]

- Stauffacher, D. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 968-981 (*Buxaminol*)
Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*Buxaminol E*)
Döpke, W. *et al.*, *Z. Chem.*, 1973, **13**, 135-136 (*N-Benzoylbuxodienine E, N-Benzoyl-O-acetyl-buxodienine E*)
Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1975, **40**, 3055-3060 (*Buxaminols A, B*)
Choudhary, M.I. *et al.*, *Tetrahedron*, 1986, **42**, 5747-5752 (*16-Acetoxybuxabenzamidienine, Buxaminol B*)
Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1989, **52**, 1319-1322; 1997, **60**, 770-774; 976-981 (*Nor-16-acetoxybuxabenzamidienine, Buxaminol C, 16-Acetoxybuxabenzamidienine*)
Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, **11**, 111-118 (*Buxus papillosa alkaloid*)
Loru, F. *et al.*, *Phytochemistry*, 2000, **54**, 951-957 (*N²⁰-Formylbuxaminol E, O-Syringoylbuxaminol E*)

Buxaminone

B-457

[114333-58-3]



C₂₆H₄₁NO 383.616

Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. solid. $[\alpha]_D^{20}$ -22 (c, 1.58 in CHCl₃). λ_{\max} 225 (ε 7760); 238 (ε 10700); 248 (ε 64600); 254 (ε 8130) (MeOH) (Derep).

16,17-Didehydro: **Buxpsine K. Buxpsine.****Buxamideine K**

[5189-69-5]

C₂₆H₃₉NO 381.6

Alkaloid from *Buxus balearica* and *Buxus sempervirens*. Mp 180-183° (176-178°). $[\alpha]_D$ +105 (c, 0.18 in CHCl₃).

16 α -Hydroxy: **Buxbarbarine K. 16-Hydroxybuxaminone**

[90468-71-6]

C₂₆H₄₁NO₂ 399.615


Alkaloid from roots of *Buxus sempervirens* (Buxaceae). Gum. $[\alpha]_D^{20}$ -54 (c, 1.56 in CHCl₃).

Tomko, J. *et al.*, *Tet. Lett.*, 1966, 915

(Buxpsine K)

Kurakina, I.O. *et al.*, *Khim. Prir. Soedin.*, 1969, 5, 406; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, 5, 337 (Buxpsine K)Mokry, P. *et al.*, *Chem. Zvesti.*, 1984, 38, 101-109 (Buxbarbarine K)Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1988, 51, 309-310 (Buxaminone)Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1992, 31, 2933-2935 (16-Hydroxybuxaminone)**Buxanaldinine****B-458**


[107259-42-7]

C₃₅H₅₀N₂O₄ 562.791

Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. $[\alpha]_D$ +12 (c, 1.39 in CHCl₃). Anomalous C-4 config.

Choudhary, M.I. *et al.*, *Tetrahedron*, 1986, 42, 5747-5752 (isol, uv, ir, pmr, ms, struct)**Buxandonine L****B-459**

20-Dimethylamino-14-methyl-9,19-cyclo-pregnan-3-one. **Buxandonine**
[17934-61-1]

C₂₄H₃₉NO 357.578

Rare *Buxus* alkaloid lacking C-4 subn. 9CI name incorrect. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 157-159° Mp 188-192°. $[\alpha]_D$ +14.

Deoxo: 3-Deoxybuxandonine


[142543-19-9]

C₂₄H₄₁N 343.595

Alkaloid from roots of *Buxus papillosa* (Buxaceae). Amorph. powder. $[\alpha]_D^{26}$ +10 (CHCl₃).

Döpke, W. *et al.*, *Pharmazie*, 1967, 22, 666; 1969, 24, 649; *CA*, 68, 87471e; 72, 43929p
Singh, C. *et al.*, *Tetrahedron*, 1977, 33, 1053-1055 (synth)Atta-ur-Rahman, *et al.*, *Tetrahedron*, 1992, 48, 3577-3584 (3-Deoxybuxandonine)**Buxanoldine****B-460**

[107259-41-6]

C₃₃H₄₈N₂O₃ 520.754

Cycloartane-type numbering used here. The C-28OH (4 α)-config. allocated here to these alkaloids is based on X-ray structures of alkaloids in this class and reverses the assignments made in the literature. Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. $[\alpha]_D$ -27.4 (c, 0.89 in CHCl₃). λ_{\max} 228 (log ε 4.07) (MeOH).

16-Ac: **Buxabenzacinine**

[149472-48-0]

C₃₅H₅₀N₂O₄ 562.791

Alkaloid from leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. $[\alpha]_D^{22}$ +55 (c, 0.036 in CHCl₃). λ_{\max} 226 (log ε 9.62) (MeOH).

28-Ac: **28-Acetylbuxanoldine**

[128351-96-2]

C₃₅H₅₀N₂O₄ 562.791

Alkaloid from leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D$ +50 (c, 1.6 in CHCl₃). Called 31-acetyl in the lit.

16,28-Di-Ac: **16,28-Diacetylbuxanoldine**

[107259-43-8]

C₃₇H₅₂N₂O₅ 604.828

Alkaloid from the roots of *Buxus sempervirens*. Gum. $[\alpha]_D^{20}$ +107 (c, 0.26 in CHCl₃). Called 16,31-di-Ac in the lit. λ_{\max} 226 (MeOH).

9,11-Dihydro, N³-Me, 16-Ac: **Buxabenzacinine**

[128741-20-8]

C₃₆H₅₄N₂O₄ 578.834

Alkaloid from *Buxus papillosa* (Buxaceae). Cryst. $[\alpha]_D^{20}$ +48 (CHCl₃).

28-Deoxy: **28-Deoxybuxanoldine**

[191101-02-7]

C₃₃H₄₈N₂O₂ 504.754

Alkaloid from *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D^{20}$ +123 (c, 0.8 in CHCl₃). Called 31-deoxy in the lit. λ_{\max} 228 (MeOH).

28-Deoxy, 16-Ac: **Buxadienine**

[123116-12-1]

C₃₅H₅₀N₂O₃ 546.792

Alkaloid from *Buxus sempervirens* (Buxaceae).

2 α -Hydroxy: **Buxirane**

Parent compd. not known.

2 α -Hydroxy, 2,16,28-tri-Ac: **2,16,28-****Triacetylbuxirane**

[912353-18-5]

C₃₉H₅₄N₂O₇ 662.865

Alkaloid from the leaves of *Buxus hyrcana*. $[\alpha]_D^{25}$ +107 (c, 0.26 in CHCl₃). λ_{\max} 227 (MeOH).

2 α -Hydroxy, 9,10-dihydro, 2,16,28-tri-Ac: **2,16,28-Triacetyl-9,10-dihydrobuxirane**


[912353-19-6]

C₃₉H₅₆N₂O₇ 664.881

Alkaloid from the leaves of *Buxus hyrcana*. $[\alpha]_D^{25}$ +89 (c, 0.39 in CHCl₃). λ_{\max} 226 (MeOH).

Choudhary, M.I. *et al.*, *Tetrahedron*, 1986, 42, 5747-5752 (Buxanoldine)Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1989, 60, 439-442 (28-Acetylbuxanoldine)Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1989, 28, 1293-1296; 1993, 32, 1059-1063 (Buxadienine, Buxabenzacinine)Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1990, 53, 319-324; 1997, 60, 770-774 (Buxabenzacinine, Deoxybuxanoldine)Ata, A. *et al.*, *Z. Naturforsch., C*, 2002, 57, 21-28 (Diacetylbuxanoldine)Meshkatsadat, M.H. *et al.*, *Z. Naturforsch., B*, 2006, 61, 201-206 (*Buxus hyrcana constiis*)**Buxapapinolamine****B-461**


[113762-73-5]

C₃₅H₄₈N₂O₅ 576.775

Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. $[\alpha]_D$ -16 (c, 1.84 in CHCl₃). Anomalous C-4 config.

Choudhary, M.I. *et al.*, *Phytochemistry*, 1988, 27, 271-274 (isol, uv, ir, pmr, ms, struct)**Buxaquamarine****B-462**

[98776-20-6]

C₂₆H₃₉NO₂ 397.6

Stereochem. revised in 2002. Alkaloid from *Buxus hyrcana*, the leaves of *Buxus*

papillosa and the roots of *Buxus sempervirens*. Gum. $[\alpha]_D^{20} +48$ (c, 0.25 in CHCl_3). λ_{max} 230 (sh); 238; 245; 253 (sh) (MeOH).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1985, **23**, 1951-1954 (*isol, uv, ir, pmr, ms, struct*)
 Ata, A. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 21-28 (*isol, pmr*)
 Babar, Z. U. *et al.*, *Steroids*, 2006, **71**, 1045-1051 (*Buxus hyrcana constit*)

Buxazine B-463

[1354-97-8]

$\text{C}_{28}\text{H}_{48}\text{N}_2\text{O}_2$ 444.699

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (Me_2CO). Mp 238-239°. $[\alpha]_D +93$ (CHCl_3).

Oxalate:

Amorph. Mp 257-260° dec.

O-Ac: Mp 238°.

Döpke, W. *et al.*, *Naturwissenschaften*, 1965, **52**, 61 (*isol, ir*)

Buxdeltine B-464

[1354-98-9]

$\text{C}_{33}\text{H}_{46}\text{N}_2\text{O}_3$ 518.738

Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. ($\text{Et}_2\text{O}/\text{Me}_2\text{CO}$). Mp 275°. $[\alpha]_D^{25} +79$ (c, 0.486 in EtOH).

Tomko, J. *et al.*, *Chem. Zvesti*, 1964, **18**, 721-731; *CA*, **62**, 4321d (*isol*)

Votický, Z. *et al.*, *CA*, 1967, **67**, 11668h (*isol*)

Buxetine B-465

[1354-99-0]

$\text{C}_{32}\text{H}_{50}\text{N}_2\text{O}_3$ 510.759


Struct. unknown. Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 263-265°. $[\alpha]_D^{27} -30$ (c, 0.816 in CHCl_3).

Tomko, J. *et al.*, *Chem. Zvesti*, 1964, **18**, 721-731; *CA*, **62**, 4321d (*isol*)

Votický, Z. *et al.*, *CA*, 1967, **67**, 11668h (*isol*)

Buxidienine I B-466

Baleabuxidienine I



$\text{C}_{24}\text{H}_{40}\text{N}_2\text{O}_2$ 388.592

Cycloartane numbering used here. The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. The parent compd., Buxidienine I, is unknown. There is uncertainty about the C-4

config. in some of these alkaloids. The accepted config. was revised in 1975 to 4*S*- (28-hydroxy), but structs. with the 4*R*- (29-hydroxy) config. continue to be published.

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*: Buxidienine F

[14777-61-8]

Mp 237°. $[\alpha]_D +8$ (CHCl_3).

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 -*Ac*: Mp 263°. $[\alpha]_D -73$.

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 - (2-methylprop-*n*oyl): *N*-Isobutyrylbaleabuxidienine F. *N*-Isobutyrylbuxidienine F

[14155-73-8]

$\text{C}_{30}\text{H}_{50}\text{N}_2\text{O}_3$ 486.737

Alkaloid from *Buxus balearica* (Buxaceae). Mp 253°. $[\alpha]_D -67$ (CHCl_3). Shown lacking the C-18 methyl group due to a drawing error.

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 -angeloyl, 16-*Ac*:

Sempervirine†

[135626-67-4]

$\text{C}_{33}\text{H}_{52}\text{N}_2\text{O}_4$ 540.785

Alkaloid from leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D^{20} +2$ (CHCl_3).

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 -benzoyl: *N*-Benzoylbaleabuxidienine F. *N*-Benzoylbuxidienine F

[14155-72-7]

$\text{C}_{33}\text{H}_{48}\text{N}_2\text{O}_3$ 520.754

Alkaloid from *Buxus balearica* (Buxaceae). Mp 291° Mp 286-288°. $[\alpha]_D -36$ (-29) (CHCl_3). Shown lacking the C-18 methyl group in one paper.

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 -benzoyl, 16-*Ac*: 16-*O*-Acetyl-*N*-benzoylbuxidienine

[126035-07-2]

$\text{C}_{35}\text{H}_{50}\text{N}_2\text{O}_4$ 562.791

Alkaloid from leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D^{20} -5$ (c, 0.2 in CHCl_3). The 4-config. is shown as the reverse of what is now known for these alkaloids.

$\text{N}^{20}, \text{N}^{20}$ -*Di-Me*, N^3 -benzoyl, 28-*Ac*: 28-*Ac*etoxy-*N*^α-benzoylbuxidienine

[104104-37-2]

[123116-13-2]

$\text{C}_{35}\text{H}_{50}\text{N}_2\text{O}_4$ 562.791

Alkaloid from leaves of *Buxus papillosa* and from *Buxus sempervirens*. Amorph. $[\alpha]_D^{20} -40$ (CHCl_3).

30-*Aldehyde*, N^{20} -*Me*: Buxalongifoline
 Unknown parent.

30-*Aldehyde*, N^{20} -*Me*, N^3 -benzoyl, 16-*Ac*: *O*-Acetyl-*N*-benzoylbuxalongifoline

[194852-53-4]

$\text{C}_{34}\text{H}_{46}\text{N}_2\text{O}_4$ 546.748

Alkaloid from the leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. $[\alpha]_D^{22} +16$ (c, 0.06 in CHCl_3). λ_{max} 229 (log ϵ 4.28); 235 (log ϵ 4.29); 245 (log ϵ 4.3); 254 (sh) (log ϵ 4.32) (MeOH).

16-*Deoxy*, $\text{N}^3, \text{N}^{20}, \text{N}^{20}$ -*Tri-Me*: 16-*Deoxy*-buxidienine C. 16-*Desoxy*buxidienine

[36127-41-0]

$\text{C}_{27}\text{H}_{46}\text{N}_2\text{O}$ 414.673

Alkaloid from *Buxus papillosa* and *Buxus madagascariensis* (Buxaceae). Cryst. (Me_2CO). Mp 200-201° Mp

183-185°. $[\alpha]_D^{25} +33$ (c, 1.0 in CHCl_3).

16-*Deoxy*, $\text{N}^3, \text{N}^{20}, \text{N}^{20}$ -*tri-Me*, N^3 - (2*S*-hydroxy-3-methylbutanoyl): Buxakarchiamine

[398477-71-9]

$\text{C}_{32}\text{H}_{54}\text{N}_2\text{O}_3$ 514.79

Alkaloid from *Buxus papillosa*. Amorph. powder. $[\alpha]_D^{25} -9$ (c, 0.22 in CHCl_3). λ_{max} 238 (log ϵ 4.2); 245 (log ϵ 4.2) (MeOH).

16-*Deoxy*, $\text{N}^3, \text{N}^{20}, \text{N}^{20}$ -*tri-Me*, N^3 - (2*R*-hydroxy-3- ζ -methylpentanoyl): Buxahejramine

[398477-72-0]

$\text{C}_{33}\text{H}_{56}\text{N}_2\text{O}_3$ 528.817

Alkaloid from *Buxus papillosa*. Amorph. powder. $[\alpha]_D^{25} -7$ (c, 0.14 in CHCl_3). Struct. drawn with missing *N*-Me group in ref. λ_{max} 239 (log ϵ 4.18); 245 (log ϵ 4.2) (MeOH).

16-*Deoxy*, $\text{N}^3, \text{N}^{20}, \text{N}^{20}$ -*tri-Me*, 28-benzoyl: *O*²⁸-Benzoyl-16-*deoxy*buxidienine C

[129225-34-9]

$\text{C}_{34}\text{H}_{50}\text{N}_2\text{O}_2$ 518.781

Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Gum. $[\alpha]_D -75$ (CHCl_3). Called 30-benzoyl in the lit.

16-*Deoxy*, *N*-tetra-*Me*: 30-*Hydroxy*buxamine A

[129225-35-0]

$\text{C}_{28}\text{H}_{48}\text{N}_2\text{O}$ 428.7

Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Amorph. $[\alpha]_D +61$ (CHCl_3). Has C-4 α hydroxymethyl group.

28-*Deoxy*, $\text{N}^{20}, \text{N}^{20}$ -*di-Me*, N^3 -benzoyl: *N*-Benzoylbuxidienine

[135626-69-6]

$\text{C}_{33}\text{H}_{48}\text{N}_2\text{O}_2$ 504.754

Alkaloid from *Buxus hyrcana* and *Buxus sempervirens*. Amorph. $[\alpha]_D^{25} +7$ (c, 1.5 in CHCl_3).

16-*Epimer*, *N*-tetra-*Me* (?): Buxaminol G

[97763-00-3]

$\text{C}_{28}\text{H}_{48}\text{N}_2\text{O}_2$ 444.699

Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Gum. $[\alpha]_D +83.33$ (CHCl_3). Little evidence for the assigned stereochem. The 4-config. is shown as the reverse of what is now known for these alkaloids. λ_{max} 237 (ϵ 6040); 245 (ϵ 5860); 253 (ϵ 4151) (MeOH).

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*Buxus balearica constits*)

Herlem, D. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1967, **264**, 798-799 (*synth*)

Shamma, M. *et al.*, *Phytochemistry*, 1973, **12**, 2051-2054 (*Desoxybuxidienine*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*config*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1985, **24**, 1398-1399; 1990, **29**, 1293-1296; 1991, **30**, 1295-1298 (*Buxaminol G*, 30-*Hydroxybuxamine A*, *O*-Benzoyl-16-*deoxy*buxidienine C, *Sempervirine*, *N*-Benzoylbuxidienine)


Atta-ur-Rahman, *et al.*, *J.C.S. Perkin I*, 1986, 919-921 (*Acetoxybenzoylbuxidienine*)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1989, **52**, 1319-1322; 1997, **60**, 976-981 (*O*-Acetyl-

N-benzoylbuxidienine, *O*-Acetyl-*N*-benzoylbuxalongifoline

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 2001, **58**, 963-968 (*Buxakarachiamine*, *Buxahejramine*)
Babar, Z. U. *et al.*, *Steroids*, 2006, **71**, 1045-1051 (*Benzoylbuxidienine*)


Buxifoliadine D B-467
[263007-68-7]



$C_{23}H_{23}NO_3$ 361.44
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 236-238°. λ_{max} 277 (log ϵ 3.32); 309 (sh) (log ϵ 3); 337 (sh) (log ϵ 2.02); 380 (log ϵ 1.72); 403 (log ϵ 1.84) (MeOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*isol*, *pmr*, *cmr*, *ms*)


Buxifoliadine E B-468
[263007-69-8]



$C_{23}H_{25}NO_5$ 395.454
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 247-249°. Racemic. λ_{max} 259 (log ϵ 4.72); 270 (log ϵ 4.59); 273 (log ϵ 4.6); 277 (log ϵ 4.59); 284 (log ϵ 4.68); 330 (sh) (log ϵ 3.85); 393 (log ϵ 3.86) (MeOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*isol*, *pmr*, *cmr*, *uv*, *ms*)


Buxifoliadine F B-469
[263007-70-1]



$C_{16}H_{11}NO_4$ 281.267
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp >280°. λ_{max} 259 (log ϵ 4.23); 379 (log ϵ 3.53); 384 (log ϵ 3.55) (MeOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*isol*, *pmr*, *cmr*, *ms*, *uv*)


Buxifoliadine G B-470
[263007-71-2]



$C_{22}H_{21}NO_4$ 363.412
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 150-152° (dec.). λ_{max} 248 (log ϵ 3.44); 277 (log ϵ 3.71); 287 (sh) (log ϵ 3.68); 399 (log ϵ 3.05) (MeOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*isol*, *pmr*, *cmr*, *ms*, *uv*)

Buxifoline B-471
*6,7,7a,8-Tetrahydro-4,10-dimethoxy-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, 9*Cl*. 3,9-Dimethoxy-1,2-methylenedioxyxynoraporphine*



$C_{19}H_{19}NO_4$ 325.363

(*R*)-form [80151-81-1]
Alkaloid from *Xylopia buxifolia* and from the leaves and stem bark of *Duguetia obovata* (Annonaceae). Noncryst.

Hydrochloride:
Cryst. (MeOH). Mp 295° dec. $[\alpha]_D$ -15 (c, 0.34 in MeOH).

N-Formyl: N-Formylbuxifoline
[89368-29-6]
 $C_{20}H_{19}NO_5$ 353.374
Minor alkaloid from the leaves of *Duguetia obovata* (Annonaceae). Cryst. (MeOH). Mp 198°. $[\alpha]_D$ -181 (c, 0.11 in CHCl₃).

N-Ac:
Cryst. (MeOH). Mp 230°. $[\alpha]_D$ -295 (c, 0.40 in CHCl₃).


N-Me: N-Methylbuxifoline. 3,9-Dimethoxy-1,2-methylenedioxyxynoraporphine
[89368-28-5]
 $C_{20}H_{21}NO_4$ 339.39

Trace alkaloid from leaves of *Duguetia obovata* (Annonaceae). Noncryst. Negative opt. rotn. in CHCl₃.

*7*S*-Hydroxy, N-Me: Polyalthine*
[67627-73-0]


$C_{20}H_{21}NO_5$ 355.39
Alkaloid from the trunk bark of *Polyalthia suaveolens*. Noncryst.; cryst. (MeOH/Et₂O) as hydrochloride. Mp 274° dec. $[\alpha]_D$ +11 (c, 0.8 in MeOH).
Cave, A. *et al.*, *Planta Med.*, 1978, **33**, 243 (*Polyalthine*)
Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551 (*uv*, *pmr*, *ms*, *struct*)
Roblot, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 862 (*isol*, *cmr*, *pmr*, *struct*, *synth*, *derivs*)

Buxiramine D B-472
Buxiramine
[25650-67-3]



$C_{27}H_{44}N_2O_2$ 428.657
Struct. not certain. Alkaloid from *Buxus sempervirens* (Buxaceae). Prisms (Me₂CO). Mp 213-215°.
Döpke, W. *et al.*, *Pharmazie*, 1969, **24**, 649; *CA*, **72**, 43929p (*isol*, *ms*)

Buxitrienine C B-473
[36151-06-1]



$C_{27}H_{44}N_2O$ 412.657
Alkaloid from *Buxus madagascariensis* (Buxaceae). Cryst. (C₆H₆/hexane). Mp 192°. $[\alpha]_D^{20}$ +57 (CHCl₃).

Deoxy: Papillotrienine
[145385-74-6]
 $C_{27}H_{44}N_2$ 396.658
Alkaloid from roots of *Buxus papillosa* (Buxaceae). Amorph. solid. $[\alpha]_D$ +60 (CHCl₃).


Deoxy, N²⁰-de-Me: N^b-Demethylpapillotrienine
[145385-75-7]
 $C_{26}H_{42}N_2$ 382.631
Isol. from roots of *Buxus papillosa* (Buxaceae). Amorph. solid. $[\alpha]_D^{20}$ +62 (CHCl₃).

Khong-Huu, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 558-560 (*Buxitrienine C*)
Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1992, **55**, 1063-1066 (*Papillotrienine*, *Demethylpapillotrienine*, *isol*, *pmr*, *cmr*)

Buxocyclamine I

B-474

4,14-Dimethyl-9,19-cyclopregnane-3,20-diamine. 3,20-Diamino-4,14-dimethyl-9,19-cyclopregnane



$C_{23}H_{40}N_2$ 344.582

In *Buxus* alkaloids the suffix letter designates the *N*-subn. pattern. The parent compd., Buxocyclamine I, is unknown.

N^{20}, N^{20} -Di-Me, N^3 -benzoyl: **Buxisine**. *N*-Benzoylbuxocyclamine F

$C_{32}H_{48}N_2O$ 476.744

Alkaloid from *Buxus sempervirens*.

N-Tetra-Me: **Buxocyclamine A**

[19642-32-1]

$C_{27}H_{48}N_2$ 400.69

Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me_2CO). Mp 187-188°. $[\alpha]_D^{25} +87$ (c, 0.15 in $CHCl_3$). 4 β -Config was assigned in the original paper, but 4 α -config is more likely by analogy with other similar alkaloids.

4,23-Didehydro, N^3, N^{20} -di-Me, N^3 -Ac:

Cyclobuxamidine

$C_{27}H_{44}N_2O$ 412.657

Alkaloid from the leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. $[\alpha]_D^{25} +24$ (c, 0.04 in $CHCl_3$). λ_{max} 220 (log ϵ 4.36) (MeOH).

Döpke, W. *et al.*, *Pharmazie*, 1968, **23**, 37-38 (Buxocyclamines)


Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997, **60**, 976-981 (Cyclobuxamidine)

Buxomeline

B-475

Methyl 2,3,6,7-tetrahydro-10,20-epoxy-19-hydroxyaspidospermidine-3-carboxylate, 9CI

[69734-96-9]



Relative configuration

$C_{21}H_{22}N_2O_4$ 366.416


Alkaloid from *Melodinus celastroides* (Apocynaceae). Noncryst. $[\alpha]_D^{25} -240$ ($CHCl_3$).

Rabaron, A. *et al.*, *Phytochemistry*, 1978, **17**, 1452 (uv, ms, pmr, struct)

Buxotrienine

B-476

[107259-40-5]



$C_{26}H_{39}NO$ 381.6


Alkaloid from the leaves of *Buxus papilosa* (Buxaceae). Amorph. $[\alpha]_D +13.5$ (c, 1.81 in $CHCl_3$).

Choudhary, M.I. *et al.*, *Tetrahedron*, 1986, **42**, 5747-5752 (isol, uv, ir, pmr, ms, struct)

Buxoxybenzamine

B-477

[126794-77-2]



$C_{35}H_{50}N_2O_5$ 578.79

Alkaloid from the leaves of *Buxus papilosa* and *Buxus sempervirens*. Shows phytotoxic activity against *Lemna minor*. Amorph. solid. $[\alpha]_D^{20} -22$ (MeOH).

O-Ac: **Buxapapilinine**

[129134-89-0]

$C_{37}H_{52}N_2O_6$ 620.828

Alkaloid from leaves of *Buxus papilosa* and *Buxus sempervirens*. Shows phytotoxic activity against *Lemna minor*. Amorph. solid. $[\alpha]_D^{20} -28$ (c, 0.1 in $CHCl_3$). $[\alpha]_D^{25} +123$ (c, 0.9 in $CHCl_3$). λ_{max} 227; 237; 245 (MeOH).

6-Deacetoxy, 16 α -acetoxy, 2-Ac: **2,16-Diacetoxy-9,11-epoxybuxamidine**

[207734-00-7]

$C_{37}H_{52}N_2O_6$ 620.828

Alkaloid from the leaves of *Buxus papillosa*. Amorph. solid. $[\alpha]_D^{20} -76$ (c, 1.2 in $CHCl_3$). λ_{max} 228 (MeOH).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1990,

31, 493-498 (Buxapapilinine)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1990,

29, 683-685 (isol, uv, ir, pmr, cmr, ms)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1998,

48, 519-528 (Diacetoxyepoxybuxamidine)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1999,

62, 665-669 (activity)


Babar, Z.U. *et al.*, *Steroids*, 2006, **71**, 1045-

1051 (Buxapapilinine)

Buxozine C

B-478

[64938-84-7]



$C_{27}H_{46}N_2O$ 414.673

Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (Et_2O). Mp 137°. $[\alpha]_D^{21} +65$ (c, 0.6 in $CHCl_3$). Redn. gives Cyclovir-obuxine C in C-917.

Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 2549-2554 (isol, ms, pmr, struct)

Votický, Z. *et al.*, *Phytochemistry*, 1977, **16**, 1860-1861 (isol, ir, ms)

Buxpapamine

B-479

[11054-79-8]

$C_{28}H_{30}N_2O$ 410.558

Steroidal (*Buxus*) alkaloid. Struct. unknown. Alkaloid from *Buxus papilosa* (Buxaceae). Cryst. (Me_2CO). Mp 205-206°. $[\alpha]_D +93$ (c, 1.28 in $CHCl_3$).

Ikram, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1968, **11**, 253-254; *CA*, **71**, 779b (isol)

Buxpapine

B-480

[11054-80-1]

$C_{27}H_{34}N_2O$ 402.578

Steroidal (*Buxus*) alkaloid. Struct. unknown. Alkaloid from *Buxus papilosa* (Buxaceae). Mp 111-112°. $[\alpha]_D +7.5$ (c, 4.8 in $CHCl_3$).

Ikram, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1968, **11**, 253-254; *CA*, **71**, 779b (isol)

Buxpapinine

B-481

$C_{37}H_{52}N_2O_6$ 620.828

Struct. unknown. Alkaloid from *Buxus papilosa* leaves (Buxaceae). Mp 253-255°. $[\alpha]_D^{25} -53$ (c, 1.24 in $CHCl_3$). Contains secondary amide and ester groups.


Ikram, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1968, **11**, 488-489; *CA*, **71**, 88432s (isol, ir, pmr)

Buxtauine M

B-482

16-Hydroxy-14-methyl-3-(methylamino)-4-methylene-9,19-cyclopregnan-20-one, 9CI. Buxtauine. Cyclomicrobuxinine. Cyclobuxoxine

[4236-73-1]



Absolute configuration

$C_{24}H_{37}NO_2$ 371.562

Alkaloid from *Buxus sempervirens*, *Buxus wallichiana*, *Buxus harlandi* and several other *Buxus* spp. (Buxaceae). Mp 178-181° (170°). $[\alpha]_D^{23} +154$ (c, 0.77 in $CHCl_3$).

Oxime: Mp 235-238°.

O,N-Di-Ac: Mp 213-215° (193-197°). $[\alpha]_D +87$ (c, 0.67 in $CHCl_3$).

N-Me: **Buxpiine K**. Cyclomicrobuxinine.

Buxpiine. *Buxippine K*

[3296-11-5]

$C_{25}H_{39}NO_2$ 385.589

Alkaloid from *Buxus sempervirens*, *Buxus hyrcana* and *Buxus harlandi* (Buxaceae). Mp 178-180° (173°). $[\alpha]_D +172$ (c, 2 in $CHCl_3$). $[\alpha]_D +159$ ($CHCl_3$).

Deoxy: **Cyclomicrobuxamine**

[128351-97-3]

$C_{24}H_{37}NO$ 355.562


Alkaloid from leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D +140$ (c, 5.8 in $CHCl_3$).

Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*,

1965, **30**, 3705-3710 (*isol, ir, uv, ms, struct*)
 Nakano, T. *et al.*, *J.C.S.*, 1965, 6688-6694 (*isol, ir, pmr, struct*)
 Votický, Z. *et al.*, *Tet. Lett.*, 1965, 3579-3584 (*isol, ir, pmr, config*)
 Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1989, **60**, 439-442 (*Cyclomicrobuxamine*)

Buxupapine**B-483**

[109305-87-5]

C₂₇H₄₆N₂ 398.674

Alkaloid from the leaves of *Buxus papilosa* (Buxaceae). Amorph. [α]_D +11 (c, 1.13 in CHCl₃).

N²⁰-De-Me: N^b-Norbuxupapine

[109305-88-6]

C₂₆H₄₄N₂ 384.647

Alkaloid from the leaves of *Buxus papilosa* (Buxaceae). Amorph. [α]_D +20 (c, 0.71 in CHCl₃).

N³-Me: N³-Methylbuxupapine

[38406-16-5]

C₂₈H₄₈N₂ 412.701


Alkaloid from the leaves of *Buxus papilosa*. Gum. [α]_D²⁰ +126 (c, 0.65 in CHCl₃). Synonym assigned in ref. is incorrect. λ_{\max} 201 (MeOH).

Choudhary, M.I. *et al.*, *J. Nat. Prod.*, 1987, **50**, 84-88 (*isol, ir, pmr, ms, struct, N²⁰-Norbuxupapine*)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, **11**, 111-118 (*N³-Methylbuxupapine*)

Buzonamine**B-484**

Decahydro-3,3-dimethyl-2H-oxireno[d]-pyrrolo[2,1-a]isoquinoline, 9CI
 [270905-32-3]



Relative Configuration


C₁₄H₂₃NO 221.342

Alkaloid from the defensive secretion of the millipede *Buzonium crassipes*.

Wood, W.F. *et al.*, *Biochem. Syst. Ecol.*, 2000, **28**, 305-312 (*isol, pmr, cmr, ms*)

Caboxine A

[53851-13-1]



Absolute Configuration

$C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from the leaves of *Cabucala fasciculata* (Apocynaceae). Mp 195-210°. $[\alpha]_D^{22}$ -68 (CHCl₃). λ_{max} 222 (log ϵ 4.45); 245 (log ϵ 4.13); 286 (log ϵ 3.43); 294 (log ϵ 3.26) (CHCl₃).

3-Epimer: **Vineridine**. *Isocaboxine A* [3489-06-3]

$C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Vinca erecta* and from *Cabucala fasciculata* (Apocynaceae). Cryst. (Me₂CO). Mp 179-180°. $[\alpha]_D^{22}$ +40.5 (c, 2.4 in CHCl₃). λ_{max} 220 (log ϵ 4.54) (EtOH).

► LR1240000

3-Epimer, nitrate: Mp 178° dec. (170°).

3-Epimer, N-oxide: **Vineridine N-oxide** [53643-15-5]

$C_{22}H_{26}N_2O_6$ 414.457
Alkaloid from *Vinca erecta* (Apocynaceae). Cryst. (CHCl₃). Mp 193-195°. $[\alpha]_D$ +20 (c, 0.68 in MeOH).

7-Epimer (?): **Reserpinine oxindole** [13483-85-7]

$C_{22}H_{26}N_2O_5$ 398.458
Trace alkaloid from leaves of *Rauwolfia vomitoria* (Apocynaceae). Off-white amorph. powder. $[\alpha]_D^{21}$ -22 (c, 0.1 in CHCl₃). C-7 config. not established.

20-Epimer: **Tetraphylline oxindole A**

[62358-36-5]
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from the bark of *Ochrosia glomerata* and from *Neisosperma glomerata*. Cryst. (MeOH). Mp 218°. $[\alpha]_D$ +20 (CHCl₃). λ_{max} 220 (log ϵ 4.45); 288 (log ϵ 3.43); 295 (log ϵ 3.28) (MeOH).

3,7-Diepimer: **Vinerine**. *Isocaboxine B* [3382-38-5]

$C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Vinca erecta* and *Cabucala fasciculata*. Cryst. (MeOH). Mp 202-203°. $[\alpha]_D$ +53 (CHCl₃). λ_{max} 220 (log ϵ 4.54) (EtOH).

3,7-Diepimer, nitrate: Mp 170° dec.

3,7-Diepimer, N-oxide: **Vinerine N-oxide** [61687-65-8]

$C_{22}H_{26}N_2O_6$ 414.457
Isol. from *Vinca erecta* (Apocynaceae). Cryst. (MeOH). Mp 246-247°.

3,7-Diepimer, N-Ac: **N-Acetylvinerine**

[3382-39-6, 33044-89-2]
 $C_{24}H_{28}N_2O_6$ 440.495
Minor alkaloid from epigeal parts of *Vinca erecta* (Apocynaceae). Mp 159-160° (152-153°). $[\alpha]_D^{25}$ -99.5 (Me₂CO).

7,20-Diepimer: **Tetraphylline oxindole B** [62358-35-4]

C-1

$C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from the leaves of *Ochrosia glomerata* and from *Neisosperma glomerata*. Mp 226°. $[\alpha]_D$ -26 (CHCl₃). λ_{max} 220 (log ϵ 4.45); 288 (log ϵ 3.43) (MeOH).

Stereoisomer (?): **Erycinine**. *Ericinine* [19775-58-7]

$C_{22}H_{26}N_2O_5$ 398.458
From *Vinca erecta* (Apocynaceae). Mp 206-207°. $[\alpha]_D^{18}$ +43.8 (Me₂CO). May be identical with *Isocaboxine A*. Stereochem. at C-3 and C-7 is not known. λ_{max} 220 (log ϵ 4.68) (EtOH).

Kasymov, Sh.Z et al., *Khim. Prir. Soedin.*, 1966, 2, 260-265; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, 2, 210-213 (*Vinerine*, *Vineridine*)

Aburakhimova, N. et al., *Khim. Prir. Soedin.*, 1968, 4, 135; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, 4, 116 (*Erycinine*)

Malikov, V.M. et al., *Khim. Prir. Soedin.*, 1970, 6, 640-641; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, 6, 662 (*N-Acetylvinerine*)

Khalmirzaev, M.M. et al., *Khim. Prir. Soedin.*, 1973, 9, 806; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, 9, 776-777 (*Vineridine N-oxide*)

Titeux, F. et al., *Phytochemistry*, 1974, 13, 1620-1621; 1975, 14, 565-568 (*Caboxine A*, *Isocaboxine A*, *isol*, *uv*, *ir*, *pmr*, *abs config*)

Titeux, F. et al., *Bull. Soc. Chim. Fr.*, 1976, 1473-1475 (*Tetraphylline oxindoles A, B*, *synth*, *uv*, *ir*, *pmr*, *ms*)

Sharipov, M.R. et al., *Khim. Prir. Soedin.*, 1976, 12, 401-402; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, 12, 355-356 (*Vinerine N-oxide*)


Amer, M.M. et al., *Phytochemistry*, 1980, 19, 1833-1836 (*Reserpinine oxindole*)

Seguin, E. et al., *J. Nat. Prod.*, 1982, 45, 738-744; 1984, 47, 687-691 (*Tetraphylline oxindoles A, B*, *isol*)

Cabudine

C-2

6,7-Dihydro-10-methoxy-7-methyl-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-4-methanol, 9CI. 5,6-Dihydro-3-hydroxymethyl-9-methoxy-6-methyl-1,2-methylenedioxy-4H-dibenzo[*de,g*]quinoline [59272-70-7]




$C_{20}H_{19}NO_4$ 337.374
Alkaloid from *Thalictrum isopyroides* (Ranunculaceae). Exhibits adrenolytic activity. Mp 184-185°.

Kurbanov, M. et al., *Dokl. Akad. Nauk Tadzh. SSR*, 1975, 18, 20; *CA*, 84, 180440j (*isol*, *uv*, *ir*, *pms*, *ms*, *struct*)

Cabufiline

[82854-07-7]



C-3

$C_{44}H_{54}N_4O_7$ 750.933
Alkaloid from the leaves of *Cabucala caudata* (Apocynaceae). Amorph. $[\alpha]_D$ -91 (c, 1.0 in EtOH).

Deepoxy: **Desoxycabufiline**

[82841-74-5]
 $C_{44}H_{54}N_4O_6$ 734.934
Alkaloid from the stem bark of *Alstonia plumosa* (Apocynaceae). $[\alpha]_D$ -114 (c, 0.9 in MeOH).

Deepoxy, O-de-Me: **Nordesoxycabufiline** [82854-85-1]

$C_{43}H_{52}N_4O_6$ 720.907
Alkaloid from the stem bark of *Alstonia plumosa* (Apocynaceae). $[\alpha]_D$ -100 (c, 0.46 in Py). Posn. of demethylation not known.


Massiott, G. et al., *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1982, 294, 579-582 (*uv*, *pmr*, *struct*)

Jacquer, M.J. et al., *Phytochemistry*, 1982, 21, 2973-2977 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *Desoxycabufiline*, *Nordesoxycabufiline*)
Subhadhirasakul, S. et al., *Chem. Pharm. Bull.*, 1994, 42, 1427-1431 (*config*, *cmr*)

Cadabicine

[99964-83-7]

C-4



$C_{25}H_{29}N_3O_4$ 435.522
CAS numbering shown. Numbering systems vary. Alkaloid from the stem bark of the famine food *Cadaba ferinosa* (luquata sigmama) and *Crataeva nurvala* (Capparidaceae). Mp 270-272° dec. *Crataeva* is not a generally recognised genus. N¹⁶, O-Di-Ac: **Cadabicine diacetate** [99964-84-8]

C₂₉H₃₃N₃O₆ 519.596Alkaloid from *Crataeva nurvala* (Cappariaceae). Rods (MeOH). Mp 265-268°.**Me ether: Cadabicine methyl ether. O-Methylcadabicine**C₂₆H₃₁N₃O₄ 449.549Alkaloid from stem bark of the famine food *Cadaba farinosa* (liquata sigma) and *Crataeva nurvala* (Cappariaceae).**26-Methoxy: Isocodonocarpine**

[126431-38-7]


C₂₆H₃₁N₃O₅ 465.548Alkaloid from root bark of *Capparis decidua* (Cappariaceae). Off-white cryst. (MeOH). Mp 220-222° dec. CAS gives incorrect name.**26-Methoxy, N¹⁶-Ac: N-Acetylisocodonocarpine**C₂₈H₃₃N₃O₆ 507.585Alkaloid from root bark of *Capparis decidua* (Cappariaceae). Amorph. powder. Mp 234-236°. Called 14-N-Ac in the lit.**26-Methoxy, N¹⁶, O-di-Ac:**

Cryst. (MeOH). Mp 205-207°.

Ahmad, V.U. et al., *Phytochemistry*, 1985, **24**, 2709; 1989, **28**, 2493 (isol, uv, ir, pmr, cmr, ms, cryst struct)Ahmad, V.U. et al., *J. Nat. Prod.*, 1987, **50**, 1186; 1992, **55**, 1509 (isol, Cadabicine, Cadabicine diacetate, 14-N-Acetylisocodonocarpine)Ahmad, V.U. et al., *Pak. J. Sci. Ind. Res.*, 1992, **35**, 475; *CA*, **119**, 17756n (Cadabicine methyl ether)**Cadambine**

[54422-49-0]


C-5



Absolute Configuration

C₂₇H₃₂N₂O₁₀ 544.557Alkaloid from the heartwood of *Anthocephalus cadamba* and the leaves of *Nauclea latifolia* (Nucleaceae). Mp 207-211°. [α]_D²⁵ -71 (MeOH).**Tetra-O-Ac:** Mp 149-151°. [α]_D²⁰ -124 (CHCl₃).**Parent acid: Cadambine acid**C₂₆H₃₀N₂O₁₀ 530.53Alkaloid from the bark of *Nauclea diderrichii*. Amorph. yellow solid. Mp 225°. [α]_D²⁰ -80 (c, 1 in MeOH). Conflicting stereochem. shown in 2005 ref.Brown, R.T. et al., *Tet. Lett.*, 1974, 1957; 1991,**32**, 1987 (uv, ir, ms, pmr, cd, struct, synth)Hotellier, F. et al., *Planta Med.*, 1979, **35**, 242 (isol)Lamidi, M. et al., *Magn. Reson. Chem.*, 2005, **43**, 427-429 (Cadambine acid)**Cadamine**

C-6

Methyl 5,7,8,13,13b,14-hexahydro-5-(hydroxymethyl)indolo[2',3':3,4]pyrido[1,2-b][2,7]naphthyridine-1-carboxylate, 9CI [60723-52-6]


Absolute Configuration

C₂₁H₂₁N₃O₃ 363.415A heteroyohimbinoind. Alkaloid from leaves of *Anthocephalus cadamba* (Rubiaceae). [α]_D²⁵ -65 (CHCl₃) (as O-Ac).**19-Epimer: Isocadamine**


[60761-52-6]

C₂₁H₂₁N₃O₃ 363.415Alkaloid from leaves of *Anthocephalus cadamba* (Rubiaceae).Brown, R.T. et al., *Tet. Lett.*, 1976, 1629-1630 (isol, uv, struct)**Cadeguomycin**

C-7

2-Amino-4,7-dihydro-4-oxo-7-β-D-ribofuranosyl-1H-pyrrolo[2,3-d]pyrimidine-5-carboxylic acid. 7-Carboxy-7-deazaguanosine [81645-08-1]C₁₂H₁₄N₄O₇ 326.265Nucleoside antibiotic. Isol. from *Streptomyces hygroscopicus*. Shows antineoplastic activity and immunostimulatory effects. Needles. Sol. MeOH, DMSO; fairly sol. H₂O; poorly sol. EtOH, hexane. Mp 231-239° dec. Log P -2.94 (calc). λ_{max} 231 (ε 14100); 272 (ε 6310); 297 (ε 6920) (0.1M HCl) (Derep). λ_{max} 225 (sh) (ε 9175); 268 (ε 9175); 282 (sh) (ε) (H₂O at pH 11) (Derep). λ_{max} 232 (ε 19680); 272 (ε 6881); 298 (ε 7610) (H₂O) (Derep).▶ LD₅₀ (mus, ipr) 200 - 600 mg/kg. UY9358980**Nitrile: 5-Cyano-7-deazaguanosine**C₁₂H₁₃N₅O₅ 307.265Isol. from tRNA^{tyr} of an *Escherichia coli* mutant. Pale yellow cryst. Mp 260-270° dec. [α]_D¹⁵ -59.1 (c, 0.1 in DMSO). Noguchi, S. et al., *Nucleic Acids Res.*, 1978, **5**, 4215 (nitrile, isol)Tanaka, H. et al., *J. Antibiot.*, 1982, **35**, 272 (isol)Wu, R.T. et al., *J. Antibiot.*, 1982, **35**, 279 (struct)Beylin, V.G. et al., *Tet. Lett.*, 1983, **24**, 4793 (synth)Yuan, B.D. et al., *J. Antibiot.*, 1985, **38**, 642 (props)Kondo, T. et al., *Tetrahedron*, 1986, **42**, 199; 207 (synth, pmr, uv, ir, nitrile)Kim, S.H. et al., *J. Antibiot.*, 1987, **40**, 1776 (pharmacol)Isono, K. et al., *J. Antibiot.*, 1988, **41**, 1711 (rev)Ramasamy, K. et al., *J.C.S. Perkin 1*, 1989, 2375 (synth, ir, pmr, uv)**Cadamine**

C-8

6-[Octahydro-8-hydroxy-1-(hydroxymethyl)-2H-quinolizin-3-yl]-2-piperidinone, 9CI [58071-45-7]C₁₅H₂₆N₂O₃ 282.382A seco-sparteine alkaloid, cf. Aphyllic acid, A-1335. Alkaloid from *Cadia purpurea* (Fabaceae).**O⁸-(Hydroxyphenylacetyl): Cadamine hydroxyphenylacetate**

[61168-58-9]

C₂₃H₃₂N₂O₅ 416.516Alkaloid from *Cadia purpurea* (Fabaceae). Full struct. not determined. The OH group is in the aromatic ring. Alkaloid not named in the lit.**O⁸-(2-Pyrrolocarbonyl): Cadamine 2-pyrrolocarboxylate**


[61166-49-2]

C₂₀H₂₉N₃O₄ 375.467Alkaloid from *Cadia purpurea* (Fabaceae). Not named in the lit.van Eijk, J.L. et al., *Planta Med.*, 1975, **28**, 139-142 (isol)van Eijk, J.L. et al., *Tet. Lett.*, 1976, **24**, 2053-2054 (struct, ms, derivs)**Caerulomycin**

C-9

4-Methoxy-[2,2'-bipyridine]-6-carboxaldehyde oxime, 8CI. 4-Methoxy-2,2'-dipyridyl-6-aldoxime. Caerulomycin†.**Caerulomycin A**

[21802-37-9]

C₁₂H₁₁N₃O₂ 229.238Antibiotic from *Streptomyces caerules* and *Nocardiosis cirriefficiens*. Active primarily against yeasts, fungi and *Entamoeba histolytica*. Needles. Sol. MeOH, acids, bases, Et₂O; fairly sol. C₆H₆, hexane; poorly sol. H₂O. Mp 175°. pK_{a1} 4.38; pK_{a2} 9.81. Log P 2.27 (calc). λ_{max} 233 (ε 37000); 260 (ε); 283 (ε 15000); 293 (sh) (ε) (EtOH) (Derep). λ_{max} 240 (ε 28000); 273 (MeOH) (Berdy). λ_{max} 240 (ε 28840) (EtOH) (Berdy).

O-Ac: [14761-57-0]
Needles (EtOH aq.). Mp 102°.

O-Me: [14761-56-9]
Mp 90°.

3-Hydroxy: Caerulomycin B

[68560-29-2]
C₁₂H₁₁N₃O₃ 245.237
Metab. of *Streptomyces caeruleus* + tryptophan. Needles (MeOH). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 215-217°. λ_{max} 249 (ε 15500); 273 (sh) (ε 10000); 328 (ε 5620) (EtOH) (Derrep). λ_{max} 249; 328 (MeOH) (Berdy).

3-Methoxy: Caerulomycin C

[68560-30-5]
C₁₃H₁₃N₃O₃ 259.264
Isol. from *Streptomyces caeruleus* + tryptophan. Prisms (EtOH). Sol. MeOH; poorly sol. H₂O. Mp 208-210°. λ_{max} 229 (ε 13500); 263 (ε 9120) (MeOH) (Derrep).

Divekar, P.V. *et al.*, *Can. J. Chem.*, 1967, **45**, 1215 (*struct*)

Ranganathan, S. *et al.*, *Can. J. Chem.*, 1969, **47**, 165 (*synth*)

McInnes, A.E. *et al.*, *Can. J. Chem.*, 1977, **55**, 4159; 1979, **57**, 3200 (*biosynth, cmr*)

Japan. Pat., 1984, 84 51 797; *CA*, **101**, 71051 (*isol*)

Chatterjee, D.K. *et al.*, *Z. Parasitenkd.*, 1984, **70**, 569 (*pharmacol*)

Vining, L. *et al.*, *Can. J. Chem.*, 1988, **66**, 191 (*biosynth*)


Mongin, F. *et al.*, *J.O.C.*, 2002, **67**, 3272-3276 (*Caerulomycin B, C, synth*)

Sammakia, T. *et al.*, *Org. Lett.*, 2002, **4**, 2385-2388 (*Caerulomycin C, synth*)

Caerulomycin D

C-10

[67857-43-6]



C₁₉H₂₁N₃O₇ 403.391


Metab. of *Streptomyces caeruleus*. Needles (EtOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 241-242°. λ_{max} 231 (ε 13800); 267 (ε 10000); 305 (sh) (ε 3310) (MeOH) (Derrep).

McInnes, A.G. *et al.*, *Can. J. Chem.*, 1978, **56**, 1836 (*isol*)

Caerulomycin E

C-11


4-Methoxy-[2,2'-bipyridine]-6-carboxaldehyde, 9CI. 6-Formyl-4-methoxy-2,2'-bipyridine
[114115-50-3]



C₁₂H₁₀N₂O₂ 214.223

Prod. by *Streptomyces caeruleus*. Prisms. Mp 83°.


Vining, L.C. *et al.*, *Can. J. Chem.*, 1988, **66**, 191 (*isol*)



Caesalpinine A

C-12

[85702-14-3]



C₂₅H₃₁N₃O₃ 421.538


Alkaloid from *Caesalpinia digyna* (Caesalpinaceae). Prisms (EtOAc). Mp 240-242°. [α]_D +16.6 (c, 0.42 in CHCl₃).

Mahato, S.B. *et al.*, *J.A.C.S.*, 1983, **105**, 4441 (*isol, cryst struct, uv, cd, ir, pmr, ms*)

Cafamycin

C-13

Kafamycin
[112303-17-0]



C₃₀H₄₁NO₄ 479.658

Polyether-type antibiotic. Analogue of Indanomycin, I-64. Isol. from a *Streptomyces* sp. Active against gram-positive bacteria. Ionophore. Sol. MeOH, bases, C₆H₆, CHCl₃; poorly sol. H₂O. Analogue of Indanomycin, I-64. λ_{max} 245 (ε 20420); 291 (ε 15850) (MeOH) (Berdy).

Murenets, N.V. *et al.*, *Antibiot. Med. Biotechnol.*, 1987, **32**, 811-814 (*isol, pmr, cd, struct*)

Antonenko, Y.N. *et al.*, *Biol. Membr.*, 1988, **5**, 1326 (*props*)

Caffaeoschizine

C-14

[1355-05-1]

C₂₀H₂₀N₂O₄ 352.389

Tentative mol. formula. Struct. unknown. Alkaloid from *Schizozygia caffaeoides* (Apocynaceae). Mp 208-212°. [α]_D²⁵ +25.6 (c, 1 in CHCl₃).

Renner, U. *et al.*, *Experientia*, 1963, **19**, 244-246 (*isol, uv, ir*)

Renner, U. *et al.*, *J. Nat. Prod.*, 1964, **27**, 406-415 (*isol*)

Caffeine, BAN, USAN

C-15

3,7-Dihydro-1,3,7-trimethyl-1H-purine-2,6-dione, 9CI. 1,3,7-Trimethylxanthine. Theine. Guaranine. Coffeine. Methyltheobromine. Cafipel. FEMA 2224. Many other names

[58-08-2]
[5743-12-4]

C₈H₁₀N₄O₂ 194.193

Component of coffee beans (*Coffea arabica*), many other *Coffea* spp., chocolate (*Theobroma cacao*), tea (*Camellia thea*), kolanut (*Cola acuminata*) and several other *Cola* spp. and several other plants (Rubiaceae, Sterculiaceae, Theaceae). Also occurs as a fungal metab. of *Claviceps sorghicola*. Stimulant used in many beverages. Flavouring ingredient. Adenosine receptor antagonist. Widely used CNS, respiratory and cardiac stimulant, main effect on the cerebral cortex. Possesses virucidal props. Chemosterilant against stored grain pests. In the horse is used to treat colic, heat stroke, circulatory disturbance and fatigue, also for horse doping. Mp 236° (phase transition at 141°). Log P -0.06 (uncertain value) (calc). Dimorphic, with β-form and α-form (only stable at high temps).

▶ Adverse systemic effects (by ingestion and other routes). Human and exp. reprod. effects. LD₅₀ (rat, orl) 192 mg/kg. EV6475000

Perchlorate: Mp 89°.

Citrate: Caffeine citrate. Cafcit [69-22-7] Used for the treatment of apnoea in premature infants. Cryst. powder.

N⁹-Me:

[916669-59-5 (iodide, monohydrate), 86180-38-3 (iodide)]

C₉H₁₃N₄O₂[⊕] 209.227

Solid (as iodide). Mp 187-190° (iodide).

[8000-95-1]

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 586A (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 710B (*ir*)

Fischer, E. *et al.*, *Ber.*, 1895, **28**, 2473; 3135 (*synth*)

Bredereck, H. *et al.*, *Chem. Ber.*, 1950, **83**, 201; 1962, **95**, 1902 (*synth*)

Blout, E.R. *et al.*, *J.A.C.S.*, 1950, **72**, 479 (*ir*)

Sutor, D.J. *et al.*, *Acta Cryst.*, 1958, **11**, 453 (*cryst struct*)

Spiteller, G. *et al.*, *Monatsh. Chem.*, 1962, **93**, 632 (*ms*)

Twanmoh, L.-M. *et al.*, *J. Het. Chem.*, 1973, **10**, 187 (*pmr*)

Nicolau, C. *et al.*, *Z. Naturforsch.*, C, 1974, **29**, 475 (*cmr*)

Suzuki, T. *et al.*, *Phytochemistry*, 1976, **15**, 1235 (*biosynth*)

Aranda, J.V. *et al.*, *J. Pediatr.*, 1977, **90**, 467-472 (*Caffeine citrate, use*)

Zubair, M.U. *et al.*, *Anal. Profiles Drug Subst.*, 1986, **15**, 71 (*rev, ir, uv, pmr, cmr, ms, occur, anal*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 921


Arnaud, M.J. *et al.*, *Prog. Drug Res.*, 1987, **31**, 273 (*rev, chem, pharmacol, tox*)

Somani, S.M. *et al.*, *Int. J. Clin. Pharmacol., Ther. Toxicol.*, 1988, **26**, 521 (*rev*)

- Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, CAK500
- Benowitz, N.L. *et al.*, *Annu. Rev. Med.*, 1990, **41**, 277 (rev. *pharmacol*)
- IARC Monog., 1991, **51**, 291 (rev. *tox*)
- Bott, K. *et al.*, *Chem. Ber.*, 1993, **126**, 1955 (*synth*)
- Sitkowski, J. *et al.*, *Spectrochim. Acta A*, 1995, **51**, 839 (*pmr, cmr, N-15 nmr*)
- Encyclopedia of Food and Color Additives*, (ed. Burdock, G.A.), CRC Press, 1997, 363
- Caffeine*, (ed. Spiller, G.A.), CRC Press, 1998, (book)
- Carrier, O. *et al.*, *Clin. Pharmacol. Ther. (St. Louis)*, 1998, **44**, 145-151 (*Caffeine citrate, metab*)
- Parliment, T.H. *et al.*, *ACS Symp. Ser.*, 2000, **754**,
- Bogo, A. *et al.*, *Phytochemistry*, 2000, **54**, 937-939 (*isol, ms*)
- Weinberg, B.A. *et al.*, *The World of Caffeine*, Routledge, 2001, (book)
- Waldvogel, S.R. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 604-605 (rev)
- Zajac, M.A. *et al.*, *Synth. Commun.*, 2003, **33**, 3291-3297 (*synth*)
- Kaskatan-Nebioglu, A. *et al.*, *J. Med. Chem.*, 2006, **49**, 6811-6818 (*methiodide*)
- Lehmann, C.W. *et al.*, *Chem. Eur. J.*, 2007, **13**, 2908-2911 (*cryst struct, dimorph*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CAK500

1-[Caffeoyloxy(carboxy)-methoxy]-1*H*-pyrrole-2,3,5-tricarboxylic acid

[133084-32-9]



$C_{18}H_{13}NO_{13}$ 451.3

Alkaloid from *Parietaria officinalis* (Urticaceae). Off-white granules. Mp 360°.

Budzianowski, J. *et al.*, *Phytochemistry*, 1990, **29**, 3299 (*isol, pmr, uv, cmr*)

Cairomycin C


[78922-12-0]

$C_{23}H_{38}N_6O_6$ 494.59

Peptide antibiotic. Struct. unknown. Contains piperazine nucleus, Lys, Gly, Val, Leu and Asp. Isol. from *Streptomyces* sp. ASC-19. Active against bacteria and fungi. Reddish-brown powder. Sol. $CHCl_3$, EtOAc, Me_2CO ; fairly sol. butanol; poorly sol. H_2O , hexane. Mp 138-140°. λ_{max} 235 ; 275 (MeOH) (Berdy).

- LD₅₀ (mus, ipr) 10 mg/kg. EV6691000
Shimi, I.R. *et al.*, *Antimicrob. Agents Chemother.*, 1981, **19**, 941-944

Caissarine B




$C_{27}H_{32}Br_4N_4O_9$ 876.187

Isol. from the Brazilian sponge *Aplysina caissara*. Glassy solid. λ_{max} 234 (ε 9000); 283 (ε 4300) (MeOH).

Saeki, B.M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 796-799; 2003, **66**, 1039 (*isol, pmr, cmr, ms*)

Caissarone

3,5,7,9-Tetrahydro-3,9-dimethyl-6-(methylimino)-8*H*-purin-8-one, 9*CI* [106145-23-7]



$C_8H_{11}N_5O$ 193.208

Alkaloid from the sea anemone *Bumodoma caissarum*. Inducer of anomalous growth in sea urchin eggs, increases muscle contraction. λ_{max} 228 (ε 21600); 302 (ε 17700) (MeOH) (Derep).

► Teratogen.

Hydrochloride:

Needles (MeOH aq.). Mp 285-290°.

Picrate:

Cryst. (EtOH). Mp 245-250°.

Zelnik, R. *et al.*, *J.C.S. Perkin 1*, 1986, 2051 (*isol, uv, ir, pmr, cmr, cryst struct*)

Fujii, T. *et al.*, *Tet. Lett.*, 1991, **32**, 97 (*synth*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1601-1607; 1867-1869 (*synth*)

Calabacine

$C_{17}H_{25}N_3O_3$ 319.403

Struct. unknown. Alkaloid from the calabar bean (*Physostigma venenosum*) (Fabaceae). Mp 138°. $[\alpha]_D^{27}$ -198 (c, 0.2 in $CHCl_3$).

Picrate:

Prisms (MeOH). Mp 215°.

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 713 (*isol*)

Calabatine

$C_{17}H_{25}NO_2$ 275.39

Struct. unknown. Alkaloid from the Calabar bean (*Physostigma venenosum*) (Fabaceae). Mp 119°. $[\alpha]_D^{24}$ -98 (c, 0.2 in $CHCl_3$).

Picrate:


Needles (MeOH). Mp 128°.

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 713 (*isol*)

C-18

Calafatimine

[77793-42-1]




$C_{38}H_{40}N_2O_7$ 636.743

Alkaloid from *Berberis buxifolia* (Berberidaceae). Cryst. (C_6H_6 /cyclohexane). Mp 180-182°. $[\alpha]_D^{20}$ -141 ($CHCl_3$).

Fajardo, V. *et al.*, *Heterocycles*, 1981, **15**, 1137 (*uv, pmr, ms, struct*)

Calafatine

6,6',7,10,12-Pentamethoxy-2,2'-dimethylberbaman, 9*CI* [73168-72-6]



$C_{39}H_{44}N_2O_7$ 652.786

Alkaloid from *Berberis buxifolia* (Berberidaceae). Needles (C_6H_6 /cyclohexane). Mp 135-137°. $[\alpha]_D^{25}$ -154 (c, 0.28 in $CHCl_3$). CAS did not assign new nos. to corrected structs. for oxides. λ_{max} 258 (log ε 3.32); 281 (log ε 3.82) (MeOH).

$N^{2'}-\alpha$ -Oxide: **Calafatine 2' α -N-oxide**

$C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Berberis buxifolia* (Berberidaceae). Amorph. $[\alpha]_D^{25}$ -48 (c, 0.17 in MeOH). Struct. revised in 1985. λ_{max} 210 (log ε 4.95); 231 (sh) (log ε 4.67); 281 (log ε 3.91) (MeOH).

$N^{2'}-\beta$ -Oxide: **Calafatine 2' β -N-oxide**

$C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Berberis buxifolia* (Berberidaceae). Amorph. $[\alpha]_D^{25}$ -19 (c, 0.14 in MeOH). Struct. revised in 1985. λ_{max} 206 (log ε 4.91); 229 (sh) (log ε 4.63); 280 (log ε 3.85) (MeOH).

O^7 -De-Me: **Berbibuxine. 7-Demethylcalafatine**

$C_{38}H_{42}N_2O_7$ 638.759

Alkaloid from *Berberis buxifolia* and *Berberis darwinii* (Berberidaceae). Amorph. $[\alpha]_D$ -228 (c, 0.19 in $CHCl_3$). CAS no. not found 10-14 CI.

[91126-81-7, 91126-80-6]

Fajardo, V. *et al.*, *Heterocycles*, 1979, **12**, 1559-1562; 1981, **15**, 1132-1139 (*uv, pmr, ms, struct*)

Leet, J.E. *et al.*, *J. Nat. Prod.*, 1983, **46**, 908-912 (*stereochem*)


Leet, J.E. *et al.*, *J.C.S. Perkin 1*, 1984, 651-652; 1985, 1565 (*isol, uv, cd, pmr, ms, oxides*)

Shamma, M. *et al.*, *Personal Communication*, (Berbibuxine, oxides)

Calafianine

[267876-68-6]

C-24



Absolute Configuration


C₂₂H₂₀Br₂N₄O₈ 628.23

Related to Aerothionin, A-158. Stereochem. revised in 2005. Isol. from the sponge *Aplysina gerardogreeni*. Solid. λ_{\max} 210 (sh); 230 (sh); 260 (sh) (EtOH). Encarnacion, R.D. *et al.*, *J. Nat. Prod.*, 2000, **63**, 874-875 (*isol, pmr, cmr*)
Ogamino, T. *et al.*, *Tet. Lett.*, 2005, **46**, 1083-1086 (*synth, struct*)
Bardhan, S. *et al.*, *Org. Lett.*, 2006, **8**, 927-930 (*synth, abs config*)

Calcaridine A

[568561-14-8]

C-25



Relative Configuration

C₂₀H₂₃N₃O₄ 369.419


Isol. from a calcareous sponge *Leucetta* sp. Yellow oil. $[\alpha]_D^{25}$ +1.6 (c, 0.12 in MeOH).

Edrada, R.A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 939-942 (*isol, pmr, cmr*)
Koswatta, P.B. *et al.*, *Org. Lett.*, 2008, **10**, 5055-5058 (*synth, config*)

Calcimycin

A 23187. Antibiotic A 23187. Calimycin [52665-69-7]

C-26

C₂₉H₃₇N₃O₆ 523.628

Polyether antibiotic. Isol. from *Streptomyces chartreusensis*. Calcium ionophore, active against gram-positive and -negative bacteria and fungi. Used as a probe for calcium-potentiated biological processes. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 186-187° (181-182°). $[\alpha]_D^{25}$ -56 (c, 0.01 in CHCl₃). Earlier (positive) opt. rotn. is incorrect. λ_{\max} 202 (ε 22200); 228 (ε 25600); 303 (ε 14500); 370 (ε 5700) (MeOH) (Derep). λ_{\max} 204 (ε 28200); 225 (ε 26200); 278 (ε 18200); 378 (ε 8200) (EtOH) (Berdy). λ_{\max} 204 (E1%/1cm 435); 226 (E1%/1cm 405); 278 (E1%/1cm 290); 380 (E1%/1cm 50) (EtOH/HCl)

(Berdy). λ_{\max} 292 (E1%/1cm 340); 353 (E1%/1cm 90) (EtOH/NaOH) (Berdy).
▶ LD₅₀ (rat, ipr) 9.2 mg/kg. LD₅₀ (mus, ipr) 10 mg/kg. DM4676000

Me ester: [58288-38-3]
Yellow solid. Mp 104-106°. $[\alpha]_D$ -11.2 (c, 0.035 in CHCl₃).

De(methylamino): **Cezomycin**

[83874-22-0]

C₂₈H₃₄N₂O₆ 494.586

Prod. by *Streptomyces chartreusis* NRRL3882 with added tryptophan. Assists in transportation of calcium across a chloroformic membrane. Shows similar antibiotic activity to Calcimycin, C-26. Poorly sol. hexane. Mp 75-77°. $[\alpha]_D^{25}$ +151 (c, 0.012 in CHCl₃).

De(methylamino), 3-hydroxy: **3-Hydroxycezozymycin**. AC 7230. Antibiotic AC 7230C₂₈H₃₄N₂O₇ 510.586

From *Dactylosporangium* sp. AC 7320. Active against gram-positive bacteria. Needles (as Na salt). Sol. MeOH, DMSO, CHCl₃; poorly sol. H₂O, hexane. Mp 300° (Na salt). $[\alpha]_D^{26}$ +328 (c, 0.6 in CHCl₃). Stereochem. not confirmed. λ_{\max} 204 (ε 30700); 257 (ε 15100); 306 (ε 21800) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 50 mg/kg.*14-Demethyl, de(methylamino)*: **11-Demethylcezozymycin**. **Frankiamide**

[726188-57-4]

C₂₇H₃₂N₂O₆ 480.56


Prod. by *Frankia* sp. strain AiPs1. Powder. $[\alpha]_D^{22}$ +192.4 (c, 0.008 in CHCl₃). Struct. of Frankiamide revised in 2003.

[79646-37-0]

Chaney, M.O. *et al.*, *J.A.C.S.*, 1974, **96**, 1932 (*ir, uv, ms, pmr, cmr, cryst struct*)Pressman, B.C. *et al.*, *Annu. Rev. Biochem.*, 1976, **45**, 501 (rev)Smith, G.O. *et al.*, *J.A.C.S.*, 1976, **98**, 1578 (*cryst struct*)Pfeiffer, D.R. *et al.*, *Ann. N.Y. Acad. Sci.*, 1978, **307**, 402 (rev)Evans, D.A. *et al.*, *J.A.C.S.*, 1979, **101**, 6789 (*synth*)Grieco, P.A. *et al.*, *J.O.C.*, 1980, **45**, 3537 (*synth*)David, L. *et al.*, *J. Antibiot.*, 1982, **35**, 1409; 1616 (*Cezomycin*)Martinez, G.R. *et al.*, *J.A.C.S.*, 1982, **104**, 1436 (*synth*)Prudhomme, M. *et al.*, *Experientia*, 1983, **39**, 256 (*Cezomycin, synth*)Zmijewski, M.J. *et al.*, *Tetrahedron*, 1983, **39**, 1255 (*biosynth*)Nakahara, Y. *et al.*, *Tetrahedron*, 1986, **42**, 6465 (*synth, ir, pmr, ms, Me ester*)Yaginuma, S. *et al.*, *J. Antibiot.*, 1987, **40**, 239-241 (*3-Hydroxycezozymycin*)Negri, D.P. *et al.*, *Tet. Lett.*, 1987, **28**, 1063 (*synth, bibl*)Ziegler, F.E. *et al.*, *J.O.C.*, 1989, **54**, 3347 (*synth*)*Synform*, 1990, **8**, 38 (rev. *synth*)Boeckman, R.K. *et al.*, *J.A.C.S.*, 1991, **113**, 5337 (*bibl, synth, Me ester, ir, pmr*)Gruetter, C.A. *et al.*, *Eur. J. Pharmacol.*, 1994, **257**, 275 (*activity*)Klika, K.D. *et al.*, *Z. Naturforsch., B*, 2003, **58**, 1210-1215 (*Frankiamide*)**Caldaphnidine A**

[871261-40-4]

C-27

C₂₃H₂₉NO₃ 367.487


Alkaloid from *Daphniphyllum calycinum*. Cryst. (MeOH). Mp 193-194°. $[\alpha]_D^{20}$ +72 (c, 0.7 in MeOH). λ_{\max} 223 (ε 3440); 297 (ε 9590) (MeOH).

Zhan, Z.-J. *et al.*, *Tetrahedron*, 2005, **61**, 11038-11045 (*isol, pmr, cmr, ms, cryst struct*)

Caldaphnidine C

[871261-42-6]

C-28

C₂₂H₃₃NO₂ 343.508


Alkaloid from *Daphniphyllum calycinum*. Amorph. powder. $[\alpha]_D^{20}$ -36.3 (c, 1 in MeOH).

Zhan, Z.-J. *et al.*, *Tetrahedron*, 2005, **61**, 11038-11045 (*isol, pmr, cmr, ms*)

Caldaphnidine D

[871261-43-7]

C-29

C₂₁H₃₅NO 317.514


Alkaloid from *Daphniphyllum calycinum*. Oil. $[\alpha]_D^{20}$ -73 (c, 1.5 in CHCl₃).

Zhan, Z.-J. *et al.*, *Tetrahedron*, 2005, **61**, 11038-11045 (*isol, pmr, cmr, ms*)

Caldaphnidine E


[871261-44-8]

C-30



$C_{30}H_{49}NO_3$ 471.722
Alkaloid from *Daphniphyllum calycinum*.
Oil. $[\alpha]_D^{20}$ -97 (c, 1.1 in $CHCl_3$).
Zhan, Z.-J. *et al.*, *Tetrahedron*, 2005, **61**,
11038-11045 (*isol*, *pmr*, *cmr*, *ms*)

Calebassine C-31
C-Toxiferine II. C-Strychnotoxine I. C-Calebassine. C-Strychnotoxine Ia. C-Curarine II. C-Calebassine A
[7257-29-6]



$C_{40}H_{48}N_4O_2^{2+}$ 616.845
V. readily forms the di-Me ether, and early refs. are confused between Calebassine and the di-Me ether with the same names being applied to both compds. Alkaloid from calabash curare, *Strychnos divaricans* and several other *Strychnos* spp. (Loganiaceae). Potent neuromuscular blocking agent (*ca.* 0.3 \times Tubocurarine). Mp 216-218° (as dipicrate). $[\alpha]_D^{24}$ +72.1 (H_2O) (as dichloride). Blue-violet colour with $Ce(SO_4)_2$, becoming carmine on standing.

► Very toxic by intravenous route (lethal dose, rabbit *ca.* 135 $\mu g/kg$). FI9000000
Di-Me ether: Mp 232-233° dec. (as dipicrate).

18-Hydroxy: **C-Alkaloid F**

[69356-54-3]
 $C_{40}H_{48}N_4O_3^{2+}$ 632.844

Alkaloid from calabash curare (Loganiaceae). Orange prisms (Me_2CO) (as dipicrate). Mp 209-210° dec. (dipicrate). Blue-violet colour with $Ce(SO_4)_2$, becoming carmine on standing.

► FI9200000
18,18'-Dihydroxy: **C-Alkaloid A. Toxiferine IV**

[6844-05-9]
 $C_{40}H_{48}N_4O_4^{2+}$ 648.844

Alkaloid from calabash curare (Loganiaceae). Mp 228-229° (as dipicrate). $[\alpha]_D$ +64 (H_2O) (as dichloride). Blue-violet colour with $Ce(SO_4)_2$, becoming carmine on standing.

► FI9100000

Wieland, H. *et al.*, *Annalen*, 1941, **547**, 156 (*isol*)

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1946, **29**, 1853; 1952, **35**, 1864; 1953, **36**, 345; 1956, **39**, 440; 1958, **41**, 673; 1961, **44**, 2211 (*isol*, *uv*, *ir*, *pmr*, *struct*)

Zürcher, A. *et al.*, *J.A.C.S.*, 1958, **80**, 1500 (*isol*)

Berlage, F. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 2650-2654 (*C-Alkaloid F, C-Alkaloid A, synth*)


Grdinic, M. *et al.*, *J.A.C.S.*, 1964, **86**, 3351

(*pmr*)

Fehlmann, M. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 303 (*cryst struct*)

Marini-Bettolo, G.B. *et al.*, *CA*, 1980, **92**, 124942c (*isol*)

Calebassine 1, 9CI C-32
[1355-07-3]




$C_{19}H_{23}N_2O_2^{\oplus}$ 311.403
Quaternary alkaloid from Calabash curare (*Strychnos* sp.) (Loganiaceae). Amorph. solid (as chloride). $[\alpha]_D^{22}$ +51.8 (c, 0.305 in EtOH) (chloride).

► EW5500000

Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2586 (*uv*, *ir*, *pmr*, *ms*, *cryst struct*)


Caledonin C-33
[171675-03-9]



$C_{26}H_{43}N_5O_2S$ 489.724
Modified peptide. *Isol.* from the marine tunicate *Didemnum rodriguezii*. Strongly binds Zn^{II} and Cu^I ions. Sol. MeOH, CH_2Cl_2 ; poorly sol. H_2O . Mp 170-172°. $[\alpha]_D^{20}$ +24 (c, 0.145 in MeOH). λ_{max} 234 (ϵ 2816) (MeOH) (Berdy).

Vázquez, M.J. *et al.*, *Tet. Lett.*, 1995, **36**, 8853-8856 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Calliactine C-34



$C_{18}H_{14}N_4O_2$ 318.334

Preferred struct. shown. An isomeric struct. cannot be ruled out.

(*S*)-form

Pigment from the sea anemone *Calliactis parasitica*. λ_{max} 267 (ϵ 12000); 272 (ϵ 11700); 299 (ϵ 9200); 454 (ϵ 8500) (MeOH) (Berdy). λ_{max} 580 (MeOH/NaOH) (Berdy).


Lederer, E. *et al.*, *Bull. Soc. Chim. Fr.*, 1940, **7**, 608-615 (*isol*)

Cimino, G. *et al.*, *Pure Appl. Chem.*, 1986, **58**, 375-386 (*rev*, *bibl*)

Cimino, G. *et al.*, *Tetrahedron*, 1987, **43**, 4023-4030 (*struct*, *uv*, *pmr*, *cmr*)

Voloshina, S. *et al.*, *Int. J. Quantum Chem.*, 2004, **100**, 1104-1113 (*abs config*)

Callichiline, 9CI C-35
[31230-09-8]



$C_{42}H_{48}N_4O_5$ 688.865
Alkaloid from *Callichilia (Hedranthera) subsessilis* and *Hedranthera barteri* (Apocynaceae). Cryst. (MeOH). Mp 208-210° Mp 228-230° (evacuated tube).


Goutarel, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1959, 893 (*isol*, *uv*, *ir*)

Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 2669 (*uv*, *ir*, *pmr*, *ms*)

Agwada, V. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 1939; 1970, **53**, 1567 (*isol*, *uv*, *ir*, *pmr*, *ms*, *ord*)

McPhail, A.T. *et al.*, *Tetrahedron*, 1983, **39**, 3629 (*cmr*, *cryst struct*)

Callimorphine C-36
[74991-73-4]




$C_{15}H_{23}NO_5$ 297.35

Ester of 2,3,5,7a-Tetrahydro-1-hydroxy-1H-pyrrolizine-7-methanol, T-188 of undetd. stereochem. "Metabolite" found in some Arctiid moths which feed as larvae on plants containing pyrrolizidine alkaloids.

Edgar, J.A. *et al.*, *Tet. Lett.*, 1980, **21**, 1383 (*isol*, *ms*, *struct*, *synth*)

L'Empereur, K.M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 360 (*isol*, *pmr*, *cmr*)

Callipeltoside A C-37
[183671-13-8]



C₃₅H₄₈ClNO₁₀ 678.218

Macrolide antibiotic. Closely related to Phorbaside A. Isol. from the sponge *Callipelta* sp. Cytotoxic agent, cell proliferation inhibitor. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ -17.6 (c, 0.04 in MeOH). λ_{max} 250 (ε 6730); 272 (ε 8030); 286 (ε 6270) (MeOH).

Zampella, A. *et al.*, *J.A.C.S.*, 1996, **118**, 11085-11088 (*isol, uv, ir, pmr, cmr*)

Trost, B.M. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 841-843 (*synth, abs config*)

Trost, B.M. *et al.*, *J.A.C.S.*, 2002, **124**, 10396-10415 (*synth, abs config*)


Paterson, I. *et al.*, *Org. Lett.*, 2003, **5**, 4477-4480 (*synth*)

Huang, H. *et al.*, *Org. Lett.*, 2004, **6**, 4383-4385 (*synth*)

Evans, D.A. *et al.*, *Tetrahedron*, 2008, **64**, 4671-4699 (*synth*)

Callophylline B C-38

[141544-39-0]


C₄₃H₅₂N₄O₈ 752.906

Alkaloid from the leaves of *Uncaria callophylla*. λ_{max} 227 ; 290 ; 300 (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 1991, **30**, 3441-3444 (*isol, pmr, cmr*)

Callosine C-39

[213912-09-5]


C₂₁H₃₁NO₇ 409.478

Alkaloid from *Senecio callosus*. Oil. [α]_D²⁰ -14.7 (c, 0.23 in MeOH). λ_{max} 207 (log ε 3.94) (MeOH).

Perez-Castorena, A.-L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1288-1291 (*isol, uv, pmr, cmr*)

Calpurnine C-40

12,13-Dihydroxylupanine
[72047-67-7]



Probable absolute configuration

C₁₅H₂₄N₂O₃ 280.366

Alkaloid from *Calpurnia aurea* subsp. *sylvatica* (Fabaceae). Cryst. (MeOH/EtOAc/Et₂O).

O¹³-2-Pyrrolicarboxylate: 13-(2-Pyrrolylcarbonyl)calpurnine

[72047-68-8]

C₂₀H₂₇N₃O₄ 373.451


Alkaloid from *Calpurnia aurea* (Fabaceae). Cryst. (MeOH/Et₂O/CH₂Cl₂).

Radema, M.H. *et al.*, *Phytochemistry*, 1979, **18**, 2063 (*isol, ms, ir*)

Vermin, W.J. *et al.*, *Acta Cryst. B*, 1979, **35**, 1839 (*cryst struct*)

Calvine C-41

[232279-19-5]



Absolute Configuration

C₁₄H₂₅NO₂ 239.357

Alkaloid from the beetles *Calvia 10-guttata* and *Calvia 14-guttata*. Yellow oil. [α]_D²⁰ +18 (c, 0.66 in CH₂Cl₂).

2-Epimer: 2-Epicalvine

[229976-47-0]

C₁₄H₂₅NO₂ 239.357

Alkaloid from *Calvia 10-guttata* and *Calvia 14-guttata*. Yellow oil. [α]_D²⁰ +8 (c, 0.58 in CH₂Cl₂).


Braekman, J.-C. *et al.*, *Eur. J. Org. Chem.*, 1999, 1749-1755 (*isol, synth, pmr, ms*)

Laurent, P. *et al.*, *Eur. J. Org. Chem.*, 2000, 2057-2062 (*synth, abs config*)

Calvet-Vitale, S. *et al.*, *Tetrahedron*, 2005, **61**, 7774-7782 (*synth*)

Szolesányi, P. *et al.*, *Tet. Lett.*, 2008, **49**, 1357-1360 (*synth*)

Calycanthine, 9CI C-42



(+)-form

C₂₂H₂₆N₄ 346.474

Various numbering schemes possible. That shown is based on numbering of the quinoline residues.

(+)-form [595-05-1]

Alkaloid from *Calycanthus floridus* (Carolina allspice), other *Calycanthus* spp., *Psychotria forsteriana* and *Pali-courea ovalis*. Shows cytotoxic activity. Platelet aggregation inhibitor. Mp 250-251° (anhyd.). [α]_D²⁰ +684. λ_{max} 252 (log ε 4.26); 310 (log ε 3.8) (MeCN). Hydrobromide (1:2): Mp 213-214°.

(-)-form [85548-42-1]

Alkaloid from Colombian poison-dart frog *Phyllobates terribilis*. Also isol. from stem bark and fruits of *Psychotria forsteriana* (Rubiaceae). Mp 245°. [α]_D²⁵ -570 (MeOH).

1,2-Didehydro: *Glomerulatine C*

[188846-23-3]

C₂₂H₂₄N₄ 344.458

Minor alkaloid from aerial parts of *Psychotria glomerulata*. λ_{max} 274 ; 304 (sh) (EtOH).

1,1',2,2'-Tetrahydro: Glomerulatine A

[188846-03-9]

C₂₂H₂₂N₄ 342.443

Alkaloid from aerial parts of *Psychotria glomerulata*. [α]_D²⁰ -466.1 (c, 0.36 in CHCl₃). λ_{max} 274 ; 304 (sh) (EtOH).

1,1',2,2'-Tetrahydro, N-de-Me: Glomerulatine B

[188846-22-2]

C₂₁H₂₀N₄ 328.416

Minor alkaloid from aerial parts of *Psychotria glomerulata*. λ_{max} 279 (MeCN).

(±)-form [16739-56-3]

Synthetic. Mp 253-258°.

meso-form

Synthetic. Mp 265-268°.

Woodward, R.B. *et al.*, *Proc. Chem. Soc., London*, 1960, 76-78 (*struct*)

Hamor, T.A. *et al.*, *J.C.S.*, 1962, 194-205 (*cryst struct*)

Hendrikson, J.B. *et al.*, *Tetrahedron*, 1964, **20**, 565-579 (*synth, ms, pmr, uv*)

Hall, E.S. *et al.*, *Tetrahedron*, 1967, **23**, 4131-4141 (*synth, ir, uv, ms, pmr*)

Beecham, A.F. *et al.*, *Nature (London)*, *Phys. Sci.*, 1973, **244**, 30-32 (*config*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1983, **39**, 41-47 ((-)-form, *isol*)

Adjibade, Y. *et al.*, *Phytochemistry*, 1992, **31**, 317-319 ((-)-form, *isol, uv, ir, pmr, cmr, ms*)

Link, J.T. *et al.*, *J.A.C.S.*, 1996, **118**, 8166-8167 (*synth*)

Solis, P.N. *et al.*, *Phytochemistry*, 1997, **44**, 963-969 (*Glomerulatinines*)


Arteaga de Garcia, L. *et al.*, *Rev. Colomb. Quim.*, 1997, **26**, 55-57; *CA*, **128**, 110442q (*activity*)

Verotta, L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 392-396 ((-)-form, *isol, pmr, cd*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1999, **121**, 7702-7703 (*synth*)

Movassaghi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 3725-3728 (*synth*)

Calcyilactone A C-43



Relative Configuration

C₂₂H₂₉NO₃ 355.476

Alkaloid from the leaves of *Daphniphyllum calycillum*. Oil. [α]_D²⁵ -128 (c, 0.62 in CHCl₃).

Di, Y.-T. *et al.*, *Tet. Lett.*, 2006, **47**, 5329-5331 (*isol, pmr, cmr*)

Calycine† C-44

C₂₃H₃₁NO₃ 369.503

Struct. unknown. Alkaloid from the leaves and bark of *Daphniphyllum calycinum* and the bark of *Daphniphyllum glaucescens* (Daphniphyllaceae). Cryst.

(Me₂CO). Mp 205° (vac.). [α]_D²⁰ +58.7 (CHCl₃).

Picrate:

Cryst. (MeOH). Mp 215-218° (vac. dec.).

Methiodide:

Cryst. (EtOH/Me₂CO). Mp 252-254° (vac. dec.).

Ethiodide:

Cryst. (EtOH/Me₂CO). Mp 245-246°.

3,5-Dinitrobenzoyl:


Cryst. (EtOH or Me₂CO). Mp 218-221° (vac. dec.).

Arthur, H.R. *et al.*, *Phytochemistry*, 1965, **4**, 627-629 (*isol, ir, pmr*)

Calyciphylline B

C-45

[596799-33-6]



Absolute Configuration

C₂₂H₃₁NO₃ 357.492
Alkaloid from *Daphniphyllum calycinum* and *Daphniphyllum subverticillatum*.
Needles (Me₂CO). Mp 178° dec. [α]_D²⁰ -70.7 (c, 0.33 in MeOH).

N-Deoxy: Deoxycalyciphylline B

[619326-74-8]

C₂₂H₃₁NO₂ 341.492

Alkaloid from the stems of *Daphniphyllum subverticillatum*. Cryst. (Me₂CO). Mp 182° dec. [α]_D²⁰ -96 (c, 1.2 in MeOH).

5-Epimer: Isocalyciphylline B

[619326-76-0]

C₂₂H₃₁NO₃ 357.492

Yellow gum. Artifact.

5-Epimer, N-deoxy: Deoxyisocalyciphylline B

[619326-75-9]

C₂₂H₃₁NO₂ 341.492

Alkaloid from the stems of *Daphniphyllum subverticillatum*. Pale solid. Mp 86-88°. [α]_D²⁰ -70.6 (c, 0.7 in MeOH).


Yang, S.-P. *et al.*, *J.O.C.*, 2003, **68**, 7961-7966 (*isol, pmr, cmr, ms, cryst struct*)

Morita, H. *et al.*, *Org. Lett.*, 2003, **5**, 2895-2898 (*isol, pmr, cmr*)

Calyciphylline C

C-46

[930774-41-7]



C₂₃H₃₁NO₃ 369.503


Alkaloid from the leaves of *Daphniphyllum calycinum*. Amorph. solid. [α]_D¹⁸ -21.5 (c, 0.2 in MeOH). λ_{max} 215 (ε 4760) (MeOH).

Saito, S. *et al.*, *Tet. Lett.*, 2007, **48**, 1587-1589 (*isol, pmr, cmr*)

Calyciphylline D

C-47

[929638-01-7]



C₂₉H₄₅NO₃ 455.679


Alkaloid from the leaves of *Daphniphyllum calycinum*. Amorph. solid. [α]_D²⁴ -25.6 (c, 0.5 in MeOH).

Saito, S. *et al.*, *Org. Lett.*, 2007, **9**, 1207-1209 (*isol, pmr, cmr*)

Calyciphylline E

C-48

[943528-89-0]



Relative Configuration

C₂₃H₃₁NO₄ 385.502


Alkaloid from the leaves of *Daphniphyllum calycinum*. Amorph. solid. [α]_D²¹ -39 (c, 0.5 in CHCl₃).

Saito, S. *et al.*, *Tet. Lett.*, 2007, **48**, 3809-3812 (*isol, pmr, cmr*)

Calyciphylline F

C-49

[943528-90-3]



Relative Configuration

C₂₂H₃₅NO₂ 345.524


Alkaloid from the leaves of *Daphniphyllum calycinum*. Amorph. solid. [α]_D¹⁶ -37 (c, 1 in CHCl₃).

Saito, S. *et al.*, *Tet. Lett.*, 2007, **48**, 3809-3812 (*isol, pmr, cmr*)

Calyciphylline G

C-50

[947730-34-9]



Relative Configuration

C₂₃H₃₀NO₂⁺ 352.496


Alkaloid from the stems of *Daphniphyllum calycinum*. Cytotoxic. Amorph. solid. [α]_D²¹ +8.5 (c, 1 in MeOH). Counterion not specified.

Saito, S. *et al.*, *Tet. Lett.*, 2007, **48**, 5693-5695 (*isol, pmr, cmr*)

Calyciphylline H

C-51

[1012789-71-7]



Relative Configuration

C₂₃H₃₁NO₃ 369.503

Alkaloid from *Daphniphyllum calycinum*. Amorph. solid. [α]_D²² +36.6 (c, 0.5 in CHCl₃).

1-Deoxy, 2-hydroxy, 3,4-dihydro: Calyciphylline I

[1012789-72-8]

C₂₃H₃₃NO₃ 371.519


Alkaloid from *Daphniphyllum calycinum*. Amorph. solid. [α]_D²² -9.2 (c, 0.5 in CHCl₃).

Saito, S. *et al.*, *Tetrahedron*, 2008, **64**, 1901-1908 (*isol, pmr, cmr*)

Calyciphylline J

C-52

[1012789-73-9]



Absolute Configuration


C₂₂H₂₉NO₃ 355.476

Alkaloid from *Daphniphyllum calycinum*. Amorph. solid. [α]_D²² -45 (c, 1 in CHCl₃).

Saito, S. *et al.*, *Tetrahedron*, 2008, **64**, 1901-1908 (*isol, pmr, cmr*)

Calycosidine

[112262-96-1]

C₃₃H₃₈N₆ 518.703

Structurally related to Hodgkinsine, H-311. Alkaloid from the stem bark of *Calycodendron milnei* (Rubiaceae). Also obt. by acid catalyzed isomerisation of Hodgkinsine. Amorph. $[\alpha]_D^{20}$ -18 (c, 1 in CHCl₃).

Libot, F. *et al.*, *J. Nat. Prod.*, 1987, **50**, 468

(isol, uv, ir, pmr, cmr, ms)


Libot, F. *et al.*, *Heterocycles*, 1988, **27**, 2381

(uv, ir, pmr, cmr, ms, struct)

Calycotomine

C-54

1,2,3,4-Tetrahydro-6,7-dimethoxy-1-isoquinolinemethanol, 9CI. 1,2,3,4-Tetrahydro-1-hydroxymethyl-6,7-dimethoxyisoquinoline



(R)-form

C₁₂H₁₇NO₃ 223.271**(R)-form** [101141-28-0]

Synthetic. Mp 149-150° (145-148°). $[\alpha]_D^{20}$ -36 (c, 1.0 in H₂O) (-28.9).

(S)-form [486-99-7]

Alkaloid from *Calycotome spinosa*, *Calycotome profliferus* and *Acacia concinna* (Fabaceae). Also obt. by resolin. Mp 149-150° (synthetic); 139-141° (natural). $[\alpha]_D^{20}$ +36 (H₂O) (synthetic). $[\alpha]_D^{20}$ +21 (H₂O) (natural). The physicochemical data of optically pure (+)-calycotomine (Brossi, *et al.*) do not correspond with those recorded for the alkaloid isolated by White, which seems to be a partial racemate.

Hydrochloride: Mp 204-205°. $[\alpha]_D^{20}$ +33 (c, 1.0 in H₂O) (opt. pure).

Benzyl ether: [204056-59-7]Syrup. $[\alpha]_D^{25}$ +19.3 (c, 1.23 in EtOH).**(±)-form** [4356-47-2]

Alkaloid from *Calycotome spinosa*, *Calycotome profliferus* and *Cytisus nigricans* (Fabaceae). Cryst. (C₆H₆ or EtOAc/petrol). Mp 139-141° (134-135°).

Hydrochloride:Cryst. (C₆H₆/MeOH). Mp 196.7-198° (192°).**Picrate:** Mp 203-204° (163-166°).**N-Ac:**

Cryst. (EtOAc). Mp 166-167°.

(ξ)-form

O⁷-De-Me, N-Me: 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-methyl-1-isoquinolinemethanol. 1,2,3,4-Tetrahydro-7-hydroxy-1-hydroxymethyl-6-methoxy-2-methylisoquinoline. **Hedycarine**


C₁₂H₁₇NO₃ 223.271

Alkaloid from *Hedycarya baudouinii* (Monimiaceae).

White, E.P. *et al.*, *N.Z. J. Sci. Technol., Sect. B*, 1944, **25**, 137; 1951, **33**, 38 (isol, struct)Důbravková, L. *et al.*, *Chem. Zvesti*, 1958, **12**, 459 (synth, uv)Battersby, A.R. *et al.*, *J.C.S.*, 1959, 1909; 1960, 1214 (synth, config)Brossi, A. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1558 (resolin)Bennington, F. *et al.*, *J.O.C.*, 1961, **26**, 194 (synth)Chatterjee, A. *et al.*, *J.O.C.*, 1962, **27**, 309 (synth, uv)Gibson, H.W. *et al.*, *J. Het. Chem.*, 1964, **1**, 251 (synth)Gupta, G.L. *et al.*, *Planta Med.*, 1971, **19**, 55 (isol)Czarnocki, Z. *et al.*, *Can. J. Chem.*, 1986, **64**, 2205-2210 (synth, pmr, ms)Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 745 (Hedycarine)Lenz, G.R. *et al.*, *Heterocycles*, 1987, **26**, 721 (synth, uv, pmr)Singh, H. *et al.*, *Indian J. Chem., Sect. B*, 1989, **28**, 802 (synth)Czarnocki, Z. *et al.*, *J. Chem. Res., Synop.*, 1992, 334 (synth)Kaufman, T.S. *et al.*, *Synth. Commun.*, 1993, **23**, 473 (synth)Suau, R. *et al.*, *Heterocycles*, 1996, **43**, 545 (synth)Morimoto, T. *et al.*, *Tetrahedron: Asymmetry*, 1998, **9**, 183-187 (S-form, synth, benzyl ether)Antri, A.E. *et al.*, *Molecules*, 2004, **9**, 650-657 (isol, pmr, cmr, cryst struct)Kaufman, T.S. *et al.*, *Synthesis*, 2005, 339-360 (rev, synth)**Calyculin A**

C-55

[101932-71-2]



Absolute Configuration

C₅₀H₈₁N₄O₁₅P 1009.181

Isol. from the marine sponge *Discodermia calyx* and from *Lamellomorpha strongylata*. Inhibitor of protein phosphatases. Apoptosis inducer. Needles (Me₂CO/Et₂O/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 247-249°. $[\alpha]_D^{25}$ +59.8 (c, 0.12 in EtOH). Major metab. The opt. rotn., uv data and yield reported in the original paper were incorr. λ_{max} 230 (€ 12000); 341 (€ 19000) (EtOH) (Derep).

O-Dephosphono: **Dephosphonocalyculin A** [191421-64-4]

C₅₀H₈₀N₄O₁₂ 929.202

Isol. from *Dysidea calyx*. Protein phosphatase inhibitor. Solid.

15-Amide: Calyculinamide A

[187991-80-6]

C₅₀H₈₃N₄O₁₆P 1027.197

Isol. from *Dysidea calyx* and *Lamellomorpha strongylata*. $[\alpha]_D^{20}$ -41 (c, 0.5 in EtOH). $[\alpha]_D^{20}$ -18 (c, 0.005 in MeOH). Isol. by two groups of workers. UV data not in agreement.

N-De-Me: N-Demethylcalyculin A

[187944-08-7]

C₄₉H₇₉N₄O₁₅P 995.155

Isol. from *Discodermia calyx*. Amorph. solid. $[\alpha]_D^{20}$ -18 (c, 0.01 in MeOH). λ_{max} 220 (€ 9000); 340 (€ 8000) (EtOH).

(9Z)-Isomer: Calyculin E

[133445-05-3]

C₅₀H₈₁N₄O₁₅P 1009.181

Isol. from *Discodermia calyx* and *Lamellomorpha strongylata*. Sol. MeOH, CHCl₃. $[\alpha]_D^{23}$ -83 (c, 0.2 in EtOH). λ_{max} 230; 325 (€ 22000) (MeOH) (Berdy). λ_{max} 227 (€ 13800); 321 (€ 16400) (EtOH) (Berdy).

(13E)-Isomer: Calyculin B

[107537-44-0]

C₅₀H₈₁N₄O₁₅P 1009.181

From *Discodermia calyx* and *Lamellomorpha strongylata*. Antitumour agent. Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{20}$ -61 (c, 0.05 in EtOH). λ_{max} 230 (€ 12000); 341 (€ 19000) (EtOH) (Derep). λ_{max} 230 (€ 18000); 341 (€ 25000) (EtOH) (Berdy).

(13E)-Isomer, 15-amide: Calyculinamide B

[188292-66-2]

C₅₀H₈₃N₄O₁₆P 1027.197

Isol. from *Lamellomorpha strongylata*. Protein phosphatase inhibitor. Amorph. solid. $[\alpha]_D^{20}$ -27 (c, 0.1 in EtOH). λ_{max} 227 (€ 26000); 335 (€ 45600) (EtOH). λ_{max} 227 (€ 26000); 335 (€ 45600) (MeOH) (Berdy).

(9Z,13E)-Isomer: Calyculin F

[133445-06-4]

C₅₀H₈₁N₄O₁₅P 1009.181

Isol. from *Discodermia calyx* and *Lamellomorpha strongylata*. Sol. MeOH, CHCl₃. $[\alpha]_D^{23}$ -33 (c, 0.2 in EtOH). λ_{max} 230; 323 (€ 24000) (MeOH) (Berdy). λ_{max} 227 (€ 15200); 321 (€ 18400) (EtOH) (Berdy).

(9Z,13E)-Isomer, 15-amide: Calyculinamide F

[187991-81-7]

C₅₀H₈₃N₄O₁₆P 1027.197

Isol. from *Dysidea calyx*. Protein phosphatase inhibitor. Amorph. solid. $[\alpha]_D^{20}$ -23 (c, 0.01 in MeOH). λ_{max} 228 (€ 11000); 236 (€ 10000); 315 (€ 14000) (EtOH). λ_{max} 228 (€ 11000); 236 (€ 10000); 315 (€ 14000) (MeOH) (Berdy).

Kato, Y. *et al.*, *J.A.C.S.*, 1986, **108**, 2780 (isol, uv, ir, pmr, cmr, cryst struct)Kato, Y. *et al.*, *J.O.C.*, 1988, **53**, 3930 (isol, pmr, cmr)Okada, A. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 2765-2771 (Calyculins E-F)Matsunaga, S. *et al.*, *Tetrahedron*, 1991, **47**, 2999Evans, D.A. *et al.*, *J.O.C.*, 1992, **57**, 1958; 1961; 1964 (synth)Evans, D.A. *et al.*, *Spec. Publ. - R. Soc. Chem.*, 1993, **119**, 117 (rev)Tanimoto, N. *et al.*, *Angew. Chem., Int. Ed.*, 1994, **33**, 673 (synth)

- Dumdei, E.J. *et al.*, *J.O.C.*, 1997, **62**, 2636 (*Calyculinamides*)
 Matsunaga, S. *et al.*, *J.O.C.*, 1997, **62**, 2640 (*Calyculinamides*)
 Matsunaga, S. *et al.*, *Tet. Lett.*, 1997, **38**, 3763 (*Dephosphocalyculin A*)
 Smith, A.B. *et al.*, *J.O.C.*, 1998, **63**, 7596-7597 (*synth*)
 Volter, K.E. *et al.*, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 717-722 (*pmr, cryst struct*)
 Smith, A.B. *et al.*, *J.A.C.S.*, 1999, **121**, 10468-10477; 10478-10486 (*synth*)

Calyculin C C-56

[107537-45-1]
 As Calyculin A, C-55 with
 R = CH₃

C₅₁H₈₃N₄O₁₅P 1023.208
 Isol. from *Discodermia calyx*. Antitumour agent. Smooth muscle contractor. Inhibitor of protein phosphatases. Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ -65 (c, 0.05 in EtOH). λ_{max} 230 (€ 12000); 341 (€ 19000) (EtOH) (Derep). λ_{max} 230 (€ 15000); 340 (€ 19000) (EtOH) (Berdy).

(13E)-form**Calyculin D**

[107447-09-6]
 From *Discodermia calyx*. Antitumour agent. Smooth muscle contractor. Inhibitor of protein phosphatases. Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ -41 (c, 0.05 in EtOH). λ_{max} 230 (€ 12000); 341 (€ 19000) (EtOH) (Derep).

(9Z)-Isomer: Calyculin G

[135212-39-4]
 C₅₁H₈₃N₄O₁₅P 1023.208
 Isol. from *Discodermia calyx*. Protein phosphatase inhibitor. Smooth muscle contractor. Sol. MeOH, CHCl₃. [α]_D²³ -81 (c, 0.1 in EtOH). λ_{max} 227 (€ 19000); 321 (€ 20800) (EtOH) (Berdy).


(9Z,13E)-Isomer: Calyculin H

[135212-40-7]
 C₅₁H₈₃N₄O₁₅P 1023.208
 Isol. from *Discodermia calyx*. Protein phosphatase inhibitor. Smooth muscle contractor. Sol. MeOH, CHCl₃. [α]_D²³ -36 (c, 0.05 in EtOH). λ_{max} 227 (€ 22800); 321 (€ 28000) (EtOH) (Berdy).

- Kato, Y. *et al.*, *J.O.C.*, 1988, **53**, 3930 (*isol, pmr, cmr, struct*)
 Matsunaga, S. *et al.*, *Tetrahedron*, 1991, **47**, 2999 (*Z-isomers, isol*)
 Scarlato, G.R. *et al.*, *J.O.C.*, 1996, **61**, 6139; 6153 (*synth*)
 Ogawa, A.K. *et al.*, *J.A.C.S.*, 1998, **120**, 12435-12442 (*synth*)

Calyculin J C-57

[187991-79-3]




C₅₀H₈₀BrN₄O₁₅P 1088.078
 Isol. from the sponge *Discodermia calyx*. Protein phosphatase inhibitor. Yellow solid. [α]_D²⁰ -10 (c, 0.08 in MeOH). λ_{max} 228 (€ 11000); 291 (€ 14000); 304 (€ 20000) (EtOH). λ_{max} 228 (€ 11000); 291 (€ 14000); 304 (€ 20000); 318 (€ 19000) (MeOH) (Berdy).

Matsunaga, S. *et al.*, *J.O.C.*, 1997, **62**, 2640 (*isol, uv, pmr, cmr*)

Calydaphninone

[929613-79-6]



C₂₃H₃₁N₄O 385.502
 Alkaloid from the leaves of *Daphniphyllum calycillum*. Powder. Mp 202-204°. [α]_D²⁰ -66.7 (c, 0.38 in CHCl₃).

Di, Y.-T. *et al.*, *Org. Lett.*, 2007, **9**, 1355-1358 (*isol, pmr, cmr, cryst struct*)

Calystegine† C-59

C₂₅H₂₃NO₆ 433.46
 Struct. unknown. Alkaloid from the Chinese drug Chin-Kuo-Lan (*Calystegia hydraceae*) (Convolvulaceae). Brown prisms (Me₂CO/Et₂O). Mp 203°.

Hydrochloride:

Prisms. Mp 230°.

Hydrobromide:

Yellow needles. Mp 232°.

Perchlorate:

Yellow needles. Mp 268°.

Nitrate:

Yellow needles. Mp 233-234°.

Picrate:

Brown needles. Mp 220°.


Methiodide:

Pale red needles. Mp 238°.

Chu, J.-H. *et al.*, *Chem. Zentralbl.*, 1960, **131**, 1859 (*isol, props*)

Calyxamine A

[199105-62-9]



C₁₂H₂₁NO 195.304
 Alkaloid from the sponge *Calyx podatypa*. Needles (CHCl₃/hexane) (as trifluoroacetate salt). λ_{max} 205 (€ 1000) (MeOH) (TFA salt).


Δ^{4,11}-Isomer: **Calyxamine B**

[150710-72-8]
 C₁₂H₂₁NO 195.304
 Alkaloid from *Calyx podatypa*. Cryst. (as trifluoroacetate salt). Possible artifact. λ_{max} 231 (€ 10400) (MeOH) (TFA salt).

Rodriguez, A.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1331-1333 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Camoensidine

[58845-84-4]



C₁₄H₂₂N₂O 234.341
 Alkaloid from the roots of *Camoensia maxima* and the seeds of *Camoensia brevicalyx* (Fabaceae). Noncryst. [α]_D²⁰ -67 (c, 1 in EtOH).

15-N-Oxide: Camoensidine N-oxide

C₁₄H₂₂N₂O₂ 250.34
 Alkaloid from the stems of *Maackia tashiroi*. Cryst. [α]_D²⁷ -57 (c, 0.79 in EtOH).

12S-Hydroxy: 12-Hydroxycamoensidine

[82041-86-9]
 C₁₄H₂₂N₂O₂ 250.34
 Alkaloid from the seeds of *Camoensia brevicalyx* (Fabaceae). Needles (MeOH). Mp 200°. [α]_D²⁰ -50 (c, 0.1 in MeOH).

11-Epimer: Tetrahydroleontidine


[35597-15-0]
 C₁₄H₂₂N₂O 234.341
 Alkaloid from stems and leaves of *Maackia amurensis* (Fabaceae). Mp 62-67°. [α]_D²⁰ -45.3. Probable stereochem.

[58845-85-5, 81979-85-3]

- Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 631; *Chem. Nat. Compd. (Engl. Transl.)*, 611 (*synth, Tetrahydroleontidine*)
 Santamaria, J. *et al.*, *Phytochemistry*, 1975, **14**, 2501 (*isol, ir, pmr, ms, struct, synth*)
 Waterman, P.G. *et al.*, *Phytochemistry*, 1982, **21**, 215 (*isol, ir, pmr, cmr, ms, deriv*)
 Kinghorn, A.D. *et al.*, *Phytochemistry*, 1982, **21**, 2269 (*isol, Tetrahydroleontidine*)
 Ohmiya, S. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1123 (*N-oxide*)

Camoensine

C-62
 1,2,3,5,6,12,13,13a-Octahydro-6,13-methano-10H-pyrido[1,2-a]pyrrole[1,2-e][1,5]diazocin-10-one, 9CI
 [58845-83-3]



C₁₄H₁₈N₂O 230.309
 Alkaloid from the roots of *Camoensia maxima* and the seeds of *Camoensia*

brevicalyx (Fabaceae). Brown amorph. solid. $[\alpha]_D^{20}$ -108 (c, 1.0 in CHCl_3). $[\alpha]_D^{20}$ -186 (c, 1 in EtOH).

Methiodide:

Cryst. + 0.5H₂O. Mp 292°.

12 α -Hydroxy: 12 α -Hydroxycamoensine

C₁₄H₁₈N₂O₂ 246.308

Alkaloid from the seeds of *Camoensia brevicalyx* (Fabaceae). Brown amorph. solid. $[\alpha]_D^{20}$ -115 (c, 1 in MeOH).

Tetrahydro: see Camoensidine, C-61

11-Epimer: Leontidine

[35721-27-8]

C₁₄H₁₈N₂O 230.309

Alkaloid from *Leontice eversmannii*, *Leontice albertii* and *Camoensia maxima* (Fabaceae, Leonticaceae). Cryst. (EtOAc). Mp 119° Mp 145°.

11-Epimer: perchlorate: Mp 278°.


Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1971, 7, 631; *Chem. Nat. Compd. (Engl. Transl.)*, 611 (isol, ms, struct, synth, bibl, Leontidine) Santamaria, J. *et al.*, *Phytochemistry*, 1975, 14, 2501 (isol, uv, ir, pmr, ms, struct, synth, Camoensine, Leontidine)

Waterman, P.G. *et al.*, *Phytochemistry*, 1982, 21, 215 (isol, uv, ir, pmr, cmr, ms, deriv)

Campedine

C-63

1-Ethyl-5-(7-hydroxy-1,3-benzodioxol-5-yl)-2-piperidinone, 9CI. 1-Ethyl-5-(3-hydroxy-4,5-methylenedioxyphenyl)-2-piperidone
[29028-18-0]



C₁₄H₁₇NO₄ 263.293

Alkaloid from seeds of *Campanula medium* (Campanulaceae). No phys. props. reported.

Döpke, W. *et al.*, *Pharmazie*, 1970, 25, 128 (isol, ir, uv)

Campestrine

C-64

C₁₃H₁₉NO₃ 237.298


Pyrrolizidine alkaloid. Struct. unknown. Alkaloid from *Senecio campestris* var. *maritimus* (Asteraceae). Mp 93°.

Blackie, J.J. *et al.*, *Pharm. J.*, 1937, 138, 102-104

Camptacumanine

C-65

[121696-54-6]



C₂₁H₁₈N₂O₄ 362.384


Anomalous indole alkaloid. The struct. does not correspond to Secologanin derivation. Minor alkaloid from seeds of *Camptotheca acuminata* (Nyssaceae).

Lin, L. *et al.*, *Huaxue Xuebao*, 1988, 46, 1207; *CA*, 111, 54155h

Camptacumotine

C-66

1,2,8,13-Tetrahydro-2-methyl-4H-indolo[2,3-a]pyrano[3,4-g]quinolizine-4,5(7H)-dione, 9CI
[121696-52-4]



C₁₉H₁₆N₂O₃ 320.347


Minor alkaloid from seeds of *Camptotheca acuminata* (Nyssaceae).

Lin, L. *et al.*, *Huaxue Xuebao*, 1988, 46, 1207; *CA*, 111, 54155h

Camptine

C-67

[56236-21-6]



C₂₀H₂₈N₂O₂ 328.453

Struct. not fully known. Alkaloid from *Strychnos camptoneura* bark (Loganiaceae). Mp 174-176°. No further ref. to this comp. to 14CI, 2001.

Garnier, J. *et al.*, *Plant. Med. Phytother.*, 1974, 8, 281-286 (isol, uv, ir, ms)

Camptinine

C-68

[56378-66-6]

C₄₃H₅₂N₄O₆ 720.907


Dimeric alkaloid prob. contg. Rutiline, R-63 residue. Struct. unknown. Alkaloid from *Strychnos camptoneura* bark (Loganiaceae). Noncryst.

Garnier, J. *et al.*, *Plant. Med. Phytother.*, 1974, 8, 281-286 (isol, uv, ms)

Camptoneurine

C-69

[38930-42-6]



C₂₂H₁₇N₃O₄ 387.394


Alkaloid from *Strychnos camptoneura*. Mp 269-270°. $[\alpha]_D^{20}$ +189 (c, 1 in CHCl_3).

Koch, M. *et al.*, *Ann. Pharm. Fr.*, 1972, 30, 299-306 (isol, uv, ir, pmr, ms, struct)

Camptothecin

C-70

4-Ethyl-4-hydroxy-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione, 9CI. CPT. NSC 94600



(S)-form

C₂₀H₁₆N₂O₄ 348.357

Log P 0.6 (calc). Many syntheses reported. Those before 1997 are not shown.

▶ Adverse effects reported when used therapeutically. Na salt produces adverse haemopoietic effects. LD₅₀ (mus, orl) 27 mg/kg. LD₅₀ (mus, ipr) 64 mg/kg.

(S)-form [7689-03-4]

Alkaloid from *Camptotheca acuminata*, *Merrilliodendron megacarpum*, *Mappia foetida*, *Ervatania heyneana*, *Nothapodytes foetida* and *Ophiorrhiza mungos* (Nyssaceae, Rubiaceae, Apocynaceae). Also a metab. of a fungus isol. from the bark of *Nothapodytes foetida*. Shows potent antineoplastic activity in exp. animals. Functions by enhancing binding of topoisomerase I to DNA, thus promoting DNA strand breaks. Used clinically in China against gastrointestinal tumours. Also shows plant growth regulatory and insect chemosterilant props. Severe side effects and rapid hydrol. at physiological pH have inhibited widespread clinical use. Also shows anti-HIV and antiprotozoal activity. Pale yellow needles (MeOH). Sol. MeOH, CHCl_3 ; poorly sol. H₂O, hexane. Mp 264-267° dec. $[\alpha]_D^{20}$ +31.3 ($\text{CHCl}_3/\text{MeOH}$, 4:1). Log P 0.6 (calc). Nonbasic. Some semisynthetic analogues, e.g. Belotecan, are marketed. λ_{max} 220 (ε 37320); 254 (ε 24230); 290 (ε 4980); 370 (ε 19900) (MeOH) (Berdy). λ_{max} 218 (ε 42700); 253 (ε 32400); 289 (ε 6460); 358 (ε 21900); 368 (ε 21900) (MeOH) (Derep).

▶ UQ0492000

O-β-D-Glucopyranoside: 20-O-β-D-Glucopyranosylcamptothecin

[246031-86-7]

C₂₆H₂₆N₂O₉ 510.499

Alkaloid from *Mostuea brunonis*. Amorph. yellow solid. $[\alpha]_D^{20}$ +23.5 (c, 0.52 in MeOH). λ_{max} 217 (log ε 4); 245 (log ε 3.8); 285 (log ε 3.2); 332 (log ε 3.3); 369 (log ε 3.5) (MeOH).

Ac: O-Acetylcamptothecin

[7688-64-4]

C₂₂H₁₈N₂O₅ 390.395


Alkaloid from stems of *Nothapodytes*

- foetida* (Icacinaeae). Exhibits potent cytotoxicity. Yellowish needles. Mp 272-275° (271-274° dec.). λ_{\max} 219 ; 253 ; 288 ; 358 ; 369 (MeOH) (Berdy).
- Hexanoyl: 20-Hexanoylcampthoecin**
[72594-33-3]
C₂₆H₂₆N₂O₅ 446.502
Minor alkaloid from the bark of *Camptotheca acuminata* (Nyssaceae). Mp 238-242°. $[\alpha]_{\text{D}}^{25}$ -26 (c, 0.0017 in CHCl₃).
- 18,19-Didehydro: 18,19-Dehydrocampthoecin**
[119403-33-7]
C₂₀H₁₄N₂O₄ 346.342
Alkaloid from the wood of *Nothapodytes foetida* (Icacinaeae). Pale yellow solid.
- 9-Hydroxy, 9-β-D-glucopyranoside: 9-β-D-Glucopyranosyloxycampthoecin**
[201805-58-5]
C₂₆H₂₆N₂O₁₀ 526.499
Alkaloid from regenerated plants of *Ophiorrhiza pumila*. λ_{\max} 262 ; 304 ; 319 ; 359 ; 372 (sh) (MeOH).
- 10-Hydroxy: 10-Hydroxycampthoecin**
[19685-09-7]
C₂₀H₁₆N₂O₅ 364.357
Minor alkaloid from *Camptotheca acuminata* (Nyssaceae). Antineoplastic agent. Cryst. + 1H₂O. Mp 268-270°. Log P 0.65 (calc).
- UQ0491700
- 10-Hydroxy, di-Ac:** Mp 270-273°.
- 10-Hydroxy, 20-deoxy: 10-Hydroxy-deoxycampthoecin**
[123914-40-9]
C₂₀H₁₆N₂O₄ 348.357
Alkaloid from the seeds of *Camptotheca acuminata* (Nyssaceae). Shows strong cytotoxicity against P-388 leukaemia cells. Log P 1.06 (calc).
- 11-Hydroxy: 11-Hydroxycampthoecin**
[68426-53-9]
C₂₀H₁₆N₂O₅ 364.357
Alkaloid from the fruits of *Camptotheca acuminata* (Nyssaceae). Has antineoplastic props. Mp 327-330° dec. $[\alpha]_{\text{D}}^{21}$ -30.7 (c, 0.326 in Py). Log P 0.65 (calc). λ_{\max} 220 (ε 24600); 253 (ε 15800); 348 (ε 10450); 367 (ε 10700); 384 (ε 104450) (MeOH) (Berdy). λ_{\max} 208 ; 265 ; 334 ; 370 ; 384 (MeOH/NaOH) (Berdy).
- 18-Hydroxy: 18-Hydroxycampthoecin**
[116139-46-9]
C₂₀H₁₆N₂O₅ 364.357
Alkaloid from seeds of *Camptotheca acuminata* (Nyssaceae). Shows strong cytotoxicity against P-388 leukaemia cells *in vitro*. Yellow cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 256-258°. $[\alpha]_{\text{D}}^{11}$ -21.4 (c, 0.1 in Py). Log P -1.39 (calc). λ_{\max} 220 ; 250 ; 290 ; 320 (MeOH) (Berdy).
- 18-Hydroxy, 20-O-β-D-glucopyranoside: 20-O-β-D-Glucopyranosyl-18-hydroxycampthoecin**
C₂₆H₂₆N₂O₁₀ 526.499
Alkaloid from the root bark of *Camptotheca acuminata*. Amorph. powder.
- $[\alpha]_{\text{D}}^{25}$ -51.3 (c, 0.01 in MeOH). λ_{\max} 216 (log ε 4.56); 255 (log ε 4.38); 290 (log ε 4.01); 359 (log ε 4.21); 370 (log ε 4.19) (MeOH).
- 7-Methoxy: 7-Methoxycampthoecin**
[80735-03-1]
C₂₁H₁₈N₂O₅ 378.384
Alkaloid from leaves of *Ophiorrhiza filistipula* (Rubiaceae). Feathery needles (MeOH). Mp 260-270° dec. $[\alpha]_{\text{D}}^{25}$ +83 (c, 0.006 in CHCl₃).
- 9-Methoxy: 9-Methoxycampthoecin**
[39026-92-1]
C₂₁H₁₈N₂O₅ 378.384
Alkaloid from *Mappia foetida*, *Ervatamia heyneana*, *Nothapodytes foetida*, *Merrilliodendron megacarpum* and *Ophiorrhiza mungos* (Rubiaceae, Apocynaceae, Icacinaceae). Antineoplastic agent. Shows v. strong activity vs. P-388 cells. Light yellow cryst. (CHCl₃/MeOH). Mp 258-260° dec. $[\alpha]_{\text{D}}^{24}$ -77.5 (c, 1 in Py). Log P 0.87 (calc). λ_{\max} 219 (ε 33900); 256 (ε 24500); 265 (sh) (ε 21400); 305 (ε 7240); 320 (ε 11200); 358 (ε 20000); 371 (sh) (ε 19500) (MeOH) (Derep). λ_{\max} 218 (ε 33100); 262 (ε 24000); 305 (ε 7080); 320 (ε 11200); 356 (ε 20000); 371 (sh) (MeOH) (Derep).
- 9-Methoxy, Ac: 20-O-Acetyl-9-methoxycampthoecin**
[39026-93-2]
Alkaloid from the stems of *Nothapodytes foetida*. Pale yellow needles (Me₂CO). Mp 231-232°. $[\alpha]_{\text{D}}^{25}$ -24 (c, 1.15 in CHCl₃).
- 9-Methoxy, 10-hydroxy, 10-O-β-D-glucopyranoside: Chaboside**
[131669-96-0]
C₂₇H₂₈N₂O₁₁ 556.525
Alkaloid from *Ophiorrhiza pumila* (Rubiaceae). Pale yellow powder. λ_{\max} 222 ; 266 ; 293 ; 328 ; 376 (sh) (MeOH) (Derep).
- 10-Methoxy: 10-Methoxycampthoecin**
[19685-10-0]
C₂₁H₁₈N₂O₅ 378.384
Alkaloid from *Camptotheca acuminata* and *Ophiorrhiza mungos* (Nyssaceae, Rubiaceae). Inhibitor of herpes virus. Yellow cryst. (Me₂CO). Mp 255-256° dec. Log P 0.87 (calc). λ_{\max} 221 (ε 40300); 265 (ε 23800); 293 (ε 5200); 312 (ε 7200); 328 (ε 9200); 365 (ε 21600); 380 (ε 24400) (MeOH) (Berdy).
- 10-Methoxy, Ac: 20-O-Acetyl-10-methoxycampthoecin**
C₂₃H₂₀N₂O₆ 420.421
Alkaloid from the root bark of *Camptotheca acuminata*. Needles (CHCl₃). Mp 275-276°. $[\alpha]_{\text{D}}^{25}$ -36.5 (c, 0.01 in MeOH). λ_{\max} 218 (log ε 4.43); 262 (log ε 4.2); 330 (log ε 2.95); 370 (log ε 4.01); 380 (log ε 4.13) (MeOH).
- 10-Methoxy, 20-hexanoyl: 20-Hexanoyl-10-methoxycampthoecin**
[72594-34-4]
C₂₇H₂₈N₂O₆ 476.528
Alkaloid from the bark of *Camptotheca acuminata* (Nyssaceae).
- 9,10-Methylenedioxy: 9,10-Methylene-dioxycampthoecin**
[143005-94-1]
C₂₁H₁₆N₂O₆ 392.367
Alkaloid from *Ophiorrhiza trichocarpum*. Bright yellow powder. $[\alpha]_{\text{D}}^{25}$ +28.8 (c, 0.2 in MeOH). λ_{\max} 224 (log ε 4.36); 274 (log ε 4.3); 338 (log ε 4.24); 374 (log ε 4) (MeOH).
- (±)-form**
Synthetic. Cryst. (MeOH/CHCl₃ or EtOAc). Mp 276-278°.
- 20-Deoxy: 20-Deoxycampthoecin**
[34141-35-0]
C₂₀H₁₆N₂O₃ 332.358
Minor alkaloid from the bark of *Camptotheca acuminata* (Nyssaceae). Cryst. (CHCl₃/MeOH/Et₂O). Mp 258-263°.
- 10-Hydroxy: [64439-81-2]**
Synthetic. Cryst. + 1H₂O (CHCl₃/MeOH). Mp 327-330° (265-268°).
- 7-Methoxy: [68520-67-2]**
Synthetic. Cryst. (CHCl₃/MeOH). Mp 295° dec.
- 10-Methoxy: [64439-80-1]**
Synthetic. Needles (MeOH/CHCl₃). Mp 272-273°.
- Wall, M.E. *et al.*, *J.A.C.S.*, 1966, **88**, 3888-3890 (*isol, uv, ir, pmr, cryst struct*)
Wani, M.C. *et al.*, *J.O.C.*, 1969, **34**, 1364-1367 (*isol, uv, ir, pmr, struct, 10-Hydroxycampthoecin, 10-Methoxycampthoecin*)
Govindachari, T.R. *et al.*, *Phytochemistry*, 1972, **11**, 3529-3531 (*9-Methoxycampthoecin*)
Schultz, A.G. *et al.*, *Chem. Rev.*, 1973, **73**, 385-405 (*rev*)
Agarwal, J.S. *et al.*, *Indian J. Chem.*, 1973, **11**, 969 (*9-Methoxycampthoecin*)
Tafur, S. *et al.*, *J. Nat. Prod.*, 1976, **39**, 261-262 (*10-Methoxycampthoecin*)
Sheriha, G.M. *et al.*, *Phytochemistry*, 1976, **15**, 505-508 (*biosynth*)
Heckendorf, A.H. *et al.*, *Tet. Lett.*, 1977, 4153-4154 (*biosynth*)
Baxmann, E. *et al.*, *Chem. Ber.*, 1978, **111**, 3403-3411 (*7-Methoxycampthoecin*)
Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1979, **42**, 475-477 (*9-Methoxycampthoecin*)
Adamovics, J.A. *et al.*, *Phytochemistry*, 1979, **18**, 1085-1086 (*20-Deoxycampthoecin, 20-Hexanoylcampthoecin*)
Wani, M.C. *et al.*, *J. Med. Chem.*, 1980, **23**, 554-560 (*20-Deoxycampthoecin, 10-Hydroxycampthoecin, 10-Methoxycampthoecin, synth*)
Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1981, 1563-1568 (*10-Methoxycampthoecin, synth*)
Hutchinson, C.R. *et al.*, *Tetrahedron*, 1981, **37**, 1047-1065 (*rev*)
Martindale, *The Extra Pharmacopoeia, 28th 29th edn.*, Pharmaceutical Press, 1982, 12515
Cai, J.-C. *et al.*, *Alkaloids (Academic Press)*, 1983, **21**, 101-137 (*rev, pharmacol*)
Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 73-89 (*rev, antineoplastic activity*)
Wall, M.E. *et al.*, *J. Med. Chem.*, 1986, **29**, 1553-1555 (*11-Hydroxycampthoecin*)
Aiyama, R. *et al.*, *Phytochemistry*, 1988, **27**, 3663-3664 (*18,19-Dehydrocampthoecin*)
Lin, L.Z. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 186-188; *CA*, **109**, 107686z (*18-Hydroxycampthoecin*)

- Lin, L. *et al.*, *Huaxue Xuebao*, 1989, **47**, 506-508; *CA*, **111**, 228962m (10-Hydroxydeoxycamptothecin)
- Lin, L.-Z. *et al.*, *J. Nat. Prod.*, 1990, **53**, 186-189 (Camptothecin, 10-Hydroxycamptothecin, *cmr*)
- Aimi, N. *et al.*, *Tet. Lett.*, 1990, **31**, 5169-5172; 1992, **33**, 5403-5404 (Chaboside)
- Ezell, E.L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1645-1650 (*pmr*, *cmr*)
- Arbain, D. *et al.*, *Aust. J. Chem.*, 1993, **46**, 977-985 (7-Methoxycamptothecin)
- Wall, M.E. *et al.*, *Chron. Drug Discovery*, 1993, **3**, 327-348 (rev)
- Beijnen, J.H. *et al.*, *J. Chromatogr.*, 1993, **617**, 111-117 (*hplc*)
- Camptothecins*, (eds. Potmesil, M. *et al.*), CRC Press, 1995, (book)
- Wall, M.E. *et al.*, *Cancer Res.*, 1995, **55**, 753-760 (rev)
- Wu, T.-S. *et al.*, *Phytochemistry*, 1995, **39**, 383-385 (O-Acetylcamptothecin)
- Bodley, A.L. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1995, **92**, 3726-3730 (*props*)
- Murata, N. *et al.*, *Synlett*, 1997, 298-300 (*synth*)
- Cao, Z. *et al.*, *Synth. Commun.*, 1997, **27**, 2013-2019 (*synth*)
- Ciufolini, M.A. *et al.*, *Tetrahedron*, 1997, **53**, 11049-11060 (*synth*, *bibl*)
- Wall, M.E. *et al.*, *Alkaloids (London)*, 1998, **50**, 509-536 (rev)
- Kitajima, M. *et al.*, *J.C.S. Perkin 1*, 1998, 389-390 (9-Glucosyloxycamptothecin)
- Chavan, S.P. *et al.*, *Tet. Lett.*, 1998, **39**, 6745-6748 (*synth*)
- Dai, J.-R. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1427-1429 (20-glucoside)
- Tagami, K. *et al.*, *Heterocycles*, 2000, **53**, 771-775 (*synth*)
- Blagg, B.S.J. *et al.*, *Tetrahedron*, 2002, **58**, 6343-6349 (*synth*, *bibl*)
- Soepenbergh, O. *et al.*, *Alkaloids (Academic Press)*, 2003, **60**, 1-50 (rev, *biochem*)
- Srinivas, K.V.N.S. *et al.*, *Biochem. Syst. Ecol.*, 2003, **31**, 85-87 (20-O-Acetyl-9-methoxycamptothecin)
- Thomas, C.J. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 1585-1604 (rev, *synth*, *sar*)
- Oberlies, N.H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 129-135 (rev, *history*)
- Twin, H. *et al.*, *Org. Lett.*, 2004, **6**, 4913-4916 (*synth*)
- Lorence, A. *et al.*, *Phytochemistry*, 2004, **65**, 2735-2749 (rev)
- Zhang, Z. *et al.*, *Planta Med.*, 2004, **70**, 1216-1221 (20-O-Acetyl-10-methoxycamptothecin, 20-Glucosyl-18-hydroxycamptothecin)
- Chavan, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 3113-3115 (*synth*)
- Yu, J. *et al.*, *Tet. Lett.*, 2004, **45**, 7247-7250 (*synth*)
- Daud, A. *et al.*, *Clin. Cancer Res.*, 2005, **11**, 3009-3016 (*karenitecin*, *clin trial*)
- Puri, S.C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1717-1719 (*funggal metab*)
- Sriram, D. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 393-412 (rev)
- Anderson, R.J. *et al.*, *Org. Lett.*, 2005, **7**, 2989-2991 (*synth*)
- Peters, R. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 498-509 (*synth*)
- Tang, C.-J. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3757-3759 (*synth*)
- Klausmeyer, P. *et al.*, *Planta Med.*, 2007, **73**, 49-52 (9,10-Methylenedioxcamptothecin)
- Zamboni, W.C. *et al.*, *Invest. New Drugs*, 2008, **26**, 399-406 (DB67, *pharmacol*)
- Verma, R.P. *et al.*, *Chem. Rev.*, 2009, **109**, (rev, *sar*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 11th edn., J. Wiley, 2004, CBB870; CBB875

Camptothecin acid C-71

α -Ethyl-9,11-dihydro- α -hydroxy-8-hydroxymethyl-9-oxoindolizino[1,2-b]quinoline-7-acetic acid, 9CI [34079-22-6]



C₂₀H₁₈N₂O₅ 366.373

(S)-form

17-Ac: 17-O-Acetylcamptothecin acid

[199581-40-3]

C₂₂H₂₀N₂O₆ 408.41

Alkaloid from *Mappia foetida*.

9-Methoxy, 17-Ac: 17-O-Acetyl-9-methoxycamptothecin acid

[199581-41-4]


C₂₃H₂₂N₂O₇ 438.436

Alkaloid from *Mappia foetida*.

Pat. Coop. Treaty (WIPO), 1997, 97 43 290; CA, **128**, 34922d

Canadine

C-72



(S)-form

C₂₁H₂₃NO₅ 369.416

Protoberberine-type numbering shown. Abs. config. of nat. alkaloid revised to (S-) in 1989.

(R)-form [124649-83-8]

Synthetic. Mp 114-116°. [α]_D -38 (c, 0.1 in CHCl₃).

(S)-form [52801-27-1]

Alkaloid from *Hydrastis canadensis* (Papaveraceae). Cryst. (Et₂O). Mp 117-118°. [α]_D +43 (c, 0.5 in CHCl₃). λ _{max} 288 (ε 78000) (MeOH).

8-Alcohol, N-de-Me: *Argenaxine*

[88114-59-4]

C₂₀H₂₃NO₅ 357.405

Alkaloid from the aerial parts of *Argemone mexicana*. Prisms. Mp 157-158°. [α]_D²⁴ +95 (c, 0.34 in CHCl₃). λ _{max} 225 (log ε 4.14); 285 (log ε 3.38) (MeOH).

8-Carboxylic acid: *Canadinic acid*

[190720-13-9]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Hydrastis canadensis* (Papaveraceae). Cryst. (EtOAc). Mp 130-132° dec. [α]_D²⁰ -112 (c, 0.1 in MeOH). λ _{max} 235 (sh) (log ε 4.14); 289 (log ε 3.81) (EtOH).

13-Oxo, 8-carboxylic acid: *Berbervirine*

[167817-66-5]

C₂₁H₂₁NO₇ 399.399

Alkaloid from whole plants of *Berberis virgetorum* (Berberidaceae). Light pink needles (CHCl₃/MeOH). Mp 202-204°. [α]_D -200 (c, 1.625 in CHCl₃). Has R-config. λ _{max} 226 (log ε 4.39); 262 (log ε 3.82); 288 (log ε 3.9); 331 (log ε 3.35) (MeOH). λ _{max} 230 (log ε 4.21); 250 (log ε 4.36); 295 (log ε 1.4); 333 (log ε 1.03) (MeOH).

(±)-form [62475-57-4]

Synthetic. Prisms (MeOH). Mp 143-143.5° (139-140°). λ _{max} 226 (log ε 4.39); 262 (log ε 3.82); 288 (log ε 3.9); 331 (log ε 3.35) (MeOH).

Gleye, J. *et al.*, *Phytochemistry*, 1974, **13**, 675-676 (*pmr*, *uv*, *ms*, *struct*)

Shamma, M. *et al.*, *Tetrahedron*, 1978, **34**, 635-640 (*config*, *synth*, *ir*, *uv*)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2685-2690 (*synth*, *uv*, *ir*, *pmr*)

Hanaoka, M. *et al.*, *Heterocycles*, 1989, **29**, 221-224 (*abs config*, *synth*)

Liu, C. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1100-1102 (*Berbervirine*)


Galeffi, C. *et al.*, *Planta Med.*, 1997, **63**, 194 (*Canadinic acid*)

Chang, Y.-C. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 521-526 (*Argenaxine*)

Canadine

C-73

5,8,13,13a-Tetrahydro-9,10-dimethoxy-6H-benzo[g]-1,3-benzodioxolo[5,6-a]quinolizine, 9CI. 9,10-Dimethoxy-2,3-(methylenedioxy)berberine, 8CI. Tetrahydroberberine [522-97-4]



(R)-form

C₂₀H₂₁NO₄ 339.39

Shows analgesic, hypotensive and papaverine-like spasmolytic activity. Potential of antimitotic effect of colchicine.

(R)-form

α -Canadine

[5096-57-1]

Alkaloid from *Corydalis tuberosa* (Papaveraceae). Mp 132°. [α]_D¹⁵ +299 (CHCl₃).

▶ LD₅₀ (mus, orl) 940 mg/kg. DR9823000

O¹⁰-De-Me, N-Me: *Escholidine*

[30389-09-4]

C₂₀H₂₂NO₄ 340.398

Principal quaternary alkaloid from the roots of *Eschscholtzia californica*, *Eschscholtzia douglasii* and *Eschscholtzia glauca*, also obt. from aerial parts of *Hunnemannia fumariaefolia* (Papaveraceae). Needles or prisms (MeOH) (as perchlorate). Mp 281-282° (perchlorate). [α]_D²² -155 (c, 0.27 in MeOH). Struct. revised in 2006.

(S)-form

β -Canadine

[2086-96-6]

Alkaloid from *Hydrastis canadensis* and several *Corydalis* spp. (Papaveraceae). Needles. Mp 134°. [α]_D²⁰ -299 (CHCl₃).

N-Me: N-Methyl- β -canadine

C₂₁H₂₄NO₄⁺ 354.425

Quaternary alkaloid from *Zanthoxylum brachyacanthum*, *Zanthoxylum veneficum* and *Fagara rhoifolia* (Rutaceae).

N-Me, chloride:

Needles (MeOH) or plates. Mp 235-237° dec Mp 240-242° (dimorph.). [α]_D²² -134.2 (c, 0.23 in H₂O). Epimers at N referred to as Canadine α - and β -methochlorides (confusing) may have been characterised.

N-Me, iodide:

Cryst. (EtOH). Mp 216-218° (resolidifies and melts at 246° dec.). [α]_D²² -115 (c, 0.3 in H₂O).

O⁹-De-Me: Nandinine. Tetrahydroberberubine

[572-76-9]

[6724-39-6]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the bark of *Nandina domestica* (Nandinaceae). Mp 195-196°. [α]_D +303 (CHCl₃). [α]_D +298 (EtOH).

O¹⁰-De-Me: Tetrahydrothalifendine

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Thalictrum fendleri* (Ranunculaceae). Mp 209-211°. [α]_D -175 (MeOH).

8-Oxo: 12,13-Dihydro-8-oxoberberine. 8-Oxotetrahydroberberine. 8-Oxocanadine. Gusanlung B

[79082-05-6]

C₂₀H₁₉NO₅ 353.374

Minor alkaloid from stems of *Cosciniun fenestratum*, also isol. from stems of *Arcangelisia gusanlung* (Menispermaceae). Yellow needles. Mp 224-226° (211-213°, 218°). [α]_D -348 (c, 0.54 in CHCl₃). The name 12,13-Dihydro-8-oxoberberine is confusing. Pmr and cmr spectra reported by Malhotra *et al* were misassigned or misinterpreted. Genus name erroneously given as Acangelisia.

8-Oxo, O⁹-de-Me: Gusanlung A. 8-Oxo-berberubine

[139220-06-7]

C₁₉H₁₇NO₅ 339.347

Alkaloid from *Berberis heteropoda* and stems of *Arcangelisia gusanlung* (Menispermaceae). Cryst. (Ac₂O). Mp 260-262° (245°-246°). [α]_D²¹ -395.5 (c, 0.27 in CHCl₃).

8-Oxo, O¹⁰-de-Me: 8-Oxotetrahydrothalifendine

[142808-33-1]

C₁₉H₁₇NO₅ 339.347

Alkaloid from stems of *Cosciniun fenestratum* (Menispermaceae). [α]_D -605 (c, 0.084 in MeOH).

(\pm)-form [29074-38-2]

Mp 169-170°.

► EX2675000

O⁹-De-Me: [17388-17-9]

Mp 186°.

8-Oxo:

Pale yellow cryst. Mp 217-218°.

Späth, E. *et al.*, *Ber.*, 1930, **63**, 3007; 1931, **64**, 1131 (*Canadine, Nandinine, isol. struct, resoln, bibl*)

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1942, **20**, 57 (*isol*)

Corrodi, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 889 (*Nandinine, abs config*)

Kuchkova, K.I. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 141; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 148 (*methochlorides*)

Giacoppello, D. *et al.*, *Tetrahedron*, 1967, **23**, 3265 (*Nandinine, synth*)

Kametani, T. *et al.*, *J.C.S. (C)*, 1969, 2036 (*synth, ir, pmr*)

Shamma, M. *et al.*, *Tetrahedron*, 1971, **27**, 727 (*Tetrahydrothalifendine*)

Kametani, T. *et al.*, *Heterocycles*, 1975, **3**, 811 (*Nandinine, synth*)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (*cmr*)

Narasimhan, N.S. *et al.*, *Tetrahedron*, 1983, **39**, 1975 (*synth*)

Bhakuni, D.S. *et al.*, *Alkaloids (Academic Press)*, 1986, **28**, 95 (*rev*)

Pyne, S.G. *et al.*, *Tet. Lett.*, 1987, **28**, 4737 (*synth*)

Hussain, R.A. *et al.*, *Heterocycles*, 1989, **29**, 2257 (*pmr, cmr*)

Malhotra, S. *et al.*, *Phytochemistry*, 1989, **28**, 1998 (*8-Oxocanadine*)

Zhang, J.-S. *et al.*, *Planta Med.*, 1991, **57**, 457 (*Gusanlungs*)

Kessar, S.V. *et al.*, *J.O.C.*, 1992, **57**, 6716 (*synth*)

Pinho, P.M.M. *et al.*, *Phytochemistry*, 1992, **31**, 1403 (*8-Oxocanadine, 8-Oxotetrahydrothalifendine*)

Yusupov, M.M. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 53; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 43 (*8-Oxoberberubine*)

Ningirawath, S. *et al.*, *Aust. J. Chem.*, 1994, **47**, 951 (*synth, Gusanlung B*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Wang, X.-L. *et al.*, *Acta Cryst. E*, 2006, **62**, 1764-1765 (*8-Oxocanadine, cryst struct*)


Chudik, S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 954-956 (*Escholidine*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TCJ800

Canavanine

C-74

O-[(Aminoiminomethyl)amino]homoserine, 9CI. 2-Amino-4-(guanidinoxy)butanoic acid



(S)-form

C₅H₁₂N₄O₃ 176.175

(S)-form

L-form

[543-38-4]

Stored in large quantities in the seeds of leguminous plants in three subfamilies.

Isol. originally from Jackbean (*Canavalia ensiformis*). Insecticidal antimetabolite props. Antineoplastic, antibacterial and antifungal agent. Cryst. (EtOH). Sol.

H₂O. Mp 184°. [α]_D²⁰ +7.9 (H₂O).

► LD₅₀ (rat, scu) 5900 mg/kg. ES7002000

Sulfate: [2219-31-0]

Cryst. (EtOH). Mp 172° dec.

Picrate: Mp 163-164°.

Tribenzoyl: Mp 86° dec.

(\pm)-form [13269-28-8]

Mp 180-182°. pK_{a1} 2.5; pK_{a2} 6.6; pK_{a3} 9.25 (25°).

Hydrochloride: Mp 190°.

Hydrochloride (1:2):

Very hygroscopic solid.

Bell, A. *et al.*, *Biochem. J.*, 1960, **75**, 618

(*occur*)

Nakatsu, S. *et al.*, *CA*, 1963, **58**, 752 (*occur*)

Yamada, Y. *et al.*, *Agric. Biol. Chem.*, 1973, **37**, 2201 (*synth*)

Rosenthal, G.A. *et al.*, *Anal. Biochem.*, 1976, **77**, 147 (*isol*)

Rosenthal, G.A. *et al.*, *Q. Rev. Biol.*, 1977, **52**, 155 (*rev*)

Boyar, A. *et al.*, *J.A.C.S.*, 1982, **104**, 1995

(*cryst struct*)

Rosenthal, G.A. *et al.*, *Phytochemistry*, 1991, **30**, 1055 (*biol props*)


Martindale. *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 12026

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AKD500

Canavanosuccinic acid

C-75

N-[[[(3-Amino-3-carboxypropoxy)amino]imino]methyl]aspartic acid, 9CI. Canavaninosuccinic acid [24764-65-6]



C₉H₁₆N₄O₇ 292.248

An intermed. in the canaline-urea cycle in plants. Purified as Ba²⁺ salt, not isol.

Walker, J.B. *et al.*, *Arch. Biochem. Biophys.*, 1955, **59**, 233


Rosenthal, G.A. *et al.*, *Anal. Biochem.*, 1975, **65**, 60

Canconrine, 9CI

C-76

Alkaloid F22

[29477-90-5]



C₃₆H₃₄N₂O₇ 606.674

Alkaloid from *Dicentra canadensis* (Papaveraceae). Orange needles (CHCl₃/MeOH). Mp 237-238°.

Hydrochloride:Cryst. (CHCl₃/MeOH). Mp 286°.**Me ether:**

Cryst. (MeOH/cyclohexane). Mp 269°.

8,14-Didehydro: Dehydrocancetrine A

[38990-01-1]

C₃₆H₃₂N₂O₇ 604.658Minor alkaloid from *Dicentra canadensis* (Papaveraceae). Yellow cryst. (MeOH). Mp 194°.**31,32-Didehydro: Dehydrocancetrine B**

[38990-02-2]


C₃₆H₃₂N₂O₇ 604.658Minor alkaloid from *Dicentra canadensis* (Papaveraceae). Red cryst. (MeOH). Mp 206°.**10-Oxo: 10-Oxocancetrine**

[68798-36-7]

C₃₆H₃₂N₂O₈ 620.657Alkaloid from *Dicentra canadensis* (Papaveraceae). λ_{max} 268 (log ε 3.35); 284 (sh) (log ε 3.34); 324 (sh) (log ε 2.9); 436 (log ε 2.65) (EtOH).Manske, R.H.F. *et al.*, *Can. J. Res.*, 1932, **7**, 258; 1938, **16B**, 81 (*isol*)Clark, G.R. *et al.*, *J.A.C.S.*, 1970, **92**, 4998 (*struct*)Rodrigo, R. *et al.*, *Can. J. Chem.*, 1972, **50**, 853; 3900 (*uv, ir, pmr, ms, struct*)MacLean, D.B. *et al.*, *Can. J. Chem.*, 1972, **50**, 862 (*Dehydrocancetrines*)Clark, G.R. *et al.*, *J.C.S. Perkin 2*, 1972, 1219 (*cryst struct, abs config*)Rodrigo, R. *et al.*, *Alkaloids (Academic Press)*, 1973, **14**, 407 (*rev*)Holland, H.L. *et al.*, *Can. J. Chem.*, 1978, **56**, 2467-2471 (*cmr*)**Candidine†**


C-77

6-(1,3-Dihydro-3-oxo-2H-indol-2-ylidene)indolo[2,1-b]quinazolin-12(6H)-one, 9CI. Indigo X. Qingdainone [97457-31-3]

C₂₃H₁₃N₃O₂ 363.375Isol. from *Candida lipolytica*, *Isatis indigotica* and *Phaius mishmensis*. Also isol. in small amts. from the urine of uraemic patients. Violet solid or dark black-brown needles. Mp 269-270°. λ_{max} 250 (log ε 4.16); 284 (log ε 4); 574 (log ε 3.88) (CHCl₃).Bergman, J. *et al.*, *Tetrahedron*, 1985, **41**, 2883-2884 (*struct*)Laatsch, H. *et al.*, *Annalen*, 1986, 1847-1853 (*isol, pmr*)Bergman, J. *et al.*, *Phytochemistry*, 1989, **28**, 3547Jao, C.-W. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1275-1279 (*isol, pmr, cmr*)**Candimine**

C-78

[24585-19-1]

C₁₈H₁₉NO₆ 345.351Alkaloid from *Hippeastrum candidum* (Amaryllidaceae). Cryst. (MeOH). Mp 218-220°. [α]_D²⁴ +220 (c, 0.2 in CHCl₃).**Perchlorate:**Cryst. (H₂O). Mp 177-179°.**Picrate:**

Cryst. (MeOH). Mp 220°.

Ac: Mp 239-241°.

Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 920; *CA*, **58**, 11416a**Caneine**

C-79


C₁₂H₂₄N₂O₃ 244.333Struct. unknown. Constit. of *Canavalia ensiformis* seeds (jack bean) (Fabaceae). Lustrous needles. Mp 188-189°. Opt. inactive.**Picrate:**

Pale yellow needles. Mp 120-121°.

Ackermann, D. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1939, **262**, 103 (*isol*)**Canelillinoxine**

C-80

[151757-05-0]

C₁₇H₁₅NO₄ 297.31Alkaloid from the stem bark of *Aniba canelilla* (Lauraceae).Oger, J.-M. *et al.*, *Can. J. Chem.*, 1993, **71**, 1128 (*isol, uv, ir, pmr, ms, struct*)**Canembine**


C-81

Raunescine†

[1355-10-8]


C₂₂H₂₈N₂O₃ 368.475Struct. unknown. Not the same as Raunescine in 17,18-Dihydroxyyohimban-16-carboxylic acid, D-687. Said to be a Yohimbine homologue. Alkaloid from *Rauwolfia canescens* (Apocynaceae). Clusters of pale-yellow needles (EtOAc).Bhattacharji, S. *et al.*, *Indian J. Pharm.*, 1956, **18**, 188-190; *CA*, **51**, 446a (*isol, ir, uv*)Bhattacharji, S. *et al.*, *J. Sci. Ind. Res.*, 1957, **16**, 97; *CA*, **53**, 8185g (*nomenclature*)**Cannabiloid B**

C-82

C₁₆H₂₁NO₁₀ 387.343Constit. of *Senecio cannabifolius*. Liq. [α]_D²⁰ +101.2 (c, 0.12 in MeOH). λ_{max} 210 (MeOH).Wu, B. *et al.*, *Pharm. Biol.*, 2006, **44**, 440-444 (*isol, pmr, cmr*)**Cannabisativine**

C-83

13-(1,2-Dihydroxyheptyl)-1,4,5,6,7,8,9,10,11,13,16,16a-dodecahydroprido[2,1-d][1,5,9]triazacyclotridecin-2(3H)-one, 9CI [57682-64-1]



Absolute Configuration


C₂₁H₃₉N₃O₃ 381.557Alkaloid from the leaves and roots of *Cannabis sativa* (Cannabidaceae). Plates (Me₂CO). Mp 167-168°. [α]_D²⁵ +55.1 (c, 0.53 in CHCl₃).**1'-Deoxy, 2'-ketone: Anhydrocannabisativine**

[65664-79-1]

C₂₁H₃₇N₃O₂ 363.542Alkaloid from the leaves and roots of *Cannabis sativa* (Cannabidaceae). [α]_D²² +18.7 (c, 0.1 in MeOH).Lotter, H.L. *et al.*, *Tet. Lett.*, 1975, 2815(*Cannabisativine, ir, pmr, ms, cryst struct*)Turner, C.E. *et al.*, *J. Pharm. Sci.*, 1976, **65**,1084 (*Cannabisativine, isol, uv, ir, pmr, ms*)Elsobly, M.A. *et al.*, *J. Pharm. Sci.*, 1978, **67**,124 (*Anhydrocannabisativine, struct*)Bailey, T.R. *et al.*, *J.A.C.S.*, 1984, **106**, 3240(*Anhydrocannabisativine, synth*)Ogawa, M. *et al.*, *Tet. Lett.*, 1984, **25**, 969(*Cannabisativine, synth, cmr*)Wasserman, H.H. *et al.*, *Tetrahedron*, 1988, **44**,3365 (*Cannabisativine, synth, ir, pmr, ms*)Hamada, T. *et al.*, *Tet. Lett.*, 1991, **32**, 1649(*Cannabisativine, synth*)Kuethe, J.T. *et al.*, *J.O.C.*, 2004, **69**, 5219-5231(*Cannabisativine, synth*)

Cannabisin A

C-84

C₃₄H₃₀N₂O₈ 594.62

Constit. of the fruit of *Cannabis sativa*.
Amorph. powder. λ_{\max} 257 (log ϵ 4.79)
(EtOH).


Hexa-Ac:

Needles. Mp 204-207°.

Sakakibara, I. *et al.*, *Phytochemistry*, 1991, **30**,
3013-3016 (*isol*, *pmr*, *cmr*)

Cannabisin B

C-85

C₃₄H₃₂N₂O₈ 596.635**(±)-form** [144506-17-2]

Isol. from fruits of *Cannabis sativa*
(Cannabaceae). Amorph. powder.

6-Me ether: Cannabisin C

[144506-18-3]

C₃₅H₃₄N₂O₈ 610.662

From fruits of *Cannabis sativa* (Can-
nabaceae). Amorph. powder.

3',6-Di-Me ether: Cannabisin D

[144506-19-4]

C₃₆H₃₆N₂O₈ 624.689

From fruits of *Cannabis sativa* (Can-
nabaceae). Needles (MeOH). Mp 165-
168°.

Hexa-Me ether:


Needles (MeOH). Mp 196-197°.

Sakakibara, I. *et al.*, *Phytochemistry*, 1992, **31**,
3219 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Cannabisin E

C-86

[163136-16-1]

C₃₆H₃₈N₂O₉ 642.704

Isol. from fruits of *Cannabis sativa*
(Cannabaceae). Amorph. powder.

Tetra-Ac:

Needles. Mp 123-125°.

7-Deoxy, 7,8-didehydro: Cannabisin F

[163136-19-4]

C₃₆H₃₆N₂O₈ 624.689


From fruits of *Cannabis sativa* (Can-
nabaceae). Amorph. powder.

Sakakibara, I. *et al.*, *Phytochemistry*, 1995, **38**,
1003 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Cannabisin G

C-87

[163136-20-7]

C₃₆H₃₆N₂O₈ 624.689

Dimer of *N*-Feruloyltyramine in H-667.
Isol. from fruits of *Cannabis sativa*
(Cannabaceae) and *Hyoscyamus niger*.
Amorph. powder.

(Z,Z)-Isomer: HyoscyamideC₃₆H₃₆N₂O₈ 624.689

Alkaloid from the seeds of *Hyoscyamus*
niger. Yellow amorph. powder.


Sakakibara, I. *et al.*, *Phytochemistry*, 1995, **38**,
1003 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Ma, C.-Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 206-
209 (*Hyoscyamide*)

Cannabisin H

C-88

[403647-08-5]

C₂₈H₃₁NO₈ 509.555


Alkaloid from the bark of *Hibiscus*
cannabinus. Yellowish oil.

Seca, A.M.L. *et al.*, *Phytochemistry*, 2001, **58**,
1219-1223 (*isol*, *pmr*, *cmr*)

Cantharimide

C-89

Hexahydro-3a,7a-dimethyl-4,7-epoxy-2H-
isoindole-1,3-dione
[76970-77-9]

C₁₀H₁₃NO₃ 195.218

Alkaloid from the Chinese blister beetle
Mylabris phalerata. Needles (EtOH). Mp
205-206°.

N-(3-Amino-1S-carboxypropyl): N-(N²-
L-Ornithyl) cantharimide
[765922-94-9]

C₁₅H₂₂N₂O₅ 310.349

Alkaloid from *Mylabris phalerata*.

Powder. Mp 157-160°. [α]_D -26.9 (c,
0.26 in MeOH aq.).

N-(4-Amino-1S-carboxybutyl): N-(N²-
L-Lysyl) cantharimide

[765922-93-8]

C₁₆H₂₄N₂O₅ 324.376

Alkaloid from *Mylabris phalerata*.

Powder. Mp 159-162°. [α]_D -23.3 (c, 0.3
in MeOH).

N-(3-Guanidino-1S-carboxypropyl): N-
(N²-L-Arginyl) cantharimide

[765922-95-0]

C₁₆H₂₄N₄O₅ 352.389

Alkaloid from *Mylabris phalerata*.

Powder. Mp 195-197°. [α]_D -21.1 (c, 2.2
in MeOH aq.).

8-Hydroxy: 8-Hydroxycanthalimide

[936025-60-4]

C₁₀H₁₃NO₄ 211.217

Alkaloid from *Mylabris phalerata*.


Powder. Mp 195-197°.

Nakatani, T. *et al.*, *Chem. Pharm. Bull.*, 2004,
52, 807-809; 2007, **55**, 92-94 (*isol*, *pmr*, *cmr*)

Canthin-6-one

C-90

6H-Indolo[3,2,1-de][1,5]naphthyridin-6-
one, 9CI
[479-43-6]

C₁₄H₈N₂O 220.23

Alkaloid from *Pentaceras australis*, *Phel-*
lodendron amurense, *Zieria smithii*, *Ai-*
lantus altissima, *Ailanthus excelsa*, some
Fagara and *Zanthoxylum* spp., *Quassia*
gabonensis, *Xanthoxylum elephantiasis*,
Hibiscus syriacus, *Zanthoxylum* sp., *Pi-*
crasma excelsa, *Brucea antidysenterica*,
Hannoa klaineana and others. Also isol.
from the mushroom *Boletus curtisii*.
Shows antimicrobial activity. Cytotoxic
to guinea pig keratinocytes. Exhibits
fungicidal effects against a wide range of
fungi. Antiinflammatory. Pale yellow
needles (Me₂CO). Mp 162.5-163.5°. λ_{\max}
250 (ϵ 13200); 259 (ϵ 12500); 267 (ϵ
11700); 300 (ϵ 8400); 347 (ϵ 8700); 362 (ϵ
14100); 381 (ϵ 11750) (EtOH) (Berdy).

Hydrochloride:

Pale yellow plates or needles (1M
HCl). Mp 244-246° dec.

Picrate:

Yellow needles (MeOH). Mp 262-264°
(softening >240°).

Methodide:

Orange-red plates (H₂O). Mp 271-273°.

3-N-Oxide: Canthin-6-one N-oxide

[60755-87-5]

C₁₄H₈N₂O₂ 236.229

Alkaloid from the wood and root bark
of *Ailanthus altissima*. Shows cytotoxic
activity against guinea pig keratino-
cytes. Pale yellow needles (MeOH).
Sol. MeOH; poorly sol. H₂O. Mp 244-
245°. λ_{\max} 246; 278; 366 (MeOH)
(Berdy). λ_{\max} 246 (ϵ 11700); 279 (ϵ
24000); 370 (ϵ 13500) (EtOH) (Berdy).

4,5-Dihydro-4,5-Dihydrocanthin-6-one


C₁₄H₁₀N₂O 222.246
Alkaloid from cell cultures of *Ailanthus altissima*. Pale yellow needles (CH₂Cl₂/petrol). Mp 128°.

Haynes, H.F. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 387 (*isol, uv, struct, oxide*)
Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1956, **6**, 86 (*isol*)
Rosenkranz, H.J. *et al.*, *Annalen*, 1966, **691**, 159 (*synth, uv, ir*)
Mitscher, L.A. *et al.*, *Heterocycles*, 1975, **3**, 7 (*synth*)
Oehl, R. *et al.*, *Chem. Ber.*, 1976, **109**, 705 (*synth, uv, ir, ms*)
Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1532; 1981, **29**, 390 (*isol, uv, ir, pmr, ms, synth, oxide*)
Campos, O. *et al.*, *Heterocycles*, 1980, **14**, 975 (*synth*)
Forgacs, P. *et al.*, *Planta Med.*, 1982, **46**, 187 (*isol*)
Anderson, L.A. *et al.*, *J. Nat. Prod.*, 1983, **46**, 374 (*isol, biochem*)
Cain, M. *et al.*, *J.A.C.S.*, 1983, **105**, 907 (*synth*)
Ohmoto, T. *et al.*, *Alkaloids (Academic Press)*, 1989, **36**, 135-170 (*rev*)
He, W. *et al.*, *Meded. Fac. Landbouwwet., Rijksuniv. Gent*, 1998, **64**, 565-569 (*activity*)
Rossler, U. *et al.*, *Tet. Lett.*, 1999, **40**, 7075-7078 (*synth*)
Czerwinski, K.M. *et al.*, *Synth. Commun.*, 2003, **33**, 1225-1231 (*synth, pmr, cmr*)
Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (*isol, pmr, cmr, ms*)
Soriano-Agatón, F. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1581-1587 (*synth, pmr, cmr*)
Suzuki, H. *et al.*, *Synthesis*, 2005, 28-31 (*synth, ir, pmr, cmr*)

Canthiumine

C-91

α -(Dimethylamino)-N-(1,2,11,12,14,15,16,16a-octahydro-1,12-dioxo-10-phenyl-5,8-etheno-10H-pyrrolo[1,2-e][1,5,8]oxadiazacyclotetradecin-11-yl)benzenepropanamide, 9CI
[26195-90-4]



Absolute Configuration


C₃₃H₃₆N₄O₄ 552.672
Alkaloid from *Canthium euryoides* (Rubiaceae). Mp 232-233°. [α]_D -254 (c, 1 in CHCl₃).

Boulvin, G. *et al.*, *Bull. Soc. Chim. Belg.*, 1969, **78**, 583; *CA*, **72**, 90691q
González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915 (*stereochem*)

Cantleyne

C-92

[30333-81-4]



Absolute Configuration

C₁₁H₁₃NO₃ 207.229
Alkaloid from *Cantleya corniculata*, a *Jasminum* sp. *Dipsacus azureus*, *Strychnos nux-vomica*, *Strychnos dinklagei*, several other *Strychnos* spp. *Alstonia angustiloba*, *Alstonia pneumatophora*, *Alstonia spatulata*, *Scaevola racemigera* (Icacinaeae, Oleaceae, Dipsacaceae, Loganiaceae, Apocynaceae, Goodeniaceae). Needles (C₆H₆). Mp 132-133° (130°). [α]_D -34 (c, 1.2 in CHCl₃). [α]_D²⁰ -40 (c, 1 in CHCl₃).

6-Epimer Isocantleyne

[143564-04-9]
C₁₁H₁₃NO₃ 207.229
Alkaloid from *Siphonostegia chinensis* (Scrophulariaceae). Cryst. Mp 124-125°.

8-(E-4-Hydroxycinnamoyloxy): 8-Hydroxycantleyne trans-p-coumarate

[119365-64-9]
C₂₀H₁₉NO₆ 369.373
Alkaloid from *Coelospermum billardieri*. Mp 180°. [α]_D²³ -121 (c, 0.9 in EtOH). Not named in the paper. λ_{max} 207 ; 224 ; 314 (EtOH).


8-(Z-4-Hydroxycinnamoyloxy): 8-Hydroxycantleyne cis-p-coumarate

[119308-96-2]
C₂₀H₁₉NO₆ 369.373
Alkaloid from *Coelospermum billardieri*. Isol. as the diacetate. Not named in the paper.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1283 (*isol, uv, ir, pmr, ms, struct*)
Sévenet, T. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 3120 (*isol, uv, ir, pmr, ms, struct, synth*)
Rakhmatullaev, T.U. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 400; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 401 (*isol, uv, ir, pmr*)
Bisset, N.G. *et al.*, *Phytochemistry*, 1974, **13**, 265 (*isol*)
Skaltsounis, A.-L. *et al.*, *Helv. Chim. Acta*, 1985, **68**, 1679 (*isol*)
Michel, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 86 (*isol*)
Ravao, T. *et al.*, *Tet. Lett.*, 1985, **26**, 837 (*cmr*)
Lopez, J.L. *et al.*, *J. Nat. Prod.*, 1988, **51**, 829 (*8-Hydroxycantleyne coumarates*)
Zhang, H.Y. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 113; *CA*, **117**, 147160v (*Isocantleyne*)
Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, Strychnos*)

Capaurine

C-93



(S)-form

C₂₁H₂₅NO₅ 371.432

(S)-form [478-14-8]
Alkaloid from *Corydalis* spp. (Papaveraceae). Stout polyhedra (MeOH). Mp 164°. [α]_D²⁴ -271 (CHCl₃).

►EX8100000

Hydrobromide:

Plates (MeOH). Mp 198-199°. [α]_D -423.

N-Oxide: **Nokoensine**

[59870-71-2]
C₂₁H₂₅NO₆ 387.432
Alkaloid from *Corydalis nokoensis* (Papaveraceae). Mp 212°. [α]_D²⁴ -180 (c, 0.58 in MeOH).

N-Me(cis-): 9-O-Methylstecepharine.

cis-N-Methylcapaurine
[179238-68-7]
C₂₂H₂₈NO₅⁺ 386.467
Quaternary alkaloid from the dried rhizomes of *Stephania cepharantha*. Amorph. powder(as perchlorate). [α]_D²⁹ -156 (c, 0.96 in MeOH) (perchlorate). λ_{max} 233 (sh) (log ε 4.24); 280 (log ε 3.48) (MeOH).

Me ether: 1,2,3,9,10-Pentamethoxytetrahydroprotoberberine.

O-Methylcapaurine. 1-Methoxytetrahydropalmatine
[6033-73-4]
C₂₂H₂₇NO₅ 385.459
Alkaloid from roots of *Stephania lincangensis* (Menispermaceae). Prisms (MeOH). Mp 152°.

O¹⁰-De-Me: Capaurimine

[30758-81-7]
C₂₀H₂₃NO₅ 357.405
Alkaloid from *Corydalis montana* and *Corydalis pallida* (Papaveraceae). Stout prisms (CHCl₃/MeOH). Mp 212°. [α]_D²⁴ -287 (c, 0.4 in CHCl₃).

O⁹-De-Me, N-Me, (cis-): Stecepharine

[267006-16-6]
C₂₁H₂₆NO₅⁺ 372.44
Quaternary alkaloid from the dried rhizomes of *Stephania cepharantha*. Amorph. powder (as perchlorate). [α]_D³⁰ -175 (c, 1.05 in MeOH). λ_{max} 230 (sh) (log ε 4.23); 282 (log ε 3.64) (MeOH).

(±)-form

Capawidine

[478-15-9]
Alkaloid from *Corydalis* spp. (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 208°.
►DS0175000

Me ether:

Needles (MeOH). Mp 208°.

O¹⁰-De-Me: [32886-78-5]

Needles (MeOH). Mp 208-210°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1942, **20**, 49 (*isol*)

Kametani, T. *et al.*, *Tet. Lett.*, 1968, 4251 (*cryst struct*)

Kametani, T. *et al.*, *J. Het. Chem.*, 1970, **7**, 491 (*synth*)

Kametani, T. *et al.*, *J.C.S.(C)*, 1971, 2396 (*Capaurimine, synth, cryst struct*)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 3185 (*pmr*)

Tani, C. *et al.*, *Yakugaku Zasshi*, 1976, **96**, 527 (*Nokoensine*)

Taira, Z. *et al.*, *Cryst. Struct. Commun.*, 1977, **6**, 755 (*Nokoensine, cryst struct*)

Sugiura, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1168 (*cmr*)


Chen, Y. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1991, **33**, 552; *CA*, **117**, 86637g (*Pentamethoxytetrahydroprotoberberine*)

Tanashi, T. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 370-373 (*Stecepharine, 9-O-Methylstecepharine*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, TDI750

Capitavine

5,7-Dihydroxy-6-(1-methylpiperidin-2-yl)flavone
[91147-11-4]



C₂₁H₂₁NO₄ 351.401

Alkaloid from the seeds of *Buchenavia capitata* (Combretaceae). Cryst. (CH₂Cl₂/MeOH). Mp 146°. [α]_D +6 (c, 0.6 in EtOH).

N-De-Me: N-Demethylcapitavine

[91147-12-5]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the fruits of *Buchenavia macrophylla* (Combretaceae). Amorph. [α]_D -11 (c, 0.34 in EtOH).

2,3-Dihydro: 2,3-Dihydrocapitavine

[91147-13-6]

C₂₁H₂₃NO₄ 353.417

Alkaloid from the fruits of *Buchenavia macrophylla* (Combretaceae). Amorph. [α]_D 0 (c, 1.49 in EtOH).

4'-Hydroxy: 4'-Hydroxycapitavine

[91147-14-7]

C₂₁H₂₁NO₅ 367.401

Alkaloid from the seeds of *Buchenavia capitata* (Combretaceae). Cryst. (MeOH/CH₂Cl₂). Mp 170-172°. [α]_D -11 (c, 0.2 in EtOH).

4'-Hydroxy, 2,3-dihydro: 2,3-Dihydro-4'-hydroxycapitavine

[91147-15-8]


C₂₁H₂₃NO₅ 369.416

Alkaloid from the seeds of *Buchenavia capitata* (Combretaceae). Amorph. [α]_D +51 (c, 1.23 in EtOH).

Ahond, A. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1984, 41 (*isol, uv, ir, pmr, cmr, ms, struct, derivs*)

Capitelline

[204001-20-7]



C₂₀H₂₄N₂O₂ 324.422


Alkaloid from *Hedyotis capitellata*. Oil. [α]_D -4 (c, 0.5 in CHCl₃). λ_{max} 240 (log ε 4.05); 290 (log ε 3.8); 354 (log ε 3.36) (EtOH).

Phuong, N.M. *et al.*, *Nat. Prod. Lett.*, 1998, 11, 93-100 (*isol, uv, ir, pmr, cmr, ms*)

C-94

Cappamensine

3-Hydroxy-7-methoxy-2-methyl-4H-1,4-benzoxazine-4-carboxaldehyde




C₁₁H₁₁NO₄ 221.212

Constit. of the roots of *Capparis sikkimensis* ssp. *formosana*. Antitumour agent. Cryst. Mp 204-206°.

Wu, J.-H. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, 13, 2223-2225 (*isol, pmr, cmr*)

Capparisine

[104899-66-3]



C₂₆H₃₁N₃O₅ 465.548

CAS numbering shown. Alkaloid from the root bark of *Capparis decidua* (Capparidaceae). Cryst. (MeOH). Mp 160-162°.

N¹⁵-Ac: 15-N-Acetylcapparisine

C₂₈H₃₃N₃O₆ 507.585

Alkaloid from root bark of *Capparis decidua* (Capparidaceae). Amorph. powder. Mp 178-180°.

N¹⁵,O-Di-Ac: [104899-65-2]

Amorph. Mp 223-224°.

26-Methoxy: Capparidisine

[100414-82-2]

C₂₇H₃₃N₃O₆ 495.574

Alkaloid from the root bark of *Capparis decidua* (Capparidaceae). Light cream-coloured cryst. (MeOH or Me₂CO aq.). Mp 180-181°.

28-Methoxy: Capparisine

[105694-42-6]

C₂₇H₃₃N₃O₆ 495.574

Alkaloid from root bark of *Capparis decidua*. Cryst. Mp 238°. Positional isomer of Capparidisine. Exists in soln. (DMSO-d₆) as slowly interconverting E- and Z-isomers.

28-Methoxy, N¹⁵,O-di-Ac: [105694-43-7]

Cryst. (MeOH). Mp 263°.

Ahmad, V.U. *et al.*, *Heterocycles*, 1985, 23, 3015 (*Capparidisine*)

Ahmad, V.U. *et al.*, *Z. Naturforsch.*, B, 1986, 41, 1033 (*Capparisine*)


Ahmad, V.U. *et al.*, *Annalen*, 1987, 161 (*Capparisine*)

Ahmad, V.U. *et al.*, *J. Nat. Prod.*, 1992, 55, 1509 (*15-N-Acetylcapparisine*)

C-96

Caprazamycin

C-98



Caprazamycin A R = -(CH₂)₂CH₃
B R = -(CH₂)₁₀CH(CH₃)₂
C R = -(CH₂)₁₁CH₃
D R = -(CH₂)₈CH(CH₃)₂
E R = -(CH₂)₁₀CH₃
F R = -(CH₂)₈CH(CH₃)₂
G R = -(CH₂)₈CH(CH₃)CH₂CH₃

Nucleoside antibiotic complex. Prod. by *Streptomyces* sp. MK730-62F2. Antituberculosis agents.

Caprazamycin A [327164-10-3]

C₅₃H₈₇N₅O₂₂ 1146.291

Powder. [α]_D²³ -1.4 (c, 0.83 in DMSO). λ_{max} 261 (ε 7400) (MeOH).

Caprazamycin B [327164-11-4]

C₅₃H₈₇N₅O₂₂ 1146.291

Powder. [α]_D²³ -2.6 (c, 0.91 in DMSO). λ_{max} 262 (ε 8000) (MeOH).

Caprazamycin C [327164-12-5]

C₅₂H₈₅N₅O₂₂ 1132.264

Powder. [α]_D²³ -1.1 (c, 1.3 in DMSO). λ_{max} 261 (ε 8300) (MeOH).

Caprazamycin D [484686-59-1]

C₅₂H₈₅N₅O₂₂ 1132.264

Powder. [α]_D²³ -3 (c, 1 in MeOH). λ_{max} 262 (ε 9200) (MeOH).

Caprazamycin E [327164-13-6]

C₅₁H₈₃N₅O₂₂ 1118.237

Powder. [α]_D²³ -5.1 (c, 0.83 in DMSO). λ_{max} 262 (ε 7700) (MeOH).

Caprazamycin F [327164-14-7]

C₅₁H₈₃N₅O₂₂ 1118.237

Powder. [α]_D²³ -4.7 (c, 0.9 in DMSO). λ_{max} 262 (ε 7600) (MeOH).

Caprazamycin G [484686-64-8]

C₅₂H₈₅N₅O₂₂ 1132.264

Powder. [α]_D²³ -4.2 (c, 1 in MeOH). λ_{max} 262 (ε 9000) (MeOH).

Igarashi, M. *et al.*, *J. Antibiot.*, 2003, 56, 580-583 (*isol*)


Igarashi, M. *et al.*, *J. Antibiot.*, 2005, 58, 327-337 (*pmr, cmr, cryst struct*)

Ichikawa, S. *et al.*, *Chem. Pharm. Bull.*, 2008, 56, 1059-1072 (*rev, synth*)

Capreomycin

C-99

α,2-Diamino-1,4,5,6-tetrahydro-4-pyrimidineacetic acid, 9CI. α-(Hexahydro-2-imino-4-pyrimidyl)glycine [21012-69-1]



Absolute Configuration

C₆H₁₂N₄O₂ 172.186

Amino acid component of Capreomycin, Tuberactinomycin N and Tuberactinomycin O. Mp 218-230°. [α]_D²⁰ +5.4 (c, 1 in H₂O). p*K*_{a1} 3; p*K*_{a2} 7.6; p*K*_{a3} 13.8 (66% DMF aq.).


Shiba, T. *et al.*, *Tet. Lett.*, 1977, 2681 (*synth, bibl*)

DeMong, D.E. *et al.*, *Tet. Lett.*, 2001, **42**, 3529-3532 (*synth*)

Yamashita, A. *et al.*, *Synth. Commun.*, 2004, **34**, 795-803 (*synth*)

Capsaicin C-100

N-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methyl-6-nonenamide, 9CI. Natural capsaicin. Axsain. Capsacutin. Capsaeni. Mioton. Zostrix. FEMA 3404 [404-86-4]



C₁₈H₂₇NO₃ 305.416

Pungent principle of various *Capsicum* spp. and *Heliopsis longipes*. Flavouring ingredient. Irritant used for desensitisation of sensory neurones. Used as a selective probe for studying neurogenic inflammation and the role of nociceptors in human physiol. Used as a counter-irritant and topical analgesic for some skin conditions. Shows anticancer, antimicrobial and antiinflammatory props. Insect antifeedant. Transient receptor potential vanilloid subfamily 1 (TRPV1) receptor agonist. Cryst. with a burning pungent taste. Mp 64-65°. Bp_{0.01} 210-220°. Log P 3.5 (uncertain value) (calc). λ_{\max} 227 (ϵ 7000); 281 (ϵ 2500) (EtOH) (Derep).

► Skin and mucous membrane irritant. LD₅₀ (mus, orl) 47.2 mg/kg. RA8530000

O- β -D-Glucopyranoside: Capsaicin glucoside

C₂₄H₃₇NO₈ 467.558

Constit. of the fruit of *Capsicum annuum* (red pepper).

Dihydro: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methylnonanamide. Dihydrocapsaicin

[19408-84-5]

C₁₈H₂₉NO₃ 307.432

Minor constit. of natural Capsaicin. Cryst. Mp 65.5-65.8°.

Dihydro, O- β -D-glucopyranoside: Dihydrocapsaicin glucoside

C₂₄H₃₉NO₈ 469.574

Constit. of the fruit of *Capsicum annuum* (red pepper).

9'-Hydroxy: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-hydroxy-8-methyl-6-nonenamide. ω -Hydroxycapsaicin

[69173-71-3]

C₁₈H₂₇NO₄ 321.416

Constit. of the fruit of *Capsicum annuum* (red pepper). Light yellow oil.

[α]_D +3.5 (c, 0.28 in CHCl₃).

(Z)-isomer: Zucapsaicin, INN, USAN [25775-90-0]

C₁₈H₂₇NO₃ 305.416

Analgesic. Cryst. (Et₂O/hexane). Mp 68.5-69.5°.

Octanoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]octanamide, 9CI

[58493-47-3]

C₁₆H₂₅NO₃ 279.378

Constit. of red pepper (*Capsicum annuum*). Component of La Jiao.

6-Methyl-4E-heptenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-6-methyl-4-heptenamide. Dinorcapsaicin.

Nornorcapsaicin

[61229-09-2]

C₁₆H₂₃NO₃ 277.363

Constit. of red pepper (*Capsicum annuum*).

Nonanoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]nonanamide, 9CI. N-Vanillylnonanamide, 8CI. Synthetic capsaicin. Pseudocapsaicin. Peltargonoyl vanillylamide. Nonivamide, INN. PSVA. FEMA 2787. HH 50. AH 23491X. 8-Nordihydrocapsaicin

[2444-46-4]

C₁₇H₂₇NO₃ 293.405

Alkaloid from *Capsicum* spp. (Solanaceae). Topical antirheumatic agent. Shows cardiovascular activity. Cryst.; odourless with a pungent burning taste. Bp_{0.05} 200-210°. Log P 3.59 (uncertain value) (calc).

6-Nonenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-6-nonenamide

[92085-23-9]

C₁₇H₂₅NO₃ 291.389

Alkaloid of *Capsicum* spp. Incorr. named in 1997 ref.

7-Methyloctanoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-7-methyloctanamide, 9CI. Nordihydrocapsaicin

[28789-35-7]

C₁₇H₂₇NO₃ 293.405

Isol. from the pungent principle of red pepper (*Capsicum annuum*). Cryst. (Et₂O/petrol). Mp 60-61°.

7-Methyl-5E-octenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-7-methyl-5-octenamide. Norcapsaicin. 7-Methyl-N-vanillyl-5-octenamide

[61229-08-1]

[62356-06-3, 112375-60-7]

C₁₇H₂₅NO₃ 291.389

Isol. from the pungent principle of red pepper (*Capsicum annuum*). Cryst. (Et₂O/hexane). Mp 42.5-44°.

Decanoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]decanamide

[31078-36-1]

C₁₈H₂₉NO₃ 307.432

Alkaloid from *Capsicum* spp. Cryst. (petrol). Mp 59-60° (synthetic).

9-Methyldecanoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-

methyldecanamide, 9CI. Homodihydrocapsaicin

[20279-06-5]

C₁₉H₃₁NO₃ 321.459

Isol. from the pungent principle of red pepper (*Capsicum annuum*). Cryst. (Et₂O/petrol). Mp 70-71°.

8-Methyl-6E-decenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methyl-6-decanamide

[71240-51-2]

C₁₉H₂₉NO₃ 319.443

Alkaloid from *Capsicum* spp.

9-Methyl-7E-decenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-methyl-7-decanamide, 9CI. Homocapsaicin. 9-Methyl-N-vanillyl-7-decanamide

[58493-48-4]

[112375-61-8, 62356-07-4]

C₁₉H₂₉NO₃ 319.443

Isol. from the pungent principle of red pepper (*Capsicum annuum*). Cryst. (Et₂O/hexane). Mp 64.5-65.5°.

10-Methyl-8E-undecenoyl analogue: N-[(4-Hydroxy-3-methoxyphenyl)methyl]-10-methyl-8-undecenamide

[71226-60-3]

C₂₀H₃₁NO₃ 333.47

Alkaloid from *Capsicum* spp. Glass.

[7553-53-9]

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 1292A; 1293A (*ir*)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **2**, 619A; 621A; 1378C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase,

1989, **3**, 1174D (*ir*)

Nelson, E.K. *et al.*, *J.A.C.S.*, 1919, **41**, 1115-

1121 (*struct*)

Crombie, L. *et al.*, *J.C.S.*, 1955, 1025-1027

(*synth, bibl*)

Kosuge, S. *et al.*, *CA*, 1963, **59**, 6724; 1964, **60**,

9827 (*Dihydrocapsaicin*)

Dirnhuber, P. *et al.*, *J. Physiol. (London)*,

1965, **178**, 41P (*Nonivamide, pharmacol*)

Friedrich, H. *et al.*, *Naturwissenschaften*, 1965,

52, 514 (*Nonivamide*)

Ger. Pat., 1966, ((Krewel-Leuffen))1 213 855;

CA, **64**, 19504h (*Nonivamide, synth*)

Leete, E. *et al.*, *J.A.C.S.*, 1968, **90**, 6837-6841

(*biosynth*)

Jentsch, K. *et al.*, *Monatsh. Chem.*, 1968, **99**,

661-663 (*Homodihydrocapsaicin*)

Bowman, W.R. *et al.*, *Chem. Comm.*, 1969,

1075-1077 (*biosynth*)

Rangoonwala, R. *et al.*, *Pharmazie*, 1969, **24**,

177 (*biosynth*)

Kosuge, S. *et al.*, *Agric. Biol. Chem.*, 1970, **34**,

248-256 (*Nordihydrocapsaicin*,

Homodihydrocapsaicin)

Refshange, C. *et al.*, *Life Sci.*, 1974, **14**, 311

(*deriv*)

Jeso, I. *et al.*, *Chem. Zvesti*, 1975, **29**, 714

(*synth*)

Todd, P. *et al.*, *J. Chromatogr. Sci.*, 1975, **13**,

577 (*Nonivamide, ilc*)

Virus, R.M. *et al.*, *Life Sci.*, 1979, **25**, 1273-

1283 (*rev, props*)

Jurenitsch, J. *et al.*, *Planta Med.*, 1979, **36**, 61-

67 (*N-Octanoylcapsaicin*, *Homocapsaicin*)

Monsereenusorn, Y. *et al.*, *Crit. Rev. Toxicol.*,

1982, **10**, 321-339 (*rev, props*)


Jurenitsch, J. *et al.*, *Sci. Pharm.*, 1982, **50**, 64

(*Nonivamide, isol*)

- Suzuki, T. *et al.*, *Alkaloids (London)*, 1983, **23**, 227-299 (rev)
- Skofitsch, G. *et al.*, *Arzneim.-Forsch.*, 1984, **34**, 154 (*Nonivamide, pharmacol*)
- Gannett, P.M. *et al.*, *J.O.C.*, 1988, **53**, 1064-1071; 6162 (*Capsaicins, synth, ir, pmr, cmr, ms*)
- Kaga, H. *et al.*, *J.O.C.*, 1989, **54**, 3477-3478 (*synth, bibl*)
- Kaga, H. *et al.*, *Synthesis*, 1989, 864-866 (*Nordihydrocapsaicin, Homodihydrocapsaicin, synth, ir*)
- Fuller, R.W. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1990, **303**, 147-156 (rev, *pharmacol*)
- Maggi, C.A. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1990, **303**, 157-166 (*pharmacol*)
- Kaga, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1992, **56**, 946-948 (*Capsacinoids, synth*)
- Cordell, G.A. *et al.*, *Ann. Pharmacother.*, 1993, **27**, 330-336 (rev)
- Janusz, J.M. *et al.*, *J. Med. Chem.*, 1993, **36**, 2595-2604 (*Zucapsaicin*)
- Lo, Y.-C. *et al.*, *Brain Res. Bull.*, 1994, **35**, 15 (*Nonivamide, pharmacol*)
- Kirby, G.W. *et al.*, *Phytochemistry*, 1994, **36**, 185-187 (*purifn, bibl*)
- Constant, H.L. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1925-1928 (*minor capsaicinoids, occur, anal*)
- Surh, Y.-J. *et al.*, *Life Sci.*, 1995, **56**, 1845-1855 (rev, *tox, metab*)
- Kaga, H. *et al.*, *Tetrahedron*, 1996, **52**, 8451-8470 (*Nordihydrocapsaicin, Homocapsaicin, synth, ir, pmr, cmr, ms*)
- Fusco, B.M. *et al.*, *Drugs*, 1997, **53**, 909-914 (rev)

- Encyclopedia of Food and Color Additives*, (ed. Burdock, G.A.), CRC Press, 1997, 1374-1375; 1979-1980 (*Capsaicin, Nonivamide*)
- Maillard, M.N. *et al.*, *Flavour Fragrance J.*, 1997, **12**, 409-413 (*minor capsaicinoids, occur, anal*)
- David, W.I.F. *et al.*, *Chem. Comm.*, 1998, 931-932 (*cryst struct*)
- Molina-Torres, J. *et al.*, *J. Ethnopharmacol.*, 1999, **64**, 241-248 (*activity*)
- Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1559
- Fenaroli's *Handbook of Flavor Ingredients*, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 819-820; 1312 (*Capsaicin, Nonivamide*)
- Merck Index, 13th edn., 2001, No. 1774 (*bibl*)
- Ochi, T. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1094-1096 (*ω-Hydroxycapsaicin*)
- Chayapathy, B. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 1854-1859 (*biosynth*)
- Higashiguchi, F. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 5948-5953 (*glucosides*)
- Remadevi, R. *et al.*, *Drugs*, 2008, **11**, 120-132 (*pharmacol*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CBF750

Capsaicinol **C-101**
[133631-97-7, 137493-36-8]




C₁₈H₂₇NO₄ 321.416

(R)-form [137493-35-7]

- Constit. of the fruit of *Capsicum frutescens* (red chilli/tabasco pepper). Antioxidant. $[\alpha]_D^{25}$ -9 (c, 1 in CHCl₃).
- Masuda, T. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 2337-2340 (*synth, pmr, abs config*)
- Kobata, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 1904-1912 (*synth, activity*)


Capsimycin **C-102**
N 461. Antibiotic N 461
[70694-08-5]



Absolute Configuration

- C₃₀H₄₀N₂O₆ 524.656
- Isol. from *Streptomyces* sp. C 49-87. Antifungal antibiotic, esp. against phyto-toxic fungi. Apoptosis inducer. Sol. MeOH, CHCl₃, Py, EtOAc; fairly sol. H₂O; poorly sol. C₆H₆, hexane. Mp 186°. $[\alpha]_D^{20}$ +196 (c, 1 in CHCl₃). pK_a 5.5 (50% THF aq.). Similar to Ikarugamycin. λ_{\max} 220 (ε 48200); 325 (ε 31200) (MeOH/HCl) (Derep). λ_{\max} 320 (ε 29800) (MeOH/NaOH) (Derep).
- LD₅₀ (mus, orl) 600-700 mg/kg. EY4930000
- Seto, H. *et al.*, *CA*, 1979, **92**, 211459 (*cryst struct*)
- Aizawa, S. *et al.*, *J. Antibiot.*, 1979, **32**, 193-196 (*isol*)


Capuramycin **C-103**
Antibiotic 446-S3-1. Antibiotic A 500359B. A 500359B
[102135-48-8]



- C₂₃H₃₁N₅O₁₂ 569.524
- Nucleoside antibiotic. Prod. by *Streptomyces griseus*. Active against *Streptococcus pneumoniae* and *Mycobacterium smegmatis*. Amorph. Mp 173-176°. $[\alpha]_D^{25}$ +99 (c, 0.5 in H₂O). λ_{\max} 214 (ε 16200); 257 (MeOH) (Derep).
- 2'-O-Carbamoyl: Antibiotic A 503083B. A 503083B
- C₂₄H₃₂N₆O₁₃ 612.549
- Prod. by *Streptomyces* sp. SANK 62799. Powder. $[\alpha]_D^{20}$ +91 (c, 0.2 in H₂O). λ_{\max} 257 (ε 10900) (H₂O).
- O-De-Me: Antibiotic A 500359G. A 500359G
- C₂₂H₂₉N₅O₁₂ 555.497
- Prod. by *Streptomyces griseus* SANK 60196. Powder. $[\alpha]_D^{20}$ +110 (c, 0.72 in H₂O). λ_{\max} 257 (ε 10000) (H₂O).
- Yamagushi, H. *et al.*, *J. Antibiot.*, 1986, **39**, 1047-1053 (*isol, struct, props, nmr, uv*)
- Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711-1739 (rev)
- Seto, H. *et al.*, *Tet. Lett.*, 1988, **29**, 2343-2346 (*struct*)

- Knapp, S. *et al.*, *J.O.C.*, 1994, **59**, 281-283 (*synth*)
- Muramatsu, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 243-252; 253-258; 259-267; 268-279 (*isol, pmr, cmr, biosynth*)
- Muramatsu, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 639-646 (*A 503083B*)
- Muramatsu, Y. *et al.*, *J. Antibiot.*, 2006, **59**, 601-606 (*prodn*)

Capuronine **C-104**
5-Ethyl-1,4,5,6,7,8,9,10-octahydro-2H-methanoazacycloundecino[5,4-b]indol-6-ol, 9CI
[57550-22-8]




Absolute Configuration

- C₁₉H₂₆N₂O 298.427
- Alkaloid from stem bark of *Capuronetta elegans* (preferred genus name *Tabernaemontana*) (Apocynaceae). Cryst. (MeOH). Mp 75°. $[\alpha]_D^{20}$ +70 (c, 1 in CHCl₃).

O-Ac:

- Cryst. (MeOH). Mp 135°. $[\alpha]_D^{20}$ -76 (c, 1 in CHCl₃).
- Chardon-Loriaux, I. *et al.*, *Tet. Lett.*, 1975, 1845 (*isol, uv, ms, pmr, cmr, struct*)

Capuvsidine **C-105**
[70031-43-5]




- C₄₀H₄₈N₄O₂ 616.845
- Alkaloid from *Capuronetta elegans* (preferred genus name *Tabernaemontana*) and *Pandaca boiteau* (Apocynaceae). $[\alpha]_D^{20}$ -7 (c, 0.71 in CHCl₃).

1',2'-Dihydro: **Dihydrocapuvsidine**
[72186-19-7]

- C₄₀H₅₀N₄O₂ 618.861
- Alkaloid from *Capuronetta elegans* and *Pandaca boiteau* (Apocynaceae). Amorph. λ_{\max} 228 (log ε 4.61); 280 (log ε 3.99); 286 (log ε 3.99); 295 (log ε 3.86) (EtOH).
- Husson, H.-P. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 1099-1102 (*uv, ir, pmr, ms, struct, dihydro*)
- Chardon-Loriaux, I. *et al.*, *Phytochemistry*, 1978, **17**, 1605-1608 (*isol, uv, ir, pmr, ms, struct*)
- Andriantsiferana, M. *et al.*, *Phytochemistry*, 1979, **18**, 911-912 (*isol, dihydro*)

Capuvosine

[57533-88-7]



C₄₀H₅₀N₄O₃ 634.86

Alkaloid from the leaves and stem bark of *Capuronetta elegans* (preferred genus name *Tabernaemontana*) (Apocynaceae). Cryst. (MeOH). Mp 236°. [α]_D²⁰ -53 (c, 1 in CHCl₃).

N-De-Me: **N-Demethylcapuvosine**

[70031-42-4]

C₃₉H₄₈N₄O₃ 620.833

Alkaloid from *Capuronetta elegans* (Apocynaceae). Amorph. [α]_D²⁰ -85 (c, 0.53 in CHCl₃).

Deoxy: **Dehydroxycapuvosine**

[70031-41-3]

C₄₀H₅₀N₄O₂ 618.861

Alkaloid from *Capuronetta elegans* and *Pandaca boiteaui* (Apocynaceae). Amorph. [α]_D²⁰ -98 (c, 0.83 in CHCl₃).

Chardon-Loriaux, I. et al., *Tet. Lett.*, 1975, 1845-1848 (uv, ir, pmr, struct)


Husson, H.-P. et al., *J. Indian Chem. Soc.*, 1978, **55**, 1099-1102 (uv, ir, pmr, ms, Dehydroxycapuvosine)

Chardon-Loriaux, I. et al., *Phytochemistry*, 1978, **17**, 1605-1608 (isol, uv, ir, pmr, ms, struct, derivs)

Caracasamide

C-107

3-(3,4-Dimethoxyphenyl)-N-[4-[[iminol(3-methyl-2-butenyl)amino]methyl]amino]butyl]-2-propenamide, **9CI** [128009-16-5]



C₂₁H₃₂N₄O₃ 388.509

Antihypertensive agent.

(**E**)-form [146269-39-8]

Constit. of *Verbesina caracasana*. Foam.

(**Z**)-form [146269-40-1]

Constit. of *Verbesina caracasana*. Foam. λ_{max} 235 ; 291 ; 316 (MeOH).


Monache, G.D. et al., *Bioorg. Med. Chem. Lett.*, 1992, **2**, 415-418 (isol, struct)

Crombie, L. et al., *J.C.S. Perkin 1*, 1992, 3179-3183 (synth, pmr)

Monache, G.D. et al., *J. Med. Chem.*, 1993, **36**, 2956-2963 (synth, uv, ir, pmr, cmr, ms, pharmacol)

Caracasandiamide

C-108



C₄₂H₆₄N₈O₆ 777.018

Deriv. of 3,4-Diphenyl-1,2-cyclobutanedicarboxylic acid. Alkaloid from *Verbesina caracasana*. Hypotensive agent. Foam.

Carmignani, M. et al., *J. Med. Chem.*, 1999, **42**, 3116-3125 (isol, synth, pharmacol, ir, pmr, cmr, ms)

Caracurine I

C-109


Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 300° (as picrate). Purple-red col. with Ce(SO₄)₂, becoming violet on standing. ▶ Highly toxic.

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1983-1992

Caracurine V, 9CI

C-110

[630-87-5]



C₃₈H₄₀N₄O₂ 584.76

Alkaloid from *Strychnos toxifera*, *Strychnos afzelii*, *Strychnos dolichothyrsa*, *Strychnos urcolata*, *Strychnos angolensis*, *Strychnos chrysophylla* and calabash curare. Potent curarising muscle relaxant. Also shows antimicrobial props. Active against gram-positive bacteria and fungi. Amorph. Purple-red col. with Ce(SO₄)₂, turning brown on standing. λ_{max} 259 ; 299 (MeOH) (Berdy).

▶ Toxic.

N^b-Oxide: **Caracurine V N-oxide**

[66067-01-4]

C₃₈H₄₀N₄O₃ 600.759

Alkaloid from *Strychnos dolichothyrsa* (Loganiaceae). Less potent muscle relaxant than Caracurine V but less toxic.

N^b,N^{b'}-Dioxide: **Caracurine V di-N-oxide** [66067-02-5]

C₃₈H₄₀N₄O₄ 616.758

Alkaloid from *Strychnos dolichothyrsa* and *Strychnos toxifera*. Less potent muscle relaxant than Caracurine V but less toxic. Active against gram-positive bacteria and fungi.

N^b,N^{b'}-Di-Me: **Caracurine V dimethosalt** C₄₀H₄₆N₄O₂[⊕] 614.829

Calabash curare alkaloid. Isom. to Toxiferine I, T-436 under mild acid conds.

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1983; 1993 (isol, uv)

Bernaer, K. et al., *Helv. Chim. Acta*, 1958, **41**, 2293 (struct, synth)

Battersby, A.R. et al., *J.C.S.*, 1960, 736 (synth)

Battersby, A.R. et al., *Proc. Chem. Soc., London*, 1961, 413 (pmr)

Verpoorte, R. et al., *J. Pharm. Sci.*, 1978, **67**, 171-174 (isol, cmr, ir, oxides)

Verpoorte, R. et al., *Planta Med.*, 1978, **33**, 237-242 (activity)

Bohlin, L. et al., *Planta Med.*, 6th Ed., 1979, **35**, 19 (isol)

Delaude, C. et al., *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (occur, *Strychnos*)


Zlotos, D.P. et al., *J. Nat. Prod.*, 2000, **63**, 864-865 (synth, pmr, cmr, abs config)

Zlotos, D.P. et al., *Eur. J. Org. Chem.*, 2004, 2375-2380 (pmr, conformn)

Caracurine II

C-111

[5516-84-7]



Absolute configuration

C₃₈H₃₈N₄O₂ 582.744

Alkaloid from *Strychnos toxifera* and calabash curare (Loganiaceae). Cryst. Mp 248-249°. [α]_D²⁶ -232 (c, 0.66 in CHCl₃). Purple col. with CeSO₄, turning brown on standing.

N^b,N^{b'}-Di-Me: **Toxiferine IX. Caracurine II dimethosalt**

C₄₀H₄₄N₄O₂[⊕] 612.813

Alkaloid from *Strychnos toxifera* (Loganiaceae). [α]_D²⁴ -106 (c, 1.02 in 50% MeOH aq.). Violet col. with Ce(IV).

King, H. et al., *J.C.S.*, 1949, 3263 (isol, deriv)

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1983; 1993

Zürcher, A. et al., *J.A.C.S.*, 1958, **80**, 1500 (isol)

Battersby, A.R. et al., *J.C.S.*, 1960, 736; 1848 (deriv)

Schroeder, H.-D. et al., *Helv. Chim. Acta*, 1961, **44**, 34 (uv, synth)

Battersby, A.R. et al., *Proc. Chem. Soc., London*, 1961, 412; 413 (pmr, uv, struct)

McPhail, A.T. et al., *Proc. Chem. Soc., London*, 1961, 416 (cryst struct)

Bourne, P.E. et al., *J. Crystallogr. Spectrosc. Res.*, 1985, **15**, 453 (cryst struct, deriv)

Caracurine III

C-112

[1355-13-1]

Struct. unknown. Alkaloid from the bark of *Strychnos toxifera* (Loganiaceae). Characterised as picrate, Mp. >300° (darkening from 210°). Purple-red col. with Ce(SO₄)₂, becoming brown-red on standing.

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1983-1992 (isol, uv)

Caracurine IV

C-113

C₂₁H₂₄N₂O₂ 336.433

Minimum formula. Struct. unknown. Alkaloid from *Strychnos toxifera* (calabash curare) (Loganiaceae). Mp 300° (as

picrate). Violet col. with $\text{Ce}(\text{SO}_4)_2$, becoming orange-brown on standing.

Asmis, H. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1983-1992 (*isol*, *uv*)

Caracurine IX C-114

Struct. unknown


Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 260° dec. (as picrate). Red-viol. col. with $\text{Ce}(\text{SO}_4)_2$, becoming brown on standing.

► Highly toxic, $\text{LD}_{50} = 1$ mg/kg.

Asmis, H. *et al.*, *Helv. Chim. Acta*, 1955, **38**, 1661-1668 (*isol*, *uv*)

Caracurine VII C-115

19,20-Didehydro-17,18-epoxycuran-17-ol, 9CI. Wieland-Gumlich aldehyde. Deacetyldiaboline
[466-85-3]



Absolute Configuration

$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_2$ 310.395

Alkaloid from *Strychnos toxifera*, *Strychnos chrysophylla*, *Strychnos dolichothyrsa*, *Strychnos kasengaensis* and other *Strychnos* spp. Prev. known as an important degradn. prod. of strychnine. Key intermed. in the biosynth. of the dimeric calabash curare alkaloids (C-curarines, C-toxiferines, etc.). Mp 211-213° dec. $[\alpha]_{\text{D}}^{22} -134$ (c, 0.52 in MeOH). Stable orange colour with $\text{Ce}(\text{SO}_4)_2$.

► Exp. reprod. effects. HI3300000

Picrate: Mp 235-237°.

N^b -Oxide: **Wieland-Gumlich aldehyde N-oxide**

[174364-49-9]

$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_3$ 326.394

Alkaloid from the root bark of *Strychnos angolensis*. $[\alpha]_{\text{D}} -14$ (c, 0.23 in MeOH).

N^a -Ac: **Diaboline**

[509-40-0]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from *Strychnos diaboli*, *Strychnos potatorum*, *Strychnos henningsii*, *Strychnos nux-vomica*, *Strychnos fendleri*, *Strychnos castelneana*, *Strychnos cathayensis*, *Strychnos pseudoquina* and some other *Strychnos* spp. (Loganiaceae). Weak convulsive and hypotensive agent. Mp 187-189°. $[\alpha]_{\text{D}}^{27} +44.8$ (c, 1 in EtOH).

► HI3162000

N^a -Ac; hydrochloride:

Cryst. + 2H₂O. Mp 238-240° dec. $[\alpha]_{\text{D}} +110.5$.

N^a -Ac, picrate: Mp 234-240°.

N^a -Ac, N^b -oxide: **Diaboline N-oxide**

[85769-35-3]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from root bark of *Strychnos potatorum*.

N^a , O-Di-Ac: **Henningsamine. Acetyldiaboline**

[2871-28-5]

$\text{C}_{23}\text{H}_{26}\text{N}_2\text{O}_4$ 394.469

Alkaloid from *Strychnos henningsii* bark, *Strychnos pungens* and *Strychnos potatorum* (Loganiaceae). Needles (EtOAc/petrol). Mp 205-206°. $[\alpha]_{\text{D}}^{19} -43.9$ (c, 1 in CHCl_3).

N^a , O-Di-Ac, picrate:

Needles (MeOH). Mp 229-231°.

N^b -Me: **Hemitoxiferine I. Alkaloid A8**

[6879-95-4]

$\text{C}_{20}\text{H}_{25}\text{N}_2\text{O}_2^{\oplus}$ 325.43

Alkaloid from *Strychnos toxifera* (Loganiaceae). Mp 300° (as chloride). $[\alpha]_{\text{D}}^{24} -43$ (c, 1 in H₂O). Stable orange colour with $\text{Ce}(\text{SO}_4)_2$.

Et ether, N^a -Ac: **Ethyldiaboline**

$\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_3$ 380.486

Alkaloid from root bark of *Strychnos castelneana* (Loganiaceae). Mp 283-285° (as picrate). Doubtless an artifact, prob. derived from EtOH present in solvent CHCl_3 . No CAS Reg. No. 1967-2001 (8CI-14CI).

Deoxy: **Desoxy-Wieland-Gumlich aldehyde. 19,20-Didehydro-17,18-epoxycuran**

$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}$ 294.396

Alkaloid from root bark of *Strychnos kasengaensis* and *Strychnos matopensis* and from stem bark of *Strychnos dolichothyrsa* (tentative identification) (Loganiaceae). No CAS Reg. No. 1967-2001 (8CI-14CI).

Deoxy, N^a -formyl: **N^a -Formyloxydeoxy-Wieland-Gumlich aldehyde**

[119308-21-3]

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_2$ 322.406

Alkaloid from root bark of *Strychnos matopensis*. $[\alpha]_{\text{D}} +39$ (c, 0.76 in CHCl_3). λ_{max} 218 ; 253 ; 282 ; 290 (MeOH).

2,16-Didehydro, N^a -Ac: **2,16-Dehydrodiaboline**

$\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_3$ 350.416

Alkaloid from *Strychnos henningsii*. No phys. props. reported. No CAS Reg. No. 1967-2001 (8CI-14CI).

3 α -Hydroxy, N^a -Ac: **3-Hydroxydiaboline**

[85769-36-4]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from upper stem bark of *Strychnos castelneana* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 218-221°. $[\alpha]_{\text{D}}^{20} +86$ (c, 0.6 in CHCl_3).

11-Methoxy: **11-Methoxy-Wieland-Gumlich aldehyde**

[36151-17-4]

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_3$ 340.421

Alkaloid from *Strychnos angolensis* (Loganiaceae).

11-Methoxy, N^a -Ac: **11-Methoxydiaboline**

[36151-16-3]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from the stem bark of *Strychnos romeu-belenii*, *Strychnos rubiginosa* and *Strychnos malacoclados*, also from *Strychnos gardneri*, *Strychnos angolensis* and *Strychnos cath-*

ayensis (Loganiaceae). Prisms (EtOAc). Mp 214-216°. $[\alpha]_{\text{D}}^{20} +20$ (c, 1 in CHCl_3).

11-Methoxy, N^a -Ac, N^b -oxide: **11-Methoxydiaboline N-oxide**

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_5$ 398.458

Alkaloid from *Strychnos malacoclados*. Details not publ. (R. Verpoorte). No CAS Reg. No. 1967-2001 (8CI-14CI).

11-Methoxy, N^a , O-di-Ac: **Condensamine. 11-Methoxyhenningsamine**

[36536-63-7]

$\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_5$ 424.496

Alkaloid from the bark of *Strychnos henningsii*, from *Strychnos holstii* var. *reticulata*, *Strychnos cocculoides* and other *Strychnos* spp. (Loganiaceae). Mp 262-265° (238-240° dec.). $[\alpha]_{\text{D}}^{21} -20$ (CHCl_3). 11-Methoxyhenningsamine and Condensamine are descr. as different alkaloids in the Delaude review, but clearly have the same struct.

11-Methoxy, Me ether: **17-O-Methyl-11-methoxy-Wieland-Gumlich aldehyde.**

19,20-Didehydro-17,18-epoxy-11,17-dimethoxycuran, 9CI

[70278-01-2]

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3$ 354.448

Alkaloid from *Strychnos angolensis* (Loganiaceae).

11-Methoxy, 19,20-dihydro, N^a -Ac: **19,20-Dihydro-11-methoxydiaboline**

[69320-14-5]

$\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4$ 384.474

Alkaloid from *Strychnos malacoclados*.

11-Methoxy, 2,16-didehydro, N^a -Ac: **2,16-Didehydro-11-methoxydiaboline. 11-Methoxy-2,16-dehydrodiaboline**

$\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_4$ 380.443

Alkaloid from *Strychnos henningsii*. No CAS Reg. No. 1967-2001 (8CI-14CI).

11-Methoxy, 12-hydroxy, N^a -Ac: **12-Hydroxy-11-methoxydiaboline**

[95260-32-5]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_5$ 398.458

Alkaloid from *Strychnos potatorum* and *Strychnos panganensis*. $[\alpha]_{\text{D}} -147$ (c, 1 in CHCl_3).

11-Methoxy, 12-hydroxy, N^a , O-di-Ac: **12-Hydroxy-11-methoxyhenningsamine**

[18797-84-7]

$\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_6$ 440.495

Alkaloid from *Strychnos henningsii*, *Strychnos pungens* and *Strychnos staudtii* (Loganiaceae). $[\alpha]_{\text{D}}^{23} -142$ (c, 0.7 in CHCl_3). V. readily hydrolysed.

17-Epimer, O¹⁷, N^a -di-Ac: **Jobertine. O-Acetyldiaboline B**

[2864-87-1]

$\text{C}_{23}\text{H}_{26}\text{N}_2\text{O}_4$ 394.469

Alkaloid from the root bark of *Strychnos jobertiana* and trunk bark of *Strychnos castelneana* (Loganiaceae). Mp 87-91° (92-94.5°). $[\alpha]_{\text{D}}^{21} +51.2$ (c, 1 in CHCl_3).

17-Epimer, 11-methoxy, O¹⁷-Me, N^a -Ac: **17-Epi-11-methoxy-17-O-methyldiaboline**

[70280-98-7]

$\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_4$ 396.485

Alkaloid from *Strychnos angolensis*

(Loganiaceae). Needles (EtOAc/Et₂O). Mp 190-192°.

17-Epimer, 11-methoxy, 12-hydroxy, N^α-Ac: **Henningsoline**. 17-Epi-12-hydroxy-11-methoxydiaboline [18797-85-8]

C₂₂H₂₆N₂O₅ 398.458

Alkaloid from the bark of *Strychnos henningsii*, *Strychnos cathayensis* and from stem bark of *Strychnos spinosa* (Loganiaceae). Prisms (Me₂CO). Mp 207-209°. [α]_D²⁰ -200 (c, 1 in CHCl₃). pK_a 7.1.

17-Epimer, 11-methoxy, 12-hydroxy, N^α-Ac, perchlorate:

Prisms (MeOH). Mp 253-254° dec.

17-Epimer, 11-methoxy, 12-hydroxy, N^α-Ac, picrate:

Prisms (MeOH aq.). Mp 167-168°.

Wieland, H. et al., *Annalen*, 1932, **494**, 191-200; 1933, **506**, 60-76 (*synth*)

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1968-1973; 1993-2001 (*isol, uv*)

Battersby, A.R. et al., *Proc. Chem. Soc., London*, 1959, 126 (*Diaboline, struct*)

Battersby, A.R. et al., *J.C.S.*, 1960, 736-741; 1848-1854 (*Hemitoxiferine*)

Deyrup, J.A. et al., *Helv. Chim. Acta*, 1962, **45**, 2266-2272 (*Diaboline, synth*)

Oecolowitz, J. et al., *Farmaco, Ed. Sci.*, 1965, **20**, 751-756 (*Condensamine*)

Grossert, J.S. et al., *J.C.S.*, 1965, 2812-2814; 2814-2818; 2818-2822 (*Henningsamine, Henningsoline, Jobertine, isol*)

Biemann, K. et al., *J.C.S.*, 1965, 2814-2818 (*Henningsamine, struct*)

Delle Monache, F. et al., *Ann. Ist. Super. Sanita*, 1967, **3**, 564; *CA*, **68**, 87440u (*Jobertine, struct*)

Spiteller-Friedmann, M. et al., *Annalen*, 1968, **712**, 179-194 (*Dehydrodiaboline, 2,16-Didehydro-11-methoxydiaboline, Henningsoline*)

Hymon, J.R. et al., *Helv. Chim. Acta*, 1969, **52**, 1564-1602 (*synth*)

Delle Monache, F. et al., *J. Nat. Prod.*, 1970, **33**, 279-283 (*Ethylidiaboline*)

Marini-Bettolo, G.B. et al., *Gazz. Chim. Ital.*, 1971, **101**, 971-980 (*11-Methoxydiaboline*)

Szabó, L. et al., *Acta Chim. Acad. Sci. Hung.*, 1977, **95**, 85-100 (*synth*)

Wenkert, E. et al., *J.O.C.*, 1978, **43**, 1099-1105 (*pmr, cmr*)

Verpoorte, R. et al., *Pharm. Weekbl.*, 1978, **113**, 1249-1261 (*19,20-Dihydro-11-methoxydiaboline*)

Bohlin, L. et al., *Planta Med.*, 1979, **35**, 19-30 (*Strychnos angolensis constits*)

Galeffi, G. et al., *Phytochemistry*, 1982, **21**, 2393-2395 (*3-Hydroxydiaboline*)

Verpoorte, R. et al., *Planta Med.*, 1982, **44**, 21-27 (*Desoxy-Wieland-Gumlich aldehyde*)

Thepenier, P. et al., *Phytochemistry*, 1984, **23**, 2659-2663; 1988, **27**, 657-659 (*Desoxy-Wieland-Gumlich aldehyde, 12-Hydroxy-11-methoxyhenningsamine, 12-Hydroxy-11-methoxydiaboline*)

Ohiri, F.C. et al., *Planta Med.*, 1984, **50**, 446-447 (*Henningsoline, isol, uv, pmr*)

Kapoor, V.K. et al., *Indian J. Chem., Sect. B*, 1988, **27**, 641-644 (*synth, uv, ir, pmr, ms, Diaboline*)

Massiot, G. et al., *Phytochemistry*, 1988, **27**, 3293-3304; 1992, **31**, 2873-2876 (*N-Formyldeoxy-Wieland-Gumlich aldehyde, Diaboline N-oxide*)

Magnus, P. et al., *J.A.C.S.*, 1993, **115**, 8116-8129 (*synth*)

Delaude, C. et al., *Bull. Soc. R. Sci. Liege*, 1995, **64**, 243-246; 1997, **66**, 183-286 (*N-oxide, isol, occur, derivs, rev*)

Caracurine VIII

C-116


Struct. unknown. Alkaloid from bark of *Strychnos toxifera* (Loganiaceae). Mp 240° dec. (as picrate). Purple red col. with Ce(SO₄)₂, becoming brown on standing. ▶ Highly toxic, LD₅₀ = 1 mg/kg.

Asmis, H. et al., *Helv. Chim. Acta*, 1955, **38**, 1661-1668 (*isol, uv*)

Carapanaubine

C-117

Isoreserpiline oxindole B. Vinine [1255-02-3]



C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Aspidosperma carapanauba*, *Rauwolfia vomitoria*, *Cabucala madagascariensis* var. *amygdalifolia* (Apocynaceae). Cryst. (EtOAc/hexane or EtOH aq.). Mp 221-223°. [α]_D²⁰ -101 (c, 1.0 in CHCl₃). λ_{max} 218 (ε 28800); 245 (sh); 280 (ε 5000); 300 (sh) (EtOH).

N-Oxide: **Carapanaubine N-oxide**

[64675-22-5]

C₂₃H₂₈N₂O₇ 444.483

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae).

3-Epimer: **Rauvoxine. Reserpiline oxindole B. 10,11-Dimethoxyuncarine F** [4062-36-6]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from the leaves of *Rauwolfia vomitoria* (Apocynaceae). Cryst. (MeOH). Mp 210°. [α]_D²⁰ +98 (c, 1.34 in CHCl₃/EtOH 199:1). λ_{max} 218 (ε 26300); 245 (sh); 280 (ε 5300); 300 (sh) (EtOH).

7-Epimer: **Isocarapanaubine. Isoreserpiline oxindole A**

[17391-09-2]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Rauwolfia vomitoria*, *Rauwolfia salicifolia* and *Ochrosia moorei* (Apocynaceae). Amorph. [α]_D²⁰ -64 (CHCl₃). λ_{max} 220 (log ε 4.52); 245 (sh) (log ε 4.19); 300 (sh) (log ε 3.91) (EtOH).

20-Epimer: **10,11-Dimethoxymitrephylline**

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Cabucala cryptophlebia*. Mp 181° (108-110°). [α]_D²⁰ -4.6 (c, 0.21 in CHCl₃).

3,7-Diepimer: **Rauvoxinine. 10,11-Dimethoxyuncarine D. Reserpiline oxindole A** [3993-09-7]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Rauwolfia vomitoria*, *Cabucala fasciculata* and *Cabucala madagascariensis* var. *amygdalifolia* (Apocynaceae). Mp 203-204°. [α]_D²⁰ +64.6 (c, 1.17 in CHCl₃/EtOH 199:1).

λ_{max} 218 (ε 24500); 245 (sh); 280 (ε 5600); 300 (sh) (EtOH).

7,19-Diepimer: **Neisosposinine**

[122923-45-9]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from the stem bark of *Neisosperma oppositifolia*. Plates (Me₂CO). Mp 228-230°. λ_{max} 215 (log ε 4.54); 245 (sh) (log ε 4.21); 302 (sh) (log ε 3.87) (EtOH).

7,20-Diepimer: **10,11-Dimethoxyisomitrephylline**

[53796-85-3]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from the aerial parts of *Cabucala madagascariensis* var. *amygdalifolia* and the leaves of *Cabucala fasciculata* (Apocynaceae). Mp 142° (115-117). [α]_D²⁰ +19.6 (c, 0.2 in CHCl₃).

7,20-Diepimer, 16α,17-dihydro: **Herbaline**

[6516-50-3]

C₂₃H₃₀N₂O₆ 430.5

Alkaloid from *Vinca herbacea* (Apocynaceae). Cryst. (EtOH). Mp 276-278° dec. [α]_D²⁰ -147 (c, 1.5 in Py). λ_{max} 215 (ε 36000); 243 (sh) (ε 9450); 273 (ε 11250); 305 (ε 9900) (EtOH).

Finch, N. et al., *J.A.C.S.*, 1963, **85**, 1520-1523 (*Carapanaubine, synth*)

Gilbert, B. et al., *J.A.C.S.*, 1963, **85**, 1523-1528 (*Carapanaubine, isol, uv, pmr, ms, struct*)

Pousset, J.-L. et al., *C. R. Hebd. Seances Acad. Sci.*, 1964, **259**, 597-600 (*Rauvoxine, Rauvoxinine, isol, pmr, struct*)

Patel, M.B. et al., *J. Pharm. Pharmacol.*, 1964, **16**, 163T-165T (*Rauvoxine, Rauvoxinine, isol, uv, ir*)

Ognyanov, I. et al., *Chem. Ber.*, 1966, **99**, 2052-2056 (*Herbaline*)

Pousset, J.-L. et al., *Bull. Soc. Chim. Fr.*, 1967, 2766-2779 (*Isocarapanaubine, Rauvoxine, Rauvoxinine, synth, ir, uv, pmr, config*)

Ognyanov, I. et al., *Chem. Comm.*, 1967, 579-581 (*Herbaline, stereochem*)

Shamma, M. et al., *J.A.C.S.*, 1967, **89**, 1739-1740 (*stereochem*)

Poisson, J. et al., *Tet. Lett.*, 1967, 1919-1923 (*config*)

Pascard-Billy, C. et al., *Bull. Soc. Chim. Fr.*, 1968, 3289-3295 (*Rauvoxinine, cryst struct*)

Kan-Fan, C. et al., *Phytochemistry*, 1972, **11**, 435 (*Dimethoxyisomitrephylline*)

Iwu, M.M. et al., *Planta Med.*, 1977, **32**, 88 (*Carapanaubine N-oxide*)

Sierra, P. et al., *Coll. Czech. Chem. Comm.*, 1982, **47**, 2912-2921 (*Isocarapanaubine*)

Verpoorte, R. et al., *Org. Magn. Reson.*, 1984, **22**, 328-335 (*cmr*)

Gunatilaka, A.A. et al., *Heterocycles*, 1989, **28**, 999-1005 (*Neisosposinine*)

Rasoanaivo, P. et al., *Fitoterapia*, 2001, **72**, 588-590 (*Dimethoxymitrephylline, Dimethoxyisomitrephylline*)

Caratagine

C-118

[1355-17-5]

Pyrrolizidine alkaloid. Struct. unknown. Alkaloid from *Solenanthes circinata*, *Solenanthes karataginis* and *Rindera oblongifolia* (Boraginaceae). No phys. props. accessible. Conts. halogen, presumably an artifact.

Akrarov, S.T. et al., *Dokl. Akad. Nauk UzSSR*, 1965, **22**, 35-38; *CA*, **63**, 16770f

(Carbamoylmethyl)trimethylammonium(1+), 8CI C-119

2-Amino-N,N,N-trimethyl-2-oxoethanaminium(1+), 9CI. Aminoacetamide trimethylbetaine. **Kalidine**

[16676-65-6]

$\text{Me}_3\text{N}^{\oplus}\text{CH}_2\text{CONH}_2$

$\text{C}_5\text{H}_{13}\text{N}_2\text{O}^{\oplus}$ 117.17

Alkaloid from the aerial parts of *Kalidium gracile* (Chenopodiaceae). Cryst. (Me_2CO aq.) (as chloride). Mp 238-239° (189-192°) (chloride).


▶ BP0715000

Barrass, B.C. *et al.*, *Br. J. Pharmacol.*, 1968, **34**, 345 (*synth*)

Batbayar, N. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 558; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 466 (*cryst struct*)

Carbazole, 9CI C-120

9-Azafluorene. Dibenzopyrrole. Diphenylenimine [86-74-8]



$\text{C}_{12}\text{H}_9\text{N}$ 167.21

Alkaloid from the root bark of *Glycosmis pentaphylla* (Rutaceae). Co-product in manuf. of Anthracene from coal tar. Used in manuf. of dyes. Plates (xylene). Spar. sol. most org. solvs.; sol. hot EtOH. Mp 246.1°. Bp 354.7° Bp₁₄₇ 200°. Sublimes readily. Exhibits practically no basic props. Sol. H_2SO_4 , reprecipitated by H_2O .

▶ LD₅₀ (mus, ipr) 200 mg/kg. FE3150000 [244-56-4, 244-58-6]

Joule, J.A. *et al.*, *Adv. Heterocycl. Chem.*, 1984, **35**, 83 (*rev*)

Hallberg, A. *et al.*, *J. Het. Chem.*, 1984, **21**, 837 (*ms, cmr*)

Chowdhury, B.K. *et al.*, *Phytochemistry*, 1987, **26**, 2138 (*isol*)

Yamaguchi, H. *et al.*, *Spectrochim. Acta A*, 1987, **43**, 1431 (*uv*)

9H-Carbazole-3-carboxaldehyde, 9CI C-121

3-Formylcarbazole. 2-Formylcarbazole (*obsol.*) [51761-07-0]

$\text{C}_{13}\text{H}_9\text{NO}$ 195.22

Alkaloid from the roots of *Clausena lansium* (wampee) and *Murraya euchrestifolia* (Rutaceae). Pale yellow needles (C_6H_6), prisms (Me_2CO /hexane). Mp 153-154°.

Hydrazide:

$\text{C}_{13}\text{H}_{11}\text{N}_3$ 209.25

Prisms. Mp 340°.

2,4-Dinitrophenylhydrazide:

Red prisms. Mp 326-328° dec.

N-Me: [21240-56-2]

$\text{C}_{14}\text{H}_{11}\text{NO}$ 209.247

Solid (hexane/EtOH). Mp 71-73°.

N-Et: [7570-45-8]

$\text{C}_{15}\text{H}_{13}\text{NO}$ 223.274

Mp 85-87°.

N-Et, methylphenylhydrazide: [75232-44-9]

$\text{C}_{22}\text{H}_{21}\text{N}_3$ 327.428

Charge-transporting agent for electrophotographic photoreceptors. Cryst. (EtOAc/EtOH). Mp 128-130°.

N-Et, diphenylhydrazide: [73276-70-7]

$\text{C}_{27}\text{H}_{23}\text{N}_3$ 389.499

Charge-transporting agent for electrophotographic photoreceptors. No phys. props. accessible.

N-Ph: [87220-68-6]

$\text{C}_{19}\text{H}_{13}\text{NO}$ 271.318

Solid (hexane/EtOH). Mp 105-107°.

N-Methoxy: N-Methoxy-3-formylcarbazole

[117592-01-5]

$\text{C}_{14}\text{H}_{11}\text{NO}_2$ 225.246

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Yellow oil. First example of an N-methoxycarbazole alkaloid.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 681A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 163C (*nmr*)

Carter, P.H. *et al.*, *J.C.S.*, 1957, 2210-2215 (*synth*)

Ger. Pat., 1980, 3019395; *CA*, **94**, 156746z (*N-Et hydrazones*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2377 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kawasaki, T. *et al.*, *Heterocycles*, 1990, **31**, 1605 (*synth, deriv*)

Li, W.-S. *et al.*, *Phytochemistry*, 1991, **30**, 343 (*isol, uv, ir, pmr, cmr, ms*)

Mochizuki, H. *et al.*, *Macromolecules*, 2003, **36**, 3457-3464 (*N-Me, N-Ph*)

9H-Carbazole-3-carboxylic acid C-122

2-Carbazolecarboxylic acid (*obsol.*) [51035-17-7]

$\text{C}_{13}\text{H}_9\text{NO}_2$ 211.22

Plates (AcOH). Mp 270-272°.

Me ester: Methyl 3-carbazolecarboxylate.

2-Carbomethoxycarbazole

[97931-41-4]

$\text{C}_{14}\text{H}_{11}\text{NO}_2$ 225.246

Alkaloid from the roots of *Clausena lansium* (wampee) (Rutaceae). Plates (MeOH). Mp 175-177°.

Et ester: 2-Carbethoxycarbazole

[51035-14-4]

$\text{C}_{15}\text{H}_{13}\text{NO}_2$ 239.273

Cryst. solid. Mp 165-168°.

Hydrazide:

$\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}$ 225.249

Prisms (EtOH). Mp 261-263°.

Nitrile: 3-Cyanocarbazole

[57102-93-9]

$\text{C}_{13}\text{H}_8\text{N}_2$ 192.22

Needles (toluene). Mp 184-185°.

N-Me: [89374-79-8]

$\text{C}_{14}\text{H}_{11}\text{NO}_2$ 225.246

Solid (Me_2CO). Mp 256-258°.

N-Me, Me ester: [89369-35-7]

$\text{C}_{15}\text{H}_{13}\text{NO}_2$ 239.273

Solid (MeOH). Mp 130-132°.

Plant, S.G.P. *et al.*, *J.C.S.*, 1934, 1142-1143 (*synth, Et ester*)

Dunlop, H.G. *et al.*, *J.C.S.*, 1939, 1945-1956 (*synth*)

Preston, R.W.G. *et al.*, *J.C.S.*, 1942, 500-504 (*nitrile*)

Carter, P.H. *et al.*, *J.C.S.*, 1957, 2210-2215 (*Me ester, hydrazide*)

Zelent, B. *et al.*, *Can. J. Chem.*, 1982, **60**, 945-956 (*synth, ir, pmr, ms*)

Moody, C.J. *et al.*, *J.C.S. Perkin 1*, 1990, 673-679 (*Et ester, synth, ir, pmr, ms*)


Li, W.-S. *et al.*, *Phytochemistry*, 1991, **30**, 343 (*isol, ester*)

Plekhanova, N.G. *et al.*, *Zh. Org. Khim.*, 2004, **40**, 402-405; *Russ. J. Org. Chem. (Engl. Transl.)*, 2004, **40**, 372-376 (*N-Me, N-Me Me ester*)

Liu, Z. *et al.*, *Tetrahedron*, 2007, **63**, 347-355 (*Me ester, synth*)

9H-Carbazole-2,7-diol, 9CI C-123

2,7-Dihydroxycarbazole



$\text{C}_{12}\text{H}_9\text{NO}_2$ 199.209

Di-Me ether: 2,7-Dimethoxycarbazole, 9CI. Clausine V [61822-18-2]

$\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262

Alkaloid from *Clausena excavata*. Mp 272° (synthetic) Mp 238-230° (natural).


Raj, K. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 371

Zelent, B. *et al.*, *J. Photochem. Photobiol., A*, 1991, **56**, 165

Wu, T.-S. *et al.*, *Phytochemistry*, 1999, **52**, 523-527 (*isol, uv, pmr, ms*)

9H-Carbazole-3-methanol C-124

3-(Hydroxymethyl)-9H-carbazole



$\text{C}_{13}\text{H}_{11}\text{NO}$ 197.236

N-Me, Ac: [94054-92-9]

$\text{C}_{16}\text{H}_{15}\text{NO}_2$ 253.3

Cryst. (hexane). Mp 73°.

N-Et: 3-(Hydroxymethyl)-9-ethyl-9H-carbazole [24301-79-9]

$\text{C}_{15}\text{H}_{15}\text{NO}$ 225.29

Cryst. (EtOH or cyclohexane/ C_6H_6). Mp 87-87.5° (75-76°).

N-Methoxy: 3-(Hydroxymethyl)-9-methoxy-9H-carbazole [142768-49-8]

$\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262


Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

[94054-93-0, 122624-68-4]

Bruck, P. *et al.*, *J.O.C.*, 1970, **35**, 2222 (*derivs*)

- Simionescu, C.I. *et al.*, *J. Polym. Sci., Polym. Chem. Ed.*, 1979, **17**, 2287 (deriv)
 Natansohn, A. *et al.*, *J. Polym. Sci., Polym. Chem. Ed.*, 1984, **22**, 3161 (deriv)
 Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 230 (isol, deriv)

Carbazomadurin A C-125
 [197703-37-0]



$C_{22}H_{27}NO_3$ 353.46
 Prod. by *Actinomadura madurae*. Neuro-
 nal cell protectant. Pale yellow powder.
 Mp 108–110° (dec.). λ_{max} 237 (€ 31700);
 250 (sh) (€ 25800); 289 (sh) (€ 7200); 299
 (€ 9400); 345 (€ 6100); 357 (€ 6600)
 (MeOH).


Kotoda, N. *et al.*, *J. Antibiot.*, 1997, **50**, 770-
 772 (isol, uv, ir, pmr, cmr)
 Knölker, H.J. *et al.*, *Chem. Comm.*, 2003, 1170-
 1171 (synth)

Carbazomadurin B C-126
 [197703-38-1]

As Carbazomadurin A, C-125 with
 R = CH_2CH_3
 $C_{23}H_{29}NO_3$ 367.487
 Prod. by *Actinomadura madurae*. Neuro-
 nal cell protectant. Pale yellow powder.
 Mp 108–110° (dec.). $[\alpha]_D^{24} +4$ (c. 0.05 in
 MeOH). λ_{max} 237 (€ 35700); 249 (sh) (€
 29100); 289 (sh) (€ 7400); 299 (€ 10400);
 347 (€ 6500); 357 (€ 7300) (MeOH).

Kotoda, N. *et al.*, *J. Antibiot.*, 1997, **50**, 770-
 772 (isol, uv, ir, pmr, cmr)


Carbazomarin A C-127
 [184427-83-6]



$C_{33}H_{31}NO_7$ 553.61
 Alkaloid from root bark of *Clausena*
excavata. Yellow powder. Mp 238–239°.
 $[\alpha]_D -27$ (c. 0.0259 in $CHCl_3$). λ_{max} 215
 (log € 4.47); 240 (log € 4.4); 253 (sh) (log €
 4.38); 267 (log € 4.58); 326 (sh) (log €
 4.12); 338 (log € 4.14) (MeOH).

Wu, T.-S. *et al.*, *Tet. Lett.*, 1996, **37**, 7819-7822
 (isol, uv, ir, pmr, cmr, ms, struct)

Carbazomycin G C-128
 1,9-Dihydro-1-hydroxy-3-methoxy-1,2-di-
 methyl-4H-carbazol-4-one, 9CI
 [115920-44-0]




$C_{15}H_{15}NO_3$ 257.288
 Isol. from *Streptovercillium ehimensense*.
 Shows moderate antifungal activity.
 Prisms. Mp 241–243° Mp 266–268°. Ra-
 cemic. λ_{max} 214 (€ 33200); 253 (€ 19800);
 272 (sh) (€ 8600); 278 (€ 7400); 340 (€
 5600) (MeOH) (Derep).

6-Methoxy: **Carbazomycin H**
 [115920-42-8]
 $C_{16}H_{17}NO_4$ 287.315
 From *Streptovercillium ehimensense*.
 Shows no significant biol. activity.
 Prisms. Sol. EtOAc, MeOH; poorly
 sol. H_2O , hexane. Mp 228–230°. Ra-
 cemic. λ_{max} 214 (€ 35700); 253 (€ 23800);
 261 (sh) (€ 20300); 292 (€ 10100); 310
 (sh) (€ 5800); 340 (€ 3500) (MeOH)
 (Derep).

Kaneda, M. *et al.*, *J. Antibiot.*, 1988, **41**, 602-
 608 (isol, pmr, cmr, ir, uv, struct)
 Hagiwara, H. *et al.*, *Tetrahedron*, 2000, **56**,
 5807-5811 (synth)
 Knölker, H.J. *et al.*, *Eur. J. Org. Chem.*, 2003,
 740-746 (synth)

Carbazokinocin C-129
 1-Alkyl-2-methyl-3H-carbazole-3,4(9H)-
 dione. CW 69. WS 63967. Antibiotic CW
 69. Antibiotic WS 63967



Carbazokinocin A - R = $CH_2CH_2CH(CH_3)CH_2CH_3$ (S-)

Carbazokinocin B - R = $(CH_2)_4CH(CH_3)_2$

Carbazokinocin C - R = $(CH_2)_6CH_3$

Carbazokinocin D - R = $(CH_2)_4CH(CH_3)CH_2CH_3$ (S-)

Carbazokinocin E - R = $(CH_2)_5CH(CH_3)_2$

Carbazokinocin F - R = $(CH_2)_6CH(CH_3)_2$

Antibiotic WS 63967A - R = $(CH_2)_4CH_3$

Antibiotic WS 63967C - R = $(CH_2)_5CH_3$

Carbazole antibiotic complex. Prod. by
Streptomyces violaceus and *Streptomyces*
griseochromogenes. Antioxidant and
 GMP phosphodiesterase inhibitor.

Carbazokinocin A
 2-Methyl-1-(3-methylpentyl)-3H-carba-
 zole-3,4(9H)-dione. WS 63967B. Anti-
 biotic WS 63967B. CW 69A. Antibiotic
 CW 69A
 [164322-78-5]

$C_{19}H_{21}NO_2$ 295.38
 Dark green powder. Mp 210–212°. λ_{max}
 240 (€ 13500); 284 (€ 11500); 453 (€ 3500)
 (MeOH/NaOH) (Derep). λ_{max} 228 (€
 14200); 264 (€ 11200); 398 (€ 2700)
 (MeOH) (Derep).

Carbazokinocin B
 2-Methyl-1-(5-methylhexyl)-3H-carba-
 zole-3,4(9H)-dione. WS 63967D. Anti-
 biotic WS 63967D. CW 69B. Antibiotic
 CW 69B
 [155233-26-4]

$C_{20}H_{23}NO_2$ 309.407
 Dark green powder. Mp 213–217°. λ_{max}
 240 (€ 13500); 284 (€ 11500); 453 (€ 3500)
 (MeOH/NaOH) (Derep). λ_{max} 228 (€
 14200); 264 (€ 11200); 398 (€ 2700)
 (MeOH) (Derep).

Carbazokinocin C
 1-Heptyl-2-methyl-3H-carbazole-
 3,4(9H)-dione. WS 63967E. Antibiotic
 WS 63967E. CW 69C. Antibiotic CW
 69C
 [155233-27-5]

$C_{20}H_{23}NO_2$ 309.407
 Dark green powder. Mp 227–228° (210-
 212°). λ_{max} 240 (€ 13500); 284 (€ 11500);
 453 (€ 3500) (MeOH/NaOH) (Derep).
 λ_{max} 228 (€ 14200); 264 (€ 11200); 398 (€
 2700) (MeOH) (Derep).

Carbazokinocin D
 2-Methyl-1-(5-methylheptyl)-3H-carba-
 zole-3,4(9H)-dione. WS 63967F. Anti-
 biotic WS 63967F. CW 69D. Antibiotic CW
 69D
 [155233-28-6]

$C_{21}H_{25}NO_2$ 323.434
 Dark green powder. Mp 208–210°. λ_{max}
 240 (€ 12100); 284 (€ 10700); 453 (€ 3200)
 (MeOH/NaOH) (Derep). λ_{max} 228 (€
 12700); 264 (€ 10600); 398 (€ 2400)
 (MeOH) (Derep).

Carbazokinocin E
 2-Methyl-1-(6-methylheptyl)-3H-carba-
 zole-3,4(9H)-dione. CW 69E. Antibiotic
 CW 69E
 [164177-49-5]

$C_{21}H_{25}NO_2$ 323.434
 Dark green powder. Mp 209–210°. λ_{max}
 240 (€ 12100); 284 (€ 10700); 453 (€ 3200)
 (MeOH/NaOH) (Derep). λ_{max} 228 (€
 12700); 264 (€ 10600); 398 (€ 2400)
 (MeOH) (Derep).

Carbazokinocin F
 2-Methyl-1-(7-methyloctyl)-3H-carba-
 zole-3,4(9H)-dione. CW 69F. Antibiotic
 CW 69F
 [164177-50-8]

$C_{22}H_{27}NO_2$ 337.461
 Dark green powder. Mp 208–210°. λ_{max}
 240 (€ 12100); 284 (€ 10700); 453 (€ 3200)
 (MeOH/NaOH) (Derep). λ_{max} 228 (€
 12700); 264 (€ 10600); 398 (€ 2400)
 (MeOH) (Derep).

Antibiotic WS 63967A
 2-Methyl-1-pentyl-3H-carbazole-
 3,4(9H)-dione. WS 63967A
 [155233-23-1]

C₁₈H₁₉NO₂ 281.354

Antibiotic WS 63967C

1-Hexyl-2-methyl-3H-carbazole-3,4(9H)-dione. *WS 63967C*
[155233-25-3]


C₁₉H₂₁NO₂ 295.38

[155233-24-2]

Japan. Pat., 1993, 93 301 857; *CA*, **121**, 7425T (Carbazquinocins, *isol. struct*)
Tanaka, M. *et al.*, *J. Antibiot.*, 1995, **48**, 326-328 (Carbazquinocins, *isol. uv, ir, pmr, cmr*)
Choshi, T. *et al.*, *J.O.C.*, 1997, **62**, 2535-2543 (Carbazquinocins C,D,E,F, *synth, ir, pmr, cmr*)
Knoelker, H.-J. *et al.*, *Synthesis*, 2002, 557-564 (Carbazquinocin C, *synth, ir, uv, pmr, cmr, ms*)

β-Carboline C-130

9H-Pyrido[3,4-b]indole, 9CI, 2,9-Diazafluorene. *Norharman*. 2-Carboline. *Carbazoline*†
[244-63-3]



C₁₁H₈N₂ 168.198

Constit. of espresso coffee. Alkaloid from *Chrysophyllum lacourtianum*, *Nocardia* sp., *Catharanthus roseus* leaves, *Lolium perenne*, *Strychnos johnsonii*, *Strychnos potatorum*, and *Festuca arundinacea*. Prod. by *Streptomyces* sp., a metab. from the New Zealand ascidian *Ritterella sigillinoides*. Also a trace constit. of the fluorescent dinoflagellate *Noctiluca miliaris*. Plant growth and enzyme inhibitor. Potentiator of benzo[a]pyrene induced mutagenicity, antiparasitic, antirypanosomal agent. A comutagen with aromatic amines active towards bacteria. Needles (EtOH). Sol. hot H₂O; spar. sol. C₆H₆, petrol. Mp 198.5°. Blue fluor. in dil. acid soln. λ_{max} 235 (ε 31000); 290 ; 350 (ε 4600) (MeOH) (Berdy).

► Exp. nephrotoxic at high doses. A comutagen. UU9350000

N²-Oxide: [24223-07-2]

C₁₁H₈N₂O 184.197
Mp 256°.

N²-Me: 2-Methyl-9H-pyrido[3,4-b]indolium. *Normelinonine F*

[17994-14-8]
[5667-11-8 , 21951-87-1]
C₁₂H₁₁N₂⁺ 183.232

Quaternary alkaloid from *Strychnos usambarensis* (Loganiaceae). Antimitotic agent. Poorly sol. hexane.

3,4-Dihydro, 9-Ac: 9-Acetyl-4,9-dihydro-3H-pyrido[3,4-b]indole, 9CI, 9-Acetyl-3,4-dihydro-β-carboline

[76528-83-1]
C₁₃H₁₂N₂O 212.251

Alkaloid from the roots of *Adhatoda vasica* (Acanthaceae). Mp 165-166°
Mp 187-188° (as hydrochloride).


1,2,3,4-Tetrahydro, 2-Me: see 1,2,3,4-Tetrahydro-2-methyl-β-carboline, T-203

[244-66-6]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 681D (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 164C (*nmr*)
Kermack, W.O. *et al.*, *J.C.S.*, 1921, **119**, 1602-1642 (*synth*)
Späth, E. *et al.*, *Ber.*, 1940, **73**, 719-723 (*synth*)
Pruckner, F. *et al.*, *Annalen*, 1943, **554**, 127-144 (*uv*)
Poindexter, E.H. *et al.*, *Phytochemistry*, 1961, **1**, 215 (*isol, uv, ir*)
Coutts, R.T. *et al.*, *Org. Mass Spectrom.*, 1970, **3**, 879 (*ms*)
Balkau, F. *et al.*, *Aust. J. Chem.*, 1973, **26**, 1501 (*pmr*)
Bush, L.P. *et al.*, *J. Chromatogr.*, 1975, **111**, 165 (*isol*)
Deumie, M. *et al.*, *J. Photochem.*, 1979, **10**, 365 (*uv*)
Inoue, S. *et al.*, *Chem. Lett.*, 1980, 297-298 (*isol, dinoflagellate*)
Bobbitt, J.M. *et al.*, *J.O.C.*, 1980, **45**, 1978 (*synth*)
Jain, M.P. *et al.*, *Phytochemistry*, 1980, **19**, 1880 (9-Ac-3,4-dihydro)
Sugimura, T. *et al.*, *Biol. React. Intermed. 2, Chem. Mech. Biol. Eff.*, Snyder, R. *et al* (ed.), (see *Adv. Exp. Med. Biol.* v136A-136B), Plenum Press, 1982, 1011 (*comutagenicity*)
Caprasse, M. *et al.*, *J. Pharm. Belg.*, 1983, **38**, 135-139 (*Normelinonine F*)
Nakagawa, M. *et al.*, *Tet. Lett.*, 1983, **24**, 2171 (*synth*)
Hiemstra, H.C. *et al.*, *Tetrahedron*, 1983, **39**, 3981 (*synth*)
Becalski, A. *et al.*, *Acta Pol. Pharm.*, 1984, **41**, 601 (*synth*)
Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 471 (*cmr*)
Atta-ur-Rahman, *et al.*, *Planta Med.*, 1985, 287 (*isol, uv, ir, pmr, cmr, ms*)
Rommelspacher, H. *et al.*, *Prog. Drug Res.*, 1985, **29**, 415 (*rev*)
Yomosa, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 921 (*isol, bibl*)
Lake, R.J. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1201 (*isol, uv, ir, pmr, cmr*)
Hagiwara, A. *et al.*, *Toxicol. Pathol.*, 1992, **20**, 197 (*tox*)
Rocca, P. *et al.*, *Tetrahedron*, 1993, **49**, 49 (*synth*)
Nakano, K. *et al.*, *Mutat. Res.*, 2000, **470**, 141-146 (*comutagenicity, pmr*)
Kast, O. *et al.*, *Synth. Commun.*, 2003, **33**, 3843-3850 (N²-oxide)
Alves, R.C. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 1832-1838 (*isol*)
Huang, W. *et al.*, *Synth. Commun.*, 2007, **37**, 2137-2143 (*synth, ir, pmr*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, NNR300

β-Carboline-1-carboxaldehyde C-131

9H-Pyrido[3,4-b]indole-1-carboxaldehyde, 9CI, 1-Formyl-β-carboline. *Kumujian C*
[20127-63-3]



C₁₂H₈N₂O 196.208


Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Yellow

needles (Me₂CO). Mp 202° (198-200°).

Jordaan, A. *et al.*, *J. S. Afr. Chem. Inst.*, 1968, **21**, 22 (*synth*)
Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 471 (*cmr*)
Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3579 (*isol, uv, ir, synth*)

β-Carboline-1-carboxylic acid C-132

9H-Pyrido[3,4-b]indole-1-carboxylic acid, 9CI
[26052-96-0]



C₁₂H₈N₂O₂ 212.207

Bright yellow cryst. Insol. most solvs. Mp 272° dec. (235°).

Me ester: 1-Methoxycarbonyl-β-carboline. 1-Carbomethoxy-β-carboline

[3464-66-2]
C₁₃H₁₀N₂O₂ 226.234

Isol. from stem bark of *Pleiocarpa mutica* and from *Nauclea diderrichii*, *Ailanthus malabarica*, *Ailanthus altissima*, *Aeschron crenata* (preferred genus name *Picrasma*), *Commelina communis* and *Picrasma* spp. (e.g. *Picrasma quassioides*) (Apocynaceae, Rubiaceae, Simaroubaceae). Active against gram-positive bacteria. Needles by subl. Mp 165-168° (160-161°). λ_{max} 246 (ε 14800); 256 (ε 18200); 257 (ε 18200); 275 (ε 20420); 301 (ε 11750); 370 (ε 6760) (EtOH).

Et ester: 1-Ethoxycarbonyl-β-carboline. 1-Carbethoxy-β-carboline. Kumujian A

[72755-19-2]
C₁₄H₁₂N₂O₂ 240.261

Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Needles (Me₂CO). Mp 123°. λ_{max} 218 (log ε 4.9); 247 (log ε 4.52); 258 (log ε 4.52); 288 (log ε 4.56); 302 (log ε 4.31); 368 (log ε 4.06) (EtOH).

Butyl ester: 1-Butoxycarbonyl-β-carboline. 1-Carbobutoxy-β-carboline

[153535-98-9]
C₁₆H₁₆N₂O₂ 268.315

Alkaloid from *Polygala tenuifolia*. Component of Yuan Zhi. Mp 95°.

Amide: 1-Carbamoyl-β-carboline. β-Carboline-1-carboxamide. Harman carboxamide

[38940-60-2]
C₁₂H₉N₃O 211.223

Isol. from *Nauclea diderrichii* and *Ailanthus malabarica* bark and roots (Simaroubaceae). Also present in root bark of *Strychnos potatorum* (Loganiaceae). Pale yellow needles (CH₂Cl₂). Mp 230° (224-226°). Erroneously claimed to be novel by Massiot *et al.* λ_{max} 214 (ε 36000); 240 (sh) (ε 17400); 244 (ε 18200); 250 (sh) (ε 17000); 270 (ε 17800); 298 (ε 10000); 366 (ε 5900) (no solvent reported).

Nitrile: 1-Cyano-β-carboline

[79960-43-3]
C₁₂H₇N₃ 193.207
Yellow needles (toluene/CHCl₃). Mp
230-232°.

- Snyder, H.R. *et al.*, *J.A.C.S.*, 1949, **71**, 527-529
(*synth*)
Kump, C. *et al.*, *Helv. Chim. Acta*, 1963, **46**,
498-505 (*Me ester, synth, uv*)
Achenbach, H. *et al.*, *J.A.C.S.*, 1965, **87**, 4177-
4181 (*Me ester, isol, ir, uv, ms, pmr*)
Sánchez, E. *et al.*, *Phytochemistry*, 1971, **10**,
2155-2159 (*Me ester, isol*)
McLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**,
1478-1485 (*Me ester, amide, isol*)
Kondo, Y. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**,
837-839 (*Me ester, isol, uv, pmr*)
Joshi, B.S. *et al.*, *Heterocycles*, 1977, **7**, 193-200
(*amide, ester*)
Yang, J.S. *et al.*, *Yaoxue Xuebao*, 1979, **14**, 167-
177 (*Me ester, activity*)
Yang, J.-S. *et al.*, *CA*, 1980, **92**, 72679a (*isol*)
Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984,
32, 3579-3583 (*esters, isol, uv, ir, synth*)
Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**,
471-473 (*cmr*)
Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**,
3378-3387 (*nitrile, synth*)
Massiot, G. *et al.*, *Phytochemistry*, 1992, **31**,
2873-2876 (*amide*)
Jin, B. *et al.*, *CA*, 1994, **120**, 240067s (*butyl
ester, isol*)
Panosyan, F.B. *et al.*, *Can. J. Chem.*, 2001, **79**,
1110-1114 (*Me ester, synth, ir, pmr*)
Kast, O. *et al.*, *Synth. Commun.*, 2003, **33**,
3843-3850 (*amide, nitrile*)

β-Carboline-3-carboxylic acid C-133

9H-Pyrido[3,4-b]indole-3-carboxylic
acid, 9CI. *Norharman*carboxylic acid
[74214-63-4]

C₁₂H₈N₂O₂ 212.207
Esters are benzodiazepine tranquilliser
antagonists. Lemon-yellow plates
(AcOH) or pale buff powder. Mp 309-
310° dec. (303-305°). Gives blue fluores-
cence in dilute mineral acids.

Monohydrate: Mp 275-283° (250°).

Me ester: 3-Methoxycarbonyl-β-carboline
[69954-48-9]
C₁₃H₁₀N₂O₂ 226.234
Prod. by *Streptomyces* sp. Ma5373.
Diazepam antagonist, tranquilliser.
Cryst. (MeCN). Mp 261-262° (248-
249°). Mp highly dependent on rate of
heating.

Et ester: 3-Ethoxycarbonyl-β-carboline

[74214-62-3]
C₁₄H₁₂N₂O₂ 240.261
Isol. from human urine and brain
tissue prob. as artifact. Alkaloid from
Picrasma quassioides (Simaroubaceae).
Cryst. (EtOAc/hexane) or yellow need-
les (MeOH). Mp 229-230° (222-224°).
λ_{max} 215 ; 242 ; 279 (no solvent
reported).

Propyl ester: [76808-18-9]

C₁₅H₁₄N₂O₂ 254.288
Cryst. (MeCN). Mp 187-188°.

Amide: [88932-13-2]

C₁₂H₉N₃O 211.223
Cryst. (MeOH). Mp 320° dec. (318-
319°).

Methylamide: *N*-Methyl-β-carboline-3-
carboxamide. β-CCA methylamide. FG
7142

[78538-74-6]
C₁₃H₁₁N₃O 225.249
Formed naturally during the combus-
tion process of many plant leaves, most
notably tobacco. Component of cigar-
ette smoke. Benzodiazepine partial
inverse receptor agonist. Anxiogenic
agent. Anorectic. Mp 265° dec.

Ethylamide: [78538-80-4]

C₁₄H₁₃N₃O 239.276
Mp 247-249° dec.

2-Me: 3-Carboxy-2-methyl-9H-pyri-
do[3,4-b]indolinium inner salt. 3-Car-
boxy-2-methyl-β-carbolinium
[228091-07-4]

C₁₃H₁₀N₂O₂ 226.234
Quaternary alkaloid from the soft
coral *Lignopsis spongiosum*. Pale yel-
low solid (MeOH). Mp 203-205°. λ_{max}
210 (log ε 4.16); 236 (log ε 4.05); 264
(log ε 4.26); 308 (log ε 3.88); 384 (log ε
3.45) (MeOH).

9-Me, *Me ester*: [82596-95-0]

C₁₄H₁₂N₂O₂ 240.261
Cryst. (MeOH). Mp 200-201°.

Tetrahydro: see 2,3,4,9-Tetrahydro-1H-
pyrido[3,4-b]indole-3-carboxylic acid,
T-224

1-Methoxy: 1-Methoxy-β-carboline-3-
carboxylic acid. *Taraxacine B*

C₁₃H₁₀N₂O₃ 242.234
Alkaloid from the aerial parts of
Taraxacum formosanum. Pale yellow
symp. λ_{max} 216 (log ε 4.3); 232 (log ε
4.3); 270 (log ε 4.4); 287 (sh) (log ε 4);
303 (log ε 3.8); 333 (log ε 3.5); 346 (log
ε 3.5) (MeOH).

Aldrich Library of FT-IR Spectra, 1st edn.,
1985, **2**, 682D (*ir*)

*Aldrich Library of 13C and 1H FT NMR
Spectra*, 1992, **3**, 167A; 167B (*nmr*)

King, H. *et al.*, *J.C.S.*, 1937, 466-473 (*synth*)
Braestrup, C. *et al.*, *Proc. Natl. Acad. Sci.
U.S.A.*, 1980, **77**, 2288-2292 (*isol, ms, pmr,
cmr*)

Petersen, E.N. *et al.*, *Eur. J. Pharmacol.*, 1982,
82, 217-221 (*methylamide, pharmacol*)

Cain, M. *et al.*, *J. Med. Chem.*, 1982, **25**, 1081-
1091 (*9-Me Me ester*)

Lippke, K.P. *et al.*, *J. Med. Chem.*, 1983, **26**,
499-503 (*synth, propyl ester, pharmacol*)

Coutts, R.T. *et al.*, *Heterocycles*, 1984, **22**, 131-
142 (*amides*)

Moody, C.J. *et al.*, *J.C.S. Perkin I*, 1984, 2895-
2901 (*synth*)

Plate, R. *et al.*, *J.O.C.*, 1986, **51**, 309-314
(*synth*)

Maclaren, J.A. *et al.*, *Aust. J. Chem.*, 1989, **42**,
813-821 (*synth, bibl*)

Koike, K. *et al.*, *Phytochemistry*, 1990, **29**,
3060-3061 (*Et ester*)

Dekhane, M. *et al.*, *Tetrahedron*, 1994, **50**,
6299-6306 (*synth*)

Cabrera, G.M. *et al.*, *J. Nat. Prod.*, 1999, **62**,
759-760 (*3-Carboxy-2-methyl-β-
carbolinium*)


Kubicki, M. *et al.*, *Acta Cryst. C*, 2001, **57**,
728-729 (*Me ester, cryst struct*)

Leu, Y.-L. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**,
599-601 (*Taraxacine B*)

Evans, A.K. *et al.*, *CNS Drug Rev.*, 2007, **13**,
475-501 (*methylamide, rev. pharmacol*)

β-Carboline-1-propanoic acid C-134

9H-Pyrido[3,4-b]indole-1-propanoic
acid, 9CI
[89915-39-9]



C₁₄H₁₂N₂O₂ 240.261

Alkaloid from the wood of *Picrasma
quassioides*. Isol. from the mushrooms
Boletus curtisii and *Cortinarius infractus*.
Needles (CHCl₃/MeOH). Mp 215°. λ_{max}
212 (log ε 4.24); 236 (log ε 4.47); 250 (sh)
(log ε 4.34); 282 (sh) (log ε 3.92); 288 (log
ε 4.1); 302 (sh) (log ε 3.64); 338 (log ε
3.57); 350 (log ε 3.6) (MeOH).

Me ester: *Infractine*. *Kumujanrine*

[91147-07-8]

C₁₅H₁₄N₂O₂ 254.288

Alkaloid from the fruiting bodies of
the gilled agaric *Cortinarius infractus*
and from the wood of *Picrasma
quassioides*. Cryst. (toluene). Mp 145-
146°.

Et ester: *Ethyl β-carboline-1-propionate*

[90686-24-1]

C₁₆H₁₆N₂O₂ 268.315

Alkaloid from the roots of *Hannoa
klaianeana* (Simaroubaceae). Whitish-
yellow needles (EtOH or C₆H₆/petrol).
Mp 125-126°.

Et ester, 2N-oxide: *Ethyl β-carboline-2N-
oxide-1-propionate*

[90686-25-2]

C₁₆H₁₆N₂O₃ 284.314

Alkaloid from the roots of *Hannoa
klaianeana* (Simaroubaceae). Whitish-
yellow needles (EtOH).

Pentyl ester: *Pentyl β-carboline-1-propio-
nate*

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from the roots of *Eurycoma
longifolia*. Yellow powder (C₆H₆). Mp
127° dec. λ_{max} 235 (log ε 3.9); 247 (log
ε 3.85); 289 (log ε 3.6); 303 (log ε 3.5);
335 (log ε 3.07); 350 (log ε 3.08); 376
(log ε 2.93) (MeOH).

N-Me:

C₁₅H₁₄N₂O₂ 254.288

Alkaloid from the mushroom *Boletus
curtisii*. Straw-yellow solid. Mp > 340°.
Zwitterionic. λ_{max} 254 (log ε 4.46); 308
(log ε 4.29); 374 (log ε 3.67) (MeOH).

6-Hydroxy, *Me ester*: 6-Hydroxyinfractine
[91147-08-9]

C₁₅H₁₄N₂O₃ 270.287

Alkaloid from the fruiting bodies of
the gilled agaric *Cortinarius infractus*.
Yellow cryst. +1H₂O (MeOH aq.). Mp
130° Mp 310-320° (double Mp).

7-Hydroxy: 7-Hydroxy-β-carboline-1-
propanoic acid

C₁₄H₁₂N₂O₃ 256.26

Alkaloid from *Eurycoma harmandiana*
and *Fontinalis squamosa*. Microcryst.
powder. Mp 245° dec.

6-Methoxy: 6-Methoxy-β-carboline-1-
propanoic acid. *Aervolanine*. *Ervolanine*

[139742-35-1]
C₁₅H₁₄N₂O₃ 270.287

Alkaloid from aerial parts of *Aerva lanata* (Amaranthaceae). Yellow needles (EtOH). Mp 194-196°.

7-Methoxy-7-Methoxy-β-carboline-1-propanoic acid

[137756-13-9]
C₁₅H₁₄N₂O₃ 270.287

Alkaloid from the roots of *Eurycoma longifolia* (Simaroubaceae). Shows antimalarial activity. Needles. Mp 160-162°.

Haynes, H.F. *et al.*, *Aust. J. Sci. Res. Ser. A*, 1952, **5**, 387 (synth)

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3579 (isol, uv, ir, synth)

Yang, J. *et al.*, *Huaxue Xuebao*, 1984, **42**, 679; *CA*, **101**, 207605a (isol, *Infrafractine*)

Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **21**, 471 (cmr)

Lumonadio, L. *et al.*, *Phytochemistry*, 1984, **23**, 453 (isol, uv, ir, pmr, ms, *struct*, *derivs*)

Steglich, W. *et al.*, *Tet. Lett.*, 1984, **25**, 2341 (isol, uv, ir, pmr, cmr, ms, *Infrafractine*, 6-Hydroxyinfractine)

Kardono, L.B.S. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1360 (7-Methoxy-β-carboline-1-propanoic acid)

Zapesochayna, G. *et al.*, *Planta Med.*, 1992, **58**, 192 (6-Methoxy-β-carboline-1-propanoic acid)

Nowak, W. *et al.*, *Annalen*, 1993, 153 (synth, *Infrafractine*, 6-Hydroxyinfractine)

Bracher, F. *et al.*, *Pharmazie*, 1995, **50**, 182 (synth)

Salm, R.F. *et al.*, *Phytochemistry*, 1998, **49**, 887-892 (7-Hydroxy-β-carboline-1-propanoic acid)


Kanchanapoom, T. *et al.*, *Phytochemistry*, 2001, **56**, 383-386 (7-Hydroxy-β-carboline-1-propanoic acid)

Kuo, P.C. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1324-1327 (pentyl ester)

Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, pmr, cmr, ms)

β-Carboline-1,3,4-trione C-135

9H-Pyrido[3,4-b]indole-1,3,4(2H,9H)-trione, 9CI, 8CI
[16641-79-5]



C₁₁H₆N₂O₃ 214.18

Alkaloid from *Picrasma quassioides* (Simaroubaceae). Yellow prisms (MeOH aq.). Mp 300°.

8-Methoxy-8-Methoxy-β-carboline-1,3,4-trione. 8-Methoxy-9H-pyrido[3,4-b]carbazole-1,3,4(2H,9H)-trione, 9CI. Picrasidine V
[131653-92-4]

C₁₂H₈N₂O₄ 244.206


Alkaloid from *Picrasma quassioides* (Simaroubaceae). Orange prisms (MeOH aq.). Mp 300°.

Member, P.A. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 249; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 208 (synth)

Koike, K. *et al.*, *Phytochemistry*, 1990, **29**, 3060 (isol, *struct*, *Picrasidine V*)

1-(β-Carbolin-1-yl)-3-hydroxy-1-propanone C-136

3-Hydroxy-1-(9H-pyrido[3,4-b]indol-1-yl)-1-propanone. 1-(3-Hydroxypropionyl)-β-carboline




C₁₄H₁₂N₂O₂ 240.261

Prod. by the marine bacterium strain Bio215. Yellowish-green solid.

Shaaban, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (isol, pmr, ms)

4-(β-Carbolin-1-yl)-4-oxobutanoic acid C-137

4-Oxo-4-(9H-pyrido[3,4-b]indol-1-yl)butanoic acid



C₁₅H₁₂N₂O₃ 268.271


Alkaloid from the roots of *Anemone altaica*. Yellow powder. Mp 234-236°. λ_{max} 220 ; 283 ; 310 ; 380 (no solvent reported).

Zou, Z.-J. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 1484-1486 (isol, pmr, cmr)

Zou, Z.-J. *et al.*, *J. Integ. Plant Biol.*, 2005, **47**, 1145-1147 (isol pmr, cmr, ms)

1-(β-Carbolin-1-yl)-3,4,5-trihydroxy-1-pentanone C-138

1-(3,4,5-Trihydroxypentanoyl)-β-carboline. **Alkaloid D†**
[180995-40-8]



C₁₆H₁₆N₂O₄ 300.313

Alkaloid from cultured hybrid cells of *Rauwolfia serpentina* x *Rhazya stricta*. Amorph. powder. [α]_D²³ -39 (c, 0.048 in MeOH). Not named in the paper. λ_{max} 217 ; 234 (sh) ; 243 ; 251 ; 260 ; 284 ; 307 ; 380 (MeOH).


Aimi, N. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1637 (isol, uv, pmr, cmr, ms, *struct*)

Kitajima, M. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2195 (synth, abs config)

Carbonarin C

[177912-42-4]

C-139



C₃₃H₂₇NO₈ 565.578


Prod. by *Aspergillus carbonarius*. Insecticidal agent. λ_{max} 228 (ε 17400); 280 (ε 18500); 410 (ε 2900) (MeOH) (Berdy).

U.S. Pat., 1996, 5 519 052; *CA*, **125**, 28297a (isol, uv, pmr, cmr)

Carbonarin D

[177912-43-5]

C-140



C₃₃H₂₇NO₈ 565.578


Prod. by *Aspergillus carbonarius*. Insecticidal agent. λ_{max} 244 (ε 17400); 285 (ε 11500); 380 (ε 28400) (MeOH) (Berdy).

U.S. Pat., 1996, 5 519 052; *CA*, **125**, 28297a (isol, uv, ir, pmr)

Carbonarin E

1,5-Dihydro-4-[(4-hydroxy-3-methoxyphenyl)methyl]-3-(4-hydroxyphenyl)-2H-pyrrol-2-one, 9CI
[177912-44-6]

C-141



C₁₈H₁₇NO₄ 311.337


Prod. by *Aspergillus carbonarius*. λ_{max} 225 (ε 10200); 280 (ε 5800); 371 (ε 1100) (MeOH) (Berdy).

U.S. Pat., 1996, 5 519 052; *CA*, **125**, 28297a

Carbonarone B

C-142

4-Hydroxy-2-oxo-6-(phenylmethyl)-1H-pyridine-3-carboxaldehyde. 6-Benzyl-1,2-dihydro-4-hydroxy-2-oxo-3-pyridinecarboxaldehyde. 6-Benzyl-3-formyl-4-hydroxy-2-pyridone. 6-Benzyl-2,4-dihydroxy-3-pyridinecarboxaldehyde
[937182-86-0]



$C_{13}H_{11}NO_3$ 229.235


Prod. by the marine-derived *Aspergillus carbonarius* WZ-4-11. Cytotoxic. Needles. Mp 209-209.5°. λ_{max} 340 (log ϵ 3.87) (MeOH).

Zhang, Y. *et al.*, *J. Antibiot.*, 2007, **60**, 153-157 (isol, pmr, cmr, cryst struct)

11-Carboxyanatoxin a

C-143

β -Oxo-9-azabicyclo[4.2.1]non-2-ene-2-propanoic acid
[923925-57-9]



Absolute Configuration

$C_{11}H_{15}NO_3$ 209.244


Isol. from *Aphanizomenon issatschenkoi* (CAWBG02).

Selwood, A.I. *et al.*, *Environ. Sci. Technol.*, 2007, **41**, 506-510

3-Carboxy-1-(carboxymethyl)pyridinium betaine

C-144

1-Carboxymethylnicotinic acid
[137591-89-0]
[148717-06-0, 148717-05-9]



$C_8H_7NO_4$ 181.148

Alkaloid from the sponge *Anthosigmella* cf. *raromicrosclera*. Cysteine protease inhibitor. λ_{max} 267 (ϵ 5000) (H_2O).

Chen, X.-M. *et al.*, *J. Mol. Struct.*, 1991, **249**, 135-140 (synth, cryst struct)


Chen, X.-M. *et al.*, *J. Crystallogr. Spectrosc. Res.*, 1993, **23**, 291-296 (cryst struct)

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 671-672 (isol, synth, uv, pmr, cmr)

5-Carboxycorynanthine

C-145

Methyl 17-hydroxy-yohimban-5,16-dicarboxylate, 9CI



Absolute Configuration

$C_{22}H_{26}N_2O_5$ 398.458

5 β -form [55624-07-2]

Alkaloid from *Adina rubescens*. Incorr. descr. in the paper as 5 α -.


Me ester: Mp 235-237°. $[\alpha]_D^{25}$ -29 (CHCl₃).

Brown, R.T. *et al.*, *Tet. Lett.*, 1974, **15**, 3429-3430 (isol, ms, pmr, cd, struct)

3-(2-Carboxyethyl)-1,4-dimethylpyridinium, 9CI

C-146

Sulcatine
[264870-65-7]



$C_{10}H_{13}NO_2$ 179.218

Zwitterionic. Isol. from the ascidian *Microcosmus vulgaris*. Shows antiproliferative activity. Amorph. solid.

Aiello, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 517-519 (isol, pmr, cmr)

3-(Carboxymethylamino)propanoic acid

C-147

N-(Carboxymethyl)- β -alanine, 9CI. N-(Carboxyethyl)glycine. Iminopropionacetic acid
[505-72-6]

$HOO^1CCH_2CH_2NHCH_2COOH$

$C_5H_9NO_4$ 147.13

Isol. from *Phaseolus radiatus* var. *typicus*. Cryst. (H_2O or EtOH aq.). Mp 195-198° (191-192°) dec.

Hydrochloride: Mp 120-122°.

Di-Et ester: [3783-61-7]

$C_9H_{17}NO_4$ 203.238

Oil. Bp 253° sl. dec. Bp₁ 98-99°.

Di-Et ester, hydrochloride: Mp 85-87°.

1-Nitrile: N-(2-Cyanoethyl)glycine, 9CI.

3-(Carboxymethylamino)propionitrile
[3088-42-4]

$C_5H_8N_2O_2$ 128.13

Cryst. Mp 193-195°.

► MB9275000

Cocker, W. *et al.*, *J.C.S.*, 1952, 1182 (deriv)

McKinney, L.L. *et al.*, *J.A.C.S.*, 1952, **74**, 1942; 5183 (synth)

Coburn, M.D. *et al.*, *J. Het. Chem.*, 1965, **2**, 308 (deriv)

Kasai, T. *et al.*, *Agric. Biol. Chem.*, 1971, **35**, 1603 (isol, synth)

Kawashiro, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 1097; 2871 (synth, ms)


2-Carboxy-1-methylpyridinium betaine

C-148

2-Carboxy-1-methylpyridinium hydroxide inner salt, 9CI. N-Methylpicolinic betaine.

Homarine

[445-30-7]



$C_7H_7NO_2$ 137.138

Alkaloid from fungus *Polyporus sulphureus*, lobsters *Homarus vulgaris* and *Homarus americanus*, Antarctic soft coral *Gersemia antarctica*, Antarctic gastropod *Marseniopsis mollis*, red alga *Pterocladia capillacea* and other spp. Shows antibacterial activity. Feeding deterrent in marine organisms. Cryst. (EtOH). Dec. on heating, deliquescent.

Hydrochloride: Mp 170-175° dec.

Picrate: Mp 158-160°.

Hantzsch, A. *et al.*, *Ber.*, 1886, **19**, 31 (synth)

Hoppe-Seyler, F. *et al.*, *Hoppe-Seyler's Z.*

Physiol. Chem., 1933, **222**, 105-115 (isol)

Green, R.W. *et al.*, *J.A.C.S.*, 1956, **78**, 4896 (uv, tautom)

Leonard, G.J. *et al.*, *Nature (London)*, 1963, **200**, 78 (isol)

Groenmeberg, T. *et al.*, *Chem. Scr.*, 1972, **2**, 17 (ms)

Sciuto, S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 322-325 (isol, red alga)

McClintock, J.B. *et al.*, *J. Chem. Ecol.*, 1994, **20**, 2539-2549 (isol, activity)

Slattery, M. *et al.*, *Marine Ecol.: Progr. Ser.*, 1997, **161**, 133-144; *CA*, **128**, 119209b (isol, activity)

Polychronopoulos, P. *et al.*, *Nat. Prod. Lett.*, 2001, **15**, 411-418 (*Homarine*, pmr, cmr, ms)


3-Carboxy-1-methylpyridinium betaine

C-149

3-Carboxy-1-methylpyridinium hydroxide inner salt. N-Methylnicotinic betaine.

Trigonelline. *Gynesis*. *Coffearin*

[535-83-1]



$C_7H_7NO_2$ 137.138

Alkaloid from fenugreek (*Trigonella foenum-graecum*) (Fabaceae), *Schumannio-phyton magnificum* (Rubiaceae), *Mappia foetida*, *Strophanthus* sp., *Cannabis sativa* and very many other spp.; also present in coffee beans, in many animals, the sponge *Calyx nicaeensis*, the soft coral *Gersemia antarctica*, and the red seaweed *Pterocladia capillacea*. Prisms + 1H₂O (EtOH)

aq.). V. sol. H₂O; sol. MeOH, EtOH; poorly sol. CHCl₃, Et₂O. Mp 218° dec. (anhyd.).

▶ LD₅₀ (rat, orl) 5000 mg/kg. YF6825000
Hydrochloride: [6138-41-6]

Cryst. (EtOH aq.). Mp 257-258° (250°, 245-250° dec.).

Picrate: Mp 205-206° (198-200°).

Gorter, K. *et al.*, *Annalen*, 1910, **372**, 237-246 (*isol*)

Späth, E. *et al.*, *Ber.*, 1944, **77**, 362-369 (*synth*)
Ackermann, D. *et al.*, *Hoppe-Seyler's Z.*

Physiol. Chem., 1961, **326**, 197-199 (*occur*)

Wang, S.Y. *et al.*, *Biochemistry*, 1968, **7**, 3740-3744 (*uv*)

Wehrli, F.W. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 229-243 (*pmr*, *cmr*)

Ghosal, S. *et al.*, *Planta Med.*, 1973, **23**, 321-329 (*isol*, *uv*, *ir*, *pmr*)

Houghton, P.J. *et al.*, *Planta Med.*, 1987, **53**, 262-264 (*isol*)

Slattery, M. *et al.*, *Marine Ecol.: Progr. Ser.*, 1997, **161**, 133-144; *CA*, **128**, 119209e (*isol*, *activity*)


Merck Index, 13th edn., 2001, No. 9765 (*bibl*)

Blunden, G. *et al.*, *Phytochemistry*, 2001, **58**, 451-454 (*occur*)

12-Carboxymonascorubramine C-150

PP-V

[321922-28-5]




C₂₃H₂₅NO₆ 411.454

Prod. by *Penicillium* sp. AZ. Violet solid.

Ogihara, J. *et al.*, *J. Biosci. Bioeng.*, 2000, **90**, 549-554; 678-680 (*isol*, *struct*, *biosynth*)

Carboxymycobactins C-151



n = 2,3,4,5,6,7,8 for Carboxymycobactins 1-7

Complex. *Isol.* from *Mycobacterium smegmatis* grown under iron-deficient conditions. Extracellular mycobactins *isol.* from *Mycobacterium avium*. Siderophores and growth factors. Similar to Mycobactin, M-787.

Carboxymycobactin 1

C₃₆H₅₁N₅O₁₂ 745.825

Incorrectly indexed in CA.

Carboxymycobactin 2

C₃₇H₅₃N₅O₁₂ 759.852

Incorrectly indexed in CA.

Carboxymycobactin 3

C₃₈H₅₅N₅O₁₂ 773.879

Incorrectly indexed in CA.

Carboxymycobactin 4

C₃₉H₅₇N₅O₁₂ 787.906

Incorrectly indexed in CA.

Carboxymycobactin 5

C₄₀H₅₉N₅O₁₂ 801.932

Incorrectly indexed in CA.

Carboxymycobactin 6

C₄₁H₆₁N₅O₁₂ 815.959

Incorrectly indexed in CA.

Carboxymycobactin 7

C₄₂H₆₃N₅O₁₂ 829.986


Incorrectly indexed in CA.

Lane, S.J. *et al.*, *Tet. Lett.*, 1995, **36**, 4129-4132; 1996, **37**, 1 (*isol*, *struct*)

Rattledge, C. *et al.*, *Microbiology (Reading, U.K.)*, 1996, **142**, 2207-2212 (*occur*, *props*)

5-[(3-Carboxypropyl)amino]-2-hydroxy-3-tridecyl-1,4-benzoquinone C-152

4-[(4-Hydroxy-3,6-dioxo-5-tridecyl-1,4-cyclohexadien-1-yl)amino]butanoic acid [915728-09-5]



C₂₃H₃₇NO₅ 407.549

Constit. of the roots of *Embelia ribes*.

Amorph. red powder. λ_{max} 207 (log ε 1.74); 317 (log ε 1.32); 500 (log ε 0.09) (MeOH).


Lin, P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1629-1632 (*isol*, *pmr*, *cmr*, *ms*)

McErlean, C.S.P. *et al.*, *J.O.C.*, 2007, **72**, 10298-10301 (*synth*, *pmr*, *cmr*)

5-Carboxystrictosidine, 8Cl C-153

Tetrahydrodeoxycordifoline

[34371-11-4]



C₂₈H₃₄N₂O₁₁ 574.583

Alkaloid from the roots of *Rhazya orientalis*, *Guettarda platypoda* and *Uncaria tomentosa*. Mp 232° (225°). [α]_D -280 (MeOH).

N⁴,O-Penta-Ac, Me ester: [34417-91-9]
C₃₉H₄₆N₂O₁₆ 798.796
Mp 200°. [α]_D²⁰ -46 (c, 1 in CHCl₃).

3,4-Didehydro: 3,4-Dehydro-5-carboxy-strictosidine. 5-Carboxy-3,4-didehydrostrictosidine

C₂₈H₃₂N₂O₁₁ 572.568

Alkaloid from *Uncaria tomentosa*. λ_{max} 239; 357 (MeOH).

Parent acid: 3α,5α-Tetrahydrodeoxycordi-

foliac acid

C₂₇H₃₂N₂O₁₁ 560.557

Alkaloid from the roots of *Rhazya orientalis* (Apocynaceae). Amorph. Not named in the paper.

[134454-28-7]

De Silva, K.T.D. *et al.*, *Chem. Comm.*, 1971, 908-909 (*struct*, *synth*)


Ferrari, F. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1150-1151 (*isol*, *pmr*, *cmr*)

Aimi, N. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 2588-2590 (*synth*)

Kitajima, M. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1410-1412; 2002, **50**, 1376-1378 (*Dehydrocarboxystrictosidine*, *isol*, *synth*, *abs config*)

5-Carboxytetrahydroalstonine C-154

[53938-11-7]




Absolute configuration

C₂₂H₂₄N₂O₅ 396.442

Alkaloid from *Adina rubescens*. Non-cryst. [α]_D²⁵ -86 (CHCl₃) (as Me ester).

Brown, R.T. *et al.*, *Tet. Lett.*, 1974, 1649-1652 (*uv*, *ir*, *ester*, *ms*, *pmr*, *struct*)

3-Carboxy-1,2,3,4-tetrahydro-β-carboline-1-propanoic acid C-155



C₁₅H₁₆N₂O₄ 288.302

Constit. of various food sources.

[136612-30-1, 126366-08-3]


Liu, Z. *et al.*, *Tet. Lett.*, 1989, **30**, 3457-3460 (*synth*)

Bailey, P.D. *et al.*, *J.C.S. Perkin 1*, 1993, 441-449 (*synth*, *pmr*, *cmr*)

Gutsche, B. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 2458-2462 (*isol*, *pmr*, *ms*)

1-Carboxytrypargine C-156

[227935-85-5]



C₁₆H₂₁N₅O₂ 315.374

Abs stereochem. not determined. Alka-

loid from a *Eudistoma* sp. Amorph. solid.
 λ_{\max} 204 (log ϵ 4); 226 (log ϵ 4.3); 274 (log
 ϵ 3.7); 282 (log ϵ 3.7); 290 (log ϵ 3.6)
 (MeOH).

Van Wagoner, R.M. *et al.*, *J. Nat. Prod.*, 1999,
62, 794-797 (*isol.*, *uv.*, *cd.*, *pmr.*, *cmr*)

Carcinine **C-157**
 3-Amino-N-[2-(1H-imidazol-4-yl)ethyl]-
 propanamide, 9CI. N- β -Alanylhistamine
 [56897-53-1]



$\text{C}_8\text{H}_{14}\text{N}_4\text{O}$ 182.225

Occurs in cardiac tissue of *Carcinus
 maenas* and other crustacea. Lowers
 blood pressure; shows antioxidant props.
 Mp 67-68° Mp 114-117°.

Hydrochloride (1:2): [57022-38-5]
 Cryst. Mp 94° Mp 195°.

N-*tert*-Butoxycarbonyl: [143885-59-0]

$\text{C}_{13}\text{H}_{22}\text{N}_4\text{O}_3$ 282.342
 Mp 147-148°.

Arnould, J.-M. *et al.*, *Comp. Biochem. Physiol.*,
C: Comp. Pharmacol., 1975, **50**, 59-66; **51**,
 301-307 (*synth*)

Arnould, J.-M. *et al.*, *Arch. Int. Physiol.*
Biochim., 1977, **85**, 339-350 (*synth*)

Nanasawa, M. *et al.*, *Bull. Chem. Soc. Jpn.*,
 1981, **54**, 1101-1104 (*synth*)

Babizhayev, M.A. *et al.*, *Biochem. J.*, 1994,
304, 509-516 (*props.*, *ir.*, *pmr.*, *cmr*)


Impellizzeri, G. *et al.*, *Eur. J. Org. Chem.*, 2000,
 1065-1076 (*N-tert*-butoxycarbonyl, *synth*)

Cardiochrysin **C-158**
 Struct. unknown. Alkaloid from the
 leaves of *Chrysophyllum perpulchrum*
 (Sapotaceae). Mp 185°.

Hydrochloride: Mp 264°. $[\alpha]_{\text{D}}$ +21 (c, 0.5
 in H_2O).

Foussard-Blanpin, O. *et al.*, *Ann. Pharm. Fr.*,
 1965, **23**, 727-731; *CA*, **65**, 1242a

Cardionidine **C-159**
 [142609-18-5]




$\text{C}_{21}\text{H}_{25}\text{NO}_5$ 371.432

The config. of the 13-OH group is *S*-.
 Alkaloid from *Delphinium cardiopetalum*
 (Ranunculaceae). Mp 310-315° dec. $[\alpha]_{\text{D}}$ -
 50 (c, 0.03 in EtOH).

Reina, M. *et al.*, *Tet. Lett.*, 1992, **33**, 1661 (*isol.*
pmr., *cmr.*, *struct*)

Caribaeorane **C-160**
 [346406-75-5]



$\text{C}_{27}\text{H}_{36}\text{N}_2\text{O}_4$ 452.592

Constit. of *Erythropodium caribaeorum*.


15-Hydroxy: **15-Hydroxycaribaeorane**
 [216317-92-9]

$\text{C}_{27}\text{H}_{36}\text{N}_2\text{O}_5$ 468.592

Constit. of *Erythropodium caribaeorum*.

Britton, R. *et al.*, *Tet. Lett.*, 2001, **42**, 2953-
 2956 (*isol.*, *pmr.*, *cmr*)

Caribaeoside **C-161**
 [259728-79-5]



$\text{C}_{35}\text{H}_{48}\text{N}_2\text{O}_{11}$ 672.771

Constit. of *Erythropodium caribaeorum*.
 Related to Eleutherobin in E-64.


Aglycone, 15-Ac: **Caribaesolin**
 [259728-87-5]

$\text{C}_{30}\text{H}_{40}\text{N}_2\text{O}_7$ 540.655

Constit. of *Erythropodium caribaeorum*.

Cinel, B. *et al.*, *Org. Lett.*, 2000, **2**, 257-260
 (*isol.*, *pmr.*, *cmr*)

Caribine **C-162**
 [74483-60-6]




$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_3$ 326.394

Alkaloid from *Hymenocallis arenicola*.
 Prisms (Me_2CO). Mp 200-204°.

Doepke, W. *et al.*, *Z. Chem.*, 1980, **20**, 26-27

Carnosadine **C-163**
 1-Amino-2-[[[(aminoiminomethyl)ami-
 no]methyl]cyclopropanecarboxylic acid,
 9CI. 1-Amino-2-(guanidinomethyl)-
 1-cyclopropanecarboxylic acid
 [93961-20-7]



$\text{C}_6\text{H}_{12}\text{N}_4\text{O}_2$ 172.186

Amino acid from red alga *Grateloupia
 carnosa*. Antiinflammatory agent.
 Hygroscopic powder. Sol. H_2O ; poorly
 sol. butanol, hexane.

Bis(*p*-hydroxyazobenzenesulfonate):
 [94061-93-5]


Mp 226-232° dec. (no clear Mp).

Wakamiya, T. *et al.*, *Tet. Lett.*, 1984, **25**, 4411-
 4412; 1986, **27**, 2143-2144 (*isol.*, *pmr.*, *cmr.*
struct., *synth*)

Stammer, H. *et al.*, *Tetrahedron*, 1990, **46**,
 2231-2254 (*rev*)

Burgess, K. *et al.*, *J.O.C.*, 1994, **59**, 2179-2185
 (*synth.*, *pmr.*, *cmr*)

Carococculine **C-164**
 7,8-Didehydro-4,7-dihydroxy-3,8-di-
 methoxy-17-methylmorphinan-6-one, 9CI
 [54302-44-2]



Absolute
 configuration


$\text{C}_{19}\text{H}_{23}\text{NO}_5$ 345.394

Alkaloid from the leaves and stems of
Cocculus carolinus (Menispermaceae).
 Rods (EtOH). Mp 219-220° dec. $[\alpha]_{\text{D}}^{26}$ -
 29.5 (c, 0.715 in CHCl_3).

Slatkin, D.J. *et al.*, *J. Nat. Prod.*, 1974, **37**, 488
 (*isol.*, *struct*)

Elsobly, M.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**,
 132 (*isol*)

Carolisterol A **C-165**
 [151171-32-3]



$\text{C}_{27}\text{H}_{47}\text{NO}_{10}\text{S}$ 577.735

Constit. of *Styracaster caroli*.

6-Ketone: **Carolisterol B**
 [151171-33-4]

$\text{C}_{27}\text{H}_{45}\text{NO}_{10}\text{S}$ 575.719

Constit. of *Styracaster caroli*.

6-Epimer, 5-deoxy: Carolisterol C

[151171-34-5]
 $C_{27}H_{47}NO_9S$ 561.736
 Constit. of *Styracaster caroli*.

De Riccardis, F. *et al.*, *Tet. Lett.*, 1993, **34**, 4381 (*isol, pmr, cmr*)

Carosidine C-166

[1355-18-6]

Bisindole alkaloid. Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Mp 263-278° gradual dec. $[\alpha]_D^{26}$ -89.8 (c, 1 in $CHCl_3$). Mol. formula not recorded.

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*isol, uv, ir*)

Carosine C-167


[1355-19-7]

$C_{46}H_{56}N_4O_{10}$ 824.969
 Bisindole alkaloid. Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Needles (CH_2Cl_2/Et_2O). Mp 214-218°. $[\alpha]_D^{26}$ +6 (c, 1 in $CHCl_3$). λ_{max} 255 (E1%/1cm 162); 294 (E1%/1cm 155) (EtOH) (Berdy).

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*isol, uv, ir*)

Carpaine, 9CI C-168

[3463-92-1]



Absolute configuration

$C_{28}H_{50}N_2O_4$ 478.714

Homologue of Azimine, A-1608. Originally considered to be a monomeric 13-membered cyclic lactone. Alkaloid from leaves of *Carica papaya* (papaya) (Caricaceae). Cardiotonic agent, CNS depressant, also showing other physiological effects (also reported to be cardiodepressant). Cubes (Et_2O). Sol. MeOH, C_6H_6 ; poorly sol. hexane. Mp 119-120° (117-118.5°). $[\alpha]_D$ +21.65 (c, 1.0 in EtOH). $[\alpha]_{230}^{220}$ -240 (c, 0.12 in MeOH). Log P 4.77 (calc). Prob. formed *in vivo* by dimerisation of Carpamic acid, from which it has been made *in vitro*.

1,2-Didehydro: Dehydrocarpaine I. 12,13-Didehydrocarpaine, 9CI

[72362-02-8]
 $C_{28}H_{48}N_2O_4$ 476.698

Alkaloid from the leaves of *Carica papaya* (papaya) (Caricaceae). Viscous oil.

1,1',2,2'-Tetradehydro: Dehydrocarpaine II. 12,13,25,26-Tetradehydrocarpaine, 9CI

[72362-03-9]
 $C_{28}H_{46}N_2O_4$ 474.682

Alkaloid from the leaves of *Carica papaya* (papaya) (Caricaceae). Viscous oil.

2-Epimer: Pseudocarpaine. ψ -Carpaine [3760-91-6]

$C_{28}H_{50}N_2O_4$ 478.714

Minor alkaloid from leaves of *Carica papaya* (papaya) (Caricaceae). Cryst. (petrol). Mp 65-68°. $[\alpha]_D^{28}$ +4.95 (c, 1.62 in EtOH). Yields carpamic and pseudocarpamic acids on hydrolysis.

2-Epimer: hydrochloride:

Cryst. (EtOH). Mp 295°.

Greshoff, M. *et al.*, *Ber.*, 1890, **23**, 3537 (*isol*)
 Govindachari, T.R. *et al.*, *J.C.S.*, 1954, 1847; 1955, 1563 (*isol, props, Carpaine, Pseudocarpaine*)
 Narasimhan, N.S. *et al.*, *Chem. Ind. (London)*, 1956, 1526 (*synth*)

Spiteller-Friedmann, M. *et al.*, *Monatsh. Chem.*, 1964, **95**, 1234 (*ms, struct*)
 Coke, J.L. *et al.*, *J.O.C.*, 1965, **30**, 3420 (*abs config*)
 Govindachari, T.R. *et al.*, *Tet. Lett.*, 1965, 1907 (*ms, pmr, struct*)

Brown, E. *et al.*, *Chem. Lett.*, 1974, 109 (*synth*)
 Corey, E.J. *et al.*, *J.A.C.S.*, 1975, **97**, 654 (*synth*)

Topuriya, L.I. *et al.*, *Khim. Prir. Soedin.*, 1978, 414; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, 354 (*isol*)

Tang, C.-S. *et al.*, *Phytochemistry*, 1979, **18**, 651 (*Dehydrocarpaines*)

Hanessian, S. *et al.*, *Tet. Lett.*, 1979, 3391 (*synth*)

Natsume, M. *et al.*, *Heterocycles*, 1980, **14**, 169 (*synth*)


Rajnikant, G. *et al.*, *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. C*, 1998, **9**, 131-137 (*cryst struct*)

Sato, T. *et al.*, *Org. Lett.*, 2003, **5**, 3839-3842 (*synth*)

Randl, S. *et al.*, *Tet. Lett.*, 2004, **45**, 1167-1169 (*synth*)

Carquinostatin A C-169

1-(2-Hydroxypropyl)-2-methyl-6-(3-methyl-2-butenyl)-3H-carbazole-3,4(9H)-dione, 9CI. CS 79A. Antibiotic CS 79A [150957-18-9]



$C_{21}H_{23}NO_3$ 337.418

Alkaloid from *Streptomyces exfoliatus*. Potent neuronal cell protecting substance. Also shows radical scavenging activity. Reddish powder. Mp 144-145°. λ_{max} 242 (ϵ 30400); 287 (ϵ 28000); 470 (ϵ 8100) (MeOH/NaOH) (Derep). λ_{max} 230 (ϵ 22200); 267 (ϵ 29700); 425 (ϵ 5400) (MeOH) (Berdy).

10S-Hydroxy: Carquinostatin B. Antibiotic CS 79B. CS 79B

[155569-69-0]
 $C_{21}H_{23}NO_4$ 353.417

Prod. by *Streptomyces exfoliatus* and *Streptomyces tendae*. Neuronal cell protecting agent. Antioxidant. Reddish-brown powder. Sol. DMSO, DMF, EtOH, EtOAc, $CHCl_3$, MeOH; poorly sol. hexane. λ_{max} 223; 248; 428

(MeOH) (Berdy).

Shin-ya, K. *et al.*, *Tet. Lett.*, 1993, **34**, 4943-4944 (*isol, uv, ir, pmr, cmr, struct*)

Japan. Pat., 1994, 94 32 778; *CA*, **121**, 7450x (*CS 79B*)


Shin-ya, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 1768-1769

(*Carquinostatin B*)

Orihara, N. *et al.*, *J. Antibiot.*, 1997, **50**, 979-981 (*biosynth*)

Knolker, H.-J. *et al.*, *Synlett*, 1999, 596-598 (*synth*)


Knoelker, H.-J. *et al.*, *Tet. Lett.*, 2000, **41**, 1171-1174 (*synth*)

Cartormin C-170

$C_{27}H_{29}NO_{13}$ 575.525

Enolised β -diketone. Constit. of *Carthamus tinctorius* (safflower). Yellow prisms (MeOH). $[\alpha]_D^{27}$ -153.4 (c, 0.01 in Py). Mp >230° dec.

Yin, H.-B. *et al.*, *Tet. Lett.*, 2000, **41**, 1955-1958 (*isol, pmr, cmr, ms, cryst struct*)

Caryolivine C-171

$C_{36}H_{34}N_2O_6$ 590.674

(R)-form [103956-43-0]

Alkaloid from *Caryomene olivascens* (Menispermaceae). $[\alpha]_D$ -46 (c, 0.08 in MeOH).

N²-Me: Berbacolorflamine

[80550-38-5]
 $C_{37}H_{37}N_2O_6^{\oplus}$ 605.709

Alkaloid from the stem and roots of *Pycnarrhena longifolia* (Menispermaceae). $[\alpha]_D^{20}$ +1000 (c, 0.004 in $CHCl_3$). No counterion indicated (prob. chloride).

(S)-form

N²-Me: Fenfangjine D. 1,3,4-Tridehydrofangchinolium [115439-62-8]

$C_{37}H_{37}N_2O_6^{\oplus}$ 605.709

Quaternary alkaloid from "Fen-Fang-Ji" (roots of *Stephania tetrandra*) (Menispermaceae). Angiotensin I-converting enzyme inhibitor. Orange amorph. powder + 1/2 H₂O (as hydroxide); orange granules (MeOH/Me₂CO) (as chloride). Mp 300°. $[\alpha]_D^{25}$ +67.8


(c, 0.116 in MeOH). The assigned struct. is the enantiomer of Berbacolorflamine but the reported opt. rotn. is much smaller.

O¹²-De-Me: Dehatridine. Stepierrine
[121256-46-0]
C₃₅H₃₂N₂O₆ 576.648
Alkaloid from the leaves of *Dehaasia triandra* (Lauraceae) and from tubers of *Stephania piirrii* (Menispermaceae). Needles (MeOH). Mp 274-276°. [α]_D²³ +98 (c, 0.1 in MeOH). [α]_D +55 (c, 0.10 in CHCl₃). Identity of Dehatridine and Stepierrine not yet establ. Mp refers to Dehatridine. λ_{\max} 202 (ε 25100); 243 (ε 15900); 282 (ε 2510); 335 (ε 1590) (EtOH) (Derep). λ_{\max} 207 (ε 100000); 263 (ε 12600); 293 (sh) (ε 3980); 373 (ε 2000) (EtOH/KOH) (Derep).

Van Beek, T.A. *et al.*, *J.O.C.*, 1982, **47**, 898 (Berbacolorflamine)
Lavault, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1148 (Caryolivine)
Ogino, T. *et al.*, *Heterocycles*, 1988, **27**, 1149 (Fengfanjine B)
Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1989, **52**, 846 (Stepierrine)
Lu, S.T. *et al.*, *Phytochemistry*, 1989, **28**, 615 (Dehatridine)
Chen, C.-K. *et al.*, *Chin. Pharm. J. (Taipei)*, 2003, **55**, 35-47 (Dehatridine)

Caseadine**C-172**

Alkaloid F35

*(S)*-formC₂₀H₂₃NO₄ 341.406***(S)*-form** [34413-12-2]

Alkaloid from *Corydalis caseana* (Papaveraceae). Mp 145°. [α]_D -393 (c, 1.22 in CHCl₃).

 α -N-Oxide: cis-Caseadine N-oxide[151563-67-6]
C₂₀H₂₃NO₅ 357.405

Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Amorph. powder. Mp 156°. [α]_D -199 (c, 0.06 in MeOH).

Me ether: Mp 186°. [α]_D -360 (c, 0.4 in CHCl₃).

O¹¹-De-Me: Caseamine. Alkaloid F33[27498-04-0]
C₁₉H₂₁NO₄ 327.379

Struct. revised in 1988. Alkaloid from *Corydalis caseana* (Papaveraceae). Major alkaloid from the whole plant of *Ceratocarpus heterocarpa* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 246-247° (257°). [α]_D -365 (c, 0.04 in CHCl₃). [α]_D -406 (c, 0.12 in CHCl₃).

O¹¹-De-Me, α -N-oxide: cis-Caseamine N-oxide[151563-66-5]
C₁₉H₂₁NO₅ 343.379


Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Pale yellowish cryst. (MeOH). Mp 244-246°. [α]_D -216 (c, 0.05 in MeOH).

(±)-form*Me ether*:Cryst. (Et₂O). Mp 165°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 153 (Caseadine, Caseamine, isol)
Chen, C.Y. *et al.*, *Tet. Lett.*, 1968, 349 (Caseamine, uv, ms)
Govindachari, T.R. *et al.*, *Heterocycles*, 1977, **6**, 1811 (struct, bibl)
Ognyanov, V. *et al.*, *Heterocycles*, 1982, **19**, 1069 (synth)
Suau, R. *et al.*, *Phytochemistry*, 1988, **27**, 1920; 1993, **34**, 559 (oxides)

Caseadinium(1+)**C-173**

[152343-37-8]


C₂₀H₂₀NO₄⁺ 338.382

Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Yellowish solid (as chloride). Mp 192° (chloride). CAS no. refers to chloride.

Suau, R. *et al.*, *Phytochemistry*, 1993, **34**, 559 (isol, uv, pmr, ms, struct)

Caseanadine**C-174**

5,8,13,13a-Tetrahydro-2,9,10-trimethoxy-6H-dibenzo[a,g]quinolizin-1-ol. 2,9,10-Trimethoxyberbin-1-ol

C₂₀H₂₃NO₄ 341.406

Unusual 1,2,9,10-tetraoxygenated substitution pattern.

***(S)*-form** [40141-47-7]

Alkaloid from *Corydalis caseana* and from the leaves, stems and flowers of *Corydalis clarkei* (Papaveraceae). Cryst. (MeOH). Mp 169-170°. [α]_D -267 (CHCl₃) (-212).

O⁹-De-Me: Clarkeanidine[99615-99-3]
C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves, stems and flowers of *Corydalis clarkei* (Papaveraceae). Cryst. (2-propanol). Mp 178-179°. [α]_D -277 (CHCl₃).

Yu, C.K. *et al.*, *Can. J. Chem.*, 1971, **49**, 124 (isol, uv, ir, pmr, ms, struct)

Rothera, M.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 802 (isol, uv, ir, pmr, cmr, ms, deriv)

Suau, R. *et al.*, *Phytochemistry*, 1988, **27**, 1920 (synth, struct, Clarkeanidine)

Casimirine**C-175**C₃₀H₃₂N₂O₅ 500.593


Struct. unknown. Alkaloid from *Casimiroa edulis* (Mexican apple). Mp 106°.

Bickern, *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1903, **241**, 166-176 (isol)

Casimiroedine**C-176**

N-[2-(1-β-D-Glucopyranosyl-1H-imidazol-4-yl)ethyl]-N-methyl-3-phenyl-2-propenamide, 9CI. N²-Cinnamoyl-N^α-methylhistamine N-glucoside
[5853-02-1]

[51013-83-3 (Z-isomer)]

C₂₁H₂₇N₃O₆ 417.461

Alkaloid from the seeds of *Casimiroa edulis*. Antineoplastic, carcinostatic agent. Mp 226.5-228° (223-224°). [α]_D²⁰ -21.3 (c, 1.2 in 5% HCl aq.) (-27, -30.7). Log P -2.06 (calc).

Picrate: Mp 110-112°.

Power, F.B. *et al.*, *J.C.S.*, 1911, 1993 (isol)
Aebi, A. *et al.*, *Helv. Chim. Acta*, 1956, **39**,

1495 (isol, ir, uv)

Djerassi, C. *et al.*, *Tetrahedron*, 1958, **2**, 168a (struct)


Raman, S. *et al.*, *Tet. Lett.*, 1962, 357 (cryst struct)

Panzica, R.P. *et al.*, *J.A.C.S.*, 1973, **95**, 8737 (uv, pmr, ms, synth, struct)

Reynolds, W.F. *et al.*, *Magn. Reson. Chem.*, 2000, **38**, 366-369 (pmr, cmr)

Casimiroin**C-177**

6-Methoxy-9-methyl-1,3-dioxolo[4,5-h]quinolin-8(9H)-one, 9CI. 4-Methoxy-1-methyl-7,8-methylenedioxy-2(1H)-quinolinone
[477-89-4]

C₁₂H₁₁NO₄ 233.223

Alkaloid from the seed and bark of *Casimiroa edulis* (Mexican apple) (Rutaceae). Cryst. (Me₂CO/hexane). Mp 202-203°. λ_{\max} 226 ; 236 ; 260 ; 300 (MeOH) (Berdy). λ_{\max} 232 ; 262 (ε 8); 465 (EtOH) (Berdy).

Kincl, F.A. *et al.*, *J.C.S.*, 1956, 4163 (isol, uv, ir)

Iriarte, J. *et al.*, *J.C.S.*, 1956, 4170 (isol)

Meisels, A. *et al.*, *J.A.C.S.*, 1957, **79**, 6328


(*struct*)

Weinstein, B. *et al.*, *Tetrahedron*, 1964, **20**, 1725 (*synth, uv, ir*)

Cassaidine

C-178

[26296-41-3]



Absolute Configuration

$C_{24}H_{41}NO_4$ 407.592

Alkaloid from the bark of *Erythrophleum guineense*, *Erythrophleum chlorostachys*, *Erythrophleum couminga* and *Erythrophleum ivorense* (Fabaceae). Cardiac stimulant, hypertensive, antiseptic agent. Local anaesthetic. Cryst. (Me₂CO/Et₂O). Mp 139.5°. [α]_D²⁰ -98 (c, 1 in EtOH). Log P 3.67 (calc).

► Toxic.

Hydrochloride: Mp 249-252°. [α]_D²² -96 (c, 0.75 in H₂O).

3-O-(3-Hydroxy-3-methylbutanoyl):

Coumidine

[22149-19-5]

$C_{29}H_{49}NO_6$ 507.709

Alkaloid from the bark of *Erythrophleum guineense* and *Erythrophleum ivorense* (Fabaceae). Cryst. (Et₂O). Mp 130-132°. [α]_D -58 (c, 0.5 in CHCl₃).

N-De-Me: **Norcassaidine**

$C_{23}H_{39}NO_4$ 393.565

Trace alkaloid detected in extracts of the bark of *Erythrophleum chlorostachys* by pmr (Fabaceae). Not indexed by CAS.

7-Ketone: **Cassaine**

[468-76-8]

$C_{24}H_{39}NO_4$ 405.576

Alkaloid from *Erythrophleum guineense*, *Erythrophleum ivorense*, *Erythrophleum suaveolens* and *Erythrophleum couminga* (Fabaceae). Strong cardiac stimulant showing digitalis-like action. Antihypertensive agent. Shows local anaesthetic props. Shows cytotoxicity against KB carcinoma cells but inactive in several other test systems. Sol. MeOH, Et₂O, C₆H₆; poorly sol. H₂O. Mp 142.5°. [α]_D²⁰ -103 (c, 1 in EtOH). Log P 3.27 (calc).

► Can cause respiratory and cardiac paralysis.

7-Ketone, *hydrochloride*:

Cryst. + 1H₂O (EtOH/butanone/Et₂O). Mp 220° (212-213°).

7-Ketone, 3-O-(3-methyl-2-butenoyl): **3-(3-Methylcrotonyl)cassaine**

[51771-63-2]

$C_{29}H_{45}NO_5$ 487.678

Alkaloid from bark of *Erythrophleum ivorense* (Fabaceae). Cryst. (Et₂O). Mp 169-170°.

7-Ketone, 3-O-(3-hydroxy-3-methylbutanoyl): **Coumingine**. β -Hydroxyisovalerylcassaine

[26241-81-6]

$C_{29}H_{47}NO_6$ 505.693

Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Cardiac stimulant, hypertensive agent, smooth muscle stimulant, inotropic agent. Shows local anaesthetic props. Cryst. (Et₂O). Mp 142°. [α]_D²⁰ -70 (c, 1 in EtOH). Log P 4.27 (calc).

► Highly toxic.

7-Ketone, 3-O-(3-hydroxy-3-methylbutanoyl); *hydrochloride*:
Needles (EtOH/Et₂O). Mp 195°.

7-Ketone, N-de-Me, 3-O-acyl: **Coumingidine**

$C_{28}H_{45}NO_6$ 491.667

Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Cryst. (Et₂O). Mp 160-161°. Acylated with a small MW hydroxyacyl group. MF not certain, may be $C_{27}H_{43}NO_6$. No further data to 2007.

7-Ketone, N-de-Me, 3-O-acyl, *hydrochloride*:

Cryst. (EtOH/Et₂O). Mp 217-219°.

7-Ketone, N-de-Me, 3-O-(3-methyl-2-butenoyl): **Ivorine**

[1748-19-2]

$C_{28}H_{43}NO_5$ 473.651

Alkaloid from the bark of *Erythrophleum ivorense* (Fabaceae). Mp 158-159°. [α]_D²⁰ -43 (c, 0.77 in CHCl₃).

7-Ketone, N-de-Me, 3-O-(3-methyl-2-butenoyl), *hydrochloride*: Mp 224-225°.

19-Hydroxy, 7-ketone: **19-Hydroxycassaine**

[51795-37-0]

$C_{24}H_{39}NO_5$ 421.576

Alkaloid from the bark of *Erythrophleum ivorense* (Fabaceae). Cryst. (CHCl₃/petrol). Mp 144-145°.

Dalma, G. *et al.*, *Helv. Chim. Acta*, 1939, **22**, 1497-1512 (*Cassaine, isol*)

Ruzicka, L. *et al.*, *Helv. Chim. Acta*, 1940, **23**, 753-764; 1941, **24**, 63-76; 1449-1458

(*Cassaidine, Cassaine, Coumingine, isol, uv*)
Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1941, **24**, 319E-352E (*Coumingidine*)

Engel, B.G. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 2364-2381 (*Cassaine, Cassaidine, isol*)

Ottinger, R. *et al.*, *Bull. Soc. Chim. Belg.*, 1965, **74**, 198-199 (*Ivorine*)

Turner, R.B. *et al.*, *J.A.C.S.*, 1966, **88**, 1766-1775 (*synth, struct*)

Clarke, R.L. *et al.*, *J.A.C.S.*, 1966, **88**, 5865-5871 (*uv, pmr, config*)

Morin, R.B. *et al.*, *Alkaloids (Academic Press)*, 1967, **10**, 298 (*Coumingidine, struct*)

Thorell, A. *et al.*, *Acta Chem. Scand.*, 1968, **22**, 2835-2844 (*Coumidine*)

Cronlund, A. *et al.*, *Acta Pharm. Suec.*, 1971, **8**, 351-360; 1973, **10**, 507-514; *CA*, **76**, 11995y; **80**, 80085w (*Cassaidine, Coumidine, 19-Hydroxycassain, 3-(3-Methylcrotonyl)cassaine*)

Friedrich-Fiechtl, J. *et al.*, *Chem. Ber.*, 1971, **104**, 3535-3548 (*ms*)

Loder, J.W. *et al.*, *Aust. J. Chem.*, 1974, **27**, 179-185 (*Norcassaidine*)

Cronlund, A. *et al.*, *Planta Med.*, 1976, **29**, 126 (*Cassaidine, Cassaine, occur*)


Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 4-6 (*pharmacol, antitumour props*)

Phoenix, S. *et al.*, *J.A.C.S.*, 2008, **130**, 13989-13995 (*Cassaine, synth*)

Cassamedine

C-179

4-Methoxy-8H-bis[1,3]benzodioxolo[6,5,4-de:5',6'-g]quinolin-8-one, 9CI [16408-75-6]



$C_{19}H_{11}NO_6$ 349.299

Alkaloid from *Cassytha americana* (*Cassytha filiformis*) and from the trunk wood of *Siparuna guianensis* (Lauraceae, Monimiaceae). Orange microcryst. solid (CHCl₃/EtOH). Mp 278°.

Demethoxy: **Cassameridine**

[16408-76-7]

$C_{18}H_9NO_5$ 319.273

Alkaloid from *Cassytha americana* and from root wood of *Litsea kawakamii* (Lauraceae). Shows antimicrobial activity. Cryst. (CHCl₃/MeOH). Mp 301-302°. λ_{max} 251 (ε 29000); 274 (ε 24500); 323 (ε 12000); 353 (ε 8100); 388 (ε 7000); 440 (ε 5400) (EtOH) (Berdy).

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2443 (*isol, uv, ir, pmr, struct*)

Cava, M.P. *et al.*, *Tetrahedron*, 1973, **29**, 2245 (*isol, uv, ir, pmr, struct, synth, deriv*)

Braz Filho, R. *et al.*, *Phytochemistry*, 1976, **15**, 1187 (*isol*)


Lu, S.T. *et al.*, *CA*, 1979, **92**, 194484d (*isol, deriv*)

Hufford, C.D. *et al.*, *J. Pharm. Sci.*, 1980, **69**, 1180-1182 (*activity*)

Cassiadinine

C-180

[104777-95-9]



$C_{16}H_{13}N_3O_4$ 311.296


Alkaloid from the flowers of *Cassia siamea* (Fabaceae). Green needles (CH₂Cl₂/petrol). Mp 80° dec. V. unstable.

Biswas, K.M. *et al.*, *Phytochemistry*, 1986, **25**, 1727 (*isol, uv, ir, pmr, ms, struct*)

Cassiarine A

C-181

[951306-03-9]



C₁₃H₁₁NO₂ 213.235

Alkaloid from the leaves of *Cassia siamea*. Antiplasmodial agent. Reddish solid. λ_{\max} 215 (ε 19000); 230 (sh) (ε 14000); 253 (ε 13600); 315 (sh) (ε 3600); 338 (ε 5000); 370 (sh) (ε 3200) (MeOH).

Morita, H. *et al.*, *Org. Lett.*, 2007, **9**, 3691-3693 (*isol, uv, pmr, cmr*)

Oshimi, S. *et al.*, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 3761-3765 (*biosynth*)


Yao, Y.-S. *et al.*, *J.O.C.*, 2008, **73**, 5221-5225 (*synth*)

Rudyanto, M. *et al.*, *Org. Lett.*, 2008, **10**, 1921-1922 (*synth*)

Cassiarine B

C-182

[951306-04-0]

C₁₈H₁₉NO₄ 313.352

Alkaloid from the leaves of *Cassia siamea*. Reddish solid. λ_{\max} 220 (ε 9400); 235 (ε 8800); 245 (ε 9400); 255 (ε 9700); 275 (ε 4400); 317 (ε 3800); 372 (ε 2500) (MeOH).


Morita, H. *et al.*, *Org. Lett.*, 2007, **9**, 3691-3693 (*isol, pmr, cmr*)

Yao, Y.-S. *et al.*, *J.O.C.*, 2008, **73**, 5221-5225 (*synth*)

Cassine

C-183

12-(5-Hydroxy-6-methyl-2-piperidinyl)-2-dodecanone, 9CI. 3-Hydroxy-2-methyl-6-(11-oxododecyl)piperidine

C₁₈H₃₅NO₂ 297.48**(-)-form** [5227-24-7]

Alkaloid from leaves of *Cassia excelsa* and *Cassia carnavale*, also *Prosopis rusciifolia*, *Prosopis alpataco* and *Prosopis sericantha* (Fabaceae). Mp 57-58°. Bp_{0.001} 90°. $[\alpha]_{\text{D}}^{25}$ -0.6 (EtOH).

Hydrochloride: Mp 173-175°.

O-(4-Hydroxy-3-methoxy-E-cinnamoyl):

O-Feruloylcassine

[1002097-72-4]

C₂₈H₄₃NO₅ 473.651

Alkaloid from the unripe fruits of *Senna spectabilis*. Pale yellow oil. $[\alpha]_{\text{D}}^{20}$ +2.9 (c, 1 in CHCl₃). λ_{\max} 236 (log ε 4.03); 326 (log ε 4.25) (MeOH).

N-Me-N-MethylcassineC₁₉H₃₇NO₂ 311.507

Alkaloid from *Prosopis* spp. (Fabaceae). Mp 110.5-111.5° (semisynthetic). $[\alpha]_{\text{D}}^{26}$ +6.5 (EtOH).

11'-ξ-Alcohol: **Carnavaleine**. 5-Hydroxy-

α,6-dimethyl-2-piperidineundecanol, 9CI. 3-Hydroxy-6-(11-hydroxydodecyl)-2-methylpiperidine

[16049-25-5]

C₁₈H₃₇NO₂ 299.496

Alkaloid from *Cassia spectabilis* (Fabaceae). Mp 60.7-61.2°.

6-Epimer: 6-Isocassine

[64474-08-4]

C₁₈H₃₅NO₂ 297.48

Alkaloid from *Cassia spectabilis* leaves (Fabaceae). Cryst. (EtOAc) (as hydrochloride). Mp 123° (hydrochloride). $[\alpha]_{\text{D}}^{25}$ +1.5 (c, 1.22 in CHCl₃).

6-Epimer, O-Ac: O-Acetyl-6-isocassine.**Spectamine B**C₂₀H₃₇NO₃ 339.517

Alkaloid from *Cassia spectabilis*.

6-Epimer, O-benzoyl: O-Benzoyl-6-isocassine. Spectamine AC₂₅H₃₉NO₃ 401.588

Alkaloid from *Cassia spectabilis* and from the flowers of *Senna spectabilis*. Inhibitor of superoxide anion production. Cytotoxic agent. Pale yellow oil. $[\alpha]_{\text{D}}^{25}$ +2.6 (c, 0.46 in EtOH). λ_{\max} 204 (log ε 5.2); 229 (log ε 5.4); 273 (log ε 4.5) (MeOH).

6-Epimer, 11'-ξ-alcohol: 6-Isocarnavaleine

[66512-85-4]

[66537-25-5]

C₁₈H₃₇NO₂ 299.496

Alkaloid from *Cassia spectabilis* (Fabaceae). Oil. Mp 123° (as hydrochloride). $[\alpha]_{\text{D}}^{25}$ -5.66 (c, 0.03 in CHCl₃).

(±)-form [57760-16-4]

Alkaloid from *Cassia excelsa* (Fabaceae). Mp 73-74°.

Hight, R.J. *et al.*, *J.O.C.*, 1964, **29**, 471; 1966, **31**, 1275 (*isol, ir, pmr, struct, config*)

Rice, W.Y. *et al.*, *J.O.C.*, 1966, **31**, 1010 (*struct, abs config*)

Lythgoe, D. *et al.*, *Tet. Lett.*, 1967, 1133 (*Carnavaleine*)

Fales, H.M. *et al.*, *J.A.C.S.*, 1970, **92**, 1590 (*ms*)

Hight, R.J. *et al.*, *Tet. Lett.*, 1970, 1803 (*ms*)

Christofidis, I. *et al.*, *Tetrahedron*, 1977, **33**, 977; 3005 (6-Isocassine, Isocarnavaleine)

Brown, E. *et al.*, *Bull. Soc. Chim. Fr.*, 1981, 281 (*synth*)

Chiale, C.A. *et al.*, *An. Asoc. Quim. Argent.*, 1982, **70**, 337 (*isol, deriv*)

Hasseberg, H.-A. *et al.*, *Annalen*, 1989, 255 (*synth, ir, pmr, cmr*)

Momose, T. *et al.*, *Tet. Lett.*, 1993, **34**, 5785 (*synth*)

Pahl, A. *et al.*, *Tetrahedron*, 1997, **53**, 7255 (*Carnavaleine, synth*)

Oetting, J. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 477-484 (*synth*)

Kamo, T. *et al.*, *Heterocycles*, 2003, **60**, 1303-1306 (6-Isocassine, Spectamines A,B)

Makabe, H. *et al.*, *Org. Lett.*, 2003, **4**, 27-29 (*synth*)

Sriphong, L. *et al.*, *Planta Med.*, 2003, **69**, 1054-1056 (6-epimer benzoyl)

Leverett, C.A. *et al.*, *J.O.C.*, 2006, **71**, 8591-8601 (*synth*)

Herdeis, C. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 524-529 (*synth*)


Viegas, C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2026-2028 (*Feruloylcassine*)

Kim, G. *et al.*, *Tet. Lett.*, 2007, **48**, 4481-4483 (*synth*)

Cassipoureamide A

C-184

4-Hydroxy-N-(2-hydroxyethyl)-1,2-dithiolane-3-carboxamide



Relative Configuration

C₆H₁₁NO₃S₂ 209.29


Alkaloid from *Cassipourea guianensis*. Amorph. powder (CHCl₃/MeOH). Mp 105-109° Mp 42-46.5° (as di-Ac). $[\alpha]_{\text{D}}^{23}$ +173.9 (c, 0.89 in CHCl₃) (di-Ac). λ_{\max} 244 (log ε 2.87); 280 (sh) (log ε 2.45); 320 (log ε 2.28) (CHCl₃) (di-Ac).

Ichimaru, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1675-1676

Cassipoureamide B

C-185

5-Hydroxy-N-(2-hydroxyethyl)-1,2,3-trithiane-4-carboxamide



Relative Configuration

C₆H₁₁NO₃S₃ 241.356

Alkaloid from *Cassipourea guianensis*. Cryst. (CHCl₃/MeOH). Mp 109.5-113°. $[\alpha]_{\text{D}}^{23}$ -97 (c, 0.71 in MeOH). λ_{\max} 206 (log ε 4.03); 223 (sh) (log ε 3.78); 267 (log ε 3.18) (MeOH).


Ichimaru, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1675-1676

Cassipourine

C-186

Dodecahydro-1H,6H-[1,2,5,6]tetrahydroino[3,4-a:8,7-a']dipyrrolizine, 9CI. Cassipurine

[14051-10-6]



Probable absolute configuration

C₁₄H₂₂N₂S₄ 346.605

Alkaloid from *Cassipourea gummiflua* twigs and leaves (Rhizophoraceae). Needles (Me₂CO). Sol. CHCl₃, Mp 212°. $[\alpha]_{\text{D}}^{20}$ -11.8 (c, 1 in CHCl₃).

Methiodide (1:2):

Prisms (H₂O). Mp 258-260°.

Cooks, R.G. *et al.*, *J.C.S.(C)*, 1967, 286 (*struct*)

Wright, W.G. *et al.*, *J.C.S.(C)*, 1967, 283 (*isol*)


Gaffner, G. *et al.*, *Acta Cryst. B*, 1969, **25**, 2114 (*cryst struct*)

Wróbel, J.T. *et al.*, *Can. J. Chem.*, 1981, **59**, 1101 (*synth*)

Cassythic acid

C-187

[1004981-48-9]



Absolute Configuration


C₂₀H₁₉NO₆ 369.373

Alkaloid from *Cassipoupa filiformis*. Brown syrup. $[\alpha]_D^{24} +105$ (c, 1 in MeOH). λ_{\max} 227 (log ϵ 4.31); 277 (log ϵ 4.06); 310 (sh) (log ϵ 3.63) (MeOH).

Tsai, T.-H. *et al.*, *J. Nat. Prod.*, 2008, **71**, 289-291 (isol, pmr, cmr)

Castanospermine**C-188**

Octahydro-1,6,7,8-indolizinetrol, 9CI. 1,6,7,8-Tetrahydroxyindolizidine [79831-76-8]

C₈H₁₅NO₄ 189.211

Alkaloid from the seeds of *Castanospermum australe* and the pods of *Alexa leiopetala* (Fabaceae). Tentatively identified in *Alexa canaracunensis*, *Alexa cowanii*, *Alexa grandiflora*, *Alexa herminiana*, *Alexa imperatricis*, *Alexa leiopetala*, *Alexa superba* and *Alexa wachenheimii*. Potent inhibitor of α - and β -glucosidases and other carbohydrate processing enzymes. Potent dicot. phytoalexin. Shows anti-HIV activity and antiinflammatory props. Large cubic cryst. (EtOH aq.). Mp 212-215° (207-210°). $[\alpha]_D +80$ (c, 0.93 in H₂O). Log P -2.25 (calc).

8-O- β -D-Glucopyranoside: [124378-36-5]

C₁₄H₂₅NO₉ 351.353

Alkaloid from the seeds of *Castanospermum australe*. $[\alpha]_D +44.1$ (c, 0.65 in H₂O).

6-Butanoyl: 6-O-Butyrylcastanospermine. **Celgosivir**, INN. MBI 3253. MDL 28574

[121104-96-9]

C₁₂H₂₁NO₅ 259.302

α -Glucosidase inhibitor, shows anti-HIV activity, antiviral agent. Used for the treatment of hepatitis C. Cryst. solid. Mp 114-115°. Log P -0.34 (calc).

6-Butanoyl; hydrochloride: **Celgosivir hydrochloride**, USAN. MDL 28574A [141117-12-6]

6-Epimer: 6-Epicastanospermine

[107244-34-8]

C₈H₁₅NO₄ 189.211

Minor alkaloid from the seeds of *Castanospermum australe* (Fabaceae). Potent inhibitor of amyloglucosidase (an exo-1,4- α -glucosidase) and a weak inhibitor of β -galactosidase (does not inhibit β -glucosidase or α -mannosidase). Solid. Mp 203-206° dec. $[\alpha]_D^{20} +2.8$ (c, 1.0 in H₂O).

6-Epimer, 7-deoxy: 7-Deoxy-6-epicastanospermine

[129724-72-7]

C₈H₁₅NO₃ 173.211

Alkaloid from *Castanospermum australe*. Mod. inhibitor of amyloglucosidase and yeast α -glucosidase. Does not inhibit α -mannosidase. Oil. $[\alpha]_D^{26} +18.3$ (c, 0.712 in MeOH).

6,7-Diepimer: 6,7-Diepicastanospermine [130983-46-9]

C₈H₁₅NO₄ 189.211

Alkaloid from seeds of *Castanospermum australe* (Fabaceae). Moderately good inhibitor of amyloglucosidase but a relatively weak inhibitor of β -glucosidase. Oil. $[\alpha]_D^{27} +42.7$ (c, 0.675 in MeOH).

Hohenschutz, L.D. *et al.*, *Phytochemistry*, 1981, **20**, 811 (isol, cryst struct, uv, pmr, cmr)

Bernotas, R.C. *et al.*, *Tet. Lett.*, 1984, **25**, 165 (synth, abs config, bibl)

Setoi, H. *et al.*, *Tet. Lett.*, 1985, **26**, 4617 (synth, abs config)

Howard, A.S. *et al.*, *Alkaloids (Academic Press)*, 1986, **28**, 287 (rev, pharmacol)

Molyneux, R.J. *et al.*, *Arch. Biochem. Biophys.*, 1986, **251**, 450 (epimer)

Nash, R.J. *et al.*, *CA*, 1986, **104**, 143181m (rev)

Hamana, H. *et al.*, *J.O.C.*, 1987, **52**, 5492 (Castanospermine, 6-Epicastanospermine, synth)

Stevens, K.L. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 1467

Nash, R.J. *et al.*, *Phytochemistry*, 1988, **27**, 1403 (isol)

Johnson, V.A. *et al.*, *Antimicrob. Agents Chemother.*, 1989, **33**, 53 (anti-HIV activity)

Winchester, B.G. *et al.*, *Biochem. J.*, 1990, **269**, 227 (anti-HIV activity)

Molyneux, R.J. *et al.*, *J. Nat. Prod.*, 1990, **53**, 609 (Deoxyepicastanospermine)

Margolin, A.L. *et al.*, *J.A.C.S.*, 1990, **112**, 2849 (synth, 6-butanoyl, pmr, ms)

Nash, R.J. *et al.*, *Phytochemistry*, 1990, **29**, 1356 (epimer)

Anzeveno, P.B. *et al.*, *Tet. Lett.*, 1990, **31**, 4321 (synth)

Bhide, R. *et al.*, *Tet. Lett.*, 1990, **31**, 4827 (synth)

Molyneux, R.J. *et al.*, *Biochemistry*, 1991, **30**, 9981 (diepimer)

Reymond, J.L. *et al.*, *J.O.C.*, 1991, **56**, 2128 (synth)

Gerspacher, M. *et al.*, *J.O.C.*, 1991, **56**, 3700 (Castanospermine, 6-Epicastanospermine, synth)

Willenborg, D.O. *et al.*, *Immunol. Cell Biol.*, 1992, **70**, 369 (pharmacol)

Mulzer, J. *et al.*, *J.O.C.*, 1992, **57**, 3194 (synth)

Burgess, K. *et al.*, *Tetrahedron*, 1992, **48**, 4045 (rev, synth, bibl)

Grassberger, V. *et al.*, *Annalen*, 1993, 379 (synth)

Pan, Y.T. *et al.*, *Arch. Biochem. Biophys.*, 1993, **303**, 134 (activity)

Hempel, A. *et al.*, *J. Med. Chem.*, 1993, **36**, 4082 (cryst struct, abs config)

Ina, H. *et al.*, *J.O.C.*, 1993, **58**, 52 (synth)

Kim, N.-S. *et al.*, *J.O.C.*, 1993, **58**, 7096 (synth)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 561

Taylor, D.L. *et al.*, *Antimicrob. Agents Chemother.*, 1994, **38**, 1780 (6-butanoyl, pharmacol)

Bartlett, M.R. *et al.*, *Immunol. Cell Biol.*, 1994, **72**, 367 (pharmacol)

Kim, N.-S. *et al.*, *Tet. Lett.*, 1994, **35**, 3489 (synth, diepimer)

Ferneaux, R.H. *et al.*, *Tetrahedron*, 1994, **50**, 2131 (synth, diepimer)

Bridges, C.G. *et al.*, *Glycobiology*, 1995, **5**, 243-247; 249-253 (6-butanoyl, pharmacol)

Kang, M.S. *et al.*, *Glycobiology*, 1996, **6**, 209-216 (6-butanoyl, metab)

Kang, S.H. *et al.*, *Chem. Comm.*, 1998, 1353-1354 (synth)

Izquierdo, I. *et al.*, *Eur. J. Org. Chem.*, 1999, 1269-1274 (1-Deoxyepicastanospermine)

Bartnicka, E. *et al.*, *Tetrahedron*, 1999, **55**, 2061-2076 (synth)

Batey, R.A. *et al.*, *Tet. Lett.*, 2000, **41**, 9935-9938 (synth, Deoxyepicastanospermine)

Zhao, H. *et al.*, *J.O.C.*, 2001, **66**, 1761-1767 (synth)

Kim, J.H. *et al.*, *Synthesis*, 2003, 2473-2478 (synth, 6-Epicastanospermine, ir, pmr, cmr)

Somfai, P. *et al.*, *Tetrahedron*, 2003, **59**, 1293-1299 (synth)

Zhang, H.-X. *et al.*, *Tetrahedron*, 2003, **59**, 2015-2020 (6-Epicastanospermine, synth)

Kato, A. *et al.*, *Tetrahedron: Asymmetry*, 2003, **14**, 325-331 (8-glucoside)


Whitby, K. *et al.*, *Antiviral Chem. Chemother.*, 2004, **15**, 141-151 (celgosivir, pharmacol)

Sorbera, L.A. *et al.*, *Drugs of the Future*, 2005, **30**, 545 (celgosivir, rev)

Zhao, Z. *et al.*, *Tetrahedron*, 2005, **61**, 8888-8894 (synth)

Karanjule, N.S. *et al.*, *J.O.C.*, 2006, **71**, 4667-4670 (synth)

Machan, T. *et al.*, *Tetrahedron*, 2008, **64**, 2725-2732 (synth)

Catacandin**C-189**

Partial structure of Catacandin A

Isol. from *Lysobacter gummosus*. Possesses good anticandidal activity. Sol. MeOH, EtOAc, bases; poorly sol. H₂O, hexane. λ_{\max} 260 (E1%/1cm 600); 335 (E1%/1cm 180) (MeOH) (Berdy). λ_{\max} 269 (E1%/1cm 600); 357 (E1%/1cm 238); 375 (E1%/1cm 180) (HCl) (Berdy). λ_{\max} 260 (E1%/1cm 760); 335 (E1%/1cm 180) (NaOH) (Berdy).

Catacandin A [100753-64-8]

C₂₉H₃₈N₂O₆ 510.629

Sol. MeOH, EtOAc, bases; poorly sol. H₂O, hexane. Main prod. λ_{\max} 205 (ϵ 14000); 268 (ϵ 25500); 358 (ϵ 13500); 375 (ϵ 11500) (MeOH/HCl) (Derep). λ_{\max} 257 (ϵ 33000); 340 (ϵ 8500) (MeOH/NaOH) (Derep). λ_{\max} 204 (ϵ 17500); 258 (ϵ 30500); 343 (ϵ 8500) (MeOH) (Derep). λ_{\max} 204 (ϵ 17500); 258 (ϵ 30500); 343 (ϵ 8500) (MeOH) (Berdy). λ_{\max} 205 (ϵ 14000); 268 (ϵ 22500); 358 (ϵ 13500); 375 (ϵ 11500) (MeOH-HCl) (Berdy). λ_{\max} 257 (ϵ 33000); 340 (ϵ 8500) (MeOH-NaOH) (Berdy). λ_{\max} 269 (E1%/1cm 600); 357 (E1%/1cm 238); 375 (E1%/1cm 180) (HCl) (Berdy). λ_{\max} 260 (E1%/1cm 760); 335 (E1%/1cm 180) (NaOH) (Berdy).

Catacandin B [100753-65-9]

C₂₉H₃₈N₂O₆ 510.629


Sol. MeOH, EtOAc; poorly sol. H₂O. Minor isoln. product isomeric with Catacandin A. λ_{\max} 215 (ϵ 17500); 324 (ϵ 10500) (MeOH/HCl) (Derep). λ_{\max} 236 (ϵ 15800); 317 (ϵ 7800) (MeOH/NaOH) (Derep). λ_{\max} 205 (ϵ 16100); 236 (ϵ 15600); 320 (ϵ 7800) (MeOH) (Derep). λ_{\max} 205 (ϵ 16100); 236 (ϵ 15600); 320 (ϵ 7800) (MeOH) (Berdy). λ_{\max} 215 (ϵ

17500); 324 (ε 10500) (MeOH-HCl) (Berdy). λ_{\max} 236 (ε 15800); 317 (ε 7800) (MeOH-NaOH) (Berdy).

Meyers, E. *et al.*, *J. Antibiot.*, 1985, **38**, 1642 (*isol, props*)

Cataline C-190

5,6,6a,7-Tetrahydro-1,2,9,10-tetra-methoxy-6-methyl-4H-dibenzo[de,g]quinolin-4-ol, 9Cl. 4-Hydroxy-1,2,9,10-tetramethoxyaporphine [38764-84-0]



$C_{21}H_{25}NO_5$ 371.432
Alkaloid from *Glaucium flavum* var. *vestitum* (Papaveraceae). Mp 183°. $[\alpha]_D^{25} +166$ (CHCl₃).

N-De-Me: Norcataline

[63598-89-0]
Synthetic. Cryst. (EtOH). Mp 160-162°. $[\alpha]_D^{25} +106.7$ (c, 1 in CHCl₃).

4-Epimer: Epicataline

[63598-92-5]
Synthetic. Mp 188-189°. $[\alpha]_D^{25} +97.3$ (c, 1 in CHCl₃).

O²-De-Me: 2,4-Dihydroxy-1,9,10-trimethoxyaporphine. Srilankine

[70509-78-3]
 $C_{20}H_{23}NO_5$ 357.405
Alkaloid from *Alseodaphne semicarpifolia* (Lauraceae). $[\alpha]_D +122$ (c, 0.18 in MeOH).


O¹-De-Me, N-de-Me: 1,4-Dihydroxy-2,9,10-trimethoxynoraporphine. 4-Hydroxywilsonirine

[107882-31-5]
 $C_{19}H_{21}NO_5$ 343.379
Alkaloid from the bark of *Popowia piscocarpa* (Annonaceae). Amorph. $[\alpha]_D^{20} +60$ (c, 0.33 in MeOH).

Ribas, I. *et al.*, *Tet. Lett.*, 1972, 2033 (*uv, ir, pmr, ms, struct*)
Hartenstein, J. *et al.*, *Angew. Chem., Int. Ed.*, 1977, **16**, 730 (*synth, pmr*)
Smolnycki, W.D. *et al.*, *Tet. Lett.*, 1978, 4617 (*Srilankine*)
Castedo, L. *et al.*, *Heterocycles*, 1982, **19**, 1591 (*pmr, cryst struct*)
Jossang, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1028 (*4-Hydroxywilsonirine*)

Cathafoline C-191

Methyl 1,2-dihydro-1-methylakuammilan-17-oate, 9Cl [50906-83-7]



Absolute Configuration

$C_{21}H_{26}N_2O_2$ 338.449
Alkaloid from *Alstonia quaternata* and *Catharanthus longifolius* (Apocynaceae). Mp 156°. $[\alpha]_D^{20} -40$ (c, 0.34 in CHCl₃).

N⁴-Oxide: Cathafoline N⁴-oxide

[154849-46-4]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Prisms (MeOH). Mp 145-155°. $[\alpha]_D^{23} -20.6$ (c, 1.02 in MeOH).

19 α ,20 α -Epoxide: Quaternoxine

[57499-01-1]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Alstonia quaternata* and *Alstonia vittensis* (Apocynaceae). Cryst. (Et₂O/MeOH). Mp 168°. $[\alpha]_D^{20} -34$ (c, 0.4 in CHCl₃).

19 ξ ,20 ξ -Epoxide: Raucubainine

[84412-95-3]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from leaves of *Rauwolfia salicifolia* (Apocynaceae). $[\alpha]_D^{20} -7$ (CHCl₃).

19 ξ ,20 ξ -Epoxide, N⁴-Me: 4-Methylraucubaininium(1+)

[123789-22-0]
 $C_{22}H_{29}N_2O_3^{\oplus}$ 369.483
Alkaloid from leaves of *Vinca minor* (Apocynaceae). Mp 228-230° dec. (as chloride). CAS Reg. No. refers to chloride.

10-Methoxy: 10-Methoxycathafoline

$C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Alstonia macrophylla*. Pale yellow oil. $[\alpha]_D -57$ (c, 0.08 in CHCl₃). λ_{\max} 204 (log ε 4.23); 249 (log ε 3.74); 315 (log ε 3.88) (EtOH).

10-Methoxy, N⁴-oxide: 10-Methoxycathafoline N⁴-oxide

$C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from the leaves of *Alstonia angustifolia* var. *latifolia*. Light yellow oil. $[\alpha]_D -32$ (c, 0.14 in CHCl₃). λ_{\max} 208 (log ε 4.16); 249 (log ε 3.79); 311 (log ε 3.37) (EtOH).

11-Methoxy: Cabucraline

[53802-11-2]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Cabucala erythrocarpa*, *Cabucala torulosa* and *Alstonia vittensis* (Apocynaceae). Cryst. (Me₂CO). Mp 148°. $[\alpha]_D^{20} -45$ (c, 0.5 in CHCl₃).

11-Methoxy, 10-formyl: Cabucraline 10-carboxaldehyde

[87596-49-4]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Alstonia plumosa* (Apocynaceae). $[\alpha]_D -33$ (c, 0.5 in CHCl₃).

11-Methoxy, N⁴-oxide: Cabucraline N⁴-oxide

[87596-48-3]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from root bark of *Alstonia plumosa* (Apocynaceae). $[\alpha]_D -37$ (c, 0.37 in CHCl₃).

11-Methoxy, 19 α ,20 α -epoxide: Caberine. 11-Methoxyquaternoxine

[53802-10-1]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from *Alstonia plumosa* and

from several *Cabucala* spp. from Madagascar (Apocynaceae).

11-Methoxy, 18-hydroxy: 18-Hydroxycabucraline

[132268-04-3]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from the stem bark of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae). $[\alpha]_D -31$ (c, 0.5 in CHCl₃).

Rasoanaivo, P. *et al.*, *Tet. Lett.*, 1973, 1425 (*uv, ir, pmr, ms, struct*)

Douzoua, L. *et al.*, *Phytochemistry*, 1974, **13**, 1994 (*isol, uv, ir, pmr, ms, Cabucraline*)

Mamatas-Kalamaras, S. *et al.*, *Phytochemistry*, 1975, **14**, 1637; 1849 (*uv, ir, pmr, ms, struct, Quaternoxine*)

Das, B.C. *et al.*, *J.O.C.*, 1977, **42**, 2785 (*cmr*)
Sierra, P. *et al.*, *Coll. Czech. Chem. Comm.*, 1982, **47**, 2912 (*Raucubainine*)

Jacquier, M.J. *et al.*, *Phytochemistry*, 1982, **21**, 2973 (*Cabucraline 10-carboxaldehyde, Cabucraline oxide, Caberine*)

Uhrin, D. *et al.*, *J. Nat. Prod.*, 1989, **52**, 637 (*4-Methylraucubaininium*)

Proksa, B. *et al.*, *Planta Med.*, 1989, **55**, 188 (*4-Methylraucubaininium*)

Morfaux, A.-M. *et al.*, *Phytochemistry*, 1990, **29**, 3345 (*18-Hydroxycabucraline*)

Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 249 (*oxide*)
Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 839-844; 2004, **65**, 603-608 (*10-Methoxycathafoline, 10-Methoxycathafoline N⁴-oxide*)

Cathalanceine

C-192

[1355-29-9]


Struct. unknown. Alkaloid from roots of *Catharanthus lanceus* (Apocynaceae). Cryst. (Me₂CO). Mp 188-190° dec. pK_a 4.5 (33% DMF aq.).

Blomster, R.N. *et al.*, *J. Nat. Prod.*, 1964, **27**, 480-485 (*isol, uv, ir*)

Cathaphylline

C-193

Methyl 2,3-didehydro-21-hydroxy-20-oxoaspido-permidine-3-carboxylate [59444-63-2]



Absolute configuration


$C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from *Catharanthus trichophyllus* (Apocynaceae). Noncryst. $[\alpha]_D^{23} -438$ (c, 0.1 in CHCl₃).

Cordell, G.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 366 (*isol, ir, uv, ms, pmr, struct*)

Catharanthamine

C-194

[78779-58-5]




$C_{46}H_{56}N_4O_9$ 808.97
Alkaloid from *Catharanthus roseus* (Apocynaceae). Shows antitumour props. λ_{max} 213 (ϵ 5750); 220 (ϵ 7800); 223 (ϵ 5100); 265 (ϵ 2200); 277 (ϵ 16000); 295 (ϵ 1100) (MeOH) (Berdy).

El-Sayed, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 289 (isol, uv, ir, pmr, cmr, ms, struct)

Catharanthine**C-195**

[2468-21-5]
[20395-98-6 (\pm)-form]



Absolute Configuration

$C_{21}H_{24}N_2O_2$ 336.433

Belongs to the opposite enantiomeric series to the other iboga alkaloids. Alkaloid from *Catharanthus ovalis* and other *Catharanthus* spp. and from *Catharanthus roseus* (Apocynaceae). Diuretic. Shows hypoglycaemic activity. Active against gram-positive bacteria and fungi. Mp 126-128°. $[\alpha]_D^{25} +29.8$ (CHCl₃). Log P 3.44 (uncertain value) (calc). λ_{max} 226; 284; 292 (MeOH).

Sulfate (2:1): Mp 164-167° dec.

Neuss, N. *et al.*, *Tet. Lett.*, 1961, 206-210 (struct)

Gorman, M. *et al.*, *J.A.C.S.*, 1965, **87**, 93-99 (pmr)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1966, 810-812; 890-891 (biosynth)

Büchi, G. *et al.*, *J.A.C.S.*, 1970, **92**, 999-1005 (synth, ms)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **39**, 2258-2266 (abs config)

Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1672-1689 (synth)

Wenkert, E. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2437-2442 (cmr)

Hernandez, N.M.R. *et al.*, *Rev. Cubana Farm.*, 1977, **11**, 249-255 (activity)

Imanishi, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4052-4059 (synth)

Raucher, S. *et al.*, *J.O.C.*, 1985, **50**, 3236-3237 (synth)

Szántay, C. *et al.*, *Tetrahedron*, 1990, **46**, 1711-1732 (synth)

Reding, M.T. *et al.*, *Org. Lett.*, 1999, **1**, 973-976 (synth)

Moisan, L. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 5334-5336 (synth)

Catharinine**C-196**

[1355-30-2]


$C_{46}H_{52}N_4O_{10}$ 820.938

Bisindole alkaloid. Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Plates (Me₂CO). Mp 231-234° dec. $[\alpha]_D^{26} +34.8$ (c, 1 in CHCl₃).

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (isol, uv, ir)

Catharine**C-197**

[1355-31-3]



Absolute configuration

$C_{46}H_{54}N_4O_{10}$ 822.953

Minor alkaloid from *Catharanthus roseus*, *Catharanthus ovalis* and *Catharanthus longifolius* (Apocynaceae). Mp 271-275° dec. $[\alpha]_D -54.2$ (CHCl₃). Could be an artifact of autoxidation.

▶ VS3630000

Rasoanaivo, P. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **279**, 75 (ir, pmr, cmr, ms, struct)

Guilhelm, J. *et al.*, *Acta Cryst. B*, 1976, **32**, 936 (cryst struct)


Kutney, J.P. *et al.*, *Can. J. Chem.*, 1979, 1682 (synth)

Langlois, N. *et al.*, *Chem. Comm.*, 1979, 582 (synth)

Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2088 (biosynth)

Catharsitoxin A**C-198**

3-Ethyl-5,6,7,8-tetrahydro-2-methylimidazo[1,2-a]pyridine
[376587-10-9]



$R^1 = -CH_2CH_3$, $R^2 = CH_3$

$C_{10}H_{16}N_2$ 164.25

Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.

Suenaga, K. *et al.*, *Tet. Lett.*, 2001, **42**, 7079-7081 (isol, synth, pmr, cmr)

Catharsitoxin B**C-199**

2-Ethyl-5,6,7,8-tetrahydro-3-methylimidazo[1,2-a]pyridine
[376587-11-0]

As Catharsitoxin A, C-198 with $R^1 = CH_3$, $R^2 = CH_2CH_3$

$C_{10}H_{16}N_2$ 164.25

Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.

Suenaga, K. *et al.*, *Tet. Lett.*, 2001, **42**, 7079-7081 (isol, pmr)

Catharsitoxin C**C-200**

5,6,7,8-Tetrahydro-3-isopropylimidazo[1,2-a]pyridine
[376587-12-1]

As Catharsitoxin A, C-198 with $R^1 = -CH(CH_3)_2$, $R^2 = H$


$C_{10}H_{16}N_2$ 164.25

Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.

Suenaga, K. *et al.*, *Tet. Lett.*, 2001, **42**, 7079-7081 (isol, pmr)

Cathedulin E3**C-201**

Cathedulin K11
[61230-22-6]



$C_{54}H_{60}N_2O_{23}$ 1105.068

Alkaloid from *Catha edulis* (Celastraceae). Cryst. (Et₂O/petrol). Mp 245-248°. $[\alpha]_D^{20} -44.8$ (c, 0.27 in CHCl₃). λ_{max} 215 (ϵ 40750); 268 (ϵ 13010); 293 (sh) (ϵ 5890) (EtOH).

O⁶-De-Ac: Cathedulin E4

[61230-20-4]

$C_{52}H_{58}N_2O_{22}$ 1063.03

Alkaloid from *Catha edulis* (Celastraceae). Amorph. solid. $[\alpha]_D^{27} -37$ (c, 0.56 in CHCl₃). λ_{max} 215 (ϵ 41000); 268 (ϵ 12040) (EtOH).

O²-De-Ac, 2-benzoyl: Cathedulin K20

[106982-89-2]

$C_{59}H_{62}N_2O_{23}$ 1167.138


Alkaloid from fresh young leaves and stems of *Catha edulis* (Celastraceae).

Baxter, R.L. *et al.*, *J.C.S. Perkin 1*, 1979, 2965-2971; 2982-2989 (*Cathedulins E3-E4*)

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1986, 531-534 (*Cathedulin K20*)

Cathedulin K17**C-202**

[106999-78-4]




C₅₉H₆₂N₂O₂₃ 1167.138

CAS Name is incorrect. Alkaloid from fresh young leaves and stems of *Catha edulis* (Celastraceae). Cryst. solid.

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1986, 531 (isol, pmr, struct)

Cathedulins K19 C-203
[106999-79-5]



Absolute Configuration

C₅₄H₅₈N₂O₂₃ 1103.052

Alkaloid from fresh young leaves and stems of *Catha edulis* (Celastraceae). Cryst. (MeOH). Mp 249-251°.

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1986, 531 (isol, pmr, cmr, struct)

Kim, T.-S. *et al.*, *Tet. Lett.*, 1993, **34**, 5535 (config)

Cathedulins Y7 C-204
[73298-55-2]

Struct. unknown. Alkaloid from Yemeni khat *Catha edulis* (Celastraceae). Obt. only in trace amt., structurally related to the other cathedulins. M⁺958, conts. acetate, nicotinoate, gallate and evoninate residues.

Baxter, R.L. *et al.*, *J.C.S. Perkin 1*, 1979, 2965-2971

Cathedulins Y9 C-205
[73298-56-3]

Struct. unknown. Trace alkaloid from Yemeni khat *Catha edulis* (Celastraceae). M⁺900, conts. acetate, nicotinoate, gallate and evoninate residues.


Baxter, R.L. *et al.*, *J.C.S. Perkin 1*, 1979, 2965-2971

Cathedulins Y10 C-206
[73298-58-5]

Struct. unknown. Trace alkaloid from Yemeni khat *Catha edulis* (Celastraceae). M⁺980, conts. acetate, nicotinoate, gallate and everninate residues.

Baxter, R.L. *et al.*, *J.C.S. Perkin 1*, 1979, 2965-2971

Cathenamines C-207
Dehydroajmalicine
[63661-74-5]



Absolute configuration

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from *Guettarda eximia* (Apocynaceae). Key intermediate in the biosynthesis of Ajmalicine, A-212 and related alkaloids. [α]_D²⁰ -52 (c, 1 in CHCl₃). Very reactive base, reduced by NaBH₄ to Tetrahydroalstonine in A-212. λ_{max} 228; 274; 280; 290 (sh) (EtOH).

17-Hydroxy, 16,17-dihydro: 16,17-Dihydro-17-hydroxycathenamamine

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Guettarda eximia* (Apocynaceae). Proposed intermed. in biosynth. of heteroyohimbine alkaloids. Inseparable mixt. of epimers. Labile, forms Cathenamamine on standing in soln. λ_{max} 228; 274; 280; 290 (EtOH).

3,19-Diepimer: 3-Iso-19-epicathenamamine

[75766-20-0]

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from cell suspension cultures of *Catharanthus ovalis* (Apocynaceae).

Stöckigt, J. *et al.*, *Chem. Comm.*, 1977, 164-166; 646-648 (biosynth)

Husson, H.P. *et al.*, *Tet. Lett.*, 1977, 1889-1892 (isol, uv, pmr, struct)

Kan-Fan, C. *et al.*, *Chem. Comm.*, 1978, 618-619 (16,17-Dihydro-17-hydroxycathenamamine)

Rüffer, M. *et al.*, *Chem. Comm.*, 1979, 1016-1018 (biosynth)

Stöckigt, J. *et al.*, *Planta Med.*, 1980, **40**, 22-30 (3-Iso-19-epicathenamamine)

Martin, S.F. *et al.*, *J.A.C.S.*, 1988, **110**, 5925-5927 (synth)


Lounasmaa, M. *et al.*, *Heterocycles*, 1997, **45**, 779-786 (synth)

Cathestatin B C-208

PF 1126B. Antibiotic PF 1126B

[162112-42-7]

[169735-72-2, 167818-23-7]

C₁₇H₂₃N₃O₆ 365.385

Prod. by *Aspergillus terricola*, *Penicillium citrinum* and the sponge-derived *Microascus longirostris*. Inhibitor of cysteine protease. Amorph. solid. Stereochemical identity with PF 1126 antibiotics not certain.

4'-Deoxy: Cathestatin A. PF 1126A.

Antibiotic PF 1126A

[162112-41-6]

[167818-22-6, 169735-71-1, 169735-77-7, 169735-75-5]

C₁₇H₂₃N₃O₅ 349.386

Prod. by *Aspergillus terricola*, *Microascus longirostris* and *Penicillium citrinum*. Inhibitor of cysteine protease. Amorph. solid. Sol. H₂O, MeOH.

Woo, J.-T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 350-352 (isol, pmr, cmr)

Japan. Pat., 1995, 95 157 479; CA, **123**, 196747z (PF 1126B)

Yu, C.-M. *et al.*, *J. Antibiot.*, 1996, **49**, 395-397 (isol, pmr, cmr)

Cathestatin C C-209

PF 1138B. Antibiotic PF 1138B

[176486-11-6]

[172940-60-2]

As Cathestatin B, C-208 with

n = 4

C₁₈H₂₅N₃O₆ 379.412

Prod. by *Chaetomium globosum* and the sponge-derived *Microascus longirostris*. Inhibitor of cysteine protease. Sol. H₂O. Stereochemical identity with PF 1138 antibiotics not certain.

4'-Deoxy: Antibiotic PF 1138A. PF 1138A

[172940-59-9]

C₁₈H₂₅N₃O₅ 363.413

Prod. by *Chaetomium globosum*. Thiol protease inhibitor.

Japan. Pat., 1995, 95 252 245; CA, **124**,

117071u (PF 1138)

Yu, C.-M. *et al.*, *J. Antibiot.*, 1996, **49**, 395 (isol, pmr, cmr)

Cathidine A C-210

[58573-03-8]

Struct. unknown. Alkaloid from *Catha edulis* (Celastraceae). Amorph. solid. Mp 73-75°. Closely related to 1,2,3,4,9,14-Hexahydroxydihydro-β-agarofuran. Anal. gave C 64.14, H 6.66, N 2.29, O 27.70%.

Cais, M. *et al.*, *Tetrahedron*, 1975, **31**, 2727-2731

Cathidine B C-211

[58573-02-7]

Struct. unknown. Alkaloid from *Catha edulis* (Celastraceae). Mp 112-118° (softens at 107°).

Cais, M. *et al.*, *Tetrahedron*, 1975, **31**, 2727-2731

Cathidine C-212


[1355-11-9]

Struct. unknown. Alkaloid from roots of *Catharanthus roseus* (Apocynaceae). Cryst. (MeOH)(as sulfate). Mp 239-245° dec. (sulfate).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (isol, uv)

Cathovaline C-213

Methyl 4-(acetyloxy)-3,6-epoxy-1-methylspidospemidine-3-carboxylate, 9CI. Cathanneine
[35470-43-0]



Absolute Configuration

C₂₄H₃₀N₂O₅ 426.511

Alkaloid from *Catharanthus ovalis* and *Catharanthus lanceus* (Apocynaceae). Mp

88-90° (76-77°). $[\alpha]_D$ -73 (c, 0.5 in CHCl_3).

O-De-Ac: Deacetylcaothovaline

[56435-45-1]
 $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_4$ 384.474

Alkaloid from *Catharanthus ovalis* (Apocynaceae). Noncryst. $[\alpha]_D$ -88 (c, 0.52 in CHCl_3).

14ξ-Hydroxy: 14-Hydroxycathovaline

[68170-67-2]
 $\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_6$ 442.511

Alkaloid from *Catharanthus ovalis* (Apocynaceae). Mp 133-134°. $[\alpha]_D$ -73 (c, 0.7 in CHCl_3).

Langlois, N. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 994; 1972, **275**, 219 (Cathovaline, 14-Hydroxycathovaline, *isol, ms, pmr, struct*)

Aynilian, G.H. et al., *Tet. Lett.*, 1972, 89; 391 (*ir, uv, pmr, ms, struct, abs config*)

Diatta, L. et al., *Bull. Soc. Chim. Fr.*, 1975, 671 (Deacetylcaothovaline)

Langlois, N. et al., *Phytochemistry*, 1979, **18**, 467 (14-Hydroxycathovaline)

Caulerpicin

C-214

[11003-42-2]

$\text{C}_{43}\text{H}_{87}\text{NO}_2$ 650.166

Consists of a mixt. of ceramides derived from 2-Amino-1,3-octadecanediol with $\text{C}_{18} \rightarrow \text{C}_{26}$ saturated fatty acids. Toxic principle, *isol.* from the green alga *Caulerpa racemosa* and other *Caulerpa* spp. Concentrates in crabs, shrimps and other organisms in the marine food chain. Cryst. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 95°.

▶ Toxic agent causing sensory numbness.


Doty, M.S. et al., *Nature (London)*, 1966, **211**, 990

Nielsen, P.G. et al., *Phytochemistry*, 1982, **21**, 1643 (*pmr, ms, struct*)

Caulersin

C-215

[196803-51-7]



$\text{C}_{21}\text{H}_{14}\text{N}_2\text{O}_3$ 342.353

Alkaloid from the alga *Caulerpa serrulata*. Bright yellow cryst. (MeOH). Mp 269-270°. λ_{max} 206 (log ϵ 4.51); 220 (log ϵ 3.62); 265 (log ϵ 2.02); 273 (log ϵ 1.1); 311 (log ϵ 1.36) (EtOH).

Su, J.-Y. et al., *J. Nat. Prod.*, 1997, **60**, 1043-1044 (*isol, uv, ir, pmr, cmr, ms*)


Wahlström, N. et al., *Tetrahedron*, 2004, **60**, 2147-2153 (*synth, pmr, cmr*)

Miki, Y. et al., *Tet. Lett.*, 2006, **47**, 5215-5218 (*synth*)

Caulidine A

C-216

3-(1,1-Dimethyl-2-propenyl)-5-(3-methyl-2-butenyl)-1H-indole. 3-(1,1-Dimethyl-2-propenyl)-5-prenylindole [820965-41-1]



$\text{C}_{18}\text{H}_{23}\text{N}$ 253.386


Alkaloid from *Isolona cauliflora*. Yellow oil.

Nkunya, M.H.H. et al., *Nat. Prod. Res.*, 2004, **18**, 253-258 (*isol, pmr, cmr, ms*)

Caulidine B

C-217

3-(1,1-Dimethyl-2-propenyl)-5-(3-oxo-1-butenyl)-1H-indole [820965-42-2]



$\text{C}_{17}\text{H}_{19}\text{NO}$ 253.343


Alkaloid from *Isolona cauliflora*. Yellow oil.

Nkunya, M.H.H. et al., *Nat. Prod. Res.*, 2004, **18**, 253-258 (*isol, pmr, cmr, ms*)

Caulindole A

C-218

[675103-47-6]



R = H

Relative Configuration

$\text{C}_{26}\text{H}_{26}\text{N}_2$ 366.505

Alkaloid from *Isolona cauliflora*. Yellow oil. $[\alpha]_D$ +13.9 (c, 0.17 in CHCl_3).

3-Epimer: Caulindole B

[675103-48-7]

$\text{C}_{26}\text{H}_{26}\text{N}_2$ 366.505

Constit. of *Isolona cauliflora*. Yellow oil.

Makangara, J.J. et al., *Phytochemistry*, 2004, **65**, 227-232 (*isol, pmr, cmr*)

Caulindole C

C-219

[675103-49-8]

As Caulindole A, C-218 with

R = -C(CH₃)₂CH=CH₂

$\text{C}_{36}\text{H}_{42}\text{N}_2$ 502.741

Alkaloid from *Isolona cauliflora*. Yellow gum.

3-Epimer: Caulindole D

[675103-50-1]

$\text{C}_{36}\text{H}_{42}\text{N}_2$ 502.741


Alkaloid from *Isolona cauliflora*. Yellow gum.

Makangara, J.J. et al., *Phytochemistry*, 2004, **65**, 227-232 (*isol, pmr, cmr*)

Caulophyllumine B

C-220

4-[2-(1-Methyl-2-piperidinyl)ethenyl]-phenol. 2-[2-(4-Hydroxyphenyl)ethenyl]-1-methylpiperidine



$\text{C}_{14}\text{H}_{19}\text{NO}$ 217.31

(S,E)-form [1009318-61-9]


Alkaloid from the roots of *Caulophyllum thalictroides* (blue cohosh). Brown powder. $[\alpha]_D^{28}$ -13.2 (c, 0.22 in MeOH). λ_{max} 265 (log ϵ 3.41) (MeOH).

Ali, Z. et al., *Phytochemistry*, 2008, **69**, 1037-1042 (*isol, pmr, cmr*)

Cavernicolin 1

C-221

[84590-04-5]



Absolute Configuration

$\text{C}_8\text{H}_7\text{Br}_2\text{NO}_3$ 324.956

Alkaloid from the sponges *Aplysina cavernicola* and *Suberea* aff. *praetensa*. λ_{max} 200 (sh) (MeOH/NaOH) (Derep). λ_{max} 257 (ϵ 8300) (MeOH) (Derep).

5-Debromo, 5-chloro: 7-Bromo-5-chloro-cavernicolin

$\text{C}_8\text{H}_7\text{BrClNO}_3$ 280.505

Metab. from *Aplysina cavernicola*. Racemic. *Isol.* as a mixt. of C(7)-epimerising monoacetates.

7-Debromo: 5-Bromocavernicolin

$\text{C}_8\text{H}_8\text{BrNO}_3$ 246.06

Isol. from the sponge *Suberea* aff. *praetensa*.

5,7-Bisdebromo, 5-chloro: 5-Chlorocavernicolin

$\text{C}_8\text{H}_8\text{ClNO}_3$ 201.609

Alkaloid from *Aplysina cavernicola* and *Suberea* aff. *praetensa*. Nearly racemic.

7-Epimer: Cavernicolin 2

[84590-06-7]

$\text{C}_8\text{H}_7\text{Br}_2\text{NO}_3$ 324.956


Alkaloid from *Aplysina cavernicola* and *Suberea* aff. *praetensa*. λ_{max} 200 (sh) (MeOH/NaOH) (Derep). λ_{max} 257 (ϵ 8300) (MeOH) (Derep).

D'Ambrosio, M. et al., *Helv. Chim. Acta*, 1984, **67**, 1484-1492 (*isol, uv, pmr, cmr, ms, struct*)

Kijjoo, A. *et al.*, *Z. Naturforsch.*, C, 2002, **57**, 732-738 (*Suberea aff. pratensis* alkaloids)

Cavidine C-222

6,6a,11,14-Tetrahydro-8,9-dimethoxy-6-methyl-12H-benzo[*a*]-1,3-benzodioxolo[4,5-*g*]quinolizine, 9CI. 2,3-Dimethoxy-13-methyl-9,10-(methylenedioxy)berbine, 8CI



C₂₁H₂₃NO₄ 353.417

(+)-form [32728-75-9]

Alkaloid from *Corydalis meifolia* and *Corydalis saxicola*. Cryst. (C₆H₆). Mp 148-149°. [α]_D +323 (c, 0.13 in CHCl₃).

13-Epimer: **Thalictrifoline**. Thalictrofoline [30342-06-4]

C₂₁H₂₃NO₄ 353.417

Alkaloid from *Corydalis thalictrifolia*. Mp 155°. [α]_D +199 (c, 0.136 in CHCl₃). [α]_D +218 (c, 0.4 in MeOH).

(±)-form [30342-07-5]

Alkaloid from *Corydalis saxicola* and *Corydalis thalictrifolia*. Cryst. (MeOH). Mp 192°.

O²-De-Me: Apocavidine

[32728-76-0]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Corydalis tuberosa*. Cryst. + MeOH (MeOH). Mp 175°. Tenaciously solvated.

O³-De-Me: Isoapocavidine

C₂₀H₂₁NO₄ 339.39

Alkaloid from whole plants of *Dactylicapnos torulosa*. Powder. Mp 180-183°.

1-Nitro, O²-de-Me: 1-Nitroapocavidine

C₂₀H₂₀N₂O₆ 384.388

Alkaloid from *Corydalis saxicola*. Amorph. yellow powder. Mp 229-231°. λ_{max} 280 (MeOH).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 111; *CA*, **37**, 4738 (*Thalictrifoline*)
Yu, C.K. *et al.*, *Can. J. Chem.*, 1970, **48**, 3673 (*isol, pmr, ms, uv*)

Ninomiya, I. *et al.*, *J.C.S. Perkin I*, 1975, 1791 (*synth, pmr*)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (*cmr*)

Iwasa, K. *et al.*, *J.O.C.*, 1981, **46**, 4744-4750 (*synth, abs config*)

Iwasa, K. *et al.*, *Tet. Lett.*, 1981, 2333 (*Thalictrifoline, synth, abs config*)

Pai, B.R. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 607 (*synth*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1986, **42**, 675-680 (*biosynth*)

Rücker, G. *et al.*, *Phytochemistry*, 1994, **36**, 519 (*Isoapocavidine*)

Li, H.L. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 367-368 (*1-Nitroapocavidine*)

Li, W. *et al.*, *Acta Cryst. E*, 2006, **62**, o3137-o3138 (*cryst struct*)

Cheng, X. *et al.*, *Chem. Biodiversity*, 2008, **5**, 1335-1344 (*isol, pmr, cmr*)

Cavincidine

C-223

Struct. unknown. Alkaloid from roots of *Catharanthus roseus* (Apocynaceae).

Sulfate: Mp 238-239° dec. pK_a 7.85 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (*isol, uv, ir*)

Cavincine

C-224

[1355-33-5]

C₂₀H₂₄N₂O₂ 324.422

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Noncryst.

Sulfate:


Irregular plates (EtOH). Mp 275-277° dec. pK_a 6.9 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (*isol, ir, uv*)

Caymandimerine

C-225

[112899-83-9]



C₄₄H₃₈N₂O₉ 738.792

Alkaloid from the stem bark of a *Zanthoxylum* sp. collected in Grand Cayman (*Zanthoxylum spinosum* or *Zanthoxylum coriaceum*). λ_{max} 228 ; 283 ; 325 (MeOH).

Ng, K.M. *et al.*, *Phytochemistry*, 1987, **26**, 3251-3254 (*isol, pmr, ms*)

Ceanothine A

C-226

[1355-37-9]

C₃₀H₄₀N₄O₄ 520.67

Struct. unknown but appears to be identical with either Sanjoinine B in F-150 or *N*-Demethyladouetine Y in A-150. Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) (Rhamnaceae). Needles (CH₂Cl₂/Et₂O). Mp 256-259°. [α]_D -256 (c, 0.5 in CHCl₃).

Warnhoff, E.W. *et al.*, *Can. J. Chem.*, 1965, **43**, 2594-2602 (*isol, pmr*)

Warnhoff, E.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1970, **28**, 162 (*struct*)

Ceanothine C

C-227

[18658-42-9]

C₂₆H₃₈N₄O₄ 470.611

Struct. uncertain. Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) (Rhamnaceae). Needles (CHCl₃/Et₂O). Mp 223-229°. [α]_D²⁵ -368 (c, 1.01 in CHCl₃).

Warnhoff, E.W. *et al.*, *Can. J. Chem.*, 1965, **43**, 2594-2602 (*isol, uv, ms, pmr*)


Fehlhaber, H.-W. *et al.*, *Z. Anorg. Allg. Chem.*, 1968, **235**, 91 (*ms, struct*)

Servis, R.E. *et al.*, *J.A.C.S.*, 1969, **91**, 5619 (*ms, struct*)

Ceanothine D

C-228

[23926-97-8]



C₂₇H₄₀N₄O₄ 484.637


A drawing error in the paper gave an incorrect and biogenetically improbable structure (3-ethyl-3-methyl in place of 3-(1-methylpropyl)) which was registered in CAS. MF also incorrect. The struct. shown is as described in the text of the paper. Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) (Rhamnaceae). Cryst. (CHCl₃/Et₂O). Mp 227-229°. [α]_D -347 (CHCl₃).

Servis, R.E. *et al.*, *J.A.C.S.*, 1969, **91**, 5619-5624 (*isol, pmr, ms, struct*)

Ceanothine E

C-229

α-(Dimethylamino)-N-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]benzenepropanamide, 9CI [23926-98-9]



C₃₄H₄₀N₄O₄ 568.714


Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) (Rhamnaceae). Cryst. (CH₂Cl₂/Et₂O). Mp 238-239°. [α]_D -285 (CHCl₃).

Servis, R.E. *et al.*, *J.A.C.S.*, 1969, **91**, 5619 (*isol, ms, struct*)

Ceciline

C-230


4-[(6-Methoxy-9H-pyrido[3,4-b]indol-1-yl)methyl]phenol, 9CI. 4-(*p*-Hydroxybenzyl)-6-methoxy-β-carboline [76525-09-2]



C₁₉H₁₆N₂O₂ 304.348
Alkaloid from the trunk wood of *Aniba santalodora*. Cryst. (EtOH). Mp 224-228° dec.

Aguiar, L.M.G. *et al.*, *Phytochemistry*, 1980, **19**, 1859-1860 (*isol, synth, uv, pmr, ms*)


Celastramycin A C-231
2,3-Dichloro-4-(3-chloro-5-hexyl-2,6-dihydroxybenzoyl)pyrrole
[491600-94-3]



C₁₇H₁₈Cl₃NO₃ 390.692
Pyrrole antibiotic. Related to Antibiotic TAN 876B, A-1283. Prod. by *Streptomyces* sp. MaB-QuH-8 (DSM 14527). Antibacterial agent. Yellow wax. λ_{max} 222 ; 312 ; 355 (sh) (MeOH).

Pullen, C. *et al.*, *Planta*, 2002, **216**, 162-167 (*isol, pmr, cmr*)


Celastromelidine C-232



C₃₈H₄₆N₄ 558.808
Alkaloid from *Melodinus celastroides* (Apocynaceae). [α]_D²⁰ +117 (c, 1 in CHCl₃).

Mehri, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 372 (*isol, struct*)


Celastromeline C-233
[135626-52-7]



C₃₈H₄₆N₄ 558.808
Alkaloid from *Melodinus celastroides* (Apocynaceae). [α]_D²⁰ +190 (c, 1 in CHCl₃).

Mehri, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 372 (*isol, struct*)

Celenamide A C-234
[74144-98-2]



R = CH₂CH(CH₃)₂, R' = OH

C₃₄H₃₆BrN₅O₈ 722.591
Alkaloid from the sponge *Cliona celata*. [α]_D²⁵ +40 (c, 1.1 in Me₂CO) (as hexa-Ac).

Stonard, R.J. *et al.*, *J.O.C.*, 1980, **45**, 3687 (*isol, uv, ir, pmr, ms, struct*)

Celenamide B C-235
[74144-99-3]

As Celenamide A, C-234 with

R = -CH(CH₃)₂, R' = OH

C₃₃H₃₄BrN₅O₈ 708.564
Metab. from the sponge *Cliona celata*. [α]_D²⁵ +22 (c, 1.1 in Me₂CO) (as hexa-Ac). Isol. and elucidated as the Hexa-Ac deriv.

Stonard, R.J. *et al.*, *J.O.C.*, 1980, **45**, 3687 (*isol, uv, ir, pmr, ms, struct*)

Celenamide C C-236
[75472-51-4]


As Celenamide A, C-234 with

R = -CH₂CH(CH₃)₂, R' = H

C₃₄H₃₆BrN₅O₇ 706.592
Minor metab. from the sponge *Cliona celata*. [α]_D²⁵ +14 (c, 0.30 in Me₂CO) (as penta-Ac). Isol. and elucidated as the Penta-Ac deriv.

Stonard, R.J. *et al.*, *Can. J. Chem.*, 1980, **58**, 2121 (*isol, ir, pmr, ms, struct*)


Celenamide D C-237
[75472-50-3]



C₃₂H₃₄N₄O₁₁ 650.641
Minor metab. from the sponge *Cliona celata*. [α]_D²⁵ -25 (c, 0.54 in Me₂CO) (as nona-Ac). Isol. and elucidated as the Nona-Ac deriv.

Stonard, R.J. *et al.*, *Can. J. Chem.*, 1980, **58**, 2121 (*isol, ir, pmr, ms, struct*)


Celenamide E C-238
[202855-03-6]



C₂₈H₂₅BrN₄O₇ 609.432
Metab. of the sponge *Cliona chilensis*. Amorph. yellow solid (MeOH aq.). Mp 212-218° dec. [α]_D²⁵ -25 (c, 0.25 in MeOH). λ_{max} 226 (log ε 4.5); 296 (log ε 4.17); 340 (log ε 4.3) (MeOH).

Palermo, J.A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 488-490 (*isol, uv, ir, pmr, cmr*)

Celesticetin C-239
[2520-21-0]



C₂₄H₃₆N₂O₉S 528.622
Isol. from *Streptomyces caelestis* NRRL2418. Active against gram-positive organisms and plant pathogens. Sol. MeOH, butanol, CHCl₃, EtOAc, Me₂CO; poorly sol. H₂O, hexane, Et₂O. [α]_D²⁴ +126.6 (c, 0.5 in CHCl₃). Amphoteric base related structurally to Lincomycin. λ_{max} 240 (ε 10200); 310 (ε 4490) (EtOH/HCl) (Derep). λ_{max} 245 (ε 6880); 338 (ε 4860) (EtOH/KOH) (Derep). λ_{max} 239 (ε 10100); 307 (ε 4440) (95% EtOH) (Derep). λ_{max} 239 (E1%/1cm 184); 307 (E1%/1cm 80.6) (EtOH) (Berdy). λ_{max} 248 (E1%/1cm 180); 341 (E1%/1cm 104) (EtOH-NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 167 mg/kg. RH6125000
Oxalate: Mp 147-152°. [α]_D²⁴ +106.6 (c, 0.5 in H₂O).

N-De-Me: N-Demethylcelesticetin
[40736-31-0]

C₂₃H₃₄N₂O₉S 514.596
Prod. by *Streptomyces caelestis* mutant (NRRL5481; NRRL2418). Active against gram-positive bacteria. Cryst. Sol. MeOH. [α]_D²⁵ +112.5 (c, 1 in H₂O) (as hydrochloride). λ_{max} 238 ; 304 (MeOH) (Berdy). λ_{max} 241 ; 332 (NaOH) (Berdy).

O⁷-De-Me: O-Demethylcelesticetin
[39032-05-8]

C₂₃H₃₄N₂O₉S 514.596
Produced by mutant *Streptomyces caelestis* NRRL5520. Shows antibiotic props. Sol. H₂O, MeOH, EtOH. [α]_D²⁵

+115 (c, 0.85 in H₂O) (as hydrochloride). λ_{\max} 240 (ϵ 10200); 310 (ϵ 4490) (EtOH/HCl) (Derrep). λ_{\max} 245 (ϵ 6880); 338 (ϵ 4860) (EtOH/KOH) (Derrep). λ_{\max} 239 (ϵ 10100); 307 (ϵ 4440) (95% EtOH) (Derrep). λ_{\max} 238; 304 (MeOH) (Berdy). λ_{\max} 242; 352 (MeOH/NaOH) (Berdy).

N,O⁷-Di-de-Me: N,O-Didemethylcelesticetin

[40736-32-1]
C₂₂H₃₂N₂O₉S 500.569
Prod. by *Streptomyces caelestis* mutant (NRRL5481). Active against gram-positive bacteria. Needles (as hydrochloride). Sol. H₂O, MeOH. $[\alpha]_D^{25} +122$ (c, 1 in H₂O) (as hydrochloride). λ_{\max} 238 (ϵ 8830); 303 (ϵ 3600) (MeOH/HCl) (Derrep). λ_{\max} 241 (ϵ 5710); 332 (ϵ 4250) (MeOH/NaOH) (Derrep). λ_{\max} 238 (ϵ 8830); 303 (ϵ 3600) (MeOH as hydrogen halide salt) (Derrep).

ar-Deoxy, ar-amino: Celesticetin C

[42715-02-6]
C₂₄H₃₇N₃O₈S 527.638
Prod. by *Streptomyces caelestis* NRRL5320. Shows antibiotic props. Cryst. (as hydrochloride). Sol. H₂O, MeOH. $[\alpha]_D^{25} +123$ (c, 1 in H₂O) (as hydrochloride). Contains a 2-aminobenzoyl residue in place of 2-hydroxybenzoyl. λ_{\max} 228 (ϵ 9910); 270 (ϵ 843); 280 (sh) (ϵ 632) (0.5N HCl) (Derrep). λ_{\max} 242 (ϵ 6320); 321 (ϵ 2850) (0.5N NaOH) (Derrep). λ_{\max} 247 (ϵ 6480); 338 (ϵ 4480) (EtOH/acid salt) (Derrep). λ_{\max} 247; 338 (EtOH) (Berdy). λ_{\max} 243; 328 (H₂O) (Berdy). λ_{\max} 228; 270 (HCl) (Berdy).


De Boer, C. *et al.*, *Antibiot. Annu.*, 1954, 831 (isol)
Hoeksema, H. *et al.*, *Antibiot. Annu.*, 1954, 837 (isol)
Hoeksema, H. *et al.*, *J.A.C.S.*, 1964, **86**, 4224; 1968, **90**, 755 (struct)
Argoudelis, A.D. *et al.*, *J. Antibiot.*, 1972, **25**, 445; 1973, **26**, 7; 131; 1974, **27**, 642 (isol, ms, deriv)

Celliamine C-240

[1398-49-8]
C₂₁H₃₅NO₂ 333.513
Alkaloid from the roots of *Helleborus viridis*. Respiratory stimulant, convulsant. Small needles (EtOH aq.). Mp 127-131° (unsharp). A misprint gives the analysis as H45. The figures given correspond to H35.

Keller, O. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1928, **266**, 545-572

Celogentin A C-241

[370089-19-3]


C₄₅H₆₃N₁₃O₉ 930.074

Related to Moroidin, M-702. Constit. of the seeds of *Celosia argentea*. Tubulin polymn. inhibitor, antimitotic agent. Solid. $[\alpha]_D^{23} -43$ (c, 0.3 in MeOH aq.). λ_{\max} 226 (log ϵ 4.3); 283 (log ϵ 3.8) (MeOH).

L-Histidine amide: Celogentin B


[370089-21-7]
C₅₁H₇₀N₁₆O₁₀ 1067.215
Constit. of the seeds of *Celosia argentea*. Solid. $[\alpha]_D^{23} -32$ (c, 0.5 in MeOH aq.). λ_{\max} 225 (log ϵ 4.2); 282 (log ϵ 3.6) (MeOH).

L-Lysyl-L-histidine amide: Celogentin D

[602308-60-1]
C₅₇H₈₂N₁₈O₁₁ 1195.389
Constit. of the seeds of *Celosia argentea*. Amorph. solid. $[\alpha]_D^{24} -33$ (c, 0.4 in MeOH aq.). λ_{\max} 226 (log ϵ 4.3); 283 (log ϵ 3.8) (MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 2001, **66**, 6626-6633 (*Celogenins A,B*)
Suzuki, H. *et al.*, *Tetrahedron*, 2003, **59**, 5307-5315 (*Celogentin D*)


Celogentin C C-242

[370089-23-9]


C₅₀H₇₀N₁₄O₁₀ 1027.191
Constit. of the seeds of *Celosia argentea*. Tubulin polymerisation inhibitor. Antimitotic agent. Solid. $[\alpha]_D^{23} -54$ (c, 0.5 in MeOH aq.). λ_{\max} 226 (log ϵ 4.2); 283 (log ϵ 3.7) (MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 2001, **66**, 6626-6633

Celogentin G C-243



Absolute Configuration

C₄₈H₆₈N₁₄O₁₀ 1001.153
Homologue of Moroidin, M-702. Constit. of the seeds of *Celosia argentea*. Tubulin polymerisation inhibitor. Amorph. solid. $[\alpha]_D^{22} -47$ (c, 1 in MeOH aq.). λ_{\max} 226 (log ϵ 4.2); 283 (log ϵ 3.7) (MeOH).

L-Aspartic acid amide: Celogentin H

C₅₂H₇₃N₁₅O₁₃ 1116.241
Constit. of the seeds of *Celosia argentea*. Tubulin polymerisation inhibitor. Amorph. solid. $[\alpha]_D^{22} -40$ (c, 0.5 in MeOH aq.). λ_{\max} 225 (log ϵ 4.2); 282


(log ϵ 3.6) (MeOH).

L-Arginine amide: Celogentin J

C₅₄H₈₀N₁₈O₁₁ 1157.34
Constit. of the seeds of *Celosia argentea*. Tubulin polymerisation inhibitor. Amorph. solid. $[\alpha]_D^{22} -38$ (c, 0.4 in MeOH aq.). λ_{\max} 225 (log ϵ 4.2); 282 (log ϵ 3.6) (MeOH).

Suzuki, H. *et al.*, *Tetrahedron*, 2003, **59**, 5307-5315 (isol, pmr, cmr, ms)

Celogentin K C-244




C₄₇H₆₈N₁₄O₁₂ 1021.141
Cyclic peptide. Related to Moroidin, M-702. Constit. of the seeds of *Celosia argentea*. Amorph. solid. $[\alpha]_D^{24} -51$ (c, 0.5 in MeOH aq.). Struct. and MF given in ref. do not match. λ_{\max} 268 (ϵ 1760) (MeOH).

Suzuki, H. *et al.*, *Tetrahedron*, 2004, **60**, 2489-2495 (isol, pmr, cmr)

Cenocladamide C-245


N-(4-Hydroxy-3,5-dimethoxycinnamoyl)- Δ^2 -piperidin-4-one. 2,3-Dihydro-1-(4-hydroxy-3,5-dimethoxycinnamoyl)-4H-pyridin-4-one



C₁₆H₁₇NO₅ 303.314
Alkaloid from the leaves of *Piper cenocladum*. Pale yellow oil. λ_{\max} 220 (log ϵ 4.7); 300 (sh) (log ϵ 4.53); 350 (log ϵ 4.67); 660 (log ϵ 0.39) (MeOH).

Dodson, C.D. *et al.*, *Phytochemistry*, 2000, **53**, 51-54 (isol, pmr, cmr)


Cepaciachelin C-246

[184028-23-7]


C₂₄H₃₂N₄O₇ 488.539
Isol. from *Burkholderia cepacia* (*Pseudomonas cepacia*). Siderophore.

Barelmann, I. *et al.*, *Z. Naturforsch., C*, 1996, **51**, 627-630 (*isol, pmr*)

Cepaciamide A C-247
[173560-56-0]



Absolute Configuration

$C_{40}H_{74}N_2O_5$ 663.035


Struct. revised in 1999. Isol. from *Pseudomonas cepacia* D-202. Fungitoxic agent. Mp 98-102°. $[\alpha]_D^{22}$ -28.5 (c, 1.08 in $CHCl_3$).

Jiao, Y. *et al.*, *Tet. Lett.*, 1996, **37**, 1039-1042 (*isol*)

Toshima, H. *et al.*, *Tet. Lett.*, 1999, **40**, 935-938; 939-942 (*synth, struct, ir, pmr, cmr*)

Toshima, H. *et al.*, *Tetrahedron*, 1999, **55**, 5793-5808 (*synth*)

Cepaciamide B C-248




$C_{39}H_{72}N_2O_5$ 649.008

Isol. from *Pseudomonas cepacia* D-202. Fungitoxic agent. $[\alpha]_D^{21}$ -26.3 (c, 1.1 in $CHCl_3$).

Toshima, H. *et al.*, *Tet. Lett.*, 1999, **40**, 935-938; 939-942 (*isol, synth, pmr, cmr, struct*)

Cephalandole A C-249
[5543-37-3]
[915396-11-1]




$C_{16}H_{10}N_2O_2$ 262.267

Structure revised in 2008. Alkaloid from *Cephalanceropsis gracilis*. Bright yellow cryst. Mp 236-237° (*synthetic*). λ_{max} 255 (log ϵ 3); 282 (log ϵ 2.9); 394 (log ϵ 3) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr*)

Mason, J.J. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1447-1450 (*synth, struct*)

Cephalandole B C-250
[915315-44-5]




$C_{17}H_{14}N_2O_3$ 294.309

Alkaloid from *Cephalanceropsis gracilis*. Amorph. yellow powder. Mp 179.5-180° (*synthetic*). λ_{max} 263 (log ϵ 2.9); 347 (log ϵ 2.8) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr*)

Mason, J.J. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1447-1450 (*synth*)

Cephalandole C C-251
[915396-12-2]




$C_{23}H_{24}N_2O_9$ 472.451

Alkaloid from *Cephalanceropsis gracilis*. Amorph. yellowish powder. $[\alpha]_D$ +39.1 (c, 0.07 in MeOH). λ_{max} 207 (log ϵ 4.2); 260 (log ϵ 3.5); 297 (log ϵ 3.4); 351 (log ϵ 3.3) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr*)

Cephalozimine A C-252
[288300-70-9]



$C_{29}H_{39}NO_{10}$ 561.628

Ester of Drupacine, D-942. Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent. Amorph. solid. $[\alpha]_D$ -42 (c, 0.9 in MeOH). λ_{max} 289 (ϵ 1900) (MeOH).

4''-Deoxy: **Cephalozimine B**

[288300-72-1]


$C_{29}H_{39}NO_9$ 545.628

Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent.

Amorph. solid. $[\alpha]_D$ -125 (c, 4.9 in MeOH). Homologue of Drupangtonine, D-943. λ_{max} 291 (ϵ 1400) (MeOH).

Morita, H. *et al.*, *Tetrahedron*, 2000, **56**, 2929-2934

Cephalozimine G C-253
[160401-51-4]



Absolute Configuration

$C_{17}H_{21}NO_4$ 303.357

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. $[\alpha]_D$ -48 (c, 1.8 in MeOH). λ_{max} 212 (ϵ 4900); 239 (ϵ 2700); 290 (ϵ 2900) (MeOH).

2-Epimer: **Cephalozimine H**

[462076-18-2]


$C_{17}H_{21}NO_4$ 303.357

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. $[\alpha]_D$ +58 (c, 0.9 in MeOH). λ_{max} 210 (ϵ 7500); 235 (ϵ 3100); 291 (ϵ 3100) (MeOH).

Isono, N. *et al.*, *J.O.C.*, 1995, **60**, 115-119 (*synth, pmr, cmr*)

Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 5489-5495 (*isol, pmr, cmr, abs config*)

Cephalimysin A C-254
[952305-75-8]




Absolute Configuration

$C_{22}H_{25}NO_6$ 399.443

Prod. by a strain of *Aspergillus fumigatus* isol. from a marine fish. Cytotoxic agent. Pale yellow oil. $[\alpha]_D$ +3.5 (c, 0.11 in EtOH). λ_{max} 208 (log ϵ 3.88); 252 (log ϵ 3.95); 277 (log ϵ 3.74) (EtOH).

Yamada, T. *et al.*, *Tet. Lett.*, 2007, **48**, 6294-6296 (*isol, pmr, cmr*)

Cephalinone A C-255
[915396-07-5]




$C_{15}H_{13}N_3O_3$ 283.286

Alkaloid from *Cephalanceropsis gracilis*. Amorph. powder. $[\alpha]_D$ -7.6 (c, 0.18 in MeOH). λ_{max} 207 (log ϵ 3.8); 255 (log ϵ 3.2); 293 (log ϵ 3); 358 (log ϵ 2.7) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr, ms*)

Cephalinone B

[915396-08-6]


C₁₈H₁₆N₂O₅ 340.335

Alkaloid from *Cephalanceropsis gracilis*. Amorph. yellow powder. $[\alpha]_D -92.5$ (c, 0.04 in CHCl₃). λ_{\max} 263 (log ϵ 3.6); 315 (log ϵ 3.5); 385 (log ϵ 3.1) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr, ms*)

Cephalinone C

[915396-09-7]


C₁₇H₁₄N₂O₃ 294.309

Alkaloid from *Cephalanceropsis gracilis*. Amorph. yellow powder. $[\alpha]_D -8.4$ (c, 0.09 in CHCl₃). λ_{\max} 265 (log ϵ 2.7); 314 (log ϵ 2.6) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr, ms*)

Cephalinone D

[915396-10-0]

C₃₂H₁₈N₄O₄ 522.519


Alkaloid from *Cephalanceropsis gracilis*. Amorph. red powder. $[\alpha]_D -191.7$ (c, 0.04 in MeOH). λ_{\max} 228 (log ϵ 4); 274 (log ϵ 3.7); 308 (log ϵ 3.6); 424 (log ϵ 3.3); 546 (log ϵ 3.5) (MeOH).

Wu, P.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1467-1470 (*isol, pmr, cmr*)

C-256

Cephalocyclidine A

[421583-14-4]



Absolute Configuration

C₁₇H₁₉NO₅ 317.341

Alkaloid from the fruit of *Cephalotaxus harringtonia* var. *nana*. Plates (MeOH/C₆H₆) (as hydrochloride). Mp 205° dec. (hydrochloride). $[\alpha]_D -36$ (c, 0.8 in MeOH). λ_{\max} 241 (ϵ 1300); 293 (ϵ 1700) (MeOH).


Kobayashi, J. *et al.*, *J.O.C.*, 2002, **67**, 2283-2286 (*isol, uv, cd, pmr, cmr, cryst struct*)

Cephalofortuneine

C-260

1,6-Didehydro-3,15,16-trimethoxy-C-homoerythrinan-2,7-diol, 9CI. Alkaloid CF 15

[68156-55-8]

C₂₀H₂₇NO₅ 361.437

Minor alkaloid from *Cephalotaxus fortunei* (Cephalotaxaceae).

O²-Et: 2-O-Ethylcephalofortuneine

[128387-09-7]

C₂₂H₃₁NO₅ 389.491

Alkaloid from *Cephalotaxus fortunei* (Cephalotaxaceae).

2-Epimer: 2-Epicephalofortuneine

[97805-96-4]

C₂₀H₂₇NO₅ 361.437

Alkaloid from *Cephalotaxus fortunei*. Component of San Jian Shan. Cryst. Mp 80-83°. $[\alpha]_D^{25} +12.1$ (c, 0.15 in CHCl₃).

2-Epimer, O²-Et: 2-Epi-2-O-ethylcephalofortuneine

[128309-01-3]

C₂₂H₃₁NO₅ 389.491


Alkaloid from *Cephalotaxus fortunei* (Cephalotaxaceae).

Ma, G. *et al.*, *Huaxue Xuebao*, 1978, **36**, 129; 1989, **31**, 137; *CA*, **89**, 193837x; **113**, 74722k (*Cephalofortuneine, 2-O-Ethylcephalofortuneine*)

Lin, W. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 283-287 (*2-Epicephalofortuneine*)

Cephalostatin 2

[116199-48-5]



R = H

C-259

C₅₄H₇₄N₂O₁₁ 927.185

Diteroidal alkaloid from the hemichordate marine worm *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Needles (EtOAc/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 350°. $[\alpha]_D +111$ (c, 0.07 in MeOH). λ_{\max} 290 (ϵ 13700); 308 (sh) (EtOH) (Derep).

14'β,15'β-Epoxyde: Cephalostatin 4

[116229-58-4]

C₅₄H₇₄N₂O₁₂ 943.185

Alkaloid from *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 350°. $[\alpha]_D +89$ (c, 0.11 in MeOH). λ_{\max} 290 (ϵ 13700); 308 (sh) (EtOH) (Derep).

9'-Deoxy: Cephalostatin 1

[112088-56-9]

C₅₄H₇₄N₂O₁₀ 911.186

Alkaloid from *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Needles (EtOAc/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 326° dec. $[\alpha]_D +102$ (c, 0.04 in MeOH). λ_{\max} 290 (ϵ 13700); 308 (sh) (EtOH) (Derep). λ_{\max} 289 (ϵ 15233) (EtOH) (Berdy).

9'-Deoxy, 8'-hydroxy, 9',11'-didehydro,

14'α,15'α-epoxyde: Cephalostatin 14

[159602-64-9]

C₅₄H₇₂N₂O₁₂ 941.169

Constit. of *Cephalodiscus gilchristi*. Cell growth inhibitor. Amorph. powder. $[\alpha]_D +80.9$ (c, 0.11 in MeOH). λ_{\max} 230 (log ϵ 3.69); 288 (log ϵ 3.9); 305 (sh) (MeOH).

26-Deoxy: Cephalostatin 17

[170591-44-3]

C₅₄H₇₄N₂O₁₀ 911.186

Constit. of *Cephalodiscus gilchristi*. Cell growth inhibitor. Amorph. solid. $[\alpha]_D +70$ (c, 0.7 in MeOH). λ_{\max} 288 (ϵ 15000); 305 (sh) (MeOH).

1α-Methoxy: Cephalostatin 10

[153698-88-5]

C₅₅H₇₆N₂O₁₂ 957.212

Alkaloid from *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Mp > 300°. $[\alpha]_D +80$ (c, 0.17 in MeOH). λ_{\max} 289 (log ϵ 4.08); 304 (sh) (MeOH).

1α-Methoxy, 9'-deoxy: Cephalostatin 18

[211050-18-9]

C₅₅H₇₆N₂O₁₁ 941.212

Alkaloid from *Cephalodiscus gilchristi*. Cell growth inhibitor. Amorph. solid. Mp > 300°. $[\alpha]_D^{25} +95$ (c, 0.06 in MeOH). λ_{\max} 288 (log ϵ 4.06); 308 (MeOH).

1'α-Methoxy: Cephalostatin 11

[153698-87-4]

C₅₅H₇₆N₂O₁₂ 957.212

Alkaloid from *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Mp > 300°. $[\alpha]_D +75$ (c, 0.13 in MeOH). λ_{\max} 288 (log ϵ 3.96); 305 (sh) (MeOH).

1'α-Methoxy, 9'-deoxy: Cephalostatin 19

[211050-19-0]

C₅₅H₇₆N₂O₁₁ 941.212

Alkaloid from *Cephalodiscus gilchristi*. Amorph. solid. Mp > 320°. $[\alpha]_D^{25} +67$ (c, 0.05 in MeOH). λ_{\max} 288 (log ϵ 3.99); 308 (MeOH).

- Pettit, G.R. *et al.*, *Chem. Comm.*, 1988, 865-867; 1440 (*Cephalostatins 2-4*)
 Pettit, G.R. *et al.*, *J.A.C.S.*, 1988, **110**, 2006 (*Cephalostatin 1*)
Pat. Coop. Treaty (WIPO), 1989, 89 08 655 (*Cephalostatins 1,2,3,4*)
 Pettit, G.R. *et al.*, *Can. J. Chem.*, 1994, **72**, 2260-2267 (*Cephalostatin 14*)
 Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1994, **57**, 52; 1998, **61**, 955-958 (*Cephalostatins 10,11,18,19*)
 Pettit, G.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 2027-2032 (*Cephalostatin 17*)
 Pettit, G.R. *et al.*, *J.O.C.*, 1995, **60**, 608-613 (*Cephalostatin 14*)

Cephalostatin 3 C-262


[116199-49-6]
 As Cephalostatin 2, C-261 with
 R = CH₃
 C₅₅H₇₆N₂O₁₁ 941.212
 Disteroidal alkaloid from the hemichordate marine worm *Cephalodiscus gilchristi*. Powerful cell growth inhibitor. Shows antitumour activity. Needles (EtOAc/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 350°. [α]_D +99 (c, 0.15 in MeOH). λ_{max} 290 (ε 13700); 308 (sh) (ε) (EtOH) (Derep).

9'-Deoxy, 8'-hydroxy, 9',11'-didehydro, 14'α,15'α-epoxide: **Cephalostatin 15**
 [159602-65-0]
 C₅₅H₇₄N₂O₁₂ 955.196
 Constit. of *Cephalodiscus gilchristi*. Shows antitumour activity. Amorph. powder. Mp > 300°. [α]_D +71.5 (c, 0.34 in MeOH). λ_{max} 202 (log ε 4.25); 228 (sh); 288 (log ε 3.96); 305 (sh) (MeOH).

Pettit, G.R. *et al.*, *Chem. Comm.*, 1988, 865-867 (*uv, ir, cmr, struct*)
Pat. Coop. Treaty (WIPO), 1989, 89 08 655 (*Cephalostatin 3*)
 Pettit, G.R. *et al.*, *Can. J. Chem.*, 1994, **72**, 2260-2267 (*Cephalostatin 15*)

Cephalostatin 5 C-263

[121071-11-2]



C₅₄H₇₂N₂O₁₀ 909.17
 Disteroidal alkaloid from the Indian Ocean (South African) marine worm *Cephalodiscus gilchristi*. Possesses lymphocytic leukaemia (murine P388) cell growth inhibitory activity (less potent than cephalostatins 1-4). Mp 350°. [α]_D +100 (c, 0.02 in MeOH). λ_{max} 289 (ε 10000); 310 (sh) (MeOH) (Derep).

Pettit, G.R. *et al.*, *Can. J. Chem.*, 1989, **67**, 1509-1513 (*isol, uv, struct*)

Cephalostatin 6 C-264


[121038-34-4]
 As Cephalostatin 5, C-263 with
 R = H

C₅₃H₇₀N₂O₁₀ 895.143
 Disteroidal alkaloid from the Indian Ocean marine worm *Cephalodiscus gilchristi*. Possesses lymphocytic leukaemia (murine P388) cell growth inhibitory activity (less potent than cephalostatins 1-4). Mp 350°. [α]_D +100 (c, 0.01 in MeOH). λ_{max} 289 (ε 10000); 310 (sh) (ε) (MeOH) (Derep).

Pettit, G.R. *et al.*, *Can. J. Chem.*, 1989, **67**, 1509-1513 (*isol, uv, pmr, cmr, struct*)

Cephalostatin 7 C-265

[138605-82-0]




C₅₄H₇₆N₂O₁₁ 929.201
 Constit. of *Cephalodiscus gilchristi*. Cytotoxic agent. Amorph. powder. Mp 315° dec. [α]_D +106 (c, 0.244 in MeOH). λ_{max} 286 (ε 17400) (MeOH).

Pettit, G.R. *et al.*, *J.O.C.*, 1992, **57**, 429-431 (*isol, pmr, cmr*)
 Jeong, J.U. *et al.*, *J.A.C.S.*, 1999, **121**, 2071-2084 (*synth*)

Cephalostatin 8 C-266

[138605-83-1]




C₅₅H₇₈N₂O₁₀ 927.229
 Constit. of *Cephalodiscus gilchristi*. Cytotoxic agent. Amorph. powder. Mp 313° (dec.). [α]_D +110 (c, 0.1 in MeOH). λ_{max} 286 (ε 20800); 310 (sh) (MeOH).

Pettit, G.R. *et al.*, *J.O.C.*, 1992, **57**, 429-431 (*isol, pmr, cmr*)

Cephalostatin 9 C-267

[138628-96-3]




C₅₄H₇₆N₂O₁₁ 929.201

Constit. of *Cephalodiscus gilchristi*. Amorph. powder. Mp 307° (dec.). [α]_D +105 (c, 0.5 in MeOH). λ_{max} 286 (ε 18600); 310 (sh) (MeOH).

Pettit, G.R. *et al.*, *J.O.C.*, 1992, **57**, 429-431 (*isol, pmr, cmr*)

Cephalostatin 13 C-268

[158371-97-2]



C₅₄H₇₆N₂O₁₃ 961.2
 Alkaloid from the marine worm *Cephalodiscus gilchristi*. Antineoplastic agent. Amorph. solid. Mp > 300°. [α]_D²⁰ +108.1 (c, 0.07 in MeOH). λ_{max} 287 (log ε 3.89); 308 (sh) (log ε 3.46) (MeOH).


1'-Deoxy: **Cephalostatin 12**

[158371-96-1]
 C₅₄H₇₆N₂O₁₂ 945.201
 Alkaloid from *Cephalodiscus gilchristi*. Antineoplastic agent. Amorph. solid. Mp > 300°. [α]_D²⁰ +157.5 (c, 0.4 in MeOH). λ_{max} 288 (log ε 4.16); 308 (sh) (log ε 3.89) (MeOH).

Pettit, G.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 1507-1512 (*isol, pmr, cmr, uv*)
 Jeong, J.U. *et al.*, *J.A.C.S.*, 1999, **121**, 2071-2084 (*synth*)

Cephalostatin 16 C-269

[170591-43-2]




C₅₄H₇₄N₂O₁₀ 911.186
 Alkaloid from the marine worm *Cephalodiscus gilchristi*. Antineoplastic agent. Amorph. powder. Mp > 300°. [α]_D +55 (c, 1.73 in MeOH). λ_{max} 288 (ε 11600); 305 (sh) (MeOH).

Pettit, G.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 2027-2032 (*isol, pmr, cmr*)

Cephalotaxidine C-270

[182889-14-1]



C₅₈H₇₄N₂O₁₉ 1103.225


Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*. Amorph. solid. $[\alpha]_D^{25}$ -172 (c, 0.10 in MeOH). First example of a dimeric *Cephalotaxus* alkaloid. λ_{\max} 291 (log ϵ 3.85) (MeOH). λ_{\max} 291 (ϵ 7080) (MeOH) (Berdy).

Takano, I. *et al.*, *Tet. Lett.*, 1996, **37**, 7053-7054 (isol, uv, ir, pmr, cmr, struct)

Cephalotaxine

C-271

[83136-88-3]



(-)-form

C₁₈H₂₁NO₄ 315.368**(+)-form**

Ac: (+)-Acetylcephalotaxine. Alkaloid CF 5

[69127-15-7]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Cephalotaxus fortunei* and *Cephalotaxus hainanensis* (Cephalotaxaceae). Mp 140°. $[\alpha]_D^{25}$ +102 (CHCl₃). $[\alpha]_D^{25}$ +130 (CHCl₃).

(-)-form [24316-19-6]

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus harringtonia* var. *drupacea*, *Cephalotaxus fortunei*, *Cephalotaxus hainanensis*, *Cephalotaxus wilsoniana*, *Cephalotaxus sinensis* and *Cephalotaxus oliveri* (Cephalotaxaceae). Mp 132-133°. $[\alpha]_D^{25}$ -204 (c, 1.8 in CHCl₃). λ_{\max} 238 (ϵ 3630); 290 (ϵ 3560) (EtOH) (Berdy).

Perchlorate: Mp 213-216° dec.

α -N-Oxide: *Cephalotaxine* α -N-oxide

[48856-88-8]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the fruit of *Cephalotaxus fortunei*. Amorph. solid. $[\alpha]_D^{21}$ -131 (c, 0.5 in CHCl₃).

β -N-Oxide: *Cephalotaxine* β -N-oxide

[488856-87-7]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the fruit of *Cephalotaxus fortunei*. Amorph. solid. $[\alpha]_D^{21}$ -221 (c, 0.5 in CHCl₃).

Ac: (-)-Acetylcephalotaxine

[24274-60-0]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Cephalotaxus fortunei* and *Cephalotaxus wilsonia* (Cephalotaxaceae). Cryst. (Et₂O). Mp 140-142°. $[\alpha]_D^{25}$ -97 (c, 2.2 in CHCl₃).

O-De-Me: Demethylcephalotaxine

[39707-71-6]

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Cephalotaxus harringtonia* var. *drupacea* and *Cephalotaxus fortunei* (Cephalotaxaceae). Also obt. by mild acid hydrol. of Cephalotaxine, C-271. Cryst. (EtOH). Mp 109-111°. $[\alpha]_D^{25}$ -110 (c, 0.28 in CHCl₃). 2-Ketone.

3-Ketone: *Cephalotaxinone*

[38750-57-1]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus fortunei* and *Cephalotaxus hainanensis* (Cephalotaxaceae). Cryst. (Et₂O). Mp 198-200° dec. $[\alpha]_D^{24}$ -146 (c, 0.63 in CHCl₃).

3,4-Didehydro, O-de-Me, 3-O- β -D-glucopyranoside: *Cephelezomine J*

C₂₃H₂₇NO₉ 461.468

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. $[\alpha]_D^{25}$ +80 (c, 0.5 in MeOH). 2-Ketone. λ_{\max} 209 (ϵ 8100); 235 (ϵ 4400); 250 (ϵ 3600); 293 (ϵ 2900); 312 (ϵ 2600) (MeOH).

4-Hydroxy: 3-Hydroxycephalotaxine. 4-Hydroxycephalotaxine

[84567-08-8]

C₁₈H₂₁NO₅ 331.368

Alkaloid from *Cephalotaxus fortunei* (Cephalotaxaceae). Pale yellow cryst. (MeOH). Mp 135-137°. $[\alpha]_D^{27}$ +120 (c, 0.025 in MeOH). Stereochem. not defined.

11-Hydroxy: 11-Hydroxycephalotaxine

[49686-55-7]

C₁₈H₂₁NO₅ 331.368

Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*, *Cephalotaxus fortunei* and *Cephalotaxus sinensis* (Cephalotaxaceae). Cryst. (MeOH). Mp 235-242° dec. $[\alpha]_D^{26}$ -139 (c, 0.56 in CHCl₃).

11 β -Hydroxy, β -N-oxide: 11 β -Hydroxycephalotaxine β -N-oxide

[488856-89-9]

C₁₈H₂₁NO₆ 347.367

Alkaloid from the fruit of *Cephalotaxus fortunei*. Amorph. solid. $[\alpha]_D^{21}$ -94 (c, 0.5 in CHCl₃).

8-Oxo: *Cephalotaxinamide*. 8-Oxocephalotaxine

[80797-04-2]

C₁₈H₁₉NO₅ 329.352

Alkaloid from *Cephalotaxus hainanensis* (Cephalotaxaceae). Mp 230°.

3-Epimer: *Epicephalotaxine*

[39707-72-7]

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Cephalotaxus fortunei* and *Cephalotaxus hainanensis* (Cephalotaxaceae). Mp 136-137°. $[\alpha]_D^{25}$ -150 (c, 0.8 in CHCl₃).

(\pm)-form [38848-21-4]

Synthetic. Cryst. (Et₂O). Mp 116-118°.

3-Ketone: [38848-26-9]

Synthetic. Cryst. (EtOAc). Mp 180-183° (178-180°).

8-Oxo: [114942-84-6]

Synthetic. Mp 230° dec Mp 258-259°.

Paudler, W.W. *et al.*, *J.O.C.*, 1963, **28**, 2194-2197; 1973, **38**, 2110-2112 (*Cephalotaxine*, *Epicephalotaxine*, *Cephalotaxinone*, *Demethylcephalotaxine*, isol, uv, ir, pmr)

Powell, R.G. *et al.*, *Tet. Lett.*, 1969, 4081-4084 (pmr, struct)

Abraham, D.J. *et al.*, *Tet. Lett.*, 1969, 4085-4086 (cryst struct)

Powell, R.G. *et al.*, *Phytochemistry*, 1972, **11**, 3317-3320 (*Acetylcephalotaxine*, *Cephalotaxinone*)

Powell, R.G. *et al.*, *J.O.C.*, 1974, **39**, 676-680 (*11-Hydroxycephalotaxine*)

Arora, S.K. *et al.*, *J.O.C.*, 1974, **39**, 1269-1271; 1976, **41**, 551-557 (cryst struct, config)

Weinreb, S.M. *et al.*, *J.A.C.S.*, 1975, **97**, 2503-2506 (*Cephalotaxine*, *Cephalotaxinone*, synth, uv, ir, pmr)

Semmelhack, M.F. *et al.*, *J.A.C.S.*, 1975, **97**, 2507-2516 (*Cephalotaxine*, *Cephalotaxinone*, synth)

Spencer, G.F. *et al.*, *J. Chromatogr.*, 1976, **120**, 335-341 (ms, glc)

Ma, G.-E. *et al.*, *Huaxue Xuebao*, 1977, **35**, 201-208; *CA*, **90**, 19013m

(*Acetylcephalotaxine*)

Schwab, J.M. *et al.*, *J.A.C.S.*, 1977, **99**, 2368-2371 (biosynth)

Weisleder, D. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 114-115 (cmr)

Xue, Z. *et al.*, *Yaoxue Xuebao*, 1981, **16**, 752-756; *CA*, **96**, 82690u (*Cephalotaxinamide*)

Ma, G.-E. *et al.*, *J. Nat. Prod.*, 1982, **45**, 585-589 (*4-Hydroxycephalotaxine*)

Yasuda, S. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 4229 (synth, ir, pmr, *Cephalotaxinamide*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1988, **53**, 3439-3450 (*Cephalotaxine*, 8-Oxocephalotaxine, synth)

Ishibashi, H. *et al.*, *Chem. Comm.*, 1990, 1436-1437 (synth)

Burkholder, T.P. *et al.*, *J.A.C.S.*, 1990, **112**, 9601-9613 (synth)

Fang, F.G. *et al.*, *J.O.C.*, 1990, **55**, 831-838 (synth)

Ikeda, M. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 276-281 (synth)

Lin, X. *et al.*, *J.A.C.S.*, 1994, **116**, 9791-9792 (synth)

Isono, N. *et al.*, *J.O.C.*, 1995, **60**, 115-119 (synth)

Nagasaka, T. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 191-194 (synth)

Ikeda, M. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 983-987 (synth)

Tietze, L.F. *et al.*, *J.A.C.S.*, 1999, **121**, 10264-10269 (synth)

Koseki, Y. *et al.*, *Org. Lett.*, 2002, **4**, 885-888 (synth)

Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 5489-5495 (*Cephelezomine J*)

Bocar, M. *et al.*, *J. Nat. Prod.*, 2003, **66**, 152-154 (*N-oxides*)

Planas, L. *et al.*, *J.O.C.*, 2004, **69**, 3087-3092 (synth)

Liu, Q. *et al.*, *J.O.C.*, 2007, **72**, 7352-7358 (synth)

Li, W.-D.Z. *et al.*, *Org. Lett.*, 2007, **9**, 1211-1214 (synth)


Hameed, A. *et al.*, *J.O.C.*, 2008, **73**, 8045-8048 (synth)

Taniguchi, T. *et al.*, *Org. Lett.*, 2008, **10**, 4129-4131 (synth)

Cepharadione A

C-272

7-Methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline-5,6(7H)-dione, 9CI [55610-01-0]

C₁₈H₁₁NO₄ 305.289

Alkaloid from *Stephania cepharantha*,

Stephania sasakii, *Piper sanctum*, *Piper auritum* (Veracruz pepper) and *Aristolochia indica*. Orange-yellow needles (CHCl₃/EtOH). Mp 340-342° dec.

N-De-Me: Norcepharadione A

[107882-42-8]
C₁₇H₉NO₄ 291.262

Alkaloid from the bark of *Oncodostigma monosperma*.

N-De-Me, N-methoxy: N-Methoxynorcepharadione A. Artabotrine

[166833-81-4]
C₁₈H₁₁NO₅ 321.289

Alkaloid from the stem bark of *Artabotrys zeylanica*. Cytotoxic. Orange-yellow rods. Mp 287-289°. λ_{max} 220 (ε 20900); 235 (ε 22390); 280 (ε 5750); 300 (ε 9800); 310 (ε 11000) (MeOH) (Berdy).

Akasu, M. *et al.*, *Tet. Lett.*, 1974, 3609 (*ir, uv, ms, pmr, struct*)

Hänsel, R. *et al.*, *J. Nat. Prod.*, 1975, **38**, 529; *Phytochemistry*, 1976, **15**, 1323 (*isol, uv, ir, ms*)

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1981, **101**, 431; *CA*, **95**, 204236c (*isol*)

Achari, B. *et al.*, *Heterocycles*, 1982, **19**, 1203 (*occur*)

Cavé, A. *et al.*, *Plant. Med. Phytother.*, 1986, **20**, 251; *CA*, **106**, 172905x (*Norcepharadione A*)


Wijeratne, E.M.K. *et al.*, *Tetrahedron*, 1995, **51**, 7877 (*Artabotrine*)

Ding, H.X. *et al.*, *Chin. Chem. Lett.*, 2006, **17**, 5-8 (*Artabotrine, synth*)

Cepharadione B

C-273

1,2-Dimethoxy-6-methyl-4H-dibenzo[de,g]quinoline-4,5(6H)-dione, 9CI [55610-02-1]



C₁₉H₁₅NO₄ 321.332

Alkaloid from the callus tissue of *Stephania cepharantha* and the woody roots of *Piper auritum* (Veracruz pepper). Also detected in the woody roots of *Piper sanctum* (Menispermaceae, Piperaceae). Possible biogenetic intermediate of Aristololactams. Orange needles (EtOH). Mp 267-268° (255-262°, 263-264°).

N-De-Me: Norcepharadione B

[57576-41-7]
C₁₈H₁₃NO₄ 307.305

Alkaloid from the callus tissue of *Stephania cepharantha* (Menispermaceae). Shows platelet aggregation inhibitory activity. Orange cryst. Mp 304-307° dec.

O²-De-Me: Aristolodione. Piperadione

[109771-09-7]
C₁₈H₁₃NO₄ 307.305

Alkaloid from the leaves and stems of *Aristolochia chinensis* and *Piper longum* (long pepper). Cryst. (CHCl₃/MeOH).

Mp 273-276° dec.

O²-De-Me, Ac:

Cryst. (C₆H₆/MeOH). Mp 244-248° dec.

O²,N-Di-de-Me: 2-Hydroxy-1-methoxy-4H-dibenzo[de,g]quinoline-4,5-dione, 9CI. 4,5-Dioxodehydroasimilobine.

Noraristolodione
[82644-81-3]
C₁₇H₁₁NO₄ 293.278

Alkaloid from the roots of *Aristolochia indica*. Red needles (CHCl₃/MeOH). Mp 310-312°. λ_{max} 246 (ε 50100); 292 (sh) (ε 13800); 305 (ε 18200); 318 (ε 19100); 459 (ε 17000) (EtOH).

7-Chloro, N-de-Me: 7-Chloro-6-demethylcepharadione B

[149682-95-1]
C₁₈H₁₂ClNO₄ 341.75

Alkaloid from *Houttuynia cordata* (Yu Xing Cao) (Saurauaceae).

1-De-methoxy, O²,N-di-de-Me: De-methoxy-4,5-dioxodehydroasimilobine

[138690-43-4]
C₁₆H₉NO₃ 263.252

Alkaloid from *Monocyclanthus vignei*. Orange needles (MeOH). Mp 283-285°.

Akasu, M. *et al.*, *Tet. Lett.*, 1974, 3609 (*uv, ir, pmr, ms, struct*)

Hänsel, R. *et al.*, *J. Nat. Prod.*, 1975, **38**, 529 (*isol, uv, ir, pmr, ms*)

Akasu, M. *et al.*, *Phytochemistry*, 1975, **14**, 1673 (*uv, ir, pmr, ms, struct, Norcepharadione B*)

Hänsel, R. *et al.*, *Phytochemistry*, 1976, **15**, 1323 (*occur*)

Saá, J.M. *et al.*, *Tet. Lett.*, 1976, 601 (*synth, uv, ir, ms*)

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1980, **100**, 337; *CA*, **93**, 95454h (*synth*)

Achari, B. *et al.*, *Heterocycles*, 1982, **19**, 1203 (*Dioxodehydroasimilobine*)

Urzua, A. *et al.*, *J. Nat. Prod.*, 1987, **50**, 305 (*Aristolodione*)

Desai, S.J. *et al.*, *Phytochemistry*, 1988, **27**, 1511 (*Aristolodione*)

Achenbach, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1331 (*Demethoxydioxodehydroasimilobine*)

Atanes, N. *et al.*, *J.O.C.*, 1991, **56**, 2984 (*synth, Cepharadione B, Norcepharadione B*)

Jong, T.T. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1993, **40**, 301; *CA*, **119**, 135624c (*7-Chloro-6-demethylcepharadione B*)


Suau, R. *et al.*, *Tetrahedron*, 1996, **52**, 11307 (*synth*)

Chia, Y.-C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1160-1163 (*activity*)

Cepharamine

C-274

7,8-Didehydro-4-hydroxy-3,7-dimethoxy-17-methylhasubanan-6-one, 9CI



C₁₉H₂₃NO₄ 329.395

(-)-form [15444-26-5]

Alkaloid from *Stephania cepharantha*

and *Stephania sasakii* (Menispermaceae). Mp 186-187°. [α]_D -248 (CHCl₃).

(±)-form [24048-61-1]

Synthetic. Oil.

Hydrobromide:

Flakes (Me₂CO). Mp 243-246° dec.

Tomita, M. *et al.*, *Tet. Lett.*, 1966, 6229 (*uv, ir, pmr, struct*)

Inubushi, Y. *et al.*, *Tet. Lett.*, 1969, 1611; *Chem. Pharm. Bull.*, 1971, **19**, 1820 (*synth, ir, pmr*)

Keely, S.L. *et al.*, *Tetrahedron*, 1970, **26**, 4729 (*synth*)

Kametani, T. *et al.*, *Chem. Ind. (London)*, 1972, 538 (*synth, ir, pmr*)


Schultz, A.G. *et al.*, *J.A.C.S.*, 1998, **120**, 8259-8260 (*synth*)

Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (*synth, rev*)

Cepharanthine

C-275

6',12'-Dimethoxy-2,2'-dimethyl-6,7-[methylenebis(oxy)]joxycanthan, 9CI. 12-O-Methylcephanoline [481-49-2]



C₃₇H₃₈N₂O₆ 606.717

Alkaloid from the root tubers of *Stephania cepharantha* and *Stephania erecta*, and from the roots of *Stephania sasakii* and *Stephania epigaeae* (Menispermaceae). Antihepatitic, radioprotective agent. Shows *in vitro* antineoplastic activity against HeLa cells. May potentiate the effects of other antineoplastic agents by restoring leucocyte count. Curarising agent. Shows antibacterial activity. Yellow amorph. powder. Mp 145-155°. [α]_D²⁰ +277 (CHCl₃). Log P 8.19 (uncertain value) (calc).

▶ LD₅₀ (mus, ipr) 125 mg/kg. FK0527000 *Methiodide*: Mp 268° (dec.).

N²-β-Oxide: Cepharanthine 2'-β-N-oxide

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from tuberous roots of *Stephania suberosa* (Menispermaceae). [α]_D +152 (c, 0.22 in MeOH).

N²-De-Me: 2-Norcepharanthine

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from tuberous roots of *Stephania suberosa* (Menispermaceae). [α]_D +318 (c, 0.25 in MeOH).

N²-De-Me: 2'-Norcepharanthine

[123854-64-8]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the tubers of *Stephania piirrii* (Menispermaceae). [α]_D +206 (c, 0.24 in CHCl₃).

O¹²-De-Me: Cepharanoline

[27686-34-6]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the root tubers of *Stephania cepharantha* (Menispermaceae). Mp 270° dec. [α]_D³⁵ +319 (CHCl₃).

O¹²-de-Me, N²-de-Me: 2-Norcepharanoline

[123854-65-9]
 $C_{35}H_{34}N_2O_6$ 578.663
 Alkaloid from the tubers of *Stephania piri* (Menispermaceae). $[\alpha]_D^{25} +257$ (c, 0.17 in $CHCl_3$).


1-Epimer, N²-de-Me: 2-Norisocepharantine

$C_{36}H_{36}N_2O_6$ 592.69
 Minor alkaloid from the tubers of *Stephania piri* (Menispermaceae). $[\alpha]_D^{25} -84$ (c, 0.25 in $CHCl_3$).

Kondo, H. *et al.*, *Ber.*, **1938**, **71**, 2553 (struct)
 Tomita, M. *et al.*, *Pharm. Bull.*, 1954, **2**, 375 (struct)
 Bick, I.R.C. *et al.*, *J.C.S.*, 1961, 1896 (pmr)
 Kunimoto, J. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 981; *CA*, **58**, 4613f (abs config)
 Battersby, A.R. *et al.*, *J.C.S.*, 1965, 2239 (ord)
 Tomita, M. *et al.*, *Tet. Lett.*, 1967, 1201 (synth, pmr)
 Tomita, M. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1678; *CA*, **72**, 107846u (Cepharanoline)
 Kunimoto, J. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1691; *CA*, **73**, 4072e (isol)
 Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592 (ms)
 Kuroda, H. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 2413-2420 (activity)
 Patra, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 424 (derivs)
 Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1989, **52**, 846 (2'-Norcepharantine, 2-Norisocepharantine, 2-Norcepharoline)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CCX550

Cephasamine

[175617-21-7]




$C_{20}H_{23}NO_5$ 357.405
 Alkaloid from tubers of *Stephania cepharantha*. Prisms (Et_2O). Mp 142-144°. $[\alpha]_D^{28} +105$ (c, 0.34 in $CHCl_3$).

Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 1996, **59**, 476 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Ceratamine A

[634151-15-8]



$C_{17}H_{16}Br_2N_4O_2$ 468.147
 Alkaloid from the marine sponge *Pseudoceratina* sp. Antimitotic agent. Yellow cryst. (MeOH). Mp 236°.


N⁶-De-Me: Ceratamine B

[634151-16-9]
 $C_{16}H_{14}Br_2N_4O_2$ 454.12
 Alkaloid from a *Pseudoceratina* sp. Yellow cryst. (MeOH). Mp 242°.

Manzo, E. *et al.*, *Org. Lett.*, 2003, **5**, 4591-4594 (isol, pmr, cmr)

Ceratocapnidine

[132160-47-5]



$C_{21}H_{23}NO_7$ 401.415
 Alkaloid from *Ceratocapnos palaestinus* (Papaveraceae). Amorph.


Me ether: Ceratocapnine

[132160-48-6]
 $C_{22}H_{25}NO_7$ 415.442
 Alkaloid from *Ceratocapnos palaestinus* (Papaveraceae). Amorph.

Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1006 (isol, struct)

Ceratonicine


$C_{21}H_{21}NO_7$ 399.399
 Alkaloid from *Ceratocapnos palaestinus* (Papaveraceae). Amorph.



Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1006 (isol, uv, ir, pmr, ms struct)

Ceratospongamide

[259794-28-0]



$C_{41}H_{49}N_7O_6S$ 767.947

C-278

C-279

C-280

Isol. as two stable conformational isomers. Isol. from the red alga *Ceratodictyon spongiosum* containing the symbiotic sponge *Signadocia symbiotica*. Anti-inflammatory agent.

cis,cis-form

Amorph. solid. $[\alpha]_D -190$ (c, 0.13 in $CHCl_3$). λ_{max} 246 (ϵ 11830) ($CHCl_3$).

trans,trans-form

Amorph. solid. $[\alpha]_D -39.2$ (c, 0.52 in $CHCl_3$). λ_{max} 246 (ϵ 11270) ($CHCl_3$).

Tan, L.T. *et al.*, *J.O.C.*, 2000, **65**, 419-425 (isol, pmr, cmr)

Yokokawa, F. *et al.*, *Synlett*, 2001, 986-988 (synth)

Kutsumura, N. *et al.*, *Bull. Chem. Soc. Jpn.*, 2002, **75**, 847-850 (synth)


Yokokawa, F. *et al.*, *Tetrahedron*, 2002, **58**, 8127-8143 (synth, conformn)

Chen, Z. *et al.*, *ARKIVOC*, 2003, **vii**, 268-285 (synth)

Doi, M. *et al.*, *Acta Cryst. C*, 2003, **59**, 323-325 (cryst struct)

Ceridimine

[102719-93-7]



$C_{31}H_{36}N_4O_2$ 496.651
 Alkaloid from the bark of *Pagiantha cerifera* (preferred genus name *Tabernaemontana*) (Apocynaceae). Cryst. (CH_2Cl_2). Mp 173-175°. $[\alpha]_D^{20} -99$ (c, 0.5 in EtOH).

N-De-Me: Demethylceridimine

[167696-88-0]
 $C_{30}H_{34}N_4O_2$ 482.624
 Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae). $[\alpha]_D -167$ (c, 1.3 in EtOH).

16-Hydroxymethyl: Hunteriatryptamine.

16-Hydroxymethylceridimine

[169626-40-8]
 $C_{32}H_{38}N_4O_3$ 526.677
 Alkaloid from leaves of *Hunteria zeylanica* (Apocynaceae). Amorph. powder.


Baudouin, G. *et al.*, *Chem. Comm.*, 1986, 3 (uv, ir, pmr, cmr, ms, struct)

Subhadhirasakul, S. *et al.*, *Heterocycles*, 1995, **41**, 2049 (Hunteriatryptamine)

Azoug, M. *et al.*, *Phytochemistry*, 1995, **39**, 1223 (Demethylceridimine)

Cermizine A

[760947-73-7]



R = CH_2COOH

C-281

C-282

C₁₉H₃₄N₂O₂ 322.49

Closely related to Phlegmarine, P-361. Alkaloid from *Lycopodium cernuum*. Amorph. solid. $[\alpha]_D^{27} +3$ (c, 1 in MeOH). Morita, H. *et al.*, *Tetrahedron*, 2004, **60**, 7015-7023 (*isol, pmr, cmr*)

Cermizine B

C-283

[760947-74-8]

As Cermizine A, C-282 with

R = H

C₁₇H₃₂N₂ 264.453


Alkaloid from *Lycopodium cernuum*. Amorph. solid. $[\alpha]_D^{27} -2$ (c, 0.6 in MeOH).

Morita, H. *et al.*, *Tetrahedron*, 2004, **60**, 7015-7023 (*isol, pmr, cmr*)

Cermizine D

C-284

[760947-78-2]



Relative Configuration

C₁₆H₃₀N₂ 250.426

Alkaloid from *Lycopodium cernuum*. Amorph. solid. $[\alpha]_D^{25} -33$ (c, 0.6 in MeOH).


Morita, H. *et al.*, *Tetrahedron*, 2004, **60**, 7015-7023 (*isol, pmr, cmr*)

Nishikawa, Y. *et al.*, *Org. Lett.*, 2008, **10**, 1987-1990 (*synth*)

Cerpegin

C-285

1,1,5-Trimethylfuro[3,4-c]pyridine-3,4(1H,5H)-dione, 9CI [129748-28-3]

C₁₀H₁₁NO₃ 193.202

Alkaloid from *Ceropegia juncea*. Cryst. (MeOH/CHCl₃). Mp 268-270°. λ_{\max} 207 (ε); 210 (ε); 216 (ε); 234 (ε); 316 (ε) (90% EtOH) (Derep).

Sivakumar, K. *et al.*, *Acta Cryst. C*, 1990, **46**, 839 (*cryst struct*)

Adibatti, N.A. *et al.*, *Phytochemistry*, 1991, **30**, 2449-2450 (*isol, uv, ir, pmr, cmr, ms*)

Kelly, T.R. *et al.*, *J.O.C.*, 1992, **57**, 6657 (*synth*)

Guillier, F. *et al.*, *Tet. Lett.*, 1992, **33**, 7355 (*synth*)

Matsuo, K. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 2091 (*synth*)

Hong, H. *et al.*, *J.O.C.*, 1996, **61**, 391 (*synth*)

Villemin, D. *et al.*, *Tet. Lett.*, 1996, **37**, 8733 (*synth*)


Matsuo, K. *et al.*, *Heterocycles*, 1997, **45**, 1191 (*synth*)

Lazaar, J. *et al.*, *Tet. Lett.*, 2005, **46**, 3811-3813 (*synth*)

Cerulenin

C-286

3-(1-Oxo-4,7-nonadienyl)oxiranecarboxamide, 9CI. 2,3-Epoxy-4-oxo-7,10-dodecadienamamide. *Helicocerin*

C₁₂H₁₇NO₃ 223.271

Antifungal antibiotic. Shows anti-HIV and antineoplastic activity. Inhibitor of the biosynth. of fatty acids and steroids. Sol. MeOH, Et₂O, CS₂, CCl₄; fairly sol. H₂O; poorly sol. hexane. Log P 0.73 (calc). λ_{\max} 207 (MeOH) (Berdy).

► LD₅₀ (mus, ivn) 154 mg/kg; LD₅₀ (mus, scu) 245 mg/kg; LD₅₀ (mus, orl) 547 mg/kg.

(+)-form [17397-89-6]

Prod. by *Cephalosporium caeruleum* and *Helicoceras oryzae*. Needles (CCl₄ or C₆H₆). Mp 93°. Bp_{0.00001} 120°. $[\alpha]_D^{16} +63$ (c, 2 in MeOH). Pharmacol. active isomer.

► LD₅₀ (mus, orl) 547 mg/kg; LD₅₀ (mus, ipr) 211 mg/kg. JR1670000

Tetrahydro: [17397-90-9]

Mp 85.5-86.5°. $[\alpha]_D^{25} +43$ (c, 0.25 in MeOH).

(-)-form [68926-46-5]

Tetrahydro: [68926-45-4]

Mp 84.5°. $[\alpha]_D^{20} -52$ (c, 1.02 in MeOH).

(±)-form [62964-96-9]Cryst. (C₆H₆/hexane). Mp 40-43°.

Sano, Y. *et al.*, *J. Antibiot., Ser. A*, 1967, **20**, 344

Awaya, J. *et al.*, *J. Antibiot.*, 1975, **28**, 824 (*nmr, biosynth*)

Boeckman, R.K. *et al.*, *J.A.C.S.*, 1977, **99**, 2805 (*synth*)

Corey, E.J. *et al.*, *Tet. Lett.*, 1977, 3847 (*synth*)

Thomas, E.W. *et al.*, *C.A.*, 1978, **89**, 42956j (*synth*)

Pougny, J.R. *et al.*, *Tet. Lett.*, 1978, 3301 (*struct, abs config*)

Omura, S. *et al.*, *Methods Enzymol.*, 1981, **72**, 520 (*rev*)

Jakubowski, A.A. *et al.*, *J.O.C.*, 1982, **47**, 1221 (*synth*)

Furukawa, J. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 1229 (*synth*)

Pal, R. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 9283 (*anti-HIV activity*)

Yoda, H. *et al.*, *Tet. Lett.*, 1991, **32**, 6771 (*synth*)

Morisaki, N. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 2945; 2954 (*synth*)

Miller, M. *et al.*, *J.O.C.*, 1993, **58**, 6779 (*synth, bibl*)

Kuhajda, F.P. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1994, **91**, 6379 (*pharmacol*)

Kedar, T.E. *et al.*, *J.O.C.*, 1996, **61**, 6121 (*synth, pmr, cmr*)

Mani, N.S. *et al.*, *J.O.C.*, 1997, **62**, 636 (*synth*)


Renard, M. *et al.*, *Tetrahedron*, 2001, **57**, 2597-2608 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ECE500

Cervinomycin A₁

C-287

AM 5344A₁. Antibiotic AM 5344A₁ [82658-23-9]

C₂₉H₂₃NO₉ 529.502

Isol. from *Streptomyces cervinus* sp. nov. and from a *Micromonospora* sp. M39.

Inhibits anaerobic bacteria and mycoplasma. Yellow powder. Fairly sol.

MeOH, C₆H₆; poorly sol. Et₂O, H₂O, hexane. Mp 240° dec. $[\alpha]_D^{23} -92$ (c, 0.05 in CHCl₃). λ_{\max} 303 (ε 27300); 376 (ε 11600); 385 (ε 11300) (CHCl₃) (Derep).

► LD₅₀ (mus, ipr) 50 mg/kg. FK8980000

8,15-Quinone: **Cervinomycin A₂**, AM 5344A₂. Antibiotic AM 5344A₂. Antibiotic 167B [82658-22-8]

C₂₉H₂₁NO₉ 527.486

From *Streptomyces cervinus* and also from a strain of *Amycolata autotrophica*.

Inhibits anaerobic bacteria and mycoplasma. Reddish-orange powder.

Sol. CHCl₃; fairly sol. MeOH, EtOAc, DMSO, CHCl₃, C₆H₆; poorly sol. Et₂O, hexane, H₂O. Mp 290° dec. $[\alpha]_D^{20} -214$ (c, 0.25 in CHCl₃). Prob. identical with Antibiotic 4181A. λ_{\max} 260 (ε 37900); 329 (ε 28800); 375 (sh) (ε 8380); 420 (ε 6590) (CHCl₃) (Derep).

► LD₅₀ (mus, ipr) 50 mg/kg. FK8980100

8,15-Quinone, O¹²-de-Me: **De-O-methyl-cervinomycin A₂**. Antibiotic 167A [113537-08-9]

C₂₈H₁₉NO₉ 513.459

Prod. by a strain of *Amycolata autotrophica*. Orange powder. Sol. CHCl₃;

fairly sol. MeOH, Py, DMSO, EtOAc; poorly sol. EtOH, hexane, H₂O. Mp 250° dec. $[\alpha]_D^{23} -129$ (c, 0.02 in CHCl₃).

Prob. identical to Antibiotic 4181B, A-1121. λ_{\max} 268 (E1%/1cm 642); 325 (E1%/1cm 528); 420 (E1%/1cm 119) (CHCl₃) (Berdy).

[83381-75-3, 85897-53-6]

Eur. Pat., 1982, 54 801; *CA*, **97**, 180143 (*isol*)

Omura, S. *et al.*, *J. Antibiot.*, 1982, **35**, 645-652 (*isol, uv, ir*)

Japan. Pat., 1983, 83 09 690; *CA*, **98**, 214134 (*isol*)

Nakagawa, A. *et al.*, *J. Antibiot.*, 1987, **40**, 301-308 (*pmr, cmr, struct*)

Rao, A.V.R. *et al.*, *Tet. Lett.*, 1988, **29**, 3991-3992; 1991, **32**, 5199-5202 (*synth*)

Kelly, T.R. *et al.*, *J.A.C.S.*, 1989, **111**, 4522-4524 (*synth*)

Yadav, J.S. *et al.*, *Pure Appl. Chem.*, 1993, **65**, 1349 (*rev, synth*)


Malkina, N.D. *et al.*, *J. Antibiot.*, 1994, **47**, 342-348 (167A, 167B)

Mehta, G. *et al.*, *Tetrahedron*, 1994, **50**, 11729-11742 (*synth*)

Cespitulactam D

C-288

[1003840-01-4]



$C_{20}H_{29}NO_2$ 315.455
Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ -52 (c, 0.08 in CH_2Cl_2). λ_{max} 226 (log ϵ 4.52) (MeOH).

N-Me: Cespitulactam H

[1003840-05-8]

 $C_{21}H_{31}NO_2$ 329.481

Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ -112 (c, 0.7 in CH_2Cl_2). λ_{max} 228 (log ϵ 4.52) (MeOH).

N-(2-Phenylethyl): Cespitulactam A

[870537-81-8]

 $C_{28}H_{37}NO_2$ 419.606

Constit. of *Cespitulactaria taeniata*.
Amorph. powder. $[\alpha]_D^{25}$ -196 (c, 1 in CH_2Cl_2). λ_{max} 217; 230; 257 (MeOH).

N-[2-(4-Hydroxyphenyl)ethyl]: Cespitulactam C

[870537-83-0]

 $C_{28}H_{37}NO_3$ 435.605

Constit. of *Cespitulactaria taeniata*.
Amorph. powder. $[\alpha]_D^{25}$ -255 (c, 0.2 in CH_2Cl_2). λ_{max} 224; 279 (MeOH).

N-[2-(1H-Indol-3-yl)ethyl]: Cespitulactam J

[1003840-07-0]

 $C_{30}H_{38}N_2O_2$ 458.642

Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ -119 (c, 1.1 in CH_2Cl_2). λ_{max} 221 (log ϵ 4.88); 274 (log ϵ 2.84) (MeOH).

6-Ac: Cespitulactam E

[1003840-02-5]

 $C_{22}H_{31}NO_3$ 357.492

Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ -193 (c, 1.37 in CH_2Cl_2). λ_{max} 228 (log ϵ 5.11) (MeOH).

6-Ketone, N-(2-phenylethyl): Cespitulactam B

[870537-82-9]

 $C_{28}H_{35}NO_2$ 417.59

Constit. of *Cespitulactaria taeniata*.
Amorph. powder. $[\alpha]_D^{25}$ -110 (c, 0.05 in CH_2Cl_2).

10 β -Hydroxy: Cespitulactam F

[1003840-03-6]

 $C_{20}H_{29}NO_3$ 331.454

Constit. of *Cespitulactaria taeniata*. Powder. $[\alpha]_D^{25}$ -156 (c, 1.27 in MeOH). λ_{max} 228 (log ϵ 4.48) (MeOH).

10 β -Hydroxy, N-Et: Cespitulactam G

[1003840-04-7]

 $C_{22}H_{33}NO_3$ 359.508

Constit. of *Cespitulactaria taeniata*.

Amorph. solid. $[\alpha]_D^{25}$ -204 (c, 1.5 in CH_2Cl_2). λ_{max} 226 (log ϵ 4.61) (MeOH).

11-Hydroxy, 7 ξ , 8, 11, 12 ξ -tetrahydro, 9, 10-didehydro, 6-ketone: Cespitulactam I

[1003840-06-9]

 $C_{26}H_{29}NO_3$ 331.454

Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ -3 (c, 0.2 in CH_2Cl_2).

 Δ^{10} -Isomer, N-[2-(4-hydroxyphenyl)ethyl]: Cespitulactam K

[1003840-08-1]

 $C_{28}H_{37}NO_3$ 435.605

Constit. of *Cespitulactaria taeniata*.
Amorph. solid. $[\alpha]_D^{25}$ +53 (c, 0.75 in CH_2Cl_2). λ_{max} 229 (log ϵ 4.89); 274 (log ϵ 3.3) (MeOH).


Shen, Y.-C. *et al.*, *Tet. Lett.*, 2005, **46**, 7893-7897 (*Cespitulactams A-C*)

Shen, Y.-C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1961-1965 (*Cespitulactams D-K*)

Cetoniacytone B

C-289

4-Amino-5-hydroxy-1-hydroxymethyl-7-oxabicyclo[4.1.0]hept-3-en-2-one, 9CI. 3-Amino-5,6-epoxy-4-hydroxy-6-hydroxymethyl-2-cyclohexen-1-one

 $C_7H_9NO_4$ 171.152

The epoxyquinomicins are structurally unrelated to the quinomicins. Minor metab. of *Actinomyces* sp. Lu 9419 isol. from *Cetonia aureata*. Solid. λ_{max} 296 (ϵ 7510) (MeOH).

N-Ac: Cetoniacytone A $C_9H_{11}NO_5$ 213.19

Minor metab. of *Actinomyces* sp. Lu 9419 from *Cetonia aureata*. Cytotoxic agent. Solid. Mp 163°. λ_{max} 284 (ϵ 10500) (MeOH).

N-(2-Hydroxybenzoyl): Epoxyquinomicin C

[200496-85-1]

 $C_{14}H_{13}NO_6$ 291.26

Prod. by *Amycolatopsis* sp. MK299-95F4. Inhibitor of histidine decarboxylase. Powder. Sol. MeOH, Me₂CO, EtOAc; fairly sol. CHCl₃; poorly sol. hexane. Mp 168-172° dec. $[\alpha]_D^{25}$ +128 (c, 1 in MeOH). λ_{max} 297 (ϵ 17430) (MeOH). λ_{max} 236; 255; 325 (ϵ 8000); 370 (MeOH) (Berdy). λ_{max} 253 (ϵ 6700); 322 (ϵ 8500) (MeOH-HCl) (Berdy). λ_{max} 234; 257; 327 (ϵ 8300); 371 (MeOH-NaOH) (Berdy).

► LD₅₀ (mus, ipr) 50 mg/kg.

N-(3-Chloro-2-hydroxybenzoyl): Epoxyquinomicin D

[200496-86-2]

 $C_{14}H_{12}ClNO_6$ 325.705

Prod. by *Amycolatopsis* sp. MK299-95F4. Powder. Sol. MeOH, Me₂CO, EtOAc; fairly sol. CHCl₃; poorly sol.

hexane. Mp 163-168° dec. $[\alpha]_D^{25}$ +142 (c, 1 in MeOH). λ_{max} 299 (ϵ 17590) (MeOH). λ_{max} 253 (ϵ 6700); 326 (ϵ 6300) (MeOH) (Berdy). λ_{max} 252 (ϵ 5700); 327 (ϵ 6500) (MeOH-HCl) (Berdy). λ_{max} 235 (ϵ 9100); 259; 324 (ϵ 5800); 376 (MeOH-NaOH) (Berdy).

1-Ketone, N-(2-hydroxybenzoyl): Epoxyquinomicin B

[175448-32-5]

 $C_{14}H_{11}NO_6$ 289.244

Prod. by *Amycolatopsis* sp. MK299-95F4. Mod. active against gram-positive bacteria and *Pasteurella* sp. Pale yellow powder. Mp 178-184° dec. $[\alpha]_D^{25}$ +32.2 (c, 0.2 in MeOH). λ_{max} 253 (sh) (ϵ 5400); 326 (ϵ 6300) (MeOH).

1-Ketone, N-(3-chloro-2-hydroxybenzoyl): Epoxyquinomicin A

[175448-31-4]

 $C_{14}H_{10}ClNO_6$ 323.689

Prod. by *Amycolatopsis* sp. MK299-95F4. Mod. active against gram-positive bacteria and *Pasteurella* sp. Pale yellow powder. Mp 168-173° dec. $[\alpha]_D^{25}$ +44.6 (c, 0.5 in MeOH). λ_{max} 255 (sh) (ϵ 5900); 325 (ϵ 8000); 370 (sh) (ϵ 2700) (MeOH).

Matsumoto, T. *et al.*, *J. Antibiot.*, 1997, **50**, 900-905; 906-911; 912-915 (*Epoxyquinomicins, isol, uv, ir, pmr, cmr, ms, activity*)

Matsumoto, T. *et al.*, *J. Antibiot.*, 2000, **53**, 637-639 (*Epoxyquinomicins, activity*)


Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 207-210 (*Epoxyquinomicin B, synth*)

Schlörke, O. *et al.*, *J. Antibiot.*, 2002, **55**, 635-642 (*Cetoniacytones*)

Mahmud, T. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1384-1391 (*rev, biosynth*)

3,6-Cevanediol

C-290

3,6-Dihydroxycevanine $C_{27}H_{45}NO_2$ 415.658

The stereochem. of the various natural stereoisomers requires clarification. In CAS, the 20 α (20 β -H)- 25 β - (25 α -H)- stereochemistries are assumed as part of the Cevane stereoparent.

(3 α , 5 α , 6 α , 25 α)-form**Petilinine**

[21851-16-1]

Alkaloid from *Petilium raddeana* (Liliaceae). Cryst. (MeOH). Mp 277-278°. $[\alpha]_D^{25}$ -9.6 (c, 1.25 in MeOH/CHCl₃ 1:1).

Hydrochloride: Mp 296-297°.

(3 β , 5 α , 6 α , 25 α)-form**Petilidine**


[22169-21-7]

Alkaloid from *Petilium raddeana* (Liliaceae). Mp 265-266°. $[\alpha]_D^{25}$ -15.5 (c, 1.2 in MeOH). There is also an alkaloid

- Stenanzidine, Mp 215-217°, $[\alpha]_D^{+5}$, from *Rhinopetalum stenatherum* (Liliaceae) which appears to have been assigned the same struct. and stereochem. as Petilidine, but the paper is not clear and the Engl. translation appears to contain errors confusing it with Stenanzine and Stenanzidine. Hydrochloride: [22169-22-8] Mp 273-274°.
- 6-Ketone: 3-Hydroxy-6-cevanone. Puqiedinone.** Zhebeirine. 25-Epiebeidinone [143120-47-2] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from bulbs of *Fritillaria puqiensis* and *Fritillaria thunbergii*. Exhibits antitussive and antitumour activity. Needles ($C_6H_{12}/EtOH/Et_2NH$). Mp 205-208°. $[\alpha]_D^{20}$ -62.3 (c, 0.2 in $CHCl_3$).
- (3 β ,5 α ,6 α ,25 β)-form**
Edwardinine. Eduardine. Isoebeidine [58116-31-7] [128351-45-1] Alkaloid from *Petilium eduardii* leaves. Cryst. (Me_2CO). Mp 255-257°. $[\alpha]_D$ +7.4 (c, 0.875 in $CHCl_3$).
- 6-Ketone: Edwardine.** Eduardine [25650-68-4] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from *Petilium eduardii* and *Fritillaria ebeiensis* var. *purpurea* (Liliaceae). Needles (hexane/ $Et_2NH/EtOH$). Mp 102-105° (Ebeidinone) Mp 247-251° (Edwardine). $[\alpha]_D$ -53.02 (MeOH). $[\alpha]_D$ -62.2 (c, 0.89 in $CHCl_3$). Same struct assigned to Edwardine and Ebeidinone. Reported Mps are widely different.
- (3 β ,5 α ,6 β ,25 α)-form**
Puqiedine [924889-55-4] Alkaloid from *Fritillaria puqiensis* bulbs. Cubes (Me_2CO). Mp 139-141°. $[\alpha]_D$ -38.4 (c, 0.1 in MeOH).
- (3 β ,5 α ,6 β ,25 β)-form**
Ebeidine. Edpetilidine [25650-70-8] Alkaloid from the bulbs of *Fritillaria ebeiensis* var. *purpurea*, *Fritillaria imperialis*, *Petilium eduardii* and *Rhinopetalum bucharicum* (Liliaceae). Needles (hexane/ $Et_2NH/EtOH$). Mp 227-228°. $[\alpha]_D$ -48.2 (c, 2.23 in Py).
- 3-O- β -D-Glucopyranoside: Edpetinosine** [207129-43-9] $C_{33}H_{55}NO_7$ 577.8 Alkaloid from *Petilium eduardii*. Cryst. ($Me_2CO/MeOH$). Mp 174-176°.
- Di-Ac:** Mp 143.5-146.5°.
- 3-Ketone: Persinine** [205586-14-7] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from *Fritillaria persica*.
- (3 β ,5 α ,20 β ,25 α)-form**
6-Ketone: Dongbeirine [151003-95-1] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from bulbs of *Fritillaria thunbergii* (Liliaceae). Mp 151-153°. $[\alpha]_D^{15}$ -51.5 (c, 0.4 in MeOH).
- (3 β ,5 α ,20 β ,25 β)-form**
6-Ketone: Dongbeirine [151003-96-2] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from bulbs of *Fritillaria thunbergii* (Liliaceae). Cryst. Mp 132-134°. $[\alpha]_D^{15}$ -50.5 (c, 0.5 in $CHCl_3$).
- (3 β ,5 α ,6 β ,17 β ,25 α)-form**
Harepermine. Persicanidine B [145307-25-1] [105814-57-1] Alkaloid from *Fritillaria persica*, *Fritillaria harelinii* and *Fritillaria imperialis*. Plates + $1H_2O$. Mp 167-169°. $[\alpha]_D^{25}$ -18 (c, 0.5 in $CHCl_3$). Stereochem. of Harepermine incorrectly shown in original ref. (1986).
- 3-O- β -D-Glucopyranoside: Hareperminoside** [144940-48-7] [105814-58-2] $C_{33}H_{55}NO_7$ 577.8 Alkaloid from *Fritillaria harelinii* and *Fritillaria persica*. Powder. Mp 168-170°. $[\alpha]_D$ -27 (c, 0.5 in MeOH).
- (3 β ,5 α ,6 β ,17 β ,25 β)-form**
Hupehenine. Delavine [98243-57-3] [84048-44-2] Alkaloid from *Fritillaria delavayi* and *Fritillaria hupehensis* (Liliaceae). Needles (MeOH). Mp 181-183°. $[\alpha]_D$ -20 (c, 0.5 in $CHCl_3$). Revision of originally assigned config.
- 3-O- β -D-Glucopyranoside: Hupeheninoside** [98985-22-9] $C_{33}H_{55}NO_7$ 577.8 Alkaloid from bulbs of *Fritillaria hupehensis* (Liliaceae). Cryst. Mp 241-244°. $[\alpha]_D^{20}$ -41 (c, 0.16 in MeOH).
- 3-Ketone: 6-Hydroxy-3-cevanone. Hupehenizine** [103805-68-1] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from *Fritillaria hupehensis* (Liliaceae).
- 3-Ketone, 6-O- β -D-glucopyranoside: Yubeiside** [157382-28-0] Alkaloid from *Fritillaria yuminensis*.
- 6-Ketone: Hupehenirine.** Delavinone. Sinpeimine A. Yubeinine [96997-98-7] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from *Fritillaria delavayi*, *Fritillaria hupehensis* and *Fritillaria valujevii* (Liliaceae). Needles (MeOH). Mp 182-184°. $[\alpha]_D$ -54 (c, 0.5 in $CHCl_3$).
- 6-Ketone: hydrochloride:** Mp 217-219°.
- 6-Ketone, 3-O- β -D-glucopyranoside: Yi-beinoside A.** Hupehenizoside [98985-24-1] $C_{33}H_{53}NO_7$ 575.784 Alkaloid from *Fritillaria pallidiflora* and from the bulbs of *Fritillaria lichuanensis*. Cryst. or needles ($CHCl_3/MeOH$). Mp 248-250° Mp 258-260°. $[\alpha]_D^{15}$ -59 (c, 0.1 in MeOH).
- (3 β ,5 α ,17 β ,22 β ,25 β)-form**
6-Ketone: Chuanbeinone [103530-47-8] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from *Fritillaria delavayi* and *Fritillaria taipaiensis*. Needles (MeOH). Mp 149-152°. $[\alpha]_D$ -62.4 (c, 0.5 in $CHCl_3$).
- 6-Ketone, 3-O- β -D-glucopyranoside: Yi-beinoside B** [150134-44-4] $C_{33}H_{53}NO_7$ 575.784 Alkaloid from *Fritillaria pallidiflora*. Cryst. (MeOH). Mp 202-204°.
- (3 β ,5 α ,6 β ,20 β ,25 α)-form**
Lichuanine [934362-24-0] Alkaloid from the bulbs of *Fritillaria lichuanensis*. Needles ($EtOAc/Me_2CO$). Mp 159-160°.
- (3 β ,5 α ,6 β ,20 β ,25 β)-form**
N-Oxide: Lichuanisinine [934362-25-1] $C_{27}H_{45}NO_3$ 431.657 Alkaloid from the bulbs of *Fritillaria lichuanensis*. Powder. Mp 148-151°.
- (3 β ,5 α ,6 β ,20 ξ ,25 ξ)-form**
Stenanzidine [94482-44-7] Alkaloid from *Rhinopetalum stenatherum* (Liliaceae). Cryst. ($MeOH/Me_2CO$). Mp 277°. $[\alpha]_D$ -17 (c, 0.7 in EtOH). Stereochem. not clear, confusion in paper. See note under Petilidine above.
- (3 β ,5 α ,6 β ,13 α ,17 β ,25 α)-form**
Persicanidine A [139757-61-2] Alkaloid from bulbs of *Fritillaria persica* (Liliaceae). Cyclic AMP phosphodiesterase inhibitor. Prisms (MeOH). Mp 208° dec. $[\alpha]_D$ -7.8 ($CHCl_3$).
- (3 β ,5 α ,6 β ,17 α ,20 β ,25 β)-form**
Forticine [471843-48-8] Alkaloid from the bulbs of *Fritillaria imperialis*. Needles. Mp 221-223°. $[\alpha]_D^{23}$ -52 (c, 0.5 in $CHCl_3$).
- (3 β ,5 α ,6 β ,17 β ,20 β ,22 β ,25 β)-form**
Songbeinine. Tortifoline [125354-97-4] Alkaloid from bulbs of *Fritillaria unibracteata* and *Fritillaria tortifolia* (Liliaceae). Cubes (MeOH). Mp 206-208° (195-197°). $[\alpha]_D^{29}$ -240 (c, 0.01 in $CHCl_3$). $[\alpha]_D$ -41 (c, 1.07 in $CHCl_3$).
- 6-Ketone: Songbeinone** [150133-32-7] $C_{27}H_{43}NO_2$ 413.642 Alkaloid from bulbs of *Fritillaria unibracteata* (Liliaceae).
- Nuriddinov, R.N. et al., *Khim. Prir. Soedin.*, 1967, **3**, 316-324; 1968, **4**, 261-262; 332-333; 1969, **5**, 333-334; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 267-273; 1968, **4**, 225-226; 282-283; 1969, **5**, 284-285 (*Petiline, Petilidine, Edpetilidine, Eduardine*) Nabiev, A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 535-536; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 570-571 (*Eduardine*) Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1976, **12**, 630-632; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 563-565 (*Eduardine*)

- Wu, J. et al., *Zhongcaoyao*, 1982, **13**, 339-342; 351; *CA*, **98**, 31403m (*Hupehenine*)
- Samikov, K. et al., *Khim. Prir. Soedin.*, 1984, **20**, 341-344; 498-500; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 318-320; 470-471 (*Stenanizidine*, *Stenanizidine*)
- Liu, Q. et al., *Yaouxue Xuebao*, 1984, **19**, 894-898; *CA*, **103**, 19851v (*Sinpeimine A*)
- Kaneko, K. et al., *Chem. Pharm. Bull.*, 1985, **33**, 2614-2617 (*Delavine*, *Delavinone*)
- Wu, J. et al., *Yaouxue Xuebao*, 1985, **20**, 372-376; *CA*, **103**, 193124v (*Hupeheninoside*)
- Min, Z. et al., *Phytochemistry*, 1986, **25**, 2008-2009 (*Harepermine*, *Hareperminoside*)
- Kaneko, K. et al., *Tet. Lett.*, 1986, **27**, 2387-2390 (*Chuanbeinone*)
- Wu, J. et al., *Zhongcaoyao*, 1986, **17**, 101-104; 1989, **20**, 530-532; *CA*, **105**, 111971; **113**, 74735 (*Hupehenirine*, *Hupehenizine*, *Hupehenine*, *Hupeheninoside*)
- Lee, P. et al., *Chem. Pharm. Bull.*, 1988, **36**, 4316-4329 (*Ebeidine*, *Ebeidinone*)
- Kitamura, Y. et al., *Chem. Pharm. Bull.*, 1989, **37**, 1514-1516 (*Tortifoline*)
- Wu, J.Z. et al., *Yaouxue Xuebao*, 1989, **24**, 600-605; *CA*, **112**, 95497
- Xu, D.M. et al., *Yaouxue Xuebao*, 1990, **25**, 795-797; 1993, **28**, 192-196 (*Yibeinosides*)
- Yu, S. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1990, **32**, 929-935; 1992, **34**, 945-949; *CA*, **116**, 55508r; **119**, 156233q (*Songbeinone*, *Songbeinone*)
- Zhang, J. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1991, **33**, 923-926; *CA*, **117**, 128151b (*Zhebeirine*)
- Ori, K. et al., *Chem. Lett.*, 1992, 163-166 (*Persicanidine A*)
- Ori, K. et al., *Phytochemistry*, 1992, **31**, 3605-3607 (*Persicanidine B*)
- Jianxing, Z. et al., *Phytochemistry*, 1993, **33**, 946-947 (*Dongbeinone*, *Dongbeirine*)
- Zhang, J.-X. et al., *Yaouxue Xuebao*, 1993, **35**, 963-967 (*Yubeimine*, *Yubeiside*)
- Lin, G. et al., *J. Nat. Prod.*, 1995, **58**, 1662-1667 (*Pugiedinone*)
- Shakirova, U.T. et al., *Khim. Prir. Soedin.*, 1997, **33**, 617-619; *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 476-478 (*Edpetinosine*)
- Bingol, F. et al., *CA*, 1998, **128**, 268255r (*Persinine*)
- Atta-ur-Rahman, et al., *Chem. Pharm. Bull.*, 2002, **50**, 1013-1016 (*Forticine*)
- Tsuda, Y. et al., *Nat. Prod. Res.*, 2004, **18**, 205-209 (*Harepermine*)
- Pi, H.-F. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 133-136; 253-257 (*Huopenhizoiside*, *Lichuanine*, *Lichuanisidine*)
- Jiang, Y. et al., *Steroids*, 2006, **71**, 843-848 (*Pugiedine*)
- Zhang, Y.-H. et al., *Chin. J. Chem.*, 2007, **25**, 1728-1731 (*Zhebeirine*)

3,4,12,14,16,17,20-Cevaneheptol C-291



$C_{27}H_{45}NO_7$ 495.655

(3 β ,4 α ,5 α ,16 β)-form

Sabine. *Neosabidine*

[6883-32-5]

Alkamine from seeds of *Schoenocaulon officinale* (Liliaceae). Shows hypotensive and brachycardic props. Mp 256-258° dec. $[\alpha]_D^{25}$ -33 (EtOH).

3-Ac: **Sabadine**. *Sabatine*

[124-80-1]

$C_{29}H_{47}NO_8$ 537.692


Minor alkaloid from seeds of *Schoenocaulon officinale* (Liliaceae). Prisms (Et₂O). $[\alpha]_D^{25}$ -11 (c, 1.93 in EtOH). Softens at 170-190°, dec. above 200°.

3,4,16-Tri-Ac: Mp 221-222° dec. (sinters from 165°). $[\alpha]_D^{25}$ +9 (c, 1.14 in CHCl₃).

Kupchan, S.M. et al., *J. Med. Chem.*, 1962, **5**, 690-714 (*Sabine*, *Sabadine*, *isol*, *ir*, *ord*, *struct*, *pharmacol*, *bibl*)

Narayanan, C.R. et al., *Prog. Chem. Org. Nat. Prod.*, 1962, **20**, 354 (*Neosabidine*)

3,6,12,14,16,20-Cevanehexol C-292



$C_{27}H_{45}NO_6$ 479.656

(3 β ,5 α ,6 α ,16 β)-form


Pingpeimine B

[82851-52-3]

Alkaloid from *Fritillaria ussuriensis* (Liliaceae). Mp 255-257°. $[\alpha]_D^{25}$ +24.9 (c, 0.08 in MeOH).

Xu, D.M. et al., *Yaouxue Xuebao*, 1988, **23**, 902-905; *CA*, **111**, 74773 (*Pingpeimine B*)

3,6,14,16,20-Cevanepentol C-293



$C_{27}H_{45}NO_5$ 463.656

(3 β ,5 α ,6 α ,16 ξ ,20 ξ)-form


Pingpeimine A

[82841-67-6]

Alkaloid from *Fritillaria ussuriensis* (Liliaceae).

Xu, D. et al., *Yaouxue Xuebao*, 1982, **17**, 355-359; *CA*, **97**, 107068h (*Pingpeimine A*)

3,6,7-Cevanetriol C-294



$C_{27}H_{45}NO_3$ 431.657

(3 α ,5 α ,6 β ,7 α ,25 α)-form


3 α -Pugiedin-7-ol

[923925-04-6]

Alkaloid from bulbs of *Fritillaria pugiensis*. Cubes (Me₂CO). Mp 267-269°. $[\alpha]_D^{20}$ -47.7 (c, 0.1 in MeOH).

Jiang, Y. et al., *Steroids*, 2006, **71**, 843-848 (*isol*, *pnr*, *cmr*)

3,6,14-Cevanetriol C-295



$C_{27}H_{45}NO_3$ 431.657

In CAS the (20 α ,25 α)-configs. are assumed as part of the cevane stereoparent.

(3 α ,5 α ,6 α ,25 α)-form

Korseveramine

[36506-67-9]

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Cryst. (MeOH). Mp 304-305°. $[\alpha]_D^{25}$ -14.63 (c, 0.823 in 10% AcOH aq.).

Hydrochloride: Mp 320-322°.

(3 α ,5 α ,6 β ,25 α)-form

Korseveriline

[21851-05-8]

Alkaloid from bulbs of *Korolkowia sewerzowii* (Liliaceae). Mp 240-243°. $[\alpha]_D^{25}$ -15 (c, 0.8 in petrol).

3-Ac: **Ceverine**. *Severine*†

[66408-50-2]

$C_{29}H_{47}NO_4$ 473.695

Alkaloid from above-ground parts of *Korolkowia sewerzowii* (Liliaceae). Mp 144-146°. $[\alpha]_D^{25}$ -20.9 (c, 0.43 in CHCl₃).

3-Ac, N-oxide: **Ceverine N-oxide**. *Severine N-oxide*

[74119-88-3]

$C_{29}H_{47}NO_5$ 489.694

Alkaloid from *Korolkowia sewerzowii*

(Liliaceae). Cryst. (MeOH). Mp 255-257°. $[\alpha]_D$ 0.

6-Ketone: 3,14-Dihydroxy-6-cevanone.

Korseverilinone

[86630-13-9]

$C_{27}H_{43}NO_3$ 429.642

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 222-223°. $[\alpha]_D$ -18.8 (c, 0.478 in $CHCl_3$).

(3 β ,5 α ,6 β ,25 α)-form

Cevedine. Sevedine

[66512-88-7]

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 212-214°. $[\alpha]_D$ -17.2 (c, 1.22 in $CHCl_3$).

N-Oxide: **Cevedine N-oxide. Sevedine N-oxide**

[74119-89-4]

$C_{27}H_{45}NO_4$ 447.657

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 241-243°. $[\alpha]_D$ -14 (c, 0.5 in EtOH).

6-Ac: **Acetylsevedine. Acetylcevedine**

[104513-83-9]

$C_{29}H_{47}NO_4$ 473.695

Alkaloid from aerial parts of *Korolkowia sewerzowii* (Liliaceae). Mp 189° (foams). $[\alpha]_D$ -36.6 (c, 1.202 in EtOH).

3,6-Di-Ac: **Diacetylcevedine**

[66512-89-8]

$C_{31}H_{49}NO_5$ 515.732

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Cryst. (Me_2CO /petrol). Mp 202-204°.

(3 β ,5 α ,6 β ,25 β)-form

Sewertzidine. Severtzidine

[61950-74-1]

Alkaloid from above-ground parts of *Korolkowia sewerzowii* (Liliaceae). Cryst. (Me_2CO). Mp 244-245°. $[\alpha]_D$ -46.4 (c, 0.69 in $CHCl_3$).

(3 β ,5 α ,6 α ,25 α)-form

Sevedamine. 6-Episevedine. Sewedamine

[107657-48-7]

Alkaloid from aerial parts of *Korolkowia sewerzowii*. Cryst. (Me_2CO). Mp 256-258°. $[\alpha]_D$ +3.6 (c, 1.25 in EtOH).

Nuriddinov, R.N. et al., *Khim. Prir. Soedin.*, 1968, 4, 258-259; 1971, 7, 773-778; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, 4, 222-223; 1971, 7, 746-750 (Korseveriline, Korseveramine)

Samikov, K. et al., *Khim. Prir. Soedin.*, 1976, 12, 367-370; 1977, 13, 673-675; 1979, 15, 823-826; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, 12, 320-322; 1977, 13, 559-560; 1979, 15, 728-730 (Sewertzidine, Cevedine, Ceverine oxide)

Abdullaeva, D.U. et al., *Khim. Prir. Soedin.*, 1977, 13, 671-673; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, 13, 557-558 (Ceverine)


Samikov, K. et al., *Khim. Prir. Soedin.*, 1981, 17, 529 (Cevedine oxide)

Kul'kova, V.V. et al., *Khim. Prir. Soedin.*, 1983, 19, 72-74; 1986, 22, 352-353; 1991, 27, 440-441; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, 19, 68-69; 1986, 22, 325-326; 1991, 27, 384-385 (Korseverilinone, Acetylcevedine, Diacetylcevedine)

Samikov, K. et al., *Khim. Prir. Soedin.*, 1986, 22, 622-624; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, 22, 585-586 (Sevedamine)

3,6,15-Cevanetriol

C-296



$C_{27}H_{45}NO_3$ 431.657

(3 α ,5 α ,6 β ,15 α)-form

Edpetsinine


[62908-12-7]

Alkaloid from *Petilium eduardii* (Liliaceae). Mp 247-248°. $[\alpha]_D$ -45.6 (c, 0.71 in MeOH).

Nabiev, A. et al., *Khim. Prir. Soedin.*, 1976, 12, 679; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, 12, 615

3,6,16-Cevanetriol

C-297



$C_{27}H_{45}NO_3$ 431.657

(3 β ,5 α ,6 β ,25 β)-form

Delafirine

[119766-89-1]

Alkaloid from bulbs of *Fritillaria delavayi* (Liliaceae). Powder. Mp 269-272°. $[\alpha]_D$ -39.9 (c, 0.50 in $CHCl_3$).

Tri-Ac: Mp 108-113°.

6-Ketone: 3,16-Dihydroxy-6-cevanone.

Delafirinone

[119766-92-6]

$C_{27}H_{43}NO_3$ 429.642

Alkaloid from bulbs of *Fritillaria delavayi* (Liliaceae). Needles (MeOH). Mp 134-136°. $[\alpha]_D$ -58.9 (c, 0.58 in $CHCl_3$).

6-Ketone, di-Ac: Mp 138-140°.

23,24-Didehydro: Cev-23-ene-3,6,16-triol.

23,24-Didehydro-3,6,16-cevanetriol.

Impericine

[471843-45-5]

$C_{27}H_{43}NO_3$ 429.642


Alkaloid from the bulbs of *Fritillaria imperialis*. Needles. Mp 195-197° dec. $[\alpha]_D^{25}$ -28 (c, 0.5 in $CHCl_3$).

Kaneko, K. et al., *Chem. Pharm. Bull.*, 1988, 36, 4700-4705 (isol, ir, pmr, cmr, ms, struct)

Atta-ur-Rahman, et al., *Chem. Pharm. Bull.*, 2002, 50, 1013-1016 (Impericine)

3,6,20-Cevanetriol

C-298



$C_{27}H_{45}NO_3$ 431.657

Log P 3.66 (uncertain value) (calc).

(3 α ,5 α ,6 α)-form

Isobaimonidine

[74184-79-5]

Minor alkaloid from *Fritillaria imperialis* (Liliaceae). Cryst. ($Me_2CO/CHCl_3$). Mp 238-239°. $[\alpha]_D^{22}$ -12 (c, 0.5 in $CHCl_3$). $[\alpha]_D$ -59.2 (c, 0.25 in $CHCl_3$).

(3 α ,5 α ,6 β)-form

Baimonidine

[73650-52-9]

Alkaloid from *Fritillaria verticillata* (Liliaceae). Mp 179-181.5°. $[\alpha]_D$ -36.4 (c, 1 in $CHCl_3$).

(3 β ,5 α ,6 α)-form

Verticine. Peimine

[23496-41-5]

Alkaloid from *Fritillaria roylei*; main alkaloid of *Fritillaria verticillata* var. *thunbergii* (Liliaceae). Antihypotensive agent. $[\alpha]_D^{16}$ -19.4 (EtOH). May be identical with Apovericine. λ_{max} 215 (EtOH) (Berdy).

► Causes paralysis to respiratory system and sensory nerves.

Hydrochloride: Mp 310-312° dec. (291-294° dec.). $[\alpha]_D^{14}$ -18.5 (H₂O).

β -N-Oxide: **Verticine N-oxide. Alkaloid F3**

[77410-34-5]

$C_{27}H_{45}NO_4$ 447.657

Isol. from the crude drug "Bai-mo" (sun-bleached lime-treated bulbs of *Fritillaria thunbergii*) (Liliaceae). Needles (EtOAc/hexane). Mp 283-288° dec. $[\alpha]_D$ +8.4 (c, 1.9 in MeOH).

3-O- β -D-Glucopyranoside: **Peiminoside**

[123937-51-9]

$C_{33}H_{55}NO_8$ 593.799

Alkaloid from *Fritillaria verticillata* var. *thunbergii* (Liliaceae). Noncryst. Mp 195° (as penta-Ac). $[\alpha]_D^{20}$ -36 (c, 0.5 in EtOH) (penta-Ac).

3-Ac: **3-O-Acetylverticine**

$C_{29}H_{47}NO_4$ 473.695

Alkaloid from the bulbs of *Fritillaria hupehensis*. Cryst. (EtOAc). Mp 193-194°. $[\alpha]_D^{20}$ -40 (c, 0.1 in MeOH).

6-Ketone: 3,20-Dihydroxy-6-cevanone.

Verticinone. Peiminine. Raddeanine.

Fritillarine

[18059-10-4]

$C_{27}H_{43}NO_3$ 429.642

Alkaloid from bulbs of *Fritillaria thunbergii* and from *Fritillaria verticillata*, *Fritillaria ebeiensis* and *Fritillaria roylei* (Liliaceae). Local anaesthetic,

antihypertensive agent. Mp 212-213°. $[\alpha]_D^{25}$ -58.3 (EtOH). $[\alpha]_D^{13}$ -54.5 (CHCl₃). Log P 3.26 (uncertain value) (calc). λ_{max} 288 (EtOH) (Berdy).

6-Ketone, N-oxide (β -): Verticinone N-oxide. Alkaloid F4
[77410-35-6]
C₂₇H₄₃NO₄ 445.641
Isol. from "Bai-mo" *Fritillaria thunbergii* (Liliaceae). Prisms (EtOAc/hexane). Mp 283-285° dec. $[\alpha]_D$ -54.5 (c, 1.8 in MeOH).

6-Ketone, 3-Ac: 3-O-Acetylverticinone
C₂₉H₄₅NO₄ 471.679
Alkaloid from the bulbs of *Fritillaria hupehensis*. Cryst. (EtOAc). Mp 113-114°. $[\alpha]_D^{20}$ -50.3 (c, 0.11 in MeOH).

(3 β ,5 α ,6 α ,22 β)-form

Wanpeinine A
[107299-20-7]
Alkaloid from the Chinese drug Wan Bei Mu (bulbs of *Fritillaria anhuiensis*) (Liliaceae). Cryst. Mp 281-283° dec. $[\alpha]_D$ -8.7.

(3 β ,5 α ,6 α ,25 α)-form

Zhebeinine
[135636-54-3]
Alkaloid from the bulbs of *Fritillaria thunbergii* (Liliaceae). Cryst. Mp 222-224°. $[\alpha]_D^{25}$ -21.3 (c, 0.5 in MeOH).
6-Ketone: Zhebeinine
[144606-87-1]
C₂₇H₄₃NO₃ 429.642
Alkaloid from *Fritillaria thunbergii* (Liliaceae). Mp 181-183°. $[\alpha]_D$ -48.8 (c, 0.03 in CHCl₃).

(3 β ,5 α ,6 α ,13 α ,22 β)-form

Siechuansine. 13-Epiwanpeinine A
[147511-59-9]
Alkaloid from bulbs of *Fritillaria siechuanica*.

(3 β ,5 α ,6 α ,22 β ,25 α)-form

3-O- β -D-Glucopyranoside: Zhebeininoside
[152695-94-8]
C₃₃H₅₃NO₈ 593.799
Alkaloid from *Fritillaria thunbergii*. Mp 226-228°. $[\alpha]_D^{15}$ -48.5 (c, 0.3 in MeOH).

(3 β ,5 α ,6 β)-form

Isoverticine
[23496-43-7]
Alkaloids from *Fritillaria verticillata* (Liliaceae). Mp 135-137°. $[\alpha]_D$ -45 (c, 1 in CHCl₃).
N-Oxide (β -): Isoverticine β -N-oxide
[866572-25-0]
C₂₇H₄₅NO₄ 447.657
Alkaloid from the bulbs of *Fritillaria wabuensis*. Mp 207-210°.

(3 β ,5 α ,6 β ,25 α)-form

Korselidine
[122405-29-2]
Alkaloid from aerial parts of *Korolkowia sewerzowii* (Liliaceae). Cryst. (Me₂CO). Mp 276-278°. $[\alpha]_D$ -53 (c, 0.3 in EtOH).

(3 β ,5 α ,6 β ,17 β ,25 α)-form

Petine
[171236-02-5]

Alkaloid from bulbs of *Petilium eduardii*. Cryst. (hexane/Me₂CO). Mp 145-147°.

N-Oxide: Petine N-oxide

[170557-06-9]
C₂₇H₄₃NO₄ 447.657
From bulbs of *Petilium eduardii*. Cryst. (Me₂CO). Mp 241-243°.

(3 β ,5 α ,17 β)-form

6-Ketone: Sipeimine. Kashmirine. Imperialine
[61825-98-7]
C₂₇H₄₃NO₃ 429.642
Alkaloid from *Fritillaria ussuriensis*, *Fritillaria meleagris*, *Fritillaria imperialis*, *Fritillaria raddeana*, *Petilium eduardii*, *Petilium raddeana*, the Chinese drug Si-pei-mu (*Fritillaria roylei*) and from *Rhinopetalum bucharicum* (Liliaceae). Smooth muscle relaxant. Transitory antihypertensive and anti-inflammatory agent. Mp 265-267° (262-265°). $[\alpha]_D$ -40.4 (CHCl₃). Log P 3.26 (uncertain value) (calc).

► FL5830000

6-Ketone, hydrochloride: [22267-72-7]
Mp 210-211°.

6-Ketone, N-oxide: Imperialine N-oxide
[62565-72-4]
C₂₇H₄₃NO₄ 445.641
Alkaloid from *Petilium eduardii* and *Petilium raddeana* (Liliaceae). Mp 266-268°. $[\alpha]_D$ -48.2 (c, 0.68 in MeOH).

6-Ketone, 3-O- β -D-glucopyranoside: Edpetiline

[32685-93-1]
[30497-28-0 (HCl salt)]
C₃₃H₅₃NO₈ 591.784
Alkaloid from *Petilium eduardii* (Liliaceae). Anti-inflammatory agent. Mp 272-276°. $[\alpha]_D$ -59.9 (MeOH). Log P 0.97 (uncertain value) (calc).

6-Ketone, 3-Ac: Mp 183-185°.

3,6-Diketone: 20-Hydroxy-3,6-cevane-dione. Sipeimone. Imperialone

[61989-75-1]
C₂₇H₄₁NO₃ 427.626
Alkaloid from *Petilium eduardii* and *Petilium raddeana* (Liliaceae). Cryst. (Me₂CO). Mp 228-231°. $[\alpha]_D$ -31.7 (petrol).

(3 β ,5 α ,17 β ,22 β)-form

6-Ketone, 3-O- β -D-glucopyranoside: Hupehemonoside
[143120-88-1]
C₃₃H₅₃NO₈ 591.784
Alkaloid from *Fritillaria hupehensis*. Amorph. powder. Mp 206-208°. $[\alpha]_D^{27}$ -48.3 (c, 0.46 in MeOH).

Chu, T.T. et al., *Huaxue Xuebao*, 1956, **22**, 205; *CA*, **51**, 445d (Sipeimone)

Chu, T.T. et al., *CA*, 1959, **53**, 7503e (Sipeimone, struct)

Morimoto, H. et al., *Chem. Pharm. Bull.*, 1960, **8**, 302-307 (Peiminoside)
Ito, S. et al., *Chem. Pharm. Bull.*, 1961, **9**, 253-255; 1963, **11**, 1337-1340 (Verticine)
Shakirov, R. et al., *CA*, 1965, **63**, 3007f; 1966, **64**, 68046 (Edpetiline)

Nuriddinov, R.N. et al., *Khim. Prir. Soedin.*, 1967, **3**, 316-324; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 267-273 (Imperialine)

Ito, S. et al., *Tet. Lett.*, 1968, 5373-5375 (Verticinone, *cryst struct*)

Nabiev, A. et al., *Khim. Prir. Soedin.*, 1976, **12**, 676-677; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 612-613 (Imperialine N-oxide)

Chatterjee, A. et al., *Tet. Lett.*, 1976, 2903-2904 (Imperialine, *cryst struct*)

Ito, S. et al., *Tet. Lett.*, 1976, 3161-3164 (Imperialine, *cryst struct, bibl, cd*)

Kutney, J.P. et al., *J.A.C.S.*, 1977, **99**, 964-966 (Verticine, *synth*)

Saidkasymov, T. et al., *CA*, 1979, **90**, 16179j (Imperialine, *pharmacol*)

Kaneko, K. et al., *Tet. Lett.*, 1979, 3737-3740 (Baimonidine, Isoverticine, *cmr*)

Kaneko, K. et al., *Chem. Pharm. Bull.*, 1980, **28**, 1345-1346 (Isobaimonidine)

Kitajima, J. et al., *Heterocycles*, 1981, **15**, 791-796 (Verticine N-oxide, Verticinone N-oxide)
Nakhatov, I. et al., *Khim. Prir. Soedin.*, 1981, **5**, 616-619 (Imperialone)

Kettmann, V. et al., *Acta Cryst. B*, 1982, **38**, 978-980 (Verticinone, *cryst struct*)

Mašterová, I. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1982, **315**, 157-164 (Isobaimonidine)

Xu, D. et al., *CA*, 1982, **97**, 107068h (*isol*)
Kettman, V. et al., *Acta Cryst. C*, 1985, **41**, 392-394 (Isobaimonidine, *cryst struct*)

Li, Q. et al., *Yaoxue Xuebao*, 1986, **21**, 767-771; *CA*, **106**, 135231d (Wanpeinine A)

Samikov, K. et al., *Khim. Prir. Soedin.*, 1989, **25**, 34-36; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 28-30 (Korselidine)

Zhang, J.X. et al., *Yaoxue Xuebao*, 1991, **26**, 231-233; 1992, **27**, 472-475 (Zhebeinine, Zhebeinone)

Wu, J.Z. et al., *Yaoxue Xuebao*, 1991, **26**, 829-835 (Hupehemonoside)

Wang, F. et al., *Chin. Chem. Lett.*, 1992, **3**, 979-980 (Siechuansine)

Zhang, J. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1993, **35**, 238-241 (Zhebeininoside)

Shekikova, U.T. et al., *Khim. Prir. Soedin.*, 1994, **30**, 524-528; *Chem. Nat. Compd. (Engl. Transl.)*, 1994, **30**, 483-486 (Petine, Petine N-oxide)

Zhang, A. et al., *Planta Med.*, 1998, **64**, 448-450 (Isoverticine, Imperialine N-oxide, *pmr, cmr, cryst struct*)

Jiang, R.-W. et al., *Acta Cryst. C*, 2001, **57**, 170-171 (Imperialine, *cryst struct*)

Chen, Q. et al., *Yaoxue Xuebao*, 2004, **39**, 348-350 (Isoverticine N-oxide)

Zhang, Y.-H. et al., *Chem. Biodiversity*, 2008, **5**, 259-266 (3-Acetylverticine, 3-Acetylverticinone)

Cevanine

C-299

C₂₇H₄₄N₂O₂ 428.657


Struct. unknown. Steroidal alkaloid (prob. Cevane type). Alkaloid from corms of *Fritillaria imperialis* (Liliaceae). Mp 272-273°. $[\alpha]_D$ +49.

Suri, J.L. et al., *Indian J. Chem.*, 1969, **7**, 1057-1058 (*isol*)

12,14,16-Cevatriene-3,6-diol

C-300

3,6-Dihydroxy- $\Delta^{12,14,16}$ -cevanene



C₂₇H₃₉NO₂ 409.611**(3β,6β)-form****Heilonine**

[128351-94-0]

Alkaloid from the Chinese herbal drug Ping-bei-mu (*Fritillaria ussuriensis*) (Liliaceae). Prisms (MeOH). Spar. sol. most org. solvs. Mp 284-286°.


Di-Ac: Mp 243-246°. [α]_D +34 (c, 1.05 in CHCl₃).

Kitamura, Y. *et al.*, *Tetrahedron*, 1989, **45**, 7281-7286 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Cev-5-ene-3,20-diol

C-301

5,6-Didehydro-3,20-cevamediol



(3α,25β)-form

C₂₇H₄₃NO₂ 413.642**(3α,25β)-form****Fritillarizine**

[76733-86-3]

Alkaloid from aerial parts of *Fritillaria verticillata* (Liliaceae). Mp 141.5-143°. [α]_D -18.6 (c, 1.0 in CHCl₃).

(3β,25β)-form**Veraflorizine**

[70598-85-5]

Alkaloid from the rhizomes of *Veratrum grandiflorum* (Liliaceae). Mp 175-176°. [α]_D -91 (c, 1 in CHCl₃).


Kaneko, K. *et al.*, *Tet. Lett.*, 1978, **19**, 4801-4804 (*Veraflorizine*)

Kaneko, K. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 3711-3713 (*Fritillarizine*)

Cev-8-ene-3,6-diol

C-302

8,9-Didehydro-3,6-cevamediol



(3α,5α,6α,25α)-form

C₂₇H₄₃NO₂ 413.642**(3α,5α,6α,25α)-form****Korseverinine**

[36506-64-6]

From *Korolkowia sewerzowii* (Liliaceae). Mp 320-322°. [α]_D +23 (c, 0.82 in 10% AcOH aq.).

6-Ketone: 3-Hydroxycev-8-en-6-one. **Korseverine**

[22223-08-1]

C₂₇H₄₁NO₂ 411.626

Alkaloid from *Korolkowia sewerzowii* bulbs (Liliaceae). Cryst. (Me₂CO). Mp 167-168°. [α]_D +82.5 (c, 1.21 in

MeOH).

6-Ketone, hydrobromide: Mp 292°.

6-Ketone, Ac: Mp 185-186°.

(3α,5α,6β,25α)-form**Korsidine**

[61989-84-2]

Alkaloid from bulbs of *Korolkowia sewerzowii* (Liliaceae). Mp 316-318°. [α]_D 0 (c, 0.89 in AcOH aq.). Probable struct.

(3β,5α,6β,25β)-form**Korsinine**

[22958-10-7]

Alkaloid from bulbs of *Korolkowia sewerzowii* (Liliaceae). Cryst. (Me₂CO). Mp 164-165°. [α]_D +105 (c, 0.77 in MeOH). Probable struct.


Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 390-392; 1969, **5**, 61-62; 1971, **7**, 767-773; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 332-333; 1969, **5**, 54-55; 1971, **7**, 740-745 (*Korseverine, Korsinine, Korseverinine*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 367-371; 780-783; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 320-322; 699-701 (*Korsidine, struct*)

Cev-8-ene-3,11-diol

C-303

8,9-Didehydro-3,11-cevamediol, 9CI

C₂₇H₄₃NO₂ 413.642**(3α,5α,11β,20α,25α)-form****Cordiline**


[97745-01-2]

Alkaloid from *Korolkowia sewerzowii* epigeal parts (Liliaceae). Cryst. (Me₂CO). Mp 286-288°.

Kulkova, V.V. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 253-255; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 237-239 (*isol, ir, pmr, ms, struct*)

Cev-8(14)-ene-1,3-diol

C-304

8,14-Didehydro-1,3-cevamediol, 9CI. 1,3-Dihydroxy-Δ⁸⁽¹⁴⁾-cevaneC₂₇H₄₃NO₂ 413.642**(1α,3β)-form****Korselimine**

[122279-80-5]


Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 272-274°. [α]_D -75 (c, 0.12 in EtOH).

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1989, **25**, 44-46; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 37-39

Cev-8(14)-ene-3,6-diol

C-305

8,14-Didehydro-3,6-cevamediol, 9CI



(3α,5α,6β,25β)-form

C₂₇H₄₃NO₂ 413.642**(3α,5α,6β,25β)-form****Seveline**

[73307-46-7]

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 145-147°. [α]_D -48.8 (c, 1.76 in EtOH/CHCl₃ 1:1).

(3β,5α,6β,25α)-form**Severzine**

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 202-204°. [α]_D -45.

(5α,6β,25β)-form*3-Ketone*: Sevelinone

[73307-48-9]

Cryst. (Me₂CO). Mp 234-236°.


Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 529-532; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 459-461 (*Seveline, Sevelinone, Seveline*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 252-253 (*Severzine*)

Cev-8(14)-ene-3,11-diol

C-306

8,14-Didehydro-3,11-cevamediol

C₂₇H₄₃NO₂ 413.642**(3β,5α,11β, 20ξ,25ξ)-form****Stenanzamine**


[94898-74-5]

Alkaloid from the aerial parts of *Rhinopetalum stenatherum* (Liliaceae). Amorph. [α]_D -22.3 (c, 0.05 in CHCl₃).

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 341-344; 399-400; *Chem. Nat. Compd.*

(Engl. Transl.), 1984, **20**, 318-320; 379-380
(isol, ir, pmr, ms, struct)

Cev-8(14)-ene-3,15-diol C-307
8,14-Didehydro-3,15-cevanediol



(3 α ,5 α ,15 β ,25 α)-form

C₂₇H₄₃NO₂ 413.642

(3 α ,5 α ,15 β ,25 α)-form
Korseveridinine
[62959-88-0]

Alkaloid of *Korolkowia sewerzowii* (Liliaceae). Mp 282-284°. [α]_D -39.5 (c, 0.58 in CHCl₃/MeOH, 1:1).

3-Ketone: Mp 122-124°.

(3 β ,5 α ,15 β ,25 α)-form
Korseveridine

[20072-14-4]


From *Korolkowia sewerzowii* (Liliaceae). Cryst. (MeOH). Mp 290-292°. [α]_D -49.3 (CHCl₃).

Hydrochloride: Mp 326°.

Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 101-106; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 86-89 (Korseveridine)

Abdullaeva, D.U. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 796-798; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 714-715 (Korseveridinine, struct)

Cev-13(17)-ene-3,6-diol C-308



C₂₇H₄₃NO₂ 413.642

(3 β ,5 α ,6 β ,25 α)-form
Ebeienine. Ziebeimine
[123116-14-3]

Alkaloid from the bulbs of *Fritillaria ebeiensis* var. *purpurea* (Liliaceae). Needles + 1H₂O (hexane/Et₂NH/EtOH). Mp 274.5-278.4°. [α]_D -2.9 (c, 0.5 in MeOH).

6-Ketone: 3-Hydroxycev-13(17)-en-6-one.

Ebeinone


[125409-58-7]

C₂₇H₄₁NO₂ 411.626

Alkaloid from bulbs of *Fritillaria imperialis* (Liliaceae). Exhibits anticholinergic activity. Brown amorph. solid. Mp 199°. [α]_D -30 (c, 0.1 in CHCl₃).

Lee, P. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 4316-4329 (isol, ir, pmr, cmr, ms, cryst struct)
Wu, J.Z. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 600-605; *CA*, **112**, 95497k (isol)
Atta-ur-Rahman, *et al.*, *Planta Med.*, 1994, **60**, 377-379 (Ebeinone)
Zhang, Y.-H. *et al.*, *Chin. J. Chem.*, 2007, **25**, 1728-1731 (isol, pmr, cmr)

Cev-5-ene-3,16,20-triol C-309
5,6-Didehydro-3,16,20-cevanetriol



C₂₇H₄₃NO₃ 429.642

(3 β ,16 β ,25 β)-form
Veramarine

[4565-85-9]

Alkaloid from *Veratrum album* ssp. *lobelianum*, *Veratrum oxyspalum* and *Veratrum nigrum* (Liliaceae). Cryst. (Me₂CO aq.) (prev. descr. as amorph.). [α]_D -77 (c, 0.7 in EtOH). [α]_D -112.7 (c, 0.22 in CHCl₃). Formerly assigned the 16 α -config.

3-Ac: Mp 254-255°. [α]_D -74 (c, 0.44 in EtOH).


Tomko, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 3320-3324 (Veramarine)

Kaneko, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1983, **48**, 2840-2843 (Veramarine, cryst struct, cmr, pmr)

Kuribayashi, T. *et al.*, *Coll. Czech. Chem. Comm.*, 1983, **48**, 2840 (cmr)

Bondarenko, N.V. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 263; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 248-249 (Veramarine, isol)

Cev-8-ene-3,6,20-triol C-310
8,9-Didehydro-3,6,20-cevanetriol, 9CI



C₂₇H₄₃NO₃ 429.642

(3 α ,5 α ,6 β ,20 β ,25 α)-form
Edpetisidine


[68676-58-4]

Alkaloid from *Petilium eduardii* (Liliaceae). Mp 257-259°. [α]_D -33.6 (c, 1.87 in

MeOH). Probable struct.

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 416-417; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 357-358 (ir, ms, pmr)

Cev-8-ene-3,6,23-triol C-311
8,9-Didehydro-3,6,23-cevanetriol



Absolute Configuration

C₂₇H₄₃NO₃ 429.642

(3 β ,5 α ,6 β)-form

Korsine

[20321-59-9]

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 236-238°. [α]_D +87.9 (c, 0.1 in EtOH). Incorr. named as 3,11,23-triol in CAS.

Hydrochloride:

Cryst. (Me₂CO). Mp 301-303°.

N-Oxide: **Korsine N-oxide**

[105256-28-8]

C₂₇H₄₃NO₄ 445.641

Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Cryst. (Me₂CO). Mp 257-259°. [α]_D -8 (c, 0.74 in EtOH).

23-Ac: **Korsinamine**

[68231-27-6]

C₂₉H₄₅NO₄ 471.679


Alkaloid from *Korolkowia sewerzowii*. Mp 155-158°. [α]_D +68.2 (c, 1.202 in CHCl₃).

Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 161-168; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 139-144 (Korsine)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 233-235; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 192-193 (Korsinamine, ir, pmr, ms)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 251; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 237 (Korsine N-oxide)

Cev-12-ene-3,6,15-triol C-312
12,13-Didehydro-3,6,15-cevanetriol



C₂₇H₄₃NO₃ 429.642

(3 β ,6 β ,15 ξ)-form
Edpetisidine


[73276-38-7]

Alkaloid from *Petilium eduardii* (Liliaceae). Cryst. (MeOH). Mp 263-265°. [α]_D -15.3 (c, 0.39 in MeOH/CHCl₃ 9:1).

Shakirov, P. *et al.*, *Khim. Prir. Soedin.*, 1979,

15, 584; *Chem. Nat. Compd. (Engl. Transl.)*,
512 (*isol, ir, ms, pmr, struct*)

Cev-5-en-3-ol C-313
5,6-Didehydrocevan-3-ol, 9CI



C₂₇H₄₃NO 397.643


**(3β,20β)-form
Shinonomenine**

[70598-84-4]

Alkaloid from the rhizomes of *Veratrum grandiflorum* (Liliaceae). Mp 95-96°. [α]_D²⁵ -90.7 (c, 0.33 in CHCl₃).

Kaneko, K. *et al.*, *Tet. Lett.*, 1978, 4801-4804;
1979, 3737-3740 (*isol, cryst struct, pmr, ms, cmr*)

Ceylanicine C-314
[119400-91-8]




C₄₁H₄₈N₄O₆ 692.853

Related to Pelankine, P-172. Alkaloid from the stems of *Petchia ceylanica* (Apocynaceae). Light yellow amorph. solid. Mp 250° dec. [α]_D²⁶ -98 (MeOH).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1988,
27, 2051 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ceylanine C-315
[119420-11-0]




C₄₂H₅₀N₄O₇ 722.88

Alkaloid from the leaves of *Petchia ceylanica* (Apocynaceae). Pale pink amorph. solid. Mp 280° dec. [α]_D²⁶ -124 (MeOH). Light sensitive.

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1988,
27, 2051 (*isol, uv, ir, pmr, cmr, ms, struct*)

Chabamide C-316
[479068-69-4]




C₃₄H₃₈N₂O₆ 570.684

Dimer of 1-Piperoylpiperidine, P-467. Alkaloid from the stems of *Piper chaba* and roots of *Piper nigrum* (pepper). Antimalarial and antituberculosis agent. Pale yellow semi-solid. [α]_D³¹ +11.6 (c, 0.15 in CHCl₃). λ_{max} 278 ; 297 ; 317 (sh) (MeOH).

Rukachaisirikul, T. *et al.*, *Planta Med.*, 2002,
68, 850-853 (*isol, pmr, cmr*)
Wei, K. *et al.*, *J.O.C.*, 2005, 70, 1164-1176
(*isol*)

Chaenorhine C-317
20-Acetyl-5-methoxy-7-oxa-15,20,24,27-tetraazatetracyclo[13.9.6.2^{8,11}.1^{2,6}]tritiaconta-2,4,6(33),8,10,12,31-heptaene-14,26-dione, 9CI. *Chenorrhine*
[40135-65-7]




C₃₁H₄₀N₄O₅ 548.681

Alkaloid from *Chaenorhinum origanifolium* and *Chaenorhinum rubifolium* (Scrophulariaceae). Cryst. (MeOH). Mp 263-268° dec. [α]_D²⁵ +46.7 (c, 1.5008 in CHCl₃/MeOH 9:1). pK_a 6.

Bernhard, H.O. *et al.*, *Helv. Chim. Acta*, 1973,
56, 1266 (*isol, uv, ord, cd, ir, pmr, ms, struct*)
Wasserman, H.H. *et al.*, *J.A.C.S.*, 1983, 105,
1697 (*synth*)

Chaenorpine C-318
[114622-14-9]



C₂₈H₃₆N₄O₄ 492.617

Alkaloid from *Chaenorhinum minus* (Scrophulariaceae). Cryst. (EtOH aq.) (as dihydrobromide). Mp 239-241° (dihydrobromide). [α]_D²² +28.2 (c, 0.22 in H₂O) (dihydrobromide).

Me ether: 18-O-Methylchaenorpine
[120374-27-8]

C₂₉H₃₈N₄O₄ 506.644

Alkaloid from the roots of *Aphelandra tetragona* (Acanthaceae). Amorph. solid. [α]_D²² -74 (c, 0.10 in MeOH).

13E-Isomer, Me ether: Iso-18-O-methylchaenorpine

[120396-81-8]


C₂₉H₃₈N₄O₄ 506.644

Alkaloid from the roots of *Aphelandra tetragona* (Acanthaceae). [α]_D²² +108.9 (c, 0.235 in MeOH).

Zhu, J. *et al.*, *Helv. Chim. Acta*, 1988, 71, 218
(*uv, ir, pmr, cmr, ms, cd, struct*)

Tawil, B.F. *et al.*, *Helv. Chim. Acta*, 1989, 72,
180 (*18-O-Methylchaenorpine, Iso-18-O-methylchaenorpine*)

Chaetochalasin A C-319
[216006-03-0]




C₂₇H₃₉NO₂ 409.611

Prod. by *Chaetomium brasiliense*. Anti-bacterial and cytotoxic agent. Cryst. (MeOH). Mp 191-193°. [α]_D²⁵ +271 (c, 0.27 in CHCl₃). λ_{max} 212 (ε 6200) (MeOH).

Oh, H. *et al.*, *Tet. Lett.*, 1998, 39, 7633-7636
(*isol, uv, ir, pmr, cmr, ms*)

Chaetocin C-320
Chetocin, 9CI
[28097-03-2]



Absolute
Configuration

x = y = 2

C₃₀H₂₈N₆O₆S₄ 696.852

Epithiodioxopiperazine antibiotic. Stereoisomeric with the series repre-

sented by Verticillin A, V-89. The original publication incorrectly shows the stereochem. of Chaetocin as a *meso*-form. The struct. has C_2 -symmetry with the bridgehead *Hs cis*- to the bond linking the two C_{15} residues. CAS numbering shown. Metab. of *Chaetomium minutum*, *Chaetomium thielavioideum* and *Farrowia* sp. Sol. Py, EtOAc, DMSO, MeOH; poorly sol. H_2O . $[\alpha]_D^{20} +379$ (c, 1 in Py). λ_{max} 203 (ϵ 50100); 240 (sh) (ϵ 14100); 301 (ϵ 5370) ($Ac_2O/MeOH$) (Derep). λ_{max} 306 (ϵ 6030) (DMSO) (Derep).
 ▶ LD_{50} (mus, ipr) 1.7 mg/kg, LD_{50} (mus, orl) 1200 mg/kg. FM3032000
 O,O'-Di-Ac: Mp 220-225° dec. $[\alpha]_D^{20} +679$ (c, 1 in $CHCl_3$).

6S-Hydroxy: Melinacidin III

[12794-84-2]
 $C_{30}H_{28}N_6O_7S_4$ 712.851
 Prod. by *Acrostalagmus cinnabarinus* var. *melinacidinus* NRRL13291 and the marine-derived *Corollospora pulchella*. Shows similar activity to Melinacidin II. Cryst. Sol. $CHCl_3$, DMF, DMSO; fairly sol. MeOH, EtOH, Me_2CO ; poorly sol. H_2O , hexane. $[\alpha]_D^{25} +776$ (c, 0.5 in $CHCl_3$). λ_{max} 241 (sh) (ϵ 15300); 300 (ϵ 5310) (MeOH) (Derep).

19-Deoxy, 6 or 6'S-hydroxy: Melinacidin II

[12794-85-3]
 $C_{30}H_{28}N_6O_6S_4$ 696.852
 Prod. by *Acrostalagmus cinnabarinus* var. *melinacidinus* NRRL13291 and *Verticillium terereri*. Shows antibacterial and antitumour activity but is highly toxic. Cryst. Sol. $CHCl_3$, DMSO, DMF; fairly sol. MeOH, EtOH, Me_2CO ; poorly sol. H_2O , hexane. $[\alpha]_D^{25} +726$ (c, 0.5 in $CHCl_3$). λ_{max} 203 (ϵ 50100); 240 (sh) (ϵ 14100); 301 (ϵ 5370) ($Ac_2O/MeOH$) (Derep). λ_{max} 306 (ϵ 6030) (DMSO) (Derep). λ_{max} 300 ($E1\%/1cm$ 73) (MeOH) (Berdy).

6S,6'S-Dihydroxy: Melinacidin IV

[37934-52-4]
 $C_{30}H_{28}N_6O_8S_4$ 728.851
 Prod. by *Acrostalagmus cinnabarinus* var. *melinacidinus* NRRL13291 and the marine-derived *Corollospora pulchella*. Shows similar activity to Melinacidin II. Cryst. $[\alpha]_D^{25} +718$ (c, 0.5 in $CHCl_3$). λ_{max} 241 (sh) (ϵ 15300); 300 (ϵ 5310) (MeOH) (Derep).

▶ FM3036000

Hauser, D. et al., *Helv. Chim. Acta*, 1970, **53**, 1061-1073 (isol, uv, cd, ir, pmr, ms, cryst struct)
 Argoudelis, A.D. et al., *J. Antibiot.*, 1972, **25**, 171-178; 1977, **30**, 468 (Melinacidins)
 Nagarajan, R. et al., *J.A.C.S.*, 1973, **95**, 7212-7222 (cd)
 Sekita, S. et al., *Chem. Pharm. Bull.*, 1980, **28**, 2428-2435 (isol)

Chaetocin B C-321

2,5-De(epidithio)-2,5-(epitritio)chaetocin, 9CI. Chetocin B
 [118101-82-9]
 As Chaetocin, C-320 with $x = 3$, $y = 2$
 $C_{30}H_{28}N_6O_6S_5$ 728.918

Epithiodioxopiperazine antibiotic. Metab. of *Chaetomium virescens* ssp. *thielavioideum*. Cytotoxic. Active against gram-positive bacteria. Sol. MeOH. Mp 225° dec. (as di-Ac). $[\alpha]_D^{20} +673.2$ (c, 0.079 in $CHCl_3$) (di-Ac). λ_{max} 209 ; 237 ; 299 (MeOH) (Berdy).

Saito, T. et al., *Chem. Pharm. Bull.*, 1988, **36**, 1942 (isol, pmr, cmr, struct)

Chaetocin C C-322


Chetocin C
 [118111-08-3]
 As Chaetocin, C-320 with $x = y = 3$

$C_{30}H_{28}N_6O_6S_6$ 760.984
 Epithiodioxopiperazine antibiotic. Metab. of *Chaetomium virescens* ssp. *thielavioideum*. Cytotoxic, active against gram-positive bacteria. Sol. MeOH. Mp 232° dec. (as di-Ac). $[\alpha]_D^{20} +658.3$ (c, 0.012 in $CHCl_3$) (di-Ac). λ_{max} 210 ; 238 ; 301 (MeOH) (Berdy).

Saito, T. et al., *Chem. Pharm. Bull.*, 1988, **36**, 1942 (isol, pmr, cmr, struct)

Chaetocochin A C-323

[912551-37-2]



$C_{36}H_{42}N_6O_6S_4$ 783.028
 Prod. by *Chaetomium cochliodes*. Amorph. powder. $[\alpha]_D^{20} +121.3$ (c, 0.1 in MeOH). λ_{max} 205 (log ϵ 3.64); 220 (log ϵ 3.49); 287 (log ϵ 2.78); 297 (log ϵ 2.77) (MeOH).


3,11a-Di-S-de-Me, 3,11a-disulfide: Chaetocochin B

[912551-38-3]
 $C_{34}H_{36}N_6O_6S_4$ 752.959
 Prod. by *Chaetomium cochliodes*. Amorph. powder. $[\alpha]_D^{20} +204$ (c, 0.11 in $CHCl_3$).

Li, G.-Y. et al., *J. Nat. Prod.*, 2006, **69**, 1374-1376 (isol, cd, pmr, cmr)

Chaetocochin C C-324

[912551-39-4]




Absolute Configuration

$C_{33}H_{36}N_6O_6S_4$ 740.948
 Prod. by *Chaetomium cochliodes*. Amorph. powder. $[\alpha]_D^{20} +206$ (c, 0.1 in $CHCl_3$). λ_{max} 205 (log ϵ 3.67); 220 (log ϵ 3.56); 287 (log ϵ 2.85); 294 (log ϵ 2.85) (MeOH).

Li, G.Y. et al., *J. Nat. Prod.*, 2006, **69**, 1374-1376

Chaetoglobosin A C-325

[50335-03-0]



Absolute Configuration

$C_{32}H_{36}N_2O_5$ 528.647
 Isoindole antibiotic. Metab. of *Chaetomium cochliodes*, *Chaetomium globosum* NRRL12718, *Chaetomium subaffine*, other *Chaetomium* spp., *Penicillium aurantiogriseum*, *Cylindroccladium floridanum* and *Ascotricha chartarum*. Cytotoxic to HeLa cells; fibrinolysis enhancer; mycotoxin; phytotoxin; microtubule protein assembly inhibitor; urokinase prodn. inducer. Pale yellow prisms (CH_2Cl_2). Sol. MeOH; poorly sol. H_2O . Mp 188° (168-170°). $[\alpha]_D -270$ (MeOH). λ_{max} 222 (ϵ 36300); 273 (ϵ 6760); 281 (ϵ 6760); 291 (ϵ 5750) (MeOH) (Derep). λ_{max} 196 (ϵ 42700); 220 (ϵ 41600); 271 (ϵ 6600); 280 (ϵ 6600); 290 (ϵ 5360) (MeOH) (Berdy). λ_{max} 223 (ϵ 40740); 274 (ϵ 6610); 282 (ϵ 6607); 292 (ϵ 5370) (EtOH) (Berdy).
 ▶ Teratogen. HA5305000

O-Ac: 19-O-Acetylchaetoglobosin A

[50939-69-0]
 $C_{34}H_{38}N_2O_6$ 570.684
 Prod. by *Chaetomium globosum*. Mycotoxin. Yellow cryst. (C_6H_6). Poorly sol. hexane. Mp 223-225°. $[\alpha]_D^{20} -304$ (c, 0.5 in $CHCl_3$). λ_{max} 196 (log ϵ 4.65); 220 (log ϵ 4.64); 271 (log ϵ 3.83); 280 (log ϵ 3.83); 289 (log ϵ 3.74) (EtOH).

20S-Alcohol: 20-Dihydrochaetoglobosin A

TAN 1142. Antibiotic TAN 1142 [149560-98-5] [147527-33-1]
 $C_{32}H_{38}N_2O_5$ 530.663
 Prod. by *Chaetomium globosum* and *Chaetomium subaffine*. Immunoregulator; mycotoxin. Cryst. ($CHCl_3/Et_2O$). Mp 145-147°. $[\alpha]_D -82.8$ (MeOH). Stereochem. relationship between 20-Dihydrochaetoglobosin A and TAN 1142 not confirmed. λ_{max} 212 (log ϵ 4.71); 221 (log ϵ 4.8); 266 (log ϵ 3.93); 272 (log ϵ 3.92); 291 (log ϵ 3.82) (MeOH).

19-Deoxy: Prochaetoglobosin III

[133613-78-2]

C₃₂H₃₆N₂O₄ 512.647

Metab. of *Chaetomium subaffine*. Pale yellow cryst. (MeOH aq.). Mp 137-139°. [α]_D -210.4 (c, 0.2 in MeOH). λ_{max} 221 (log ε 4.7); 280 (log ε 3.87); 290 (log ε 3.79) (MeOH).

19-Deoxy, 20ξ-alcohol: 20-Dihydroprotochaetoglobosin III

[149439-84-9]

C₃₂H₃₈N₂O₄ 514.663

Prod. by *Chaetomium subaffine*. Proposed biosynthetic intermed. to Chaetoglobosin A. Oil. [α]_D -60 (c, 0.06 in MeOH). λ_{max} 222 (log ε 4.63); 291 (log ε 3.72) (MeOH).

19-Deoxy, 20-deoxo: Prochaetoglobosin IV

[137604-97-8]

C₃₂H₃₈N₂O₃ 498.664

Metab. of *Chaetomium subaffine*. Cryst. (CHCl₃). Mp 256-258°. [α]_D -54.5 (c, 0.02 in MeOH). λ_{max} 222 (log ε 4.45); 291 (log ε 3.67) (MeOH).

21,22-Dihydro: Penochalasin F

[345642-82-2]

C₃₂H₃₈N₂O₅ 530.663

Prod. by a marine *Penicillium* sp. Cytotoxic. Oil. [α]_D -80 (c, 0.13 in CHCl₃). λ_{max} 220 (log ε 4.68); 240 (sh) (log ε 4.28); 280 (log ε 4.17); 291 (log ε 4.1) (EtOH).

21,22-Dihydro, 19-ketone: Chaetoglobosin C

[50645-76-6]

C₃₂H₃₆N₂O₅ 528.647

Prod. by *Chaetomium cochliodes*, *Chaetomium globosum* NRRL12718 and other organisms. Cytotoxic to HeLa cells. Mycotoxin. Phycotoxin. Leaflets (Me₂CO). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 259-261°. [α]_D -30 (c, 0.1 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 220 (ε 47860); 248 (ε 9800); 281 (ε 6610); 290 (ε 5490) (EtOH) (Berdy).

21,22-Dihydro, 19-ketone, 20β-alcohol: Chaetoglobosin F

[55945-75-0]

C₃₂H₃₈N₂O₅ 530.663

Prod. by *Chaetomium cochliodes*, *Chaetomium subaffine* and *Chaetomium globosum* NRRL12718. Cytotoxic to HeLa cells; mycotoxin. Leaflets (C₆H₆). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 177-178°. [α]_D -69 (c, 0.10 in CHCl₃). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 222; 276; 283; 292 (EtOH) (Berdy).

19-Epimer, 21,22-dihydro: Penochalasin E

[345642-81-1]

C₃₂H₃₈N₂O₅ 530.663

Prod. by a marine *Penicillium* sp. Cytotoxic. Oil. [α]_D +73 (c, 0.14 in CHCl₃). λ_{max} 221 (log ε 4.43); 241 (sh) (log ε 3.97); 280 (log ε 3.84); 291 (log ε 3.77) (EtOH).

Umeda, M. *et al.*, *Experientia*, 1975, **31**, 435-438 (*isol*)

Silverton, J.V. *et al.*, *Acta Cryst. B*, 1978, **34**, 588-593 (*cryst struct*)

Probst, A. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 2056-2064; 2065-2077 (*19-Ac*)

Sekita, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1609-1617; 1618-1628; 1629-1638; 1983, **31**, 490-498 (*isol, uv, pmr, cmr, struct, Chaetoglobosin F*)

Oikawa, H. *et al.*, *J.C.S. Perkin 1*, 1992, 2949-2953; 2955-2959 (*Prochaetoglobosins*)

Japan. Pat., 1992, 92 360 691; *CA*, **118**, 25339g (*TAN 1142*)

Oikawa, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 628-631 (*20-Dihydroprotochaetoglobosin II, 20-Dihydrochaetoglobosin A*)

Shinohara, C. *et al.*, *J. Antibiot.*, 2000, **53**, 762-768 (*activity*)


Iwamoto, C. *et al.*, *Tetrahedron*, 2001, **57**, 2997-3004 (*Penochalasin*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., *J. Wiley*, 2000, CDG750

Chaetoglobosin B

C-326

[50335-04-1]

C₃₂H₃₆N₂O₅ 528.647

Prod. by *Chaetomium cochliodes*, *Chaetomium rectum*, *Chaetomium mollipilum* and *Chaetomium globosum*. Cytotoxic to HeLa cells. Mycotoxin. Pale yellow needles (C₆H₆). Sol. MeOH; poorly sol. H₂O. Mp 186-187°. [α]_D -176 (c, 0.1 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 222 (ε 43650); 274 (ε 7940); 281 (ε 7950); 290 (ε 6760) (EtOH) (Berdy).

19-Ac: 19-O-Acetylchaetoglobosin B

[80375-18-4]

C₃₄H₃₈N₂O₆ 570.684

Prod. by *Chaetomium globosum* and a marine-derived *Acremonium* sp. (MST-MF588a). Mycotoxin. Yellow needles (C₆H₆). Poorly sol. hexane. Mp 154-157°. [α]_D²⁰ -148 (c, 0.5 in CHCl₃). λ_{max} 221 (log ε 4.64); 272 (log ε 3.83); 279 (log ε 3.83); 289 (log ε 3.74) (EtOH).

21,22-Dihydro: Chaetoglobosin O

[52645-09-7]

C₃₂H₃₈N₂O₅ 530.663

Prod. by *Cylindrocladium floridanum*. Phytotoxin. Mycotoxin. Oil. [α]_D -168 (CHCl₃).

21,22-Dihydro, 19-ketone: Chaetoglobosin G

[65773-98-0]

C₃₂H₃₆N₂O₅ 528.647

Prod. by *Chaetomium cochliodes* and *Chaetomium subaffine* ATCC22132. Cytotoxic to HeLa cells. Mycotoxin. Leaflets (MeOH). Mp 251-253°. [α]_D +89 (c, 0.10 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 222; 275; 282; 291 (EtOH) (Berdy).

21,22-Dihydro, 19-ketone, 20β-alcohol: Chaetoglobosin E

[55945-74-9]

C₃₂H₃₈N₂O₅ 530.663

Prod. by *Chaetomium cochliodes*, *Chaetomium subaffine* ATCC22132, *Chaetomium mollipilum* and *Chaetomium globosum*. Cytotoxic to HeLa cells. Needles (MeOH). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 279-280°. [α]_D +158 (c, 0.1 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 221 (ε 56200); 275 (ε 7080); 281 (ε 7080); 291 (ε 6800) (EtOH) (Berdy).

19-Deoxy: Prochaetoglobosin III

[146426-37-1]

C₃₂H₃₆N₂O₄ 512.647

Metab. of *Chaetomium subaffine*. Pale yellow cryst. (CHCl₃). Mp 153-155°. [α]_D -15.3 (c, 0.06 in CHCl₃). λ_{max} 222 (log ε 4.43); 282 (log ε 3.66); 291 (log ε 3.6) (MeOH).

19-Epimer, 21,22-dihydro: Penochalasin H

[345642-84-4]

C₃₂H₃₈N₂O₅ 530.663

Prod. by a marine *Penicillium* sp. Powder. Mp 180-182°. [α]_D -72.7 (c, 0.18 in CHCl₃). λ_{max} 220 (log ε 4.16); 243 (sh) (log ε 3.97); 281 (log ε 3.52); 290 (log ε 3.5) (EtOH).

Umeda, M. *et al.*, *Experientia*, 1975, **31**, 435-438 (*Chaetomium globosum constitis*)

Sekita, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **30**, 1609-1617; 1618-1628; 1629-1638 (*isol, ir, uv, pmr, ms, struct, Chaetoglobosins E,G*)

Probst, A. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 2056-2064 (*19-Ac*)

Oikawa, H. *et al.*, *J.C.S. Perkin 1*, 1992, 2949-2953; 2955-2959 (*Prochaetoglobosin III*)


Ichihara, A. *et al.*, *Biosci., Biotechnol., Biochem.*, 1996, **60**, 360-361 (*Chaetoglobosin O*)

Iwamoto, C. *et al.*, *Tetrahedron*, 2001, **57**, 2997-3004 (*Penochalasin H*)

Chaetoglobosin D

C-327

[55945-73-8]



C₃₂H₃₆N₂O₅ 528.647

Prod. by *Chaetomium cochliodes* and *Chaetomium globosum*. Cytotoxic. Pale yellow prisms (CH₂Cl₂). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 216°. [α]_D -269 (c, 0.1 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 221 (ε 43650); 273 (ε 9162); 281 (ε 9160); 290 (ε 7590) (EtOH) (Berdy).

19-Ac: 19-O-Acetylchaetoglobosin DC₃₄H₃₈N₂O₆ 570.684

Prod. by *Chaetomium globosum* and a marine-derived *Acremonium* sp. (MST-MF558a). Yellow prisms (CHCl₃). Poorly sol. hexane. Mp 239-241°. [α]_D -176 (c, 0.5 in CHCl₃). λ_{max} 221 (log ε 4.64); 272 (log ε 3.83); 279 (log ε 3.83); 289 (log ε 3.74) (EtOH). λ_{max} 224 (ε 43600); 272 (ε 6760); 279 (ε 6760); 289 (ε 5500) (MeOH) (Berdy).

21,22-Dihydro, 19-ketone: Isochaetoglobosin D

[83530-07-8]

C₃₂H₃₆N₂O₅ 528.647

Prod. by *Chaetomium globosum*. Mp 264-266° dec. [α]_D +41 (c, 0.1 in MeOH). λ_{max} 222 (log ε 4.53); 275 (log ε 3.82); 281 (log ε 3.82); 291 (log ε 3.76) (EtOH).

21,22-Dihydro, 19-ketone, 20β-alcohol: Chaetoglobosin F_{ex}

[149457-95-4]

C₃₂H₃₈N₂O₅ 530.663

Prod. by *Chaetomium subaffine*. Cryst. Mp 151-153°. [α]_D +63.8 (c, 0.16 in MeOH). λ_{max} 221 (log ε 4.59); 274 (log ε 3.78); 283 (log ε 3.8); 291 (log ε 3.72) (MeOH).

Umeda, M. *et al.*, *Experientia*, 1975, **31**, 435 (isol)

Probst, A. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 2056-2064 (19-Ac)


Sekita, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1609; 1618; 1629 (ir, uv, pmr, ms, struct)

Oikawa, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 628-631 (Chaetoglobosin F_{ex})

Kanokmedhakul, S. *et al.*, *Planta Med.*, 2002, **68**, 834-836 (Isochaetoglobosin D)

Chaetoglobosin J

[65745-47-3]

C₃₂H₃₆N₂O₄ 512.647

Prod. by *Chaetomium cochliodes*, *Chaetomium globosum* and *Chaetomium subaffine*. Cytotoxic to HeLa cells. Pale yellow prisms. Mp 149-151°. [α]_D +41 (c, 0.1 in MeOH). λ_{max} 222 (ε 36300); 273 (ε 6760); 281 (ε 6760); 291 (ε 5750) (MeOH) (Derep). λ_{max} 224; 270; 280; 290 (EtOH) (Berdy).

6β,7β-Epoxyde: see Chaetoglobosin A, C-325

21,22-Dihydro: Penochalasin GC₃₂H₃₈N₂O₄ 514.663

Prod. by a marine *Penicillium* sp. Cytotoxic. Powder. Mp 124-126°. [α]_D -143.6 (c, 0.19 in CHCl₃). λ_{max} 220 (log ε 4.42); 240 (sh) (log ε 3.82); 280 (log ε 3.8) (EtOH).

21,22-Dihydro, 19-ketone: Isochaetoglobosin J

[146426-38-2]

C₃₂H₃₆N₂O₄ 512.647

Metab. of *Chaetomium subaffine*. Pale yellow cryst. (CHCl₃). Mp 117-119°. [α]_D -20 (c, 0.07 in MeOH). λ_{max} 221 (log ε 4.59); 291 (log ε 3.74) (MeOH).

19-Deoxy: Prochaetoglobosin II

[133625-26-0]

C₃₂H₃₆N₂O₃ 496.648

Metab. of *Chaetomium subaffine*. Pale yellow cryst. (MeOH). Mp 187-189°. [α]_D -204.2 (c, 0.4 in MeOH). λ_{max} 221 (log ε 4.45); 273 (log ε 3.59) (MeOH).

19-Deoxy, 20ξ-alcohol: 20-Dihydroprochaetoglobosin II

[149439-83-8]

C₃₂H₃₈N₂O₃ 498.664

Metab. of *Chaetomium subaffine*. Proposed biosynth. intermediate of Chaetoglobosin A, C-325. Oil. [α]_D -68.6 (c, 0.07 in MeOH). λ_{max} 221 (log ε 4.34); 291 (log ε 3.38) (MeOH).

20-Deoxo: Chaetoglobosin TC₃₂H₃₈N₂O₃ 498.664

Prod. by *Chaetomium globosum*. Solid. [α]_D²⁰ -80 (c, 0.1 in MeOH). λ_{max} 220 (log ε 4.39); 283 (log ε 3.58); 290 (log ε 3.54) (MeOH).

20-Deoxo, 19-deoxy: Prochaetoglobosin I

[133613-77-1]

C₃₂H₃₈N₂O₂ 482.664

Metab. of *Chaetomium subaffine*. Cryst. (MeOH). Mp 214-216°. [α]_D -101.3 (c, 0.3 in MeOH). λ_{max} 223 (log ε 4.45); 291 (log ε 3.5) (MeOH).

Sekita, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1609; 1618; 1629 (isol, ir, uv, pmr, ms, struct)

Oikawa, H. *et al.*, *J.C.S. Perkin 1*, 1992, 2949; 2955 (isol, uv, ir, pmr, cmr, ms, biosynth, derivs)

Oikawa, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 628-631 (20-Dihydroprochaetoglobosin II)


Iwamoto, C. *et al.*, *Tetrahedron*, 2001, **57**, 2997-3004 (Penochalasin G)

Jiao, W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1722-1725 (isol, pmr, cmr)

Chaetoglobosin K

[72509-61-6]

C-329

C₃₄H₄₀N₂O₅ 556.7

Cytochalasin antibiotic. Obt. from cultures of *Diplodia macrospora*. Plant growth inhibitor, toxic to chickens. Mycotoxin. Yellow prisms (Me₂CO). Mp 235-240° Mp 264-266°. λ_{max} 221 (ε 43700); 273 (ε 9120); 281 (ε 9120); 290 (ε 7590) (EtOH) (Derep). λ_{max} 219 (ε 44800) (EtOH) (Berdy).

▶ JF7954000

19-Ketone: Chaetoglobosin M

[119212-28-1]

C₃₄H₃₈N₂O₅ 554.685

Isol. from *Diplodia macrospora* DM7. Mycotoxin. Sl. yellow gum. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²² -18.3 (c, 0.92 in CH₂Cl₂). λ_{max} 230 (ε 12300); 290 (ε 3980) (CH₂Cl₂) (Derep).

Deepoxy, 6,7-didehydro: Chaetoglobosin 540C₃₄H₄₀N₂O₄ 540.701

Prod. by a marine-derived *Phomopsis asparagi* incubated with Jaspamide, J-26. Pale yellow powder. [α]_D²⁵ -118 (c, 0.009 in MeOH). λ_{max} 221 (log ε 4.43); 289 (log ε 3.36) (MeOH).

Deepoxy, 6,7-didehydro, 20β-alcohol: Chaetoglobosin 542C₃₄H₄₂N₂O₄ 542.717

Prod. by a marine-derived *Phomopsis asparagi* incubated with Jaspamide, J-26. Cytotoxic. Amorph. powder. [α]_D²⁵ -44 (c, 0.01 in MeOH). λ_{max} 222 (log ε 4.21); 281 (log ε 3.43); 290 (log ε 3.35) (MeOH).

Deepoxy, 6,7-didehydro, 19-deoxy, 20-deoxo: Chaetoglobosin 510C₃₄H₄₂N₂O₂ 510.718

Prod. by a marine-derived *Phomopsis asparagi* incubated with Jaspamide, J-26. Amorph. powder. [α]_D²⁵ -100 (c, 0.02 in MeOH). λ_{max} 222 (log ε 4.37); 281 (log ε 3.57); 290 (log ε 3.5) (MeOH).

Deepoxy, 6,12-didehydro, 7β-hydroxy: Chaetoglobosin L

[83481-23-6]


C₃₄H₄₀N₂O₅ 556.7

Obt. from cultures of *Diplodia macrospora*. Yellow gum. Not obt. completely pure.

Cutler, H.G. *et al.*, *CA*, 1980, **92**, 141619 (props)


- Cutler, H.G. *et al.*, *J. Agric. Food Chem.*, 1980, **28**, 139 (*isol, cryst struct*)
 Springer, J.P. *et al.*, *Tet. Lett.*, 1980, **21**, 1905 (*cryst struct*)
 Probst, A. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 1543 (*isol, pmr, struct*)
 Cox, R.H. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 405 (*pmr, cmr, use*)
 Spöndlin, C. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 1881 (*deriv*)
 Christian, O.E. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1592-1597 (*Chaetoglobosins 510,540,542*)

Chaetoglobosin P C-330
 [191403-93-7]



- $C_{33}H_{38}N_2O_5$ 542.674
 Isol. from the fungus *Discosia* sp. TCF 9535. Pale yellow gum.
 19-Ketone, 21,22-dihydro: **Chaetoglobosin N**
 [156980-59-5]
 $C_{33}H_{38}N_2O_5$ 542.674
 Isol. from *Phomopsis leptostromiformis* on sweet corn (*Zea mays*). Mycotoxin. Mp 205°.
 Cherton, J.C. *et al.*, *Analisis*, 1994, **22**, 210-216; 217-221 (*deriv, isol, pmr, cmr*)
 Donoso, R. *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 49-54 (*isol, pmr, cmr*)

Chaetoglobosin Q C-331




- $C_{32}H_{38}N_2O_6$ 546.662
 Prod. by *Chaetomium globosum*. Pale yellow solid. $[\alpha]_D^{20}$ -100 (c, 0.1 in MeOH). λ_{max} 217 (log ϵ 4.81); 289 (log ϵ 4.05) (MeOH).
 6-Epimer: **Chaetoglobosin R**
 $C_{32}H_{38}N_2O_6$ 546.662

- Prod. by *Chaetomium globosum*. Pale yellow solid. $[\alpha]_D^{20}$ -100 (c, 0.1 in MeOH). λ_{max} 220 (log ϵ 4.46); 282 (log ϵ 3.73) (MeOH).

- Jiao, W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1722-1725 (*pmr, cmr*)

Chaetoglobosin U C-332




Absolute Configuration

- $C_{32}H_{36}N_2O_5$ 528.647
 Prod. by *Chaetomium globosum* IFB-E019. Cytotoxic. Pale yellow powder. Mp 167-168°. $[\alpha]_D^{20}$ -17 (c, 0.18 in $CHCl_3$). λ_{max} 222 (log ϵ 0.48); 281 (log ϵ 0.09) ($CHCl_3$).

- Ding, G. *et al.*, *J. Nat. Prod.*, 2006, **69**, 302-304 (*isol, pmr, cmr, ms*)

Chaetoindicin A C-333




Relative Configuration

- $C_{16}H_{19}NO_4$ 289.33
 Prod. by *Chaetomium indicum*. Cryst. (MeOH). Mp 231-232°. $[\alpha]_D^{20}$ +8.5 (c, 0.1 in MeOH). λ_{max} 213 (log ϵ 4.29); 248 (log ϵ 4.2) (MeOH).

- Li, G.-Y. *et al.*, *Org. Lett.*, 2006, **8**, 3613-3615 (*isol, pmr, cmr, ms*)

Chaetoindicin B C-334




Relative Configuration

- $C_{16}H_{19}NO_4$ 289.33
 Prod. by *Chaetomium indicum*. Cryst. (MeOH). Mp 255-256°. $[\alpha]_D^{20}$ -7.5 (c, 0.1 in MeOH). λ_{max} 210 (log ϵ 4.36); 244 (log ϵ 4.2); 285 (log ϵ 3.92); 309 (log ϵ 3.96) (MeOH).

- 5-Epimer: **Chaetoindicin C**
 $C_{16}H_{19}NO_4$ 289.33
 Prod. by *Chaetomium indicum*. Amorph. powder. $[\alpha]_D^{20}$ +9.6 (c, 0.1 in MeOH). λ_{max} 211 (log ϵ 4.35); 245 (log ϵ 4.21); 285 (log ϵ 3.91); 310 (log ϵ 3.95) (MeOH).

- Li, G.-Y. *et al.*, *Org. Lett.*, 2006, **8**, 3613-3615 (*isol, pmr, cmr, ms, cryst struct*)

Chaetomin C-335
Chetomin
 [1403-36-7]



Absolute Configuration

- $C_{31}H_{30}N_6O_6S_4$ 710.879
 Metab. of *Chaetomium cochliodes*, *Chaetomium globosum*, *Chaetomium seminudum*, *Chaetomium subglobosum* and *Chaetomium tenuissimum*. Antibacterial agent and immunosuppressant. Cytotoxic. Powder. Sol. Me_2CO , EtOAc, $CHCl_3$, C_6H_6 , Py, dioxan; fairly sol. MeOH, EtOH, Et_2O ; poorly sol. H_2O , hexane. Mp 218-220°. $[\alpha]_D^{25}$ +278 (c, 0.12 in $CHCl_3$). λ_{max} 276 (log ϵ 3.8); 287 (log ϵ 3.83); 295 (log ϵ 3.82) (EtOH).

- LD₅₀ (rat, orl) 75 mg/kg. FM3038000
 O,O'-Di-Ac: Mp 175-177°.

- N-Me, O,O'-di-Ac: Mp 112-114°.

Ring D trithia homologue: **Chetoseminudin A**

- $C_{31}H_{30}N_6O_6S_5$ 742.945
 Prod. by *Chaetomium seminudum* 72-S-204-1. Immunosuppressant. Amorph. solid. $[\alpha]_D^{24}$ +279 (c, 0.05 in $CHCl_3$). λ_{max} 277 (log ϵ 3.88); 285 (log ϵ 3.89); 295 (log ϵ 3.87) (EtOH).

Ring G trithia homologue: **Chetomin B**

- Chaetomin B**
 $C_{31}H_{30}N_6O_6S_5$ 742.945
 Prod. by the marine-derived *Chaetomium* sp. Gö 100/9. Amorph. solid. Mp > 200°. $[\alpha]_D^{20}$ +270 (c, 0.1 in $CHCl_3$). λ_{max} 205 (sh) (log ϵ 4.18); 290 (log ϵ 3.53) (MeOH).

- Safe, S. *et al.*, *J.C.S. Perkin 1*, 1972, 472 (*isol, uv, pmr, cd, struct*)

- McInnes, A.G. *et al.*, *J.A.C.S.*, 1976, **98**, 6741 (*cmr, nmr, struct*)

- Brewer, D. *et al.*, *J.C.S. Perkin 1*, 1978, 1248 (*nmr*)


- Jeu, W.C. *et al.*, *Can. J. Microbiol.*, 1983, **29**, 1399 (*props*)

- Fujimoto, H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 98-102; 526 (*isol, pmr, cmr, Chetoseminudin A*)

- Schlörke, O. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*marine, isol*)

- Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 601

Chaetominine C-336
 [918659-56-0]



Absolute Configuration

$C_{22}H_{18}N_4O_4$ 402.409
Prod. by *Chaetomium* sp. IFB-E015.
Cytotoxic. Cryst. (MeOH). Mp 161-163°.
[α]_D²⁰ -70 (c, 0.48 in MeOH). λ_{max} 205
(ϵ 38700); 225 (sh) (ϵ 28700) (MeOH).

Jiao, R.H. *et al.*, *Org. Lett.*, 2006, **8**, 5709-5712
(*isol, cd, pmr, cmr, ms, cryst struct*)
Snider, B.B. *et al.*, *Org. Lett.*, 2007, **9**, 4913-
4915 (*synth*)
Toumi, M. *et al.*, *Org. Lett.*, 2008, **10**, 5027-
5030 (*synth*)

Chairamidine C-337


$C_{22}H_{26}N_2O_4$ 382.458
Struct. unknown. Alkaloid from the bark of
Remijia purdieana (Rubiaceae).
Amorph. powder + 1H₂O. Mp 126-128°
dec. (anhyd.). [α]_D¹⁵ +7.3 (c, 3 in EtOH).
Hesse, O. *et al.*, *Annalen*, 1884, **225**, 211

Chairamine C-338

[1398-56-7]
 $C_{22}H_{26}N_2O_4$ 382.458
Struct. unknown. Alkaloid from *Remijia*
purdieana (Rubiaceae). Needles or prisms
+ 1H₂O (EtOH). Mp 140° (hydrate) Mp
233° (anhyd.). Resembles Mitragnanine,
M-657 and Corynantheine, C-689.
Hesse, O. *et al.*, *Annalen*, 1884, **225**, 211
Raymond-Hamet, M. *et al.*, *C. R. Hebd.*
Seances Acad. Sci., 1958, **247**, 1387-1389
(*uv*)

Chaksine C-339

[486-53-3]



$C_{22}H_{38}N_6O_4$ 450.58
Struct. revised in 1985 (formerly thought
to be C₁₁). Conts. 2 tautomeric amino-
imidazole systems. Alkaloid from the
seeds, leaves and roots of *Cassia absus*
(Fabaceae). Antiinflammatory agent.
Active against gram-positive bacteria.
▶ LD₅₀ (mus, scu) 80-120 mg/kg.
JG7526000

Hydrochloride (1:2):
Cryst. (MeOH). Mp 175°.


Dipicrate:

Cryst. (MeOH). Mp 236°.

Siddiqui, S. *et al.*, *Proc. - Indian Acad. Sci.*,
Sect. A, 1935, **2**, 421; *CA*, **30**, 1799 (*isol*)
Gupta, K.C. *et al.*, *Indian J. Med. Res.*, 1953,
41, 459-460 (*activity*)
Wiesner, K. *et al.*, *J.A.C.S.*, 1958, **80**, 1521 (*ir*)
Krishna Rao, R.V. *et al.*, *J. Nat. Prod.*, 1979,
42, 299 (*isol*)
Voelter, W. *et al.*, *Angew. Chem., Int. Ed.*, 1985,
24, 959 (*cryst struct, ir, abs config*)
Ahmad, V.U. *et al.*, *Pak. J. Pharm. Sci.*, 1988,
1, 5; *CA*, **109**, 129413k (*pmr, cmr*)

Chalciporone C-340

2-Methyl-7-(7-oxo-1,3-nonadienyl)-2H-
azepine
[112448-74-5]



$C_{16}H_{21}NO$ 243.348
Pungent principle from fruit-bodies of
the mushroom *Chalciporus piperatus*. Sl.
yellow oil. Sol. MeOH, EtOAc, Et₂O;
poorly sol. H₂O. [α]_D²² -452 (c, 1.3 in
Et₂O). Spontaneously isomerises to Iso-
chalciporone, e.g. in CDCl₃ soln. at r.t.
 λ_{max} 252 (ϵ 21000); 260 (sh); 270 (sh);
315 (sh) (EtOH) (Derep).

$\Delta^{2,3:4,5}$ -Isomer: **Isochalciporone**. 7-
Methyl-2-(7-oxo-1,3-nonadienyl)-3H-
azepine
[112448-72-3]


$C_{16}H_{21}NO$ 243.348
Constit. of *Chalciporus piperatus*. Yel-
low oil. Sol. MeOH, Et₂O, EtOAc;
poorly sol. H₂O. Nonpungent. λ_{max}
241 (ϵ 28000); 246 (sh); 292 (ϵ 14700);
328 (sh) (EtOH) (Derep).

$\Delta^{2,3:4,5}$ -Isomer, 12,13-didehydro: 7-
Methyl-2-(7-oxo-1,3,5-nonatrienyl)-
3H-azepine. **Dehydroisochalciporone**
 $C_{16}H_{19}NO$ 241.332
Nonpungent constit. of *Chalciporus*
piperatus. Intense yellow oil. Sol.
MeOH, Et₂O, EtOAc; poorly sol. H₂O.
 λ_{max} 226 (ϵ 21800); 285 (sh); 338 (ϵ
23000) (EtOH) (Derep).

Sterner, O. *et al.*, *Tetrahedron*, 1987, **43**, 1075
(*isol, uv, ir, pmr, cmr, ms, struct*)
Spiteller, P. *et al.*, *J.A.C.S.*, 2001, **123**, 4837-
4838 (*biosynth*)

Chalcostrobamine C-341

1-(3-Hydroxy-8-methyl-8-azabicy-
clo[3.2.1]oct-2-en-2-yl)-3-phenyl-2-pro-
pen-1-one, 9CI
[75638-72-1]




Absolute
Configuration

$C_{17}H_{19}NO_2$ 269.343
Enolised β -diketone. Alkaloid from the
leaves of *Knightsia strobilina* (Protaceae).
Viscous liq. [α]_D²⁰ +12 (CHCl₃) (natural).
[α]_D +165 (c, 1.1 in CHCl₃) (92%ee)
(synthetic).
Lounasmaa, M. *et al.*, *Phytochemistry*, 1980,
19, 953-955 (*uv, ir, pmr, ms, struct*)
Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 1983, **46**,
429 (*synth*)
Majewski, M. *et al.*, *J.O.C.*, 1995, **60**, 5825-
5830 (*synth, abs config*)

Chaloridone C-342

7-Methyl-1,3-dioxolo[4,5-a]furo[2,3-
c]acridin-12(7H)one, 9CI. N-Methyl-
2,3-methylenedioxyfuro[2,3-c]acridone
[114216-83-0]




$C_{17}H_{11}NO_4$ 293.278
Alkaloid from the roots of *Ruta chale-*
pensis. λ_{max} 224 (log ϵ 4.4); 240 (log ϵ
4.5); 270 (log ϵ 4.8); 320 (log ϵ 4.4); 387
(log ϵ 3.8) (MeOH).

Ulubelen, A. *et al.*, *Phytochemistry*, 1988, **27**,
650-651 (*isol, pmr, cmr, ms*)

Chamaedrone C-343

5,6,7,8-Tetrahydro-3,7-dimethoxy-2-
methyl-5-octyl-4(1H)-quinolinone
[954378-11-1]




Absolute
Configuration

$C_{20}H_{33}NO_3$ 335.486
Related to Antidesmone, A-1325. Alka-
loid from the roots of *Melochia chamae-*
drys. Yellowish viscous oil. [α]_D²⁵ -3.8 (c,
0.12 in CHCl₃).

Dias, G.O.C. *et al.*, *Planta Med.*, 2007, **73**, 289-
292 (*isol, pmr, cmr*)

Chamobtusine A C-344

[956377-80-3]




Relative
Configuration

$C_{20}H_{31}NO_2$ 317.47
Alkaloid from *Chamaecyparis obtusa* cv.
tetragon. Pale yellow cryst. (MeOH aq.).
Mp 148-150°. [α]_D²⁴ -220.1 (c, 0.24 in
MeOH). λ_{max} 248 (log ϵ 2.48); 322 (log ϵ
1.87) (MeOH).

Zhang, Y.-M. *et al.*, *Org. Lett.*, 2007, **9**, 4579-
4581 (*isol, pmr, cmr, ms, cryst struct*)

Chandrananimycin C C-345

[664355-11-7]



$C_{17}H_{16}N_2O_3$ 296.325

Prod. by a marine bacteria *Actinomadura* sp. M045 and *Halomonas* sp. GWS-BW-H8hM. Antibiotic. Orange solid. λ_{\max} 243 (log ϵ 4.49); 282 (log ϵ 4.19); 366 (log ϵ 4.16); 433 (log ϵ 4.2) (MeOH).

Maskey, R.P. et al., *J. Antibiot.*, 2003, **56**, 622-629 (*isol, ir, pmr, cmr*)

Bitzer, J. et al., *J. Antibiot.*, 2006, **59**, 86-92 (*isol, pmr, ms*)

Chandrine

C-346

$C_{25}H_{30}N_2O_8$ 486.521

Struct. unknown. Alkaloid from roots of *Rauwolfia serpentina* (Apocynaceae). Antiarrhythmic agent. Light-brown needles or plates (1,2-dichloroethane or butanol). Mp 230-231°. Strong greenish-blue fluor. in soln.

Hydrochloride: Mp 185°.

Nitrate: Mp 220-221°.

Picrate: Mp 180°.


Rakshit, B. et al., *Indian Pharm.*, 1954, **9**, 226-227; **10**, 84-88; *CA*, **49**, 4938; 10329a (*isol*)

MacLeod, D.P. et al., *Arch. Int. Pharmacodyn. Ther.*, 1962, **138**, 437-450; *CA*, **58**, 846c (*pharmacol*)

Channaine

C-347

[1355-47-1]



$C_{32}H_{38}N_2O_6$ 546.662

Alkaloid from *Sceletium tortuosum* and *Sceletium strictum* (Aizoaceae). Mp 179-180°. Opt. inactive. Prob. an artifact derived from dimerisation of normesembrenone (following racemisation).


Abou-Donia, A. et al., *Chem. Comm.*, 1978, 1078 (*isol, cryst struct, bibl*)

Jeffs, P.W. et al., *Alkaloids (Academic Press)*, 1981, **19**, 18 (*rev*)

Chanoclavine I

C-348

Chanoclavine



(-)-form

$C_{16}H_{20}N_2O$ 256.347

(-)-form [2390-99-0]

Alkaloid from *Claviceps purpurea* and from several spp. in the genera *Argyreaia*, *Stictocardia*, *Rivea* and *Ipomoea* (Convolvulaceae). Prisms (Me₂CO or MeOH). Mp 220-222° dec. $[\alpha]_D^{20}$ -240 (c, 1.0 in Py).

O- β -D-Fructofuranoside: *Chanoclavine I fructoside* [126432-33-5]

$C_{22}H_{30}N_2O_6$ 418.489

Alkaloid from submerged cultures of *Claviceps fusiformis* strain W1 supplemented with Chanoclavine I.

O- $[\beta$ -D-Fructofuranosyl-(2 \rightarrow 1)- β -D-fructofuranoside]: *Chanoclavine I fructosylfructoside* [126432-34-6]

$C_{28}H_{40}N_2O_{11}$ 580.631

Alkaloid from submerged cultures of *Claviceps fusiformis* strain W1 supplemented with Chanoclavine I.

N-Ac:

Prisms (MeOH). Mp 226-228°. $[\alpha]_D^{20}$ -80 (c, 0.5 in Py).

O,N-Di-Ac:

Needles + 1C₆H₆ (C₆H₆/petrol). Mp 174-175°. $[\alpha]_D^{20}$ -55 (c, 0.9 in Py).

N-De-Me: *Norchanoclavine I*

[51743-69-2]

$C_{15}H_{18}N_2O$ 242.32

Naturally occurring ergot alkaloid.

8R,9-Dihydro: *Dihydrochanoclavine I*

[29077-45-0]

$C_{16}H_{22}N_2O$ 258.363

Alkaloid from *Claviceps paspali* and from the sclerotia of two rye ergot strains.

8S,9-Dihydro: *Isodihydrochanoclavine I*

[29077-44-9]

$C_{16}H_{22}N_2O$ 258.363

Alkaloid from *Claviceps paspali* and from the sclerotia of two rye ergot strains.

18-Aldehyde: *Chanoclavine I aldehyde*

[25800-42-4]

$C_{16}H_{18}N_2O$ 254.331

Alkaloid from *Claviceps purpurea*. λ_{\max} 223 ; 275 ; 283 ; 293 (petrol/EtOH).

18-Carboxylic acid: *Chanoclavine I acid*

[25437-30-3]

$C_{16}H_{18}N_2O_2$ 270.33

Alkaloid from the seeds of *Ipomoea violacea* (Convolvulaceae). Cryst. (MeOH). Mp 245-247° dec.

(Z)-Isomer: *Isochanoclavine I*

[1150-44-3]

$C_{16}H_{20}N_2O$ 256.347

Alkaloid from *Claviceps purpurea* (Convolvulaceae). Cryst. (Me₂CO or 2-propanol). Mp 181°. $[\alpha]_D^{20}$ -216 (c, 0.50 in Py).

10-Epimer: *Chanoclavine II*

[1466-08-6]

$C_{16}H_{20}N_2O$ 256.347

Alkaloid from *Claviceps purpurea* and several *Argyreaia*, *Stictocardia*, *Rivea* and *Ipomoea* spp. (Convolvulaceae). Prisms (Me₂CO). Mp 174°. $[\alpha]_D^{20}$ -332

(c, 0.50 in Py).

10-Epimer: *hydrochloride*:

Needles (EtOH). Mp 247°. $[\alpha]_D^{20}$ -271 (c, 0.5 in 50% EtOH aq.).

10-Epimer, N-de-Me: *Norchanoclavine II*

[51704-40-6]

$C_{15}H_{18}N_2O$ 242.32

Trace alkaloid from *Claviceps* strain SD 58. Noncryst. λ_{\max} 223 ; 274 ; 283 ; 294 (MeOH).

(\pm)-form [57550-19-3]

Synthetic. Mp 185-186° (sealed tube) Mp 194-195° dec.

N-De-Me: Synthetic. Mp 182-183°.

18-Carboxylic acid: Synthetic. Powder (EtOH aq.). Mp 289-290° dec.

(Z)-Isomer: [76109-09-6]

Synthetic. Mp 162-167° Mp 200-201° dec.

10-Epimer: (\pm)-*Chanoclavine II*

[1150-43-2]

$C_{16}H_{20}N_2O$ 256.347

Alkaloid from *Claviceps purpurea*, several *Argyreaia* spp. and *Ipomoea violacea* (Convolvulaceae). Cryst. (CHCl₃ or Me₂CO). Mp 179°.

10-Epimer, N-de-Me: Synthetic. Mp 208-210° dec.

Hofmann, A. et al., *Helv. Chim. Acta*, 1957, **40**, 1358-1373 (*isol, uv, ir, struct*)

Stauffer, D. et al., *Helv. Chim. Acta*, 1964, **47**, 2186-2194 (*isol, ir, pmr, config, struct, Chanoclavine I, Chanoclavine II, Isochanoclavine I*)

Acklin, W. et al., *Chem. Comm.*, 1966, 799-800 (*stereochem, Isochanoclavine I*)

Voigt, R. et al., *Pharmazie*, 1970, **25**, 272 (*Dihydrochanoclavine I, Isodihydrochanoclavine I*)

Cassady, J.M. et al., *J. Nat. Prod.*, 1973, **36**, 390-396 (*Norchanoclavine II*)

Chao, J.-M. et al., *Phytochemistry*, 1973, **12**, 2435-2440 (*occur*)

Plieninger, H. et al., *Chem. Ber.*, 1976, **109**, 2140-2147 (*synth, pmr, ms*)

Floss, H.G. et al., *Tetrahedron*, 1976, **32**, 873-912 (*biosynth, rev, Norchanoclavine I*)

Choong, T.-C. et al., *Tet. Lett.*, 1977, 3137-3138 (*Chanoclavine I acid*)

Kozikowski, A.P. et al., *J.A.C.S.*, 1980, **102**, 4265-4267 (*synth*)

Natsume, M. et al., *Heterocycles*, 1981, **16**, 375-379 (*synth*)

Oppolzer, W. et al., *Tetrahedron*, 1983, **39**, 3695-3705 (*synth, Isochanoclavine I*)

Somei, M. et al., *Chem. Pharm. Bull.*, 1986, **34**, 948-950 (*synth*)

Kren, V. et al., *FEMS Microbiol. Lett.*, 1986, **37**, 31-34 (*Chanoclavine I aldehyde*)

Somei, M. et al., *Heterocycles*, 1987, **26**, 895-898; 1990, **31**, 1919-1921; 1997, **45**, 1263-1266 (*Chanoclavine I, Chanoclavine I acid, Chanoclavine II, Norchanoclavine II, synth*)

Kiguchi, T. et al., *Heterocycles*, 1989, **28**, 19-22 (*Isochanoclavine I, synth*)

Flieger, M. et al., *J. Nat. Prod.*, 1990, **53**, 171-175 (*fructofuranosides*)

Ninomiya, I. et al., *J.C.S. Perkin I*, 1991, 3275-3285 (*synth, Chanoclavine I, Isochanoclavine I*)

Kardos, N. et al., *Tetrahedron: Asymmetry*, 1994, **5**, 1525-1533 (*synth*)


Yokoyama, Y. et al., *Tet. Lett.*, 1996, **37**, 9309-9312 (*synth*)

Yamada, F. et al., *Heterocycles*, 2007, **72**, 599-620 (*synth*)

Chartellamide A

C-349

[111268-63-4]

Absolute
ConfigurationC₂₀H₁₂Br₄N₄O₂ 659.956

Alkaloid from the marine bryozoan *Chartella papyracea* and from *Flustra papyracea*. Sol. MeOH, EtOH; poorly sol. H₂O. λ_{max} 212 (log ε 4.28); 230 (log ε 4.41); 258 (log ε 4.03); 294 (log ε 3.76); 303 (log ε 3.74); 318 (log ε 3.53) (EtOH).

6-Bromo: Chartellamide B

[111268-64-5]

C₂₀H₁₁Br₃N₄O₂ 738.852


From *Chartella papyracea*. Sol. MeOH, EtOH; poorly sol. H₂O. λ_{max} 213 (log ε 3.7); 231 (log ε 3.81); 260 (log ε 3.46); 293 (log ε 3.19); 303 (log ε 3.15); 318 (log ε 2.85) (EtOH).

Anthoni, U. *et al.*, *J.O.C.*, 1987, **52**, 5638-5639 (*isol, uv, ir, cd, pmr, cmr, ms, struct*)

Chartelline A

C-350

[96845-55-5]

C₂₀H₁₃Br₄ClN₄O 680.418

Alkaloid from the marine bryozoan *Chartella papyracea* and from *Flustra papyracea*. Cryst. (EtOAc). Sol. MeOH, CHCl₃. Mp 214-216° dec. [α]_D²⁰ -421 (c, 0.038 in EtOH). Shows imidazole tautomerism in soln. λ_{max} 230 (ε 38900); 243 (ε 37200) (EtOH) (Derep). λ_{max} 230 (ε 38900) (MeOH) (Berdy).

17-Debromo: Chartelline B

[110271-21-1]

C₂₀H₁₄Br₃ClN₄O 601.521

Alkaloid from *Chartella papyracea* and *Flustra papyracea*. [α]_D²⁰ -339.1 (c, 0.023 in EtOH). λ_{max} 225 (ε 27500); 235 (ε 26900) (EtOH) (Derep).

17,18-Didebromo: Chartelline C

[110271-22-2]

C₂₀H₁₅Br₂ClN₄O 522.625

Alkaloid from *Chartella papyracea* and *Flustra papyracea*. [α]_D²⁰ -217.4 (c, 0.023 in EtOH). λ_{max} 227 (ε 25700); 234 (ε 25700) (EtOH) (Derep).

3-Dechloro, 3-methoxy: Methoxydechloro-chartelline A

[110271-23-3]

C₂₁H₁₆Br₄N₄O₂ 675.999

Alkaloidal artifact from *Chartella papyracea* and *Flustra papyracea*. [α]_D²⁰ -196.3 (c, 0.055 in EtOH). λ_{max} 230 (ε 31622); 245 (ε 25700) (EtOH) (Berdy).

Chevolot, L. *et al.*, *J.A.C.S.*, 1985, **107**, 4542-4543 (*isol, uv, ir, pmr, cmr, cryst struct*)
 Anthoni, U. *et al.*, *J.O.C.*, 1987, **52**, 4709-4712 (*derivs, isol*)
 Baran, P.S. *et al.*, *J.A.C.S.*, 2006, **128**, 14028-14029 (*Chartelline C, synth*)

Chatinine

C-351


[1390-84-7]

Struct. unknown. Alkaloid from the roots of *Valeriana officinalis*. Mp 211-212° (as hydrochloride) Mp 97-98° (as picrate).

Goris, A. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1921, **172**, 1059-1061 (*isol*)

Chelamidine

C-352

10-Hydroxyhomochelidonine

(+)-form

R¹ = R² = MeC₂₁H₂₃NO₆ 385.416**(+)-form [134177-17-6]**

Alkaloid from the roots of *Chelidonium majus* (Papaveraceae). Prisms (CHCl₃/EtOH). Mp 225-226°. [α]_D²¹ +123 (c, 0.50 in CHCl₃). λ_{max} 259 (log ε 3.3); 287 (log ε 3.9) (no solvent reported).

5-Deoxy: α-Homochelidonine. Homochelidonine

[476-33-5]

C₂₁H₂₃NO₅ 369.416

Alkaloid from the roots of *Chelidonium majus* (Papaveraceae). Pharmacol. said to resemble that of Chelidonine, C-357. Sol. MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. H₂O. Mp 169-170° Mp 182° Mp 192-193.5°. [α]_D +116 (CHCl₃).

(±)-form [102419-38-5]*5-Deoxy: [70191-85-4]*

Synthetic. Mp 170-171° Mp 192-193.5°.

Späth, E. *et al.*, *Ber.*, 1931, **64**, 1123-1127 (*Homochelidonine, struct*)
 Bersch, H.W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1958, **291**, 491-492 (*Homochelidonine, ir, config*)
 Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 3697-3704 (*Chelamidine*)
 Ninomiya, I. *et al.*, *Heterocycles*, 1977, **7**, 137-141 (*Homochelidonine, synth*)
 Hanaoka, M. *et al.*, *Tet. Lett.*, 1985, **26**, 5163-5166 (*Homochelidonine, synth, pmr*)
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)
 Yoshida, M. *et al.*, *Tet. Lett.*, 2002, **43**, 6751-6753 (*Homochelidonine, synth*)
 McManus, H.A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 433-436 (*Homochelidonine, synth*)

Chelamine

C-353

10-Hydroxychelidonine

As Chelamidine, C-352 with

R¹, R² = -CH₂-C₂₀H₁₉NO₆ 369.373**(+)-form [11028-61-8]**

Alkaloid from the roots of *Chelidonium majus* (Papaveraceae), also from *Sarcocapnos enneaphylla* and *Sarcocapnos saetabensis* (Papaveraceae). Prisms (EtOH). Mp 203-204°. [α]_D²⁰ +107 (c, 0.28 in CHCl₃).

O¹⁰.Me: 10-MethoxychelidonineC₂₁H₂₁NO₆ 383.4

This struct. was assigned to an alkaloid *isol.* from *C. majus* in 1924 but this has been shown to be a mixt. of Chelamine, Chelamidine, C-352 and Homochelidonine.

(±)-form [107651-17-2]

Synthetic. Mp 246-247°.

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 3697; 1977, **42**, 2686 (*isol, uv, ir, struct*)

Hanaoka, M. *et al.*, *Chem. Lett.*, 1986, 739 (*synth, pmr, stereochem*)

Tojo, E. *et al.*, *Phytochemistry*, 1991, **30**, 1005 (*isol*)


Blanco, O. *et al.*, *Phytochemistry*, 1991, **30**, 2071 (*isol*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1994, **59**, 429 (*Methoxychelidonine*)

Chelelactam

C-354

[112899-82-8]

C₂₅H₂₄N₂O₅ 432.475


Alkaloid from the stem bark of a *Zanthoxylum* sp. collected in Grand Cayman (*Zanthoxylum spinosum* or *Zanthoxylum coriaceum*) (Rutaceae).

Ng, K.M. *et al.*, *Phytochemistry*, 1987, **26**, 3251 (*isol, uv, ir, pmr, cmr, ms, struct*)

Chelerythridimerine

C-355

13,13'-(2-Oxotrimethylene)bis[12,13-dihydrochelerythrine], 8Cl. *1,3-Bis(8-hydrochelerythrinyl)acetone*. *1,3-Bis(6-hydrochelerythrinyl)acetone*. *1,3-Bis(11-hydrochelerythrinyl)acetone*. Alkaloid A† [23006-09-9]



C₄₅H₄₀N₂O₉ 752.819

Minor alkaloid from *Bocconia arborea*, also obt. by the base-catalysed condensation of Chelerythrine and acetonedicarboxylic acid (Papaveraceae). Plates (CHCl₃/Me₂CO). Mp 304°.

N-De-Me: Toddalidimerine. 1,3-(8-Hydrochelerythrinyl-8'-hydro-N-norchelerythrinyl)acetone

[81421-65-0]

C₄₄H₃₈N₂O₉ 738.792

Alkaloid from the roots of *Toddalia asiatica* (Rutaceae). Cryst. (CH₂Cl₂/Et₂O). Mp 307°. [α]_D²⁰ +60 (c, 2 in CHCl₃). Stereochem. and site of N-methylation unknown; may be the deriv. of a diastereoisomer of Chelerythridimerine.

MacLean, D.B. et al., *Can. J. Chem.*, 1969, **47**, 1951 (*ms, struct, synth*)

Sharma, P.N. et al., *Phytochemistry*, 1981, **20**, 2781 (*Toddalidimerine*)


Chelerythrine

C-356

1,2-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridinium(1+), 9CI.

7,8-Dimethoxy-10-methyl-2',3'-methylenedioxy-1,2-benzophenanthridinium(1+). Toddaline

[34316-15-9]



C₂₁H₁₈NO₄[⊕] 348.377

Several numbering systems have been used. Alkaloid from a wide variety of genera in the Papaveraceae (*Argemone*, *Bocconia*, *Chelidonium*, *Dicranostigma*, *Eschscholtzia*, *Glaucium*, *Hunnemannia*, *Hylomecon*, *Macleaya*, *Papaver*, *Platystemon*, *Sanguinaria*, *Stylomecon*, *Stylophorum*), Rutaceae (*Fagara*, *Toddalia*, *Zanthoxylum*), Papaveraceae (*Corydalis*, *Dicentra*), Sapindaceae (*Pteridophyllum*) and Hypecoaceae (*Hypecoum*). Potent cytotoxic agent functioning by DNA intercalation and uncoupling of oxidative phosphorylation. Inhibitor of rat liver aminotransferases. Antimicrobial and antiinflammatory agent recommended for use against oral infections. Temporary hypertensive agent in mice, rabbits and cats. Analgesic. Antineoplastic agent. Prolongs sleep. Antiparasitic, hepatotoxic. Log P 0.11 (uncertain value) (calc). λ_{max} 228 (ε 28200); 273 (ε 33900); 282 (ε 38900); 315 (sh) (ε 16200); 321 (ε 16600); 347 (sh) (ε 6200); 400 (ε 1400) (EtOH).

Chloride: [3895-92-9]

Cryst. + 1H₂O (dil. HCl). Mp 213-214° dec. (193°, 202-203°, 207°).

Picrate: Mp 236°.

N-De-Me: 1,2-Dimethoxy[1,3]benzodioxolo[5,6-c]phenanthridine, 9CI.

Norchelerythrine. O-Methyldecarine.

Des-N-methylchelerythrine

[6900-99-8]

C₂₀H₁₅NO₄ 333.343

Alkaloid from *Argemone albiflora*, *Argemone mexicana*, *Argemone polyanthemus*, *Chelidonium japonicum*, *Toddalia aculeata*, *Toddalia asiatica*, *Zanthoxylum ailanthoides*, *Zanthoxylum arnottianum*, *Zanthoxylum cuspidatum*, *Zanthoxylum nitidum*, *Zanthoxylum bungeanum* and *Zanthoxylum ovalifolium* (Papaveraceae, Rutaceae). Needles (CH₂Cl₂/hexane or CHCl₃/MeOH). Mp 220-221° (211-213°, 215-216°, 217-221°). λ_{max} 211 (log ε 4.3); 243 (log ε 4.57); 256 (log ε 4.54); 276 (log ε 4.67); 289 (sh) (log ε 4.51); 324 (log ε 4.18); 338 (sh) (log ε 4); 363 (sh) (log ε 3.56); 385 (log ε 3.49) (EtOH).

O⁹-De-Me: NK 109

[143201-31-4]

C₂₀H₁₆NO₄[⊕] 334.351

DNA topoisomerase II inhibitor for the treatment of drug resistant cancer. Orange solid + 2H₂O (as sulfate). Mp 256° dec. (sulfate dihydrate). CAS no. refers to sulfate.

O⁹-De-Me, N-de-Me: 2-Methoxy[1,3]benzodioxolo[5,6-c]phenanthridin-1-ol, 9CI. **Isodecarine**

[66855-60-5]

C₁₉H₁₃NO₄ 319.316

Alkaloid from the root bark of *Zanthoxylum integrifolium*. Cryst. (CHCl₃). Mp 234° (225-227°). λ_{max} 243 (log ε 4.54); 257 (sh) (log ε 4.53); 275 (log ε 4.59); 320 (sh) (log ε 4.01); 380 (sh) (log ε 3.35) (MeOH).

O¹⁰-De-Me: **Fagaridine.** *Isogagaridine*

[51059-64-4]

[149998-48-1]

C₂₀H₁₆NO₄[⊕] 334.351

Alkaloid from roots or root bark of *Fagara xanthoxyloides*, *Fagara tessmannii*, *Fagara macrophylla* and *Zanthoxylum nitidum* (Rutaceae). Inhibitor of DNA topoisomerase I. Yellow needles (as chloride). Mp 231-233° (chloride). Struct. of Fagaridine revised in 1998. Prev. assigned as the 9-hydroxy-10-methoxy isomer. Isogagaridine, reported in 1993, was actually Fagaridine assigned correctly for the first time. λ_{max} 228 (log ε 4.55); 284 (log ε 4.65); 322 (sh) (log ε 4.14) (EtOH).

O¹⁰-De-Me, N-de-Me: 1-Methoxy[1,3]benzodioxolo[5,6-c]phenanthridin-2-ol, 9CI. 9-Methoxy-10-hydroxy-2,3-methylenedioxybenzophenanthridine.

Decarine

[54354-62-0]

C₁₉H₁₃NO₄ 319.316

Alkaloid from the bark of *Zanthoxylum arnottianum* and *Zanthoxylum microcarpum*, and from the stem bark of *Zanthoxylum decaryi* and *Zanthoxylum viride* (Rutaceae). Orange needles (CHCl₃/MeOH or MeOH/HCl). Mp 248-251° (244-246°). λ_{max} 249 (log ε

4.54); 257 (log ε 4.55); 277 (log ε 4.67); 285 (sh) (log ε 4.52); 326 (log ε 4.2); 335 (sh) (log ε 4.11); 384 (log ε 3.46) (EtOH). λ_{max} 253 (log ε 4.31); 297 (log ε 4.37); 330 (log ε 4.06); 384 (log ε 3.46) (EtOH/NaOH).

O¹⁰-De-Me, N-de-Me, O-Ac: **Decarine acetate.** O-Acetyldecarine

[54354-63-1]

C₂₁H₁₅NO₅ 361.353

Alkaloid from stem bark of a *Zanthoxylum* spp. (*Zanthoxylum spinosum* or *Zanthoxylum coriaceum*). Prisms (petrol). Mp 210-212°. λ_{max} 239; 266; 273; 286; 320 (MeOH). λ_{max} 259; 314; 341 (MeOH/NaOH).

Di-O-de-Me: 1,2-Dihydroxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridinium(1+), 9CI. 7,8-Dihydroxy-2,3-methylenedioxy-5-methylbenzo[c]phenanthridinium(1+), 9,10-Demethylenesanguinarine

[70617-22-0]

C₁₉H₁₄NO₄[⊕] 320.324

Alkaloid from *Macleaya* sp. (Papaveraceae). Mp 226-227° Mp 217.5-220° (as trifluoroacetate). Artifact of the isoln. process of Sanguinarine, S-56. CAS no. refers to trifluoroacetate.

Di-O-de-Me, 9,10-quinone: 9,10-De-methylene-9,10-dehydrosanguinarine

[70617-24-2]

C₁₉H₁₂NO₄[⊕] 318.308

Isol. from *Macleaya* sp. (Papaveraceae). Cryst. (CHCl₃) (as trifluoroacetate). Mp 183° (trifluoroacetate). Artifact of the isoln. procedure. CAS no. refers to trifluoroacetate.

12-Hydroxy: **12-Hydroxychelerythrine**

[131984-75-3]

C₂₁H₁₈NO₅[⊕] 364.377

Alkaloid from *Eschscholtzia californica* (Papaveraceae). Orange cryst. (MeOH/HCl) (as chloride). Mp 244-246° (chloride).

8-Methoxy, N-de-Me: 1,2,13-Trimethoxy[1,3]benzodioxolo[5,6-c]phenanthridine. **8-Methoxynorchelerythrine**

C₂₁H₁₇NO₅ 363.369

Alkaloid from the roots of *Zanthoxylum nitidum*. Amorph. solid. Mp 213-224°.

8-Methoxy, O⁹-de-Me, N-de-Me: 2,13-Dimethoxy[1,3]benzodioxolo[5,6-c]phenanthridin-1-ol. **8-Methoxyisodecarine**

C₂₀H₁₅NO₅ 349.342

Alkaloid from the rhizomes of *Zanthoxylum nitidum*. Yellow powder. Mp 203-214°. λ_{max} 237 (log ε 4.4); 287 (log ε 4.49); 328 (sh) (log ε 4.35); 355 (log ε 3.92); 400 (log ε 3.07); 476 (log ε 3.12) (EtOH).

12-Methoxy: 1,2,4-Trimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridinium(1+), 9CI. **Chellitine**

[55950-32-8]

C₂₂H₂₀NO₅[⊕] 378.404

Alkaloid from a variety of genera in

the Papaveraceae (*Chelidonium*, *Eschscholtzia*, *Glaucium*, *Hunnemannia*, *Hylomecon*, *Macleania*, *Sanguinaria*) and Papaveraceae (*Dicentra*). Mp 197-198° (184-186°) (as chloride).

5-Ethoxy: 6-Ethoxy-1,2-dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridinium(1+), 9CI. 12-Ethoxy-7,8-dimethoxy-5-methyl-2,3-methylene-dioxybenzo[c]phenanthridinium(1+).

5-Ethoxychelerythrine. Alkaloid C† [79559-55-0]

C₂₃H₂₂NO₅[⊕] 392.43

Alkaloid from *Zanthoxylum nitidum* (Rutaceae). Inhibits the growth of Ehrlich ascites tumours in mice.

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1955, **20**, 21-26; 1960, **25**, 1667-1675; 1968, **33**, 1619-1623 (*isol*, *pmr*, *ms*, *Chelilutine*)

Bailey, A.S. et al., *J.C.S.*, 1956, 4535-4543 (*synth*, *uv*)

Govindachari, T.R. et al., *Indian J. Chem.*, 1967, **5**, 280 (*N-Norchelerythrine*)

Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 3420-3444 (*uv*, *Chelilutine*)

Šmula, V. et al., *Can. J. Chem.*, 1972, **50**, 1544-1547 (*N-Norchelerythrine*, *synth*)

Haisová, K. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 3312-3320 (*N-Norchelerythrine*)

Torto, F.G. et al., *Phytochemistry*, 1973, **12**, 2315-2317 (*uv*, *pmr*, *ms*, *ir*, *Fagaridine*)

Kessar, S.V. et al., *Indian J. Chem.*, 1974, **12**, 323 (*synth*)

Vaquette, J. et al., *Phytochemistry*, 1974, **13**, 1257-1259 (*Decarine*, *uv*, *ir*, *pmr*, *ms*)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1977, **25**, 3120-3121; 1978, **26**, 864-873 (*Chelilutine*, *synth*, *struct*)

Lasskaya, O.E. et al., *Khim. Prir. Soedin.*, 1978, **14**, 764-767; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 650-652 (*9,10-Demethylenesanguinarine*)

Adad-Mensah, I. et al., *Planta Med.*, 1979, **35**, 94 (*isol*, *Fagaridine*)

Wang, M.-H. et al., *Yaoxue Tongbao*, 1981, **16**, 48; *CA*, **95**, 192260r (*5-Ethoxychelerythrine*)

Šmidrkal, J. et al., *Coll. Czech. Chem. Comm.*, 1984, **49**, 1412-1420 (*synth*)

Krane, B.D. et al., *J. Nat. Prod.*, 1984, **47**, 1-43 (*rev*, *Isodecarine*)

Hanaoka, M. et al., *J.C.S. Perkin 1*, 1986, 2253-2256 (*synth*, *pmr*)

Ng, K.M. et al., *Phytochemistry*, 1987, **26**, 3251-3254 (*Decarine acetate*)

Hanaoka, M. et al., *Chem. Pharm. Bull.*, 1989, **37**, 857-858 (*Chelilutine*, *synth*)

Tanahashi, T. et al., *J. Nat. Prod.*, 1990, **53**, 579 (*12-Hydroxychelerythrine*)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2002-2006 (*synth*)

Martin, G. et al., *J.O.C.*, 1992, **57**, 5907-5911 (*synth*)

Fang, S.-D. et al., *J.O.C.*, 1993, **58**, 5025-5027 (*Isofagaridine*)

Cho, W.J. et al., *Arch. Pharmacol. Res.*, 1996, **19**, 240 (*synth*, *Isofagaridine*)

Kanzawa, F. et al., *Br. J. Cancer*, 1997, **76**, 571-581 (*NK 109*)

Harayama, T. et al., *Heterocycles*, 1998, **48**, 1989-1992; 2003, **59**, 293-301 (*Chelerythrine*, *Fagaridine*, *Decarine*, *synth*)

Nakanishi, T. et al., *J. Nat. Prod.*, 1998, **61**, 1263-1267; 1999, **62**, 864-867 (*Fagaridine*, *NK 109*, *synth*, *pmr*, *cmr*, *struct*)

Nakanishi, T. et al., *J.O.C.*, 1998, **63**, 4235-4239 (*NK 109*)

Marek, R. et al., *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Harayama, T. et al., *J.C.S. Perkin 1*, 2001, 523-528 (*N-Norchelerythrine*, *synth*, *bibl*)

Seckárová, P. et al., *Magn. Reson. Chem.*, 2002, **40**, 147-152 (*pmr*, *cmr*)

Martin, M.T. et al., *Fitoterapia*, 2005, **76**, 590-593 (*Norchelerythrine*, *Decarine*, *pmr*, *cmr*)

Chen, J.-J. et al., *Planta Med.*, 2005, **71**, 470-475 (*Isodecarine*)

Hu, J. et al., *Chem. Biodiversity*, 2006, **3**, 990-995 (*8-Methoxynorchelerythrine*)

Dvorák, Z. et al., *Heterocycles*, 2006, **68**, 2403-2422 (*rev*)


Styskala, J. et al., *Heterocycles*, 2007, **73**, 769-775 (*Isodecarine*, *synth*)

Cui, X.-G. et al., *Helv. Chim. Acta*, 2008, **91**, 155-158 (*8-Methoxyisodecarine*)

Chelidonine, 9CI

Stylophorine. *Diphylline*

C-357



(+)-form

C₂₀H₁₉NO₅ 353.374

Diphylline was the racemate. Shows mod. *in vivo* cytotoxic activity vs. P388 leukaemia and Walker carcinosarcoma cells. Strong spasmolytic agent, smooth muscle antispasmodic and relaxant, hypotensive, cholagogic and choleric agent. Has been used against cholecystitis and hepatitis and as bile duct antiseptic. Has also been used, together with α -alocryptopine in treating exanthemas in infants. *C. majus* is a major European folk remedy against papillomas and warts. Log P 0.68 (uncertain value) (calc).

► Toxic, a CNS depressant.

(+)-form [476-32-4]

Alkaloid from the roots of *Chelidonium majus* and *Stylophorum diphyllum*, and the aerial parts of *Glaucium fimbriigerum*. Chelidonine (with no stereochemical designation) has also been isol. from *Symphoricarpos albus*, *Dicranostigma franchetianum*, *Dicentra spectabilis*, *Hylomecon vernalis*, *Sarcocapnos enneaphylla* and *Glaucium vitellinum* (Papaveraceae, Caprifoliaceae). Mp 135-136°. [α]_D²³ +115 (c, 0.5 in EtOH). λ _{max} 238 (log ϵ 3.96); 289 (log ϵ 3.9) (MeOH). λ _{max} 206 (log ϵ 4.85) (EtOH).

► FL9450000

N-De-Me: (+)-Norchelidonine. N-De-methylchelidonine

C₁₉H₁₇NO₅ 339.347

Alkaloid from *Chelidonium majus* (Papaveraceae). Amorph. [α]_D +74 (c, 0.07 in CHCl₃). [α]_D +92.5 (c, 0.08 in MeOH). Incorrectly indexed in CA. λ _{max} 238 (log ϵ 3.95); 289 (log ϵ 3.92) (MeOH).

Ac: Mp 173-174° (165-166°). [α]_D²⁰ +110 (c, 0.93 in CHCl₃).

8-Oxo: Oxychelidonine

C₂₀H₁₇NO₆ 367.357

Alkaloid from *Chelidonium majus* (Papaveraceae). Needles (AcOH or CHCl₃). Mp 285°. [α]_D +102.5 (CHCl₃/EtOH).

14-Epimer: 14-Epichelidonine

[136173-03-0]

C₂₀H₁₉NO₅ 353.374

Alkaloid from *Sarcocapnos saetabensis* (Papaveraceae). Prisms (EtOH). Mp 196°. λ _{max} 209; 240; 292 (EtOH).

(-)-form

Alkaloid from the roots and above-ground parts of *Glaucium flavum*, and from the above-ground parts of *Glaucium corniculatum* (Papaveraceae). Prisms (EtOH aq.). Mp 135-136°. [α]_D²² -112 (c, 0.2 in EtOH) (c, 0.5 in Py).

N-De-Me: (-)-Norchelidonine

[6901-00-4]

C₁₉H₁₇NO₅ 339.347

Alkaloid from the roots and above-ground parts of *Glaucium flavum*, the roots of *Glaucium flavum* var. *vestitum*, and the above-ground parts of *Glaucium flavum* var. *fulvum* (Papaveraceae). Prisms (CHCl₃/EtOH). Mp 198-199°. [α]_D²² -112 (c, 0.40 in EtOH). [α]_D -100 (c, 0.50 in CHCl₃). λ _{max} 239 (log ϵ 3.86); 288 (log ϵ 3.91) (EtOH).

N-De-Me, N-Ac:

Cryst. (EtOH). Mp 195-196°.

N-De-Me, N,O-di-Ac: Mp 140-155° (efferv.). [α]_D²² -100 (c, 0.39 in CHCl₃).

(±)-form

Diphyllin†

[20267-87-2]

Alkaloid from the roots of *Stylophorum diphyllum* and *Chelidonium majus*, the aerial parts of *Glaucium corniculatum* ssp. *refractum*, and the above-ground parts of *Glaucium elegans* (Papaveraceae). Cryst. (EtOH or CHCl₃/EtOH). Mp 218-220° (215-216°). λ _{max} 236 (log ϵ 3.92); 288 (log ϵ 3.89) (94% EtOH).

N-De-Me: [33458-80-9]

Synthetic. Mp 212-217°. λ _{max} 236 (log ϵ 3.92); 288 (log ϵ 3.89) (94% EtOH).

Gadamer, J. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1924, **262**, 578-589 (*Oxychelidonine*, *isol*, *struct*)

Manske, R.H.F. et al., *Can. J. Res., Sect. B*, 1942, **20**, 53-56 (*isol*)

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1957, **22**, 279-285; 1959, **24**, 3601-3605; 1965, **30**, 3697-3704 (*Chelidonine*, *Norchelidonine*, *isol*, *uv*, *struct*)

Šantavý, F. et al., *Coll. Czech. Chem. Comm.*, 1960, **25**, 1344-1349 (*Norchelidonine*, *ir*, *struct*, *abs config*)

Snatzke, G. et al., *Tetrahedron*, 1970, **26**, 5013-5028 (*Norchelidonine*, *uv*, *cd*, *abs config*)

Battersby, A.R. et al., *J.C.S. Perkin 1*, 1975, 1147-1156 (*biosynth*)

Ghanbarpour, A. et al., *J. Nat. Prod.*, 1978, **41**, 472 (*isol*, *pmr*, *ms*)

Szauffer, M. et al., *Phytochemistry*, 1978, **17**, 1446-1447 (*isol*, *uv*, *pmr*, *ms*)

Takao, N. et al., *Tet. Lett.*, 1979, 495-496 (*cryst struct*, *abs config*)

Cushman, M. et al., *J.O.C.*, 1980, **45**, 5067-5073 (*synth*, *pmr*)


Oppolzer, W. et al., *Helv. Chim. Acta*, 1983, **66**, 1119-1128 (*Norchelidonine*, *synth*, *uv*, *ir*, *pmr*, *ms*)

Hanaoka, M. et al., *Chem. Lett.*, 1986, 739-742 (*synth*)

- Blanco, O. *et al.*, *Phytochemistry*, 1991, **30**, 2071-2074 (14-Epichelidonine)
 Kadan, G. *et al.*, *Planta Med.*, 1992, **58**, 477 (+)-Norchelidonine
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (N-15 nmr)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CDL000

Cheliensisamine C-358

2-Acetyl-2-amino-4-hydroxyindeno[1,2-b]oxet-3(2H)-one, 9CI
 [337363-34-5]




C₁₂H₉NO₄ 231.207

Alkaloid from the bark of *Goniotalamus cheliensis*.

Gu, Z. *et al.*, *CA*, 2001, **134**, 323467n (isol)

Cheliensisine C-359

[332378-20-8]




C₂₀H₂₀N₂O₅ 368.388

Alkaloid from *Goniotalamus cheliensis*.

Gu, Z. *et al.*, *CA*, 2001, **134**, 277887z (isol)

Chelirubine C-360

5-Methoxy-13-methyl[1,3]benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridinium(1+), 9CI. Bocconine. 12-Methoxysanguinarine
 [18203-11-7]



C₂₁H₁₆NO₅[⊕] 362.361

Several numbering systems have been used. Alkaloid from a wide variety of genera in the Papaveraceae (*Bocconia*, *Chelidonium*, *Dicranostigma*, *Eschscholtzia*, *Glaucium*, *Hunnemannia*, *Hylomecon*, *Hypecoum*, *Macleaya*, *Papaver*, *Platystemon*, *Sanguinaria*, *Stylophorum*) and the Papaveraceae (*Dicentra*). Purple needles (dil. HCl) (as chloride). Mp 299-302° dec. (282-283°) (chloride).

O-De-Me: 12-Hydroxysanguinarine
 [131984-74-2]

C₂₀H₁₄NO₅[⊕] 348.334

Alkaloid from *Eschscholtzia californica*

(Papaveraceae). Dark red cryst. (MeOH/HCl aq.) (as chloride). Mp 230-231° dec. (chloride).

5-Hydroxy: 5-Hydroxychelirubine. 12-Hydroxychelirubine
 [132023-26-8]

C₂₁H₁₆NO₆[⊕] 378.36

Alkaloid from *Eschscholtzia californica* (Papaveraceae). Pale red cryst. (MeOH/HCl) (as chloride). Mp 301-304° dec. (chloride). Turns colourless at ca. 200°.

5-Methoxy: 5,7-Dimethoxy-13-methyl[1,3]benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridinium(1+).

Macarpine

[23594-80-1]

C₂₂H₁₈NO₆[⊕] 392.387

Alkaloid from *Eschscholtzia douglasii*, *Eschscholtzia lobbii*, *Eschscholtzia glauca*, *Macleaya cordata*, *Macleaya microcarpa* and *Stylophorum diphyllum* (Papaveraceae). Cytotoxic. Red needles (dil. HCl) (as chloride). Mp 283-285° (chloride). Struct. revised in 1981.

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1955, **20**, 198; 356; 1961, **26**, 1839; 2933; 1963, **28**, 2530; 1965, **30**, 887; 1967, **32**, 4420; 1968, **33**, 1619; 1972, **37**, 2804; 1976, **41**, 2429 (*Chelirubine*, *Macarpine*, *isol*, *pmr*, *uv*, *struct*)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 864 (*struct*, *synth*, *pmr*)

Takao, N. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 473 (*synth*, *biosynth*)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 3335 (*synth*)

Tanahashi, T. *et al.*, *J. Nat. Prod.*, 1990, **53**, 579 (5-Hydroxychelirubine, 5-Hydroxysanguinarine)


Ishikawa, T. *et al.*, *Tetrahedron*, 1995, **51**, 8447 (*Macarpine*, *synth*)

Dostal, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1998, **63**, 1045-1055 (*pmr*, *ms*, *struct*)
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (N-15 nmr)

Seckárová, P. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 147-152 (*pmr*, *cmr*)

Chelonin A C-361

2-(3-Indolyl)-6-(3,4,5-trimethoxyphenyl)morpholine
 [133985-26-9]



Relative configuration

C₂₁H₂₄N₂O₄ 368.432


Alkaloid from the marine sponge *Chelonaplysilla* sp. Possesses anti-inflammatory and antibacterial activities. Cryst. (MeOH). Mp 182°. [α]_D -11.7 (c, 0.32 in CHCl₃). λ_{max} 213 (ε 37100); 272 (ε 6300); 278 (ε 6400); 288 (ε 5100) (MeOH) (Derep).

Bobzin, S.C. *et al.*, *J.O.C.*, 1991, **56**, 4403 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Somei, M. *et al.*, *Heterocycles*, 1995, **41**, 5 (*synth*)

Chelonin B C-362

3-Bromo-α-[[[2-(1H-indol-3-yl)ethyl]amino]methyl]-4-methoxybenzenemethanol, 9CI. 3-[2-[[2-(3-Bromo-4-methoxyphenyl)-2-hydroxyethyl]amino]ethyl]indole. N-(3-Bromo-β-hydroxy-4-methoxyphenethyl)tryptamine
 [133985-27-0]



C₁₉H₂₁BrN₂O₂ 389.291

λ_{max} 208 (ε 29000); 221 (ε 30600); 280 (ε 6400); 288 (sh) (MeOH) (Derep).

(S)-form

Alkaloid from the marine sponge *Chelonaplysilla* sp. Shows antibacterial activity. Solid. Mp 260° dec.

5-Bromo: **Bromochelonin B**

[134004-59-4]

C₁₉H₂₀Br₂N₂O₂ 468.187


From the sponge *Chelonaplysilla* sp. and *Dendrilla* sp. Shows antibacterial activity. Gum. [α]_D +3.7 (c, 0.27 in DMSO). λ_{max} 208 (ε 29000); 221 (ε 30600); 279 (sh); 287 (ε 2500); 299 (sh) (DMSO) (Derep).

Bobzin, S.C. *et al.*, *J.O.C.*, 1991, **56**, 4403-4407 (*isol*, *pmr*, *cmr*, *struct*)

Lawrence, N.J. *et al.*, *Tet. Lett.*, 2001, **42**, 7671-7674 (*synth*, *abs config*)

Chelonin C C-363

4-[6-(3,4-Dimethoxyphenyl)-2-morpholinyl]phenol, 9CI. 2-(4-Hydroxyphenyl)-6-(3,4-dimethoxyphenyl)morpholine
 [133985-28-1]



Relative configuration


C₁₈H₂₁NO₄ 315.368

Alkaloid from the marine sponge *Chelonaplysilla* sp. Solid. Mp 238° dec. [α]_D +5.8 (c, 0.53 in CHCl₃). λ_{max} 210 (ε 18200); 236 (ε 12900); 279 (ε 3800); 295 (sh) (MeOH/NaOH) (Derep). λ_{max} 208 (ε 15100); 227 (ε 13800); 277 (ε 3600) (MeOH) (Derep).

Bobzin, S.C. *et al.*, *J.O.C.*, 1991, **56**, 4403 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Chenoalbicin C-364


[848478-85-3]



C₃₆H₃₄N₂O₇ 606.674Alkaloid from the roots of *Chenopodium album*. Oil. Racemic. λ_{max} 196 (log ε 4.2); 227 (log ε 3.1); 283 (log ε 2.6) (MeOH).Cutillo, F. *et al.*, *Chem. Biodiversity*, 2004, **1**, 1579-1583 (*isol*, *pmr*, *cmr*)**Cheratamine**


C-365

[92664-88-5]

C₃₆H₃₄N₂O₇ 606.674Alkaloid from the stems of *Cocculus pendulus* (Menispermaceae). [α]_D²⁵ +190 (c, 0.33 in MeOH).Hussain, S.F. *et al.*, *Tetrahedron*, 1984, **40**, 2513 (*isol*, *uv*, *ir*, *pmr*, *ms*, *cd*, *struct*)**Cherianoine**

C-366

7-Hydroxy-6,8-dimethoxy-2-methyl-1(2H)-isoquinolinone

C₁₂H₁₃N₁O₄ 235.239Alkaloid from *Annona cherimola*. Needles. Mp 122-124°. λ_{max} 220 (log ε 4.51); 260 (log ε 4.41); 270 (log ε 3.35); 330 (log ε 3.39) (MeOH).Chen, C.-Y. *et al.*, *Phytochemistry*, 2001, **56**, 753-757**Cherimoline†**


C-367

[196958-70-0]

C₁₂H₇N₁O₂ 197.193Struct. originally assigned has been shown to be incorrect. Alkaloid from the stems of *Annona cherimola*. Powder. Mp 203-205°.Chen, C.-Y. *et al.*, *Tet. Lett.*, 1997, **38**, 6247-6248 (*isol*)Ajana, W. *et al.*, *Tetrahedron*, 1998, **54**, 4405-4412**Cherionaine**


C-368

[202589-82-0]

C₄₀H₄₄N₂O₁₁ 728.794Alkaloid from the stems of *Annona cherimola*. Amorph. powder. Opt. inactive (*meso*-compd.). λ_{max} 220 ; 290 ; 317 (no solvent reported).Chen, C.-Y. *et al.*, *Tet. Lett.*, 1998, **39**, 407-410 (*isol*, *pmr*, *cmr*, *ms*)**Cherylline**

C-369

1,2,3,4-Tetrahydro-4-(4-hydroxyphenyl)-6-methoxy-2-methyl-7-isoquinolinol, 9CI. Crinine†


C₁₇H₁₉N₁O₃ 285.342**(R)-form** [25515-34-8]

Synthetic. Mp 214-215°.


Di-Me ether:Cryst. (Et₂O/petrol). Mp 87-89°. [α]_D^{24.9} +21.58 (c, 1.01 in MeOH).**(S)-form** [23367-61-5]Alkaloid from *Crinum moorei* and *Crinum powellii* var. *alba* (Amaryllidaceae). Cryst. (Me₂CO). Mp 217-218° (213-214° dec.). [α]_D²² -89 (c, 0.60 in MeOH) (-69).*Hydrochloride*:Cryst. (Me₂CO). Mp 238-239°.*Di-Me ether*:Cryst. (Et₂O/petrol). Mp 87-88°. [α]_D^{24.9} -21.65 (c, 0.965 in MeOH).**(±)-form** [26996-80-5]Synthetic. Needles (CHCl₃ or CHCl₃/MeOH). Mp 216-218° (209-212°).*Hydrochloride*:Cryst. (EtOH/Et₂O). Mp 185° Mp 240-243° (double Mp).*Di-Me ether, hydrochloride*:Cryst. (MeOH/Et₂O). Mp 228-229°.Boit, H.-G. *et al.*, *Chem. Ber.*, 1954, **87**, 1704 (*isol*)Brossi, A. *et al.*, *J.O.C.*, 1970, **35**, 1100; 3559 (*isol*, *uv*, *ord*, *cd*, *struct*, *config*, *pmr*, *ms*)Schwartz, M.A. *et al.*, *J.O.C.*, 1971, **36**, 1827 (*synth*, *uv*, *ir*, *pmr*, *ms*)Kametani, T. *et al.*, *Tetrahedron*, 1975, **31**, 235 (*synth*, *ir*, *pmr*, *ms*)Hart, D.J. *et al.*, *J.A.C.S.*, 1978, **100**, 1548 (*synth*, *uv*, *ir*, *pmr*, *ms*)Irie, H. *et al.*, *Chem. Lett.*, 1980, 875 (*synth*)
Kessar, S.V. *et al.*, *Chem. Comm.*, 1981, 1074 (*synth*)Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1982, 2935 (*synth*, *pmr*)Hara, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3107 (*synth*, *ir*, *pmr*)Katakawa, J. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3928 (*synth*)Couture, A. *et al.*, *J.C.S. Perkin 1*, 1999, 789-794 (*synth*)Ruchirawat, S. *et al.*, *Heterocycles*, 2001, **55**, 635-640 (*synth*)Lebrun, S. *et al.*, *Org. Biomol. Chem.*, 2003, 1701-1706 (*synth*)**Chestnutamide**

C-370

11,11a-Dihydro-1H-pyrrolizino[3,2-b]quinoline-3,10(2H,5H)-dione, 9CI [359762-42-8]


C₁₄H₁₂N₂O₂Alkaloid from the flowers of *Castanea mollissima*.Wang, S. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 89-93 (*isol*)**Chetoseminudin C**

C-371

C₁₆H₁₉N₃O₃S₂ 365.476Prod. by *Chaetomium seminudum* 72-S-204-1. Pale yellow solid. λ_{max} 274 (log ε 3.72); 281 (log ε 3.74); 290 (log ε 3.7) (EtOH).*N*⁴-Me: **Chetoseminudin B**C₁₇H₂₁N₃O₃S₂ 379.503Prod. by *Chaetomium seminudum* 72-S-204-1. Pale yellow solid. [α]_D²⁴ +40 (c, 0.34 in CHCl₃). λ_{max} 274 (log ε 3.73); 281 (log ε 3.76); 290 (log ε 3.69) (EtOH).Fujimoto, H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 98-102; 526 (*Chetoseminudins*)**Chetracin A**

C-372

2,5:2',5'-Dide(epidithio)-2,5:2',5'-bis(e-pitetrathio)-6,6'-dihydroxychetocin, 9CI. Chaetracin A [99615-92-6]



Absolute configuration

C₃₀H₂₈N₆O₈S₈ 857.115

Diketopiperazine antibiotic. Prod. by *Chaetomium nigricolor* CBS-542.63/ ATCC 32394 (*Chaetomium abuense*) and *Chaetomium retardatum*. Mycotoxin. Shows strong cytotoxicity to HeLa cells. Active against gram-positive bacteria. Amorph. powder. Sol. MeOH, EtOAc; poorly sol. CHCl₃, H₂O, hexane. Mp 248–251° dec. $[\alpha]_D^{20} +723.5$ (c, 0.0012 in CHCl₃). λ_{\max} 242 (ε 21400); 306 (ε 5010) (EtOH) (Derep).


Tri-Ac:

Cryst. + 2H₂O. Mp 260° dec. $[\alpha]_D^{20} +830$ (c, 0.018 in CHCl₃).

Saito, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 1942–1956 (*isol, uv, ir, pmr, cmr, cd, cryst struct*)

Chicamycin A C-373

Antibiotic BBM 2040A. BBM 2040A [89675-37-6]

C₁₄H₁₈N₂O₅ 294.307

Anthramycin-type antibiotic. Isol. from *Streptomyces albus*. Possesses antitumour activity and weak activity against some gram-positive and acid-fast bacteria. Needles. Sol. MeOH, Py, butanol; fairly sol. EtOAc, H₂O, Me₂CO; poorly sol. CHCl₃, hexane. Mp 161–163° dec. $[\alpha]_D^{26} +350$ (c, 0.5 in Py). λ_{\max} 221 (ε 19200); 260 (sh) (ε 7900); 290 (sh) (ε 2800); 320 (ε 1200) (MeCN/HCl) (Derep). λ_{\max} 230 (ε 18000); 254 (sh) (ε 15100); 287 (ε 14000); 317 (ε 10100) (MeCN/NaOH) (Derep). λ_{\max} 223 (ε 23800); 232 (sh) (ε 21600); 258 (sh) (ε 7000); 322 (ε 3900) (MeCN) (Derep). λ_{\max} 223; 237; 262; 317 (MeOH) (Berdy). λ_{\max} 223 (ε 24900); 323 (ε 3900) (MeCN) (Berdy).

► LD₅₀ (mus, ipr) 28 mg/kg. UY8534500


Konishi, M. *et al.*, *J. Antibiot.*, 1984, **37**, 191; 200 (*isol, struct*)

Wong, H. *et al.*, *J. Antibiot.*, 1984, **37**, 300

Wong, H. *et al.*, *J. Med. Chem.*, 1985, **28**, 388 (*synth*)

Chicamycin B C-374

BBM 2040B. Antibiotic BBM 2040B [89675-39-8]

C₁₃H₁₄N₂O₄ 262.265

Anthramycin antibiotic. Isol. from *Streptomyces albus*. Active against tumours. Weakly inhibits gram-positive bacteria. Amorph. powder + 1H₂O. Sol.

MeOH, Py, butanol; fairly sol. Me₂CO, EtOAc, H₂O; poorly sol. CHCl₃, hexane. Mp 134–136° dec. $[\alpha]_D^{26} +552$ (c, 0.5 in Py). Chicamycin A is not a natural product but the MeOH adduct of Chicamycin B. λ_{\max} 222 (ε 16600); 260 (sh) (ε 7100); 290 (sh) (ε 2900); 323 (ε 1900) (CH₃CN/HCl) (Derep). λ_{\max} 234 (ε 17900); 253 (ε 17300); 288 (ε 12600); 318 (ε 11300) (CH₃CN/NaOH) (Derep). λ_{\max} 223 (ε 19400); 237 (ε 17800); 262 (ε 7400); 317 (ε 2900) (MeOH) (Derep). λ_{\max} 223; 237; 262; 317 (MeOH) (Berdy). λ_{\max} 232 (ε 20200); 318 (ε 3000) (MeCN) (Berdy).


► LD₅₀ (mus, ipr) 57 mg/kg. UY8545500

Konishi, M. *et al.*, *J. Antibiot.*, 1984, **37**, 191; 200 (*isol, uv, ir, pmr, cmr*)

Kaneko, T. *et al.*, *J. Antibiot.*, 1984, **37**, 300 (*synth*)

Chilenine C-375

12b-Hydroxy-9,10-dimethoxy-6H-1,3-dioxolo[4,5-h]isoindolo[1,2-b][3]benzazepine-8,13(5H,12bH)-dione, 9Cl. 1,1,4-Dioxo-2-hydroxy-7,8-(methylenedioxy)-12,13-dimethoxyaporphoeadane

C₂₀H₁₇NO₇ 383.357

(±)-form [71700-15-7]

Trace alkaloid from the stems and above-ground wood of *Berberis empetrifolia* (Berberidaceae). Prisms (MeOH). Mp 155° (114.5–116°).

Me ether:

Prisms (MeOH). Mp 147–148°.

Deoxy: 13-Deoxychilenine

[95456-41-0]

C₂₀H₁₇NO₆ 367.357

Alkaloid from stems and twigs of *Berberis actinacantha* (Berberidaceae). Amorph.

Moniot, J.L. *et al.*, *J.O.C.*, 1979, **44**, 4343 (*synth, ir, pmr*)

Fajardo, V. *et al.*, *Tet. Lett.*, 1982, **23**, 39 (*isol, uv, ir, pmr, ms*)

Dorn, C.R. *et al.*, *J.O.C.*, 1984, **49**, 2642 (*synth, ir, pmr, ms*)

Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957 (13-Deoxychilenine)

Fang, F.G. *et al.*, *Tet. Lett.*, 1989, **30**, 2747 (*synth*)

Kessar, S.V. *et al.*, *Indian J. Chem., Sect. B*, 1991, **30**, 299 (*synth, ir, pmr, ms*)

Ishibashi, H. *et al.*, *Tet. Lett.*, 1995, **36**, 6733 (*synth*)

Yoda, H. *et al.*, *Tet. Lett.*, 2002, **43**, 4667–4669 (*synth*)

Koseki, Y. *et al.*, *Heterocycles*, 2003, **59**, 527–540 (*synth*)


Honda, T. *et al.*, *Tet. Lett.*, 2005, **46**, 6823–6825 (*synth*)

Fuwa, H. *et al.*, *Heterocycles*, 2008, **76**, 521–539 (*synth*)

Kim, G. *et al.*, *Tet. Lett.*, 2008, **49**, 2391–2392 (*synth*)

Chillanamine C-376

[89412-84-0]


C₃₇H₄₂N₂O₇ 626.748

Alkaloid from *Berberis buxifolia* (Berberidaceae).

Leet, J.E. *et al.*, *J. Nat. Prod.*, 1983, **46**, 908–912 (*isol, uv, cd, pmr, ms, struct*)

Chilocorine A C-377

Chilocorine [154071-72-4]



Relative Configuration


C₂₆H₃₄N₂O 390.567

Alkaloid from the ladybird beetle *Chilocorus cacti*.

McCormick, K.D. *et al.*, *Tetrahedron*, 1994, **50**, 2365–2372 (*isol, uv, pmr, cmr, ms, struct*)

Chilocorine B C-378

[166990-22-3]



Relative Configuration


C₂₆H₃₄N₂O 390.567

Alkaloid from the coccinellid beetle *Chilocorus cacti*. Cryst. (CH₂Cl₂/MeOH, 1:1). Mp not reported.

Shi, X. *et al.*, *Tetrahedron*, 1995, **51**, 8711–8718 (*isol, pmr, cmr, ms, cryst struct*)

Chilocorine C C-379

[206443-44-9]

C₂₆H₃₆N₂O₂ 408.583


Config. shown is tentative. Alkaloid from the beetle *Chilocorus cacti*.

Huang, Q. *et al.*, *J. Nat. Prod.*, 1998, **61**, 598-601 (*isol, pmr, cmr, cryst struct*)

Chilocorine D

C-380

[479073-23-9]



Relative Configuration

C₂₆H₃₄N₂O₂ 406.567


Alkaloid from the coccinellid beetle *Chilocorus renipustulatus*. Oil. [α]_D²⁵ +43 (c, 0.14 in CH₂Cl₂). λ_{max} 204 (ε 15750) (no solvent reported).

Laurent, P. *et al.*, *Tet. Lett.*, 2002, **43**, 7465-7467 (*isol, pmr, cmr*)

Chiloenine

C-381

Methyl [6-[2-(dimethylamino)ethyl]-9-hydroxy-8-methoxy-1-oxonaphtho[1,2-c]furan-3(1H)-ylidene]acetate, 9CI [88176-95-8]

C₂₀H₂₁NO₆ 371.389

A secoisoquinoline alkaloid. Alkaloid from *Berberis buxifolia* (whole plant, no leaves). Minor constit. of *Berberis actinacantha* (branches without leaves) (Berberidaceae). Yellow-orange amorph. solid.

7,8-Dihydro: **Chiloenammine**

[88176-96-9]


C₂₀H₂₃NO₆ 373.405

Alkaloid from *Berberis actinacantha* and *Berberis buxifolia* (Berberidaceae). Amorph. yellow solid. Opt. inactive.

Shamma, M. *et al.*, *Chem. Comm.*, 1983, 799 (*uv, ir, pmr, ms, struct*)

Chimonanthine

C-382



(±)-form

C₂₂H₂₆N₄ 346.474

CAS numbering shown. Abs. configs. revised in 1999 and 2004.

(+)-form [85610-66-8]

Alkaloid from Colombian poison-dart frog *Phylllobates terribilis*. Also isol. from *Argostemma yappii*. Mp 171-172°. [α]_D²⁵ +280 (MeOH). The isolate named as Isochimonanthine has been shown to be a 1:1 mixt. of the (+)- and meso-forms of Chimonanthine.

(-)-form [5545-89-1]

Alkaloid from *Chimonanthus fragrans*, *Psychotria forsteriana* and *Calycanthus floridus* (Carolina allspice) (Calycanthaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 188-189°. [α]_D -329. λ_{max} 247 (ε 13600); 303 (ε 5600) (EtOH) (Berdy). λ_{max} 239 ; 294 (EtOH-HCl) (Berdy).

N-Me: Calycanthidine

[5516-85-8]

C₂₃H₂₈N₄ 360.501

Alkaloid from the seeds of *Calycanthus floridus* (Carolina allspice) (Calycanthaceae). Mp 142°. [α]_D²⁰ -317 (EtOH).

N⁸,N^{8'}-Di-Me: Folicanthine

[6879-55-6]

C₂₄H₃₀N₄ 374.528

Alkaloid from *Calycanthus floridus* (Carolina allspice) and *Calycanthus occidentalis* (Californian allspice) (Calycanthaceae). Mp 118-119°. Bp_{0.1} 150-170°. [α]_D -364 (EtOH).

N⁸,N^{8'}-Di-Me, picrate: Mp 179°.**N⁸,N^{8'}-Di-Me, N¹-de-Me: Chimonanthidine**C₂₃H₂₈N₄ 360.501

Alkaloid from the seeds of *Chimonanthus praecox*. Amorph. powder. [α]_D²⁰ -285 (c, 0.05 in EtOH). λ_{max} 208 ; 245 ; 300 (MeOH).

(±)-form [4147-36-8]

Synthetic. Mp 183-185°.

N⁸,N^{8'}-Di-Me: Synthetic. Mp 168-169°.**meso-form** [4147-37-9]

Alkaloid from *Argostemma yappii*, *Calycanthus floridus* (Carolina allspice) and *Psychotria forsteriana*. Mp 199-202° (176°). λ_{max} 248 (ε 13800); 305 (ε 4900) (EtOH).

N-De-Me: N-Demethyl-meso-chimonanthineC₂₁H₂₄N₄ 332.447

Alkaloid from *Psychotria lyciflora*. Amorph. powder. λ_{max} 245 (log ε 4); 303 (log ε 3.6) (EtOH).

Grant, I.J. *et al.*, *Proc. Chem. Soc., London*, 1962, 148 (*Folicanthine, cryst struct*)

Hino, T. *et al.*, *Tet. Lett.*, 1963, 1757-1760;

1978, 4913 (*Folicanthine, synth*)

Hendrikson, J.B. *et al.*, *Tetrahedron*, 1964, **20**,

565-579 (*synth, ms, pmr*)

Grant, I.J. *et al.*, *J.C.S.*, 1965, 5678-5696 (*cryst struct*)

O'Donovan, D.G. *et al.*, *J.C.S. (C)*, 1966,

1570-1572 (*biosynth*)

Hall, E.S. *et al.*, *Tetrahedron*, 1967, **23**,

4131-4141 (*meso-form, synth, isol, uv, ir, pmr, ms*)

Kirby, G.W. *et al.*, *J.C.S. (C)*, 1969, 1916-1919 (*biosynth*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1983, **39**, 41-47 ((±)-form, *isol*)

Adjibade, Y. *et al.*, *Phytochemistry*, 1992, **31**, 317-319 (*meso-form*)

Fang, C.L. *et al.*, *J.A.C.S.*, 1994, **116**, 9480-9486 (*Folicanthine, synth*)

Verotta, L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 392-396 (*isol, pmr, cd*)

Jannic, V. *et al.*, *J. Nat. Prod.*, 1999, **62**, 838-843 (*N-Demethyl-meso-chimonanthine*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1999, **121**, 7702-7703 (*synth, abs config*)

Takayama, H. *et al.*, *Tetrahedron*, 2004, **60**, 893-900 (*Calycanthidine, abs config*)


Matsuda, Y. *et al.*, *Heterocycles*, 2005, **65**, 1031-1033 (*Isochimonanthine*)

Movassaghi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **47**, 3725-3728 (*synth*)

Chinese bitter-sweet alkaloid I

C-383

Methyl tetrahydro-2-(1H-pyrrol-2-yl)-2H-1,3-oxazine-4-carboxylate



R = H

C₁₀H₁₄N₂O₃ 210.232

Alkaloid from the seeds of *Celastrus angulatus*. Oil. [α]_D²⁰ -0.12 (c, 0.45 in CHCl₃). λ_{max} 240 ; 280 (CHCl₃).

Yin, W.-P. *et al.*, *Phytochemistry*, 1999, **52**, 1731-1734

Chinese bitter-sweet alkaloid II

C-384

Methyl tetrahydro-5-(2-methylheptyl)-2-(1H-pyrrol-2-yl)-2H-1,3-oxazine-4-carboxylate

As Chinese bitter-sweet alkaloid I, C-383 with

R = -CH₂CH(CH₃)(CH₂)₄CH₃(ξ-)C₁₈H₃₀N₂O₃ 322.447


Alkaloid from the seeds of *Celastrus angulatus*. Yellowish liq. λ_{max} 240 ; 279 (CHCl₃).

Yin, W.-P. *et al.*, *Phytochemistry*, 1999, **52**, 1731-1734

Chinese bitter-sweet alkaloid III

C-385

7-Acetyloxy-2,9a-diethyloctahydro-6,9-dimethyl-7-propyl-2H-pyrido[1,2-a]pyrazine-1-carboxaldehyde [396068-94-3]



Relative Configuration


C₂₀H₃₆N₂O₃ 352.516

Alkaloid from the leaves of *Celastrus angulatus*. Yellow oil.

Yin, W.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 183-189 (*isol, pmr, cmr, ms*)

Chingazumianine

C-386

C₂₃H₂₃Cl₂NO₆ 480.343


Alkaloid from *Corydalis koidzumiana*.
Needles (CHCl₃/MeOH). Mp 195-196°. λ_{\max} 220 (log ϵ 4.57); 260 (log ϵ 4.33); 302 (log ϵ 4.2) (MeOH).

Chan, S.-C. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 2993-2999

Chingchengenamide A

C-387

7-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4-heptadienamide, 9CI. 7-(3,4-Methylenedioxyphenyl)-2,4-heptadienoic acid isobutylamide [139906-29-9]

C₁₈H₂₃NO₃ 301.385

Isol. from the rhizomes of *Asarum chingchengense* (Aristolochiaceae) and from leaves and stems of *Piper falconeri* (Piperaceae). Exhibits significant insecticidal activity. Needles (Me₂CO/petrol). Mp 126-128°.

5'-Methoxy: **Chingchengenamide B**. 7-(7-Methoxy-1,3-benzodioxol-5-yl)-N-(2-methylpropyl)-2,4-heptadienamide. 7-(3-Methoxy-4,5-methylenedioxyphenyl)-2,4-heptadienoic acid isobutylamide [139906-30-2]

C₁₉H₂₅NO₄ 331.411

Isol. from the rhizomes of *Asarum chingchengense* (Aristolochiaceae). Needles (Me₂CO/petrol). Mp 128.5-130°.

(Z,Z)-Isomer:

C₁₈H₂₃NO₃ 301.385

Alkaloid from *Piper hispidum*.
Amorph. solid. λ_{\max} 260 (MeOH).

Zhuhui, D. *et al.*, *Phytochemistry*, 1991, **30**, 3797 (isol, uv, ir, pmr, cmr, ms, struct)

Parmar, V.S. *et al.*, *Indian J. Chem., Sect. B*, 1993, **32**, 392 (isol, uv, ir, pmr, cmr, ms)

Navickiene, H.M.D. *et al.*, *Phytochemistry*, 2000, **55**, 621-626 (Z,Z-isomer)

Chinpeimine

C-388

C₂₇H₄₃NO₂ 413.642

Steroidal alkaloid. Struct. unknown. Alkaloid from the Chinese drug Chin-peimu. Mp 247-248°. $[\alpha]_D^{25}$ -21.58 (CHCl₃). λ_{\max} 288 nm.


Chloroaurate: Mp 156-158°.

Ac: Mp 178°.

Chu, T.-T. *et al.*, *CA*, 1951, **51**, 445

Chippiine

[98776-12-6]

C₂₃H₃₀N₂O₅ 414.5

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Rather unstable and dec. slowly to 3-Hydroxyconopharyngine in C-620. λ_{\max} 207; 228; 282; 301; 306; 310 (sh) (MeOH).

10,11-Bis-(demethoxy): **10,11-Demethoxy-chippiine**

[162229-96-1]

C₂₁H₂₆N₂O₃ 354.448

Minor alkaloid from bark of *Tabernaemontana markgrafiana* (Apocynaceae). Brown oil. $[\alpha]_D$ -26 (c, 0.02 in CHCl₃).

3-Epimer, 10-demethoxy, 19S-hydroxy:

Dippinine A

[223532-42-1]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Tabernaemontana corymbosa* leaves. Light yellow oil. $[\alpha]_D$ +19 (c, 0.67 in CHCl₃). λ_{\max} 231 (log ϵ 4.27); 280 (log ϵ 3.67); 300 (log ϵ 3.61) (EtOH).

3-Epimer, bis(demethoxy), 19-oxo: **Dippinine B**

[409063-45-2]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the stems of *Tabernaemontana corymbosa*. Light yellow oil. $[\alpha]_D$ -54 (c, 0.06 in CHCl₃). λ_{\max} 226 (log ϵ 4.18); 285 (log ϵ 3.65); 294 (log ϵ 3.61) (EtOH).

Van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400-423 (*Chippiine*)


Nielsen, H.B. *et al.*, *Phytochemistry*, 1994, **37**, 1729-1735 (*10,11-Demethoxychippiine*)

Kam, T.-S. *et al.*, *Heterocycles*, 2001, **55**, 2405-2412 (*Dippinines*)

Chiriquitoxin

C-390

11-(Aminocarboxymethyl)tetrodotoxin, 9CI [61132-15-8]

C₁₃H₂₀N₄O₁₀ 392.322

Toxin from skin and eggs of *Atelopus chiriquiensis*. Amorph. solid. Sol. H₂O. $[\alpha]_D^{25}$ -17.3 (c, 0.075 in AcOH aq.).


▶ FM6102600

Yotsu, M. *et al.*, *Tet. Lett.*, 1990, **31**, 3187-3190 (isol, pmr, cmr, struct)

Absolute Configuration

Chisitine 1

[771563-40-7]



R = MeSCO-

C₁₉H₂₇N₃O₃S 377.507

Alkaloid from the leaves of *Chisocheton weinlandii*. Mp 89-90°.

Tzouros, M. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1411-1425

Chisitine 2

C-392

[771563-41-8]

As Chisitine 1, C-391 with

R = PhCO-

C₂₄H₂₉N₃O₃ 407.511


Alkaloid from the leaves of *Chisocheton weinlandii*. Mp 102-103°.

Tzouros, M. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1411-1425

Chivosazole E

C-393

[195881-74-4]



Absolute Configuration

C₄₆H₆₅NO₁₂ 824.019

Macrolide antibiotic. Prod. by *Sorangium cellulosum*. Cytotoxic agent. $[\alpha]_D^{25}$ -153.2 (c, 0.4 in MeOH). λ_{\max} 265 (sh); 270 (log ϵ 4.98); 277 (sh); 332 (log ϵ 4.42) (MeOH).

3'-Me ether: **Chivosazole D**

[195881-73-3]

C₄₇H₆₇NO₁₂ 838.046

Prod. by *Sorangium cellulosum*. Cytotoxic agent. $[\alpha]_D^{25}$ -144.3 (c, 0.5 in MeOH). λ_{\max} 265 (sh); 270 (log ϵ 4.98); 277 (sh); 333 (log ϵ 4.42) (MeOH).

20-Me ether: **Chivosazole C**

[195881-72-2]

C₄₇H₆₇NO₁₂ 838.046

Prod. by *Sorangium cellulosum*. Cytotoxic agent. $[\alpha]_D^{25}$ -86.4 (c, 1 in MeOH). λ_{\max} 264 (sh); 270 (log ϵ 4.97); 276 (sh); 331 (log ϵ 4.39) (MeOH).

2',3'-Di-Me ether: **Chivosazole B**

[195881-71-1]



C₄₈H₆₉NO₁₂ 852.073

Prod. by *Sorangium cellulosum*. Cytotoxic agent. $[\alpha]_D^{25}$ -186.2 (c, 0.9 in MeOH). λ_{\max} 264 (sh); 270 (log ϵ 5.12); 276 (sh); 334 (log ϵ 4.53)

(MeOH).

2',3',20-Tri-Me ether: Chivosazole A
[169181-40-2]C₄₉H₇₁NO₁₂ 866.099Prod. by *Sorangium cellulosum*. Cytotoxic agent. Sol. Me₂CO, EtOAc, CH₂Cl₂, DMSO, MeOH; fairly sol. Et₂O; poorly sol. hexane, H₂O. [α]_D²² -113.6 (c, 0.7 in MeOH). λ_{max} 264 (sh); 270 (log ε 4.89); 276 (sh); 332 (log ε 4.3) (MeOH).**2',3',20-Tri-Me ether, 6E-isomer: Chivosazole A₁**
[195881-70-0]C₄₉H₇₁NO₁₂ 866.099Prod. by *Sorangium cellulosum*. Cytotoxic agent. [α]_D²² -158.1 (c, 0.4 in MeOH). λ_{max} 265 (sh); 271 (log ε 4.92); 277 (sh); 337 (log ε 4.49) (MeOH).**Aglycone, 20-Me ether: Chivosazole F**
[195881-75-5]C₄₁H₅₇NO₈ 691.903Prod. by *Sorangium cellulosum*. Cytotoxic agent. [α]_D²² -4.8 (c, 1 in MeOH). λ_{max} 270 (log ε 4.25); 277 (sh); 332 (log ε 4.25) (MeOH).Irschik, H. *et al.*, *J. Antibiot.*, 1995, **48**, 962-966 (*isol*)Jansen, R. *et al.*, *Liebigs Ann./Recl.*, 1997, 1725-1732 (*isol*, *uv*, *pmr*, *cmr*, *ms*)Janssen, D. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 4898-4901 (*abs config*)**Chloptosin**

[258843-09-3]

C₆₈H₉₄Cl₂N₁₈O₁₈ 1522.506Dimeric cyclohexapeptide antibiotic. Similar to Himastatin. Prod. by *Streptomyces* sp. MK498-98F14. Apoptosis inducer. Active against gram-positive bacteria incl. MRSA. Cryst. (EtOAc/Me₂CO). [α]_D²⁵ -16 (c, 0.25 in CHCl₃). Mp >260°. λ_{max} 213 (ε 98570); 256 (ε 23710); 299 (sh) (ε 7900) (MeOH).Umezawa, K. *et al.*, *J.O.C.*, 2000, **65**, 459-463 (*isol*, *ir*, *pmr*, *cmr*, *ms*)Hong, W.X. *et al.*, *Org. Lett.*, 2006, **8**, 4919-4922 (*partial synth*, *cryst struct*)**Chlorobiocin****C-395***Chlorobiocin. Antibiotic RP 18631. Antibiotic 2562A. RP 18631*
[39868-96-7]C₃₅H₃₇ClN₂O₁₁ 697.137

Coumermycin-type antibiotic. Prod. by

Streptomyces hygroscopicus, *Streptomyces roseochromogenus* var. *oscitans*, *Streptomyces albocinerescens* and *Streptomyces griseovariabilis*. Active against gram-positive and -negative bacteria. Mp 204-206°. [α]_D²⁰ -68 (c, 1.0 in EtOH). Log P 5.71 (calc). λ_{max} 275 (ε 30900); 307 (sh) (ε 14000); 337 (ε 30200) (CHCl₃) (Derrep).▶ LD₅₀ (mus, orl) 2200 mg/kg. UX9377000**5-Demethyl: Antibiotic 2562B**


[75207-21-5]

C₃₄H₃₅ClN₂O₁₁ 683.11From *Streptomyces griseovariabilis*.Active against gram-negative organisms. Sol. MeOH, CHCl₃, bases; poorly sol. H₂O, hexane, C₆H₆. Mp 218-220°. [α]_D¹⁸ -7.1 (c, 0.7 in EtOH). Log P 5.21 (calc). λ_{max} 225 (E1%/1cm 252); 263 (E1%/1cm 332); 308 (E1%/1cm 108) (EtOH) (Berdy).Ninet, L. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1972, **275**, 455 (*isol*)Dolak, J. *et al.*, *J. Antibiot.*, 1973, **26**, 121 (*struct*)Berger, J. *et al.*, *J. Chromatogr. Libr.*, 1978, 15 (*rev*)Gauze, G.F. *et al.*, *Antibiotiki (Moscow)*, 1979, **24**, 643 (*deriv*)Lysenkova, L.N. *et al.*, *Antibiotiki (Moscow)*, 1980, **25**, 483Sebek, O.K. *et al.*, *J. Antibiot.*, 1984, **37**, 136Eustáquio, A.S. *et al.*, *Chem. Biol.*, 2003, **10**, 279-288 (*biosynth*)**N-(4-Chlorobutyl)butanamide****C-396**

[329270-31-7]

H₃CCH₂CH₂CONHCH₂CH₂CH₂CH₂ClC₈H₁₆ClNO 177.673Alkaloid from *Aloe sabaea*. Oil.Blitzke, T. *et al.*, *Phytochemistry*, 2000, **55**, 979-982 (*isol*)D'hooghe, M. *et al.*, *Org. Prep. Proced. Int.*, 2003, **35**, 501-507 (*synth*)**3-Chloro-9H-carbazole, 9CI****C-397**

[2732-25-4]

C₁₂H₈ClN 201.655

Isol. from bovine urine. Silvery cryst. (EtOH or AcOH). Mp 201-202°.

Picrate: Mp 142-143°.*N-Ac*: [10336-16-0]C₁₄H₁₀ClNO 243.692

Needles (petrol) or plates (EtOH). Mp 124-125°.

N-Benzoyl: [189188-71-4]C₁₉H₁₂ClNO 305.763

Cryst. (EtOH). Mp 221-222°.

N-Me: [33268-90-5]C₁₃H₁₀ClN 215.681


Cryst. (hexane). Mp 40-42°.

N-Ph: [193686-61-2]C₁₈H₁₂ClN 277.752

Needles (EtOH). Mp 93-94°.

Ullmann, F. *et al.*, *Annalen*, 1904, **332**, 82-104 (*synth*)Buu-Hoi, N.P. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1954, **73**, 197-202 (*synth*)Matevosyan, R.O. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 3186-3193; *J. Gen. Chem. USSR (Engl. Transl.)*, 1960, **33**, 3155-3160 (*synth*)Lopatinskii, V.P. *et al.*, *CA*, 1966, **64**, 19542 (*synth, derivs*)Chen, H.J. *et al.*, *J.A.C.S.*, 1971, **93**, 5102 (*uv, nmr*)Akermark, B. *et al.*, *J.O.C.*, 1975, **40**, 1365 (*synth, uv*)Luk, K.-C. *et al.*, *J. Nat. Prod.*, 1983, **46**, 852-861 (*isol*)Dodsworth, D.J. *et al.*, *J. Het. Chem.*, 1988, **25**, 167-171 (*cmr*)Bonesi, S.M. *et al.*, *J. Het. Chem.*, 1997, **34**, 877-889; 891-900; 2004, **41**, 161-171 (*synth, pmr, cmr, ms, derivs*)Liu, Z. *et al.*, *Tetrahedron*, 2007, **63**, 347-355 (*synth*)**6-Chloro-β-carboline****C-398***6-Chloro-9H-pyrido[3,4-b]indole, 9CI. 6-Chloronorharman*


[30684-46-9]

C₁₁H₇ClN₂ 202.642Cryst. (CHCl₃). Mp 238-240° Mp 270-271°.

▶ Greater comutagenic activity with aromatic amines than that of β-Carboline, C-130.

N²-Me: 6-Chloro-2-methyl-β-carboline**Nostocarboline**C₁₂H₁₀ClN₂ 217.677Alkaloid from the freshwater cyanobacterium *Nostoc* 78-12A.

Potent cholinesterase inhibitor. Bright yellow needles (MeOH) (as iodide).

Mp > 215° (iodide). Counterion of natural compd. was not determined. λ_{max} 254 (log ε 3.68); 307 (log ε 3.46); 385 (log ε 2.81) (MeOH) (iodide).Ho, B.-T. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1445 (*synth*)Nakano, K. *et al.*, *Mutat. Res.*, 2000, **47**, 141-146 (*synth, comutagenicity, pmr, cmr*)Ponce, M.A. *et al.*, *J. Het. Chem.*, 2003, **40**, 419-426 (*synth, pmr, cmr, ms*)Becher, P.G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1793-1795 (*Nostocarboline*)**7-Chloro-β-carboline****C-399***7-Chloro-9H-pyrido[3,4-b]indole. 7-Chloronorharman*C₁₁H₇ClN₂ 202.642*N-Me: 7-Chloro-9-methyl-β-carboline.***Bauerine A**

[156312-09-3]


C₁₂H₉ClN₂ 216.669

Alkaloid from the terrestrial blue-green alga *Dichothrix baueriana*. Exhibits antiviral activity. Needles (hexane/Me₂CO). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 109–110°. λ_{max} 246 (ε 38300); 281 (ε 8200); 292 (ε 12300); 345 (ε 4500); 359 (ε 5500) (MeOH) (Berdy).

Larsen, L.K. et al., *J. Nat. Prod.*, 1994, **57**, 419 (isol, uv, pmr, cmr, struct)Dantale, S.W. et al., *Tetrahedron*, 2003, **59**, 5507–5514 (*Bauerine A*, synth)Pohl, B. et al., *Synth. Commun.*, 2007, **37**, 1273–1280 (*Bauerine A*, synth)

[2-Chloro-2-[4-chloro-5-hydroxy-3-methyl-3-(4-methyl-3-pentenyl)cyclohexylidene]ethyl]carbonimidic dichloride, 9CI C-400

6,14-Dichloro-5-hydroxy-3(14),9-axinysadien-15-yl carbonimidic dichloride [64789-88-4]


C₁₆H₂₃Cl₄NO 387.175

Isol. from the sponge *Pseudaxinysa pitys*. Oil. [α]_D²⁰ +36 (c, 1.1 in CHCl₃).

Wratten, S.J. et al., *J.A.C.S.*, 1977, **99**, 7367–7368 (isol, uv, pmr, cmr)

3-Chloro-11,12-dihydroindolo[2,3-a]carbazole, 9CI C-401

Tjipanazole I [139083-26-4]

C₁₈H₁₁ClN₂ 290.751

Minor alkaloid from the blue-green alga *Tolypothrix tjipanasensis*. λ_{max} 259 (ε 49600); 268 (ε 48500); 289 (ε 28400); 329 (ε 27200); 361 (ε 4430) (MeOH/CHCl₃) (Derep).

N¹¹-β-D-Xylopyranosyl: *Tjipanazole F1* [139083-23-1]

C₂₃H₁₉ClN₂O₄ 422.867

Minor alkaloid from *Tolypothrix tjipanasensis*. [α]_D²⁰ +14.9 (c, 1.0 in CHCl₃/MeOH 1:1). Opt. rotn. refers to a mixt. with F2. λ_{max} 258 (ε 46200); 271 (ε 43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).

N¹²-β-D-Xylopyranosyl: *Tjipanazole F2* [139083-24-2]

C₂₃H₁₉ClN₂O₄ 422.867

Minor alkaloid from *Tolypothrix tjipanasensis*. λ_{max} 258 (ε 46200); 271 (ε

43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).

N¹¹-α-L-Rhamnopyranosyl: *Tjipanazole C3*

[139083-17-3]

C₂₄H₂₁ClN₂O₄ 436.894

Minor constit. of *Tolypothrix tjipanasensis*. λ_{max} 258 (ε 46200); 271 (ε 43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).

N¹²-α-L-Rhamnopyranosyl: *Tjipanazole C4*

[139083-18-4]

C₂₄H₂₁ClN₂O₄ 436.894

Minor constit. of *Tolypothrix tjipanasensis*. λ_{max} 258 (ε 46200); 271 (ε 43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).

N¹¹-(6-Deoxy-β-D-gulopyranosyl):

Tjipanazole C1

[139083-15-1]

C₂₄H₂₁ClN₂O₄ 436.894

Minor alkaloid from blue-green alga *Tolypothrix tjipanasensis*. [α]_D²⁰ +18.1 (c, 1.1 in CHCl₃). Opt. rotn. refers to a mixt. with other *Tjipanazole C* components. λ_{max} 258 (ε 46200); 271 (ε 43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).

N¹²-(6-Deoxy-β-D-gulopyranosyl):

Tjipanazole C2

[139083-16-2]

C₂₄H₂₁ClN₂O₄ 436.894

Minor constit. of *Tolypothrix tjipanasensis*. λ_{max} 258 (ε 46200); 271 (ε 43300); 291 (ε 21200); 328 (ε 25700); 349 (ε 6810); 366 (ε 4690) (MeOH) (Derep).


Bonjouklian, R. et al., *Tetrahedron*, 1991, **47**,

7739 (isol, uv, pmr, struct)

Kuethe, J.T. et al., *Org. Lett.*, 2003, **5**, 3721–3723 (synth)

4-Chloro-5,6-dihydroxy-1H-indole C-402

4-Chloro-1H-indole-5,6-diol

C₈H₆ClNO₂ 183.594

N-Sulfonic acid: 4-Chloro-5,6-dihydroxy-1H-indole-1-sulfonic acid. *Ancorinolate C*

[473740-09-9]

C₈H₆ClNO₃S 263.658

Alkaloid from the sponge *Ancorina* sp. Powder (as Na salt). λ_{max} 216 (log ε 4.14); 272 (log ε 3.78); 302 (log ε 3.64) (MeOH) (Na salt).

N,O⁶-Disulfo: *Ancorinolate A*

[473740-07-7]

C₈H₆ClNO₈S₂ 343.722


Alkaloid from the sponge *Ancorina* sp.

Powder (as di-Na salt). λ_{max} 220 (log ε 4.31); 271 (log ε 3.86); 300 (log ε 3.56) (MeOH) (di-Na salt).

Meragelman, K.M. et al., *J.O.C.*, 2002, **67**, 6671–6677 (*Ancorinolates*)

2-Chloro-7-ethyl-6,8-dihydroxy-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, 9CI C-403

[208707-45-3]

C₁₃H₁₀ClNO₄ 279.679

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 245; 293; 336 (MeCN aq.).

6-Me ether: 2-Chloro-7-ethyl-8-hydroxy-6-methoxy-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, 9CI

C₁₄H₁₂ClNO₄ 293.706

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 248; 291; 341 (MeCN aq.).

1-Chloro: 1,2-Dichloro-7-ethyl-6,8-dihydroxy-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, 9CI

[208707-49-7]

C₁₃H₉Cl₂NO₄ 314.124


Prod. by *Streptomyces rimosus*. Powder. λ_{max} 245; 293; 336 (MeCN aq.).

Pat. Coop. Treaty (WIPO), 1998, 98 25 931; CA, **129**, 52003e

Trew, S.J. et al., *J. Antibiot.*, 2000, **53**, 1–11 (isol)

2-Chloro-7-ethyl-8-hydroxy-6-methyl-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, 9CI C-404

[208707-51-1]


C₁₄H₁₂ClNO₃ 277.706

Prod. by *Streptomyces rimosus*.

Pat. Coop. Treaty (WIPO), 1998, 98 25 931; CA, **129**, 52003e

Chlorofusin C-405

[329363-06-6]



Absolute Configuration

C₆₃H₉₉ClN₁₂O₁₉ 1363.997

Azaphilone-peptide antibiotic. Prod. by *Microdochium caespitosum*. p53-MDM2 antagonist. Yellow solid. λ_{\max} 202 (ε 28910); 346 (ε 9890); 402 (ε 12870) (MeOH).

Duncan, S.J. *et al.*, *J.A.C.S.*, 2001, **123**, 554-560 (*isol*, *pmr*, *cmr*)


Duncan, S.J. *et al.*, *Tet. Lett.*, 2002, **43**, 1075-1078 (*biosynth*)

Woon, E.C.Y. *et al.*, *J.O.C.*, 2007, **72**, 5146-5151 (*synth*)

Clark, R.C. *et al.*, *J.A.C.S.*, 2008, **130**, 12355-12369 (*synth*, *abs config*)

5-Chloro-2-hydroxy-6-methoxybenzoxazole C-406

5-Chloro-6-methoxy-2(3H)-benzoxazolone. 5-Chloro-6-methoxy-2-benzoxazolone




$C_8H_6ClNO_3$ 199.593

Isol. from *Zea mays*.

Kato-Noguchi, H. *et al.*, *Phytochemistry*, 1998, **49**, 433-435

(3-Chloro-4-hydroxyphenyl)acetic acid C-407

3-Chloro-4-hydroxybenzeneacetic acid, 9CI
[33697-81-3]



$C_8H_7ClO_3$ 186.594

Prod. by the fungi *Marasmius palmivorus* and *Xylaria* sp. FRR 5657.

Amorph. yellow solid. Mp 108-109°. λ_{\max} 219 (log ε 3.66); 282 (log ε 3.31) (MeOH). λ_{\max} 219 (log ε 3.68); 249 (log ε 3.72); 301 (log ε 3.47) (MeOH/NaOH).

Me ester: [57017-95-5]

$C_9H_9ClO_3$ 200.621

Bp_{0.15} 106-108°.

Amide: (3-Chloro-4-hydroxyphenyl)acetamide

$C_8H_8ClNO_2$ 185.609

Prod. by *Xylaria* sp. FRR 5657. Yellow cryst. Mp 160-162°. λ_{\max} 205 (log ε 4.08); 231 (sh) (log ε 3.59); 282 (log ε 3.18) (MeOH). λ_{\max} 206 (log ε 4.27); 248 (log ε 3.98); 299 (log ε 3.62) (MeOH/NaOH).

[94613-72-6, 52803-46-0]

Kuchar, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 1723-1735 (*synth*, *ester*)


D'Ambrosio, M. *et al.*, *Helv. Chim. Acta.*, 1984, **67**, 1484-1492 (*synth*)

Gribble, G.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1996, **68**, 201 (*occur*)

Davis, R.A. *et al.*, *Tet. Lett.*, 2005, **46**, 919-921 (*Xylaria* sp. FRR 5657 *metabs*)

3-Chloro-4-hydroxy-2-piperidinone C-408

[174204-83-2]




$C_5H_8ClNO_2$ 149.576

Alkaloid from *Piper hancei*.

Narui, T. *et al.*, *Nat. Med. (Tokyo)*, 1995, **49**, 438-441 (*isol*, *struct*)

3-Chloro-1H-indole, 9CI C-409

[16863-96-0]



C_8H_6ClN 151.595

Principal odorous component of the marine hemichordate *Ptychodera flava laysanica*. Mp 95.5-96° dec.

N-Benzoyl:

$C_{15}H_{10}ClNO$ 255.703

Prisms (EtOH). Mp 97-99°.


Pappalardo, G. *et al.*, *Gazz. Chim. Ital.*, 1958, **88**, 1147-1169 (*synth*, *uv*)

Higa, T. *et al.*, *Naturwissenschaften*, 1975, **62**, 395-396 (*isol*)

Brennan, M.P. *et al.*, *Heterocycles*, 1986, **24**, 2879-2885 (*synth*, *ir*, *pmr*, *cmr*)

4-Chloro-1H-indole-3-acetic acid C-410

[2519-61-1]



$C_{10}H_8ClNO_2$ 209.631

Auxin from the seeds of *Pisum sativum* (pea) and isol. from *Pinus sylvestris* (Scotch pine). Cryst. (1,2-dichloroethane/EtOH). Mp 179-180°.

Me ester: [19077-78-2]

$C_{11}H_{10}ClNO_2$ 223.658

Auxin from the seeds of *Pisum sativum* (pea). Cryst (EtOAc/hexane). Mp 120-121°.

Hansch, C. *et al.*, *J.A.C.S.*, 1951, **73**, 3518 (*synth*)

Marumo, S. *et al.*, *Nature (London)*, 1968, **219**, 959 (*isol*)

Magnus, V. *et al.*, *Phytochemistry*, 1997, **46**, 675-681 (*isol*)

Katayama, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 808-815 (*synth*, *ir*, *pmr*, *ms*)

Chloriodoacetic acid, 9CI C-411

[53715-09-6]

CICHICOOH

$C_2H_2ClIO_2$ 220.394

(+)-form

$[\alpha]_D +0.9$ (H₂O).

NH₄ salt: $[\alpha]_D +23.5$ (H₂O).

(±)-form

Constit. of *Asparagopsis taxiformis*.

Leaflets. V. sol. H₂O, mod. sol. petrol.

Mp 90°.

Phenyl ester:

$C_8H_6ClIO_2$ 296.491

Mp 110°.

Amide: Chloriodoacetamide

[62872-35-9]

C_2H_3ClINO 219.409

Constit. of *Asparagopsis taxiformis*.

Needles (H₂O or C₆H₆). Mp 140°.

Crompton, H. *et al.*, *J.C.S.*, 1923, **123**, 576-577 (*synth*)

McMath, A.M. *et al.*, *J.C.S.*, 1927, 537-543 (*synth*, *resoln*)

Seyferth, D. *et al.*, *J. Organomet. Chem.*, 1974, **71**, 335-346 (*synth*)


Woolard, F.X. *et al.*, *Tetrahedron*, 1976, **32**, 2843-2846 (*amide*, *isol*)

Woolard, F.X. *et al.*, *Phytochemistry*, 1979, **18**, 617-620 (*isol*)

5-Chloro-6-methoxy-1-methyl-1H-indole-2,3-dione, 9CI C-412

5-Chloro-6-methoxy-1-methylisatin

[16979-73-0]



$C_{10}H_8ClNO_3$ 225.631


Metab. of *Micromonospora carbonacea*. Deep red cryst. Mp 254-255°. λ_{\max} 269 (ε 27700); 274 (ε 26800); 316 (ε 6700); 425 (ε 900) (MeOH) (Berdy).

Reimann, H. *et al.*, *Chem. Ind. (London)*, 1967, 2173

Omote, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1969, **42**, 3016 (*synth*)

4-(Chloromethyl)-2,16-dihydro-16-(methoxycarbonyl)condyfolanium, 9CI C-413

[39020-70-7]



Absolute Configuration


$C_{21}H_{26}ClN_2O_2^{\oplus}$ 373.901

Doubtless an artifact. Alkaloid from *Craspidospermum verticillatum* (Apocynaceae). Mp 266-270° dec. (as chloride). $[\alpha]_D +507$ (c, 1.1 in EtOH) (chloride). CAS no. refers to chloride.

Besselièvre, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1972, 1477 (*uv*, *pmr*, *struct*)

4-Chloromethylene-1,2,3,4-tetrahydro-8-methyl-6-quinolinol C-414

4-Chloromethylene-1,2,3,4-tetrahydro-6-hydroxy-8-methylquinoline


C₁₁H₁₂ClNO 209.675**(E)-form**

O-(2,4-Di-O-methyl-β-D-xylopyranoside): [501373-67-7]

C₁₈H₂₄ClNO₅ 369.844Alkaloid from a Puerto Rican *Lyngbya majuscula*. Amorph. solid. [α]_D²² -20.8 (c, 0.12 in CHCl₃). λ_{max} 212 (log ε 4.58); 241 (log ε 4.29); 269 (log ε 3.9) (MeOH).Nogle, L.M. et al., *J. Nat. Prod.*, 2003, **66**, 217-220 (isol, pmr, cmr, ms)**2-Chloro-1,6-phenazinediol**, C-415

9CI

2-Chloro-1,6-dihydroxyphenazine [93768-40-2]

C₁₂H₇ClN₂O₂ 246.652Isol. from *Streptosporangium* sp. 1656. Shows a broad spectrum of antifungal activity *in vitro* against dermatophytes and *Candida* spp. Reddish-purple cryst. (CH₂Cl₂). Mp 252° dec. λ_{max} 265 (ε 23000); 296 (ε 65000) (MeOH/NaOH) (Derep). λ_{max} 275 (ε 71200); 377 (ε 4950); 440 (ε 3400) (MeOH) (Derep).

▶ SG1579700

Patel, M. et al., *J. Antibiot.*, 1984, **37**, 943 (isol, uv, ir, pmr, cmr)**6-Chloro-2-quinoxaline-carboxylic acid** C-416

[29821-63-4]

C₉H₅ClN₂O₂ 208.603

Cryst. (EtOH). Mp 223-224°.

Me ester: [29821-64-5]C₁₀H₇ClN₂O₂ 222.63Cryst. (CCl₄). Mp 147.5-148.5°.*Chloride*:C₉H₄Cl₂N₂O 227.049

Mp 103-103.5°.


1,4-Dioxide: **Antibiotic MSD 819**. MSD 819

[22587-15-1]


C₉H₅ClN₂O₄ 240.602Isol. from *Streptomyces ambofaciens*. Shows a broad spectrum of activity against gram-positive and -negative bacteria. Bright yellow cryst. Sol.DMSO, DMF; fairly sol. H₂O, MeOH; poorly sol. butanol, hexane. Dec. on heating, no def. Mp. λ_{max} 244 (ε 22900); 259 (ε 23300); 366 (ε 12200); 383 (ε 12400) (pH 7 phosphate buffer) (Derep). λ_{max} 244 (E1%/1cm 950); 259 (E1%/1cm 966); 366 (E1%/1cm 505); 383 (E1%/1cm 515) (pH 7 buffer) (Berdy).1,4-Dioxide, *Me ester*: [22587-18-4]C₁₀H₇ClN₂O₄ 254.629Yellow cryst. (CHCl₃/hexane). Mp 166-170°.Stapley, E.O. et al., *Antimicrob. Agents Chemother.*, 1968, 249 (isol)Miller, T.W. et al., *J. Antibiot.*, 1969, **22**, 293 (isol)Moreno, H.R. et al., *J. Med. Chem.*, 1970, **13**, 1005 (synth, uv)**Chlorpromazine**, BAN, INN, C-417

USAN

2-Chloro-N,N-dimethyl-10H-phenothiazine-10-propanamine, 9CI. 2-Chloro-10-(3-dimethylaminopropyl)phenothiazine, 8CI. Thorazine. Wintermin. Contomin. Chlordelazin. Fenactil. Aminazin. Many other names [50-53-3]

C₁₇H₁₉ClN₂S 318.869Claimed to be isol. from the Eastern folk medicine Mumiyo. Used in photometric detn. of Au(III). Tranquilliser, antiemetic. Used in treatment of schizophrenia. Oily liq. Insol. H₂O. Bp_{0.8} 200-205°. Log P 5.29 (calc).▶ Range of adverse effects from therapeutic usage reported: antimuscarinic effects; hypotension; extrapyramidal disorders; hypersensitivity reactions; reprod. effects; and skin reactions incl. phototoxic effects. Human and exp. teratogen. LD₅₀ (rat, orl) 142 mg/kg. Can cause contact dermatitis. SN8925000Khalikov, S.K. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2003, **39**, 22-25 (isol)**Choisyne** C-418

1,2-Dihydro-4,10-dimethoxy-α,α-dimethylidifuro[2,3-b:3',2'-f]quinoline-2-methanol, 9CI. Shuazine

C₁₈H₁₉NO₅ 329.352**(R)-form**7-Demethoxy: **Demethoxychoisyne**. *Demethoxychoisyne*C₁₇H₁₇NO₄ 299.326Alkaloid from the leaves of *Choisya ternata*. Cryst. (Me₂CO). Mp 184-186°. [α]_D -43.8 (c, 0.6 in CHCl₃) (64% ee).**(±)-form** [18556-07-5]Alkaloid from leaves of *Choisya ternata*. Needles (EtOH). Mp 189-190°.*Ac*: Mp 231-232°.Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1975 (isol, pmr, ms, struct)Grundon, M.F. et al., *J.C.S. Perkin 1*, 1974, 2181; 1975, 302 (biosynth)Boyd, D.R. et al., *Org. Biomol. Chem.*, 2007, **5**, 2983-2991 (abs config, *Demethoxychoisyne*)**Choline** C-419

2-Hydroxy-N,N,N-trimethylethanaminium(1+), 9CI. (2-Hydroxyethyl)trimethylammonium. Amanitin†. Arachine. Araquine. Bilinearine. Sincalin. Sinkalin [62-49-7]

Me₃N⁺CH₂CH₂OHC₅H₁₄NO⁺ 104.172

Occurs free and combined in many animal and vegetable prods., e.g. bile, brain, yolk of egg, hops, belladonna, strophanthus, and the seeds, leaves and stems of many other plants. Constit. of Lecithin. Acetylcholine precursor. Present free and as esters in the hypobranchial glands of molluscs. Lipotropic agent. Nutritional supplement.

▶ FZ9625000

Hydroxide: [123-41-1]C₅H₁₅NO₂ 121.179Syrup, cryst. with difficulty. V. sol. H₂O, EtOH; insol. Et₂O. Strong base, absorbs CO₂ from air. Forms cryst. addn. compds. with H₂PtCl₆ etc.

▶ GA4025500

Chloride: **Choline chloride**, INN. *Biocolina*. *Hepacholine*. *Lipotril* [67-48-1]C₅H₁₄ClNO 139.624

Lipotropic agent used to treat liver disorders. Deliquescent cryst.

▶ LD₅₀ (rat, orl) 3400 mg/kg. KH2975000*Iodide*: [17773-10-3]C₅H₁₄INO 231.076

Mp 272-274°.

Perchlorate: [17876-30-1]C₅H₁₄ClNO₅ 203.622

Cryst. (EtOH). Mp 273°.

Dihydrogen citrate: *Neurotropan*

[77-91-8] Lipotropic agent. Granules. Mp 105-107.5°.

▶ GA2582000

Theophylline salt (1:1): *Choline theophyllinate*, BAN, INN. *Cholinophylline*. **Oxytriphyllyne**, USAN. *Oxytrimethyllyne*. *Teperine* [4499-40-5]C₈H₂₀NO₆P 257.223

Smooth muscle relaxant, bronchodilator and diuretic. Granules.

▶ LD₅₀ (rat, orl) 600 mg/kg. GA7420000*Bitartrate*: [87-67-2]

Hygroscopic solid. Mp 151-153°.

- O-Nitrate, perchlorate: **Nitricholine perchlorate, INN**. Choline perchlorate nitrate [7009-91-8]
C₅H₁₃ClN₂O₇ 248.62
Cholinergic agent. Mp 185-186°.
- Sulfate (inner salt): N,N,N-Trimethyl-2-(sulfoxy)ethanaminium hydroxide inner salt, 9CI. Cholinesulfuric acid [4858-96-2]
C₅H₁₃NO₄S 183.228
Present in *Penicillium* spp., *Aspergillus sydowi*, *Porphyra umbilicalis*, *Ptilota pectinata* and *Dermatisicum thunbergii*. Cryst. Mp 300° dec.
- Salt with 2-[4-(4-chlorobenzoyl)phenoxy]-2-methylpropanoic acid (1:1): **Choline fenofibrate, INN**. ABT35 [856676-23-8] Antihyperlipidaemic agent. Cryst. Mp 209-211°.
- O-Ac: see Acetylcholine(1+), A-47
- O-Propanoyl: **Propionylcholine** [5072-54-8]
[2494-55-5, 2365-13-1]
C₈H₁₈NO₂[⊕] 160.236
Isol. from *Viscum album* (Viscaceae).
- O-Butanoyl: **Butyrylcholine** [3922-86-9]
[2963-78-2 (chloride), 2494-56-6 (iodide)]
C₉H₂₀NO₂[⊕] 174.263
Mp 108-111° (as chloride) Mp 93-94° (as iodide).
- O-(2-Methyl-2-propenyl): **Methacryloylcholine** [39316-06-8, 5039-78-1 (chloride)]
C₉H₁₈NO₂[⊕] 172.247
Monomer. Polymers are used as coagulants in sewage treatment.
- O-(3-Methyl-2-butenyl): **Seneciolycholine**. β,β-Dimethylacrylylcholine [20284-79-1]
C₁₀H₂₀NO₂[⊕] 186.274
Alkaloid from the hypobranchial body of the marine snail *Thais floridana*. Also isol. in salt form from the gastropod *Mancinella keineri*. Sol. H₂O. Mp 160° (as iodide).
- O-Pentanoyl: [2963-75-9 (iodide)]
C₁₀H₂₂NO₂[⊕] 188.289
Cryst. (as iodide). Mp 107-108° (iodide).
- O-Dodecanoyl: [25234-60-0 (chloride)]
C₁₇H₃₆NO₂[⊕] 286.477
Mp 196° (as chloride).
- O-Tetradecanoyl: [4277-89-8 (chloride)]
C₁₉H₄₀NO₂[⊕] 314.531
Mp 195° (as chloride).
- O-(2-Pentadecenyl): [116613-92-4]
C₂₀H₄₀NO₂[⊕] 326.542
Constit. of toxic skin secretions of the trunkfish *Anoplocapros lenticularis*. E- and Z-isomers detected. Poss. artifact derived from homologue of Pahutoxin, P-12. Genus name incorr. given as *Anoplocapros*.
- O-(2-Hexadecenyl):
C₂₁H₄₂NO₂[⊕] 340.568
Constit. of the toxic skin secretions of the trunkfish *Anoplocapros lenticularis*,

Aracana aurita, *Aracana ornata*, *Lactoria fornasini*, *Ostracion cubicus*, *Rhinesomas reipublicae*, *Strophuichthys inermis* and *Strophuichthys robustus*. E- and Z-isomers detected. Poss. artifact.

O-(2-Heptadecenyl):
C₂₂H₄₄NO₂[⊕] 354.595
Constit. of the toxic skin secretions of the trunkfish *Anoplocapros lenticularis*, *Aracana ornata*, *Lactoria fornasini*, *Ostracion cubicus* and *Strophuichthys robustus*. E- and Z-isomers detected. Poss. artifact.

O-(2-Octadecenyl):
C₂₃H₄₆NO₂[⊕] 368.622
Constit. of toxic skin secretions of the trunkfish *Anoplocapros lenticularis* and *Aracana ornata*. E- and Z-isomers detected. Poss. artifact.

O-Benzoyl:
[2964-09-2, 17518-43-3 (iodide)]
C₁₂H₁₈NO₂[⊕] 208.28
Cryst. (as iodide). Mp 247° (iodide). Chloride also known.

O-(4-Hydroxybenzoyl): **(4-Hydroxybenzoyl)choline** [5094-31-5]
C₁₂H₁₈NO₃[⊕] 224.279
Alkaloid from white mustard (*Sinapis alba*).

O-(3,4-Dimethoxybenzoyl): **Hesperaline** [15797-38-3]
[95210-18-7]
C₁₄H₂₂NO₄[⊕] 268.332
Alkaloid from seeds of damask violet (*Hesperis matronalis* Brassicaceae). Mp 182-183° (as iodide).

O-(4-Hydroxy-3,5-dimethoxybenzoyl): see Sinapine, S-308

O-(3-Hydroxy-4-methoxycinnamoyl): **Isoferuloylcholine** [28504-22-5]
C₁₅H₂₂NO₄[⊕] 280.343
Isol. from seeds of *Sibara virginica*. Mp 152-153° (as chloride). λ_{max} 218 (ε 14200); 230 (sh) (ε 100000); 300 (sh) (ε 132000); 325 (ε 16000) (H₂O).

O-(4-Hydroxy-3-methoxycinnamoyl): **Feruloylcholine** [85927-25-9]
[22513-68-4, 31545-00-3]
C₁₅H₂₂NO₄[⊕] 280.343
Constit. of garlic mustard (*Alliaria officinalis*) and *Cleome pungens*. Mp 203-205° (as chloride). λ_{max} 236 (log ε 4.05); 334 (log ε 4.31) (no solvent reported).

3-(4-Hydroxyphenyl)propanoyl: **3-(4-Hydroxyphenyl)propanoylcholine** C₁₄H₂₂NO₃[⊕] 252.333
Constit. of the flowers of *Polyscias murrayi*. Amorph. solid. λ_{max} 218 (sh) (log ε 3.52); 279 (log ε 2.9) (EtOH).

[55357-38-5, 23038-04-2, 64681-08-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 395D; 2, 292B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 616C; 859C (nmr)


Renshaw, R.R. et al., J.A.C.S., 1910, 32, 128 (synth)

- Fourneau, E. et al., Bull. Soc. Chim. Fr., 1914, 15, 548 (esters, synth)
- U.S. Pat., 1951, 2 575 611; CA, 46, 1722g (synth, Ferrocyclinate)
- Coates, G.E. et al., J.C.S., 1953, 2519
- Tammelin, L.E. et al., Acta Chem. Scand., 1956, 10, 145-146 (esters, synth)
- Parker, J.M. et al., J. Pharmacol. Exp. Ther., 1956, 118, 359 (pharmacol, Oxtriphyllyne)
- Cornacchione, G. et al., Boll. Chim. Farm., 1957, 96, 335 (Nitricholine perchlorate)
- Kyi, Z.-Y. et al., Br. J. Pharmacol., 1957, 12, 38 (iodide, esters)
- Roth, F.E. et al., J. Pharmacol. Exp. Ther., 1957, 121, 487 (pharmacol, Oxtriphyllyne)
- U.S. Pat., 1957, 2 776 287; CA, 51, 8150 (synth, Oxtriphyllyne)
- Whitaker, V.P. et al., Biochem. J., 1959, 71, 32-34 (3-methyl-2-butenoyl)
- Org. Synth., Coll. Vol., 4, 1963, 84 (O-benzoyl)
- Harper, S.H. et al., Chem. Ind. (London), 1966, 419 (sulfate)
- Gmelin, R. et al., Arch. Pharm. (Weinheim, Ger.), 1967, 300, 176 (Hesperaline)
- Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, 1, 142 (use)
- Bird, P.R. et al., J.C.S. (C), 1967, 447 (phosphate)
- Gmelin, R. et al., Phytochemistry, 1970, 9, 667 (isol, Isoferuloylcholine)
- Möhrle, H. et al., Arch. Pharm. (Weinheim, Ger.), 1973, 306, 489 (synth, Isoferuloylcholine)
- Byrn, S.R. et al., J.O.C., 1976, 41, 2283 (conformn, ir, cryst struct)
- Baker, J.T. et al., Tet. Lett., 1976, 1233-1234 (Choline, Seneciolycholine, occur, molluscs)
- Zeisel, S.H. et al., Annu. Rev. Nutr., 1981, 1, 195 (rev)
- Clausen, S. et al., Phytochemistry, 1982, 21, 917 (4-Hydroxybenzoylcholine)
- Larsen, L.M. et al., Phytochemistry, 1983, 22, 219-222 (Feruloylcholine)
- Stokr, J. et al., Coll. Czech. Chem. Comm., 1987, 52, 1256 (iodide)
- Negwer, M. et al., Organic-Chemical Drugs and their Synonyms, 6th edn., Akademie-Verlag, 1987, 368 (synonyms)
- Goldberg, A.S. et al., Toxicol., 1988, 26, 651-663 (alkenoyl derivs, occur)
- Kirk-Othmer Encycl. Chem. Technol., 4th edn., Wiley, 1991, 6, 199 (rev)
- Frydenvang, K. et al., Acta Cryst. C, 1992, 48, 1343; 1994, 50, 617 (cryst struct)
- Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1037; 1317
- Buchanan, M.S. et al., Phytochemistry, 2005, 66, 481-485 (3-4-Hydroxyphenylpropanoylcholine)
- U.S. Pat., 2005, 2005 148 594; CA, 143, 103273 (choline fenofibrate, synth, pharmacol)
- Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold, 1992, CMF000; CMF500; CMF750

Chondocurine

C-420

Tubocurine. Chondocurine

C₃₆H₃₈N₂O₆ 594.706

Diastereoisomeric with Curine, C-812.

(+)-form [477-58-7]

Alkaloid from the stem and bark of *Chondodendron tomentosum* and from the roots of *Cyclea madagascariensis* (Menispermaceae). Antiinflammatory props. Hepatoprotective in mice. Mp 232-234°. $[\alpha]_D^{24} +105$ (c, 0.9 in Py). $[\alpha]_D^{18} +173$ (c, 0.5 in CHCl_3). $[\alpha]_D^{24} +200$ (c, 0.5 in 0.1M HCl). Log P 7.58 (uncertain value) (calc).

N²-De-Me: Nor-N^b-chondocurine

[96738-71-5]

C₃₅H₃₆N₂O₆ 580.679

Alkaloid from Peruvian curare (Menispermaceae). Cryst. (EtOAc). Mp 159-161°. $[\alpha]_D^{20} -242$ (c, 0.5 in CHCl_3).

N²-Me: Tubocurarine. Tubarine

[57-95-4]

C₃₇H₄₁N₂O₆[⊕] 609.741

Quaternary alkaloid from Amazon curare (*Chondodendron tomentosum*) (Menispermaceae). The name Tubocurarine has been applied both to the C₃₇ monoquaternary salts and to the C₃₈ diquaternary (tetra-N-Me) salts.

N²-Me, chloride hydrochloride: Tubocurarine chloride, BAN, INN, JAN, USAN.

Amelizol. Many other names

[57-94-3]

[6989-98-6, 6533-76-2]

C₃₇H₄₂Cl₂N₂O₆ 681.654

Potassium (BK_{Ca}, SK_{Ca}) channel blocker. Acts as a nondepolarising muscle relaxant producing paralysis. Used mainly as an adjuvant to anaesthesia. Paralysis from therapeutic use is reversible. Overdose causes apnoea. Mp 274-275°. $[\alpha]_D^{22} +215$ (H₂O).

► LD₅₀ (rat, ims) 0.5 mg/kg. LD₅₀ (mus, ivn) 0.13 mg/kg. Very toxic by intramuscular, intravenous, and other routes of administration. Powerful curarising poison. YO4025000

N²-Me, iodide hydroiodide:C₃₇H₄₂I₂N₂O₆ 864.558Mp 263-265° dec. $[\alpha]_D^{23} +140$ (H₂O).**N²,N²-Di-Me: Chondocurarine. Chondrocurarine. Curarine**

[6880-94-0]

C₃₈H₄₄N₂O₆[⊕] 624.775

Quaternary alkaloid from the stem and bark *Cyclea tomentosum* (Menispermaceae).

N²,N²-Di-Me, dichloride:C₃₈H₄₄Cl₂N₂O₆ 695.681

Amorph. $[\alpha]_D^{23} +188$ (c, 1.08 in H₂O). $[\alpha]_D^{23} +195$ (c, 0.85 in MeOH).

N²,N²-Di-Me, diiodide:C₃₈H₄₄I₂N₂O₆ 878.584

Cryst. + 1CHCl₃ (MeOH/CHCl₃). Mp 277-280° dec. $[\alpha]_D^{23} +150$ (c, 0.71 in H₂O). $[\alpha]_D^{23} +170$ (c, 0.5 in MeOH).

O¹²-Me: [601468-47-7]C₃₇H₄₀N₂O₆ 608.733

Isol. from root of *Cissampelos mucronata*.

Di-Me ether, N²,N²-di-Me, diiodide: Metocurine iodide, USAN. Metubine iodide. Auxoperan. Curane B.

Diamethine. Mecostrin

[7601-55-0]

C₄₀H₄₈I₂N₂O₆ 906.638

Skeletal muscle relaxant. Mp 266°.

 $[\alpha]_D^{24} +160$ (H₂O).► LD₅₀ (rat, ipr) 0.37 mg/kg. YO3700000**Di-Me ether, N²,N²-di-Me, dichloride:****Dimethyltubocurarinum chloride. Di-****methyltubocurarine. Dimeberine chlor-****ide. Mecostrin chloride. Metocurine****chloride. Many other names**

[33335-58-9]

C₄₀H₄₈Cl₂N₂O₆ 723.734

Synthetic. Intramuscular relaxant, anticholinergic agent. Mp 236° dec. $[\alpha]_D^{25} +195$ (c, 0.5 in H₂O).

(-)-form [31503-97-6]

Alkaloid from the stems of *Chondodendron toxicoferum* (Menispermaceae) and roots of *Cissampelos mucronata*. Pure compd. could not be isol., phys. props. could not be determined.

N²-Me: (-)-Tubocurarine

[22154-22-9]

C₃₇H₄₁N₂O₆[⊕] 609.741

From *Chondodendron tomentosum* (Menispermaceae). Minute needles + 5H₂O (H₂O) (as chloride hydrochloride). Mp 275° dec. (chloride hydrochloride). $[\alpha]_D^{20} -258$ (c, 0.38 in H₂O) (chloride hydrochloride) (anhyd.). This appears to have been isolated only once and is the abnormal enantiomer. Much less powerful pharmacologically than the (+)-form.

O¹²-Me: Hayatidine

[16543-77-4]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the root and leaves of *Cissampelos pareira* (Menispermaceae). Mp 179-180°. $[\alpha]_D -109$ (Py).

[22260-42-0]

Dutcher, J.D. *et al.*, *J.A.C.S.*, 1946, **68**, 419-424; 1952, **74**, 2221-2225 (*isol*, Chondocurine, Chondocurarine)

Bick, I.R.C. *et al.*, *J.C.S.*, 1953, 3893-3896; 1961, 1896-1903 (*struct*, *pmr*)

Boissier, J.R. *et al.*, *J. Nat. Prod.*, 1965, **28**, 191 (*isol*)

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 2239-2247 (*ord*)

Bhatnagar, A.K. *et al.*, *Experientia*, 1967, **23**, 242-243 (*Hayatidine*)

Cava, M.P. *et al.*, *Phytochemistry*, 1969, **8**, 2341-2343 (*-)-Tubocurine*)

Everett, A.J. *et al.*, *Chem. Comm.*, 1970, 1020-1021 (*(+)-Tubocurarine*, *(+)-Chondocurine*, *struct*)

Baldas, J. *et al.*, *J.C.S. Perkin I*, 1972, 599-601 (*ms*)

Sobell, H.M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1972, **69**, 2212-2215 (*cryst struct*)

Hughes, R. *et al.*, *Br. J. Anaesth.*, 1976, **48**, 969-974 (*Metocurine, pharmacol*)

Koike, L. *et al.*, *J.O.C.*, 1981, **46**, 2385-2389 (*cmr*)

Lemli, J. *et al.*, *Planta Med.*, 1985, 68-69 (*Nor-N^b-chondocurine*)

Kondo, Y. *et al.*, *Int. J. Immunopharmacol.*, 1992, **14**, 1181-1186 (*pharmacol*)

Kondo, Y. *et al.*, *Biochem. Pharmacol.*, 1993, **48**, 1861-1863; 1887-1892 (*pharmacol*)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1322

Tshibangu, J.N. *et al.*, *Phytochem. Anal.*, 2003, **14**, 13-22 (*(+)-O¹²-Me, isol, hplc*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, COF825; DUM000

Chondriamide A

C-421

3-(1H-Indol-3-yl)-N-[2-(1H-indol-3-yl)ethenyl]-2-propenamide, 9CI

[142677-09-6]

C₂₁H₁₇N₃O 327.385

Alkaloid from the red alga *Chondria* sp. Cytotoxic. Shows antiviral activity. Anthelmintic agent. Yellow cryst. (MeOH aq.). Mp 193-194°. λ_{max} 208 (ε 27700); 226 (ε 31200); 278 (ε 12900); 364 (ε 26500) (MeOH) (Derep).

(10'Z)-Isomer: Chondriamide C

[218774-24-4]

C₂₁H₁₇N₃O 327.385

Alkaloid from *Chondria atropurpurea*. Anthelmintic agent. Yellow powder. Mp 230.5-232°. λ_{max} 208 (log ε 4.43); 275 (log ε 3.7); 358 (log ε 4.33) (MeOH).

7-Hydroxy: Chondriamide B

[142677-10-9]

C₂₁H₁₇N₃O₂ 343.384

From *Chondria* sp. Cytotoxic. Shows mild antifungal activity. Yellow cryst. (MeOH aq.). Mp 208-209°. λ_{max} 208 (ε 27700); 226 (ε 31200); 278 (ε 12900); 364 (ε 26500) (MeOH) (Derep). λ_{max} 206 (ε 26154); 222 (ε 25134); 274 (ε 10923); 368 (ε 21846) (MeOH) (Berdy).


Palermo, J.A. *et al.*, *Tet. Lett.*, 1992, **33**, 3097-3100 (*isol, uv, ir, pmr, cmr, ms, struct*, Chondriamide B)

Davyt, D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1560-1563 (*Chondriamide C, isol, pmr, cmr*)

Wang, X. *et al.*, *J.O.C.*, 2001, **66**, 8215-8221 (*synth*)

Chondrochloren A

C-422



Absolute Configuration

C₂₇H₄₀ClNO₇ 526.068

Prod. by *Chondromyces crocatus* strain Cm c5. Amorph. solid. $[\alpha]_D^{21} -82.4$ (c, 0.9 in MeOH). λ_{max} 205 (log ε 4.47); 224 (log ε 4.47); 278 (log ε 4.38) (MeOH).


2-O-De-Me, 2-Et ether: Chondrochloren BC₂₈H₄₂ClNO₇ 540.095

Prod. by *Chondromyces crocatus* strain Cm c5. Amorph. solid. $[\alpha]_D^{21} -81.7$ (c, 0.84 in MeOH). λ_{max} 204 (log ε 4.4); 224 (log ε 4.42); 278 (log ε 4.32) (MeOH).

Jansen, R. *et al.*, *Eur. J. Org. Chem.*, 2003, 2684-2689 (*isol, pmr, cmr, ms*)

Chrestifoline A C-423

1-Methoxy-4-[(1-methoxy-9H-carbazol-3-yl)methyl]-3-methyl-9H-carbazole, 9CI [129748-50-1]




C₂₈H₂₄N₂O₂ 420.51
Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)
Itoigawa, M. et al., *J. Nat. Prod.*, 2000, **63**, 893-897 (activity)

Chrestifoline B C-424

3,11-Dihydro-11-[(1-methoxy-9H-carbazol-3-yl)methyl]-3,3,5-trimethylpyrano[3,2-a]carbazole, 9CI [129748-51-2]




C₃₂H₂₈N₂O₂ 472.585
Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)

Chrestifoline C C-425

3,11-Dihydro-11-[(1-methoxy-9H-carbazol-3-yl)methyl]-3,5-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole, 9CI [129748-52-3]




C₃₇H₃₆N₂O₂ 540.704
Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil. [α]_D²⁵ -5.6 (c, 0.054 in CHCl₃).

Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)

Chromoviridan

[215675-75-5]



C₄₁H₂₈N₆O 620.712
Prod. by *Chromobacterium violaceum*. Green pigment (Me₂CO/hexane). Mp >230°. λ_{max} 338 (ε 19300); 388 (ε 16600); 490 (ε 18400); 562 (ε 27800); 644 (ε 53000) (MeOH aq./pH 2).

Deoxy: Deoxychromoviridan

[215675-77-7]


C₄₁H₂₈N₆ 604.712

Prod. by *Chromobacterium violaceum*. Green pigment (Me₂CO/hexane). Mp >230°. λ_{max} 489 (ε 13400); 641 (ε 59600) (MeOH aq./pH 2). λ_{max} 328 (ε 12610); 387 (ε 10580); 563 (ε 27600) (MeOH aq./pH 12).

Momen, A.Z.M.R. et al., *J.C.S. Perkin 1*, 1998, 3087-3092 (isol, cmr, pmr, ms, ir, uv, deoxy)

Chromoxymycin

FR 900447. Antibiotic 900447 [101156-09-6]



C₄₉H₆₀N₄O₁₄ 929.031

Anthraquinone antibiotic related to Hedamycin. Prod. by *Streptomyces rubropurpureus* and *Streptomyces libani-rubropurpureus*. Active against gram-positive bacteria, leukaemia and melanoma. Yellow needles + 4H₂O. Sol. MeOH; fairly sol. MeOH, Me₂CO; poorly sol. Et₂O, hexane, CHCl₃. Mp 85° dec. [α]_D²³ +291 (c, 0.675 in H₂O). λ_{max} 272 (ε 37100); 330 (sh) (ε 7500) (H₂O/HCl) (Derep). λ_{max} 243 (ε 25500); 278 (ε 32500); 410 (ε 2900) (H₂O/NaOH) (Derep). λ_{max} 276 (ε 36200); 340 (sh) (ε 7400) (H₂O) (Derep).

▶ LD₅₀ (mus, ivn) 1000 mg/kg. GB9140000
Kawai, Y. et al., *Tet. Lett.*, 1985, **26**, 3273

C-426

(struct, nmr, ir)

Hori, Y. et al., *J. Antibiot.*, 1986, **39**, 6; 12 (isol, uv, ir, pmr, cmr)

Chrycentrine

C-428

C₁₈H₁₅NO₅ 325.32


Struct. unknown. Alkaloid from *Dicentra chrysantha*. Mp 216°.

Manske, R.H.F. et al., *Can. J. Res., Sect. B*, 1937, **15**, 274-277; *CA*, **31**, 6663³

Chrysopentamine

C-429

[688796-87-4]



C₃₅H₃₈N₅O₂[⊕] 560.718

Related to Strychnopentamine, S-601.


Alkaloid from the leaves of *Strychnos usambarensis*. Antiplasmodial agent.

Amorph. orange powder (as chloride). λ_{max} 231 (log ε 3.37); 290 (log ε 2.92); 325 (log ε 2.67); 469 (log ε 3.12) (MeOH) (chloride).

Frédérich, M. et al., *Planta Med.*, 2004, **70**, 72-76 (isol, uv, cd, pmr, cmr)

Chrysophysarin A

C-430



C₂₁H₃₀N₃O₂[⊕] 356.487

Mesomeric imidazolium system.

(S)-form [262601-40-1]

Prod. by the slime mould *Physarum polycephalum*. Amorph. golden-yellow powder. λ_{max} 262 (log ε 0.4); 341 (log ε 1.28); 355 (log ε 1.55); 372 (log ε 1.15) (MeOH).


Eisenbarth, S. et al., *Tetrahedron*, 2000, **56**, 363-365

Chrysopine

C-431

9,10,11-Trihydroxy-2-oxo-1,7-dioxo-4-azaspiro[5.5]undecane-3-propanamide, 9CI


[177031-46-8]



C₁₁H₁₈N₂O₇ 290.272
Isol. from *Agrobacterium tumefaciens*.
Crown gall opine.

Vaudequin-Dransart, V. *et al.*, *CA*, 1995, **123**,
29164a

Chrysopterin C-432
2-Amino-1,5-dihydro-7-methyl-4,6-pteridin-
edione, 9CI. 7-Methylxanthopterin
[492-10-4]




C₇H₇N₅O₂ 193.165
Pigment of wings of *Gonepteryx rhamni*.
Yellow needles.

Schöpf, C. *et al.*, *Annalen*, 1933, **507**, 266;
1936, **524**, 49

Tschesche, R. *et al.*, *Chem. Ber.*, 1951, **84**, 641
Williams, V.P. *et al.*, *J. Het. Chem.*, 1973, **10**,
827 (ms)

Chrysopticine C-433
[200884-09-9]



C₂₁H₂₆N₂O₂ 338.449
Alkaloid from *Hedyotis chrysopticha*
(Rubiaceae). Yellow prisms (MeOH/
AcOH). Mp 160-161°. [α]_D²⁵ +26 (c, 0.05
in MeOH). λ_{max} 255; 309; 376 (MeOH).

N-De-Me (9H): **Cyclocapitelline**
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from *Hedyotis capitellata*.
Amorph. yellow solid. [α]_D²⁵ +43 (c, 0.5
in CHCl₃). λ_{max} 242 (log ε 4.3); 289
(log ε 4.07); 338 (log ε 3.59) (EtOH).

5'-Epimer: **Isochrysopticine**
[259821-82-4]

C₂₁H₂₆N₂O₂ 338.449
Alkaloid from *Hedyotis capitellata*.
Amorph. yellow solid (as monohy-
drochloride). [α]_D²⁵ -110 (c, 0.5 in
MeOH) (monohydrochloride). λ_{max}
210 (log ε 4.13); 250 (log ε 4.05); 311
(log ε 3.95) (EtOH) (monohydrochlor-
ide).

5'-Epimer, N-de-Me: **Isocyclocapitelline**
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from *Hedyotis capitellata*.
Yellow prisms (Me₂CO). Mp 199-200°.
[α]_D²⁵ -75 (c, 0.5 in CHCl₃). λ_{max} 242
(log ε 4.36); 290 (log ε 4.14); 339 (log ε
3.62) (EtOH).

Peng, J.-N. *et al.*, *Phytochemistry*, 1997, **46**,
1119-1121 (isol, uv, ir, pmr, cmr, ms)

Phuong, N.M. *et al.*, *Phytochemistry*, 1999, **52**,
1725-1729 (Isochrysopticine, Cyclocapitelline,
Isocyclocapitelline)


Mahboobi, S. *et al.*, *Monatsh. Chem.*, 2000,
131, 383-392 (synth, uv, pmr, cmr, cryst
struct)

Zhang, J.-X. *et al.*, *Tet. Lett.*, 2000, **41**, 2211-
2213 (synth)

Wang, G.X. *et al.*, *Chin. Chem. Lett.*, 2001, **12**,
199-202 (synth)

Volz, F. *et al.*, *Org. Biomol. Chem.*, 2007, **5**,
1519-1521 (Isochrysopticine,
Isocyclocapitelline, synth)

Chuanfumine C-434



C₂₂H₃₅NO₅ 393.522
Alkaloid from roots of *Aconitum carmi-
chaeli*.

Wei, X.-Y. *et al.*, *Zhiwu Xuebao (Acta Bot.
Sin.)*, 1990, **2**, 57

Yunusov, M.S. *et al.*, *Nat. Prod. Rep.*, 1993, **10**,
471 (rev)

Chuan-wu base A C-435
[1355-57-3]

C₂₃H₃₇NO₆ 423.548
Struct. unknown. Tertiary base with
2OH, 2OMe and one N-Et groups.
Alkaloid from the Chinese drug Chuan-
wu (*Aconitum carmichaeli*, Ranuncula-
ceae). Mp 111°. [α]_D²⁵ 0 (CHCl₃).

Ch'en, Y. *et al.*, *Yaoxue Xuebao*, 1965, **12**, 435-
439; *CA*, **63**, 16400c


Chuan-wu base B C-436
[1355-58-4]

C₃₂H₃₅NO₄ 497.633
Struct. unknown. Alkaloid from the
Chinese drug Chuan-wu (*Aconitum car-
michaeli*, Ranunculaceae). Mp 185°.
[α]_D^{25,3} 0 (CHCl₃).

Ch'en, Y. *et al.*, *Yaoxue Xuebao*, 1965, **12**, 435-
439; *CA*, **63**, 16400c

Cichorine C-437

2,3-Dihydro-6-hydroxy-4-methoxy-5-
methyl-1H-indol-1-one, 9CI. 6-Hydro-
xy-4-methoxy-5-methylphthalimidine
[114090-43-6]



C₁₀H₁₁NO₃ 193.202
Metab. of *Aspergillus silvaticus*. Also
from *Alternaria cichorii*. Phytotoxin.
Cryst. powder. Mp 217° subl. λ_{max} 212
(log ε 3.95); 252 (log ε 3.29); 295 (log ε
2.98) (EtOH).

O-(3-Methyl-2-butenyl): **Zinnimidine**
[148717-77-5]
C₁₅H₁₉NO₃ 261.32
Metab. of *Alternaria cichorii* and *Al-
ternaria porri*. Phytotoxin. Needles.
Mp 136-138°. λ_{max} 213 (ε 53700); 252
(ε 10964); 292 (ε 4677) (MeOH)
(Berdy).

O-(3-Methyl-2-butenyl), N-(2-hydro-
xyethyl): **Porritoxin**
[143114-82-3]
C₁₇H₂₃NO₄ 305.373
Isol. from the fungi *Alternaria porri*
and *Alternaria solani*. Phytotoxin. In-
hibits lettuce seedling growth at 10ppm
concentration. Needles. Mp 115-116°.
Struct. revised in 2002. λ_{max} 215 (ε
53700); 255 (ε 117500); 292 (ε 4000)
(MeOH) (Berdy).

O-(3-Methyl-2-butenyl), N-(2-sul-
foethyl): **Porritoxinsulfonic acid**
C₁₇H₂₃NO₇S 385.437
Isol. from *Alternaria porri*. Phytotoxin.
Amorph. solid. Mp 199.5-201.7°. λ_{max}
218 (log ε 4.3); 254 (log ε 3.8); 289 (log
ε 3.4) (MeOH).

O-(4-Hydroxy-3-methyl-2Z-butenyl): **Z-
Hydroxyzinnimidine**
C₁₅H₁₉NO₄ 277.319
Metab. of *Alternaria cichorii*. Solid
(CHCl₃/MeOH, 1:1). Mp not reported.

Kawahara, N. *et al.*, *Chem. Pharm. Bull.*, 1988,
36, 398-400 (Cichorine)

Suemitsu, R. *et al.*, *Phytochemistry*, 1992, **31**,
2325-2326; 1995, **38**, 495-497 (Porritoxin,
Zinnimidine)

Stierle, A. *et al.*, *Phytochemistry*, 1993, **32**,
1145-1149 (Cichorine, Zinnimidine, Z-
Hydroxyzinnimidine)


Horiuchi, M. *et al.*, *J. Nat. Prod.*, 2002, **65**,
1204-1205 (Porritoxin, pmr, cmr, struct)

Horiuchi, M. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 2003, **67**, 1580-1583
(Porritoxinsulfonic acid)

Cornella, I. *et al.*, *J.O.C.*, 2004, **69**, 2191-2193
(Porritoxin, synth)

Moreau, A. *et al.*, *J.O.C.*, 2006, **71**, 3303-3305
(Porritoxin, synth)

Ciliatamide A C-438
[1008161-75-8]



C₂₆H₃₉N₃O₃ 441.612
Constit. of *Aptios ciliata*. Yellow oil.
[α]_D²⁰ +40 (c, 0.05 in MeOH). λ_{max} 206 (ε
29000) (MeOH).

N¹-Deacyl, N¹-octanoyl: **Ciliatamide B**
[1008161-87-2]
C₂₄H₃₇N₃O₃ 415.575
Constit. of *Aptios ciliata*. Yellow oil.
[α]_D²⁰ +55 (c, 0.1 in MeOH). λ_{max} 206 (ε
15000) (MeOH).


Nakao, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 469-
472 (isol, pmr, cmr)

Lewis, J.A. *et al.*, *Org. Lett.*, 2008, **10**, 4545-
4548 (synth, config)

Ciliatamide C

C-439

[1008161-89-4]

C₂₅H₃₇N₃O₃ 427.586

Constit. of *Aaptos ciliata*. Yellow oil. $[\alpha]_D^{26} +74$ (c, 0.1 in MeOH). λ_{\max} 203 (ε 13600) (MeOH).


Nakao, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 469-472 (*isol*, *pmr*, *cmr*)

Lewis, J.A. *et al.*, *Org. Lett.*, 2008, **10**, 4545-4548 (*synth*, *config*)

Cimiciduphytine

C-440

[138842-99-6]



Absolute Configuration

C₃₈H₄₀N₄O₇ 664.757


Alkaloid from *Haplophyton cimicidum* (Apocynaceae). Small plates (EtOH). Mp 287-290° dec. λ_{\max} 223 (ε 42000); 260 (ε 16000); 290 (ε 3000) (EtOH).

Adesomoju, A.A. *et al.*, *Heterocycles*, 1991, **32**, 1461-1462 (*isol*, *uv*, *ir*, *ms*, *pmr*, *struct*)

Cimiciphytine

C-441

[68353-31-1]



Absolute Configuration

C₃₇H₄₂N₄O₇ 654.761

Alkaloid from *Haplophyton cimicidum*. Prod. in low yield by reduct. of Haplophytine, H-58 (Apocynaceae). Needles (EtOH). Mp 210-212°. $[\alpha]_D -146$ (c, 0.1 in CHCl₃). λ_{\max} 222 (ε 44000); 270 (ε 17400); 315 (ε 2300) (EtOH).

O¹¹-De-Me: Norcimiciphytine

[68353-32-2]

C₃₆H₄₀N₄O₇ 640.735

Alkaloid from *Haplophyton cimicidum* (Apocynaceae). Mp 240° dec. $[\alpha]_D -115$ (c, 0.086 in CHCl₃). λ_{\max} 222 (ε 26750); 265 (ε 9604) (EtOH).

O¹²-De-Me: Crooksine

[135574-52-6]

C₃₆H₄₀N₄O₇ 640.735

Alkaloid from *Haplophyton crooksii* (Apocynaceae). Possesses antiacetyl-

cholinesterase activity. Mp 335° dec.


Lakshmiathan, M.V. *et al.*, *Heterocycles*, 1978, **9**, 1009-1013 (*Cimiciphytine*, *Norcimiciphytine*)

Mroue, M. *et al.*, *Phytochemistry*, 1991, **30**, 1741-1744 (*Crooksine*)

Cimilophytine

C-442

[86527-32-4]

C₃₈H₄₂N₄O₈ 682.772

Alkaloid from *Haplophyton cimicidum*. Cryst. (CHCl₃/EtOH). Mp 325° dec. $[\alpha]_D^{20} -84.9$ (EtOH). λ_{\max} 228 (32800); 266 (15900); 300 (4000) (EtOH).

Adesomoju, A.A. *et al.*, *J.O.C.*, 1983, **48**, 3015-3017 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)


Cinachyramine

C-443

2,5,6,8,9,9b-Hexahydro-2-methyl-3H,4H,7H-1,2,3a,6a-tetraazaphenylene, 9CI

[879490-03-6]

[879490-05-8 (trifluoroacetate)]

C₁₀H₁₈N₄ 194.279


Constit. of *Cinachyrella* sp. Oil (as trifluoroacetate salt).

Shimogawa, H. *et al.*, *Tet. Lett.*, 2006, **47**, 1409-1411 (*isol*, *cd*, *pmr*, *cmr*)

Cinchonamine

C-444

[482-28-0]



Absolute Configuration

C₁₉H₂₄N₂O 296.411

Alkaloid from *Remijia purdieana* and *Cinchona pubescens* (Rubiaceae). Shows no antimalarial props. Mp 185-186°. $[\alpha]_D^{25} +128$ (c, 0.3 in EtOH).

Nitrate: Mp 227° dec.

Methiodide: Mp 208-209°.

Aldehyde: Cinchonaminal

[29560-34-7]

C₁₉H₂₂N₂O 294.396

Prob. intermed. in biosynth. of cinchona alkaloids. Not yet characterised as

a nat. prod.

10-Methoxy: 10-Methoxycinchonamine

[83852-64-6]

C₂₀H₂₆N₂O₂ 326.438

Minor alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae).

Arnaud, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1881, **93**, 593; 1883, **97**, 174 (*isol*)

Goutarel, R. *et al.*, *Helv. Chim. Acta*, 1950, **33**, 150 (*ir*, *uv*, *struct*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1959, **81**, 1474 (*config*)

Sawa, Y.K. *et al.*, *Tetrahedron*, 1970, **26**, 2923 (*config*)

Grethe, G. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2268 (*synth*, *uv*, *ir*, *pmr*, *ms*)


Mulder-Krieger, T. *et al.*, *Pharm. Weekbl., Sci. Ed.*, 1982, 91 (*cmr*)

Mulder-Krieger, T. *et al.*, *Planta Med.*, 1982, **46**, 19; 1984, **50**, 17 (*10-Methoxycinchonamine*)

Cinchonaminone

C-445

2-(3-Ethenyl-4-piperidiny)-1-[3-(2-hydroxyethyl)-1H-indol-2-yl]ethanone, 9CI [121053-50-7]



Absolute Configuration

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from *Cinchona succirubra*.


Monoamine oxidase inhibitor. Pale yellow powder + 1H₂O. $[\alpha]_D^{25} +10$ (c, 0.35 in EtOH). λ_{\max} 237 (log ε 4.14); 312 (log ε 4.23) (EtOH).

Mitsui, N. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 363-366 (*isol*, *pmr*, *cmr*, *ms*)

Cinchonidine

C-446

Cinchonan-9-ol, 9CI. α -Quinidine [485-71-2]



Absolute Configuration

C₁₉H₂₂N₂O 294.396

Stereoisomeric with Cinchonine, C-447. Alkaloid from *Cinchona tucujensis*, main base of *Cinchona succirubra*, present in many other *Cinchona* spp., also in *Remijia* spp., and the leaves of *Olea europaea* and *Ligustrum vulgare* (Rubiaceae, Oleaceae). Resolving agent, chiral co-catalyst. Antimalarial agent. Mp 210.5° (205°). $[\alpha]_D -110$ (EtOH). λ_{\max} 283 (ε 5190); 301 (ε 3940); 314 (ε 3000) (MeOH).

▶ LD₅₀ (rat, ipr) 206 mg/kg. GD2975000

Hydrochloride (1:2): Mp 242°.

Hydroiodide: Mp 215-220°.

O-Benzoyl: Mp 208-211° (183°).

N¹-Benzyl: [69257-04-1]

$C_{26}H_{29}N_2O^{\oplus}$ 385.528
Chiral phase-transfer catalyst (as chloride). Mp 210° dec. (chloride). $[\alpha]_D^{20}$ -180 (c, 1.3 in H_2O) (chloride).

Ketone: Cinchonidinone. *Cinchonan-9-one.* *Cinchoninone* [14509-68-3]
 $C_{19}H_{20}N_2O$ 292.38
Shown to be a nat. prod. in *Cinchona ledgeriana* (Rubiaceae).

10,11-Dihydro: Hydrocinchonidine. *10,11-Dihydrocinchon-9-ol, 9CI.* *10,11-Dihydrocinchonidine.* *Cinchamidine* [485-64-3]
 $C_{19}H_{24}N_2O$ 296.411
Minor base normally accompanying cinchonidine in *Cinchona* spp. (Rubiaceae). Antimalarial. Leaflets (EtOH). Mp 232° (229°). $[\alpha]_D$ -98.4 (EtOH).

10,11-Dihydro; hydrochloride:
Prisms + 2 H_2O . V. sol. H_2O . Mp 202.3° (anhyd.). $[\alpha]_D$ -98.4.


[524-61-8, 5948-98-1]

Goutarel, R. et al., *Helv. Chim. Acta*, 1950, **33**, 150-164 (*ir, uv*)
Lyle, G.G. et al., *Tetrahedron*, 1967, **23**, 3253-3263 (*config*)
Kamath, B.R. et al., *Indian J. Chem.*, 1968, **6**, 510-511 (*uv*)
Battersby, A.R. et al., *Chem. Comm.*, 1971, 30-31; 31-32 (*biosynth*)
Schneider, G. et al., *Planta Med.*, 1972, **22**, 109-116 (*isol*)
Moreland, C.G. et al., *J.O.C.*, 1974, **39**, 2413-2416 (*cmr*)
Oleksyn, B. et al., *Acta Cryst. B*, 1982, **38**, 1832-1834 (*cryst struct*)
Ihara, M. et al., *Chem. Comm.*, 1986, 573-574 (*synth, 10,11-dihydro*)
Bürgi, T. et al., *J.A.C.S.*, 1998, **120**, 12920-12926 (*pmr, conformm*)
Olsen, R.A. et al., *J.A.C.S.*, 2006, **128**, 15594-15595 (*pmr, conformm*)

Cinchonine

C-447

Cinchonan-9-ol, 9CI
[118-10-5]



Absolute Configuration

$C_{19}H_{22}N_2O$ 294.396
Stereoisomeric with Cinchonidine, C-446. Alkaloid from *Cinchona officinalis* and all *Cinchona* spp. and from some *Remijia* spp., e.g. *Remijia pedunculata* and *Remijia purdieana*, and from the leaves of *Olea europaea* and *Ligustrum vulgare* (Rubiaceae, Oleaceae). Resolving agent, chiral cocatalyst. Used in precipitation and separation of W(VI) and as a 1% soln. in dil. HNO_3 for detn. of Bi. Antimalarial. Cryst. Mp 264° Mp 255°. $[\alpha]_D$ +229 (EtOH). Log P 2.64 (calc). λ_{max} 284 (ε 5270); 301 (ε 3860); 314 (ε 3020) (MeOH). λ_{max} 225 (log ε 3.9); 284 (sh) (log ε 3.2); 314 (sh) (log ε 4) (EtOH).
▶ LD₅₀ (rat, ipr) 152 mg/kg. GD3500000
Hydrochloride: [5949-11-1]

[24302-67-8]
Mp 217-218°.
Sulfate: [5949-16-6]
Mp 206°.
▶ GD3800000

N-Me:
 $C_{20}H_{25}N_2O^{\oplus}$ 309.43
Mp 269-270° (as iodide).
O-Benzoyl: Mp 106-107°.
N-Benzyl: [69221-14-3]
 $C_{26}H_{29}N_2O^{\oplus}$ 385.528
Chiral phase transfer catalyst (as chloride). Mp 256° dec. (chloride).
10,11-Dihydro: Hydrocinchonine. (9S)-*10,11-Dihydrocinchon-9-ol, 9CI.* *10,11-Dihydrocinchonine.* *Cinchotine.* *ψ-Cinchonine.* *Pseudocinchonine.* *Cinchonifine* [485-65-4]
 $C_{19}H_{24}N_2O$ 296.411
Minor congener of Cinchonine from *Cinchona* spp., *Remijia* spp. and *Olea europaea* (Rubiaceae, Oleaceae). Antimalarial. Prisms. Mp 268-269°. $[\alpha]_D^{25}$ +204.5 (EtOH). Log P 3.12 (calc).

10,11-Dihydro; hydrochloride: Mp 220-221°.

N-[4-(Trifluoromethyl)benzyl]: N-[4-(Trifluoromethyl)benzyl]cinchoninium [95088-20-3]
[205441-34-5]
 $C_{27}H_{28}F_3N_2O^{\oplus}$ 453.526
Chiral phase transfer catalyst. Mp 245°. Comly. available as bromide.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 864C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 439A (*nmr*)

Hesse, O. et al., *Ber.*, 1871, **4**, 818-820; 1883, **16**, 58-63 (*isol*)

Rabe, P. et al., *Ber.*, 1908, **41**, 62-70 (*struct*)

Potratz, A.H. et al., *Anal. Chem.*, 1949, **21**, 1276-1279 (*detn, Bi*)

Goutarel, R. et al., *Helv. Chim. Acta*, 1950, **33**, 150-164 (*ir, uv*)

Lyle, G.G. et al., *Tetrahedron*, 1967, **23**, 51; 3253-3263 (*uv, ord, config, pmr*)

Kamath, B.R. et al., *Indian J. Chem.*, 1968, **6**, 510-511 (*uv*)

Battersby, A.R. et al., *Chem. Comm.*, 1971, 30-31; 31-32 (*biosynth*)

Schneider, G. et al., *Planta Med.*, 1972, **22**, 109-116 (*isol, uv*)

Norwitz, G. et al., *Anal. Chim. Acta*, 1974, **69**, 59-68 (*use*)

Moreland, C.G. et al., *J.O.C.*, 1974, **39**, 2413-2416 (*cmr*)

Oleksyn, B. et al., *Acta Cryst. B*, 1979, **35**, 440-444 (*cryst struct*)

Ihara, M. et al., *Chem. Comm.*, 1986, 573-574 (*synth*)

Hughes, O.L. et al., *J.O.C.*, 1987, **52**, 4745-4752 (*trifluoromethylbenzyl bromide*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1988, **13**, 264; 1989, **14**, 326-327; 1990, **15**, 340 (*trifluoromethylbenzyl bromide, use*)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **7**, 5156-5158 (*trifluoromethylbenzyl bromide, use*)


Ruiz-Mesia, L. et al., *J. Agric. Food Chem.*, 2005, **53**, 1921-1926 (*isol, pmr, cmr, ms*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CMP925

Cinchophyllamine

C-448

3β,17β-Cinchophylline
[10438-16-1]
[1355-62-0]



$C_{31}H_{36}N_4O_2$ 496.651

CAS gives different systematic names to Cinchophyllamine and Cinchophylline, which is a general name for the various stereoisomers of this struct. Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against gram-positive bacteria. Mp 143-145°. $[\alpha]_D$ +8 (c, 0.7 in $CHCl_3$). λ_{max} 207 (log ε 4.31); 228 (log ε 4.38); 281 (log ε 3.93) (EtOH).

▶ RC1210000

18,19-Dihydro: 18,19-Dihydrocinchophyllamine
[10438-18-3]

$C_{31}H_{38}N_4O_2$ 498.667

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against gram-positive bacteria. Noncryst.

3-Epimer: 3-Epicinchophyllamine. *17-Epiisocinchophyllamine.* *3α,17β-Cinchophylline*

$C_{31}H_{36}N_4O_2$ 496.651

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against gram-positive bacteria. Cryst. (CH_2Cl_2). Mp 159-160°. $[\alpha]_D$ +19 (c, 0.6 in $CHCl_3$). λ_{max} 228 (log ε 4.71); 282 (log ε 4.23); 296 (log ε 4.18); 308 (log ε 3.91) (EtOH).

3-Epimer, 4',17-didehydro: Dehydroisocinchophyllamine. *17,4'-Dehydro-3α-cinchophylline*

[77500-49-3]

$C_{31}H_{34}N_4O_2$ 494.635

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against gram-positive bacteria. Noncryst. $[\alpha]_D$ +75.5 (c, 0.75 in $CHCl_3$). λ_{max} 216 (log ε 4.63); 297 (log ε 4.07); 309 (log ε 4.05) (EtOH).

3-Epimer, 4',5',6',17-tetradehydro: Tetradehydroisocinchophyllamine. *17,4',5',6'-Tetradehydro-3α-cinchophylline*
[77500-48-2]

$C_{31}H_{32}N_4O_2$ 492.619

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Noncryst. $[\alpha]_D$ +15 (c, 0.36 in $CHCl_3$). λ_{max} 218 (log ε 4.52); 231 (log ε 4.6); 248 (sh) (log ε 4.3); 259 (sh) (log ε 4.15); 290 (log ε 4.14); 298 (log ε 4.25) (EtOH).

17-Epimer: 17-Epicinchophyllamine.

3β,17α-Cinchophylline

[77646-14-1]

$C_{31}H_{36}N_4O_2$ 496.651

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against

gram-positive bacteria. Noncryst. $[\alpha]_D$ -87.5 (c, 0.6 in CHCl_3). λ_{max} 227 (log ϵ 4.66); 282 (log ϵ 4.17); 296 (log ϵ 4.11); 308 (log ϵ 3.81) (EtOH).

3,17-Diepimer: *Isocinchophyllamine*.

3\alpha,17\alpha-Cinchophylline. *3-Epi-17-epi-cinchophyllamine*

[10438-17-2]
 $\text{C}_{31}\text{H}_{36}\text{N}_4\text{O}_2$ 496.651

Alkaloid from the leaves of *Cinchona ledgeriana* (Rubiaceae). Active against gram-positive bacteria. Cryst. (CH_2Cl_2). Mp 144°. $[\alpha]_D$ +7 (c, 0.6 in CHCl_3). λ_{max} 228 (log ϵ 4.74); 282 (log ϵ 4.26); 297 (log ϵ 4.2); 308 (log ϵ 3.96) (EtOH) (Berdy).

3,17-Diepimer, *N-Ac*: Mp 238°. $[\alpha]_D$ +121 (c, 0.35 in CHCl_3).

3,17-Diepimer, *18,19-dihydro*: Mp 163°. $[\alpha]_D$ +21 (c, 0.35 in CHCl_3).

Potier, P. et al., *Bull. Soc. Chim. Fr.*, 1966, 2309-2319 (*Cinchophyllamine*, *Isocinchophyllamine*, *uv. ir. pmr. ms*)

Guilhelm, J. et al., *Acta Cryst. B*, 1974, **30**, 742-747 (*Isocinchophyllamine*, *cryst. struct.*)


Zeche, M. et al., *Phytochemistry*, 1980, **19**, 2451 (*Cinchona ledgeriana constits*)

Carou, C. et al., *Planta Med.*, 1988, **54**, 409-412 (*activity*)

Cinereain

C-449

10,11-Dihydro-8-(1-methylethyl)-11-(2-methylpropylidene)-6H-oxepino[2,3-d]pyrazino[1,2-a]pyrimidine-6,9(8H)-dione, *9CI*
 [117013-51-1]



$\text{C}_{18}\text{H}_{21}\text{N}_3\text{O}_3$ 327.382


Isol. from *Botrytis cinerea* ATCC 64157 on sunflower seed. Plant growth regulator. Phytotoxin. Ruby-red cryst. (Me_2CO). Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 201-203°. λ_{max} 238 (ϵ 10471); 276 (ϵ 8710); 371 (ϵ 1230) (EtOH) (Berdy).

Cutler, H.G. et al., *Agric. Biol. Chem.*, 1988, **52**, 1725-1733 (*isol. struct. props*)

Cinnabaramide D

C-450

[744200-69-9]



Absolute Configuration

$\text{C}_{19}\text{H}_{31}\text{NO}_6$ 369.457

Related to Lactacystin, L-6 and Salinosporamide A, S-28. Prod. by *Streptomyces* sp. strain JS360. Needles. $[\alpha]_D$ 0. λ_{max} 202 (log ϵ 4.05) (MeOH).

(*19*→*3*)-Lactone: *Cinnabaramide B*
 [744200-67-7]

$\text{C}_{19}\text{H}_{29}\text{NO}_5$ 351.442

Prod. by *Streptomyces* sp. strain JS360. Amorph. solid. $[\alpha]_D$ -140 (c, 0.5 in MeOH). λ_{max} 201 (log ϵ 4.12); 225 (log ϵ 3.54) (MeOH).

12-Deoxy: *Cinnabaramide E*

[744200-72-4]

$\text{C}_{19}\text{H}_{31}\text{NO}_5$ 353.458

Prod. by *Streptomyces* sp. strain JS360. Needles. $[\alpha]_D^{20}$ -23 (c, 0.1 in MeOH). λ_{max} 201 (log ϵ 4) (MeOH).

12-Deoxy, (*19*→*3*)-lactone: *Cinnabaramide A*

[744200-66-6]

$\text{C}_{19}\text{H}_{29}\text{NO}_4$ 335.442

Prod. by *Streptomyces* sp. strain JS360. Cryst. $[\alpha]_D^{20}$ -93 (c, 0.5 in MeOH). λ_{max} 202 (log ϵ 4.24); 227 (log ϵ 3.68) (MeOH).

4,12-Dideoxy, (*19*→*3*)-lactone: *Cinnabaramide C*

[744200-68-8]

$\text{C}_{19}\text{H}_{29}\text{NO}_3$ 319.443

Prod. by *Streptomyces* sp. strain JS360. Amorph. solid. $[\alpha]_D^{20}$ -80 (c, 0.5 in MeOH). λ_{max} 202 (log ϵ 4.19); 223 (log ϵ 3.76) (MeOH).


Stadler, M. et al., *J. Nat. Prod.*, 2007, **70**, 246-252 (*isol. pmr. cmr. cryst. struct.*)

Ma, G. et al., *Org. Lett.*, 2007, **9**, 2143-2146 (*synth.*)

Cinnabaramide F

C-451

[744215-21-2]



$\text{C}_{24}\text{H}_{38}\text{N}_2\text{O}_7\text{S}$ 498.639

Related to Lactacystin, L-6 and Salinosporamide A, S-28. Prod. by *Streptomyces* sp. strain JS360. Proteasome inhibitor. Amorph. solid. $[\alpha]_D^{20}$ +37 (c, 0.5 in MeOH). λ_{max} 202 (log ϵ 4.15); 234 (log ϵ 3.7) (MeOH).

Me ester: *Cinnabaramide G*

[744215-22-3]

$\text{C}_{25}\text{H}_{40}\text{N}_2\text{O}_7\text{S}$ 512.666


Prod. by *Streptomyces* sp. strain JS360. Proteasome inhibitor. Amorph. solid. $[\alpha]_D^{20}$ +24 (c, 0.5 in MeOH). λ_{max} 202 (log ϵ 4.16); 230 (log ϵ 3.71) (MeOH).

Stadler, M. et al., *J. Nat. Prod.*, 2007, **70**, 246-252 (*isol. pmr. cmr.*)

Cinnabarine

C-452

2-Amino-9-hydroxymethyl-3-oxo-3H-phenoxazine-1-carboxylic acid, *9CI*. *Polystictin*
 [146-90-7]



$\text{C}_{14}\text{H}_{10}\text{N}_2\text{O}_5$ 286.243

Pigment of *Trametes cinnabarina*, *Coriolus sanguineus*, *Polystictus cinnabarinus* and *Pycnoporus sanguineus*. Active against *Staphylococcus aureus* and fungi. Orange-yellow cryst. or red needles (Py or PhNO_2 /anisole). Mp 320° (300° dec.).

Me ester:

Orange cryst. Mp 252° dec.

O-Ac: *O-Acetylcinnabarine*

[133076-09-2]

$\text{C}_{16}\text{H}_{12}\text{N}_2\text{O}_6$ 328.281

Isol. from *Pycnoporus sanguineus*. Red needles (Me_2CO /petrol). Mp 250-252° dec.

O-Ac, Me ester:

Orange cryst. Mp 234° dec.

N-Me, Me ester: [133056-32-3]

$\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_5$ 314.297

Isol. from *Pycnoporus sanguineus*. Red cryst. Mp 182° dec.

Me ether:

Red needles (Me_2CO aq. or EtOAc/petrol). Mp 183-185° Mp 200-202° dec. (variable).

I''-Aldehyde: *2-Amino-9-formyl-3-oxo-3H-phenoxazine-1-carboxylic acid*, *9CI*
 [133056-33-4]

$\text{C}_{14}\text{H}_8\text{N}_2\text{O}_5$ 284.228

Isol. from *Pycnoporus sanguineus*. Mp 228° dec. (as Me ester). Isol. as Me ester.

I''-Carboxylic acid: *2-Amino-3-oxo-3H-phenoxazine-1,9-dicarboxylic acid*, *9CI*. *Cinnabarinic acid*
 [606-59-7]

$\text{C}_{14}\text{H}_8\text{N}_2\text{O}_6$ 300.227

Minor constit. of *Pycnoporus sanguineus*. Major component of free pigment of red kangaroo (*Megaleia rufa*) hair. Dark brown cryst. (Py). Dec. >300° without melting.

I'-Aldehyde, *I''-carboxylic acid*: *2-Amino-1-formyl-3-oxo-3H-phenoxazine-9-carboxylic acid*, *9CI*. *Tramesanguin*
 [34083-17-5]

$\text{C}_{14}\text{H}_8\text{N}_2\text{O}_5$ 284.228

Pigment from the fungus *Trametes cinnabarina*. Yellow (?) cryst. (Py). Dec. without melting >250°.

I'-Aldehyde, *I''-carboxylic acid, Me ester*:
 Orange cryst. ($\text{C}_6\text{H}_6/\text{CHCl}_3$). Mp 245-250°.

Gripenberg, J. et al., *Acta Chem. Scand.*, 1951, **5**, 590; 1957, **11**, 1485; 1958, **12**, 603 (*struct.*)

Lemberg, R. et al., *Aust. J. Exp. Biol. Med. Sci.*, 1952, **30**, 271 (*isol.*)

Cavill, G.W.K. et al., *Proc. Chem. Soc., London*, 1957, 346 (*isol. struct.*)

Gripenberg, J. et al., *Acta Chem. Scand.*, 1963, **17**, 703 (*Tramesanguin*)

Subba Rao, P.V. et al., *Biochem. J.*, 1965, **95**, 628 (*synth. Cinnabarinic acid*)

Subba Rao, P.V. et al., *Arch. Biochem. Biophys.*, 1966, **115**, 27 (*biosynth. Cinnabarinic acid*)

Schäfer, W. et al., *Tet. Lett.*, 1968, 2161 (*synth.*)
 Schäfer, W. et al., *Tetrahedron*, 1972, **28**, 5261 (*uv.*)


Prinz, W. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1977, **358**, 1161 (*synth. Cinnabarinic acid*)

Achenbach, H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 3 (*derivs.*)

Giurg, M. *et al.*, *Synth. Commun.*, 2007, **37**, 1779-1789 (*Cinnabarinic acid, synth*)

Cinnamolaurine C-453

4-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)methyl]phenol, 9CI. 1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-6,7-methylenedioxyisoquinoline



C₁₈H₁₉NO₃ 297.353

(R)-form [25866-03-9]

Alkaloid from the bark of an unidentified New Guinea *Cinnamomum* sp. (Lauraceae). Prisms (EtOH). Mp 212-213° dec. [α]_D²⁵ -100 (c, 0.75 in EtOH). λ_{max} 287 (log ε 3.72) (no solvent reported).

Hydrochloride: Mp 230-233° dec.

***N*-De-Me: Norcinnamolaurine**

[34168-00-8]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the bark of an unidentified New Guinea *Cinnamomum* sp. (Lauraceae). Needles (EtOH). Mp 197-198°. [α]_D²⁵ +55 (c, 0.4 in EtOH). λ_{max} 287 (log ε 3.78) (no solvent reported).

Me ether: O-Methylcinnamolaurine

[108210-74-8]

C₁₉H₂₁NO₃ 311.38

Alkaloid from *Hedyccarya angustifolia* (Monimiaceae). Gum. Opt. rotn. not recorded.

(S)-form***Me ether: Doryafranine***

[4727-86-0]

C₁₉H₂₁NO₃ 311.38

Major alkaloid from *Doryphora sassafras* (Monimiaceae). Plates (hexane). Mp 92-94°. [α]_D²⁷ +41.5 (c, 2.12 in CHCl₃). Abs. config. not explicitly stated. May be identical with *O*-Methylcinnamolaurine.

Me ether, hydrochloride: Mp 195-197°.

Gharbo, S.A. *et al.*, *J. Nat. Prod.*, 1965, **28**, 237-244 (*Doryafranine*)


Gellert, E. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2095-2099 (*R*-form, *Norcinnamolaurine, isol, uv, pmr, struct, synth*)

Marsaioli, A.J. *et al.*, *Phytochemistry*, 1978, **17**, 1655-1658 (*cmr, Doryafranine*)

Gunawardana, Y.A.G.P. *et al.*, *Heterocycles*, 1987, **26**, 447-456 (*O-Methylcinnamolaurine*)

3-[2-(Cinnamoylamino)ethyl]-3-hydroxy-2-indolinone C-454

N-[2-(2,3-Dihydro-3-hydroxy-2-oxo-1*H*-indol-3-yl)ethyl]-3-phenyl-2-propenamide, 9CI



C₁₉H₁₈N₂O₃ 322.363


(±)-(E)-form [79087-89-1]

Alkaloid from the leaves of *Cinnamomum triplinervis* (Lauraceae). Cryst. (MeOH/C₆H₆). Mp 162-164°.

Ripperger, H. *et al.*, *Phytochemistry*, 1981, **20**, 1453-1454 (*isol, uv, ir, pmr, cmr, ms, struct*)

***N*^α-Cinnamoylhistamine** C-455

N-[2-(1*H*-Imidazol-4-yl)ethyl]-3-phenyl-2-propenamide, 9CI. *N*-(2-Imidazol-4-ylethyl) cinnamamide, 8CI



(*E*)-1*H*-form

C₁₄H₁₅N₃O 241.292

(E)-1*H*-form [53813-77-7]

Alkaloid from *Acacia argentea* (formerly *Acacia leptostachya*), *Acacia polystachya*, *Acacia spirorbis*, *Acacia longifolia*, *Argyrodendron peralatum* (preferred genus name *Croton*), *Glochidion* sp. (probably *Glochidion philippicum*), *Dolichochele sphaerica* and *Lycium cestroides* leaves collected in autumn (Fabaceae, Euphorbiaceae, Solanaceae). Needles (H₂O). Mp 179-180.5°.

Hydrochloride: Mp 189-190° (187-189°).

Picrate: Mp 165-166°.

***N*¹-Me: trans-*N*^α-Cinnamoyl-*N*¹-methylhistamine**

[126661-93-6]

C₁₅H₁₇N₃O 255.319

Isol. from *Lycium cestroides* (Solanaceae).

4''-Hydroxy, 3''-methoxy: (4-Hydroxy-3-methoxycinnamoyl)histamine. *N*^α-*Feruloyl*histamine

C₁₅H₁₇N₃O₃ 287.318

Alkaloid from the roots of an *Ephedra* sp. (Ephedraceae). Shows weak hypotensive activity. Mp 160-162°.

(Z)-1*H*-form [95906-73-3]

Major alkaloid from the leaves of *Lycium cestroides* collected in autumn.

***N*¹-Me: cis-*N*^α-Cinnamoyl-*N*¹-methylhistamine**

[126661-92-5]

C₁₅H₁₇N₃O 255.319

Isol. from *Lycium cestroides* (Solanaceae).

(E)-3*H*-form

Trace const. of *Lycium cestroides* (Solanaceae). Intramolecularly H bonded; referred to as the 'quasibicyclic isomer'. Readily isomerised to the normal 1*H*-isomer.

(Z)-3*H*-form

Isol. from *Lycium cestroides* (Solanaceae).

Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1964, **17**, 375 (*isol, pmr, ms, synth*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 555; 1969, **22**, 1309 (*isol*)

Rosenberg, H. *et al.*, *J. Nat. Prod.*, 1974, **37**, 313 (*biosynth*)

Repke, D.B. *et al.*, *J. Nat. Prod.*, 1975, **38**, 101 (*isol, uv, pmr, ms*)

Poupat, C. *et al.*, *Phytochemistry*, 1975, **14**, 1881 (*isol*)

Hikino, H. *et al.*, *Planta Med.*, 1983, **48**, 108 (*N-Feruloylhistamine*)

Chiale, C.A. *et al.*, *An. Asoc. Quim. Argent.*, 1984, **72**, 569; *CA*, **102**, 163700s (*Z-isomer*)

Chiale, C.A. *et al.*, *Phytochemistry*, 1990, **29**, 688 (*3*H*-forms, derivs*)

***N*-Cinnamoylputrescine** C-456

H₂NCH₂CH₂CH₂CH₂NH

COCH=CHPh

C₁₃H₁₈N₂O 218.298

(E)-form

N'-(2-Methylpropanoyl): *N*¹-Cinnamoyl-*N*⁴-(2-methylpropanoyl)putrescine. *Secopiriferine* [349642-86-0]

C₁₇H₂₄N₂O₂ 288.389

Alkaloid from *Aglaia gracilis*. Cryst. (CHCl₃). Mp 161-161.5°. λ_{max} 218; 274 (MeOH).

N'-(2*S*-Methylbutanoyl): *N*¹-Cinnamoyl-*N*⁴-(2-methylbutanoyl)putrescine. *Secoodorine* [349642-87-1]

C₁₈H₂₆N₂O₂ 302.416

Alkaloid from *Aglaia gracilis*. Cryst. (CHCl₃). Mp 135-137° (natural) Mp 146.5-147° (synthetic). [α]_D²⁰ +11 (c, 0.3 in CHCl₃). λ_{max} 218; 274 (MeOH).

N'-Tigloyl: *Grandiamide B. N*¹-Cinnamoyl-*N*⁴-tigloylputrescine [279678-94-3]

C₁₈H₂₄N₂O₂ 300.4

Alkaloid from *Aglaia grandis*. Cryst. (hexane/EtOH). Mp 100.5-101.5°.

N'-(3-Hydroxy-2*ξ*-methylpropanoyl): *Secoisopiriferinol*

C₁₇H₂₄N₂O₃ 304.388

Alkaloid from the leaves of *Aglaia spectabilis*. [α]_D²⁰ +21 (c, 0.2 in MeOH). λ_{max} 216 (log ε 4.31); 222 (sh) (log ε 4.19); 276 (log ε 4.24) (MeOH aq.).

N'-[3-(Methylthio)-2*E*-propenoyl]: *Secopyrrolotenin*

C₁₇H₂₂N₂O₂S 318.439

Alkaloid from the leaves of *Aglaia tenuicaulis*. λ_{max} 217 (log ε 4.2); 221 (sh) (log ε 4.17); 274 (log ε 4.53) (MeOH aq.).

N'-(4-Hydroxy-2*ξ*-methylbutanoyl): *Secoisodorinol*

C₁₈H₂₆N₂O₃ 318.415

Alkaloid from the leaves of *Aglaia spectabilis*. Needles. Mp 131-132°. [α]_D²⁰ +12 (c, 0.4 in MeOH). λ_{max} 216 (log ε 4.12); 222 (sh) (log ε 4.04); 274 (log ε 4.25) (MeOH aq.).

N'-(4-Hydroxy-2-methyl-2*E*-butenoyl): *N*¹-Cinnamoyl-*N*⁴-(4-hydroxytigloyl)-putrescine. *Aglairubine. Dasyclamide*

[373644-07-6]

[218144-40-2]

C₁₈H₂₄N₂O₃ 316.399

Alkaloid from *Aglaia dasyclada* and *Aglaia rubiginosa*. Needles (MeOH). Mp 116-117°.

N'-(3-Hydroxy-2-methylenebutanoyl):

Grandiamide C

[279678-95-4]

C₁₈H₂₄N₂O₃ 316.399Alkaloid from *Aglaiia grandis*.

Amorph. powder. Mp 104.5-106.5° (synthetic). Probably racemic.

N'-(3R,4-Dihydroxy-2-methylenebutanoyl): **Grandiamide D**

[955123-03-2]

C₁₈H₂₄N₂O₄ 332.399Alkaloid from the leaves of *Aglaiia gigantea*. Amorph. solid. [α]_D²⁰ +200 (c, 0.47 in MeOH). λ_{max} 217 (ε 14950); 222 (ε 12290); 277 (ε 16610) (MeOH).

N'-Benzoyl: N-[4-[1-(1-Oxo-3-phenyl-2-propenyl)amino]butyl]benzamide. N-Benzoyl-N'-cinnamoylputrescine. **Haplamiidine**. Pyrimidatine

[64223-54-7]

C₂₀H₂₂N₂O₂ 322.406Alkaloid from *Aglaiia pyramidata* and *Haplophyllum latifolium*. Mp 139-140° Mp 173-174°. Haplamiidine (to which lower Mp refers) was not specifically assigned the (E)-config., but is presumably identical with Pyrimidatine.

4''-Hydroxy: 4-Hydroxycinnamoylputrescine. **4-Coumaroylputrescine**

[34136-53-3]

C₁₃H₁₈N₂O₂ 234.297Alkaloid from *Pennisetum americanum* (pearl millet), *Triticum vulgare*, *Zea mays* (sweet corn), *Persea gratissima* (avocado), *Salix* sp., *Lycopersicon esculentum* (tomato), *Nicotiana tabacum* and a *Petunia* hybrid (Poaceae, Lauraceae, Salicaceae, Solanaceae). Faint yellow needles + 1H₂O (H₂O). Mp 182-183.5°.

4''-Hydroxy, N'-(4-hydroxybenzoyl):

Glochidiatusamide. *Glochidiatusin*C₂₀H₂₂N₂O₄ 354.405Alkaloid from *Thyrocarpus glochidiatus*. Cryst. Mp 280-282°. λ_{max} 228; 255; 292; 310 (MeOH).

3'',4''-Dihydroxy: N-(4-Aminobutyl)-3-(3,4-dihydroxyphenyl)-2-propanamide.

N-(3,4-Dihydroxycinnamoyl)-1,4-butanediamine. **Caffeoylputrescine**. **Paucine**

[29554-26-5]

[59862-95-2, 26148-06-1]

C₁₃H₁₈N₂O₃ 250.297Alkaloid from *Nicotiana tabacum*, *Pentaclethra macrophylla*, *Salix* sp., *Persea gratissima* and a *Petunia* hybrid (Solanaceae, Fabaceae, Salicaceae, Lauraceae). Also isol. from elicitor-treated cell suspension cultures and fungus-infected leaves of *Solanum tuberosum* (Solanaceae). Cryst. + 2H₂O (as hydrochloride). Mp 247-250° dec. (hydrochloride). λ_{max} 219; 235; 295; 320 (H₂).

3''-Methoxy, 4''-hydroxy: **Subaphylline**. *Feruloylputrescine*

[501-13-3]

C₁₄H₂₀N₂O₃ 264.324Alkaloid from *Ananas comosus* (pineapple), *Pennisetum*, *Triticum*, *Gomphrena*, *Salix*, *Persea* and other spp. (Bromeliaceae, Poaceae, Amaranthaceae, Salicaceae, Lauraceae, Solana-ceae, Rutaceae, Chenopodiaceae). Also isol. from elicitor-treated cell suspension cultures and fungus-infected leaves of *Solanum tuberosum* (Solanaceae). Cryst. (C₆H₆ or MeOH). Mp 171.5-172°.

▶GD6981000

3''-Methoxy, 4''-hydroxy, hydrochloride: Cryst. + 1H₂O. Mp 151-152°.

3''-Methoxy, 4''-hydroxy, N'-(4-hydroxybenzoyl): **N-trans-Feruloyl-N'-(4-hydroxybenzoyl)putrescine**

C₂₁H₂₄N₂O₅ 384.431Alkaloid from the roots of *Paris verticillata*. Yellowish gum. λ_{max} 247 (log ε 3.63); 317 (log ε 3.45) (MeOH).

4''-Methoxy, N,N',N'-tri-Me: [29802-83-3]

Synthetic. Pale yellow viscous oil. Bp₃ 239-243°.

3'',5''-Dimethoxy, 4''-hydroxy: **Sinapoylputrescine**. N-(3,4-Dihydroxy-5-methoxycinnamoyl)-1,4-butanediamine

[70185-57-8]

C₁₅H₂₂N₂O₄ 294.35Alkaloid from *Ananas comosus* (pineapple) and *Lilium* sp. (Bromeliaceae, Liliaceae).**(Z)-form**

4''-Hydroxy, N,N',N'-tri-Me: N-[4-(Dimethylamino)butyl]-3-(4-hydroxyphenyl)-N-methyl-2-propenamide. N-[4-(Dimethylamino)butyl]-p-hydroxy-N-methylcinnamamide. N'-(4-Hydroxycinnamoyl)-N,N,N'-trimethylputrescine

[30891-99-7]

C₁₆H₂₄N₂O₂ 276.378Alkaloid from the leaves of *Kniphofia flavovirens*, *Kniphofia foliosa* and *Kniphofia tuckii* (Liliaceae). Mp 60-65°.

4''-Methoxy, N,N',N'-tri-Me: N,N,N'-Trimethyl-N'-(4-methoxycinnamoyl)-putrescine

[29802-82-2]

C₁₇H₂₆N₂O₂ 290.405Alkaloid from *Kniphofia flavovirens*, *Kniphofia foliosa* and *Kniphofia tuckii* (Liliaceae). Oil. Bp₆ 250-256° (bath).

Ryabinin, A.A. et al., *Dokl. Akad. Nauk SSSR*, 1949, **67**, 513-510; 1951, **76**, 689-692; *CA*, **44**, 1455h; **45**, 8479d (*Subaphylline*)

Wheaton, T.A. et al., *Nature (London)*, 1965, **206**, 620-621 (*Subaphylline*)

Ripperger, H. et al., *J. Prakt. Chem.*, 1970, **312**, 449-455 (*Kniphofia alkaloids*)

Hollerbach, A. et al., *Monatsh. Chem.*, 1970, **101**, 141-156 (*Paucine*)

Mizusaki, S. et al., *Phytochemistry*, 1971, **10**, 1347-1348 (*Paucine*, 4-Coumaroylputrescine, *Subaphylline*, *isol*, *synth*)

Buta, J.G. et al., *Phytochemistry*, 1972, **11**, 1188-1189 (*Paucine*)

Martin-Tanguy, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1973, **276**, 1433-1435 (*Subaphylline*, *uv*, *synth*)

Mbadiwe, E.I. et al., *Phytochemistry*, 1973, **12**, 2546 (*Caffeoylputrescine*)

Cabanne, F. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1976, **282**, 1959-1962 (*Paucine*, *uv*)

Nesmelova, E.F. et al., *Khim. Prir. Soedin.*, 1977, **13**, 427; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 365-366 (*Haplamiidine*)

Martin-Tanguy, J. et al., *Phytochemistry*, 1978, **17**, 1927-1928 (*Paucine*, *Subaphylline*, *Sinapoylputrescine*, *occur*)

Saifah, E. et al., *J. Nat. Prod.*, 1993, **56**, 473-477 (*Pyrimidatine*)

Keller, H. et al., *Phytochemistry*, 1996, **42**, 389-396 (*Paucine*, *Subaphylline*, *isol*, *pnr*)

Inada, A. et al., *Phytochemistry*, 2000, **53**, 1091-1095 (*Grandiamides*)

Chaidir, et al., *J. Nat. Prod.*, 2001, **64**, 1216-1220 (*Dasyclamide*)

Greger, H. et al., *Phytochemistry*, 2001, **57**, 57-64 (*Secoisodorine*, *Secopiriferine*)

Seger, C. et al., *Monatsh. Chem.*, 2002, **133**, 97-100 (*Aglairubine*)

Detterbeck, R. et al., *Tetrahedron*, 2002, **58**, 6887-6893 (*Aglaiia bisamides*, *synth*)

Luo, Y. et al., *Nat. Prod. Res.*, 2006, **20**, 1063-1066 (*Glochidiatusamide*)


Duong, T.N. et al., *J. Nat. Prod.*, 2007, **70**, 1640-1643 (*Dasyclamide*, *Grandiamide D*)

Lee, K.H. et al., *Molecules*, 2008, **13**, 41-45 (*N-Feruloyl-N'-hydroxybenzoylputrescine*)

Greger, H. et al., *Phytochemistry*, 2008, **69**, 928-938 (*Secoisodorinol*, *Secoisopiriferinol*, *Secopyrrolotenin*)

1-Cinnamoyl-1H-pyrrole C-457


1-(1-Oxo-3-phenyl-2-propenyl)-1H-pyrrole, 9CI. Cinnamic acid pyrrolide

C₁₃H₁₁NO 197.236**(E)-form** [252248-89-8]Alkaloid from *Piper argyrophyllum*. Needles (EtOAc/petrol). Mp 108°. λ_{max} 297 (MeOH).

Gupta, S. et al., *Indian J. Chem., Sect. B*, 1999, **38**, 823-827 (*isol*, *pnr*, *cmr*)

1-Cinnamoylpyrrolidine, 8CI C-458

1-(1-Oxo-3-phenyl-2-propenyl)pyrrolidine, 9CI. Cinnamic acid pyrrolidide [19202-21-2]

C₁₃H₁₅NO 201.268**(E)-form** [52438-21-8]Alkaloid from the roots of *Piper methysticum* (kava) (Piperaceae). Needles (petrol). Mp 101-103° (94-95°).

Achenbach, H. et al., *Chem. Ber.*, 1970, **103**, 2535 (*isol*, *uv*, *ir*, *ms*, *synth*, *struct*)

Engel, P. et al., *Z. Kristallogr., Kristallphys., Kristallchem.*, 1974, **139**, 207 (*cryst struct*)

Huang, Y.Z. et al., *Tet. Lett.*, 1987, **28**, 2159 (*synth*)

Rajeswari, S. et al., *Tet. Lett.*, 1987, **28**, 5099 (*synth*)


Son, T. et al., *Bull. Chem. Soc. Jpn.*, 1988, **61**, 1251 (*synth*)

Lindsay, C.M. et al., *J.C.S. Perkin I*, 1988, 569 (*synth*)

Ohta, A. et al., *Heterocycles*, 1991, **32**, 965 (*synth*)

Cinodine I, 9CI**C-459**

Glycocinmaspermicidin B. Antibiotic BM 123₇₁. BM 123₇₁. Antibiotic LL-BM 123₇₁. LL-BM 123₇₁ [60830-76-4]



C₃₇H₅₉N₁₃O₁₃ 893.952

Glycolipid antibiotic. Isol. from *Nocardia* spp. Active against gram-positive and -negative bacteria and exp. infections in mice. DNA gyrase inhibitor. Sol. H₂O; fairly sol. MeOH; poorly sol. Et₂O, hexane, butanol. λ_{max} 286 (ε 22500) (H₂O) (Derep). λ_{max} 286 (E1%/1cm 225) (MeOH) (Berdy).

► LD₅₀ (mus, scu) 88 mg/kg. GE3753000

Hydrochloride: Cinodine hydrochloride. CL 98984 [68782-58-1]

Amorph. powder. [α]_D²⁵ +55 (c, 0.8 in H₂O). Dec. at ca. 200°.

Tresner, H. et al., *J. Antibiot.*, 1978, **31**, 394-397 (*isol*)

Martin, J.H. et al., *J. Antibiot.*, 1978, **31**, 398-404 (*isol, props*)


Ellestad, G.A. et al., *J.A.C.S.*, 1978, **100**, 2515-2524 (*struct*)

Chiu, S.H.L. et al., *J. Antibiot.*, 1984, **37**, 1000-1006; 1079-1081 (*biosynth, cmr*)

Osborne, M.S. et al., *Antimicrob. Agents Chemother.*, 1990, **34**, 1450-1452 (*activity*)

Cinodine II, 9CI**C-460**

LL-BM 123₇₂. Antibiotic LL-BM 123₇₂. BM 123₇₂. Antibiotic BM 123₇₂. Glycocinmaspermicidin C [60830-75-3]



C₃₇H₅₉N₁₃O₁₃ 893.952

Glycolipid antibiotic. Isol. from *Nocardia* sp. Active against gram-positive and -negative bacteria and exp. infections in mice. Amorph. powder (as hydrochloride). Sol. H₂O; fairly sol. MeOH; poorly sol. butanol, hexane. Mp 200° dec. (hydrochloride). [α]_D²⁵ +60 (c, 0.85 in H₂O). λ_{max} 286 (ε 22500) (H₂O) (Derep). λ_{max} 286 (E1%/1cm 220) (MeOH) (Berdy). λ_{max} 286 (E1%/1cm 220) (HCl) (Berdy). λ_{max} 286 (E1%/1cm 220) (NaOH) (Berdy).

► CB9626010


Ellestad, G.A. et al., *J.A.C.S.*, 1978, **100**, 2515 (*struct*)

U.S. Pat., 1979, 4 154 925; *CA*, **91**, 138848t (*manuf*)

Chiu, S.H.L. et al., *J. Antibiot.*, 1984, **37**, 1000; 1079 (*biosynth*)

Circinamide**C-461**

[190271-96-6]




C₁₈H₃₄N₄O₅ 386.49

From the cyanobacterium *Anabaena circinalis* isol. from Lake Kasumigaura, Japan. Papain inhibitor. Cryst. Sol. MeOH. [α]_D²⁰ +15.5 (c, 0.1 in MeOH). λ_{max} 250 (ε 530) (MeOH). λ_{max} 250 (ε 530) (MeOH) (Berdy).

Shin, H.J. et al., *Tetrahedron*, 1997, **53**, 5747-5754 (*isol, uv, pmr, cmr*)

Circinatin**C-462**

[133507-94-5]




C₂₀H₃₁N₃O₇ 425.481

Metab. of the plant pathogenic fungus *Periconia circinata*. Cryst. (H₂O). Mp 165°. [α]_D +29 (c, 0.9 in H₂O).

Macko, V. et al., *Experientia*, 1990, **46**, 1206 (*isol*)

Macko, V. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 9574-9578 (*config*)

Circumdatin A**C-463**

C₂₁H₁₉N₃O₅ 393.398

Structure revised in 2008.

(S)-form [223130-52-7]

Prod. by *Aspergillus ochraceus* and the marine-derived *Aspergillus ostianus* 01F313. Orange-red solid. [α]_D²² -421 (c, 0.018 in EtOH). λ_{max} 238 (sh) (log ε 3.84); 290 (sh) (log ε 3.33); 357 (log ε 3.3) (EtOH).

5-Demethoxy: Circumdatin B


[223130-57-2]

C₂₀H₁₇N₃O₄ 363.372

Prod. by *Aspergillus ochraceus* and *Aspergillus ostianus* 01F313. Orange-red solid. [α]_D²² -163 (c, 0.04 in EtOH). λ_{max} 284 (sh) (log ε 2.73); 358 (log ε 2.76) (EtOH).

Rahbaek, L. et al., *J.O.C.*, 1999, **64**, 1689-1692 (*isol, uv, cd, pmr, cmr*)

Ookura, R. et al., *J.O.C.*, 2008, **73**, 4245-4247 (*isol, pmr, cmr, cryst struct*)

Circumdatin C**C-464**

C₁₇H₁₃N₃O₃ 307.308

(S)-form [223130-61-8]

Prod. by a terrestrial strain of *Aspergillus ochraceus* and a marine-derived *Exophiala* sp. MFC353-1. Solid. [α]_D²² -75 (c, 0.16 in MeOH). λ_{max} 229 (log ε 4.31); 272 (log ε 3.85); 312 (log ε 3.5) (MeOH).

Deoxy: Circumdatin F

[232922-23-5]

C₁₇H₁₃N₃O₂ 291.309

Prod. by a terrestrial strain of *Aspergillus ochraceus*. Solid. [α]_D -18.9 (c, 0.1 in MeOH).

Deoxy, 15-hydroxy: Circumdatin G

[326597-48-2]

C₁₇H₁₃N₃O₃ 307.308

Prod. by a marine strain of *Aspergillus ochraceus* and a marine-derived *Exophiala* sp. MFC353-1. Powder. [α]_D -21.7 (c, 0.2 in MeOH). λ_{max} 215 (log ε 2.58); 235 (log ε 2.64); 276 (log ε 1.96); 278 (log ε 2.17); 327 (log ε 1.98) (MeOH).

Rahbaek, L. et al., *J. Nat. Prod.*, 1999, **62**, 904-905 (*Circumdatin F*)

Rahbaek, L. et al., *J.O.C.*, 1999, **64**, 1689-1692 (*Circumdatin C*)


Dai, J.-R. et al., *J. Nat. Prod.*, 2001, **64**, 125-126 (*Circumdatin G*)

Witt, A. et al., *J.O.C.*, 2001, **66**, 2784-2788 (*synth*)

Snider, B. et al., *Tetrahedron*, 2001, **57**, 3301-3307 (*Circumdatin F, synth*)

Liu, J.-F. et al., *J.O.C.*, 2005, **70**, 10488-10493 (*Circumdatin F, synth*)

Zhang, D. et al., *J. Antibiot.*, 2008, **61**, 40-42 (*Exophiala constits*)

Circumdatin E**C-465**

C₂₀H₁₇N₃O₄ 363.372

(S)-form [232922-22-4]

Prod. by a terrestrial strain of *Aspergillus ochraceus* and the marine-derived *Aspergillus ostianus* 01F313. $[\alpha]_D^{25}$ -90 (c, 0.007 in MeOH). λ_{\max} 244 (log ϵ 4); 339 (log ϵ 3.16) (MeOH).

15-Deoxy: Circumdatin H

$C_{20}H_{17}N_3O_3$ 347.373

Prod. by *Aspergillus ochraceus* and *Aspergillus ostianus* 01F313. Amorph. solid. $[\alpha]_D^{25}$ -26.3 (c, 0.08 in MeOH). λ_{\max} 230 (log ϵ 2.04); 276 (log ϵ 2.61); 329 (log ϵ 3.08) (MeOH).

5-Methoxy: Circumdatin D

[232922-21-3]

$C_{21}H_{19}N_3O_5$ 393.398

Prod. by *Aspergillus ochraceus* and *Aspergillus ostianus* 01F313. Solid. $[\alpha]_D^{25}$ -129 (c, 0.02 in MeOH). λ_{\max} 247 (log ϵ 4.28); 285 (log ϵ 3.83); 339 (log ϵ 3.5) (MeOH).

5-Methoxy, 15-deoxy: Circumdatin J

[1031388-27-8]

$C_{21}H_{19}N_3O_4$ 377.399

Prod. by a marine strain of *Aspergillus ostianus* 01F313. Amorph. solid. λ_{\max} 287 (ϵ 17000); 330 (ϵ 7600); 343 (ϵ 6000) (EtOH).

Rahbaek, L. *et al.*, *J. Nat. Prod.*, 1999, **62**, 904-905 (*Circumdatins D,E*)


López-Gresa, M.P. *et al.*, *J. Antibiot.*, 2005, **58**, 416-419 (*Circumdatin H*)

Ookura, R. *et al.*, *J.O.C.*, 2008, **73**, 4245-4247 (*Circumdatin J*)

Cissaglaberrimine

C-466

3-Hydroxy-1,2-methylenedioxy-noraporphine
[188845-73-0]



$C_{17}H_{15}NO_3$ 281.31

(ξ)-form


Alkaloid from leaves and stems of *Cissampelos glaberrima*. Red-brown amorph. powder. Mp 203-205° dec. λ_{\max} 225 ; 242 ; 278 ; 300 (sh) (MeOH).

Barbosa-Filho, J.M. *et al.*, *Phytochemistry*, 1997, **44**, 959-961 (*isol, pmr, cmr*)

Cissampentin

C-467

[147451-90-9]




$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from *Cissampelos mucronata*, aerial parts of *Cissampelos fasciculata* and roots of *Cissampelos mucronata* (Menispermaceae). Exhibits significant activity as a repellent to the leafcutter ant, and limited antifungal activity. Yellow oil. $[\alpha]_D^{25}$ 0 (c, 0.002 in CH_2Cl_2).

Galinis, D.L. *et al.*, *Tetrahedron*, 1993, **49**, 1337-1342 (*isol, pmr, cmr, ms, struct*)
Tshibangu, J.N. *et al.*, *Phytochem. Anal.*, 2003, **14**, 13-22 (*isol, hplc*)

Cisetin

C-468



$C_{23}H_{33}NO_4$ 387.518

Tetramic acid deriv. Enolised triketone. Related to Equisetin, E-142. Prod. by the fungus OS 50185. Antibacterial agent. Powder. λ_{\max} 200 ; 235 ; 291 (MeOH aq.).

6',8a'-Diepimer, N-de-Me: Paecilosetin

$C_{22}H_{31}NO_4$ 373.491

Prod. by *Paecilomyces farinosus*. Antibacterial agent. Pale yellow solid. $[\alpha]_D^{20}$ -398 (c, 0.1 in MeOH). λ_{\max} 227 (log ϵ 3.95); 250 (log ϵ 3.91); 286 (log ϵ 4.08) (MeOH).


Boros, C. *et al.*, *J. Antibiot.*, 2003, **56**, 862-865 (*Cisetin*)

Lang, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 810-811 (*Paecilosetin*)

Citbismine B

C-469

[169055-77-0]



$C_{36}H_{34}N_2O_{11}$ 670.671

Alkaloid from roots of Marsh grapefruit (*Citrus paradisi*) and Hirado-buntan (*Citrus grandis*) (Rutaceae). Yellow cubes (Me_2CO). Mp 336-344°. $[\alpha]_D$ 0 ($CHCl_3$).

N^{10'}-Me: Citbismine C

[169055-79-2]

$C_{37}H_{36}N_2O_{11}$ 684.698

From roots of *Citrus paradisi* (grapefruit) and *Citrus grandis* (pummelo). Yellow cubes (Me_2CO). Mp 314-326°. Opt. inactive.

O^{6'}-De-Me, N^{10'}-Me: Citbismine E

[173429-82-8]

$C_{36}H_{34}N_2O_{11}$ 670.671

From roots of *Citrus paradisi* (grapefruit) and *Citrus grandis* (pummelo). Pale yellow oil. Opt. inactive.

6-Deoxy, O⁵-de-Me: Citbismine A

[161068-61-7]

$C_{35}H_{32}N_2O_{10}$ 640.645

Alkaloid from roots of *Citrus paradisi* (Marsh grapefruit) and *Citrus grandis* (pummelo) (Rutaceae). Yellow cubes (DMSO). Mp 335-336°. $[\alpha]_D$ 0 (DMSO).

6-Deoxy, O⁵-de-Me, N^{10'}-Me: Citbismine F

$C_{36}H_{34}N_2O_{10}$ 654.672

Alkaloid from *Citrus paradisi* (Marsh grapefruit). Yellow cubes. Mp 330-341° (dec.). $[\alpha]_D$ 0 ($CHCl_3$). λ_{\max} 222 (log ϵ 3.87); 266 (log ϵ 4.28); 275 (log ϵ 4.27); 334 (log ϵ 3.77); 390 (log ϵ 3.41) (EtOH).

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1548; 1995, **43**, 1340 (*Citbismine A, Citbismine B, isol, uv, ir, pmr, cmr, ms, crystal structure*)


Ju-ichi, M. *et al.*, *Heterocycles*, 1996, **42**, 237 (*Citbismine E*)

Takemura, Y. *et al.*, *Heterocycles*, 1999, **51**, 851-855 (*Citbismine F*)

Citbismine D

C-470

[173429-81-7]



$C_{40}H_{38}N_2O_{11}$ 722.747


Alkaloid from roots of Marsh grapefruit (*Citrus paradisi*). Pale yellow oil. Opt. inactive.

Ju-ichi, M. *et al.*, *Heterocycles*, 1996, **42**, 237 (*isol, uv, ir, pmr, ms, struct*)

Citorellamine

C-471

[99102-23-5]



$C_{22}H_{24}Br_2N_4S$ 536.332


Initially considered to be a C_{11} monomer. Alkaloid from the tunicate *Polycitrella mariae*. Exhibits cytotoxicity and potent antimicrobial activity. Needles (MeOH). Mp 210° dec. λ_{\max} 225 (ϵ 18000); 285 (ϵ 3000); 291 (ϵ 2800) (MeOH) (Derep).

Roll, D.M. *et al.*, *Tet. Lett.*, 1985, **26**, 4303-4306 (*uv, ir, pmr, cmr, ms*)

Moriarty, R.M. *et al.*, *Tet. Lett.*, 1987, **28**, 749-752 (*pmr, cmr, ms, struct, synth*)

Citracridone III C-472

3,12-Dihydro-6,10,11-trihydroxy-3,3,12-trimethyl-7H-pyrano[2,3-c]acridin-7-one, 9CI
[139219-98-0]



C₁₉H₁₇NO₅ 339.347

Acridone numbering shown. Alkaloid from *Citrus yuko* (Rutaceae). Yellow cubes (CH₂Cl₂). Mp 135-140°.

5-Me ether: Citracridone I

[81525-61-3]

C₂₀H₁₉NO₅ 353.374

Alkaloid from the root bark of *Citrus depressa*, *Citrus sinensis* var. *brasiliensis* (navel orange) and *Citrus grandis* f. *hakunikuyu* (Rutaceae). Orange plates (Me₂CO). Mp 275-278°.

5-Me ether, N-de-Me: Acrifoline†. N-Demethylcitracridone I

[172335-13-6]

C₁₉H₁₇NO₅ 339.347

Alkaloid from root bark and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow amorph. powder.

6-Me ether: 3,12-Dihydro-6,11-dihydroxy-10-methoxy-3,3,12-trimethyl-7H-pyrano[2,3-c]acridin-7-one, 9CI. 2',2'-Dimethyl(pyrano-5',6':3,4)-1,5-dihydroxy-6-methoxy-10-methylacridone

[90846-44-9]

C₂₀H₁₉NO₅ 353.374

Alkaloid from the roots of *Citrus decumana* (pummelo) (Rutaceae). Cryst. (MeOH). Mp 260°.

5,6-Di-Me ether: Citracridone II

[81525-62-4]

C₂₁H₂₁NO₅ 367.401

From *Citrus depressa* and *Citrus grandis* f. *hakunikuyu* root barks (Rutaceae). Yellow needles (Et₂O). Mp 161-163°.

1',2'-Dihydro, 1',2'-dihydroxy (trans-), 5-Me ether: Dihydroxycitracridone I

[161043-20-5]

C₂₀H₂₁NO₇ 387.388

Alkaloid from roots of *Citrus paradisi* (Rutaceae). Yellow cubes. Mp 235-238°. [α]_D +107 (c, 0.056 in EtOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 895; 901 (*isol, uv, ir, pmr, ms, struct*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1983, **22**, 1493 (*isol*)

Basa, S.C. *et al.*, *J. Nat. Prod.*, 1984, **47**, 325 (6-Me ether)


Ju-ichi, M. *et al.*, *Heterocycles*, 1991, **32**, 1781 (*isol, pmr, cmr, struct*)

Takemura, Y. *et al.*, *Heterocycles*, 1995, **41**, 187-190 (*Dihydroxycitracridone I*)

Ono, T. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1629-1631 (*Acrifoline*)

Citreamicin α C-473

LL-E19085α. Antibiotic LL-E19085α [122535-63-1]



R¹ = COCH₂CH(CH₃)₂

R² = CH₃

C₃₆H₃₁NO₁₂ 669.64

Quinonoid antibiotic. Struct. revised in 1990. Prod. by *Micromonospora citrea* NRRL 18351. Feed additive. Active against gram-positive bacteria and anaerobes. Coccidiostatic agent. Sol. CHCl₃, Me₂CO, CH₂Cl₂; fairly sol. MeCN, EtOAc, MeOH, EtOH; poorly sol. H₂O, hexane. [α]_D²⁵ -58 (c, 0.295 in CHCl₃). λ_{max} 223 (ε 42400); 255 (ε 38900); 320 (ε 32500); 384 (ε 9030); 410 (ε 6750) (MeCN) (Derep). λ_{max} 222 (ε 38200); 240 (ε 30100); 255 (ε 30100); 321 (ε 24500); 384 (ε 8050) (MeOH) (Berdy). λ_{max} 224 (ε 27700); 255 (ε 24900); 328 (ε 16400); 420 (ε 4230) (MeOH/HCl) (Berdy). λ_{max} 217 (ε 76100); 340 (ε 19100); 399 (ε 12900) (MeOH/NaOH) (Berdy).

O²-De-Me: Citreamicin ζ

[128969-89-1]

C₃₅H₂₉NO₁₂ 655.614

From *Micromonospora citrea* NRRL 18351. Feed additive. Sol. CHCl₃, CH₂Cl₂, Me₂CO; fairly sol. MeCN, EtOH, EtOAc, MeOH; poorly sol. H₂O, hexane. λ_{max} 223 (ε 42400); 255 (ε 38900); 320 (ε 32500); 384 (ε 9030); 410 (ε 6750) (MeCN) (Derep). λ_{max} 258; 324; 380; 420 (MeOH) (Berdy).

Maiese, W.M. *et al.*, *J. Antibiot.*, 1989, **42**, 846-851 (*isol, props*)

Eur. Pat., 1990, 353381 (LL-E 19085α)

Carter, G.T. *et al.*, *J. Antibiot.*, 1990, **43**, 504-512 (*isol, pmr, cmr, struct*)

Carter, G.T. *et al.*, *J.C.S. Perkin I*, 1991, 2215-2219 (*biosynth*)

Citreamicin β C-474

[128999-30-4]

As Citreamicin α, C-473 with

R¹ = -COCH(CH₃)₂, R² = CH₃

C₃₅H₂₉NO₁₂ 655.614

Quinone antibiotic. Prod. by *Micromonospora citrea*. Sol. CHCl₃, Me₂CO, CH₂Cl₂; fairly sol. MeCN, MeOH, EtOH, EtOAc; poorly sol. H₂O, hexane. λ_{max} 223 (ε 42400); 255 (ε 38900); 320 (ε 32500); 384 (ε 9030); 410 (ε 6750) (MeCN) (Derep).

Carter, G.T. *et al.*, *J. Antibiot.*, 1990, **43**, 504-512 (*isol, pmr, cmr, struct*)

Citreamicin γ C-475

[128999-31-5]

As Citreamicin α, C-473 with

R¹ = Ac, R² = CH₃

C₃₃H₂₅NO₁₂ 627.56

Prod. by *Micromonospora citrea*. No phys. props. reported. Sol. CHCl₃, Me₂CO, CH₂Cl₂; fairly sol. MeCN, MeOH, EtOAc, EtOH; poorly sol. H₂O, hexane. λ_{max} 223 (ε 42400); 255 (ε 38900); 320 (ε 32500); 384 (ε 9030); 410 (ε 6750) (MeCN) (Derep). λ_{max} 258; 325; 385 (MeOH) (Berdy).

Carter, G.T. *et al.*, *J. Antibiot.*, 1990, **43**, 504-512 (*isol, pmr, cmr, struct*)

Citreamicin η C-476

[128969-88-0]

As Citreamicin α, C-473 with

R¹ = H, R² = CH₃


C₃₁H₂₃NO₁₁ 585.523

Quinone antibiotic. Prod. by *Micromonospora citrea*. Sol. CHCl₃, Me₂CO, CH₂Cl₂; fairly sol. MeCN, EtOAc, MeOH, EtOH; poorly sol. H₂O, hexane. [α]_D²⁵ -63 (c, 0.19 in DMSO). λ_{max} 223 (ε 42000); 255 (ε 39000); 320 (ε 32000); 384 (ε 9030); 410 (MeCN) (Derep). λ_{max} 223 (ε 37800); 302 (ε 21750); 321 (ε 21700) (MeOH) (Berdy).

Carter, G.T. *et al.*, *J. Antibiot.*, 1990, **43**, 504-512 (*isol, pmr, cmr, struct*)

Citreoazopyrone C-477


[190783-70-1]



C₁₃H₁₈N₂O₅ 282.296

Metab. of a hybrid strain derived from *Penicillium citreo-viride*. Topoisomerase I inhibitor. Oil. Shows small negative opt. rotn.

Kosemura, S. *et al.*, *Tet. Lett.*, 1997, **38**, 3025 (*isol, ir, pmr, cmr*)

Citridone A C-478

Relative Configuration


C₁₉H₁₉NO₂ 293.365

Prod. by *Penicillium* sp. FKI-1938. Potentiator of miconazole antifungal activity. Needles. Mp 172-175°. [α]_D²⁵ -1.6 (c, 0.1 in MeOH). λ_{max} 205 (ε 9800); 246 (ε 10700) (MeOH).

Fukuda, T. *et al.*, *J. Antibiot.*, 2005, **58**, 309-314; 315-321 (*isol, pmr, cmr, activity*)

Citridone B

C-479

Relative
ConfigurationC₁₉H₂₁NO₅ 343.379


Prod. by *Penicillium* sp. FKI-1938. Potentiator of miconazole antifungal activity. Needles. Mp 168-170°. $[\alpha]_D^{25} +102.4$ (c, 0.1 in MeOH). Isol. as a 3:2 mixt. with Citridone B' to which data refers. λ_{\max} 207 (ε 21000); 233 (ε 18900) (MeOH).

3-Epimer: Citridone B'C₁₉H₂₁NO₅ 343.379Prod. by *Penicillium* sp. FKI-1938.

Fukuda, T. *et al.*, *J. Antibiot.*, 2005, **58**, 309-314; 315-321 (*isol, pmr, cmr, activity*)

Citridone C

C-480

Relative
ConfigurationC₁₉H₂₁NO₄ 327.379


Prod. by *Penicillium* sp. FKI-1938. Potentiator of miconazole antifungal activity. Pale yellow amorph. solid. $[\alpha]_D^{25} -74.4$ (c, 0.1 in MeOH). λ_{\max} 203 (ε 16900); 233 (ε 11800) (MeOH).

Fukuda, T. *et al.*, *J. Antibiot.*, 2005, **58**, 309-314; 315-321 (*isol, pmr, cmr, activity*)

Citridone D

C-481

[929102-15-8]

Relative
ConfigurationC₁₉H₁₉NO₄ 325.363


Prod. by *Penicillium* sp. FKI-1938. Potentiator of miconazole antifungal activity. Pale yellow powder. $[\alpha]_D^{25} +73.5$ (c, 0.1 in MeOH). λ_{\max} 203 (ε 24000); 232 (ε 17000) (MeOH).

Fukuda, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 1659-1661 (*isol, pmr, cmr*)

Citridinadin A

C-482

[774233-49-7]



Absolute Configuration

C₃₅H₅₂N₄O₆ 624.819

Prod. by *Penicillium citrinum* (strain N-059) isol. from the alga *Actinotrichia fragilis*. Cytotoxic. Oil. $[\alpha]_D^{19} -17$ (c, 0.4 in MeOH). λ_{\max} 224 (ε 9000); 230 (sh); 249 (ε 9600); 265 (sh); 335 (ε 3100) (MeOH).

14-De(acyloxy): Citrinadin B

[869803-70-3]

C₂₈H₃₉N₃O₄ 481.634

Prod. by *Penicillium citrinum* (strain N-059). Pale yellow solid. $[\alpha]_D^{20} +8$ (c, 1 in MeOH). λ_{\max} 203 (ε 12500); 223 (ε 7600); 248 (ε 9800); 333 (ε 3100) (MeOH).

Tsuda, M. *et al.*, *Org. Lett.*, 2004, **6**, 3087-3089 (*isol, pmr, cmr*)

Mugishima, T. *et al.*, *J.O.C.*, 2005, **70**, 9430-9435 (*abs config*)

Citriodorol

C-483


Struct. unknown. Alkaloid from *Eucalyptus citriodora*. Antibacterial agent. Brownish-yellow powder. Mp 115-117°. $[\alpha]_D^{25} +12.5$ (c, 2 in MeOH).

Satwalekar, S.S. *et al.*, *J. Indian Inst. Sci.*, 1957, **39**, 195-212; *CA*, **52**, 1548f

Citropone A

C-484

[98496-33-4]

C₂₂H₁₉NO₆ 393.395


Alkaloid from the root bark of *Citrus grandis* f. *buntan* and *Citrus natsudaiddai* (Rutaceae). Red needles (Me₂CO). Mp 280-282°. Citropone A and Citropone B, C-485 are the first examples of naturally occurring homoacridone alkaloids. λ_{\max} 212 (ε 30900); 270 (ε 29500); 282 (sh) (ε 28800); 307 (sh) (ε 30200); 324 (ε 30900) (MeOH/NaOMe) (Derep). λ_{\max} 215 (ε 18200); 230 (sh) (ε 17800); 272 (ε 24500); 287 (sh) (ε 24000); 330 (sh) (ε 12300); 410 (ε 6920) (MeOH) (Derep).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1881 (*isol, pmr, cmr, uv, cryst struct*)

Citropone B

C-485

[98496-34-5]

C₂₃H₂₃NO₆ 409.438

Alkaloid from the root bark of *Citrus grandis* f. *buntan* (Rutaceae). Orange needles (Me₂CO). Mp 192-194°. λ_{\max} 210 (ε 40700); 252 (ε 22900); 325 (ε 30200) (MeOH/NaOMe) (Derep). λ_{\max} 217 (ε


28800); 230 (sh) (ε 25700); 265 (ε 25700); 309 (ε 19500); 330 (sh) (ε 16600); 407 (ε 9770) (MeOH) (Derep).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1881 (*isol, uv, pmr, cmr, cryst struct*)

Citropone C

C-486

7-Acetyl-6,7-dihydro-1,7-dihydroxy-3-methoxy-5H-cyclohepta[b]quinoline-8,11-dione, 9CI
[131815-28-6]

C₁₇H₁₅NO₆ 329.309


Alkaloid from the roots and stem bark of several hybrids of *Citrus* spp. (Rutaceae). Yellow prisms. Mp 198-200° dec. $[\alpha]_D^{25} +7.14$ (c, 0.056 in CHCl₃). λ_{\max} 206; 248; 283; 335; 341; 396 (MeOH) (Derep).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1881 (*isol, uv, pmr, cmr, struct*)

Cividiclavine

C-487

6,6',8,8'-Tetramethyl-1,13'-biergolin-14-ol, 9CI
[84826-93-7]


C₃₂H₃₈N₄O 494.678Prod. by *Penicillium citreo-viride*.

Vining, L.C. *et al.*, *FEMS Symp.*, 1982, **13**, 243-251; *CA*, **98**, 103970u (*isol, struct*)

Cladioxazole

C-488

[602297-42-7]

Relative
ConfigurationC₁₆H₂₇NO 249.395

Alkaloid from the soft coral *Cladiella* sp. Oil.

Ata, A. *et al.*, *Tet. Lett.*, 2003, **44**, 6951-6953 (*isol, pmr, cmr*)

Clathculine A

C-489

H₂C=CHCH₂CH₂CH=CHC≡C(CH₂)₉NMeCH₂CH₂NHMe

C₂₁H₃₈N₂ 318.545

(Z)-form [300811-99-8]

Isol. from the sponge *Clathrina* aff. *reticulum*. Isol. as a mixt. with Clathru-line B. λ_{\max} 234 (ϵ 6300); 246 (ϵ 7600); 285 (ϵ 2800) (EtOH).

16,17-Dihydro: Clathraline B

[300812-19-5]


C₂₁H₄₀N₂ 320.56Isol. from *Clathrina* aff. *reticulum*.

Rudi, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1434-1436 (*isol, pmr, cmr, uv*)

Clathramide A

C-490

[182888-51-3]

C₁₃H₁₇BrN₄O₃ 357.206

Neutral form shown. Exists as zwitterion with +ve charge distributed onto the two imidazole N's. Alkaloid from the Caribbean sponge *Agelas clathrodes*. Exhibits moderate antifungal activity. Amorph. solid. $[\alpha]_D^{25}$ -5 (c, 0.001 in MeOH). λ_{\max} 208 (ϵ 14900); 230 (ϵ 13000); 265 (ϵ 10900) (MeCN).

N-De-Me: Clathramide C

[200264-71-7]

C₁₂H₁₅BrN₄O₃ 343.18

Alkaloid from *Agelas dispar*. Amorph. solid. $[\alpha]_D^{25}$ -6 (c, 0.001 in MeOH). λ_{\max} 208 (ϵ 14900); 230 (ϵ 13000); 265 (ϵ 10900) (MeCN).

8-Epimer: Clathramide B

[183072-78-8]

C₁₃H₁₇BrN₄O₃ 357.206

From *Agelas clathrodes*. Exhibits moderate antifungal activity. Amorph. solid. $[\alpha]_D^{25}$ +11 (c, 0.001 in MeOH). λ_{\max} 208; 230 (ϵ 13000); 265 (ϵ 10900) (MeCN).

8-Epimer, N-de-Me: Clathramide D

[200264-72-8]

C₁₂H₁₅BrN₄O₃ 343.18

Alkaloid from *Agelas dispar*. Serotonin receptor antagonist. λ_{\max} 208 (ϵ 14900); 230 (ϵ 13000); 265 (ϵ 10900) (MeCN) (Berdy).


Cafieri, F. *et al.*, *Tetrahedron*, 1996, **52**, 13713-13720 (*isol, uv, ir, pmr, cmr, struct*)

Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 122-125 (*isol, uv, ir, pmr, cmr, ms*)

Clathridine A

C-491

[122759-55-1]

C₁₆H₁₅N₅O₄ 341.326

Metab. from the marine sponge *Clathrina clathrus* and the nudibranch *Notodoris gardineri*. Cryst. (CHCl₃). Fairly sol. MeOH; poorly sol. butanol, hexane. Mp

260-262° dec. λ_{\max} 285 (ϵ 5400); 371 (ϵ 14700) (MeOH) (Derep). λ_{\max} 284; 359 (MeOH/NaOH) (Berdy).

Zn complex: Clathridine Zn

[122780-90-9]

C₃₂H₂₈N₁₀O₈Zn 746.025

Metab. of *Clathrina clathrus* and a *Leucetta* sp. Yellow microcryst. (Et₂O/CHCl₃). Fairly sol. MeOH; poorly sol. butanol, hexane. Mp 158-160°. λ_{\max} 285 (ϵ 9500); 361 (ϵ 22700); 372 (ϵ 22500); 399 (sh) (ϵ 13300) (MeOH) (Derep).

Ciminiello, P. *et al.*, *Tetrahedron*, 1989, **45**, 3873-3878 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ciminiello, P. *et al.*, *Tetrahedron*, 1990, **46**, 4387-4392 (*Zn-Clathridine, cryst struct*)


Alvi, K.A. *et al.*, *Tetrahedron*, 1993, **49**, 329-336 (*isol*)

Ohta, S. *et al.*, *Tet. Lett.*, 2000, **41**, 4623-4627 (*synth*)

Clathridine B

C-492

[146556-28-7]

C₂₄H₂₁N₅O₆ 475.46


Metab. from the nudibranch *Notodoris gardineri*. Yellow oil. λ_{\max} 285; 377 (MeOH).

Alvi, K.A. *et al.*, *Tetrahedron*, 1993, **49**, 329 (*isol, uv, pmr, cmr, ms, struct*)

Clathridine C

C-493

[152273-86-4]


C₁₅H₁₅N₅O₃ 313.315

Alkaloid from the sponge *Leucetta* sp. Weakly cytotoxic. Yellow plates (DMSO). Mp 255-257°. Related to Clathridine A, C-491.

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1993, **46**, 1229 (*isol, uv, ir, pmr, cmr, ms, struct*)

Clathridine-9-N-(2-sulfoethyl)imine

C-494

C₁₈H₂₀N₆O₆S 448.459


Alkaloid from the marine sponge *Leucetta microraphis*. Yellow needles. Mp 275-277° dec. λ_{\max} 232 (ϵ 9530); 278 (ϵ 8610); 382 (ϵ 13700) (MeOH) (Derep).

He, H.-Y. *et al.*, *J.O.C.*, 1992, **57**, 2176-2178 (*isol, uv, ir, pmr, cmr, cryst struct*)

Clathrotropine

C-495

[502686-87-5]

C₁₇H₂₄N₂O₃ 304.388


Alkaloid from the bark of *Clathrotropis glaucophylla*. Cryst. $[\alpha]_D^{25}$ -146 (c, 0.1 in EtOH). λ_{\max} 205 (log ϵ 2.67); 234 (log ϵ 2.79); 309 (log ϵ 2.87) (MeOH).

Sagen, A.-L. *et al.*, *Phytochemistry*, 2002, **61**, 975-978 (*isol, pmr, cmr, ms*)

Clathrymine A

C-496

6-Carboxy-1,2,3,4-tetrahydro-7-phenylquinolinolizinium inner salt, 9CI
[175669-25-7]

C₁₆H₁₆NO₂⁺ 254.308

Exists as zwitterion (COO⁻). Alkaloid from the Indo-Pacific sponge *Clathria basilana*. Viscous amber oil. Readily decarboxylates to Clathrymine B. λ_{\max} 241 (ϵ 5500); 291 (ϵ 4500) (MeOH).

Decarboxy: 1,2,3,4-Tetrahydro-1-phenylquinolinolizinium(1+), 9CI. Clathrymine B

[175669-26-8]

C₁₅H₁₆N⁺ 210.298

Viscous brown oil. Counterion not specified.


Sperry, S. *et al.*, *Tet. Lett.*, 1996, **37**, 2389-2390 (*isol, uv, ir, pmr, cmr, struct*)

Daab, J.C. *et al.*, *Monatsh. Chem.*, 2003, **134**, 573-583 (*Clathrymine B, synth*)

Clathrynamide A

C-497

11-Bromo-3,12-dihydroxy-2,2,4,10-tetramethyl-4,6,8,10,14,16-eicosahexaen-18-ynamide
[160638-12-0]



Absolute Configuration

C₂₄H₃₂BrNO₃ 462.426

Isol. from the marine sponges *Clathria* sp. and *Psammoclemma* sp. Antifungal agent. Pale yellow oil. [α]_D²³ +149 (c, 0.022 in MeOH). λ_{max} 247 (ε 19500); 309 (ε 29500); 321 (ε 38000); 336 (ε 28800) (MeOH).

N-(4-Hydroxy-1-methylpentyl): **Clathrynamide B**

[152273-73-9]

C₃₀H₄₄BrNO₄ 562.586

Isol. from a *Clathria* sp. Gum. [α]_D²⁵ +76 (c, 0.0033 in MeOH). λ_{max} 267 (ε 17400); 307 (ε 23400); 321 (ε 30200); 337 (ε 23400) (no solvent reported).

N-(1-Methyl-4-oxopentyl): **Clathrynamide C**

[152273-74-0]

C₃₀H₄₂BrNO₄ 560.57

Isol. from a *Clathria* sp. Gum. λ_{max} 268 (ε 25700); 307 (ε 36300); 321 (ε 38000); 337 (ε 29500) (no solvent reported).

Debromo: **Debromoclathrynamide A**

[602306-79-6]

C₂₄H₃₃NO₃ 383.53

Isol. from a *Psammoclemma* sp. Pale yellow oil. [α]_D²³ +159 (c, 0.02 in MeOH). λ_{max} 263 (sh) (log ε 4.4); 272 (log ε 4.42); 283 (log ε 4.36); 297 (log ε 4.53); 310 (log ε 4.64); 324 (log ε 4.54) (MeOH).

6E-Isomer: **(6E)-Clathrynamide A**

[603129-66-4]

C₂₄H₃₂BrNO₃ 462.426

Isol. from a *Psammoclemma* sp. Pale yellow oil. [α]_D²³ -58 (c, 0.02 in MeOH). λ_{max} 268 (log ε 3.84); 294 (sh) (log ε 3.83); 308 (log ε 4.09); 322 (log ε 4.24); 338 (log ε 4.16) (MeOH).

(4E,6E)-Isomer, debromo: **(4E,6E)-Debromoclathrynamide A**

[602302-49-8]

C₂₄H₃₃NO₃ 383.53

Isol. from a *Psammoclemma* sp. Pale yellow oil. [α]_D²⁴ -80 (c, 0.004 in MeOH). λ_{max} 272 (log ε 4.02); 284 (log ε 3.99); 298 (log ε 4.11); 312 (log ε 4.22); 327 (log ε 4.15) (MeOH).


Ohta, S. et al., *Tet. Lett.*, 1993, **34**, 5935-5938 (isol)

Ojika, M. et al., *Biosci., Biotechnol., Biochem.*, 2003, **67**, 1568-1573 (abs config, *Psammoclemma constits*)

Clausamine B

C-498

[205116-60-5]



C₁₉H₁₇NO₃ 307.348

(±)-form

Alkaloid from *Clausena anisata*. Oil. λ_{max} 205 ; 222 ; 237 ; 248 ; 269 ; 278 (sh) ; 310 (sh) ; 321 ; 334 (MeOH).

O-De-Me: **Clausamine A**

[205116-59-2]

C₁₈H₁₅NO₃ 293.321

Alkaloid from *Clausena anisata*. Pale yellow powder. λ_{max} 204 ; 222 ; 238 ; 248 ; 269 ; 278 (sh) ; 310 (sh) ; 322 ; 335 (MeOH).

1',2'-Dihydro, 1'-hydroxy: **Clausamine C**

[205116-61-6]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Clausena anisata*. Pale yellow oil. λ_{max} 204 ; 223 ; 238 ; 249 ; 270 ; 278 (sh) ; 312 (sh) ; 322 ; 336 (MeOH).


Ito, C. et al., *Chem. Pharm. Bull.*, 1998, **46**, 344-346 (*Clausamines A - C*)

Lebold, T.P. et al., *Org. Lett.*, 2008, **10**, 997-1000 (synth)

Clausamine F

C-499

[256222-02-3]



C₁₉H₁₉NO₄ 325.363

Alkaloid from *Clausena anisata*. Pale yellow oil. λ_{max} 223 ; 242 ; 272 ; 324 ; 341 (MeOH).

1-Me ether: **Clausamine E**

[256222-01-2]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Clausena anisata*. Oil. λ_{max} 224 ; 239 ; 273 ; 322 ; 337 (MeOH).

3'-Hydroperoxide, 1-Me ether: **Clausamine G**

[256222-03-4]

C₂₀H₂₁NO₅ 355.39


Alkaloid from *Clausena anisata*. Pale yellow oil. λ_{max} 223 ; 236 ; 270 ; 313 ; 321 ; 333 (MeOH).

Ito, C. et al., *J. Nat. Prod.*, 2000, **63**, 125-128 (*Clausamines E-G*)

Clausaniline

C-500

2-Hydroxy-7-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-2-hydroxy-7-prenylcarbazole [58523-27-6]



C₁₈H₁₇NO₂ 279.338


Alkaloid from the roots of *Clausena anisata* (Rutaceae). Silky yellow cryst. Mp 154-156°. Isomer of Heptaphylline, H-125.

Okorie, D.A. et al., *Phytochemistry*, 1975, **14**, 2720 (isol, uv, ir, pmr, ms, struct)

ζ-Clausenamide

C-501

5,6-Dihydro-5-hydroxy-3-methyl-6-phenyl-3-benzazocin-4(3H)-one, 9CI [136173-85-8]



C₁₈H₁₇NO₂ 279.338

Alkaloid from *Clausena lansium* (wampee) (Rutaceae).


Yang, M. et al., *Chin. Chem. Lett.*, 1991, **2**, 291; *CA*, **115**, 155059 (isol, struct)

Yang, L. et al., *Org. Lett.*, 2007, **9**, 1387-1390 (synth)

Clausenamide

C-502

3-Hydroxy-5-(hydroxyphenylmethyl)-1-methyl-4-phenyl-2-pyrrolidinone, 9CI. *Lansimide 4†* [119616-14-7]



C₁₈H₁₉NO₃ 297.353

λ_{max} 257 (ε 500) (MeOH) (Berdy).

(±)-form [103541-15-7]

Isol. from the leaves of *Clausena lansium* (wampee) (Rutaceae). Needles. Mp 239-240°.

Di-Ac:

Cubes. Mp 165-167°.

1'-Me ether: **Lansimide 3†**

[119518-25-1]

C₁₉H₂₁NO₃ 311.38

Constit. of *Clausena lansium* (wampee). Spasmodic agent.

1',5-Diepimer: **Neoclausenamide**

[114528-82-4]

Isol. from *Clausena lansium* (wampee) (Rutaceae). Cubes. Mp 205-216°.

Lakshmi, V. et al., *Indian Drugs*, 1988, **26**, 105 (*Lansimides*)

Yang, M. et al., *Phytochemistry*, 1988, **27**, 445 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Rao, E.C. et al., *Chin. Chem. Lett.*, 1994, **5**, 267-268 (synth)

Huang, D.F. et al., *Chin. Chem. Lett.*, 1994, **5**, 371-372 (synth)


Wang, J.-Q. et al., *J.C.S. Perkin 1*, 1996, 209-212 (synth, cryst struct, *Neoclausenamide*)

Cappi, M.W. et al., *Chem. Comm.*, 1998, 1159-1160 (synth)

Yang, L. *et al.*, *Org. Lett.*, 2008, **10**, 2461-2464 (synth)

Clausenamine A C-503

6,6',8,8'-Tetramethoxy-3,3'-dimethyl-[2,2'-bi-9H-carbazole]-1,1'-diol. 2,2'-Bi(1-hydroxy-6,8-dimethoxy-3-methylcarbazole) [184427-49-4]



C₃₀H₂₈N₂O₆ 512.561
Alkaloid from stem bark of *Clausena excavata*. Shows antiplasmodial activity. Granules. Mp 278-280°. [α]_D²¹ +142.2 (c, 0.75 in CHCl₃). λ_{max} 218 (sh); 235; 257 (sh); 307; 343 (sh) (MeOH).


Wu, T.-S. *et al.*, *Tet. Lett.*, 1996, **37**, 7819-7822 (isol, uv, ir, pmr, cmr, ms, struct)

Zhang, A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1021-1023 (synth)

Lin, G. *et al.*, *Tetrahedron*, 2000, **56**, 7163-7171 (synth, activity)

Clausenapin C-504

1-Methoxy-3-methyl-2-(3-methyl-2-butenyl)-9H-carbazole, 9CI [54313-57-4]




C₁₉H₂₁NO 279.381
Alkaloid from the leaves of *Clausena heptaphylla* (Rutaceae). Cryst. (hexane). Mp 101°.

Joshi, B.S. *et al.*, *Indian J. Chem.*, 1974, **12**, 437 (synth)

Bhattacharyya, P. *et al.*, *Chem. Ind. (London)*, 1984, 301 (isol, uv, ir, pmr)

Clausenaquinone A C-505


7-Hydroxy-3-methoxy-6-methyl-1H-carbazole-1,4-(9H)-dione, 9CI [159959-56-5]



C₁₄H₁₁NO₄ 257.245
Alkaloid from stem bark of *Clausena excavata* (Rutaceae). Cytotoxic. Platelet aggregation inhibitor. Brown powder. Mp 290°. λ_{max} 200; 227; 265; 294; 320; 419 (MeOH) (Berdy).

Wu, T.-S. *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 2395 (isol, pmr, uv, activity)

Clausevatine E C-506
[215377-69-8]



C₁₈H₁₇NO₅ 327.336
Alkaloid from the root bark of *Clausena excavata*. Yellowish granules. Mp 208-212°. [α]_D -92.4 (c, 0.05 in MeOH). λ_{max} 213 (sh); 222; 240; 251; 271; 282; 315 (sh); 325; 339 (MeOH).

1'-Epimer: **Clausevatine F** [215377-70-1]
C₁₈H₁₇NO₅ 327.336
Alkaloid from the root bark of *Clausena excavata*. Granules. Mp 162-164°. [α]_D -199 (c, 0.02 in MeOH). λ_{max} 202; 222; 241; 252; 271; 281; 314 (sh); 326; 339 (MeOH).


1'-Deoxy: **Clausevatine D** [215377-68-7]
C₁₈H₁₇NO₄ 311.337
Alkaloid from the root bark of *Clausena excavata*. Yellow granules. Mp 241-244°. [α]_D -5.7 (c, 0.93 in MeOH). λ_{max} 202; 223; 241; 251; 272; 278 (sh); 325; 340 (MeOH).

1'-Deoxy, 1',2'-didehydro: **Clausevatine G** [215377-72-3]
C₁₈H₁₅NO₄ 309.321
Alkaloid from the root bark of *Clausena excavata*. Yellow granules. Mp >280°. λ_{max} 207; 223; 264 (sh); 274; 287; 338; 353 (MeOH).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1459-1461 (Clausevatines D-G)
Lebold, T.P. *et al.*, *Org. Lett.*, 2008, **10**, 997-1000 (synth)

Clausine D C-507

1-Hydroxy-4-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde. 3-Formyl-1-hydroxy-4-prenylcarbazole [142846-95-5]



C₁₈H₁₇NO₂ 279.338
Alkaloid from stem bark of *Clausena excavata* (Rutaceae). Exhibits significant antiplatelet aggregation activity. Brown powder. Mp 300°. λ_{max} 225 (ε 18600); 242 (ε 21400); 253 (sh) (ε 18600); 276 (ε 25100); 289 (ε 19100); 351 (ε 10000) (possibly MeOH, not reported) (Derep).

Me ether: **Ekeberginine**. 1-Methoxy-4-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-1-methoxy-4-prenylcarbazole [101242-43-7]
C₁₉H₁₉NO₂ 293.365
Alkaloid from the stem bark of *Ekebergia senegalensis* (Meliaceae). Mp 230-231°. Spectral data reported refer to the N-Me deriv.

Me ether, N-Me: Mp 155-157°.

Carboxylic acid, Me ester: **Clausine F** [142846-96-6]
C₁₉H₁₉NO₃ 309.364
Alkaloid from the stem bark of *Clausena excavata* (Rutaceae). Exhibits significant antiplatelet aggregation activity. Powder. Mp 200-202°. λ_{max} (solvent not reported) (Derep). λ_{max} 225 (sh) (ε 20400); 244 (sh) (ε 28200); 251 (sh) (ε 28800); 269 (ε 38900); 313 (ε 7950); 323 (ε 7240); 340 (ε 5890) (possibly MeOH, not reported) (Derep).

Carboxylic acid, Me ether, Me ester: **Clausamine D** C₂₀H₂₁NO₃ 323.391
Alkaloid from *Clausena anisata*. Powder. λ_{max} 222; 240; 248; 256; 269; 310; 321; 335 (MeOH).


Lontsi, D. *et al.*, *Tet. Lett.*, 1985, **26**, 4249-4252 (Ekeberginine)

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 1069-1071 (Clausines)

Ito, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 125-128 (Clausamine D)

Clausine W C-508

[192706-30-2]



Relative Configuration


C₁₈H₁₇NO₅ 327.336
Alkaloid from the root bark of *Clausena excavata* (Rutaceae). Yellow cryst. (Me₂CO). Mp 178-179°. [α]_D -3.4 (c, 1 in MeOH). λ_{max} 202 (log ε 4.35); 224 (sh) (log ε 4.34); 237 (sh) (log ε 4.45); 243 (log ε 4.48); 250 (sh) (log ε 4.47); 287 (sh) (log ε 4.48); 301 (log ε 4.65); 349 (log ε 4.04) (MeOH).

4'-Deoxy: **Clausine T** [192706-49-3]
C₁₈H₁₇NO₄ 311.337
Alkaloid from *Clausena excavata* (Rutaceae). Yellowish cryst. (Me₂CO). Mp 155-158°. [α]_D -82.1 (c, 0.03 in MeOH). λ_{max} 202; 242; 249; 274; 285 (sh); 302; 345 (MeOH).

Wu, T.-S. *et al.*, *Heterocycles*, 1997, **45**, 969-973 (isol, uv, ir, pmr, cmr)

Clauszoline A

[185508-02-5]



C₂₃H₂₃NO₃ 361.44

Alkaloid from stem bark of *Clausena excavata*. Pale yellow powder. λ_{max} 204 ; 240 ; 265 ; 306 ; 375 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2231 (*isol, uv, ir, pmr, ms, struct*)

Clauszoline B

[185508-03-6]

As Clauszoline A, C-509 with

R = H


C₁₈H₁₅NO₃ 293.321

Alkaloid from stem bark of *Clausena excavata*. Yellow oil. λ_{max} 204 ; 242 ; 252 (sh) ; 264 ; 291 (sh) ; 304 ; 376 (sh) (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2231 (*isol, uv, ir, pmr, ms, struct*)

Clauszoline H

2,11-Dihydro-9-methoxy-2,2,8-trimethylpyrano[2,3-a]carbazole, 9CI [187110-70-9]




C₁₉H₁₉NO₂ 293.365

Alkaloid from *Clausena excavata*. Powder. λ_{max} 230 ; 258 ; 288 ; 310 ; 323 ; 348 ; 363 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 48-52 (*isol, uv, pmr*)

Clavamine

[129121-68-2]



C₃₅H₆₀N₁₂O₉ 792.934

Isol. from the venom of the Joro spider *Nephila clavata*. Shows insecticidal props. [α]_D¹⁵ -12.6 (c, 0.94 in H₂O).


Yoshioka, M. *et al.*, *Biog. Amines*, 1990, **7**, 375 (*isol*)

Teshima, T. *et al.*, *Tetrahedron*, 1990, **46**, 3819 (*synth*)

C-509

Clavepictine B

6-(1,3-Decadienyl)octahydro-4-methyl-2H-quinolizin-3-ol, 9CI [132621-84-2]



Absolute Configuration

C₂₀H₃₅NO 305.503

Alkaloid from the tunicate *Clavelina picta*. Cytotoxic agent. Cryst. Mp 70-72°. [α]_D +27.1 (c, 0.03 in CH₂Cl₂). λ_{max} 230 (ε 18000) (EtOH) (Derep).

O-Ac: **Clavepictine A**

[132621-83-1]

C₂₂H₃₇NO₂ 347.54

Alkaloid from *Clavelina picta*. Cytotoxic agent. Oil. [α]_D -75.6 (c, 0.7 in CH₂Cl₂). λ_{max} 230 (ε 19000) (EtOH) (Derep).

Raub, M.F. *et al.*, *J.A.C.S.*, 1991, **113**, 3178-3180 (*isol, struct*)

Toyooka, N. *et al.*, *J.O.C.*, 1996, **61**, 4882-4883 (*synth*)

Ha, J.D. *et al.*, *J.O.C.*, 1997, **62**, 4550-4551 (*synth*)


Ha, J.D. *et al.*, *J.A.C.S.*, 1999, **121**, 10012-10020 (*synth*)

Toyooka, N. *et al.*, *Tetrahedron*, 1999, **55**, 15209-15224 (*synth, abs config*)

Xu, S. *et al.*, *Org. Lett.*, 2006, **8**, 3179-3182 (*synth*)

Clavicipitic acid I

trans-Clavicipitic acid [33062-26-9]



Absolute Configuration

C₁₆H₁₈N₂O₂ 270.33

Prod. by *Claviceps* strain SD58 and by *Claviceps fusiformis* strain 139/2/IG. Plates (EtOH). Mp 264° (262° dec.). Isol. only as a mixt. with Clavicipitic acid II, approx. 1:1 from *C. fusiformis* and with Clavicipitic acid I predominating in the case of *C.* strain SD58.

10-Epimer: **Clavicipitic acid II**. cis-Clavicipitic acid [72690-85-8]

C₁₆H₁₈N₂O₂ 270.33

Prod. by *Claviceps fusiformis* and *Claviceps* strain SD58 (see above).

Robbers, J.E. *et al.*, *Tet. Lett.*, 1969, 1857 (*isol, uv, ir, pmr, ms, biosynth*)

Bajwa, R.S. *et al.*, *Phytochemistry*, 1975, **14**, 735 (*biosynth*)

King, G.S. *et al.*, *J.C.S. Perkin 1*, 1977, 2099 (*isol, pmr, ms, cd, struct*)

Robbers, J.E. *et al.*, *J.O.C.*, 1980, **45**, 1117 (*cryst struct, biosynth*)

C-514

Muratake, H. *et al.*, *Heterocycles*, 1983, **20**, 1963 (*synth, ir, pmr*)

Kozikowski, A.P. *et al.*, *J.O.C.*, 1984, **49**, 2310 (*synth, uv, ir, ms*)

Boyles, D.A. *et al.*, *J.O.C.*, 1988, **53**, 5128 (*synth*)

Semmelhack, M.F. *et al.*, *Tet. Lett.*, 1993, **34**, 5051 (*synth*)

Somei, M. *et al.*, *Heterocycles*, 1994, **37**, 719 (*synth*)

Iwao, M. *et al.*, *Tetrahedron*, 1997, **53**, 51 (*synth*)

Shinohara, H. *et al.*, *Tetrahedron*, 1999, **55**, 10989-11000 (*synth*)


Yokoyama, Y. *et al.*, *Eur. J. Org. Chem.*, 2004, 1244-1253 (*synth*)

Ku, J.-M. *et al.*, *J.O.C.*, 2007, **72**, 8115-8118 (*synth*)

Clavizepine

[142697-01-6 ((±)-form Me ether), 154968-81-7 ((±)-form)]

C-515



C₁₉H₂₁NO₄ 327.379

λ_{max} 209 (ε 40700); 262 (ε 6760); 306 (ε 6460) (MeOH/NaOH) (Derep). λ_{max} 207 (ε 25100); 250 (ε 2510); 295 (ε 4170) (MeOH) (Derep).

(S)-form [109305-77-3]

Isol. from *Corydalis claviculata* (Papaveraceae). Cryst. (MeOH). Mp 234-235°. [α]_D -40 (c, 0.05 in MeOH). The first naturally occurring dibenzopyranazepine alkaloid.

Boente, J.M. *et al.*, *Tet. Lett.*, 1986, **27**, 4077 (*uv, pmr, cmr, ms, struct*)


Vásquez, R. *et al.*, *Synlett*, 1994, 433 (*synth*)

Ishibashi, H. *et al.*, *Tetrahedron*, 1994, **50**, 10215 (*synth*)

De la Fuente, M.C. *et al.*, *J.O.C.*, 1996, **61**, 5818 (*synth*)

Clavosine A

C-516



C₆₀H₁₀₁N₄O₂₀P 1229.447

Isol. from the sponge *Myriastra clavosa*. Cytotoxic agent and inhibitor of protein phosphatases. Powder. [α]_D -5 (c, 0.36 in CH₂Cl₂). λ_{max} 228 (ε 20920); 332 (ε 23270) (MeOH).

(2E)-Isomer: Clavosine BC₆₀H₁₀₁N₄O₂₀P 1229.447


Isol. from *Myriastra clavosa*. Cytotoxic agent and inhibitor of protein phosphatases. Powder. $[\alpha]_D^{25}$ -3.2 (c, 0.62 in CH₂Cl₂). λ_{\max} 228 (ε 18570); 328 (ε 23000) (MeOH).

(2E,6Z)-Isomer: Clavosine CC₆₀H₁₀₁N₄O₂₀P 1229.447

Isol. from *Myriastra clavosa*. Cytotoxic agent and inhibitor of protein phosphatases. Powder. $[\alpha]_D^{25}$ -31.7 (c, 0.12 in CH₂Cl₂). λ_{\max} 228 (ε 20300); 318 (ε 19960) (MeOH).

Fu, X. *et al.*, *J.O.C.*, 1998, **63**, 7957-7963**Clavulazine****C-517**


7,8-Dihydro-5H-pyrano[3,4-b]pyrazine-7-methanol, 9CI
[220525-28-0]

C₈H₁₀N₂O₂ 166.179**(S)-form**


Alkaloid from the soft coral *Clavularia viridis*. Needles (EtOAc/hexane). Mp 101-102°. $[\alpha]_D^{25}$ -99.4 (c, 0.17 in CHCl₃). λ_{\max} 272 (log ε 3.89); 277 (log ε 3.89); 310 (log ε 3.06) (EtOH).

Watanabe, K. *et al.*, *Heterocycles*, 1998, **49**, 269-274 (isol, pmr, cmr)Shen, Y.-C. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2004, **51**, 1421-1424 (isol, pmr, cmr, ms)Kumar, S.P. *et al.*, *Tet. Lett.*, 2006, **47**, 7149-7151 (synth)**Clavulazol A****C-518**

[851591-69-0]

C₈H₁₂N₂O₃ 184.194

Alkaloid from the soft coral *Clavularia viridis*. Amorph. solid. $[\alpha]_D^{25}$ -22.8 (c, 1 in MeOH). λ_{\max} 277 (log ε 3.89); 311 (log ε 3.01) (no solvent reported).


Shen, Y.-C. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2004, **51**, 1421-1424 (isol, pmr, cmr, ms)**Clavulazol B****C-519**C₈H₁₀N₂O₂ 166.179**(-)-form** [851591-70-3]

Alkaloid from the soft coral *Clavularia viridis*. Amorph. solid. $[\alpha]_D^{25}$ -58.9 (c, 1 in MeOH). λ_{\max} 272 (log ε 3.88); 278 (log

ε 3.86); 309 (log ε 3.29) (no solvent reported).

Shen, Y.-C. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2004, **51**, 1421-1424 (isol, pmr, cmr, ms)**Clavulinin**


[185559-59-5]

C-520C₁₆H₂₁NO₄ 291.346

Constit. of *Clavularia inflata*. Needles. Mp 190-191°.

Su, J. *et al.*, *Chin. Sci. Bull.*, 1996, **41**, 1877-1880; *CA*, **126**, 87252a (isol, pmr, cmr)**Clazamycin****C-521**

2-Chloro-2,3-dihydro-5-imino-1H-pyrrolizin-7a(5H)-ol

**(2S,7aR)-form**C₇H₉ClN₂O 172.614

Exists in aq. soln. as a mixt. of two epimers, Clazamycin A and B, the ratio of which is pH dependent.

(2S,7aR)-form**Clazamycin A**

[71806-55-8]

Isol. from *Streptomyces* MF 990-BF4.

Antibiotic with weak antibacterial activity, inhibits leukaemia L-1210 in mice.

Cryst. (as hydrochloride). Sol. H₂O, MeOH. Mp 110° dec. (hydrochloride).

$[\alpha]_D^{25}$ -56 (c, 1.0 in H₂O). λ_{\max} 212 (ε 11800); 250 (ε 2600) (HCl SALT/H₂O) (Derep). λ_{\max} 212 (ε 11800); 250 (ε 2600) (H₂O) (Berdy).

▶ LD₅₀ (mus, ivn) 50 - 100 mg/kg.**(2S,7aS)-form****Clazamycin B. U 54702D. U 54703.***Antibiotic 354. Antibiotic U 54702D.**Antibiotic U 54703*

[71774-49-7]

Isol. from *Streptomyces* MF 990-BF4and *Streptomyces puniceus doliceus*. Possesses weak antibacterial activity. Hygroscopic powder (as hydrochloride). $[\alpha]_D^{24}$

+96 (c, 1.0 in H₂O). λ_{\max} 212 (ε 11800); 250 (ε 2600) (HCl SALT/H₂O) (Derep).

▶ UY8427350


[69136-00-1, 68859-24-5]

Horiuchi, Y. *et al.*, *J. Antibiot.*, 1979, **32**, 762Nakamura, H. *et al.*, *J. Antibiot.*, 1979, **32**, 765 (cryst struct)Dolak, L.A. *et al.*, *J. Antibiot.*, 1980, **33**, 83 (struct)Hori, M. *et al.*, *J. Antibiot.*, 1984, **37**, 260 (props)Buechter, D.D. *et al.*, *J. Nat. Prod.*, 1987, **50**,

360

Cleavamine**C-522**

5-Ethyl-1,4,7,8,9,10-hexahydro-2H-3,7-methanoazacycloundecino[5,4-b]indole, 9CI

C₁₉H₂₄N₂ 280.412**(R)-form** [1674-01-7]

Degradn. prod. of Catharanthine, C-195, Vinblastine, V-110 and Leurosine, L-162. Needles (MeOH). Mp 117-119°. $[\alpha]_D^{26}$ +73 (CHCl₃).

Methodide:

Cryst. (MeOH/Et₂O/petrol). Mp 244-245° dec.

15,20-Dihydro: (+)-Dihydrocleavamine

[1966-42-3]

C₁₉H₂₆N₂ 282.428Alkaloid from *Pandaca boiteau*(Apocynaceae). Mp 135°. $[\alpha]_D^{25}$ +68 (CHCl₃).**15,20-Dihydro, 20S-hydroxy: (+)-Velbanamine**

[29951-44-8]

Fission prod. of Vinblastine, V-110 and Vincristine, V-128. $[\alpha]_D^{25}$ +56.2 (CHCl₃).

(S)-form [57550-18-2]

Obt. by dehydration of Capuronine, C-104. $[\alpha]_D^{20}$ -19 (c, 1 in CHCl₃).

15,20-Dihydro: (-)-Dihydrocleavamine

[1805-75-0]

C₁₉H₂₆N₂ 282.428Alkaloid from *Pandaca boiteau*(Apocynaceae). Noncryst. $[\alpha]_D^{25}$ -87 (CHCl₃).**15,20-Dihydro, 20R-hydroxy: Velbanamine**

[62246-22-4]

[34964-89-1, 19637-92-4, 11034-65-4]

C₁₉H₂₆N₂O 298.427

Trace alkaloid from leaves and twigs of

Tabernaemontana eglanulosa(Apocynaceae). Mp 139-141° (125-130°) (synthetic). $[\alpha]_D^{25}$ -49.3 (CHCl₃).**(±)-form** [31679-35-3]

Mp 113-114.5°.

Cameran, N. *et al.*, *Acta Cryst.*, 1964, **17**, 384 (cryst struct)Kutney, J.P. *et al.*, *Can. J. Chem.*, 1965, **43**, 1545 (synth, uv, ir, pmr)Gorman, M. *et al.*, *J.A.C.S.*, 1965, **87**, 93 (synth)Büchi, G. *et al.*, *J.A.C.S.*, 1968, **92**, 999

(Velbanamine, synth, bibl)

Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1975, **58**,

1672 (synth)

Chardon-Loriaux, I. *et al.*, *Tet. Lett.*, 1975,

1845 (synth)

Wenkert, E. *et al.*, *Helv. Chim. Acta*, 1976, **59**,

2711 (cmr)

Wenkert, E. *et al.*, *Heterocycles*, 1979, **12**, 1439

(cmr)

Takano, S. *et al.*, *Tet. Lett.*, 1980, 3697


(Velbanamine, synth)

Imanishi, T. *et al.*, *Chem. Pharm. Bull.*, 1981,**29**, 901; 1983, **31**, 1183 (synth, uv, pmr, ms)Takano, S. *et al.*, *Chem. Lett.*, 1982, 733

(synth)


Van Beek, T.A. *et al.*, *Tetrahedron*, 1984, **40**, 737 (*Velbanamine, isol, uv, ms*)
 Danieli, B. *et al.*, *Tet. Lett.*, 2000, **41**, 3489-3492 (*synth, Dihydrocleavamine*)
 Kanada, R.M. *et al.*, *Tet. Lett.*, 2001, **42**, 7311-7313 (*Dihydrocleavamine, synth*)
 Amat, M. *et al.*, *Org. Lett.*, 2003, **5**, 3139-3142 (*Dihydrocleavamine, synth*)

Clemaïne C-523
 [171784-07-9]



C₁₈H₁₆NO₂[⊕] 278.33
 Counterion not specified. Presumably exists as internal salt with ionised OH group. Quaternary alkaloid from *Clematis purpurea* var. *hybrida*. Pale yellow oil.
 Sayed, H.M. *et al.*, *Indian J. Chem., Sect. B*, 1995, **34**, 1111-1113

Cleomycin B₂ C-524
 [76741-88-3]




Peptide antibiotic. Isol. from *Streptomyces verticillus*. Active against gram-positive and -negative bacteria and tumours. Sol. H₂O. Occurs complexed with Cu. λ_{max} 244 ; 294 (MeOH) (Berdy).

Hydrochloride:
 Pale-yellow amorph. powder. Mp 185-188° dec.

Cu complex:
 Blue powder. Mp 197-201°.
 Umezawa, H. *et al.*, *J. Antibiot.*, 1980, **33**, 1079 (*struct, nmr*)

3,14-Clerodadiene-13,18-diol C-525



C₂₀H₃₄O₂ 306.487

(ent-5α,13E)-form
Sagittariol
 [56497-92-8]
 Constit. of *Sagittaria sagittifolia* (arrowhead). Cryst. (C₆H₆/hexane). Mp 109°. [α]_D +41 (CHCl₃).
18-(2-Pyrrolecarbonyl): Nakamurol D
 [174756-41-3]

C₂₅H₃₇NO₃ 399.572
 Constit. of *Agelas nakamurai*. Pale yellow oil. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D²⁵ -8.5 (c, 1.2 in CHCl₃).

13-Epimer, Deacetylrigidusol
 [329689-18-1]
 C₂₀H₃₄O₂ 306.487
 Constit. of *Haplopappus rigidus*. Gum. [α]_D²⁴ -21.1 (c, 1.21 in CHCl₃).

13-Epimer, 18-Ac: Rigidusol
 [329688-58-6]
 C₂₂H₃₆O₃ 348.525
 Constit. of *Haplopappus rigidus*. Gum. [α]_D²⁴ -31.8 (c, 0.11 in CHCl₃).


Sharma, S.C. *et al.*, *Phytochemistry*, 1975, **14**, 1055; 1984, **23**, 1194 (*Sagittariol*)
 Shoji, N. *et al.*, *J. Nat. Prod.*, 1996, **59**, 448-450 (*Nakamurol D*)
 Morales, G. *et al.*, *Phytochemistry*, 2000, **55**, 863-866 (*Rigidusol, Deacetylrigidusol*)

Clinacoside C C-526

[212000-76-5]
 MeS(O)CH=CHCONHCH₂CH₂OGlc
 C₁₂H₂₁NO₈S 339.366
 Constit. of *Clinacanthus nutans*. Oil. [α]_D²¹ +60 (c, 0.4 in Py). λ_{max} 252 (log ε 3.64) (MeOH).

Teshima, K.-I *et al.*, *Phytochemistry*, 1998, **48**, 831-835 (*isol, pmr, cmr*)

Clionamide C-527




C₁₉H₁₈BrN₃O₄ 432.273

(S)-form [68835-91-6]
 Major metab. from the sponge *Cliona celata*. Chelate forming. Unstable yellow powder. [α]_D +32.1 (c, 2.12 in MeOH). λ_{max} 227 (ε 31000); 291 (ε 11000); 296 (ε 12000); 311 (sh) (ε 9400) (MeOH) (De-rep).

Tetra-Ac: Tetraacetylclionamide
 [68857-44-3]
 C₂₇H₂₆BrN₃O₈ 600.422
 Isol. from *Cliona celata*. Cryst. (THF/diisopropyl ether). Mp 209-211°. [α]_D +45 (c, 0.7 in Me₂CO).

Andersen, R.J. *et al.*, *Can. J. Chem.*, 1979, **57**, 2325 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Stonard, R.J. *et al.*, *J.O.C.*, 1980, **45**, 3687 (*isol, deriv*)

Cliviahaksine C-528
 [85967-04-0]




C₁₇H₁₉NO₅ 317.341
 Alkaloid from *Clivia miniata* (Amaryllidaceae). Fine prisms (Me₂CO). Mp 160-165°. [α]_{Hg}²⁰ -60 (c, 0.05 in CHCl₃).

O-(2,6-Dimethyl-3-pyridinecarbonyl): Cliviaaline
 [83482-59-1]
 C₂₅H₂₆N₂O₆ 450.49
 Trace alkaloid from *Clivia miniata* (Amaryllidaceae). Fine needles (Me₂CO). Mp 200-201°.

Döpke, W. *et al.*, *Z. Chem.*, 1982, **22**, 310 (*Cliviaaline*)
 Döpke, W. *et al.*, *Z. Chem.*, 1983, **23**, 102 (*Cliviahaksine*)


Clivisyaline C-529
Cliviasyaline (incorr.)
 [79038-08-7]



C₁₇H₂₁NO₅ 319.357
 Alkaloid from *Clivia miniata* (Amaryllidaceae). Fine prisms (Me₂CO). Mp 140-145°. [α]_{Hg}²⁴ +125 (c, 0.1 in CHCl₃).

Döpke, W. *et al.*, *Z. Chem.*, 1981, **21**, 223-224 (*isol, uv, ir, pmr, ms, struct*)


Clivojuline C-530
 [77410-45-8]



Absolute Configuration

C₁₈H₂₁NO₄ 315.368
 Evidently a secolycorenine alkaloid. Alkaloid from *Clivia miniata* (Amaryllidaceae). Fine needles (Me₂CO). Mp 130-135° dec. [α]_{Hg}²⁰ +25 (c, 0.1 in CHCl₃).
 Döpke, W. *et al.*, *Heterocycles*, 1981, **16**, 529 (*uv, pmr, ms, struct*)

Clivonidine C-531
 [86667-23-4]




Absolute Configuration

C₁₇H₁₇NO₄ 299.326
 Alkaloid from the total herb of *Clivia miniata* (Amaryllidaceae). Platelets

(EtOH). Mp 238-239°.

Ali, A.A. *et al.*, *J. Nat. Prod.*, 1983, **46**, 350-352 (*isol, uv, ir, pmr, ms, struct*)**Clivonine****C-532**

[477-16-7]

C₁₇H₁₉NO₅ 317.341Alkaloid from the rhizomes of *Clivia miniata* (Amaryllidaceae). Prisms (EtOAc). Mp 199-200° (193-194°). [α]_D²³ +41.24 (c, 1.11 in CHCl₃).**Hydrochloride:**Prisms (MeOH/Et₂O). Mp 282-287° dec.**Picrate:** Mp 250-254° dec.**Ac:** Mp 194-196°.**Propanoyl: Poetinatine**

[52657-07-5]

C₂₀H₂₃NO₆ 373.405Alkaloid from *Narcissus poeticus* var. *ornatus* (Amaryllidaceae). Prisms (Me₂CO). Mp 212-213°. [α]_D +50 (c, 0.15 in CHCl₃).**O-(3-ξ-Hydroxybutanoyl): Clivatine**

[1355-66-4]

C₂₁H₂₅NO₇ 403.431Alkaloid from *Clivia miniata* (Amaryllidaceae). Mp 166-169° (159-161°). [α]_D²⁵ +52 (c, 0.2 in CHCl₃).**O-(3-Hydroxybutanoyl): Nobilisitine B**C₂₁H₂₅NO₇ 403.431Alkaloid from *Clivia nobilis*. Needles (MeOH). Mp 205-207°. Epimer of Clivatine at the acyl side-chain. Config. not detd. λ_{max} 232 (log ε 4.21); 266 (log ε 3.66); 305 (log ε 3.65) (MeOH).**O-(3-Oxobutanoyl): Clivacetine**

[74730-02-2]

C₂₁H₂₃NO₇ 401.415Alkaloid from *Clivia miniata* (Amaryllidaceae). Prisms (EtOAc). Mp 152-155°. [α]_D²⁴ +53.8 (c, 0.67 in CHCl₃).**O-(5-Ethoxycarbonyl-2,6-dimethyl-3-pyridinecarbonyl): Cliviamartine**

[76983-01-2]

C₂₈H₃₀N₂O₈ 522.554Alkaloid from *Clivia miniata* (Amaryllidaceae). Needles (Me₂CO). Mp 172-175°. [α]_D²⁰ +45 (c, 0.2 in CHCl₃). Ester of Clivonine, C-532.**5a-Epimer: Cliviasine**

[19504-94-0]

Alkaloid from *Clivia miniata* (Amaryllidaceae). Mp 195-197°. Opt. rotn. not reported.**5a-Epimer, perchlorate:** Mp 273-276°.**5,5a-Diepimer: Clividine**


[29073-38-9]

C₁₇H₁₉NO₅ 317.341Alkaloid from *Clivia miniata*. Mp 195-197°. [α]_D²² -75 (c, 0.2 in CHCl₃).**3a,5a-Diepimer: Nobilisine**

[115587-27-4]

C₁₇H₁₉NO₅ 317.341Alkaloid from *Clivia nobilis* (Amaryllidaceae). Plates (MeOH). Mp 275-277°.**3a,5,5a,11c-Tetraepimer: Nobilisitine A**C₁₇H₁₉NO₅ 317.341Alkaloid from *Clivia nobilis*. Cryst. Mp 186-187°. λ_{max} 232 (log ε 4.28); 265 (log ε 3.67); 305 (log ε 3.71) (MeOH).Briggs, C.K. *et al.*, *J.A.C.S.*, 1956, **78**, 2899 (*isol*)Boit, H.-G. *et al.*, *Naturwissenschaften*, 1961, **48**, 603 (*Clivatine*)Döpke, W. *et al.*, *Tet. Lett.*, 1967, 451; 1970, 3245 (*Clivonine, Clividine, ir, pmr, cd, ms, struct*)Döpke, W. *et al.*, *Pharmazie*, 1970, **25**, 700 (*Cliviasine*)Jeffs, P.W. *et al.*, *Tetrahedron*, 1971, **27**, 5065 (*pmr, struct*)Irie, H. *et al.*, *Chem. Comm.*, 1973, 302 (*synth, Clividine*)Döpke, W. *et al.*, *Z. Chem.*, 1974, **14**, 57 (*Poetinatine*)Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 1827 (*Clivonine, Clivacetine, Clivatine*)Döpke, W. *et al.*, *Z. Chem.*, 1980, **20**, 374 (*Cliviamartine*)Jeffs, P.W. *et al.*, *J. Nat. Prod.*, 1988, **51**, 549 (*Nobilisine*)Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591 (*cd*)Evidente, A. *et al.*, *Phytochemistry*, 1999, **51**, 1151-1155 (*Nobilisitines*)**Clivorine****C-533**

[33979-15-6]



Absolute Configuration

C₂₁H₂₇NO₇ 405.447Cyclic otonecine diester. Alkaloid from *Ligularia clivorum* and *Ligularia hodgsonii*. Cryst. + 1H₂O or solvated tetrahedra (EtOAc or Me₂CO). Mp 148-150°. [α]_D²⁴ +79 (c, 0.713 in CHCl₃). [α]_D²⁴ +49 (c, 0.713 in MeOH).

▶ VT5800000

Picrate:


Yellow needles (EtOH). Mp 173-175°.

20,21-Dihydro, 20ξ-hydroxy: Hodgsonine

[218143-68-1]

C₂₁H₂₉NO₈ 423.462Alkaloid from *Ligularia hodgsonii*.Klásek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 2512-2522 (*isol, uv, ir*)Birnbaum, K.B. *et al.*, *Acta Cryst. B*, 1972, **28**, 2825-2833 (*cryst struct, abs config*)Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918-3935 (*cd*)Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*pmr*)Lin, G. *et al.*, *Rapid Commun. Mass Spectrom.*, 1998, **12**, 1445-1456 (*Hodgsonine*)Lin, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 857-863 (*isol, pmr, cmr*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CMV950**Clovamide****C-534**

N-[3-(3,4-Dihydroxyphenyl)-1-oxo-2-propenyl]-3-hydroxytyrosine, 9CI. N-(3,4-Dihydroxycinnamoyl)DOPA. N-Caffeoyl DOPA




(S,E)-form

C₁₈H₁₇NO₇ 359.335**(S,E)-form** [53755-02-5]Isol. from *Trifolium pratense* (red clover) and from the bark of *Dalbergia melanoxylon*. Also from cocoa liquor *Theobroma cacao*. Natural antioxidant. Amorph. powder or oil. Mp 125-130°. [α]_D¹⁸ -30.9 (c, 0.5 in H₂O).**Tetra-Me ether, Me ester:** [64803-83-4] Mp 152-153°. [α]_D²² +137 (c, 1 in CHCl₃).**(S,Z)-form** [53755-03-6]Isol. from *Trifolium pratense* (red clover) and from the bark of *Dalbergia melanoxylon*. Oil. [α]_D¹⁹ +28.7 (c, 0.57 in H₂O). **Tetra-Me ether, Me ester:** [64803-85-6] Syrup. [α]_D¹⁶ +83 (c, 1 in CHCl₃).

[120094-83-9]

Yoshihara, T. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 1107; 1977, **41**, 1679 (*isol*)Van Heerden, F.R. *et al.*, *Phytochemistry*, 1980, **19**, 2125 (*isol, cmr, synth*)Robiot, G. *et al.*, *J. Nat. Prod.*, 1988, **51**, 596 (*synth*)Sanbongi, C. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 454-457 (*S,E-form, isol cacao, hplc, pmr, cmr*)**CNS 2103****C-535**

N-[5-[[3-[[3-[[3-(3-Aminopropyl)amino]propyl]amino]propyl]amino]pentyl]-4-hydroxy-1H-indole-3-acetamide, 9CI [144923-45-5]



m = 5, n = 2

C₂₄H₄₂N₆O₂ 446.635Alkaloid from the water spider *Dolomedes okefinokensis*. Ca channel antagonist. Powder.McCormick, K.D. *et al.*, *Tetrahedron*, 1993, **49**, 11155-11168 (*isol, uv, pmr, ms*)**CNS 2104****C-536**

N-[3-[[3-[[3-[[3-(5-Aminopentyl)amino]propyl]amino]propyl]amino]propyl]-4-hydroxy-1H-indole-3-acetamide, 9CI [144923-46-6]

As CNS 2103, C-535 with m = 3, n = 4

C₂₄H₄₂N₆O₂ 446.635


Minor alkaloid from the water spider *Dolomedes okefinokensis*. Ca channel antagonist. Powder.

Pat. Coop. Treaty (WIPO), 1992, 92 14 709; *CA*, **118**, 824b (*isol*)

McCormick, K.D. *et al.*, *Tetrahedron*, 1993, **49**, 11155-11168 (*occur*)

Cocaine, BAN, USAN C-537

Methyl 3-(benzoyloxy)-8-methyl-8-azabicyclo[3.2.1]octane-2-carboxylate, 9CI. Benzoylcegonine methyl ester. Ecgonine methyl ester benzoate

C₁₇H₂₁NO₄ 303.357

α- and β-Cocaines are nonalkaloid struct. isomers. Adrenergic receptor antagonist. Potent CNS stimulant, bactericidal agent. Mydriatic agent. Topical anaesthetic. Drug of abuse. Log P 2.72 (calc).

(-)-form [50-36-2]

Principal alkaloid of *Erythroxylum* spp., esp. *Erythroxylum coca*, *Erythroxylum monogynum* (Erythroxylaceae). Prisms (EtOH). Mp 98°. Bp_{0.1} 187-188°. [α]_D -16 (CHCl₃). [α]_D -30 (MeOH). Pharmacol. active enantiomer.

- ▶ Abuse can lead to habituation or addiction. Adverse human CNS effects include distorted perception, convulsions, general anaesthesia. Eye irritant. LD₅₀ (mus, orl) 99 mg/kg. YM2800000

Hydrochloride: Cocaine hydrochloride, JAN, USAN

[53-21-4]

Mp 200-202° (197°). [α]_D²⁰ -71.9 (c, 2 in H₂O).

- ▶ Drug of abuse. Abuse can lead to habituation or addiction. LD₅₀ (mus, orl) 96 mg/kg. Exp. reprod. and teratogenic effects. YM3050000

Nitrate:Deliquescent cryst. + H₂O. Mp 58-63°.**(+)-Tartrate:**Prisms + H₂O. Mp 114-115°. [α]_D -41.2 (H₂O).**Picrate:**

Yellow powder.

Methiodide: Mp 169°.**2-Epimer: Pseudococaine. ψ-Cocaine.****Delcaine**

[478-73-9]

C₁₇H₂₁NO₄ 303.357

Minor alkaloid obt. prob. as an artifact by the ready base-cat. epimerisation of Cocaine during isol., e.g. from *Erythroxylum maxillense* (Erythroxylaceae). Prismatic cryst. Mp 46-47°.

2-Epimer, hydrochloride:Elongated leaflets (EtOH). Mp 205-210°. [α]_D +43 (H₂O).**N-De-Me: Norcocaine**

[18717-72-1]

C₁₆H₁₉NO₄ 289.33


Primary photoprod. of, and metab. prod. of cocaine. Shows strong physiol. activity comparable with that of Cocaine and is responsible for most of its toxicity *in vivo*. Mp 80-82°.

(+)-formSynthetic. Mp 96-98°. [α]_D²⁴ +15.5 (c, 1 in CHCl₃).**(±)-form [21206-60-0]**

Mp 79-80°.

Willstätter, R. *et al.*, *Annalen*, 1923, **434**, 111 (*synth, resoln*)Kovács, O. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 892 (*config*)Gabe, E.J. *et al.*, *Acta Cryst.*, 1963, **16**, 796 (*cryst struct*)Ashley, R. *et al.*, *Cocaine, its History, Uses and Effects*, Warner Books, N.Y., 1975,Wallace, G.J.E. *et al.*, *Anal. Chem.*, 1976, **48**, 34 (*glc*)*Cocaine: Chem., Biol., Clin., Soc. Treat.**Aspects*, (Mule, S.J., Ed.), CRC Press, 1976,Stenberg, V.I. *et al.*, *J. Het. Chem.*, 1976, **13**, 363 (*synth, pmr, Norcocaine*)Baker, J.K. *et al.*, *J. Het. Chem.*, 1978, **15**, 165 (*cmr*)Tufariello, J.J. *et al.*, *J.A.C.S.*, 1979, **101**, 2435 (*synth, pmr*)Hrynchuk, R.J. *et al.*, *Can. J. Chem.*, 1983, **61**, 481 (*cryst struct*)Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1986, **15**, 151 (*rev, uv, ir, pmr, cmr, ms, anal*)Lewin, A.H. *et al.*, *J. Het. Chem.*, 1987, **24**, (*synth, bibl*)Turner, C.E. *et al.*, *Cocaine: An Annotated Bibliography*, Research Institute of Pharmaceutical Sciences, Mississippi, 1988, (*book*)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1006Zhu, N. *et al.*, *Acta Cryst. C*, 1994, **50**, 2067 (*cryst struct, Norcocaine*)Fischman, M.W. *et al.*, *Handb. Exp. Pharmacol.*, 1996, **118**, 159 (*rev, pharmacol, tox*)Lin, R. *et al.*, *J.O.C.*, 1998, **63**, 4069-4078 (*synth, bibl*)Mans, D.M. *et al.*, *Org. Lett.*, 2004, **6**, 3305-3308 (*synth*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CNE750; CNF000**Cocculine****C-538***1,6-Didehydro-3-methoxyerythrinan-15-ol, 9CI*

[27675-39-4]

C₁₇H₂₁NO₂ 271.358

Alkaloid from the leaves of *Cocculus laurifolius*, the fruit of *Cocculus carolinus* and the rhizomes of *Cocculus trilobus* (Menispermaceae). Hypotensive agent which acts *via* ganglionic blocking. Cryst. (Me₂CO). Mp 217-218°. [α]_D +271.1 (MeOH).

Hydrochloride: Mp 222-223°.*Nitrate:* Mp 196-197°.**N-Me: Pachygonine**

[75667-86-6]

[75667-87-7]

C₁₈H₂₄NO₂[⊕] 286.393

Alkaloid from the roots of *Pachygone ovata* (Menispermaceae). Needles (MeOH/EtOAc)(as hydroxide). Mp 265-267° dec.(hydroxide). [α]_D +196.6 (c, 4.7 in MeOH) (as hydroxide).

Me ether: Cocculidine

[27675-40-7]

[78463-74-8]

C₁₈H₂₃NO₂ 285.385

Alkaloid from the leaves of *Cocculus laurifolius*. Hypotensive agent, ganglion blocker. Cryst. (petrol). Mp 93-95° (86-87°). [α]_D +250.9 (CHCl₃).

Me ether, N-oxide: Cocculidine N-oxide

[206559-17-3]

C₁₈H₂₃NO₃ 301.385

Alkaloid from the leaves of *Cocculus laurifolius*.

Me ether, N-Me:C₁₉H₂₆NO₂[⊕] 300.42

Mp 241-242° (as iodide).

O-De-Me: Coclafine


[142713-49-3]

C₁₆H₁₉NO₂ 257.332

Alkaloid from leaves of *Cocculus laurifolius* (Menispermaceae). Cryst. (Me₂CO). Mp 264-266° (Me₂CO). [α]_D +255 (c, 0.15 in MeOH).

Yunusov, S. *et al.*, *Zh. Obshch. Khim.*, 1950, **20**, 368; *CA*, **44**, 6582g (*isol*)Yunusov, S. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 74; *Chem. Nat. Compd. (Engl. Transl.)*, 69 (*pmr, ms, struct*)Razakov, R. *et al.*, *Chem. Comm.*, 1974, 150 (*cryst struct, abs config*)Razakov, R. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1974, 218 (*Cocculidine*)Elsohly, M.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 132 (*isol, uv, ir, pmr, ms*)McPhail, A.T. *et al.*, *Tet. Lett.*, 1976, 485 (*pmr*)McPhail, A.T. *et al.*, *J.C.S. Perkin 2*, 1977, 1156 (*cryst struct*)Ju-ichi, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 563; 1981, **29**, 396 (*isol, synth, pmr, ms, ir*)Bhakuni, D.S. *et al.*, *J.C.S. Perkin 1*, 1978, 618 (*biosynth*)Bhat, S.V. *et al.*, *J. Nat. Prod.*, 1980, **43**, 588 (*Pachygonine, isol, uv, pmr, ms, struct*)Madhusudan, K.P. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 907 (*ms*)Ziyayev, R. *et al.*, *Khim. Prir. Soedin.*, 1991, 84-86; *Chem. Nat. Compd. (Engl. Transl.)*, 73-74 (*Coclafine*)Tsakadze, D. *et al.*, *CA*, 1998, **128**, 306258d (*Cocculidine N-oxide*)**Cocculitine****C-539***1,6-Didehydro-3,15-dimethoxyerythrinan-2-ol, 9CI*

[64996-74-3]

C₁₈H₂₃NO₃ 301.385

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Cryst. (EtOAc). Mp 142-143°. $[\alpha]_D^{25} +93$ (c, 0.4 in MeOH).

Ac:

Cryst. (EtOAc). Mp 135°.

O¹⁵-De-Me: **Cocculitinine**

[77795-07-4]

C₁₇H₂₁NO₃ 287.358

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae).

2-Ketone: **Cocculidinone**

[77795-06-3]

C₁₈H₂₁NO₃ 299.369

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Oil.

Singh, A.N. *et al.*, *J. Nat. Prod.*, 1977, **40**, 322 (isol, uv, ir, pmr, ms, struct)


Bhakuni, D.S. *et al.*, *Tetrahedron*, 1980, **36**, 3107 (isol, ms, uv, ir, struct, derivs)

Madhusudan, K.P. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 907 (ms)

Cocculolidine

C-540

[13497-04-6]



C₁₅H₁₉NO₃ 261.32

Alkaloid from leaves and rhizomes of *Cocculus trilobus* and the fruit of *Cocculus carolinus* (Menispermaceae). Has insecticidal properties. Antifeedant.

Prismatic cryst. (CCl₄). Mp 144-146°. $[\alpha]_D^{25} +273$ (c, 1 in CHCl₃).

Hydrochloride: Mp 247-251° dec.

Wada, K. *et al.*, *Agric. Biol. Chem.*, 1967, **31**, 452; 1968, **32**, 1187 (isol, uv, ir, pmr, ms, struct, stereochem)

Wada, K. *et al.*, *J. Agric. Food Chem.*, 1968, **16**, 471 (isol)


Elsöhly, M.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 132 (isol, uv, ir, pmr, ms)

Ju-ichi, M. *et al.*, *Yakugaku Zasshi*, 1978, **98**, 886 (isol)

Kawasaki, T. *et al.*, *Tet. Lett.*, 2001, **42**, 8003-8006 (synth)

Coccuorbiculatine A

C-541



C₃₈H₄₂N₂O₆²⁺ 622.76

(S)-form

Quaternary alkaloid from the stems of *Cocculus orbiculatus*. Powder (CHCl₃/MeOH). $[\alpha]_D^{25} +110.2$ (c, 0.3 in MeOH).


Counterion not specified. λ_{\max} 218 ; 228 ; 279 ; 304 (MeOH).

Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1056-1060 (isol, pmr, cmr, ms)

Coccutrine

C-542

1,6-Didehydro-3,17-dimethoxyerythrinan-15-ol, 9CI
[59553-89-8]



C₁₈H₂₃NO₃ 301.385

Alkaloid from the rhizomes of *Cocculus trilobus* (Menispermaceae). Needles. Mp 263-265°. $[\alpha]_D +232$ (MeOH).

McPhail, A.T. *et al.*, *Tet. Lett.*, 1976, 485 (pmr, ms, struct)


McPhail, A.T. *et al.*, *J.C.S. Perkin 2*, 1977, 1156 (cryst struct)

Ju-ichi, M. *et al.*, *Yakugaku Zasshi*, 1978, **98**, 886; *CA*, **89**, 215622c (isol)

Cocconvine

C-543

1,2,6,7-Tetrahydro-3-methoxyerythrinan-15-ol, 9CI



C₁₇H₁₉NO₂ 269.343

Natural-form [61445-80-5]

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Mp 137-138°.

Me ether: **Cocconvinine**. O-Methylcocconvine

[64543-55-1]

C₁₈H₂₁NO₂ 283.369

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Needles (hexane). Mp 103-104°.

(±)-form

Me ether: [78463-73-7]

Synthetic. Oil.

Me ether, picrate:

Yellow prisms (EtOH). Mp 166-167°.

Singh, A.N. *et al.*, *Experientia*, 1976, **32**, 1368 (uv, ir, pmr, ms, struct, stereochem)

Singh, A.N. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 388 (Cocconvinine)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1980, **36**, 2153; 3107 (isol, biosynth, uv, ir, pmr, struct)

Ju-ichi, M. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 396 (Cocconvinine, Cocconvinine, synth, ir, pmr, ms)


Madhusudan, K.P. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 907 (ms)

Sano, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **65**, 94 (synth)

Cochinchistemonine

C-544

[930775-40-9]



C₂₂H₃₅NO₆ 409.522

Alkaloid from the roots of *Stemona cochinchinensis*. Needles. Mp 176-180°. $[\alpha]_D^{20} +9$ (c, 0.16 in CHCl₃). λ_{\max} 232 (log ϵ 6.95) (MeOH).

12-Ketone: **Cochinchistemoninone**

[949092-26-6]

C₂₂H₃₃NO₆ 407.506

Alkaloid from the roots of *Stemona cochinchinensis*. Yellow-brown gum.

$[\alpha]_D^{20} +39$ (c, 0.28 in MeOH). λ_{\max} 229 (log ϵ 3.91) (MeOH).

Wang, Y.-Z. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1356-1359 (Cochinchistemoninone)


Lin, L.-G. *et al.*, *Tet. Lett.*, 2007, **48**, 1559-1561 (Cochinchistemonine)

Cochinnicin I

C-545

RP 59451. Antibiotic RP 59451

[143728-97-6]



C₄₆H₄₇N₇O₁₂ 889.917

Depsipeptide antibiotic. Prod. by a *Microbispora* sp. Endothelin antagonist.

$[\alpha]_D^{23} -10$ (c, 0.1 in MeOH). Identity with RP 59451 not confirmed. λ_{\max} 212 (ϵ 39600); 230 (sh) (ϵ 15300); 269 (ϵ 15600) (MeOH) (Derep).

5''-Chloro: **Cochinnicin III**

[143728-99-8]

C₄₆H₄₆ClN₇O₁₂ 924.362

Prod. by a *Microbispora* sp.

ATCC55140. Endothelin antagonist.

Antiasthmatic, antihypertensive and antiischemic agent. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{23} -10$ (c, 0.1 in MeOH). λ_{\max} 213 (ϵ 44600); 230 (sh) (ϵ 19200); 275 (ϵ 20200) (MeOH) (Derep).

2-Epimer: **Cochinnicin V**

[146876-02-0]

C₄₆H₄₇N₇O₁₂ 889.917

Prod. by a *Microbispora* sp.

ATCC55140. Endothelin antagonist.

Antihypertensive agent. $[\alpha]_D^{23} +20$ (c, 0.1 in MeOH). λ_{\max} 212 (ϵ 39600); 230 (sh) (ϵ 15300); 269 (ϵ 15600) (MeOH) (Derep).

2-Epimer, 5''-chloro: **Cochinnicin II**. RP

55185. Antibiotic RP 55185

[114919-77-6]

C₄₆H₄₆ClN₇O₁₂ 924.362

Prod. by a *Microbispora* sp. Endothelin antagonist. $[\alpha]_D^{23} +20$ (c, 0.1 in MeOH). Identity with RP 55185 not confirmed. λ_{\max} 213 (ε 44600); 230 (sh) (ε 19200); 275 (ε 20200) (MeOH) (Derep).

2-Epimer, 17-hydroxy, 5'-chloro: Cochlinicin IV

[146874-41-1]

C₄₆H₄₆ClN₇O₁₃ 940.361


Prod. by a *Microbispora* sp. ATCC55140. Endothelin antagonist. Antihypertensive agent. $[\alpha]_D^{23} +30$ (c, 0.1 in MeOH). λ_{\max} 213 (ε 44600); 230 (sh) (ε 19200); 275 (ε 20200) (MeOH) (Derep).

[143728-98-7]

Lam, Y.K.T. *et al.*, *J. Antibiot.*, 1992, **45**, 1709; 1717; 1792 (*isol, pmr, cmr, struct, props*)

Cochlearenine**C-546**

[914475-10-8]


C₂₂H₃₅NO₄ 377.523

Alkaloid from *Aconitum cochleare* and *Delphinium linearilobum*. Antioxidant. Amorph. solid. $[\alpha]_D^{20} -35.5$ (c, 0.5 in CHCl₃). λ_{\max} 203 (log ε 4.2); 226 (sh) (log ε 3); 284 (log ε 2.5) (MeOH).

Kolak, U. *et al.*, *Phytochemistry*, 2006, **67**, 2170-2175 (*isol, pmr, cmr, ms*)

Cochliodinol**C-547**

2,5-Dihydroxy-3,6-bis[5-(3-methyl-2-butenyl)-1H-indol-3-yl]-2,5-cyclohexadiene-1,4-dione, 9CI
[11051-88-0]

C₃₂H₃₀N₂O₄ 506.6


Benzoquinone antibiotic. Metab. of *Chaetomium globosum*, *Chaetomium cochliodes*, *Chaetomium elatum* and *Chaetomium abuense*. Possesses fungistatic and fungicidal activities. Purple cryst. Sol. MeOH, Et₂O; poorly sol. H₂O, hexane. Mp 213-215° (208° dec.). λ_{\max} 278 (ε 31600); 484 (ε 3980) (CHCl₃) (Derep). λ_{\max} 279 (ε 30900); 471 (ε 4470) (EtOH) (Derep). λ_{\max} 278 (ε 30900); 471 (ε 3550) (MeOH) (Berdy). λ_{\max} 278 (ε 31620); 484 (ε 3980) (CHCl₃) (Berdy). λ_{\max} 282 (ε 26300); 470 (ε 3800) (EtOH) (Berdy).

► LD₅₀ (mus, ipr) 200 mg/kg, LD₅₀ (mus, ipr) 40.8 mg/kg, LD₅₀ (mus, scu) 147 mg/kg, LD₅₀ (mus, orl) 221 mg/kg.

DK3529000

Di-Me ether: Mp 208-210° dec.Jerram, W.A. *et al.*, *Can. J. Chem.*, 1975, **53**,727 (*uv, ir, pmr, cmr, ms, struct*)Jerram, W.A. *et al.*, *Acta Cryst. B*, 1977, **33**,293 (*cryst struct*)Taylor, A. *et al.*, *Phytochemistry*, 1978, **17**,1045 (*biosynth*)Brewer, D. *et al.*, *Can. J. Microbiol.*, 1984, **30**,1068 (*props*)Hörcher, U. *et al.*, *Annalen*, 1986, 1765 (*synth*)Saito, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**,1942 (*isol*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties**of Industrial Materials*, 8th edn., Van*Nostrand Reinhold*, 1992, CNF159**Cochliodinone****C-548**

3,6-Bis[5-(3-methyl-2-butenyl)-1H-indol-3-yl]furo[3,2-b]furan-2,5-dione, 9CI
[56489-12-4]

C₃₂H₂₈N₂O₄ 504.584

Metab. of *Chaetomium cochliodes* and *Chaetomium globosum*. Red-orange needles (toluene). Mp 275-276°. Shows orange fluor. in soln.

Di-N-Ac:Yellow needles (Ac₂O). Mp 35°.


Brewer, D. *et al.*, *Can. J. Microbiol.*, 1968, **14**, 861-866 (*isol*)

Jerram, W.A. *et al.*, *Can. J. Chem.*, 1975, **53**,727-737 (*synth, struct, pmr, cmr, uv*)**Coclamine****C-549**C₁₉H₂₃NO₃ 313.396

Struct. unknown. Alkaloid from *Cocculus laurifolius*. Needles. Mp 140-142° Mp 237-240° (hydrobromide). $[\alpha]_D -245$ (MeOH) (as hydrobromide).

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1956, **76**,1048-1050; *CA*, **51**, 1542i (*isol*)**Coclifoline****C-550**C₁₉H₂₇NO₃ 317.427

Struct. unknown. Alkaloid from *Cocculus laurifolius*. Mp 158-159° (as picrate).

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1956, **76**,1048-1050; *CA*, **51**, 1542i (*isol*)**Coclobine****C-551**C₃₇H₃₈N₂O₆ 606.717

Alkaloid from *Cocculus trilobus* (Menispermaceae).

(R)-form*O*⁷-*De-Me*: **1,2-Dehydro-2-norlimacusine**

[103976-35-8]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from *Caryomene olivascens* (Menispermaceae). $[\alpha]_D -94$ (c, 0.16 in MeOH).

(S)-form


Alkaloid from *Cocculus trilobus* (Menispermaceae). Amorph. $[\alpha]_D^{20} +123$ (c, 0.69 in CHCl₃).

*O*¹²-*De-Me*: **12-O-Demethylcoclobine**C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the stem bark of *Guateria guianensis* (Annonaceae). $[\alpha]_D^{20} +220$ (c, 0.25 in CHCl₃).

Ito, K. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1163;*CA*, **71**, 124748k (*isol, uv, pmr, ms, struct*)Lavault, M. *et al.*, *Chem. Pharm. Bull.*, 1986,**34**, 1148 (*1,2-Dehydro-2-norlimacusine*)Berthou, S. *et al.*, *J. Nat. Prod.*, 1989, **52**, 95*(12-O-Demethylcoclobine)***Cocculinine****C-552**

[54370-90-0]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the leaves and stems of *Cocculus pendulus* (Menispermaceae). Antineoplastic agent. Cryst. (CHCl₃/MeOH). Mp 260-263°. $[\alpha]_D +312$ (c, 0.50 in CHCl₃). Log P 7.89 (uncertain value) (calc).

Di-Ac: Mp 217°. $[\alpha]_D +225.9$ (c, 0.47 in CHCl₃).

N-De-Me: **N-Norcocculinine**C₃₄H₃₂N₂O₆ 564.637

Alkaloid from leaves and stems of *Cocculus pendulus* (Menispermaceae). Amorph. powder. Mp 250° dec. $[\alpha]_D +294$. Site of demethylation unknown.

Di-N-de-Me: **Cocsilinine. 2,2'-Bisnorcocculinine**C₃₃H₃₀N₂O₆ 550.61

Alkaloid from leaves and stems of *Cocculus pendulus* (Menispermaceae). Amorph. powder. Mp 295° dec.

*O*⁶-*Me*: **6'-O-Methylcocculinine**C₃₆H₃₆N₂O₆ 592.69

Alkaloid from leaves and stems of *Cocculus pendulus* (Menispermaceae). Mp 168-170°. $[\alpha]_D +307$.

*O*⁶-*Me*, *N*²-*de-Me*: **Pendine**

[59114-65-7]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the leaves and stems of *Cocculus pendulus* (Menispermaceae). Cryst. (MeOH/Et₂O). Mp 170-171°. $[\alpha]_D +275$ (c, 1.0 in CHCl₃).

*O*⁶-*Me*, *N-de-Me*: **Cocsilone. 6'-O-Methyl-N-norcocculinine**C₃₅H₃₄N₂O₆ 578.663

Alkaloid from leaves and stems of *Cocculus pendulus* (Menispermaceae). Amorph. powder. Mp 257° dec. $[\alpha]_D +297$. Site of demethylation not determined but presumably *N*² as the props.

do not correspond with those of Pendine.

O¹²-Me: N-Methylgilletine

Mp 156-157°. [α]_D³⁰ +310 (c, 0.41 in CHCl₃).

O¹²-Me, N²-de-Me: Gilletine. Alkaloid TGLA

[52038-20-7]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the leaves of *Triclisia gilletii* (Menispermaceae). Rosettes of needles (CHCl₃/Et₂O). Mp 174-176°. [α]_D²⁸ +294.3 (c, 0.56 in MeOH).

Di-Me ether: Obt. by methylation of Cocculinine, Gilletine or Pendine. Cryst. (EtOAc/MeOH). Mp 145-146° Mp 173-175° Mp 201-203°. [α]_D +300 (CHCl₃) (193.0).

Di-Me ether, N²-De-Me: Pendilinine. O-Methylpendine

[142617-86-5]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from leaves and stems of *Cocculus pendulus* (Menispermaceae). Amorph. yellow powder. Mp 280° dec. [α]_D +253 (CHCl₃). [α]_D +286 (CHCl₃) (natural).

O⁶-De-Me, O⁶-Me: Pendulinine

[59114-69-1]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the leaves and stems of *Cocculus pendulus* (Menispermaceae). Cryst. (CHCl₃/MeOH). Mp 272-273°. [α]_D +285 (c, 0.5 in CHCl₃).

O⁶-De-Me, O⁶-Me, di-Ac: Mp 210-211°. [α]_D +203.3 (CHCl₃).

Stereoisomer, O¹²-Me, N²-de-Me, N²-oxide: Isogilletine N-oxide

[77431-57-3]

C₃₅H₃₄N₂O₇ 594.663

Alkaloid from the leaves of *Triclisia gilletii* (Menispermaceae). Mp 218-220° dec. [α]_D²⁹ +216 (c, 0.94 in CHCl₃/MeOH 9:1). Stereochem. unknown.

Joshi, P.P. *et al.*, *Indian J. Chem.*, 1974, **12**, 517 (*Cocculinine, uv, ir, pmr, ms, struct*)


Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 1 (*Gilletine, isol*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1975, **31**, 2575 (*Pendine, Pendulinine, Cocculinine*)

Dwuma-Badu, D. *et al.*, *Heterocycles*, 1978, **9**, 995 (*Gilletine, uv, ir, pmr, ms, struct*)

Owusu, P.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 61 (*Isogilletine N-oxide*)


Jain, S. *et al.*, *Fitoterapia*, 1991, **62**, 391-395 (*6'-O-Methylcocculinine, Cocsiline, N-Norcocculinine, Cocsilinine, Pendine, Pendilinine, Pendulinine*)

Cocsupendine**C-553**

C₃₄H₃₀N₂O₆ 562.621

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +116.6 (MeOH).

Atta-ur-Rahman, *et al.*, *Pure Appl. Chem.*, 1986, **58**, 663 (*uv, ir, pmr, ms, struct*)

Codamine**C-554**

C₂₀H₂₅NO₄ 343.422

(R)-form [21052-10-8]

Cryst. (petrol). Mp 127-128°. [α]_D¹⁷ -68.8 (c, 1 in EtOH).

Hydrochloride:

Cryst. (CHCl₃/Et₂O). Mp 120-125°. [α]_D¹⁸ -56 (c, 0.5 in H₂O).

Picrate:

Cryst. (2-propanol). Mp 147-149°. [α]_D²⁴ -84.4 (c, 1 in CHCl₃).

Methiodide:

Cryst. (2-propanol). Mp 200-200.5°. [α]_D³⁰ -100 (c, 1 in EtOH).

(S)-form [21040-59-5]

Minor constit. of opium, of *Bongardia chrysogonum*, *Guatteria chrysopetala* (Annonaceae), and *Argemone grandiflora*. Cryst. (petrol). Mp 126-127°. [α]_D¹⁵ +66.1 (c, 1 in EtOH).

Hydrochloride:

Cryst. (CHCl₃/Et₂O). Mp 123°. [α]_D¹⁸ +59 (c, 0.5 in H₂O). No CAS Reg No to CA 125.

Picrate: Mp 147-149°.

Methiodide:

Cryst. (2-propanol). Mp 200.5-201°. [α]_D +97.6 (c, 1 in EtOH).

N-oxide (S-): Codamine cis-N-oxide

[112468-25-4]

C₂₀H₂₅NO₅ 359.421

Alkaloid from trunk bark of *Duguetia spixiana* (Annonaceae). Noncryst. [α]_D 0 (c, 0.16 in CHCl₃).

Me ether: see Laudanosine, L-69

(±)-form [5977-85-5]

Cryst. (petrol). Mp 106-108°.

Brockmann-Hanssen, E. *et al.*, *J. Pharm. Sci.*, 1965, **54**, 1531 (*isol*)

Cassels, B.K. *et al.*, *Tetrahedron*, Suppl. 8, 1966, 485-490 (*synth, resoln*)

Battersby, A.R. *et al.*, *J.C.S. (C)*, 1968, 210-216 (*synth, resoln, O-benzyl*)


Benn, M.H. *et al.*, *Phytochemistry*, 1972, **11**, 461-464 (*isol*)

Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 1302-1306 (*pmr*)

Debourges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664-673; 852-859 (*N-oxide*)

Codeine, BAN, USAN**C-555**

7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-ol, 9CI. Methylmorphine. Codecept. Kodein. Tussipan [76-57-3]



Absolute Configuration

C₁₈H₂₁NO₃ 299.369

Opium alkaloid (*Papaver somniferum*) (content ca. 1%), also obt. synthetically by methylation of Morphine, M-704 (Papaveraceae). Relatively nonaddictive analgesic. Antitussive agent. Possesses anaesthetic props. Prisms (Et₂O or C₆H₆), octahedra or rhombic prisms + H₂O (H₂O). Sol. EtOH, CHCl₃, Me₂CO; spar. sol. CCl₄, H₂O. Mp 155°. [α]_D -137.8 (EtOH). Log P 0.82 (uncertain value) (calc). Also used as complex with sulfonated styrene-divinylbenzene copolymer (Codeine polistirex, USAN).

▶ Human and exp. reprod. effects. Exp. teratogen. LD₅₀ (rat, orl) 427 mg/kg. QD0893000

Hydrochloride: [1422-07-7]

Prisms + 2H₂O (H₂O). Mp 287° (as dihydrate). [α]_D^{22.5} -108.2 (H₂O).

▶ LD₅₀ (rat, orl) 750 mg/kg. QD1050000

Sulfate salt: Codeine sulfate, USAN

[6854-40-6]

[1420-53-7]

Needles + 3H₂O. Component of Copavin.

Phosphate salt: Codeine phosphate, BAN, JAN, USAN. Brochodine. Codeinfos.

Codeisan. Codeophen. Codlin. Duraspán. Paveral. Sedantole. Solcodein.

Vixaton

[41444-62-6]

[52-28-8]

C₁₈H₂₁NO₆P 378.341

Needles + ½H₂O. Component of Proval and Migraleve.

Compd. with calcium bis(2-acetylsalicylate): Paxidine

[81648-77-3]

N-Oxide: Codeine N-oxide. Codeigene.

Codeine aminoxyde. Genocodein

[3688-65-1]

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Platelets (H₂O). Mp 231-232°.

▶ QD1260000

N-Oxide; hydrochloride:

Cryst. Mp 219-220°. [α]_D²⁰ -105.8 (c, 2 in H₂O).

6-O-β-D-Glucuronopyranoside: [20736-11-2]

C₂₄H₂₉NO₉ 475.494

Mp 274-275° dec. (hemihydrate). [α]_D²⁰ -224 (c, 1 in H₂O).

Ac: [6703-27-1]

Prisms (Et₂O). Mp 133.5°.

▶ LD₅₀ (mus, scu) 99 mg/kg. QD0960000

6-Ketone: 7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-one.**Codeinone**

[467-13-0]

C₁₈H₁₉NO₃ 297.353


Naturally occurring alkaloid in *Papaver* spp. (Papaveraceae); precursor of Codeine and Morphine, M-704. Prisms (EtOAc). Mp 181.5-182.5°. [α]_D²⁰ -205 (EtOH). Subl. under reduced pressure.

▶ LD₅₀ (mus, scu) 11 mg/kg. QD2169000 [6059-47-8, 5913-76-8]

- Findlay, S.P. *et al.*, *J.A.C.S.*, 1950, **72**, 3247 (*Codeinone, synth*)
- Bentley, K.W. *et al.*, *Chemistry of the Morphine Alkaloids*, Oxford Univ. Press, 1954, 57-97 (*isol, props, uv, synth*)
- Kartha, G. *et al.*, *Acta Cryst.*, 1962, **15**, 326-333 (*cryst struct, abs config*)
- Brochmann-Hanssen, E. *et al.*, *J. Pharm. Sci.*, 1964, **53**, 1549-1550 (*glc, chromatog*)
- Batterham, T.J. *et al.*, *Aust. J. Chem.*, 1965, **18**, 1799-1806 (*pmr*)
- Wheeler, D.M.S. *et al.*, *J.A.C.S.*, 1967, **89**, 4494-4501 (*ms*)
- DeAngelis, G.G. *et al.*, *Tetrahedron*, 1969, **25**, 5099-5112 (*ord*)
- Parker, H.I. *et al.*, *J.A.C.S.*, 1972, **94**, 1276-1282 (*biosynth*)
- Weller, D.D. *et al.*, *J. Med. Chem.*, 1976, **19**, 1171-1175 (*synth*)
- Carroll, F.I. *et al.*, *J.O.C.*, 1976, **41**, 996-1001 (*cmr*)
- Phillipson, J.D. *et al.*, *Phytochemistry*, 1976, **15**, 1297-1301 (*oxide*)
- Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1981, **10**, 93 (*rev, phosphate*)
- Whittle, B.J.R. *et al.*, *Gut*, 1981, **22**, 798 (*Paxidine*)
- Moos, W.H. *et al.*, *J.O.C.*, 1983, **48**, 227-238 (*synth*)
- White, J.D. *et al.*, *Tetrahedron*, 1983, **39**, 2393-2397 (*synth, ir, pmr*)
- Quiding, H. *et al.*, *Eur. J. Clin. Pharmacol.*, 1986, **30**, 673 (*metab*)
- Canfield, D.V. *et al.*, *Acta Cryst. C*, 1987, **43**, 977-979 (*cryst struct*)
- Ishida, T. *et al.*, *Drug Metab. Dispos.*, 1991, **19**, 895 (*Codeinone, metab*)
- Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1069
- Nair, M.M. *et al.*, *Spectrosc. Lett.*, 1997, **30**, 497-505 (*pmr, cmr*)
- White, J.D. *et al.*, *J.O.C.*, 1999, **64**, 7871-7884 (*synth*)
- Rukhman, I. *et al.*, *Tetrahedron*, 2001, **57**, 1083-1092 (*6-glucuronoside*)
- Kolev, T. *et al.*, *Acta Cryst. E*, 2006, **62**, 255-257 (*Codeinone, cryst struct*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CNF500; CNG500; CNG675; CNF750; CNG750

Codonocarpine C-556

4-Hydroxy-26-methoxy-2-oxa-11,15,20-triazatricyclo[22.2.2.1^{3,7}]nonacosane-3,5,7(29),8,22,24,26,27-octaene-10,21-dione, 9CI
[33442-48-7]



C₂₆H₃₁N₃O₅ 465.548
Alkaloid from the bark of *Codonocarpus australis* (Gyrostemoniaceae). Pale-yellow rosettes of needles (MeOH). Mp 187° dec.

Hydrochloride:

Needles (HCl aq.). Mp 212-214°.

N,O-Di-Ac:

Cryst. (Me₂CO). Mp 169-171°.

N¹⁵-Me: **N-Methylcodonocarpine**
[38146-55-3]

C₂₇H₃₃N₃O₅ 479.575

Alkaloid from the bark of *Codonocarpus australis* (Gyrostemoniaceae). Pale-yellow needles (MeOH/C₆H₆). Mp 167-171° (164-166°). Opt. inactive. 15-Posn. (central N-atom) numbered acc. to CAS.


N¹⁵-Me, O-Ac:

Cryst. (Me₂CO). Mp 177-179°.

- Doskotch, R.W. *et al.*, *Chem. Comm.*, 1971, 300 (*uv, ir, pmr, struct*)
- Pilewski, N.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 186 (*isol, uv, ir, deriv*)
- Doskotch, R.W. *et al.*, *Tetrahedron*, 1974, **30**, 3229; 3237 (*struct, synth, pmr*)
- Humora, M.J. *et al.*, *Tet. Lett.*, 1980, **21**, 3971 (*synth*)
- Fujita, E. *et al.*, *Pure Appl. Chem.*, 1981, **53**, 1141 (*synth*)
- Nagao, Y. *et al.*, *Heterocycles*, 1982, **17**, 537 (*synth*)

Codonopsine C-557

2-(3,4-Dimethoxyphenyl)-1,5-dimethyl-3,4-pyrrolidinediol, 9CI. 3,4-Dihydroxy-2-(3,4-dimethoxyphenyl)-1,5-dimethylpyrrolidine
[26989-20-8]



C₁₄H₂₁NO₄ 267.324

Related to Radicamine A, R-10. Alkaloid from the epigeal parts of *Codonopsis clematidea*. Shows hypotensive action in exptl. animals. Shows antibiotic activity. Cryst. (Me₂CO/MeOH). Mp 150-151°. [α]_D²⁰ -16 (MeOH). Isomeric struct. originally assigned.

Methiodide:

Cryst. (MeOH). Mp 207-208°.

3'-Demethoxy: **Codonopsinine**. 2-(4-Methoxyphenyl)-1,5-dimethyl-3,4-pyrrolidinediol, 9CI. 3,4-Dihydroxy-2-(4-methoxyphenyl)-1,5-dimethylpyrrolidine
[32490-07-6]

C₁₃H₁₉NO₃ 237.298


Alkaloid from *Codonopsis clematidea* (Campanulaceae). Cryst. (MeOH). Mp 169-170°. [α]_D²⁰ -8.8 (c, 0.1 in MeOH). MeO group originally thought to be in the m-posn. Several stereoisomers synthesised.

- Matkhalikova, S.F. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 30; 606; 607; 1971, **7**, 210; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 24; 528; 530; 1971, **7**, 207 (*isol, ms, struct*)
- Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 495; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 488 (*stereochem, pmr*)
- Iida, H. *et al.*, *J.O.C.*, 1987, **52**, 1956 (*synth, pmr, ms, abs config*)
- Wang, C.-L.J. *et al.*, *J.O.C.*, 1991, **56**, 4341 (*synth*)
- Yoda, H. *et al.*, *Tet. Lett.*, 1996, **37**, 5531 (*synth, Codonopsinine*)
- Oliveira, D.F. *et al.*, *Tet. Lett.*, 1999, **40**, 2083-2086 (*synth*)
- Severino, E.A. *et al.*, *Org. Lett.*, 2000, **2**, 3039 (*activity, synth*)

- Goti, A. *et al.*, *Org. Lett.*, 2003, **5**, 4235-4238 (*synth*)
- Tashkhodzhaev, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **4**, 618-619 (*cryst struct, abs config*)
- Chandrasekhar, S. *et al.*, *Tet. Lett.*, 2005, **46**, 3127-3129 (*Codonopsinine, synth*)
- Reddy, J.S. *et al.*, *J.O.C.*, 2007, **72**, 2224-2227 (*Codonopsinine, synth*)

Coelenteramide C-558

[50611-86-4]



C₂₅H₂₁N₃O₃ 411.459

The light-emitter in *Aequorea*, *Cavernularia obesa* and other bioluminescent coelenterates. Blue fluorescent compd.

4'-Deoxy: **Renilla Oxyluciferin**

[50909-85-8]

[145022-31-7]


C₂₅H₂₁N₃O₂ 395.46

Chemiluminescent agent from *Renilla reniformis*.

- Hori, K. *et al.*, *Biochemistry*, 1973, **12**, 4463; 1975, **14**, 2371-2376 (*Renilla Oxyluciferin*)
- Shimomura, O. *et al.*, *Chem. Lett.*, 1975, 247 (*isol*)
- Shimomura, O. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1975, **72**, 1546 (*isol*)

Coelenteramine C-559

4-[5-Amino-6-phenylmethyl]pyrazinyl]-phenol, 9CI. 2-Amino-3-benzyl-5-(4-hydroxyphenyl)pyrazine
[37156-84-6]



C₁₇H₁₅N₃O 277.325

Isol. from jellyfish *Aequorea*. Luminescent agent. Yellow needles. Mp 217-219°.


N-Ac:

Yellow needles. Mp 199-200°.

- Shimomura, O. *et al.*, *Biochemistry*, 1972, **11**, 1602
- Kishi, Y. *et al.*, *Tet. Lett.*, 1972, 2747 (*synth*)
- Hori, K. *et al.*, *Chem. Comm.*, 1973, 492

Coelenterazine C-560

6-(4-Hydroxyphenyl)-2-[(4-hydroxyphenyl)methyl]-8-(phenylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one, 9CI. **Oplophorus Luciferin**. *Watasenia Praeluciferin*
[55779-48-1]



C₂₆H₂₁N₃O₃ 423.47Bioluminescence factor isol. from various marine sources incl. *Aequorea victoria*. Yellow cryst. Mp 176-181° dec.**4',4''-Di-O-sulfate: Watasenia Luciferin**


[61369-27-5]

C₂₆H₂₁N₃O₉S₂ 583.599Isol. from *Watasenia scintillans*.**4'-Deoxy: 6-(4-Hydroxyphenyl)-2,8-bis(phenylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one, 9CI. Renilla Luciferin**


[50909-86-9]

C₂₆H₂₁N₃O₂ 407.471Isol. from *Renilla reniformis* and other marine animals. Bioluminescence factor.**1'',7'-Didehydro: Dehydrocoelenterazine.***Watasenia Dehydropreluciferin*

[62541-09-7]

C₂₆H₁₉N₃O₃ 421.454Isol. from the liver of *Watasenia scintillans*. Bioluminescence factor.Dark red cryst. (Et₂O). Dec. at ca. 250°.Inoue, S. *et al.*, *Tet. Lett.*, 1976, **34**, 2971; 1977,2685 (*Watasenia Luciferin, Coelenterazine*)Inoue, S. *et al.*, *Chem. Lett.*, 1977, 259(*Dehydrocoelenterazine*)Inoue, S. *et al.*, *Chem. Lett.*, 1980, 299 (*synth*)Shimomura, O. *et al.*, *Comp. Biochem.**Physiol.*, B: *Comp. Biochem.*, 1980, 65; 435 (*occur*)Teranishi, K. *et al.*, *Bull. Chem. Soc. Jpn.*,1990, **63**, 3132 (*synth, conformn*)Kakoi, H. *et al.*, *Phytochemistry*, 1998, **48**,1669-1672 (*synth*)Adamczyk, M. *et al.*, *Org. Prep. Proced. Int.*,2001, **33**, 477-485 (*synth, ir, pmr*)Wu, C. *et al.*, *Tetrahedron*, 2001, **57**, 9575-9583(*biochem, bibl*)Teranishi, K. *et al.*, *Bioorg. Chem.*, 2007, **35**,82-111 (*rev*)**Coelibactin****C-561**Partial struct. shown. Compd. predicted genomically to be prod. by *Streptomyces coelicolor* A3(2). Siderophore.Bentley, S.D. *et al.*, *Nature (London)*, 2002, **417**, 141-147 (*occur*)**Coelobillardierine****C-562****6,7-Dihydro-7-methyl-5H-2-pyrindin-5-ol, 9CI**

[119365-15-0]



Relative Configuration

C₉H₁₁NO 149.192Alkaloid from stems and leaves of *Coelospermum billardieri*. Mp 156°. [α]_D²³ -9 (c, 0.7 in CHCl₃).**Ketone: 6,7-Dihydro-7-methyl-5H-pyrindin-5-one, 9CI. Coelosperminone**

[119308-94-0]

C₉H₉NO 147.176Alkaloid from stems and leaves of *Coelospermum billardieri*. Oil. [α]_D²³ +19 (c, 1 in CHCl₃).**7,8-Didehydro: 6,7-Dihydro-7-methylene-5H-2-pyrindin-5-ol, 9CI. 7,8-Dehydro-coelobillardierine**

[119308-95-1]

C₉H₉NO 147.176Alkaloid from stems and leaves of *Coelospermum billardieri*. Mp 140°. [α]_D²³ +41 (c, 1 in CHCl₃).**Stereoisomer (?): Leptorhabine**

[55727-36-1]

C₉H₁₁NO 149.192Alkaloid from the above-ground parts of *Leptorhabdos parviflora* (Scrophulariaceae). Liq. [α]_D²⁰ +10 (c, 1.015 in CHCl₃). Stereochem. undefined. The racemate has also been reported.**(±)-form****Ketone: Aucubinine B**

[129893-31-8]

Microbial transformation prod.

Ketone, 8-hydroxy: 6,7-Dihydro-7-(hydroxymethyl)-5H-2-pyrindin-5-one, 9CI. Aucubinine A


[129722-96-9]

C₉H₉NO₂ 163.176

Microbial transformation prod.

Kadyrov, Kh.A. *et al.*, *Khim. Prir. Soedin.*,1974, **10**, 683; 1975, **11**, 269; *Chem. Nat.**Compd. (Engl. Transl.)*, 1974, **10**, 711; 1975,**11**, 286 (*Leptorhabine*)Lopez, J.L. *et al.*, *J. Nat. Prod.*, 1988, **51**, 829(*Coelobillardierine, Coelosperminone, Dehydrocoelobillardierine*)Hattori, M. *et al.*, *Phytother. Res.*, 1990, **4**, 66(*Aucubinines*)Baghdikian, B. *et al.*, *Planta Med.*, 1999, **65**,164-166 (*Aucubinine B*)Yang, X.-X. *et al.*, *Chin. J. Chem.*, 2003, **21**,970-971 (*Aucubinine B, synth*)**Cohirsine****C-563**


[113807-36-6]

C₂₀H₂₅NO₄ 343.422Alkaloid from *Cocculus hirsutus* (Menispermaceae). Gum. [α]_D²⁶ +147 (CHCl₃).**O²,O³-Di-de-Me: Shaheenine**


[138704-13-9]

C₁₈H₂₁NO₄ 315.368Alkaloid from *Cocculus hirsutus* (Menispermaceae). Gum. [α]_D²⁵ +125 (CHCl₃).Ahmad, V.U. *et al.*, *Tetrahedron*, 1987, **43**,5865 (*isol, uv, ir, pmr, cmr, ms, struct*)Rasheed, T. *et al.*, *Fitoterapia*, 1991, **62**, 157(*Shaheenine*)**Cohirsinine****C-564****1,2,3,5,6,7-Hexahydro-11-hydroxy-2,12-dimethoxydibenzo[b,i]quinolizin-9(14H)-one, 9CI**


[135091-06-4]

C₁₉H₂₃NO₄ 329.395Alkaloid from *Cocculus hirsutus*. [α]_D²⁵ +136 (CHCl₃). λ_{max} 213; 240; 303 (MeOH).Ahmad, V.U. *et al.*, *Phytochemistry*, 1991, **30**, 1350-1351 (*isol, pmr, cmr, ms*)**Cohirsitine****C-565**

[148717-82-2]

C₂₄H₂₉NO₇ 443.496Alkaloid from aerial parts of *Cocculus hirsutus* (Menispermaceae). [α]_D²⁷ +147 (CHCl₃).Ahmad, V.U. *et al.*, *Fitoterapia*, 1992, **63**, 308(*isol, uv, ir, pmr, cmr, ms, struct*)**Cohirsitinine****C-566**


[142717-67-7]



Relative configuration

C₁₈H₂₃NO₃ 301.385A ring-contracted analogue of Cohirsinine, C-564. Alkaloid from whole plants of *Cocculus hirsutus* (Menispermaceae). Gum. [α]_D²⁵ +51 (CDCl₃).Ahmad, V.U. *et al.*, *J. Nat. Prod.*, 1992, **55**, 237(*isol, uv, ir, pmr, cmr, ms, struct*)**Colchibiphenylene****C-567****N-(6,7-Dihydro-4,9-dihydroxy-3,10,11-trimethoxy-5H-dibenzo[a,c]cyclohepten-5-yl)acetamide, 9CI**

[129724-65-8]



C₂₀H₂₃NO₆ 373.405**(S)-form**

Alkaloid from *Colchicum ritchii* (Liliaceae). Amorph. [α]_D²⁵ -25.5 (c, 0.094 in MeOH). Mixt. of conformers in soln.

O³-Me: Androbiphenylene

[126223-60-7]

C₂₁H₂₅NO₆ 387.432

Alkaloid from the bulbs of *Colchicum ritchii* and *Androcymbium palaestinum* (Liliaceae). Amorph. [α]_D²⁵ -48 (c, 0.1 in MeOH).

Tojo, E. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1163 (*Androbiphenylene*)


Al-Tel, T.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 623 (*isol, pmr, struct*)

Brecht, R. *et al.*, *Liebigs Ann./Recl.*, 1997, 2275-2279 (*synth, Androbiphenylene*)

Colchiciline

6-Hydroxycolchicine

[61036-87-1]

C₂₂H₂₅NO₇ 415.442

Alkaloid from the seeds of *Colchicum latifolium*, *Colchicum doerfleri*, *Colchicum autumnale*, and *Colchicum visianii*; commercial samples of colchicine may contain colchiciline as a minor impurity (Liliaceae). Yellow cryst. (EtOAc). Mp 170-171° dec. [α]_D²² -121 (c, 1.93 in CHCl₃).

O-Ac:

Cryst. (EtOAc). Mp 150-151°.

N-De-Ac, N-formyl: Gloriosamine D

[1006585-05-2]

C₂₁H₂₃NO₇ 401.415

Alkaloid from *Gloriosa rothschildiana*. Amorph. solid. [α]_D²⁵ -95 (c, 0.06 in CHCl₃). λ_{\max} 233 ; 244 ; 353 (EtOH).

N-De-Ac, N-(hydroxyacetyl): Gloriosamine C

[1006585-04-1]

C₂₂H₂₅NO₈ 431.441

Alkaloid from *Gloriosa rothschildiana*. Amorph. solid. [α]_D²⁴ -156 (c, 0.09 in CHCl₃). λ_{\max} 245 ; 355 (EtOH).

O²-De-Me: 2-Demethylcolchicine

[61036-88-2]

C₂₁H₂₃NO₇ 401.415

Alkaloid from the seeds of *Colchicum latifolium* (Liliaceae). Amorph. Tentative struct. based on colour reactions.

Isomer: Alkaloid CC 12

[5083-61-4]

[1348-81-8]

C₂₂H₂₅NO₇ 415.442

Alkaloid from seeds of *Colchicum cornigerum* (Liliaceae) and *Colchicum latifolium*. Yellowish needles (Me₂CO). Mp 197-199°. [α]_D²² -45 (c, 0.242 in CHCl₃). [α]_D²² -83 (c, 0.51 in MeOH). A hydroxycolchicine of unknown regiochemistry.

Saleh, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413-3423 (*Alkaloid CC 12*)

Cross, A.D. *et al.*, *Coll. Czech. Chem. Comm.*, 1966, **31**, 374-378 (*Alkaloid CC 12, pmr*)

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3146-3156; 1977, **42**, 1571-1580 (*Colchiciline, 2-Demethylcolchicine, isol, uv, ir, pmr, cmr, ms, struct*)

Iorio, M.A. *et al.*, *Heterocycles*, 1980, **14**, 625 (*isol, ms*)


Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1982, **47**, 2258 (*cd*)

Kitajima, M. *et al.*, *Tet. Lett.*, 2008, **49**, 257-260 (*Gloriosamines C,D*)

Colchicine, JAN, USAN

C-569

Colchiceine methyl ether. Artrichine. Colchisol. Colsaloid. NSC 757. Many other names

C₂₂H₂₅NO₆ 399.443

Colchicine has one chiral centre (7*S*- in the natural enantiomer) and an axis of chirality. The barrier to interconversion of the atropisomers is approx. 23 kcal mol⁻¹ which is too low to allow separate characterisation for colchicine itself, although isomers of various close relatives have been isolated, and de(acetamido)colchicine can be resolved. The abs. config. of the chiral axis is (*aR*) as shown (axial chirality), which was incorrectly described for some time as *S*- using an incorrect application of the Cahn-Ingold-Prelog rules. Log P -0.21 (uncertain value) (calc).

(+)-form [75520-89-7]Synthetic. Cryst. (H₂O). Mp 139-141°.[α]_D²⁵ +131 (c, 0.87 in CHCl₃).

▶ GH0705000

(-)-form [64-86-8]

Chief alkaloid from *Colchicum autumnale*, many other *Colchicum* spp., several *Merendera* spp., *Gloriosa superba* and others (Liliaceae). Functions by binding to tubulin and arresting cell division at metaphase. Also inhibits many cell transport systems. Antimitotic agent used to induce polyploidy in plant breeding and for studying cell division. Toxicity has precluded exploitation of these props. for antineoplastic use. Potential HIV inhibitor. Antiinflammatory agent. Used for relief of pain in acute gout. Sol. MeOH, C₆H₆, H₂O; fairly sol. Et₂O; poorly sol. hexane. Mp 155-157°. [α]_D²² -121 (CHCl₃). Log P -0.21 (uncertain value) (calc). Pharmacol. active isomer. λ_{\max} 243 (ε 18600); 350 (ε 16700) (EtOH) (Berdy).

▶ Gastrointestinal effects when used therapeutically. Other adverse effects reported incl. a lethal dose of 7 mg in humans. Severe eye irritant. *Colchicum*

spp. bulbs are freq. cause of accidental poisoning. Exp. reprod. and teratogenic effects. LD₅₀ (mus, orl) 6 mg/kg; LD₅₀ (mus, ipr) 2 mg/kg. GH0700000

Tetrachloroaurate: Mp 209°.

(±)-form [54192-66-4]

Mp 277-279°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 361C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1374C (nmr)

Dewar, M.J.S. *et al.*, *Nature (London)*, 1945, **155**, 141 (*struct*)

King, M.V. *et al.*, *Acta Cryst. B*, 1952, **5**, 437-440 (*cryst struct*)

Corrodi, H. *et al.*, *Helv. Chim. Acta*, 1955, **38**, 2030-2033 (*abs config*)

Schreiber, J. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 540-597 (*synth, uv, ir*)

Wilson, J.M. *et al.*, *Tetrahedron*, 1963, **19**, 2225-2231 (*ms*)

Ulubelen, A. *et al.*, *Planta Med.*, 1978, **34**, 216-217 (*isol, uv, ir, pmr, ms*)

Hufford, C.D. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 50-56 (*cmr*)

Wyatt, D.K. *et al.*, *Anal. Profiles Drug Subst.*, 1981, **10**, 139-182 (*ir, uv, pmr, cmr, anal*)

Rösner, M. *et al.*, *J. Med. Chem.*, 1981, **24**, 257-261 (*(+)-form, synth*)

Evans, D.A. *et al.*, *J.A.C.S.*, 1981, **103**, 5813-5821 (*synth*)

Capraro, H.-G. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 1-70 (*rev, pharmacol*)

Boger, D.L. *et al.*, *J.O.C.*, 1985, **50**, 3425-3428 (*synth*)

Dumont, R. *et al.*, *J.O.C.*, 1986, **51**, 2515-2521 (*synth*)

Brossi, A. *et al.*, *Med. Res. Rev.*, 1988, **8**, 77-94 (*rev, props*)

Brossi, A. *et al.*, *FEBS Lett.*, 1990, **262**, 5-7 (*cd, abs config*)

Lee, K.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 632-637 (*anti-HIV activity*)

Muzaffar, A. *et al.*, *Pharmacol. Ther.*, 1991, **49**, 105-109 (*rev*)

Levy, M. *et al.*, *Pharmacotherapy (Carlisle, Mass.)*, 1991, **11**, 196; 1994, **12**, 171 (*pharmacol, tox, rev*)

Textbook of Adverse Drug Reactions, 4th edn., (ed. Davies, D.M.), Oxford University Press, 1991,

Banwell, M.G. *et al.*, *J.C.S. Perkin 1*, 1992, 1415-1426 (*synth*)

Hood, R.L. *et al.*, *J. Emergency Med.*, 1994, **12**, 171 (*tox, rev*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 8951 (*synonyms*)

Maier, U.H. *et al.*, *Tet. Lett.*, 1997, **38**, 7357-7360 (*biosynth*)

Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1998, 2979-2987; 2989-2994; 2995-3001; 3003-3009 (*biosynth, bibl*)

Le Hello, C. *et al.*, *Alkaloids (Academic Press)*, 1999, **53**, (*rev, pharmacol*)

Berg, U. *et al.*, *Helv. Chim. Acta*, 1999, **82**, 323-325 (*stereochem*)

Brossi, A. *et al.*, *Helv. Chim. Acta*, 1999, **82**, 1223-1224 (*stereochem*)

Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 393

Lee, J.C. *et al.*, *Tetrahedron*, 2000, **56**, 10175-10184 (*synth, bibl*)

Graening, T. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 3230-3256 (*rev, synth*)

Graening, T. *et al.*, *Org. Lett.*, 2005, **7**, 4317-4329 (*synth*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, CNG830

Colchiethanamine

C-570

1,2,3,4-Tetrahydro-1-[2-(4-hydroxyphenyl)ethyl]-7-methoxy-2-methyl-6-isoquinolinol, 9CI



C₁₉H₂₃NO₃ 313.396

(S)-form [129724-55-6]

Alkaloid from the bulbs of *Colchicum szovitsii* (Liliaceae). [α]_D +10 (c, 0.2 in MeOH).

O^{4'}-Me: Colchiethine

[129724-56-7]

C₂₀H₂₅NO₃ 327.422


Alkaloid from the bulbs of *Colchicum szovitsii* (Liliaceae). [α]_D +8 (c, 0.2 in MeOH).

Tojo, E. et al., *J. Nat. Prod.*, 1990, **53**, 634 (isol, pmr, struct)

Colchifoline

C-571

[74515-40-5]



C₂₂H₂₅NO₇ 415.442

Minor alkaloid from commercial samples of Colchicine. Isol. from *Colchicum autumnale* and *Androcymbium gramineum*. Cryst. (EtOAc). Mp 151-152°. [α]_D²² -149 (c, 0.94 in CHCl₃).

▶ AC4010000

O²-De-Me: 2-Demethylcolchifoline

[70553-72-9]

C₂₁H₂₃NO₇ 401.415

Isol. from the leaves of *Colchicum autumnale* (Liliaceae).

O²-De-Me, O²-Ac: 2-Acetyl-2-demethylcolchifoline

[70553-74-1]

C₂₃H₂₅NO₈ 443.452

Noncryst. [α]_D²³ -112 (c, 0.52 in CHCl₃).

O²-De-Me, di-Ac:

Cryst. (EtOAc). Mp 229-232°. [α]_D²³ -109 (c, 0.68 in CHCl₃).

O¹⁰-De-Me: Colchifoleine

[78517-64-3]

C₂₁H₂₃NO₇ 401.415

Cryst. (MeOH aq.). Mp 227-228°. [α]_D²² -254 (c, 0.65 in CHCl₃).

Sedmera, P. et al., *Heterocycles*, 1979, **12**, 337-342 (2-Demethylcolchifoline)

Iorio, M.A. et al., *Heterocycles*, 1980, **14**, 625-630 (ms)


Iorio, M.A. et al., *Can. J. Chem.*, 1981, **59**,

283-284 (synth, pmr)

Ellington, E. et al., *Biochem. Syst. Ecol.*, 2003, **31**, 715-722 (isol)

Colchilutine

C-572



C₁₉H₂₃NO₃ 313.396


Alkaloid from *Colchicum luteum* (Liliaceae). Mp 190-191°. [α]_D -83 (CHCl₃).

Shakirov, R. et al., *Khim. Prir. Soedin.*, 1996, **32**, 410-512; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 386-512

Colchiritchine

C-573

[111509-14-9]



C₂₀H₂₁NO₅ 355.39

Alkaloid from *Colchicum ritchii*.

Amorph. [α]_D +207 (c, 0.15 in MeOH). λ_{max} 213 (log ε 4.41); 241 (log ε 4.08); 279 (log ε 3.7) (MeOH).


Freyer, A.J. et al., *J. Nat. Prod.*, 1987, **50**, 684-689 (isol, uv, ir, pmr, ms, struct)

Collismycin

C-574

4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-carboxaldehyde oxime. *SF* 2738. *Antibiotic SF* 2738

[149759-19-3]



(E)-form

C₁₃H₁₃N₃O₂S 275.331

(E)-form

Collismycin A. *SF* 2738A. *Antibiotic SF* 2738A

[158792-24-6]

Prod. by *Streptomyces* sp. MQ22 and *Streptomyces* sp. *SF* 2738. Inhibitor of dexamethasone-glucocorticoid receptor binding. Antimicrobial and antitumour agent. Cryst. Mp 174-176° (170-172° dec.). Similar to Caerulomycin, C-9. λ_{max}

244 (ε 26400) (MeOH) (Derep).

S-Oxide: Pyrisulfoxin A. *BS* 75A. *Antibiotic BS* 75A

[187337-07-1]

C₁₃H₁₃N₃O₃S 291.33

Prod. by *Streptomyces californicus* sp. BS75A. Antitumour agent. Powder or yellow powder. Mp 178-180°. λ_{max} 243 (ε 23500); 288 (ε 14500) (MeOH).

(Z)-form

Collismycin B. *SF* 2738B. *Antibiotic SF* 2738B

[158792-25-7]

Prod. by *Streptomyces* sp. MQ22. Inhibitor of dexamethasone-glucocorticoid receptor binding. Cytotoxic agent. Cryst. Mp 148-159° dec. (140-142°). λ_{max} 244 (ε 26400) (MeOH) (Derep).

Japan. Pat., 1993, 93 78 322; *CA*, **119**, 135507 (isol, pmr, cmr, ir, uv)

Shindo, K. et al., *J. Antibiot.*, 1994, **47**, 1072-1074 (isol)

Gomi, S. et al., *J. Antibiot.*, 1994, **47**, 1385-1394 (isol, pmr, cmr)

Japan. Pat., 1997, 97 12 550; *CA*, **126**, 170489d (BS 75A)

Tsuge, N. et al., *J. Antibiot.*, 1999, **52**, 505-507 (*Pyrisulfoxin A*)


Stadler, M. et al., *Arch. Pharm. (Weinheim, Ger.)*, 2001, **334**, 143-147 (isol, pmr)

Collismycin C

C-575

4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-methanol. 6-Hydroxymethyl-4-methoxy-5-(methylthio)-2,2'-bipyridine. *Antibiotic SF* 2738C. *SF* 2738C

[161504-74-1]



R = CH₂OH

C₁₃H₁₄N₂O₂S 262.332

Prod. by *Streptomyces* sp. *SF* 2738.

Active against gram-positive and -negative bacteria, fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 104-106°. Similar to Caerulomycin, C-9. λ_{max} 206 (E1%/1cm 720); 219 (E1%/1cm 719); 292 (E1%/1cm 538) (MeOH) (Berdy). λ_{max} 211 (E1%/1cm 646); 227 (E1%/1cm 583); 260 (E1%/1cm 544); 272 (E1%/1cm 492); 306 (E1%/1cm 470) (MeOH/HCl) (Berdy). λ_{max} 215 (E1%/1cm 1260); 291 (E1%/1cm 550) (MeOH/NaOH) (Berdy).

I'-Aldehyde, oxime: see Collismycin, C-574

Gomi, S. et al., *J. Antibiot.*, 1994, **47**, 1385-1394 (isol, uv, ir, pmr, cmr, props)

Collismycin D

C-576

4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-carbonitrile. 6-Cyano-4-methoxy-5-(methylthio)-2,2'-bipyridine. *SF* 2738D. *Antibiotic SF* 2738D

[161504-75-2]

As Collismycin C, C-575 with

R = -CN

C₁₃H₁₁N₃OS 257.315

Prod. by *Streptomyces* sp. SF 2738.
Active against gram-positive and -negative bacteria, fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 123-125°. λ_{max} 206 (E1%/1cm 868); 236 (ε 1200); 288 (E1%/1cm 591) (MeOH) (Berdy).

S-Oxide: Pyrisulfoxin BC₁₃H₁₁N₃O₂S 273.315

Prod. by *Streptomyces californicus*. Powder. Mp 163-165°. λ_{max} 243 (ε 17900); 288 (ε 12700) (MeOH).
Gomi, S. et al., *J. Antibiot.*, 1994, **47**, 1385-1394 (isol, uv, ir, pmr, cmr, props)
Tsuge, N. et al., *J. Antibiot.*, 1999, **52**, 505-507 (Pyrisulfoxin B)

Collismycin E

C-577

Antibiotic SF 2738E. SF 2738E

[161504-76-3]

As Collismycin C, C-575 with R = -CH₂NHAcC₁₅H₁₇N₃O₂S 303.384


Prod. by *Streptomyces* sp. SF 2738.
Active against gram-positive and -negative bacteria, fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 123-125°. λ_{max} 206 (E1%/1cm 759); 287 (E1%/1cm 476) (MeOH) (Berdy).

Gomi, S. et al., *J. Antibiot.*, 1994, **47**, 1385-1394 (isol, uv, ir, pmr, cmr, props)

Collismycin F

C-578

7-Methoxy-5-(2-pyridinyl)isothiazolo[4,5-b]pyridine, 9CI. Antibiotic SF 2738F. SF 2738F
[161504-77-4]

C₁₂H₉N₃OS 243.289


Prod. by *Streptomyces* sp. SF 2738.
Active against gram-positive and -negative bacteria, fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 145-148°. λ_{max} 203 (E1%/1cm 472); 248 (E1%/1cm 930); 252 (E1%/1cm 925); 272 (E1%/1cm 572) (MeOH) (Berdy).

Gomi, S. et al., *J. Antibiot.*, 1994, **47**, 1385 (isol, uv, ir, pmr, cmr, props)

Colubricidin A

C-579

[257939-57-4]

C₉₆H₁₅₄Cl₃N₅O₄₂ 2156.639Macrolide antibiotic. Prod. by *Strepto-*

myces sp. LL-C13122. Active against gram-positive bacteria. Yellowish powder. [α]_D²⁵ -7 (c, 0.5 in MeOH). λ_{max} 280 (sh); 375 (no solvent reported).

Kong, F. et al., *Tet. Lett.*, 1999, **40**, 9219-9223

Columellarine

C-580

[11023-69-1]

C₂₁H₂₅N₄O₄ 355.433


Struct. unknown. No further work to 2001. Alkaloid from *Baometra columellaris*. Mp 150-153°. [α]_D²² -201 (c, 0.937 in CHCl₃).

Pijewska, L. et al., *Coll. Czech. Chem. Comm.*, 1967, **32**, 158-170 (isol, uv)

Communesin A

C-581

[148439-45-6]



Relative Configuration

R = COCH₃C₂₈H₃₂N₄O₂ 456.586

Communesin alkaloids have been isolated from *Penicillium* spp. by two different groups of workers simultaneously, and various Communesin names have been given to different but closely similar compounds. See also under Communesin B, C-582. Metab. from the mycelium of a strain of *Penicillium* sp. isolated from the marine alga *Enteromorpha intestinalis*. Also prod. by the terrestrial *Penicillium expansum* Link MK-57. Cytotoxic and insecticidal agent. Amorph. powder. Mp 194-196° Mp > 300°. [α]_D²⁰ -174 (c, 1.34 in CHCl₃). λ_{max} 208 (ε 53700); 247 (ε 12300); 268 (ε 11500); 290 (sh) (ε 3980); 315 (ε 2880) (EtOH) (Derep).

N-De-Me: **Communesin E**†. **Communesin D**†

[727416-58-2]

C₂₇H₃₀N₄O₂ 442.56

Prod. by *Penicillium expansum* Link MK-57. Insecticidal agent. Amorph. powder. Mp 250° dec. [α]_D²⁰ -156 (c, 0.11 in CHCl₃). λ_{max} 243 (ε 9700) (MeOH).

21,22-Deepoxy, 21,22-didehydro: **Communesin F**. **Communesin E**†

[727416-60-6]

C₂₈H₃₂N₄O 440.587

Prod. by *Penicillium expansum* Link MK-57. Insecticidal agent. Amorph. powder. Mp 144-147°. [α]_D²⁰ -264 (c, 0.34 in CHCl₃). λ_{max} 268 (ε 10900) (MeOH).

Numata, A. et al., *Tet. Lett.*, 1993, **34**, 2355-2358 (isol, ir, pmr, cmr)

Hayashi, H. et al., *Biosci., Biotechnol., Biochem.*, 2004, **68**, 753-756 (isol, pmr, cmr, ms)

Wigley, L.J. et al., *Phytochemistry*, 2006, **67**, 561-569 (biosynth)

May, J.A. et al., *Tetrahedron*, 2006, **62**, 5262-5271 (biosynth, struct)

Yang, J. et al., *J.A.C.S.*, 2007, **129**, 13794-13795 (Communesin F, synth)

Communesin B

C-582

Nomofungin

[148439-46-7]

As Communesin A, C-581 with R = H₂C=CHCH=CHCH₂CO-

C₃₂H₃₆N₄O₂ 508.662

Struct. of Nomofungin revised in 2003. See comment under Communesin A, C-581. Metab. from the mycelium of a strain of *Penicillium* sp. isolated from the marine alga *Enteromorpha intestinalis* and from the terrestrial *Penicillium expansum* Link MK-57. Isol. from an unidentified fungus from the bark of *Ficus microcarpa* (Nomofungin). Cytotoxic, mycotoxin and insecticide. Amorph. powder. Mp 165-170° (152-154°). [α]_D²⁰ -74.9 (c, 1.5 in CHCl₃). [α]_D²² +8.7 (c, 0.2 in CHCl₃). λ_{max} 208 (ε 45700); 250 (sh) (ε 32400); 266 (ε 38900); 315 (ε 3090) (EtOH) (Derep).

N-De-Me: **Communesin C**†
[648413-35-8]

C₃₁H₃₄N₄O₂ 494.635

Prod. by a *Penicillium* sp. isolated from the marine sponge *Axinella verrucosa*. [α]_D²⁰ -30 (c, 0.04 in MeOH). λ_{max} 206; 271 (MeOH).

N-De-Me, N¹⁵-formyl: **Communesin D**†. **Communesin C**†

[648413-36-9]

C₃₂H₃₄N₄O₃ 522.646

Prod. by a *Penicillium* sp. isolated from *Axinella verrucosa* and by the terrestrial *Penicillium expansum* Link MK-57. Amorph. powder. Mp 190-195°. [α]_D²⁰ +23.3 (c, 0.04 in MeOH). [α]_D²⁰ +150 (c, 0.14 in CHCl₃). λ_{max} 206; 267 (MeOH).

Numata, A. et al., *Tet. Lett.*, 1993, **34**, 2355-2358 (isol, pmr, cmr, struct)

Ratnayake, A.S. et al., *J.O.C.*, 2001, **66**, 8717-8721; 2003, **68**, 1640 (Nomofungin)

May, J.A. et al., *Tet. Lett.*, 2003, **44**, 1203-1205 (struct)

Hayashi, H. et al., *Biosci., Biotechnol., Biochem.*, 2004, **68**, 753-756 (isol, pmr, cmr, ms)

Jadulco, R. et al., *J. Nat. Prod.*, 2004, **67**, 78-81 (isol, pmr, cmr, ms)

Wigley, L.J. et al., *Phytochemistry*, 2006, **67**, 561-569 (biosynth)

May, J.A. et al., *Tetrahedron*, 2006, **62**, 5262-5271 (biosynth, struct)

Communesin G

C-583

[848950-21-0]

As Communesin A, C-581 with

R = COCH₂CH₃C₂₉H₃₄N₄O₂ 470.613

Prod. by *Penicillium rivulum* (IBT 24420). Amorph. powder (MeCN aq.). Mp 162-166°. [α]_D²⁵ -157 (c, 0.02 in MeOH). λ_{max} 207 (log ε 4.59); 248 (log ε 3.92); 268 (log ε 3.94); 316 (log ε 3.39) (MeOH).

Dalsgaard, P.W. et al., *J. Nat. Prod.*, 2005, **68**, 258-261 (isol, cd, pmr, cmr)


Communesin H C-584

[848950-22-1]
As Communesin A, C-581 with
R = COCH₂CH₂CH₃
C₃₀H₃₆N₄O₂ 484.64
Prod. by *Penicillium rivulum* (IBT 24420).
Amorph. powder (MeCN aq.). Mp 143-
147°. [α]_D²⁵ -167 (c, 0.02 in MeOH). λ_{\max}
208 (log ϵ 4.68); 248 (log ϵ 4); 268 (log ϵ
4.02); 316 (log ϵ 3.48) (MeOH).

Dalsgaard, P.W. *et al.*, *J. Nat. Prod.*, 2005, **68**,
258-261 (*isol, cd, pmr, cmr*)

Comoramide A C-585

[217449-19-9]




C₃₄H₄₈N₆O₆S 668.856

Cyclic peptide alkaloid. Isol. from *Didemnum molle*. Cytotoxic agent. Amorph.
powder. [α]_D +0.5 (c, 0.17 in MeOH).

Rudi, A. *et al.*, *Tetrahedron*, 1998, **54**, 13203-
13210 (*isol, pmr, cmr, ms*)

Comoramide B C-586

[217449-22-4]




C₃₄H₅₀N₆O₇S 686.871

Cyclic peptide alkaloid. Isol. from *Didemnum molle*. Cytotoxic agent. Amorph.
powder. [α]_D -100 (c, 0.05 in MeOH).

Rudi, A. *et al.*, *Tetrahedron*, 1998, **54**, 13203-
13210 (*isol, pmr, cmr, ms*)

Comosine C-587

3-Epi-6,7-dihydrochelhammeridine. 6,7-
Dihydrohomoerythraline
[31689-96-0]



C₁₉H₂₃NO₃ 313.396

Config. revised in 1986. The name
Comosine was proposed for this alkaloid
in 1986. Alkaloid from the leaves of
Phelline comosa (Phellinaceae). Non-
cryst. [α]_D +75 (c, 1.5 in CHCl₃).

Hydrochloride:

Cryst. (MeOH/Et₂O). Mp 260°. [α]_D
+100 (c, 1.7 in EtOH).

3-Epimer: 3-Epicomosine. 6,7-Dihy-
drochelhammeridine. Alkaloid A†
[21030-73-9]

C₁₉H₂₃NO₃ 313.396

Alkaloid from *Schelhammera pedun-
culata* (Liliaceae). Gum. [α]_D -100 (c,
0.37 in CHCl₃).

3-Epimer: picrate:

Yellow prisms (MeOH). Mp 188-189°.

Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1969, **22**,
2187 (*isol, epimer*)


Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**,
2219 (*uv, pmr, ms, struct, epimer*)

Langlois, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970,
3535 (*isol, uv, ir, pmr, ms, struct*)

Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**,
515 (*synth, config*)

Comosivine C-588

O-Methylathrocupressine. Alkaloid 5
[31690-00-3]



Absolute
Configuration

C₂₁H₂₉NO₄ 359.464

Alkaloid from *Phelline comosa*, *Athro-
taxis cupressoides* and *Lagarostrobos
colensoi* (*Dacrydium colensoi*). Cryst.
(hexane). Mp 100-101°. [α]_D +91 (c, 1.5
in CHCl₃).

1 α ,6 α -Epoxide: 1,6-Epoxycomosivine

[123064-73-3]

C₂₁H₂₉NO₅ 375.464

Alkaloid from the leaves of *Phelline
comosa* var. *robusta* (Phellinaceae). Mp
136°. [α]_D +103 (c, 0.55 in CHCl₃).

O³-De-Me: Robustivine

[123064-72-2]

C₂₀H₂₇NO₄ 345.438

Alkaloid from the leaves of *Phelline
comosa* var. *robusta* (Phellinaceae). Mp
234° (as hydrochloride). [α]_D +103.7 (c,
0.7 in CHCl₃).

O³-De-Me, 1 α ,6 α -epoxide: 1,6-Epoxyro-
bustivine

[749864-97-9]

C₂₀H₂₇NO₅ 361.437

Alkaloid from leaves of *Phelline
comosa*. [α]_D +99 (c, 0.68 in CHCl₃).
 λ_{\max} 207 (ϵ 35700); 283 (ϵ 1275)
(EtOH).

O-De-Me: Holidinine

[95066-36-7]

C₂₀H₂₇NO₄ 345.438

Trace alkaloid from the leaves of
Phelline sp. aff. *Phelline lucida* (Phelli-
naceae) and *Lagarostrobos colensoi*

(*Dacrydium colensoi*). Cryst. (Et₂O).
Mp 164-165°. [α]_D +91 (c, 1.03 in
CHCl₃). Posn. of OH not definitely
establ.

O¹⁸-De-Me: Athrocupressine

[93888-46-1]

C₂₀H₂₇NO₄ 345.438

Alkaloid from the twigs and foliage of
Athrotaxis cupressoides and *Athrotaxis
selaginoides* (Taxodiaceae). Cryst.
(Me₂CO). Mp 152-153°. [α]_D¹⁹ +102.2
(c, 0.43 in CHCl₃).

O³,O¹⁶-Di-de-Me: Robusticene

[128508-54-3]

C₁₉H₂₅NO₄ 331.411

Minor alkaloid from leaves of *Phelline
comosa* var. *robusta* (Phellinaceae). [α]_D
+100 (c, 0.83 in CHCl₃).

O¹⁶,O¹⁸-Di-de-Me: 18-De-O-methylholi-
nidine

[749864-99-1]

C₁₉H₂₅NO₄ 331.411

Alkaloid from leaves of *Phelline co-
mosa*. Mp 140°. [α]_D +111 (c, 0.54 in
MeOH). λ_{\max} 208 (ϵ 29600); 276 (ϵ
1242) (EtOH).

2 α -Hydroxy: 2-Hydroxycomosivine

[138909-10-1]

C₂₁H₂₉NO₅ 375.464

Alkaloid from the heartwood of *Phel-
line comosa* (Phellinaceae). Dull white
cryst. (EtOH). Mp 196-197°. [α]_D²⁵ +93
(c, 1.5 in EtOH).

2 α -Methoxy: 2-Methoxycomosivine

[155272-47-2]

C₂₂H₃₁NO₅ 389.491

Alkaloid from stems of *Dysoxylum
lenticellare* (Meliaceae). Brown oil.

Langlois, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970,
3535-3543 (*Comosivine*)

Langlois, N. *et al.*, *Heterocycles*, 1984, **22**,
2453-2457; 1990, **30**, 659-664 (*Holidinine,
Robusticene*)

Panichanun, S. *et al.*, *Tetrahedron*, 1984, **40**,
2677-2684; 2685-2689 (*Comosivine*)

Pusset, J. *et al.*, *Phytochemistry*, 1989, **28**,
1298-1300 (*Robustivine, 1,6-
Epoxycomosivine, Holidinine*)


Aladesanmi, A.J. *et al.*, *Phytochemistry*, 1991,
30, 3497-3498; 1994, **35**, 1361-1362 (*2-
Hydroxycomosivine, 2-Methoxycomosivine*)

Bloor, S.J. *et al.*, *Phytochemistry*, 1996, **41**, 801-
802 (*Holidinine, Comosivine*)

Langlois, N. *et al.*, *C. R. Chim.*, 2004, **7**, 51-55
(*1,6-Epoxyrobustivine, 18-De-O-
methylholidinine*)

Compactinervine C-589

Methyl 2,16-didehydro-19,20-dihydroxy-
curan-17-oate, 9CI
[2270-73-7]



Absolute
Configuration

C₂₀H₂₄N₂O₄ 356.421

Alkaloid from the bark of *Aspidosperma
compactinervium*, also from stem bark of
Alstonia lanceolata (Apocynaceae). Sol-
vated needles (EtOH aq.). Mp 110-120°
Mp 235-245° dec. (double Mp). [α]_D²⁵ -515

(c, 0.55 in EtOH). $[\alpha]_D$ -640 (Py). λ_{\max} 237 (log ϵ 3.97); 297 (log ϵ 3.95); 331 (log ϵ 4.15) (EtOH).

Hydrochloride:

Prisms (MeOH/Me₂CO). Mp 191-192° dec.

Picrate:

Yellow prisms (Me₂CO). Mp 219-220° dec.

O,O-Di-Ac: Mp 208° dec. $[\alpha]_D$ -623 (CHCl₃).

11-Methoxy: Alstovine. 11-Methoxycompactinervine
[57576-38-2]

C₂₁H₂₆N₂O₅ 386.447

Alkaloid from the trunk bark of *Alstonia vitiensis* (Apocynaceae). Shows antitumour activity. Fine needles (EtOH). Mp 168°. $[\alpha]_D^{20}$ -502 (c, 1.0 in EtOH). λ_{\max} 261 (log ϵ 3.9); 307 (log ϵ 3.67); 331 (log ϵ 3.99) (EtOH).

11-Methoxy, 19-ketone: Lagumidine
[154849-52-2]

C₂₁H₂₄N₂O₅ 384.431

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. $[\alpha]_D^{26}$ -144.2 (c, 0.48 in MeOH).

12-Methoxy: 12-Methoxycompactinervine
[79659-68-0]

C₂₁H₂₆N₂O₅ 386.447

Alkaloid from the leaves of *Alstonia lenormandii* var. *minutifolia*, the stem bark of *Alstonia lenormandii* var. *lenormandii* and from *Alstonia lanceolata* (Apocynaceae). Cryst. (MeOH) or amorph. solid. Mp 155°. $[\alpha]_D$ -482 (c, 1 in CHCl₃) (-204). Struct. revised in 1986, formerly considered to be 10-methoxy. λ_{\max} 235 ; 288 ; 337 (no solvent reported).

Djerassi, C. *et al.*, *Experientia*, 1963, **19**, 467-469 (*isol, uv, pmr, struct, abs config*)

Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141-1166 (*isol, uv, ord, struct*)

Mamatas-Kalamaras, S. *et al.*, *Phytochemistry*, 1975, **14**, 1637-1639 (*Alstovine*)

Vercauteren, J. *et al.*, *Phytochemistry*, 1981, **20**, 1411-1413 (*10-Methoxycompactinervine*)

Ravao, T. *et al.*, *Phytochemistry*, 1982, **21**, 2160-2161 (*Alstovine, 11-Methoxycompactinervine*)

Verpoorte, R. *et al.*, *Planta Med.*, 1983, **48**, 283 (*isol*)


Legseir, B. *et al.*, *Phytochemistry*, 1986, **25**, 1735-1738 (*12-Methoxycompactinervine*)

Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 249-252; 253-257 (*Lagumidine*)

Complanadine A

C-590

[317802-77-0]



C₃₂H₄₂N₄ 482.711

Dimer of Lycodine, L-324. Alkaloid from *Lycopodium complanatum*. Cytotoxic. $[\alpha]_D^{24}$ +14 (c, 0.3 in MeOH). λ_{\max} 251 (ϵ 12000); 290 (ϵ 14000) (no solvent reported).

N,1 α ,2,3-Tetrahydro: Complanadine D
[919769-28-1]

C₃₂H₄₆N₄ 486.742

Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{21}$ -32 (c, 1 in MeOH). λ_{\max} 209 (ϵ 15100); 272 (ϵ 3900) (MeOH).

6-Oxo: Complanadine B

[849096-84-0]

C₃₂H₄₀N₄O 496.694

Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{24}$ -13 (c, 0.5 in MeOH). λ_{\max} 211 (ϵ 8000); 268 (ϵ 12000) (MeOH).

Kobayashi, J. *et al.*, *Tet. Lett.*, 2000, **41**, 9069-9073 (*Complanadine A*)


Morita, H. *et al.*, *Tetrahedron*, 2005, **61**, 1955-1960 (*Complanadine B*)

Ishuchi, K. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 413-417 (*Complanadine D*)

Complanadine C

C-591

[919769-27-0]



C₃₂H₄₂N₂O₃ 502.695


Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{22}$ -9 (c, 0.2 in MeOH). λ_{\max} 223 (ϵ 4900); 347 (ϵ 1900) (MeOH).

Ishuchi, K. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 413-417 (*isol, pmr, cmr*)

Complanatin†

C-592

4-[(Aminoiminomethyl)(3-methyl-2-butenyl)amino]butanoic acid, 9CI. N-(3-Carboxypropyl)-N-(3-methyl-2-butenyl)-guanidine. 4-(N¹-Prenylguanido)butanoic acid
[142287-95-4]



C₁₀H₁₉N₃O₂ 213.279


Isol. from seeds of *Astragalus complanatus* (Fabaceae).

Xue, Z. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 691; *CA*, **117**, 44537u (*isol, cryst struct*)

ComX_{RO-E-2} pheromone

C-593

[863658-74-6]



C₄₈H₆₆N₈O₁₀ 915.097

Isol. from *Bacillus subtilis* RO-E-2. Quorum-sensing peptide pheromone.


Okada, M. *et al.*, *Nat. Chem. Biol.*, 2005, **1**, 23-24 (*isol, pmr, struct*)

Okada, M. *et al.*, *Tetrahedron*, 2006, **62**, 8907-8918 (*synth*)

ComX_{RO-H-1} pheromone

C-594

[1010806-86-6]



C₅₁H₇₄N₈O₁₀S 991.259


Isol. from *Bacillus mojavensis* RO-H-1. Quorum-sensing peptide pheromone. Extremely acid labile.

Okada, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 1807-1810 (*isol, synth, ms, struct*)

Conarrhimine

C-595

3-Amino-23-norcon-5-enine. 19,20-Epiminopregn-5-en-3-amine
[468-39-3]



C₂₁H₃₄N₂ 314.513

Minor alkaloid from the bark of *Holarhena antidysenterica* (Apocynaceae). Cryst. (EtOH or by subl.). Mp 229-230°. $[\alpha]_D$ +31 (c, 1 in CHCl₃). Difficult to separate, forms a eutectic with Holarhimine (See 3,20-Diaminopregn-5-en-18-ol, D-279).

N³-Me: Conimine

[468-41-7]

C₂₂H₃₆N₂ 328.54

Alkaloid from *Holarhena antidysenterica* (Apocynaceae). Mp 130°. $[\alpha]_D$ -30 (EtOH).

N³-Me; hydrochloride: Mp 318-320° dec.

N³-Me, picrate: Mp 140-141°.

N²³-Me: Conamine. Con-5-enin-3-amine

[468-34-8]

C₂₂H₃₆N₂ 328.54Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 97.5-101.5°. [α]_D -21 (CHCl₃).N³,N³-Di-Me: **Conessimine**

[631-05-0]

C₂₃H₃₈N₂ 342.567Minor alkaloid from *Holarrhena febrifuga* and *Holarrhena antidysenterica* (Apocynaceae). Mp 100° (88-92°). [α]_D -22.3 (CHCl₃).N³,N³-Di-Me; dihydrochloride: Mp 342-344°.N³,N²³-Di-Me: **Isoconessimine**. *Kurchine*. *Norconessimine*

[468-36-0]

C₂₃H₃₈N₂ 342.567Alkaloid from *Funtumia elastica*, *Holarrhena febrifuga*, *Holarrhena antidysenterica* and *Wrightia tomentosa*. Cryst. (Me₂CO), cryst. + 2H₂O (dioxan). Mp 92°. Bp_{0.7} 238-240°. [α]_D +30 (c, 1 in EtOH).N³,N²³-Di-Me; dihydrochloride: [63550-88-9]
Mp 335°.

▶ GK7621150

N³,N³,N²³-Tri-Me: see Conessimine, C-601N²³-Hydroxy, N³,N³-di-Me: **Regholarrhene D**

[128718-00-3]

C₂₃H₃₈N₂O 358.566Alkaloid from stem bark of *Holarrhena antidysenterica* (Apocynaceae). Amorph. solid (EtOAc). Mp 255-257°.5 α ,6-Dihydro, N²³-Me, N³-Ac: **Malouphyllamine**

[7050-49-9]

C₂₄H₄₀N₂O 372.593Alkaloid from the leaves of *Malouetia bequaertiana* (Apocynaceae). Mp 219-220°. [α]_D +42 (CHCl₃).5 α ,6-Dihydro, N³,N²³-di-Me: **Dihydroisoconessimine**

[7050-30-8]

C₂₃H₄₀N₂ 344.582Alkaloid from the bark of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (Me₂CO). Mp 97-98.5°.18-Oxo, N³,N³-di-Me: **Antidysentericine**


[285550-89-2]

C₂₃H₃₆N₂O 356.55Constit. of *Holarrhena antidysenterica*. Cryst. Mp 118-119°. [α]_D²² -24.5 (c, 0.1 in CHCl₃). λ_{\max} 222 (log ϵ 6.5) (MeOH).3-Epimer, N²³-Me: **3-Epiconamine**

[14152-35-3]

C₂₂H₃₆N₂ 328.54Alkaloid from the bark of *Holarrhena antidysenterica*. Cryst. Mp 95-100°. [α]_D +27 (+11) (EtOH). Could not be recryst.Haworth, R.D. *et al.*, *J.C.S.*, 1932, 631-634; 1953, 1102-1109 (*Isoconessimine*, *Dihydroisoconessimine*, *isol*)Siddiqui, S. *et al.*, *J. Indian Chem. Soc.*, 1934, 9, 283-291 (*Isoconessimine*, *Conimine*, *Conessimine*)Tschesche, R. *et al.*, *Chem. Ber.*, 1954, 87, 1719-1725 (*Conessine*, *Conessimine*, *Conimine*)Hora, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1961, 26, 2217-2228 (*Conamine*, *synth*)Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 641-646 (*Malouphyllamine*)Lábler, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, 28, 2015-2020 (*Conessimine*, *synth*)Černý, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, 29, 1591-1597 (*Conarrhimine*, *Dihydroisoconessimine*, *3-Epiconamine*)Dadoun, H. *et al.*, *Ann. Pharm. Fr.*, 1973, 31, 237-247 (*isol*, *Conessine*, *Isoconessimine*, *Conessimine*)Bhutani, K.K. *et al.*, *Phytochemistry*, 1990, 29, 969-972 (*Regholarrhene D*)Siddiqui, B.S. *et al.*, *Heterocycles*, 1995, 41, 267-276 (*isol*, *in pmr, cmr, uv*)Kumar, A. *et al.*, *Fitoterapia*, 2000, 71, 101-104 (*Antidysentericine*)Zirih, G.N. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, 15, 2637-2640 (*Conessine*, *Isoconessimine*)**Concavine**

[870474-81-0]

Relative
ConfigurationC₂₂H₃₅NO 329.525Prod. by *Clitocybe concava*. Oil. [α]_D +8.4 (c, 0.14 in MeOH).Arnone, A. *et al.*, *Tet. Lett.*, 2005, 46, 8037-8039 (*isol*, *pmr, cmr*)**Conchairamidine**

C-597

C₂₂H₂₆N₂O₄ 382.458Struct. unknown. Alkaloid from bark of *Remijia purdieana* (Rubiaceae). Oil; cryst. + 1H₂O. Mp 114-115° (anhyd.). [α]_D¹⁵ -60 (c, 3 in EtOH).Hesse, O. *et al.*, *Annalen*, 1884, 225, 211**Conchairamine**

C-598


C₂₂H₂₆N₂O₄ 382.458Struct. unknown. Alkaloid from bark of *Remijia purdieana* (Rubiaceae). Thick prisms + 1H₂O + 1EtOH (Et₂OH). Mp 120° (anhyd.). [α]_D¹⁵ +68.4 (c, 2 in EtOH).Hesse, O. *et al.*, *Annalen*, 1884, 225, 211**Concusconine**

C-599

[1398-82-9]

C₂₃H₂₆N₂O₄ 394.469Struct. unknown. Alkaloid from the bark of *Remijia purdieana* (Rubiaceae). Monoclinic needles + 1H₂O (EtOH). Mp 144° Mp 206-208° (double Mp). [α]_D¹⁵ +40.8 (c, 2 in EtOH). Conts. 2 OMe groups.Hesse, O. *et al.*, *Annalen*, 1884, 225, 211Howard, B.F. *et al.*, *J. Soc. Chem. Ind.*, London, 1909, 28, 53-57**Condaline A**

C-600

 α -(Methylamino)-N-[7-(1-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]benzenepropanamide
[497172-91-5]Absolute
ConfigurationC₃₃H₃₈N₄O₄ 554.688Stereochemistry of the various alkaloids requires clarification. The phys. props. reported for Condaline A are almost identical with those for the previously known Aralionine B below. Alkaloid from the root bark of *Condalia buxifolia*. Needles (CHCl₃/MeOH). Mp 115-116°. [α]_D²⁵ -73 (c, 0.08 in MeOH).3-Epimer: **Scutianine M**

[872200-75-4]

C₃₃H₃₈N₄O₄ 554.688Alkaloid from the root bark of *Scutia buxifolia*. Powder. Mp 257-259°. [α]_D²⁵ +120 (c, 0.02 in CHCl₃).7'-Epimer, 2''-N-Me: **Adouetine Y**

[19542-38-2]

C₃₄H₄₀N₄O₄ 568.714Alkaloid from *Waltheria americana*, *Ceanothus americanus* and *Discaria americana*. Cryst. (MeOH or CHCl₃/Et₂O). Mp 292°. [α]_D -230 (CHCl₃/MeOH 9:1).Stereoisomer (1), 2''-N-Me: **Scutianine L**

[202394-60-3]

C₃₄H₄₀N₄O₄ 568.714Alkaloid from *Scutia buxifolia*. Cryst. (MeOH/diisopropyl ether). Mp 122-123°. [α]_D²⁵ -72 (c, 2.4 in CHCl₃). Assigned (4S,7S)-config., with unknown C-3 and C-7'-configs.Stereoisomer (2): **Alkaloid AM 2**. **AM 2**

[151379-40-7]

C₃₃H₃₈N₄O₄ 554.688Alkaloid from leaves and branches of *Antidesma montana* (Euphorbiaceae). Fine needles (MeOH). Mp 257-258°. No stereochem. determined. Reported Mp is identical with that of Scutianine M. Stated in 2006 review to be identical with Aralionine B, but this appears to be in error.Stereoisomer (3): **Aralionine B**


[29432-61-9]

C₃₃H₃₈N₄O₄ 554.688Minor alkaloid from the leaves of *Araliorhamnus vaginatus* (Rhamnaceae) and from *Antidesma montana*. Needles (C₆H₆/petrol). Mp 103-105°. [α]_D²⁰ -73 (c, 0.1 in MeOH). No stereochem. determined.Pais, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 1145-1148 (*Adouetine Y*)

- Servis, R.E. *et al.*, *J.A.C.S.*, 1969, **91**, 5619 (*Adouetine Y, isol, ms*)
 Tschesche, R. *et al.*, *Chem. Ber.*, 1970, **103**, 2501-2504 (*Aralionine B*)
 Arbain, D. *et al.*, *Phytochemistry*, 1993, **33**, 1263-1266 (*Alkaloid AM 2*)
 Morel, A.F. *et al.*, *Phytochemistry*, 1998, **47**, 125-129; 2002, **61**, 561-566; 2005, **66**, 2571-2576 (*Scutianine L, Condaline A, Scutianine M*)
 Giacomelli, S.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 997-999 (*Adouetine Y, isol*)

Conessine, INN **C-601**

N,N-Dimethylcon-5-enin-3-amine. 3-(*Dimethylamino*)con-5-enine, 8*Cl*. 3-(*Dimethylamino*-5-conanene. *Wrightine. Roquessine. Neriine* [546-06-5]



$C_{24}H_{40}N_2$ 356.593
 Alkaloid from *Holarrhena antidysenterica* and many other *Holarrhena* spp., from *Wrightia tomentosa* and from *Funtumia elastica* (Apocynaceae). Narcotic to frogs but not to mammals. Antibacterial, antineoplastic agent. Has local anaesthetic props. but causes local necrosis on injection. Shows strong antifungal activity. Mp 123-124°. $[\alpha]_D^{20}$ -1.9 (CHCl₃). $[\alpha]_D^{20}$ +25.3 (EtOH). Log P 5.29 (uncertain value) (calc). Source of 18-substd. pregnanes.

▶ GK7620000

Hydrochloride (1:2):

Cryst. + 1H₂O. Mp 340°. $[\alpha]_D^{20}$ +9.3 (H₂O).

Hydrobromide (1:2): *Norine*

[5913-82-6] Formerly used to treat amoebic dysentery. Mp 340° dec. $[\alpha]_D^{20}$ +7 (c, 5 in H₂O).

▶ LD₅₀ (mus, orl) 390 mg/kg. GK7620000

Dipicrate: Mp 222-224°.

5 α ,6-Dihydro: **Dihydroconessine**

[33514-74-8]

$C_{24}H_{42}N_2$ 358.609

Isol. from the methylated alkaloid mixt. from *Holarrhena antidysenterica*; detected by ms and is prob. a minor alkaloid constit. (Apocynaceae).

7 α -Hydroxy: **7 α -Hydroxyconessine**

[7347-55-9]

$C_{24}H_{40}N_2O$ 372.593

Minor alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 176-178°. $[\alpha]_D^{20}$ -61 (c, 0.95 in CHCl₃). The presence of 7 β -Hydroxyconessine was also demonstrated chromatographically.

3-Epimer: **Conkuessine. Concuessine**

[14152-37-5]

$C_{24}H_{40}N_2$ 356.593

Alkaloid from the bark of *Holarrhena antidysenterica* (Apocynaceae). Mp 93-94° (86.5-87.5°). $[\alpha]_D$ +7 (CHCl₃). $[\alpha]_D$ +18 (EtOH).

3-Epimer, 5 α ,6-dihydro: **Dihydroconkuessine. Dihydroconcuessine**

[6869-90-5]

$C_{24}H_{42}N_2$ 358.609

Alkaloid from the bark of *Holarrhena antidysenterica*. Component of Zhi Xie Mu Pi. Needles (Me₂CO). Mp 93-94°. $[\alpha]_D^{21}$ +48 (c, 0.06 in CHCl₃).

Pyman, F.L. *et al.*, *J.C.S.*, 1919, **115**, 163-166 (*isol*)

Stephenson, R.P. *et al.*, *Br. J. Pharmacol.*, 1948, **3**, 237-245 (*pharmacol*)

Haworth, R.D. *et al.*, *J.C.S.*, 1953, 1102-1109 (*uv, ir, struct*)

Tschesche, R. *et al.*, *Chem. Ber.*, 1962, **95**, 1144-1154; 1964, **97**, 2316-2325 (*Conkuessine, 7 α -Hydroxyconessine*)

Marshall, J.A. *et al.*, *J.A.C.S.*, 1962, **84**, 1485-1486 (*synth*)

Stork, G. *et al.*, *J.A.C.S.*, 1962, **84**, 2018-2020 (*synth*)

Barton, D.H.R. *et al.*, *J.C.S.*, 1962, 622-631 (*synth*)

Labler, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 2345-2355 (*Dihydroconkuessine*)

Wolters, B. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1964, **297**, 748-759 (*activity*)

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 1566-1574 (*use*)

Johnson, W.S. *et al.*, *Tetrahedron, Suppl.*, No. 8, 1966, 541-601 (*synth*)

De Lajudie, P. *et al.*, *Eur. J. Med. Chem. (Chim. Ther.)*, 1970, **5**, 129-137 (*activity*)

Drake, A.F. *et al.*, *Tetrahedron*, 1977, **33**, 937-949 (*cd*)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 4772

Jiang, B. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 2543-2546 (*synth*)


Zirih, G.N. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2637-2640 (*isol, pmr, cmr*)

Lannang, A.M. *et al.*, *Acta Cryst. E*, 2008, **63**, o4398 (*cryst struct*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, DOX000

Conglobatin**C-602**

[72263-05-9]



Absolute Configuration

$C_{28}H_{38}N_2O_6$ 498.618

Dilactone antibiotic. Struct. revised in 1984. Prod. by *Streptomyces conglobatus* ATCC 31005. Shows some immunosuppressant activity but in general weak biol. activity. Cryst. (Et₂O/hexane). Sol. MeOH, EtOAc, Et₂O; poorly sol. hexane, H₂O. Mp 125°. $[\alpha]_D$ -44 (c, 1 in CHCl₃). λ_{max} 214 (ε 43800) (EtOH) (Derep).


▶ JG7521800

Westley, J.W. *et al.*, *J. Antibiot.*, 1979, **32**, 874-877 (*isol, cryst struct*)

Schrengelberger, C. *et al.*, *Tet. Lett.*, 1984, **25**, 5881-5884 (*synth*)

Conhydrine**C-603**

α -Ethyl-2-piperidinemethanol, 9*Cl*. 2-(1-Hydroxypropyl)piperidine. 1-(2-Piperidinyl)-1-propanol. α -Conhydrine [63401-12-7]



$C_8H_{17}NO$ 143.228

(+)-form [495-20-5]

Alkaloid from hemlock (*Conium maculatum*) (Apiaceae). Leaflets (Et₂O). Mp 121°. Bp₇₂₀ 224.5°. $[\alpha]_D$ +10 (c, 10 in EtOH).

▶ Extremely poisonous. GK8000000

N-Ac: Bp₃ 133-135°.

O,N-Dibenzoyl: Mp 133-134°. $[\alpha]_D$ -13 (c, 3 in CHCl₃).

N-Me: Bp₁₁ 94-95°.

(-)-form [18209-37-5]

Synthetic. $[\alpha]_D$ -9.3 (95% EtOH).

(±)-form [3238-62-8]

Synthetic. Mp 99-100°.

N-Me; methiodide: Mp 178-179°.

Wertheim, T. *et al.*, *Annalen*, 1856, **100**, 328 (*isol*)

Chemnitius, F. *et al.*, *J. Prakt. Chem.*, 1928, **118**, 25 (*isol*)

Galinovsky, F. *et al.*, *Monatsh. Chem.*, 1948, **79**, 426 (*synth*)

Marion, L. *et al.*, *Alkaloids (Academic Press)*, 1949, **1**, 218 (*rev*)

Hill, R.K. *et al.*, *J.A.C.S.*, 1958, **80**, 1609 (*config*)

Govindachari, T.R. *et al.*, *J.C.S.*, 1958, 1306 (*synth*)

Fodor, G. *et al.*, *Can. J. Chem.*, 1969, **47**, 4393 (*abs config, pmr*)

Craig, J.C. *et al.*, *Tetrahedron*, 1978, **34**, 501 (*cd*)

Stork, G. *et al.*, *Tet. Lett.*, 1979, 771 (*synth*)

Shono, T. *et al.*, *Tet. Lett.*, 1983, **24**, 4577 (*synth*)

Pilard, S. *et al.*, *Tet. Lett.*, 1984, **25**, 1555 (*synth*)

Masaki, Y. *et al.*, *Tet. Lett.*, 1989, **30**, 6395 (*synth*)

Agami, C. *et al.*, *Tet. Lett.*, 2000, **41**, 4113-4116 (*synth*)

Enders, D. *et al.*, *Tetrahedron: Asymmetry*, 2002, **13**, 285-291 (*synth*)

Kandula, S.V. *et al.*, *Tet. Lett.*, 2003, **44**, 1957-1958 (*synth*)

Nagata, K. *et al.*, *Heterocycles*, 2005, **66**, 107-109 (*synth*)

Pandey, S.K. *et al.*, *Tet. Lett.*, 2005, **46**, 4091-4093 (*synth*)

Chang, M.-Y. *et al.*, *Tetrahedron*, 2006, **62**, 10843-10848 (*synth*)


Voituriez, A. *et al.*, *J.O.C.*, 2007, **72**, 5358-5361 (*synth*)

Rodriguez, D. *et al.*, *Tet. Lett.*, 2008, **49**, 6866-6869 (*synth*)

Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 514

Conhydrinone C-604

1-(2-Piperidinyl)-1-propanone, 9CI. 2-Propanoylpiperidine. 1'-Oxoconiine [97073-23-9]



C₈H₁₅NO 141.213

(±)-form [38726-76-0]

Alkaloid from *Conium maculatum* (hemlock) and *Semnostachya menglaensis* (preferred genus name *Strobilanthes*).

Hydrochloride: [38726-77-1]

Mp 249-250°.

Hydrobromide: [38726-78-2]

Mp 228-229°.

Leete, E. et al., *J.A.C.S.*, 1972, **94**, 5472 (*isol, struct, synth*)


Tressl, R. et al., *J. Agric. Food Chem.*, 1985, **33**, 924-928 (*occur*)

Naef, R. et al., *J. Agric. Food Chem.*, 2005, **53**, 9161-9164 (*isol, synth, pmr, cmr, ms*)

Conicaquinone A

C-605

[519154-45-1]



R¹ = -CH₂CH₂CH=C(CH₃)₂, R² = H

C₁₈H₁₉NO₄S 345.418

Alkaloid from the ascidian *Aplidium conicum*. Cytotoxic. λ_{max} 261 (ε 16600); 298 (ε 7400) (MeOH).

Aiello, A. et al., *Eur. J. Org. Chem.*, 2003, 898-900 (*isol, pmr, cmr*)

Conicaquinone B

C-606

[519154-46-2]

As Conicaquinone A, C-605 with

R¹ = H, R² = -CH₂CH₂CH=C(CH₃)₂

C₁₈H₁₉NO₄S 345.418


Isol. from the ascidian *Aplidium conicum*. Cytotoxic. λ_{max} 263 (ε 14700); 300 (ε 8050) (MeOH).

Aiello, A. et al., *Eur. J. Org. Chem.*, 2003, 898-900 (*isol, pmr, cmr*)

Conioidine A

C-607

[154887-98-6]



C₂₂H₃₉NO₃ 365.555

Alkaloid from *Chamaesaracha conioides* (Solanaceae). Exhibits DNA-specific KB cell cytotoxicity. Oil. [α]_D²⁵ +48.2 (c, 0.2 in MeOH). λ_{max} 210 (ε 7400) (MeOH) (Berdy).

4'-Hydroxy: Conioidine B

[154887-99-7]


C₂₂H₃₉NO₄ 381.554

Alkaloid from *Chamaesaracha conioides* (Solanaceae). Exhibits DNA-specific KB cell cytotoxicity. Oil. [α]_D²⁵ +20.9 (c, 0.1 in MeOH). λ_{max} 211 (ε 7200) (MeOH) (Berdy).

Chan, G.W. et al., *J. Nat. Prod.*, 1993, **56**, 708 (*isol, uv, ir, pmr, cmr, ms, struct*)

Coniosetin

C-608



Absolute Configuration

C₂₅H₃₅NO₄ 413.556

Tetramic acid deriv. Related to Equisetin, E-142. Prod. by *Coniochaeta ellipsoidea* DSM 13856. Active against gram-positive bacteria and fungi. Powder. [α]_D²³ -319 (c, 0.1 in MeOH). λ_{max} 232 (ε 19000); 287 (ε 5500) (MeOH).

Segeth, M.P. et al., *J. Antibiot.*, 2003, **56**, 114-122 (*isol, cd, uv, pmr, cmr, ms*)

Coniothyriomycin

C-609

[141805-68-7]



C₁₃H₁₂ClNO₅ 297.694


Metab. of the fungus *Coniothyrium* sp. Exhibits antifungal and herbicidal activities. Cryst. (Me₂CO). Sol. MeOH, Me₂CO; fairly sol. Et₂O, CH₂Cl₂; poorly sol. H₂O, hexane. Mp 153°. λ_{max} 208 (ε 4406); 280 (ε 3417) (MeOH) (Berdy).

Krohn, K. et al., *Annalen*, 1992, 787 (*isol, pmr, cmr*)

Conkurchine

C-610

23-Norcona-5,18(22)-dienin-3-amine, 9CI. Irehline. Concurchine [3792-62-9]



Absolute Configuration

C₂₁H₃₂N₂ 312.497

Alkaloid from *Holarrhena antidysenterica*, *Holarrhena febrifuga*, *Holarrhena crassifolia*, *Funtumia elastica*, *Wrightia tomentosa* and *Malouetia arborea* (Apocynaceae). Shows hypotensive and vasodilatory activity. Mp 151°. [α]_D -51 (c, 1 in CHCl₃).

N-Formyl: N-Formylconkurchine

N-Formylirehline

[36132-43-1]

C₂₂H₃₂N₂O 340.508

Alkaloid from the leaves of *Holarrhena crassifolia* (Apocynaceae). Cryst. (Me₂CO). Mp 247°. [α]_D -50 (CHCl₃).

N³-Me: Conessidine

[6877-20-9]

C₂₂H₃₄N₂ 326.524

Alkaloid from *Holarrhena antidysenterica*, some other *Holarrhena* spp. and from *Wrightia tomentosa* (Apocynaceae). Mp 123°. [α]_D¹⁸ -61 (c, 0.85 in CHCl₃).

N³-Me, diperchlorate: Mp 243° dec.

5α,6-Dihydro: 23-Norcon-18(22)-enin-3-amine

Wrightiamine A

[640266-38-2]

C₂₁H₃₄N₂ 314.513

Alkaloid from the leaves of *Wrightia javanica*. Cytotoxic. Amorph. solid. [α]_D²⁵ -14 (c, 0.2 in MeOH).

Tschesche, R. et al., *Chem. Ber.*, 1954, **87**, 1719-1725; 1956, **89**, 1288-1295

(*Conessidine*)

Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1964, 1555-1563; 1564-1566 (*Conessidine, isol, ir, pmr, struct*)

Sóti, F. et al., *Tet. Lett.*, 1967, **8**, 1437-1441 (*Conkurchine, isol*)

Einhorn, J. et al., *Phytochemistry*, 1972, **11**, 769-777 (*N-Formylconkurchine*)

Dadoun, H. et al., *Ann. Pharm. Fr.*, 1973, **31**, 237-247 (*Conessidine, Conkurchine, isol*)


Einhorn, J. et al., *Bull. Soc. Chim. Fr.*, 1973, 301-303 (*Conessidine, synth*)

Kawamoto, S. et al., *Chem. Pharm. Bull.*, 2003, **51**, 737-739 (*Wrightiamine A*)

Conodiparine A

C-611

[213748-27-7]



C₄₄H₅₄N₄O₇ 750.933

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Amorph. pale yellow powder. [α]_D -34 (c, 0.71 in CHCl₃).

λ_{\max} 227 (log ϵ 4.92); 286 (log ϵ 4.32); 296 (log ϵ 4.35) (EtOH). λ_{\max} 227; 286; 296 (MeOH) (Berdy).

N-De-Me: Conodiparine E

[486444-25-1]

$C_{43}H_{52}N_4O_7$ 736.906

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Amorph. pale yellow powder. $[\alpha]_D$ -101 (c, 0.07 in $CHCl_3$). λ_{\max} 223 (log ϵ 4.23); 285 (log ϵ 3.64); 294 (log ϵ 3.66) (EtOH).

19'-Ketone: Conodiparine C

[213748-29-9]

$C_{44}H_{52}N_4O_7$ 748.917

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Amorph. pale yellow powder. $[\alpha]_D$ -27 (c, 0.5 in $CHCl_3$). λ_{\max} 227 (log ϵ 4.63); 286 (log ϵ 4.02); 296 (log ϵ 4.05) (EtOH).

Kam, T.-S. et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 1693-1696 (isol, uv, pmr, cmr)

Kam, T.-S. et al., *J. Nat. Prod.*, 2003, 66, 11-16 (isol, pmr, cmr, ms)

Conodiparine B

C-612

[213748-28-8]

As Conodiparine A, C-611 with

C-3 to C-12' linkage

$C_{44}H_{54}N_4O_7$ 750.933

Alkaloid from the leaves of *Tabernaemontana corymbosa*. MDR reversing agent. Amorph. pale yellow powder. $[\alpha]_D$ -64 (c, 0.93 in $CHCl_3$). λ_{\max} 224 (log ϵ 4.9); 286 (log ϵ 4.32); 293 (log ϵ 4.3) (EtOH) (Berdy).

N-De-Me: Conodiparine F

[486444-26-2]

$C_{43}H_{52}N_4O_7$ 736.906

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Amorph. pale yellow powder. $[\alpha]_D$ -73 (c, 0.32 in $CHCl_3$). λ_{\max} 224 (log ϵ 4.86); 286 (log ϵ 4.29); 294 (log ϵ 4.28) (EtOH).

11-O-De-Me: Cononitarine A

[486444-29-5]

$C_{43}H_{52}N_4O_7$ 736.906

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Pale yellow oil. $[\alpha]_D$ -52 (c, 0.11 in $CHCl_3$). λ_{\max} 223 (log ϵ 4.78); 285 (log ϵ 4.19); 294 (log ϵ 4.25) (EtOH).

19-Ketone: Conodiparine D

[213748-30-2]

$C_{44}H_{52}N_4O_7$ 748.917

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Amorph. pale yellow powder. $[\alpha]_D$ -42 (c, 0.9 in $CHCl_3$). λ_{\max} 224 (log ϵ 4.17); 286 (log ϵ 4.13); 296 (log ϵ 4.12) (EtOH).

19-Deoxy, 11-O-de-Me: Cononitarine B

[486444-30-8]

$C_{43}H_{52}N_4O_6$ 720.907

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Pale yellow oil. $[\alpha]_D$ -43 (c, 0.1 in $CHCl_3$). λ_{\max} 223 (log ϵ 4.52); 285 (log ϵ 3.92); 294 (log ϵ 3.89) (EtOH).


Kam, T.-S. et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 1693-1696 (isol, uv, pmr, cmr)

Kam, T.-S. et al., *J. Nat. Prod.*, 2003, 66, 11-16 (isol, pmr, cmr, ms)

Conodirinine A

[605674-08-6]

C-613



$C_{44}H_{52}N_4O_7$ 748.917

Related to Conodiparine A, C-611.

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Light yellowish oil. $[\alpha]_D$ -88 (c, 0.03 in $CHCl_3$). λ_{\max} 222 (log ϵ 4.56); 287 (log ϵ 4); 294 (log ϵ 4.01) (EtOH).

Kam, T.-S. et al., *Helv. Chim. Acta*, 2003, 86, 122-126 (isol, pmr, cmr)

Conodirinine B

[605674-09-7]

C-614

As Conodirinine A, C-613 with

C-3 to C-12' linkage

$C_{44}H_{52}N_4O_7$ 748.917


Related to Conodiparine B, C-612. Alkaloid from the leaves of *Tabernaemontana corymbosa*. Light yellowish oil. $[\alpha]_D$ -60 (c, 0.23 in $CHCl_3$). λ_{\max} 222 (log ϵ 4.62); 286 (log ϵ 4.05); 293 (log ϵ 4.03) (EtOH).

Kam, T.-S. et al., *Helv. Chim. Acta*, 2003, 86, 122-126 (isol, pmr, cmr)

Conoduramine

[2580-82-7]

C-615



$C_{43}H_{52}N_4O_5$ 704.908

Numbering systems vary, esp. which monomer is numbered with primes. Alkaloid from *Conopharyngia durissima*, *Tabernaemontana pachysiphon*, *Tabernaemontana chippii*, *Peschiera laeta*, *Gabunia odoratissima* (preferred genus name *Tabernaemontana*), *Tabernaemontana elegans*, *Tabernaemontana holstii* and *Tabernaemontana johnstonii*. Antineoplastic agent, shows cytotoxic activity vs. mouse lymphocyte P-388 and human nasopharynx KB cells. Also shows strong activity against gram-positive and weak activity against gram-negative bacteria.

Leishmanicide. Needles (MeOH/ Me_2CO). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 215-217° dec. $[\alpha]_D^{22}$ -77.5 (c, 1 in $CHCl_3$). Log P 7.36 (uncertain value) (calc). λ_{\max} 228; 285; 294 (MeOH) (Berdy).

N-De-Me: Gabunamine. N-Demethylconoduramine

[66086-99-5]

$C_{42}H_{50}N_4O_5$ 690.881

Alkaloid from the stem bark of *Tabernaemontana johnstonii* (Apocynaceae). Shows cytotoxic activity against P-388 lymphocytic leukaemia and human nasopharynx KB cells.

O-De-Me: Demethylconoduramine

[95066-49-2]

$C_{42}H_{50}N_4O_5$ 690.881

Minor alkaloid from *Tabernaemontana pachysiphon* stem bark and root bark (Apocynaceae). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. λ_{\max} 224; 286; 294 (MeOH) (Berdy). λ_{\max} 285; 294; 323 (MeOH/NaOH) (Berdy).

19,20ξ-Epoxyde: 19,20-Epoxyconoduramine

[66138-60-1]

$C_{43}H_{52}N_4O_6$ 720.907

Isol. from the stem bark of *Tabernaemontana johnstonii* (Apocynaceae).

19,20α-Dihydro: Tabernaeelegantine B

[58850-42-3]

$C_{43}H_{54}N_4O_5$ 706.923

Alkaloid from the root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (Me $_2$ CO). Mp 199°. $[\alpha]_D^{20}$ +14.4 ($CHCl_3$).

19,20β-Dihydro: Tabernaeelegantine D

[58865-84-2]

$C_{43}H_{54}N_4O_5$ 706.923

Alkaloid from root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (MeOH). Mp 206°. $[\alpha]_D^{20}$ +11.3 ($CHCl_3$).

3'-Hydroxy: 3'-Hydroxyconoduramine

[93627-69-1]

$C_{43}H_{52}N_4O_6$ 720.907

A major alkaloid from the root bark of *Tabernaemontana chippii*. Isol. as epimeric mixt., 3'R:3'Sca. 2:1.

3'-Hydroxy, 19,20α-dihydro: 3'-Hydroxy-tabernaeelegantine B

[103763-53-7]

$C_{43}H_{54}N_4O_6$ 722.923

Alkaloid from *Tabernaemontana elegans* (Apocynaceae). Mixt. of 3'-epimers.

19'R-Hydroxy: 19'R-Hydroxyconoduramine

[155637-19-7]

$C_{43}H_{52}N_4O_6$ 720.907

Alkaloid from leaves of *Tabernaemontana subglobosa* (Apocynaceae). Amorph. powder. $[\alpha]_D^{17}$ -78.5 (c, 1.0 in $CHCl_3$).

19'S-Hydroxy: 19'S-Hydroxyconoduramine

[602279-67-4]

$C_{43}H_{52}N_4O_6$ 720.907

Alkaloid from the leaves and stem bark of *Tabernaemontana corymbosa*. Light yellow oil. $[\alpha]_D$ -43 (c, 0.6 in $CHCl_3$).

λ_{\max} 227 (log ϵ 4.22); 288 (log ϵ 3.62); 295 (log ϵ 3.64) (EtOH).

Demethoxy: Ervahanine B

[80981-94-8]

C₄₂H₅₀N₄O₄ 674.881

Alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae). Amorph. $[\alpha]_{\text{D}}^{18}$ -123 (c, 0.95 in CHCl₃).

Demethoxy, 3'-hydroxy, N-de-Me: 3'-Hydroxy-N⁴-demethylervahanine B

[99257-57-5]

C₄₁H₄₈N₄O₅ 676.854

Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Active against gram-positive bacteria. Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. λ_{\max} 232 (ϵ 47800); 288 (ϵ 15200) (EtOH) (Berdy).

Demethoxy, 3'-oxo: Ervahimine B

[126398-89-8]

C₄₂H₄₈N₄O₅ 688.865

Minor alkaloid from roots of *Ervatamia hainanensis* (Apocynaceae).

3'-S-Cyano, 19,20 α -dihydro: Tabernaegantinine D

[72542-43-9]

C₄₄H₅₃N₅O₅ 731.933

Alkaloid from the root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (MeOH). Mp 259° dec.

Renner, U. et al., *Helv. Chim. Acta*, 1959, **42**, 1572-1581 (*isol, uv, ir*)

Renner, U. et al., *Tet. Lett.*, 1964, 283-287 (*struct*)

Bombardelli, E. et al., *J.C.S. Perkin 1*, 1976, 1432 (*Tabernaegantines B,D*)

Kingston, D.G.I. et al., *J. Pharm. Sci.*, 1978, **67**, 249-251 (*Gabunamine, 19',20'-Epoxyconoduramine*)

Danieli, B. et al., *J.C.S. Perkin 1*, 1980, 601 (*Tabernaegantinine D*)

Feng, X.-Z. et al., *J. Nat. Prod.*, 1981, **44**, 670-675; 1989, **52**, 928-933 (*Ervahanine B, Ervahimine B*)

Van Beek, T.A. et al., *Phytochemistry*, 1984, **23**, 1771-1778 (*Demethylconoduramine*)

Van Beek, T.A. et al., *J. Nat. Prod.*, 1985, **48**, 400-423 (*3'-Hydroxyconoduramine*)

Perera, P. et al., *Phytochemistry*, 1985, **24**, 2097-2104 (*3'-Hydroxy-N⁴-demethylervahanine B*)

Van der Heijden, R. et al., *Planta Med.*, 1986, 144 (*3'-Hydroxytabernaegantinine B*)


Takayama, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 280-284 (*19'R-Hydroxyconoduramine*)

Kam, T.-S. et al., *Phytochemistry*, 2003, **63**, 625-629 (*19'S-Hydroxyconoduramine*)

Conodurine

C-616

[2665-57-8]

C₄₃H₅₂N₄O₅ 704.908

Numbering systems vary, esp. which ring

system has primed locants. Alkaloid from *Conopharyngia durissima* (*Voacanga africana*), *Tabernaemontana pachysyphon*, *Tabernaemontana chippii*, *Peschiera laeta*, *Gabunia odoratissima* (preferred genus name *Tabernaemontana*) and *Tabernaemontana johnstonii*. Antineoplastic agent, shows cytotoxic activity vs. mouse lymphocyte P-388 and human nasopharynx KB cells. Also shows strong activity against gram-positive and weak activity against gram-negative bacteria. Antiparasitic agent. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 222-225°. $[\alpha]_{\text{D}}$ -101 (CHCl₃). Log P 7.36 (uncertain value) (calc). λ_{\max} 225 (log ϵ 4.78); 285 (log ϵ 4.19); 292 (log ϵ 4.16) (no solvent reported).

N-De-Me: Gabunine. N-Demethylconodurine

[1357-30-8]

C₄₂H₅₀N₄O₅ 690.881

Alkaloid from *Gabunia odoratissima*, *Tabernaemontana von heurkii* and *Tabernaemontana holstii* (Apocynaceae). Shows cytotoxic activity vs. mouse P-388 carcinoma cells. Antiparasitic agent. Needles (MeOH or EtOH). Mp 244-246° dec. $[\alpha]_{\text{D}}^{20}$ -105 (c, 0.62 in CHCl₃). λ_{\max} 226; 287; 295 (EtOH) (Berdy).

19,20 α -Dihydro: Tabernaegantinine A.

Alkaloid T

[58850-41-2]

C₄₃H₅₄N₄O₅ 706.923

Alkaloid from the root bark of *Tabernaemontana elegans*, *Hazunta costata* and from *Hazunta modesta* (Apocynaceae). Cryst. (MeOH). Mp 231°. $[\alpha]_{\text{D}}$ -31.8 (CHCl₃). λ_{\max} 224 (log ϵ 4.7); 285 (log ϵ 4.14); 293 (log ϵ 4.1) (EtOH).

19,20 α -Dihydro, 19'R-hydroxy: 19'R-Hydroxytabernaegantinine A. Alkaloid X

[79555-02-5]

C₄₃H₅₄N₄O₆ 722.923

Alkaloid from the stem bark of *Hazunta costata* (Apocynaceae). Cryst. (MeOH). Mp 268°. $[\alpha]_{\text{D}}$ -33 (c, 1 in CHCl₃). λ_{\max} 226 (ϵ 46650); 228 (ϵ 45725); 286 (ϵ 13000); 295 (ϵ 12500) (EtOH).

19,20 β -Dihydro: Tabernaegantinine C

[58865-83-1]

C₄₃H₅₄N₄O₅ 706.923

Alkaloid from root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (MeOH). Mp 171°. $[\alpha]_{\text{D}}^{20}$ -36.8 (CHCl₃). λ_{\max} 224 (log ϵ 4.76); 285 (log ϵ 4.18); 293 (log ϵ 4.16) (EtOH).

3'-Hydroxy: 3'-Hydroxyconodurine

[93627-70-4]

C₄₃H₅₂N₄O₆ 720.907

Major alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Approx. equal mixt. of 3'-epimers.

3'-Oxo: 3'-Oxoconodurine

[64192-90-1]

C₄₃H₅₀N₄O₆ 718.891

Alkaloid from the roots of *Tabernaemontana holstii* (Apocynaceae). Mp 217-218°.

19'R-Hydroxy: 19'R-Hydroxyconodurine

[155637-18-6]

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from leaves of *Tabernaemontana subglobosa* (Apocynaceae). Amorph. powder. $[\alpha]_{\text{D}}^{12}$ -96.5 (c, 1.0 in CHCl₃). λ_{\max} 224; 285; 293 (EtOH).

19'S-Hydroxy: 19'S-Hydroxyconodurine

[602279-66-3]

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from the leaves and stem bark of *Tabernaemontana corymbosa*. Light yellowish oil. $[\alpha]_{\text{D}}$ -69 (c, 0.12 in CHCl₃). λ_{\max} 223 (log ϵ 4.84); 286 (log ϵ 4.28); 293 (log ϵ 4.27) (EtOH).

19'S-Hydroxy, N-de-Me: Conodurinine

[603066-10-0]

C₄₂H₅₀N₄O₆ 706.88

Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Light yellow oil. $[\alpha]_{\text{D}}$ -55 (c, 0.47 in CHCl₃). λ_{\max} 224 (log ϵ 4.14); 286 (log ϵ 3.55); 293 (log ϵ 3.54) (EtOH).

Demethoxy: Ervahanine C

[80981-95-9]

C₄₂H₅₀N₄O₄ 674.881

Alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae). Amorph. $[\alpha]_{\text{D}}^{18}$ -62 (c, 0.39 in CHCl₃).

3'-Methoxy, 19,20 β -dihydro: 3'-Methoxytabernaegantinine C

[103763-54-8]

C₄₄H₅₆N₄O₆ 736.95

Alkaloid from *Tabernaemontana elegans* (Apocynaceae).

16'-Decarbomethoxy, 3'-hydroxy: 16'-Decarbomethoxy-3'-hydroxyconodurine. 3'-Hydroxy-16'-decarbomethoxyconodurine

[98798-63-1]

C₄₁H₅₀N₄O₄ 662.87

Trace alkaloid from *Tabernaemontana chippii* (Apocynaceae). Isol. as a mixt. of 3'R:3'S = 3:2.

Renner, U. et al., *Helv. Chim. Acta*, 1959, **42**, 1572-1581 (*isol, uv, ir*)

Renner, U. et al., *Tet. Lett.*, 1964, 283-287 (*pmr, struct*)

Cava, M.P. et al., *Tet. Lett.*, 1965, 931-935 (*Gabunine*)

Bombardelli, E. et al., *J.C.S. Perkin 1*, 1976, 1432-1438 (*Tabernaegantine A*)

Kingston, D.G.I. et al., *J. Pharm. Sci.*, 1977, **66**, 1135-1138; 1978, **67**, 249-251 (*Gabunine, 3-Oxoconodurine*)

Bui, A.-M. et al., *Phytochemistry*, 1977, **16**, 703-706 (*Alkaloid X*)

Urrea, M. et al., *Bull. Soc. Chim. Fr.*, 1981, **Part 2**, 147-149 (*19'R-Hydroxytabernaegantinine A*)

Feng, X.-Z. et al., *J. Nat. Prod.*, 1981, **44**, 670-675 (*Ervahanine C*)

Van Beek, T.A. et al., *J. Nat. Prod.*, 1985, **48**, 400-423 (*3-Hydroxyconodurine, 3-Hydroxy-16-decarbomethoxyconodurine*)

van der Heijden, R. et al., *Planta Med.*, 1986, 144-147 (*3'-Methoxytabernaegantinine C*)

Takayama, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 280-284 (*19'R-Hydroxyconodurine*)


Munoz, V. et al., *Planta Med.*, 1994, **60**, 455-459 (*isol, Gabunine, pharmacol*)

Medeiros, W.L.B. et al., *Magn. Reson. Chem.*, 1999, **37**, 676-681 (*pmr, cmr*)

Kam, T.-S. et al., *Phytochemistry*, 2003, **63**, 625-629 (*cmr, Conodurinine, 19S-Hydroxyconodurine*)

Conodutarine A

[486444-27-3]

C₄₂H₅₂N₄O₅ 692.897

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Pale yellow oil. $[\alpha]_D^{25}$ -51 (c, 0.18 in CHCl₃). λ_{\max} 227 (log ϵ 4.85); 285 (log ϵ 4.26); 293 (log ϵ 4.25) (EtOH).

19'-Ketone: Conodutarine B

[486444-28-4]


C₄₂H₅₀N₄O₅ 690.881

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Pale yellow oil. $[\alpha]_D^{25}$ -45 (c, 0.14 in CHCl₃). λ_{\max} 227 (log ϵ 4.87); 285 (log ϵ 4.31); 293 (log ϵ 4.28) (EtOH).

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2003, **66**, 11-16 (isol, pmr, cmr, ms)

Conofoline

Pedunculine†
[168922-21-2]

C₄₃H₅₂N₄O₇ 736.906

Pedunculine and Conofoline have not been compared but are probably identical. Mp uv and opt. rotn. refer to Conofoline. Alkaloid from leaves of *Ervatamia peduncularis* and *Tabernaemontana divaricata*. Purple amorph. solid. Mp 196-198°. $[\alpha]_D^{25}$ -97 (c, 0.725 in CHCl₃). V. difficult to crystallise. λ_{\max} 205 (log ϵ 3.16); 230 (log ϵ 3.68); 313 (log ϵ 3.76); 339 (log ϵ 3.84) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 1995, **40**, 313-316 (Conofoline)

Zèches-Hanrot, M. *et al.*, *Phytochemistry*, 1995, **40**, 587-591 (Pedunculine)

Chen, W. *et al.*, *Z. Kristallogr. - New Cryst. Struct.*, 1998, **213**, 183-184 (Conofoline, *cryst struct*)

ConollineC₁₃H₂₀N₂O 220.314

Prob. a lupine alkaloid. Struct. unknown. Alkaloid from *Ammodendron conollyi* (Fabaceae). Mp 192.5-193.5°. Cooccurs with Anagryne, A-970; and piperidine alkaloids. Opt. inactive, contains no OH or OMe groups.


C-617

Hydrochloride: Mp 180-182°.**Hydroiodide:** Mp 195-196°.**Perchlorate:** Mp 197-198°.

Proskurnina, N.F. *et al.*, *Zh. Obshch. Khim.*, 1949, **19**, 1396-1402; *J. Gen. Chem. USSR (Engl. Transl.)*, 1949, **19**, 1399

Conopharyngine†

[76-98-2]

C₂₃H₃₀N₂O₄ 398.501

Alkaloid from the stem and root bark of *Conopharyngia durissima* (Apocynaceae). Cryst. (Et₂O/petrol). Mp 141-143°. $[\alpha]_D^{25}$ -40.5 (c, 1 in CHCl₃). λ_{\max} 225 (log ϵ 4.47); 304 (log ϵ 4.05) (no solvent reported).

▶ GL1400000

3-Hydroxy: 3-Hydroxyconopharyngine

[95066-50-5, 93627-64-6 (R-form)]

C₂₃H₃₀N₂O₅ 414.5

Minor alkaloid from the root bark and stem bark of *Tabernaemontana pachysiphon* (Apocynaceae) and from *Tabernaemontana chippii* and *Tabernaemontana arborea*. Active against gram-positive bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. Isol. as a mixt. of epimers to which the spectroscopic data refers. However Cabezas *et al* assigned the 3*R*-config. to their material from *T. arborea*. λ_{\max} 223; 273 (sh); 302; 306; 311 (MeOH).

3-Oxo: 3-Oxoconopharyngine. 19-Oxocopharyngine

[20078-89-1]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Conopharyngia jollyana* (Apocynaceae). Cryst. (EtOAc). $[\alpha]_D^{20}$ -48 (c, 0.3 in CHCl₃). λ_{\max} 225 (log ϵ 4.4); 304 (log ϵ 4.02) (EtOH).

19*S*-Hydroxy: 19-Hydroxyconopharyngine

[16790-93-5]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from bark of *Conopharyngia jollyana* and *Conopharyngia durissima*. Also from *Tabernaemontana crassa* and *Tabernaemontana divaricata* (Apocynaceae). Mp 124-126° (as Ac). $[\alpha]_D^{26}$ -36.4 (c, 1.62 in CHCl₃). λ_{\max} 226 (log ϵ 4.42); 304 (log ϵ 3.93) (no solvent reported).

Renner, U. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 1572-1581 (isol, uv, ir, *struct*)

Hootele, C. *et al.*, *Chimia*, 1967, **21**, 133-134; 1968, **22**, 245-246 (19-Hydroxyconopharyngine, 3-Oxoconopharyngine)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 3350-3352 (19-Hydroxyconopharyngine)

Van Beek, T.A. *et al.*, *Phytochemistry*, 1984, **23**, 1771-1778 (3-Hydroxyconopharyngine)

Van Beek, T.A. *et al.*, *J. Ethnopharmacol.*, 1985, **14**, 315-318 (3-Hydroxyconopharyngine, activity)

Van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400-423 (3-Hydroxyconopharyngine)


Cabezas, J. *et al.*, *CA*, 1987, **107**, 4325 (3-Hydroxyconopharyngine)

Federici, E. *et al.*, *Planta Med.*, 2000, **66**, 93-95 (isol, cmr)

Conophyllidine

[152406-44-5]

C-621

C₄₄H₅₀N₄O₉ 778.9

Alkaloid from *Tabernaemontana divaricata* (single flower variety) (Apocynaceae). Farnesyl transferase inhibitor. Amorph.

▶ LD₅₀ (mus, ipr) 32.3 mg/kg.**14',15'-Epoxide: Conophylline. Polyervine**

[142741-24-0]

C₄₄H₅₀N₄O₁₀ 794.9

Alkaloid from the leaves of *Tabernaemontana divaricata* (single flower variety) and twigs and leaves of *Tabernaemontana glandulosa* (Apocynaceae). Also isol. as Polyervine from leaves of *Ervatamia polyneura*. Light yellow prisms (EtOAc). Mp 181-184° Mp ca.° 200 dec. $[\alpha]_D^{25}$ -133 (c, 0.7 in CHCl₃). Conophylline and Polyervine have not been compared but are prob. identical.

14',15'-Dihydro, 14',15'-β-dihydroxy:**Conophyllinine**C₄₄H₅₂N₄O₁₁ 812.915

Alkaloid from the leaves of *Tabernaemontana divaricata* (double flower variety). Light yellowish oil. $[\alpha]_D^{25}$ -109 (c, 0.09 in CHCl₃). λ_{\max} 204 (log ϵ 3.43); 239 (log ϵ 3.23); 311 (log ϵ 3.25); 334 (log ϵ 3.31) (EtOH).

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1865 (Conophyllidine, Conophylline, isol, uv, pmr, cmr, *cryst struct*)

Achenbach, H. *et al.*, *Phytochemistry*, 1994, **37**, 1737 (Conophyllidine, Conophylline, isol, ir, cd)


Clivio, P. *et al.*, *Phytochemistry*, 1995, **40**, 953 (Polyervine)

Kam, T.S. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 1292-1297 (Conophyllinine)

Conisculine

[219829-73-9]

C-622

C₂₀H₂₉NO₅ 363.453

Alkaloid from the roots of *Convolvulus siculus*. λ_{\max} 250 (MeOH aq.).

2'-Deoxy: Consabatine

[219829-74-0]

$C_{20}H_{29}NO_4$ 347.453

Alkaloid from *Convolvulus sabatius* ssp. *mauritanicus* and *Merremia quinata*. λ_{\max} 255 (MeOH aq.).

2'-Deoxy, 4'- ξ -alcohol: 4'-Dihydroconsabatine

$C_{20}H_{31}NO_4$ 349.469

Alkaloid from *Merremia quinata*. Oil.

Jenett-Siems, K. *et al.*, *Phytochemistry*, 1998, **49**, 1449-1451 (*Consiculine, Consabatine*)

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (*4'-Dihydroconsabatine*)

Consolidine† C-623

[1391-11-3]

$C_{33}H_{49}NO_9$ 603.751

Struct. unknown. Alkaloid from *Delphinium consolida* (Ranunculaceae). Prismatic needles (MeOH). Mp 153-157° (rapid htg.). $[\alpha]_D^{25} +64$ (c, 0.36 in MeOH).


Hydrochloride:

Wedge-shaped prisms (MeOH). Mp 240° (rapid htg.). $[\alpha]_D^{25} +33$ (c, 0.589 in 90% MeOH aq.).

Marion, L. *et al.*, *J.A.C.S.*, 1947, **69**, 2010-2014

Constrictosine C-624

3,10-Dihydroxydibenz[c,g]azecine-13(14H)-one, 9CI
[196086-14-3]



$C_{17}H_{13}NO_3$ 279.295

Alkaloid from *Aristolochia constricta* (Aristolochiaceae). Amorph. yellow solid. Mp 200-243°. λ_{\max} 224 (log ϵ 3.1); 256 (log ϵ 3.24); 282 (log ϵ 4.06); 302 (log ϵ 3.86) (MeOH).

3-Me ether: 3-O-Methylconstrictosine

[195971-30-3]

$C_{18}H_{15}NO_3$ 293.321

Alkaloid from *Aristolochia constricta*. Amorph. yellow solid. Mp 215-239°. λ_{\max} 224 (log ϵ 3.32); 256 (log ϵ 3.38); 282 (log ϵ 4.16); 302 (log ϵ 3.92) (MeOH).

Di-Me ether: 3,10-Di-O-methylconstrictosine

[195971-24-5]

$C_{19}H_{17}NO_3$ 307.348

Alkaloid from *Aristolochia constricta*. Amorph. yellow solid. Mp 189-195°. λ_{\max} 224 (log ϵ 4.12); 256 (log ϵ 4.36); 282 (log ϵ 4.96); 302 (log ϵ 4.72) (MeOH).

5,6-Dihydro: 5,6-Dihydroconstrictosine

[195971-28-9]

$C_{17}H_{15}NO_3$ 281.31

Alkaloid from *Aristolochia constricta*.

Amorph. yellow solid. Mp 187-192°. λ_{\max} 256 (log ϵ 2.36); 282 (log ϵ 3.28) (MeOH).

5,6-Dihydro, di-Me ether: 5,6-Dihydro-3,10-di-O-methylconstrictosine

[195971-26-7]


$C_{19}H_{19}NO_3$ 309.364

Alkaloid from *Aristolochia constricta*. Amorph. yellow solid. Mp 186-192°. λ_{\max} 256 (log ϵ 4.4); 282 (log ϵ 4.82) (MeOH).

Rastrelli, L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1065-1069 (*isol, uv, pmr, cmr*)

Convolutamine A C-625

[161099-50-9]



$C_{13}H_{18}Br_3NO_2$ 460.003

Alkaloid from the marine bryozoan *Amathia convoluta*. Exhibits cytotoxicity against murine P388 lymphocytic leukaemia. Oil. Racemic. λ_{\max} 212 (ϵ 27900) (MeOH) (Berdy).

N-De-Me: Convolutamine C

[161099-52-1]

$C_{12}H_{16}Br_3NO_2$ 445.976

From *Amathia convoluta*. Oil. Stereochem. unknown.

N-De(2-hydroxypropyl): Convolutamine F

[243858-10-8]

$C_{10}H_{12}Br_3NO$ 401.923

Alkaloid from *Amathia convoluta*. Oil. $[\alpha]_D^{20} +24.3$ (c, 0.4 in $CHCl_3$). λ_{\max} 213 (ϵ 21800) (MeOH).

4-Debromo: Convolutamine B

[161099-51-0]

$C_{13}H_{15}Br_2NO_2$ 381.107

From *Amathia convoluta*. Exhibits cytotoxicity against murine P388 lymphocytic leukaemia. Oil. Racemic. λ_{\max} 211 (ϵ 20100) (MeOH) (Berdy).

4-Debromo, N-de(2-hydroxypropyl): Convolutamine G

[243858-11-9]

$C_{10}H_{13}Br_2NO$ 323.027

Alkaloid from *Amathia convoluta*. Oil. λ_{\max} 206 (ϵ 8910); 229 (ϵ 1930) (MeCN).


Zhang, H. *et al.*, *Chem. Lett.*, 1994, 2271-2274 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kamano, Y. *et al.*, *Coll. Czech. Chem. Comm.*, 1999, **64**, 1147-1153 (*Convolutamines F,G*)

Hashima, H. *et al.*, *Bioorg. Med. Chem.*, 2000, **8**, 1757-1766 (*synth*)

Convolutamine D C-626

[161099-53-2]




$C_{13}H_{16}Br_3NO_2$ 457.987

Alkaloid from the marine bryozoan *Amathia convoluta*. Exhibits cytotoxicity against murine P388 lymphocytic leukaemia. Oil.

Zhang, H. *et al.*, *Chem. Lett.*, 1994, 2271-2274 (*isol, pmr, cmr, struct*)

Convolutamine E C-627

[161099-54-3]




$C_{15}H_{20}Br_3NO_3$ 502.04

Alkaloid from the marine bryozoan *Amathia convoluta*. Oil.

Zhang, H. *et al.*, *Chem. Lett.*, 1994, 2271 (*isol, pmr, cmr, struct*)

Convolutamine H C-628

N-Methyl-2-(2,3,4-tribromo-5,6-dimethoxyphenyl)ethylamine, 2,3,4-Tribromo-5,6-dimethoxy-N-methylbenzeneethanamine, 9CI
[443355-92-8]




$C_{11}H_{14}Br_3NO_2$ 431.949

Alkaloid from the marine bryozoan *Amathia convoluta*. Pale yellow oil. λ_{\max} 217 (log ϵ 4.4) (MeOH).

Narkowicz, C.K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 938-941 (*isol, pmr, cmr, ms*)

Convolutamydines C-629



Absolute Configuration

Convolutamidine A; R = CH_2COCH_3
B; R = CH_2CH_2Cl
C; R = CH_3
D; R = $CH=CH_2$
E; R = CH_2CH_2OH

Convolutamidine A

4,6-Dibromo-1,3-dihydro-3-hydroxy-3-(2-oxopropyl)-2H-indol-2-one, 9CI
[163564-85-0]
[184587-77-7]

$C_{11}H_9Br_2NO_3$ 363.005

Alkaloid from the marine bryozoan

Amathia convoluta. Cell differentiation and growth inhibitor. Amorph. solid (MeOH). Mp 190-195°. $[\alpha]_D^{26} +27.4$ (c, 0.06 in MeOH) (natural). $[\alpha]_D^{20} +48.2$ (c, 0.2 in MeOH) (synthetic). λ_{\max} 224 (ε 21600); 262 (ε 5200); 291 (ε 3100) (MeOH) (Berdy).

Convolutamydine B

4,6-Dibromo-3-(2-chloroethyl)-1,3-dihydro-3-hydroxy-2H-indol-2-one, 9CI [163564-86-1]

C₁₀H₈Br₂ClNO₂ 369.439

Alkaloid from the marine bryozoan *Amathia convoluta*. Amorph. solid (Me₂CO). Mp 225-227°. $[\alpha]_D^{25} +18.1$ (c, 0.42 in MeOH).

Convolutamydine C

4,6-Dibromo-1,3-dihydro-3-hydroxy-3-methyl-2H-indol-2-one, 9CI [163564-87-2]

C₉H₇Br₂NO₂ 320.968

Alkaloid from the marine bryozoan *Amathia convoluta*. Amorph. solid (Me₂CO). Mp 175-180°. $[\alpha]_D^{25} +32.4$ (c, 0.03 in MeOH).

Convolutamydine D

4,6-Dibromo-3-ethenyl-1,3-dihydro-3-hydroxy-2H-indol-2-one, 9CI [163564-94-1]

C₁₀H₇Br₂NO₂ 332.979

Alkaloid from the marine bryozoan *Amathia convoluta*. Cell differentiation and growth inhibitor. Amorph. solid. $[\alpha]_D^{26} +14$ (c, 0.04 in MeOH). λ_{\max} 220 (ε 7488); 290 (ε 1475) (MeOH) (Berdy).

Convolutamydine E

4,6-Dibromo-1,3-dihydro-3-hydroxy-3-(2-hydroxyethyl)-2H-indol-2-one, 9CI [243858-13-1]

C₁₀H₉Br₂NO₃ 350.994

Alkaloid from the marine bryozoan *Amathia convoluta*. Oil. λ_{\max} 223 (ε 16870); 298 (ε 850) (MeCN).

Kamano, Y. *et al.*, *Tet. Lett.*, 1995, **36**, 2783-2784 (*Convolutamydine A*, *isol*)

Zhang, H. *et al.*, *Tetrahedron*, 1995, **51**, 5523-5528 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cd*, *struct*)

Miah, S. *et al.*, *J.C.S. Perkin I*, 1997, 2405-2412 (*synth*, *Convolutamydine C*)

Kamano, Y. *et al.*, *Coll. Czech. Chem. Comm.*, 1999, **64**, 1147-1163 (*Convolutamydine E*)

Jnaneshwara, G.K. *et al.*, *Synth. Commun.*, 1999, **29**, 3627-3633 (*Convolutamydine A*)

Takayama, H. *et al.*, *Tetrahedron*, 1999, **55**, 6841-6846 (*abs config*)

Kawasaki, T. *et al.*, *Tetrahedron*, 2004, **60**, 3493-3503 (*synth*)

Nakamura, T. *et al.*, *Org. Lett.*, 2006, **8**, 677-679 (*synth*)

Luppi, G. *et al.*, *Tetrahedron*, 2006, **62**, 12017-12024 (*synth*)

Cravotto, G. *et al.*, *Tetrahedron: Asymmetry*, 2006, **17**, 3070-3074; 2007, **18**, 298


(*Convolutamydine A*, *synth*, *abs config*)

Malkov, A.V. *et al.*, *Org. Lett.*, 2007, **9**, 5473-5476 (*synth*)

Convolutindole A

C-630

2,4,6-Tribromo-1,7-dimethoxy-N,N-dimethyl-1H-indole-3-ethanamine. 2,4,6-Tribromo-3-(2-dimethylaminoethyl)-1,7-dimethoxyindole [443356-86-3]



C₁₄H₁₇Br₃N₂O₂ 485.013

Alkaloid from the marine bryozoan *Amathia convoluta*. Amorph. solid. Mp 61.5-62.5°. λ_{\max} 235 (log ε 4.7); 289 (log ε 4.1); 305 (log ε 4) (MeOH).

Narkowicz, C.K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 938-941 (*isol*, *pmr*, *cmr*, *ms*)

Convolvine

C-631

C₁₀H₁₆N₂ 164.25


Struct. unknown. Alkaloid from *Convolvulus pseudocantabricus*. Mp 200-202° (as dipicrate). Bp 250-260°.

Orekhov, A.P. *et al.*, *Ber., B*, 1935, **68**, 814-819 (*isol*)

Convolidine

C-632

3,4-Dimethoxybenzoic acid 1,2-ethanediylybis(8-azabicyclo[3.2.1]octane-8,3-diy) ester, 9CI [50656-81-0]



C₃₄H₄₄N₂O₈ 608.73

Alkaloid from *Convolvulus subhirsutus* (*Convolvulus pseudocantabricus*) and *Evolvulus sericeus* var. *holosericeus* (Convolvulaceae). Mp 192-193°. Artifact resulting from condensation of Convolvine, C-633 with C₂H₄Cl₂. Stated by Orechhoff to be opt. active, but this must have been erroneous.

Orechhoff, A. *et al.*, *Ber.*, 1935, **68**, 814 (*isol*)


Yusunov, S.Yu. *et al.*, *Dokl. Akad. Nauk SSSR*, 1958, **10**, 17; *CA*, **53**, 18391 (*struct*)

Consuelo Fonseca, L. *et al.*, *CA*, 1973, **79**, 15798w (*isol*)

Convolvine

C-633

8-Azabicyclo[3.2.1]oct-3-yl 3,4-dimethoxybenzoate, 9CI. Veratroylnortropine. 3α-Veratroyloxynortropene. 3α-(3,4-Dimethoxybenzoyloxy)nortropene [537-30-4]



C₁₆H₂₁NO₄ 291.346

Alkaloid from *Convolvulus pseudocantabricus*, *Convolvulus erinaceus*, *Convolvulus lineatus*, *Convolvulus subhirsutus* and *Evolvulus sericeus* var. *holosericeus* (Convolvulaceae). Mp 114-115°. Opt. inactive (*meso*-).

Hydrochloride: Mp 260-261° dec.

N-Formyl: **Confoline**. N-Formylconvolvine [76971-33-0]

C₁₇H₂₁NO₅ 319.357

Alkaloid from roots of *Convolvulus subhirsutus* (Convolvulaceae). Cryst. (Me₂CO). Mp 140-141°.

N-Ac: **Convolicine**. 3α-Veratroyloxy-N-acetylnortropene

[72994-86-6]

C₁₈H₂₃NO₅ 333.383

Alkaloid from the epigeal parts of *Convolvulus krauseanus* (Convolvulaceae). Cryst. (Et₂O). Mp 144-145°.

N-Me: **Convolumine**. Veratroyltropine [500-56-1]

C₁₇H₂₃NO₄ 305.373

Alkaloid from *Convolvulus pseudocantabricus*, *Convolvulus lineatus* and *Convolvulus subhirsutus* (Convolvulaceae). Mp 114-115°.

N-Me, N-oxide: **Convolumine N-oxide** [97534-16-2]

C₁₇H₂₃NO₅ 321.372

Alkaloid from *Convolvulus krauseanus*. Cryst. Mp 118-119°.

N-Isopropyl: **Convosine**

[107373-72-8]

C₁₉H₂₇NO₄ 333.427

Alkaloid from the roots of *Convolvulus subhirsutus* (Convolvulaceae). Cryst. (Me₂CO). Mp 103-104° (synthetic).

N-Hydroxy: **Convolvine**. 3α-Veratroyloxy-N-hydroxynortropene

[89783-61-9]

C₁₆H₂₁NO₅ 307.346

Alkaloid from *Convolvulus krauseanus* (Convolvulaceae). Mp 184-185°.

O⁴-De-Me: **Convolidine**. 3α-Vanilloxyloxynortropene

[63911-32-0]

C₁₅H₁₉NO₄ 277.319

Alkaloid from *Convolvulus krauseanus* (roots and epigeal parts) and from *Convolvulus subhirsutus* (Convolvulaceae). Cryst. (MeOH). Mp 214-215°.

O⁴-De-Me, N-formyl: **Confolidine**

C₁₆H₁₉NO₅ 305.33

Alkaloid from *Convolvulus subhirsutus*. Mp 178-179°.

O⁴-De-Me, N-Me: **Phyllalbine**. Vanillyltropate

[4540-25-4]


C₁₆H₂₁NO₄ 291.346

Alkaloid from the roots of *Phyllanthus discoides*, above-ground parts of *Convolvulus krauseanus* and roots of *Convolvulus subhirsutus* (Euphorbiaceae, Convolvulaceae). Cryst. (MeOH). Mp 209-210°.

O⁴-De-Me, N-Me, hydrochloride: Cryst. (MeOH). Mp 254°.

O⁴-De-Me, N-Me, N-oxide: **Phyllalbine N-oxide**

[954099-40-2]

C₁₆H₂₁NO₅ 307.346Alkaloid from *Convolvulus subhirsutus*.
Amorph. solid.**5'-Methoxy-3 α -(3,4,5-Trimethoxybenzoyloxy)nortropane**C₁₇H₂₃NO₅ 321.372Alkaloid from the root bark of *Erythroxyllum zambesiaceae* (Erythroxyllaceae). Plates (EtOH aq.) (as picrate). Mp 235° dec. (softens at 225°) (picrate).**5'-Methoxy, N-Me: Tropan-3 α -yl 3,4,5-trimethoxybenzoate. O-(3,4,5-Trimethoxybenzoyl) tropine**C₁₈H₂₅NO₅ 335.399Alkaloid from root bark of *Erythroxyllum monogynum* (Erythroxyllaceae). Fine needles (EtOH) (as hydrobromide). Mp 212° dec. (hydrobromide).Orchoff, A. et al., *Ber.*, 1935, **68**, 814 (*struct. Convolvine, Convolamine*)Yunusov, S.Yu. et al., *Dokl. Akad. Nauk SSSR*, 1958, **10**, 17; *CA*, **53**, 18391Parello, J. et al., *Bull. Soc. Chim. Fr.*, 1963, 2787 (*Phyllalbine*)Israilov, I. et al., *CA*, 1965, **63**, 7346f (*isol. Convolvine, Convolamine*)Consuela Fonseca, L. et al., *CA*, 1973, **79**, 15798w (*isol*)Agar, J.T.H. et al., *J.C.S. Perkin I*, 1976, 1550 (*Tropanyl trimethoxybenzoate*)Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1977, **13**, 290; 1979, **15**, 527; 1983, **19**, 749; 1985, **21**, 275; 1986, **22**, 618; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 255; 1979, **15**, 457; 1983, **19**, 712; 1985, **21**, 261; 1986, **22**, 581 (*Convolidine, Convolicine, Convoline, Convolamine N-oxide, Convosine*)Sharova, S.F. et al., *Khim. Prir. Soedin.*, 1980, **16**, 672; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 487 (*Confoline, Phyllalbine*)El-Iman, Y.M.A. et al., *Phytochemistry*, 1987, **26**, 2385 (*Trimethoxybenzoyloxynortropane*)Razzakov, N.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 54-55 (*Confolidine*)Gapparov, A.M. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 291-292 (*Phyllalbine N-oxide*)**Copiamycin****C-634***Niphithricin A. Niphithricin I. NSC 110326*
[11078-23-2]C₅₄H₉₅N₃O₁₇ 1058.354Macrolide antibiotic. Isol. from *Streptomyces hygroscopicus* var. *crystallogenes* and *Streptomyces violaceoniger*. Active against gram-positive bacteria and fungi. Shows synergistic activity with imidazole antifungals. Cryst. (MeOH). Mp 144° dec. [α]_D^{25.5} +14.4 (MeOH). Interconverts with Niphithricin B on standing in soln.Related to Niphimycin. λ_{\max} 205 (ϵ 15850); 220 (ϵ 20600) (MeOH) (Derep).

▶GL5300000

N-De-Me: Neocopiamycin A

[89989-28-6]

C₅₃H₉₃N₃O₁₇ 1044.328From *Streptomyces hygroscopicus* var. *crystallogenes*. Antifungal, less toxic than Copiamycin. Hygroscopic powder. Sol. MeOH, butanol, Py, DMSO; poorly sol. H₂O, hexane, Me₂CO. Mp 134-137° dec. [α]_D²⁵ +23 (c, 1 in MeOH). λ_{\max} 205 (ϵ 15850); 220 (ϵ 20600) (MeOH) (Derep).▶LD₅₀ (mus, ivn) 30-70 mg/kg; LD₅₀ (mus, ipr) 1450-1550 mg/kg. GL5310000**21-O-Demalonyl: Demalonylcopiamycin**

[84959-52-4]

C₅₁H₉₃N₃O₁₄ 972.307From *Streptomyces hygroscopicus* var. *crystallogenes*. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 222 (E1%_{1cm} 1737) (EtOH) (Berdy).**15-Me ether, 21-O-demalonyl: Demalonylmethylcopiamycin**

[101923-21-1]

C₅₂H₉₅N₃O₁₄ 986.334From *Streptomyces hygroscopicus* var. *crystallogenes*. Prisms (MeOH aq.). Sol. MeOH, DMSO; poorly sol. CHCl₃, H₂O, hexane. Mp 140-143°. [α]_D +11 (MeOH). λ_{\max} 215 (MeOH) (Berdy).**34,35E-Didehydro: Antibiotic TMC 34***TMC 34*

[170244-28-7]

C₅₄H₉₃N₃O₁₇ 1056.339Prod. by a *Streptomyces* sp. Antifungal agent. Sol. DMSO, Py, MeOH, 1-propanol, EtOH; fairly sol. EtOAc, Me₂CO, CHCl₃; poorly sol. H₂O. Mp 146-148°. [α]_D²⁰ +18 (c, 0.3 in MeOH). λ_{\max} 205 (ϵ 17600) (MeOH) (Berdy).**13-Deoxy: Neocopiamycin B. 13-Deoxycopiamycin**


[113189-06-3]

C₅₄H₉₅N₃O₁₆ 1042.355Prod. by *Streptomyces hygroscopicus* var. *crystallogenes*. Powder. Dec. at 124-126°.

[80450-11-9]

Arai, T. et al., *J. Antibiot., Ser. A*, 1965, **18**, 63 (*isol*)Seigi, K. et al., *Appl. Microbiol.*, 1971, **21**, 986 (*props*)Fiedler, H.P. et al., *J. Antibiot.*, 1981, **34**, 1107 (*isol*)Uno, J. et al., *Antimicrob. Agents Chemother.*, 1983, **24**, 552-559 (*activity*)Gassmann, P. et al., *Helv. Chim. Acta*, 1984, **67**, 696 (*struct*)Arai, T. et al., *J. Antibiot.*, 1984, **37**, 103 (*isol, uv, ir, cmr*)Takesako, K. et al., *J. Antibiot.*, 1985, **38**, 1363 (*derivis*)Fukai, T. et al., *Heterocycles*, 1986, **24**, 3351; 1988, **27**, 2333 (*deriv, isol*)Kohn, J. et al., *J. Antibiot.*, 1995, **48**, 1173 (*TMC 34*)Fukai, T. et al., *J. Antibiot.*, 1999, **52**, 340-344 (*Neocopiamycin B*)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CNH800**Coprogen****C-635**


[31418-71-0]

C₃₅H₅₃FeN₆O₁₃ 821.683Isol. from bacteria and fungi incl. *Neurospora crassa* and *Penicillium* spp. Siderophore. Microbiol. iron transport compd. Growth factor for coprophilic fungi. Phytotoxin. Red cryst. (EtOH). Sol. H₂O, MeOH. Mp 205° dec. λ_{\max} 435 (ϵ 2809) (MeOH) (Berdy). λ_{\max} 217 (ϵ 28840); 250 (ϵ 16600); 434 (ϵ 2820); 440 (ϵ 2951) (H₂O) (Berdy). λ_{\max} 213; 254; 450 (EtOH aq.) (Berdy).**N-De-Ac: Coprogen B**

[12705-44-1]

C₃₃H₅₁FeN₆O₁₂ 779.646From *Fusarium* spp., *Neurospora crassa* and other fungi. Siderophore. Sol. H₂O, MeOH. λ_{\max} 220; 250; 440 (ϵ 2754) (H₂O) (Berdy).

[30315-65-2, 81559-20-8, 1391-13-5]

Hesseltine, C.W. et al., *J.A.C.S.*, 1952, **74**, 1362 (*isol*)Pidacks, C. et al., *J.A.C.S.*, 1953, **75**, 6064 (*spectra*)*Fr. Pat.*, 1964, 1 355 923; *CA*, **62**, 5856 (*synth*)Diekmann, H. et al., *Arch. Microbiol.*, 1970, **73**, 65 (*Coprogen B*)Keller-Schierlein, W. et al., *Helv. Chim. Acta*, 1970, **53**, 2035 (*isol, struct*)Dell, A. et al., *Biomed. Mass Spectrom.*, 1982, **9**, 158 (*ms*)Wong, G.B. et al., *J.A.C.S.*, 1983, **105**, 810 (*isol, cd, config*)Matzanke, B.F. et al., *Eur. J. Biochem.*, 1987, **162**, 643Ohra, J. et al., *Biosci., Biotechnol., Biochem.*, 1995, **59**, 1782 (*isol*)**Coproverdin†****C-636***Methyl 1-formyl-4,9-dihydro-1,9-dihydroxy-4-oxo-1H-carbazole-8-carboxylate, 9CI. Methyl 8-formyl-8,9-dihydro-8,9-dihydroxy-5-oxo-5H-carbazole-1-carboxylate*
[437702-23-3]C₁₅H₁₁NO₆ 301.255Alkaloid from a New Zealand tunicate. Antitumour agent. Yellow oil. [α]_D²⁰ -8 (c,


0.36 in EtOH). λ_{\max} 208 (ϵ 20000); 270 (ϵ 6700); 302 (ϵ 4800); 382 (ϵ 16000) (EtOH).

Urban, S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1371-1373 (*isol*, *pmr*, *cmr*, *ms*)

Copsomycin

C-637

[129898-72-2]




Thiazole peptide antibiotic. Related to Nosiheptide, N-319. Struct. of chromophore shown. Prod. by a *Streptomyces* sp. Active against gram-positive bacteria.

Alekseeva, L.E. *et al.*, *Antibiot. Khimioter.*, 1990, **35**, 12-13 (*pmr*, *cmr*)

Coptisine

C-638

6,7-Dihydrobis[1,3]benzodioxolo[5,6-a:4',5'-g]quinolinizinium, 9Cl. 2,3:9,10-Bismethylenedioxyprotoberberine [3486-66-6]



$C_{19}H_{14}NO_4^{\oplus}$ 320.324

Quaternary alkaloid from *Coptis japonica* (Ranunculaceae) and many other spp., mostly in the Papaveraceae. Antineoplastic agent. Antiinflammatory agent. Uterine contractant. λ_{\max} 226 (ϵ 24550); 239 (ϵ 22400); 264 (ϵ 19500); 356 (ϵ 19950) (MeOH) (Berdy).

Hydroxide: [521-25-5]

Mp 216-218°.

Chloride: [6020-18-4]

Orange needles. Mp 300° dec.

Iodide: [6020-84-4]

Mp 280°.

Späth, E. *et al.*, *Ber.*, 1929, **62**, 1029 (*synth*)

Huang-Minlon, *et al.*, *Ber.*, 1936, **69**, 1737 (*isol*)

Jewers, K. *et al.*, *J.C.S. Perkin 2*, 1972, 1393 (*cmr*)

Pavelka, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3157 (*uv*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1991, **56**, 1116 (*isol*)


Ghee, T.T. *et al.*, *J. Nat. Prod.*, 1991, **54**, 143 (*isol*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (N-15 *nmr*)

Corallistine†

C-639

[121098-71-3]



$C_{10}H_{13}N_5OS$ 251.312

Isol. from the New Caledonian sponge *Corallistes fulvodesmus*. Light yellow cryst. (MeOH). Mp 192° dec.

Debitus, C. *et al.*, *Tet. Lett.*, 1989, **30**, 1535-1538 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cryst struct*)

Coranicine

C-640

[1355-99-3]

$C_{16}H_{17}NO_4$ 287.315

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Ammodarichis coranica*, *Crinum erubescens* and *Hippeastrum bifida* (Amaryllidaceae). Amorph. $[\alpha]_D^{21} +115$ (c, 0.195 in EtOH). $[\alpha]_D^{21} +156$ (c, 0.198 in $CHCl_3$).

Perchlorate:

Cryst. (Me₂CO). Mp 192-194° dec.

Methiodide:

Cryst. (MeOH aq.). Mp 245-247° dec.

Di-Ac:

Needles (Et₂O/petrol). Mp 118-119°.

$[\alpha]_D^{21} +90$ (c, 0.212 in EtOH). $[\alpha]_D^{21} +80$ (c, 0.215 in $CHCl_3$).

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 1307-1317 (*isol*, *uv*, *ir*)

Wildman, W.C. *et al.*, *Pharmazie*, 1967, **22**,

725; *CA*, **69**, 19337m (*isol*)


Wildman, W.C. *et al.*, *J.A.C.S.*, 1968, **90**, 6439-6446 (*ms*)

Wildman, W.C. *et al.*, *J.O.C.*, 1968, **33**, 3749-3753 (*isol*)

Cordifoline

C-641

[25455-00-9]



Absolute configuration

$C_{28}H_{30}N_2O_{12}$ 586.551

Alkaloid from the heartwood of *Adina cordifolia* (Rubiaceae). Noncryst. Mp 91-93° (as tetra-*O*-Ac, Me ester). $[\alpha]_D^{20} -144$ (c, 0.11 in MeOH) (tetra-Ac, Me ester).

Deoxy: **Deoxycordifoline**. *Desoxycordifoline*

[21451-50-3]

$C_{28}H_{30}N_2O_{11}$ 570.552

Alkaloid from the heartwood of *Adina cordifolia* and *Adina rubescens* (Rubiaceae). Cryst. (Me₂CO/MeOH). Mp 178-180°. $[\alpha]_D^{25} -212$ (H₂O). Lacks the phenolic OH group. λ_{\max} 239 (log ϵ 4.58); 268 (log ϵ 4.54); 336 (log ϵ 3.7); 349 (log ϵ 3.71) (EtOH).

Deoxy, *parent acid*: **Desoxycordifolinic acid**

[88839-94-5]

$C_{27}H_{28}N_2O_{11}$ 556.525

Alkaloid from the heartwood of *Naucllea diderrichii* (Rubiaceae). Brown rosettes + 1H₂O (MeCN aq.). Mp 206-208° dec. $[\alpha]_D -45.7$ (c, 0.126 in MeOH).

Decarboxy: see Lyaloside, L-316

Brown, R.T. *et al.*, *Chem. Comm.*, 1967, 453 (*isol*, *uv*, *pmr*, *ms*)

Merlini, L. *et al.*, *Gazz. Chim. Ital.*, 1968, **98**, 974-977 (*Cordifoline*, *Deoxycordifoline*, *uv*, *pmr*, *ms*)

Brown, R.T. *et al.*, *Phytochemistry*, 1978, **17**, 1686-1687 (*Cordifoline*, *Deoxycordifoline*, *isol*, *uv*, *ir*, *pmr*, *ms*)

Adeoye, A.O. *et al.*, *Phytochemistry*, 1983, **22**, 2097-2098 (*Desoxycordifolinic acid*)


Brandt, V. *et al.*, *Phytochemistry*, 1999, **51**, 1171-1176 (*Deoxycordifoline*, *uv*, *pmr*, *cmr*, *ms*)

Cardoso, C.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1882-1885 (*cmr*)

Cordizine

C-642

[863203-35-4]



Absolute Configuration

$C_{22}H_{33}NO$ 327.509

Alkaloid from *Delphinium corymbosum*. Cryst. (MeOH). Mp 122-124°.


Salimov, B.T. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 579-581 (*isol*, *pmr*, *ms*)

Cordobine

C-643

(-)-*Antioquine*

[121960-20-1]



Alkaloids covered by this entry (1*R*,1'*S*') are enantiomeric with those in Rodiasine, R-114 (1*S*,1'*R*') and diastereomeric with those in Guattaguanine, G-212 (1*S*,1'*S*'). Cordobine is the enantiomer of (+)-Antioquine in R-114. Alkaloid from an unidentified *Crematosperma* sp. and *Guatteria boliviana* (Annonaceae). Exhibits antiparasitic and cytotoxic activities. Amorph. $[\alpha]_D^{20} -170$ (c, 0.2 in $CHCl_3$).

O^{12'}-*Me*: **Monterine**

[121892-85-1]

$C_{38}H_{42}N_2O_6$ 622.76

Alkaloid from the stem bark of an unidentified *Crematosperma* sp. (Annonaceae). Amorph. $[\alpha]_D^{20} -135$ (c, 0.4 in $CHCl_3$).

Di-Me ether: **Granjine**

[121959-06-6]

$C_{39}H_{44}N_2O_6$ 636.786

Alkaloid from the stem bark of an unidentified *Crematosperma* sp. (Annonaceae). Amorph. $[\alpha]_D^{20}$ -63 (c, 0.6 in CHCl_3).


Saez, J. *et al.*, *Can. J. Chem.*, 1989, **67**, 275 (isol, struct)

Jossang, A. *et al.*, *J.O.C.*, 1996, **61**, 3023 (conformn)

Mahiou, V. *et al.*, *Phytochemistry*, 2000, **54**, 709-716 (isol, cd, pmr, cmr, ms)

Cordrastine C-644

6,7-Dimethoxy-3-(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)-1(3H)-isobenzofuranone, 9CI



$\text{C}_{22}\text{H}_{25}\text{NO}_6$ 399.443

Poss. alkaloid from *Corydalis aurea* (Papaveraceae). Fine needles ($\text{CHCl}_3/\text{MeOH}$). Mp 196° (prev. darkening). Struct. not certain, stereochem. not detd. Cordrastine is, however, well characterised as a synthetic compd. and the Mp of the alkaloid corresponds with synthetic Cordrastine I.

(1R,9R)-form

(-)-Cordrastine I

[34408-08-7]

Synthetic. Cryst. ($\text{CHCl}_3/\text{MeOH}$ or EtOH). Mp 196° (189-190 $^\circ$). $[\alpha]_D$ -99 (c, 1 in CHCl_3).

(1R,9S)-form

(-)-Cordrastine II

[34408-07-6]

Synthetic. Cryst. (Et₂O/petrol). Mp 90° . $[\alpha]_D$ -10 (c, 1.0 in CHCl_3).

Hydrobromide: Mp 212-213 $^\circ$. $[\alpha]_D$ +188 (c, 1 in MeOH).

(1RS,9RS)-form

(±)-Cordrastine I

[51547-89-8]

Synthetic. Cryst. (MeOH). Mp 156-157 $^\circ$.

(1RS,9SR)-form

(±)-Cordrastine II

[51547-90-1]

Synthetic. Cryst. (MeOH). Mp 119° .

Picrate: Mp 208° (202 $^\circ$) dec.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 81 (isol)

Haworth, W. *et al.*, *J.C.S.*, 1950, 1777

Teitel, S. *et al.*, *J.O.C.*, 1972, **37**, 1879 (synth, pmr, cd, uv)

Smula, V. *et al.*, *Can. J. Chem.*, 1973, **51**, 3287 (synth, ir, pmr)

Kametani, T. *et al.*, *J. Chem. Soc. Pak.*, 1976, 1221 (synth)

de Silva, S.O. *et al.*, *Can. J. Chem.*, 1979, **57**, 1598 (synth)


Shono, T. *et al.*, *J.O.C.*, 1983, **48**, 1621 (synth)

Orito, K. *et al.*, *Heterocycles*, 1987, **26**, 3077 (synth)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 869 (synth)

Cordyformamide C-645

[930783-39-4]




$\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_6$ 356.334

Related to Xanthocillin X, X-7 and Melanocin A, M-154. Isol. from *Cordyceps brunnearubra* BCC 1395. Antimalarial agent. Powder. λ_{max} 204 (log ϵ 4.49); 348 (log ϵ 4.46) (MeOH).

Isaka, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 656-658 (isol, pmr, cmr)

Cordypyridone A C-646

3-(2-Ethenyl-2,4,6-trimethylcyclohexyl)-1,4-dihydroxy-2(1H)-pyridinone, 9CI. 8-Methylpyridoxatin



Absolute Configuration

$\text{C}_{16}\text{H}_{23}\text{NO}_3$ 277.363

Prod. by the fungi OS-F61800 and *Cordyceps nipponica*. Induces erythropoietin gene expression. Plates. Mp 261-263 $^\circ$ dec. $[\alpha]_D^{25}$ -11 (c, 0.5 in MeOH). λ_{max} 216 (ϵ 34000); 289 (ϵ 6850) (MeOH).

Atropisomer: Cordypyridone B


$\text{C}_{16}\text{H}_{23}\text{NO}_3$ 277.363

Prod. by *Cordyceps nipponica*. Prisms (Me₂CO). Mp 252-255 $^\circ$ dec. $[\alpha]_D^{26}$ +164 (c, 0.14 in MeOH). λ_{max} 212 (log ϵ 4.54); 246 (log ϵ 3.61); 288 (log ϵ 3.72) (MeOH).

Cai, P. *et al.*, *J. Nat. Prod.*, 1999, **62**, 397-399 (isol, uv, ir, pmr, cmr, ms)

Isaka, M. *et al.*, *J.O.C.*, 2001, **66**, 4803-4808 (isol, uv, ir, pmr, cmr, ms)

Cordypyridone D C-647



Relative Configuration

$\text{C}_{17}\text{H}_{25}\text{NO}_4$ 307.389

Similar to Fusaricide, F-225. Prod. by *Cordyceps nipponica*. Cryst. Mp 206-209 $^\circ$ dec. $[\alpha]_D^{26}$ +249 (c, 0.14 in MeOH). λ_{max} 218 (log ϵ 4.47); 293 (log ϵ 3.75) (MeOH).

Deoxy: Cordypyridone C


$\text{C}_{17}\text{H}_{25}\text{NO}_3$ 291.389

Prod. by *Cordyceps nipponica*. Cryst. Mp 150-151 $^\circ$ dec. $[\alpha]_D^{24}$ +243 (c, 0.06 in MeOH). λ_{max} 218 (log ϵ 4.41); 292 (log ϵ 3.67) (MeOH).

Isaka, M. *et al.*, *J.O.C.*, 2001, **66**, 4803-4808

Coriainenine A C-648

[182011-67-2]



$\text{C}_{28}\text{H}_{58}\text{N}_4\text{O}_2$ 482.792

Alkaloid from the sponge *Clathrina coriacea*. Yellow oil (as per-Ac). $[\alpha]_D$ -6.1 (c, 0.24 in CHCl_3) (per-Ac).

Didehydro: Coriainenine B

[182265-33-4]


$\text{C}_{28}\text{H}_{56}\text{N}_4\text{O}_2$ 480.776

Alkaloid from *Clathrina coriacea*. Yellow oil (as per-Ac). $[\alpha]_D$ -6.3 (c, 0.08 in CHCl_3) (per-Ac). Posn. of double bond not known, but is located between the other 2 double bonds.

Casapullo, A. *et al.*, *J.O.C.*, 1996, **61**, 7415-7419

Coriainenine D C-649

[182011-68-3]



$\text{C}_{30}\text{H}_{58}\text{N}_4\text{O}_2$ 506.814

Alkaloid from the sponge *Clathrina coriacea*. Yellow oil (as per-Ac). $[\alpha]_D$ -10 (c, 0.36 in CHCl_3).

Dihydro: Coriainenine C

[182149-41-3]


$\text{C}_{30}\text{H}_{60}\text{N}_4\text{O}_2$ 508.83

Alkaloid from *Clathrina coriacea*. Yellow oil (as per-Ac). $[\alpha]_D$ -6.2 (c, 0.16 in CHCl_3). Posn. of one double bond not determined.

Casapullo, A. *et al.*, *J.O.C.*, 1996, **61**, 7415-7419

Coriainenine E C-650

[182149-43-5]




$\text{C}_{29}\text{H}_{58}\text{N}_4\text{O}_2$ 494.803

Posn. of one double bond not determined. Alkaloid from the sponge *Clathrina coriacea*. Yellow oil.

Casapullo, A. *et al.*, *J.O.C.*, 1996, **61**, 7415-7419 (isol, struct)

Corialstonidine

[155416-41-4]


C₂₂H₂₆N₂O₅ 398.458

Stereochem. not reported but prob. the same as for Corialstonine, C-652. Alkaloid from *Alstonia coriacea*. Shows minor antimalarial activity (ca. 10% of quinine). No phys. props. reported.

Phillipson, J.D. *et al.*, *J. Ethnopharmacol.*, 1993, **40**, 41-45 (*isol*)

Corialstonine

[113145-60-1]


C₂₃H₂₆N₂O₅ 410.469

Alkaloid from the stem bark of *Alstonia coriacea* (Apocynaceae). Shows minor antimalarial activity (approx. 10% that of quinine). Amorph. [α]_D +102 (c, 1 in CHCl₃).

Cherif, A. *et al.*, *Phytochemistry*, 1989, **28**, 667-670 (*isol, uv, ir, pmr, cmr, ms, struct*)

Corlumine

6-(1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI



(+)-form

C₂₁H₂₁NO₆ 383.4

Diastereoisomer of Adlumine, A-146.

(+)-form [485-51-8]

Alkaloid from *Corydalis scouleri*, *Corydalis govaniana*, *Corydalis nobilis*, *Corydalis severtzovii*, *Corydalis sibirica* and *Dicentra cucullaria* (Papaveraceae).

GABA antagonist. Convulsive agent. Mp 162° (159°). [α]_D²⁶ +77 (c, 1.13 in CHCl₃).

▶ GM3500000

O⁷-De-Me: Corlumidine

[25344-54-1]

C₂₀H₁₉NO₆ 369.373

C-651

Alkaloid from *Corydalis scouleri*.Cryst. (CHCl₃ or MeOH). Mp 236°.[α]_D²³ +80 (c, 0.4 in CHCl₃).**Di-O-de-Me: Papraïne**

[125263-86-7]

C₁₉H₁₇NO₆ 355.346

Alkaloid from whole plant of *Fumaria indica* (Papaveraceae). Amorph. solid. [α]_D²⁴ +25.

(-)-form [79082-64-7]

Alkaloid from *Fumaria parviflora* (Papaveraceae). Mp 158-160° dec. (synthetic). [α]_D²⁶ -73 (c, 1.32 in CHCl₃) (synthetic). Natural alkaloid obt. as an oil, characterised by CD.

(±)-form [64397-12-2]

Synthetic. Mp 175-176°.

[130480-92-1]

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1936, **14**, 325; 347 (*isol, Corlumine, Corlumidine*)

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1961, **39**, 1801 (*isol, uv, ir, pmr*)

Safe, S. *et al.*, *Can. J. Chem.*, 1964, **42**, 160 (*pmr, config*)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 2328 (*isol*)

Yunusov, M.S. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 61; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 54 (*isol*)

Snatzke, G. *et al.*, *Tetrahedron*, 1969, **25**, 5059 (*ord, abs config*)

Shamma, M. *et al.*, *Tet. Lett.*, 1974, 2339 (*pmr, synth*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 612; 811; 1978, **14**, 103; 465; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 642; 826; 1978, **14**, 82; 395 (*isol, cd*)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (*cmr*)

Nalliah, B.C. *et al.*, *Can. J. Chem.*, 1979, **57**, 1546 (*synth*)

Blaskó, G. *et al.*, *J. Nat. Prod.*, 1981, **44**, 475 (*isol*)

Huber, I.M.P. *et al.*, *Helv. Chim. Acta*, 1987, **70**, 1944 (*synth, ir, pmr, cmr, ms*)


Atta-ur-Rahman, *et al.*, *Heterocycles*, 1989, **29**, 1091 (*Papraïne*)

Seeger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**, 882-886 (*pmr, cmr*)

C-652

Cornigerine

C-654

C₂₁H₂₁NO₆ 383.4**(S)-form** [6877-25-4]

Alkaloid from *Colchicum cornigerum*, *Colchicum cilicicum*, *Iphigenia bechuani-ca*, *Iphigenia indica*, *Iphigenia pallida*, *Androcymbium melanthioides* and *Camp-torrhiza strumosa*. Reported to exhibit similar biol. activity to Colchicine, C-569. Prisms (EtOAc/Et₂O). Sol. MeOH, CHCl₃, Mp 268-270°. [α]_D²⁰ -150 (c, 0.631 in CHCl₃). [α]_D²⁰ -233 (c, 0.728 in MeOH).

▶ AC8587000

O¹⁰-De-Me: Cornigereine

Alkaline saponification prod. of Cornigerine. Yellow prisms (EtOAc or Me₂CO). Mp 168-170°. [α]_D²² -222 (c, 0.758 in CHCl₃). [α]_D²² -245 (c, 0.700 in MeOH).

4-Methoxy: Gloriosamine A

[1006585-02-9]

C₂₂H₂₃NO₇ 413.426

Alkaloid from *Gloriosa rothschildiana*. Amorph. solid. [α]_D²⁵ -35 (c, 0.1 in CHCl₃). λ _{max} 218; 235; 351 (EtOH).

4-Methoxy, N-de-Ac, N-(hydroxyacetyl): Gloriosamine B

[1006585-03-0]

C₂₂H₂₃NO₈ 429.426

Alkaloid from *Gloriosa rothschildiana*. Amorph. solid. [α]_D²⁵ -60 (c, 0.1 in CHCl₃). λ _{max} 219; 235; 351 (EtOH).

El-Hamidi, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 2111 (*isol, uv, ir*)

Cross, A.D. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 1187 (*pmr, ms*)

Kaul, J.L. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 1689 (*isol*)

Rösner, M. *et al.*, *J.O.C.*, 1981, **46**, 3686 (*struct, ir, pmr, synth*)


Potěšilová, H. *et al.*, *Planta Med.*, 1985, 344 (*isol*)

Kitajima, M. *et al.*, *Tet. Lett.*, 2008, **49**, 257-260 (*Gloriosamines A, B*)

Cornucervine

C-655

[31948-48-8]

C₁₇H₂₉NO₅ 327.42


Ester of Trachelanthamidine in H-629.

Alkaloid from *Phalaenopsis cornu-cervi* (Orchidaceae). Oil. [α]_D²² -4.3 (c, 1.6 in EtOH).

Brandänge, S. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 349; 1973, **27**, 3668 (*isol, struct, abs config*)

Coronafacic acid

C-656

C₁₂H₁₆O₃ 208.257**(+)-form** [62251-98-3]

The unnatural racemic form also prepd. A component of Coronatine. Also isol. from *Pseudomonas coronofaciens* and *Pseudomonas syringae*. Prisms (diisopropyl ether). Mp 142-143°. [α]_D²⁰ +119.1 (c, 3.3 in MeOH).

L-Serine amide: N-CoronafacylserineC₁₅H₂₁NO₅ 295.335Prod. by *Pseudomonas syringae*.

Chlorosis inducer.

L-Threonine amide: N-Coronafacoyl-threonine

C₁₆H₂₃NO₅ 309.361
Prod. by *Pseudomonas syringae*.
Chlorosis inducer.

L-Isoleucine amide: N-Coronafacoyl-isoleucine

[73344-10-2]
C₁₈H₂₇NO₄ 321.416
Prod. by *Pseudomonas syringae* var. *glycinea*. Chlorosis inducer. Intermediate in biosynth. of Coronatine. Powder. Mp 53-55°. [α]_D²⁵ +44 (c, 2.25 in MeOH). λ_{max} 227; 265; 275; 294; 300 (MeOH) (Berdy).

L-Alloisoleucine amide: N-Coronafacoyl-alloisoleucine

[99630-78-1]
C₁₈H₂₇NO₄ 321.416
Prod. by *Pseudomonas syringae* var. *glycinea*. Chlorosis inducer. Intermediate in Coronatine biosynth.

L-Valine amide: N-Coronafacoylvaline

[91793-99-6]
C₁₇H₂₅NO₄ 307.389
Prod. by *Pseudomonas syringae*. Chlorosis inducer. λ_{max} 227; 265; 275; 294; 300 (MeOH) (Chem.).

3a-Epimer: [87335-74-8]
Prisms. Mp 142-143°. [α]_D²⁰ -119 (c, 0.86 in MeOH).

7a-Epimer: [62251-99-4]
Formed in hydrol. of Coronatine. Mp 141-142°.

(±)-form [66322-23-4]
Mp 130-132°.

[92843-35-1]

Shiraishi, K. *et al.*, *Agric. Biol. Chem.*, 1979, **43**, 1753-1757 (*amino acid amides, synth, bibl*)

Ohira, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 1902-1907 ((±)-*form, synth, resolu*)

Liu, H.J. *et al.*, *Can. J. Chem.*, 1984, **62**, 1747-1750 ((±)-*form, synth*)

Mitchell, R.E. *et al.*, *Phytochemistry*, 1984, **23**, 791-793; 1985, **24**, 2716-2717 (*amino acid amides, isol*)

Hoelder, S. *et al.*, *Synlett*, 1996, 505-506 (*synth*)

Toshima, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 752-753 ((+)-*form, synth*)

Nara, S. *et al.*, *Tetrahedron*, 1997, **53**, 9509-9524 ((+)-*form, synth*)

Mitchell, R.E. *et al.*, *Phytochemistry*, 1998, **49**, 1579-1583 (*amides*)


Sono, M. *et al.*, *Tet. Lett.*, 2000, **41**, 5115-5118 ((+)-*form, synth*)

Mehta, G. *et al.*, *J.C.S. Perkin 1*, 2001, 1153-1161 ((+)-*form, synth*)

Moreau, B. *et al.*, *J.O.C.*, 2007, **72**, 1235-1240 ((±)-*form, synth*)

Coronaridine**C-657**

Carbomethoxyibogamine
[467-77-6]



Absolute
Configuration

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Tabernaemontana coronaria* (*Ervatamia coronaria*), *Tabernaemontana divaricata*, *Tabernanthe iboga*, *Tabernanthe pubescens* and many other spp. in Apocynaceae. Noncryst. Mp 235° dec. (as hydrochloride). [α]_D²⁵ -8.5 (c, 1 in MeOH) (hydrochloride). λ_{max} 244 (ε 35500); 284 (ε 7850); 292 (ε 6780) (MeOH). λ_{max} 225; 285; 293 (EtOH).

5,6-Didehydro: 5,6-Dehydrocoronaridine

[162229-93-8]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from bark of *Tabernaemontana markgrafiana* (Apocynaceae). Amorph. Unstable; dec. in CHCl₃.

3R-Hydroxy: Eglantine. 3-Hydroxycoronaridine

[53508-36-4]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from *Gabunia eglanulosa* (preferred genus name *Tabernaemontana*), *Tabernanthe iboga* and *Muntafara sessilifolia* (preferred genus name *Tabernaemontana*) (Apocynaceae). Shows strong antibiotic activity. Noncryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D -57 (CHCl₃). Struct. revised in 1985 (originally described as 3,6-Oxidocoronaridine). Some isolates of 3-hydroxycoronaridine were of undetd. C3-config, but their props. were closely similar to those of Eglantine. λ_{max} 228 (log ε 4.41); 285 (log ε 3.9); 292 (log ε 3.86) (MeOH).

3R-Methoxy: 3R-Methoxycoronaridine

[162229-94-9]
C₂₂H₂₈N₂O₃ 368.475
Alkaloid from bark of *Tabernaemontana markgrafiana* (Apocynaceae). Amorph. light yellow solid. [α]_D -44 (c, 0.3 in CHCl₃).

3R-Ethoxy: 3-Ethoxycoronaridine

[74627-73-9]
C₂₃H₃₀N₂O₃ 382.502
Alkaloid from *Tabernanthe glandulosa* (Apocynaceae). [α]_D²⁰ -59. 3-Config. undetd. but prob. R-. λ_{max} 225 (log ε 4.51); 286 (log ε 3.92); 292 (log ε 3.88) (no solvent reported).

18-Hydroxy: Albifloranine

[77431-58-4]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from the stem bark of *Tabernaemontana albiflora* (Apocynaceae). Cryst. (MeOH). Mp 192-194°. [α]_D²⁰ -210 (c, 1 in CHCl₃).

19R-Hydroxy: 19-Epiheyneanine

[57511-56-5]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from *Pandaca mocquersyia* var. *pendula* and the root bark of *Peschiera affinis* (Apocynaceae). Cryst. (cyclohexane). Mp 170-172°. [α]_D²⁴ -46 (CHCl₃). λ_{max} 226 (ε 26900); 280 (ε 5500); 285 (ε 6200); 293 (ε 5300) (EtOH).

19S-Hydroxy: Heyneanine

[4865-78-5]
[76129-69-6]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from *Ervatamia heyneana*

(preferred genus name *Tabernaemontana*), *Tabernaemontana sananho*, *Conopharyngia jollyana* and *Pandaca* spp. (Apocynaceae). Shows mod. cytotoxicity vs. mouse P388 carcinoma cells. Mp 105-107° Mp 160-162° (dimorph.). [α]_D³⁰ -19 (c, 0.8 in CHCl₃). λ_{max} 224 (log ε 4.71); 284 (log ε 4.33); 292 (log ε 4.27) (MeOH).

3-Ethoxy, 19R-hydroxy: 3-Ethoxy-19-epiheyneanine

[437764-35-7]
C₂₃H₃₀N₂O₄ 398.501
Alkaloid from *Tabernaemontana corymbosa*. Oil. Isol. as a 1:1 mixt. of 3R- and 3S-isomers. λ_{max} 225 (log ε 3.91); 285 (log ε 3.29); 292 (log ε 3.23) (EtOH).

3-Ethoxy, 19S-hydroxy: 3-Ethoxyheyneanine

[437764-34-6]
C₂₃H₃₀N₂O₄ 398.501
Alkaloid from *Tabernaemontana corymbosa*. Oil. Isol. as a 1:1 mixt. of 3R- and 3S-isomers. λ_{max} 225 (log ε 3.94); 285 (log ε 3.33); 292 (log ε 3.26) (EtOH).

18,19R-Dihydroxy: 18,19R-Dihydroxycoronaridine

[167696-84-6]
C₂₁H₂₆N₂O₄ 370.447
Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae). [α]_D -6 (c, 0.3 in MeOH). λ_{max} 213; 228 (sh); 275 (sh); 285; 294 (MeOH).

3-Oxo: Eglanuloline. 3-Oxocoronaridine

[53508-33-1]
C₂₁H₂₄N₂O₃ 352.432
Alkaloid from *Tabernanthe dichotoma*, *Gabunia eglanulosa* and *Muntafara sessilifolia* (Apocynaceae). Noncryst. [α]_D -71 (CHCl₃). Struct. revised in 1985; originally descr. as 6-Hydroxy-3-oxocoronaridine. An alkaloid given this struct. was earlier detected by ms in *Conopharyngia jollyana* (Hooteele *et al.*, 1968), but it is not clear that it is the same as Eglanuloline.

3-Oxo, 19R-hydroxy: 3-Oxo-19-epiheyneanine

[131670-18-3]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from the leaves and stem bark of *Ervatamia polyneura* (Apocynaceae). [α]_D -49 (c, 0.5 in CHCl₃).

3-Oxo, 19ξ-hydroxy: 19-Hydroxy-3-oxocoronaridine

C₂₁H₂₄N₂O₄ 368.432
Alkaloid detected by ms in *Conopharyngia jollyana* (Apocynaceae).

5-Oxo: 5-Oxocoronaridine

[76129-70-9]
C₂₁H₂₄N₂O₃ 352.432
Alkaloid from the root bark of *Tabernaemontana divaricata* (Apocynaceae). Mp 272-275°. [α]_D²⁵ -10.7 (Py). λ_{max} 229; 285; 292 (MeOH).

6-Oxo: 6-Oxocoronaridine

[76129-71-0]
C₂₁H₂₄N₂O₃ 352.432
Alkaloid from the root bark of *Tabernaemontana divaricata* (Apocynaceae).

Mp 262–267°. $[\alpha]_D^{25}$ -33.8 (MeOH). λ_{\max} 218 ; 255 ; 260 (sh) ; 268 (sh) ; 335 (MeOH).

6-Oxo, 5-hydroxy: 5-Hydroxy-6-oxocoronaridine

[76129-72-1]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the root bark of *Tabernaemontana divaricata* (Apocynaceae). Mp 285–288°. $[\alpha]_D^{25}$ +43.8 (Py). λ_{\max} 215 ; 248 ; 263 (sh) ; 267 (sh) ; 310 (MeOH).

5,6-Dioxo: 5,6-Dioxocoronaridine

[909120-04-3]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from the roots of *Ervatamia hainanensis*.

10-Methoxy: see Voacangine, V-170

11-Methoxy: see Isovoacangine, I-344

10,11-Dimethoxy: see Conopharyngine, C-620

3S-Cyano: 3S-Cyanocoronaridine

[746642-97-7]

C₂₂H₂₅N₃O₂ 363.458

Alkaloid from the stem bark of *Tabernaemontana divaricata*. Cytotoxic. Light yellow oil. $[\alpha]_D$ -64 (c, 1.72 in CHCl₃). λ_{\max} 225 (log ϵ 4.68); 276 (log ϵ 4.02); 285 (log ϵ 4.06); 293 (log ϵ 4.01) (EtOH).

(±)-form [14853-22-6]

19-Hydroxy: (±)-19-Hydroxycoronaridine

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the root bark of *Tabernaemontana divaricata* (Apocynaceae). Mp 159°. Identity with (±)-Heyneanine not clearly establ. λ_{\max} 230 ; 285 ; 294 (MeOH).

Gorman, M. et al., *J.A.C.S.*, 1960, **82**, 1142–1145 (isol, struct)

Govindachari, T.R. et al., *Tet. Lett.*, 1965, 3873–3878 (Heyneanine, isol, uv, ir, pmr, ms, struct)

Govindachari, T.R. et al., *Chem. Comm.*, 1966, 97 (struct)

Hirai, S. et al., *Chem. Comm.*, 1968, 1016–1017 (synth, ir, uv)

Hootele, C. et al., *Chimia*, 1968, **22**, 245–246 (3-Oxocoronaridine, 19-Hydroxy-3-oxocoronaridine)

Kutney, J.P. et al., *J.A.C.S.*, 1970, **92**, 1708–1712; 1973, **95**, 5407–5409 (synth, cryst struct, config)

Le Men, J. et al., *Bull. Soc. Chim. Fr.*, 1974, 1369–1372 (Eglantine, Eglantulosine)

Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1974, **39**, 2258–2266 (abs config, bibl)

Panas, J.M. et al., *Phytochemistry*, 1975, **14**, 1120–1122 (Eglantulosine, isol, ms)

De Bellefont, M. et al., *Phytochemistry*, 1975, **14**, 1649–1652 (19-Epiheyneanine)

Wenkert, E. et al., *Helv. Chim. Acta*, 1976, **59**, 2437 (cmr)

Matos, F.J.A. et al., *Phytochemistry*, 1976, **15**, 551–553 (19-Epiheyneanine)

Achenbach, H. et al., *Phytochemistry*, 1980, **19**, 716–717; 2185–2188 (3-Ethoxycoronaridine)

Rastogi, K. et al., *Phytochemistry*, 1980, **19**, 1209–1212 (*Tabernaemontana divaricata constits*)

Gunasekara, S.P. et al., *Phytochemistry*, 1980, **19**, 1213–1218 (*Coronaridine, Heyneanine, isol, activity*)

Kan, C. et al., *Planta Med.*, 1981, **41**, 72–74 (*Albifloranine*)

Le Men-Olivier, L. et al., *Bull. Soc. Chim. Fr.*, 1985, 94–97 (*Eglantine, struct, pmr, cmr*)

Okuyama, E. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2075–2079 (5-(2-

Oxopropyl)coronaridine)

Bornmann, W.G. et al., *J.O.C.*, 1992, **57**, 1752–1760 (synth)

Nielsen, H.B. et al., *Phytochemistry*, 1994, **37**, 1729–1735 (5,6-Dehydrocoronaridine, 3R-Methoxycoronaridine)

Azoug, M. et al., *Phytochemistry*, 1995, **39**, 1223–1228 (18,19-Dihydroxycoronaridine)

Bandarage, U.K. et al., *Tetrahedron*, 1999, **55**, 9405–9424 (*Albifloranine, synth*)

Kuehne, M.E. et al., *Tetrahedron*, 2001, **57**, 2085–2094 (synth)

Kam, T.-S. et al., *J. Nat. Prod.*, 2002, **65**, 669–672 (3-Ethoxyheyneanine, 3-Ethoxy-19-epiheyneanine)


Kam, T.-S. et al., *Chem. Biodiversity*, 2004, **1**, 646–656 (3S-Cyanocoronaridine)

Huang, J.P. et al., *Chin. Chem. Lett.*, 2006, **17**, 779–782 (5,6-Dioxocoronaridine)

Pereira, P.S. et al., *Quim. Nova*, 2008, **31**, 20–24 (pmr, cmr)

Coronaridine hydroxyindolenine C-658

[16671-16-2]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Conopharyngia durissima*, from the root bark of *Tabernaemontana divaricata*, from bark of *Tabernaemontana markgrafiana* and roots of *Tabernaemontana quadrangularis* (Apocynaceae). Mp 113–117° (98–100°). $[\alpha]_D$ -8 (EtOH). $[\alpha]_D^{25}$ -16 (MeOH). λ_{\max} 225 (log ϵ 4.45); 271 (log ϵ 3.79); 285 (log ϵ 3.95); 295 (log ϵ 3.92); 314 (sh) (log ϵ 3.48) (MeOH).

Hydrochloride:

Cryst. (MeOH/Me₂CO). Mp 278–279°.

19S-Hydroxy: Heyneanine hydroxyindolenine

[80151-98-0]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from whole plant of *Ervatamia coronaria* var. *plena* (Apocynaceae). Amorph. powder. Epimeric 9-OH config. shown in the paper. λ_{\max} 223 ; 282 (MeOH).

10-Methoxy, 19R-hydroxy: 19-Epivoacristine hydroxyindolenine

[18646-17-8]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Tabernaemontana calcarea*. Oil. $[\alpha]_D$ +6 (c, 0.36 in CHCl₃). λ_{\max} 220 (log ϵ 3.92); 283 (log ϵ 4.12); 292 (log ϵ 3.76); 302 (log ϵ 3.62); 316 (log ϵ 3.51) (MeOH).

10-Methoxy, 19S-hydroxy: Voacristine hydroxyindolenine

[15215-86-8]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from the root bark of *Ervatamia dichotoma* (Apocynaceae).

Cryst. (C₆H₆/petrol). Mp 176–179° dec. $[\alpha]_D^{26}$ -22 (c, 0.51 in CHCl₃). 7-Config. unspecified. λ_{\max} 230 (c 12380); 268 (c 4400); 291 (c 4780); 300 (sh) (c 4410); 314 (c 3810) (EtOH).

10,11-Dimethoxy: Jollyanine†. Conopharyngine hydroxyindolenine

[16671-17-3]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from *Conopharyngia jollyana* and *Tabernaemontana cumminsii* (Apocynaceae). Cryst. (MeOH). Mp 161–163°. $[\alpha]_D^{22}$ -56 (c, 0.6 in CHCl₃). 7-Config. undefined. λ_{\max} 217 (c 13800); 238 (c 13300); 300 (c 4700) (no solvent reported).

10,11-Dimethoxy, 3-hydroxy: 3-Hydroxyconopharyngine hydroxyindolenine

[93627-66-8]

C₂₃H₃₀N₂O₆ 430.5

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Mixt. of epimers. Ratio 3R:3S = 1:1. Probably an artifact. λ_{\max} 232 ; 298 ; 306 (sh) ; 328 (sh) (MeOH).

7-Epimer, 10-methoxy: Voacangine hydroxyindolenine

[3464-63-9]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from *Voacanga africana*, *Tabernaemontana markgrafiana*, *Tabernaemontana quadrangularis* and *Peschiera campestris* (Apocynaceae). Shows mod.-weak cytotoxic activity. Cryst. (Et₂O) or amorph. yellow-green solid. Mp 135–137° (129–132°). $[\alpha]_D$ -26 (c, 1.1 in CHCl₃) (-18). λ_{\max} 228 (log ϵ 4.44); 272 (log ϵ 3.78); 286 (log ϵ 3.79); 293 (log ϵ 3.78); 313 (sh) (log ϵ 3.68) (MeOH).

Hootele, C. et al., *Bull. Soc. Chim. Belg.*, 1967, **76**, 303–307 (*Jollyanine*)

Das, B.C. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1967, **264**, 1765–1767 (isol, ir, uv, ms, struct)

Schnoes, H.K. et al., *J.O.C.*, 1968, **33**, 1225–1227 (*Voacristine hydroxyindolenine*)

Thomas, D.W. et al., *Tetrahedron*, 1968, **24**, 4223–4231 (*Voacangine hydroxyindolenine*)

Hwang, B. et al., *J.O.C.*, 1969, **34**, 412–415 (*Voacristine hydroxyindolenine, synth*)

Crooks, P.A. et al., *J. Pharm. Pharmacol.*, 1970, **22**, 471–472 (*Jollyanine*)

Harmouche, A. et al., *Ann. Pharm. Fr.*, 1976, **34**, 31–35 (pmr)

Rastogi, K. et al., *Phytochemistry*, 1980, **19**, 1209–1212 (isol, uv, ir, pmr, ms)

Achenbach, H. et al., *Z. Naturforsch., B*, 1980, **35**, 219–225 (*Coronaridine hydroxyindolenine, Voacangine hydroxyindolenine, isol, uv, ir, ms, cd, abs config*)

Van Beek, T.A. et al., *J. Nat. Prod.*, 1985, **48**, 400–428 (3-Hydroxyconopharyngine hydroxyindolenine)

Gower, A.E. et al., *Phytochemistry*, 1986, **25**, 2908–2910 (*Voacangine hydroxyindolenine*)

Sharma, P. et al., *J. Nat. Prod.*, 1988, **51**, 528–531 (*Heyneanine hydroxyindolenine*)


Nielsen, H.B. et al., *Phytochemistry*, 1994, **37**, 1729–1735 (*Voacangine hydroxyindolenine*)

Madinaveitia, A. et al., *Helv. Chim. Acta*, 1998, **81**, 1645–1653 (*Voacangine hydroxyindolenine, abs config*)

Chaturvedula, V.S.P. et al., *J. Nat. Prod.*, 2003, **66**, 528–531 (19-Epivoacristine hydroxyindolenine)

Pereira, P.S. *et al.*, *Quim. Nova*, 2008, **31**, 20-24 (pmr, cmr)

Correantine A C-659
[164178-50-1]




$C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from leaves of *Psychotria correae* (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} +211$ (c, 0.31 in MeOH).

3-Epimer: 3-Isocorreantine A
 $C_{22}H_{26}N_2O_4$ 382.458
Formed *in vitro* by enzymatic hydrolysis of Dolichantoside in S-583.

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1537 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Gerasimenko, I. *et al.*, *Eur. J. Biochem.*, 2002, **269**, 2204-2213 (biosynth, 3-Isocorreantine A)

Correantine B C-660
[164178-51-2]




$C_{21}H_{22}N_2O_3$ 350.416
Alkaloid from leaves of *Psychotria correae* (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} -81$ (c, 0.18 in MeOH).

20-Epimer: 20-Epicorreantine B
[164322-84-3]

$C_{21}H_{22}N_2O_3$ 350.416
Alkaloid from leaves of *Psychotria correae* (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} -102$ (c, 0.10 in MeOH). Poss. artifact.

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1537 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Correantine C C-661
[164178-49-8]




$C_{21}H_{22}N_2O_3$ 350.416
Alkaloid from leaves of *Psychotria correae*

reea (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} -177$ (c, 0.2 in MeOH).

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1537 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Correantoside C-662
[164178-48-7]




$C_{27}H_{32}N_2O_8$ 512.558
Alkaloid from leaves and roots of *Psychotria correae* (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} -83$ (c, 0.32 in MeOH).

10-Hydroxy: 10-Hydroxycorreantoside
[164032-50-2]

$C_{27}H_{32}N_2O_9$ 528.558
Alkaloid from leaves of *Psychotria correae* (*Cephaelis correae*) (Rubiaceae). Amorph. $[\alpha]_D^{25} -43$ (c, 0.11 in MeOH).

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1537 (isol, uv, ir, pmr, cmr, ms, cd, struct)


Corsutine C-663
[866776-05-8]



$C_{21}H_{27}NO_7$ 405.447
Alkaloid from *Cocculus hirsutus*. λ_{max} 270 (MeOH).

Yadav, S.B. *et al.*, *Indian J. Chem., Sect. B*, 2005, **44**, 212-214 (isol, pmr, struct)


Cortamidine oxide C-664



$C_{13}H_{17}N_3O_4S_2$ 343.427
Isol. from *Cortinarius* sp. $[\alpha]_D^{22} -10$ (c, 0.03 in MeOH). Possible artifact. λ_{max} 190 ; 240 (MeOH).

Nicholas, G.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 341-344

Cortistatin B C-665
[882976-96-7]



$C_{30}H_{36}N_2O_4$ 488.625
Alkaloid from the sponge *Corticium simplex*. Anti-angiogenic agent. $[\alpha]_D^{20} +15.6$ (c, 0.27 in MeOH). λ_{max} 224 (€ 44400) (MeOH).

16-Ketone: Cortistatin C
[882976-97-8]

$C_{30}H_{34}N_2O_4$ 486.61
Alkaloid from *Corticium simplex*. $[\alpha]_D^{20} -45$ (c, 0.71 in MeOH). λ_{max} 222 (€ 49800) (MeOH).

17 α -Hydroxy, 16-ketone: Cortistatin D
[882976-98-9]

$C_{30}H_{34}N_2O_5$ 502.609
Alkaloid from *Corticium simplex*. $[\alpha]_D^{20} -37.1$ (c, 0.45 in MeOH). λ_{max} 219 (€ 48900) (MeOH).

16-Deoxy: Cortistatin A
[882976-95-6]

$C_{30}H_{36}N_2O_3$ 472.626
Alkaloid from *Corticium simplex*. $[\alpha]_D^{20} +30.1$ (c, 0.56 in MeOH). λ_{max} 219 (€ 45600) (MeOH).

1,2,16-Trideoxy, 1,2-didehydro: Cortistatin J
[944804-62-0]

$C_{30}H_{34}N_2O$ 438.611
Alkaloid from *Corticium simplex*. Exhibits cytostatic props. Powder. $[\alpha]_D^{20} -54$ (c, 0.26 in $CHCl_3$). λ_{max} 223 (€ 46200); 269 (€ 3000); 280 (€ 8500); 293 (€ 21200) (MeOH).


Aoki, S. *et al.*, *J.A.C.S.*, 2006, **128**, 3148-3149 (isol, pmr, cmr, cryst struct)

Aoki, S. *et al.*, *Tet. Lett.*, 2007, **48**, 4485-4488 (*Cortistatin J*)

Shenvi, R.A. *et al.*, *J.A.C.S.*, 2008, **130**, 7241-7243 (*Cortistatin A*, synth)

Lee, H.M. *et al.*, *J.A.C.S.*, 2008, **130**, 16864-16866 (*Cortistatin A*, synth)

Cortistatine F C-666
[941271-08-5]




$C_{32}H_{50}N_2O$ 478.76
Isol. from the sponge *Corticium simplex*. Powder. $[\alpha]_D^{20} +142.7$ (c, 0.2 in $CHCl_3$). λ_{max} 269 (€ 20900); 280 (€ 27800); 292 (€ 21700) (MeOH).

Watanabe, Y. *et al.*, *Tetrahedron*, 2007, **63**, 4074-4079 (isol, pmr, cmr)

Cortistatine G

[941271-09-6]

C-667

C₃₁H₄₂N₂O 458.686Isol. from the sponge *Corticium simplex*.Powder. [α]_D²⁰ -52.7 (c, 0.88 in CHCl₃).λ_{max} 245 (ε 22600) (MeOH).**22,23-Dihydro: Cortistatine H**

[941271-10-9]

C₃₁H₄₄N₂O 460.701Isol. from *Corticium simplex*. Powder.[α]_D²⁰ -57.8 (c, 0.34 in CHCl₃). λ_{max} 242

(ε 20200) (MeOH).

22,23,24ξ,25,26,27,28,29ξ-Octahydro, N-Me: Cortistatine E

[941271-07-4]


C₃₂H₅₂N₂O 480.776Isol. from *Corticium simplex*. Powder.[α]_D²⁰ -45 (c, 0.5 in CHCl₃). λ_{max} 242 (ε 22500) (MeOH).Watanabe, Y. et al., *Tetrahedron*, 2007, **63**,

4074-4079 (isol, pmr, cmr)

Cortistatine L

[944804-65-3]

C-668

C₃₀H₃₆N₂O₂ 456.627Alkaloid from *Corticium simplex*. [α]_D²⁰ -28.9 (c, 0.2 in CHCl₃). λ_{max} 224 (ε 55900) (MeOH).**2-Deoxy: Cortistatine K**

[944804-64-2]

C₃₀H₃₆N₂O 440.627Alkaloid from *Corticium simplex*.Aoki, S. et al., *Tet. Lett.*, 2007, **48**, 4485-4488

(isol, pmr, cmr, ms)


Coruscine

[1356-01-0]

C-669

C₁₈H₂₃NO₅ 333.383Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine corusca* and *Brunsdonna tubergenii* (Amaryllidaceae). Prisms (Me₂CO). Mp 170°. [α]_D²⁵ +70 (c, 0.5 in CHCl₃).**Hydroiodide:**Prisms (H₂O). Mp 179-180° dec.**Methiodide:**Prisms (H₂O). Mp 311° dec.**Methyl perchlorate:**Rhomb (H₂O). Mp 278-280° dec.Boit, H.-G. et al., *Chem. Ber.*, 1957, **90**, 369-373 (isol, ir)Boit, H.-G. et al., *Naturwissenschaften*, 1960, **47**, 159 (isol)**Corycavamine**

C-670

4,6,7,14-Tetrahydro-5,14-dimethylbis[1,3]benzodioxolo[4,5-c:5',6'-g]azecin-13(5H)-one, 9CI. Corycavine*(R)*-formRR' = —CH₂—C₂₁H₂₁NO₅ 367.401

The name Corycavamine was applied to the racemate.

(R)*-form** [521-85-7]Alkaloid from root nodules of *Corydalis tuberosa* (Papaveraceae). Mp 148-149°.[α]_D²⁰ +167 (CHCl₃).(S)*-form** [144606-40-6]

Obt. by resoln. of the racemate. Needles.

Mp 145-146°. [α]_D²⁵ -170 (c, 1.3 in CHCl₃).***(±)*-form** [521-87-9]Alkaloid from *Corydalis tuberosa*, *Corydalis incisa* and several other *Corydalis* spp. (Papaveraceae). Intermed. in biosynth. of Corynoline, C-692. Cardioactive agent. Plates (EtOH). Mp 221-222°.**Hydrochloride:** Mp 219°.**Methiodide:** Mp 220°.Gardmer, J. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1902, **240**, 81 (isol)Späth, E. et al., *Ber.*, 1927, **60**, 1892 (struct)Kametani, T. et al., *Phytochemistry*, 1971, **10**, 1881 (isol)Nonaka, G. et al., *Yakugaku Zasshi*, 1973, **93**, 87 (isol)Takao, N. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2859 (biosynth)Iwasa, K. et al., *J.O.C.*, 1981, **46**, 4744 (abs config)Kamiguchi, M. et al., *Helv. Chim. Acta*, 1987, **70**, 1482 (cryst struct, pmr)Kamiguchi, M. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1992, **325**, 585 (cd, abs config, resoln)**Corycavidine**

C-671

5,7,8,15-Tetrahydro-3,4-dimethoxy-6,15-dimethylbenzo[e][1,3]dioxolo[4,5-k][3]benzazecin-14(6H)-one, 9CI

As Corycavamine, C-670 with


R = R' = Me

C₂₂H₂₅NO₅ 383.443***(+)*-form** [521-93-7]Alkaloid from the root nodules of *Corydalis tuberosa* and the tubers of *Corydalis cava* (Papaveraceae). Prisms (MeOH). Mp 213-214°. [α]_D²⁵ +200 (c, 0.24 in CHCl₃).***(±)*-form** [6018-34-4]Alkaloid from the tubers of *Corydalis**cava* (Papaveraceae). Cryst. (MeOH or CHCl₃/Et₂O). Mp 192-194°.**Methiodide:**Cryst. (Me₂CO). Mp 198°.v. Bruchhausen, F. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1925, **263**, 570 (struct)Govindachari, T.R. et al., *J. Sci. Ind. Res., Sect. B*, 1957, **16**, 506 (synth, uv)Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261 (isol, uv, ir, ms)**Corycidine**

C-672

C₁₁H₁₁NO₄ 221.212Struct. unknown. Alkaloid from *Corydalis stewartii* (Papaveraceae). Mp 290-291° dec.Ikram, M. et al., *Pak. J. Sci. Ind. Res.*, 1966, **9**, 34-37; *CA*, **67**, 117031k**Corydaine**

C-673

7,8-Dihydro-8'-hydroxy-6-methylspiro[1,3-dioxolo[4,5-g]isoquinoline-5(6H),7'-[7H]indeno[4,5-d][1,3]dioxol]-6'(8'H)-one, 9CIC₂₀H₁₇NO₆ 367.357***(+)*-form** [31456-80-1]Alkaloid from *Corydalis paczkoskii* and *Corydalis vaginans* (Papaveraceae). Cryst. (EtOH or Et₂O). Mp 184° Mp 189-189.5°. [α]_D²² +145 (c, 1.3 in CHCl₃). λ_{max} 236 (log ε 4.49); 290 (log ε 4.04); 314 (log ε 4.03) (EtOH).**8-Epimer: Sibiricine**

[24181-66-6]

C₂₀H₁₇NO₆ 367.357Alkaloid from *Corydalis sibirica*, *Corydalis ledebouriana* and *Corydalis paniculigera* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 225°. Opt. rotn. not recorded. λ_{max} 205 (log ε 4.8); 240 (log ε 3.94); 291 (log ε 3.91); 313 (sh) (log ε 3.99) (no solvent reported).***(±)*-form** [64397-09-7]

Synthetic. Cryst. (MeOH). Mp 127-129°.


8-Epimer: [64397-10-0]

Synthetic. Cryst. (MeOH). Mp 223-225°.

Manske, R.H.F. et al., *Can. J. Chem.*, 1969, **47**, 3585-3588 (*Sibiricine*, isol, ir, uv, pmr, struct)Baisheva, Kh.Sh. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 465-467 (isol, uv, ir, pmr, ms)Yu, C.K. et al., *Can. J. Chem.*, 1971, **49**, 3025-3037 (*Sibiricine*, ms)Fesenko, D.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 157-159 (struct)Margvelashvili, N.N. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 118-120 (isol, uv, ir, pmr, ms)Nalliah, B.C. et al., *Can. J. Chem.*, 1977, **55**, 922-924; 1979, **57**, 1545-1549 (*Sibiricine*, *Corydaine*, synth)

- McLean, S. *et al.*, *Can. J. Chem.*, 1977, **55**, 924-925 (*synth*)
 Hughes, D.W. *et al.*, *Can. J. Chem.*, 1977, **55**, 3304-3311 (*Sibiricine, Corydaine, cmr*)
 Nasirov, S.M. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 640-645 (*Sibiricine, cryst struct*)
 Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 2621-2623 (*Sibiricine, Corydaine, synth, ir, pmr*)
 Kessar, S.V. *et al.*, *Chem. Comm.*, 1994, 1327-1328 (*Sibiricine, Corydaine, synth*)

Corydalic acid C-674
 [50802-23-8]



C₂₁H₂₁NO₆ 383.4

(+)-form


- Liq. [α]_D +24.3 (c, 2.28 in CHCl₃).
Me ester: Corydalic acid methyl ester
 C₂₂H₂₃NO₆ 397.427
 Alkaloid from *Corydalis incisa* (Papaveraceae). Mp 140-141°. [α]_D +85.4 (c, 0.68 in CHCl₃).

(±)-form

- Me ester:* [88610-31-5]
 Synthetic. Cryst. (CHCl₃/hexane). Mp 144-147°.
 Nonaka, G. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 1020 (*ir, uv, ms, pmr, isol, struct*)
 Yagi, A. *et al.*, *Phytochemistry*, 1977, **16**, 1197 (*biosynth*)
 Iwasa, K. *et al.*, *Tet. Lett.*, 1981, 2333 (*abs config*)
 Cushman, M. *et al.*, *J.O.C.*, 1984, **49**, 1278 (*synth, pmr*)
 Clark, R.D. *et al.*, *J.O.C.*, 1989, **54**, 1174 (*synth*)
 Kamiguchi, M. *et al.*, *J.C.S. Perkin 2*, 1997, 631-636 (*pmr, cryst struct*)

Corydaline C-675

5,8,13,13a-Tetrahydro-2,3,9,10-tetra-methoxy-13-methyl-6H-dibenzo[a,g]quinolizine, 9CI. 2,3,9,10-Tetramethoxy-13-methylberbine, 8CI. *Corydalis A* [518-69-4]
 [6018-35-5 ((±)-form)]



C₂₂H₂₇NO₄ 369.46

Alkaloid from *Corydalis tuberosa* and many other *Corydalis* spp. (Papaveraceae). Shows analgesic and antirheumatic props. Prisms (EtOH). Mp 135°. [α]_D²⁵ +235 (EtOH).

- LD₅₀ (mus, ivn) 136 mg/kg.
 O²-De-Me: **Isocorybulbine**

[22672-74-8]
 [27313-92-4]
 C₂₁H₂₅NO₄ 355.433
 Alkaloid from *Corydalis cava* (Papaveraceae). Cryst. (EtOH), Mp 179-180°. [α]_D²⁰ +299.8 (c, 1.04 in CHCl₃).

O³-De-Me: **Corybulbine**. *Corydalis G* [518-77-4]
 C₂₁H₂₅NO₄ 355.433
 Alkaloid from the root nodules of *Corydalis tuberosa* (*cava*) and from a few other *Corydalis* spp. (Papaveraceae). Mp 242°. [α]_D +303.3 (CHCl₃). Has been confused with Corydalmine, C-677.

O³-De-Me, hydrochloride: Mp 245-250°.


O¹⁰-De-Me: **Yuanhunine**
 [104387-15-7]
 C₂₁H₂₅NO₄ 355.433
 Alkaloid from tubers of yuanhu (*Corydalis turtchaninovi* f. *yanhusuo*) (Papaveraceae). Cryst. Mp 166-168°. [α]_D²³ +229.7 (c, 0.16 in EtOH).

O³,O¹⁰-Di-de-Me: **Corydalidzine**
 [50656-80-9]
 [59555-26-5 ((±)-form)]
 C₂₀H₂₃NO₄ 341.406
 Alkaloid from *Corydalis koidzumiana* (Papaveraceae). Mp 209-210° (*in vacuo*). [α]_D²³ +333 (MeOH).

- Späth, E. *et al.*, *Ber.*, 1925, **58**, 1274 (*Corybulbine, Isocorybulbine, struct*)
 Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1956, **34**, 1 (*isol*)
 Jeffs, P.W. *et al.*, *Experientia*, 1965, **21**, 690 (*Isocorybulbine, abs config*)
 Nakuto, S. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 1017 (*Isocorybulbine, synth*)
 Tani, C. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 313 (*Corydalidzine*)
 Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (*cmr*)
 Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 3185 (*Corybulbine, config, ir, pmr*)
 Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 1151 (*synth, pmr*)
 Holland, H.L. *et al.*, *Can. J. Chem.*, 1979, **56**, 1588-1597 (*biosynth*)
 Iwasa, K. *et al.*, *J.O.C.*, 1981, **46**, 4744 (*Isocorybulbine, Corybulbine, Corydalidzine, synth, abs config*)
 Fu, X. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 447-453 (*Yuanhunine*)
 Ribár, B. *et al.*, *Acta Cryst. C*, 1992, **48**, 1864 (*cryst struct*)
 Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 399-404 (*synth*)
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr*)

Corydalisol C-676

5-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)methyl]-1,3-benzodioxole-4-methanol, 9CI. 1,2,3,4-Tetrahydro-1-(2-hydroxymethyl-3,4-methylenedioxybenzyl)-2-methyl-6,7-methylenedioxyisoquinoline



(R)-form

C₂₀H₂₁NO₅ 355.39
 Protoberberine-type numbering shown.

(R)-form [55934-47-9]
 Alkaloid from *Corydalis incisa* (Papaveraceae). Prisms (MeOH). Mp 160-161°. [α]_D²¹ +21.4 (c, 0.28 in CHCl₃).

8-Aldehyde: **Aobamine**
 [59614-37-8]
 C₂₀H₁₉NO₅ 353.374
 Alkaloid from *Corydalis ochotensis* var. *raddeana* (Papaveraceae). Syrup. Unstable. No opt. rotn. reported, abs. config. not detd.; may correspond to (S)-Corydalisol.

8-Carboxylic acid: **Coryximine**
 [127460-61-1]
 C₂₀H₁₉NO₆ 369.373
 Alkaloid from bulbs of *Corydalis hsuowensis* (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 197-198°. [α]_D²³ +147.7 (c, 0.5 in CHCl₃).

(S)-form [86688-04-2]
 Alkaloid from *Hypecoum procumbens* (Hypecoaceae). [α]_D²³ -18 (c, 0.11 in MeOH).


(±)-form [61665-07-4]
 Synthetic. Mp 127-128° Mp 147-148°.

8-Aldehyde: [64598-67-0]
 Synthetic. Cryst. (MeOH). Mp 168-168.5°.

- Nonaka, G. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 294 (*uv, ir, pmr, ms, struct*)
 Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 390 (*isol, ir, pmr, ms, struct, Aobamine*)
 Shamma, M. *et al.*, *Tetrahedron*, 1978, **34**, 635 (*synth, ir, pmr, Corydalisol, Aobamine*)
 Gözler, T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 414 (*isol, uv, ir, ms, cd, struct*)
 Gözler, B. *et al.*, *J.C.S. Perkin 1*, 1983, 2431 (*synth, uv, ir, pmr, ms*)
 Iwasa, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 998 (*synth, ir, pmr, cmr*)
 Rozwadowska, M.D. *et al.*, *Tetrahedron*, 1988, **44**, 1221 (*synth, uv, ir, pmr, ms, Corydalisol, Aobamine*)
 Zhou, J. *et al.*, *Planta Med.*, 1991, **57**, 156 (*Coryximine*)
 Chrzanoska, M. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1998, **72**, 2399-2404 (*synth*)

Corydalmine C-677

5,8,13,13a-Tetrahydro-2,3,9-trimethoxy-6H-dibenzo[a,g]quinolizin-10-ol, 9CI. *Kikemanine. Cycemanine. Schefferine*



C₂₀H₂₃NO₄ 341.406

Kikemanine and *Schefferine* were until 1977 thought to have isomeric structs.

(S)-form [30413-84-4]
 Alkaloid from *Corydalis pallida* and

Stephania glabra (Papaveraceae, Menispermaceae). Plates (CHCl₃/MeOH). Mp 177-178° (174-175°). [α]_D²² -303 (c, 1.14 in EtOH).

α-N-Oxide: Corydalmine α-N-oxide

[250686-94-3]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Corydalis tashiroi*.

Prisms (CHCl₃/MeOH). Mp 205-207°. [α]_D²⁶ -28.2 (c, 0.31 in CHCl₃). λ_{max} 206 (log ε 4.71); 227 (sh) (log ε 4.1); 284 (log ε 3.71) (EtOH).

β-N-Oxide: Corydalmine β-N-oxide

[250686-95-4]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Corydalis tashiroi*.

Prisms (CHCl₃/MeOH). Mp 176-178°. [α]_D²⁴ -75.3 (c, 0.22 in CHCl₃). λ_{max} 209 (log ε 4.58); 228 (sh) (log ε 4.17); 284 (log ε 3.74) (EtOH).

N-Me: N-Methylcorydalmine

[81010-29-9]

C₂₁H₂₆NO₄[⊕] 356.441

Quaternary alkaloid from *Stephania elegans* (Menispermaceae). Mp 190° (as chloride).

Me ether: see Tetrahydropalmatine, T-212

[95462-84-3]

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2785-2789 (*isol*)

Kametani, T. *et al.*, *J.C.S. (C)*, 1970, 1060-1064 (*Kikemanine, isol, pmr, cd*)

Brochmann-Hanssen, E. *et al.*, *J.O.C.*, 1977, **42**, 3588-3591 (*Schefferine, struct, bibl*)

Singh, R.S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 664-667 (*N-Methylcorydalmine*)

Ohiri, F.C. *et al.*, *Planta Med.*, 1983, **49**, 162-164 (*pmr*)

Chen, J.-J. *et al.*, *Planta Med.*, 1999, **65**, 643-647 (*N-oxides*)


Bianchi, D.A. *et al.*, *Can. J. Chem.*, 2000, **78**, 1165-1169 (*synth*)

Blanchfield, J.T. *et al.*, *Phytochemistry*, 2003, **63**, 711-720 (*isol, pmr, cmr*)

Corydamine

C-678

[49870-84-0]



C₂₀H₁₈N₂O₄ 350.373

Alkaloid from *Corydalis incisa* (Papaveraceae). Mp 255-257° (235-239°) dec. (as hydrochloride). May form a mono- and a dihydrochloride.

N-Formyl: N-Formylcorydamine

[50506-34-8]

C₂₁H₁₈N₂O₅ 378.384

Alkaloid from *Corydalis incisa* (Papaveraceae). Mp 159.5-160.5°.

Nonaka, G. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 1410 (*ir, uv, pmr, ms, struct*)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 1587 (*ir, uv, pmr, ms, struct, isol, synth*)


Yagi, A. *et al.*, *Phytochemistry*, 1977, **16**, 1197 (*biosynth*)

Seger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**, 882-886 (*pmr, cmr*)

Corydecumbine

C-679

[154722-69-7]



C₂₀H₁₉NO₆ 369.373


Alkaloid from tubers of *Corydalis decumbens* (Papaveraceae). Creamy-white cryst. Mp 214°. [α]_D -142.3 (c, 0.62 in CHCl₃).

Basnet, P. *et al.*, *Heterocycles*, 1993, **36**, 2205 (*isol, pmr, cmr, struct*)

Corydendramine A

C-680

6-(1,3,5,9-Dodecatetraenyl)-2-methyl-3-piperidinol [291764-44-8]



C₁₈H₂₉NO 275.433

Defence compd. *isol.* from the marine hydroid *Corydendrium parasiticum*. Pale yellow oil. [α]_D -24.3 (c, 0.17 in MeOH). λ_{max} 258 (ε 24200); 268 (ε 32700); 278 (ε 25600) (MeOH).

7',8'-Didehydro(E-), 5',6'-dihydro: 6-(1,3,7,9-Dodecatetraenyl)-2-methyl-3-piperidinol. **Corydendramine B**

[291764-45-9]

C₁₈H₂₉NO 275.433

Defence compd. *isol.* from *Corydendrium parasiticum*. Amorph. powder. [α]_D +83.7 (c, 0.08 in MeOH). λ_{max} 236 (ε 40400) (MeOH).

Lindquist, N. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1290-1291 (*Corydendramines A, B*)

Corydicine

C-681

C₁₉H₁₇NO₅ 339.347

Struct. unknown. Alkaloid from *Corydalis stewartii* (Papaveraceae). Mp 181-184°.

Hydrochloride: Mp 244-245° dec.

Picrate: Mp 268-270° dec.


Chloroplatinate: Mp 212-213° dec.

Ikram, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1966, **9**, 34-37; *C.A.*, **67**, 117031k

Corydine

C-682

1-Hydroxy-2,10,11-trimethoxyaporphine. *Glaucentrine*



C₂₀H₂₃NO₄ 341.406

Log P 2.69 (uncertain value) (calc).

(S)-form [476-69-7]

Alkaloid from *Corydalis tuberosa* and *Dicentra eximia* (Papaveraceae), also from the Annonaceae, Lauraceae, Liliaceae, Menispermaceae, Ranunculaceae, Rutaceae, Berberidaceae, Papaveraceae and Monimiaceae. Shows irritant and respiratory stimulant and CNS depressant activities plus some antineoplastic activity. Adrenolytic and weak cholinergic agent. Shows antimicrobial activity. Mp 149°. [α]_D²⁵ +204 (c, 0.6 in EtOH). Pharmacol. active isomer.

▶ CE1057000

N-Oxide: Corydine N-oxide

[74804-29-8]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Glaucium fimbriigerum* (Papaveraceae).

N-Me: N-Methylcorydine

[4668-04-6]

C₂₁H₂₆NO₄[⊕] 356.441

Alkaloid from *Zanthoxylum nigrescens* (Rutaceae). Mp 190-200° dec. (as iodide). [α]_D²⁸ +154 (c, 0.20 in 50% EtOH).

N-De-Me: 1-Hydroxy-2,10,11-trimethoxyaporphine. Norcorydine

[26931-78-2]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Annona squamosa* (sugar apple), *Popowia* sp. cf. *cyanocarpa*, *Stephania dinklagei*, *Laurelia philippiana*, *Glaucium fimbriigerum* and *Xylopiopsis danguyella* (Annonaceae, Menispermaceae, Monimiaceae, Papaveraceae). Shows antiplasmodial and antiamebic activities. Noncryst. Mp 249-251° (as hydrochloride). [α]_D +156 (MeOH). [α]_D²³ +190 (c, 0.2 in CHCl₃).

N-De-Me, N,O-di-Ac:

Amorph. [α]_D¹⁶ +266 (c, 0.76 in CHCl₃).

6a,7-Didehydro: Dehydrocorydine

[74799-12-5]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Glaucium fimbriigerum* (Papaveraceae). Amorph.

4R-Hydroxy: Glaufidine. Glaufiridine

[71609-80-8]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Glaucium fimbriigerum* (Papaveraceae). [α]_D +182 (c, 0.4 in MeOH).

4S-Hydroxy: Epiglaufidine

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Glaucium fimbriigerum* (Papaveraceae). $[\alpha]_D^{25} +198$ (c, 0.5 in MeOH).

(±)-form

N-De-Me: [60661-35-0]

Alkaloid from the leaves of *Xylopia pancheri* (Annonaceae).

Go, J. *et al.*, *Yakugaku Zasshi*, 1929, **49**, 801 (isol)

Shamma, M. *et al.*, *Tet. Lett.*, 1965, 1509 (synth, uv, struct)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 363 (Norcorydine)

Bakuni, D.S. *et al.*, *Phytochemistry*, 1972, **11**, 1819 (isol, uv, ms)

Shamma, M. *et al.*, *Phytochemistry*, 1973, **12**, 1505 (isol, pmr, uv, ms)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 6 (Norcorydine)

Nieto, M. *et al.*, *Planta Med.*, 1976, **30**, 48 (± Norcorydine)

Israelov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 104 (Glaufidine)

Castedo, L. *et al.*, *Heterocycles*, 1980, **14**, 1135 (Dehydrocorydine)

Karimova, S.U. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 224; 1983, **18**, 493; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 177 (Isocorydine N-oxide, Dehydrocorydine, Glaufidine, Epiglaufidine)


Hoquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551 (Norcorydine, occur)

Urzúa, A. *et al.*, *Phytochemistry*, 1982, **21**, 773 (Norcorydine, isol, ir, pmr, ms)

Villar, A. *et al.*, *Farm. Tijdschr. Belg.*, 1984, **61**, 300 (activity)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1985, **48**, 646 (Glaufidine, Epiglaufidine, config)

Wright, C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1638-1640 (activity)

Corydinemethine**C-683**


$C_{21}H_{25}NO_4$ 355.433

Alkaloid from *Berberis cretica* (whole plant) (Berberidaceae). Amorph.

Ross, S.A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 159 (isol, uv, ir, pmr, ms, struct)

Corydione**C-684**

1,2-Dimethoxy-6-methyl-4H-benzo[1,3]benzodioxolo[5,6-g]quinoline-4,5(6H)-dione, 9CI. **4,5-Dioxodehydro-*nantenine***. Alkaloid Cm-4 [72007-91-1]



$C_{20}H_{15}NO_6$ 365.342

Alkaloid from *Corydalis bulbosa* and *Corydalis marshalliana*, and from the dried fruits of *Nandina domestica* (Papaveraceae, Nandiniaceae). Red needles (EtOH or EtOH/Et₂O). Mp 273-275°.


Kunitomo, J. *et al.*, *Shoyakugaku Zasshi*, 1979, **33**, 84; *CA*, **91**, 207404j (isol, struct, synth, uv, ms)

Kiryakov, H.G. *et al.*, *Planta Med.*, 1980, **39**, 210; 1981, **41**, 298; **43**, 51 (isol, uv, ir, pmr, ms)

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1980, **100**, 337; *CA*, **93**, 95454h (struct)

Corygovanine**C-685**

5,8,14,14a-Tetrahydro-3-methoxy-6H-benzo[a][1,3]benzodioxolo[5,6-g]quinolin-2-ol, 9CI. **Pseudocheilanthifoline** [70031-37-7]

**(S)-form**

$C_{19}H_{19}NO_4$ 325.363

(S)-form [63729-95-3]

Alkaloid from the leaves and stems of *Corydalis govaniana* (Papaveraceae). Cryst. (MeOH). Mp 134-136°. $[\alpha]_D^{25} -368.2$. λ_{max} 212; 232 (sh); 287 (MeOH). λ_{max} 215; 288; 294 (sh) (MeOH/NaOH).

(±)-form


Synthetic. Mp 138-140°.

Mehra, K. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 844-888 (isol, uv, ir, pmr, ms, struct, synth)

Moullis, C. *et al.*, *Org. Magn. Reson.*, 1978, **11**, 398-400 (cmr)

Corymotine**C-686**

5,8,13,13a-Tetrahydro-2,3,9,10-tetramethoxy-13,13-dimethyl-6H-dibenzo[a,g]quinolizine, 9CI. **Corybrachylobine**



$C_{23}H_{29}NO_4$ 383.486

(S)-form [115569-75-0]


Alkaloid from the tubers of *Corydalis remota* and *Corydalissolida* ssp. *brachyloba* (Papaveraceae). Prisms (MeOH). Mp 148-149°. $[\alpha]_D^{25} -188$ (c, 0.087 in CHCl₃).

Fu, X. *et al.*, *J. Nat. Prod.*, 1988, **51**, 262 (isol, uv, ir, pmr, cmr, ms, struct)

Sener, B. *et al.*, *J. Chem. Soc. Pak.*, 1991, **13**, 63; *CA*, **116**, 3548j

Corynantheidinaline**C-687**

[132922-57-7]



$C_{21}H_{22}N_2O_3$ 350.416

MF and struct. revised in 2001. Alkaloid from the v. young leaves of *Mitragyna speciosa* (Rubiaceae). Dark orange cryst. (EtOH). Mp 197-199° dec.

9-Methoxy: Mitragynaline

[132943-52-3]

$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae). Dark orange cryst. (EtOH). Mp 210-221° dec. $[\alpha]_D^{24} -4.3$ (c, 0.34 in CHCl₃). Struct. revised in 2001. λ_{max} 224 (sh) (log ϵ 4.39); 264 (log ϵ 4.3); 348 (log ϵ 3.97); 456 (sh) (log ϵ 4.62); 486 (log ϵ 4.83) (MeOH).

Houghton, P.J. *et al.*, *Phytochemistry*, 1991, **30**, 347-350 (isol, uv, ir, pmr, cmr, ms, struct)


Takayama, H. *et al.*, *Tet. Lett.*, 2001, **42**, 1741-1743 (pmr, cmr, struct)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 916-928 (rev)

Corynantheidinalinic acid**C-688**

15,16-Didehydro-14,16-diformylcorynan-17-oic acid, 9CI

[133019-99-5]



$C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from the young leaves of *Mitragyna speciosa* (Rubiaceae). Dark yellow cryst. (EtOH). Mp 213-214° dec.

9-Methoxy: Mitragynalinic acid

[132943-53-4]


$C_{22}H_{24}N_2O_5$ 396.442

Alkaloid from the young leaves of *Mitragyna speciosa* (Rubiaceae). Dark yellow cryst. (EtOH). Mp 221-224° dec.

Houghton, P.J. *et al.*, *Phytochemistry*, 1991, **30**, 347 (isol, ir, ms, pmr, cmr, struct)

Corynantheine**C-689**

[18904-54-6]




Absolute Configuration

- $C_{22}H_{26}N_2O_3$ 366.459
Alkaloid from the bark of *Pseudocinchona africana* (preferred genus name *Corynanthe*) and from *Mitragyna parvifolia*, *Rauwolfia canescens* and *Uncaria rhynchophylla*. Sympatholytic agent. Cryst. (EtOH aq.). Mp 164°. $[\alpha]_D^{20} +28.8$ (c, 1 in MeOH). Occurs together with Dihydrocorynantheine which caused confusion in the early lit. First obt. pure in 1953.
- LD₅₀ (mus, ivn) 35 mg/kg. VS3675000
Hydrochloride: Mp 167-188° (synthetic). $[\alpha]_D^{20} +37$ (c, 0.14 in MeOH) (synthetic).
- O¹⁷-De-Me: **Demethylcorynantheine**. Alkaloid RMB 1
[18786-21-5]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from the stem bark of *Rauwolfia mombasiana* (Apocynaceae).
- 18,19-Dihydro: **Dihydrocorynantheine**
[50439-68-4]
[4684-43-9]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Corynanthe yohimbe*, *Pseudocinchona africana*, *Cephalanthus occidentalis*, *Uncaria rhynchophylla*, *Uncaria callophylla* and *Uncaria gambier* (Apocynaceae, Rubiaceae). Cryst. (MeOH aq. or Et₂O). Mp 105°. $[\alpha]_D^{20} +36.2$ (c, 0.123 in MeOH). λ_{max} 227 ; 283 ; 291 (EtOH).
- GN1038000
18,19-Dihydro, N⁴-oxide: **Dihydrocorynantheine N-oxide**
[55137-72-9]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from *Uncaria tomentosa* (Rubiaceae). No phys. props. reported.
- 18,19-Dihydro, O¹⁷-de-Me: **Demethyl-dihydrocorynantheine**. Alkaloid RMB 7
[5027-70-3]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from the stem bark of *Rauwolfia mombasiana* (Apocynaceae). Yellow amorph. powder. λ_{max} 226 ; 275 (sh) ; 293 (MeOH).
- 10-Hydroxy, 18,19-dihydro: **Isogambirine**. 10-Hydroxydihydrocorynantheine
[142934-41-6]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from leaves of *Uncaria callophylla* (Rubiaceae). λ_{max} 226 ; 240 ; 279 (EtOH).
- 3-Epimer: **Hirsuteine**. Δ^{18} -Hirsutine
[35467-43-7]
 $C_{22}H_{26}N_2O_3$ 366.459
Alkaloid from *Mitragyna parvifolia*, *Mitragyna hirsuta*, and from the genus *Uncaria* (Rubiaceae). Mp 92-94°.
- GN1039000
3-Epimer, 18,19-dihydro: **Hirsutine**[†]
[7729-23-9]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Mitragyna hirsuta*, *Mitragyna parvifolia*, *Mitragyna stipulosa*, *Cephalanthus occidentalis*, and several *Uncaria* spp., e.g. *Uncaria rhynchophylla* (Rubiaceae). Ganglionic blocking agent showing long-lasting neuro blocking. Cryst. (Et₂O). Mp 101°. $[\alpha]_D^{23} +68.6$ (c, 0.32 in CHCl₃). λ_{max} 226 (log
- € 4.64); 282 (log € 3.84); 290 (log € 3.79) (EtOH).
- VS3660000
3-Epimer, 18,19-dihydro, N⁴-oxide: **Hirsutine N-oxide**
[55176-58-4]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from *Uncaria tomentosa* (Rubiaceae). No phys. props. reported.
- 20-Epimer, 18,19-dihydro: **Corynantheidine**[†]
[23407-35-4]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Rauwolfia canescens*, *Pseudocinchona africana*, *Mitragyna speciosa* and several *Uncaria* spp. Antihypertensive, sympatholytic agent. Mp 117° (125°). $[\alpha]_D -171$ (c, 1 in MeOH) (-142). λ_{max} 225 (log € 4.59); 280 (log € 3.93) (no solvent reported).
- 20-Epimer, 18,19-dihydro, hydrochloride: Mp 213° (hydrate). $[\alpha]_D^{15} -128$ (MeOH).
- 20-Epimer, 18,19-dihydro, O-de-Me: **Demethylcorynantheidine**. Alkaloid RMB 5
[71657-70-0]
 $C_{21}H_{26}N_2O_3$ 354.448
Trace alkaloid from the stem bark of *Rauwolfia mombasiana* (Apocynaceae). Yellow amorph. powder. Probable struct.; insufficient material for full characterisation. λ_{max} 227 ; 275 (sh) ; 283 ; 291 (MeOH).
- 20-Epimer, 9-methoxy, 18,19-dihydro: see Mitragynine, M-657
- 3,20-Diepimer: **Epiallocorynantheine**
[57820-49-2]
 $C_{22}H_{26}N_2O_3$ 366.459
Alkaloid from *Uncaria attenuata* ssp. *bulusanensis* (Rubiaceae). λ_{max} 225 ; 245 (sh) ; 284 ; 291 (EtOH).
- 3,20-Diepimer, 18,19-dihydro: **3-Isocorynantheidine**
[7729-22-8]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Mitragyna speciosa* and *Uncaria* spp. (Rubiaceae). Mp 221-222° (as perchlorate).
- (±)-form [25920-83-6]
Synthetic. Mp 225-226°.
Hydrochloride:
Synthetic. Mp 176-179°.
- 20-Epimer, 18,19-dihydro: Synthetic. Noncryst., cryst. (as perchlorate). Mp 241-242° (perchlorate).
- Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1949, 509-515 (synth)
Janot, M.-M. et al., *Helv. Chim. Acta*, 1951, **34**, 1207-1210 (struct)
Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1953, 1033-1038 (Corynantheidine, uv, ir)
Goutarel, R. et al., *Helv. Chim. Acta*, 1953, **43**, 337-340 (purifn)
Van Tamelen, E.E. et al., *J.A.C.S.*, 1957, **79**, 6426-6430; 1969, **91**, 7349-7359 (synth, config, Dihydrocorynantheine)
Bartlett, M.F. et al., *J.A.C.S.*, 1962, **84**, 622-630 (Corynantheidine, config)
Weisbach, J.A. et al., *Tet. Lett.*, 1965, 3457-3463 (Corynantheidine, synth)
Shellard, E.J. et al., *J. Pharm. Pharmacol.*, 1966, **18**, 553-555 (Hirsutine, uv, ir, pmr, struct)
- Merlini, L. et al., *Tet. Lett.*, 1967, 1571-1574 (Dihydrocorynantheine, isol)
Lee, C.M. et al., *Tetrahedron*, 1967, **23**, 375-385 (Dihydrocorynantheine, Corynantheidine, uv, cd, ord)
Trager, W.F. et al., *Tetrahedron*, 1967, **23**, 1043-1047 (Hirsutine, uv, ord, cd, abs config)
Autrey, R.L. et al., *J.A.C.S.*, 1968, **90**, 4917-4923 (synth, bibl)
Szántay, C. et al., *Chem. Ber.*, 1969, **102**, 3963-3973 (Corynantheidine, synth)
Beckett, A.H. et al., *Tetrahedron*, 1969, **25**, 5961-5969 (Corynantheidine, Hirsutine, Dihydrocorynantheine, ms)
Shellard, E.J. et al., *Planta Med.*, 1972, **21**, 382-392; 1978, **34**, 253-263 (Hirsuteine, 3-Isocorynantheidine)
Phillipson, J.D. et al., *Phytochemistry*, 1973, **12**, 1507; 1975, **14**, 1855-1863 (Hirsuteine, Epiallocorynantheine)
Hemingway, S.R. et al., *J. Pharm. Pharmacol., Suppl.*, 1974, **26**, 113P (Dihydrocorynantheine oxide, Hirsutine oxide)
Phillipson, J.D. et al., *J. Chromatogr.*, 1975, **105**, 163-178 (Hirsuteine, 3-Isocorynantheidine, tlc, glc, ms)
Bárczai-Beke, M. et al., *Tetrahedron*, 1976, **32**, 1153-1159 (Dihydrocorynantheine, synth, ir, bibl)
Fujii, T. et al., *Heterocycles*, 1977, **7**, 149-153 (Dihydrocorynantheine, synth)
Sakai, S. et al., *Chem. Pharm. Bull.*, 1978, **26**, 2596-2598 (Corynantheidine, synth, pmr)
Iwu, M.M. et al., *Planta Med.*, 1979, **36**, 208-212 (Demethylcorynantheine, Demethyl-dihydrocorynantheine, Demethylcorynantheidine)
Le Xuan, P. et al., *Chromatographia*, 1980, **13**, 693-697 (tlc)
Brown, R.T. et al., *Chem. Comm.*, 1984, 847-848 (Hirsutine, synth)
Goh, S.H. et al., *Phytochemistry*, 1985, **24**, 880-881 (Dihydrocorynantheine, isol, uv, ir, pmr, cmr, ms)
Szántay, C. et al., *Alkaloids (Academic Press)*, 1986, **27**, 131-268 (rev)
Naito, T. et al., *Heterocycles*, 1987, **26**, 1739-1742 (Hirsuteine, synth)
Kam, T.-S. et al., *Phytochemistry*, 1992, **31**, 2031-2034 (Isogambirine)
Lounasmaa, M. et al., *Heterocycles*, 1998, **49**, 445-450 (Hirsutine, 3-Isocorynantheidine, synth)
Tietze, L.F. et al., *Angew. Chem., Int. Ed.*, 1999, **38**, 2045-2047 (Hirsutine, Dihydrocorynantheine, synth)
Yu, S. et al., *J.A.C.S.*, 2000, **122**, 7827-7828 (Corynantheidine, synth)
Staerk, D. et al., *Planta Med.*, 2000, **66**, 531-536 (Corynantheidine, Corynantheine, Dihydrocorynantheine, isol, pmr, cmr)
Merck Index, 13th edn., 2001, No. 2572 (props, bibl)
Deiters, A. et al., *J.O.C.*, 2006, **71**, 6547-6561 (Hirsutine, synth)

Coryneine

C-690

3,4-Dihydroxy-N,N,N-trimethylbenzenethanaminium(1+), 9CI. (3,4-Dihydroxyphenethyl) trimethylammonium, 8CI. Dopamine methosalt [7224-66-0]



$C_{11}H_{18}NO_2^+$ 196.269

Alkaloid from *Stetsonia coryne*, *Desmodium triflorum*, *Aconitum carmichaeli*, *Fagara hyemalis* and *Alhagi pseudalhagi* (Cactaceae, Fabaceae, Ranunculaceae, Rutaceae). Sympathomimetic agent.

Chloride: [13075-92-8]
C₁₁H₁₈ClNO₂ 231.721
Mp 263° dec. (201°).

Iodide: [52336-53-5]
C₁₁H₁₈INO₂ 323.173
Mp 205°.


[55-61-8, 458-39-9]

Barger, G. *et al.*, *J.C.S.*, 1910, 2253 (*synth*)
Reti, L. *et al.*, *CA*, 1935, 29, 2961 (*isol*)
Buck, J.S. *et al.*, *J.A.C.S.*, 1938, 60, 1788 (*synth*)
Kuck, A.M. *et al.*, *Phytochemistry*, 1967, 6, 1541 (*isol*)
Agurell, S. *et al.*, *J. Nat. Prod.*, 1971, 34, 183 (*isol*)
Ghosal, S. *et al.*, *J. Pharm. Sci.*, 1973, 62, 1555 (*isol, uv, pmr*)
Barlow, R.B. *et al.*, *Br. J. Pharmacol.*, 1976, 57, 517; 1980, 69, 597 (*Pharmacol*)
Konno, C. *et al.*, *Planta Med.*, 1979, 35, 150 (*isol*)

Corynolamine

C-691

[67998-93-0]



C₂₂H₂₃NO₆ 397.427

(±)-form

Alkaloid from *Corydalis incisa* (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 198-199°.

Di-Ac:


Needles (MeOH/Me₂CO). Mp 191-192°.

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1978, 26, 1880; 1979, 27, 2194 (*isol, uv, ir, pmr, cmr, ms, struct*)

Corynoline

C-692

5b,6,7,12b,13,14-Hexahydro-5b,13-dimethyl[1,3]benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridin-6-ol, 9CI



C₂₁H₂₁NO₅ 367.401

(+)-form [18797-79-0]

Alkaloid from *Corydalis incisa* and *Corydalis solida* ssp *tauricola*. Cell adhesion inhibitor. Shows antifungal activity. Cryst. (Me₂CO/Et₂O or MeOH/Et₂O). Mp 180-181° (178-179°). [α]_D²⁵ +132 (c, 2.64 in CHCl₃) (+116.4).

O-Sulfate: Corynoline O-sulfate

[74045-63-9]
C₂₁H₂₁NO₈S 447.465
Alkaloid from *Corydalis incisa* (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 253-254°. [α]_D²³ +67 (c, 0.19 in MeOH).

5α-Hydroxy: 5-Hydroxycorynoline

C₂₁H₂₁NO₆ 383.4
Alkaloid from *Corydalis bulleyana* whole plants (Papaveraceae). Plates (MeOH). Mp 149-150°. [α]_D¹⁸ +98.6 (EtOH).

5α-Hydroxy, O-formyl: 5-Formyloxycorynoline

C₂₂H₂₁NO₇ 411.41
Alkaloid from *Corydalis bulleyana* whole plants (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 209-211°. [α]_D²⁰ +15 (MeOH).

8-Oxo: 8-Oxocorynoline

[55739-71-4]
C₂₁H₁₉NO₆ 381.384
Minor alkaloid from *Corydalis incisa* (Papaveraceae). Mp 295°.

8-Oxo, Ac: O-Acetyl-8-oxocorynoline. 8-Oxoacetylcorynoline

[104513-81-7]
C₂₃H₂₁NO₇ 423.421
Alkaloid from whole plants of *Corydalis bulleyana*. Prisms (CHCl₃/MeOH). Mp 156-158°. [α]_D²⁰ +92.1 (MeOH). λ_{max} 207 (log ε 4.51); 285 (log ε 3.75); 319 (log ε 3.56) (EtOH).

8,9-Didehydro, N-de-Me, O-Ac: Coryncine

[255835-77-9]
C₂₂H₁₉NO₆ 393.395
Alkaloid from *Corydalis incisa*. Pale yellow powder. Mp 210-212°. [α]_D²¹ +186.7 (c, 0.3 in CHCl₃). λ_{max} 262 (log ε 4.18); 269 (log ε 4.17); 294 (log ε 4.07) (CHCl₃).

14-Epimer: Isocorynoline. 14-Epicorynoline

[51151-82-7]
C₂₁H₂₁NO₅ 367.401
Alkaloid from *Corydalis incisa* and from *Corydalis bungeana* (Papaveraceae). Needles (MeOH); prisms (CHCl₃/MeOH). Mp 234-235° (231-232°). [α]_D²⁵ +125 (c, 0.12 in MeOH). [α]_D²⁵ +136 (c, 0.26 in CHCl₃). Shown in one paper as the enantiomer (6-Epiisocorynoline) but the phys. props of the two isolates are in good agreement.

14-Epimer, Ac: Acetylisocorynoline

[42881-67-4]
C₂₃H₂₃NO₆ 409.438
Alkaloid from *Corydalis incisa*. Needles (MeOH). Mp 210-211° (204-206°). [α]_D²⁸ +62.8 (c, 0.33 in CHCl₃).

(-)-form [74163-86-3]

Alkaloid from *Corydalis taliensis*. Also obt. by resoln. of (±)-Corynoline (Papaveraceae). Mp 178-179°. [α]_D²⁰ -116.4 (c, 2.5 in CHCl₃).

(±)-form [68035-45-0]

Alkaloid from *Corydalis incisa* and *Corydalis bungeana* (Papaveraceae). Cryst. (MeOH or Et₂O/MeOH). Mp 218-220°

(216-217°).

Ac: Acetylcorynoline

C₂₃H₂₃NO₆ 409.438
Alkaloid from *Corydalis incisa* (Papaveraceae). Shows antifungal activity. Plates (MeOH). Mp 159-160° (153-155°).

5α-Hydroxy: (±)-5-Hydroxycorynoline

[55739-69-0]
Minor alkaloid from *Corydalis incisa* (Papaveraceae). Prisms (CHCl₃/MeOH). Mp 245-246.5° (238-240°).

8-Oxo: [82950-95-6]

Synthetic. Mp 310-312°.

6-Epimer: 6-Epicorynoline

[51095-71-7]
C₂₁H₂₁NO₅ 367.401
Minor alkaloid from *Corydalis incisa* (Papaveraceae). Cryst. (MeOH or CHCl₃/MeOH). Mp 195-196.5° (182.5-183°).

13-Epimer: 6-Epiisocorynoline

Synthetic. Mp 185-186° Mp 238-240°.

14-Epimer: [83607-67-4]

Synthetic. Cryst. (MeOH). Mp 174-176° (204-205°).

14-Epimer, Ac: Synthetic. Mp 178-180°.

Tani, C. *et al.*, *Yakugaku Zasshi*, 1962, 82, 594 (*Corynoline, Isocorynoline, isol*)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1963, 11, 1306; 1973, 21, 1096; 1976, 24, 2859; 1978, 26, 1880 (*Corynoline, Isocorynoline, isol, uv, ir, pmr, cmr, struct*)

Kametani, T. *et al.*, *Phytochemistry*, 1971, 10, 1881 (*Acetylcorynoline, Isocorynoline*)

Nonaka, G. *et al.*, *Yakugaku Zasshi*, 1973, 93, 87 (*Acetylcorynoline, Acetylisocorynoline*)

Nonaka, G. *et al.*, *Chem. Pharm. Bull.*, 1975, 23, 521 (*8-Oxocorynoline, 5-Hydroxycorynoline, 6-Epicorynoline*)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1976, 24, 2859; 1978, 26, 1880 (*isol, pmr, cmr, biosynth*)

Iwasa, K. *et al.*, *Phytochemistry*, 1979, 18, 1725 (*sulfate*)

Takao, N. *et al.*, *Tetrahedron*, 1979, 35, 1099; 1977 (*isol, resoln, abs config, cryst struct, ms*)

Ninomiya, I. *et al.*, *J.C.S. Perkin 1*, 1980, 212 (*synth*)

Cushman, M. *et al.*, *J.A.C.S.*, 1983, 105, 2873 (*Corynoline, Isocorynoline, 8-Oxocorynoline, synth, ir, pmr, ms*)

Hong, H. *et al.*, *Planta Med.*, 1986, 193 (*O-Acetyl-8-oxocorynoline, 5-Hydroxycorynoline, 5-Formyloxycorynoline*)

Zeng, W. *et al.*, *Phytochemistry*, 1988, 27, 599 (*13-Epicorynoline*)

Hanaoka, M. *et al.*, *Tet. Lett.*, 1988, 29, 6621 (*synth, pmr*)

Sener, B. *et al.*, *Planta Med.*, 1990, 56, 510 (*Tauricoline*)


Ma, W.G. *et al.*, *Fitoterapia*, 1999, 70, 258-265 (*activity, Coryncine*)

Kamigauchi, M. *et al.*, *Bioorg. Med. Chem.*, 2005, 13, 1867-1872 (*cryst struct, sar*)

Corynoloxine

C-693

11-Deoxy-6,11-epoxy-13-methylchelidonium, 9CI
[31470-65-2]



Absolute Configuration

C₂₁H₁₉NO₅ 365.385Alkaloid from *Corydalis incisa* (Papaveraceae). Needles (CHCl₃/MeOH). Mp 209-210°.

Picrate: Mp 188°.

12R-Hydroxy: **12R-Hydroxycorynoloxine**

[55739-73-6]

[55781-77-6]


C₂₁H₁₉NO₆ 381.384Alkaloid from *Corydalis incisa*. Mp 216-218°. Racemate.

[152323-60-9]

Tani, C. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 594; *CA*, **57**, 4758f (*isol*)Naruto, S. *et al.*, *Tet. Lett.*, 1968, 1705 (*pmr*)Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 247 (*pmr, struct, synth*)Kametani, T. *et al.*, *Phytochemistry*, 1971, **10**, 1881 (*isol, ms*)Kim, D.K. *et al.*, *Arch. Pharmacol. Res.*, 2000, **23**, 589-591 (*Corynoloxine, 12-Hydroxycorynoloxine*)**Corypalmine****C-694**

Tetrahydroajatrorrhizine. Discretinine

[13063-54-2]



(R)-form

C₂₀H₂₃NO₄ 341.406

Shows strong papaverine-like activity.

(R)-formAlkaloid from *Corydalis cava* (Papaveraceae). Mp 235-236°. [α]_D¹⁶ +280 (CHCl₃).N-Me: **N-Methylcorypalmine**

[169218-54-6]

C₂₁H₂₆NO₄[⊕] 356.441Quaternary alkaloid from *Berberis iliensis*. Mp 230-231° (as iodide). [α]_D +127.3 (c, 0.14 in MeOH) (iodide).**(S)-form** [6018-40-2]Alkaloid from *Dicentra oregana* and several *Corydalis* spp. (Papaveraceae).Cryst. (CHCl₃/MeOH). Mp 230° (246°). [α]_D¹⁷ -274 (c, 0.2 in CHCl₃).N-Me (cis-): **α-Hainanine**

[79523-84-5]

[79549-51-2]


C₂₁H₂₆NO₄[⊕] 356.441Quaternary alkaloid from *Cyclea hainanensis*.*Me ether*: see Tetrahydropalmatine, T-212**(±)-form** [27313-86-6]

Synthetic. Mp 215-217° (207°).

Späth, E. *et al.*, *Ber.*, 1923, **56**, 875 (*isol*)Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1939, **17**, 51 (*isol*)Govindachari, T.R. *et al.*, *Chem. Ber.*, 1959, **92**, 1654 (*synth, ir*)Zhang, X.X. *et al.*, *CA*, 1981, **95**, 192262t (*α-Hainanine*)Hussain, R.A. *et al.*, *Heterocycles*, 1989, **29**, 2257 (*pmr, cmr*)Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 83-84; *Chem. Nat. Compd. (Engl.**Transl.*), 1993, **29**, 69-70 (*N-Methylcorypalmine*)**Coryphanthine****C-695**

β-Methoxy-N,N,N-trimethylbenzeneethanaminium, 9CI


[85553-38-4]

C₁₂H₂₀NO[⊕] 194.296**(S)-form** [88147-90-4]Quaternary alkaloid from *Coryphantha greenwoodii* (Cactaceae).*Chloride*: [88147-92-6]Hygroscopic needles (MeOH/Me₂CO). Mp 161-163° (sealed tube). [α]_D +62 (c, 2.3 in H₂O).*Iodide*: [88147-91-5]Mp 180-181°. [α]_D +12.2 (c, 2 in H₂O).


[88198-81-6, 88147-93-7]

Meyer, B.N. *et al.*, *J. Nat. Prod.*, 1983, **46**, 688 (*isol, struct, spectra, synth*)**Coryphidine****C-696***Coriphidine*

[359436-27-4]

C₃₁H₄₄N₂O₃ 492.7Alkaloid from *Aconitum koreanum* (Ranunculaceae). Cryst. (MeOH). Mp 247-249°. Species name given as coreanum in the ref. λ_{max} 205 (log ε 3.95) (EtOH).Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*, 1992, **28**, 243-246; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 209-212 (*isol, uv, ir, pmr, ms*)**Coryphine****C-697***Corifine*

[142735-24-8]



Absolute Configuration

C₃₁H₄₂N₂O₂ 474.685Alkaloid from epigeal parts of *Aconitum koreanum* (Ranunculaceae). Mp 199-200°. [α]_D²⁰ +150 (c, 0.4 in MeOH).


Species name spelt as coreanum in the ref.

Perchlorate:

Cryst. (EtOH aq. or MeOH). Mp 225-226°.

Yusipova, I.M. *et al.*, *Khim. Prir. Soedin.*, 1991, 396-403; *Chem. Nat. Compd. (Engl. Transl.)*, 1991, **27**, 343-349 (*isol, pmr, cmr, ms, cryst struct*)**Coryrutine****C-698**

[104736-02-9]


C₂₂H₂₅NO₇ 415.442Coryrutine appears to be identical with *N*-Methylhydrasteine in *N*-35 in the lactol form, but this requires confirmation. Alkaloid from *Corydalis rutifolia* (Papaveraceae).*1-Oxo*: **Oxocoryrutine**

[156953-76-3]

C₂₂H₂₃NO₈ 429.426Alkaloid from whole plants of *Dactylicapnos torulosa* (Papaveraceae). Powder. Mp 173-175°. [α]_D²⁰ 0 (c, 1.0 in MeOH). Appears to be identical to *N*-Methylxohydrasteine in *N*-35 drawn in the lactol form but this requires confirmation. λ_{max} 230 (sh) (log ε 4.89); 294 (log ε 4.67); 320 (sh) (log ε 4.61) (MeOH).Şener, B. *et al.*, *CA*, 1986, **105**, 206252d (*Coryrutine*)Rücker, G. *et al.*, *Phytochemistry*, 1994, **36**, 519-523 (*Oxocoryrutine*)**Corysamine****C-699**

6,7-Dihydro-13-methylbis[1,3]benzodioxolo[5,6-a:4',5'-g]quinolinizinium, 9CI. 13-Methylcoptisine

[30243-28-8]


C₂₀H₁₆NO₄[⊕] 334.351Alkaloid from *Corydalis incisa*, *Corydalis cava*, *Corydalis lutea*, *Meconopsis napaulensis*, *Papaver*, *Glaucium*, *Argemone*, *Eschscholtzia* and *Bocconia* spp. (Papaveraceae). Alcohol dehydrogenase inhibitor.*Chloride*: [11028-77-6]C₂₀H₁₆ClNO₄ 369.803

Trihydrate. Mp 230° (210-211°).

Iodide: [96422-46-7]
 $C_{20}H_{16}INO_4$ 461.255
 Mp 300°.

Tani, C. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 598; 748 (*isol, struct*)
 Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1981, **46**, 2587; 1984, **49**, 704; 1318; 1985, **50**, 854; 2299 (*isol*)
 Hanaoka, M. *et al.*, *Chem. Comm.*, 1985, 1257 (*synth*)


Corystewartine C-700
 [119777-72-9]



$C_{21}H_{21}NO_5$ 367.401
 Alkaloid from the whole plant of *Corydalis stewartii* (Papaveraceae). $[\alpha]_D^{25} +82$ (c, 0.12 in $CHCl_3$).

13-Deoxy, 13,15-didehydro: **Ochotensidine** [119777-71-8]
 $C_{21}H_{19}NO_4$ 349.385
 Alkaloid from the whole plant of *Corydalis stewartii* (Papaveraceae). $[\alpha]_D^{25} +44$ (c, 0.13 in MeOH).
 Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1136 (*isol, uv, pmr, ms, struct, cd*)

Corytenchirine C-701




$C_{21}H_{25}NO_4$ 355.433

(-)-**form** [56974-41-5]
 Alkaloid from *Corydalis ochotensis* (Papaveraceae). Prisms (MeOH). Mp 246-247°. $[\alpha]_D^{24} -299$ (c, 1.0 in $CHCl_3$).

(+)-**form**
O²-De-Me: Hemiargyrine
 $C_{20}H_{23}NO_4$ 341.406
 Alkaloid from *Croton hemiargyreus*. Amorph. solid. Mp 164-166°. $[\alpha]_D^{25} +198$ (c, 0.1 in MeOH). λ_{max} 207; 228 (sh); 287 (MeOH).

(±)-**form** [60594-54-9]
 Synthetic. Prisms (MeOH). Mp 203-205°.
 Lu, S.T. *et al.*, *J.C.S. Perkin I*, 1976, 63 (*uv, ms, pmr*)
 Kametani, T. *et al.*, *J.C.S. Perkin I*, 1976, 1218 (*synth, ir, pmr*)
 Amaral, A.C.F. *et al.*, *Phytochemistry*, 1988, **47**, 1445-1447 (*Hemiargyrine, isol, uv, ir, pmr, cmr, ms*)

Corytuberine C-702
 1,11-Dihydroxy-2,10-dimethoxyaporphine



$C_{19}H_{21}NO_4$ 327.379
 Analeptic, antineoplastic agent. Respiratory stimulant, vagus depressant, increases salivation. Log P 2.42 (uncertain value) (calc). λ_{max} 208 (ε 31600); 226 (ε 25100); 276 (ε 5010); 318 (ε 5010) (EtOH/KOH) (Derep). λ_{max} 222 (ε 25100); 268 (ε 10000); 306 (ε 3160) (EtOH) (Derep).

(S)-**form** [517-56-6]
 Alkaloid from a variety of genera in the Papaveraceae (*Corydalis*, *Dicentra*, *Glaucium*, *Papaver*, *Dicranostigma*, *Eschscholtzia*), Lauraceae (*Mezilaurus*) and Menispermaceae (*Stephania*). Mp 240°. $[\alpha]_D^{25} +286$ (c, 0.1 in EtOH). Pharmacol. active isomer.


Di-Ac: Mp 72°.
N-Me: Magnoflorine. Thalictrine. Esholine. Escholine. Corytuberine methosalt [2141-09-5]
 $C_{20}H_{24}NO_4^{(+)}$ 342.414
 Alkaloid from *Magnolia obovata* (Magnoliaceae) and *Aquilegia hybrida* (Ranunculaceae), also from the Annonaceae, Berberidaceae, Euphorbiaceae, Menispermaceae, Papaveraceae, Rutaceae, Aristolochiaceae, and Rhamnaceae. Weak curarising and hypotensive agent. Shows antiyeast activity. Mp 248-249° (as iodide). $[\alpha]_D^{20} +193$ (c, 0.20 in MeOH) (iodide). Iodide shows grinding-induced polymorphism. Many lit. references where Magnoflorine has supposedly been identified may be in error because of the similarity in props. to *N,N*-Dimethylindcarpine in T-248. λ_{max} 227 (log ε 4.68); 275 (log ε 3.99); 315 (log ε 3.96) (H_2O) (iodide).

N-De-Me: 1,11-Dihydroxy-2,10-dimethoxynoraporphine. Muricinine
 $C_{18}H_{19}NO_4$ 313.352
 Alkaloid from *Annona muricata* (soursop) (Annonaceae). Tentative struct., not well characterised.

(±)-**form** [71630-07-4]
 Mp 242°. *Hydrochloride*: Mp 250° dec.
 Dobbie, J.J. *et al.*, *J.C.S.*, 1893, **63**, 485-488 (*isol*)
 Gulland, J.M. *et al.*, *J.C.S.*, 1928, 1834-1838 (*synth*)
 Späth, E. *et al.*, *Ber.*, 1931, **64**, 2038-2048 (*synth*)
 Manske, R.H.F. *et al.*, *Alkaloids (Academic Press)*, 1954, **4**, 142-143 (*Muricinine*)
 Nakano, T. *et al.*, *Chem. Pharm. Bull.*, 1954, **2**, 329-334 (*Magnoflorine*)
 Tomita, M. *et al.*, *Yakugaku Zasshi*, 1957, **77**, 157 (*Magnoflorine*)
 Albonico, S.M. *et al.*, *J.C.S. (C)*, 1966, 1340-1342 (*ord, abs config*)


Jackson, A.H. *et al.*, *J.C.S. (C)*, 1966, 2222-2229 (*pmr*)
 Slavic, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 3514-3520 (*Esholine*)
 Dominguez, X.A. *et al.*, *Phytochemistry*, 1974, **13**, 680 (*Magnoflorine*)
 Kametani, T. *et al.*, *J.A.C.S.*, 1977, **99**, 3805-3808 (*synth*)
 Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80-85 (*cd*)
 Tsai, I.L. *et al.*, *Kaohsiung J. Med. Sci.*, 1989, **5**, 132-145 (*Magnoflorine, activity*)
 Hoard, M.S. *et al.*, *Phytochemistry*, 1996, **43**, 1129-1133 (*Magnoflorine, polymorphism*)
 Barbosa-Filho, J.M. *et al.*, *Phytochemistry*, 1997, **44**, 959-961 (*Magnoflorine, isol, pmr, cmr*)
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*Magnoflorine, N-15 nmr*)

Coryzeylamine C-703
 [164176-11-8]



$C_{44}H_{52}N_4O_7$ 748.917
 Alkaloid from leaves of *Hunteria zeylanica* (Apocynaceae). Amorph. powder. $[\alpha]_D^{25} -24$ (c, 0.25 in MeOH).
 Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1957 (*isol, uv, pmr, cmr, struct*)

Coscinamide A C-704
 [298196-72-2]



$C_{20}H_{14}BrN_3O_2$ 408.254
 Isol. from the marine sponge *Coscino-derma* sp. Yellow solid. $[\alpha]_D -1.9$ (c, 1.3 in MeOH). λ_{max} 207 (log ε 4.47); 256 (log ε 4.12); 335 (log ε 3.96) (MeOH).


Debromo: Coscinamide B [298196-73-3]
 $C_{20}H_{15}N_3O_2$ 329.357
 Isol. from *Coscino-derma* sp. Orange solid. Mp 228-230° dec. $[\alpha]_D +23.6$ (c, 0.83 in MeOH). λ_{max} 207 (log ε 4.53); 229 (log ε 4.29); 267 (log ε 4.26); 347 (log ε 4.17) (MeOH).

7-Hydroxy: Coscinamide C [298196-74-4]

C₂₀H₁₄BrN₃O₃ 424.253
Isol. from *Coscinoderma* sp. Yellow solid. [α]_D -9.9 (c, 0.14 in MeOH). λ_{max} 213 (log ε 4.09); 261 (log ε 3.77); 363 (log ε 3.6) (MeOH).

Bokesch, H.R. *et al.*, *Tet. Lett.*, 2000, **41**, 6305-6308 (*Coscinamides*, *isol*, *pmr*, *cmr*)
Chakrabarty, M. *et al.*, *Synthesis*, 2003, 2011-2014 (*Coscinamide B*, *synth*)


Costinone A C-705



C₁₇H₁₃NO₆ 327.293
Alkaloid from *Isatis costata*. Pale yellow cryst. (EtOH). Mp 207-208°. [α]_D +110 (c, 0.12 in MeOH). Opt. rotn. unaccounted for.

Fatima, I. *et al.*, *Heterocycles*, 2006, **68**, 1421-1428 (*isol*, *pmr*, *cmr*, *ms*)

Costinone B C-706




C₁₈H₁₅NO₅ 325.32
Alkaloid from *Isatis costata*. Pale yellow cryst. (EtOH). Mp 195-197°. [α]_D +99.7 (c, 0.12 in MeOH). λ_{max} 210 (log ε 4.47); 230 (log ε 4.1); 279 (log ε 2.5) (MeOH).

Fatima, I. *et al.*, *Heterocycles*, 2006, **68**, 1421-1428 (*isol*, *pmr*, *cmr*, *ms*)

Cotarnoline C-707

7,8-Dihydro-4-hydroxy-6-methyl-1,3-dioxolo[4,5-g]isoquinolinium(1+), 9CI. 3,4-Dihydro-8-hydroxy-2-methyl-6,7-methylenedioxyisoquinolinium(1+)
[525-13-3]




C₁₁H₁₁NO₃ 205.213
CAS name refers to iminium salt form. Alkaloid from *Papaver pseudo-orientale* and other *Papaver* spp. (Papaveraceae). Amorph. orange solid.

Hydrochloride: [33329-37-2]
Mp 263°. Exists as the colourless

inium salt.
Goeber, B. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1966, **299**, 196
Sariyar, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1302 (*isol*, *uv*, *ms*, *pmr*)

Cotinine, INN C-708

1-Methyl-5-(3-pyridinyl)-2-pyrrolidinone, 9CI
[75202-09-4]



C₁₀H₁₂N₂O 176.218
Viscous oil. Bp₆ 210-211°. Log P -0.47 (uncertain value) (calc).

(S)-form [486-56-6]
Alkaloid from leaf tobacco (*Nicotiana tabacum*), also detected in *Duboisia hopwoodii* (Solanaceae). Antidepressant. Nicotine metabolite, used as a biomarker for exposure to cigarette smoke. Shows behavioural effects in animals. Mp 40-43°. Bp 210-211°. V. hygroscopic. Prob. artifact of autoxidn. of nicotine. ▶LD₅₀ (mus, orl) 1604 mg/kg. LD₅₀ (mus, ipr) 930 mg/kg. GN1925500

Fumarate (2:1): *Cotinine fumarate*, *USAN*. *Scotinine* [5695-98-7]


Picrate: Mp 102-103°.

(±)-form [15569-85-4]
Synthetic. Yellow oil. Bp 210-211°. *Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 294C (*nmr*)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1537C (*ir*)
Pinner, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1893, **231**, 378 (*synth*)
Frankenburg, W.G. *et al.*, *J.A.C.S.*, 1957, **79**, 149 (*isol*, *struct*, *ir*)
Testa, B. *et al.*, *Mol. Pharmacol.*, 1973, **9**, 10 (*conformm*)
Leete, E. *et al.*, *J.O.C.*, 1976, **41**, 3438; 1978, **43**, 2860 (*synth*, *ms*)
Tsujiino, Y. *et al.*, *Agric. Biol. Chem.*, 1979, **43**, 871 (*synth*)
Acheson, R.M. *et al.*, *J.C.S. Perkin 1*, 1980, **2**, 579 (*synth*)
Nishida, T. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 434 (*cmr*)
Luanratana, O. *et al.*, *Phytochemistry*, 1982, **21**, 449 (*isol*)
Benowitz, N.L. *et al.*, *Clin. Pharmacol. Ther. (St. Louis)*, 1983, **34**, 604 (*metab*)
Kyerematen, G.A. *et al.*, *Life Sci.*, 1983, **32**, 551 (*metab*)
Moriarty, R.M. *et al.*, *Tet. Lett.*, 1988, **29**, 6913 ((±)-form, *synth*)
Anderson, I.G. *et al.*, *Analyst (London)*, 1991, **116**, 691 (*gc*)
Sato, T. *et al.*, *Heterocycles*, 1992, **33**, 139 (*synth*)
Deutsch, J. *et al.*, *J. Chromatogr.*, 1992, **579**, 93 (*gc-ms*)
McAdams, S.A. *et al.*, *J. Chromatogr.*, 1993, **615**, 148 (*gc-ms*)
Keenan, R.M. *et al.*, *Clin. Pharmacol. Ther. (St. Louis)*, 1994, **55**, 581 (*pharmacol*)
Sato, T. *et al.*, *J.C.S. Perkin 1*, 1995, 1115 (*synth*)

Terry, A.V. *et al.*, *CNS Drug Rev.*, 2005, **11**, 229-252 (*rev*)

Coulteroberbinone C-709

[246262-28-2]




C₂₁H₂₀NO₆⁺ 382.392
Ref. config. only detd. Quaternary alkaloid from *Romneya coulteri*. Powder (MeOH/CHCl₃/Et₂O) (as chloride). Mp 212-213° (chloride). [α]_D -53 (c, 0.4 in MeOH) (chloride). λ_{max} 240 (log ε 4.19); 290 (log ε 3.95); 324 (log ε 3.75) (MeOH) (chloride).

Valpuesta, M. *et al.*, *Phytochemistry*, 1999, **51**, 1157-1160 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Coumaperine C-710

1-[5-(4-Hydroxyphenyl)-1-oxo-2,4-pentadienyl]piperidine, 9CI. N-[5-(4-Hydroxyphenyl)-2,4-pentadienyl]piperidine. 5-(4-Hydroxyphenyl)-2,4-pentadienoic acid piperidide
[76733-91-0]



C₁₆H₁₉NO₂ 257.332
Isol. from the fruits of pepper (*Piper nigrum*) (Piperaceae). Cryst. (Me₂CO). Mp 199.5-200.5°.

Ac: [76733-93-2]
Needles (Et₂O). Mp 127-128°.


3'-Methoxy: *Feruperine* [77795-15-4]
C₁₇H₂₁NO₃ 287.358
Alkaloid from pepper (*Piper nigrum*) (Piperaceae). Antioxidant. Pale yellow needles (C₆H₆). Mp 159°.

3'-Methoxy, 4',5'-dihydro: *Dihydroferuperine* [77795-17-6]
C₁₇H₂₃NO₃ 289.374
Alkaloid from pepper (*Piper nigrum*) (Piperaceae). Cryst. (petrol). Mp 78°.

Nakatani, N. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 2831 (*isol*, *ms*, *uv*, *ir*, *pmr*, *struct*, *synth*)
Inatani, R. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 667 (*Feruperine*, *Dihydroferuperine*)

Coumermycin A₁ C-711


Coumermycin, *USAN*. *Coumamycin*, *INN*. *Notomycin A₁*. *Sugordomycin D_{1a}*. *BU 620*. *NSC 107412*. *Antibiotic BU 620*. *Ro 5-4645/10*
[4434-05-3]
[87901-11-9 (Na salt)]



C₅₅H₅₉N₅O₂₀ 1110.093
 Isol. from *Streptomyces rishiriensis* and other *Streptomyces* spp. Active against gram-positive and -negative bacteria and mycobacteria. Sol. MeOH, Me₂CO, THF, Py, bases, dioxan, DMF, EtOAc, butanol, EtOH; fairly sol. CHCl₃, C₆H₆; poorly sol. hexane, CCl₄, H₂O, acids. Mp 258-260° dec. [α]_D²⁰ -141 (c, 1.0 in 75% Me₂CO aq.). Also used as Na salt (Coumermycin sodium, USAN). λ_{max} 280 (ε 44200); 345 (ε 44300) (EtOH/HCl) (Derep). λ_{max} 280 (ε 62200); 308 (ε 34200) (EtOH/NaOH) (Derep). λ_{max} 280 (ε 59400); 336 (ε 42200) (EtOH) (Derep).

- ▶ LD₅₀ (mus, ivn) 25 mg/kg, LD₅₀ (mus, ipr) 159 mg/kg, LD₅₀ (mus, scu) 380 mg/kg, LD₅₀ (mus, orl) 2000 mg/kg. UX9375000
- Berger, J. et al., *Antimicrob. Agents Chemother.*, 1965, 778 (synth)
- Kawaguchi, H. et al., *J. Antibiot., Ser. A*, 1965, 18, 11 (struct)
- Wick, A.E. et al., *Tetrahedron*, 1976, 32, 2057 (config)
- Berger, J. et al., *J. Chromatogr. Libr.*, 1978, 15, 101 (rev)
- Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 19000
- Neu, H.C. et al., *Antimicrob. Agents Chemother.*, 1984, 25, 687 (props)
- Claridge, C.A. et al., *Drugs Pharm. Sci.*, 1984, 22, 413 (rev)
- Olson, S.H. et al., *Tet. Lett.*, 2003, 44, 61-63 (synth)
- Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CNV500

Coumermycin A₂ C-712
Notomycin A₂. Sugordomycin D_{1d}
 [3130-60-7]



C₅₃H₅₅N₅O₂₀ 1082.039
 Isol. from *Streptomyces rishiriensis* and other *Streptomyces* spp. Sol. MeOH, bases, EtOAc; fairly sol. CHCl₃, C₆H₆; poorly sol. hexane, H₂O. λ_{max} 267 (ε 50000); 340 (ε 42000) (EtOH) (Derep).


- Berger, J. et al., *Antimicrob. Agents Chemother.*, 1965, 778 (isol)
- Kawaguchi, H. et al., *J. Antibiot., Ser. A*, 1965, 18, 220 (struct)
- Berger, J. et al., *J. Chromatogr. Libr.*, 1978, 15, 101 (rev)
- Claridge, C.A. et al., *Drugs Pharm. Sci.*, 1984, 22, 413 (rev)

Coumingaine C-713

C₃₂H₅₁NO₈ 577.757
 Struct. unknown. *Erythrophleum* alkaloid. Poorly descr. in the lit. Alkaloid from *Erythrophleum couminga* (Fabaceae). Cardiac stimulant showing digitalis-like activity. Local anaesthetic, hypertensive agent. Amorph.

- ▶ Toxic.
- Chen, K.K. et al., *J. Am. Pharm. Assoc.*, 1938, 27, 9-16


Coyhaiquine C-714
 [85643-88-5]



C₂₆H₂₇NO₅ 433.503
 Alkaloid from *Berberis empetrifolia* (Berberidaceae). [α]_D²⁵ +28 (c, 0.02 in MeOH). The first known oxidized proaporphine-benzylisoquinoline alkaloid.

- 1''-Aldehyde: *Coyhaiquinine* [101242-45-9]
- C₂₆H₂₅NO₅ 431.487
- Constit. of *Berberis empetrifolia*.
- Fajardo, V. et al., *Chem. Comm.*, 1982, 1350 (isol, uv, pmr, ms, cd, struct)
- Fajardo, V. et al., *Bol. Soc. Chil. Quim.*, 1985, 30, 51; *CA*, 104, 145455c (*Coyhaiquinine*)

Crambescidin 359 C-715
 [259733-99-8]



C₂₁H₃₃N₃O₂ 359.511
 Alkaloid from the sponge *Monanchora unguiculata*. Gum. [α]_D²⁵ -9 (c, 0.18 in CH₂Cl₂).


- 16S-Hydroxy: *16-Hydroxycrambescidin 359* [956142-72-6]
- C₂₁H₃₃N₃O₃ 375.51
- Alkaloid from *Monanchora unguifera*. Cryst. [α]_D²⁵ -3 (c, 0.1 in MeOH). λ_{max} 208 (ε 6334) (MeOH).
- Braekman, J.C. et al., *J. Nat. Prod.*, 2000, 63, 193-196 (*Crambescidin 359*)
- Nagasawa, K. et al., *Org. Lett.*, 2002, 4, 177-180 (synth)
- Aron, Z.D. et al., *J.A.C.S.*, 2005, 127, 3380-3390 (synth)
- Hua, H.-M. et al., *Tetrahedron*, 2007, 63, 11179-11188 (*16β-Hydroxycrambescidin 359*)
- Moore, C.G. et al., *Tetrahedron*, 2007, 63, 11771-11780 (synth)

Crambescidin 431 C-716

[259734-00-4]
 As Crambescidin 359, C-715 with R = -COOEt
 C₂₄H₃₇N₃O₄ 431.574
 Alkaloid from the sponge *Monanchora unguiculata*. Gum. [α]_D²⁵ +12 (c, 0.19 in CH₂Cl₂).

- Parent acid: *Crambescidin acid* [147664-30-0]
- C₂₂H₃₃N₃O₄ 403.52
- Alkaloid from *Monanchora unguiculata*. [α]_D²⁵ -19 (c, 0.04 in MeOH). λ_{max} 231 (log ε 3.2); 268 (log ε 2.84) (MeOH).
- Braekman, J.C. et al., *J. Nat. Prod.*, 2000, 63, 193-196 (*Crambescidin 431*)
- Meragelman, K.M. et al., *J. Nat. Prod.*, 2004, 67, 1165-1167 (*Crambescidin acid*)
- Aron, Z.D. et al., *J.A.C.S.*, 2005, 127, 3380-3390 (synth)

Crambescidin 816 C-717
 [135257-45-3]




C₄₅H₈₀N₆O₇ 817.163
 Alkaloid from the sponges *Crambe crambe* and *Batzella* sp. Possesses antiviral and cytotoxic props. Ca antagonist, ichthyotoxin. Oil. [α]_D²⁵ -20.14 (c, 0.4 in MeOH).

- 13-Deoxy: *Crambescidin 800* [135257-46-4]
- C₄₅H₈₀N₆O₆ 801.164
- From the sponges *Crambe crambe* and *Batzella* sp. Possesses antiviral and cytotoxic props. Ichthyotoxin. Oil.
- 13-Deoxy, 13,14,15-triepipimer: *Isocrambescidin 800* [151121-78-7]
- C₄₅H₈₀N₆O₆ 801.164
- Alkaloid from *Crambe crambe*. Ichthyotoxin. Oil. [α]_D²⁵ -48 (c, 0.53 in MeOH).

- Jares-Erijman, E.A. et al., *J.O.C.*, 1991, 56, 5712-5715; 1993, 58, 4805-4808 (isol, pmr, cmr, struct, derivs)
- Berlinck, R.G.S. et al., *J. Nat. Prod.*, 1993, 56, 1007-1015 (isol)
- Coffey, D.S. et al., *J.A.C.S.*, 1999, 121, 6944-6945; 2000, 122, 4893-4903; 4904-4914 (synth, *Isocrambescidin 800*)

Crambescidin 826 C-718
 [628727-32-2]



C₄₇H₈₂N₆O₆ 827.201
 Alkaloid from the sponge *Monanchora*

sp. Inhibitor of HIV-1 fusion. Glassy solid. $[\alpha]_D^{20}$ -7.7 (c, 0.09 in MeOH). λ_{\max} 207 (log ϵ 4.04); 274 (log ϵ 3.24) (MeOH).

Chang, L.C. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1490-1494 (*isol*, *pmr*, *cmr*)

Crambescidin 830 C-719

[135257-47-5]

As Crambescidin 816, C-717 with $n = 16$

$C_{46}H_{82}N_6O_7$ 831.19

Alkaloid from the sponge *Crambe crambe*. Possesses antiviral and cytotoxic props. Oil.

Jares-Erijman, E.A. *et al.*, *J.O.C.*, 1991, **56**, 5712 (*isol*, *pmr*, *cmr*, *struct*)

Crambescidin 844 C-720

[135283-73-7]

As Crambescidin 816, C-717 with $n = 17$

$C_{47}H_{84}N_6O_7$ 845.217


Alkaloid from the sponge *Crambe crambe*. Possesses antiviral and cytotoxic props. Oil. $[\alpha]_D^{25}$ -10.32 (c, 0.19 in MeOH).

Jares-Erijman, E.A. *et al.*, *J.O.C.*, 1991, **56**, 5712 (*isol*, *pmr*, *cmr*, *struct*)

Crambescin A† C-721

Crambine A

[132210-62-9]



$n = 10$

$C_{24}H_{44}N_6O_2$ 448.651

Major component of bis-guanidine complex where in addition $n = 9, 11$ and 12 . Alkaloids from the marine sponge *Crambe crambe*. Glassy solid. $[\alpha]_D +2$ (c, 0.7 in MeOH). λ_{\max} 204 (ϵ 10600); 286 (ϵ 5500) (MeOH) (Derep).

Lower homologue ($n = 4$): **Antibiotic Sch 575948**. *Sch 575948*

$C_{18}H_{32}N_6O_2$ 364.49


Isol. from the marine sponge *Ptilocaulis spiculifer*. Antibacterial agent.

Berlinck, R.G.S. *et al.*, *Tet. Lett.*, 1990, **31**, 6531-6534 (*Crambine A*, *isol*, *pmr*, *struct*)
Yang, S.W. *et al.*, *J. Antibiot.*, 2003, **56**, 970-972 (*Sch 575948*)

Crambescin B C-722

Crambine B

[132210-63-0]



$x = 8, y = 7$

$C_{25}H_{48}N_6O_3$ 480.693

Major component of bis-guanidine com-

plex where in addition $x + y = 14, 16, 17$. Struct. revised in 1993; originally descr. as having $x = 10, y = 5$. Alkaloids from the sponge *Crambe crambe*. Cytotoxic against L1210 murine leukaemia cells. Cell reagent-inhibitor. Ichthyotoxin. Glassy solid. $[\alpha]_D +52$ (c, 0.9 in MeOH).

Berlinck, R.G.S. *et al.*, *Tet. Lett.*, 1990, **31**, 6531-6534 (*isol*, *pmr*)


Snider, B.B. *et al.*, *J.O.C.*, 1992, **57**, 2526-2528 (*stereochem*)

Jares-Erijman, E.A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2186-2188 (*struct*)

Crambescin C1 C-723

Crambine C1

[142605-06-9]



$x = 8, y = 7$

$C_{25}H_{48}N_6O_3$ 480.693

Major homologue of bis-guanidine complex where in addn. $x + y = 14, 16$ and 17 . Struct revised in 1993; originally descr. as the compd. with $x = 10, y = 5$. Alkaloid from the Mediterranean sponge *Crambe crambe*. Ichthyotoxic agent. Glassy solid. Sol. MeOH, $CHCl_3$. $[\alpha]_{404} +13.7$ (c, 0.8 in MeOH). λ_{\max} 207 (ϵ 4600); 279 (ϵ 2200) (MeOH) (Derep).

Berlinck, R.G.S. *et al.*, *J. Nat. Prod.*, 1992, **55**, 528-532 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cd*)
Jares-Erijman, E.A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2186-2188 (*struct*)

Crambescin C2 C-724

Crambine C2

[142605-07-0]

As Crambescin C1, C-723 with $x + y = 14$

$C_{24}H_{46}N_6O_3$ 466.666


Originally descr. as the compd. with $x = 10, y = 4$, but this is doubtful in view of the revised struct. of Crambescin C1.

Alkaloid from the sponge *Crambe crambe*. Ichthyotoxic agent. Glassy solid. Sol. MeOH, $CHCl_3$. $[\alpha]_D -30$ (c, 0.2 in MeOH). λ_{\max} 204 (ϵ 10600); 286 (ϵ 5500) (MeOH) (Derep). λ_{\max} 203 (ϵ 6000); 281 (ϵ 2500) (MeOH) (Berdy).

Berlinck, R.G.S. *et al.*, *J. Nat. Prod.*, 1992, **55**, 528-532 (*isol*, *uv*, *pmr*)
Jares-Erijman, E.A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2186-2188 (*nomencl*)

Crambidine C-725

[151271-96-4]



Absolute Configuration

$C_{45}H_{79}N_6O_6^{\oplus}$ 800.156

Alkaloid from the marine sponge *Crambe crambe*. Shows ichthyotoxic props. Glassy solid (as per-Ac, chloride).


Berlinck, R.G.S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1007-1015 (*isol*, *uv*, *pmr*, *cmr*)

Overman, L.E. *et al.*, *J.A.C.S.*, 2005, **127**, 15652-15658 (*synth*, *abs config*)

Crassanine C-726

Conopharyngine oxindole

[16790-92-4]



Relative Configuration


$C_{23}H_{30}N_2O_5$ 414.5

Minor alkaloid from *Tabernaemontana crassa* (Apocynaceae). Plates ($CHCl_3$). Mp 190-191°. $[\alpha]_D^{26} +21.4$ (c, 0.013 in EtOH). λ_{\max} 210 (log ϵ 4.37); 275 (log ϵ 3.68); 302 (log ϵ 3.55) (no solvent reported).

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 3350-3552 (*isol*, *uv*, *ir*, *ms*, *pmr*, *struct*)

Crassifolazonine C-727


[96910-80-4]



$C_{19}H_{23}NO_4$ 329.395

Alkaloid from *Corydalis claviculata* (Papaveraceae). Cryst. (hexane/ C_6H_6). Mp 160-162°. $[\alpha]_D^{20} -50$ (c, 0.06 in $CHCl_3$).

Boente, J.M. *et al.*, *Heterocycles*, 1985, **23**, 1069 (*uv*, *pmr*, *cmr*, *struct*, *synth*)

Crassifoline C-728

(R)-form

$C_{19}H_{23}NO_4$ 329.395

(R)-form

N-Me: Isotembetarine

[186502-92-1]

$C_{20}H_{26}NO_4^{\oplus}$ 344.43

Quaternary alkaloid from roots of *Zanthoxylum nitidum*. Cryst. (Me_2CO /hexane) (as perchlorate). Mp 172° (perchlorate). $[\alpha]_D^{25} +3.5$ (c, 0.43 in

MeOH).

(S)-form [87098-80-4]

Alkaloid from *Sarcocapnos crassifolia* and *Corydalis claviculata* (Papaveraceae). Oil; cryst. (EtOH) (as perchlorate). Mp 228-230° (as perchlorate). $[\alpha]_D^{22} +37.6$ (c, 0.1 in EtOH). The first 7,8,3',4'-tetraoxygenated benzyloquinoline from a natural source.

(±)-form [31278-84-9]


Synthetic. Needles (Et₂O/petrol). Mp 61-63°.

- Jackson, A.H. *et al.*, *J.C.S. Perkin 1*, 1974, 1911 (*synth*, *pmr*)
 Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2303 (*ir*, *pmr*, *ms*)
 Guinaudeau, H. *et al.*, *Heterocycles*, 1984, **22**, 107 (*isol*)
 Blaschke, G. *et al.*, *Phytochemistry*, 1985, **24**, 585 (*isol*, *uv*, *pmr*, *cmr*, *ms*, *cd*, *config*)
 Moriyasu, M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 299 (*Isotembetarine*)
 Rodrigues, J.A.R. *et al.*, *J.O.C.*, 2004, **69**, 2920-2928 (*synth*)

Crassifoline methine

C-729

3-[2-(Dimethylamino)ethyl]-2-[2-(3-hydroxy-4-methoxyphenyl)ethenyl]-6-methoxyphenol, 9CI. 2-(Dimethylaminoethyl)-3',6-dihydroxy-4',5-dimethoxystilbene [133084-00-1]

C₂₀H₂₅NO₄ 343.422


Alkaloid from *Corydalis claviculata* (Papaveraceae).

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1280 (*isol*, *struct*)

Craugsodine

C-730

5-[[[2-(4-Hydroxyphenyl)ethyl]imino]-methyl]-2-methoxyphenol, 9CI. N-(3-Hydroxy-4-methoxybenzylidene)-4'-hydroxyphenethylamine [102686-18-0]

C₁₆H₁₇NO₃ 271.315

Isol. from the flower-stem fluid of *Crinum augustum* (Amaryllidaceae). Possible direct precursor of Amaryllidaceae alkaloids. Light-brown solid. Mp 144-146°.

4'-O-De-Me, O^{3'}-Me: **Isocraugsodine**

[116528-01-9]
 [116528-00-8, 116527-99-2]
 C₁₆H₁₇NO₃ 271.315

Alkaloid from the fruits of *Crinum asiaticum* (Amaryllidaceae).

Considered a direct biogenetic precursor of the Amaryllidaceae alkaloids. Orange-yellow needles (MeOH). Mp 220°. Exists as 3 isomers in soln. (*E*-form ⇌ quinone methide ⇌ *Z*-form).

Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1986, 28 (*Craugsodine*)

Ghosal, S. *et al.*, *Phytochemistry*, 1988, **27**, 1849 (*Isocraugsodine*)

Creatine

C-731

N-(Aminoiminomethyl)-N-methylglycine, 9CI. Methylguanidinoacetic acid [57-00-1]

HN=C(NH₂)NMeCH₂COOHC₄H₉N₃O₂ 131.134

Constit. of blood, muscle and other tissues. Prisms + 1H₂O. Mod. sol. hot H₂O, spar. sol. EtOH, insol. Et₂O. Mp 303°.

Hydrate: [6020-87-7]

Mp 292° dec.

Hydrobromide: [15366-30-0]

Cryst. (EtOH/Et₂O). Mp 150-151°.

Picrate:

Needles (H₂O). Mp 218-220°.

N,N-Di-Ac:

C₈H₁₃N₃O₄ 215.208

Needles (EtOH). Mp 165°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 785B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1280B (*nmr*)

Studel, H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1921, **112**, 54 (*synth*)

King, H. *et al.*, *J.C.S.*, 1930, 2374 (*synth*)


Yoshizaki, K. *et al.*, *Biochim. Biophys. Acta*, 1981, **678**, 283 (*pmr*)

Tsuchiya, M. *et al.*, *Int. J. Mass Spectrom. Ion Phys.*, 1983, **46**, 355 (*ms*)

Creatinine

C-732

2-Amino-1,5-dihydro-1-methyl-4H-imidazol-4-one, 9CI. 2-Amino-1-methyl-2-imidazol-4-one [60-27-5]

C₄H₇N₃O 113.119

Tautomeric: the amino and imino (2 stereoisomers) forms are of similar energy. Constit. of urine, muscle, etc. Metabolite of analytical importance in clinical chemistry, particularly as indicator of renal function. Rhombic prisms. Mp 305° dec. Forms stable addn. compd. with ZnCl₂. Undergoes ring-opening to Creatine, C-731 in alkaline soln. (Jaffé reaction).

Picrate:

Needles. Mp 220-221°.

[62708-52-5]

Aldrich Library of NMR Spectra, 2nd edn.,

1983, **2**, 489C (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 618A (*ir*)

Greenwald, I. *et al.*, *J. Biol. Chem.*, 1929, **81**, 73 (*synth*)

King, H. *et al.*, *J.C.S.*, 1930, 2377 (*synth*)

Org. Synth., Coll. Vol., **1**, 1932, 172 (*synth*)

Kenyon, G.L. *et al.*, *J.A.C.S.*, 1971, **93**, 5552 (*struct*)

Rowley, G.L. *et al.*, *J. Het. Chem.*, 1972, **9**, 203 (*synth*)

Butler, A.R. *et al.*, *J.C.S. Perkin 2*, 1985, 1465 (*cmr*, *tautom*)


Kotsyubynsky, D. *et al.*, *Magn. Reson.*

Chem., 2004, **42**, 1027-1036 (*pmr*, N-15 *nmr*)

Crebanine

C-733

6,7,7a,8-Tetrahydro-9,10-dimethoxy-7-methyl-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, 9CI. 8,9-Dimethoxy-1,2-methylenedioxyaporphine

C₂₀H₂₁NO₄ 339.39**(R)-form** [25127-29-1]

Alkaloid from *Stephania capitata*, *Stephania sasakii* and *Stephania venosa* (Menispermaceae). Mp 115-118°. $[\alpha]_D -61$ (c, 0.44 in CHCl₃).

►DE5376000

N-Oxide: **Crebanine N-oxide**

[102719-85-7]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Stephania succifera* (Menispermaceae). Mp 132-134°. $[\alpha]_D -66$.

N-De-Me, N-methoxycarbonyl: **8,9-Dimethoxy-N-methoxycarbonyl-1,2-methylenedioxyaporphine**

C₂₁H₂₁NO₆ 383.4

Alkaloid from the stems of *Fissistigma bracteolatum*. Powder. Mp 190-192°.

$[\alpha]_D^{20} -265.2$ (c, 0.01 in CHCl₃). λ_{max} 218 (log ε 4.47); 239 (log ε 4.09); 282 (log ε 4.24); 322 (log ε 3.56) (no solvent reported).

O⁸-De-Me, N-de-Me: **8-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine. Norannuradhapurine**

[83694-79-5]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the bark and leaves of *Polyalthia acuminata* (Annonaceae). Amorph. Annuradhapurine appears to be unknown.

O⁹-De-Me: **9-Hydroxy-8-methoxy-1,2-methylenedioxyaporphine. Stesakine**

[77784-23-7]

Alkaloid from the stem and root of *Stephania sasakii* and the seeds of *Stephania cepharantha* (Menispermaceae). Mp 188-190°. $[\alpha]_D -78.7$ (CHCl₃).

O⁹-De-Me, O-β-D-glucopyranoside: **Stesakine O-β-D-glucopyranoside**

C₂₅H₂₉NO₉ 487.505Alkaloid from the seeds of *Stephania cepharantha*. Amorph. powder. [α]_D²⁷ -74.5 (c, 0.2 in MeOH).**6a,7-Didehydro: Dehydrocrebanine**

[77784-22-6]

C₂₀H₁₉NO₄ 337.374Alkaloid from the stem and root of *Stephania sasakii* and from the seeds of *Stephania cepharantha*. Also present in *Stephania venosa* (Menispermaceae). Pale yellow needles (EtOH). Mp 152-153°.**►E5378000****6a,7-Didehydro, O⁹-de-Me: Dehydrostesakine**

[77784-24-8]

C₁₉H₁₇NO₄ 323.348Alkaloid from the stem and roots of *Stephania sasakii* (Menispermaceae). Cryst. (Me₂CO). Mp 201-203°.**4S-Hydroxy: 4-Hydroxycrebanine**

[79559-57-2]

C₂₀H₂₁NO₅ 355.39Alkaloid from *Stephania sasakii* (Menispermaceae). Needles (Me₂CO). Mp 191-192°. [α]_D²⁵ -90.2 (c, 0.266 in CHCl₃).**7R-Hydroxy: Sukhodianine**

[82413-17-0]

C₂₀H₂₁NO₅ 355.39Alkaloid from tuberous roots of *Stephania venosa* (Menispermaceae). Amorph.**7R-Hydroxy, N-oxide (β-): Sukhodianine β-N-oxide**

[113145-65-6]

C₂₀H₂₁NO₆ 371.389Alkaloid from the rhizomes of *Stephania venosa* (Menispermaceae). Amorph. [α]_D -13 (c, 0.07 in MeOH).**7R-Acetoxy: O-Acetylsukhodianine**

[98604-29-6]

C₂₂H₂₃NO₆ 397.427Alkaloid from the leaves of *Stephania venosa* (Menispermaceae). [α]_D²⁵ -68 (c, 0.06 in CHCl₃).**(±)-form**

Synthetic. Stout needles (petrol). Mp 123-123.5°.


Hydriodide:

Needles (MeOH). Mp 250° dec.


4R-Hydroxy: Synthetic. Cryst. (C₆H₆). Mp 195-196°.**4S-Hydroxy:** Synthetic. Amorph.Govindachari, T.R. *et al.*, *J.C.S.*, 1958, 983 (*synth, uv, bibl*)Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1969, 89, 1691; *CA*, 73, 4072e (*isol*)Kunitomo, J. *et al.*, *Phytochemistry*, 1980, 19, 2735 (*Dehydrocrebanine, Stesakine, Dehydrostesakine*)Guinaudeau, H. *et al.*, *Chem. Comm.*, 1981, 1118 (*Dehydrocrebanine*)Kumimoto, J. *et al.*, *Chem. Pharm. Bull.*, 1981, 29, 2251 (*4-Hydroxycrebanine*)Pharadai, K. *et al.*, *Heterocycles*, 1981, 15, 1067 (*cd, pmr, abs config*)Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1981, 101, 951 (*Stesakine, Dehydrostesakine*)Kunitomo, J. *et al.*, *Heterocycles*, 1982, 19, 1883 (*cd, ord, abs config*)Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1982, 45, 355 (*Sukhodianine*)Zarga, M.H.A. *et al.*, *J. Nat. Prod.*, 1982, 45, 471 (*Norannuradhapurine*)Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1985, 33, 5245 (*synth, uv, ir, pmr, ms, 4-Hydroxycrebanine*)Pharadai, K. *et al.*, *J. Nat. Prod.*, 1985, 48, 658 (*Acetylsukhodianine*)Xue, Z. *et al.*, *Yaoyue Xuebao*, 1986, 21, 223; *CA*, 105, 3560e (*Crebanine N-oxide*)Charles, B. *et al.*, *J. Nat. Prod.*, 1987, 50, 1113 (*Sukhodianine β-N-oxide*)Bartley, J.P. *et al.*, *Phytochemistry*, 1994, 36, 1327-1331 (*isol, pmr, cmr*)Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 2000, 63, 477-479 (*Stesakine glucoside*)Deng, Y. *et al.*, *Chin. Chem. Lett.*, 2002, 13, 862-864 (*N-de-Me N-methoxycarbonyl*)Blanchfield, J.T. *et al.*, *Phytochemistry*, 2003, 63, 711-720 (*isol, pmr, cmr*)**Crenulatin†**

[343782-11-6]


C-734

C₁₆H₂₁NO₂ 259.347Alkaloid from stems of *Limonia crenulata*. Amorph. powder. [α]_D²⁴ -12.5 (c, 0.7 in MeOH). λ_{\max} 223 (log ϵ 4.54); 281 (log ϵ 3.87); 289 (log ϵ 3.79) (MeOH).Niu, X.-M. *et al.*, *Chin. Chem. Lett.*, 2001, 12, 243-244 (*isol, pmr, cmr*)Niu, X.-M. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, 3, 299-311 (*isol, pmr, cmr*)**Crepidamine**

C-735

1-(Octahydro-6-hydroxy-7-methyl-6-phenyl-5-indolizyl)-2-propanone, 9CI [50906-93-9]C₁₈H₂₅NO₂ 287.401Alkaloid from *Dendrobium crepidatum* (Orchidaceae). Needles (Et₂O). Mp 107.5-109°. Opt. inact. (racemic).Elander, M. *et al.*, *Acta Chem. Scand.*, 1973, 27, 1907 (*isol, uv, ord, ir, pmr, ms, struct*)**Crepidine**

C-736


1-(Decahydro-4,6-dihydroxy-4,7-dimethyl-6-phenyl-1H-pyrrolo[2,1,5-de]quinolizin-3-yl)ethanone, 9CI [34443-74-8]

Relative configuration

C₂₁H₂₉NO₃ 343.465Alkaloid from *Dendrobium crepidatum* (Orchidaceae). Needles (EtOH). Mp 221-222°. [α]_D²⁴ -82 (c, 0.43 in MeOH).**Methodide:** Mp 240-242° dec. [α]_D²⁴ -17 (c, 1.04 in MeOH).Kierkegaard, P. *et al.*, *Acta Chem. Scand.*, 1970, 24, 3757 (*cryst struct*)Pilotti, A.-M. *et al.*, *Acta Cryst. B*, 1971, 27, 887 (*cryst struct*)Elander, M. *et al.*, *Acta Chem. Scand.*, 1973, 27, 1907 (*isol, ir, uv, pmr, ms, ord*)**Criamide A**

C-737

[169181-26-4]


C₃₅H₅₆N₈O₅ 668.878Peptide antibiotic. Isol. from the sponge *Cymbastela* sp. Cytotoxic agent. Microtubule formation inhibitor. Amorph. solid. Sol. MeOH. [α]_D²⁵ +97 (c, 0.02 in MeOH). Related to Hemiasterlin, H-98.**N¹-Me: Criamide B**

[169181-27-5]

C₃₆H₅₈N₈O₅ 682.905Isol. from *Cymbastela* sp. Cytotoxic agent. Amorph. solid. Sol. MeOH.Coleman, J.E. *et al.*, *Tetrahedron*, 1995, 51, 10653-10662 (*isol, pmr, cmr*)**Cribrochalinamine oxide A**

C-738

[152273-70-6]



n = 3

C₂₁H₃₆N₂O 332.528Alkaloid from the marine sponge *Cribrachalina* sp. Shows antifungal activity. Sol. MeOH, Et₂O; poorly sol. H₂O. λ_{\max} 255 (ϵ 2100); 270 (ϵ 1900) (MeOH) (Derep).Matsunaga, S. *et al.*, *Tet. Lett.*, 1993, 34, 5953 (*isol, uv, pmr, cmr, struct*)**Cribrochalinamine oxide B**

C-739

[152273-71-7]

As Cribrochalinamine oxide A, C-738 with


n = 5, $\Delta^{13,14}$ (Z-)C₂₃H₃₈N₂O 358.566Alkaloid from the marine sponge *Cribrachalina* sp. H₂O. λ_{\max} 255 (ϵ 2100); 270 (ϵ 1900) (MeOH) (Derep). λ_{\max} 255 (ϵ 1700); 271 (ϵ 1500) (MeOH) (Berdy).

Matsunaga, S. *et al.*, *Tet. Lett.*, 1993, **34**, 5953
(*isol, uv, pmr, cmr, struct*)

Cribrostatin 6

C-740

[518056-45-6]



$C_{15}H_{14}N_2O_3$ 270.287

Isol. from the marine sponge *Cribrochalina* sp. Cancer cell growth inhibitor. Antibacterial agent. Dark blue needles (Me₂CO). Mp 169-171°. λ_{max} 203 (ε 26760); 266 (ε 24430); 323 (ε 5600); 552 (ε 1480) (no solvent reported).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 544-547 (*isol, uv, pmr, cmr, ms, cryst struct*)

Nakahara, S. *et al.*, *Heterocycles*, 2004, **63**, 2355-2362; 2006, **68**, 515-520 (*synth, pmr, cmr*)


Pettit, R.K. *et al.*, *J. Med. Microbiol.*, 2004, **53**, 61-65 (*activity*)

Markey, M.D. *et al.*, *J.O.C.*, 2008, **73**, 7441-7443 (*synth*)

Crinafolidine

C-741

[106549-07-9]




$C_{19}H_{23}NO_6$ 361.394

Alkaloid from the mature fruits of *Crinum latifolium* (Amaryllidaceae). Has antitumour props. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O.

Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1986, 312-313 (*isol, uv, ir, pmr, struct*)

Crinamabine

C-742



Absolute Configuration

$C_{16}H_{19}NO_5$ 305.33


Alkaloid from the bulbs of *Crinum amabile*. Mp 235-238°. $[\alpha]_D^{25}$ +35 (c, 0.09 in MeOH). Dec. at 250°. λ_{max} 206 (log ε 4.08); 238 (sh) (log ε 3.21); 292 (log ε 3.25) (MeOH).

Pham, L.H. *et al.*, *Phytochemistry*, 1998, **48**, 371-376 (*isol, cd, uv, ir, pmr, cmr, ms*)

Crinamidine

C-743

Crinalbine. 1,2-Epoxy Powelline



$C_{17}H_{19}NO_5$ 317.341

Crinalbine was the (+)-form and Crinamidine the (-)-form.

(+)-form

Alkaloid from *Crinum powellii* var. *album* (Amaryllidaceae). Mp 235-236° dec. $[\alpha]_D^{25}$ +23 (c, 0.2 in CHCl₃). One reference erroneously gives a -ve specific rotn.

(-)-form [6793-66-4]

Alkaloid from *Crinum powellii*, *Crinum moorei* and several other spp. in the Amaryllidaceae. Mp 235-236° dec. $[\alpha]_D^{23}$ -24 (c, 0.6 in CHCl₃).

Picrate: Mp 131-132°.

Me ether: **Undulatin**[†]

[6882-09-3]

$C_{18}H_{21}NO_5$ 331.368

Alkaloid from *Nerine undulata* and other *Nerine* spp. (Amaryllidaceae). Mp 151-152°. $[\alpha]_D^{22}$ -46 (c, 0.5 in CHCl₃).

Me ether, perchlorate: Mp 229.5-230.5° dec.

6α-Hydroxy: 6α-Hydroxycrinamidine

$C_{17}H_{19}NO_6$ 333.34

Alkaloid from the bulbs of *Ammod-charis tinneana*. Mp 254-256°. $[\alpha]_D^{20}$ +26 (c, 0.45 in MeOH). The 6-OH is *trans*-to the ethylene bridge (6*R*).

6α-Hydroxy, O³-Me: 6-Hydroxyundulatin

$C_{18}H_{21}NO_6$ 347.367

Alkaloid from the bulbs of *Ammod-charis tinneana*. Mp 113-116°. $[\alpha]_D^{20}$ +8.4 (c, 0.53 in MeOH).

6ξ-Hydroxy: 6-Hydroxycrinamidine

$C_{17}H_{19}NO_6$ 333.34

Alkaloid from *Crinum latifolium*. Very probably identical to the 6α-epimer above.

1,2-Diepimer, 3-ketone: Tubispacine

$C_{17}H_{17}NO_5$ 315.325

Alkaloid from the bulbs of *Zephyranthes tubispatha* (Amaryllidaceae). Fine prisms (Me₂CO). Mp 197-199°. $[\alpha]_D^{24}$ -145 (c, 0.3 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1954, **87**, 1704 (*isol*)

Warnhoff, E.W. *et al.*, *J.A.C.S.*, 1960, **82**, 1472 (*Undulatin*)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 498 (*isol*)

Fales, H.M. *et al.*, *J.O.C.*, 1961, **26**, 181 (*struct*)

Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1965, **298**, 704-708 (*Tubispacine*)

Zetta, L. *et al.*, *J.C.S. Perkin 2*, 1973, 1180 (*cmr*)

Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 961-965 (*Undulatin*)

Viladomat, F. *et al.*, *Phytochemistry*, 1996, **43**, 1379 (*cmr*)


Vo, T.B.H. *et al.*, *CA*, 1998, **128**, 292716n (6-Hydroxycrinamidine)

Machocho, A. *et al.*, *Phytochemistry*, 1999, **51**, 1185-1191 (6-Hydroxycrinamidine, 6-Hydroxyundulatin)

Crinasiadine

C-744

[1,3]Dioxolo[4,5-*j*]phenanthridin-6(5H)-one, 9*CI*. 8,9-Methylenedioxy-6-phenanthridone
[40141-86-4]



$C_{14}H_9NO_3$ 239.23

Alkaloid from the flowering bulbs of *Crinum asiaticum* (Amaryllidaceae). Exhibits bacteriostatic and tumour-inhibiting activity. Cryst. (DMF). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 333° (276-278° dec.). λ_{max} 248 (log ε 4.33); 252 (log ε 4.3); 268 (log ε 4.01); 285 (log ε 4.03); 298 (log ε 3.98); 309 (log ε 3.88); 324 (log ε 3.62); 338 (log ε 3.52) (MeOH).

N-Me: N-Methylcrinasiadine

[40141-98-8]

$C_{15}H_{11}NO_3$ 253.257

Alkaloid from bulbs and leaves of *Lapiedra martinii* (Amaryllidaceae). Yellow cryst. (Me₂CO). Mp 250-255° (244-245°). λ_{max} 240 (log ε 3.37); 284 (log ε 3.39); 294 (log ε 3.39); 300 (log ε 3.36); 315 (log ε 3.34); 340 (log ε 3.15) (MeOH).

N-(3-Methoxycarbonylpropyl): Phamine

[167033-99-0]

$C_{19}H_{17}NO_5$ 339.347

Alkaloid from bulbs of *Hippeastrum equestre*. Needles (Me₂CO). Mp 142-143°.

4-Hydroxy: Arolycoricidine

[39954-31-9]

$C_{14}H_9NO_4$ 255.229

Alkaloid from the bulbs of *Lycoris sanguinea* (Amaryllidaceae). Mp 295° dec. (281-283°). Prob. artifact of extraction.

4-Acetoxy:

Cryst. (DMF/MeOH). Mp 297-298°.

4-Methoxy: Mp 292-300° dec.**4,8-Dihydroxy: Narciprimine. Arolycoricidinol**

[28233-43-4]

$C_{14}H_9NO_5$ 271.229

Alkaloid from the bulbs of *Lycoris sanguinea* and *Narcissus* spp. (Amaryllidaceae). Small yellowish needles (AcOH or EtOH). Mp 300-320° dec. Artifact prod. from Narciclasine, N-37 during acid extraction. λ_{max} 233 (log ε 4.38); 257 (log ε 4.72); 274 (log ε 4.24); 295 (log ε 3.96); 322 (log ε 3.67); 337 (log ε 3.86); 353 (log ε 3.89) (EtOH).

4,8-Dihydroxy, di-O-Ac:


Cryst. (MeOH). Mp 247°.

4,8-Dimethoxy:
Amorph. Mp 320°.

- Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 1860-1864 (*Narciprimine, synth*)
 Piozzi, F. *et al.*, *Tetrahedron*, 1968, **24**, 1119-1131 (*Narciprimine, isol, uv, ir, pmr*)
 Mondon, A. *et al.*, *Chem. Ber.*, 1970, **103**, 2729-2743; 1972, **105**, 3726-3747 (*Narciprimine, synth, uv, ir, pmr, ms*)
 Takagi, S. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 617-622; *C.A.*, **81**, 74924y (*Narciprimine, Arolycoricidine, isol*)
 Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1985, 100-101 (*isol, uv, ir, ms*)
 Suau, R. *et al.*, *Phytochemistry*, 1990, **29**, 1710-1712 (*N-Methylcrinasiatine*)
 Banwell, M.G. *et al.*, *Aust. J. Chem.*, 1994, **47**, 2235-2254 (*synth*)
 Banwell, M.G. *et al.*, *Chem. Comm.*, 1995, 2551-2553 (*N-Methylcrinasiatine, synth*)
 Döpke, W. *et al.*, *Planta Med.*, 1995, **61**, 564-566 (*Phamine*)
 Shao, H.W. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 493-496 (*synth*)

Crinasiatine

C-745



$C_{22}H_{17}NO_4$ 359.381


(±)-form [97682-69-4]

Alkaloid from the flowering bulbs of *Crinum asiaticum* (Amaryllidaceae). Exhibits bacteriostatic and tumour-inhibiting activity. Light brown powder. Sol. MeOH, EtOAc. Dec. at ca. 270° without melting. λ_{max} 233; 249; 264; 282; 295; 305; 335 (MeOH). λ_{max} 245; 251; 267; 294; 308; 324; 338; 345 (MeOH/NaOH). O-Ac: Prisms (MeOH/Me₂CO). Mp 245-248° dec.

Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1985, 100 (*isol, uv, ir, pmr, ms, struct*)

Crinatine†

C-746



$C_{16}H_{19}NO_5$ 305.33

Not the same as Crinatine, C-747 but they are given the same CAS no. Alkaloid isol. from *Crinum oliganthum*. Trimino, Z. *et al.*, *Rev. Cubana Quim.*, 1997, **9**, 119-122 (*Crinatine*)

Crinatine†

C-747

$C_{18}H_{21}NO_5$ 331.368

Amaryllidaceae alkaloid. Struct. unknown. No further reports to 2007. Isol.


from the bulbs of *Crinum natans* (Amaryllidaceae). Mp 262°.

Onyiriuka, O.S. *et al.*, *Isr. J. Chem.*, 1978, **17**, 185-192 (*isol, ms*)

Crinine†

C-748

Vittatine. Crinidine



$C_{16}H_{17}NO_3$ 271.315

(+)-form [510-69-0]

Alkaloid from *Hippeastrum vittatum* and a number of other spp. in the Amaryllidaceae. Shows weak analgesic activity in mice. Tachycardic agent in dogs. Mp 207-208°. $[\alpha]_D^{25} +26$ (c, 0.5 in CHCl₃) (+38). ▶ Toxic, LD₅₀ 10 mg/kg (dog).

Me ether: (+)-Buphanisine

[124093-57-8]

$C_{17}H_{19}NO_3$ 285.342

Alkaloid from *Sternbergia sicula* (Amaryllidaceae). Amorph. $[\alpha]_D +18$ (c, 2.1 in CHCl₃).

6-Hydroxy, O³-Me: 6-Hydroxy-(+)-buphanisine

[80665-69-6]

[80665-70-9]

$C_{17}H_{19}NO_4$ 301.341

Alkaloid from the bulbs of *Pancreatium sickenbergeri*. Amorph. yellow solid. $[\alpha]_D -6.2$ (c, 0.77 in MeOH). Isol. as a mixt. of 6-epimers. λ_{max} 221 (log ϵ 2.04); 249 (log ϵ 0.67); 304 (log ϵ 1.09) (MeOH).

3-Epimer: Epicrinine. Epivittatine

[25375-48-8]

Alkaloid from the bulbs of *Nerine bowdenii* and *Crinum erubescens* (Amaryllidaceae). Also isol. from *Boophone flava* bulbs (Amaryllidaceae). Cryst. (CHCl₃/Me₂CO or by subl.). Mp 209-210°. $[\alpha]_D^{22} +136$. $[\alpha]_{436}^{22} +329$ (c, 1.03 in CHCl₃).

3-Epimer, Me ether: Epibuphanisine

[6835-79-6]

$C_{17}H_{19}NO_3$ 285.342

Alkaloid from the bulbs of *Ammocharis coranica* and *Boophone flava* (Amaryllidaceae). Prisms (Et₂O). Mp 123-125°. $[\alpha]_D^{21} +133$ (c, 0.219 in EtOH). $[\alpha]_D^{21} +141$ (c, 0.266 in CHCl₃).

3-Epimer, Me ether, perchlorate:

Prisms (Me₂CO). Mp 244-246° dec.

$[\alpha]_D^{21} +88$ (c, 0.211 in EtOH).

(-)-form [510-67-8]

Alkaloid from *Crinum moorei*, several other *Crinum* spp. and from several other spp. in the Amaryllidaceae. Mp 208-210°. $[\alpha]_D^{20} -23$ (c, 0.5 in CHCl₃). λ_{max} 240 (log ϵ 3.52); 296 (log ϵ 3.72) (EtOH).

Perchlorate: Mp 135-137°.

Ac: Krepowine

[93452-26-7]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Crinum powelli* var.

krelagei and from the bulbs of *Crinum bulbispermum*. Mp 145-146° (138-139°). $[\alpha]_D^{20} +68$ (c, 0.11 in EtOH).

Me ether: (-)-Buphanisine

[468-22-4]

$C_{17}H_{19}NO_3$ 285.342

Alkaloid from the bulbs of *Boophone fischeri* (Amaryllidaceae). Prisms (Et₂O). Mp 122-124°. $[\alpha]_D^{20} -26$ (c, 0.76 in EtOH).

3-Ketone: Oxocrinine

[31106-06-6]

$C_{16}H_{15}NO_3$ 269.299

Alkaloid from the bulbs of *Crinum americanum*. Intermed. in the biosynth. of Amaryllidaceae alkaloids. Needles (MeOH). Mp 185-186°. $[\alpha]_D -24.6$ (c, 1.0 in CHCl₃).

1,2-Dihydro: Elwesine. Crinan-3-ol. Dihydrocrinine

[10438-97-8]

$C_{16}H_{19}NO_3$ 273.331

Alkaloid from the bulbs of *Galanthus elwesii* (Amaryllidaceae). Mp 218-219°. $[\alpha]_D^{25} -32$ (c, 0.4 in CHCl₃). $[\alpha]_D -21$ (c, 0.7 in CHCl₃) (synthetic).

1,2-Dihydro, Ac: Mp 149-150°.

1β,2β-Epoxyde: Flexinine

[509-88-6]

$C_{16}H_{17}NO_4$ 287.315

Alkaloid from the bulbs of *Nerine flexuosa*, *Crinum amabile* and *Crinum erubescens* (Amaryllidaceae). Long prisms (EtOH). Mp 232-234° (221-222°). $[\alpha]_D^{25} -12.7$ (c, 0.82 in CHCl₃). Epoxide function is *cis*- to the bridge and *trans*- to the OH-group. λ_{max} 205 (log ϵ 4.27); 240 (sh) (log ϵ 3.49); 294 (log ϵ 3.64) (MeOH).

1β,2β-Epoxyde, perchlorate:

Needles (H₂O). Mp 250° dec.

1β,2β-Epoxyde, Ac:

Cryst. (MeOH). Mp 206-207°.

1β,2β-Epoxyde, Me ether: Augustine

[79659-60-2]

$C_{17}H_{19}NO_4$ 301.341

Alkaloid from leaves, bulbs and roots of *Crinum augustum*. Prisms (C₆H₆). Mp 174-176°. $[\alpha]_D^{25} -46.25$ (c, 0.8 in EtOH). λ_{max} 205 (log ϵ 4.34); 239 (log ϵ 3.56); 295 (log ϵ 3.74) (MeOH).

6-Hydroxy: 6-Hydroxycrinine

[80665-68-5]

[80665-67-4]

$C_{16}H_{17}NO_4$ 287.315

Alkaloid from *Crinum augustum* (Amaryllidaceae). Prisms (MeOH). Mp 268-270°. $[\alpha]_D^{25} +13.58$ (c, 0.5 in EtOH). Isol. as an inseparable mixt. of C6-epimers.

6-Hydroxy, O³-Me: 6-Hydroxy(-)-buphanisine

[80665-69-6, 80665-70-9]

$C_{17}H_{19}NO_4$ 301.341

Isol. from *Crinum augustum* (Amaryllidaceae). Prisms (MeOH). Mp 126-128°. $[\alpha]_D^{25} +40.04$ (c, 0.5 in EtOH). Insep. mixt. of 6-epimers.

3-Epimer:

Synthetic. Cryst. (Me₂CO). Mp 209-209.5°. $[\alpha]_D^{27} -142$. $[\alpha]_{436}^{27} -343$ (c, 1.03 in CHCl₃).

3-Epimer, Me ether

Synthetic. Prisms (Et₂O). Mp 124-126°. [α]_D²² -139 (c, 0.168 in CHCl₃).

3-Epimer, 1,2-dihydro: Epielwesine

[10279-81-9]
Synthetic. Prisms (Me₂CO aq.). Mp 101-103°. [α]_D -10.6 (c, 0.7 in CHCl₃). Was stated to be a minor alkaloid of *G. elwesii*, prob. incorrectly (Mp 187-188.5°, opt. inactive).

(±)-form [7679-55-2]

Synthetic. Mp 173-175°.

Me ether: [111958-06-6]

Synthetic. Cryst. (hexane). Mp 100-101.5°.

3-Ketone: [7700-86-9]

Synthetic. Powder (EtOAc). Mp 170-172°.

1,2-Dihydro: [33531-72-5]

Synthetic. Prisms (Me₂CO). Mp 227-230° (216-220°).

3-Epimer: Synthetic. Mp 239° (235.5-237°).**3-Epimer, 1,2-dihydro: [32209-87-3]**

Synthetic. Cryst. (C₆H₆/cyclohexane). Mp 187-188.5° (182-184°).

Boit, H.G. *et al.*, *Chem. Ber.*, 1954, **87**, 1704-1707; 1957, **90**, 369-373 ((-)-form, *Flexinine, isol*)

Renz, J. *et al.*, *Helv. Chim. Acta*, 1955, **38**, 1209-1222 (*Buphanisine*)

Mason, L.H. *et al.*, *J.A.C.S.*, 1955, **77**, 1253-1256 ((-)-form, *isol, uv*)

Wildman, W.C. *et al.*, *J.A.C.S.*, 1956, **78**, 4180-4181; 1958, **80**, 2567-2575 (*Epicrinine, synth, struct*)

Fales, H.M. *et al.*, *J.A.C.S.*, 1960, **82**, 3368-3375 (*Buphanisine, struct*)

Jeffs, P.W. *et al.*, *J.C.S.*, 1960, 1090-1094

(*config, Buphanisine, Epicrinine*)

Fales, H.M. *et al.*, *J.O.C.*, 1961, **26**, 181-193

(*Flexinine, struct*)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1961, **48**, 406 (*Elwesine, Krepowine, isol, struct*)

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 1307-1317 (*Epibuphanisine, isol, struct, synth*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902-4912 (*ms*)

Muxfeldt, H. *et al.*, *J.A.C.S.*, 1966, **88**, 3670-3671 (*synth*)

Whitlock, H.W. *et al.*, *J.A.C.S.*, 1967, **89**, 3600-3606 (*synth*)

Wildman, W.C. *et al.*, *Pharmazie*, 1967, **22**, 725; *CA*, **69**, 19337m (*Epicrinine, Flexinine*)

Irie, H. *et al.*, *J.C.S. (C)*, 1968, 1802-1804

(*Elwesine, Epicrinine, synth*)

DeAngelis, G.G. *et al.*, *Tetrahedron*, 1968, **24**, 5469-5481 (*ord, cd*)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 1488-1491 (*Epicrinine, synth*)

Zetta, L. *et al.*, *J.C.S. Perkin 2*, 1973, 1180-1184 (*cmr*)

Ali, A.A. *et al.*, *Phytochemistry*, 1981, **20**, 1121-1123; 1731-1733; 1986, **25**, 2399-2401

(*Augustine, 6-Hydroxycrinine, 6-Hydroxybuphanisine, Oxocrinine*)

Frahm, A.W. *et al.*, *Phytochemistry*, 1981, **20**, 1735-1738 (*Augustine, struct, pmr, cmr*)

Sanchez, I.H. *et al.*, *J.A.C.S.*, 1983, **105**, 7640-7643 (*Oxocrinine, Elwesine, Epielwesine, synth*)

Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3015-3022 (*isol, ir, pmr, Krepowine*)

Overman, L.E. *et al.*, *Helv. Chim. Acta*, 1985, **68**, 745-749 (*synth*)

Martin, S.F. *et al.*, *J.O.C.*, 1988, **53**, 3184-3190 (*synth, Buphanisine, pmr, cmr, ms*)

Pabuçcuoglu, V. *et al.*, *J. Nat. Prod.*, 1989, **52**, 785-791 ((+)-*Buphanisine*)

Ishibashi, H. *et al.*, *J.O.C.*, 1991, **56**, 95-102

(*Elwesine, Epielwesine, synth*)

Burk, R.M. *et al.*, *Heterocycles*, 1993, **35**, 205-225 (*Epielwesine, synth*)

Matsumura, Y. *et al.*, *Tetrahedron*, 1993, **49**, 8503-8512 (*Elwesine, synth*)

Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 307-311; 961-965 (*Crinine, Epicrinine, Buphanisine, Epibuphanisine*)

Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591-6600 (*cd*)

Pearson, W.H. *et al.*, *J.O.C.*, 1998, **63**, 3607-3617 (*synth*)

Pham, L.H. *et al.*, *Phytochemistry*, 1998, **48**, 371-376 (*Crinine, Augustine, Buphanisine, Flexinine*)

Machocho, A. *et al.*, *Phytochemistry*, 1999, **51**, 1185-1191 (*Flexinine, isol, ir, cd, pmr, cmr, ms*)

Abou-Donia, A.H. *et al.*, *Planta Med.*, 2002, **68**, 379-381 (*6-Hydroxy-(+)-buphanisine*)

Kodama, S. *et al.*, *Tetrahedron*, 2004, **60**, 4901-4907 (*Epicrinine, Oxocrinine, synth*)


Bru, C. *et al.*, *Tetrahedron*, 2006, **62**, 9043-9048 (*synth*)

Bohno, M. *et al.*, *Tetrahedron*, 2007, **63**, 6977-6989 (*synth*)

Tam, N.T. *et al.*, *J.O.C.*, 2008, **73**, 6258-6264 (*synth*)

Crisinine

[161068-63-9]



C₁₇H₂₁NO₆ 335.356

Alkaloid from bulbs of *Crinum asiaticum* var. *sinicum*.

Tang, R.J. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 855 (*isol*)

Lewis, J.R. *et al.*, *Nat. Prod. Rep.*, 1996, **13**, 171

Crisinosine


C₁₇H₂₁NO₄ 303.357

Amaryllidaceae alkaloid. Struct. unknown. Isol. from an unidentified variety of *Crinum powellii* (Amaryllidaceae). Mp 195°. [α]_D²¹ +210 (c, 0.2 in EtOH).

Döpke, W. *et al.*, *Pharmazie*, 1965, **20**, 586; *CA*, **63**, 18649g (*isol*)

Criophylline

[52659-53-7]



Relative Configuration

C₄₀H₄₆N₄O₄ 646.828

The position of the interunit link is not proven. Major alkaloid from the leaves of *Crioceras dipladeniiflorus* (Apocynaceae). Cryst. (Me₂CO or MeOH). Mp 276-279° dec. [α]_D -176 (c, 1.04 in CHCl₃). λ_{\max} 214 (log ϵ 4.26); 228 (log ϵ 4.08); 257 (log ϵ 3.93); 299 (log ϵ 4.06); 327 (log ϵ 4.28) (EtOH).

De(methoxycarbonyl): Mp 260°. [α]_D -63 (c, 0.74 in CHCl₃).


Cavé, A. *et al.*, *Tet. Lett.*, 1973, 5081 (*uv, pmr, cmr, struct*)

Bruneton, J. *et al.*, *Phytochemistry*, 1974, **13**, 1963-1967 (*isol, uv, ms*)

Cripowellin A

C-752

[196814-62-7]



C₂₅H₃₁NO₁₂ 537.519


Alkaloid from the bulbs of *Crinum powellii*. Powerful insecticidal agent. Amorph. solid. [α]_D -44 (c, 1 in MeOH). λ_{\max} 206 ; 292 (heptane/EtOH).

Velten, R. *et al.*, *Tet. Lett.*, 1998, **39**, 1737-1740 (*isol, uv, ir, pmr, cmr, ms*)

Cripowellin B

C-753

[196812-55-2]



C₂₅H₃₃NO₁₁ 523.536

Alkaloid from the bulbs of *Crinum powellii*. Powerful insecticidal agent. Amorph. solid. [α]_D²⁰ -64 (c, 1 in MeOH). λ_{\max} 206 ; 292 (heptane/EtOH).

Velten, R. *et al.*, *Tet. Lett.*, 1998, **39**, 1737-1740 (*isol, uv, ir, pmr, cmr, ms*)

Crispanine

C-754

Alkaloid NB

[1356-13-4]

C₁₈H₂₁NO₅ 331.368

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Nerine crispa* and *Nerine bowdenii* (Amaryllidaceae). Mp 139-141°. [α]_D²⁴ +78 (c, 0.2 in CHCl₃).

Perchlorate: Mp 221° dec.

Picrate: Mp 201° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109; 1962, **49**, 469 (*isol, ir*)

Crispine†

C-755

C₁₈H₂₃NO₆ 349.383

Amaryllidaceae alkaloid. Struct. unknown. *Isol.* from the bulbs of *Nerine undulata* (Amaryllidaceae). Cryst. (Me₂CO). Mp 275° dec. [α]_D²² -96 (c, 0.25 in CHCl₃).

Perchlorate:

Prisms. Mp 268-269° dec.

Picrate: Mp 258° dec.

Methoperchlorate:


Cryst. (H₂O). Mp 282-283° dec.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129-1134 (*isol*)

Crispine A†

C-756

1,2,3,5,6,10b-Hexahydro-8,9-dimethoxy-pyrrolo[2,1-a]isoquinoline



(R)-form

C₁₄H₁₉NO₂ 233.31

(R)-form [474304-16-0]

Alkaloid from *Carduus crispus*. Needles. Mp 87-89°. [α]_D²⁵ +91 (MeOH).

N,5,6,10b-Tetrahydro-2,3-Dihydro-8,9-dimethoxy-1H-pyrrolo[2,1-a]isoquinolinium. *Carcrisine A*. **Crispine B** [474304-17-1]

C₁₄H₁₆NO₂⁺ 230.286

Quaternary alkaloid from *Carduus crispus*. Cytotoxic. Plates (as chloride). Mp 213-215° (206-208°)(chloride). λ_{max} 237 ; 260 ; 308 (MeOH) (chloride).

(±)-form

Cryst. (Et₂O/petrol). Mp 89°.

Orito, K. *et al.*, *Heterocycles*, 1988, **27**, 2403-2412 (*synth, pmr*)

Zhang, Q. *et al.*, *Tetrahedron*, 2002, **58**, 6795-6798 (*isol, pmr, cmr*)

Xie, W.D. *et al.*, *Chin. Chem. Lett.*, 2004, **15**, 1057-1059 (*Carcrisine A*)

Meyer, N. *et al.*, *Eur. J. Org. Chem.*, 2006, 3997-4002 (*synth*)

Allin, S.M. *et al.*, *J.O.C.*, 2007, **72**, 8972-8975 (*synth*)


King, F.D. *et al.*, *Tetrahedron*, 2007, **63**, 2053-2056 (*synth*)

Szawalko, J. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 406-413 (*synth, cryst struct, abs config*)

Crispine C

C-757

Carcrisine B
[474304-18-2]



R = H

C₁₅H₂₀N₄O₂ 288.349

Alkaloid from *Carduus crispus*. Light yellow cryst. (MeOH). Mp 208-210°. λ_{max} 230 ; 266 ; 278 ; 312 ; 325 (MeOH).

Zhang, Q. *et al.*, *Tetrahedron*, 2002, **58**, 6795-6798 (*isol, pmr, cmr, ms*)

Xie, W.D. *et al.*, *Chin. Chem. Lett.*, 2004, **15**, 1057-1059 (*isol, pmr, cmr*)

Crispine D

C-758

[474304-19-3]

As Crispine C, C-757 with

R = -(CH₂)₄NHC(NH₂)=NH

C₂₀H₃₁N₇O₂ 401.511


Alkaloid from *Carduus crispus*. Needles. Mp 107-110°.

Zhang, Q. *et al.*, *Tetrahedron*, 2002, **58**, 6795-6798 (*isol, pmr, cmr*)

Crispine E

C-759

[474304-20-6]



C₁₅H₂₄N₄O₂ 292.38


Alkaloid from *Carduus crispus*. Needles. Mp 130-133°.

Zhang, Q. *et al.*, *Tetrahedron*, 2002, **58**, 6795-6798 (*isol, pmr, cmr*)

Czarnocki, S.J. *et al.*, *Tetrahedron*, 2008, **64**, 3176-3182 (*synth*)

Croaegyptine

C-760



C₁₇H₂₅NO₆ 339.388


Isol. as an insep. mixt. with Crosemperine, C-769. Alkaloid from *Crotalaria aegyptiaca* (Fabaceae).

Roeder, E. *et al.*, *Phytochemistry*, 1993, **34**, 1421

Croalbidine

C-761

[41714-30-1]



C₁₈H₂₉NO₇ 371.43

Absolute configuration

Cyclic diester of the rare base croalbinecine (see 2,7-Dihydroxy-1-hydroxy-methylpyrrolizidine, D-597) with (+)-trichodesmic acid. Alkaloid from *Crotalaria albida* (Fabaceae). Cryst. (EtOH or Py). Mp 208-209°.

Hydrochloride: Mp 154-155°.

Picrate: Mp 224-225°.

Sawhney, R.S. *et al.*, *Indian J. Chem.*, 1973, **11**, 88 (*isol*)


Sawhney, R.S. *et al.*, *Aust. J. Chem.*, 1974, **27**, 1805 (*abs config*)

Crobarbatine

C-762

12-Hydroxy-19,20-dinorcrotalanan-11,15-dione, 9CI

[49679-23-4]



Probable absolute configuration

C₁₅H₂₁NO₅ 295.335

Cyclic ester of Retronecine in T-188 with crobarbatic acid (unique to this alkaloid). Stereochem. of the necic acid component is tentative. Alkaloid from *Crotalaria barbata* (Fabaceae). Cryst. (C₆H₆). Mp 142-143°.

Picrate: Mp 237°.

Puri, S.C. *et al.*, *Experientia*, 1973, **29**, 390 (*isol, pmr, ms, struct*)


Huang, J. *et al.*, *J.A.C.S.*, 1981, **103**, 861 (*synth*)

Jang, D.-P. *et al.*, *Org. Lett.*, 2001, **3**, 983-985 (*Crobarbatic acid, synth*)

Crocacin A

C-763

Methyl N-[6-[(7,9-dimethoxy-3,6,8-trimethyl-1-oxo-11-phenyl-2,4,10-undecatrienyl)amino]-1-oxo-2,5-hexadienyl]glycinate, 9CI
[157698-34-5]



Relative Configuration

C₃₁H₄₂N₂O₆ 538.683

Prod. by *Chondromyces crocatus*. Inhibits a wide spectrum of yeasts and molds and a few gram-positive bacteria. Sol. MeOH, CHCl₃, EtOAc, Me₂CO; fairly sol. Et₂O; poorly sol. hexane, H₂O. [α]_D²⁰ +109.6 (c, 1 in MeOH). λ_{max} 213 ; 219 (sh) ; 254 (log ε 4.54) ; 261 ; 275 (sh) ; 282 (log ε 4.36) ; 291 (log ε 4.33) ; 298 (sh) (MeOH). λ_{max} 213 ; 254 (ε 38994) ; 261 ; 282 (ε 23014) ; 291 (ε 21429) (MeOH) (Berdy).


Parent acid: **Crocacin B**

C₃₀H₄₀N₂O₆ 524.656

Prod. by *Chondromyces crocatus*. [α]_D²² +99 (c, 0.5 in MeOH). λ_{max} 254 (log ε 4.48) ; 266 ; 276 (sh) ; 283 (log ε 4.31) ; 293 ; 300 (sh) (MeOH).

5,6-Dihydro- Crocacin DC₃₁H₄₄N₂O₆ 540.698Prod. by *Chondromyces pediculatus*.[α]_D²² +109.6 (c, 0.6 in MeOH). λ_{max} 255 (log ε 4.55); 262; 276; 280; 290 (sh) (MeOH).**N¹⁰-De(aminoalkyl)- Crocacin C**C₂₂H₃₁NO₃ 357.492Prod. by *Chondromyces crocatus*. [α]_D²² +52.2 (c, 0.3 in MeOH). λ_{max} 254 (log ε 4.64); 266; 277; 284; 293 (sh) (MeOH).Kunze, B. *et al.*, *J. Antibiot.*, 1994, **47**, 881-886 (*isol, uv, ir, props*)*Ger. Pat.*, 1995, 4 324 672; *CA*, **122**, 237917eJansen, R. *et al.*, *Eur. J. Org. Chem.*, 1999,1085-1089 (*isol, uv, ir, pmr, cmr, ms*)Dias, L.C. *et al.*, *Org. Lett.*, 2001, **3**, 3951-3954(*Crocacin C, synth*)Chakraborty, T.K. *et al.*, *Tetrahedron*, 2001,**57**, 9461-9467 (*Crocacin C, synth*)Chakraborty, T.K. *et al.*, *Tet. Lett.*, 2002,**43**, 2645-2648; 2003, **44**, 4989-4992(*synth*)Dias, L.C. *et al.*, *J.O.C.*, 2005, **70**, 2225-2234(*Crocacin D, synth*)Yadav, J.S. *et al.*, *Tet. Lett.*, 2007, **48**, 145-148(*Crocacin C, synth*)Sirasani, G. *et al.*, *J.O.C.*, 2008, **73**, 6386-6388(*Crocacin C, synth*)De Oliveira, L.G. *et al.*, *Quim. Nova*, 2008, **31**,854-871 (*rev, synth*)Feutrill, J.T. *et al.*, *Tetrahedron*, 2008, **64**, 4880-4895 (*synth*)**Crocandine**

[72855-83-5]

C₁₆H₂₅NO₅ 311.377Cyclic diester of Turneforicidine in T-188 with fulvic acid. Alkaloid from the seeds of *Crotalaria candidans* (Fabaceae). Needles (MeOH). Mp 244-246°. [α]_D +130 (c, 1% in MeOH).**Stereoisomer: Isocrocandine**

[72903-70-9]


C₁₆H₂₅NO₅ 311.377Alkaloid from seeds of *Crotalaria candidans* (Fabaceae). Cryst. (EtOAc). Mp 172-174°. [α]_D +36 (c, 1 in MeOH). Cyclic diester of Turneforicidine in T-188 with Chromaduric acid.Siddiqi, M.A. *et al.*, *Phytochemistry*, 1979, **18**, 1413 (*isol, ir, pmr, ms, struct*)**Croceocurine**

C-765


Struct. unknown

Prob. a strychnos-type alkaloid. Alkaloid from the bark of an unidentified South American plant (prob. *Viburnaceae*). Bright-orange. Forms a cryst. picrate (no Mp reported).Meyer, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1214-1218 (*isol, uv*)**Crociflorinine**

Krokiflorinine

C₂₂H₂₇NO₅ 385.459Alkaloid from *Colchicum kesselringii*.Mp 209-210°. [α]_D -205.**N-Oxide: Crociflorine. Krokiflorine**C₂₂H₂₇NO₆ 401.458Alkaloid from *Colchicum kesselringii*.Mp 229-230°. [α]_D -255 (CHCl₃).**Methiodide:** Mp 230°.Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**,410-512; 761-863; *Chem. Nat. Compd. (Engl.**Transl.*), 1996, **32**, 386-512; 737-858 (*rev*)**Cronaburmine**

C-767

14,19-Dihydro-12-(1-hydroxyethyl)-17,20-dinorcrotalaran-11,15-dione, 9CI
[71295-32-4]C₁₇H₂₅NO₅ 323.388Alkaloid from *Crotalaria nana* seeds (Fabaceae). Mp 133-134°. Hydrol. gives Retronecine in T-188 and cronaburmic acid (unique to this alkaloid).**Hydrochloride:** [71327-59-8]


Mp 250°.

Picrate: [71327-60-1]

Mp 233° dec.

Siddiqi, M.A. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 1132 (*isol, ms, pmr, struct*)**Cropodine**


[83601-85-8]

C₁₆H₂₅NO₆ 327.377Cyclic diester of Turneforicidine with monocrotalic acid. Alkaloid from the pericarps of *Crotalaria candidans* (Fabaceae). Cryst. (Me₂CO). Mp 226-228°. [α]_D²⁵ +70 (c, 0.4 in MeOH).Haksar, C.N. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 492 (*isol, ir, pmr, ms, struct*)

C-768

Crosemperine


C-769

14,19-Dihydro-8,12-dihydroxy-4,19-dimethyl-11,15-dioxocrotalaranium, 9CI
[30785-56-9]C₁₉H₂₉NO₆ 367.441Cyclic diester of Otonecine, O-136 with incanic acid, found formerly in *Incanine* in T-488. Alkaloid from *Crotalaria semperflorens* (Fabaceae). Mp 117-118°. [α]_D²⁹ +45 (CHCl₃). [α]_D²⁰ +2.2 (c, 1.5 in EtOH).▶ Toxic, LD₅₀ 32 mg/kg. GP9030800**Hydrochloride:** Mp 180° dec.**Methiodide:** Mp 208-209° dec.


[15401-71-5]

Atal, C.K. *et al.*, *Aust. J. Chem.*, 1967, **20**, 805 (*isol, pmr*)Sharma, R.K. *et al.*, *J. Med. Chem.*, 1968, **11**, 620 (*pharmacol, tox*)**Crotafoline**

C-770

12-Hydroxy-14-methyl-18-norsenecionan-8,11,16-trione, 9CI
[38494-87-0]C₁₈H₂₅NO₆ 351.399Otonecine cyclic diester. Alkaloid from *Crotalaria laburnifolia* (Fabaceae). Cryst. (Me₂CO). Mp 176-182° dec.Crout, D.H.G. *et al.*, *J.C.S. Perkin 1*, 1972, 1602 (*isol, pmr, ir, struct*)**Crotalarine**

C-771

14,19-Dihydro-12,13-dihydroxy-14-methylcrotalaran-11,15-dione, 9CI. Croburhine
[53937-97-6]

C₁₈H₂₇NO₆ 353.414

Cyclic ester of Retronecine in T-188 and Crotalaric acid. Alkaloid from *Crotalaria burhia* and *Crotalaria aegyptiaca* (Fabaceae). Cryst. (C₆H₆). Mp 167-168°. [α]_D^{18.5} -78 (c, 0.7 in EtOH). Evidence for the stereochem. of the necic acid portion is unconvincing. Crotalarine and Crobruhine do not appear to have been compared but their props. are identical.


Hydrochloride:Cryst. (EtOH/Et₂O). Mp 208-209°.**Methiodide:**Cryst. (MeOH/Et₂O). Mp 201-203°.**Picrate:**

Cryst. (EtOH). Mp 240° dec.

Ali, M.A. *et al.*, *Pak. J. Sci. Ind. Res.*, 1973, **16**, 227 (*isol, pmr, struct*)Rao, P.G. *et al.*, *Indian J. Chem.*, 1975, **13**, 835 (*isol, ms, pmr*)Roeder, E. *et al.*, *Phytochemistry*, 1993, **34**, 1421 (*isol, ir, pmr, cmr, ms, struct*)**Crotalarine lactone**

C-772

[152344-01-9]


C₁₈H₂₇NO₆ 353.414

Alkaloid from *Crotalaria aegyptiaca* (Fabaceae). Powder. [α]_D^{18.5} -12.25 (c, 0.25 in EtOH).

Roeder, E. *et al.*, *Phytochemistry*, 1993, **34**, 1421 (*isol, ir, pmr, cmr, ms, struct*)**Crotaleschenine**

C-773


[120154-95-2]

C₁₆H₂₃NO₅ 309.361

Erroneously identified as Crispatine when first isol. from the same source. Alkaloid from *Crotalaria leschenaultii* (Fabaceae). Mp 142-143°. [α]_D²⁰ -44.5 (EtOH). Adopts 2 different conformns. in the solid state.

Smith, L.W. *et al.*, *Aust. J. Chem.*, 1988, **41**, 429 (*pmr, cmr, ms, cryst struct*)**Crotananine**

[71295-28-8]


C₁₇H₂₅NO₅ 323.388

Struct. revised in 1993. Previously assigned a 12-membered macrocyclic diester struct. Alkaloid from *Crotalaria nana* (Fabaceae). Needles (EtOAc). Mp 174-175°. [α]_D²³ -80 (c, 1.02 in MeOH).

Siddiqui, M.A. *et al.*, *Phytochemistry*, 1978, **17**, 2143 (*isol, pmr, ms*)Sharma, S.D. *et al.*, *Cryst. Res. Technol.*, 1993, **28**, 945 (*cryst struct*)Robins, D.J. *et al.*, *Nat. Prod. Rep.*, 1995, **12**, 413**Crotanecine**

C-775

2,3,5,7a-Tetrahydro-7-(hydroxymethyl)-1H-pyrrolizine-1,2-diol, 9CI. 6,7-Dihydroxy-1-hydroxymethyl-1,2-dehydro-8 α -pyrrolizidine [5096-50-4]



Absolute Configuration

C₈H₁₃NO₃ 171.196

Necine base from Anacrotine, A-967 and Madurensine, M-36. Prisms (Et₂O/EtOH). Mp 202-203.5° (188-190°, 193-197°). [α]_D +34.5 (c, 0.055 in EtOH). Abs. config. derives from cryst. struct. studies of the alkaloids.

2-Epimer: Uspallatinecine

[98303-06-1]

C₈H₁₃NO₃ 171.196

Necine base from Uspallatine in A-967. CAS numbering.


Atal, C.K. *et al.*, *Tet. Lett.*, 1966, 537Mattocks, A.R. *et al.*, *J.C.S. (C)*, 1968, 325 (*synth*)Yadav, V.K. *et al.*, *Heterocycles*, 1984, **22**, 2735 (*synth*)Pestchanker, M.J. *et al.*, *Phytochemistry*, 1985, **24**, 1622-1624 (*Uspallatinecine*)Buchanan, J.G. *et al.*, *J.C.S. Perkin 1*, 1987, 2377 (*synth*)Bennett, R.B. *et al.*, *Tet. Lett.*, 1990, **31**, 5437 (*synth*)Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*pmr*)Denmark, S.E. *et al.*, *J.A.C.S.*, 1997, **119**, 125 (*synth*)

C-774

Crotonine†

C-776

1-[5-(2-Furanyl)pyrazinyl]-1,2,4-butanetriol. 2-Furanyl-5-(1,2,4-trihydroxybutyl)pyrazine [917370-00-4]


C₁₂H₁₄N₂O₄ 250.254

Alkaloid from the leaves of *Croton tiglium*. Analgesic agent. Yellow solid. [α]_D²⁰ -6.8 (c, 0.25 in MeOH). λ_{\max} 205 ; 280 ; 335 (MeOH).

Wu, X.-A. *et al.*, *J. Asian Nat. Prod. Res.*, 2007, **9**, 437-441 (*isol, pmr, cmr, ms*)**Cruentarene A**

C-777

[533925-98-3]



Absolute Configuration


C₃₃H₅₁NO₈ 589.768

Prod. by *Byssosvorax cruenta*. Antifungal agent. Cytotoxic. Amorph. solid. [α]_D²² -3.4 (c, 13.5 in MeOH). λ_{\max} 216 (log ϵ 4.26); 265 (log ϵ 3.98); 302 (log ϵ 3.66); 346 (log ϵ 1.94) (MeOH).

Jundt, L. *et al.*, *Eur. J. Org. Chem.*, 2006, 5036-5044 (*isol, pmr, cmr, cryst struct*)Kunze, B. *et al.*, *J. Antibiot.*, 2006, **59**, 664-668 (*isol, activity*)Vintonyak, V.V. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 5209-5211 (*synth*)Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 9275-9278 (*synth*)**Cruentarene B**

C-778

[533925-99-4]



Absolute Configuration

C₃₃H₅₁NO₈ 589.768

Prod. by *Byssosvorax cruenta*. Amorph. solid. [α]_D²² -9.1 (c, 6.6 in MeOH). λ_{\max} 210 (log ϵ 4.36); 215 (log ϵ 4.38); 267 (log ϵ 4.14); 301 (log ϵ 3.77) (MeOH).

Jundt, L. *et al.*, *Eur. J. Org. Chem.*, 2006, 5036-5044 (*isol, cd, pmr, cmr*)Kunze, B. *et al.*, *J. Antibiot.*, 2006, **59**, 664-668 (*isol*)Chakraborty, T.K. *et al.*, *J.O.C.*, 2008, **73**, 3578-3581 (*synth*)**Cruentine A†**

C-779

[1356-15-6]

C₁₈H₂₅NO₅ 335.399

Pyrrolizidine alkaloid. Struct. unknown.

Alkaloid from *Senecio cruentus* (Asteraceae). Mp 218-220°. $[\alpha]_D^{17.7}$ -94.1 (CHCl₃). On hydrol. gives retronecine and a necic acid, Mp 139-141°.

Methiodide: Mp 228-231°.

Picrate: Mp 185-187°.

Chu, Y.-L. *et al.*, *Yaoxue Xuebao*, 1964, **11**, 168-173; *CA*, **61**, 1904d

Cruentine B

C-780

[1356-16-7]

C₁₈H₂₅NO₆ 351.399

Pyrrrolizidine alkaloid. Struct. unknown. Alkaloid from *Senecio cruentus* (Asteraceae). Mp 200-202°. $[\alpha]_D^{17.6}$ -63.4 (CHCl₃). Hydrol. gives retronecine and a necic acid, Mp 177-179°.


Picrate: Mp 170-172°.

Chu, Y.-L. *et al.*, *Yaoxue Xuebao*, 1964, **11**, 168-173; *CA*, **61**, 1904d

Cryptochine

C-781

[147127-62-6]

C₁₉H₂₅NO₃ 315.411

Diastereoisomeric with Amuroline, A-957. Alkaloid from stem bark of *Cryptocarya chinensis* (Lauraceae). Amorph. solid. $[\alpha]_D^{24}$ +40 (c, 0.48 in MeOH).

*O*¹-*De-Me*: **1-O-Demethylcryptochine**. *1-Hydroxycryptochine*
C₁₈H₂₃NO₃ 301.385
Alkaloid from *Cryptocarya chinensis*. Needles (Me₂CO). Mp 117-119°. $[\alpha]_D$ +65.3 (c, 0.32 in CHCl₃). λ_{\max} 227 (sh) (log ϵ 3.83); 288 (log ϵ 3.27) (MeOH).

4-Ketone: Isoamuronine. *8,9-Dihydro-N-methylstepharine*

C₁₉H₂₃NO₃ 313.396

Alkaloid from *Cryptocarya chinensis*. Yellow powder. λ_{\max} 250 (sh) (log ϵ 3.22); 286 (log ϵ 3.75) (MeOH).

4-Ketone, N-de-Me: N-Demethylisamuronine. *8,9-Dihydrostepharine*

C₁₈H₂₁NO₃ 299.369

Alkaloid from *Cryptocarya chinensis*. Needles (MeOH). $[\alpha]_D$ +141.4 (c, 0.005 in MeOH). λ_{\max} 224 (log ϵ 3.64); 249 (sh) (log ϵ 3.45); 285 (log ϵ 3.15) (MeOH).

4-Ketone, O¹-De-Me: N-Methylcrotosparinine

[23179-62-6]

C₁₈H₂₁NO₃ 299.369

Alkaloid from *Croton sparsiflorus* (Euphorbiaceae). Cryst. (EtOAc). Mp 160-161°. $[\alpha]_D$ +244 (c, 0.92 in CHCl₃).

4-Ketone, O¹-de-Me, N-de-Me: Crotosparinine

[23179-61-5]

C₁₇H₁₉NO₃ 285.342

Alkaloid from *Croton sparsiflorus* (Euphorbiaceae). Cryst. (EtOAc). Mp 184-185°. $[\alpha]_D$ +215 (c, 2.37 in CHCl₃). λ_{\max} 228 (log ϵ 4.28); 285 (log ϵ 3.1) (no solvent reported).

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1970, **9**, 2573-2580; 1974, **13**, 2767-2769

(*Crotosparinine, N-Methylcrotosparinine, isol, uv, ir, pmr, struct, biosynth*)

Casagrande, C. *et al.*, *J.C.S. Perkin 1*, 1975, 1659-1663 (*stereochem*)

Colombo, A. *et al.*, *J.C.S. Perkin 2*, 1976, 1218-1221 (*N-Me, cryst struct*)


Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 227 (*Cryptochine, 1-O-Demethylcryptochine*)

Wu, T.S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1404-1407 (*Isoamuronine, N-Demethylisoamuronine*)

Cryptadine A

C-782

[959153-14-1]



Relative Configuration

C₃₀H₄₉N₃O 467.737

Alkaloid from *Lycopodium cryptomerinum*. Amorph. solid. $[\alpha]_D^{19}$ -50 (c, 0.8 in MeOH).

7-Epimer: Cryptadine B

[959153-15-2]

C₃₀H₄₉N₃O 467.737


Alkaloid from *Lycopodium cryptomerinum*. Amorph. solid. $[\alpha]_D$ -69 (c, 0.2 in MeOH).

Koyama, K. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 7803-7808 (*isol, cd, pmr, cmr*)

Cryptaustoline

C-783

5,6,12,12a-Tetrahydro-2-hydroxy-3,9,10-trimethoxy-7-methylindolo[2,1-a]isoquinolinium(I⁺), 9CI



(-)-form

C₂₀H₂₄NO₄⁺ 342.414

Abs. config. revised in 1992.

(-)-form [26754-21-2]

Alkaloid from the bark of *Cryptocarya bowiei* and *Cryptocarya oubatchensis* (Lauraceae). λ_{\max} 216 (log ϵ 4.72); 222 (sh) (log ϵ 4.21); 288 (log ϵ 4.03) (EtOH).

Chloride:

C₂₀H₂₄ClNO₄ 377.867

Amorph. $[\alpha]_D$ -150 (c, 0.45 in EtOH).

Iodide: [58939-31-4]

C₂₀H₂₄INO₄ 469.318

Prisms (MeOH). Mp 214° dec. $[\alpha]_D^{20}$ -151 (c, 0.4 in EtOH).

Me ether:

C₂₁H₂₆NO₄⁺ 356.441

Cryst. + 2H₂O (as iodide). Mp 153-155° (iodide). $[\alpha]_D^{20}$ -175 (c, 0.4 in EtOH).

*O*⁹-*De-Me: Litcubine*

[172924-22-0]

C₁₉H₂₂NO₄⁺ 328.387

Alkaloid from the roots of *Litsea cubeba* (mountain pepper). Amorph. solid (as perchlorate). $[\alpha]_D^{25.5}$ -111 (c, 1.0 in MeOH). Abs. config. incorrectly assigned in the paper, which ignores the 1992 revision of the abs. config. of Cryptaustoline. λ_{\max} 233 (sh) (log ϵ 4.06); 287 (log ϵ 3.96) (MeOH). λ_{\max} 249 (log ϵ 4.14); 304 (log ϵ 3.99) (MeOH + NaOH).

*O*¹⁰-*De-Me: Mangochinine*. *10-De-O-methylcryptaustoline*

[189686-29-1]

C₁₉H₂₂NO₄⁺ 328.387

Alkaloid from *Magnolia officinalis* and bark of *Manglietia chingii*. Light-grey needles (EtOH/Et₂O) (as hydroxide). Mp 162-164° (hydroxide). $[\alpha]_D$ -164 (c, 0.4 in MeOH) (hydroxide). λ_{\max} 211 (log ϵ 4.27); 287 (log ϵ 3.83) (MeOH) (hydroxide).

*O*⁹,*O*¹⁰-*Di-de-Me: Litcubinine*

[172924-24-2]

C₁₈H₂₀NO₄⁺ 314.36

Alkaloid from the roots of *Litsea cubeba* (mountain pepper). Amorph. solid (as perchlorate). $[\alpha]_D^{25.5}$ -144 (c, 0.5 in MeOH). Abs. config. incorrectly assigned, see above. λ_{\max} 234 (sh) (log ϵ 4.03); 290 (log ϵ 3.99) (MeOH). λ_{\max} 248 (log ϵ 4.1); 305 (log ϵ 4.08) (MeOH/NaOH).

(±)-form [17138-35-1]

[17138-16-8]

Prisms + ½H₂O (EtOH aq.) (as iodide). Mp 260° (242-243°, 254-255°) (iodide).

Me ether:

C₂₁H₂₆NO₄⁺ 356.441

Cryst. + ½H₂O (EtOH aq.) (as iodide). Mp 243-245° dec. (iodide).

Ewing, J. *et al.*, *Aust. J. Chem.*, 1953, **6**, 78-85 (*isol, struct*)

Hughes, G.K. *et al.*, *Aust. J. Chem.*, 1953, **6**, 315-317 (*synth*)

Kametani, T. *et al.*, *J.C.S. (C)*, 1967, 2208-2212 (*synth, ms*)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 1498-1502; 1973, **21**, 766-769 (*synth, pmr, ms, struct*)

Ninomiya, I. *et al.*, *Heterocycles*, 1977, **6**, 1855-1860 (*synth*)

Takano, S. *et al.*, *Heterocycles*, 1987, **26**, 1483-1485; 1487-1489 (*synth*)

Leboeuf, M. *et al.*, *Can. J. Chem.*, 1989, **67**, 947-952 (*isol, uv, pmr, ms*)

Yasuda, S. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1682-1683 (*synth*)

Meyers, A.I. *et al.*, *J.A.C.S.*, 1992, **114**, 8483-8489 (*abs config*)


Lee, S.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 80-82 (*Litcubine, Litcubinine*)

Moriyasu, M. *et al.*, *Nat. Med. (Tokyo)*, 1996, **50**, 413-416 (*10-Demethylcryptaustoline*)

Qiu, S.-X. *et al.*, *Tet. Lett.*, 1998, **39**, 4167-4170 (*Mangochinine*)

Cryptocin


[264913-57-7]

C₂₁H₃₁NO₄ 361.48

Tetramic acid deriv. Prod. by *Cryptosporiopsis* cf. *quercina*. Antimycotic agent. Cryst. (as Na salt). Dec. at 175-180° (Na salt).

Li, J.Y. *et al.*, *Org. Lett.*, 2000, **2**, 767-770**Cryptoheptine**

[160568-17-2]


C₁₇H₁₄N₂O 262.31

There are doubts about the proposed struct. Alkaloid from roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Exhibits antibacterial activity against gram-negative and gram-positive bacteria. Yellowish amorph. solid.

Paulo, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1485-1491 (*isol, uv, ir, pmr, cmr, ms, struct*)Zhang, P. *et al.*, *J. Nat. Prod.*, 2000, **63**, 643-645 (*struct*)**Cryptolepicarboline**

C-786

5-Methyl-11-(9H-pyrido[3,4-b]indol-9-yl)-5H-quindoline, 9CI
[171090-86-1]

C₂₇H₁₈N₄ 398.466


Alkaloid from roots of *Cryptolepis sanguinolenta*. Yellow amorph. solid. λ_{max} 225 (log ε 4.74); 275 (log ε 4.63); 282 (log ε 4.64); 308 (log ε 3.97); 341 (log ε 4); 357 (sh) (log ε 4.16); 373 (log ε 4.4); 394 (sh) (log ε 3.81); 450 (log ε 3.59) (MeOH).

Sharaf, M.H.M. *et al.*, *Magn. Reson. Chem.*, 1995, **33**, 767 (*isol, uv, ir, pmr, cmr, ms, struct*)

C-784

Cryptomisrine

Di-10H-quindolin-11-ylmethanone, 9CI
[158453-41-9]


C₃₁H₁₈N₄O 462.509

Alkaloid from roots of *Cryptolepis sanguinolenta*. Orange residue. λ_{max} 227 (log ε 4.85); 273 (log ε 4.9); 348 (log ε 4.19); 368 (log ε 4.18); 459 (log ε 3.94) (MeOH).

Sharaf, M.H.M. *et al.*, *J. Het. Chem.*, 1996, **33**, 789-797 (*isol, uv, ir, pmr, cmr, struct*)Arzel, E. *et al.*, *Tetrahedron*, 1999, **55**, 12149-12156 (*synth, ir, pmr, cmr*)**Cryptophorine**

C-788

6-(1,3,5,7-Decatrienyl)-1,2-dimethyl-3-piperidinol, 9CI
[56022-14-1]

C₁₇H₂₇NO 261.406

Alkaloid from *Bathiorhannus cryptophorus* (Rhamnaceae). Mp 116-118°. [α]_D²⁵ -61 (c, 1 in CHCl₃).

Ac: Mp 103-104°.

7',8'-Dihydro, Me ether: 6-(1,3,5-Decatrienyl)-3-methoxy-1,2-dimethylpiperidine


C₁₈H₃₁NO 277.449

Alkaloid from *Microcos paniculata*. Insecticidal agent. Cryst. (CH₂Cl₂). Mp 52-53°. [α]_D²⁵ +29.2. Relative config. determined. λ_{max} 268 (EtOH).

Bruneton, J. *et al.*, *Tet. Lett.*, 1975, 739 (*isol, uv, pmr, ms, struct*)Bandara, K.A.N.P. *et al.*, *Phytochemistry*, 2000, **54**, 29-32 (*Microcos alkaloid, isol*)Nakatani, Y. *et al.*, *Tetrahedron*, 2006, **62**, 160-165 (*Microcos alkaloid, synth*)**Cryptophorinine**

C-789

6-(5-Hydroxy-1,3,7-decatrienyl)-1,2-dimethyl-3-piperidinol, 9CI. 3-Hydroxy-6-(5-hydroxy-1,3,7-decatrienyl)-1,2-dimethylpiperidine
[56022-17-4]


C₁₇H₂₉NO₂ 279.422

Minor alkaloid from leaves of *Bathiorhannus cryptophorus* (Rhamnaceae). [α]_D²⁵ -68 (c, 1.3 in CHCl₃).

Bruneton, J. *et al.*, *Tet. Lett.*, 1975, 739 (*isol, ir, pmr, ms, struct*)**Cryptopine**

C-790

4,6,7,13-Tetrahydro-9,10-dimethoxy-5-methylbenzo[e]1,3-dioxolo[4,5-1][2]benzazecin-12(5H)-one, 9CI. Cryptocavine. Thalisyprine
[482-74-6]

C₂₁H₂₃NO₅ 369.416

Alkaloid from a variety of genera in the Papaveraceae (*Corydalis*, *Dicentra*), Papaveraceae (*Argemone*, *Meconopsis*, *Papaver*, *Stylomecon*) and Ranunculaceae (*Thalictrum*). Possesses uterotropic props. Prisms (Py/EtOH). Mp 221-223°.

▶ Highly toxic. DF4935300

Perchlorate: Mp 226-228°.

Picrate: Mp 163°.

O³-De-Me: **Izmirine**

[89117-94-2]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Fumaria parviflora* (Papaveraceae).

13-Oxo: **13-Oxocryptopine**. 14-Oxocryptopine
[15215-66-4]


C₂₁H₂₁NO₆ 383.4

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Cryst. (Me₂CO). Mp 185-186°. The 13-(or 14-) position is the methylene carbon adjoining the carbonyl group.

Haworth, R.D. *et al.*, *J. C. S.*, 1926, 1769 (*synth*)
Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 2479 (*ms*)Ma, J.C.N. *et al.*, *Can. J. Chem.*, 1965, **43**, 1849 (*pmr*)Hanus, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 1759 (*13-Oxocryptopine*)Hruban, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 3414 (*uv*)Hall, S.R. *et al.*, *Acta Cryst. B*, 1968, **24**, 346 (*cryst struct*)Brochmann-Hanssen, E. *et al.*, *Planta Med.*, 1970, **18**, 366 (*isol, deriv*)Nakashima, T.T. *et al.*, *Org. Magn. Reson.*, 1973, **5**, 9 (*cmr*)Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1983, **46**, 934 (*Izmirine*)Seger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**, 882-886 (*pmr, cmr*)Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 522**Cryptopleuridine**

C-791

11,12,13,14,14a,15-Hexahydro-6-methoxy-9H-1,3-dioxolo[6,7]phenanthro[9,10-b]quinolizin-12-ol, 9CI
[27303-58-8]



Probable structure

C₂₃H₂₃NO₄ 377.439

The alternative structure with the methylenedioxy-group at positions 6 and 7 and the methoxy-group at position 3 could not be excluded. Alkaloid from the bark of *Cryptocarya pleurosperma* (Lauraceae). Needles (C₆H₆). Mp 196-197°. [α]_D²⁰ +90 (c, 0.13 in CHCl₃).


Ac:

Needles (Me₂CO/CHCl₃). Mp 268-269°. [α]_D²⁰ +74 (c, 0.15 in CHCl₃).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 353 (*isol, uv, ir, pmr, ms, struct*)

Cryptoleurine C-792

11,12,13,14,14a,15-Hexahydro-2,3,6-trimethoxy-9H-phenanthro[9,10-b]quinoline, 9CI



(R)-form

C₂₄H₂₇NO₃ 377.482**(R)-form [482-22-4]**

Constit. of *Cryptocarya pleurosperma*, *Cryptocarya laevigata*, *Boehmeria cylindrica*, *Boehmeria platyphylla*, *Boehmeria candula*, *Boehmeria caudata* and *Cissus rheiifolia* (Lauraceae, Urticaceae, Vitidaceae). Shows antiyeast activity, antiviral activity against Herpes virus, and potent cytotoxic action vs. human nasopharyngeal cell cultures. Highly active inhibitor of protein synthesis. Cryst. (C₆H₆). Mp 197-198°. [α]_D¹⁸ -106 (c, 1.52 in CHCl₃). pK_a 7.55 (70% MeOH aq.). Isol. from *B. cylindrica* and *B. platyphylla* as a partial racemate. λ_{max} 282 (MeOH) (Berdy).

Hydrochloride: Mp 262° dec.

Hydrobromide: Mp 258-260° dec.

15R-Hydroxy: **15R-Hydroxycryptoleurine**

C₂₄H₂₇NO₄ 393.482

Alkaloid from the roots of *Boehmeria pannosa*. Yellow powder. Mp 201-203°. [α]_D²² -56.4 (c, 0.05 in MeOH). λ_{max} 262 (log ε 1.34) (MeOH).

(S)-form [87302-53-2]

Synthetic. Mp 196-197°. [α]_D²³ +106 (c, 1.0 in CHCl₃).

(±)-form [23365-52-8]

Synthetic. Shows antifungal and antibacterial activity. Needles (Me₂CO). Mp 201-202°.

Gellert, E. *et al.*, *Aust. J. Chem.*, 1954, **7**, 113; 1956, **9**, 489; 1978, **31**, 2095 (*isol, uv, ir, struct, ord, cd, abs config*)

Bradsher, C.K. *et al.*, *J.A.C.S.*, 1957, **79**, 3287; 1958, **80**, 930 (*synth, uv*)

Marchini, P. *et al.*, *Can. J. Chem.*, 1958, **36**, 581 (*synth, uv*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1968, **21**, 2579 (*isol*)

Farnsworth, N.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1805 (*isol, biochem*)

Paton, J.M. *et al.*, *J.C.S. (C)*, 1969, 1309 (*synth, uv, ms*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 353 (*pmr*)

Saxton, J.E. *et al.*, *Alkaloids (London)*, 1971, **1**, 86 (*rev*)

Krmpotic, E. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 1508 (*pharmacol*)

Kotani, E. *et al.*, *Tetrahedron*, 1974, **30**, 3027 (*synth, uv, ir, pmr, ms*)

Japan. Pat., 1975, 75 59 398; *CA*, **83**, 131809g (*synth*)

Herbert, R.B. *et al.*, *Chem. Comm.*, 1978, 794 (*synth*)

Iida, H. *et al.*, *Tet. Lett.*, 1981, **22**, 1913 (*synth*)

Söllhuber, M. *et al.*, *Heterocycles*, 1982, **19**, 347 (*cmr*)

Cragg, J.E. *et al.*, *J.C.S. Perkin 1*, 1982, 2487 (*synth*)

Al-Shamma, A. *et al.*, *Phytochemistry*, 1982, **21**, 485 (*isol, pharmacol*)

Saifah, E. *et al.*, *J. Nat. Prod.*, 1983, **46**, 353 (*isol*)

Bremmer, M.L. *et al.*, *J.O.C.*, 1983, **48**, 3661 (*synth*)

Buckley, T.F. *et al.*, *J.O.C.*, 1983, **48**, 4222 (*synth, uv, ir, pmr, ms*)

Iwao, M. *et al.*, *Tetrahedron*, 1983, **39**, 1955 (*synth*)

Iida, H. *et al.*, *J.O.C.*, 1984, **49**, 2412 (*synth, pmr, ms*)

Grieco, P.A. *et al.*, *J.O.C.*, 1988, **53**, 3325 (*synth, ir, pmr*)

Suzuki, H. *et al.*, *J.O.C.*, 1995, **60**, 6114 (*synth*)

Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1996, **118**, 12082 (*synth*)

Lebrun, S. *et al.*, *Tetrahedron*, 1999, **55**, 2659-2670 (*synth*)


Kim, S. *et al.*, *J.O.C.*, 2004, **69**, 3144-3149 (*synth*)

Fürstner, A. *et al.*, *Chem. Eur. J.*, 2006, **12**, 7398-7410 (*synth*)

Cai, X.F. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1095-1097 (*15-Hydroxycryptoleurine*)

Cryptoleurospermine C-793

[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl](3-hydroxy-4-methoxyphenyl)ethanedione, 9CI [27263-58-7]


C₂₀H₂₁NO₆ 371.389

Alkaloid from the bark of *Cryptocarya pleurosperma* (Lauraceae). Needles (Me₂CO). Mp 188-190°.

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 353 (*isol, uv, ir, pmr, ms, struct*)

Cryptoquinoline C-794

5-Methyl-11-(10H-quinolin-10-yl)-5H-quinoline, 9CI [160568-18-3]

C₃₁H₂₀N₄ 448.526

Alkaloid from roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Active against gram-positive bacteria. Dark green or orange amorph. powder. Artifact.


Paulo, A. *et al.*, *J. Ethnopharmacol.*, 1994, **44**, 127-130 (*activity*)

Paulo, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1485 (*isol, uv, ir, ms, struct*)

Cimanga, K. *et al.*, *Tet. Lett.*, 1996, **37**, 1703 (*pmr, cmr*)

Cryptospirolepine C-795

[154888-00-3]


C₃₄H₂₄N₄O 504.59

Alkaloid from roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Light brownish-pink cryst. (EtOH).

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1993, **56**, 653 (*isol, uv, ir, pmr, cmr, ms, struct*)

Cryptostyline I C-796

1-(1,3-Benzodioxol-5-yl)-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, 9CI. 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-(3,4-methylenedioxyphenyl)isoquinoline



(R)-form

R,R' = -CH₂-C₁₉H₂₁NO₄ 327.379

(R)-form

Alkaloid from *Cryptostylis erythroglossa* (Orchidaceae). Mp 101-102°. $[\alpha]_D^{22}$ -56 (c, 0.4 in CHCl₃).

Hydrobromide:

Cryst. (EtOH). Mp 239-240°. $[\alpha]_D$ -15.5 (MeOH).

(S)-form [22324-79-4]

Alkaloid from *Cryptostylis fulva* (Orchidaceae). Fine needles (Et₂O). Mp 101-102°. $[\alpha]_D$ +56 (c, 2.7 in CHCl₃).

Hydrobromide:

Cryst. (EtOH/Et₂O). Mp 239-240°. $[\alpha]_D$ +15.6 (MeOH).

(±)-form [20071-04-9]

Synthetic. Cryst. (Et₂O). Mp 117-118°.

(ξ)-form

1,2-Didehydro: 1-(3,4-Methylenedioxyphenyl)-6,7-dimethoxy-2-methyl-3,4-dihydroisoquinolinium (1+)

[52703-74-9, 18780-58-0]

C₁₉H₂₀NO₄⁺ 326.371

Quaternary alkaloid from *Cryptostylis erythroglossa* (Orchidaceae). Mp 207-208° (as bromide).

1,2,3,4-Tetrahydro: 6,7-Dimethoxy-2-methyl-1-(3,4-methylenedioxyphenyl)-isoquinolinium (1+)

C₁₉H₁₈NO₄⁺ 324.355

Quaternary alkaloid from *Cryptostylis erythroglossa* (Orchidaceae). Amorph. solid (as chloride); cryst. (MeOH aq.) (as picrate). Mp 218-223° (picrate). CAS no. refers to chloride.

Leander, K. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 244; 1973, **27**, 710 (*isol, uv, pmr, ms, synth, cryst struct, didehydro*)

Brossi, A. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 1564 (*cd, ord, abs config, uv, pmr*)

Westin, L. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 2305 (*cryst struct, abs config*)

Agurell, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1974, **28**, 239-243; 1175-1179 (*isol, biosynth, tetrahydro*)

Venkov, A.P. *et al.*, *Synthesis*, 1982, 486 (*synth*)
Ruchirawat, S. *et al.*, *Synth. Commun.*, 2003, **33**, 621-625 (*synth, ir, pmr, cmr, ms*)

Cryptostyline II C-797

1-(3,4-Dimethoxyphenyl)-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, 9CI

As Cryptostyline I, C-796 with R = R' = Me

C₂₀H₂₅NO₄ 343.422

(R)-form [33755-92-9]

Alkaloid from *Cryptostylis erythroglossa* (Orchidaceae). Cryst. (Et₂O). Mp 116-117°. $[\alpha]_D^{22}$ -58 (c, 0.4 in CHCl₃).

Hydrobromide:

Cryst. (EtOH). Mp 230-231°. $[\alpha]_D$ -22 (MeOH).

5'-Methoxy: 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-(3,4,5-trimethoxyphenyl)isoquinoline, 9CI. Cryptostyline III

[33033-86-2]

C₂₁H₂₇NO₅ 373.448

Alkaloid from *Cryptostylis erythroglossa* (Orchidaceae). Mp 128-130° (124-125°). $[\alpha]_D^{22}$ -52 (c, 0.3 in CHCl₃).

$[\alpha]_D$ -77.5 (CHCl₃).

5'-Methoxy, hydrobromide:

Cryst. (EtOH/Et₂O). Mp 245-246°.

$[\alpha]_D$ -21.2 (MeOH).

(S)-form [22324-82-9]

Alkaloid from *Cryptostylis fulva* (Orchidaceae). Needles (Et₂O). Mp 117-118°. $[\alpha]_D^{25}$ +58 (c, 0.28 in CHCl₃).

Hydrobromide:

Cryst. (EtOH/Et₂O). Mp 230-231°. $[\alpha]_D$ +21.5 (MeOH).

5'-Methoxy: [22325-16-2]

Alkaloid from *Cryptostylis fulva* (Orchidaceae). Needles (Et₂O). Mp 126-129° (120-121°). $[\alpha]_D^{25}$ +51 (c, 0.15 in CHCl₃). $[\alpha]_D^{19}$ +73.8 (c, 0.454 in CHCl₃).

5'-Methoxy, hydrobromide:

Cryst. (EtOH). Mp 245-246°. $[\alpha]_D$ +21.7 (MeOH).

(±)-form [22329-48-2]

Synthetic. Cryst. + 1H₂O (MeOH aq.), needles (Et₂O). Mp 94° (monohydrate) Mp 103-104° (anhyd.).

5'-Methoxy: [22324-83-0]

Synthetic. Cryst. (Et₂O or EtOH). Mp 141-142° (137-138°).

Müller, A. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1967, **52**, 261 (*synth, uv, ir*)

Leander, K. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 244; 1973, **27**, 710 (*isol, uv, pmr, ms, abs config, synth*)

Brossi, A. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 1564 (*cd, ord, abs config, synth, uv, pmr*)
Kametani, T. *et al.*, *J. Het. Chem.*, 1973, **10**, 451 (*Cryptostyline III, synth, cd*)

Blount, J.F. *et al.*, *Tetrahedron*, 1973, **29**, 31 (*synth, cd, abs config, cryst struct*)


Agurell, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1974, **28**, 239 (*isol*)

Venkov, A. *et al.*, *Synthesis*, 1982, 486 (*synth*)
Ruchirawat, S. *et al.*, *Synth. Commun.*, 2003, **33**, 621-625 (*synth, ir, pmr, cmr, ms, 5'-methoxy*)

Umetsu, K. *et al.*, *Tet. Lett.*, 2008, **49**, 2722-2725 (*synth*)

Cryptowoline C-798

5,6,13,13a-Tetrahydro-2-hydroxy-3-methoxy-7-methyl[1,3]dioxolo[5,6]indolo[2,1-a]isoquinolinium (1+), 9CI



C₁₉H₂₀NO₄⁺ 326.371

Revised abs. shown, by analogy with the revised abs. config. of Cryptaustoline, C-783.

(-)-form [15583-51-4]

Alkaloid from the bark of *Cryptocarya bowiei* and *Cryptocarya phyllostemon* (Lauraceae). Antineoplastic agent.

Chloride:

C₁₉H₂₀ClNO₄ 361.824

Cryst. (CHCl₃). Mp 188-190°. $[\alpha]_D$ -200 (c, 1 in EtOH).

Iodide:

C₁₉H₂₀INO₄ 453.276

Off-white prisms (MeOH). Mp 245-246° dec. $[\alpha]_D^{20}$ -186 (c, 0.4 in EtOH).

Me ether:

C₂₀H₂₂NO₄⁺ 340.398

Prisms (H₂O)(as iodide). Mp 227° (iodide). $[\alpha]_D^{20}$ -179 (c, 0.4 in EtOH) (as iodide).

O-De-Me, O²-Me: Cryptowolidine

[122992-81-8]

C₁₉H₂₀NO₄⁺ 326.371

Alkaloid from the stem bark of *Cryptocarya phyllostemon* (Lauraceae). Amorph. (as chloride). Negative opt. rotn. Abs. config. not determined but prob. as shown following the revised abs. config. for Cryptowoline.

13β-Hydroxy: Cryptowolinol

[122992-79-4]

C₁₉H₂₀NO₅⁺ 342.371

Alkaloid from *Cryptocarya oubatchensis* and *Cryptocarya phyllostemon* (Lauraceae). Cryst. (Me₂CO) (as chloride). Mp 218-220° (chloride). $[\alpha]_D$ -152 (c, 1 in EtOH). Revised abs. config. following the revised abs. config. for Cryptowoline.

(±)-form [17233-95-3]

Chloride: [17138-17-9]

Synthetic. Cryst. + ¹/₃ H₂O (EtOH aq.). Mp 255-257° dec.

Iodide: [7695-55-8]

Synthetic. Mp 150-151° dec Mp 245-246° dec.

[123050-06-6, 123050-05-5]

Ewing, J. *et al.*, *Aust. J. Chem.*, 1953, **6**, 78-85 (*isol, struct*)

Kametani, T. *et al.*, *J.C.S.(C)*, 1967, 2208-2212 (*synth, ms*)


Benington, F. *et al.*, *J.O.C.*, 1967, **32**, 1050-1053 (*synth*)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 766-769 (*synth*)

Leboeuf, M. *et al.*, *Can. J. Chem.*, 1989, **67**, 947-952 (*Cryptowolinol, Cryptowolidine*)

Crystamidine C-799

1,2,6,7,10,11-Hexadehydro-3-methoxy-15,16-[methylenebis(oxy)]erythrinan-8-one, 9CI. 10,11-Dehydro-8-oxoerythraline



C₁₈H₁₅NO₄ 309.321

(+)-form [58779-39-8]

Alkaloid from the leaves of *Erythrina crista-galli* and the seeds of *Erythrina brucei* (Fabaceae). Pale yellow oil. $[\alpha]_D^{23}$ +840 (c, 0.5 in CHCl₃). Possibly an artifact.

(±)-form


Synthetic. Gum.

Ito, K. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 52 (*isol, uv, ir, pmr, ms, struct*)

- Tsuda, Y. *et al.*, *Heterocycles*, 1984, **22**, 2255
(*synth*, *uv*, *ir*, *pmr*)
Dagne, E. *et al.*, *Phytochemistry*, 1984, **23**, 449
(*isol*)
Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**,
965 (*synth*)

Cuanzine**C-800**

[53492-09-4]

Absolute
ConfigurationC₂₂H₂₆N₂O₅ 398.458


Stereochem. revised in 1990. Alkaloid from the root bark of *Voacanga chalotiana* (Apocynaceae). Cryst. (C₆H₆). Mp 196°. [α]_D²² -11 (c, 1 in CHCl₃). [α]_D³⁰ (c, 2 in Py).

▶ JW4791800

- Bombardelli, E. *et al.*, *Tetrahedron*, 1974, **30**, 4141 (*uv*, *ir*, *ms*, *pmr*, *cmr*)
Palmisano, G. *et al.*, *J.O.C.*, 1990, **55**, 2182; 1991, **56**, 2380 (*pmr*, *cryst struct*, *synth*)
Ortuno, J.-C. *et al.*, *Tet. Lett.*, 1991, **32**, 4491 (*synth*)
Sóti, F. *et al.*, *Tetrahedron*, 1994, **50**, 8209 (*synth*)

Cucubalactam**C-801**

5-(1,1,2-Trimethylpropenylperoxy)-2-pyrrolidinone

C₁₀H₁₇NO₃ 199.249**(±)-form**


Constit. of *Cucubalus baccifer*. Amorph. powder (CHCl₃). Mp 32-33°.

- Cheng, Y.-X. *et al.*, *Heterocycles*, 2001, **55**, 1943-1949 (*isol*, *pmr*, *cmr*)

Cucumopine**C-802**

4,6-Dicarboxy-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-4-propanoic acid, 9CI

[110342-24-0]

C₁₁H₁₃N₃O₆ 283.24

Isol. from grapevine crown gall tumours and carrot hairy-root cultures. Amorph. powder + 1H₂O. Sol. H₂O. Mp 205-210° dec. [α]_D²² -33 (c, 0.37 in H₂O). λ_{\max} 218 (ε 5900) (H₂O) (Berdy).

Tri-Me ester:

Amorph. [α]_D²² -50 (c, 0.71 in CHCl₃).

4-Epimer: Mikimopine


[120293-52-9]

C₁₁H₁₃N₃O₆ 283.24

Alkaloid from hairy roots of tobacco induced by *Agrobacterium rhizogenes*. Amorph. powder. Sol. H₂O. [α]_D -98 (c, 0.48 in H₂O).

- Davioud, E. *et al.*, *Heterocycles*, 1988, **27**, 2423 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *synth*, *struct*)

- Isogai, A. *et al.*, *Phytochemistry*, 1990, **29**, 3131 (*Mikimopine*)

Culantraminol**C-803****(-)-form**C₃₂H₄₈N₂O₃ 508.743**(-)-form**

Alkaloid from *Zanthoxylum avicennae*.

Oil. [α]_D²⁴ -112.2 (c, 0.25 in CHCl₃).

Hydrobromide:

Cryst. (EtOH/EtOAc). Mp 296-298°.

*N-Oxide: Culantraminol N-oxide*C₃₂H₄₈N₂O₄ 524.742

Alkaloid from *Zanthoxylum avicennae*. Oil. [α]_D²⁵ -256.1 (c, 0.2 in CHCl₃). Isol. as a mixt. of the two possible *N*-oxide regioisomers.

7-Deoxy, 7,8-didehydro: Culantramine

[76129-63-0]

C₃₂H₄₆N₂O₂ 490.728

Alkaloid from *Zanthoxylum avicennae*. Oil. [α]_D²² -196.7 (c, 1 in CHCl₃). Struct. revised in 1985.

*7-Deoxy, 7,8-didehydro, N-oxide: Culantramine N-oxide*C₃₂H₄₆N₂O₃ 506.727

Alkaloid from *Zanthoxylum avicennae*. Oil. [α]_D²⁵ -200.8 (c, 0.5 in CHCl₃). Isol. as a mixt. of the two possible *N*-oxides.

(±)-form [103805-67-0]

Alkaloid from the leaves of *Zanthoxylum procerum* and *Zanthoxylum culantrillo* (Rutaceae). Noncryst.

7-Deoxy, 7,8-didehydro: [103823-32-1]

Alkaloid from the leaves of *Zanthoxylum culantrillo* and *Zanthoxylum procerum* (Rutaceae). Noncryst.

*7-Deoxy, 7,8-didehydro, N,N'-di-de-Me:***N,N'-Didemethylculantramine**

[129743-94-8]

C₃₀H₄₂N₂O₂ 462.674

Alkaloid from the leaves of *Zanthoxylum coriaceum* (Rutaceae). Amorph.

(ξ)-form*4-Epimer, 7-deoxy, 7,8-didehydro: Isoculantramine*

[76162-23-7]

C₃₂H₄₆N₂O₂ 490.728

Alkaloid from stems of *Zanthoxylum culantrillo*.

5-Epimer: 5-Epiculantraminol

[103882-04-8]

C₃₂H₄₈N₂O₃ 508.743

Minor alkaloid from the leaves of *Zanthoxylum procerum* (Rutaceae). Noncryst. Opt. activity not measured.

4,5-Diepimer: Alloculantraminol

[103882-03-7]

C₃₂H₄₈N₂O₃ 508.743

Minor alkaloid from leaves of *Zanthoxylum procerum* (Rutaceae). Oil. Opt. activity not measured.

- Swinehart, J.A. *et al.*, *Phytochemistry*, 1980, **19**, 1219-1223 (*Culantramine*)

- Schroeder, D.R. *et al.*, *Tetrahedron*, 1985, **41**, 4309-4320 (*isol*, *pmr*, *cmr*, *ms*, *struct*, *synth*, *epimers*)


- Marcos, M. *et al.*, *Phytochemistry*, 1990, **29**, 2315 (*Didemethylculantramine*)

- Miao, Z. *et al.*, *Youji Huaxue*, 1993, **13**, 145-149 (*Isoculantramine*)

- Thuy, T.T. *et al.*, *Phytochemistry*, 1999, **50**, 903-907 (*isol*, *N-oxides*)

Cularicine**C-804**

2,3,13,13a-Tetrahydro-1-methyl-1H-[1,3]dioxolo[7,8][1]benzoxepino[2,3,4-ij]isoquinolin-6-ol, 9CI

**(S)-form**C₁₈H₁₇NO₄ 311.337

Biogenetic (benzylisoquinoline-type) numbering shown.

(S)-form [2271-08-1]

Alkaloid from *Corydalis claviculata* (Papaveraceae). Prisms (MeOH). Mp 185°. [α]_D²² +295 (c, 0.96 in CHCl₃).

N-De-Me: Norcularicine

[89461-93-8]

C₁₇H₁₅NO₄ 297.31

Alkaloid from *Corydalis claviculata* (Papaveraceae). [α]_D²⁵ +216 (c, 0.06 in MeOH).

Me ether: O-Methylcularicine

[87578-91-4]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Corydalis claviculata* (Papaveraceae). Oil. [α]_D +283 (c, 0.3 in EtOH).

Me ether, hydrochloride:

Cryst. (MeOH). Mp 267°.

4R-Hydroxy: Corycularicine

[133137-53-8]

C₁₈H₁₇NO₅ 327.336

Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. [α]_D²⁵ +136 (c, 0.11 in MeOH).

(±)-form [40705-18-8]

Synthetic. Cryst. (Et₂O). Mp 155-156°.

Forms Me₂CO and MeOH solvates, Mp 81-82° and 95-96° resp.

Hydrochloride:

Needles (MeOH/Et₂O). Mp 255-257° dec.

N-*De-Me*: Synthetic. Cryst. (MeOH). Mp 101°.

Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1965, **43**, 989 (*isol, struct*)

Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2197 (*abs config*)

Noguchi, I. *et al.*, *Can. J. Chem.*, 1975, **53**, 125 (*synth, ir, pmr, ms*)


Allais, D.P. *et al.*, *J. Nat. Prod.*, 1983, **46**, 881; 1990, **53**, 1280 (*Norcularicine, Corycularicine*)

Blaschke, G. *et al.*, *Z. Naturforsch., C*, 1983, **38**, 670 (*O-Methylcularicine*)

Blaschke, G. *et al.*, *Phytochemistry*, 1985, **24**, 585 (*O-Methylcularicine*)

Cularine**C-805**

2,3,12,12a-Tetrahydro-6,9,10-trimethoxy-1-methyl-1H-[1]benzoxepino[2,3,4-ij]-isoquinoline, 9CI

*(S)-form*

C₂₀H₂₃NO₄ 341.406

Biogenetic (benzylisoquinoline-type) numbering shown.

(R)-form

Synthetic. Mp 113-114.5°. [α]_D -280 (c, 0.5 in MeOH).

N-*De-Me*: Obt. by resoln. of (±)-cularimine. Needles (Et₂O). Mp 101-102°. [α]_D -262.9 (c, 2.13 in MeOH).

(S)-form [479-39-0]

Alkaloid from *Corydalis claviculata*, *Dicentra cucullaria*, *Dicentra eximia*, *Dicentra formosa* and *Dicentra oregana* (Papaveraceae). Pharmacology of Cularine-group alkaloids little studied. Shows local anaesthetic and hypotensive activity. Cryst. (Et₂O or MeOH). Mp 115°. [α]_D²⁵ +285 (c, 0.8 in MeOH).

Hydrochloride: Mp 207°.

N-α-Oxide: cis-Cularine N-oxide

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Ceratocarpus heterocarpa*. Amorph. powder. Mp 122-124°. [α]_D +202 (c, 0.081 in MeOH). λ_{max} 230 (sh) (log ε 4.09); 284 (log ε 3.74) (MeOH).

N-Me: N-Methylcularine

[38852-37-8]

C₂₁H₂₆NO₄⁺ 356.441

Quaternary alkaloid from aerial parts of *Sarcocapnos baetica* ssp. *integrifolia* (Papaveraceae). Amorph. solid (as iodide). Mp 205° (iodide, synthetic). [α]_D +200 (c, 0.42 in MeOH). CAS no. refers to iodide.

N-*De-Me*: **Cularimine**. Alkaloid F30 [479-42-5]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Dicentra eximia* (Papaveraceae). Needles (Et₂O). Mp 101-102°. [α]_D +259.5 (c, 0.94 in MeOH).

N-De-Me, N-benzoyl:

Cryst. (MeOH/Et₂O). Mp 174°.

O⁷-De-Me: Cularidine. Alkaloid F10

[5140-50-1]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Corydalis claviculata* and *Dicentra cucullaria* (Papaveraceae). Needles (MeOH/Et₂O). Mp 157°. [α]_D +292 (MeOH).

O⁷-De-Me, perchlorate:

Cryst. (MeOH). Mp 297°.

O⁷-De-Me, N-α-oxide: cis-Cularidine N-oxide

[122555-12-8]

C₁₉H₂₁NO₅ 343.379

Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Amorph. solid. Mp 258-260°. [α]_D +240 (c, 0.075 in MeOH).

O⁷-De-Me, N-β-oxide: trans-Cularidine N-oxide

[122620-83-1]

C₁₉H₂₁NO₅ 343.379

Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Amorph. solid. Mp 200-202°. [α]_D +201 (c, 0.06 in MeOH).

O⁷-De-Me, N-De-Me: Norcularidine

[89240-43-7]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. [α]_D²⁵ +216 (c, 0.06 in MeOH).

O¹²-De-Me: Celtine

[91106-26-2]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Cryst. (EtOH). Mp 94-96°. [α]_D +181 (c, 0.08 in MeOH).

O¹³-De-Me: Enneaphylline

[104420-84-0]

Alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Prisms (EtOH). Mp 205-207°. [α]_D²⁵ +256 (c, 0.8 in EtOH). Abs. config. shown as (S-) but not explicitly assigned. Incorr. descr. as Cularine in one paper.

O⁷, O¹²-Di-de-Me: Celtisine

[91106-25-1]

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Cryst. (EtOH). Mp 158-160°. [α]_D +212 (c, 0.25 in MeOH).

O⁷, O¹³-Di-de-Me: Culacorine. Breoganine

[16209-79-3]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Corydalis claviculata* and *Sarcocapnos crassifolia* (Papaveraceae). Cryst. (EtOH). Mp 249-250°. [α]_D +278 (c, 0.057 in MeOH). [α]_D²⁵ +188 (c, 0.08 in MeOH).

4R-Hydroxy: Limousamine

[87853-59-6]

C₁₉H₂₁NO₅ 343.379

Alkaloid from *Corydalis claviculata* (Papaveraceae). [α]_D²⁵ +185 (c, 0.074 in MeOH).

(±)-form [28111-18-4]

Synthetic. Mp 134-135° (119°, 113-114°).

N-Me:

Cryst. + 1H₂O (MeOH/Et₂O). Mp 218°.

N-De-Me: [2512-67-6]

Synthetic. Mp 141-142°.

O⁷-De-Me: [50453-79-7]

Synthetic. Needles (MeOH). Mp 201-203°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 81; 1940, **18**, 97 (*isol, Cularine, Cularidine*)

Manske, R.H.F. *et al.*, *J.A.C.S.*, 1950, **72**, 55 (*struct, Cularine, Cularidine*)

Ohashi, M. *et al.*, *J.A.C.S.*, 1963, **85**, 2807 (*ms*)

Kametani, T. *et al.*, *J.C.S.*, 1963, 4289; 1964, 4142; 1965, 5565 (*synth, resoln, Cularine, Cularidine*)

Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1965, **43**, 989; 1966, **44**, 561 (*isol, struct, Cularidine*)

Bhacca, N.S. *et al.*, *Tetrahedron*, 1966, **22**, 1467 (*uv, ord, pmr, abs config*)

Ishiwata, S. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1850 (*synth, pmr*)

Kametani, T. *et al.*, *J. Het. Chem.*, 1970, **7**, 339 (*synth, pmr, Cularidine*)

Kametani, T. *et al.*, *Bioorg. Chem.*, 1971, **1**, 40 (*synth*)

Kametani, T. *et al.*, *Chem. Comm.*, 1971, 352; 1972, 1072 (*biosynth, cryst struct, abs config*)

Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2197 (*abs config*)

Iida, H. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 1001 (*synth, pmr, Cularidine*)

Jackson, A.H. *et al.*, *J.C.S. Perkin 1*, 1974, 1911 (*synth, pmr, ms, Cularine, Cularidine*)

Hagaman, E.W. *et al.*, *Org. Magn. Reson.*, 1976, **8**, 389 (*cmr*)

Shono, T. *et al.*, *Tet. Lett.*, 1981, **22**, 2385 (*synth*)

Allais, D.P. *et al.*, *Heterocycles*, 1983, **20**, 2055 (*Limousamine, isol, uv, pmr, ms, cd, struct*)

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1983, **46**, 881 (*Culacorine*)

Guinaudeau, H. *et al.*, *Heterocycles*, 1984, **22**, 107 (*Norcularidine*)

Boente, J.M. *et al.*, *Tet. Lett.*, 1984, **25**, 1829 (*Culacorine, Celtisine*)

De Lera, A.R. *et al.*, *Heterocycles*, 1986, **24**, 2219 (*Limousamine, synth*)

Castedo, L. *et al.*, *Heterocycles*, 1987, **26**, 29; 29 (*N-Methylcularine, Enneaphylline*)

Rodriguez de Lera, A. *et al.*, *J. Het. Chem.*, 1987, **24**, 95 (*synth, Celtisine, Enneaphylline*)


De Sousa, J.D.F. *et al.*, *J.A.C.S.*, 1994, **116**, 9745 (*synth*)

Suau, R. *et al.*, *Heterocycles*, 1995, **41**, 2575 (*Cularidine N-oxides*)

Garcia, A. *et al.*, *Tetrahedron*, 1995, **51**, 8585 (*synth*)

Suau, R. *et al.*, *Phytochemistry*, 1996, **43**, 1389 (*oxide*)

Rodrigues, J.A.R. *et al.*, *J.O.C.*, 2004, **69**, 2920-2928 (*synth*)

Curacautine**C-806***(S)-form*

C₃₉H₄₂N₂O₉ 682.769

(R)-form

10-Demethoxy, O¹²-de-Me: **Sindamine**
[84435-34-7]
C₃₇H₃₈N₂O₈ 638.716
Minor alkaloid from the roots of *Berberis lycium* (Berberidaceae). [α]_D²⁵ +38 (c, 0.04 in MeOH) (as Ac).

10-Demethoxy, α'-carboxylic acid: **Pycmanilline**
[115439-17-3]
C₃₈H₄₀N₂O₉ 668.742
Alkaloid from the roots and stems of *Pycnarrhena manillensis* (Menispermaceae). Also obt. by KMnO₄ oxidation of Phaeanthine, P-301. Mp 254-255°. [α]_D²² +33 (c, 0.61 in CHCl₃).

(S)-form [89412-86-2]

Alkaloid from *Berberis buxifolia* (Berberidaceae).

α'-Carboxylic acid, Me ester: **Talcamine**
[89412-85-1]
C₄₀H₄₄N₂O₁₀ 712.795
Alkaloid from *Berberis buxifolia* (Berberidaceae). [α]_D²⁵ -2 (c, 0.29 in MeOH).


(ξ)-form

10-Demethoxy: **Secoisotetrandrine**
[172924-26-4]
C₃₈H₄₀N₂O₈ 652.743
Alkaloid from leaves of *Laurelia sempervirens* (Peruvian nutmeg) (Monimiaceae). [α]_D²⁰ +2.73 (c, 1.1 in CHCl₃). Abs. config. shown as *S*- in the lit., but apparently without evidence.

Leet, J.E. et al., *Heterocycles*, 1982, **19**, 2355-2360 (*Sindamine*, *isol*, *uv*, *cd*, *ir*, *pmr*, *ms*, *struct*)
Leet, J.E. et al., *J. Nat. Prod.*, 1983, **46**, 908-912 (*Curacutine*, *Talcamine*, *isol*, *uv*, *ir*, *ms*, *cd*, *struct*)
Regalado, J.C. et al., *Heterocycles*, 1987, **26**, 2573-2578 (*Pycmanilline*, *uv*, *ir*, *pmr*, *ms*, *cd*, *struct*)
Schmeda-Hirschmann, G. et al., *Phytochemistry*, 1996, **41**, 339-341 (*Secoisotetrandrine*)

Curacin A**C-807**

[155233-30-0]



C₂₃H₃₅NOS 373.602
Isol. from the marine cyanobacterium *Lyngbya majuscula*. Antimitotic agent. Also possesses exceptional brine shrimp toxic and antiproliferative activities. Shows herbicidal props. Sol. MeOH, CHCl₃; fairly sol. hexane; poorly sol. H₂O. [α]_D +62 (c, 1.10 in CHCl₃) (a higher value of +86 was originally reported). Unstable when stored neat. λ_{max} 242 (hexane) (Berdy).

7Z-Isomer: Curacin B

[157319-51-2]
C₂₃H₃₅NOS 373.602
Isol. from the marine cyanobacterium

Lyngbya majuscula. Antimitotic and antiinflammatory agent. Immunosuppressant. Possesses brine shrimp toxic and antiproliferative activities. [α]_D²⁵ +62 (c, 0.84 in CHCl₃). λ_{max} 242 (ε 22000) (hexane) (Berdy).

9Z-Isomer: Curacin C

[164454-35-7]
C₂₃H₃₅NOS 373.602
From *Lyngbya majuscula*. Potent antimitotic agent. Sol. MeOH, CHCl₃. [α]_D²⁵ +56 (c, 0.15 in CHCl₃). λ_{max} 242 (ε 25000) (MeOH) (Berdy).

Gerwick, W.H. et al., *J.O.C.*, 1994, **59**, 1243 (*isol*, *pmr*, *cmr*, *struct*)

U.S. Pat., 1994, 5 324 739; *CA*, **121**, 149068c (*isol*, *pmr*)

Yoo, H.D. et al., *J. Nat. Prod.*, 1995, **58**, 1961 (*Curacins B and C*)

Nagle, D.G. et al., *Tet. Lett.*, 1995, **36**, 1189 (*abs config*)

Wipf, P. et al., *J.O.C.*, 1996, **61**, 6556 (*synth*)

White, J.D. et al., *J.A.C.S.*, 1997, **119**, 103 (*synth*, *abs config*)

Hoemann, M.Z. et al., *Tetrahedron*, 1997, **53**, 11087 (*synth*)

Muir, J.C. et al., *Tet. Lett.*, 1998, **39**, 2861-2864 (*synth*)

Wipf, P. et al., *Curr. Pharm. Des.*, 2004, **10**, 1417-1437 (*rev*)

Curacin D**C-808**

[164454-36-8]
As Curacin A, C-807 with

R = H


C₂₂H₃₃NOS 359.575

Absolute configuration has not been confirmed. Isol. from *Lyngbya majuscula*. Antimitotic agent, tubulin polymerisation inhibitor. Pale yellow oil. [α]_D +33 (c, 0.14 in CHCl₃). The name Curacin D also refers to a semisynthetic stereoisomer of Curacin A. λ_{max} 224 (ε 9000) (hexane).

Marquez, B. et al., *Phytochemistry*, 1998, **49**, 2387-2389 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

C-Curarine**C-809**

18,18'-Dideoxy-2,2'-epoxytoxiferine I,
9CI. C-Curarine I
[7168-64-1]



C₄₀H₄₄N₄O₂[⊕] 596.814
Produced by photooxidation of C-Dihydrotoxiferine. Alkaloid from calabash curare, *Strychnos divaricans* and some other *Strychnos* spp. (Loganiaceae). Powerful curarising agent largely responsible for the paralytic effect of amazonian arrow poisons. Mp 308-309° (as dipicrate). [α]_D +72 (H₂O) (as

dichloride). Blue col. with Ce(SO₄)₂, turning chrome green on standing.

▶ LD_{Lo} (mus, ivn) 0.05 mg/kg. V. toxic by intravenous route. GS3675000

Di-N-de-Me: Bisor-C-curarine

C₃₈H₃₈N₄O 566.744
Minor alkaloid from *Strychnos dolichothyrsa* and from *Strychnos matopensis* and *Strychnos minfiensis* (Loganiaceae).

18-Hydroxy: C-Alkaloid G

C₄₀H₄₄N₄O₂[⊕] 612.813
Alkaloid from calabash curare and *Strychnos solimoesana* (Loganiaceae). Mp 272° (as picrate). Blue col. with Ce(SO₄)₂, becoming green on standing.

18,18'-Dihydroxy: C-Alkaloid E

C₄₀H₄₄N₄O₃[⊕] 628.813
Alkaloid from calabash curare, *Strychnos froesii*, *Strychnos solimoesana* and *Strychnos tomentosa* (Loganiaceae). Mp 285-286° (as picrate). Blue col. with Ce(SO₄)₂, becoming green on standing.

Wieland, H. et al., *Annalen*, 1938, **536**, 68; 1941, **547**, 140 (*isol*)

Schmid, H. et al., *Helv. Chim. Acta*, 1952, **35**, 1864; 1953, **36**, 102; 345 (*isol*, *synth*)

Bernauer, K. et al., *Helv. Chim. Acta*, 1957, **40**, 1999; 1958, **41**, 1202 (*synth*)

Zürcher, A. et al., *J.A.C.S.*, 1958, **80**, 1500 (*isol*)

Berlage, F. et al., *Helv. Chim. Acta*, 1959, **42**, 2650-2654 (*C-Alkaloid G*)

Nagyváry, J. et al., *Tetrahedron*, 1961, **14**, 138 (*uv*, *ir*, *pmr*, *struct*)

Grdicin, M. et al., *J.A.C.S.*, 1964, **86**, 3357 (*pmr*)


Jones, N.D. et al., *Chem. Comm.*, 1972, 805 (*cryst struct*)

Verpoorte, R. et al., *Org. Magn. Reson.*, 1977, **9**, 567 (*cmr*)

Massiot, G. et al., *Phytochemistry*, 1988, **27**, 3293-3304 (*isol*, *cmr*)

Curassavine**C-810**

Trachelanthamidine curassavate
[68385-70-6]

C₁₆H₂₉NO₄ 299.409

Ester of trachelanthemidine with 2-hydroxy-2-(1-hydroxyethyl)-3-methylpentanoic (Curassavic acid, Homoviridifloric acid). Alkaloid from *Heliotropium curassavicum* (Boraginaceae). Pale-yellow gum. [α]_D +0.9 (c, 0.32 in EtOH).

N-Oxide: Curassavine N-oxide

[68385-71-7]
C₁₆H₂₉NO₅ 315.409
Alkaloid from *Heliotropium curassavicum* (Boraginaceae). Mp 123-125° Mp 186-188° (dimorph.).

1-Epimer: Heliotropcurassavine

[82398-74-1]
C₁₆H₂₉NO₄ 299.409

Minor alkaloid of *Heliotropium curassavicum* (Boraginaceae). Pale-yellow gum. $[\alpha]_D^{25}$ -14.9 (c, 0.0037 in CHCl_3).

1,2-Didehydro: Curassavinine. (-)-Supinidine curassavate

[84306-92-3]

$\text{C}_{16}\text{H}_{27}\text{NO}_4$ 297.394

Minor alkaloid from *Heliotropium curassavicum* (Boraginaceae). Gum. Probable struct. Ester of Supinidine, S-644. Obt. only as a mixt. with Coromandalinine and Heliopinine in A-669.


Subramanian, P.S. *et al.*, *Aust. J. Chem.*, 1980, **33**, 1357 (*isol, ir, pmr, ms, struct*)

Mohanraj, S. *et al.*, *Phytochemistry*, 1982, **21**, 1775 (*Helicourassavine, Curassavinine*)

Curicycleatjine

C-811

[131984-84-4]



$\text{C}_{37}\text{H}_{36}\text{N}_2\text{O}_7$ 620.701

Alkaloid from *Cyclea atjehensis* (Menispermaceae). Amorph. $[\alpha]_D$ -193 (c, 0.27 in CHCl_3). $[\alpha]_D$ -122 (c, 0.27 in MeOH).

12-Me ether: Curicycleatjenine

[131984-83-3]

$\text{C}_{38}\text{H}_{38}\text{N}_2\text{O}_7$ 634.727

Alkaloid from *Cyclea atjehensis* (Menispermaceae). $[\alpha]_D$ -120 (c, 0.3 in CHCl_3). $[\alpha]_D$ -101 (c, 0.3 in MeOH).

1-Epimer: Isocuricycleatjine

[131984-86-6]

$\text{C}_{37}\text{H}_{36}\text{N}_2\text{O}_7$ 620.701

Alkaloid from *Cyclea atjehensis* (Menispermaceae). $[\alpha]_D$ -173 (c, 0.13 in CHCl_3). $[\alpha]_D$ -149 (c, 0.13 in MeOH).

1-Epimer, 12-Me ether: Isocuricycleatjenine

[131984-85-5]

$\text{C}_{38}\text{H}_{38}\text{N}_2\text{O}_7$ 634.727


Alkaloid from *Cyclea atjehensis* (Menispermaceae). $[\alpha]_D$ -238 (c, 0.18 in CHCl_3). $[\alpha]_D$ -191 (c, 0.18 in MeOH).

Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 553 (*isol, uv, cd, pmr, ms, struct*)

Curine

C-812

Bebeerine. Pelosine



$\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706

Diastereoisomeric with Chondocurine, C-420. Antimalarial agent. Shows muscle relaxant props. Log P 7.58 (uncertain value) (calc).

(+)-form [477-60-1]

Alkaloid from *Chondodendron candicans*, *Chondodendron microphyllum*, *Nectandra*

rodioei and *Radix pareirae bravae*. Cryst. (MeOH). Mp 215°. $[\alpha]_{\text{Hg}}^{20}$ +345.7 (c, 0.4 in 1M HCl).

Hydrochloride (1:2): Mp 271° (259-260°).

O¹²-Me: 12-O-Methyl(+)-curine

[6879-67-0]

$\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733

Alkaloid from the roots of *Cissampelos pareira* and the foliage of *Cyclea hainanensis* (Menispermaceae). Cryst. (MeOH/Py). Mp 164°. $[\alpha]_D$ +273 (c, 0.7 in CHCl_3).

O¹²-Me, picrate: Mp 204-205°.

O¹²-Me, Ac: Mp 204-205°.

O¹²-Me, N², N^{2'}-di-Me: Mp 200° dec. (as diiodide).

Di-Me ether: Mp 115-120°. $[\alpha]_D$ +266 (c, 0.97 in CHCl_3).

Di-Me ether, N², N^{2'}-di-Me: Mp 245-250° dec. (as diiodide). $[\alpha]_D$ +139 (c, 0.44 in 1:9 EtOH aq.).

(-)-form

Chondodendrine. Aristolochine†

[436-05-5]

Alkaloid from *Aristolochia indica*, *Chondodendron platyphyllum*, *Chondodendron tomentosum*, *Chondodendron toxicoferum*, *Cissampelos pareira*, *Cissampelos mucronata*, *Isolona pilosa*, *Pleogyne cunninghamii*, *Radix pareirae bravae*, *Paracyclea ohaiana*, *Cyclea barbata*, *Cyclea hainanensis*, *Stephania epigaeae* (Aristolochiaceae, Menispermaceae) and *Isolona ghesquierina*. Cryst. (MeOH). Mp 213-214° (C_6H_6 solvate 165-166°). $[\alpha]_D^{23}$ -284 (c, 0.34 in CHCl_3). $[\alpha]_D^{23}$ -331 (c, 0.4 in Py). $[\alpha]_D^{23}$ -286 (c, 0.4 in 0.1M HCl). More pharmacol. active isomer.

Hydrochloride (1:2): Mp 271-273° dec.

1:1 Complex with (-)-Tubocurine:

Toxicoferine

[12578-01-7]

$\text{C}_{72}\text{H}_{76}\text{N}_4\text{O}_{12}$ 1189.412

Isol. from stems of *Chondodendron toxicoferum* (Menispermaceae). Cryst. ($\text{CHCl}_3/\text{Me}_2\text{CO}$). Mp 286°. $[\alpha]_D$ -263 (1M HCl/EtOH). Behaves as a single compd.

N², N^{2'}-Di-Me: Mp 249-250° dec. $[\alpha]_D^{24}$ -135 (c, 1.18 in MeOH).

O⁷-Me: Chondrofoline. 7-O-Methyl(-)-curine

[31944-97-5]

$\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733

Alkaloid from the leaves of *Chondodendron platyphyllum*, *Isolona ghesquierina* and *Uvaria ovata* (Menispermaceae, Annonaceae). Antiplasmodial agent. Plates ($\text{CHCl}_3/\text{Et}_2\text{O}$ or Et_2O). Mp 136-140°. $[\alpha]_D^{20}$ -257 (c, 0.10 in 0.1M HCl). $[\alpha]_D^{20}$ -238 (c, 1.13 in CHCl_3). Struct. revised in 2000. λ_{max} 232 (log ϵ 4.57); 281 (log ϵ 3.97) (no solvent reported).

O¹²-Me: 12-O-Methyl(-)-curine

[59685-16-4]

$\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733

Alkaloid from the stem bark of *Guatteria megalophylla* (Annonaceae). Mp 162-164°. $[\alpha]_D^{20}$ -303 (c, 0.63 in CHCl_3).

O¹²-Me, Ac: Mp 214-215°.

Di-Me ether: O,O-Dimethyl(-)-curine.

Paraquileginine

[1812-55-1]

$\text{C}_{38}\text{H}_{42}\text{N}_2\text{O}_6$ 622.76

Alkaloid from the stem bark of *Guatteria megalophylla* (Annonaceae). Mp 133-136° (116-121°).

O⁶-De-Me: Cycleacurine. 6'-O-De-methyl(-)-curine

[38849-84-2]

$\text{C}_{35}\text{H}_{36}\text{N}_2\text{O}_6$ 580.679

Alkaloid from the roots of *Cyclea peltata* (Menispermaceae). Cryst. + 2H₂O (MeCN). Mp 205-208°. $[\alpha]_D^{25}$ -202 (c, 1.0 in MeOH).

O⁶-De-Me; hydrobromide (1:2):

Cryst. + 1H₂O (EtOH aq.). Mp 293-296°.

(±)-form

Hayatine

[26057-51-2]

Alkaloid from the roots and leaves of *Cissampelos pareira* and from the foliage of *Cyclea hainanensis* (Menispermaceae). Mp 298-303° dec.

Hydrochloride (1:2):

Prismatic rods (MeOH/Et₂O). Mp 286° dec.

Dipicrate: Mp 234-235° dec. (shrinks at 217-220°).

N, N'-Di-Me: Daijising

[26057-50-1]

$\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_6^{\oplus}$ 624.775

Alkaloid from *Cissampelos pareira* (Menispermaceae). Muscle relaxant. Mp 281° dec. (synthetic) (as diiodide).

O¹²-Me: Hayatinine

[16595-09-8]

$\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733

Alkaloid from the roots of *Cissampelos pareira* (Menispermaceae). Mp 231-232°.

Krishnaswamy, P.R. *et al.*, *J. Indian Chem. Soc.*, 1935, **12**, 476-485; **14**, 39-41 (*Aristolochine*)

King, H. *et al.*, *J.C.S.*, 1940, 737-746 (*isol, Chondrofoline*)

Dutcher, J.D. *et al.*, *J.A.C.S.*, 1946, **68**, 419-424 (*isol*)

Anet, F.A.L. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1950, **3**, 346-349 (*isol*)

Bick, I.R.C. *et al.*, *J.C.S.*, 1953, 3893-3896; 1960, 1896-1903; 2402-2407 (*isol, pmr, struct*)

Kidd, D.A.A. *et al.*, *J.C.S.*, 1954, 669-677 (*struct*)

Bhattacharji, S. *et al.*, *J. Sci. Ind. Res., Sect. B*, 1956, **15**, 363 (*Hayatine*)

Boissier, J.R. *et al.*, *J. Nat. Prod.*, 1965, **28**, 191 (*Hayatine*)

Haynes, L.J. *et al.*, *J.C.S.(C)*, 1966, 615-617 (*12-O-Methyl(+)-curine*)

Milne, G.W.A. *et al.*, *J.C.S.(C)*, 1966, 1966-1968 (*Hayatine*)

Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1967, **5**, 655 (*Chondodendrine*)

Bhatnagar, A.K. *et al.*, *J.O.C.*, 1967, **32**, 819-820 (*Hayatine*)

Cava, M.P. *et al.*, *Phytochemistry*, 1969, **8**, 2341 (*Curine, Toxicoferine, isol, struct*)


Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 599-602 (*Curine, Chondrofoline, pmr, ms, struct*)

Chowdhury, A.R. *et al.*, *Sci. Cult.*, 1972, **38**, 358-359; *CA*, **78**, 82058x (*Hayatine*)

- Kupchan, S.M. *et al.*, *J.O.C.*, 1973, **38**, 1846-1852 (*Cycleacurine*)
- Hocquemiller, R. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1977, **285**, 447-450 (*isol*)
- Panichpol, K. *et al.*, *Phytochemistry*, 1977, **16**, 621-622 (*Chondrofoline*)
- Chen, Z.-L. *et al.*, *CA*, 1981, **94**, 127204r (*Daijisong*)
- Koike, L. *et al.*, *J.O.C.*, 1981, **46**, 2385-2389 (*Curine, Chondrofoline, pmr, cmr*)
- Zhang, X.-X. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1981, **23**, 216-224; *CA*, **95**, 192262t (*Hayatine*)
- Felix, J.P. *et al.*, *Biochemistry*, 1992, **31**, 11793-11800 (*Chondodendrine, pharmacol*)
- Pedersen, S.E. *et al.*, *J. Biol. Chem.*, 1995, **270**, 31141-31150 (*Chondodendrine, pharmacol*)
- Martindale, *The Extra Pharmacopoeia*, 32nd edn., *Pharmaceutical Press*, 1999, 1322 (*Tubocurarine*)
- Mambu, L. *et al.*, *Planta Med.*, 2000, **66**, 537-540 (*isol, pmr, cmr, N-15 nmr*)
- Tshibangu, J.N. *et al.*, *Phytochem. Anal.*, 2003, **14**, 13-22 (*Chondodendrine, isol, hplc*)

Curryangine C-813

9a,10,11,12,13,13a-Hexahydro-2,9,9,12-tetramethyl-1,12-epoxy-9H-indolo[3,2,1-de]phenanthridine, 9CI. Curryangine. Mahanimbidine. Murrayazoline



C₂₃H₂₅NO 331.457

(+)-form

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Needles (Me₂CO). Mp 276-278°. [α]_D²⁵ +2.25 (c, 0.4 in CHCl₃).

2'-Hydroxy: Murrayazolinol

[125287-09-4]

C₂₃H₂₅NO₂ 347.456

Minor alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 290°. The OH-config. does not appear to have been detd.

2'-Hydroxy, picrate: [125409-38-3]

Cryst. (C₆H₆). Mp 335°.

7-Hydroxy: Murrayamine E

[172617-68-4]

C₂₃H₂₅NO₂ 347.456

Alkaloid from leaves of *Murraya euchrestifolia* (Rutaceae). Prisms (MeOH). Mp 275-276° dec. [α]_D²⁵ +39.68 (c, 0.133 in CHCl₃).

(±)-form [25488-37-3]

Alkaloid from the leaves and stem bark of *Murraya koenigii* (curryleaf tree) and the leaves of *Murraya exotica* (Rutaceae). Also obt., together with Curryangine, by heating Mahanimbidine, M-46 at 200°. Cubes (C₆H₆). Mp 266° (260-262°).

Dutta, N.L. *et al.*, *Indian J. Chem.*, 1969, **7**, 1061; 1168 (*synth, uv, pmr*)

Kureel, S.P. *et al.*, *Tet. Lett.*, 1969, 3857 (*uv, ir, pmr, ms*)

Bordner, J. *et al.*, *Experientia*, 1972, **28**, 1406 (*uv, cryst struct*)

Narasimhan, N.S. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 430 (*synth, uv, ir, pmr*)

Bhattacharyya, P. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 308 (*uv, ir*)

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4132 (*isol, uv, ir, pmr, ms*)


Bhattacharyya, L. *et al.*, *J. Indian Chem. Soc.*, 1989, **66**, 140 (*Murrayazolinol*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1995, **40**, 1817 (*Murrayamine E*)

Ueno, A. *et al.*, *Org. Lett.*, 2008, **10**, 1999-2002 (*synth*)

Curryangine C-814

1,2,3,4,5,13-Hexahydro-5,7-dimethyl-2-(1-methylethenyl)-1,5-methanooxocino[3,2-a]carbazole, 9CI. Cyclomahanimbine. Murrayazolidine. Curryanine [25488-33-9]



C₂₃H₂₅NO 331.457

Alkaloid from the leaves and stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Needles (hexane). Mp 137° Mp 146°. [α]_D 0 (CHCl₃). [α]_D³⁰ +20 (CHCl₃). Probably an artifact. Obt. by brief acid or prolonged heat treatment of Mahanimbidine. λ_{max} 242 (log ε 4.83); 256 (log ε 4.61); 305 (log ε 4.41); 330 (log ε 3.87) (MeOH).

N-Me: Mp 169-170°.

1',2'-Dihydro: 1,2,3,4,5,13-Hexahydro-5,7-dimethyl-2-(1-methylethyl)-1,5-methanooxocino[3,2-a]carbazole. Exozoline

[70561-79-4]

C₂₃H₂₇NO 333.472

Alkaloid from the leaves of *Murraya exotica* (Rutaceae). Cryst. (C₆H₆/petrol). Mp 180-182° (136-137°, 146°). The lower Mps. refer to semisynthetic dihydrocurryanine. However it was reported that Exozoline and dihydrocurryanine were identical. λ_{max} 242 (log ε 4.47); 256 (log ε 4.18); 306 (log ε 3.97) (EtOH).

1',2'-Dihydro, 1'-hydroxy, stereoisomer: Murrayazolinine

[49620-01-1]

C₂₃H₂₇NO₂ 349.472

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) and from *Murraya euchrestifolia*. Mp 184°. Stereochem. not detd. λ_{max} 240 (ε 3160); 253 (ε 1740); 258 (ε 1620); 304 (ε 1150) (EtOH).

11-Hydroxy: Murrayamine D

[172548-90-2]

C₂₃H₂₅NO₂ 347.456

Alkaloid from leaves of *Murraya euchrestifolia* (Rutaceae). Oil. λ_{max} 215; 239; 266; 315; 324 (MeOH).

12-Methoxy: Murrayamine H

[183130-90-7]

C₂₄H₂₇NO₂ 361.483

Alkaloid from leaves of *Murraya euchrestifolia*. Syrup. λ_{max} 221; 241; 262; 303; 307 (MeOH).

Kureel, S.P. *et al.*, *Tet. Lett.*, 1969, 3857-3862 (*Cyclomahanimbine, uv, ir, pmr, ms, struct*)

Chakraborty, D.P. *et al.*, *Chem. Ind. (London)*, 1970, 593-594; 1973, 322-323; 1974, 260 (*Murrayazolidine, Murrayazolinine, uv, ir, pmr, struct, synth*)

Bandaranayake, W.M. *et al.*, *J.C.S. Perkin 1*, 1974, 998-1007 (*Cyclomahanimbine, struct, pmr, synth*)

Narasimhan, N. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 430-433 (*Curryangine, uv, ir, pmr, struct, synth*)


Ganguly, S.N. *et al.*, *Phytochemistry*, 1978, **17**, 1816-1817 (*Exozoline*)

Reisch, J. *et al.*, *Phytochemistry*, 1992, **31**, 2877-2879 (*Murrayazolidine, cryst struct*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1995, **40**, 1817-1819; 1996, **43**, 785-789 (*Murrayamine D, Murrayamine H, Murrayazolinine*)

Curromycin A C-815

[97412-76-5]



C₃₈H₅₅N₃O₁₀ 713.867

Isol. from *Streptomyces hygroscopicus*. Active against some kinds of gram-positive bacteria and shows cytotoxic props. It has a narrower antibacterial spectrum than Oxazolomycin, O-155 to which it is related. Yellow amorph. powder. Sol. MeOH, EtOAc. Mp 103-105°. [α]_D²⁵ +39 (c, 0.115 in MeOH). λ_{max} 229 (ε 19800); 267 (sh) (ε 16300); 275 (ε 19800); 285 (sh) (ε 15100) (MeOH) (Derep). λ_{max} 228; 275 (MeOH) (Berdy).

30-Demethoxy: 30-Demethoxycurromycin A. Curromycin B. IM 8443T. Antibiotic IM 8443T

[97412-77-6]

C₃₇H₅₃N₃O₉ 683.84

From *Streptomyces hygroscopicus*. Activity resembles that of Curromycin A. Yellow amorph. powder. Mp 106-109°. [α]_D²² +35 (c, 0.1 in MeOH).


Ogura, M. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 1909

Ogura, M. *et al.*, *J. Antibiot.*, 1985, **38**, 669 (*isol, uv, ir, cmr, pmr, ms*)

Okabe, T. *et al.*, *J. Antibiot.*, 1985, **38**, 964 (*isol*)

Curvupallide A C-816

[164269-18-5]



C₁₄H₁₇NO₅ 279.292

Alkaloid from the fungus *Curvularia pallidescens*. Herbicide. Oil. λ_{max} 230; 262 (MeOH) (Berdy).

7-Epimer: *Curvupallide B*

[164454-39-1]

C₁₄H₁₇NO₅ 279.292From *Curvularia pallescens*. Herbicide.Oil. $[\alpha]_D^{27}$ -92.7 (c, 1.00 in MeOH).**7-Epimer, demethoxy: *Curvupallide C***

[164269-19-6]

C₁₃H₁₅NO₄ 249.266From *Curvularia pallescens*. Herbicide.

Mp 168-170° dec.


Abraham, W.-R. *et al.*, *Tetrahedron*, 1995, **51**, 4947-4952 (isol, uv, ir, pmr, cmr, ms, struct)**Cuscamidine****C-817**Struct. unknown. Alkaloid from bark of *Cinchona pelletierana* (Rubiaceae).

Amorph. Forms amorphous salts.

Hesse, O. *et al.*, *Annalen*, 1880, **200**, 302**Cuscohygrine****C-818****1,3-Bis(1-methyl-2-pyrrolidinyl)-2-propanone, 9CI. Cuskhygrine. Bellaradine. Hellaradine**

[454-14-8]

[46727-05-3]

C₁₃H₂₄N₂O 224.345

Stereochem. not fully established. Natural Cuscohygrine is either the *meso*-form (illus.) or a mixt. of *meso*- and (\pm)-forms. Like other related alkaloids it appears to racemise and epimerise readily. It is unknown in opt. active form but opt. active dihydrocuscohygrine (see Dihydrocuscohygrine, D-413) is known which suggests that the opt. active form may be present in the plant. The correct struct. was assigned by Liebermann in 1896 but for some years an incorrect alternative was put forward by Hess. Hellaradine appears to be a misprint in *CA* for Bellaradine. Alkaloid from *Atropa belladonna* (roots only), *Hyoscyamus niger*, *Datura* spp., *Erythroxyllum coca* and other *Erythroxyllum* spp., *Solanum*, *Cyphomandra*, *Margaranthus*, *Acnistus*, *Salpichroa*, *Scopolia*, *Solandra*, *Convolvulus*, *Mandragora*, *Anthocercis* and *Lycianthes* spp. (Solana-ceae). Inhibitor of 2,4-dinitrofluorobenzene-induced hypersensitivity in mice. Several cuscohygrine-contg. plants are used in folk medicine as sedatives or narcotics. Oil; cryst. + 3½H₂O. d_4^{16} 0.98. Mp 40-41° (hydrate). $n_D^{18.4}$ 1.4845.

Methiodide: Mp 244°.**Oxime:** Mp 53-54°.


[72407-34-2]

Liebermann, C. *et al.*, *Ber.*, 1889, **22**, 675; 1896, **29**, 2050 (isol, struct)Späth, E. *et al.*, *Monatsh. Chem.*, 1948, **79**, 119 (synth)Rapoport, H. *et al.*, *J.O.C.*, 1949, **14**, 664 (synth, struct)Anet, E. *et al.*, *Nature (London)*, 1949, **163**, 289 (synth)Galinsky, F. *et al.*, *Monatsh. Chem.*, 1951, **82**, 551; 1953, **84**, 798 (synth, stereochem, bibl)El-Olemy, M.M. *et al.*, *J. Nat. Prod.*, 1966, **29**, 58 (synth)O'Donovan, D.G. *et al.*, *J.C.S. Perkin I*, 1969, 223-226 (biosynth)McGraw, B.A. *et al.*, *Phytochemistry*, 1978, **17**, 257; 1979, **18**, 189 (biosynth)Narasimhan, N.J. *et al.*, *CA*, 1980, **92**, 42185u (synth)Evans, W.C. *et al.*, *Phytochemistry*, 1980, **19**, 2351 (occur)Turner, C.E. *et al.*, *Phytochemistry*, 1981, **20**, 1403 (isol, stereochem, bibl)Leete, E. *et al.*, *J.A.C.S.*, 1982, **104**, 1403 (biosynth)Leete, E. *et al.*, *Phytochemistry*, 1983, **22**, 699 (biosynth)Ghani, A. *et al.*, *Indian J. Pharm. Sci.*, 1985, **47**, 127; *CA*, **104**, 3308q (occur)Stapper, C. *et al.*, *J.O.C.*, 2002, **67**, 6456-6460 (synth)**Cusconidine****C-819**Struct. unknown. Alkaloid from bark of *Cinchona pelletierana* (Rubiaceae).

Amorph. Forms amorphous salts.

Hesse, O. *et al.*, *Annalen*, 1880, **200**, 302**Cusconine****C-820**C₂₃H₂₆N₂O₄ 394.469

Struct. unknown. Alkaloid from bark of *Cinchona pelletierana* (Rubiaceae). Leaflets + 2H₂O (Et₂O). Mp 110° (anhyd.). $[\alpha]_D^{15}$ -54.3 (c, 2 in EtOH). Cooccurs with Aricine, A-1423 and Cusconidine, C-819. Forms amorph. salts.

Hesse, O. *et al.*, *Annalen*, 1877, **185**, 296**Cuspareine****C-821**C₂₀H₂₅NO₂ 311.423**(S)-form**

Alkaloid from the bark of *Galipea officinalis* (Rutaceae). Active against *Mycobacterium tuberculosis*. Mp 56°. $[\alpha]_D^{20}$ -20.4 (EtOH). λ_{max} 259 (log ϵ 4.02); 311 (log ϵ 3.43) (CHCl₃).

Methiodide: Mp 156°.**3'-O-De-Me: Galipeine**C₁₉H₂₃NO₂ 297.396


Alkaloid from the bark of *Galipea officinalis*. Oil. $[\alpha]_D$ -13.6 (CHCl₃). λ_{max} 259 (log ϵ 3.75); 311 (log ϵ 3.32) (CHCl₃).

(±)-form

Mp 32-35°.

Schläger, J. *et al.*, *Monatsh. Chem.*, 1950, **81**, 714 (uv, struct, synth)Rakotoson, J.H. *et al.*, *Planta Med.*, 1998, **64**, 762-763 (isol, uv, pmr, cmr, ms)Jacquemon-Collet, I. *et al.*, *Phytochemistry*, 1999, **51**, 1167-1169 (*Galipeine*)Houghton, P.J. *et al.*, *Planta Med.*, 1999, **65**, 250-254 (isol, pmr, cmr, activity)O'Byrne, A. *et al.*, *Tetrahedron*, 2008, **64**, 8079-8072 (synth)**Cusparine****C-822**

2-[2-(1,3-Benzodioxol-5-yl)ethyl]-4-methoxyquinoline, 9CI. 2-[2-(3,4-Methylenedioxyphenyl)ethyl]-4-methoxyquinoline
[529-92-0]

C₁₉H₁₇NO₃ 307.348

Alkaloid from bark of *Galipea officinalis* (Rutaceae). Antispasmodic agent. Active against *Mycobacterium tuberculosis*. Mp 91-92°. Log P 4.25 (calc).

Hydrochloride: Mp 189-191°.Späth, E. *et al.*, *Ber.*, 1924, **57**, 1243 (synth)Fournet, A. *et al.*, *Can. J. Chem.*, 1989, **67**, 2116-2118 (isol, pmr)Houghton, P.J. *et al.*, *Planta Med.*, 1999, **65**, 250-254 (activity)**Cyanamide, 9CI****C-823****Carbamionitrile. Carbamic acid nitrile.****Hydrogen cyanamide. Carbimide†.****Carbodiimide†**

[420-04-2]

H₂NC≡N (C_s)CH₂N₂ 42.04


Behaves chemically as the diimide HN=C=NH. Manuf. from limestone, coke and N₂. Synth. from CaCN₂ + H₂O/H₂SO₄. Isol. from *Robinia pseudo-acacia*, *Vicia cracca* and *Vicia villosa*. Aldehyde dehydrogenase inhibitor. Shows herbicidal props. Reportedly in use as a weed killer, e.g. on tobacco plants. White cryst. (C₆H₆/Et₂O). Sol. H₂O, EtOH, Et₂O, CHCl₃, C₆H₆, Me₂CO. Mp 46°. Bp 260° dec. Bp₁₉ 140°. pK_a -1.2 (H₂O). Steam-volatile; hydrol. by H⁺ or OH⁻ to urea. Converted by dil. OH⁻ into Cyanoguanidine. At 122°, trimerises to 1,3,5-Triazine-2,4,6-triamine.

► Fl. p. 141°. May dec. violently >50° in contact with H₂O, acids or alkalis. Max storage temp. 27° recommended. Eye, skin and mucous membrane irritant. Can cause dermatitis. LD₅₀ (rat, orl) 125 mg/kg. Exp. reprod. effects. OES: long-term 2 mg m⁻³. GS5950000

Ruppert, I. *et al.*, *Tet. Lett.*, 1977, 1987 (cmr)
Denner, L. *et al.*, *Acta Cryst. C*, 1988, **44**, 1979 (cryst struct)Kirk-Othmer *Encycl. Chem. Technol.*, 4th edn., Wiley, 1991, **7**, 736 (rev)Kamo, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 2310-2312 (occur)Kamo, T. *et al.*, *Phytochemistry*, 2008, **69**, 1166-1172 (occur)

Cyanocycline D

[152104-53-5]

Absolute
ConfigurationC₂₃H₂₇N₅O₅ 453.497


Artifact prod. from Cyanonaphthridinomycin, C-830 during workup. Isol. from a cyanide-treated broth of *Streptomyces lusitanus*. Active against a variety of bacteria. Cryst. (EtOAc/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 112-114°. λ_{max} 204 ; 268 ; 370 (MeCN) (Berdy).

Gould, S.J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1239-1245 (*isol, uv, ir, pmr, cmr, struct*)

5-(12-Cyano-6-dodeceny)-2-pyrrolicarboxaldehyde

C-825

13-(5-Formyl-1H-pyrrol-2-yl)-7-tridecenitrile, 9CI

C₁₈H₂₆N₂O 286.416**(Z)-form** [75234-01-4]


Metab. of the marine sponge *Laxosuberites* sp.

Sterle, D.B. *et al.*, *J.O.C.*, 1980, **45**, 4980 (*isol, uv, ir, pmr, cmr, ms, struct*)

N-[1-Cyano-2-(4-hydroxyphenyl)ethenyl]formamide, 9CI

C-826

2-(Formamido)-3-(4-hydroxyphenyl)-2-propenenitrile

C₁₀H₈N₂O₂ 188.185**(Z)-form** [957062-51-0]


Prod. by *Burkholderia mallei*. Cryst.

Brady, S.F. *et al.*, *J.A.C.S.*, 2007, **129**, 12102-12103 (*isol, pmr, cryst struct*)

5-(23-Cyano-23-hydroxy-6-tricosenyl)-1H-pyrrole-2-carboxaldehyde

C-827

24-(5-Formyl-1H-pyrrol-2-yl)-2-hydroxy-18-tetracosenenitrile, 9CI

C₂₉H₄₈N₂O₂ 456.71**(Z)-form** [75234-02-5]

Metab. of the marine sponge *Laxosuberites* sp. Mp 38-40°. Unusually stable to normal isol. and storage procedures. λ_{max} 297 (ε 16100) (MeCN) (Derep).

23-Deoxy: 5-(23-Cyano-6-tricosenyl)-1H-pyrrole-2-carboxaldehyde [290817-49-1]

C₂₉H₄₈N₂O 440.711

Isol. from the sponge *Mycale tenuispiculata*. Mp 56-58°. λ_{max} 297 (ε 15300) (MeCN).


Sterle, D.B. *et al.*, *J.O.C.*, 1980, **45**, 4980-4982 (*isol, uv, ir, pmr, cmr, ms, struct*)

Venkatesham, U. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1318-1320 (23-deoxy)

5-Cyano-6-methoxyindolo[2,3-a]carbazole

C-828

11,12-Dihydro-6-methoxyindolo[2,3-a]carbazole-5-carbonitrile, 9CI [131926-78-8]

C₂₀H₁₃N₃O 311.342

Isol. from the blue-green alga, *Nostoc sphaericum*. Shows varied cytotoxic activities. Amorph. λ_{max} 234 ; 252 ; 290 ; 348 ; 354 ; 372 (THF) (Berdy).

N¹¹-Me: 5-Cyano-6-methoxy-11-methylindolo[2,3-a]carbazole [131926-77-7]

C₂₁H₁₅N₃O 325.369


From *Nostoc sphaericum*. Cytotoxic. Amorph. Mp 230° dec. λ_{max} 234 (ε 25500); 252 (ε 28300); 290 (ε 66800); 338 (ε 15000); 354 (ε 10000); 372 (ε 11100) (THF) (Derep).

Knübel, G. *et al.*, *J. Antibiot.*, 1990, **43**, 1236 (*isol, pmr, cmr*)

3-Cyanomethyl-3-hydroxy-oxindole

C-829

2,3-Dihydro-3-hydroxy-2-oxo-1H-indole-3-acetonitrile, 9CI. 3-Cyanomethyl-3-hydroxy-1H-indol-2(3H)-one

*(S)*-formC₁₀H₈N₂O₂ 188.185**(S)-form** [137761-24-1]

Alkaloid from *Brassica oleracea* (Brassicaceae) inoculated with *Pseudomonas cichorii*. [α]_D -37.1 (c, 0.21 in MeOH).

(±)-form [137761-25-2]
Mp 162-163°.

(ξ)-form

N-β-D-Glucopyranosyl: [872851-13-3]

C₁₆H₁₈N₂O₇ 350.327

Alkaloid from the seeds of *Rheum maximowiczii*. [α]_D²¹ +38.5 (c, 2.8 in MeOH).

Hallmann, G. *et al.*, *Chem. Ber.*, 1962, **95**, 1138 (*synth*)

Monde, K. *et al.*, *Phytochemistry*, 1991, **30**, 2915 (*isol, uv, ir, pmr, ms*)


Monde, K. *et al.*, *Tet. Lett.*, 2003, **44**, 6017-6020 (*synth, cd, abs config*)

Komakine, N. *et al.*, *Nat. Med. (Tokyo)*, 2005, **59**, 45-48 (N-glucosyl)

Cyanonaphthridinomycin C-830

Cyanocycline A. Antibiotic 49A

[82423-05-0]

Absolute
ConfigurationC₂₂H₂₆N₄O₅ 426.471

Quinone-type antibiotic. Isol. from *Streptomyces flavogriseus*. DNA intercalator. Shows broad spectrum antimicrobial and antineoplastic activity. Stabilised and more active form of Naphthridinomycin A, N-32. Orange-red needles. [α]_D +76 (c, 0.5 in MeOH). pK_a 6.6. Log P -0.77 (uncertain value) (calc). Browns at 163°. Dec. at 168-170°. λ_{max} 263 (ε 9460) (MeOH/HCl) (Derep). λ_{max} 283 (ε 10400) (MeOH/NaOH) (Derep). λ_{max} 268 (ε 11000) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 10 mg/kg. GS9693000

O-De-Me: Cyanocycline F. Naphthocyanidin [82475-12-5]

C₂₁H₂₄N₄O₅ 412.444

Prod. semisynthetically from Antibiotic SF 1739, A-1254 or Cyanocycline A. Broad spectrum antibiotic with antineoplastic activity. Orange-red cryst. Mp 160°. [α]_D²⁵ +61 (c, 1 in CHCl₃). [α]_D +76 (c, 0.5 in MeOH). Log P -1.03 (uncertain value) (calc). λ_{max} 270 (EtOH).

▶ LD₅₀ (mus, ivn) 24 mg/kg. QL0527000

N-De-Me: Cyanocycline B. N-Demethylcyanocycline A

[152053-14-0]

C₂₁H₂₄N₄O₅ 412.444

Isol. from a cyanide-treated broth of *Streptomyces lusitanus*. Active against a variety of bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 210 ; 266 ; 375 (MeCN) (Berdy).

Hydroquinone: Cyanocycline C

[152075-38-2]

C₂₂H₂₈N₄O₅ 428.487


Isol. from a cyanide-treated broth of *Streptomyces lusitanus*. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 98-100° (as di-Me ether).

[73051-89-5]

- Japan. Pat., 1979, ((Chugai))79 126 793; CA, **93**, 68655t (isol)
- Zmijewski, M.J. et al., *J. Antibiot.*, 1982, **35**, 524-526 (synth, props)
- Itoh, I. et al., *J. Antibiot.*, 1982, **35**, 642-644 (isol)
- Hayashi, T. et al., *J. Antibiot.*, 1982, **35**, 771-777; 1983, **36**, 1228-1235 (isol, ur, ir, pmr, props)
- Hayashi, T. et al., *J.C.S. Perkin 2*, 1983, 335-343 (cryst struct)
- Fukuyama, T. et al., *J.A.C.S.*, 1987, **109**, 1587-1589 (synth)
- Hill, G.C. et al., *J. Med. Chem.*, 1991, **34**, 2079-2088 (DNA binding-conformn)
- Fukuyama, T. et al., *Adv. Heterocycl. Nat. Prod. Synth.*, 1992, **2**, 189-249 (rev, synth)
- Gould, S.J. et al., *J. Nat. Prod.*, 1993, **56**, 1239-1245 (Cyanocyclines B,C)
- Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, COL125

Cycastioamide C-831

3-(2-Carboxy-5-oxo-1-piperidinyl)thioalanyl-glycine, 9CI
[192801-72-2]




C₁₁H₁₇N₃O₅S 303.338
Alkaloid from the seeds of *Cycas revoluta* (Cycadaceae). Amorph. powder.

Pan, M. et al., *Phytochemistry*, 1997, **45**, 517-519 (isol, pmr, cmr)

Cyclamen C-832

10-Hydroxy-19-methylazacycloeicos-3,5,7,11,13,15,17-heptaen-2-one, 9CI
[155645-45-7]




C₂₀H₂₅NO₂ 311.423
Prod. by *Streptomyces* sp. MHW 846. Myeloperoxidase inhibitor and anti-inflammatory agent. Yellow powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁵ +3840 (DMF). λ_{max} 213 (ε 416); 276 (ε 2642); 285 (ε 3228) (MeOH) (Berdy). λ_{max} 228 (ε 294); 275 (ε 2262); 285 (ε 2812) (MeOH/HCl) (Berdy). λ_{max} 215 (ε 368); 276 (ε 2300); 285 (ε 2834) (MeOH/NaOH) (Berdy).

Ger. Pat., 1994, 4 231 289; CA, **121**, 7441v (isol)
Nazaré, M. et al., *Chem. Eur. J.*, 2001, **7**, 3363-3376 (synth)

Cyclanoline
Cissamine. Cissampine

C-833



(7R,13aS)-form

C₂₀H₂₄NO₄⁺ 342.414**(7R,13aS)-form****β-Cyclanoline**

[63527-13-9]

Quaternary alkaloid from *Stephania tetrandra* and *Cissampelos pareira* (Menispermaceae).

Chloride: [63527-14-0]
Cryst. Mp 205°. [α]_D²³ -150.3 (c, 0.1 in MeOH).

(7S,13aS)-form**α-Cyclanoline**

Chloride: [17472-50-3]

Mp 215-220°. [α]_D²⁶ -129 (c, 1.00 in CHCl₃).

▶DR9848000

Iodide: [17472-51-4]

Mp 185° dec.

Answer, F. et al., *Experientia*, 1968, **24**, 999 (uv, ms, pmr)


Yoshikawa, K. et al., *Chem. Lett.*, 1975, 961 (cmr)

Martin, H.J. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1977, **310**, 314 (isol, ms, uv)

Marek, R. et al., *Magn. Reson. Chem.*, 2002, **40**, 687-692 (N-15 nmr)

Cyclaneonine

C-834



(1R,1'R)-form

C₃₈H₄₂N₂O₆ 622.76**(1R,1'R)-form** [151380-29-9]

Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae). Cytotoxic. Amorph. powder. [α]_D²⁶ -119 (c, 1.28 in CHCl₃). λ_{max} 211 (ε 72443); 275 (ε 5248) (MeOH) (Berdy).

(1R,1'S)-form**Isocyclaneonine**

[151380-30-2]

Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae). Cytotoxic. Amorph. powder. [α]_D²⁶ +5.1 (c, 0.12 in CHCl₃). λ_{max} 211 (ε 64545); 275 (ε 5495) (MeOH) (Berdy).

(1S,1'S)-form [116520-07-1]

Alkaloid from stems of *Cyclea racemosa* (Menispermaceae). Cytotoxic. Amorph. powder. Mp 96-97°. [α]_D +376.8 (c, 0.5 in CHCl₃). λ_{max} 204 (ε 83000); 275 (ε 4180) (EtOH) (Berdy).


Lai, S. et al., *Yaoxue Xuebao*, 1988, **23**, 356-360; CA, **109**, 125816w

Wang, X.-K. et al., *Phytochemistry*, 1993, **33**, 1249-1252 (isol, uv, ir, pmr, ms, struct)

Cyclanone

C-835

Methylisochondodendrine. 7,7'-O,O-Di-methylisochondodendrine
[518-94-5]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from *Chondodendron tomentosum*, *Cissampelos insularis*, *Cissampelos pareira*, *Cleistopholis staudtii*, *Cyclea tonkinensis*, *Epinetrum cordifolium*, (preferred genus name *Albertisia*), *Epinetrum mangelotii*, *Epinetrum villosum*, *Heraclium wallichi*, *Limaciopsis loangensis*, *Paracylea ochiaiana*, *Stephania capitata*, *Stephania cepharantha*, *Stephania rotunda*, *Stephania elegans*, *Synclisia scabrida* and others (Menispermaceae, Annonaceae, Apiaceae). Shows antiinflammatory and anaesthetic props. Curarising agent. Shows selective antiplasmodial activity. Cryst. (Et₂O). Mp 280° (268-271°). [α]_D²² -15.9 (c, 0.69 in CHCl₃). Log P 7.8 (uncertain value) (calc). λ_{max} 232 (sh) (log ε 4.87); 276 (log ε 3.89); 285 (sh) (log ε 3.83) (MeOH).

N²-Oxide: Cyclanone N²-oxide

[85805-55-6]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from the stems of *Synclisia scabrida* (Menispermaceae). [α]_D²⁵ -7.6 (c, 0.38 in MeOH). The nmr spectra of two synthetic diastereoisomeric N-oxides were significantly different from those of the natural product which suggests that its struct. is incorr.

N^{2'}-De-Me: N^{2'}-Desmethylecyclanone

[83730-51-2]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the rhizomes of *Stephania glabra* (Menispermaceae). Mp 102-103°. [α]_D³² -165 (c, 0.29 in CHCl₃).

O⁷-De-Me: Norcyclanone

[478-63-7]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from *Chondodendron tomentosum*, *Cyclea insularis*, *Epinetrum cordifolium*, *Epinetrum mangelotii* and others (Menispermaceae). Needles (Me₂CO). Mp 245-246°. [α]_D²⁴ -22.5 (c, 1 in EtOH).

O⁷, O^{7'}-Di-de-Me: Isochondodendrine.*Isobebeerine. Isendryl. Isodendril.**O⁷, O^{7'}-Didemethylecyclanone*

[477-62-3]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Chondodendron candicans*, *Chondodendron limacifolium*, *Chondodendron microphyllum*, *Chondodendron platyphyllum*, *Chondodendron tomentosum*, *Chondodendron toxicoferum*, *Cissampelos mucronata*, *Cissampelos pareira*, *Cyclea barbata*, *Cyclea insularis*, *Cyclea madagascariensis*, *Cyclea peltata*, *Cyclea hainanensis*, *Cleistopholis staudtii*, *Epinetrum cordifolium*, *Epinetrum manganotii*, *Epinetrum illosum*, *Guateria megalophylla*, *Heracleum wallichii*, *Isolona pilosa*, *Paracyclea ochiatiana*, *Pleogyne cunninghamii*, *Sciadotenia toxifera*, *Stephania hernandifolia*, *Stephania elegans* and others (Menispermaceae, Annonaceae, Apiaceae). Antiplasmodial agent, sedative, anti-neoplastic agent. Mp 305° (288°, 275°) dec. $[\alpha]_D^{22} +120$ (c, 0.24 in 0.1M HCl). $[\alpha]_D^{22} +50$ (c, 0.965 in Py). $[\alpha]_D^{20} -29$ (c, 1.3 in CHCl₃). $[\alpha]_D^{20} -31$ (c, 0.33 in CHCl₃). Log P 7.32 (uncertain value) (calc). Exhibits anomalous solvent-dependent opt. activity. λ_{max} 211 (log ϵ 4.72); 231 (sh) (log ϵ 4.58); 278 (log ϵ 3.73); 285 (log ϵ 3.72) (EtOH).

► LD₅₀ (mus, ipr) 1001 mg/kg. GT8815000
O⁷,O⁷-Di-de-Me, sulfate: [32434-24-5]
Mp 291-292° dec. $[\alpha]_D +194$ (c, 1.3 in 1M H₂SO₄ aq.). $[\alpha]_D +50$ (c, 0.034 in 1M H₂SO₄ aq.). Opt. rotn. concn. dependent.

O⁷,O⁷-Di-de-Me, N²,N²-di-Me:
C₃₈H₄₄N₂O₆[⊕] 624.775
Mp 280° dec. (as diiodide). $[\alpha]_D^{24} +87$ (c, 0.7 in H₂O).

3,4-Didehydro: 3,4-Dehydrocycleanine
[189354-09-4]
C₃₈H₄₀N₂O₆ 620.744

Alkaloid from *Stephania cepharantha*.
Cryst. (Me₂CO). Mp 259-261° (dec.).
 $[\alpha]_D^{24} +79$ (c, 0.65 in CHCl₃). λ_{max} 340 (log ϵ 3.97) (MeOH).

1-Epimer: see Sciadanine, S-148

(±)-form [10438-10-5]
Synthetic. Mp 222-225°.

N²,N²-Di-de-Me: 2,2'-Dimethylcycleaninium
[75846-14-9]
C₄₀H₄₈N₂O₆[⊕] 652.829
Skeletal muscle relaxant. Antihypertensive agent. Used as dibromide, to which CAS no. refers.

► GT8820000

Dutcher, J.D. et al., *J.A.C.S.*, 1946, **68**, 419-424
(*Isochondodendrine*, isol)

Kidd, D.A.A. et al., *J.C.S.*, 1954, 669-677
(*struct*)

Jeffreys, J.A.D. et al., *J.C.S.*, 1956, 4451-4455
(*Isochondodendrine*, *struct*, *props*)

Kikuchi, T. et al., *Yakugaku Zasshi*, 1958, **78**,
1408-1412; 1959, **79**, 262-265; *CA*, **53**, 7219f;
13188d (*Norcycleanine*)

Bick, I.R.C. et al., *J.C.S.*, 1961, 1896-1903
(*pmr*)

Battersby, A.R. et al., *J.C.S.*, 1965, 2239-2247
(*ord*)

Debray, M. et al., *Ann. Pharm. Fr.*, 1966, **24**,
551-558; *CA*, **66**, 28942v (*Norcycleanine*)

Tomita, M. et al., *Chem. Pharm. Bull.*, 1968,
16, 62-69 (*synth*, *pmr*)

Baldas, J. et al., *J.C.S. Perkin 1*, 1972, 599-601
(*ms*)

Dwuma-Badu, D. et al., *Phytochemistry*, 1975,
14, 2520-2521 (*isol*, *uv*, *ir*, *pmr*, *ms*,
Isochondodendrine)

Galeffi, C. et al., *Gazz. Chim. Ital.*, 1978, **108**,
97-100 (*Isochondodendrine*, *isol*, *pmr*)

Marsaioli, A.J. et al., *Phytochemistry*, 1978, **17**,
1655-1658 (*cmr*)

Scheinmann, F. et al., *Phytochemistry*, 1980,
19, 1837-1840 (*isol*, *ir*, *pmr*, *cmr*, *ms*,
conformn)

Singh, R.S. et al., *J. Nat. Prod.*, 1981, **44**, 664-
667 (*Cycleanine*, *Isochondodendrine*, *isol*, *uv*,
ir, *pmr*, *ms*)

Bhakuni, D.S. et al., *J. Nat. Prod.*, 1982, **45**,
407-411 (*N-Desmethylcycleanine*)

Ohiri, F.C. et al., *Planta Med.*, 1983, **47**, 87-89
(*Cycleanine N²-oxide*)

Bhakuni, D.S. et al., *Tetrahedron*, 1983, **39**,
4003-4010 (*biosynth*)

Jin, G. et al., *Chin. Med. J. (Beijing, Engl.
edn.)*, 1984, **97**, 877-884 (*rev*, *pharmacol*,
methobromide)

Kashiwaba, N. et al., *Chem. Pharm.
Bull.*, 1997, **45**, 470-475 (3,4-
Dehydrocycleanine)

Kanyinda, B. et al., *J. Nat. Prod.*, 1997, **60**,
1121-1124 (*isol*, *pmr*, *cmr*)

Kashiwaba, N. et al., *J. Nat. Prod.*, 1998, **61**,
253-255 (*N-oxides*, *synth*, *pmr*, *cmr*)


Angerhoffer, C.K. et al., *J. Nat. Prod.*, 1999,
62, 59-66 (*activity*)

Mambu, L. et al., *Planta Med.*, 2000, **66**,
537-540 (*isol*, *pmr*, *cmr*, *N-15 nmr*,
activity)

Cycleatjehine

C-836

[128582-52-5]

C₃₆H₃₄N₂O₆ 590.674

New subgroup of bisbenzylisoquinolines incorporating the unusual methyleneoxy bridge (see also Warifteine, W-9 and derivs.). Alkaloid from the leaves of *Cyclea atjehensis* (Menispermaceae). Shows selective antiplasmodial activity. Amorph. $[\alpha]_D +321$ (c, 0.23 in CHCl₃). $[\alpha]_D +252$ (c, 0.23 in MeOH).

O¹²-Me: *Cycleatjehinine*
[128562-90-3]

C₃₇H₃₆N₂O₆ 604.701

Alkaloid from the leaves of *Cyclea atjehensis* (Menispermaceae). Shows selective antiplasmodial activity. Mp 218°. $[\alpha]_D +391$ (c, 0.25 in CHCl₃). $[\alpha]_D +352$ (c, 0.25 in MeOH).

Di-Me ether: $[\alpha]_D +272$ (c, 0.11 in MeOH). $[\alpha]_D +266$ (c, 0.175 in CHCl₃).

Tantisewie, B. et al., *Tetrahedron*, 1990, **46**, 325
(*isol*, *uv*, *pmr*, *ms*, *cd*, *struct*)


Parvez, M. et al., *Acta Cryst. C*, 1991, **47**, 448
(*cryst struct*, *Cycleatjehinine*)

Angerhoffer, C.K. et al., *J. Nat. Prod.*, 1999,
62, 59-66 (*activity*)

Cyclizidine

C-837

M 146791. Antibiotic M 146791. L 892-4.
Antibiotic L 892-4
[84393-28-2]

C₁₇H₂₅NO₃ 291.389

Indolizidine antibiotic. Prod. by *Streptomyces* sp. NCIB 11649. Shows immunostimulatory activity. Needles (EtOAc), cryst. (Et₂O). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 184° (176-178°). $[\alpha]_D^{23.5} -46.3$ (c, 2 in MeOH). Log P 1.62 (calc). λ_{max} 251 (ϵ 28400) (EtOH) (Derep).

6-Ac: [84393-27-1]

Cryst. Mp 63-65°. $[\alpha]_D^{22.5} -83.6$ (c, 2.0 in MeOH).

Freer, A.A. et al., *Chem. Comm.*, 1982, 1160
(*isol*, *struct*)


Leeper, F.J. et al., *Chem. Comm.*, 1987, 505
(*biosynth*, *pmr*, *cmr*)

Zhang, H. et al., *CA*, 1993, **119**, 67534w (*isol*)

Cyclo(alanyl-4-hydroxyprolyl)

C-838

Hexahydro-7-hydroxy-3-methylpyrrolo[1,2-a]pyrazine-1,4-dione, 9CI

C₈H₁₂N₂O₃ 184.194

(3S,7R,8aR)-form

Prod. by a strain of *Ruegeria* sp. isol. from the sponge *Suberites domuncula*. Amorph. solid. $[\alpha]_D^{25} +17.8$ (c, 0.001 in MeOH).

(3S,7R,8aS)-form

Prod. by a strain of *Ruegeria* sp. from *Suberites domuncula*. Also isol. from *Streptomyces fradiae*. Amorph. solid. $[\alpha]_D^{25} -4.2$ (c, 0.002 in MeOH).


Hoffmeister, D. et al., *Chem. Biol.*, 2000, **7**,
821-831 (*isol*)

Mitova, M. et al., *J. Nat. Prod.*, 2004, **67**,
1178-1181 (*isol*, *pmr*, *cmr*, *ms*)

Cyclo(alanylprolyl)

C-839

Hexahydro-3-methylpyrrolo[1,2-a]pyrazine-1,4-dione, 9CI





(3S,8aS)-form

C₈H₁₂N₂O₂ 168.195**(3R,8aS)-form***D,L*-form. (3*R*-cis)-form
[36238-64-9]
Mp 139-142°.**(3S,8aR)-form***L,D*-form. (3*S*-cis)-form
[19943-29-4]From *Beauveria bassiana*. Mp 162-166°.
[α]_D²⁰ -85.**(3S,8aS)-form***L,L*-form. (3*S*-trans)-form
[36357-32-1]Isol. from cocoa, *Pestalotia palmarum*,
Panax notoginseng, root of *Psammosilene*
tunicoides and the Caribbean sponge
Tedania ignis. Also prod. by *Streptomyces*
spp. Metab. of *Pseudoalteromonas halo-*
planktis, *Penicillium janthinellium* and
Cordyceps militaris. Mp 153-156°. [α]_{Hg}²⁰ -
160 (c, 1 in EtOH).




[65556-33-4, 19943-28-3]

Westley, J.W. *et al.*, *Anal. Chem.*, 1968, **40**,
1888-1890 (glc, chromatog, pmr)Siemion, I.Z. *et al.*, *Org. Magn. Reson.*, 1971,
3, 545-550 (synth, pmr)Young, P.E. *et al.*, *J.A.C.S.*, 1976, **98**, 5365-
5371 (synth, pmr, cmr)Siemion, I.Z. *et al.*, *Org. Magn. Reson.*, 1976,
8, 432-435 (cmr)Izumiya, N. *et al.*, *Pept. Chem.*, 1977, **15**, 49-54
(pharmacol)Pancoska, P. *et al.*, *Coll. Czech. Chem. Comm.*,
1979, **44**, 1296-1311 (cd)Cotrait, M. *et al.*, *Cryst. Struct. Commun.*,
1979, **8**, 819-822 (cryst struct)Grove, J.F. *et al.*, *Phytochemistry*, 1981, **20**,
815-816 (3*S,8aR*-form, isol)Schmitz, F.J. *et al.*, *J.O.C.*, 1983, **48**, 3941-3945
(isol, pmr, ms)Langhammer, M. *et al.*, *Fresenius' Z. Anal.*
Chem., 1986, **324**, 5-8 (ms)Zhang, Y. *et al.*, *Zhongguo Yaowu Huaxue*
Zazhi, 2002, **12**, 208-209; *CA*, **139**, 210500
(*L,L*-form, isol)Ma, X. *et al.*, *Shenyang Yaoke Daxue Xuebao*,
2003, **20**, 255-257 (*L,L*-form, isol)Stark, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**,
7222-7231 (isol, ms)Carls, K.L. *et al.*, *J. Phys. Chem. A*, 2005, **109**,
5463-5470 (cd)Mitova, M. *et al.*, *Mar. Biotechnol.*, 2005, **7**,
523-531 (*L,L*-form, isol)**Cycloanchinopeptolide C****C-840**


[160433-76-1]

C₃₃H₄₄N₁₀O₈ 708.773Stereochem. at C-5, C-4', C-5 and
around the cyclobutane ring is relative.
Alkaloid from the Mediterranean marine
sponge *Anchinoe tenacior*. [α]_D +18.6 (c,
0.5 in MeOH). λ_{max} 215 (ε 21000); 282 (ε
8080) (MeOH) (Berdy).Casapullo, A. *et al.*, *J. Nat. Prod.*, 1994, **57**,
1227 (isol, uv, pmr, cmr, cd, struct)**Cyclo(arginylpropyl)****C-841**3-[3-(Aminoiminomethyl)propyl]hexahy-
dropyrrolo[1,2-a]pyrazine-1,4-dione. 3-
(3-Guanidinypropyl)hexahydropyrro-
lo[1,2-a]pyrazine-1,4-dioneC₁₁H₁₉N₅O₂ 253.303**(3S,8aR)-form***L,D*-form. Antibiotic CI 4. CI 4
[173524-51-1]Prod. by the marine-derived *Pseudomo-*
nas sp. Chitinase inhibitor. Powder. [α]_D²⁰
+42 (c, 0.2 in H₂O).**(3S,8aS)-form***L,L*-form. *Verpamide A*
[74838-83-8]Isol. from *Axinella vacoleti*. [α]_D²⁴ -69.2 (c,
0.4 in MeOH).

Monohydrochloride: [115181-61-8]

Mp 245-250°. [α]_D²⁵ -72 (c, 1 in H₂O).**8,8a-Didehydro: Verpamide B**C₁₁H₁₇N₅O₂ 251.288Isol. from *Axinella vacoleti*. Oil (as for-
mate salt). [α]_D²⁴ -5.7 (c, 0.32 in MeOH)
(formate). Prob. config. is (3*S*-).Ishibashi, N. *et al.*, *Agric. Biol. Chem.*, 1988,
52, 819-827 (synth)Izumida, H. *et al.*, *J. Antibiot.*, 1996, **49**, 76-80
(Antibiotic CI 4)Vergne, C. *et al.*, *Org. Lett.*, 2006, **8**, 2421-2424
(*Verpamides*)**Cyclobrassinin****C-842**4,9-Dihydro-2-(methylthio)-1,3-thiazi-
no[6,5-b]indole, 9CI
[105748-58-1]C₁₁H₁₀N₂S₂ 234.345Isol. from Chinese cabbage (*Brassica*
campestris ssp. *pekinensis*) (Brassicaceae)
heads inoculated with *Pseudomonas ci-*
chorii. Phytoalexin. Mp 136-137°. First
report of the isol. of sulfur-containing
phytoalexins (see also Brassinin, B-284).
λ_{max} 204 (ε 24700); 227 (ε 34000); 284 (ε
7830); 294 (ε 8170) (MeOH) (Berdy).
λ_{max} 220 (ε 19200); 270 (ε 13200) (EtOH)
(Berdy).*S*-Oxide: 4,9-Dihydro-2-(methylsulfinyl)-
1,3-thiazino[6,5-b]indole, 9CI. *Cyclo-*
brassinin sulfoxide
[128722-96-3]C₁₁H₁₀N₂OS₂ 250.345Isol. from leaves of brown mustard
Brassica juncea (Brassicaceae). Phytoa-
lexin. Pale yellow grains. Mp 188-190°.**9-N-Methoxy: Sinalbin B**C₁₂H₁₂N₂OS₂ 264.372Isol. from *Sinapis alba* (white mus-
tard). Phytoalexin. λ_{max} 231 (log ε4.13); 275 (log ε 3.8) (CH₂Cl₂).9-*N*-Methoxy, *S*-oxide: *Sinalbin A*
Isol. from *Sinapis alba* (white mus-
tard). Phytoalexin. λ_{max} 231 (log ε
4.33); 280 (log ε 3.96) (CH₂Cl₂).Takasugi, M. *et al.*, *Chem. Comm.*, 1986, 1077
(isol, synth, struct)Devys, M. *et al.*, *Phytochemistry*, 1990, **29**,
1087 (isol, ir, pmr, ms, struct, oxide)Kutschy, P. *et al.*, *Tetrahedron*, 1998, **54**, 3549-
3566 (synth)Pedras, M.S.C. *et al.*, *Phytochemistry*, 2000, **53**,
161-176 (rev)Pedras, M.S.C. *et al.*, *Phytochemistry*, 2000, **55**,
213-216 (*Sinalbins*)**Cyclobrassinone****C-843**2-Methoxy-1,3-thiazino[6,5-b]indol-
4(9H)-one, 9CI
[156953-82-1]C₁₁H₈N₂O₂S 232.262The name Cyclobrassinone is misleading.
Isol. from *Brassica oleracea* var. *gongy-*
lodes (kohlrabi). Phytoalexin.Gross, D. *et al.*, *Z. Naturforsch., C*, 1994, **49**,
281-285 (isol)Kutschy, P. *et al.*, *Tetrahedron*, 2002, **58**, 9029-
9039 (synth)**Cyclobuxaphylamine****C-844**4,4,14-Trimethyl-3-(methylamino)-9,19-
cyclopregna-7,17(20)-dien-16-one, 9CIC₂₅H₃₇NO 367.573**(E)-form** [113762-74-6]Alkaloid from the leaves of *Buxus papilosa*
(Buxaceae). [α]_D -30 (c, 1.89 in CHCl₃).**(Z)-form** [113762-75-7]Alkaloid from the leaves of *Buxus papilosa*
(Buxaceae). [α]_D -76 (c, 1.11 in CHCl₃).Choudhary, M.I. *et al.*, *Phytochemistry*, 1988,
27, 271 (isol, uv, ir, pmr, ms, struct)**Cyclobuxargentine G****C-845**

[62874-53-7]



Probable structure


C₂₅H₄₂N₂O 386.62

Rare Buxus alkaloid lacking a C-4 substit. Minor alkaloid from *Buxus sempervirens* var. *argentea* (Buxaceae). Cryst. (MeOH/CH₂Cl₂). Mp 283°. [α]_D²³ -51 (c, 1.4 in CHCl₃).

Kuchkova, K.I. et al., *Chem. Zvesti*, 1976, **30**, 174-178; *CA*, **87**, 2362q (*isol, ir, ms, struct*)

Cyclobuxine I C-846

3,20-Diamino-4-methylene-14-methyl-9,19-cyclopregnan-16-ol, 9CI

C₂₃H₃₈N₂O 358.566

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Cyclobuxine I, is unknown.

N³,N²⁰-Di-Me: **Cyclobuxine D**. *Cyclobuxine*

[2241-90-9]

C₂₅H₄₂N₂O 386.62

Alkaloid from *Buxus sempervirens*, *Buxus hyrcana*, *Buxus wallichiana*, *Buxus microphylla* and *Buxus harlandi* (Buxaceae). Shows antiinflammatory and antihypertensive activity. Mp 245-247°. [α]_D²³ +98 (CHCl₃).

▶ LD₅₀ (mus, scu) 300 mg/kg. GZ0590020

N³,N²⁰-Di-Me; hydrobromide: Mp 288-292° dec.

N³,N²⁰-Di-Me, N³,N²⁰,O-tri-Ac: Mp 283-285° dec. [α]_D²⁴ +10 (CHCl₃).

N³,N³,N²⁰-Tri-Me: **Cyclobuxine B**

[5232-38-2]

C₂₆H₄₄N₂O 400.646

Alkaloid from *Buxus sempervirens* (Buxaceae). Prisms (MeOH). Mp 230-233°. [α]_D²⁵ +119 (c, 1.0 in CHCl₃).

N-Tetra-Me: *Cyclobuxine A*

Semisynthetic. Mp 204-205°. [α]_D²⁵ +99 (CHCl₃).

4β,23-Dihydro: 3,20-Diamino-4,14-dimethyl-9,19-cyclopregnan-16-ol. *Cyclobuxamine I*

4β,23-Dihydro, N²⁰-Me: **Cyclobuxamine H**. *Cyclobuxamine*

[6866-90-6]

C₂₄H₄₂N₂O 374.609

Alkaloid from *Buxus sempervirens* and *Buxus balearica* (Buxaceae). HIV reverse transcriptase (HIV-rt) inhibitor. Mp 209-211° dec. [α]_D²⁴ +67 (c, 1.08 in CHCl₃). [α]_D +30 (CHCl₃).

4β,23-Dihydro, N²⁰-Me, O,N³,N²⁰-tri-Ac: Mp 261-263° dec.

4β,23-Dihydro, N-tetra-Me: *Cyclobuxamine A*

[58700-02-0]

Semisynthetic. Mp 217-218°. [α]_D²⁴ +33 (c, 0.5 in CHCl₃).

4β,23-Dihydro, 6,7-didehydro, N-tetra-Me: **Cyclobullatine A**

[58672-78-9]

C₂₇H₄₆N₂O 414.673

Alkaloid from *Buxus sempervirens* var. *bullata* (Buxaceae). Cryst. (Me₂CO). Mp 275°. [α]_D -99 (c, 0.46 in EtOH).

3-Epimer, N³,N²⁰-di-Me: **Pseudocyclobuxine D**

[54357-48-1]

C₂₅H₄₂N₂O 386.62

Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (EtOH). Mp 229-231°. [α]_D +89.6 (CHCl₃).

Schlittler, E. et al., *Helv. Chim. Acta*, 1949, **32**, 2209-2226; 2226-2240 (*Cyclobuxine D*)

Brown, K.S. et al., *J.A.C.S.*, 1964, **86**, 4414-4424; 4420-4433; 4424-4430 (*Cyclobuxines, Cyclobuxamines*)

Votický, Z. et al., *Chem. Zvesti*, 1969, **23**, 702-703 (*Cyclobuxine B*)

Burnell, R.H. et al., *Phytochemistry*, 1972, **11**, 1853-1854 (*Cyclobuxine D*)

Khodzhaev, B.U. et al., *Khim. Prir. Soedin.*, 1973, **9**, 755-760; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 723-726 (*Pseudocyclobuxine D*)

Votický, Z. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 3055-3060; 1977, **42**, 541-547 (*Cyclobuxine D, Cyclobuxamine H, Cyclobullatine A*)


Abramson, D. et al., *Phytochemistry*, 1977, **16**, 1935-1937 (*biosynth*)

Saidkasimov, T. et al., *CA*, 1979, **91**, 235j (*pharmacol*)

Sun, H.D. et al., *J. Nat. Prod.*, 1996, **59**, 525-528 (*Cyclobuxine D, activity*)

Cyclobuxomicreimine C-847

[145194-13-4]

C₂₅H₃₇NO 367.573


Alkaloid from leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. [α]_D²⁰ +40 (CHCl₃).

Atta-ur-Rahman, et al., *Phytochemistry*, 1992, **31**, 2933-2935 (*isol, uv, ir, pmr, ms, struct*)

Cyclobuxophylline O C-848

3-Amino-4,4,14-trimethyl-9,19-cyclopregn-17(20)-en-16-one, 9CI

[79205-49-5]

C₂₄H₃₇NO 355.562

Alkaloid from *Buxus sempervirens* leaves (Buxaceae). Cryst. (Me₂CO). Mp 219-

222°. [α]_D²² -61.5 (c, 1.1. in CHCl₃).

N-Ethoxycarbonyl: **Buxene O**. *Buxene*

[24663-11-4]

C₂₇H₄₁NO₃ 427.626

Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 202-204°.

N-Me: *Buxenone*. *Buxenone M*. **Cyclobuxophylline M**. *Cyclobuxophyllinine*. *Cyclobuxophylline M*

[1053-21-0]

C₂₅H₃₉NO 369.589

Alkaloid from *Buxus microphylla* and *Buxus sempervirens* (Buxaceae). Needles (petrol). Mp 181-182° (174°). [α]_D -51 (c, 0.59 in CHCl₃). This alkaloid is called Cyclobuxophyllinine or Cyclobuxophylline M in the literature, but the systematic designation is Cyclobuxophylline M.

N-Me, N-Ac:

Needles (EtOH). Mp 234-235°. [α]_D -114 (c, 1.12 in CHCl₃).

N-Me, N-benzoyl: N-Benzoylcyclobuxophylline M. *Buxanine*. **Buxanine M**

[16049-32-4]

C₃₂H₄₃NO₂ 473.697

Alkaloid from *Buxus sempervirens* (Buxaceae). Prisms (Me₂CO). Mp 196-199°. [α]_D -38 (c, 0.2 in CHCl₃).

N-Me, N-ethoxycarbonyl: N-Methylbuxene M. N-Methylbuxene

[24663-12-5]

C₂₈H₄₃NO₃ 441.653

Minor alkaloid from *Buxus sempervirens* (Buxaceae). Mp 180-182°. [α]_D -104 (CHCl₃).

N,N-Di-Me: **Cyclobuxophylline K**. *Cyclobuxophylline*

[10088-21-8]

C₂₆H₄₁NO 383.616

Alkaloid from *Buxus microphylla* and *Buxus sempervirens*. Needles (Me₂CO or EtOH). Mp 194-196° Mp 201-203°. [α]_D -72 (c, 0.52 in CHCl₃) (-67). λ_{max} 244 (MeOH).

1,2-Didehydro, N-Me: **Buxithienine M**

[79205-50-8]

C₂₅H₃₇NO 367.573

Alkaloid from leaves of *Buxus sempervirens* var. *rotundifolia* (Buxaceae). Cryst. (Me₂CO/MeOH). Mp 166-169°. [α]_D²⁵ -75.6 (c, 1.2 in CHCl₃).

3-Epimer, N-Me: **Buximicrophylline B**

[15626-25-2]

C₂₅H₃₉NO 369.589

Alkaloid from *Buxus microphylla*. Needles. Mp 149-151°. λ_{max} 249 (log ε 2.77) (CCl₄).

[11011-82-8]

Nakano, T. et al., *J.C.S.(C)*, 1966, 1412

(*Cyclobuxophylline K, M*)

Döpke, W. et al., *Pharmazie*, 1966, **21**, 643; 769

Döpke, W. et al., *Tet. Lett.*, 1969, 4423

(*Buxene O*)

Vassová, A. et al., *Pharmazie*, 1970, **25**, 363

(*N-Methylbuxene M*)

Kuchkova, K.I. et al., *Chem. Zvesti*, 1976, **30**,

174 (*isol*)

Huong, L. et al., *Coll. Czech. Chem. Comm.*,

1981, **46**, 1425 (*isol, ir, pmr*)

Desai, M.C. et al., *Tetrahedron*, 1981, **37**, 2935


(*synth*)

- Du, J. *et al.*, *J. Asian Nat. Prod. Res.*, 1999, **1**, 239-244 (*Buxmicrophylline B*)
 Ata, A. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 21-28 (*Cyclobuxophylline K, isol, pmr*)
 Liu, H.-F. *et al.*, *Acta Cryst. E*, 2007, **63**, 2194-2195 (*Cyclobuxophylline M, cryst struct*)

Cyclobuxoviramine

C-849

[123853-65-6]




$C_{25}H_{39}NO$ 369.589
 Alkaloid from the leaves of *Buxus papillosa*. $[\alpha]_D^{20}$ -35.

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1989, **28**, 2848-2850 (*isol, pmr, cmr*)

Cyclobuxoviridine L

C-850

20-(Dimethylamino)-4,4,14-trimethyl-9,19-cyclopregn-1-en-3-one, 9CI. *Cyclobuxoviridine*
 [10088-20-7]



$C_{26}H_{41}NO$ 383.616
 Alkaloid from *Buxus sempervirens*, *Buxus papillosa*, *Buxus hyrcana* and *Buxus microphylla* (Buxaceae). Plates (petrol). $[\alpha]_D^{23}$ -20 (-16) ($CHCl_3$). One report gives an opt. rotn. of +16°, presumably due to a misprint.

N-De-Me: *Cyclobuxoviridine B. Cyclobuxoviricine*
 [100477-90-5]

$C_{25}H_{39}NO$ 369.589
 Alkaloid from leaves of *Buxus papillosa* and from *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D^{10}$ -54 ($CHCl_3$). Originally isol. by Arigoni *et al.*, under the name Cyclobuxoviridine B, but details were unpubl. Later independently isol. and called Cyclobuxoviricine but with incomplete stereochem. and it is not clear if the two are identical. Phys. props. given refer to Cyclobuxoviricine.

Nakano, T. *et al.*, *J.C.S. (C)*, 1966, 1412-1421 (*isol, uv, pmr, struct*)


Kuchkova, K.I. *et al.*, *Chem. Zvesti*, 1976, **30**, 174-178; *CA*, **87**, 2362q (*isol, ms, pmr*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1985, **24**, 3082-3083 (*Cyclobuxoviricine*)

Choudhary, M.I. *et al.*, *J. Nat. Prod.*, 1987, **50**, 84-88 (*isol, uv, ir, pmr, ms*)

Cyclobuxoxazine C*Cyclobuxoxazine*

[3294-83-5]



$C_{27}H_{46}N_2O_2$ 430.673

C-4 config. revised in 1975. Minor alkaloid from *Buxus microphylla* and *Buxus wallichiana* (Buxaceae). Cryst. (EtOH). Mp 245-246° (234-236°). $[\alpha]_D$ +48 (EtOH). $[\alpha]_D$ +29 ($CHCl_3$).

N,O-Di-Ac:

Needles (Me_2CO /hexane). Mp 240°. $[\alpha]_D$ -35 ($CHCl_3$).

N-Me: *Cyclobuxoxazine A. N-Methylcyclobuxoxazine*

[3676-46-8]

$C_{28}H_{48}N_2O_2$ 444.699

Alkaloid from *Buxus rolfei* (Buxaceae). Mp 197° Mp 201-202° (synthetic).

16-Deoxy, N-Me: *Desoxycyclobuxoxazine A*

[34437-03-1]

$C_{28}H_{48}N_2O$ 428.7

Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (MeOH). Mp 161-162°. $[\alpha]_D$ +56 (c, 0.2 in $CHCl_3$).

11-Oxo: *Baleabuxoxazine C. Cyclobuxoxazine C*

[14155-74-9]

$C_{27}H_{44}N_2O_3$ 444.656

Alkaloid from *Buxus balearica* (Buxaceae). Mp 292°. $[\alpha]_D$ +116 ($CHCl_3$).

Nakano, T. *et al.*, *J.C.S.*, 1965, 4537-4542 (*isol, pmr, struct, synth*)

Khuong-Huu-Lainé, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 1216-1221 (*Cyclobuxoxazine A*)

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1966, **22**, 3321-3327 (*Baleabuxoxazine C*)

Vassová, A. *et al.*, *Pharmazie*, 1970, **25**, 363-365; *CA*, 1970, **73**, 127748n (*isol*)


Härtel, R. *et al.*, *Tet. Lett.*, 1971, 2741-2742 (*Desoxycyclobuxoxazine A*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*cmr, config*)

Cyclocarbamide A

C-852

3-Methyl-N-(5a,6a,7,8-tetrahydro-4-methyl-1,5-dioxo-1H,5H-pyrrolo[1,2-c][1,3]oxazepin-3-yl)butanamide, 9CI. 7,8,9,9a-Tetrahydro-3-[(3-methyl-1-oxobutyl)amino]-4-methyl-1H,5H-pyrrolo[1,2-c][1,3]oxazepine-1,5-dione. *Herbicidal substance 1328-2*
 [102719-89-1]



$C_{14}H_{20}N_2O_4$ 280.323

Prod. by *Streptovorticillium* sp. Herbicide,

C-851

plant growth regulator. Prisms ($EtOAc$ /hexane). Mp 109-110°. $[\alpha]_D$ -259 (c, 0.5 in MeOH). λ_{max} 282 (ε 10400); 328 (ε 12000) (MeOH/NaOH) (Derep). λ_{max} 274 (ε 9800) (MeOH) (Derep). λ_{max} 274 (MeOH/HCl) (Berdy).

N-Hexanoyl analogue: N-(5a,6a,7,8-Tetrahydro-4-methyl-1,5-dioxo-1H,5H-pyrrolo[1,2-c][1,3]oxazepin-3-yl)hexanamide, 9CI. 7,8,9,9a-Tetrahydro-3-[(1-oxohexyl)amino]-4-methyl-1H,5H-pyrrolo[1,2-c][1,3]oxazepine-1,5-dione. *Cyclocarbamide B. Herbicidal substance 1328-3*
 [102719-90-4]

$C_{15}H_{22}N_2O_4$ 294.35

From *Streptovorticillium* sp. Herbicide, plant growth regulator. $[\alpha]_D$ -150 (c, 0.03 in MeOH). λ_{max} 282 (ε 10400); 328 (ε 12000) (MeOH/NaOH) (Derep). λ_{max} 274 (ε 9800) (MeOH) (Derep). λ_{max} 275 (MeOH/HCl) (Berdy).


Isogai, A. *et al.*, *Tet. Lett.*, 1986, **27**, 1161 (*isol, ir, uv, cryst struct, pmr, cmr*)

Japan. Pat., 1987, 87 72 691; *CA*, **107**, 38125k (*isol*)

Cyclocelabenzine

C-853

4,5,6,7,8,9,12,13-Octahydro-8-phenyl-2,13-methano-2H-2,7,11-benzotriazacyclopentadecine-1,10(3H,11H)-dione, 9CI
 [70535-05-6]



Absolute Configuration

$C_{23}H_{27}N_3O_2$ 377.485

Alkaloid from *Maytenus mossambicensis* var. *mossambicensis* (Celastraceae). Mp 180-183°. $[\alpha]_D^{25}$ +30.2 (c, 0.83 in $CHCl_3$).

N-Ac: Mp 260-265°. $[\alpha]_D^{22}$ +22.1 (c, 0.362 in $CHCl_3$).


Wagner, H. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 739 (*isol, uv, ir, pmr, cmr, ms, struct*)

Schultz, K. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 1295 (*synth, cryst struct, abs config*)

Cyclocinamide A

C-854

[194427-76-4]



Absolute Configuration


$C_{29}H_{33}BrClN_9O_8$ 750.992

Cyclic peptide antibiotic. Isol. from the sponge *Psammocinia* aff. *bulbosa*. Cytotoxic. Amorph. solid. $[\alpha]_D$ +29 (c, 0.1 in MeOH).

7,14-Diepimer, 36-chloro: **Cyclocinamide B**
[936720-48-8]
C₂₉H₃₂BrCl₂N₉O₈ 785.437
Isol. from the sponge *Corticium* sp. [α]_D²¹
+9.6 (c, 0.033 in MeOH). λ_{\max} 201 (log ϵ
3.53); 274 (log ϵ 2.43) (MeOH).

Clark, W.D. *et al.*, *J.A.C.S.*, 1997, **119**, 9285-
9286 (isol, pmr, cmr)
Grieco, P.A. *et al.*, *Tet. Lett.*, 1998, **39**, 8925-
8928 (synth, abs config)
Laird, D.W. *et al.*, *J. Nat. Prod.*, 2007, **70**, 741-
746 (Cyclocinamide B)
Rubio, B.K. *et al.*, *J. Nat. Prod.*, 2008, **71**,
1475-1478 (isol, abs config)


Cycloclausenamide **C-855**
5-Methyl-3,7-diphenyl-2-oxo-5-azabicyclo[2.2.1]heptan-6-one, 9CI
[103541-16-8]



Absolute
Configuration

C₁₈H₁₇NO₂ 279.338
Obt. as a pair of C-2 epimers. Isol. from
the leaves of *Clausena lansium* (wampee).
Prisms. Mp 164-166°. [α]_D^{24.5} -40 (c, 0.255
in MeOH).
Yang, M. *et al.*, *Phytochemistry*, 1988, **27**, 445-
450 (isol, uv, ir, pmr, cmr, ms, cryst struct)
Zhao, B. *et al.*, *Yaouxue Xuebao*, 2001, **36**, 373-
376 (cryst struct, conformn)

Cycloclavine **C-856**
[26057-57-8]




Absolute
congruation

C₁₆H₁₈N₂ 238.332
Alkaloid from the seeds of *Ipomoea*
hildebrandtii. Cryst. (MeOH or by subl.).
Mp 165-166°. [α]_D²⁰ +63 (c, 1 in CHCl₃).
[α]_D²⁰ +39 (c, 1 in Py).

Methiodide:
Cryst. (EtOH). Mp 246° dec. [α]_D²⁰ +30
(c, 0.9 in 50% EtOH aq.).

Stauffer, D. *et al.*, *Tetrahedron*, 1969, **25**,
5879-5887 (isol, uv, ir, pmr, cryst struct)
Ince, M. *et al.*, *Tetrahedron*, 2008, **64**, 2924-
2929 (synth)


Cyclodercitine **C-857**
[125236-61-5]



C₁₉H₁₄N₃S[⊕] 316.406
Struct. revised in 1992. Related to *N*-
Deacetylkuanoniamine D, D-93. Alka-
loid from the marine sponges *Dercitus* sp.
and *Stelletta* sp. Inhibits proliferation of
P388 murine leukaemia cells *in vitro*. Blue
powder (as chloride). Mp 298° (chloride).
 λ_{\max} 245 (ϵ 13800); 307 (ϵ 16900); 361 (ϵ
3900) (MeOH) (Derep).


9,10-Didehydro, *N*-de-Me: **Dehydronor-**
cyclodercitine. Nordehydrocycloderciti-
tine
[141224-24-0]
C₁₈H₉N₃S 299.355
Isol. from the ascidian *Aplidium* cf.
cratiferum. Yellow solid. λ_{\max} 256 (log
 ϵ 4.37); 293 (log ϵ 4.11); 338 (log ϵ
3.63); 360 (log ϵ 3.67) (MeOH).
Gunawardana, G.P. *et al.*, *Tet. Lett.*, 1989, **30**,
4359-4362 (isol, uv, pmr, cmr)
Gunawardana, G.P. *et al.*, *J.O.C.*, 1992, **57**,
1523-1526 (struct)
Agrawal, M.S. *et al.*, *Nat. Prod. Res.*, 2007, **21**,
782-786 (Dehydronorcyclodercitine)

Cyclodidemnamide **C-858**
[170894-38-9]




C₃₄H₄₃N₇O₅S₂ 693.89
Cyclic peptide antibiotic. Stereochem.
revised in 1998. Isol. from *Didemnum*
molle. Weak cytotoxic agent. Solid. Sol.
MeOH. Mp 114-118°. [α]_D +128.8 (c, 2.6
in MeOH). λ_{\max} 202 (log ϵ 4.46); 248 (log
 ϵ 4.04) (MeOH).
Toske, S.G. *et al.*, *Tet. Lett.*, 1995, **36**, 8355-
8358 (isol, uv, ir, pmr, cmr)
Norley, M.C. *et al.*, *Tet. Lett.*, 1998, **39**, 3087-
3090 (synth, struct)
Boden, C.D.J. *et al.*, *J.C.S. Perkin 1*, 2000, 883-
888 (synth, struct)

Cyclodidemnamide B **C-859**
[446821-87-0]




C₃₂H₄₇N₇O₆S₂ 689.899
Isol. from *Didemnum molle*. Amorph.
solid. [α]_D +8.6 (c, 0.28 in CHCl₃).
Arrault, A. *et al.*, *Tet. Lett.*, 2002, **43**, 4041-
4044 (isol, synth, pmr, cmr)

Cyclodidemnerinol **C-860**
[276673-10-0]



C₃₈H₆₆N₂O₁₉S₃ 951.139
Isol. from the ascidian *Didemnum gutta-*
tum. HIV-1 integrase inhibitor. Oil (as
tri-Na salt). [α]_D -26.6 (tri-Na salt). λ_{\max}
210 (ϵ 4500) (MeOH).
Mitchell, S.S. *et al.*, *Org. Lett.*, 2000, **2**, 1605-
1607

Cyclo[2-(1,1-dimethyl-2-propenyl)tryptophyl]prolyl **C-861**
Deoxybrevianamide E
[34610-68-9]



C₂₁H₂₅N₃O₂ 351.447
Nomenclature confused; many different
names possible. The name Deoxybreviana-
midamide E is a misnomer. Isol. from
Aspergillus ustus and *Penicillium italicum*.
Redn. prod. of Brevianamide E, B-290.
Powder (C₆H₆). [α]_D²² -59 (c, 1.2 in
CHCl₃).


12,13-Didehydro: **Cyclo(2,3-didehydro-**
prolyl)[2-(1,1-dimethyl-2-propenyl)-
tryptophyl]. 12,13-Dehydroprolyl-2-
(1,1-dimethylallyl)tryptophyl diketopi-
perazine
[41222-64-4]
C₂₁H₂₃N₃O₂ 349.432
Metab. of *Aspergillus ustus* and *Peni-*
cillium italicum. Powder. [α]_D²² -38 (c,
1.3 in CHCl₃).

[53892-61-8]

Birch, A.J. *et al.*, *Tetrahedron*, 1970, **26**, 2329
(synth, uv, ir, pmr)
Steyn, P.S. *et al.*, *Tetrahedron*, 1973, **29**, 107
(isol, deriv)
Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1981, 959
(synth, uv, ir, pmr)
Ritchie, R. *et al.*, *Tetrahedron*, 1981, **37**, 4295
(synth, uv, ir, pmr)
Schkeryantz, J.M. *et al.*, *J.A.C.S.*, 1999, **121**,
11964-11975 (synth)
Cole, R.J. *et al.*, *Handbook of Toxic*
Fungal Metabolites, Academic Press, 1981,
453

Cyclodopa C-862

2,3-Dihydro-5,6-dihydroxy-1H-indole-2-carboxylic acid, 9CI. *Leucodopachrome* [18791-20-3]



C₉H₉NO₄ 195.174

(S)-form [18766-67-1]

Degradn. prod. of Betanidin, B-110. Intermed. in formation of 2,3,5,6-Tetrahydro-5,6-dioxo-1H-indole-2-carboxylic acid from Levodopa in A-745. $[\alpha]_D^{25}$ -92 (c, 1.65 in 20% HCl).

5-O-β-D-Glucopyranoside: **Cyclodopa glucoside**

[71242-23-4]
C₁₅H₁₉NO₉ 357.316

Constit. of red beet cabbage (*Beta vulgaris* var. *rubra*). Prob. intermed. in biosynth. of Betanidin, B-110. Mp 159.5-162.5°. $[\alpha]_D^{20.5}$ -110.7 (H₂O, pH 3). pK_{a1} 1.58; pK_{a2} 4.75; pK_{a3} 9.42.

Tri-Ac, Me ester: Mp 94-96° (after drying). $[\alpha]_D^{20}$ -61 (c, 1.03 in CHCl₃).

N-(4-Hydroxy-E-cinnamoyl), 6-O-β-D-glucopyranoside: **Oleracein A** [872100-54-4]

C₂₄H₂₅NO₁₁ 503.462

Alkaloid from *Portulaca oleracea* (purslane). Yellow powder. λ_{max} 304 (sh) (log ε 4.01); 336 (log ε 4.07) (MeOH).

N-(4-Hydroxy-3-methoxy-E-cinnamoyl), 6-O-β-D-glucopyranoside: **Oleracein B** [872100-55-5]

C₂₅H₂₇NO₁₂ 533.488

Alkaloid from *Portulaca oleracea* (purslane). Yellow powder.

N-[4-(β-D-Glucopyranosyloxy)-E-cinnamoyl], 6-O-β-D-glucopyranoside: **Oleracein C** [872100-56-6]

C₃₀H₃₅NO₁₆ 665.604

Alkaloid from *Portulaca oleracea* (purslane). Yellow powder. $[\alpha]_D^{26}$ -83.7 (c, 0.35 in H₂O). λ_{max} 306 (log ε 4.32); 335 (log ε 4.28) (MeOH).

N-[4-(β-D-Glucopyranosyloxy)-3-methoxy-E-cinnamoyl], 6-O-β-D-glucopyranoside: **Oleracein D** [872100-57-7]

C₃₁H₃₇NO₁₇ 695.63

Alkaloid from *Portulaca oleracea* (purslane). Yellow powder. $[\alpha]_D^{26}$ +263.8 (c, 0.15 in H₂O). λ_{max} 239 (log ε 4.13); 295 (log ε 4.13); 335 (log ε 4.2) (H₂O).

(±)-form

Tri-Ac, Me ester: Mp 157.5-158.5°.

(ξ)-form

O⁵,N,N-Tri-Me: [165329-95-3]

C₁₂H₁₅NO₄ 237.255


Metab. of the Caribbean sponge *Pseudoceratina crassa*. $[\alpha]_D^{25}$ -12 (c, 0.001 in MeOH). Zwitterion.

Wyler, H. et al., *Helv. Chim. Acta*, 1962, **45**, 638-640; 1968, **51**, 1476-1494 (*S-form, synth. isol. struct*)

Wyler, H. et al., *Helv. Chim. Acta*, 1984, **67**, 1348-1355 (*5-glucoside*)

Ciminiello, P. et al., *J. Nat. Prod.*, 1995, **58**, 689-696 (*tri-Me*)

Xiang, L. et al., *Phytochemistry*, 2005, **66**, 2595-2601 (*Oleraceins A-D*)

**14,18-Cyclo-21,23-epoxyyap-
tirucallane-3,7,21,24,25-pentol** C-863

(3α,7α,21R,23R,24S)-form

C₃₀H₅₀O₆ 506.721

(3β,7α,21R,23R,24S)-form

21-Me ether, 3-O-(2-methylaminobenzoyl): [413615-73-3]

C₃₉H₅₉NO₇ 653.898

Constit. of *Raulinoa echinata*. Amorph. solid. Mp 144-145°. $[\alpha]_D^{25}$ +15.5 (c, 0.01 in CH₂Cl₂). λ_{max} 222 (log ε 4.16); 253 (log ε 3.78); 353 (log ε 3.64) (MeOH).

(3β,7α,21S,23R,24S)-form

21-Me ether, 3-O-(2-methylaminobenzoyl): [413614-52-5]

Constit. of *Raulinoa echinata*. Amorph. solid. Mp 218-220°. $[\alpha]_D^{25}$ +25.5 (c, 0.03 in CH₂Cl₂). λ_{max} 224 (log ε 4.01); 254 (log ε 3.6); 354 (log ε 3.44) (MeOH).

(3β,7α,21ξ,23R,24S)-form

3-O-(2-Methylaminobenzoyl): [413615-74-4]


C₃₈H₅₇NO₇ 639.871

Constit. of *Raulinoa echinata*. Amorph. solid. Mp 176-178°. $[\alpha]_D^{25}$ +13.1 (c, 0.01 in CH₂Cl₂). λ_{max} 222 (log ε 4.07); 254 (log ε 3.64); 354 (log ε 3.5) (MeOH).

Biavatti, M.W. et al., *J. Nat. Prod.*, 2002, **65**, 562-565 (*Raulinoa echinata* constits)

Cyclofoveoglin C-864

[948312-51-4]



C₃₈H₃₈N₂O₈ 650.727


Alkaloid from the leaves of *Aglaiia foveolata*. Pale yellow powder. $[\alpha]_D^{20}$ -51.7 (c, 0.41 in CHCl₃). λ_{max} 203 (log ε 4.74);

214 (sh) (log ε 4.64) (MeOH).

Salim, A.A. et al., *Tetrahedron*, 2007, **63**, 7926-7934 (*isol, pmr, cmr*)

Cyclo(glycylprolyl) C-865

Hexahydropyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. 3,6-Dioxohexahydropyrrolo[1,2-a]pyrazine (incorr.). Cyclo(prolylglycyl). 1,4-Diazabicyclo[4.3.0]nonane-2,5-dione [19179-12-5]



C₇H₁₀N₂O₂ 154.168

(S)-form

L-form

[3705-27-9]

Constit. of roasted cocoa nibs and coffee. Prod. by cultures of *Fusarium* spp. Isol. from the echinoderm *Luidia clathrata*. Constit. of *Pinellia pedatisecta*. Prod. by the marine bacteria *Ruegeria* sp. SDC-1 and *Pseudoalteromonas luteoviolacea*. Cryst. (MeOH/Me₂CO). Mp 216-218°. $[\alpha]_D^{20}$ -217.4 (c, 7.5 in H₂O).

(±)-form [62057-47-0]

Cryst. (MeOH/Me₂CO). Mp 216-218°.

[96193-26-9]

Vičar, J. et al., *Coll. Czech. Chem. Comm.*, 1972, **37**, 4060; 1973, **38**, 1940; 1957 (*synth. ir, pmr, bibl*)

Blaha, K. et al., *Coll. Czech. Chem. Comm.*, 1972, **37**, 4660 (*synth*)

White, E.P. et al., *N.Z. J. Sci.*, 1972, **15**, 178; *CA*, 77, 98731b (*isol*)

Blaha, K. et al., *Tet. Lett.*, 1972, 1437 (*synth*)
Pettit, G.R. et al., *Experientia*, 1973, **29**, 521 (*isol, ms, struct*)

Von Dreele, R.B. et al., *Acta Cryst. B*, 1975, **31**, 966 (*cryst struct*)

Johnstone, R.A.W. et al., *J.C.S. Perkin 2*, 1975, 1297 (*ms*)

Siemion, I.Z. et al., *Org. Magn. Reson.*, 1976, **8**, 432 (*cmr*)

Qin, W. et al., *Zhongcaoyao*, 1984, **15**, 490; *CA*, **102**, 109774f (*isol*)

de Costa, B.R. et al., *J. Med. Chem.*, 1993, **36**, 2311 (*synth, pmr*)


Jiang, Z. et al., *Nat. Prod. Lett.*, 2000, **14**, 435-440 (*isol, pmr, cmr*)

Mitova, M. et al., *J. Nat. Prod.*, 2004, **67**, 1178-1181 (*isol*)

Stark, T. et al., *J. Agric. Food Chem.*, 2005, **53**, 7222-7231 (*isol, ms*)

Cycloguineense A C-866

[486449-08-5]



C₂₃H₂₉NO₃ 367.487
Alkaloid from *Piper guineense*.

1,8a-Diepimer: Cycloguineense B


[486449-09-6]

C₂₃H₂₉NO₃ 367.487

Alkaloid from *Piper guineense*.

Adesina, S.K. et al., *Pharmazie*, 2003, **58**, 423-425 (isol, ms)

Cyclohaliclونamines C-867



Cyclohaliclونamide	A	n = 1
"	B	n = 2
"	C	n = 3
"	D	n = 4
"	E	n = 5

Alkaloids from the sponge *Haliclona* sp.

Cyclohaliclونamine A

C₃₀H₄₄N₂²⁺ 432.691

Cyclohaliclونamine B

C₄₅H₆₆N₃³⁺ 649.037

Cyclohaliclونamine C

C₆₀H₈₈N₄⁴⁺ 865.382


Cyclohaliclونamine D

Cyclohaliclونamine E

Teruya, T. et al., *J. Nat. Prod.*, 2006, **69**, 135-137 (isol, pmr, cmr, ms)

Cyclohexylamine C-868

Cyclohexanamine, 9CI. Aminocyclohexane. Hexahydroaniline [108-91-8]



C₆H₁₃N 99.175


Isol. from *Toddalia asiatica*. Gut microflora metabolite of Cyclohexylsulfamic acid. Corrosion inhibitor; food contaminant arising from its use as a boiler water additive. Liq. with strong amine odour. Misc. H₂O, org. solvs. d₂₅²⁵ 0.87. Fp -18. Bp 134° Bp₇₁ 65° Bp₂₀ 36.4°. n_D²⁵ 1.4565. Strong base. Forms azeotrope with H₂O contg. 55.8% H₂O, Bp 96.4°.

► Flammable, fl. p. 21/31°, autoignition temp. 260°. Vapour irritates eyes and respiratory tract. Liquid is corrosive to skin and eyes. Toxic by skin absorption causing gastrointestinal effects. LD₅₀ (rat, oral) 156 mg/kg. LD₅₀ (rbt, skin) 277 mg/kg. Exp. reprod. and teratogenic effects. OES: long-term 10 ppm (Sk). GX0700000

Tsai, I.-L. et al., *Phytochemistry*, 1998, **48**, 1377-1382 (isol, pmr)

2-Cyclohexylethylamine C-869

Cyclohexaneethanamine, 9CI. (2-Aminoethyl)cyclohexane [4442-85-7]



C₈H₁₇N 127.229

Alkaloid from *Acacia rigidula*. Bp 188-189° Bp₂₅ 83-85°.

Hydrochloride: [5471-55-6]

Cryst. (EtOH aq.). Mp 258°.

N-Me: *N-Methyl-2-cyclohexylethylamine*

[62141-38-2]

C₉H₁₉N 141.256

Alkaloid from *Acacia rigidula*. Oil. Bp₇ 71-72°.

N-Me, hydrochloride:

Cryst. (2-propanol/Et₂O). Mp 171-172°.

Zenitz, B.L. et al., *J.A.C.S.*, 1947, **69**, 1117-1121 (synth)

Kindler, K. et al., *Annalen*, 1948, **560**, 215-221; 1961, **644**, 23-30 (synth)

Wilson, K.R. et al., *J.O.C.*, 1959, **24**, 1046-1051 (synth)


Scheinbaum, M.L. et al., *J.O.C.*, 1970, **35**, 2785-2790 (synth)

New, J.S. et al., *Synthesis*, 1983, 388-389 (synth, ir, pmr, cmr)

Clement, B.A. et al., *Phytochemistry*, 1998, **49**, 1377-1380 (isol)

Cyclo(homoleucylpropyl) C-870

Hexahydro-3-(3-methylbutyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Homoleucylpropyl anhydride



(3*S*,8*aR*)-form

C₁₂H₂₀N₂O₂ 224.302

(3*S*,8*aR*)-form [62835-71-6]

Found in the sclerotia and saprophytic culture of ergot fungus *Claviceps purpurea*.

(3*ξ*,8*aξ*)-form [118243-97-3]


Isol. from liquid cultures of *Alternaria alternata* and prod. by a *Streptomyces* sp.

Ohmomo, S. et al., *Nippon Nogei Kagaku Kaishi*, 1976, **50**, 543-546; *CA*, **86**, 190303x (isol, *Claviceps*, synth)

Stierle, A.C. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 8008-8011 (isol, *Alternaria*)

Cyclo(4-hydroxypropylleucyl) C-871

Hexahydro-7-hydroxy-3-(2-methylpropyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Hydroxyproline leucine anhydride [308368-99-2]



(3*R*,7*R*,8*aS*)-form

C₁₁H₁₈N₂O₃ 226.275

(3*R*,7*R*,8*aS*)-form

D,L-form

[208850-33-3]

Isol. from a marine bacterium associated with a *Palythoa* sp. Solid. [α]_D²⁵ +14.7 (c, 0.34 in MeOH). λ_{max} 230 (MeOH).

(3*S*,7*R*,8*aR*)-form [205648-55-1]

Metab. of the marine yeast *Aureobasidium pullulans* and two marine bacteria associated with the sponge *Ircinia variabilis*. Amorph. solid. [α]_D²⁵ +40.5 (c, 1 in MeOH).

(3*S*,7*R*,8*aS*)-form

7-Hydroxyguacidin W. L-L-form

[115006-86-5]

Isol. from rabbit skin tissue. Also from a marine bacterium associated with a *Palythoa* sp. and two marine bacteria associated with the sponge *Ircinia variabilis*. Plant growth regulator. Mp 178-179°. [α]_D²⁸ -148.2 (c, 1 in H₂O). λ_{max} 232 (MeOH).

Ienaga, K. et al., *Tet. Lett.*, 1987, **28**, 1285-1286 (3*S*,7*R*,8*aS*-form, isol, pmr, cmr, struct)

Shigemori, H. et al., *J. Nat. Prod.*, 1998, **61**, 696-698 (3*S*,7*R*,8*aR*-form, isol, ir, pmr, cmr)


Cronan, J.M. et al., *Nat. Prod. Lett.*, 1998, **11**, 271-278 (*Palythoa* sp. isol, uv, pmr, cmr)

De Rosa, S. et al., *Biomol. Eng.*, 2003, **20**, 311-316 (*Ircinia variabilis* constits)

Mitova, M. et al., *Z. Naturforsch., C*, 2003, **58**, 740-745 (*Ircinia variabilis* constiti)

Cyclo(2-hydroxypropylphenylalanyl) C-872

Hexahydro-8a-hydroxy-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI



(3*S*,8*aR*)-form

C₁₄H₁₆N₂O₃ 260.292

(3*S*,8*aR*)-form

D,L-form

Prod. by the marine-derived *Chromocleista* sp. strain R721. [α]_D²⁵ -26.1 (c, 0.03 in MeOH). λ_{max} 198 (log ε 3.5); 257 (log ε 0.3) (MeOH).

8a-Deoxy, 8,8a-didehydro: 2,3,6,7-Tetrahydro-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI

C₁₄H₁₄N₂O₂ 242.277

Prod. by *Chromocleista* sp. strain R721. $[\alpha]_D^{25}$ -31.9 (c, 0.05 in MeOH). Dec. prod. of parent. λ_{\max} 200 (log ϵ 3.7); 257 (log ϵ 0.5); 330 (log ϵ 0.2) (MeOH).


(3S,8aS)-form*L,L*-form

Prod. by the marine-derived *Chromocleista* sp. strain R721. λ_{\max} 200 (log ϵ 3.7); 257 (log ϵ 0.4) (MeOH).

Park, Y.C. *et al.*, *J. Nat. Prod.*, 2006, **69**, 580-584 (isol, pmr, cmr)

Cyclo(4-hydroxypropylphenylalanyl) C-873

Hexahydro-7-hydroxy-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. 3-Benzylhexahydro-7-hydroxypyrrolo[1,2-a]pyrazine-1,4-dione



$C_{14}H_{16}N_2O_3$ 260.292

(3R,7S,8aR)-form [631922-77-5]

Prod. by the marine bacterial strains CF-20 and C-148 obt. from the mollusc *Pecten maximus*. Active against *Vibrio anguillarum*. $[\alpha]_D$ +7.3 (c, 0.02 in MeOH). Error in struct. diag. in ref.

(3S,7R,8aR)-form [205648-54-0]

Metab. of the marine yeast *Aureobasidium pullulans* and two strains of microorganism isol. from the marine sponge *Ircinia variabilis*. Amorph. solid. $[\alpha]_D^{24}$ +34.7 (c, 1 in MeOH). λ_{\max} 208 (ϵ 28000); 281 (ϵ 2900) (MeOH).

(3S,7R,8aS)-form [118477-06-8]

Metab. of an undescr. Fijian marine sponge (Jaspidae), the marine bacterium *Pseudoalteromonas luteoviolacea* and two strains of microorganism isol. from the marine sponge *Ircinia variabilis*. Plant growth regulator. Viscous oil. $[\alpha]_D^{20}$ -6.7 (c, 0.016 in MeOH).

Adamczeski, M. *et al.*, *J.A.C.S.*, 1989, **111**, 647-654 (isol, ir, pmr, cmr, ms, struct, Jaspidae constit)

Shigemori, H. *et al.*, *J. Nat. Prod.*, 1998, **61**, 696-698 (*Aureobasidium pullulans* constit)

Cronan, J.M. *et al.*, *Nat. Prod. Lett.*, 1998, **11**, 271-278 (*3S,7R,8aS*-form, isol)


Jiang, Z. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 435-440 (*Pseudoalteromonas luteoviolacea* constit)

De Rosa, S. *et al.*, *Biomol. Eng.*, 2003, **20**, 311-316 (*Ircinia variabilis* constit)

Fdhila, F. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1299-1301 (*3R,7S,8aR*-form, isol)

Cyclo(4-hydroxypropyltyrosyl) C-874

Hexahydro-7-hydroxy-3-[(4-hydroxyphenyl)methyl]pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI



(3S,7R,8aR)-form

$C_{14}H_{16}N_2O_4$ 276.291

(3S,7R,8aR)-form [813461-20-0]

Prod. by a *Ruegeria* sp. isol. from the sponge *Suberites domuncula*. $[\alpha]_D$ +82.3 (c, 0.001 in MeOH).


(3S,7R,8aS)-form [813461-21-1]

Prod. by a *Ruegeria* sp. isol. from *Suberites domuncula*. $[\alpha]_D$ -10.4 (c, 0.003 in MeOH).

Mitova, M. *et al.*, *Mar. Biotechnol.*, 2004, **6**, 95-103 (isol)

Cyclo(isoleucylprolyl) C-875

Hexahydro-3-(1-methylpropyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI [58917-56-9]



$C_{11}H_{18}N_2O_2$ 210.275

(1'R,3R,8aS)-form*L,D*-allo-form

[61117-55-3]

Cryst. (hexane/Et₂O/Me₂CO). Mp 107-108°.

(1'S,3R,8aS)-form*L,D*-form

[61117-54-2]

Cryst. (Me₂CO). Mp 143-145°.

(1'S,3S,8aR)-form [162830-28-6]

Isol. from the sponge *Calyx* cf. *podatypa*. Amorph. powder. $[\alpha]_D^{23}$ +98 (c, 0.1 in EtOH). Config. of nat. prod. revised in 1998.

(1'S,3S,8aS)-form [57089-60-8]

Isol. from *Calyx* cf. *podatypa*. Prod. by marine-derived *Pseudomonas aeruginosa* and *Vibrio parahaemolyticus*. Constit. of *Schizandra chinensis*. Amorph. solid. $[\alpha]_D^{20}$ -173.5 (c, 0.16 in EtOH).

(1'ε,3R,8aR)-form [631922-76-4]

Prod. by the marine bacterial strains CF-20 and C-148 isol. from cultures of the larvae of the mollusc *Pecten maximus*. Active against *Vibrio anguillarum*. $[\alpha]_D$ +168.1 (c, 0.13 in EtOH).

Bycroft, B.W. *et al.*, *Chem. Comm.*, 1975, 988-989 (synth)

Young, P.E. *et al.*, *J.A.C.S.*, 1976, **98**, 5358-5364; 5365-5371 (synth, conformn)

Suzuki, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 233-237 (synth)

Adamczeski, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 201-208 (isol, pmr, cmr)

Jayatilake, G.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 293-296 (isol)

Bull, S.D. *et al.*, *J.C.S. Perkin 1*, 1998, 2313-2320 (synth, pmr)

Ginz, M. *et al.*, *J. Agric. Food Chem.*, 2000, **48**, 3528-3532 (isol, coffee)


Fdhila, F. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1299-1301 (isol, pmr, cmr)

Stark, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 7222-7231 (*1S,3S,8aS*-form, isol, ms)

Pedras, M.S.C. *et al.*, *Z. Naturforsch., C*, 2005, **60**, 717-722 (isol, synth, pmr, cmr)

Cyclo(leucylprolyl)**C-876**

Hexahydro-3-(2-methylpropyl)pyrrolo[1,2-a]pyrazine-1,4-dione. Prolylleucyl-diketopiperazine. Gancidin W. Maculosin 6 [5654-86-4]



$C_{11}H_{18}N_2O_2$ 210.275

(3R,8aR)-form*D,D*-form

[162872-80-2]

Prod. by the marine bacterial strains CF-20 and C-148 obt. from molluscs. Amorph. solid. $[\alpha]_D$ +128.3 (c, 0.11 in EtOH).

(3R,8aS)-form [36238-67-2]

Cryst. Sol. MeOH, EtOAc. Mp 148-149° (119-122°). $[\alpha]_D^{25}$ -105.5 (c, 0.5 in EtOH).

(3S,8aR)-form [32510-93-3]

Isol. from the sponges *Calyx* cf. *podatypa* and *Stelletta clavosa*. Solid. $[\alpha]_D^{25}$ +35 (c, 0.11 in EtOH).

(3S,8aS)-form*L,L*-form

[2873-36-1]

Prod. by *Streptomyces gancidicus*, *No-cardia restricta*, *Candida albicans*, *Guignardia loricata* and several *Ceratomyxystis* spp. Also from the sponges *Calyx* cf. *podatypa* and *Tedania ignis*. Shows antitumour and antifungal props. Phytotoxin associated with fungal diseases of trees. Cryst. Mp 168-172° (158-159°). $[\alpha]_D^{25}$ -133 (c, 1 in EtOH) (lit. gives a temp. range). $[\alpha]_D^{21}$ -144 (c, 0.5 in H₂O).

▶ LD₅₀ (mus, ivn) 80 mg/kg. UY8708800

[43041-29-8, 19943-30-7]

Kodaira, Y. *et al.*, *Agric. Biol. Chem.*, 1961, **25**, 261

Nitecki, D.E. *et al.*, *J.O.C.*, 1968, **33**, 864 (synth)

Siemion, I.Z. *et al.*, *Org. Magn. Reson.*, 1971, **3**, 545 (synth, nmr)

Vicar, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 4060; 1973, **38**, 1940 (synth, ir, pmr)

Karle, I.L. *et al.*, *J.A.C.S.*, 1972, **94**, 81 (cryst struct)

Takahashi, K. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 927

Bycroft, B.W. *et al.*, *Chem. Comm.*, 1975, 988 (synth)

Davies, D.B. *et al.*, *J.C.S. Perkin 2*, 1976, 187 (conformn)

Pancoska, P. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **44**, 1296 (cd)

Jain, T.C. *et al.*, *Heterocycles*, 1977, **7**, 341 (isol, struct, uv, ms)

Bjoerkman, S. *et al.*, *J. Med. Chem.*, 1979, **22**, 931 (synth)

Suzuki, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 233 (synth)


Schmidtz, F.J. *et al.*, *J.O.C.*, 1983, **48**, 3941 (isol, ms, pmr, synth, bibl)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1986, **64**, 904 (isol)

Kricheldorf, H.R. *et al.*, *Magn. Reson. Chem.*, 1986, **24**, 21 (cmr)

- Adamczeski, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 201 (*isol, pmr, cmr, bibl*)
 Bull, S.D. *et al.*, *J.C.S. Perkin 1*, 1998, 2313-2320 (*config, biosynth*)
 Fdhila, F. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1299-1301 (*isol, pmr, cmr*)
 Wegerski, C.J. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 5631-5637 (*isol*)
 Stark, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 7222-7231 (*L,L-form, isol, ms*)
 Pedras, M.S.C. *et al.*, *Z. Naturforsch., C*, 2005, **60**, 717-722 (*isol, synth, pmr, cmr*)
 Li, Z. *et al.*, *Biochem. Syst. Ecol.*, 2008, **36**, 230-234 (*isol, pmr, cmr*)

Cyclomarazine A C-877
 [1019858-65-1]




$C_{23}H_{31}N_3O_3$ 397.516
 Related to Cyclo(leucyltryptophyl). Prod. by *Salinospora arenicola* CNS-205. Oil. $[\alpha]_D -9$ (c, 0.19 in MeOH). λ_{max} 225 (log ϵ 3.7); 285 (log ϵ 3) (MeOH).

N-De-Me: Cyclomarazine B
 [1019858-66-2]

$C_{22}H_{29}N_3O_3$ 383.489
 Prod. by *Salinospora arenicola* CNS-205. Oil. $[\alpha]_D -16$ (c, 0.03 in MeOH). λ_{max} 224 (log ϵ 3.6); 284 (log ϵ 3) (MeOH).

Schultz, A.W. *et al.*, *J.A.C.S.*, 2008, **130**, 4507-4516 (*isol, cd, pmr, cmr*)


Cyclomegistine C-878



$C_{18}H_{19}NO_7$ 361.351
 Alkaloid from the bark of *Sarcomelicope megistophylla*. Pale yellow cryst. (Me₂CO). Mp 146°. Opt. inactive. λ_{max} 318 (log ϵ 2.46); 332 (log ϵ 2.52) (MeOH).

Fokialakis, N. *et al.*, *Tet. Lett.*, 2001, **42**, 5323-5325 (*isol, uv, pmr, cmr, cryst struct*)

Cyclomegistine B C-879



$C_{17}H_{19}NO_6$ 333.34


(ξ)-form

Alkaloid from the bark of *Sarcomelicope*

megistophylla. λ_{max} 318 (log ϵ 2.49); 332 (log ϵ 2.55) (MeOH).

Mitaku, S. *et al.*, *Fitoterapia*, 2007, **78**, 169-170 (*isol, pmr, cmr, ms*)

Cyclo(methionylpropyl) C-880
Hexahydro-3-[2-(methylthio)ethyl]pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Cyclo(-prolylmethionyl)



$C_{10}H_{16}N_2O_2S$ 228.315

(3S,8aS)-form

L-L-form

[53049-06-2]


Prod. by *Pseudomonas aeruginosa* isol. from an Antarctic sponge. Amorph. solid. $[\alpha]_D^{20} -82.2$ (c, 0.14 in EtOH).

Padmanabhan, V.M. *et al.*, *Acta Cryst. C*, 1987, **43**, 349-351 (*cryst struct*)

Jayatilake, G.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 293-296 (*isol, synth, pmr, cmr*)

Cyclomethoxazine B C-881

[6908-24-3]



Absolute Configuration

$C_{28}H_{48}N_2O_2$ 444.699

C-4 config. revised in 1975. Alkaloid from *Buxus rolfiei* (Buxaceae). Mp 195°. $[\alpha]_D +54$ (c, 1 in CHCl₃).

N²⁰-Me: Buxmicrophylline D

[256236-37-0]

$C_{29}H_{50}N_2O_2$ 458.726

Alkaloid from *Buxus microphylla*. Needles. Mp 228-230°.


Khuong-Huu-Lainé, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 1216-1221 (*isol, pmr, ms, struct*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*cmr, struct*)

Du, J. *et al.*, *J. Asian Nat. Prod. Res.*, 1999, **1**, 239-244 (*Buxmicrophylline D*)

Cyclomicrobuxeine K C-882

3-(Dimethylamino)-14-methyl-4-methylene-9,19-cyclopregn-16-en-20-one, 9CI. Cyclomicrobuxeine. Anhydrobuxpiene [4409-26-1]



$C_{25}H_{37}NO$ 367.573

Alkaloid from *Buxus microphylla* (Buxaceae). Needles (MeOH). Mp 141-142°. $[\alpha]_D +126$ (c, 1.20 in CHCl₃). Possibly an artifact.

N-De-Me: Norcyclomicrobuxeine

[14747-70-7]

$C_{24}H_{35}NO$ 353.547

Alkaloid from leaves of *Buxus papillosa* (Buxaceae). Amorph. $[\alpha]_D +34$ (c, 1.90 in CHCl₃).

N-De-Me, N-formyl: N-Formylnorcyclomicrobuxeine
(incorr.)

[112926-18-8]

$C_{25}H_{35}NO_2$ 381.557

Alkaloid from leaves of *Buxus papillosa* (Buxaceae). Amorph. $[\alpha]_D +16$.

4β,23-Dihydro: 3-(Dimethylamino)-4,14-dimethyl-9,19-cyclopregn-16-en-20-one, 9CI. Cyclobuxomicreine K. Cyclobuxomicreine

[7669-79-6]

$C_{25}H_{39}NO$ 369.589

Alkaloid from *Buxus microphylla* (Buxaceae). Needles (EtOH). Mp 195-197°. $[\alpha]_D +37$ (c, 1.62 in CHCl₃). Possibly an artifact.


Nakano, T. *et al.*, *J.C.S. (C)*, 1966, 1412-1421 (*Cyclomicrobuxeine K, Cyclobuxomicreine*)

Choudhary, M.I. *et al.*, *J. Nat. Prod.*, 1987, **50**, 84-88 (*Norcyclomicrobuxeine*)

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1987, **53**, 496-497 (*N-Formylnorcyclomicrobuxeine*)

Cyclomicrophylline I C-883

3,20-Diamino-16-hydroxy-4,14-dimethyl-9,19-cyclopregn-6-ene-4-methanol, 9CI



$C_{24}H_{40}N_2O_2$ 388.592

The suffix letter in *Buxus* alkaloids systemically designates the *N*-subn. pattern. The parent compd., Cyclomicrophylline I, is unknown. C-4 configs. in this group of alkaloids revised in 1975. C-4 configs. reversed in 1975.

N²⁰,N²⁰-Di-Me, N³-benzoyl: 3-Benzoylcyclomicrophylline F. Buxidine F[†].

Buxidine. Buxizine

[18050-99-2]

$C_{33}H_{48}N_2O_3$ 520.754


Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 154-157°. $[\alpha]_D^{25} +67.5$ (c, 0.2 in CHCl₃). Mol. formula twice revised before struct. settled. Buxizine is a misprint. The name *N³-Benzoylcyclomicrophylline F* is recommended to avoid confusion.

N²⁰,N²⁰-Di-Me, N³-benzoyl, O¹⁶-Ac:

- Buxandrine F.** *Buxandrine. Semperviramidine*
[18051-00-8]
C₃₅H₅₀N₂O₄ 562.791
Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 289-290° dec. Mol. formula incorrectly given as C₃₅H₅₂N₂O₄ in one paper. Semperviramidine (amorph., [α]_D²⁰ + 33°), is shown with the opposite C-4 config. but this appears to be incorrect which would make it identical with Buxandrine F. (See comment above).
- N³,N³,N²⁰-Tri-Me: Cyclomicrophylline B.** *Cyclobalebuxine*
[3556-11-4]
C₂₇H₄₆N₂O₂ 430.673
Alkaloid from *Buxus microphylla*, *Buxus balearica* and *Buxus malayana* (Buxaceae). Cryst. (EtOH). Mp 254° (251-252°). [α]_D -65 (CHCl₃).
- N³,N³,N²⁰-Tri-Me, N²⁰,O¹⁶-di-Ac:** Mp 202-203°. [α]_D -151 (CHCl₃).
- N³,N²⁰,N²⁰-Tri-Me: Cyclomicrophylline C**
[3296-09-1]
C₂₇H₄₆N₂O₂ 430.673
Alkaloid from *Buxus microphylla* and *Buxus sempervirens* (Buxaceae). Mp 283-284°. [α]_D -40 (CHCl₃).
- N³,N²⁰,N²⁰-Tri-Me, N³-benzoyl: Cyclomicrosine.** *N-Benzoylcyclomicrophylline C. Cyclomicrosine C*
[10088-26-3]
C₃₄H₅₀N₂O₃ 534.781
Alkaloid from *Buxus microphylla* (Buxaceae). Needles (Me₂CO/hexane). Mp 282-284°. [α]_D -33 (CHCl₃).
- N-Tetra-Me: Cyclomicrophylline A**
[2124-69-8]
C₂₈H₄₈N₂O₂ 444.699
Alkaloid from *Buxus microphylla*, *Buxus balearica*, *Buxus sempervirens* and *Buxus malayana* (Buxaceae). Mp 232-233°. [α]_D²⁰ -40 (CHCl₃).
- N-Tetra-Me, 31-Ac: 31-Acetylcyclomicrophylline A**
[135626-68-5]
C₃₀H₅₀N₂O₃ 486.737
Alkaloid from *Buxus sempervirens* (Buxaceae). [α]_D -92 (CHCl₃).
- N-Tetra-Me, O¹⁶-benzoyl: Cyclomicrophyllidine A**
[3296-10-4]
C₃₅H₅₂N₂O₃ 548.807
Alkaloid from *Buxus microphylla* (Buxaceae). Amorph. [α]_D -160 (CHCl₃).
- N-Tetra-Me, dibenzoyl:**
Amorph. Mp 105-113°.
- 6,7-Dihydro: 3,20-Diamino-16-hydroxy-4,14-dimethyl-9,19-cyclopregnan-4-methanol.** *Dihydrocyclocyclomicrophylline I*
- 6,7-Dihydro, N³,O¹⁶-di-Ac, N²⁰-formyl: Cycloprotobuxinamine**
[188947-70-8]
C₂₉H₄₆N₂O₅ 502.693
Alkaloid from *Buxus microphylla* (Buxaceae).
- 6,7-Dihydro, N²⁰,N²⁰-di-Me: Dihydrocyclocyclomicrophylline F.** *Cyclobuxidine F*
[14155-69-2]
C₂₆H₄₆N₂O₂ 418.662
Alkaloid from *Buxus microphylla* (Buxaceae). Mp 260°. [α]_D +4.6 (CHCl₃).
- 6,7-Dihydro, N²⁰,N²⁰-di-Me, N³-tigloyl, O¹⁶-Ac: Hyrcamine**
C₃₃H₅₄N₂O₄ 542.801
Alkaloid from the leaves of *Buxus hyrcana*. Amorph. solid. [α]_D²⁴ +30 (c, 0.07 in CHCl₃). λ_{max} 202 (log ε 2.72) (MeOH).
- 6,7-Dihydro, N²⁰,N²⁰-di-Me, N³-benzoyl: Buxepidine.** *N³-Benzoyldihydrocyclocyclomicrophylline F*
[17934-60-0]
C₃₃H₅₀N₂O₃ 522.77
Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 292-294° (278-279°). [α]_D -20 (CHCl₃). [α]_D +19.
- 6,7-Dihydro, N³,N²⁰,N²⁰-tri-Me: Dihydrocyclocyclomicrophylline C.** *Cyclorolfiebuxine C*
C₂₇H₄₈N₂O₂ 432.688
Alkaloid from *Buxus rolfiei* (Buxaceae). Cryst. (Me₂CO). Mp 256° Mp 265° (synth.). Methylation with HCHO gives Cyclobuxoxazine A.
- 6,7-Dihydro, N-tetra-Me: Dihydrocyclocyclomicrophylline A.** *Cyclorolfiebuxine A. N-Methylcyclorolfiebuxine C*
[4003-53-6]
C₂₈H₅₀N₂O₂ 446.715
Alkaloid from *Buxus microphylla* (Buxaceae). Needles (MeOH). Mp 271-272° (266°). [α]_D +37 (CHCl₃).
- 6,7-Dihydro, N-tetra-Me, O¹⁶-benzoyl: Dihydrocyclocyclomicrophyllidine A**
[3668-22-2]
C₃₅H₅₄N₂O₃ 550.823
Alkaloid from *Buxus microphylla* (Buxaceae). Amorph. [α]_D -33 (CHCl₃).
- 6,7-Dihydro, 16-ketone: 3,20-Diamino-4-hydroxymethyl-4,14-dimethyl-9,19-cyclopregnan-16-one.** *Buxazidine I*
- 6,7-Dihydro, 16-ketone, N³,N³,N²⁰-tri-Me: Buxazidine B**
[15208-57-8]
C₂₇H₄₆N₂O₂ 430.673
Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 234-236°. [α]_D -31 (c, 0.1 in CHCl₃). C-4 config. revised here from original assignment.
- 4-Epimer, N²⁰,N²⁰-di-Me, N³-(2-methylpropanoyl): Buxmicrophylline C**
[256236-36-9]
C₃₀H₅₀N₂O₃ 486.737
Alkaloid from *Buxus microphylla*. Needles. Mp 229-230°. C-4 stereo. in doubt.
- Stereoisomer, 6,7-dihydro, N-tetra-Me: Isodihydrocyclocyclomicrophylline A**
C₂₈H₅₀N₂O₂ 446.715
Alkaloid from *Buxus sempervirens*. Cryst. (EtOH). Mp 215-217°. [α]_D -64.3 (c, 0.527 in CHCl₃). Uncertain struct. Suggested to be epimeric with Dihydrocyclocyclomicrophylline A at C-4 and/or C-20. CAS No. not found.
[1355-00-6]
Herlem-Gaulier, D. et al., *Bull. Soc. Chim. Fr.*
- 1965, 657-668 (*Cyclocyclomicrophylline B, ir, ms, pmr, struct*)
Nakano, T. et al., *J.C.S.*, 1965, 4512-4537 (*Buxus microphylla constits*)
Khuong-Huu-Lainé, F. et al., *Bull. Soc. Chim. Fr.*, 1966, 758-762; 1216-1221 (*Cyclocyclomicrophylline B. Dihydrocyclocyclomicrophylline C, isol, ms, ir, pmr*)
Nakano, T. et al., *J.C.S.(C)*, 1966, 1412-1421 (*Cyclocyclomicrosine*)
Döpke, W. et al., *Naturwissenschaften*, 1967, 54, 200 (*Buxazidine B*)
Döpke, W. et al., *Tet. Lett.*, 1967, 4247-4249 (*Buxidine F, Buxandrine F*)
Sangare, M. et al., *Tet. Lett.*, 1975, 1791-1794 (*cmr, config*)
Atta-ur-Rahman, et al., *Phytochemistry*, 1988, 27, 2367-2368; 1991, 30, 1295-1298 (*31-Acetylcyclocyclomicrophylline A, Buxandrine F*)
Khodzhev, B.U. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1994, 29, 812-813 (*Isodihydrocyclocyclomicrophylline A*)
Du, J. et al., *Zhivü Xuebao (Acta Bot. Sin.)*, 1996, 38, 483-488; *CA*, 126, 274768k (*Cycloprotobuxinamine*)
Du, J. et al., *J. Asian Nat. Prod. Res.*, 1999, 1, 239-244 (*Buxmicrophylline C*)
Choudhary, M.I. et al., *Chem. Biodiversity*, 2006, 3, 1039-1052 (*Hyrcamine*)


Cyclomicurarine L C-884

20-(Dimethylamino)-16-hydroxy-4,4,14-trimethyl-9,19-cyclopregnan-3-one, 9CI.
Cyclomicurarine
[7671-39-8]



- C₂₆H₄₃NO₂ 401.631
Alkaloid from *Buxus microphylla* and *Buxus sempervirens*. Plates (MeOH). Mp 209-211°. [α]_D²⁰ -3 (c, 0.56 in EtOH).
- N-De-Me: N-Demethylcyclomicurarine L.** *N-Demethylcyclomicurarine*
C₂₅H₄₁NO₂ 387.604
Alkaloid from the roots of *Buxus sempervirens*. [α]_D²⁰ -23 (c, 0.26 in CHCl₃). λ_{max} 203 (MeOH).
- 1,2-Didehydro: 16-Hydroxycyclobuxoviridine L.** *N^b-Methylcyclobuxoviridine*
C₂₆H₄₁NO₂ 399.615
Alkaloid from the leaves of *Buxus hyrcana*. Amorph. solid. [α]_D²⁴ -53 (c, 0.06 in CHCl₃). λ_{max} 222 (log ε 4.2); 268 (log ε 3.6) (MeOH).
- 1,2-Didehydro, N-de-Me: Cyclobuxoviridine**
[102092-24-0]
C₂₅H₃₉NO₂ 385.589
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. powder. [α]_D²⁶ -41.2 (CHCl₃).
- Nakano, T. et al., *J.C.S.(C)*, 1966, 1412-1421 (*isol, struct, ir, uv, pmr, ord*)
Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1986, 49, 106 (*Cyclobuxoviridine*)
Ata, A. et al., *Z. Naturforsch., C*, 2002, 57, 21-28 (*N-de-Me*)

Choudhary, M.I. *et al.*, *Chem. Biodiversity*, 2006, **3**, 1039-1052 (*N^b-Methylcyclobuxoviricine*)

Cyclomyrsinol**C-885**

$C_{20}H_{30}O_8$ 398.452

5,14-Bis(3-pyridinecarbonyl), 8-(2-methylpropanoyl), 3,10,15-tri-Ac: [213920-23-1]

$C_{42}H_{48}N_2O_{14}$ 804.846

Constit. of *Euphorbia seguieriana*.

Cryst. Mp 255°. λ_{max} 230 (log ϵ 1.5); 255 (log ϵ 1.13); 260 (log ϵ 1.3) (EtOH).

8,14-Bis(3-pyridinecarbonyl), 3-propionyl, 5,10,15-tri-Ac: [171864-16-7]

$C_{41}H_{46}N_2O_{14}$ 790.819


Constit. of *Euphorbia seguieriana*. $[\alpha]_D^{20}$ +12.5 (c, 0.2 in $CHCl_3$).

Jeske, F. *et al.*, *Phytochemistry*, 1995, **40**, 1743 (*isol, pmr, cmr*)

Öksüz, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1198-1201 (*isol, pmr, cmr*)

Cycloneosamandione**C-886**

[3148-28-5]



$C_{19}H_{29}NO_2$ 303.444

Alkaloid from skin glands of *Salamandra maculosa*. Mp 118-119°. $[\alpha]_D^{21}$ -207.6 (Me_2CO).


Hydrochloride: Mp 272-273°.

Picrate: Mp 225-226°.

Schöpf, C. *et al.*, *Annalen*, 1960, **633**, 127-156 (*isol, ir, struct*)

Habermehl, G. *et al.*, *Angew. Chem., Int. Ed.*, 1963, **2**, 157 (*struct*)

Oka, K. *et al.*, *J.A.C.S.*, 1977, **99**, 3859-3860 (*synth, bibl*)

Cycloroidin**C-887**

Absolute Configuration

$C_{11}H_{11}Br_2N_5O$ 389.049

Alkaloid from the sponge *Agelas oroides*. Pale yellow solid. $[\alpha]_D$ -12 (c, 0.02 in MeOH).


Fattorusso, E. *et al.*, *Tet. Lett.*, 2000, **41**, 9917-9922 (*isol, pmr, cmr*)

Papeo, G. *et al.*, *Tet. Lett.*, 2005, **46**, 8635-8638 (*synth*)

Patel, J. *et al.*, *Tet. Lett.*, 2006, **47**, 5561-5563 (*synth*)

Cyclopapilosine I**C-888**

3,20-Diamino-4,14-dimethyl-9,19-cyclo-pregnane-4-methanol



$C_{24}H_{42}N_2O$ 374.609

The suffix letter in *Buxus* alkaloids systematically defines the *N*-subn. pattern. The parent compd., Cyclopapilosine I, is unknown.

N^3, N^{20} -Di-Me: **Cyclopapilosine D**

[51059-61-1]

$C_{26}H_{46}N_2O$ 402.662

Alkaloid from *Buxus papillosa* (Buxaceae). Cryst. (Me_2CO). Mp 233-235°. $[\alpha]_D^{25}$ +54 (c, 0.86 in $CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me, N^3 -(4-hydroxy-3,5-dimethoxybenzoyl): **Buxakashmiramine**

[398477-70-8]

$C_{36}H_{56}N_2O_5$ 596.849

Alkaloid from *Buxus papillosa*.

Amorph. powder. $[\alpha]_D^{25}$ -3 (c, 0.62 in $CHCl_3$). λ_{max} 211 (log ϵ 4.46); 260 (log ϵ 3.8) (MeOH).


Shamma, M. *et al.*, *Phytochemistry*, 1973, **12**, 2051-2054 (*isol*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*struct*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 2001, **58**, 963-968 (*Buxakashmiramine*)

Cyclopenin**C-889**

4-Methyl-3'-phenylspiro[3H-1,4-benzodiazepine-3,2'-oxirane]-2,5(1H,4H)-dione, 9CI
[19553-26-5]



$C_{17}H_{14}N_2O_3$ 294.309

(-)-form [20007-87-8]

Metab. of *Penicillium cyclopium* and *Penicillium corymbiferum*. Needles (EtOAc/petrol). Sol. acids, bases, EtOH, MeOH; poorly sol. H_2O . Mp 183-184°. $[\alpha]_D^{20}$ -291 (c, 1.2 in MeOH). λ_{max} 211 (ϵ 37200); 290 (ϵ 2060) (MeOH) (Berdy).

3'-Hydroxy: **Cyclopenol**

[20007-85-6]

$C_{17}H_{14}N_2O_4$ 310.309

From *Penicillium cyclopium* and *Penicillium viridicatum*. Prisms ($C_6H_6/EtOAc$). Sol. MeOH, C_6H_6 ; poorly sol. H_2O . Mp 215° dec. $[\alpha]_D^{20}$ -309 (c, 1.3 in MeOH). λ_{max} 285 (ϵ 3740) (MeOH) (Berdy).

(±)-form [19357-57-4]

Mp 194-195°.

White, J.D. *et al.*, *Tetrahedron*, 1970, **26**, 233 (*synth, bibl*)

Richter, H. *et al.*, *Pharmazie*, 1974, **29**, 506 (*synth*)

Kirby, G.W. *et al.*, *J.C.S. Perkin I*, 1976, 1564 (*biosynth, bibl*)

Rhee, R.P. *et al.*, *J.O.C.*, 1977, **42**, 3650 (*synth, bibl*)

Ohmomo, S. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 1929 (*isol, pmr*)


Ishikura, M. *et al.*, *J.O.C.*, 1982, **47**, 2456 (*synth, ir, pmr*)

Kusano, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2559-2568 (*isol, uv, pmr, cmr, ms*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 849; 853

Cyclopeptine**C-890**

3,4-Dihydro-4-methyl-3-(phenylmethyl)-1H-1,4-benzodiazepine-2,5-dione, 9CI, 3-Benzyl-3,4-dihydro-4-methyl-1H-1,4-benzodiazepine-2,5-dione



(S)-form

$C_{17}H_{16}N_2O_2$ 280.326

(S)-form [50886-63-0]

Prod. by *Penicillium cyclopium*, *Penicillium expansum* and *Penicillium puberulum*. Mp 95-98°.

3,10-Didehydro: **Dehydrocyclopeptine**

[31965-37-4]

$C_{17}H_{14}N_2O_2$ 278.31

Prod. by *Penicillium cyclopium*, *Penicillium expansum* and *Penicillium puberulum*. Cryst. (MeOH). Mp 246-247.5°. Achiral.

4'-Methoxy: 3,4-Dihydro-3-(4-methoxybenzyl)-4-methyl-1H-1,4-benzodiazepine-2,5-dione. **4'-Methoxycyclopeptine**

$C_{18}H_{18}N_2O_3$ 310.352

Prod. by *Penicillium cf. simplicissimum*. Amorph. solid. Mp 72-75°. $[\alpha]_D^{20}$ -90.2 (c, 0.1 in EtOH). Exists as 2 conformational isomers. λ_{max} 219 (ϵ 55000); 283 (ϵ 5900) (EtOH).

3,10-Epoxyde: see Cyclopenin, C-889

(±)-form [65027-11-4]

Prisms (Me_2CO /hexane). Mp 100.5-103°. [19553-22-1, 65002-54-2]

Framm, J. *et al.*, *Eur. J. Biochem.*, 1973, **37**, 78 (*isol*)


El-Azouny, A. *et al.*, *Pharmazie*, 1977, **32**, 318 (*synth*)

Ishikura, M. *et al.*, *J.O.C.*, 1982, **47**, 2456 (*synth*)

- Gerlach, M. *et al.*, *Phytochemistry*, 1985, **24**, 1935 (*synth*)
 Solov'eva, T.F. *et al.*, *Mikrobiologiya*, 1992, **61**, 395 (*isol*)
 Kusano, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2559-2568 (*4'-Methoxycyclopeptide*)

Cyclo(phenylalanylprolyl) C-891

Hexahydro-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. 3-Benzylhexahydropyrrolo[1,2-a]pyrazine-1,4-dione, 8CI. Phenylalanylproline anhydride [14705-60-3]



(3R,8aR)-form

C₁₄H₁₆N₂O₂ 244.293**(3R,8aR)-form***D-D-form*

[5654-85-3]

Prod. by the marine bacterial strains CF-20 and C-148 obt. from molluscs.

Amorph. solid. $[\alpha]_D^{25} +88.7$ (c, 0.22 in EtOH).

(3R,8aS)-form [26488-24-4]

Prod. by *Alternaria alternata* and *Streptomyces rochei*. Phytotoxic. Mp 150-153°. $[\alpha]_D^{25} -94.2$ (c, 0.2 in H₂O).

(3S,8aR)-form [32021-26-4]

Isol. from the sponge *Calyx cf. podatypa*. Prod. by *Aspergillus flavipes*. Cryst. (EtOAc). Mp 150-151°. $[\alpha]_D^{20} +99$ (c, 0.23 in H₂O).

(3S,8aS)-form*L-L-form*

[3705-26-8]

Prod. by *Alternaria alternata*, *Candida albicans*, *Penicillium bilaii* and *Rosellinia necatrix*. Isol. from the sponge *Stelletta clavosa*. Shows broad spectrum antibacterial and gastrointestinal cell maturation enhancing activity. Mp 133°. $[\alpha]_D -89$ (c, 0.3 in H₂O).

[43041-16-3, 43041-30-1, 52645-32-6]

Ott, H. *et al.*, *Tetrahedron*, 1963, **19**, 1675 (*synth*)

Lingappa, B.T. *et al.*, *Science (Washington, D.C.)*, 1969, **163**, 192 (*isol*)

Poisel, H. *et al.*, *Chem. Ber.*, 1973, **106**, 3408 (*synth*)

Vicar, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 1940 (*synth, pmr*)

Ramani, R. *et al.*, *Acta Cryst. B*, 1976, **32**, 1051 (*cryst struct*)

Young, P.E. *et al.*, *J.A.C.S.*, 1976, **98**, 5365 (*pmr, cmr*)

Suzuki, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 233 (*synth*)

Mazza, F. *et al.*, *Acta Cryst. C*, 1984, **40**, 1974 (*cryst struct*)

Sleeckx, J.J.M. *et al.*, *Bull. Soc. Chim. Belg.*, 1985, **94**, 187 (*pmr, conformm*)

Sterle, A.C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 8008 (*isol*)

Adamczeski, M. *et al.*, *J. Nat. Prod.*, 1995, **38**, 201-208 (*isol, pmr, cmr*)

Bull, S.D. *et al.*, *J.C.S. Perkin 1*, 1998, 2313-2320 (*config, biosynth*)

Graz, M. *et al.*, *Pharmazie*, 1999, **54**, 772 (*activity*)

Fdhila, F. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1299-1301 (*isol, pmr, cmr*)

Wegerski, C.J. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 5631-5637 (*isol*)


Stark, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 7222-7231 (*L,L-form, isol, ms*)

Tullberg, M. *et al.*, *Tetrahedron*, 2006, **62**, 7484-7491 (*synth, pmr, cmr*)

Li, Z. *et al.*, *Biochem. Syst. Ecol.*, 2008, **36**, 230-234 (*isol, pmr, cmr*)

Cyclopiamide**C-892**

7,8-Dihydro-2,7,7-trimethyl-1H-isoindolo[4,5,6-cd]indole-1,9(2H)dione, 9CI [126382-02-3]


C₁₆H₁₄N₂O₂ 266.299

Metab. of *Penicillium cyclopium*. Mycotoxin. Pale yellow solid (CHCl₃/hexane). Sol. MeOH, Et₂O, CHCl₃; poorly sol. H₂O. Mp 157-158°. λ_{max} 225 (ε 8750); 240 (ε 7400); 268 (ε 10200); 341 (ε 828); 358 (ε 1090); 406 (ε 700) (MeOH) (Berdy).

Holzappel, C.W. *et al.*, *Phytochemistry*, 1990, **29**, 639 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Cyclopiamine A**C-893**

[72361-74-1]



Relative configuration

C₂₆H₃₃N₃O₅ 467.564

Metab. of *Penicillium cyclopium* and *Penicillium urticae*. Noncryst.; cryst. (as hydrobromide). Mp 234-236° (hydrobromide). λ_{max} 230 (ε 15850); 261 (ε 10300); 344 (ε 5630) (MeOH) (Berdy).

► Toxic.

22-Epimer: Cyclopiamine B

[72401-58-2]

C₂₆H₃₃N₃O₅ 467.564


Metab. from *Penicillium cyclopium*, *Penicillium urticae* and *Aspergillus caespitosus*. Mp 245-246°. λ_{max} 232 (ε 16050); 260 (ε 10650); 345 (ε 5740) (MeOH) (Berdy).

Bond, R.F. *et al.*, *J.C.S. Perkin 1*, 1979, 1751 (*isol, uv, cd, ir, pmr, ms, struct*)

 α -Cyclopiazonic acid**C-894**

[18172-33-3]

[83136-88-3]



Relative Configuration

C₂₀H₂₀N₂O₃ 336.39

Enolised β -triketone. CAS numbering shown. Toxic alkaloid from *Penicillium cyclopium*, *Penicillium camemberti*, *Penicillium duclauxii* MF2831, *Aspergillus versicolor*, *Aspergillus flavus*, *Aspergillus oryzae* ATCC20235 and *Aspergillus tamaraii* IAM1465. Antioxidant. Mycotoxin. Inhibits sarcoplasmic reticulum Ca²⁺-ATPase and ATP-dependent Ca²⁺ transport. Pharmacol. tool for studying intracellular Ca²⁺ homeostasis. Exerts negative inotropic effects on adult myocardium. Mp 245-246°. $[\alpha]_D^{18} -74$ (c, 1 in CHCl₃). λ_{max} 225 (ε 39800); 253 (ε 16600); 275 (sh) (ε 19100); 284 (ε 20400); 292 (sh) (ε 17400) (MeOH) (Derep). λ_{max} 224 (ε 32360); 282 (ε 17780); 290 (ε 14125) (MeOH) (Berdy). λ_{max} 223 (ε 39400); 281 (ε 19300) (EtOH) (Berdy). λ_{max} 222 (ε 40800); 252 (ε 16300); 281 (ε 19300) (EtOH/NaOH) (Berdy).

► Exp. reprod. and teratogenic effects.

Tremorgenic toxin. Adverse effects incl. muscle incoordination, hypokinesia, convulsions; LD₅₀ (mus, ipr) 2.3 mg/kg. LD₅₀ (rat, orl) 36 mg/kg. UY8587000

Hydrazone: Mp 189-190°.*1-Hydroxy, 2-Me: Speridine A*

[566943-74-6]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from a marine-derived *Aspergillus tamaraii*. Pale yellow solid. $[\alpha]_D^{18} -79$ (c, 1 in CHCl₃). Exists in 1-oxo-2NH-form. λ_{max} 210 (ε 15000); 253 (ε 7700); 280 (ε 6400) (MeOH).

Holzappel, C.W. *et al.*, *Tetrahedron*, 1968, **24**, 2101-2119 (*isol, uv, ir, ms, pmr, cd, struct*)

Holzappel, C.W. *et al.*, *Phytochemistry*, 1971, **10**, 351-358 (*biosynth*)

Purchase, I.F.H. *et al.*, *Toxicol. Appl. Pharmacol.*, 1971, **18**, 114-123 (*tox*)

Steyn, P.S. *et al.*, *Chem. Comm.*, 1975, 465-466 (*biosynth*)

De Jesus, A.E. *et al.*, *J.C.S. Perkin 1*, 1981, 3292-3294 (*biosynth*)

Chalmers, A.A. *et al.*, *Chem. Comm.*, 1982, 1367-1368 (*biosynth*)

Pohland, A.E. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 2220-2284 (*uv, ir, pmr, ms, cd*)

Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2153-2156 (*synth*)


Kozikowski, A.P. *et al.*, *J.A.C.S.*, 1984, **106**, 6873-6874 (*synth*)

Nishie, K. *et al.*, *Food Chem. Toxicol.*, 1985, **23**, 831-839; 1069-1076 (*tox, metab*)

Muratake, H. *et al.*, *Heterocycles*, 1985, **23**, 1111 (*synth*)


- Norred, W.P. *et al.*, *J. Agric. Food Chem.*, 1988, **36**, 113-116
- Seidler, N.W. *et al.*, *J. Biol. Chem.*, 1989, **264**, 17816-17823 (*pharmacol*)
- Agata, N. *et al.*, *Br. J. Pharmacol.*, 1993, **108**, 571-572 (*pharmacol*)
- Bradburn, N. *et al.*, *Phytochemistry*, 1994, **35**, 817 (*activity*)
- Plenge-Tellechea, F. *et al.*, *J. Biol. Chem.*, 1997, **272**, 2794-2800 (*pharmacol*)
- Schwinger, R.H.G. *et al.*, *J. Pharmacol. Exp. Ther.*, 1997, **283**, 286-292 (*pharmacol*)
- Tsuda, M. *et al.*, *Tetrahedron*, 2003, **59**, 3227-3230 (*Speradine A*)
- Haskins, C.M. *et al.*, *Chem. Comm.*, 2005, 3162-3164 (*synth*)
- Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 497
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CQD000

Cyclopiazonic acid imine **C-895**
[31008-94-3]




- $C_{20}H_{21}N_3O_2$ 335.405
- Isol. from *Penicillium cyclopium* and *Aspergillus versicolor*. Cryst. (MeOH). Mp 277-278°. $[\alpha]_D^{20}$ -11.1. Relatively nontoxic compared with cyclopiazonic acid. λ_{max} 224 (ε 33000); 293 (ε 22380) (MeOH) (Berdy).
- Holzappel, C.W. *et al.*, *Tetrahedron*, 1970, **26**, 5239 (*isol, uv, ir, ms*)
- Ohmomo, S. *et al.*, *Nippon Nogei Kagaku Kaishi*, 1973, **47**, 57 (*isol, pmr*)
- Riley, R.T. *et al.*, *Toxicol. Appl. Pharmacol.*, 1992, **114**, 261-267 (*pharmacol*)
- Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 501

Cyclopiperstachine **C-896**
2-(1,3-Benzodioxol-5-yl)-1,2,4a,5,6,7,8,8a-octahydro-N-(2-methylpropyl)-1-naphthalenecarboxamide, 9CI




- $C_{22}H_{29}NO_3$ 355.476
- (±)-*form* [58514-07-1]
- Alkaloid from *Piper trichostachyon* (Piperaceae). Needles (CH₂Cl₂/hexane). Mp 220°.
- Joshi, B.S. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 2295 (*isol, uv, ir, pmr, ms, struct, synth*)

Cycloprodigosin **C-897**
4,5,6,7-Tetrahydro-3-[[3-methoxy-5-(1H-pyrrol-2-yl)-2H-pyrrol-2-ylidene]methyl]-1,4-dimethyl-2H-isoindole, 9CI [86797-91-3]



- $C_{20}H_{23}N_3O$ 321.421
- Pyrrole antibiotic. Revised struct. Minor red pigment from cultures of the aerobic marine bacterium *Alteromonas rubra*. Also isolated from *Pseudoalteromonas denitrificans* and *Beneckea gazogenes*. Shows antibacterial, antifungal, immunosuppressant, antiarteriosclerotic and antiosteoporotic activities. λ_{max} 216 (ε 8660); 275 (sh) (ε 6440); 296 (ε 12200); 371 (ε 7750); 382 (sh) (ε 7240); 510 (sh) (ε 54400); 541 (ε 132000) (EtOH, pH 2.9) (Derep). λ_{max} 226 (ε 10800); 289 (ε 9820); 337 (ε 7590); 471 (ε 41600); 539 (ε 17200) (EtOH, pH 7.4) (Derep). λ_{max} 257 (ε 7950); 281 (ε 9520); 336 (ε 7860); 469 (ε 42400) (EtOH, pH 11) (Derep). λ_{max} 544 (ε 110000) (CHCl₃/HCl) (Berdy).
- Laatsch, H. *et al.*, *Tet. Lett.*, 1983, **24**, 2701 (*isol, uv, pmr, ms, struct*)
- Gerber, N.N. *et al.*, *Tet. Lett.*, 1983, **24**, 2797 (*cmr, pmr*)
- Wassermann, H.H. *et al.*, *Tet. Lett.*, 1984, 1387 (*synth*)
- Japan. Pat., 1998, 98 80 293; CA, **128**, 294012d (*isol, activity*)

Cyclo(prolylprolyl) **C-898**
Octahydro-5H,10H-dipyrrolo[1,2-a:1',2'-d]pyrazine-5,10-dione, 9CI. Prolylproline anhydride. Octahydroprocoll [6708-06-1]




- $C_{10}H_{14}N_2O_2$ 194.233
- (5aR,10aR)-*form*
D-D-form
[53990-71-9]
Mp 141-143°. $[\alpha]_D^{30}$ +147.5 (c, 1 in H₂O).
- (5aR,10aS)-*form*
L-D-form
[53990-72-0]
Mp 193-195°. Opt. inactive (*meso*-).
- (5aS,10aS)-*form*
L-L-form
[19943-27-2]
Constit. of roasted cocoa nibs and roasted malt; also from the tubers of *Pinellia pedatisecta*. Prod. by *Aspergillus fumigatus* and *Pseudoalteromonas haloplanktis*. Mp 143-144°. $[\alpha]_D^{30}$ -149.5 (c, 1 in

- H₂O).
- (5aRS,10aRS)-*form*
(±)-*form*
Cryst. (EtOH). Mp 149-150°.
[19943-26-1]

- Rothe, M. *et al.*, *Angew. Chem., Int. Ed.*, 1972, **11**, 293 (*synth*)
- Eguchi, C. *et al.*, *Bull. Chem. Soc. Jpn.*, 1974, **47**, 2277-2282 (*synth*)
- Kralj, B. *et al.*, *Biomed. Mass Spectrom.*, 1975, **2**, 215-218 (*synth, ms*)
- Benedetti, E. *et al.*, *Cryst. Struct. Commun.*, 1975, **4**, 641-645 (*cryst struct*)
- Kricheldorf, H.R. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 52-58 (*synth, cmr, N-15 nmr*)
- Ueda, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1983, **56**, 568-572 (*synth*)
- Qin, W. *et al.*, *Zhongcaoyao*, 1984, **15**, 490-492; CA, **102**, 109774f (*isol*)
- Valentine, B. *et al.*, *Int. J. Pept. Protein Res.*, 1985, **25**, 56-68 (*O-17 nmr*)
- Stark, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 7222-7231 (*L,L-form, isol, ms*)
- Furtado, N.A.J.C. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 1448-1453 (*isol, pmr, cmr*)
- Mitova, M. *et al.*, *Mar. Biotechnol.*, 2005, **7**, 523-531 (*L,L-form, isol*)

Cyclo(prolyltryptophyl) **C-899**
Hexahydro-3-(1H-indol-3-ylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Cyclo(tryptophanylprolyl). Cyclo(tryptophylprolyl) [67889-75-2]




- $C_{16}H_{17}N_3O_2$ 283.329
- (3R,8aS)-*form*
D-L-form
[41222-71-3]
Cryst. (Me₂CO). Mp 204-206°. $[\alpha]_D^{22}$ -101 (c, 1.78 in AcOH).
- (3S,8aR)-*form*
L-D-form
[73136-47-7]
Cryst. (Me₂CO/petrol). Mp 191-193°. $[\alpha]_D^{20}$ +120 (c, 1.3 in AcOH).
- (3S,8aS)-*form*
L-L-form. Brevianamide F
[38136-70-8]
Prod. by *Penicillium brevi-compactum*, *Penicillium piscarium* and *Alteromonas distincta* He169. Also prod. by a *Vibrio* sp. isol. from the sponge *Hyrtios altum*. Shows broad spectrum antibacterial and gastrointestinal cell maturation enhancing activity. Mp 173-175°. $[\alpha]_D^{24}$ -99 (c, 1.2 in AcOH). λ_{max} 220 (ε 11000); 276 (ε 1900); 281 (ε 2000); 290 (ε 1700) (MeOH) (Berdy).
- 8,8a-Didehydro: 12,13-Dehydroprolyl-tryptophyldiketopiperazine. Cyclo(dehydroprolyltryptophyl)
[290348-28-6]
 $C_{16}H_{15}N_3O_2$ 281.313

Prod. by *Penicillium piscarium*.

- Birch, A.J. *et al.*, *Tetrahedron*, 1972, **28**, 2999 (isol, uv, ir, ms, pmr, synth)
 Steyn, P.S. *et al.*, *Tetrahedron*, 1973, **29**, 107
 Sammes, P.G. *et al.*, *J.C.S. Perkin 1*, 1979, 3048 (synth, bibl)
 Kobayashi, M. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 2449-2451 (*Vibrio*, isol)
 Grant, G.D. *et al.*, *J. Chem. Crystallogr.*, 1999, **29**, 435-447 (cryst struct)
 Graz, M. *et al.*, *Pharmazie*, 1999, **54**, 772 (activity)
 Kozlovskii, A.G. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 2000, **36**, 317-321 (isol)
 Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (rev, synth, biosynth)
 Shabaan, M. *et al.*, *Dissertation*, Univ. of Gottingen, 2004, (*Alteromonas*, isol)

Cyclo(prolyltyrosyl) C-900

Hexahydro-3-(4-hydroxybenzyl)pyrrolo[1,2-a]pyrazine-1,4-dione
 [74345-23-6]



C₁₄H₁₆N₂O₃ 260.292

(3S,8aR)-form

L,D-form

[61117-56-4]

Isol. from two strains of microorganisms isol. from the marine sponge *Ircinia variabilis* and the fungus *Aspergillus flavipes*. Cryst. (Et₂O/Me₂CO). Mp 218-220°.

(3S,8aS)-form

L,L-form. *Maculosin 1*. ML 1532II. A 19C. Antibiotic A 19C. Antibiotic ML 1532II

[4549-02-4]

Prod. by *Alternaria alternata*, *Aspergillus flavipes*, *Streptomyces gelaticus*, *Streptomyces* sp. ML1532, *Fusarium nivale*, a fungal growth on *Zinnia elegans* and *Bacillus licheniformis*. Isol. from the sponges *Jaspis digonoxea* and *Tedania anhelans*, a marine-derived *Pseudomonas aeruginosa*, marine *Penicillium* sp., marine bacteria *Vibrio parahaemolyticus* and *Pseudoalteromonas haloplanktis* and pathogenic fungus *Pestalotia palmarum*. Present in coml. peptone. Cytotoxic. Inhibits SV-40 transformed cells and germination. Plant growth regulator. Possesses pesticidal props. Plates. Mp 152-154° (140°). [α]_D -148 (c, 1 in H₂O).

(3ξ,8aξ)-form

3'-Nitro: Hexahydro-3-[(4-hydroxy-3-nitrophenyl)methyl]pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Cyclo(prolyl-3-nitrotyrosyl). *Pyricularamide*
 [134876-76-9]
 [125034-08-4]


C₁₄H₁₅N₃O₅ 305.29

Isol. from *Pyricularia oryzae* and the marine-derived *Flavobacterium* sp. T436. Plant growth inhibitor. Bright yellow needles (Me₂CO). Mp 183-185° dec. [α]_D -84.4 (c, 0.66 in MeOH).

- Tamura, S. *et al.*, *Agric. Biol. Chem.*, 1964, **28**, 650-652 (*3S,8aS*-form, isol)
 Tatsuno, T. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 1498-1500 (*3S,8aS*-form, isol)
 Madison, V. *et al.*, *J.A.C.S.*, 1976, **98**, 5365-5371 (synth, cd, pmr, cmr)
 Kamikawa, T. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 691-692 (*3S,8aS*-form, isol)
 Munekata, M. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 2613-2618 (*3S,8aS*-form, isol)
 Sasaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4435-4443 (*3S,8aS*-form, synth)
 Stierle, A.C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 8008-8011 (*Maculosin 1*)
 Boldyrev, A.A. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1989, **94**, 237 (*Pyricularamide*, pharmacol)
 Sviridov, S.I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 691-696 (*Pyricularamide*)
 Bavykina, N.I. *et al.*, *Zh. Obshch. Khim.*, 1990, **60**, 170-175; *J. Gen. Chem. USSR (Engl. Transl.)*, 1990, **60**, 148-152 (*Pyricularamide*, synth)
 Milne, P.J. *et al.*, *J. Crystallogr. Spectrosc. Res.*, 1992, **22**, 643-649 (cryst struct, conformm)
 Barrow, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 471-476 (*Aspergillus flavipes* constit)
 Rudi, A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 829-836 (isol, pmr, cmr)
 Jayatilake, G.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 293-296 (isol, pmr, cmr)
 De Rosa, S. *et al.*, *Biomol. Eng.*, 2003, **20**, 311-316 (*Ircinia variabilis* constit)
 Schuhmann, I. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*Pyricularamide*, isol, marine)
 Mitova, M. *et al.*, *Mar. Biotechnol.*, 2005, **7**, 523-531 (*L,L*-form, isol)

Cyclo(prolylvalyl) C-901

Hexahydro-3-(1-methylethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, 9CI. Hexahydro-3-isopropylpyrrolo[1,2-a]pyrazine-1,4-dione. Prolylvaline anhydride. *Maculosine 5*. Antibiotic K 73. K 73
 [5654-87-5]



C₁₀H₁₆N₂O₂ 196.249

Diketopiperazine antibiotic.

(3R,8aR)-form

D,D-form

Prod. by the marine bacterial strains CF-20 and C-148 obt. from molluscs. Active against *Vibrio anguillarum*. [α]_D +120.1 (c, 0.1 in EtOH).

(3R,8aS)-form

D,L-form

[27483-18-7]

Cryst. (EtOAc/hexane). Mp 154-156°.
 [α]_D²⁰ -96.4 (c, 0.25 in H₂O).

(3S,8aR)-form

L,D-form

[156617-55-9]

Prod. by *Aspergillus* sp. F70609 and *Aspergillus flavipes*. Specific β-glucosidase inhibitor. Cryst. (EtOAc/hexane). Mp 147-149°. [α]_D²⁰ +88 (c, 1 in H₂O).

(3S,8aS)-form

L,L-form

[2854-40-2]

Prod. by *Aspergillus*, *Streptomyces*, *Oospora* spp. and *Pinnellia pedatisecta*. Alkaloid from the Caribbean sponges *Calyx* cf. *podatypa* and *Tedania ignis*. Also from the alga *Scenedesmus* sp., the fungi *Rhizoctonia solani* and *Rosellinia necatrix*, and root of *Psammosilene tunicoides*. Shows antibiotic and phytotoxic props. Needles (butanol). Mp 189.5° (169-172°). [α]_D²⁰ -157 (c, 1 in CHCl₃). [α]_D²⁰ -180.5 (c, 1 in EtOH).

Luedemann, G. *et al.*, *J.O.C.*, 1961, **26**, 4128-4130 (isol)

Siemion, I.Z. *et al.*, *Org. Magn. Reson.*, 1971, **3**, 545-550 (synth, nmr)

Ogura, H. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2474-2477 (isol)

Munekata, M. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 2613-2618 (isol)

Schmidtz, F.J. *et al.*, *J.O.C.*, 1983, **48**, 3941-3945 (isol, pmr, ms, synth, bibl)

Giralt, E. *et al.*, *Synthesis*, 1985, 181-184 (synth)

Stierle, A.C. *et al.*, *Experientia*, 1988, **44**, 1021 (isol)

Barrow, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 471-476 (*L,D*-form, isol)

Adamczeski, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 201-208 (isol, pmr, cmr)

Jayatilake, G.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 293-296 (isol, cmr)

Bull, S.D. *et al.*, *J.C.S. Perkin 1*, 1998, 2313-2320 (configs, biosynth)

Kwon, O.S. *et al.*, *J. Antibiot.*, 2001, **54**, 179-181 (*L,D*-form, isol, pmr, cmr)


Fdhila, F. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1299-1301 (isol, pmr, cmr)

Ding, Z. *et al.*, *Zhongguo Zhongyao Zazhi*, 2003, **28**, 337-339; *CA*, **142**, 215155 (isol, pmr, cmr)

Pedras, M.S.C. *et al.*, *Z. Naturforsch., C*, 2005, **60**, 717-722 (isol, synth, pmr, cmr)

Cycloprotobuxine I C-902

4,4,14-Trimethyl-9,19-cyclopregnane-3,20-diamine, 9CI



R = H

C₂₄H₄₂N₂ 358.609

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Cycloprotobuxine I, is unknown. Log P 5.88 (uncertain value) (calc).

N^3, N^{20} -Di-Me: **Cycloprotobuxine D**
[2255-38-1]
 $C_{26}H_{46}N_2$ 386.663
Alkaloid from *Buxus balearica*,
Buxus harlandi and *Buxus sempervirens*
(Buxaceae). Cryst. (EtOH). Mp 140°.
[α]_D +112 (c, 1 in $CHCl_3$).

N^3, N^{20} -Di-Me, N^{20} -Ac: **N-Acetylcycloprotobuxine D**
[16974-77-9]
 $C_{28}H_{48}N_2O$ 428.7
Alkaloid from *Buxus sempervirens*
(Buxaceae). Mp 221-224°. [α]_D +53
($CHCl_3$).

N^{20}, N^{20} -Di-Me: **Cycloprotobuxine F**
[36151-05-0]
 $C_{26}H_{46}N_2$ 386.663
Alkaloid from *Buxus madagascariensis*
(Buxaceae). Has been used as a synthetic material for holothurinogenins. Needles (Me_2CO). Mp 163°. [α]_D +42
($CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me: **Cycloprotobuxine C**
[1936-70-5]
 $C_{27}H_{48}N_2$ 400.69
Alkaloid from *Buxus sempervirens*,
Buxus harlandi, *Buxus microphylla* and
other *Buxus* spp. (Buxaceae). Mp 212°
(200-202°). [α]_D +76 (c, 1.09 in
 $CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me, N^3 -Ac: **N-Acetylcycloprotobuxine C**
[16975-12-5]
 $C_{29}H_{50}N_2O$ 442.727
Alkaloid from *Buxus sempervirens*
(Buxaceae). Mp 222-235°.

N^3, N^{20}, N^{20} -Tri-Me, N^3 -benzoyl: **N-Benzoylcycloprotobuxine C**
[16974-72-4]
 $C_{34}H_{52}N_2O$ 504.798
Alkaloid from *Buxus sempervirens*.
Cryst. (C_6H_6). Mp 231-233°. [α]_D²⁶ +32
(c, 1.0 in $CHCl_3$).

N^3, N^3, N^{20}, N^{20} -Tetra-Me: **Cycloprotobuxine A**
[2278-38-8]
 $C_{28}H_{50}N_2$ 414.716
Alkaloid from *Buxus sempervirens*,
Buxus microphylla, *Buxus malayana*
and *Buxus balearica* (Buxaceae). Plates
(Me_2CO). Mp 207-208°. [α]_D +75 (c, 1
in $CHCl_3$) (+31).

►GZ0585000

6,7-Didehydro, N^3, N^{20}, N^{20} -tri-Me: **Cyclobuxupaline C**
[51059-60-0]
 $C_{27}H_{46}N_2$ 398.674
Alkaloid from *Buxus papilosa* (Buxaceae). Cryst. (Me_2CO). Mp 111-113°.
[α]_D²⁵ -37 (c, 1.0 in $CHCl_3$).

6,7-Didehydro, N^3, N^{20}, N^{20} -tri-Me, N^3 -benzoyl: **Buxeridine C**. **Buxeridine**
[17337-35-8]
 $C_{34}H_{50}N_2O$ 502.782
Alkaloid from *Buxus sempervirens*
leaves (Buxaceae). Prisms (Me_2CO).
Mp 208-210°. [α]_D +14 ($CHCl_3$).
Struct. not certain.

20-Epimer, N^3, N^{20}, N^{20} -tri-Me, N^3 -Ac: **Buxaline C**

[16975-11-4]
 $C_{29}H_{50}N_2O$ 442.727
Alkaloid from *Buxus sempervirens*
(Buxaceae). Mp 230-232°. [α]_D +29.41
($CHCl_3$). Tentative struct.

3,20-Diepimer, N^3, N^{20}, N^{20} -tri-Me: **1-Cycloprotobuxine C**
[56687-61-7]
 $C_{27}H_{48}N_2$ 400.69
Alkaloid from *Buxus sempervirens*. Mp
195-197°. [α]_D -62 (c, 0.25 in $CHCl_3$).
Structure proof tenuous; (20*R*)-config.
unprecedented.

Calame, J.P. *et al.*, *Chimia*, 1964, **18**, 185
(*struct.*, *Cycloprotobuxine D*)
Herlem-Gaulier, D. *et al.*, *Bull. Soc. Chim. Fr.*,
1965, 657-668 (*isol.*, *Cycloprotobuxine A*,
Cycloprotobuxine C)
Nakano, T. *et al.*, *J.C.S.*, 1965, 6688-6694
(*Cycloprotobuxine C*, *isol.*, *pmr.*, *struct.*)
Nakano, T. *et al.*, *J.C.S. (C)*, 1966, 1412-1421
(*isol.*, *Cycloprotobuxine C*)
Khuong-Huu-Laine, F. *et al.*, *Tetrahedron*,
1966, **22**, 3321-3327 (*isol.*, *Cycloprotobuxine A*,
Cycloprotobuxine C, *Cycloprotobuxine D*)
Döpke, W. *et al.*, *Naturwissenschaften*, 1967,
54, 200 (*Buxeridine C*)
Kupchan, S.M. *et al.*, *Tetrahedron*, 1967, **23**,
4563-4586 (*isol.*, *pmr.*, *N-Acetylcycloprotobuxine D*)
Khuong-Huu, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, 558-560
(*Cycloprotobuxine F*)
Shamma, M. *et al.*, *Phytochemistry*, 1973, **12**,
2051-2054 (*Cyclobuxupaline C*)
Khodzhaev, B.U. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 281-282; 1976, **12**,
500-501 (*l-Cycloprotobuxine C*)
Khodzhaev, B.U. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 795-797 (*Buxaline C*)
Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794
(*Cycloprotobuxine F*, *cmr*)
Singh, C. *et al.*, *Tetrahedron*, 1977, **33**, 1053-1055
(*Cycloprotobuxine F*, *Cycloprotobuxine A*, *synth*)
Khodzhaev, B.U. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2003, **39**, 52-53 (*N-Benzoylcycloprotobuxine C*)

Cycloprotobuxoline I C-903

3,20-Diamino-9,19-cyclopregnan-2-ol
As Cycloprotobuxine I, C-902 with
R = OH

$C_{24}H_{42}N_2O$ 374.609

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Cycloprotobuxoline I, is unknown.


N^3, N^{20} -Di-Me, N^3 -benzoyl: **N-Benzoylcycloprotobuxoline D**
[16974-80-4]
 $C_{33}H_{50}N_2O_2$ 506.77
Alkaloid from *Buxus sempervirens*
(Buxaceae). Mp 236-238° dec. [α]_D²⁶ +42 (c, 1.05 in $CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me, N^3 -benzoyl: **N-Benzoylcycloprotobuxoline C**
[16974-74-6]
 $C_{34}H_{52}N_2O_2$ 520.797
Alkaloid from *Buxus sempervirens*
(Buxaceae). Mp 252-255° dec. [α]_D²⁷ +43 (c, 0.85 in $CHCl_3$).

Kupchan, S.M. *et al.*, *Tetrahedron*, 1967, **23**, 4563

Cyclorocaglamide C-904

[496956-76-4]




$C_{30}H_{29}NO_9$ 547.56

Flavogline compd. Constit. of the twigs of *Aglaia oligophylla*. Amorph. solid.
[α]_D²⁰ +180.5 (c, 1.25 in EtOH).

Bringmann, G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 80-85 (*isol.*, *cd.*, *pmr.*, *cmr.*, *ms*)

Cyclorolfeine C-905

16-Hydroxy-4,4,14-trimethyl-3-methylamino-9,19-cyclopregnan-20-one
[6908-25-4]



$C_{25}H_{41}NO_2$ 387.604


Alkaloid from *Buxus rolfei* (Buxaceae). Cryst. (Me_2CO). Mp 253°. [α]_D +119 (c, 1 in $CHCl_3$).

N-Ac:
Cryst. (Me_2CO). Mp 235°. [α]_D -34 (c, 1 in $CHCl_3$).

Khuong-Huu-Lainé, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 1216-1221 (*isol.*, *pmr.*, *ms.*, *ir.*, *struct.*)

Cyclorolfoxazine C-906

[7044-75-9]




$C_{26}H_{41}NO_3$ 415.615

C-4 config. revised in 1975. Alkaloid from *Buxus rolfei* (Buxaceae). Cryst. (Me_2CO). Mp 239°. [α]_D +106 (c, 1 in $CHCl_3$).


Khuong-Huu-Lainé, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 1216-1221 (*isol.*, *ir.*, *ms.*, *pmr.*, *struct.*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (*config.*)

Cycloshermilamine D C-907
[272118-05-5]C₂₁H₁₆N₄OS 372.45

Alkaloid from the tunicate *Cystodytes violatinctus*. Amorph. yellow powder. λ_{\max} 208 (log ϵ 4.18); 215 (log ϵ 4.16); 262 (log ϵ 4.04); 299 (log ϵ 3.88); 391 (log ϵ 3.35); 480 (log ϵ 3.19) (MeOH).

Koren-Goldschlager, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 830-831 (*isol. uv, pmr*)

Cycloshermilamine D C-908
[501691-51-6]C₂₁H₂₉NO₃ 343.465


Isol. from the Australian sponge *Spongia* sp. Wine-coloured oil. $[\alpha]_D^{25}$ -18 (c, 0.12 in CHCl₃). λ_{\max} 211 (log ϵ 3.01); 313 (log ϵ 2.84) (EtOH).

Utkina, N.K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1263-1265 (*abs config*)

Utkina, N.K. *et al.*, *Tet. Lett.*, 2003, **44**, 101-102 (*isol. pmr, cmr*)

Cyclostachine A C-909

1-[2-(1,3-Benzodioxol-5-yl)-1,2,4a,5,6,7,8,8a-octahydro-1-naphthalenyl]carbonyl]pyrrolidine, 9CI

C₂₂H₂₇NO₃ 353.46(±)-**form** [57110-32-4]

Alkaloid from the stems of *Piper trichostachyon* (Piperaceae). Cryst. (Et₂O/hexane). Mp 136-138°. λ_{\max} 235 (ϵ 4460); 287 (ϵ 4170) (EtOH) (Berdy).


1,8a-Diepimer: **Cyclostachine B**

[58514-08-2]

C₂₂H₂₇NO₃ 353.46

Alkaloid from *Piper trichostachyon* (Piperaceae). Needles (CH₂Cl₂/hexane). Mp 135-136°.

Joshi, B.S. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 2295 (*isol. uv, ir, pmr, cmr, ms, struct, synth*)

Cyclostelletamines C-910

Cyclostelletamine A, x = 3, y = 3
 B, x = 3, y = 4
 C, x = 4, y = 4
 D, x = 3, y = 5
 E, x = 4, y = 5
 F, x = 5, y = 5
 G, x = 2, y = 3
 H, x = 1, y = 3
 I, x = 1, y = 4
 K, x = 1, y = 5
 L, x = 2, y = 5

Muscarinic receptor binding inhibitors.

Cyclostelletamine A [156953-85-4]C₃₄H₅₆N₂ 492.83

Alkaloid from the marine sponge *Stelletta maxima*. Sol. MeOH, butanol. λ_{\max} 266 (MeOH) (Berdy).

Cyclostelletamine B [156953-87-6]C₃₅H₅₈N₂ 506.857

From *Stelletta maxima*. Sol. MeOH, butanol. λ_{\max} 268 (MeOH) (Berdy).

Cyclostelletamine C [156953-91-2]

[181212-11-3]

C₃₆H₆₀N₂ 520.883

From *Stelletta maxima*. Amorph. solid (EtOAc/hexane) (as bis(trifluoromethanesulfonate)). Sol. MeOH, butanol. Mp 123-124° (bis(trifluoromethanesulfonate)). λ_{\max} 266 (ϵ 6320) (MeOH).

Cyclostelletamine D [156953-89-8]C₃₆H₆₀N₂ 520.883

From *Stelletta maxima*. Sol. MeOH, butanol. λ_{\max} 267 (MeOH) (Berdy).

7',8'Z-Didehydro: **Dehydrocyclostelletamine D**

[706784-72-7]

C₃₆H₅₈N₂ 518.868

Isol. from *Xestospongia* sp. Amorph. solid. CAS No. refers to trifluoroacetic acid salt.

Cyclostelletamine E [156953-93-4]C₃₇H₆₂N₂ 534.91

From *Stelletta maxima*. Sol. MeOH, butanol. λ_{\max} 268 (MeOH) (Berdy).

7',8'Z-Didehydro: **Dehydrocyclostelletamine E**

[706784-74-9]

C₃₇H₆₀N₂ 532.894

Isol. from a *Xestospongia* sp. Amorph. solid. CAS No. refers to trifluoroacetic acid salt. λ_{\max} 204 (ϵ 19000); 215 (sh) (ϵ 9400); 269 (ϵ 8600); 274 (sh) (ϵ 7200) (MeOH).

Cyclostelletamine F [156953-95-6]C₃₈H₆₄N₂ 548.937

From *Stelletta maxima*. Sol. MeOH, butanol. λ_{\max} 267 (MeOH) (Berdy).

Cyclostelletamine G [708255-98-5]C₃₃H₅₄N₂ 478.803

Isol. from a *Xestospongia* sp. and from *Pachychalina* sp. Amorph. solid. CAS No. refers to trifluoroacetic acid salt. λ_{\max} 203 (ϵ 13000); 216 (sh) (ϵ 6400); 268 (ϵ 4700); 274 (sh) (ϵ 4000) (MeOH).

Cyclostelletamine HC₃₂H₅₂N₂ 464.776

From *Pachychalina* sp.

Cyclostelletamine IC₃₃H₅₄N₂ 478.803

From *Pachychalina* sp.

Cyclostelletamine KC₃₄H₅₆N₂ 492.83

From *Pachychalina* sp.

Cyclostelletamine LC₃₅H₅₈N₂ 506.857

From *Pachychalina* sp.

Fusetani, N. *et al.*, *Tet. Lett.*, 1994, **35**, 3967-3970 (*Cyclostelletamines A-F, isol. uv, pmr, struct*)

Fusetani, N. *et al.*, *Tetrahedron*, 1996, **52**, 10849-10860 (*synth, Cyclostelletamine C, pmr, cmr, uv*)

Wanner, M.J. *et al.*, *Eur. J. Org. Chem.*, 1998, 889-895 (*synth*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (*rev*)

Baldwin, J.E. *et al.*, *Tetrahedron*, 1998, **54**, 13655-13680 (*synth*)

Oku, N. *et al.*, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 2617-2620 (*Cyclostelletamine G, Dehydrocyclostelletamines*)


De Oliveira, J.H.H.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1685-1689 (*Cyclostelletamines G-L*)

Grube, A. *et al.*, *Eur. J. Org. Chem.*, 2006, 1285-1295 (*Cyclostelletamine H, I, K, L, synth, ms*)

De Oliveira, J.H.H.L. *et al.*, *Mar. Drugs*, 2006, **4**, 1-8 (*activity*)

Cyclostrychnine

[141565-04-0]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448


Stereochem. at two chiral centres not shown in the lit. It is prob. as shown here (and in CAS), by analogy with related alkaloids. Alkaloid from *Strychnos henningii* (Loganiaceae). $[\alpha]_D$ -23 (c, 0.77 in CHCl₃). λ_{\max} 207 ; 250 ; 280 ; 288 (MeOH).

Massiot, G. *et al.*, *Phytochemistry*, 1991, **30**, 3449-3456 (*isol. uv, ir, pmr, cmr, ms, struct*)

Hayaoni, S. *et al.*, *J.O.C.*, 2006, **71**, 5574-5585 (*synth, oxide, cryst struct*)

Cyclosuffrobuxine K C-912

3-(Dimethylamino)-14-methyl-4-methylene-9,19-cyclopregn-17(20)-en-16-one.
Cyclosuffrobuxine
[7671-41-2]



C₂₅H₃₇NO 367.573

Alkaloid from *Buxus microphylla* (Buxaceae). Needles (MeOH). Mp 167-172°. [α]_D²⁰ -92 (c, 0.42 in CHCl₃). Possible artifact.

N-De-Me: Cyclosuffrobuxinine M. *Cyclosuffrobuxinine*
[7759-55-9]

C₂₄H₃₅NO 353.547

Alkaloid from *Buxus microphylla* (Buxaceae). Needles (Et₂O or petrol). Mp 144-149°. [α]_D²⁰ -67 (c, 0.64 in CHCl₃). Note difference in nomenclature between this and its N-Me analogue (Cyclosuffrobuxine), which is presumably accidental. Poss. artifact resulting from deamination of a 20-amino compd. during isol.

4β,23-Dihydro: Cyclobuxosuffrine K. *Cyclobuxosuffrine*
[10088-22-9]

C₂₅H₃₉NO 369.589

Alkaloid from *Buxus microphylla* (Buxaceae). Needles (EtOH). Mp 201-204°. [α]_D²⁰ -62 (c, 0.52 in CHCl₃). Poss. an artifact.

17Z-Isomer, N-de-Me: trans-Cyclosuffrobuxinine M
[38774-68-4]

C₂₄H₃₅NO 353.547

Alkaloid from *Buxus sempervirens* and *Buxus longifolia* (Buxaceae). [α]_D²⁰ -47 (CHCl₃). Possible artifact.


Nakano, T. *et al.*, *J.C.S. (C)*, 1966, 1412-1421 (*Cyclosuffrobuxine K, Cyclosuffrobuxinine M*)

Votický, Z. *et al.*, *Chem. Zvesti.*, 1972, **26**, 376-380 (*trans-Cyclosuffrobuxinine M*)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997, **60**, 976-981 (*trans-Cyclosuffrobuxinine M*)

Cyclothiazomycin C-913

[133352-26-8]



C₅₉H₆₄N₁₈O₁₄S₇ 1473.729


Cyclic peptide antibiotic. Prod. by a *Streptomyces* sp. Renin inhibitor. Light-brown amorph. powder. Sol. H₂O; fairly sol. MeOH, DMSO; poorly sol. Me₂CO, hexane. Mp 210° dec. [α]_D²⁶ +46 (c, 0.69 in H₂O). λ_{max} 222 (ε 8700); 270 (ε 4100); 284 (sh); 300 (sh); 320 (sh) (MeOH) (Derep).

Aoki, M. *et al.*, *J. Antibiot.*, 1991, **44**, 582 (*isol. pmr, cmr, ir, uv, props*)

Aoki, M. *et al.*, *Tet. Lett.*, 1991, **32**, 217; 221 (*struct*)

Cyclotryprostatin A C-914

[111468-06-5]



Relative Configuration

C₂₂H₂₅N₃O₅ 411.457

Prod. by *Aspergillus fumigatus* BM939. Mammalian cell cycle inhibitor; microtubule polymerisation inhibitor. Pale yellow cryst. powder. Mp 180-185° dec. [α]_D²³ +104.3 (c, 0.07 in CHCl₃). Closely related to Fumitremorgin C, F-190. λ_{max} 226 (ε 27660); 259 (ε 7230); 265 (sh) (ε 6860); 294 (ε 7070); 303 (sh) (ε 5510) (MeOH).

13-Me ether: Cyclotryprostatin B

[184305-67-7]

C₂₃H₂₇N₃O₅ 425.483

From *Aspergillus fumigatus* BM939. Mammalian cell cycle inhibitor; microtubule polymerisation inhibitor. Pale yellow cryst. Mp 159-165° dec. [α]_D²⁴ +95.7 (c, 0.36 in CHCl₃). λ_{max} 224 (ε 41690); 262 (ε 7180); 267 (ε 7180); 295 (ε 8290); 303 (sh) (ε 6590) (MeOH).

Demethoxy, 12-epimer: Cyclotryprostatin C

[111427-97-5]

C₂₁H₂₃N₃O₄ 381.43

From *Aspergillus fumigatus* BM939. Mammalian cell cycle inhibitor; microtubule polymerisation inhibitor. Needles. Mp 221-223° dec. [α]_D²⁵ +23.4 (c, 1.0 in CHCl₃). λ_{max} 224 (ε 19680); 272 (ε 6570); 280 (sh) (ε 6325); 290 (sh) (ε 5220) (MeOH).

Demethoxy, 12-epimer, 13-ketone: Cyclotryprostatin D

[184305-68-8]

C₂₁H₂₁N₃O₄ 379.415

From *Aspergillus fumigatus* BM939. Mammalian cell cycle inhibitor. Pale yellow amorph. powder. [α]_D²⁶ +88 (c, 0.10 in CHCl₃). λ_{max} 216 (ε 18290); 240 (sh) (ε 12320); 247 (ε 15310); 264 (ε 9720); 267 (sh) (ε 9570); 301 (ε 8550) (MeOH).


Cui, C.-B. *et al.*, *Tetrahedron*, 1997, **53**, 59-72 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kondoh, M. *et al.*, *J. Antibiot.*, 1998, **51**, 801-804 (*activity*)

Afiyatullo, S.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 615-617 (*isol, pmr, cmr*)

Cycloviroboxeine I C-915

3,20-Diamino-4,4,14-trimethyl-9,19-cyclopregn-6-en-16-ol, 9CI



C₂₄H₄₀N₂O 372.593

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Cycloviroboxeine I, is unknown.

N³,N³,N²⁰-Tri-Me: Cycloviroboxeine B
[5672-25-3]

C₂₇H₄₆N₂O 414.673

Alkaloid from *Buxus malayana* and *Buxus sempervirens* (Buxaceae). Mp 203° (198-200 dec.). [α]_D²⁰ -75 (c, 1 in CHCl₃).

N³,N³,N²⁰-Tri-Me, N²⁰-formyl: N-Formylcycloviroboxeine B. *N-Formylcycloviroboxeine B (incorr.)*, 9CI
[30416-90-1]

C₂₈H₄₆N₂O₂ 442.684

Alkaloid from *Buxus malayana* (Buxaceae). Cryst. (Me₂CO). Mp 290°. [α]_D²⁰ -157. Mol. formula incorrectly given as C₂₈H₄₈N₂O₂ in the paper. Reported pmr spectrum shows olefinic protons.

N³,N³,N²⁰-Tri-Me, N,O-Di-Ac: Mp 172°.
[α]_D²⁰ -153 (c, 1 in CHCl₃).

N³,N³,N²⁰-Tri-Me, O-tigloyl: O-Tigloylcycloviroboxeine B

[16974-75-7]

C₃₂H₅₂N₂O₂ 496.775

Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 178-183°. [α]_D²⁷ -150 (c, 1.03 in CHCl₃).

N³,N³,N²⁰-Tri-Me, O-(4-hydroxycinnamoyl): Cyclomalayanine. Cyclomalayanine B

C₃₆H₅₂N₂O₃ 560.818

Alkaloid from *Buxus malayana* (Buxaceae). Mp 170°. [α]_D²⁰ -61 (c, 1 in CHCl₃).

N-Tetra-Me: Cycloviroboxeine A

[7727-91-5]

C₂₈H₄₈N₂O 428.7

Alkaloid from *Buxus malayana* and *Buxus sempervirens* (Buxaceae). Needles or prisms (Me₂CO). Mp 220°. [α]_D²⁰ -87 (c, 1.0 in CHCl₃).


Khuong-Huu-Lainé, F. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 758-762 (*Cycloviroboxeines A,B, isol, struct, pmr, ms*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1966, **31**, 608-610 (*isol, struct, Cycloviroboxeine B*)

Khuong-Huu-Lainé, F. *et al.*, *Ann. Pharm. Fr.*, 1970, **28**, 211-222 (*N-Formylcycloviroboxeine B, isol, ir, ms, pmr*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791-1794 (cmr)


Cycloviobuxine F C-916
[194852-52-3]



$C_{35}H_{48}N_2O_4$ 560.775
Alkaloid from the leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. $[\alpha]_D^{22} +7$ (c, 0.1 in $CHCl_3$). λ_{max} 230 (log ϵ 4.31); 238 (log ϵ 4.32); 246 (log ϵ 4.4); 254 (log ϵ 4.35) (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997, **60**, 976-981 (isol, uv, ir, pmr, cmr, ms)

Cycloviobuxine I C-917
3,20-Diamino-4,4,14-trimethyl-9,19-cyclopropan-16-ol, 9CI



$C_{24}H_{42}N_2O$ 374.609

The suffix letter in *Buxus* alkaloids systematically designates the *N*-subn. pattern. The parent compd., Cycloviobuxine I, is unknown.

N^3, N^{20} -Di-Me: **Cycloviobuxine D. Buxine**
[860-29-7]

$C_{26}H_{46}N_2O$ 402.662

Alkaloid from *Buxus wallichiana*, *Buxus sempervirens argentea*, *Buxus sempervirens* and *Buxus microphylla* (Buxaceae). Antiarrhythmic agent. Mp 205-210° Mp 221°.

N^3, N^{20} -Di-Me, N^3, N^{20} , O-tri-Ac: Mp 231°.

N^3, N^{20} -Di-Me, O-(3-hydroxy-4-methoxybenzoyl): **O-Vanilloylcycloviobuxine D**
[31687-96-4]

$C_{34}H_{52}N_2O_4$ 552.796

Alkaloid from *Buxus malayana* (Buxaceae). Cryst. (EtOAc). Mp 210°. $[\alpha]_D$ 0 ($CHCl_3$). Incorr. referred to as O-Vanilloylcycloviobuxine D in the paper.

N^{20}, N^{20} -Di-Me: **Cycloviobuxine F**
[100676-13-9]

$C_{26}H_{46}N_2O$ 402.662

Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 224-226°. $[\alpha]_D$ +52.71 (c, 0.7 in $CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me: **Cycloviobuxine C**

[30276-35-8]

$C_{27}H_{48}N_2O$ 416.689

Alkaloid from *Buxus malayana*, *Buxus microphylla* and *Buxus sempervirens* (Buxaceae). Mp 201°. $[\alpha]_D$ +65 ($CHCl_3$).

N^3, N^{20}, N^{20} -Tri-Me, N^3 -Ac: [30276-39-2]
Mp 216°. $[\alpha]_D$ -7 ($CHCl_3$).

N^3, N^3, N^{20}, N^{20} -Tetra-Me: **Cycloviobuxine A**, **N,N'-Dimethylcycloviobuxine D**
Semisynthetic. Needles (Me_2CO). Mp 241-242°.

N^3 -Isopropylidene, N^{20}, N^{20} -di-Me: **Buxmicrophylline A**
[188948-30-3]

$C_{29}H_{50}N_2O$ 442.727

Alkaloid from *Buxus microphylla* (Buxaceae). Mp 215-217°. $[\alpha]_D^{23} +53.1$ (c, 0.73 in $CHCl_3$).

Me ether, N^3, N^{20} -di-Me: **O-Methylcycloviobuxine D**
[108195-58-0]

$C_{27}H_{48}N_2O$ 416.689

Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (EtOH). Mp 231-233°. $[\alpha]_D$ +83.52 (c, 0.903 in $CHCl_3$).

1,2-Didehydro, N^3, N^3, N^{20} -tri-Me: **Buxasamarine**
[194852-54-5]

$C_{27}H_{46}N_2O$ 414.673

Alkaloid from the leaves of *Buxus longifolia* (Buxaceae). Amorph. solid. $[\alpha]_D^{25} +23$ (c, 0.1 in $CHCl_3$).

6,7-Didehydro: see Cycloviobuxine I, C-915

11-Oxo, N^{20}, N^{20} -di-Me, N^3 -benzoyl:

Buxarine F. Buxarine

[11011-80-6]

$C_{33}H_{48}N_2O_3$ 520.754

Struct. not certain. Alkaloid from *Buxus sempervirens* (Buxaceae). Prisms (Me_2CO). Mp 210-212°. $[\alpha]_D$ +98 ($CHCl_3$).

11-Oxo, N^{20}, N^{20} -di-Me, N^3 -benzoyl, dihydrochloride: Mp 257-260°.

3-Epimer, N-tetra-Me: Semisynthetic.

Cryst. (Me_2CO). Mp 300°. $[\alpha]_D$ +50 (c, 2.22 in $CHCl_3$). Presumed 3 α -epimer of Cycloviobuxine A. Attempted interconversion was unsuccessful.

3-Epimer, 11,12-didehydro: 3,20-Diamino-4,4,14-trimethyl-9,19-cyclopropan-11-en-16-ol. **Cyclokoreanine A**

$C_{24}H_{40}N_2O$ 372.593

Unknown as nat. prod.

3-Epimer, 11,12-didehydro, N^3, N^3, N^{20} -tri-Me: **Cyclokoreanine B**

[10413-97-5]

$C_{27}H_{46}N_2O$ 414.673

Alkaloid from *Buxus koreana* and *Buxus microphylla* var. *sinica* (Buxaceae). Mp 235-236°. $[\alpha]_D$ +109 (c, 1.65 in $CHCl_3$). Struct. not conclusively proven. Unusual 3 α -config. for *Buxus* alkaloid.

3-Epimer, 11,12-didehydro, N-tetra-Me:

N-Methylcyclokoreanine B

[10283-44-0]

$C_{28}H_{48}N_2O$ 428.7

Semisynthetic. Cryst. (Me_2CO). Mp 240-241°. $[\alpha]_D$ +92 (c, 0.85 in $CHCl_3$).

Nakano, T. *et al.*, *J.C.S. (C)*, 1966, 1805-1810 (synth, **Cyclokoreanine B**)

Döpke, W. *et al.*, *Pharmazie*, 1966, **21**, 643;

CA, **66**, 65668f (**Buxarine F**)

Khuong-Huu, F. *et al.*, *Ann. Pharm. Fr.*, 1970, **28**, 211-222 (*Buxus malayana constitis*)

Vassová, A. *et al.*, *Pharmazie*, 1970, **25**, 363-365; *CA*, 1970, **73**, 127748n (**Cycloviobuxine D**)

Bauerová, O. *et al.*, *Pharmazie*, 1973, **28**, 212-214 (**Cyclokoreanine B**)

Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*,

1975, **40**, 3055-3060 (**Cycloviobuxine C**)

Abramson, D. *et al.*, *Phytochemistry*, 1977, **16**,

1935-1937 (biosynth)

Khodzhaev, B.U. *et al.*, *Khim. Prir. Soedin.*,

1980, **16**, 130-131 (**Cycloviobuxine D**)

Hu, S. *et al.*, *CA*, 1981, **95**, 108438z

(pharmacol)

Khodzhaev, B.U. *et al.*, *Khim. Prir. Soedin.*,

1985, **21**, 718-719; 1986, **22**, 799; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 679-680;

1986, **22**, 743-744 (**Cycloviobuxine F**, **O-Methylcycloviobuxine-D**)

Du, J. *et al.*, *Zhiv. Xuebao (Acta Bot. Sin.)*,

1996, **38**, 483-488; *CA*, **126**, 274768k

(**Buxmicrophylline A**, *cryst struct*)


Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997,

60, 976-981 (**Buxasamarine**)

Liu, H.-F. *et al.*, *Acta Cryst. E*, 2006, **62**,

o4517-o4518 (synth)

3,5'-Cycloxanthosine C-918
[10380-93-5]



$C_{10}H_{10}N_4O_5$ 266.213


Alkaloid from the marine sponge *Erylus* sp. Cryst. (H_2O). Mp 310-314°. $[\alpha]_D$ -18 (c, 0.02 in DMSO). λ_{max} 236 (log ϵ 3.91); 266 (log ϵ 3.99) (H_2O).

Hampton, A. *et al.*, *J.O.C.*, 1967, **32**, 1688-1691 (synth)

Chen, G.S. *et al.*, *Nucleosides Nucleotides*, 2004, **23**, 347-359 (synth)

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1689-1691 (isol, pmr, cmr, ms)

Cycloxazoline C-919
Trisoxazoline. Westiellamide
[131998-54-4]



C₂₇H₄₂N₆O₆ 546.665


Isol. from the ascidian *Lissoclinum bis-tratum* and *Westiellopsis prolifica*. Cytotoxic. Amorph. [α]_D +30 (c, 0.1 in MeOH).

Prinsep, M.R. *et al.*, *J. Nat. Prod.*, 1992, **55**, 140-142 (*isol, pmr, cmr, struct*)

Hambley, T.W. *et al.*, *Tetrahedron*, 1992, **48**, 341-348 (*isol, pmr, cmr, struct*)

Wipf, P. *et al.*, *Tetrahedron*, 2000, **56**, 9143-9150 (*synth*)

Cycloboxine I **C-920**
3,20-Diamino-4,4,14-trimethyl-9,19-cyclo-pregnan-11-one

C₂₄H₄₀N₂O 372.593

The suffix letter in *Buxus* alkaloids unambiguously designates the *N*-subn. pattern. The parent compd., Cycloboxine I, is unknown.

N²⁰,N²⁰-Di-Me, N³-(2-methylpropa-noyl): N³-Isobutyrylcycloboxine F. *Baleabuxine* [2988-97-8]

C₃₀H₅₀N₂O₂ 470.737

Alkaloid from leaves of *Buxus balearica* (Buxaceae). Hemihydrate. Mp 258-259°. [α]_D +115 (CHCl₃).

N²⁰,N²⁰-Di-Me, N³-benzoyl: N³-Benzoyl-cycloboxine F. *Buxatine* [16974-67-7]

C₃₃H₄₈N₂O₂ 504.754


Alkaloid from *Buxus sempervirens* (Buxaceae). Needles (Me₂CO). Mp 214-217° dec. [α]_D +112 (c, 0.2 in CHCl₃) (+90).

Herlem-Gaulier, D. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 657-668; 1966, 3478-3486 (*Baleabuxine*)

Döpke, W. *et al.*, *Naturwissenschaften*, 1967, **54**, 249; *CA*, **67**, 82292s (*Buxatine*)

Kupchan, S.M. *et al.*, *Tetrahedron*, 1967, **23**, 4563-4586 (*N³-Benzoylcycloboxine F*, *isol, pmr*)

Cycloboxoline I **C-921**
3,20-Diamino-28-hydroxy-4,4,14-tri-methyl-9,19-cyclopregnan-11-one

C₂₄H₄₀N₂O₂ 388.592

In *Buxus* alkaloids the suffix letter unambiguously designates the subn.

pattern. The parent compd., Cycloboxoline I, is unknown.

N²⁰,N²⁰-Di-Me, N³-benzoyl: N-Benzoyl-cycloboxoline F

C₃₃H₄₈N₂O₃ 520.754

Alkaloid from *Buxus sempervirens* (Buxaceae). Cryst. (Me₂CO). Mp 255-256°. [α]_D +76 (c, 0.84 in CHCl₃).

N²⁰,N²⁰-Di-Me, N³-benzoyl, O-Ac: N-Benzoyl-O-acetylcycloboxoline F [16974-76-8]


C₃₅H₅₀N₂O₄ 562.791

Alkaloid from *Buxus sempervirens* (Buxaceae). Mp 216-218°. [α]_D +114 (c, 1.0 in CHCl₃).

Kupchan, S.M. *et al.*, *Tetrahedron*, 1967, **23**, 4563 (*isol, pmr*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, 1791 (*config*)

Cyclozoanthamine **C-922**
[159509-35-0]

C₃₀H₄₁NO₆ 511.657

The abs. config. is prob. as shown, by analogy with Norzoanthamine, N-317. Alkaloid from *Zoanthus* sp. Cytotoxic agent. Oil.

Japan. Pat., 1994, 94 199 867; *CA*, **122**, 9777d

Fukuzawa, S. *et al.*, *Heterocycl. Commun.*, 1995, **1**, 207-214


Cygnine **C-923**

C₁₉H₂₂N₂O₃ 326.394

Struct. unknown. MF given in abstract is C₁₀H₂₂N₂O₃ - this is prob. incorr. (see Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1497-1500). Alkaloid from *Gastrolobium calycinum* (Fabaceae).

Mann, E.A. *et al.*, *Proc. R. Soc. London, B*, 1907, **79**, 485-491 (*isol*)

Cylindramide **C-924**
[147362-39-8]

C₂₇H₃₄N₂O₅ 466.576

Relative Configuration

Tetramic acid antibiotic. Shows enol tautomerism over the dioxopyrrolidine-carbonyl system. Isol. from the sponge *Halichondria cylindrata*. Cytotoxic agent. Wax. [α]_D²⁵ +167 (c, 0.4 in MeOH). λ_{\max} 265 (€ 22000) (MeOH).

17,29-Didehydro, Mg salt (2:1): *Geodin A*

C₅₄H₆₂MgN₄O₁₀ 951.41

Isol. from the sponge *Geodia* sp. Nematocide. Amorph. solid. Mp 173° dec. [α]_D²⁰ +179 (c, 1 in DMSO). Stereochem. not fully determined. Mg coordinated to the dioxopyrrolidine-carbonyl chromophore with charge/bond delocalisation. λ_{\max} 264 (€ 19100) (DMSO).

[153175-60-1]

Kanazawa, S. *et al.*, *Tet. Lett.*, 1993, **34**, 1065-1068 (*isol, pmr, cmr*)


Capon, R.J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1256-1259 (*Geodin A*)

Cramer, N. *et al.*, *Chem. Eur. J.*, 2006, **12**, 2488-2503 (*synth, pmr, cmr*)

Hart, A.C. *et al.*, *J.A.C.S.*, 2006, **128**, 1094-1095 (*synth*)

Cylindricine B **C-925**

[152273-68-2]



Relative configuration


C₁₉H₃₂ClNO 325.921

Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

Blackman, A.J. *et al.*, *Tetrahedron*, 1993, **49**, 8645-8656 (*isol, pmr, cmr, ms, cryst struct*)

Liu, J.F. *et al.*, *J.O.C.*, 1999, **64**, 8263-8266 (*synth*)

Cylindricine C **C-926**



(-)-form

C₁₉H₃₃NO₂ 307.475

(-)-form [156953-77-4]

Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

I'-Ac: *Cylindricine E*

[156953-79-6]

C₂₁H₃₅NO₃ 349.512

From *Clavelina cylindrica*. Light yellow oil.

Me ether: *Cylindricine D*

[156953-78-5]

C₂₀H₃₅NO₂ 321.502

From *Clavelina cylindrica*. Light yellow oil.

1'-Deoxy, 1'-thiocyanato: Cylindricine F
[156953-80-9]

$C_{20}H_{32}N_2OS$ 348.552

Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

1'-Deoxy, 1'-chloro: Cylindricine A
[152273-89-7]

$C_{19}H_{32}ClNO$ 325.921

Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

7-Deoxy, 10-oxo, 8,9-didehydro, 1'-deoxy, 1'-chloro: Cylindricine K
[164740-28-7]

$C_{19}H_{30}ClNO$ 323.905

Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

(+)-form

Deoxy: Lepadiformine A

[155944-27-7]

$C_{19}H_{35}NO$ 293.492

Alkaloid from the ascidians *Clavelina lepadiformis* and *Clavelina moluccensis*. Exhibits moderate cytotoxicity. Oil. $[\alpha]_D^{28}$ -15 (c, 0.37 in MeOH) (synthetic). Struct. revised in 2002. λ_{max} 206 (log ϵ 2.74); 310 (log ϵ 1.56) (EtOH).

Blackman, A.J. *et al.*, *Tetrahedron*, 1993, **49**, 8645-8656 (*Cylindricine A*, *isol*, *pmr*, *cmr*, *ms*, *cryst* *struct*)

Li, C. *et al.*, *Aust. J. Chem.*, 1994, **47**, 1355-1361; 1995, **48**, 955-965 (*Cylindricines C-F and K*)

Biard, J.F. *et al.*, *Tet. Lett.*, 1994, **35**, 2691-2694 (*Lepadiformine A*)

Snider, B.B. *et al.*, *J.O.C.*, 1997, **62**, 5630-5633 (*Cylindricines*, *synth*)

Werner, K.M. *et al.*, *J.O.C.*, 1999, **64**, 686-687 (*synth*)

Molander, G.A. *et al.*, *J.O.C.*, 1999, **64**, 5183-5187 (*Cylindricine C*, *synth*)

Liu, J.F. *et al.*, *J.O.C.*, 1999, **64**, 8263-8266 (*Cylindricine A*, *synth*)

Greshock, T.J. *et al.*, *Org. Lett.*, 2001, **3**, 3511-3514 (*synth*)

Sun, P. *et al.*, *J.O.C.*, 2002, **67**, 4337-4345 (*synth*, *abs* *config*)

Kibayashi, C. *et al.*, *Bull. Chem. Soc. Jpn.*, 2003, **76**, 2059-2074 (*synth*, *abs* *config*)

Trost, B.M. *et al.*, *Org. Lett.*, 2003, **5**, 4599-4602 (*synth*)

Canesi, S. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 4336-4338 (*synth*)

Arai, T. *et al.*, *Tet. Lett.*, 2004, **45**, 5921-5924 (*synth*)

Abe, H. *et al.*, *J.A.C.S.*, 2005, **127**, 1473-1480 (*synth*, *abs* *config*)

Wang, J. *et al.*, *Heterocycles*, 2006, **70**, 423-459 (*synth*)

Sauviat, M.-P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 558-562 (*isol*, *pmr*, *cmr*, *activity*)

Lee, M. *et al.*, *Org. Lett.*, 2006, **8**, 745-748 (*synth*)

Swidorski, J.J. *et al.*, *Org. Lett.*, 2006, **8**, 777-780 (*synth*, *abs* *config*, *bibl*)

Schär, P. *et al.*, *Org. Lett.*, 2006, **8**, 1569-1571 (*synth*)

Caldwell, J.J. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2631-2634 (*synth*)


Mihara, H. *et al.*, *Heterocycles*, 2007, **72**, 421-438 (*Cylindricine C*, *synth*)

Lygo, B. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 3085-3090 (*Lepadiformine A*, *synth*)

Flick, A.C. *et al.*, *Org. Lett.*, 2008, **10**, 1871-1874 (*Cylindricine C*, *synth*)

Cylindricine G

[156953-81-0]



Relative Configuration


$C_{18}H_{28}N_2OS$ 320.498

Abs. config. not determined, but redrawn here to correspond with Lepadiformine A in C-926. Alkaloid from the ascidian *Clavelina cylindrica*. Light yellow oil.

Li, C. *et al.*, *Aust. J. Chem.*, 1994, **47**, 1355-1361 (*isol*, *pmr*, *cmr*, *ms*, *struct*)

Cylindricine J

[164740-27-6]



$C_{20}H_{32}N_2O_2S$ 364.551

Alkaloid from the ascidian *Clavelina cylindrica*. Oil.

Thiocyanate: [404850-27-7]

$C_{20}H_{32}N_2O_2S$ 364.551


Alkaloid from the stems of *Indigofera longiracemosa*. Cryst. (CH_2Cl_2 /MeOH). Mp 181-182°. $[\alpha]_D$ -52.7 (c, 0.31 in $CHCl_3$). Has -SCN replacing -NCS. λ_{max} 218 (log ϵ 2.8); 229 (log ϵ 2.96); 247 (log ϵ 3.29); 258 (log ϵ 3.37) (MeOH).

Li, C. *et al.*, *Aust. J. Chem.*, 1995, **48**, 955-965 (*isol*, *ir*, *pmr*, *cmr*, *ms*)

Thangadurai, D. *et al.*, *Nat. Prod. Lett.*, 2001, **15**, 287-290 (*thiocyanate*)

Cylindrocarine

C-929



$C_{21}H_{28}N_2O_3$ 356.464

(-)-form [22222-79-3]

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Mp 204-205°. $[\alpha]_D^{20}$ -280 (c, 0.0025 in MeOH).

N-Formyl: N-Formylcylindrocarine

[22222-81-7]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Mp 161-162°. $[\alpha]_D^{20}$ -140 (c, 0.01 in MeOH).

N-Ac: Cylindrocarpine

C-927

[6858-95-3]

$C_{23}H_{30}N_2O_4$ 398.501

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Mp 120-121.5°. $[\alpha]_D^{27}$ -122 (c, 0.72 in $CHCl_3$).

N-Propanoyl: Homocylindrocarpine

[17391-26-3]

$C_{24}H_{32}N_2O_4$ 412.528

Alkaloid from *Tabernaemontana amygdalaeifolia* (Apocynaceae). Bp_{0.001} 120°. $[\alpha]_D^{30}$ -82 (c, 1.7 in $CHCl_3$).

N-Benzoyl: N-Benzoylcylindrocarine

[22222-82-8]

$C_{28}H_{32}N_2O_4$ 460.572

Alkaloid in *Aspidosperma cylindrocarpon* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -131 (c, 0.075 in MeOH).

N-Cinnamoyl: Cylindrocarpine

[6793-70-0]

[58581-12-7]

$C_{30}H_{34}N_2O_4$ 486.61

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Mp 168-169°. $[\alpha]_D^{21}$ -181 (c, 0.4 in $CHCl_3$).

N-Me: N-Methylcylindrocarine

[22226-36-4]

$C_{22}H_{30}N_2O_3$ 370.491

Alkaloid from *Aspidosperma cylindrocarpon*. Noncryst. $[\alpha]_D^{20}$ -110 (c, 0.001 in MeOH).

5-Oxo, N-Ac: 5-Oxocylindrocarpine. 10-Oxocylindrocarpine

[16531-05-8]

$C_{23}H_{28}N_2O_5$ 412.485

Alkaloid from *Tabernaemontana amygdalaeifolia* (Apocynaceae). Mp 217-218° (213°). $[\alpha]_D^{30}$ -64 (c, 0.38 in $CHCl_3$) (-39).

Demethoxy, N-Ac: Demethoxycylindrocarpine. N-Acetyldemethoxycylindrocarpine

[17391-27-4]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from *Tabernaemontana amygdalaeifolia* (Apocynaceae). Bp_{0.001} 120°. $[\alpha]_D^{30}$ -49 ($CHCl_3$).

20ξ-Hydroxy: 20-Hydroxycylindrocarine

[22222-83-9]

$C_{21}H_{28}N_2O_4$ 372.463

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -300 (c, 0.004 in MeOH).

20ξ-Hydroxy, N-formyl: N-Formyl-20-hydroxycylindrocarine

[22226-30-8]

$C_{22}H_{28}N_2O_5$ 400.474

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -250 (c, 0.002 in MeOH).

20ξ-Hydroxy, N-Ac: N-Acetyl-20-hydroxycylindrocarine

[22226-31-9]

$C_{23}H_{30}N_2O_5$ 414.5

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Cubes (MeOH). Mp 207-210°. $[\alpha]_D^{20}$ -400 (c, 0.005 in MeOH).

20ξ-Hydroxy, N-Benzoyl: N-Benzoyl-20-hydroxycylindrocarine

[23334-95-4]

$C_{28}H_{32}N_2O_5$ 476.571

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Cubes (MeOH).

$[\alpha]_D^{20}$ -150 (c, 0.013 in MeOH).

20 ξ -Hydroxy, N-cinnamoyl: N-Cinnamoyl-20-hydroxycylindrocarine
[22226-29-5]

C₃₀H₃₄N₂O₅ 502.609

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Needles (MeOH). Mp 123-124°. $[\alpha]_D^{20}$ -384 (c, 0.0042 in MeOH).

20 ξ -Hydroxy, N-(3-phenylpropanoyl): N-Dihydrocinnamoyl-20-hydroxycylindrocarine
[22344-67-8]

C₃₀H₃₆N₂O₅ 504.625

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -278 (c, 0.0057 in MeOH).

(±)-form [58566-52-2]

Synthetic. Noncryst.

N-Ac: [58566-53-3]

Synthetic. Mp 178-179°.

Demethoxy, N¹-Ac: (±)-Demethoxycylindrocarpine

Alkaloid from *Aspidosperma cylindrocarpon* (Apocynaceae). Needles. Mp 160-162°.

Achenbach, H. *et al.*, *Tet. Lett.*, 1967, 1793 (5-Oxocylindrocarpine)

Achenbach, H. *et al.*, *Z. Naturforsch., B*, 1967, **22**, 955 (*derivs*)

Milborrow, B.V. *et al.*, *J.C.S. (C)*, 1969, 417 (*isol, uv, ir, ms, pmr, struct, Cylindrocarine, 20-Hydroxycylindrocarine derivs, N-Acetyldemethoxycylindrocarine*)


Gebreyesus, T. *et al.*, *J.C.S. Perkin I*, 1972, 849 (5-Oxocylindrocarpine)

Lawson, G. *et al.*, *Tetrahedron*, 1977, **33**, 1641 (*synth*)

Cylindrospermopsin

C-930

[143545-90-8]



Absolute Configuration

C₁₅H₂₁N₅O₇S 415.426

Exists as prototropic zwitterion. C-7 config. revised in 2001. Alkaloid from the cyanobacteria *Aphanizomenon ovalisporum*, *Cylindrospermopsis raciborskii* and *Umezakia natans*. Potent hepatotoxin, glutathione synthesis inhibitor. Off-white microcryst. $[\alpha]_D$ -31 (c, 0.1 in H₂O). λ_{max} 262 (ε 5800); 290 (sh) (ε 2100) (H₂O) (Derep).

▶ LD₅₀ (mus, ipr) 0.2 mg/kg, LD₅₀ (mus, ipr) 2.1 mg/kg.

7-Deoxy: 7-Deoxycylindrospermopsin

[221554-69-4]

C₁₅H₂₁N₅O₆S 399.427

Alkaloid from *Cylindrospermopsis raciborskii*.

7-Epimer: 7-Epicylindrospermopsin

[265652-18-4]

C₁₅H₂₁N₅O₇S 415.426

Alkaloid from *Aphanizomenon ovalisporum*. Hepatotoxin. Amorph. solid. $[\alpha]_D^{25}$ -20.5 (c, 0.06 in H₂O).

Ohtani, I. *et al.*, *J.A.C.S.*, 1992, **114**, 7941-7942 (*isol, uv, pmr, cmr, ms*)

Norris, R.L. *et al.*, *Environ. Toxicol.*, 1999, **14**, 163-165 (7-Deoxycylindrospermopsin)

Banker, R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 387-389 (7-epimer)

Xie, C. *et al.*, *J.A.C.S.*, 2000, **122**, 5018-5024 (*synth*)

Burgoyne, D.L. *et al.*, *J.O.C.*, 2000, **65**, 152-156 (*biosynth*)

Heintzelman, G.R. *et al.*, *J.A.C.S.*, 2002, **124**, 3939-3945 (*synth, abs config*)

White, J.D. *et al.*, *J.O.C.*, 2005, **70**, 1963-1977 (*synth, abs config*)


Looper, R.E. *et al.*, *Tetrahedron*, 2006, **62**, 4549-4562 (*synth, abs config*)

Mihali, T.K. *et al.*, *Appl. Environ. Microbiol.*, 2008, **74**, 716-722 (*biosynth*)

Cynodine

C-931

[50656-84-3]



C₂₃H₂₃N₃O₃ 389.453

Alkaloid from the leaves of *Cynometra ananta*, and from the stem bark and seeds of *Cynometra hankei* (Fabaceae). Mp 155°. $[\alpha]_D$ +15.

N¹-De-Me: N¹-Demethylcynodine

[81345-39-3]

C₂₂H₂₁N₃O₃ 375.426

Minor alkaloid from the stem bark of *Cynometra hankei*. Amorph. brown gum.


Khuong-Huu, F. *et al.*, *Tet. Lett.*, 1973, 1757 (*struct*)

Waterman, P.G. *et al.*, *Phytochemistry*, 1981, **20**, 2765 (*isol, uv, ir, pmr, ms, deriv*)

Cynolujine

C-932

[85644-17-3]



C₁₆H₁₉N₃O₂ 285.345


Major alkaloid from the leaves of *Cynometra lujae* (Fabaceae). Cryst. (Me₂CO). Mp 108°. $[\alpha]_D^{20}$ -6 (c, 1 in CHCl₃).

Tchissambou, L. *et al.*, *Tetrahedron*, 1982, **38**, 2687 (*isol, ir, pmr, cmr, struct*)

Cynometrine

C-933

[50656-83-2]



Absolute Configuration

C₁₆H₁₉N₃O₂ 285.345

Alkaloid from the leaves of *Cynometra ananta* and *Cynometra lujae*, and from the stem bark and seeds of *Cynometra hankei* (Fabaceae). Needles (CHCl₃). Mp 211°. $[\alpha]_D^{20}$ -27.1 (c, 0.1 in CHCl₃). λ_{max} 258 (ε 257); 264 (ε 214) (EtOH) (Derep).

N¹-De-Me: N¹-Demethylcynometrine

[81345-38-2]

C₁₅H₁₇N₃O₂ 271.318

Alkaloid from the stem bark and seeds of *Cynometra hankei*. Needles (CHCl₃). Mp 204°. $[\alpha]_D^{20}$ -56.2 (c, 0.1 in CHCl₃).

Khuong-Huu, F. *et al.*, *Tet. Lett.*, 1973, 1757-1760 (*uv, ir, pmr, cmr, ms, struct*)

Chiaroni, A. *et al.*, *J. Chem. Res., Synop.*, 1981, 182; *J. Chem. Res., Miniprint*, 1981, 2116-2145 (*cryst struct, abs config*)


Waterman, P.G. *et al.*, *Phytochemistry*, 1981, **20**, 2765-2767 (*isol, uv, ir, pmr, cmr, ms, N¹-de-Me*)

Tchissambou, L. *et al.*, *Tetrahedron*, 1982, **38**, 2687-2695 (*isol, cmr, synth*)

Fishwick, C.W.G. *et al.*, *Tet. Lett.*, 1996, **37**, 3915-3918 (*synth*)

Cynthichlorine

C-934



C₁₁H₈Cl₃NO₃ 308.547

(±)-form


Isol. from the tunicate *Cynthia savignyi*. Antifungal and antibacterial agent. Cytotoxic.

Abourriche, A. *et al.*, *Farmaco*, 2003, **58**, 1351-1354 (*isol, pmr*)

Cypholophine

C-935

4-[2-(3,4-Dimethoxyphenyl)ethyl]-5-methyl-1H-imidazole-2-butanol, 9CI
[26482-10-0]



C₁₈H₂₆N₂O₃ 318.415

Major alkaloid from the leaves of *Cypholophus friesianus* (Urticaceae). Needles (Me₂CO or EtOAc). Mp 126-127°.


O-Ac: O-Acetylcypholophine

[26482-11-1]

C₂₀H₂₈N₂O₄ 360.452

Minor alkaloid from the leaves of *Cypholophus friesianus* (Urticaceae). Gum.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1971, **24**, 857 (*isol, uv, ir, ms, pmr, synth, struct, deriv*)

Aspergillus oryzae Cysteine protease inhibitors C-936

CPI 1	R ¹ = 4-OH-PhCH ₂ , R ² = -NH(CH ₂) ₃ NH ₂
CPI 2	R ¹ = -CH(CH ₃)CH ₂ CH ₃ , R ² = NH ₂
CPI 3	R ¹ = -CH(CH ₃)CH ₂ CH ₃ , R ² = -CH ₂ NH ₂
CPI 4	R ¹ = -CH(CH ₃)CH ₂ CH ₃ , R ² = -NH(CH ₂) ₃ NH ₂
CPI 5	R ¹ = -PhCH ₂ , R ² = -NH(CH ₂) ₃ NH ₂

Tentative structs. assigned to CPI 1 and CPI 5. Prod. by *Aspergillus oryzae*. Cysteine protease inhibitors.

CPI 1

C₂₀H₃₀N₄O₆ 422.48
Shows antiallergic props. Syrup. Sol. H₂O, MeOH, DMSO. [α]_D²⁵ +153.8 (c, 1 in H₂O). λ_{max} 198 (ε 69200); 219 (sh) (ε 33800); 274 (ε 3700); 280 (sh) (ε 3300) (H₂O). λ_{max} 198 (ε 69200); 219; 274 (ε 3700); 280 (H₂O) (Berdy).

CPI 2

C₁₄H₂₅N₃O₅ 315.369
Shows antiallergic props. Powder. Sol. H₂O, DMSO, MeOH. [α]_D²⁵ +47 (c, 0.2 in H₂O). λ_{max} 198 (ε 23100) (H₂O).

CPI 3

C₁₅H₂₇N₃O₅ 329.395
Shows antiallergic props. Powder. Sol. H₂O, DMSO, MeOH. [α]_D²⁵ +86.5 (c, 0.2 in H₂O). λ_{max} 201 (ε 28800) (H₂O).


CPI 4

Kojistatin A. Antibiotic 460A
[183069-12-7]
C₁₇H₃₂N₄O₅ 372.464
Syrup. [α]_D²⁵ +57 (c, 0.2 in H₂O). λ_{max} 197 (ε 27000) (H₂O).

CPI 5

C₂₀H₃₀N₄O₅ 406.481
Syrup. Sol. H₂O, MeOH, DMSO. [α]_D²⁵ +90.5 (c, 0.04 in H₂O). λ_{max} 196 (ε 65800); 204 (ε 60000); 251 (ε 812); 257 (ε 851) (H₂O). λ_{max} 196 (ε 65800); 204 (ε 60000); 251 (ε 812); 257 (ε 851); 267 (ε 696) (MeOH) (Berdy).

Sato, N. *et al.*, *Biosci., Biotechnol., Biochem.*, 1996, **60**, 1747-1748 (*Kojistatin A*)
Yamada, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 907-914 (*isol. uv, ir, pmr, cmr*)

Cystothiazole E C-937
[214215-11-9]


Absolute Configuration

C₁₇H₂₂N₂O₂S₂ 350.505
Prod. by *Cystobacter fuscus* strain AJ-13278. Antifungal agent. Oil. [α]_D²⁴ +17.8 (c, 0.2 in CHCl₃). λ_{max} 220 (ε 23900); 247 (ε 21600); 311 (ε 11100) (MeOH).

Suzuki, Y. *et al.*, *Tetrahedron*, 1998, **54**, 11399-11404 (*isol. uv, ir, pmr, ms*)
Bach, T. *et al.*, *Chem. Eur. J.*, 2002, **8**, 5585-5592 (*synth*)
Takayama, H. *et al.*, *Heterocycles*, 2007, **71**, 75-85 (*synth*)

Cytisine C-938

1,2,3,4,5,6-Hexahydro-1,5-methano-8H-pyrido[1,2-a][1,5]diazocin-8-one, 9CI. Baptitoxine. Citisine. Sophorine†. Ulexine. Cytiton. Tabex



C₁₁H₁₄N₂O 190.244
Log P -0.94 (calc).

(-)-form [485-35-8]

Alkaloid from *Cytisus laburnum*, many other *Cytisus* spp., *Baptisia*, *Genista*, *Laburnum*, *Sophora*, *Thermopsis*, *Ammodendron*, *Anagyris* and *Euchresta* spp. (Fabaceae). Shows nicotine-like CNS activity. Antiinflammatory agent. Shows psychoactive props. Has been used as respiratory stimulant in the former USSR. Common cause of poisoning of humans and animals by *Cytisus laburnum*. Sol. H₂O, EtOH, C₆H₆; poorly sol. Et₂O, hexane. Mp 155° subl. [α]_D⁷ -119 (H₂O). Pharmacol. active isomer.

► Highly toxic. LD₅₀ (mus, orl) 101 mg/kg; LD₅₀ (mus, ivn) 1.3 mg/kg. HA4025000
Hydrochloride: [6047-01-4]
Mp 218°.

► HA4200000

Perchlorate: Mp 270°.
N^b-Oxide: Cytisine N-oxide. 12-Hydroxycytisine

[138922-22-2]
C₁₁H₁₄N₂O₂ 206.244
Alkaloid from *Sophora exigua* (Fabaceae). Pale yellow needles. Mp 238° dec. [α]_D²⁵ -209 (c, 0.11 in EtOH).

N-Formyl: N-Formyleytisine

[53007-06-0]
C₁₂H₁₄N₂O₂ 218.255
Alkaloid from *Thermopsis chinensis*, *Sophora tomentosa*, *Echinosophora koreensis* and *Euchresta* spp. (Fabaceae). Mp 170-172°. [α]_D²⁰ -233 (c, 0.43 in EtOH).

N-Ac: N-Acetylcytisine

[6018-52-6]
C₁₃H₁₆N₂O₂ 232.282
Alkaloid from *Sophora tomentosa* and *Thermopsis alterniflora* (Fabaceae). Mp 210-213° (208°). [α]_D²⁶ -208 (c, 0.182 in EtOH).

N-Acetoxy: N-Acetoxyeytisine

[145364-52-9]
C₁₃H₁₆N₂O₃ 248.281
Constit. of *Petteria ramentacea*.

N-Me: N-Methylcytisine. Caulophylline

[486-86-2]
C₁₂H₁₆N₂O 204.271

Alkaloid from *Cytisus laburnum*, *Genista* spp., *Leontice* spp., *Spartium junceum* and very many other spp. and genera in the Fabaceae. Prisms (EtOH). Mp 137°. [α]_D -221.6 (H₂O).

► HA4400000

N-Et: N-Ethylcytisine

[83728-92-1]
C₁₃H₁₈N₂O 218.298
Alkaloid from fresh flowers of *Euchresta koreensis* (Fabaceae). Needles (C₆H₆/hexane). Mp 112°. [α]_D²⁷ -216.7 (c, 0.31 in EtOH).

N-Acetamidomethyl: N-(N-Acetylaminomethyl)cytisine

C₁₄H₁₉N₃O₂ 261.323
Alkaloid from *Maackia hupehensis*. Oil. [α]_D²³ -91 (c, 0.48 in EtOH).

N-(2-Hydroxyethyl): N-(2-Hydroxyethyl)cytisine

C₁₃H₁₈N₂O₂ 234.297
Alkaloid from above-ground parts of *Sophora alopecuroides* (Fabaceae). Mp 63-65°. [α]_D²⁰ -187 (c, 0.725 in EtOH).

N-(2-Chloroethyl): Alternidine

[642-49-9]
C₁₃H₁₇ClN₂O 252.743
Artifact derived from use of ClCH₂CH₂Cl as solvent. Mp 105-107°. [α]_D -190.5.

N-(3-Butenyl): Rhombifoline

[529-78-2]
C₁₅H₂₀N₂O 244.336
Alkaloid from *Thermopsis rhombifolia*, *Thermopsis lanceolata*, *Ammodendron longiracemosum*, *Clathrotropis brachypetala*, *Sophora* spp., *Genista carinalis* and *Genista tinctoria* (Fabaceae). Yellowish oil. Bp_{0.2} 120° (bath). [α]_D²⁵ -232.4 (c, 2.13 in EtOH).

N-(3-Butenyl), hydrochloride: Mp 256-258° dec.*N-(3-Butenyl), picrate*: Mp 226° dec.*N-(3-Oxobutyl): N-(3-Oxobutyl)cytisine*

[64408-08-8]
C₁₅H₂₀N₂O₂ 260.335
Alkaloid from aerial parts of *Euchresta koreensis* (Fabaceae). Plates (C₆H₆/petrol). Mp 118°. [α]_D²¹ -211.6 (c, 0.219 in EtOH).

► Toxic, LD₅₀ (mus) 71 mg/kg.*N-Methoxycarbonyl: 12-Methoxycarbonylcytisine*

[125109-97-9]
C₁₃H₁₆N₂O₃ 248.281
Constit. of *Leontice watereri* (Fabaceae) and *Petteria ramentacea*. Possible artifact.

N-Ethoxycarbonyl: 12-Ethoxycarbonylcytisine

[132216-16-1]
C₁₄H₁₈N₂O₃ 262.308
Constit. of *Laburnum watereri* and *Spartium junceum* (Fabaceae). Possible artifact.

N-(Carboxymethyl): see 12-Cytisineacetic acid, C-939*N-(2-Oxo-1-pyrrolidinylmethyl): N-(2-Oxo-1-pyrrolidinylmethyl)cytisine*

C₁₆H₂₁N₃O₂ 287.361

Alkaloid from *Maaackia hupehensis*.
Cryst. (CHCl₃). Mp 169-170°. [α]_D²³ -167 (c, 0.36 in EtOH).

11,12-Didehydro- 11,12-Dehydrocytisine

C₁₁H₁₂N₂O 188.229

Constit. of *Petteria ramentacea*. Tentative struct.

12,13-Didehydro- 12,13-Dehydrocytisine

C₁₁H₁₂N₂O 188.229

Constit. of *Petteria ramentacea*. Tentative struct.

11-Oxo- 11-Oxocytisine

C₁₁H₁₂N₂O₂ 204.228

Alkaloid from the leaves of *Sophora secundiflora* (Fabaceae). Needles (Et₂O/MeOH). Mp 272-275°. [α]_D¹² +6.7 (c, 0.45 in EtOH).

Tetrahydro: see Tetrahydrocytisine, T-149

(±)-form

Mp 147°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 1324B (nmr)

Ing, H.R. *et al.*, *J.C.S.*, 1931, 2195; 1932, 2778 (isol, struct)

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, 21, 144 (*Rhombifoline*, isol)

Cockburn, W.F. *et al.*, *Can. J. Chem.*, 1952, 30, 92 (*Cytisine*, *Rhombifoline*, isol, struct)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1956, 89, 792; 1975, 108, 1043 (synth, ir, uv, cmr)

Pakudina, Z.P. *et al.*, *CA*, 1960, 54, 6783 (*Alternidine*)

Shakirov, T.T. *et al.*, *Khim. Prir. Soedin.*, 1970, 6, 723; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, 6, 733 (isol)

Ishbaev, A.I. *et al.*, *Khim. Prir. Soedin.*, 1972, 8, 328; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, 8, 322 (ord)

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1973, 9, 59; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, 9, 52 (*N*-(2-Hydroxyethyl)cytisine)

Ohmiya, S. *et al.*, *Phytochemistry*, 1974, 13, 643; 1016 (*N*-Formylcytisine, *N*-Acetylcytisine)

Ruenitz, P.C. *et al.*, *J. Het. Chem.*, 1977, 14, 423 (ms)

Murakoshi, I. *et al.*, *Phytochemistry*, 1977, 16, 1460; 1982, 21, 1470; 2385 (isol, ms, pmr, synth, 3-Oxobutylcytisine, *N*-Ethylcytisine)

Murakoshi, I. *et al.*, *Phytochemistry*, 1986, 25, 2000 (*11-Oxocytisine*)

Freer, A.A. *et al.*, *Acta Cryst. C*, 1987, 43, 1119 (cryst struct)

Greinwald, R. *et al.*, *Phytochemistry*, 1990, 29, 3553 (*12-Ethoxycarbonylcytisine*)

Greinwald, R. *et al.*, *Z. Naturforsch., C*, 1990, 45, 1085 (*12-Ethoxycarbonylcytisine*)

Gazaliev, A.M. *et al.*, *Khim. Prir. Soedin.*, 1991, 27, 301; *Chem. Nat. Compd. (Engl. Transl.)*, 1991, 27, 259 (rev, bibl)

Takamatsu, S. *et al.*, *Phytochemistry*, 1991, 30, 3793 (*Cytisine N-oxide*)

Veen, G. *et al.*, *Phytochemistry*, 1992, 31, 3487 (*Dehydrocytisines*, *N*-Acetoxycytisine, *12-Methoxycarbonylcytisine*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 12886

Wang, Y.-H. *et al.*, *Chem. Pharm. Bull.*, 1999, 47, 1308-1310 (*N*-acetamidomethyl, *N*-2-oxopyrrolidinylmethyl)

O'Neill, B.T. *et al.*, *Org. Lett.*, 2000, 2, 4201-4204 (synth)

Coe, J.W. *et al.*, *Org. Lett.*, 2000, 2, 4205-4208 (synth)

Botuha, C. *et al.*, *Org. Biomol. Chem.*, 2004, 2, 1825-1826 (synth)

Danieli, B. *et al.*, *Org. Lett.*, 2004, 6, 493-496 (synth)

Honda, T. *et al.*, *J.O.C.*, 2005, 70, 499-504 (synth)


Gray, D. *et al.*, *Angew. Chem., Int. Ed.*, 2006, 45, 2419-2423 (synth)

Stead, D. *et al.*, *Tetrahedron*, 2007, 63, 1885-1897 (rev, synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CQL500; CQL750

12-Cytisineacetic acid C-939

1,5,6,8-Tetrahydro-8-oxo-1,5-methano-2H-pyrido[1,2-a][1,5]diazocine-3(4H)-acetic acid, 9CI
[72362-04-0]



C₁₃H₁₆N₂O₃ 248.281

Alkaloid from *Euchresta japonica* (Fabaceae). Cryst. (EtOH aq.). Mp 234-235° dec. [α]_D¹⁵ -200.1 (c, 0.34 in H₂O). Isol. as the Me and Et esters, which are almost certainly artifacts of methylation or ethylation.

Me ester: [72362-05-1]

C₁₄H₁₈N₂O₃ 262.308

Cryst. (hexane). Mp 107-109°. [α]_D¹⁵ -174.2 (c, 0.19 in EtOH).

Et ester:

C₁₅H₂₀N₂O₃ 276.335

Oil.

Amide: 12-Cytisineacetamide

[138867-34-2]

C₁₃H₁₇N₃O₂ 247.296

Alkaloid from *Sophora exigua* (Fabaceae). Oil. [α]_D²⁵ -204 (c, 0.17 in EtOH).


Ohmiya, S. *et al.*, *Phytochemistry*, 1979, 18, 649-650; 1985, 24, 2707-2708 (isol, ms, pmr, struct, synth)

Ohmiya, S. *et al.*, *Chem. Pharm. Bull.*, 1980, 28, 546-551 (isol)

Takamatsu, S. *et al.*, *Phytochemistry*, 1991, 30, 3793-3795 (amide)

5-(12-Cytisinylmethyl)-6-hydroxylupanine C-940

12-(6-Hydroxy-5-lupaninylmethyl)cytisine
[278617-93-9]




C₂₇H₃₈N₄O₃ 466.622

Alkaloid from *Maaackia amurensis*. Oil. [α]_D²³ -31.5 (c, 0.26 in EtOH).

Wang, Y.-H. *et al.*, *Chem. Pharm. Bull.*, 2000, 48, 641-645 (isol, synth, pmr, cmr, ms)

Cytoblastin

[137109-42-3]



Absolute configuration

C₂₈H₃₄N₄O₄ 490.601

Lactam antibiotic. Prod. by *Streptovorticillium eurodicum*. Immunomodulator. Tumour promoter. Powder. Sol. MeOH, DMSO; fairly sol. Me₂CO, CHCl₃, EtOAc; poorly sol. H₂O, hexane. Mp 177-182° dec. [α]_D²¹ -110 (c, 0.49 in MeOH). λ _{max} 202 (ε 114800); 225 (ε 91200); 283 (ε 21400); 290 (sh) (ε 16200); 300 (sh) (ε 16200) (MeOH) (Derep).


Kumagai, H. *et al.*, *J. Antibiot.*, 1991, 44, 1029 (isol, pmr, cmr, struct)

Moreno, O.A. *et al.*, *J.A.C.S.*, 1996, 118, 8180 (synth, abs config)

Moreno, O.A. *et al.*, *Bioorg. Med. Chem.*, 1998, 6, 1243-1254 (synth, abs config)

Cytochalasin V

[144306-98-9]



C₃₁H₃₅NO₇ 533.62


Metab. from *Phoma exigua* var. *heteromorpha*. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁵ +26.9 (c, 0.66 in CHCl₃). λ _{max} 296 (ε 12880) (MeOH) (Berdy).

Evidente, A. *et al.*, *Tetrahedron*, 1992, 48, 6317 (isol, uv, ir, pmr, cmr, ms, struct)

Cytochalasin B

Phomin

[14930-96-2]



C₂₉H₃₇NO₅ 479.615

Metab. of *Helminthosporium dematioides* *Phoma exigua* var. *heteromorpha*, *Phoma semeniperda* and other toxigenic fungi. Antineoplastic agent. Has antibiotic props. Cytokinesis-blocking prop. used in micronucleus assays for detecting numerical and structural chromosome changes. Inhibits lipid droplet formation. Induces morphological changes on macrophages. Active against phytopathogenic fungi. Needles (Me₂CO). Mp 218–221°. [α]_D²⁵ +83 (MeOH). Log P 2.54 (uncertain value) (calc). λ_{max} 213 (ε 26300); 219 (ε 20900); 258 (sh) (ε 490); 264 (sh) (ε 302); 267 (sh) (ε 209) (EtOH) (Derep).

► LD₅₀ (rat, ipr) 11 mg/kg. Exp. teratogen. Mutagen. RO0205000

O⁷-Ac: **7-O-Acetylcytochalasin B**

C₃₁H₃₉NO₆ 521.652

Alkaloid from *Phoma exigua* var. *heteromorpha*. Active against phytopathogenic fungi. Toxic to brine shrimp. [α]_D²⁵ -2.97 (c, 0.37 in CHCl₃).

Di-Ac:

Amorph. [α]_D +98 (EtOH).

20-Ketone: **Cytochalasin A. Dehydropho-**
min

[14110-64-6]

Metab. of *Helminthosporium dematioides* and a *Phoma* sp. Shows antibiotic props. Blocks the process of cytokinesis in animal cells without interfering with cell division. Needles (Me₂CO). [α]_D +92 (EtOH). Log P 2.8 (uncertain value) (calc). λ_{max} 213 (ε 26300); 219 (ε 20900); 258 (sh) (ε 490); 264 (sh) (ε 302); 267 (sh) (ε 209) (EtOH) (Derep).

20-Ketone, Ac: Mp 176–181°. [α]_D +18 (CHCl₃).

2'-Hydroxy: **Cytochalasin Z₄**

C₂₉H₃₇NO₆ 495.614

Prod. by *Phoma exigua* var. *heteromorpha*. Amorph. solid. λ_{max} 276 (log ε 3.13) (MeOH).

4'-Hydroxy: **Cytochalasin Z₅**

C₂₉H₃₇NO₆ 495.614

Prod. by *Phoma exigua* var. *heteromorpha*. Amorph. solid. λ_{max} 278 (log ε 3.02) (MeOH).

20-Deoxy, 19ξ-hydroxy: **Cytochalasin Z₃**

[450398-32-0]

C₂₉H₃₇NO₅ 479.615

Prod. by *Pyrenophora semeniperda*.

Phytotoxin. Amorph. solid.

A⁶-Isomer, 7-deoxy: **Cytochalasin T**

[139687-54-0]

C₂₉H₃₇NO₄ 463.616

Metab. of *Phoma exigua* var. *heteromorpha*. Mycotoxin, toxic to brine shrimp. [α]_D²⁵ -32 (c, 0.39 in CHCl₃).

A⁶-Isomer, 7-deoxy, 12-hydroxy: **Cytochalasin Z₂**

[450398-31-9]

C₂₉H₃₇NO₅ 479.615

Prod. by *Pyrenophora semeniperda*.

Amorph. solid.

A⁶-Isomer, 7,20-dideoxy, 4'-hydroxy: **Cytochalasin Z₁**

[450398-30-8]

C₂₉H₃₇NO₄ 463.616

Prod. by *Pyrenophora semeniperda*.

Amorph. solid.

Aldrich Library of Infrared Spectra, 3rd edn.,

1981, 1111B; 1111C (ir)

Aldridge, D.C. et al., *J.C.S. (C)*, 1967, 1667

(isol, ir, uv, ms, nmr, struct)

McLaughlin, G.M. et al., *Chem. Comm.*, 1970,

1398 (cryst struct)

Rothweiler, W. et al., *Helv. Chim. Acta*, 1970,

53, 696 (isol, struct, ir, uv, ms, nmr)

Turner, W.B. et al., *Biochem. J.*, 1972, **127**, 1P

(rev)

Graf, W. et al., *Helv. Chim. Acta*, 1974, **57**,

1801 (cmr, biosynth)

Masamune, S. et al., *J.A.C.S.*, 1977, **99**, 6756

(synth)

Stork, G. et al., *J.A.C.S.*, 1978, **100**, 7775

(synth, ir, pmr)

Kim, M.Y. et al., *J.O.C.*, 1981, **46**, 5383 (synth)

Pohland, A.E. et al., *Pure Appl. Chem.*, 1982,

54, 2220 (uv, ir, pmr, ms)

Stork, G. et al., *J.A.C.S.*, 1983, **105**, 5510

(synth)

Ackermann, J. et al., *Helv. Chim. Acta*, 1984,

67, 254 (synth)

Capasso, R. et al., *Phytochemistry*, 1991, **30**,

3945 (7-Acetylcytochalasin B, *Cytochalasin T*)

Ellard, S. et al., *Mutagenesis*, 1993, **8**, 317 (use)

Fenech, M. et al., *Mutat. Res.*, 1993, **285**, 35

(micronucleus assay, rev)

Obara, K. et al., *Eur. J. Pharmacol.*, 1994, **255**,

139 (pharmacol)

Evidente, A. et al., *Nat. Toxins*, 1998, **5**, 228–

233; *CA*, **129**, 199012r (activity)

Namatame, I. et al., *J. Antibiot.*, 2000, **53**, 19–

25 (activity)

Evidente, A. et al., *Phytochemistry*, 2002, **60**,

45–53 (*Cytochalasins Z*)

Evidente, A. et al., *J. Nat. Prod.*, 2003, **66**,

1540–1544 (*Cytochalasins Z₄, Z₅*)

Haidle, A.M. et al., *Proc. Natl. Acad. Sci.*

U.S.A., 2004, **101**, 12048–12053 (synth)

Cole, R.J. et al., *Handbook of Toxic Fungal*

Metabolites, Academic Press, 1981, 276; 281

Lewis, R.J. et al., *Sax's Dangerous Properties*


of Industrial Materials, 8th edn., Van

Nostrand Reinhold, 1992, CQM125

Cytochalasin C

C-944

[22144-76-9]



C₃₀H₃₇NO₆ 507.625

Metab. of *Metarhizium anisopliae*, *Bipolaris neergaardii*, *Phoma herbarum* and *Hypoxylon terricola*. Shows antibiotic and cytotoxic props. similar to Cytochalasin B, C-943. Needles (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 260–264°. [α]_D²⁸ -14.7 (c, 0.82 in dioxan). Log P 1.54 (uncertain value) (calc). λ_{max} 286 (ε 330) (MeOH) (Derep).

► HA5300500

O-De-Ac: **Deacetylcytochalasin C**

C₂₈H₃₅NO₅ 465.588

Metab. of *Metarhizium anisopliae*. [α]_D²⁵ +27.6 (c, 0.25 in EtOH).

5β,6β-Epoxyde: **Cytochalasin N†**

C₃₀H₃₇NO₇ 523.625

Metab. of *Hypoxylon terricola*. Hair-like needles (MeOH). Mp 272°. [α]_D²³ -4 (c, 0.5 in MeOH).

5β,6β-Epoxyde, O⁷-Ac:

Rosettes of needles (cyclohexane/

Me₂CO). Mp 187–192°.

19β,20β-Epoxyde: **19β,20β-Epoxycytochalasin C**

C₃₀H₃₇NO₇ 523.625

Metab. of *Xylaria hypoxylon*. [α]_D -300 (c, 0.04 in CHCl₃).

19,20-Epoxyde: **19,20-Epoxycytochalasin C**

C₃₀H₃₇NO₇ 523.625

Prod. by the fungus *Xylaria obovata*. [α]_D -6.8 (c, 0.25 in CHCl₃). The large difference in opt. rotations implies that this is not identical with the 19β,20β-epoxyde from *X. obovata*.

19,20-Epoxyde, O-de-Ac: **Deacetyl-19,20-epoxycytochalasin C**

C₂₈H₃₅NO₆ 481.588

Prod. by *Xylaria obovata*. [α]_D +11.6 (c, 0.25 in CHCl₃).

5β,6β:19β,20β-Diepoxyde: **19,20-Epoxycytochalasin N**

C₃₀H₃₇NO₈ 539.624

Metab. of *Xylaria hypoxylon*. [α]_D -55 (c, 0.04 in CHCl₃).

5β,6α-Dihydro, 7-ketone, O-de-Ac:

C₂₈H₃₅NO₅ 465.588

Metab. of *Metarhizium anisopliae*. [α]_D²⁵ -29.5 (c, 0.2 in EtOH).

Aldrich Library of Infrared Spectra, 3rd edn.,

1981, 1111D (ir)

Aldridge, D.C. et al., *J.C.S. (C)*, 1969, 923

(isol, struct, nmr)

Clark, D.A. et al., *Diss. Abstr. Int.*, **B**, 1979, **40**,

243 (synth)

Edwards, R.L. et al., *J.C.S. Perkin 1*, 1989, **57**

(*Cytochalasin N*)

Abate, D. et al., *Phytochemistry*, 1997, **44**, 1443

(19,20-Epoxycytochalasin C, Deacetyl-19,20-

epoxycytochalasin C)

Espada, A. et al., *Tetrahedron*, 1997, **53**, 6485

(epoxides)

Fujii, Y. et al., *J. Nat. Prod.*, 2000, **63**, 132–135

(*Deacetylcytochalasin C*)


Cole, R.J. et al., *Handbook of Toxic Fungal*

Metabolites, Academic Press, 1981, 294

Cytochalasin E

C-945

[36011-19-5]



C₂₈H₃₃NO₇ 495.571

Metab. of *Rosellinia necatrix* and *Aspergillus clavatus*. Also from the marine-derived *Spicaria elegans*. Antibiotic and cytotoxic agent. Fibroblast inhibitor, lipid droplet formation inhibitor, cholesterol ester synthesis inhibitor. Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O. Mp 206-208° dec. $[\alpha]_D^{25}$ -25.6 (MeOH). Log P 1.25 (uncertain value) (calc). λ_{max} 252 (ε 5370); 257 (ε 6025); 263 (ε 3890) (MeOH) (Berdy).

▶ LD₅₀ (rat, orl) 9.1 mg/kg; LD₅₀ (rat, ipr) 2.6 mg/kg. Exp. teratogen. HA5360000

4'-Methoxy-4'-Methoxycytochalasin E. **Phenochalasin B**. Antibiotic Mer-WF 1726. Mer-WF 1726 [207679-46-7]

C₂₉H₃₅NO₈ 525.597

Prod. by *Daldinia concentrica*, *Phomopsis* sp. FT-0211 and *Libertella* sp. Mer-WF 1726. Cytotoxic agent. Powder. Mp 130-132°. $[\alpha]_D^{22}$ -5.7 (c, 0.53 in MeOH). Contains a 4-methoxyphenyl group. λ_{max} 200 (ε 24500); 225 (ε 14400); 275 (ε 3200); 283 (ε 2900) (MeOH).

4'-Hydroxy, 6,7-deepoxy, 6,7-didehydro: **Phenochalasin A**

C₂₈H₃₃NO₇ 495.571

Prod. by *Phomopsis* sp. FT-0211. Powder. Mp 143-145°. $[\alpha]_D^{22}$ -4.3 (c, 0.53 in MeOH). λ_{max} 200 (ε 27800); 225 (ε 10900); 285 (ε 3000); 300 (ε 2100) (MeOH).

Aldrich Library of Infrared Spectra, 3rd ed., 1981, 1111F (ir)

Aldridge, D.C. et al., *Chem. Comm.*, 1973, 551 (struct, pmr, cmr)

Büchi, G. et al., *J.A.C.S.*, 1973, **95**, 5423 (struct, ms, ir, pmr, cryst struct)

Kajimoto, T. et al., *Chem. Pharm. Bull.*, 1989, **37**, 2212 (pmr, cmr)

Japan. Pat., 1998, 98 114 776; *CA*, **129**, 15357c (Mer-WF 1726)

Tomoda, H. et al., *J. Antibiot.*, 1999, **52**, 851-856; 857-861 (*Phenochalasin*s)

Quang, D.N. et al., *J. Nat. Prod.*, 2002, **65**, 1869-1874 (*Phenochalasin B* isol)

Liu, R. et al., *J. Nat. Prod.*, 2006, **69**, 871-875 (*Spicaria* constit, isol, pmr, cmr)


Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 266

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CQM250

Cytochalasin F

C-946

[36084-18-1]



C₂₉H₃₇NO₅ 479.615

Minor metab. of *Helminthosporium*

dematioideum, *Phoma exigua* and *Geniculosporium* sp. Exhibits algicidal props. Inhibits photosynthesis. Active against phytopathogenic fungi. Oil. $[\alpha]_D^{20}$ -52 (c, 0.5 in CHCl₃) (-33.1). Log P 2.84 (uncertain value) (calc). λ_{max} 213 (ε 26300); 219 (ε 20900); 258 (sh) (ε 490); 264 (sh) (ε 302); 267 (sh) (ε 209) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 2.6 mg/kg. RO0206000

20-Deoxy, 19 ζ -hydroxy: **Cytochalasin Z₆**

C₂₉H₃₇NO₅ 479.615

Prod. by *Phoma exigua* var. *heteromorphia*. Amorph. solid.

Aldridge, D.C. et al., *Chem. Comm.*, 1972, 148; 1973, 551 (isol, struct, pmr, cmr)

Büchi, G. et al., *J.A.C.S.*, 1973, **95**, 5423 (struct, ms, ir, pmr)

Stork, G. et al., *J.A.C.S.*, 1983, **105**, 5510 (synth)

Capasso, R. et al., *Phytochemistry*, 1991, **37**, 3945-3950 (isol)

Evidente, A. et al., *Nat. Toxins*, 1998, **5**, 228-233; *CA*, **129**, 199012r (activity)

Koenig, G.M. et al., *J. Nat. Prod.*, 1999, **62**, 155-157 (isol, pmr, cmr, activity)


Evidente, A. et al., *J. Nat. Prod.*, 2003, **66**, 1540-1544 (*Cytochalasin Z₆*)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 286

Cytochalasin G

C-947

[54874-57-6]



C₂₉H₃₄N₂O₄ 474.599

Prod. by an unidentified *Nigrosabulum* spp. and *Pseudeurotium zonatum*. Shows antibiotic and cytotoxic props. Needles (MeOH). Mp 255-257°. $[\alpha]_D^{24}$ -99 (c, 0.35 in MeOH). Log P 2.3 (uncertain value) (calc). λ_{max} 220 (log ε 4.98); 273 (log ε 4.32); 290 (log ε 4.27) (MeOH).

19 α ,20 α -Epoxy: **Cytochalasin Z**

[460352-11-8]

C₂₉H₃₂N₂O₅ 488.582

Prod. by *Pseudeurotium zonatum*. Solid. $[\alpha]_D^{20}$ -20 (c, 0.02 in MeOH). λ_{max} 220 (log ε 4.88); 273 (log ε 4.24); 290 (log ε 4.2) (MeOH).

Cameron, A.F. et al., *J.C.S. Perkin 2*, 1974, 1741-1744 (isol, ir, uv, ms, pmr, cryst struct)

Probst, A. et al., *Helv. Chim. Acta*, 1981, **64**, 2056 (cmr)

Dyke, H. et al., *J.C.S. Perkin 1*, 1989, 525 (synth)


Feng, Y. et al., *J. Nat. Prod.*, 2002, **65**, 1274-1277 (*Cytochalasin Z*)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 272

Cytochalasin K†

C-948

[79648-72-9]



C₃₂H₃₇NO₆ 531.647

Not to be confused with Cytochalasin K, C-949. Prod. by *Chalara microspora*.

Shows antibiotic and cytotoxic props.

Causes inhibition of cell movement.

Amorph. solid. $[\alpha]_D^{25}$ -177 (EtOH). λ_{max} 232 (ε 11300) (EtOH) (Derep). λ_{max} 232 (ε 11300) (EtOH) (Berdy).


Fex, T. et al., *Tet. Lett.*, 1981, 2703 (isol, ir, uv, pmr, cmr)

Cytochalasin K†

C-949

[460085-82-9]

[81657-79-6]



C₂₈H₃₃NO₇ 495.571

Related to Cytochalasin E, C-945 and not to be confused with Cytochalasin K, C-948. Prod. by *Aspergillus clavatus* MCR1181 and *Mycotypha* sp. UMF-006.

Also isol. from the marine-derived *Spicaria elegans*. Mycotoxin. Shows antibiotic and cytotoxic props. Cryst. (hexane/Me₂CO). Sol. MeOH, C₆H₆; poorly sol. H₂O, hexane. Mp 246-248°. λ_{max} 257 (MeOH) (Berdy).

$\Delta^{6,12}$ -Isomer: [460085-83-0]

C₂₈H₃₃NO₇ 495.571

Prod. by *Mycotypha* sp. UMF-006.

4'-Methoxy: **Scoparasin A**

C₂₉H₃₅NO₈ 525.597

Prod. by *Eutypella scoparia* PSU-D44. Amorph. solid. $[\alpha]_D^{29}$ -52.7 (c, 0.17 in MeOH). Conts. *p*-methoxyphenyl residue. λ_{max} 224 (log ε 3.63); 276 (log ε 2.82); 283 (log ε 2.76) (MeOH).

4'-Methoxy, 5 β ,6 β -dihydro, 6 β -hydroxy: **Scoparasin B**

C₂₉H₃₇NO₉ 543.613

Prod. by *Eutypella scoparia* PSU-D44. Amorph. solid. $[\alpha]_D^{29}$ -52.7 (c, 0.17 in MeOH). λ_{max} 224 (log ε 3.63); 276 (log

ε 2.82); 283 (log ε 2.76) (MeOH).

Steyn, P.S. *et al.*, *J.C.S. Perkin 1*, 1982, 541-544 (*isol, struct*)

Takamatsu, S. *et al.*, *J. Antibiot.*, 2002, **55**, 585-592 (*A^{6,12}-isomer*)


Pongcharoen, W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 856-858 (*Scoparasin*)

Liu, R. *et al.*, *J. Nat. Prod.*, 2006, **69**, 871-875 (*isol, pmr, cmr*)

Cytochalasin L

C-950

[79637-87-9]



C₃₂H₃₇NO₇ 547.647


Prod. by *Chalara microspora*. Shows antibiotic and cytotoxic props. Amorph. solid. [α]_D²⁵ -165 (EtOH).

Fex, T. *et al.*, *Tet. Lett.*, 1981, 2703 (*isol, ir, uv, pmr, cmr*)

Cytochalasin M

C-951

[79648-73-0]



C₃₀H₃₇NO₆ 507.625


Secondary metab. of the fungus *Chalara microspora*. Shows antibiotic and cytotoxic props. Cryst. (Me₂CO aq.). Mp 161-162°. [α]_D²⁵ +18.7 (EtOH). λ_{max} 235 nm (EtOH). λ_{max} 235 (ε 11300) (EtOH) (Derep). λ_{max} 235 (ε 11300) (EtOH) (Berdy).

Albertsson, J. *et al.*, *Acta Chem. Scand., Ser. B*, 1981, **35**, 707 (*cryst struct*)

Fex, T. *et al.*, *Tet. Lett.*, 1981, **22**, 2703 (*isol*)

Cytochalasin O†

C-952



Absolute Configuration

C₃₀H₃₉NO₇ 525.641

Metab. of the fungus *Hypoxyylon terricola*. Needles (toluene/MeOH). Mp

258-265°. [α]_D²³ -39.3 (c, 1.0 in MeOH).

6-Epimer: Cytochalasin P†

C₃₀H₃₉NO₇ 525.641

Metab. of *Hypoxyylon terricola*. Needles (Me₂CO/petrol). Mp 169-173°.

[α]_D²³ -35.8 (c, 1.0 in MeOH).

6,18-Diepimer, 17-deoxo: Cytochalasin P†. Cytochalasin P_{pho}

[108050-27-7]

C₃₀H₄₁NO₆ 511.657

Prod. by *Phomopsis* sp. Cytotoxic mycotoxin. Cryst. (CHCl₃). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 117-118°. [α]_D -116 (MeOH). λ_{max} 208 (ε 15200) (MeOH) (Berdy).

[119179-33-8]

Tomioka, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 902 (*Cytochalasin P_{pho}*)


Edwards, R.L. *et al.*, *J.C.S. Perkin 1*, 1989, 57 (*isol, ir, pmr, cmr, struct*)

Merifield, E. *et al.*, *J.C.S. Perkin 1*, 1999, 3269-3283 (*synth, config*)

Cytochalasin O†. Cytochalasin O_{pho}

C-953

[108050-26-6]



C₂₈H₃₇NO₄ 451.605

Prod. by *Phomopsis* sp. Cytotoxic mycotoxin. Needles (Me₂CO/hexane). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 187-188°. [α]_D +59.6 (MeOH). λ_{max} 206 (ε 20800) (MeOH) (Berdy).

O²¹-Ac: Cytochalasin N†. Cytochalasin N_{pho}

[108050-28-8]

C₃₀H₃₉NO₅ 493.624


Prod. by *Phomopsis* sp. Cytotoxic mycotoxin. Powder (Me₂CO). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 253-254°. [α]_D +85.4 (MeOH). λ_{max} 208 (ε 19800) (MeOH) (Berdy).

Tomioka, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 902 (*isol, struct*)

Cytochalasin Q

C-954

[119143-44-1]



C₃₀H₃₇NO₆ 507.625

Metab. of the fungus *Hypoxyylon terricola*. Fine silky needles + 1 Me₂CO (Me₂CO/petrol). Mp 145-147°. [α]_D²³ -94.5 (c, 1.0 in CHCl₃) (solvate). Isomerises almost instantaneously to Cytochalasin C, C-944 at r.t. in the presence of a trace of acid.

13β,14β-Epoxyde: Cytochalasin R

[119144-87-5]

C₃₀H₃₇NO₇ 523.625

Metab. of the fungus *Hypoxyylon terricola*. Glistening rods + 1 MeCN (MeCN). Mp 159-167°. [α]_D²³ -49.3 (c, 1.0 in MeCN) (solvate).

19β,20β-Epoxyde: 19,20-Epoxycytochalasin Q

[156098-31-6]

C₃₀H₃₇NO₇ 523.625

Metab. of *Xylaria obovata* and other *Xylaria* spp. Cytotoxic agent. Shows antimalarial activity. Sol. MeOH, CHCl₃, Mp 266-268°. [α]_D -73 (c, 1.1 in CHCl₃).

19β,20β-Epoxyde, O-de-Ac: Deacetyl-19,20-epoxycytochalasin Q. C 5507. Antibiotic C 5507

[156616-84-1]

[139441-74-0]

C₂₈H₃₅NO₆ 481.588

Metab. of *Xylaria obovata* and a bacterial eumycote sp. C5507. Cytotoxic agent, plant growth regulator. Mp 119-121°.

13β,14β:19β,20β-Diepoxide: 19,20-Epoxycytochalasin R

C₃₀H₃₇NO₈ 539.624

Metab. of *Xylaria hypoxyylon*. [α]_D -60 (c, 0.05 in CHCl₃).

18-Deoxy, 19β,20β-epoxyde: 19,20-Epoxy-18-deoxycytochalasin Q

C₃₀H₃₇NO₆ 507.625

Metab. of *Xylaria hypoxyylon*. [α]_D -360 (c, 0.05 in CHCl₃).

18-Deoxy, 19,20-epoxyde: Xylobovatin

[189371-38-8]

C₃₀H₃₇NO₆ 507.625

Isol. from the fungus *Xylaria obovata*. [α]_D²⁷ +44.5 (c, 1.3 in MeOH). This is presumably a stereoisomer of the 19β,20β-epoxyde (note very different opt. rotn.), but assigned same stereochem. in CA.

18-Deoxy, 13β,14β:19β,20β-diepoxide: 18-Deoxy-19,20-epoxycytochalasin R.

19,20-Epoxy-18-deoxycytochalasin R

C₃₀H₃₇NO₇ 523.625

Metab. of *Xylaria hypoxyylon*.

Edwards, R.L. *et al.*, *J.C.S. Perkin 1*, 1989, 57-65 (*isol, ir, pmr, cmr, struct*)

Japan. Pat., 1991, 91 251 584; CA, **116**, 127016d (*Antibiotic C 5507*)

Dagne, E. *et al.*, *Tetrahedron*, 1994, **50**, 5615-5620 (*Epoxycytochalasin Q*)

Abate, D. *et al.*, *Phytochemistry*, 1997, **44**, 1443-1448 (*Xylobovatin*)


Espada, A. *et al.*, *Tetrahedron*, 1997, **53**, 6485-6492 (*Xylaria hypoxyylon constits*)

Isaka, M. *et al.*, *Planta Med.*, 2000, **66**, 473-475 (*isol, activity*)

Cytochalasin S
[121987-40-4]

C-955

Evidente, A. *et al.*, *Nat. Toxins*, 1996, **4**, 53-57




C₂₈H₃₉NO₅ 469.62
Prod. by *Phomopsis* sp. Powder (hexane/Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 149-151°. [α]_D²⁵ -62.9 (MeOH). λ_{max} 208 (ε 13370) (MeOH). λ_{max} 208 (ε 13370) (MeOH) (Berdy).

Izawa, Y. *et al.*, *Tetrahedron*, 1989, **45**, 2323 (isol, uv, ir, pmr, cmr, ms)

Cytochalasin U†
[144279-59-4]

C-956




C₃₁H₃₇NO₇ 535.636
Metab. from *Phoma exigua* var. *heteromorpha*. [α]_D²⁵ -71.7 (c, 0.75 in CHCl₃).

Evidente, A. *et al.*, *Tetrahedron*, 1992, **48**, 6317-6324 (isol, uv, ir, pmr, cmr, ms, struct)

Cytochalasin W
[177985-19-2]


C-957



C₃₀H₃₇NO₇ 523.625
Enolised β,γ-dioxoaldehyde system. Prod. by *Phoma exigua* var. *heteromorpha*. [177985-20-5]

Cytochalasin Y
[460352-10-7]

C-958



C₂₉H₃₆N₂O₅ 492.614
Isol. from *Pseudeurotium zonatum*. Solid. [α]_D²⁰ +47.3 (c, 0.02 in MeOH). λ_{max} 220 (log ε 4.95); 273 (log ε 4.11); 290 (log ε 4.09) (MeOH).

19-Deoxy, 18-ketone: Cytochalasin X
[460352-09-4]


C₂₉H₃₄N₂O₄ 474.599
Isol. from *Pseudeurotium zonatum*. Solid. [α]_D²⁰ +13.2 (c, 0.02 in MeOH). λ_{max} 220 (log ε 4.94); 273 (log ε 4.22); 290 (log ε 4.18) (MeOH).

Feng, Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1274-1277 (isol, ir, pmr, cmr)

Cytochalasin Z₇

C-959

7,17-Dihydroxy-16,18-dimethyl-10-phenyl-22-oxa[12]cytochalasa-6(12),13,19-triene-1,21-dione



C₂₈H₃₅NO₅ 465.588
Metab. of the marine-derived fungus *Spicaria elegans*. Needles (MeOH). Mp 194-196°. [α]_D²⁵ +58.3 (c, 0.08 in MeOH). λ_{max} 236 (log ε 3.31) (MeOH).

17-Deoxy, 18α-hydroxy:

C₂₈H₃₅NO₅ 465.588
Isol. from a *Daldinia* sp. Amorph. solid. [α]_D +67.8 (c, 0.27 in MeOH).

Δ⁵-Isomer: Cytochalasin Z₈

C₂₈H₃₅NO₅ 465.588
Metab. of the marine-derived fungus *Spicaria elegans*. Needles (MeOH). Mp

215-217°. [α]_D²⁵ +68.2 (c, 0.08 in MeOH). λ_{max} 234 (log ε 3.47) (MeOH).


Δ⁵-Isomer, 17-deoxy, 18α-hydroxy: Cytochalasin Z₉

C₂₈H₃₅NO₅ 465.588
Isol. from *Spicaria elegans*. Needles (MeOH). Mp 252-254°. [α]_D²⁵ +70.4 (c, 0.1 in MeOH). λ_{max} 231 (log ε 3.37) (MeOH).

Buchanan, M.S. *et al.*, *Phytochemistry*, 1996, **41**, 821-828 (*Daldinia constii*)
Liu, R. *et al.*, *J. Nat. Prod.*, 2006, **69**, 871-875 (isol, cd, pmr, cmr)

Cytomycin
Saitomycin
[2005-98-3]

C-960



C₁₇H₂₃N₇O₅ 405.413


Nucleoside-type antibiotic. Degradn. prod. of Blasticidin S by microbes such as *Pseudomonas marginalis*, *Pseudomonas ovalis*, *Streptomyces griseochromogenes* and *Fusarium oxysporum*. Shows antitumour activity. Sol. H₂O; fairly sol. MeOH; poorly sol. Me₂CO, hexane. Mp 237-239° dec. λ_{max} 266 (ε 7450) (pH 9) (Derep). λ_{max} 274 (ε 13600) (pH 3 H₂O) (Derep).

▶ LD₅₀ (mus, ipr) 2000 mg/kg. HA5375000
Otake, N. *et al.*, *Agric. Biol. Chem.*, 1966, **30**, 132
Yamaguchi, I. *et al.*, *Agric. Biol. Chem.*, 1972, **36**, 1719 (isol)

Cytosine

C-961

4-Amino-2-(1H)-pyrimidinone, 9CI. 4-Amino-2-hydroxypyrimidine
[71-30-7]



C₄H₅N₃O 111.103

1H-Form predominates. Constit. of DNA of plants and animals. Obt. from hydrol. of nucleic acids. Plates + 5H₂O (H₂O). Mp 320-325° dec. pK_{a1} 4.6; pK_{a2} 12.16 (25°).

▶ UW7350150

4-N-Ac: [14631-20-0]
C₆H₇N₃O₂ 153.14

Cryst. (AcOH). Mp 325° (dec.). Mp >300°.

4-N-Me: *N*-Methylcytosine, 8CI. (4-Methylamino)-2(1H)-pyrimidinone [6220-47-9]
C₅H₇N₃O 125.13
Cubes (Me₂CO aq.). Mp 270°.

1H-form

1-β-D-Arabinofuranosyl: see Cytosine arabinoside, C-962

1-β-L-Ribofuranosyl: *L*-Cytidine [26524-60-7]
C₉H₁₃N₃O₅ 243.219
Cryst. (EtOH). Mp 207-208°. [α]_D²⁰ -40 (c, 1.0 in H₂O). λ_{max} 229 (sh) (ε 6900); 271 (ε 8100) (H₂O).

1-α-D-Xylofuranosyl: [77172-20-4]
C₉H₁₃N₃O₅ 243.219
Cryst. (EtOH). Mp 264-266° dec.

1-β-D-Xylofuranosyl: [3530-56-1]
C₉H₁₃N₃O₅ 243.219
Cryst. (EtOH aq.). Mp 240-242°. [α]_D²⁰ -20 (c, 0.75 in DMSO).

▶ UW7370800

1-N-β-D-Glucuronopyranosyl: **Cytosylglucuronic acid** [59862-05-4]
C₁₀H₁₃N₃O₇ 287.229
Metab. of *Streptomyces griseochromogenes*.

1-N-(3-Deoxy-β-D-galacturonopyranosyl):
C₁₀H₁₃N₃O₆ 271.229
Prod. by *Streptomyces griseochromogenes*. Solid.

1-Me: 4-Amino-1-methyl-2(1H)-pyrimidinone, 9CI. 1-Methylcytosine [1122-47-0]
C₅H₇N₃O 125.13
Cryst. (H₂O). Mp 260-265° Mp 300°.

OH-form

Me ether: 4-Amino-2-methoxypyrimidine, 8CI [3289-47-2]
C₅H₇N₃O 125.13
Cryst. (H₂O). Mp 170-171°. pK_a 5.3 (20°).

3H-form

3-Me: 6-Amino-1-methyl-2(1H)-pyrimidinone, 9CI. 3-Methylcytosine [4776-08-3]
C₅H₇N₃O 125.13
Yellow cryst. (EtOH aq.). Mp 213°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 827C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 383B (nmr)

Wheeler, H.L. et al., *Am. Chem. J.*, 1903, **29**, 429 (4-N-Ac)

Hilbert, G.E. et al., *J.A.C.S.*, 1935, **57**, 552 (synth)

Flynn, E.H. et al., *J.A.C.S.*, 1953, **75**, 5871 (synth, 1-Me)

Kenner, G.W. et al., *J.C.S.*, 1955, 855 (synth, 1-Me)

Kokko, J.P. et al., *J.A.C.S.*, 1962, **84**, 1042 (pmr)

Katritzky, A.R. et al., *J.C.S.*, 1963, 3046 (synth, 3-Me)

Barker, D.L. et al., *Acta Cryst.*, 1964, **17**, 1581 (cryst struct)

Matthews, F.S. et al., *Nature (London)*, 1964, **201**, 179 (cryst struct, 1-Me)

Rice, J.M. et al., *J.A.C.S.*, 1965, **87**, 4569 (ms)

Hill, J.A. et al., *J.C.S.*, 1965, 1515 (synth)

Sakai, T.T. et al., *J. Het. Chem.*, 1968, **5**, 849 (synth, pmr, uv, 1-Me)

David, S. et al., *Bull. Soc. Chim. Fr.*, 1969, 816 (synth)

Maehr, H. et al., *J. Het. Chem.*, 1972, **9**, 1389 (synth, pmr, uv, 3-Me)

Coletta, F. et al., *J. Magn. Reson.*, 1976, **22**, 453 (cmr, pmr)

Yamauchi, K. et al., *J.O.C.*, 1976, **41**, 3691 (synth, 1-Me)

Mandel, N.S. et al., *Acta Cryst. B*, 1977, **33**, 1079 (cryst struct)

Ward, A.D. et al., *J. Med. Chem.*, 1977, **20**, 88 (4-N-Ac)

Srikrishnan, T. et al., *Acta Cryst. B*, 1978, **34**, 1730 (cryst struct, 3-Me)

Angibeaud, P. et al., *Carbohydr. Res.*, 1980, **78**, 195 (4-N-Ac, uv, pmr)

Mathlouth, M. et al., *Carbohydr. Res.*, 1986, **146**, 1 (ir, Raman)

Gosselin, G. et al., *J. Med. Chem.*, 1986, **29**, 203 (derivis)

Guo, J. et al., *J.A.C.S.*, 1991, **113**, 5898-5899 (Cytosylglucuronic acid)

Okabe, M. et al., *J.O.C.*, 1991, **56**, 4392 (4-N-Ac, pmr)

Khutova, B.M. et al., *Khim. Geterotsikl. Soedin.*, 1991, 512 (4-N-Ac)

Gould, I.R. et al., *Spectrochim. Acta A*, 1992, **48**, 811 (ir, tautom)

Kwiatkowski, J.S. et al., *J. Phys. Chem.*, 1996, **100**, 941 (ir, struct, tautom)

Florian, J. et al., *J. Phys. Chem.*, 1996, **100**, 5578 (ir, Raman, tautom)


Zhang, Q. et al., *J. Nat. Prod.*, 1998, **61**, 648-651 (galactouronoside)

Moyroud, E. et al., *Tetrahedron*, 1999, **55**, 1277-1284 (β-L-ribofuranosyl, pmr)

Sivets, G.G. et al., *Synthesis*, 2002, 253-259 (β-L-ribofuranosyl, uv)

Cytosine arabinoside**C-962**

4-Amino-1-arabinofuranosyl-2(1H)-pyrimidinone, 9CI. Arabinosylcytosine. 4-Amino-1-arabinofuranosyl-2-oxo-1,2-dihydropyrimidine. Arabinofuranosylcytosine. Cyclocide



C₉H₁₃N₃O₅ 243.219

[116459-64-4]

Tougaard, P. et al., *Acta Cryst. B*, 1974, **30**, 86 (cryst struct)


Hruska, F.E. et al., *Can. J. Chem.*, 1982, **60**, 3026 (pmr, cmr)

Creasey, W.A. et al., *Antibiotics (N.Y.)*, 1983, **6**, 12 (rev)

Takahashi, A. et al., *Chem. Pharm. Bull.*, 1992, **40**, 1313 (isol, pmr)

Cytosporone A**C-963**

3,5-Dihydroxy-2-(1-oxooctyl)benzeneacetic acid. 3,5-Dihydroxy-2-octanoylphenylacetic acid [97902-31-3]



C₁₆H₂₂O₅ 294.347

Prod. by *Cytospora* sp., *Diaporthe* sp. and a *Phoma* sp. Cryst. Mp 130°. λ_{max} 219; 270 (MeCN).

Et ester: Cytosporone B

C₁₈H₂₆O₅ 322.4

Prod. by *Cytospora* sp., *Diaporthe* sp. and *Dothiorella* sp. λ_{max} 219; 268 (MeCN).

2',3',4',5'-Tetrahydro(E,E)-, Et ester:

Cytosporone F

[1004540-64-0]

C₁₈H₂₂O₅ 318.369

Prod. by *Paraphaeosphaeria quadrisepitata*. Pale brown oil. λ_{max} 208 (log ε 8.32); 288 (log ε 7.47) (EtOH).

2',3',4',5'-E-Tetrahydro, 2'ξ,3'ξ-epoxide,

Et ester: Cytosporone I

[1004540-67-3]

C₁₈H₂₂O₆ 334.368

Prod. by *Paraphaeosphaeria quadrisepitata*. Amorph. solid. λ_{max} 201 (log ε 5.29); 217 (log ε 4.22); 277 (log ε 4.33); 309 (log ε 5.39) (EtOH).

6'ξ-Hydroxy, Et ester: Dothiorelone B

[849758-67-4]

C₁₈H₂₆O₆ 338.4

Prod. by the mangrove fungus *Dothiorella* sp. Cytotoxic.

7'ξ-Hydroxy, Et ester: Dothiorelone A

[849758-66-3]

C₁₈H₂₆O₆ 338.4

Prod. by *Dothiorella* sp. Cytotoxic.

7'ξ-Hydroxy, 2',3',4',5'-tetrahydro(E,E)-,

Et ester: Cytosporone H

[1004540-66-2]

C₁₈H₂₂O₆ 334.368

Prod. by *Paraphaeosphaeria quadrisepitata*. Yellow oil. [α]_D²⁵ -13 (c, 0.13 in MeOH). Abs. conf. not clear from ref. λ_{max} 204 (log ε 5.23); 298 (log ε 5.09); 305 (log ε 5.55) (EtOH).

8'-Hydroxy, Et ester: Dothiorelone C

[849758-68-5]

C₁₈H₂₆O₆ 338.4

Prod. by *Dothiorella* sp. Cytotoxic.

8'-Hydroxy, 2',3',4',5'-tetrahydro(E,E)-,

Et ester: Cytosporone G

[1004540-65-1]

C₁₈H₂₂O₆ 334.368

Prod. by *Paraphaeosphaeria quadrisepitata*. Pale brown oil. λ_{max} 206 (log ε 7.44); 287 (log ε 7.58) (EtOH).

Voblikova, V.D. et al., *Chem. Nat. Compd.*

(*Engl. Transl.*), 1985, **21**, 362-365 (isol)


Brady, S.F. et al., *Org. Lett.*, 2000, **2**, 4043-4046 (isol, pmr, cmr)

Xu, Q. et al., *Acta Oceanol. Sin.*, 2004, **23**, 541-547; *CA*, **142**, 388784y (*Dothiorelones*)

Paranagama, P.A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1939-1945 (*Cytosporones F-I*)
 Singh, M.P. *et al.*, *Mar. Drugs*, 2007, **5**, 71-84
 (*isol, activity*)

Cytosaxone**C-964**

5-(Hydroxymethyl)-4-(4-methoxyphenyl)-2-oxazolidinone, 9CI
 [220736-25-4]



Absolute
 Configuration

C₁₁H₁₃NO₄ 223.228

Prod. by *Streptomyces* sp. RK95-31.
 Cytokine modulator. Immunostimulant.
 Cryst. Mp 118-121°. [α]_D²³ -71 (c, 0.1 in MeOH). The other stereoisomers have


been synthesised. λ_{\max} 225 (ϵ 19600); 277 (ϵ 2710); 284 (sh) (ϵ 2300) (MeOH).

Takeya, H. *et al.*, *J. Antibiot.*, 1998, **51**, 1126-1128 (*isol, activity*)
 Seki, M. *et al.*, *Eur. J. Org. Chem.*, 1999, 2965-2967 (*synth*)
 Takeya, H. *et al.*, *J.O.C.*, 1999, **64**, 1052-1053 (*uv, ir, cd, pmr, cmr, cryst struct*)
 Hammeršak, Z. *et al.*, *Synthesis*, 2001, 1989-1992 (*synth, stereoisomers*)
 Madhan, A. *et al.*, *Tetrahedron: Asymmetry*, 2001, **12**, 2009-2011 (*synth, pmr, cmr*)
 Carter, P.C. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 1237-1239 (*synth, sar*)
 Kumar, A.R. *et al.*, *Synth. Commun.*, 2003, **33**, 2907-2916 (*synth, ir, pmr, cmr*)
 Davies, S.G. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 1549-1553 (*synth*)
 Milicevic, S. *et al.*, *Tet. Lett.*, 2004, **45**, 955-957 (*synth*)
 Miyata, O. *et al.*, *Tetrahedron*, 2004, **60**, 3893-3914 (*synth*)

Sugiyama, S. *et al.*, *Tetrahedron: Asymmetry*, 2004, **15**, 3149-3153 (*synth, stereoisomers*)
 Asano, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2005, **69**, 145-148 (*synth*)
 Giorgio, E. *et al.*, *J.O.C.*, 2005, **70**, 6557-6563 (*ord, abs config*)
 Tokic-Vujosevic, Z. *et al.*, *Synth. Commun.*, 2005, **35**, 435-447 (*synth, epimer*)
 Smitha, G. *et al.*, *Synth. Commun.*, 2006, **36**, 1795-1800 (*epimer, synth, ir, pmr, cmr*)
 Paraskar, A.S. *et al.*, *Tetrahedron*, 2006, **62**, 5756-5762 (*synth*)
 Kim, I.S. *et al.*, *Tetrahedron*, 2006, **62**, 9349-9358 (*synth*)
 Mishra, R.K. *et al.*, *Org. Lett.*, 2007, **9**, 575-578 (*synth*)
 Narina, S.V. *et al.*, *Tet. Lett.*, 2007, **48**, 65-68 (*synth*)
 Grajewska, A. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 803-813 (*rev, synth*)
 Pham, V.-T. *et al.*, *Heterocycles*, 2008, **75**, 2817-2823 (*synth*)

Dactylamide A

[398143-15-2]



Absolute Configuration


C₂₅H₂₈N₄O₄ 448.521

Isol. from *Aplysia dactylomela*. Amorph. solid. Mp 204-210° dec. $[\alpha]_D^{20} +7$ (c, 0.68 in MeOH aq.). λ_{max} 220 (log ϵ 4.34); 272 (log ϵ 3.56); 286 (log ϵ 3.53); 290 (log ϵ 3.45) (MeOH).

Appleton, D.R. *et al.*, *Tetrahedron*, 2001, **57**, 10181-10189 (*isol, pmr, cmr, uv*)

Dactylamide B

[398143-16-3]


C₂₄H₂₇N₄O₂⁺ 403.503

Isol. from *Aplysia dactylomela*. Pale yellow solid. Mp 160-163° dec. $[\alpha]_D^{20} +160$ (c, 0.4 in MeOH). Counterion not specified. λ_{max} 219 (log ϵ 4.44); 282 (log ϵ 3.85); 292 (log ϵ 3.83); 313 (log ϵ 3.85) (MeOH).

Appleton, D.R. *et al.*, *Tetrahedron*, 2001, **57**, 10181-10189 (*isol, pmr, cmr, uv*)

Dactylicapnosine

[168434-29-5]



Relative Configuration


C₄₂H₃₈N₂O₁₄ 794.767

Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Mp 135-136°. $[\alpha]_D^{20} 0$ (c, 0.3 in MeOH).

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 299 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Dactylicapnosinine

[168434-28-4]



Relative Configuration


C₂₁H₁₉NO₇ 397.384

Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Needles (CHCl₃/EtOAc, 2:1). Mp 184.5-186°. $[\alpha]_D^{20} 0$ (c, 0.1 in MeOH).

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 299 (*isol, uv, ir, pmr, cmr, ms, struct*)

Dactylidine

[168434-27-3]


C₄₀H₃₀N₂O₈²⁺ 666.686

Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Orange powder (as dichloride). Mp 300° (dichloride).

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 299 (*isol, uv, ir, pmr, cmr, struct*)

Dactylyne

[168434-26-2]


C₂₂H₂₅N₂O₅⁺ 397.45

Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Needles (MeOH) (as chloride). Mp 110° dec. (chloride). $[\alpha]_D^{20} 0$ (c, 1.0 in MeOH) (chloride).

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 299 (*isol, uv, ir, pmr, cmr, ms, struct*)

Daechuine S3

[123089-20-3]

C₃₄H₅₃N₅O₆ 627.823

Alkaloid from the Daechu tree (*Zizyphus jujuba* var. *inermis*) and from *Paliurus*


ramossissimus. Mp 192-194°. $[\alpha]_D^{30} -454$ (c, 0.76 in MeOH).

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443-448

Lee, S.-S. *et al.*, *Phytochemistry*, 2001, **58**, 1271-1276 (*isol, cd, uv, pmr, cmr, ms*)

Daechuine S7

[123089-22-5]


C₂₈H₄₂N₄O₅ 514.664

Alkaloid from the stem bark of the Daechu tree (*Zizyphus jujuba* var. *inermis*) (Rhamnaceae). Mp 158°. $[\alpha]_D -648.3$.

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443

Daechuine S8-1

[123089-23-6]


C₃₄H₅₃N₅O₆ 627.823

Alkaloid from the stem bark of the Daechu tree (*Zizyphus jujuba* var. *inermis*) (Rhamnaceae). Mp 185-188°. $[\alpha]_D -218.2$.

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443

Dakaramine

[173075-47-3]


C₁₅H₂₄I₂N₂O 502.176

Metab. from the Senegalese sponge *Ptilocaulis spiculifer*.


Diop, M. *et al.*, *J. Nat. Prod.*, 1996, **59**, 271-272 (*isol, uv, ir, pmr, cmr, struct*)

Dalomycin T

[879920-36-2]

C₂₈H₂₉NO₉ 523.538Metab. of *Streptomyces venezuelae* ISP 5230.Syvitski, R.T. *et al.*, *Org. Lett.*, 2006, **8**, 697-700 (*isol*, *pmr*, *cmr*)**Dambullin**


[160896-51-5]

C₁₇H₂₃NO₄S 337.439Alkaloid from leaves of *Glycosmis angustifolia* (Rutaceae). Mp 145-147°.**N-Me: Methyldambullin**

[160896-52-6]

C₁₈H₂₅NO₄S 351.466Alkaloid from leaves of *Glycosmis angustifolia*. Exhibits moderate antifungal activity. Mp 70-73°.**12-Hydroxy: Sakambullin**C₁₇H₂₃NO₅S 353.438Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 111-113°. λ_{max} 224 (sh); 275 (MeOH).**12-Methoxy: O-Methylsakambullin.***Methoxydambullin*C₁₈H₂₅NO₅S 367.465Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 136-138°. λ_{max} 227 (sh); 275 (MeOH).Greger, H. *et al.*, *Phytochemistry*, 1994, **37**, 1305-1310 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)
Hinterberger, S. *et al.*, *Tetrahedron*, 1998, **54**, 487-496 (*synth*)Hofer, O. *et al.*, *Phytochemistry*, 2000, **54**, 207-213 (*Sakambullin*)**Damirone A**

1,3,4,5-Tetrahydro-1,5-dimethylpyrrolo[4,3,2-de]quinoline-7,8-dione, 9CI [138683-66-6]

C₁₂H₁₂N₂O₂ 216.239Alkaloid from the sponges *Damiria* sp. and *Zyzzya* sp. Purple solid. Mp 240-242° dec. λ_{max} 242 (ε 12600); 347 (ε 10200); 518 (ε 912) (MeOH) (*Derep*).

D-11

λ_{max} 214 (ε 9000); 245 (ε 21000); 347 (ε 7200); 516 (ε 500) (MeOH) (*Derep*).**N¹-De-Me: Damirone B**

[138683-67-7]

C₁₁H₁₀N₂O₂ 202.212Alkaloid from the sponges *Damiria* sp. and *Zyzzya fuliginosa*. Purple solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 250°. λ_{max} 242 (ε 7080); 350 (ε 4470); 498 (ε 257) (MeOH) (*Derep*). λ_{max} 242 (ε 29500); 347 (ε 19500) (MeOH) (*Berdy*). λ_{max} 242 (ε 19000); 347 (ε 19300) (MeOH/NaOH) (*Berdy*). λ_{max} 243; 348; 492 (EtOH) (*Berdy*).**N⁵-De-Me: 6-Dechlorobatzelline C**

[138683-68-8]

C₁₁H₁₀N₂O₂ 202.212Isol. from *Zyzzya fuliginosa*.**N¹,N⁵-Di-de-Me: Damirone C**


[157669-64-2]

C₁₀H₈N₂O₂ 188.185Alkaloid from the sponge *Zyzzya fuliginosa*. Red-brown solid.**N¹,N⁵-Di-de-Me, N¹-β-D-ribofuranosyl:****N¹-β-D-Ribofuranosyldamirone C**C₁₅H₁₆N₂O₆ 320.301Alkaloid from the sponge *Strongylo-desma aliwaliensis*. Red amorph. solid. [α]_D²⁵ +5 (c, 0.06 in MeOH). λ_{max} 246 (ε 14100); 334 (ε 7990); 526 (ε 650) (MeOH).**6-Bromo, N⁵-de-Me: Makaluvone**

[146555-84-2]

C₁₁H₉BrN₂O₂ 281.108Isol. from the sponge *Zyzzya fuliginosa*. Grey solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 246 (ε 9800); 330 (ε 6300) (MeOH/NaOH) (*Derep*). λ_{max} 246 (ε 10100); 330 (ε 5500) (MeOH) (*Derep*).**6-Bromo, N¹,N⁵-di-de-Me: 6-Bromo-1,3,4,5-tetrahydropyrrolo[4,3,2-de]quinoline-7,8-dione, 9CI. Makaluvamine O**C₁₀H₇BrN₂O₂ 267.082Alkaloid from the sponge *Smenospongia* sp. Purple solid. Mp 282-283° dec. λ_{max} 222 (log ε 3.81); 244 (log ε 3.28); 328 (log ε 2.94); 362 (sh) (log ε 2.86); 542 (log ε 2.06) (MeOH). λ_{max} 244 (log ε 4.01); 337 (log ε 3.75) (MeOH).Stierle, D.B. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1131 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *cmr*)Baumann, C. *et al.*, *Angew. Chem., Int. Ed.*, 1993, **105**, 1087 (*synth*)Radisky, D.C. *et al.*, *J.A.C.S.*, 1993, **115**, 1632 (*isol*, *Damirone B*)Sadanandan, E.V. *et al.*, *Tet. Lett.*, 1993, **34**, 2405 (*synth*)Yamada, F. *et al.*, *Heterocycles*, 1995, **41**, 1905 (*synth*)Schmidt, E.W. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1861 (*Damirone C*)Roue, N. *et al.*, *J. Pharm. Belg.*, 1995, **50**, 94 (*synth*)Peat, A.J. *et al.*, *J.A.C.S.*, 1996, **118**, 1028 (*synth*)Roberts, D. *et al.*, *J.O.C.*, 1997, **62**, 568 (*synth*)
Hu, J.F. *et al.*, *J. Nat. Prod.*, 2002, **65**, 476-480 (*Makaluvamine O*)Tasdemir, D. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 914-922 (*Makaluvamine O*)Keyzers, R.A. *et al.*, *Tet. Lett.*, 2004, **45**, 9415-9418 (*N-Ribofuranosyldamirone C*)Dijoux, M.-G. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 6035-6044 (*isol*, *cmr*, *activity*)**Danaidal**


2,3-Dihydro-1H-pyrrolizine-7-carboxaldehyde, 9CI. 1-Formyl-6,7-dihydro-5H-pyrrolizine [27628-46-2]

C₈H₉NO 135.165Isol. from hair pencil and wing scent organ secretions of the butterflies *Danaus affinis affinis*, *Utetheisa lotrix*, *Phragmatobia fuliginosa* and *Pyrrharcia isabella*. Cryst. solid. Mp 59-60°.

▶ UY7760000

7R-Hydroxy: Hydroxydanaidal. 2,3-Dihydro-1-hydroxy-1H-pyrrolizine-7-carboxaldehyde, 9CI. 1-Formyl-7-hydroxy-6,7-dihydro-5H-pyrrolizine [28379-58-0]C₈H₉NO₂ 151.165Found in butterflies *Danaus hamatus hamatus*, *Euploea tulliola tulliola* and *Euploea sylvester sylvester* and moths. [α]_D²⁵ -140. Prod. in the butterfly from Heliotrine, H-95 with inversion of stereochem.**7-Oxo: 2,3-Dihydro-1-oxo-1H-pyrrolizine-7-carboxaldehyde. 1-Formyl-6,7-dihydro-7-oxo-5H-pyrrolizine. 7-Oxodanaidal**Alkaloid from the roots of *Ligularia cymbulifera*. Amorph. solid. [α]_D²¹ -7 (c, 0.7 in CHCl₃). Opt. rotn. unaccounted for. λ_{max} 270 (log ε 1.28); 306 (log ε 1.6) (CHCl₃).Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1869 (*synth*)Edgar, J.A. *et al.*, *Experientia*, 1971, **27**, 761; 1972, **28**, 627 (*isol*)Pizzorno, M.T. *et al.*, *Chem. Ind. (London)*, 1978, 349 (*synth*)Schneider, D. *et al.*, *Science (Washington, D.C.)*, 1982, **215**, 1264 (*isol*)Komai, H. *et al.*, *Agric. Biol. Chem.*, 1983, **47**, 157 (*isol*)Bell, T.W. *et al.*, *Experientia*, 1984, **40**, 713 (*abs config*, *Hydroxydanaidal*)Röder, E. *et al.*, *Annalen*, 1986, 1645 (*synth*, *ir*, *ms*, *pmr*, *cmr*)Krasnoff, S.B. *et al.*, *J. Chem. Ecol.*, 1987, **13**, 807Liu, C.-M. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 308-316 (*7-Oxodanaidal*)**Dankastatin A**

[1011269-08-1]



Absolute Configuration

C₂₄H₃₅Cl₂NO₃ 488.45Prod. by *Gymnascella dankaliensis* OUPS-N134. Cytotoxic. Powder. Mp 169-171°. [α]_D²² +114.4 (c, 0.18 in CHCl₃). λ_{max} 266 (log ε 4.45) (EtOH).4,8,9-Triepimer, demethoxy: **Dankastatin B**


[1011269-09-2]
 $C_{23}H_{33}Cl_2NO_4$ 458.423

Prod. by *Gymnascella dankaliensis*
 OUPS-N134. Powder. Mp 90-92.5°.
 $[\alpha]_D^{22}$ -157.4 (c, 0.18 in $CHCl_3$). λ_{max}
 265 (log ϵ 4.45) (EtOH).

Amagata, T. *et al.*, *J. Nat. Prod.*, 2008, **71**, 340-345 (isol, cd, pmr, cmr, ms)

Danomycin**D-16**

[11005-96-2]



Danomycin A, R = $C_{12}H_{20}NO_7(OMe)(OOCNH_2)$
 Danomycin B, R = $C_{12}H_{20}NO_7(OMe)(OH)$

Fe complex antibiotic containing a disaccharide of unknown struct. Contains two components. Prod. by *Streptomyces albaduncas*. Active against gram-positive bacteria and weakly against gram-negative bacteria and mycobacteria. Side-phore. Reddish-brown cryst. Sol. H_2O , phenol; fairly sol. MeOH, EtOH- H_2O ; poorly sol. butanol, hexane. Mp 135-138° dec. λ_{max} 270 (E1%/1cm 48); 325 (E1%/1cm 12.6); 430 (E1%/1cm 15.6) (H_2O) (Berdy). λ_{max} 325 (E1%/1cm 18); 450 (E1%/1cm 13) (HCl) (Berdy). λ_{max} 325 (E1%/1cm 20); 420 (E1%/1cm 25) (NaOH) (Berdy).

► LD₅₀ (mus, ivn) 3250 mg/kg. HB4777000

Danomycin B

$C_{42}H_{70}FeN_4O_{20}$ 1006.877
 Sol. H_2O , MeOH. λ_{max} 420 (E1%/1cm 20) (H_2O) (Berdy).

Des-Ferri deriv.:

Beige powder. Mp 122-125° dec.

Aglycone: Danoxamine

[105031-30-9]

$C_{27}H_{46}FeN_5O_{11}$ 672.534

Red-brown powder. Mp ca. ° 150 (dec.).

[104381-50-2, 104673-04-3, 105031-29-6]

Tsukiura, H. *et al.*, *J. Antibiot., Ser. A*, 1964, **17**, 39 (isol, props)


Japan. Pat., 1965, 65 13 796; *CA*, **63**, 17092 (isol, props)

Huber, P. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 236 (struct)

Roosenberg, J.M. *et al.*, *J.O.C.*, 2000, **65**, 4833-4838 (*Danoxamine, synth*)

Danuphylline**D-17**

[213478-78-5]



Absolute Configuration

$C_{24}H_{26}N_2O_8$ 470.478

Alkaloid from the leaves of *Kopsia dasyrachis*. Amorph. $[\alpha]_D$ -30 (c, 0.07 in $CHCl_3$). λ_{max} 214 (log ϵ 3.95); 225 (log ϵ 4.13); 248 (log ϵ 3.66); 283 (log ϵ 3.03); 293 (log ϵ 2.99) (no solvent reported).

De(methylenedioxy): Danuphylline A

$C_{23}H_{26}N_2O_6$ 426.468

Alkaloid from the leaves of *Kopsia officinalis*. Amorph. powder. $[\alpha]_D^{25}$ +28.9 (c, 0.36 in $CHCl_3$). λ_{max} 243; 279; 286 ($CHCl_3$).

Kam, T.S. *et al.*, *Tet. Lett.*, 1998, **39**, 5823-5826 (isol, uv, pmr, cmr)

Suzuki, K. *et al.*, *J. Antibiot.*, 1999, **52**, 460-465 (activity)


Kam, T.S. *et al.*, *Phytochemistry*, 1999, **52**, 959-963 (cryst struct)

Kam, T.S. *et al.*, *Tetrahedron*, 1999, **55**, 1457-1468 (synth)

Zhou, H. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 515-519 (*Danuphylline A*)

Daphcalyic acid**D-18**

[666174-79-4]



$C_{22}H_{31}NO_3$ 357.492

Alkaloid from the seeds of *Daphniophyllum calycinum*. Cryst. (MeOH). Mp 124-125°. $[\alpha]_D^{22}$ -33 (c, 0.3 in MeOH).

6'-Ester with Geniposidic acid: Daphmacalycinosidine A

[666174-77-2]


$C_{38}H_{51}NO_{12}$ 713.82

Alkaloid from the seeds of *Daphniophyllum calycinum*. Cryst. (MeOH). Mp 134-136°. $[\alpha]_D^{22}$ -16 (c, 0.6 in MeOH).

El Bitar, H. *et al.*, *Tet. Lett.*, 2004, **45**, 515-518; 2027-2028 (isol, pmr, cmr)

Daphcalyine**D-19**

[496836-44-3]



Relative Configuration


$C_{23}H_{33}NO_3$ 371.519

Alkaloid from the stem bark of *Daphniophyllum calycinum*. Cytotoxic. Amorph. solid. $[\alpha]_D$ -73 (c, 0.8 in MeOH). λ_{max} 210 (ϵ 2700) (MeOH).

Jossang, A. *et al.*, *J.O.C.*, 2003, **68**, 300-304 (isol, pmr, cmr)

Daphcalycinosidine B**D-20**

[666174-78-3]



$C_{40}H_{57}NO_{13}$ 759.889

Alkaloid from the seeds of *Daphniophyllum calycinum*. Cryst. (MeOH). Mp 201-202°. $[\alpha]_D^{22}$ +6 (c, 0.3 in MeOH).

Demethoxy, 2,3-didehydro: Daphmacropodosidine B

[944454-27-7]

$C_{39}H_{53}NO_{12}$ 727.847

Alkaloid from the fruit of *Daphniophyllum macropodum*. Amorph. powder (MeOH). Mp 196-200°. λ_{max} 309 (log ϵ 2.52); 378 (log ϵ 2.26) (MeOH).

3'-Ester isomer: Daphmacropodosidine A

[944454-26-6]

$C_{40}H_{57}NO_{13}$ 759.889


Alkaloid from the fruit of *Daphniophyllum macropodum*. Cryst. (MeOH). Mp 176-178°. $[\alpha]_D^{29}$ -43.5 (c, 0.84 in MeOH). λ_{max} 307 (log ϵ 2.73); 374 (log ϵ 2.42) (MeOH).

El Bitar, H. *et al.*, *Tet. Lett.*, 2004, **45**, 515-518; 2027-2028 (*Daphcalycinosidine B*)

Kong, N.-C. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 972-976 (*Daphmacropodosidines A,B*)

Daphlongeramine A**D-21**

[937719-93-2]




$C_{27}H_{35}NO_5$ 453.577

Alkaloid from the fruit of *Daphniophyllum longeracemosum*. Amorph. solid. $[\alpha]_D^{25}$ -39.4 (c, 0.14 in $CHCl_3$). λ_{max} 241 (log ϵ 3.56) (no solvent reported).


Li, C. *et al.*, *Tet. Lett.*, 2007, **48**, 2737-2740 (isol, pmr, cmr)

Daphlongeranine A


[944453-93-4]

Relative
ConfigurationC₂₄H₃₁NO₅ 413.513Alkaloid from the fruit of *Daphniphyllum longercemosum*. Light yellow cryst. (petrol/Me₂CO). Mp 227-229°. [α]_D²² +142.7 (c, 0.16 in CHCl₃).Li, C.-S. *et al.*, *Org. Lett.*, 2007, 9, 2509-2512 (isol, pmr, cmr, cryst struct)**Daphlongeranine B**

[944453-94-5]

Relative
ConfigurationC₂₂H₂₉NO₄ 371.475Alkaloid from the fruit of *Daphniphyllum longercemosum*. Needles (petrol/Me₂CO). Mp 241-243°. [α]_D¹⁷ +47.2 (c, 0.41 in CHCl₃).Li, C.-S. *et al.*, *Org. Lett.*, 2007, 9, 2509-2512 (isol, pmr, cmr)**Daphmacrine**

[19775-48-5]

Absolute
ConfigurationC₃₂H₄₉NO₄ 511.743Alkaloid from *Daphniphyllum macropodium* (Daphniphyllaceae). Noncryst.*Hydrobromide*:Cryst. (CHCl₃/Me₂CO). Mp 300°. [α]_D +30.1 (c, 1.79 in MeOH).*Lactol*: **Daphmacropodine**

[39729-21-0]

C₃₂H₅₁NO₄ 513.759Alkaloid from *Daphniphyllum macropodium* (Daphniphyllaceae). Mp 214-215°. [α]_D +4.9 (c, 1.11 in CHCl₃). The lactone carbonyl of Daphmacrine is reduced to OH (stereochem. undetermined).*Lactol*; *hydrobromide*:Cryst. (Me₂CO). Mp 215-218°.*7S-Hydroxy, lactol*: **Calyciphylline M**C₃₂H₅₁NO₅ 529.759

D-22

Alkaloid from *Daphniphyllum calycinum*. Amorph. solid. [α]_D¹⁷ +1.1 (c, 1 in CHCl₃).Nakano, T. *et al.*, *Chem. Comm.*, 1968, 600


(isol, ir, pmr, ms, cryst struct)

Gibbons, C.S. *et al.*, *J.C.S. (B)*, 1969, 840



(cryst struct)

Yang, S.-P. *et al.*, *Helv. Chim. Acta*, 2006, 89, 2783-2788 (*Daphmacropodine*)Mu, S.-Z. *et al.*, *Chem. Biodiversity*, 2007, 4, 129-138 (*Daphmacropodine*)Saito, S. *et al.*, *Tetrahedron*, 2008, 64, 1901-1908 (*Calyciphylline M*)**Daphmanidine A**

[462117-89-1]

Absolute
ConfigurationC₂₅H₃₃NO₅ 427.539Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Cytotoxic. Amorph. solid. [α]_D -62 (c, 0.5 in MeOH). λ_{\max} 283 (ε 2200) (MeOH).*14,15-Didehydro*: **Daphmanidine E**C₂₅H₃₁NO₅ 425.524Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Amorph. solid. [α]_D²⁰ +11 (c, 0.5 in MeOH). λ_{\max} 296 (ε 8200) (MeOH).*14,15-Didehydro, 9β,10β-epoxide*: **Daphmanidine F**C₂₅H₃₁NO₆ 441.523Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Amorph. solid. [α]_D²⁰ +25 (c, 1 in MeOH). λ_{\max} 235 (ε 6200); 291 (ε 4200) (MeOH).Kobayashi, J. *et al.*, *J.O.C.*, 2002, 67, 6546-6549 (*Daphmanidine A*)Morita, H. *et al.*, *J. Nat. Prod.*, 2006, 69, 418-420 (*Daphmanidines E,F*)**Daphmanidine B**

[462117-90-4]

Absolute
ConfigurationC₂₅H₃₅NO₆ 445.555Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Cytotoxic. Amorph. solid. [α]_D -15 (c, 0.2 in MeOH).Kobayashi, J. *et al.*, *J.O.C.*, 2002, 67, 6546-6549 (isol, pmr, cmr)**Daphmanidine C**Relative
Configuration

D-25

C₂₅H₃₃NO₇ 459.538Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Amorph. solid. [α]_D -15 (c, 0.1 in MeOH). λ_{\max} 250 (ε 12000) (MeOH).*15-Hydroxy, 2-ketone*: **Daphmanidine D**C₂₅H₃₁NO₈ 473.522Alkaloid from the leaves of *Daphniphyllum teijsmanii*. Amorph. solid. [α]_D -18 (c, 0.3 in MeOH). λ_{\max} 246 (ε 13000) (MeOH).Morita, H. *et al.*, *Org. Lett.*, 2005, 7, 459-462 (isol, pmr, cmr)**Daphnarcine**


[1356-42-9]

C₁₆H₁₇NO₄ 287.315Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Narcissus* sp. (Amaryllidaceae). Prisms (Me₂CO/MeOH). Mp 258-260° dec. [α]_D²⁵ +40 (c, 0.1 in DMF). *Picrate*: Mp 246° dec.Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, 90, 2197-2202 (isol)

D-28

Daphnezomine A


[247078-42-8]

Absolute
ConfigurationC₂₂H₃₅NO₃ 361.523Zwitterionic. Alkaloid from *Daphniphyllum humile*. Needles. Mp 218-220°. [α]_D -30 (c, 0.1 in CHCl₃/MeOH).*Me ester*: **Daphnezomine B**

[247078-43-9]

C₂₃H₃₇NO₃ 375.55Alkaloid from *Daphniphyllum humile*. Powder. [α]_D -31 (c, 0.3 in CHCl₃).Morita, H. *et al.*, *J.O.C.*, 1999, 64, 7208-7212 (isol, ir, pmr, cmr, cd, cryst struct)**Daphnezomine C**

[251111-54-3]

C₃₀H₄₅NO₄ 483.69Alkaloid from *Daphniphyllum humile*.

D-30


Solid. $[\alpha]_D^{24}$ -94 (c, 0.3 in CHCl_3). λ_{max} 260 (ϵ 4500) (MeOH).

Morita, H. *et al.*, *Tetrahedron*, 1999, **55**, 12549-12556

Daphnezomine D

D-31

[251111-55-4]



$\text{C}_{32}\text{H}_{49}\text{NO}_5$ 527.743

Alkaloid from *Daphniphyllum humile*.


Solid. $[\alpha]_D^{24}$ -151 (c, 0.4 in CHCl_3). λ_{max} 260 (ϵ 7000) (MeOH).

Morita, H. *et al.*, *Tetrahedron*, 1999, **55**, 12549-12556

Daphnezomine F

D-32

[276255-94-8]



$\text{C}_{27}\text{H}_{35}\text{NO}_8$ 501.575

Alkaloid from *Daphniphyllum humile*.

Solid. $[\alpha]_D$ -29 (c, 0.5 in CHCl_3). λ_{max} 215 (ϵ 3700); 260 (ϵ 700) (MeOH).

7-Deoxo, 6,7-didehydro: **Daphnezomine G**

[276255-96-0]

$\text{C}_{27}\text{H}_{35}\text{NO}_7$ 485.576

Alkaloid from *Daphniphyllum humile*.


Solid. $[\alpha]_D$ -93 (c, 0.4 in CHCl_3). λ_{max} 215 (ϵ 5600); 255 (ϵ 2000) (MeOH).

Morita, H. *et al.*, *J.O.C.*, 2000, **65**, 3432-3442 (synth, config, pmr, cmr, ms)

Daphnezomine K

D-33

[38966-00-6]



Absolute Configuration

$\text{C}_{23}\text{H}_{33}\text{NO}_5$ 403.517

Alkaloid from *Daphniphyllum humile*.

Amorph. solid. $[\alpha]_D^{23}$ -15 (c, 0.3 in MeOH).

4,21-Di-Ac: **Yuzurimine**. **Macrodaphnidine**

[17819-76-0]

$\text{C}_{27}\text{H}_{37}\text{NO}_7$ 487.592

Alkaloid from the leaves and bark of

Daphniphyllum macropodum and the fruits of *Daphniphyllum teijsmanni* (Daphniphyllaceae). Weak CNS depressant. Cryst. ($\text{Me}_2\text{CO}/\text{Et}_2\text{O}$). Mp 150-152°. $[\alpha]_D$ +8.3 (c, 1.3 in CHCl_3). pK_a 7.4 (80% MCS).

4,21-Di-Ac, hydrochloride:

Cryst. ($\text{Me}_2\text{CO}/\text{MeOH}$). Mp 231-234°.

14,15-Didehydro, 4,21-di-Ac: **Daphtenidine D**

$\text{C}_{27}\text{H}_{35}\text{NO}_7$ 485.576

Alkaloid from the leaves of *Daphniphyllum teijsmanni*. Amorph. solid.

$[\alpha]_D^{23}$ +36 (c, 1 in MeOH). λ_{max} 301 (ϵ 11060) (MeOH).

17-Oxo, 4,21-di-Ac: **17-Oxoyuzurimine**

[949171-73-7]

$\text{C}_{27}\text{H}_{35}\text{NO}_8$ 501.575

Alkaloid from the leaves of *Daphniphyllum macropodum*. Oil. $[\alpha]_D^{23}$ +14 (c, 0.26 in CHCl_3). λ_{max} 248 (log ϵ 3.39) (MeOH).

Parent acid, 4,21-di-Ac: **Macropodumine H**

[949172-78-5]

$\text{C}_{26}\text{H}_{35}\text{NO}_7$ 473.565

Alkaloid from the leaves of *Daphniphyllum macropodum*. Amorph. powder. $[\alpha]_D^{23}$ +14.5 (c, 0.47 in CHCl_3).

1-Deoxy, 4,21-di-Ac: **Deoxyyuzurimine**

[38966-09-5]

$\text{C}_{27}\text{H}_{37}\text{NO}_6$ 471.592

Alkaloid from the leaves of *Daphniphyllum humile* (Daphniphyllaceae). Plates (cyclohexane). Mp 132-134°.

1-Deoxy, 4,21-di-Ac, N-oxide: **Macrodaphnine**

[24148-90-1]

$\text{C}_{27}\text{H}_{37}\text{NO}_7$ 487.592

Alkaloid from *Daphniphyllum macropodum* (Daphniphyllaceae). Cryst. ($\text{CHCl}_3/\text{Et}_2\text{O}$). Mp 180-181.5°.

1-Deoxy, 1(N)-dehydro, 21-Ac: **Daphnezomine J**

[276870-44-1]

$\text{C}_{25}\text{H}_{34}\text{NO}_5^{\oplus}$ 428.547

Quaternary alkaloid from *Daphniphyllum humile*. Amorph. solid. $[\alpha]_D^{23}$ -20 (c, 0.9 in MeOH). Counterion not specified.

4-Deoxy, 21-Ac: **Yuzurimine E**. **Daphniglaucine J**

[752252-76-9]

$\text{C}_{25}\text{H}_{35}\text{NO}_5$ 429.555

Alkaloid from the seeds of *Daphniphyllum calycinum* and the leaves of *Daphniphyllum glaucescens*. Amorph. solid. $[\alpha]_D^{22}$ -33 (c, 0.3 in MeOH). $[\alpha]_D^{24}$ -17 (c, 0.4 in MeOH).

21-Deoxy: **Macrodaphniphyllamine**. **Deacetylyuzurimine A**

[17807-72-6]

$\text{C}_{23}\text{H}_{33}\text{NO}_4$ 387.518

Alkaloid from *Daphniphyllum macropodum* and *Daphniphyllum yunnanense*. Cryst. ($\text{CHCl}_3/\text{hexane}$). Mp 152-153°.

$[\alpha]_D$ -51.7 (c, 1.25 in CHCl_3).

21-Deoxy, hydrochloride:

Plates ($\text{MeOH}/\text{Et}_2\text{O}$). Mp 251-254° (sealed tube).

21-Deoxy, 4-Ac: **Yuzurimine A**

[17232-64-3]

$\text{C}_{25}\text{H}_{35}\text{NO}_5$ 429.555

Minor alkaloid from the bark and leaves of *Daphniphyllum macropodum* (Daphniphyllaceae). Needles ($\text{MeOH}/\text{Et}_2\text{O}$) (as hydrochloride). Mp 249-242° (sealed tube) (hydrochloride). Lacks the primary acetoxy group.

1,4-Dideoxy: **Yuzurimine B**

[17232-65-4]

$\text{C}_{23}\text{H}_{33}\text{NO}_3$ 371.519

Alkaloid from the leaves and bark of *Daphniphyllum macropodum* and the fruits of *Daphniphyllum teijsmanni* (Daphniphyllaceae). Cryst. (MeOH) (as hydrochloride). Mp 282-284.5° (sealed tube) (hydrochloride). λ_{max} 210 (ϵ 7230) (EtOH).

1,4-Dideoxy, 21-Ac: **Macrodaphniphyllidine**

[26548-56-1]

$\text{C}_{25}\text{H}_{35}\text{NO}_4$ 413.556

Alkaloid from *Daphniphyllum macropodum* (Daphniphyllaceae). Cryst. ($\text{Me}_2\text{CO}/\text{MeOH}$) (as hydrobromide).

Mp 305-306° (hydrobromide). $[\alpha]_D$ +3.9 (c, 1.11 in MeOH) (hydrobromide).

1,4-Dideoxy, 21-Ac, N-oxide: **Daphniglaucine K**

[753450-67-8]

$\text{C}_{25}\text{H}_{35}\text{NO}_5$ 429.555

Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. $[\alpha]_D^{25}$ -13 (c, 0.7 in MeOH).

1,4-Dideoxy, parent acid: **Yuzurimic acid B**

[752252-77-0]

$\text{C}_{22}\text{H}_{31}\text{NO}_3$ 357.492

Alkaloid from the seeds of *Daphniphyllum calycinum*. Cryst. (MeOH). Mp 253-255°. $[\alpha]_D^{22}$ +11 (c, 0.3 in MeOH).

1,4-Dideoxy, 14,15-didehydro, N-oxide: **Desacetyldaphnijsmine**. **Deacetyldaphnijsmine**

[55855-03-3]

$\text{C}_{23}\text{H}_{31}\text{NO}_4$ 385.502

Minor alkaloid from the fruits of *Daphniphyllum teijsmanni* (Daphniphyllaceae). Cryst. (hexane/ EtOAc). Mp ca.° 200 dec.

1,4-Dideoxy, 14,15-didehydro, 21-Ac, N-oxide: **Daphnijsmine**

[55855-02-2]

$\text{C}_{25}\text{H}_{33}\text{NO}_5$ 427.539

Minor alkaloid from the fruits of *Daphniphyllum teijsmanni* (Daphniphyllaceae). Cryst. (hexane/ EtOAc). Mp 205-207°.

1,21-Dideoxy: **Yunnandaphnine B**

[917955-53-4]

$\text{C}_{23}\text{H}_{33}\text{NO}_3$ 371.519

Alkaloid from *Daphniphyllum yunnanense*. Powder. $[\alpha]_D^{25}$ +0.8 (c, 0.2 in CHCl_3).

4,21-Dideoxy: **Yunnandaphnine A**

[917955-52-3]

$\text{C}_{23}\text{H}_{33}\text{NO}_3$ 371.519

Alkaloid from *Daphniphyllum yunnanense*. Powder. $[\alpha]_D^{25}$ -56 (c, 0.2 in CHCl_3).

4,21-Dideoxy, 14,15-didehydro: **Calycinine A**

$\text{C}_{23}\text{H}_{31}\text{NO}_3$ 369.503

Alkaloid from *Daphniphyllum calycinum*. Mp 159-161°.

Trideoxy: Yunnandaphnine D

[917955-55-6]

C₂₃H₃₃NO₂ 355.519

Alkaloid from *Daphniphyllum yunnanense*. Powder. $[\alpha]_D^{25}$ -45.2 (c, 0.21 in CHCl₃).

4-Epimer, 1,21-dideoxy: Yunnandaphnine C

[917955-54-5]

C₂₃H₃₃NO₃ 371.519

Alkaloid from *Daphniphyllum yunnanense*. Powder. $[\alpha]_D^{25}$ -24 (c, 0.22 in CHCl₃).

Sakabe, N. *et al.*, *Tet. Lett.*, 1966, **7**, 963-964 (*Yuzurimine, isol*)

Sakurai, H. *et al.*, *Tet. Lett.*, 1966, **7**, 6309-6314; 1967, 2883-2888 (*Yuzurimine, Yuzurimine B, uv, ir, pmr, ms, cryst struct*)

Nakano, T. *et al.*, *Tet. Lett.*, 1967, **8**, 4791-4797 (*Macrodaphnine, Macrodaphniphyllamine, Macrodaphniphyllidine*)

Sasaki, K. *et al.*, *J.C.S. (B)*, 1971, 1565-1568 (*abs config*)

Niwa, H. *et al.*, *Tet. Lett.*, 1972, **13**, 2697-2700 (*isol*)

Irikawa, H. *et al.*, *Tetrahedron*, 1972, **28**, 3727-3738 (*Yuzurimine, Yuzurimine B, Deoxyyuzurimine*)

Yamamura, S. *et al.*, *Tet. Lett.*, 1974, 2849-2852 (*Daphnijsmine, Deacetyldaphnijsmine*)

Toda, M. *et al.*, *Tetrahedron*, 1974, **30**, 2683-2688 (*isol*)

Yamamura, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 2120-2123 (*cmr*)

Hao, X.-J. *et al.*, *Yunnan Zhiwu Yanjiu*, 1993, **15**, 205-207 (*Calycinine A*)

Morita, H. *et al.*, *Tetrahedron*, 2000, **56**, 2641-2646 (*Daphnezomines J-K*)

El Bitar, H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1094-1099 (*Yuzurimine E, Yuzurimic acid B*)

Takatsu, H. *et al.*, *Tetrahedron*, 2004, **60**, 6279-6284 (*Daphniglaucines*)


Di, Y.-T. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1745-1748 (*Yunnandaphnines A-D, Macrodaphniphyllamine*)

Kubota, T. *et al.*, *Tetrahedron*, 2006, **62**, 4743-4748 (*Daphntenidine D*)

Li, Z.-Y. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 1353-1359 (*17-Oxoyuzurimine, Macropodumine H*)

Daphnezomine L**D-34**

[474535-58-5]



Absolute Configuration

C₂₂H₃₃NO₂ 343.508

Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Solid. $[\alpha]_D^{24}$ -137 (c, 0.1 in MeOH).

7,N-Dihydro: Paxdaphnidine B

[676365-62-1]

C₂₂H₃₅NO₂ 345.524

Alkaloid from *Daphniphyllum paxianum*. Small needles (MeOH). Mp 121-122°. $[\alpha]_D^{20}$ -50.5 (c, 0.58 in MeOH). Exists as zwitterion.

7,N-Dihydro, Me ester: Calyciphylline KC₂₃H₃₇NO₂ 359.551

Alkaloid from *Daphniphyllum calycinum*. Amorph. solid. $[\alpha]_D^{19}$ -38.1 (c, 1 in CHCl₃).


Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 6637-6641 (*Daphnezomine L*)

Zhan, Z.-J. *et al.*, *J.O.C.*, 2004, **69**, 1726-1729 (*Paxdaphnidine B*)

Saito, S. *et al.*, *Tetrahedron*, 2008, **64**, 1901-1908 (*Calyciphylline K*)

Daphnezomine M**D-35**

[474535-59-6]



Absolute Configuration

C₂₂H₃₅NO₂ 345.524

Alkaloid from the stems of *Daphniphyllum humile*. Solid. $[\alpha]_D^{24}$ -21 (c, 0.2 in MeOH).

1,4-Didehydro: Daphnezomine N

[474535-60-9]


C₂₂H₃₃NO₂ 343.508

Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Solid. $[\alpha]_D^{24}$ -113 (c, 0.3 in MeOH).

Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 6637-6641 (*isol, pmr, cmr*)

Daphnezomine O**D-36**

[474535-61-0]



Absolute Configuration


C₂₁H₃₃NO₂ 331.497

Lower homologue of Methyl homodaphniphyllate. Alkaloid from the stems of *Daphniphyllum humile*. Solid. $[\alpha]_D^{24}$ -22 (c, 0.3 in MeOH).

Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 6637-6641 (*isol, pmr, cmr*)

Daphnezomine P**D-37**

[572889-48-6]

R = CH₃C₄₀H₅₇NO₁₃ 759.889

Alkaloid from the fruit of *Daphniphyllum humile*. Amorph. solid. $[\alpha]_D^{+10}$ (c, 1 in MeOH). λ_{max} 232 (ε 10000) (MeOH).

Parent acid: Caldaphnidine F

[871261-45-9]

C₃₉H₅₅NO₁₃ 745.862

Alkaloid from *Daphniphyllum calycinum*. Amorph. powder. $[\alpha]_D^{20}$ +21 (c, 1 in H₂O).

Morita, H. *et al.*, *Tetrahedron*, 2003, **59**, 3575-3579 (*Daphnezomine P*)

Zhan, Z.-J. *et al.*, *Tetrahedron*, 2005, **61**, 11038-11045 (*Caldaphnidine F*)

Daphnezomine Q**D-38**

[572889-49-7]

As Daphnezomine P, D-37 with

R = H

C₃₉H₅₅NO₁₃ 745.862

Alkaloid from the fruit of *Daphniphyllum humile*. Amorph. solid. $[\alpha]_D^{+10}$ (c, 0.6 in MeOH). λ_{max} 233 (ε 10000) (MeOH).

Morita, H. *et al.*, *Tetrahedron*, 2003, **59**, 3575-3579 (*isol, pmr, cmr*)

Daphnicadine**D-39**

[1356-43-0]

C₂₂H₂₇NO₂ 337.461

Struct. unknown. Alkaloid from the seeds of *Daphniphyllum calycinum* (Daphniphyllaceae). Rhombic cryst. Mp 285-287°.

Hydrobromide: Mp 217-219°.

Perchlorate: Mp 184-186°.

Tartrate: Mp 142-144°.

Fang, S.-T. *et al.*, *Huaxue Xuebao*, 1964, **30**, 270-274; *CA*, **61**, 15035e (*isol*)

Daphnicaline**D-40**

[1356-44-1]

C₂₁H₂₉NO₂ 327.466

Struct. unknown. Alkaloid from the seeds of *Daphniphyllum calycinum* (Daphniphyllaceae). Syrup.

Hydrobromide:

Needles. Mp 314°.

Hydroiodide:

Prisms. Mp 316-318°.

Perchlorate:

Leaflets. Mp 304°.

Methiodide:

Needles. Mp 277-281°.

Fang, S.-T. *et al.*, *Huaxue Xuebao*, 1964, **30**, 270-274; *CA*, **61**, 15035e (*isol*)

Daphnicamine**D-41**

[1356-45-2]

C₂₁H₃₁NO₂ 329.481

Struct. unknown. Alkaloid from the seeds of *Daphniphyllum calycinum* (Daphniphyllaceae). Needles + 2H₂O (MeOH). Mp 221-222°.

Perchlorate:


Needles. Mp 310-312°.

Methodide:

Prisms. Mp 254-258°.

Fang, S.-T. *et al.*, *Huaxue Xuebao*, 1964, **30**, 270-274; *CA*, **61**, 15035e (*isol*)**Daphnicyclidine A****D-42**

[385384-22-5]



Absolute Configuration

 $C_{22}H_{25}NO_4$ 367.444Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.**N,4-Dehydro: Daphnicyclidine B**


[384357-80-6]

 $C_{22}H_{24}NO_4^{\oplus}$ 366.436Quaternary alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.**2-Hydroxy: Daphnicyclidine C**

[385384-23-6]

 $C_{22}H_{25}NO_5$ 383.443Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.Kobayashi, J. *et al.*, *J.A.C.S.*, 2001, **123**, 11402-11408 (*isol, uv, cd, pmr, cmr, cryst struct*)**Daphnicyclidine D****D-43**

[385384-24-7]



Absolute Configuration

 $C_{23}H_{27}NO_4$ 381.471Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.**N,4-Dehydro: Daphnicyclidine E**

[385384-25-8]

 $C_{23}H_{26}NO_4^{\oplus}$ 380.463Quaternary alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.**2-Hydroxy: Daphnicyclidine F**


[385384-26-9]

 $C_{23}H_{27}NO_5$ 397.47Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.**De(methoxycarbonyl), 2-hydroxy: Daphnicyclidine G**

[385384-27-0]

 $C_{21}H_{25}NO_3$ 339.433Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.Kobayashi, J. *et al.*, *J.A.C.S.*, 2001, **123**, 11402-11408 (*isol, uv, cd, pmr, cmr*)**Daphnicyclidine H****D-44**


[385384-29-2]



Absolute Configuration

 $C_{23}H_{29}NO_5$ 399.486Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid.Kobayashi, J. *et al.*, *J.A.C.S.*, 2001, **123**, 11402-11408 (*isol, uv, cd, pmr, cmr*)**Daphnicyclidine J****D-45**


[408306-44-5]




Absolute Configuration

 $C_{23}H_{25}NO_5$ 395.454Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid. $[\alpha]_D^{20}$ -15 (c, 0.3 in MeOH). λ_{max} 206 (ε 15600); 245 (ε 6100); 320 (ε 6300); 330 (ε 6200) (MeOH).Morita, H. *et al.*, *J.O.C.*, 2002, **67**, 2278-2282 (*isol, pmr, cmr*)**Daphnicyclidine K****D-46**


[408306-45-6]



Relative Configuration

 $C_{23}H_{27}NO_6$ 413.469Alkaloid from the stems of *Daphniphyllum humile*. Cytotoxic. Amorph. solid. $[\alpha]_D^{20}$ -246 (c, 0.3 in MeOH). λ_{max} 245 (ε 9500); 320 (ε 11500) (MeOH).Morita, H. *et al.*, *J.O.C.*, 2002, **67**, 2278-2282 (*isol, pmr, cmr*)**Daphnicyclidine L****D-47** $C_{23}H_{29}NO_6$ 415.485Zwitterionic. Closely related to Daphnicyclidine H, D-44. Alkaloid from the stem bark of *Daphniphyllum macropodum*. Brown cryst. (CHCl₃MeOH). Mp 235-238°. $[\alpha]_D^{20}$ -117.3 (c, 0.3 in DMSO).Gan, X.-W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1255-1259 (*isol, pmr, cmr, cryst struct*)**Daphniglaucine B****D-48**

[543701-04-8]




Relative Configuration

 $C_{24}H_{34}NO_4$ 400.537Alkaloid from the leaves of *Daphniphyllum glaucescens*. Cytotoxic. Solid. $[\alpha]_D^{20}$ -30 (c, 0.6 in MeOH).**Demethoxy: Daphniglaucine A**

[543701-03-7]

 $C_{23}H_{32}NO_3$ 370.511Alkaloid from the leaves of *Daphniphyllum glaucescens*. Solid. $[\alpha]_D^{20}$ -51 (c, 1 in MeOH).Kobayashi, J. *et al.*, *Org. Lett.*, 2003, **5**, 1733-1736 (*isol, pmr, cmr*)**Daphniglaucine C****D-49**

[692752-40-2]



Relative Configuration

 $C_{23}H_{33}NO_5$ 403.517Alkaloid from the leaves of *Daphniphyllum glaucescens*. Cytotoxic. Amorph. solid. $[\alpha]_D^{20}$ -14 (c, 0.7 in MeOH).**Parent acid: Daphnimacropodine B**

[949880-83-5]


 $C_{22}H_{31}NO_5$ 389.491Alkaloid from the fruit of *Daphniphyllum macropodum*. Amorph. powder. $[\alpha]_D^{20}$ -30.1 (c, 0.15 in MeOH).**Parent acid, ε-lactone: Daphnimacropodine C**

[949880-84-6]

 $C_{22}H_{29}NO_4$ 371.475Alkaloid from the fruit of *Daphniphyllum macropodum*. Amorph. powder. $[\alpha]_D^{20}$ -49.4 (c, 0.3 in Me₂CO).Morita, H. *et al.*, *Tet. Lett.*, 2004, **45**, 901-904 (*Daphniglaucine C*)Kong, N.-C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1348-1351 (*Daphnimacropodines B,C*)

Daphnigraciline

[57362-24-0]

Absolute
ConfigurationR = -CH₂CH₃C₂₃H₃₅NO₄ 389.534Alkaloid from leaves and bark of *Daphniphyllum gracile* (Daphniphyllaceae). Cryst. (hexane/Et₂O). Mp 76-78°.**Me ether: Yuzurine**

[54370-35-3]

C₂₄H₃₇NO₄ 403.561Alkaloid from the bark and leaves of *Daphniphyllum macropodum* and from the leaves of *Daphniphyllum gracile* (Daphniphyllaceae). Viscous liq.; pale-yellow needles (C₆H₆/MeOH) (as methiodide). Mp 229-230° dec. (methiodide).**14,15-Didehydro: Daphgraciline**

[73861-50-4]

C₂₃H₃₃NO₄ 387.518Alkaloid from the bark of *Daphniphyllum gracile* (Daphniphyllaceae). Viscous liq.**11-Oxo: Oxodaphnigraciline**

[57426-46-7]

C₂₃H₃₃NO₅ 403.517Alkaloid from the leaves of *Daphniphyllum gracile* (Daphniphyllaceae). Cryst. (hexane/Et₂O). Mp 107-109°.**11S-Hydroxy, 14,15-didehydro: Hydroxy-daphgraciline**

[73861-52-6]

C₂₃H₃₃NO₅ 403.517Alkaloid from the bark of *Daphniphyllum gracile* (Daphniphyllaceae). Viscous liq.**2-Deoxy, 2,3-didehydro: Dehydrodaphnigraciline**

[57362-27-3]

C₂₃H₃₃NO₃ 371.519Alkaloid from the fruit of *Daphniphyllum oldhamii*. Amorph. solid. [α]_D²⁶ -52 (c, 0.62 in CHCl₃).**Parent acid, Me ether: Yuzuric acid**C₂₃H₃₅NO₄ 389.534Alkaloid from the fruit of *Daphniphyllum oldhamii*. Amorph. solid. [α]_D²⁰ -23.6 (c, 0.6 in MeOH).**2-Epimer, 11-oxo: Epioxodaphnigraciline**

[57361-73-6]

C₂₃H₃₃NO₅ 403.517Alkaloid from leaves of *Daphniphyllum gracile* (Daphniphyllaceae). Cryst. (hexane/Et₂O). Mp 102-104°.Yamamura, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 1836-1840 (*Daphnigraciline*, *Oxodaphnigraciline*, *Epioxodaphnigraciline*)Yamamura, S. *et al.*, *Chem. Lett.*, 1980, 393-396 (*Daphgraciline*, *Hydroxydaphgraciline*)Gan, L.-S. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2395-2400 (*Yuzuric acid*)Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 564-569 (*Dehydrodaphnigraciline*)**Daphnigracine**O³-Demethyl-17-methyluzurine, 9CI

[57362-22-8]

As Daphnigraciline, D-50 with R = -CH(CH₃)₂C₂₄H₃₇NO₄ 403.561Alkaloid from the leaves of *Daphniphyllum gracile* (Daphniphyllaceae). Viscous liq.; cryst. (as methiodide). Mp 198-199° dec. (methiodide).**Me ether (?): Daphnezomine R**

[572889-50-0]

C₂₅H₃₉NO₄ 417.587Alkaloid from the fruit of *Daphniphyllum humile*. Amorph. solid. [α]_D -14 (c, 1 in MeOH). Not interrelated with Daphnigracine. 15-Config. undefined, so could be a stereoisomer of the Me ether.**Me ether, parent acid: Daphnezomic acid**C₂₄H₃₇NO₄ 403.561Alkaloid from the fruit of *Daphniphyllum oldhamii*. Amorph. solid. [α]_D²⁰ -24.9 (c, 0.45 in MeOH).**14,15-Didehydro: Daphgracine**

[73861-49-1]


C₂₄H₃₅NO₄ 401.545Alkaloid from the bark of *Daphniphyllum gracile* (Daphniphyllaceae). Viscous liq.**2-Deoxy, 2,3-didehydro: Dehydrodaphnigracine**C₂₄H₃₅NO₃ 385.545Alkaloid from the fruit of *Daphniphyllum longeracemosum*. Pale yellow needles (as methiodide). Mp 243-245° dec. (methiodide).**11R-Hydroxy: 11-Hydroxydaphnigracine**C₂₄H₃₇NO₅ 419.56Alkaloid from the fruit of *Daphniphyllum longeracemosum*. Amorph. solid. [α]_D²¹ -13.7 (c, 0.81 in MeOH). Called 13-hydroxy in the lit.**11-Oxo: Oxodaphnigracine**

[57362-23-9]


C₂₄H₃₅NO₅ 417.544Alkaloid from the leaves of *Daphniphyllum gracile*. Cryst. (hexane/Et₂O). Mp 116-117°.Yamamura, S. *et al.*, *Chem. Lett.*, 1975, 923; 1980, 393 (*Daphgracine*, *Daphnigracine*)Yamamura, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 1836 (*Daphnigracine*, *Oxodaphnigracine*)Morita, H. *et al.*, *Tetrahedron*, 2003, **59**, 3575-3579 (*Daphnezomine R*)Li, L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1457-1462 (*11-Hydroxydaphnigracine*, *Dehydrodaphnigracine*)Gan, L.-S. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2395-2400 (*Daphnezomic acid*)**Daphnilactone A**

[38210-98-9]

[111795-09-6 ((±)-form)]

Absolute
Configuration**D-51**C₂₃H₃₅NO₂ 357.535Minor alkaloid from the bark and leaves of *Daphniphyllum macropodum* (Daphniphyllaceae). Needles (C₆H₆/hexane). Mp 194.5-195.5°.Sasaki, K. *et al.*, *J.C.S. Perkin 2*, 1972, 1411 (*cmr, cryst struct*)Sasaki, K. *et al.*, *Tet. Lett.*, 1972, 1275 (*pmr*)Toda, M. *et al.*, *Tetrahedron*, 1972, **28**, 1477 (*isol, ir, pmr, ms*)Ruggeri, R.B. *et al.*, *J.A.C.S.*, 1989, **111**, 1530 (*synth*)Heathcock, C.H. *et al.*, *J.O.C.*, 1992, **57**, 2585 (*synth*)**Daphnilactone B**

[38826-56-1]

D-53Absolute
ConfigurationC₂₂H₃₁NO₂ 341.492A major alkaloid from the fruits of *Daphniphyllum macropodum*, *Daphniphyllum teijsmanni* and *Daphniphyllum humile* (Daphniphyllaceae). Plates + 1/2 C₆H₆ (C₆H₆/hexane). Mp 92-94°.**N-Oxide: Daphnezomine I. Daphnilactone B N-oxide**

[276870-42-9]

C₂₂H₃₁NO₃ 357.492Alkaloid from *Daphniphyllum humile*. Amorph. solid. [α]_D²³ -22 (c, 0.6 in CHCl₃).**10,17-Didehydro, 9R,15-dihydro: Iso-daphnilactone B**

[62074-39-9]

C₂₂H₃₁NO₂ 341.492Alkaloid from the leaves of *Daphniphyllum humile* (Daphniphyllaceae). Viscous liq. Mp 196-198° (as methiodide).**Parent hydroxyacid: 2-Deoxydaphnezomine S**

[59630-48-7]

C₂₂H₃₃NO₃ 359.508Alkaloid from the fruits of *Daphniphyllum teijsmanni* (Daphniphyllaceae). Plates (Me₂CO aq.). Mp 247-248° dec. Exists as zwitterion.**Parent hydroxyacid, Me ester: Daphni-macropodine D**

[949880-85-7]

C₂₃H₃₅NO₃ 373.534Alkaloid from the fruit of *Daphniphyllum macropodum*. Gum. [α]_D²⁰ -22.5 (c, 0.2 in MeOH).**Parent hydroxyacid, amide: Macropodumine I**

[949172-79-6]

C₂₂H₃₄N₂O₂ 358.523Alkaloid from the leaves of *Daphniphyllum macropodum*. Oil. [α]_D²⁰ -22.1 (c, 0.38 in MeOH).**2-Hydroxy: Daphnezomine H**

[276870-41-8]


C₂₂H₃₁NO₃ 357.492Alkaloid from *Daphniphyllum humile*. Amorph. solid. [α]_D²³ -59 (c, 0.4 in CHCl₃).

2-Hydroxy, parent hydroxyacid: Daphnezomine S


[572889-51-1]

C₂₂H₃₃NO₄ 375.507Alkaloid from the fruit of *Daphniphyllum humile*. Amorph. solid. [α]_D -7 (c, 1 in MeOH). Zwitterionic.Niwa, H. *et al.*, *Tet. Lett.*, 1972, 2697-2700; 1973, 2129-2132 (*isol, ir, pmr, struct, biosynth*)Sasaki, K. *et al.*, *Acta Cryst. B*, 1973, **29**, 547-553 (*cryst struct*)Toda, M. *et al.*, *Tetrahedron*, 1974, **30**, 2683-2698 (*isol, ir, pmr, ms, struct*)Yamamura, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 839 (*hydroxyacid*)Yamamura, S. *et al.*, *Chem. Lett.*, 1976, 1381 (*Isodaphnilactone B*)Morita, H. *et al.*, *Tetrahedron*, 2000, **56**, 2641-2646; 2003, **59**, 3575-3579 (*Daphnezomines*)Li, Z.Y. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 1353-1359 (*Macropodumine I*)Kong, N.C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1348-1351 (*Daphnimacropodine D*)**Daphnilongeranine A****D-54**

[874201-05-5]

C₂₃H₂₉NO₄ 383.486Alkaloid from *Daphniphyllum longeracemosum*. Gum. [α]_D -52 (c, 0.55 in CHCl₃). λ_{\max} 291 (log ϵ 3.33) (MeOH).Yang, S.-P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 79-82 (*isol, pmr, cmr*)**Daphnilongeranine C****D-55**

[750649-07-1]



Absolute Configuration

C₂₂H₂₉NO₃ 355.476Alkaloid from *Daphniphyllum longeracemosum*. Amorph. powder. [α]_D -22 (c, 0.08 in MeOH). λ_{\max} 213 (log ϵ 3.08) (MeOH).**Me ester: Daphniyunnine A**C₂₃H₃₁NO₃ 369.503Alkaloid from stems and leaves of *Daphniphyllum yunnanense*. Cryst. (Me₂CO). Mp 171-172°. [α]_D -130.5 (c, 0.11 in CHCl₃). λ_{\max} 210 (log ϵ 3.96) (MeOH).**Me ester, N-oxide: Calyciphylline A**

[596799-30-3]

C₂₃H₃₁NO₄ 385.502Alkaloid from the leaves of *Daphniphyllum calycinum*. Solid. [α]_D -51 (c, 1 in MeOH).**6'-Ester with Geniposidic acid: Daphcalycinosidine C**

[752252-75-8]

C₃₈H₄₉NO₁₂ 711.805Alkaloid from the seeds of *Daphniphyllum calycinum*. Amorph. solid. [α]_D -15 (c, 0.6 in MeOH).**14,15-Didehydro, Me ester: Longistylumphylline A**

[857672-34-5]

C₂₃H₂₉NO₃ 367.487Alkaloid from *Daphniphyllum longistylum*. Oil. [α]_D -78 (c, 0.8 in CHCl₃). λ_{\max} 295 (log ϵ 4) (MeOH).**21-Hydroxy, Me ester: Daphniglaucine D**

[753450-62-3]

C₂₃H₃₁NO₄ 385.502Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. [α]_D -21 (c, 0.5 in MeOH).**21-Hydroxy, Me ester, N-oxide: Daphniglaucine G**

[753450-65-6]

C₂₃H₃₁NO₅ 401.502Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. [α]_D -40 (c, 0.5 in MeOH).**21-Acetoxy, Me ester, N-oxide: Daphniglaucine H**

[753450-66-7]

C₂₅H₃₃NO₆ 443.539Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. [α]_D -11 (c, 0.2 in MeOH).**Conformational isomer, 21-hydroxy, Me ester, N-oxide: Daphniglaucine E**


[753450-63-4]

C₂₃H₃₁NO₅ 401.502Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. [α]_D -23 (c, 0.7 in MeOH).**Conformational isomer, 21-acetoxy, Me ester, N-oxide: Daphniglaucine F**

[753450-64-5]

C₂₅H₃₃NO₆ 443.539Alkaloid from the leaves of *Daphniphyllum glaucescens*. Amorph. solid. [α]_D -18 (c, 0.2 in MeOH).Morita, H. *et al.*, *Org. Lett.*, 2003, **5**, 2895-2898 (*Calyciphylline A*)El Bitar, H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1094-1099 (*Daphcalycinosidine C*)Takatsu, H. *et al.*, *Tetrahedron*, 2004, **60**, 6279-6284 (*Daphniglaucines*)Chen, X. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 854-860 (*Longistylumphylline A*)Yang, S.P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 79-82 (*Daphnilongeranin C*)Zhang, H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 553-557 (*Daphniyunnine A*)**Daphnilongerine****D-56**


[910453-13-3]



Relative Configuration

R = -CH(CH₃)₂C₂₄H₃₅NO₄ 401.545Alkaloid from the fruit of *Daphniphyllum longeracemosum*. Amorph. solid. [α]_D +65 (c, 1 in MeOH).**Lower homologue (R = CH₂CH₃):****Daphnioldhanine K**C₂₃H₃₃NO₄ 387.518Alkaloid from the fruit of *Daphniphyllum oldhami*. Amorph. solid. [α]_D +32.2 (c, 0.45 in CHCl₃).Li, L. *et al.*, *Tet. Lett.*, 2006, **69**, 6259-6262 (*isol, pmr, cmr*)Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 564-569 (*isol, pmr, cmr*)**Daphnilongertone****D-57**


[911101-75-2]



Relative Configuration


C₂₃H₃₃NO₄ 387.518Alkaloid from the fruit of *Daphniphyllum longeracemosum*. Amorph. solid. [α]_D -18.1 (c, 0.82 in MeOH).Li, L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1457-1462 (*isol, pmr, cmr*)**Daphnimacrine****D-58**C₂₇H₄₁NO₄ 443.625Struct. unknown. Alkaloid from the bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Amorph. Mp 75-84°. No further reports, and not referred to in modern investigations of this sp.Yagi, S. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1910, **20**, 117-130; *CA*, **4**, 1981 (*isol*)Sakabe, N. *et al.*, *Tet. Lett.*, 1966, **7**, 963-964 (*Daphniphyllum macropodum constits*)Nakano, N. *et al.*, *Tet. Lett.*, 1967, **8**, 4791-4797 (*Daphniphyllum macropodum constits*)**Daphnimacropodine A****D-59**

[949880-81-3]

C₂₂H₃₁NO₃ 357.492Alkaloid from the fruit of *Daphniphyllum macropodum*. Gum. [α]_D +50.5 (c, 1 in Me₂CO).Kong, N.-C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1348-1351 (*isol*)

Daphnine

[80531-83-5]

C₃₇H₃₂N₂O₇ 616.669


Alkaloid from the bark and leaves of *Daphnandra repandula* (Monimiaceae). Orange-yellow cryst. Unique 7,7'-ether link.

Guilhem, J. *et al.*, *Chem. Comm.*, 1981, 1007 (cryst struct)

Howard, A.S. *et al.*, *J. Chem. Res., Synop.*, 1993, 262 (struct)

Daphnioldhanine A

[906647-30-1]

C₂₃H₃₅NO₄ 389.534

Alkaloid from *Daphniphyllum oldhami*. Cryst. (petrol/diethylamine). Mp 118°. [α]_D¹⁸ -30.2 (c, 1.07 in CHCl₃).

4R-Hydroxy, 21-Ac: Daphnioldhanine B [906647-31-2]


C₂₅H₃₇NO₆ 447.57

Alkaloid from *Daphniphyllum oldhami*. Needles (petrol/CHCl₃). Mp 204°. [α]_D¹⁷ -39.1 (c, 0.57 in CHCl₃).

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1065-1069 (isol, pmr, cmr, cryst struct)

Daphnioldhanine C

[906647-32-3]


C₂₂H₃₃NO₄ 375.507

Related to Daphnilactone B, D-53. Alkaloid from *Daphniphyllum oldhami*. Amorph. solid. [α]_D¹⁴ -20.3 (c, 0.85 in CHCl₃).

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1065-1069 (isol, pmr, cmr)

Daphnioldhanine F

D-60

C₃₀H₄₉NO₃ 471.722**(+)-form****26-Ac: Daphnilongeridine**

[922522-15-4]

C₃₂H₅₁NO₄ 513.759

Alkaloid from *Daphniphyllum longeracemosum*. Amorph. powder. [α]_D²⁴ +45 (c, 0.27 in MeOH).

(-)-form [952482-12-1]

Alkaloid from the roots of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²⁵ -48.6 (c, 0.6 in CHCl₃).

26-Ac: Daphnioldhanine G

[952482-13-2]

C₃₂H₅₁NO₄ 513.759

Alkaloid from the roots of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²⁶ -52 (c, 0.34 in CHCl₃).

25-Ketone: Daphnioldhanine D

[952482-10-9]

C₃₀H₄₇NO₃ 469.706

Alkaloid from the roots of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²⁶ -22.8 (c, 0.82 in CHCl₃).

25-Ketone, 26-Ac: Daphnioldhanine E

[952482-11-0]

C₃₂H₄₉NO₄ 511.743

Alkaloid from the roots of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²⁰ -16.1 (c, 0.83 in CHCl₃).

N,1-Didehydro, 25-ketone, 26-Ac: Daphnioldhanine IC₃₂H₄₇NO₄ 509.728

Alkaloid from the roots of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²¹ -135.1 (c, 0.29 in CHCl₃).


Yang, S.-P. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 2783-2788 (*Daphnilongeridine*)

Mu, S.-Z. *et al.*, *Chem. Biodiversity*, 2007, **4**, 129-138 (*Daphnioldhanines D-G*)

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 564-569 (*Daphnioldhanine I*)

Daphnioldhanine J

[1007392-49-5]



Relative Configuration

D-63

C₂₂H₃₁NO₃ 357.492


Alkaloid from the leaves of *Daphniphyllum oldhami*. Amorph. solid. [α]_D²⁵ -5.6 (c, 0.15 in CHCl₃).

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 564-569 (isol, pmr, cmr)

Daphnipaxianine A

D-65

[956243-00-8]

C₂₁H₂₇NO₃ 341.449

Alkaloid from *Daphniphyllum paxianum*. Cryst. (Me₂CO). Mp 175°. [α]_D²² -93.8 (c, 0.32 in CHCl₃). λ_{max} 208 (log ε 3.41); 243 (log ε 3.88) (CHCl₃).

10-Epimer: Daphnipaxianine B

[956243-02-0]

C₂₁H₂₇NO₃ 341.449


Alkaloid from *Daphniphyllum paxianum*. Amorph. powder. [α]_D²² -99.7 (c, 0.3 in CHCl₃). λ_{max} 205 (log ε 3.48); 251 (log ε 3.81) (CHCl₃).

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1628-1631 (isol, pmr, cmr, cryst struct)

Daphnipaxianine D

D-66

[956243-07-5]

C₂₅H₃₉NO₄ 417.587


Alkaloid from *Daphniphyllum paxianum*. Amorph. solid. [α]_D²⁵ -23 (c, 1 in MeOH).

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1628-1631 (isol, pmr, cmr)

Daphnipaxinine

D-67

[693780-84-6]




Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Related to Daphnicyclidine A, D-42. Alkaloid from the stems of *Daphniphyllum paxianum*. Cryst. (Me₂CO). Mp 207° dec. [α]_D²⁰ +148 (c, 0.28 in MeOH). λ_{max} 295 (log ε 4.46); 371 (log ε 4.27) (MeOH).

Yang, S.-P. *et al.*, *Org. Lett.*, 2004, **6**, 1401-1404 (isol, pmr, cmr, ms)

Daphniphylline D-68
Daphniphyllamine
[15007-67-7]



Absolute Configuration

C₃₂H₄₉NO₅ 527.743

One of the main alkaloids from *Daphniphyllum macropodum* and *Daphniphyllum teijsmannii*. Also isol. from the bark of *Daphniphyllum gracile* and from the dried calyx and bract of *Hibiscus sabdariffa* (Daphniphyllaceae, Malvaceae). Non-cryst.; needles (CHCl₃/Et₂O) (as hydrochloride). Mp 238-240° (sealed tube) (hydrochloride).

N-Oxide: Daphnezomine E. *Daphniphylline N-oxide*
[251111-56-5]

C₃₂H₄₉NO₆ 543.742

Alkaloid from *Daphniphyllum humile*. Solid. [α]_D²⁴ +31 (c, 1.4 in CHCl₃).

O-De-Ac: Daphnilongeranine D

[42521-95-9]

C₃₀H₄₇NO₄ 485.706

Alkaloid from *Daphniphyllum longercemosum*. Amorph. powder. [α]_D²⁰ +50 (c, 0.23 in MeOH). λ_{max} 202 (log ε 2.46) (MeOH).

O-De-Ac, 14-ketone, 22ξ-alcohol: Daphniphyllidine

[50764-62-0]

C₃₀H₄₇NO₄ 485.706

Alkaloid from the leaves and bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Mp 264° dec. (as methiodide). Isomer of Deacetyldaphniphylline.

7S-Hydroxy: Daphnioldhanine H

C₃₂H₄₉NO₆ 543.742

Alkaloid from the leaves of *Daphniphyllum oldhami*. Amorph. powder. [α]_D²³ +11.4 (c, 0.34 in CHCl₃).

Deacetoxy: Codaphniphylline. *Desacetoxydaphniphylline*

[14694-15-6]

C₃₀H₄₇NO₃ 469.706

Alkaloid from the bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Cryst. (CHCl₃/Et₂O) (as hydrochloride). Mp 266-267° (sealed tube) (as hydrochloride). [α]_D +4.2 (c, 2.40 in CHCl₃).

Deacetoxy, 7S-hydroxy: Daphnimacropine
[20249-89-2]

C₃₀H₄₇NO₄ 485.706

Alkaloid from the bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Mp 306-307° (as methiodide).

Sakabe, N. *et al.*, *Tet. Lett.*, 1966, 965-968

(*Daphniphylline, cryst struct*)

Kamijo, N. *et al.*, *Tet. Lett.*, 1966, 2889-2892

(*Daphnimacropine, cryst struct*)

Nakano, T. *et al.*, *Tet. Lett.*, 1967, 4791-4797

(*Daphniphylline*)

Irikawa, H. *et al.*, *Tetrahedron*, 1968, **24**, 5691-

5700 (*Daphniphylline, Codaphniphylline*)

Toda, M. *et al.*, *Nippon Kagaku Zasshi*, 1970,

91, 103; *C.A.* **73**, 22137j (*Daphniphylline,*

Codaphniphylline, isol)

Sasaki, K. *et al.*, *J.C.S. (B)*, 1971, 1565-1568

(*Daphniphylline, abs config*)

Toda, M. *et al.*, *Tet. Lett.*, 1973, 797-798

(*Daphniphyllidine*)

Suzuki, K.T. *et al.*, *Tet. Lett.*, 1973, 799-802

(*Daphniphylline, Codaphniphylline, biosynth*)

Toda, M. *et al.*, *Tetrahedron*, 1974, **30**, 2683-

2688 (*Daphniphylline*)

Yamamura, S. *et al.*, *Alkaloids (Academic*

Press), 1975, **15**, 41-81 (rev)

Yamamura, S. *et al.*, *Chem. Lett.*, 1980, 393-

396 (*Daphniphylline*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1995, **60**, 1120-

1130 (*Codaphniphylline*)

Morita, H. *et al.*, *Tetrahedron*, 1999, **55**,

12549-12556 (*Daphnezomine E*)


Yang, S.-P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 79-82

(*Daphnilongeranine D*)

Mu, S.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 564-

569 (*Daphnioldhanine H*)

Daphniyunnine D D-69
[881388-90-5]



C₂₁H₂₇NO₃ 341.449

Alkaloid from the stems and leaves of *Daphniphyllum yunnanense*. Cytotoxic. Amorph. powder. [α]_D²⁰ -140 (c, 0.12 in CHCl₃). λ_{max} 213 (log ε 3.92); 230 (log ε 3.9) (MeOH).

15-Deoxy: Daphnilongeranine B

[874201-06-6]

C₂₁H₂₇NO₂ 325.45

Alkaloid from *Daphniphyllum longercemosum*. Amorph. powder. [α]_D²⁰ -159 (c, 0.1 in MeOH). λ_{max} 244 (log ε 3.32) (MeOH).

17-Deoxy: Daphnipaxianine C

[956243-05-3]

C₂₁H₂₉NO₂ 327.466

Alkaloid from *Daphniphyllum paxianum*. Amorph. powder. [α]_D²² -153.2 (c, 0.26 in CHCl₃).

15-Epimer: Daphniyunnine E

[881388-91-6]

C₂₁H₂₇NO₃ 341.449

Alkaloid from the stems and leaves of *Daphniphyllum yunnanense*. Amorph. powder. [α]_D²⁰ -28.8 (c, 0.13 in CHCl₃). λ_{max} 245 (log ε 3.9) (MeOH).

Yang, S.-P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 79-82

(*Daphnilongeranine B*)


Zhang, H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 553-

557 (*Daphniyunnines D,E*)

Mu, S.Z. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1628-

1631 (*Daphnipaxianine C*)

Daphtenidine A D-70




C₃₁H₄₇NO₄ 497.717

Alkaloid from the leaves of *Daphniphyllum teijsmannii*. Amorph. solid. [α]_D²³ -3 (c, 0.1 in MeOH).

Kubota, T. *et al.*, *Tetrahedron*, 2006, **62**, 4743-4748 (isol, pmr, cmr)

Daphtenidine B D-71




C₂₄H₃₇NO₂ 371.562

Alkaloid from the leaves of *Daphniphyllum teijsmannii*. Amorph. powder. [α]_D²³ +97 (c, 0.1 in MeOH).

Kubota, T. *et al.*, *Tetrahedron*, 2006, **62**, 4743-4748 (isol, pmr, cmr)

Daphtenidine C D-72




C₂₇H₃₅NO₇ 485.576

Alkaloid from the leaves of *Daphniphyllum teijsmannii*. Amorph. solid. [α]_D²³ -106 (c, 1 in MeOH).

Kubota, T. *et al.*, *Tetrahedron*, 2006, **62**, 4743-4748 (isol, pmr, cmr)

Dapiramicin B D-73
[90044-18-1]



C₂₁H₂₉N₅O₁₁ 527.487

Nucleoside antibiotic. Isol. from *Micromonospora* sp. SF-1917. Active against sheath blight in rice plants; less effective than Dapiramicin A. Needles + 1H₂O (MeOH aq.). Sol. AcOH, DMF; fairly sol. MeOH, H₂O; poorly sol. Me₂CO, Et₂O, CHCl₃, EtOAc. Mp 241–243°. [α]_D²⁰ -37.6 (c, 1 in 50% AcOH). λ_{\max} 228 (ε 21700); 250 (ε 22500); 305 (ε 7670) (0.1N NaOH) (Derep). λ_{\max} 227 (ε 42400); 290 (ε 7270) (H₂O) (Derep).

6'-Deoxy: Epidapiramicin A

[90044-19-2]

C₂₁H₂₉N₅O₁₀ 511.488

Isol. from *Micromonospora* sp. Shows activity against sheath blight in rice plants. Less effective than Dapiramicin A. Needles. Mp 222–224°. [α]_D²⁰ -14.4 (c, 1.0 in AcOH). Artifact. λ_{\max} 228 (ε 47500); 292 (ε 8000) (H₂O).


1'-Epimer, 6'-deoxy: Dapiramicin A. SF 1917. Antibiotic SF 1917

[67298-15-1]

C₂₁H₂₉N₅O₁₀ 511.488

Isol. from *Micromonospora* sp. SF-1917. Effective against sheath blight in rice plants caused by *Rhizoctonia solani*. Needles. Sol. AcOH, DMF; fairly sol. MeOH, H₂O; poorly sol. Me₂CO, Et₂O, CHCl₃, EtOAc. Mp 220–222°. [α]_D²⁰ +117 (c, 1.0 in MeOH). λ_{\max} 228 (ε 21700); 250 (ε 22500); 305 (ε 7670) (0.1N NaOH) (Derep). λ_{\max} 228 (ε 40200); 289 (ε 8640) (H₂O) (Derep).

- LD₅₀ (mus, ivn) 400 200 mg/kg, LD₅₀ (mus, ipr) 1000 500 mg/kg. UY9105600
Nishizawa, N. *et al.*, *J. Antibiot.*, 1984, **37**, 1-5 (isol, uv, ir, pmr, props)
Ohno, H. *et al.*, *Tet. Lett.*, 2006, **47**, 5747-5750 (synth)

Darlingianine**D-74**

(E,E)-form

C₁₇H₂₃NO 257.375

Rel. config. only known.

(E,E)-form [65600-88-6]

Alkaloid from the upper stems and leaves of *Darlingia darlingiana* (Proteaceae). Straw-coloured rhombic cryst. (EtOH). Mp 93.5°. [α]_D¹⁹ +62 (CHCl₃).

Ac:

Cryst. (EtOH). Mp 82.5°.

Methiodide:

Cryst. (MeOH). Mp 123–126°.

3,4-Dihydro: Dihydrodarlingianine

[73045-39-3]

C₁₇H₂₅NO 259.391

Isol. from *Darlingia darlingiana* (Proteaceae). Straw-coloured cryst. (EtOH). Mp 72–74°. [α]_D¹⁹ +34 (CHCl₃).

Tetrahydro: Tetrahydrodarlingianine

[73045-40-6]

C₁₇H₂₇NO 261.406

Isol. from *Darlingia darlingiana* (Proteaceae). Noncryst. Mp 178–181° (as picrate). [α]_D¹⁹ +22 (CHCl₃).

2-Ketone: Dehydrodarlingianine

[73045-41-7]

C₁₇H₂₁NO 255.359

Isol. from *Darlingia darlingiana* (Proteaceae). Pale-yellow plates (petrol). Mp 54–55°. [α]_D¹⁹ +53 (CHCl₃).

(E,Z)-form**Isodarlingianine**


[73069-58-6]

Alkaloid from the upper stems and leaves of *Darlingia darlingiana* (Proteaceae). Straw-coloured cryst. (petrol). Mp 50–52°. [α]_D¹⁹ +47 (CHCl₃). Isomerises to Darlingianine when refluxed with iodine in xylene.

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 2523 (isol, uv, ir, pmr, cmr, ms, struct, derivs)

Darlingine**D-75**

6,7,8,9-Tetrahydro-2,3,10-trimethylcyclohepta[b]pyran-5,8-imin-4(5H)-one, 9CI [58471-10-6]

C₁₃H₁₇NO₂ 219.283

Authors' numbering shown. Alkaloid from the upper stems and leaves of *Darlingia darlingiana*, the leaves, stems and roots of *Darlingia ferruginea*, and from the upper leaves, flowers and flowering stems of *Bellendena montana* (Proteaceae). Needles (Et₂O). Mp 166–167°. [α]_D¹⁹ +104 (CHCl₃).

Hydrochloride: Mp 180–182°.

N-Oxide: Darlingine N-oxideC₁₃H₁₇NO₃ 235.282

Alkaloid from *Darlingia darlingiana*. [α]_D²⁷ +16.8 (c, 0.67 in MeOH). λ_{\max} 216 (log ε 4); 258 (log ε 4) (MeOH).

2,3-Dihydro: 2,3-Dihydrodarlingine

[72362-48-2]

[72401-76-4]

C₁₃H₁₉NO₂ 221.299

Alkaloid from the upper leaves, flowers and flowering stems of *Bellendena montana* (Proteaceae). Oil. Mp 191–194° (as picrate). [α]_D¹⁹ +68 (CHCl₃).

5,11-Dihydro: 5,11-Dihydrodarlingine

[73088-64-9] (picrate)

C₁₃H₁₉NO₂ 221.299

Alkaloid from the upper stems and leaves of *Darlingia darlingiana* (Proteaceae). Noncryst. Mp 165–168° (as picrate). [α]_D¹⁹ +37 (CHCl₃). CAS no. refers to picrate.

10R-Hydroxy: 10-HydroxydarlingineC₁₃H₁₇NO₃ 235.282Alkaloid from *Triunia erythrocarpa*

(Proteaceae). Solid. [α]_D²⁷ +12 (c, 0.3 in CHCl₃). Ref. states C-10 config. as (R)- but shows diag. with (S)-config. λ_{\max} 216 (log ε 4); 258 (log ε 4) (MeOH).

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 1827–1840; 2523–2536; 2537–2543 (isol, uv, ir, pmr, cmr, ms, struct, derivs)

Lounasmaa, M. *et al.*, *Planta Med.*, 1983, **48**, 56–58 (synth, 2,3-Dihydrodarlingine)


Majewski, M. *et al.*, *J.O.C.*, 1995, **60**, 5825–5830 (synth, abs config)

Katavic, P.L. *et al.*, *Phytochemistry*, 1999, **52**, 529–531 (isol, pmr, cmr, N-oxide)

Butler, M.S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 688–689 (10-Hydroxydarlingine)

Darlinine**D-76**

[73069-56-4]

C₁₅H₂₁NO 231.337

Abs. config. not detd. Alkaloid from the upper stems and leaves of *Darlingia darlingiana* (Proteaceae). Straw-coloured cryst. (EtOH). Mp 59–61°. [α]_D¹⁹ +75 (CHCl₃).

2-Epimer: Epidarlinine

[73045-38-2]

C₁₅H₂₁NO 231.337

Isol. from *Darlingia darlingiana* (Proteaceae). Straw-coloured cryst. (EtOH). Mp 66–68°. [α]_D¹⁹ +59 (CHCl₃).

2-Ketone: Dehydrodarlinine

[73045-42-8]


C₁₅H₁₉NO 229.321

Isol. from *Darlingia darlingiana* (Proteaceae). Pale-yellow plates (petrol). Mp 42–44°. [α]_D¹⁹ +64 (CHCl₃).

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 2523 (isol, uv, ir, pmr, ms, struct, derivs)

Darlucin A**D-77**

[162341-15-3]


C₁₉H₁₆N₂O₃ 320.347

Prod. by *Sphaerellopsis filum* (preferred name *Eudarlucia caricis*, also known as *Darlucina filum*). Active against bacteria and fungi. Shows weak cytotoxic props. Oil. Similar to Xanthocillin X, X-7. λ_{\max} 225 (ε 24500); 283 (ε 4570) (MeOH) (Derep). λ_{\max} 225 (ε 24550); 283 (ε 4571) (MeOH) (Berdy).

Zapf, S. *et al.*, *J. Antibiot.*, 1995, **48**, 36 (isol, uv, ir, pmr, cmr, ms)

Darlucin B

[162341-16-4]

C₁₉H₂₀N₂O₃ 324.379


Prod. by *Sphaerellopsis filum* (preferred name *Eudarlucia caricis*, also known as *Darlucia filum*). Active against bacteria and fungi. Shows weak cytotoxic props. Oil. λ_{max} 226 (ε 12900); 275 (sh) (ε 3630) (MeOH) (Derep). λ_{max} 226 (ε 12882); 275 (ε 3630) (MeOH) (Berdy).

Zapf, S. et al., *J. Antibiot.*, 1995, **48**, 36 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Darvasamine

D-79

(5β,11α)-Matridin-11-one, 9CI. Tetrahydro-neosoporphamine [36284-98-7]

C₁₅H₂₄N₂O 248.367

Config. uncertain. Stereoisomer of Matrine, M-121, Sophoridine, S-380, Isomatrine, I-244 and Allomatrine, A-628. The phys. props. of the natural alkaloid Darvasamine do not coincide with the semisynthetic tetrahydroneosoporphamine of known config. Alkaloid from *Leontice darvasica* and from tubers of *Gymnospermium alberti* (Leonticaceae). Cryst. (Et₂O). Mp 102° (natural) Mp 152-154° (synthetic). [α]_D²⁰ -21 (c, 1.9 in EtOH) (synthetic).

Perchlorate:

Cryst. (Me₂CO). Mp 265° dec.

Methiodide:

Cryst. (Me₂CO). Mp 303-305° dec.

13ξ-Hydroxy: Albertamine

[54383-31-2]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from epigeal parts of *Leontice albertii* (Leonticaceae). Mp 190-192°. [α]_D +11.4 (c, 0.4 in EtOH).

13-Hydroxy; perchlorate: Mp 119-120°.

13-Hydroxy; picrate:

Cryst. (EtOH). Mp 84-85°.

Zunnunzhanov, S. et al., *Khim. Prir. Soedin.*, 1971, **7**, 851; *Chem. Nat. Compd. (Engl. Transl.)*, 838 (*isol*)

Sadykov, B. et al., *Khim. Prir. Soedin.*, 1974, **10**, 377; *Chem. Nat. Compd. (Engl. Transl.)*, 374 (*Albertamine*)

Monakhova, T.E. et al., *Khim. Prir. Soedin.*, 1974, **10**, 472; *Chem. Nat. Compd. (Engl. Transl.)*, 477 (*synth*)

Ibragimov, B.T. et al., *Khim. Prir. Soedin.*, 1979, **15**, 588; *Chem. Nat. Compd. (Engl. Transl.)*, 517 (*cryst struct*)

Kurbanov, M. et al., *CA*, 1982, **97**, 141727g (*isol*)

Ibragimov, B.T. et al., *Khim. Prir. Soedin.*, 1982, **18**, 71; *Chem. Nat. Compd. (Engl. Transl.)*, 66 (*rev*)


Darvasine

D-80

5,17-Didehydromatridin-15-one, 9CI†.

5,17-Dehydroisomatrine

[24533-37-7]

C₁₅H₂₂N₂O 246.352

Stereoisomeric with Isoleontalbine, I-241 and Leontalbine, L-93. Abs. config. not well established. Alkaloid from epigeal parts of *Leontice darvasica*, also isol. from tubers of *Gymnospermium alberti* (Leonticaceae). Acicular cryst. Mp 145-146°.

Perchlorate: Mp 249-250°.

Picrate:

Cryst. (EtOH). Mp 231° dec.

Iskandarov, S. et al., *Khim. Prir. Soedin.*, 1969, **5**, 132; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 115 (*isol, struct*)

Zunnunzhanov, A. et al., *Khim. Prir. Soedin.*, 1974, **10**, 373; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 371 (*isol, struct, abs config*)

Kurbanov, M. et al., *CA*, 1982, **97**, 141727g (*isol*)


Darvasoline

D-81

11-Hydroxymatridin-15-one, 9CI†.

11-Hydroxymatrine

[52484-77-2]

C₁₅H₂₄N₂O₂ 264.367

Abs. config. not clear. Stereoisomeric with Leontismine, L-99. Alkaloid from epigeal parts of *Leontice darvasica* (Leonticaceae). Cryst. (Et₂O). [α]_D +28.


Perchlorate: Mp 235-237°.

Methiodide: Mp 247-249°.

Zunnunzhanov, A. et al., *Khim. Prir. Soedin.*, 1974, **10**, 115; *Chem. Nat. Compd. (Engl. Transl.)*, 131 (*isol, struct*)

Daryamide A

[918785-41-8]



Absolute Configuration

C₁₈H₂₆N₂O₅ 350.414

Prod. by the marine-derived *Streptomyces* strain CNQ-085. Pale yellow powder. Mp 153-154°. [α]_D²⁵ -6.9 (c, 0.25 in MeOH). λ_{max} 271 (log ε 4.22); 285 (log ε 4.23) (MeOH).

5-Deoxy, 5,6-didehydro: **Daryamide B**

[918785-42-9]

C₁₈H₂₄N₂O₄ 332.399

Prod. by *Streptomyces* strain CNQ-085. Dark yellow oil. [α]_D²⁵ -42.8 (c, 0.51 in MeOH). λ_{max} 278 (log ε 4.78) (MeOH).

5-Deoxy, 5,6-didehydro, N-deacyl, N-(6-methyl-2E,4E-heptadienoyl): **Daryamide C**

[918785-43-0]

C₁₇H₂₂N₂O₄ 318.372


Prod. by *Streptomyces* strain CNQ-085. Dark yellow oil. [α]_D²⁵ -17.4 (c, 0.075 in MeOH). λ_{max} 273 (log ε 4.19) (MeOH).

Asolkar, R.N. et al., *J. Nat. Prod.*, 2006, **69**, 1756-1759 (*isol, pmr, cmr*)

Dasycarine

D-83

4,5,8-Trimethoxy-3-(3-methyl-2-but-enyl)-2(1H)-quinolinone. 4,5,8-Trimethoxy-3-prenyl-2(1H)-quinolone

C₁₇H₂₁NO₄ 303.357

Alkaloid from *Dictamnus dasycarpus*. Needles (Me₂CO). Mp 172-173°.

Chen, J. et al., *Chin. Chem. Lett.*, 2000, **11**, 707-708


Dasycarpidone

D-84

Dasycarpidan-1-one, 9CI

[2825-08-3]

[18700-27-1 (±)-form]



Absolute Configuration

C₁₇H₂₀N₂O 268.358

Alkaloid from the bark of *Aspidosperma dasycarpon* (Apocynaceae). Noncryst. [α]_D²⁶ +64.7 (c, 1.02 in CHCl₃). λ_{max} 237

(log ϵ 4.15); 316 (log ϵ 4.29) (EtOH).

N-De-Me: N-Nordascarpidone. De-N-methylascarpidone
[3620-81-3]

$C_{16}H_{18}N_2O$ 254.331

Alkaloid from the bark of *Aspidosperma dasycarpon* (Apocynaceae). Cryst. (CHCl₃). Mp 208-210°. λ_{max} 238 (log ϵ 4.15); 317 (log ϵ 4.23) (EtOH).

1 β -Alcohol: Dascarpidol

[2744-46-9]

$C_{17}H_{22}N_2O$ 270.374

Alkaloid from the bark of *Aspidosperma dasycarpon* (Apocynaceae). Mp 118-122°. $[\alpha]_D^{27}$ -54 (c, 1.03 in EtOH). λ_{max} 220 (log ϵ 4.54); 282 (log ϵ 3.89); 290 (log ϵ 3.81) (EtOH).

3-Epimer: 3-Epidascarpidone

[16530-89-5]

[18688-38-5 (\pm)-form]

$C_{17}H_{20}N_2O$ 268.358

Alkaloid from *Aspidosperma subincanum* (Apocynaceae). Noncryst.

Ohashi, M. *et al.*, *Experientia*, 1964, **20**, 363-364 (*uv. struct.*, *Dascarpidol*)

Joule, J.A. *et al.*, *J.C.S.*, 1964, 2777-2790 (*ms*)

Joule, J.A. *et al.*, *Tetrahedron*, 1965, **21**, 1717-1734 (*Dascarpidone*, *Nordascarpidone*, *Dascarpidol*)

Gaskell, A.J. *et al.*, *Chem. Ind. (London)*, 1967, 1089-1090 (*config.*, *epimer*, *pmr*)

Shamma, M. *et al.*, *Tet. Lett.*, 1967, 2489-2492 (*Dascarpidol*, *struct*)

Jackson, A. *et al.*, *J.C.S. (C)*, 1969, 2738-2747 (*synth*)

Dolby, L.J. *et al.*, *J.O.C.*, 1970, **35**, 3843-3845 (*synth*)

Kametani, T. *et al.*, *J.O.C.*, 1971, **36**, 1291-1293 (*synth*, *uv.*, *ir.*, *pmr.*, *ms.*, *epimer*)

Bonjoch, J. *et al.*, *Chem. Comm.*, 1991, 1687-1688 (*Dascarpidone*, *Dascarpidol*, *Nordascarpidone*, *synth.*, *cmr*)

Gràcia, J. *et al.*, *J.O.C.*, 1994, **59**, 3939-3951 (*Dascarpidone*, *Nordascarpidone*, *Dascarpidol*, *synth*)

Blechert, S. *et al.*, *Synthesis*, 1995, 592-604 (3-*Epidascarpidone*, *synth*)

Forns, P. *et al.*, *Tetrahedron*, 1996, **52**, 3563-3574 (*epimer*, *synth*)


Sato, M. *et al.*, *Chem. Comm.*, 1997, 765-766 (*synth.*, *abs config*)

Uludag, N. *et al.*, *J. Het. Chem.*, 2006, **43**, 585-591 (*synth.*, *epimer*)

Dascarpine

[10550-82-0]

D-85



Relative
Configuration

$C_{20}H_{35}N_3$ 317.517

Stereoisomeric with Templetine, T-68, Ormosanine, O-115 and Piptanthine, P-476. CAS gives an absolute config. but there does not seem to be any evidence in the lit. for this. Major alkaloid from *Ormosia dasycarpa* (Fabaceae). Oil. Bp_{0.5} 160-165°. $[\alpha]_D^{20}$ +12.5 (c, 1.1 in EtOH).

Hydrochloride:

Needles + 1H₂O (EtOH/Et₂O or EtOAc). Mp 205-207° dec.

Perchlorate:

Needles + 3H₂O (EtOH/Et₂O). Mp 240° dec.

N-Ac:

Plates (EtOH). Mp 155-157°.


Clarke, R.T. *et al.*, *J.C.S.*, 1960, 41-42; 1963, 535-539 (*isol*)

Deslongchamps, P. *et al.*, *Can. J. Chem.*, 1966, **44**, 2539-2551 (*struct*)

Dasyrachine

[237078-15-8]

D-86



Absolute
Configuration

$C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from *Kopsia dasyrachis*. $[\alpha]_D^{20}$ +17 (c, 0.06 in CHCl₃). λ_{max} 221 (log ϵ 4.34); 243 (log ϵ 3.93); 281 (log ϵ 3.06) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 159-169 (*isol.*, *uv.*, *pmr.*, *cmr.*, *ms*)

Daucine

$C_{11}H_{18}N_2$ 178.277

D-87


Struct. unknown. Alkaloid from the leaves of *Daucus carota* (carrot) (Apiaceae). Oil. Bp 240-250°. $[\alpha]_D^{20}$ +7.74 (c, 0.645 in Et₂O).

Pictet, A. *et al.*, *Ber.*, 1907, **40**, 3771 (*isol*)

Dauricine

6,6'-Di-O-methylauricoline
[524-17-4]

D-88



$C_{38}H_{44}N_2O_6$ 624.775

Alkaloids covered by this entry are enantiomeric with those covered by Thalibrine, T-309 (*S,S*-config.) and diastereoisomeric with those covered by Berbaminine, B-98 (1*R*,1'*S*-config.) and Grisabine, G-177 (1*S*,1'*R*-config.). Alkaloid from the rhizomes of *Menispermum dauricum*, the subterranean parts of *Polypodium canadense*, and from *Polypodium nitidissima* (Menispermaceae, Annonaceae). Antiarrhythmic, antihypertensive agent. Shows antiinflammatory and anaesthetic props. Weak curarising agent. Platelet aggregation inhibitor. Mp 115°. $[\alpha]_D^{20}$ -139 (MeOH). Log P 6.67 (calc).

► LD₅₀ (mus, orl) 1180 mg/kg; LD₅₀ (mus, ipr) 185 mg/kg; LD₅₀ (mus, ipr) 125-175 mg/kg. SM9490000

N,N'-Di-Me:

Needles (MeOH) (as diiodide). Mp 201° (diiodide). $[\alpha]_D^{20}$ -114 (c, 0.348 in H₂O).

N²-De-Me: N-Demethylauricine

[146763-55-5]

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloid from rhizomes of *Menispermum dauricum* (Menispermaceae).

N²-De-Me: N'-Demethylauricine

[34302-34-6]

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloid from the rhizomes of *Menispermum canadense* (Menispermaceae). Unstable oil. $[\alpha]_D^{27}$ -98 (MeOH).

Me ether: O-Methylauricine

[2202-17-7]

$C_{39}H_{46}N_2O_6$ 638.802

Alkaloid from the bark of *Colubrina asiatica* and *Popowia* cf. *cyanocarpa* (Rhamnaceae, Annonaceae). Active against gram-positive bacteria. Shows activity against Walker 256 carcinoma. Brittle foam. $[\alpha]_D^{27}$ -128 (c, 0.06 in MeOH).

► LD₅₀ (mus, ipr) 125 mg/kg. NX5018450

Me ether, 2-N-oxide: O-Methylauricine

2-N-oxide

[107882-03-1]

$C_{39}H_{46}N_2O_7$ 654.802

Alkaloid from the bark and leaves of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D^{20}$ -138 (c, 0.5 in MeOH). Isol. as an inseparable mixt. with the 2'-N-oxide (ratio 10:7 or 7:10).

Me ether, 2'-N-oxide: O-Methylauricine

2'-N-oxide

[107882-02-0]

$C_{39}H_{46}N_2O_7$ 654.802

Alkaloid from the bark and leaves of *Popowia pisocarpa* (Annonaceae). Amorph.

O⁶-De-Me: Dauricinoline

[30984-80-6]

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloid from rhizomes of *Menispermum dauricum* (Menispermaceae). Muscle relaxant. Amorph. powder. $[\alpha]_D^{21}$ -94.6 (MeOH).

O⁶-De-Me, O¹²-Me: Popidine

[107882-00-8]

$C_{38}H_{44}N_2O_6$ 624.775

Alkaloid from the bark and leaves of *Popowia pisocarpa* (Annonaceae). Amorph.

O⁶-De-Me: Daurinoline

[2831-75-6]

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloid from the rhizomes of *Menispermum canadense* and *Menispermum dauricum* (Menispermaceae). Muscle relaxant. Mp 95-98°. $[\alpha]_D^{27}$ -114 (MeOH). λ_{max} 210 (c 26400); 285 (c 6750) (MeOH) (Berdy).

O⁶-De-Me, O¹²-Me: Popisidine

[107881-99-2]


$C_{38}H_{44}N_2O_6$ 624.775

Alkaloid from the bark and leaves of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D^{20}$ -128 (c, 1 in MeOH). Isol. as an inseparable mixt. with Popidine (approx. ratio 1:2).

- O⁷-De-Me: Daurisoline**
[70553-76-3]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from *Menispermum dauricum* (Menispermaceae). Muscle relaxant. Mp 96-102°. [α]_D²⁰ -129 (c, 0.65 in MeOH). Log P 6.4 (calc).
▶ NX5995000
- O⁷-De-Me, N²-de-Me: Pampulhamine.**
2'-N-Nordaurisoline. 7'-O-Methylpedroamine
[110300-71-5]
C₃₆H₄₀N₂O₆ 596.722
Alkaloid from the stems of *Abuta pahni* (Menispermaceae) and leaves of *Aristolochia gigantea* (Aristolochiaceae). [α]_D -58 (c, 0.18 in MeOH). Opt. rotn. refers to Pampulhamine.
- O⁷-De-Me, N,N'-di-de-Me: 7'-O-Methyl-lindoldhamine**
[88524-57-6]
C₃₅H₃₈N₂O₆ 582.695
Alkaloid from *Polyalthia nitidissima* (Annonaceae). [α]_D +21 (c, 0.2 in MeOH).
- O⁷-De-Me, O¹²-Me: 7'-O-Methylcuspidaline**
[94410-10-3]
C₃₈H₄₄N₂O₆ 624.775
Alkaloid from the leaves of *Aristolochia elegans* (Aristolochiaceae). Amorph. powder. [α]_D -105 (c, 0.001 in CHCl₃).
- O⁷-De-Me, O¹²-Me, N²-de-Me: Geraldoamine. 7,12-Di-O-methylpedroamine. 12-O-Methylpampulhamine**
[112523-89-4]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from leaves of *Aristolochia gigantea*. Amorph. powder. [α]_D -65 (c, 0.1 in MeOH).
- O⁷-De-Me: Isodaurisoline**
[88524-56-5]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from *Polyalthia nitidissima* (Annonaceae). Mp 105-115°. [α]_D -150 (c, 0.6 in MeOH).
- O⁷-De-Me, N,N'-di-de-Me: 7-O-Methyl-lindoldhamine**
[88524-58-7]
C₃₅H₃₈N₂O₆ 582.695
Alkaloid from *Polyalthia nitidissima* (Annonaceae). [α]_D +21 (c, 0.2 in MeOH).
- O⁷-De-Me, O¹²-Me: Popisine**
[107882-17-7]
C₃₈H₄₄N₂O₆ 624.775
Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. [α]_D -148 (c, 1 in MeOH).
- O⁶,O⁶-Di-de-Me: Dauricoline**
[29550-42-3]
C₃₆H₄₀N₂O₆ 596.722
Alkaloid from rhizomes of *Menispermum dauricum* (Menispermaceae). Muscle relaxant. Amorph. powder. [α]_D²⁰ -105 (MeOH).
- O⁶,O⁶-Di-de-Me, O¹²-Me: Popisonine**
[107882-01-9]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from the bark and leaves of *Popowia pisocarpa* (Annonaceae).
Amorph. [α]_D -118 (c, 1 in MeOH).
- O⁶,O⁷-Di-de-Me: Dauricoline**
[256642-22-5]
C₃₆H₄₀N₂O₆ 596.722
Alkaloid from the rhizomes of *Menispermum dauricum*.
- O⁶,O⁷-Di-de-Me, O¹²-Me: Popisopine**
[107913-38-2]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. [α]_D -119 (c, 1 in MeOH).
- O⁷,O⁷-Di-de-Me: Guattegaumerine. Dauriciline. N,N'-Dimethylindoldhamine**
[21446-35-5]
C₃₆H₄₀N₂O₆ 596.722
Alkaloid from *Polyalthia nitidissima* and *Guatteria gaumeri* (Annonaceae). Also isol. from *Menispermum dauricum* (Menispermaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D -90 (c, 0.7 in CHCl₃).
- O⁷,O⁷-Di-de-Me, N²-de-Me: 2'-N-Methylindoldhamine**
C₃₅H₃₈N₂O₆ 582.695
Alkaloid from the stems of *Abuta pahni* (Menispermaceae). [α]_D -47 (c, 0.17 in MeOH).
- O⁷,O⁷-Di-de-Me, N²-de-Me: Pedroamine. 2-N-Methylindoldhamine**
[110320-85-9]
C₃₅H₃₈N₂O₆ 582.695
Alkaloid from the stems of *Abuta pahni* (Menispermaceae) and leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. powder. [α]_D -74 (c, 0.1 in MeOH). Enantiomer of Northalibroline, in T-309. Opt. rotn. refers to Pedroamine.
- O⁷,O⁷-Di-de-Me, N²,N²-di-de-Me: Lindoldhamine**
[60342-37-2]
C₃₄H₃₆N₂O₆ 568.668
Alkaloid from the leaves of *Lindera oldhamii*, the stems of *Albertisia papuana*, and from *Polyalthia nitidissima* (Lauraceae, Menispermaceae, Annonaceae). Fine needles (EtOH/Me₂CO/CHCl₃). Mp 183-186°. [α]_D³³ +35 (c, 1.0 in EtOH).
- O⁷,O⁷-Di-de-Me, O¹²-Me: Cuspidaline**
[10410-53-4]
C₃₇H₄₂N₂O₆ 610.749
Alkaloid from *Limacia cuspidata* and *Limacia oblonga* (Menispermaceae). Oil. [α]_D -48 (CHCl₃).
- O⁷,O⁷-Di-de-Me, O¹²-Me, N,N'-di-de-Me: Costaricine. 12-O-Methylindoldhamine**
[176772-29-5]
C₃₅H₃₈N₂O₆ 582.695
Alkaloid from trunk bark and roots of *Nectandra salicifolia*. Shows antiplasmodial activity. Yellow amorph. powder. [α]_D +46.4 (c, 0.248 in CHCl₃).
- Manske, R.H.F. et al., *Can. J. Res., Sect. B*, 1943, **21**, 17 (*Dauricine, isol*)
Inubushi, Y. et al., *Yakugaku Zasshi*, 1952, **72**, 762; *CA*, **47**, 6430d (*Dauricine, struct*)
Tomita, M. et al., *Chem. Pharm. Bull.*, 1955, **3**, 449 (*O-Methylauricine, synth*)
- Manske, R.H.F. et al., *Chem. Pharm. Bull.*, 1965, **13**, 1476 (*Dauricine, pmr*)
Tomita, M. et al., *Yakugaku Zasshi*, 1965, **85**, 456; 1967, **87**, 793; 1560; 1970, **90**, 1178; 1182 (*Cuspidaline, Daurinoline, Dauricoline, Dauricinoline*)
Johns, S.R. et al., *Aust. J. Chem.*, 1970, **23**, 363 (*O-Methylauricine, isol, pmr, ms, config*)
Tschesche, R. et al., *Phytochemistry*, 1970, **9**, 1683 (*O-Methylauricine, isol, uv, ir, pmr, ms*)
Doskotch, R.W. et al., *J. Nat. Prod.*, 1971, **34**, 292 (*Demethylauricine, Daurinoline*)
Baldas, J. et al., *J.C.S. Perkin I*, 1972, 592 (*ms, derivs*)
Kuroda, H. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2413-2420 (*Dauricine, activity*)
Lu, S.T. et al., *Heterocycles*, 1976, **4**, 1073 (*Lindoldhamine, uv, pmr, ms, struct*)
Zheng, X.W. et al., *CA*, 1979, **91**, 27216r (*Daurisoline*)
Jossang, A. et al., *Planta Med.*, 1983, **49**, 20 (*Dauricine, Lindoldhamine, O-Methylindoldhamines, Isodaurisoline*)
Dehaussy, H. et al., *Planta Med.*, 1983, **49**, 25 (*Guattegaumerine*)
El-Sebakhy, N. et al., *Phytochemistry*, 1984, **23**, 2706 (*7'-O-Methylcuspidaline*)
Jossang, A. et al., *J. Nat. Prod.*, 1986, **49**, 1018 (*Dauricine oxides, Popisonine, Popisidine, Popidine, Popisopine, Popisine*)
Cortes, D. et al., *J. Nat. Prod.*, 1987, **50**, 910 (*Pedroamine, Pampulhamine, Geraldoamine*)
Duté, P. et al., *Phytochemistry*, 1987, **26**, 2136 (*N-Methylindoldhamines, Nordaurisoline*)
Kong, R. et al., *Chin. J. Chem.*, 1991, **9**, 275; *CA*, **117**, 48977k (*Daurisoline, abs config*)
Pan, X.P. et al., *Yaoxue Xuebao*, 1991, **26**, 387; 1992, **27**, 788; *CA*, **115**, 131958; **118**, 165183e (*Daurilicine, N-Demethylauricine*)
Zeng, G.Q. et al., *Yaoxue Xuebao*, 1993, **14**, 329 (*Dauricine, pharmacol*)
Böhlke, M. et al., *J. Nat. Prod.*, 1996, **59**, 576 (*Costaricine*)
Pan, X.P. et al., *CA*, 2000, **132**, 134721p (*Dauricoline*)

Daurioxoisoporphine B D-89


6-Amino-4,5,9-trimethoxy-7H-dibenzo[4,5-f]quinolin-7-one



- C₁₉H₁₆N₂O₄ 336.346
Alkaloid from the rhizomes of *Menispermum dauricum*. Cytotoxic agent. Amorph. yellow powder. λ_{max} 219 (log ε 2.8); 255 (log ε 3.2); 350 (log ε 4.5); 430 (log ε 2.6); 455 (log ε 4.1) (MeOH).
- 4-Demethoxy, N⁶-Me: Daurioxoisoporphine C**
C₁₉H₁₆N₂O₃ 320.347
Alkaloid from the rhizomes of *Menispermum dauricum*. Amorph. yellow powder. λ_{max} 215 (log ε 4.32); 250 (log ε 3.42); 310 (log ε 2.43); 360 (log ε 2.42); 406 (log ε 3.34); 420 (log ε 4.33); 458 (log ε 3.43) (MeOH).

Yu, B.-W. et al., *J. Nat. Prod.*, 2001, **64**, 968-970

Dauriporphine **D-90**
4,5,6,9-Tetramethoxy-7H-dibenzo[de,h]-quinolin-7-one, 9CI. *Bianfugenine* [88142-60-3]




$C_{20}H_{17}NO_5$ 351.358
Alkaloid from the vines of *Menispermum dauricum* (Menispermaceae). Yellow needles (MeOH). Mp 167° (159-161.5°).

O⁶-De-Me: Dauriporphinoline
[100009-82-3]
 $C_{19}H_{15}NO_5$ 337.331
Alkaloid from the rhizomes of *Menispermum dauricum* (Menispermaceae).

2,3-Dihydro: 2,3-Dihydrodauriporphine
 $C_{20}H_{19}NO_5$ 353.374
Alkaloid from the roots of *Menispermum dauricum*. Yellow needles (hexane/EtOAc). Mp 141-142°. λ_{max} 219 (log ϵ 4.35); 271 (log ϵ 4.44); 329 (log ϵ 3.78) (no solvent reported).

Takani, M. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3091 (*isol, uv, ir, pmr, ms, struct*)
Kunitomo, J. et al., *Chem. Pharm. Bull.*, 1985, **33**, 2778 (*synth, uv, ir, pmr, ms*)
Hou, C. et al., *Yaoxue Xuebao*, 1985, **20**, 112; *CA*, **103**, 3691g
Zhao, S. et al., *Zhongguo Yaoke Daxue Xuebao*, 1989, **20**, 312; *CA*, **113**, 55823c (*Dauriporphinoline*)
Sugimoto, Y. et al., *Phytochemistry*, 1999, **52**, 1431-1435 (*Dauriporphine, 2,3-Dihydrodauriporphine*)

Davallioside A **D-91**
[131747-15-4]




$C_{25}H_{29}NO_{12}$ 535.504
Constit. of the rhizomes of *Davallia mariesii*. Amorph. solid. $[\alpha]_D^{23} +10.2$ (MeOH). λ_{max} 230 (sh) (log ϵ 4.37); 281

(log ϵ 3.79) (no solvent reported).

1''-Epimer: Davallioside B
[131831-19-1]
 $C_{25}H_{29}NO_{12}$ 535.504
Constit. of the rhizomes of *Davallia mariesii*. Amorph. solid. $[\alpha]_D^{23} -114.6$ (MeOH).

Cui, C.-B. et al., *Chem. Pharm. Bull.*, 1990, **38**, 2620 (*isol, pmr, cmr*)

N-Deacetylcolchicine **D-92**
Desacetylcolchicine



$C_{20}H_{23}NO_5$ 357.405
See also refs. given under Colchicine, C-569.

(S)-form [3476-50-4]
[102419-91-0]
Alkaloid from *Kreysigia multiflora*. Antineoplastic agent. Cryst. (EtOAc). Mp 125-127°. $[\alpha]_D -148$ (c, 1.1 in $CHCl_3$).

Tartrate: Cytostal. NSC 36354. NCI 1136. SKF 250
[49720-72-1]
[27963-65-1, 3145-41-3, 36191-19-2]
Mp 242-244° dec. $[\alpha]_D^{22} -176.2$ (c, 0.9 in H_2O). λ_{max} 245 (log ϵ 4.53); 353 (log ϵ 4.36) (no solvent reported).

N-Formyl: Gloriosine. N-Deacetyl-N-formylcolchicine. Substance B
[7411-12-3]
 $C_{21}H_{23}NO_6$ 385.416
Alkaloid from *Androcymbium melanthioides*, *Bulbocodium vernum*, *Camptorrhiza strumosa*, *Colchicum autumnale* and several other *Colchicum* spp., *Gloriosa superba* and some other *Gloriosa* spp., *Iphigenia bechuanica*, *Iphigenia indica*, *Iphigenia pallida*, *Kreysigia multiflora*, *Littonia modesta*, *Merendera caucasica* and other *Merendera* spp., *Ornithoglossum glaucum*, *Ornithoglossum viride* and *Sandersonia aurantiaca* (Liliaceae). Pale yellow cryst. (EtOAc/Et₂O). Mp 264-267° dec. $[\alpha]_D^{22} -171.2$ (c, 1.0863 in $CHCl_3$).

▶ GH0794000
N-Ac: see Colchicine, C-569
N-Me: **Demecolcine, BAN, INN. N-Methyldeacetylcolchicine. Colchamine. Colcemid. Kolkamin. Omain. Santavy's Substance F. Ciba 12669A. NSC 3096. Alkaloid F†**
[477-30-5]
 $C_{21}H_{25}NO_5$ 371.432

Alkaloid from *Colchicum speciosum*, several other *Colchicum* spp., *Merendera jolantae* (Liliaceae), *Merendera persica* and *Androcymbium melanthioides*. Antineoplastic and antimetabolic agent. Antitubulin agent which induces apoptosis. Mp 186°. $[\alpha]_D^{23} -129$

(c, 1.02 in $CHCl_3$). Log P 0.6 (calc). Colchicerine, Mp 187-187.5° from *C. speciosum* was a 1:1 molecular complex of (S)-Demecolcine and (S)-Colchicine. λ_{max} 245 (c 35500); 355 (c 17400) (MeOH). λ_{max} 234; 350 (EtOH).

▶ LD₅₀ (mus, ori) 25.5 mg/kg; LD₅₀ (mus, ipr) 35 mg/kg. Exp. reprod. and teratogenic effects. Mutagenic props. GH0800000

N-Me, N-formyl: **N-Formyldemecolcine. Alkaloid CC 19**

[14686-61-4]
 $C_{22}H_{25}NO_6$ 399.443
Alkaloid from the seeds of *Colchicum cornigerum* (Liliaceae). Pale yellow prisms (MeOH/Et₂O). $[\alpha]_D^{22} -189$ (c, 1.0192 in $CHCl_3$). $[\alpha]_D^{22} -202$ (c, 0.79 in EtOH).

▶ GH0795000

N-Me, N-Ac: **N-Methylcolchicine**
 $C_{23}H_{27}NO_6$ 413.469
Mp 199-200° Mp 233-235° (double Mp).

N-(3-Oxobutanoyl): **N-Deacetyl-N-3-oxobutyrylcolchicine. N-Acetoacetyl-N-deacetylcolchicine**
 $C_{24}H_{27}NO_7$ 441.48
Minor alkaloid from the seeds of *Colchicum autumnale* (Liliaceae). Cryst. (EtOAc/Et₂O). Mp 222-223°. $[\alpha]_D^{25} -194$ (c, 1.07 in $CHCl_3$).

N-Me, N-ethoxycarbonyl: **N-Ethoxycarbonyldemecolcine**
 $C_{24}H_{29}NO_7$ 443.496
Alkaloid from *Colchicum cilicicum* (Liliaceae).

N,N-Di-Me: **N-Methyl-demecolcine. Alkaloid CC 5**
[7336-44-9]
 $C_{22}H_{27}NO_5$ 385.459
Alkaloid from seeds, leaves, stems and flowers of *Colchicum cornigerum* (Liliaceae) and *Kreysigia multiflora*. Pale yellow prisms (EtOAc/Et₂O). Mp 208-210° (205-208°). $[\alpha]_D^{22} -104$ (c, 0.678 in $CHCl_3$). $[\alpha]_D^{20} -111$ (c, 1.78 in $CHCl_3$).

▶ GH0792000

O²-De-Me, N-formyl: **N-Deacetyl-2-demethyl-N-formylcolchicine**
[57866-22-5]
 $C_{20}H_{21}NO_6$ 371.389
Alkaloid from the leaves and flowers of *Colchicum autumnale* (Liliaceae). Noncryst. λ_{max} 243 (log ϵ 4.46); 355 (log ϵ 4.2) (EtOH). λ_{max} 247 (log ϵ 4.45); 290 (log ϵ 3.86); 356 (log ϵ 4.19) (EtOH/NaOH).

O²-De-Me, N-Ac: **2-Demethylcolchicine. 2-Desmethylcolchicine. Alkaloid L5†. Alkaloid CC 6. Substance E₁**
[7336-36-9]
 $C_{21}H_{23}NO_6$ 385.416
Alkaloid from *Androcymbium gramineum*, *Androcymbium melanthioides*, *Colchicum autumnale* and some other *Colchicum* spp., *Bulbocodium vernum*, *Gloriosa superba*, *Littonia modesta* and *Merendera robusta* (Liliaceae). Cryst. ($CHCl_3$). Mp 110-140° Mp 178-180° (double Mp). Does not occur in fresh plant extracts. Present only in dried

material, attached in an unknown manner to cell constituents, and liberated enzymatically. λ_{\max} 243 (log ϵ 4.46); 355 (log ϵ 4.2) (EtOH). λ_{\max} 247 (log ϵ 4.45); 290 (sh) (log ϵ 3.86); 356 (log ϵ 4.19) (EtOH/NaOH).

▶ GH0880000

O²-De-Me, N-Me: **2-Demethyldemecolcicine**. Substance S. Alkaloid CC 7 [7359-91-3]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Colchicum autumnale*, *Colchicum hierosolymitanum*, *Colchicum speciosum*, *Colchicum latifolium*, other *Colchicum* spp., *Merendera persica*, *Merendera robusta* (Liliaceae) and *Androcymbium melanthioides*. Pale yellow plates (MeOH/Et₂O). Mp 136-138°. $[\alpha]_{\text{D}}^{22}$ -119 (c, 1.00 in CHCl₃). λ_{\max} 241 (log ϵ 4.49); 355 (log ϵ 4.19) (EtOH). λ_{\max} 244 (log ϵ 4.6); 287 (log ϵ 3.96); 355 (log ϵ 4.13) (EtOH/NaOH).

O³-De-Me: **3-Demethyl-N-deacetylcolchicine**. Substance U

C₁₉H₂₁NO₅ 343.379

Alkaloid from the corms, seeds and flowers of *Colchicum autumnale* (Liliaceae). The phys. props. of the free base have not been reported.

O³-De-Me, N-formyl: **N-Deacetyl-3-demethyl-N-formylcolchicine**

[18172-26-4]

C₂₀H₂₁NO₆ 371.389

Alkaloid from the seeds of *Colchicum latifolium*, *Colchicum autumnale*, *Colchicum luteum*, also occurs in *Gloriosa superba* (Liliaceae). Cryst. (EtOAc). Mp 263-267°. $[\alpha]_{\text{D}}^{25}$ -180 (c, 0.30 in CHCl₃). λ_{\max} 232 (log ϵ 4.48); 245 (sh) (log ϵ 4.44); 357 (log ϵ 4.19) (EtOH). λ_{\max} 225 (log ϵ 4.52); 245 (log ϵ 4.51); 279 (log ϵ 3.99); 374 (log ϵ 4) (EtOH/NaOH).

O³-De-Me, N-Ac, O³- α -D-glucopyranoside: **1'-Epicolchicoside**

[404866-95-1]

C₂₇H₃₃NO₁₁ 547.558

Alkaloid from the seeds of *Gloriosa superba*. Mp 188-190°. $[\alpha]_{\text{D}}^{25}$ -317 (c, 0.88 in H₂O). Not named in the lit. λ_{\max} 243 (log ϵ 4.74); 355 (log ϵ 4.65) (MeOH).

O³-De-Me, N-Ac, O³- β -D-glucopyranoside: **Colchicoside**

[477-29-2]

C₂₇H₃₃NO₁₁ 547.558

Alkaloid from seeds of *Colchicum autumnale* and *Gloriosa superba* (Liliaceae). Antiinflammatory agent. Cryst. (EtOH). Mp 192-195° Mp 216-218° (block). Log P -3.61 (uncertain value) (calc). λ_{\max} 242 (ϵ 32200); 346 (ϵ 19400) (EtOH).

▶ GH0982000

O³-De-Me, N-Ac: **3-Demethylcolchicine**. Substance C

[7336-33-6]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Androcymbium gramineum*, *Androcymbium melanthioides*, *Colchicum autumnale* and several other *Colchicum* spp., *Bulbocodium vernum*, *Gloriosa superba* and some other

Gloriosa spp., *Merendera caucasica* and other *Merendera* spp., *Littonia modesta*, *Ornithoglossum glaucum* var.

grandiflorum, *Ornithoglossum viride* and *Sandersonia aurantiaca* (Liliaceae). Antineoplastic agent. Prisms (CHCl₃). Mp 176-182° (180-190 and 275-280°, double Mp). $[\alpha]_{\text{D}}^{22}$ -130.7 (c, 2.0118 in CHCl₃). Log P -0.45 (uncertain value) (calc). Does not occur in fresh plant extracts. Present only in dried material, attached in an unknown manner to cell constituents, and liberated enzymatically. λ_{\max} 243 (log ϵ 4.46); 355 (log ϵ 4.19) (EtOH). λ_{\max} 251 (log ϵ 4.46); 286 (log ϵ 3.86); 340 (sh) (log ϵ 3.86); 395 (log ϵ 4.06) (EtOH/NaOH).

▶ GH0880500

O³-De-Me, O³, N-di-Ac:

Fine needles (EtOAc/Et₂O). Mp 227-228°. $[\alpha]_{\text{D}}^{20}$ -93 (c, 0.822 in CHCl₃).

O³-De-Me, N-Me: **3-Demethyldemecolcicine**

[2034-50-6]

C₂₀H₂₃NO₅ 357.405

Alkaloid from the corms and leaves of *Colchicum cornigerum* (Liliaceae), other *Colchicum* spp. and *Androcymbium melanthioides*. Mp 220-222°. $[\alpha]_{\text{D}}^{22}$ -128 (c, 0.888 in CHCl₃). λ_{\max} 244 (log ϵ 4.47); 356 (log ϵ 4.2) (EtOH). λ_{\max} 235 (log ϵ 4.44); 250 (log ϵ 4.47); 291 (log ϵ 3.87); 318 (sh) (log ϵ 3.82); 394 (log ϵ 4.01) (EtOH/NaOH).

O³-De-Me, N,N-di-Me: **3-Demethyl-N-methyldemecolcicine**

[929683-21-6]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Colchicum tauri*.

O¹⁰-De-Me: **N-Deacetylcolchicine**. *De-sacetylcolchicine*

[3482-37-9]

C₁₉H₂₁NO₅ 343.379

Alkaloid from the above-ground parts of *Merendera robusta* (Liliaceae).

Cryst. + $\frac{1}{2}$ H₂O (EtOH aq.). Mp 155° (151-153°). $[\alpha]_{\text{D}}$ -185 (c, 1 in CHCl₃).

▶ GH0960000

O¹⁰-De-Me, N-Ac: **Colchicine**. **10-Demethylcolchicine**

[477-27-0]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Colchicum autumnale*, *Colchicum kesselringii*, *Colchicum luteum*, *Merendera robusta* and *Merendera jolantae* (Liliaceae). Used as 0.02M CHCl₃ soln. as an extractive indicator in the EDTA titration of Cu(II) (pH 4.5). Needles (EtOAc). Sol. CHCl₃. Mp 175-177°. $[\alpha]_{\text{D}}^{20}$ -256 (c, 1.005 in CHCl₃). λ_{\max} 243 (log ϵ 4.47); 352 (log ϵ 4.24) (EtOH). λ_{\max} 241 (log ϵ 4.49); 259 (sh) (log ϵ 4.4); 352 (log ϵ 4.24); 367 (sh) (log ϵ 4.29); 403 (log ϵ 4.04) (EtOH/NaOH).

▶ GH0520000

O¹⁰-De-Me, O¹⁰, N-di-Ac: Mp 122-124°. $[\alpha]_{\text{D}}^{20}$ -260 (c, 0.759 in CHCl₃).

O¹⁰-De-Me, N-Me: **Demecolcicine**. **N-Deacetyl-N-methylcolchicine**. *Colchameine*. Substance T_a

[518-11-6]

C₂₀H₂₃NO₅ 357.405

Alkaloid from the corms of *Colchicum autumnale*, and from *Merendera robusta* and *Merendera jolantae* (Liliaceae). Yellow cryst. (MeOH). Mp 133-135°. $[\alpha]_{\text{D}}^{20}$ -211 (c, 0.72 in CHCl₃). $[\alpha]_{\text{D}}$ -228 (c, 1.24 in CHCl₃).

▶ GH0600000

1,2-Di-O-de-Me, N-Ac: **1,2-Didemethylcolchicine**

[78231-83-1]

C₂₀H₂₁NO₆ 371.389

Alkaloid from seeds of *Gloriosa superba* (Liliaceae). Amorph. yellow solid. Mp 238°. $[\alpha]_{\text{D}}$ -148 (c, 0.167 in DMF).

2,3-Di-O-de-Me: **N-Deacetyl-2,3-didemethylcolchicine**. **2,3-Didemethyl-N-deacetylcolchicine**

[57866-20-3]

C₁₈H₁₉NO₅ 329.352

Alkaloid from *Gloriosa superba* (Liliaceae). λ_{\max} 245 ; 355 (log ϵ 4.15) (EtOH). λ_{\max} 240 ; 258 ; 297 ; 356 (sh) ; 405 (EtOH/NaOH).

2,3-Di-O-de-Me, N-Ac: **2,3-Didemethylcolchicine**

[57866-21-4]

C₂₀H₂₁NO₆ 371.389

Alkaloid from the seeds of *Colchicum latifolium* and from *Gloriosa superba* (Liliaceae). Amorph. Mp 205°. $[\alpha]_{\text{D}}$ -230 (c, 0.23 in MeOH).

2,3-Di-O-de-Me, N-Me: **2,3-Didemethyl-demecolcicine**

C₁₉H₂₁NO₅ 343.379

Alkaloid from *Colchicum brachyphyllum*. Yellowish powder. $[\alpha]_{\text{D}}^{25}$ -90 (c, 0.1 in MeOH). λ_{\max} 241 (ϵ 11800) (MeOH).

O², O¹⁰-Di-de-Me, N-Ac: **2-Demethylcolchicine**

[33530-04-0]

C₂₀H₂₁NO₆ 371.389

Alkaloid from the corms of *Gloriosa superba* and epigeal parts of *Merendera robusta* (Liliaceae). Cryst. (EtOH). Mp 259-261°. $[\alpha]_{\text{D}}$ -230 (c, 2.1 in CHCl₃). λ_{\max} 236 (log ϵ 4.78); 280 (sh) (log ϵ 4.1); 343 (log ϵ 4.49); 380 (log ϵ 4.35) (EtOH/NaOH).

O³, O¹⁰-Di-de-Me, N-Ac: **3-Demethylcolchicine**. Alkaloid L5†

[7336-34-7]

C₂₀H₂₁NO₆ 371.389

Alkaloid from *Colchicum luteum* and from the corms of *Gloriosa superba* (Liliaceae). Amorph. Mp 179-183°. λ_{\max} 242 (log ϵ 4.71); 258 (log ϵ 4.64); 266 (log ϵ 4.49); 302 (log ϵ 3.94); 368 (log ϵ 4.46); 397 (log ϵ 4.45) (EtOH/NaOH).

O³, O¹⁰-Di-de-Me, di-Ac:

Cryst. (EtOAc/Et₂O). Mp 119-122°.

$[\alpha]_{\text{D}}$ -170 (c, 0.64 in CHCl₃).

(±)-form [102491-73-6]

Cryst. (MeOH/Et₂O). Mp 158-160°.

N-Me: [102491-77-0]

Cryst. (MeOH/Et₂O). Mp 178-179°.

O¹⁰-De-Me: [73307-44-5]

Synthetic. Yellow cryst. (EtOH). Mp 244-246°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 874A (nmr)

Šantavý, F. et al., *Coll. Czech. Chem. Comm.*, 1950, **15**, 552-569; 1953, **18**, 710-716; 1954, **19**, 141-152; 805-816; 1289-1301 (Demethylcolchicines, isol, struct)

Šantavý, F. et al., *Pharm. Acta Helv.*, 1950, **25**, 248 (Demecolcine, isol)

Kiselev, V.V. et al., *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1952, **87**, 227-228; *CA*, **48**, 695i (Colchicerine)

Šantavý, F. et al., *Helv. Chim. Acta*, 1953, **36**, 1319-1324; 1954, **37**, 18-34 (struct, ir)

Fabian, J. et al., *Bull. Soc. Chim. Fr.*, 1955, 1455-1463 (Colchicoside)

Corradi, H. et al., *Helv. Chim. Acta*, 1957, **40**, 193-199 (Colchiceine, resoh)

Santavý, F. et al., *Coll. Czech. Chem. Comm.*, 1959, **24**, 2237-2246 (2-Demethyl-demecolcine)

Hrbek, J. et al., *Coll. Czech. Chem. Comm.*, 1962, **27**, 255-267 (Gloriosine)

Saleh, M. et al., *Coll. Czech. Chem. Comm.*, 1963, **28**, 3413-3423 (N-Methyl-demecolcine, 3-Demethyl-demecolcine)

Wilson, J.M. et al., *Tetrahedron*, 1963, **19**, 2225-2231 (ms)

Potěšilová, H. et al., *Coll. Czech. Chem. Comm.*, 1969, **34**, 2128-2133; 3540-3552 (2,3-Didemethylcolchicine, N-Formyl-demecolcine)

Severini-Ricca, G. et al., *Gazz. Chim. Ital.*, 1969, **99**, 133-151 (pmr)

Chommadov, B.Ch. et al., *Khim. Prir. Soedin.*, 1970, **6**, 82-88; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 77-82 (3-Demethylcolchicine)

Zweig, M.H. et al., *J. Pharmacol. Exp. Ther.*, 1972, **182**, 344-350 (pharmacol)

Turdikulov, K. et al., *Khim. Prir. Soedin.*, 1972, **8**, 247-248; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 247-248 (N-Deacetylcolchicine, N-Deacetylcolchiceine, isol, uv)

Kupchan, S.M. et al., *J. Nat. Prod.*, 1973, **36**, 338-340 (3-Demethylcolchicine, isol, pmr, ms)

Margulis, T.N. et al., *J.A.C.S.*, 1974, **96**, 899-902 (Demecolcine, cryst struct)

Thakur, R.S. et al., *Planta Med.*, 1975, **28**, 201-209 (N-Deacetylcolchicine-2,3-didemethylcolchicine)

Potesilová, H. et al., *Coll. Czech. Chem. Comm.*, 1976, **41**, 3146-3156 (N-Deacetyl-3-demethyl-N-formylcolchicine, 2,3-Didemethylcolchicine)

Silverton, J.V. et al., *Acta Cryst. B*, 1979, **35**, 2800-2803 (Colchiceine, cryst struct)

Capraro, H.-G. et al., *Helv. Chim. Acta*, 1979, **62**, 965-970 (Demecolcine, synth)

Hufford, C.D. et al., *J. Pharm. Sci.*, 1979, **68**, 1239-1243 (ir, pmr, cmr)

Malichová, V. et al., *Planta Med.*, 1979, **36**, 119-127 (Demethylcolchicines, N-deacetyl-3-demethyl-N-formylcolchicine)

Hufford, C.D. et al., *Helv. Chim. Acta*, 1980, **63**, 50-56 (cmr)

Rösner, M. et al., *J. Med. Chem.*, 1981, **24**, 257-261 (Colchiceine, Demethylcolchicines, synth)

Evans, D.A. et al., *J.A.C.S.*, 1981, **103**, 5813-5821 (Colchiceine, synth, ir, pmr)

Rösner, M. et al., *J.O.C.*, 1981, **46**, 3686-3688 (2,3-Didemethylcolchicine, synth)

Hrbek, J. et al., *Coll. Czech. Chem. Comm.*, 1982, **47**, 2258-2279 (cd)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 1829

Šantavý, F. et al., *Coll. Czech. Chem. Comm.*, 1983, **48**, 2989-2993 (N-Deacetyl-N-3-oxobutyrylcolchicine)

Kaushik, N.K. et al., *Microchem. J.*, 1985, **31**, 329-331 (Colchiceine, detn, Cu)

Dumont, R. et al., *J.O.C.*, 1986, **51**, 2515-2521 (racemates, unnatural enantiomers, 2-demethyl derivs, synth)

Yoshida, K. et al., *Agric. Biol. Chem.*, 1988, **52**, 593-594 (Colchicoside, isol, ir, pmr, cmr, struct)

Chommadov, B. et al., *Khim. Prir. Soedin.*, 1991, **27**, 67-71; *Chem. Nat. Compd. (Engl. Transl.)*, 1991, **27**, 58-61 (2-Demethylcolchiceine)

Chaudhuri, P.K. et al., *J. Nat. Prod.*, 1993, **56**, 1174-1176 (1,2-Didemethylcolchicine)

Sherwood, S.W. et al., *Exp. Cell Res.*, 1994, **215**, 373 (Demecolcine, pharmacol)

Lebeau, L. et al., *Synth. Commun.*, 1997, **27**, 293-296 (synth, S-form)

Suri, O.P. et al., *Nat. Prod. Lett.*, 2001, **15**, 217-219 (1'-Epicolchicoside)


Alali, F.Q. et al., *J. Nat. Prod.*, 2005, **68**, 173-178 (2,3-Didemethyl-demecolcine)

Alali, F.Q. et al., *Nat. Prod. Commun.*, 2006, **1**, 95-99 (3-Demethyl-N-methyl-demecolcine)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, DAN300; ADE000; DAN375; MIW500

N-Deacetylkuanoniamine D D-93

[202932-14-7]



$C_{18}H_{14}N_4S$ 318.401
Alkaloid from the sponge *Oceanapia* sp. Cytotoxic agent. DNA intercalator. Amorph. orange powder. λ_{max} 201 (log ϵ 4.14); 239 (log ϵ 4.33); 264 (log ϵ 4.11); 295 (log ϵ 4.08); 304 (log ϵ 4.05); 342 (log ϵ 3.82); 355 (log ϵ 3.85) (MeOH). λ_{max} 201 (ϵ 13800); 223 (ϵ 26900); 295 (ϵ 12000); 342 (ϵ 6600); 365 (ϵ 3550) (MeOH) (Berdy).

N²-Ac: Kuanoniamine D

[133401-13-5]
 $C_{20}H_{16}N_4OS$ 360.439
Alkaloid from a tunicate and its mollusc predator *Chelynotus semperi*. Also isol. from a purple tunicate tentatively identified as *Cystodytes* sp. Topoisomerase II inhibitor, chelating agent., adenosine and GABA receptor binding agent; cytotoxic agent; insecticide, toxic to brine shrimp. Amorph. yellow powder (CHCl₃). Mp 300°. λ_{max} 206 (ϵ 14100); 240 (ϵ 16600); 270 (ϵ 11700); 306 (ϵ 18600); 360 (ϵ 5620); 526 (ϵ 2450) (MeOH/HCl) (Derep). λ_{max} 206 (ϵ 17400); 240 (ϵ 20900); 264 (ϵ 15800); 294 (sh); 344 (ϵ 7590); 358 (ϵ 7410); 452 (ϵ 2400) (MeOH) (Derep). λ_{max} 206; 240; 270; 360; 526 (MeOH/NaOH) (Berdy).

N²-Propanoyl: Kuanoniamine C. Dercitamide

[133401-12-4]
 $C_{21}H_{18}N_4OS$ 374.465
Alkaloid from a tunicate and its mollusc predator *Chelynotus semperi*.

Also isol. from a purple tunicate tentatively identified as *Cystodytes* sp. and from a deep water marine sponge *Stelletta* sp. Cytotoxic. Inhibits proliferation of P388 murine leukaemia cells in vitro. Also displays immunosuppressive activity. Amorph. yellow powder (CHCl₃). Mp 300° (192°). λ_{max} 206 (ϵ 14100); 240 (ϵ 16600); 270 (ϵ 11700); 306 (ϵ 18600); 360 (ϵ 5620); 526 (ϵ 2450) (MeOH/HCl) (Derep). λ_{max} 245 (ϵ 13800); 307 (ϵ 16900); 361 (ϵ 3900); 541 (ϵ 1800) (MeOH) (Derep). λ_{max} 206 (ϵ 17400); 240 (ϵ 20900); 264 (ϵ 15800); 294 (sh); 344 (ϵ 7590); 358 (ϵ 7410); 452 (ϵ 2400) (MeOH) (Derep).

N²-(2-Methylpropanoyl): Kuanoniamine E

[445471-59-0]
 $C_{22}H_{20}N_4OS$ 388.492
Alkaloid from a Singaporean ascidian. Yellow gum. λ_{max} 204 (log ϵ 4.27); 241 (log ϵ 4.52); 265 (log ϵ 4.3); 296 (sh) (log ϵ 3.97); 346 (log ϵ 4.07); 362 (log ϵ 4.08); 452 (log ϵ 3.67) (MeOH).

N²-(2-Methylbutanoyl): Kuanoniamine F

[445471-61-4]
 $C_{23}H_{22}N_4OS$ 402.519
Alkaloid from a Singaporean ascidian. Yellow gum. $[\alpha]_D^{+105}$ (c, 0.18 in MeOH). λ_{max} 206 (log ϵ 4.26); 242 (log ϵ 4.49); 262 (log ϵ 4.29); 296 (sh) (log ϵ 3.98); 346 (log ϵ 4.04); 363 (log ϵ 4.05); 455 (log ϵ 3.38) (MeOH).

N²-(3-Methylbutanoyl): Kuanoniamine B

[133401-11-3]
 $C_{23}H_{22}N_4OS$ 402.519
Alkaloid from a tunicate and its mollusc predator *Chelynotus semperi*. Cytotoxic agent. Amorph. yellow powder (CHCl₃). Mp 300°. λ_{max} 204 (ϵ 17400); 240 (ϵ 20900); 264 (ϵ 15100); 294 (sh) (ϵ 7760); 344 (ϵ 7410); 360 (ϵ 7410); 450 (ϵ 2750) (MeOH) (Derep).

N²-(3-Methyl-2-butenoyl): Dehydrokuanoniamine B

[158734-24-8]
 $C_{23}H_{20}N_4OS$ 400.503
Alkaloid from *Cystodytes* sp. Antitumour agent; DNA intercalator; topoisomerase II inhibitor. Orange solid. λ_{max} 237 (ϵ 50970); 261 (ϵ 30120); 344 (ϵ 16410); 359 (ϵ 16950); 452 (ϵ 6050) (MeOH).

N²-Me: Dercitamine

[125236-60-4]
 $C_{19}H_{16}N_4S$ 332.428
Alkaloid from *Stelletta* sp. Inhibits proliferation of P388 murine leukaemia cells in vitro. Also displays immunosuppressive activity. Orange solid. Mp 135°. Regiochem. revised in 1992. Genus name incorr. given as *Stelletta*. λ_{max} 245 (ϵ 13800); 307 (ϵ 16900); 361 (ϵ 3900) (MeOH) (Derep).

N²,N²-Di-Me: Nordercitin

[125236-59-1]
 $C_{20}H_{18}N_4S$ 346.455
Alkaloid from a deep water marine sponge *Stelletta* sp. Inhibits proliferation of P388 murine leukaemia cells in vitro. Also displays immunosuppres-

sive activity. Yellow solid. Mp 176°. λ_{\max} 245 (ϵ 13800); 307 (ϵ 16900); 361 (ϵ 3900) (MeOH) (Derep).

Gunawardana, G.P. *et al.*, *Tet. Lett.*, 1989, **30**, 4359-4362 (*Dercitamine*, *Nordercitin*)

Carroll, A.R. *et al.*, *J.O.C.*, 1990, **55**, 4426-4431 (*Kuanoniamines A-D*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Bishop, M.J. *et al.*, *J.A.C.S.*, 1992, **114**, 10081-10082 (*synth*)

Gunawardana, G.P. *et al.*, *J.O.C.*, 1992, **57**, 1523-1526 (*Kuanoniamine D*, *Dercitamine*, *Nordercitin*, *struct*)

Eldredge, G.S. *et al.*, *J. Med. Chem.*, 1994, **37**, 3819-3827 (*Dehydrokuanoniamine B*)

Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1995, **117**, 12460-12469 (*synth*)

Eder, C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 301-305 (*Deacetylkuanoniamine D*)


Salomon, C.E. *et al.*, *Mar. Biol. (Berlin)*, 2001, **139**, 313 (*biosynth*)

Nilar, *et al.*, *J. Nat. Prod.*, 2002, **65**, 1198-1200 (*Kuanoniamines E-F*)

Sidebottom, N.P.J. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1198-1200 (*Kuanoniamine E,F*)

O-Deacetyl-1,2,4,5-tetrahydro-2-hydroxy-1-methylkuanoniaminium(1+)

[870717-13-8 (chloride)]




$C_{22}H_{27}N_2O_4^+$ 383.466

Quaternary alkaloid from the leaves of *Alstonia scholaris*. Powder (as chloride). Mp 229-232° (chloride). $[\alpha]_D^{18}$ -80.1 (c, 0.2 in $CHCl_3$) (chloride).

Zhou, H. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 2508-2512 (*isol*, *pmr*, *cmr*)

Deacylbengazole C



Absolute Configuration

$C_{13}H_{18}N_2O_7$ 314.294

O^{10} -Tridecanoyl: **Bengazole C**

[147512-35-4]

$C_{26}H_{42}N_2O_8$ 510.626

Isol. from a *Jaspis* sp. Anthelmintic agent. Oil. λ_{\max} 217 (MeOH aq.).

O^{10} -Tetradecanoyl: **Bengazole A**

[112549-08-3]

$C_{27}H_{44}N_2O_8$ 524.653

Isol. from a *Jaspidae* marine sponge from Fiji. Exhibits anthelmintic activity. Antifungal agent. Viscous oil. $[\alpha]_D^{20}$ +5 (c, 0.107 in MeOH). λ_{\max} 209 (ϵ 1400) (MeOH) (Derep).

O^{10} -(12-Methyltridecanoyl): **Bengazole D**

[147362-22-9]

$C_{27}H_{44}N_2O_8$ 524.653

Isol. from a *Jaspis* sp. Anthelmintic agent. Oil. λ_{\max} 217 (MeOH aq.).

O^{10} -Pentadecanoyl: **Bengazole E**

[147362-23-0]

$C_{28}H_{46}N_2O_8$ 538.68

Isol. from a *Jaspis* sp. Anthelmintic agent. Oil. λ_{\max} 217 (MeOH aq.).

O^{10} -(12-Methyltetradecanoyl): **Bengazole B₁**

[147391-82-0]

$C_{28}H_{46}N_2O_8$ 538.68

Isol. from a Great Barrier Reef sponge *Jaspis* sp. Inseparable 1:5 mixt. with Bengazole B.

O^{10} -(13-Methyltetradecanoyl): **Bengazole B**

[112549-09-4]

$C_{28}H_{46}N_2O_8$ 538.68

Isol. from a *Jaspidae* marine sponge from Fiji and from an Australian *Jaspis* sp. (as an inseparable mixt. with Bengazole B₁). Exhibits anthelmintic and antifungal activity. Viscous oil. $[\alpha]_D^{20}$ +4.7 (c, 0.024 in MeOH). λ_{\max} 209 (ϵ 1400) (MeOH) (Derep).

O^{10} -Hexadecanoyl: **Bengazole G**

[147362-24-1]

$C_{29}H_{48}N_2O_8$ 552.707

Isol. from a *Jaspis* sp. Anthelmintic agent. Oil. λ_{\max} 217 (MeOH aq.).

O^{10} -(13-Methylpentadecanoyl): **Bengazole F**

[147391-83-1]

$C_{29}H_{48}N_2O_8$ 552.707

Isol. from a *Jaspis* sp. Anthelmintic agent. Oil. λ_{\max} 217 (MeOH aq.).

10-Deoxy: see Bengazole Z, B-61

Adamczeski, M. *et al.*, *J.A.C.S.*, 1988, **110**, 1598 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Molinski, T.F. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1 (*biochem*)

Searle, P.A. *et al.*, *J.O.C.*, 1996, **61**, 4073-4079 (*abs config*)


Mulder, R.J. *et al.*, *J.O.C.*, 1999, **64**, 4995-4998; 2000, **65**, 8126 (*synth*)

Chittari, P. *et al.*, *Heterocycles*, 2003, **59**, 465-472 (*synth*)

Bull, J.A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 5515-5538 (*Bengazole A, B*, *synth*)

Deacylconvolutamide

3-(3,4-Dibromo-5-hydroxyphenyl)-3-hydroxy-2-pyrrolidinone



$C_{10}H_9Br_2NO_3$ 350.994

N-Tetradecanoyl: **Convolutamide A**

[158182-21-9]

$C_{24}H_{35}Br_2NO_4$ 561.353

Isol. from the marine bryozoan *Amathia convoluta*. Exhibits cytotoxicity against L1210 murine leukaemia and KB human epidermoid carcinoma cells. Amorph. solid. $[\alpha]_D^{20}$ -6 (c, 0.4 in

$CHCl_3$). Opt. rotn. and biol. activity refer to a 1:1.7 inseparable mixt. with Convolutamide B. λ_{\max} 225 (ϵ 34000) (MeOH) (Berdy).

N-(9Z-Hexadecenoyl): **Convolutamide B**

[158182-22-0]

$C_{26}H_{37}Br_2NO_4$ 587.391

From *Amathia convoluta*. See statement under Convolutamide A above.

N-Hexadecenoyl: **Convolutamide C**

[158182-23-1]

$C_{26}H_{39}Br_2NO_4$ 589.406

From *Amathia convoluta*. Shows no cytotoxicity. Amorph. solid. $[\alpha]_D^{20}$ -5.1 (c, 0.4 in $CHCl_3$). Opt. rotn. refers to a 1.8:1 inseparable mixt. with Convolutamide D.

N-(9Z-Octadecenoyl): **Convolutamide D**

[158182-24-2]

$C_{28}H_{41}Br_2NO_4$ 615.444

From *Amathia convoluta*. See statement under Convolutamide C above.

N-Octadecenoyl: **Convolutamide E**

[158182-25-3]

$C_{28}H_{43}Br_2NO_4$ 617.46

From *Amathia convoluta*. Shows no cytotoxicity. Amorph. solid. $[\alpha]_D^{20}$ -25 (c, 0.1 in $CHCl_3$). Opt. rotn. refers to a 7.9:1 inseparable mixt. with Convolutamide F.

N-(8-Eicosenoyl): **Convolutamide F**


[158182-26-4]

$C_{30}H_{45}Br_2NO_4$ 643.498

From *Amathia convoluta*. See statement under Convolutamide E above. Geometry of side chain double bond not assigned.

Zhang, H. *et al.*, *Tetrahedron*, 1994, **50**, 10201-10206 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Deacylcystodytin



$C_{17}H_{13}N_3O$ 275.309

N²-Ac: **Cystodytin J**

[158734-25-9]

$C_{19}H_{15}N_3O_2$ 317.346

Isol. from a *Cystodytes* sp. Cytotoxic agent. Inhibitor of topoisomerase II. DNA intercalator. Yellow solid. λ_{\max} 218 (ϵ 12414); 277 (ϵ 6033); 300 (ϵ 7491); 318 (ϵ 8123); 359 (ϵ 1708); 382 (ϵ 1716); 538 (ϵ 1423) (MeOH) (Berdy).

N²-(3-Methyl-2-butenoyl): **Cystodytin A**

[113321-71-4]

$C_{22}H_{19}N_3O_2$ 357.411

Isol. from *Cystodytes dellechiaiei*. Topoisomerase II inhibitor. Calcium release agent. Yellow cryst. Sol. MeOH, toluene, $CHCl_3$; poorly sol. H_2O . Mp 181-183°. Inseparable mixt. with Cystodytin B. λ_{\max} 214 (ϵ 37000); 274 (ϵ 30000); 384 (ϵ 11000) (MeOH) (De-

rep). λ_{\max} 226 (€ 35000); 272 (€ 26000); 380 (€ 11500) (MeOH) (Derep).

$N^{2'}-(3\text{-Hydroxy-3-methylbutanoyl})$:

Cystodytin C

[113351-76-1]

$C_{22}H_{21}N_3O_3$ 375.426

Isol. from *Cystodytes dellechiaiei*. Calcium release agent. Light yellow cryst. Sol. MeOH, $CHCl_3$, toluene; poorly sol. H_2O . Mp 257-259°. λ_{\max} 214 (€ 37000); 274 (€ 30000); 384 (€ 11000) (MeOH) (Derep). λ_{\max} 226 (€ 35000); 272 (€ 26000); 380 (€ 11500) (MeOH) (Derep). λ_{\max} 228 (€ 29900); 272 (€ 29100); 380 (€ 11800) (MeOH) (Berdy).

$N^{2'}\text{-Tigloyl}$: **Cystodytin B**

[113351-75-0]

$C_{22}H_{19}N_3O_2$ 357.411

Isol. from *Cystodytes dellechiaiei*. Shows calcium release activity. Sol. MeOH, $CHCl_3$, toluene; poorly sol. H_2O . Obt. only in admixture with Cystodytin A. λ_{\max} 214 (€ 37000); 274 (€ 30000); 384 (€ 11000) (MeOH) (Derep). λ_{\max} 226 (€ 35000); 272 (€ 26000); 380 (€ 11500) (MeOH) (Derep).

$1'\text{-Hydroxy}$, $N^{2'}-(3\text{-methyl-2-butanoyl})$:

Cystodytin D

[141544-60-7]

$C_{22}H_{19}N_3O_3$ 373.41

Isol. from *Cystodytes dellechiaiei*. Sol. MeOH; poorly sol. hexane, H_2O . λ_{\max} 214 (€ 37000); 274 (€ 30000); 384 (€ 11000) (MeOH) (Derep). λ_{\max} 226 (€ 35000); 272 (€ 26000); 380 (€ 11500) (MeOH) (Derep).

$1'\text{-Hydroxy}$, $N^{2'}\text{-tigloyl}$: **Cystodytin E**

[141544-61-8]

$C_{22}H_{19}N_3O_3$ 373.41

Alkaloid from *Cystodytes dellechiaiei*. λ_{\max} 214 (€ 37000); 274 (€ 30000); 384 (€ 11000) (MeOH) (Derep). λ_{\max} 226 (€ 35000); 272 (€ 26000); 380 (€ 11500) (MeOH) (Derep).

$1'\text{-}(9\text{-Octadecenoyloxy})$, $N^{2'}-(3\text{-methyl-2-butanoyl})$: **Cystodytin H**

[141544-64-1]

$C_{40}H_{51}N_3O_4$ 637.861

Alkaloid from *Cystodytes dellechiaiei*. Sol. MeOH; poorly sol. hexane, H_2O . λ_{\max} 225 (€ 37000); 273 (€ 25000); 382 (€ 10000) (MeOH) (Derep).

$1'\text{-}(9\text{-Octadecenoyloxy})$, $N^{2'}\text{-tigloyl}$:

Cystodytin I

[141657-37-6]

$C_{40}H_{51}N_3O_4$ 637.861

Alkaloid from *Cystodytes dellechiaiei*. λ_{\max} 225 (€ 37000); 273 (€ 25000); 382 (€ 10000) (MeOH) (Derep).

$1'\text{-Methoxy}$, $N^{2'}\text{-Ac}$: **Cystodytin K**

[496909-76-3]

$C_{20}H_{17}N_3O_3$ 347.373

Isol. from *Lissoclinum notti*. Yellow solid. $[\alpha]_D^{20}$ -292 (c, 0.06 in MeOH). λ_{\max} 207 (log € 4.57); 272 (log € 4.17); 382 (log € 3.67) (MeOH/trifluoroacetate).

$1'\text{-Methoxy}$, $N^{2'}-(3\text{-methyl-2-butanoyl})$:

Cystodytin F

[141544-62-9]

$C_{23}H_{21}N_3O_3$ 387.437

Alkaloid from *Cystodytes dellechiaiei*. Sol. MeOH; poorly sol. hexane, H_2O . λ_{\max} 225 (€ 30000); 272 (€ 23000); 380 (€ 11000) (MeOH) (Derep).

$1'\text{-Methoxy}$, $N^{2'}\text{-tigloyl}$: **Cystodytin G**

[141544-63-0]

$C_{23}H_{21}N_3O_3$ 387.437

Alkaloid from *Cystodytes dellechiaiei*. λ_{\max} 225 (€ 30000); 272 (€ 23000); 380 (€ 11000) (MeOH) (Derep).

Kobayashi, J. et al., *J.O.C.*, 1988, **53**, 1800-1804 (*Cystodytins A-C*, isol, uv, ir, pmr, cmr, ms, struct)

Kobayashi, J. et al., *J. Nat. Prod.*, 1991, **54**, 1634-1638 (*Cystodytins D-I*)

Ciufolini, M.A. et al., *J.A.C.S.*, 1991, **113**,

8016-8024 (*Cystodytins A,B*, synth)

Eldredge, G.S. et al., *J. Med. Chem.*, 1994, **37**,

3819-3827 (*Cystodytin J*)

Ciufolini, M.A. et al., *Tet. Lett.*, 1995, **36**,


4709-4712 (*Cystodytin A*, synth)

Appleton, D.R. et al., *Tetrahedron*, 2002, **58**,

9779-9783 (*Cystodytin K*)

Deacylisariotin

D-98



$C_9H_{13}NO_5$ 215.205

$N\text{-}(2E\text{-Decenoyl})$: **Isariotin D**

[952703-99-0]

$C_{19}H_{29}NO_6$ 367.441

Isol. from *Isaria tenuipes* BCC 7831. Amorph. solid. $[\alpha]_D^{27}$ -37 (c, 0.1 in MeOH). λ_{\max} 212 (log € 4.03) (MeOH).

$N\text{-}(7\text{-Oxo-}2E\text{-dodecenoyl})$: **Isariotin C**

[952703-98-9]

$C_{21}H_{31}NO_7$ 409.478

Isol. from *Isaria tenuipes* BCC 7831. Amorph. solid. $[\alpha]_D^{27}$ -25 (c, 0.09 in MeOH). λ_{\max} 211 (log € 4.2) (MeOH).

$N\text{-}(12\text{-Hydroxy-}2E\text{-dodecenoyl})$: **Isariotin A**

[952703-96-7]

$C_{21}H_{33}NO_7$ 411.494

Isol. from *Isaria tenuipes* BCC 7831. Amorph. solid. $[\alpha]_D^{27}$ -45 (c, 0.09 in MeOH). λ_{\max} 213 (log € 4.16) (MeOH).

$N\text{-}(11\text{-Carboxy-}2E\text{-undecenoyl})$: **Isariotin B**

[952703-97-8]

$C_{21}H_{31}NO_8$ 425.478

Isol. from *Isaria tenuipes* BCC 7831. Amorph. solid. $[\alpha]_D^{26}$ -23 (c, 0.07 in MeOH). λ_{\max} 211 (log € 4.24) (MeOH).

7-Me ether , $N\text{-}(2E\text{-decenoyl})$: **Antibiotic TK 57-164B**. TK 57-164B

[745050-51-5]

$C_{20}H_{31}NO_6$ 381.468

Isol. from *Isaria* sp. TK-57. Stereochem. not determined.

7-Me ether , $N\text{-}(2E\text{-dodecenoyl})$: **Antibio-**

tic TK 57-164A. TK 57-164A

[745050-50-4]

$C_{22}H_{35}NO_6$ 409.522

Isol. from *Isaria* sp. TK-57. Stereochem. not determined.

Pat. Coop. Treaty (WIPO), 2004, 04 74 269;

CA, **141**, 224074 (TK 57-164)

Haritakun, R. et al., *J. Nat. Prod.*, 2007, **70**,


1478-1480 (isol, pmr, cmr)

3-Deamino-3-hydroxysolanocapsine

D-99

16,23-Epoxy-16,28-secosolanidane-3,23-diol, 9CI. **3-Desamino-3-hydroxysolanocapsine**

[35865-62-4]



$C_{27}H_{45}NO_3$ 431.657

Alkaloid from roots of *Solanum aculeatum* (Solanaceae). Needles (MeOH aq.). Mp 204°. $[\alpha]_D^{25}$ +20.1 (c, 1 in $CHCl_3$).

23-Me ether: Aculeamine

[91377-17-2]

$C_{28}H_{47}NO_3$ 445.684

Alkaloid from roots of *Solanum aculeatum* (Solanaceae). Needles (MeOH aq.). Mp 205-207°. $[\alpha]_D^{25}$ +50.8 (c, 0.9 in $CHCl_3$).

23-Et ether: 3-Deamino-22-O-ethyl-3-hydroxysolanocapsine. **3-Desamino-22-O-ethyl-3-hydroxysolanocapsine**

[92070-74-1]

$C_{29}H_{49}NO_3$ 459.711

Isol. from roots of *Solanum aculeatum* (Solanaceae). Needles (Me_2CO aq.). Mp 183-185°. $[\alpha]_D^{25}$ +45.2 (c, 0.4 in $CHCl_3$). Artifact.

16,23,25-Triepimer: Solanocardinol. **Pimpifolidine**

[138665-46-0]

$C_{27}H_{45}NO_3$ 431.657

Alkaloid from *Lycopersicon pimpinellifolium* (currant tomato) and *Solanum neocardenasii* (Solanaceae). Cryst. (Me_2CO aq.). Mp 200-203°. $[\alpha]_D^{20}$ -1.9 (c, 1.05 in Py). Solanocardinol and Pimpifolidine not compared. Phys. props. refer to Pimpifolidine.

16,22,23,25-Tetraepimer: 22-Isopimpifolidine

[152322-51-5]



$C_{27}H_{45}NO_3$ 431.657

Alkaloid from roots of *Lycopersicon pimpinellifolium* (currant tomato). Needles (Me_2CO). Mp 200-204°. $[\alpha]_D^{20}$ -13.6 (c, 0.77 in Py).


16,22,23,25-Tetraepimer: 3-O- $[\beta\text{-D-glucopyranosyl-(1}\rightarrow\text{2)}\text{-}[\beta\text{-D-xylopyranosyl-(1}\rightarrow\text{3)}\text{-}[\beta\text{-D-glucopyranosyl-(1}\rightarrow\text{4)}\text{-}\beta\text{-D-galactopyranoside}]$: Lycoperside

H

[675828-28-1]

C₅₀H₈₃NO₂₂ 1050.199Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. [α]_D²⁵ -29.8 (c, 1.2 in MeOH).Coll, F. *et al.*, *Phytochemistry*, 1983, **22**, 2099-2100; 1984, **23**, 883-885 (*isol, pmr, cmr*)Kubschabsky, L. *et al.*, *CA*, 1984, **101**, 111265p (*cryst struct*)Osman, S.F. *et al.*, *Phytochemistry*, 1991, **30**, 3161-3163 (*Solanocardinol*)Ripperger, H. *et al.*, *Phytochemistry*, 1994, **35**, 813-815 (*Pimpifolidine, 22-Isopimpifolidine*)Yahara, S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 500-502 (*Lycoperside H*)**Debenzoylaralione A****D-100***Desbenzoylaralione A*
[21761-50-2]Absolute
ConfigurationC₂₇H₃₄N₄O₄ 478.59Isol. from stem bark of *Araliorhamnus vaginatus*. Mp 101-104°. [α]_D²⁰ +100 (c, 0.16 in MeOH).Tschesche, R. *et al.*, *Chem. Ber.*, 1969, **102**, 50-63 (*Debenzoylaralione A*)Gournelis, D.G. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1998, **75**, 1 (*rev*)**N-Debenzoyllucentamycin A****D-101**

Absolute Configuration

C₂₁H₃₈N₆O₄ 438.569N^{2'}-(3-Methylbutanoyl): **Lucentamycin C**
[949009-88-5]C₂₆H₄₆N₆O₅ 522.687Prod. by *Nocardiopsis lucentensis* (strain CNR-712). Yellow oil. [α]_D²⁵ -32 (c, 0.25 in MeOH), λ_{max} 206 (log ε 4.2); 217 (sh) (log ε 3.19) (MeOH).N^{2'}-(3-Methyl-2-butenoyl): **Lucentamycin D**
[949009-89-6]C₂₆H₄₄N₆O₅ 520.671Prod. by *Nocardiopsis lucentensis* (strain CNR-712). Yellow oil. [α]_D²⁵ -36 (c, 0.25 in MeOH), λ_{max} 206 (log ε 4.2); 217 (sh) (log ε 4) (MeOH).N^{2'}-Benzoyl: **Lucentamycin A**
[949009-86-3]C₂₈H₄₂N₆O₅ 542.677Prod. by the marine-derived *Nocar-**diopsis lucentensis* (strain CNR-712).Cytotoxic. Yellow oil. [α]_D²⁵ -6.3 (c, 0.17 in MeOH), λ_{max} 206 (log ε 4.1); 217 (sh) (log ε 3.9); 300 (log ε 2.7) (MeOH).Cho, J.Y. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1321-1328 (*isol, pmr, cmr*)**Debilic acid****D-102***8-Methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-acetic acid, 9CI. 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthreneacetic acid*
[475-85-4]C₁₈H₁₃NO₇ 355.303Alkaloid from the roots of *Aristolochia debilis*. Also assumed to be present in the stems of *Aristolochia manshuriensis* (Aristolochiaceae). Yellow needles. Mp 300° dec Mp 350°.*Me ester*: Mp 260°.Ku, Y.-T. *et al.*, *Kexue Tongbao (Chin. edn.)*,1957, 761; *CA*, **53**, 12256g (*isol, struct*)Tseng, K.-F. *et al.*, *Yaohue Xuebao*, 1958, **6**, 33; 316; *CA*, **53**, 15225a; **54**, 453i (*isol, struct*)Ruecker, G. *et al.*, *Planta Med.*, 1975, **27**, 68**2,8-Decadiene-4,6-diyonic acid****D-103***Matricaria acid. Matricaric acid*
H₃CCH=CHC≡CC≡CCH=CHCOOHC₁₀H₈O₂ 160.172**(2E,8E)-form**

Mp 173° dec.

Me ester: trans,trans-*Matricaria ester*
[820-96-2]C₁₁H₁₀O₂ 174.199Constit. of *Matricaria tenuifolia*, *Polyporus anthracophilus* and *Grindelia paludosa*. Mp 63°.**(2E,8Z)-form**

Mp 83-90°.

Me ester: trans-*Matricaria ester*
[23180-59-8]Constit. of *Matricaria inodora*, *Erigeron karwinskyanus* and *Amellus strigosus*. Mp 2°.*2-Methylpropylamide*: N-(2-Methylpropyl)-2,8-decadiene-4,6-diyonic acid isobutylamide
[113235-92-0]C₁₄H₁₇NO 215.294Isol. from underground parts of *Achillea wilhelmsii* (Asteraceae). Mp 85-89°.*3-Methylbutylamide*: N-(3-Methylbutyl)-2,8-decadiene-4,6-diyonic acid isopentylamide
[113262-63-8]C₁₅H₁₉NO 229.321Isol. from *Achillea wilhelmsii* (Asteraceae). Mp 88-91°.*2-Phenylethylamide*: N-(2-Phenylethyl)-2,8-decadiene-4,6-diyonic acid 2-phenylethylamide
[113262-64-9]C₁₈H₁₇NO 263.338Isol. from *Achillea wilhelmsii* (Asteraceae). Mp 108-111°.**(2Z,8E)-form***Me ester*: [25019-41-4]Constit. of *Solidago virgaurea* and *Erigeron* spp.**(2Z,8Z)-form** [13030-78-9]Isol. from fresh roots of *Anthemis ruffoliana*. Mp 98-99°. λ_{max} 246; 259; 315; 337 (MeOH) (Berdy).*Me ester*: *Matricaria ester*
[928-36-9]Constit. of *Matricaria inodora* and *Erigeron* spp. Widely distributed in the Asteraceae. Mp 37°.**(2ξ,8ξ)-form***2-Methylpropylamide*, N-Me: N-Methyl-N-(2-methylpropyl)-2,8-decadiene-4,6-diyonic acid
[37064-13-4]C₁₅H₁₉NO 229.321Isol. from the roots of *Anacyclus pyrethrum* (Asteraceae). Oil. λ_{max} 328, 306.5, 287 nm.*8,9-Epoxyde*: 8,9-Epoxy-2-decene-4,6-diyonic acid
C₁₀H₈O₃ 176.171Isol. as Me ester from aerial parts of *Erigeron philadelphicus*. Oil (as Me ester).Bruun, T. *et al.*, *Acta Chem. Scand.*, 1951, **5**, 1244 (*synth*)Sørensen, J.S. *et al.*, *Acta Chem. Scand.*, 1952, **6**, 883; 1953, **7**, 1375; 1954, **8**, 26; 34 (*occur*)Bu'Lock, J.D. *et al.*, *J.C.S.*, 1957, 1607 (*isol, struct*)Bell, I. *et al.*, *J.C.S.*, 1958, 1313 (*synth*)Bohlmann, F. *et al.*, *Chem. Ber.*, 1963, **96**, 1485; 1964, **97**, 1193; 1965, **98**, 369; 1966, **99**, 1642; 2096; 1967, **100**, 611; 1982, **21**, 167 (*isol, pmr*)Jente, R. *et al.*, *Chem. Ber.*, 1972, **105**, 1694 (*isol, struct, uv, ms*)Bohlmann, F. *et al.*, *Phytochemistry*, 1982, **21**, 167 (*isol*)Jakupovic, J. *et al.*, *Phytochemistry*, 1986, **25**, 1223 (*epoxides*)Greger, H. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1100 (*amides, isol, struct, ir, uv, ms, pmr, cmr*)**2,4-Decadienoic acid****D-104**

[92015-76-4]

H₃C(CH₂)₄CH=CHCH=CHCOOHC₁₀H₁₆O₂ 168.235**(E,E)-form** [30361-33-2]Mp 49-50°. Bp_{0.3} 121-124°.*2-Methylpropylamide*: see 2,4-Decadienoic acid isobutylamide, D-106*2-Methylbutylamide*: N-(2-Methylbutyl)-2,4-decadienamide. 2,4-Decadienoic acid 2-methylbutylamide
C₁₅H₂₇NO 237.384

Alkaloid from *Piper sarmentosum*. Oil. $[\alpha]_D^{20}$ -5 (c, 0.12 in EtOH). λ_{\max} 261 (no solvent reported).

3-Methylbutylamide: 2,4-Decadienoic acid isopentylamide

[78910-26-6]
C₁₅H₂₇NO 237.384

Isol. from underground parts of *Achillea wilhelmsii* (Asteraceae). Mp 63-65°.

2-Phenylethylamide: 2,4-Decadienoic acid 2-phenylethylamide

[83170-20-1]
C₁₈H₂₅NO 271.402

Isol. from underground parts of *Achillea wilhelmsii* (Asteraceae). Mp 96-98°.

Pyrrolidide: 1-(1-Oxo-2,4-decadienyl)-pyrrolidine, 9CI. 2,4-Decadienoic acid pyrrolidide. 1-(2,4-Decadienyl)pyrrolidine. Sarmentine

[78910-33-5]
C₁₄H₂₃NO 221.342

Constit. of fruits of pepper (*Piper nigrum*) and cha-plu (*Piper sarmentosum*) (Piperaceae). Shows larvicidal activity. Oil.

Piperidide: 1-(1-Oxo-2,4-decadienyl)piperidine, 9CI. 2,4-Decadienoic acid piperidide

[42997-42-2]
C₁₅H₂₅NO 235.369

Alkaloid from *Achillea nabelikii*, *Achillea sudetica* and other *Achillea* spp. (Asteraceae). Oil.

(4-Hydroxyphenyl)ethylamide: see N-(4-Hydroxyphenethyl)-2,4-decadienamide, H-668

(2E,4Z)-form

Phycodioic acid. Stillingia acid
[544-48-9]

Constit. of Stillingia oil. Also from *Phycomyces blakesleeanus*, *Streptomyces viridochromogenes* and *Agromyces* sp. Herbicide. Bp_{0.5} 122°.

Amide: 2,4-Decadienamide. Amidenin
[149764-39-6]
C₁₀H₁₇NO 167.25

Isol. from *Amycolatopsis* sp. Plant-growth regulator. Needles. Mp 88-90°. λ_{\max} 261 (€ 21900) (MeOH) (Derep).

Piperidide: Neopellitorine B
[88855-36-1]

C₁₅H₂₅NO 235.369

Alkaloid from *Artemisia dracunculus*. Yellow oil. Indexed as 4E-isomer in CA. λ_{\max} 257 (MeOH).

Hofer, O. et al., *Tetrahedron*, 1986, **42**, 2707-2716 (piperidide, cmr)

Greger, H. et al., *J. Nat. Prod.*, 1987, **50**, 1100 (2-phenylethylamide, isopentylamide)

Likhitwitayawuid, K. et al., *Tetrahedron*, 1987, **43**, 3689 (Sarmentine)

Kiuchi, F. et al., *Chem. Pharm. Bull.*, 1988, **36**, 2452 (pyrrolidide)

Bari, S.S. et al., *J. Indian Chem. Soc.*, 1990, **67**, 995 (synth, Sarmentine)

Kanbe, K. et al., *Biosci., Biotechnol., Biochem.*, 1993, **57**, 1261 (Amidenin)

Strunz, G.M. et al., *Can. J. Chem.*, 1996, **74**, 419-432 (piperidide, pyrrolidide, synth)

Stoehr, J.R. et al., *Planta Med.*, 1999, **65**, 175-177 (2-methylbutylamide)

Saadali, B. et al., *Phytochemistry*, 2001, **58**, 1083-1086 (Neopellitorine B)

6,8-Decadienoic acid D-105

H₃CCH=CHCH=CH(CH₂)₄COOH
C₁₀H₁₆O₂ 168.235

(6Z,8E)-form

N-(2-Methylpropyl)amide: N-(2-Methylpropyl)-6,8-decadienamide. N-Isobutyl-6,8-decadienamide. Hydrospilanthol
[94450-20-1]
C₁₄H₂₅NO 223.358

Alkaloid from the flowers of *Acmella ciliata*.

Martin, R. et al., *Phytochemistry*, 1984, **23**, 1781-1783 (Hydrospilanthol)

2,4-Decadienoic acid isobutylamide D-106

Pellitorine. N-(2-Methylpropyl)-2,4-decadienamide, 9CI. N-Isobutyl-2,4-decadienamide, 8CI. Pyretrin

H₃C(CH₂)₄CH=CHCH=CHCONHCH₂CH(CH₃)₂
C₁₄H₂₅NO 223.358

(E,E)-form [18836-52-7]

Constit. of *Fagara xanthoxyloides*, *Piper sylvaticum*, *Piper nepalense* and several other *Piper* spp., *Anacyclus pyrethrum*, *Achillea millefolium* (yarrow) and *Asiasarum heterotropoides* (Rutaceae, Piperaceae, Asteraceae, Aristolochiaceae). Produces intense formication and local anaesthesia of the mucous membranes. Insecticide. Needles (petrol). Mp 75° (69°) Mp 90-95°.

► Severe irritant.

6-Hydroxy: 6-Hydroxy-N-(2-methylpropyl)-2,4-decadienamide, 9CI. 6-Hydroxy-N-isobutyl-2,4-decadienamide. 6-Hydroxypellitorine
[133640-43-4]
C₁₄H₂₅NO₂ 239.357

Constit. of *Anacyclus monanthos*.

(2E,4Z)-form

Alkaloid from *Artemisia dracunculus*. Yellow oil. λ_{\max} 257 (MeOH).

Crombie, L. et al., *J.C.S.*, 1955, 1007 (synth)
Bowden, K. et al., *J.C.S.*, 1963, 3503 (isol, uv, ir)

Loder, J.W. et al., *Aust. J. Chem.*, 1969, **22**, 1531 (isol, uv, ir, ms, synth)

Banerji, A. et al., *Experientia*, 1974, **30**, 223 (isol, struct)

Mahanta, P.K. et al., *J. Pharm. Sci.*, 1974, **63**, 1160 (isol, uv, ir)

Dasgupta, S. et al., *Indian J. Chem., Sect. B*, 1979, **17**, 538 (isol, uv, ir, pmr, ms)

Ono, N. et al., *Chem. Lett.*, 1980, 1365 (synth)

Nokami, J. et al., *Tet. Lett.*, 1980, **21**, 4455 (synth)

Yasusa, I. et al., *Chem. Pharm. Bull.*, 1981, **29**, 564 (isol, uv, ir, pmr, cmr, ms)

Tamura, Y. et al., *Tet. Lett.*, 1981, **22**, 1343 (synth)

Sato, T. et al., *Tet. Lett.*, 1981, **22**, 2375 (synth, ir, pmr)

Bari, S.S. et al., *J. Indian Chem. Soc.*, 1990, **67**, 995 (synth)

Ahmed, A. et al., *Pharmazie*, 1990, **45**, 941 (6-Hydroxypellitorine)

Kaga, H. et al., *Synlett*, 1994, 607 (synth)

Strunz, G.M. et al., *Can. J. Chem.*, 1996, **74**, 419 (synth)


Abarbri, M. et al., *Synth. Commun.*, 1998, **28**, 239-249 (synth, ir, pmr, cmr, ms)

Saadali, B. et al., *Phytochemistry*, 2001, **58**, 1083-1086 (2E,4Z-isomer)

Ley, J.P. et al., *Eur. J. Org. Chem.*, 2004, 5135-5140; 2005, 618 (synth)

5-(1,3-Decadienyl)octahydroindolizine D-107

5-(1,3-Decadienyl)indolizidine. Piclavine B
[142609-25-4]



C₁₈H₃₁N 261.45

Alkaloid from the tunicate *Clavelina picta*. Exhibits antimicrobial props. Oil. $[\alpha]_D$ +33.5 (c, 1 in CH₂Cl₂). Obt. as a mixt. of stereoisomers. λ_{\max} 232 (€ 12500) (EtOH) (Derep). λ_{\max} 232 (€ 12500) (MeOH) (Berdy).

5',6'-Didehydro: 5-(1,3,5-Decatrienyl)octahydroindolizidine. 5-(1,3,5-Decatrienyl)indolizidine. Piclavine C
[142609-26-5]

C₁₈H₂₉N 259.434

Isol. from *Clavelina picta*. Exhibits antimicrobial props. Pale yellow oil. $[\alpha]_D$ +36 (c, 5 in CH₂Cl₂). Obt. as a mixt. of stereoisomers. λ_{\max} 262 (€ 11800); 267 (€ 15000); 281 (€ 11800) (EtOH) (Derep). λ_{\max} 262 (€ 11800); 267 (€ 15000); 281 (€ 11800) (MeOH) (Berdy).

Raub, M.F. et al., *Tet. Lett.*, 1992, **33**, 2257-2260 (isol, struct)

2,8-Decadien-6-ynoic acid D-108

H₃CCH=CHC≡CCH₂CH₂CH=CHCOOH

C₁₀H₁₂O₂ 164.204

(2Z,8Z)-form


2-Phenylethylamide: 2,8-Decadien-6-ynoic 2-phenylethylamide
[96917-23-6]

C₁₈H₂₁NO 267.37

Constit. of *Salmea scandens*. Oil.

Bohlmann, F. et al., *Phytochemistry*, 1985, **24**, 595-596 (isol)

Decahydro-2,5-bis(4-pentenyl)quinoline D-109



(2R*,4aR*,5S*,8aS*)-form

C₁₉H₃₃N 275.476

(2R,4aR,5S,8aS)-form**Dendrobates Alkaloid 275B**

Alkaloid from skin extracts of *Dendrobates auratus*, *Dendrobates granuliferus* and *Dendrobates pumilio*.

(2R,4aS,5R,8aR)-form**Dendrobates Alkaloid 275B'**

Alkaloid from skin extracts of *Dendrobates* spp.

(2R,4aS,5R,8aS)-form**5-epi-trans-275B**

Alkaloid from extracts of virgin queens of *Solenopsis azteca*.

(2R,4aS,5S,8aR)-form**5-epi-cis-275B'**


Alkaloid from extracts of virgin queens of a myrmecine ant *Solenopsis azteca*.

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (pmr, cmr, occur)

Toyooka, N. *et al.*, *J.O.C.*, 2002, **67**, 6078-6081 (synth)

Decahydro-10-[(3-hydroxy-2,4,6-octadecatrienoyl)amino]-1,11a,11b-trihydroxy-2H-pyrazino[1,2-a]pyrrolo[2,1-c]pyrazine-3,6,9-trione

[212070-43-4]




$C_{28}H_{42}N_4O_8$ 562.662

Enolised β -diketone. Prod. by *Sporormiella minimoides* MF 5867. Antifungal agent. Off-white solid. $[\alpha]_D^{20} +10$ (c, 0.65 in MeOH). λ_{max} 283 (ϵ 9900) (MeOH). *U.S. Pat.*, 1998, 5 801 172; *CA*, **129**, 215776a

7-[5-(Decahydro-4a-hydroxy-1,2,5,5-tetramethyl-1-naphthalenyl)-3-methyl-2-pentenyl]-3,7-dihydro-2,3-dimethyl-6H-purin-6-one, 9CI

[90162-26-8]



$C_{27}H_{42}N_4O_2$ 454.654


Isol. from the marine sponge *Agelas mauritiana*. Cryst. (C_6H_6) (as Ac). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 95-98° (Ac). Seemingly an artifact; the struct. of the true sponge metab. is unknown. λ_{max} 221 (ϵ 9400); 267 (ϵ 3300); 344 (ϵ 200) (MeOH) (Berdy).

Nakatsu, T. *et al.*, *Tet. Lett.*, 1984, **25**, 935 (isol, uv, pmr, ms, cryst struct)

Decahydro-2-methyl-5-propylquinoline

D-112

2-Methyl-5-propyldecahydroquinoline



$C_{13}H_{25}N$ 195.347

(2R,4aR,5R,8aS)-form**Alkaloid 195J**


[232613-57-9]

Alkaloid from the ant *Solenopsis picea*.

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1999, **25**, 1179-1193

Decahydro-8-(1-piperidinyl-methyl)isoquinoline Dihydrochoberine

D-113



$C_{15}H_{28}N_2$ 236.4

Alkaloid from the aerial parts of *Nitraria sibirica*. Cryst. (EtOH). Mp 251-252°.


Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 556-558 (isol, pmr, ms)

Decahydro-2,5,8-trimethyl-2H-1-oxa-4,7,9b-triazaphenalene, 9CI

D-114

Base T

[54352-32-8]



$C_{12}H_{23}N_3O$ 225.333

Alkaloid from *Lycopodium thyooides*. Cryst. (hexane). Mp 101.5-102°. Opt. inactive. Transparent to uv. Poss. artifact, though it has been stated that the related compd. Dodecahydro-2,5,8-trimethyl-1,4,7,9b-tetraazaphenalene, D-888 is not.

Stereoisomer(?): *Lycopodium Alkaloid C*

[29066-78-2]

Cryst. + $\frac{1}{2} H_2O$. Mp 55-57°. There appears to be no evidence for isolation of this compd. The name is as reported in the CAS Index Guide.

[29066-78-2]

Braekman, J.C. *et al.*, *Phytochemistry*, 1974, **13**, 2519

Decanoic acid

D-115

Capric acid. Caprinic acid. Decylic acid. Decoic acid (obso.). FEMA 2364

[334-48-5]

$H_3C(CH_2)_8COOH$

$C_{10}H_{20}O_2$ 172.267

Widespread in plant oils, esp. leaf oils and as glycerides in seed oils. Used in corrosion inhibitors and surfactants and in extraction-preconcentration of Pb. Salts (Na, K, Mg, Ca, Al) used as binders, emulsifiers and anticaking agents in food manuf. Needles. Sol. EtOH, Et₂O, Me₂CO, C₆H₆, $CHCl_3$, alkalis; poorly sol. H_2O . d_{40}^{20} 0.89. Mp 31.5°. Bp 268-270° Bp₁₁ 148-150°. n_D^{40} 1.4288.

► Fl. p. >66°. Skin irritant. LD₅₀ (mus, ivn) 129 mg/kg. HD9100000

2-Methylpropylamide: N-(2-Methylpropyl)decanamide, 9CI. **Decanoic acid isobutylamide**

[73785-31-6]

$C_{14}H_{29}NO$ 227.389

Trace constit. of *Cissampelos glaberrima* roots. Solid (EtOAc). Unstable at r.t. Struct. based on ms data only.

[22620-93-5, 1002-62-6, 13747-30-3, 13040-18-1, 30673-38-2]


Rosario, S.L. *et al.*, *Planta Med.*, 1996, **62**, 376 (isobutylamide)

5-Decanoyl-2-nonylpyridine

D-116

1-(6-Nonyl-3-pyridinyl)-1-decanone, 9CI

[149682-94-0]



$C_{24}H_{41}NO$ 359.594

Alkaloid from *Houttuynia cordata* (Yu Xing Cao).


Jong, T.T. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1993, **40**, 301-303 (isol)

Proebstle, A. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 235-240 (isol)

Decarbomethoxyapocuanzine

D-117

[53492-10-7]



$C_{20}H_{22}N_2O_2$ 322.406


Alkaloid from the root bark of *Voacanga chalotiana* (Apocynaceae). Cryst. (EtOAc). Mp 196°. $[\alpha]_D^{22} -132$ ($CHCl_3$).

Bombardelli, E. *et al.*, *Experientia*, 1974, **30**, 979 (uv, pmr, ms, struct)

Sóti, F. *et al.*, *Tetrahedron*, 1994, **50**, 8209 (synth)

16'-Decarbomethoxy-19,20-dihydroconoduramine

D-118

C₁₉H₂₈N₂ 284.444C₄₁H₅₂N₄O₃ 648.887

Alkaloid from *Ervatamia officinalis*. Indexed by CAS, apparently incorrectly, as 16'-Decarbomethoxy-19,20-dihydrovoacamine in T-8.

20-Epimer: 16'-Decarbomethoxy-20-epi-19,20-dihydroconoduramineC₄₁H₅₂N₄O₃ 648.887Alkaloid from *Ervatamia officinalis*.


Possibly the same as 16'-Decarbomethoxy-19,20-dihydrovoacamine in T-8.

Huang, L. *et al.*, *Zhongcaoyao*, 1997, **28**, 323-326; *CA*, **128**, 203013r (*isol*)

16-Decarbomethoxy-19,20-dihydrobasine

D-119

16-De(methoxycarbonyl)-19,20-dihydrovoacamine, 8CI
[19779-70-5]



Absolute Configuration

C₁₉H₂₄N₂O 296.411

Alkaloid from *Rauwolfia discolor* (Apocynaceae). Cryst. (MeOH). Mp 100-110°. $[\alpha]_D^{25}$ -20 (c, 1.5 in CHCl₃). λ_{max} 238 (ε 15170); 313 (ε 22330) (95% EtOH).

Methiodide: Mp 250°.

20-Epimer: 16-Decarbomethoxy-20-epi-19,20-dihydrovoacamine. 20-Epi-16-decarbomethoxy-19,20-dihydrovoacamine

[203929-19-5]

C₁₉H₂₄N₂O 296.411Alkaloid from *Ervatamia officinalis*.


Combes, G. *et al.*, *Phytochemistry*, 1968, **7**, 477-483 (*uv, ir, pmr, ms, struct*)

Huang, L. *et al.*, *Zhongcaoyao*, 1997, **28**, 323-326; *CA*, **128**, 203013r (*20-epimer*)

Decarbomethoxytetrahydrosecodine

D-120

2-Ethyl-3-[2-(3-ethylpiperidino)ethyl]indole, 9CI, 8CI
[20785-62-0]



(R)-form

(R)-form

Alkaloid from the leaves of *Tabernaemontana cumminsii* and whole plants of *Haplophyton crooksii* (Apocynaceae). Also found in *Aspidosperma marcgravianum* and *Rhazya stricta* (Apocynaceae). Exhibits antiacetylcholinesterase activity. Oil or amorph. powder. $[\alpha]_D$ +90 (CHCl₃). Specific rotn. only reported for the *Rhazya stricta* alkaloid (+90) but abs. config. not assigned. However, synthetic (+)-enantiomer ($[\alpha]_D$ +11.8) has R-config.

Picrate: Mp 137-138°.

16-Oxo: Crooksidine

[149301-48-4]

C₁₉H₂₆N₂O 298.427

Alkaloid from whole plants of *Haplophyton crooksii* (Apocynaceae). Exhibits antiacetylcholinesterase activity. Glass. $[\alpha]_D$ +27.6 (c, 0.205 in CHCl₃) (natural). $[\alpha]_D^{31}$ +7.8 (c, 0.6 in CHCl₃) (synthetic).

19-Oxo: 19-Oxodecarbomethoxytetrahydrosecodine

[88607-53-8]

C₁₉H₂₆N₂O 298.427

Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph. Abs. config. not certain.


(S)-form

Synthetic. $[\alpha]_D^{22}$ -10.8 (c, 0.45 in CHCl₃). Crooks, P.A. *et al.*, *Chem. Comm.*, 1968, 1210 (*isol, uv, ms, synth, struct*)
Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694 (*19-Oxodecarbomethoxytetrahydrosecodine*)
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1991, **30**, 1285 (*isol, uv, ir, pmr, cmr, ms*)
Mroue, M.A. *et al.*, *Phytochemistry*, 1993, **33**, 217 (*Crooksidine*)
Aclinou, P. *et al.*, *Nat. Prod. Lett.*, 1994, **5**, 197 (*Crooksidine, synth*)
Palmisano, G. *et al.*, *Tetrahedron: Asymmetry*, 1995, **6**, 1229 (*synth*)
Sakagami, H. *et al.*, *Synlett*, 1996, 163 (*Crooksidine, synth, abs config*)
Amat, M. *et al.*, *Tetrahedron: Asymmetry*, 1996, **7**, 3091 (*synth*)

16-Decarbomethoxyvoacamine pseudoindoxyl

D-121

[139022-94-9]


C₄₁H₅₀N₄O₄ 662.87

Alkaloid from leaves and root bark of *Ervatamia hirta* (Apocynaceae). $[\alpha]_D$ -14 (c, 0.5 in MeOH).

Clivio, P. *et al.*, *Phytochemistry*, 1991, **30**, 3785 (*isol, uv, ir, pmr, cmr, ms, struct*)

2-Decarboxybetanidin

D-122

2-DecarboxybetanidinC₁₇H₁₇N₂O₆[⊕] 345.331**5-O-β-D-Glucopyranoside: 2-Decarboxybetanin. 2-Decarboxybetanin**C₂₃H₂₇N₂O₁₁[⊕] 507.473Constit. of *Beta vulgaris*.**5-O-(6-O-Malonyl-β-D-glucopyranoside): 2-Decarboxyphylloactin. 2-Decarboxyphylloactin**C₂₆H₂₉N₂O₁₄[⊕] 593.52Constit. of *Beta vulgaris*.

Kobayashi, N. *et al.*, *Phytochemistry*, 2001, **56**, 429-436 (*isol, pmr, ms*)
Wybraniec, S. *et al.*, *Tet. Lett.*, 2006, **47**, 1725-1728 (*pmr, cmr*)

2,4,8-Decatrienoic acid

D-123

H₃CCH=CHCH₂CH₂CH=CHCH=CHCOOHC₁₀H₁₄O₂ 166.219**(2E,4E,8Z)-form**

2-Methylpropylamide: N-Isobutyl-2,4,8-decatrienamamide. N-(2-Methylpropyl)-2,4,8-decatrienamamide, 9CI. 2,4,8-Decatrienoic acid isobutylamide
[52657-13-3]

C₁₄H₂₃NO 221.342

Alkaloid from the roots of *Achillea millefolium* (yarrow) (Asteraceae). Constit. of *Otanthus maritimus*. Oil.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1974, **107**, 1038-1043 (*isol, uv, ir, pmr, ms, struct*)

2,6,8-Decatrienoic acid

D-124

H₃CCH=CHCH=CHCH₂CH₂CH=CHCOOHC₁₀H₁₄O₂ 166.219

Powerful insecticide. Log P 3.15 (calc).

(2E,6Z,8E)-form**Spilanthic acid**

[94450-21-2]

N-(2-Methylpropyl)amide: N-(2-Methylpropyl)-2,6,8-decatrienamamide, 9CI. N-Isobutyl-2,6,8-decatrienamamide.

Affinine[†]. *Spilanthol*

[25394-57-4]

C₁₄H₂₃NO 221.342Alkaloid from the roots of *Heliopsis*

longipes, *Spilanthes oleraceae* and *Spilanthes acmella*. Shows inhibiting activity against *Escherichia coli* and *Saccharomyces cerevisiae*. Pale-yellow viscous oil. Mp 23°. Bp_{0.5} 165° Bp_{0.001} 120-125°. n_D²⁵ 1.5135. In the earlier lit. *H. longipes* was incorrectly identified as *Erigeron affinis*.

► HE1800000

(all-E)-form

N-(2-Methylpropyl)amide: [76361-77-8]
Mp 91.5-92.5°. Bp_{0.25} 145°.

Asano, M. *et al.*, *Ber.*, 1932, **65**, 1602 (*isol*)
Gokhale, V.G. *et al.*, *J. Indian Chem. Soc.*, 1945, **22**, 250 (*isol*)

Acree, F. *et al.*, *J.O.C.*, 1945, **10**, 236; 449 (*isol*, *uv*, *struct*)

Jacobson, M. *et al.*, *J.O.C.*, 1947, **12**, 731

Jacobson, M. *et al.*, *J.A.C.S.*, 1955, **77**, 2461 (*synth*)

Jacobson, M. *et al.*, *Chem. Ind. (London)*, 1957, 50 (*struct*, *synth*)

Crombie, L. *et al.*, *J.C.S.*, 1963, 4970 (*ir*, *uv*, *synth*)

Correa, J. *et al.*, *Org. Magn. Reson.*, 1971, **3**, 1 (*pmr*)


Yasuda, I. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2251 (*isol*, *synth*, *pmr*, *cmr*, *ms*)

Martin, R. *et al.*, *Phytochemistry*, 1984, **23**, 1781-1783 (*isol*)

Molina-Torres, J. *et al.*, *J. Ethnopharmacol.*, 1999, **64**, 241-248 (*Affinine*, *activity*)

2,4,6-Decatrienoic acid dehydropiperidide D-125

1,2,3,4-Tetrahydro-1-(1-oxo-2,4,6-decatrienyl)pyridine, 9CI



C₁₅H₂₁NO 231.337

(2E,4E,6Z)-form [43110-67-4]

Alkaloid from *Achillea nabelikii*, *Achillea millefolium* (yarrow) and *Achillea crithmifolia* (Asteraceae). Yellow cryst. (Et₂O/petrol). Mp 55°.

6,7-Dihydro: **2,4-Decadienoic acid dehydropiperidide**

[81427-11-4]

C₁₅H₂₃NO 233.353

Alkaloid from *Achillea crithmifolia* (Asteraceae). Gum.

8,9-Didehydro: **2,4,6,8-Decatetraenoic acid dehydropiperidide**

[43110-68-5]

C₁₅H₁₉NO 229.321

Alkaloid from *Achillea nabelikii* and *Achillea millefolium* (yarrow) (Asteraceae). Yellowish oil. Not. obt. completely pure.

[42997-42-2, 42997-44-4]


Bohlmann, F. *et al.*, *Chem. Ber.*, 1973, **106**, 1328 (*isol*, *uv*, *pmr*, *struct*, *synth*)

Greger, H. *et al.*, *Phytochemistry*, 1981, **20**, 2579 (*isol*, *deriv*)

Strunz, G.M. *et al.*, *Can. J. Chem.*, 1996, **74**, 419-432 (*synth*, *deriv*)

Decatromicin A

D-126



C₄₅H₅₇ClN₂O₁₀ 821.405

Similar to Kijanimicin and Pyrrolosporin A, P-969. Prod. by *Actinomadura* sp. MK73-NF4. Active against gram-positive bacteria incl. MRSA. Powder. Mp 223-225° (dec.). [α]_D²⁶ +2 (c, 1.3 in MeOH). λ_{max} 271 (log ε 4.47) (MeOH).

4''-Chloro: **Decatromicin B**

C₄₅H₅₆Cl₂N₂O₁₀ 855.85

Prod. by *Actinomadura* sp. MK73-NF4. Active against gram-positive bacteria incl. MRSA. Powder. Mp 202-206° (dec.). [α]_D²⁶ -9 (c, 1.3 in MeOH). λ_{max} 269 (log ε 4.46) (MeOH).

5''-Chloro: **Antibiotic BE 45722A. BE 45722A**

[196402-69-4]

C₄₅H₅₆Cl₂N₂O₁₀ 855.85

Prod. by *Actinomadura* sp. A 45722. Antibacterial agent. Stereochem. not confirmed.

4'',5''-Dichloro: **Antibiotic BE 45722B. BE 45722B**

[196402-70-7]

C₄₅H₅₅Cl₃N₂O₁₀ 890.295

Prod. by *Actinomadura* sp. A 45722. Antibacterial agent. Stereochem. not confirmed.

4'',5''-Dichloro, N''-Me: **Antibiotic BE 45722C. BE 45722C**

[196402-71-8]

C₄₆H₅₇Cl₃N₂O₁₀ 904.322

Prod. by *Actinomadura* sp. A 45722. Antibacterial agent. Stereochem. not confirmed.

Japan. Pat., 1997, 97 227 587; CA, **127**,


261794m (BE 45722)

Momose, I. *et al.*, *J. Antibiot.*, 1999, **52**, 781-786; 787-796

Decaturin A

D-127

[509095-70-9]



C₃₀H₃₅NO₆ 505.61

Related to Oxalicine A, O-148.

Prod. by *Penicillium decaturensis*.

Cryst. (MeCN aq.). Mp 160-162°. [α]_D +32 (c, 0.5 in CH₂Cl₂). λ_{max} 205 (ε 16000); 235 (ε 16000); 335 (ε 5800) (MeOH).

23-Deoxy: **Decaturin C**

[849339-59-9]

C₃₀H₃₅NO₅ 489.61

Prod. by *Penicillium thiersii*.

Cryst. Mp 171-174°. [α]_D +167

(c, 0.2 in CH₂Cl₂). λ_{max} 235 (ε 15000); 270 (ε 5900); 325 (ε 1600) (MeOH).

23-Deoxy, 15β-hydroxy: **Decaturin B**

[509095-72-1]

C₃₀H₃₅NO₆ 505.61

Prod. by *Penicillium thiersii*. Antiinsectan.

Cryst. (CH₂Cl₂). Mp 235-237°.

[α]_D +124 (c, 0.2 in CH₂Cl₂). λ_{max} 233 (ε 19000); 270 (ε 7800); 317 (ε 5500) (MeOH).


Zhang, Y. *et al.*, *Org. Lett.*, 2003, **5**, 773-776 (*isol*, *pmr*, *cmr*, *ms*)

Li, C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 319-322 (*Decaturin C*)

Decaturin D

D-128

[849339-60-2]



Absolute Configuration

C₃₀H₃₅NO₄ 473.611

Related to Oxalicine A, O-148. Prod. by

Penicillium thiersii. Antiinsectan. Oil.

[α]_D +58 (c, 0.1 in CH₂Cl₂). λ_{max} 233 (ε 14000); 270 (ε 5100); 327 (ε 3100) (CH₂Cl₂).

Li, C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 319-322 (*isol*, *pmr*, *cmr*, *ms*)

Hosoe, S. *et al.*, *Tet. Lett.*, 2006, **47**, 4425-4428 (*synth*, *abs config*)

2-Decene-4,6-diyonic acid D-129

Lachnophyllic acid

H₃CCH₂CH₂C≡CC≡CCH=

CHCOOH

C₁₀H₁₀O₂ 162.188

(ξ)-form

N-Methyl-N-(2-methylpropyl)amide: N-Methyl-N-(2-methylpropyl)-2-decene-4,6-diyynamide

[37064-12-3]

C₁₅H₂₁NO 231.337

Isol. from the roots of *Anacyclus pyrethrum*. Oil. Stereochem. unspecified. λ_{max} 252; 264; 281; 299 (no solvent reported).

[114825-67-1, 54497-59-5, 114825-66-0]

Jente, R. *et al.*, *Chem. Ber.*, 1972, **105**, 1694 (*Anacyclus pyrethrum constii*)

2-Decene-6,8-diyonic acid, 9CI D-130

$\text{H}_3\text{C}\equiv\text{CC}\equiv\text{CCH}_2\text{CH}_2\text{CH}=\text{CHCOOH}$
 $\text{C}_{10}\text{H}_{10}\text{O}_2$ 162.188

(Z)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2-decene-6,8-diyonamide, 9CI. 2-Decene-6,8-diyonic acid isobutylamide [96602-66-3]

$\text{C}_{14}\text{H}_{19}\text{NO}$ 217.31

Isol. from the roots of *Spilanthes oleracea*. Oil.

2-Phenylethylamide: N-(2-Phenylethyl)-2-decene-6,8-diyonamide [75872-73-0]

$\text{C}_{18}\text{H}_{19}\text{NO}$ 265.354

Isol. from *Spilanthes alba*. Oil.

2-Phenylethenylamide, (E-): N-(2-Phenylethenyl)-2-decene-6,8-diyonamide [75872-76-3]

$\text{C}_{18}\text{H}_{17}\text{NO}$ 263.338

Isol. from *Spilanthes alba*. Oil.

2-Phenylethenylamide, (Z-): [75883-36-2] Isol. from *Spilanthes alba*. Oil.

Bohlmann, F. et al., *Phytochemistry*, 1980, **19**, 1535; 1985, **24**, 595 (isol, struct, ir, ms)

Greger, H. et al., *Monatsh. Chem.*, 1985, **116**, 273 (isol, struct, uv, ir, pmr, cmr)

2-Decene-4,6,8-triynoic acid D-131

Dehydromatricaric acid
 $\text{H}_3\text{CC}\equiv\text{CC}\equiv\text{CC}\equiv\text{CCH}=\text{CHCOOH}$
 $\text{C}_{10}\text{H}_6\text{O}_2$ 158.156

(E)-form [7199-97-5]

Isol. from *Solidago virga-aurea*.

Me ester: Dehydromatricaria ester [692-94-4]

$\text{C}_{11}\text{H}_8\text{O}_2$ 172.183

Occurs in *Matricaria inodora* and *Matricaria oreades* and many other Asteraceae spp. Also isol. from culture medium of *Pleurotus ulmarius*. Light-yellow cryst. (Et₂O/petrol). Mp 105-106°.

3-Methylbutylamide: 2-Decene-4,6,8-triynoic acid isopentylamide [113235-91-9]

$\text{C}_{15}\text{H}_{17}\text{NO}$ 227.305

Isol. from the underground parts of *Achillea wilhelmsii* (Asteraceae). Cryst. Dec. to blue polymers on htg.

2-Phenylethylamide: N-(2-Phenylethyl)-2-decene-4,6,8-triynamide. 2-Decene-4,6,8-triynoic acid 2-phenylethylamide [113235-93-1]

$\text{C}_{18}\text{H}_{15}\text{NO}$ 261.323

Isol. from the underground parts of *Achillea wilhelmsii* (Asteraceae). Cryst. Dec. to blue polymer on htg.

2-Methylpropylamide: Dehydromatricaric acid isobutylamide [37064-10-1]

$\text{C}_{14}\text{H}_{15}\text{NO}$ 213.279

Isol. from *Achillea spinulifolia*, *Achillea ptarmica*, *Achillea impatiens*, *Achillea sibirica*, *Anacychus pyrethrum* and *Cladanthus arabisicus*. Cryst. (CCl₄). Mp

133-139° Mp 144.5-145.5°

(Z)-form [23050-77-3]

Isol. from *Solidago virga-aurea*. Mp 150° dec.

Me ester: [2739-57-3]

Occurs in *Artemisia vulgaris* (mugwort) and many other Asteraceae. Light-yellow needles (petrol). Mp 114-115° (112°).

[43110-85-6, 7329-73-9]

Stauholt, K. et al., *Acta Chem. Scand.*, 1950, **4**, 1567 (isol)

Christensen, P.K. et al., *Acta Chem. Scand.*, 1952, **6**, 602 (synth)

Sorensen, J.S. et al., *Acta Chem. Scand.*, 1954, **8**, 26 (isol)

Bell, I. et al., *J.C.S.*, 1958, 1313 (synth, Me ester)

Gardner, J.N. et al., *J.C.S.*, 1960, 691 (isol)

Bohlmann, F. et al., *Chem. Ber.*, 1962, **95**, 1742; 1973, **106**, 1328 (isol, struct, synth)

Hodge, P. et al., *J.C.S. (C)*, 1966, 1216 (bibl)

Karrer, W. et al., *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 770

Kobayashi, A. et al., *Agric. Biol. Chem.*, 1975, **39**, 911 (synth)

Kawazu, K. et al., *Agric. Biol. Chem.*, 1977, **41**, 223

Greger, H. et al., *Phytochemistry*, 1978, **17**, 86; 1982, **21**, 1071 (isol, struct)

Bohlmann, F. et al., *Phytochemistry*, 1979, **18**, 1736; 1980, **19**, 841; 2655 (isol)

Greger, H. et al., *J. Nat. Prod.*, 1987, **50**, 1100 (amides, isol, ir, uv, ms, pmr)

Japan. Pat., 1991, 03 287532; *CA*, **116**, 181120w (isol)

Lu, T. et al., *Phytochemistry*, 1993, **32**, 1483 (cryst struct)

Taha, A.A. et al., *Phytochemistry*, 2000, **55**, 921-926 (isol, pmr)

2-Decenoic acid

D-132

[3913-85-7]

$\text{H}_3\text{C}(\text{CH}_2)_6\text{CH}=\text{CHCOOH}$

$\text{C}_{10}\text{H}_{18}\text{O}_2$ 170.251

(E)-form

FEMA 3913

[334-49-6]

Constit. of essential oil of *Achasma walang* (preferred genus name

Etilingera). Also occurs in pear,

capsicum and black tea. Mosquito

repellant. Comly. available flavour

ingredient. Liq. or solid with fresh,

sweet fruity-fatty odour. Mp 12°. Bp₁₅

165° Bp_{4.5} 148-149°.

2-Methylpropylamide: N-(2-Methylpropyl)-2-decenamide. **2-Decenoic acid isobutylamide**

[73785-32-7]

$\text{C}_{14}\text{H}_{27}\text{NO}$ 225.373


Trace constit. of *Cissampelos glaberrima* roots. Solid (EtOAc).

Unstable at r.t. Struct. based on ms data only.

Rosario, S.L. et al., *Planta Med.*, 1996, **62**, 376 (isobutylamide)

5-(2-Decenyl)octahydroindolizine D-133

5-(2-Decenyl)indolizidine. **Piclavine A**



(2'E,5R,8aR)-form

$\text{C}_{18}\text{H}_{33}\text{N}$ 263.465

Shows antimicrobial props. Oil. $[\alpha]_{\text{D}} +15$ (c, 1 in CH_2Cl_2). Data given is for the mixt. of isomers.

(2'E,5R,8aR)-form

Piclavine A₁

[142609-24-3]

Isol. from the tunicate *Clavelina picta*. $[\alpha]_{\text{D}}^{25} -5.6$ (c, 0.84 in CH_2Cl_2).

(2'E,5R,8aS)-form

Piclavine A₃

[142696-98-8]

From *Clavelina picta*. $[\alpha]_{\text{D}}^{26} -73.6$ (c, 1.3 in CH_2Cl_2).

(2'Z,5R,8aR)-form

Piclavine A₂

[142696-99-9]

From *Clavelina picta*. $[\alpha]_{\text{D}}^{27} +4$ (c, 0.21 in CH_2Cl_2).

(2'Z,5R,8aS)-form

Piclavine A₄

[142697-00-5]

From *Clavelina picta*. $[\alpha]_{\text{D}}^{27} -76.4$ (c, 0.63 in CH_2Cl_2).

Raub, M.F. et al., *Tet. Lett.*, 1992, **33**, 2257-2260 (isol, struct)

Jefford, C.W. et al., *Helv. Chim. Acta*, 1995, **78**, 1511-1524 (synth)

Takahata, H. et al., *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1799-1801 (synth)

McAlonan, H. et al., *Tet. Lett.*, 2000, **41**, 5411-5414 (synth)

N-(3-Decenyl)sulfamic acid D-134

$\text{H}_3\text{C}(\text{CH}_2)_5\text{CH}=\text{CHCH}_2\text{CH}_2\text{NHSO}_3\text{H}$

$\text{C}_{10}\text{H}_{21}\text{NO}_3\text{S}$ 235.347

(Z)-form [1040630-91-8]


Isol. from *Daphnia pulex*. Kairomone.

Yasumoto, K. et al., *Chem. Pharm. Bull.*, 2008, **56**, 133-136 (isol, pmr, cmr)

Decumbenine


D-135

8-(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)furo[3,4-e]-1,3-benzodioxol-6(8H)-one, 9CI [76733-83-0]




C₂₀H₁₇NO₆ 367.357Alkaloid from *Corydalis decumbens* (Papaveraceae).Chu, T.-Y. *et al.*, *CA*, 1981, **94**, 109170x (*isol. uv, ir, pmr*)**Decumbenine B** **D-136**

5-(1,3-Dioxolo[4,5-f]isoquinolin-8-yl)-1,3-benzodioxole-4-methanol, 9CI. 3-(2-Hydroxymethyl-3,4-methylenedioxyphenyl)-5,6-methylenedioxyisoquinoline [164991-68-8]

C₁₈H₁₃NO₅ 323.304Alkaloid from roots of *Corydalis decumbens* (Papaveraceae). Needles (EtOAc/petrol). Mp 222-224°.Zhang, J.-S. *et al.*, *Phytochemistry*, 1995, **39**, 435 (*isol. uv, ir, pmr, cmr, ms, struct*)Chiba, K. *et al.*, *J.C.S. Perkin 1*, 1998, 2939-2942 (*synth*)Xu, X.-Y. *et al.*, *Tetrahedron*, 1998, **54**, 14179-14188 (*synth, pmr, cmr*)Roesch, K.R. *et al.*, *J.O.C.*, 2002, **67**, 86-94 (*synth*)Wada, Y. *et al.*, *Eur. J. Org. Chem.*, 2007, 4320-4327 (*synth*)**Decursivine** **D-137**


[474780-76-2]



Relative Configuration

C₂₀H₁₆N₂O₄ 348.357Similar to Serotobenine, S-252. Alkaloid from the leaves of *Rhaphidophora decursiva*. Antimalarial agent. Light yellow prisms (Me₂CO). [α]_D²⁰ +299 (c, 0.02 in MeOH), λ_{max} 207 (log ε 4.61); 235 (log ε 4.08); 289 (log ε 3.2) (MeOH).Zhang, H. *et al.*, *Pharm. Biol.*, 2002, **40**, 221-224 (*isol. pmr, cmr, ms*)Leduc, A.B. *et al.*, *Eur. J. Org. Chem.*, 2007, 237-240 (*synth*)**Decussine** **D-138**

1,2,3,8-Tetrahydro-1,8-dimethyl-1,7b,10-triazabenzof[5,6]cyclohepta[1,2,3-jk]fluorene, 9CI

C₂₀H₁₉N₃ 301.39**(R)-form** [75375-52-9]Alkaloid from stem bark of *Strychnos dale*, *Strychnos floribunda* and *Strychnos elaeocarpa* (Loganiaceae). Shows muscle-relaxant activity. Yellow prisms (MeOH). Mp 203-205°. Intense blue-green to yellow-green fluor. Brown col. with Ce(IV).

▶ XX8470000

3α,14-Dihydro: **Mostueine**. 3,14-Dihydrodecussine

[76328-62-6]

C₂₀H₂₁N₃ 303.406Alkaloid from *Strychnos dale*, *Strychnos decussata* and *Mostuea brunonis* (Loganiaceae). Shows weak muscle-relaxant activity. Brown cryst. (EtOH). Mp 78-82°. [α]_D +196. Dehydrogenates to Decussine on storage. The Mp given refers to Dihydrodecussine and the opt. rotn. to Mostueine. Incorrect rel. config. originally assigned.5,6-Didehydro: **Rouhamine**C₂₀H₁₇N₃ 299.374Alkaloid from stem bark of *Strychnos decussata* and from *Strychnos floribunda* (Loganiaceae). Readily formed by dec. of Decussine, D-138. Yellow needles (MeOH/Et₂O). Mp 205-210°. Red. col. with Ce(IV).10-Hydroxy, 3,14-dihydro: **10-Hydroxy-3,14-dihydrodecussine**


[79034-10-9]

C₂₀H₂₁N₃O 319.405Alkaloid from stem bark of *Strychnos decussata*. Pale brown solid.

[79082-57-8, 87585-02-2, 87585-01-1]

Onanga, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1980, 191 (*Mostueine, isol*)Rolfsen, W.N.A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 415-421 (*3,14-Dihydrodecussine, 10-Hydroxy-3,3,4-dihydrodecussine*)Verpoorte, R. *et al.*, *Planta Med.*, 1981, **42**, 32-36 (*Rouhamine, isol. uv, ms, struct*)McGee, L.R. *et al.*, *Tet. Lett.*, 1984, **25**, 2115 (*synth*)Rey, A.W. *et al.*, *Can. J. Chem.*, 1992, **70**, 2922-2928 (*Mostueine, synth, abs config*)**2-Decyl-4-hydroxyquinoline** **D-139**

2-Decyl-4-(1H)-quinolinone. 2-Decyl-4-quinolinol, 9CI [857758-85-1]

C₁₉H₂₇NO 285.428Metab. of *Pseudomonas aeruginosa*.

N-Oxide: 2-Decyl-4(1H)-quinolinone N-oxide

C₁₉H₂₇NO₂ 301.428Metab. of *Pseudomonas aeruginosa*. No CAS no. found to 2007.


1',2'-Didehydro(E-): 2-(1-Decenyl)-4(1H)-quinolinone. 2-(1-Decenyl)-4-hydroxyquinoline

C₁₉H₂₅NO 283.413Metab. of *Pseudomonas aeruginosa*. No CAS no. found to 2007.

1',2'-Didehydro(Z-):

C₁₉H₂₅NO 283.413Metab. of *Pseudomonas aeruginosa*. No CAS no. found to 2007.Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur, oxide, 1',2'-didehydro, ms*)**Deethylbophyllidine** **D-140**


Methyl 2,3-didehydro-E,20,21-trinoraspido-spermidine-3-carboxylate, 9CI [74170-69-7]



Absolute Configuration

C₁₈H₂₀N₂O₂ 296.368Alkaloid from the bark of *Tabernaemontana albiflora* and from *Anacamptis disticha*. Amorph. [α]_D +444 (c, 1 in CHCl₃). λ_{max} 226 (4.15); 295 (log ε 4.08); 326 (4.44) (EtOH).

[77080-77-4 (±)-form]

Kan, C. *et al.*, *Tet. Lett.*, 1980, **21**, 55-58 (*ir, pmr, ms, struct*)Kuehne, M.E. *et al.*, *J.O.C.*, 1981, **46**, 2002-2009 (*synth, uv, ir, pmr, ms*)Barsi, M.-C. *et al.*, *Chem. Comm.*, 1985, 88-89 (*synth*)Catena, J. *et al.*, *Tet. Lett.*, 1994, **35**, 4433-4436 (*synth, cmr*)Fernández, J.-C. *et al.*, *Chem. Comm.*, 1995, 2317-2318 (*synth*)Bonjoch, J. *et al.*, *J.O.C.*, 1996, **61**, 7106-7115 (*synth*)**O-Deethyltalaroconvolutin C** **D-141**C₃₂H₄₅NO₅ 523.711

Parent compd. not isol.


4'-Et ether: **Talaroconvolutin C**C₃₄H₄₉NO₅ 551.765Prod. by *Talaromyces convolutus*.Antifungal agent. Amorph. powder. [α]_D²⁰ -56 (c, 0.03 in CHCl₃). Possible artifact. λ_{max} 224 (log ε 3.87); 262 (log ε 3.52) (cyclohexane).4',5'-Isopropylidene: **Talaroconvolutin D**C₃₅H₄₉NO₅ 563.776Prod. by *Talaromyces convolutus*.Antifungal agent. Amorph. powder. [α]_D²⁰ -34 (c, 0.04 in EtOH). Possible artifact. λ_{max} 226 (log ε 3.92); 276 (log ε 3.44); 326 (log ε 2.79) (EtOH).Suzuki, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 768-772

Deferoxamine, INN, USAN D-142

N'-[5-[4-[5-(Acetylhydroxyamino)-pentyl]amino]-1,4-dioxobutyl]hydroxyamino]pentyl]-*N*-(5-aminopentyl)-*N*-hydroxybutanediamide, 9CI.

Desferrioxamine, BAN, Desferrin. NSC 527604

[70-51-9]



$\text{C}_{25}\text{H}_{48}\text{N}_6\text{O}_8$ 560.69

Isol. from *Streptomyces pilosus*. Used as 4mM aq. soln. for photometric detn. of V(I) (λ_{max} 480 nm, ϵ 3150, pH 1.3), Fe(III). Immunosuppressive agent. Siderophore. Shows anti-HIV and antineoplastic activity. Chelating agent for iron mobilisation with low oral toxicity; used in the treatment of β -thalassaemia and as an antidote to iron poisoning; also chelates aluminium, of interest with respect to treatment of Alzheimer's disease. Cryst. + $1\text{H}_2\text{O}$ (EtOH aq.). Sol. H_2O . Mp 138-140°. $\text{pK}_{\text{a}1}$ 8.39; $\text{pK}_{\text{a}2}$ 9.03; $\text{pK}_{\text{a}3}$ 9.7; $\text{pK}_{\text{a}4}$ 11 (20°, 0.1M NaNO₃).
 ▶ Long term treatment may cause visual and hearing disturbances. LD₅₀ (mus, orl) 1340 mg/kg. UG5300000

Hydrochloride: Deferoxamine hydrochloride, USAN, Ba 29837
 [1950-39-6]
 Mp 172-175°.

Methanesulfonate: Desferrioxamine mesilate, BAN, JAN, Desferal. Ba 33112.
 Ciba 33112
 [138-14-7]
 Cryst. (EtOH aq.). Mp 148-149°.


▶ UG5310000

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118; 2129; 1963, **46**, 1385 (*isol, synth*)
 Prelog, V. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 631 (*synth*)
 Bownern, N. *et al.*, *Aust. J. Biol. Sci.*, 1984, **62**, 743-754 (*activity*)
 Luterotti, S. *et al.*, *Acta Pharm. Jugosl.*, 1986, **36**, 341; *CA*, **105**, 237518z (*dem, Fe*)
 Luterotti, S. *et al.*, *Analyst (London)*, 1986, **111**, 1163 (*detn, V*)
 Bergeron, R.J. *et al.*, *J.O.C.*, 1988, **53**, 3131 (*synth, bibl*)
 Allain, P. *et al.*, *Drugs of Today (Barcelona)*, Suppl.A, 1992, **28**, 89 (*rev*)
 Hider, R.C. *et al.*, *Chemistry of Iron*, (Silver, J, ed.) Blackie, 1993, 275 (*use*)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 676
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DAK200; DAK300

Deformylcoryzeamine

D-143

[164230-53-9]




$\text{C}_{43}\text{H}_{52}\text{N}_4\text{O}_6$ 720.907

Alkaloid from leaves of *Hunteria zeylanica* (Apocynaceae). Amorph. powder. $[\alpha]_{\text{D}}^{25}$ -90.4 (c, 0.24 in MeOH).

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1957 (*isol, pmr, cmr, struct*)

N-Deformyldichotamine

D-144



Absolute Configuration

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_3$ 340.421
 Mp 197-198°.

N-Formyl: Dichotamine

[6793-25-5]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Vallislea dichotoma* (Apocynaceae). Cryst. (EtOAc or MeOH aq.). Mp 262-263° dec. (254-257°). $[\alpha]_{\text{D}}$ -116 (CHCl₃).

O-De-Me, N-Ac: 18-Oxohaplocidine

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph.

O-De-Me, N-propanoyl: Cimicine

[7096-82-4]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Haplophyton cimicidum* (Apocynaceae). Cryst. (EtOH). Mp 229-231°. $[\alpha]_{\text{D}}$ +113 (CHCl₃).

11-Methoxy, O¹²-de-Me, N-propanoyl: Cimicidine

[7096-81-3]

$\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_5$ 412.485

Insecticidal alkaloid from *Haplophyton cimicidum* (Apocynaceae). Mp 268-270° dec. $[\alpha]_{\text{D}}^{25}$ +23 (c, 1 in CHCl₃).

11-Methoxy, O¹²-de-Me, N-propanoyl, hydrochloride: Mp 247-249° dec.

10,11-Dimethoxy, N-propanoyl: 18-Oxo-O-methylaspidoalbine

$\text{C}_{25}\text{H}_{32}\text{N}_2\text{O}_6$ 456.538

Alkaloid from *Aspidosperma exalatum* (Apocynaceae). Cryst. (MeOH). Mp 184-185°. $[\alpha]_{\text{D}}^{31}$ +96 (CHCl₃).

10, 11-Dimethoxy, O¹²-de-Me, N-propanoyl: 18-Oxoaspidoalbine

[11040-97-4]

$\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_6$ 442.511

Alkaloid from *Aspidosperma exalatum* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 208-209°.

Snyder, H.R. *et al.*, *J.A.C.S.*, 1954, 4601 (*Cimicine, isol*)

Holker, J.S.E. *et al.*, *J.O.C.*, 1959, **24**, 314 (*Dichotamine, isol, uv, ir*)

Brown, K.S. *et al.*, *Tet. Lett.*, 1963, 1731 (*Dichotamine, uv, pmr, ms, struct*)

Brown, K.S. *et al.*, *J.A.C.S.*, 1966, **88**, 4984 (*18-Oxoaspidoalbine, 18-Oxo-O-methylaspidoalbine*)

Cava, M.P. *et al.*, *Can. J. Chem.*, 1973, **51**, 3102 (*Cimicine, Cimicidine*)


Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694 (*18-Oxohaplocidine*)

Deformyltalbotinic acid

D-145

methyl ester

Methyl 2,3,3a,4,5,6-hexahydro-5-[1-(hydroxymethyl)-1-propenyl]-1H-indolo[3,2,1-de][1,5]naphthyridine-6-carboxylate, 9CI
 [30809-33-7]



$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_3$ 340.421

Alkaloid from the leaves of *Pleiocarpa talbotii* (Apocynaceae). Needles (MeOH/Et₂O). Mp 240° dec. $[\alpha]_{\text{D}}^{22}$ +111 (c, 0.181 in MeOH).


N⁶,O-Di-Ac:

Cryst. (Et₂O/pentane). Mp 99-100°.

Pinar, M. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 15; 1973, **56**, 2719 (*isol, uv, ms, ir, pmr, struct, synth*)

Sambucus nigra Degraded cyanogenic glycosides

D-146



$\text{C}_{16}\text{H}_{19}\text{NO}_8$ 353.328

Degraded cyanogenic glucoside. Isol. from *Sambucus nigra* (elderberry). $[\alpha]_{\text{D}}$ +15 (c, 0.6 in CHCl₃). λ_{max} 221 (ϵ 3150); 278 (ϵ 2170) (EtOH).

2-Epimer:

$\text{C}_{16}\text{H}_{19}\text{NO}_8$ 353.328

Isol. from *Sambucus nigra* (elderberry). $[\alpha]_{\text{D}}$ -21 (c, 0.3 in CHCl₃). λ_{max} 225 (ϵ 3160); 276 (ϵ 2190) (EtOH).

2'-Epimer:

$\text{C}_{16}\text{H}_{19}\text{NO}_8$ 353.328


Isol. from *Sambucus nigra* (elderberry). $[\alpha]_{\text{D}}$ -3 (c, 0.5 in CHCl₃). λ_{max} 224 (ϵ 3130); 279 (ϵ 2130) (EtOH).

Della Greca, M. *et al.*, *Tet. Lett.*, 2000, **41**, 6507-6510

Della Greca, M. *et al.*, *Nat. Prod. Res.*, 2003, **17**, 177-181 (*synth*)

Dehassiline

D-147



$\text{C}_{19}\text{H}_{23}\text{NO}_4$ 329.395

The synthetic prod. was shown to be different from the nat. prod. (1997). Nmr data suggests that the struct. needs to be re-examined (1998). Trivial name is based

on an incorrect spelling (*Dehassia*) of the genus.


(S)-form [142717-64-4]

Alkaloid from bark of *Dehaasia kurzii* (Lauraceae). Amorph. solid. $[\alpha]_D^{27} +147.3$ (c, 0.01M in CHCl_3).

Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1991, **62**, 261 (*isol, uv, ir, pmr, cmr, cd, struct*)
Takaba, K. *et al.*, *Heterocycles*, 1997, **45**, 1111
Watanabe, A. *et al.*, *Heterocycles*, 1998, **48**, 1623-1630 (*synth, pmr, cmr*)

Dehatrine **D-148**

1',2'-Didehydro-2'-demethylpceanthine, 8CI



$\text{C}_{37}\text{H}_{38}\text{N}_2\text{O}_6$ 606.717

(R)-form

Alkaloid from the trunk of *Dehaasia triandra*. Also *isol.* from the Indonesian medicinal plant *Beilschmiedia madang* (Lauraceae). Shows antimalarial and antimicrobial props. Needles + H_2O (EtOH). Mp 158-160° (146-148°). $[\alpha]_D^{27} +27$ (c, 1.0 in CHCl_3). Exists as a 1:1 mixt. of two rotamers in the solid state. λ_{max} 223 (ε 31600); 281 (ε 6310); 310 (ε 3160) (MeOH) (Derap).

(ξ)-form

3',4'-Didehydro: Fangchirine

Appears to be currently unknown as an alkaloid.

3',4'-Didehydro, α'-oxo: Oxofangchirine

[102516-53-0]

$\text{C}_{37}\text{H}_{34}\text{N}_2\text{O}_7$ 618.685

Alkaloid from the roots of *Stephania tetrandra* (Menispermaceae).

Inubushi, Y. *et al.*, *J.C.S.(C)*, 1969, 1547 (*synth*)

Hu, T. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 29; *CA*, **104**, 221975q (*Oxofangchirine*)

Tsai, I.L. *et al.*, *Kaohsiung J. Med. Sci.*, 1989, **5**, 131-145 (*activity*)

Lu, S.-T. *et al.*, *Phytochemistry*, 1989, **28**, 615 (*isol, uv, ir, pmr, ms, struct*)


Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 997 (*isol, ir, pmr, cmr, cryst struct*)

Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*pmr, cmr*)

Dehatriphine **D-149**

9-O-(8-Isocorydiny)-N-methylaurotetanine


[173268-83-2]



$\text{C}_{40}\text{H}_{44}\text{N}_2\text{O}_8$ 680.796

The first ether-linked bisaporphine. Alkaloid from leaves of *Dehaasia triandra*. Amorph. powder. $[\alpha]_D^{23} +13$ (c, 1.0 in MeOH).

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 55 (*isol, uv, ir, cd, pmr, cmr, ms, struct*)

Dehydroapocavidine **D-150**

$\text{C}_{20}\text{H}_{18}\text{NO}_4^{\oplus}$ 336.367

Quaternary alkaloid from *Corydalis cava*. Orange-yellow clusters (MeOH)(as iodide). Mp 300-308° dec. (iodide).

Me ether: Dehydrocavidine

[57392-71-9]

$\text{C}_{21}\text{H}_{20}\text{NO}_4^{\oplus}$ 350.393

Quaternary alkaloid from *Corydalis saxicola* and *Corydalis meifolia*. Mp 273-275° (counterion not specified). λ_{max} 269 (log ε 4.39); 350 (log ε 4.33) (EtOH).

O³-De-Me, 2-Me ether: Dehydroisoapocavidine

$\text{C}_{20}\text{H}_{18}\text{NO}_4^{\oplus}$ 336.367

Alkaloid from *Corydalis saxicola*. Red needles (as hydroxide). Mp 281-283° dec. (hydroxide). λ_{max} 278 (log ε 4.31); 347 (log ε 3.69) (MeOH) (hydroxide).

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261 (*isol, uv, struct*)

Ke, M. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1982, **24**, 289-291; *CA*, **97**, 13341n (*Dehydrocavidine*)

Bhakuni, D.S. *et al.*, *J. Indian Chem. Soc.*, 1984, **61**, 1016 (*Dehydroapocavidine, Dehydrocavidine*)


Halbsguth, C. *et al.*, *Planta Med.*, 2003, **69**, 305-309 (*isol*)

Li, H.-L. *et al.*, *Chem. Biodiversity*, 2008, **5**, 777-783 (*Dehydroisoapocavidine*)

Cheng, X. *et al.*, *Chem. Biodiversity*, 2008, **5**, 1335-1344 (*Dehydrocavidine*)

Dehydroazasirosterol **D-151**

[145401-26-9]




$\text{C}_{47}\text{H}_{59}\text{NO}_6$ 733.986

Metab. of a *Sirococcus* sp. Amorph. powder.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1992, **70**, 1905 (*isol, pmr, cmr*)

Dehydrobatzelladine C **D-152**

[763082-99-1]



$\text{C}_{27}\text{H}_{46}\text{N}_6\text{O}_2$ 486.699


Alkaloid from the sponge *Monanchora arbuscula*. Pale brown gum. λ_{max} 206 (ε 8500); 258 (ε 7500); 303 (ε 1600) (MeOH).

Braekman, J.C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 193-196

Collins, S.K. *et al.*, *Org. Lett.*, 2004, **6**, 1253-1255 (*synth*)

Dehydrobufotenine **D-153**

[17232-69-8]



$\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$ 202.255

Main constit. of the secretion of the toad *Bufo marinus* and from *Bufo bufo bufo*.

► Convulsant lethal dose (mus, scu) 6 mg/kg. UY9454800

Hydrochloride: Mp 237-238° dec.

O-Hydrogen sulfate inner salt: Bufothionine

[475-39-8]

$\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}_4\text{S}$ 282.32

Constit. of the Japanese toad. Mp 250° dec.

Märki, F. *et al.*, *J.A.C.S.*, 1961, **83**, 3341 (*uv, pmr, struct*)

Robinson, B. *et al.*, *Proc. Chem. Soc., London*, 1961, 310 (*pmr, struct*)

Daly, J.W. *et al.*, *Science (Washington, D.C.)*, 1967, **156**, 970 (*tox*)


Gannon, W.F. *et al.*, *Tet. Lett.*, 1967, 1531 (*synth*)

Honda, K. *et al.*, *Acta Cryst. C*, 1991, **47**, 1506 (*cryst struct, Bufothionine*)

Peat, A.J. *et al.*, *J.A.C.S.*, 1996, **118**, 1028 (*synth*)

Dehydrocorydaline **D-154**

[30045-16-0]



$\text{C}_{22}\text{H}_{24}\text{NO}_4^{\oplus}$ 366.436

Quaternary alkaloid from *Corydalis*

ambigua and several other *Corydalis* spp. and from *Berberis floribunda* (Papaveraceae, Berberidaceae). Shows strong gastric antisecretory and antiulcer activity. Uterine contractant. Mp 237° (as nitrate).

▶ DR9868000

O²-De-Me: 13-Methylcolumbamine


[137760-67-9]

C₂₁H₂₂NO₄[⊕] 352.409Alkaloid from *Corydalis solida* ssp. *brachyloba*.**O³-De-Me: Dehydrocorybulbine**


[59870-72-3]

C₂₁H₂₂NO₄[⊕] 352.409Quaternary alkaloid from *Corydalis turtchanovii* f. *yahusuo*. Yellow solid (as chloride). Mp 188-190° (chloride).Späth, E. *et al.*, *Ber.*, 1921, **54**, 3074 (*struct*)Koepfli, J.B. *et al.*, *J.C.S.*, 1928, 2989 (*synth*)Tani, C. *et al.*, *Yakugaku Zasshi*, 1970, **90**, 407 (*Dehydrocorybulbine*)Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261 (*isol*)Bhakuni, D.S. *et al.*, *Alkaloids (Academic Press)*, 1986, **28**, 168 (*pharmacol*)Sener, B. *et al.*, *J. Chem. Soc. Pak.*, 1991, **13**, 63; *CA*, **116**, 3548j (*13-Methylcolumbamine*)**Dehydrocrambine A***Dehydrocrambescin A*


[628727-34-4]

C₂₄H₄₂N₆O₂ 446.635Alkaloid from the sponge *Monanchora* sp. Brown gum. [α]_D²⁰ -12.1 (c, 0.09 in MeOH). λ_{max} 206 (log ε 3.83); 259 (log ε 3.53); 289 (log ε 3.18) (MeOH).Chang, L.C. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1490-1494 (*isol, pmr, cmr*)**Dehydrocentrine**

[19843-03-9]

C₂₀H₁₉NO₄ 337.374Alkaloid from *Ocotea macropoda*, *Cissampelos pareira*, *Stephania brachyandra* and *Glaucium vitellinum* (Lauraceae, Menispermaceae, Papaveraceae). Golden-yellow needles (CHCl₃). Mp 218°.**O¹⁰-De-Me: Dehydrophanostenine**

[79642-39-0]

C₁₉H₁₇NO₄ 323.348Alkaloid from *Stephania sasakii* (Menispermaceae). Mp 198-200°.Cava, M.P. *et al.*, *Tet. Lett.*, 1968, 2437 (*uv, pmr, struct*)Dwuma-Badu, D. *et al.*, *Phytochemistry*, 1975, **14**, 2520 (*isol, uv, ir, pmr, ms*)Shafiee, A. *et al.*, *J. Nat. Prod.*, 1978, **41**, 657; 1979, **42**, 174 (*isol*)Kunimoto, J. *et al.*, *Yakugaku Zasshi*, 1981, **101**, 431-436 (*Dehydrophanostenine*)Castedo, L. *et al.*, *An. Quim., Ser. C*, 1982, **78**, 103 (*cmr*)Chen, Y. *et al.*, *Zhongcaoyao*, 1982, **13**, 1; *CA*, **97**, 178723d (*isol*)**Dehydrodopamine****D-157***2-(3,4-Dihydroxyphenyl)ethenamine. Dihydrodopamine*C₈H₉NO₂ 151.165**(E)-form***N-Ac: N-[2-(3,4-Dihydroxyphenyl)ethenyl]acetamide, 9CI. N-Acetyl-1,2-didehydrodopamine. N-Acetyldehydrodopamine*

[83104-76-1]

C₁₀H₁₁NO₃ 193.202

Precursor in insect cuticular sclerotization. Cryst. (AcOH aq.). Mp 197-198°.

N-β-Alanyl: 3-Amino-N-[2-(3,4-dihydroxyphenyl)ethenyl]propanamide, 9CI. 1,2-Didehydro-N-β-alanyldopamine. N-β-Alanyldidehydrodopamine


[158018-58-7]

C₁₁H₁₄N₂O₃ 222.243

Precursor in insect cuticular sclerotization. No phys. props. reported.

Ramamurthy, B. *et al.*, *Synthesis*, 1987, 523 (*N-Ac, synth*)Dali, H. *et al.*, *Org. Prep. Proced. Int.*, 1988, **20**, 191-195 (*N-Ac, synth*)Ricketts, D. *et al.*, *J. Biol. Chem.*, 1994, **269**, 22217-22221 (*N-β-alanyl, synth, biochem*)**Dehydroevodiamine****D-158***5,7,8,13-Tetrahydro-14-methyl-5-oxoindolo[2',3':3,4]pyrido[2,1-b]quinazolinium hydroxide inner salt, 9CI. 13,13b-Dehydroevodiamine*

[67909-49-3]

C₁₉H₁₅N₃O 301.347CAS numbering shown. Alkaloid from the unripe fruit of *Evodia rutaecarpa* (Rutaceae). Mp 189°.*Hydrochloride:*Yellow cryst. + 1/4H₂O. Mp 208-209°.*2-Methoxy:*C₂₀H₁₇N₃O₂ 331.373Alkaloid from the stem bark of *Araliopsis tabouensis*. Amorph. yellow powder (as hydrochloride). Represented as the alternative Δ^{5(13a)} form. λ_{max} 212 (log ε 1.91); 284 (log ε 0.49); 362 (log ε 0.7); 376 (log ε 0.35) (MeOH) (hydrochloride).*10-Methoxy: Hortiamine. 8,14-Dihydro-10-methoxy-14-methylindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI*


[477-40-7]

C₂₀H₁₇N₃O₂ 331.373Alkaloid from the bark of *Hortia arborea* and *Hortia braziliiana* (Rutaceae). Hypotensive agent. Red-orange needles (C₆H₆ or CHCl₃/C₆H₆). Mp 209° dec.*10-Methoxy; hydrochloride:*Cryst. + 1H₂O (EtOH aq.). Mp 243° dec. Sublimation at 230-270°/0.5 mm yields Hortiacine, H-361.*10-Methoxy; methiodide:*

Cryst. (EtOH aq.). Mp 208-209°.

Pachter, I.J. *et al.*, *J.A.C.S.*, 1960, **82**, 5187;1961, **83**, 535 (*Hortiamine, isol, uv, struct, synth*)Danieli, B. *et al.*, *Heterocycles*, 1978, **9**, 803 (*synth*)King, C.L. *et al.*, *J. Nat. Prod.*, 1980, **43**, 577 (*isol, ir, pmr*)Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth*)Christopher, E. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2914-2918 (*2-methoxy*)**Dehydroglaucine****D-159***5,6-Dihydro-1,2,9,10-tetramethoxy-6-methyl-4H-dibenzo[de,g]quinoline, 9CI*

[22212-26-6]

C₂₁H₂₃NO₄ 353.417Alkaloid from *Glaucium flavum* (incl. varieties), heartwood and discoloured sapwood of *Liriodendron tulipifera* and *Thalictrum* sp. Antimicrobial agent. Pale yellow prisms (EtOH). Mp 133-134°. λ_{max} 264 (ε 47000); 293 (ε 17200); 341 (ε 10600) (EtOH).*N-De-Me: 6a,7-Dehydronorglaucine*

[39945-38-5]

Synthetic. Cryst. (Et₂O/MeOH). Mp 180-183°.*O¹-De-Me: Dehydrothaliporphine*

[41823-69-2]

Synthetic. Mp 157-160° dec.

O²-De-Me: Dehydropredicentrine

[91794-10-4]

C₂₀H₂₁NO₄ 339.39Isol. from *Polyalthia cauliflora*. Cryst. (Me₂CO). Mp 198-199°. λ_{max} 215 (log

ε 4.14); 243 (sh) (log ε 4.28); 262 (log ε 4.49); 270 (sh) (log ε 4.45); 294 (sh) (log ε 4.04); 329 (log ε 3.89); 380 (sh) (log ε 3.31) (no solvent reported).

O¹⁰-De-Me: Dehydrolirioferine

[75652-88-9]

Synthetic. Cryst. (EtOAc). Mp 140-142° dec. Unstable.

O¹,O⁹-Di-de-Me: Dehydroisoboldine

[53729-86-5]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Nandina domestica* (Nandinaceae).

O²,O⁹-Di-de-Me: Dehydroboldine

[91599-23-4]

C₁₉H₁₉NO₄ 325.363

Isol. from *Peumus boldus* (boldo). Mp 178-179°. λ_{max} 263 (log ε 4.72); 327 (log ε 4.06) (EtOH).

7-Hydroxy: 7-Hydroxydehydroglauicine

[218629-68-6]

C₂₁H₂₃NO₅ 369.416

Alkaloid from *Annona purpurea* (soncoya). Antiplatelet aggregation agent. Green needles (CHCl₃). Mp 242-243°. λ_{max} 222 ; 260 ; 335 (EtOH).

7-Hydroxy, 4,5-didehydro, N-de-Me, N-formyl: N-Formyl-4,5,6a,7-tetra-dehydro-7-hydroxyglauicine

[959850-98-7]

C₂₁H₁₉NO₆ 381.384

Alkaloid from the bark of *Friesodielsia obovata*. Yellow cryst. Mp 198-200°. λ_{max} 217 ; 237 ; 283 ; 308 ; 319 (MeOH).

Δ^{6,6a}-Isomer, N-de-Me: 6,6a-Dehydronorglauicine

[52309-70-3]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Glaucium flavum* (Papaveraceae). First member of a small subgroup of aporphine-type alkaloids with a conjugated azomethine group. λ_{max} 242 (sh) (ε 28300); 260 (ε 47200); 270 (ε 43400); 335 (ε 10800); 381 (ε 3330) (MeOH).

Kiryakov, H.G. *et al.*, *Chem. Ind. (London)*, 1968, 1807-1808 (uv, ir, pmr, struct)

Cava, M.P. *et al.*, *J.O.C.*, 1970, **35**, 175-179 (synth, uv)

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 1149-1153; *CA*, **82**, 14024r (Dehydroisoboldine)

Hufford, C.D. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 789-792 (isol, uv, pmr)

Chen, C.-L. *et al.*, *Phytochemistry*, 1976, **15**, 547-550; 1161-1167 (isol, pmr, ms)

Castedo, L. *et al.*, *Heterocycles*, 1980, **14**, 1135-1138 (O¹-de-Me, O¹⁰-de-Me, synth)

Castedo, L. *et al.*, *An. Quim., Ser. C*, 1982, **78**, 103-107 (7-Me, synth, cmr)

Jossang, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 504-513 (Dehydropredicetrine)

Urzua, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 525-526 (Dehydroboldine)

Lenz, G.R. *et al.*, *J.C.S. Perkin 1*, 1984, 1273-1277 (6,6a-Dehydronorglauicine, synth)


Gupta, S. *et al.*, *Synth. Commun.*, 1989, **19**, 393-401 (synth)

Chang, F.-R. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1457-1461 (7-Hydroxydehydroglauicine)

Joseph, C.C. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1009-1015 (Formyltetradehydro-7-hydroxynorglauicine)

Dehydroisolongistrobine

[23507-01-9]



C₁₇H₁₇N₃O₃ 311.34

Alkaloid from the aerial parts of *Macrorungia longistrobis* (preferred genus name *Anisotes*) (Acanthaceae). Needles (Me₂CO/hexane). Mp 131°. λ_{max} 258 (ε 17350) (EtOH). λ_{max} 235 (ε 14350) (1% HCl/EtOH).

Dihydro: Isolongistrobine

[23544-52-7]

C₁₇H₁₉N₃O₃ 313.355

Alkaloid from the aerial parts of *Macrorungia longistrobis* (preferred genus name *Anisotes*) (Acanthaceae). Yellowish needles (Me₂CO/hexane). Mp 134-139° (132-136°). A hydroxylactam having one of the maleimide carbonyl groups reduced to OH. λ_{max} 253 (ε 16750) (EtOH). λ_{max} 235 (ε 14400) (1% HCl/EtOH).


Arndt, R.R. *et al.*, *Tetrahedron*, 1969, **25**, 2767-2799 (Dehydroisolongistrobine, Isolongistrobine, isol, ir, uv, pmr, ms)

Wuonola, M.A. *et al.*, *Tetrahedron*, 1976, **32**, 1085-1095 (Dehydroisolongistrobine, Isolongistrobine, struct, synth, ir, uv, pmr, ms)

5,6-Dehydrolupanine

D-161

5,6-Didehydrolupanine, 8CI. 2-Oxo-Δ⁵-dehydroparteine. Alkaloid BNC 1
[32101-29-4]



C₁₅H₂₂N₂O 246.352

Alkaloid from *Lupinus densiflorus* and *Lupinus bicolor* ssp. *microphyllus* and from *Sophora secundiflora* seeds (mescal beans)(Fabaceae). Liq. Unstable to hydrolysis.

11-Epimer: 5,6-Dehydro-α-isolupanine

C₁₅H₂₂N₂O 246.352

Alkaloid from *Lupinus bicolor* ssp. *microphyllus* (Fabaceae).

Cho, Y.D. *et al.*, *Arch. Mass Spectral Data*, 1971, **2**, 328 (ms)

Cho, Y.D. *et al.*, *Can. J. Chem.*, 1971, **49**, 265 (isol, uv, ms, pmr, struct)

Hatfield, G.M. *et al.*, *J. Nat. Prod.*, 1977, **40**, 374; 1980, **43**, 164 (isol)


Kinghorn, A. *et al.*, *Planta Med.*, 1980, **38**, 280 (isol, ms)

D-160

Dehydronapelline

Dehydroglucicine

[110064-72-7]



C₂₂H₃₁NO₃ 357.492

Alkaloid from *Aconitum flavum*, *Aconitum japonicum* and *Aconitum yesoense* (Ranunculaceae). Mp 103.5-105°. [α]_D¹⁶ +78.3 (c, 0.48 in EtOH).

12-Ac: Subdesculine. 12-Acetyldehydroglucicine

[62517-45-7]

C₂₄H₃₃NO₄ 399.529

Alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Mp 134-135° dec. (as perchlorate). [α]_D¹⁹ +7.3 (c, 0.3 in CHCl₃).

15-Ac: Dehydroglucidusculine

[99815-76-6]

C₂₄H₃₃NO₄ 399.529

Alkaloid from the roots of *Aconitum yesoense* var. *macroysesense* (formerly known as *Aconitum lucidusculum*) (Ranunculaceae). Mp 186-189°. [α]_D¹¹ +2.6 (c, 1.0 in EtOH). 15-Config. shown as α- in the paper, but Lucidusculine, in N-30X, with which Dehydroglucidusculine was correlated, has 15β-config.

Di-Ac: 12-Acetyldehydroglucidusculine

[112579-73-4]

C₂₆H₃₅NO₅ 441.566

Alkaloid from the rhizomes of *Aconitum yesoense* var. *macroysesense* (Ranunculaceae). Amorph. [α]_D +9.3 (c, 0.43 in EtOH).

N-de-Et, 15-Ac: N-Deethyldehydroglucidusculine

[99815-75-5]

C₂₂H₂₉NO₄ 371.475

Alkaloid from the roots of *Aconitum yesoense* var. *macroysesense* (Ranunculaceae). Amorph. powder. [α]_D¹⁷ -9.6 (c, 0.27 in EtOH).

12-Ketone: Songoramine

[23179-78-4]

C₂₂H₂₉NO₃ 355.476

Alkaloid from *Aconitum karakolicum* and roots of *Aconitum songoricum* (Ranunculaceae). Cryst. (Me₂CO). Mp 211-212°. [α]_D²⁶ -44.2 (c, 0.27 in CHCl₃). λ_{max} 295 (log ε 2.6) (no solvent reported).

12-Ketone, 15-Ac: 15-Acetylsongoramine

[76971-23-8]

C₂₄H₃₁NO₄ 397.513

Alkaloid from roots of *Aconitum songoricum* (Ranunculaceae).

12-Ketone, N-de-Et: Norsongoramine. N-Deethylsongoramine

[78982-16-8]

$C_{20}H_{25}NO_3$ 327.422
Alkaloid from *Delphinium thamarae*.
Cryst. (Me₂CO). Mp 286-288°.

12-Epimer: 12-Epidehydronapelline

[116197-11-6]
 $C_{22}H_{31}NO_3$ 357.492
Alkaloid from above-ground parts of *Aconitum napellus* ssp. *castellanum* (Ranunculaceae) and from *Aconitum liangshanium*. Resin or amorph. solid. $[\alpha]_D^{23} +45$ (c, 0.17 in EtOH). $[\alpha]_D^{23} +56.8$ (c, 1.02 in CHCl₃).

12-Epimer, 12-Ac: 12-Epiacetyldehydronapelline

[116171-99-4]
 $C_{24}H_{33}NO_4$ 399.529
Alkaloid from above-ground parts of *Aconitum napellus* ssp. *castellanum* (Ranunculaceae). Resin. $[\alpha]_D +25$ (c, 0.22 in EtOH).

12-Epimer, 15-Ac: 12-Epidehydro-lucidusculine

[137121-73-4]
 $C_{24}H_{33}NO_4$ 399.529
Alkaloid from *Aconitum liangshanium* (Ranunculaceae). Amorph. $[\alpha]_D^{23} -9.6$ (c, 1.2 in CHCl₃).

Yusunov, M.S. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 95-99 (Songoramine)

Zhamierashvili, M.G. et al., *CA*, 1972, **97**, 107014n (15-Acetylsongoramine)

Beshitaishvili, L.V. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 156-157 (Norsongoramine)

Wada, K. et al., *Heterocycles*, 1985, **23**, 2473-2477 (Dehydro-lucidusculine, N-Deethyldehydro-lucidusculine)

Chen, Z. et al., *Heterocycles*, 1987, **26**, 1455-1460 (Dehydronapelline)

Bando, H. et al., *Heterocycles*, 1987, **26**, 2623-2637 (12-Acetyldehydro-lucidusculine)

Bando, H. et al., *Chem. Pharm. Bull.*, 1988, **36**, 1604-1606 (Subdesculine)


De la Fuente, G. et al., *Heterocycles*, 1988, **27**, 1109-1113 (12-Epidehydronapelline, 12-Epiacetyldehydronapelline)

Takayama, H. et al., *Chem. Pharm. Bull.*, 1991, **39**, 1644-1646 (12-Epidehydro-lucidusculine)

Csupor, D. et al., *Helv. Chim. Acta*, 2006, **89**, 2981-2986 (Songoramine, synth)

Dehydronormacrostomine D-163

Preininger alkaloid
[115107-84-1]




$C_{23}H_{22}N_2O_4$ 390.438
Alkaloid from *Papaver macrostomum* (Papaveraceae). Mp 193-195°.

Mnatsakanyan, V.A. et al., *Coll. Czech. Chem. Comm.*, 1977, **42**, 1421 (isol, uv, ms, struct)

Mahboobi, S. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 275 (synth)

Dehydronuciferine D-164

[7630-74-2]



$C_{19}H_{19}NO_2$ 293.365
Alkaloid from the leaves of *Nelumbo nucifera* (East Indian lotus), *Colubrina faralaotra* ssp. *faralaotra* and *Colubrina faralaotra* subsp. *trichocarpa* (Nelumbo-naceae, Rhamnaceae). Cryst. (EtOH). Mp 130-131°.

N-De-Me: Dehydronornuciferine

[92664-95-4]
 $C_{18}H_{17}NO_2$ 279.338
Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae). Off-yellow flakes. Mp 149.5-150.5°.

N-De-Me, N-formyl: N-Demethyl-N-formyldehydronuciferine

[111017-06-2]
 $C_{19}H_{17}NO_3$ 307.348
Alkaloid from the drug Sinomeni Caulis et Rhizoma (root, root-stalk and stems of *Sinomenium acutum*) (Menispermaceae). Needles (MeOH). Mp 140.5°.

► Mutagen.**O¹-De-Me: Dehydroiridinine**

$C_{18}H_{17}NO_2$ 279.338
Alkaloid from *Annona purpurea*. Anti-platelet aggregation agent. Green needles (CHCl₃). Mp 189-190°. λ_{max} 216 ; 253 ; 312 (EtOH).

O²-De-Me, O²- α -L-rhamnopyranoside:

6a,7-Dehydrofloripavidine
 $C_{24}H_{27}NO_6$ 425.48
Alkaloid from the aerial parts of *Papaver fugax*. Amorph. solid. λ_{max} 251 ; 263 ; 320 ; 396 (MeOH).

Cava, M.P. et al., *J.O.C.*, 1970, **35**, 175 (synth, uv)

Kunitomo, J. et al., *Phytochemistry*, 1973, **12**, 699 (occur)

Guinaudeau, H. et al., *Planta Med.*, 1975, **27**, 304; 1976, **29**, 54; **30**, 201 (isol)

Castedo, L. et al., *An. Quim., Ser. C*, 1982, **78**, 103 (cmr)

Lenz, G.R. et al., *J.C.S. Perkin 1*, 1984, 1273 (Dehydronornuciferine, synth, uv, ir, pmr, cmr)

Cortes, D. et al., *J. Nat. Prod.*, 1986, **49**, 878 (Dehydronornuciferine, isol)

Nozaka, T. et al., *Chem. Pharm. Bull.*, 1987, **35**, 2844 (N-Demethyl-N-formyldehydronuciferine)


Atanes, N. et al., *J.O.C.*, 1991, **56**, 2984 (synth, Dehydronornuciferine)

Chang, F.R. et al., *J. Nat. Prod.*, 1998, **61**, 1457-1461 (Dehydroiridinine)

Sari, A. et al., *Nat. Prod. Res.*, 2004, **18**, 265-268 (6a,7-Dehydrofloripavidine)

Dehydropuntarenine D-165

[141290-70-2]




$C_{21}H_{17}NO_6$ 379.368
Alkaloid from roots of *Berberis actinacantha* (Berberidaceae).

Rahimizadeh, M. et al., *J. Sci., Islamic Repub. Iran*, 1990, **1**, 364; *CA*, **116**, 231903b

Dehydroermerine D-166

Dehydroaporheine
[36285-03-7]



$C_{18}H_{15}NO_2$ 277.322
Alkaloid from *Nelumbo nucifera* (East India lotus), *Stephania sasakii*, *Stephania micrantha*, *Stephania kwangsiensis*, *Papaver glaucum*, *Papaver rhoeas* (corn poppy), *Papaver spicatum* var. *spicatum*, *Papaver spicatum* var. *luschanii*, *Liriodendron tulipifera*, *Colubrina faralaotra* ssp. *faralaotra* and *Colubrina faralaotra* ssp. *sinuata* (Nelumbonaceae, Menispermaceae, Papaveraceae, Magnoliaceae, Rhamnaceae). Mp 88-89°.

N-De-Me: Dehydroanonaine

[41679-82-7]
 $C_{17}H_{13}NO_2$ 263.295
Alkaloid from the leaves of *Nelumbo nucifera* (East India lotus) (Nelumbonaceae).

N-De-Me, N-formyl: N-Formyldehydroanonaine

[165900-12-9]
 $C_{18}H_{13}NO_3$ 291.306
Alkaloid from woody stems of *Telictoxicum krukovii* (Menispermaceae). Cryst. (hexane/EtOAc). Mp 185-186°.

N-De-Me, N-Ac: N-Acetyldehydroanonaine

[132646-11-8]
 $C_{19}H_{15}NO_3$ 305.332
Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper). Prisms (MeOH). Mp 151-153°.

4,5-Didehydro: Didehydroermerine. Didehydroaporheine

$C_{18}H_{13}NO_2$ 275.306
Alkaloid from *Papaver urbanianum* (Papaveraceae). Amorph.


Kunitomo, J. et al., *Phytochemistry*, 1973, **12**, 699 (Dehydroanonaine)

Preininger, V. et al., *Planta Med.*, 1973, **23**, 233 (Didehydroermerine)

- Guinaudeau, H. *et al.*, *Planta Med.*, 1975, **27**, 304; 1976, **30**, 201 (*isol*)
 Ziyaev, R. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 715; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 602 (*isol, uv, ms*)
 Min, Z. *et al.*, *Yaouxue Xuebao*, 1980, **15**, 532; *CA*, **94**, 117773m (*isol*)
 Phillipson, J.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 296 (*occur*)
 Sariyar, G. *et al.*, *Plant. Med. Phytother.*, 1981, **15**, 160; *CA*, **96**, 119030g (*isol*)
 Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1981, **101**, 431; *CA*, **95**, 204236c (*isol*)
 Min, Z. *et al.*, *Yaouxue Xuebao*, 1981, **16**, 557; *CA*, **97**, 3595m (*isol*)
 Atanes, N. *et al.*, *J.O.C.*, 1991, **41**, 1425 (*Dehydroanonaine, synth*)
 Menachery, M.D. *et al.*, *Heterocycles*, 1995, **41**, 1425 (*N-Formyldehydroanonaine*)
 Chen, I.-S. *et al.*, *Phytochemistry*, 1996, **42**, 217 (*N-Acetyldehydroanonaine*)

Dehydrosaulatine D-167

[141290-71-3]



$C_{22}H_{21}NO_6$ 395.411
 Alkaloid from roots of *Berberis actinacantha* (Berberidaceae).

Rahimizadeh, M. *et al.*, *J. Sci., Islamic Repub. Iran*, 1990, **1**, 364; *CA*, **116**, 231903b


11,12-Dehydrosparteine D-168

11,12-Didehydrosparteine, *5,6-Didehydro- α -isosparteine*, *5,6-Dehydro- α -isosparteine* [2130-67-8]



$C_{15}H_{24}N_2$ 232.368
 Alkaloid identified in stems of *Templetonia egena*, prob. also present in *Lupinus polyphyllus* and *Sarothamnus scoparius* (Fabaceae). Also (descr. as 5,6-didehydro- α -isosparteine) detected spectroscopically in *Acosmium panamense*. The alkaloid from *L. polyphyllus* was originally identified as the Δ^{12} -isomer but this is prob. erroneous.


Wink, M. *et al.*, *Planta Med.*, 1981, **43**, 342-352 (*occur, ms*)
 Balandrin, M.F. *et al.*, *Heterocycles*, 1982, **19**, 1931-1934 (*5,6-Didehydro- α -isosparteine*)
 Kinghorn, A.D. *et al.*, *Phytochemistry*, 1982, **21**, 2269-2275 (*isol, ms, synth, bibl*)

2,3-Dehydro- β -stachydrine D-169

$C_7H_{11}NO_2$ 141.169
 Isol. from the red alga *Pterocladia capillacea*. Off-white amorph. powder.
 Sciuto, S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 322

Dehydrothalicarpine D-170

Thalictrucarpine
 [7224-94-4]



$C_{41}H_{46}N_2O_8$ 694.823
 Alkaloid from *Thalictrum minus* ssp. *elatum*, the roots of *Thalictrum dasycarpum* and the root bark of *Hernandia ovigera* (Ranunculaceae, Hernandiaceae). Cryst. (EtOAc, MeOH or MeOH/Et₂O). Mp 186-187°. $[\alpha]_D^{25} +55$ (c, 0.44 in CHCl₃). Possibly an artifact prod. from Thalicarpine, T-312 by aerial oxidn.

Methiodide (1:2):

Cryst. + 1H₂O (MeOH). Mp 167-171°.

*O*⁷-*De-Me: Dehydrothalmelatine*, *6'-(6a,7-Didehydro-1,2,10-trimethoxyaporphin-9-yl)oxy*codamine [16624-99-0]

$C_{40}H_{44}N_2O_8$ 680.796

Alkaloid from *Thalictrum minus* ssp. *elatum* and from the bark of *Hernandia peltata* (Ranunculaceae). Cryst. + 1 H₂O (MeOH). Mp 126-128°. $[\alpha]_D^{19} +31.9$ (c, 0.15 in CHCl₃). Poss. an artifact of aerial oxidn.

Dutschewska, H.B. *et al.*, *Chem. Ind. (London)*, 1966, 770 (*uv, pmr, struct*)

Dutschewska, H.B. *et al.*, *Chem. Ber.*, 1967, **100**, 3135 (*isol, struct, Dehydrothalicarpine, Dehydrothalmelatine*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1968, **33**, 1052 (*isol, uv, pmr, struct, synth*)


Cava, M.P. *et al.*, *Tetrahedron*, 1972, **28**, 4299; *J.O.C.*, 1975, **40**, 3601 (*synth*)

Yang, T.-H. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1976, **23**, 29 (*isol*)

Chalandre, M.C. *et al.*, *Can. J. Chem.*, 1986, **64**, 123 (*isol, deriv*)

Dehydrothalictricavine D-171

5,6-Dihydro-9,10-dimethoxy-13-methylbenzo[g]-1,3-benzodioxolo[5,6-a]quinolinium, *9CI* [38691-92-8]



$C_{21}H_{20}NO_4^{\oplus}$ 350.393
 Alkaloid from tubers of *Corydalis cava* and aerial parts of *Corydalis solida*.

Chloride: [54260-72-9]

$C_{21}H_{20}ClNO_4$ 385.846

Yellow needles (H₂O). Mp 185-195° dec.

Iodide: [2220-12-4]

$C_{21}H_{20}INO_4$ 477.298


Yellow cryst. (MeOH). Mp 215-218°.

Pavelka, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3654 (*synth, uv, ir*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261; 1985, **50**, 2299 (*isol, uv*)

Dehydrotryptamine D-172

2-(1H-Indol-3-yl)ethenamine, *3-(2-Aminovinyl)-1H-indole*, *Didehydrotryptamine*



$C_{10}H_{10}N_2$ 158.202

(E)-form

N^α-*Formyl*: *N-[2-(1H-Indol-3-yl)ethenyl]formamide*, *N-Formyldehydrotryptamine* [957062-44-1]

$C_{11}H_{10}N_2O$ 186.213

Prod. by *Pseudomonas aeruginosa*. Not fully characterised.

N^α,*N*^α,*N*^α-*Tri-Me*: *2-(1H-Indol-3-yl)-N,N,N-trimethylethenaminium*, *Conicamine* [648437-96-1]

$C_{13}H_{17}N_2^{\oplus}$ 201.291


Alkaloid from the tunicate *Aplidium conicum*. Histamine antagonist. Counterion not specified.

Aiello, A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4481-4483 (*Conicamine*)

Brady, S.F. *et al.*, *J.A.C.S.*, 2007, **129**, 12102-12103 (*N-Formyldehydrotryptamine*)

Dehydrovoachalotine D-173

Methyl 6,17-epoxy-1-methylsarpagan-16-carboxylate, *9CI* [18783-45-4]



Absolute Configuration

$C_{22}H_{24}N_2O_3$ 364.443

Alkaloid from *Voacanga chalotiana* (Apocynaceae). Cryst. (C₆H₆). Mp 239-239.5°. $[\alpha]_D^{22} +124$ (c, 0.9 in MeOH). pK_a 4.5 (80% MCS aq.). λ_{max} 225 (ε 44000); 283 (ε 7200); 290 (ε 6000) (no solvent reported).

17ξ-Hydroxy: 17-Hydroxydehydrovoachalotine

[123225-55-8]


$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the leaves and roots of *Alstonia undulata* (Apocynaceae). Amorph. $[\alpha]_D +93$ (c, 0.5 in CHCl₃).

Tirions, G. *et al.*, *Chimia*, 1968, **22**, 87-88 (*isol, uv, ir, pmr, ms, struct*)

Nouls, J.C. *et al.*, *Tet. Lett.*, 1968, 2731-2734 (*pmr, config*)
 Bombardelli, E. *et al.*, *Phytochemistry*, 1976, **15**, 2021-2022 (*cmr*)
 Morfaux, A.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1989, **309**, 33-36 (17-Hydroxydehydrovoachalotine)


2-Dehydroxy-*O*²-buxafuranamine **D-174**
 [142674-56-4]



C₃₅H₄₈N₂O₄ 560.775
 Alkaloid from leaves of *Buxus hildebrandtii* (Buxaceae). Amorph. solid. [α]_D²⁰ +130 (c, 1.7 in CHCl₃).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1992, **34**, 157-171 (*isol, uv, ir, pmr, ms, struct*)

Dehydroxylopine **D-175**
 6,7-Dihydro-10-methoxy-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, 9CI, 9-Methoxydehydroanonaine
 [113425-60-8]




C₁₈H₁₅NO₃ 293.321
 Alkaloid from *Xylopiella vieillardii* (Annonaceae). Amorph.

N-Me: Dehydroisolaureline

[65967-04-6]
 C₁₉H₁₇NO₃ 307.348
 Alkaloid from *Liriodendron tulipifera* (Magnoliaceae). Mp 143-145°.

Ziyayev, R. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 715; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 602 (*Dehydroisolaureline*)
 Jossang, A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 466 (*Dehydroxylopine*)

Delagoenine **D-176**
 [221001-70-3]



R¹ = R² = OMe

C₁₉H₂₅NO₅ 347.41
 Alkaloid from the bulbs of *Crinum*

delagoense. Mp 120-122°. [α]_D²⁰ +34.2 (c, 0.48 in MeOH).

Nair, J.J. *et al.*, *Phytochemistry*, 1998, **49**, 2539-2543 (*isol, ir, cd, pmr, cmr, ms*)

Delagoensine **D-177**


As Delagoenine, D-176 with R¹R² = -OCH₂O-

C₁₈H₂₁NO₅ 331.368
 Alkaloid from the bulbs of *Crinum delagoense*. Mp 132-134°. [α]_D²⁰ +28.2 (c, 0.47 in MeOH).

Nair, J.J. *et al.*, *Phytochemistry*, 1998, **49**, 2539-2543 (*isol, ir, cd, pmr, cmr, ms*)

Delamide **D-178**


Methyl 2-[(4-amino-2-methyl-1,4-dioxobutyl)amino]benzoate, 9CI
 [151805-38-8]



C₁₃H₁₆N₂O₄ 264.28
 Alkaloid from *Delphinium omeiense*. Component of Fu Zi.

Zhang, C.Y. *et al.*, *Fitoterapia*, 1993, **64**, 188-189

Delaminomycin A **D-179**
 [149779-38-4]



C₂₉H₄₃NO₆ 501.662
 Prod. by *Streptomyces albus*. Immunomodulator. Active against gram-positive bacteria. Cell adhesion inhibitor, extracellular matrix receptor antagonist. Powder + 1H₂O. Sol. MeOH, DMSO; poorly sol. hexane, H₂O, EtOAc. Related to Lydicamycin, L-353. λ_{max} 232 (ε 29000); 288 (ε 7110) (MeOH) (Derep). λ_{max} 232 (E1%/1cm 728); 288 (E1%/1cm 208) (MeOH) (Berdy).

5'-Me ether: Delaminomycin B

[149779-39-5]
 C₃₀H₄₅NO₆ 515.689
 Prod. by *Streptomyces albus*. Immunomodulator. Active against gram-positive bacteria. Cell adhesion inhibitor, extracellular matrix receptor antagonist. Powder. Sol. MeOH, DMSO; poorly sol. EtOAc, hexane, H₂O. λ_{max} 232 (ε 29000); 288 (ε 7110) (MeOH) (Derep).


5'-Deoxy: Delaminomycin C

[149779-40-8]
 C₂₉H₄₃NO₅ 485.662
 Prod. by *Streptomyces albus*. Immunomodulator. Active against gram-

positive bacteria. Cell adhesion inhibitor, extracellular matrix receptor antagonist. Powder. Sol. MeOH, DMSO; poorly sol. EtOAc, H₂O, hexane. λ_{max} 232 (ε 29000); 288 (ε 7110) (MeOH) (Derep).

Ueno, M. *et al.*, *J. Antibiot.*, 1993, **46**, 719; 979; 1020; 1156; 1390 (*isol, pmr, cmr, struct, activity*)


Delatisine **D-180**
 [141591-00-6]



C₂₀H₂₅NO₃ 327.422
 Alkaloid from the seeds of *Delphinium elatum* (Ranunculaceae). Cryst. (Me₂CO). Mp 274.5-276.5°. [α]_D²⁵ +8.6 (c, 0.21 in CHCl₃).

Ross, S.A. *et al.*, *Tetrahedron*, 1991, **47**, 9585-9598 (*isol, pmr, cmr, cryst struct*)

Delavaine **D-181**
Delavayine
 [27989-72-6]



Absolute configuration

C₂₀H₂₃NO₅ 357.405
 See also Delavaine A, D-182. Alkaloid from *Stephania delavayi* (Menispermaceae). Mp 149-150°. [α]_D -240 (CHCl₃).

N-De-Me: N-Nordelavaine

C₁₉H₂₁NO₅ 343.379
 Alkaloid from the tuberous roots of *Stephania suberosa* (Menispermaceae). Amorph. [α]_D²⁵ -187 (c, 0.61 in CHCl₃).

16-Oxo: 16-Oxodelavaine

[38146-58-6]
 C₂₀H₂₁NO₆ 371.389
 Alkaloid from *Stephania delavayi*. Mp 221-222°. [α]_D -180 (CHCl₃). Rel. config. only detd.

Fadeeva, I.I. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 784; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 756 (*isol, struct*)


Il'inskaya, T.N. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 129; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 134 (*isol, uv, ir, pmr, struct, deriv*)

Inubushi, Y. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 393 (*ir, pmr, struct*)

Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 2391 (*N-Nordelavaine*)

Delavaine A†

[109291-57-8]

R = CH(CH₃)CH₂COOMeC₃₈H₅₄N₂O₁₁ 714.851

See also Delavaine B, D-183. Not to be confused with Delavaine, D-181. Alkaloid from the roots of *Delphinium delavayi* var. *pogonanthum*. Also isol. from *Delphinium tricorne* as Alkaloid B (a mixt. with Delavaine B, D-183). Amorph. [α]_D²⁹ +39.4 (c, 0.8 in CHCl₃).

Et ester analogue: **Gyalanine A**

[133744-29-3]

C₃₉H₅₆N₂O₁₁ 728.878Alkaloid from the roots of *Delphinium gyalanum* (Ranunculaceae).Butyl ester analogue: **Potandine A**

[161068-58-2]

C₄₁H₆₀N₂O₁₁ 756.932Alkaloid from the roots of *Delphinium potaninii*. Amorph. powder. [α]_D¹⁰ +15.5 (c, 1 in MeOH).16-Demethoxy: **16-Demethoxydelavaine A**C₃₇H₅₂N₂O₁₀ 684.825Alkaloid from *Delphinium cuneatum*. Isol. as a mixt. with 16-Demethoxydelavaine B, in D-183.Parent acid: **Shawurensine**

[953392-29-5]

C₃₇H₅₂N₂O₁₁ 700.825Alkaloid from the aerial parts of *Delphinium shawurensense*. Amorph. solid.Pelletier, S.W. et al., *Heterocycles*, 1986, **24**, 1853-1865 (*Delavaine A*)Wang, F. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1990, **32**, 733-736; *CA*, **114**, 203508 (*Gyalanine A*)Pu, H. et al., *Yaoxue Xuebao*, 1994, **29**, 689-692; *CA*, **122**, 122898 (*Potandine A*)Khairitdinova, E.D. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 572-574 (16-Demethoxydelavaine A)Gu, D.Y. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 298-301 (*Shawurensine*)**Delavaine B****D-183**

[109314-15-0]



As Delavaine A, D-182 with

R = CH₂CH(CH₃)COOMeC₃₈H₅₄N₂O₁₁ 714.851

See also Delavaine A, D-182. Not to be confused with Delavaine, D-181. Alkaloid from the roots of *Delphinium delavayi* var. *pogonanthum*. Also from *Delphinium tricorne* as a mixt. with Delavaine A, D-182. Amorph. [α]_D²⁹ +31.7 (c, 0.8 in CHCl₃).

Et ester analogue: **Gyalanine B**


[133744-30-6]

D-182C₃₉H₅₆N₂O₁₁ 728.878Alkaloid from the roots of *Delphinium gyalanum* (Ranunculaceae).16-Demethoxy: **16-Demethoxydelavaine B**C₃₇H₅₂N₂O₁₀ 684.825Alkaloid from *Delphinium cuneatum*. Isol. as a mixt. with 16-Demethoxydelavaine A, in D-182.Pelletier, S.W. et al., *Heterocycles*, 1986, **24**, 1853-1865 (*Delavaine B*)Wang, F. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1990, **32**, 733-736; *CA*, **114**, 203508 (*Gyalanine B*)Khairitdinova, E.D. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 572-574 (16-Demethoxydelavaine B)**Delavayine A†****D-184**C₁₉H₂₈NO₂⁺ 302.436Alkaloid from *Incarvillea delavayi*. Antinociceptive agent. Yellow powder. [α]_D²² -5.1 (c, 0.9 in Py). Counterion not specified.Nakamura, M. et al., *Phytochemistry*, 2000, **53**, 253-256**Delavayine B†****D-185**R = -CH(COO⁻)CH₂PhC₂₁H₂₅NO₄ 355.433Alkaloid from *Incarvillea delavayi*. Off-white powder. [α]_D²⁴ -6.5 (c, 1.8 in Py). Zwitterionic.Nakamura, M. et al., *Chem. Pharm. Bull.*, 2000, **48**, 1826-1827**Delavayine C†****D-186**

As Delavayine B, D-185 with

R = -(CH₂)₃COO⁻C₁₆H₂₃NO₄ 293.362Alkaloid from *Incarvillea delavayi*. Off-white powder. [α]_D²⁴ -72.3 (c, 0.5 in MeOH). Zwitterionic.Nakamura, M. et al., *Chem. Pharm. Bull.*, 2000, **48**, 1826-1827**Delboxine****D-187**

[95066-32-3]

C₂₄H₃₇NO₇ 451.559Alkaloid from the roots of *Delphinium bonvalotii*. Mp 200-202°. [α]_D¹⁸ +43.5 (c, 0.1 in CHCl₃).O⁸-De-Me: **Tuguacnitine**

[93414-10-9]

C₂₃H₃₅NO₇ 437.532Alkaloid from the roots of *Aconitum sibiricum*. Cryst. (EtOAc/Et₂O). Mp 197-199°.O¹⁴-De-Me: **14-O-Demethyldeboxine**C₂₃H₃₅NO₇ 437.532Alkaloid from *Consolida orientalis*. Amorph. solid. [α]_D²⁵ +43.3 (c, 0.5 in CHCl₃).O⁸,O¹⁴-Di-de-Me: **14-O-Demethyltuguacnitine**C₂₂H₃₃NO₇ 423.505Alkaloid from the roots of *Delphinium stapeliosum*. Cryst. (Me₂CO/Et₂O). Mp 208-210°. [α]_D²⁰ +59.5 (c, 0.34 in CHCl₃).O⁸,O¹⁴-Di-de-Me, 14-ketone: **Hohenackeridene**

[194224-78-7]

C₂₂H₃₁NO₇ 421.489Alkaloid from aerial parts of *Aconitella hohenackeri* (*Consolida hohenackeri*). Cryst. (hexane/EtOAc). Mp 224-225°. [α]_D +39.5 (c, 0.38 in MeOH).O⁶,O⁸,O¹⁴-Tri-de-Me, O¹-Me: **Tiantaishansine**

[959843-30-2]

C₂₂H₃₃NO₇ 423.505Alkaloid from the roots of *Delphinium tiantaishanense*. Amorph. powder. Mp 94-96°. [α]_D²⁰ +35.4 (c, 0.84 in CHCl₃).6-Demethoxy, 7-deoxy, O⁸-de-Me: **Monticamine**

[81047-05-4]

C₂₂H₃₃NO₅ 391.506Alkaloid from *Aconitum monticola*. Cryst. (Et₂O/Me₂CO). Mp 163-164°. [α]_D +4.6-Demethoxy, O⁸-de-Me: **Monticoline**

[81037-22-1]

C₂₂H₃₃NO₆ 407.506Alkaloid from *Aconitum monticola* and from the aerial parts of budding *Aconitum karakolicum*. Cryst. (Me₂CO). Mp 166-167°. [α]_D +15 (CHCl₃).6-Demethoxy, 9 β -hydroxy, 7-deoxy, O⁸-de-Me: **Excelsine†**

[41645-62-9]

C₂₂H₃₃NO₆ 407.506Alkaloid from the roots of *Aconitum excelsum*. Mp 103-105°.6-Demethoxy, 9 β -hydroxy, 7-deoxy, O⁸-de-Me, 1-Ac: **1-Acetylexcelsine. Akiradine**

[142748-49-0]

C₂₄H₃₅NO₇ 449.543Alkaloid from *Aconitum kirinense*. Cryst. (Et₂O). Mp 108-110°.6-Demethoxy, 9 β -hydroxy, 7-deoxy, O⁸-de-Me, 8-Ac: **8-Acetylexcelsine**

[142735-46-4]

C₂₄H₃₅NO₇ 449.543Alkaloid from epigeal parts of *Aconitum kirinense*. Amorph.

6-Demethoxy, 9 β -hydroxy, O⁸-de-Me: Sinomontanine G

[872977-65-6]

C₂₂H₃₃NO₇ 423.505Alkaloid from the roots of *Aconitum sinomontanum*. [α]_D¹⁷ -32 (c, 0.5 in CHCl₃).**6-Epimer, 7-deoxy, O⁸-de-Me: Kiritine**

[161068-75-3]

C₂₃H₃₅NO₆ 421.533Alkaloid from *Aconitum kirinense*.Nasirov, S.M. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 206; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 184 (*Excelsine, cryst struct*)Ametova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1981, **446**; *Chem. Nat. Compd. (Engl. Transl.)*, **345** (*Monticamine, Monticoline*)Jiang, Q.P. *et al.*, *Heterocycles*, 1985, **23**, 11 (*Delboxine*)Chung, B.S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1074-1077 (*Tuguaconitine*)Nishanov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1991, **258**; *Chem. Nat. Compd. (Engl. Transl.)*, **222** (*8-Acetylexcelsine*)Feng, F. *et al.*, *CA*, 1995, **122**, 128605f (*Kiritine*)Almanza, G. *et al.*, *Phytochemistry*, 1997, **45**, 1079 (*Hohenackeridine*)Tshebaeva, U.T. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1999, **35**, 692 (*Akiradine*)Shrestha, P.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 2-5 (*14-Demethyltuguaconitine*)Alva, A. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 2110-2119 (*14-Demethyldelboxine*)Peng, C.-S. *et al.*, *Youji Huaxue*, 2005, **25**, 1235-1239; *CA*, **144**, 103949n (*Sinomontanine G*)Li, J. *et al.*, *Molecules*, 2007, **12**, 353-360 (*Tiantaishansine*)Kang, C.K. *et al.*, *Arch. Pharmacol. Res.*, 2008, **31**, 259-263 (*Tuguaconitine, cryst struct*)**Delflexine****D-188**


[11045-95-7]

C₂₄H₃₉NO₆ 437.575Struct. unknown. Alkaloid from *Delphinium flexuosum* (Ranunculaceae). Mp 191-192°.Brutko, L.I. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 21-23; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 16**Delfrenine****D-189**

[11045-96-8]

C₂₇H₃₁NO₆ 465.545Struct. unknown. Alkaloid from *Delphinium freynii* (Ranunculaceae). Mp 246-247°. Unsaturated benzoate ester containing no methoxy or methylenedioxy groups. Saponification gives benzoic acid an amino acid characterised as its Me ester (C₂₁H₃₁NO₆).Brutko, L.I. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 21-23; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 16**Delnudine, 9CI****D-190**


[26503-54-8]



Absolute configuration

C₂₀H₂₅NO₃ 327.422Alkaloid from the seeds of *Delphinium denudatum* (Ranunculaceae). Mp 235-237°. pK_a 7.8 (MCS).Birnbaum, K.B. *et al.*, *Tet. Lett.*, 1969, 5245 (*cryst struct*)Götz, M. *et al.*, *Tet. Lett.*, 1969, 5335 (*isol, uv, ir, pmr*)Birnbaum, K.B. *et al.*, *Acta Cryst. B*, 1971, **27**, 1169 (*cryst struct*)**Delorine****D-191**C₂₂H₂₉NO₅ 387.475Struct. unknown. Alkaloid from the seeds of *Delphinium orientale* (Ranunculaceae). Mp 227-229°. [α]_D +54.7 (c, 2.34 in CHCl₃).*Hydrochloride*: Mp 227-229°.Platonova, T.F. *et al.*, *Med. Promst. SSSR*, 1963, **17**, 19-20; *CA*, **59**, 6723g (*isol, uv*)**Delphatisine B****D-192**

[1003600-47-2]

C₂₄H₃₃NO₄ 399.529Alkaloid from *Delphinium chrysotrichum*. Amorph. powder. [α]_D²⁷ -2 (c, 1 in CHCl₃).He, Y.Q. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 545-547 (*isol, pmr, cmr*)**Delphoccine****D-193**

[1356-58-7]

Struct. unknown. Alkaloid contaminating samples of Deltaline (see Dictyocarpine, in M-307) from *Delphinium occidentale* (Ranunculaceae). No other information recorded.Carmack, M. *et al.*, *J.A.C.S.*, 1958, **80**, 497**Delphoorine****D-194**


[11054-87-8]

C₂₀H₂₈NO₆ 378.444Struct. unknown. Alkaloid from the seeds of *Delphinium orientale* (Ranunculaceae). Mp 276-278°.Boyadzhieva, M. *et al.*, *Farmatsiya (Sofia)*, 1968, **18**, 5-6; *CA*, **71**, 19513g**Delpyrine****D-195**

[11045-98-0]

C₄₉H₈₂N₂O₁₇ 971.19Struct. unknown. Alkaloid from *Delphinium pyramidatum* (Ranunculaceae). Amorph. Mp 76°. [α]_D +58. Contains 2 OH groups and saponifies to an uncharacterised acid and an amorph. base (C₄₆H₇₄N₂O₅).Brutko, L.I. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 21-23; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 16**Delsemine****D-196**

[6887-42-9]

Delsemine A; R = CH(CH₃)CH₂CONH₂ (S-) Delsemine B; R = CH₂CH(CH₃)CONH₂ (S-)Artifacts. Formerly isol. as a pair of regioisomers, later characterised separately. Alkaloid from *Delphinium semibarbatum*, *Delphinium oreophilum*, *Delphinium tricornis*, *Delphinium formosum*, *Consolida ambigua* (*Delphinium ajacis*) and *Delphinium delavayi* var. *pogonanthum* (Ranunculaceae). Cryst. (70% EtOH).
▶AR5530000**Delsemine A** [66878-11-3]C₃₇H₅₃N₃O₁₀ 699.84Amorph. solid. [α]_D³⁰ +368 (c, 0.7 in CHCl₃).O¹⁴-De-Me, 14-(2-methylpropanoyl):**Giraldine G**

[740816-09-5]

C₄₀H₅₇N₃O₁₁ 755.904Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder. Mp 108-110°. [α]_D²⁰ +35.4 (c, 0.42 in CHCl₃).O¹⁴-De-Me, 14-(2S-methylbutanoyl):**Giraldine H**

[740816-10-8]

C₄₁H₅₉N₃O₁₁ 769.931Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder. Mp 122-124°. [α]_D²⁰ +34.6 (c, 0.35 in CHCl₃).**Delsemine B** [66841-98-3]C₃₇H₅₃N₃O₁₀ 699.84Amorph. solid. [α]_D³⁰ +28.2 (c, 0.6 in CHCl₃).O¹⁴-De-Me: **Trifoliolasine B**

[740815-34-3]

C₃₆H₅₁N₃O₁₀ 685.813Alkaloid from *Delphinium trifoliolatum*. Amorph. powder. Mp 103-105°. [α]_D²⁰ +36.6 (c, 0.48 in CHCl₃).O¹⁴-De-Me, 14-(2-methylpropanoyl):**Trifoliolasine C**

[740815-35-4]

C₄₀H₅₇N₃O₁₁ 755.904Alkaloid from *Delphinium trifoliolatum*. Amorph. powder. Mp 117-118°. [α]_D²⁰ +24 (c, 0.3 in CHCl₃).Yunusov, S. *et al.*, *Zh. Obshch. Khim.*, 1951, **21**, 967-973; *J. Gen. Chem. USSR (Engl. Transl.)*, 1951, **21**, 1059-1065; *CA*, **46**, 516 (*isol*)Pelletier, S.W. *et al.*, *Tet. Lett.*, 1977, 2735-2736 (*occur, struct*)Pelletier, S.W. *et al.*, *J.A.C.S.*, 1981, **103**, 6536-6538 (*config*)Pelletier, S.W. *et al.*, *Heterocycles*, 1986, **24**, 1853-1865 (*Delsemines A,B, isol, ir, pmr, cmr, ms, synth*)

Zhou, X.-L. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 381-383; 456-458 (*Trifoliosines*, *Geraldines*)

Demanyl phosphate D-197

Phosphoric acid mono[2-(dimethylamino)ethyl] ester, 9CI. Mono-2-dimethylaminoethanol dihydrogen phosphate. Phosphoryldimethylaminoethanol. Phosphoryldimethylcholamine. Phosphoryldimethylethanolamine. PDMEA.

Panclar. Pantos
[6909-62-2]
Me₂NCH₂CH₂OP(O)(OH)₂

C₄H₁₂NO₄P 169.117
Isol. from a choline-requiring mutant of *Neurospora crassa*. Psychotonic agent. Cryst. + 1H₂O (EtOH). Mp 78-81° Mp 175-176° (anhyd.).

Cherbuliez, E. *et al.*, *Helv. Chim. Acta*, 1956, **41**, 1168-1175 (*synth*)

Wolf, B. *et al.*, *J. Biol. Chem.*, 1959, **234**, 1068-1071 (*synth, isol*)


Ansell, G.B. *et al.*, *Biochem. J.*, 1966, **98**, 303-310 (*biochem*)

Taguchi, Y. *et al.*, *J.O.C.*, 1975, **40**, 2310-2313 (*synth*)

Miyazaki, H. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 763-769 (*metab*)


Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 12624

Andriamampandry, C. *et al.*, *Biochem. J.*, 1989, **264**, 555-561 (*biochem*)

Demecolcinone D-198

C₁₉H₁₉NO₅ 341.363
Alkaloid from *Colchicum brachyphyllum*. Yellowish powder. [α]_D²³ +242 (c, 0.3 in MeOH). λ_{max} 232 (ε 18735) (MeOH).

Alali, F.Q. *et al.*, *J. Nat. Prod.*, 2005, **68**, 173-178 (*isol, pmr, cmr*)

3-Demethoxyerythratidinone D-199

(+)-form

C₁₈H₂₁NO₃ 299.369
Unique *Erythrina* alkaloid lacking a 3-methoxy group.

(+)-form [41431-29-2]

Alkaloid from the leaves of *Erythrina lithosperma* (Fabaceae). Cryst. (C₆H₆/petrol). Mp 111-112°. [α]_D +325 (c, 0.249 in CHCl₃).

Picrate: Mp 200-202°.

(-)-form

[α]_D²⁰ -236 (c, 0.8 in CHCl₃) (72% ee).

(±)-form

Synthetic. Mp 105-108° (101-102°).

Picrate: Mp 250-252°.

Barton, D.H.R. *et al.*, *J.C.S. Perkin 1*, 1973, 874 (*isol, uv, ir, pmr, ms, struct*)

Tanaka, H. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 1578 (*synth, ir, pmr, ms*)

Danishefsky, S.J. *et al.*, *J.A.C.S.*, 1987, **109**, 917 (*synth*)

Irie, H. *et al.*, *Heterocycles*, 1989, **29**, 1033 (*synth*)

Wasserman, H.H. *et al.*, *J.O.C.*, 1989, **54**, 5843 (*synth*)

Ishibashi, H. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 907 (*synth, ir, pmr*)

Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1462; 1994, **42**, 204 (*synth, ir, pmr*)

Hosoi, S. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 3115 (*synth*)

Cassayre, J. *et al.*, *Tet. Lett.*, 1998, **39**, 8995-8998 (*synth*)


Allin, S.M. *et al.*, *Tet. Lett.*, 2004, **45**, 5493-5496 (*synth*)

Wang, Q. *et al.*, *Org. Lett.*, 2006, **8**, 601-604 (*synth*)

Gao, S. *et al.*, *Org. Lett.*, 2006, **8**, 2373-2376 (*synth*)

3-Demethoxy-2,3-methylene-dioxyerythroline D-200

Methyl 1,6-didehydro-16-methoxy-2,3-[methylenebis(oxy)]erythrinan-15-carboxylate, 9CI
[89560-98-5]



C₂₀H₂₃NO₅ 357.405
Alkaloid from the leaves of *Hyperbaena columbica*. Cryst. (MeOH). Mp 94-97°. [α]_D²⁵ +229 (c, 0.56 in CHCl₃). λ_{max} 215 (log ε 4.55); 238 (sh) (log ε 4.07); 305 (log ε 3.54) (EtOH).


Parent acid, amide: 15-Amido-3-demethoxy-2,3-methylenedioxyerythroline

C₁₉H₂₂N₂O₄ 342.394

Alkaloid from the leaves of *Hyperbaena valida*. Cryst. (CH₂Cl₂/MeOH). Mp 219-220°. [α]_D +228 (c, 0.1 in CHCl₃). λ_{max} 224 (log ε 4.95); 301 (log ε 4.61); 330 (log ε 4.1) (EtOH).

Ripperger, H. *et al.*, *Phytochemistry*, 1983, **22**, 2603-2605 (*isol, ord, pmr, ms*)

Freyer, A.J. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1514-1516 (*amide*)


7-Demethyldeoxynupharidine D-201

C₁₄H₂₁NO 219.326
Alkaloid from the scent glands of *Castor fiber*. Oil. [α]_D²⁰ -85 (c, 0.5 in CHCl₃).

Maurer, B. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 1169

Demethylneodrupacine D-202

[80791-43-1]




C₁₇H₁₉NO₅ 317.341
Alkaloid from *Cephalotaxus hainanensis* (Cephalotaxaceae). Mp 197-200°.

Xue, Z. *et al.*, *Yaoxue Xuebao*, 1981, **16**, 752; *CA*, **96**, 82690u

Huang, L. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 157 (*rev*)

7-Demethylpicridin A₁ D-203

SN 198D. Antibiotic SN 198D
[155148-30-4]



C₂₄H₃₅NO₄ 401.545
Prod. by *Streptomyces* sp. SN-198. Yellowish oil. Related to Picridin, P-421.


4'-O-α-L-Rhamnopyranoside: 7-*Demethyl-4'-rhamnopicridin A₁*. SN 198B. Antibiotic SN 198B
[155148-29-1]

C₃₀H₄₅NO₈ 547.687
Prod. by *Streptomyces* sp. SN-198. Amorph. powder.

Kimura, K. *et al.*, *J. Antibiot.*, 1996, **49**, 697 (*isol, pmr*)

Demethylsecologanol D-204

[872360-10-6]



C₁₆H₂₄O₁₀ 376.36
Constit. of *Ophiorrhiza liukuensis*. Amorph. solid. [α]_D²³ -108.3 (c, 0.06 in MeOH). λ_{max} 233 (MeOH).

Me ester: *Secologanol*

[72463-81-1]

C₁₇H₂₆O₁₀ 390.386

Constit. of *Gentiana verna*.

7-O-(3-Pyridinecarbonyl), *Me ester*:

Tripterospermumcin A

[929642-20-6]

C₂₃H₂₉NO₁₁ 495.482

Constit. of *Tripterospermum chinense*.
Amorph. powder. $[\alpha]_D^{20}$ -147.6 (c, 0.25
in MeOH). λ_{\max} 227 (log ϵ 4.03)
(MeOH).

7-Aldehyde, Me ester: **Secologanin**.

Loniceroside[†]

[19351-63-4]

C₁₇H₂₄O₁₀ 388.371

Constit. of *Strychnos nux-vomica*, *La-
thraea clandestina* and *Lonicera mor-
rowii*. Precursor of indole alkaloids.
Oil. Sol. MeOH; poorly sol. hexane.
 $[\alpha]_D$ -96 (MeOH). λ_{\max} 235 (ϵ 9100)
(MeOH) (Berdy).

Battersby, A.R. *et al.*, *J.C.S. (C)*, 1969, 1187
(*Secologanin, synth*)

Souzu, I. *et al.*, *Tet. Lett.*, 1970, 191
(*Secologanin, isol*)

Guarnaccia, R. *et al.*, *J.A.C.S.*, 1974, **96**, 7079
(*Secologanin, biosynth*)

Bisset, N.G. *et al.*, *Phytochemistry*, 1974, **13**,
265 (*Secologanin, isol*)

Takeda, Y. *et al.*, *Chem. Pharm. Bull.*, 1976,
24, 79 (*biosynth*)

Nakane, M. *et al.*, *J.O.C.*, 1980, **45**, 4233
(*Secologanin, synth*)

Brown, R.T. *et al.*, *Chem. Comm.*, 1986, 1818
(*Secologanin, synth*)

Inoue, K. *et al.*, *Phytochemistry*, 1989, **28**, 2971
(*biosynth*)


Yamamoto, H. *et al.*, *Phytochemistry*, 2000, **53**,
7-12 (*biosynth*)

Zhu, K.-C. *et al.*, *Helv. Chim. Acta*, 2007, **90**,
291-296 (*Tripterospermumcin A*)

Demissidine

D-205

Solanidan-3-ol, 9CI. *Solanine D*. *Dihy-
drosolanidine T*
[474-08-8]



C₂₇H₄₅NO 399.659

Product of hydrol. of Demissine, obt. by
hydrol. of many *Solanum* extracts
(Solanaceae). Shows antifungal activity.
Mp 219-220°. $[\alpha]_D^{18}$ +28 (MeOH).

► WF0193000

O- $[\beta$ -D-Glucopyranosyl-(1→2)- $[\beta$ -D-xy-
lopyranosyl-(1→3)]- $[\beta$ -D-glucopyrano-
syl-(1→4)- β -D-galactopyranoside]:

Demissine

[6077-69-6]

C₅₀H₈₃NO₂₀ 1018.2

Alkaloid from *Solanum demissum*,
Solanum depexum, *Solanum horowitzii*,
Solanum jamesii, *Solanum polyade-
nium*, *Solanum punae*, *Solanum schrei-
teri*, *Solanum acaule*, *Solanum
chacoense*, *Solanum commersonii*, *Sol-
anum cartilobum*, *Solanum juzepczukii*
and *Lycopersicon pimpinellifolium*.
Cholinesterase inhibitor. Repellent to
the Colorado beetle *Leptinotarsa de-
cemlineata*. Mp 305-308° dec. $[\alpha]_D$ -20
(Py).

3-O- $[\beta$ -D-Glucopyranosyl-(1→2)- $[\beta$ -D-
glucopyranosyl-(1→3)]- β -D-glucopyr-
anosyl-(1→4)- β -D-galactopyranoside]:

Commersonine

[60776-42-3]

C₅₁H₈₅NO₂₁ 1048.227

Alkaloid from leaves of *Solanum cha-
coense* and *Solanum commersonii* (Sol-
anaceae). Cryst. solid. Mp 230-232°.
 $[\alpha]_D^{25}$ -17 (Py).

Glycoside: β -Demissine

Isol. from a mutant of *Lycopersicon
pimpinellifolium* (Solanaceae).

O-Ac: Mp 194°.

5,6-Didehydro, 3-O- $[\beta$ -D-glucopyranosyl-
(1→2)- $[\beta$ -D-glucopyranosyl-(1→3)]-
 β -D-glucopyranosyl-(1→4)- β -D-galac-
topyranoside]: **Dehydrocommersonine**
[65428-74-2]

C₅₁H₈₃NO₂₁ 1046.211

Alkaloid from root cultures of *Sola-
num chacoense* (Solanaceae).

3-Epimer:

C₂₇H₄₅NO 399.659

Alkaloid from *Solanum tuberosum*
(potato). Needles (Me₂CO). Mp 215-
217° (210-211°). $[\alpha]_D$ +31 (CHCl₃).

**5-Epimer: 3 β -Allosolanidan-3-ol. 5-Epi-
demissidine**

[78513-80-1]

C₂₇H₄₅NO 399.659

Alkaloid from *Solanum tuberosum*
(potato) (Solanaceae). Mp 219-220°.
 $[\alpha]_D$ +26 (CHCl₃).

Prelog, V. *et al.*, *Helv. Chim. Acta*, 1944, **27**,
390 (*struct*)

Kuhn, R. *et al.*, *Chem. Ber.*, 1947, **80**, 406;
1957, **90**, 203 (*Demissine*)

Aslanov, K.A. *et al.*, *CA*, 1963, **59**, 6725f (3-
epimer, isol)

Schreiber, K. *et al.*, *Annalen*, 1964, **672**, 232-
235 (3-*epimer, synth*)

Adam, C. *et al.*, *Tetrahedron*, 1964, **20**, 1719
(*synth*)

Wolters, B. *et al.*, *Planta Med.*, 1966, **14**, 1392-
1401 (*activity*)

Höhne, E. *et al.*, *Tetrahedron*, 1966, **22**, 673
(*cryst struct*)

Osman, S.F. *et al.*, *Phytochemistry*, 1976, **15**,
1065 (*Commersonine*)


Radeglia, R. *et al.*, *Tet. Lett.*, 1977, 903 (*cmr*)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996,
32, 244; *Chem. Nat. Compd. (Engl. Transl.)*,
1996, **32**, 234 (*Allosolanidanol*)

Dendridine A

D-206

3,3'-Bis(2-aminoethyl)-5,5'-dibromo-4,4'-
bi-1H-indole-7,7'-diol



C₂₀H₂₀Br₂N₄O₂ 508.212


Isol. from the marine sponge *Dictyoden-
drilla* sp. Antibacterial and antifungal
agent. Yellowish solid. λ_{\max} 282 (ϵ 5700);
292 (ϵ 5800); 303 (ϵ 4000) (no solvent
reported).

Tsuda, M. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1277-
1278 (*isol*)

Dendrine

D-207

Methyl 12-oxodendrobane-2-acetate, 9CI
[2651-57-2]



Absolute
Configuration

C₁₉H₂₉NO₄ 335.442

Minor alkaloid from *Dendrobium nobile*
(Orchidaceae). Mp 191-192°. $[\alpha]_D^{11}$ -114
(c, 0.85 in CHCl₃). $[\alpha]_D^{24}$ -81 (c, 0.14 in
CHCl₃).


Inubushi, Y. *et al.*, *Tet. Lett.*, 1965, 2723 (*ir*,
pmr, ms, struct)

Graneli, I. *et al.*, *Acta Chem. Scand.*, 1970, **24**,
1108 (*synth, abs config*)

Dendroamide A

D-208

[176666-84-5]



R¹ = CH(CH₃)₂, R² = CH₃

C₂₁H₂₄N₆O₄S₂ 488.59

Isol. from the terrestrial blue-green alga
Stigonema dendroideum. Exhibits multi-
drug-resistance reversing activity. $[\alpha]_D$
+40.5 (c, 3.50 in CH₂Cl₂). Related to
Bistratamide A, B-223. λ_{\max} 224 (ϵ
28000) (MeOH). λ_{\max} 244 (ϵ 28000)
(MeOH) (Berdy).

Ogino, J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 581
(*isol, uv, ir, pmr, cmr, ms*)

Xia, Z. *et al.*, *J.O.C.*, 2001, **66**, 3459-3466
(*synth*)

Bertram, A. *et al.*, *Heterocycles*, 2002, **58**, 521-
561 (*synth*)

You, S.-L. *et al.*, *J.O.C.*, 2003, **68**, 9506-9509
(*synth*)

Yonezawa, Y. *et al.*, *Heterocycles*, 2005, **65**, 95-
105 (*synth*)

Matsumoto, T. *et al.*, *Tetrahedron*, 2007, **63**,
8571-8575 (*synth*)

Dendroamide B

D-209

[176666-85-6]

As Dendroamide A, D-208 with


$R^1 = \text{CH}_3$, $R^2 = -\text{CH}_2\text{CH}_2\text{SMe}$
 $\text{C}_{21}\text{H}_{24}\text{N}_6\text{O}_4\text{S}_3$ 520.656
 Isol. from the terrestrial blue-green alga
Stigonema dendroideum. $[\alpha]_{\text{D}} +44.5$ (c,
 1.53 in CH_2Cl_2). λ_{max} 224 (ϵ 31400)
 (MeOH).

S⁹-Oxide: Dendroamide C
 [176666-86-7]
 $\text{C}_{21}\text{H}_{24}\text{N}_6\text{O}_5\text{S}_3$ 536.656
 From *Stigonema dendroideum*. $[\alpha]_{\text{D}}$
 +26 (c, 3.75 in CH_2Cl_2). Mixt. of
 sulfoxide diastereoisomers. λ_{max} 224 (ϵ
 30900) (MeOH) (Berdy).

Orgino, J. et al., *J. Nat. Prod.*, 1996, **59**, 581
 (isol, uv, ir, pmr, cmr, ms, struct)

Dendrobine**D-210**

Dendroban-12-one, 9CI
 [2115-91-5]
 [30646-45-8 (\pm)-form]



Absolute
 Configuration

$\text{C}_{16}\text{H}_{25}\text{NO}_2$ 263.379

Numbering systems vary. Main alkaloid
 from *Dendrobium* spp., esp. *Dendrobium*
nobile (Orchidaceae). Shows antipyretic,
 hypotensive and convulsant activity.
 Cryst. (Et_2O). Mp 134-136°. $[\alpha]_{\text{D}}^{14}$ -48.4
 (c, 1.89 in MeOH).

N-Oxide: Dendrobine N-oxide

[40072-57-9]
 $\text{C}_{16}\text{H}_{25}\text{NO}_3$ 279.378
 Alkaloid from *Dendrobium nobile*
 (Orchidaceae). Needles + H_2O
 ($\text{Me}_2\text{CO}/\text{Et}_2\text{O}$). Mp 160-180° dec. $[\alpha]_{\text{D}}^{23}$
 -34 (c, 0.27 in MeOH).

**N-De-Me: N-Demethyldendrobine. Mu-
 bironine B**

$\text{C}_{15}\text{H}_{23}\text{NO}_2$ 249.352
 Alkaloid from *Dendrobium* Snowflake
 'Red Star'. Solid. $[\alpha]_{\text{D}}^{23}$ -15 (c, 1.6 in
 MeOH).

N-Me: N-Methyldendrobine

$\text{C}_{17}\text{H}_{28}\text{NO}_2^{\oplus}$ 278.414
 Quaternary alkaloid from *Dendrobium*
nobile (Orchidaceae). Prisms (Me_2CO);
 needles ($\text{Me}_2\text{CO}/\text{Et}_2\text{O}$)(as iodide). Mp
 253° (235-238°, 245-246°)(iodide).

**N-(3-Methyl-2-butenyl): N-Isopentenyl-
 dendrobine**

[40072-62-6] (bromide)
 $\text{C}_{21}\text{H}_{34}\text{NO}_2^{\oplus}$ 332.505
 Quaternary alkaloid from *Dendrobium*
nobile (Orchidaceae). Noncryst. (as
 bromide). $[\alpha]_{\text{D}}^{23}$ -33 (c, 1.4 in MeOH).
 CAS no. refers to bromide.

2ξ-Hydroxy: 2-Hydroxydendrobine

[29414-86-6]
 $\text{C}_{16}\text{H}_{25}\text{NO}_3$ 279.378
 Alkaloid from *Dendrobium findlaya-*

num (Orchidaceae). Cryst. (hexane).
 Mp 103-105°. $[\alpha]_{\text{D}}^{26}$ -45 (c, 0.73 in
 CHCl_3).

6-Hydroxy: Dendramine

[7668-75-9]
 $\text{C}_{16}\text{H}_{25}\text{NO}_3$ 279.378
 Minor alkaloid from *Dendrobium no-*
bile. Also isol. from *Dendrobium hil-*
debrandii (Orchidaceae). Mp 186-188°.
 $[\alpha]_{\text{D}}^{27}$ -27 (CHCl_3).

**2-Oxo (lactam): Mubironine A. 2-Oxo-
 dendrobine**

$\text{C}_{16}\text{H}_{23}\text{NO}_3$ 277.363
 Alkaloid from *Dendrobium* Snowflake
 'Red Star'. Solid. $[\alpha]_{\text{D}}^{23}$ -9 (c, 0.4 in
 MeOH).

**2-Oxo (lactam), 3-hydroxy: 3-Hydroxy-
 2-oxodendrobine**

[99616-00-9]
 $\text{C}_{16}\text{H}_{23}\text{NO}_4$ 293.362
 Alkaloid from the fresh stems of
Dendrobium nobile (Orchidaceae). Fine
 needles (Et_2O). Mp 210-211°. $[\alpha]_{\text{D}}^{20}$ -
 52.7 (c, 0.3 in EtOH).

[62624-23-1 8-Epimer]

Inubushi, Y. et al., *Chem. Ind. (London)*, 1964,
 1689 (abs config)

Inubushi, Y. et al., *Chem. Pharm. Bull.*, 1964,
12, 1175; 1966, **14**, 668 (*N-*
Methyldendrobine, 6-Hydroxydendrobine)

Inubushi, Y. et al., *Tetrahedron*, 1964, **20**, 2007
 (ir, pmr, struct)

Okamoto, T. et al., *Chem. Pharm. Bull.*, 1966,
14, 676 (*Dendrobine, 6-Hydroxydendrobine,*
ms, pmr)

Graneli, I. et al., *Acta Chem. Scand.*, 1970, **24**,
 1209 (isol, ir, pmr, ms, struct, 2-
Hydroxydendrobine)

Elander, M. et al., *Acta Chem. Scand.*, 1971,
25, 717 (cd)

Hedman, K. et al., *Acta Chem. Scand.*, 1972,
26, 3177 (derivs, isol, ir, pmr, config, synth)

Yamada, K. et al., *J.A.C.S.*, 1972, **94**, 8278
 (synth)

Inubushi, Y. et al., *Chem. Pharm. Bull.*, 1974,
22, 349 (synth, ir, pmr)

Kende, A.S. et al., *J.A.C.S.*, 1974, **96**, 4332
 (synth)

Corbella, A. et al., *Chem. Comm.*, 1975, 288
 (biosynth)

Suzuki, M. et al., *Chem. Lett.*, 1975, 611
 (synth, *Dendrobine, 2-Hydroxydendrobine*)

Roush, W.R. et al., *J.A.C.S.*, 1978, **100**, 3599;
 1980, **102**, 1390 (synth, ir, pmr, ms)

Wang, H. et al., *J. Nat. Prod.*, 1985, **48**, 796 (*3-*
Hydroxy-2-oxodendrobine)

Martin, S.F. et al., *J.O.C.*, 1989, **54**, 265; 1991,
56, 642 (synth)

Lee, C.H. et al., *J.A.C.S.*, 1992, **114**, 4089
 (synth)

Mori, M. et al., *J.O.C.*, 1992, **57**, 3519 (synth)
 Mori, M. et al., *Chem. Lett.*, 1993, 213 (synth)

Uesaka, N. et al., *J.O.C.*, 1994, **59**, 5633
 (synth)

Sha, C.K. et al., *J.A.C.S.*, 1997, **119**, 4130
 (synth)

Cassayre, J. et al., *J.A.C.S.*, 1999, **121**, 6072-
 6073 (synth)


Padwa, A. et al., *Org. Lett.*, 2000, **2**, 3233
 (activity, synth)

Morita, H. et al., *Tetrahedron*, 2000, **56**, 5801-
 5805 (*Mubironines*)

Cassayre, J. et al., *J. Organomet. Chem.*, 2001,
624, 316-326 (synth, ir, pmr, cmr)

Dendrochrysin**D-211**

*2-[3-(1-Methyl-2-pyrrolidinyl)-2-oxopro-
 pyl]-1-(1-oxo-3-phenyl-2-propenyl)pyrro-
 lidine*, 9CI. *N-Cinnamoylnorcusciphygrine*
 [51019-45-5]



(*E*)-form
 Absolute
 configuration

$\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_2$ 340.464

(*E*)-form [50347-38-1]

Alkaloid from *Dendrobium chrysanthum*
 (Orchidaceae). Viscous oil. $[\alpha]_{\text{D}}^{22}$ -11 (c,
 0.81 in CHCl_3).


(*Z*)-form [50347-52-9]

Alkaloid from *Dendrobium chrysanthum*.
 Viscous oil. $[\alpha]_{\text{D}}^{22}$ -19 (c, 1.92 in CHCl_3).

Ekevg. U. et al., *Acta Chem. Scand.*, 1973, **27**,
 1982 (isol, ir, uv, pmr, ms, cd, abs config,
 synth)

Dendrocrepine**D-212**

*1,3-Bis(octahydro-6-hydroxy-7-methyl-6-
 phenyl-5-indoliziny)-2-propanone*, 9CI
 [51020-39-4]



Relative
 configuration

$\text{C}_{33}\text{H}_{44}\text{N}_2\text{O}_3$ 516.722

Alkaloid from *Dendrobium crepidatum*
 (Orchidaceae). Needles (Et_2O). Mp 158-
 163° dec. Opt. inactive (racemic).


Elander, M. et al., *Acta Chem. Scand.*, 1973,
27, 1907 (isol, uv, ord, ir, pmr, ms, struct)

Pilotti, A.M. et al., *Acta Cryst. B*, 1973, **29**,
 1563 (cryst struct)

Leete, E. et al., *Tet. Lett.*, 1978, 5163 (cmr)

Dendrodoine**D-213**

[75351-10-9]



$\text{C}_{13}\text{H}_{12}\text{N}_4\text{OS}$ 272.33


Isol. from the marine tunicate *Dendrodo-*
grossularia. Cytotoxic. Cryst. (EtOAc).
 Sol. Et_2O ; poorly sol. H_2O . Mp 280-285°
 Mp 192-197° (as Ac). λ_{max} 250 (ϵ 15000);

270 (ε 12000); 300 (ε 8000) (MeOH) (Derep).

Heitz, S. *et al.*, *Tet. Lett.*, 1980, **21**, 1457-1458 (ir, pmr, ms, cryst struct)
Hogan, I.T. *et al.*, *Tetrahedron*, 1984, **40**, 681 (synth, ir, pmr, cmr, ms)

Dendronobilin A**D-214**

[572881-21-1]



Relative Configuration


C₁₉H₂₉NO₃ 319.443

Related to Dendrobine, D-210. Alkaloid from the stems of *Dendrobium nobile*. Cryst. Mp 85-87°. [α]_D²⁰ -69 (c, 0.1 in MeOH). Struct. not shown in ref. but confirmed by personal communication.

Liu, Q.F. *et al.*, *Chin. Chem. Lett.*, 2003, **14**, 278-279 (isol, pmr, cmr)
Zhao, W.M. *et al.*, *Personal Communication*,

Dendrowardine**D-215**

[50834-48-5]



Absolute Configuration

C₁₉H₃₂NO₄[⊕] 338.466


Quaternary alkaloid from *Dendrobium wardianum* (Orchidaceae). Cryst. (Me₂CO)(as chloride). Mp 168-172° (chloride). [α]_D²⁵ -28 (c, 1.1 in MeOH) (chloride).

Blomqvist, L. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 1439 (isol, ir, pmr, cd, struct)

Dendroxine**D-216**

14,10-(Methyleneoxy)dendroban-12-one, 9CI

[7668-77-1]



Absolute Configuration

C₁₇H₂₅NO₃ 291.389

Minor alkaloid from *Dendrobium nobile* (Orchidaceae). Mp 114-115°. [α]_D -30.1 (EtOH). pK_a 4.5 (50% EtOH).

N-(3-Methyl-2-butenyl): **N-Isopentenyl-dendroxine**

[34146-40-2]

C₂₂H₃₄NO₃[⊕] 360.515

Quaternary alkaloid from *Dendrobium nobile*, *Dendrobium friedricksianum* and *Dendrobium hildebrandii* (Orchidaceae). Hygroscopic needles + 1EtOH (Me₂CO/Et₂O/EtOH)(as chloride). Mp 108-120° dec. (as chloride). [α]_D²² -48 (c, 0.54 in MeOH) (chloride). CAS no. refers to chloride.

4-Hydroxy: 4-Hydroxydendroxine

[38762-85-5]

C₁₇H₂₅NO₄ 307.389

Minor alkaloid from *Dendrobium nobile* (Orchidaceae). Plates + ½ H₂O. Mp 208-210°.

6-Hydroxy: 6-HydroxydendroxineC₁₇H₂₅NO₄ 307.389

Minor alkaloid from *Dendrobium nobile* (Orchidaceae). Amorph. [α]_D²² +26 (c, 0.71 in MeOH).

6-Hydroxy, N-(3-methyl-2-butenyl): 6-Hydroxy-N-isopentenyl-dendroxine

[34222-69-0]


C₂₂H₃₄NO₄[⊕] 376.515

Quaternary alkaloid from *Dendrobium nobile*, *Dendrobium friedricksianum* and *Dendrobium hildebrandii* (Orchidaceae). Needles + 1EtOH (EtOH) (as chloride). Mp 144-156° dec. (chloride). [α]_D²² -30 (c, 0.51 in MeOH) (chloride). CAS no. refers to chloride.

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1966,**14**, 672; 676 (ir, pmr, ms, struct)Hedman, K. *et al.*, *Acta Chem. Scand.*, 1971,**25**, 1142 (derivs, isol, ir, pmr, struct)Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1972,**20**, 418 (4-Hydroxydendroxine, ir, pmr, ms, struct)**Densiflorine[†]****D-217**

7-[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]-6H-indeno[4,5-d]-1,3-dioxole-6,8(7H)-dione, 9CI


[94607-90-6]

C₂₂H₂₃NO₆ 397.427

Alkaloid from *Fumaria densiflora* (Papaveraceae).

Sener, B. *et al.*, *Int. J. Crude Drug Res.*, 1984,**22**, 79 (isol, pmr, struct)**Densiflorine[†]****D-218**

[88607-49-2]

C₂₀H₁₅NO₆ 365.342


Alkaloid from the aerial parts of *Fumaria densiflora* (Papaveraceae). Cryst. (EtOH).

Mp 249-251°.

Popova, M.E. *et al.*, *Planta Med.*, 1983, **48**, 272 (isol, uv, ir, pmr, cmr, ms, struct)

Densinine**D-219**

N-[2-(2-Acetyl-4,5-dimethoxyphenyl)ethyl]acetamide, 9CI. N,2-Diacetyl-4,5-dimethoxyphenethylamine [57621-03-1]

C₁₄H₁₉NO₄ 265.308

Alkaloid from the leaves of *Berberis densiflora*. Cryst. (EtOH or C₆H₆). Mp 128-129° (124-125°). λ_{max} 231 (ε 24100); 274 (ε 8750); 304 (ε 5500) (EtOH).

Org. Synth., 1977, **56**, 3-7 (synth, uv, pmr)Orito, K. *et al.*, *Heterocycles*, 1988, **27**, 2403-2412 (synth, ir, pmr)Khamidov, I.I. *et al.*, *Khim. Priir. Soedin.*,1997, **33**, 420-423; *Chem. Nat. Compd.*(Engl. Transl.), 1997, **33**, 323-325 (isol, uv,

ir, ms)

Denudatine**D-220**

[1356-67-8]

C₂₃H₃₅NO₅ 405.533

Struct. unknown. Alkaloid from the crude Indian drug Judwan (roots of *Delphinium denudatum*) (Ranunculaceae). Lustrous light brown prismatic cryst. (EtOH). Mp 273°. [α]_D²¹ +31.56 (EtOH).

Picrate:

Yellow micro-needles (MeOH). Mp 100°.

Reineckate:


Microprisms (Me₂CO). Mp 190°.

Singh, N. *et al.*, *J. Pharm. Pharmacol.*, 1962, **14**, 288-293 (isol)

Denudatine**D-221**

16,17-Didehydro-21-ethyl-4-methyl-7,20-cyclooctidane-11,15-diol, 9CI

[26166-37-0]



Absolute configuration

C₂₂H₃₃NO₂ 343.508

Alkaloid from the roots of *Delphinium denudatum*. Also isol. from *Aconitum jinyangense* (Ranunculaceae). Mp 248-249°. [α]_D²¹ +0.15 (EtOH).

▶GT9330000

Hydroiodide:

Cryst. (60% EtOH). Mp 267-268°.

Picrate:

Yellow cryst. (MeOH). Mp 215°.

O¹⁵-Ac: **Jynosine**. 15-Acetyldenudatine [80665-71-0]

C₂₄H₃₅NO₃ 385.545Alkaloid from *Aconitum jinyangense* (Ranunculaceae).

Di-Ac:

Microcryst. powder (abs. EtOH). Mp 128-130°.


11-Deoxy: **Gymnandine**

[150036-84-3]

C₂₂H₃₃NO 327.509Alkaloid from aerial parts of *Aconitum gymnandrum* (Ranunculaceae).Singh, N. et al., *J. Pharm. Pharmacol.*, 1962, **14**, 288 (isol, uv, ir, struct)Götz, M. et al., *Tet. Lett.*, 1969, 4369 (struct)Brisse, F. et al., *Tet. Lett.*, 1969, 4373 (cryst struct)Wright, L.H. et al., *Chem. Comm.*, 1970, 359 (cryst struct)Brisse, F. et al., *Acta Cryst. B*, 1971, **27**, 1047 (cryst struct)Chen, D. et al., *Yaoxue Xuebao*, 1981, **16**, 748; *CA*, **96**, 65677c (isol, struct, Jynosine)Ding, L.S. et al., *Yaoxue Xuebao*, 1993, **28**, 188; *CA*, **119**, 156207j (Gymnandine)Batsuren, D. et al., *Heterocycles*, 1998, **49**, 327-341 (isol, pmr, cmr)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DAP875**Deoxyamphimedine**


D-222

10-Methyl-8-oxo-8H-benzo[b]pyridido[4,3,2-de][1,8]phenanthrolium. Deoxyamphimedine [340774-96-1]


C₁₉H₁₂N₃O⁺ 298.323Quaternary alkaloid from two *Xestospongia* spp. Cytotoxic. Yellow-brown amorph. solid. Counterion not specified. λ_{max} 206 (log ε 3.92); 244 (log ε 3.93); 294 (sh) (log ε 3.52); 388 (log ε 3.51); 478 (log ε 2.59) (MeOH).Tasdemir, D. et al., *J.O.C.*, 2001, **66**, 3246-3248**Deoxycordifoline lactam**

D-223

[50764-63-1]


C₂₇H₂₆N₂O₁₀ 538.51Alkaloid from the heartwood of *Adina rubescens* (Rubiaceae). Mp 165-168° (as tetra-Ac). [α]_D -135 (MeOH) (tetra-Ac).Brown, R.T. et al., *Tet. Lett.*, 1973, 841 (uv, ms, struct)**Deoxydehydrochorismic acid** D-224

3-(1-Carboxyvinyl)benzoic acid [16929-37-6]

C₁₀H₈O₅ 208.17Isol. from *Mycena crocata*. Cryst. (MeOH/Et₂O). Mp 180-181°. λ_{max} 288 (log ε 3) (MeOH).1-Amide, 3'-Me ester: Methyl deoxydehydrochorismicamide. **Antibiotic NP 25301**. NP 25301C₁₁H₁₁NO₄ 221.212Prod. by *Streptomyces* sp. UMA-044. Cell adhesion inhibitor. Amorph. solid (CHCl₃). Mp 95-97°. λ_{max} 238; 261 (sh); 284 (MeOH).Ife, R.J. et al., *J.C.S. Perkin 1*, 1976, 1776-1783 (synth)Buchanan, M.S. et al., *Z. Naturforsch., C*, 1999, **54**, 463-468 (isol)Zhang, Q. et al., *J. Antibiot.*, 2003, **56**, 673-681 (N 25301)**Deoxymalyngamide C**

D-225

[96845-20-4]

C₂₄H₃₈ClNO₄ 440.021Minor constit. of a shallow-water variety of the marine blue-green alga *Lyngbya majuscula*. Oil. [α]_D -23 (c, 6.6 in EtOH). λ_{max} 235 (ε 6300) (EtOH) (Derep).4α,9α-Epoxyde: **Malyngamide C**

[70622-52-5]

C₂₄H₃₈ClNO₅ 456.021Metab. of a shallow-water variety of the marine blue-green alga *Lyngbya majuscula*. Oil. [α]_D^{23.5} -19.6 (c, 1.4 in CHCl₃). [α]_D -27.4 (c, 5.8 in EtOH).4α,9α-Epoxyde, Ac: **Malyngamide C acetate**

[96845-19-1]

C₂₆H₄₀ClNO₆ 498.058Minor constit. of *Lyngbya majuscula*. Oil. [α]_D -32.4 (c, 1.4 in EtOH).Deoxy: **Malyngamide K**. Dideoxymalyngamide C

[96845-21-5]

C₂₄H₃₈ClNO₃ 424.022Isol. from *Lyngbya majuscula*. [α]_D -8.4 (c, 0.3 in CHCl₃). λ_{max} 223 (ε 8900) (MeOH).Deoxy, 4α,9α-epoxyde: **8'-Deoxymalyngamide C**


[132362-43-7]

C₂₄H₃₈ClNO₄ 440.021Isol. from a shallow-water variety of a blue-green alga, *Lyngbya* sp. Yellow oil. [α]_D²⁰ +5.8 (c, 0.8 in CHCl₃). Erroneously named 8'-Deacetoxy-

lyngamide C in the paper based on an incorr. struct. shown for Malyngamide C.


Cardellina, J.H. et al., *Phytochemistry*, 1978, **17**, 2091-2095 (isol, pmr, ms)Ainslie, R.D. et al., *J.O.C.*, 1985, **50**, 2859-2862 (Malyngamide C, Malyngamide C acetate, Deoxymalyngamide C)Wright, A.D. et al., *J. Nat. Prod.*, 1990, **53**, 845-861 (Deoxymalyngamide C)Wu, M. et al., *Tetrahedron*, 1997, **53**, 15983-15990 (Malyngamide K)**6-Deoxy-8-methylbostryco-**

D-226

din
9-Hydroxy-7-methoxy-3,8-dimethylbenz[*g*]isoquinoline-5,10-dioneC₁₆H₁₃NO₄ 283.2835-Deoxy-7-methyl acc. to alternative numbering. Isol. from cultures of spore-derived mycobionts of the lichen *Haematomma* sp. Yellow powder. Mp 149-153°. λ_{max} 246 (log ε 1.7); 286 (log ε 1.55); 325 (log ε 0.38); 420 (log ε 0.49) (CHCl₃).Moriyasu, Y. et al., *Phytochemistry*, 2001, **58**, 239-241 (isol, pmr, cmr, ms)Van, T.N. et al., *Tetrahedron*, 2005, **51**, 2295-2300 (synth)**Deoxynupharidine**

D-227

4-(3-Furanyl)octahydro-1,7-dimethyl-2H-quinolizine [1143-54-0]



Absolute Configuration

C₁₅H₂₃NO 233.353Alkaloid from *Nuphar japonicum*, *Nuphar luteum* ssp. *variegatum*, *Nuphar pumila* and from the scent glands of the Canadian beaver *Castor fiber*. Mp 21-22°. Bp₈ 112-115°. [α]_D -112.5.

Hydrochloride: Mp 262°.

Picrate: Mp 153°.

N-Oxide (R-): **Nupharidine**

[468-89-3]

C₁₅H₂₃NO₂ 249.352Alkaloid from *Nuphar japonicum* and *Nuphar pumila* (Nymphaeaceae). Mp 222°. [α]_D +14.5 (H₂O).

N-Oxide, picrate: Mp 176°.

3,4-Didehydro: **Dehydrodeoxynupharidine**C₁₅H₂₁NO 231.337Minor alkaloid from the roots of *Nuphar japonicum* (Nymphaeaceae). Unstable liq.; needles or prisms

(MeOH)(as perchlorate). Mp 159-161° (perchlorate). Bp₃ 125° (bath). $[\alpha]_{\text{D}}^{20}$ +130.1 (CHCl₃) (perchlorate).

9,10-Didehydro: Nupharopumiline

[63947-66-0]
C₁₅H₂₁NO 231.337

Alkaloid from the rhizomes of *Nuphar pumila* (Nymphaeaceae). Cryst. (CCl₄). Mp 195-197°. $[\alpha]_{\text{D}}^{20}$ +27 (CHCl₃).

7-Hydroxy: Nupharolutine

[38681-18-4]

C₁₅H₂₃NO₂ 249.352

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Cryst. (MeOH/Me₂CO). Mp 96-98°. $[\alpha]_{\text{D}}^{20}$ -105 (CHCl₃).

8 α -Hydroxy: Isocastoramine

[60125-06-6]

C₁₅H₂₃NO₂ 249.352

Alkaloid from the scent glands of the Canadian beaver *Castor fiber*. Needles (CH₂Cl₂/pentane). Mp 109°. $[\alpha]_{\text{D}}^{20}$ -124 (c, 1.0 in CHCl₃).

9 ξ -Hydroxy: Nupharolidine

[24181-81-5]

C₁₅H₂₃NO₂ 249.352

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Mp 110°. Stereochem. not certain.

9-Hydroxy, hydrochloride: Mp 240-245° dec.

12-Hydroxy: Castoramine

[6874-86-8]

Main basic constit. from the scent glands of the Canadian beaver (*Castor fiber* and *Castor canadensis*). Mp 64.5-65.5°. Bp_{0.08} 125-135°. $[\alpha]_{\text{D}}^{20}$ -80 (c, 1 in CHCl₃).

6 ξ ,7-Dihydroxy: Deoxynupharidine-6,7-diol

C₁₅H₂₃NO₃ 265.352

Alkaloid detected in the rhizomes of *Nuphar luteum* subsp. *macrophyllum* (Nymphaeaceae). Sticky oil. Uncertain at present whether this alkaloid is a true plant metab. Dimerizes slowly to 6,7-Oxidodeoxynupharidine dimer, O-161 when stored neat at 0°.

1-Epimer: 1-Epideoxynupharidine

C₁₅H₂₃NO 233.353

Alkaloid from the scent glands of *Castor fiber*. Oil. $[\alpha]_{\text{D}}^{20}$ -36 (c, 0.5 in CHCl₃).

7-Epimer: 7-Epideoxynupharidine

C₁₅H₂₃NO 233.353

Alkaloid from the rhizomes of *Nuphar luteum* subsp. *variegatum*, *Nuphar pumila* (Nymphaeaceae) and the scent glands of *Castor fiber*. Oil. $[\alpha]_{\text{D}}^{25}$ -89 (c, 3.5 in 95% in EtOH). $[\alpha]_{\text{D}}^{20}$ -75 (c, 1.0 in CHCl₃).

7-Epimer, hydrochloride:

Cryst. (MeOH/Et₂O). Mp 255-258°. $[\alpha]_{\text{D}}^{25}$ -39.7 (c, 0.79 in 95% in EtOH).

7-Epimer, N-oxide: 7-Epinupharidine

[29487-76-1]

C₁₅H₂₃NO₂ 249.352

Alkaloid from the rhizomes of *Nuphar pumila* (Nymphaeaceae). Mp 199-202° (synthetic).

7-Epimer, 6 ξ ,7-dihydroxy: 7-Epideoxynu-

pharidine-6,7-diol

C₁₅H₂₃NO₃ 265.352

Alkaloid detected in the rhizomes of *Nuphar luteum* ssp. *macrophyllum* (Nymphaeaceae). Mp 176-177°. $[\alpha]_{\text{D}}^{25}$ -64 (c, 0.5 in CHCl₃) (extrapolated; shows mutarotation). Uncertain at present whether this is a true plant metab.

1,7-Diepimer: 1-Epi-7-epideoxynupharidine

C₁₅H₂₃NO 233.353

Alkaloid from the scent glands of *Castor fiber*. Oil. $[\alpha]_{\text{D}}^{20}$ -70 (c, 1.0 in CHCl₃).

Kotake, M. *et al.*, *Annalen*, 1957, **606**, 148 (struct)

Valenta, Z. *et al.*, *Tet. Lett.*, 1959, No. **12**, 1 (*Castoramine*, *isol*, *ir*, *pmr*, *struct*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1961, **94**, 3151 (*synth*, *ir*)

Bohlmann, F. *et al.*, *Tetrahedron*, 1963, **19**, 195 (*Castoramine*, *synth*)

Arata, Y. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1394; 1965, **13**, 907; 1968, **16**, 2074 (*Dehydrodeoxynupharidine*)

Achmatowicz, O. *et al.*, *Tet. Lett.*, 1964, 927 (*ms*)

Wróbel, J.T. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1969, **43**, 997 (*Nupharolidine*)

Oda, K. *et al.*, *J.C.S. (B)*, 1970, 1450 (*cryst struct*)

Wong, C.F. *et al.*, *J.O.C.*, 1970, **35**, 517 (*config*)

Wong, C.F. *et al.*, *Phytochemistry*, 1970, **9**, 659 (*7-Epideoxynupharidine*)

LaLonde, R.T. *et al.*, *J.A.C.S.*, 1971, **93**, 2501; 1972, **94**, 8522 (*7-Epinupharidine, diols*)

Wróbel, J.T. *et al.*, *Can. J. Chem.*, 1972, **50**, 1831 (*Nupharolutine*)

Martin, T.I. *et al.*, *Can. J. Chem.*, 1974, **52**, 2705 (*pmr*)

LaLonde, R.T. *et al.*, *Can. J. Chem.*, 1975, **53**, 1714 (*cmr*)

Maurer, B. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 1169 (*Deoxynupharidine, Epideoxynupharidines, Castoramine, Isocastoramine*)

Yasuda, S. *et al.*, *Heterocycles*, 1977, **6**, 391 (*Epideoxynupharidines, synth*)

Peura, P. *et al.*, *Phytochemistry*, 1977, **16**, 1122 (*Nupharopumiline*)

Szychowski, J. *et al.*, *Chem. Ind. (London)*, 1978, 273 (*synth*)

Hwang, Y.C. *et al.*, *J.O.C.*, 1985, **50**, 2719 (*synth, pmr, cmr, ms*)


Miyazawa, M. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 1059-1063 (*Nupharolutine, Castoramine, isol, pmr, cmr, ms*)

Moran, W.J. *et al.*, *Org. Lett.*, 2003, **5**, 3427-3429 (*synth*)

Katoh, M. *et al.*, *Heterocycles*, 2006, **69**, 193-216 (*synth*)

Deoxypumiliotoxin 251H D-228

6-(Hexahydro-8-methyl-6(5H)-indolizinyldiene)-5-methyl-2-hexanol, 9CI [164301-81-9]



Relative configuration


C₁₆H₂₉NO 251.411

Trace alkaloid from skin extracts of the dendrobatid frog *Epipedobates tricolor*.

Jain, P. *et al.*, *J. Nat. Prod.*, 1995, **58**, 100 (*isol, ir, pmr, ms, struct*)

Deoxyrubroflavin D-229

2-[4-Hydroxy-2,6-bis(methylthio)phenyl]diazene-carboxamide. 4-Hydroxy-2,6-bis(methylthio)phenyl-1-azofornamide. 2,6-Bis(methylthio)-1,4-benzoquinone 1-semicarbazone



C₉H₁₁N₃O₂S₂ 257.337

The parent compd. and its S-oxides show solv.-dependent tautomerism between semicarbazone and azophenol forms. Minor pigment from the puffball mushroom *Calvatia rubro-flava*. Red powder. Mp 168-169°. λ_{max} 212 (log ϵ 4.06); 232 (log ϵ 4.18); 253 (log ϵ 4.16); 278 (log ϵ 4.05); 310 (sh) (log ϵ 3.58); 387 (log ϵ 3.97) (MeOH).

3-S-Oxide (S-): Rubroflavin. CC 3.

Compound CC 3

[169758-58-1]

C₉H₁₁N₃O₃S₂ 273.336

Isol. from *Calvatia craniformis*. Major pigment from *Calvatia rubro-flava*. Cytotoxic agent. Red needles (MeOH) or red needles + 1H₂O. Mp 192-194° (184-185° dec.). $[\alpha]_{\text{D}}^{25}$ -1740 (c, 0.4 in Py). $[\alpha]_{\text{D}}^{22}$ -2180 (c, 0.07 in MeOH). λ_{max} 248 (log ϵ 4.27); 285 (log ϵ 4.28); 355 (log ϵ 4.1) (MeOH). λ_{max} 214 (log ϵ 4.01); 243 (log ϵ 3.92); 252 (log ϵ 3.59); 275 (log ϵ 3.65); 295 (log ϵ 3.61); 435 (log ϵ 4.03); 460 (sh) (log ϵ 3.97) (MeOH).

3,3-S-Dioxide: Craniformin†. CCU 18.

Compound CCU 18

[169758-57-0]

C₉H₁₁N₃O₄S₂ 289.336

Isol. from *Calvatia craniformis*. Cytotoxic agent. Red plates (MeOH). Mp 192-194°. The original authors used the name Craniformin for this dioxide, but Fugmann *et al* apply it to the trioxide below.

3,5-S-Dioxide: Oxyrubroflavin

C₉H₁₁N₃O₄S₂ 289.336

Minor pigment from *Calvatia rubro-flava*. Amorph. orange-red solid. $[\alpha]_{\text{D}}$ -840 (c, 0.1 in MeOH). Isol. as a 1:1 mixt. of 5R- and 5S-diastereomers. Dec. at 200-210°. Prob. an artifact derived by non-enzymic oxidn. λ_{max} 197 (log ϵ 4.19); 245 (log ϵ 3.98); 280 (sh) (log ϵ 3.88); 352 (log ϵ 4.01); 455 (sh) (log ϵ 3.12) (MeOH).

3S,5,5-S-Trioxide: Craniformin†

C₉H₁₁N₃O₅S₂ 305.335


Isol. from *Calvatia rubro-flava*. Orange-red powder (synthetic). Mp 168-170°. $[\alpha]_{\text{D}}^{20}$ -1034 (MeOH). λ_{max} 198 (log ϵ 4.29); 253 (log ϵ 3.96); 258 (sh)

(log ϵ 3.77); 345 (log ϵ 3.96); 430 (log ϵ 3.67) (MeOH).

Takaishi, Y. *et al.*, *Phytochemistry*, 1997, **45**, 997-1001 (*isol, uv, ir, pmr, cmr, ms*)
 Fugmann, B. *et al.*, *Eur. J. Org. Chem.*, 2001, 3097-3104 (*isol, uv, ir, cd, ms, pmr, cmr, abs config*)

Deplancheine D-230

3-Ethylidene-1,2,3,4,6,7,12,12b-octa-hydroindolo[2,3-a]quinolizine, 9CI



C₁₇H₂₀N₂ 252.358

A bisnor-corynanthe alkaloid. Abs. config. revised in 1986.

(R)-form [74559-69-6]

Alkaloid from *Alstonia deplanchei* (Apocynaceae). Cryst. (Et₂O). Mp 115°. [α]_D²⁰ +56 (c, 1 in CHCl₃).

10-Methoxy-10-Methoxydeplancheine

[80787-52-6]

C₁₈H₂₂N₂O 282.385

Alkaloid from the leaves of *Alstonia lanceolifera* (Apocynaceae). Cryst. (Me₂CO). Mp 75°. [α]_D +33 (c, 0.9 in CHCl₃).

(S)-form

Synthetic. Cryst. (Et₂O/hexane). Mp 140.5-141.5°. [α]_D²⁰ -52 (c, 1.0 in CHCl₃). [α]_D -64.9 (c, 1.0 in EtOH).


(±)-form [56491-03-3]

Synthetic. Cryst. (MeOH or Et₂O). Mp 140-142° Mp 161.5°.

Thielke, D. *et al.*, *Chem. Ber.*, 1975, **108**, 1791 (*synth, uv, ir, pmr, ms*)
 Besselièvre, R. *et al.*, *Tet. Lett.*, 1980, **21**, 63 (*isol, synth*)
 Ashcroft, W.R. *et al.*, *Tet. Lett.*, 1980, **21**, 2341 (*synth*)
 Hämeilä, M. *et al.*, *Acta Chem. Scand., Ser. B*, 1981, **35**, 217 (*synth, ir, pmr, ms*)
 Petitfrère-Auvray, N. *et al.*, *Phytochemistry*, 1981, **20**, 1987 (10-Methoxydeplancheine)
 Overman, L.E. *et al.*, *J.O.C.*, 1982, **47**, 5297 (*synth, uv, ir, pmr, cmr*)
 Calabi, L. *et al.*, *Tet. Lett.*, 1982, **23**, 2139 (*synth*)
 Lesma, G. *et al.*, *J.C.S. Perkin I*, 1984, 1593 (*synth*)
 Meyers, A.I. *et al.*, *J.O.C.*, 1986, **51**, 3108 (*synth, ir, pmr, cmr, abs config*)
 Jokela, R. *et al.*, *Planta Med.*, 1987, **53**, 386 (*synth, ir, pmr, cmr, ms*)
 Mandal, S.B. *et al.*, *J.O.C.*, 1988, **53**, 4236 (*synth*)
 Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2822 (*synth*)
 Rosenmund, P. *et al.*, *Annalen*, 1990, 233; 1992, 1321 (*synth*)
 Sankar, P.J. *et al.*, *Heterocycles*, 1991, **32**, 1109 (*synth*)
 Itoh, T. *et al.*, *Heterocycles*, 2001, **55**, 1165-1171 (*synth*)
 Allin, S.M. *et al.*, *J.O.C.*, 2005, **70**, 357-359 (*synth*)
 Sydorenko, N. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 2140-2144 (*synth*)
 Chang, M.-Y. *et al.*, *Tetrahedron*, 2005, **61**, 585-591 (*synth*)

Dercitine

[115141-47-4]



C₂₁H₂₀N₄S 360.482

Regiochemistry of the thiazole moiety revised in 1992. Alkaloid from a deep water marine sponge *Dercitus* sp. and *Stelletta* sp. Exhibits antitumour, antiviral and immunosuppressive activities. Hygroscopic deep violet powder. Mp 168°. λ_{\max} 245 (ϵ 13800); 307 (ϵ 16900); 361 (ϵ 3900); 541 (ϵ 1800) (MeOH) (Derep). λ_{\max} 234 (ϵ 13800); 245 (ϵ 14500); 307 (ϵ 19500); 354 (ϵ 4500); 515 (ϵ 12600); 261 (ϵ 11750); 293 (ϵ 14450); 405 (ϵ 6000); 527 (ϵ 2050); 558 (ϵ 1750); 610 (ϵ 1550) (MeOH/NaOH) (Berdy).

► UV1198950

Gunawardana, G.P. *et al.*, *J.A.C.S.*, 1988, **110**, 4856 (*isol, pmr, cmr, ms*)
 Burres, N.S. *et al.*, *Cancer Res.*, 1989, **49**, 5267-5274 (*activity*)
 Bishop, M.J. *et al.*, *J.A.C.S.*, 1992, **114**, 10081 (*synth*)
 Gunawardana, G.P. *et al.*, *J.O.C.*, 1992, **57**, 1523 (*struct*)
 Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1995, **117**, 12460 (*synth*)

Desdamethine


U 11994. Antibiotic U 11994

C₈H₁₄N₂OS 186.277

Pyrrolidine antibiotic. Struct. unknown. Prod. by *Streptomyces caelestis* with Methionine. Mainly active against gram-positive bacteria. Sol. H₂O, DMF; fairly sol. MeOH; poorly sol. Et₂O, hexane. Meyer, C.E. *et al.*, *Antimicrob. Agents Chemother.*, 1965, 850 (*isol, ir*)

Deserpideine

[2671-59-2]



C₃₂H₃₆N₂O₈ 576.645

Alkaloid from *Rauwolfia nitida* (Apocynaceae). Antihypertensive agent, tranquilliser. Mp 149-152°. [α]_D²⁵ -133 (c, 1 in Py). [α]_D²⁵ -108 (CHCl₃). Log P 2.87 (calc).

Hydrochloride: [15215-04-0]

Cryst. (MeOH). Mp 277° dec. [α]_D²⁵ -98.5 (c, 0.5 in CHCl₃).

D-231

11-Methoxy- **Raujemidine**. 19-Dehydroreserpine

[549-70-2]

C₃₃H₃₈N₂O₉ 606.671

Alkaloid from *Rauwolfia canescens* (*R. tetraphylla*) (Apocynaceae). Shows about half the antihypertensive activity of Reserpine, R-52; of no practical importance. Square plates + 1H₂O (MeOH). Mp 144-150°. [α]_D²⁵ -88 (CHCl₃). pK_a 5.53 (40% MeOH aq.).

11-Methoxy; perchlorate: Mp 253-257°.

11-Methoxy, N⁴-oxide: **Raujemidine N-oxide**

C₃₃H₃₈N₂O₁₀ 622.671

Alkaloid from *Rauwolfia canescens* (Apocynaceae). Mp 215-217°. Details unpubl. CAS no. not found to 2007.


Ulshafer, P.R. *et al.*, *J.O.C.*, 1956, **21**, 923 (*isol, Raujemidine*)

Smith, E. *et al.*, *J.A.C.S.*, 1967, **89**, 2469-2476 (*isol, struct*)

Deserpidine, BAN, INN D-234

11-Desmethoxyreserpine. Canescine†.

Harmonyl. Raunormine. Recanescine. Reserpidine. Lilly 22641 [131-01-1]



C₃₂H₃₈N₂O₈ 578.661

Alkaloid from *Rauwolfia canescens* and several other *Rauwolfia* spp., *Tonduzia longifolia* and *Ochrosia vieillardii* (Apocynaceae). Antihypertensive, neuroleptic tranquilliser agent. Trimorphic. Mp 228-232° (220-222° dec.). [α]_D²⁴ -137 (CHCl₃). [α]_D -118 (c, 1 in CHCl₃). Log P 3.35 (calc).

► Exp. reprod. effects. LD₅₀ (mus, orl) 500 mg/kg. ZG0875000

Hydrochloride:

Cryst. + 1MeOH (MeOH). Mp 241-243° dec.

19,20-Didehydro: see Deserpideine, D-233

Schlittler, E. *et al.*, *Experientia*, 1955, **11**, 64-65 (*isol*)

Carletti, A. *et al.*, *Experientia*, 1955, **11**, 98-99 (*pharmacol*)

Jurg, A. *et al.*, *J. Pharmacol. Exp. Ther.*, 1955, **114**, 10-13 (*pharmacol*)

Klohs, M.W. *et al.*, *J.A.C.S.*, 1955, **77**, 4084-4087 (*isol, struct*)

Neuss, N. *et al.*, *J.A.C.S.*, 1955, **77**, 4087-4090 (*isol, uv, ir, struct*)

Cronheim, G. *et al.*, *Proc. Soc. Exp. Biol. Med.*, 1955, **89**, 21-23 (*pharmacol*)

Weichet, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1961, **26**, 1529-1536 (*synth*)

Ben-David, M. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1968, **171**, 274-284 (*pharmacol*)

Creveling, C.R. *et al.*, *J. Med. Chem.*, 1968, **11**, 596-598 (*activity*)


- Levin, R.H. *et al.*, *J.O.C.*, 1973, **38**, 1983-1986 (cmr)
 Timmins, P. *et al.*, *Phytochemistry*, 1974, **13**, 1997 (isol)
 Pantarotto, C. *et al.*, *Adv. Mass Spectrom. Biochem. Med.*, 1977, **2**, 351 (ms)
 Szantay, C. *et al.*, *Heterocycles*, 1977, **7**, 155-160 (synth)
 Sakai, S. *et al.*, *Heterocycles*, 1978, **10**, 67-71 (synth)
 Szantay, C. *et al.*, *Annalen*, 1983, 1292-1309 (synth)
 Miyata, O. *et al.*, *Heterocycles*, 1984, **22**, 1041-1044 (synth)
 Lounasmaa, M. *et al.*, *Heterocycles*, 1985, **23**, 371-375 (pmr)
 Naito, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 901-906 (synth)
 Baxter, E.W. *et al.*, *J.A.C.S.*, 1990, **112**, 7682-7692 (synth)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 353
 Varchi, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1629-1631 (synth)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, RDF000

Desferroferrithiocin D-235

4,5-Dihydro-2-(3-hydroxy-2-pyridinyl)-4-methyl-4-thiazolecarboxylic acid, 9CI.

Desferrithiocin

[76045-30-2]



C₁₀H₁₀N₂O₃S 238.267

(S)-form

Isol. from *Streptomyces antibioticus* as Fe complex. Mp 90-92°. [α]_D +30.1 (c, 1.01 in MeOH). λ_{\max} 200 (ε 15150); 308 (ε 6200); 382 (ε 2140) (H₂O) (Berdy).

Fe complex: **Ferrithiocin**

[76082-64-9]

Metab. of *Streptomyces antibioticus*.

Dark red-brown powder. Mp 160° dec.

[α]_D -578 (c, 0.0064 in H₂O).

[105635-69-6]

Naegeli, H.U. *et al.*, *Helv. Chim. Acta*, 1980,

63, 1400

Mulqueen, G.C. *et al.*, *Tetrahedron*, 1993, **49**,

5359 (synth)

Bergeron, R.J. *et al.*, *J. Med. Chem.*, 1994, **37**,


2889 (metab)

Desformocorymine D-236

Methyl 3-hydroxy-1-methyl-2,4(1H)-cyclo-3,4-secoakammilan-17-oate, 9CI.

Deformylcorymine

[21290-54-0]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Hunteria eburnea*, *Hunteria elliptii* and *Hunteria zeylanica* (Apocynaceae). Also obt. from Corymine, in D-410 on treatment with KOH aq. Mp 170-171°. [α]_D²⁴ -90 (EtOH).

Kiang, A.K. *et al.*, *Proc. Chem. Soc., London*, 1962, 298

Kompiš, I. *et al.*, *Coll. Czech. Chem. Comm.*,

1968, **33**, 4328 (synth, uv, ir, ms, ord)

Morfaux, A.M. *et al.*, *Phytochemistry*, 1978,


17, 167 (isol)

Subhadhirasakul, S. *et al.*, *Chem. Pharm.*

Bull., 1994, **42**, 2645

Desmodimine D-237

[150036-83-2]



C₁₂H₁₅NO₄ 237.255

Alkaloid from aerial parts of *Desmodium styracifolium* (Fabaceae).


Yang, J.S. *et al.*, *Yaouxue Xuebao*, 1993, **28**, 197;

CA, **119**, 156209m

Desmotrichinin D-238

5-[(3,5-Dihydroxyphenoxy)methyl]-3-pyridinol

[691394-44-2]



C₁₂H₁₁NO₄ 233.223

Alkaloid from the pseudobulb of *Desmotrichum fimbriatum* (preferred genus name *Flickingeria*). Yellow cryst. (CHCl₃/MeOH). Mp 195-198°. [α]_D²⁰ +25 (c, 0.1 in MeOH). λ_{\max} 276 (log ε 5.2); 335 (log ε 6.92); 418 (log ε 7.2) (MeOH).


Ali, A. *et al.*, *Pharmazie*, 2003, **58**, 762-763

(isol, pmr, cmr, ms)

Dethiotetrakis(methylthio)-chetomin D-239

2,5:2',5'-Dide(epidithio)-2,2',5',5'-tetrakis(methylthio)chetomin, 9CI. Dethiotetra(methylthio)chetomin

[84260-71-9]




Absolute Configuration

C₃₅H₄₂N₆O₆S₄ 771.017

Prod. by *Chaetomium cochliodes* and *Chaetomium globosum*. Antimicrobial agent. Cryst. (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 191-193°. [α]_D +174 (CHCl₃). λ_{\max} 212 (ε 72200); 288 (ε 10000); 296 (ε 10500) (MeOH) (Berdy).

Kikuchi, T. *et al.*, *Chem. Pharm. Bull.*, 1982,

30, 3846 (isol, uv, ir, pmr, cmr, struct)

Detoxinine D-240

C₇H₁₃NO₄ 175.184

(-)-form [54963-44-9]

Structural component of Detoxin D₁.

[α]_D²³ -4.1 (c, 0.5 in H₂O).

(±)-form

Prisms. Mp 200° dec.

Haüsler, J. *et al.*, *Annalen*, 1983, 982 (synth)

Ohfuno, Y. *et al.*, *Tet. Lett.*, 1984, **25**, 4133

(synth)

Ewing, W.R. *et al.*, *Tetrahedron*, 1986, **42**, 2421

(synth, ir, pmr)

Li, W.-R. *et al.*, *Heterocycles*, 1993, **36**, 359


(rev)

Flögel, O. *et al.*, *Chem. Eur. J.*, 2003, **9**, 1405-

1415 (synth)

Diabroticin A D-241

[154163-98-1]



C₁₆H₂₈N₆O₆ 400.434

Isol. from *Bacillus subtilis*. Insecticide.

[α]_D²³ -6.7 (c, 2 in MeOH). λ_{\max} 275 (no solvent reported).

1'-Deoxy: **Diabroticin B**


[154163-96-9]

C₁₆H₂₈N₆O₅ 384.434

Isol. from *Bacillus subtilis*. Insecticide.

Stonard, R.J. *et al.*, *ACS Symp. Ser.*, 1994,

551, 25-36 (isol, pmr, cmr, ms)

3,5-Diacyl-1,4-dihydro-4-nonylpyridines D-242**3,5-Didecanoyl-1,4-dihydro-4-nonylpyridine**

1,1'-(1,4-Dihydro-4-nonyl-3,5-pyridinediyl)bis[1-decanone], 9CI

1,2-Diacylglyceryl-**3-(*O*-carboxyhydroxymethyl)choline**

[158528-00-8]

C₃₄H₆₁NO₂ 515.861Alkaloid from *Houttuynia cordata* (Yu Xing Cao).**3-Decanoyl-5-dodecanoyl-1,4-dihydro-4-nonylpyridine**


1-[1,4-Dihydro-4-nonyl-5-(1-oxododecyl)-3-pyridinyl]-1-dodecanone, 9CI

[158528-01-9]


C₃₆H₆₅NO₂ 543.915Alkaloid from *Houttuynia cordata* (Yu Xing Cao).**3,5-Didodecanoyl-1,4-dihydro-4-nonylpyridine**

1,1'-(1,4-Dihydro-4-nonyl-3,5-pyridine-diyl)bis[1-dodecanone], 9CI


[158528-02-0]

C₃₈H₆₉NO₂ 571.969Alkaloid from *Houttuynia cordata* (Yu Xing Cao).Proebstle, A. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 235-240**1,2-Diacylglyceryl-****D-243****3-(*O*-carboxyhydroxymethyl)choline**
DGCC


R and R' are long chain acyl residues. Betaine lipid present in some plants and algae.

Kato, M. *et al.*, *Phytochemistry*, 1996, **42**, 1341**1,2-Diacylglyceryl-****D-244*****O*-2'-(hydroxymethyl)-*N,N,N*-trimethyl-β-alanine**
DGTA

R and R' are long chain acyl residues. Isol. from various marine algae.

Kato, M. *et al.*, *Phytochemistry*, 1996, **42**, 1341Vaskovsky, V.E. *et al.*, *Phytochemistry*, 1996, **42**, 1347**4-*O*-(1,2-Diacylglyceryl)-****D-245*****N,N,N*-trimethylhomoserine**
DGTS

R and R' are long-chain acyl residues. Betaine lipid present in fungi, lichens, most ferns, some higher plants and green algae.

Sato, N. *et al.*, *Phytochemistry*, 1984, **23**, 1625-1627 (bibl)**2,5-Dialkylpyrrolidines**Dembitsky, V.M. *et al.*, *Prog. Lipid Res.*, 1996, **35**, 1-51 (rev)Kuenzler, K. *et al.*, *Phytochemistry*, 1997, **46**, 883-892 (occur)Vaskovsky, V.E. *et al.*, *Phytochemistry*, 1998, **47**, 755-760 (occur)Rozentsvet, O.A. *et al.*, *Phytochemistry*, 2000, **54**, 401-407 (occur)**2,5-Dialkylpyrrolidines****D-246**R, R' = Alkyl groups, various dehydro analogues also found. Components of the defence secretions of the ants *Solenopsis* and *Monomorium* spp.**2-Ethyl-5-pentylpyrrolidine**C₁₁H₂₃N 169.3091,2-Didehydro: 5-Ethyl-3,4-dihydro-2-pentyl-2H-pyrrole, 9CI. 2-Ethyl-5-pentyl-Δ¹-pyrroline

[61772-94-9]

C₁₁H₂₁N 167.294From *Solenopsis punctaticeps*.1,5-Didehydro: 2-Ethyl-3,4-dihydro-5-pentyl-2H-pyrrole, 9CI. 5-Ethyl-2-pentyl-Δ¹-pyrroline

[61772-95-0]

C₁₁H₂₁N 167.294From *Solenopsis punctaticeps*.**2-Ethyl-5-heptylpyrrolidine** [61772-96-1]C₁₃H₂₇N 197.363From *Solenopsis punctaticeps*.1,2-Didehydro: 2-Ethyl-5-heptyl-Δ¹-pyrrolineC₁₃H₂₅N 195.347From *Solenopsis punctaticeps*.1,5-Didehydro: 2-Ethyl-5-heptyl-3,4-dihydro-2H-pyrrole, 9CI. 5-Ethyl-2-heptyl-Δ¹-pyrroline

[61772-93-8]

C₁₃H₂₅N 195.347From *Solenopsis punctaticeps*.**2-Ethyl-5-undecylpyrrolidine**C₁₇H₃₅N 253.47Constit. of the venom of *Monomorium* sp.**2-Ethyl-5-(12-tridecenyl)pyrrolidine, 9CI**C₁₉H₃₇N 279.508Constit. of the venom of *Monomorium* sp.**2-Butyl-5-propylpyrrolidine**C₁₁H₂₃N 169.309Isol. from the venom of *Myrmecaria melanogaster*. Obt. as a mixt. of *cis*- and *trans*-isomers.**2-Butyl-5-pentylpyrrolidine****Dendrobates Alkaloid 197B**

[19501-71-4]

C₁₃H₂₇N 197.363From *Solenopsis punctaticeps* and *Monomorium* sp. Also isol. from skin extracts of *Dendrobates histrionicus*.**2-Butyl-5-heptylpyrrolidine** [61772-92-7]C₁₅H₃₁N 225.417From *Solenopsis fugax*, *Solenopsis punctaticeps*, *Monomorium latinode* and cultures of *Streptomyces longispororuber*. The *S. longispororuber* isolate was the (2*R*,5*R*)-enantiomer.***N*-Me: 2-Butyl-5-heptyl-1-methylpyrrolidine, 9CI**

[83688-93-1]

C₁₆H₃₃N 239.443Venom from *Monomorium latinode*.1,5-Didehydro: 2-Butyl-5-heptyl-3,4-dihydro-2H-pyrrole, 9CI. 5-Butyl-2-heptyl-Δ¹-pyrroline

[83688-87-3]

C₁₅H₂₉N 223.401Venom from *Monomorium latinode*.**2-Butyl-5-nonylpyrrolidine**

8'',9''-Didehydro: 2-Butyl-5-(8-nonenyl)pyrrolidine

[120091-01-2]

C₁₇H₃₃N 251.454From *Monomorium* sp.**2-Hexyl-5-pentylpyrrolidine** [73427-38-0]C₁₅H₃₁N 225.417From *Solenopsis molesta* and *Solenopsis texanus*.

5',6'-Didehydro: 2-(5-Hexenyl)-5-pentylpyrrolidine, 9CI

[58623-41-9]

C₁₅H₂₉N 223.401Venom from *Monomorium pharaonis*.**2-Hexyl-5-nonylpyrrolidine** [81943-71-7]C₁₉H₃₉N 281.524From *Monomorium* spp.

5',6',8'',9''-Tetradehydro: 2-(5-Hexenyl)-5-(8-nonenyl)pyrrolidine

[81943-70-6]

C₁₉H₃₅N 277.492Component of venom from *Monomorium* sp. and from *Chelaner antarcticus*. The *C. antarcticus* isolate was shown to have the *trans*-config.5',6',8'',9''-Tetradehydro, *N*-Me: 2-(5-Hexenyl)-1-methyl-5-(8-nonenyl)pyrrolidineC₂₀H₃₇N 291.519Venom from *Monomorium* sp.**2-Heptyl-5-(5-hexenyl)pyrrolidine**C₁₇H₃₃N 251.454From *Monomorium* sp.**2-(5-Hexenyl)-5-nonylpyrrolidine** [100594-88-5]C₁₉H₃₇N 279.508From *Monomorium* sp.*N*-Me: 2-(5-Hexenyl)-1-methyl-5-nonylpyrrolidineC₂₀H₃₉N 293.535From *Monomorium* sp.1,2,8'',9''-Tetradehydro: 2-(5-Hexenyl)-5-(8-nonenyl)-Δ¹-pyrrolineC₁₉H₃₃N 275.476From *Monomorium* sp.1,5,8'',9''-Tetradehydro: 5-(5-Hexenyl)-2-(8-nonenyl)-Δ¹-pyrrolineC₁₉H₃₃N 275.476

From *Monomorium* sp.

[116558-88-4, 116558-86-2, 114640-33-4, 114640-32-3, 114640-29-8, 114640-28-7, 103833-64-3, 103833-14-3, 95018-46-5, 95018-45-4, 81943-78-4, 131682-66-1, 75372-27-9, 71732-78-0, 71732-77-9, 75372-32-6]

Pedder, D.J. *et al.*, *Tetrahedron*, 1976, **32**, 2275
Jones, T.H. *et al.*, *Tet. Lett.*, 1979, 1031-1034
(2-Hexyl-5-pentylpyrrolidine)

Blum, M.S. *et al.*, *Naturwissenschaften*, 1980, **67**, 144

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1982, **8**, 285
Jones, T.H. *et al.*, *Tetrahedron*, 1982, **38**, 1949-1958 (rev)

Attygalle, A.B. *et al.*, *Chem. Soc. Rev.*, 1984, **13**, 245

Daly, J.W. *et al.*, *J. Nat. Prod.*, 1986, **49**, 265
(Alkaloid 197B)

Jones, T.H. *et al.*, *J.O.C.*, 1986, **51**, 2712

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 35
(isol)

Jones, T.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 375
(isol, pmr, cmr)

Bäckvall, J.-E. *et al.*, *J.O.C.*, 1990, **55**, 826
(synth)

Machinaga, N. *et al.*, *J.O.C.*, 1991, **56**, 1386
(synth)

Bloch, R. *et al.*, *Tetrahedron: Asymmetry*, 1994, **5**, 745-750 (Alkaloid 197B, synth)


Kumagai, K. *et al.*, *J. Antibiot.*, 2000, **53**, 467-473 (2-Butyl-5-heptylpyrrolidine)

Davis, F.A. *et al.*, *J.O.C.*, 2006, **71**, 2779-2786
(Alkaloid 197B, synth)

Jones, T.H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 160-168 (2-Butyl-5-propylpyrrolidine)

2,4-Dialkylthiazoles

D-247



Maillard prods. present in cooked food volatiles.

2-Isopropyl-4-methylthiazole

4-Methyl-2-(1-methylethyl)thiazole, 9CI, FEMA 3555

[15679-13-7]

C₇H₁₁NS 141.237

Present in durian (*Durio zibethinus*), red tomatoes, yeast extract and coriander seed oil. Liq. with camphorous, nutty odour. d₄²⁰ 1.01. Bp₁₀ 59-62°. n_D²⁰ 1.4989.

4-Isopropyl-2-methylthiazole

2-Methyl-4-(1-methylethyl)thiazole, 9CI [32272-52-9]

C₇H₁₁NS 141.237

2,4-Diethylthiazole, 9CI [32272-49-4]

C₇H₁₁NS 141.237

2-Butyl-4-methylthiazole, 9CI [76572-47-9]

C₈H₁₃NS 155.263

4-Butyl-2-methylthiazole, 9CI [41981-69-5]

C₈H₁₃NS 155.263

2-Ethyl-4-propylthiazole, 9CI [41981-67-3]

C₈H₁₃NS 155.263

Liq. Mp 96° (as picrate). Bp₁₅ 97.5-98°

4-Ethyl-2-propylthiazole, 9CI [41981-68-4]

C₈H₁₃NS 155.263

4-Methyl-2-pentylthiazole, 9CI [96693-92-4]

C₉H₁₅NS 169.29

2,4-Dipropylthiazole, 9CI [41981-74-2]

C₉H₁₅NS 169.29

2-Isopropyl-4-propylthiazole

2-(1-Methylethyl)-4-propylthiazole, 9CI [87116-69-6]

C₉H₁₅NS 169.29

4-Ethyl-2-pentylthiazole, 9CI [96693-88-8]

C₁₀H₁₇NS 183.317

2-Butyl-4-propylthiazole, 9CI [96693-89-9]

C₁₀H₁₇NS 183.317

4-Ethyl-2-hexylthiazole, 9CI [96693-85-5]

C₁₁H₁₉NS 197.344

2-Pentyl-4-propylthiazole, 9CI [96693-86-6]

C₁₁H₁₉NS 197.344

4-Ethyl-2-heptylthiazole, 9CI [96693-84-4]

C₁₂H₂₁NS 211.371

4-Ethyl-2-octylthiazole, 9CI [96693-83-3]

C₁₃H₂₃NS 225.397

2-Octyl-4-propylthiazole, 9CI [96693-82-2]

C₁₄H₂₅NS 239.424

Beraud, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 2072-2074 (synth)

Asinger, F. *et al.*, *Annalen*, 1964, **672**, 156-178 (synth)

Vincent, E.J. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 3524-3530 (pmr)

Babadjanian, A. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 4878-4880 (synth)

Roussel, C. *et al.*, *Bull. Soc. Chim. Fr.*, 1971, 1087-1092 (synth)

Buttery, R.G. *et al.*, *J. Agric. Food Chem.*, 1973, **21**, 488-490 (synth, ms)

Vitzthum, O.G. *et al.*, *J. Food Sci.*, 1974, **39**, 1210-1215 (occur, coffee)

Haag, A. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 511-524 (ms)

Loser, B. *et al.*, *Food Technol. (Chicago)*, 1978, **32**, 60-70 (tox)

Ger. Pat., 1978, 2 804 077; CA, **76**, 163587s (use)

Ho, C.T. *et al.*, *Lebensm.-Wiss. Technol.*, 1982, **15**, 340-342 (glc, occur, tomato)

Ho, C.T. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 1384-1386 (synth, ms, occur, peanut)

Ho, C.T. *et al.*, *J. Food Sci.*, 1983, **48**, 1570-1571 (synth, props, ms)

Ho, C.T. *et al.*, *Perfum. Flavor.*, 1984, **9**, 15-18 (synth, odour)

Carlin, J.T. *et al.*, *J. Am. Oil Chem. Soc.*, 1986, **63**, 1031-1036 (occur)

Lamparsky, D. *et al.*, *Perfum. Flavor.*, 1988, **13**, 17-25 (occur, coriander)

Ames, J.M. *et al.*, *Flavour Fragrance J.*, 1992, **7**, 89-103 (occur, yeast)


ACS Symp. Ser., 1995, 147-159 (occur, yeast extract)

Weenan, H. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3291-3293 (glc, occur, durian)

Elmore, J.S. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 3595-3602 (occur)

Swaine, R.L. *et al.*, *Perfum. Flavor.*, 1997, **22**, 57-58; **22(5)**, 60; 62; 64 (occur)

1,4-Diamino-2-butanol D-248 2-Hydroxyputrescine [539-59-3]



C₄H₁₂N₂O 104.152

(R)-form

[40894-78-8 (dihydrochloride)]

Mp 249-250° (as dihydrochloride). [α]_D²⁵ -3.3 (c, 0.05 in H₂O) (dihydrochloride).

N,N'-Di-E-cinnamoyl: N¹,N⁴-Di-trans-cinnamoyl-2-hydroxyputrescine

[1003323-66-7]

C₂₂H₂₄N₂O₃ 364.443

Isol. from *Pholiota spumosa*. Cryst.

[α]_D²⁵ +3 (c, 0.5 in MeOH). λ_{max} 204 (ε 9200); 275 (ε 10600) (MeOH).

N,N'-Di-E-cinnamoyl, O-(3S-hydroxy-3-methylglutaryl): Pholiotic acid

[1003869-80-4]

C₂₈H₃₂N₂O₇ 508.57

Isol. from *Pholiota spumosa*. Waxy solid. [α]_D²⁵ +18.5 (c, 0.4 in MeOH). λ_{max} 216 (ε 6800); 274 (ε 9000) (MeOH).

(S)-form [24177-21-7]

[26097-54-1 (dihydrochloride)]

Occurs in *Pseudomonas* spp. Mp 249-250° (as dihydrochloride). [α]_D²⁵ +3.21 (c, 0.05 in H₂O) (dihydrochloride).

1-N-(4-Hydroxycinnamoyl): 1-Coumaroyl-2-hydroxyputrescine

[24177-22-8]

C₁₃H₁₈N₂O₃ 250.297

Alkaloid from *Triticum aestivum*

(wheat) (Poaceae). Geom. of double bond not defined. λ_{max} 293; 316 (H₂O) (Berdy). λ_{max} 309; 363 (NaOH) (Berdy).

(Berdy).

1-N-(4-Hydroxy-3-methoxycinnamoyl):

1-Feruloyl-2-hydroxyputrescine

[24177-48-8, 24177-49-9]

C₁₄H₂₀N₂O₄ 280.323

Alkaloid from *Triticum aestivum*

(wheat) (Poaceae). [α]_D²⁴ -20 (c, 0.7 in CHCl₃) (as tri-Ac). Mixt. of E- and Z-forms.

(±)-form [13094-22-9]

Dihydrochloride: [6210-86-2]

Mp 231°.

Dipicrate: Mp 255° dec.

Macholan, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 2074 (synth)

Stoessel, A. *et al.*, *Tet. Lett.*, 1969, **10**, 2807-2810 (1-Coumaroyl-2-hydroxyputrescine, 1-Feruloyl-2-hydroxyputrescine)

Tobari, J. *et al.*, *J. Biol. Chem.*, 1971, **246**, 1262-1265 (S-form, isol, synth, spectra)

Kullnig, R.K. *et al.*, *J. Biol. Chem.*, 1973, **248**, 2487-2488 (R-form, S-form, synth)

Tice, C.M. *et al.*, *J.O.C.*, 1983, **48**, 5043-5048 (1-Coumaroyl-2-hydroxyputrescine, synth)

Clericuzio, M. *et al.*, *Eur. J. Org. Chem.*, 2007, 5551-5559 (dicinnamate, Pholiotic acid)

1,12-Diamino-4,8-diazadodecane **D-249**

N-[3-[(3-Aminopropyl)amino]propyl]-1,4-butanediamine, 9CI. 1,5,9,14-Tetraazatetradecane. *Thermospermine*. PA334 [70862-11-2]



$\text{C}_{10}\text{H}_{26}\text{N}_4$ 202.342

Prod. by *Acidothermus* spp., *Agrobacterium* sp. and other bacteria. Also isol. from *Vicia sativa* and from the venom of the spider *Hololena curta*, and occurs in *Halocynthia roretzi* and *Anthocidaris crassispina*.

N^4 -Me: N-[3-[(3-Aminopropyl)methylamino]propyl]-1,4-butanediamine, 9CI. N^4 -Methylthermospermine [142808-35-3]

$\text{C}_{11}\text{H}_{28}\text{N}_4$ 216.369

Alkaloid from various Fabaceae. No phys. props. reported.

N^4 -Hydroxy: [864812-10-2]

$\text{C}_{10}\text{H}_{26}\text{N}_4\text{O}$ 218.342

Isol. from venom of the spider *Hololena curta*.

Ganem, B. et al., *Methods Enzymol.*, 1983, **94**, 416-418 (synth)

Samejima, K. et al., *Chem. Pharm. Bull.*, 1984, **32**, 3428-3435 (synth)

Hamana, K. et al., *FEMS Microbiol. Lett.*, 1990, **67**, 267-273; **71**, 71-76 (occur)

Hamana, K. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur)

Hamana, K. et al., *J. Biochem. (Tokyo)*, 1991, **109**, 444-449 (occur)

Hamana, K. et al., *Phytochemistry*, 1991, **30**, 3319-3322; 1992, **31**, 1410-1412 (occur, N^3 -Me)

Tzouros, M. et al., *Toxicon*, 2005, **46**, 350-354 (*Hololena curta* constits)

1,13-Diamino-4,9-diazatridecane **D-250**

1,5,10,15-Tetraazapentadecane. N^1 -(3-Aminopropyl)homospermidine

[86812-44-4]



$\text{C}_{11}\text{H}_{28}\text{N}_4$ 216.369

Constit. of the seeds of *Vicia sativa*. Also prod. by various bacteria incl. *Acidothermus* sp., *Agrobacterium* sp., *Thermoleophilum album* and *Thermoleophilum minutum*. Residue present in Glycerin A, G-131.

Samejima, K. et al., *Chem. Pharm. Bull.*, 1984, **32**, 3428-3435 (synth)

Hamana, K. et al., *Can. J. Microbiol.*, 1990, **36**, 567-572; 1991, **37**, 350-354 (synth, biosynth)

Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747 (*Thermoleophilum* constits)

1,13-Diamino-5,9-diazatridecane **D-251**

$\text{N,N}''$ -1,3-Propanediylbis[1,4-butanediamine]. 1,6,10,15-Tetraazapentadecane.

Canavalmine. PA434

[70862-15-6]



$\text{C}_{11}\text{H}_{28}\text{N}_4$ 216.369

Isol. from the seeds of *Canavalia gladiata* (sword bean). Prod. by the thermophilic

eubacteria *Bacillus schlegelii*, *Hydrogenobacter thermophilus*, *Thermoleophilum album* and *Thermoleophilum minutum*.

Viscous oil. Bp_{0.01} 135-140°.

[107886-53-3, 107886-65-7]

Fujihara, S. et al., *Biochem. Biophys. Res. Commun.*, 1982, **107**, 403 (isol, synth, ir)

Aikens, D. et al., *Biophys. Chem.*, 1983, **17**, 67 (cmr)

Samejima, K. et al., *Chem. Pharm. Bull.*, 1984, **32**, 3428 (synth)

Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747 (*thermophilic eubacteria* constit)

Schwesinger, R. et al., *Chem. Ber.*, 1994, **127**, 2435 (synth, pmr)

4,4'-Diaminodibutylamine, **D-252**
8CI

N-(4-Aminobutyl)-1,4-butanediamine, 9CI. 1,9-Diamino-5-azanonane. sym-Homospermidine. 1,6,11-Triazaundecane

[4427-76-3]



$\text{C}_8\text{H}_{21}\text{N}_3$ 159.274

Constit. of *Santalum album* (sandalwood) and present in legume root nodules. Also isol. from organs of the Japanese newt and from roots of water hyacinth *Eichhornia crassipes*. Occurs in *Anthocidaris crassispina* and *Stichopus japonicus*. Prod. by the thermophilic eubacteria, *Hydrogenobacter thermophilus*. Oil.

Hydrochloride (1:2): Mp 283-285° dec.

Hydrochloride (1:3): [189340-78-1]

Cryst. (EtOH aq.).

N-Tetra-Me: see Solamine, S-340

[138656-54-9]

Kuttan, R. et al., *Biochemistry*, 1971, **10**, 361-365 (isol, synth)

Yamamoto, S. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3315-3318 (isol)

Hamana, K. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur)

Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747 (*thermophilic eubacteria* constits)

Ramaswamy, S. et al., *CA*, 1992, **116**, 55605v (cryst struct)

Bergeron, R.J. et al., *J. Med. Chem.*, 1997, **40**, 1475-1494 (synth, pmr)


Parker, L.L. et al., *Tetrahedron*, 2003, **59**, 10165-10171 (synth, pmr, cmr)

2,4-Diamino-5,6-dihydroxypyrimidine **D-253**

2,6-Diamino-5-hydroxy-4(1H)-pyrimidinone, 9CI. 2,4-Diamino-5,6-pyrimidinediol. Divicine

[60337-65-7]

[60337-65-7]



$\text{C}_4\text{H}_6\text{N}_4\text{O}_2$ 142.117

Pale yellow microcryst. Mp 300° dec.

5-O-Arabinyopyranoside: *Charine*

[165171-52-8]

$\text{C}_9\text{H}_{14}\text{N}_4\text{O}_6$ 274.233

Alkaloid from the unripe fruit of *Momordica charantia* (bitter melon). Needles (MeOH). Mp 180-182°. λ_{max} 239 (no solvent reported).

5-O- β -D-Glucopyranoside: *Vicine*†. *Vicio-side*

[152-93-2]

$\text{C}_{10}\text{H}_{16}\text{N}_4\text{O}_7$ 304.259

Isol. from vetch (*Vicia sativa*), *Vicia faba* and other legumes. Mp 243-244° dec. $[\alpha]_{\text{D}}^{25}$ -11.7 (0.2M NaOH).

Bendich, A. et al., *Biochim. Biophys. Acta*, 1953, **12**, 462 (isol, struct)

Davoll, J. et al., *J.C.S.*, 1956, 2124 (synth)

Chesterfield, J.H. et al., *J.C.S.*, 1964, 1001 (synth)

Brown, E.G. et al., *Phytochemistry*, 1972, **11**, 3203

Dutta, P.K. et al., *Indian J. Chem., Sect. B*, 1981, **20**, 669 (isol, pmr, cmr)


Kunesh, N. et al., *Annalen*, 1994, 1059 (synth, *Vicine*)

El-Gengaihi, S. et al., *Pharmazie*, 1995, **50**, 361-362 (*Charine*)

4,5-Diamino-2,6-dihydroxypyrimidine **D-254**

5,6-Diamino-2,4(1H,3H)-pyrimidinedione, 9CI. 5,6-Diamino-2,4-pyrimidinediol. 5,6-Diaminouracil

[3240-72-0]



$\text{C}_4\text{H}_6\text{N}_4\text{O}_2$ 142.117

Two intermediate mono-NH tautomers also possible. Prod. by a riboflavin-free mutant of *Aspergillus nidulans*. Cryst. (as dihydrochloride).

1-Ph:

$\text{C}_{10}\text{H}_{10}\text{N}_4\text{O}_2$ 218.215

Yellow-brownish cryst. (Py aq.). Mp 232-233°.

Di-NH-form

1,3-Di-Me: [5440-00-6]

$\text{C}_6\text{H}_{10}\text{N}_4\text{O}_2$ 170.171

Used as 0.1M aq. soln. for photo-metric detn. of Co. Long needles (H_2O). Hydrol on standing; turns yellow. Mp 209° Mp 275° dec.

[32014-70-3, 50996-16-2, 50787-05-8]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 812D (ir)

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 704D (nmr)

Org. Synth., Coll. Vol., **4**, 1963, 247

Sadique, J. et al., *Naturwissenschaften*, 1966, **53**, 282 (isol)

Ruzička, E. et al., *Mikrochim. Acta*, 1968, 938 (detn, Co)

Khmelnitskii, R.A. et al., *CA*, 1974, **81**, 62685h (ms, struct)

3,3'-Diaminodipropylamine, **D-255**
8CI

N-(3-Aminopropyl)-1,3-propanediamine, 9CI. Bis(3-aminopropyl)amine.

Dipropylentriamine. 1,7-Diamino-4-azaheptane. Iminobispropylamine. 1,5,9-Triazanonane. Norspermidine. *Caldine* [56-18-8]



$\text{C}_6\text{H}_{17}\text{N}_3$ 131.22

Metab. of various plant spp. and bacteria, e.g. the thermophilic eubacteria *Thermoleophilum album*, *Thermoleophilum minutum*, *Hydrogenobacter thermophilus* and *Bacillus schlegelii*. Isol. from the green alga *Codium fragile*. Occurs in *Anthocidaris crassispina*, *Stichopus japonicus*, *Crassostrea gigas*, *Anthocynthia roretzi* and *Tapes philippinarum*. Colorimetric reagent for nitro compds. Cross-linking agent for epoxy resins. Liq. Sol. H₂O, polar org. solvs. d_{20}^{20} 0.93. Mp -14°. Bp 240° Bp₂ 100°. n_D^{20} 1.4810.

► Skin and severe eye irritant. LD₅₀ (rat, orl) 810 mg/kg. LD₅₀ (skn, rbt) 110 mg/kg. JL9450000

Hydrochloride (1:3):

Cryst. (MeOH aq. or EtOH/Et₂O). Mp 270° (259°).

Tripicrate:

Cryst. (H₂O). Mp 226-227°.

N³,N^{3'}-Bis(2,3-dihydroxybenzoyl):

[88381-35-5]

C₂₀H₂₅N₃O₆ 403.434

Isol. from *Vibrio fluvialis*. Siderophore. λ_{\max} 248 (E1%/1cm 12.8); 316 (E1%/1cm 8.1) (MeOH) (Berdy).

N¹-Me: N-(3-Aminopropyl)-N-methyl-1,3-propanediamine. 3,3'-Diamino-N-methylpropylamine. N,N-Bis(3-aminopropyl)methylamine

[105-83-9]

C₇H₁₉N₃ 145.247

Monomer for polyamides and polyureas. Liq. Bp₁₂ 112-114°.

► Skin and severe eye irritant. LD₅₀ (rat, orl) 1540 mg/kg. JL9625000

N¹,N³,N^{3'}-Tri-Me: N,N'-Dimethyl-N-[3-(methylamino)propyl]-1,3-propanediamine. N-Methyl-3,3'-bis(methylamino)propylamine

[123-70-6]

C₉H₂₃N₃ 173.301

No phys. props. reported.

N³,N³,N^{3'},N^{3'}-Tetra-Me: N'-[3-(Dimethylamino)propyl]-N,N'-trimethyl-1,3-propanediamine. 9CI. 3,3'-Bis(dimethylamino)-N-methyldipropylamine. 8CI.

[6711-48-4]

C₁₀H₂₅N₃ 187.328

Mp -78°. Bp₂₀ 128-131°. n_D^{20} 1.4490.

► JM1925000

N-Penta-Me: N-[3-(Dimethylamino)propyl]-N,N',N'-trimethyl-1,3-propanediamine. 9CI. 3,3'-Bis(dimethylamino)-N-methyldipropylamine. 8CI.

[3855-32-1]

C₁₁H₂₇N₃ 201.354

Catalyst for polyurethane foam production; auxiliary for lithiation reactions. Liq. Bp₁₁ 100-102°.

[27708-70-9, 72864-15-4]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 309C; 310B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 500B; 500C; 501A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 397A; 397B (ir)

Wiedeman, D.F. et al., *J.A.C.S.*, 1945, 67, 1994-1996 (synth)

Terent'ev, A.P. et al., *Zh. Obshch. Khim.*, 1950, 20, 1073-1078; *CA*, 44, 9350a (synth)

Marxer, A. et al., *Helv. Chim. Acta*, 1951, 34, 924-931 (N-penta-Me)

Israel, M. et al., *J. Med. Chem.*, 1964, 7, 710-716 (synth)

Schrier, M. et al., *Mikrochim. Acta*, 1965, 53, 1091-1097; 1967, 55, 218-222 (use)

Ger. Pat., 1978, 2 739 353; *CA*, 89, 111389r (N-penta-Me)

Bergeron, R.J. et al., *Synthesis*, 1981, 732-733 (synth)

Canadian Pat., 1982, 1 132 292; *CA*, 98, 55043v (N-penta-Me)

Murahashi, S. et al., *J.A.C.S.*, 1983, 105, 5002-5011 (synth, pmr, ir)

Dagnall, S.P. et al., *J.C.S. Perkin 2*, 1984, 435-440 (cmr)

Hamana, K. et al., *J. Biochem. (Tokyo)*, 1985, 97, 1595-1601; 1991, 109, 444-449 (occur)

Tanaka, Y. et al., *CA*, 1988, 109, 232044q (crosslinking)

Carboni, B. et al., *Tet. Lett.*, 1988, 29, 1279-1282 (synth)

Hamana, K. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, 100, 59-62 (occur, hplc, gc)

Hamana, K. et al., *Biochem. J.*, 1992, 284, 741-747 (thermophilic eubacteria constits)


Yamamoto, S. et al., *J. Biochem. (Tokyo)*, 1993, 113, 538-544 (bisdihydroxybenzoyl)

Thurner, A. et al., *Synth. Commun.*, 1998, 28, 443-449 (N-penta-Me, use)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AIX250; TDP750; BGU750

2,4-Diamino-3-hydroxybenzoic acid

D-256



C₇H₈N₂O₃ 168.152

N²,N⁴-Di-Me, Me ester: Damascinine. Methyl 3-hydroxy-2,4-bis(methylamino)benzoate. 9CI

[28917-01-3]


C₁₀H₁₄N₂O₃ 210.232

Alkaloid from the seeds of *Nigella damascena* (Ranunculaceae). Mp 75-79°.

Doepke, W. et al., *Pharmazie*, 1970, 25, 69 (ir, pmr, ms)

3,20-Diamino-18-hydroxy-pregna-5-ene-11,16-dione

D-257



C₂₁H₃₂N₂O₃ 360.495

Tentative struct.

(3β,20S)-form

Holarricine. 11,16-Dioxoholarrhimine

[84294-76-8]

Alkaloid from the seeds of *Holarrhena*

antidysenterica (Apocynaceae). Mp 350-351°. $[\alpha]_D^{27}$ -31 (EtOH).


N-Tetra-Me: Mp 192-193°.

N-Tetra-Me, O-Ac: Mp 265-266°.

Siddiqui, S. et al., *Pak. J. Sci. Ind. Res.*, 1981, 24, 167

3,20-Diaminopregna-4,14-dien-2-ol

D-258



C₂₁H₃₄N₂O 330.512

(2β,3β,20S)-form

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl): Salignamide B

[208835-29-4]

C₂₈H₄₄N₂O₂ 440.668

Alkaloid from *Sarcococca saligna*.

Gum. $[\alpha]_D^{30}$ +48 (c, 0.04 in MeOH).


λ_{\max} 215 (log ε 3.65); 224 (log ε 3.67);

228 (log ε 3.66) (MeOH).

Atta-ur-Rahman, et al., *Nat. Prod. Lett.*, 1998, 11, 297-304 (isol, uv, pmr, cmr, ms)

3,20-Diaminopregna-4,16-dien-2-ol

D-259



C₂₁H₃₄N₂O 330.512

(2α,3β,20S)-form

N²⁰,N²⁰-Di-Me, N³-tigloyl: Salignamide D

[426814-98-4]

C₂₈H₄₄N₂O₂ 440.668


Alkaloid from *Sarcococca saligna*.

Amorph. solid. $[\alpha]_D^{25}$ +8 (c, 0.3 in CHCl₃). λ_{\max} 231 (log ε 3.63); 254 (log ε 3.66) (MeOH).

Atta-ur-Rahman, et al., *Helv. Chim. Acta*, 2002, 85, 678-688 (isol, ir, pmr, cmr)

3,20-Diaminopregna-2,16-dien-4-one

D-260




C₂₁H₃₂N₂O 328.497**(5 α ,20S)-form**N²⁰, N²⁰-di-Me, N³-tigloyl: 16-Dehydro-sarcovagine D. **Sarcovagine C**C₂₈H₄₂N₂O₂ 438.652Alkaloid from the roots of *Sarcococca vagans* (Buxaceae). Cryst. Mp 160-161°. [α]_D²⁵ +5.4 (c, 0.06 in CHCl₃).16 α ,17 α -Epoxide, N²⁰-Me, N³-tigloyl: **16,17-Epoxysarcovagine D**

[497164-36-0]

C₂₇H₄₀N₂O₃ 440.625Alkaloid from the leaves of *Sarcococca coriacea*. Amorph. yellow solid. [α]_D²⁵ +24 (c, 0.12 in CHCl₃). λ_{\max} 223 (log ϵ 5.2) (MeOH).Zou, Z.-M. et al., *Phytochemistry*, 1997, **46**, 1091-1093 (*Sarcovagine C*)Kalauni, S.K. et al., *Chem. Pharm. Bull.*, 2002, **50**, 1423-1426 (*16,17-Epoxysarcovagine D*)**3,20-Diaminopregna-5,14-dien-4-one**


D-261

C₂₁H₃₂N₂O 328.497**(3 β ,20S)-form**N²⁰, N²⁰-Di-Me, N³-angeloyl: **Sarsaligne-none**

[199922-56-0]

C₂₈H₄₂N₂O₂ 438.652Alkaloid from *Sarcococca saligna* (Buxaceae). Powder. Mp 20°. [α]_D²⁵ -113.6 (c, 0.1 in MeOH). Doubtful mp. given in ref. λ_{\max} 237 (log ϵ 2.9); 257 (log ϵ 2.9) (MeOH).Atta-ur-Rahman, et al., *Phytochemistry*, 1997, **46**, 771-775 (*isol, uv, ir, pmr, cmr, ms*)**3,20-Diaminopregnane-2,4-diol**

D-262

C₂₁H₃₈N₂O₂ 350.543

CA gives incorrect names for Axillarines C-F.

(2 α ,3 β ,4 β ,5 α ,20S)-formN²⁰, N²⁰-Di-Me, N³-tigloyl, 4-Ac: **Sarcovagine B**

[192385-46-9]

C₃₀H₅₀N₂O₄ 502.736Alkaloid from *Sarcococca vagans*. Cryst. + 1/3 H₂O. Mp 205-206°. [α]_D²⁵+19.6 (c, 0.06 in CHCl₃).N²⁰, N²⁰-Di-Me, N³-(3-methyl-2-butenoyl), 4-Ac: **Hookerianamide B**

[771533-16-5]

C₃₀H₅₀N₂O₄ 502.736Alkaloid from *Sarcococca hookeriana*.Cryst. Mp 224-226°. [α]_D²⁴ +19.2 (c, 0.05 in MeOH). λ_{\max} 210 (log ϵ 4.8); 221 (log ϵ 4.4) (MeOH).N²⁰, N²⁰-Di-Me, N³-phthalimide, 4-Ac: 4-Acetoxy-20-(dimethylamino)-3-phthalimidopregnan-2-ol. 20-(Dimethylamino)-2-hydroxy-3-phthalimidopregnan-4-yl acetate

[214402-22-9]

C₃₃H₄₆N₂O₅ 550.737Alkaloid from *Pachysandra procumbens*. Needles. Mp 269°. [α]_D²⁰ +48 (c, 0.03 in CHCl₃). λ_{\max} 228 (log ϵ 4.19); 243 (log ϵ 3.94); 293 (log ϵ 3.33) (MeOH).**(2 β ,3 β ,4 β ,5 α ,20S)-form**N²⁰, N²⁰-Di-Me, N³-tigloyl: **Sarcovagine A**

[192385-45-8]

C₂₈H₄₈N₂O₃ 460.699Alkaloid from *Sarcococca vagans*.Cryst. + 1/2 H₂O. Mp 277-279°. [α]_D²⁵ +21.2 (c, 0.11 in CHCl₃).N²⁰, N²⁰-Di-Me, N³-tigloyl, 4-Ac: **Axillarine F**

[128255-12-9]

C₃₀H₅₀N₂O₄ 502.736Alkaloid from *Pachysandra axillaris*(Buxaceae). Cryst. Mp 241-244°. [α]_D²² +29.5 (c, 0.398 in CHCl₃).N²⁰, N²⁰-Di-Me, N³-benzoyl: **Axillarine E**

[128255-09-4]

C₃₀H₄₆N₂O₃ 482.705Alkaloid from *Pachysandra axillaris*

(Buxaceae). Cryst. Mp 285-290°.

N²⁰, N²⁰-Di-Me, N³-benzoyl, 4-Ac: **Axillarine C**

[128255-10-7]

C₃₂H₄₈N₂O₄ 524.742Alkaloid from *Pachysandra axillaris*(Buxaceae). Cryst. Mp 272-274°. [α]_D²² +22.4 (c, 0.981 in CHCl₃).N²⁰, N²⁰-Di-Me, N³-benzoyl, 2,4-di-Ac: **Axillarine D**, 2,4-Diacetoxypachysamine D


[128255-11-8]

C₃₄H₅₀N₂O₅ 566.779Alkaloid from *Pachysandra axillaris*(Buxaceae) and *Sarcococca saligne*.Cryst. Mp 223-225°. [α]_D²² +11.6 (c, 0.433 in CHCl₃). [α]_D +83 (c, 0.05 in MeOH).5 α ,6 α -Epoxide, N²⁰, N²⁰-di-Me, N³-tigloyl, 4-Ac: **Salignarine A**

[299203-53-5]

C₃₀H₄₈N₂O₅ 516.72Alkaloid from *Sarcococca saligna*.Brown gum. [α]_D²⁰ -60 (c, 0.1 in CHCl₃). λ_{\max} 208 (log ϵ 2.4) (MeOH).Chiu, M. et al., *J. Nat. Prod.*, 1992, **55**, 25-28 (*Axillarines C-F, struct*)Yu, S.S. et al., *Chin. Chem. Lett.*, 1997, **8**, 511-514 (*Sarcovagines*)Cheng, L.C. et al., *J. Nat. Prod.*, 1998, **61**, 1257-1262 (*phthalimido deriv*)Atta-ur-Rahman, et al., *Nat. Prod. Lett.*, 1998,**11**, 297-304 (*Diacetoxypachysamine D*)Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000,**63**, 1364-1368 (*Salignarine A*)Choudhary, M.I. et al., *Helv. Chim. Acta*,2004, **87**, 1099-1108 (*Hookerianamide B*)**3,20-Diaminopregnane-4,12-diol**

D-263

C₂₁H₃₈N₂O₂ 350.543**(3 α ,4 β ,5 α ,12 β ,20S)-form**N³, N²⁰, N²⁰-Tri-Me, N³-benzoyl, 4-Ac: **Paxillarine B**, 12 β -Hydroxypachysandrine A


[145022-90-8]

C₃₃H₅₀N₂O₄ 538.769Alkaloid from whole plants of *Pachysandra axillaris*. Prisms (Me₂CO/CH₂Cl₂). Mp 234-237°. [α]_D²⁵ +65.2 (c, 0.357 in CHCl₃).**(3 β ,4 β ,5 α ,12 β ,20S)-form**N³, N²⁰, N²⁰-Tri-Me, N³-benzoyl, 4-Ac: **Axillarine B**

[128254-96-6]

C₃₃H₅₀N₂O₄ 538.769Alkaloid from *Pachysandra axillaris*.Qiu, M. et al., *CA*, 1990, **113**, 55838m(*Axillarine B*)Qiu, M. et al., *Chem. Pharm. Bull.*, 1996, **44**, 2015-2019 (*Paxillarine B*)**3,20-Diaminopregnane-4,16-diol**

D-264

C₂₁H₃₈N₂O₂ 350.543**(3 α ,4 β ,5 α ,16 β ,20S)-form**N³, N²⁰, N²⁰-Tri-Me, N³-benzoyl, 4-Ac: **Paxillarine A**, 16 β -Hydroxypachysandrine A


[145022-89-5]

C₃₃H₅₀N₂O₄ 538.769Alkaloid from whole plants of *Pachysandra axillaris*. Prisms (Me₂CO). Mp263-265°. [α]_D²⁵ +77.7 (c, 0.515 inCHCl₃).**(3 β ,4 β ,5 α ,16 β ,20S)-form**N³, N²⁰, N²⁰-Tri-Me, N³-benzoyl, 4-Ac: **Axillarine A**

[128269-64-7]

C₃₃H₅₀N₂O₄ 538.769Alkaloid from *Pachysandra axillaris*.Qiu, M. et al., *CA*, 1990, **113**, 55838m


(Axillarine A)

Qiu, M. et al., *Chem. Pharm. Bull.*, 1996, **44**, 2015-2019 (Paxillarine A)**3,20-Diaminopregnan-2-ol** D-265C₂₁H₃₈N₂O 334.544**(2β,3β,5α,20S)-form**

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl), 2-Ac: **Hookerianamide C** [771533-17-6]
C₃₀H₅₀N₂O₃ 486.737
Alkaloid from *Sarcococca hookeriana*. Cholinesterase inhibitor. Cryst. Mp 240-242°. [α]_D²⁴ -136 (c, 0.1 in MeOH). λ_{max} 220 (log ε 3.6) (MeOH).

N²⁰,N²⁰-Di-Me, N³-benzoyl: **2-Hydroxyepipachysamine D** [426814-99-5]
C₃₀H₄₆N₂O₂ 466.706
Alkaloid from *Sarcococca saligna*. Amorph. solid. [α]_D²⁵ +52 (c, 0.21 in CHCl₃). λ_{max} 221 (log ε 2.7); 269 (log ε 3.78) (MeOH).

Atta-ur-Rahman, et al., *Helv. Chim. Acta*, 2002, **85**, 678-688 (2-Hydroxyepipachysamine D)
Choudhary, M.I. et al., *Helv. Chim. Acta*, 2004, **87**, 1099-1108 (Hookerianamide C)

3,20-Diaminopregnan-4-ol D-266
3,20-Diamino-4-hydroxypregnananeC₂₁H₃₈N₂O 334.544**(3α,4α,5α,20S)-form**

N³,N²⁰,N²⁰-Tri-Me: 20-(Dimethylamino)-3-(methylamino)pregnan-4-ol, 9Cl. **Pachysandrine C** [6801-30-5]
C₂₄H₄₄N₂O 376.624
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 214-215°. [α]_D -38 (CHCl₃).

N³,N²⁰,N²⁰-Tri-Me, O-(3-methyl-2-butenoyl): **Pachysandrine D** [6911-18-8]
C₂₉H₅₀N₂O₂ 458.726
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 184-185°. [α]_D +2 (CHCl₃).

(3α,4β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-benzoyl: **Epipachysandrine A** [5874-19-1]
C₃₀H₄₆N₂O₂ 466.706
Minor alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO/CH₂Cl₂). Mp 290-292°. [α]_D +12 (MeOH/CHCl₃ 1:1). Not epimeric with Pachysandrine A.

N³,N²⁰,N²⁰-Tri-Me, O⁴-benzoyl: **Pachysanaximine A** [128255-15-2]
C₃₁H₄₈N₂O₂ 480.732
Alkaloid from *Pachysandra axillaris* (Buxaceae). Mp 195-196°. [α]_D²² +12.4 (c, 0.28 in CHCl₃).

N³,N²⁰,N²⁰-Tri-Me, N³-benzoyl, O-Ac: **Pachysandrine A** [6879-28-3]
C₃₃H₅₀N₂O₃ 522.77
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 235-236°. [α]_D¹³ +80 (CHCl₃).

▶CU8702340

N³,N²⁰,N²⁰-Tri-Me, N³-(3-methyl-2-butenoyl): **O-Desacetyl pachysandrine B**. O-Deacetyl pachysandrine B [15027-88-0]
C₂₉H₅₀N₂O₂ 458.726
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 184-185° Mp 201-202°.

N³,N²⁰,N²⁰-Tri-Me, N³-(3-methyl-2-butenoyl), O-Ac: **Pachysandrine B** [6879-29-4]
C₃₁H₅₂N₂O₃ 500.763
Alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO/CH₂Cl₂). Mp 187-189°. [α]_D¹⁹ +93.4 (CHCl₃).


N³,N³,N²⁰,N²⁰-Tetra-Me: 3,20-Bis(dimethylamino)pregnan-4-ol. **O,N-Desacyl-N-methyl pachysandrine A** [15027-71-1]
C₂₅H₄₆N₂O 390.651
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 126-150°.

(3β,4β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl), O-Ac: **Vaganine A** [144028-94-4]
C₃₀H₅₀N₂O₃ 486.737
Alkaloid from *Sarcococca vagans* (Buxaceae). [α]_D²⁵ +119 (c, 0.2 in MeOH). λ_{max} 204 (log ε 3.8) (MeOH).
N²⁰,N²⁰-Di-Me, N³-tigloyl, O-Ac: **Sarcovagine C** [192385-48-1]
C₃₀H₅₀N₂O₃ 486.737
Alkaloid from *Sarcococca vagans*. Cryst. + ½ H₂O. Mp 192-194°. [α]_D¹³ -9 (c, 0.12 in CHCl₃).


Kikuchi, T. et al., *Tet. Lett.*, 1964, **5**, 1817-1823; 1969, **10**, 1679-1682 (*Pachysandra terminalis* constits, synth)
Tomita, M. et al., *Chem. Pharm. Bull.*, 1967, **15**, 193-207; 207-213; 577-581 (*Pachysandrines A-D*, Epipachysandrine A)
Shojiro, N. et al., *Yakugaku Zasshi*, 1967, **87**, 215-227; *CA*, **67**, 32888 (*Pachysandra terminalis* constits)

Kikuchi, T. et al., *Chem. Pharm. Bull.*, 1971, **19**, 1886-1892 (*Pachysandrines*, Epipachysandrine A, synth)
Qiu, M. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1990, **32**, 626-630; *CA*, **114**, 118538g (*Pachysanaximine A*)
Qiu, M. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1993, **35**, 885; *CA*, **121**, 78257a (*Vaganine A*)
Yu, S.S. et al., *Chin. Chem. Lett.*, 1997, **8**, 511-514 (*Sarcovagine C*)
Atta-ur-Rahman, et al., *Phytochemistry*, 1997, **46**, 771-775 (*Vaganine A*)

3,20-Diaminopregnan-11-ol D-267C₂₁H₃₈N₂O 334.544**(3β,11ξ,20S)-form**

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl): **11-Hydroxyepipachysamine E** [312962-53-1]
C₂₈H₄₈N₂O₂ 444.699
Alkaloid from *Sarcococca brevifolia*. Amorph. solid. [α]_D -15 (c, 0.25 in CHCl₃). Sign of opt. rotation unclear in ref. λ_{max} 199 (EtOH).

Jayasinghe, U.L.B. et al., *Nat. Prod. Lett.*, 2000, **14**, 293-298

3,20-Diaminopregnan-12-ol D-268C₂₁H₃₈N₂O 334.544**(3α,5α,12β,20S)-form**


N³,N²⁰,N²⁰-Tri-Me, N³-(3-methyl-2-butenoyl): [282726-64-1]
C₂₉H₅₀N₂O₂ 458.726
Alkaloid from *Pachysandra procumbens*. Needles. Mp 179°. [α]_D²⁰ +61.4 (c, 0.04 in CHCl₃). λ_{max} 214 (log ε 4.35) (MeOH).

N³,N²⁰,N²⁰-Tri-Me, N³-benzoyl, 12-Ac: Alkaloid from *Pachysandra procumbens*. Needles. Mp 190°. [α]_D²⁰ +35 (c, 0.16 in CHCl₃). No CAS no. found to 15Cl. λ_{max} 218 (log ε 4.26); 242 (log ε 3.76) (MeOH).

Chang, L.C. et al., *Tetrahedron*, 2000, **56**, 3133-3138 (*N*-3-methyl-2-butenoyl, *N*-benzoyl, isol, uv, ir, pmr, cmr, ms)

3,20-Diaminopregnan-18-ol

D-269

C₂₁H₃₆N₂O₂ 348.528C₂₁H₃₈N₂O 334.544

(3β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-Ac: 3-Acetamido-20-(dimethylamino)pregnan-18-ol. **Malouphyllinine**. *Malouphyllinol*

[7050-52-4]

C₂₅H₄₄N₂O₂ 404.635

Alkaloid from *Malouetia bequaertiana* (Apocynaceae). Mp 257°. [α]_D -11 (CHCl₃).

18-Aldehyde: 3,20-Diaminopregnan-18-al

18-Aldehyde, N²⁰,N²⁰-di-Me, N³-Ac: 3-Acetamido-20-(dimethylamino)pregnan-18-al. **Malouphylline**

[7050-51-3]

C₂₅H₄₂N₂O₂ 402.619


Alkaloid from *Malouetia bequaertiana* leaves (Apocynaceae). Cryst. (Me₂CO). Mp 261°. [α]_D +32 (CHCl₃).

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 648-651 (*Malouphylline*)

Khuong-Huu-Lainé, F. *et al.*, *Ann. Pharm. Fr.*, 1965, 395-409 (*Malouphyllinine*)

3,20-Diaminopregnan-5-ene-15,18-diol

D-270

C₂₁H₃₆N₂O₂ 348.528

(3β,15β,20S)-form

N³,N²⁰,N²⁰-Tri-Me, 18-O-(3,4-dimethyl-2-pentenoyl): 20-(Dimethylamino)-18-ikemaoyl-3-(methylamino)pregn-5-ene-15,18-diol. 3-(Methylamino)-15,18-dihydroxy-18-ikemaoyl-20-(dimethylamino)pregn-5-ene

[91147-26-1]


C₃₁H₅₂N₂O₃ 500.763

Alkaloid from the stem bark of *Didymeleles madagascariensis* (Didymelaceae). [α]_D +4 (c, 1.5 in CHCl₃).

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1984, 71-76 (*isol, pmr, ms*)

3,20-Diaminopregnan-5-ene-16,18-diol

D-271



(3β,16α,20S)-form

N³,N²⁰,N²⁰-Tri-Me: 20-(Dimethylamino)-3-(methylamino)pregn-5-ene-16,18-diol

[113857-62-8]


C₂₄H₄₂N₂O₂ 390.608

Alkaloid from the root bark of *Didymeleles perrieri* (Didymelaceae). [α]_D -2 (c, 0.68 in EtOH).

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1987, 877 (*isol, ir, pmr, cmr, ms, struct*)

3,20-Diaminopregnan-14-ene-2,4-diol

D-272

C₂₁H₃₆N₂O₂ 348.528

(2β,3β,4β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-tigloyl: **Salonine A**

C₂₈H₄₆N₂O₃ 458.683

Alkaloid from *Sarcococca saligna*. Cholinesterase inhibitor. Yellow gum. [α]_D²⁰ +60 (c, 0.03 in MeOH). λ_{max} 212 (log ε 2.66) (MeOH).

N²⁰,N²⁰-Di-Me, N³-tigloyl, 4-Ac: **Saligenamide C**

[426814-97-3]

C₃₀H₄₈N₂O₄ 500.72


Alkaloid from *Sarcococca saligna*. Pale yellow gum. [α]_D²⁵ +90 (c, 0.1 in CHCl₃). λ_{max} 209 (log ε 3.1) (MeOH).

Atta-ur-Rahman, *et al.*, *Helv. Chim. Acta.*, 2002, **85**, 678-688 (*Saligenamide C*)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Res.*, 2003, **17**, 235-241 (*Salonine A*)

3,20-Diaminopregnan-16-ene-2,4-diol

D-273

C₂₁H₃₆N₂O₂ 348.528

(2α,3β,4β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-tigloyl, 4-Ac: **Sarcovagennine B**

[200868-70-8]

C₃₀H₄₈N₂O₄ 500.72

Alkaloid from the roots of *Sarcococca vagans* (Buxaceae). Cryst. Mp 283-284°. [α]_D²⁵ +5.9 (c, 0.1 in CHCl₃).

(2β,3β,4β,5α,20S)-form

N²⁰-Me, N³-tigloyl, 2,4-di-Ac: **Nepapakistamine A**

[353228-47-4]

C₃₁H₄₈N₂O₅ 528.731

Alkaloid from *Sarcococca coriacea*. Cryst. (CHCl₃). Mp 162-163°. [α]_D²³ +18.7 (c, 0.09 in CHCl₃). λ_{max} 203 (log ε 4.8) (MeOH).

N²⁰,N²⁰-Di-Me, N³-tigloyl: **Sarcovagennine A**

[200868-69-5]

C₂₈H₄₆N₂O₃ 458.683

Alkaloid from the roots of *Sarcococca vagans* (Buxaceae). Cryst. Mp 260-261°. [α]_D²⁵ +9.4 (c, 0.05 in CHCl₃).

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl): **Hookerianamide A**

[771533-15-4]

C₂₈H₄₆N₂O₃ 458.683

Alkaloid from *Sarcococca hookeriana*. Cholinesterase inhibitor. Cryst. Mp 212-214°. [α]_D²⁴ -53 (c, 0.06 in MeOH).

16α,17α-Epoxyde, N²⁰-Me, N³-tigloyl, 2,4-di-Ac: **16,17-Epoxynepapakistamine A**

[497162-05-7]

C₃₁H₄₈N₂O₆ 544.73

Alkaloid from the leaves of *Sarcococca coriacea*. Cryst. (CHCl₃). Mp 119-120°. [α]_D²⁵ +14 (c, 0.07 in CHCl₃). λ_{max} 209 (log ε 5.3) (MeOH).

Zou, Z.-M. *et al.*, *Phytochemistry*, 1997, **46**, 1091-1093 (*Sarcovagennines*)


Kalauni, S.K. *et al.*, *J. Nat. Prod.*, 2001, **64**, 842-844 (*Nepapakistamine A*)

Kalauni, S.K. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1423-1426 (*16,17-Epoxynepapakistamine A*)

Choudhary, M.I. *et al.*, *Helv. Chim. Acta.*, 2004, **87**, 1099-1108 (*Hookerianamide A*)

3,20-Diaminopregnan-2-en-4-ol

D-274

C₂₁H₃₆N₂O 332.528

(4β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl): **Vagennine B**

[156430-95-4]

C₂₈H₄₆N₂O₂ 442.684

Alkaloid from *Sarcococca vagans* (Buxaceae).

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl), 4-Ac: **Vagennine C**

[156430-96-5]

C₃₀H₄₈N₂O₃ 484.721

From *Sarcococca vagans* (Buxaceae).

N²⁰,N²⁰-Di-Me, N³-benzoyl:

C₃₀H₄₄N₂O₂ 464.69


Alkaloid from *Pachysandra procumbens*. Needles. Mp 189°. [α]_D²⁰ +40 (c, 0.06 in CHCl₃). λ_{max} 220 (log ε 3.66); 269 (log ε 3.43) (MeOH).

N^{20}, N^{20} -Di-Me, N^3 -benzoyl, 4-Ac:
 $C_{32}H_{46}N_2O_3$ 506.727
 Alkaloid from *Pachysandra procumbens*. Needles. Mp 179°. $[\alpha]_D^{20}$ +48.2 (c, 0.04 in $CHCl_3$). λ_{max} 224 (log ϵ 3.94); 264 (log ϵ 3.73) (MeOH).

4-Ketone, N^{20}, N^{20} -di-Me, N^3 -tigloyl:
Sarcovagine D
 [192385-49-2]
 $C_{28}H_{44}N_2O_2$ 440.668
 Alkaloid from *Sarcococca vagans*. Mp 170-172°. $[\alpha]_D^{25}$ +39.5 (c, 0.07 in $CHCl_3$).

4-Ketone, N^{20}, N^{20} -di-Me, N^3 -benzoyl:
Axillaridine A
 [128255-16-3]
 $C_{30}H_{42}N_2O_2$ 462.674
 Alkaloid from whole plants of *Pachysandra axillaris* (Buxaceae). Mp 223-224°. $[\alpha]_D^{25}$ +51.3 (c, 0.517 in $CHCl_3$).
 Chiu, M.H. et al., *Phytochemistry*, 1992, **31**, 2571-2572 (*Axillaridine A*)
 Qiu, M. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1993, **35**, 885; *CA*, **121**, 78257a (*Vaganine B*)
 Yu, S.S. et al., *Chin. Chem. Lett.*, 1997, **8**, 511-514 (*Sarcovagine D*)
 Chang, L.C. et al., *J. Nat. Prod.*, 1998, **61**, 1257-1262 (*Pachysandra procumbens deriv*)
 Chang, L.C. et al., *Tetrahedron*, 2000, **56**, 3133-3138 (*20-N-di-Me-3-N-benzoyl*)

3,20-Diaminopregn-4-en-2-ol D-275




$C_{21}H_{36}N_2O$ 332.528

(2 β ,3 β ,20S)-form

N^{20}, N^{20} -Di-Me, N^3 -tigloyl: **2-Hydroxy-salignarine E**. Warsine A
 [657411-70-6]
 $C_{28}H_{46}N_2O_2$ 442.684
 Alkaloid from *Sarcococca saligna*.
 Cholinesterase inhibitor. Yellow gum.
 $[\alpha]_D^{20}$ +38.9 (c, 0.3 in MeOH). λ_{max} 211 (log ϵ 1.32); 229 (log ϵ 2.76) (MeOH).

Zaheer-ul-Haq, et al., *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4375-4380 (*isol*)
 Atta-ur-Rahman, et al., *Helv. Chim. Acta*, 2004, **87**, 439-448 (*isol, pmr, cmr, ms*)

3,20-Diaminopregn-5-en-2-ol D-276



$C_{21}H_{36}N_2O$ 332.528


(2 β ,3 β ,20S)-form

N^{20}, N^{20} -Di-Me, N^3 -tigloyl: **Salignarine B**

[299203-54-6]
 $C_{28}H_{46}N_2O_2$ 442.684
 Alkaloid from *Sarcococca saligna*.
 Amorph. powder. $[\alpha]_D^{20}$ +94 (c, 0.2 in $CHCl_3$). λ_{max} 212 (log ϵ 1.8) (MeOH).

N^{20}, N^{20} -Di-Me, N^3 -(3-methyl-2-buteno-nyl): **Salignarine C**
 [299203-55-7]
 $C_{28}H_{46}N_2O_2$ 442.684
 Alkaloid from *Sarcococca saligna*.
 Powder. $[\alpha]_D^{20}$ -12 (c, 0.2 in $CHCl_3$).
 λ_{max} 211 (log ϵ 1.9) (MeOH).
 Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000, **63**, 1364-1368

3,20-Diaminopregn-5-en-4-ol D-277



$C_{21}H_{36}N_2O$ 332.528


(3 β ,4 β ,20S)-form

N^{20}, N^{20} -Di-Me, N^3 -tigloyl: **Salignarine F**
 [657411-74-0]
 $C_{28}H_{46}N_2O_2$ 442.684
 Alkaloid from *Sarcococca saligna*.
 Cholinesterase inhibitor. Yellow gum.
 $[\alpha]_D^{20}$ -71 (c, 0.01 in MeOH). λ_{max} 203 (log ϵ 2.6) (MeOH).

Atta-ur-Rahman, et al., *Helv. Chim. Acta*, 2004, **87**, 439-448 (*isol, pmr, cmr, ms*)

3,20-Diaminopregn-5-en-15-ol D-278

3,20-Diamino-15-hydroxypregn-5-ene



$C_{21}H_{36}N_2O$ 332.528

(3 β ,15 α ,20S)-form

N^{20}, N^{20} -Di-Me, N^3 -Ac: **N-Acetyl-15 α -hydroxyirehdiamine F**. 15 α -Hydroxy-N-acetylirehdiamine F
 [91147-23-8]
 $C_{25}H_{42}N_2O_2$ 402.619
 Alkaloid from the stem bark of *Didymeleles madagascariensis* (Didymela-ceae). Cryst. (Et₂O). Mp 266°. $[\alpha]_D$ -35 (c, 0.82 in $CHCl_3$).

(3 β ,15 β ,20S)-form

N^{20}, N^{20} -Di-Me, N^3 -Ac: **N-Acetyl-15 β -hydroxyirehdiamine F**. 15 β -Hydroxy-N-acetylirehdiamine F
 [91147-22-7]
 $C_{25}H_{42}N_2O_2$ 402.619
 Alkaloid from the stem bark of *Didymeleles madagascariensis* (Didymela-ceae). Cryst. (Et₂O). Mp 249-250°. $[\alpha]_D$

+91 (c, 1.14 in $CHCl_3$).


N^3, N^{20}, N^{20} -Tri-Me: **20-(Dimethylamino)-3-(methylamino)pregn-5-en-15-ol**. **15 β -Hydroxyirehdiamine H**
 [91147-25-0]

$C_{24}H_{42}N_2O$ 374.609
 Alkaloid from the stem bark of *Didymeleles madagascariensis* (Didymela-ceae). Cryst. (Me₂CO). Mp 206°. $[\alpha]_D$ +8 (c, 0.57 in $CHCl_3$).

Sánchez, V. et al., *Bull. Soc. Chim. Fr., Part II*, 1984, 71-76 (*Didymeleles madagascariensis constits*)

3,20-Diaminopregn-5-en-18-ol D-279

3,20-Diamino-18-hydroxypregn-5-ene



$C_{21}H_{36}N_2O$ 332.528

(3 α ,20S)-form

Holarrhidine

[468-32-6]
 Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 180-181°. $[\alpha]_D$ -23 ($CHCl_3$).

(3 β ,20S)-form

Holarrhimine

[468-31-5]
 Alkaloid from *Holarrhena antidysenterica*, *Holarrhena mitis* and *Holarrhena febrifuga* (Apocynaceae). Mp 183°. $[\alpha]_D$ -14 ($CHCl_3$).

Hydrochloride: Mp 345° dec. $[\alpha]_D^{25}$ -22.8 (MeOH).

N^3 -Me: N^3 -Methylholarrhimine

[27741-48-6]
 $C_{22}H_{38}N_2O$ 346.555
 Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 163-164° (160-163°). $[\alpha]_D$ -19 ($CHCl_3$).

N^{20} -Me: N^{20} -Methylholarrhimine

[95221-60-6]
 [124223-73-0 (hydrochloride)]
 $C_{22}H_{38}N_2O$ 346.555
 Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 165-166°. $[\alpha]_D$ -19 ($CHCl_3$).

N^{20}, N^{20} -Di-Me, N^3 -Ac: **5,6-Dehydromalouphyllinol**

[91147-24-9]
 $C_{25}H_{42}N_2O_2$ 402.619
 Alkaloid from the stem bark of *Didymeleles cf. madagascariensis* (Didymela-ceae), prev. obt. semisynthetically from Holarrhimine. Cryst. (Et₂O). Mp 244-245°. $[\alpha]_D$ -41 (c, 0.35 in $CHCl_3$).

N^3, N^3, N^{20}, N^{20} -Tetra-Me: **N-Tetramethylholarrhimine**. Regholarrhine F
 [10308-37-9]

$C_{25}H_{44}N_2O$ 388.635
 Alkaloid from *Holarrhena antidysenterica* and *Malouetia tamaquarina* (Apocynaceae). Mp 233-235° (227-229°). $[\alpha]_D$ -35 (c, 1 in $CHCl_3$).

Regholarrenine F (from *H. antidy-senterica*, Mp 210-212°) appears identical with the prev. known *N*-Tetramethylholarrhimine.

N-Tetra-Me; hydrochloride: Mp 315-316° dec.

*N*²⁰-Acetoxy: **Holadysenterine** [949087-26-7]

C₂₃H₃₈N₂O₃ 390.565

Alkaloid from the stem bark of *Holarrhena antidy-senterica*, Amorph. powder. Mp 218-220°. [α]_D²⁵ -14.4 (c, 0.23 in MeOH).

Siddiqui, S. et al., *J. Indian Chem. Soc.*, 1932, **9**, 553-563 (*Holarrhimine*)

Cerný, V. et al., *Chem. Listy*, 1955, **49**, 723-730; 1389-1394; 1956, **50**, 1126-1133; 1957, **51**, 2351-2355; **52**, 2344-2350 (*Holarrhidine*, *Holarrhimine*)

Tschesche, R. et al., *Chem. Ber.*, 1958, **91**, 1504-1511 (*N*³-Methylholarrhimine, *N*²⁰-Methylholarrhimine, *N*-Tetramethylholarrhimine)

Labler, L. et al., *Coll. Czech. Chem. Comm.*, 1959, **24**, 370-377; 378-383 (*Holarrhidine*)

Khuong-Huu, Q. et al., *Bull. Soc. Chim. Fr.*, 1965, 3035-3040 (occur, bibl)

Sóti, F. et al., *Tet. Lett.*, 1967, 1437-1441 (*N*-Tetramethylholarrhimine)


Dadoun, H. et al., *Ann. Pharm. Fr.*, 1973, **31**, 237-247 (*Holarrhimine*)

Sánchez, V. et al., *Bull. Soc. Chim. Fr.*, 1984, 71-76 (5,6-Dehydromalouphyllinol)

Bhutani, K.K. et al., *Phytochemistry*, 1990, **29**, 969-972 (*Regholarrenine F*)

Kumar, N. et al., *Chem. Pharm. Bull.*, 2007, **55**, 912-914 (*Holadysenterine*)

3,20-Diaminopregn-16-en-4-ol D-280 ol



C₂₁H₃₆N₂O 332.528

(3β,4β,5α,20S)-form

*N*²⁰,*N*²⁰-Di-Me, *N*³-(3-methyl-2-butenoyl), 4-Ac: **Vaganine D**


[353228-46-3]

C₃₀H₄₈N₂O₃ 484.721

Alkaloid from *Sarcococca coriacea*. Amorph. solid (CHCl₃). Mp 220°. [α]_D²⁵ -21.4 (c, 0.14 in CHCl₃). λ_{max} 206 (log ε 4.2) (MeOH).

Kalauni, S.K. et al., *J. Nat. Prod.*, 2001, **64**, 842-844 (*Vaganine D*)

3,20-Diaminopregn-2-en-4-one D-281 one



C₂₁H₃₄N₂O 330.512

(5α,20S)-form

*N*²⁰-Me, *N*³-tigloyl: **Phulchowkiamide A** [771500-78-8]

C₂₇H₄₂N₂O₂ 426.641

Alkaloid from *Sarcococca hookeriana*. Cholinesterase inhibitor. Gum. [α]_D²⁴ +48 (c, 0.1 in MeOH). λ_{max} 230 (log ε 3.9) (MeOH).

*N*²⁰,*N*²⁰-Di-Me, *N*³-formyl: **Hookeriana-mide H** [957466-14-7]


C₂₄H₃₈N₂O₂ 386.576

Alkaloid from *Sarcococca hookeriana*. Amorph. solid. [α]_D²⁵ +138 (c, 0.3 in MeOH). λ_{max} 224 (log ε 3.3); 252 (log ε 2.7) (MeOH).

Choudhary, M.I. et al., *Helv. Chim. Acta*, 2004, **87**, 1099-1108 (*Phulchowkiamide A*)

Devkota, K.P. et al., *Chem. Pharm. Bull.*, 2007, **55**, 1397-1401 (*Hookerianamide H*)

3,20-Diaminopregn-5-en-4-one D-282 one



C₂₁H₃₄N₂O 330.512

(3β,20S)-form

*N*²⁰-Me, *N*³-(3-methyl-2-butenoyl):

[218604-86-5]

C₂₇H₄₂N₂O₂ 426.641

Alkaloid from *Sarcococca brevifolia*. Amorph. solid. Mp 145-147°. [α]_D²⁵ -8 (c, 0.29 in CHCl₃). λ_{max} 194 (log ε 4.11) (MeOH).

*N*²⁰,*N*²⁰-Di-Me, *N*³-angeloyl: 3-(Ange-loylamino)-20-(dimethylamino)-5-pregnen-4-one. **Sarsalignone** [199922-55-9]

C₂₈H₄₄N₂O₂ 440.668

Alkaloid from *Sarcococca saligna* (Buxaceae). Powder. Mp 278°. [α]_D²⁵ -106.4 (c, 0.1 in MeOH). λ_{max} 237 (log ε 2.9); 257 (log ε 2.9) (MeOH).

*N*²⁰,*N*²⁰-Di-Me, *N*³-(3-methyl-2-butenoyl): **Epipachysamine E 5-en-4-one**

[218604-81-0]

C₂₈H₄₄N₂O₂ 440.668

Alkaloid from *Sarcococca brevifolia*. Amorph. solid. Mp 150-151°. [α]_D²⁵ +6 (c, 0.2 in CHCl₃). Possible incorrect sign of opt. rotation. λ_{max} 194 (log ε 4.11) (MeOH).

Atta-ur-Rahman, et al., *Phytochemistry*, 1997, **46**, 771-775 (isol, uv, ir, pmr, cmr, ms)

Jayasinghe, U.L.B. et al., *Nat. Prod. Lett.*, 1998, **12**, 103-109 (isol, struct)

1,20-Diamino-4,8,12,16-tetraazaecicosane D-283

4,8,12,16-Tetraazaecicosane-1,20-diamine, 9CI. 1,5,7,13,17,22-Hexaazadocosane.

Homocaldohexamine. PA33334

[133416-04-3]

H₂N(CH₂)₄NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₄NH₂

C₁₆H₄₀N₆ 316.532

Prod. by *Thermus thermophilus* HB8, *Marinithermus hydrothermalis* and other thermophilic spp. Isol. from the venom of the spider *Hololena curta*.

*N*⁴-Hydroxy: [864812-17-9]

C₁₆H₄₀N₆O 332.532

Isol. from venom of the spider *Hololena curta*.

Hamana, K. et al., *J. Biochem. (Tokyo)*, 1991, **109**, 444 (isol)

Tzouros, M. et al., *Toxicol.*, 2005, **40**, 350-354 (isol, *N*-hydroxy)

1,20-Diamino-4,8,12,17-tetraazaecicosane D-284

4,8,12,17-Tetraazaecicosane-1,20-diamine, 9CI. 1,5,9,13,18,22-Hexaazadocosane.

Thermohexamine. PA33343

[143085-73-8]

H₂N(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₄NH(CH₂)₃NH₂

C₁₆H₄₀N₆ 316.532

Isol. from the thermophilic bacterium *Bacillus schlegelii* and the venom of the spider *Hololena curta*.

*N*¹,*N*²¹-Di-Ac: [864812-19-1]

C₂₀H₄₄N₆O₂ 400.607

Isol. from the venom of the spider *Hololena curta*.

4-*N*-Hydroxy: [864812-16-8]

C₁₆H₄₀N₆O 332.532

Isol. from the venom of the spider *Hololena curta*.

Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747 (*Bacillus schlegelii* const. isol, struct)

Tzouros, M. et al., *Toxicol.*, 2005, **46**, 350-354 (*Hololena curta* const. isol)

1,20-Diamino-4,8,13,17-tetraazaecicosane D-285

4,8,13,17-Tetraazaecicosane-1,20-diamine, 9CI. 1,5,9,14,18,22-Hexaazadocosane.

Homothermohexamine

[143085-74-9]

H₂N(CH₂)₃NH(CH₂)₃NH(CH₂)₄NH(CH₂)₃NH(CH₂)₃NH₂

C₁₆H₄₀N₆ 316.532

Prod. by a thermophilic bacterium *Bacillus schlegelii* and isol. from the venom of the spider *Agelenopsis aperta*.

Hamana, K. et al., *Biochem. J.*, 1992, **284**, 741-747

Hamana, K. et al., *Microbios*, 1993, **75**, 23-32 (isol)

Tzouros, M. et al., *Toxicol.*, 2005, **46**, 350-354 (isol)

1,24-Diamino-5,10,15,20-tetraazatetracosane D-286

5,10,15,20-Tetraazatetracosane-1,24-diamine, 9CI. 1,6,11,16,21,26-Hexaazahexacosane.

Homohexamine

[104235-48-5]

H₂NCH₂(CH₂)₃NH(CH₂)₄NH(CH₂)₄¹⁰-



$\text{C}_{20}\text{H}_{48}\text{N}_6$ 372.639

Constit. of *Vicia sativa*.

N^5 -(4-Aminobutyl): N^5 -(4-Aminobutyl)-**homohexamine**

[139035-39-5]

$\text{C}_{24}\text{H}_{57}\text{N}_7$ 443.761

Constit. of *Vicia sativa*.

N^{10} -(4-Aminobutyl): N^{10} -(4-Aminobutyl)-**homohexamine**

[139035-40-8]

$\text{C}_{24}\text{H}_{57}\text{N}_7$ 443.761


Constit. of *Vicia sativa*.

Hamana, K. *et al.*, *Phytochemistry*, 1991, **30**, 3319-3322 (occur)

1,17-Diamino-4,9,13-triazaheptadecane D-287

1,5,10,14,19-Pentaazanonadecane. N' -(3-Aminopropyl)canavalmine. PA3434

[129225-31-6]



$\text{C}_{14}\text{H}_{35}\text{N}_5$ 273.464

Constit. of *Canavalia gladiata* (sword bean).

Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (synth, cmr)


Matsuzaki, S. *et al.*, *Phytochemistry*, 1990, **29**, 1311-1312 (isol)

1,16-Diamino-4,8,12-triazahexadecane D-288

N' -[3-[3-[3-(3-Aminopropyl)amino]propyl]amino]propyl]-1,4-butanediamine, 9CI. 4,8,12-Triazahexadecane-1,16-diamine. 1,5,9,13,18-Pentaazaoctadecane.

Homocaldopentamine. PA3334

[84807-66-9]



$\text{C}_{13}\text{H}_{33}\text{N}_5$ 259.437

Prod. by *Thermomicrobium roseum*, *Thermus thermophilus* and other thermophilic spp. Also isol. from venom of the spider *Hololena curta*. Cryst. (as pentahydrochloride).

N^4 -Hydroxy: [864812-12-4]

$\text{C}_{13}\text{H}_{33}\text{N}_5\text{O}$ 275.437

Isol. from the venom of the spider *Hololena curta*.

[107886-58-8]

Oshima, T. *et al.*, *J. Biochem. (Tokyo)*, 1983, **93**, 1455-1456 (isol, struct)

Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (synth, cmr)

Hamana, K. *et al.*, *FEMS Microbiol. Lett.*, 1990, **68**, 27-30; 31-34 (isol)


Tzouros, M. *et al.*, *Toxicon*, 2005, **46**, 350-354 (isol, N-hydroxy)

1,16-Diamino-4,8,13-triazahexadecane D-289

N' -(3-Aminopropyl)- N' -[3-[3-(3-aminopropyl)amino]propyl]-1,4-butanediamine, 9CI. 1,5,9,14,18-Pentaazaoctadecane.

Thermopentamine. PA3343

[102203-44-1]



$\text{C}_{13}\text{H}_{33}\text{N}_5$ 259.437

Prod. by *Bacillus schlegelii*, *Thermomicrobium roseum*, *Thermus thermophilus* and other thermophilic spp. Occurs in *Anthodiaris crassispina* and venom of the spider *Agelenopsis aperta*.

N^4 -Hydroxy: [864812-11-3]

$\text{C}_{13}\text{H}_{33}\text{N}_5\text{O}$ 275.437

Isol. from venom of the spider *Hololena curta*.

[102203-43-0]

Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (synth, cmr)

Hamana, K. *et al.*, *FEMS Microbiol. Lett.*, 1990, **68**, 27-30; 31-34 (isol)

Hamana, K. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (occur, hplc)

Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-747 (*Bacillus schlegelii* constit)

Tzouros, M. *et al.*, *Toxicon*, 2005, **46**, 350-354 (isol, N-hydroxy)

1,19-Diamino-6,11,16-triaza-nonadecane D-290

N' -(4-Aminobutyl)- N' -[4-[4-(4-aminobutyl)amino]butyl]-1,4-butanediamine, 9CI. 1,6,11,16,21-Pentaazadocosane. **Homopentamine**

[15518-43-1]



$\text{C}_{16}\text{H}_{39}\text{N}_5$ 301.518

Constit. of *Vicia sativa*.

N^5 -(4-Aminobutyl): N^5 -(4-Aminobutyl)-**homopentamine**

[139035-36-2]

$\text{C}_{20}\text{H}_{48}\text{N}_6$ 372.639

Constit. of *Vicia sativa*.

N^{10} -(4-Aminobutyl): N^{10} -(4-Aminobutyl)-**homopentamine**

[139035-37-3]

$\text{C}_{20}\text{H}_{48}\text{N}_6$ 372.639

Constit. of *Vicia sativa*.

$\text{N}^5, \text{N}^{10}$ -Bis(4-aminobutyl): $\text{N}^5, \text{N}^{10}$ -**Bis(4-aminobutyl)homopentamine**

[139035-41-9]

$\text{C}_{24}\text{H}_{57}\text{N}_7$ 443.761

Constit. of *Vicia sativa*.

$\text{N}^5, \text{N}^{15}$ -Bis(4-aminobutyl): $\text{N}^5, \text{N}^{15}$ -**Bis(4-aminobutyl)homopentamine**

[139035-42-0]

$\text{C}_{24}\text{H}_{57}\text{N}_7$ 443.761

Constit. of *Vicia sativa*.

[107886-48-6]


Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032 (synth, cmr)

Hamana, K. *et al.*, *Phytochemistry*, 1991, **30**, 3319 (occur)

1,18-Diamino-5,9,14-triazaoctadecane D-291

1,6,10,15,20-Pentaazaeicosane. N' -(4-Aminobutyl)canavalmine. PA4344

[129225-32-7]




$\text{C}_{15}\text{H}_{37}\text{N}_5$ 287.491

Constit. of *Canavalia gladiata* (sword bean).

Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (synth, cmr)

Matsuzaki, S. *et al.*, *Phytochemistry*, 1990, **29**, 1311-1312 (isol)

2,27-Diamino-3,26,28-trihydroxy-11-octacosanone D-292



(2R,3R,26R,27R)-form

$\text{C}_{28}\text{H}_{58}\text{N}_2\text{O}_4$ 486.777

(2R,3R,26R,27R)-form

Rhizochalinin C

26-O-β-D-Galactopyranoside: **Rhizochalin C**

$\text{C}_{34}\text{H}_{68}\text{N}_2\text{O}_9$ 648.919

Constit. of *Rhizochalina incrustata*.

Amorph. solid. $[\alpha]_D^{25}$ -14 (c, 0.09 in MeOH).

(2R,3R,26R,27S)-form

Oceanin

[276879-24-4]

Glass. $[\alpha]_D$ +14.7 (c, 0.3 in MeOH).

26-O-β-D-Glucopyranoside: **Oceanapiside**

[251319-53-6]

$\text{C}_{34}\text{H}_{68}\text{N}_2\text{O}_9$ 648.919

Isol. from the sponge *Oceanapia philippensis*. Antifungal agent. Amorph. solid. $[\alpha]_D$ -5.5 (c, 1.2 in MeOH).

Struct. revised in 2007.

(2S,3S,26R,27S)-form

Calyxinin

[392688-08-3]

Gum. $[\alpha]_D$ -2.2 (c, 0.23 in MeOH).

28-O-β-D-Glucopyranoside: **Calyxoside**

[392688-06-1]

$\text{C}_{34}\text{H}_{68}\text{N}_2\text{O}_9$ 648.919

Isol. from the sponge *Calyx* sp. Gel-like substance. $[\alpha]_D^{25}$ -15.8 (c, 0.32 in MeOH).

Nicholas, G.M. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1678-1681 (*Oceanapiside*, isol)

Nicholas, G.M. *et al.*, *J.A.C.S.*, 2000, **122**, 4011-4019 (*abs config*)

Zhou, B.-N. *et al.*, *Tetrahedron*, 2001, **57**, 9549-9554 (*Calyxoside*)

Makarievva, T.N. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1991-1998 (*Rhizochalin C*, *Oceanapiside*, struct)



Diatom polyamines

D-293



Isol. from the cell walls of diatoms such as *Cylindrotheca fusiformis* and *Nitzschia angularis*. Involved in the species-specific patterning of diatom biosilica.


Kröger, N. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 14133-14138 (isol, struct)

1,13-Diazacyclotetracosane-2,14-dione, 9CI**Clausenlactam**
[95264-91-8]C₂₂H₄₂N₂O₂ 366.586Constit. of the leaves of *Clausena excavata*.Bosch, I. *et al.*, *Tet. Lett.*, 1993, **34**, 4671-4674 (*synth*)Shang, L. *et al.*, *CA*, 1994, **120**, 101930e (*isol*)**Diazaquinomycin B**OM 704B. Antibiotic OM 704B
[87614-39-9]C₂₀H₂₄N₂O₄ 356.421Quinonoid antibiotic. *Isol.* as minor component from *Streptomyces* sp. OM-704. Active against gram-positive bacteria. Inhibits thymidylate synthase and mycoplasmas. Needles. Mp 300°. λ_{max} 277 (ε 20100); 310 (sh) (ε 14800); 325 (sh) (ε 11750); 356 (ε 6230); 373 (ε 6050) (MeOH).**Quinone: Diazaquinomycin A. OM 704A.**
Antibiotic OM 704A
[87614-40-2]C₂₀H₂₂N₂O₄ 354.405*Isol.* from *Streptomyces* sp. OM-704 and *Streptomyces* sp. GW48/1497. Active against gram-positive bacteria. Inhibits folate metabolism. Deep red needles. Fairly sol. MeOH, EtOAc, DMSO, dioxan, CHCl₃; poorly sol. H₂O, hexane, C₆H₆. Mp 291-295°. λ_{max} 250 (sh); 260 (sh); 278 (sh); 286 (ε 21700); 309 (ε 9760); 321 (ε 8950); 367 (ε 4130); 490 (ε 1150) (MeOH).▶ LD₅₀ (mus, ipr) 100 mg/kg. CB9517000
[84419-65-8]Omura, S. *et al.*, *J. Antibiot.*, 1982, **35**, 1425; 1985, **38**, 1016; 1025 (*isol, struct, props*)
Japan. Pat., 1983, 83 205 497; *CA*, **100**, 119343 (*isol*)Omura, S. *et al.*, *Tet. Lett.*, 1983, **24**, 3643 (*isol, uv, pmr*)Kelly, T.R. *et al.*, *Tet. Lett.*, 1988, **29**, 3545 (*synth*)Perez, J.M. *et al.*, *Tetrahedron*, 2000, **56**, 4575-4583 (*synth*)

D-294

Diazaquinomycin C

D-296


C₂₂H₂₆N₂O₄ 382.458Prod. by *Streptomyces* sp. GW48/1497. Brick-red solid. λ_{max} 291 (log ε 4.49); 383 (log ε 3.72); 527 (log ε 3.32) (MeOH).Maskey, R.P. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 137-142**Diazeprinomicin**

D-297

5,10-Dihydro-4,6,8-trihydroxy-10-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-11H-dibenzo[*b,e*][1,4]diazepin-11-one, 9CI. Antibiotic BU 4664L. BU 4664L. ECO 4601


[733035-26-2]

[179981-40-9 (BU 4664L)]

C₂₈H₃₄N₂O₄ 462.588Structure of BU 4664L revised in 2005. Prod. by *Micromonospora* spp. (strain DPJ12, strain 046Eco-11 and strain M990-6). Antimicrobial, antiinflammatory and antitumour agent. 5-Lipoxygenase inhibitor. Mp 184-185° (BU 4664L). λ_{max} 212; 230; 298 (MeOH aq.). λ_{max} 211 (ε 65800) (MeOH). λ_{max} 208 (ε 44300); 553 (ε 9100) (MeOH/NaOH).Charan, R.D. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1431-1433 (*isol, pmr, cmr, ms*)Igarashi, Y. *et al.*, *J. Antibiot.*, 2005, **58**, 350-352 (BU 4664L)McAlpine, J.B. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1585-1590 (*biosynth*)**Diazonamide B**

[131703-15-6]

D-298

R = CH₃

Absolute Configuration

C₃₅H₂₅BrCl₂N₆O₄ 744.43Struct. revised in 2001. Alkaloid from the marine ascidian *Diazona chinensis*. Cytotoxic. Amorph. λ_{max} 205 (ε 110000); 222 (ε 4700); 350 (sh) (MeOH/NaOH) (Derep). λ_{max} 215 (ε 21000); 281 (sh); 290 (ε 4900); 297 (sh) (MeOH) (Derep). λ_{max} 220 (ε 110000); 292 (ε 4700) (MeOH/NaOH) (Berdy).**Debromo, N²-(2S-hydroxy-3-methylbutanoyl): Diazonamide A**
[131727-01-0]C₄₀H₃₄Cl₂N₆O₆ 765.651Alkaloid from *Diazona chinensis*. Cytotoxic. Glass. [α]_D -217.3 (c, 8.8 in MeOH). λ_{max} 204 (ε 101000); 296 (ε 4500); 350 (sh) (MeOH/NaOH) (Derep). λ_{max} 216 (ε 19200); 280 (sh); 290 (ε 5300); 298 (sh) (MeOH) (Derep).**Debromo, N²-(2S-amino-3-methylbutanoyl): Diazonamide C**

[1021691-30-4]

C₄₀H₃₅Cl₂N₇O₅ 764.666Alkaloid from a *Diazona* sp. Cytotoxic. Pale yellow solid. [α]_D²⁵ -24.1 (c, 0.1 in MeOH).**Homologue (R = CH₂CH₃), debromo:****Diazonamide E**

[1021691-36-0]

C₃₆H₂₈Cl₂N₆O₄ 679.56Alkaloid from a *Diazona* sp. Cytotoxic. Pale yellow solid. [α]_D²⁵ -56.8 (c, 0.02 in MeOH).**Homologue (R = CH₂CH₃), debromo,****14-chloro: Diazonamide D**


[1021691-33-7]

C₃₆H₂₇Cl₃N₆O₄ 714.005Alkaloid from a *Diazona* sp. Cytotoxic. Pale yellow solid. [α]_D²⁵ -33.5 (c, 0.1 in MeOH).Lindquist, N. *et al.*, *J.A.C.S.*, 1991, **113**, 2303-2304 (*isol, struct*)Li, J. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 4770-4773 (*struct*)Ritter, T. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 2489-2495 (*rev*)Burgett, A.W.G. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 4961-4966 (*synth*)Nicolaou, K.C. *et al.*, *J.A.C.S.*, 2004, **126**, 12888-12896; 12897-12906; 15316 (*synth*)Lachia, M. *et al.*, *Nat. Prod. Rep.*, 2008, **25**, 227-253 (*rev, synth*)Fernández, R. *et al.*, *Tet. Lett.*, 2008, **49**, 2283-2285 (*Diazonamides C-E*)**Dibenzo[*b,j*]dipyrido[4,3,2-*de*:2',3',4'-*gh*][1,10]phenanthroline, 9CI**

D-299

Eilatrin

[120154-96-3]




C₂₄H₁₂N₄ 356.386
Alkaloid from the Red Sea tunicate *Eudistoma* sp. and by *Cystodytes* sp. Topoisomerase inhibitor, Ni chelating agent. Bright yellow cryst. (CHCl₃/MeOH/H₂O). Mp 310°. λ_{max} 305 (ε 39800); 440 (ε 21500) (MeOH/HCl) (Derep). λ_{max} 242 (ε 48200); 286 (ε 36700); 363 (ε 11500); 388 (ε 21000); 408 (ε 30400); 434 (ε 27000) (MeOH) (Derep). λ_{max} 240 (ε 47600); 275 (ε 35400); 284 (ε 37700); 295 (ε 24800); 367 (ε 13200); 385 (ε 21700); 407 (ε 30500); 432 (ε 27100) (EtOH) (Berdy).

Rudi, A. et al., *Tet. Lett.*, 1988, **29**, 6655-6656 (uv, ir, pmr, cmr, ms, cryst struct)
Rudi, A. et al., *J.O.C.*, 1989, **54**, 5331-5337 (isol, uv, ir, pmr, cmr, ms)
Nakahara, S. et al., *Heterocycles*, 1993, **36**, 1139-1144 (synth)
Gellerman, G. et al., *Tet. Lett.*, 1993, **34**, 1827-1830 (synth)

3,6-Dibenzyl-2,5-dihydroxy-pyrazine D-300

3,6-Bis(phenylmethyl)-2,5-pyrazinediol.
3,6-Dibenzyl-5-hydroxy-2(1H)-pyrazinone



C₁₈H₁₆N₂O₂ 292.337

Mono-Me ether: 5-Methoxy-3,6-bis(phenylmethyl)-2(1H)-pyrazinone, 9CI. 2,5-Dibenzyl-3-hydroxy-6-methoxy-pyrazine
[132213-65-1]

C₁₉H₁₈N₂O₂ 306.363

Isol. from fruiting bodies of the mushroom *Albatrellus confluens* (edibility unknown). Promotes melanin synthesis by B16 melanoma cells. Needles. Mp 159-161°.

2-Me ether, N⁴-oxide: 5-Methoxy-3,6-bis(phenylmethyl)-2(1H)-pyrazinone 4-oxide, 9CI. 2,5-Dibenzyl-3-hydroxy-6-methoxy-pyrazine 1-oxide. **Emeheterone**
[117333-12-7]


C₁₉H₁₈N₂O₃ 322.363

Isol. from *Emericella heterothallica*. Needles (C₆H₆). Sol. MeOH, CHCl₃. Mp 215-217°. Struct. revised in 1990. λ_{max} 228 (ε 28200); 276 (ε 7000); 352 (ε 9800) (MeOH) (Berdy).

Kawahara, N. et al., *Phytochemistry*, 1988, **27**, 3022 (isol, uv, ir, pmr, cmr, ms, Emeheterone)
Ohta, A. et al., *Heterocycles*, 1990, **31**, 1655 (synth, struct, Emeheterone)
Kawagishi, H. et al., *Phytochemistry*, 1996, **42**, 547 (isol, pmr, cmr, cryst struct, Me ether)

2,5-Dibenzylpiperazine D-301

2,5-Bis(phenylmethyl)hexahydropyrazine



C₁₈H₂₂N₂ 266.385

(2S,5S)-form [79982-83-5]
Bp 130-134° approx. [α]_D²⁰ -9.98 (c, 1.492 in EtOH).

Hydrochloride (1:2): [79982-84-6]
Cryst. (EtOH/Et₂O). [α]_D²⁰ -29.65 (c, 2.705 in H₂O).

N,N'-Di-Me: 2,5-Dibenzyl-1,4-dimethylpiperazine
[81536-08-5]

C₂₀H₂₆N₂ 294.439

Alkaloid from the leaves of *Zanthoxylum arborescens* (Rutaceae). Cryst. (EtOAc). Mp 123-124.5°. [α]_D²³ +118 (c, 0.054 in EtOH).

Nagel, U. et al., *Z. Naturforsch., B*, 1974, **36**, 578 (synth)

Grina, J.A. et al., *J.O.C.*, 1982, **47**, 2648 (isol, uv, pmr, cmr, ms, struct, synth)

Takeuchi, S. et al., *Heterocycles*, 1990, **31**, 2073 (synth)

N,N'-Dibenzylurea D-302

N,N'-Bis(phenylmethyl)urea, 9CI
[1466-67-7]

PhCH₂NHCONHCH₂Ph

C₁₅H₁₆N₂O 240.304

Alkaloid from the roots of *Pentadiplandra brazzeana*. Needles (EtOH). Sol. EtOH, AcOH; insol. H₂O. Mp 172-173°.

4''-Methoxy: N-Benzyl-N'-(4-methoxybenzyl)urea
[188911-54-8]

C₁₆H₁₈N₂O₂ 270.33

Alkaloid from the roots of *Pentadiplandra brazzeana*. Cryst. (hexane/EtOAc). Mp 127-129°. λ_{max} 201 (log ε 4.45); 223 (log ε 4.05); 275 (log ε 3.16); 282 (log ε 3.1) (MeOH).

4'',4'''-Dimethoxy: N,N'-Bis(4-methoxybenzyl)urea. N,N'-Di(4-methoxybenzyl)urea
[93731-94-3]

C₁₇H₂₀N₂O₃ 300.357

Alkaloid from *Pentadiplandra brazzeana*. Cryst. (hexane/EtOAc). Mp 171-173°.

Boivin, P.A. et al., *Can. J. Chem.*, 1954, **32**, 242-248 (synth, ir)

Wiley, R.H. et al., *J.A.C.S.*, 1954, **76**, 311 (4'',4'''-dimethoxy, synth)

Wolman, Y. et al., *J.O.C.*, 1962, **27**, 1902 (synth)

Argabright, P.A. et al., *J.O.C.*, 1967, **32**, 3261-3262 (synth)

Vorbrüggen, H. et al., *Chem. Ber.*, 1984, **117**, 1523-1541 (synth)

Atanassova, I.A. et al., *Synth. Commun.*, 1989, **19**, 147-153 (synth, 4'',4'''-dimethoxy, ir)

Leung, M.-K. et al., *J.O.C.*, 1996, **61**, 4175-4179 (synth)

Patonay, T. et al., *Synth. Commun.*, 1996, **26**, 4253 (synth, ir, pmr)

Tsopmo, A. et al., *J. Nat. Prod.*, 1999, **62**, 1435-1436 (4''-methoxy, 4'',4'''-dimethoxy, isol, pmr, cmr, ms)

Hikita, T. et al., *Macromolecules*, 2002, **35**, 6202-6209 (synth, ir, pmr, cmr)

Perveen, S. et al., *Synth. Commun.*, 2005, **35**, 1663-1674 (synth, ir, uv, pmr, ms)

Mizuno, T. et al., *Synthesis*, 2006, 2825-2830 (synth, ir, pmr, cmr, ms)

Arturo, E. et al., *Synthesis*, 2007, 1096-1102 (synth)

Dibromoacetic acid, 9CI, D-303

8CI

[631-64-1]

Br₂CHCOOH

C₂H₂Br₂O₂ 217.845

Deliquescent cryst. Sol. EtOH, Et₂O. Mp 48°. Bp 232-234° dec. Bp₁₅ 75°. pK_a 1.48 (25°).

Me ester: [6482-26-4]

C₃H₄Br₂O₂ 231.871

Bp 182-184°.

Et ester: [617-33-4]

C₄H₆Br₂O₂ 245.898

Constit. of *Asparagopsis armata*. Synthetic reagent. d₂₀²⁰ 1.9. Bp 194° Bp₇₄ 121°. n_D¹⁵ 1.5017.

Bromide:

C₂HBr₃O 280.741

Fuming liq. Bp 194°.

Amide: Dibromoacetamide

[598-70-9]

C₂H₃Br₂NO 216.86

Constit. of *Asparagopsis taxiformis*. Mp 156°.

Nitrile: Dibromoacetoneitrile. Dibromocyanomethane

[3252-43-5]

C₂HBr₂N 198.845

Used with aldehydes and ketones to prepare α-bromoesters. Bp₂₄ 67-69°.

▶AL8450000

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 509A; 849A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 792B; 1367A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 807B (ir)

Genvesse, M.P. et al., *Bull. Soc. Chim. Fr.*, 1892, **7**, 364-366 (synth)

Org. Synth., Coll. Vol., **4**, 1963, 254-256 (nitrile)

Parrot, J. et al., *Bull. Soc. Chim. Fr.*, 1964, 1063-1069 (synth)

Netherlands Pat., 1965, 6 401 521; *CA*, **64**, 3360 (synth)

Elvidge, J.A. et al., *J.C.S. (C)*, 1968, 2188-2198 (use, nitrile)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1969, **2**, 195; 1975, **5**, 186 (use)

Legris, C. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **278**, 77-79 (use, nitrile)


Woolard, F.X. et al., *Tetrahedron*, 1976, **32**, 2843-2846 (amide, isol)

McConnell, O. et al., *Phytochemistry*, 1977, **16**, 367-374 (Et ester, isol)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **3**, 1543-1544 (nitrile, use)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **4**, 2423-2424 (Et ester, use)

4,6-Dibromo-2,3-bis(methylthio)-1H-indole D-304



C₁₀H₉Br₂NS₂ 367.128

Isol. from the red algae *Laurencia brongiartii* and *Laurencia grevilleana*. Mp 108.5-110°.

2-S-Oxide: 4,6-Dibromo-2-(methylsulfinyl)-3-(methylthio)-1H-indole. **Itomanindole A**

[119340-95-3]

C₁₀H₉Br₂NOS₂ 383.127

Isol. from *Laurencia brongiartii*.

Cryst. (CHCl₃). Mp 134-136°. [α]_D +8 (c, 1.56 in CHCl₃). Partial racemate.

3-S-Oxide: 4,6-Dibromo-3-(methylsulfinyl)-2-(methylthio)-1H-indole. **Itomanindole B**

[119340-97-5]

C₁₀H₉Br₂NOS₂ 383.127

Isol. from *Laurencia brongiartii*.

Cryst. (CHCl₃/MeOH). Mp 104-106°. [α]_D -38 (c, 0.33 in CHCl₃/MeOH 1:1). Partial racemate.

2,3-Di-S-oxide: 4,6-Dibromo-2,3-bis(methylsulfinyl)-1H-indole

C₁₀H₉Br₂NO₂S₂ 399.127

Isol. from *Laurencia brongiartii*. Mp 132-134°. λ_{max} 230 (log ε 4.4); 310 (log ε 4) (MeOH).

N-Me, 2-S-oxide: Mp 117.5-118.5°. [α]_D -6.9 (c, 0.54 in CHCl₃).

5-Bromo: 4,5,6-Tribromo-2,3-bis(methylthio)-1H-indole

[128351-86-0]

C₁₀H₈Br₃NS₂ 446.024

Isol. from *Laurencia brongiartii* and *Laurencia grevilleana*.

5-Bromo, 2-S-oxide: 4,5,6-Tribromo-2-(methylsulfinyl)-3-(methylthio)-1H-indole

C₁₀H₈Br₃NOS₂ 462.023


Isol. from *Laurencia brongiartii*. Mp 142-144°. λ_{max} 234 (log ε 4.6); 315 (log ε 4.2) (MeOH).

Christoffersen, C. et al., *Alkaloids (Academic Press)*, 1985, **24**, 25

Tanaka, J. et al., *Tetrahedron*, 1989, **45**, 7301-7310 (isol, pmr, cmr, cryst struct)

El-Gamal, A.A. et al., *J. Nat. Prod.*, 2005, **68**, 815-817 (isol, pmr, cmr, ms)

2,3-Dibromo-4-(3-bromo-2-nitrophenyl)-1H-pyrrole, 9CI **D-305**
Bromonitrin B. Bromopyrrolnitrin B
[23305-91-1]



C₁₀H₅Br₃N₂O₂ 424.874

Pyrrole antibiotic. Isol. from *Pseudomonas pyrrolnitrica* + NH₄Br. Effective against *Trichophyton interdigitale* and *Penicillium chrysogenum*. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 252 (EtOH) (Berdy).

2-Debromo: 3-Bromo-4-(3-bromo-2-nitrophenyl)-1H-pyrrole, 9CI. **Bromonitrin A. Bromopyrrolnitrin A**
[23305-90-0]

C₁₀H₆Br₂N₂O₂ 345.978

Isol. from *Pseudomonas pyrrolnitrica*

or *Pseudomonas acidula*, both with NH₄Br. Effective against *Trichophyton interdigitale* and *Penicillium chrysogenum*. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 252; 275 (EtOH) (Berdy).

3'-Debromo, 5'-bromo: 2,3-Dibromo-4-(5-bromo-2-nitrophenyl)-1H-pyrrole, 9CI. **Bromonitrin C. Bromopyrrolnitrin C**
[32736-23-5]

C₁₀H₅Br₃N₂O₂ 424.874

Isol. from *Pseudomonas pyrrolnitrica* + NH₄Br. Effective against *Trichophyton interdigitale* and *Penicillium chrysogenum*. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 252; 325 (EtOH) (Berdy).


Ajisaka, M. et al., *Agric. Biol. Chem.*, 1969, **33**, 294-295 (Bromonitrins A-C, isol, struct)

Japan. Pat., 1972, 72 14 916; CA, **77**, 60049x (isol)

Van Pee, K.H. et al., *J. Antibiot.*, 1983, **36**, 1735-1742 (Bromonitrins A, B, biosynth, props)

2,7-Dibromo-9H-carbazole, D-306
9CI

[136630-39-2]



C₁₂H₇Br₂N 325.002

Alkaloid from the cyanobacterium *Kyrthuthrix maculans*. Prisms (hexane/C₆H₆). Mp 198-203° Mp 236-238°.


Yamato, T. et al., *J.O.C.*, 1991, **56**, 6248-6250 (synth, ir, pmr)

Lee, S.-C. et al., *Phytochemistry*, 1999, **52**, 537-540 (isol, ir, pmr, cmr, ms)

Dierschke, F. et al., *Synthesis*, 2003, 2471-2472 (synth, pmr, cmr)

3,6-Dibromo-9H-carbazole, D-307
9CI

[6825-20-3]



C₁₂H₇Br₂N 325.002

Alkaloid from the cyanobacterium *Kyrthuthrix maculans*. Prisms or needles (EtOH). Sol. CHCl₃, C₆H₆, spar. sol. petrol. Mp 212-213°.

N-Benzoyl:

C₁₉H₁₁Br₂NO 429.11

Mp 213-214°.

N-Me: [58246-82-5]

C₁₃H₉Br₂N 339.029

Plates. Mp 141°.

N-Et: [33255-13-9]

C₁₄H₁₁Br₂N 353.056

Needles (EtOH). Mp 140-141°.

N-Ph: [57103-20-5]

C₁₈H₁₁Br₂N 401.1

Needles (EtOH). Mp 66-67°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 680B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 162B (nmr)

McLintock, J. et al., *J.C.S.*, 1927, 1214 (synth) Lopatinskii, V.P. et al., *CA*, 1966, **65**, 2203e (synth)

Smith, K. et al., *Tetrahedron*, 1992, **48**, 7479-7488 (synth, pmr, cmr)

Polivin, Y.N. et al., *Izv. Akad. Nauk, Ser. Khim.*, 1993, 234-235 (synth)

Park, M. et al., *Tetrahedron*, 1998, **54**, 12707-12714 (N-Et, N-Ph, synth, pmr, cmr)


Lee, S.-C. et al., *Phytochemistry*, 1999, **52**, 537-540 (isol, pmr, cmr, ms)

Bellon, M. et al., *Cryst. Growth Des.*, 2005, **5**, 1443-1450 (synth, cryst struct)

Ponce, M.B. et al., *Helv. Chim. Acta*, 2006, **89**, 1123-1139 (N-Me, synth, pmr, ms)

2,7-Dibromo-3-chloro-1H-indole, D-308
9CI

3-Chloro-2,7-dibromoindole
[68234-20-8]




C₈H₄Br₂ClN 309.387

Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 86-87°.

Brennan, M.R. et al., *Tet. Lett.*, 1978, **19**, 1637-1640 (isol, pmr, struct)

Erickson, K.L. et al., *Synth. Commun.*, 1981, **11**, 253-259 (synth, ir, pmr, ms)

5,5'-Dibromo-4,4'-dichloro-7,7'-dimethoxy-3,3'-bi-1H-indole, D-309
3,3'-Bis(5-bromo-4-chloro-7-methoxy-1H-indole)




C₁₈H₁₂Br₂Cl₂N₂O₂ 519.018

Isol. from the green alga *Chaetomorpha basiretorsa*. Amorph. powder.

Shi, D.Y. et al., *Chin. Chem. Lett.*, 2005, **16**, 777-780 (isol, pmr, cmr, ms)

4,7-Dibromo-2,3-dichloro-1H-indole, D-310
9CI

[68234-22-0]




C₈H₃Br₂Cl₂N 343.832

Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 142.5-143°.

Brennan, M.R. *et al.*, *Tet. Lett.*, 1978, 1637
(*isol*, *pmr*, *struct*)
Ohta, T. *et al.*, *Heterocycles*, 1989, **29**, 1663
(*synth*)


7-(2,3-Dibromo-4,5-dihydroxybenzyl)-3,7-dihydro-1H-purine-2,6-dione **D-311**
[929006-57-5]



$C_{12}H_8Br_2N_4O_4$ 432.028
Isol. from *Rhodomela confervoides*.
Amorph. powder.

Ma, M. *et al.*, *J. Nat. Prod.*, 2007, **70**, 337-341
(*isol*, *pmr*, *cmr*, *ms*)

N-(2,3-Dibromo-4,5-dihydroxybenzyl)-N'-(3-methoxycarbonylpropyl)urea **D-312**
Methyl N'-(2,3-dibromo-4,5-dihydroxybenzyl) γ-ureidobutyrate




$C_{13}H_{16}Br_2N_2O_5$ 440.088
Constit. of *Rhodomela confervoides*.
Amorph. powder. Mp 155-158°.

$N^{4'}-(2,3-Dibromo-4,5-dihydroxybenzyl):$
 N,N' -Bis(2,3-dibromo-4,5-dihydroxybenzyl)-N-(3-methoxycarbonylpropyl)urea
 $C_{20}H_{20}Br_4N_2O_7$ 720.003
Constit. of *Rhodomela confervoides*.
Brown gum.

Ma, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 206-210
(*isol*, *pmr*, *cmr*, *ms*)

2,4-Dibromo-3,6-dihydroxyphenylacetic acid **D-313**
2,4-Dibromo-3,6-dihydroxybenzeneacetic acid, 9CI



$C_8H_6Br_2O_4$ 325.941

Amide: 2,4-Dibromo-3,6-dihydroxyphenylacetamide. 2,4-Dibromo-3,6-dihydroxybenzeneacetamide, 9CI
[55895-97-1]

$C_8H_7Br_2NO_3$ 324.956
Constit. of *Verongia aurea*. Sol.

MeOH, Et₂O; poorly sol. H₂O. Mp 180-182°. λ_{max} 250 (ϵ 4000); 280 (ϵ 2900); 368 (ϵ 7200) (0.1M NaOH).
 λ_{max} 222 (ϵ 5400) (MeOH).

Krejcarek, G.E. *et al.*, *Tet. Lett.*, 1975, 507 (*uv*, *ir*, *ms*, *cryst struct*)
Krohn, K. *et al.*, *Tet. Lett.*, 1975, 4667 (*uv*, *nmr*, *synth*)

3,5-Dibromo-2,4-dihydroxyphenylacetic acid **D-314**

3,5-Dibromo-2,4-dihydroxybenzeneacetic acid, 9CI
[37677-03-5]

$C_8H_6Br_2O_4$ 325.941

4-Me ether, amide: 3,5-Dibromo-2-hydroxy-4-methoxyphenylacetamide
[28495-12-7]

$C_9H_9Br_2NO_3$ 338.983

Constit. of the sponges *Aplysina aerophoba*, *Aplysina thiona* and *Psammaphysilla purpurea*. Cryst. (CHCl₃). Mp 174-176° (166-167°). λ_{max} 250 (sh) (ϵ 1610); 290 (ϵ 2500); 314 (sh) (ϵ 450) (MeOH) (Derep).

Fattorusso, E. *et al.*, *J.C.S. Perkin 1*, 1972, 16-18 (*synth*)


Chang, C.W.J. *et al.*, *Tet. Lett.*, 1977, **18**, 4005-4007 (*synth*)

Kelecom, A. *et al.*, *Ann. Acad. Bras. Cinc.*, 1979, **51**, 639 (*4-Me ether amide*)

Cruz, F. *et al.*, *J. Nat. Prod.*, 1990, **53**, 543-548 (*4-Me ether, amide*)

6,7-Dibromo-2,4-dihydroxyquinoline **D-315**

6,7-Dibromo-4-hydroxy-2(1H)-quinolinone, 9CI. *6,7-Dibromo-2,4-quinolinediol*
[376598-45-7]




$C_9H_5Br_2NO_2$ 318.952

Isol. from the sponge *Hyrtilis erecta*.
Yellow powder. λ_{max} 216 (ϵ 37400) (MeOH).

Aoki, S. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1372-1374 (*isol*, *pmr*, *cmr*)

3,5-Dibromo-4-ethoxy-1-hydroxy-4-methoxy-2,5-cyclohexadiene-1-acetamide **D-316**

[41841-34-3]



Relative
Configuration

$C_{11}H_{15}Br_2NO_4$ 385.052


Related to Aplysintetal A, A-1353. Alkaloid from an *Aplysina* sp. and *Pseudoceratina purpurea*. Originally thought to be an artifact.

Andersen, R.J. *et al.*, *Tet. Lett.*, 1973, **14**, 1175-1178 (*isol*)

Kijjoa, A. *et al.*, *Z. Naturforsch., B*, 2005, **60**, 904-908 (*isol*)

4,5-Dibromo-2-furancarboxylic acid, 9CI **D-317**

4,5-Dibromo-2-furoic acid, 8CI. *4,5-Dibromopyromucic acid*
[2434-03-9]



$C_5H_2Br_2O_3$ 269.877

Cryst. (H₂O). Mp 170.5-171.5°.

tert-Butyl ester: [54113-43-8]

$C_9H_{10}Br_2O_3$ 325.984

Syrup.

Amide: 4,5-Dibromo-2-furancarboxamide

$C_5H_3Br_2NO_2$ 268.892

Isol. from the marine sponge *Phakellia conulus* from the Andaman Islands. Needles (MeOH aq.). Mp 142-143°. λ_{max} 289 (ϵ 35000) (EtOH).

[54113-41-6, 54113-42-7]

Divald, S. *et al.*, *J.O.C.*, 1976, **41**, 2835-2846 (*synth*, *pmr*)

Sarma, N.S. *et al.*, *Indian J. Chem., Sect. B*, 1990, **29**, 771-772 (*amide*)


Karminski-Zamola, G. *et al.*, *Heterocycles*, 1994, **38**, 759-767 (*synth*)

Bury, P. *et al.*, *Tetrahedron*, 1994, **50**, 8793-8808 (*synth*, *ir*, *pmr*, *ms*)

Muratake, H. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 799-806 (*tert-butyl ester*)

3,5-Dibromo-1-hydroxy-4,4-dimethoxy-2,5-cyclohexadiene-1-acetamide, 9CI **D-318**

[24742-01-6]



$C_{10}H_{13}Br_2NO_4$ 371.025


Related to Aplysintetal A, A-1353. Isol. from the sponges *Aplysina* sp., *Verongia fistularis* and *Verongia cauliformis*. Prisms (EtOAc). Fairly sol. Et₂O, EtOAc. Mp 191°.

Sharma, G.M. *et al.*, *J. Antibiot., Ser. A*, 1967, **20**, 200-203 (*isol*)

Sharma, G.M. *et al.*, *J.O.C.*, 1970, **35**, 2823-2826 (*ir*, *pmr*, *struct*)

Toscano, R.A. *et al.*, *Acta Cryst. C*, 1992, **48**, 2235-2237 (*cryst struct*)

7,9-Dibromo-10-hydroxy-8-methoxy-6-methyl-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxylic acid **D-319**




C₁₁H₁₁Br₂NO₅ 397.02**(5R*,10S*)-form**Amide, N-(2-carboxyethyl): **Purpuroceratic acid A**

[870071-25-3]

C₁₄H₁₆Br₂N₂O₆ 468.098Alkaloid from *Pseudoceratina purpurea*. Gum. [α]_D²⁰ -27.7 (c, 0.55 in MeOH).Amide, N-(3-carboxypropyl): **Purpuroceratic acid B**

[870071-26-4]

C₁₅H₁₈Br₂N₂O₆ 482.125Alkaloid from *Pseudoceratina purpurea*.Kijjoo, A. et al., *Z. Naturforsch., B*, 2005, **60**, 904-908 (isol, pmr, cmr)**7,9-Dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxylic acid, 9CI** D-320**(5R,6S)-form**C₁₀H₉Br₂NO₅ 382.993**(5R,10S)-form** [168480-66-8]Isol. from the sponge *Pseudoceratina* sp. Antibacterial agent. λ_{max} 224 (ε 19000); 285 (ε 9400) (MeOH).**(5S,10R)-form***Me ester*: [179798-47-1]

[160116-56-3]

C₁₁H₁₁Br₂NO₅ 397.02Alkaloid from sponges *Aplysina archeri* and *Verongula* sp.

Amide, N-(3-carboxypropyl): [945408-72-0]

[167469-52-5]

C₁₀H₁₀Br₂N₂O₄ 382.008Alkaloid from the Okinawan marine sponge *Psammaphysilla pura* and from the Caribbean sponge *Verongula* sp. Oil. [α]_D²⁴ +86 (c, 0.19 in MeOH).

Amide, N-(3-carboxypropyl): [945408-72-0]

C₁₄H₁₆Br₂N₂O₆ 468.098Alkaloid from *Aplysina fulva*.Amorph. solid. [α]_D²³ +140 (c, 0.04 in MeOH). λ_{max} 223 (log ε 3.93); 281 (log ε 3.63) (MeOH).

Amide, N-[4-(methoxycarbonylamino)-2-oxobutyl]: [223580-93-6]

C₁₆H₁₉Br₂N₃O₇ 525.15Alkaloid from *Aplysina cauliformis*. λ_{max} 229 (ε 18500); 280 (ε 10600) (MeOH).

Amide, N-[4-(methoxycarbonylamino)-3-oxobutyl]: [223580-91-4]


C₁₆H₁₉Br₂N₃O₇ 525.15Alkaloid from *Aplysina cauliformis*. λ_{max} 227 (ε 19000); 281 (ε 10500)

(MeOH).

Ciminiello, P. et al., *J. Nat. Prod.*, 1994, **57**, 1564-1569 (isol)Aiello, A. et al., *Biochem. Syst. Ecol.*, 1995, **23**, 377-381 (isol)Kobayashi, J. et al., *Chem. Pharm. Bull.*, 1995, **43**, 403-407 (isol, uv, cd, ir, pmr, cmr)Ciminiello, P. et al., *J. Nat. Prod.*, 1999, **62**, 590-593 (*Aplysina cauliformis* constits)Rogers, E.W. et al., *J. Nat. Prod.*, 2007, **70**, 1191-1194 (*Aplysina fulva* constit)**(3,5-Dibromo-4-hydroxyphenyl)acetic acid** D-321

3,5-Dibromo-4-hydroxybenzeneacetic acid, 9CI

[24744-58-9]

C₈H₆Br₂O₃ 309.942Constit. of the red alga *Halopytis incurvus*. Cryst. (H₂O). Mp 195-196°.*Et ester*: [24744-59-0]C₁₀H₁₀Br₂O₃ 337.995

Cryst. (hexane). Mp 105°.

Amide: 3,5-Dibromo-4-hydroxyphenylacetamide

[17194-96-6]

C₈H₇Br₂NO₂ 308.957Isol. from the sponges *Aplysina fistularis* and *Verongia archeri*. Antimicrobial agent. Mp 190-191°.

Nitrile: 3,5-Dibromo-4-hydroxybenzeneacetonitrile. 3,5-Dibromo-4-hydroxybenzyl cyanide. 2,6-Dibromo-4-(cyanomethyl)phenol


[73348-20-6]

C₈H₅Br₂NO 290.942Isol. from the sponge *Verongia aurea*. Antibacterial agent. Pale yellow solid. Mp 176.5°.*Me ether, nitrile*: 3,5-Dibromo-4-methoxybenzeneacetonitrile. 3,5-Dibromo-4-methoxybenzyl cyanide. 2,6-Dibromo-4-(cyanomethyl)anisole

[55027-78-6]

C₉H₇Br₂NO 304.968Isol. from *Verongia aurea*. Antibacterial agent.Sharma, G.M. et al., *J.O.C.*, 1970, **35**, 2823-2826 (synth)Chantraine, J.M. et al., *Phytochemistry*, 1973, **12**, 1793-1796 (isol)Yamada, Y. et al., *Chem. Lett.*, 1974, 1399-1400 (synth, amide)Chib, J.S. et al., *J. Pharm. Sci.*, 1978, **67**, 264-265 (isol, amide)Goo, Y.M. et al., *CA*, 1980, **92**, 169100t (nitriles)D'Ambrosio, M. et al., *Helv. Chim. Acta*, 1984, **67**, 1484-1492 (synth)Goo, Y.M. et al., *Arch. Pharmacol. Res.*, 1985, **8**, 21-30 (amide, isol)Boehlow, T.R. et al., *J.O.C.*, 2001, **66**, 3111-3118 (nitrile, synth, ir, pmr, cmr)Nuñez, C.V. et al., *Biochem. Syst. Ecol.*, 2008, **36**, 283-296 (amide, isol)**2-(3,5-Dibromo-4-hydroxyphenyl)ethylamine** D-322

4-(2-Aminoethyl)-2,6-dibromophenol, 9CI. 3,5-Dibromo-4-hydroxybenzeneethanamine. 3,5-Dibromotyramine [134755-34-3]

C₈H₉Br₂NO 294.973

N,N,N-Tri-Me: 3,5-Dibromo-4-hydroxy-N,N,N-trimethylbenzeneethanaminium, 9CI. 3,5-Dibromo-4-hydroxyphenethyltrimethylammonium. 3,5-Dibromo-N,N,N-trimethyltyramine [73414-57-0]

C₁₁H₁₆Br₂NO⁺ 338.062Isol. from the sponge *Verongia fistularis*. Dual adrenergic agent. Needles (MeOH) (? as chloride). Poorly sol. hexane. Mp 230° dec. (? chloride).*Me ether*: (3,5-Dibromo-4-methoxyphenyl)ethylamine. 3,5-Dibromo-4-methoxybenzeneethanamine, 9CIC₉H₁₁Br₂NO 309Alkaloid from a *Eudistoma* sp.

Amorph. solid (as trifluoroacetate).

λ_{max} 208 (log ε 4.9); 224 (log ε 4.3); 284 (log ε 3) (MeOH) (as trifluoroacetate).*Me ether, N,N,N-tri-Me*: 3,5-Dibromo-4-methoxy-N,N,N-trimethylbenzeneethanaminium. 3,5-Dibromo-N,N,N,O-tetramethyltyramine

[160116-58-5]

[160116-59-6]


C₁₂H₁₈Br₂NO⁺ 352.088Metab. of the Caribbean sponge *Verongula* sp. and a *Pseudoceratina* sp. Amorph. solid (as trifluoroacetate). CAS no. refers to trifluoroacetate. λ_{max} 277 (ε 1260) (EtOH) (as trifluoroacetate).

[73414-58-1, 13062-88-9]

Benington, F. et al., *J.O.C.*, 1958, **23**, 1979-1984 (synth)Hollenbeak, K.H. et al., *CA*, 1980, **92**, 180749g (isol)Ciminiello, P. et al., *J. Nat. Prod.*, 1994, **57**, 1564-1569 (*N,N,N,O-tetra-Me*)Aiello, A. et al., *Biochem. Syst. Ecol.*, 1995, **23**, 377-381 (*N,N,N,O-tetra-Me*)Van Wagoner, R.M. et al., *J. Nat. Prod.*, 1999, **62**, 794-797 (isol, synth, uv, pmr, cmr, *Me ether*)Fu, X. et al., *J. Nat. Prod.*, 1999, **62**, 1072-1073 (isol, pmr, cmr, *N-tri-Me*)Schoenfeld, R.C. et al., *Bioorg. Med. Chem. Lett.*, 2002, **12**, 823-825 (synth)**3-(3,5-Dibromo-4-hydroxyphenyl)-2-propenoic acid** D-323

3,5-Dibromo-4-hydroxycinnamic acid

[119405-33-3]



C₉H₆Br₂O₃ 321.953**(E)-form** [56926-78-4]

Mp 244°.

4-O-(3-Dimethylaminopropyl)-3,5-Dibromo-4-[3-(dimethylamino)propoxy]-cinnamic acid

[134276-56-5]

C₁₄H₁₇Br₂NO₃ 407.101Isol. from the sponge *Pseudoceratina crassa*. Cryst. Mp 193-194° dec. λ_{max} 223 (ε 13700); 267 (ε 11000) (MeOH) (Berdy).

► Genotoxic.

4-O-(3-Dimethylaminopropyl), Et ester: [134276-55-4]

C₁₆H₂₁Br₂NO₃ 435.155Isol. from *Pseudoceratina crassa*.

Antibacterial agent. Flakes. Mp 67°.


λ_{max} 230 (ε 11200); 282 (ε 9500)

(MeOH) (Berdy).


► Genotoxic.

Gupta, S.K. *et al.*, *J. Indian Chem. Soc.*, 1988, **65**, 187 (*synth*)Kassühlke, K.E. *et al.*, *Tetrahedron*, 1991, **47**, 1809-1814 (*isol. synth. derivs*)**2-(3,5-Dibromo-2-hydroxyphenyl)-3,4,5-tribromopyrrole** **D-324**2,4-Dibromo-6-(3,4,5-tribromo-1H-pyrrol-2-yl)phenol, 9CI. **Pentabromopseudilin**

[10245-81-5]



C₁₀H₄Br₅NO 553.668Pyrrole antibiotic. Isol. from the marine bacterium *Pseudomonas bromoutilis* and as the major metab. of yellow and off-white strains of *Chromobacterium*. Also from another *Chromobacterium* sp. and *Alteromonas luteo-violaceus*. Active against gram-positive bacteria. Inhibitor of esterases, aminopeptidases, elastase, urokinase and alkaline phosphatase. Cryst. (C₆H₆/hexane). Sol. Me₂CO, butanol, EtOAc, Et₂O; fairly sol. CHCl₃, MeOH; poorly sol. H₂O, hexane. Mp 130-170° dec. λ_{max} 286; 308 (MeOH) (Berdy). λ_{max} 284; 293; 355 (MeOH/NaOH) (Berdy).► LD₅₀ (mus, ivn) 25 - 75 mg/kg, LD₅₀ (mus, scu) 250 - 350 mg/kg, LD₅₀ (mus, ipr) 50 mg/kg.

Me ether: Mp 124-125°.

Chloro analogue: 2,4-Dichloro-6-(2,3,4-trichloro-1H-pyrrol-5-yl)phenol. 2-(3,5-Dichloro-2-hydroxyphenyl)-3,4,5-trichloropyrrole. **Pentachloropseudilin**. A 15104Y. *Antibiotic A 15104 Y* [69640-38-6]C₁₀H₄Cl₅NO 331.411Prod. by *Actinoplanes* sp. ATCC33002. Cryst. (petrol). Sol. bases, CHCl₃, MeOH; poorly sol. H₂O, acids. Mp 126-130°. λ_{max} 284; 309 (EtOH). λ_{max} 282 (ε 9200); 310 (ε 7600) (MeOH) (Berdy). λ_{max} 330 (MeOH-NaOH) (Berdy).► LD₅₀ (mus, ipr) 5.33 mg/kg.Burkholder, P.R. *et al.*, *Appl. Microbiol.*, 1966, **14**, 649 (*isol. struct*)Hanessian, S. *et al.*, *J.A.C.S.*, 1966, **88**, 4509 (*synth. ms*)Lovell, F.M. *et al.*, *J.A.C.S.*, 1966, **88**, 4510 (*cryst. struct*)ApSimon, J.W. *et al.*, *Chem. Ind. (London)*, 1973, 275 (*synth*)Andersen, R.J. *et al.*, *Mar. Biol. (Berlin)*, 1974, **27**, 281 (*isol. uv. pmr. ms*)Cavalleri, B. *et al.*, *Curr. Microbiol.*, 1978, **1**, 319-324 (*chloro analogue, isol*)ApSimon, J.W. *et al.*, *J.C.S. Perkin 1*, 1978, 1588-1594 (*chloro analogue, synth*)Faulkner, D.J. *et al.*, *Top. Antibiot. Chem.*, 1978, **2**, 9Laatsch, H. *et al.*, *Annalen*, 1989, 863 (*synth. bibl*)Hanefeld, U. *et al.*, *J.O.C.*, 1994, **59**, 3604 (*biosynth*)Xu, Z. *et al.*, *J.O.C.*, 1998, **63**, 5031-5041 (*synth*)**6,6'-Dibromoindigotin** **D-325**6-Bromo-2-(6-bromo-1,3-dihydro-3-oxo-2H-indol-2-ylidene)-1,2-dihydro-3H-indol-3-one, 9CI. 6,6'-Dibromoindigo. *Tyrian purple*. C.I. Natural Violet 1. CI 75800 [19201-53-7]C₁₆H₈Br₂N₂O₂ 420.059Isol. from molluscs *Murex trunculus*, *Purpura* spp., *Dicathais orbita*, *Mancinella keineri*, *Ptychodera flava laysanica* and others. Important dyestuff in ancient times, but has never been prod. commercially. Violet cryst. (1,2-dichlorobenzene). Mp > 300°.

N,N'-Di-Ac: Mp 306°.


Monobromo: 6-Bromoindigotin. 6-Bromoindigo [139582-54-0]

C₁₆H₉BrN₂O₂ 341.163Constit. of glandular secretions of *Murex trunculus*. Black solid with copper lustre (ethyl benzoate).Sachs, F. *et al.*, *Ber.*, 1904, **37**, 1861-1874 (*synth*)Friedländer, P. *et al.*, *Ber.*, 1909, **42**, 765-770 (*struct*)Baker, J.T. *et al.*, *Tet. Lett.*, 1968, 43-46 (*biosynth*)Fouquet, H. *et al.*, *Angew. Chem., Int. Ed.*, 1971, **10**, 816-817 (*rev*)Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 227-230 (*isol*)Süsse, P. *et al.*, *Naturwissenschaften*, 1979, **66**, 110 (*cryst. struct*)Larsen, S. *et al.*, *Acta Chem. Scand., Ser. A*, 1980, **34**, 171-176 (*cryst. struct*)Voss, G. *et al.*, *Chem. Ber.*, 1989, **122**, 1199-1201 (*synth. ir. bibl*)McGovern, P.E. *et al.*, *Acc. Chem. Res.*, 1990, **23**, 152-158 (*rev*)Wouters, J. *et al.*, *J. Soc. Dyers Colour.*, 1991, **107**, 266-269 (*hplc*)Kirk-Othmer *Encycl. Chem. Technol.*, 4th edn., Wiley, 1991, **8**, 559 (*bibl*)Michel, R.H. *et al.*, *J. Soc. Dyers Colour.*, 1992, **108**, 145-150 (*occur. 6-debromo*)Ghireti, F. *et al.*, *Experientia*, 1994, **50**, 802-807 (*bibl*)Koren, Z.C. *et al.*, *Isr. J. Chem.*, 1995, **35**, 117-124 (*hplc*)Clark, R.J.H. *et al.*, *Nouv. J. Chim.*, 1999, **23**, 323-328 (6-debromo, *synth. ir*)Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-739 (*rev*)Imming, P. *et al.*, *Synth. Commun.*, 2001, **31**, 3721-3727 (*synth*)Tanoue, Y. *et al.*, *The Fischer Indole Synthesis*, 2001, **67**, 726-729 (*synth*)**3,6-Dibromo-1H-indole, 9CI** **D-326**
[74076-56-5]C₈H₅Br₂N 274.942Isol. from the ascidian *Distaplia regina*. Probable constit. of the hemichordates *Ptychodera flava* and *Glossobalanus* sp. Shows antibacterial activity. Oil. λ_{max} 224 (log ε 5.26); 282 (log ε 4.59); 293 (log ε 4.48) (hexane).Higa, T. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1980, **65**, 525-530 (*isol*)Qureshi, A. *et al.*, *Nat. Prod. Lett.*, 1999, **13**, 59-62 (*isol. pmr. cmr. activity*)**4,6-Dibromo-1H-indole, 9CI** **D-327**
Glossobalol
[99910-50-6]C₈H₅Br₂N 274.942Alkaloid from the acorn worm *Glossobalanus* sp. Mp 60-61°.

Ac:

C₁₀H₇Br₂NO 316.979

Mp 108-108.5° (102-104°).

Higa, T. *et al.*, *Experientia*, 1985, **41**, 1487 (*isol. ir. pmr. ms. struct*)Martin, P. *et al.*, *Tet. Lett.*, 1987, **28**, 1645 (*synth*)Ohta, T. *et al.*, *Heterocycles*, 1987, **26**, 2817 (*synth*)**Dibromoisophakellin** **D-328**
Dibromocantharelline**(+)-form**C₁₁H₁₁Br₂N₅O 389.049**(+)-form** [101481-34-9]Metab. from the New Caledonian sponge *Pseudaxinyssa cantharella*. Cryst. +

2H₂O + 1MeOH (MeOH). Mp 260°. [α]_D²⁰ +95 (c, 1.0 in MeOH).

2-Debromo: Monobromoisophakellin

[425668-66-2]

C₁₁H₁₂BrN₅O 310.153

Isol. from *Agelas* sp. λ_{max} 276 (log ε 3.84) (MeOH).

11α-Hydroxy, 12α-chloro: 12-Chloro-11-hydroxydibromoisophakellin

[389120-65-4]

C₁₁H₁₀Br₂ClN₅O₂ 439.493

Isol. from the sponge *Axinella brevisityla*. [α]_D²³ +51 (c, 0.41 in MeOH). λ_{max} 207 (log ε 4.4); 242 (sh) (log ε 3.8) (MeOH).

(-)-form [104758-96-5]

Metab. from the marine sponge *Acanthella carteri*. Cryst. (MeOH) (as hydrochloride). Mp 275-277° (hydrochloride). [α]_D -101 (c, 0.56 in MeOH).

N¹-Me: Dibromo-N¹-methylisophakellin

N¹-Methyldibromoisophakellin

[373604-86-5]

C₁₂H₁₃Br₂N₅O 403.076

Isol. from the sponge *Stylissa caribica*. λ_{max} 288 (log ε 3.92) (H₂O).

[105088-20-8]

De Nanteuil, G. *et al.*, *Tetrahedron*, 1985, **41**, 6019-6033 (isol, uv, ir, pmr, cmr, ms, cd, cryst struct, Dibromocantharelline, (+)-form)

Fedoreyev, S.A. *et al.*, *Tet. Lett.*, 1986, **27**, 3177-3180 (uv, ir, pmr, cmr ms, cd, cryst struct, (-)-form)

Assmann, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1345-1347 (*N*-Me)

Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1576-1578

(Chlorohydroxydibromoisophakellin)


Wiese, K.J. *et al.*, *Tet. Lett.*, 2002, **43**, 5135-5136 (synth)

Assmann, M. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 153-156 (Monobromoisophakellin)

Meyer, S.W. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1524-1529 (pmr, cmr, N-15 nmr)

5,7-Dibromo-6-methoxy-1H-indole, 9CI D-329

[58933-47-4]



C₉H₇Br₂NO 304.968


Alkaloid from the hemichordate *Ptychodera flava laysanica*.

Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 227-230 (isol, pmr, struct)

2-(2,4-Dibromo-5-methoxyphenyl)ethylamine D-330

2,4-Dibromo-5-methoxybenzeneethanamine, 9CI

[121135-06-6]




C₉H₁₁Br₂NO 309

First report of a naturally occurring brominated β-phenylethylamine. Isol. from the bryozoan *Amathia wilsoni*. Possible biosynthetic precursor to the Amathamide alkaloids. Oil.

Blackman, A.J. *et al.*, *J. Nat. Prod.*, 1989, **52**, 436-438 (isol, ir, pmr, cmr, ms, struct)

3-(3,5-Dibromo-4-methoxyphenyl)-2-methoxy-N-methylpropylamine D-331

3,5-Dibromo-β,4-dimethoxy-N-methylbenzenepropanamine



C₁₂H₁₇Br₂NO₂ 367.08


(ξ)-form

Isol. from a *Pachychalina* sp. [α]_D²⁹ -6.4 (c, 0.001 in MeOH). λ_{max} 283 (log ε 4.09) (MeOH).

De Oliveira, M.F. *et al.*, *Planta Med.*, 2006, **72**, 437-441 (isol, pmr, cmr, ms)

4,6-Dibromo-2-methyl-1H-indole, 9CI D-332

[99910-51-7]



C₉H₇Br₂N 288.969

Alkaloid from the acorn worm *Glossobalanus* sp. Mp 96-96.5°.

Ac:

C₁₁H₉Br₂NO 331.006


Mp 146-146.5° (141-143.5°).

Higa, T. *et al.*, *Experientia*, 1985, **41**, 1487 (isol, ir, pmr, ms, struct)

Ohta, T. *et al.*, *Heterocycles*, 1987, **26**, 2817 (synth)

N-[2-[3,5-Dibromo-4-[3-(3-methyl-1-oxo-2-butenyl)amino]propoxy]phenyl]ethyl]-4-hydroxybenzeneacetamide, 9CI D-333

1,3-Dibromo-5-[2-[(4-hydroxyphenyl)acetamido]ethyl]-2-[3-(3-methyl-2-butenamido)propoxy]benzene [160666-52-4]



C₂₄H₂₈Br₂N₂O₄ 568.304

Metab. from the Caribbean sponge *Iotrochota birotulata*.

Monoiodo analogue: N-[2-[3-Bromo-5-iodo-4-[3-[(3-methyl-1-oxo-2-butenyl-

l)amino]propoxy]phenyl]ethyl]-4-hydroxybenzeneacetamide, 9CI

[160666-53-5]

C₂₄H₂₈BrIN₂O₄ 615.305

Metab. from *Iotrochota birotulata*.

Diiodo analogue: N-[2-[3,5-Diiodo-4-[3-[(3-methyl-1-oxo-2-butenyl)amino]propoxy]phenyl]ethyl]-4-hydroxybenzeneacetamide, 9CI

[160666-54-6]


C₂₄H₂₈I₂N₂O₄ 662.305

Metab. from *Iotrochota birotulata*.

Costantino, V. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1552 (isol, ir, pmr, cmr, struct)

4,6-Dibromo-2-methylthio-1H-indole D-334

[128351-88-2]




C₉H₇Br₂NS 321.035

Alkaloid from the Okinawan red alga *Laurencia brongiartii*. Cryst. (hexane/CCl₄). Mp 59-61°.

Tanaka, J. *et al.*, *Tetrahedron*, 1989, **45**, 7301 (isol, ir, pmr, ms, struct)

5-[3,5-Dibromo-4-[(2-oxo-5-oxazolidinyl)methoxy]phenyl]-2-oxazolidinone, 9CI D-335

[54448-72-5]



C₁₃H₁₂Br₂N₂O₅ 436.056

(5R,5'R)-form [120442-03-7]

Isol. from sponge *Aplysina aerophoba*. Mp 220-222°. [α]_D -33 (c, 1.1 in MeOH).

(5R,5'S)-form

Isol. from the sponges *Aplysina lacunosa* and *Verongula rigida* and ascidian *Clavelina oblonga*. Mp 220-222°. [α]_D -6.5 (c, 0.44 in MeOH). [α]_D²⁴ -9.2 (c, 0.25 in MeOH).

(5S,5'R)-form

Isol. from the sponges *Aplysina fistularis* (*Verongia fistularis*) forma *fulva*, *Aplysina lacunosa* and *Aplysina cauliformis*. Mp 226-227° (222-225°). [α]_D²⁰ +7.1 (c, 0.35 in MeOH).

2-Ac: [474670-22-9]

C₁₅H₁₄Br₂N₂O₆ 478.093

Isol. from *Suberea* aff. *praetensa*.

2,2'-Di-Ac: [474670-23-0]

C₁₇H₁₆Br₂N₂O₇ 520.131

Isol. from *Suberea* aff. *praetensa*.

(5RS,5'SR)-form

Isol. from the sponges *Aplysina cauliformis* and *Aplysina fistularis*. Mp 247-249°.

(5ξ,5'ξ)-form

O-Ac:

C₁₅H₁₄Br₂N₂O₆ 478.093Isol. from *Suberea* aff. *praetensa* sponge. Derived from a hydroxyoxazolidine tautomer.


O,O-Di-Ac:

C₁₇H₁₆Br₂N₂O₇ 520.131Isol. from a *Suberea* sponge. Derived from the bis(hydroxyoxazolidine) tautomer.Borders, D.B. *et al.*, *Tet. Lett.*, 1974, 2709-2712 (*isol, uv, ir, pmr, cmr, ms, struct*)Gopichand, Y. *et al.*, *Tet. Lett.*, 1979, 3921-3924 (*isol*)Makarieva, T.N. *et al.*, *Comp. Biochem. Physiol. B: Comp. Biochem.*, 1981, **68**, 481-484 (*occur*)Norte, M. *et al.*, *Tetrahedron*, 1988, **44**, 4973 (*5R,5'R-form, isol, cryst struct, abs config*)Kijjioa, A. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 732-738 (*Suberea acetates*)Kossuga, M.H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1879-1881 (*Clavelina oblonga* *constit, isol, cmr*)Rogers, E.W. *et al.*, *J. Nat. Prod.*, 2005, **68**, 891-896 (*stereochem*)

Dibromophakellstatin

D-336

[185750-71-4]




Absolute Configuration

C₁₁H₁₀Br₂N₄O₂ 390.034Alkaloid from the Indian ocean sponge *Phakellia mauritiana*. Antineoplastic agent. Cryst. (toluene/MeOH). Mp 245° (dec.).Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1997, **60**, 180-183 (*isol, pmr, cmr, ms, cryst struct*)Wiese, K.J. *et al.*, *Tet. Lett.*, 2002, **43**, 5135-5136 (*synth*)Poullennec, K.G. *et al.*, *J.A.C.S.*, 2003, **125**, 6344-6345 (*synth*)Cheung, R. *et al.*, *Org. Lett.*, 2004, **6**, 3881-3884 (*synth*)Jacquot, D.E.N. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 2295-2298 (*synth*)Feldman, K.S. *et al.*, *Org. Lett.*, 2005, **7**, 929-931 (*synth*)Zöllinger, M. *et al.*, *J.O.C.*, 2006, **71**, 9431-9439 (*synth*)Atodiresci, I. *et al.*, *Chirality*, 2007, **19**, 542-549 (*electronic cd*)

2,3-Dibromo-1H-pyrrole, 9CI

D-337


[69624-10-8]

C₄H₃Br₂N 224.882Metab. from a marine sponge *Agelas* sp. Extremely unstable.Audebert, P. *et al.*, *Synth. Met.*, 1986, **15**, 9 (*ms, pmr*)Tada, H. *et al.*, *Chem. Lett.*, 1988, 803 (*isol, ir, pmr, cmr, ms*)

2,5-Dibromo-1H-pyrrole-3-carboxylic acid

D-338


[270084-77-0]

C₅H₃Br₂NO₂ 268.892Isol. from *Spongia oblique*.Xu, S.-H. *et al.*, *Youji Huaxue*, 2000, **20**, 248-250; *CA*, **132**, 345562a (*isol*)

4,5-Dibromo-1H-pyrrole-2-carboxylic acid, 9CI

D-339

[34649-21-3]

C₅H₃Br₂NO₂ 268.892Constit. of the sponges *Agelas oroides*, *Agelas clathrodes*, *Agelas conifera* and *Agelas flabelliformis*. Possesses potent *in vitro* immunosuppressive activity. Feeding deterrent. Mp 148° subl.*Me ester: Methyl 4,5-dibromo-1H-pyrrole-2-carboxylate, 9CI*

[937-16-6]

C₆H₅Br₂NO₂ 282.919Constit. of *Agelas oroides*, *Phakellia fusca* and a *Lissodendoryx* sp. Mp 161° (155°).*Amide: 4,5-Dibromo-1H-pyrrole-2-carboxamide, 9CI*

[34649-20-2]

C₅H₄Br₂N₂O 267.907Constit. of *Acanthella carteri*, *Agelas mauritiana*, *Agelas oroides*, *Phakellia fusca* and *Pseudoceratina purpurea*. Larval metamorphosis-promoting compd. Feeding deterrent for ascidians. Mp 164-166°. λ_{max} 233 (ε 6200); 273 (ε 11800) (MeOH) (Berdy). λ_{max} 296 (ε 18800) (MeOH/NaOH) (Berdy).*(Methoxymethyl)amide: 4,5-Dibromo-N²-methoxymethyl-1H-pyrrole-2-carboxamide*

[219782-98-6]

C₇H₈Br₂N₂O₂ 311.96Isol. from the sponge *Homaxinella* sp.*Nitrile: 4,5-Dibromo-1H-pyrrole-2-carbonitrile, 9CI. 5-Cyano-2,3-dibromopyrrole*

[34649-19-9]

C₅H₂Br₂N₂ 249.892Constit. of *Agelas oroides*. Mp 172-173°.*N-Me: 4,5-Dibromo-1-methyl-1H-pyrrole-2-carboxylic acid, 9CI*

[66067-06-9]

C₆H₃Br₂NO₂ 282.919Constit. of an unidentified orange marine sponge collected in the Marshall Islands. Cryst. (CHCl₃). Mp 153°.*N-Me. Me ester: [1198-71-6]*C₇H₇Br₂NO₂ 296.946Isol. from sponge *Agelas mauritiana*. Mp 87-90°. Poss. artifact.*N-Et: 4,5-Dibromo-1-ethyl-1H-pyrrole-2-carboxylic acid*

[150314-24-2]


C₇H₇Br₂NO₂ 296.946Isol. from a sponge *Axinella* sp. Cryst. Mp 141-143°. λ_{max} 205 (ε 10600); 239 (ε 8500); 275 (ε 12300) (MeOH).*N-2-Propenyl: 1-Allyl-4,5-dibromo-1H-pyrrole-2-carboxylic acid*

[150314-25-3]

C₈H₇Br₂NO₂ 308.957Isol. from a sponge *Axinella* sp. Amorph. cream-coloured solid. Mp 128-129°. λ_{max} 205 (ε 12800); 238 (ε 10600); 273 (ε 14000) (MeOH).Rinkes, I.J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1941, **60**, 303-308 (*ester, synth*)Anderson, H.J. *et al.*, *Can. J. Chem.*, 1965, **43**, 409-414 (*ester, synth*)Hodge, P. *et al.*, *J.C.S.*, 1965, 459-470 (*synth, uv, ir*)Forenza, S. *et al.*, *Chem. Comm.*, 1971, 1129-1130 (*Agelas oroides* *constit*)Chevolot, L. *et al.*, *Heterocycles*, 1977, **7**, 891-894 (*amide*)Schmitz, F.J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 47-53 (*ester, isol, ir, pmr, cmr, ms*)Utkina, N.K. *et al.*, *Khim. Prir. Soedin.*, 1985, 578; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 547-548 (*isol*)Fathi-Afshar, R. *et al.*, *Can. J. Chem.*, 1988, **66**, 45-50 (*N-Me Me ester, isol, pmr, cmr*)Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1989, **52**, 757-761 (*isol*)Barrow, R.A. *et al.*, *Nat. Prod. Lett.*, 1993, **1**, 243-250 (*isol, Axinella derivis*)Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 501-503 (*amide, isol*)Ponasiak, J.A. *et al.*, *Tet. Lett.*, 1996, **37**, 6041-6044 (*synth*)Tsukamoto, S. *et al.*, *Tetrahedron*, 1996, **52**, 8181-8186 (*amide, isol*)Umeyama, A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1433-1434 (*methoxymethylamide*)Koenig, G.M. *et al.*, *Planta Med.*, 1998, **64**, 443-447 (*isol, pmr, cmr*)Tasdemir, D. *et al.*, *Bioorg. Med. Chem.*, 2008, **15**, 6834-6845 (*isol, pmr, cmr*)

4,5-Dibromo-1H-pyrrole-2,3-dicarboxylic acid

D-340


C₆H₃Br₂NO₄ 312.902*3-Nitrile, Me ester: 2,3-Dibromo-4-cyano-5-methoxycarbonylpyrrole*

[866410-40-4]

C₇H₄Br₂N₂O₂ 307.929Constit. of the sponge *Phakellia fusca*.Xu, S. *et al.*, *Tianran Chanwu Yanjiu Yu Kaifa*, 2003, **15**, 393-395; *CA*, **143**, 363646n (*isol*)

3,4-Dibromo-1H-pyrrole-2,5-dione, 9CI D-341

Dibromomaleimide
[1122-10-7]



$C_4HBr_2NO_2$ 254.865

Alkaloid from the sponge *Axinella brevistyla*. Antifungal and cytotoxic agent. Yellow needles (xylene or $CHCl_3$). Mp 230° (226°). λ_{max} 236 (log ϵ 4.1); 243 (log ϵ 4.1); 251 (sh) (log ϵ 3.9); 303 (log ϵ 3.1) (MeOH).

N-Me: 3,4-Dibromo-1-methyl-1H-pyrrole-2,5-dione. 2,3-Dibromo-N-methylmaleimide
[3005-27-4]
 $C_5H_3Br_2NO_2$ 268.892
Pale yellow needles (H_2O). Mp 121° ($114-116^\circ$).

N-Et:
 $C_6H_5Br_2NO_2$ 282.919
Cryst. (EtOH). Mp $93-94^\circ$.

N-Butyl: [181864-30-2]
 $C_8H_9Br_2NO_2$ 310.973
Cryst. (MeOH). Mp $75-77^\circ$.

N-Benzyl: [91026-00-5]
 $C_{11}H_7Br_2NO_2$ 344.99
Mp $117-117.5^\circ$ ($101-103^\circ$).

N-Ph: 3,4-Dibromo-1-phenyl-1H-pyrrole-2,5-dione. 2,3-Dibromo-N-phenylmaleimide
[65833-14-9]
 $C_{10}H_5Br_2NO_2$ 330.963
Yellow solid. Mp $166-168^\circ$.

Khotinsky, E. *et al.*, *Ber.*, 1904, **37**, 2798-2802 (*N-Me*)

Ruggli, P. *et al.*, *Helv. Chim. Acta*, 1920, **3**, 493-514 (*synth*)

Elvidge, J.A. *et al.*, *J.C.S.*, 1957, 2466-2472 (*synth*)

Martin, E.L. *et al.*, *J.O.C.*, 1961, **26**, 2032-2037 (*N-Ph*)

Scharf, H.D. *et al.*, *Chem. Ber.*, 1965, **98**, 764-780 (*N-Me*, *synth*, *ir*, *uv*)

Joyce, R.P. *et al.*, *J.O.C.*, 1987, **52**, 1177-1185 (*N-benzyl*)

Chen, W. *et al.*, *Acta Cryst. C*, 1997, **53**, 631-633 (*N-butyl*, *cryst struct*)


Choi, D.-S. *et al.*, *J.O.C.*, 1998, **63**, 2646-2655 (*N-benzyl*, *N-Ph*)

Tsakamoto, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1576-1578 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Shorunov, S.V. *et al.*, *Russ. J. Org. Chem. (Engl. Transl.)*, 2006, **42**, 1490-1497 (*N-butyl*)

4,5-Dibromo-1H-pyrrole-2-methanol D-342

2,3-Dibromo-5-hydroxymethylpyrrole



$C_5H_5Br_2NO$ 254.909

Me ether: 2,3-Dibromo-5-(methoxy-

methyl)-1H-pyrrole, 9CI

[115502-60-8]


$C_6H_7Br_2NO$ 268.935

Metab. from a marine sponge *Agelas* sp. Extremely unstable.

Tada, H. *et al.*, *Chem. Lett.*, 1988, 803 (*isol*, *pmr*, *cmr*, *ms*, *struct*)

5,6-Dibromotryptamine D-343

5,6-Dibromo-1H-indole-3-ethanamine, 9CI. 3-(2-Aminoethyl)-5,6-dibromoindole [41115-69-9]



$C_{10}H_{10}Br_2N_2$ 318.01

Alkaloid from the Caribbean sponge *Polyfibrospongia maynardii*. Shows *in vitro* but not *in vivo* activity against gram-negative as well as gram-positive bacteria. Mp $110-120^\circ$. λ_{max} 232 (ϵ 29200); 297 (ϵ 3700); 307 (ϵ 3300) (MeOH) (Berdy).

N^b-Me: 5,6-Dibromo-N^b-methyltryptamine. 5,6-Dibromo-N-methyl-1H-indole-3-ethanamine, 9CI [41115-68-8]

$C_{11}H_{12}Br_2N_2$ 332.037
Isol. from *Polyfibrospongia maynardii*. Shows *in vitro* but not *in vivo* activity against gram-negative as well as gram-positive bacteria. Mp $132-134^\circ$. λ_{max} 232 (ϵ 29200); 297 (ϵ 3700); 307 (ϵ 3300) (MeOH) (Berdy).

N^b,N^b-Di-Me: 5,6-Dibromo-N^b,N^b-dimethyltryptamine. 5,6-Dibromo-N,N-dimethyl-1H-indole-3-ethanamine, 9CI [72853-80-6]

$C_{12}H_{14}Br_2N_2$ 346.064
Constit. of the marine sponges *Smenospongia echina*, *Smenospongia aurea* and *Verongula gigantea*. Shows antimicrobial props. Cryst. (MeOH aq.). Sol. MeOH, EtOAc; poorly sol. H_2O . Mp $113-115^\circ$. λ_{max} 230 (ϵ 40100); 285 (ϵ 4900); 300 (ϵ 3300) (MeOH) (Derrep). λ_{max} 207 (ϵ 9600); 232 (ϵ 30800); 291 (ϵ 6300) (EtOH) (Berdy).


Van Lear, G.E. *et al.*, *Tet. Lett.*, 1973, 299-300 (*uv*, *pmr*, *ms*, *N^b-Me*, *struct*)

Djura, P. *et al.*, *J.O.C.*, 1980, **45**, 1435-1441 (5,6-Dibromo-N^b,N^b-dimethyltryptamine)

Tymiak, A.A. *et al.*, *Tetrahedron*, 1985, **41**, 1039-1047 (5,6-Dibromo-N^b,N^b-dimethyltryptamine)

5,5''-Dicapsaicin D-344

[134386-10-0]



$C_{36}H_{52}N_2O_6$ 608.817

Oxidn. prod. of Capsaicin, C-100. Alkaloid from *Capsicum annum* var. *annuum*. Antioxidant.

6'',7''-Dihydro: 6'',7''-Dihydro-5,5''-dicapsaicin

$C_{36}H_{54}N_2O_6$ 610.832

Constit. of the fruit of *Capsicum annum*. Antioxidant. Light yellow oil.

Bernal, M.A. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3085-3089 (*synth*, *ms*)


Henderson, D.E. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 2563-2570 (*synth*, *ms*)

Ochi, T. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1094-1096 (*isol*, *pmr*, *cmr*)

Martínez-Juárez, V.M. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 972-979 (*isol*, *pmr*, *cmr*)

Dicarprine C D-345

2-Amino-5,7,9-dodecatrien-3-ol [67951-12-6]



$C_{12}H_{21}NO$ 195.304

Alkaloid from *Dicarpellum pronyensis* (preferred genus name *Salacia*) (Celastraceae). Noncryst.

N-Me: Dicarprine B

[67951-11-5]

$C_{13}H_{23}NO$ 209.331

Alkaloid from *Dicarpellum pronyensis* (Celastraceae). Noncryst.

O,N-Di-Me: Dicarprine A

[67951-10-4]


$C_{14}H_{25}NO$ 223.358

Main alkaloid from *Dicarpellum pronyensis* (Celastraceae). Noncryst. Mp $119-120^\circ$ as tartrate. $[\alpha]_D^{+11}$ (c, 1 in MeOH) (tartrate).

Adéoti, B. *et al.*, *Phytochemistry*, 1978, **17**, 831 (*isol*, *uv*, *ms*, *pmr*, *struct*)

Dicentrine D-346

9,10-Dimethoxy-1,2-methylenedioxyaporphine. N,O-Dimethylactinodaphnine. *Ex-imine*†



$C_{20}H_{21}NO_4$ 339.39

Log P 2.86 (uncertain value) (calc).

(*R*)-form [28832-07-7]

Alkaloid from a Brazilian *Duguetia* sp. (Annonaceae). Shows antineoplastic activity. Mp 169° . $[\alpha]_D -53$ (c, 1 in $CHCl_3$).

N-De-Me: (-)-Nordicentrine

[151601-88-6]

$C_{19}H_{19}NO_4$ 325.363

Alkaloid from tubers of *Stephania pierrei* (Menispermaceae). Mp 248°

dec. $[\alpha]_D^{20}$ -34 (c, 0.2 in MeOH). λ_{\max} 220 (ϵ 26300); 231 (ϵ 23440); 272 (ϵ 10715); 281 (ϵ 14790) (MeOH) (Berdy).

O⁹-De-Me: (-)-Cassythicine

[104641-05-6]
C₁₉H₁₉NO₄ 325.363
Alkaloid from the leaves of *Annona glabra* (Annonaceae) and from *Stephania epigeae*. Cryst. (MeOH). Mp 210-211°. $[\alpha]_D$ -61 (c, 0.58 in CHCl₃). λ_{\max} 221 (ϵ 25700); 282 (ϵ 4677); 309 (ϵ 12580) (MeOH) (Berdy).

O¹⁰-De-Me: 10-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine. Phanostenine

[25368-02-9]
C₁₉H₁₉NO₄ 325.363
Alkaloid from *Stephania sasakii* (Menispermaceae). Mp 126-128°. $[\alpha]_D$ -39 (c, 0.48 in CHCl₃). The CAS reg. no. refers to the (S)-enantiomer but the alkaloid prob. has the (R)-config. λ_{\max} 234 (ϵ 14450); 273 (ϵ 19050); 318 (ϵ 3630) (MeOH) (Berdy).

(S)-form [517-66-8]

Alkaloid from a variety of genera in the Lauraceae (*Cassytha*, *Laurus*, *Lindera*, *Ocotea*, *Litsea*), Menispermaceae (*Cocculus*, *Cissampelos*, *Stephania*), and Papaveraceae (*Dicentra*, *Glaucium*). Shows sedative and analgesic props. Mp 168-169°. $[\alpha]_D$ +57 (c, 1 in EtOH). λ_{\max} 281 (log ϵ 4.18); 305 (log ϵ 4.23) (95% EtOH).

►CE0455000

Hydrochloride: Mp 169°.

N-Me: N-Methyldicentrinium

[209323-82-0]
C₂₁H₂₄NO₅⁺ 354.425
Alkaloid from *Dicentra peregrina*. Mp 224° (as iodide) (synthetic). Counterion not specified for natural compd.

N-De-Me: 9,10-Dimethoxy-1,2-methylenedioxyaporphine. (+)-Nordicentrine

[25394-59-6]
C₁₉H₁₉NO₄ 325.363
Alkaloid from the trunk and leaves of *Lindera oldhamii*, the leaves of *Litsea salicifolia*, and the stem bark of *Guatteria scandens* (Lauraceae, Annonaceae). Mp 254-255° dec. $[\alpha]_D^{25}$ +31 (c, 0.65 in MeOH).

O⁹-De-Me: 9-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine. Cassythicine. N-Methylactinodaphnine

[5890-28-8]
C₁₉H₁₉NO₄ 325.363
Alkaloid from *Cassytha melantha*, *Cassytha glabella*, *Cassytha americana* (*Cassytha filiformis*), *Hernandia cordigera*, *Laurus nobilis* (bay laurel), *Litsea kawakamii*, *Litsea laurifolia*, *Litsea glutinosa* var. *glabraria*, *Neolitsea sericea* and *Ocotea brachybotra* (Lauraceae, Hernandiaceae). Shows antimicrobial activity. Cryst. + 1CHCl₃ (CHCl₃). Mp 210-212°. $[\alpha]_D$ +62 (c, 0.43 in CHCl₃). λ_{\max} 283 (log ϵ 4.23); 307 (log ϵ 4.27) (95% EtOH).

O⁹-De-Me, N-de-Me: 9-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine. Actinodaphnine

[517-69-1]
C₁₈H₁₇NO₄ 311.337
Alkaloid from a variety of genera in the Hernandiaceae (*Hernandia*, *Illigera*), Lauraceae (*Actinodaphne*, *Cassytha*, *Laurus*, *Litsea*, *Neolitsea*) and Annonaceae (*Guatteria*). Shows antimicrobial activity. Needles (EtOH). Mp 210-211°. $[\alpha]_D$ +39 (c, 1 in CHCl₃).

O⁹-De-Me, N-de-Me, hydrochloride: Mp 280-281° dec. $[\alpha]_D^{20}$ +8.75 (H₂O).

O⁹-De-Me, N-de-Me, N-methoxycarbonyl: Cathafliline. N-Methoxycarbonylactinodaphnine

[196099-31-7]
C₂₀H₁₉NO₆ 369.373
Alkaloid from *Cassytha filiformis* (Lauraceae). Amorph. brown solid (CHCl₃). Mp 118-120°. $[\alpha]_D$ +142.4 (c, 0.1 in CHCl₃). λ_{\max} 236 ; 286 ; 305 (EtOH).

O¹⁰-De-Me, N-de-Me: 10-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine. Litseferine. Litseglutine A

[60142-18-9]
C₁₈H₁₇NO₄ 311.337
Alkaloid from *Litsea glutinosa* and *Litsea sebifera*. Cryst. (Py). Mp 258°. $[\alpha]_D^{24}$ +36.6 (c, 0.13 in MeOH). λ_{\max} 218 ; 282 ; 306 (MeOH).

O¹⁰-De-Me, N-de-Me, N,O-di-Ac: Powder. Mp 254°.
6a,7-Didehydro: see Dehydrodicentrine, D-156

4S-Hydroxy: 4-Hydroxydicentrine

[72170-12-8]
C₂₀H₂₁NO₅ 355.39
Alkaloid from leaves of *Ocotea minarum* (Lauraceae). Mp 210°. $[\alpha]_D$ +60 (c, 0.5 in CHCl₃).

7-Hydroxy: see Duguetine, D-948

(±)-form [26110-43-0]

Mp 178-179°. **Hydrochloride:** Mp 263-265° dec. **N-Me:** Mp 228-229° (as iodide). **O⁹-De-Me:** [38849-64-8] Synthetic. Shows antimicrobial activity. Cryst. (MeOH). Mp 117° (111-113°).

O⁹-De-Me, N-de-Me: [58072-86-9] Synthetic. Brown needles (Me₂CO). Mp 195-198°.

O¹⁰-De-Me: Synthetic. Plates (MeOH). Mp 209-210°.

O¹⁰-De-Me, hydrochloride: Needles (EtOH). Mp 255° dec.


Asahina, Y. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1909, **247**, 201-212 (isol)
Haworth, R.D. et al., *J.C.S.*, 1925, **127**, 2018-2023; 1926, **128**, 29 (synth, resoln)
Ghose, T.P. et al., *Helv. Chim. Acta*, 1934, **17**, 919-930 (*Actinodaphnine*)
Tomita, M. et al., *Yakugaku Zasshi*, 1957, **77**, 1011-1015; 1015-1018 (*Cassythicine*, *Phanostenine*, *synth*)
Craig, J.C. et al., *Tetrahedron*, 1965, **21**, 395-399 (ord, uv)
Johns, S.R. et al., *Aust. J. Chem.*, 1966, **19**, 2339-2345 (*Actinodaphnine*, *Cassythicine*)
Nakasato, T. et al., *Yakugaku Zasshi*, 1966, **86**, 129-134 (*Cassythicine*)
Cava, M.P. et al., *J.O.C.*, 1968, **33**, 2443-2446 (*Actinodaphnine*, *Cassythicine*, *uv*)

Kunimoto, J. et al., *Yakugaku Zasshi*, 1969, **89**, 1691-1695 (*Phanostenine*)
Yang, T.H. et al., *J. Chin. Chem. Soc. (Taipei)*, 1971, **18**, 133-136; *CA*, **77**, 16567r ((-)-*Cassythicine*)
Tewari, S. et al., *Phytochemistry*, 1972, **11**, 1149-1152 (*Cassythicine*, *isol*, *pmr*, *ms*)
Lu, S.T. et al., *Yakugaku Zasshi*, 1972, **92**, 910-917 (*isol*, *pmr*, *ir*)
Cava, M.P. et al., *Tetrahedron*, 1973, **29**, 2245-2249 (*synth*)
Premila, M.S. et al., *Indian J. Chem.*, 1975, **13**, 945-946 (*Actinodaphnine*, *synth*)
Sivakumaran, M. et al., *Indian J. Chem., Sect. B*, 1976, **14**, 150 (*Litseferine*)
Chen, I.S. et al., *J. Chin. Chem. Soc. (Taipei)*, 1977, **24**, 41-44; *CA*, **87**, 35908a ((+)-*Nordicentrine*)
O'Brien, J.P. et al., *Heterocycles*, 1978, **11**, 347-350 (*synth*)
Vecchietti, V. et al., *Farmaco, Ed. Sci.*, 1979, **34**, 829-840 (*4-Hydroxydicentrine*)
Rastogi, R.C. et al., *Phytochemistry*, 1980, **19**, 998-999 ((+)-*Nordicentrine*)
Ringdahl, B. et al., *J. Nat. Prod.*, 1981, **44**, 80-85 (*cd*)
Hocquemiller, R. et al., *J. Nat. Prod.*, 1983, **46**, 335 ((+)-*Nordicentrine*)
Hara, H. et al., *Chem. Pharm. Bull.*, 1986, **34**, 1946-1949 (*Dicentrine*, *Cassythicine*, *synth*, *pmr*)
Soicke, H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 149-152 ((±)-*Cassythicine*)
Tsai, I.L. et al., *Kaohsiung J. Med. Sci.*, 1989, **5**, 131-145 (*Actinodaphnine*, *activity*)
Likhitwitayawuid, K. et al., *J. Nat. Prod.*, 1993, **56**, 1468-1478 ((-)-*Nordicentrine*, (-)-*Cassythicine*)
Wu, Y.C. et al., *Phytochemistry*, 1997, **46**, 181-184 (*Cathafliline*)
Konishi, T. et al., *Nat. Med. (Tokyo)*, 1998, **52**, 47-53 (*N-Methyldicentrinium*)
Ma, W.G. et al., *Fitoterapia*, 2000, **71**, 527-534 (*cmr*)
Stévigny, C. et al., *Planta Med.*, 2002, **68**, 1042-1044 (*Actinodaphnine*, *pmr*, *cmr*)
Yang, J.-H. et al., *Helv. Chim. Acta*, 2005, **88**, 2523-2526 (*Litseglutine A*)

Dicentrinone

D-347

Oxodicentrine
[16408-78-9]



C₁₉H₁₃NO₅ 335.315
Alkaloid from *Ocotea macropoda* and *Lindera oldhamii*, also from the genera *Guatteria* and *Glaucium* (Lauraceae, Annonaceae, Papaveraceae). Topoisomerase I inhibitor. Yellow needles (CHCl₃/EtOH). Mp 300° dec. λ_{\max} 213 (log ϵ 4.59); 254 (log ϵ 4.75); 271 (log ϵ 4.47); 311 (log ϵ 4); 349 (log ϵ 4.04); 400 (log ϵ 3.97) (EtOH).

O⁹-De-Me: Machigline
[87264-30-0]
C₁₈H₁₁NO₅ 321.289
Alkaloid from leaves of *Machilus glaucescens* (Lauraceae). Red needles

(CHCl₃/MeOH). Mp 315° dec.Cava, M.P. *et al.*, *Tetrahedron*, 1971, **27**, 2639 (isol, uv, ir, pmr)Lu, S.-T. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 910 (isol)Talapatra, B. *et al.*, *J. Indian Chem. Soc.*, 1982, **59**, 1364 (*Machigline*)Wu, Y.C. *et al.*, *Planta Med.*, 1989, **55**, 163-165 (isol)Zhou, B.-N. *et al.*, *J. Nat. Prod.*, 2000, **63**, 217-221 (isol, pmr, cmr, activity)**Dichloroacetic acid, 9CI** D-348*Dichloroethanoic acid*

[79-43-6]


Cl₂CHCOOHC₂H₂Cl₂O₂ 128.942Constit. of the red alga *Asparagopsis taxiformis*. Has fungicidal props. Corrosive liq. Misc. H₂O, EtOH. d₄²⁰ 1.56. Mp 5-6° (11°). Bp 194° Bp₂₀ 102°. n_D²² 1.4659.► Fl. p. >66°. Corrosive and irritating to all tissues. LD₅₀ (rbt, skn) 510 mg/kg. AG6125000*Amide: Dichloroacetamide*

[683-72-7]


C₂H₃Cl₂NO 127.957Constit. of the red alga *Marginisporium aberrans*. Preservative. Shows antimicrobial props. Cryst. Mp 98°. Bp₇₄₅ 234°. Steam-volatile, subl.

► AB6475000

[13425-80-4]

Org. Synth., Coll. Vol., 3, 1955, 260 (*amide*)
Ohta, K. *et al.*, *Phytochemistry*, 1977, **16**, 1085-1086 (*amide, isol*)
Pellegata, R. *et al.*, *Synthesis*, 1985, 517 (*amide*)**2,10-Dichloro-6,11-cyclo-3(15),7(14)-farnesadien-1-yl carbonimidic dichloride** D-349*2-Chloro-5-(3-chloro-2,2-dimethyl-6-methylenecyclohexyl)-3-methylenepentyl carbonimidic dichloride*
[321657-93-6]C₁₆H₂₃Cl₄N 371.175Constit. of *Stylotella aurantium*. Oil. [α]_D²⁵ +4.5 (c, 0.2 in CHCl₃). λ_{max} 205 (log ε 3.6) (MeOH).Musman, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 111-113 (isol, pmr, cmr)**3,8-Dichloro-11,12-dihydroindolo[2,3-a]carbazole** D-350*Tjipanazole D*

[139083-25-3]

C₁₈H₁₀Cl₂N₂ 325.196Alkaloid from blue-green alga *Tolypothrix tjipanansensis* and *Fischerella ambiguu*. λ_{max} 259 (ε 63100); 268 (sh) (ε 54100); 291 (ε 25500); 331 (ε 31000); 345 (sh) (ε 7300); 366 (ε 4030) (MeOH) (Derep).*N-β-D-Xylopyranosyl: Tjipanazole B*

[139083-21-9]

C₂₃H₁₈Cl₂N₂O₄ 457.312Alkaloid from *Tolypothrix tjipanansensis*. [α]_D -4.9 (c, 1.03 in CHCl₃). [α]_D +10.5 (c, 0.95 in CHCl₃/MeOH 1:1). λ_{max} 261 (ε 61000); 294 (ε 23200); 333 (ε 31300); 354 (ε 7340); 371 (ε 4800) (MeOH) (Derep). λ_{max} 259 (ε 59400); 292 (ε 24800); 330 (ε 30400); 349 (ε 8140); 366 (ε 4850) (MeOH) (Berdy).*N-(6-Deoxy-β-D-gulopyranosyl): Tjipanazole A1*


[139083-13-9]

C₂₄H₂₀Cl₂N₂O₄ 471.338Major alkaloid from the blue-green alga *Tolypothrix tjipanansensis*. Exhibits antifungal activity. [α]_D +9.1 (c, 1.0 in CHCl₃). λ_{max} 261 (ε 61000); 294 (ε 23200); 333 (ε 31300); 354 (ε 7340); 371 (ε 4800) (MeOH) (Derep).*N-α-L-Rhamnopyranosyl: Tjipanazole A2*

[139083-14-0]

C₂₄H₂₀Cl₂N₂O₄ 471.338From *Tolypothrix tjipanansensis*. Exhibits antifungal activity. [α]_D +25.12 (c, 1.0 in CHCl₃). λ_{max} 261 (ε 61000); 294 (ε 23200); 333 (ε 31300); 354 (ε 7340); 371 (ε 4800) (MeOH) (Derep).*N-β-D-Glucopyranosyl: Tjipanazole E*



[139083-22-0]

C₂₄H₂₀Cl₂N₂O₅ 487.338Minor alkaloid from *Tolypothrix tjipanansensis*. [α]_D +63.8 (c, 1.0 in CHCl₃/MeOH 1:1). λ_{max} 261 (ε 61000); 294 (ε 23200); 333 (ε 31300); 354 (ε 7340); 371 (ε 4800) (MeOH) (Derep). λ_{max} 259 (ε 54400); 293 (ε 21700); 332 (ε 26900); 350 (ε 7360); 368 (ε 4460) (MeOH) (Berdy).Bonjouklian, R. *et al.*, *Tetrahedron*, 1991, **47**, 7739 (isol, uv, pmr, cmr, struct)Kuethe, J.T. *et al.*, *Org. Lett.*, 2003, **5**, 3721-3723 (*synth*)**7,8-Dichloro-1-hydroxy-9-methyl-β-carboline** D-351*7,8-Dichloro-9-methyl-9H-pyrido[3,4-b]indol-1-ol*, *7,8-Dichloro-2,9-dihydro-9-methylpyrido[3,4-b]indol-1-one*. *Bauerine C*
[156312-11-7]C₁₂H₈Cl₂N₂O 267.113Alkaloid from terrestrial blue-green alga *Dichothrix baueriana*. Solid (EtOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 254-258° dec. λ_{max} 250 (ε 43900); 290 (ε 4370); 301 (ε 2740); 334 (ε 4040); 347 (ε 6000); 362 (ε 4630) (MeOH) (Berdy).Larsen, L.K. *et al.*, *J. Nat. Prod.*, 1994, **57**,

419-421 (isol, uv, pmr, cmr, struct)

Pohl, B. *et al.*, *Synth. Commun.*, 2007, **37**, 1273-1280 (*synth, pmr, cmr*)Lingam, Y. *et al.*, *Synth. Commun.*, 2007, **37**, 4313-4318 (*synth, pmr*)**1,2-Dichloro-8-hydroxy-6-methyl-7-propyl-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, 9CI** D-352


[208707-52-2]

C₁₅H₁₃Cl₂N₂O₃ 326.178Prod. by *Streptomyces rimosus*.*Pat. Coop. Treaty (WIPO)*, 1998, 98 25 931; *CA*, **129**, 52003e**6,6'-Dichloroindigotin** D-353C₁₆H₈Cl₂N₂O₂ 331.157*N-(4-Amino-4,6-dideoxy-β-D-glucopyranosyl): Akashin A*
[428516-77-2]C₂₂H₁₉Cl₂N₃O₅ 476.315Prod. by *Streptomyces* sp. GW 48/1497. Antitumour agent. [α]_D²⁵ +2560 (c, 0.00125 in MeOH). Config. of sugar residue not detd. λ_{max} 241 (log ε 4.33); 290 (log ε 4.23); 619 (log ε 4.01) (MeOH).*N-(4-Acetamido-4,6-dideoxy-β-D-glucopyranosyl): Akashin B*

[428517-17-3]

C₂₄H₂₁Cl₂N₃O₆ 518.352Prod. by *Streptomyces* sp. GW 48/1497. [α]_D²⁵ +2840 (c, 0.0014 in MeOH). λ_{max} 244 (log ε 4.34); 291 (log ε 4.28); 618 (log ε 4.03) (MeOH).Maskey, R.P. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 597-599 (isol, uv, pmr, cmr, ms)Maskey, R.P. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 137-142 (*struct*)**Dichlorolissoclimide** D-354

[138935-82-7]

C₂₀H₂₉Cl₂N₂O₄ 418.359Constit. of tunicate *Lissoclinum voeltzkowi*. Tyrosine metab. and 5-aminolevu-

linic acid dehydrase inhibitor. Immunosuppressant. Mp 210°. $[\alpha]_D^{20}$ -20. $[\alpha]_D^{20} +30$ (c, 0.2 in MeOH). No comment made on conflicting optical rotations. λ_{\max} 242 (ε 76) (EtOH).

7-Ac: **Haterumaimide A**

[335155-25-4]
C₂₂H₃₁Cl₂NO₅ 460.396

Isol. from a *Lissoclinium* sp. Cytotoxic agent. $[\alpha]_D^{29} +31.3$ (c, 0.13 in MeOH).

3-Dechloro: **Chlorolissoclimide**

[148717-91-3]
C₂₀H₃₀ClNO₄ 383.914

Constit. of *Lissoclinium voeltzkowi*. Highly cytotoxic. Glass.

3-Dechloro, 7-Ac: **Haterumaimide N**

[913545-22-9]
C₂₂H₃₂ClNO₅ 425.951

Constit. of a *Lissoclinium* sp. Oil. $[\alpha]_D^{32} +59.7$ (c, 0.79 in MeOH). λ_{\max} 215 (log ε 3.6) (MeOH).

3-Dechloro, 12-deoxy: **Haterumaimide P**

[913545-24-1]
C₂₀H₃₀ClNO₃ 367.915

Constit. of a *Lissoclinium* sp. Oil. $[\alpha]_D^{29} +68.8$ (c, 0.11 in MeOH). λ_{\max} 210 (log ε 3.6) (MeOH).

3-Dechloro, 12-deoxy, 12,13-didehydro (E-), 7-Ac: **Haterumaimide O**

[913545-23-0]
C₂₂H₃₀ClNO₄ 407.936

Constit. of a *Lissoclinium* sp. Oil. $[\alpha]_D^{28} +16$ (c, 0.08 in MeOH). λ_{\max} 235 (log ε 3.9) (MeOH).

2,3-Didechloro: **Haterumaimide Q**

[913545-25-2]
C₂₀H₃₁NO₄ 349.469

Constit. of a *Lissoclinium* sp. Oil. $[\alpha]_D^{28} +36$ (c, 0.19 in CHCl₃). λ_{\max} 204 (log ε 3.6) (MeOH).

Malochet-Grivois, C. et al., *Tet. Lett.*, 1991, **32**, 6701-6702 (*isol, pmr, cmr*)

Biard, J.-F. et al., *Nat. Prod. Lett.*, 1994, **4**, 43-50 (*Chlorolissoclimide*)


Toupet, L. et al., *J. Nat. Prod.*, 1996, **59**, 1203-1204 (*cryst struct*)

Uddin, M.J. et al., *Heterocycles*, 2001, **54**, 1039-1048 (*Haterumaimide A*)

Uddin, J. et al., *Bioorg. Med. Chem.*, 2006, **14**, 6954-6961 (*Haterumaimides N-Q*)

7,8-Dichloro-9-methyl-β-carboline **D-355**

7,8-Dichloro-9-methylpyrido[3,4-b]indole. **Bauerine B** [156312-10-6]



C₁₂H₈Cl₂N₂ 251.114


Alkaloid from terrestrial blue-green alga *Dichothrix baueriana*. Shows antiviral activity. Solid (MeOH/hexane). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 163-164°. λ_{\max} 246 (ε 38300); 281 (ε 8200); 292 (ε 12300); 345 (ε 4500); 359 (ε 5500) (MeOH) (Berdy).

Larsen, L.K. et al., *J. Nat. Prod.*, 1994, **57**,

419-421 (*isol, uv, pmr, cmr, struct*)
Rocca, P. et al., *Synth. Commun.*, 1995, **25**, 2901-2908 (*synth*)
Pohl, B. et al., *Synth. Commun.*, 2007, **37**, 1273-1280 (*synth, pmr, cmr*)

2,3-Dichloro-4-nitro-1H-pyrrole **D-356**

Pyrrrolomycin A. SF 2080A. Antibiotic SF 2080A [79763-01-2]



C₄H₂Cl₂N₂O₂ 180.977

Pyrrrole antibiotic. Isol. from *Actinosporangium vitaminophilum*. Antifungal agent. Yellow cryst. (C₆H₆). Sol. MeOH, dioxan, Me₂CO; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 211-213°. Log P 2.16 (calc). λ_{\max} 270 (ε 7830); 318 (ε 3020) (MeOH/HCl) (Derrep). λ_{\max} 323 (ε 11300) (MeOH/NaOH) (Derrep). λ_{\max} 268 (ε 7400); 269 (E1%/1cm 490); 318 (E1%/1cm 170); 320 (E1%/1cm 230); 323 (ε 11400) (MeOH) (Berdy).
▶ LD₅₀ (mus, ipr) 10 - 30 mg/kg. UX9459000

Fr. Pat., 1981, 2 472 611; *CA*, **96**, 4942
Ezaki, N. et al., *J. Antibiot.*, 1981, **34**, 1363-1365; 1569 (*isol, struct, synth, pharmacol*)
Ratnayake, A.S. et al., *J. Nat. Prod.*, 2008, **71**, 1923-1926 (*biosynth*)

1,15-Dichloro-1,14-pentadecadiene-3,12-diyn-8-amine **D-357**

8-Amino-1,15-dichloro-1,14-pentadecadiene-3,12-diyne
ClCH=CHC≡C(CH₂)₃
CH(NH₂) (CH₂)₃C≡CCH=CHCl
C₁₅H₁₉Cl₂N 284.227

(1E,14E)-form

N-Ac: 8-Acetamido-1,15-dichloro-1,14-pentadecadiene-3,12-diyne. **Taveuniamide H**

C₁₇H₂₁Cl₂NO 326.264

Isol. from a mixed assemblage of *Lyngbya majuscula* and *Schizothrix* sp. λ_{\max} 238 (ε 10300) (EtOH).

1-Chloro, N-Ac: 8-Acetamido-1,1,15-trichloro-1,14-pentadecadiene-3,12-diyne. **Taveuniamide I**

C₁₇H₂₀Cl₃NO 360.709

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25} -2$ (c, 0.4 in CHCl₃). λ_{\max} 242 (ε 37500) (EtOH).

1-Chloro, 1,2,3,3,4,4-hexahydro, N-Ac: 8-Acetamido-1,15,15-trichloro-1-pentadecen-3-yne. **Taveuniamide F**

C₁₇H₂₆Cl₃NO 366.757

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25} -5.2$ (c, 0.4 in CHCl₃). λ_{\max} 238 (ε 9100) (EtOH).

1,15-Dichloro, N-Ac: 8-Acetamido-1,1,15,15-tetrachloro-1,14-pentadecadiene-3,12-diyne. **Taveuniamide J**

C₁₇H₁₉Cl₄NO 395.154

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. λ_{\max} 244 (ε 37700) (EtOH).

1,15-Dichloro, 1,2-dihydro, N-Ac: 8-Acetamido-1,1,15,15-tetrachloro-1-pentadecen-3,12-diyne. **Taveuniamide G**

C₁₇H₂₁Cl₄NO 397.17

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25} +0.3$ (c, 0.04 in CHCl₃). λ_{\max} 244 (ε 21100) (EtOH).

(1E,14Z)-form

3Z,4-Dihydro, N-Ac: 8-Acetamido-1,15-dichloro-1,3,14-pentadecatrien-12-yne. **Taveuniamide K**

C₁₇H₂₃Cl₂NO 328.28


Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25} +2.4$ (c, 0.1 in CHCl₃). λ_{\max} 242 (ε 18200) (EtOH).

Williamson, R.T. et al., *Tetrahedron*, 2004, **60**, 7025-7033 (*isol, pmr, cmr*)

Dichomine

D-358

[89647-74-5]



C₁₉H₂₄N₂O 296.411


Unusual indole alkaloid. Can theoretically be derived from a Secodine precursor by Diels-Alder type cyclisation. The VX5400 classification given here is provisional. Alkaloid from the leaves of *Tabernaemontana dichotoma* and leaves and twigs of *Tabernaemontana eglanulosa* (Apocynaceae).

Perera, P. et al., *Planta Med.*, 1983, **49**, 232 (*uv, pmr, cmr, ms, cd, struct*)

Dichotine, 9CI

D-359

[27530-76-3]



C₂₂H₂₆N₂O₆ 414.457

Related to Condylocarpan alkaloids but with an additional C₂ unit of unknown origin. Alkaloid from the bark of *Vallesia dichotoma* (Apocynaceae). Cryst. (Me₂CO). Mp 211-213°. $[\alpha]_D +88$ (CHCl₃). pK_a 7.2 (MeCN).

Ac: [29474-87-1]

Cryst. (Me₂CO). Mp 247-248°.

11-Methoxy: **11-Methoxydichotine** [29537-51-7]

C₂₃H₂₈N₂O₇ 444.483Alkaloid from *Vallesia dichotoma* bark (Apocynaceae). Cryst. (Me₂CO). Mp 120-121°. [α]_D²⁰ +11 (CHCl₃). pK_a 7.2 (MeCN).


11-Methoxy, Ac: [29474-86-0]

Cryst. (Me₂CO). Mp 189-190°.

[27530-77-4]

Ling, N.C. et al., *J.A.C.S.*, 1970, **92**, 222; 6019 (isol, uv, pmr, ms, struct)**Dichotomide II****D-360**

[755036-57-8]

C₁₈H₁₇N₃O₄ 339.35Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. [α]_D¹⁹ +7.2 (c, 0.3 in CHCl₃). λ_{max} 226 (log ε 4.02); 284 (log ε 4.07); 324 (log ε 3.85) (MeOH).Sun, B. et al., *J. Nat. Prod.*, 2004, **67**, 1464-1469 (isol, pmr, cmr, ms)Omura, K. et al., *Chem. Pharm. Bull.*, 2008, **56**, 237-238 (struct)**Dichroidine****D-361**C₁₈H₂₅N₃O₃ 331.414Struct. unknown. Alkaloid from *Dichroa febrifuga* (Hydrangeaceae). Mp 213°.Jang, C.S. et al., *Nature (London)*, 1948, **161**, 400-401**N,N'-Dicinnamoylputrescine****D-362**Ph³CH=CHCO¹NH(CH₂)₄NHCOC-4NHCOCH=CHPhC₂₂H₂₄N₂O₂ 348.444**(E,E)-form**

NSC 107212

[37946-56-8]

Prod. by the mushroom *Pholiota spumosa*. Cryst. (EtOAc/MeOH). Mp 249-253° (247-248°).4'',4''''-Dihydroxy: N,N'-1,4-Butanediyl-bis[3-(4-hydroxyphenyl)-2-propenamide], 9CI. N,N'-Bis(4-hydroxycinnamoyl)-1,4-butanediamine. N¹,N⁴-Di-p-coumaroylputrescine [37946-59-1]C₂₂H₂₄N₂O₄ 380.443Alkaloid from *Dianthus caryophyllus*, *Helianthus annuus* (sunflower), *Nicotiana tabacum*, *Pyrus communis* (pear), *Rubus idaeus* (raspberry) and *Vicia faba*. Also isol. from corn bran (*Zea mays*). Antioxidant. λ_{max} 220; 229; 294; 310 (MeOH).3''',3''''',4'',4''''-Tetrahydroxy: N,N'-Bis(3,4-dihydroxycinnamoyl)-1,4-butanediamine. N¹,N⁴-Dicaffeoylputrescine [60422-23-3]C₂₂H₂₄N₂O₆ 412.441Alkaloid from *Helianthus annuus* (sunflower), *Pyrus communis* (pear), *Nicotiana tabacum* and *Salix* spp.3''',3''''-Methoxy, 4'',4''''-dihydroxy: N¹-p-Coumaroyl-N⁴-feruloylputrescine [380302-96-5]C₂₃H₂₆N₂O₅ 410.469Isol. from corn bran (*Zea mays*).Powder. λ_{max} 220; 229; 294; 310 (MeOH).3''',3''''-Dimethoxy, 4'',4''''-dihydroxy: N,N'-Bis(4-hydroxy-3-methoxycinnamoyl)-1,4-butanediamine. N¹,N⁴-Diferuloylputrescine. Terrestribisamide [42369-86-8]


[91000-13-4]

C₂₄H₂₈N₂O₆ 440.495Alkaloid from *Ananas comosus* (pineapple), *Triticum vulgare*, *Gomphrena globosa*, *Dianthus caryophyllus*, *Tribulus terrestris*, *Vicia faba*, *Lycopersicon esculentum* (tomato), *Nandina domestica*, *Petunia* sp. and *Nicotiana tabacum*. Also isol. from corn bran (*Zea mays*).

Lipoxygenase inhibitor. Antioxidant. Mp 143-145°.

3''',3''''',5''',5''''-Tetramethoxy, 4'',4''''-dihydroxy: N,N'-Bis(4-hydroxy-3,5-dimethoxycinnamoyl)-1,4-butanediamine. N¹,N⁴-Disinapoylputrescine [70185-58-9]C₂₆H₃₂N₂O₈ 500.547Alkaloid from *Lilium* sp.2',3'-Dihydro, N¹-Ac: N¹-Acetyl-N²-cinnamoyl-N¹-dihydrocinnamoylputrescine. Edulimide [217481-11-3]C₂₄H₂₈N₂O₃ 392.497Alkaloid from *Aglaia edulis*. λ_{max} 282 (log ε 3.91) (MeOH).Martin-Tanguy, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1973, **276**, 1433-1435 (*Dicoumaroylputrescine*, *Diferuloylputrescine*)Cabanne, F. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1976, **282**, 1959-1962 (*Dicaffeoylputrescine*)Martin-Tanguy, J. et al., *Phytochemistry*, 1978, **17**, 1927-1928 (occur)Brader, G. et al., *J. Nat. Prod.*, 1998, **61**, 1482-1490 (*Edulimide*)Wu, T.S. et al., *Phytochemistry*, 1999, **50**, 1411-1415 (*Terrestribisamide*)Clericuzio, M. et al., *Croat. Chem. Acta*, 2004, **77**, 605-611 (isol, cryst struct)Choi, S.W. et al., *J. Agric. Food Chem.*, 2007, **55**, 3920-3925 (corn bran constits)**Dicrotaline****D-363**

13-Hydroxy-17,19,20-trinorcrotalanan-11,15-dione, 9CI [480-87-5]

C₁₄H₁₉NO₅ 281.308

Cyclic ester of Retronecine in T-188 with 3-Hydroxy-3-methylpentanedioic acid.

Alkaloid from *Crotalaria dura* and *Crotalaria globifera* (Fabaceae). Mp 170° (dec.). Unstable.


▶ Carcinogenic.

Hydrochloride: Mp 258-260°. Unstable.Ac: **Acetyldicrotaline**


[88205-13-4]

C₁₆H₂₁NO₆ 323.345Minor alkaloid from leaves of *Crotalaria lachnosema* (Fabaceae).Marais, J.S.C. et al., *CA*, 1945, **39**, 4116 (isol) Adams, R. et al., *J.A.C.S.*, 1953, **75**, 2377 (struct)Brown, K. et al., *J.C.S. Perkin 1*, 1982, 2079 (synth)Niwa, H. et al., *Bull. Chem. Soc. Jpn.*, 1988, **61**, 3017 (synth)Mattocks, A.R. et al., *Phytochemistry*, 1988, **27**, 3289 (*Acetyldicrotaline*)Rycroft, D.S. et al., *Magn. Reson. Chem.*, (Spec. Issue), 1992, **30**, S42 (pmr, cmr, config, *Acetyldicrotaline*)Sax, N.I. et al., *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 568**Dictyodendrine A****D-364**

[510709-68-9 (Na salt)]


C₄₃H₃₄N₂O₁₁S 786.814Alkaloid from the sponge *Dictyodendrilla verongiformis*. Telomerase inhibitor.Amorph. red solid (as Na salt). [α]_D¹⁹ -4.6 (c, 0.01 in MeOH/0.3M NaClO₄) (Na salt). λ_{max} 208 (ε 55200); 227 (ε 56600); 328 (ε 18500); 480 (ε 3950) (MeOH).Warabi, K. et al., *J.O.C.*, 2003, **68**, 2765-2770 (isol, pmr, cmr, ms)**Dictyodendrine B****D-365**

[510709-69-0 (Na salt)]

C₄₁H₃₀N₂O₁₀S 742.761Alkaloid from the sponge *Dictyodendrilla verongiformis*. Telomerase inhibitor.Amorph. yellow solid (as Na salt). λ_{max} 228 (ε 34700); 305 (sh); 397 (ε 9010) (MeOH) (Na salt).Warabi, K. et al., *J.O.C.*, 2003, **68**, 2765-2770 (isol, pmr, cmr, ms)

Fürstner, A. *et al.*, *J.A.C.S.*, 2006, **128**, 8087-8094 (*synth*)

Dictyodendrine C **D-366**
[510709-70-3 (Na salt)]




$C_{34}H_{24}N_2O_9S$ 636.638
Alkaloid from the sponge *Dictyodendrilla verongiformis*. Telomerase inhibitor. Greenish-yellow solid (as Na salt). λ_{max} 238 (ϵ 18300); 277 (ϵ 10500); 302 (ϵ 8600); 415 (ϵ 6540) (MeOH) (Na salt).

4'-O-sulfate: Dictyodendrine D
[510709-71-4 (di-Na salt)]
 $C_{34}H_{24}N_2O_{12}S_2$ 716.702
Alkaloid from *Dictyodendrilla verongiformis*. Telomerase inhibitor. Greenish-yellow solid (as Na salt). λ_{max} 243 (ϵ 12500); 276 (ϵ 6610); 305 (ϵ 6230); 353 (ϵ 5520); 416 (ϵ 4960) (MeOH) (Na salt).

Warabi, K. *et al.*, *J.O.C.*, 2003, **68**, 2765-2770 (*isol, pmr, emr, ms*)
Fürstner, A. *et al.*, *J.A.C.S.*, 2006, **128**, 8087-8094 (*synth*)

Dictyolomide A **D-367**
2-(3-Hexenyl)-1,2,3,4-tetrahydro-6H-pyr-
ido[1,2-a]quinolin-6-one, 9CI
[168434-18-2]




$C_{19}H_{23}NO$ 281.397
Alkaloid from stem bark of *Dictyoloma peruviana* (Rutaceae). Shows antileishmanial activity. $[\alpha]_D +21$ (c, 0.6 in $CHCl_3$).

1'-Hydroxy, 3',4'-dihydro: Dictyolomide B
[168434-19-3]
 $C_{19}H_{25}NO_2$ 299.412
Alkaloid from stem bark of *Dictyoloma peruviana* (Rutaceae). Shows antileishmanial activity. $[\alpha]_D +32$ (c, 0.9 in $CHCl_3$).

6-Methoxy: 6-Methoxydictyolomide A
 $C_{20}H_{25}NO_2$ 311.423
Alkaloid from the leaves of *Dictyoloma vandellianum*. Yellow powder. $[\alpha]_D +22$ (c, 0.7 in $CHCl_3$). λ_{max} 240 ; 331 ; 347 (MeOH).

Lavaud, C. *et al.*, *Phytochemistry*, 1995, **40**, 317-320 (*Dictyolomide A*)
Sartor, C.F.P. *et al.*, *Phytochemistry*, 2003, **63**, 185-192 (*6-Methoxydictyolomide A*)

Dictyolucidine **D-368**
17-Methyl-3-methylamino-D-homoandros-
tane-17,17a-diol
[2328-79-2]



$C_{22}H_{39}NO_2$ 349.556
Alkaloid from *Dictyophleba lucida* (Apocynaceae). Cryst. (Et₂O). Mp 198°. $[\alpha]_D^{20} +30$ ($CHCl_3$).


N-Ac:
Cryst. (Me₂CO). Mp 278°. $[\alpha]_D +9$ ($CHCl_3$).

N-Me: Dictyolucidamine
[2449-83-4]
 $C_{23}H_{41}NO_2$ 363.582
Alkaloid from *Dictyophleba lucida* (Apocynaceae). Mp 205°. $[\alpha]_D +6$ ($CHCl_3$).

N-Me, O-Ac:
Cryst. (EtOH). Mp 228-229°. $[\alpha]_D +30$ ($CHCl_3$).

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 3472 (*isol, struct, ir, pmr, ms, synth*)

Dictyoquinazole A **D-369**
3-(2-Hydroxymethyl-4-methoxyphenyl)-
6-methoxy-4(3H)-quinazolinone
[477203-69-3]



$C_{17}H_{16}N_2O_4$ 312.324
Alkaloid from the edible mushroom *Dictyophora indusiata*. Neuroprotective agent. Pale yellow oil. λ_{max} 224 (log ϵ 3.9); 274 (log ϵ 3.43); 319 (log ϵ 2.78) (MeOH).

N¹,2-Dihydro, N¹-formyl: 1-Formyl-2,3-dihydro-3-(2-hydroxymethyl-4-methoxyphenyl)-6-methoxy-4(1H)-quinazolinone. Dictyoquinazole C
[477203-71-7]
 $C_{18}H_{18}N_2O_5$ 342.351


Alkaloid from *Dictyophora indusiata*. Pale yellow oil. $[\alpha]_D +60$ (c, 0.03 in MeOH). Exists as a mixt. of rotamers. λ_{max} 228 (log ϵ 3.9); 319 (log ϵ 2.9) (MeOH).

4-Deoxy, N¹,2-dihydro, N¹-formyl: 1-Formyl-1,2,3,4-tetrahydro-3-(2-hydroxymethyl-4-methoxyphenyl)-6-methoxyquinazolinone. Dictyoquinazole B
[476488-21-8]
 $C_{18}H_{20}N_2O_4$ 328.367

Alkaloid from *Dictyophora indusiata*. Pale yellow oil. $[\alpha]_D -80$ (c, 0.05 in MeOH). Exists as a mixt. of rotamers. λ_{max} 206 (log ϵ 3.7); 239 (log ϵ 3.85); 290 (log ϵ 3.2) (MeOH).

Lee, I.-K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1769-1772 (*isol, pmr, emr*)
Oh, C.H. *et al.*, *Synth. Commun.*, 2007, **37**, 3311-3317 (*synth, pmr, Dictyoquinazols B, C*)

Dictysine **D-370**
Dictyzine. Dictizine
[67256-05-7]



$C_{21}H_{33}NO_3$ 347.497
Alkaloid from *Delphinium dictyocarpum*, *Delphinium corymbosum* and *Delphinium brunonianum* (Ranunculaceae).

15-O-(3,4-Dimethoxybenzoyl): 15-Veratroyldictysine
[852446-99-2]
 $C_{30}H_{41}NO_6$ 511.657
Alkaloid from *Aconitum variegatum*. Amorph. solid. $[\alpha]_D^{20} +79.6$ (c, 0.72 in $CHCl_3$).

15-O-(3,4-Dimethoxybenzoyl), 17-Ac: 17-Acetyl-15-veratroyldictysine
[852381-88-5]
 $C_{32}H_{43}NO_7$ 553.694
Alkaloid from *Aconitum variegatum*. Amorph. solid. $[\alpha]_D^{25} +69.1$ (c, 0.81 in $CHCl_3$).

N-De-Me, N-Et: N-Ethyl-N-demethyl-dictysine
[863192-72-7]
 $C_{22}H_{35}NO_3$ 361.523
Alkaloid from *Delphinium corymbosum*. Unpubl. information (1999).

16,17-Methylene ether: Corumdizine
[863203-33-2]
 $C_{22}H_{33}NO_3$ 359.508
Alkaloid from *Delphinium corymbosum* (Ranunculaceae). Cryst. (hexane). Mp 104-105°.

16,17-Methylene ether, N-de-Me, N-Et: Corumdizine
[863203-31-0]
 $C_{23}H_{35}NO_3$ 373.534
Alkaloid from *Delphinium corymbosum* (Ranunculaceae). Amorph.

16,17-O-Isopropylidene: Dictysine acetone
[82413-15-8]
 $C_{24}H_{37}NO_3$ 387.561
Alkaloid from *Delphinium dictyocarpum* and *Delphinium corymbosum* (Ranunculaceae).

15-Ketone: 15-Dehydrodictysine
[74119-95-2]
 $C_{21}H_{31}NO_3$ 345.481
Alkaloid from *Delphinium dictyocarpum* (Ranunculaceae). Mp 145° (as acetone). $[\alpha]_D -58$ ($CHCl_3$) (acetone).

19-Oxo, 15-O-(3,4-dimethoxybenzoyl), 17-Ac: 17-Acetyl-15-veratroyl-19-oxo-dictysine
[852381-89-6]
 $C_{32}H_{41}NO_8$ 567.678

Alkaloid from *Aconitum variegatum*. Amorph. solid. $[\alpha]_D^{25} +49.3$ (c, 0.15 in CHCl_3).

Salimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 812-817; 1982, **18**, 86-91; 1985, **21**, 95-98; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 718-722; 1982, **18**, 81-85; 1985, **21**, 91-94 (*isol, struct*)

Dzhakhangirov, F.N. *et al.*, *Dokl. Akad. Nauk SSSR*, 1982, 37-38; *CA*, **99**, 151861b (*pharmacol*)

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 230-235; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 211-215 (*cryst struct*)

Deng, W. *et al.*, *Heterocycles*, 1986, **24**, 869-872 (*occur*)

Joshi, B.S. *et al.*, *Tetrahedron*, 1991, **47**, 4299-4316 (*pmr, cmr*)


Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 410-512; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 386-512 (*Corundizine, Corundizimine*)

Shrestha, P.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1574-1576 (*pmr, cmr, ms*)

Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 837-846 (*Aconitum variegatum alkaloids*)

3,5-Didecanoylpyridine D-371

1,1'-(3,5-Pyridinediyl)bis-*I*-decanone, 9CI
[149682-93-9]




$\text{C}_{25}\text{H}_{41}\text{NO}_2$ 387.604

Alkaloid from *Houttuynia cordata* (Yu Xing Cao).

Jong, T.T. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1993, **40**, 301-303 (*isol*)

Proebstle, A. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 235-240 (*isol*)

6,8-Di-1-decenyl-2,3-dihydro-5,7-dimethyl-1H-indolizinium D-372



$\text{C}_{30}\text{H}_{50}\text{N}^{\oplus}$ 424.732


(Z,Z)-form

Quaternary alkaloid from the wood of *Aniba panurensis*. Antifungal agent. Yellow oil (as TFA salt). λ_{max} 284 (log ϵ 3.6) (MeOH) (as trifluoroacetate salt).

Klausmeyer, P. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1732-1735 (*isol, pmr, cmr, ms*)

6a,7-Didehydro-7-hydroxy-1,2-dimethoxyaporphine D-373

[252564-91-3]



$\text{C}_{19}\text{H}_{19}\text{NO}_3$ 309.364

Alkaloid from the stem bark of *Enantia chlorantha*. Brown powder (MeOH). Mp 258-260°.

N-De-Me: 6a,7-Didehydro-7-hydroxy-1,2-dimethoxyaporphine
[252564-92-4]


$\text{C}_{18}\text{H}_{17}\text{NO}_3$ 295.337

Alkaloid from the stem bark of *Enantia chlorantha*. Black powder (MeOH). Mp 256-257°.

Wafu, P. *et al.*, *Fitoterapia*, 1999, **70**, 157-160 (*isol, pmr, cmr*)

16,17-Didehydro-17-hydroxy-16-(hydroxymethyl)corynan-5-carboxylic acid, 9CI

RCNSI
[93597-92-3]




$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_4$ 370.447

Tentative struct. Alkaloid from *Rauwolfia caffra* stem bark (Apocynaceae). Yellow amorph. powder. Unstable.

Nasser, A.M.A.G. *et al.*, *J. Ethnopharmacol.*, 1984, **11**, 99; *CA*, **102**, 3224m (*isol, uv, ms, struct*)

1,3a-Didehydro-8-hydroxy-ptilocaulin D-375

[191353-14-7]



$\text{C}_{15}\text{H}_{23}\text{N}_3\text{O}$ 261.366

The name is not an accurate representation of the struct (cf. Ptilocaulin, P-773). Alkaloid from the sponges *Batzella* sp. and *Monanchora unguifera*. Antiplasmodial agent. Oil. $[\alpha]_D +39.3$ (c, 0.28 in MeOH). λ_{max} 239 ; 246 ; 309 (MeOH).

8-Epimer:

$\text{C}_{15}\text{H}_{23}\text{N}_3\text{O}$ 261.366


Alkaloid from *Monanchora unguifera*. Antiplasmodial agent. Isol. as a mixt. with its 8-epimer.

Patil, A.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 704-707 (*isol, pmr, cmr*)

Hua, H.-M. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 6461-6464 (*isol, pmr, cmr*)

3,14-Didehydro-19-methyl-normalindine D-376

[292159-40-1]




$\text{C}_{20}\text{H}_{19}\text{N}_3$ 301.39

Alkaloid from *Ophiorrhiza kunstleri*. Yellow needles (EtOAc/petrol). Mp 128-130° dec. λ_{max} 217 (ϵ 18300); 254 (ϵ 7800); 368 (ϵ 16200) (MeOH).

Dachriyanus, *et al.*, *Aust. J. Chem.*, 2000, **53**, 221-224

1,2-Didehydropseudoaspidospermidine D-377

1,2-Dehydro- ψ -aspidospermidine. Δ^1 -Pseudoaspidospermidine. epi- Δ^1 - ψ -Aspidospermidine
[90702-16-2]



Absolute Configuration

$\text{C}_{19}\text{H}_{24}\text{N}_2$ 280.412

A major alkaloid from leaves and twigs of *Tabernaemontana eglanulosa* and from *Pandaca boiteaui*. $[\alpha]_D^{20} +209$ (c, 0.44 in CHCl_3). λ_{max} 221 ; 265 ; 270 ; 328 ; 345 (MeOH).

14,15-Didehydro: 14,15-Anhydrocapuronidine
[70031-39-9]

$\text{C}_{19}\text{H}_{22}\text{N}_2$ 278.396

Minor alkaloid from leaves and stem bark of *Capuronetta elegans* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. $[\alpha]_D^{20} -54$ (c, 0.72 in CHCl_3). λ_{max} 222 (log ϵ 4.24); 262 (log ϵ 3.77) (EtOH).

14,15-Didehydro, 1,2- ζ -dihydro: 14,15-Anhydrodihydrocapuronidine
[70031-40-2]

$\text{C}_{19}\text{H}_{24}\text{N}_2$ 280.412

Minor alkaloid from the leaves and stem bark of *Capuronetta elegans* (Apocynaceae). Amorph. $[\alpha]_D^{20} +40$ (c, 1.12 in CHCl_3). λ_{max} 226 (log ϵ 3.86); 244 (log ϵ 3.74); 292 (log ϵ 3.44) (EtOH).

15 α -Hydroxy: Capuronidine
[57533-92-3]

$\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}$ 296.411

Alkaloid from stem bark of *Capuronetta elegans* (Apocynaceae). Amorph. $[\alpha]_D^{20} +220$ (c, 1 in CHCl_3). λ_{max} 220 (log ϵ 4.29); 260 (log ϵ 3.6) (EtOH).

20-Hydroxy: 1,2-Didehydro-20-hydroxy-pseudoaspidospermidine. 20-Hydroxy-1,2-dehydro- Ψ -aspidospermidine
[90865-28-4]

C₁₉H₂₄N₂O 296.411

Minor alkaloid from the leaves and twigs of *Tabernaemontana eglandulosa* (Apocynaceae). Oil.

20-Epimer: 1,2-Didehydro-7-ethyl-20,21-dinoraspido-spermidine. Δ¹-Pseudoaspido-spermidine. Δ¹-ψ-Aspidospermidine [56423-79-1]

C₁₉H₂₄N₂ 280.412

Alkaloid from *Tabernaemontana eglandulosa*, *Pandaca boiteaui* and *Pandaca eusepala* (Apocynaceae). Noncryst. [α]_D²⁵ +153 (CHCl₃). λ_{max} 220 ; 258 ; 270 ; 290 (sh) (MeOH).

20-Epimer, dihydro: [26251-93-4]

Alkaloid from *Pandaca boiteaui*. Mp 89°. [α]_D²⁵ +60 (CHCl₃).

Quirin, F. et al., *Phytochemistry*, 1975, **14**, 812-813 (Δ¹-ψ-Aspidospermidine)

Chardon-Loriaux, I. et al., *Tet. Lett.*, 1975, 1845-1848 (*Capuronidine*)


Chardon-Loriaux, I. et al., *Phytochemistry*, 1978, **17**, 1605-1608 (*Anhydrocapuronidine*, *Anhydrodihydrocapuronidine*)

Andriantsiferana, M. et al., *Phytochemistry*, 1979, **18**, 911-912 (*Pandaca boiteaui* constits)

Van Beek, T.A. et al., *Tetrahedron*, 1984, **40**, 737-748 (*Tabernaemontana eglandulosa* constits)

19,20-Didehydroyohimbine D-378

Methyl 19,20-didehydro-17-hydroxy-yohimban-16-carboxylate, 9CI. 19-Dehydroyohimbine [95727-43-8]



Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Aspidosperma pyricollum* (Apocynaceae). Mp 254° dec. [α]_D²⁷ +106 (c, 0.53 in Py).

O-Ac: O-Acetyl-19,20-didehydroyohimbine. 19,20-Dehydro-O-acetylyohimbine [74713-13-6]

C₂₃H₂₆N₂O₄ 394.469

Alkaloid from leaves of *Alstonia angustifolia*. [α]_D²⁵ +75 (c, 0.1 in MeOH). Config. at C-17 not detd. Could be the C-17 epimer, to which the CAS No. refers. λ_{max} 227 ; 283 ; 290 (MeOH).

16-Epimer: 19,20-Didehydro-α-yohimbine. 19,20-Dehydro-α-yohimbine [60142-21-4]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the seeds of *Aspidosperma oblongum* and from *Rauwolfia obscura* (tentative stereochem.) (Apocynaceae). Amorph. [α]_D²⁵ +90 (c, 0.64 in MeOH).

17-Epimer: 19,20-Didehydro-β-yohimbine. 19,20-Dehydro-β-yohimbine [4109-62-0]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from seeds of *Aspidosperma*

oblongum (Apocynaceae). Amorph. [α]_D²⁵ -45 (c, 0.42 in Py).

Arndt, R.R. et al., *Experientia*, 1965, **21**, 566-568 (*isol, uv*)

Timmins, P. et al., *Phytochemistry*, 1976, **15**, 733-735 (16-epimer, *isol, uv*)


Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 708-722 (*isol, uv, ir, pmr, ms, cd, struct, epimers*)

Miyata, O. et al., *Heterocycles*, 1984, **22**, 2719-2722 (*synth*)

Ghedira, K. et al., *Phytochemistry*, 1988, **27**, 3955-3962 (*Ac, isol*)

Didemnimide A D-379

3-(1H-Imidazol-4-yl)-4-(1H-indol-3-yl)-1H-pyrrole-2,5-dione, 9CI [186143-93-1]

C₁₅H₁₀N₄O₂ 278.27

Alkaloid from the Caribbean mangrove ascidian *Didemnum conchyliatum*. Potent fish feeding deterrent. Irregular orange needles (MeCN aq.). Mp 234-235°. λ_{max} 430 (ε 3000) (MeCN).

N¹⁵-Me: Didemnimide E

[219828-98-5]

C₁₆H₁₂N₄O₂ 292.296

From *Didemnum granulatum*. Amorph. orange solid.

N¹⁷-Me: Didemnimide C

[186144-21-8]

C₁₆H₁₂N₄O₂ 292.296

From *Didemnum conchyliatum*. Potent fish feeding deterrent. Dark orange needles (MeCN aq.). Mp > 300°. λ_{max} 420 (ε 5700) (MeCN).

6-Bromo: Didemnimide B

[186144-09-2]

C₁₅H₉BrN₄O₂ 357.166

From *Didemnum conchyliatum*. Potent fish feeding deterrent. Fine light orange needles (MeCN aq.). Mp > 300°. λ_{max} 420 (ε 4300) (MeCN).

6-Bromo, N¹⁷-Me: Didemnimide D

[186144-28-5]

C₁₆H₁₁BrN₄O₂ 371.193

From *Didemnum conchyliatum*. Potent fish feeding deterrent. Small dark orange needles (MeCN aq.). Mp > 250° dec. More active than other Didemnimides. λ_{max} 416 (ε 4500) (MeCN).

Vervoort, H.C. et al., *J.O.C.*, 1997, **62**, 1486-1490 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Berlinck, R.G.S. et al., *J.O.C.*, 1998, **63**, 9850-9856 (*Didemnimide E*)


Terpin, A. et al., *Tetrahedron*, 1998, **54**, 1745-1752 (*Didemnimide C, synth*)

Piers, E. et al., *J.O.C.*, 2000, **65**, 530-535 (*Didemnimide C, synth*)

Fan, H. et al., *Chem. Rev.*, 2008, **108**, 264-287 (*rev*)

Didemnoline A D-380

7-Bromo-9-[[4-(methylthio)-1H-imidazol-3-yl]methyl]-9H-pyrido[3,4-b]indole, 9CI [168434-22-8]

C₁₆H₁₃BrN₄S 373.275

Alkaloid from the marine ascidian *Didemnum* sp. Exhibits moderate cytotoxicity toward human epidermoid carcinoma (KB) cells. Also shows antimicrobial activity. λ_{max} 240 (ε 38700); 290 (ε 13500); 358 (ε 5000) (MeOH) (Berdy).

S-Oxide: Didemnoline C

[168434-23-9]

C₁₆H₁₃BrN₄OS 389.275

From *Didemnum* sp. Cytotoxic. Shows antimicrobial activity. [α]_D²⁵ +97.2 (c, 0.1 in DMSO). *Isol.* as a ca. 10:1 inseparable mixture with Didemnoline D. λ_{max} 239 (ε 37200); 290 (ε 14000); 358 (ε 4000) (MeOH) (Berdy). λ_{max} 243 ; 277 (ε 34000); 288 (ε 2400); 342 ; 352 (ε 4000) (EtOH) (Berdy).

Debromo: Didemnoline B

[168434-24-0]

C₁₆H₁₄N₄S 294.379

From *Didemnum* sp. Moderately cytotoxic. λ_{max} 240 (ε 36800); 290 (ε 13500); 358 (ε 5000) (MeOH) (Berdy). λ_{max} 216 (ε 33000); 238 (ε 39800); 290 (ε 14100); 358 (ε 4900) (EtOH) (Berdy).

Debromo, S-oxide: Didemnoline D

[168434-25-1]


C₁₆H₁₄N₄OS 310.379

From *Didemnum* sp. Not obtained pure. Schumacher, R.W. et al., *Tetrahedron*, 1995, **51**, 10125 (*isol, uv, ir, pmr, cmr, ms, struct*)

Schumacher, R.W. et al., *Tetrahedron*, 1999, **55**, 935-942 (*synth*)

Didmolamide A D-381

[528815-53-4]



Absolute Configuration

C₂₅H₂₆N₆O₄S₂ 538.65

Isol. from the ascidian *Didemnum molle*. Oil. [α]_D²⁵ -35.7 (c, 0.6 in MeOH).

Stereoisomer (?): Banyascyclamide A

[501903-99-7]

C₂₅H₂₆N₆O₄S₂ 538.65

Isol. from *Nostoc* sp. TAU strain IL-235. Amorph. solid. [α]_D²⁵ -19.3 (c, 0.12 in MeOH). Stereochem. partly deter-

mined. May be identical with Didmolamide A. λ_{\max} 218 (ϵ 17300); 238 (ϵ 17900) (MeOH).


Ploutno, A. *et al.*, *Tetrahedron*, 2002, **58**, 9949-9957 (*Banyascyclamide A*)

Rudi, A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 575-577 (*Didmolamide A*)

You, S.-L. *et al.*, *Tet. Lett.*, 2005, **46**, 2567-2570 (*synth*)

Didmolamide B

[528815-55-6]

D-382

Absolute Configuration

 $C_{25}H_{28}N_6O_5S_2$ 556.665

Isol. from the ascidian *Didemnum molle*. Oil. $[\alpha]_D^{25}$ -216 (c, 0.11 in MeOH).


Rudi, A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 575-577 (*isol, pmr, cmr, ms*)

You, S.-L. *et al.*, *Tet. Lett.*, 2005, **46**, 2567-2570 (*synth*)

Bertram, A. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1541-1553 (*synth*)

Didymeline

3-Hydroxy-9,10-seco-1,3,5,8(14)-conatetraen-9-one

D-383 $C_{22}H_{29}NO_2$ 339.477

Secosteroid alkaloid. Alkaloid from the leaves of *Didymele madagascariensis* (Didymelaceae). Cryst. (MeOH). Sol. H_2O . Mp 165°. $[\alpha]_D^{20}$ +60 (c, 0.43 in EtOH). λ_{\max} 240 (ϵ 15400); 280 (ϵ 11100) (H_2O) (Berdy).

8,14-Dihydro- 3-Hydroxy-9,10-seco-1,3,5-conatrien-9-one. 3-Hydroxy-9-oxo-9,10-seco-1,3,5-conatriene


 $C_{22}H_{31}NO_2$ 341.492

Alkaloid from the leaves of *Didymele madagascariensis* (Didymelaceae). $[\alpha]_D^{20}$ +100 (c, 0.08 in EtOH).

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1987, 877 (*isol, uv, ir, pmr, cmr, ms, cd, cryst struct*)

Dielsine†

[125581-66-0]


D-384 $C_{37}H_{38}N_2O_7$ 622.716

Alkaloid from bark of *Daphnandra dielsii* (Monimiaceae). Fine yellow needles + $\frac{1}{2}CHCl_3$ ($CHCl_3$). Mp 185-190° dec. $[\alpha]_D^{27}$ +243 ($CHCl_3/MeOH$).

Critchett, C.D. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2043-2046 (*isol, uv, ir, pmr, ms, struct*)

Dielsinol

4-(Hydroxymethyl)-1H-indeno[1,2-b]pyridine-2,5-dione, 9CI [104669-01-4]

D-385 $C_{13}H_9NO_3$ 227.219

Isomeric struct. originally proposed. Alkaloid from the trunkwood of *Guatteria dielsiana* (Annonaceae). Yellow-greenish cryst. Mp 252-254°. λ_{\max} 253 (log ϵ 4.26); 269 (sh) (log ϵ 4.11); 283 (log ϵ 4.03); 333 (log ϵ 3.77) (EtOH).

Deoxy: 4-Methyl-1H-indeno[1,2-b]pyridine-2,5-dione, 9CI. **Dielsine†** [104696-14-2]

 $C_{13}H_9NO_2$ 211.22

Alkaloid from trunkwood of *Guatteria dielsiana* (Annonaceae). Yellow needles (Me_2CO or AcOH). Mp 254-256° natural Mp 325° synthetic. The struct. assigned to this alkaloid is erroneous but has not yet been corrected. Consequently, the struct. of dielsinol may also be in error. No further work to 2006. λ_{\max} 243 (sh) (log ϵ 4.15); 256 (log ϵ 4.18); 262 (sh) (log ϵ 4.17); 272 (sh) (log ϵ 4.04); 282 (log ϵ 3.56); 341 (log ϵ 3.5) (EtOH) (natural). λ_{\max} 231 (log ϵ 4.38); 249 (log ϵ 4.19); 291 (log ϵ 4.13); 301 (log ϵ 4.23); 334 (log ϵ 3.88); 349 (log ϵ 3.96); 425 (log ϵ 3.05) (EtOH) (synthetic).


Goulart, M.O.F. *et al.*, *Phytochemistry*, 1986, **25**, 1691-1695 (*isol, uv, ir, pmr, ms*)

Tadić, D. *et al.*, *Phytochemistry*, 1987, **26**, 1551-1552 (*struct*)

Bracher, F. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1992, **325**, 645-648 (*deoxy, synth*)

Dienomycin C

3-Methyl-2-(4-phenyl-1,3-butadienyl)-4-piperidinol [27542-56-9]

D-386 $C_{16}H_{21}NO$ 243.348

From *Streptomyces* strain MC67-Cl. Shows weak antibiotic activity against

mycobacteria. Needles ($MeOH/Me_2CO$). Sol. $MeOH$, Me_2CO , EtOH, DMSO, $CHCl_3$; poorly sol. H_2O , Et_2O , C_6H_6 , EtOAc. Mp 130-131°. $[\alpha]_D^{25}$ +72 (c, 1.5 in MeOH). λ_{\max} 211 (ϵ 16000); 220 (ϵ 15000); 227 (ϵ 15000); 234 (ϵ 11000); 280 (sh) (ϵ 42000); 287 (ϵ 46000); 297 (sh) (ϵ 35000); 307 (sh) (ϵ 20000) (MeOH) (Derep). λ_{\max} 211 (ϵ 16000); 220 (ϵ 15000); 227 (ϵ 15000); 234 (ϵ 12000); 287 (ϵ 50000) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 45 mg/kg. TN7750000

Hydrochloride:

Cryst. (EtOAc/MeOH aq.). Mp 252-253°. $[\alpha]_D^{26}$ +65 (c, 1 in MeOH).

O-Ac: Dienomycin B

[27542-55-8]

 $C_{18}H_{23}NO_2$ 285.385

From *Streptomyces* strain MC67-Cl. Cryst. (EtOAc/MeOH). Sol. $MeOH$, DMSO, acids, $CHCl_3$, EtOH; fairly sol. H_2O ; poorly sol. C_6H_6 , Et_2O , EtOAc. λ_{\max} 211 (ϵ 14800); 220 (ϵ 14700); 227 (ϵ 14700); 234 (ϵ 10500); 280 (sh) (ϵ 40900); 287 (ϵ 44100); 297 (sh) (ϵ 33600); 307 (sh) (ϵ 19600) (MeOH) (Derep). λ_{\max} 210 (ϵ 14500); 218 (ϵ 14600); 226 (ϵ 14600); 232 (ϵ 10500); 287 (ϵ 42000) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 360 mg/kg. TN7760000

O-Ac; hydrochloride:

Cryst. (EtOAc/MeOH aq.). Mp 280-281° (sealed tube). $[\alpha]_D^{20}$ +80 (c, 1 in MeOH).

O-(2-Methylpropanoyl): **Dienomycin A** [27542-54-7]

 $C_{20}H_{27}NO_2$ 313.439

From *Streptomyces* strain MC67-Cl. Cryst. (EtOAc/MeOH). Sol. $MeOH$, acids, EtOH, DMSO, $CHCl_3$; fairly sol. H_2O ; poorly sol. C_6H_6 , Et_2O , EtOAc. pK_a 8.8. λ_{\max} 211 (ϵ 14800); 220 (ϵ 14700); 227 (ϵ 14700); 234 (ϵ 10500); 280 (sh) (ϵ 40900); 287 (ϵ 44100); 297 (sh) (ϵ 33600); 307 (sh) (ϵ 19600) (MeOH) (Derep). λ_{\max} 211 (ϵ 14800); 220 (ϵ 14700); 227 (ϵ 14700); 234 (ϵ 10500); 287 (ϵ 44100) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 90 mg/kg. NQ5430000


O-2-Methylpropanoyl; hydrochloride:

Cryst. (EtOAc/MeOH aq.). Mp 212-214°. $[\alpha]_D^{20}$ +85 (c, 1 in MeOH).

Umezawa, S. *et al.*, *J. Antibiot., Ser. A*, 1970, **23**, 20; 28

Ripoche, I. *et al.*, *Eur. J. Org. Chem.*, 1999, **37**, 1517-1521 (*synth, pmr, cmr, abs config*)

Comins, D.L. *et al.*, *Tet. Lett.*, 1999, **40**, 217-218 (*synth*)

11,13:11,16-Diepoxy-16,17-dihydro-11,12-secohetisan-2-ol, 9CI**D-387**

C₂₀H₂₇NO₃ 329.438**2 α -form** [80249-79-2]

Alkaloid from *Aconitum hepterothphyllum* and *Consolidia glandulosa*. Insect antifeedant. Cryst. Mp 278-280°. [α]_D²⁸ +9.1 (c, 1.4 in CH₂Cl₂).

Pelletier, S.W. *et al.*, *Chem. Comm.*, 1981, 327-329 (*synth, pmr, cmr, cryst struct*)

González-Coloma, A. *et al.*, *Chem. Biodiversity*, 2004, **1**, 1327-1335 (*isol, activity*)

Diethanolamine **D-388**

2,2'-Iminobisethanol, 9CI. Bis(2-hydroxyethyl)amine. 2,2'-Iminodiethanol. 2,2'-Dihydroxydiethylamine. Diolamine [111-42-2]

HOCH₂CH₂NHCH₂CH₂OHC₄H₁₁NO₂ 105.136


Manuf. by reaction of oxirane with ammonia. Isol. from *Senecio dunedin* and *Echinops exaltatus*. Industrial basic solv. Crosslinking agent for epoxy resins and polyurethanes. Base used in pharmaceuticals etc. Deliquescent prisms. V. sol. H₂O, EtOH, Me₂CO; insol. Et₂O, C₆H₆. Mp 28°. Bp 270° Bp₁₀ 154-155°. pK_a 8.91 (25°, H₂O). Nonvolatile in steam. Strong base.

► Fl. p. 138° (oc), autoignition temp. 662°. Severe eye and mild skin irritant. LD₅₀ (rat, orl) 710 mg/kg. OES: long-term 3ppm. KL2975000

Brown, L.S.R. *et al.*, *J. Nat. Prod.*, 1986, **49**, 910-912 (*isol*)

2,5-Diethyl-3,6-dimethylpyrazine **D-389**

[18903-30-5]

C₁₀H₁₆N₂ 164.25

Isol. from an arctic marine bacterium. Liq. Bp 214-217° Bp₂ 65°.

Octahydrate: Mp 48-50°.

Asinger, F. *et al.*, *Monatsh. Chem.*, 1959, **90**, 402-416 (*synth*)

Baltes, W. *et al.*, *J. Agric. Food Chem.*, 1987, **35**, 340-346 (*occur, coffee, ms*)

Rizzi, G.P. *et al.*, *J. Agric. Food Chem.*, 1988, **36**, 349-352 (*synth, ms, pmr, uv*)

Silwar, R. *et al.*, *Z. Lebensm.-Unters. -Forsch.*, 1992, **195**, 112-119 (*occur*)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, **2**, 318-353 (*isol*)

2,6-Diethyl-3,5-dimethylpyrazine **D-390**

[18940-74-4]

C₁₀H₁₆N₂ 164.25

Isol. from an arctic marine bacterium. Characterised spectroscopically.

Baltes, W. *et al.*, *J. Agric. Food Chem.*, 1987, **35**, 340-346 (*occur, coffee, ms*)


Rizzi, G.P. *et al.*, *J. Agric. Food Chem.*, 1988, **36**, 349-352 (*occur, ms, uv*)

Silwar, R. *et al.*, *Z. Lebensm.-Unters. -Forsch.*, 1992, **195**, 112-119 (*occur*)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, **2**, 318-353 (*isol*)

8,10-Diethyllobelidol **D-391**

α, α' -Diethyl-1-methyl-2,6-piperidine-diethanol. 2,6-Bis(2-hydroxybutyl)-1-methylpiperidine [110250-13-0]

C₁₄H₂₉NO₂ 243.389

The alkaloids covered by this entry are not stereochemically homologous as some are opt. active and some inactive. However, since the known information is fragmentary they are merged here and known data is noted. Abs. configs. are unknown. Minor alkaloid from *Lobelia inflata* (Campanulaceae). Opt. inactive.

Hexachloroplatinate: Mp 203-204° dec.

Picrate: [125826-49-5]

Needles (AcOH/Et₂O). Mp 138°.

Monoketone: **8,10-Diethyllobelionol**. 2-(2-Hydroxybutyl)-1-methyl-6-(2-oxobutyl)piperidine. Alkaloid D2b [139406-22-7]

[90980-50-0 (HCl)]

C₁₄H₂₇NO₂ 241.373

Alkaloid from *Lobelia syphilitica* (Campanulaceae). Mp 120-121° (as hydrochloride). 2,6-*cis*-. Laevorotatory hydrochloride. The nat. alkaloid may be the *trans*-form (alkali used during isol.).

Monoketone, N-de-Me: **8,10-Diethylnorlobelionol**. 2-(2-Hydroxybutyl)-6-(2-oxobutyl)piperidine. Alkaloid C1 [803649-66-3]

[96932-55-7 (HCl)]

C₁₃H₂₅NO₂ 227.346

Alkaloid from *Lobelia syphilitica* (Campanulaceae). Mp 183-184° (as hydrochloride). 2,6-*cis*-. No opt. rotn. reported.

Diketone, N-de-Me: **8,10-Diethylnorlobelidone**. 2,6-Bis(2-oxobutyl)piperidine. Alkaloid D1†

[500545-69-7]

[700116-78-5 (HCl)]

C₁₃H₂₃NO₂ 225.33

Alkaloid from *Lobelia syphilitica* (Campanulaceae). Mp 183-184° (as hydrochloride). *cis*-. No opt. rotn. reported.

3,4-Didehydro: **3,4-Dehydro-8,10-diethyllobelidol**. Alkaloid C3. 1,2,5,6-Tetrahydro-2,6-bis(2-hydroxybutyl)-1-methylpyridine. 2,6-Bis(2-hydroxybutyl)-1-methyl- Δ^3 -piperidine. α, α' -Diethyl-1,2,5,6-tetrahydro-1-methyl-2,6-pyridinediethanol, 9CI [105694-50-6]

C₁₄H₂₇NO₂ 241.373

Alkaloid from *Lobelia syphilitica* and *Lobelia berlandieri* (Campanulaceae). Oil. Mp 128° (as hydrochloride). [α]_D²⁵ -114 (EtOH). Both isolates 2,6-*trans*-, but the two isolates need not have the same stereochem.

3,4-Didehydro, 2'-ketone: **6-(2-Hydroxybutyl)-1-methyl-2-(2-oxobutyl)- Δ^3 -piperidine**. 1-[1,2,5,6-Tetrahydro-6-(2-hydroxybutyl)-1-methyl-2-pyridinyl]-2-butanone, 9CI. 1,2,5,6-Tetrahydro-6-(2-hydroxybutyl)-1-methyl-2-(2-oxobutyl)pyridine [105694-52-8]

C₁₄H₂₅NO₂ 239.357

Alkaloid from the toxic plant *Lobelia berlandieri* (Campanulaceae).

[4748-03-2, 96998-45-7]

Schöpf, C. *et al.*, *Annalen*, 1957, **608**, 88-127 (3,4-dihydro, *isol, struct, biosynth, nomencl*)


Tschesche, R. *et al.*, *Chem. Ber.*, 1961, **94**, 3327-3336 (*Lobelia syphilitica constits*)

Williams, H.J. *et al.*, *J. Agric. Food Chem.*, 1987, **35**, 19-22 (3,4-didehydro 2'-ketone)

2,3-Diethyl-5-methylpyrazine, 8CI **D-392**

FEMA 3336

[18138-04-0]

C₉H₁₄N₂ 150.223

Found in galbanum oil. Aroma constit. of many foods. Flavouring agent. Bp₁₂ 79-80°. n_D²⁰ 1.4980.

U.S. Pat., 1975, 3 924 015; CA, **84**, 164364g (*synth*)

Masuda, H. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 944 (*occur*)


Mihara, S. *et al.*, *J. Chromatogr.*, 1987, **402**, 309 (*glc*)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 819-820 (*use, occur*)

3,5-Diethyl-2-methylpyrazine **D-393**

FEMA 3916

[18138-05-1]

C₉H₁₄N₂ 150.223

Constit. of galbanum oil. Flavouring agent.


Kato, S. *et al.*, *Agric. Biol. Chem.*, 1970, **34**, 1826 (*synth*)

Burrell, J.W.K. *et al.*, *Chem. Ind. (London)*, 1970, 1409 (*isol*)

Shigematsu, H. *et al.*, *Agric. Biol. Chem.*, 1972, **36**, 1631 (*synth*)

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 410

6,8-Diethyloctahydro-5-propyl-1*H*-indolizine, 9CI D-394
6,8-Diethyl-5-propylindolizidine. *Indolizidine 223A*. Dendrobates *Alkaloid 223A*† [185417-25-8]



C₁₅H₂₉N 223.401

(5*R*,6*R*,8*R*,8*aS*)-form

Alkaloid from the skin extract of *Dendrobates pumilio*. Stereochem. revised in 2002.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 2-5 (*isol*)


Toyooka, N. *et al.*, *Org. Lett.*, 2002, **4**, 1715-1717 (*synth, rel config*)

Pu, X. *et al.*, *J.O.C.*, 2003, **68**, 4400-4405 (*synth*)

Davis, F.A. *et al.*, *J.A.C.S.*, 2005, **127**, 8398-8407 (*synth*)

Zhu, W. *et al.*, *Org. Lett.*, 2005, **7**, 705-708 (*synth*)

5,8'-Diferulic acid D-395
4,4'-Dihydroxy-3,5'-dimethoxy- α ,3'-bicinamic acid. 5,8'-Dehydrodiferulic acid



C₂₀H₁₈O₈ 386.357

Dehydrodimer of 3-(4-Hydroxy-3-methoxyphenyl)-2-propenoic acid.

(*E,E*)-form [160097-30-3]

Prod. by *Aspergillus unilateralis*. Constit. of the cell walls of various Poaceae. Fine cryst. Mp 160-165° dec.

Bis[2-(4-hydroxyphenyl)ethylamide]:

C₃₆H₃₆N₂O₈ 624.689

Constit. of the leaves of *Aptenia cordifolia*. Amorph. powder. λ_{\max} 246 (log ϵ 3.2); 315 (log ϵ 2.6); 336 (log ϵ 0.9) (MeOH).

Ralph, J. *et al.*, *J.C.S. Perkin 1*, 1994, 3485-3498 (*isol, pmr, cmr, synth*)


Micard, V. *et al.*, *Phytochemistry*, 1997, **44**, 1365-1368 (*isol*)

Rouau, X. *et al.*, *Phytochemistry*, 2003, **63**, 899-903 (*isol*)

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 123-129 (*isol*)

DellaGreca, M. *et al.*, *Tetrahedron*, 2006, **62**, 2877-2882 (*bis-4-hydroxyphenethylamide*)

Difforine D-396
4-(17-Norsarpagan-16-yl)-3-buten-2-one, 9CI
[102719-87-9]




Absolute Configuration

C₂₂H₂₄N₂O 332.444

Alkaloid from the leaves of *Vinca difformis* (Apocynaceae). Cryst. (Et₂O). Mp 242-244°. λ_{\max} 226; 272; 281 (sh) (EtOH).

Garnier, J. *et al.*, *Planta Med.*, 1986, **52**, 66-67 (*isol, uv, ir, pmr, cmr, ms, struct*)

Difforlemenine D-397
[96681-59-3]



Absolute Configuration


C₂₂H₂₄N₂O₅ 396.442

Struct. revised in 1994. Alkaloid from the leaves of *Vinca difformis* and from twigs and leaves of *Tabernaemontana glandulosa* (Apocynaceae). Cryst. (Me₂CO). Mp 238-240° (236-238°). $[\alpha]_D$ -47 (c, 0.52 in MeOH).

Garnier, J. *et al.*, *Tet. Lett.*, 1985, **26**, 1513 (*uv, ir, pmr, ms*)

Achenbach, H. *et al.*, *Phytochemistry*, 1994, **37**, 1737 (*isol, uv, ir, pmr, cmr, cd, ms, struct*)

Difforlemenitine D-398
[163181-61-1]



Absolute Configuration

C₂₂H₂₆N₂O₆ 414.457

Alkaloid from twigs and leaves of *Tabernaemontana glandulosa* (Apocynaceae). Cryst. (Me₂CO). Mp 126-131°. $[\alpha]_D$ +0.2 (c, 0.42 in CHCl₃).

19-Epimer: 19-Epidifforlemenitine


[162289-00-1]

C₂₂H₂₆N₂O₆ 414.457

Alkaloid from twigs and leaves of *Tabernaemontana glandulosa* (Apocynaceae). Cryst. (Me₂CO). Mp 142-145°. $[\alpha]_D$ +2.3 (c, 0.83 in CHCl₃).

Achenbach, H. *et al.*, *Phytochemistry*, 1994, **37**, 1737 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

Digracine D-399
[216444-10-9]



C₃₄H₃₄N₂O₆ 566.652

Alkaloid from *Galanthus gracilis*. Amorph. solid. $[\alpha]_D$ -74 (c, 0.17 in MeOH). λ_{\max} 232 (sh) (log ϵ 4.29); 293 (log ϵ 3.99) (MeOH).

Noyan, S. *et al.*, *Heterocycles*, 1998, **48**, 1777-1791 (*isol, uv, cd, ir, pmr, cmr, ms*)

1,4-Diguanidinobutane D-400
N,N''-1,4-Butanediylbisguanidine, 9CI.

Arcaine

[544-05-8]

[36587-93-6 (sulfate)]

HN=C(NH₂)NH(CH₂)₄NHC(NH₂)=NH

C₆H₁₆N₆ 172.233

Occurs in marine bivalve *Arca noae* and in *Panus tigrinus*. Also from the marine polychaete worm *Audouinia tentaculata*. Lowers blood sugar of mammals. Needles (as sulfate salt). Mp 291° dec. (sulfate).

Picrate:

Needles or fine powder. Mp 251-254° dec.

Kutscher, F. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1931, **199**, 27; **203**, 132 (*isol*)

Sugino, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1942, **17**, 126; *CA*, **41**, 4455 (*synth*)

Roche, J. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1965, **260**, 7023 (*isol*)

Boldt, A. *et al.*, *Phytochemistry*, 1971, **10**, 731 (*biosynth*)

Thailambal, V.G. *et al.*, *Acta Cryst. C*, 1985, **41**, 105 (*cryst struct*)

1,5-Diguanidinopentane D-401
N,N'-1,5-Pentanediybisguanidine, 9CI.

1,1'-Pentamethylenediguanidine, 8CI.

Diamidinocadaverine. Audouine

[5070-04-2]

HN=C(NH₂)NH(CH₂)₅NHC(NH₂)=NH

C₇H₁₈N₆ 186.259

Alkaloid in the polychaete worm *Audouinia tentaculata*. Pale-yellow oil. Sol. H₂O.

Sulfate:

Needles. Mp 317-318°.



Picrate:

Yellow needles. Mp 226-227°.

[52780-74-2, 58585-48-1, 14279-79-9, 38142-20-0]

Roche, C. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1965, **260**, 7023



Robin, Y. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 3965

3,5-Di-5-hexenylpyrrolizidine D-402C₁₉H₃₃N 275.476**(3*R**,5*R**)-form [120091-06-7]**Isol. from the venom of the ant *Mono-*
morium smithii.Jones, T.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 375
(*isol*, *pmr*, *cmr*, *struct*)**Dihomosedinone D-403**1-[6-(2-Hydroxy-2-phenylethyl)-1-
methyl-2-piperidinyl]-2-pentanone, 9CI.
2-(2-Hydroxy-2-phenylethyl)-1-methyl-6-
(2-oxopentyl)piperidine
[145774-75-0]Absolute
ConfigurationC₁₉H₂₉NO₂ 303.444Trace alkaloid from *Sedum acre* (Cras-
sulaceae). Cryst. (hexane). Mp 67-68°.
[α]_D²⁰ -42 (c, 0.3 in CHCl₃).**6-Epimer: 6-Epidihomosedinone**


[145307-62-6]

C₁₉H₂₉NO₂ 303.444Alkaloid from *Sedum acre*.**Stereoisomer (?): 10-Phenyl-8-propyllo-
beliolone. 8-Propyl-10-phenyllobeliolone**


[66976-95-2]

C₁₉H₂₉NO₂ 303.444Alkaloid from *Sedum acre*. May be
identical with Dihomosedinone, 6-
Epidihomosedinone or a mixture.Francis, L.P.S. *et al.*, *Planta Med.*, 1977, **32**,
268 (10-Phenyl-8-propyllobeliolone)Durant, A. *et al.*, *Can. J. Chem.*, 1992, **70**,
2722 (Dihomosedinone, 6-
Epidihomosedinone)**Dihydroavicine D-404**5,6-Dihydro-5-methyl-1,3-benzodioxo-
lo[5,6-c][1,3]dioxolo[4,5-j]phenanthri-
dine, 9CI
[34490-82-9]C₂₀H₁₅NO₄ 333.343Alkaloid from the roots of *Toddalia*
asiatica and the root bark of *Zanthoxy-*
lum avicennae. Also prod. (together with
avicine, A-1566, when the latter is
liberated from its salts (Rutaceae). Prisms
(EtOH or C₆H₆/Et₂O). Mp 212-213°.*Hydrochloride*: Mp 255-258° (vac.,
block).8-Oxo: 5-Methyl-1,3-benzodioxolo[5,6-
c][1,3]dioxolo[4,5-j]phenanthridin-
6(5H)-one, 9CI. Oxyavicine
[7097-06-5]
C₂₀H₁₃NO₅ 347.326
Prod. by disproportionation of Avi-
cine, A-1566 when it is liberated from
its salts. Plates (CHCl₃/EtOH). Mp
278-283° (257-259° and 275-277°,
double Mp).Arthur, H.R. *et al.*, *J.C.S.*, 1959, 4007 (*synth*,
uv)Gopinath, K.W. *et al.*, *Tetrahedron*, 1961, **14**,
322 (*synth*, *deriv*)Hruban, L. *et al.*, *Coll. Czech. Chem. Comm.*,
1970, **35**, 3420 (*uv*)Ishii, H. *et al.*, *Tet. Lett.*, 1971, 2429 (*pmr*)Fish, F. *et al.*, *Phytochemistry*, 1975, **14**, 841
(*isol*, *ms*)Ninomiya, I. *et al.*, *J.C.S. Perkin I*, 1975, 762
(*synth*, *ir*, *deriv*)Sharma, P.N. *et al.*, *Indian J. Chem., Sect. B*,
1979, **17**, 299 (*isol*, *ms*)**3,4-Dihydro-2*H*-1,3-benzox-
azin-2-one, 9CI D-405****Luteanine**†
[1125-85-5]C₈H₇NO₂ 149.149Alkaloid extracted from *Reseda lutea*
(Resedaceae). Needles (H₂O). Mp 189-
191°.O'Sullivan, D.G. *et al.*, *J.C.S.*, 1957, 2916-2920
(*ir*)Hoover, F.W. *et al.*, *J.O.C.*, 1963, **28**, 1825
(*synth*)Nakhatov, I.K. *et al.*, *Khim. Prir. Soedin.*,
1977, **13**, 424; *Chem. Nat. Compd. (Engl.*
Transl.), 1977, **13**, 362 (*isol*, *ir*, *pmr*, *ms*,
struct)**Dihydrocadambine D-406**

[54483-84-0]

C₂₇H₃₄N₂O₁₀ 546.573Alkaloid from the heartwood of *Antho-*
cephalus cadamba, the leaves of *Nauclea*
latifolia and from *Nauclea diderrichii*
(Rubiaceae). Shows hypotensive activity
in laboratory animals. Amorph. [α]_D²⁵ -40
(MeOH).

▶ MT0588000

Penta-O-Ac: Mp 197-199°. [α]_D²⁵ -135
(MeOH).**3-Epimer: 3β-Dihydrocadambine**
[61989-80-8]C₂₇H₃₄N₂O₁₀ 546.573Alkaloid from the leaves of *Antho-*
cephalus cadamba (Rubiaceae).
Amorph. solid (as penta-Ac). [α]_D²⁵ -69(CHCl₃).Brown, R.T. *et al.*, *Tet. Lett.*, 1974, 1957; 1976,
2723 (*epimers*, *isol*, *struct*)Dimitrienko, G.I. *et al.*, *Tet. Lett.*, 1974, 1961
(*uv*, *ir*, *pmr*, *struct*)Hotellier, F. *et al.*, *Planta Med.*, 1979, **35**, 242
(*isol*)Aisaka, K. *et al.*, *Planta Med.*, 1985, 424**Dihydrochelerythrine D-407**12,13-Dihydro-1,2-dimethoxy-12-
methyl[1,3]benzodioxolo[5,6-c]phenan-
thridine, 9CI. Toddalinine. Dihydrotodda-
line
[6880-91-7]C₂₁H₁₉NO₄ 349.385Numbering systems vary. Alkaloid widely
distributed in the Papaveraceae (e.g.*Argemone mexicana*, *Chelidonium majus*,
Glaucium flavum var. *vestitum*, *Glauca-*
ium vitellinum), Rutaceae (e.g. *Fagara chaly-*
bea, *Fagara holstii*, *Fagara semiarticula-*
tum, *Fagara rubescens*, *Fagara*
macrophylla, *Toddalia aculeata*, *Toddalia*
asiatica, *Zanthoxylum tsihanimposa*,
Zanthoxylum senegalense, *Zanthoxylum*
coriaceum, *Zanthoxylum elephantiasis*)
and Papaveraceae (*Corydalis ledebouri-*
ana). Cryst. (EtOH, C₆H₆, or CHCl₃/
MeOH). Mp 169-171° (161-162°, 164-
165°). Probably an artifact. λ_{max} 230 (log
ε 4.62); 282 (log ε 4.72); 325 (sh) (log ε
4.28) (MeOH).**5-Methoxy: 5-Methoxydihydrocheleryth-**
rine. 12-Methoxydihydrochelerythrine†
C₂₂H₂₁NO₅ 379.412Alkaloid from *Bocconia integrifolia*
(Papaveraceae). V. pale yellow powder.
Mp 173-174.5°. [α]_D²⁰ -15 (c, 0.2 in
CHCl₃). Numbering systems vary. λ_{max}
219 (log ε 4.68); 288 (log ε 4.7); 323
(log ε 4.24); 355 (sh) (EtOH).**8-Hydroxy: 8-Hydroxydihydrocheleryth-**
rine

[4070-42-2]

[55438-12-5 (Ac)]

C₂₁H₁₉NO₅ 365.385Alkaloid from the roots of *Toddalia*
asiatica (Rutaceae). Cryst. (CH₂Cl₂/
Et₂O)(as Ac). Mp 145° (Ac).**8-Methoxy: 8-Methoxydihydrocheleryth-**
rine. Angoline. 8-O-Methyldihydroche-
lerythrine. Alkaloid P61. Chelerythrine
pseudomethanolate
[21080-31-9]C₂₂H₂₁NO₅ 379.412Alkaloid from *Bocconia arborea*, *Tod-*
dalia asiatica, *Hunnemannia fumariae-*
folia, *Fagara angolensis*, *Fagara*
tessmannii, *Fagara viridis*, *Fagara cha-*
lybea, *Fagara lepreurii*, *Fagara macro-*
phylla and *Fagara xanthoxyloides*
(Papaveraceae, Rutaceae). Cryst.

(CHCl₃/MeOH). Mp 210° (176°). [α]_D 0. Prob. artifact. λ_{max} 230 (ε 33000); 289 (ε 49300); 325 (ε 16000) (no solvent reported).

8-Methoxy, O¹⁰-de-Me: 9-Demethyl-8-methoxydihydrochelerythrine
[126234-22-8]

C₂₁H₁₉NO₅ 365.385

Alkaloid from *Zanthoxylum nitidum* (Rutaceae). Assigned R-config.

8-Ethoxy: 8-Ethoxydihydrochelerythrine. Chelerythrine pseudoethanolate. Alkaloid A₂. Artarine

[28974-35-8]

[125226-39-3]

C₂₃H₂₃NO₅ 393.438

Alkaloid from root bark of *Fagara xanthoxyloides* and *Fagara rubescens* and from roots of *Hunnemannia fumariaefolia* (Rutaceae, Papaveraceae). Cryst. (EtOH). Mp 200-202° Mp 239-242° (double Mp).

[α]_D 0. λ_{max} 228 (log ε 4.53); 284 (log ε 4.67); 320 (sh) (log ε 4.18) (95% EtOH).

12-Methoxy: 12-Methoxydihydrochelerythrine†. Dihydrochelilutine

[65427-95-4]

C₂₂H₂₁NO₅ 379.412

Minor alkaloid detected in the roots of *Chelidonium majus* (Papaveraceae).

Cryst. (CHCl₃/MeOH). Mp 132-135° (block) Mp 136-137° (capillary). λ_{max} 230 (log ε 4.52); 280 (log ε 4.56); 325 (log ε 4.21) (MeOH).

8-Oxo: 1,2-Dimethoxy-12-methyl[1,3]-benzodioxolo[5,6-c]phenanthridin-13(12H)-one. Oxychelerythrine. 8-Oxo-chelerythrine

[28342-33-8]

C₂₁H₁₇NO₅ 363.369

Alkaloid from *Zanthoxylum nitidum* and the bark of *Zanthoxylum arnotianum* (Rutaceae). Cryst. (MeOH). Mp 199-201°. λ_{max} 242 (ε 33900); 282 (sh) (ε 40800); 375 (ε 8500) (EtOH).

► FL9220000

8-Oxo, O¹⁰-de-Me: 2-Hydroxy-1-methoxy-12-methyl[1,3]-benzodioxolo[5,6-c]phenanthridin-13(12H)-one. 9-Demethyloxychelerythrine

C₂₀H₁₅NO₅ 349.342

Alkaloid from the root bark of *Zanthoxylum integrifolium*. Needles (CHCl₃/MeOH). Mp 224-226°. λ_{max} 207 (log ε 4.21); 240 (log ε 4.24); 280 (sh) (log ε 4.24); 287 (log ε 4.33); 327 (sh) (log ε 3.81); 375 (sh) (log ε 3.55) (EtOH).

Slavková, L. *et al.*, *Chem. Listy*, 1955, **49**, 1546-1549; *CA*, **50**, 4990h (isol)

Govindachari, T.R. *et al.*, *J.C.S.*, 1956, 769-771 (isol, synth)

Scheuer, P.J. *et al.*, *J.O.C.*, 1962, **27**, 1472-1473 (isol, synth)

Torto, F.G. *et al.*, *Tet. Lett.*, 1966, 181-183 (8-ethoxy, uv, ir, pmr, struct)

Desai, P.D. *et al.*, *Indian J. Chem.*, 1967, **5**, 41-42 (struct)

Slávic, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 1619-1623 (ms)

Fonzes, L. *et al.*, *Phytochemistry*, 1968, **7**, 1889-1890 (Angoline, uv, ir, pmr, ms)

MacLean, D.B. *et al.*, *Can. J. Chem.*, 1969, **47**, 1951-1956 (pmr, struct, 8-methoxy)

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1435-1439 (pmr)

Dadson, B.A. *et al.*, *J.C.S. Perkin I*, 1976, 146-147 (occur)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 2686-2693 (Dihydrochelerythrine, Dihydrochelilutine, isol, uv, synth)

Ishii, H. *et al.*, *Yakugaku Zasshi*, 1977, **97**, 890; *CA*, **87**, 197250g (Oxychelerythrine)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 864-873 (Dihydrochelilutine, struct)

Sharma, P.N. *et al.*, *Indian J. Chem., Sect. B*, 1979, **17**, 299-300 (Angoline)

Swinehart, J.A. *et al.*, *Phytochemistry*, 1980, **19**, 1219-1223 (isol)

Khusainova, Kh.Sh. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 670; *CA*, **96**, 65673y (isol)

Sharma, P.N. *et al.*, *Phytochemistry*, 1982, **21**, 252-253 (8-hydroxy)

Hanaoka, M. *et al.*, *J.C.S. Perkin I*, 1986, 2253-2256 (synth)

Chen, Y. *et al.*, *Huaxue Xuebao*, 1989, **47**, 1048; *CA*, **112**, 175616 (9-Demethyl-8-methoxydihydrochelerythrine)

Chen, Y. *et al.*, *Jiegou Huaxue*, 1989, **8**, 95-98; *CA*, **112**, 88707a (Artarine, cryst struct)

Oechslin, S.M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 519-524 (12-Methoxydihydrochelerythrine)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (N-15 nmr)

Harayama, T. *et al.*, *Synthesis*, 2001, 444-450 (12-Methoxydihydrochelerythrine, synth)


Watanabe, T. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 3024-3032 (12-Methoxydihydrochelerythrine, synth)

Le, T.N. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 118-120 (Oxychelerythrine, synth)

Chen, J.-I. *et al.*, *Planta Med.*, 2005, **71**, 470-475 (9-Demethyl-8-oxychelerythrine)

Dihydrochelerythrinyl-8-acetaldehyde **D-408**

12,13-Dihydro-1,2-dimethoxy-12-methyl[1,3]-benzodioxolo[5,6-c]phenanthridine-13-acetaldehyde, 9CI. Alkaloid ZT₂
[51876-09-6]



C₂₃H₂₁NO₅ 391.423

Various numbering schemes used. Probably an artifact. Alkaloid from the roots of *Glaucium flavum* var. *vestitum* and the bark of *Zanthoxylum tsihanimposa* (Papaveraceae, Rutaceae). Cryst. (CHCl₃/MeOH). Mp 206-210°. Opt. inactive.

2'-Carboxylic acid: 8-Carboxymethyl-dihydrochelerythrine

C₂₃H₂₁NO₆ 407.422

Alkaloid from the stem bark of a *Zanthoxylum* sp. collected in Grand Cayman (*Zanthoxylum spinosum* or *Zanthoxylum coriaceum*). Also reported to occur in *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae).

Decaudain, N. *et al.*, *Phytochemistry*, 1974, **13**, 505 (uv, ir, pmr, struct)


Castedo, L. *et al.*, *Heterocycles*, 1981, **16**, 533 (occur)

Ng, K.M. *et al.*, *Phytochemistry*, 1984, **26**, 3251 (8-Carboxymethyl-dihydrochelerythrine)

Dihydrocoptisine

D-409

[53777-78-9]



C₁₉H₁₅NO₄ 321.332

Alkaloid from seeds of *Fumaria indica*. Large yellow prisms (Me₂CO). Mp 194-196° dec. (175-179°).

1,8ξ-Dihydroxy: 1,8-Dihydroxy-2,3:9,10-bis(methylenedioxy)protoberberine

C₁₉H₁₅NO₆ 353.331

Alkaloid from the roots of *Thalictrum delavayi*. Brownish powder. [α]_D²⁰ +8.8 (c, 1.8 in CHCl₃). λ_{max} 214; 283; 378 (MeOH).

8-Oxo: 2,3:9,10-Bis(methylenedioxy)-8-oxo protoberberine. 8-Oxocoptisine

C₁₉H₁₃NO₅ 335.315

Alkaloid from *Coptis japonica* and *Thalictrum delavayi*. Yellow needles. Mp 282-284°. λ_{max} 224; 340; 377; 395 (sh) (MeOH).

Haworth, R.D. *et al.*, *J.C.S.*, 1926, 1769 (synth)

Jeffs, P.W. *et al.*, *J.O.C.*, 1975, **40**, 644 (synth, pmr)

Castedo, L. *et al.*, *Heterocycles*, 1986, **24**, 5 (synth)

Tripathi, Y.C. *et al.*, *Pharmazie*, 1987, **42**, 745 (isol)


Li, M. *et al.*, *Planta Med.*, 2001, **67**, 189-190 (Bis(methylenedioxy)oxoprotoberberine, Dihydroxybis(methylenedioxy)-protoberberine)

Min, Y.D. *et al.*, *Arch. Pharmacol. Res.*, 2006, **29**, 757-761 (8-Oxocoptisine)

Dihydrocorymine

D-410

Methyl 3,17-dihydroxy-1-methyl-2,4(1H)-cyclo-3,4-secoakummilan-16-carboxylate, 9CI
[65725-01-1]



Absolute Configuration

C₂₂H₂₈N₂O₄ 384.474

Appears to be unknown as a nat. alkaloid. Mp 222°. [α]_D -151 (c, 1 in MeOH). λ_{max} 258; 315 (no solvent reported).

Di-Ac: [65758-40-9]

Noncryst. [α]_D -94 (c, 1 in CHCl₃).

17-Aldehyde, 17→3 hemiacetal: Corymine. Corymine hemiacetal

[6472-42-0]
 $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from *Hunteria corymbosa*,
Hunteria umbellata and *Hunteria*
eburnea (Apocynaceae). Cryst. (propanol
 aq.). Mp 189-192° (182°). $[\alpha]_D^{18}$
 +27 (CHCl₃). $[\alpha]_D^{18}$ +48 (solv. not
 given). 17*S*-config. λ_{max} 215 (log ϵ
 4.25); 258 (log ϵ 4.04); 324 (log ϵ 3.65)
 (no solvent reported).

17-Aldehyde, hemiacetal, hydrobromide:
 Cryst. (H₂O). Mp 210°.

17-Aldehyde, hemiacetal, O-Ac:

O-Acetylcorymine

$C_{24}H_{28}N_2O_5$ 424.496

Alkaloid from *Hunteria umbellata*
 (Apocynaceae). Cryst. (propanol). Mp
 194-195°. $[\alpha]_D^{20}$ +0.4. The uv spectrum
 is identical to that reported for Cory-
 mine, which is unlikely. λ_{max} 215 (log
 ϵ 4.25); 258 (log ϵ 4.04); 324 (log ϵ 3.65)
 (no solvent reported).

17-Aldehyde, hemiacetal, N-de-Me: N-
 Demethylcorymine

[162585-01-5]

$C_{21}H_{24}N_2O_4$ 368.432

Alkaloid from leaves of *Hunteria*
zeylanica (Apocynaceae). Amorph.
 powder. $[\alpha]_D^{28}$ +12 (c, 1.25 in CHCl₃).

10-Methoxy, 3-deoxy: **Cabumamine**

[55857-18-6]

$C_{23}H_{30}N_2O_4$ 398.501

Alkaloid from *Cabucala erythrocarpa*
 (Apocynaceae). Mp 245-248°. $[\alpha]_D$ -
 110 (CHCl₃). λ_{max} 252; 330 (MeOH).

3-Epimer, 3-Epidihydrocorymine

[65758-39-6]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from leaves of *Hunteria*
elliotii and from *Hunteria congolana*
 (Apocynaceae). Needles (Me₂CO). Mp
 148°. $[\alpha]_D$ -70 (c, 1 in MeOH). λ_{max} 213
 ; 255; 315 (MeOH).

3-Epimer, N-de-Me: N-Demethylchita-
 mine. Norechitamine

[60048-88-6]

$C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from the root bark of *Alstonia*
scholaris (Apocynaceae). Off-white
 cryst. Mp 238-240°.

3-Epimer, N-de-Me, N-oxide: N-De-
 methylchitamine N-oxide. Norechita-
 mine N-oxide

[118957-11-2]

$C_{21}H_{26}N_2O_5$ 386.447

Alkaloid from the roots of *Winchia*
calophylla (syn. *Alstonia pachycarpa*)
 (Apocynaceae). Needles (MeOH). Mp
 216-219°. $[\alpha]_D^{20}$ -60 (c, 0.46 in EtOH).

3-Epimer, N-de-Me, N⁴-Me: **Echitamine**.
 Ditaine

[6871-44-9]

[6878-36-0]

$C_{22}H_{29}N_2O_4$ 385.482

Quaternary alkaloid from *Alstonia*
congensis, *Alstonia scholaris*, *Alstonia*
angustiloba, *Alstonia spectabilis*,
Alstonia verticillata, *Alstonia gilletii*,
Alstonia spatulata and *Alstonia*
neriifolia (Apocynaceae). DNA
 binding agent. Needles or short prisms
 (H₂O) (as chloride). Sol. H₂O, EtOH,

Et₂O; poorly sol. hexane. Mp 274° dec.
 (chloride). $[\alpha]_D^{15}$ -58 (c, 1 in H₂O)
 (chloride). λ_{max} 235 (ϵ 8510); 295 (ϵ
 3550) (MeOH).

► Highly toxic orally.

3-Epimer, 3-Ac: **3-Epidihydrocorymine 3-
 acetate**

[69734-92-5]

$C_{24}H_{30}N_2O_5$ 426.511

Alkaloid from *Hunteria zeylanica* and
Hunteria congolana. Cryst. (Et₂O). $[\alpha]_D$
 -76 (c, 0.7 in CHCl₃). λ_{max} 213 (log ϵ
 4.22); 253 (log ϵ 3.93); 308 (log ϵ 3.45)
 (MeOH).

3-Epimer, 17-Ac: **3-Epidihydrocorymine
 17-acetate**. O¹⁷-Acetyl-3-epidihydrocor-
 ymine

[82570-78-3]

$C_{24}H_{30}N_2O_5$ 426.511

Alkaloid from *Hunteria zeylanica*
 (Apocynaceae). Mp 260°. $[\alpha]_D$ -76 (c,
 0.7 in CHCl₃). λ_{max} 217 (log ϵ 4.33);
 260 (log ϵ 4.35); 317 (log ϵ 3.88)
 (MeOH).

3-Epimer, 17-Ac, N-de-Me: **17-O-Acetyl-
 N-demethylchitamine**

[123064-74-4]

$C_{23}H_{28}N_2O_5$ 412.485

Alkaloid from stem bark of *Alstonia*
glaucescens (Apocynaceae). Amorph.

3-Epimer, 17-Ac, N-de-Me, N⁴-Me: **17-
 O-Acetylcorymine**

[132937-97-4]

$C_{24}H_{31}N_2O_5$ 427.519

Alkaloid from the bark of *Alstonia*
scholaris (Apocynaceae).

3-Epimer, di-Ac: **Di-O-acetyl-3-epidithy-
 drocorymine**

[69761-86-0]

$C_{26}H_{32}N_2O_6$ 468.549

Alkaloid from *Hunteria congolana*
 seeds (Apocynaceae). Mp 192°. $[\alpha]_D$ -
 48 (c, 0.5 in CHCl₃).

3-Epimer, parent acid, N-de-Me, N⁴-Me:
Echitaminic acid

[7163-44-2]

$C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from the bark of *Alstonia*
glaucescens and *Alstonia scholaris*.
 Cryst. (MeOH). Mp 218-220°. $[\alpha]_D$ -
 42.3 (c, 0.11 in MeOH). Zwitterionic.

16-Epimer, 3-deoxy, N-de-Me: **Vinco-
 phylline**

[958247-23-9]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from the stem bark of *Kopsia*
singaporensis. Pale yellow oil. $[\alpha]_D^{25}$
 +102 (c, 0.1 in CHCl₃). λ_{max} 206 (log ϵ
 4.36); 241 (log ϵ 3.75); 296 (log ϵ 3.3)
 (EtOH).

Hamilton, J.A. et al., *J.C.S.*, 1962, 5061-5075
 (*Echitamine, cryst struct*)

Kiang, A.K. et al., *Proc. Chem. Soc., London*,
 1962, 298-300 (*Corymine, Dihydrocorymine,*
struct)

Bevan, C.W.L. et al., *Chem. Ind. (London)*,
 1965, 603-604 (*Corymine, cryst struct*)

Talapatra, S.K. et al., *J. Indian Chem. Soc.*,
 1967, **44**, 639-640 (*Echitamine, isol*)

Bevan, C.W.L. et al., *Tetrahedron*, 1967, **23**,
 3809 (*Corymine, Acetylcorymine, isol, pmr,*
ms, uv)

Kompiš, I. et al., *Coll. Czech. Chem. Comm.*,
 1968, **33**, 4328-4336 (*Corymine, ord*)

Roy, S.K. et al., *J. Indian Chem. Soc.*, 1968, **45**,
 21-22 (*Echitamine, isol*)

Govindachari, T.R. et al., *J. Indian Chem.*
Soc., 1968, **45**, 945-957 (*Echitamine, rev, uv,*
ir)

Douzoua, L. et al., *Phytochemistry*, 1974, **13**,
 1994-1995 (*Cabumamine, isol, uv, ir, pmr, ms*)

Mansour, M. et al., *Phytochemistry*, 1974, **13**,
 2861-2863 (*Cabumamine, struct, abs config*)

Morfaux, A.M. et al., *Phytochemistry*, 1978,
17, 167 (*3-Epidihydrocorymine, isol, uv, ir,*
pmr, ms, struct)

Vercauteren, J. et al., *Bull. Soc. Chim. Fr., Part*
2, 1982, 291-296 (*3-*

Acetylepithydrocorymine)

Lavaud, C. et al., *Phytochemistry*, 1982, **21**,
 445-447 (*17-Acetylepithydrocorymine*)

Chen, W.M. et al., *Planta Med.*, 1988, **54**, 480-
 481 (*N-Demethylchitamine N-oxide*)

Yamauchi, T. et al., *Phytochemistry*, 1990, **29**,
 3321-3325 (*17-Acetylcorymine*)

Subhadhirasakul, S. et al., *Chem. Pharm.*
Bull., 1994, **42**, 2645-2646 (*N-*

Demethylcorymine)

Keavpradub, N. et al., *Phytochemistry*, 1994,
37, 1745-1749 (*17-Acetyl-N-*

demethylchitamine, Echitaminic acid)


Salim, A.A. et al., *J. Nat. Prod.*, 2004, **67**,
 1591-1594 (*Echitaminic acid*)

Subramaniam, G. et al., *J. Nat. Prod.*, 2007,
70, 1783-1789 (*Vincophylline*)

Sax, N.I. et al., *Dangerous Properties of*
Industrial Materials, 5th edn., Van Nostrand
 Reinhold, 1979, 630

**Dihydrocorynantheine pseu- D-411
 doindoxyl**

[57800-43-8]



$C_{22}H_{28}N_2O_4$ 384.474


Alkaloid from *Uncaria attenuata* (Ru-
 biaceae). Yellow.

Phillipson, J.D. et al., *Phytochemistry*, 1975,
14, 1855 (*isol, ms, struct, synth*)

Dihydrocorynantheol D-412

[2270-72-6]

[23107-08-6 ((±)-form)]



$C_{19}H_{26}N_2O$ 298.427

Alkaloid from *Aspidosperma marcgra-
 vianum*, *Aspidosperma auriculatum*, *Am-
 sonia tabernaemontana*, *Mitragyna*
parvifolia, *Hunteria zeylanica*, *Ochrosia*
moorei, *Rhazya stricta* and others
 (Apocynaceae, Rubiaceae). Shows

activity against gram-positive bacteria. Cryst. (MeOH aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 185–186°. [α]_D²⁷ -18 (c, 0.92 in CHCl₃). [α]_D²³ -37 (c, 0.938 in Py). λ_{\max} 226 (log ϵ 4.56); 281 (log ϵ 3.87); 290 (sh) (log ϵ 3.8) (EtOH).

N⁴-Oxide: Dihydrocorynantheol N-oxide
[52992-34-4]
C₁₉H₂₆N₂O₂ 314.427

Alkaloid from the leaves of *Mitragyna parvifolia* and the stem bark of *Aspidosperma marcgravianum* (Rubiaceae, Apocynaceae). Cream-coloured substance.

N⁴-Me: Dihydrocorynantheol N-metho salt
C₂₀H₂₉N₂O[⊕] 313.462
Quaternary alkaloid from the root and stem bark of *Hunteria eburnea* (Apocynaceae). Cryst. (MeOH/Et₂O) (as chloride). Mp 272–273° dec. (296–297°) (chloride). [α]_D²¹ +63.3 (c, 0.366 in 72.5% MeOH aq.). [α]_D +101 (solv. unspecified). CAS no. not found to 2001.

Ac: Mp 125–128°.

Ac; hydrochloride:

Yellow needles (MeOH/Et₂O). Mp 253–260° dec.

18,19-Didehydro: Corynantheol

[23443-70-1]

C₁₉H₂₄N₂O 296.411

Alkaloid from the stem bark and leaves of *Rauwolfia cumminsii* and from the leaves of *Rauwolfia mombasiana* (Apocynaceae). Cryst. (MeOH). Mp 212–213°. [α]_D -61 (Py).

18,19-Didehydro, aldehyde: Corynantheal
[572-78-1]

C₁₉H₂₂N₂O 294.396

Alkaloid from the leaves of *Rauwolfia cumminsii* (Apocynaceae). Cryst. (C₆H₆/CH₂Cl₂). Mp 226–229° (230° subl.). [α]_D -48.5 (Py).

19,20-Didehydro: see Geissoschizol, G-33

10-Hydroxy: 10-Hydroxydihydrocorynantheol

C₁₉H₂₆N₂O₂ 314.427

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). Needles + ¼H₂O (50% MeOH aq.). Mp 225–226° dec. [α]_D²³ +6.4 (c, 2.061 in EtOH). λ_{\max} 224; 280; 298 (sh); 310 (sh) (EtOH). λ_{\max} 277; 297 (sh); 307 (sh); 319 (sh); 343 (sh) (EtOH/NaOH).

10-Hydroxy, N⁴- β -Me: Ochrosandwine

[20361-97-1]

C₂₀H₂₉N₂O₂[⊕] 329.461

Quaternary alkaloid from *Ochrosia sandwichensis* (Apocynaceae). Cryst. (H₂O or MeOH/C₆H₆) (as chloride). Mp 288–289° dec. (evac. tube) (chloride) Mp 261–262° (block) (chloride). λ_{\max} 267 (sh) (log ϵ 3.87); 275 (log ϵ 3.92); 299 (log ϵ 3.65); 309 (sh) (log ϵ 3.58) (MeOH) (chloride).

10-Hydroxy, 18,19-didehydro, N⁴- α -Me:

10-Hydroxy-N⁴-methylcorynantheol

[116521-74-5]

C₂₀H₂₇N₂O₂[⊕] 327.445

Quaternary alkaloid from the bark of

Strychnos usambarensis (Loganiaceae).

19S-Hydroxy: 19S-Hydroxydihydrocorynantheol

[155310-31-9]

C₁₉H₂₆N₂O₂ 314.427

Minor alkaloid from roots of *Antirhea portoricensis* (Rubiaceae). [α]_D²⁵ -3 (c, 0.6 in CDCl₃).

10-Methoxy: 10-Methoxydihydrocorynantheol. Alkaloid AD-IV

[15266-55-4]

C₂₀H₂₈N₂O₂ 328.453

Alkaloid from *Aspidosperma* sp. (No. 119070), *Aspidosperma discolor*, *Ochrosia confusa*, *Ochrosia moorei*, *Ochrosia oppositifolia*, *Neisosperma glomerata*, *Neisosperma kilneri* and others (Apocynaceae). Cryst. (Me₂CO, Me₂CO/Et₂O or MeOH). Mp 167–168° dec. [α]_D²³ -1.4 (c, 1.48 in CHCl₃). [α]_D²⁴ -16.3 (c, 1.55 in Py). λ_{\max} 227 (log ϵ 4.38); 280 (log ϵ 3.86); 293 (sh) (log ϵ 3.82) (EtOH).

10-Methoxy, hydrochloride:

Cryst. (MeOH/Et₂O). Mp 256–261° dec.

10-Methoxy, picrate:

Cryst. (MeOH/Et₂O). Mp 145–147° dec.

10-Methoxy, N⁴- α -oxide: 10-Methoxydihydrocorynantheol 4 α -N-oxide

[77794-99-1]

C₂₀H₂₈N₂O₃ 344.453

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{\max} 278; 295 (sh); 305 (sh) (EtOH).

10-Methoxy, N⁴- β -oxide: 10-Methoxydihydrocorynantheol 4 β -N-oxide

[77794-98-0]

C₂₀H₂₈N₂O₃ 344.453

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{\max} 280; 294 (sh); 307 (sh) (EtOH).

10-Methoxy, 18,19-didehydro: 10-Methoxycorynantheol

[56053-12-4]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from leaves of *Ochrosia alyxioides* (Apocynaceae). Noncryst. [α]_D²⁰ +5 (c, 0.2 in MeOH).

10-Methoxy, 18,19-didehydro, N⁴- α -Me:

10-Methoxycorynantheol α -N-metho salt

[92446-38-3]

C₂₁H₂₉N₂O₂[⊕] 341.472

Quaternary alkaloid from the bark of *Neisosperma glomerata* (Apocynaceae). Cryst. (MeOH) (as chloride). Mp 237° (chloride). [α]_D²⁰ +45 (c, 1 in MeOH).

10-Methoxy, 18,19-didehydro, N⁴- β -Me:

10-Methoxycorynantheol β -N-metho salt

[55322-92-4]

C₂₁H₂₉N₂O₂[⊕] 341.472

Quaternary alkaloid from the bark of *Ochrosia nakaiana* (Apocynaceae).

11-Methoxy: Ochroianine. 11-Methoxydihydrocorynantheol

[55322-94-6]

[126783-55-9 ((\pm)-form)]

C₂₀H₂₈N₂O₂ 328.453

Alkaloid from the bark of *Ochrosia miana* (Apocynaceae). Mp 162–169° dec. (synthetic). [α]_D²⁰ -15 (c, 1 in EtOH). [α]_D²⁸ -28 (c, 1.00 in EtOH) (synthetic). λ_{\max} 228 (c, 37100); 270 (c, 4840); 298 (c, 6180) (EtOH).

10,11-Dimethoxy: Ochroprosinine

[37687-31-3]

[107887-35-4 ((\pm)-form)]

C₂₁H₃₀N₂O₃ 358.48

Alkaloid from the bark of *Ochrosia oppositifolia*, *Ochrosia vieillardii* and *Ochrosia moorei* and from the bark and leaves of *Neisosperma glomerata* (Apocynaceae). Amorph. [α]_D²⁰ -18 (c, 1 in CHCl₃). λ_{\max} 227 (c, 28400); 282 (sh) (c, 5630); 299 (c, 9420); 303 (c, 9490); 309 (sh) (c, 8120) (EtOH).

3-Epimer, N⁴- β -Me: Lercheine. 3-Epidihydrocorynantheol N-metho salt

[146075-41-4 (chloride)]

C₂₀H₂₉N₂O[⊕] 313.462

Quaternary alkaloid from leaves of *Lerchea bracteata* (Rubiaceae). Mp 289–291° (as chloride). [α]_D²³ +100 (c, 0.2 in MeOH) (chloride).

3-Epimer, 18,19-didehydro, stereoisomer:

Normelinonine B

[84024-87-3]

C₁₉H₂₄N₂O 296.411

Alkaloid from the root bark of *Strychnos nux-vomica* (Loganiaceae). Stereochem. only partly known. λ_{\max} 231; 273; 279; 281; 289 (MeOH).

3-Epimer, 18,19-didehydro, N⁴-Me, stereoisomer: Melinonine B

[6801-38-3]

C₂₀H₂₇N₂O[⊕] 311.446

Quaternary alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Needles (MeOH or H₂O) (as chloride). Mp 311° dec. (305–306°) (chloride). [α]_D²¹ -14.8 (c, 0.272 in 27.5% MeOH aq.) (chloride). Stereochem. only partly known.

20-Epimer: Corynantheidol

[483-27-2]

[4695-99-2 ((\pm)-form)]

C₁₉H₂₆N₂O 298.427

Alkaloid from the leaves of *Mitragyna parvifolia* (Rubiaceae). Cryst. (CH₂Cl₂/petrol). Mp 183–186° dec. [α]_D²² -99.1 (c, 0.464 in Py).

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1951, 588–602 (*Corynantheal, synth, ir*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 29–45 (*Melinonine B, isol, uv*)

Van Tamelen, E.E. *et al.*, *Chem. Ind. (London)*, 1956, 793 (*Corynantheal, stereochem*)

Vamvacas, C. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 1793–1808 (*Dihydrocorynantheol N-metho salt, Corynantheidol, Melinonine B*)

Gilbert, B. *et al.*, *J.O.C.*, 1962, **27**, 4702–4704 (*Dihydrocorynantheol, isol, uv, ir*)

Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 1445–1449 (*Dihydrocorynantheol N-metho salt*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1965, **87**, 5461–5467 (*Corynantheidol, synth*)

Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141–1166 (*10-Methoxydihydrocorynantheol, isol, uv, ir, ms*)

Jordan, W. *et al.*, *Tetrahedron*, 1965, **21**, 3731–3740 (*Ochrosandwine, isol, uv, ir, pmr, struct*)

Dastoor, N.J. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 213-231 (*Dihydrocorynantheol*, 10-Methoxydihydrocorynantheol, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Ziegler, F.E. *et al.*, *Tet. Lett.*, 1969, 1097-1100 (*Dihydrocorynantheol*, 3-Epidihydrocorynantheol *N*-metho salt, *synth*)

Sawa, Y.K. *et al.*, *Tetrahedron*, 1969, **25**, 5329-5337 (*10-Hydroxydihydrocorynantheol*, *Ochrosandwine*, *Dihydrocorynantheol*, *synth*, *uv*, *pmr*, *cd*)

Peube-Locou, N. *et al.*, *Phytochemistry*, 1972, **11**, 2109-2111 (*Ochroprosinine*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Shellard, E.J. *et al.*, *Planta Med.*, 1973, **24**, 13-17; 341-352; 1974, **25**, 172-174 (*Corynantheidol*, *Dihydrocorynantheol N*-oxide, *isol*, *uv*, *pmr*, *ms*, *ord*, *biosynth*)

Preaux, N. *et al.*, *Phytochemistry*, 1974, **13**, 2607-2609 (*Ochromianine*, *uv*, *ir*, *pmr*, *struct*)

Sakai, S. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 1274; *CA*, **82**, 43633u (*10-Methoxycorynantheol β-N-methosalt*)

Djakouré, L.A. *et al.*, *Tetrahedron*, 1975, **31**, 2247-2254 (*Corynantheol*, *Corynantheal*, *synth*, *ir*, *pmr*, *ms*)

Iwu, M.M. *et al.*, *Planta Med.*, 1978, **33**, 232-236; 360-364 (*Corynantheol*, *Corynantheal*, *isol*)

Ahond, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 193-199 (*Ochroprosinine*, 10-Hydroxydihydrocorynantheol, 10-Methoxydihydrocorynantheol *N*-oxides)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1981, 3168-3175 (*synth*, *ir*, *ms*)

Başer, K.H.C. *et al.*, *Phytochemistry*, 1982, **21**, 1423-1429 (*Normelinone B*)

Imanishi, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 1551-1560 (*Corynantheidol*, *synth*, *ir*, *pmr*, *ms*)

Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694-707 (3,4,5,6-Tetradehydrodihydrocorynantheol)

Brown, R.T. *et al.*, *Chem. Comm.*, 1984, 847-848 (*synth*)

Seguin, E. *et al.*, *J. Nat. Prod.*, 1984, **47**, 687-691 (*10-Methoxycorynantheol α-N-methosalt*)

Danieli, B. *et al.*, *J.C.S. Perkin I*, 1984, 1237-1240 (*synth*, *uv*, *ir*, *cmr*, *ms*)

Suzuki, T. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1584-1588 (*synth*)

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1986, **52**, 73-74 (*cmr*)

Quetin-Leclercq, J. *et al.*, *Phytochemistry*, 1988, **27**, 1923-1926 (*10-Hydroxy-N^b-methylcorynantheol*)

Boughandjioua, N. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1107-1112 (*10-Methoxycorynantheol*)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1200-1204; 1991, **33**, 75-78 (*Ochromianine*, *Ochroprosinine*, *synth*, *cd*, *struct*, *abs config*)

Lounasmaa, M. *et al.*, *Heterocycles*, 1990, **31**, 1351-1358 (*Corynantheidol*, *synth*)

Ihara, M. *et al.*, *J.C.S. Perkin I*, 1990, 2771-2773 (*synth*)

Ohba, M. *et al.*, *Heterocycles*, 1991, **32**, 319 (*synth*)

Beard, R.L. *et al.*, *J.O.C.*, 1991, **56**, 2091-2096 (*Corynantheidol*, *synth*)

Arbain, D. *et al.*, *J.C.S. Perkin I*, 1992, 3039-3042 (*Lercheine*)

Diez, A. *et al.*, *Tet. Lett.*, 1993, **34**, 733-736 (*synth*)

Weniger, B. *et al.*, *J. Nat. Prod.*, 1994, **57**, 287-290 (*19S-Hydroxydihydrocorynantheol*)

Meittinen, J. *et al.*, *Planta Med.*, 1996, **62**, 42-45 (*10-Methoxydihydrocorynantheol*, 10-Methoxycorynantheol, *synth*)


Tosaka, A. *et al.*, *Heterocycles*, 2006, **70**, 153-159 (*synth*)

Deiters, A. *et al.*, *J.O.C.*, 2006, **71**, 6547-6561 (*synth*)

Itoh, T. *et al.*, *Org. Lett.*, 2006, **8**, 1533-1535 (*synth*)

Dihydrocuscohygrine D-413

1-Methyl-α-[1-methyl-2-pyrrolidinyl]-methyl]-2-pyrrolidineethanol, 9CI. 1,3-Bis(1-methyl-2-pyrrolidinyl)-2-propanol [80408-57-7]



Relative configuration


C₁₃H₂₆N₂O 226.361
Alkaloid from leaves of *Erythroxylum coca* and other *Erythroxylum* spp. Yellow oil. [α]_D²⁴ -105.5 (c, 1.67 in Me₂CO). The opt. activity of the nat. alkaloid fixes its rel. stereochem. as shown. It corresponds to the unknown opt. active form of cuscohygrine in C-818.

[58131-40-1, 80408-55-5, 80408-56-6, 80366-09-2]

Turner, C.E. *et al.*, *Phytochemistry*, 1981, **20**, 1403-1405 (*isol*, *synth*, *ir*, *pmr*, *ms*)
El-Imam, Y.M.A. *et al.*, *Phytochemistry*, 1985, **24**, 2285-2289 (*occur*)
Stapper, C. *et al.*, *J.O.C.*, 2002, **67**, 6456-6460 (*synth*)
Yamauchi, T. *et al.*, *J.O.C.*, 2008, **73**, 9484-9487 (*synth*, *pmr*, *cmr*)

Dihydrocycloakagerine D-414

[113141-71-2]




C₂₀H₂₄N₂O 308.422
Alkaloid from the root bark and stem bark of *Strychnos johnsonii* (Loganiaceae). [α]_D -239 (c, 0.33 in MeOH).

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

5,12-Dihydrocycloocta[1,2-b:5,6-b']diindole-6,13-dicarboxylic acid, 9CI D-415

Caulerpinic acid. *Caulerpin diacid* [28056-19-1]



C₂₂H₁₄N₂O₄ 370.364
Struct. revised in 1978, formerly assigned as a dibenzophenazine. Alkaloid from the green alga *Caulerpa racemosa*. Dark brown cryst. Mp 256° subl.
Mono-Me ester: **Monomethyl caulerpinate**

[137761-17-2]
C₂₃H₁₆N₂O₄ 384.39
Alkaloid from *Caulerpa racemosa*. Orange-yellow needles. Mp 158-161°.


Di-Me ester: Caulerpin

[26612-48-6]
C₂₄H₁₈N₂O₄ 398.417
Pigment from green algae *Caulerpa serrulata*, *Caulerpa racemosa*, *Caulerpa sertularioides*, *Caulerpa taxifolia*, *Caulerpa cupressoides*, *Caulerpa scalpelliformis*, *Caloglossa lepreurii* and *Laurencia majuscula*. Plant growth regulator, phycotoxin, vermifuge. Red prisms or needles (Et₂O or Me₂CO). Fairly sol. Et₂O, Me₂CO; poorly sol. H₂O. Mp 317°. λ_{max} 222 (ε 50000); 270 (ε 27000); 292 (ε 29000); 317 (ε 35000) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) 200 - 400 mg/kg, LD₅₀ (mus, orl) 2000 - 4000 mg/kg.
N,N'-Di-Me, di-Me ester: [65079-49-4]
Red needles (CHCl₃/petrol). Mp 262-263°.

Aguilar-Santos, S. *et al.*, *J.C.S.(C)*, 1970, 842-843 (*isol*, *uv*, *ir*, *pmr*, *ms*)
Maiti, B.C. *et al.*, *J. Chem. Res., Synop.*, 1978, 126-127; *J. Chem. Res., Miniprint*, 1978, 1683-1693 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)
Schwede, J.G. *et al.*, *Phytochemistry*, 1987, **26**, 155-158 (*occur*, *Caulerpin*)
Anjaneyulu, A.S.R. *et al.*, *Phytochemistry*, 1991, **30**, 3041-3042 (*mono Me-ester*)
Govenkar, M.B. *et al.*, *Phytochemistry*, 2000, **54**, 979-981 (*Caulerpin*, *isol*, *cmr*)


1,3-Dihydrocyclopentapyridin-2-one D-416



C₇H₆N₂O 134.137
Tentative struct. *Isol.* from a sponge *Haliclona* sp. Shows antibiotic props. Mp 260-265°. No CAS no. assigned. No further work reported to 2001.
Stempien, M.F. *et al.*, *CA*, 1977, **86**, 95905 (*isol*, *struct*)

1,2-Dihydro-1,2-dihydroxy- D-417

1,2,3,12-Tetrahydro-1,2-dihydroxy-6-methoxy-3,3,12-trimethyl-7H-pyrano[2,3-c]acridin-7-one, 9CI. 1,2-Dihydroxy-1,2-dihydroacronycine



(1*R*,2*R*)-form

C₂₀H₂₁NO₅ 355.39

(1*R*,2*R*)-form
(-)-cis-form

[107882-25-7]

Alkaloid from the stem bark of *Sarcomelicope glauca* (Rutaceae). Needles (MeOH). Mp 232-234°. $[\alpha]_D^{20}$ -10 (c, 0.2 in MeOH) (natural). $[\alpha]_D$ -37.8 (c, 0.5 in MeOH) (synthetic).

N-De-Me: **1,2-Dihydro-1,2-dihydroxy-N-demethylacronycine**

[110883-41-5]

C₁₉H₁₉NO₅ 341.363

Alkaloid from the leaves of *Sarcomelicope dogniensis* (Rutaceae). Yellow amorph. compd. $[\alpha]_D^{20}$ -2 (c, 0.25 in MeOH).

(1R*,2S*)-form

(-)-trans-form

[107882-15-5]

Alkaloid from the stem bark of *Sarcomelicope glauca* (Rutaceae). Noncryst. $[\alpha]_D^{20}$ -6 (c, 0.7 in MeOH). λ_{\max} 230 (ε 8510); 269 (sh) (ε 16600); 276 (ε 20000); 298 (ε 5500); 320 (ε 3720); 384 (ε 3310) (MeOH) (Derep).

2-Deoxy, N-de-Me: **N-Desmethyl-1,2-dihydro-1-hydroxyacronycine**. 1-Hydroxy-1,2-dihydro-N-desmethylacronycine

[110883-42-6]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the leaves of *Sarcomelicope dogniensis* (Rutaceae). Light yellow prisms (MeOH). Mp 212-214°. $[\alpha]_D^{20}$ +4 (c, 0.25 in MeOH).

2-Deoxy, 1-ketone, N-de-Me: **1-Oxo-1,2-dihydro-N-desmethylacronycine**

[110883-43-7]

C₁₉H₁₇NO₄ 323.348

Alkaloid from the leaves of *Sarcomelicope dogniensis* (Rutaceae). Yellow amorph. solid.

2-Deoxy, 1-ketone, N-de-Me, N-hydroxy: **N-Demethyl-1,2-dihydro-N-hydroxy-1-oxoacronycine**. 1-Oxo-1,2-dihydro-N-demethyl-N-hydroxyacronycine

[110883-44-8]

C₁₉H₁₇NO₅ 339.347

Alkaloid from leaves of *Sarcomelicope dogniensis*. Amorph. λ_{\max} 211 (log ε 4.13); 257 (log ε 4.29); 278 (log ε 4.21); 306 (log ε 3.86); 381 (log ε 3.79) (MeOH).

Mitaku, S. et al., *J. Nat. Prod.*, 1986, **49**, 1091 (isol, uv, ir, pmr, ms, struct)


Mitaku, S. et al., *Heterocycles*, 1987, **26**, 2057-2063 (derivs)

Mitaku, S. et al., *Nat. Prod. Lett.*, 1995, **7**, 219 (deriv)

Costes, N. et al., *J. Nat. Prod.*, 1999, **62**, 490-492 (synth, abs config)

4,5-Dihydro-1,9-dihydroxy- D-418**2,11-dimethoxy-7-oxoaporphine**

4,5-Dihydro-1,9-dihydroxy-2,11-dimethoxy-7H-dibenzo[de,g]quinolin-7-one


C₁₈H₁₅NO₅ 325.32

Alkaloid from the leaves and stems of *Milusa cuneata*. Purple-black cryst. (MeOH). Mp >320°. λ_{\max} 216 ; 282 ; 339 (MeOH).

Chen, B. et al., *Nat. Prod. Res.*, 2003, **17**, 397-402 (isol, pmr, cmr, ms)

2,3-Dihydro-5,6-dihydroxy- D-419**1,1-dimethylindolium(1+)**

[113679-36-0]


C₁₀H₁₄NO₂[⊕] 180.226

Isol. from a deep water marine sponge *Dercitus* sp. Cryst. + 1H₂O (MeOH/CHCl₃) (as chloride). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 244° (chloride). CAS no. refers to chloride. λ_{\max} 206 (ε 12200); 289 (ε 3700) (MeOH) (Berdy). λ_{\max} 209 (ε 16500); 248 (ε 5800); 303 (ε 5100) (MeOH/NaOH) (Berdy).

Kohmoto, S. et al., *Experientia*, 1988, **44**, 85 (isol, uv, ir, pmr, cmr, ms, struct)

3,4-Dihydro-3,4-dihydroxy- D-420**2-(hydroxymethyl)-2H-pyrrole**

3,4-Dihydro-2-(hydroxymethyl)-2H-pyrrole-3,4-diol, 9CI



(2R,3R,4R)-form

C₅H₉NO₃ 131.131**(2R,3R,4R)-form**

Nectrisine. FR 900483. Antibiotic FR 900483

[108692-47-3]

Isol. from *Nectria lucida*. Hypoglycaemic and immunostimulant. Inhibits α-glucosidase and α-mannosidase. Powder. Mp 75° dec. $[\alpha]_D^{23}$ +22 (c, 0.55 in H₂O). λ_{\max} 320 (ε 100) (H₂O) (Derep).

Japan. Pat., 1987, 87 36 355; *CA*, **107**, 5742 (isol, props)

Shibata, T. et al., *J. Antibiot.*, 1988, **41**, 296-301 (isol, struct)

Chen, S.-H. et al., *Tet. Lett.*, 1990, **31**, 2229-2232 (synth)

Kayakiri, H. et al., *Chem. Pharm. Bull.*, 1991, **39**, 2807-2812 (pmr, cmr, synth)

Kogoshi, N. et al., *Heterocycles*, 1998, **49**, 63-66 (synth)

Kim, Y.J. et al., *Tetrahedron*, 1999, **55**, 8353-8364 (synth, activity)

Bosco, M. et al., *Tet. Lett.*, 2001, **42**, 7949-7952 (synth)


Hulme, A.N. et al., *Tet. Lett.*, 2003, **44**, 7649-7653 (synth)

Merino, P. et al., *Eur. J. Org. Chem.*, 2008, 2929-2947 (synth)

Ribes, C. et al., *J.O.C.*, 2008, **73**, 7779-7782 (synth)

3,4-Dihydro-6,7-dihydroxy- D-421**isoquinoline**

3,4-Dihydro-6,7-isoquinolinediol

C₉H₉NO₂ 163.176

6-Me ether, N-Me: **3,4-Dihydro-7-hydroxy-6-methoxy-2-methylisoquinolinium(1+)**, 9CI. **Pycnarrhine**. 1,2-Dehydrocorypalline

[72142-82-6]

C₁₁H₁₄NO₂[⊕] 192.237

Alkaloid from *Pycnarrhena longifolia*, *Corydalis ophiocarpa*, *Corydalis stricta*, *Arcangelisia flava* and *Xylopiavieillardii*.

6-Me ether, N-Me, chloride: [69734-98-1] C₁₁H₁₄ClNO₂ 227.69

Glistening yellow platelets (MeOH/Me₂CO). Mp 187° (dec.) (sinters at 155°).

6-Me ether, N-Me, iodide: [21885-30-3]

C₁₁H₁₄INO₂ 319.142

Yellow cryst. (MeOH). Mp 216-218° (211-212.5°).

7-Me ether, N-Me: **3,4-Dihydro-6-hydroxy-7-methoxy-2-methylisoquinolinium(1+)**, 9CI. **Isopycnarrhine**

[107882-22-4]

C₁₁H₁₄NO₂[⊕] 192.237

Alkaloid from the bark of *Popowia piscarpa* (Annonaceae). Oil. Prob. isol. as the internal salt with the phenolic OH group (≡ hydroxide).

Di-Me ether: **3,4-Dihydro-6,7-dimethoxyisoquinoline**, 9CI. **Dehydroheliamine**

[3382-18-1]

C₁₁H₁₃NO₂ 191.229

Trace alkaloid from *Carnegiea gigantea* (Cactaceae). Cryst. (EtOH/Et₂O) (as hydrochloride). Mp 194-196° (hydrochloride).

Di-Me ether, N-Me: **3,4-Dihydro-6,7-dimethoxy-2-methylisoquinolinium(1+)**

C₁₂H₁₆NO₂[⊕] 206.264

Alkaloid from the roots of *Xylopiaparviflora*. Cryst. (as perchlorate). Mp 184.5-187.5° (perchlorate). λ_{\max} 248 (log ε 3.92); 309 (log ε 3.59); 361 (log ε 3.52) (MeOH) (perchlorate).

Akabori, S. et al., *Bull. Chem. Soc. Jpn.*, 1926, **1**, 125 (*Pycnarrhine*, synth)

Doskotch, R.W. et al., *Tetrahedron*, 1969, **25**, 469 (*Pycnarrhine*, synth)

Brossi, A. et al., *Org. Prep. Proced. Int.*, 1970, **2**, 281 (*Pycnarrhine*, synth)

Siwon, J. et al., *Phytochemistry*, 1981, **20**, 323 (*Pycnarrhine*, isol, uv, pmr, cmr, ms, synth)


Verpoorte, R. et al., *J. Nat. Prod.*, 1982, **45**, 582 (*Pycnarrhine*, isol, uv, pmr, cmr, ms)

Ordaz, C. et al., *Phytochemistry*, 1983, **22**, 2101 (*Dehydroheliamine*)

Jossang, A. et al., *J. Nat. Prod.*, 1986, **49**, 1028 (*Isopycnarrhine*)

Nishiyama, Y. et al., *Phytochemistry*, 2004, **65**, 939-944 (*di-Me ether N-Me*)

3,4-Dihydro-4,5-dihydroxy-3-methoxy-4-(4-methoxyphenyl)-2(1H)-quinolinone D-422
[184046-65-9]



Relative Configuration

C₁₇H₁₇NO₅ 315.325

Prod. by *Penicillium* sp. NTC-47 and *Penicillium* cf. *simplicissimum*. Insecticidal agent. Prisms (MeOH). Mp 208-210°. [α]_D²⁰ -55 (c, 0.02 in MeOH). λ_{max} 225 (ε 35000); 280 (ε 7000); 296 (ε 8400).

5-Deoxy-3,4-Dihydro-4-hydroxy-3-methoxy-4-(4-methoxyphenyl)-2(1H)-quinolinone [183854-01-5]

C₁₇H₁₇NO₄ 299.326

Prod. by *Penicillium* sp. NTC-47, *Penicillium* cf. *simplicissimum* and the marine-derived *Penicillium janczewskii*. Insecticidal agent. Needles (MeOH). Mp 76-79°. [α]_D²⁰ -62 (c, 0.3 in MeOH). λ_{max} 229 (ε 11000); 254 (ε 6500); 281 (ε 2800) (MeOH).

5-Deoxy, O³-de-Me: 3,4-Dihydro-3,4-dihydroxy-4-(4-methoxyphenyl)-2(1H)-quinolinone. Yaequinolone A₂ C₁₆H₁₅NO₄ 285.299

Prod. by *Penicillium* sp. FKI-2140 and the marine-derived *Penicillium janczewskii*. Insecticidal agent. Pale yellow powder. [α]_D¹⁵ -4.2 (c, 0.5 in MeOH). [α]_D²³ -50.9 (c, 0.1 in EtOH). λ_{max} 208 (ε 43000); 227 (ε 18900); 250 (ε 11300); 283 (ε 5100) (EtOH).

3-Epimer, 5-deoxy, O³-de-Me: Yaequinolone A₁

C₁₆H₁₅NO₄ 285.299

Prod. by *Penicillium* sp. FKI-2140 and the marine-derived *Penicillium janczewskii*. Insecticidal agent. Pale yellow powder. [α]_D¹⁵ -12.9 (c, 0.7 in MeOH). [α]_D²³ -32.2 (c, 0.1 in EtOH). λ_{max} 209 (ε 40800); 226 (ε 19200); 250 (ε 12700); 280 (ε 4600) (EtOH).


Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 914-916 (*isol, uv, ir, pmr, cmr*)

Kusano, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2559-2568 (*isol, uv, pmr, cmr, ms*)

He, J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1397-1399 (*isol, pmr, cmr*)

Uchida, R. *et al.*, *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*Yaequinolones*)

2,3-Dihydro-3,5-dihydroxy-2-oxo-1H-indole-3-acetic acid D-423
5-Hydroxydioxindole-3-acetic acid. 3,5-Dihydroxy-2-indolinone-3-acetic acid [61935-03-3]



(R)-form

C₁₀H₉NO₅ 223.185**(R)-form**

Constit. of rice bran. [α]_D²¹ +6 (c, 1 in MeOH).

5-O-β-D-Glucopyranoside: [218604-37-6] C₁₆H₁₉NO₁₀ 385.327

Constit. of rice bran. Powder. [α]_D²⁰ -25.8 (c, 0.2 in H₂O). λ_{max} 210 (ε 16400); 259 (ε 6600); 305 (ε 1200) (MeOH).

5-O-[β-D-Glucopyranosyl-(1→4)-β-D-glucopyranoside]: [218604-48-9]

C₂₂H₂₉NO₁₅ 547.469
Constit. of rice bran. Powder. [α]_D²⁰ -30.7 (c, 0.3 in H₂O). λ_{max} 210 (ε 19000); 259 (ε 7400); 305 (ε 1300) (MeOH).

(S)-form

5-O-β-D-Glucopyranoside: [218604-43-4]

Constit. of rice bran. Powder. [α]_D²⁰ -44.1 (c, 1.2 in H₂O). λ_{max} 210 (ε 16400); 259 (ε 6600); 305 (ε 1200) (MeOH).

5-O-[β-D-Glucopyranosyl-(1→4)-β-D-glucopyranoside]: [218604-54-7]

Constit. of rice bran. Powder. [α]_D²⁰ -46.5 (c, 1.3 in H₂O). λ_{max} 210 (ε 19000); 259 (ε 7400); 305 (ε 1300) (MeOH).

(±)-form [63389-29-7]

Pale yellow powder.

Me ester: [57061-18-4]

C₁₁H₁₁NO₅ 237.212

Constit. of rice bran. Needles. Mp 78-80°. Possibly an artifact.


Kinashi, H. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 2465 (*isol*)

Suzuki, Y. *et al.*, *Phytochemistry*, 1977, **16**, 635 (*isol*)

Tateishi, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 1870-1874 (*glycosides*)

2,3-Dihydro-3,7-dihydroxy-2-oxo-1H-indole-3-acetic acid D-424

3,7-Dihydroxy-2-indolinone-3-acetic acid



(R)-form

C₁₀H₉NO₅ 223.185**(R)-form**

7-O-β-D-Glucopyranoside: Zeanoside C [120293-55-2]

C₁₆H₁₉NO₁₀ 385.327

Isol. from immature sweet corn kernels *Zea mays* (Poaceae). Fine cryst. (EtOH). Mp 162-163°. [α]_D¹⁹ -162.1 (c, 0.1 in H₂O).

(S)-form

7-O-β-D-Glucopyranoside: Zeanoside A [113202-68-9]


C₁₆H₁₉NO₁₀ 385.327

Isol. from sweet corn kernels *Zea mays* (Poaceae). Fine needles (EtOH). Mp

197-198°. [α]_D -36.8 (c, 0.1 in H₂O).

Tateishi, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 3445-3447; 1988, **52**, 3231-3233; 1989, **53**, 2545-2551 (*isol, uv, ir, pmr, cmr, ms, struct*)

3,4-Dihydro-6,8-dimethoxy-1,3-dimethylisoquinoline, 9CI D-425



(S)-form

C₁₃H₁₇NO₂ 219.283**(S)-form** [104292-08-2]

Alkaloid from bark of *Ancistrocladus tectorius* (Ancistrocladaceae). Mp 202° (197°) (as hydrobromide). [α]_D²⁰ -141 (c, 1.0 in MeOH).

[104292-09-3]


Bringmann, G. *et al.*, *Angew. Chem., Int. Ed.*, 1986, **25**, 913 (*synth*)

Montagnac, A. *et al.*, *Phytochemistry*, 1995, **39**, 701 (*isol*)

6,7-Dihydro-2,3-dimethyl-5H-cyclopentapyrazine, 9CI D-426

FEMA 3917

[38917-63-4]

C₉H₁₂N₂ 148.207

Formed from cysteine and rhamnose under roasting conditions. Found in *Cyperus esculentus*. Mp 25-27°. Bp₁₀ 102-104°.

N-Oxide: [61928-81-2]

C₉H₁₂N₂O 164.207

Hygroscopic cryst. Mp 36-39°. Bp_{0.001} 85°.

Flament, I. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 610-619; 1976, **59**, 2308-2313; 2314-2326 (*synth, occur, pmr, ms, 1-oxide*)

Pittet, A.O. *et al.*, *J. Agric. Food Chem.*, 1974, **22**, 273-279 (*synth*)

Flament, I. *et al.*, *Bull. Soc. Chim. Belg.*, 1979, **88**, 941-950 (*synth, pmr, ms*)

Silwar, R. *et al.*, *Z. Lebensm.-Unters. -Forsch.*, 1992, **195**, 112-119 (*N-oxide*)


Cantelejo, M.J. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 1853-1860 (*occur, Cyperus esculentus*)

Fenaroli's Handbook of Flavor Ingredients, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 421 (*occur, use*)

Dihydro-4,6-dimethyl-4H-1,3,5-dithiazine D-427

[51647-36-0]

[82989-68-2]

C₅H₁₁NS₂ 149.281

(3R,5S)-form


cis-form

[82989-67-1]

Constit. of *Acacia latescens*.Kruse, H.P. *et al.*, *Nahrung*, 1982, **26**, 369-376 (occur)Peerzada, N. *et al.*, *Sulfur Lett.*, 2000, **23**, 185-192 (isol, synth)**4,5-Dihydro-2,4(5)-dimethyl-1H-imidazole, 9CI**

2,4-Dimethyl-2-imidazoline, 8CI

[930-61-0]

C₅H₁₀N₂ 98.147**(±)-form**

Component of crosslinking agent for epoxy resins. Rubber vulcanisation accelerator.

Liq. Bp₁₅ 107.5-108°. Irritant.


Picrate:

Solid. Mp 141-141.5°.

(ξ)-form

Volatile component of rainbow trout tissue.



[73006-98-1]

Ohashi, M. *et al.*, *Org. Mass Spectrom.*, 1968, **1**, 703 (ms)Hiatt, M.H. *et al.*, *Anal. Chem.*, 1983, **55**, 506 (isol, glc)Butler, R.N. *et al.*, *J.C.S. Perkin 1*, 1989, 155 (synth)**5,6-Dihydro-3,6-dioxo-1,2,4-triazine-4(3H)-carboxylic acid, 9CI**C₄H₃N₃O₄ 157.085

Me ester: [118040-43-0]

C₅H₅N₃O₄ 171.112Alkaloid from the seed coat of *Butea monosperma* (Fabaceae). Shows anti-fertility props.Porwal, M. *et al.*, *Natl. Acad. Sci. Lett. (India)*, 1988, **11**, 81; *C.A.*, **110**, 21066f (isol)**4,5-Dihydro-2,5-diphenyloxazole**2,5-Diphenyl-2-oxazoline. *Oxytriphine*

[22020-69-5]


C₁₅H₁₃NO 223.274**(S)-form** [87443-39-8]Alkaloid from aerial parts of *Oxytropis trichophysa*. Light yellow oil. Mp 38-40°.[α]_D +116 (c, 1.34 in CHCl₃). [α]_D +162.3 (c, 4.7 in CHCl₃) (synthetic). Confusion over abs. config. The isolation paper says the nat. prod. is *S*- but draws the *R*-form. CAS indexes it as *R*- but the opt. rotn. by comparison with synthetic material shows that it is *S*-.Meyers, A.I. *et al.*, *Tetrahedron*, 1983, **39**, 1991-1999 (*S*-form, synth)
Akhmedzhanova, V.I. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 873-876; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 778-780
Vorbruggen, H. *et al.*, *Tetrahedron*, 1993, **49**, 9353-9372 (synth, ir, pmr)
Badiang, J.G. *et al.*, *J.O.C.*, 1996, **61**, 2484-2487 (synth)
Minakata, S. *et al.*, *Chem. Comm.*, 2007, 3279-3281 (synth)**Dihydroerysodine**1,6-Didehydro-3,15-dimethoxyerythrinan-16-ol, 9CI
[7236-36-4]C₁₈H₂₃NO₃ 301.385Alkaloid from *Cocculus laurifolius* (Menispermaceae). Needles (Me₂CO). Mp 208-209°. [α]_D +224.39 (MeOH).Me ether: *Dihydroerysotrine*

[63694-44-0]


Synthetic, not known as a natural alkaloid. Pale-yellow viscous oil; yellow microscopic needles (as picrate). Mp 165-167° (picrate).

O¹⁵-De-Me, O¹⁶-Me: 1,6-Didehydro-3,16-dimethoxyerythrinan-15-ol, 9CI. *Dihydroerysovine*
[63694-43-9]C₁₈H₂₃NO₃ 301.385Alkaloid from the rhizomes of *Cocculus trilobus* (Menispermaceae). Oil. [α]_D +223 (CHCl₃).Prelog, V. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 453; 1956, **39**, 498 (synth, uv)Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1956, **4**, 225 (isol, uv, ir, struct)Millington, D.S. *et al.*, *J.A.C.S.*, 1974, **96**, 1909 (synth, pmr)Ju-ichi, M. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 533; 1978, **26**, 563 (*Dihydroerysodine*, *Dihydroerysovine*, isol, ir, pmr, ms, struct, synth)**2,3-Dihydro-1-formyl-2-hydroxy-5-methoxy-6-methyl-1H-indole**


2,3-Dihydro-2-hydroxy-5-methoxy-6-methyl-1H-indole-1-carboxaldehyde. 1-Formyl-2-hydroxy-5-methoxy-6-methylindoline

C₁₁H₁₃NO₃ 207.229Alkaloid from *Corydalis saxicola*. Yellow cryst. Mp 188-190°. λ_{max} 262 (log ε 3.17) (MeOH).

2-Deoxy: 2,3-Dihydro-1-formyl-5-methoxy-6-methyl-1H-indole. 2,3-Dihydro-5-methoxy-6-methyl-1H-indole-1-carboxaldehyde. 1-Formyl-5-methoxy-6-methylindoline

C₁₁H₁₃NO₂ 191.229Alkaloid from *Corydalis saxicola*. Yellow cryst. Mp 165-167°. λ_{max} 264 (log ε 3.29) (MeOH).Li, H.-L. *et al.*, *Chem. Biodiversity*, 2008, **5**, 777-783 (isol, pmr, cmr, ms)**6,7-Dihydro-4H-furo[2,3-c]azepine-4,8(5H)-dione***Fuscaïne*
[223678-23-7]C₈H₇NO₃ 165.148Isol. from the Chinese sponge *Phakellia fusca*. Cytotoxic.Zeng, L.-M. *et al.*, *CA*, 1999, **130**, 309248c**Dihydrogambirtannine**

Methyl 15,16,17,18,19,20-hexadecahydro-yohimban-16-carboxylate, 9CI

C₂₁H₂₀N₂O₂ 332.401**(R)-form** [2447-63-4]Synthetic. Cryst. + 1/2 EtOH (EtOH). Mp 174-176°. [α]_D²⁵ +288 (Py).**(S)-form** [25425-11-0]Alkaloid from *Uncaria gambier* (Gambir tannin) (Rubiaceae). Soft yellow cryst. (Et₂O/hexane). Mp 163°. [α]_D²⁰ -270 (c, 0.088 in CHCl₃).De(methoxycarbonyl): *Decarbomethoxy-dihydrogambirtannine*. 15,17,19-Yohimbatriene. Demethoxycarbonyldihydrogambirtannine

[15026-41-2]

C₁₉H₁₈N₂ 274.365Constit. of the leaves of *Ochrosia lifuana* and *Ochrosia miana*, and from fruits of *Strychnos usambarensis* (Apocynaceae, Loganiaceae). Mp 192-194° (racemate). [α]_D²⁰ -254 (c, 1 in EtOH).**(±)-form** [20361-83-5]Synthetic. Cryst. (C₆H₆/hexane). Mp 155-160° Mp 176-178° (175°) (dimorph.). *Hydrochloride*:

Cryst. (MeOH/Me₂CO). Mp 248-250° (244-245°) dec.

3,14-Didehydro: Gambirtannine

[24987-88-0]

C₂₁H₁₈N₂O₂ 330.385

Alkaloid from *Uncaria gambier* (*Ouroparia gambier*) (Gambir tannin). Orange needles. Mp 150-153° dec. Readily dec. Prob. artifact of autoxidation, not detected in the fresh plant.

3,14-Didehydro, 21-oxo: Oxogambirtannine

[18110-97-9]

C₂₁H₁₆N₂O₃ 344.369

Alkaloid from *Uncaria gambier* (Rubiaceae). Bright-yellow needles with yellow-green fluor. (CHCl₃/hexane). Cryst. (MeOH). Mp 208-209° Mp 228°.

De(methoxycarbonyl): [61825-78-3]

Pale yellow prisms or cream needles. Mp 191-194°.

(ξ)-form

De(methoxycarbonyl), 15,16,17,18-tetrahydro: **Demethoxycarbonyl-3,4,15,16,17,18-hexahydrogambirtannine**

[113122-69-3]

C₁₉H₂₂N₂ 278.396

Alkaloid from *Strychnos johnsonii*. [α]_D -138 (MeOH). λ_{max} 227 ; 274 ; 283 ; 290 (MeOH).

Smith, E. *et al.*, *J.A.C.S.*, 1964, **86**, 2083-2084 (*R-form*, *synth*)

Taylor, W.I. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1966, 1141-1143 (*Oxogambirtannine*, *isol*)

Merlini, L. *et al.*, *Tetrahedron*, 1967, **23**, 3129-3145 (*Gambirtannine*, *Dihydrogambirtannine*, *Oxogambirtannine*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *abs config*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1968, **90**, 5251-5256 (*synth*)

Beisler, J.A. *et al.*, *Tetrahedron*, 1970, **26**, 1961-1965 (*synth*, *Oxogambirtannine*)

Puebe-Locou, N. *et al.*, *Phytochemistry*, 1973, **12**, 199-200 (*Decarbomethoxydihydrogambirtannine*)

Irie, H. *et al.*, *Chem. Comm.*, 1975, 63 (*Oxogambirtannine*, *synth*)

Ninomiya, I. *et al.*, *J.C.S. Perkin 1*, 1976, 1865-1868 (*Decarbomethoxydihydrogambirtannine*)

Markaryan, E.A. *et al.*, *CA*, 1977, **86**, 140317 (*synth*)

Kan-Fan, C. *et al.*, *Tet. Lett.*, 1980, **21**, 1463-1466 (*S-form*, *synth*)


Chatterjee, A. *et al.*, *Synthesis*, 1981, 818-820 (*synth*, *Decarbomethoxydihydrogambirtannine*)

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*Demethoxycarbonylhexahydrogambirtannine*)

Cobas, A. *et al.*, *J.O.C.*, 1992, **57**, 6765-6769 (*synth*, *Decarbomethoxydihydrogambirtannine*)

3,4-Dihydro-5-(5-hexenyl)-2-nonyl-2H-pyrrole D-435

2-(5-Hexenyl)-5-nonyl-Δ¹-pyrroline [116902-76-2]




C₁₉H₃₅N 277.492

Ant venom alkaloid from european *Monomorium* species. Exhibits a v. broad insecticidal action. Bp_{0.1} 137°.

Bacos, D. *et al.*, *Tet. Lett.*, 1988, **29**, 3061 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *synth*, *struct*)

3,4-Dihydro-3-hydroxy-1,4-benzoxazin-2-one D-436

Acanthamine



C₈H₇NO₃ 165.148

(ξ)-form

O-β-D-Glucopyranoside: Acanthaminoside

C₁₄H₁₇NO₈ 327.29

Alkaloid from *Acanthus arboreus*. Needles (MeOH). Mp 213-214°. λ_{max} 250 ; 275 ; 283 (MeOH).

N-Hydroxy: 3,4-Dihydro-3,4-dihydroxy-1,4-benzoxazin-2-one. 4-Hydroxy-acanthamine


C₈H₇NO₄ 181.148

Alkaloid from *Acanthus arboreus*. Yellow needles (MeOH). Mp 135°. λ_{max} 253 ; 279 ; 290 (MeOH).

Amer, M.E. *et al.*, *J. Braz. Chem. Soc.*, 2004, **15**, 262-266 (*isol*, *pmr*, *cmr*, *ms*)

3,4-Dihydro-8-hydroxy-6,7-dimethoxyisoquinoline D-437

8-Hydroxy-6,7-dimethoxy-3,4-dihydroisoquinolinium inner salt



C₁₁H₁₃NO₃ 207.229

Alkaloid from *Lophophora williamsii* (Cactaceae). Mp 159-165°. Exists as inner salt.


N-Me, inner salt: 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-2-methylisoquinolinium inner salt

C₁₂H₁₅NO₃ 221.255

Alkaloid from *Lophophora williamsii* (Cactaceae). Mp 95-104°.

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 745 (*uv*, *ir*, *pmr*, *rev*)

1,3-Dihydro-3-hydroxy-4,9-dimethoxy-3-methyl-2H-benz[*f*]indol-2-one D-438



C₁₅H₁₅NO₄ 273.288

(S)-form [1040378-12-8]


Alkaloid from the roots of *Goniotalamus cheliensis*. Yellow powder. [α]_D²⁵ -37.9 (c, 0.1 in MeOH).

Jiang, M.M. *et al.*, *Chin. Chem. Lett.*, 2008, **19**, 302-304 (*isol*, *pmr*, *cmr*)

3,4-Dihydro-8-hydroxy-6,7-dimethoxy-1-methylisoquinoline D-439

3,4-Dihydro-6,7-dimethoxy-1-methyl-8-isoquinolinol, 9CI

[31241-40-4]



C₁₂H₁₅NO₃ 221.255

Alkaloid from *Lophophora williamsii* (Cactaceae). Cryst. (C₆H₆/CHCl₃). Mp 173-175°.

N-Me, inner salt: 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinolinium inner salt


C₁₃H₁₇NO₃ 235.282

Alkaloid from *Lophophora williamsii* (Cactaceae).

Kapadia, G.J. *et al.*, *J.A.C.S.*, 1970, **92**, 6943 (*synth*, *pmr*, *ms*)

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 745 (*uv*, *pmr*, *ms*, *rev*)

2,3-Dihydro-2-(1-hydroxyethyl)-2-methylquinazolin-4(1H)-one D-440



(1*R**, 2*R**)-form

C₁₁H₁₄N₂O₂ 206.244

(1*R*, 2*R*)-form

Prod. by *Streptomyces* sp. GW23/1540. Amorph. solid. [α]_D²⁰ -17.5 (c, 0.2 in MeOH). λ_{max} 253 (log ε 3.51); 350 (log ε 3.25) (MeOH).


(1*R*, 2*S*)-form

Prod. by *Streptomyces* sp. GW23/1540. Amorph. solid. [α]_D²⁰ +47 (c, 0.2 in MeOH). λ_{max} 252 (log ε 3.61); 348 (log ε 3.33) (MeOH).

Maskey, R.P. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1131-1134 (*isol*, *pmr*, *cmr*, *ms*)

3-(2,3-Dihydro-3-hydroxy-1*H*-indol-3-yl)-2-oxopropanoic acid D-441

2,3-Dihydro-3-hydroxy-α-oxo-1*H*-indole-3-propanoic acid



C₁₁H₁₁NO₄ 221.212


(ξ)-form

Prod. by the marine-derived *Halomonas* sp. RK377. Amorph. solid.

Liang, L. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*isol*, *pmr*, *cmr*, *ms*)

2,3-Dihydro-4-hydroxy-1H-isoindol-1-one, 9CI **D-442**

4-Hydroxyphthalimidine. 4-Hydroxy-1-isoindolinone
[366453-21-6]



C₈H₇NO₂ 149.149

Alkaloid from the antlion (larva of *Myrmeleontidae* sp.). Pale yellow needles (MeOH). Mp 271° dec.

N-(2-Hydroxyethyl): *2,3-Dihydro-4-hydroxy-2-(2-hydroxyethyl)-1H-isoindol-1-one. 4-Hydroxy-2-(2-hydroxyethyl)-1-isoindolinone*
[919079-22-4]


C₁₀H₁₁NO₃ 193.202

Alkaloid from the antlion. Needles (MeOH).

Nakatani, T. *et al.*, *J. Nat. Med. (Tokyo)*, 2006, **60**, 261-263 (*isol*, *pmr*, *cmr*, *ms*)

1,3-Dihydro-3-hydroxy-4-methoxy-3-methyl-2H-indol-2-one, 9CI **D-443**

3-Hydroxy-4-methoxy-3-methylindole
[110011-50-2]




C₁₀H₁₁NO₃ 193.202

Alkaloid from the roots of *Capparis tomentosa* (Capparidaceae). Cryst. (EtOAc). Mp 230°.

Dekker, T.G. *et al.*, *Phytochemistry*, 1987, **26**, 1845 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-8-hydroxy-6-methoxy-1,3-dimethylisoquinoline **D-444**

4-(3,4-Dihydro-8-hydroxy-6-methoxy-1,3-dimethylisoquinolin-5-yl)-8-methoxy-3-methyl-1-naphthalenol. 3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-6-methoxy-1,3-dimethyl-8(2H)-isoquinolinone
[180342-44-3]



C₂₄H₂₅NO₄ 391.466

Inseparable mixt. of (3*S*,5*R*) and (3*S*,5*S*) atropisomers. Alkaloid from leaves and twigs of *Ancistrocladus tectorius*. [α]_D -90.3 (c, 0.39 in MeOH). Exists exclusively in the 8-oxo-form. λ_{max} 233 (log ε 4.6); 265 (log ε 3.7); 309 (log ε 3.9); 322 (log ε 3.9); 337 (log ε 3.9); 383 (log ε 4.3) (EtOH).

4'-Me ether: 3,4-Dihydro-5-(4,5-dimethoxy-2-methyl-1-naphthalenyl)-8-hydroxy-6-methoxy-1,3-dimethylisoquinoline. 3,4-Dihydro-5-(4,5-dimethoxy-2-methyl-1-naphthalenyl)-6-methoxy-1,3-dimethyl-8(2H)-isoquinolinone
[180342-43-2]

C₂₅H₂₇NO₄ 405.493


From *Ancistrocladus tectorius*. [α]_D -5.8 (c, 0.24 in MeOH). Exists exclusively in the 8-oxo-form. λ_{max} 232 (log ε 4.4); 270 (log ε 3.6); 306 (log ε 3.8); 322 (log ε 3.8); 335 (log ε 3.8); 384 (log ε 4.2) (EtOH).

8-Me ether: 3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-6,8-dimethoxy-1,3-dimethylisoquinoline
[180342-42-1]

C₂₅H₂₇NO₄ 405.493

From *Ancistrocladus tectorius*. [α]_D +41.7 (c, 0.29 in MeOH). λ_{max} 227 (log ε 4.5); 280 (log ε 3.9); 309 (log ε 3.9); 322 (log ε 3.9); 337 (log ε 3.8) (EtOH).

Manfredi, K.P. *et al.*, *J. Nat. Prod.*, 1996, **59**, 854-859 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cd*, *struct*)

3,4-Dihydro-3-hydroxy-1-methyl-4,5-diphenyl-2(1H)-pyridinone, 9CI **D-445**

C₁₈H₁₇NO₂ 279.338

(3*R*,4*S*)-form Homoclausenamidine

[136112-76-0]


Alkaloid from *Clausena lansium* (wampee) (Rutaceae).

Yang, M. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 291; *CA*, **115**, 155059t (*isol*, *struct*)

Yang, L. *et al.*, *Org. Lett.*, 2008, **10**, 2461-2464 (*synth*)

6,7-Dihydro-4-(hydroxy-methyl)-2-(*p*-hydroxyphenethyl)-7-methyl-5*H*-2-pyridinium(1+) **D-446**

[15794-93-1]




C₁₈H₂₂NO₃[⊕] 284.377

Minor alkaloid from the roots of *Valeriana officinalis* (valerian) (Valerianaceae). Mp 220-223° dec. (as chloride). [α]_D²² +19.3 (MeOH) (as trifluoroacetate).

Torssell, K. *et al.*, *Acta Chem. Scand.*, 1967, **21**, 53 (*isol*, *ir*, *pmr*, *ms*, *struct*)

6,7-Dihydro-3-hydroxy-methyl-8(5*H*)-indolizinsonone Polygonatine A **D-447**

[400715-63-1]



C₉H₁₁NO₂ 165.191

Alkaloid from the rhizomes of *Polygonatum sibiricum*. Needles (Me₂CO). Mp 101-103°. λ_{max} 206 (log ε 2.8); 298 (log ε 3.3) (no solvent reported).

Et ether: 3-(Ethoxymethyl)-6,7-dihydro-8(5H)-indolizinsonone, 9CI. Polygonatine B
[215662-61-6]

C₁₁H₁₅NO₂ 193.245

Alkaloid from *Polygonatum kingianum* and *Polygonatum sibiricum*. Pale yellow syrup. λ_{max} 214 (log ε 3); 323 (log ε 3.2) (no solvent reported).

Butyl ether: 3-(Butoxymethyl)-6,7-dihydro-8(5H)-indolizinsonone. Kinganone
[744198-33-2]

C₁₃H₁₉NO₂ 221.299

Alkaloid from the rhizomes of *Polygonatum kingianum*. Yellow oil.

Sun, L. *et al.*, *CA*, 1999, **130**, 2168n (*Polygonatine B*)


Wang, Y.-F. *et al.*, *Planta Med.*, 2003, **69**, 1066-1068 (*Kinganone*)

Sun, L.R. *et al.*, *J. Asian Nat. Prod. Res.*, 2005, **7**, 127-130 (*Polygonatine A*)

Dinsmore, A. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1032-1037 (*synth*)

1,5-Dihydro-5-hydroxy-3-methyl-1-(5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, 9CI **D-448**


5-Hydroxy-3-methyl-1-(2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one
[142750-27-4]



C₉H₁₂N₂O₃ 196.205

Alkaloid from bulbs of *Lilium candidum* (Liliaceae). Mp 176-178°.

Deoxy: 1,5-Dihydro-3-methyl-1-(5-oxo-2-pyrrolidinyl)-2H-pyrrol-2-one, 9CI. 3-Methyl-1-(2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one
[114020-02-9]

C₉H₁₂N₂O₂ 180.206Isol. from the petals of *Lilium candidum* (Liliaceae). Mp 167-169°.Haladová, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1988, **53**, 157 (deoxy)Eisenreichová, E. *et al.*, *Phytochemistry*, 1992, **31**, 1084-1085 (isol, pmr, ms, struct)**1,5-Dihydro-5-hydroxy-3-methyl-2H-pyrrol-2-one, 9CI** D-449*5-Hydroxy-3-methyl-3-pyrrolin-2-one. Jatropham. Jatrophine†. Lilidine* [37772-60-4]C₅H₇NO₂ 113.116

Log P -1.28 (uncertain value) (calc).

(R)-form [50656-76-3]

Alkaloid from epigeal parts of *Jatropha macrorrhiza*, from bulbs of *Lilium hansonii* (Euphorbiaceae, Liliaceae). Shows anti-neoplastic props. Marginally active against P-388 lymphocytic leukaemia. Cryst. (MeCN), needles (CHCl₃/MeOH). Sol. H₂O, butanol, Me₂CO, MeOH; fairly sol. EtOAc, CHCl₃; poorly sol. Et₂O, C₆H₆, hexane. Mp 131-132° (119-123°). [α]_D²⁵₄₉ -62 (H₂O). [α]_D²⁵ -76.2 (MeOH) (-26.3). λ_{max} 230 (ε 1150) (MeOH).

O-β-D-Glucopyranoside: [109028-41-3]

[109028-42-4 (tetra-Ac)]

C₁₁H₁₇NO₇ 275.258

Alkaloid from the bulbs of *Lilium hansonii* and *Lilium candidum* (Liliaceae). Needles (CHCl₃/MeOH). Sol. MeOH. Mp 183-185°. [α]_D -22 (c, 0.2 in MeOH).

O-(6-O-p-Hydroxycinnamoyl-β-D-glucopyranoside): *Jatropham 5-(6-p-coumaroylglucoside)*

[174205-03-9]

C₂₀H₂₃NO₉ 421.403

Alkaloid from fresh bulbs of *Lilium martagon*. Pale yellow amorph. solid. [α]_D²⁹ -26.3 (c, 0.19 in MeOH).

O-[β-D-Glucopyranosyl-(1→3)-β-D-glucopyranoside]: [145042-03-1]

C₁₇H₂₇NO₁₂ 437.4

Alkaloid from bulbs of *Lilium hansonii* (Liliaceae). Amorph. powder. [α]_D²⁹ -70 (c, 0.10 in MeOH).

(±)-form [79703-33-6]

Cryst. (EtOAc/petrol). Mp 120-121° (115-118°) (108°).

Me ether. (±)-5-O-Methyljatropham

[109050-99-9]

C₆H₉NO₂ 127.143

Alkaloid from the bulbs of *Lilium hansonii* (Liliaceae). Syrup. Possible artifact.

Wiedhopf, J.R.M. *et al.*, *J. Pharm. Sci.*, 1973, **62**, 1206 (isol, uv, ir, pmr)

Farina, F. *et al.*, *Synthesis*, 1973, 167 (synth)

Yakushijin, K. *et al.*, *Heterocycles*, 1980, **14**, 1073; 1981, **16**, 1157 (struct, synth, ir, uv, ms, pmr, cmr)

Nagasaka, T. *et al.*, *Heterocycles*, 1981, **16**, 1987 (synth, ir, uv, pmr, cmr)

Pinder, A.R. *et al.*, *Alkaloids (London)*, 1982, **12**, 36 (nomencl)

Farina, F. *et al.*, *Heterocycles*, 1984, **22**, 1733

Abdullaev, N.D. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 692; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 576 (isol, pmr, cmr, struct)

Shimomura, H. *et al.*, *Phytochemistry*, 1987, **26**, 582 (isol, uv, ir, pmr, cmr, cd, abs config, derivs)

Ori, K. *et al.*, *Phytochemistry*, 1992, **31**, 2769 (glucosylglucoside)


Dittami, J.P. *et al.*, *Tet. Lett.*, 1995, **36**, 4201 (synth)

Mucaji, P. *et al.*, *Coll. Czech. Chem. Comm.*, 1996, **61**, 1662-1664 (5-glucoside)

Satou, T. *et al.*, *Phytochemistry*, 1996, **41**, 1225 (coumaroylglucoside)

N-[5-[(1,6-Dihydro-2-hydroxy-5-octanamido-6-oxo-3-pyridyl)imino]-1,2,5,6-tetrahydro-2,6-dioxo-3-pyridyl]octanamide, 8CI D-450


[16236-90-1]

H₃C(CH₂)₆CONHC₂₆H₃₇N₅O₆ 515.608

Pigment from *Pseudomonas lemmonieri*. Glistening bronze cryst. Fairly sol.

AcOH, DMF; poorly sol. MeOH, acids, H₂O, hexane. Mp 288°. λ_{max} 627 nm (log ε 4.96) (DMSO). λ_{max} 595; 627 (DMSO) (Berdy). λ_{max} 635 (Py) (Berdy). λ_{max} 625 (DMF) (Berdy).

Knackmuss, H.J. *et al.*, *Chem. Ber.*, 1967, **100**, 2537 (synth, uv, ir, pmr)

2,3-Dihydro-3-hydroxy-2-oxo-1H-indole-3-acetic acid, 9CI D-451*Dioxindole-3-acetic acid*C₁₀H₉NO₄ 207.185**(ξ)-form***Me ester: Coixlactam B*


[205656-36-6]

C₁₁H₁₁NO₄ 221.212

Constit. of *Coix lachryma-jobi* var. *mayuen* (adlay bran), *Hibiscus moscheutos* and rice bran. Prob. intermed. in biosynth. of 2,6-Dihydroxy-4-quinolinecarboxylic acid, D-672 from 1H-Indole-3-acetic acid, I-81. Amorph. powder. λ_{max} 209 (ε 22500); 254 (ε 5080); 293 (ε 1380) (MeOH). λ_{max} 211 (ε 74700); 253 (ε 5110) (MeOH/NaOH aq.).

Kinashi, H. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 2465-2470 (isol, pmr, ms)

Lee, M.-Y. *et al.*, *Food Chem. Toxicol.*, 2008, **46**, 1933-1939 (isol)


2,3-Dihydro-4-hydroxy-2-oxo-1H-indole-3-acetic acid D-452*4-Hydroxyoxindole-3-acetic acid*C₁₀H₉NO₄ 207.185

Nitrile: 2,3-Dihydro-4-hydroxy-2-oxo-1H-indole-3-acetonitrile. 4-Hydroxyoxindole-3-acetonitrile


C₁₀H₈N₂O₂ 188.185

Alkaloid from *Isatis indigotica*. Cryst. (EtOAc). Mp 150-152°.

Li, B. *et al.*, *Yaoxue Xuebao*, 2000, **35**, 508-510

2,3-Dihydro-5-hydroxy-2-oxo-1H-indole-3-acetic acid, 9CI D-453*5-Hydroxyoxindole-3-acetic acid*C₁₀H₉NO₄ 207.185*Me ester:*C₁₁H₁₁NO₄ 221.212

Isol. from rice bran. Amorph. powder. Kinashi, H. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 2465

4,5-Dihydro-2-(2-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic acid, 9CI D-454*4-Methylaeruginic acid*C₁₁H₁₁NO₃S 237.279

Trivial name is misleading.

(S)-form [189820-27-7]

Prod. by *Streptomyces* sp. KCTC 9303. Cytotoxic agent. Yellow powder. Sol. MeOH, EtOAc, H₂O, Me₂CO; poorly sol. hexane, CHCl₃. λ_{max} 215 (ε 27900); 250 (ε 14250); 312 (ε 5378) (MeOH).

Ryoo, I.-J. *et al.*, *J. Antibiot.*, 1997, **50**, 256-258 (isol, uv, ir, cd, pmr, cmr)


Bergeron, R.J. *et al.*, *J. Med. Chem.*, 1999, **42**, 2432-2440 (synth, abs config, cd)

Martin, G.E. *et al.*, *J. Nat. Prod.*, 2000, **63**, 543-585 (N-15 nmr)

4,5-Dihydro-2-(2-hydroxyphenyl)-4-oxazolecarboxylic acid, 9CI

D-455

2-(2-Hydroxyphenyl)-2-oxazoline-4-carboxylic acid
[37592-62-4]



$C_{10}H_9NO_4$ 207.185

(-)-form

Me ester: [199173-66-5]
[37592-61-3, 199540-89-1]
 $C_{11}H_{11}NO_4$ 221.212
Prod. by *Actinomadura* sp. MJ502-77F8.
[α]_D²⁰ -50.1 (c, 1.5 in $CHCl_3/MeOH$). λ_{max}
238 ; 247 ; 253 (sh) ; 306 (MeOH).


(±)-form

Cryst. ($CHCl_3$). Mp 157-158°.
Me ester:
Needles (petrol). Mp 71-72°.
Black, D.St.C. et al., *Aust. J. Chem.*, 1972, **25**,
1797-1810 (*synth*)
Sasaki, T. et al., *J. Antibiot.*, 1997, **50**, 881-883
(*isol, uv, ir, pmr, cmr, activity*)

5,6-Dihydro-3-[2-(4-hydroxyphenyl)-2-oxoethyl]-2(1H)-pyridinone, 9CI

D-456

[71886-38-9]




$C_{13}H_{13}NO_3$ 231.251

Isol. from an algae-infested Caribbean sponge *Halichondria melanodocia*. Cryst. ($CHCl_3/MeOH$). Mp 235-235.5°.

Gopichand, Y. et al., *J.O.C.*, 1979, **44**, 4995
(*isol, uv, ir, pmr, ms, struct*)

3,4-Dihydro-6-(4-hydroxyphenyl)-2H-1,4-thiazine D-457

4-(3,4-Dihydro-2H-1,4-thiazin-6-yl)phenol, 9CI



$C_{10}H_{11}NOS$ 193.269

S,S-Dioxide: 4-(3,4-Dihydro-1,1-dioxido-2H-1,4-thiazin-6-yl)phenol, 9CI. 3,4-Dihydro-6-(4-hydroxyphenyl)-1,1-dioxo-2H-1,4-thiazine


[154887-08-8]

$C_{10}H_{11}NO_3S$ 225.268
Alkaloid from the sponge *Anchinoe tenacior*. λ_{max} 274 (ϵ 8100) (MeOH).

Casapullo, A. et al., *Tet. Lett.*, 1994, **35**, 2421-2422 (*isol, uv, pmr*)

5,6-Dihydro-5-hydroxy-2(1H)-pyridinone D-458

3,4-Dehydro-5-hydroxy-2-piperidone



$C_5H_7NO_2$ 113.116


(+)-form

Alkaloid from *Piper sintonense*. Brownish oil. [α]_D +55.7 (c, 0.2 in MeOH). λ_{max} 209 (log ϵ 4.24); 247 (sh) (log ϵ 3.52) ($CHCl_3$).

Chen, J.-J. et al., *Helv. Chim. Acta*, 2003, **86**, 2058-2064 (*isol, pmr, cmr*)

1,3-Dihydro-3-hydroxy-2H-pyrrol-2-one, 9CI D-459

[79463-75-5]



$C_4H_5NO_2$ 99.089

Mp 98-100°.

O- β -D-Glucopyranoside: *Pisatocide*

[18814-39-6]
 $C_{10}H_{15}NO_7$ 261.231
Isol. from seedlings of *Pisum sativum* (peas). Mp 104-107°. [α]_D²⁰ -49 (H_2O).


Me ether: 1,3-Dihydro-3-methoxy-2H-pyrrol-2-one
[40090-78-6]
 $C_5H_7NO_2$ 113.116
Solid by subl. Mp 50-52°.

[18814-40-9]

Kocourek, J. et al., *Arch. Biochem. Biophys.*, 1967, **121**, 531-532 (*isol, struct*)
Abramovitch, R.A. et al., *J.A.C.S.*, 1976, **98**, 1478-1486 (*Me ether*)

1,5-Dihydro-5-hydroxy-2H-pyrrol-2-one, 9CI D-460

Dihydromaleimide. *Isosuccinimide*
[34085-09-1]



$C_4H_5NO_2$ 99.089

No evidence for tautomerism from form shown.

(R)-form [87710-47-2]

Alkaloid from the shoots of dwarf pea (*Pisum sativum*) (Fabaceae). Prisms. Mp 103-105°. [α]_D²³ -12.6.

O- β -D-Glucopyranoside: *Dihydromaleimide* β -D-glucoside. *Isosuccinimide* β -D-glucoside
[26696-59-3]

$C_{10}H_{15}NO_7$ 261.231

Alkaloid from *Pisum sativum* (peas) (Fabaceae). Prisms. Mp 185-188°. [α]_D^{22.5} -73.7. The identity (structural and stereochemical) of the various isolates of this glucoside is not certain. May be identical with Pisatocide, in D-459. No further information to 2007.

(±)-form

Cryst. (Me_2CO). Mp 102°.

Me ether: 1,5-Dihydro-5-methoxy-2H-pyrrol-2-one, 9CI
[34128-18-2]
 $C_5H_7NO_2$ 113.116
Mp 38°. Bp₉ 95°.

de Mayo, P. et al., *Chem. Ind. (London)*, 1962, 1576-1577 (*synth*)


Liu, T.-Y. et al., *Plant Physiol.*, 1970, **45**, 424-428; *CA*, **73**, 41981w (*isol, glucoside*)

Farina, F. et al., *Synthesis*, 1973, 167-168 (*synth*)

Masuko, M. et al., *Phytochemistry*, 1983, **22**, 1278-1280 (*isol, uv, ir, pmr, cmr, cd, struct, glucoside, bibl*)

3,4-Dihydro-6-hydroxy-2(1H)-quinolinone, 9CI D-461

[54197-66-9]



$C_9H_9NO_2$ 163.176

Isol. from Ayurvedic processed commercial root tubers of *Aconitum ferox*. Solid.

Me ether: [54197-64-7]
 $C_{10}H_{11}NO_2$ 177.202
Needles (hexane/EtOAc). Mp 144-146°.

Jones, G.H. et al., *J. Med. Chem.*, 1987, **30**, 295-303 (*Me ether, synth*)

Sakamoto, T. et al., *J. Het. Chem.*, 1988, **25**, 1279-1281 (*Me ether, synth, ir, pmr*)


Hanuman, J.B. et al., *Nat. Prod. Lett.*, 1993, **3**, 227-231 (*isol*)

Iyobe, A. et al., *Chem. Pharm. Bull.*, 2001, **49**, 822-829 (*Me ether, synth, ir, pmr*)

Occhiato, E.G. et al., *J. Med. Chem.*, 2004, **47**, 3546-3560 (*synth, pmr*)

7,8-Dihydroimidazo[1,5-c]pyrimidin-5(6H)-one, 9CI D-462

5,6,7,8-Tetrahydro-5-oxoimidazo[1,5-c]pyrimidine. 5,6,7,8-Tetrahydroimidazo[1,5-c]pyrimidin-5-one
[14509-66-1]




C₆H₇N₃O 137.141Metab. from the Caribbean sponge *Aplysina fistularis* f. *fulva*. Cryst. (EtOH). Mp 220-222°.

Methiodide: [51720-37-7]

Cryst. Mp 228-230° dec.


Mechoulam, R. *et al.*, *Tetrahedron*, 1967, **23**, 239 (*synth*, *pmr*, *ir*, *use*)Durant, C.G. *et al.*, *J. Med. Chem.*, 1976, **19**, 923 (*synth*)Jairam, R. *et al.*, *J.O.C.*, 1992, **57**, 4136 (*synth*)Ciminiello, P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 705 (*isol*, *pmr*, *cmr*)**6,7-Dihydro-8(5H)-indolizone, 9CI** D-463

[54906-44-4]

C₈H₉NO 135.165Constit. of Virginia tobacco (*Nicotiana tabacum*). Mp 34°.Wahlberg, I. *et al.*, *Phytochemistry*, 1977, **16**, 1233-1235 (*isol*)Bond, T.J. *et al.*, *J.C.S. Perkin 1*, 1993, 2241-2242 (*synth*)Barton, D.H.R. *et al.*, *Tet. Lett.*, 1994, **35**, 9157-9158 (*synth*)Dinsmore, A. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1032-1037 (*synth*)**11,12-Dihydroindolo[2,3-a]carbazole, 9CI** D-464

Indolo[2,3-a]carbazole

[60511-85-5]

C₁₈H₁₂N₂ 256.306

Residue present in a number of microbial and marine alkaloids. Cryst. (EtOH). Mp 370°.

N-Me:

C₁₉H₁₄N₂ 270.333


Prisms (EtOH). Mp 244-245°.

N,N'-Di-Me:

C₂₀H₁₆N₂ 284.36Needles (Me₂CO aq.). Mp 209°.N-(6-Deoxy-β-D-gulopyranosyl): **Tjipanazole G1**C₂₄H₂₂N₂O₄ 402.449Minor alkaloid from the blue-green alga *Tolypothrix tjipanasensis*. Protein kinase C activator. Exhibits antifungal activity. [α]_D +9.1 (c, 1.0 in CHCl₃).λ_{max} 256 (ε 35400); 270 (ε 32800); 287 (ε 18800); 324 (ε 20000); 343 (ε 5800); 359 (ε 4080) (MeOH) (Derep).N-α-L-Rhamnopyranosyl: **Tjipanazole G2**C₂₄H₂₂N₂O₄ 402.449Minor constit. of *Tolypothrix tjipanasensis*. λ_{max} 256 (ε 35400); 270 (ε 32800); 287 (ε 18800); 324 (ε 20000); 343 (ε 5800); 359 (ε 4080) (MeOH) (Derep).Mann, F.G. *et al.*, *J.C.S.*, 1958, 1525 (*synth*, *derivs*)Hünig, S. *et al.*, *Annalen*, 1976, 1090 (*uv*)Bonjouklian, R. *et al.*, *Tetrahedron*, 1991, **47**, 7739 (*Tjipanazoles*)**2,3-Dihydro-1H-indol-5-ol, 9CI** D-465

2,3-Dihydro-5-hydroxy-1H-indole. 5-Hydroxyindoline

[172078-33-0]

C₈H₉NO 135.165Alkaloid from *Phoebe chekiangensis* (Lauraceae).

Hydrochloride: [92818-38-7]

Off-white solid.

N-Ac: [4770-32-5]

C₁₀H₁₁NO₂ 177.202

Needles (EtOAc). Mp 249-250°.

Me ether: 2,3-Dihydro-5-methoxy-1H-indole, 9CI. 5-Methoxyindoline

[21857-45-4]

C₉H₁₁NO 149.192

Oil.

Benzyl ether: [92818-36-5]


C₁₅H₁₅NO 225.29

Oil. Waxy solid at 0°.


Benzyl ether; hydrochloride: [92818-37-6] Mp 188-190°.

Hunt, R.R. *et al.*, *J.C.S. (C)*, 1966, 344-345 (*N-Ac*)Young, T.E. *et al.*, *J.O.C.*, 1984, **49**, 4833-4838 (*synth*, *pmr*, *cmr*, *ir*)Hegde, V.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 1207-1212 (*isol*, *pmr*, *cmr*)Gangjee, A. *et al.*, *J. Med. Chem.*, 1997, **40**, 479-485 (*Me ether*)**5a,6-Dihydro-3-(1H-indol-3-yl)-1,4-dimethoxy-10bH-benzofuro[2,3-b]indole-2,10b-diol, 9CI** D-466

[78279-83-1]

C₂₄H₂₀N₂O₅ 416.432Metab. of *Aspergillus terreus* var. *africanus*. Brownish powder + 1/2 C₆H₆ (C₆H₆). Mp 131-140°. [α]_D²⁵ +51.2 (c, 0.13 in dioxan).**O²-Me: 5a,6-Dihydro-3-(1H-indol-3-yl)-1,2,4-trimethoxy-10bH-benzofuro[2,3-b]indol-10b-ol, 9CI**


[78279-82-0]

C₂₅H₂₂N₂O₅ 430.459Metab. of *Aspergillus terreus* var. *africanus*. Sl. brownish powder (CCl₄). Mp 134-138°. [α]_D¹⁸ +34.3 (c, 0.11 in EtOH).Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1005 (*pmr*, *struct*)**3,4-Dihydro-3-(1H-indol-3-yl)methyl-1H-1,4-benzodiazepine-2,5-dione, 9CI** D-467C₁₈H₁₅N₃O₂ 305.335

(S)-form [101409-14-7]


Isol. from *Aspergillus flavipes*. Solid. Mp 236-240°. [α]_D +138.8 (c, 0.5 in MeOH).Barrow, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 471 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *cd*, *struct*)**5,6-Dihydro-3-[2-(1H-indol-3-yl)-2-oxoethyl]-2(1H)-pyridinone, 9CI** D-468

[71886-40-3]

C₁₅H₁₄N₂O₂ 254.288Isol. from an algae-infested Caribbean sponge *Halichondria melanodocia*. Cryst. (MeOH/CHCl₃). Mp 194-194.5°.Gopichand, Y. *et al.*, *J.O.C.*, 1979, **44**, 4995-4997 (*isol*, *uv*, *ir*, *pmr*, *ms*)Johnson, A.-L. *et al.*, *Tetrahedron*, 2006, **62**, 10815-10820 (*synth*)**3,4-Dihydro-1(2H)-isoquinolinone, 9CI** D-469

3,4-Dihydroisocarbostyryl. 1-Oxo-1,2,3,4-tetrahydroisoquinoline. 3,4-Dihydro-1-isoquinolinol. 3,4-Dihydro-1-hydroxyisoquinoline

[1196-38-9]



C₉H₉NO 147.176*NH*-form predominates. Cryst. (MeOH aq.). Mp 58°. Bp_{0.1} 125-127°.**NH-form**

N-Ac: [74315-06-3]

C₁₁H₁₁NO₂ 189.213

Cryst. (EtOAc/hexane). Mp 97-98°.

N-Benzoyl: [75435-41-5]


C₁₆H₁₃NO₂ 251.284Bp₂ 210°.

N-Me: [6772-65-2]

C₁₀H₁₁NO 161.203Liq. Bp₆ 157° Bp_{0.4} 118°.

N-Hydroxy: 3,4-Dihydro-2-hydroxy-1(2H)-isoquinolinone, 9CI


[116526-30-8]

C₉H₉NO₂ 163.176Prod. by *Streptomyces griseus* 2002-104. Forms Fe chelates. Siderophore. Allergy inhibitor. Eubacterial growth promoter. λ_{max} 480 (H₂O+FeCl₃) (Berdy).Schneider, W. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1958, **291**, 560-566 (*synth*, N-Me)Davies, R.V. et al., *J.C.S. Perkin I*, 1978, 180-184 (*synth*, *ir*)Misra, R.N. et al., *Bioorg. Med. Chem. Lett.*, 1991, **1**, 295-298 (*synth*, N-hydroxy)Graefe, U. et al., *J. Basic Microbiol.*, 1994, **35**, 351-355 (*isol*, N-hydroxy)Norman, M.H. et al., *J. Med. Chem.*, 1994, **37**, 2552-2563 (*synth*, *pmr*)Kawase, M. et al., *Chem. Pharm. Bull.*, 1997, **45**, 1248-1253 (N-acyl derivs, *ir*, *pmr*, *cmr*)Koltunov, K. et al., *J.O.C.*, 2002, **67**, 8943-8951 (*synth*, *pmr*, *cmr*)Cai, C. et al., *Synth. Commun.*, 2005, **35**, 349-356 (*synth*, *pmr*)**5,6-Dihydro-N-[3-(7-methoxy-1,3-benzodioxol-5-yl)-2-propenoyl]-2(1H)-pyridinone** D-470N-(3-Methoxy-4,5-methylenedioxy-cinnamoyl)-Δ³-pyridin-2-one. 5,6-Dihydro-1-(3-methoxy-4,5-methylenedioxy-cinnamoyl)-2(1H)-pyridinone [130263-11-5]C₁₆H₁₅NO₅ 301.298Alkaloid from the stems of *Piper arborescens* (Piperaceae). Pale yellow needles (MeOH). Mp 157-158°. λ_{max} 225 (ε 57543); 341 (ε 52480) (MeOH) (Berdy).

2',3'-Dihydro: 5,6-Dihydro-1-[3-(7-methoxy-1,3-benzodioxol-5-yl)-1-oxopropyl]-2(1H)-pyridinone, 9CI [133034-10-3]

C₁₆H₁₇NO₅ 303.314Alkaloid from the leaves of *Piper arborescens*. Cytotoxic. Needles. Mp 80-81°. λ_{max} 220 (ε 3240); 242 (ε 2100); 285 (ε 350) (MeOH) (Berdy).Duh, C.-Y. et al., *J. Nat. Prod.*, 1990, **53**, 1575 (*isol*, *pmr*, *cmr*, *struct*, *deriv*)Duh, C.-Y. et al., *Phytochemistry*, 1990, **29**, 2689 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)**1,3-Dihydro-3-(3-methyl-2-butenylidene)-2H-indol-2-one, 9CI** D-471

3-(3-Methyl-2-butenylidene)-2-indolinone



(E)-form

C₁₃H₁₃NO 199.252


(E)-form [67987-50-2]

Pigment from the rhizomes of *Cimicifuga dahurica* (Ranunculaceae). Yellow needles (hexane/EtOAc). Mp 200-203°.

(Z)-form [67987-49-9]

Pigment from the rhizomes of *Cimicifuga dahurica* (Ranunculaceae). Yellow plates (hexane/EtOAc). Mp 213-214°.6-Methoxy: 6-Methoxy-3-(3-methyl-2-butenylidene)-2-indolinone. **Soulieotine** C₁₄H₁₅NO₂ 229.278Alkaloid from the rhizomes of *Souliea vaginata*. Red needles (MeOH). Mp 201-203°. [α]_D²⁰ +1.4 (c, 0.07 in CHCl₃/MeOH). Text. in ref. incorrectly states E-config., but the struct. was drawn correctly. λ_{max} 210 (log ε 4.4); 262 (log ε 3.94); 282 (log ε 3.09) (MeOH).Baba, K. et al., *Chem. Pharm. Bull.*, 1981, **29**, 2182 (*isol*, *uv*, *ir*, *pmr*, *cryst struct*, *synth*)Zhou, L. et al., *Heterocycles*, 2005, **65**, 1409-1414 (*Soulieotine*)**2,3-Dihydro-2-methylindole, 9CI** D-472

2-Methylindoline [6872-06-6]



(R)-form

C₉H₁₁N 133.193

▶ NM1926350

(R)-form [22160-13-0]

Oil. [α]_D²⁰ +11 (c, 2 in C₆H₆) (97.2% ee).

N-Ac:

C₁₁H₁₃NO 175.23Cryst. (petrol). Mp 89°. [α]_D +59.6 (EtOH).

N-Benzoyl:

C₁₆H₁₅NO 237.301Cryst. (EtOH aq.). Mp 119°. [α]_D +37 (EtOH).

(S)-form [22160-09-4]

Liq. with blue fluor. Bp 228-229°. [α]_D²⁰ -16.3 (c, 0.5 in CHCl₃) (99% ee). [α]_D²³ -11 (c, 0.11 in C₆H₆) (94% ee).

Hydrochloride:

Cryst. (C₆H₆). Mp 58°. [α]_D +1.7 (H₂O).

N-Ac:

Cryst. (petrol). Mp 89°. [α]_D -61.9 (EtOH).

N-Benzoyl:

Cryst. (EtOH). Mp 119°. [α]_D -37.1 (EtOH).

(±)-form [138380-84-4]

Constit. of the essential oil of *Nepeta nuda*. Bp 228-229° Bp₂₀ 116-116.5°.

Picrate:

Yellow cryst. (C₆H₆). Mp 151°.

N-Formyl: 2,3-Dihydro-2-methyl-1H-indole-1-carboxaldehyde, 9CI. N-Formyl-2-methylindoline [63378-52-9]

C₁₀H₁₁NO 161.203

Needles (petrol). Mp 62-63°.

N-Ac:

Cryst. (petrol). Mp 55-56°.

N-Benzoyl:

Cryst. (EtOH). Mp 91.5°.

N-Benzenesulfonyl:

Cryst. (EtOH). Mp 90°.

N-Nitroso:

C₉H₁₀N₂O 162.191

Yellow cryst. (petrol). Mp 54-55°.

N-Amino: 2,3-Dihydro-2-methyl-1H-indol-1-amine. 1-Amino-2,3-dihydro-2-methyl-1H-indole. 1-Amino-2-methylindoline [31529-46-1]


C₉H₁₂N₂ 148.207

Intermed. for synth. of Indapamide. Cryst. Mp 45°. Bp 255°. Unstable in air.


Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 655C (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 1178D (*ir*)*Aldrich Library of Infrared Spectra*, 3rd edn., 1981, 791G (*ir*)Kost, A.N. et al., *Zh. Obshch. Khim.*, 1959, **29**, 3820-3824; *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 3782-3786 (N-amine)Karger, B.L. et al., *Anal. Chem.*, 1967, **39**, 228-230 (*glc*, *resoln*)von Koch-Pomeranz, U. et al., *Helv. Chim. Acta*, 1977, **60**, 768-797 (*synth*)Kikugawa, Y. et al., *Chem. Pharm. Bull.*, 1978, **26**, 108-110 (*synth*)Fanso-Free, S.N.Y. et al., *J.A.C.S.*, 1979, **101**, 1549-1553 (N-15 *nmr*)Rogovik, V.M. et al., *Zh. Prikl. Khim. (Leningrad)*, 1981, **54**, 1356-1359 (*synth*)Kikugawa, Y. et al., *Synthesis*, 1982, 785-787 (*synth*, N-formyl, *ir*, *pmr*)De Pooter, H.L. et al., *Phytochemistry*, 1987, **26**, 2311-2314 (*occur*)Tripathi, S.R. et al., *Indian J. Phys., B*, 1989, **63**, 240-244; 346-363 (*ir*, *Raman*, *uv*)Peyrot, L. et al., *J. Het. Chem.*, 2001, **38**, 885-893 (N-amino, *ir*, *uv*, *pmr*, *cmr*, *ms*)Krasnov, V.P. et al., *Tetrahedron: Asymmetry*, 2003, **14**, 1985-1988 (*resoln*, R-form)Arp, F.O. et al., *J.A.C.S.*, 2006, **128**, 13368-13368; s1-s31 (S-form, *synth*, N-Ac)Gotor-Fernández, V. et al., *Tetrahedron: Asymmetry*, 2006, **17**, 2558-2564 (S-form, *synth*)

2,3-Dihydro-1-(2-methyl-3-oxodecanoyl)pyrrole D-473


1-(2,3-Dihydro-1H-pyrrol-1-yl)-2-methyl-1,3-decanedione. N-(2-Methyl-3-oxodecanoyl)-2-pyrroline

C₁₅H₂₅NO₂ 251.368**(+)-form**Prod. by *Penicillium brevicompactum*. Exhibits anti-juvenile hormone activity. [α]_D²⁰ +27 (c, 0.07 in CHCl₃).Moya, P. et al., *J.O.C.*, 1998, **63**, 8530-8535 (isol, synth)**1-(2,5-Dihydro-4-methyl-5-oxo-1H-pyrrol-2-yl)-4-methyl-5-oxoproline, 9CI** D-474

1-(3-Methyl-2-oxo-5-pyrrolinyl)-4-methyl-5-oxo-2-pyrrolidinedicarboxylic acid [145042-04-2]

C₁₁H₁₄N₂O₄ 238.243Alkaloid from bulbs of *Lilium hansonii* (Liliaceae). Syrup. [α]_D²⁷ +40.9 (c, 0.43 in MeOH).**Me ester:**Syrup. [α]_D²⁶ +40 (c, 0.30 in MeOH).Ori, K. et al., *Phytochemistry*, 1992, **31**, 2767-2775 (isol, ir, pmr, cmr, ms, struct)**4,5-Dihydro-2-(1-methylpropyl)thiazole** D-475


2-sec-Butyl-2-thiazoline [56367-27-2]



(R)-form

C₇H₁₃NS 143.252**(R)-form**Oil. Bp₁₁ 69-71°. [α]_D²² -21.3 (c, 1 in CHCl₃) (91.6% ee).**(S)-form**Oil. Bp₈ 61-63°. [α]_D²² +21.6 (c, 1 in CHCl₃) (92.6% ee).**(ξ)-form**Pheromone of the common mouse *Mus musculus*. Preorbital gland secretion of grey duiker *Sylvicapra grimmia*. Androgen-dependent substance eliciting male aggressive behaviour.Fomum, Z.T. et al., *Tet. Lett.*, 1975, 1101 (synth)Novotny, M. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1985, **82**, 2059Harvey, S. et al., *J. Chem. Ecol.*, 1989, **15**, 2061Burger, B.V. et al., *J. Chem. Ecol.*, 1990, **16**, 397Tashiro, T. et al., *Eur. J. Org. Chem.*, 1999, 2167-2173 (synth, ir, pmr, cmr)**2,3-Dihydro-3-methyl-1H-pyrrole** D-476


β-Methylpyrroline

C₅H₉N 83.133**(+)-form**Alkaloid from black pepper *Piper nigrum* (Piperaceae).**Hydrochloride:**V. hygroscopic, long needles. [α]_D +2.77 (c, 11.22 in H₂O).**Picrate:**


Yellow powder or prisms. Mp 165°.

Pictet, A. et al., *Helv. Chim. Acta*, 1927, **10**, 593-595 (isol, struct)**3,4-Dihydro-3-methyl-2H-pyrrole-2-carboxylic acid** D-477


4-Methyl-1-pyrroline-5-carboxylic acid

C₆H₉NO₂ 127.143**(2R,3R)-form**Residue present in Pyrrolysine. Oil (as Me ester). [α]_D⁰ +7.9 (c, 3.2 in CHCl₃) (Me ester).Hao, B. et al., *Chem. Biol.*, 2004, **11**, 1317-1324 (synth)**7,8-Dihydro-3-methylpyrrolo[1,2-a]pyrimidin-2(6H)-one, 9CI** D-478

[76884-47-4]

C₈H₁₀N₂O 150.18Alkaloid from the roots of *Glycyrrhiza uralensis* (Chinese licorice) (Fabaceae).Ger. Pat., 1980, 3 017 625; *C.A.*, **94**, 121592 (synth)Han, Y.N. et al., *Arch. Pharmacol. Res.*, 1990, **13**, 103; *C.A.*, **113**, 227985 (isol)**4,5-Dihydro-2-methylthiazole, 9CI** D-479

2-Methyl-2-thiazoline, 8CI [2346-00-1]

C₄H₇NS 101.172Sex pheromone of cockroach *Nauphoeta cinerea*. Also from various cooked foods. Reagent used in synth. of aldehydes. Pale yellow liq. Mp -101°. Bp 144.5-145°.

▶ XJ4261470

Picrate: [5243-50-5]

Yellow needles (EtOH). Mp 171-172°.


N-Me: 4,5-Dihydro-2,3-dimethylthiazolium(I+)

C₅H₁₀NS⁺ 116.207

Needles (EtOH) (as iodide). Mp 235-237° (iodide).

N-Et: 3-Ethyl-4,5-dihydro-2-methylthiazolium(I+)

[54654-71-6]

C₆H₁₂NS⁺ 130.234Needles (MeOH/Me₂CO) (as iodide). Mp 191-193° (iodide). CAS no. refers to iodide.Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 824A (ir)Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1333A (nmr)Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 793C (ir)Wenker, H. et al., *J.A.C.S.*, 1935, **57**, 1079 (synth)Brooker, L.G.S. et al., *J.A.C.S.*, 1936, **58**, 662 (N-Et, synth)Kuhn, R. et al., *Annalen*, 1954, **590**, 55 (synth)Meyers, A.I. et al., *J.O.C.*, 1975, **40**, 2021 (use)Suzuki, N. et al., *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3155 (synth, ir, pmr)Mille, G. et al., *J. Mol. Struct.*, 1978, **50**, 247 (ir, Raman)Chen, B.C. et al., *Helv. Chim. Acta*, 1983, **66**, 1537 (N-15 nmr)Laduranty, J. et al., *Bull. Soc. Chim. Fr.*, 1989, 850 (synth)Sirugue, D. et al., *J. Chem. Ecol.*, 1992, **18**, 2261 (occur)**3,5-Dihydro-3-methyl-2H-thiopyrano[4,3,2-cd]indole-2-carboxylic acid, 9CI** D-480

(2R,3S)-form

C₁₂H₁₁NO₂S 233.29**(2R,3S)-form****Chuangxinmycin.** Chuanghsinmycin [63339-68-4]Isol. from *Streptomyces nigrifaciens* MK219. Active against gram-positive and -negative bacteria. Sol. MeOH, DMF, bases, Py, THF, dioxan, Et₂O, EtOH, Me₂CO; fairly sol. CHCl₃; poorly sol. hexane, H₂O, CCl₄, C₆H₆. Mp 186-

189°. $[\alpha]_D^{20}$ -29 (EtOH aq.). λ_{\max} 228 (ε 36800); 285 (sh) (ε 12000); 295 (ε 12600); 306 (ε 12500) (MeOH).

► LD₅₀ (mus, orl) 1770 mg/kg; LD₅₀ (mus, ivn) 600 mg/kg; LD₅₀ (mus, ipr) 875 mg/kg. XN3876900

Me ester: [83602-19-1]

Cryst. (CH₂Cl₂/petrol). Mp 145-146°.

(2S,3R)-form [114819-47-5]

Mp 181-184°.

(2R,3R)-form [115304-71-7]

Mp 120-125°. $[\alpha]_D^{20}$ -34 (EtOH aq.).

(2S,3S)-form [83602-20-4]

Mp 144-147°. $[\alpha]_D^{20}$ +54.5 (EtOH aq.).

[113531-66-1, 119242-31-8, 118918-28-8, 65058-55-1]

Hsu, H.-C. et al., *CA*, 1980, **93**, 2678 (cryst struct)

Kozikowski, A.P. et al., *J.A.C.S.*, 1982, **104**, 7622-7626 (synth, pmr, config)

Japan. Pat., 1983, 83 76 098; *CA*, **99**, 156830

Murase, M. et al., *Chem. Pharm. Bull.*, 1987, **35**, 2656-2660 (synth, bibl)

Guo, X. et al., *Yaoxue Xuebao*, 1987, **22**, 671 (synth, resolt)

Murase, M. et al., *Heterocycles*, 1990, **30**, 905 (synth)

Dickens, M.J. et al., *J.C.S. Perkin 1*, 1992, 323 (synth, bibl, cryst struct)

Ishibashi, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 271 (synth)


Kato, K. et al., *Tet. Lett.*, 1997, **38**, 1805 (synth)

Kato, K. et al., *Tetrahedron*, 2001, **57**, 10055-10062 (synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CML820 (323)

1-[4,9-Dihydro-2-(methylthio)-1,3-thiazino[6,5-b]indol-4-yl]-2-propanone, 9CI

[113866-43-6]



C₁₄H₁₄N₂OS₂ 290.409


Isol. from Chinese cabbage (*Brassica campestris* ssp. *pekinensis*, Brassicaceae) inoculated with the bacterium *Pseudomonas cichorii*. Amorph. solid. λ_{\max} 202 (ε 18900); 227 (ε 28000); 282 (ε 6490); 294 (ε 6690) (MeOH) (Berdy).

Takasugi, M. et al., *Bull. Chem. Soc. Jpn.*, 1988, **61**, 285 (isol, uv, ir, pmr, cmr, ms, struct)

Dihydrontidine

D-482

12,13-Dihydro-2,3-dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridine, 9CI
[13063-06-4]



C₂₁H₁₉NO₄ 349.385

Several numbering schemes in use. Alkaloid from the roots of *Toddalia asiatica* and *Zanthoxylum nitidum*, and from the root bark of *Zanthoxylum myriacanthum* (Rutaceae). Prisms (EtOH). Mp 221-223° (208-211°, 216-217°). λ_{\max} 228 (ε 42660); 278 (ε 36300); 310 (ε 20900) (MeOH) (Derep).

Hydrochloride:

Cryst. + 4H₂O. Mp 215-216° (dec., vac.).

8-Oxo: 2,3-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridin-13(12H)-one. 8,9-Dimethoxy-2,3-methylenedioxybenzo[c]phenanthridin-6(5H)-one. **Oxynitidine**

[548-31-2]

C₂₁H₁₇NO₅ 363.369

Alkaloid from the roots of *Zanthoxylum nitidum*, the wood of *Zanthoxylum inerme* (= *Fagara boninensis*), and the bark of *Zanthoxylum cuspidatum* and *Zanthoxylum ailanthoides* (Rutaceae). Antineoplastic agent. Needles (EtOH). Mp 284-285°. Log P 2.8 (calc). λ_{\max} 251 (ε 38900); 277 (ε 52500); 288 (ε 64600); 320 (ε 15800); 333 (ε 15100); 367 (ε 4270) (EtOH) (Derep).

8-Oxo, O¹⁰-de-Me: **Oxyterihanine**

[95066-54-9]

C₂₀H₁₅NO₅ 349.342

Minor alkaloid from *Zanthoxylum nitidum* (*Fagara nitida*) (Rutaceae). Mp 300°.

8-Oxo, O¹¹-de-Me: **Oxyisoterihanine**

[111154-44-0]

Synthetic. Needles (CHCl₃/MeOH). Mp 300°.

8ξ-Methoxy: 7,8-Dihydro-8-methoxynitidine

[41349-33-1]

C₂₂H₂₁NO₅ 379.412

Alkaloid from *Fagara macrophylla* and *Zanthoxylum nitidum*. Shows antineoplastic activity. Yellow prisms (MeOH). Mp 234-236°. $[\alpha]_D^{20}$ -8 (c, 0.01 in CHCl₃). Log P 4.32 (uncertain value) (calc). λ_{\max} 235 (ε 19500); 270 (ε 34700); 292 (ε 32400); 328 (ε 25700) (MeOH) (Derep).

11-Demethoxy, 12-methoxy, 8-oxo: **Turraeanthine B**

C₂₁H₁₇NO₅ 363.369

Alkaloid from the stem bark of *Turraeanthus africanus*. Brown cryst. (MeOH). Mp 270-272°.

Arthur, H.R. et al., *J.C.S.*, 1959, 1840; 4010 (*Dihydrontidine*, *Oxynitidine*, *isol*, *struct*, *synth*, *uv*)

Gopinath, K.W. et al., *J.C.S.*, 1959, 4012 (*synth*, *uv*)

Gopinath, K.W. et al., *Indian J. Chem.*, 1963, **1**, 99 (*Oxynitidine*, *synth*)

Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 3420 (*uv*)

Torto, F.G. et al., *Phytochemistry*, 1970, **9**, 911 (*pmr*, *ms*)

Ishii, H. et al., *Tet. Lett.*, 1971, 2429 (*pmr*)

Ishii, H. et al., *Yakugaku Zasshi*, 1972, **92**, 118; 1976, **96**, 1458; 1983, **103**, 279; 1984, **104**, 1030 (*Oxynitidine*, *Oxyterihanine*)

Kametani, T. et al., *J. Het. Chem.*, 1973, **10**, 31 (*Oxynitidine*, *synth*)

Ninomiya, I. et al., *J.C.S. Perkin 1*, 1975, 762 (*synth*)

Waterman, P.G. et al., *Phytochemistry*, 1975, **14**, 2530 (*isol*)

Begley, W.J. et al., *J.C.S. Perkin 1*, 1977, 2324 (*Dihydrontidine*, *Oxynitidine*, *synth*, *pmr*, *ms*)

Sharma, P.N. et al., *Indian J. Chem., Sect. B*, 1979, **17**, 299 (*isol*, *uv*, *pmr*)

Huang, Z.-X. et al., *Huaxue Xuebao*, 1980, **38**, 535; *CA*, **94**, 99773e (*isol*)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1987, **35**, 2717 (*Oxyterihanine*, *synth*, *ir*, *pmr*)

Hanaoka, M. et al., *Heterocycles*, 1987, **26**, 1499 (*Oxyterihanine*, *ir*, *pmr*, *synth*)

Wall, M.E. et al., *J. Nat. Prod.*, 1987, **50**, 1095 (*7,8-Dihydro-8-methoxynitidine*)

Clark, R.D. et al., *J.O.C.*, 1988, **53**, 2378; 5192 (*Oxynitidine*, *synth*, *pmr*)

De Moura, N.F. et al., *Phytochemistry*, 1997, **46**, 1443-1446 (*isol*, *pmr*, *cmr*)

Le, T.N. et al., *Chem. Pharm. Bull.*, 2006, **54**, 476-480 (*Oxynitidine*, *synth*)


Vardamides, J.C. et al., *Chem. Pharm. Bull.*, 2006, **54**, 1034-1036 (*Turraeanthine B*)

Le, T.N. et al., *Arch. Pharmacol. Res.*, 2008, **31**, 6-9 (*Oxyterihanine*, *synth*)

Yang, G. et al., *Chem. Biodiversity*, 2008, **5**, 1718-1722 (*7,8-Dihydro-8-methoxynitidine*)

1-(2,3-Dihydro-2-oxo-3-furanyl)-5-(hydroxymethyl)-1H-pyrrole-2-carboxaldehyde, 9CI

[112663-86-2]



C₁₀H₉NO₄ 207.185

Constit. of the roots of *Pisum sativum* seedlings. Cell cycle regulator.


Lynn, D.G. et al., *J.A.C.S.*, 1987, **109**, 5858 (*isol*, *props*)

Evans, L.S. et al., *Phytochemistry*, 1987, **26**, 2891 (*isol*, *props*)

2,3-Dihydro-2-oxo-1H-imidazole-4-carboxylic acid, 9CI

2-Oxo-4-imidazolinecarboxylic acid.

Glyoxalone-4-carboxylic acid
[39828-47-2]



C₄H₄N₂O₃ 128.087

Prod. from the venom of *Crotalus adamanteus*. Mp 251°. pK_a 3.37.

Et ester: [71123-17-6]

C₆H₈N₂O₃ 156.141

Plates (EtOH aq.). Mp 255°.

Otter, B.A. et al., *J.O.C.*, 1968, **33**, 3593 (*synth*, *spectra*)


Cooper, A.J.L. et al., *J. Biol. Chem.*, 1973, **248**, 8499 (*biosynth*)

Stanovnik, B. et al., *Heterocycles*, 1979, **12**, 761 (*synth*)

2,3-Dihydro-2-oxo-1H-indole-3-acetic acid, 9CI

D-485

2-Hydroxy-1H-indole-3-acetic acid. Oxindole-3-acetic acid. 2-(2-Oxo-3-indoliny-1)acetic acid
[2971-31-5]

C₁₀H₉NO₃ 191.186

No mention of stereoisomers in lit. Enantiomers presumably interconvert through the enol tautomer. Isol. from *Ribes rubrum*, *Brassica* spp. and *Helianthus annuus*. Prod. of catabolism of 1H-Indole-3-acetic acid, I-81 in *Zea mays* (sweetcorn). Needles (Me₂CO/C₆H₆). Mp 146°.

Me ester: [110425-83-7]
[61989-29-5]

C₁₁H₁₁NO₃ 205.213

Isol. from rice bran. Prob. intermed. in biosynth. of 2,6-Dihydroxy-4-quinolinecarboxylic acid, D-672 from 1H-Indole-3-acetic acid, I-81. Amorph. solid or cryst. (Me₂CO or MeOH). Mp 187-188° (171-173°). Dimorphic. Amorph. solid with lower mp obt. by rapid cooling of solns. and cryst. obt. by slow cooling.

Et ester: [40940-16-7]

C₁₂H₁₃NO₃ 219.24


Cryst. (petrol). Mp 94-98°.

β-D-Glucopyranosyl ester: [945774-92-5]

C₁₆H₁₉NO₈ 353.328Metab. of 1H-Indole-3-acetic acid, I-81 in *Arabidopsis* sp.Julian, P.L. et al., *J.A.C.S.*, 1953, **75**, 5305-5309 (synth)Hinman, R.L. et al., *J.O.C.*, 1964, **29**, 1206-1215 (synth)Kinashi, H. et al., *Agric. Biol. Chem.*, 1976, **40**, 2465-2470 (isol)Nonhebel, H.M. et al., *J. Biol. Chem.*, 1985, **260**, 12685-12689 (biosynth)Takase, S. et al., *Tetrahedron*, 1986, **42**, 5879-5886 (synth, ir, pmr)Lower, P. et al., *J.C.S. Perkin 1*, 1987, 753-757 (synth, pmr)Rivera-Becerril, E. et al., *Heterocycles*, 2006, **68**, 1459-1466 (Me ester, cryst struct)Kai, K. et al., *Phytochemistry*, 2007, **68**, 1651-1663 (glucosyl ester)**2-(1,3-Dihydro-3-oxo-2H-indol-2-ylidene)-7-hydroxycyclopent[gl]indol-3(2H)-one**

D-486

[868837-50-7]

C₁₉H₁₀N₂O₃ 314.3


Isol. from *Escherichia coli* DH10B with inserted genes for toluene-4-monooxygenase.

McClay, K. et al., *Appl. Environ. Microbiol.*, 2005, **71**, 5476-5483 (isol)

α-(1,3-Dihydro-3-oxo-2H-indol-2-ylidene)-1H-indole-3-acetonitrile, 9CI

D-487

2-[Ciano(3-indolyl)methylene]-3-indolone
[359013-25-5]

C₁₈H₁₁N₃O 285.304


Alkaloid from the roots of *Isatis indigotica*. Purple needles. Mp 213-215°. λ_{max} 274 ; 505 (MeOH).

Chen, W.S. et al., *Chin. Chem. Lett.*, 2001, **12**, 501-502 (isol, pmr, cmr)

2,3-Dihydro-1-oxo-1H-pyrrolizine-3-carboxylic acid, 9CI

D-488

4-Oxo-1-azabicyclo[3.3.0]octa-5,7-diene-2-carboxylic acid

C₈H₇NO₃ 165.148

Metab. of *Streptomyces olivaceus*. Oil (as Me ester).


[80314-99-4 ; 80314-96-1]

Box, S.J. et al., *Tet. Lett.*, 1981, **22**, 3293 (isol, struct, synth)

1,5-Dihydro-1-[1-oxo-3-(3,4,5-trimethoxyphenyl)propyl]-2H-pyrrol-2-one, 9CI

D-489

[124727-05-5]


C₁₆H₁₉NO₅ 305.33

Alkaloid from the aerial parts of *Piper demeraranum* (Piperaceae). Needles (petrol/Me₂CO). Mp 150-151°.

Maxwell, A. et al., *J. Nat. Prod.*, 1989, **52**, 891 (isol, uv, ir, pmr, cmr, ms, struct)

5,6-Dihydro-2-(1,3-pentadienyl)-4(1H)-pyridinone

D-490

C₁₀H₁₃NO 163.219*(E,E)-form***Streptazone D**Prod. by *Streptomyces* sp. FORM5.


Yellow solid. Mp 105-109°. λ_{max} 288 (log ε 4.34); 361 (log ε 4.11) (MeOH).

Puder, C. et al., *J. Nat. Prod.*, 2000, **63**, 1258-1260

Dihydroperaksine

D-491

19,20-Dihydro-21-methyl-18-norsarpagan-17,19-diol, 9CI. Dihydrovomifoline
[26263-40-1]



Absolute Configuration

C₁₉H₂₄N₂O₂ 312.411

Different numbering systems used. Alkaloid from leaves of *Rauwolfia caffra* (Apocynaceae). Prisms by subl. Mp 290-291° (part subl. at 280°). [α]_D²¹ +40.8 (c, 1 in Py). pK_a 7.75 (50% MeOH aq.).

17-Aldehyde, 17→19 hemiacetal: see Peraksine, P-241

19-Aldehyde, 17-Ac: **O-Acetylpreperaksine**. Preperaksine O-acetate

[73221-32-6]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the stem bark of *Rauwolfia volkensii*. Off-white powder. [α]_D²⁵ +45 (c, 0.1 in CHCl₃). Stereochem. not determined. λ_{max} 226 (log ε 4.24); 283 (log ε 3.74); 290 (log ε 3.68) (MeOH).

20,21-Diepimer: **20R,21S-Dihydroperaksine**

[16100-84-8]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from hairy root cultures of *Rauwolfia serpentina*. Cryst. (MeOH). Mp 208-212°. Subl. 192. [α]_D²⁵ +35.3 (c, 0.12 in Py). λ_{max} 225 (log ε 4.49); 279 (log ε 3.83); 289 (log ε 3.73) (MeOH).

20,21-Diepimer, 17-aldehyde: **20R,21S-Dihydroperaksin-17-al**

[451478-46-9]

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from hairy root cultures of *Rauwolfia serpentina*. Powder. [α]_D²⁵ +50 (c, 0.07 in MeOH). λ_{max} 224 (log ε 4.21); 279 (log ε 3.55); 289 (log ε 3.45) (MeOH).

20,21-Diepimer, 10-hydroxy: **20R,21S-Dihydro-10-hydroxyperaksine**

[451478-47-0]


C₁₉H₂₄N₂O₃ 328.41

Alkaloid from hairy root cultures of *Rauwolfia serpentina*. Cryst. (MeOH). Mp 323-335°. [α]_D²⁵ +42.6 (c, 0.15 in Py). λ_{\max} 226 (log ϵ 4.14); 277 (log ϵ 3.71) (MeOH).

[11015-98-8]

Kiang, A.K. *et al.*, *Tetrahedron*, 1966, **22**, 3293-3300 (*synth*, *ms*)Akinloye, B.A. *et al.*, *Planta Med.*, 1979, **37**, 361-366 (*O-Acetylpreperakine*)Nasser, A.M.A.G. *et al.*, *Phytochemistry*, 1983, **22**, 2297-2300 (*isol*)Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1456-1460 (*rev*)Sheludko, Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1006-1010 (*20,21-diepipimers*)**5,6-Dihydro-3-phenyl-4H-pyrrolo[1,2-b]pyrazole, 9CI** **D-492**

Withasomnine. *Newbouldine*†
[10183-74-1]

C₁₂H₁₂N₂ 184.24

The struct. of Newbouldine (Mp 215-217°) *isol.* by Houghton *et al.* and claimed to be novel corresponds with that of Withasomnine. The large discrepancy in Mps is noteworthy. See also Newbouldine, N-185. Alkaloid from the roots of *Withania somnifera* (Solanaceae) and root bark of *Newbouldia laevis* (Bignoniaceae). Cryst. (petrol). Mp 117-118°.

Hydrochloride:

Hygroscopic cryst. Mp 115-117°.

Picrate:Cryst. (H₂O). Mp 170-173°.

4'-Hydroxy: 4'-Hydroxywithasomnine
[155416-37-8]

C₁₂H₁₂N₂O 200.24

Alkaloid from root bark of *Newbouldia laevis* (Bignoniaceae). Cryst. (Me₂CO). Mp 238-239°. Hydroxylated in the phenyl ring.

4'-Methoxy: 4'-Methoxywithasomnine

C₁₃H₁₄N₂O 214.266

Alkaloid from the root bark of *Newbouldia laevis*. Methoxylated in the phenyl ring. λ_{\max} 253 (log ϵ 4.33) (MeOH).

Schröter, H.-B. *et al.*, *Tetrahedron*, 1966, **22**, 2895 (*isol*, *uv*, *ir*, *ms*, *pmr*, *struct*)Morimoto, T. *et al.*, *Tet. Lett.*, 1968, 5707 (*synth*)Onaka, T. *et al.*, *Tet. Lett.*, 1968, 5711 (*synth*)O'Donovan, D.G. *et al.*, *Tet. Lett.*, 1970, 3637 (*biosynth*)Onaka, T. *et al.*, *CA*, 1972, **77**, 48328e (*synth*)Ranganathan, D. *et al.*, *Synth. Commun.*, 1985, **15**, 259 (*synth*, *ir*, *pmr*, *ms*)Guzmán-Pérez, A. *et al.*, *Synth. Commun.*, 1991, **21**, 1667 (*synth*)Adesanya, S.A. *et al.*, *Phytochemistry*, 1994, **35**, 1053 (*4'-Hydroxywithasomnine*)Houghton, P.J. *et al.*, *Phytochemistry*, 1994, **35**, 1602Kulinkovich, O. *et al.*, *Tet. Lett.*, 1996, **37**, 1095 (*synth*)Aladesanmi, A.J. *et al.*, *Planta Med.*, 1998, **64**, 90-91 (*4'-Methoxywithasomnine*)**3,4-Dihydro-5-propanoyl-2H-pyrrole** **D-493**


1-(3,4-Dihydro-2H-pyrrol-5-yl)-1-propanone, *9CI*. *2-Propionyl-1-pyrroline*
[133447-37-7]

C₇H₁₁NO 125.17

Formed by thermal treatment of proline and glucose mixtures. Constit. of freshly popped corn aroma. Bp₇₀ 115°. Odour threshold 0.02 ng/L in air.

Schieberle, P. *et al.*, *J. Agric. Food Chem.*, 1991, **39**, 1141-1144 (*detn*, *synth*, *pmr*, *ms*)Hofman, T. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 2721-2726 (*occur*, *ms*)Fuganti, C. *et al.*, *Tetrahedron*, 2007, **63**, 4762-4767 (*synth*, *pmr*, *cmr*)**1,5-Dihydro-2,4,6(3H)-pteridinetrione, 9CI** **D-494**

2,4,6-(1H,3H,5H)-Pteridinetrione. *6-Oxohumazine*
[2577-35-7]

C₆H₄N₄O₃ 180.123

Isol. from honey bee (*Apis mellifera*).

1-Me: [50996-37-7]C₇H₆N₄O₃ 194.149

Yellow-brown cryst. (H₂O). Mp 335-338° dec.

3-Me: [58947-87-8]C₇H₆N₄O₃ 194.149

Yellow-brown cryst. Mp 350°.

7-Me:C₇H₆N₄O₃ 194.149

Yellow-brown cryst. (H₂O). Mp 340°.

1,3-Di-Me: [61846-18-2]C₈H₈N₄O₃ 208.176

Yellowish cryst. (H₂O). Mp 300°.

1,7-Di-Me:C₈H₈N₄O₃ 208.176

Cryst. (H₂O). Mp 315° dec.

3,7-Di-Me:C₈H₈N₄O₃ 208.176


Mp 340°.

1,3,5-Tri-Me:C₉H₁₀N₄O₃ 222.203

Yellow needles (H₂O). Mp 180-181°.

Pfleiderer, W. *et al.*, *Chem. Ber.*, 1957, **90**, 2604; 2617 (*synth*)McNutt, W.S. *et al.*, *Biochemistry*, 1962, **1**, 1161 (*metab*)Dustmann, J.H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1971, **352**, 1599 (*isol*)**5,6-Dihydro-2(1H)-pyridinone** **D-495**


5,6-Dihydro-2(1H)-pyridone, *8CI*. *3,4-Dehydro-2-piperidone*. *5-Amino-2-pentenoic acid γ -lactam*. *5,6-Dihydropyridin-2-ol*
[6052-73-9]

C₅H₇NO 97.116

Alkaloid from *Bunchosia pseudonitida* and *Piper sintonense*. Constit. of tobacco smoke. Cryst. (Et₂O/petrol). Mp 63-65°. λ_{\max} 209 (log ϵ 4.24); 247 (sh) (log ϵ 3.52) (EtOH).

Schumacher, J.N. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 310 (*occur*)Boll, P.M. *et al.*, *Tetrahedron*, 1984, **40**, 171 (*synth*)Calle, A.A. *et al.*, *CA*, 1986, **104**, 126581 (*isol*)Hua, D.H. *et al.*, *J.O.C.*, 1994, **59**, 5084 (*synth*, *pmr*, *cmr*)Chen, J.-J. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2058-2064 (*isol*, *pmr*, *cmr*, *ms*)Simon, L. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 1682-1701 (*synth*, *ir*, *pmr*, *cmr*)**3,4-Dihydro-5-(2-pyridinyl)-2H-pyrrole-2-carboxylic acid** **D-496**

2-(2-Pyridyl)- Δ^1 -pyrroline-5-carboxylic acid. *Proferrorosamine A*. *Pyrimine*
[26927-08-2]

*(R)-form*C₁₀H₁₀N₂O₂ 190.201

Prod. by *Erwinia rhapontici* and *Pseudomonas roseus fluorescens*.

*(R)-form**NH₄ salt*: [100016-79-3][α]_D²⁵ -37 (c, 4 in H₂O).*NH₄ salt; hydrochloride (1:2)*:

Needles (Me₂CO aq.). [α]_D²⁴ -26 (c, 5 in 1M HCl). Sinters at 155°, dec. at 179°.

Fe complex (3:1): Neopurpuratin

[80124-41-0]

C₃₀H₂₈FeN₆O₆ 624.435

Prod. by *Streptomyces propurpuratus*.

Purple-red pigment. Sol. H₂O, EtOH,

MeOH. λ_{\max} 277 (E1%/1cm 388); 355

(E1%/1cm 60); 558 (E1%/1cm 168)

(H₂O) (Berdy).*(S)-form*

Prod. by *Pseudomonas* GH. Exists in this form >pH2.

Fe complex (3:1): Ferropyrimine


Prod. by *Pseudomonas* GH. Purple-red pigment.

*(±)-form**Fe complex (3:1): Ferrorosamine A*

[59568-22-8]
 $C_{30}H_{28}FeN_6O_6$ 624.435
 From *Erwinia rhapontici* and *Pseudomonas roseus fluorescens*. Red powder.
 Shiman, R. et al., *Biochemistry*, 1965, **4**, 2233 (isol, struct)
 Helbling, A.M. et al., *Helv. Chim. Acta*, 1976, **59**, 938 (synth)
 Feistner, G. et al., *Curr. Microbiol.*, 1983, **8**, 239 (isol)
 Handa, A. et al., *Agric. Biol. Chem.*, 1985, **49**, 2227 (*Neopurpuratin*)
 Ohshima, M. et al., *J. Ferment. Technol.*, 1985, **63**, 79 (*Neopurpuratin*)
 Vande Woestyne, M. et al., *Appl. Environ. Microbiol.*, 1991, **57**, 949 (isol)

Dihydro-2,4(1H,3H)-pyrimidinedione, 9CI D-497

Hydrouracil, 8CI. *Dihydrouracil*. *Hexahydro-2,4-dioxypyrimidine*
 [504-07-4]



$C_4H_6N_2O_2$ 114.104
 Component of transfer ribonucleic acid (tRNA). Cryst. (Me₂CO). Mp 275-276°.


l-Nitroso: [16813-36-8]

$C_4H_5N_3O_3$ 143.102

► Exp. carcinogen. LD₅₀ (rat, ipr) 850 mg/kg. MX9280000

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1319C (nmr)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 811B (ir)
 Fischer, E. et al., *Ber.*, 1901, **34**, 3751 (synth)
 Rohrer, D. et al., *Chem. Comm.*, 1968, 746 (cryst struct)
 Elad, D. et al., *Chem. Comm.*, 1968, 879 (synth)
 Takemoto, K. et al., *Chem. Lett.*, 1972, 767 (synth)
 Sax, N.I. et al., *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 867
 Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NJY000

4,8-Dihydropyrimido[5,4-e]-1,2,4-triazine-3,5,7(6H)-trione D-498



$C_5H_3N_5O_3$ 181.11

Many tautomers possible.

6,8-Di-Me: 2,8-Dihydro-6,8-dimethylpyrimido[5,4-e]-1,2,4-triazine-3,5,7(6H)-trione. *Fervenuone*

[18969-83-0]
 $C_7H_7N_5O_3$ 209.164
 Mp 260-261° (anhyd.). Forms a monohydrate and a monoethanolate.

► UW7910000

2,6,8-Tri-Me: 2,8-Dihydro-2,6,8-trimethylpyrimido[5,4-e]-1,2,4-triazine-3,5,7(6H)-trione. *2-Methylfervenuone*. *Antibiotic MSD 92*. *Antibiotic 2096C*. 2096C. *MSD 92*
 [22712-32-9]

$C_8H_9N_5O_3$ 223.191

Isol. from *Streptomyces* spp. Broad-spectrum antibiotic. Inhibitor of protein tyrosine phosphatase. Yellow plates (EtOH, H₂O or EtOAc). Sol. MeOH, Et₂O; fairly sol. H₂O; poorly sol. hexane. Mp 92-93° Mp 157-158° Mp 174-175° Mp 181-182° (tetramorph.). λ_{max} 240 (ε 19000); 280 (sh) (ε 1940); 415 (ε 3410) (H₂O) (Derep). λ_{max} 218 (ε 12100); 288 (ε 6530); 413 (ε 1004) (MeOH) (Derep). λ_{max} 238; 417 (MeOH-THCONH₂) (Berdy).

► LD₅₀ (mus, ipr) 1.2 mg/kg. UW7960000

4,6,8-Tri-Me: 4,8-Dihydro-4,6,8-trimethylpyrimido[5,4-e]-1,2,4-triazine-3,5,7(6H)-trione, 9CI. *4-Methylfervenuone*
 [22712-42-1]

$C_8H_9N_5O_3$ 223.191

Shows antibiotic props. Pale yellow plates (EtOH). Mp 218-220°.

Miller, T.W. et al., *Antimicrob. Agents Chemother.*, 1963, 58-62 (2-*Methylfervenuone*, isol)

Taylor, E.C. et al., *J.O.C.*, 1975, **40**, 2321 (synth, bibl)

Ichiba, M. et al., *J.O.C.*, 1978, **43**, 469 (synth, bibl)

Wang, H. et al., *J. Nat. Prod.*, 2000, **63**, 1641-1646 (2-*Methylfervenuone*, isol, pmr, cmr, N-15 nmr, activity)

2,3-Dihydro-1H-pyrrole, 9CI D-499

2-Pyrroline, 8CI

[638-31-3]



C_4H_7N 69.106

N-Ac: [23105-58-0]

C_6H_9NO 111.143

Odorous constit. of cooked rice and of pandan rampeh (*Pandanus amaryllifolius*). Oil.

N-Ph: 2,3-Dihydro-1-phenyl-1H-pyrrole.

l-Phenyl-2-pyrroline

$C_{10}H_{11}N$ 145.204

Cryst. Mp 166-167.5°.

N-tert-*Butoxycarbonyl*:

$C_9H_{15}NO_2$ 169.223

Wittig, G. et al., *Annalen*, 1955, **594**, 1 (*N*-Ph)

Sato, S. et al., *CA*, 1969, **71**, 21828 (deriv)

Leblanc, R. et al., *Tet. Lett.*, 1969, 2441 (deriv)

Westheimer, F.H. et al., *J.O.C.*, 1971, **36**, 1570 (deriv)

Buttery, R.G. et al., *Chem. Ind. (London)*,

1983, 478 (isol, deriv)

Marais, W. et al., *Synth. Commun.*, 1998, **28**, 3681-3691 (*N*-Ac, *N*-benzoyl)
Org. Synth., 2008, **85**, 64-71 (*N*-Boc)

3,4-Dihydro-2H-pyrrole, 9CI D-500

l-Pyrroline. *FEMA 3898*

[5724-81-2]



C_4H_7N 69.106

Male sex pheromone of Mediterranean fruit fly *Ceratitis capitata*. Present in clam and squid. Primary odour constit. of sperm. Stable in dil. soln., trimerises readily.

N-Oxide: [24423-88-9]

C_4H_7NO 85.105

Cyclic nitron used in intramol. 1,3-dipolar cycloadditions. Bp_{0.1} 74-76°.

Amoore, J.E. et al., *J. Chem. Ecol.*, 1975, **1**, 299

Poisel, H. et al., *Monatsh. Chem.*, 1978, **109**, 925 (synth, pmr)

Baker, R. et al., *Chem. Comm.*, 1985, 824

Bock, H. et al., *Chem. Ber.*, 1987, **120**, 1961 (synth, pe, bibl)

Guillemin, J.-C. et al., *Tetrahedron*, 1988, **44**, 4447 (synth, pmr)

Murahashi, S. et al., *J.O.C.*, 1990, **55**, 1736-1744 (*N*-oxide, synth, ir, pmr)


Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **6**, 4389-4390 (*N*-oxide, use)

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 1568

3,4-Dihydro-2H-pyrrole-2-carboxylic acid, 9CI D-501

l-Pyrroline-5-carboxylic acid, 8CI. *l*,5-Didehydropyrroline

[2906-39-0]



(*S*)-form

$C_5H_7NO_2$ 113.116

Dimerises in solution. Cyclic anhydride of 2-Amino-5-oxopentanoic acid.

(*S*)-form

L-form

[64199-88-8]

Intermed. in biosynth. and biodegradn. of Proline.

Et ester: [172879-74-2]

$C_7H_{11}NO_2$ 141.169

[α]_D²⁵ +14.1 (c, 8.5 in CHCl₃).

(±)-form [23141-14-2]

Cryst. (H₂O). Mp 140-142° dec.

Et ester: Liq. Bp₁₇ 110°.

Vogel, H.J. et al., *J.A.C.S.*, 1952, **74**, 109-112 (synth, bibl)


Osugi, K. et al., *Yakugaku Zasshi*, 1958, **78**,

1332-1338; *CA*, **53**, 8109 (synth, ester)

Strecker, H.J. et al., *Methods Enzymol.*, Part B, 1971, **17**, 254-258 (synth)

- Mezl, V.A. *et al.*, *Anal. Biochem.*, 1976, **74**, 430-440 (*synth*)
 Smith, R.J. *et al.*, *Anal. Biochem.*, 1977, **82**, 170-176 (*synth*)
 De March, F. *et al.*, *Chem. Comm.*, 1995, 2097-2098 (*Et ester, synth*)

2,3-Dihydro-1H-pyrrolo[2,1-c][1,4]benzodiazepine-5,11(10H,11aH)-dione, 9CI **D-502**
Cycloanthranilylproline




C₁₂H₁₂N₂O₂ 216.239

(R)-form [154802-75-2]
 Prisms (EtOAc). Mp 221-222°. [α]_D²⁵ -491 (c, 0.2 in CHCl₃).

(S)-form [18877-34-4]

Alkaloid from the roots of *Isatis indigotica*. Isol. from the myxomycete *Fuligo candida*. Cytotoxic. Cryst. (Me₂CO/hexane). Mp 229° (223-225°). [α]_D²⁵ +567 (c, 1 in DMF). [α]_D²⁰ +505 (c, 0.1 in MeOH).
 Wright, W.B. *et al.*, *J. Med. Chem.*, 1978, **21**, 1087 (*synth*)
 Mori, M. *et al.*, *Tetrahedron*, 1986, **42**, 3793-3806 (*synth, ir, pmr, cmr*)
 Feigel, M. *et al.*, *Z. Naturforsch., B*, 1990, **45**, 258-266 (*synth, pmr, cmr, cryst struct*)
 Kamal, A. *et al.*, *J.O.C.*, 1991, **56**, 2237 (*synth*)
 Fryer, R.I. *et al.*, *Med. Chem. Res.*, 1993, **3**, 183 (*R-form*)
 Wu, X. *et al.*, *Planta Med.*, 1997, **63**, 55-57 (*isol*)
 Nakatani, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 368-370 (*isol, pmr, cmr*)
 Tang, H. *et al.*, *Eur. J. Org. Chem.*, 2008, 126-135 (*S-form, synth, pmr*)

2,3-Dihydropyrrolo[2,1-b]quinazolin-9(1H)-one, 9CI **D-503**
Deoxyvasicinone
 [530-53-0]



C₁₁H₁₀N₂O 186.213

Major alkaloid from the foliage of *Peganum nigellastrum*. Also present in *Macklinaya macrosciadia*, the seeds and above-ground parts of *Peganum harmala* and the roots of *Adhatoda vasica* (Zygophyllaceae, Araliaceae, Acanthaceae). Claimed to show cholinergic activity. Needles (C₆H₆/petrol). Mp 110-111°. λ_{max} 206 (ε 32260); 224 (ε 34670); 256 (ε 9550); 302 (ε 4790); 313 (ε 3980) (MeOH) (Berdy).

N^d-Oxide: *Deoxyvasicinone N-oxide*
 [168781-18-8]

C₁₁H₁₀N₂O₂ 202.212
 Alkaloid from above-ground parts of *Nitraria komarovii* (Zygophyllaceae). Cryst. (Me₂CO). Mp 152-153°.

9ξ-Alcohol: 1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazolin-9-ol. *Peganol*. 9-Hydroxydeoxypeganine
 [36101-54-9]

C₁₁H₁₂N₂O 188.229

Alkaloid from *Peganum harmala* (Zygophyllaceae). Cryst. (MeOH). Mp 178-180°.

9ξ-Alcohol, N¹⁰-oxide: *Peganol N-oxide*
 [335373-42-7]

C₁₁H₁₂N₂O₂ 204.228

Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (EtOH). Mp 182-183°. λ_{max} 207 (log ε 4.08); 270 (log ε 3.62) (EtOH).

2,3-Didehydro, 9ξ-alcohol: 9-Hydroxy-2,3-didehydrodeoxypeganine
 [119459-56-2]

C₁₁H₁₀N₂O 186.213

Alkaloid from aerial parts of *Galium aparine* collected at the flowering period. Amorph. [α]_D -44.

3-Hydroxy: see Vasicinone, V-38

5-Hydroxy: 5-Hydroxydeoxyvasicinone.

2,3-Dihydro-5-hydroxypyrrolo[2,1-b]quinazolin-9(1H)-one, 9CI
 [35214-95-0]

C₁₁H₁₀N₂O₂ 202.212

Metab. of *Klebsiella pneumoniae* var. *oxytoca*. Mp 158-160°. Solns. rapidly turn brown in light.

Petersen, S. *et al.*, *Annalen*, 1959, **623**, 166

(*synth, 5-Hydroxydeoxyvasicinone*)

Chatterjee, A. *et al.*, *Phytochemistry*, 1968, **7**, 307 (*isol, synth*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1971, **24**, 223 (*isol*)

Telezhenetskaya, M.V. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 849; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 835 (*Peganol*)

Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4387 (*synth, 5-Hydroxydeoxyvasicinone*)

Johne, S. *et al.*, *J. Prakt. Chem.*, 1977, **319**, 919

(*synth, cmr*)

Mohr, N. *et al.*, *Annalen*, 1981, 1515 (*isol, uv, ir, pmr, cmr, ms, struct, 5-Hydroxydeoxyvasicinone*)

Al-Shamma, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 745 (*isol, uv, ir, pmr, ms*)

Johne, S. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 129 (*pharmacol, rev*)

Sener, B. *et al.*, *Gazi Univ. Eczacilik Fak. Derg.*, 1988, **5**, 33 (*9-Hydroxy-2,3-didehydrodeoxypeganine*)

Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 580; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 509; 2000, **36**, 76-78 (*Deoxyvasicinone N-oxide, Peganol N-oxide*)

Turgunov, K.K. *et al.*, *Khim. Prir. Soedin.*, 1995, **31**, 849-854; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 714-718 (*cryst struct*)


Mhaske, S.B. *et al.*, *J.O.C.*, 2002, **66**, 9038-9040 (*synth*)

Liu, J.-F. *et al.*, *Org. Lett.*, 2005, **7**, 3363-3366 (*synth, pmr, cmr*)

Hamid, A. *et al.*, *Tet. Lett.*, 2006, **47**, 1777-1781 (*synth*)

Bowman, W.R. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 103-113 (*synth*)

2-(3,4-Dihydro-2H-pyrrol-5-yl)pyridine **D-504**
 2-(1-Pyrrolin-2-yl)pyridine, 8CI. 2-(2-Pyridyl)-1-pyrroline. *Apoferrosamine*
 [4593-27-5]




C₉H₁₀N₂ 146.191

Produced by dist. of Ferrerosamine, a red Fe pigment prod. by adding Fe²⁺ to cultures of *Bacillus roseus fluorescens*. Poorly sol. hexane. Mp 55-55.5°. Bp_{0.3} 65°.

Pouteau-Thouvenot, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 3238 (*synth*)

Stevens, R.V. *et al.*, *J.A.C.S.*, 1968, **90**, 5576 (*synth*)

5,10-Dihydro-11H-quinolin-11-one, 9CI **D-505**
 5H,10H-Indolo[3,2-b]quinolin-11-one. *Quindolinone*
 [80289-15-2]



C₁₅H₁₀N₂O 234.257

Minor alkaloid from *Cryptolepis sanguinolenta* (Asclepiadaceae). Cryst. + 1H₂O (EtOH). Mp >° 360.

Hydrochloride: [80289-16-3]

Yellow needles (MeOH). Mp >330°.

5-Me: 5,10-Dihydro-5-methyl-11H-quinolin-11-one. *Hydroxycryptolepine*.

Cryptolepinone

[160113-29-1]

[178884-07-6]

C₁₆H₁₂N₂O 248.284

From roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Active against gram-positive bacteria.

Anticomplementary agent. Yellowish amorph. solid. Artifact. Exhibits ketonol tautomerism. λ_{max} 233; 272; 311; 326; 405; 471 (MeOH) (Berdy). λ_{max} 326; 346; 447; 487 (MeOH/HCl) (Berdy). λ_{max} 232; 272; 312; 326; 367; 406; 455 (MeOH/NaOH) (Berdy).

Görlitzer, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1981, **314**, 852-861 (*synth, ir, uv, pmr*)

Crouch, R.C. *et al.*, *J. Het. Chem.*, 1995, **32**, 1077 (*pmr, cmr, N-15 nmr, struct*)

Paulo, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1485 (*Hydroxycryptolepine*)

Cimanga, K. *et al.*, *Planta Med.*, 1996, **62**, 22-27 (*activity*)

Cooper, M.M. *et al.*, *Tet. Lett.*, 1996, **37**, 4283-4286 (*Hydroxycryptolepine, synth*)

Sharaf, M.H.M. *et al.*, *J. Het. Chem.*, 1998, **35**, 1365-1369 (*Cryptolepinone, pmr, cmr, N-15 nmr*)


Martin, G.E. *et al.*, *J. Nat. Prod.*, 1998, **61**, 555-559 (*Cryptolepinone*)

Fort, D.M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1528-1530 (*Cryptolepinone, synth*)

Radl, S. *et al.*, *J. Het. Chem.*, 2000, **37**, 855-862
(*synth*, *ir*, *uv*, *pmr*)

5,6-Dihydro-11H-quinindolin-11-one, 9CI

Quinindolin-11(5H)-one, 8CI. 5H,6H-*Indolo*[2,3-b]quinolin-11-one
[13220-19-4]



C₁₅H₁₀N₂O 234.257

Alkaloid from the leaves of *Justicia betonica*. Amorph. pink powder. Mp 234° (possibly incorr.) Mp > 360°.


Kikumoto, R. *et al.*, *Tetrahedron*, 1966, **22**, 3337-3343 (*synth*, *ir*)

Bergman, J. *et al.*, *Tetrahedron*, 2003, **59**, 1033-1048 (*synth*, *ir*, *pmr*, *cmr*)

Subbaraju, G.V. *et al.*, *J. Nat. Prod.*, 2004, **67**, 461-462 (*isol*, *pmr*, *cmr*)

1,4-Dihydroquinoline

[50624-35-6]



C₉H₉N 131.177

Plates (Et₂O/petrol). Mp 46-47°. Polymerises at r.t., rel. stable in Et₂O at -15° under N₂.

N-Me: 1,4-Dihydro-1-methylquinoline

C₁₀H₁₁N 145.204

Prod. by a marine-derived *Pseudomonas aeruginosa*. Antibacterial agent (incl. MRSA). Yellow powder (EtOAc). Mp 280°. [α]_D²⁵ -46.1 (MeOH). Opt. rotation unaccounted for, struct. therefore dubious.


Birch, A.J. *et al.*, *Tet. Lett.*, 1974, 2395 (*synth*, *pmr*, *ir*)

Forrest, T.P. *et al.*, *Can. J. Chem.*, 1985, **63**, 412 (*synth*, *pmr*)

Uzair, B. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 1326-1331 (*N*-Me, *isol*, *pmr*, *cmr*)

Dihydrorutaecarpine

8,13,13b,14-Tetrahydroindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI. 13b,14-Dihydrorutaecarpine
[59863-00-2]



C₁₈H₁₅N₃O 289.336

Alkaloid from the root bark of *Zanthoxylum flavum* and the ripe fruits of *Evodia rutaecarpa* (Rutaceae). Mp

168° Mp 214° Mp 226-228°. [α]_D²⁸ -564 (c, 0.355 in DMF).

*N*¹⁴-Formyl: 14-Formyldihydrorutaecarpine. 7,8,13,13b-Tetrahydro-5-oxoindolo[2',3':3,4]pyrido[2,1-b]quinazolin-14(5H)-carboxaldehyde, 9CI
[68353-23-1]

C₁₉H₁₅N₃O₂ 317.346

Alkaloid from the ripe fruits of *Evodia rutaecarpa* (Rutaceae). Mp 285-287° (282-283°). [α]_D²⁸ +260 (c, 0.535 in DMF).


Waterman, P.G. *et al.*, *Phytochemistry*, 1976, **15**, 578 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Kamikado, T. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 1515

Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth*)

Dihydrosanguilutine

5,6-Dihydro-5-methyl-2,3,7,8,10-pentamethoxybenzo[*c*]phenanthridine
[56296-86-7]



C₂₃H₂₅NO₅ 395.454

Struct. revised in 1978. Alkaloid from the roots of *Sanguinaria canadensis* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 155-157° (154-155°).

Kim, D.K. *et al.*, *Phytochemistry*, 1975, **14**, 834 (*occur*, *uv*, *pmr*, *ms*)


Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 2686 (*synth*, *uv*)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 864 (*struct*)

Hanaoka, M. *et al.*, *J.C.S. Perkin 1*, 1987, 677 (*synth*, *uv*, *pmr*, *ms*)

Dihydrosanguinarine

13,14-Dihydro-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*i*]phenanthridine, 9CI. 5,6-Dihydro-5-methyl-2,3:7,8-bis(methylenedioxy)benzo[*c*]phenanthridine
[3606-45-9]



C₂₀H₁₅NO₄ 333.343

Several numbering systems have been used. Alkaloid from a wide variety of genera in the Papaveraceae (*Argemone*, *Bocconia*, *Chelidonium*, *Eschscholtzia*, *Glaucium*, *Hypocoum*, *Macleaya*, *Papaver*, *Romneya*), Papaveraceae (*Corydalis*, *Dicentra*), and Sapindaceae (*Pteridophyllum*). Cryst. (CHCl₃/MeOH or C₆H₆/EtOH). Mp 195-196° (188-189°, 191-193°).

8-Hydroxy: 8-Hydroxydihydrosanguinarine

C₂₀H₁₅NO₅ 349.342

Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Yellow powder. Mp 170° dec. [α]_D²⁰ -9 (c, 0.4 in MeOH).

8-Methoxy: 8-Methoxydihydrosanguinarine. *Sanguinarine pseudomethanolate*
[72401-54-8]
[49702-53-6]

C₂₁H₁₇NO₅ 363.369

Probably an artifact. Alkaloid from *Corydalis tashiroi*, *Fumaria indica* and *Hunnemannia fumariaefolia*. Mp 216-219° Mp 220-222°. [α]_D²⁵ -232.5 (c, 0.13 in CHCl₃).

8-Ethoxy: Ethoxydihydrosanguinarine. *Sanguinarine pseudoethanolate*
[28342-31-6]

C₂₂H₁₉NO₅ 377.396

Alkaloid from the roots of *Hunnemannia fumariaefolia* (Papaveraceae). Mp 207-209°. Prob. artifact.

12-Hydroxy: 12-Hydroxydihydrosanguinarine
[131984-76-4]

C₂₀H₁₅NO₅ 349.342

Alkaloid from *Eschscholtzia californica* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 208-211° dec.

12-Methoxy: Dihydrochelirubine.

Dihydrobocconine

[28342-26-9]

C₂₁H₁₇NO₅ 363.369

Minor alkaloid from the roots of *Chelidonium majus* (Papaveraceae). Needles (MeOH). Mp 206-207° (198-199°).

12-Methoxy, 5-hydroxy: 5-Hydroxydihydrochelirubine

[131984-77-5]

C₂₁H₁₇NO₆ 379.368

Alkaloid from *Eschscholtzia californica* (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 219-221° dec.

5,12-Dimethoxy: Dihydromacarpine. 5-Methoxydihydrochelirubine

[77785-12-7]

C₂₂H₁₉NO₆ 393.395

Alkaloid from cell cultures of *Eschscholtzia californica* (Papaveraceae). Needles (Et₂O). Mp 178-179°.

8-Oxo: 13-Methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*i*]phenanthridin-14(13H)-one. 5-Methyl-2,3:7,8-bis(methylenedioxy)benzo[*c*]phenanthridin-6(5H)-one. Oxosanguinarine.

Alkaloid C₇. Oxosanguinarine

[548-30-1]

C₂₀H₁₃NO₅ 347.326

Alkaloid from a wide variety of genera in the Papaveraceae (*Bocconia*, *Chelidonium*, *Dicranostigma*, *Eschscholtzia*, *Glaucium*, *Hypocoum*, *Macleaya*, *Papaver*, *Sanguinaria*), Papaveraceae (*Corydalis*, *Dicentra*, *Fumaria*), and Sapindaceae (*Pteridophyllum*). Mp 300-308° dec Mp 347-349° dec Mp 360-362° Mp 366-368°.

8-Oxo, imide: 8-Iminosanguinarine

[78118-34-0]

C₂₀H₁₄N₂O₄ 346.342

Alkaloid from the roots of *Glaucium flavum* var. *vestitum* (Papaveraceae).

Needles + 1H₂O (EtOH/CHCl₃). Mp 241-242°. Artifact.

8-Oxo, N-de-Me: N-Demethyloxysanguinarine

C₁₉H₁₁NO₅ 333.3

Alkaloid from the aerial parts of *Argemone mexicana*. Amorph. powder. λ_{max} 222 (log ε 4.02); 283 (log ε 3.36); 327 (log ε 4.13) (EtOH).

8-Oxo, 5,6-dihydro, N-de-Me: N-Demethyl-5,6-dihydrooxysanguinarine [65427-94-3]

C₁₉H₁₃NO₅ 335.315

Minor alkaloid from roots of *Chelidonium majus* (Papaveraceae). Yellow needles (CHCl₃/MeOH). Mp 304-306° (block).

8-Oxo, 12-methoxy: Oxychelirubine.

Oxybocconine

[55950-31-7]

C₂₁H₁₅NO₆ 377.353

Alkaloid from the roots of *Glauconium flavum* var. *vestitum* (Papaveraceae). Pale yellow needles (CHCl₃/Me₂CO or CHCl₃/MeOH). Mp 307-308° (295-296°).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 140 (isol)

Slavková, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1956, **21**, 211 (isol)

Dominguez, X.A. *et al.*, *Can. J. Chem.*, 1965, **43**, 679 (Oxosanguinarine)

MacLean, D.B. *et al.*, *Can. J. Chem.*, 1969, **47**, 1951 (pmr, struct)

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1435 (Dihydroxysanguinarine, Oxychelirubine, synth, pmr)

Ikuta, A. *et al.*, *Phytochemistry*, 1974, **13**, 2175; 1976, **15**, 577 (occur)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 2859 (biosynth)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 2686 (Dihydrochelirubine, N-Demethyl-5,6-dihydrooxysanguinarine)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 864 (Dihydrochelirubine, Oxychelirubine, occur, ir, pmr, struct)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1978, **41**, 145-150 (8-Methoxydihydroxysanguinarine, Ethoxydihydroxysanguinarine)

Shamma, M. *et al.*, *J.O.C.*, 1978, **43**, 2852 (Oxosanguinarine, synth, uv, ir)

Pandey, V.B. *et al.*, *Phytochemistry*, 1979, **18**, 695-696 (8-Methoxydihydroxysanguinarine, Oxosanguinarine)

Takao, N. *et al.*, *Heterocycles*, 1981, **16**, 221 (Dihydromacarpine, synth)

Castedo, L. *et al.*, *Heterocycles*, 1981, **16**, 533 (8-Iminosanguinarine, Oxychelirubine)

Berlin, J. *et al.*, *Z. Naturforsch., C*, 1983, **38**, 346 (Dihydromacarpine)

Smidrkal, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1984, **49**, 1412 (Oxosanguinarine, synth)

Hanaoka, M. *et al.*, *Chem. Lett.*, 1986, 739 (synth)

Tanahashi, T. *et al.*, *Tet. Lett.*, 1988, **29**, 5625 (synth)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 3335 (Dihydromacarpine, synth)

Tanahashi, T. *et al.*, *J. Nat. Prod.*, 1990, **53**, 579 (10-Hydroxydihydroxysanguinarine, Dihydromacarpine, 12-Hydroxydihydrochelirubine)

Williams, R.D. *et al.*, *Phytochemistry*, 1993, **32**, 719-723 (Oxosanguinarine, pmr)

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 299 (8-Hydroxydihydroxysanguinarine)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (N-15 nmr)

Chang, Y.-C. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 521-526 (N-Demethyloxysanguinarine)

Le, T.N. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 476-480 (Oxosanguinarine, synth)

Dihydroshihunine

D-511



(R)-form

C₁₂H₁₅NO₂ 205.256

(R)-form

Epidihydroshihunine

Alkaloid from the fruits of *Rhoiptelea chiliantha*. Needles + 1/4 H₂O. Mp 221-223°. [α]_D²⁵ -257.4 (c, 0.2 in CHCl₃).

6'-Methoxy: 6'-Methoxyepidihydroshihunine

C₁₃H₁₇NO₃ 235.282

Alkaloid from the fruits of *Rhoiptelea chiliantha*. Needles + 3/4 H₂O. Mp 178-180°. [α]_D¹⁵ -189.4 (c, 0.5 in CHCl₃).

(S)-form [83807-63-0]

Alkaloid from *Banisteriopsis caapi* (Malpighiaceae). Mp 187-189°. [α]_D²⁵ +234.7 (c, 0.542 in CHCl₃).


Kawanishi, K. *et al.*, *J. Nat. Prod.*, 1982, **45**, 637-639 (isol, uv, ir, pmr, cmr, ms, struct, synth)

Jiang, Z.H. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 737-740 (Epidihydroshihunine, 6'-Methoxyepidihydroshihunine)

N-(Dihydrocinapoyl)tyramine

D-512

4-Hydroxy-N-[2-(4-hydroxyphenylethyl)-3,5-dimethoxybenzenepropanamide, 9CI



C₁₉H₂₃NO₅ 345.394


Constit. of the seeds of *Annona squamosa* (sugar apple). Wax.

Yang, Y.-L. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1392-1399 (isol, pmr, cmr, ms)

5,6-Dihydro-2,3,7,8-tetra-

D-513

methoxy-5-methylbenzo[c]phenanthridine-6-acetic acid



C₂₄H₂₅NO₆ 423.465

(S)-form [473267-10-6]


Alkaloid from the roots of *Corydalis flabellata*. Amorph. powder (MeOH/C₆H₆). Mp 240-241°. [α]_D²⁰ +19.7 (c, 0.02 in MeOH). λ_{max} 227 (log ε 4.5); 280 (log ε 4.6); 323 (log ε 4.2) (MeOH).

Koul, S. *et al.*, *Planta Med.*, 2002, **68**, 262-265 (isol, pmr, cmr, ms)

5,6-Dihydro-2,3,7,8-tetra-methoxy-5-methylbenzo[c]phenanthridin-6-one

D-514

5,6-Dihydro-2,3,7,8-tetramethoxy-5-methyl-6-oxobenzo[c]phenanthridine



C₂₂H₂₁NO₅ 379.412


Alkaloid from the roots of *Corydalis flabellata*. Needles (CHCl₃/C₆H₆). Mp 190-191°. λ_{max} 235 (log ε 4.5); 288 (log ε 4.6); 335 (log ε 4.8) (MeOH).

Koul, S. *et al.*, *Planta Med.*, 2002, **68**, 262-265 (isol, pmr, cmr, ms)

6,7-Dihydro-1,3,6,6-tetra-methyl-8(5H)-isoquinolinone

D-515

[55713-38-7]



C₁₃H₁₇NO 203.283

Alkaloid from Burley tobacco (*Nicotiana tabacum*) isol. by glc. Mp 34-35°.


Demole, E. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 523 (isol, ir, pmr, ms, synth)

Swiss Pat., 1977, 585 523; CA, **86**, 16814a

3-(4,5-Dihydro-2-thiazolecarbonyl)-1H-indole

D-516

(4,5-Dihydro-2-thiazolyl)-1H-indole-3-yl-methanone, 9CI [244295-62-3]



C₁₂H₁₀N₂OS 230.29


Prod. by *Paracoccus* sp. Uv absorbent.

Japan. Pat., 1999, 99 269 175; CA, **131**, 242087z

1,2-Dihydrothiazolo[2',3':2,3]pyrimido[4,5-*b*]indole-1-carboxylic acid, 9CI

D-517

Cabbage identification factor 1. CIF 1 [238757-90-9]

C₁₃H₉N₃O₂S 271.299Alkaloid found on the leaf surfaces of *Brassica oleracea* cv. *botrytis* (cauliflower). Oviposition stimulant for the cabbage root fly, *Delia radicum*.

Glycine amide: CIF 2

[238757-91-0]

C₁₅H₁₂N₄O₃S 328.351Alkaloid found on the leaf surfaces of *Brassica oleracea* cv. *botrytis* (cauliflower). Oviposition stimulant for *Delia radicum*.

6-Methoxy: CIF 3

[338799-24-9]


C₁₄H₁₁N₃O₃S 301.325Alkaloid from the roots of *Brassica napus* var. *napobrassica* (swede). Oviposition stimulant for *Delia radicum*. Incorrect position of methoxy given in CAS.Hurter, J. *et al.*, *Phytochemistry*, 1999, **51**, 377-382 (*isol*, *pmr*, *cmr*, *ms*)De Jong, R. *et al.*, *Chemoecology*, 2000, **10**, 205-209 (CIF 3)**2,3-Dihydro-2-thioxo-4(1H)-pyrimidinone, 9CI**

D-518

2-Thiouracil. 4-Hydroxy-2-mercaptopyrimidine. Thiouracil. Deracil

[141-90-2]

[156-82-1]

C₄H₄N₂OS 128.154


6 Tautomers theoretically possible; *NH*-form (*illus.*) prob. predominates. Present in *Escherichia coli* t-RNA. Used as 0.3% soln. in Me₂CO/HCl (pH 1) for photometric detn. of Pd(II) (λ_{\max} 428 nm). Shows antibacterial and antiviral props. by alterations to protein synth. Formerly used as an antithyroid agent. Antidote for Hg poisoning. Prisms (H₂O or EtOH). Sol. Me₂CO, EtOH, H₂O. Mp 340° dec.

► Exp. carcinogen, human and exp. teratogen. LD₅₀ (mus, orl) 3900 mg/kg. Possible human carcinogen (IARC 2B). YR1575000

Still, I.W.J. *et al.*, *Can. J. Chem.*, 1978, **56**, 725 (*cmr*)Hirota, K. *et al.*, *J.O.C.*, 1978, **43**, 1193 (*synth*)
Katritzky, A.R. *et al.*, *J.C.S. Perkin 2*, 1989, 1499; 1507; 1990, 871 (*pe*, *tautom*, *bibl*)
IARC Monog. (*Web*),**1,5-Dihydro-3,4,5-trihydroxy-2H-pyrrol-2-one**

D-519

2,3-Dihydroxydihydromaleimide


C₄H₅NO₄ 131.088

(S)-form [955008-77-2]

Alkaloid from the leaves of *Arum palaestinum*. Amorph. powder.El-Desouky, S.K. *et al.*, *Arch. Pharmacol Res.*, 2007, **30**, 927-931 (*isol*, *pmr*, *cmr*, *ms*)**3,4-Dihydro-*ar*-trimethoxy-1-methylisoquinoline**

D-520


[93474-23-8]

C₁₃H₁₇NO₃ 235.282Alkaloid detected in *Pachycereus weberi* (Cactaceae) by tandem mass spectrom.Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109-114 (*occur*)**1,4-Dihydro-4,6,7-trimethyl-9H-imidazo[1,2-*a*]purin-9-one, 9CI**


D-521

7-Methylwybe

[96881-39-9]


C₁₀H₁₁N₅O 217.23Found in tRNA of highly thermophilic archaeobacteria. Monohydrate. Mp 300°. Milloskey, J.A. *et al.*, *Nucleic Acids Res.*, 1987, **15**, 683Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 284; 1990, **38**, 2656 (*synth*, *pmr*)**2,7-Dihydro-2,5,7-trimethyl-5H-imidazo[4,5-*e*]-1,2,4-triazine-3,6-dione**

D-522

Antibiotic 2096D. 2096DC₇H₉N₅O₂ 195.18Prod. by *Streptomyces* sp. IM 2096. Pale yellow solid. Possible degradation prod. Wang, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1641-1646**3,9-Dihydro-1,3,9-trimethyl-8-nitroso-1H-purine-2,6-dione, 9CI**


D-523

1,3,9-Trimethyl-8-nitrosoisoxanthine [153534-80-6]


C₈H₉N₅O₃ 223.191The first isoxanthine (isocaffeine) deriv. from a natural source. Isol. from the sea cucumber *Cucumaria frondosa*. Powder. Mp 273° dec.Yayli, N. *et al.*, *J. Nat. Prod.*, 1994, **57**, 84-89 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)**5,8-Dihydro-1,8,8-trimethyl-1H-pyrano[3,4-*b*]pyridine-4,6-dione, 9CI**

D-524

[911366-95-5]


C₁₁H₁₃NO₃ 207.229Alkaloid from the aerial parts of *Stapelia hirsuta*. Needles. Mp 178-180°. λ_{\max} 234; 322 (MeOH).Shabana, M. *et al.*, *Turk. J. Chem.*, 2006, **30**, 89-92 (*isol*, *pmr*, *cmr*, *ms*)**5,6-Dihydro-3,6,6-trimethyl-7H-2-pyridin-7-one, 9CI**

D-525

5,6-Dihydro-3,6,6-trimethyl-7H-cyclopenta[*c*]pyridin-7-one, 9CI [55713-43-4]C₁₁H₁₃NO 175.23Alkaloid from Virginia tobacco (*Nicotiana tabacum*).Wahlberg, I. *et al.*, *Phytochemistry*, 1977, **16**, 1233-1235 (*isol*)**2,5-Dihydro-2,4,5-trimethylthiazole, 9CI**

D-526

2,4,5-Trimethyl-3-thiazoline, 8CI [60633-24-1]

(2*R*,5*R*)-formC₆H₁₁NS 129.226

Constit. of the faeces of red fox (*Vulpes vulpes*). Flavour constit. of cooked meats. Bp₁₂ 57°.

Picrate:

Cryst. (MeOH). Mp 117-118°.

[134281-37-1, 134281-36-0]

Thiel, M. *et al.*, *Annalen*, 1960, **638**, 174

(*synth*)

Mussinan, C.J. *et al.*, *ACS Symp. Ser.*, 1976, **26**, 133 (*occur*)

Hwang, S.S. *et al.*, *J. Agric. Food Chem.*, 1986, **34**, 538 (*synth, occur*)


Sullivan, T.P. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 363; 379 (*isol*)

Dihydroxanthommatin D-527

α -Amino-3-carboxy-1,5-dihydroxy- γ -oxo-12H-pyrido[3,2-a]phenoxazine-11-butanoic acid, 9CI

[25705-16-2]

[81624-03-5]



C₂₀H₁₅N₃O₈ 425.354

(S)-form

Found in eyes and skin of *Loligo vulgaris*, *Sepia officinalis* and other molluscs, also in heads of flies *Musca domestica* and bees *Apis mellifera*, and in lepidopteran wings and excreta. Red cryst. Mp >350°.

5-O-Sulfate: Ommatin D

[28991-26-6]

C₂₀H₁₅N₃O₁₁S 505.418

Occurs in lepidopteran wings and excreta. Isol. from the secretions of the butterfly *Vanessa urticae*. Red-brown powder.

5-O- β -D-Glucopyranoside: Rhodommatin

C₂₆H₂₅N₃O₁₃ 587.496

Occurs in lepidopteran wings and excreta. Isol. from the secretions of *Vanessa urticae*. Red cryst. Mp >350°.

[121727-27-3]

Butenandt, A. *et al.*, *Annalen*, 1957, **607**, 207-214 (*Ommatin D*)

Butenandt, A. *et al.*, *Hoppe-Seyler's Z.*

Physiol. Chem., 1960, **321**, 258-275


(*Ommatin D, Rhodommatin*)

Bolognese, A. *et al.*, *J. Het. Chem.*, 1988, **25**, 1247-1250

1,3-Dihydroxyacridone D-528

1,3-Dihydroxy-9(10H)-acridinone, 9CI

[20324-10-1]



C₁₃H₉NO₃ 227.219

Yellow needles (Me₂CO/EtOH). Mp 345° dec. (355° dec., 370°).

3-Ac: [28333-05-3]

C₁₅H₁₁NO₄ 269.256

Yellow needles (Me₂CO/EtOAc). Mp 286-287°.

1,3-Di-Ac: [28333-04-2]

C₁₇H₁₃NO₅ 311.293

Buff prisms (Me₂CO/EtOAc). Mp 205-207°.

N-Me: 1,3-Dihydroxy-10-methylacridone

[28333-02-0]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the roots of *Citrus tamurana* and *Citrus natsudaikai* (Rutaceae). Light yellow cubes (CH₂Cl₂). Mp 292.5-295° dec. (>295°).

1-Me ether, N-Me: 3-Hydroxy-1-methoxy-10-methylacridone. Tegerrardin A

C₁₅H₁₃NO₃ 255.273

Alkaloid from the stem bark of *Teclea gerrardii*. Pale yellow powder. Mp 158-159°.

3-Me ether: 1-Hydroxy-3-methoxyacridone

[91998-87-7]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the bark of *Fagara macrophylla*. Mp 300-301° (synthetic).

3-Me ether, N-Me: 1-Hydroxy-3-methoxy-10-methylacridone

[13161-83-6]

C₁₅H₁₃NO₃ 255.273

Alkaloid from the bark of *Esenbeckia littoralis*, the roots of *Glycosmis mauritiana* and *Ruta graveolens* (rue), and the root and stem bark of *Fagara lepreurii* and *Fagara rubescens* (Rutaceae). Yellow cryst. (EtOAc/petrol or EtOAc/C₆H₆). Mp 164-165° Mp 174-176°.

3-Me ether, N-Me, 1-Ac: 1-Acetoxy-3-methoxy-10-methylacridone

[75533-56-1]

C₁₇H₁₅NO₄ 297.31

Alkaloid from the bark of *Esenbeckia littoralis*. Cryst. (EtOAc/petrol). Mp 219-220°.

Di-Me ether: 1,3-Dimethoxyacridone

[27221-40-5]

C₁₅H₁₃NO₃ 255.273

Alkaloid from leaves of *Bauerella simplicifolia* (preferred genus name *Acronychia*) (Rutaceae). Cryst. (EtOAc), green cryst. (EtOH). Mp 256-257° Mp 286-287°.

Di-Me ether, N-Me: 1,3-Dimethoxy-10-methylacridone

[13082-10-5]

C₁₆H₁₅NO₃ 269.299

Alkaloid from *Bauerella simplicifolia* (preferred genus name *Acronychia*), *Diphasia angolensis*, *Oricia suaveolens*, *Teclea verdoorniana*, *Vepris pilosa*, *Vepris ampody* and *Acronychia baueri* (Rutaceae). Needles (EtOAc, CHCl₃/petrol or C₆H₆/cyclohexane). Mp 165°. λ_{\max} 261 (log ϵ 4.66); 267 (log ϵ 4.67); 290 (log ϵ 4.15); 380 (log ϵ 3.94) (EtOH).

Di-Me ether, N-Me, picrate:

Yellow needles (MeOH). Mp 203-205°.

O³-(3-Methyl-2-butenyl), N-Me: 3-Hydroxy-10-methyl-1-prenyloxyacridone.

Tegerrardin B

C₁₉H₁₉NO₃ 309.364

Alkaloid from the stem bark of *Teclea gerrardii*. Pale yellow gum.

O³-(3-Methyl-2-butenyl), N-Me: 1-Hydroxy-10-methyl-3-prenyloxyacridone.

Vebilicine

[177857-30-6]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the leaves of *Vepris bilocularis*. Yellow cryst. (Et₂O/petrol). Mp 145-147°.

Lamberton, J.A. *et al.*, *Aust. J. Chem.*, 1953, **6**, 66-77 (*di-Me ether N-Me*)

Bowie, J.H. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1179-1193 (*ms*)

Hlubucek, J. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1881-1889 (*N-Me, 1,3-di-Me N-Me*)

Oh, C.S. *et al.*, *J. Het. Chem.*, 1970, **7**, 261-267 (*N-Me*)

Kan-Fan, C. *et al.*, *Phytochemistry*, 1970, **9**, 1283-1291 (*1,3-di-Me ether N-Me, isol*)

Fish, F. *et al.*, *Phytochemistry*, 1971, **10**, 3322-3324 (*3-Me ether N-Me, isol, uv, ir*)

Scharlemann, W. *et al.*, *Z. Naturforsch., B.*, 1972, **27**, 806-808 (*3-Me ether N-Me, isol, uv*)

Bandaranayake, W.M. *et al.*, *J.C.S. Perkin 1*, 1974, 998-1007 (*synth*)

Adams, J.H. *et al.*, *J.C.S. Perkin 1*, 1976, 2089-2093 (*di-Me ether, synth, uv*)

Fish, F. *et al.*, *Planta Med.*, 1978, **33**, 228-231 (*di-Me ether N-Me, isol*)

Begenthal, D. *et al.*, *Phytochemistry*, 1979, **18**, 161-163 (*cmr*)

Tillequin, F. *et al.*, *J. Nat. Prod.*, 1980, **43**, 498-502 (*1,3-di-Me ether, 1,3-di-Me ether N-Me*)

Dreyer, D.L. *et al.*, *Phytochemistry*, 1980, **19**, 941-944 (*3-Me ether N-Me, 3-Me ether N-Me 1-Ac, di-Me ether N-Me, isol, uv, pmr, ms, synth*)

Ju-ichi, M. *et al.*, *Heterocycles*, 1987, **26**, 1873 (*N-Me, isol, uv, ir, pmr, cmr, ms, struct, deriv*)

Bahar, M.H. *et al.*, *Indian J. Chem., Sect. B.*, 1987, **26**, 782-783 (*3-Me ether N-Me, synth, uv, ir, pmr, ms*)

Reisch, J. *et al.*, *Annalen*, 1991, 685-689 (*N-Me, 3-Me ether N-Me, 1,3-di-Me ether N-Me, synth, ir, pmr, cmr*)

Brader, G. *et al.*, *Phytochemistry*, 1996, **42**, 881-884 (*Vebilicine*)

Spatafora, C. *et al.*, *Phytochem. Anal.*, 1997, **8**, 139-142 (*1-Hydroxy-3-methoxyacridone*)

Waffo, A.F.K. *et al.*, *Phytochemistry*, 2007, **68**, 663-667 (*Tegerrardin A, B*)

1,7-Dihydroxyacridone D-529

1,7-Dihydroxy-9(10H)-acridinone, 9CI

[112649-95-3]

C₁₃H₉NO₃ 227.219

Alkaloid from the aerial parts of *Boronia lanceolata*. Amorph. yellow solid. Mp 278-285°. λ_{\max} 240 (sh); 259; 267; 308; 323; 414 (MeOH).

N-Me: 1,7-Dihydroxy-10-methylacridone

[65582-55-0]

C₁₄H₁₁NO₃ 241.246

Constit. of roots of *Boenninghausenia albiflora* (Rutaceae). Yellow needles (Me₂CO). Mp 308° dec. λ_{\max} 244 (sh); 262; 272; 314 (sh); 328 (sh); 410 (MeOH).


Rózsa, Z. *et al.*, *Phytochemistry*, 1978, **17**, 169-170 (*N-Me, uv, pmr, ms*)

Begenthal, D. *et al.*, *Phytochemistry*, 1979, **18**, 161-163 (*N-Me, cmr*)

Ahsan, M. *et al.*, *Phytochemistry*, 1993, **33**, 1507-1510 (*isol, pmr, cmr, ms*)

1,8-Dihydroxyacridone **D-530**

1,8-Dihydroxy-9(10H)-acridinone, 9CI
[151077-55-3]



C₁₃H₉NO₃ 227.219


Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae), also from sp. aff. *Samadera bidwillii*. Yellow needles (CHCl₃/MeOH). Mp 250°. λ_{max} 251; 258 (sh); 310 (sh); 322; 408; 426 (MeOH).

N-Me: 1,8-Dihydroxy-10-methylacridone
[130897-51-7]

C₁₄H₁₁NO₃ 241.246

Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae). Yellow needles (CHCl₃/MeOH). Mp 235-237°. λ_{max} 251; 262 (sh); 320; 413 (MeOH).

Ahsan, M. et al., *Phytochemistry*, 1993, **33**, 1507 (isol, uv, ir, pmr, cmr, ms, struct)
Gibbons, S. et al., *Phytochemistry*, 1997, **44**, 1109 (isol)

1,2-Dihydroxyaporphine**D-531***(R)-form*

C₁₇H₁₇NO₂ 267.327

(R)-form

O¹-Me: 2-Hydroxy-1-methoxyaporphine. Floribundine. N-Methylasimilobine. O-Nornuciferine

[3153-55-7]

C₁₈H₁₉NO₂ 281.354

Alkaloid from *Nelumbo nucifera* (East India lotus) (Nelumbonaceae) and from several genera in the Annonaceae (*Asimina*, *Xylopia*), Menispermaceae (*Stephania*), Papaveraceae (*Papaver*) and Rhamnaceae (*Colubrina*). Active against gram-positive bacteria. Mp 195-196°. [α]_D²⁰ -220 (c, 0.40 in CHCl₃). λ_{max} 231 (log ε 4.07); 272 (log ε 4); 314 (log ε 3.29) (no solvent reported).

O¹-Me, O-β-D-glucopyranoside: N-Methylasimilobine glucoside

[267886-73-7]

C₂₄H₂₉NO₇ 443.496

Alkaloid from the seeds of *Stephania cepharantha*. Amorph. powder. [α]_D²⁷ -116.1 (c, 0.6 in MeOH).

O¹-Me, O-α-L-rhamnopyranoside: Floripavidine

[62867-48-5]

C₂₄H₂₉NO₆ 427.496

Alkaloid from *Papaver floribundum* (*Papaver fugax*), *Papaver cylindricum*, *Papaver armeniacum* and *Papaver persicum* (*Papaver tauricola*) (Papavera-

ceae). Mp 241-242°. [α]_D -156 (c, 1.6 in MeOH). λ_{max} 273 (log ε 4.25); 310 (log ε 3.45) (no solvent reported).

O¹-Me, N-de-Me: 2-Hydroxy-1-methoxynoraporphine. Asimilobine†

[6871-21-2]

C₁₇H₁₇NO₂ 267.327

Alkaloid from a wide variety of genera in the Annonaceae (*Anaxagorea*, *Annona*, *Asimina*, *Melodorum*, *Desmos*, *Guatteria*, *Hexalobus*, *Mitrella*, *Monanthotaxis*, *Popowia*, *Polyalthia*, *Scheffermitra*, *Xylopia*, *Uvaria*), Aristolochiaceae (*Aristolochia*), Magnoliaceae (*Liriodendron*, *Magnolia*), Monimiaceae (*Laurelia*), Nelumbonaceae (*Nelumbo*), Lauraceae (*Ocotea*) and Rhamnaceae (*Zizyphus*). Serotonin receptor antagonist. Shows antimicrobial activity. Mp 177-179°. [α]_D¹⁴ -213 (c, 0.64 in CHCl₃). λ_{max} 274 (log ε 4.21); 308 (log ε 3.51) (no solvent reported).

O¹-Me, N-de-Me, O-β-D-glucopyranoside: Asimilobine 2-glucoside

[151601-87-5]

C₂₃H₂₇NO₇ 429.469

Alkaloid from tubers of *Stephania pierrei* (Menispermaceae). Mp 158°. [α]_D²⁰ -107 (c, 0.1 in MeOH).

O¹-Me, N-de-Me, N-Ac: N-Acetylasimilobine

[60031-90-5]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the heartwood and discoloured sapwood of *Liriodendron tulipifera* (Magnoliaceae). Needles (CHCl₃). Mp 281-283°. [α]_D²⁵ -405 (c, 0.595 in Py).

O¹-Me, N-de-Me, N-carbamoyl: N-Carbamoylasimilobine

[83459-47-6]

C₁₈H₁₈N₂O₃ 310.352

Alkaloid from the stem bark and root bark of *Hexalobus crispiflorus* (Annonaceae). Cryst. (MeOH). Mp 218° dec. [α]_D²⁰ -373 (c, 0.06 in MeOH). λ_{max} 273 (log ε 4.08); 305 (log ε 3.43) (MeOH).

O¹-Me, N-de-Me, N-methoxycarbonyl: Romucosine A

[356048-23-2]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Rollinia mucosa* (biri-ba). Amorph. brown powder. [α]_D²⁵ -105 (c, 0.1 in CHCl₃). λ_{max} 237 (log ε 4.4); 285 (log ε 4.22); 307 (log ε 4.05) (MeOH).

O¹-Me, N-de-Me, N-ethoxycarbonyl, O-glucopyranoside: Kamaline

[157207-94-8]

C₂₆H₃₁NO₉ 501.532

Alkaloid from whole plants of *Stephania venosa* (Menispermaceae). Cryst. (MeOH/CHCl₃). Mp 158-160°. [α]_D²⁰ -285 (c, 0.107 in CHCl₃).

O²-Me: [37082-15-8]

Synthetic. Mp 167-169°.

O²-Me, hydrochloride:

Cryst. + ½ H₂O. Mp 220° dec.

O²-Me, N-de-Me: 1-Hydroxy-2-methoxynoraporphine. (-)-Caaverine

[6899-64-5]

C₁₇H₁₇NO₂ 267.327

Alkaloid from *Symplocos celastriana* and *Ocotea glaziovii* (Symplocaceae, Lauraceae). Mp 208-210° dec. [α]_D²⁵ -89 (c, 1 in MeOH). λ_{max} 272 (log ε 4.25); 310 (sh) (log ε 3.7) (no solvent reported).

▶RB6009000

Di-Me ether: 1,2-Dimethoxyaporphine. Nuciferine. Sanjoinine E

[475-83-2]

C₁₉H₂₁NO₂ 295.38

Alkaloid from *Nelumbo lutea* (Nelumbonaceae) and *Colubrina faralaotra* (Rhamnaceae), also from the Araceae, Berberidaceae, Lauraceae, Menispermaceae, Papaveraceae, Magnoliaceae and Annonaceae. Inhibitor of adenylate cyclase. Shows strong sedative activity. Antispasmodic agent. Active against gram-positive bacteria. Mp 165.5°. [α]_D²⁰ -164 (c, 1 in EtOH). λ_{max} 230 (ε 18300); 272 (ε 15400); 310 (ε 2100) (95% EtOH).

▶LD₅₀ (rat, orl) 280 mg/kg. CE0350000

Di-Me ether, hydrochloride: Mp 241° (in vacuo).

Di-Me ether, N-Me:

C₂₀H₂₄NO₂⁺ 310.415

Mp 177-178° (as iodide).

Di-Me ether, N-de-Me: 1,2-Dimethoxynoraporphine. N-Nornuciferine. N-Desmethylnuciferine. N-De-

methylnuciferine. N-Desmethylnuciferine. Sanjoinine Ia.

Daechualkaloid E

[4846-19-9]

C₁₈H₁₉NO₂ 281.354

Alkaloid from *Nelumbo lutea* (Nelumbonaceae) and *Colubrina faralaotra* (Rhamnaceae), also from the Annonaceae, Magnoliaceae, Menispermaceae and Monimiaceae. Antispasmodic agent, serotonin receptor antagonist. Mp 128-129°. [α]_D²⁵ -145 (c, 0.98 in EtOH). λ_{max} 230 (ε 20500); 272 (ε 16800); 310 (ε 2200) (95% EtOH).

Di-Me ether, N-de-Me, N-formyl: N-Formylnornuciferine

[107633-68-1]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae). Negative opt. rotn. λ_{max} 212 (log ε 4.48); 226 (sh) (log ε 4.32); 269 (log ε 4.26); 312 (sh) (log ε 3.74) (EtOH).

Di-Me ether, N-de-Me, N-Ac: N-Acetylnornuciferine

[1942-03-6]

C₂₀H₂₁NO₃ 323.391

Alkaloid from the heartwood of *Liriodendron tulipifera* (Magnoliaceae). Cryst. (MeOH). Mp 229-230°. [α]_D²⁵ -406 (c, 0.65 in CHCl₃).

6a,7-Didehydro, di-Me ether: see Dehydronuciferine, D-164

(S)-form

O²-Me: 1-Hydroxy-2-methoxyaporphine. Lirididine. N-Methylcaaverine

[54383-28-7]

C₁₈H₁₉NO₂ 281.354

Alkaloid from *Liriodendron tulipifera*, *Papaver persicum*, *Papaver tauricola*

and *Papaver armeniacum* (Magnoliaceae, Papaveraceae). Mp 214-215°. λ_{\max} 271 (log ϵ 4.12); 312 (log ϵ 3.67) (no solvent reported).

O²-Me, N-de-Me: (+)-Caaverine

[54383-30-1]
C₁₇H₁₇NO₂ 267.327
Alkaloid from *Liriodendron tulipifera* (Magnoliaceae). $[\alpha]_{\text{D}}^{25}$ +95 (c, 0.12 in MeOH).

Di-Me ether, N-de-Me: [20454-22-2]

Alkaloid from *Liriodendron tulipifera*. $[\alpha]_{\text{D}}^{20}$ +140 (c, 0.18 in EtOH).

Di-Me ether, N-de-Me, N-methoxycarbonyl: N-Methoxycarbonyl-N-nornuciferine

[356042-11-0]
C₂₀H₂₁NO₄ 339.39
Alkaloid from *Rollinia mucosa* (biri-ba). Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +165 (c, 0.05 in CHCl₃). λ_{\max} 235 (log ϵ 4.23); 285 (log ϵ 4.2); 307 (log ϵ 3.36) (MeOH).

3-Chloro, O¹-Me, N-de-Me, N-methoxycarbonyl: Romucosine B

[356048-24-3]
C₁₉H₁₈ClNO₄ 359.808
Alkaloid from *Rollinia mucosa* (biri-ba). Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +153 (c, 0.05 in CHCl₃). Opposite abs. config. to Romucosine A. λ_{\max} 245 (log ϵ 4.43); 285 (log ϵ 4.25) (MeOH).

(±)-form

O²-Me: [22222-86-2]

Alkaloid from the leaves of *Ocotea glaziovii* (Lauraceae). Cryst. (Et₂O/hexane). Mp 156°.

► **CE1040000**

O²-Me, N-Me:

Prisms (MeOH/Et₂O). Mp 234-236° dec.

Di-Me ether, hydrochloride: [5868-18-8]
Mp 258° dec.

► **CE0340000**

Di-Me ether, N-de-Me: [54750-04-8]
Synthetic. Mp 124-125°.

► **RB5916300**

O¹-Me, N-de-Me, hydrochloride: Mp 260-262°.

Di-Me ether, N-de-Me, N-Ac: [29424-85-9]
Synthetic. Rosettes (C₆H₆/hexane), needles (MeOH). Mp 232-233°.

Gulland, J.M. *et al.*, *J.C.S.*, 1928, 581-591 (*Nuciferine, synth*)

Konvalova, R. *et al.*, *Ber.*, 1935, **68**, 2277-2282 (*Floribundine, isol*)

Arthur, H.R. *et al.*, *J.C.S.*, 1959, 2306 (*Nuciferine, isol*)

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 469-473; 1965, **85**, 77-82 (*Floribundine, Asimilobine*)

Kupchan, S.M. *et al.*, *Tetrahedron*, 1963, **19**, 227-232 (*N-Nornuciferine*)

Gilbert, B. *et al.*, *J.A.C.S.*, 1964, **86**, 694-696 (*Liridinine, Caaverine, synth, uv, pmr, ord*)

Tschesche, R. *et al.*, *Tetrahedron*, 1964, **20**, 1435-1447 (*(-)-Caaverine*)

Kunimoto, J.I. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1141-1142 (*Floribundine, Nuciferine, isol*)

Baxter, I. *et al.*, *J.C.S.*, 1965, 4014-4024 (*N-Acetylnornuciferine*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 363-368; 423-426 (*Asimilobine, N-Nornuciferine*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1971, **36**, 2413-2418 (*Nuciferine, synth, uv*)

Guinaudeau, H. *et al.*, *Phytochemistry*, 1971, **10**, 1963-1966 (*Nuciferine, isol, ir, pmr, ms*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 157-176 (*activity*)

Ziyaev, R. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 760-763; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 727-729 (*(+)-Caaverine, Liridinine, isol, uv, ir, pmr, ms*)

Cassagrande, C. *et al.*, *Farmaco, Ed. Sci.*, 1975, **30**, 479-480; *C.A.*, **83**, 93873p (*Caaverine, (±)-Liridinine*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1975, **38**, 276-338 (*rev, Asimilobine*)

Guinaudeau, H. *et al.*, *Planta Med.*, 1975, **27**, 304-318 (*Floribundine*)

Israelov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 799-801; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 716-717 (*Floripavidine*)

Chen, C.L. *et al.*, *Phytochemistry*, 1976, **15**, 1161-1167 (*N-Acetylasimilobine*)

Hufford, C.D. *et al.*, *Phytochemistry*, 1976, **15**, 1169-1171 (*N-Acetylasimilobine, N-Acetylnornuciferine*)

Jackman, L.M. *et al.*, *J. Nat. Prod.*, 1979, **42**, 437-449 (*cmr*)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80-85 (*cd*)

Phillipson, J.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 296-307 (*Floripavidine*)

Phillipson, J.D. *et al.*, *Planta Med.*, 1981, **41**, 105-118 (*Liridinine, isol, uv, ms*)

Achenbach, H. *et al.*, *Annalen*, 1982, 1623-1633 (*N-Carbamoylasimilobine*)

Cortes, D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 878-884 (*N-Formylnornuciferine*)

Villar, A. *et al.*, *Planta Med.*, 1986, **57**, 556-557 (*activity*)

Simeon, S. *et al.*, *Pharmazie*, 1989, **44**, 593-597 (*Asimilobine, N-Methylasimilobine, activity*)

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443 (*Nuciferine, N-Nornuciferine, isol*)

Ramana, M.M.V. *et al.*, *Tet. Lett.*, 1996, **37**, 1671-1674 (*N-Nornuciferine, synth*)

Hasrat, J.A. *et al.*, *Phytomedicine*, 1997, **4**, 133-140 (*activity, Nornuciferine, Asimilobine*)

Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 2000, **63**, 477-479 (*N-Methylasimilobine glucoside*)


Kuo, R.-Y. *et al.*, *Phytochemistry*, 2001, **57**, 421-425 (*Romucosines, N-Methoxycarbonylnornuciferine*)

Lafrance, M. *et al.*, *Eur. J. Org. Chem.*, 2007, 811-825 (*Nuciferine, synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NOE500

2-(2,5-Dihydroxybenzamido)benzoic acid

D-532



C₁₄H₁₁NO₅ 273.245

Amide, 5'-O-β-D-allopyranoside: [853234-82-9]

C₂₀H₂₂N₂O₉ 434.402


Alkaloid from *Rabdosia rubescens*.

Liu, J. *et al.*, *CA*, 2005, **143**, 40889z (*Rabdosia rubescens amide*)

4,5-Dihydroxy-1,2-benzenedicarboxylic acid, 9CI

D-533

4,5-Dihydroxyphthalic acid, 8CI. Catechol-4,5-dicarboxylic acid. Normetahepimic acid
[63958-66-7]



C₈H₆O₆ 198.132

Degradn. prod. of humic acid. Prisms (H₂O). Mp 175° (→ anhydride).

Di-Me ester: [66323-03-3]

C₁₀H₁₀O₆ 226.185
Needles (petrol/EtOAc). Mp 141.5-142.5°.

Di-Et ester:

C₁₂H₁₄O₆ 254.239
Needles (H₂O). Mp 152°.

Dinitrile: 4,5-Dicyano-1,2-benzenediol
[300853-66-1]

C₈H₄N₂O₂ 160.132
Needles (H₂O). Mp 285°.

Mono-Me ether, imide, N-Me: 6-Hydroxy-5-methoxy-2-methyl-1H-isoindole-1,3(2H)-dione. 6-Hydroxy-5-methoxy-N-methylphthalimide

C₁₀H₉NO₄ 207.185

Alkaloid from the roots of *Menispermum dauricum*. Light yellow needles (CHCl₃). Mp 213-215°. λ_{\max} 216 (log ϵ 3.04); 240 (log ϵ 3.49); 298 (log ϵ 0.62); 349 (log ϵ 0.54) (no solvent reported).

Di-Me ether: 4,5-Dimethoxy-1,2-benzenedicarboxylic acid. 4,5-Dimethoxyphthalic acid. Veratrole-4,5-dicarboxylic acid. Metahepimic acid. m-Hepimic acid

[577-68-4]
C₁₀H₁₀O₆ 226.185

Degradn. prod. of many alkaloids. Isol. from poppy straw (*Papaver somniferum*). Needles (H₂O). Mp 174-175°.

Di-Me ether, di-Me ester: [17078-61-4]

C₁₂H₁₄O₆ 254.239
Constit. of soil humus. Mp 88-89°.

Di-Me ether, anhydride: 5,6-Dimethoxy-1,3-isobenzofurandione, 9CI. m-Hepimic anhydride

[4821-94-7]
C₁₀H₈O₅ 208.17

Cryst. (diisopropylether). Mp 175-176°.

Di-Me ether, imide: 5,6-Dimethoxy-1H-isoindole-1,3(2H)-dione, 9CI. m-Hepimide

[4764-20-9]
C₁₀H₉NO₄ 207.185

Needles (C₆H₆). Mp 317-320°.

Di-Me ether, imide, N-Me: 5,6-Dimethoxy-2-methyl-1H-isoindole-1,3(2H)-dione. 5,6-Dimethoxy-N-methylphthalimide

[92288-92-1]
C₁₁H₁₁NO₄ 221.212

Isol. from the trunk bark of *Hernandia nymphaeifolia*. Prisms (CHCl₃/MeOH). Mp 149-151°.

Dibenzyl ether, dinitrile: [206995-45-1]

C₂₂H₁₆N₂O₂ 340.381

Needles (C₆H₆). Mp 183.5-184.5°.

Di-Ph ether, dinitrile: 4,5-Diphenoxy-1,2-dicyanobenzene

[147699-63-6]

C₂₀H₁₂N₂O₂ 312.327

Cryst. (MeOH). Mp 148-149°.

Bruck, O. *et al.*, *Ber.*, 1901, **34**, 2741-2747 (*synth*)

Mason, F.A. *et al.*, *J.C.S.*, 1914, **105**, 2013-2024 (*di-Me ether imide*)

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1945, **28**, 722-740 (*isol, di-Me ether*)

Ogner, G. *et al.*, *Can. J. Chem.*, 1971, **49**, 1053-1063 (*di-Me ether di-Me ester, ir, ms*)

Anderson, D.R. *et al.*, *J.O.C.*, 1978, **43**, 2726-2730 (*di-Me ester, di-Me ether imide, synth, ir, pmr*)

Chen, J.J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (*5,6-Dimethoxy-N-methylphthalimide*)

Hu, M. *et al.*, *J. Med. Chem.*, 1998, **41**, 1789-1802 (*dibenzyl ether dinitrile, synth, ir, pmr, ms*)

Cabezon, B. *et al.*, *Eur. J. Org. Chem.*, 2000, 2767-2775 (*dinitrile*)


Drager, A.S. *et al.*, *J.O.C.*, 2000, **65**, 2257-2260 (*di-Me ester, synth, pmr, cmr, ms*)

Plater, M.J. *et al.*, *J.C.S. Perkin 1*, 2002, 91-96 (*di-Ph ether dinitrile, synth, uv, ir, pmr, cmr*)

Zhang, X. *et al.*, *Phytochemistry*, 2004, **65**, 929-932 (*6-Hydroxy-5-methoxy-N-methylphthalimide*)

5,7-Dihydroxy-2H-benz[b]-isoindeole-4,9-dione, 9CI

D-534



C₁₂H₇NO₄ 229.192

7-Me ether: 5-Hydroxy-7-methoxy-2H-benz[b]isoindeole-4,9-dione, 9CI. *Azamonosporascone*

C₁₃H₉NO₄ 243.218

Prod. by *Monosporascus cannonballus*. Yellow cryst. Mp 279-283°. λ_{max} 223 (log ε 3.92); 250 (log ε 3.84); 272 (log ε 3.85); 396 (log ε 3.94) (MeOH).


Stipanovic, R.D. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 4109-4112; 4113-4120 (*isol, pmr, cmr, ms*)

2,4-Dihydroxybenzoic acid, 9CI

D-535

β-Resorcylic acid, 8CI. *Resorcinol-4-carboxylic acid*. FEMA 3798

[89-86-1]



C₇H₆O₄ 154.122

Constit. of *Pterocarpus santalinus* and *Adenanthera pavonina*. Gives colour reaction with Fe; fluorescence reaction with B. Monomer for phenolic resins used as ion exchangers. Taste modifier in foods. Solid (CHCl₃). Mp 152° subl Mp 175° Mp 218-219° (213° rapid heat). pK_{a1}

3.11; pK_{a2} 8.55; pK_{a3} 14 (25°).

► Exp. teratogenic effects. VH3708050

Nitrile: 2,4-Dihydroxybenzonitrile. 4-Cyano-1,3-benzenediol. 4-Cyanoresorcinol [64419-24-5]

C₇H₅NO₂ 135.122

Mp 175°.

Nitrile, 2-O-β-D-glucopyranoside: *Amiroside*

[258277-63-3]

C₁₃H₁₅NO₇ 297.264

Constit. of *Veronica polita*.

Zaghloul, M.G. *et al.*, *CA*, 2000, **132**, 148988t (*Amiroside*)

3,4-Dihydroxybenzoic acid D-536

Protocatechuic acid, 8CI. *Carbohydroquinonic acid*. *Catechol-4-carboxylic acid*.

Hypogallic acid

[99-50-3]

C₇H₆O₄ 154.122

Can be extracted from lignin. Isol. in free state from various higher plants, e.g. *Fagopyrum* and *Alnus* spp. and some *Allium* spp. Antioxidant, free radical scavenger, dietary chemopreventive agent (inhibits development of neoplasms in animal models), inhibits LDL oxidation, platelet aggregation inhibitor. Shows cytostatic activity. Needles + 1H₂O (H₂O). Mp 195-196° (anhyd.). pK_{a1} 4.49; pK_{a2} 8.64; pK_{a3} 1 (25°).

► UL0560000

3-Me ether, amide: 4-Hydroxy-3-methoxybenzamide

[19072-58-3]

[179951-61-2 (hydrochloride)]


C₈H₉NO₃ 167.164

Constit. of *Naravelia zeylanica*. Cryst. (EtOH aq.). Mp 179-182° (153-154°).

Jaroszewski, J.W. *et al.*, *Nat. Prod. Rep.*, 2005, **19**, 291-294 (*4-Hydroxy-3-methoxybenzamide*)

2,5-Dihydroxy-2H-1,4-benzoxazin-3(4H)-one

D-537



C₈H₇NO₄ 181.148

(R)-form

2-O-β-D-Glucopyranoside: [381722-83-4]

C₁₄H₁₇NO₉ 343.29

Constit. of *Acanthus ilicifolius*.


Amorph. powder. [α]_D²⁶ +95 (c, 0.4 in DMSO).

Kanchanapoom, T. *et al.*, *Phytochemistry*, 2001, **58**, 637-640 (*isol, pmr, cmr*)

2,7-Dihydroxy-2H-1,4-benzoxazin-3(4H)-one, 9CI

D-538

[69804-59-7]



C₈H₇NO₄ 181.148

(R)-form

Identified in sweet corn extracts. Sol. MeOH, H₂O, CHCl₃, λ_{max} 254 ; 281 (MeOH) (Berdy). λ_{max} 253 ; 279 (H₂O) (Berdy).

2-O-β-D-Glucopyranoside: *DHBOA-Glc* [28512-70-1]

C₁₄H₁₇NO₉ 343.29

Constit. of *Acanthus ilicifolius*, *Coix lachryma-jobi* (Job's tears) and *Lamium galeobdolon*. Cryst. (MeOH). Mp 267-269°. [α]_D²⁵ +22.3 (c, 0.6 in Py). λ_{max} 263 (log ε 4); 286 (sh) (log ε 3.84) (EtOH).

7-Me ether: 2-Hydroxy-7-methoxy-2H-1,4-benzoxazin-3(4H)-one. *HMBOA* [17359-53-4]

C₉H₉NO₄ 195.174

Constit. of *Coix lachryma-jobi* (Job's tears), wheat and sweet corn (*Zea mays*). Cryst. Mp 198-199.5° (179-181°). λ_{max} 258 (log ε 4.06); 286 (sh) (log ε 3.78) (EtOH).

7-Me ether, 2-O-β-D-glucopyranoside: *HMBOA-Glc*

[17622-26-3]

C₁₅H₁₉NO₉ 357.316

Constit. of the roots of *Coix lachryma-jobi* (Job's tears) and *Aphelandra* spp. Cryst. (MeOH). Mp 250-251° (230-235°). [α]_D²⁰ +38.4 (MeOH). λ_{max} 205 (ε 12340); 259 (ε 9141); 282 (sh) (ε 7100) (MeOH).

Di-Me ether: 2,7-Dimethoxy-2H-1,4-benzoxazin-3(4H)-one

[131761-45-0]

C₁₀H₁₁NO₄ 209.201

Cryst. Mp 149-150°.

N-Hydroxy, O⁷-Me, 2-O-β-D-glucopyranoside: *DIMBOA-Glc*

[15893-52-4]

[147802-16-2, 147802-15-1, 157194-49-5]

C₉H₉NO₅ 211.174

Isol. from wheat, in which it is present mainly as glucoside. Appears to be a natural aphicide, insecticide and fungicide. Involved in the *in vivo* detoxification of herbicides, e.g. Simazine. Antialgal agent. Pink needles. Mp 168-169° (138°).

N-Hydroxy, O⁷-Me, 2-O-β-D-glucopyranoside: *DIMBOA-Glc*

[113565-32-5]

[157241-65-1, 18607-79-9]

C₁₅H₁₉NO₁₀ 373.316

Isol. from sweet corn (*Zea mays*) and *Aphelandra* spp. Needles (EtOH). Mp 262-263° dec.

N-Methoxy, 7-Me ether: 2-Hydroxy-4,7-dimethoxy-2H-1,4-benzoxazin-3(4H)-one, 9CI. *HDMBOA*

[149182-67-2]

C₁₀H₁₁NO₅ 225.201

Isol. from sweet corn (*Zea mays*).

N-Methoxy, 7-Me ether, 2-O-β-D-glucopyranoside: *HDMBOA-Glc*

[113565-33-6]

[29845-81-6]

C₁₆H₂₁NO₁₀ 387.343

Constit. of the roots of *Coix lachryma-joba* (Job's tears) and from wheat and

sweet corn (*Zea mays*). Pale yellow cryst. Mp 143-145°. $[\alpha]_D^{21} +22$ (c, 0.25 in H₂O). λ_{\max} 262 (log ϵ 3.97); 285 (sh) (log ϵ 3.84) (EtOH).

8-Methoxy, 7-Me ether, 2-O- β -D-glucopyranoside: [40246-09-1]

C₁₆H₂₁NO₁₀ 387.343

Isol. from sweet corn (*Zea mays*).

Wahlroos, O. et al., *Acta Chem. Scand.*, 1959, **13**, 1906 (isol, struct)

Reimann, J.E. et al., *Biochemistry*, 1964, **3**, 847 (biosynth)

Hofman, J. et al., *Tet. Lett.*, 1969, 5001 (isol, uv, glucoside)

Woodward, M.D. et al., *Plant Physiol.*, 1979, **63**, 9; *CA*, **90**, 147969q (isol)

Argandoña, V.H. et al., *Phytochemistry*, 1980, **19**, 1665; 1981, **20**, 673 (isol)

Nagao, T. et al., *Phytochemistry*, 1985, **24**, 2959-2962 (isol, uv, pmr, abs config)

Atkinson, J. et al., *J.O.C.*, 1991, **56**, 1788 (deriv, synth, pmr, ms, ir)

Bravo, H.R. et al., *J. Agric. Food Chem.*, 1996, **44**, 1569 (DIMBOA, activity)

Kluge, M. et al., *Tetrahedron*, 1996, **52**, 10389-10398 (synth)

Kluge, M. et al., *Carbohydr. Res.*, 1997, **298**, 147 (synth)

Glawischnig, E. et al., *Phytochemistry*, 1997, **45**, 715 (biosynth)

Larsen, E. et al., *J. Agric. Food Chem.*, 2000, **48**, 2556-2558 (isol)


Baumeler, A. et al., *Phytochemistry*, 2000, **53**, 213-222 (HMBOA, HMBOA-Glc)

Alipieva, K.I. et al., *Phytochemistry*, 2003, **64**, 1413-1417 (DHBOA-Glc, cmr)

Macias, F. et al., *J. Agric. Food Chem.*, 2006, **54**, 991-1000 (DIMBOA, isol, uv, pmr, cmr)

2-[2-(2,5-Dihydroxybenzoyl)amino]-5-hydroxybenzoic acid D-539

N-(2,5-Dihydroxybenzoyl)-5-hydroxyanthranilic acid



C₁₄H₁₁NO₆ 289.244

2'-O- β -D-Glucopyranoside: *Yokonoside*

[53823-12-4]

[61377-88-6]

C₂₀H₂₁NO₁₁ 451.386

Isol. from the roots of *Aconitum japonicum*. Powder. Mp 205-210°.

2'-O- β -D-Glucopyranoside, Me ester:

C₂₁H₂₃NO₁₁ 465.413

Isol. from callus cultures of *Delphinium stephisagria*. Amorph. $[\alpha]_D^{25} -11.2$ (c, 1.9 in MeOH). λ_{\max} 215 (log ϵ 4.39); 237 (log ϵ 4); 280 (log ϵ 3.8); 319 (log ϵ 3.7) (MeOH).

5'-Me ether, 2'-O- β -D-glucopyranoside, Me ester:

C₂₂H₂₅NO₁₁ 479.44

Isol. from callus cultures of *Delphinium stephisagria*. Amorph. solid. $[\alpha]_D^{25} -14.2$ (c, 0.22 in MeOH). λ_{\max} 216 (log ϵ 4.6); 241 (log ϵ 4.17); 281 (log ϵ 3.95); 322 (log ϵ 3.88) (MeOH).

5'-Deoxy, 2'-O- β -D-glucopyranoside, Me

ester:

C₂₁H₂₃NO₁₀ 449.413

Isol. from callus cultures of *Delphinium stephisagria*. Amorph. $[\alpha]_D^{25} -22.9$ (c, 0.25 in MeOH). λ_{\max} 207 (log ϵ 4.31); 236 (log ϵ 4); 281 (log ϵ 3.82); 355 (log ϵ 3.5) (MeOH).

6-Methoxy, 2'-O- β -D-glucopyranoside,

Me ester:

C₂₂H₂₅NO₁₂ 495.439


Isol. from callus cultures of *Delphinium stephisagria*. Amorph. solid. $[\alpha]_D^{25} -27.9$ (c, 0.21 in MeOH). λ_{\max} 213 (log ϵ 4.5); 240 (log ϵ 4); 305 (log ϵ 3.8) (MeOH).

Kosuge, T. et al., *Chem. Pharm. Bull.*, 1976, **24**, 176 (*Yokonoside*)

Naruta, S. et al., *Yakugaku Zasshi*, 1976, **96**, 945-957 (*Yokonoside*, synth)

Diaz, J.G. et al., *Phytochemistry*, 2005, **66**, 723-739 (isol, pmr, cmr, ms)

8-(3,4-Dihydroxybenzyl)-2,3,9,10-berbintetrol D-540



C₂₄H₂₃NO₆ 421.449

(8*R*,13*aS*)-form


Javaberine B

[387868-70-4]

Alkaloid from the roots of *Talinum paniculatum*. Powder (as hexa-Ac). $[\alpha]_D^{26} +8$ (c, 0.8 in CHCl₃) (hexa-Ac). λ_{\max} 268 (log ϵ 3.3) (MeOH) (hexa-Ac).

Shimoda, H. et al., *Heterocycles*, 2001, **55**, 2043-2050

8-(3,4-Dihydroxybenzyl)-2,3,10,11-berbintetrol D-541



C₂₄H₂₃NO₆ 421.449

(8*R*,13*aS*)-form

Javaberine A

[387868-69-1]


Alkaloid from the roots of *Talinum paniculatum*. Powder (as hexa-Ac). $[\alpha]_D^{24} +5$ (c, 0.9 in CHCl₃). λ_{\max} 287 (log ϵ 3.9) (no solvent reported).

Shimoda, H. et al., *Heterocycles*, 2001, **55**, 2043-2050

3,4-Dihydroxy-2,5-bis(hydroxymethyl)pyrrolidine D-542

3,4-Dihydroxy-2,5-pyrrolidinedimethanol, 9CI

[59920-31-9]



(2*R*,3*R*,4*R*,5*R*)-form

C₆H₁₃NO₄ 163.173

(2*R*,3*R*,4*R*,5*R*)-form

2,5-Dideoxy-2,5-imino-D-mannitol

Alkaloid from *Albizia myriophylla*, *Baphia nitida*, *Connarus ferrugineus*, *Derris elliptica*, *Lonchocarpus sericeus* and *Stemona tuberosa*. Also isol. from a *Cylindrospermum* sp. Glycosidase inhibitor. Cryst. (EtOH/Me₂CO/EtOAc). Mp 115-117°. $[\alpha]_D^{20} +56.4$ (c, 7 in H₂O).

3-O- β -D-Glucopyranoside: [865486-85-7]

C₁₂H₂₃NO₉ 325.315

Alkaloid from *Albizia myriophylla* and *Baphia nitida*. Powder. $[\alpha]_D +7.2$ (c, 0.26 in H₂O).

1'-O- β -D-Fructofuranoside:

C₁₂H₂₃NO₉ 325.315

Alkaloid from the leaves of *Baphia nitida*. Powder. $[\alpha]_D -3.1$ (c, 1.1 in H₂O).

(2*R*,3*R*,4*R*,5*S*)-form

2,5-Dideoxy-2,5-imino-D-glucitol

Alkaloid from the roots of *Stemona tuberosa*. Sl. yellow powder. Mp 139-142.5° (132-136°). $[\alpha]_D +26.1$ (c, 0.84 in H₂O).

(2*S*,3*R*,4*R*,5*S*)-form

2,5-Dideoxy-2,5-imino-L-itol, 9CI

[105015-44-9]

Mp 161-162°. $[\alpha]_D +14.3$ (c, 0.93 in H₂O).

(2*S*,3*R*,4*S*,5*S*)-form

L-alto-form. 2,5-Dideoxy-2,5-imino-L-altritol

Yellow oil. C-3 and C-4 are pseudoasymmetric.

Hydrochloride: [152785-78-9]

Mp 191-192°. $[\alpha]_D^{22} -51$ (c, 1.1 in H₂O).

(2*R*,3*R*,4*S*,5*R*)-form

allo-form. 2,5-Dideoxy-2,5-iminoallitol

[368424-96-8]

Hygroscopic gum. Obt. only in admixture with the D-alto isomer. Called D-allo- in the ref. although it is a meso-compd.

(2*R*,3*S*,4*R*,5*S*)-form

all-cis-form. 2,5-Dideoxy-2,5-iminogalactitol

Oil(?). Opt. inactive (meso-).

Welter, A. et al., *Phytochemistry*, 1976, **15**, 747 (isol, ir, pmr, cmr, ms, struct)

Lamotte-Brasseur, J. et al., *Acta Cryst. B*, 1977, **33**, 409 (cryst struct)

Fleet, G.W.J. et al., *Tet. Lett.*, 1985, **26**, 1469 (synth, abs config)

Shing, T.K.M. et al., *Chem. Comm.*, 1987, 262 (synth)


Hung, R.R. et al., *J.O.C.*, 1991, **56**, 3849 (synth)

Liu, K.K.C. et al., *J.O.C.*, 1991, **56**, 6280 (synth)

- Zou, W. *et al.*, *Carbohydr. Res.*, 1993, **242**, 311 (synth)
- Legler, G. *et al.*, *Carbohydr. Res.*, 1993, **250**, 67 (synth)
- Wang, Y. *et al.*, *Angew. Chem., Int. Ed.*, 1994, **33**, 1242 (synth, cryst struct)
- Watson, A.A. *et al.*, *Phytochemistry*, 1997, **46**, 255 (isol, props)
- Huwe, C.M. *et al.*, *Synthesis*, 1997, 61 (synth, pmr, cmr, L-altro)
- Fechter, M.H. *et al.*, *Carbohydr. Res.*, 1999, **319**, 55-62 (D-galacto, D-altro, L-altro, allo, synth)
- Dondoni, A. *et al.*, *J.O.C.*, 2002, **67**, 7203-7214 (gluco, manno, altro, allo, synth, pmr)
- Wrodnigg, T.M. *et al.*, *Monatsh. Chem.*, 2002, **133**, 393-426 (rev. isol, synth, activity)
- Jüttner, F. *et al.*, *J. Phycol.*, 2003, **39**, 26-32 (*Cylindrospermum*, isol)
- Garcia, A.L.L. *et al.*, *Tet. Lett.*, 2003, **44**, 1553-1557 (all-R-form, synth)
- Asano, N. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1238-1242 (isol, pmr, cmr)
- Trost, B.M. *et al.*, *Chem. Eur. J.*, 2006, **12**, 6607-6620 (all-R-form, synth)
- Merino, P. *et al.*, *Eur. J. Org. Chem.*, 2008, 2929-2947 (synth)
- Donohoe, T.J. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 3896-3898 (synth)
- Kato, A. *et al.*, *Phytochemistry*, 2008, **69**, 1261-1265 (1'-fructoside)

1,2-Dihydroxy-3,5-bis(hydroxymethyl)-1H-pyrrolizidine D-543

Hexahydro-1,2-dihydroxy-1H-pyrrolizine-3,5-dimethanol, 9CI



C₉H₁₇NO₄ 203.238

(1S,2R,3R,5R,7aR)-form D-543

Hyalanthacin B₁
[240117-29-7]

Alkaloid from *Hyalanthoides non-scripta* and *Scilla campanulata*. [α]_D +41.3 (c, 1.04 in H₂O).

(1S,2R,3R,5S,7aR)-form D-543


Hyalanthacin B₂
[240117-31-1]

Alkaloid from *Scilla campanulata*. [α]_D +41.3 (c, 0.36 in H₂O).

- Kato, A. *et al.*, *Carbohydr. Res.*, 1999, **316**, 95-103 (isol, pmr, cmr)
- Sengoku, T. *et al.*, *Tetrahedron*, 2008, **64**, 8052-8058 (synth, abs config)

1,9-Dihydroxycanthin-6-one D-544

1,9-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI



C₁₄H₈N₂O₃ 252.229

9-Me ether: 1-Hydroxy-9-methoxycanthin-6-one

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the roots of *Eurycoma longifolia*. Yellow powder (MeOH). Mp 235-237°. λ_{max} 274 (log ε 3.05); 283 (log ε 3.15); 358 (log ε 3.11); 374 (log ε 3.06) (MeOH).

Kuo, P.C. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1324-1327 (1-Hydroxy-9-methoxycanthin-6-one)

1,11-Dihydroxycanthin-6-one D-545

1,11-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one

C₁₄H₈N₂O₃ 252.229

O¹-Me: 11-Hydroxy-1-methoxycanthin-6-one
[109521-83-7]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the wood of *Brucea antidysenterica*. Cytotoxic. Yellow needles (CHCl₃/Et₂O). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 237-240° dec. λ_{max} 347 (ε 11220); 370 (ε 8130) (MeOH) (Berdy).

O¹¹-Me: 1-Hydroxy-11-methoxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one.

1-Hydroxy-11-methoxycanthin-6-one

[97915-43-0]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the root bark of *Brucea antidysenterica* (Simaroubaceae). Yellow needles (CHCl₃/Et₂O/hexane). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 220-223° dec. λ_{max} 244 (log ε 3.83); 275 (sh) (log ε 3.59); 282 (log ε 3.83); 320 (sh) (log ε 3.72); 348 (log ε 3.98); 362 (log ε 4.08); 380 (log ε 4.05) (no solvent reported). λ_{max} 282 (ε 6760); 348 (ε 9550); 362 (ε 12020); 380 (ε 11220) (MeOH) (Berdy).

Di-Me ether: 1,11-Dimethoxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one.

1,11-Dimethoxycanthin-6-one

[103839-24-3]

C₁₆H₁₂N₂O₃ 280.282

Alkaloid from the stem of *Brucea antidysenterica* (Simaroubaceae). Yellow cryst. (MeOH/CHCl₃). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 217-220° dec. λ_{max} 253 (ε 8740); 271 (ε 8600); 280 (ε 8400); 355 (ε 16400) (EtOH) (Berdy).

Harris, A. *et al.*, *Planta Med.*, 1985, **51**, 151-153 (11-Me ether, isol, uv, pmr, ms, struct)

Fukamiya, N. *et al.*, *J. Nat. Prod.*, 1986, **49**, 428-434 (di-Me ether, isol, uv, ir, pmr, cmr, ms, struct)

Fukamiya, N. *et al.*, *Planta Med.*, 1987, **53**, 140-143 (mono-Me ethers)

4,5-Dihydroxycanthin-6-one D-546

4,5-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI

C₁₄H₈N₂O₃ 252.229

4-Me ether: 5-Hydroxy-4-methoxycanthin-6-one. **Nigakinone**

[18110-86-6]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the stems of *Picrasma ailanthoides*, the wood of *Picrasma excelsa* (Jamaican quassia) and the heartwood of *Picrasma quassioides* (Simaroubaceae). Active against gram-positive bacteria. Yellow needles

(MeOH). Mp 223-224° dec.

4-Me ether, N-oxide: 5-Hydroxy-4-methoxy-6-canthinone 3-N-oxide

[129722-98-1]

C₁₅H₁₀N₂O₄ 282.255

Alkaloid from the wood of *Quassia amara* (Surinam quassia) (Simaroubaceae). Red-orange powder (EtOH). Mp 300°.

4-Me ether, Ac:

Needles (MeOH). Mp 194° dec.

5-Me ether: 4-Hydroxy-5-methoxycanthin-6-one. **Picrasidine Q**

[101219-61-8]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the root wood of *Picrasma quassioides* and *Picrasma excelsa* and from cell suspension cultures of *Brucea javanica* (Simaroubaceae). Pale yellow needles (CHCl₃/MeOH). Mp 286-289°. The struct. originally reported was revised by Liu *et al.* to 4-Hydroxy-3-methylcanthine-5,6-dione. X-ray anal. has shown that the struct. originally proposed was correct. The compd. isol. by Liu *et al.* should be reassigned as 4,5-Dihydroxycanthin-6-one, D-546. λ_{max} 247 (log ε 4.24); 270 (sh) (log ε 3.97); 288 (log ε 3.98); 299 (sh) (log ε 3.95); 340 (sh) (log ε 3.82); 370 (sh) (log ε 3.85); 385 (log ε 3.98) (EtOH).

Di-Me ether: 4,5-Dimethoxycanthin-6-one. **Methylnigakinone**

[18110-87-7]

C₁₆H₁₂N₂O₃ 280.282

Alkaloid from the stems and wood of *Picrasma ailanthoides*, the heartwood of *Picrasma quassioides* and trunk bark of *Odyendea gabonensis* (Simaroubaceae). Active against gram-positive bacteria. Pale yellow needles (MeOH). Mp 147.3-147.5°.

Di-Me ether, picrate:

Bright yellow prisms (EtOH). Mp 185-186° dec.

Inamoto, N. *et al.*, *Bull. Chem. Soc. Jpn.*, 1961, **34**, 888-889 (*Methylnigakinone*)

Kondo, Y. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 837-839 (*Nigakinone*, isol, pmr, struct)

Wagner, H. *et al.*, *Planta Med.*, 1979, **36**, 113-119 (*Nigakinone*, isol)

Yang, J.S. *et al.*, *Yaoxue Xuebao*, 1979, **14**, 167-177 (activity)

Forgacs, P. *et al.*, *Planta Med.*, 1982, **46**, 187-189 (*Methylnigakinone*)

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4901-4902 (*Picrasidine Q*, isol, uv, ir, pmr, ms)

Barbetti, P. *et al.*, *Planta Med.*, 1990, **56**, 216-217 (4-Me ether N-oxide)

Li, H.Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1807-1811 (*Picrasidine Q*, cryst struct)

5,9-Dihydroxycanthin-6-one D-547

5,9-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI

C₁₄H₈N₂O₃ 252.229

λ_{max} 214 (sh) (log ε 3.8); 245 (log ε 3.4); 265 (sh) (log ε 3.29); 275 (log ε 3.41); 330 (log ε 3.15) (MeOH).

Di-Me ether: 5,9-Dimethoxycanthin-6-one
[155861-54-4]

C₁₆H₁₂N₂O₃ 280.282

Alkaloid from the bark of *Eurycoma longifolia* (Simaroubaceae). Needles.

Mp 300°.

Mitsunaga, K. *et al.*, *Phytochemistry*, 1994, **35**, 799-802 (*isol, uv, ir, pmr, ms, struct*)**5,11-Dihydroxycanthin-6-one** D-548*5,11-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI*C₁₄H₈N₂O₃ 252.229*Di-Me ether: 5,11-Dimethoxycanthin-6-one*

[152592-77-3]

C₁₆H₁₂N₂O₃ 280.282Alkaloid from cell suspension cultures of *Brucea javanica* (Simaroubaceae). λ_{\max} 228 (log ϵ 3.88); 319 (log ϵ 3.55); 344 (sh) (log ϵ 3.5); 358 (log ϵ 3.21); 375 (log ϵ 3.69) (MeOH).Cheng, K.C.S. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1993, **40**, 403-405; *CA*, **120**, 101909e**8,9-Dihydroxycanthin-6-one** D-549*8,9-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one*C₁₄H₈N₂O₃ 252.229*9-Me ether: 8-Hydroxy-9-methoxycanthin-6-one*

[143257-71-0]

C₁₅H₁₀N₂O₃ 266.256Alkaloid from the stem bark of *Picrolemma granatensis* (Simaroubaceae). Yellow needles. Mp 198-201°. λ_{\max} 222 (log ϵ 3.78); 264 (log ϵ 3.45); 273 (log ϵ 3.42); 325 (log ϵ 3.38); 339 (log ϵ 3.38); 406 (log ϵ 2.85) (EtOH).Rodrigues Fo, E. *et al.*, *Phytochemistry*, 1992, **31**, 2499-2501 (*isol, uv, ir, pmr, cmr, ms, struct*)**9,10-Dihydroxycanthin-6-one** D-550*9,10-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI*C₁₄H₈N₂O₃ 252.229*9-Me ether: 10-Hydroxy-9-methoxycanthin-6-one*


[155861-52-2]

C₁₅H₁₀N₂O₃ 266.256Alkaloid from the bark of *Eurycoma longifolia* (Simaroubaceae). Yellow needles. Mp 255-257°. λ_{\max} 230 (sh) (log ϵ 3.02); 282 (log ϵ 3.17); 304 (sh) (log ϵ 2.63); 360 (log ϵ 2.75); 375 (log ϵ 2.73) (MeOH).*Di-Me ether: 9,10-Dimethoxycanthin-6-one*

[155861-51-1]

C₁₆H₁₂N₂O₃ 280.282Alkaloid from the bark of *Eurycoma longifolia* (Simaroubaceae). Yellow needles. Mp 213-215°. λ_{\max} 232 (log ϵ 3.87); 270 (sh) (log ϵ 4.05); 280 (log ϵ 4.17); 300 (log ϵ 3.61); 310 (log ϵ 3.56); 359 (log ϵ 3.79); 375 (log ϵ 3.75) (MeOH).Mitsunaga, K. *et al.*, *Phytochemistry*, 1994, **35**, 799-802 (*isol, uv, ir, pmr, cmr, ms, struct*)**10,11-Dihydroxycanthin-6-one** D-551*10,11-Dihydroxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one*C₁₄H₈N₂O₃ 252.229*10-Me ether: 11-Hydroxy-10-methoxycanthin-6-one*

[155861-53-3]

C₁₅H₁₀N₂O₃ 266.256Alkaloid from the wood of *Eurycoma longifolia* (Simaroubaceae). Yellow needles. Mp 235-237°. λ_{\max} 218 (log ϵ 4.02); 245 (log ϵ 3.57); 265 (log ϵ 3.33); 320 (log ϵ 3.24); 394 (log ϵ 3.24) (MeOH).Mitsunaga, K. *et al.*, *Phytochemistry*, 1994, **35**, 799-802 (*isol, uv, ir, pmr, ms, struct*)**1,6-Dihydroxy-9H-carbazole-3-carboxaldehyde** D-552*3-Formyl-1,6-dihydroxycarbazole*C₁₃H₉NO₃ 227.219*6-Me ether: 1-Hydroxy-6-methoxy-9H-carbazole-3-carboxaldehyde. 3-Formyl-1-hydroxy-6-methoxycarbazole. Clausine I*

[182261-94-5]

C₁₄H₁₁NO₃ 241.246Alkaloid from stem bark of *Clausena excavata*. Powder (Me₂CO). Mp 222-224°. λ_{\max} 204 (log ϵ 4.24); 223 (log ϵ 4.14); 242 (log ϵ 4.14); 255 (log ϵ 4.09); 278 (log ϵ 4.23); 296 (log ϵ 4.17); 341 (log ϵ 3.85); 353 (log ϵ 3.86) (MeOH).*Di-Me ether: 1,6-Dimethoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-1,6-dimethoxy-9H-carbazole. 6-Methoxymurrayanine*

[132922-59-9]

C₁₅H₁₃NO₃ 255.273Alkaloid from the roots of *Clausena lansium* (wampee) (Rutaceae). Prisms (Me₂CO/hexane). Mp 231-233°. λ_{\max} 239 (log ϵ 4.35); 251 (sh) (log ϵ 4.2); 277 (log ϵ 4.4); 294 (log ϵ 4.36); 335 (log ϵ 3.85); 349 (log ϵ 3.85) (MeOH).Li, W.-S. *et al.*, *Phytochemistry*, 1991, **30**, 343-346 (*di-Me ether*)Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 133-140 (*Clausine I*)Bernal, P. *et al.*, *Synthesis*, 2007, 1943-1948 (*di-Me ether, synth*)**1,7-Dihydroxy-9H-carbazole-3-carboxaldehyde** D-553*3-Formyl-1,7-dihydroxy-9H-carbazole*C₁₃H₉NO₃ 227.219*1-Me ether: 7-Hydroxy-1-methoxy-9H-carbazole-3-carboxaldehyde. 3-Formyl-7-hydroxy-1-methoxy-9H-carbazole. Clausine Q*C₁₄H₁₁NO₃ 241.246Alkaloid from *Clausena excavata*. Brown powder (Me₂CO). Mp 85-87°. λ_{\max} 201; 242; 252 (sh); 286; 297 (sh); 325 (sh); 338 (MeOH).Wu, T.-S. *et al.*, *Phytochemistry*, 1999, **52**, 523-527 (*isol, uv, pmr, ms*)**1,8-Dihydroxy-9H-carbazole-3-carboxaldehyde** D-554*3-Formyl-1,8-dihydroxycarbazole*C₁₃H₉NO₃ 227.219*Di-Me ether: 1,8-Dimethoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-1,8-dimethoxycarbazole. Clausenal*

[162857-93-4]

C₁₅H₁₃NO₃ 255.273Alkaloid from leaves of *Clausena heptaphylla* (Rutaceae). Exhibits antimicrobial activity against both gram-positive and -negative bacteria. Antifungal. Needles (C₆H₆/petrol). Mp 198°. λ_{\max} 231 (ϵ 28183); 257 (ϵ 12300); 261 (ϵ 12590); 327 (ϵ 5012) (EtOH) (Berdy).Chakraborty, A. *et al.*, *Phytochemistry*, 1995, **38**, 787**2,6-Dihydroxy-9H-carbazole-3-carboxaldehyde** D-555*3-Formyl-2,6-dihydroxycarbazole*C₁₃H₉NO₃ 227.219*6-Me ether: 2-Hydroxy-6-methoxy-9H-carbazole-3-carboxaldehyde. 3-Formyl-2-hydroxy-6-methoxycarbazole. Lansine*

[74606-99-8]

C₁₄H₁₁NO₃ 241.246Alkaloid from the leaves of *Clausena lansium* (wampee) and the stem bark of *Micromelum hirsutum*. Mp 225-226°. λ_{\max} 233; 248; 309; 340; 368 (MeOH).*Di-Me ether: 2,6-Dimethoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-2,6-dimethoxycarbazole. Glycozolidal*

[51971-09-6]

C₁₅H₁₃NO₃ 255.273Alkaloid from the roots of *Glycosmis pentaphylla* (Rutaceae). Mp 185°.Sharma, R.B. *et al.*, *Chem. Ind. (London)*, 1980, 158 (*synth*)Prakash, D. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 1075-1076 (*Lansine, isol, uv, ir, pmr, ms, struct*)Bhattacharyya, P. *et al.*, *J. Nat. Prod.*, 1985, **48**, 465-466 (*di-Me ether*)Ma, C. *et al.*, *Planta Med.*, 2005, **71**, 261-267 (*Lansine, isol, cmr*)**2,7-Dihydroxy-9H-carbazole-3-carboxaldehyde, 9CI** D-556*3-Formyl-2,7-dihydroxycarbazole. Clausine O*

[250259-35-9]

C₁₃H₉NO₃ 227.219Alkaloid from *Clausena excavata*. Yellowish needles. Mp >280°. λ_{\max} 200; 223; 240; 252 (sh); 291 (sh); 301; 321 (sh); 340 (MeOH).*7-Me ether: 2-Hydroxy-7-methoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-2-hydroxy-7-methoxycarbazole*

[119736-83-3]

C₁₄H₁₁NO₃ 241.246Alkaloid from the root bark of *Clausena harmandiana* and root of *Murraya siamensis*. Cryst. (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 226-227°. λ_{\max} 224

(ϵ 18200); 240 (ϵ 18200); 294 (sh) (ϵ 29510); 300 (ϵ 38000); 338 (ϵ 7760) (MeOH).

Di-Me ether: 2,7-Dimethoxy-9H-carbazole-3-carboxaldehyde. 3-Formyl-2,7-dimethoxycarbazole

[132160-50-0]
C₁₅H₁₃NO₃ 255.273

Alkaloid from the roots of *Murraya siamensis* and stem bark of *Clausena excavata*. Mp 219–220°.

Chaichantipiyuth, C. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1285–1288 (7-Me ether, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Ruangrungsi, N. *et al.*, *J. Nat. Prod.*, 1990, **53**, 946–952 (7-Me ether, *di-Me ether*, *isol*, *pmr*, *cmr*, *struct*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 133–140; 1999, **52**, 523–527 (*di-Me ether*, *Clausine O*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 48–52 (*Clauszoline J*)

Yenjai, C. *et al.*, *Planta Med.*, 2000, **66**, 277–281 (*activity*, *Clausine H*)

Kataeva, O. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 3099–3101 (*synth*)

2,8-Dihydroxy-9H-carbazole-3-carboxaldehyde, 9CI

6-Formyl-1,7-dihydroxycarbazole. *Clauszoline M*

[187110-72-1]
C₁₃H₉NO₃ 227.219

Alkaloid from *Clausena excavata*. Pale yellow powder. λ_{\max} 217; 240; 275; 290; 356 (MeOH).

8-Me ether: 2-Hydroxy-8-methoxy-9H-carbazole-3-carboxaldehyde, 9CI. 6-Formyl-7-hydroxy-1-methoxycarbazole.

Clausine A


[80054-00-8]
C₁₄H₁₁NO₃ 241.246

Alkaloid from stem bark of *Clausena excavata*. Yellowish needles (Me₂CO). Mp 184–186°. λ_{\max} 202 (log ϵ 4.27); 218 (log ϵ 4.33); 241 (log ϵ 4.52); 268 (sh) (log ϵ 4.42); 275 (log ϵ 4.53); 288 (log ϵ 4.38); 352 (log ϵ 4.04) (MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 1427–1429 (*Clausine A*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 48–52 (*isol*, *uv*, *pmr*)

1,6-Dihydroxy-9H-carbazole-3-carboxylic acid



C₁₃H₉NO₄ 243.218

6-Me ether, Me ester: 3-Carbomethoxy-1-hydroxy-6-methoxycarbazole. *Clausine G*

[186002-62-0]
C₁₅H₁₃NO₄ 271.272

Alkaloid from stem bark of *Clausena excavata*. Granules (Me₂CO). Mp >280°. λ_{\max} 224; 272; 282; 316 (sh); 353 (sh) (MeOH).

Wu, T.S. *et al.*, *Phytochemistry*, 1996, **43**, 1427–1429 (*Clausine G*)

1,7-Dihydroxy-9H-carbazole-3-carboxylic acid

C₁₃H₉NO₄ 243.218

Me ester: 3-Carbomethoxy-1,7-dihydroxy-9H-carbazole. *Clausine R*

[250259-37-1]
C₁₄H₁₁NO₄ 257.245

Alkaloid from *Clausena excavata*. Yellowish needles (Me₂CO). Mp 178–181°. λ_{\max} 214; 241; 252; 282; 320 (sh); 333 (MeOH).

Wu, T.S. *et al.*, *Phytochemistry*, 1999, **52**, 523–527 (*isol*, *uv*, *pmr*, *ms*)

2,7-Dihydroxy-9H-carbazole-3-carboxylic acid

C₁₃H₉NO₄ 243.218

7-Me ether, Me ester: *Clausine TY*

[1011725-88-4]
C₁₅H₁₃NO₄ 271.272

Alkaloid from the stem bark of *Clausena excavata*. Cytotoxic. Brownish powder. Mp 220–222°.

Di-Me ether: 2,7-Dimethoxy-9H-carbazole-3-carboxylic acid. *Clausine K*.

Clauszoline J

[182261-96-7]
C₁₅H₁₃NO₄ 271.272

Alkaloid from stem bark of *Clausena excavata*. Brownish powder (Me₂CO). Mp 250–256°. λ_{\max} 240; 276; 308; 318 (MeOH).

Di-Me ether, Me ester: *Clausine H*.

Clauszoline C

[182261-90-1]
C₁₆H₁₅NO₄ 285.299

From stem bark of *Clausena excavata*. Shows antiplasmodial activity. Grey needles (Me₂CO). Mp 192–194°. λ_{\max} 205; 222; 246; 282; 309; 320; 336 (sh) (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2231–2235; 1997, **45**, 48–52 (*Clauszoline S*)

Wu, T.S. *et al.*, *Phytochemistry*, 1996, **43**, 133–140 (*Clausines H, K*)


Yenjai, C. *et al.*, *Planta Med.*, 2000, **66**, 277–281 (*Clausine H*: *activity*)

Kataeva, O. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 3099–3101 (*Clausines H, K*, *synth*)

Taufiq-Yap, Y.H. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 810–813 (*Clausine TY*)

4,8-Dihydroxy- β -carboline-1-carboxylic acid

4,8-Dihydroxypyrido[3,4-b]indole-1-carboxylic acid



C₁₂H₈N₂O₄ 244.206


Di-Me ether, Me ester: 1-Carbomethoxy-4,8-dimethoxy- β -carboline

[123828-65-9]
C₁₅H₁₄N₂O₄ 286.287

Alkaloid from the leaves of *Ailanthus altissima* (Simaroubaceae). Pale-yellow needles. Mp 184–185°.

Souleles, C. *et al.*, *Planta Med.*, 1989, **55**, 286 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

3,25-Dihydroxycevan-6-one



C₂₇H₄₃NO₃ 429.642

(3 β ,5 α ,17 β ,22 β)-form

Taipeienine. *Taipeienine*


[151183-22-1]

Alkaloid from bulbs of *Fritillaria taipiensis* var. *ningriensis* (Liliaceae). Needles (Me₂CO). Mp 120–122°. $[\alpha]_D^{25}$ +12.9 (c, 0.3 in MeOH).

Hu, C.H. *et al.*, *Yaoxue Xuebao*, 1993, **28**, 516–521; *CA*, **119**, 245534f (*isol*, *struct*)

Feng, R. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 383–384; *CA*, **121**, 153343f (*isol*, *ir*, *cmr*, *struct*)

6,12-Dihydroxycevan-3-one, 9CI



C₂₇H₄₃NO₃ 429.642

(5 α ,6 β ,25 α)-form

Sevedinine


[849802-77-3]

Alkaloid from the aerial parts of *Korolkowia severtzovii*. Cryst. (Me₂CO). Mp 233–235°. $[\alpha]_D$ -13.6 (c, 0.4 in CHCl₃).

Abdullaeva, D.U. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 394–397 (*isol*, *pmr*, *cmr*)

2,3-Dihydroxycycloarta-22,24-dien-26-oic acid

2,3-Dihydroxy-9,19-cyclolanost-22,24-dien-26-oic acid



C₃₀H₄₆O₄ 470.691**(2α,3β,22E,24E)-form**


N-(Tetrahydro-4,5-dimethyl-2-oxo-3-furanyl)amide, 3-O-[α-L-rhamnopyranosyl-(1→2)-O-β-D-glucopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Mussaendoside P** [157536-45-3]

C₆₀H₉₅NO₂₃ 1198.403

Constit. of *Mussaenda pubescens*. Amorph. powder. [α]_D²⁵ +7 (c, 0.5 in Py). λ_{max} 265 (MeOH).

Zhao, W.M. et al., *Chin. Chem. Lett.*, 1994, 5, 309Zhao, W. et al., *J. Nat. Prod.*, 1994, 57, 1613 (isol, pmr, cmr)**1,4-Dihydroxy-2-cyclopentene-1-carboxylic acid**

D-565



(1R,4R)-form

C₆H₈O₄ 144.127**(1R,4R)-form**

Nitrile, 1-O-β-D-glucopyranoside: **Volkenin**. Epitetracycline B [66575-40-4]

C₁₂H₁₇NO₇ 287.269

Isol. from *Adenia volkensii*. Syrup. [α]_D²² +27 (c, 0.7 in MeOH).

Nitrile, 1-O-β-D-glucopyranoside, penta-Ac: [115014-72-7] Mp 124-124.5°. [α]_D²² +41.5 (c, 0.4 in MeOH).

(1R,4S)-form

Nitrile, 1-O-β-D-glucopyranoside: **Taraktophyllin**. Epipassicoriacin [110115-55-4]

C₁₂H₁₇NO₇ 287.269

Isol. from *Taraktogenos heterophylla* (preferred genus name *Hydnocarpus*), *Hydnocarpus anthelmintica* and *Passiflora coriacea*. Syrup. [α]_D²³ -75 (c, 1 in MeOH).

Nitrile, 1-O-β-D-glucopyranoside, penta-Ac: [110114-82-4] Mp 126-127°. [α]_D²³ -48 (c, 0.5 in MeOH).

Nitrile, 1-O-[α-L-Rhamnopyranosyl-(1→6)-β-D-glucopyranoside]: [117557-46-7]

C₁₈H₂₇NO₁₁ 433.411

Isol. from seeds of *Hydnocarpus pentandra*. Amorph. powder. [α]_D²⁴ -84 (c, 0.5 in MeOH).

(1S,4R)-form

Nitrile, 1-O-β-D-glucopyranoside: **Epivolkenin**. *Passicoriacin* [109905-56-8]

C₁₂H₁₇NO₇ 287.269

Isol. from *Taraktogenos heterophylla*, *Hydnocarpus anthelmintica* and *Passiflora coriacea*. Syrup. [α]_D²³ +43 (c, 1 in MeOH).

Nitrile, 1-O-β-D-glucopyranoside, penta-

Ac: [110037-31-5] Mp 131-132°. [α]_D²³ +23 (c, 0.5 in MeOH).

Nitrile, 1-O-β-D-glucopyranoside, 4-O-(6-deoxy-β-D-gulopyranoside): **Passibiflorin**

[97564-61-9]

C₁₈H₂₇NO₁₁ 433.411

Isol. from *Passiflora biflora*. Noncryst. Mp 160-161° (as hexa-Ac). [α]_D²⁰ -7 (c, 0.5 in MeOH). Rare occurrence of Antiarose (6-deoxygulose) residue.

Nitrile, 1-O-β-D-glucopyranoside, 4-O-(2,6-dideoxy-β-D-xylo-hexopyranoside): **Passicapsin**

[82829-54-7]

C₁₈H₂₇NO₁₀ 417.412

Isol. from *Passiflora capsularis*. Cryst. (EtOAc). Mp 124-126° (116-119°). [α]_D²⁴ +8 (c, 0.56 in MeOH). Rare occurrence of boivoinose residue, not prev. encountered except in cardenolides.

Nitrile, 1-O-β-D-glucopyranoside, 4-O-(6-deoxy-β-D-allopyranoside): **Passitriasciatin**

[97564-60-8]

C₁₈H₂₇NO₁₁ 433.411

Isol. from *Passiflora trifasciata*. Noncryst. Mp 150-151° (as hepta-Ac). [α]_D²² +6.4 (c, 0.5 in MeOH). Revised struct. (1991).

Nitrile, 1-O-[α-L-Rhamnopyranosyl-(1→6)-β-D-glucopyranoside]:

117557-45-6

Isol. from seeds of *Hydnocarpus pentandra*.

Amorph. powder. [α]_D²⁴ -4.4 (c, 0.6 in MeOH).

Amide: 1,4-Dihydroxy-2-cyclopentene-1-carboxamide

[133071-13-3]

C₆H₉NO₃ 143.142

Isol. from the leaves of *Lindackeria dentata*. Cryst. (MeOH aq.). Mp 143-144° (synthetic). [α]_D +120 (c, 0.13 in MeOH) (natural). [α]_D +275 (c, 0.2 in MeOH) (synthetic).

(1S,4S)-form

Nitrile, 1-O-β-D-glucopyranoside: **Tetracycline B**. *Barterin*

[34323-07-4]

Isol. from *Adenia*, *Barteria*, *Turnera* and *Tetraphaëa* (preferred genus name *Passiflora*) spp. Needles (MeOH/EtOAc). Mp 169-170°. [α]_D²⁶ -75 (c, 0.5 in MeOH). [α]_D²⁵ -35.6 (c, 1 in H₂O).

Nitrile, 1-O-β-D-glucopyranoside, penta-Ac: [115014-71-6]

Mp 114-115°. [α]_D²¹ -53.5 (c, 0.5 in MeOH).

Nitrile, 1-O-β-D-glucopyranoside, 4-O-sulfate: **Tetracycline B sulfate**

[85758-30-1]

C₁₂H₁₇NO₁₀S 367.333

Isol. from *Passiflora caerulea* (blue passion flower) and other *Passiflora* spp. Glassy syrup.

(1ξ,4ξ)-form

Nitrile, 1-O-[β-D-rhamnopyranosyl-(1→6)-β-D-glucopyranoside], 4-O-sulfate:

Passicoccin

[99694-22-1]

C₁₈H₂₇NO₁₄S 513.476Isol. from *Passiflora coccinea*.

Amorph. yellow solid. Unusual occurrence of D-rhamnose.

[121783-15-1, 133576-30-4, 99694-23-2, 85799-11-7, 109906-27-6, 97564-59-5, 85758-31-2]

Russell, G.B. et al., *Phytochemistry*, 1971, 10, 1373-1377 (*Tetracycline B*)

Spencer, K.C. et al., *Planta Med.*, 1981, 43, 175-178 (*Tetracycline B*)

Siegler, D.S. et al., *Phytochemistry*, 1982, 21, 2277-2285 (*Tetracycline B sulfate*)

Spencer, K.C. et al., *Phytochemistry*, 1985, 24, 2615-2617; 1987, 26, 1661-1663 (*Passicoccin*, *Taraktophyllin*, *Epivolkenin*)

Jaroszewski, J.W. et al., *Acta Chem. Scand., Ser. B*, 1987, 41, 410-421 (*Volkenin*, *Tetracycline B*)

Jaroszewski, J.W. et al., *Tetrahedron*, 1987, 43, 2349-2354 (*Taraktophyllin*, *Epivolkenin*)

Jaroszewski, J.W. et al., *Planta Med.*, 1988, 54, 333-357 (*rutinosides*)

Olafsdottir, E.S. et al., *Acta Chem. Scand.*, 1989, 43, 51-55 (*Passibiflorin*, *Passicapsin*)

Olafsdottir, E.S. et al., *Phytochemistry*, 1989, 28, 127-132; 1991, 30, 867-869 (*Volkenin*, *Epivolkenin*, *Tetracycline B*, *Taraktophyllin*, *Passitriasciatin*)


Siegler, D.S. et al., *Phytochemistry*, 1989, 28, 931-932 (*Passicoriacin*, *Epicoriacin*, *Epitetracycline B*)

Olafsdottir, E.S. et al., *J.O.C.*, 1991, 56, 2650-2655 (*amide*, *synth*)

Jaroszewski, J.W. et al., *J.O.C.*, 2004, 70, 1001-1003 (*amide*, *isol*)

4,5-Dihydroxy-2-decenoic acid

D-566



(2E,4R*,5S*)-form

C₁₀H₁₈O₄ 202.25**(2E,4R*,5S*)-form**

2-Methylpropylamide: 4,5-Dihydroxy-N-(2-methylpropyl)-2-decenamide, 9CI.

N-Isobutyl-4,5-dihydroxy-2-decenamide. **Sylvamide**

[83905-60-6]

C₁₄H₂₇NO₃ 257.372

Alkaloid from the seeds of *Piper sylvaticum* (Piperaceae). Plates (Me₂CO/petrol). Mp 143-144°. [α]_D²¹ -2 (EtOH).

(2E,4RS,5RS)-form

Piperidine: 1-(4,5-Dihydroxy-1-oxo-2-decenyloxy)piperidine. **4,5-Dihydroxy-2-decenoic acid piperidine**

C₁₅H₂₇NO₃ 269.383


Alkaloid from the roots of *Piper nigrum* (pepper). Oil.

(2E,4RS,5SR)-form

Piperidine: Alkaloid from the roots of *Piper nigrum* (pepper). Oil.

Banerji, A. et al., *Phytochemistry*, 1982, 21, 1321-1323; 1983, 22, 1028-1030 (*Sylvamide*)

Wei, K. et al., *J. Nat. Prod.*, 2004, 67, 1005-1009 (*piperidides*)

1,2-Dihydroxydibenz[cd,f]indol-4(5H)-one, 9CI D-567C₁₅H₉NO₃ 251.241

CAS numbering shown. Alternative (phenanthrene) numbering frequently encountered, in which 1,2- are 4,3- respectively.

N-Methoxy: Piperumbellactam CC₁₆H₁₁NO₄ 281.267

Alkaloid from *Piper umbellatum* (pariparoba). Amorph. powder. λ_{\max} 203; 226; 246; 279; 313; 378 (MeOH).

1-Me ether: Aristolactam A II

[53948-07-5]

C₁₆H₁₁NO₃ 265.268

Alkaloid from the roots of *Aristolochia argentina* and *Aristolochia indica* (Aristolochiaceae). Cytotoxic. Shows platelet aggregation inhibitory activity. Cryst. (AcOH). Mp 271°.

1-Me ether, N-methoxy: Piperlactam S

[188546-49-8]

C₁₇H₁₃NO₄ 295.294

Alkaloid from leaves and stems of *Piper puberulum*. Yellow needles. Mp 242-244°. A drawing error in CAS (vol 121) erroneously shows this to be identical with Sauristolactam. λ_{\max} 195; 206; 232; 248; 274; 284; 315; 380 (MeOH).

2-Me ether: 1-Hydroxy-2-methoxydibenz[cd,f]indol-4(5H)-one, 9CI. Piperolactam A. Aristolactam F I

[112501-42-5]

C₁₆H₁₁NO₃ 265.268

Alkaloid from roots of *Piper longum* (long pepper), from *Piper attenuatum*, *Piper boehmerifolium*, *Piper hamiltonii* and *Pararistolochia flos-avis*. Shows platelet aggregation inhibitory activity. Cryst. (C₆H₆/MeOH). Mp 303-306° dec. (271-273°).

2-Me ether, N-hydroxy: N-Hydroxypiperolactam A. Piperumbellactam B

Alkaloid from *Piper umbellatum* (pariparoba). Amorph. powder. λ_{\max} 194; 203; 226; 247; 274; 280; 316; 378 (MeOH).

O¹,N-Di-Me: Sauristolactam

[128533-02-8]

C₁₇H₁₃NO₃ 279.295

Alkaloid from *Saururus cernuus* (Saururaceae) and *Piper puberulum* (Piperaceae). Pale yellow prisms (CHCl₃/hexane). Mp 290°.

O²,N-Di-Me: N-Methylpiperolactam A

[144549-43-9]

C₁₇H₁₃NO₃ 279.295

Alkaloid from stems of *Piper ribesoides*. Amorph. solid. Mp 205-207°.

Di-Me ether: Aristolactam B II. Cepharanone B. Alkaloid Y

[53948-09-7]

C₁₇H₁₃NO₃ 279.295

Alkaloid from the roots of *Aristolochia argentina*, bark of *Schefferomitra subaequalis*, callus tissue of *Stephania cepharantha* and stem bark of *Goniotalamus velutinus*. Shows platelet aggregation inhibitory activity. Cryst. (MeOH or Me₂CO). Mp 264-265° (247-250°). λ_{\max} 232 (log ϵ 4.68); 264 (log ϵ 4.61); 276 (log ϵ 4.67); 287 (log ϵ 4.66); 318 (log ϵ 4.03); 384 (log ϵ 4.01) (MeOH).

Di-Me ether, N-hydroxy: N-Hydroxyaristolactam BII. Piperumbellactam AC₁₇H₁₃NO₄ 295.294

Alkaloid from *Piper umbellatum* (pariparoba). Amorph. powder. λ_{\max} 196; 209; 231; 242; 274; 283; 311; 379 (MeOH).

O,O,N-Tri-Me: Caldensine

[74575-73-8]

C₁₈H₁₅NO₃ 293.321

Alkaloid from the roots of *Piper caldense*. Amorph. yellow solid.

Methylene ether: Benzo[f]-1,3-benzodioxolo[6,5,4-cd]indol-5(6H)-one. 10-Amino-3,4-methylenedioxy-1-phenanthrenecarboxylic acid lactam. Aristolactam II. Cepharanone A

[55610-00-9]

C₁₆H₉NO₃ 263.252

Alkaloid from the callus tissue of *Stephania cepharantha* and from *Aristolochia argentina*. Cytotoxic. Pale yellow amorph. solid (DMF). Mp 308-310° (297-298°). λ_{\max} 215; 232; 265; 277; 288; 328; 341; 376; 393 (MeOH) (Berdy).

► DE5350000

Methylene ether, N-β-D-glucopyranosyl: Cepharanone A N-β-D-glucoside

[139163-25-0]

C₂₂H₁₉NO₈ 425.394

Alkaloid from roots of *Aristolochia cinnabarina* (Aristolochiaceae). Yellow amorph. powder. $[\alpha]_D^{25}$ -3.54 (c. 0.22 in DMSO).

Methylene ether, N-hydroxy: N-Hydroxyaristolactam IIC₁₆H₉NO₄ 279.251

Alkaloid from *Piper umbellatum* (pariparoba).

Methylene ether, N-methoxy: N-Methoxyaristolactam II. Piperumbellactam DC₁₇H₁₁NO₄ 293.278

Alkaloid from *Piper umbellatum* (pariparoba). Amorph. powder. λ_{\max} 197; 209; 231; 243; 276; 282; 311; 380 (MeOH).

Gellert, E. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2477 (*Aristolactam BII*)

Crohare, R. *et al.*, *Phytochemistry*, 1974, **13**, 1957 (*Aristolactam AII*)

Akasu, M. *et al.*, *Tet. Lett.*, 1974, 3609 (*uv, ir, pmr, ms, struct*)

Dyke, S.F. *et al.*, *Phytochemistry*, 1978, **17**, 599 (*Aristolactam BII*)

Achari, B. *et al.*, *Heterocycles*, 1982, **19**, 1203 (*occur*)

Priestap, H.A. *et al.*, *Phytochemistry*, 1985, **24**, 849 (*isol, uv, ir, pmr, ms*)

Sun, N.J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 843 (*Piperolactam A*)

Desai, S.J. *et al.*, *Phytochemistry*, 1988, **27**, 1511 (*Piperolactam A*)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 1990, **53**, 309 (*Sauristolactam*)

Ruangrungsi, N. *et al.*, *Phytochemistry*, 1992, **31**, 2397 (*N-Methylpiperolactam A*)

Hong, L. *et al.*, *Phytochemistry*, 1994, **37**, 237 (*Cepharanone A glucoside*)

Wang, E.C. *et al.*, *Heterocycles*, 1996, **43**, 969 (*Aristolactam BII*)

Wu, Q. *et al.*, *Phytochemistry*, 1997, **44**, 727 (*Piperlactam S*)

Couture, A. *et al.*, *Synlett*, 1997, 1475-1477 (*synth*)

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1006-1009 (*activity*)

Chia, Y.-C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1160-1163 (*isol, activity*)

Júnior, E.L.C. *et al.*, *Pharm. Biol.*, 2003, **41**, 216-218 (*Caldensine*)

Tabopda, T.K. *et al.*, *Phytochemistry*, 2008, **69**, 1726-1731 (*Piperumbellactams A-C*)

1,3-Dihydroxydibenz[cd,f]indol-4(5H)-one D-568C₁₅H₉NO₃ 251.241**Di-Me ether: 1,3-Dimethoxydibenz[cd,f]indol-4(5H)-one, 9CI. 10-Amino-2,4-dimethoxy-1-phenanthrenecarboxylic acid lactam**

[226953-00-0]


C₁₇H₁₃NO₃ 279.295

Alkaloid from the bark of *Goniotalamus griffithii*.

Hu, Z. *et al.*, *Zhongcaoyao*, 1999, **30**, 81-83; *CA*, **131**, 29813b


5,8-Dihydroxy-4,7-dimethoxy-2,6-dimethylisoquinolinium(1+) D-569

[113585-57-2]

C₁₃H₁₆NO₅⁺ 250.274

Isol. from *Myxococcus xanthus*. Red oil (as formate). CAS no. refers to formate. λ_{\max} 267 (ϵ 17000); 341 (ϵ 4360); 425 (ϵ 5000) (MeOH/HCl).

Trowitzsch-Kienast, W. *et al.*, *Annalen*, 1988, **483** (*isol, pmr, cmr, struct*)

9,10-Dihydroxy-2,3-dimethoxytetrahydroprotoberberine D-570C₁₉H₂₁NO₄ 327.379

(ξ)-form

N-Me: *Haitinosporine*

[214075-11-3]

C₂₀H₂₄NO₄[⊕] 342.414

Quaternary alkaloid from *Tinospora hainanensis*. Component of Hai Nan Qing Niu Dan. Cryst. Mp 242-245° dec. Counterion not specified.

Di-Me ether: see Tetrahydropalmatine, T-212

O²-De-Me: 2,9,10-Trihydroxy-3-methoxytetrahydroprotuberberine

[949933-14-6]

C₁₈H₁₉NO₄ 313.352


Alkaloid from *Corydalis saxicola*. Needles (MeOH). Mp 232-234° (182-184°). Two different melting points given in ref. λ_{max} 280 (MeOH).

Guo, Y. et al., *Yaouxue Xuebao*, 1998, **33**, 350-354 (*Haitinosporine*)

Li, H.-L. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 173-175 (*Trihydroxymethoxytetrahydroprotuberberine*)

3,4-Dihydroxy-1,2-dimethylcarbazole D-571

1,2-Dimethyl-9H-carbazole-3,4-diol

C₁₄H₁₃NO₂ 227.262

3-Me ether: 4-Hydroxy-3-methoxy-1,2-dimethylcarbazole. 3-Methoxy-1,2-dimethyl-9H-carbazol-4-ol, 9CI.

Carbazomycin B

[75139-38-7]

C₁₅H₁₅NO₂ 241.289

From *Streptomyces* sp. H1051-MY-10 and *Streptovorticillium ehimense*. Enzyme Inhibitor, free radical scavenger. Active against phytopathogenic fungi and weakly active against gram-positive bacteria and yeasts. Pale yellow prisms (C₆H₆/hexane or EtOAc/hexane). Sol. MeOH, C₆H₆; fairly sol. CCl₄; poorly sol. H₂O. Mp 160-162°. λ_{max} 224 (ε 38000); 244 (ε 46800); 290 (ε 16100); 329 (ε 5300); 340 (ε 5800) (90% MeOH) (Derep).

► FE6360000

Di-Me ether: 3,4-Dimethoxy-1,2-dimethyl-9H-carbazole, 9CI. **Carbazomycin A**

[75139-39-8]

C₁₆H₁₇NO₂ 255.316

Isol. from *Streptovorticillium* sp. H1051-MY-10 and *Streptovorticillium ehimense*. Active against *Staphylococcus aureus*, weakly active against fungi and yeast. Pale-yellow needles (EtOAc/hexane), plates (CH₂Cl₂/hexane). Sol. MeOH, C₆H₆, CCl₄. Mp 51-52.5° (natural) Mp 143-146° (synthetic). λ_{max} 224 (ε 38000); 244 (ε 46800); 290 (ε 16100); 329 (ε 5300); 340 (ε 5800) (90% MeOH) (Derep). λ_{max} 242 (ε 51000); 273 (ε 36000); 293 (ε 20700); 327 (ε 4700); 340 (ε 4600) (MeOH) (Berdy).

► FE6206000

Sakano, K. et al., *J. Antibiot.*, 1980, **33**, 683; 961

Kaneda, M. et al., *Heterocycles*, 1981, **15**, 993 (*cryst struct, pmr, cmr*)

Yamasaka, K. et al., *J. Antibiot.*, 1983, **36**, 552 (*biosynth, cmr*)

Knoelker, H.-J. et al., *Chem. Comm.*, 1989, 1468 (*synth*)

Moody, C.J. et al., *J.C.S. Perkin 1*, 1989, 376; 2463 (*synth*)

Kaneda, M. et al., *J. Antibiot.*, 1990, **43**, 1623 (*biosynth*)

Knoelker, H.-J. et al., *Helv. Chim. Acta*, 1993, **76**, 2500 (*synth*)


Hibino, S. et al., *Heterocycles*, 1993, **35**, 441 (*synth*)

Clive, D.L.J. et al., *J.O.C.*, 1993, **58**, 2442 (*synth*)

Beccalli, E.M. et al., *Tetrahedron*, 1996, **52**, 3029 (*synth*)

Knoelker, H.-J. et al., *Tet. Lett.*, 1999, **40**, 6915-6918 (*synth*)

Crich, D. et al., *Tetrahedron*, 2004, **60**, 1513-1516 (*Carbazomycin B, synth*)

7,9-Dihydroxy-8,10-dimethyl-2,4-dodecadienoic acid D-572C₁₄H₂₄O₄ 256.341**(2Z,4E,7S,8S,9R,10S)-form**

7-O-(E-Cinnamoyl), amide: **Basiliskamide B**

[474410-87-2]

C₂₃H₃₁NO₄ 385.502

Prod. by a marine *Bacillus laterosporus*. Solid. [α]_D²⁵ -12 (MeOH). λ_{max} 262 (ε 43000) (MeOH).

9-O-(E-Cinnamoyl), amide: **Basiliskamide A**. Antibiotic TAN 1771B. TAN 1771B

[170034-41-0]

[474410-77-0]

C₂₃H₃₁NO₄ 385.502


Prod. by a *Bacillus* sp. and a marine *Bacillus laterosporus*. Antifungal agent. Solid. [α]_D²⁵ -78 (MeOH). λ_{max} 204 (ε 19400); 216 (ε 22700); 222 (ε 21600); 262 (ε 40000) (MeOH) (Berdy).

Japan. Pat., 1995, 206 797; CA, **123**, 306561u (TAN 1771B)

Barsby, T. et al., *J. Nat. Prod.*, 2002, **65**, 1447-1451 (*Basiliskamides*)

Lipomi, D.J. et al., *Org. Lett.*, 2004, **6**, 3533-3536 (*synth*)

Yadav, J.S. et al., *Tet. Lett.*, 2008, **49**, 5427-5430 (*synth*)


7,18-Dihydroxy-16,18-dimethyl-10-phenyl-22-oxa[12]cytochalsa-6(12),13,19-triene-1,21-dione D-573

Relative configuration

C₂₈H₃₅NO₅ 465.588

Claimed to be first example of a 22-oxa[12]cytochalsan, but see Rosellichalasin, R-128. Metab. from an unidentified *Daldinia* fungus. Amorph. solid. [α]_D +67.8 (c, 0.27 in MeOH).

Buchanan, M.S. et al., *Phytochemistry*, 1996, **41**, 821 (*isol, pmr, cmr, ms, struct*)

7,9-Dihydroxy-8,10-dimethyl-2,4-tridecadienoic acid D-574

Absolute Configuration

C₁₅H₂₆O₄ 270.368**(2Z,4E,7S,8S,9R,10S)-form**9-O-Cinnamoyl(E-), amide: **Antibiotic**

YM 47522. YL 03709B-A. YM 47522.

Antibiotic YL 03709B-A. TAN 1771A.

Antibiotic TAN 1771A

[171964-61-7]

[182698-42-6, 170034-40-9]


C₂₄H₃₃NO₄ 399.529Prod. by *Bacillus* sp. YL-03709B.

Antifungal agent. Gum. [α]_D²⁵ -106.1 (c, 0.3 in MeOH). λ_{max} 217 (ε 19000); 222 (ε 18000); 263 (ε 31000) (MeOH).

Japan. Pat., 1995, 95 206 797; CA, **123**, 306561u (*isol*)

Sugawara, T. et al., *J. Antibiot.*, 1996, **49**, 340-344; 345-348 (*isol, uv, ir, pmr, cmr, props*)

Ermolenko, M.S. et al., *Tet. Lett.*, 1996, **37**, 6711-6712 (*synth, abs config*)

3,6-Dihydroxy-4,5-diphenyl-2-piperidinone D-575C₁₇H₁₇NO₃ 283.326N-Me: 3,6-Dihydroxy-1-methyl-4,5-diphenyl-2-piperidinone. **Lansamide 3.****Lansimide 3†**

[211504-98-2]

C₁₈H₁₉NO₃ 297.353Constit. of *Clausena lansium* (wampee). Spasmolytic agent. Needles. Mp

236-237°. Racemic. It appears that this is a struct. revision for the compound formerly assigned a pyrrolidinone struct. and isol. from a sp. referred to as *Clausena lansium*. However this is not clear from the paper.



N,O²-Di-Me: 3-Hydroxy-6-methoxy-1-methyl-4,5-diphenyl-2-piperidinone.

Lansamide 4. **Lansimide 4†**

[211505-00-9]


C₁₉H₂₁NO₃ 311.38Constit. of *Clausena lansium* (wampee). Spasmolytic agent. Needles. Mp

187-188°. Racemic.


Lakshmi, V. et al., *Indian J. Chem., Sect. B*, 1998, 37, 422-424**6,11-Dihydroxy-2,7,9-dodecatrienoic acid** D-576(2*E*,6*RS*,7*E*,9*E*,11*RS*)-formC₁₂H₁₈O₄ 226.272**(2*E*,6*RS*,7*E*,9*E*,11*RS*)-form**(2-Hydroxy-2-methylpropyl)amide: **ZP-amide F**C₁₆H₂₇NO₄ 297.394Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 230 (log ε 5) (MeOH).6-Ketone, (2-hydroxy-2-methylpropyl)amide: **ZP-amide B**C₁₆H₂₅NO₄ 295.378Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 271 (log ε 4.66) (MeOH).11-Ketone, (2-hydroxy-2-methylpropyl)amide: **ZP-amide A**C₁₆H₂₅NO₄ 295.378Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 270 (log ε 4.63) (MeOH).**(2*E*,6*RS*,7*E*,9*E*,11*SR*)-form**(2-Hydroxy-2-methylpropyl)amide: **ZP-amide E**C₁₆H₂₇NO₄ 297.394Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 230 (log ε 5.04) (MeOH).Hatano, T. et al., *Phytochemistry*, 2004, 65, 2599-2604 (isol, pmr, cmr)**8,11-Dihydroxy-2,4,9-dodecatrienoic acid** D-577C₁₂H₁₈O₄ 226.272**(2*E*,4*E*,8*ξ*,9*E*,11*ξ*)-form**

2-Methylpropylamide: 8,11-Dihydroxy-N-(2-methylpropyl)-2,4,9-dodecatrienoamide

[1017577-33-1]

C₁₆H₂₇NO₃ 281.394Alkaloid from *Spilanthes scallimorpha*.Li, G.-P. et al., *J. Integ. Plant Biol.*, 2007, 49, 1608-1610 (isol)**10,11-Dihydroxy-2,6,8-dodecatrienoic acid** D-578(2*E*,6*Z*,8*E*,10*R**,11*R**)-formC₁₂H₁₈O₄ 226.272**(2*E*,6*Z*,8*E*,10*R*,11*R*)-form**(2-Hydroxy-2-methylpropyl)amide: **ZP-amide D**C₁₆H₂₇NO₄ 297.394Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 230 (log ε 4.71) (MeOH).**(2*E*,6*Z*,8*E*,10*R*,11*S*)-form**(2-Hydroxy-2-methylpropyl)amide: **ZP-amide C**C₁₆H₂₇NO₄ 297.394Alkaloid from the pericarp of *Zanthoxylum piperitum*. Unstable syrup. λ_{max} 233 (log ε 4.69) (MeOH).Hatano, T. et al., *Phytochemistry*, 2004, 65, 2599-2604 (isol, pmr, cmr)**1-(1,2-Dihydroxyethyl)-β-carboline-3-carboxylic acid** D-579

1-(1,2-Dihydroxyethyl)-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI

C₁₄H₁₂N₂O₄ 272.26**(S)-form****Dichotomine B**

[755036-41-0]

Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. [α]_D²⁷ -19 (c, 1 in MeOH). λ_{max} 237 (log ε 3.91); 269 (log ε 4.04) (MeOH).1'-O-β-D-Glucopyranoside: **Glucodichotomine B**C₂₀H₂₂N₂O₉ 434.402Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. [α]_D²⁷ -28.2 (c, 0.2 in MeOH). λ_{max} 218 (log ε 4.3); 239 (log ε 4.35); 270 (log ε 4.51) (MeOH).Me ester: **Dichotomine C**[†]


[755036-45-4]

C₁₅H₁₄N₂O₄ 286.287Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Antiallergic agent. Yellow powder. [α]_D²⁷ -16.6 (c, 0.5 in MeOH). λ_{max} 283 (log ε 3.56) (MeOH).Butyl ester: **Dichotomine D**[†]

[755036-50-1]

C₁₈H₂₀N₂O₄ 328.367Alkaloid from the roots of *Stellaria dichotoma* var. *lanceolata*. Yellow powder. [α]_D²⁷ -1.8 (c, 0.75 in CHCl₃). λ_{max} 283 (log ε 3.66) (MeOH).Morikawa, T. et al., *Chem. Pharm. Bull.*, 2004, 52, 1194-1199 (*Glucodichotomine B*)Sun, B. et al., *J. Nat. Prod.*, 2004, 67, 1464-1469 (isol, ir, pmr, cmr, ms)Omura, K. et al., *Chem. Pharm. Bull.*, 2008, 56, 237-238 (*Dichotomine C*, synth)**2-(1,2-Dihydroxyethyl)-3,4-dihydroxy-5-(hydroxymethyl)pyrrolidine** D-580

3,4-Dihydroxy-α-(hydroxymethyl)-2,5-pyrrolidinedimethanol, 9CI, 2,5-Dideoxy-2,5-iminoheptitol

(1'*R*,2*S*,3*R*,4*R*,5*R*)-formC₇H₁₅NO₅ 193.199**(1'*R*,2*S*,3*R*,4*R*,5*R*)-form** [173142-29-5] Amorph.**(1'*S*,2*R*,3*R*,4*R*,5*R*)-form** [173142-30-8] Amorph.**(1'*S*,2*R*,3*S*,4*R*,5*R*)-form**

2,5-Dideoxy-2,5-imino-D-glycero-D-taloheptitol [146399-47-5]

Cryst. (MeOH/CHCl₃) (as hydrochloride). Mp 148-149° (hydrochloride). [α]_D²⁰ +26.9 (c, 1 in H₂O).**(1'*S*,2*S*,3*S*,4*S*,5*R*)-form** [196494-58-3]Oil. [α]_D²² -18.2 (c, 1 in H₂O).**(1'*ξ*,2*R*,3*R*,4*R*,5*R*)-form**

2,5-Dideoxy-2,5-imino-glycero-D-mannoheptitol

[197390-30-0]

Alkaloid from *Hyacinthoides non-scripta*, *Hyacinthus orientalis* and *Scilla sibirica*. Inhibitor of β-glucosidases. [α]_D²⁵ +31.5 (c, 0.4 in H₂O). C-1' config. not determined.

2'-O-β-D-Xylopyranoside: [240117-28-6]

C₁₂H₂₃NO₉ 325.315Alkaloid from *Hyacinthoides non-scripta*, *Muscari armeniacum* and *Scilla campanulata*. [α]_D -8.7 (c, 0.94 in H₂O).

2'-O-Apiofuranoside: [197390-31-1]


C₁₂H₂₃NO₉ 325.315Alkaloid from *Hyacinthoides non-scripta*. Not isol. pure.**(1'*ξ*,2*R*,3*S*,4*R*,5*S*)-form**

2,5-Dideoxy-2,5-imino-glycero-D-galactohexitol

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. [α]_D +33.6 (c, 0.19 in H₂O).Myerscough, P.M. et al., *Tetrahedron*, 1992, 48, 10177-10190 (synth, pmr, cmr)Watson, A.A. et al., *Phytochemistry*, 1997, 46, 255-259 (isol, pmr, cmr)Lee, R.E. et al., *Tet. Lett.*, 1997, 38, 6733-6736 (synth, pmr)Asano, N. et al., *J. Nat. Prod.*, 1998, 61, 625-628 (isol)Kato, A. et al., *Carbohydr. Res.*, 1999, 316, 95-103 (2'-xyloside)Takebayashi, M. et al., *J.O.C.*, 1999, 64, 5280-5291 (synth, pmr)Yamashita, T. et al., *J. Nat. Prod.*, 2002, 65, 1875-1881 (*Scilla sibirica* alkaloids)

4',7-Dihydroxyflavanone D-581

2,3-Dihydro-7-hydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one. **Liquiritigenin**



C₁₅H₁₂O₄ 256.257
Log P 2.35 (uncertain value) (calc).

(S)-form [578-86-9]

Isol. from *Centrobium*, *Dalbergia*, *Medicago*, *Umtiza*, *Zollernia*, *Onobrychis*, *Myroxyton*, *Peltogyne*, *Glycyrrhiza*, *Cicer* and *Platymiscium* (all Fabaceae). Also in *Citrus*. Several glycosides, particularly the rutinoid and neohesperidoid, are important in influencing citrus fruit flavour. Antiinflammatory agent. Cryst. Mp 203-205°. [α]_D²⁵ -36.2 (c, 0.09 in MeOH). Pharmacol. active isomer.

4'-O-[1H-Indole-3-ylcarbonyl-(\rightarrow 5)- β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Licorice glycoside E**

C₃₅H₃₅NO₁₄ 693.66


Constit. of *Glycyrrhiza uralensis* (Chinese licorice). Amorph. pale yellow powder. [α]_D -52.9 (c, 1 in MeOH). λ_{\max} 206 (log ϵ 4.68); 223 (log ϵ 4.22); 280 (log ϵ 3.83); 285 (log ϵ 3.8) (MeOH).

[25826-69-1, 32274-71-8]

Hatano, T. et al., *Phytochemistry*, 1998, **47**, 287-293 (*Licorice glycosides*)

Dihydroxygirinimbine D-582

[97400-67-4]



C₁₈H₁₉NO₃ 297.353


Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Needles. Mp 189-190°. [α]_D -4 (MeOH).

Furukawa, H. et al., *Heterocycles*, 1985, **23**, 1391 (*uv, ir, pmr, struct*)

Knölker, H.J. et al., *Tet. Lett.*, 1996, **37**, 7947 (*synth*)

2-(3,6-Dihydroxyheptyl)-3,4-dihydroxy-5-hydroxymethylpyrrolidine D-583

2-(3,6-Dihydroxyheptyl)-5-hydroxymethyl-3,4-pyrrolidinediol, 9CI




C₁₂H₂₅NO₅ 263.333

(2R,3R,3'ε,4R,5R,6'ε)-form [240117-27-5]

Alkaloid from *Hyacinthoides non-scripta* and *Scilla campanulata*. [α]_D +42.8 (c, 0.6 in H₂O).

Kato, A. et al., *Carbohydr. Res.*, 1999, **316**, 95-103

6,13-Dihydroxyhetisan-11-one D-584

C₂₀H₂₅NO₃ 327.422

The C-13 configs. in these alkaloids are ambiguously drawn and described in the lit. Configs. given here are as given in CAS but these may not be reliable.

(13R)-form**Spirasine XIII**

[115610-47-4]


Minor alkaloid from the roots of *Spiraea japonica* var. *fortunei* (Rosaceae). Cryst. (CH₂Cl₂). Mp 188-189°. [α]_D¹⁷ +25.7 (c, 0.93 in CHCl₃).

(13S)-form**Spirasine XII**

[115610-46-3]

Minor alkaloid from roots of *Spiraea japonica* (Rosaceae). Needles (CH₂Cl₂/MeOH). Mp 226-228°. [α]_D¹⁷ +17.9 (c, 1.2 in CHCl₃).

Sun, F. et al., *J. Nat. Prod.*, 1988, **51**, 50-53 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

9,13-Dihydroxyhetisan-2-one D-585

C₂₀H₂₅NO₃ 327.422


(13S)-form**9-Ac: Fissumine**

[152606-54-7]

C₂₂H₂₇NO₄ 369.46

Alkaloid from aerial parts of *Delphinium fissum* ssp. *anatolicum* (Ranunculaceae). Amorph. [α]_D²⁰ -33.8 (c, 0.4 in CHCl₃).

Ulubelen, A. et al., *Phytochemistry*, 1993, **34**, 1165 (*isol, ir, pmr, cmr, ms, struct*)

13,15-Dihydroxyhetisan-2-one D-586

C₂₀H₂₅NO₃ 327.422

(13R,15β)-form**Venusone**


[145237-03-2]

Alkaloid from the aerial parts of *Delphinium venulosum* (Ranunculaceae). [α]_D²² +27.3 (c, 0.20 in MeOH).

Ulubelen, A. et al., *Phytochemistry*, 1992, **31**, 3239 (*isol, ir, pmr, cmr, ms, struct*)

3,4-Dihydroxy-2-(1-hydroxyethyl)-5-(hydroxymethyl)pyrrolidine D-587

3,4-Dihydroxy- α -methyl-2,5-pyrrolidinedimethanol



C₇H₁₅NO₄ 177.2

(1'ξ,2R,3R,4R,5R)-form


2,5,7-Trideoxy-2,5-imino-D-glycero-D-manno-heptitol

Alkaloid from the bulbs of *Scilla sibirica*. Inhibitor of β -galactosidase and β -glucosidase. Solid. [α]_D +22 (c, 0.39 in H₂O).

Yamashita, T. et al., *J. Nat. Prod.*, 2002, **65**, 1875-1881 (*isol, pmr, cmr*)

3,4-Dihydroxy-2-(1-hydroxyethyl)pyrrolidine D-588

2-(1-Hydroxyethyl)-3,4-pyrrolidinediol



C₆H₁₃NO₃ 147.174

(1'R,2R,3R,4S)-form

3-O-[α -D-Glucuronopyranosyl-(1 \rightarrow 2)-3,4-di-O-acetyl- α -L-arabinopyranosyl-(1 \rightarrow 4)]-3-O-methyl- α -D-glucopyranoside]:

C₂₈H₄₅NO₂₀ 715.658


Isol. from the cyanobacterium *Anabaena* sp. Plates (EtOH aq.). Mp 211-213° dec. [α]_D²⁶ +156.2 (c, 1 in MeOH aq.).

Thammana, S. et al., *J. Nat. Prod.*, 2006, **69**, 365-368 (*isol, pmr, cmr, cryst struct*)

3,4-Dihydroxy-3-hydroxy-methylbutanoic acid D-589
(HOCH₂)₂C(OH)CH₂COOH
C₅H₁₀O₅ 150.131

Nitrile, 4-O-β-D-glucopyranoside (3R-):
[148965-97-3]
[148966-04-5]
C₁₁H₁₉NO₈ 293.273
Constit. of barley (*Hordeum vulgare*).
Pourmohseni, H. et al., *Phytochemistry*, 1993,
33, 295-297

6,7-Dihydroxy-1-hydroxy-methylisoquinoline D-590
1-Hydroxymethyl-6,7-isoquinolinediol



C₁₀H₉NO₃ 191.186

7-Me ether: 6-Hydroxy-7-methoxy-1-isoquinolinemethanol. 6-Hydroxy-1-hydroxymethyl-7-methoxyisoquinoline
[367947-68-0]

C₁₁H₁₁NO₃ 205.213

Alkaloid from the sponge *Haliclona* sp.
Needles (EtOH). Mp 202-204°. λ_{max}
239 (log ε 4.35); 253 (sh) (log ε 3.87);
258 (log ε 3.73); 279 (log ε 3.56); 325
(log ε 3.32); 363 (log ε 3.31) (MeOH).


6,7-Di-Me ether, N-oxide: 1-Hydroxy-methyl-6,7-dimethoxyisoquinoline N-oxide
C₁₂H₁₃NO₄ 235.239
Alkaloid from the seeds of *Calycotome villosa* ssp. *intermedia*. Mp 209-211°. λ_{max} 217 ; 259 ; 304 ; 338 ; 352 (MeOH).

Rashid, M.A. et al., *J. Nat. Prod.*, 2001, **64**, 1249-1250 (7-Me ether, isol)

Saito, N. et al., *Chem. Pharm. Bull.*, 2004, **52**, 282-286 (7-Me ether, synth, pmr, cmr, ms)

El Antri, A. et al., *Fitoterapia*, 2004, **75**, 774-778 (6,7-di-Me ether N-oxide)

3,4-Dihydroxy-2-(hydroxy-methyl)-5-methylpyrrolidine D-591
2-(Hydroxymethyl)-5-methyl-3,4-pyrrolidinediol, 9CI



(2R,3R,4R,5R)-form

C₆H₁₃NO₃ 147.174

(2R,3R,4R,5R)-form

1,2,5-Trideoxy-2,5-imino-D-glucitol
[147060-64-8]

Alkaloid from seeds of *Angylocalyx pynaertii*. β-Mannosidase inhibitor.
Wax. [α]_D²³ +26.2 (c, 1.1 in MeOH) (natural). [α]_D²³ +42.9 (c, 0.72 in MeOH) (synthetic).

(2R,3R,4S,5R)-form

1,2,5-Trideoxy-2,5-imino-D-altritol
Alkaloid from the bark of *Angylocalyx pynaertii*. [α]_D +41.8 (c, 0.32 in H₂O).

(2S,3R,4S,5R)-form

2,5-Dideoxy-2,5-imino-D-fucitol. 1,2,5-Trideoxy-2,5-imino-D-galactitol
Alkaloid from the bark and pods of *Angylocalyx pynaertii*. [α]_D -11.1 (c, 0.28 in H₂O).

(2S,3S,4S,5R)-form

1,2,5-Trideoxy-2,5-imino-L-glucitol
Alkaloid from the bark and pods of *Angylocalyx pynaertii*. [α]_D +23.3 (c, 0.64 in H₂O).


Molyneux, R.J. et al., *J. Nat. Prod.*, 1993, **56**, 1356-1364 (isol, synth, pmr, cmr, ms)

Asano, N. et al., *Eur. J. Biochem.*, 2001, **268**, 35-41 (isol, pmr, cmr)

Yasuda, K. et al., *J. Nat. Prod.*, 2002, **65**, 198-202 (isol, pmr, cmr)

Merino, P. et al., *Eur. J. Org. Chem.*, 2008, 2929-2947 (synth)

1,2-Dihydroxy-3-hydroxy-methyl-5-methyl-1H-pyrrolidine D-592
Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizine-1,2-diol, 9CI



(1R,2R,3R,5R,7aR)-form

C₉H₁₇NO₃ 187.238

(1R,2R,3R,5R,7aR)-form

Hyacinthacine A₃

[268209-90-1]

Alkaloid from the bulbs of *Muscari armeniacum*. [α]_D +19.2 (c, 0.43 in H₂O).

(1R,2S,3R,5R,7aR)-form

Hyacinthacine A₅

[479348-16-8]

Alkaloid from the bulbs of *Scilla sibirica*. Powder. [α]_D -39.4 (c, 0.3 in H₂O).

(1S,2R,3R,5R,7aR)-form

Hyacinthacine A₆

[479348-17-9]

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. [α]_D +16.3 (c, 0.22 in H₂O).

(1S,2R,3S,5S,7aR)-form

Hyacinthacine A₄

[479348-15-7]

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. [α]_D -49.5 (c, 0.48 in H₂O).

(1S,2R,3R,5S,7aR)-form

Hyacinthacine A₇

[479348-18-0]

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. [α]_D -51.8 (c, 0.45 in H₂O).

Asano, N. et al., *Tetrahedron: Asymmetry*, 2000, **11**, 1-8 (isol, pmr, cmr)


Yamashita, T. et al., *J. Nat. Prod.*, 2002, **65**, 1875-1881 (isol, pmr, cmr)

Izquierdo, I. et al., *Eur. J. Org. Chem.*, 2007, 6078-6083 (synth, abs config)

Izquierdo, I. et al., *Tetrahedron*, 2008, **64**, 4613-4618 (synth, abs config)

3,4-Dihydroxy-2-(hydroxy-methyl)pyrrolidine D-593

1,4-Dideoxy-1,4-iminopentitol, 11CI. 2-(Hydroxymethyl)-3,4-pyrrolidinediol, 12CI. DAB-1
[98632-49-6]



(2R,3R,4R)-form

C₅H₁₁NO₃ 133.147

(2R,3R,4R)-form

1,4-Dideoxy-1,4-imino-D-arabinitol

[100937-52-8]

Isol. from *Arachniodes standishii* and *Angylocalyx* spp. α-Glucosidase inhibitor. Hygroscopic oil. [α]_D²⁰ +7.8 (c, 0.46 in H₂O).

Hydrochloride: [100991-92-2]

Cryst. (Me₂CO aq.) or solid (MeOH/Et₂O). Mp 113-115° (110-112°). [α]_D²⁰ +37.9 (c, 0.53 in H₂O) (+31.1).

1'-O-β-D-Glucopyranoside: 1,4-Dideoxy-1,4-imino-5-O-β-D-glucopyranosyl-D-arabinitol

C₁₁H₂₁NO₈ 295.289

Alkaloid from the pods of *Angylocalyx pynaertii*. [α]_D +40.1 (c, 0.16 in H₂O).

4-O-β-D-Glucopyranoside:

C₁₁H₂₁NO₈ 295.289

Alkaloid from leaves of *Morus bombycis* (Moraceae). Powder. [α]_D -29.8 (c, 0.73 in H₂O).

N-(2-Hydroxyethyl): 3,4-Dihydroxy-1-(2-hydroxyethyl)-2-(hydroxymethyl)-pyrrolidine

C₇H₁₅NO₄ 177.2

Alkaloid from the pods of *Angylocalyx pynaertii*. [α]_D +54.7 (c, 0.38 in H₂O).

N-(3-Amino-3-oxopropyl): 3,4-Dihydroxy-2-hydroxymethyl-1-pyrrolidine-propanamide

C₈H₁₆N₂O₄ 204.225

Alkaloid from *Morus alba* (white mulberry). [α]_D -53.7 (c, 0.41 in H₂O).

N¹,2,3,4-Tetrabenzyl: 2,3,5-Tri-O-benzyl-1,4-benzylimino-1,4-dideoxy-D-arabinitol

C₃₃H₃₅NO₃ 493.644

[α]_D +25.8 (c, 2.0 in CHCl₃).

(2R,3R,4S)-form

1,4-Dideoxy-1,4-imino-D-ribitol

[105990-41-8]

Alkaloid from roots of *Morus alba* (white mulberry) (Moraceae). Solid. [α]_D +42 (c, 0.53 in H₂O).

Hydrochloride: [117781-12-1]

Cryst. Mp 128-132°. [α]_D²⁰ +57.6 (c, 0.5 in H₂O).

(2R,3S,4R)-form

1,4-Dideoxy-1,4-imino-D-lyxitol

[100937-51-7]

α-Galactosidase inhibitor. [α]_D²⁰ -15.8 (c, 0.14 in H₂O).

Hydrochloride: [100991-93-3]

Cryst. Mp 159-161°. [α]_D²⁰ +19.8 (c, 0.45 in H₂O).

(2R,3S,4S)-form*1,4-Dideoxy-1,4-imino-D-xylitol*

[97058-12-3]

Alkaloid from the bark of *Angylocalyx pynaerti*.*Hydrochloride*: [101399-04-6]Cryst. (EtOH/Me₂CO). Mp 127-129°.[α]_D²⁰ +9.8 (c, 0.6 in H₂O).**(2S,3R,4R)-form***1,4-Dideoxy-1,4-imino-L-xylitol, 2,5-Dideoxy-2,5-imino-D-xylitol*

[105990-42-9]

Oil. [α]_D³¹ -4.1 (c, 1.0 in H₂O).*Hydrochloride*: [114488-33-4]Cryst. (EtOH/Me₂CO). Mp 128-129°.[α]_D²⁰ -9.9 (c, 0.71 in H₂O).*N*¹,2,3,4-Tetrabenzyl: 2,3,5-Tri-O-benzyl-*1,4-benzylimino-1,4-dideoxy-L-xylitol*

[134307-35-0]

C₃₃H₃₅NO₃ 493.644[α]_D²⁰ +28.7 (c, 2.0 in CHCl₃). [α]_D²²+30.5 (c, 0.95 in CHCl₃).**(2S,3R,4S)-form***1,4-Dideoxy-1,4-imino-L-lyxitol*

[129568-47-4]

Hydrochloride: [129568-48-5]Cryst. Mp 155-156°. [α]_D²⁰ -18.4 (c, 0.22in H₂O).**(2S,3S,4R)-form***1,4-Dideoxy-1,4-imino-L-ribitol*


[105990-40-7]

Pale yellow solid. Mp 119-120°. [α]_D²⁰ -53.9 (c, 0.6 in H₂O).*Hydrochloride*: [117770-01-1]Cryst. Mp 126-131°. [α]_D²⁰ -59 (c, 0.59in H₂O).**(2S,3S,4S)-form***1,4-Dideoxy-1,4-imino-L-arabinitol*

[100937-53-9]

[α]_D²⁰ -12 (c, 0.21 in MeOH).*Hydrochloride*: [100991-91-1]Cryst. (Me₂CO aq.). Mp 107-111°.[α]_D²⁵ -34.6 (c, 0.37 in H₂O).Furukawa, J. *et al.*, *Phytochemistry*, 1985, **24**,593 (*isol, pmr, cmr, ms, ord, config*)Nash, R.J. *et al.*, *Phytochemistry*, 1985, **24**,1620 (*isol, pmr, cmr, ms, struct*)*Japan. Pat.*, 1986, 86 118 360 (*1,4-Dideoxy-1,4-**imino-D-arabinitol*)Fleet, G.W.J. *et al.*, *Tetrahedron*, 1986, **42**,5685; 1987, **43**, 3083; 1988, **44**, 2637; 2649(*synth, ir, pmr*)Setol, H. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**,3995 (*synth, ir, pmr*)Ikota, N. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**,1087; 3399 (*synth*)Wehmer, V. *et al.*, *Angew. Chem., Int. Ed.*,1990, **29**, 1169 (*synth*)Duréault, A. *et al.*, *J. Carbohydr. Chem.*, 1990,**9**, 121-123 (*2S,3R,4S-form, synth*)Buchanan, J.G. *et al.*, *J.C.S. Perkin 1*, 1990,699 (*synth*)Meng, Q. *et al.*, *Helv. Chim. Acta*, 1991, **74**,445-450 (*2S,3R,4R-form, tetrabenzyl, synth,**pmr*)Van der Klein, P.A.M. *et al.*, *Synth. Commun.*,1992, **22**, 1763 (*synth*)Asano, N. *et al.*, *Carbohydr. Res.*, 1994, **253**,235; **259**, 243 (*1,4-Dideoxy-1,4-imino-2-O-β-**D-glucopyranosyl-D-arabinitol, 1,4-Dideoxy-**1,4-imino-D-ribitol*)Griffart-Brunet, D. *et al.*, *Tet. Lett.*, 1994, **35**,2889 (*synth*)Blanco, M.-J. *et al.*, *J.O.C.*, 1996, **61**, 4748(*2R,3S,4R-form, synth, ir, pmr, cmr*)Huang, Y. *et al.*, *J.O.C.*, 1997, **62**, 372-376(*synth, pmr, ir, cmr*)Watson, A.A. *et al.*, *Phytochemistry*, 1997, **46**,255 (*isol, props*)Huwe, C.M. *et al.*, *Synthesis*, 1997, 61 (*synth,**pmr, cmr*)Hulme, A.N. *et al.*, *J.C.S. Perkin 1*, 2000,1837-1841 (*2R,3R,4R-form, synth, pmr, cmr*)Lee, B.W. *et al.*, *Synthesis*, 2000, 1305-1309(*2S,3R,4R-form*)Asano, N. *et al.*, *Eur. J. Biochem.*, 2001, **268**,35-41 (*1,4-Dideoxy-1,4-imino-D-xylitol, isol*)Asano, N. *et al.*, *J. Agric. Food Chem.*, 2001,**49**, 4208-4213 (*N-propanamide*)Lombardo, M. *et al.*, *J.O.C.*, 2001, **66**, 1264-1268 (*1,4-Dideoxy-1,4-imino-L-arabinitol*)Yasuda, K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 198-202 (*1,4-Dideoxy-1,4-imino-5-**glucopyranosylarabinitol, 1-hydroxyethyl**deriv*)El-Ashry, E.H. *et al.*, *Carbohydr. Res.*, 2003,**338**, 2265-2290 (*rev. synth*)Ayad, T. *et al.*, *Eur. J. Org. Chem.*, 2003, 2903-2910 (*1,4-dideoxy-1,4-imino-D-arabinitol*)Koh, D.W. *et al.*, *J. Med. Chem.*, 2003, **46**,4322-4332 (*2R,3R,4S-form, synth, pmr, cmr*)Dhavale, D.D. *et al.*, *Org. Biomol. Chem.*,2005, **3**, 3720-3726 (*2S,3R,4R-form*)Lauritsen, A. *et al.*, *Org. Biomol. Chem.*, 2006,**4**, 2898-2905 (*2R,3R,4R-form, 2S,3R,4R-**form, synth, pmr, cmr, ms, tetrabenzyl*)Kim, I.S. *et al.*, *Org. Lett.*, 2006, **8**, 4101-4104(*synth*)Merino, P. *et al.*, *Eur. J. Org. Chem.*, 2008,2929-2947 (*synth*)**1,2-Dihydroxy-3-hydroxy-****methyl-1H-pyrrolizidine**

D-594

*Hexahydro-3-hydroxymethyl-1H-pyrroli-**zine-1,2-diol, 9CI*

(1R,2R,3R,7aR)-form

C₈H₁₅NO₃ 173.211**(1R,2R,3R,7aR)-form*****Hyacinthacine A₂***


[268209-89-8]

Alkaloid from *Muscari armeniacum*. [α]_D+20.1 (c, 0.44 in H₂O).**(1S,2R,3R,7aR)-form*****Hyacinthacine A₁***

[268209-88-7]

Alkaloid from *Muscari armeniacum*.Glycosidase inhibitor. [α]_D +38.2 (c, 0.23in H₂O).Asano, N. *et al.*, *Tetrahedron: Asymmetry*,2000, **11**, 1-8 (*isol, pmr, cmr*)Cardona, F. *et al.*, *Tet. Lett.*, 2003, **44**, 2315-2318 (*Hyacinthacine A₂, synth*)Desvergnès, S. *et al.*, *J.O.C.*, 2005, **70**, 1459-1462 (*Hyacinthacine A₂, synth*)Donohoe, T.J. *et al.*, *J.O.C.*, 2005, **70**, 7297-7304 (*Hyacinthacine A₁, synth*)Chabaud, L. *et al.*, *Org. Lett.*, 2005, **7**, 2587-2590 (*Hyacinthacine A₁, synth*)Dewi-Wülfing, P. *et al.*, *Eur. J. Org. Chem.*,2006, 1852-1856 (*Hyacinthacine A₂, synth*)Izquierdo, I. *et al.*, *Eur. J. Org. Chem.*, 2007,6078-6083 (*Hyacinthacine A₁, synth*)Chandrasekhar, S. *et al.*, *J.O.C.*, 2008, **73**,7826-7828 (*Hyacinthacine A₁, synth*)Reddy, P.V. *et al.*, *Org. Biomol. Chem.*, 2008, **6**,1170-1172 (*Hyacinthacine A₁, synth*)**1,2-Dihydroxy-7-hydroxy-****methylpyrrolizidine**


D-595

*Hexahydro-6,7-dihydroxy-1H-pyrrolizine-**1-methanol, 7-Hydroxymethyl-1,2-pyrroli-**zidinediol*C₈H₁₅NO₃ 173.211**(1R*,2R*,7R*,7aR*)-form***2-Angeloyl: 6-Angeloyloxyplatynecine*C₁₃H₂₁NO₄ 255.313Alkaloid from *Senecio nemorensis*.

Yellow oil.

Christov, V. *et al.*, *Nat. Prod. Res.*, 2005, **19**,389-392 (*isol, pmr, cmr, ms*)**1,7-Dihydroxy-1-hydroxy-****methylpyrrolizidine**

D-596

*1-Hydroxymethyl-1,7-pyrrolizidinediol.**Hexahydro-1,7-dihydroxy-1H-pyrrolizine-**1-methanol*

(1S,7R,7aS)-form

C₈H₁₅NO₃ 173.211**(1S,7R,7aS)-form*****Helibractinecine***

[156475-63-7]

Alkaloid from aerial parts of *Heliotrop-**ium bracteatum* (Boraginaceae). Gum.[α]_D²⁰ -18.68 (c, 0.67 in EtOH).*O*⁷-Angeloyl: ***Heliscabine***C₁₃H₂₁NO₄ 255.313Alkaloid from *Heliotropium scabrum*(Boraginaceae). Gum. [α]_D²⁵ -19.5 (c,

0.1 in EtOH).

(1R,7S,7aS)-form***Helibracteatinecine***Necine base obt. by hydrol. of *Helibrac-**teatine* and *Helibracteatine*. Gum. [α]_{Hg}²⁵

+12.1 (c, 0.6 in EtOH).

*O*⁷-Angeloyl: ***Helibracteatinine***

[169626-42-0]

C₁₃H₂₁NO₄ 255.313From *Heliotropium bracteatum* (Bora-*ginaceae*). Pale yellow gum. [α]_{Hg}²⁵

+10.85 (c, 0.64 in EtOH).

*O*⁷-Angeloyl: ***Helibracteatine***

[169869-88-9]

C₁₃H₂₁NO₄ 255.313From *Heliotropium bracteatum* (Bora-*ginaceae*). Brown gum. [α]_{Hg}²⁵ +3.64 (c,

0.33 in EtOH).

(1R,7R,7aR)-form


N-Me, O⁷, O^{1'}-diangeloyl: **7,9-Diangeloyl-1-hydroxy-N-methylplatynecium (1+)** [135574-54-8]
C₁₉H₃₀NO₅⁺ 352.45
Alkaloid from *Senecio integrifolius* var. *fauriri* (Asteraceae). Fine needles (Me₂CO/EtOAc) (as chloride). Mp 143° (chloride). [α]_D²⁵ -60.6 (c, 1 in EtOH).

Roeder, E. *et al.*, *Phytochemistry*, 1991, **30**, 1734 (7,9-Diangeloyl-1-hydroxy-N-methylplatynecinium)

Lakshmanan, A.J. *et al.*, *Phytochemistry*, 1994, **36**, 245; 1995, **39**, 473; **40**, 291 (*Helibracteatine*, *Helibracteatinine*)

2,7-Dihydroxy-1-hydroxymethylpyrrolizidine D-597

Hexahydro-7-(hydroxymethyl)-1H-pyrrolizine-1,6-diol, 9CI. 7-Hydroxymethyl-1,6-pyrrolizidinediol



(1*S*,2*R*,7*R*,8*R*)-form

C₈H₁₅NO₃ 173.211

(1*S*,2*R*,7*R*,8*R*)-form

Rosmarinicine

[520-61-6]

Necine base from the alkaloid Rosmarinicine, R-133. Stout prisms (Py). Mp 171-172°. [α]_D²⁵ -118.5.

Picrate: Mp 175°.

Tri-Ac:

Needles (EtOH)(as picrate). Mp 138-139.5° (picrate).

(1*R*,2*S*,7*R*,8*R*)-form

Croalbinecine

[41690-66-8]

Necine base from the alkaloid Croalbinecine, C-761. Gum.

Hydrochloride:

Cryst. (EtOH/Me₂CO). Mp 165-166°.

Picrate:

Cryst. (EtOAc/CHCl₃). Mp 155-156°.

O⁹-Angeloyl: *Helifoline*

[80787-56-0]

C₁₃H₂₁NO₄ 255.313

Alkaloid from *Heliotropium ovalifolium* (Boraginaceae). Cubes (Me₂CO). Mp 131-132°. [α]_D²⁵ +25.4 (c, 0.0006 in EtOH).

Richardson, M.F. *et al.*, *J.C.S.*, 1943, 452 (*synth*, *struct*)

Dry, L.J. *et al.*, *J.C.S.*, 1955, 59 (*struct*)

Sawney, R.S. *et al.*, *Indian J. Chem.*, 1973, **11**, 88 (*isol*, *pmr*, *ms*)

Sawney, R.S. *et al.*, *Aust. J. Chem.*, 1974, **27**, 1805 (*struct*, *pmr*)

Mohanraj, S. *et al.*, *Phytochemistry*, 1981, **20**, 1991 (*Helifoline*)

Rüeger, H. *et al.*, *Heterocycles*, 1983, **20**, 1331 (*synth*)

Tatsuta, K. *et al.*, *J.A.C.S.*, 1983, **105**, 3653 (*synth*)

Goti, A. *et al.*, *Eur. J. Org. Chem.*, 2000, 3633-3645 (*synth*)


Ahn, J.-B. *et al.*, *J.O.C.*, 2000, **65**, 9249-9251 (*synth*)

Goti, A. *et al.*, *Org. Lett.*, 2001, **3**, 1367-1369 (*synth*)

Akai, S. *et al.*, *Chem. Comm.*, 2005, 2369-2371 (*synth*)

5,6-Dihydroxy-1*H*-indole-2-carboxylic acid, 9CI D-598

[4790-08-3]



C₉H₇NO₄ 193.159

Degradn. prod. of melanin. Likely precursor in biosynth. of eumelanins. Cryst. Mp 240°.

Et ester: [113370-04-0]

C₁₁H₁₁NO₄ 221.212

Yellow prisms (EtOH). Mp 172°.

Di-Me ether: 5,6-Dimethoxy-1*H*-indole-2-carboxylic acid

[88210-96-2]

C₁₁H₁₁NO₄ 221.212

Plates (C₆H₆). Mp 202-203° dec.

H₂SO₄ → orange soln.

Di-Me ether, *Me ester*: [28059-24-7]

C₁₂H₁₃NO₄ 235.239

Cryst. Mp 167-168°.

Perkin, W.H. *et al.*, *J.C.S.*, 1926, 360

Raper, H.S. *et al.*, *Biochem. J.*, 1927, **21**, 89

Oxford, A.E. *et al.*, *J.C.S.*, 1927, 420

Benigni, J.D. *et al.*, *J. Het. Chem.*, 1965, **2**, 387 (*synth*, *uv*)

d'Ischia, M. *et al.*, *Tet. Lett.*, 1985, **26**, 2801 (*isol*)


Chioccare, F. *et al.*, *Synth. Commun.*, 1987, **17**, 1815 (*synth*)

Palumbo, P. *et al.*, *Tetrahedron*, 1987, **43**, 4203 (*biochem*)

Coowar, D. *et al.*, *J. Med. Chem.*, 2004, **47**, 6270-6282 (*di-Me ether* *Me ester*)

6,7-Dihydroxy-3-isocyano-2*H*-1-benzopyran-2-one D-599

6,7-Dihydroxy-3-isocyanocoumarin [957062-50-9]




C₁₀H₅NO₄ 203.154

Prod. by *Pseudomonas aeruginosa*. Cryst.

Brady, S.F. *et al.*, *J.A.C.S.*, 2007, **129**, 12101-12102

6,7-Dihydroxy-1,3,4(2*H*)-isoquinolinetriene D-600



C₉H₅NO₅ 207.142


Di-Me ether, *N-Me*: 6,7-Dimethoxy-2-methyl-1,3,4(2*H*)-isoquinolinetriene

C₁₂H₁₁NO₅ 249.223

Alkaloid from the roots of *Menispermum dauricum*. Yellow powder. Mp 185-189°. λ_{max} 211 (log ε 1.85); 228 (log ε 1.38); 256 (log ε 1.3); 363 (log ε 0.26) (no solvent reported).

Zhang, X. *et al.*, *Phytochemistry*, 2004, **65**, 929-932 (*isol*, *pmr*, *cmr*, *ms*)

6,9-Dihydroxy-13-labdene-16,15-lactam D-601



C₂₀H₃₃NO₃ 335.486

(6β,8α,9β)-form

6-*Ac*: *Vitexlactam A*


[459167-05-6]

C₂₂H₃₅NO₄ 377.523

Alkaloid from the fruit of *Vitex agnus-castus*. Cryst. Mp 207°. [α]_D²⁴ -10.7 (c, 0.42 in CHCl₃).

Li, S.-H. *et al.*, *Tet. Lett.*, 2002, **43**, 5131-5134 (*isol*, *pmr*, *cmr*, *ms*, *ir*, *cryst struct*)

3,18-Dihydroxy-8,22,24-trien-26-oic acid D-602



C₃₀H₄₆O₄ 470.691

(3β,22*E*,24*E*)-form

N-(*Tetrahydro-4,5-dimethyl-2-oxo-3-furanyl*)amide, 3-*O*-[α-*L*-rhamnopyranosyl-(1→2)-β-*D*-glucopyranosyl-(1→2)-[α-*L*-rhamnopyranosyl-(1→4)]-β-*D*-glucopyranoside]: *Mussaendoside Q* [161407-77-8]


C₆₀H₉₅NO₂₃ 1198.403

Constit. of *Mussaenda pubescens*.

Amorph. powder. [α]_D¹⁴ +25 (c, 0.14 in CHCl₃).

Zhao, W. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1613-1618 (*isol*, *pmr*, *cmr*)


3,15-Dihydroxy-5,12-lathyradien-14-one D-603



C₂₀H₃₀O₃ 318.455

(2β,3β,5E,12E,15β)-form

5α,6α-Epoxyde, 3-(3-pyridinecarbonyl),
15-Ac: [88202-64-6]
C₂₈H₃₅NO₆ 481.588
Constit. of *Euphorbia characias*.
Seip, E.H. et al., *Phytochemistry*, 1983, **22**,
1791-1795 (*Euphorbia characias* constits)


3,4-Dihydroxylupanine**D-604**C₁₅H₂₄N₂O₃ 280.366**(3β,4α)-form****Lebeckianine**

[125263-79-8]
Alkaloid from *Lebeckia* spp. (Fabaceae).
Pale-brown oil. [α]_D¹⁸ +61 (c, 0.82 in
CHCl₃).

4-Angeloyl: Sessilifoline

[126381-87-1]
C₂₀H₃₀N₂O₄ 362.468
Alkaloid from *Pearsonia cajanifolia*
ssp. *cryptantha* (Fabaceae). [α]_D²² -72 (c,
1.6 in CHCl₃).

Van Wyk, B.E. et al., *Biochem. Syst. Ecol.*,
1989, **17**, 225-229 (*Lebeckianine*)
Verdoorn, G.H. et al., *Phytochemistry*, 1990,
29, 1297-1302 (*Sessilifoline*)

3,13-Dihydroxylupanine**D-605**C₁₅H₂₄N₂O₃ 280.366**(3β,13α)-form**

Alkaloid from the seeds of *Cytisus*
scoparius (Fabaceae). Needles (CH₂Cl₂/
hexane). Mp 194-196°. [α]_D¹⁵ -7.4 (c, 0.59
in EtOH).

13-Angeloyl: Cajanifoline


[126381-86-0]
C₂₀H₃₀N₂O₄ 362.468
Alkaloid from *Pearsonia cajanifolia*
ssp. *cryptantha* (Fabaceae). Mp 87-91°.
[α]_D²² -11 (c, 1.4 in CHCl₃).

13-Tigloyl: 3-Hydroxy-13-tigloyloxylupanine

[156755-38-3]
C₂₀H₃₀N₂O₄ 362.468
Alkaloid from seedlings of *Cytisus*
scoparius (Fabaceae). Amorph. solid.
[α]_D -8.7 (c, 0.013 in MeOH).

Murakoshi, I. et al., *Phytochemistry*, 1986, **25**,
521 (*isol, pmr, cmr, ms, struct, abs config*)
Verdoorn, G.H. et al., *Phytochemistry*, 1990,
29, 1297 (*Cajanifoline*)

Saito, K. et al., *Phytochemistry*, 1994, **36**, 309
(*3β-Hydroxy-13α-tigloyloxylupanine*)

4,13-Dihydroxylupanine**D-606****(4α,13β)-form**C₁₅H₂₄N₂O₃ 280.366**(4α,13β)-form****O¹³-Me: 4-Hydroxy-13-methoxylupanine**

C₁₆H₂₆N₂O₃ 294.393
Minor alkaloid from bark of *Acos-*
mium panamense.

(4β,13α)-form

Minor alkaloid from the leaves of *Cal-*
purnia aurea ssp. *aurea* (Fabaceae).

O¹³-(2-Pyrrolocarbonyl): Digitine

[105798-93-4]
C₂₀H₂₇N₃O₄ 373.451
Minor alkaloid from the leaves of
Calpurnia aurea ssp. *aurea* (Fabaceae).

(4ξ,13ξ)-form [81149-31-7]

Minor alkaloid from seeds of *Lupinus*
mutabilis and seeds of *Sarothamnus*
scoparius (Fabaceae). Tentative struct.

13-Angeloyl: 13-(Angeloyloxy)-4-hydroxylupanine

[86632-27-1]
C₂₀H₃₀N₂O₄ 362.468
Minor alkaloid from seeds of *Lupinus*
mutabilis (Fabaceae).

O¹³-(3,4-Dimethoxybenzoyl): Catalauverine

C₂₄H₃₂N₂O₆ 444.527
Alkaloid from *Sarothamnus catalauni-*
cus (Fabaceae). Mp 154°. [α]_D²⁵ +328 (c,
0.5 in MeOH). Tentative struct.

O¹³-(3,4,5-Trimethoxybenzoyl): Catalaudesmine

C₂₅H₃₄N₂O₇ 474.553
Alkaloid from *Sarothamnus catalauni-*
cus (Fabaceae). Mp 176°. [α]_D²² +320 (c,
0.3 in MeOH). Tentative struct.

Faugeras, G. et al., *An. Chim.*, 1972, **68**, 811
(*Catalauverine, Catalaudesmine*)


Wink, M. et al., *Planta Med.*, 1981, **43**, 342
(*occur, ms*)

Hatzold, T. et al., *J. Agric. Food Chem.*, 1983,
31, 934 (*isol*)

Asres, K. et al., *Planta Med.*, 1986, 302 (*isol, ir,*
pmr, ms, cd, struct,

Dihydroxylupanine, Digitine)

Veitch, N.C. et al., *Phytochemistry*, 1997, **45**,
847 (*4-Hydroxy-13-methoxylupanine*)

8,13-Dihydroxylupanine**D-607**C₁₅H₂₄N₂O₃ 280.366**(8α,13α)-form [138680-23-6]**

Alkaloid from *Pearsonia sessilifolia* ssp.
marginata (Fabaceae). [α]_D²² +29 (c, 2.4 in
CHCl₃).

13-Angeloyl: Cryptanthine


[129349-79-7]
C₂₀H₃₀N₂O₄ 362.468
Alkaloid from *Pearsonia cajanifolia*
ssp. *cryptantha* (Fabaceae). Mp 108-
110°. [α]_D²³ +117 (c, 1.4 in CHCl₃).

Verdoorn, G.H. et al., *Phytochemistry*, 1990,
29, 1297 (*Cryptanthine*)

Verdoorn, G.H. et al., *Phytochemistry*, 1991,
30, 3631 (*isol*)

10,13-Dihydroxylupanine**D-608**

[61166-47-0]

Relative
configurationC₁₅H₂₄N₂O₃ 280.366

Alkaloid from *Cadia purpurea* and *Cal-*
purnia aurea (Fabaceae).

O¹³-(2-Pyrrolocarbonyl): 10-Hydroxy-13-(2-pyrrolocarbonyloxy)lupanine

[61166-48-1]


C₂₀H₂₇N₃O₄ 373.451Alkaloid from *Calpurnia purpurea*.

van Eijk, J.L. et al., *Tet. Lett.*, 1976, **17**, 2053-
2054 (*isol, ms*)

Radema, M.H. et al., *Phytochemistry*, 1979,
18, 2063-2064 (*isol*)

3-(2,6-Dihydroxy-4-methoxybenzoyl)pyridine**D-609**

(2,6-Dihydroxy-4-methoxyphenyl)-3-pyri-
dylmethanone, 9CI. **Duckeine**
[55051-81-5]


C₁₃H₁₁NO₄ 245.234

Alkaloid from the trunk wood of *Aniba*
duckei (Lauraceae). Yellow cryst.
(EtOH). Mp 243-245°.

Correa, D. de B. et al., *Phytochemistry*, 1975,
14, 271 (*isol, uv, ir, pmr, ms, struct*)

5,7-Dihydroxy-2-[[1-(4-methoxy-2-oxo-2H-pyran-6-yl)-2-phenylethyl]amino]-1,4-naphthoquinone**D-610**

[1021857-44-2]


C₂₄H₁₉NO₇ 433.417Prod. by a marine-based *Aspergillus niger*

EN-13. Amorph. red powder. Mp 118–120°. $[\alpha]_D^{25} +54.9$ (c, 0.19 in CHCl_3). λ_{max} 203 (log ϵ 3.94); 239 (log ϵ 3.08); 266 (log ϵ 3.05); 477 (log ϵ 1.77) (CHCl_3).

Zhang, Y. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 951-953 (*isol*, *pmr*, *cmr*)

7,18-Dihydroxy-4-methylaconit-8(15)-en-14-one D-611

7-Hydroxy-4-(hydroxymethyl)aconit-8(15)-en-14-one



$\text{C}_{19}\text{H}_{25}\text{NO}_3$ 315.411

5 β -form

O⁷-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Talassicumine B**
[160791-10-6]


$\text{C}_{31}\text{H}_{38}\text{N}_2\text{O}_5$ 518.652

Alkaloid from roots of *Aconitum talassicum* (Ranunculaceae). Amorph. powder.

Yue, J. *et al.*, *Phytochemistry*, 1994, **37**, 1467 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

1,8-Dihydroxy-3-methylbenzo[b]phenanthridine-7,12-dione, 9CI D-612

[124903-84-0]



$\text{C}_{18}\text{H}_{11}\text{NO}_4$ 305.289

Prod. by *Streptomyces viridochromogenes*. Antitumour agent. Mp 234-235°.

1-O-(2,3,6-Trideoxy-3-methylamino- α -D-ribo-hexopyranoside): **Phenanthroviridin**
[124903-83-9]


$\text{C}_{25}\text{H}_{24}\text{N}_2\text{O}_6$ 448.474

Prod. by *Streptomyces viridochromogenes*. Antitumour agent. Mp 196–198°. $[\alpha]_D^{20} +62$ (c, 0.518 in MeOH).

Eur. Pat., 1989, 304 400; *CA*, **112**, 75295 (*isol*)
Gore, M.P. *et al.*, *J.O.C.*, 1991, **56**, 2289 (*synth*)
De Frutos, O. *et al.*, *Eur. J. Org. Chem.*, 2001, 163-171 (*synth*)

1,6-Dihydroxy-3-methyl-9H-carbazole D-613

3-Methyl-9H-carbazole-1,6-diol



$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235

6-Me ether: 1-Hydroxy-6-methoxy-3-methyl-9H-carbazole. **Clausenof**
[168434-21-7]

$\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262

Alkaloid from stem bark of *Clausena anisata* (Rutaceae). Highly active against gram-positive and gram-negative bacteria and fungi. Cryst. (C_6H_6). Mp 139°. λ_{max} 228 (ϵ 20000); 253 (ϵ 23000); 303 (ϵ 10900); 356 (ϵ 4000) (EtOH) (Berdy).

6-Me ether, O-Ac: [168293-30-9]

$\text{C}_{16}\text{H}_{15}\text{NO}_3$ 269.299

Needles (C_6H_6 /petrol). Mp 132°.

Di-Me ether: 1,6-Dimethoxy-3-methyl-9H-carbazole. **Clausenine**
[168434-20-6]

$\text{C}_{15}\text{H}_{15}\text{NO}_2$ 241.289

From stem bark of *Clausena anisata* (Rutaceae). Shows low activity against gram-negative bacteria and fungi. Cryst. (C_6H_6 /petrol). Mp 151°. λ_{max} 226 (ϵ 22000); 242 (ϵ 21000); 299 (ϵ 10000); 340 (ϵ 2760); 354 (ϵ 2630) (EtOH) (Berdy).

Chakraborty, A. *et al.*, *Phytochemistry*, 1995, **40**, 295 (*Clausenol*, *Clausenine*, *isol*, 6-Me ether Ac, *uv*, *ir*, *pmr*, *cmr*, *synth*, *cryst struct*)
Bernal, P. *et al.*, *Synthesis*, 2007, 1943-1948 (*Clausenine*, *synth*)

1,7-Dihydroxy-6-methyl-9H-carbazole D-614

6-Methyl-9H-carbazole-1,7-diol

$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235

Di-Me ether: 1,7-Dimethoxy-6-methyl-9H-carbazole. **Clausine P**
[82463-77-2]

$\text{C}_{15}\text{H}_{15}\text{NO}_2$ 241.289

Alkaloid from *Clausena excavata*. Yellow oil. λ_{max} 212; 236 (sh); 239; 251 (sh); 261 (sh); 290 (sh); 298; 317; 330 (MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1999, **52**, 523-527 (*di-Me ether*, *isol*, *uv*, *pmr*, *ms*)

2,3-Dihydroxy-6-methyl-9H-carbazole D-615

6-Methyl-9H-carbazole-2,3-diol

$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235

Methylene ether: 6-Methyl-2,3-methylenedioxy-carbazole. **Clausenalene**
[155961-89-0]

$\text{C}_{14}\text{H}_{11}\text{NO}_2$ 225.246

Alkaloid from the stem bark of *Clausena heptaphylla* (Rutaceae). Needles (C_6H_6 /petrol). Mp 224-225° dec. λ_{max} 236 (ϵ 40000); 312 (ϵ 26200); 336 (ϵ 12000); 353 (ϵ 13800) (EtOH) (Berdy).

Bhattacharyya, P. *et al.*, *Phytochemistry*, 1993, **33**, 248 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *synth*, *struct*)

2,5-Dihydroxy-3-methyl-9H-carbazole D-616

3-Methyl-9H-carbazole-2,5-diol, 9CI

$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235

5-Me ether: 2-Hydroxy-5-methoxy-3-methyl-9H-carbazole. **Carbalexin A**
[371159-83-0]

$\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262

Stress metab. of *Glycosmis parviflora* and *Glycosmis pentaphylla*. Cryst. (CHCl_3). Mp 178-180°. λ_{max} 239; 293; 317; 330 (MeOH).

Di-Me ether: 2,5-Dimethoxy-3-methyl-9H-carbazole. **Glybomine A**

$\text{C}_{15}\text{H}_{15}\text{NO}_2$ 241.289

Alkaloid from the stems of *Glycosmis arborea*. Oil. λ_{max} 209; 239; 264 (sh); 294; 316; 330 (MeOH).

Pacher, T. *et al.*, *Phytochemistry*, 2001, **58**, 129-135 (*Carbalexin A*)

Ito, C. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1488-1491 (*Glybomine A*)

2,6-Dihydroxy-3-methyl-9H-carbazole D-617

3-Methyl-9H-carbazole-2,6-diol

$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235

2-Me ether: 6-Hydroxy-2-methoxy-3-methyl-9H-carbazole. **Glycozolidol**
[58005-42-8]

$\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262

Alkaloid from the roots of *Glycosmis pentaphylla* (Rutaceae). Active against *Staphylococcus aureus*, *Bacillus firmis*, *Agrobacterium tumefaciens* and others. Cryst. (CHCl_3). Sol. MeOH, C_6H_6 ; poorly sol. H_2O . Mp 240°. λ_{max} 232 (ϵ 31500); 260 (ϵ 12580); 308 (ϵ 19500) (EtOH) (Berdy).

2-Me ether, O-Ac:

$\text{C}_{16}\text{H}_{15}\text{NO}_3$ 269.299

Cryst. (C_6H_6 /petrol). Mp 194°.

6-Me ether: 2-Hydroxy-6-methoxy-3-methyl-9H-carbazole. **Carbalexin C**
 $\text{C}_{14}\text{H}_{13}\text{NO}_2$ 227.262

Stress metab. of *Glycosmis parviflora* and *Glycosmis pentaphylla*. Phytoalexin. Oil. λ_{max} 231; 263 (sh); 310 (MeOH).

Di-Me ether: 2,6-Dimethoxy-3-methyl-9H-carbazole, 9CI. **Glycozolidine**
[29093-41-2]

$\text{C}_{15}\text{H}_{15}\text{NO}_2$ 241.289

Alkaloid from the root bark of *Glycosmis pentaphylla* and the roots of *Glycosmis mauritiana* (Rutaceae). Cryst. (C_6H_6 /MeOH). Mp 166-167° (161-162°).

Islam, A. *et al.*, *Chem. Comm.*, 1972, 537 (*synth*, *pmr*)

Anwer, F. *et al.*, *Indian J. Chem.*, 1972, **10**, 959 (*struct*)

Rastogi, K. *et al.*, *Phytochemistry*, 1980, **19**, 945 (*isol*)

Bhattacharyya, P. *et al.*, *Phytochemistry*, 1985, **24**, 882 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Iwao, M. *et al.*, *Heterocycles*, 1993, **36**, 1483 (*synth*, *Glycozolidine*)

Chakravarty, A.K. *et al.*, *Phytochemistry*, 1999, **50**, 1263-1266 (*isol*, *pmr*, *cmr*, *Glycozolidine*)

Pacher, T. *et al.*, *Phytochemistry*, 2001, **58**, 129-135 (*Carbalexin C*)

2,8-Dihydroxy-3-methyl-9H-carbazole D-618

3-Methyl-9H-carbazole-2,8-diol

$\text{C}_{13}\text{H}_{11}\text{NO}_2$ 213.235


8-Me ether: 2-Hydroxy-8-methoxy-3-methyl-9H-carbazole. **Carbalexin B**

[74059-97-5]
C₁₄H₁₃NO₂ 227.262
Stress metab. of *Glycosmis parviflora*
and *Glycosmis pentaphylla*. Needles
(CHCl₃). Mp 199-201° dec. λ_{max} 235 ;
239 (sh) ; 251 (sh) ; 299 ; 317 (sh) ; 330
(MeOH).

Mester, I. et al., *Annalen*, 1980, 241-245 (synth,
pmr, cmr)
Pacher, T. et al., *Phytochemistry*, 2001, 58, 129-
135 (isol)

2,7-Dihydroxy-6-methyl-9H-carbazole-1-carboxaldehyde D-619

1-Formyl-2,7-dihydroxy-6-methylcarbazole



C₁₄H₁₁NO₃ 241.246

2-Me ether: 7-Hydroxy-2-methoxy-6-methyl-9H-carbazole-1-carboxaldehyde. 1-Formyl-7-hydroxy-2-methoxy-6-methylcarbazole. **Murrayaline B** [139726-42-4]

C₁₅H₁₃NO₃ 255.273
Alkaloid from the stem bark of *Murraya euchrestifolia*. Yellow prisms (Me₂CO). Mp 240-242°. λ_{max} 223 ; 259 (sh) ; 303 ; 380 (MeOH).

Di-Me ether: 2,7-Dimethoxy-6-methyl-9H-carbazole-1-carboxaldehyde. 1-Formyl-2,7-dimethoxy-6-methylcarbazole. **Murrayaline A**. *Murrayaline* [104778-01-0]

C₁₆H₁₅NO₃ 269.299


Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Pale-yellow prisms (Me₂CO). Mp 248-250°.

Furukawa, H. et al., *Chem. Pharm. Bull.*, 1986, 34, 2672-2675 (*Murrayaline A*)

Ito, C. et al., *Chem. Pharm. Bull.*, 1991, 39, 2525-2528 (*Murrayaline B*)

6,7-Dihydroxy-1-methyl-β-carboline D-620

1-Methyl-9H-pyrido[3,4-b]indole-6,7-diol



C₁₂H₁₀N₂O₂ 214.223

Di-Me ether: 6,7-Dimethoxy-1-methyl-9H-pyrido[3,4-b]indole, 9CI. 6,7-Dimethoxy-1-methyl-β-carboline. **Roeharmin** [131984-70-8]

C₁₄H₁₄N₂O₂ 242.277
Alkaloid from *Roemeria hybrida* (Papaveraceae).

IR, 2,3,4-Tetrahydro, di-Me ether: 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methyl-β-carboline. **1,2,3,4-Tetrahydroroeharmin**

[131984-71-9]
C₁₄H₁₈N₂O₂ 246.308
Alkaloid from *Roemeria hybrida*
(Papaveraceae). [α]_D -4 (c, 0.1 in
CHCl₃) (natural). [α]_D -18 (c, 1.04 in
MeOH) (>98% op). Nat. prod. probably
partially racemic.

Gözler, B. et al., *J. Nat. Prod.*, 1990, 53, 740
(isol, uv, ms, pmr, struct)

Reddy, M.S. et al., *Tet. Lett.*, 1994, 35, 5413
(synth)

7,8-Dihydroxy-1-methyl-β-carboline D-621

1-Methyl-9H-pyrido[3,4-b]indole-7,8-diol

C₁₂H₁₀N₂O₂ 214.223

O⁷-Me, 8-O-β-D-glucopyranoside: **Ruine**. 8-Glucosyloxyharmine

[32472-23-4]

C₁₉H₂₂N₂O₇ 390.392

Alkaloid from seedlings of *Peganum harmala* (Zygophyllaceae). Mp 227-229°.

3,4-Dihydro, O⁷-Me, 8-O-β-D-Glucopyranoside: **Dihydroruine**

[52898-19-8]

C₁₉H₂₄N₂O₇ 392.408

Alkaloid from callus cultures of *Peganum harmala* (Zygophyllaceae). Easily autoxidised to Ruine.


Nettleship, L. et al., *Phytochemistry*, 1971, 10, 231 (*Ruine*)

McKenzie, E. et al., *Phytochemistry*, 1975, 14, 273 (*Dihydroruine*)

Gohar, A. et al., *Mansoura J. Pharm. Sci.*, 1994, 10, 109; *CA*, 121, 251119t (8-Glucosyloxyharmine)

2,4-Dihydroxy-3-methylenebutanoic acid D-622

2-Hydroxy-3-hydroxymethyl-3-butenic acid



C₅H₈O₄ 132.116

(S)-form

Nitrile: 2-Hydroxy-3-hydroxymethyl-3-butenenitrile, 9CI

[29768-67-0]

C₅H₇NO₂ 113.116

Constit. of *Cardiospermum halicacabum* and other Sapindaceae. Component of *Jia Ku Gua*.

Nitrile, 2-O-β-D-glucopyranoside:

Cardiospermin

[54525-10-9]

C₁₁H₁₇NO₇ 275.258

Isol. from *Cardiospermum hirsutum* and *Heterodendron oleaefolium*.

Nitrile, 2-O-β-D-glucopyranoside, 5-O-sulfate: **Cardiospermin 5-sulfate**

[78856-87-8]

C₁₁H₁₇NO₁₀S 355.322

Isol. from *Cardiospermum grandiflor-*

um.

Nitrile, 5-(4-hydroxybenzoyl), 2-O-β-D-glucopyranoside: **Cardiospermin 5-(4-hydroxybenzoate)**

[60444-99-7]

C₁₈H₂₁NO₉ 395.365

Constit. of *Sorbaria arborea* and *Sorbaria sorbifolia* var. *stellipila*. Powder.

Nitrile, 5-(4-hydroxycinnamoyl), 2-O-β-D-glucopyranoside: **Cardiospermin 5-p-coumarate**

[79197-19-6]

C₂₀H₂₃NO₉ 421.403

Constit. of *Sorbaria arborea*.

[84976-22-7, 73621-97-3]

Seigler, D.S. et al., *Phytochemistry*, 1974, 13, 2330 (*isol*)

Huebel, W. et al., *Phytochemistry*, 1975, 14, 2723 (*isol*)

Huebel, W. et al., *Tet. Lett.*, 1979, 45, 4395 (*sulfate*)

Huebel, W. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1981, 314, 609 (*cmr*)

Nahrstedt, A. et al., *Phytochemistry*, 1981, 20, 1309-1310 (*Cardiospermin esters*)

Spitzer, V. et al., *Phytochemistry*, 1996, 42, 1357-1360 (*isol, nitrile*)


Kim, D.K. et al., *Chem. Pharm. Bull.*, 2000, 48, 1766-1767 (5-4-hydroxybenzoate)

3,7-Dihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI D-623

2,7-Dihydroxy-1-methyl-4-azafluorenone.

2,7-Dihydroxyonychine

[161196-95-8]



C₁₃H₉NO₃ 227.219

Alkaloid from roots of *Piptostigma fugax* (Annonaceae). Orange cryst. Mp 300°.

3-Me ether: 7-Hydroxy-3-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 7-Hydroxy-2-methoxy-1-methyl-4-aza-9-fluorenone. 7-Hydroxy-2-methoxyonychine

[161196-96-9]

C₁₄H₁₁NO₃ 241.246

Alkaloid from stem bark of *Piptostigma fugax* (Annonaceae). Yellow needles. Mp 300°.

Di-Me ether: [161196-94-7]

Orange needles. Mp 204-206°.

Achenbach, H. et al., *Phytochemistry*, 1995, 38, 1037 (*isol, uv, ir, pmr, cmr, ms, struct*)

6,7-Dihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one D-624

7,8-Dihydroxy-1-methyl-4-azafluoren-9-one

C₁₃H₉NO₃ 227.219

6-Me ether: 7-Hydroxy-6-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 7-Hydroxy-8-methoxy-1-methyl-4-azafluoren-9-one. **Macondine**

[111316-31-5]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the stem bark of

Oxandra cf. *major* (Annonaceae).
Amorph.

6-*Me ether*, *Ac*: Mp 173°.

Arango, G.J. *et al.*, *Phytochemistry*, 1987, **26**, 2093 (*isol, uv, ir, pmr, cmr, ms, struct*)

7,8-Dihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one

6,7-Dihydroxy-1-methyl-4-aza-9-fluorenone, 6,7-Dihydroxyonychine

C₁₃H₉NO₃ 227.219

7-*Me ether*: 8-Hydroxy-7-methoxy-1-methyl-5H-indeno[1,2-b]pyridin-5-one, 6-Hydroxy-7-methoxyonychine. **Oncodine**

[122890-42-0]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the stem bark of *Oncodostigma monosperma* (Annonaceae). Orange-yellow needles (Me₂CO). Mp 187-188°. λ_{max} 213 (log ε 4); 226 (sh) (log ε 3.93); 255 (log ε 4.03); 316 (log ε 3.96); 369 (log ε 3.7) (EtOH/HCl). λ_{max} 226 (sh) (log ε 3.71); 253 (log ε 4.06); 268 (sh) (log ε 3.87); 324 (log ε 4); 370 (sh) (log ε 3.46); 480 (log ε 3.22) (EtOH/NaOH). λ_{max} 210 (log ε 3.95); 240 (sh) (log ε 3.96); 249 (log ε 4); 264 (sh) (log ε 4); 282 (log ε 4.03); 300 (log ε 3.96); 330 (log ε 3.42); 350 (sh) (log ε 3.28) (EtOH).

8-*Me ether*: 7-Hydroxy-8-methoxy-1-methyl-5H-indeno[1,2-b]pyridin-5-one, 7-Hydroxy-6-methoxyonychine. **Isooncocode**

[122890-43-1]

C₁₄H₁₁NO₃ 241.246

Alkaloid from *Polyalthia longifolia* (Annonaceae). Yellow powder. λ_{max} 218 (log ε 3.45); 264 (log ε 3.78); 293 (log ε 3.39); 314 (log ε 3.3); 328 (log ε 3.28); 340 (sh) (log ε 3.12) (MeOH). λ_{max} 225 (log ε 3.46); 284 (log ε 3.58); 318 (log ε 3.56); 343 (log ε 3.28); 354 (log ε 3.23) (MeOH/NaOH).

Di-Me ether: 7,8-Dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 6,7-Dimethoxyonychine. **Polyfothine**

[122908-91-2]

C₁₅H₁₃NO₃ 255.273

Alkaloid from the bark of *Polyalthia longifolia* (Annonaceae). Yellow solid (CHCl₃/petrol) or pale yellow powder. Mp 188-190°. λ_{max} 220 (log ε 3.43); 268 (log ε 3.79); 293 (log ε 3.38); 314 (log ε 3.32); 328 (log ε 3.3) (MeOH).

Bou-Abdallah, E. *et al.*, *J. Nat. Prod.*, 1989, **52**, 273-278 (*Oncodine, isol, Isooncocode, synth*)

Cassels, B.K. *et al.*, *J. Nat. Prod.*, 1989, **52**, 420 (*cmr*)

Chakrabarty, M. *et al.*, *Indian J. Chem., Sect. B*, 1990, **29**, 394 (*isol, struct*)

Wu, Y.-C. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1327-1331 (*Isooncocode, Polyfothine, isol*)

8,9-Dihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI

5,6-Dihydroxy-1-methyl-4-azafluorenone

C₁₃H₉NO₃ 227.219

8-*Me ether*: 9-Hydroxy-8-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 5-Hydroxy-6-methoxy-1-methyl-4-aza-9-fluorenone. **Isoursuline**. 5-Hydroxy-6-methoxyonychine

[112368-57-7]

C₁₄H₁₁NO₃ 241.246

Alkaloid from *Polyalthia stenopetala*, *Unonopsis spectabilis* and *Oxandra xylopioides* (Annonaceae). Shows mild cytotoxic effect. Orange amorph. solid or yellowish prisms (petrol). Mp 140-142°. This struct. has been erroneously attributed to Oxylopine below.

8-*Me ether*, 9-*Ac*: [101899-50-7]

Pale yellow microcryst. Mp 195°.

9-*Me ether*: 8-Hydroxy-9-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. **Ursuline**. Oxylopine

[111316-34-8]

C₁₄H₁₁NO₃ 241.246

Alkaloid from *Unonopsis spectabilis*, stem bark of *Oncodostigma monosperma* and *Oxandra xylopioides*. Yellow needles (CH₂Cl₂) or amorph. solid. Mp 158-160°.

9-*Me ether*, 8-*Ac*:

Microcryst. Mp 175-177°.

Di-Me ether: [112368-58-8]

Yellow microcryst. Mp 144°.

5-*Alcohol*, 8-*Me ether*: 8-Methoxy-4-methyl-5H-indeno[1,2-b]pyridine-5,9-diol. 5-Hydroxy-6-methoxy-1-methyl-4-azafluoren-9-ol. **Polylongine**

[122279-87-2]

C₁₄H₁₃NO₃ 243.262

Alkaloid from the leaves of *Polyalthia longifolia* (Annonaceae). Pale yellowish needles (CHCl₃). Mp 148-151°. [α]_D -1.6 (c, 0.1 in CHCl₃).

[101899-49-4]

Zhang, J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 800-806 (*isol, Isoursuline*)

Tadic, D. *et al.*, *Heterocycles*, 1988, **27**, 407-421 (*8-Me ether, 9-Me ether, synth, pmr, uv, ms*)

Laprévôt, O. *et al.*, *J. Nat. Prod.*, 1988, **51**, 555-561 (*Unonopsis spectabilis constits*)

Wu, Y.-C. *et al.*, *Heterocycles*, 1989, **29**, 463 (*Polylongine*)


Koyama, J. *et al.*, *Heterocycles*, 1989, **29**, 1649-1654 (*Ursuline, Isoursuline, synth*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1033 (*rev*)

Chen, C.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1475-1478 (*Isoursuline, activity*)

6,7-Dihydroxy-1-methylisoquinoline

1-Methyl-6,7-isoquinolinediol



C₁₀H₉NO₂ 175.187

▶NW8220000

6-*Me ether*: 6-Methoxy-1-methyl-7-isoquinolinol, 9CI. 7-Hydroxy-6-methoxy-1-methylisoquinoline

[4594-05-2]

C₁₁H₁₁NO₂ 189.213

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Needles (CHCl₃/MeOH). Mp 121-123°.

Di-Me ether: 6,7-Dimethoxy-1-methylisoquinoline, 9CI. **Isosalsolidine**. *Nigellimine*

[4594-02-9]

C₁₂H₁₃NO₂ 203.24

Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. Trace constit. in seeds of *Nigella sativa* (black cummin). Cryst. (Et₂O/EtOAc). Mp 118-119°.

Di-Me ether, N-oxide: **Nigellimine N-oxide**

[96562-85-5]

C₁₂H₁₃NO₃ 219.24

Minor alkaloid from the seeds of *Nigella sativa* (black cummin). Amorph.

3,4-Dihydro, *di-Me ether*: 3,4-Dihydro-6,7-dimethoxy-1-methylisoquinoline.

Dehydrosalsolidine

C₁₂H₁₅NO₂ 205.256

Alkaloid from *Carnegiea gigantea*. Mp 201-202° (195-197° as hydrochloride).

1,2,3,4-Tetrahydro: see 1,2,3,4-

Tetrahydro-6,7-dihydroxy-1-methylisoquinoline, T-162

Pummangura, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 277 (*Dehydrosalsolidine*)


Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109 (*Isosalsolidine, occur*)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1985, **23**, 953 (*Nigellimine oxide*)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1992, **55**, 676 (*Isosalsolidine*)

Chen, J.J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (*6-Me ether*)

3,5-Dihydroxy-6-methyl-7-(2-methyl-4-thiazolyl)-6-heptenoic acid



C₁₂H₁₇NO₄S 271.337

Prod. by *Sorangium cellulosum*. Precursor of Epothilones. Amorph. solid. [α]_D²² +9 (c, 0.3 in MeOH). λ_{max} 210 (ε 15900); 248 (ε 12200) (MeOH).

Homologue (R = CH₃): 3,5-Dihydroxy-2,6-dimethyl-7-(2-methyl-4-thiazolyl)-6-heptenoic acid

C₁₃H₁₉NO₄S 285.363

Prod. by *Sorangium cellulosum*.


Precursor of Epothilones. Amorph. powder. [α]_D²² +2 (c, 0.6 in MeOH). λ_{max} 212 (ε 16000); 248 (ε 12100) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

4-(5,7-Dihydroxy-2-methyl-4-oxo-4H-1-benzopyran-6-yl)-2-piperidinone, 9CI

D-629

4-(5,7-Dihydroxy-2-methyl-6-chroman-yl)-2-piperidone
[69735-30-4]



$C_{15}H_{15}NO_5$ 289.287

Alkaloid from root bark of *Schumannio-phyton problematicum* (Rubiaceae). Cryst. (EtOH). Mp 300-303°.

*O*⁷-Me: [69735-32-6]
Mp 262-263°.

N-Me: 4-(5,7-Dihydroxy-2-methyl-4-oxo-4H-1-benzopyran-6-yl)-1-methyl-2-piperidinone

[69735-31-5]

$C_{16}H_{17}NO_5$ 303.314


Alkaloid from *Schumannio-phyton problematicum* (Rubiaceae). Cryst. (EtOH/ C_6H_6). Mp 309-313°.

Schlittler, E. et al., *Tet. Lett.*, 1978, 2911 (*isol, uv, ir, pmr, ms, struct*)

4,5-Dihydroxy-1-methyl-3-oxo-2-(trichloromethyl)-3H-indolum(1+)

D-630

[942477-70-5]



$C_{10}H_7Cl_3NO_3^{\oplus}$ 295.528


Alkaloid from the roots of *Zanthoxylum nitidum*. Yellow needles (MeOH) (as chloride). Possible artifact. λ_{max} 225 (log ϵ 3.73); 285 (log ϵ 3.55); 300 (log ϵ 3.68) (MeOH) (chloride).

Hu, J. et al., *Helv. Chim. Acta*, 2007, **90**, 720-722 (*isol, pmr, cmr*)

2,5-Dihydroxy-3-methylpyrazine

D-631

5-Hydroxy-3-methyl-2(1H)-pyrazinone, 9CI. 3-Methyl-2,5-pyrazinediol. PB4 [102694-22-4]



$C_5H_6N_2O_2$ 126.115


Isol. from rice inoculated with *Pseudallescheria boydii*. Inhibitor of monoamine oxidase.

Maebayashi, Y. et al., *CA*, 1986, **105**, 3147a (*isol*)

3,4-Dihydroxy-2-methylpyridine

D-632

3-Hydroxy-2-methyl-4(1H)-pyridinone, 9CI. 2-Methyl-3,4-pyridinediol [17184-19-9]



$C_6H_7NO_2$ 125.127

Prod. by *Streptomyces* sp. MJ 210. Cryst. (H_2O). Mp 285-288°.

Hydrochloride: [17184-18-8]

Prisms (EtOAc/MeOH). Mp 185°.

***O*³- β -D-Glucopyranoside: Innovanamine**

[15266-34-9]

$C_{12}H_{17}NO_7$ 287.269

Isol. from leaves of *Evodiopanax innovans* (preferred genus name *Gamblea*). Prisms + 2 H_2O (H_2O). Mp 116-118°. $[\alpha]_D^{25}$ -61.6 (c, 10 in H_2O).

***O*³-Ac: 3-Acetoxy-2-methyl-4(1H)-pyridinone**

[17184-20-2]

$C_8H_9NO_3$ 167.164

Cryst. (EtOAc). Mp 205-207°.

3-Me ether: 3-Methoxy-2-methyl-4(1H)-pyridinone, 9CI

[76015-11-7]

$C_7H_9NO_2$ 139.154

Cryst. ($Me_2CO/MeOH$). Mp 155-156°.

3-Benzyl ether: [61160-18-7]

$C_{13}H_{13}NO_2$ 215.251

Prisms (EtOH). Mp 162-163°.

NH-form

N-Me: *Deferiprone*, BAN, INN. 3-Hydroxy-1,2-dimethyl-4(1H)-pyridinone, 9CI. Kelfer. CP 20 [30652-11-0]

$C_7H_9NO_2$ 139.154

Chelating agent. Used in the treatment of thalassaemia. Metal poisoning antidote. Launched 1995 (India). Cryst. ($MeOH/Et_2O$). Mp 260° dec. Log P -0.67 (calc).

► UU7785940

N-Me, hydrochloride: [150630-21-0]
Powder (EtOH/ Et_2O). Mp 189-190°.

N-Benzyl: [30652-22-3]

$C_{13}H_{13}NO_2$ 215.251

Solid ($Et_2O/EtOAc$). Mp 203-205°.

N-Ph: 3-Hydroxy-2-methyl-1-phenyl-4(1H)-pyridinone, 9CI

[49744-73-2]

Used as soln. in 1,2-dichloroethane or dioxan for extraction-sepn. of In, as 0.01M soln. in $CHCl_3$ for extraction-photometric detn. of V(*V*) (λ_{max} 625 nm, ϵ 5600 in $CHCl_3$); as 7mM soln. in $CHCl_3$ for extraction-sepn. of Ga, U and extraction-photometric detn. of Fe(III) (λ_{max} 470 nm, ϵ 6300 in $CHCl_3$) and W (λ_{max} 406 nm, ϵ 10600 in $CHCl_3$). Cryst. pK_{a1} 3.02; pK_{a2} 9.56.

Di-OH-form

4-Me ether, N-Me: [181647-53-0]

$C_8H_{12}NO_2^{\oplus}$ 154.188

Solid (EtOAc) (as trifluoromethanesulfonate salt). Mp 105-106° (trifluoromethanesulfonate). CAS no. refers to the salt.

Dibenzyl ether: [127234-69-9]

$C_{20}H_{19}NO_2$ 305.376

Cryst. ($CHCl_3$ /petrol). Mp 85-87°.

Dibenzyl ether, N-oxide: [127234-70-2]

$C_{20}H_{19}NO_3$ 321.375

Mp 127-129°.

[123742-61-0]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 721B (*nmr*)

Yasue, M. et al., *Chem. Pharm. Bull.*, 1966, **14**, 1443-1444 (*Innovanamine, isol*)

Tamhina, B. et al., *J. Inorg. Nucl. Chem.*, 1974, **36**, 1855-1857; 1976, **38**, 1505-1507 (*synth, N-Ph, detn, U*)

Harris, R.L.N. et al., *Aust. J. Chem.*, 1976, **29**, 1329-1334 (*synth*)

Tamhina, B. et al., *Microchem. J.*, 1977, **22**,

144-148 (*detn, W*)

Hwang, D.R. et al., *J. Pharm. Sci.*, 1980, **69**,

1074-1076 (*synth*)

Kontoghiorghes, G.J. et al., *Arzneim.-Forsch.*, 1987, **37**, 1099-1102 (*Deferiprone, pharmacol*)

Kontoghiorghes, G.J. et al., *Inorg. Chim. Acta*,

1987, **136**, L11-L12 (*synth*)

Nelson, W.O. et al., *Can. J. Chem.*, 1988, **66**,

123-131 (*Deferiprone, synth*)

Ishii, H. et al., *Bull. Chem. Soc. Jpn.*, 1989, **62**,

1817-1821 (*detn, In*)

Hider, R.C. et al., *J. Chem. Res., Synop.*, 1990,

316-317; *J. Chem. Res., Miniprint*, 1990,

2520-2555 (*Deferiprone, cryst struct*)

Bergeron, R.J. et al., *Blood*, 1992, **79**, 1882

(*Deferiprone, pharmacol*)

Dobbin, P.S. et al., *J. Med. Chem.*, 1993, **36**,

2448-2458 (*N-Me*)

Japan. Pat., 1993, 93 146 298; *CA*, **119**, 115500f

(*isol*)

Rumbo, A. et al., *J.O.C.*, 1996, **61**, 6114-6120

(*N-Me, Me ether*)

Fox, R.C. et al., *Synth. Commun.*, 1999, **29**,

989-1001 (*N-Me*)

Piyamongkol, S. et al., *Tetrahedron*, 2001, **57**,

3479-3486 (*3-benzyl ether, 3,4-dibenzyl ether*)

Ma, Y. et al., *J. Med. Chem.*, 2004, 6349-6362

(*Me ether*)


Barral, K. et al., *J. Med. Chem.*, 2006, **49**, 43-

50 (*benzyl ether*)

2,4-Dihydroxy-5-methylquinazoline

D-633

5-Methyl-2,4(1H,3H)-quinazolinone, 9CI. 5-Methyl-2,4-quinazolinediol [52570-39-5]



$C_9H_8N_2O_2$ 176.174


Prod. by *Streptomyces* sp. GW2/577. Mp 272-278°.

DeGraw, J.I. et al., *J. Med. Chem.*, 1974, **17**, 762-764 (*synth*)


Maskey, R.P. et al., *J. Nat. Prod.*, 2004, **67**, 1131-1134 (*isol, pmr, cmr, ms*)

1,9-Dihydroxy-4-(methylthio)canthin-6-one D-634


1,9-Dihydroxy-4-methylthio-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI

C₁₅H₁₀N₂O₃S 298.322**S-Oxide(S-): Curtisin**C₁₅H₁₀N₂O₄S 314.321Alkaloid from the mushroom *Boletus curtisii*. Pale yellow solid. [α]_D²¹ -149 (c, 0.08 in MeOH). Mp >340°. λ_{\max} 239 (sh) (log ϵ 4.14); 257 (sh) (log ϵ 4.16); 303 (log ϵ 4.16); 416 (log ϵ 4.09) (MeOH).Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (*isol, uv, cd, ir, pmr, cmr, ms*)**5,8-Dihydroxy-2-naphthalenecarboxylic acid** D-635

5,8-Dihydroxy-2-naphthoic acid

C₁₁H₈O₄ 204.182**Di-Me ether, amide:** 5,8-Dimethoxy-2-naphthalenecarboxamide



[955367-98-3]

C₁₃H₁₃NO₃ 231.251Prod. by *Streptovercillium morookaense*.Feng, N. *et al.*, *Nat. Prod. Commun.*, 2007, 2, 151-153 (*di-Me ether amide*)**3,4-Dihydroxy-10-nitro-1-phenanthrenecarboxylic acid** D-636C₁₅H₉NO₆ 299.239**4-Me ether, Me ester:** Methyl 3-hydroxy-4-methoxy-10-nitro-1-phenanthrenecarboxylate

[128397-29-5]

C₁₇H₁₃NO₆ 327.293Alkaloid from the stems of *Aristolochia liukinensis* (Aristolochiaceae). Pale yellow needles (CHCl₃/MeOH). Mp 164-165°. λ_{\max} 214 (ϵ 31600); 257 (ϵ 50100); 295 (ϵ 15800); 356 (ϵ 6310); 374 (ϵ 6310) (MeOH) (Derep).**Di-Me ether, Me ester:** Methyl 3,4-dimethoxy-10-nitro-1-phenanthrenecarboxylate. **Ariskanin A.** *Aristolochic acid***BII methyl ester**

[128397-31-9]

C₁₈H₁₅NO₆ 341.32Constit. of *Aristolochia manshuriensis* and *Aristolochia kankauensis*. Cryst. (MeOH) or yellowish needles (CHCl₃). Mp 123-124°.**Methylene ether:** see 3,4-Methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-447Mizuno, M. *et al.*, *J. Nat. Prod.*, 1990, 53, 179 (*isol, uv, ir, pmr, cmr, ms, struct*)Houghton, P.J. *et al.*, *Phytochemistry*, 1991, 30, 253 (*isol*)Wu, T.S. *et al.*, *Phytochemistry*, 1994, 36, 1063Lou, F.C. *et al.*, *Yaoxue Xuebao*, 1995, 30, 588; *CA*, 123, 25141n (*Aristolochic acid BII*)**4,5-Dihydroxy-2-octenoic acid** D-637C₈H₁₄O₄ 174.196**(4R,5R)-(E)-form****(2-Methylpropyl)amide:** 4,5-Dihydroxy-N-(2-methylpropyl)-2-octenamamide. 4,5-Dihydroxy-N-isobutyl-2-octenamamideC₁₂H₂₃NO₃ 229.319Alkaloid from the roots of *Piper nigrum* (pepper). Oil.Wei, K. *et al.*, *J. Nat. Prod.*, 2004, 67, 1005-1009 (*isol, pmr, cmr*)**3,4-Dihydroxy-6-oxo-1-cyclohexene-1-acetic acid** D-638C₈H₁₀O₅ 186.164**(3R*,4R*)-form****(+)-trans-form****Nitrile:** 3,4-Dihydroxy-6-oxo-1-cyclohexene-1-acetonitrileC₈H₉NO₃ 167.164Constit. of the whole plant of *Aquilegia ecalcarata*. Needles. Mp 91-92°.[α]_D²⁷ +27 (c, 0.2 in MeOH). λ_{\max} 214 (MeOH).Chen, S.-B. *et al.*, *Planta Med.*, 2002, 68, 554-556 (*isol, pmr, cmr, ms*)**3,5-Dihydroxy-2-[[1-oxo-3-(3,4,5-trihydroxyphenyl)-2-propenyl]-amino]benzoic acid, 9CI** D-639C₁₆H₁₃NO₈ 347.281**(E)-form****Miriamide**

[153698-89-6]

Isol. from eggs of the large white cabbage butterfly *Pieris brassicae*. Host-marking pheromone.**5-O- β -Glucopyranoside:** **Miriamide 5-glucoside**


[153566-00-8]

C₂₂H₂₃NO₁₃ 509.423From eggs of *Pieris brassicae*. Host-marking pheromone.**5-Deoxy:** **5-Deoxymiriamide**

[153566-01-9]

C₁₆H₁₃NO₇ 331.281From eggs of *Pieris brassicae*. Host-marking pheromone.Blaakmeer, A. *et al.*, *J. Nat. Prod.*, 1994, 57, 90 (*isol, uv, pmr, cmr, struct, synth*)**2,3-Dihydroxy-1-phenazinecarboxylic acid** D-640

[71670-84-3]

C₁₃H₈N₂O₄ 256.217Isol. from *Pseudomonas aureofaciens*.Römer, A. *et al.*, *Tet. Lett.*, 1979, 509 (*uv, ir, pmr, struct*)**2,6-Dihydroxy-1-phenazinecarboxylic acid** D-641

[79417-74-6]

C₁₃H₈N₂O₄ 256.217Minor pigment from *Pseudomonas aureofaciens*.Römer, A. *et al.*, *Z. Naturforsch., B*, 1981, 36, 1037 (*isol, uv, ir, pmr, ms, struct*)**2,9-Dihydroxy-1-phenazinecarboxylic acid** D-642

[23448-75-1]

C₁₃H₈N₂O₄ 256.217Phenazine antibiotic. Prod. by *Pseudomonas phenazinium* and *Pseudomonas fluorescens*. Mp 270°.**Di-Me ether, Me ester:** [23448-74-0]C₁₆H₁₄N₂O₄ 298.298

Mp 148-153°.

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1969, 6, 297 (*isol*)Byng, G.S. *et al.*, *J. Gen. Microbiol.*, 1976, 97, 57 (*isol*)**3,6-Dihydroxy-1-phenazinecarboxylic acid** D-643**Phenacein**

[94818-84-5]

C₁₃H₈N₂O₄ 256.217

Phenazine antibiotic. Prod. by *Streptomyces tanashiensis-zaomyeticus*. Angiotensin converting enzyme inhibitor. Shows weak antimicrobial activity. Hypotensive agent. Fine orange needles + $\frac{1}{2}$ H₂O (MeOH aq.). Sol. DMF, bases; fairly sol. MeOH, AcOH, MeCN, CHCl₃; poorly sol. H₂O, hexane, acids. Mp 310°. λ_{\max} 215; 268; 395; 466 (MeOH/HCl) (Derep). λ_{\max} 212; 293; 420; 530 (MeOH/NaOH) (Derep). λ_{\max} 218 (ϵ 5120); 268 (ϵ 21500); 385 (ϵ 2050); 460 (sh) (MeOH) (Derep).

Di-Me ether, Me ester:

C₁₆H₁₄N₂O₄ 298.298
Needles. Mp 181-182°.

Bush, K. *et al.*, *J. Antibiot.*, 1984, **37**, 1308
(*isol, props*)

Liu, W.-C. *et al.*, *J. Antibiot.*, 1984, **37**, 1313
(*struct, synth*)

4,9-Dihydroxy-1,6-phenazinedicarboxylic acid D-644

C₁₄H₈N₂O₆ 300.227


Di-Me ester: Dimethyl 4,9-dihydroxy-1,6-phenazinedicarboxylate

[66997-72-6]
C₁₆H₁₂N₂O₆ 328.281

Isol. from a strain of *Pseudomonas cepacia* ATCC 17460. Sol. MeOH, CHCl₃, C₆H₆, bases; poorly sol. H₂O, hexane. λ_{\max} 276; 368; 448 (MeOH/HCl) (Berdy). λ_{\max} 350; 365; 450 (MeOH).

Korth, H. *et al.*, *J. Gen. Microbiol.*, 1978, **104**, 299-303 (*isol, uv, ir, pmr, struct*)

2-(3,4-Dihydroxyphenethyl)-quinoline D-645



C₁₇H₁₅NO₂ 265.311

Di-Me ether: 2-[2-(3,4-Dimethoxyphenyl)ethyl]quinoline, 9CI. 2-(3,4-Dimethoxyphenethyl)quinoline

[95279-34-8]
C₁₉H₁₉NO₂ 293.365

Alkaloid from the stem bark of *Galipea longiflora* (Rutaceae). Amorph.

Methylene ether: 2-[2-(1,3-Benzodioxol-5-yl)ethyl]quinoline, 9CI. 2-(3,4-Methylenedioxyphenethyl)quinoline

[124902-95-0]
C₁₈H₁₅NO₂ 277.322

Alkaloid from *Galipea longiflora* and *Galipea bracteata* (Rutaceae). Shows molluscicidal and plant growth inhibitory activity. Cryst. (MeOH or MeOH aq.). Mp 68-70° (65°).


Fournet, A. *et al.*, *Can. J. Chem.*, 1989, **67**, 2116 (*isol, uv, pmr, ms, struct*)

Vieira, P.C. *et al.*, *Phytochemistry*, 1990, **29**, 813 (*isol, ir, pmr, cmr, ms, struct*)

Caron, S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 138
(*synth, pmr, cmr, ms*)

(2,4-Dihydroxyphenyl)acetic acid, 8CI D-646

2,4-Dihydroxybenzeneacetic acid
[614-82-4]



C₈H₈O₄ 168.149

Constit. of the venom of the spider *Nephila maculata*. Cryst. (C₆H₆/Et₂O). Mp 116-118°.

Nitrile: (2,4-Dihydroxyphenyl)acetone-trile. 4-(Cyanomethyl)-1,3-benzenediol
[57576-34-8]

C₈H₇NO₂ 149.149

Constit. of *Erica scoparia*. Cryst. Mp 130-131°.

Nitrile, 2-O-β-D-glucopyranoside: Ehretioside B
[156368-84-2]

C₁₄H₁₇NO₇ 311.291


Constit. of *Ehretia philippinensis* and *Semiaquilegia adoxoides*. Needles (MeOH). Mp 201-202°. $[\alpha]_D^{28}$ -78.5 (c, 0.6 in MeOH).

Ballester, A. *et al.*, *Phytochemistry*, 1975, **14**, 1667-1668 (*isol, synth, nitrile*)
Simpol, L.R. *et al.*, *Phytochemistry*, 1994, **36**, 91 (*Ehretioside B*)

Su, Y.-F. *et al.*, *J. Asian Nat. Prod. Res.*, 2005, **7**, 171-174 (*Ehretioside B*)

3-(3,4-Dihydroxyphenyl)-2-hydroxy-2-methylpropanoic acid D-647

α,3,4-Trihydroxy-α-methylbenzenepranoic acid, 9CI



C₁₀H₁₂O₅ 212.202

(ξ)-form

3'-Me ether, amide: 2-Hydroxy-3-(4-hydroxy-3-methoxyphenyl)-2-methylpropanamide

[214967-88-1]


C₁₁H₁₅NO₄ 225.244

Constit. of *Cordia macleodii*.

El-Sayed, N.H. *et al.*, *Rev. Latinoam. Quim.*, 1998, **26**, 30-35

3-(3,4-Dihydroxyphenyl)-2-hydroxypropanoic acid D-648

α,3,4-Trihydroxybenzenepranoic acid, 9CI. 3-(3,4-Dihydroxyphenyl)lactic acid. Danshensu. Danshensuan A
[23028-17-3]



(R)-form

C₉H₁₀O₅ 198.175

(R)-form [76822-21-4]

Constit. of *Coptidis Rhizoma* (root of *Coptis chinensis*) and *Salvia miltiorrhiza*. Coronary vasodilator. Prisms (EtOAc/C₆H₆). Mp 84-87°. $[\alpha]_D^{18}$ +10.8 (c, 3.7 in MeOH).

Amide: 2-Hydroxy-3-(3,4-dihydroxyphenyl)propanamide. 3-(3,4-Dihydroxyphenyl)lactamide

C₉H₁₁NO₄ 197.19

Alkaloid from the rhizomes of *Salvia miltiorrhiza*. Amorph. yellow powder. $[\alpha]_D^{20}$ +15.8 (c, 0.1 in MeOH). λ_{\max} 208 (log ϵ 4.04); 280 (log ϵ 3.54) (MeOH).


[69339-96-4, 67810-33-7, 67920-52-9, 42085-50-7, 81075-52-7]

Choi, J.S. *et al.*, *Fitoterapia*, 2001, **72**, 30-34
(*amide*)

2-(3,4-Dihydroxyphenyl)-2-oxoacetic acid D-649

3,4-Dihydroxy-α-oxobenzeneacetic acid, 9CI. (3,4-Dihydroxyphenyl)glyoxylic acid, 8CI. 3,4-Dihydroxybenzoylformic acid

[10118-81-7]



C₈H₆O₅ 182.132

Isol. from cultures of *Polyporus tumulosus*. Needles (CHCl₃/petrol), yellow needles + 1H₂O (AcOH/petrol). Mp 92° (colourless form) Mp 159° (yellow form).

3'-Me ether, amide: 2-(4-Hydroxy-3-methoxyphenyl)-2-oxoacetamide

C₉H₉NO₄ 195.174

Isol. from the heartwood of *Populus euphratica*. Amorph. yellow powder. Mp 105-107°. λ_{\max} 203; 231; 289; 315 (MeOH).

Di-Me ether: 3,4-Dimethoxy-α-oxobenzeneacetic acid. Veratroylformic acid

C₁₀H₁₀O₅ 210.186

Needles (C₆H₆). Mp 138-139°.

Di-Me ether, nitrile:

C₁₀H₉NO₃ 191.186


Needles (petrol). Mp 116-117°.

Barger, G. *et al.*, *J.C.S.*, 1909, **95**, 552 (*synth*)
Kindler, K. *et al.*, *Ber.*, 1943, **76**, 308 (*synth*)
Glennie, D.W. *et al.*, *J.A.C.S.*, 1955, **77**, 2409
(*synth, spectra*)

Crowden, R.K. *et al.*, *Aust. J. Chem.*, 1961, **14**, 475 (*isol*)

Luo, J.-R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2008, **44**, 6-9 (3'-Me ether amide)

5-(3,4-Dihydroxyphenyl)-2-phenyloxazole **D-650**



C₁₅H₁₁NO₃ 253.257

Di-Me ether: 5-(3,4-Dimethoxyphenyl)-2-phenyloxazole. *Balsoxine* [70216-32-9]

C₁₇H₁₅NO₃ 281.31

Alkaloid from *Amyris balsamifera* (Rutaceae). Sol. H₂O. Mp 99-100°.

▶ LD₅₀ (mus, ivn) 5000 - 15000 mg/kg.

Methylene ether: 5-(1,3-Benzodioxol-5-yl)-2-phenyloxazole. 5-(3,4-Methylenedioxyphenyl)-2-phenyloxazole. *Texamine*

[115070-71-8]

C₁₆H₁₁NO₃ 265.268

Isol. from the roots of *Amyris texana* (Rutaceae). Cryst. (EtOAc/hexane). Mp 139-141°.

Burke, B. *et al.*, *Heterocycles*, 1979, **12**, 349-351 (*Balsoxine*)

Huth, A. *et al.*, *Annalen*, 1984, 641-648 (*synth*)
Dominguez, X.A. *et al.*, *Heterocycles*, 1988,


27, 35-38 (*Texamine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Sengupta, S. *et al.*, *Indian J. Chem., Sect. B*, 2002, **41**, 854-855 (*Texamine*, *synth*, *pmr*, *cmr*)

Hodgetts, K.J. *et al.*, *Org. Lett.*, 2002, **4**, 2905-2907 (*synth*)

2,3-Dihydroxy-3-phenylpropanoic acid, 9CI **D-651**

α,β-Dihydroxybenzenepropanoic acid, 9CI. 3-Phenylglyceric acid, 8CI. *α,β-Dihydroxyhydrocinnamic acid* [5695-95-4]



(2R,3R)-form

C₉H₁₀O₄ 182.176

(2R,3R)-form

(-)-erythro-form

[70094-91-6]

Cryst. Mp 97-98°. [α]_D²⁰ -25.6 (H₂O).

2-Phenylethylamide: *Secodemethylclausenamide*

[140848-74-4]

[151670-53-0, 151670-52-9, 157366-45-5]

C₁₇H₁₉NO₃ 285.342

Alkaloid from *Clausena lansium* (wampee) (Rutaceae). Mp 151-152°.

[α]_D²⁸ +38 (c, 0.3 in MeOH). Abs.

config. revised in 1994. λ_{max} 255 (log ε 2.6) (MeOH).

2-Phenylethylamide, N-Me: *Secoclause-namide*

[140848-73-3]

[157332-69-9]

C₁₈H₂₁NO₃ 299.369

Alkaloid from *Clausena lansium*

(wampee) (Rutaceae). Mp 145-147°.

[α]_D²² -65.8 (c, 0.2 in CHCl₃). Abs.

config. revised in 1994. λ_{max} 256 (log ε 2.8) (MeOH).


[65870-48-6, 73610-80-7, 60209-23-6, 65870-46-4, 65870-49-7]

Yang, M.H. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 775-776 (*Secoclause-namides*)

Du, J.F. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 179-180; 269-270 (*Secoclause-namides*)

3-(3,4-Dihydroxyphenyl)propanoic acid **D-652**

3,4-Dihydroxybenzenepropanoic acid, 9CI. 3,4-Dihydroxyhydrocinnamic acid, 8CI. *Hydrocaffeic acid* [1078-61-1]



C₉H₁₀O₄ 182.176

Occurs in spores of *Lycopodium clavatum* and also in higher plants. Leaflets (H₂O). Mp 139°. pK_{a1} 4.56; pK_{a2} 9.36; pK_{a3} 11.6 (30°, 0.1M NaClO₄).

▶ MW5143500

Di-Me ether, amide: 3-(3,4-Dimethoxyphenyl)propanamide [14773-41-2]

C₁₁H₁₅NO₃ 209.244

Alkaloid from *Piper arboricola*. Needles (C₆H₆). Mp 120-121°.

Di-Me ether, 2-methylpropylamide: 3-(3,4-Dimethoxyphenyl)propanoic acid 2-methylpropylamide. 3,4-Dimethoxyhydrocinnamic isobutylamide [210689-26-2]

C₁₅H₂₃NO₃ 265.352

Alkaloid from *Melicope melanophloia*.

He, Q. *et al.*, *CA*, 1982, **97**, 11684e (*Di-Me ether amide*)


Latif, Z. *et al.*, *Biochem. Syst. Ecol.*, 1998, **26**, 467-468 (*Di-Me ether isobutylamide*)

3-(3,4-Dihydroxyphenyl)-2-propenoic acid, 9CI **D-653**

3,4-Dihydroxycinnamic acid, 8CI. *Caffeic acid*

[331-39-5]

[2316-26-9]



(E)-form

C₉H₈O₄ 180.16

Log P 0.82 (calc).

▶ GD8950000

(ξ)-form

Di-Me ether, 2-methylpropylamide: 3,4-Dimethoxycinnamic acid isobutylamide

N-(3,4-Dimethoxycinnamoyl)isobutylamide [102934-28-1]

C₁₅H₂₁NO₃ 263.336

Constit. of *Piper amalago*. Config. not determined.

[110993-57-2 (Na salt, ns), 342431-60-1 (K salt)]

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (*di-Me ether isobutylamide*)

3-(3,5-Dihydroxyphenyl)-2-propenoic acid, 9CI **D-654**

3,5-Dihydroxycinnamic acid, 8CI [28374-93-8]

C₉H₈O₄ 180.16

Isol. from peach buds. Plant growth inhibitor. Needles (H₂O). Mp 245-246°.

Me ester, O-β-D-glucopyranoside:

C₁₆H₂₀O₉ 356.329

Constit. of the flowers of *Moricandia arvensis*. Powder. Mp 195-197°. [α]_D²⁵ +100 (c, 0.2 in MeOH). λ_{max} 215 (log ε 3.9); 230 (log ε 3.1); 290 (log ε 4) (MeOH).

3-Me ether, *Me ester*, 5-O-β-D-glucopyranoside: *Linusitamarin*

C₁₇H₂₂O₉ 370.355

Constit. of *Linum usitatissimum* (flax). Amorph. powder.

Di-Me ether: 3,5-Dimethoxycinnamic acid [16909-11-8]

C₁₁H₁₂O₄ 208.213

Needles (H₂O). Mp 175-176°. pK_{a1} 4.36 (25°).

Di-Me ether, pyrrolidide: N-(3,5-Dimethoxycinnamoyl)pyrrolidine. 3,5-Dimethoxycinnamic acid pyrrolidide [102934-29-2]

C₁₅H₁₉NO₃ 261.32

Alkaloid from *Piper amalago*.

3-Me ether, 5-O-(4-O-methyl-β-D-glucopyranoside): [934008-84-1]

C₁₇H₂₂O₉ 370.355

Isol. from *Cordyceps cicadae*. Amorph. powder. λ_{max} 216 (log ε 3.86); 225 (log ε 3.8); 281 (log ε 3.85); 308 (log ε 3.76) (MeOH).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 184A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1057B (nmr)

Mauthner, F. *et al.*, *J. Prakt. Chem.*, 1925, **110**, 126 (*synth*)

Altree-Williams, S. *et al.*, *Aust. J. Plant Physiol.*, 1975, **2**, 105; *CA*, **83**, 40204 (*isol*)

Mukherjee, J. *et al.*, *Indian J. Chem.*, 1975, **13**, 859 (*deriv*)

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (*pyrrolidide*)


Luyengi, L. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2012 (*Linusitamarin*)

Braham, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 517-522 (*Me ester glucoside*)

Zhang, S.-W. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 404-410 (*3-Me ether 4-methylglucoside*)

N-(2,5-Dihydroxyphenyl)pyridinium(1+) **D-655**

[33354-74-4]



C₁₁H₁₀NO₂[⊕] 188.205


Isol. from the leaves of *Punica granatum* (pomegranate). Needles (as chloride). Mp 193° (chloride, monohydrate). CAS no. refers to chloride.

Nawwar, M.A.M. *et al.*, *Phytochemistry*, 1994, **37**, 1175-1177 (*isol*, *pmr*, *cmr*)

Schmidt, A. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 541-546 (*struct*)

2,3-Dihydroxy-4-phenylquinoline D-656

3-Hydroxy-4-phenyl-2(1H)-quinolinone. Viridicatin
[129-24-8]

C₁₅H₁₁NO₂ 237.257

Isol. from *Penicillium viridicatum* and *Penicillium cyclopium*. Active against *Mycobacterium tuberculosis* at dilution 1:15,000. Needles (MeOH). Sol. bases; fairly sol. acids, MeOH, Et₂O; poorly sol. H₂O. Mp 268°. λ_{max} 224 (ε 37400); 240 (ε 18000); 298 (ε 7100); 310 (ε 7000); 320 (ε 9600); 332 (ε 8200) (EtOH) (Berdy). λ_{max} 228 (ε 28400); 253 (ε 16400); 295 (ε 7400); 334 (ε 12100); 344 (ε 13700) (EtOH-NaOH) (Berdy). λ_{max} 229 (ε 24000); 254 (ε 19000); 342 (ε 12800) (NaOH) (Berdy).

Ac:

C₁₇H₁₃NO₃ 279.295

Prisms (EtOH aq.). Mp 200-201°.

Me ether: 3-Methoxy-4-phenyl-2(1H)-quinolinone. O-Methylviridicatin

C₁₆H₁₃NO₂ 251.284Metab. of *Penicillium puberulum*.

Prisms (MeOH). Mp 248-249°.

3'-Hydroxy: 3-Hydroxy-4-(3-hydroxyphenyl)-2(1H)-quinolinone, 9CI. Viridicatol

C₁₅H₁₁NO₃ 253.257

From *Penicillium viridicatum*. Cryst. (EtOAc). Mp 280°. λ_{max} 226; 284; 316; 329 (MeOH) (Berdy).

Cunningham, K.G. *et al.*, *Biochem. J.*, 1953, **53**, 328

Eistert, B. *et al.*, *Z. Naturforsch., B.*, 1962, **17**, 202 (*synth*)

Birkinshaw, J.H. *et al.*, *Biochem. J.*, 1963, **89**, 196 (*deriv*)

Luckner, M. *et al.*, *Tet. Lett.*, 1964, 1987 (*deriv*)

Rapoport, H. *et al.*, *J.A.C.S.*, 1969, **91**, 6083

McCamish, M. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 241 (*ms*)

Wilson, B.J. *et al.*, *Microb. Toxins*, (Liegler, A. *et al* Eds.), VI, Academic Press, 1971, 510 (*rev*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 834

2,4-Dihydroxy-3-phenylquinoline D-657

4-Hydroxy-3-phenyl-2(1H)-quinolinone, 9CI
[14933-29-0]

C₁₅H₁₁NO₂ 237.257

Pale yellow solid.

NH-form

N-Me: 4-Hydroxy-1-methyl-3-phenyl-2(1H)-quinolinone. Arboricine†
[519-66-4]

C₁₆H₁₃NO₂ 251.284

Alkaloid from the leaves of *Glycosmis arborea*. Cryst. Mp 226° (222°).

Chakravarti, D. *et al.*, *J.C.S.*, 1953, 3337-3340 (*N-Me, isol*)

Bowman, R.E. *et al.*, *J.C.S.*, 1959, 444-447 (*N-Me, synth*)

Stadlbauer, W. *et al.*, *Monatsh. Chem.*, 1980, **111**, 1005-1013 (*N-Me, synth, pmr*)

Coppola, G.M. *et al.*, *Synth. Commun.*, 1985, **15**, 135-139 (*N-Me, synth, cmr*)


Bezuglyi, P.A. *et al.*, *Khim. Geterotsikl. Soedin.*, 1992, 522-524; *CA*, **117**, 251593z (*N-Me, synth*)

Klásek, A. *et al.*, *J. Het. Chem.*, 2003, **40**, 747-752 (*N-Me, synth*)

Moon, B.S. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 4952-4959 (*synth, pmr, cmr*)

4,5-Dihydroxy-2-phenylquinoline D-658

5-Hydroxy-2-phenyl-4(1H)-quinolinone, 9CI. 2-Phenyl-4,5-quinolinediol

C₁₅H₁₁NO₂ 237.257

NH-form

N-Me: 5-Hydroxy-1-methyl-2-phenyl-4(1H)-quinolinone
[19843-07-3]

C₁₆H₁₃NO₂ 251.284

Minor alkaloid from leaves of *Lunasia quercifolia* (Rutaceae). Pale yellow needles (MeOH). Mp 174-176°.

5-Me ether: 4-Hydroxy-5-methoxy-2-phenylquinolinone

C₁₆H₁₃NO₂ 251.284

Yellow needles (EtOAc). Mp 154-155°.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1968, **21**, 1389 (*isol, pmr, ms, struct*)

Venturella, P. *et al.*, *Gazz. Chim. Ital.*, 1970, **100**, 678 (*synth*)

Sato, S. *et al.*, *J. Het. Chem.*, 1999, **36**, 1345-1347 (*5-Me ether*)

4,6-Dihydroxy-2-phenylquinoline D-659

6-Hydroxy-2-phenyl-4(1H)-quinolinone. 2-Phenyl-4,6-quinolinediol
[147197-84-0]

C₁₅H₁₁NO₂ 237.257

Red powder + 0.9 AcOH (AcOH). Mp 243-245°.

NH-form

O,N-Di-Me: 6-Methoxy-1-methyl-2-phenyl-4(1H)-quinolinone, 9CI. Eduline
[6878-08-6]

C₁₇H₁₅NO₂ 265.311

Alkaloid from seeds of *Casimiroa edulis* (Mexican apple) and leaves of *Skimmia japonica* (Rutaceae). Plates (Me₂CO/Et₂O). Mp 187-188°.

O,N-Di-Me, perchlorate:

Needles (Me₂CO/Et₂O). Mp 250-252°.

O,N-Di-Me, picrate:

Golden-yellow plates (MeOH). Mp 225-227° dec.

OH-form

Di-Me ether: 4,6-Dimethoxy-2-phenylquinolinone
[22680-65-5]

C₁₇H₁₅NO₂ 265.311

Mp 118°.

Dziewoński, K. *et al.*, *CA*, 1935, **29**, 1092 (*synth*)

Kinl, F.A. *et al.*, *J.C.S.*, 1956, 4163 (*isol, uv, Eduline*)

Beyerman, H.C. *et al.*, *CA*, 1961, **55**, 10488b (*struct, synth, Eduline*)

Dreyer, D.L. *et al.*, *J.O.C.*, 1968, **33**, 3577 (*isol, pmr, Eduline*)

Boyd, D.R. *et al.*, *J.C.S.(C)*, 1970, 556 (*isol, pmr, ms, Eduline*)

Sato, S. *et al.*, *J. Het. Chem.*, 1999, **36**, 1189-1193 (*synth, ir, pmr*)

4,7-Dihydroxy-2-phenylquinoline D-660

7-Hydroxy-2-phenyl-4(1H)-quinolinone. 2-Phenyl-4,7-quinolinediol

C₁₅H₁₁NO₂ 237.257

Mp 200-201°.

NH-form

7-Me ether, N-Me: 7-Methoxy-1-methyl-2-phenyl-4(1H)-quinolinone. Eduline
[483-51-2]

C₁₇H₁₅NO₂ 265.311

Alkaloid from the bark of *Lunasia quercifolia*, *Lunasia amara* and *Casimiroa edulis* (Mexican apple). Needles (EtOH). Mp 200-201°.

7-Me ether, N-Me, hydrochloride:

Needles (5% HCl). Mp 248.5° dec.

7-Me ether, N-Me, picrate:


Yellow needles (EtOH). Mp 220-220.5° (191-192°).

Johnstone, R. *et al.*, *Aust. J. Chem.*, 1958, **11**, 562 (*isol, uv, ir, struct, synth*)

Sondheimer, F. *et al.*, *J.O.C.*, 1958, **23**, 762 (*isol, ir, struct*)

4,5-Dihydroxy-2-piperidinecarboxylic acid D-661

4,5-Dihydroxypipercolic acid

(2*S*,4*R*,5*R*)-formC₆H₁₁NO₄ 161.157(2*S*,4*R*,5*R*)-form [59246-13-8]

Isol. from *Julbernardia paniculata*, *Julbernardia globifera* (Fabaceae) and *Brachystegia speciformis*.

4-Sulfate: 4,5-Dihydroxypipercolic acid 4-sulfate. Cribronic acid

[552866-14-5]

C₆H₁₁NO₇S 241.221

Isol. from the sponge *Cribrochalina olemda*. Glutamate receptor agonist. Fine needles (MeOH aq.). [α]_D¹⁸ -16.3 (c, 0.24 in H₂O).

(2S,4R,5S)-form [59284-78-5]

Constit. of *Calliandra haematocephala* and *Derris elliptica* (Fabaceae). Mp 239-243° dec. $[\alpha]_D^{25}$ -22.9 (c, 2 in H₂O).

(2S,4S,5S)-form [59246-14-9]

From *Calliandra haematocephala*, *Derris elliptica*, *Isobertinia tomentosa* and *Isobertinia angolensis* (Fabaceae). Cubes (as hydrochloride). Mp 195-196° (hydrochloride). $[\alpha]_D^{20}$ +24.4 (c, 0.64 in 2M HCl) (hydrochloride).

(2ξ,4ξ,5ξ)-form

N-Me: 4,5-Dihydroxy-1-methyl-2-piperidinecarboxylic acid. **Glabin†** [35024-30-7]
C₇H₁₃NO₄ 175.184
Isol. from seeds of *Pongamia glabra* (Fabaceae). Cryst. (EtOH/H₂O/Et₂O). Mp 290-292° dec. $[\alpha]_D^{20}$ -54.82 (5M HCl).

[126641-66-5, 114297-04-0, 63357-05-1, 35024-31-8, 38146-56-4, 126643-94-5]

Kumar, P.S.J. et al., *Tet. Lett.*, 1971, 4451-4454 (*Glabin*)

Evrard, G. et al., *Cryst. Struct. Commun.*, 1972, 1, 215-217; 1974, 3, 61-63 (*cryst struct*)

Marlier, M. et al., *Phytochemistry*, 1972, 11, 2597-2599; 1976, 15, 183-185 (*Calliandra haematocephala*, *Derris elliptica* constits)

Shewry, P.R. et al., *Phytochemistry*, 1976, 15, 1981-1983 (*2S,4R,5R-form*, *isol*, *bibl*)


Bashyal, B.P. et al., *Tetrahedron*, 1987, 43, 415-422; 423-430 (*synth*, *ir*, *pmr*, *ms*)

Thieme, M. et al., *Synthesis*, 2000, 2051-2059 (*synth*, *pmr*)

Sakai, R. et al., *J. Nat. Prod.*, 2003, 66, 784-787 (*Cribronic acid*)

5,7-Dihydroxy-8-(2-piperidinyl)flavone D-662

N,O-Bis(demethyl)buchenavianine [91147-19-2]



C₂₀H₁₉NO₄ 337.374

Alkaloid from the fruits of *Buchenavia macrophylla* (Combretaceae). Cryst. (CHCl₃/MeOH). Mp 214-215°. λ_{\max} 213 (ε 25100); 268 (ε 20300) (MeOH) (Berdy). λ_{\max} 212; 271; 325 (MeOH-NaOH) (Berdy). λ_{\max} 214; 267; 330 (MeOH-HCl) (Berdy).

N-Me: O-Demethylbuchenavianine

[91147-18-1]

C₂₁H₂₁NO₄ 351.401

Major alkaloid from the fruits of *Buchenavia macrophylla* and *Buchenavia capitata* (Combretaceae). Shows anti-HIV activity. Cryst. (MeOH). Mp 229-231°. $[\alpha]_D$ 0 (c, 1 in CHCl₃). $[\alpha]_D$ -31 (c, 0.54 in MeOH). Log P 2.88 (calc). λ_{\max} 210 (ε 31400); 273 (ε 21800); 326 (ε 7100) (MeOH) (Berdy). λ_{\max} 213; 275; 346 (MeOH-NaOH)

(Berdy). λ_{\max} 214; 269; 308 (MeOH-HCl) (Berdy).

5-Me ether: N-Demethylbuchenavianine

[91147-17-0]

C₂₁H₂₁NO₄ 351.401

Alkaloid from the leaves of *Buchenavia macrophylla* (Combretaceae). Cryst. (MeOH). Mp 278-280°. $[\alpha]_D$ +9 (c, 0.7 in MeOH).

O⁵,N-Di-Me: Buchenavianine


[91147-16-9]

C₂₂H₂₃NO₄ 365.428

Major alkaloid from the leaves of *Buchenavia macrophylla*, also from *Buchenavia capitata*. Cryst. (MeOH). Mp 183-185°. $[\alpha]_D$ -4 (c, 2.09 in EtOH). λ_{\max} 214 (ε 31400); 243 (ε 8500); 272 (ε 13500); 336 (ε 6290) (MeOH) (Berdy). λ_{\max} 215; 240; 277; 361 (MeOH/NaOH) (Berdy). λ_{\max} 214; 247; 269; 336 (MeOH/HCl) (Berdy).

Ahond, A. et al., *Bull. Soc. Chim. Fr.*, Part II, 1984, 41 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *derivs*)

Beutler, J.A. et al., *J. Nat. Prod.*, 1992, 55, 207 (*Buchenavianine*, *isol*, *activity*)

1,3-Dihydroxy-2-prenyl-acridone D-663

C₁₈H₁₇NO₃ 295.337

3-Me ether, N-Me: 1-Hydroxy-3-methoxy-2-(3-methyl-2-butenyl)-10-methylacridone. 1-Hydroxy-3-methoxy-10-methyl-2-prenylacridone

[75652-64-1]

C₂₀H₂₁NO₃ 323.391

Alkaloid from the roots of *Glycosmis mauritiana* (Rutaceae). Deep yellow needles. Mp 134-136°.

3-Me ether, N-Me, 1-Ac: Mp 115-116°.

Rastogi, K. et al., *Phytochemistry*, 1980, 19, 945 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

1,3-Dihydroxy-4-prenyl-acridone D-664**1,3-Dihydroxy-4-(3-methyl-2-butenyl)-9(10H)-acridinone**

C₁₈H₁₇NO₃ 295.337

N-Me: 1,3-Dihydroxy-10-methyl-4-(3-methyl-2-butenyl)-9(10H)-acridinone. 1,3-Dihydroxy-10-methyl-4-prenylacridone

C₁₉H₁₉NO₃ 309.364

Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Orange needles (Me₂CO). Mp 168-169°.

3-Me ether, N-Me: 3-O-Methylglycoctrine II

[82354-37-8]

C₂₀H₂₁NO₃ 323.391

Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Orange needles (Me₂CO). Mp 134-135°.

Hlubucek, J. et al., *Aust. J. Chem.*, 1970, 23, 1881 (*synth*, *ir*, *uv*, *pmr*)

Wu, T.-S. et al., *Heterocycles*, 1982, 19, 1047 (*uv*, *ir*, *pmr*, *ms*, *struct*)


Wu, T.-S. et al., *J.C.S. Perkin 1*, 1983, 1681 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *deriv*)

Anand, R.C. et al., *J.C.S. Perkin 1*, 1991, 2339 (*synth*)

Anand, R.C. et al., *Chem. Comm.*, 1996, 199 (*synth*)

2-(1,2-Dihydroxypropyl)-3,4-dihydroxy-5-(hydroxymethyl)-pyrrolidine D-665

2,5-Imino-1,3,4,6,7-octanepentol



C₈H₁₇NO₅ 207.226

(1'S,2R,2'ξ,3R,4R,5R)-form [710948-62-2]

Alkaloid from the bulbs of *Scilla peruviana*. Syrup. $[\alpha]_D$ +13.3 (c, 1.23 in H₂O).


1'-Deoxy: 3,4-Dihydroxy-2-(hydroxymethyl)-5-(2-hydroxypropyl)pyrrolidine

[710948-61-1]

C₈H₁₇NO₄ 191.227

Alkaloid from the bulbs of *Scilla peruviana*. Syrup. $[\alpha]_D$ +40.8 (c, 0.54 in H₂O). Relative config. only determined.

Asano, N. et al., *J. Nat. Prod.*, 2004, 67, 846-850 (*isol*, *pmr*, *cmr*)

5-(1,2-Dihydroxypropyl)-2-pyridinecarboxylic acid D-666

C₉H₁₁NO₄ 197.19

(1'R,2'S)-form

Antibiotic CJ 15335. CJ 15335

Amorph. powder. Config. shown as (1'S,2'S)-form in ref.

Me ester: Antibiotic CJ 14877. CJ 14877

C₁₀H₁₃NO₄ 211.217

Prod. by *Marasmiellus* sp. CL21624. Cytokine prodn. inhibitor. Amorph. powder. $[\alpha]_D^{24}$ +20 (c, 0.13 in MeOH). λ_{\max} 230 (ε 9500); 270 (ε 5800) (MeOH).

Me ester, 2'-Ac: Antibiotic CJ 14897. CJ 14897

C₁₂H₁₅NO₅ 253.254


Prod. by *Marasmiellus* sp. CL21624. Cytokine prodn. inhibitor. Amorph. powder. $[\alpha]_D^{24}$ +27.1 (c, 0.17 in MeOH). λ_{\max} 230 (ε 8200); 270 (ε 4400) (MeOH).

Ichikawa, K. et al., *J. Antibiot.*, 2001, 54, 703-709 (*isol*, *uv*, *cd*, *pmr*)

6-(6,8-Dihydroxy-9-puriny)-suberosanone

D-667

[864514-59-0]

 $C_{20}H_{26}N_4O_3$ 370.45


Alkaloid from *Subergorgia suberosa*. Powder. $[\alpha]_D^{20} +28$ (c, 0.2 in $CHCl_3$). C-2 stereochem. revised in 2008. λ_{max} 212; 264 (MeOH).

Qi, S.-H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1288-1289; 2008, **71**, 716-718 (*isol*, *pmr*, *cmr*)

2,4-Dihydroxy-3-pyridine-carboxylic acid

D-668

1,2-Dihydro-4-hydroxy-2-oxo-3-pyridine-carboxylic acid. *2,4-Dihydroxynicotinic acid*

 $C_6H_5NO_4$ 155.11Cryst. (H_2O). Mp 182° dec.

4-Me ether, nitrile: 1,2-Dihydro-4-methoxy-2-oxo-3-pyridinecarbonitrile. *3-Cyano-4-methoxy-2-pyridone*. *3-Cyano-2-hydroxy-4-methoxypyridine*. **N-De-methylricinine**

[21642-98-8]

 $C_7H_6N_2O_2$ 150.137

Alkaloid from *Ricinus communis*. Hepatoprotective agent. Pale yellow needles (MeOH). Mp 293-294°.

NH-form

N-Me, nitrile: 1,2-Dihydro-4-hydroxy-1-methyl-2-oxo-3-pyridinecarbonitrile, *9CI*. *3-Cyano-4-hydroxy-1-methyl-2-pyridone*. **Ricininic acid**. **O-Demethylricinine**

[520-78-5]

 $C_7H_6N_2O_2$ 150.137Alkaloid from *Ricinus communis*.Needles (H_2O). Mp 293-294°.

O⁴,N-Di-Me, nitrile: see Ricinine, R-96
Schroeter, G. *et al.*, *Ber.*, 1932, **65**, 432-445 (*synth*, *derivs*)

Skursky, L. *et al.*, *J. Biol. Chem.*, 1969, **244**, 3238-3242 (*Me ether nitrile*)


Lee, H.J. *et al.*, *Phytochemistry*, 1972, **11**, 965-973 (*Ricininic acid*)

Yano, S. *et al.*, *Heterocycles*, 1993, **36**, 145-148 (*Me ether nitrile*)

2,6-Dihydroxy-4-pyrimidine-carboxylic acid

D-669

1,2,3,6-Tetrahydro-2,6-dioxo-4-pyrimidine-carboxylic acid, *9CI*. **Orotic acid**, **BAN**, **INN**. *Uracil-6-carboxylic acid*. *Vitamin B₁₃*. *Whey factor*. *Many other names* [65-86-1]

 $C_5H_4N_2O_4$ 156.098

Intermediate tautomers also possible. Exists as the dioxo tautomer in the cryst. state. Occurs in milk and other biol. systems. Key compd. involved in biosynth. of nucleic acid pyrimidine bases. Forms metal complexes. Used in photometric detn. of Zn. Salts used as dietary and mineral supplements. Shows bacteriostatic and cytostatic props. Uricosuric agent. Cryst. + H_2O (H_2O). Mp 322-325° Mp 345-346°. pK_{a1} 1.8; pK_{a2} 9.55 (25°). Log P -0.75 (calc). Also used as Fe^{2+} salt (Ferrous orotate, JAN).

► LD_{50} (mus, orl) 2000 mg/kg. RM3180000

Me ester: [6153-44-2] $C_6H_6N_2O_4$ 170.124Cryst. Mp 240-242° Mp 249°. pK_{a1} 7.93 (25°).*Et ester*: [1747-53-1] $C_7H_8N_2O_4$ 184.151Cryst. (H_2O). Mp 207-208°.*Propyl ester*: [4450-26-4] $C_8H_{10}N_2O_4$ 198.178

Cryst. Mp 234-236°.

Isopropyl ester: [4450-03-7] $C_8H_{10}N_2O_4$ 198.178

Cryst. Mp 217-218°.

Butyl ester: [22754-37-6] $C_9H_{12}N_2O_4$ 212.205Cryst. (H_2O). Mp 185-186° (182°).*Choline ester: Choline orotate. Cholergol.**Rufai B*

[24381-49-5]

 $C_{10}H_{17}N_3O_5$ 259.261

Used in treatment of liver disorders.

Amide: Orotamide. 6-Carbamoyluracil

[769-97-1]

 $C_5H_5N_3O_3$ 155.113Cryst. + H_2O . Mp 373°.*Dimethylamide: Orotic acid dimethyl-**amide. Orotosans S*

[3687-44-3]

 $C_7H_9N_3O_3$ 183.166

Used to treat liver disorders. Cryst. (EtOH). Mp 249°.

Nitrile: 6-Cyanouracil

[79457-91-3]

 $C_5H_3N_3O_2$ 137.098

Solid (EtOAc). Mp 260° dec.

NH-form

N¹-Me: 1,2,3,6-Tetrahydro-1-methyl-2,6-dioxo-4-pyrimidinecarboxylic acid. 1-Methylorotic acid, 8CI

[705-36-2]

 $C_6H_6N_2O_4$ 170.124Cryst. (H_2O). Mp 316-323° (>300°). pK_{a1} 1; pK_{a2} 10.52.*N¹-Me, Me ester*: [24766-55-0] $C_7H_8N_2O_4$ 184.151

Cryst. (MeOH). Mp 140-141° Mp 203-210°.

N¹-Me, amide: [55643-21-5] $C_6H_7N_3O_3$ 169.14Cryst. (H_2O). Mp 300°.*N¹-Propyl*: [118738-50-4] $C_8H_{10}N_2O_4$ 198.178

Cryst. (EtOH). Mp 233°.

N¹-Benzyl: [5971-86-8] $C_{12}H_{10}N_2O_4$ 246.222Cryst. (H_2O or EtOH). Mp 226-227.5° dec. (218-219° dec.).*N¹-Ph*: [723-76-2] $C_{11}H_8N_2O_4$ 232.195

Solid. Mp 288-290°.

N³-Me: 1,2,3,6-Tetrahydro-3-methyl-2,6-dioxo-4-pyrimidinecarboxylic acid

[2013-90-3]

 $C_6H_6N_2O_4$ 170.124

Cryst. (MeOH). Mp 257° (sealed tube).

N³-Et: [31555-24-5] $C_7H_8N_2O_4$ 184.151

Cryst. (MeOH). Mp 250-251° (sealed tube).

N¹,N³-Di-Me: 1,2,3,6-Tetrahydro-1,3-dimethyl-2,6-dioxo-4-pyrimidinecarboxylic acid

[4116-38-5]

 $C_7H_8N_2O_4$ 184.151Cryst. or needles (Et₂O). Mp 153-154°.*N¹,N³-Di-Me, Me ester*: [4116-39-6] $C_8H_{10}N_2O_4$ 198.178Cryst. or needles (Et₂O). Mp 77-78°.*N¹,N³-Dibenzyl*: $C_{19}H_{16}N_2O_4$ 336.346

Cryst. (EtOH aq.). Mp 194-196°.

N¹,N³-Di-Me, amide: [2019-20-7] $C_7H_9N_3O_3$ 183.166

Needles (MeOH). Mp 238-239°.

N¹,N³-Di-Me, nitrile: 6-Cyano-1,3-dimethyluracil

[49846-86-8]

 $C_7H_7N_3O_2$ 165.151

Needles (EtOH). Mp 168-170°.

[154-85-8, 24598-73-0, 5266-20-6, 50887-69-9, 22454-86-0]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 834A (*ir*)

Sadtler Standard C-13 NMR Spectra, 16598 (*cmr*)

Bachstsz, M. *et al.*, *Ber.*, 1931, **64**, 2683 (*I-Me*)

Nyc, J.F. *et al.*, *J.A.C.S.*, 1947, **69**, 1382 (*synth*)

Deghenghi, R. *et al.*, *Can. J. Chem.*, 1960, **38**, 1255 (*synth*)

Kokko, J.P. *et al.*, *J.A.C.S.*, 1962, **84**, 1042 (*pmr*)


Japan. Pat., 1963, 63 23 180; *CA*, **60**, 2964f (*dimethylamide*)

Curran, W.V. *et al.*, *J.O.C.*, 1966, **31**, 201 (*alkyl derivs, synth*)

Rambacher, P. *et al.*, *Angew. Chem., Int. Ed.*, 1968, **7**, 383 (*synth*)

- Takusagawa, F. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 2011 (*cryst struct*)
- Senda, S. *et al.*, *J.O.C.*, 1975, **40**, 353; 1979, **44**, 970 (*derivs, synth, 1-Me*)
- Ivin, B.A. *et al.*, *Zh. Org. Khim.*, 1976, **12**, 1836; *J. Org. Chem. USSR (Engl. Transl.)*, 1976, **12**, 1802 (*synth*)
- Walther, H. *et al.*, *Pharmazie*, 1979, **34**, 309 (*pharmacol*)
- Cihak, A. *et al.*, *Orotic Acid*, MTP: Lancaster, U.K., 1980, (*book*)
- Falk, M. *et al.*, *Pharmazie*, 1985, **40**, 377 (*rev, props*)
- Ullmann's Encycl. Ind. Chem.*, 5th Ed., VCH, Weinheim, 1985, **A12**, 156 (*synth*)
- Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 261
- Mirzoyan, V.S. *et al.*, *Khim. Geterotsikl. Soedin.*, 1990, 520; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1990, 446 (*ms*)
- Párkányi, C. *et al.*, *Struct. Chem.*, 1992, **3**, 277 (*uv*)
- Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1396
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, OJV500

4,5-Dihydroxy-1H-pyrrole-3-carboxylic acid D-670



C₅H₅NO₄ 143.099

4-Me ether, Me ester: **Methyl 5-hydroxy-4-methoxy-1H-pyrrole-3-carboxylate**


[128922-71-4]

C₇H₉NO₄ 171.152

Alkaloid from the stems of *Gnetum montanum* (Gnetaceae). Mp 204-206° (as hydrochloride).

Zhou, J. *et al.*, *Zhivwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 878-882; *CA*, **113**, 94757d (*isol, deriv*)

4,8-Dihydroxy-2-quinoline-carbothioic acid D-671



C₁₀H₇NO₃S 221.236

4-Me ether: 8-Hydroxy-4-methoxy-2-quinolinecarbothioic acid

[76995-83-0]


C₁₁H₉NO₃S 235.263

Prod. by *Pseudomonas fluorescens*. λ_{max} 218 (log ε 4.27); 268 (log ε 4.4); 355 (log ε 3.25); 417 (log ε 3.13) (MeOH).

Neuenhaus, W. *et al.*, *Z. Naturforsch., B*, 1980, **35**, 1569-1571 (*isol, uv, synth*)

2,6-Dihydroxy-4-quinoline-carboxylic acid D-672

1,2-Dihydro-6-hydroxy-2-oxo-4-quinoline-carboxylic acid. 2,6-Dihydroxycinchoninic acid. β-Acid [4363-99-9]



C₁₀H₇NO₄ 205.17

Mp 326° (browns at 300°).

Me ester: [66416-75-9]

C₁₁H₉NO₄ 219.196

Alkaloid from the aleurone layer of rice *Oryza sativa* cv. *heugjinmi*. Antioxidant. Pale yellow needles. Mp >320°. λ_{max} 242 (log ε 4.8); 280 (log ε 4.4); 382 (log ε 4.2) (MeOH).

Sahashi, Y. *et al.*, *Biochem. Z.*, 1927, **189**, 208

Makino, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1944, **19**, 95

Chung, H.S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1579-1580 (*Me ester, isol*)

2,8-Dihydroxy-4-quinoline-carboxylic acid D-673

1,2-Dihydro-8-hydroxy-2-oxo-4-quinoline-carboxylic acid. 2,8-Dihydroxycinchoninic acid. 8-Hydroxy-2(1H)-quinolinone-4-carboxylic acid. **Zeanic acid** [30536-55-1]

C₁₀H₇NO₄ 205.17

Isol. from corn steep liquor. Growth regulator. Needles (CHCl₃/MeOH). Mp 340° dec.

O-β-D-Glucopyranoside: **Zeanoside B**

[113202-67-8]

C₁₆H₁₇NO₉ 367.312

Isol. from immature corn kernels (*Zea mays*) (Poaceae). Sl. yellow needles. Mp 297-302°. [α]_D²³ -69.4 (c, 0.1 in NH₄OH).

Me ether: 1,2-Dihydro-8-methoxy-2-oxo-4-quinolinecarboxylic acid. 2-Hydroxy-8-methoxy-4-quinolinecarboxylic acid [37749-17-0]

C₁₁H₉NO₄ 219.196

Alkaloid from *Cynoglossum gansuense*. Yellowish powder. Mp 310° dec.

Matsushima, M. *et al.*, *Agric. Biol. Chem.*, 1970, **34**, 1430

Fukumi, H. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 768

Tateishi, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 3445 (**Zeanoside B**)

Shibata, H. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 849; 2545 (**Zeanoside B**)

Jin, Y.-P. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 776-781 (*Me ether, isol*)

3,4-Dihydroxy-2-quinoline-carboxylic acid D-674

1,4-Dihydro-3-hydroxy-4-oxo-2-quinoline-carboxylic acid. 3,4-Dihydroxyquinaldinic acid. 3-Hydroxy-4(1H)-quinolinone-2-carboxylic acid [33925-79-0]

C₁₀H₇NO₄ 205.17

Constit. of the Mediterranean sponge

Aplysinaerophoba. Mp 261-262° (253-254°) dec. λ_{max} 247 (ε 29100); 256 (ε 28800); 300 (ε 1740); 321 (ε 2110); 365 (ε 8580); 382 (ε 7580) (MeOH).

Coppini, D. *et al.*, *Gazz. Chim. Ital.*, 1950, **80**, 36 (*synth*)

Fattorusso, E. *et al.*, *Gazz. Chim. Ital.*, 1971, **101**, 104 (*isol, uv, ir, struct*)

4,6-Dihydroxy-2-quinoline-carboxylic acid D-675

1,4-Dihydro-6-hydroxy-4-oxo-2-quinoline-carboxylic acid, 9CI. 4,6-Dihydroxyquinaldinic acid. 6-Hydroxykynurenic acid

C₁₀H₇NO₄ 205.17

Alkaloid from tobacco (*Nicotiana tabacum*), *Thapsia villosa* and *Ginkgo biloba* (ginkgo). Appears to be widespread in the Solanaceae, also found in Scrophulariaceae, Lamiaceae and other families. Mp 287° dec.

6-Me ether: 1,4-Dihydro-6-methoxy-4-oxo-2-quinolinecarboxylic acid. 4-Hydroxy-6-methoxy-2-quinolinecarboxylic acid. 6-Methoxykynurenic acid [52980-06-0]

C₁₁H₉NO₄ 219.196

Alkaloid from stems of *Ephedra pachyclada* ssp. *sinatica*. Cryst. (MeOH). Mp 298-300° dec. (294-295° dec.).

[52980-07-1, 3778-29-8]

Macnicol, P.K. *et al.*, *Biochem. J.*, 1968, **107**, 473 (*isol, struct, synth, ir, pmr, uv, occur*)

Hall, C.M. *et al.*, *J. Med. Chem.*, 1974, **17**, 685 (*synth, 6-Methoxykynurenic acid*)

Mendez, J. *et al.*, *Phytochemistry*, 1975, **14**, 1136 (*isol*)

Schennen, A. *et al.*, *Planta Med.*, 1986, 235 (*isol, cmr*)

Starratt, A.N. *et al.*, *Phytochemistry*, 1996, **42**, 1477 (*isol, 6-Methoxykynurenic acid*)

4,7-Dihydroxy-2-quinoline-carboxylic acid D-676

1,4-Dihydro-7-hydroxy-4-oxo-2-quinoline-carboxylic acid. 4,7-Dihydroxyquinaldinic acid

C₁₀H₇NO₄ 205.17

7-Me ether: 1,4-Dihydro-7-methoxy-4-oxo-2-quinolinecarboxylic acid, 9CI. 7-Methoxy-4-hydroxy-2-quinolinecarboxylic acid. **Ephedralone** [77474-33-0]

C₁₁H₉NO₄ 219.196

Alkaloid from *Ephedra alata* whole plant (Ephedraceae). Prisms (MeOH aq.). Mp 268°.

Nawwar, M.A.M. *et al.*, *Phytochemistry*, 1985, **24**, 878-879 (*isol, uv, ir, pmr, cmr, ms, struct*)

4,8-Dihydroxy-2-quinoline-carboxylic acid D-677

1,4-Dihydro-8-hydroxy-4-oxo-2-quinoline-carboxylic acid. 4,8-Dihydroxyquinaldinic acid. **Xanthurenic acid**. Xanthuric acid [59-00-7]

C₁₀H₇NO₄ 205.17

Metab. of tryptophan. Present in human urine as metab. of Kynurenic acid in cases of vitamin B₆ deficiency. Occurs as a moulting inhibitor in crustaceans. Constit. of *Trididemnum* sp. Yellow cryst.

(H₂O). Mp 286° (297°). Forms a monohydrate. λ_{\max} 243 (ϵ 30000); 342 (ϵ 6500) (pH 6.95, H₂O).

► UZ9275000

8-O- β -D-Glucopyranoside: **Cardinalic acid**[†]

[97451-32-6]
C₁₆H₁₇NO₉ 367.312

Tryptophan metab. of *Drosophila melanogaster*.

Me ester: [5934-38-3]

C₁₁H₉NO₄ 219.196

Yellow cryst. (MeOH). Mp 262°.

Dibenzoyl, Me ester:

C₂₅H₁₇NO₆ 427.412

Mp 171°.

4-Me ether: 8-Hydroxy-4-methoxy-2-quinolinecarboxylic acid, 9CI. **Quinolobactin**

[28027-14-7]

C₁₁H₉NO₄ 219.196

Alkaloid from *Pseudomonas fluorescens*. Siderophore. Mp 248°. λ_{\max} 211 (log ϵ 4.48); 249 (log ϵ 4.56); 315 (log ϵ 3.45); 348 (log ϵ 3.57) (MeOH).

8-Me ether: 4-Hydroxy-8-methoxy-2-quinolinecarboxylic acid

[2929-14-8]

C₁₁H₉NO₄ 219.196

Metab. of tryptophan. Isol. from human urine. Possible endogenous carcinogen in man. Mp 240-241°.

► Exp. carcinogen. UZ9550000

[121691-37-0]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 870D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 449B (nmr)

Mebane, A.D. et al., *J.A.C.S.*, 1951, 73, 3520 (synth, uv)

Furst, A. et al., *J.O.C.*, 1951, 16, 412-414 (synth, uv)

Price, J.M. et al., *J. Biol. Chem.*, 1956, 223, 699-704 (8-Me ether, isol, synth)

Neuhaus, W. et al., *Z. Naturforsch., B*, 1980, 35, 1569-1571 (Quinolobactin)

Suzuki, M. et al., *Chem. Pharm. Bull.*, 1984, 32, 2340-2343 (8-Me ether, biochem)

Ferre, J. et al., *J. Biol. Chem.*, 1985, 260, 7509-7514; 1990, 265, 7407-7412 (Cardinalic acid)

De Silva, E.D. et al., *Tet. Lett.*, 1992, 33, 2917-2920 (Trididemnum, isol)


Okabe, N. et al., *Acta Cryst. C*, 1996, 52, 663-665 (cryst struct)

Mossialos, D. et al., *Appl. Environ. Microbiol.*, 2000, 66, 487-492 (Quinolobactin)

Moon, B. et al., *Tetrahedron*, 2000, 56, 9057-9062 (occur)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DNC200; HLT500

5,13-Dihydroxy-4,6,10,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadiene-3,7-dione 4-(3,11-Dihydroxy-2,6,10,12-tetramethyl-9,13-dioxo-1,5-pentadecadienyl)-2-methylthiazole




C₂₃H₃₅NO₄S 421.6

Epithilone fragment. Isol. from recombinant *Myxococcus xanthus*. Yellow oil. $[\alpha]_D^{25}$ -6.8 (c, 0.026 in MeOH). λ_{\max} 212 (ϵ 26700); 246 (ϵ 20000) (MeOH).

Starks, C.M. et al., *J. Nat. Prod.*, 2003, 66, 1313-1317 (isol, pmr, cmr)

5,13-Dihydroxy-2,4,6,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadien-3-one D-679




C₂₃H₃₇NO₃S 407.616

Prod. by *Sorangium cellulosum*. Precursor of Epithilones. Amorph. solid. $[\alpha]_D^{25}$ -28 (c, 0.4 in MeOH). λ_{\max} 211 (ϵ 16100); 247 (ϵ 12100) (MeOH).

Hardt, I. et al., *J. Nat. Prod.*, 2001, 64, 847-856

Dihydroxytriangularine D-680




C₁₈H₂₇NO₇ 369.414

Alkaloid from *Alkanna tinctoria* (Boraginaceae). $[\alpha]_D^{20}$ +14.7 (CHCl₃).

Röder, E. et al., *Phytochemistry*, 1984, 23, 2125 (isol, ir, ms, pmr, struct)

N-(2,4-Dihydroxy-3,5,6-trimethylbenzoyl)glycine, 9CI Antibiotic USF 406C. USF 406C [190782-86-6] D-681



C₁₂H₁₅NO₅ 253.254

Prod. by *Mortierella* sp. USF-406. Antioxidant. Needles (hexane/EtOAc). Mp 192°. λ_{\max} 207 (log ϵ 4.59); 258 (log ϵ 3.76) (MeOH).

Me ester: **Antibiotic USF 406D. USF 406D**

[190782-87-7]


C₁₃H₁₇NO₅ 267.281

Prod. by *Mortierella* sp. USF-406. Antioxidant. Needles (hexane/EtOAc). Mp 162°.

Hirota, A. et al., *Biosci., Biotechnol., Biochem.*, 1997, 61, 647-650 (isol, struct)

2,4-Dihydroxy-3,3,4-trimethyl-5-oxo-2-pyrrolidinecarboxylic acid D-682

2,4-Dihydroxy-3,3,4-trimethyl-5-oxoproline, 9CI. 2,4-Dihydroxy-3,3,4-trimethylpyroglutamic acid [143785-44-8]




C₈H₁₃NO₅ 203.194

Residue present in the lipopolysaccharide of *Vibrio anguillarum* V-123. On acid hydrolysis of the polysaccharide, undergoes isom. to a lactone.

Eguchi, H. et al., *Carbohydr. Res.*, 1992, 231, 147 (struct, pmr, cmr)

4,5-Dihydroxytryptamine D-683
3-(2-Aminoethyl)-1H-indole-4,5-diol, 9CI [42241-03-2]



C₁₀H₁₂N₂O₂ 192.217

N^b-Tricosanoyl: **4,5-Dihydroxy-N^b-tricosanoyltryptamine**

C₃₃H₅₆N₂O₃ 528.817

Constit. of the seeds of *Annona atemoya* (custard apple).

N^b-Tetracosanoyl: **4,5-Dihydroxy-N^b-tetracosanoyltryptamine**. 4,5-Dihydroxy-N^b-lignoceroyltryptamine

C₃₄H₅₈N₂O₃ 542.844

Constit. of the seeds of *Annona atemoya* (custard apple).

N^b-Pentacosanoyl: **4,5-Dihydroxy-N^b-pentacosanoyltryptamine**

C₃₅H₆₀N₂O₃ 556.871

Constit. of the seeds of *Annona atemoya* (custard apple).

N^b-Heptacosanoyl: **N^b-Heptacosanoyl-4,5-dihydroxytryptamine**

C₃₇H₆₄N₂O₃ 584.924

Constit. of the seeds of *Annona atemoya* (custard apple).

Wu, Y.-C. et al., *J. Nat. Prod.*, 2005, 68, 406-408 (N-Acyltryptamines)

5,6-Dihydroxytryptamine D-684
3-(2-Aminoethyl)-1H-indole-5,6-diol, 9CI [5090-36-8]

C₁₀H₁₂N₂O₂ 192.217

Mp 92-95° (as formate salt). Hygroscopic and unstable to light and air.

5-Me ether: 6-Hydroxy-5-methoxytryptamine

Off-white cryst. (EtOH/Et₂O) (as formate salt). Mp 207-208°.

5-*Me ether*, N^b-*Ac*: **6-Hydroxymelatonin**
[2208-41-5]
C₁₃H₁₆N₂O₃ 248.281
Metab. of Melatonin in humans. Cryst.
(EtOH/hexane). Mp 174-175°.

5-*Me ether*, N^b-*Ac*, 6-*sulfate*: [2208-40-4]
C₁₃H₁₆N₂O₆S 328.345
Amorph. powder (EtOAc/hexane).

6-*Me ether*: 5-*Hydroxy-6-methoxytryptamine*
Mp 71-73° Mp 136-138° (dimorphic).

Dibenzyl ether:

Orange-red cryst. (EtOH aq.) (as picrate). Mp 183-185° dec.

Schlossberger, H.G. *et al.*, *Annalen*, 1963, **662**, 132-138 (*synth*)


Benigni, J.D. *et al.*, *J. Het. Chem.*, 1965, **2**, 387-392 (*synth*, 5-*Me ether*, 6-*Me ether*, 6-*Hydroxymelatonin*)

Leone, A.M. *et al.*, *J. Pineal Res.*, 1988, **5**, 367-371 (5-*Me ether*, N-*Ac sulfate*)

Karam, O. *et al.*, *Tet. Lett.*, 2003, **44**, 1511-1513 (6-*Hydroxymelatonin*, *synth*, *pmr*)

4,8-Dihydroxy-2,9-undecadienedioic acid D-685

3,7-Dihydroxy-1,8-nonadiene-1,9-dicarboxylic acid



C₁₁H₁₆O₆ 244.244

(2Z,4E,8E,9Z)-form

Diamide: 4,8-Dihydroxy-2,9-undecadienediamide

[382138-86-5]


C₁₁H₁₈N₂O₄ 242.274

Isol. from an unidentified marine actinomycete.

Smelcerovic, A.A. *et al.*, *Hem. Ind.*, 2001, **55**, 399-401 (*isol*)

4,8-Dihydroxy-1-vinyl-β-carboline D-686

1-Ethenyl-9H-pyrido[3,4-b]indole-4,8-diol, 9CI



C₁₃H₁₀N₂O₂ 226.234

4-*Me ether*: 8-*Hydroxy-4-methoxy-1-vinyl-beta-carboline*. **Picrasidine I**. 8-*Hydroxydehydrocrenatinine*
[100234-59-1]

C₁₄H₁₂N₂O₂ 240.261

Alkaloid from the bark of *Picrasma quassioides* and *Picrasma javanica* (Simaroubaceae). Plates (CHCl₃/MeOH). Mp 240-241° dec. The alkaloid from *Picrasma javanica* was originally assigned the isomeric 5-hydroxy struct. and named 5-Hydroxydehydrocrenatinine (Arbain *et al.*, 1987).

4-*Me ether*, *Ac*:

Needles (Me₂CO). Mp 180-183°.

Di-Me ether: 1-Ethenyl-4,8-dimethoxy-9H-pyrido[3,4-b]indole. 4,8-Dimethoxy-1-vinyl-β-carboline. **Dehydrocrenatinidine**. *Kumujian G*
[65236-62-6]

C₁₅H₁₄N₂O₂ 254.288

Alkaloid from root and bark of *Ailanthus malabarica* and heartwood of *Picrasma quassioides*. Brownish-yellow plates. Mp 157°.

1',2'-*Dihydro*, 4-*Me ether*: 1-Ethyl-8-hydroxy-4-methoxy-β-carboline. 1-Ethyl-4-methoxy-9H-pyrido[3,4-b]indol-8-ol, 9CI. **Picrasidine J**. 8-*Hydroxycrenatinine*
[100234-62-6]

C₁₄H₁₄N₂O₂ 242.277

Alkaloid from the bark of *Picrasma quassioides* and *Picrasma javanica* (Simaroubaceae). Pale-yellow prisms (CHCl₃/MeOH). Mp 212-213°. Erroneously assigned the isomeric 5-hydroxy struct. when first isol. from *P. javanica* and named 5-Hydroxycrenatinine.

1',2'-*Dihydro*, 4-*Me ether*, *Ac*:

Needles (Me₂CO). Mp 193-195°.

1',2'-*Dihydro*, *di-Me ether*: 1-Ethyl-4,8-dimethoxy-9H-pyrido[3,4-b]indole. 1-Ethyl-4,8-dimethoxy-β-carboline. **Crenatinidine**
[30467-79-9]

C₁₅H₁₆N₂O₂ 256.304

Alkaloid from the bark of *Picrasma crenata* and from the bark and roots of *Ailanthus malabarica*. Mp 157-158°.

Sanchez, E. *et al.*, *Phytochemistry*, 1971, **10**, 2155 (*Crenatinidine*)

Joshi, B.S. *et al.*, *Heterocycles*, 1977, **7**, 193 (*Dehydrocrenatinidine*)


Yang, J.S. *et al.*, *Yaoxue Xuebao*, 1979, **14**, 167; *CA*, **92**, 72679a (*Kumujian G*)

Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 471-473 (*cmr*)

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3356 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Arbain, D. *et al.*, *Aust. J. Chem.*, 1987, **40**, 1527; 1990, **43**, 433 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)

17,18-Dihydroxyhimbaban-16-carboxylic acid D-687



C₂₀H₂₄N₂O₄ 356.421

Me ester: 18-**Hydroxyyohimbine**

[69862-18-6]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Rauwolfia cumminsii*, *Rauwolfia macrophylla*, *Rauwolfia mombasiana*, *Rauwolfia volkensii* and *Rauwolfia vomitoria* (probably) (Apocynaceae). A hydroxyyohimbine of undetd. struct. was isol. from *R.*

suaveolens. The identification of all samples is not well established.

16-*Epimer*, *Me ester*: 18-**Hydroxy-α-yohimbine**
[74081-22-4]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Rauwolfia mombasiana* roots. Stereochem. tentative.

3,16,18-Triepimer, *Me ester*: 18-**Hydroxyepialloyohimbine**. 18-*Hydroxyyohimbine* (*incorr.*)
[81703-06-2]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Rauwolfia nitida* root bark (Apocynaceae). Parent alcohol of the common ester alkaloids Reserpine, R-52, Deserpidine, D-234, etc.

3,16,18-Triepimer, O¹⁷-*Me*: **Deserpidic acid**. **Deserpidinic acid**. **Recanescic acid**. **Canescic acid**. **Raunormic acid**
[439-67-8]

C₂₁H₂₆N₂O₄ 370.447

Mp 270-273° (245-249°).

3,16,18-Triepimer, O¹⁷-*Me*, *Me ester*: **Methyl deserpidate**

[54290-61-8]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from roots of *Rauwolfia obscura*, *Rauwolfia canescens* and *Rauwolfia volkensii* (Apocynaceae). Mp 271-276° (as nitrate). [α]_D²⁴ -78 (c, 1 in Py) (nitrate).

3,16,18-Triepimer, 17-O-(3,4,5-trimethoxybenzoyl), *Me ester*: **Isoraunesine**

[483-07-8]

C₃₁H₃₆N₂O₈ 564.634

Alkaloid from *Rauwolfia canescens*, also detected in *Rauwolfia ligustrina* (Apocynaceae). Cryst. (MeOH aq.). Mp 241-242°. [α]_D -70 (c, 1 in CHCl₃).

3,16,18-Triepimer, 18-O-(3,4,5-trimethoxybenzoyl), *Me ester*:

Raunesine[†]

[117-73-7]

C₃₁H₃₆N₂O₈ 564.634

Alkaloid from *Rauwolfia canescens* and *Rauwolfia tetraphylla*, also detected in *Rauwolfia ligustrina* (Apocynaceae). Cryst. (MeOH aq.). Mp 160-170° (monohydrate). [α]_D -74 (c, 1 in CHCl₃).

3,16,18-Triepimer, 18-O-(3,4,5-trimethoxybenzoyl), *Me ester*, nitrate:
Mp 223-225°. [α]_D -80 (c, 1 in 5M AcOH).

3,16,18-Triepimer, O¹⁷-*Me*, 18-O-(3,4,5-trimethoxybenzoyl), *Me ester*: see Deserpidine, D-234

Stereoisomer, O¹⁷-*Me*, 18-O-(2-acetamidobenzoyl), *Me ester*: **Desanthraserpidine**

[123690-06-2]

C₃₁H₃₅N₃O₆ 545.634

Alkaloid from *in vitro* hairy-root cultures of *Catharanthus trichophyllus*. Amorph. Tentative struct. No stereochem. determined.

Stereoisomer, O¹⁷-*Me*, 18-O-(2-acetamido-4,5-dimethoxybenzoyl), *Me ester*: **Dimethoxydesanthraserpidine**

[123690-07-3]


C₃₃H₃₉N₃O₈ 605.686

Alkaloid from *in vitro* hairy-root cultures of *Catharanthus trichophyllus*. Amorph. Tentative struct. No stereochem. determined.

- Hosansky, N. *et al.*, *J. Am. Pharm. Assoc.*, 1955, **44**, 639-640 (*Raunescine*, *Isoraunescine*, *isol*, *uv*)
 Klohs, M.W. *et al.*, *J.A.C.S.*, 1955, **77**, 4084-4087 (*Methyl deserpidate*, *struct*)
 Huebner, C.F. *et al.*, *J.A.C.S.*, 1957, **79**, 250-251 (*Raunescine*, *Isoraunescine*, *struct*)
 Van Tamelen, E.E. *et al.*, *J.A.C.S.*, 1957, **79**, 5256-5262 (*Raunescine*, *Isoraunescine*, *struct*)
 Majumdar, S.P. *et al.*, *Phytochemistry*, 1973, **12**, 1167-1169 (*Rauwolfia suaveolens* *constit*)
 Timmins, P. *et al.*, *Phytochemistry*, 1974, **13**, 1997 (*Methyl deserpidate*)
 Iwu, M.M. *et al.*, *Phytochemistry*, 1978, **17**, 1651-1654 (*Me ester*)
 Sabri, N.N. *et al.*, *Phytochemistry*, 1978, **17**, 2023-2026 (*Me ester*)
 Iwu, M.M. *et al.*, *Planta Med.*, 1980, **38**, 260-263 (*isol*, *18-Hydroxy-α-yohimbine*)
 Amer, M.A. *et al.*, *Phytochemistry*, 1981, **20**, 2569-2573 (*18-Hydroxyepialloyohimbine*, *Me ester*)
 Szantay, C. *et al.*, *Annalen*, 1983, 1278-1291 (*Me ester*, *Raunescine*, *Isoraunescine*, *synth*)
 Davioud, E. *et al.*, *Phytochemistry*, 1989, **28**, 1383-1387 (*Desantraserpidine*, *Dimethoxydesantraserpidine*)

3,17-Dihydroxyzenkerine D-688

16,17,19,20-Tetrahydrocuran-3,17-diol, 9CI



Absolute Configuration


C₁₉H₂₂N₂O₂ 310.395

Enolised aldehyde. Alkaloid from the leaves of *Strychnos pseudoquina*. Pale yellow gum.

Da Silva, M.A. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 881-885 (*isol*, *pmr*, *cmr*)

3,3-Di-1H-indol-3-yl-2-butanone D-689

[28926-40-1]

C₂₀H₁₈N₂O 302.375


Isol. from the marine bacterium *Vibrio parahaemolyticus* Bio249. Pale yellow solid. Mp 197° (*synthetic*). λ_{max} 274 (sh) (log ε 2.91); 281 (log ε 2.94); 289 (log ε 2.89) (MeOH).

Zhungietu, G.I. *et al.*, *CA*, 1970, **73**, 25230u (*synth*)

Veluri, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1520-1523 (*isol*, *pmr*, *cmr*, *ms*)


3,4-Di-1H-indol-3-yl-2(5H)-furanone, 9CI D-690

[244295-63-4]

C₂₀H₁₄N₂O₂ 314.343

Prod. by *Paracoccus* sp. UV absorbent.

Japan. Pat., 1999, 99 269 175; *CA*, **131**, 242087s


2,2-Di-1H-indol-3-yl-4-methylpentanoic acid D-691C₂₂H₂₂N₂O₂ 346.428

Isol. from *Escherichia coli* supplemented with indole.

Garbe, T.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 596-598

4-[(Di-1H-indol-3-yl)methyl]-phenol, 9CI D-692

7,7-Bis(3-indolyl)-p-cresol. 4-Hydroxyphenyl-di-3-indolylmethane [151358-47-3]


C₂₃H₁₈N₂O 338.408

Isol. from a strain of the bacterium *Vibrio* sp. obtained from a marine sponge *Hyatella* sp. Exhibits antimicrobial activity. Gum. λ_{max} 224 (ε 52500); 282 (ε 11500) (MeOH) (*Derep*).

Oclarit, J.M. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 309-312 (*isol*, *pmr*, *cmr*, *uv*, *ir*)

3,3-Di-1H-indol-3-yl-1,2-propanediol, 9CI D-693


3,3-(2,3-Dihydroxypropyl)diindole [4531-83-3]

C₁₉H₁₈N₂O₂ 306.363

Alkaloid from the clavicipitaceous fungus *Balansia epichloë* which parasitises on pasture grasses. Also prod. by *Streptomyces cerevisiae* mutants. Red cryst. Mp 90-91°. λ_{max} 221 (log ε 4.88); 275 (log ε 3.97); 282 (log ε 4.01); 291 (log ε 3.96) (MeOH).

Lingens, F. *et al.*, *Biochim. Biophys. Acta*, 1967, **148**, 70-83 (*isol*, *ir*, *synth*)

Porter, J.K. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 88-93 (*isol*, *uv*, *pmr*, *ms*, *struct*, *synth*)

2,2-Di-1H-indol-3-ylpropanoic acid D-694C₁₉H₁₆N₂O₂ 304.348

Isol. from *Escherichia coli* supplemented with indole. Orange-yellow powder. Mp 109-114°. Turns blue on standing. λ_{max} 222 (log ε 4.75); 281 (log ε 4.04); 290 (log ε 3.99); 388 (log ε 2.02); 479 (log ε 1.23) (MeOH).

Garbe, T.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 596-598

Diiodoacetic acid D-695

[598-89-0]

I₂CHCOOHC₂H₂I₂O₂ 311.846

Constit. of *Asparagopsis taxiformis*. Pale-yellow needles (CHCl₃). Mod. sol. H₂O. Mp 110-111°.

Amide: *Diiodoacetamide*

[5875-23-0]

C₂H₃I₂NO 310.861

Constit. of *Asparagopsis taxiformis*.

Cryst. (H₂O). Mp 201-202° dec. (softens at 198°).

[82665-91-6]

Clarke, L. *et al.*, *J.A.C.S.*, 1914, **36**, 1899-1908 (*synth*)


Cobb, R.L. *et al.*, *J.O.C.*, 1958, **23**, 1368-1405 (*synth*)

Woolard, F.X. *et al.*, *Tetrahedron*, 1976, **32**, 2843-2846 (*amide*, *isol*)

Woolard, F.X. *et al.*, *Phytochemistry*, 1979, **18**, 617-620 (*isol*)

3,6-Diiodo-9H-carbazole D-696

[57103-02-3]

C₁₂H₇I₂N 419.003

Alkaloid from the cyanobacterium

Kyrtuthrix maculans. Leaflets (EtOH).
Mp 202-204°.

N-Ac:

C₁₄H₉I₂NO 461.04
Needles (EtOH or C₆H₆). Mp 224-225°.

N-Me: [90338-06-0]

C₁₃H₉I₂N 433.03
Needles (hexane) or rods (Me₂CO).
Mp 187° (181-182°).

N-Benzoyl: [96853-15-5]

C₁₉H₁₃I₂N 509.127
Needles (hexane). Mp 169-172°.

N-Ph: [57103-21-6]

C₁₈H₁₁I₂N 495.101
Needles (hexane). Mp 179-180°.

Tucker, S.H. *et al.*, *J.C.S.*, 1926, 546-553; 1927, 1214-1221 (*synth*)

Ambrose, J.F. *et al.*, *J. Electrochem. Soc.*, 1975, **122**, 876-894 (*synth*, 9-Ph)

Beginn, C. *et al.*, *Macromol. Chem. Phys.*, 1994, **195**, 2353-2370 (*synth*, *pmr*, *cmr*)

Lee, S.-C. *et al.*, *Phytochemistry*, 1999, **52**, 537-540 (*isol*, *pmr*, *cmr*, *ms*)


Bonesi, S.M. *et al.*, *J. Het. Chem.*, 2001, **38**, 77-87 (*synth*, *ir*, *pmr*, *cmr*, *ms*)

Monge, M.E. *et al.*, *J. Het. Chem.*, 2002, **39**, 933-941 (9-Me, 9-Ph, 9-benzyl)

Filimonov, V.D. *et al.*, *Zh. Org. Khim.*, 2003, **39**, 924-929; *Russ. J. Org. Chem. (Engl. Transl.)*, 2003, **39**, 875-880 (*synth*, *ms*)

(3,5-Diiodo-4-methoxyphenyl)ethylamine D-697

3,5-Diiodo-4-methoxybenzenethanamine
[89631-86-7]



C₉H₁₁I₂NO 403.001

Metab. of an unidentified didemnid tunicate. Isol. from the ascidian *Didemnum rubeum*. Mildly cytotoxic, antifungal. Cryst. (petrol). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 55-57°.

Hydrochloride:

Cryst. (CHCl₃). Mp 213-215°.


Jatzkewitz, H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1951, **287**, 43-46 (*synth*)

Sesin, D.F. *et al.*, *Tet. Lett.*, 1984, **25**, 403-404 (*isol*, *pmr*, *cmr*, *synth*)

Ford, P.W. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1051-1053 (*isol*)

7,20-Diisocyanoisocycloamphilectane D-698

7,20-Diisocyanoadociane
[60197-58-2]



C₂₂H₃₂N₂ 324.508

Isol. from a sponge *Adocia* sp., from the Great Barrier reef and *Cymbastela hoo-*

peri. Potent *in vitro* antimalarial activity. Shows antimycobacterial and cytotoxic activities. Cryst. (hexane). Mp 109-110°. [α]_D²⁵ +47.4 (c, 0.7 in CH₂Cl₂).

7-Isothiocyante: 20-Isocyanato-7-isothiocyanoisocycloamphilectane
[175861-78-6]

C₂₂H₃₂N₂S 356.574

Constit. of *Cymbastela hooperi*. Shows antimycobacterial, antimalarial and cytotoxic activities. Oil. [α]_D²⁵ +23.3 (c, 0.15 in CHCl₃). Has -NCS replacing -NC at C-7. λ_{max} 247 (ε 1250) (MeOH) (Berdy).

7-Isocyanate: 7-Isocyanato-20-isocyanoisocycloamphilectane
[175861-79-7]

C₂₂H₃₂N₂O 340.508

Constit. of *Cymbastela hooperi*. Oil. [α]_D²⁵ +36.1 (c, 0.75 in CHCl₃). Has -NCO replacing -NC at C-7.

20-Isocyanate: 20-Isocyanato-7-isocyanoisocycloamphilectane. 20-Isocyanato-7-isocyanoadociane
[175861-80-0]

C₂₂H₃₂N₂O 340.508

Isol. from the tropical marine sponge *Cymbastela hooperi*. Shows potent *in vitro* antimalarial activity. Oil. [α]_D²⁵ +37 (c, 0.58 in CHCl₃). Has -NCO replacing -NC at C-20.

[175861-77-5]

Baker, J.T. *et al.*, *J.A.C.S.*, 1976, **98**, 4010 (*isol*, *cryst struct*, *pmr*)

Corey, E.J. *et al.*, *J.A.C.S.*, 1987, **109**, 287-289 (*synth*)

Fookes, C.J.R. *et al.*, *J.C.S. Perkin 1*, 1988, 1003 (*cryst struct*, *biosynth*)

Koenig, G.M. *et al.*, *Magn. Reson. Chem.*, 1995, **33**, 694 (*pmr*, *cmr*)

Koenig, G.M. *et al.*, *J.O.C.*, 1996, **61**, 3259-3267 (*isol*, *ir*, *pmr*, *cmr*, *uv*, *ms*, *bibl*, *activity*)


Koenig, G.M. *et al.*, *Planta Med.*, 2000, **66**, 337-342 (*activity*)

Simpson, J.S. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 939-948 (*biosynth*)

Fairweather, K.A. *et al.*, *Org. Lett.*, 2006, **8**, 3395-3398 (*synth*)

2,5-Diisopropyl-3-methoxy-pyrazine D-699

3-Methoxy-2,5-bis(1-methylethyl)pyrazine, 9CI
[87386-79-6]




C₁₁H₁₈N₂O 194.276

Prod. by *Chondromyces crocatus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*, *synth*, *pmr*, *cmr*)

2,5-Diisopropylpyrazine D-700

2,5-Bis(1-methylethyl)pyrazine, 9CI
[24294-83-5]



C₁₀H₁₆N₂ 164.25

Prod. by *Chondromyces crocatus* and *Paenibacillus polymyxa*. Also prod. by microorganisms living on rotting pineapple (*Ananas sativus*). Attractant of the pineapple beetle *Carpophilus humeralis*. Bp_{0.001} 24-25.5°.

Zbiral, E. *et al.*, *Annalen*, 1969, **727**, 231-233 (*synth*)

Ohta, A. *et al.*, *Heterocycles*, 1977, **6**, 1881-1887 (*pmr*)

Matsuo, M. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 172-179 (*cmr*)

Zilkowski, B.W. *et al.*, *J. Chem. Ecol.*, 1999, **25**, 229-252 (*isol*, *activity*)

Beck, H.C. *et al.*, *FEMS Microbiol. Lett.*, 2003, **220**, 67-73 (*isol*, *pmr*, *cmr*, *ms*)

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*, *synth*, *pmr*, *cmr*)

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 5013-5018 (*isol*, *synth*, *pmr*, *cmr*, *ms*)

2,6-Diisopropylpyrazine D-701

2,6-Bis(1-methylethyl)pyrazine
[222048-85-3]

C₁₀H₁₆N₂ 164.25

Prod. by *Chondromyces crocatus*.


α-Hydroxy: 2-(1-Hydroxy-1-methylethyl)-6-isopropylpyrazine. α,α-Dimethyl-6-(1-methylethyl)pyrazine-methanol, 9CI
[870544-24-4]

C₁₀H₁₆N₂O 180.249

Prod. by *Chondromyces crocatus*.

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*, *synth*, *pmr*, *cmr*)

Aspergillus niger Diketo-piperazine dimer D-702



C₃₃H₃₈N₆O₄ 582.701


Similar to Antibiotic WIN 64745, A-1310. Prod. by *Aspergillus niger*. Powder. [α]_D +132 (c, 0.004 in MeOH). λ_{max} 240; 300 (MeCN aq.).

Ovenden, S.P.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 2093-2095 (*isol*, *pmr*, *cmr*)

Dilemmaone A

[205990-55-2]

D-703



C₁₄H₁₅NO₂ 229.278

Alkaloid probably from the sponge *Ectyonancora flabellata*. Off-white solid.

Mp 146-148°. Genus name freq. spelt Ectyonanchora. λ_{\max} 250 (ϵ 8060); 310 (ϵ 4100) (CHCl₃).

O-De-Me: Dilemmaone B

[205990-56-3]
C₁₃H₁₃NO₂ 215.251

Alkaloid probably from *Ectyonanchora flabellata*. Buff solid. Mp 168-170°. λ_{\max} 250 (ϵ 8420); 310 (ϵ 4230) (CHCl₃).

5-Hydroxy: Dilemmaone C


[205990-57-4]
C₁₄H₁₅NO₃ 245.277

Alkaloid probably from *Ectyonanchora flabellata*. Pale yellow solid. Mp 187-190°. λ_{\max} 250 (ϵ 7950); 310 (ϵ 12330) (CHCl₃).

Beukes, D.R. et al., *J. Nat. Prod.*, 1998, **61**, 699-701 (isol, uv, ir, pmr, cmr, ms)

Dimethamine D-704

3-(3,4-Dihydro-12-methyl-5-cytisyl)-12-methylcytisine. 3',4'-Dihydro-3,5'-bi(N-methylcytisine)
[37551-60-3]



Absolute Configuration

C₂₄H₃₂N₄O₂ 408.542
Alkaloid from epigeal parts of *Thermopsis alterniflora* (Fabaceae). Cryst. (MeOH/Me₂CO). Sol. H₂O. Mp 216-217° dec. $[\alpha]_D^{25}$ +143 (c, 0.58 in MeOH).


Hydrochloride (1:2): Mp 217-218°.

Dipicrate: Mp 218-219° dec.

Iskandarov, S. et al., *Khim. Prir. Soedin.*, 1972, **8**, 218; *Chem. Nat. Compd. (Engl. Transl.)*, 216 (isol, ir, ms, pmr, struct)

3,4-Dimethoxybenzaldehyde D-705

N-(3,4-dimethoxybenzoyl)hydrazide
3,4-Dimethoxybenzoic acid [(3,4-dimethoxyphenyl)methylene]hydrazide, 9CI. 3,4-Dimethoxybenzoic acid 3,4-dimethoxybenzylidenehydrazide. Veratrylidenehydrazide
[102017-12-9]



C₁₈H₂₀N₂O₅ 344.366
Constit. of the leaves of *Wedelia biflora*. Antifungal agent. Orange powder. Mp


140°. λ_{\max} 320 (log ϵ 4.5); 368 (log ϵ 4.56); 376 (log ϵ 4.57) (no solvent reported).

Wilson, C.V. et al., *J.A.C.S.*, 1948, **70**, 1901-1903 (synth)

Miles, D.H. et al., *Phytochemistry*, 1993, **32**, 1427-1429 (isol, pmr, ms)

3,4-Dimethoxybenzoic acid D-706

Veratric acid. Protocatechuic acid dimethyl ether
[93-07-2]



C₉H₁₀O₄ 182.176
Isol. from *Verbascum thapsus*, *Schoenocaulon officinale*, *Lithospermum* sp., *Stephania abyssinica* and other plants. Widespread in glycosides and alkaloids in esterified form. Isol. from *Aristolochia cucurbitifolia* as the Na salt. Needles (H₂O or AcOH aq.), rhombs by subl. Mp 181-182°. pK_a 4.44 (25°).

► DG8598750

Me ester: [2150-38-1]
C₁₀H₁₂O₄ 196.202
Cryst. (EtOH aq.). Mp 59-60°. Bp 300° Bp₂ 130-131°.

4-Chlorobutyl ester: 4-Chlorobutyl veratrate
[69788-75-6]
C₁₃H₁₇ClO₄ 272.728
Liq. Bp_{0.5} 175-184°.

Chloride: [3535-37-3]
C₉H₉ClO₃ 200.621
Mp 71-73°. Bp_{0.5} 124°.

Amide: [1521-41-1]
C₉H₁₁NO₃ 181.191
Cryst. (EtOH). Mp 164°.

► CV4367000

2-Methylpropylamide: N-Isobutyl-3,4-dimethoxybenzamide. N-(3,4-Dimethoxybenzoyl)isobutylamine
[6302-41-6]
C₁₃H₁₉NO₃ 237.298
Alkaloid from *Piper amalago*. Mp 120-121°.

Anilide: [1522-67-4]
C₁₅H₁₅NO₃ 257.288
Cryst. (EtOH). Mp 154° Mp 162-163°. Mps 141-142 to 172-173 have been recorded.

Nitrile: **4-Cyano-1,2-dimethoxybenzene**
[2024-83-1]
C₉H₉NO₂ 163.176
Mp 68-69°.

Anhydride: [24824-54-2]
C₁₈H₁₈O₇ 346.336
Cryst. (EtOAc or C₆H₆). Mp 124-125°.

[3943-77-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 212A; 299C; 350C; 456D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1110B; 1259A; 1349A; 1532C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1369A; 1418B (ir)

Matsmoto, K.U. et al., *Ber.*, 1878, **11**, 122-141 (synth, Me ester)

Meyer, H. et al., *Monatsh. Chem.*, 1901, **22**, 415-442 (chloride, amide, synth)

Allan, J. et al., *J.C.S.*, 1926, 2334-2336 (anhydride, synth)

Freudenberg, K. et al., *Ber.*, 1941, **74**, 1001-1002 (Me ester, synth)

Wittmer, F.B. et al., *J.O.C.*, 1945, **10**, 527 (synth)

Belg. Pat., 1962, 609 490; *CA*, **59**, 516h (chlorobutyl ester, synth)

Grammaticakis, P. et al., *Bull. Soc. Chim. Fr.*, 1964, 848-858 (amide, synth)

Locksley, H.D. et al., *J.C.S.(C)*, 1970, 392-398 (chloride, synth)

Wiegand, W. et al., *Helv. Chim. Acta*, 1976, **59**, 949 (synth)

de Pascual, T.J. et al., *An. Quim.*, 1978, **74**, 311 (isol)

Suzuki, H. et al., *Chem. Lett.*, 1983, 449-452 (anilide, synth)

Achenbach, H. et al., *Planta Med.*, 1986, **52**, 12-18 (isobutylamide)

Terada, S. et al., *Chem. Pharm. Bull.*, 1987, **35**, 2437-2442 (2-methylpropylamide, synth, ir, pmr)


Suzuki, H. et al., *Synthesis*, 1992, 641 (nitrile)

Wu, T.S. et al., *Chem. Pharm. Bull.*, 2000, **48**, 1006-1009 (isol, Na salt)

Daskiewicz, J.-B. et al., *J. Med. Chem.*, 2005, **48**, 2790-2804 (anhydride)

5,6-Dimethoxy-2(3H)-benzoxazolone, 9CI D-707

5,6-Dimethoxy-2-benzoxazolinone. Peristrophamide
[57601-62-4]




C₉H₉NO₄ 195.174

Alkaloid from *Peristrophe roxburghiana*. Mp 221-222°. λ_{\max} 232 ; 291 (MeOH).

Qin, J.P. et al., *Yaoxue Xuebao*, 1999, **34**, 599-603; *CA*, **132**, 219502 (isol, pmr, cmr)

6,7-Dimethoxy-1(3H)-benzoxazolone D-708

2-Hydroxy-6,7-dimethoxybenzoxazole



C₉H₉NO₄ 195.174

Constit. of the dried tissues of *Zea mays* (sweet corn). Amber cryst. (H₂O). Mp 180°.

4-Chloro-4-Chloro-6,7-dimethoxy-2-benzoxazolinone

C₉H₈ClNO₄ 229.619


Isol. from light-grown sweet corn shoots (*Zea mays*). Auxin-inhibiting substance.

Klun, J.A. et al., *J. Agric. Food Chem.*, 1970, **18**, 663 (isol, struct, synth)

Anai, T. et al., *Phytochemistry*, 1996, **42**, 273-275 (deriv)

4,8-Dimethoxy- β -carboline D-709

4,8-Dimethoxy-9H-pyrido[3,4-b]indole, 9CI. **Picrasidine P** [99964-78-0]




$C_{13}H_{12}N_2O_2$ 228.25

Alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae). Needles (Me_2CO). Mp 198°.

Ohmoto, T. et al., *Chem. Pharm. Bull.*, 1985, **33**, 3847 (isol, uv, ir, pmr, ms, struct)

6,8-Dimethoxy-1,3-dimethylisoquinoline, 9CI D-710

[66178-59-4]



$C_{13}H_{15}NO_2$ 217.267

Alkaloid from bark of *Ancistrocladus tectorius* (Ancistrocladaceae). Needles (Et_2O or cyclohexane). Mp 91° (71-73°, 65-67°).

3,4-Dihydro: see 3,4-Dihydro-6,8-dimethoxy-1,3-dimethylisoquinoline, D-425

1''-Hydroxy: **3-Hydroxymethyl-6,8-dimethoxy-1-methylisoquinoline**

[165816-68-2]

$C_{13}H_{15}NO_3$ 233.266

Alkaloid from bark of *Ancistrocladus tectorius* (Ancistrocladaceae). Cryst. ($MeOH/CH_2Cl_2$). Mp 164°.


Hirota, T. et al., *Chem. Pharm. Bull.*, 1978, **26**, 245 (synth, pmr)

Rizzacasa, M.A. et al., *Aust. J. Chem.*, 1990, **43**, 79 (synth, pmr)

Montagnac, A. et al., *Phytochemistry*, 1995, **39**, 701 (isol, deriv)

5,8-Dimethoxy-2-(3-methoxyphenyl)-3-propyl-4(1H)-quinolinone D-711

4-Hydroxy-5,8-dimethoxy-2-(3-methoxyphenyl)-3-propylquinoline [1010070-87-7]



$C_{21}H_{23}NO_4$ 353.417

Alkaloid from the fruit of *Casimiroa edulis*. Cryst. ($CHCl_3$). Mp 130-131°. λ_{max} 267 (€ 22965); 304 (€ 12220); 330 (€ 9580) (EtOH).

2'-Methoxy: 2-(2,3-Dimethoxyphenyl)-5,8-dimethoxy-3-propyl-4(1H)-quinolinone. 2-(2,3-Dimethoxyphenyl)-4-hydroxy-5,8-dimethoxy-3-propylquinoline [1010070-88-8]

[1048701-05-8]

$C_{22}H_{25}NO_5$ 383.443


Alkaloid from the fruit of *Casimiroa edulis*. Cryst. ($CHCl_3$). Mp 120-121°.

λ_{max} 285 (€ 14450); 310 (€ 8145); 325 (€ 6595) (EtOH).

Awaad, A.S. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 576-580 (isol, pmr, cmr)

5,6-Dimethoxy-2-(3-methoxyphenyl)-4(1H)-quinolinone, 9CI D-712

4-Hydroxy-5,6-dimethoxy-2-(3-methoxyphenyl)quinoline [437659-23-9]



$C_{18}H_{17}NO_4$ 311.337

Alkaloid from the leaves of *Casimiroa edulis*. Amorph. powder. Mp 111-114°. λ_{max} 270 (€ 14300); 302 (€ 8140); 324 (€ 6590) (MeOH).

4'-Methoxy: 2-(3,4-Dimethoxyphenyl)-5,6-dimethoxy-4(1H)-quinolinone, 9CI. 2-(3,4-Dimethoxyphenyl)-4-hydroxy-5,6-dimethoxyquinoline [437659-24-0]

$C_{19}H_{19}NO_5$ 341.363

Alkaloid from the leaves of *Casimiroa edulis*. Amorph. powder. Dec. >100°. λ_{max} 266 (€ 22960); 302 (€ 12215); 332 (€ 9580) (MeOH).

2',6'-Dimethoxy: 5,6-Dimethoxy-2-(2,3,6-trimethoxyphenyl)-4(1H)-quinolinone. 4-Hydroxy-5,6-dimethoxy-2-(2,3,6-trimethoxyphenyl)quinoline [437659-25-1]

$C_{20}H_{21}NO_6$ 371.389


Alkaloid from the fruit and leaves of *Casimiroa edulis*. Needles (hexane/EtOAc). Mp 156-157°. λ_{max} 250 (€ 11160); 256 (€ 11145); 328 (€ 5470) (MeOH).

Khaleel, A.E.M. et al., *Monatsh. Chem.*, 2002, **133**, 183-187 (isol, pmr, cmr, ms)

Awaad, A.S. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 576-580 (isol)

5,7-Dimethoxy-4-(4-methoxyphenyl)-2(1H)-quinolinone, 9CI D-713

[132905-26-1]



$C_{18}H_{17}NO_4$ 311.337

Alkaloid from the roots of *Chiococca alba* (Rubiaceae). Mp 300°.

8-Hydroxy: **8-Hydroxy-5,7-dimethoxy-4-(4-methoxyphenyl)-2(1H)-quinolinone, 9CI**

[132905-27-2]

$C_{18}H_{17}NO_5$ 327.336


Alkaloid from the roots of *Chiococca alba* (Rubiaceae). Cryst. ($CH_2Cl_2/EtOAc$). Mp 237-239°.

El Abbadi, N. et al., *Planta Med.*, 1989, **55**, 603 (isol)

Kitahara, Y. et al., *Heterocycles*, 1990, **31**, 2085 (synth, ir, uv, pmr, cmr)

7,9-Dimethoxy-2,3-methylenedioxybenzo[c]phenanthridine D-714

[252894-93-2]




$C_{20}H_{15}NO_4$ 333.343

Alkaloid from the stem bark of *Zanthoxylum myriacanthum*. Cryst. ($CHCl_3$). Mp 217-219°.

Sukari, M.A. et al., *Fitoterapia*, 1999, **70**, 197-199 (isol, ir, pmr, cmr)

2,4-Dimethoxy-7,8-methylenedioxy-3-prenylquinoline D-715

6,8-Dimethoxy-7-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-h]quinoline, 9CI. **Isopteleprenine** [17232-49-4]



$C_{17}H_{19}NO_4$ 301.341

Alkaloid from stems of *Orixa japonica* (Rutaceae). Oil. Bp_{0.05} 160-170° (bath).

2'R,3'-Epoxide: **Preorixine** [17232-51-8]

$C_{17}H_{19}NO_5$ 317.341

Alkaloid from stems of *Orixa japonica*. $[\alpha]_D^{25}$ -7.5 (c, 0.8 in $CHCl_3$).

2',3'-Dihydroxy, 2',3'-dihydro: see Orixine, O-113

4'3'-Isomer, 2'-hydroxy: 3-(2-Hydroxy-3-methyl-3-butenyl)-2,4-dimethoxy-7,8-methylenedioxyquinoline. **Isoptelefolidine** [156953-83-2]

$C_{17}H_{19}NO_5$ 317.341


Alkaloid from stems of *Orixa japonica* (Rutaceae). Oil. $[\alpha]_D^{27}$ +9.8 (c, 0.21 in $CHCl_3$).

Bowman, R.M. et al., *J.C.S.(C)*, 1967, 2368 (synth)

Funayama, S. et al., *Phytochemistry*, 1994, **36**, 525 (Isopteleprenine, Isoptelefolidine)

Funayama, S. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1885-1889 (*Preorixine*)


3,6-Dimethoxy-1-methyl-2-phenyl-4(1H)-quinolinone, 9CI **D-716**
Japonine
[30426-61-0]



$C_{18}H_{17}NO_3$ 295.337
Alkaloid from the leaves of *Orixa japonica* (Rutaceae). Prisms (Et₂O/MeOH). Mp 143°.

Hä-Huy-Kê, *et al.*, *Phytochemistry*, 1970, **9**, 2199 (*isol, uv, ir, pmr, ms, struct*)
Venturella, P. *et al.*, *Heterocycles*, 1976, **4**, 1089 (*synth*)


3-(1,1-Dimethoxy-2-oxoethoxy)-9,10-dihydro-1-methoxy-9-oxo-4-acridinecarboxaldehyde, 9CI **D-717**
1-Methoxy-3-(2-methylpropanal-2-oxo)acridin-9-one-4-carbaldehyde
[110883-40-4]



$C_{19}H_{17}NO_5$ 339.347
Alkaloid from the leaves of *Sarcomelicope dogntiensis* (Rutaceae). Yellow amorph. solid.

Mitaku, S. *et al.*, *Heterocycles*, 1987, **26**, 2057 (*uv, ir, pmr, struct*)

3,4-Dimethoxyphenethylamine, 8CI **D-718**
3,4-Dimethoxybenzeneethanamine, 9CI. Homoveratrylamine. 4-(β-Aminoethyl)-veratrole. O,O-Dimethyldopamine
[120-20-7]



$C_{10}H_{15}NO_2$ 181.234
Isol. from *Desmodium tiliaefolium*, *Pachycereus pecten-aboriginum*, *Trichocereus peruvianus* and other *Trichocereus* spp., *Echinocereus merkeri* and mescal (*Lophophora williamsii*) (Fabaceae, Cactaceae). Cryst. (C₆H₆/petrol). Mp 124°. Bp₁₅ 188°.

▶ SH2300000

Hydrochloride: [635-85-8]
Mp 154-155°.

▶ LD₅₀ (rat, ipr) 146 mg/kg. SH3000000
N-Formyl: [14301-36-1]
 $C_{11}H_{15}NO_3$ 209.244
Mp 40-42°. Bp_{0,01} 170°.

N-Ac: *N-Acetylhomoveratrylamine. N-Acetyl-3,4-dimethoxyphenethylamine*
[6275-29-2]
 $C_{12}H_{17}NO_3$ 223.271
Alkaloid from *Berberis sibirica* (Berberidaceae). Cryst. (C₆H₆). Mp 104-105°.

N-(2-Hydroxy-5-methoxybenzoyl): *Taiwanamide A*
[1000991-96-7]
 $C_{18}H_{21}NO_5$ 331.368
Alkaloid from the stems of *Piper taiwanense*. Needles (hexane). Mp 104-105°. λ_{max} 215 (log ε 4.81); 280 (log ε 4.1); 325 (log ε 4.15) (MeOH).

N-(2,5-Dimethoxybenzoyl): *Taiwanamide B*
[1000991-97-8]
 $C_{19}H_{23}NO_5$ 345.394
Alkaloid from the stems of *Piper taiwanense*. Oil. λ_{max} 217 (log ε 4.77); 281 (log ε 4.02); 314 (log ε 4.03) (MeOH).

N-(3,4-Dimethoxy-E-cinnamoyl): *Rubemamine*
[121817-65-0]
 $C_{21}H_{25}NO_5$ 371.432
Alkaloid from the stem bark of *Zanthoxylum rubescens* (Rutaceae). Mp 121-122°.

N-Me: *3,4-Dimethoxy-N-methylphenethylamine*
[3490-06-0]
 $C_{11}H_{17}NO_2$ 195.261
Isol. from *Pilosocereus guerreronis*, *Coryphantha* spp., *Echinocereus merkeri* and *Ariocarpus* spp. (Cactaceae). Oil. Bp_{0,8} 120°.

N-Me; hydrochloride: [13078-76-7]
Cryst. (EtOH). Mp 134-135°.

▶ SH3237000

N-Me, N-(3,4-dimethoxy-E-cinnamoyl): *Rubemamide*
[121817-66-1]
 $C_{22}H_{27}NO_5$ 385.459
Alkaloid from the stem bark of *Zanthoxylum rubescens* (Rutaceae).

N,N-Di-Me: *N,N-Dimethyl-3,4-dimethoxyphenethylamine*
[3490-05-9]
 $C_{12}H_{19}NO_2$ 209.288
Alkaloid from *Echinocereus merkeri*, *Echinocereus cinerascens*, *Pilosocereus guerreronis* and *Ariocarpus scapharostus* (Cactaceae). Mp 193-196° (as hydrochloride).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1292B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 619B; 619C; 620A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1175A (*ir*)

Org. Synth., Coll. Vol., **3**, 1955, 720 (*synth*)
Viel, C. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 431-440 (*synth*)

Agurell, S. *et al.*, *J. Pharm. Sci.*, 1969, **58**, 1413-1414 (*isol, struct, ms, derivs*)

Ghosal, S. *et al.*, *Phytochemistry*, 1973, **12**, 193-197 (*isol*)

Ranieri, R. *et al.*, *J. Nat. Prod.*, 1976, **39**, 172-174 (*isol, N-Me*)

Bruhn, J.G. *et al.*, *J. Nat. Prod.*, 1976, **39**, 175-177 (*isol*)

Lindgren, J.E. *et al.*, *J. Nat. Prod.*, 1976, **39**, 464-466 (*isol, struct, derivs*)

Pardanani, J.H. *et al.*, *J. Nat. Prod.*, 1977, **40**, 585-590 (*isol*)

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9-18 (*rev, occur*)

Bruhn, J.G. *et al.*, *Phytochemistry*, 1977, **16**, 622-623 (*isol, derivs*)

Kihara, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 155-160 (*synth*)

Mata, R. *et al.*, *J. Pharm. Sci.*, 1980, **69**, 94 (*isol*)

Adesina, S.K. *et al.*, *Phytochemistry*, 1989, **28**, 839 (*Rubemamine, Rubemamide*)

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 424-428; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 361 (*N-Acetylhomoveratrylamine, isol*)


Org. Synth., 1999, **76**, 133-141 (*N,N-di-Me, synth, ir, pmr*)

Chen, I.-S. *et al.*, *Fitoterapia*, 2007, **78**, 414-419 (*Taiwanamides A, B*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van

Nostrand Reinhold, 1992, DOE200; DOI400

N-[2-(2,4-Dimethoxyphenyl)ethenyl]-3-pyridinecarboxamide, 9CI **D-719**
N-(2,4-Dimethoxystyryl)-3-pyridinecarboxamide



$C_{16}H_{16}N_2O_3$ 284.314

(E)-form [69142-85-4]

Alkaloid from *Amyris plumieri* (Rutaceae). Mp 159-160°.

α,β-Dihydro: Mp 112-113°.


(Z)-form [95690-61-2]

Alkaloid from leaves and twigs of *Amyris plumieri* (Rutaceae). Pale-yellow plates (MeOH). Mp 124.5-125.5°.

Burke, B.A. *et al.*, *Tet. Lett.*, 1978, 2723 (*isol, uv, ir, pmr, struct*)

Burke, B.A. *et al.*, *Heterocycles*, 1985, **23**, 257 (*isol, uv, ir, pmr, struct*)

7,8-Dimethoxyplatydesmine **D-720**



$C_{17}H_{21}NO_5$ 319.357

(R)-form [142741-27-3]

Alkaloid from *Dictamnus dasycarpus* and *Dutaillaea baudouinii*. λ_{max} 230 ; 238 (sh) ; 262 ; 283 ; 307 ; 320 (EtOH).

(S)-form

Alkaloid from *Melicope semecarpifolia*.

[α]_D²³ -10.3 (c, 0.16 in CHCl₃).

N-Me: *Veprisinium*

[79808-98-3]

$C_{18}H_{24}NO_5^{\oplus}$ 334.391

Quaternary alkaloid from the stem bark of *Vepris louisii* (Rutaceae). Shows antibacterial props. λ_{\max} 205 (ϵ 24700); 222 (ϵ 46700); 255 (ϵ 49500); 327 (ϵ 15400) (MeOH) (Derep).

N-Me, chloride:

Amorph. powder. $[\alpha]_D^{21}$ -14.2 (c, 0.6 in MeOH).

N-Me, perchlorate:

Needles (MeOH/Et₂O). Mp 171-172°. $[\alpha]_D^{21}$ -11.8 (c, 0.60 in MeOH).

(ξ)-form

Alkaloid from *Melicope semecarpifolia*.

1'-Deoxy, 1',2'-didehydro, N-Me: 2,3-Dihydro-4,7,8-trimethoxy-9-methyl-2-(1-methylethenyl)furo[2,3-b]quinolinium, 9CI. **Isoptelefolonium** [60593-03-5]

C₁₈H₂₂NO₄⁺ 316.376

Alkaloid from the leaves of *Ptelea trifoliata* (Rutaceae). Mp 146-148° (as perchlorate). $[\alpha]_D^{20}$ +9.83 (c, 0.006 in MeOH).

Rideau, M. et al., *Phytochemistry*, 1979, **18**, 155 (*Isoptelefolonium*)

Ayafor, J.F. et al., *Planta Med.*, 1982, **44**, 139-142 (*isol, uv, ir, pmr, ms, cd, struct*)


Muyard, F. et al., *Phytochemistry*, 1992, **31**, 1087-1089 (*isol, uv, ir, pmr, ms, struct*)

Wu, J.F. et al., *Chin. J. Phys. (Taipei)*, 1994, **46**, 447-455 (*isol*)

Chen, I.-S. et al., *J. Nat. Prod.*, 2001, **64**, 1143-1147 (*isol, pmr, cmr*)

(2,4-Dimethoxy-3-prenylcinnamoyl)piperidine D-721

1-[3-[2,4-Dimethoxy-3-(3-methyl-2-butenyl)phenyl]-1-oxo-2-propenyl]piperidine, 9CI. 2,4-Dimethoxy-3-(3-methyl-2-butenyl)cinnamic acid piperidide



C₂₁H₂₉NO₃ 343.465


(E)-form [88053-60-5]

Alkaloid from the stemwood of *Excoecaria agallocha* (Euphorbiaceae). Oil.

Prakash, S. et al., *Phytochemistry*, 1983, **22**, 1836 (*isol, uv, ir, pmr, ms, struct, synth*)

2,3-Dimethoxypyrido[2,3-b]pyrazine, 9CI D-722

[103518-15-6]




C₉H₉N₃O₂ 191.189

Isol. from the lipid extract of the spring-tail *Tetradontophora bielansensis*. Needles. Mp 125-125.5°.

Buděšínský, M. et al., *Coll. Czech. Chem. Comm.*, 1986, **51**, 956 (*isol, ms, pmr, struct*)

4,7-Dimethoxy-1-vinyl- β -carboline D-723

1-Ethenyl-4,7-dimethoxy-9H-pyrido[3,4-b]indole [33867-64-0]



C₁₅H₁₄N₂O₂ 254.288

Alkaloid from *Perriera madagascariensis* (Simaroubaceae). Mp 145-147°. λ_{\max} 226 (log ϵ 4.37); 255 (log ϵ 4.4); 284 (log ϵ 4.36); 311 (log ϵ 4.1); 351 (log ϵ 3.93) (no solvent reported).

Bourguignon-Zylber, N. et al., *Eur. J. Med. Chem. (Chim. Ther.)*, 1970, **5**, 396; *CA*, **75**, 31236j

Krebs, H.C. et al., *J. Nat. Prod.*, 1997, **60**, 1183-1185 (*isol, uv, pmr, cmr*)

4,9-Dimethoxy-1-vinyl- β -carboline D-724

1-Ethenyl-4,9-dimethoxy-9H-pyrido[3,4-b]indole, 9CI. **Picrasidine D** [88142-62-5]


C₁₅H₁₄N₂O₂ 254.288

Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Needles (Me₂CO). Mp 191-192°.

Ohmoto, T. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3198; 1984, **32**, 3579 (*isol, uv, ir, pmr, ms, struct, nomencl*)

Dimethyl N²-creatininylphosphate D-725

[145644-08-2]



C₆H₁₂N₃O₄P 221.152

Isol. from the sponge *Ulosa ruetzleri*. Mp 154-155°.

Van Wagenen, B.C. et al., *J.O.C.*, 1993, **58**, 335 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Dimethylamine D-726

N-Methylmethanamine, 9CI

[124-40-3]

Me₂NH

C₂H₇N 45.084

Found in higher plants, fungi and bacteria. Rubber vulcanisation accelerator, tanning agent. Reagent used in the ms determination of the location of ethylenic bonds. Aminating agent in the manuf. of ion-exchange resins. Gas at r.t. V. sol. H₂O, sol. EtOH, Et₂O. d₄²⁰ 0.68. Mp -96°. Bp 7°. pK_a 3.28 (25°).

► Extremely flammable, fl. p. -6°, auto-ignition temp. 400°. Eye, skin and respiratory tract irritant (gas or aq. solns.) Prolonged exposure can cause dermatitis and conjunctivitis. IP8750000

Hydrochloride: Dimethylammonium chloride

[506-59-2]

C₂H₈ClN 81.545

Volatile cryst. Sol. H₂O, EtOH, CHCl₃; insol. Et₂O. Mp 171°.

► IQ0220000

DMSO complex (1:1): Dimsulf

C₄H₁₃NOS 123.219

Powerful absorbent for H₂S. Distillable liq.

[17000-01-0]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 294B; 294C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 470A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 378C (*ir*)

Werner, E.A. et al., *J.C.S.*, 1917, **111**, 850 (*synth*)

Ger. Pat., 1929, 468 895; *CA*, **23**, 846 (*synth*)

U.S. Pat., 1948, 2 456 599; *CA*, **43**, 3440g (*manuf*)

Gohlke, R.S. et al., *Acta Chim. Hung.*, 1962, **34**, 1281 (*ms*)

Suhr, H. et al., *Chem. Ber.*, 1963, **96**, 1720 (*pmr*)

Serban, S. et al., *Rev. Chim. (Bucharest)*, 1963, **14**, 451; *CA*, **60**, 5097b (*synth*)

Audier, H. et al., *Bull. Soc. Chim. Fr.*, 1964, 3034 (*use*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1979, 7, 119 (*use*)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 846-847

Chemical Hazards of the Workplace, 2nd edn., (eds. Proctor, N.H. et al.), J.B. Lippincott, 1988, 207

Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn., (ed. Snyder, R.), Elsevier, Volume 2, 1990, 73 (*tox*)


Bretherick, L. et al., *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 0887

Luxon, S.G. et al., *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 492

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DOQ800; DOR000; DOR600

3-Dimethylaminoacetyl-5-methoxyindole D-727

2-(Dimethylamino)-1-(5-methoxy-1H-indol-3-yl)ethanone, 9CI. 5-Methoxy-N,N-dimethyl- β -oxotryptamine [42883-65-8]




C₁₃H₁₆N₂O₂ 232.282

Alkaloid from the aerial parts of *Melicope leptococca* (Rutaceae).

Skaltsounis, A.L. et al., *J. Nat. Prod.*, 1983, **46**, 732-735 (*isol, uv, ir, pmr, ms, struct*)

2-(Dimethylamino)-1,5-dihydro-5-(1H-indol-3-yl)-5-(2-oxo-propyl)-4H-imidazol-4-one, 9CI
[205117-53-9]



C₁₆H₁₈N₄O₂ 298.344


Isol. from the marine tunicate *Dendrodoa grossularia*. Powder. Mp 265-266°. [α]_D -15 (c, 0.14 in MeOH).

Loukaci, A. et al., *J. Nat. Prod.*, 1998, **61**, 519-522 (isol, pmr, cmr)

Hupp, C.D. et al., *Org. Lett.*, 2008, **10**, 3737-3739 (synth, cryst struct)

α -(Dimethylamino)-3,5-diiodo-N-[5-[[3-(3-iodo-4-methoxyphenyl)-2-(methylamino)-1-oxopropyl]aminopentyl]-4-methoxybenzenepropanamide, 9CI

3-(3,5-Diiodo-4-methoxyphenyl)-2-dimethylamino-3'-(3-iodo-4-methoxyphenyl)-2'-methylamino-N,N'-(1,5-pentanediy)bispropanamide
[149196-93-0]



C₂₈H₃₉I₃N₄O₄ 876.354

Alkaloid from the colonial ascidian *Aplidium* sp. Glutathione reductase inhibitor. Gum. [α]_D -0.74 (c, 0.24 in CHCl₃). λ_{\max} 210 (ϵ 40000); 222 (ϵ 39000); 278 (ϵ 9200) (EtOH) (Berdy).

N²⁹-Me: 3-(3,5-Diiodo-4-methoxyphenyl)-3'-(3-iodo-4-methoxyphenyl)-N,N'-(1,5-pentanediy)bis[2-dimethylaminopropanamide]

[149196-91-8]

C₂₉H₄₁I₃N₄O₄ 890.381

From *Aplidium* sp. Cytotoxic. Shows inhibition of glutathione reductase activity. Needles (CHCl₃/EtOAc). Mp 135-137°. [α]_D -0.23 (c, 0.51 in CHCl₃).

4-Iodo, N²⁹-Me: N,N'-(1,5-Pentanediy)bis[3-(3,5-diiodo-4-methoxyphenyl)-2-dimethylaminopropanamide]

[149196-92-9]

C₂₉H₄₀I₄N₄O₄ 1016.277

From *Aplidium* sp. Cytotoxic. Shows inhibition of glutathione reductase activity. Needles (CHCl₃/EtOAc). Mp 114-116°. [α]_D -0.2 (c, 0.35 in CHCl₃). λ_{\max} 207 (ϵ 32000); 275 (ϵ 8400) (EtOH) (Berdy).

Carroll, A.R. et al., *Aust. J. Chem.*, 1993, **46**, 825-832 (isol, uv, ir, pmr, cmr, ms, struct)

2-(Dimethylamino)ethanol D-730

Dimethyl-2-hydroxyethylamine. N-Dimethylethanolamine. **Deanol**, **BAN**.

Norcholine. **Bimanol**

[108-01-0]

Me₂NCH₂CH₂OH

C₄H₁₁NO 89.137

Isol. from a *Neurospora crassa* strain. Residue present in the alkaloids Cassaine in C-178 and Cassaidine, C-178. Choline precursor. CNS stimulant. Antidepressant. d_4^{20} 0.89. Bp 135°. Log P -0.58 (calc).

► Skin and eye irritant. Flammable, fl. p. 41° (oc), autoignition temp. 295°.

KK6125000

4-(Acetylamino)benzoate salt (1:1):

Deaner. *Diforene*. *Deanol acetamidobenzoate*. Many other names

[3635-74-3]

Antidepressant. Cryst. (EtOH/EtOAc). Mp 159-161.5°.

► LD₅₀ (mus, orl) 39.8 mg/kg. KK6300000

N-Acetyl-L-glutamate salt (1:1): *Deanol aceglumate*, *INN*. *Demanol aceglumate*. *Cleregil*. *Dardanin*. *Otrun*. *Risaratarun*. *Rischiaril*

[3342-61-8]

Antidepressant.

► LZ9800000

Picrate: Mp 96-97°.

O-Phosphate:

C₄H₁₂NO₄P 169.117

Isol from *Neurospora crassa*. Mp 175-176°.

Ac: [1421-89-2]

C₆H₁₃NO₂ 131.174

Bp₈₀ 86-88°.

2-Propenoyl: [2439-35-2]

C₇H₁₃NO₂ 143.185

Bp₁₂ 64°. n_D^{20} 1.4380.

► AS8578000

O-Benzoyl: 2-(Dimethylamino)ethyl benzoate

[2208-05-1]

C₁₁H₁₅NO₂ 193.245

d 1.01, Mp 144° (as hydrochloride).

Bp₂₀ 155-159° Bp₂ 107-108°. n_D^{20} 1.5077.

O-Cinnamoyl: 2-Dimethylaminoethyl cinnamate

[4067-25-8]

[35241-68-0, 35241-67-9]

C₁₃H₁₇NO₂ 219.283

Alkaloid from leaves of *Erythrophleum chlorostachys* (Fabaceae). Gum; cryst. (EtOH)(as hydrobromide). Mp 130° (as hydrobromide)(synthetic).

Me ether: 2-Methoxy-N,N-dimethylethanolamine, 9CI. 2-Methoxy-N,N-dimethylethylamine

[3030-44-2]

C₅H₁₃NO 103.164

Isol. from *Didemnum ligulum*. Bp₇₅₇ 101°.

Et ether: 2-Ethoxy-N,N-dimethylethanolamine, 9CI

[26311-17-1]

C₆H₁₅NO 117.191

Liq. d_4^{20} 0.81. Mp 160-165° (as methiodide) Mp 119-121° (as picrate). Bp 121°.

[29870-28-8]

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 342A (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **1**, 545A; 1073C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 432A; 734A (ir)

Knorr, L. et al., *Ber.*, 1901, **34**, 3482 (synth)

Fränkel, S. et al., *Ber.*, 1918, **51**, 1660 (synth, derivs)

U.K. Pat., 1957, 879 259; CA, **56**, 14080 (derivs) Schaezle, E. et al., *Helv. Chim. Acta*, 1959, **42**, 1708 (benzoyl)

Wolf, B. et al., *J. Biol. Chem.*, 1959, **234**, 1068 (phosphate)

Fr. Pat., 1964, M2487; CA, **61**, 12085c (deriv)

Griffin, W.J. et al., *Phytochemistry*, 1971, **10**, 2793 (cinnamoyl)

Štokr, J. et al., *Coll. Czech. Chem. Comm.*, 1987, **52**, 1256 (pmr, cmr, conformn)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 244

Jaszay, Z.M. et al., *Synthesis*, 1989, 745 (benzoyl)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1223

Kossuga, M.H. et al., *Quim. Nova*, 2007, **30**, 1194-1202 (*Me ether*, isol, pmr, cmr)

Luxon, S.G. et al., *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 493

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DOY800; DOZ000


4-[2-(Dimethylamino)ethyl]imidazole, 8CI

D-731

N,N-Dimethyl-1H-imidazole-4-ethanamine, 9CI. N^z,N^z-Dimethylhistamine.

Hippospongine†

[673-46-1]



C₇H₁₃N₃ 139.2

The application of the name Hippospongine to this compound is tentative. Alkaloid from *Echinocereus blanckii*, *Echinocereus triglochidiatus*, *Echinocereus triglochidiatus neomexicanus*, *Casimiroa edulis* seeds (Cactaceae, Rutaceae), the sponges *Geodia gigas* and *Ianthella* sp., and from skin extracts of amphibians, e.g. *Leptodactylus pentadactylus labyrinthicus* and *Nyctimystes disrupta* and from basidiomycete *Coprinus comatus*.

► NI4840000

Hydrochloride (1:2): Mp 182-184°.

Dipicrate: Mp 229-230°.

Ackermann, D. et al., *Z. Biol. (Munich)*, 1924, **82**, 278 (isol)

Garforth, B. et al., *J.C.S.*, 1935, 489 (synth)

Huebner, C.F. et al., *J.A.C.S.*, 1949, **71**, 3942 (synth)

List, P.H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1958, **291**, 502 (isol)

Major, R.T. et al., *J.O.C.*, 1958, **23**, 1564 (isol)

Ling, J.S.L. et al., *J. Pharmacol. Exp. Ther.*, 1958, **122**, 44A (pharmacol)


Erspamer, V. et al., *Experientia*, 1963, **19**, 346 (occur)

Ingle, P.H.B. et al., *J. Pharm. Pharmacol.*, 1963, **15**, 620 (synth)

Erspamer, V. et al., *Arch. Biochem. Biophys.*, 1964, **105**, 620 (occur)


- German, V.F. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 495 (*isol, pmr, ms*)
 Roseghini, M. *et al.*, *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1976, **54**, 31 (*occur*)
 Roseghini, M. *et al.*, *Z. Naturforsch., C*, 1976, **31**, 118 (*occur*)
 Ferrigni, N.R. *et al.*, *J. Ethnopharmacol.*, 1982, **5**, 359 (*isol*)
 Wagner, H. *et al.*, *Planta Med.*, 1982, **45**, 95 (*isol*)
 Romero, M.L. *et al.*, *J. Chromatogr.*, 1983, **281**, 245 (*hplc*)
 Ferrigni, N.R. *et al.*, *Rev. Latinoam. Quim.*, 1984, **14**, 131; *CA*, **101**, 23790a (*pmr, cmr*)

2-(Dimethylamino)-5-(1H-indol-3-yl)-4H-imidazol-4-one, 9CI
 [103590-22-3]




- $C_{13}H_{12}N_4O$ 240.264
 First report of a 4H-imidazol-4-one as a nat. prod. Isol. from the tunicate *Dendrodoa grossularia*. Orange needles (MeOH). Mp 356-358°. Sensitive to light in soln. λ_{max} 213 (ϵ 15300); 261 (ϵ 7140); 284 (ϵ 6010); 343 (ϵ 5120); 430 (ϵ 5360) (EtOH) (Derep).
 Guyot, M. *et al.*, *Tet. Lett.*, 1986, **27**, 2621-2622 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)
 Dalkafouki, A. *et al.*, *Tet. Lett.*, 1991, **32**, 5325 (*synth*)

3-[(Dimethylamino)methyl]-5,7-dimethoxyindole
 5,7-Dimethoxy-N,N-dimethyl-1H-indole-3-methanamine, 9CI. 5,7-Dimethoxygramine
 [24450-47-3]



- $C_{13}H_{18}N_2O_2$ 234.297
 Alkaloid from the roots of *Phalaris aquatica* (Poaceae). Cryst. (C_6H_6 /petrol or EtOH aq.). Mp 121-122° (119-120°).
 Crohare, R. *et al.*, *J. Het. Chem.*, 1970, **7**, 729 (*synth, pmr*)
 Mulvena, D.P. *et al.*, *Phytochemistry*, 1983, **22**, 2885 (*isol, pmr*)

3-[(Dimethylamino)methyl]-5-hydroxyindole
 3-[(Dimethylamino)methyl]-1H-indol-5-ol, 9CI. 5-Hydroxygramine
 [83662-85-5]



- $C_{11}H_{14}N_2O$ 190.244
 Mp 198-199°.
 Hydrochloride:
 Polyhedra ($CHCl_3$). Sl. sol. H_2O , MeOH, EtOH; v. sol. Et_2O . Mp 198-199° (same value as reported for free base; possibly erroneous).
 Me ether: 3-(Dimethylaminomethyl)-5-methoxyindole. 5-Methoxygramine [16620-52-3]
 $C_{12}H_{16}N_2O$ 204.271
 Alkaloid detected in the roots of *Phalaris aquatica* (Gramineae). Needles (C_6H_6). Mp 128° (124-125°).

- Me ether, picrate:
 Amber prisms (C_6H_6 /EtOH). Mp 168° (164-165°).
 Et ether: 3-(Dimethylaminomethyl)-5-ethoxyindole
 $C_{13}H_{18}N_2O$ 218.298
 Needles (Me_2CO aq.). Mp 146°.
 Et ether, hydrochloride:
 Prisms (EtOH/ Et_2O). Mp 150°.
 Benzyl ether: [1453-97-0]
 $C_{18}H_{20}N_2O$ 280.369
 Cryst. (C_6H_6 /hexane). Mp 138.5-139.5°. pK_a 16.9.

- N-Methoxy, Me ether: 1,5-Dimethoxy-N,N-dimethyl-1H-indole-3-methanamine. 3-[(Dimethylamino)methyl]-1,5-dimethoxyindole, 8CI. 1,5-Dimethoxygramine [16620-50-1]
 $C_{13}H_{18}N_2O_2$ 234.297
 Major alkaloid from *Gymnacranthera paniculata* (Myristicaceae). Oil.
 N-Methoxy, Me ether, picrate:
 Red or golden-yellow cryst. (EtOH). Mp 154-155°.


- ar, N-Dimethoxy, Me ether: 3-(Dimethylaminomethyl)trimethoxyindole. ar-Tri-methoxygramine
 $C_{14}H_{20}N_2O_3$ 264.324
 Minor alkaloid from *Gymnacranthera paniculata* (Myristicaceae). Not. obt. free of the major component. A 1,ar,ar-trimethoxy compd. of undetd. regiochemistry.

- Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 662C (Me ether, ir)
 Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 134A (Me ether, nmr)
 Davies, R.E. *et al.*, *J.C.S.*, 1947, 191 (Et ether)
 Bell, J.B. *et al.*, *J.O.C.*, 1948, **13**, 547 (Me ether, synth)
 Cook, J.W. *et al.*, *J.C.S.*, 1951, 1203 (Me ether)
 Ek, A. *et al.*, *J.A.C.S.*, 1954, **76**, 5579 (benzyl ether, ir)
 Troxler, F. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 2073 (synth)
 Koo, J. *et al.*, *J.O.C.*, 1959, **24**, 179 (benzyl ether)
 Marchand, B. *et al.*, *Chem. Ber.*, 1962, **95**, 577 (synth)
 Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1737-1742 (1,5-Dimethoxygramine, Trimethoxygramine, Me ether, isol, uv, struct, synth, pmr, ms)
 Hojo, T. *et al.*, *J. Chromatogr.*, 1982, **247**, 157 (chromatog)
 Mulvena, D.P. *et al.*, *Phytochemistry*, 1983, **22**, 2885 (Me ether, occur)

3-[(Dimethylamino)methyl]-7-hydroxyindole
 3-[(Dimethylamino)methyl]-1H-indol-7-ol, 9CI. 7-Hydroxygramine
 $C_{11}H_{14}N_2O$ 190.244

- Me ether: 7-Methoxy-N,N-dimethyl-1H-indole-3-methanamine, 9CI. 3-[(Dimethylamino)methyl]-7-methoxyindole. 7-Methoxygramine [46388-50-5]
 $C_{12}H_{16}N_2O$ 204.271
 Alkaloid from the roots of *Phalaris aquatica* (Poaceae). Cryst. (C_6H_6 /petrol). Mp 107-109°.
 Mulvena, D.P. *et al.*, *Phytochemistry*, 1983, **22**, 2885 (*isol, pmr, struct*)

3-[(Dimethylamino)methyl]-indole
 N,N-Dimethyl-1H-indole-3-methanamine, 9CI. Gramine. Donaxine. Doranine
 [87-52-5]



- $C_{11}H_{14}N_2$ 174.245
 Alkaloid from *Arundo donax*, *Phalaris arundinacea*, some *Lupinus* spp. and barley shoots. Also from *Acer saccharinum*, *Acer rubrum*, *Brassica oleracea* and *Desmodium pulchellum*. Antifouling agent, barnacle repellent. Mp 138-139°. pK_a 16 (25°/NH, aq. KOH).
 ▶ LD₅₀ (mus, ipr) 122 mg/kg. NL7525000
 Picrate: Mp 144-145°.

- N-Oxide: Gramine N-oxide [17206-03-0]
 $C_{11}H_{14}N_2O$ 190.244
 Alkaloid from *Arundo donax* (Poaceae). Noncryst. Mp 122° (H_2O_2 solvate).
 N²-Me: Gramine methosalt [5457-31-8]
 $C_{12}H_{17}N_2^{\oplus}$ 189.28
 Alkaloid from flowers of *Arundo donax* (Poaceae). Mp 168-169° (iodide). CAS no. refers to iodide.


- N⁴-(1,1-Dimethyl-2-propenyl): 3-(Dimethylaminomethyl)-1-(1,1-dimethyl-2-propenyl)-1H-indole [150226-14-5]
 $C_{16}H_{22}N_2$ 242.363
 Prod. by *Penicillium daleae*. 5-HT receptor agonist, shows antimigraine props. Sol. MeOH. λ_{max} 223 (ϵ 12100); 273 (ϵ 4890); 279 (ϵ 5090); 289 (ϵ 4120) (MeOH) (Derep). λ_{max} 223 (ϵ 12100); 273 (ϵ 4890); 279 (ϵ 5090); 289 (ϵ 4120) (MeOH) (Berdy). λ_{max} 219 (E1%/1cm 500); 267 (E1%/1cm 118); 279 (E1%/1cm 114); 290 (E1%/1cm 93) (EtOH) (Berdy).
 [6170-37-2]

- Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 131B (nmr)
 Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 660B (ir)

- Kühn, H. *et al.*, *Ber.*, 1937, **70**, 567 (*synth*)
 Marion, L. *et al.*, *J.A.C.S.*, 1951, **73**, 305 (*ir*)
 Bild, N. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 1885 (*ms, oxide*)
 Jamieson, W.D. *et al.*, *Phytochemistry*, 1970, **9**, 2029 (*ms*)
 Ghosal, S. *et al.*, *Phytochemistry*, 1971, **10**, 2852 (*Gramine methosalt*)
 Leete, E. *et al.*, *Phytochemistry*, 1975, **14**, 471 (*biosynth*)
 U.S. Pat., 1993, 5 233 050; *CA*, **119**, 179344 (*isol, deriv*)
 Lam, Y.K.T. *et al.*, *J. Antibiot.*, 1994, **47**, 724 (*isol, deriv*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DYC000

1-[2-(Dimethylamino)-1-oxo-3-phenylpropyl]-2,3,3a,13a,14,15,16,18a-octahydro-8-methoxy-5,9-metheno-9H-dipyrrolo[3,2-b:1',2'-e][1,5,8]oxadiazacyclopentadecine-13,18(1H,12H)-dione, 9CI

[147471-64-5]




C₃₀H₃₆N₄O₅ 532.638

Alkaloid from flowers of *Sphaeranthus indicus* (Asteraceae). Mp 75°.

Chughtai, M.I.D. *et al.*, *Sci. Int. (Lahore)*, 1992, **4**, 151; *CA*, **118**, 230141f (*isol*)

2-Dimethylamino-1-phenyl-1-propanol

α -[1-(Dimethylamino)ethyl]benzenemethanol, 9CI
 [17605-71-9]



(1R,2S)-form

C₁₁H₁₇NO 179.261

See also references under 2-(Methylamino)-1-phenyl-1-propanol, M-386. Log P 1.3 (calc).

▶ DO4620000

(1R,2S)-form

(-)-erythro-form. *N-Methylephedrine*. *Methylephedrine*, BAN. *Metheph*. *Methy-F*

[552-79-4]

Alkaloid from *Ephedra sinica* and *Ephedra vulgaris* (Ephedraceae). Analeptic. Needles (MeOH). Mp 87-88°. [α]_D -29.2 (MeOH).

Hydrochloride:

Tablets (EtOH). Sol. H₂O, EtOH; spar. sol. Me₂CO. Mp 188-189°. [α]_D -29.8 (H₂O).

Methiodide:

Tablets (EtOH). Mp 212-213°. [α]_D -21.8.

Picrate:

Cryst. (EtOH). Mp 144°.

(1S,2S)-form

(+)-threo-form. *Methyl-ψ-ephedrine*. *Methylisoephedrine*. *Methylpseudoephedrine*

[51018-28-1]

Alkaloid from Ma Huang. Cryst. (petrol). Sol. most org. solvs.; spar. sol. H₂O. Mp 30°. [α]_D +48.1.

Methiodide:

Cryst. (EtOH). Mp 216-217°.

Picrate:

Cryst. (EtOH). Mp 152-153°.

(1RS,2SR)-form

(±)-erythro-form

Cryst. (petrol or MeOH). Mp 63.5-64.5°.

Hydrochloride: [18760-80-0]

Cryst. (Me₂CO). Mp 207-208°.

▶ LD₅₀ (mus, orl) 758 mg/kg. DO5100000

[1201-56-5, 14222-20-9, 21067-55-0, 39263-91-7, 62560-55-8, 42151-56-4]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 318D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 579C: 580C (*nmr*)

Smith, S. *et al.*, *J.C.S.*, 1927, 2056

Nagai, W.N. *et al.*, *Annalen*, 1929, **470**, 157 (*synth*)

Leithe, W. *et al.*, *Ber.*, 1932, **65**, 660

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*rev, occur*)

Shono, T. *et al.*, *Tet. Lett.*, 1983, **24**, 4577 (*synth*)

Blagg, J. *et al.*, *Chem. Comm.*, 1985, 653 (*synth*)

Miyano, S. *et al.*, *J.O.C.*, 1985, **50**, 4350 (*resoln*)

Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1249


Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **5**, 3496-3501 (*use*)

Kim, H.K. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1382-1385 (*pmr, anal*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MJU750; MNN350

3-(3-Dimethylamino-propyl)-1(3H)-isobenzofuranone

Piervardine. 3-(3-Dimethylaminopropyl)phthalide
 [24282-25-5]



C₁₃H₁₇NO₂ 219.283

(S)-form

Alkaloid from *Dendrobium pierardii* (Orchidaceae). Liq.; cryst. (as hydro-

chloride). Mp 185-186° (hydrochloride). [α]_D²³ -69 (c, 1.2 in CHCl₃).

Elander, M. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 2177; 1971, **25**, 721 (*isol, spectra, struct, synth*)

N-[4-(Dimethylarsinoyl)butanoyl]aminoethylsulfonic acid

N-[4-(Dimethylarsinoyl)butanoyl]taurine

[142732-42-1]

Me₂As(O)CH₂CH₂CH₂CONHCH₂CH₂-SO₃H


C₈H₁₈AsNO₅S 315.221

Constit. of the kidney of *Tridacna maxima*. Needles. Mp 215-216°.

Francesconi, K.A. *et al.*, *J.C.S. Perkin I*, 1992, 1349 (*isol, synth*)

6,6-Dimethyl-2-azaspiro[4.4]-non-1-ene

Polyzonimine



C₁₀H₁₇N 151.251

(S)-form [55811-47-7]

Occurs in defence secretion of the millipede *Polyzonium rosalbum*. Oil. Bp₁₀ 81°. [α]_D²⁰ +3.26 (CHCl₃).

Smolanoff, J. *et al.*, *Science (Washington, D.C.)*, 1975, **188**, 734 (*isol, struct, pmr, synth*)


Sugahara, T. *et al.*, *Chem. Comm.*, 1984, 214 (*synth*)

Takagi, Y. *et al.*, *J. Braz. Chem. Soc.*, 2000, **11**, 578-583 (*synth, abs config*)

Mori, K. *et al.*, *Tet. Lett.*, 2000, **41**, 6623-6625 (*synth*)

4,4-Dimethyl-15-azasterol

4,4-Dimethyl-15-aza-D-homoergosta-8,14,24(28)-trien-3-ol, 9CI. Geotrichum Alkaloid A 25822A. A 25822A. *Antibiotic A 25822A*
 [50686-99-2]



(3β,5α)-form

C₃₀H₄₉NO 439.723

Steroidal alkaloid from cultures of *Geotrichum flavo-brunneum*. Also from *Calcarisporium thermophilum*. Antifungal agent. Sol. MeOH, CHCl₃; fairly sol. C₆H₆, CCl₄; poorly sol. H₂O, hexane. Mp 147° (133-136°). [α]_D²⁵ -72 (c, 1.15 in MeOH). λ _{max} 240 (ε 12300); 270 (sh) (EtOH). λ _{max} 277 (13400) (EtOH/HCl).


Chamberlin, J.W. *et al.*, *J. Antibiot.*, 1974, **27**, 992; 1975, **28**, 102 (*isol, pmr, uv, ir, struct*)

Dolle, R.E. *et al.*, *Chem. Comm.*, 1988, 133 (*synth*)

Chrisp, P. *et al.*, *Z. Naturforsch., C*, 1990, **45**, 179 (*isol*)

1,1'-Dimethyl-3,3'-bipyridinium(2+), 9CI D-743

N,N'-Dimethyl-3,3'-bipyridyl(2+). *Metacquat* [41491-84-3]



$C_{12}H_{14}N_2^{2+}$ 186.256

Quaternary alkaloid from an arrow poison of South-East Asia. Exhibits curare-like activity.

Diiodide: [63095-07-8]

$C_{12}H_{14}I_2N_2$ 440.065

Yellow cryst. Mp 300° (278-279° dec.).

Methosulfate (1:2): [118535-24-3]

$C_{14}H_{20}N_2O_8S_2$ 408.453

Cryst. (EtOH). Mp 186-190°.


Lukes, R. *et al.*, *Chem. Listy*, 1958, **52**, 759-762 (*synth*)

Singh, S.K. *et al.*, *Z. Naturforsch., B*, 1988, **43**, 778-784 (*synth*)

Solling, T.I. *et al.*, *Acta Chem. Scand.*, 1998, **52**, 372-373 (*isol, synth, uv, pmr, cmr*)

2,7-Dimethyl-5,10-[2]butenophenazine-11,14-dione, 9CI D-744

[945539-89-9]




$C_{18}H_{14}N_2O_2$ 290.321

Prod. by a marine-derived *Bacillus* sp. Cytotoxic. Yellow needles (MeOH). Mp 186.7-187.4°. $[\alpha]_D^{25}$ -19.8 (c, 0.05 in $CHCl_3$). λ_{max} 240 (log ϵ 3.72); 281 (log ϵ 3.46); 289 (log ϵ 3.41); 348 (log ϵ 3.39); 364 (log ϵ 3.48); 384 (log ϵ 3.31) (MeOH).

Li, D. *et al.*, *Arch. Pharmacol. Res.*, 2007, **30**, 552-555 (*isol, pmr, cmr*)

2,9-Dimethyl-2,9-diazatricyclo[10.2.2.2^{5,8}]octadeca-5,7,12,14,15,17-hexaene-3,10-diol, 9CI D-745

N,α',N',α'-Cyclodi(4-methylaminobenzeneethanol) [124190-18-7]




$C_{18}H_{22}N_2O_2$ 298.384

Alkaloid from peelings of *Citrus unshiu* (satsuma mandarin) (Rutaceae).

Matsubara, Y. *et al.*, *CA*, 1990, **112**, 73757; 113, 17474 (*isol, pmr, cmr, ms, struct*)

4,6-Dimethyl-2,4-dodecadienoic acid D-746



$C_{14}H_{24}O_2$ 224.342

(2E,4E,6R)-form [152984-48-0]

Oil. $[\alpha]_D$ -63.9 (c, 2.1 in CH_2Cl_2).

Amide: 4,6-Dimethyl-2,4-dodecadienamide. *Gymnamide*

$C_{14}H_{25}NO$ 223.358

Prod. by *Gymnascella dankaliensis* *isol.* from a *Halichondria* sp. Oil. $[\alpha]_D^{26}$ -47.6 (c, 0.76 in $CHCl_3$). λ_{max} 228 (sh) (log ϵ 3.71); 261 (log ϵ 4.24) (EtOH).

(2E,4E,6S)-form [138875-82-8]

Prod. by a *Phomopsis* sp. Oil. $[\alpha]_D$ +63.3 (c, 1 in CH_2Cl_2). λ_{max} 265 (no solvent reported).

11-Hydroxy: 11-Hydroxy-4,6-dimethyl-2,4-dodecadienoic acid [253351-46-1]

$C_{14}H_{24}O_3$ 240.342

Prod. by a *Phomopsis* sp. Stereochem. not fully determined. λ_{max} 266 (no solvent reported).

Wipf, P. *et al.*, *J.O.C.*, 1993, **58**, 7195-7203 (*synth, pmr, cmr*)

Wrigley, S.K. *et al.*, *J. Antibiot.*, 1999, **52**, 862-872 (*isol, pmr, cmr*)

Gurjar, M.K. *et al.*, *Heterocycles*, 2000, **53**, 143-149 (*synth*)

Amagata, T. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1384-1388 (*Gymnamide*)

Dimethylformamide, 9CI D-747

Formyldimethylamine. DMF

[68-12-2]

$HCONMe_2$

C_3H_7NO 73.094

Constit. of *Astragalus cibarius*. Manuf. by carbonylation of dimethylamine or by reaction between methyl formate and dimethylamine. Widely used aprotic solv. and reagent in org. synth. and chemical analysis. Liq. Misc. H_2O , EtOH, Et_2O , C_6H_6 , $CHCl_3$. $d_4^{22.4}$ 0.95. Mp -61°. Bp 153° Bp₃₉ 76°. $n_D^{22.4}$ 1.4294. Dielectric constant ϵ 36.71. Viscosity 0.802 cP.


► Flammable, fl. p. 55°, autoignition temp. 440°. Possible human carcinogen. May increase skin absorption of other carcinogens. Skin and eye irritant. Symptoms of occup. exposure include anorexia, abdominal pain, nausea, headache, dizziness. Hepatotoxic. LD₅₀ (rat, orl) 2800 mg/kg. Exp. reprod. and teratogenic effects. Reacts violently with many materials. OES: long-term 10 ppm; short-term 20 ppm (Sk). LQ2100000

[4472-41-7]

Stermitz, F.R. *et al.*, *Phytochemistry*, 1972, **11**, 1117 (*occur*)

1,3-Dimethylguanine D-748

1,2,3,7-Tetrahydro-2-imino-1,3-dimethyl-6H-purin-6-one [224801-13-2]



$C_7H_9N_5O$ 179.181

Isol. from the ascidian *Botrylloides leachi*. Solid. λ_{max} 201 (log ϵ 3.7); 263 (log ϵ 3.3) (MeOH).

7H-form

7-N-Me: 1,2,3,7-Tetrahydro-2-imino-1,3,7-trimethyl-6H-purin-6-one. **1,3,7-Trimethylguanine**

[110025-83-7]

$C_8H_{11}N_5O$ 193.208

Isol. from *Eudistoma maculosum* and *Latrunclia brevis*. Pale-yellow needles (MeOH/ Et_2O). Mp 192-194°. λ_{max} 213 (ϵ 7900); 288 (ϵ 1600) (pH 13) (Derep). λ_{max} 264 (ϵ 2000) (pH 7) (Derep).


Perry, N.B. *et al.*, *J. Nat. Prod.*, 1987, **50**, 307-308 (*1,3,7-Trimethylguanine*)

Lindsay, B.S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 638-639 (*isol, uv, pmr, cmr, ms*)

Berry, Y. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 479-483 (*1,3,7-Trimethylguanine*)

2,4(5)-Dimethylimidazole D-749

[930-62-1]



$C_5H_8N_2$ 96.132

Probable artifact from the flowers of *Sophora flavescens*. Constit. of tobacco smoke. Cryst. Mp 92° (*in vacuo*). Bp_{0.02} 118-120°. pK_a 8.36 (25°).

Hydrochloride: [70807-88-4]

Hygroscopic needles (EtOH/ Et_2O). Mp 205°.

3H-form

3-Me: 1,2,5-Trimethyl-1H-imidazole

[1739-81-7]

$C_6H_{10}N_2$ 110.158

Yellow liq. Bp 220-240°.

3-Me; picrate: Mp 208-209°.

[126311-05-5, 1689-36-7]

Grindley, R. *et al.*, *J.C.S.*, 1927, 3134 (*3-Me*)

Weidenhagen, R. *et al.*, *Ber.*, 1935, **68**, 1960 (*synth*)

Scheinbaum, M.L. *et al.*, *Tet. Lett.*, 1971, 2205 (*synth*)

Casey, M. *et al.*, *Chem. Comm.*, 1982, 714 (*synth*)

Murakoshi, I. *et al.*, *Phytochemistry*, 1982, **21**, 2379 (*isol*)

Watanabe, T. *et al.*, *J. Het. Chem.*, 1983, **20**, 1277 (*synth*)
 Garcia, M.L.S. *et al.*, *J.C.S. Perkin 2*, 1983, 1391 (*nqr*)
 Lipshutz, B.H. *et al.*, *J.O.C.*, 1983, **48**, 3745 (*synth*)

4,5-Dimethylimidazole D-750 [2302-39-8]

C₅H₈N₂ 96.132
 Plates. Mp 120°. pK_a 8.28 (25°, 0.1 M KCl). Several earlier syntheses have given the formate salt having the same Mp. (see D'Sa *et al.*)

Hydrochloride: [53316-51-1]
 Cryst. Mp 305° dec.

N-Benzoyl:
 C₁₂H₁₂N₂O 200.24
 Plates (petrol). Mp 74-75°.

N-Me: 1,4,5-Trimethyl-1H-imidazole
 [20185-22-2]
 C₆H₁₀N₂ 110.158
 Mp 46°. Bp₁₃ 118°.

N-Me, hydrochloride:
 Cryst. + 1H₂O. Mp 80° (hydrate) Mp 199° (anhyd.).

N-Me, methiodide: Mp 158°.

N-Me, N'-oxide: [29982-09-0]
 C₆H₁₀N₂O 126.158
 Cryst. (CH₂Cl₂/Et₂O). Mp 164-166°.

1,3-Dibenzyl: 1,3-Dibenzyl-4,5-dimethylimidazolium (1+). **Lepidiline A**
 [596093-98-0]
 C₁₉H₂₁N[⊕] 277.388

Alkaloid from the roots of *Lepidium meyenii* (maca). Needles (Me₂CO) (as chloride). Mp 231-233° (chloride). λ_{max} 200 (log ε 3.33); 258 (log ε 2.29); 278 (log ε 2.12) (MeOH (chloride)).

[86027-00-1]

Fargher, R.G. *et al.*, *J.C.S.*, 1919, **115**, 217-260 (*synth*)

Weidenhagen, R. *et al.*, *Ber.*, 1937, **70**, 570-583 (*synth*)

Beak, P. *et al.*, *Tetrahedron*, 1969, **25**, 3287-3295 (*I-Me, synth, ir, uv, pmr*)

Wegner, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1974, **307**, 972-975 (*synth*)


Sattler, H.J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1975, **308**, 795-798 (*I-Me, cmr*)

D'Sa, A. *et al.*, *J. Het. Chem.*, 1991, **28**, 1819-1820 (*synth, pmr, cmr*)

Mloston, G. *et al.*, *Helv. Chim. Acta*, 1998, **81**, 1585-1595 (*I-Me 3-oxide, synth, ir, pmr, cmr*)

Cui, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1101-1103 (*Lepidiline A, isol, pmr, cmr, cryst struct*)

2,4-Dimethyl-1H-indole, 9CI D-751 [10299-61-3]



C₁₀H₁₁N 145.204
 Alkaloid from fruit bodies of the unpalatable mushrooms *Tricholoma sciodes* and *Tricholoma virgatum*. Liq. Bp_{0.5} 94-

96°. λ_{max} 223 (ε 19200); 268 (ε 4400); 271 (ε 4400); 273 (ε 4300); 278 (ε 2900) (EtOH).

Picrate:

Cryst. Mp 164.5°.

Marion, L. *et al.*, *Can. J. Res., Sect. B*, 1947, **25**, 1 (*synth*)

Gassman, P.G. *et al.*, *J.A.C.S.*, 1974, **96**, 5495 (*synth*)


McDonald, B.G. *et al.*, *J.C.S. Perkin 1*, 1975, 1446 (*synth*)

Bard, R.R. *et al.*, *J.O.C.*, 1980, **45**, 1546 (*synth*)

Garlaschelli, L. *et al.*, *Tetrahedron*, 1994, **50**, 3571-3574 (*isol, uv, ir, pmr, cmr, ms*)

Söderberg, B.C. *et al.*, *J.O.C.*, 1999, **64**, 9731-9734 (*synth*)


3,4-Dimethyl-7-(methylamino)-5,8-isoquinolinedione D-752 3,4-Dimethyl-7-(methylamino)-5,8-isoquinolinequinone



C₁₂H₁₂N₂O₂ 216.239
 Prod. by the marine-derived *Streptomyces* sp. B1848. Red solid. λ_{max} 232; 273; 339; 447 (MeOH).

Shaaban, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (*isol, pmr, cmr, ms*)


3,6-Dimethyl-1-(3-methyl-2-butenyl)-9H-carbazole D-753 3,6-Dimethyl-1-prenyl-9H-carbazole



C₁₉H₂₁N 263.382
 Alkaloid from the root bark of *Murraya koenigii*. Cryst. (Me₂CO). Mp 168°. λ_{max} 223 (log ε 4.56); 235 (log ε 4.55); 242 (log ε 4.55); 253 (log ε 4.55); 275 (log ε 4.55); 327 (log ε 4.53); 341 (log ε 4.55); 357 (log ε 4.54); 381 (log ε 3.5) (EtOH).

Begum, N.A. *et al.*, *J. Indian Chem. Soc.*, 2005, **82**, 165-171 (*isol, uv, pmr, cmr*)

2,5-Dimethyl-3-(3-methyl-2-butenyl)pyrazine, 9CI D-754 3-Isopentyl-2,5-dimethylpyrazine. 2,5-Dimethyl-3-prenylpyrazine



C₁₁H₁₆N₂ 176.261
 Isol. from the ant *Iridomyrmex humilis*. Prod. by a marine bacterium.

2',3'-Dihydro: 2,5-Dimethyl-3-(3-methylbutyl)pyrazine, 9CI
 [18433-98-2]
 C₁₁H₁₈N₂ 178.277
 Isol. from the ants *Anochetus kempfi*, *Anochetus mayri*, *Mesoponera castanea* and *Mesoponera castaneicolor*. Prod. by various marine bacteria.

Cairll, G.W.K. *et al.*, *Aust. J. Chem.*, 1974, **27**, 879


Fales, H.M. *et al.*, *Tetrahedron*, 1988, **44**, 5045 (*deriv*)

Jones, T.H. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1343-1345 (*isol, deriv*)

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*marine isol*)

Dickschat, J.S. *et al.*, *J. Chem. Ecol.*, 2005, **31**, 925-947 (*marine isol*)


2,5-Dimethyl-3-(2-methylbutyl)pyrazine D-755 [72668-36-1]



C₁₁H₁₈N₂ 178.277
 Prod. by the marine bacterium *Sulfobacter pontiacus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol, synth, pmr, cmr, ms*)


α,α-Dimethyl-5-(3-methyl-1-oxo-2-butenyl)-1H-indole-3-propanal, 9CI D-756 3-(2-Formyl-2-methylpropyl)-5-(3-methyl-1-oxo-2-butenyl)-1H-indole [170128-45-7]



C₁₈H₂₁NO₂ 283.369
 Alkaloid from stem bark of *Isolona maitlandii*. Oil. Poss. artifact. λ_{max} 215 (log ε 4.33); 267 (log ε 4.54); 310 (log ε 4.17) (MeOH).

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **40**, 967-973 (*isol, uv, ir, pmr, cmr, ms, struct*)

2,5-Dimethyl-3-(2-methylpropyl)pyrazine D-757 3-Isobutyl-2,5-dimethylpyrazine [32736-94-0]




C₁₀H₁₆N₂ 164.25

Prod. by various marine bacteria.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*, *synth*, *pmr*, *cmr*, *ms*)

2,5-Dimethyl-3-(methylthio)pyrazine

[59021-08-8]




C₇H₁₀N₂S 154.235

Prod. by the marine bacterium *Sulfitorbacter pontiacus*. Pale yellow liq.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*, *synth*, *pmr*, *cmr*, *ms*)

3-(3,7-Dimethyl-2,6-octadienyl)-1H-indole



C₁₈H₂₃N 253.386

(*E*)-form

3-Geranylindole

[102720-32-1]


Alkaloid from *Monodora angolensis*.

Mirand, C. *et al.*, *Tet. Lett.*, 1985, **26**, 3985-3988 (*synth*)

Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (*isol*)

2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-4(1H)-quinolinone

2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-4-quinolinol. 2-(3,7-Dimethyl-2,6-octadienyl)-4-hydroxy-3-methylquinoline



C₂₀H₂₅NO 295.424

(*E*)-form

Antibiotic CJ 13136. *CJ 13136*

[189372-40-5]

Prod. by *Pseudonocardia* sp. CL38489.

Active against *Helicobacter pylori*. Glass. Sol. MeOH, EtOAc, Me₂CO; poorly sol. H₂O. λ_{max} 213 (ε 25400); 242 (ε 26300); 322 (ε 9200); 335 (ε 9400) (MeOH).

N-Me: 2-(3,7-Dimethyl-2,6-octadienyl)-1,3-dimethyl-4(1H)-quinolinone. *Antibiotic CJ 13217*. *CJ 13217*

[189372-43-8]

C₂₁H₂₇NO 309.45

Prod. by *Pseudonocardia* sp. CL38489. Glass. Sol. MeOH, Me₂CO, EtOAc;

poorly sol. H₂O. λ_{max} 214 (ε 16000); 247 (ε 15700); 331 (ε 6400); 344 (ε 7000) (MeOH).

N-(Methylthiomethyl): 2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-1-(methylthiomethyl)-4(1H)-quinolinone. *Antibiotic CJ 13536*. *CJ 13536*

[189372-45-0]

C₂₂H₂₉NOS 355.543

Prod. by *Pseudonocardia* sp. CL38489.

Glass. Sol. MeOH, EtOAc, Me₂CO; poorly sol. H₂O. λ_{max} 212 (ε 38100); 247 (ε 42800); 311 (ε 20600); 344 (ε 23900) (MeOH).

6'ξ,7'ξ-Epoxyde. N-Me: *Antibiotic CJ 13564*. *CJ 13564*

[189372-48-3]

C₂₁H₂₇NO₂ 325.45

Prod. by *Pseudonocardia* sp. CL38489.

Glass. Sol. MeOH, Me₂CO, EtOAc; poorly sol. H₂O. [α]_D²⁵ -17.5 (c, 0.04 in MeOH). λ_{max} 214 (ε 38200); 246 (ε 39700); 332 (ε 15700); 344 (ε 15800) (MeOH).


Dekker, K.A. *et al.*, *J. Antibiot.*, 1998, **51**, 145-152 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

4-(3,7-Dimethyl-2,6-octadienyl)-1H-pyrrole-2-carboxylic acid, 9CI

4-Geranyl-1H-pyrrole-2-carboxylic acid.

Pyrolostatin. EC 40. *Antibiotic EC 40*

[144314-68-1]



C₁₅H₂₁NO₂ 247.336

Prod. by *Streptomyces chrestomyceticus*. Lipid peroxidation inhibitor. Pale yellow powder. Mp 121-124° dec. λ_{max} 237 (ε 8500); 271 (ε 16100) (prob. MeOH, not reported) (Derep).


Japan. Pat., 1992, 92 95 069; *CA*, **117**, 210728d (EC 40)

Kato, S. *et al.*, *J. Antibiot.*, 1993, **46**, 892 (*isol*, *pmr*, *cmr*, *struct*, *props*)

Fumoto, Y. *et al.*, *J.O.C.*, 1999, **64**, 6518-6521 (*synth*, *pmr*, *cmr*)

2-(3,7-Dimethyl-2,6-octadienyl)-4(1H)-quinolinone

2-(3,7-Dimethyl-2,6-octadienyl)-4-hydroxyquinoline



C₁₉H₂₃NO 281.397

(*E*)-form

Antibiotic CJ 13565. *CJ 13565*

[189372-51-8]

Prod. by *Pseudonocardia* sp. CL38489.

Active against *Helicobacter pylori*. Glass. Sol. MeOH, Me₂CO, EtOAc; poorly sol. H₂O. λ_{max} 213 (ε 37100); 238 (ε 34700); 316 (ε 13100); 328 (ε 12700) (MeOH). λ_{max} 213 (ε 37100); 238 (ε 34700); 316 (ε 13100); 328 (ε 12700) (MeOH) (Berdy).

N-Me: 2-(3,7-Dimethyl-2,6-octadienyl)-1-methyl-4(1H)-quinolinone. *Antibiotic CJ 13566*. *CJ 13566*

[189372-53-0]

C₂₀H₂₅NO 295.424

Prod. by *Pseudonocardia* sp. CL38489.

Glass. Sol. MeOH, Me₂CO, EtOAc; poorly sol. H₂O. λ_{max} 214 (ε 33400); 242 (ε 31500); 323 (ε 14200); 335 (ε 14700) (MeOH). λ_{max} 214 (ε 33400); 242 (ε 31500); 325 (ε 14200); 335 (ε 14700) (MeOH) (Berdy).

1'ξ-Hydroxy, N-Me: 2-(1-Hydroxy-3,7-dimethyl-2,6-octadienyl)-1-methyl-4(1H)-quinolinone. *Antibiotic CJ 13567*. *CJ 13567*

[189372-55-2]

C₂₀H₂₅NO₂ 311.423

Prod. by *Pseudonocardia* sp. CL38489.

Glass. Sol. MeOH, Me₂CO, EtOAc; poorly sol. H₂O. [α]_D²⁵ +66.3 (c, 0.57 in MeOH). λ_{max} 214 (ε 38000); 243 (ε 37000); 324 (ε 16400); 337 (ε 17500) (MeOH). λ_{max} 214 (ε 38000); 243 (ε 37000); 324 (ε 16400) (MeOH) (Berdy).

Δ'ξ-Isomer, 3'ξ-hydroxy, N-Me: 2-(3-Hydroxy-3,7-dimethyl-1,6-octadienyl)-1-methyl-4(1H)-quinolinone. *Antibiotic CJ 13568*. *CJ 13568*

[189372-57-4]


C₂₀H₂₅NO₂ 311.423

Prod. by *Pseudonocardia* sp. CL38489.

Glass. Sol. MeOH, EtOAc, Me₂CO; poorly sol. H₂O. [α]_D²⁵ -51.4 (c, 0.03 in MeOH). λ_{max} 219 (ε 23500); 252 (ε 38600); 330 (sh); 339 (ε 15200) (MeOH). λ_{max} 219 (ε 23500); 252 (ε 38600); 339 (ε 15200) (MeOH) (Berdy).

Dekker, K.A. *et al.*, *J. Antibiot.*, 1998, **51**, 145-152 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

4,8-Dimethyl-15-oxo-16-aza-18-norandrost-13-ene-4-carboxylic acid



C₂₀H₂₉NO₃ 331.454

(4β,5α)-form

N-(2-Phenylethyl): *Haumanamide* [141266-06-0]

C₂₈H₃₇NO₃ 435.605

Alkaloid from a *Spongia* sp. Cytotoxic agent. Powder. [α]_D -163.2 (c, 0.13 in CHCl₃). λ_{max} 250 (ε 340) (EtOH) (Derep).

N-(2-Imidazol-5-ylethyl): **Spongolactam A**

C₂₅H₃₅N₃O₃ 425.57

Alkaloid from a *Spongia* sp. Powder. [α]_D²⁵ -37 (c, 0.04 in MeOH). λ_{max} 208 (ε 16000); 238 (sh) (ε 3700) (MeOH).

N-Carboxymethyl: **Spongolactam C**

C₂₂H₃₁NO₅ 389.491

Alkaloid from a *Spongia* sp. Powder. [α]_D²⁷ -30 (c, 0.02 in MeOH). λ_{max} 215 (ε 13000); 243 (sh) (ε 3300) (MeOH).

Pham, A.T. et al., *Tet. Lett.*, 1992, **33**, 1147-1148 (*Haumanamide*)

Mori, D. et al., *J.O.C.*, 2007, **72**, 7190-7198 (*Spongolactams A, C*)

4,8-Dimethyl-17-oxo-16-aza-18-norandrost-13-ene-4-carboxylic acid D-764

C₂₀H₂₉NO₃ 331.454

(4β,5α)-form

N-(2-Imidazol-5-ylethyl): **Spongolactam B**

[952514-16-8]


C₂₅H₃₅N₃O₃ 425.57

Isol. from a *Spongia* sp. Powder. [α]_D²⁵ -12 (c, 0.03 in MeOH). λ_{max} 208 (ε 13000); 238 (sh) (ε 3400) (MeOH).

Mori, D. et al., *J.O.C.*, 2007, **72**, 7190-7198 (*isol, pmr, cmr*)

4-[2-(3,5-Dimethyl-2-oxo-cyclohexyl)-2-hydroxyethyl]-2,6-piperidinedione, 9CI D-765

[17280-60-3]



(1R,3R,5S,αR)-form

C₁₅H₂₃NO₄ 281.351

Glutarimide antibiotic.

(1R,3R,5S,αR)-form

Isocycloheximide

[6746-42-5]

Isol. from *Streptomyces griseus*. Active against *Saccharomyces pastorianus*. Possesses 30% of Cycloheximide activity. Sol. H₂O, CHCl₃, MeOH; poorly sol. hexane. Mp 101-102°. [α]_D²³ +36 (c, 10 in MeOH). λ_{max} 287 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 500 mg/kg. MA4375100

(1R,3R,5S,αS)-form

α-Epiisocycloheximide. Neoisocycloheximide

[94799-82-3]

Isol. from Nongkang 101. Antifungal agent. Cryst. (EtOAc/Et₂O). Mp 130-131°.

(1R,3S,5S,αR)-form

Naramycin B

[642-81-9]

From *Streptomyces naraensis* and *Streptomyces griseus*. Active against fungi and yeasts. Possesses herbicidal props. Plates. Sol. MeOH, H₂O, CHCl₃; fairly sol. EtOAc; poorly sol. hexane. Mp 113°.

[α]_D²² +55.8 (c, 5 in MeOH). λ_{max} 292 (ε 31) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 70 mg/kg. QN4555000

(1S,3S,5S,αR)-form

Cycloheximide, USAN. Cicloheximide,

INN. Actidione. Acti-Aid. Naramycin.

Naramycin A. NSC 185. U 4527

[66-81-9]

From *Streptomyces griseus*. Antifungal agent. Active against leaf blight and rust diseases. Now superseded. Rodent repellent. Cryst. (H₂O). Mp 119.5-121°. [α]_D²⁹ -33.8 (c, 1 in CHCl₃). pK_a 11.2. Log P -0.49 (calc). Unstable in alkali.

► Skin irritant. LD₅₀ (mus, ipr) 100 mg/kg; LD₅₀ (rat, orl) 2 mg/kg. Exp. reprod. and teratogenic effects. MA4375000

Ac: Mp 150.2°. [α]_D²⁵ +24.6 (c, 3.7 in MeOH).

4-Nitrobenzoyl: Mp 215-220° dec.

α-Ketone: Dehydroactidione. Dehydrocycloheximide

[363-27-9]

C₁₅H₂₁NO₄ 279.335

From *Streptomyces noursei* and other *Streptomyces* spp. Cryst. (EtOH). Mp 178°. [α]_D²⁰ -29.8 (c, 2 in dioxan).

1,6-Didehydro: Inactone

C₁₅H₂₁NO₄ 279.335

From *Streptomyces griseus*. Needles. Sol. H₂O, MeOH, CHCl₃; fairly sol. C₆H₆. Mp 116°. [α]_D -55 (H₂O). Only 1.2% as biol. active as Cycloheximide. λ_{max} 331 (ε 70) (EtOH) (Berdy).

3R-Hydroxy: Streptovitacin C₂

[103957-20-6, 103957-21-7]

C₁₅H₂₃NO₅ 297.35

Glutarimide-type antibiotic. Prod. by *Streptomyces griseus*. Shows antitumour activity. Cryst. Sol. MeOH, EtOAc, H₂O; fairly sol. CHCl₃, Et₂O; poorly sol. hexane. Mp 91-96°. λ_{max} 236; 263 (H₂O) (Berdy).

4ξ-Hydroxy: Streptovitacin B

[634-50-4]

[11040-33-8]

C₁₅H₂₃NO₅ 297.35

Glutarimide-type antibiotic. Produced by *Streptomyces griseus*. Shows antitumour activity. Cryst. Sol. H₂O, EtOAc, MeOH; fairly sol. CHCl₃, Et₂O; poorly sol. hexane. Mp 124-128°. λ_{max} 232; 264 (H₂O) (Berdy).

4ξ-Hydroxy, di-Ac: Mp 155-158°.

5R-Hydroxy: Streptovitacin A. 5-Hydroxy-

xy-cycloheximide. Resactin. NSC 39147. U 9361. Antibiotic U 9361

[523-86-4]

C₁₅H₂₃NO₅ 297.35

Glutarimide-type antibiotic. Produced by *Streptomyces griseus*. Egg-laying inhibitor of red spider mites in fruit tree pest control. Also used in aphid control. Shows antitumour activity. Cryst. Sol. H₂O, EtOAc, MeOH; fairly sol. CHCl₃, Et₂O; poorly sol. hexane. Mp 156-159°. λ_{max} 232; 244; 262 (H₂O) (Berdy).

► MA4900000

5R-Acetoxy: Streptovitacin E73. SF

2441. Antibiotic SF 2441. 5-Acetoxy-cycloheximide. 4-Acetoxy-cycloheximide

[2885-39-4]

C₁₇H₂₅NO₆ 339.388

Isol. from *Streptomyces albulus*. Cryst. Mp 140° Mp 165-168°. [α]_D²⁵ -8.8 (c, 1 in MeOH).

► MA5075000

5R-Acetoxy, α-Ac:

Needles (CH₂Cl₂/Et₂O). Mp 177-178°.

(1R,3S,5S,αS)-form

Mp 98.5-99.5°. [α]_D²⁰ +74.7 (c, 0.55 in MeOH).

(1S,3S,5S,αS)-form

Mp 106-107°. [α]_D²⁰ -23.5 (c, 2.5 in MeOH).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 798C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1305C (*nmr*)

Whiffen, A.J. et al., *J.A.C.S.*, 1947, **69**, 474 (*isol*)

Kornfeld, E.C. et al., *J.A.C.S.*, 1949, **71**, 150 (*struct*)

Raul, R. et al., *Bull. Soc. Chim. Fr.*, 1955, 1316 (*isol*)

Eble, T.E. et al., *Antibiot. Annu.*, 1958, 535 (*Streptovitacins, purifn. props*)

Herr, R.R. et al., *J.A.C.S.*, 1959, **81**, 2595 (*Streptovitacins, struct*)

Rao, K.V. et al., *J.A.C.S.*, 1960, **82**, 1127; 1129 (*Streptovitacin E73*)

Suzuki, M. et al., *Chem. Pharm. Bull.*, 1963, **11**, 582; 589 (*synth*)

Starkovsky, N.A. et al., *Tet. Lett.*, 1964, 919 (*abs config*)

Rouzaud, J. et al., *Bull. Soc. Chim. Fr.*, 1965, 2030 (*Streptovitacins, pmr, stereochem*)

Johnson, F. et al., *J.A.C.S.*, 1966, **88**, 149 (*synth*)

U.S. Pat., 1967, 3 305 554; CA, **66**, 98487z (*Streptovitacins, struct, manuf*)

Johnson, F. et al., *J.O.C.*, 1968, **33**, 904 (*Streptovitacin A, struct*)

Sayers, J. et al., *J.A.C.S.*, 1977, **99**, 3848 (*cryst struct*)

Jeffs, P.W. et al., *J.A.C.S.*, 1981, **103**, 6185 (*biosynth, cmr*)

Berg, D. et al., *Z. Naturforsch., C*, 1982, **37**, 1100 (*isol, props*)

Kudo, S. et al., *Agric. Biol. Chem.*, 1984, **48**, 2315; 2739; 1985, **49**, 2919 (*synth, stereochem, bibl*)

Jost, J.L. et al., *Drugs Pharm. Sci.*, 1984, **22**, 531 (*rev*)

Ye, D. et al., *CA*, 1985, **102**, 92566 (*isol*)

Dangerous Prop. Ind. Mater. Rep., 1989, **9**, 55 (*tox*)

Perry, N.B. et al., *Magn. Reson. Chem.*, 1989, **27**, 624 (*pmr, conformn*)

Kondo, H. et al., *Agric. Biol. Chem.*, 1990, **54**, 1531 (*Streptovitacins, synth*)

Pesticide Manual, 9th edn., 1991, No. 3600


Martindale. The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1125

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van

Nostrand Reinhold, 1992, CPE750

3,5-Dimethyl-2-pentylpyrazine D-766

[50888-62-5]




C₁₁H₁₈N₂ 178.277

Constit. of the glands of various ants incl. *Brachyponera* sp. and *Streblognathus* sp.

Longhurst, C. *et al.*, *J. Insect Physiol.*, 1978, **24**, 833-837 (*isol*)
 Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1998, **24**, 125-134 (*isol, ms*)

2,5-Dimethyl-3-(2-phenylethenyl)pyrazine, 9CI **D-767**
2,5-Dimethyl-3-styrylpyrazine



C₁₄H₁₄N₂ 210.278

(E)-form [54290-13-0]


Component of defence secretion of ant *Iridomyrmex humilis*. Oil. Bp₃ 117-122°. Converts into (*Z*)-form on exposure to light.

(Z)-form

Component of the defence secretion of *Iridomyrmex humilis*. Oil. Bp₃ 105-118°. Possible artifact.

Cavill, G.W.K. *et al.*, *Aust. J. Chem.*, 1974, **27**, 879-889 (*isol, uv, pmr*)
 Akita, Y. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1447-1458 (*synth, (E)-form, pmr*)

2,6-Dimethyl-4-phenyl-3,5-pyridinedicarboxylic acid **D-768**



C₁₅H₁₃NO₄ 271.272

Di-Et ester:


C₁₉H₂₁NO₄ 327.379

Alkaloid from the rhizomes of *Jatropha elliptica*. Multidrug resistance reversal agent. Powder (hexane). Mp 60-61°.

Marquez, B. *et al.*, *Phytochemistry*, 2005, **66**, 1804-1811 (*di-Et ester, isol, pmr, cmr, ms*)

3,6-Dimethyl-2,5-piperazine-dione, 9CI **D-769**

Lactimide. Alanine cyclic anhydride. Cyclo(alanylalanyl). Alanine anhydride. Dialanine dianhydride. Dialanine anhydride [5625-46-7]



(3*R*,6*R*)-form

C₆H₁₀N₂O₂ 142.157

► TL6380150

(3*R*,6*R*)-form

D-D-form. (-)-cis-form

N,N'-Di-Ph: [174390-96-6]

C₁₈H₁₈N₂O₂ 294.352

Prisms (toluene). Mp 188-189°. [α]_D²⁰ -30.8.

(3*S*,6*S*)-form

L-L-form. (+)-cis-form

[5845-61-4]

Constit. of the roots of *Psammosilene tunicoides*. Cryst. (H₂O). Mp 272° (289°). [α]_D²⁰ +29.1 (EtOH aq.). [α]_D²⁵ +26.4 (c, 1.3 in H₂O).

N-Me: 1,3,6-Trimethyl-2,5-piperazine-dione

[59042-86-3]

C₇H₁₂N₂O₂ 156.184

Mp 116-117°. [α]_D²⁵ +63.7 (c, 3 in CHCl₃).

(3*R,6*R**)-form**

N-Me, N'-E-cinnamoyl: *Nigerazine A. I 639A. Antibiotic I 639A*

[92008-50-9]

C₁₆H₂₂N₂O 258.363

Prod. by *Aspergillus niger*. Inhibits plant root growth. Oil. Sol. MeOH, C₆H₆, Me₂CO, CHCl₃; fairly sol. H₂O. λ_{max} 280 (ε 21900) (MeOH) (Berdy).

(3*R*,6*RS*)-form

(±)-cis-form

[72904-45-1]

Needles or leaflets. Sol. H₂O. Mp 279°.

(3*R*,6*SR*)-form

trans-form

[35590-65-9]

Cryst. or plates (H₂O) (dimorph.). Opt. inactive (*meso*-).

N-(2*E*,4*E*-Hexadienoyl): *Nigragilline*

[24779-38-2]

C₁₃H₂₂N₂O 222.33

Alkaloid from *Aspergillus phoenicis* and other spp. in the *Aspergillus niger* group. Insecticide. Viscous liq. Sol. CHCl₃, Et₂O. Mp 179-179.5° (as picrate). [α]_D²⁴ +107.4 (c, 2.5 in CHCl₃). λ_{max} 262 (ε 26200) (MeOH) (Berdy).

N-Me, N'-E-cinnamoyl: 1,2,5-Trimethyl-4-(1-oxo-3-phenyl-2-propenyl)piperazine. 1-Cinnamoyl-2,4,5-trimethylpiperazine. *Nigerazine B. I 639B. Antibiotic I 693B*

[85982-75-8]

[87980-22-1]

C₁₆H₂₂N₂O 258.363

Prod. by *Aspergillus niger* and *Alternaria brassicae*. Shows plant growth inhibitory props. Oil. [α]_D¹⁹ +103.2 (c, 1.1 in MeOH). Not a *meso*-compd. owing to the differing N groups. Abs. config. not determined. λ_{max} 280 (MeOH).

► TM2483300

[23927-13-1]

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **1**, 1249A (*nmr*)

Kopple, K.D. *et al.*, *J.A.C.S.*, 1956, **78**, 6199 (*synth*)

Zahn, H. *et al.*, *Annalen*, 1957, **605**, 212 (*synth*)

Augustin, M. *et al.*, *Annalen*, 1967, **705**, 185 (*synth*)

Caesar, F. *et al.*, *Pharm. Acta Helv.*, 1969, **44**, 676 (*Nigragilline*)

Sletton, E. *et al.*, *J.A.C.S.*, 1970, **92**, 172 (*synth*)

Tul'chinski, V.M. *et al.*, *Khim. Prir. Soedin.*, 1973, 779; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, 745 (*ir*)

Hagler, A.T. *et al.*, *J.A.C.S.*, 1974, **96**, 5327 (*cryst struct*)

Kralj, B. *et al.*, *Biomed. Mass Spectrom.*, 1975, **2**, 215 (*ms*)

Kawabata, Y. *et al.*, *Makromol. Chem.*, 1975, **176**, 2797 (*synth*)

Schöllkopf, V. *et al.*, *Angew. Chem.*, 1979, **91**, 922 (*synth*)

Kricheldorf, H.R. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 52 (*cmr*)

Iwamoto, T. *et al.*, *Agric. Biol. Chem.*, 1983, **47**, 739; 1985, **49**, 3323 (*Nigerazines*)

Japan. Pat., 1983, 83 146 289; *CA*, **100**, 4788u (*Nigerazines*)

Cheam, T.C. *et al.*, *Spectrochim. Acta A*, 1988, **44**, 185 (*cryst struct, ir*)

Dahiya, J.S. *et al.*, *Phytochemistry*, 1991, **30**, 2825-2828 (*Nigerazine B*)

Bazuik, V.A. *et al.*, *Synthesis*, 1992, 449 (*synth*)

D'Angeli, F. *et al.*, *J.O.C.*, 1996, **61**, 1252


(*N,N-Di-Ph synth, pmr, cryst struct*)

Ding, Z. *et al.*, *Zhongcaoyao*, 2000, **31**, 803-805; *CA*, **134**, 323448 (*isol*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 798 (*Nigragilline*)

2,6-Dimethylpiperidine **D-770**

Lupetidine, 8CI. Nanofin, INN. Nanophyn [504-03-0]



(2*R*,6*R*)-form

C₇H₁₅N 113.202

Log P 1.59 (calc).

► Highly flammable, fl. p. 16°. OK5775000

(2*R*,6*R*)-form [14231-78-8]

Alkaloid from *Nanophyton erinaceum* (Chenopodiaceae). Ganglion blocking agent. Oil. [α]_D -43.02 (neat).

Hydrochloride: [5072-45-7]

Mp 243-244° Mp 280-282°. [α]_D²² +12.4 (c, 0.3 in EtOH).

► OK5785000

N-Me: 1,2,6-Trimethylpiperidine

C₈H₁₇N 127.229

Obt. from *Nanophyton erinaceum* (Chenopodiaceae).

(2*S*,6*S*)-form [40250-84-8]

Oil. [α]_D +12.5 (c, 0.5 in EtOH).

(2*RS*,6*RS*)-form

(±)-trans-form

[10066-29-2]

Oil. Bp 136-138°.

Hydrochloride:

Cryst. (EtOAc/AcOH). Mp 236-238°.

Picrate: Mp 144-146°.

N-Me: Bp 154-155°.

N-Me, picrate: Mp 244-245°.

(2*RS*,6*SR*)-form

cis-form

[766-17-6]

Oil. Bp 127-128°. pK_a 11.07 (25°).

N-Me: Bp₁₂ 50-55°. pK_a 10.49 (25°).

N-Me, picrate: Mp 225°.

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 360C (*ir*)

Aldrich Library of 13C and 1H FT NMR


Spectra, 1992, **1**, 568B (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 451D (*ir*)

- Marcuse, A. *et al.*, *Ber.*, 1899, **32**, 2528 (*synth*)
 Adkins, H. *et al.*, *J.A.C.S.*, 1934, **56**, 2427 (*synth*)
 Kuzovkov, A.D. *et al.*, *Zh. Obshch. Khim.*, 1950, **20**, 1524 (*isol*)
U.S. Pat., 1957, 2 813 100; *CA*, **52**, 6414f (*synth*)
 Vidal-Beretervide, K. *et al.*, *J. Pharmacol. Exp. Ther.*, 1966, **152**, 181 (*pharmacol*)
 Hill, R.K. *et al.*, *J.O.C.*, 1966, **31**, 3451 (*abs config*)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1969, **2**, 156 (*use*)
 Kawazoe, Y. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 429 (*synth, pmr*)
 Lecomte, C. *et al.*, *Cryst. Struct. Commun.*, 1975, **4**, 477 (*cryst struct*)
 House, H.O. *et al.*, *J.O.C.*, 1976, **41**, 863 (*synth, deriv*)
 Duthaler, R.O. *et al.*, *J.A.C.S.*, 1977, **99**, 8406 (*synth*)
 Frazer, R.R. *et al.*, *Tet. Lett.*, 1982, **23**, 4195 (*cmr*)
 Erra-Balsells, R. *et al.*, *Aust. J. Chem.*, 1988, **41**, 103 (*pmr, cmr, ms*)
 Freville, S. *et al.*, *Tetrahedron*, 1997, **53**, 8447 (*synth*)
 Adamo, M.F.A. *et al.*, *Synth. Commun.*, 1999, **29**, 1747-1756 (*2S,6S-form, synth*)

8-(1,1-Dimethyl-2-propenyl)-4-methoxyfuro[2,3-*b*]quinolin-7-ol, 9CI

8-(1,1-Dimethylallyl)confusameline
 [116987-98-5]




$C_{17}H_{17}NO_3$ 283.326
 Alkaloid from the roots of *Haplophyllum glabrinum* (Rutaceae). Needles (Me_2CO). Mp 118-119°.

Rózsza, Zs. *et al.*, *Phytochemistry*, 1988, **27**, 2369 (*isol, uv, pmr, ms, struct*)

2,5-Dimethyl-3-propylpyrazine, 9CI

[18433-97-1]



$C_9H_{14}N_2$ 150.223
 Minor component of secretion of the ant *Iridomyrmex humilis*.

Cavill, G.W.K. *et al.*, *Aust. J. Chem.*, 1974, **27**, 879-889 (*occur*)

3,5-Dimethyl-2-propylpyrazine, 9CI

[32350-16-6]


$C_9H_{14}N_2$ 150.223
 Volatile constit. of roasted coconut, cooked shrimp and roast beef. Present in cocoa. Characterised spectroscopically.

Flament, I. *et al.*, *Helv. Chim. Acta*, 1971, **54**,

1911-1913 (*synth, ms, occur, beef*)
U.S. Pat., 1975, 3 881 025; *CA*, **78**, 109541p (*synth, ms*)
 Jinap, S. *et al.*, *J. Sci. Food Agric.*, 1998, **77**, 441-448 (*occur, cocoa*)

2,5-Dimethylpyrazine, 9CI

FEMA 3272
 [123-32-0]



$C_6H_8N_2$ 108.143
 Component of urinary signalling pheromone in the tree shrew *Tupaia belangeri* and the pine vole *Microtus pinetorum*. Prod. by *Chondromyces crocatus*. Isol. from an arctic marine bacterium. Used as flavour additive and odorant in foods. Liq. or solid with potato chip flavour. Sol. H_2O , EtOH, Et_2O . Mp 15°. Bp 155°. pK_{a1} 1.85; pK_{a2} -4.6 (27°, H_2SO_4 aq.). pK_{a1} 1.99; pK_{a2} -4.42 (25°, H_2O). Steam-volatile. Odour threshold 1800 ppb in H_2O .

► Fl. p. 64° (oc). LD₅₀ (rat, orl) 1020 mg/kg. UQ2800000

Picrate: Mp 157°.

Mono-oxide: [6890-37-5]

$C_6H_8N_2O$ 124.142
 Needles (C_6H_6). Mp 105-108°. Subl.0.001 100.

1,4-Dioxide: [6890-38-6]

$C_6H_8N_2O_2$ 140.141
 Needles ($CHCl_3$). Mp 280° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 840D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 398C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1556C (*ir*)

Newbold, G.T. *et al.*, *J.C.S.*, 1947, 1183-1185 (*1-oxide, 1,4-dioxide*)

Langdon, W.K. *et al.*, *Ind. Eng. Chem. Prod. Res. Dev.*, 1964, **3**, 8 (*synth*)

Wilen, S.H. *et al.*, *Chem. Ind. (London)*, 1969, **8**, 237 (*synth*)

Paudler, W.W. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 513 (*ms*)

Paudler, W.W. *et al.*, *Org. Magn. Reson.*, 1971, **3**, 217 (*nmr*)

Oertel, R.P. *et al.*, *Anal. Chem.*, 1972, **44**, 1589 (*Raman*)

Bus, J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1973, **92**, 123 (*ir*)

Gumbley, S.J. *et al.*, *J. Het. Chem.*, 1985, **22**, 1143 (*props*)

von Stralendorff, F. *et al.*, *J. Chem. Ecol.*, 1987, **13**, 655

Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, DTU600

Boyer, M.L. *et al.*, *J. Chem. Ecol.*, 1989, **15**, 649

Eiermann, U. *et al.*, *Chem. Ber.*, 1990, **123**, 523 (*uv*)


Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 874 (*use, occur*)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, **2**, 318-353 (*marine isol*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DTU600

2,6-Dimethylpyridine, 9CI

D-775
 2,6-Lutidine, 8CI. *FEMA* 3540
 [108-48-5]



C_7H_9N 107.155

Manuf. by reaction of acetone, formaldehyde and ammonia. Constit. of coal tar and shale oil. Present in tea. Flavouring agent. Intermed. for pharmaceuticals. Liq. Sol. EtOH, Et_2O , cold H_2O ; less sol. hot H_2O . d^{20}_4 0.92. Mp -6°. Bp 139-141° (145.6-145.8°). n^{20}_D 1.4977. pK_a 6.6 (25°).

► Flammable, fl. p. 33°. Toxic. OK9700000

Hydrochloride: [15439-85-7]

Cryst. (EtOH). Mp 230-231° dec.

Picrate: [2798-38-1]

Yellow needles (EtOH). Mp 163-164°.

N-Oxide: Ivin

[1073-23-0]

C_7H_9NO 123.154

Liq. Bp₁₈ 115-119°.

N-Oxide; hydrochloride: [6890-58-0]

Mp 219.5°.

[27175-64-0]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 740B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 252C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1520B (*ir*)

Biddiscombe, D.P. *et al.*, *J.C.S.*, 1954, 1957 (*synth, props*)

Červinka, O. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 567 (*synth*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, **1**, 626 (*use*)

Green, J.H.S. *et al.*, *Spectrochim. Acta A*, 1970, **26**, 2139 (*ir, Raman*)

Hejda, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 3035 (*ms*)

Schuster, I.I. *et al.*, *J.O.C.*, 1979, **44**, 2658 (*cmr*)

Bedford, G.R. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 637 (*pmr*)

Yajima, H. *et al.*, *J. Phys. Chem.*, 1986, **90**, 2589 (*uv, fluorescence*)

Org. Synth., *Coll. Vol.*, **6**, 1988, 611


Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 192

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 876-877

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DTU089

2,6-Dimethyl-3,5-pyridine-dicarboxylic acid, 9CI

D-776
 2,6-Dimethylidimicotinic acid, 2,6-Lutidine-3,5-dicarboxylic acid
 [2602-36-0]



$C_9H_9NO_4$ 195.174

Spar. sol. EtOH, Et_2O , H_2O . Mp 316°.

Di-Me ester: [27525-74-2]

$C_{11}H_{13}NO_4$ 223.228

Cryst. (EtOH aq.). Mp 101-102°
(Et₂O/pentane). Bp_{1.3} 131°.

Et ester: [61258-25-1]
C₁₁H₁₃NO₄ 223.228
Needles (H₂O). Mp 131°.

Di-Et ester: [1149-24-2]
C₁₃H₁₇NO₄ 251.282
Alkaloid from the leaves of *Viburnum tinus*. Needles. Mp 73°. Bp 301-302°
Bp₁₆ 179-180°.

Hydrazide: Reagent for pptn. of Progesterone.

Dinitrile: 3,5-Dicyano-2,6-dimethylpyridine
[1656-95-7]
C₉H₇N₃ 157.174
Cryst. (EtOH aq.). Mp 119-120° (115-117°).

Dibenzamide: [55275-98-4]
C₂₁H₁₉N₃O₂ 345.4
Mp 292-294° (278-280°).

Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 673B (nmr)

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 1344E (ir)

Meyer, H. et al., *Monatsh. Chem.*, 1914, 35, 208
Org. Synth., Coll. Vol., 2, 1943, 214-216 (*di-Et ester*)

Braude, E.A. et al., *J.C.S.*, 1960, 3249-3257 (*di-Me ester*)

Lukes, R. et al., *Coll. Czech. Chem. Comm.*, 1961, 26, 1422-1428 (*dinitrile*)

Kurz, J.L. et al., *J.A.C.S.*, 1961, 83, 584-588 (*di-Me ester, synth*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, 1, 626 (*hydrazide, use*)


Patzold, F. et al., *Synth. Commun.*, 1992, 22, 281-288 (*di-Me ester, di-Et ester, dinitrile*)

Eynde, J.J.V. et al., *Tetrahedron*, 1995, 51, 6511-6516 (*di-Et ester, bibl*)

Mohamed, M.A. et al., *Phytochemistry*, 2005, 66, 2780-2786 (*di-Et ester, isol, pmr, cmr*)

1,3-Dimethyl-2,4(1*H*,3*H*)-pyrimidinedione, 9CI

1,3-Dimethyl-2,6-dioxypyrimidine. 1,3-Dimethyluracil
[874-14-6]



C₆H₈N₂O₂ 140.141
Pheromone of the ponerine ant *Megaponera foetens*. Prisms (Et₂O/EtOH). Mp 121-122°.

Yamauchi, K. et al., *J.C.S. Perkin 1*, 1973, 391-392 (*synth, ir, uv*)

De Member, J.R. et al., *J.A.C.S.*, 1975, 97, 6240-6245 (*cmr*)

Harada, S. et al., *Tet. Lett.*, 1976, 2321-2322 (*synth*)

Banerjee, A. et al., *Acta Cryst. B*, 1977, 33, 90-94 (*cryst struct*)

Stolarski, R. et al., *Z. Naturforsch., C*, 1977, 32, 894-900 (*pmr, cmr*)

Iza, N. et al., *J. Mol. Struct.*, 1988, 175, 31-36 (*uv, tautom*)


Janssen, E. et al., *J. Chem. Ecol.*, 1995, 21, 1947-1955 (*isol*)

Rodriguez, J. et al., *Chem. Eur. J.*, 1999, 5, 3549-3561 (*synth, pmr*)

Zajac, M.A. et al., *Synth. Commun.*, 2003, 33, 3291-3297 (*synth*)

2,4-Dimethyl-1*H*-pyrrole-3-carboxylic acid

[17106-13-7]



C₇H₉NO₂ 139.154
Mp 183° dec. p*K*_a 6.13.

Me ester: [52459-90-2]
C₈H₁₁NO₂ 153.18
Cryst. (MeOH aq.). Mp 105-106°.

Et ester: [2199-51-1]
C₉H₁₃NO₂ 167.207
Mp 78-79°. Bp 291° Bp₃₅ 181°.

α-L-Rhamnopyranosyl ester:

C₁₃H₁₉NO₆ 285.296
Prod. by *Streptomyces griseoviridis*.
Mp 90° dec. [α]_D²⁰ -22 (c, 0.1 in MeOH).
λ_{max} 205 (log ε 4.1); 221 (log ε 3.81);
233 (log ε 3.88); 252 (log ε 3.62); 262
(log ε 3.64) (MeOH).

Knorr, L. et al., *Ber.*, 1902, 35, 3007 (*synth*)


Clezy, P.S. et al., *Aust. J. Chem.*, 1975, 28, 1589

Puzicha, G. et al., *J. Het. Chem.*, 1990, 27, 2117 (*synth*)

Grond, S. et al., *Eur. J. Org. Chem.*, 2000, 929-937 (*rhamnopyranosyl ester*)

2-(3,4-Dimethyl-1*H*-pyrrol-1-yl)-3-methylbutanoic acid

*1-(1-Carboxy-2-methylpropyl)-3,4-dimethyl-1*H*-pyrrole. PP-Val*



C₁₁H₁₇NO₂ 195.261


(S)-form

Constit. of mixed purees of onion and garlic (*Allium cepa* and *Allium sativum*).
Needles. Mp 70-72°. [α]_D²⁴ -6.5 (c, 0.2 in MeOH). λ_{max} 230 (ε 5200) (MeOH). λ_{max} 230 (ε 5900) (MeOH/HCl). λ_{max} 220 (ε 4900) (MeOH/NaOH).

Imai, S. et al., *J. Agric. Food Chem.*, 2006, 54, 843-847 (*isol, uv, pmr, cmr*)

2-(3,4-Dimethyl-1*H*-pyrrol-1-yl)propanoic acid

*1-(1-Carboxyethyl)-3,4-dimethyl-1*H*-pyrrole. PP-Ala*



664

C₉H₁₃NO₂ 167.207


(S)-form

Constit. of mixed purees of onion and garlic (*Allium cepa* and *Allium sativum*).
Needles. Mp 38-40°. [α]_D²⁴ +22 (c, 0.2 in MeOH). λ_{max} 228 (ε 6300) (MeOH). λ_{max} 228 (ε 6300) (MeOH/HCl). λ_{max} 222 (ε 5600) (MeOH/NaOH).

Imai, S. et al., *J. Agric. Food Chem.*, 2006, 54, 843-847 (*isol, uv, pmr, cmr*)

2,4-Dimethylquinazoline, 9CI

[703-63-9]



C₁₀H₁₀N₂ 158.202

Metab. of the bacterium *Pseudomonas aeruginosa*. Cryst. + 2H₂O. Mp 72°. Bp₇₁₃ 249°. p*K*_a 3.58 (20°).

Picrate:

Yellow cryst. (EtOH aq.). Mp 170° dec.

1-Oxide: [37920-81-3]

C₁₀H₁₀N₂O 174.202
Mp 103°.

Bogert, M.T. et al., *J.A.C.S.*, 1924, 46, 1932 (*synth*)


Kövendi, A. et al., *Chem. Ber.*, 1965, 98, 1049 (*synth*)

Mann, S. et al., *Arch. Mikrobiol.*, 1967, 56, 324

Batterham, T.J. et al., *J.C.S. (B)*, 1967, 892 (*ms*)

Kant, J. et al., *J. Het. Chem.*, 1985, 22, 1313 (*synth*)

2,2-Dimethyl-4(1*H*,3*H*)-quinazolinone



C₁₀H₁₂N₂O 176.218


Prod. by *Cytophaga marinoflava* sp. AM13.1. Amorph. solid.

Shaaban, M. et al., *J. Nat. Prod.*, 2002, 65, 1660-1663 (*isol*)

Shabaan, M. et al., *Dissertation*, Univ. of Göttingen, 2004, (*isol, pmr, ms*)

2,4-Dimethylquinolizidine

*Octahydro-2,4-dimethyl-2*H*-quinolizine, 9CI*



C₁₁H₂₁N 167.294


*(2*R**,4*S**,9*aS**)-form*

Cermizine C

[760947-75-9]

Alkaloid from *Lycopodium cernuum*.Amorph. solid. $[\alpha]_D^{25} +4$ (c, 0.8 in MeOH).Morita, H. *et al.*, *Tetrahedron*, 2004, **60**, 7015-7023 (isol, pmr, cmr)Snider, B.B. *et al.*, *J.O.C.*, 2007, **72**, 1039-1042 (synth)**5-Dimethylsulfonio 2-amino-4-hydroxypentanoate** **D-784**


(4-Amino-4-carboxy-2-hydroxybutyl)dimethylsulfonium hydroxide inner salt, 9CI [117845-16-6]

C₇H₁₅NO₃S 193.266Isol. from the alga *Lophocladia lallemandi*.Sciuto, S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1017**N,N-Dimethyltryptamine** **D-785**

N,N-Dimethyl-1H-indole-3-ethanamine, 9CI. 3-(2-Dimethylaminoethyl)indole.

Nigerine. DMT

[61-50-7]

C₁₂H₁₆N₂ 188.272Alkaloid from *Mimosa hostilis*, *Acacia* spp., *Arundo donax*, *Desmodium* spp., *Phalaris* spp., *Banisteriopsis argentea*, *Psychotria* spp., *Viola peruviana*, *Zanthoxylum* spp. and others (Fabaceae, Poaceae, Malpighiaceae, Rubiaceae, Myristicaceae, Rutaceae). Shows affinity for 5HT_{1D}-receptor sites. Shows hallucinogenic props. Drug of abuse. Mp 48-49° (39-44°). Log P 1.82 (calc). λ_{max} 226 ; 275 (sh) ; 279 ; 284 ; 293 (EtOH).▶ LD₅₀ (mus, ipr) 47 mg/kg. NL7350000**Picrate:**

Yellow (stable) or red (metastable) cryst. Mp 171-172°.

N^b-Oxide: N,N-Dimethyltryptamine N-oxide

[948-19-6]

C₁₂H₁₆N₂O 204.271Alkaloid from *Desmodium pulchellum*, *Desmodium triflorum* and other *Desmodium* spp., also *Piptadenia peregrina* and *Piptadenia pulchellum* (Fabaceae). Hydrate. Mp 123-128°.**N^b-Oxide, picrate:**

Cryst. (EtOH). Mp 178-183°.

N^b-Me: N,N,N-Trimethyltryptamine

[17333-56-1]

C₁₃H₁₉N[⊕] 203.307Alkaloid from roots of *Zanthoxylum nitidum*. Isol. from the skin of the frog *Litoria moorei*. Mp 215-216° (as iodide).**N^b-Chloromethyl: N-Chloromethyl-N,N-dimethyltryptamine**

[941674-03-9]

C₁₃H₁₈ClN[⊕] 237.751Alkaloid from *Acacia confusa*.Amorph. solid. λ_{max} 217 (log ϵ 4.15); 280 (log ϵ 3.44); 290 (sh) (log ϵ 3.37) (MeOH).**1-Methoxy: 1-Methoxy-N,N-dimethyl-1H-indole-3-ethanamine. 3-[2-(Dimethylamino)ethyl]-1-methoxyindole. 1-Methoxy-N,N-dimethyltryptamine.****Lepedamine**


[4335-93-7]

C₁₃H₁₈N₂O 218.298Alkaloid from the leaves of *Lepedeza bicolor* var. *japonica* (Fabaceae). Oil. Bp_{0.28} 113-114° Bp_{0.08} 100-106°. λ_{max} 223 (log ϵ 4.49); 278 (log ϵ 3.66); 291 (log ϵ 3.68) (MeOH).**1-Methoxy, hydrochloride:**Needles (C₆H₆/MeOH). Mp 163-164° dec.**1-Methoxy, picrate:**


Yellow prisms (MeOH). Mp 160-162° dec.

Fish, M.S. *et al.*, *J.A.C.S.*, 1955, **77**, 5892-5895 (isol, oxide)Hochstein, F.A. *et al.*, *J.A.C.S.*, 1957, **79**, 5735-5736 (isol)Pachter, I.J. *et al.*, *J.O.C.*, 1959, **24**, 1285-1287 (isol)Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1964, **17**, 1301-1304 (isol)Morimoto, H. *et al.*, *Annalen*, 1965, **682**, 212-218 (*Lepedamine*, isol, uv, ir, pmr, struct)Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1965, **18**, 433-434 (isol)Ghosal, S. *et al.*, *J.O.C.*, 1966, **31**, 2284-2288 (isol)Falkenberg, G. *et al.*, *Acta Cryst. B*, 1972, **28**, 3075-3083 (cryst struct)Couch, M.W. *et al.*, *Anal. Biochem.*, 1972, **50**, 612-622 (ms)Roseghini, M. *et al.*, *Z. Naturforsch., C*, 1976, **31**, 118-120 (*Trimethyltryptamine*)Acheson, R.M. *et al.*, *J.C.S. Perkin 1*, 1978, 1117-1125 (*Lepedamine*, synth)Grina, J.A. *et al.*, *J.O.C.*, 1982, **47**, 2648-2651 (isol, uv, pmr, cmr)Sitaram, B.R. *et al.*, *Biochem. Pharmacol.*, 1987, **36**, 1509-1512 (metab)Glennon, R.A. *et al.*, *Drug Dev. Res.*, 1991, **22**, 25-36 (sar)Strassman, R.J. *et al.*, *Arch. Gen. Psychiatry*, 1994, **51**, 85-97; 98-108 (pharmacol, human)Moriyasu, M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 299-301 (*Trimethyltryptamine*)Somei, M. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 87-96 (*Lepedamine*, synth)Buchanan, M.S. *et al.*, *Magn. Reson. Chem.*, 2007, **45**, 359-361 (pmr, cmr, N-chloromethyl)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, DPF600**Dinklageine†**


[91668-36-9]

D-786C₁₉H₂₃NO₅ 345.394Alkaloid from the leaves of *Strychnos dinklagei* (Loganiaceae). Needles (CHCl₃). Mp 201-204°. $[\alpha]_D^{20} +60$ (c, 1 in MeOH).Skaltsounis, A.L. *et al.*, *Tet. Lett.*, 1984, **25**, 2783-2786 (uv, pmr, cmr, ms, struct, synth, abs config)**Dinklageine†****D-787**C₃₆H₃₈N₂O₆ 594.706Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the root and stem of *Stephania dinklagei* (Menispermaceae). Mp 285°. $[\alpha]_D -24.4$ (CHCl₃).Paris, R.A. *et al.*, *Ann. Pharm. Fr.*, 1955, **13**, 200-204; *CA*, **49**, 11959e (isol)**19,23-Dinorcona-****D-788****1,3,5(10),14-tetraen-3-ol**

3-Hydroxy-19-norcona-1,3,5(10),14-tetraene. 18,20-Imino-19-norpregna-1,3,5(10),14-tetraen-3-ol [91147-31-8]

C₂₀H₂₅NO 295.424Alkaloid from the stem bark of *Didymopanax cf. madagascariensis* (Didymelaceae). No phys. props. reported. λ_{max} 282 ; 286 (sh) (no solvent reported).Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr.*, Part II, 1984, 71-76 (isol, ir, uv, pmr, cmr)**Diolmycin A1****D-789**

1-(4-Hydroxyphenyl)-4-(1H-indol-3-yl)-2,3-butanediol, 9CI [150408-69-8]

C₁₈H₁₉NO₃ 297.353Prod. by a *Streptomyces* sp. Anticoccidial agent. Powder. $[\alpha]_D^{25} -8$ (c, 0.1 in MeOH). λ_{max} 222 (ϵ 33300); 280 (ϵ 5900); 290 (ϵ 4500) (MeOH) (Derep).**2'-Epimer: Diolmycin A2**

[150408-70-1]


C₁₈H₁₉NO₃ 297.353Prod. by a *Streptomyces* sp. Anticoccidial agent. Powder. Sol. MeOH, EtOH, MeCN, DMSO; poorly sol. CHCl₃, hexane. $[\alpha]_D^{25} -12$ (c, 0.1 in MeOH). λ_{max} 222 (ϵ 33300); 280 (ϵ

5900); 290 (ϵ 4500) (MeOH) (Derep).
[150407-38-8]

Tabata, N. *et al.*, *J. Antibiot.*, 1993, **46**, 756-761; 762-769 (*isol, synth, pmr, cmr, props*)
Fernandes, R.A. *et al.*, *Tetrahedron*, 2002, **58**, 1223-1227 (*synth, bibl*)

Dioncolactone A **D-790**


1,2,3,4-Tetrahydro-9-hydroxy-8-methoxy-2,4-dimethyl-6H-naphtho[1',2':4,5]pyrano[3,2-h]isoquinolin-6-one, 9CI
[135626-75-4]



Absolute Configuration

$C_{23}H_{21}NO_4$ 375.423
Alkaloid from *Triphyophyllum peltatum* (Dioncophyllaceae). Yellow solid. $[\alpha]_D^{25}$ -64 (c, 0.26 in $CHCl_3$).

Bringmann, G. *et al.*, *Phytochemistry*, 1991, **30**, 1691 (*isol, ir, pmr, ms, struct*)
Bringmann, G. *et al.*, *Tetrahedron*, 1999, **55**, 423-432 (*synth*)

Dioncophyllacine A **D-791**

$C_{26}H_{27}NO_4$ 417.504


(±)-**form** [146471-70-7]

Alkaloid from leaves of *Triphyophyllum peltatum* (Dioncophyllaceae). Needles (MeOH). Mp 177-179°.

Bringmann, G. *et al.*, *Phytochemistry*, 1992, **31**, 4015 (*isol, ir, pmr, ms, cryst struct*)

Dioncophylline D **D-792**

[853579-15-4]



$C_{24}H_{23}NO_3$ 373.451


Unlike Dioncophylline A, D-793, the biaryl axis is not stereogenic at normal temperatures. Alkaloid from the leaves of *Ancistrocladus benomensis*. Amorph.

solid (MeOH). Mp 212°. λ_{max} 231 (log ϵ 1.62); 315 (log ϵ 0.41); 375 (log ϵ 0.17) (MeOH).

Bringmann, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 686-690 (*isol, pmr, cmr, ms*)

Dioncophylline A **D-793**

7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-1,3-dimethyl-8-isoquinolinol, 9CI. Triphyophylline
[60142-17-8]



Absolute Configuration

$C_{24}H_{27}NO_3$ 377.482
Alkaloid from *Dioncophyllum thollonii* and *Triphyophyllum peltatum*. Shows antifungal and molluscicidal props. Insect antifeedant and growth retardant. Yellow cryst. (Me_2CO). Mp 215°. $[\alpha]_D^{25}$ -14 (c, 1 in $CHCl_3$).

N-Me: N-Methylidioncophylline A. N-Methyltryphyophylline

[135683-53-3]
[68727-49-1]
 $C_{25}H_{29}NO_3$ 391.509

Alkaloid from *Dioncophyllum thollonii* and *Triphyophyllum peltatum*. Cryst. (Me_2CO). Mp 193° (185°). $[\alpha]_D^{20}$ +70 (c, 1 in $CHCl_3$).

Me ether: O-Methylidioncophylline A. O-Methyltryphyophylline

$C_{25}H_{29}NO_3$ 391.509

Alkaloid from the stems of *Triphyophyllum peltatum*. Noncryst. $[\alpha]_D^{20}$ -30 (c, 1.2 in $CHCl_3$).

Me ether, N-Ac: Mp 177°. $[\alpha]_D^{20}$ +87 (c, 1.22 in $CHCl_3$).

O⁵-De-Me: 5'-O-Demethylidioncophylline A

[220436-87-3]
 $C_{23}H_{25}NO_3$ 363.455

Alkaloid from the root bark of *Triphyophyllum peltatum*. Needles (MeOH). Mp 233°. $[\alpha]_D^{20}$ -11.1 (c, 0.01 in $CHCl_3$).

O⁵-De-Me, O⁸-Me: 5'-O-Demethyl-8-O-methylidioncophylline A

[158705-45-4, 158664-19-8]
 $C_{24}H_{27}NO_3$ 377.482

Minor alkaloid from stem bark of *Triphyophyllum peltatum*. $[\alpha]_D^{20}$ +37.1 (c, 0.30 in $CHCl_3$).

1,2-Didehydro, Me ether: O-Methyl-1,2-didehydrotryphyophylline

[64996-70-9]
 $C_{25}H_{27}NO_3$ 389.493

Alkaloid from the stems of *Triphyophyllum peltatum*. Amorph. $[\alpha]_D^{20}$ 0 (solvent not specified).

1,2,3,4-Tetrahydro: Dioncophylline A
[148810-30-4]

$C_{24}H_{23}NO_3$ 373.451
Semisynthetic.

1,2,3,4-Tetrahydro, Me ether: O-Methyltetrahydrotryphyophylline
[68727-51-5]

$C_{25}H_{25}NO_3$ 387.477

Alkaloid from *Dioncophyllum thollonii* and *Triphyophyllum peltatum*. Cryst. (EtOH). Mp 166°. $[\alpha]_D^{20}$ 0 ($CHCl_3$).

1''-Hydroxy: 5'-O-Methylidioncopoline A. Habropetaline A

$C_{24}H_{27}NO_4$ 393.482

Alkaloid from *Habropetalum dawei* and *Triphyophyllum peltatum*. Antimalarial agent. Mp 225°. $[\alpha]_D^{25}$ -16.5 (c, 0.1 in MeOH).

1''-Hydroxy, O⁵-de-Me: 1,2,3,4-Tetrahydro-7-[5-hydroxy-2-(hydroxymethyl)-4-methoxy-1-naphthalenyl]-1,3-dimethyl-8-isoquinolinol. Dioncopoline A. Triphypeltine

[60158-81-8]

$C_{23}H_{25}NO_4$ 379.455

Alkaloid from *Dioncophyllum thollonii* and *Triphyophyllum peltatum*. Cryst. (MeOH). Mp 241°. $[\alpha]_D^{20}$ -96 (c, 0.5 in Py).

3-Epimer: Isotriphyophylline

[68780-11-0]

Minor alkaloid from *Triphyophyllum peltatum* and *Dioncophyllum thollonii*. Cryst. (Me_2CO). Mp 256°. $[\alpha]_D^{20}$ -22 (c, 1.2 in $CHCl_3$). Prob. abs. config.

Atropisomer: 7-Epidioncophylline A

[180324-35-0]

$C_{24}H_{27}NO_3$ 377.482

Alkaloid from *Ancistrocladus barteri*.

Atropisomer, N-Me: 7-Epi-N-methylidioncophylline A. N-Methyl-7-epidioncophylline A

[135683-54-4]

$C_{25}H_{29}NO_3$ 391.509

Alkaloid from *Ancistrocladus abbreviatus*. Cryst. (Me_2CO). Mp 230°. $[\alpha]_D^{25}$ -6 (c, 1.12 in $CHCl_3$).

Atropisomer, O⁵-de-Me, O⁸-Me: 5'-O-Demethyl-7-epi-8-O-methylidioncophylline A. 5'-O-Demethyl-8-O-methyl-7-epidioncophylline A

[135683-54-4]

$C_{24}H_{27}NO_3$ 377.482

Minor alkaloid from stem bark of *Triphyophyllum peltatum*. Amorph. powder. $[\alpha]_D^{20}$ -41.5 (c, 0.30 in $CHCl_3$).

Atropisomer, 1,2,3,4-tetrahydro: ent-Dioncophylline A

[853575-57-2]

$C_{24}H_{23}NO_3$ 373.451

Alkaloid from *Ancistrocladus benomensis*. Solid. Mp 265°.

Atropisomer, 1,2,3,4-tetrahydro, O⁵-de-Me: 5'-O-Demethyl-ent-dioncophylline A

[853575-58-3]

$C_{23}H_{21}NO_3$ 359.424

Alkaloid from *Ancistrocladus benomensis*.


Bruneton, J. *et al.*, *Phytochemistry*, 1976, **15**, 817 (*Dioncophylline A, Dioncopoline A*)

Lavault, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1977, **285**, 167; 1978, **287**, 129 (*derivs, uv, ir, pmr, struct, config*)

- Bringmann, G. *et al.*, *Phytochemistry*, 1981, **30**, 1691 (*Dioncopeltine A*)
- Bringmann, G. *et al.*, *Tet. Lett.*, 1984, **25**, 2537; 1990, **31**, 639; 643 (*O-Methyltetrahydrotriphophylline*)
- Bringmann, G. *et al.*, *Acta Cryst. C*, 1991, **47**, 1703 (*cryst struct*)
- Bringmann, G. *et al.*, *Annalen*, 1991, 803 (*stereochem*)
- Bringmann, G. *et al.*, *Phytochemistry*, 1994, **36**, 1057; 1998, **49**, 1667-1673 (*5'-O-Demethyl-8-O-methyldioncophylline A*, *5'-O-Demethyl-8-O-methyl-7-epi-dioncophylline A*, *5'-Dimethyldioncophylline A*)
- Bringmann, G. *et al.*, *J. Nat. Prod.*, 1997, **60**, 342 (*props*)
- Bringmann, G. *et al.*, *Magn. Reson. Chem.*, 1997, **35**, 397-301 (*pmr, config*)
- Bringmann, G. *et al.*, *Phytochemistry*, 1997, **45**, 1283 (*cryst struct, abs config*)
- Bringmann, G. *et al.*, *Synthesis*, 1999, 525-558 (*rev, synth*)
- Bringmann, G. *et al.*, *Tetrahedron*, 1999, **55**, 423-432 (*synth*)
- Bringmann, G. *et al.*, *Phytochemistry*, 2003, **62**, 345-349 (*Habropetaline A*)
- Bringmann, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 686-690 (*ent-Dioncophyllines*)
- Bringmann, G. *et al.*, *Tetrahedron*, 2007, **63**, 1755-1761 (*biosynth*)

Dioncophylline B**D-794**

[140367-82-4]

 $C_{23}H_{25}NO_3$ 363.455

Unlike Dioncophylline A, D-793, the biaryl axis is not stereogenic at normal temperatures. Alkaloid from the root bark of *Triphyophyllum peltatum* (Dioncophyllaceae). Antifungal agent, insecticide, antifedant. Needles + $\frac{1}{2}$ MeOH. $[\alpha]_D^{20}$ -37.6 (c, 0.37 in $CHCl_3$).

1-Epimer: 1-Epidioncophylline B

[216757-24-3]

[288846-61-7 (*Dioncophylline D*)] $C_{23}H_{25}NO_3$ 363.455

Alkaloid from *Triphyophyllum peltatum*. Amorph. yellow solid (MeOH). Mp 234-236°. $[\alpha]_D^{25}$ +15.5 (c, 0.1 in EtOH). Struct. revised and renamed in 2000; formerly given the name Dioncophylline D.

1-Epimer, 8-Me ether: 8-O-Methyl-1-epi-dioncophylline B $C_{24}H_{27}NO_3$ 377.482

Alkaloid from *Triphyophyllum peltatum*. Amorph. yellow solid (MeOH). Mp 228-232°. $[\alpha]_D^{25}$ +21 (c, 0.05 in $CHCl_3$). Struct. revised and renamed in 2000.


Bringmann, G. *et al.*, *Phytochemistry*, 1991, **30**, 3845; 1998, **49**, 1151-1155; 2000, **54**, 337-346 (*isol, ir, pmr, cmr, struct*)

Bringmann, G. *et al.*, *Synthesis*, 2000, 1843-1847 (*synth*)

Bringmann, G. *et al.*, *Tetrahedron*, 2001, **57**, 1253-1259 (*synth*)

Dioncophylline C**D-795**

[146471-75-2]

 $C_{23}H_{25}NO_3$ 363.455

The biaryl axis is stereogenic (cf. Dioncophylline A, D-793). Alkaloid from root bark of *Triphyophyllum peltatum* (Dioncophyllaceae). Very potent anti-malarial agent. Mp 246° dec. $[\alpha]_D^{20}$ +19.2 (c, 0.52 in $CHCl_3$).

Bringmann, G. *et al.*, *Phytochemistry*, 1992, **31**, 4019 (*isol, ir, pmr, cd, ms, struct*)


Francois, G. *et al.*, *J. Ethnopharmacol.*, 1995, **46**, 115-120 (*isol, activity*)

Francois, G. *et al.*, *Ann. Trop. Med. Parasitol.*, 1996, **90**, 115-123 (*activity*)

Bringmann, G. *et al.*, *Tetrahedron*, 1998, **54**, 497-512 (*synth*)

Dioncophyllinol B**D-796**

[207676-81-1]

 $C_{23}H_{25}NO_4$ 379.455

Alkaloid from the leaves of *Triphyophyllum peltatum*. Yellow solid. Mp 209°. $[\alpha]_D^{25}$ +17 (c, 0.05 in $CHCl_3$). Struct. of isolate revised in 2000. Prev. named Dioncophyllinol D.

8-Me ether: 8-O-Methyldioncophyllinol B

[288846-62-8]

 $C_{24}H_{27}NO_4$ 393.482


Alkaloid from *Triphyophyllum peltatum*. Yellow solid. $[\alpha]_D^{25}$ -12.8 (c, 0.7 in $CHCl_3$).

Bringmann, G. *et al.*, *Heterocycles*, 1998, **47**, 985-990 (*isol, ir, pmr, cmr*)

Bringmann, G. *et al.*, *Phytochemistry*, 2000, **54**, 337-346 (*isol, struct*)

Dioscoretine**D-797**

[128637-87-6]

 $C_{13}H_{23}NO_3$ 241.33

Alkaloid from the tubers of the famine food *Dioscorea dumetorum* (Dioscorea-


ceae). Hypoglycaemic principle.

► LD₅₀ 580mg/kg (mice). CL5602600

Iwu, M.M. *et al.*, *Planta Med.*, 1990, **56**, 119-120; 264-267 (*isol, cmr, struct, activity*)

Dioscorine**D-798**

2,4'-Dimethylspiro[2-azabicyclo[2.2.2]octane-5,2'-[2H]pyran]-6'-(3'H)-one, 9CI
[3329-91-7]



Absolute Configuration

 $C_{13}H_{19}NO_2$ 221.299

Alkaloid from the tubers of *Dioscorea hirsuta* and the wild yam *Dioscorea hispida* (Dioscoreaceae). Analeptic agent, CNS stimulant. Mp 43.5° (34-35°). $[\alpha]_D^{18}$ -35 (c, 3.4 in $CHCl_3$). Log P 0.76 (calc).

► LD₅₀ (mus, ipr) 60 mg/kg. Exp. convulsant effects. JG5787500

Hydrochloride: Mp 210-211° dec.

Picrate: Mp 187° dec.

N-Oxide: Dioscorine N-oxide

[171432-58-9, 171595-79-2]

 $C_{13}H_{19}NO_3$ 237.298

Alkaloid from *Dioscorea hispida* (Dioscoreaceae).

4'S,5'-Dihydro: Dihydrodioscorine

[96552-66-8]

 $C_{13}H_{21}NO_2$ 223.314

Alkaloid from the tubers of the famine food *Dioscorea dumetorum* and *Dioscorea sanziborensis* (Dioscoreaceae). Oil. Sol. H₂O. $[\alpha]_D^{18}$ -42.2 (c, 3.4 in $CHCl_3$). Dihydrodioscorine cannot be obt. by hydrogenation of Dioscorine. λ_{max} 208 (MeOH) (Berdy).

► Convulsive poison; LD₅₀ (mus, ipr) 0.65 mg/kg.

4'S,5'-Dihydro, hydrochloride: Mp 209°.

4'S,5'-Dihydro, picrate: Mp 189°.

Pinder, A.R. *et al.*, *J.C.S.*, 1952, 2236 (*isol, uv, ir*)

Broadbent, J.L. *et al.*, *Br. J. Pharmacol.*, 1958, **13**, 213 (*pharmacol*)

Bevan, C.W.L. *et al.*, *Chem. Ind. (London)*, 1958, 103 (*Dihydrodioscorine, isol*)

Davies, W.A.M. *et al.*, *Chem. Ind. (London)*, 1961, 1410 (*struct*)

Correia Alves, A. *et al.*, *CA*, 1963, **59**, 6452d (*Dihydrodioscorine, isol*)

Beecham, A.F. *et al.*, *Tet. Lett.*, 1969, 3745 (*config*)


Leete, E. *et al.*, *Phytochemistry*, 1977, **16**, 1704 (*biosynth, cmr*)

Corley, D.G. *et al.*, *Tet. Lett.*, 1985, **26**, 1615 (*Dihydrodioscorine, isol, struct, abs config*)

Goh, S.-H. *et al.*, *Malays. J. Sci.*, 1994, **15**, 23; *CA*, **124**, 25562b (*Dioscorine N-oxide*)

Dioxamine**D-799**

3-(1,3-Benzodioxol-5-yl)-N-[2-(1,3-benzodioxol-5-yl)ethyl]-2-propenamide, 9CI. 3,4-Methylenedioxybenzylidene 3,4-methylenedioxyphenylethylamide [119060-89-8]



$C_{19}H_{17}NO_5$ 339.347

Alkaloid from the roots and pericarps of *Zanthoxylum rubescens* (Rutaceae). Powder. Mp 144-146°.

N-Me: Dioxamide

[119060-88-7]

 $C_{20}H_{19}NO_5$ 353.374


Alkaloid from roots and pericarps of *Zanthoxylum rubescens* (Rutaceae). Amorph.

Adesina, S.K. *et al.*, *Pharmazie*, 1988, **43**, 517; *CA*, **110**, 92013x

Adesina, S.K. *et al.*, *Phytochemistry*, 1989, **28**, 839 (isol, uv, ir, pmr, cmr, ms, synth, struct)

Dioxinoacrimarine A**D-800**

[158182-14-0]




$C_{29}H_{23}NO_8$ 513.503

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) and Dancy tangerine (*Citrus tangerina*)] (Rutaceae). Yellow cubes (Me₂CO). Mp 179-181°. [α]_D +18 (c, 0.1 in CHCl₃).

Takemura, Y. *et al.*, *Heterocycles*, 1994, **38**, 1937 (isol, uv, ir, pmr, cmr, ms, struct)

Dioxocularine**D-801**

[107602-50-6]



$C_{20}H_{17}NO_6$ 367.357

The trivial name is misleading (Cularine, C-805 is 1,α-dihydro). Trace alkaloid from *Corydalis claviculata* (Papaveraceae). Red cryst. (EtOH). Mp 212-214°.

Boente, J.M. *et al.*, *Heterocycles*, 1986, **24**, 3359 (uv, ir, pmr, cmr, ms, struct)


García, A. *et al.*, *Tetrahedron*, 1995, **51**, 8585 (synth)

Suau, R. *et al.*, *Tet. Lett.*, 1996, **37**, 9357

(synth)

4,5-Dioxodehydrocrebanine**D-802**


9,10-Dimethoxy-7-methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline-5,6(7H)-dione, 9CI [77784-21-5]



$C_{20}H_{15}NO_6$ 365.342

Alkaloid from the stem and root of *Stephania sasakii*. Also obt. by air oxidation of dehydrocrebanine in the presence of alkali (Menispermaceae). Orange needles (EtOH). Mp 278-280°.

Kunitomo, J. *et al.*, *Phytochemistry*, 1980, **19**, 2735 (isol, uv, ir, pmr, ms, struct, synth)

5,10-Dioxo-1,6-diazacyclo-decane-2,7-dicarboxylic acid**D-803***γ*-Cyclic diglutamic acid

$C_{10}H_{14}N_2O_6$ 258.23

(S,S)-form [35009-41-7]

[869485-13-2 (di-Na salt)]

Isol. from the pupae of *Bombyx mori*.


Also prod. by *Trichophyton mentagrophytes*. Powder (as di-Na salt). Mp 240° dec. (di-Na salt). [α]_D -13.8 (c, 1.7 in H₂O) (di-Na salt).

Wöllmann, H. *et al.*, *Pharmazie*, 1977, **32**, 185-186 (isol)

Tanaka, R. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 3055-3062 (isol, cd, pmr, cmr)

4-(2,5-Dioxo-4-imidazolidinyl)carbamic acid**D-804**

5-Carboxyamino-2,4-imidazolidinedione

**(R)-form**

$C_4H_5N_3O_4$ 159.101

(R)-form [244172-47-2]Constit. of *Cistanche deserticola*.


Amide: see Allantoin, A-625

Xu, Z. *et al.*, *J. Chin. Pharm. Sci.*, 1999, **8**, 61-63 (isol, cryst struct)

10,22-Dioxokopsane**D-805**

Kopsan-10,22-dione, 9CI. 10-Oxokopsanone

[3703-90-0]



$C_{20}H_{20}N_2O_2$ 320.39

Alkaloid from the stem bark of *Pleiocarpa mutica* and from root bark of *Alstonia venenata* (Apocynaceae). Needles (MeOH), cryst. (EtOAc). Mp 307-308° (298°). [α]_D +123 (c, 0.635 in CHCl₃). [α]_D +156 (c, 0.32 in CHCl₃).

N-Ac:

Cryst. (EtOAc). Mp 254-255°.

N-Me: N-Methyl-10,22-dioxokopsane

[3703-91-1]

 $C_{21}H_{22}N_2O_2$ 334.417

Alkaloid from *Pleiocarpa mutica* (Apocynaceae). Cubes (MeOH). Mp 269°. [α]_D +74 (c, 0.61 in MeOH).

N-Methoxycarbonyl: N-Methoxycarbonyl-10,22-dioxokopsane

[3901-67-5]

 $C_{22}H_{22}N_2O_4$ 378.427

Alkaloid from *Pleiocarpa mutica* (Apocynaceae). Needles (CHCl₃/petrol). Mp 264-265°. [α]_D +110 (c, 1.31 in CHCl₃).

22R-Alcohol: Epikopsan-22-ol-10-lactam.

Epikopsanol-10-lactam. 22-Hydroxy-10-kopsanone

[6698-63-1]

[23495-95-6]

 $C_{20}H_{22}N_2O_2$ 322.406

Alkaloid from *Aspidosperma duckei* and *Aspidosperma verbascifolium* (Apocynaceae). Mp 242-245° dec.

22R-Alcohol, N⁴,O-di-Ac: Mp 285-287°.[α]_D +91.5 (c, 1.18 in CHCl₃).**14,15-Didehydro: 14,15-Didehydro-10,22-dioxokopsane. Kopsorinine**


[768388-31-4]

 $C_{20}H_{18}N_2O_2$ 318.374

Alkaloid from *Kopsia fruticosa*. Light yellow oil. $[\alpha]_D^{25} +14$ (c, 0.35 in CHCl_3). λ_{max} 220 (log ϵ 4.49); 245 (log ϵ 3.95); 280 (log ϵ 3.39) (EtOH).

- Achenbach, H. *et al.*, *J.A.C.S.*, 1965, **87**, 4944-4950 (*Pleiocarpa mutica constits*)
 Kump, C. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 1237-1243 (*synth, uv, ir, ms*)
 Ferreira Filho, J.M. *et al.*, *J.C.S.(C)*, 1966, 1260-1266 (*Epikopsanol-10-lactam*)
 Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*, 1969, **78**, 63-68 (*Aspidosperma verbascifolium constits*)
 Chatterjee, A. *et al.*, *Indian J. Chem., Sect. B*, 1979, **17**, 651 (*5,22-Dioxokopsane*)
 Magnus, P. *et al.*, *J.A.C.S.*, 1984, **106**, 2105-2114 (*synth, pmr*)
 Kam, T.-S. *et al.*, *Phytochemistry*, 2004, **65**, 2119-2122 (*Kopsorinine*)


Dioxolamycin D-806
 [91432-46-1]



$\text{C}_{11}\text{H}_{15}\text{NO}_6$ 257.243
 Isol. from *Streptomyces filamentosus*. Shows moderate cytostatic activity against L-1210 cells. Cryst. (EtOAc). Sol. MeOH, CHCl_3 ; poorly sol. C_6H_6 , hexane, H_2O . Mp 214-215°. $[\alpha]_D^{25} -53.1$ (c, 0.36 in MeOH). λ_{max} 218 (ϵ 8130) (MeOH) (Derrep). λ_{max} 218 (E1%/1cm 316) (MeOH) (Berdy).

- LD_{50} (mus, ivn) 200 - 400 mg/kg. DF4912775
 Zhu, B. *et al.*, *J. Antibiot.*, 1984, **37**, 673 (*isol, uv, ir, pmr, cmr*)

Dioxolide A D-807
 3a,4-Dihydro-2-hydroxymethyl-1,3-benzodioxole-5-carboxamide, 9CI
 [176226-75-8]



$\text{C}_9\text{H}_{11}\text{NO}_4$ 197.19
 Prod. by *Streptomyces tendae*. λ_{max} 308 (MeOH) (Berdy).


O-Ac: **Dioxolide B**
 [176226-76-9]
 $\text{C}_{11}\text{H}_{13}\text{NO}_5$ 239.227
 Prod. by *Streptomyces tendae*. λ_{max} 308 (MeOH) (Berdy).

3a,4-Didehydro: **Dehydrodioxolide A**
 [176226-77-0]
 $\text{C}_9\text{H}_9\text{NO}_4$ 195.174
 Prod. by *Streptomyces tendae*. λ_{max} 258; 295 (MeOH). λ_{max} 258; 295 (MeOH) (Berdy).

3a,4-Didehydro, O-Ac: **Dehydrodioxolide B**
 [176226-78-1]
 $\text{C}_{11}\text{H}_{11}\text{NO}_5$ 237.212
 Prod. by *Streptomyces tendae*. λ_{max} 258; 295 (MeOH) (Berdy).

Blum, S. *et al.*, *J. Basic Microbiol.*, 1996, **36**, 19-25 (*isol*)


N-(2,6-Dioxo-3-piperidinyl)-benzeneacetamide, 9CI D-808
 3-(N-Phenylacetylamino)piperidine-2,6-dione. **Antineoplaston A10**
 [77658-84-5]



$\text{C}_{13}\text{H}_{14}\text{N}_2\text{O}_3$ 246.265

- (S)-form [91531-30-5]
 Isol. from human urine and plasma. Antineoplastic agent. Inhibits human hepatocellular carcinoma *in vitro* and *in vivo* in combination with Diamminedichloroplatinum(II). Cryst. (MeOH).
 ► LD_{50} (mus, ipr) 10.33 mg/kg. [158930-26-8, 108929-54-0]
Eur. Pat., 1983, 69 232; *CA*, **98**, 137623 (*synth, pharmacol*)
 Burzynski, S.R. *et al.*, *Drugs Exp. Clin. Res.*, 1984, **10**, 113; 1986, **12**, 11; 37; 57; 1987, **13**, 77 (*pharmacol*)
 Xu, W. *et al.*, *Yiyao Gongye*, 1988, **19**, 198; *CA*, **110**, 75258 (*synth*)
 Tsuda, H. *et al.*, *Jpn. J. Cancer Res.*, 1992, **83**, 527 (*pharmacol*)
 Michalska, D. *et al.*, *Spectrochim. Acta A*, 1993, **49**, 303 (*ir, Raman, bibl*)
 Martindale, *The Extra Pharmacopoeia*, 31st edn., Pharmaceutical Press, 1996, 543


6-(6,8-Dioxo-1-purinyl)suberosanone D-809
 [1015765-87-3]



$\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}_3$ 370.45
 Alkaloid from *Subergorgia suberosa*. Powder. $[\alpha]_D^{20} +27.3$ (c, 0.2 in CHCl_3). λ_{max} 212; 265 (MeOH).

Qi, S.-H. *et al.*, *J. Nat. Prod.*, 2008, **71**, 716-718 (*isol, pmr, cmr*)

10,17-Dioxosparteine D-810
 Alkaloid B3
 [52717-73-4]




Absolute Configuration

$\text{C}_{15}\text{H}_{22}\text{N}_2\text{O}_2$ 262.351
 Alkaloid from *Lupinus hartwegii* whole plants (Fabaceae). Liq.

Anderson, J.N. *et al.*, *J.O.C.*, 1976, **41**, 3441-3444 (*isol, struct, ms*)

Dioxybrassinin D-811
 Methyl [(2,3-dihydro-3-hydroxy-2-oxo-1H-indol-3-yl)methyl]carbamdithioate, 9CI. **Dioxybrassinin**




$\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_2\text{S}_2$ 268.36

(S)-form [133761-64-5]
 Alkaloid from *Brassica oleracea* (Brassicaceae) inoculated with *Pseudomonas cichorii*. $[\alpha]_D -7.6$ (c, 0.6 in MeOH).

(±)-form [137821-31-9]
 Mp 164-166°. Powell, R.G. *et al.*, *Experientia*, 1991, **47**, 304 (*isol, pmr, cmr, ms, cryst struct*)
 Monde, K. *et al.*, *Phytochemistry*, 1991, **30**, 2915 (*isol, uv, ir, pmr, ms, struct, synth*)
 Monde, K. *et al.*, *Tet. Lett.*, 2003, **44**, 6017-6020 (*synth, cd, abs config*)

Dipepine D-812
 1',2,2',3,3',9'-Hexahydro[1,9'-bipyrrrolo[2,1-b]quinazolin]-9(1H)-one, 9CI
 [53416-72-1]




$\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}$ 356.426
 Alkaloid from *Peganum harmala* (Zygophyllaceae). Cryst. (Me_2CO). Mp 221-223°. The first known dimeric quinazolinone alkaloid. λ_{max} 226 (log ϵ 4.48); 277 (log ϵ 4.14); 305 (log ϵ 4.05); 317 (log ϵ 3.89) (CCl_4).

9-Hydroxy: **Dipeginol**
 $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_2$ 372.426
 Alkaloid from *Peganum harmala*. Prisms (MeOH). Mp 243° dec. λ_{max} 225 (log ϵ 4.46); 276 (log ϵ 4.11); 303 (log ϵ 4.04); 317 (log ϵ 3.82) (CCl_4).

Zharekeev, B.K. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 264; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 282 (*uv, ir, cmr, ms, struct*)

Faskhutdinov, M.F. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 602-605 (*Dipeginol*)

2,6-Di-4-pentenylpiperidine D-813C₁₅H₂₇N 221.385**(2RS,6RS)-form**(±)-trans-form
[127629-05-4]Alkaloid from the venom of the ant *Monomorium delagoense*. Shows insecticidal props.**4',5'-Dihydro: trans-2-(4-Pentenyl)-6-pentylpiperidine**

[127629-04-3]

C₁₅H₂₉N 223.401Alkaloid from *Monomorium delagoense*. Shows insecticidal and repellent props.**(2RS,6SR)-form**

(±)-cis-form

[127629-03-2]


Alkaloid from *Monomorium delagoense*. Shows insecticidal and repellent props.**4',5'-Dihydro: cis-2-(4-Pentenyl)-6-pentylpiperidine**

[127629-02-1]

C₁₅H₂₉N 223.401Alkaloid from *Monomorium delagoense*. Shows insecticidal and repellent props.Jones, T.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 429
(*isol, struct, synth*)**Diphenazithionin**


D-814

[185843-29-2]

C₃₀H₁₈N₄O₉S 610.56Isol. from *Streptomyces griseus* ISP 5236. Antioxidant. Dark green powder. λ_{max} 260; 305; 375; 442 (no solvent reported).Hosoya, Y. *et al.*, *Tet. Lett.*, 1996, **37**, 9227
(*isol, uv, pmr, cmr, cryst struct*)**Diphenyl ether-3,3',5,5'-tetracarboxylic acid**

D-815

5,5'-Oxybis-1,3-benzenedicarboxylic acid

C₁₆H₁₀O₉ 346.25**3,3'-Diamide: 3,3'-Oxybis[5-(aminocarbonyl)benzoic acid]. 3,3'-Bis(aminocarbonyl)-5,5'-dicarboxydiphenyl ether**

[177334-43-9]

C₁₆H₁₂N₂O₇ 344.28Prod. by *Fusarium oxysporum*.


Siderophore.

Wijesundera, R.L.C. *et al.*, *CA*, 1996, **125**, 5167u**4,8-Diphenyl-5H-1,3-dioxolo[4,5-d]azepine-5,7(6H)-dione, 9CI**

D-816


Ustalic acid imide

[470696-80-1]

C₁₉H₁₃NO₄ 319.316Isol. from *Tricholoma ustale*. Cryst. Mp 239-242°.Sano, Y. *et al.*, *Chem. Comm.*, 2002, 1384-1385
(*isol, pmr, cmr*)**2,3-Diphenyl-1H-pyrrole, 9CI**

D-817

[26093-30-1]

C₁₆H₁₃N 219.285Constit. of the stems of *Gnetum montanum*. Needles (EtOH aq.). Mp 127°.

N-Me:

C₁₇H₁₅N 233.312

Cryst. Mp 96-96.5°.

N-Benzyl:

C₂₃H₁₉N 309.41

Cryst. Mp 116-117°.

N-Ph: 1,2,3-Triphenyl-1H-pyrrole

[53646-86-9]

C₂₂H₁₇N 295.383

Prisms (EtOAc/hexane). Mp 181-182°.

Severin, T. *et al.*, *Chem. Ber.*, 1971, **104**, 2856-2863 (*synth*)Zhou, J. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 878-882 (*isol, pmr, cmr*)Katritzky, A.R. *et al.*, *J.O.C.*, 1994, **59**, 4551-4555 (*synth, pmr, cmr*)Di Santo, R. *et al.*, *Synth. Commun.*, 1995, **25**, 795 (*synth*)Katritzky, A.R. *et al.*, *J. Het. Chem.*, 1997, **34**, 1379-1381 (*N-Ph*)Romashin, Y.N. *et al.*, *Chem. Comm.*, 1999, 447-448 (*N-Me, N-Benzyl, synth, pmr*)***N,N'*-Diphenylurea, 9CI**

D-818

Carbanilide

[102-07-8]

[26763-63-3]

PhNHCONHPh

C₁₃H₁₂N₂O 212.251Isol. from coconut milk (*Cocos nucifera*).

Cytokinin. Needles (EtOAc); prisms. Mp 246-248° (239-240°). Bp 260°.

▶FD9800000

Mono-N-Ac:

C₁₅H₁₄N₂O₂ 254.288

Mp 106°.

N-Me: N-Methyl-N,N'-diphenylurea

[612-01-1]

C₁₄H₁₄N₂O 226.277

Cryst. (EtOH or xylene). Mp 106°. Bp 203-205° dec.

N,N'-Di-Me: N,N'-Dimethyl-N,N'-diphenylurea. Centralite II

[611-92-7]

C₁₅H₁₆N₂O 240.304

Explosion regulator. Mp 122°.

▶FE0600000

N-Et: N-Ethyl-N,N'-diphenylurea

[64544-71-4]

C₁₅H₁₆N₂O 240.304

Prisms (EtOH). Mp 91°.

N-Et, N'-Me: N-Ethyl-N'-methyl-N,N'-diphenylurea. Centralite III

[4474-03-7]

C₁₆H₁₈N₂O 254.331

Explosion regulator. Mp 58-59°.

N,N'-Di-Et: N,N'-Diethyl-N,N'-diphenylurea. Carbamate. Centralite I. Ethyl centralite

[85-98-3]

C₁₇H₂₀N₂O 268.358

Explosion regulator. Cryst. (EtOH).


Mp 79° (72-73°). Bp 327°.

▶Fl. p. 150°. FE0350000

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 387A (*ir*)Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1418C; 1420C; 1421A (*nmr*)Org. Synth., Coll. Vol., 1, 1932, 453-455 (*synth*)Shantz, E.M. *et al.*, *J.A.C.S.*, 1952, **74**, 6133-6135; 1955, **77**, 6351-6353 (*isol*)Baker, J.W. *et al.*, *J.C.S.*, 1957, 4652-4662(*synth*)Novacek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 1712-1718 (*synth*)Deshapande, S.V. *et al.*, *Acta Cryst. B*, 1968, **24**, 1396-1397 (*cryst struct*)Knobler, Y. *et al.*, *Isr. J. Chem.*, 1971, **9**, 165-175 (*synth*)Norikov, E.G. *et al.*, *Zh. Prikl. Spektrosk.*, 1971, **14**, 833 (*uv*)Siti, M.P. *et al.*, *J.O.C.*, 1979, **44**, 3017-3022(*cmr, N-15 nmr*)Ayyangar, N.R. *et al.*, *Chem. Ind. (London)*, 1988, 599-600 (*synth, bibl, derivs*)Laufer, D.A. *et al.*, *Org. Prep. Proced. Int.*, 1989, **21**, 771-776 (*synth*)Cooper, C.F. *et al.*, *Synth. Commun.*, 1995, **25**, 2467-2474 (*synth*)Jimenez Blanco, J.L. *et al.*, *Synthesis*, 1999, 1907-1914 (*synth*)Hikita, T. *et al.*, *Macromolecules*, 2002, **35**, 6202-6209 (*synth, ir, pmr, cmr*)Artamkina, G.A. *et al.*, *Russ. J. Org. Chem. (Engl. Transl.)*, 2002, **38**, 538-545 (*synth, ir, pmr*)Tran, K. *et al.*, *Org. Prep. Proced. Int.*, 2004, **36**, 71-73 (*synth, ir, pmr, cmr*)Perveen, S. *et al.*, *Synth. Commun.*, 2005, **35**, 1663-1674 (*synth, ir, uv, pmr, ms*)Mizuno, T. *et al.*, *Synthesis*, 2007, 2825-2830; 3135-3140 (*synth, ir, pmr, cmr, ms*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CBM250; DRB200; DJC400

Dipiperamide A

[432041-19-5]

C₃₄H₃₈N₂O₆ 570.684

Stereochem. revised in 2005 to a *meso*-stereoisomer (centre of symmetry). Dipiperamide B was formerly assigned this stereochem. but now needs to be revised. Alkaloid from white pepper, *Piper nigrum*. Inhibitor of cytochrome P450 3A4 activity. Mp 175-176° (synthetic). λ_{max} 203 (log ε 4.9); 238 (sh) (log ε 4.4); 287 (log ε 4) (EtOH).

Diastereoisomer: Dipiperamide B

[432041-20-8]

C₃₄H₃₈N₂O₆ 570.684


Alkaloid from white pepper, *Piper nigrum*. Racemic. λ_{max} 204 (log ε 4.8); 237 (sh) (log ε 4.4); 273 (sh) (log ε 4.3); 289 (sh) (log ε 4.2) (EtOH).

Tsukamoto, S. *et al.*, *Tetrahedron*, 2002, **58**, 1667-1671 (*isol*, *pmr*, *cmr*, *ms*)

Takahashi, M. *et al.*, *Tet. Lett.*, 2005, **46**, 57-50 (*synth*, *stereochem*)

Dipiperamide C

[432041-21-9]


C₃₃H₃₆N₂O₆ 556.657

Alkaloid from white pepper, *Piper nigrum*. Inhibitor of cytochrome P450 3A4 activity. Racemic. λ_{max} 207 (log ε 4.6); 268 (log ε 4.3); 278 (sh) (log ε 4.2); 308 (log ε 4.1); 325 (sh) (log ε 3.8) (EtOH).

Tsukamoto, S. *et al.*, *Tetrahedron*, 2002, **58**, 1667-1671 (*isol*, *pmr*, *cmr*, *ms*)

Dipiperamide D

[500757-85-7]



Relative Configuration


D-819C₃₆H₄₀N₂O₆ 596.722

Alkaloid from white pepper, *Piper nigrum*. Inhibitor of cytochrome P450 3A4 activity. λ_{max} 209 (log ε 4.6); 266 (log ε 4.3) (EtOH).

Tsukamoto, S. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 2981-2985 (*isol*, *pmr*, *cmr*)

Dipiperamide E

[500757-86-8]



Relative Configuration


D-822C₃₄H₃₈N₂O₆ 570.684

Alkaloid from white pepper, *Piper nigrum*. Inhibitor of cytochrome P450 3A4 activity. λ_{max} 206 (log ε 3.8); 242 (sh) (log ε 4.1); 266 (sh) (log ε 4); 289 (log ε 4.2) (EtOH).

Tsukamoto, S. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 2981-2985 (*isol*, *pmr*, *cmr*)

Diplamine

[123794-30-9]

**D-823**C₂₀H₁₇N₃O₂S 363.439

Related to Deacylcystodytin, D-97. Alkaloid from the tunicate *Diplosoma* sp. Exhibits antitussive and antitumour activity. Burnt-orange solid. Mp 202-204° dec. λ_{max} 263 (ε 24900); 300 (ε 14700); 377 (ε 6000) (MeOH) (Derep).

N-De-Ac, N-(3-methylbutanoyl): Lissoclin A

[158761-12-7]

C₂₃H₂₃N₃O₂S 405.52

Alkaloid from the tropical ascidian *Lissoclinum* sp. and *Didemnum* sp. Orange needles (MeOH). Mp 202-204°, λ_{max} 263 (ε 28000); 300 (ε 17000); 383 (ε 6800); 450 (ε 4900) (MeOH) (Berdy).

N-De-Ac, N-tigloyl: Lissoclin B. (Demethylthio)cystodytin B

[158761-13-8]

C₂₃H₂₁N₃O₂S 403.504

Alkaloid from *Lissoclinum* sp. and *Didemnum* sp. Orange solid. λ_{max} 263 (ε 28000); 300 (ε 17000); 383 (ε 6800); 450 (ε 4900) (MeOH) (Berdy).

De(methylthio), 5-methylthio: Isodipla-**mine**

[496909-75-2]

C₂₀H₁₇N₃O₂S 363.439

Alkaloid from *Lissoclinum notti*. Green solid. Mp 208-210°. λ_{max} 215 (log ε 4.68); 274 (log ε 4.16); 320 (log ε 4.01); 406 (log ε 4) (MeOH/TFA).

Charyulu, G.A. *et al.*, *Tet. Lett.*, 1989, **30**, 4201-4202 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Szczepankiewicz, B.G. *et al.*, *J.O.C.*, 1994, **59**, 3512-3513 (*synth*)


Searle, P.A. *et al.*, *J.O.C.*, 1994, **59**, 6600-6605 (*Lissoclins*)

Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1995, **117**, 12460 (*synth*)

Appleton, D.R. *et al.*, *Tetrahedron*, 2002, **58**, 9779-9783 (*Isodiplamine*)

Dippinine C

[223532-43-2]



Absolute Configuration


D-824C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the stems of *Tabernaemontana corymbosa*. Light yellow oil. [α]_D +19 (c, 0.07 in CHCl₃). λ_{max} 228 (log ε 4.46); 285 (log ε 3.91); 294 (log ε 3.86) (EtOH).

11-Methoxy, 3-ketone: Dippinine DC₂₃H₂₆N₂O₅ 410.469

Alkaloid from the leaves of *Tabernaemontana corymbosa*. Light yellow oil. [α]_D -152 (c, 0.06 in CHCl₃). λ_{max} 235 (log ε 4.35); 280 (log ε 4.05); 290 (log ε 4.01) (EtOH).

Kam, T.-S. *et al.*, *Heterocycles*, 1999, **51**, 345-348; 2001, **55**, 2405-2412 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriaconta-7,10,22,25-tetraene-15,30-dione**D-825**C₃₂H₅₄N₂O₄ 530.79

Constit. of the pupal defensive secretions of the ladybird beetle *Subcoccinella vigintiquatuorpuntata* (*Subcoccinella 24-punctata*).

7,8-Dihydro: 5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriaconta-7,10,25-triene-15,30-dioneC₃₂H₅₆N₂O₄ 532.805

Isol. from *Subcoccinella vigintiquatuorpuntata*.

7,8,22,23-Tetrahydro: 5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriaconta-10,25-diene-15,30-dioneC₃₂H₅₈N₂O₄ 534.821

Isol. from *Subcoccinella vigintiquatuorpunctata*.

Schroeder, F.C. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1998, **95**, 13387-13391

Gronquist, M.R. et al., *J.O.C.*, 2001, **66**, 1075-1081

Diptaline **D-826**

[11-(Methylsulfinyl)undecyl]urea, 9CI.

Diphthaline

[94898-72-3]

H₂NCONH(CH₂)₁₁SOMe

C₁₃H₂₈N₂O₂S 276.442

Alkaloid from the seeds of *Diptychocarpus strictus* (Brassicaceae). Oil. [α]_D -10.2 (MeOH).

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1984, **400**; *Chem. Nat. Compd. (Engl. Transl.)*, 380 (isol, ir, struct)

Diptamine **D-827**

N-(1-Methylethyl)-N'-[7-(methylsulfinyl)heptyl]urea, 9CI. Diptamine

[75069-56-6]

(H₃C)₂CHNCONH(CH₂)₇SOMe

C₁₂H₂₆N₂O₂S 262.416

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Cryst. Mp 87-89°. [α]_D -42 (MeOH).

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1983, **19**, 660; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 631 (isol, ir, pmr, ms)

Diptocarpaine **D-828**

[6-(Methylsulfinyl)hexyl]urea, 9CI.

Diptocarpaine

[58985-19-6]

H₂NCONH(CH₂)₆SOMe

C₈H₁₈N₂O₂S 206.308

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Mp 124-125°. [α]_D -80.33 (EtOH).

N²-Isopropyl: **Diptocarpamine**. N-(1-Methylethyl)-N'-[6-(methylsulfinyl)hexyl]urea, 9CI. Diptocarpamine

[58985-20-9]

C₁₁H₂₄N₂O₂S 248.389

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Mp 100-101°. [α]_D²⁶ -58.21.

S-Deoxo: [6-(Methylthio)hexyl]urea.

Deoxydiptocarpaine. Deoxydiptocarpaine

[62580-21-6]

C₈H₁₈N₂OS 190.309

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Cryst. (Me₂CO/MeOH). Mp 118-119°.

N²-Isopropyl, S-deoxo: **Deoxydiptocarpamine**. Deoxydiptocarpamine. N-Isopropyl-N'-[6-(methylthio)hexyl]urea

[62784-29-6]

C₁₁H₂₄N₂OS 232.389

Alkaloid from *Diptychocarpus strictus*. Cryst. Mp 108-109°.

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1975, **11**, 762-765; 1976, **12**, 674-675; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 784-786; 1976, **12**, 609-610 (isol, ir, pmr, ms, struct)

Abdilatimov, O. et al., *Khim. Prir. Soedin.*, 1978, **14**, 535-536; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 463-464

(Deoxydiptocarpaine)

Abdilatimov, O. et al., *Khim. Prir. Soedin.*, 1980, **16**, 365-370; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 274-278 (ms)

Tolstikova, O.V. et al., *Khim. Prir. Soedin.*, 1989, **25**, 232-236; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 199-202

(synth)

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1996, **32**, 952-954; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 926-927

(Deoxydiptocarpamine)

Diptocarpidine **D-829**

N,N'-Bis[6-(methylsulfinyl)hexyl]urea, 9CI. Diptocarpidine

[58985-21-0]

MeSO(CH₂)₆NHCONH(CH₂)₆SOMe

C₁₅H₃₂N₂O₃S₂ 352.561

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Mp 135-136°. [α]_D²⁵ -70.5 (MeOH).

S-Deoxy: **Diptocarpiline**. N-[6-(Methylsulfinyl)hexyl]-N'-[6-(methylthio)hexyl]urea, 9CI

[68231-35-6]

C₁₅H₃₂N₂O₂S₂ 336.562

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Mp 95-97°. [α]_D²⁸ -53.2 (MeOH).

S,S'-Dideoxy: N,N'-Bis[6-(methylthio)hexyl]urea, 9CI. N,N'-Di(6-methylthiohexyl)urea

[68231-36-7]

C₁₅H₃₂N₂O₂S₂ 320.563

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Liq.

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1975, **11**, 762; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 784 (isol)

Abdilatimov, O. et al., *Khim. Prir. Soedin.*, 1978, **14**, 223; 1980, **16**, 363; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 183; 1980, **16**, 271 (uv, ir, pmr, ms, struct, deriv)

Diptocarpilidine **D-830**

7-(Methylsulfinyl)heptanenitrile, 9CI.

Diptocarpilidine

[75272-86-5]

[139450-85-4]

NC(CH₂)₆SOMe

C₈H₁₅NOS 173.279

Alkaloid from *Diptychocarpus strictus* (Brassicaceae). Antihypoxic agent. Oil. Bp₄ 193-194°. [α]_D -49 (CHCl₃). λ_{max} 206 (log ε 3.15) (no solvent reported).

S-Deoxo: 7-(Methylthio)heptanenitrile

[75272-78-5]

C₈H₁₅NS 157.279

Constit. of *Alyssum minimum* and *Wasabia japonica* (Japanese horseradish) and other conifers. Antifungal agent.

Aripova, S.F. et al., *Khim. Prir. Soedin.*, 1984, **20**, 84-86; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 79-81 (isol, pmr, ms)

Tolstikov, A.G. et al., *Khim. Prir. Soedin.*, 1991, **27**, 261-263; *Chem. Nat. Compd.*

(*Engl. Transl.*), 1991, **27**, 225-226 (synth, cmr)


Kumagai, H. et al., *Biosci., Biotechnol., Biochem.*, 1994, **58**, 2131-2135 (deriv, isol)

Pedras, M.S.C. et al., *Phytochemistry*, 1998, **49**, 1959-1965 (isol, ir, pmr, cmr, ms)

Di-2-pyridinyl disulfide **D-831**

2,2'-Dithiobispyridine, 9CI. 2,2'-Dipyridyl disulfide. Aldrithiol 2

[2127-03-9]



C₁₀H₈N₂S₂ 220.319

Reagent for peptide coupling and macrocyclic lactonisation of ω-hydroxy acids and selective reagent for detecting thiols. Used as a 1mM soln. in EtOH for photometric detn. of S²⁻. Cryst. (petrol). Sol. H₂O, EtOH, Me₂CO, C₆H₆. Mp 57-58°.

N,N'-Dioxide: **Dipyrrithione**, **BSI**, **INN**, **USAN**. *Omadine disulfide*. *Omadine-DS*. *OMDS*

[3696-28-4]

C₁₀H₈N₂O₂S₂ 252.317

Prod. by *Cortinarius* sp. Antibacterial, antifungal agent. Cryst. (MeOH). Mp 205-206° Mp 235-237°. Log P -2.31 (calc).

► Exp. reprod. and teratogenic effects. UT2965000

N,N'-Dioxide, MgSO₄ complex: **Bispyrrithione magsulfex**, **USAN**. *1,1'-dioxide-O,O',S,S']sulphato(2-)-O]magnesium*, 9CI. *Omadine MDS*

[67182-81-4]

C₁₀H₈MgN₂O₆S₃ 372.686

Antibacterial, antidandruff and antifungal agent.

[43143-11-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 760B; 914D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 283C (nmr)

Marckwald, W. et al., *Ber.*, 1900, **33**, 1556-1566 (synth)

Räth, C. et al., *Annalen*, 1931, **487**, 105-119 (synth)

Canadian Pat., 1954, ((*Dominion Rubber*))501 851; *CA*, **50**, 6740 (deriv)

U.S. Pat., 1956, ((*Olin Mathieson*))2 742 476; *CA*, **50**, 16877c (deriv)

Humphrey, R.E. et al., *Anal. Chem.*, 1971, **43**, 140-142 (detn, S²⁻)

Ger. Pat., 1973, ((*Olin Corp*))2 248 880; *CA*, **79**, 45658 (bispyrrithione magsulfex)

Corey, E.J. et al., *J.A.C.S.*, 1974, **96**, 5614-5616 (use)

Adams, M.D. et al., *Toxicol. Appl. Pharmacol.*, 1976, **36**, 523 (bispyrrithione magsulfex)


Keats, N.G. et al., *J. Het. Chem.*, 1977, **14**, 231-233 (ms)

Grimshaw, C.E. et al., *J.A.C.S.*, 1979, **101**, 1521-1532 (use)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1980, **8**, 214 (use)

- Mitoma, C. *et al.*, *Fundam. Appl. Toxicol.*, 1983, **3**, 256 (*bispyrithione magsulfex, metab*)
- Johnson, D.E. *et al.*, *Fundam. Appl. Toxicol.*, 1984, **4**, 81 (*bispyrithione magsulfex*)
- Kato, E. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 486-495 (*synth*)
- Barton, D.H.R. *et al.*, *Tetrahedron*, 1988, **44**, 7385-7392; 1991, **47**, 6127-6138 (*synth, pmr, cmr, dioxide*)
- Pesticide Manual*, 9th edn., 1991, No. 5325
- Encyclopaedia of Reagents for Organic Synthesis*, (ed. Paquette, L.A.), Wiley, 1995, **4**, 2277-2280 (*use*)
- Stoyanov, S. *et al.*, *J. Het. Chem.*, 1996, **33**, 927-931 (*uv*)
- Martindale, The Extra Pharmacopoeia*, 31st edn., Pharmaceutical Press, 1996, 1085
- Bodige, S.G. *et al.*, *Chem. Comm.*, 1997, 1669-1670 (*dioxide, cryst struct*)
- Nicholas, G.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 341-344 (*dioxide, isol*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DXN300

3,5-Di-2-pyrrolidinylpyridine D-832



$C_{13}H_{19}N_3$ 217.313

(S,S)-(?) -form


N,N'-Di-Me: 3,5-Bis(1-methyl-2-pyrrolidinyl)pyridine

$C_{15}H_{23}N_3$ 245.367

Alkaloid from the roots of *Nicotiana tabacum*.

Wei, X. *et al.*, *Life Sci.*, 2005, **78**, 495-505 (*isol, synth, pmr, cmr, ms*)

5H,10H-Dipyrrolo[1,2-a:1',2'-d]pyrazine-5,10-dione, 9CI Pyrocoll [484-73-1]



$C_{10}H_6N_2O_2$ 186.17

Obt. from gelatine. Prod. by the alkaliphilic *Streptomyces* sp. AK 409. Exhibits various antimicrobial activities. Plates (AcOH), cryst. (C_6H_6). Mp 272-273° (sealed tube). Gives 1H-Pyrrole-2-carboxylic acid, P-944 with KOH.

Ciamician, G. *et al.*, *Monatsh. Chem.*, 1880, **1**, 279; *Ber.*, 1884, **17**, 105

Hale, W.J. *et al.*, *J.A.C.S.*, 1916, **38**, 1065


Mold, J.D. *et al.*, *Tob. Int.*, 1960, **4**, 130; *CA*, **54**, 25597

Johnson, A.W. *et al.*, *J.C.S. Perkin I*, 1972, 2681

Boatman, R.J. *et al.*, *J.O.C.*, 1976, **41**, 3050

Dieter, A. *et al.*, *J. Antibiot.*, 2003, **56**, 639-646 (*isol, synth, pmr, cmr, ms, activity*)


Discarene C D-834 [357641-18-0]



$C_{29}H_{35}N_3O_4$ 489.613

Not to be confused with Discarine C, D-837. Only relative config. at C-3/C-4 was determined. Alkaloid from the bark of *Discaria americana*. Powder. Mp 297° dec. $[\alpha]_D^{20}$ -51.7 (c, 0.2 in MeOH/ $CHCl_3$). Giacomelli, S.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 997-999 (*isol, ir, pmr, cmr, struct*)

Discarene D D-835 [357641-20-4]




$C_{32}H_{33}N_3O_4$ 523.63

Not to be confused with Discarine D, D-838. Only rel. config. at C-3/C-4 was determined. C-7-Config. incorrectly given as *R*- in a 2006 review. Alkaloid from the bark of *Discaria americana*. Amorph. powder ($CHCl_3$). $[\alpha]_D^{25}$ -176 (c, 0.2 in MeOH/ $CHCl_3$).

Giacomelli, S.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 997-999 (*isol, pmr, cmr, struct*)

Giacomelli, S.R. *et al.*, *Phytochemistry*, 2004, **95**, 933-937 (*occur*)

Discarine B D-836 [36211-11-7]



Absolute Configuration

$C_{33}H_{43}N_5O_4$ 573.734

Alkaloid from *Discaria longispina*, *Discaria americana*, *Discaria febrifuga*, *Ceanothus sanguineus*, and *Ceanothus*

integerrimus (Rhamnaceae). Shows anti-bacterial props. Cryst. ($CHCl_3/Et_2O$). Mp 235-236°. $[\alpha]_D$ -172 (c, ca. 0.1 in $CHCl_3$).

N-De-Me: Discarine I. N-Demethyldiscarine B

[105708-60-9]

$C_{32}H_{41}N_5O_4$ 559.707

Alkaloid from the root bark of *Discaria febrifuga* (Rhamnaceae). Mp 140°. $[\alpha]_D^{25}$ -149 (c, 0.1 in MeOH). May be identical with Homoamericine, in Texensine, T-300. λ_{max} 227; 270; 278; 289 (MeOH).

10,11-Dihydro:

Cryst. (MeOH aq.). Mp 263-263.5°.

10,11-Dihydro, 11-hydroxy: Discarine K [109175-41-9]

$C_{33}H_{45}N_5O_5$ 591.749

Alkaloid from the roots of *Discaria febrifuga* (Rhamnaceae). Mp 237°. $[\alpha]_D^{20}$ -62 (MeOH). No stereochem. determined.

González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915-916 (*stereochem, pmr*)

Mascaretti, O.A. *et al.*, *Phytochemistry*, 1972, **11**, 1133-1137 (*isol, uv, ir, pmr, ms, struct*)

Chang, C.-J. *et al.*, *Phytochemistry*, 1974, **13**, 1273-1277 (*pmr*)

Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 220-227; 663-668 (*isol, ms*)


Pais, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869-1872 (*cmr*)

Hennig, P. *et al.*, *Z. Naturforsch., B*, 1986, **41**, 1180-1185 (*Discarine I*)

Voelter, W. *et al.*, *Z. Naturforsch., B*, 1987, **42**, 467 (*Discarine K*)

Giacomelli, S.R. *et al.*, *Phytochemistry*, 2004, **65**, 933-937 (*occur*)

Discarine C D-837 [88607-21-0]



$C_{31}H_{42}N_4O_4$ 534.697

No stereochem. determined. Alkaloid from *Discaria americana* and *Discaria febrifuga* (Rhamnaceae). Phys. constants not recorded.

Digel, M. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1983, **364**, 1641-1643 (*uv, ir, pmr, ms, struct*)

Giacomelli, S.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 997-999 (*occur*)

Giacomelli, S.R. *et al.*, *Phytochemistry*, 2004, **65**, 933-937 (*occur*)

Discarine D† D-838 [88644-36-4]

As Discarine C, D-837 with

R = -CH₂PhC₃₄H₄₀N₄O₄ 568.714

Stereochem. not determined. Alkaloid from the root bark of *Discaria americana* and *Discaria febrifuga* (Rhamnaceae). Mp 212°. [α]_D²⁰ -148 (c, 0.1 in CHCl₃).

Stereoisomer: **Crenatine A**[†]

[52801-20-4]

C₃₄H₄₀N₄O₄ 568.714


Alkaloid from *Discaria crenata* (Rhamnaceae). Mp 223°. [α]_D -292.58 (CHCl₃).

Silva, M. *et al.*, *Phytochemistry*, 1974, **13**, 861-863 (*Crenatine A*)

Digel, M. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1983, **364**, 1641-1643 (*Discarine D*)
Giacomelli, S.R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 997-999 (*occur*)

Discarine E


[98263-18-4]

D-839C₂₈H₄₄N₄O₄ 500.68

Alkaloid from the bark of *Discaria febrifuga* and from *Discaria longispina* (Rhamnaceae). Mp 270-273°. [α]_D²⁵ +236 (c, 0.5 in AcOH).

Morel, A. *et al.*, *Chimia*, 1985, **39**, 98 (*isol, ms, pmr, uv, struct*)Machado, E.C. *et al.*, *J. Nat. Prod.*, 1995, **58**, 548 (*isol, ir, pmr, cmr, ms*)**Discarine M**

[714249-70-4]

D-840

Absolute Configuration

C₂₆H₃₇N₅O₄ 455.596

Only rel. config. of C-3/C-4 was determined. Alkaloid from the bark of *Discaria americana*. Amorph. powder. [α]_D²⁰ -176.7 (c, 0.2 in MeOH/CHCl₃).

N-Deacyl, N-(*E*-cinnamoyl): **Sanjoine-nine**

[107446-80-0]


C₂₉H₃₅N₃O₄ 489.613

Alkaloid from the seeds of *Zizyphus vulgaris* var. *spinousus* and from *Zizy-*

phus lotus. Needles (CHCl₃/MeOH). Mp 281-282°. [α]_D -272.5 (c, 1.6 in Py). Gross. struct. only determined, may not be the same stereochem. as that of Discarine M. The *Z. lotus* isolate reported to be racemic. λ_{\max} 224 (log ϵ 4.38); 281 (log ϵ 4.39) (MeOH).

Han, B.H. *et al.*, *Phytochemistry*, 1990, **29**, 3315-3319 (*Sanjoineine*)Abu-Zarga, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 504-511 (*(±)-Sanjoineine*)Giacomelli, S.R. *et al.*, *Phytochemistry*, 2004, **65**, 933-937 (*Discarine M*)**Discarine N**

[714249-71-5]

D-841

Absolute Configuration

C₃₂H₃₃N₃O₅ 539.63

Only rel. config. of C-3/C-4 was determined. Alkaloid from the bark of *Discaria americana*. Powder. Mp 233-235°. [α]_D²⁰ +98.1 (c, 0.09 in MeOH/CHCl₃).

Stereoisomer(?): **Scutianene C**. *Scutianene D*


[55857-24-4]

C₃₂H₃₃N₃O₅ 539.63

Alkaloid from *Scutia buxifolia*. Cryst. (MeOH). Mp 232-234°. [α]_D +203 (c, 0.1 in CHCl₃/MeOH). No stereochem. determined. Not to be confused with Scutianine C, S-166. λ_{\max} 207 (log ϵ 4.43); 218 (log ϵ 4.39); 275 (log ϵ 4.32) (no solvent reported).

Gonzalez, S. *et al.*, *Phytochemistry*, 1974, **13**, 2865-2869 (*Scutianene C*)Giacomelli, S.R. *et al.*, *Phytochemistry*, 2004, **65**, 933-937 (*Discarine N*)Morel, A.F. *et al.*, *Phytochemistry*, 2005, **66**, 2571-2576 (*Scutianene D*)**Discobahamin A**

[155547-93-6]

D-842

R = H

C₄₇H₆₅N₉O₁₁ 932.084

Cyclic peptide antibiotic. Constit. of the marine sponge *Discodermia* sp. Antifungal agent. Pale yellow gum. Sol. MeOH. [α]_D²⁴ -29 (c, 0.5 in MeOH). λ_{\max} 203 (ϵ

1400); 220 (ϵ 10200); 267 (ϵ 6800); 305 (ϵ 1500) (MeOH) (Berdy).

Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 79 (*isol, pmr, cmr*)**Discobahamin B****D-843**

[155547-94-7]


As Discobahamin A, D-842 with

R = -CH₃C₄₈H₆₇N₉O₁₁ 946.111

Cyclic peptide antibiotic. Constit. of the marine sponge *Discodermia* sp. Antifungal agent. Pale yellow gum. Sol. MeOH. [α]_D²⁴ -31 (c, 0.1 in MeOH). λ_{\max} 203 (ϵ 1400); 220 (ϵ 10200); 267 (ϵ 6800); 305 (ϵ 1500) (MeOH) (Berdy).

Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 79 (*isol, pmr, cmr*)**Discodermide**

[134458-00-7]

D-844


Relative Configuration

C₂₇H₃₄N₂O₆ 482.575

Tetramic acid deriv. Isol. from the sponge *Discodermia dissoluta*. Antifungal and antitumour agent. Mp ca.° 200 dec. [α]_D +97.5 (c, 0.2 in CHCl₃/MeOH). λ_{\max} 238 (ϵ 16500); 313 (ϵ 9650) (MeOH).

Gunasekera, S.P. *et al.*, *J.O.C.*, 1991, **56**, 4830-4833 (*isol, uv, pmr, cmr*)**Discodermindole**

[133523-28-1]

D-845C₁₁H₁₀Br₂N₄ 358.035

Alkaloid from the sponge *Discodermia polydiscus*. Cytotoxic. Viscous oil. [α]_D²⁰ -27 (c, 1 in MeOH). λ_{\max} 224 (ϵ 34000); 282 (ϵ 5700); 292 (ϵ 6100); 300 (ϵ 5000) (MeOH) (Derep).


6-Hydroxy: **6-Hydroxydiscodermindole**C₁₁H₁₀Br₂N₄O 374.034

Isol. from *Discodermia polydiscus*. [α]_D²¹ -41.6 (c, 0.1 in MeOH). λ_{\max} 215 (log ϵ 4.34); 260 (log ϵ 3.94); 310 (log ϵ 4.79) (MeOH).

Sun, H.H. *et al.*, *J.O.C.*, 1991, **56**, 4307-4308 (*isol, uv, ir, pmr, cmr, ms*)Cohen, J. *et al.*, *Pharm. Biol.*, 2004, **42**, 59-61 (*6-Hydroxydiscodermindole*)

Discogouattine**D-846**

6,7,7a,8-Tetrahydro-10,12-dimethoxy-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline, 9Cl. 9,11-Dimethoxy-1,2-methylenedioxyynoraporphine. O-Methylcalycinine

C₁₉H₁₉NO₄ 325.363**(R)-form** [89368-38-7]

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Noncryst. Negative opt. rotn. in MeOH.

O⁹-De-Me: **Isocalycinine**. 9-Hydroxy-11-methoxy-1,2-methylenedioxyynoraporphine

[91174-09-3]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Beige cryst. (MeOH). Mp 219°. [α]_D -147 (c, 0.53 in Py).

O¹¹-De-Me: **Calycinine**. 6,7,7a,8-Tetrahydro-10-methoxy-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-12-ol, 9Cl. 11-Hydroxy-9-methoxy-1,2-methylenedioxyynoraporphine. Fissoldine

[70420-58-5]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the leaves of *Duguetia calycina* and the leaves and stem bark of *Duguetia obovata*. Also isol. from *Fissistigma oldhamii* (Annonaceae). Cryst. (Et₂O). Mp 156°. [α]_D -145 (c, 0.53 in CHCl₃).

O¹¹-De-Me; hydrobromide:

Light-greyish needles (EtOH). Mp 276-277°. [α]_D²⁵ -99.1 (c, 0.45 in EtOH).

O¹¹-De-Me; N,O-Di-Ac:

Noncryst. [α]_D -313 (c, 0.55 in CHCl₃).

O¹¹-De-Me, N-Me: **N-Methylcalycinine**. 11-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine

[86537-66-8]

C₁₉H₁₉NO₄ 325.363

Trace alkaloid from the stem bark of *Duguetia obovata* (Annonaceae). Noncryst. Negative opt. rotn. in CHCl₃.

O¹¹-De-Me, N-Me, 6-hydroxy: see Spixianine, S-469

(S)-form

O¹¹-De-Me: **Fissistigine A**

[83685-17-0]

C₁₈H₁₇NO₄ 311.337

Alkaloid from *Fissistigma oldhamii* (Annonaceae).

Lu, S.-T. et al., *Heterocycles*, 1983, **20**, 813 (uv, ir, pmr, ms, struct, Calycinine)


Roblot, F. et al., *J. Nat. Prod.*, 1983, **46**, 862 (uv, ir, pmr, cmr, ms, struct, Discogouattine, Calycinine)

Xu, C. et al., *Zhongcaoyao*, 1983, **14**, 148; *C.A.* **99**, 19659n (isol, uv, ir, pmr, ms, struct, Fissistigine A)

Hocquemiller, R. et al., *J. Nat. Prod.*, 1984, **47**, 353 (isol, uv, ir, pmr, ms, struct, Isocalycinine)

Discokiolide A**D-847**

[141266-02-6]

C₅₃H₇₀N₈O₁₃ 1027.182

Depsipeptide antibiotic. Isol. from marine sponge *Discodermia kiiensis*. Cytotoxic agent. [α]_D²⁵ -52.3 (c, 1 in MeOH) (as Me ester). λ_{max} 251 (ε 19500) (MeOH).

Tada, H. et al., *Chem. Lett.*, 1992, 431-434 (isol, struct)

Discokiolide B**D-848**

[141266-03-7]

As Discokiolide A, D-847 with

R = -CH(CH₃)CH₂CH₃C₅₄H₇₂N₈O₁₃ 1041.209

Depsipeptide antibiotic. Isol. from the marine sponge *Discodermia kiiensis*. Cytotoxic. [α]_D²⁵ -43.5 (c, 1 in MeOH) (as Me ester). λ_{max} 251 (ε 19500) (MeOH) (Derep).

A⁷-Isomer: **Discokiolide D**

[141266-05-9]

C₅₄H₇₂N₈O₁₃ 1041.209

Isol. from *Discodermia kiiensis*. Cytotoxic. [α]_D²⁵ -49.5 (c, 0.6 in MeOH) (as Me ester). λ_{max} 258 (ε 15900) (MeOH) (Derep).

Demethoxy: **Discokiolide C**

[141266-04-8]


C₅₃H₇₀N₈O₁₂ 1011.182

Isol. from *Discodermia kiiensis*. Cytotoxic. [α]_D²⁵ -28.7 (c, 0.6 in MeOH) (as Me ester). λ_{max} 252 (ε 18200) (MeOH) (Derep).

Tada, H. et al., *Chem. Lett.*, 1992, 431 (isol, struct)

Discorhabdin V**D-849**

[721395-09-1]

C₁₈H₁₇BrN₃O₂⁺ 387.255Alkaloid from the sponge *Tsitsikamma*

pedunculata. Dark green solid (as trifluoroacetate salt).

14-Bromo, 1-hydroxy: **14-Bromo-1-hydroxydiscorhabdin V**

[721395-38-6]


C₁₈H₁₆Br₂N₃O₃⁺ 482.151

Alkaloid from *Tsitsikamma flavus* and *Tsitsikamma pedunculata*. Brown solid (as trifluoroacetate salt).

Antunes, E.M. et al., *J. Nat. Prod.*, 2004, **67**, 1268-1276 (isol, cd, pmr, cmr)

Discorhabdin A**D-850****Prianosin A**

[112515-41-0]

C₁₈H₁₄BrN₃O₂S 416.297

This group of alkaloids exist in nature in two enantiomeric forms, depending upon the location of their natural source. Closely related to Discorhabdin C, D-851. Alkaloid from the marine sponge *Prianos melanos* and *Latrunculia brevis*. Exhibits potent antineoplastic activity. Green solid (as hydrochloride). Mp 300°. [α]_D²⁴ +248 (c, 0.19 in CHCl₃). [α]_D +400 (c, 0.05 in MeOH) (hydrochloride). λ_{max} 335 (ε 14000); 473 (ε 1000) (MeOH/KOH) (Derep). λ_{max} 248 (ε 29500); 353 (ε 11000); 430 (sh); 567 (ε 900) (MeOH) (Derep).

4,5-Didehydro: **Discorhabdin B**

[115439-61-7]

C₁₈H₁₂BrN₃O₂S 414.282

Isol. from a *Latrunculia* sp. Cytotoxic agent. Green solid (hydrochloride). Sol. DMSO; fairly sol. MeOH, H₂O. Mp > 360° (hydrochloride). [α]_D +400 (c, 0.2 in MeOH) (hydrochloride). λ_{max} 248 (ε 30600); 309 (ε 10800); 357 (ε 10600); 567 (ε 1100) (MeOH). λ_{max} 231 (ε 22500); 306 (ε 14700) (MeOH/NaOH) (Berdy).

16,17-Didehydro: **Prianosin B**

[116302-35-3]

C₁₈H₁₂BrN₃O₂S 414.282

Alkaloid from *Prianos melanos*. Cytotoxic agent. Red cryst. Sol. EtOAc, MeOH, CHCl₃; poorly sol. H₂O. Mp 250-251° dec. [α]_D³⁰ +360 (c, 0.1 in CHCl₃). λ_{max} 228 (ε 17800); 263 (ε 15000); 410 (sh); 430 (ε 11200) (MeOH) (Derep).

4,5,16,17-Tetrahydro: **Discorhabdin Q**

[224447-29-4]

C₁₈H₁₀BrN₃O₂S 412.266

Alkaloid from the sponges *Latrunculia purpurea* and *Zyzzya* spp. Orange solid. [α]_D -452.4 (c, 0.004 in CHCl₃). [α]_D -904 (c, 0.01 in MeOH). λ_{max} 222 (ε 28200); 428 (ε 9900) (MeOH).

Debromo, 4,5-didehydro: Discorhabdin I
[663597-85-1]
C₁₈H₁₃N₃O₂S 335.386
Formerly given synonym Discorhabdin G which confuses it with Discorhabdin G in Discorhabdin C, D-851. Isol. from *Latrunculia* sp. Cytotoxic. Green solid (as trifluoroacetate salt). [α]_D²⁰ +540 (c, 0.05 in MeOH) (as trifluoroacetate salt).


Debromo, 4,5-didehydro, 1ξ,2ξ-epoxide: Discorhabdin R
[308355-78-4]
C₁₈H₁₃N₃O₃S 351.385
Alkaloid from the sponges *Latrunculia* sp. and *Negombata* sp. Green solid. [α]_D²⁰ +161 (c, 0.1 in MeOH). λ_{max} 202 (ε 7600); 255 (ε 9000); 262 (ε 9200); 287 (ε 5400); 326 (ε 3500); 368 (ε 3800); 567 (ε 600) (MeOH).

Kobayashi, J. *et al.*, *Tet. Lett.*, 1987, **28**, 4939-4942 (*uv, ir, pmr, cmr, cryst struct, abs config*)
Cheng, J. *et al.*, *J.O.C.*, 1988, **53**, 4621-4624 (*Prianosins*)
Perry, N.B. *et al.*, *Tetrahedron*, 1988, **44**, 1727-1734 (*Discorhabdin B*)
Dijoux, M.-G. *et al.*, *J. Nat. Prod.*, 1999, **62**, 636-637 (*Discorhabdin Q*)
Ford, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1527-1528 (*Discorhabdin R*)
Tohma, H. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 348-350 (*Discorhabdin A, synth*)
Tohma, H. *et al.*, *J.A.C.S.*, 2003, **125**, 11235-11240 (*Discorhabdin A, synth*)
Reyes, F. *et al.*, *J. Nat. Prod.*, 2004, **67**, 463-465 (*Discorhabdin I*)
Harayama, Y. *et al.*, *Curr. Org. Chem.*, 2005, **9**, 1567-1588 (*rev, synth*)
Antunes, E.M. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 62-72 (*rev*)
Grkovic, T. *et al.*, *J.O.C.*, 2008, **73**, 9133-9136 (*isol, cd, pmr, cmr, abs config*)
Wada, Y. *et al.*, *Tetrahedron*, 2009, **65**, 1059-1062 (*Prianosin B, synth*)

Discorhabdin C

D-851

[105372-81-4]



C₁₈H₁₃Br₂N₃O₂ 463.128
Pigment from a sponge of the genus *Latrunculia*. Sol. DMSO; fairly sol. MeOH. Mp 360° (as hydrochloride). λ_{max} 245 (ε 28500); 351 (ε 10000); 545 (ε 500) (MeOH/HCl) (Derep). λ_{max} 337 (ε 13000); 481 (ε 1500) (MeOH/KOH) (Derep). λ_{max} 245 (ε 28500); 250 (ε 14600); 351 (ε 10000); 355 (ε 7700); 545 (ε 500) (MeOH) (Berdy).

► Highly cytotoxic; LD₅₀ (mus, ipr) 2 mg/kg.

13-N-Me: Discorhabdin P

[219566-63-9]
C₁₉H₁₅Br₂N₃O₂ 477.154
Isol. from a sponge of the *Batzella* sp.

Cytotoxic agent. Mp >° 360 (blackens at 162°). λ_{max} 200 (log ε 4.26); 245 (log ε 4.28); 337 (log ε 4); 488 (log ε 3.49) (MeOH).

3-Alcohol: 3-Dihydrodiscorhabdin C

C₁₈H₁₅Br₂N₃O₂ 465.143
Isol. from *Tsitsikamma pedunculata*. Red solid (as trifluoroacetate). λ_{max} 239 (ε 23700); 351 (ε 6030); 551 (ε 740) (MeOH) (trifluoroacetate).

7,8-Didehydro, 3-alcohol: 7,8-Didehydro-3-dihydrodiscorhabdin C. 3-Dihydro-7,8-didehydrodiscorhabdin C

[721395-08-0]
C₁₈H₁₃Br₂N₃O₂ 463.128
Isol. from the sponges *Tsitsikamma flavus* and *Tsitsikamma pedunculata*. Olive-green solid.

14-Bromo: 14-Bromodiscorhabdin C

[182928-53-6]
C₁₈H₁₂Br₃N₃O₂ 542.024
Isol. from the South African latrunculiid sponge *Tsitsikamma pedunculata* and another sponge. Exhibits antimicrobial activity. λ_{max} 242; 317; 374 (MeOH) (Berdy).

14-Bromo, 3-alcohol: 14-Bromo-3-dihydrodiscorhabdin C

[182928-55-8]
C₁₈H₁₄Br₃N₃O₂ 544.04
Isol. from *Tsitsikamma flavus*, *Tsitsikamma pedunculata* and another sponge. Exhibits antimicrobial activity. λ_{max} 242; 317; 374 (MeOH) (Berdy).

14-Bromo, 7,8-didehydro, 3-alcohol: 14-Bromo-7,8-didehydro-3-dihydrodiscorhabdin C. 14-Bromo-3-dihydro-7,8-didehydrodiscorhabdin C

[721395-37-5]
C₁₈H₁₂Br₃N₃O₂ 542.024
Isol. from *Tsitsikamma pedunculata*. Green solid.

2-Debromo: Discorhabdin E

[159308-95-9]
C₁₈H₁₄BrN₃O₂ 384.232
Alkaloid from the New Zealand sponge *Latrunculia* cf. *bocagei*. Red solid (as trifluoroacetate). λ_{max} 201 (ε 14500); 244 (ε 18000); 360 (ε 7300); 551 (ε 1000) (MeOH) (Berdy).

2-Debromo, 7,8-didehydro, 1,2-dihydro: Discorhabdin G†

[172961-01-2]
C₁₈H₁₄BrN₃O₂ 384.232
Not to be confused with Discorhabdin G in Discorhabdin A, D-850. Alkaloid from the Antarctic sponge *Latrunculia apicalis*. Antimicrobial agent. Green pigment (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ +27 (c, 0.063 in MeOH). λ_{max} 210 (ε 7500); 250 (ε 8100); 322 (ε 3200); 402 (ε 2300); 610 (ε 600) (MeOH).

Perry, N.B. *et al.*, *J.O.C.*, 1986, **51**, 5476-5478 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)
Nishiyama, S. *et al.*, *Tet. Lett.*, 1991, **32**, 4151 (*synth*)

Kita, Y. *et al.*, *J.A.C.S.*, 1992, **114**, 2175 (*synth*)
Copp, B.R. *et al.*, *J.O.C.*, 1994, **59**, 8233-8238 (*Discorhabdin E*)

Tao, X.L. *et al.*, *Tetrahedron*, 1994, **50**, 2017 (*synth*)

Yang, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1596-1599 (*Discorhabdin G*)

Sadanandan, E.V. *et al.*, *J.O.C.*, 1995, **60**, 1800 (*synth*)

Hooper, G.J. *et al.*, *Tet. Lett.*, 1996, **37**, 7135 (*14-Bromodiscorhabdin C, 14-Bromodihydrodiscorhabdin C*)

Roberts, D. *et al.*, *J.O.C.*, 1997, **62**, 568 (*synth*)
Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1999, **62**, 173-175 (*Discorhabdin P*)

Aubart, K.M. *et al.*, *J.O.C.*, 1999, **64**, 16-22 (*synth*)


Antunes, E.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1268-1276 (*14-Bromo-3-dihydro-7,8-didehydrodiscorhabdin C, 3-Dihydro-7,8-didehydrodiscorhabdin C*)

Harayama, Y. *et al.*, *Curr. Org. Chem.*, 2005, **9**, 1567-1588 (*rev, synth*)

Discorhabdin D

D-852

Prianosin D
[115384-95-7]



C₁₈H₁₄N₃O₂S⁺ 336.393
Alkaloid from the New Zealand sponge *Latrunculia brevis* and from the Okinawan marine sponge *Prianos melanos*. Exhibits significant *in vivo* antitumor and antineoplastic activity and antimicrobial action. Also induces Ca²⁺ release from sarcoplasmic reticulum. Green solid. Mp 300°. [α]_D²⁰ +344 (c, 0.01 in MeOH). Unstable in soln. λ_{max} 250 (ε 18100); 284 (ε 11100); 325 (ε 6600); 392 (ε 6950) (MeOH) (Derep). λ_{max} 262 (ε 30900); 290 (ε 15500); 368 (ε 9550) (MeOH/KOH) (Derep). λ_{max} 248 (ε 22400); 281 (ε 14100); 320 (ε 8510); 395 (ε 8910); 584 (ε 692) (MeOH) (Derep).

iR-Hydroxy: Discorhabdin L

[663597-84-0]
C₁₈H₁₄N₃O₃S⁺ 352.393
Alkaloid from *Latrunculia brevis*. Cytotoxic. Green solid (as TFA salt). Mp >250°. Blackens at 160° (TFA salt). Optically inactive at sodium D-line wavelength. λ_{max} 268 (log ε 3.75); 310 (sh); 352 (log ε 3.59); 550 (log ε 2.65) (MeOH) (TFA salt).

1ξ-Amino: 1-Aminodiscorhabdin D

[721395-40-0]
C₁₈H₁₅N₄O₂S⁺ 351.408
Isol. from *Latrunculia bellae*. Dark green solid.

1ξ-(Carboxymethylamino): Discorhabdin N

[721395-10-4]
C₂₀H₁₇N₄O₄S⁺ 409.445
Isol. from *Latrunculia bellae*. Red-brown solid.


1ξ-Methoxy: 1-Methoxydiscorhabdin D

[721395-39-7]
C₁₉H₁₆N₃O₃S⁺ 366.42

Isol. from *Latrunculia bellae*. Brown solid.


- Perry, N.B. *et al.*, *J.O.C.*, 1988, **53**, 4127-4128 (isol, uv, ir, pmr, cmr)
 Cheng, J.F. *et al.*, *J.O.C.*, 1988, **53**, 4621-4624 (isol, uv, ir, pmr, cmr, ms, cd)
 Kobayashi, J. *et al.*, *Tet. Lett.*, 1991, **32**, 1227-1228 (struct)
 Reyes, F. *et al.*, *J. Nat. Prod.*, 2004, **67**, 463-465 (Discorhabdin L)
 Antunes, E.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1268-1276 (Discorhabdin N, 1-Aminodiscorhabdin D, 1-Methoxydiscorhabdin D)
 Antunes, E.M. *et al.*, *Nat. Prod. Rep.*, 2005, **22**, 62-72 (rev)

Discorhabdin H D-853
 [721395-11-5]



C₂₅H₂₃N₆O₄S[⊕] 535.627
 Alkaloid from the sponge *Strongyloides algoensis*. Dark green solid.
 Antunes, E.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1268-1276 (isol, pmr, cmr)

Discorhabdin T D-854
 [649761-80-8]



C₂₀H₁₄BrN₃O₂S 440.32
 Isol. from the sponge *Batzella* sp. Cytotoxic. Dark orange solid. λ_{max} 225 (log ε 4.11); 245 (log ε 3.97); 305 (log ε 3.65); 412 (log ε 3.55); 432 (log ε 3.65) (MeOH).


7,8-Dihydro: Discorhabdin S
 [649761-79-5]

C₂₀H₁₆BrN₃O₂S 442.335
 Isol. from a *Batzella* sp. Cytotoxic. Dark orange solid. λ_{max} 225 (log ε 4.54); 260 (log ε 4.27); 305 (log ε 4.04); 400 (log ε 3.94); 422 (log ε 4.14) (MeOH).

16,17-Dihydro: Discorhabdin U
 [649761-81-9]
 [1078709-79-1 (±)-form]


C₂₀H₁₆BrN₃O₂S 442.335
 Isol. from a *Batzella* sp. Cytotoxic. Dark orange solid. λ_{max} 205 (log ε 4.42); 242 (log ε 4.22); 287 (log ε 4.01); 340 (log ε 3.95); 425 (log ε 3.12) (MeOH).
 Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1615-1617 (isol, pmr, cmr)

Discorhabdin W D-855
 [872199-41-2]



C₃₆H₂₂Br₂N₆O₄S₂ 826.548
 Isol. from a New Zealand *Latrunculia* sp. [α]_D²⁰ +220 (c, 0.05 in MeOH). λ_{max} 203; 241; 301; 444 (TFA/MeCN aq.).
 Lang, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1796-1798 (isol, pmr, cmr)

Discretamine D-856
Aequaline




C₁₉H₂₁NO₄ 327.379
 (S)-form [1356-73-6]

Alkaloid from *Xylopia discreta*, *Xylopia buxifolia*, *Desmos tiebaghiensis*, *Duguetia calycina*, *Schefferomitra subaequalis* and *Mitrella kentii* (Annonaceae). Cryst. (CHCl₃/MeOH). Mp 221-224° dec. (232° dec.). [α]_D²³ -368 (c, 1.01 in Py). λ_{max} 212 (log ε 4.5); 225 (sh) (log ε 4.1); 284 (log ε 3.79) (EtOH).
Di-Me ether: see Tetrahydropalmatine, T-212

(±)-form [55934-50-4]

Synthetic. Mp 208-212° dec.
 Schmutz, J. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 335-343 (isol, uv)
 Gellert, E. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2477-2482 (isol, uv, ir, pmr, ms, struct)
 Ellis, J. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2735-2736 (isol)
 Richter, W.J. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 209-211 (ms, struct)
 Chiang, H.-C. *et al.*, *J.O.C.*, 1977, **42**, 3190-3194 (synth, pmr, ms)
 Brochmann-Hanssen, E. *et al.*, *J.O.C.*, 1977, **42**, 3588-3591 (struct)
 Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551-556 (occur)
 Leboeuf, M. *et al.*, *J. Nat. Prod.*, 1982, **45**, 617-623 (isol, uv, pmr, ms)
 Kaewamatawong, R. *et al.*, *J. Nat. Med. (Tokyo)*, 2007, **61**, 349-350 (isol, pmr, cmr)


Disorazole C₁ D-857
 [158181-52-3]



C₄₄H₅₈N₂O₁₀ 774.95
 Prod. by *Sorangium cellulosum*. Highly cytotoxic and active against fungi. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. [α]_D²² -124.8 (c, 0.6 in MeOH). λ_{max} 272; 351 (MeOH) (Berdy).

Jansen, R. *et al.*, *Annalen*, 1994, 759 (isol, uv, pmr, cmr)
 Hillier, M.C. *et al.*, *Tet. Lett.*, 2000, **41**, 2821-2824 (synth)
 Wipf, P. *et al.*, *J.A.C.S.*, 2004, **126**, 15346-15347 (synth)


Disorazole C₂ D-858
 [158181-53-4]



C₄₄H₅₈N₂O₁₀ 774.95
 Isol. from bacterium *Sorangium cellulosum*. Cytotoxic, fungicidal. [α]_D²² -140 (c, 0.1 in MeOH).

Jansen, R. *et al.*, *Annalen*, 1994, 759 (isol, uv, pmr, cmr, struct)

Disorazole F₁ D-859



C₄₃H₅₄N₂O₉ 742.908
 Gross stereochem. only publ. so far for most members of this series. See also Disorazole C₁, D-857 which is the member of the series for which stereochem. is best known. Prod. by *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. [α]_D²² +16 (c, 0.1 in MeOH). λ_{max} 254; 264 (sh); 271 (ε 49500); 282; 320; 339 (sh) (MeOH).

O-De-Me: Disorazole F₂

C₄₂H₅₂N₂O₉ 728.881
 From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly

sol. H₂O, hexane. $[\alpha]_D^{22}$ -66.2 (c, 0.7 in MeOH). λ_{\max} 263 (sh); 271 (ε 43650); 280; 338 (sh); 351 (ε 41000); 366 (sh) (MeOH).

9R,10R-Epoxyde: **Disorazole A₇**

C₄₃H₅₄N₂O₁₀ 758.907

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -211 (c, 0.4 in MeOH). λ_{\max} 252; 266 (sh); 274 (ε 40830); 281; 298; 307; 311 (sh) (MeOH).

9R,10S-Epoxyde: **Disorazole A₁**

C₄₃H₅₄N₂O₁₀ 758.907

Major metab. from the bacterium *Sorangium cellulosum*. Cytotoxic and fungicidal agent. Mp 172°. $[\alpha]_D^{22}$ -77 (c, 0.75 in MeOH). Comps. of the Disorazole complex are inactive against yeasts and bacteria. λ_{\max} 264 (sh); 272; 278 (sh); 305 (sh) (MeOH) (Derep). λ_{\max} 251; 265 (sh); 272 (ε 55600); 298; 306; 321 (sh) (MeOH).

9R,10S-Epoxyde, O-de-Me: **Disorazole A₂**

C₄₂H₅₂N₂O₁₀ 744.88

From *Sorangium cellulosum*. Cytotoxic, fungicidal. $[\alpha]_D^{22}$ -95.3 (c, 0.8 in MeOH). λ_{\max} 252; 264 (sh); 272 (ε 51400); 281; 298; 306; 321 (sh) (MeOH).

9R,10R:9'R,10'S-Diepoxyde: **Disorazole E₂**

C₄₃H₅₄N₂O₁₁ 774.906

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -220 (c, 0.5 in MeOH). λ_{\max} 277 (sh); 288 (ε 27200); 297; 308; 312 (sh) (MeOH).

9R,10S:9'R,10'S-Diepoxyde: **Disorazole E₁**

C₄₃H₅₄N₂O₁₁ 774.906

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 194°. $[\alpha]_D^{22}$ +51.6 (c, 0.5 in MeOH). λ_{\max} 291 (sh); 297 (ε 29000); 306; 321 (sh) (MeOH).

9R,10S:11R,12S-Diepoxyde: **Disorazole H**

C₄₃H₅₄N₂O₁₁ 774.906

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -64.4 (c, 0.3 in MeOH). λ_{\max} 265 (sh); 273 (ε 46450); 282; 306; 321 (sh) (MeOH).

9R,10S-Epoxyde, 16'-ketone: **Disorazole I**

C₄₃H₅₂N₂O₁₀ 756.891

From *Sorangium cellulosum*. Cytotoxic, fungicidal. $[\alpha]_D^{22}$ -95.8 (c, 0.5 in MeOH). λ_{\max} 254; 264 (sh); 272 (ε 51400); 281; 298; 306; 321 (sh) (MeOH).

9,10-Dihydroxy, 9,10-dihydro: **Disorazole D₁**

C₄₃H₅₆N₂O₁₁ 776.922

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -103.6 (c, 0.6 in MeOH). λ_{\max} 266 (sh); 272 (ε 42800); 282; 304 (sh) (MeOH).

9,10-Dihydroxy, 9,10-dihydro, O¹⁰-Me:

Disorazole D₄

C₄₄H₅₈N₂O₁₁ 790.949

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +27.4 (c, 0.1 in MeOH). λ_{\max} 264 (sh); 273 (ε 60400); 282; 302 (sh) (MeOH).

Epimer, 9,10-dihydroxy, 9,10-dihydro:

Disorazole D₂

C₄₃H₅₆N₂O₁₁ 776.922

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -67 (c, 0.6 in MeOH). Epimeric at C-9 and/or C-10. λ_{\max} 252; 265 (sh); 272 (ε 105000); 281; 304 (sh) (MeOH).

Demethoxy, 9,10:9',10'-diepoxyde, 5',6'-Z-didehydro: **Disorazole B₁**

[158252-69-8, 158181-50-1]

C₄₂H₅₀N₂O₁₀ 742.864

Metab. from the bacterium *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +64.7 (c, 0.5 in MeOH/CH₂Cl₂, 1:1). λ_{\max} 288 (ε 62661); 304 (MeOH) (Berdy).

Demethoxy, 9',10'-dihydroxy, 9,10-epoxyde 5',6'-Z-didehydro, 9',10'-dihydro:

Disorazole B₄

[158181-51-2]

C₄₂H₅₂N₂O₁₁ 760.88

From *Sorangium cellulosum*. Cytotoxic, fungicidal. $[\alpha]_D^{22}$ -29.2 (c, 0.1 in MeOH).

Demethoxy, 9,9',10,10'-tetrahydroxy, 5',6'-Z-didehydro, 9,9',10,10'-tetrahydro: **Disorazole B₂**

[158181-49-8]

C₄₂H₅₄N₂O₁₂ 778.895

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -91.7 (c, 0.6 in MeOH). λ_{\max} 282 (ε 48417); 304 (MeOH) (Berdy).

5E-Isomer, 9R,10S-epoxyde: **Disorazole A₆**

C₄₃H₅₄N₂O₁₀ 758.907

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -436 (c, 0.1 in MeOH). λ_{\max} 265 (sh); 274; 282 (ε 65160); 298; 307; 325 (sh) (MeOH).

7Z-isomer, 9R,10R:9'R,10'S-diepoxyde: **Disorazole E₃**

C₄₃H₅₄N₂O₁₁ 774.906

From *Sorangium cellulosum*. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +24.5 (c, 0.2 in MeOH). λ_{\max} 278; 289; 297 (ε 45100); 307; 319 (sh) (MeOH).

7'Z-Isomer, demethoxy, 9,10:9',10'-diepoxyde, 5',6'-Z-didehydro: **Disorazole B₃**

C₄₂H₅₀N₂O₁₀ 742.864

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -140 (c, 0.4 in MeOH). λ_{\max} 289 (ε 59156); 308 (MeOH) (Berdy).

9'E-Isomer, 9R,10S-epoxyde: **Disorazole A₃**

C₄₃H₅₄N₂O₁₀ 758.907

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +101 (c, 0.6 in MeOH). λ_{\max} 274 (ε 57676); 282; 301 (MeOH) (Berdy).

toxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +81.3 (c, 0.1 in MeOH). λ_{\max} 254; 266 (sh); 274 (ε 73800); 283; 298; 306; 322 (sh) (MeOH).

11'E-Isomer, 9R,10S-epoxyde: **Disorazole A₄**

C₄₃H₅₄N₂O₁₀ 758.907

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +181 (c, 0.1 in MeOH). λ_{\max} 252; 263 (sh); 272 (ε 56400); 281; 298; 306; 321 (sh) (MeOH).

11'E-Isomer, 9,10-dihydroxy, 9,10-dihydro: **Disorazole D₃**

C₄₃H₅₆N₂O₁₁ 776.922

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ -16.7 (c, 0.15 in MeOH). λ_{\max} 262 (sh); 271 (ε 59800); 286; 293; 304 (sh) (MeOH).

9E,11E-Isomer: **Disorazole F₃**

C₄₃H₅₄N₂O₉ 742.908

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +29.2 (c, 0.1 in MeOH). λ_{\max} 252; 264 (sh); 271 (ε 38500); 281; 320 (sh); 351 (ε 40500); 365 (sh) (MeOH).

9'E,11'E-Isomer, 9R,10S-epoxyde: **Disorazole A₅**

C₄₃H₅₄N₂O₁₀ 758.907

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +21.5 (c, 0.5 in MeOH). λ_{\max} 242; 252; 263 (sh); 271 (ε 53300); 280; 298; 305; 321 (sh) (MeOH).

9'E,11'E-isomer, 10-methoxy, 9-hydroxy, 9,10-dihydro: **Disorazole D₅**

C₄₄H₅₈N₂O₁₁ 790.949

From *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +64.9 (c, 0.1 in MeOH). λ_{\max} 252; 262 (sh); 270 (ε 75500); 280 (ε 58500); 300 (sh) (MeOH).


Jansen, R. et al., *Annalen*, 1994, 759 (isol, uv, pmr, cmr, ms)

Irschik, H. et al., *J. Antibiot.*, 1995, 48, 31 (isol, ir, props)

Disorazole G₁

[158181-59-0]

D-860




C₄₃H₅₄N₂O₁₀ 758.907

Metab. of bacterium *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH; poorly sol. H₂O, hexane. $[\alpha]_D^{22}$ +101 (c, 0.6 in MeOH). λ_{\max} 274 (ε 57676); 282; 301 (MeOH) (Berdy).

Jansen, R. *et al.*, *Annalen*, 1994, 759 (isol, uv, pmr, cmr, struct)


Disorazole G₂ **D-861**
[158181-60-3]



C₄₃H₅₄N₂O₁₀ 758.907
Prod. by bacterium *Sorangium cellulosum*. Cytotoxic, fungicidal. $[\alpha]_D^{25}$ -161 (c, 0.15 in MeOH).

Jansen, R. *et al.*, *Annalen*, 1994, 759 (isol, uv, pmr, cmr, struct)


Disorazole G₃ **D-862**
[158181-61-4]



C₄₃H₅₆N₂O₁₁ 776.922
Isol. from bacterium *Sorangium cellulosum*. Cytotoxic, fungicidal. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. $[\alpha]_D^{25}$ -47.4 (c, 0.6 in MeOH). λ_{\max} 252; 274; 282 (MeOH) (Berdy).

Jansen, R. *et al.*, *Annalen*, 1994, 759 (isol, uv, pmr, cmr, struct)

Dispacamide A **D-863**
[177744-33-1]



C₁₁H₁₁Br₂N₅O₂ 405.048

Closely related to Tauroacidin A, T-39. Major alkaloid from the Caribbean sponges *Agelas dispar* and *Agelas clathrodes*. Exhibits remarkable antihistamine activity. Amorph. solid.

2-Debromo-Dispacamide B. 2-Debromodispacamide A
[177744-34-2]
C₁₁H₁₂BrN₅O₂ 326.152
Major alkaloid from the sponges *Agelas conifera* and *Agelas longissima*. Shows remarkable antihistamine activity.

9-Hydroxy-Dispacamide C
[197448-22-9]
C₁₁H₁₁Br₂N₅O₃ 421.048
Alkaloid from *Agelas* spp. λ_{\max} 227 (ε 11200); 272 (ε 12500) (MeOH).

2-Debromo-9-hydroxy-Dispacamide D. Mukanadin A
[197448-24-1]
C₁₁H₁₂BrN₅O₃ 342.152
Alkaloid from *Agelas* spp. and *Axinella verrucosa*.

2-Debromo-9-methoxy-9-Methoxydispacamide B. 9-O-Methylmukanadin A
C₁₂H₁₄BrN₅O₃ 356.178
Isol. from *Axinella verrucosa*.

Bis(debromo)-Debromodispacamide B
C₁₁H₁₃N₅O₂ 247.256
Alkaloid from *Agelas mauritiana*.

Bis(debromo)-9-hydroxy-Debromodispacamide D
C₁₁H₁₃N₅O₃ 263.255
Alkaloid from *Agelas mauritiana*. Racemic.

Cafieri, F. *et al.*, *Tet. Lett.*, 1996, **37**, 3587-3590 (isol, uv, ir, pmr, cmr)

Cafieri, F. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 2283-2288 (isol, pmr, cmr)

Lindel, T. *et al.*, *Tet. Lett.*, 1997, **38**, 8935-8938 (synth)

Uemoto, H. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1581-1583 (Mukanadin A)


Fresneda, P.M. *et al.*, *Tet. Lett.*, 2001, **42**, 851-854 (synth)

Hoffmann, H. *et al.*, *Synthesis*, 2003, 1753-1783 (rev)

Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 17-24 (9-Methoxydispacamide B)

Vergne, C. *et al.*, *Org. Lett.*, 2008, **10**, 493-496 (Debromodispacamides B,D)


Dispegatrine **D-864**
10,10',17,17'-Tetrahydroxy-4,4'-dimethyl-9,9'-bisarparanium, 9CI
[102488-56-2]



C₄₀H₄₈N₄O₄⁺ 648.844
Alkaloid from the roots of *Rauwolfia verticillata* var. *hainanensis* (Apocynaceae). Has α-adrenergic blocking props. Cryst. $[\alpha]_D^{23}$ +230 (c, 0.1 in MeOH). Mp >280° dec.

Lin, M. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 114-118

Dispyrin **D-865**
[934995-58-1]




C₁₈H₂₃Br₂N₃O₂ 473.207
Alkaloid from *Agelas dispar*. λ_{\max} 293 (log ε 2.4) (MeOH).

Piña, I.C. *et al.*, *J. Nat. Prod.*, 2007, **70**, 613-617 (isol, pmr, cmr, ms)

Yoshida, M. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 1362-1363 (synth)

Kennedy, J.P. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1783-1786 (synth)

Distamycin A **D-866**
Stallimycin, INN. Herperal. Dst 3. FI 6426. Distamycin 3
[636-47-5]



C₂₂H₂₇N₉O₄ 481.513

Peptide antibiotic. Closely related to Netropsin, N-180. Isol. from *Streptomyces distallicus* NRRL2886. Antiviral agent, used topically to treat herpes simplex infections. Sol. H₂O, MeOH; poorly sol. butanol, hexane. Mp 154-156°. Log P -1.45 (uncertain value) (calc). λ_{\max} 237 (ε 30000); 302 (ε 37000) (EtOH) (as hydrogen halide salt) (Derep). λ_{\max} 236 (E1%/1cm 591); 305 (E1%/1cm 723) (H₂O) (Berdy).

▶ LD₅₀ (mus, ipr) 500 mg/kg; LD₅₀ (mus, ivn) 75 mg/kg. WZ7108000

Hydrochloride: Stallimycin hydrochloride, USAN

[6576-51-8]

Cryst. (dil. HCl). Mp 186-189°.

▶ LD₅₀ (mus, ipr) 160 mg/kg. WZ7110000

N'-De-Me: 1-Nordistamycin A

[85407-06-3]

C₂₁H₂₅N₉O₄ 467.486

Synthetic analogue. Shows improved antiviral activity and similar toxicity to parent. Mp 150° (darkens).

Arcamone, F. *et al.*, *Nature (London)*, 1964, **203**, 1064 (isol, struct, synth)


Arcamone, F. *et al.*, *Gazz. Chim. Ital.*, 1967, **97**, 1097; 1110 (isol, struct, uv, ir, nmr, synth)

Hahn, F.E. *et al.*, *Antibiotics*, (Corcoran, J.W. *et al.* Eds.), Springer, N.Y., 1975, **3**, 79 (rev)

Bialer, M. *et al.*, *Tetrahedron*, 1978, **34**, 2389 (pmr, ir, ms, synth)

Grehn, L. *et al.*, *J. Med. Chem.*, 1983, **26**, 1042 (derivs)

- Bialer, M. *et al.*, *Dev. Mol. Virol.*, 1984, **4**, 143 (props, rev)
 Lown, J.W. *et al.*, *J.O.C.*, 1985, **50**, 3774 (synth, bibl)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 555
 Boger, D.L. *et al.*, *J.A.C.S.*, 2000, **122**, 6382-6384 (synth)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DXG600; SLI300


Distichamine**D-867**

Relative Configuration

- $C_{18}H_{19}NO_5$ 329.352
 Alkaloid isol. from the bulbs of *Boophone disticha* (Amaryllidaceae).
 Large prisms (Me₂CO). Mp 161-162°. $[\alpha]_D^{21}$ -56 (c, 0.478 in CHCl₃).
 Methiodide:
 Cryst. (MeOH). Mp 223-226° dec.
 Hauth, H. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 491 (isol, uv, ir)
 Steenkamp, P.A. *et al.*, *Thesis*, Univ. of Johannesburg, 2005, (struct)

Distomadine A**D-868**

[549521-28-0]



R = H

- $C_{16}H_{14}N_4O_4$ 326.311
 Isol. from the ascidian *Pseudodistoma aureum*. Yellow powder. Mp 232° dec. $[\alpha]_D^{20}$ +47 (c, 0.3 in MeOH). Fluorescent.
 Pearce, A.N. *et al.*, *Tet. Lett.*, 2003, **44**, 3897-3899 (isol, pmr, cmr)

Distomadine B**D-869**


[549521-30-4]

As Distomadine A, D-868 with

R = COOH

 $C_{17}H_{14}N_4O_6$ 370.321Isol. from the ascidian *Pseudodistoma aureum*.


- Pearce, A.N. *et al.*, *Tet. Lett.*, 2003, **44**, 3897-3899 (isol, pmr, cmr)

Cortinarius Disulfide**D-870** $C_{16}H_{26}N_4O_6S_2$ 434.537Isol. from *Cortinarius* sp. Possible artifact.

- Nicholas, G.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 341-344 (isol, pmr, cmr, ms)

Dithermamine**D-871**

3',4'-Dihydro-3,5'-bithermopsine [38948-08-2]

 $C_{30}H_{40}N_4O_2$ 488.672Alkaloid from the above-ground parts of *Thermopsis lanceolata* (Fabaceae). Mp 235°. $[\alpha]_D$ +121.8 (c, 1.0 in CHCl₃). Dec. on dist. to Thermopsine, T-363.


Perchlorate: Mp 211-212°.

Picrate: Mp 203-204°.

- Vinogradova, V.I. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 463; 1972, **8**, 87; *Chem. Nat. Compd. (Engl. Transl.)*, 82; 440 (isol, ir, ms, struct)

Dithicolchicine**D-872** $C_{22}H_{25}NO_4S_2$ 431.576Struct. unknown. Alkaloid from the seeds of *Colchicum autumnale* (Liliaceae). Yellow plates (MeOH). Mp 265-267°. $[\alpha]_D$ -366 (c, 1.00 in CHCl₃). Purification by chromatography gave a prod. with no carbonyl group and no reaction with conc. HCl. Material was never isolated again and is possibly an artifact.

- Bellet, P. *et al.*, *Ann. Pharm. Fr.*, 1955, **13**, 84-87; *CA*, **50**, 4107h (isol)


Dithyreanitrile**D-873**7-Methoxy- α,α -bis(methylthio)-1H-indole-3-acetonitrile, 9CI [128717-80-6] $C_{13}H_{14}N_2OS_2$ 278.398Alkaloid from the seeds of *Dithyrea wislizenii* (Brassicaceae). Insect antifeedant. Off-white cryst. Mp 135°.

- Powell, R.G. *et al.*, *Experientia*, 1991, **47**, 304 (isol, pmr, cmr, ms, cryst struct)

- Mantus, E.K. *et al.*, *Tet. Lett.*, 1993, **34**, 1085 (synth)

Ditryptophenaline**D-874**

[64947-43-9]



Absolute Configuration

 $C_{42}H_{40}N_6O_4$ 692.816Diketopiperazine dimer. Prod. by *Aspergillus flavus* var. *columnaris* MRC1174. Inhibits Substance P binding. Tachykinin antagonist. Cryst. (CH₂Cl₂/MeOH). Mp 204-205°. $[\alpha]_D^{23.5}$ -330 (c, 52 in CH₂Cl₂). There is some uncertainty about the abs. config. of the nat. prod. The original isolate was reported to have a CD spectrum corresp. to the enantiomeric config., but this appears to have been an error in the publication. λ_{max} 244 (ε 15250); 303 (ε 6200) (EtOH).

N-1'-(2-Phenylethyl): 1'-(2-Phenylethylene)dityryptophenaline [159355-77-8]

 $C_{50}H_{46}N_6O_4$ 794.951From *Aspergillus flavus* SC1661. Mp 183-186°. $[\alpha]_D$ -125 (c, 0.05 in CHCl₃). λ_{max} 221 (ε 33800); 244 (ε 11100); 324 (ε 14690); 338 (ε 14180) (MeOH) (Berdy).

- Springer, J.P. *et al.*, *Tet. Lett.*, 1977, 2403-2406 (isol, struct)

- Maes, C.M. *et al.*, *J.C.S. Perkin 1*, 1986, 861-866 (isol, pmr, abs config)

- Barrow, C.J. *et al.*, *J.O.C.*, 1993, **58**, 6016-6019 (isol, activity)

- Barrow, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1239-1244 (1'-(2-


Phenylethylene)dityryptophenaline, activity)

- Overman, L.E. *et al.*, *J.A.C.S.*, 2001, **123**, 9465-9467 (synth)

- Movassaghi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1485-1487 (synth)

Divarine**D-875**


[158848-31-8]

 $C_{39}H_{44}N_4O_2$ 600.802Alkaloid from *Strychnos divaricans* (Loganiaceae). Needles (MeOH/EtOAc). Mp 240-242° dec.

Mukherjee, R. *et al.*, *Heterocycles*, 1994, **38**, 1965 (*isol, uv, pmr, cmr, struct*)

Dnacin A₁ **D-876**

Antibiotic C 14482A₁. C 14482A₁ [76828-83-6]



R = CN

C₂₀H₂₃N₅O₄ 397.433

The abs. config. is likely to be the opposite of that illus. See note under Naphthyrindinomycin A, N-32. Isol. from *Nocardia* sp. C-14482 (N-1001) and *Actinosynnema pretiosum*. Phosphatase inhibitor, DNA biosynth. inhibitor. Dark red needles. Sol. CHCl₃, MeOH, DMSO; fairly sol. EtOH, EtOAc, H₂O, butanol; poorly sol. hexane. Mp 300° dec. [α]_D^{24.5} +150 (c, 0.05 in EtOH). λ_{max} 213 (ε 22300); 281 (ε 9000); 496 (ε 2100) (MeOH) (Derep). λ_{max} 214 (E1%/1cm 503); 281 (E1%/1cm 209); 498 (E1%/1cm 46) (EtOH) (Berdy). λ_{max} 213; 281; 496 (MeOH/NaOH) (Berdy).

► LD₅₀ (mus, ivn) 5 mg/kg.

[69913-37-7]

Tanida, S. *et al.*, *J. Antibiot.*, 1980, **33**, 1443 (*isol*)

Muroi, M. *et al.*, *J. Antibiot.*, 1980, **33**, 1449 (*uv*)

Hida, T. *et al.*, *J. Antibiot.*, 1994, **47**, 917 (*pmr, cmr, struct*)

Dnacin B₁ **D-877**

Antibiotic C 14482B₁. C 14482B₁ [76828-84-7]

As Dnacin A₁, D-876 with

R = OH

C₁₉H₂₄N₄O₅ 388.422

Quinone antibiotic. Isol. from *Nocardia* sp. C-14482 (N-1001). Active against gram-positive and -negative bacteria and weakly active against fungi. Dark-red or reddish-brown needles. Mp 300° dec. [α]_D²⁰ +50 (c, 0.06 in CHCl₃). Strain also prod. C 14482B₂ and C 14482B₃. λ_{max} 213 (ε 22300); 281 (ε 9000); 496 (ε 2100) (MeOH) (Derep).

► JQ5811000

[76050-36-7]

Tanida, S. *et al.*, *J. Antibiot.*, 1980, **33**, 1443 (*isol*)

Muroi, M. *et al.*, *J. Antibiot.*, 1980, **33**, 1449 (*uv*)

Tanida, S. *et al.*, *Antimicrob. Agents Chemother.*, 1982, **22**, 735 (*props*)

Hida, T. *et al.*, *J. Antibiot.*, 1994, **47**, 917 (*pmr, cmr, props*)

2,4,10-Docosatrienoic isobutylamide **D-878**

C₂₆H₄₇NO 389.663

(E,E,Z)-form**Filifiline**

[62726-18-5]


Constit. of the fruits of *Piper officinarum* (Javanese long pepper). Has antifertility props. Cryst. (petrol). Mp 66-66.5°.

Gupta, O.P. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 912 (*isol*)

Gupta, O.P. *et al.*, *Indian J. Pharm. Sci.*, 1978, **40**, 113 (*use*)

6-Docosenoic acid, 9CI **D-879**

[116802-23-4]



C₂₂H₄₂O₂ 338.573

(Z)-form [82683-06-5]**Amide: 6-Docosenamamide**

[145701-07-1]

C₂₂H₄₃NO 337.588

Isol. from *Asimina parviflora*. Mp 42-44°.

[82683-25-8]

Ratnayake, S. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1462-1467 (*amide, isol*)

13-Docosenoic acid **D-880**

[25378-26-1]



C₂₂H₄₂O₂ 338.573

Present in lipids of *Physalia physalis* (Portuguese-man-of-war). Antineoplastic agent. Log P 9.86 (uncertain value) (calc).

(E)-form**Brassicidic acid**

[506-33-2]

Cryst. (EtOH). Mp 61.5°. Bp₃₀ 282°

Bp₁₀ 256°.

Me ester: [7439-44-3]

C₂₃H₄₄O₂ 352.599

Mp 34-35°.

Et ester: [116723-91-2]

C₂₄H₄₆O₂ 366.626

Mp 30.5°.

Nitrile:

C₂₂H₄₁N 319.573

Mp 22°. Bp₁₇ 257°.

Anhydride:

C₄₄H₈₂O₃ 659.13

Mp 64°.

(Z)-form**Erucic acid**

[112-86-7]

[28929-01-3]

An important vegetable fatty acid, mainly confined to seed fats of the families Brassicaceae and Tropaeolaceae (e.g. rape, mustard and wallflower seeds). Occurs widely in marine organisms. Cryst. (MeCN). Mp 33.5°. Pharmacol. active isomer. Undergoes solid-state phase transitions.

Me ester: [1120-34-9]

Oil. Bp₅ 221-222°. Ir v 1744 cm⁻¹.

Et ester: [37910-77-3]

Bp₅ 229-230°.

Chloride: [7459-29-2]

C₂₂H₄₁ClO 357.018

Mp 14°.

Amide: 13-Docosenamamide. Newtron S [112-84-5]

C₂₂H₄₃NO 337.588

Isol from the cerebrospinal fluid of humans, cats and rats. Antiblocking agent/lubricant for polymers. Sleep inducer. Mp 94° (65-66°).

Nitrile: [73170-89-5]

Bp 238-240°.

Anhydride: [103213-60-1]

Mp 47.5-48°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 501D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 784A; 1221C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 588C (*ir*)

Holde, D. *et al.*, *Ber.*, 1923, **56**, 2052-2058

(*anhydrides*)

Org. Synth., Coll. Vol., 2, 1943, 258 (*synth*)

Bowman, R.E. *et al.*, *J.C.S.*, 1950, 177 (*synth*)

Lakizo, V.I. *et al.*, *CA*, 1968, **69**, 2475g (*synth*)

Niewiadomski, H. *et al.*, *Przem. Chem.*, 1969, **48**, 659; *CA*, **72**, 78; 293 (*rev*)

Chang, S.P. *et al.*, *J. Am. Oil Chem. Soc.*, 1972, **49**, 422 (*synth*)

Mitcham, D. *et al.*, *J. Am. Oil Chem. Soc.*, 1973, **50**, 446 (*ir*)

Stillway, L.W. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1976, **53**, 535-537

(*Physalia physalis consti*)

Barrette, J.P. *et al.*, *J. Assoc. Off. Anal. Chem.*, 1976, **59**, 855 (*glc*)

Yamada, K. *et al.*, *J.O.C.*, 1978, **43**, 2076

(*synth*)

Kaneko, F. *et al.*, *Acta Cryst. C*, 1992, **48**, 1060; 1993, **49**, 1232 (*cryst struct, erucic acid*)


Painuly, P. *et al.*, *J. Chromatogr.*, 1992, **590**, 139 (*purifn*)

Kling, M.R. *et al.*, *J.C.S. Perkin 1*, 1993, 1183 (*synth, ir, ms*)

Lerner, R. *et al.*, *Science (Washington, D.C.)*, 1995, **268**, 1506 (*isol, amide*)

3,5-Dodecadien-2-amine **D-881**

2-Amino-3,5-dodecadiene




C₁₂H₂₃N 181.32

(±)-(3E,5Z)-form [134387-52-3]

Constit. of the ascidian *Pseudodistoma novaezelandiae*. λ_{max} 233 (MeOH) (Berdy).

Perry, N.B. *et al.*, *Aust. J. Chem.*, 1991, **44**, 627-633 (*isol, uv, pmr, cmr*)

2,4-Dodecadiene-8,10-diynoic acid, 9CI **D-882**

C₁₂H₁₂O₂ 188.226

(2E,4E)-form

2-Methylpropylamide: [18679-23-7]

Isol. from *Achillea* sp. Cryst. (Et₂O/petrol). Mp 146°.

(2E,4Z)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2,4-dodecadiene-8,10-diyamide, **9CI**. 2,4-Dodecadiene-8,10-diyonic acid isobutylamide. N-Isobutyl-2,4-dodecadiene-8,10-diyamide
[113817-69-9]

C₁₆H₂₁NO 243.348

Isol. from *Echinacea purpurea* and *Echinacea angustifolia* (Asteraceae). Cryst. (hexane). Mp 74°.

2-Methylbutylamide: N-(2-Methylbutyl)-2,4-dodecadiene-8,10-diyamide, **9CI**. 2,4-Dodecadiene-8,10-diyonic acid 2-methylbutylamide
[117505-95-0]

C₁₇H₂₃NO 257.375

Isol. from roots of *Echinacea purpurea* (Asteraceae). Oil.

(2Z,4E)-form

2-Methylpropylamide: [13894-70-7]
Isol. from *Echinacea purpurea*, *Echinacea angustifolia* and *Achillea* sp. Cryst. (petrol). Mp 86°. λ_{max} 258 (4.44) (MeOH).

[52997-50-9]

Bohlmann, F. *et al.*, *Chem. Ber.*, 1966, **99**, 3197; 1967, **100**, 3861; 1973, **106**, 1328; 1974, **107**, 2120 (*isol, struct, synth, uv, pmr, ir*)

Bauer, R. *et al.*, *Phytochemistry*, 1988, **27**, 2339; 1989, **28**, 505 (*isol, uv, ir, pmr, ms, struct*)

Chen, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 773-776 (*isol, pmr, cmr, ms*)

Kraus, G.A. *et al.*, *Molecules*, 2006, **11**, 758-767 (2Z,4E-isobutylamide, *synth*)

2,4-Dodecadienoic acid D-883

H₃C(CH₂)₆CH=CHCH=CHCOOH

C₁₂H₂₀O₂ 196.289

Probable constit. of *Sebastiania linguistrina* seed oil.

(2E,4E)-form [24738-48-5]

Cryst. (petrol). Mp 49-51°.

Me ester: [24738-46-3]

C₁₃H₂₂O₂ 210.316

Bp_{0.3} 106-108°.

2-Methylpropylamide: 2,4-Dodecadienoic acid isobutylamide
[24738-51-0]

C₁₆H₂₉NO 251.411

Alkaloid from *Anacyclus pyrethrum* and *Leucocyclus formosus* (Asteraceae) (in admixture with homologues). Mp 89-90°.

Pyrrolidide: 2,4-Dodecadienoic acid pyrrolidide. 1-(2,4-Dodecadienoyl)pyrrolidide
[117137-69-6]

C₁₆H₂₇NO 249.395

Constit. of pepper (*Piper nigrum*) (Piperaceae). Shows larvicidal activity. Not obt. completely pure.

Holman, R.T. *et al.*, *J. Am. Oil Chem. Soc.*, 1955, **32**, 356 (*isol*)

Burden, R.S. *et al.*, *J.C.S. (C)*, 1969, 2477 (*isol, synth*)

Greger, H. *et al.*, *Phytochemistry*, 1981, **20**, 2579 (*isol, ms, struct*)

Kiuchi, F. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2452 (*pyrrolidide*)

Abarbri, M. *et al.*, *Synthesis*, 1996, 82 (*synth, ir, pmr, cmr, ms*)

2,8-Dodecadienoic acid D-884

H₃C(CH₂)₂CH=CH(CH₂)₄CH=CHCOOH

C₁₂H₂₀O₂ 196.289

(2E,8E)-form

Large plates. Mp 34-35°. Bp_{0.8} 140°.

S-Benzylthiuronium salt:
Plates (EtOAc). Mp 159°.

2-Methylpropylamide:
Needles (petrol). Mp 53°.

(2Z,8E)-form

Bp₁ 127-129°.

S-Benzylthiuronium salt:
Plates (EtOAc). Mp 148°.

2-Methylpropylamide: Bp_{0.6} 154° Bp_{0.03} 142°.

(2ξ,8ξ)-form

2-Methylpropylamide: N-Isobutyl-2,8-dodecadienamamide. **Herculin**
C₁₆H₂₉NO 251.411
Isol. from *Zanthoxylum clava-herculis* (Hercules' club). Insecticide. Pungent needles (hexane). Mp 59-60°. Struct. doubtful.

Jacobson, M. *et al.*, *J.A.C.S.*, 1948, **70**, 4234 (*isol, struct*)

Raphael, R.A. *et al.*, *J.C.S.*, 1950, 115; 1951, 2693 (*synth*)

Crombie, L. *et al.*, *J.C.S.*, 1952, 2997 (*synth, struct*)

N-(3,6-Dodecadienyl)-sulfamic acid D-885

[1040630-93-0]

H₃C(CH₂)₄CH=CHCH₂CH=CHCH₂CH₂NHSO₃H


C₁₂H₂₃NO₃S 261.385

(Z,Z)-form

Isol. from *Daphnia pulex*. Kairomone.

Yasumoto, K. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 133-136 (*isol, pmr*)

Dodecahydro-2-methylpyrido[2,1,6-de]quinolizine D-886



(2α,3αα,6αβ,9αβ)-(3aS,9aS)-form

C₁₃H₂₃N 193.331

(2α,3αα,6αβ,9αβ)(3aS,9aS)-form Hippodamine

[53990-36-6]

[59685-36-8 ((±)-form)]

Alkaloid from the defence secretion of the ladybugs *Hippodamia convergens*, *Hippodamia caseyi*, *Anisosticta 19-punctata* and *Chauliognathus pulchellus*. Oil. Small and solvent-dependent opt. rotn. which could not be reliably measured.

N-Oxide: Convergin

[41701-36-4]

[59685-37-9 ((±)-form)]

C₁₃H₂₃NO 209.331

Constit. of the defence secretion of *Hippodamia convergens* and *Hippodamia caseyi*. Oil; solid (as hydrochloride). Hydrochloride chars without melting above 215°.

(2α,3αβ,6αα,9αβ)-form

Precoccinelline

[38211-56-2]

Alkaloid from the defence secretion of the ladybugs *Coccinella transversoguttata*, *Coccinella septempunctata*, *Chauliognathus pulchellus* and *Hippodamia convergens*. Oil. Mp 195-197° (as picrate). Meso-stereoisomer.

N-Oxide: Coccinelline

[34290-97-6]

C₁₃H₂₃NO 209.331

Constit. of the defence secretion of *Coccinella transversoguttata* and *Coccinella septempunctata*. Cryst. (Me₂CO). Mp 204° dec. Opt. inactive. Chars >215°.

(2α,3αβ,6αβ,9αβ)-form

Myrrhine

[58207-39-9]

Alkaloid from the defence secretion of the ladybug *Myrrha octodecimguttata*. Noncryst. Meso-stereoisomer.

N-Oxide:

Needles (CH₂Cl₂). Dec. >150° without melting.

Karlsson, R. *et al.*, *Chem. Comm.*, 1972, 626 (*Precoccinelline, cryst struct*)

Tursch, B. *et al.*, *Tet. Lett.*, 1974, 409

(*Hippodamine, cryst struct, abs config*)

Tursch, B. *et al.*, *Tetrahedron*, 1975, **31**, 1541

(*Myrrhine, Coccinelline, struct, biosynth*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1976, **54**, 473; 1494; 1807 (*Hippodamine, Myrrhine, Coccinelline, Precoccinelline, synth, isol, ir, pmr, ms, oxide*)

Stevens, R.V. *et al.*, *J.A.C.S.*, 1979, **101**, 7032

(*Coccinelline, synth*)

Mueller, R.M. *et al.*, *J.O.C.*, 1984, **49**, 2217

(*Hippodamine, Myrrhine, Precoccinelline, synth, ir, cmr, pmr, ms*)

Yue, C. *et al.*, *Tetrahedron*, 1994, **50**, 3139

(*Precoccinelline, synth*)


Lebrun, B. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 60-64 (*pmr, cmr*)

Laurent, P. *et al.*, *Insect Biochem. Mol. Biol.*, 2002, **32**, 1017-1023 (*biosynth*)

Rejzek, M. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 73-83 (*synth*)

Gerasyuto, A.I. *et al.*, *Org. Lett.*, 2006, **8**, 4899-4902 (*synth*)

1,1a,2,4,4a,5,5a,6,6a,7,8,8a-Dodecahydro-2aH-3-thia-4-azacyclopenta[mno]pentaleno[2,1,6,5-fghi]-aceanthrylene-2a-carboxylic acid, 9CI




C₁₉H₁₉NO₂S 325.431

3,5,8-Trimethyldecyl ester: [87055-59-2]

C₃₂H₄₅NO₂S 507.779Isol. from *Darius* crude oil.Masohan, A. et al., *J. Indian Chem. Soc.*, 1983, **60**, 392-395 (isol, struct)**Dodecahydro-2,5,8-trimethyl-1,4,7,9b-tetraazaphenalene, 9CI**

D-888

Tricotonyltetramine

(2 α ,3 $\alpha\alpha$,5 α ,6 $\alpha\alpha$,8 α ,9 $\alpha\alpha$)-formC₁₂H₂₄N₄ 224.348**(2 α ,3 $\alpha\alpha$,5 α ,6 $\alpha\alpha$,8 α ,9 $\alpha\alpha$)-form***Bongardamine*

[221677-44-7]

Alkaloid from *Bongardia chrysogonum*. Cryst. (Me₂CO). Mp 203°. [α]_D²⁵ -7.7 (c, 0.3 in MeOH). λ_{\max} 205 (log ϵ 2.57) (MeOH).**(2 α ,3 $\alpha\beta$,5 α ,6 $\alpha\beta$,8 α ,9 $\alpha\beta$)-form***Base A*

[54352-30-6]

Alkaloid from *Lycopodium thyoides* and *Sophora alopecuroides* (Lycopodiaceae, Fabaceae), prev. known synthetically. Cryst. (Me₂CO). Mp 105° (101-103°). Opt. inactive. Said not to be an artifact by Monakhova et al, but other workers are uncertain.*Hydrate*: Mp 95-96°.*Hydrochloride* (1:2):

Cryst. (EtOH), Mp 289° dec.

Dipicrate: Mp 222-224°.**(2 α ,3 $\alpha\beta$,5 α ,6 $\alpha\beta$,8 β ,9 $\alpha\beta$)-form***Base B[†]*

[54352-31-7]

Isol. from *Lycopodium thyoides* (Lycopodiaceae). Mp 65°. Poss. artifact.Delépine, M. et al., *C. R. Hebd. Seances Acad. Sci.*, 1943, **216**, 649; 785 (synth)Manakhova, T.E. et al., *Khim. Prir. Soedin.*, 1974, **10**, 752; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 771 (isol, pmr, ms, struct)Braekman, J.C. et al., *Phytochemistry*, 1974, **13**, 2519 (isol, pmr, ms, struct)Atta-ur-Rahman, et al., *Phytochemistry*, 1999, **50**, 333-336 (*Bongardamine*)**Dodecanoic acid, 9CI**

D-889

Lauric acid. Laurostearic acid. FEMA 2614

[143-07-7]

H₃C(CH₂)₁₀COOHC₁₂H₂₄O₂ 200.32Occurs as glyceride in coconut oil and palm kernel oil. Widespread in plants, esp. in the Arecaceae and Lauraceae. Flavour ingredient, defoamer, lubricant. Cryst. (EtOH). d^{20} 0.88. Mp 44°. Bp₁₀₀ 225° Bp_{0.65} 141-142° (lit. gives a pressure range). pK_a 5.3 (20°).▶ LD₅₀ (rat, orl) 12000 mg/kg. OE9800000*Amide: Dodecanamide. Diamid Y*

[1120-16-7]

C₁₂H₂₅NO 199.336Isol. from *Rhizoclonium hieroglyphicum*. Cryst. Mp 110°. Bp_{12.5} 199-200°.

[37811-72-6, 2224-49-9]

Dembitsky, V.M. et al., *Phytochemistry*, 2000, **54**, 965-967 (isol, amide)**2,4,6,8,10-Dodecapentaene-dioic acid**

D-890

[6790-22-3]

HOOCCH=CHCH=CHCH=CHCH=CHCH=CHCOOH

C₁₂H₁₂O₄ 220.224**(all-E)-form***1-L-Aspartic acid-L-isoleucine diamide:**Boletocrocin C*C₂₂H₂₈N₂O₈ 448.472Isol. from the fruit bodies of *Boletus laetissimus*.**(all- ξ)-form***Di-Me ester*: [77185-99-0]C₁₄H₁₆O₄ 248.278

Deep yellow prisms. Mp 224.5-225.5°.

Subl. ca. 150.

Di-Et ester: [1441-60-7]

[6071-61-0]

C₁₆H₂₀O₄ 276.332

Mp 157.5-159.5°.

[101417-57-6]

Weedon, B.C.L. et al., *J.C.S.*, 1954, 4168-4174 (ester, synth)Kucherov, V.F. et al., *CA*, 1961, **55**, 24560i

(ester, synth)

Boschelli, D. et al., *Tet. Lett.*, 1985, **26**, 5239-

5242 (ester, synth)

Kahner, L. et al., *Phytochemistry*, 1998, **49**,1693-1697 (*Boletocrocin C*)**2,4,8,10-Dodecatetraenoic acid**

D-891

H₃CCH=CHCH=CHCH₂CH₂CH=CHCH=CHCOOHC₁₂H₁₆O₂ 192.257**(2E,4E,8E,10E)-form***2-Methylpropylamide*: [543-44-2]

[13894-71-8 (unspecified stereo)]

Mp 104-106°. Synthetic.

(2E,4E,8E,10Z)-form*2-Methylpropylamide*: [75947-32-9]C₁₆H₂₅NO 247.38Constit. of *Echinacea angustifolia*, *Echinacea purpurea* and *Spilanthes mauritiana*. Mp 78-79°. λ_{\max} 260 (ϵ 30000) (EtOH).**(2E,4E,8Z,10E)-form***2-Methylpropylamide*: [75917-90-7]Constit. of *Acmella ciliata*, *Asarum**forbesii*, *Echinacea angustifolia*, *Echinacea purpurea* and *Salmea scandens*. Needles (hexane). Mp 63°.*2-Methylbutylamide*: [99617-47-7]C₁₇H₂₇NO 261.406Constit. of *Acmella ciliata*.**(2E,4E,8Z,10Z)-form***2-Methylpropylamide*: [77448-63-6]Constit. of *Echinacea angustifolia*, *Echinacea purpurea* and *Salmea scandens*.**(2E,4Z,8Z,10E)-form***2-Methylpropylamide*: [99631-25-1]Constit. of *Acmella ciliata*.**(2E,4Z,8Z,10Z)-form***2-Methylpropylamide*: [851881-29-3]Constit. of *Asarum forbesii*. Needles (hexane). Mp 63°. λ_{\max} 236 ; 262 (MeOH).Herz, W. et al., *Phytochemistry*, 1985, **24**, 173-174 (*Salmea scandens* constits)Martin, R. et al., *Phytochemistry*, 1985, **24**,2295-2300 (*Acmella ciliata* constits)Jondiko, I.J.O. et al., *Phytochemistry*, 1986, **25**,2289-2290 (*Spilanthes mauritiana* constit)Bauer, R. et al., *Phytochemistry*, 1988, **27**,2339-2342 (*Echinacea purpurea* constits)Zhang, F. et al., *J. Asian Nat. Prod. Res.*, 2005,**7**, 1-5 (*Asarum forbesii* constits)Matovic, N. et al., *Org. Biomol. Chem.*, 2007,**5**, 169-174 (isol, synth, pmr, cmr)**2,4,8,11-Dodecatetraenoic acid**

D-892

H₂C=CHCH₂CH=CHCH₂CH₂CH=CHCH=CHCOOHC₁₂H₁₆O₂ 192.257**(2E,4E,8Z,11E)-form***2-Methylpropylamide: 2,4,8,11-Dodecate-**traenoic acid isobutylamide*

[117824-00-7]

C₁₆H₂₅NO 247.38Isol. from the aerial parts of *Brachycome ciliocarpa* (Asteraceae). Oil. Genus name given as the alternative spelling *Brachycome*.Zdero, C. et al., *Phytochemistry*, 1988, **27**, 2984

(isol, ir, pmr, ms, struct)

2,6,8,10-Dodecatetraenoic acid

D-893

H₃CCH=CHCH=CHCH=CHCH₂CH₂CH=CHCOOHC₁₂H₁₆O₂ 192.257**(all-E)-form***2-Methylpropylamide: β -Sanshool*

[10076-00-3]

Prod. by isomerisation of α -Sanshool.Constit. of *Zanthoxylum piperitum*

(Japanese pepper tree). Insecticide.

Cryst. (hexane). Mp 100-101°.

2-Hydroxy-2-methylpropylamide: Hydroxy- β -sanshool

[97465-69-5]

C₁₆H₂₅NO₂ 263.379Constit. of *Zanthoxylum bungeanum*and *Zanthoxylum piperitum* (Japanese

pepper tree). Powder.

(2E,6Z,8E,10E)-form [18744-21-3]

Cryst. (hexane). Mp 93.5-95°. Material contaminated with some *all-E*-isomer.

2-Methylpropylamide: N-Isobutyl-2,6,8,10-dodecatetraenamide. **α -Sanshool.** *Neoherculin*. *Echinacein* [504-97-2]

C₁₆H₂₅NO 247.38

Constit. of *Echinacea angustifolia* roots and *Zanthoxylum* spp. Insecticide. Cryst. (petrol). Mp 69° (61-63°). Unstable in air.

2-Hydroxy-2-methylpropylamide: Hydroxy- α -sanshool

[83883-10-7]

[645-70-5 (Sanshoamide)]

C₁₆H₂₅NO₂ 263.379

Constit. of *Zanthoxylum piperitum* (Japanese pepper tree) and *Zanthoxylum planispinum* (Rutaceae). Unstable oil. Probably the major constit. of Sanshoamide isol. as a mixt. in 1951. λ_{\max} 260 (ε 22400); 269 (ε 29200); 280 (ε 23200) (EtOH).

[119719-30-1]

Aihara, T. *et al.*, *Yakugaku Zasshi*, 1951, **71**, 1112 (*Sanshoamide*)

Crombie, L. *et al.*, *J.C.S.*, 1952, 2997; 1955, 995; 1957, 2760 (*isol, uv, ir, struct*)

Crombie, L. *et al.*, *Nature (London)*, 1954, **174**, 833

Sonnet, P.E. *et al.*, *J.O.C.*, 1969, **34**, 1147 (*synth, uv*)

Yasuda, I. *et al.*, *Phytochemistry*, 1982, **21**, 1295 (*isol, pmr, cmr, ms, occur*)

Mizutani, K. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2362 (*isol, ir, pmr, cmr*)

Nakamura, N. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2647 (*ir*)

2,4,6-Dodecatrienoic acid **D-894**

H₃C(CH₂)₄CH=CHCH=CHCH=CHCOOH

C₁₂H₁₈O₂ 194.273

(2E,4E,6E)-form

Et ester: [80387-34-4]

C₁₄H₂₂O₂ 222.327

Oil.

Amide: [149764-42-1]

C₁₂H₁₉NO 193.288

Isol. from an actinomycete *Amycolatopsis* sp. Herbicide. Sol. MeOH, hexane, Me₂CO, EtOAc, CHCl₃; fairly sol. H₂O. λ_{\max} 296 (MeOH) (Berdy).

(2E,4E,6Z)-form

Constit. of the latex of *Euphorbia lathyris*.

Amide: [149764-40-9]

Isol. from an actinomycete *Amycolatopsis* sp. Herbicide. Sol. MeOH, Me₂CO, CHCl₃, EtOAc, hexane; fairly sol. H₂O. λ_{\max} 297 (MeOH) (Berdy).

(2Z,4E,6E)-form

Constit. of the latex of *Euphorbia lathyris*. Isol. as Me ester.

[77761-51-4, 77761-50-3]

Warnaar, F. *et al.*, *Phytochemistry*, 1981, **20**, 89 (*isol, ms*)

Sakai, T. *et al.*, *J.O.C.*, 1982, **47**, 1101 (*synth, ester, ir, pmr, cmr, ms*)

Japan. Pat., 1993, 05 977 86; *CA*, **119**, 133461y (*isol, amide*)

2,4,8-Dodecatrienoic acid **D-895**

H₃CCH₂CH₂CH=CHCH₂CH₂CH=CHCH=CHCOOH

C₁₂H₁₈O₂ 194.273

(2E,4E,8Z)-form

2-Methylpropylamide: 2,4,8-Dodecatrienoic acid isobutylamide

[117505-96-1]

C₁₆H₂₇NO 249.395

Isol. from the roots of *Echinacea purpurea* (Asteraceae). Oil.

Bauer, R. *et al.*, *Phytochemistry*, 1988, **27**, 2339 (*isol, uv, ir, pmr, ms, struct*)

2,4,10-Dodecatrien-8-ynoic acid **D-896**

H₃CCH=CHC≡CCH₂CH₂CH=CHCH=CHCOOH

C₁₂H₁₄O₂ 190.241

(2E,4E,10E)-form

2-Methylpropylamide: 2,4,10-Dodecatrien-8-ynoic acid isobutylamide

[113836-87-6]

C₁₆H₂₃NO 245.364

Constit. of the roots of *Echinacea purpurea*. Oil. λ_{\max} 232; 261 (MeCN aq.).

(2E,4Z,10Z)-form

2-Methylpropylamide: [120727-33-5]

C₁₆H₂₃NO 245.364

Constit. of the roots of *Echinacea angustifolia*. Oil. λ_{\max} 231 (sh); 261 (MeCN aq.).

(2Z,4E,10Z)-form

2-Methylpropylamide: Constit. of the roots of *Echinacea angustifolia*. Viscous oil. λ_{\max} 257 (log ε 4.1) (MeOH).

Bauer, R. *et al.*, *Phytochemistry*, 1988, **27**, 2339-2342; 1989, **28**, 505-508 (*isol, uv, ir, pmr, ms*)

Chen, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 773-776 (*isol, pmr, cmr, ms*)

2-Dodecene-8,10-diynoic acid, 9CI **D-897**

H₃CC≡CC≡C(CH₂)₄CH=CHCOOH

C₁₂H₁₄O₂ 190.241

(E)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2-dodecene-8,10-diyamide, 9CI.

2-Dodecene-8,10-diynoic acid isobutylamide

[120727-29-9]

C₁₆H₂₃NO 245.364

Isol. from the roots of *Echinacea angustifolia*. Cryst. (hexane). Mp 104°. λ_{\max} 211 (ε 10600) (MeCN aq.).

2-Methylbutylamide: N-(2-Methylbutyl)-2-dodecene-8,10-diyamide, 9CI

[120727-31-3]

C₁₇H₂₅NO 259.391

Isol. from the roots of *Echinacea angustifolia*. Oil.


Bauer, R. *et al.*, *Phytochemistry*, 1989, **28**, 505-508 (*isol, pmr, ms*)

Wu, L. *et al.*, *Phytochemistry*, 2004, **65**, 2477-2484 (*synth, occur*)

2-Dodecyl-4-hydroxyquinoline **D-898**

2-Dodecyl-4(1H)-quinolinone. 2-Dodecyl-4-quinolinol, 9CI

[80554-65-0]



C₂₁H₃₁NO 313.482

Alkaloid from *Boronia ternata* var. *elongata* and *Ruta graveolens*. Metab. of *Pseudomonas aeruginosa*. Amorph. powder. Mp 70°. λ_{\max} 215 (log ε 4.15); 239 (log ε 4.48); 321 (log ε 4.06); 334 (log ε 4.08) (MeOH).

N-Oxide: [161424-22-2]

C₂₁H₃₁NO₂ 329.481

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(E)-: 2-(1E-Dodeceny)-4(1H)-quinolinone. 2-(1E-Dodeceny)-4-hydroxyquinoline

C₂₁H₂₉NO 311.466

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(Z)-: 2-(1Z-Dodeceny)-4(1H)-quinolinone. 2-(1Z-Dodeceny)-4-hydroxyquinoline

C₂₁H₂₉NO 311.466

Metab. of *Pseudomonas aeruginosa*.

NH-form

N-Me: 2-Dodecyl-1-methyl-4(1H)-quinolinone

[182055-94-3]

C₂₂H₃₃NO 327.509

Trace alkaloid from fruits of *Evodia rutaecarpa*. Cryst. Mp 75-76°. λ_{\max} 212 (log ε 4.44); 238 (log ε 4.48); 321 (log ε 4.16); 333 (log ε 4.17) (EtOH).


Tang, Y.-Q. *et al.*, *Phytochemistry*, 1996, **43**, 719-722 (*N-Me, isol, pmr, cmr, ms*)

Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur, oxide, 1',2'-didehydro*)

Agier, C. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 698-703 (*isol, pmr, cmr, ms*)

3-Dodecylpyridine **D-899**

[59652-37-8]




C₁₇H₂₉N 247.423

Isol. from egg-tar. Bp_{0.8} 141°. λ_{\max} 258; 264; 271 (MeOH) (Berdy).

- Tsuji, K. *et al.*, *Yakugaku Zasshi*, 1976, **96**, 479-483; *CA*, **85**, 99075t (*isol, synth*)
 Kozintsev, S.I. *et al.*, *Dokl. Akad. Nauk BSSR*, 1990, **34**, 620-622; *CA*, **114**, 42514 (*synth*)

Doisuthine **D-900**
 N-Methylantranilic acid N-methylphenethylamide
 [173867-26-0]



$C_{17}H_{20}N_2O$ 268.358
 Alkaloid from leaves of *Glycosmis ovoides*.


N^{amide}-*De-Me*: **Glycoamide A**. N-Methylantranilic acid phenethylamide
 $C_{16}H_{18}N_2O$ 254.331
 Alkaloid from *Glycosmis cochinchinensis*. Oil. λ_{max} 211 (log ϵ 4.43); 256 (log ϵ 4); 342 (log ϵ 3.65) (MeOH).

4'-Methoxy: **Methoxydoisuthine**. N-Methylantranilic acid N-methyl-4-methoxyphenethylamide
 [173907-01-2]
 $C_{18}H_{22}N_2O_2$ 298.384
 From leaves of *Glycosmis ovoidea*.

4'-Methoxy, *N*^{amide}-*de-Me*: **Glycoamide B**. N-Methylantranilic acid 4-methoxyphenethylamide
 $C_{17}H_{20}N_2O_2$ 284.357
 Constit. of *Glycosmis cochinchinensis*. Oil. λ_{max} 219; 259; 279; 341 (MeOH).

- Hofer, O. *et al.*, *Annalen*, 1995, 1789 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1491-1493 (*Glycoamides*)

Dolabellin **D-901**
 [165467-49-2]



$C_{24}H_{32}Cl_2N_2O_8S_2$ 611.563
 Isol. from *Dolabella auricularia* and *Lyngbya majuscula*. Cytotoxic. Oil. Sol. MeOH, EtOAc; poorly sol. hexane. $[\alpha]_D^{28}$ -7.3 (c, 0.34 in $CHCl_3$). λ_{max} 204 (ϵ 29400); 236 (ϵ 14600) (MeOH).


Sone, H. *et al.*, *J.O.C.*, 1995, **60**, 4774-4781 (*isol, synth uv, ir, pmr, cmr*)

Dolastatin 1 **D-902**
 [79394-15-3]

Thiazole-peptide antibiotic. Struct. unknown. Isol. from *Dolabella auricularia* and *Aplysia pulmonica*. Cytotoxic agent. Powder. λ_{max} 218 (MeOH).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 482-485

Dolastatin I **D-903**
 [192800-84-3]



$C_{24}H_{32}N_6O_5S$ 516.62
 Isol. from *Dolabella auricularia*. Cytotoxic agent. Amorph. powder. $[\alpha]_D^{30}$ -50 (c, 0.06 in $CHCl_3$). λ_{max} 198 (ϵ 27000); 220 (ϵ 20000) (MeCN).


Sone, H. *et al.*, *Tetrahedron*, 1997, **53**, 8149-8154 (*isol, uv, ir, pmr, cmr*)
 Kigoshi, H. *et al.*, *Tetrahedron*, 1999, **55**, 12301-12308 (*synth*)

Dolastatin 2 **D-904**
 [79394-16-4]

Thiazole-peptide antibiotic. Struct. unknown. Isol. from *Dolabella auricularia* and *Aplysia pulmonica*. Cytotoxic agent. Powder. λ_{max} 217 (MeOH).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 482-485

Dolastatin 3 **D-905**
 [80387-90-2]



R = CH_3

$C_{29}H_{40}N_8O_6S_2$ 660.817
 Cyclic peptide antibiotic. Isol. from the Indian Ocean sea hare *Dolabella auricularia*, sponge *Aplysia pulmonica* and *Lyngbya majuscula*. Shows potent anti-neoplastic props. Inhibitor of HIV 1 integrase. Amorph. solid. Mp 133-137°. $[\alpha]_D^{25}$ -35.5 (c, 0.09 in MeOH). λ_{max} 206 (ϵ 13940); 208; 238 (ϵ 8960) (MeOH) (Berdy).

Homologue (R = $-CH_2CH_3$): **Homodolastatin 3**

[261373-24-4]
 $C_{30}H_{42}N_8O_6S_2$ 674.844
 Isol. from *Lyngbya majuscula*. Amorph. solid. λ_{max} 240 (ϵ 7000) (MeOH).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 482-485 (*isol*)
 Pettit, G.R. *et al.*, *J.A.C.S.*, 1982, **104**, 905-907; 1987, **109**, 7581-7582 (*uv, ir, pmr, cmr, ms, struct, bibl, synth*)
 Schmidt, U. *et al.*, *Angew. Chem., Int. Ed.*, 1984, **23**, 725-726 (*struct*)


- Hamada, Y. *et al.*, *Tet. Lett.*, 1984, **25**, 5303-5306 (*synth*)
 Bredenkamp, M.W. *et al.*, *Annalen*, 1990, 871-875 (*synth, conformn*)
 Holzapfel, C.W. *et al.*, *Tetrahedron*, 1990, **46**, 649-660 (*synth*)
 Pettit, G.R. *et al.*, *Tetrahedron*, 1993, **49**, 9151
 Mitchell, S.S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 279-282 (*Homodolastatin 3*)

Dolastatin 4 **D-906**
 [79394-18-6]

Thiazole-peptide antibiotic. Struct. unknown. Isol. from *Dolabella auricularia* and *Aplysia pulmonica*. Cytotoxic agent. Powder.

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 482-485

Dolastatin 10 **D-907**
 NSC 376128
 [110417-88-4]



R = H

$C_{42}H_{68}N_6O_6S$ 785.102
 Peptide antibiotic. Isol. from *Dolabella auricularia* and *Symploca* sp. VP642. Antitumour and antimetabolic agent. Shows fungicidal activity. Tubulin polymerisation inhibitor. Amorph. powder (MeOH/ CH_2Cl_2). Sol. MeOH, $CHCl_3$. Mp 107-112°. $[\alpha]_D^{29}$ -68 (c, 0.01 in MeOH). λ_{max} 203 (log ϵ 4.39); 216 (ϵ 20200); 242 (ϵ 3610); 245 (log ϵ 3.63) (MeOH) (Derep).

Homologue (R = CH_3): **Symplostatin 1**


[212007-18-6]
 $C_{43}H_{70}N_6O_6S$ 799.129
 Isol. from the cyanobacterium *Symploca hydnoides*. Antitumour agent. Microtubule assembly inhibitor. $[\alpha]_D$ -45 (c, 1.6 in MeOH). λ_{max} 209 (ϵ 20420); 245 (ϵ 5430) (MeOH).

Pettit, G.R. *et al.*, *J.A.C.S.*, 1987, **109**, 6883-6885 (*isol, struct*)
 Pettit, G.R. *et al.*, *J.A.C.S.*, 1989, **111**, 5463-5465 (*synth, abs config*)
 Shioiri, T. *et al.*, *Tetrahedron*, 1993, **49**, 1913-1924 (*synth*)
 Pettit, G.R. *et al.*, *Tetrahedron*, 1993, **49**, 9151-9170 (*isol, bibl*)
 Pettit, G.R. *et al.*, *J.O.C.*, 1994, **59**, 6127-6130 (*synth, cryst struct*)
 Roux, F. *et al.*, *Tetrahedron*, 1994, **50**, 5345-5360 (*synth*)
 Alattia, T. *et al.*, *Tetrahedron*, 1995, **51**, 2593-2604 (*pmr, cmr, conformn*)
 Pettit, G.R. *et al.*, *J.C.S. Perkin 1*, 1996, 853-858; 859-863 (*synth*)
 Pettit, R.K. *et al.*, *Antimicrob. Agents Chemother.*, 1998, **42**, 2961-2965 (*Dolastatin 10, activity*)
 Harrigan, G.G. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1075-1077 (*Symplostatin 1*)
 Munro, M.H.G. *et al.*, *J. Biotechnol.*, 1999, **70**, 15-25 (*rev*)

- Vaishampayan, U. *et al.*, *Clin. Cancer Res.*, 2000, **6**, 1293-1301; 4205-4208 (*pharmacol*)
 Luesch, H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 907-910 (*isol, activity, Symplostatin I*)
 Hoffman, M.A. *et al.*, *Gynecol. Oncol.*, 2003, **89**, 95-98 (*clin trial*)
 Mooberry, S.L. *et al.*, *Int. J. Cancer*, 2003, **104**, 512-521 (*Symplostatin I, pharmacol*)

Dolastatin 18**D-908**

[190181-65-8]


C₃₅H₄₆N₄O₄S 618.839

Isol. from *Dolabella auricularia*. Cytotoxic agent. Powder. [α]_D²⁵ -2.3 (c, 0.1 in MeOH).

- Pettit, G.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 827-832 (*isol, pmr, cmr*)
 Pettit, G.R. *et al.*, *J.O.C.*, 2004, **69**, 4019-4022 (*synth*)

Dolastatin E**D-909**

[165967-00-0]



Absolute configuration

C₂₁H₂₆N₆O₄S₂ 490.606


Isol. from the sea hare *Dolabella auricularia*. Cytotoxic agent. Powder. [α]_D²⁷ -22 (c, 0.2 in MeOH). λ_{max} 205 (ε 26600); 213 (sh) (ε 23800); 240 (sh) (ε 13800) (MeOH) (Derep).

[170720-11-3, 170720-12-4]

- Ojika, M. *et al.*, *Tet. Lett.*, 1995, **36**, 5057-5058; 5059-5062 (*isol, synth, pmr, cmr, abs config*)

Dolichocurine**D-910**

17,18-Deepoxycaracurine II, 9CI
 [69347-05-3]

C₃₈H₄₀N₄O 568.76

- Minor alkaloid from *Strychnos dolichochothyrsa* (Loganiaceae). Mp 300° (as hydrochloride). Stereochem. not establ., that shown is probable by analogy with Caracurine II, C-111.
 Verpoorte, R. *et al.*, *Planta Med.*, 1982, **44**, 21-27 (*isol, ir, ms, pmr, struct*)


Dolichothine**D-911**

[69345-27-3]

- Dimeric caracurine-type alkaloid cooccurring with Dolichocurine, D-910. Struct. unknown. Minor alkaloid from *Strychnos dolichochothyrsa* (Loganiaceae).
 Verpoorte, R. *et al.*, *Planta Med.*, 1982, **44**, 21-27

Domoic acid**D-912**

2-Carboxy-4-(5-carboxy-1-methyl-1,3-hexadienyl)-3-pyrrolidineacetic acid, 9CI
 [14277-97-5]

C₁₅H₂₁NO₆ 311.334

- Constit. of the red algae *Chondria armata* and *Alsidium corallinum*. Also isol. from toxin contaminated edible mussels (*Mytilus edulis*) containing diatom *Nitzschia multiseries* (formerly *Nitzschia pungens*). Ionotropic glutamate (kainate) receptor agonist. Neurotoxin. Amnesic shellfish poison. Vermifuge, insecticide. Needles + 2H₂O. Sol. H₂O, AcOH; fairly sol. MeOH, EtOH; poorly sol. Me₂CO, C₆H₆. Mp 213° dec. [α]_D¹² -109.6 (H₂O). λ_{max} 242 (ε 24300) (H₂O at pH 2) (Derep). λ_{max} 242 (ε 26100) (H₂O pH 7) (Derep). λ_{max} 242 (ε 17500) (H₂O) (Berdy).

▶ Toxic, LD₅₀ 10mg/kg (mice). UX9665100**(1'E)-Isomer: Isodomoic acid E**

[126872-96-6]

C₁₅H₂₁NO₆ 311.334

Isol. from mussel (*Mytilus edulis*). [α]_D²⁵ -19.5 (H₂O).

(3'Z)-Isomer: Isodomoic acid D

[101977-26-8]

C₁₅H₂₁NO₆ 311.334

Isol. from mussel (*Mytilus edulis*). [α]_D²⁵ -72 (H₂O). λ_{max} 213 (ε 5030) (MeOH) (Berdy).

(1'E,3'Z)-Isomer: Isodomoic acid F

[127761-30-2]

C₁₅H₂₁NO₆ 311.334

Isol. from mussel (*Mytilus edulis*). [α]_D²⁵ -85 (H₂O).

5'-Epimer: 5'-Epidomoic acidC₁₅H₂₁NO₆ 311.334


Isol. from mussel and contaminating microalga *Nitzschia pungens*. [α]_D²⁴ -48.8 (c, 0.18 in H₂O). λ_{max} 242 (no solvent reported).

- Daigo, K. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 350; 353; 356; 365; *CA*, **53**, 14218d (*isol, algae*)
 Takemoto, T. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 874 (*pmr, struct*)
 Impellizzeri, G. *et al.*, *Phytochemistry*, 1975, **14**, 1549 (*isol, algae*)
 Ohfuné, Y. *et al.*, *J.A.C.S.*, 1982, **104**, 3511-3513 (*synth, struct*)

- Thibault, P. *et al.*, *Biomed. Mass Spectrom.*, 1989, **18**, 373 (*ms*)
 Wright, J.L.C. *et al.*, *Can. J. Chem.*, 1989, **67**, 481; 1990, **68**, 22 (*Domoic acid, Isodomoic acids, isol, mussels*)
 Falk, M. *et al.*, *Can. J. Chem.*, 1989, **67**, 1421 (*uv*)
 Tryphonas, L. *et al.*, *Toxicol. Pathol.*, Pt. 2, 1990, **18**, 165 (*rev, tox*)
 Walter, J.A. *et al.*, *Can. J. Chem.*, 1992, **70**, 1156 (*pmr, cmr*)
 Walter, J.A. *et al.*, *Can. J. Chem.*, 1994, **72**, 430-436 (*5'-Epidomoic acid*)
 Parsons, A.F. *et al.*, *Tetrahedron*, 1996, **52**, 4149-4174 (*rev*)
Food Sci. Technol., Seafood and Freshwater Toxins, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (*revis*)
 Clayden, J. *et al.*, *Tetrahedron*, 2005, **61**, 5713-5724 (*rev*)

Domoilactone A**D-913**

[101899-46-1]

C₁₅H₂₁NO₇ 327.333

Isol. from the red alga *Chondria armata*. λ_{max} 200 (ε 5700) (MeOH) (Derep).

4,5-Diepimer: Domoilactone B

[101977-27-9]


C₁₅H₂₁NO₇ 327.333

From *Chondria armata*. λ_{max} 200 (ε 5700) (MeOH) (Derep).

- Maeda, M. *et al.*, *Tet. Lett.*, 1987, **28**, 633 (*isol, struct*)

Donasine**D-914**

[1017237-81-8]


C₂₃H₂₈N₄O₃ 408.499

Alkaloid from the rhizomes of *Arundo donax*. Cryst. (CH₂Cl₂/MeOH). Mp 201-202°. λ_{max} 215; 278 (MeOH).

- Jia, A.-L. *et al.*, *J. Asian Nat. Prod. Res.*, 2008, **10**, 105-109 (*isol, pmr, cmr, cryst struct*)

Donaxanine**D-915**

[178493-90-8]


C₁₂H₁₄N₂O₂ 218.255

Alkaloid from *Arundo donax*, also obt. by condensation of Donaxaridine, D-916 with formaldehyde. Cryst. (Me₂CO). Mp 162-164°.

Khuzhaev, V.U. et al., *Khim. Prir. Soedin.*, 1995, **31**, 728-730; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 610-611 (isol, ir, pmr, struct, synth)

Donaxaridine**D-916**

[62209-18-1]



(R)-form

C₁₁H₁₄N₂O₂ 206.244

Struct. revised in 1995. A synthesis reported in 1997 was of the former struct. shown to be incorrect.

(R)-formN²-Me: **Chimonamidine**C₁₂H₁₆N₂O₂ 220.271

Alkaloid from the seeds of *Chimonanthus praecox*. Amorph. powder. [α]_D¹⁹ -12.6 (c, 0.06 in EtOH) (nat.). [α]_D¹⁹ -178 (synthetic). Partial racemate. Obt. as a mixt. of enantiomers with R-isomer predominant. λ_{max} 202 ; 246 ; 300 (MeOH).

(S)-formN²-Me: [α]_D +171.**(±)-form**

Alkaloid from *Arundo donax* (Poaceae). Also formed by treatment of Donaxarine, D-917 with KOH. Mp 175-176°.

N-Ac: Mp 185-186°.

N²-Me:

Needles (EtOAc). Mp 213-215°.

Ubaidullaev, K.A. et al., *Khim. Prir. Soedin.*, 1976, **12**, 553; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 499 (Donaxaridine, isol, uv, ir, pmr, ms)

Khuzhaev, V.U. et al., *Khim. Prir. Soedin.*, 1995, **31**, 720-727; 1996, **32**, 217-220; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 604-609; 1996, **32**, 190-193 (Donaxaridine, *cryst struct, ms*)


Khuzhaev, V.U. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 516-517 (isol, pmr, cmr)

Takayama, H. et al., *Tetrahedron*, 2004, **60**, 893-900 (Chimonamidine, *isol, synth, pmr, cmr*)

Kawasaki, T. et al., *Tetrahedron*, 2004, **60**, 3493-3503 (*synth*)

Donaxarine**D-917**

[62209-19-2]



Relative Configuration


C₁₃H₁₆N₂O₂ 232.282

Struct. revised in 1995. Alkaloid from *Arundo donax* (Poaceae). Mp 218-220°. Opt. inactive.

Khuzhaev, V.U. et al., *Khim. Prir. Soedin.*, 1995, **31**, 720-727; 1996, **32**, 217-220; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 604-609; 1996, **32**, 190-193 (*cryst struct, ms*)

Donglingine**D-918**

[955938-88-2]



Absolute Configuration

C₁₅H₁₉N₃O₅ 321.332

Alkaloid from the aerial parts of *Rabdosia rubescens*.

Feng, W.-S. et al., *Zhongguo Tianran Yaowu*, 2007, **5**, 92-95; *CA*, **147**, 498887v (*isol, pmr, cmr*)

Donine**D-919**

Diethyl 4,4'-(methylenediimino)bisbenzoate, 9CI
[74763-68-1]


C₁₉H₂₂N₂O₄ 342.394

Alkaloid from *Arundo donax* (Poaceae). Cryst. (MeOH). Mp 126-128° (natural) Mp 189° (synthetic). λ_{max} 201 (log ε 4.76); 246 (log ε 4.62) (no solvent reported).

Giumanini, A.G. et al., *J. Phys. Chem.*, 1987, **329**, 1087-1103 (*synth, ir, pmr, ms*)
Khuzhaev, V.U. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 194-197 (*isol, uv, pmr, cmr*)

Dopamine, BAN, INN**D-920**

4-(2-Aminoethyl)-1,2-benzenediol, 9CI.
2-(3,4-Dihydroxyphenyl)ethylamine. 3,4-Dihydroxyphenethylamine. Hydroxytyramine. Oxytyramine. Cardiosteril. Dopastat. Intropin. NSC 169105. ASL 279.
Many other names
[51-61-6]

C₈H₁₁NO₂ 153.18

Occurs in animals, esp. in the brain and nervous system, and in several higher plants, such as broom (*Cytisus scoparius*), banana (*Musa sapientum*) and *Hermidium alipes* (preferred genus name *Mirabilis*), and in the marine alga *Monostroma fuscum*. Also isol. from marine animals (e.g. *Metridium* sp.). Adrenergic, sympathomimetic agent.

Antihypotensive, cardiotonic, antiparkinsonian agent. Central neurotransmitter and precursor of noradrenaline. Log P 0.02 (calc). Autoxidises very readily as free base.

▶ LD₅₀ (rat, ipr) 163 mg/kg. Exp. reprod. and teratogenic effects. UX1088000

Hydrochloride: **Dopamine hydrochloride**, JAN, USAN
[62-31-7]

Mp 240-241° dec. (>220° dec.).

▶ LD₅₀ (rat, ipr) 597 mg/kg. Exp. reprod. and teratogenic effects. UX1092000

Hydrobromide: [645-31-8]

Mp 218-220°.

▶ SH2290150

O³-β-D-Glucopyranoside: **Dopamine 3-glucoside**

[50908-96-8]

C₁₄H₂₁NO₇ 315.322

Alkaloid from seeds of *Entada pursaetha* (Fabaceae). Semicryst. brown solid. [α]_D²⁰ -62 (c, 2 in H₂O).

O⁴-β-D-Glucopyranoside: **Dopamine 4-glucoside**

[50909-01-8]

C₁₄H₂₁NO₇ 315.322

Isol. from *Manduca sexta* and *Papilio xuthus*. Cryst. [α]_D²⁰ -66 (c, 2 in H₂O).

N-Ac: N-(3,4-Dihydroxyphenethyl)acetamide, 8CI

[2494-12-4]

C₁₀H₁₃NO₃ 195.218

Constit. of the cast-off shells of the cicada, *Cryptotympana* sp. Precursor of sclerotised insect cuticle. No phys. props. reported. Mp 78-80°. Component of Zentai.

N-Ac, 3-O-β-glucopyranoside: [86512-21-2]

C₁₆H₂₃NO₈ 357.36

Isol. from insects *Mantis religiosa*, *Tenodera angustipennis*, *Manduca sexta* and *Statilia maculata*. No phys. props. reported.

N-Ac, 4-O-β-D-glucopyranoside: [29625-66-9]

C₁₆H₂₃NO₈ 357.36

Isol. from insects *Calliphora erythrocephala*, *Bombyx mori* and *Manduca sexta*.

N-Carboxyacetyl, 3-O-β-D-glucopyranoside: **N-Malonyldopamine 3-glucoside**

[86502-98-9]

C₁₇H₂₃NO₁₀ 401.369

Isol. from *Hierodula patellifera* and *Tenodera angustipennis*. No phys. props. reported.

N-[3-(Carboxyacetylamino)propanoyl], 3-O-β-D-glucopyranoside: **N-(N-Malonyl-β-alanyl)dopamine 3-glucoside**

[96735-98-7]

C₂₀H₂₈N₂O₁₁ 472.448

Isol. from *Hierodula patellifera* and *Tenodera angustipennis*. No phys. props. reported.

N-(4-Hydroxy-E-cinnamoyl): [103188-46-1]

C₁₇H₁₇NO₄ 299.326

Constit. of *Atraphaxis spinosa*.

N-(3,4-Dihydroxy-E-cinnamoyl): **Caffeodymine**
 [103188-49-4]
 [105955-00-8]
 $C_{17}H_{17}NO_5$ 315.325
 Constit. of *Theobroma cacao*, *Capsicum annuum* (bell pepper) and *Lycium chinense* (wolfberry).
 Cyclooxygenase inhibitor. No phys. props. reported.

N-(4-Methoxy-Z-cinnamoyl): N-(4-Methoxycinnamoyl)dopamine. **Lyciumide A**
 [245343-48-0]
 $C_{18}H_{19}NO_4$ 313.352
 Alkaloid from *Lycium barbarum*. Gum.

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): N-trans-Feruloyldopamine. **Tuberosine B†**
 [142350-99-0]
 $C_{18}H_{19}NO_5$ 329.352
 Alkaloid from *Allium tuberosum* (Chinese chives) and *Atraphaxis spinosa*.

N-Jasmonoyl: **N-Jasmonoyldopamine**
 $C_{20}H_{27}NO_4$ 345.438
 Constit. of the flowers of *Vicia faba*.

N-Me: see Epinine, E-98

N,N-Di-Me: 4-[2-(Dimethylamino)ethyl]-1,2-benzenediol. N,N-Dimethyl-3,4-dihydroxyphenethylamine. N,N-Dimethyldopamine
 [21581-37-3]
 [13075-91-7, 50309-53-0]
 $C_{10}H_{15}NO_2$ 181.234
 Alkaloid from *Acacia rigidula*. Cryst. (EtOH). Mp 156°.

N-Tri-Me: see Coryneine, C-690

N-Benzyl:
 $C_{15}H_{17}NO_2$ 243.305
 Cryst. powder (EtOH/Et₂O) (as hydrochloride). Mp 87° Mp 180° (double Mp) (hydrochloride).

O³-Me: see 4-Hydroxy-3-methoxyphenethylamine, H-566

Di-Me ether: see 3,4-Dimethoxyphenethylamine, D-718

Dibenzyl ether: [1699-56-5]
 $C_{22}H_{23}NO_2$ 333.429
 Cryst. (EtOAc/EtOH) (as hydrochloride). Mp 133° (hydrochloride). CAS no. refers to hydrochloride.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 1294A; 1294B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 622B; 622C (nmr)

Buck, J.S. et al., *J.A.C.S.*, 1931, 53, 2192-2200 (N-benzyl, N-benzyl O,O-di-Me)

Forbes, E.J. et al., *J.C.S.*, 1955, 3926-3932 (O,O-dibenzyl)

Waalkes, T.P. et al., *Science (Washington, D.C.)*, 1958, 127, 648-650 (*Musa sapientum* constit)

Karlson, P. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1962, 327, 86-94 (N-Ac, N-Ac-4-glucoside, isol)

Keppe, T. et al., *J. Med. Chem.*, 1965, 8, 368-374 (uv)

Tocher, R.D. et al., *Can. J. Bot.*, 1966, 44, 605-608 (*Monostroma fuscum* constit)

Seiler, N. et al., *Fresenius' Z. Anal. Chem.*, 1970, 252, 127-136 (ms, N-Ac)

Lundström, J. et al., *Acta Chem. Scand.*, 1971, 25, 3489-3499 (biosynth)

Milne, G.W. et al., *Anal. Chem.*, 1973, 45, 1952-1954 (ms)

Larsen, P.O. et al., *Phytochemistry*, 1973, 12, 2243-2247 (synth, glucosides)

Lambert, F. et al., *Org. Magn. Reson.*, 1975, 7, 266-273 (pmr)

Lenicque, P.M. et al., *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1977, 56, 31-34 (Metridium senile constit)

Smith, T.A. et al., *Phytochemistry*, 1977, 16, 9-18 (occur, bibl)

Scratchley, G.A. et al., *J. Chromatogr.*, 1979, 169, 313-319 (hplc)

Carter, J.E. et al., *Anal. Profiles Drug Subst.*, 1982, 11, 257-272 (rev, uv, ir, pmr, cmr, ms, anal)

Yago, M. et al., *Insect Biochem.*, 1984, 14, 7-9; 487-489 (N-malonyl-3-glucoside, N-malonyl-β-alanyl 3-glucoside)

Quantitative Analysis of Catecholamines and Related Compounds, (ed. Krstulovic, A.M.), Ellis Horwood, 1986, (book)

Carey, R.M. et al., *Recent Prog. Horm. Res.*, 1986, 42, 251-296 (rev)

Peripheral Actions of Dopamine, (eds. Bell, C. et al.), Macmillan, 1988, (book)

Eur. Pat., 1989, ((Mitsui))333 522; CA, 112, 138764w (synth, pmr, N-Ac)

Ishizaki, Y. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1990, 97, 563 (isol, 4-glucoside)

Mueller, D.D. et al., *Bioconj. Chem.*, 1993, 4, 47-53 (pmr, cmr)

Dopamine Receptors and Transporters, (ed. Niznik, H.B.), M. Dekker, 1994, (book)

Volin, P. et al., *J. Chromatogr., B*, 1994, 655, 121-126 (hplc)

El-Gamal, A.A. et al., *Nat. Med. (Tokyo)*, 1994, 48, 304-306; CA, 122, 209765k (N-cinnamoyl derivs)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 1113 (synonyms)

CNS Neurotransmitters and Neuromodulators: Dopamine, (ed. Stone, T.W.), CRC Press, 1996, (book)

Martindale, The Extra Pharmacopoeia, 31st edn., Pharmaceutical Press, 1996, 1574

Hay, M. et al., *J. Chromatogr., B*, 1997, 703, 15-23 (hplc)

Zou, C. et al., *Chin. Chem. Lett.*, 1999, 10, 131-132 (Lyciumide A)

Noda, N. et al., *Chem. Pharm. Bull.*, 2000, 48, 1749-1752 (isol, N-Ac)

Merck Index, 13th edn., 2001, No. 3455 (bibl)


Kramell, R. et al., *J. Nat. Prod.*, 2005, 68, 1345-1349 (N-Jasmonoyldopamine)

Park, J.B. et al., *J. Agric. Food Chem.*, 2007, 55, 2171-2175 (caffedymine, biol, bibl)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, DYC400; DYC600

Dopaxanthin D-921

4-[[[1-Carboxy-2-(3,4-dihydroxyphenyl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI
 [71199-31-0]



$C_{18}H_{18}N_2O_8$ 390.349


Isol. from *Glottiphyllum longum*. Also occurs in *Conophytum* spp., *Portulaca* spp. and in the Cactaceae. Intermed. in biosynth. of betacyanins from DOPA. Amorph. orange solid. λ_{max} 488 (ε 41800) (no solvent reported).

Impellizzeri, G. et al., *Phytochemistry*, 1973, 12, 2293 (isol)

Wyller, H. et al., *Helv. Chim. Acta*, 1979, 62, 1330-1339 (synth, biosynth)

Doriasenine D-922

[120091-13-6]




$C_{18}H_{25}NO_6$ 351.399

Diester of Retronecine in T-188. Alkaloid from *Senecio doria* (Asteraceae). Oil. $[\alpha]_D^{20} +6$ (EtOH).

Röder, E. et al., *Phytochemistry*, 1988, 27, 4000 (isol, ir, pmr, cmr, ms, struct)

Dorriginocin A D-923

[158446-29-8]



$C_{27}H_{41}NO_8$ 507.623

Glutarimide antibiotic. Prod. by *Streptomyces platensis* ssp. *rosaceus*. Strictly a shunt metab. of Isomigrastatin, I-245. Antifungal agent, yeast mating pheromone. Oil. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{20} +91$ (c. 0.1 in MeOH). λ_{max} 209 (ε 11000) (MeOH/NaOH). λ_{max} 205 (ε 9300) (MeOH).

13-Epimer: 13-Epidorriginocin A

$C_{27}H_{41}NO_8$ 507.623


Prod. by *Streptomyces platensis*. Strictly a shunt metab. of Isomigrastatin, I-245. $[\alpha]_D^{25} +14.6$ (c. 0.13 in MeOH).

Karwowski, J.P. et al., *J. Antibiot.*, 1994, 47, 862-869; 870-874; 875-880 (isol, pmr, cmr)

Ju, J. et al., *J.A.C.S.*, 2005, 127, 1622-1623 (biosynth)

Dorriginocin B D-924

[158446-30-1]



$C_{27}H_{41}NO_8$ 507.623


Glutarimide antibiotic. Prod. by *Streptomyces platensis* ssp. *rosaceus*. Strictly a shunt metab. of Isomigrastatin, I-245. Antifungal agent. Oil. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{20} +16$ (c,

0.3 in MeOH). λ_{\max} 209 (ϵ 11000) (MeOH/NaOH). λ_{\max} 205 (ϵ 9300) (MeOH).

Karwowski, J.P. *et al.*, *J. Antibiot.*, 1994, **47**, 862-869; 870-874; 875-880 (*isol*, *pmr*, *cmr*)
Ju, J. *et al.*, *J.A.C.S.*, 2005, **127**, 1622-1623 (*biosynth*)

Doryflavine

[54140-26-0]



$C_{16}H_{11}NO_4$ 281.267

Alkaloid from the bark of *Doryphora sassafras* (Monimiaceae). Golden-yellow plates (MeOH). Mp 339°. For a tentative struct. see Mix, *et al.* λ_{\max} 230; 251; 260; 277; 290; 320; 400 (MeOH) (Berdy). λ_{\max} 244; 263; 295; 345; 430 (MeOH/NaOH) (Berdy).

Tri-Ac:

Needles (MeOH/EtOH). Mp 218-219°.

Di-Me ether: Mp 219-221°.

Chen, C.R. *et al.*, *J. Nat. Prod.*, 1974, **37**, 493-500 (*isol*, *uv*, *ir*, *pmr*, *ms*)

Mix, D.B. *et al.*, *J. Nat. Prod.*, 1982, **45**, 657-666 (*rev*)

Doryphorine

[1399-14-0]


$C_{18}H_{21}NO_4$ 315.368

Struct. unknown. Alkaloid from the bark of *Doryphora sassafras* (Monimiaceae). Amorph. grey powder. Mp 115-117°.

Petrie, J.M. *et al.*, *Proc. Linn. Soc. N.S.W.*, 1913, **37**, 139; *CA*, **7**, 2092

Dovyalicin A**D-927**

1-[4-(Benzoylamino)butyl]hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1H)-one [597578-55-7]



Absolute Configuration

$C_{24}H_{31}N_3O_2$ 393.528

Alkaloid from *Dovyalis abyssinica* and *Dovyalis macrocalyx*. Gum. $[\alpha]_D^{25}$ -12.4 (c, 0.45 in $CHCl_3$).

N^4 -Debenzoyl, N^4 -Ac: **Dovyalicin B**

[597578-56-8]

$C_{19}H_{29}N_3O_2$ 331.457

Alkaloid from *Dovyalis abyssinica* and *Dovyalis macrocalyx*. Gum. $[\alpha]_D^{25}$ -23.5 (c, 0.39 in $CHCl_3$).

N^5 -De-Me: **Dovyalicin E**

[911697-89-7]

$C_{23}H_{29}N_3O_2$ 379.501

Alkaloid from *Dovyalis abyssinica*.


Gum. $[\alpha]_D^{25}$ -11.7 (c, 0.42 in $CHCl_3$).

Staerk, D. *et al.*, *Org. Lett.*, 2003, **5**, 2793-2796 (*isol*, *pmr*, *cmr*)

Rasmussen, B. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1300-1304 (*isol*)

Dovyalicin D

[597578-58-0]



$C_{25}H_{31}N_3O_4$ 437.538

Alkaloid from *Dovyalis macrocalyx*.

Gum. Racemic.

Deoxy: **Dovyalicin C**

[597578-57-9]

$C_{25}H_{31}N_3O_3$ 421.538

Alkaloid from *Dovyalis macrocalyx*.


Gum. $[\alpha]_D^{25}$ -1.1 (c, 0.79 in $CHCl_3$).

Staerk, D. *et al.*, *Org. Lett.*, 2003, **5**, 2793-2796 (*Dovyalicins C-D*, *isol*, *pmr*, *cmr*)

Rasmussen, B. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1300-1304 (*Dovyalicin C*, *isol*)

Dragmacidin A**D-929**

3,3'-(1-Methyl-2,5-piperazinediyl)bis[6-bromo-1H-indole], 9CI. 2,5-Bis(6-bromo-3-indolyl)-1-methylpiperazine [128364-31-8]



$C_{21}H_{20}Br_2N_4$ 488.224

Alkaloid from the marine sponges *Hexadella* sp., *Dragmacidon* spp., *Spongosorites* spp. and from tunicate *Didemnum candidum*. Cytotoxic. Shows antifungal, antiviral and antiinflammatory activity. Pale yellow powder.

N^4 -Me: **Dragmacidin B**

[128629-37-8]

$C_{22}H_{22}Br_2N_4$ 502.251

Alkaloid from the sponge *Hexadella* sp. Pale yellow powder.

7'-Bromo, 4'-hydroxy: **Dragmacidin.**

Biemnidin

[114582-72-8]

$C_{21}H_{19}Br_3N_4O$ 583.119

Alkaloid from the deep-water sponge *Dragmacidon* sp. Cytotoxic agent.

Powder. $[\alpha]_D^{25}$ -3 (c, 13.2 in Me_2CO). λ_{\max} 220 (ϵ 52600); 275 (ϵ 11700); 286 (sh) (ϵ 10900); 293 (sh) (ϵ 10100) (MeOH) (Derep).

N -De-Me: 3,3'-(2,5-Piperazinediyl)-bis[6-bromo-1H-indole]. 2,5-Bis(6-bromo-3-indolyl)piperazine. **Dragmacidin C**

[135048-54-3]

$C_{20}H_{18}Br_2N_4$ 474.197

Alkaloid from the marine tunicate *Didemnum candidum*. Opaque glass. $[\alpha]_D$ 0.

Eur. Pat., 1988, 284 337; *CA*, **111**, 50423 (*Biemnidin*)

Kohmoto, S. *et al.*, *J.O.C.*, 1988, **53**, 3116-3118 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Morris, D.A. *et al.*, *Tetrahedron*, 1990, **46**, 715-720 (*isol*, *pmr*, *struct*)

Fahy, E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 564-569 (*Dragmacidin C*)

Jiang, B. *et al.*, *J.O.C.*, 1994, **59**, 6823-6827 (*Dragmacidin, synth*)

Whitlock, C.R. *et al.*, *Tet. Lett.*, 1994, **35**, 371-374 (*Dragmacidin B, synth*)

Kawasaki, T. *et al.*, *Org. Lett.*, 2000, **2**, 3027-3029 (*synth*)

Miyake, F.Y. *et al.*, *Org. Lett.*, 2000, **2**, 3185-3187 (*Dragmacidin B, synth*)


Kawasaki, T. *et al.*, *Tet. Lett.*, 2002, **43**, 4245-4248 (*Dragmacidin B, Dragmacidin C, synth*)

Garg, N.K. *et al.*, *Tet. Lett.*, 2005, **46**, 2423-2426 (*Dragmacidin B, synth*)

Anstiss, M. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 4135-4143 (*synth*)

Dragmacidin D**D-930**

[142979-34-8]



$C_{25}H_{21}BrN_7O_2^{\oplus}$ 531.391

Alkaloid from a deep-water marine sponge *Spongosorites* sp. Exhibits antimicrobial activity. Also inhibits the *in vitro* growth of P388 murine and A589 human lung tumour cell lines, and *in vitro* replication of feline leukaemia virus. Yellow solid. $[\alpha]_D +12$ (c, 0.95 in EtOH). λ_{\max} 213 (ϵ 43000); 270 (ϵ 14400); 383 (ϵ 20700) (EtOH) (Berdy). λ_{\max} 214 (ϵ 54000); 280 (ϵ 16400); 452 (ϵ 19946) (EtOH-HCl) (Berdy).


Wright, A.E. *et al.*, *J.O.C.*, 1992, **57**, 4772 (*isol*, *uv*, *pmr*, *cmr*, *struct*)

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2002, **61**, 660-662 (*isol*, *cmr*)

Garg, N.K. *et al.*, *J.A.C.S.*, 2002, **124**, 13179-13184 (*synth*)

Dragmacidin E**D-931**

[206061-75-8]



Relative Configuration

$C_{25}H_{20}BrN_7O_2$ 530.383

Tautomeric with the hydroxypyrazine form. *Isol.* from the sponge *Spongosorites*


sp. Inhibitor of serine-threonine protein phosphatases. Yellow solid. $[\alpha]_D^{25}$ -34 (c, 0.9 in EtOH). Fluorescent. λ_{max} 211 (ε 46200); 281 (ε 7530); 427 (ε 27700) (EtOH). λ_{max} 427 (ε 26400) (EtOH/HCl). λ_{max} 410 (ε 26100) (EtOH/NaOH).

Capon, R.J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 660-662 (*isol, uv, pmr, cmr*)

Drarmacidin F

D-932

[239121-79-0]



C₂₅H₂₀BrN₇O₃ 546.382

Alkaloid from the sponge *Halicortex* sp. Antiviral agent. Pale yellow solid. Mp > 260°. $[\alpha]_D^{25}$ -159 (c, 0.4 in MeOH). λ_{max} 207 (ε 14000); 274 (ε 6460); 280 (ε 6430); 388 (ε 9600) (MeOH).


Cutignano, A. *et al.*, *Tetrahedron*, 2000, **56**, 3743-3748 (*isol*)

Garg, N.K. *et al.*, *J.A.C.S.*, 2004, **126**, 9552-9553; 2005, **127**, 5970-5978 (*synth*)

Drarmacidonamine A

D-933

3-Carboxyhyrtiomanzamine
[835612-74-3]



C₁₉H₁₇N₄O₄S[⊕] 397.434

Counterion not specified. Charge is delocalised. Related to Hyrtiomanzamine, H-781. Alkaloid from the sponge *Drarmacidon* sp. Amorph. yellow solid. λ_{max} 221 ; 254 ; 317 (MeOH).

10-Deoxo: **Drarmacidonamine B**

[835612-84-5]


C₁₉H₁₉N₄O₃S[⊕] 383.45

Alkaloid from *Drarmacidon* sp. Brownish oil. λ_{max} 222 ; 242 ; 282 (MeOH).

Pedpradab, S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 2113-2116 (*isol*)

Drazepinone

D-934



Relative Configuration

C₁₉H₁₉NO₂ 293.365


Prod. by *Drechslera siccans* isol. from the seeds of *Lolium perenne*. Phytotoxin. Oil. $[\alpha]_D^{25}$ +7.1 (c, 0.2 in CHCl₃). λ_{max} 215 (sh) ; 248 (log ε 4.05); 275 (sh) ; 310 (sh) (MeCN).

Evidente, A. *et al.*, *Phytochemistry*, 2005, **66**, 715-721 (*isol, pmr, cmr, ms*)

Dregamine

D-935

Methyl 19,20-dihydro-3-oxovobasan-17-oate, 9CI. Epitabernaemontanine
[2299-26-5]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Voacanga drepei*, *Tabernaemontana elegans*, *Ervatamia coronaria*, *Ervatamia orientalis* and several other spp. in the Apocynaceae. Antiinflammatory agent. Shows weak local anaesthetic, convulsant and respiratory stimulant props. Cryst. (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 106-108° Mp 137-141° Mp 181-183°. Log P 3.3 (calc). λ_{max} 237 ; 313 (MeOH) (Berdy).

► ZB1635000

Hydrochloride:

Cryst. (EtOH). Mp 249-250° (238-240°).

Methiodide:

Prisms (MeOH). Mp 215-217°. $[\alpha]_D^{23}$ -63.4 (MeOH).

3R-Alcohol: **Dregaminol**

[56691-99-7]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Tabernaemontana elegans* (Apocynaceae).

3R-Alcohol, Me ether: **Dregaminol methyl ether**

[103763-52-6]

C₂₂H₃₀N₂O₃ 370.491

Alkaloid from *Tabernaemontana elegans* (Apocynaceae).

Parent acid: **Dregaminic acid**

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the stems of *Ervatamia flabelliformis*. Powder (MeOH) (as Na salt). $[\alpha]_D^{25}$ +24.1 (c, 0.2 in EtOH) (Na salt). Isol. as Na salt, Sodium dregamine.

20-Epimer: **Tabernaemontanine**. 20-Epi-dregamine

[2134-98-7]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid present in *Ervatamia orientalis*, *Tabernaemontana* spp., *Voacanga globosa*, *Muntafara sessilifolia* (preferred genus name *Tabernaemontana*) and several other spp. in the Apocynaceae. Active against gram-positive bacteria. Cryst. (MeOH or Et₂O). Mp 216-218° (204-206°). $[\alpha]_D^{22}$ -58 (c, 1 in CHCl₃).

20-Epimer, 3R-alcohol: **Tabernaemontaninol**

[56692-00-3]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Tabernaemontana elegans* whole plants (Apocynaceae).

Neuss, N. *et al.*, *Experientia*, 1959, **15**, 414 (*isol, uv, ir*)

Renner, U. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 2186 (*struct*)

Ferrari, G. *et al.*, *Phytochemistry*, 1971, **10**, 439 (*isol, ms, pmr*)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 929 (*cd, abs config*)

Husson, A. *et al.*, *Tetrahedron*, 1973, **29**, 3095 (*config*)

Knox, J.R. *et al.*, *Aust. J. Chem.*, 1975, **28**, 1843 (*struct*)

Talapatra, B. *et al.*, *Phytochemistry*, 1975, **14**, 1652 (*isol*)

Ahond, A. *et al.*, *J.O.C.*, 1976, **41**, 1878 (*cmr*)

Hernandez, N.M.R. *et al.*, *Rev. Cubana Farm.*, 1977, **11**, 249-255 (*activity*)

Kutney, J.P. *et al.*, *J.A.C.S.*, 1978, **100**, 938 (*synth*)


Van der Heijden, R. *et al.*, *Planta Med.*, 1986, 144 (*derivis*)

Liang, S. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 239-243 (*Sodium dregamine*)

Drimentine A

D-936

[204398-90-3]



C₃₂H₄₅N₃O₂ 503.726


Prod. by Actinomycete strain MST-8651. Antibiotic. Cryst. Mp 130-132°. $[\alpha]_D^{25}$ -209.6 (c, 0.006 in CHCl₃). λ_{max} 240 (ε 9445); 297 (ε 4167) (EtOH).

Pat. Coop. Treaty (WIPO), 1998, 98 09 968; CA, **128**, 217535z (*isol, cd, pmr, cmr*)

Drimentine C

D-937

[204398-92-5]



C₃₁H₄₁N₃O₂ 487.684

Prod. by Actinomycete strain MST-8651. Antibiotic. Cryst. Mp 108-110°. $[\alpha]_D^{25}$ -275.9 (c, 0.006 in CHCl₃). λ_{max} 240 (ε 6078); 297 (ε 2157) (EtOH).

3,25-Didehydro: **Drimentine B**


[204398-91-4]

$C_{31}H_{39}N_3O_2$ 485.668
Prod. by Actinomycete strain MST-8651. Antibiotic. Cryst. Mp 158-160°. $[\alpha]_D^{25}$ -195.2 (c, 0.007 in $CHCl_3$). λ_{max} 243 (ε 12420); 273 (sh) (ε 6667); 295 (sh) (ε 4091) (EtOH).

Pat. Coop. Treaty (WIPO), 1998, 98 09 968; CA, 128, 217535z (isol, cd, pmr, cmr)

Drimentine D D-938

[204398-93-6]




$C_{32}H_{45}N_3O_2$ 503.726
Prod. by Actinomycete strain MST-8651. Antibiotic. Cryst. Mp 141-145°. $[\alpha]_D^{25}$ -24.2 (c, 0.006 in $CHCl_3$). λ_{max} 235 (sh) (ε 3146); 273 (ε 1326); 280 (sh) (ε 1236) (EtOH).

Pat. Coop. Treaty (WIPO), 1998, 98 09 968; CA, 128, 217535z (isol, cd, pmr, cmr)

Drimentine E D-939

[204398-94-7]




$C_{31}H_{39}N_3O_2$ 485.668
Prod. by Actinomycete strain MST-8651. Antibiotic. Cryst. Mp 260-262°. $[\alpha]_D^{25}$ +17.6 (c, 0.01 in $CHCl_3$). λ_{max} 226 (sh) (ε 8235); 268 (sh) (ε 5441) (EtOH).

Pat. Coop. Treaty (WIPO), 1998, 98 09 968; CA, 128, 217535z (isol, cd, pmr, cmr)

Drimiopsine A† D-940

2,7,8,9-Tetramethoxy-11H-[1]benzopyrano[4,3-c]isoquinolin-4-ol
[1045704-45-7]



$C_{20}H_{19}NO_6$ 369.373

Alkaloid from *Drimiopsis barteri*.


1-Deoxy, 2-hydroxy: **Drimiopsine B†**

[1045704-46-8]
 $C_{20}H_{19}NO_6$ 369.373
Alkaloid from *Drimiopsis barteri*.

Ngamga, D. et al., Nat. Prod. Commun., 2008, 3, 769-777 (isol, pmr, ms)

Drosopterin D-941

[33466-46-5]



Relative configuration

$C_{15}H_{16}N_{10}O_2$ 368.357
Red eye pigment from *Drosophila melanogaster*. Red cryst. powder. Mp 350° dec. pK_{a1} 9.33; pK_{a2} 8.27; pK_{a3} 1.24; pK_{a4} 0.45 (20°, H_2O). Opt. active. All attempts to crystallise Drosopterin or its stereoisomers have been unsuccessful.

Stereoisomer: **Neodrosopterin**

[56711-41-2]

 $C_{15}H_{16}N_{10}O_2$ 368.357

Pigment from *Drosophila melanogaster*.

Enantiomer: **Isodrosopterin**

[33466-47-6]

 $C_{15}H_{16}N_{10}O_2$ 368.357

Pigment from *Drosophila melanogaster*.

Schlobach, H. et al., Helv. Chim. Acta, 1972, 55, 2518-2525; 2525-2533; 2533-2540; 2541-2545 (isol, uv, ord, cd, pmr, synth)


Rokos, K. et al., Chem. Ber., 1975, 108, 2728-2736 (Neodrosopterin)

Theobald, N. et al., Chem. Ber., 1978, 111, 3385-3402 (uv, pmr, cd, struct)

Pfleiderer, W. et al., J. Het. Chem., 1992, 29, 583-605 (rev)

Drupacine D-942

[49686-57-9]



Absolute configuration

$C_{18}H_{21}NO_5$ 331.368
Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*, *Cephalotaxus fortunei*, *Cephalotaxus hainanensis*, *Cephalotaxus wilsoniana* and *Cephalotaxus sinensis* (Cephalotaxaceae). Mp 70-72°. $[\alpha]_D^{30}$ -137 (c, 0.79 in $CHCl_3$).

[24268-97-1, 125712-90-5]


Powell, R.G. et al., J.O.C., 1974, 39, 676 (isol, uv, ir, ms, pmr, struct)

Weisleder, D. et al., Org. Magn. Reson., 1980, 13, 114 (cmr)

Burkholder, T.P. et al., J.A.C.S., 1990, 112, 9601 (synth)

Drupangtonine D-943

[181133-84-6]




Absolute Configuration

$C_{28}H_{37}NO_9$ 531.602
Ester of Drupacine, D-942. Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*. Strongly inhibits the growth of P-388 leukaemia cells. Oil. $[\alpha]_D$ -24 (c, 0.06 in MeOH). λ_{max} 291 (ε 3020) (MeOH) (Berdy).

Takano, I. et al., Bioorg. Med. Chem. Lett., 1996, 6, 1689 (isol, uv, cd, pmr, cmr, struct)

Dryadodaphnine D-944

6',7'-Dimethoxy-2,2'-dimethylthalam-6,12-diol, 9CI
[22559-06-4]



$C_{36}H_{38}N_2O_6$ 594.706

The alkaloids covered by this entry (2*S*,2'*R*-) belong to the enantiomeric series compared with those covered by Lauberine, L-67 (2*R*,2'*S*-) and are diastereomeric with those covered by Thalmine, T-341 (2*S*,2'-). Alkaloid from the bark of *Dryadodaphne novoguineensis* (Monimiaceae). Foam. $[\alpha]_D$ +390 (MeOH).

*O*⁶-Me: 6-O-Methyldryadodaphnine.

Thalifortine

[129785-17-7]

 $C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the whole plant of *Thalictrum fortunei*. Mp 143-145°. $[\alpha]_D^{25}$ +271.4 (c, 0.37 in MeOH). λ_{max} 215 (log ε 4.61); 280 (log ε 3.91) (MeOH).

*O*¹²-Me: 12-O-Methyldryadodaphnine.

Dryadine

[22559-05-3]

 $C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the bark of *Dryadodaphne novoguineensis* (Monimiaceae). Needles ($CHCl_3$ /MeOH). Mp 249-251° dec. $[\alpha]_D^{15}$ +486 (c, 1.5 in $CHCl_3$).

Baldas, J. et al., Tet. Lett., 1968, 6315 (pmr)


Bick, I.R.C. et al., J.C.S.(C), 1969, 1627 (isol, uv, pmr, ord, ms, struct)

Baldas, J. et al., J.C.S. Perkin 1, 1972, 597 (ms)

Wu, Z. et al., *Zhivw Xuebao (Acta Bot. Sin.)*, 1990, **32**, 210-214; *CA*, **113**, 168973 (Thalifortine)

Dubinidine**D-945**

2-(2,3-Dihydro-4-methoxyfuro[2,3-b]quinolin-2-yl)-1,2-propanediol, 9CI [22964-77-8]



$C_{15}H_{17}NO_4$ 275.304

Alkaloid from *Haplophyllum dubium* leaves (Rutaceae). CNS depressant. Cryst. (Me₂CO). Mp 132-133°. [α]_D^{26.5} -62.95.

Hydrochloride: Mp 195-196°.

Picrate: Mp 155-156°.

2'-Ac: **Dubinine**

[23092-72-0]

$C_{17}H_{19}NO_5$ 317.341

Alkaloid from the leaves of *Haplophyllum dubium* (Rutaceae). Cryst. (MeOH or EtOH). Mp 185-186°. [α]_D -73.1 (Me₂CO).

2'-Ac; hydrochloride: Mp 170-171°.

Di-Ac: Mp 108-109°.

2'-Deoxy: see Platydesmine, P-515


Yunusov, S.Y. et al., *Zh. Obshch. Khim.*, 1955, **25**, 2009; *CA*, **50**, 9435i (isol)

Bessonova, I.A. et al., *Khim. Prir. Soedin.*, 1969, **5**, 29; 1970, **6**, 446; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 22; 1970, **6**, 455 (struct, ms)

Grundon, M.F. et al., *Tet. Lett.*, 1971, 4727 (synth, bibl)

Duguenaine**D-946**

9,10-Dihydro-5H,7H-benzo[f][1,3]dioxolo[6,7]isoquinolo[8,1,2-hij][3,1]benzoxazine, 9CI [80550-24-9]



$C_{19}H_{15}NO_3$ 305.332

Modified aporphine alkaloid. Alkaloid from *Duguetia calycina* (Annonaceae). Cryst. (MeOH). Mp 167-170°.

11-Methoxy: **Duguecalyne**

[80550-25-0]

$C_{20}H_{17}NO_4$ 335.359

Alkaloid from *Duguetia calycina* (Annonaceae). Cryst. (Me₂CO). Mp 202°.


Roblot, F. et al., *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1981, **293**, 373 (uv, pmr, cmr, ms, struct)

Lenz, G.R. et al., *J.C.S. Perkin 1*, 1984, 1273 (synth, Duguenaine)

Atanes, N. et al., *J.O.C.*, 1991, **56**, 2984 (synth, Duguenaine)

Dugespixine**D-947**

[96910-76-8]



$C_{19}H_{17}NO_3$ 307.348


Doubts have been expressed regarding the struct. Alkaloid from the bark of *Duguetia spixiana* (Annonaceae). Amorph.

Debourges, D. et al., *J. Nat. Prod.*, 1985, **48**, 310; 1987, **50**, 664 (isol, uv, ir, pmr, cmr, ms, struct)

Atanes, N. et al., *Heterocycles*, 1987, **26**, 1183

Duguetine**D-948**

7-Hydroxy-1,2-methylenedioxy-9,10-dimethoxyaporphine. 7-Hydroxydicentrine [31984-61-9]



Absolute Configuration

$C_{20}H_{21}NO_5$ 355.39

Alkaloid from an unidentified Brazilian *Duguetia* sp. (Annonaceae). Mp 149-150°. [α]_D -41 (c, 1 in EtOH).

7-Epimer: **Dasymachaline**

[105814-54-8]

$C_{20}H_{21}NO_5$ 355.39

Alkaloid from the leaves of *Desmos dasymachus*. Amorph. [α]_D²⁶ -47 (c, 0.34 in CHCl₃). Rather unstable, dec. on standing.

7-Epimer, α -N-oxide: **Dasymachaline α -N-oxide**

[374106-25-9]

$C_{20}H_{21}NO_6$ 371.389

Alkaloid from the roots of *Desmos dasymachus*. Trypanocidal agent.

7-Epimer, N-de-Me: **N-Demethylasymachaline**. 7-Hydroxynordicentrine

[209323-83-1]

$C_{19}H_{19}NO_5$ 341.363

Alkaloid from *Dicentra peregrina*.

Casagrande, C. et al., *Farmaco, Ed. Sci.*, 1970, **25**, 442; *CA*, **73**, 38472a (*Duguetine*)


Chan, K.C. et al., *Phytochemistry*, 1986, **25**, 1999 (*Dasymachaline*)

Konishi, T. et al., *Nat. Med. (Tokyo)*, 1998, **52**, 47-53 (*7-Hydroxynordicentrine*)

Asaruddin, M.R. et al., *Nat. Med. (Tokyo)*, 2001, **55**, 149-151 (*Dasymachaline N-oxide*)

Duguevalline**D-949**

4,10,12-Trimethoxy-8H-benzo[g]benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. 3,9,11-Trimethoxy-1,2-methylenedioxyoaporphine [853226-55-8]



$C_{20}H_{15}NO_6$ 365.342


Alkaloid from the stem bark of *Duguetia vallicola* and stems of *Desmos chinensis*. Orange needles (CHCl₃). Mp 202-203° (198-200°). λ_{max} 205 (log ϵ 3.01); 214 (log ϵ 3.09); 223 (log ϵ 3.73); 289 (log ϵ 3.04); 385 (log ϵ 2.11) (MeOH).

Pérez, E. et al., *Lett. Org. Chem.*, 2004, **1**, 102-104 (isol, pmr, cmr)

Liu, X.-T. et al., *Zhongguo Tianran Yaowu*, 2004, **2**, 205-207; *CA*, **143**, 40886w (isol, pmr, cmr)

Duguevanine**D-950**

6,7,7a,8-Tetrahydro-4,10-dimethoxy-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-12-ol, 9CI. 11-Hydroxy-3,9-dimethoxy-1,2-methylenedioxyaporphine



$C_{19}H_{19}NO_5$ 341.363

(R)-form [89368-33-2]

A major alkaloid from the leaves and stem bark of *Duguetia obovata* (Annonaceae). Cryst. (Me₂CO). Mp 165°. [α]_D -127 (c, 0.45 in CHCl₃). [α]_D -175 (c, 1.62 in Py).

N-Formyl: **N-Formylduguevanine**

[89368-32-1]

$C_{20}H_{19}NO_6$ 369.373

Minor alkaloid from the stem bark of *Duguetia obovata* (Annonaceae). Cryst. (Me₂CO). Mp 255°. [α]_D -358 (c, 0.47 in CHCl₃).

N,O-Di-Ac:

Cryst. (Me₂CO). Mp 221°. [α]_D -330 (c, 0.72 in CHCl₃).

N-Me: **N-Methyluguevanine**

[89368-34-3]


C₂₀H₂₁NO₅ 355.39

Minor alkaloid from the stem bark of *Duguetia obovata* (Annonaceae). Non-cryst. Negative opt. rotn. in CHCl₃.

Roblot, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 862 (isol, uv, ir, pmr, cmr, ms, struct)

Dumetorine**D-951**

5,6-Dihydro-4-methyl-6-[(1-methyl-2-piperidinyl)methyl]-2H-pyran-2-one, 9CI [96552-67-9]

C₁₃H₂₁NO₂ 223.314

Alkaloid from the tubers of the famine food *Dioscorea dumetorum*. Oil. [α]_D²⁸ +40 (c, 0.021 in CHCl₃).


Corley, D.G. *et al.*, *Tet. Lett.*, 1985, **26**, 1615-1618 (isol, pmr, cmr, abs config)

Amarasekara, A.S. *et al.*, *Tet. Lett.*, 1987, **28**, 3151-3154 (synth)

Rückert, A. *et al.*, *Tet. Lett.*, 2006, **47**, 7977-7981 (synth)

Duocarmycin A**D-952**

DC 88A. Antibiotic DC 88A [118292-34-5]

C₂₆H₂₅N₃O₈ 507.499

Isol. from *Streptomyces* sp. DO-88. Active against gram-positive and -negative bacteria and tumours. Yellow powder. Mp 147-148°. [α]_D²² +282 (c, 0.1 in MeOH). λ_{max} 215 (ε 50000); 298 (ε 22000); 333 (ε 27000); 430 (sh) (ε 3500) (0.01N HCl) (Derep). λ_{max} 215 (ε 53000); 257 (sh) (ε 24000); 325 (ε 30000); 455 (ε 4600) (0.01N NaOH) (Derep). λ_{max} 215 (ε 55000); 310 (ε 18000); 358 (ε 28000); 425 (ε 8000) (MeOH) (Derep).

▶ GZ2233200

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3728 (uv, ir, pmr, cmr)

Takahashi, I. *et al.*, *J. Antibiot.*, 1988, **41**, 1915 (isol, struct)

Fukuda, Y. *et al.*, *Tet. Lett.*, 1990, **31**, 6699 (synth)

Fukuda, Y. *et al.*, *Bioorg. Med. Chem. Lett.*, 1992, **2**, 755 (synth)

Nakatani, K. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2255 (rev, synth)

Fukuda, Y. *et al.*, *Tetrahedron*, 1994, **50**, 2793; 2809 (synth)

Boger, D.L. *et al.*, *Acc. Chem. Res.*, 1995, **28**, 20 (rev)

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 378 (isol, uv, ir, pmr, cmr, abs config)


Boger, D.L. *et al.*, *Chem. Rev.*, 1997, **97**, 787-828 (rev, synth)

Boger, D.L. *et al.*, *J.A.C.S.*, 1997, **119**, 311 (synth)

Yamada, K. *et al.*, *J.A.C.S.*, 2003, **125**, 6630-6631 (synth)

Duocarmycin C₁**D-953**

Pyrindamycin B. DC 89A₁. Antibiotic DC 89A₁. SF 2582B. Antibiotic SF 2582B [118292-35-6]

C₂₆H₂₆ClN₃O₈ 543.959

Isol. from *Streptomyces* spp. Active against gram-positive and -negative bacteria and tumours. Orange or yellow powder. Mp 173-175° dec. [α]_D²² -128 (c, 0.2 in MeOH). λ_{max} 210 (ε 53800); 248 (sh) (ε 19500); 322 (ε 30600); 406 (ε 5380); 410 (sh) (MeOH) (Derep).

▶ UY9454200

Bromo analogue: Duocarmycin B₁. DC 89B₁. Antibiotic DC 89B₁ [124325-93-5]

C₂₆H₂₆BrN₃O₈ 588.411
Isol. from *Streptomyces* sp. DO-89. Active against gram-positive and -negative bacteria and tumours. Yellow powder. Sol. MeOH, EtOAc, CHCl₃, DMSO; poorly sol. H₂O, hexane. Mp 148-149°. [α]_D²⁰ -113.5 (c, 0.2 in MeOH). λ_{max} 208 (ε 38000); 329 (ε 17800); 410 (ε 3200) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) .37 mg/kg, LD₅₀ (mus, ipr) 200 mg/kg.

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3728 (struct)

Ichimura, M. *et al.*, *J. Antibiot.*, 1988, **41**, 1285 (isol)

Ohba, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1515 (isol, pmr, cmr, struct)


Ogawa, T. *et al.*, *J. Antibiot.*, 1989, **42**, 1299 (*Duocarmycin B₁*)

Boger, D.L. *et al.*, *Acc. Chem. Res.*, 1995, **28**, 20 (rev)

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 378 (isol, pmr, cmr, ir, uv, abs config)

Duocarmycin C₂**D-954**

Pyrindamycin A. DC 89A₂. Antibiotic DC 89A₂. SF 2582A. Antibiotic SF 2582A [118292-36-7]

C₂₆H₂₆ClN₃O₈ 543.959

Isol. from *Streptomyces* spp. Active against gram-positive and -negative bacteria and tumours. Yellow needles. Mp 256-257° (235-237° dec.). [α]_D²² -51 (c, 0.2 in MeOH). λ_{max} 208 (ε 39600); 248 (sh) (ε 16000); 299 (ε 20000); 337 (ε 28000); 434

(ε 5000) (MeOH) (Derep).

▶ DF4936660

Bromo analogue: Duocarmycin B₂ [124325-94-6]

C₂₆H₂₆BrN₃O₈ 588.411

Prod. by *Streptomyces* sp. Antitumour agent. Orange cryst. Sol. MeOH, EtOAc, CHCl₃, DMSO; poorly sol. H₂O, hexane. Mp 214-215°. [α]_D²⁰ -57.5 (c, 0.2 in MeOH). λ_{max} 208 (ε 45000); 248 (sh) (ε 19100); 298 (ε 20300); 337 (ε 31500); 434 (ε 4400) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) .25 mg/kg.

Hydroxy analogue: Pyrindamycin C. SF 2582C. Antibiotic SF 2582C [125600-37-5]

C₂₆H₂₇N₃O₉ 525.514

Prod. by *Streptomyces* sp. SF2582. Antitumour agent. Orange cryst. (CHCl₃/MeOH). Sol. Py, DMSO; fairly sol. MeOH; poorly sol. EtOAc, H₂O, CHCl₃, hexane. [α]_D -54 (MeOH). Has -OH replacing -Cl. λ_{max} 208 (ε 845); 297 (E1%/1cm 473); 336 (E1%/1cm 690); 432 (E1%/1cm 93) (MeOH) (Berdy).

[134233-13-9, 130060-75-2, 134233-12-8]

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3728; 1995, **43**, 378 (struct, uv, ir, pmr, cmr, abs config)


Ohba, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1515 (isol, pmr, cmr, struct)

Ogawa, T. *et al.*, *J. Antibiot.*, 1989, **42**, 1299 (*Duocarmycin B₂*)

Ichimura, M. *et al.*, *J. Antibiot.*, 1991, **44**, 1045 (prodn)

U.S. Pat., 1991, 4 994 578; CA, **116**, 19720v (SF 2582)

Boger, D.L. *et al.*, *Acc. Chem. Res.*, 1995, **28**, 20 (rev)


Duocarmycin D**D-955**C₂₆H₂₅N₃O₉ 523.498**(R)-(?)-form** [169181-35-5]

Isol. from *Streptomyces* sp. DO-89. Potent antitumour antibiotic. Yellow powder. [α]_D -10 (c, 0.2 in MeOH). λ_{max} 350 (ε 37000); 425 (ε 5000) (MeOH).

Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 378 (isol, uv, ir, pmr, cmr, struct)

Duocarmycin SA**D-956**

DC 113. Antibiotic DC 113 [130288-24-3]




C₂₅H₂₃N₃O₇ 477.473

Prod. by *Streptomyces* sp. D0113. Exhibits antibacterial and antitumour activities. Pale yellow powder. Sol. MeOH, DMSO, EtOAc; poorly sol. H₂O, hexane. $[\alpha]_D^{24} +180$ (c, 0.1 in MeOH). λ_{\max} 235 (sh) (ϵ 21000); 316 (ϵ 16000); 367 (ϵ 27000) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 0.143 mg/kg.Ichimura, M. *et al.*, *J. Antibiot.*, 1990, **43**, 1037-1038 (isol)Boger, D.L. *et al.*, *J.A.C.S.*, 1992, **114**, 10056-10058; 1993, **115**, 9025-9036 (synth)Boger, D.L. *et al.*, *Acc. Chem. Res.*, 1995, **28**, 20-29 (rev)Yasuzawa, T. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 378 (isol, uv, ir, pmr, cmr, abs config)Muratake, H. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1064-1066; 1998, **46**, 400-412; 559-571 (synth, abs config)Boger, D.L. *et al.*, *Chem. Rev.*, 1997, **97**, 787-828 (rev, synth)Fukuda, Y. *et al.*, *Tet. Lett.*, 1997, **38**, 7207-7208 (synth)Yamada, K. *et al.*, *J.A.C.S.*, 2003, **125**, 6630-6631 (synth)Tichenor, M.S. *et al.*, *J.A.C.S.*, 2006, **128**, 15683-15696 (synth)**Duricaulic acid****D-957**


2,3-Dihydro-5-methoxy-6-methyl-7-[(3-methyl-2-butenyl)oxy]-1-oxo-4(1H)-isoindolecarboxylic acid [99307-44-5]

C₁₆H₁₉NO₅ 305.33

Isol. from *Aspergillus duricaulis*. Biologically inactive. Cryst. (Py/toluene). Mp 220° dec.

Achenbach, H. *et al.*, *Z. Naturforsch., B*, 1985, **40**, 1219 (isol, props)**Dutadrupine****D-958**

[80151-78-6]


C₁₇H₁₅NO₃ 281.31

Alkaloid from the stem bark of *Dutailyea drupacea* (Rutaceae). Needles (CHCl₃). Mp 107-109°.

Baudouin, G. *et al.*, *J. Nat. Prod.*, 1981, **44**, 546 (isol, uv, ir, pmr, ms, struct)**Dynemicin A**

BU 3420T

[124412-57-3]

C₃₀H₁₉NO₉ 537.481

Anthraquinone antibiotic. Prod. by *Micromonospora chersina* and *Micromonospora globosa*. Possesses potent antibacterial activity. Also shows anti-neoplastic props. DNA cleaving agent. Violet powder. Sol. DMF, DMSO, dioxan; fairly sol. CHCl₃, EtOAc, MeOH; poorly sol. hexane, H₂O. Mp 208-210° dec. $[\alpha]_D^{24} +270$ (c, 0.01 in DMF). λ_{\max} 239 (ϵ 24900); 283 (sh) (ϵ 5400); 530 (sh) (ϵ 5400); 569 (ϵ 10800); 599 (ϵ 10100) (MeOH) (Derep). λ_{\max} 239 (ϵ 24900); 240 (ϵ 8300); 395 (ϵ 2600); 568 (ϵ 4200); 569 (ϵ 10800); 597 (ϵ 4100); 599 (ϵ 10100) (MeOH) (Berdy). λ_{\max} 239 (ϵ 8300); 394 (ϵ 4000); 563 (ϵ 3600); 600 (ϵ 3300) (MeOH-HCl) (Berdy). λ_{\max} 216 (ϵ 15500); 264 (ϵ 23000); 394 (ϵ 3200); 597 (ϵ 10200); 641 (ϵ 10500) (MeOH-NaOH) (Berdy).

▶ KC9300000

Aldehyde: **Dynemicin C**

[143250-23-1]

C₃₀H₁₉NO₈ 521.482Prod. by *Micromonospora chersina*.

Antitumour agent. Purple powder. Sol. DMSO, dioxan, MeCN, DMF; fairly sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 286 (ϵ 6800); 570 (ϵ 8200); 600 (ϵ 7800) (MeOH). λ_{\max} 239; 290; 568; 598 (EtOH) (Berdy).

12-Deoxy: **Deoxydynemicin A**


[130640-33-4]

C₃₀H₁₉NO₈ 521.482

From *Micromonospora globosa*. Highly active against gram-positive bacteria. Red-brown cryst. (DMF/CHCl₃). λ_{\max} 234 (ϵ 12600); 257 (sh) (ϵ 10200); 291 (sh) (ϵ 5370); 393 (sh) (ϵ 1860); 507 (sh) (ϵ 3470); 544 (ϵ 4470); 587 (sh) (ϵ 3550) (MeOH) (Derep).

Konishi, M. *et al.*, *J. Antibiot.*, 1989, **42**, 1449; 1991, **44**, 1300; 1306 (isol, struct, props)Shiomi, K. *et al.*, *J. Antibiot.*, 1990, **43**, 1000 (Deoxydynemicin)Konishi, M. *et al.*, *J.A.C.S.*, 1990, **112**, 3715 (cryst struct)Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1991, **30**, 1387 (rev)Eur. Pat., 1992, 484 856; CA, **117**, 149440h (Dynemicin C)Tokiwa, Y. *et al.*, *J.A.C.S.*, 1992, **114**, 4107 (biosynth)Myers, A.G. *et al.*, *Chem. Biol.*, 1995, **2**, 33 (synth, abs config)Danishefsky, S.J. *et al.*, *J.O.C.*, 1996, **61**, 16 (rev)**D-959**Shair, M.D. *et al.*, *J.A.C.S.*, 1997, **118**, 9509 (synth)Myers, A.G. *et al.*, *J.A.C.S.*, 1997, **119**, 6072 (synth)Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (synth, rev)**Dynemicin L****D-960**

[127032-74-0]

C₃₀H₂₂ClNO₉ 575.958Prod. by *Micromonospora chersina*

ATCC 53710. Potent antibacterial and cytotoxic agent. Blue amorph. powder. Sol. DMSO, dioxan; fairly sol. EtOAc, MeOH, EtOH; poorly sol. H₂O, hexane. Mp 222-225°. $[\alpha]_D^{25} -820$ (c, 0.01 in MeOH). λ_{\max} 242 (ϵ 47000); 450 (sh); 612 (ϵ 18900); 654 (ϵ 20500) (MeOH/NaOH) (Derep). λ_{\max} 241 (ϵ 48100); 454 (ϵ 2400); 594 (ϵ 18000); 639 (ϵ 17900) (MeOH) (Derep). λ_{\max} 241 (ϵ 47900); 453 (ϵ 1800); 594 (ϵ 17700); 637 (ϵ 17500) (MeOH/HCl).

4a-Dechloro, 4a-hydroxy: **Dynemicin N**


[127003-55-8]

C₃₀H₂₃NO₁₀ 557.512From *Micromonospora chersina* ATCC

53710. Blue amorph. powder. Sol. DMSO, dioxan; fairly sol. EtOAc, EtOH, MeOH; poorly sol. H₂O, hexane. Mp 253-256°. $[\alpha]_D^{25} -200$ (c, 0.01 in MeOH). λ_{\max} 241 (ϵ 53100); 452 (ϵ 4300); 592 (ϵ 24500); 639 (ϵ 25200) (MeOH) (Berdy). λ_{\max} 243 (ϵ 56900); 450 (sh); 605 (ϵ 18300); 651 (ϵ 28300) (MeOH/NaOH) (Berdy). λ_{\max} 241 (ϵ 52700); 452 (ϵ 3900); 592 (ϵ 24100) (MeOH/HCl) (Berdy).

Konishi, M. *et al.*, *J. Antibiot.*, 1991, **44**, 1300-1305 (isol, props, abs config)**Dynemicin P****D-961**

[138370-13-5]

C₂₈H₂₁NO₉ 515.475Anthraquinone antibiotic. Prod. by *Micromonospora chersina*.

Active against gram-positive bacteria. Cytotoxic. Blue powder. Sol. DMSO; fairly sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 210 (ϵ 18100); 241 (ϵ 30700); 333 (ϵ 2630); 556 (ϵ 13300); 632 (ϵ 14000) (MeOH) (Berdy).

5-Me ether: **Dynemicin M**

[127003-54-7]

C₂₉H₂₃NO₉ 529.502

Prod. by *Micromonospora chersina*. Blue amorph. powder. Sol. DMSO, dioxan; fairly sol. EtOAc, MeOH, EtOH; poorly sol. H₂O, hexane. Mp 238–240°. [α]_D²⁷ -2460 (c, 0.01 in MeOH). λ_{\max} 243 (ε 44200); 450 (sh); 607 (ε 19200); 655 (ε 22400) (MeOH/NaOH) (Derep). λ_{\max} 241 (ε 41700); 453 (ε 1500); 589 (ε 17000); 633 (ε 17200) (MeOH) (Derep). λ_{\max} 241 (ε 41300); 453 (ε 1400); 589 (ε 16900); 638 (ε 17000) (MeOH/HCl) (Berdy).

5-Oxo: Dynemicin Q

[138370-14-6]

C₂₈H₁₉NO₆ 513.459

Prod. by *Micromonospora chersina*. Active against gram-positive bacteria. Cytotoxic. Blue powder. Sol. DMSO; fairly sol. MeOH, C₆H₆; poorly sol. H₂O. Exists in 4-enol form. λ_{\max} 217 (ε 15900); 243 (ε 20800); 335 (ε 3070); 588 (ε 13600); 634 (ε 13700) (MeOH) (Derep).

3'-Hydroxy, 5-Me ether: Dynemicin O

[138370-12-4]


C₂₉H₂₃NO₁₀ 545.501

From *Micromonospora chersina*. Active against gram-positive bacteria. Potent antibacterial and cytotoxic agent. Blue powder. Sol. DMSO; fairly sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 212 (ε 13800); 241 (ε 20600); 283 (ε 5900); 335 (ε 2360); 589 (ε 11600); 633 (ε 11500) (MeOH) (Derep).

Miyoshi-Saitoh, M. *et al.*, *J. Antibiot.*, 1991, **44**, 1037-1044; 1300-1305 (*isol, pmr, cmr, cd, props*)

Dysazecine

[85547-18-8]

D-962C₂₁H₂₅NO₄ 355.433


Minor alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae). Cryst. (EtOH) (as picrate). Mp 217–219° (picrate).

Aladesanmi, A.J. *et al.*, *J. Nat. Prod.*, 1983, **46**, 127 (*isol, uv, pmr, cmr, ms, cd, struct*)

Tanaka, H. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 2063 (*synth*)

Dyshomerythrine**D-963**

3-Epi-18-methoxyxyschelhammericine
[91897-64-2]

C₂₀H₂₅NO₄ 343.422

Alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae) and *Lagarostrobos colensoi* (*Dacrydium colensoi*). [α]_D²⁵ +84 (c, 0.1 in EtOH). Separated with difficulty from 3-Epischelhammerine in S-126. Spectroscopic data were recorded for enriched mixtures.

Picrate: Mp 115–118°.

2 α -Hydroxy: 2-Hydroxydyshomerythrine
[138909-11-2]

C₂₀H₂₅NO₅ 359.421

Alkaloid from heartwood of *Phelline comosa* (Phellinaceae). Cryst. Mp 188–190° dec. [α]_D²⁵ +176 (c, 1.5 in EtOH).

2 α -Methoxy: 2-Methoxydyshomerythrine. 3-Epi-2,18-dimethoxyxyschelhammericine
[117610-39-6]

C₂₁H₂₇NO₅ 373.448

Alkaloid from stems of *Dysoxylum lenticellare* (Meliaceae). Brownish gum. [α]_D²⁵ +76.

Aladesanmi, A.J. *et al.*, *J. Chem. Res., Synop.*, 1984, 108-109; *J. Chem. Res., Miniprint*, 1984, 1001-1039 (*pmr, ms, uv, cryst struct*)


Aladesanmi, A.J. *et al.*, *Tetrahedron*, 1988, **44**, 3749-3756 (*2 α -Methoxydyshomerythrine*)

Aladesanmi, A.J. *et al.*, *Phytochemistry*, 1991, **30**, 3497 (*2 α -Hydroxydyshomerythrine*)

Bloor, S.J. *et al.*, *Phytochemistry*, 1996, **41**, 801-802 (*isol*)

Dysibetaine**D-964**

[247166-12-7]



Relative
Configuration

C₉H₁₆N₂O₄ 216.236

Isol. from the marine sponge *Dysidea herbacea*. [α]_D²⁰ -7.3 (c, 0.26 in H₂O).

Sakai, R. *et al.*, *Tet. Lett.*, 1999, **40**, 6941-6944 (*isol, pmr, cmr, cryst struct*)

Snider, B.B. *et al.*, *Org. Lett.*, 2001, **3**, 1761-1763 (*synth*)


Wardrop, D.J. *et al.*, *Chem. Comm.*, 2004, 1230-1231 (*synth*)

Langlois, N. *et al.*, *J.O.C.*, 2004, **69**, 7558-7564 (*synth*)

Isaacson, J. *et al.*, *Org. Lett.*, 2008, **10**, 1461-1463 (*synth*)

Dysibetaine PP**D-965**

[673498-51-6]



Relative
Configuration

C₁₂H₁₈N₂O₃ 238.286


Isol. from the marine sponge *Dysidea herbacea*. Amorph. solid. [α]_D¹⁸ -70.3 (c, 0.24 in H₂O).

Sakai, R. *et al.*, *J.O.C.*, 2004, **69**, 1180-1185 (*isol, pmr, cmr*)

Ijzendoorn, D.R. *et al.*, *Org. Lett.*, 2006, **8**, 239-242 (*synth*)

Dysidamide C**D-966**

5-(3,3-Dichloro-2-methylpropyl)-4-hydroxy-3,3-dimethyl-2-pyrrolidinone, 9CI
[133738-40-6]



Absolute
Configuration

C₁₀H₁₇Cl₂NO₂ 254.155

Constit. of the Red Sea sponge *Lamellodysidea herbacea*. Needles. [α]_D -41 (c, 1 in MeOH).

N-(4,4-Dichloro-3-methylbutanoyl): Dysidamide B

[133738-39-3]

C₁₅H₂₃Cl₄NO₃ 407.162

Constit. of *Lamellodysidea herbacea*. [α]_D +6 (c, 0.16 in CHCl₃).

N-(4,4,4-Trichloro-3 ζ -methylbutanoyl): Dysidamide E

C₁₅H₂₂Cl₃NO₃ 441.607

Constit. of *Lamellodysidea herbacea*. Glass. [α]_D -3.5 (c, 0.3 in CH₂Cl₂).

3'-Chloro: Dysidamide F

C₁₀H₁₆Cl₃NO₂ 288.6

Constit. of *Lamellodysidea herbacea*. Needles. Mp 171–172°. [α]_D -36 (c, 0.84 in CH₂Cl₂).

3'-Chloro, N-(4,4-dichloro-3 ζ -methylbutanoyl): Dysidamide D

C₁₅H₂₂Cl₅NO₃ 441.607

Constit. of *Lamellodysidea herbacea*. Glass. [α]_D -4 (c, 0.15 in CH₂Cl₂).

3'-Chloro, N-(4,4,4-trichloro-3S-methylbutanoyl): 4-Hydroxy-3,3-dimethyl-1-(4,4,4-trichloro-3-methyl-1-oxobutyl)-5-(3,3,3-trichloro-2-methylpropyl)-2-pyrrolidinone, 9CI. Dysidamide

[117694-99-2]

C₁₅H₂₁Cl₆NO₃ 476.052

Alkaloid from the Red Sea sponge *Lamellodysidea herbacea*. Neurotoxin. Needles (petrol). Mp 133° (123–124°). [α]_D -16.1 (c, 2.76 in CH₂Cl₂).

3'-Chloro, 4-ketone: Dysidamide G

C₁₀H₁₄Cl₃NO₂ 286.584

Constit. of *Lamellodysidea herbacea*. Needles. Mp 109°. [α]_D -38.5 (c, 0.75 in CH₂Cl₂). Data refers to mixt. with 5-epimer.

3'-Chloro, 4-ketone, N-(4,4,4-trichloro-3 ζ -methylbutanoyl): Dysidamide H

C₁₅H₁₉Cl₆NO₃ 474.036

Constit. of *Lamellodysidea herbacea*. Glass. [α]_D -62.8 (c, 0.6 in CH₂Cl₂).

5-Epimer, 3'-chloro, 4-ketone: 5-Epidysidamide G

C₁₀H₁₄Cl₃NO₂ 286.584

Constit. of *Lamellodysidea herbacea*. Isol. as a mixt. with 5-epimer.


Carmely, S. *et al.*, *Aust. J. Chem.*, 1990, **43**, 1881-1883 (*Dysidamide*)

Isaacs, S. *et al.*, *J. Nat. Prod.*, 1991, **54**, 83-91 (*isol, pmr, cmr, ms*)

Sauleau, P. *et al.*, *Tetrahedron*, 2005, **61**, 955-963 (*isol, pmr, cmr*)

Dysidazirine

D-967

*(R,E)*-formC₁₉H₃₃NO₂ 307.475***(R,E)*-form** [113507-74-7]

Isol. from Fijian marine sponge, *Dysidea fragilis*. Low melting solid. Sol. MeOH, CHCl₃. [α]_D²³ -165 (c, 0.5 in MeOH). λ_{max} 222 (ε 16600) (MeOH) (Berdy).

► Cytotoxic.

***(S,E)*-form** [163013-15-8]

From *Dysidea fragilis*. Oil. [α]_D +47.2 (c, 1.08 in CHCl₃). Erroneous MF in paper.

***(R,Z)*-form** [172140-90-8]

Minor constit. of *Dysidea fragilis*. Oil. [α]_D²² -93.5 (c, 0.16 in CHCl₃).

Molinski, T.F. *et al.*, *J.O.C.*, 1988, **53**, 2103

(*isol, uv, pmr, ms, abs config*)

Salomon, C.E. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1463 (*isol, uv, ir, pmr, cmr, ms, cd*)


Davis, F.A. *et al.*, *J.A.C.S.*, 1995, **117**, 3651

(*synth*)

Skepper, C.K. *et al.*, *Org. Lett.*, 2008, **10**, 5269-5271 (*synth*)

Dysideaproline A

D-968

C₁₉H₂₇Cl₄N₃O₂S 503.318

Similar to Dysidenin, D-972. Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +16.5 (c, 0.02 in MeOH). λ_{max} 213 (ε 8295); 239 (ε 4400) (MeOH).

N-De-Me: Dysideaproline CC₁₈H₂₅Cl₄N₃O₂S 489.291

Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +23.5 (c, 0.02 in MeOH). λ_{max} 212 (ε 6210); 239 (ε 3920) (MeOH).

9-Dechloro: Dysideaproline FC₁₉H₂₈Cl₃N₃O₂S 468.873

Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +18.8 (c, 0.02 in MeOH). λ_{max} 215 (ε 6040); 236 (ε 4135) (MeOH).

11-Dechloro: Dysideaproline DC₁₉H₂₈Cl₃N₃O₂S 468.873

Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +28.8 (c, 0.02 in MeOH). λ_{max} 215 (ε 5800); 237 (ε 4095) (MeOH).

9,9-Bis(dechloro): Dysideaproline EC₁₉H₂₉Cl₂N₃O₂S 434.428

Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +45.5 (c, 0.02 in MeOH). λ_{max} 214 (ε 5500); 235 (ε 4335) (MeOH).

15-Methyl: Dysideaproline BC₂₀H₂₉Cl₄N₃O₂S 517.345


Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +37.5 (c, 0.02 in MeOH). λ_{max} 214 (ε 5885); 240 (ε 3270) (MeOH).

Harrigan, G.G. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1133-1138

Dysideapyrrolidone

D-969

[151805-42-4]

C₁₇H₂₂Cl₆N₂O₄ 531.088


Isol. from the marine sponge *Dysidea herbacea* (Dictyoceratida). Feeding deterrent. Prisms. Mp 165-166°. [α]_D +16.6 (c, 0.4 in CHCl₃). λ_{max} 215 (ε 5100); 238 (ε 14000) (CHCl₃). λ_{max} 215 (ε 5100); 238 (ε 14000) (MeOH) (Berdy).

Unson, M.D. *et al.*, *J.O.C.*, 1993, **58**, 6336 (*isol, uv, ir, pmr, cmr*)

Dysideathiazole

D-970

4,4,4-Trichloro-3-methyl-N-[4,4,4-trichloro-3-methyl-1-(2-thiazolyl)butyl]butanamide, 9CI
[151805-43-5]

C₁₃H₁₆Cl₆N₂O₂S 461.064

Isol. from the marine sponge *Dysidea herbacea* (Dictyoceratida). Feeding deterrent. Needles. Mp 176-177°. [α]_D -71.8 (c, 2 in CHCl₃). λ_{max} 202 (ε 6865); 241 (ε 5900) (CHCl₃). λ_{max} 202 (ε 6865); 241 (ε 5900) (MeOH) (Berdy).

N-Me: N-Methyldysideathiazole

[151805-44-6]

C₁₄H₁₈Cl₆N₂O₂S 475.091

Isol. from *Dysidea herbacea*. Feeding deterrent. Needles. Mp 96°. [α]_D -108.3 (c, 2 in CHCl₃). λ_{max} 205 (ε 6530); 243 (ε 4325) (CHCl₃). λ_{max} 205 (ε 6530); 243 (ε 4325) (MeOH) (Berdy).

10-Dechloro: 10-Dechlorodysideathiazole

[151805-45-7]

C₁₃H₁₇Cl₅N₂O₂S 426.62

Isol. from *Dysidea herbacea*. Feeding deterrent. Oil. [α]_D -57.5 (c, 0.6 in CHCl₃). λ_{max} 245 (ε 4040) (CHCl₃). λ_{max} 244 (ε 4040) (MeOH) (Berdy).

10-Dechloro, N-Me: 10-Dechloro-N-methyldysideathiazole
[151805-46-8]

C₁₄H₁₉Cl₅N₂O₂S 440.646

Isol. from *Dysidea herbacea*. Feeding deterrent. Prisms. Mp 118-119°. [α]_D -98.9 (c, 0.5 in CHCl₃). λ_{max} 245 (ε 4040) (CHCl₃). λ_{max} 244 (ε 4040) (MeOH) (Berdy).

9,10-Bisdechloro, N-Me: 9,10-Didechloro-N-methyldysideathiazole

[151805-47-9]

C₁₄H₂₀Cl₄N₂O₂S 406.202


Isol. from *Dysidea herbacea*. Feeding deterrent. Oil. [α]_D -79.9 (c, 3.2 in CHCl₃). λ_{max} 214 (ε 4270); 241 (ε 4880) (CHCl₃). λ_{max} 214 (ε 4270); 241 (ε 4880) (MeOH) (Berdy).

[151908-88-2]

Unson, M.D. *et al.*, *J.O.C.*, 1993, **58**, 6336-6343 (*isol, uv, ir, pmr, cmr, cryst struct*)

Dysidenamide

D-971

C₁₅H₂₃Cl₆N₃O₃ 506.081


Isol. from *Lyngbya majuscula*. Amorph. solid. [α]_D -40.3 (c, 0.96 in CH₂Cl₂). Related to Dysidenin, D-972.

Jimenez, J.I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 200-203

Dysidenin

D-972

5,5,5-Trichloro-4-methyl-2-[methyl(4,4,4-trichloro-3-methyl-1-oxobutyl)amino]-N-[1-(2-thiazolyl)ethyl]pentanamide, 9CI
[65647-65-6]

C₁₇H₂₃Cl₆N₃O₂S 546.17

Metab. of the sponge *Dysidea herbacea* from the Great Barrier Reef; also isol. in small amt. from a specimen from the New Guinea coast. Shows antibiotic props. Ichthyotoxin. Iodine transport inhibitor. Needles (hexane). Sol. MeOH, CHCl₃. Mp 98-99°. [α]_D²¹ -98 (c, 0.5 in CHCl₃). λ_{max} 240 (ε 3980) (MeOH) (Derep).

N-De-Me: Nordysidenin

[331970-05-9]

C₁₆H₂₁Cl₆N₃O₂S 532.143

Isol. from *Lyngbya majuscula*.

Amorph. solid. [α]_D -56.7 (c, 0.76 in CH₂Cl₂).

N-De-Me, N¹²-Me: **Pseudodysidenin**
[331970-03-7]
[383898-07-5]
C₁₇H₂₃Cl₆N₃O₂S 546.17
Isol. from *Lyngbya majuscula*. Cryst.
[α]_D²⁰ -96.9 (c, 0.03 in CHCl₃).

5-Epimer: **Isodysidenin**
[67528-34-1]
C₁₇H₂₃Cl₆N₃O₂S 546.17
Toxic constit. of *Dysidea herbacea*
from New Guinea. Ichthyotoxin. Iodine transport inhibitor. Amorph. solid. Sol. MeOH, CHCl₃. [α]_D²² +47 (c, 0.88 in CHCl₃). λ_{max} 240 (ε 3980) (MeOH) (Derep). λ_{max} 238 (ε 3311) (MeOH) (Berdy).

13-Epimer: **Neodysidenin**
[294638-50-9]
C₁₇H₂₃Cl₆N₃O₂S 546.17
Isol. from *Dysidea herbacea*. [α]_D²⁰ -52.1 (c, 0.16 in CHCl₃). λ_{max} 241 (ε 4300) (MeOH).

13-Demethyl: **13-Demethyldysidenin**
[81801-19-6]
C₁₆H₂₁Cl₆N₃O₂S 532.143
From *Dysidea herbacea* (Great Barrier Reef). Gum. [α]_D²⁰ -97 (c, 1.23 in CHCl₃). λ_{max} 240 (ε 3980) (MeOH) (Derep).

13-Demethyl, 5-epimer: **13-Demethylisodysidenin**
[81754-76-9]
C₁₆H₂₁Cl₆N₃O₂S 532.143
From *Dysidea herbacea* (Great Barrier Reef) and *Oscillatoria spongelliae*. Antihypertensive agent. Gum. [α]_D²⁰ +52 (c, 2.6 in CHCl₃). λ_{max} 242 (ε 14000) (MeOH) (Berdy).

13-Demethyl, 5-epimer, 9-dechloro: **9-Monodechloro-13-demethylisodysidenin**
[81754-77-0]
C₁₆H₂₂Cl₅N₃O₂S 497.698
From *Dysidea herbacea* (Great Barrier Reef). Gum. [α]_D²⁰ +69 (c, 0.27 in CDCl₃).

13-Demethyl, 5-epimer, 11-dechloro: **11-Monodechloro-13-demethylisodysidenin**
[81747-68-4]
C₁₆H₂₂Cl₅N₃O₂S 497.698
From *Dysidea herbacea* (Great Barrier Reef). Gum. [α]_D²⁰ +85 (c, 1.04 in CHCl₃).

13-Demethyl, 5-epimer, 9,11-bisdechloro: **9,11-Didechloro-13-demethylisodysidenin**
[189297-59-4]
C₁₆H₂₃Cl₄N₃O₂S 463.253
From *Dysidea herbacea*. [α]_D²⁰ +174 (c, 0.04 in CHCl₃).


13-Demethyl, 13-isopropyl: [383898-06-4]
C₁₉H₂₇Cl₆N₃O₂S 574.223
From *Dysidea herbacea*.

Kazlauskas, R. et al., *Tet. Lett.*, 1977, 3183 (isol, ms, ir, uv, pmr, cmr, struct)
Charles, C. et al., *Tet. Lett.*, 1978, 1519 (isol, ir, pmr, struct, Isodysidenin)
Erickson, K.L. et al., *Aust. J. Chem.*, 1982, 35, 31 (derivs)
Biskupiak, J.E. et al., *Tet. Lett.*, 1984, 25, 2935 (abs config)

Dumdei, E.J. et al., *Aust. J. Chem.*, 1997, 50, 139-144 (isol, derivs, pmr, cmr, ms)
MacMillan, J.B. et al., *Org. Lett.*, 2000, 2, 2721-2723 (Neodysidenin)
Dumrongchai, N. et al., *ACGC Chem. Res. Commun.*, 2001, 13, 17-22; *CA*, 2002, 136, 67123n (13-de-Me-13-isopropyl)
Jimenez, J.I. et al., *J. Nat. Prod.*, 2001, 64, 200-203 (Nordysidenin, Pseudodysidenin)

Dysidine† D-973

1,5-Dihydro-4-methoxy-5-(1-methylethyl)-1-(6,6,6-trichloro-3-methoxy-5-methyl-1-oxo-2-hexenyl)-2H-pyrrol-2-one, 9CI
[63079-71-0]



Absolute Configuration


C₁₆H₂₂Cl₃NO₄ 398.712
Constit. of the marine sponge *Dysidea herbacea* and prod. by *Streptomyces galilaeus* ATCC 31534. Shows antibiotic props. Needles (hexane). Sol. MeOH, acids, C₆H₆, DMSO, CHCl₃; poorly sol. H₂O, hexane. Mp 127-129°. [α]_D²⁵ +141 (c, 1 in CHCl₃). λ_{max} 226 (ε 15850); 264 (ε 26920) (EtOH) (Derep). λ_{max} 228 (E1%/1cm 570); 258 (E1%/1cm 375); 288 (E1%/1cm 145); 433 (E1%/1cm 160) (MeOH) (Berdy). λ_{max} 239 (E1%/1cm 588); 288 (E1%/1cm 160); 315 (E1%/1cm 80); 525 (E1%/1cm 175) (MeOH/NaOH) (Berdy).
▶ LD₅₀ (mus, ipr) 40 - 80 mg/kg.

(-)-form
Synthetic. Needles (hexane). Mp 130°. [α]_D²⁰ -139 (c, 0.3 in CHCl₃).

(ξ)-form
[91279-96-8]
Trace constit. of *Carteriospongia flabellifera*.
[91279-96-8]

Hofheinz, W. et al., *Helv. Chim. Acta*, 1977, 60, 660 (isol, uv, cd, ir, pmr, cmr, ms, abs config, cryst struct)
Köhler, H. et al., *Helv. Chim. Acta*, 1984, 67, 1783 (synth)
Williard, P.G. et al., *J.O.C.*, 1984, 49, 3489 (synth, ir, pmr, cmr)
Schmitz, F.J. et al., *J. Nat. Prod.*, 1988, 51, 745-748; 1112 (isol)


Dysiherbaine D-974



C₁₂H₂₀N₂O₇ 304.299
Zwitterionic amino acid. Isol. from the sponge *Dysidea herbacea*. Potent and subtype-selective agonist for ionotropic glutamate receptors. Neurotoxin. [α]_D²⁶ -3.5 (c, 0.4 in H₂O).
Sakai, R. et al., *J.A.C.S.*, 1997, 119, 4112 (isol, ir, pmr, cmr)
Masaki, H. et al., *J.A.C.S.*, 2000, 122, 5216-5217 (synth)
Snider, B.B. et al., *Org. Lett.*, 2000, 2, 635-638 (synth)
Sasaki, M. et al., *Tet. Lett.*, 2000, 41, 3923-3926; 2007, 48, 5697-5700 (synth)
Phillips, D. et al., *J.O.C.*, 2002, 67, 3194-3201 (synth)
Takahashi, K. et al., *Chem. Comm.*, 2007, 4158-4160 (synth)

Dysinosin A D-975

[477708-72-8]



Absolute Configuration

C₂₆H₄₄N₆O₁₀S 632.734
Isol. from an Australian sponge of the family Dysideidae. Inhibitor of Factor VIIa and thrombin. Amorph. solid.

3''R-Chloro: **Chlorodysinosin A**
[548443-14-7]
C₂₆H₄₃ClN₆O₁₀S 667.179
Isol. from a *Dysidea* sp. Potent inhibitor of thrombin, Factor VIIa and Factor Xa. Amorph. solid.

Carroll, A.R. et al., *J.A.C.S.*, 2002, 124, 13340-13341 (*Dysinosin A*, struct)
Hanessian, S. et al., *J.A.C.S.*, 2002, 124, 13342-13343 (*Dysinosin A*, synth)
Pat. Coop. Treaty (WIPO), 2003, 03 51 831; *CA*, 139, 47155 (*Chlorodysinosin A*)
Hanessian, S. et al., *J.A.C.S.*, 2006, 128, 10491-10495; 11727-11728 (*Chlorodysinosin A*, synth)

Dysinosin C D-976

As Dysinosin A, D-975 with
R = -CH(CH₃)₂

C₂₅H₄₂N₆O₁₀S 618.707
Isol. from the sponge *Lamellodysidea chlorea*. Inhibitor of factor VIa and thrombin. Amorph. solid. Stereochem. not confirmed. λ_{max} 202 (log ε 3.92) (MeOH).


O-Desulfo: **Dysinosin D**
C₂₅H₄₂N₆O₇ 538.643

Isol. from *Lamellodysidea chlorea*. Amorph. solid.

14-O-α-D-Glucoopyranoside: **Dysinosin B**
C₃₁H₅₂N₆O₁₅S 780.849
Isol. from *Lamellodysidea chlorea*. Amorph. solid. [α]_D²⁵ +72 (c, 0.02 in MeOH). λ_{max} 203 (log ε 4.2) (MeOH).
Carroll, A.R. et al., *J. Nat. Prod.*, 2004, 67, 1291-1294 (isol, pmr, cmr)

Dysithiazolamide


D-977

Absolute
ConfigurationC₁₈H₂₇Cl₄N₃O₂S 491.307Isol. from an unidentified *Dysidea* sp.
[α]_D²⁵ -35 (c, 0.08 in CH₂Cl₂).Ardá, A. *et al.*, *Tetrahedron*, 2005, **61**, 10093-
10098 (*isol, pmr, cmr*)Ardá, A. *et al.*, *Org. Lett.*, 2008, **10**, 2175-2178
(*synth*)

Dysoxyline


D-978

1-[2-(1,3-Benzodioxol-5-yl)ethyl]-
1,2,3,4-tetrahydro-6,7-dimethoxy-2-
methylisoquinoline, 9Cl. 1,2,3,4-Tetrahy-
dro-6,7-dimethoxy-2-methyl-1-[2-(3,4-
methylenedioxyphenyl)ethyl]isoquinoline


*(S)*-formC₂₁H₂₅NO₄ 355.433**(S)**-form [85547-16-6]Alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae). [α]_D²⁵ +22 (c, 0.34 in EtOH). λ_{max} 230 (sh) (ε 13000); 286 (ε 7800) (EtOH).*Picrate*: Mp 159-161°.**(±)**-form [156558-41-7]Synthetic. Prisms (EtOH). Mp 107-108°. λ_{max} 229 (log ε 4.06); 284 (log ε 3.88) (MeOH).Aladesanmi, A.J. *et al.*, *J. Nat. Prod.*, 1983, **46**, 127-131 (*isol, uv, pmr, ms, cmr, cd, struct*)
Nimgirawath, S. *et al.*, *Aust. J. Chem.*, 1994, **47**, 957-962 (*synth*)

Ebeietinone


[143114-79-8]

C₂₇H₄₃NO₃ 429.642Alkaloid from bulbs of *Fritillaria ebeiensis* var. *purpurea* (Liliaceae). Necdles (Me₂CO). Mp 199-203°. [α]_D -53.5 (c, 0.24 in CHCl₃).Li, P. *et al.*, *Phytochemistry*, 1992, **31**, 2190-2191 (*isol, ir, pmr, cmr, ms, struct*)**Ebracteolatinoside A**

[225921-25-5]

C₁₈H₁₉NO₁₁ 425.348Struct. information obt. from CAS. Alkaloid from the roots of *Euphorbia ebracteolata*. Component of Yue Xian Da Ji.Wang, W. *et al.*, *Zhongcaoyao*, 1999, **30**, 1-3; *CA*, **131**, 2778g**Eburenine**

1,2-Didehydroaspidospermidine, 8CI. 1,2-Dehydroaspidospermidine [19751-76-9]

C₁₉H₂₄N₂ 280.412**(+)-form** [56245-51-3]Alkaloid from *Amsonia tabernaemontana*, *Hunteria eburnea*, *Rhazya stricta*, *Vinca minor*, *Aspidosperma neblinae* and other *Aspidosperma* spp. (Apocynaceae). Bp_{0.01} 150° approx. [α]_D²⁰ +243 (EtOH).**N^b-Oxide: 1,2-Dehydroaspidospermidine N-oxide**

[104736-04-1]

C₁₉H₂₄N₂O 296.411Alkaloid from the roots of *Rhazya stricta* (Apocynaceae). [α]_D +120 (CHCl₃).12-Hydroxy: 1,2-Didehydro-12-hydroxyaspidospermidine. 12-Hydroxy-Δ¹-aspidospermidine

E-1

[857349-22-5]


C₁₉H₂₄N₂O 296.411Alkaloid from *Schinopsis lorentzii*.**(-)-form**Alkaloid from *Pleiocarpa tubicina* (Apocynaceae). Bp_{0.01} 140°. [α]_D²³ -212 (EtOH).

[65377-84-6]

Biemann, K. *et al.*, *Tet. Lett.*, 1961, 484 (*isol, ms*)Smith, G.F. *et al.*, *J.C.S.*, 1963, 4002 (*uv, ir*)Bycroft, B.W. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1147 (*isol*)Ban, Y. *et al.*, *Tetrahedron*, 1983, **39**, 3657 (*synth*)Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1986, **25**, 1779 (*N-oxide*)Azzam, S.M. *et al.*, *CA*, 2005, **143**, 112300k (1,2-Didehydro-12-hydroxyaspidospermidine)**Eburine**

Methyl aspidospermidine-3-carboxylate, 9CI

[19634-20-9]



Absolute Configuration

C₂₁H₂₈N₂O₂ 340.464Alkaloid from *Hunteria eburnea* (Apocynaceae). Noncryst. [α]_D -18 (CHCl₃).**N-Methoxycarbonyl: Eburicine**

[19634-21-0]


C₂₃H₃₀N₂O₄ 398.501From *Hunteria eburnea* (Apocynaceae). Noncryst. [α]_D -61 (CHCl₃).

[104270-92-0, 104270-93-1]

Olivier, L. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1970, **270**, 1667Yoshida, K. *et al.*, *Tetrahedron*, 1985, **41**, 5495 (*synth, Eburicine*)**Eburnamenine**

Vincamenine

[38199-30-3]

C₁₉H₂₂N₂ 278.396**(+)-form** [517-30-6]Alkaloid from *Hunteria eburnea*, *Vinca minor* and *Amsonia ciliata* (Apocynaceae). Amorph. [α]_D +183 (CHCl₃).**Picrate:**Cryst. (EtOH). Mp 186-196°. [α]_D²⁵ +183 (CHCl₃).**Methiodide:**Cryst. (H₂O). Mp 274° dec.

16,17-Dihydro: Dihydroeburnamenine. Eburnan

E-5

[47122-74-7]

C₁₉H₂₄N₂ 280.412Alkaloid from roots and leaves of *Rhazya stricta* (Apocynaceae).**(-)-form**Synthetic. Pale-yellow amorph. powder. [α]_D -151 (c, 0.16 in CHCl₃).14,15-Didehydro: Δ¹⁴-Vincamenine

[112219-48-4]

C₁₉H₂₀N₂ 276.38Isol. from aerial parts of *Melodinus celastroides*. [α]_D²² -45 (c, 1 in CHCl₃). λ_{max} 226 (log ε 4.17); 262 (log ε 3.78); 302 (log ε 3.32); 312 (log ε 3.32) (EtOH).11-Methoxy, 14,15-didehydro: 11-Methoxy-Δ¹⁴-vincamenine

[90357-61-2]


C₂₀H₂₂N₂O 306.407Alkaloid from the stem bark and aerial parts of *Melodinus guillauminii* (Apocynaceae). The abs. config. is unusual for alkaloids lacking a C-16 COOME group.**(±)-form** [62960-98-9]Alkaloid from *Vinca minor* (Apocynaceae). Amorph.

16,17-Dihydro: Mp 99-101°.

Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 2197-2199 (*struct, uv, cd*)Trojáněk, J. *et al.*, *Chem. Ind. (London)*, 1965, 1261 (*struct*)Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 3833-3847 (*ord, abs config*)Zèches, M. *et al.*, *Phytochemistry*, 1984, **23**, 171-174 (11-Methoxy-Δ¹⁴-vincamenine)Takano, S. *et al.*, *J.C.S. Perkin 1*, 1985, 305-309 (*synth, ir, pmr, ms*)Baasou, S. *et al.*, *Ann. Pharm. Fr.*, 1987, **45**, 49-56 (Δ¹⁴-Vincamenine)Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1991, **30**, 1285-1293 (Dihydroeburnamenine)**Eburnamine**

E-6

Pleiocarpinidine. Hunteriline. Vincanol. Torondel. INN. RGH4406

**(+)-form**C₁₉H₂₄N₂O 296.411

Both enantiomers are natural alkaloids. Eburnamine was the (-)-enantiomer and Vincanol the (+)-form. Many syntheses reported; only those appearing in the lit. since 1990 are given here.

(+)-form [19877-89-5]Alkaloid from *Kopsia* spp. and *Melodinus celastroides*. Antihypertensive and cerebrotonic agent. Needles (EtOH). Mp 178-180°. [α]_D +96 (c, 0.10 in CHCl₃).**Me ether:** [59630-36-3]

Mp 106°.

16-Ketone: Vinburnine, INN. Vincamone.

(-)-Eburnamonine. Eburnal. Eburnoxine. Eburnoxin. Eubron. Luvenil. Monil.


- Scleramin. Tensiplex. CH 846. Cervox-an*
[4880-88-0]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Vinca minor* and *Voa-
canga africana* (Apocynaceae). Drug
for stimulation of muscle activity.
Cerebrotonic. Mp 173-174°. [α]_D²⁰ -102
(CHCl₃). Log P 3.77 (uncertain value)
(calc). Note that the (-) enantiomer
corresponds to (+)-Eburnamine and
vice-versa. λ_{max} 242 (log ε 4.28); 268
(log ε 4); 294 (log ε 3.68); 301 (log ε
3.68) (MeOH). λ_{max} 241 (ε 19800); 295
(ε 10200); 303 (ε 4800) (EtOH).
- ▶ LD₅₀ (mus, ipr) 265 mg/kg. YY8575570
- 14,15-Didehydro: Δ¹⁴-Vincanol. 14,15-
Didehydrovincanol**
[81781-82-0]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Voaacanga africana*,
Melodinus insulae-pinorum and *Melo-
dinus celastroides*. λ_{max} 230 (log ε 3.99);
280 (log ε 3.39) (EtOH).
- 14,15-Didehydro, 16-ketone: Δ¹⁴-Vincam-
one. 14,15-Didehydrovincamone**
[74075-57-3]
C₁₉H₂₀N₂O 292.38
Alkaloid from *Voaacanga africana* and
Melodinus celastroides.
- 11-Methoxy, 16-ketone: 11-Methoxye-
burnamonine**
[4800-93-5]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from *Vinca minor* (Apocyna-
ceae). Needles (Me₂CO). Mp 169-170°.
[α]_D²² -107 (c, 0.15 in CHCl₃). λ_{max} 247
(log ε 4.33); 279 (log ε 4.07) (MeOH).
- 11-Methoxy, 14,15-didehydro: 11-Meth-
oxy-Δ¹⁴-vincanol**
[90357-62-3]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from the stem bark and aerial
parts of *Melodinus guillauminii* (Apoc-
ynaceae). [α]_D²⁰ +111 (c, 0.4 in CHCl₃).
- 16-Epimer (-)-Isoeburnamine**
[19877-90-8]
C₁₉H₂₄N₂O 296.411
Alkaloid from the roots of *Kopsia
officinalis* (Apocyanaceae). Mp 216°.
[α]_D²⁰ -106 (c, 0.68 in CHCl₃).
- 16-Epimer, 14,15-didehydro, Me ether: 16-
Epi-O-methyl-Δ¹⁴-vincanol. O-Methyl-
16-epi-Δ¹⁴-vincanol**
[256387-86-7]
C₂₀H₂₄N₂O 308.422
Alkaloid from *Voaacanga africana* and
Melodinus celastroides. λ_{max} 226 (log ε
3.95); 278 (log ε 3.39) (EtOH).
- (-)-form** [473-99-4]
Alkaloid from several genera in the
Apocynaceae notably in *Hunteria eburnea*,
Vinca minor and *Amsonia tabernaemontana*.
Cryst. (EtOH). Mp 180-181°.
Bp_{0.01} 140° subl. [α]_D²⁰ -98 (CHCl₃). λ_{max}
229 (ε 33800); 276 (sh) (ε 8400); 282 (ε
8100) (EtOH).
- Me ether: O-Methyleburnamine**
[78184-82-4]
C₂₀H₂₆N₂O 310.438
Alkaloid from *Haplophyton cimidum*
and bark of *Hunteria zeylanica*
(Apocynaceae). Mp 181°. Drawn in
one paper as *O-Methylisoeburnamine*.
- Et ether: O-Ethyleburnamine**
[77123-12-7]
C₂₁H₂₈N₂O 324.465
Alkaloid from the bark of *Hunteria
zeylanica*, and from *Kopsia larutensis*.
also obt. by cryst. of Eburnamine from
EtOH (Apocynaceae). Cryst. (EtOH).
Mp 116-117°. [α]_D²⁰ -107 (c, 0.48 in
CHCl₃). Poss. artifact. Descr. in the
paper as *O-Ethyleburnamine* but
drawn as *O-Ethylisoeburnamine*.
- 16-Ketone: Eburnamonine. Huntericine**
[474-00-0]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Hunteria eburnea*, *Am-
sonia tabernaemontana*, *Vinca minor*
and others in the Apocynaceae. Mp
183°. [α]_D²⁶ +89 (CHCl₃).
- 16-Ketone, N⁴-oxide: Eburnamonine N⁴-
oxide**
[71658-66-7]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from leaves of *Kopsia laru-
tensis* (Apocynaceae). Mp 119-121°.
[α]_D²⁰ +56 (c, 0.35 in CHCl₃).
- 18-Hydroxy: Eburnaminol**
[138615-18-6]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from the bark and stems of
Kopsia larutensis (Apocynaceae).
Amorph. [α]_D²⁰ -54 (c, 0.17 in CHCl₃).
Revised struct.
- 19R-Hydroxy: 19R-Hydroxyeburnamine**
[220054-52-4]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from *Kopsia dasyrachis*. Mp
246-248°. [α]_D²⁰ +111 (c, 0.09 in CHCl₃).
λ_{max} 201 (log ε 3.86); 229 (log ε 4.02);
283 (log ε 3.42); 291 (log ε 3.31) (EtOH).
- 19-Oxo: 19-Oxoeburnamine**
[184870-19-7]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from stems of *Kopsia pauci-
flora*. Amorph. powder. [α]_D²⁰ +83 (c,
0.058 in CHCl₃). λ_{max} 202 (log ε 3.81);
229 (log ε 4); 282 (log ε 3.39); 292 (log ε
3.25) (EtOH).
- 11,12-Dimethoxy, 16-ketone: Dimethox-
yeburnamonine**
[19775-49-6]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from *Vinca minor* (Apocyna-
ceae). Mp 220°.
- 16-Epimer: Isoeburnamine. Hunteridine.
16-Epieburnamine**
[4201-84-7]
Alkaloid from *Hunteria eburnea*, *Hun-
teria zeylanica*, *Amsonia sinensis*, *Vinca
minor*, *Kopsia larutensis* and *Haplo-
phyton cimidum* (Apocynaceae).
Cryst. (MeOH). Mp 217-220°. [α]_D²⁰
+111 (CHCl₃). λ_{max} 229 (log ε 4.54);
283 (log ε 3.92); 290 (log ε 3.82) (EtOH).
- 16-Epimer, Me ether: O-Methylisoebur-
namine**
[78184-83-5]
C₂₀H₂₆N₂O 310.438
Alkaloid from the bark of *Haplophyton
zeylanica* (Apocynaceae). Not obt. free
of *O-Methyleburnamine*. Referred to
- by the authors as *O-Methylisoeburna-
mine* but the struct. illus. in the paper
apparently shows it to be *O-Methyle-
burnamine*.
- 16-Epimer, 14,15-didehydro: Δ¹⁴-Isoebur-
namine. 14,15-Didehydroisoeburnamine**
[5083-11-4]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Amsonia sinensis*
(Apocynaceae). Cryst. Mp 175-176°.
[α]_D²⁸ +258.3 (c, 0.04 in CHCl₃).
- 16-Epimer, 19R-hydroxy: 19R-Hydroxyyi-
soeburnamine**
[220654-55-7]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from *Kopsia dasyrachis*. [α]_D²⁰
-16 (c, 0.18 in CHCl₃). λ_{max} 203 (log ε
4.39); 229 (log ε 4.53); 282 (log ε 3.74);
292 (log ε 3.83) (EtOH).
- (±)-form** [2934-73-8]
Mp 178-181°.
- 16-Ketone: Vincanorine. (±)-Eburnamo-
nine**
[2580-88-3]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Vinca minor* (Apocyna-
ceae). Cryst. (MeOH). Mp 203-204°.
- 16-Ketone, perchlorate:**
Cryst. (dioxan). Mp 243-245°.
- Aldrich Library of FT-IR Spectra, 1st edn.*,
1985, **2**, 1066C (ir)
- Aldrich Library of 13C and 1H FT NMR
Spectra*, 1992, **3**, 613B (nmr)
- Plat, M. et al., *Bull. Soc. Chim. Fr.*, 1962, 1082-
1088 (ms)
- Cava, M.P. et al., *Chem. Ind. (London)*, 1963,
1242-1243 (*O-Methyleburnamine*)
- Bartlett, M.F. et al., *J.O.C.*, 1963, **28**, 2197-2199
(*Eburnamine, Isoeburnamine, isol, uv, cd*)
- Trojánek, J. et al., *Coll. Czech. Chem. Comm.*,
1964, **29**, 433-446 (*Eburnamonine, uv, ir, synth*)
- Trojánek, J. et al., *Chem. Ind. (London)*, 1965,
1261 (cd, uv)
- Mokry, J. et al., *Coll. Czech. Chem. Comm.*,
1967, **32**, 2523-2531 (*(±)-Eburnamonine,
Isoeburnamine, Eburnamine, isol, uv, ms*)
- Bláha, K. et al., *Coll. Czech. Chem. Comm.*,
1968, **33**, 3833-3847 (*ord, abs config*)
- Döpke, W. et al., *Pharmazie*, 1968, **23**, 99
(*Dimethoxyeburnamonine*)
- Döpke, W. et al., *Tet. Lett.*, 1968, 1805-1806
(*11-Methoxyeburnamonine*)
- Chiesi, V.A. et al., *Cryst. Struct. Commun.*,
1973, **2**, 599-603 (*Vincamone, cryst struct*)
- De Angelis, L. et al., *Drugs of Today
(Barcelona)*, 1978, **14**, 160-164
(*Eburnamonine, rev*)
- Toth, G. et al., *J.C.S. Perkin 2*, 1980, 701-703;
1985, 1319-1322 (*cd, pmr, cmr*)
- Arambewela, L.S.R. et al., *Phytochemistry*,
1981, **20**, 349-350 (*Me ether, Et ether*)
- Rodier, N. et al., *Acta Cryst. B*, 1982, **38**, 863-
867 (*14,15-Didehydrovincanol: cryst struct*)
- Feng, X.Z. et al., *Planta Med.*, 1983, **48**, 280-
282 (*(-)-Isoeburnamine*)
- Feng, X.Z. et al., *J. Nat. Prod.*, 1984, **47**, 117-
122 (*pmr*)
- Zèches, M. et al., *Phytochemistry*, 1984, **23**,
171-174 (*11-Methoxy-Δ¹⁴-vincanol*)
- Baassou, S. et al., *Ann. Pharm. Fr.*, 1987, **45**,
49-56 (*Melodinus celastroides constits*)
- Sado, P.A. et al., *J. Chromatogr.*, 1988, **434**,
157-167 (*Vincamone, hplc*)
- Node, M. et al., *J.O.C.*, 1990, **55**, 517 (*synth*)
- Drago, F. et al., *Pharmacol. Biochem. Behav.*,
1990, **37**, 53 (*Eburnamonine, pharmacol*)
- Awang, K. et al., *Phytochemistry*, 1991, **30**,
3164-3167 (*Eburnaminol*)

- Liu, H.M. *et al.*, *Planta Med.*, 1991, **57**, 566-568 (*Δ^1 -Isoeburnamine, cryst struct*)
- Karvinen, E. *et al.*, *Heterocycles*, 1992, **34**, 1773-1782 (*(\pm)-Eburnamonine, synth*)
- Kam, T.-S. *et al.*, *Phytochemistry*, 1992, **31**, 2936-2938 (*Eburnamonine N-oxide*)
- Lounasmaa, M. *et al.*, *Heterocycles*, 1993, **36**, 751-760 (*synth, Eburnaminol*)
- Kam, T.S. *et al.*, *Phytochemistry*, 1993, **33**, 921-924 (*(-)-O-Ethyleburnamine, (+)-Isoeburnamine, isol, cryst struct, abs config, bibl*)
- Kaufman, M.D. *et al.*, *J.O.C.*, 1994, **59**, 7197-7198 (*(\pm)-Eburnamonine, synth*)
- Palmisano, G. *et al.*, *Tetrahedron*, 1994, **50**, 9487-9494 (*(-)-Eburnamonine, synth*)
- Kam, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 1385-1387; 1999, **51**, 159-169 (*19-Oxoeburnamine, 19-Hydroxyeburnamine*)
- Schultz, A.G. *et al.*, *J.O.C.*, 1997, **62**, 6855-6861 (*Eburnamonine, synth*)
- Kam, T.S. *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 293-298 (*19-Hydroxyeburnamine, 19-Hydroxyisoeburnamine*)
- Pegnemb, D.E. *et al.*, *Fitoterapia*, 1999, **70**, 446-448 (*Δ^1 -Vincanol, 16-Epi-O-methyl- Δ^1 -vincanol*)
- Grieco, P.A. *et al.*, *J.O.C.*, 1999, **64**, 7586-7593 (*(\pm)-Eburnamonine, synth, pmr, cmr*)
- Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1644 (*Vincamone*)
- Ghosh, A.K. *et al.*, *J.O.C.*, 2000, **65**, 5433-5435 (*(\pm)-Eburnamonine, synth, pmr, cmr*)
- Wee, A.G.H. *et al.*, *J.O.C.*, 2002, **66**, 8935-8943 (*Eburnamonine, synth*)
- Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 2764-2770 (*synth*)

Eburnaphylline

E-7

Methyl 17,20-epoxy-19,20-dihydro-18-hydroxysarpagan-16-carboxylate, 9CI [35594-10-6]



Absolute Configuration

$C_{21}H_{24}N_2O_4$ 368.432

Alkaloid from the leaves of *Hunteria eburnea* (Apocynaceae). Mp 237°. $[\alpha]_D^{25} +15.4$ (Py).

O-Ac:

Cryst. (Et₂O). Mp 215°. $[\alpha]_D +43$ (CHCl₃).


Morfaux, A.-M. *et al.*, *Bull. Soc. Chim. Fr.*, 1971, 3967-3973 (*ir, uv, ms, pmr, struct*)

Morfaux, A.M. *et al.*, *Tet. Lett.*, 1973, 1939-1941 (*struct, abs config*)

Ecgonine

E-8

3-Hydroxy-8-methyl-8-azabicyclo[3.2.1]octane-2-carboxylic acid, 9CI. *3-Hydroxytryptane-2-carboxylic acid*, 8CI



$C_9H_{15}NO_3$ 185.222

► Highly toxic by inhalation, allergen.

(-)-form [481-37-8]

Obt. by hydrol. of Cocaine, C-537. Mp 205°. $[\alpha]_D -45.4$ (H₂O).

Me ester: Methylecgonine. Ecgonine methyl ester

[7143-09-1]
 $C_{10}H_{17}NO_3$ 199.249

Minor alkaloid from *Erythroxylum coca* and other *Erythroxylum* spp. (Erythroxylaceae). Anaesthetic. Oil (synthetic). $d_{20} 1.15$. $Bp_{0.2}$ 84-88°. $n_D^{20} 1.4890$. Log P 0.13 (calc). Poorly characterised as a natural alkaloid.

Me ester; hydrochloride: [38969-40-3] Mp 212° dec. (199-200°) (synthetic).

O-Benzoyl: **Benzoylecgonine**

[519-09-5]
 $C_{16}H_{19}NO_4$ 289.33

Alkaloid from *Erythroxylum* spp. Major metab. of Cocaine. Mp 195°. $[\alpha]_D -63.3$ (H₂O).

O-Benzoyl, *Me ester*: see Cocaine, C-537

O-(E-Cinnamoyl), *Me ester: Cinnamoylcocaine. Cinnamoylecgonine methyl ester* [521-67-5]

Alkaloid from *Erythroxylum coca* and *Erythroxylum monogynum*, first obt. synthetically (Erythroxylaceae). Does not show significant mydriatic or anaesthetic props. Mp 121°. The name Cinnamoylcocaine is somewhat misleading.

O-(Z-Cinnamoyl), *Me ester: (Z)-Cinnamoylcocaine*

[50763-21-8]
 $C_{19}H_{23}NO_4$ 329.395

Minor alkaloid from *Erythroxylum* spp., found in Cocaine samples (Erythroxylaceae).

O-(2-Furancarboxyl), *Me ester*: [188948-53-0]

$C_{15}H_{19}NO_5$ 293.319

Trace alkaloid of *Erythroxylum coca* (Erythroxylaceae).

O-(3-Furancarboxyl), *Me ester*: [188948-51-8]

$C_{15}H_{19}NO_5$ 293.319

Trace alkaloid of *Erythroxylum coca*.

O-(2-Pyrrololecarbonyl), *Me ester*: [188948-54-1]

$C_{15}H_{20}N_2O_4$ 292.334

Trace alkaloid of *Erythroxylum coca* (Erythroxylaceae).

O-(3-Pyridinecarbonyl), *Me ester: Nicotinoylecgonine methyl ester*

[188948-52-9]

$C_{16}H_{20}N_2O_4$ 304.345

Trace alkaloid of *Erythroxylum coca* (Erythroxylaceae).

2-Epimer, *Me ester: Methylpseudoecgonine. Pseudoecgonine methyl ester*

$C_{10}H_{17}NO_3$ 199.249

Alkaloid from *Datura stramonium*, *Merremia aegyptia* and *Merremia gemella*.

(\pm)-form

Mp 203° (anhyd.). α - and β -Ecgonines are synthetic struct. isomers.

[56053-46-4, 56053-45-3]

Willstätter, R. *et al.*, *Annalen*, 1923, **434**, 111 (*synth*)

Chopra, R.N. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1938, **276**, 340 (*isol, Cinnamoylcocaine*)

Hegnauer, R. *et al.*, *Pharm. Acta Helv.*, 1960, **35**, 43; *CA*, **54**, 17791f (*ester*)

Bazilevskaya, G.I. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 2088; *CA*, **55**, 6511d (*synth, ester*)

Moore, J.M. *et al.*, *J. Assoc. Off. Anal. Chem.*, 1973, **56**, 1199 (*Cinnamoylcocaines*)

Baker, J.K. *et al.*, *J. Het. Chem.*, 1978, **15**, 165 (*cmr*)

Tufariello, J.J. *et al.*, *J.A.C.S.*, 1979, **101**, 2435 (*synth, pmr*)

Moore, J.M. *et al.*, *J. Forensic Sci.*, 1997, **42**, 246-255

Martinet, F. *et al.*, *Synth. Commun.*, 1997, **27**, 3485-3490 (*Benzoylecgonine*)

Singh, S. *et al.*, *Synth. Commun.*, 1997, **27**, 4003-4012 (*Methylecgonine*)

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (*Methylpseudoecgonine*)

Pedersoli, S. *et al.*, *Spectrosc. Lett.*, 2008, **41**, 101-103 (*Benzoylecgonine, pmr, cmr*)


Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 639

Echiguanine B

E-9

2-Amino-N-(3-aminopropyl)-4,7-dihydro-4-oxo-1H-pyrrolo[2,3-d]pyrimidine-5-carboxamide, 9CI

[137319-26-7]



$C_{10}H_{14}N_6O_2$ 250.26


Prod. by *Streptomyces* sp. Phosphatidylinositol kinase inhibitor. Amorph. solid. Sol. H₂O, DMSO, MeOH; poorly sol. Me₂CO, CHCl₃, EtOAc, hexane. Dec. at 270-280°. λ_{max} 217 (ϵ 15200); 250 (sh) (ϵ 10400); 281 (ϵ 7160) (0.1M HCl) (Derep). λ_{max} 256 (ϵ 9630); 292 (ϵ 7160) (0.1M NaOH) (Derep). λ_{max} 220 (ϵ 14700); 227 (ϵ 14700); 252 (sh) (ϵ 9300); 296 (ϵ 8800) (H₂O) (Derep).

Nishioka, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1321 (*isol*)

Echihumiline

E-10

[174285-73-5]



$C_{20}H_{31}NO_7$ 397.467

Alkaloid from *Echium humile* and *Echium vulgare*. Oil. $[\alpha]_D +10$ (c, 0.1 in EtOH).

N-Oxide: **Echihumiline N-oxide**

[176391-52-9]

$C_{20}H_{31}NO_8$ 413.467

Alkaloid from *Echium humile*. Gum.

3'-Ac, N-oxide: **3'-O-Acetylechihumiline N-oxide**

$C_{22}H_{33}NO_9$ 455.504

Alkaloid from *Onosma leptantha*. Oil. $[\alpha]_D^{25} +35$ (c, 0.2 in MeOH).


El-Shazly, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 310-313 (*occur*)

- El-Shazly, A. *et al.*, *Phytochemistry*, 1996, **42**, 225-230 (*isol*, *pmr*, *cmr*, *ms*, *struct*)
 Kretsi, O. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 3136-3140 (*3'-Ac-N-oxide*)

Echinidine

E-11

[520-68-3]

C₂₀H₃₁NO₇ 397.467

Stereochem. at C-3' not established.
 Alkaloid from *Echium plantagineum*, (Russian comfrey) and from *Symphytum orientale* and *Symphytum tuberosum* (Boraginaceae). Also from *Echium rawolfii* and *Echium horridum*. Hepatotoxic. Shows antimicrobial activity. Glass or gum. [α]_D¹⁸ +13.4 (EtOH).

▶ EM9252420

Picrate:Yellow needles (H₂O). Mp 142-143°.**N-Oxide: Echinidine N-oxide**

[41093-89-4]

C₂₀H₃₁NO₈ 413.467

Alkaloid from *Symphytum caucasicum* and *Echium angustifolium*. Prisms (Me₂CO/MeOH). Mp 165° dec.

3'-Ac: 3'-Acetylechimidine

[78184-80-2]

C₂₂H₃₃NO₈ 439.505


Alkaloid from *Echium vulgare*. Oil. Stereochem. at C-3' not established.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1956, **9**, 512; 1965, **18**, 1605; 1980, **33**, 1105 (*isol*, *pmr*)Ulubelen, A. *et al.*, *Phytochemistry*, 1977, **16**, 499 (*isol*)Culvenor, C.C.J. *et al.*, *Experientia*, 1980, **36**, 377 (*tox*)Huizing, H.J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1985, **318**, 476 (*cmr*)Sarg, T. *et al.*, *Fitoterapia*, 1992, **63**, 466-468 (*isol*, *oxide*)El-Shazly, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 310-313 (*3'-Acetylechimidine*)El-Shazly, A. *et al.*, *Z. Naturforsch.*, C, 1999, **54**, 295-300 (*isol*, *activity*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EAC500**Echinatine†**

E-12

Echinatine

[480-83-1]

C₁₅H₂₅NO₅ 299.366

Ester of Heliotridine in T-188 with Viridifloric acid. Diastereoisomeric with Indicine, I-68 and Intermedine, I-160. Alkaloid from *Echium vulgare*, *Rindera echinata*, *Lithospermum canescens*, *Cynoglossum amabile*, *Cynoglossum creticum*,

Eupatorium maculatum, *Eupatorium cannabinum* and others. Also from *Anchusa* spp., poss. as a stereoisomer (Boraginaceae, Asteraceae). Cryst. Mp 109-110°. [α]_D +12.8. Cryst. with difficulty, formerly obt. only as gum.

▶ EK7790000

Picolonate: Mp 205-207° dec.**N-Oxide: Echinatine N-oxide**

[20267-93-0]

C₁₅H₂₅NO₆ 315.366

Alkaloid from *Lindelofia macrostyla* (Boraginaceae).

O⁷-Ac: 7-Acetylechinatine

[56317-18-1]

C₁₇H₂₇NO₆ 341.403

Alkaloid from *Lindelofia spectabilis* (Boraginaceae).

7-O-(3-Hydroxy-3-methylbutanoyl): CanescenineC₂₀H₃₃NO₇ 399.483

Alkaloid from the aerial parts of *Lithospermum canescens*.

7-O-(3-Hydroxy-3-methylbutanoyl), 3'-Ac: Acetylcanescenine

[612541-68-1]

C₂₂H₃₅NO₈ 441.52

Alkaloid from the aerial parts of *Lithospermum canescens*.

3'-O-(4-Hydroxycinnamoyl): Cynoglossamine

[120193-39-7]

C₂₄H₃₁NO₇ 445.511

Alkaloid from *Cynoglossum creticum* (Boraginaceae). Gum. [α]_D²⁷ -4.9 (c, 0.71 in CHCl₃).

3'-Epimer: Rinderine

[6029-84-1]

C₁₅H₂₅NO₅ 299.366

Alkaloid from *Rindera baldschuanica* and *Eupatorium serotinum* (Boraginaceae, Asteraceae). Cryst. (Me₂CO). Mp 100-101°. [α]_D²⁰ +24.6 (EtOH). Ester of Heliotridine in T-188 with Trachelanthic acid.

▶ VJ7600000

3'-Epimer; hydrochloride:

Cryst. (EtOH). Mp 152-153°.

3'-Epimer; hydrobromide:Cryst. (Me₂CO/Et₂O). Mp 123-124°.**3'-Epimer, 7-Ac: 7-O-Acetylinderine**C₁₇H₂₇NO₆ 341.403

Alkaloid from *Eupatorium fortunei* and *Symphyti Radix* (root of *Symphytum officinale*). Gum. [α]_D +15.9 (c, 0.1 in MeOH).

3'-Epimer, 7-O-(3-hydroxy-3-methylbutanoyl): Canescine†

[612820-62-9]

C₂₀H₃₃NO₇ 399.483

Alkaloid from the aerial parts of *Lithospermum canescens*.

3'-Epimer, 7-O-(3-hydroxy-3-methylbutanoyl), 3'-Ac: Acetylcanescine

[612541-67-0]

C₂₂H₃₅NO₈ 441.52

Alkaloid from the aerial parts of *Lithospermum canescens*.


3'-Epimer, O⁷-Me: see Heliotrine, H-95**2',3'-Diepimer: Isoechinatine**C₁₅H₂₅NO₅ 299.366

Alkaloid from *Cynoglossum furcatum*. Pale yellow gum. [α]_D²⁰ +4.2 (c, 0.6 in EtOH).

Men'shikov, G.P. *et al.*, *CA*, 1955, **49**, 5496 (*isol*, *struct*)Akramov, S.T. *et al.*, *CA*, 1962, **57**, 16676; 1964, **61**, 11005 (*isol*, *struct*, *Rinderine*)Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1965, **18**, 1625-1637 (*pmr*)Locock, R.A. *et al.*, *J. Nat. Prod.*, 1966, **29**, 201-205 (*isol*, *Rinderine*)Pedersen, G. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 249-256 (*ms*)Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918-3935 (*cd*)Suri, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 505-507 (*7-Acetylechinatine*)Pedersen, E. *et al.*, *Phytochemistry*, 1975, **14**, 2086-2087 (*isol*)Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173-1184 (*cmr*)Sadykov, Yu.D. *et al.*, *CA*, 1985, **102**, 218363s (*isol*, *oxide*)Glinski, J.A. *et al.*, *Tet. Lett.*, 1985, **26**, 2857-2860 (*synth*)Gable, R.W. *et al.*, *Acta Cryst. C*, 1988, **44**, 1478-1481 (*cryst struct*)Asibal, C.F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 109-118 (*Cynoglossamine*)Liu, K. *et al.*, *Phytochemistry*, 1992, **31**, 2573-2574 (*7-O-Acetylinderine*)Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*rev*, *pmr*)Wiedenfeld, H. *et al.*, *Z. Naturforsch.*, C, 2003, **58**, 173-176 (*Canescine*, *Canescenine*)Ravikumar, R. *et al.*, *Indian J. Chem., Sect. B*, 2004, **43**, 406-409 (*Isoechinatine*)**Echinine†**

E-13

1,2-Dihydro-3-methoxy-1-methyl-4-quinolinol, 9CI. 1,2-Dihydro-4-hydroxy-3-methoxy-1-methylquinoline
 [28789-22-2]

C₁₁H₁₃NO₂ 191.229

Alkaloid from the seeds of *Echinops ritro*. Component of Lou Lu. Oil.

Doepke, W. *et al.*, *Pharmazie*, 1969, **24**, 782**Echinobetaine A**

E-14


MeOCH₂CH(COO[⊖])CH₂N[⊕]Me₃C₈H₁₇NO₃ 175.227

Isol. from the sponge *Echinodictyum* sp. Nematocide. [α]_D²² -49 (c, 0.6 in MeOH).

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 179-182 (*isol*, *synth*)**Echinobetaine B**

E-15

4-(Carboxymethoxymethyl)-1,3-di-methyl-1H-imidazolium hydroxide inner salt




C₈H₁₂N₂O₃ 184.194

Positive charge delocalised over the two Ns. Isol. from the sponge *Echinodictyum* sp. Nematocidal agent. [α]_D +30 (c, 0.6 in MeOH) (as TFA salt). λ_{\max} 214 (ϵ 2340) (MeOH) (TFA salt).

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 118-122 (*isol, synth, pmr, cmr*)

Echinoclathrine A E-16
[223774-66-1]

R = -(CH₂)₈CH₃C₂₂H₃₀N₂O₂ 354.491

Alkaloid from the Okinawan sponge *Echinoclathria* sp. Immunosuppressant. Amorph. solid (hexane/EtOAc). Mp 143-144°. λ_{\max} 256 (ϵ 7500) (MeOH).

Kitamura, A. *et al.*, *Tetrahedron*, 1999, **55**, 2487-2492 (*isol, uv, ir, pmr, cmr, ms*)

Echinoclathrine C E-17

[223775-02-8]

As Echinoclathrine A, E-16 with

R = -(CH₂)₁₁CH₂SHC₂₅H₃₆N₂O₂S 428.638

Alkaloid from the Okinawan sponge *Echinoclathria* sp. Amorph. solid (hexane/EtOAc). Mp 121-122°. λ_{\max} 259 (ϵ 11700) (MeOH).

S-Ac: **Echinoclathrine B**


[223774-86-5]

C₂₇H₃₈N₂O₃S 470.675

Alkaloid from an *Echinoclathria* sp. Immunosuppressant. Amorph. solid (hexane/EtOAc). Mp 135-136°. λ_{\max} 257 (ϵ 4200) (MeOH).

Kitamura, A. *et al.*, *Tetrahedron*, 1999, **55**, 2487-2492 (*isol, uv, ir, pmr, cmr, ms*)


Echinophyllin A E-18
[276690-69-8]

C₂₈H₃₇NO₄ 451.605

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -53.4 (c, 0.73 in MeOH).

Kobayashi, J. *et al.*, *Tet. Lett.*, 2000, **41**, 2939-2943 (*isol, pmr, cmr*)


Echinophyllin B E-19
[276690-70-1]

C₃₄H₄₆N₂O₆ 578.747

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -26.3 (c, 0.25 in MeOH).

Kobayashi, J. *et al.*, *Tet. Lett.*, 2000, **41**, 2939-2943 (*isol, pmr, cmr*)


Echinophyllin C E-20
[310433-44-4]

C₂₀H₂₉NO₃ 331.454

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -25.9 (c, 0.22 in MeOH). λ_{\max} 213 (log ϵ 3.85) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1576-1579 (*isol, pmr, cmr*)


Echinophyllin D E-21
[310433-45-5]

C₂₈H₃₇NO₄ 451.605

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -23.6 (c, 0.3 in EtOH). λ_{\max} 213 (log ϵ 4.03); 280 (log ϵ 3.2) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1576-1579 (*isol, pmr, cmr*)


Echinophyllin E E-22
[310433-47-7]

C₃₄H₄₆N₂O₆ 578.747

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -62.7 (c, 0.3 in EtOH). λ_{\max} 210 (log ϵ 4.02); 291 (log ϵ 3.45); 316 (log ϵ 3.45) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1576-1579 (*isol, pmr, cmr*)


Echinophyllin F E-23
[310433-49-9]

C₂₀H₂₉NO₃ 331.454

Constit. of *Echinodorus macrophyllus*. Amorph. solid. [α]_D -34.5 (c, 0.17 in MeOH). λ_{\max} 213 (log ϵ 4.01) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1576-1579 (*isol, pmr, cmr*)


Echinoramine I E-24
 α -Amino- β -(aminocarbonyl)-1,4-dihydro-4-methoxy-1-methyl-4-quinolinepropanoic acid, 9CI
[59669-20-4]

C₁₅H₁₉N₃O₄ 305.333Alkaloid from *Echinops* spp.

Pham Thanh, Ky. *et al.*, *Biochem. Physiol. Pflanz.*, 1976, **169**, 461 (*isol*)

Echinoserine

[167324-03-0]


C₅₁H₆₈N₁₂O₁₄S₂ 1137.302

Non-cyclic form of Quinomycin A, Q-58 but not a biosynth. precursor. Prod. by *Streptomyces tendae*. Biol. less active than Quinomycin A, Q-58. λ_{\max} 242; 320 (MeOH) (Berdy).

Blum, S. *et al.*, *J. Antibiot.*, 1995, **48**, 619 (*isol. struct. props*)

Echinosporamycin

[754983-47-6]



Relative Configuration


C₃₅H₂₅N₃O₁₂ 679.595

Prod. by *Micromonospora echinospora* ssp. *echinospora* LL-P175. Active against gram-positive bacteria incl. MRSA. Amorph. red powder. λ_{\max} 248; 276 (sh); 325; 470; 490 (sh) (MeCN aq.).

He, H. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1385-1391 (*isol. pmr, cmr*)

Echinosulfone A

[246037-89-8]


C₁₇H₁₀Br₂N₂O₄S 498.151

Isol. from the sponge *Echinodictyum* sp. Orange oil. λ_{\max} 221 (ϵ 42000); 250 (sh) (ϵ 14000); 280 (ϵ 17000); 320 (sh) (ϵ 9000) (MeOH).

Ovenden, S.P.B. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1246-1249 (*isol. ir, pmr, cmr, uv*)

Echinosulfonic acid C

[246037-85-4]



E-25

C₁₉H₁₄Br₂N₂O₆S 558.203

Isol. from the sponge *Echinodictyum* sp. Antibacterial agent. Orange oil. λ_{\max} 222 (ϵ 47000); 276 (ϵ 8800) (MeOH).

1''-Me ether: Echinosulfonic acid B

[246037-60-5]

C₂₀H₁₆Br₂N₂O₆S 572.23

Isol. from *Echinodictyum* sp. and *Psammoclemma* sp. Orange oil. λ_{\max} 223 (ϵ 39000); 276 (ϵ 8000) (MeOH).

1''-Et ether: Echinosulfonic acid A

[246036-86-2]

C₂₁H₁₈Br₂N₂O₆S 586.257

Isol. from *Echinodictyum* sp. Orange oil. λ_{\max} 223 (ϵ 42000); 276 (ϵ 8000) (MeOH).

1''-Deoxy: Echinosulfonic acid DC₁₉H₁₄Br₂N₂O₅S 542.204

Isol. from the sponge *Psammoclemma* sp. Cytotoxic. Brown amorph. solid. λ_{\max} 285 (MeCN/trifluoroacetic acid).

Ovenden, S.P.B. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1246-1249 (*isol. ir, pmr, cmr, uv*)

Rubnov, S. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 75-79; 2006, **20**, 517 (*Echinosulfonic acid D*)

Echinoxolinone

E-29

C₁₀H₁₀N₂O₂ 190.201

Struct. unknown. Originally considered as 3-(2-Hydroxyethyl)-4(3*H*)-quinazolinone but props. of authentic synthetic material differ from those reported for natural Echinoxolinone. The isomeric struct., 1-(2-Hydroxyethyl)-4(1*H*)-quinazolinone has also been excluded by synthesis. Alkaloid from aerial parts of *Echinops echinatus* (Asteraceae). Pale-yellow needles (CHCl₃/MeOH). Mp 150°.

7-Hydroxy: 7-Hydroxyechinoxolinone

[125386-83-6]

C₁₀H₁₀N₂O₃ 206.201

Minor alkaloid from flowers of *Echinops echinatus* (Asteraceae). Amorph. solid. Doubtful structural assignment (see statement under Echinoxolinone above).

Chaudhuri, P.K. *et al.*, *Phytochemistry*, 1987, **26**, 587-589

Reisch, J. *et al.*, *J. Nat. Prod.*, 1989, **52**, 404-407


Chaudhuri, P.K. *et al.*, *J. Nat. Prod.*, 1992, **55**, 249-250 (*7-Hydroxyechinoxolinone*)

Echitamidine

E-30

Methyl 2,16-didehydro-19-hydroxycuran-17-oate, 9CI. 19,20-Dihydro-19-hydroxyakuammicine

[38681-90-2]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the bark of *Alstonia congensis*, also from *Alstonia angustiloba*, *Alstonia pneumatophora* and *Winchia calophylla*. Hypotensive agent. Plates

(Et₂O). Mp 135° (softens at 122°).

Hydrochloride: Mp 179° dec.

Picrate: Mp 218-219°.

N^b-Oxide: Echitamidine N^b-oxideC₂₀H₂₄N₂O₄ 356.421

Alkaloid from stem bark of *Alstonia glaucescens* (Apocynaceae). Needles (EtOAc/MeOH). Mp 187-188°. $[\alpha]_D^{20}$ -475 (c, 0.17 in MeOH).

N^b-Oxide, 19-O-β-D-glucopyranoside: Echitamidine N^b-oxide 19-glucoside

[761445-63-0]

C₂₆H₃₄N₂O₉ 518.563

Alkaloid from *Alstonia scholaris*. Amorph. light yellow powder. $[\alpha]_D^{25}$ -148.4 (c, 0.35 in MeOH). λ_{\max} 246 (log ϵ 3.35); 293 (log ϵ 3.56) (MeOH).

N^a-Formyl: N^a-Formylechitamidine

[90706-49-3]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from stem bark of *Alstonia boonei* (Apocynaceae). Cryst. (EtOAc/hexane). Mp 171-173°. $[\alpha]_D^{25}$ -163 (c, 0.8 in EtOH).

Parent acid, N^b-Me: 17-Carboxy-N^b-methylechitamidineC₂₀H₂₅N₂O₃[⊕] 341.429

Quaternary alkaloid from the stem bark of *Winchia calophylla*. Amorph. solid (as chloride). $[\alpha]_D^{20}$ -207 (c, 0.52 in H₂O) (chloride). λ_{\max} 219 (log ϵ 3.92); 287 (log ϵ 3.81); 314 (log ϵ 3.71) (H₂O) (chloride).

12-Hydroxy, N^b-Me: N^b-Methylscholaricine

[132923-07-0]

C₂₁H₂₇N₂O₃[⊕] 371.455

Alkaloid from leaves of *Alstonia scholaris*. Solid. $[\alpha]_D^{25}$ -232.7 (c, 0.85 in MeOH). Counterion not specified.

12-Methoxy: Scholarine

[90762-31-5]

[78897-55-9 ((±)-form)]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Alstonia scholaris* (Apocynaceae). Cryst. (C₆H₆/Me₂CO). Mp 70°. $[\alpha]_D^{20}$ -421.3 (EtOH).

Scholarine was also reported as a racemate from the same plant (and from the same department) as a brown amorph. solid, Mp. 205-206°. This information appears unreliable. The uv spectrum for this amorphous material also differs considerably from that for semisynthetic (-)-Scholarine reported by Oguakwa. λ_{\max} 212 (log ϵ 4.7); 240 (log ϵ 4.57); 288 (log ϵ 4.11); 334 (log ϵ 4.58) (no solvent reported) (synthetic material). λ_{\max} 235 (log ϵ 3.98); 291 (log ϵ 3.7); 335 (log ϵ 4.05) (EtOH).

12-Methoxy, N^b-oxide: Scholarine N^b-oxideC₂₁H₂₆N₂O₅ 386.447

Alkaloid from the leaves of *Alstonia scholaris*. $[\alpha]_D$ -280 (c, 0.534 in CHCl₃). Strictly the name Scholarine has been applied to the racemate (see below). λ_{\max} 232 (log ϵ 3.95); 284 (log ϵ 3.51); 334 (log ϵ 3.78) (EtOH).

12-Methoxy, N^a-formyl: N^a-Formyl-12-methoxyechitamidine

[90706-51-7]

Alkaloid from stem bark of *Alstonia boonei* (Apocynaceae). Cryst. (EtOAc/hexane). Mp 185-186°. $[\alpha]_D^{25}$ -133.3 (c, 0.8 in EtOH).

12-Methoxy, parent acid, N^b-Me: 17-Carboxy-12-methoxy-N^b-methylechitamide

C₂₁H₂₇N₂O₄[⊕] 371.455
Quaternary alkaloid from the stem bark of *Winchia calophylla*. Amorph. solid (as chloride). $[\alpha]_D^{20}$ -133 (c, 0.46 in H₂O) (chloride). λ_{\max} 208 (log ϵ 4.04); 280 (log ϵ 3.63); 318 (log ϵ 3.67) (H₂O) (chloride).

20-Epimer: 19 ξ -Hydroxy-19,20R-dihydroakummicine. 20-Epi-19 ξ -echitamide

[90318-84-6]
C₂₀H₂₄N₂O₃ 340.421
Minor alkaloid from *Alstonia angustifolia*, *Alstonia pneumatophora*, *Alstonia glaucescens* and *Alstonia undulifolia* (Apocynaceae). Amorph. 19-Config. uncertain.

20-Epimer, N^b-Me: 19-Epialstogustine

[125249-28-7]
C₂₁H₂₇N₂O₃[⊕] 355.456
Quaternary alkaloid from the stem bark of *Alstonia angustifolia* (Apocynaceae). Cryst. (1-propanol) (as chloride). Mp 249° (chloride). $[\alpha]_D^{22}$ -467 (c, 1.05 in MeOH).

20-Epimer, 12-hydroxy: Scholaricine

[99694-90-3]
C₂₀H₂₄N₂O₄ 356.421
Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Mp 180° dec. $[\alpha]_D$ -200 (CHCl₃).

19,20-Diepimer: N^b-Demethylalstogustine

[125276-63-3]
C₂₀H₂₄N₂O₃ 340.421
Alkaloid from the stem bark of *Alstonia angustifolia* and *Winchia calophylla*. Mp 161-163°. $[\alpha]_D^{22}$ -442.4 (c, 0.55 in EtOH).

19,20-Diepimer, N^b-oxide: N^b-Demethylalstogustine N-oxide

[125205-50-7]
C₂₀H₂₄N₂O₄ 356.421
Alkaloid from the stem bark of *Alstonia angustifolia* (Apocynaceae). Mp 233.7-233.8°. $[\alpha]_D^{22}$ -429.3 (c, 0.73 in CHCl₃).

19,20-Diepimer, N^b-Me: Alstogustine

[125227-44-3]
C₂₁H₂₇N₂O₃[⊕] 355.456
Quaternary alkaloid from stem bark of *Alstonia angustifolia* (Apocynaceae). Cryst. (EtOH) (as chloride). Mp 231.2-231.7° (chloride). $[\alpha]_D^{22}$ -442 (c, 0.90 in MeOH).

19,20-Diepimer, 12-hydroxy: 19-Epischolaricine

[132923-06-9]
C₂₀H₂₄N₂O₄ 356.421
Alkaloid from leaves of *Alstonia scholaris*. Prisms (MeOH). Mp 210-230°. $[\alpha]_D^{28}$ -322.4 (c, 0.17 in MeOH).

19,20-Diepimer, 12-methoxy: N^b-Demethyl-12-methoxyalstogustine

C₂₁H₂₆N₂O₄ 370.447
Alkaloid from the stem bark of

Winchia calophylla. Amorph. solid. $[\alpha]_D^{20}$ -442 (c, 0.48 in CHCl₃). λ_{\max} 211 (log ϵ 4.27); 290 (log ϵ 3.75); 333 (log ϵ 4.18) (H₂O).

Goodson, J.A. *et al.*, *J.C.S.*, 1932, 2626-2630 (isol)

Raymond-Hamet, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1951, **233**, 560-562 (uv)
Djerassi, C. *et al.*, *Tet. Lett.*, 1962, **5**, 653-659 (struct, ir, ms, pmr)

Banerji, A. *et al.*, *Phytochemistry*, 1981, **20**, 540-542 ((\pm)-Scholaricine)
Oguakwa, J.U. *et al.*, *Gazz. Chim. Ital.*, 1983, **113**, 533-535 (*Alstonia boonei* constits)
Banerji, J. *et al.*, *Indian J. Chem., Sect. B*, 1984, **23**, 455 (Scholaricine)

Zeches, M. *et al.*, *Tet. Lett.*, 1984, **25**, 659-662 (cmr, config, cryst struct)
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1985, **24**, 2771-2773 (Scholaricine)

Hu, W. *et al.*, *Phytochemistry*, 1989, **28**, 1963-1966 (Alstogustine, 19-Epialstogustine)

Hu, W.-L. *et al.*, *Planta Med.*, 1989, **55**, 463-464 (N^b-Demethylalstogustine, N^b-Demethylalstogustine N-oxide)

Yamauchi, T. *et al.*, *Phytochemistry*, 1990, **29**, 3547-3552 (19-Epischolaricine, N^b-Methylscholaricine)

Keawpradub, N. *et al.*, *Phytochemistry*, 1994, **37**, 1745-1749 (oxide)

Bonjoch, J. *et al.*, *J.A.C.S.*, 1997, **119**, 7230-7240 (synth)

Kam, T.-S. *et al.*, *Phytochemistry*, 1997, **45**, 1303-1305 (Scholaricine N^b-oxide)


Salim, A.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1591-1594 (N-oxide, N-oxide glucoside, N-Demethylalstogustine)

Gan, L.-S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 18-22 (*Winchia calophylla* alkaloids)

Echiumine

[633-16-9]

E-31



C₂₀H₃₁NO₆ 381.468

Alkaloid from *Echium plantagineum*, *Amsinckia intermedia*, *Cryptantha leiocarpa* and *Cryptantha clevelandii* (Boraginaceae). Cryst. (petrol) (also descr. as a brown gum). Mp 99-100°. $[\alpha]_D$ +14.4 (c, 2.02 in EtOH).

Picrate:

Cryst. (EtOH aq.). Mp 131-132°.

2''R,3''R-Epoxyde: 2'',3''-Epoxyechiumine

[150072-70-1]
C₂₀H₃₁NO₇ 397.467

Alkaloid from *Cryptantha leiocarpa* and *Cryptantha clevelandii* (Boraginaceae). Brown gum.

2'',3''-Dihydro, 2''R*,3''S*-dihydroxy:

threo-2'',3''-Dihydroxyechiumine
[150072-69-8]

C₂₀H₃₃NO₈ 415.483

Alkaloid from *Cryptantha leiocarpa* and *Cryptantha clevelandii* (Boraginaceae). Brown gum.

2'',3''-Dihydro, 2''R*-hydroxy, 3''S*-

chloro: erythro-3''-Chloro-2''-hydroxyechiumine

[150044-60-3]
C₂₀H₃₂ClNO₇ 433.928

Alkaloid from *Cryptantha leiocarpa* and *Cryptantha clevelandii* (Boraginaceae). Brown gum.

2''E-Isomer: Myoscorpine

[82535-76-0]
C₂₀H₃₁NO₆ 381.468

Alkaloid from aerial parts of *Myosotis scorpioides*. Isol. only as a mixt. with symphytine. Ester of retronecine with tiglic and trachelanthic acids.

2''E-Isomer, N-oxide: Myoscorpine N-oxide

C₂₀H₃₁NO₇ 397.467
Alkaloid from *Echium piniana* (Boraginaceae). Gum. $[\alpha]_D$ +4.1 (c, 1 in EtOH).

2''E-Isomer, 1''-hydroxy: Hydroxymyoscorpine

[126642-92-0]
C₂₀H₃₁NO₇ 397.467
Alkaloid from roots of *Lithospermum erythrorhizon* (Boraginaceae). $[\alpha]_D$ +2.2 (c, 0.2 in EtOH).

7-Epimer: 7-Angelylheliotridine trachelanthate

C₂₀H₃₁NO₆ 381.468
Minor alkaloid from *Heliotropium supinum* (Boraginaceae). Diester of Heliosupine with angelic and trachelanthic acids. Obt. only as a mixt. with 7-Angelylheliotridine viridiflorate.

3'-Epimer: Symlandine

[74410-74-5]
C₂₀H₃₁NO₆ 381.468
Alkaloid from leaves of *Symphytum x uplandicum* (hybrid of *Symphytum officinale* and *Symphytum asperum*) (Boraginaceae). Gum. $[\alpha]_D$ +4.4 (c, 0.3 in CHCl₃).

3'-Epimer, 2'E-isomer: Symphytine

[22571-95-5]
C₂₀H₃₁NO₆ 381.468
Alkaloid from dried roots of *Symphytum officinale* and from aerial parts of *Myosotis scorpioides* (Boraginaceae). Glass or oil. $[\alpha]_D^{24}$ +3.65 (c, 4.28 in EtOH).

▶EM9252500

3',7-Diepimer: 7-Angelylheliotridine viridiflorate. 7-Angelyl-9-iridiflorylheliotridine

C₂₀H₃₁NO₆ 381.468
Minor alkaloid from *Heliotropium supinum* (Boraginaceae). Diester of Heliosupine with angelic and viridifloric acids. Obt. only as a mixt. with 7-Angelylheliotridine trachelanthate.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1956, **9**, 512; 1959, **12**, 694; 1966, **19**, 1955; 1980, **33**, 1105 (*Echiumine*, *Symlandine*)

Furuya, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 2512 (*Symphytine*)

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (uv)

Furuya, T. *et al.*, *Phytochemistry*, 1971, **10**, 2217 (*Symphytine*, isol)

Resch, J. *et al.*, *J. Nat. Prod.*, 1982, **45**, 358 (*Myoscorpine*, *Symphytine*)


Glinski, J.A. *et al.*, *Tet. Lett.*, 1985, **26**, 2857 (synth)

- Roeder, E. *et al.*, *Phytochemistry*, 1990, **29**, 690; 1991, **30**, 3107 (*Hydroxymyoscorpine, Myoscorpine N-oxide*)
 Stermitz, F.R. *et al.*, *Phytochemistry*, 1993, **33**, 383 (*Epoxyechiumine, 2'',3''-Dihydroxyechiumine, 3''-Chloro-2''-hydroxyechiumine*)
 Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*rev, pmr*)
 Kim, N.-C. *et al.*, *J. Nat. Prod.*, 2001, **64**, 251-253 (*Symplandine, Symphytine, isol, pmr, cmr*)

Echiupinine

[138590-57-5]

E-32

C₂₀H₃₁NO₆ 381.468Alkaloid from *Echium pininana* (Boraginaceae). Gum. [α]_D -15 (c, 0.1 in EtOH).**N-Oxide: Echiupinine N-oxide**

[138590-58-6]

C₂₀H₃₁NO₇ 397.467Alkaloid from *Echium pininana* (Boraginaceae). Gum. [α]_D -8.5 (c, 1 in EtOH).**4'''-Hydroxy: Pycnanthine†**

[176391-55-2]


C₂₀H₃₁NO₇ 397.467Alkaloid from *Echium humile*. Oil. [α]_D +4 (c, 0.1 in EtOH).**3'-Epimer: Symviridine**

[145774-76-1]

C₂₀H₃₁NO₆ 381.468Alkaloid from roots of *Symphytum officinale*, *Symphytum asperum* and *Symphytum x uplandicum*. Gum. [α]_D -8 (c, 0.1 in EtOH).Roeder, E. *et al.*, *Phytochemistry*, 1991, **30**, 3107-3110; 1992, **31**, 4041-4042 (*Echiupinine, Symviridine*)El-Shazly, A. *et al.*, *Phytochemistry*, 1996, **42**, 225-230 (*Pycnanthine*)**Ecliptalbine**

[215805-00-8]

E-33


C₂₇H₃₉NO₂ 409.611Alkaloid from *Solanum surinamense*. Powder (MeOH). Mp 275-278°. [α]_D²⁶ -77 (c, 0.41 in MeOH). Original plant source incorrectly identified in 1998. λ_{max} 203 (log ε 4.1); 290 (log ε 3.8) (MeOH). λ_{max} 203 (ε 12600); 290 (ε 6300) (MeOH) (Berdy).

- Abdel-Kader, M.S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1202-1208; 2000, **63**, 1184 (*isol, uv, pmr, cmr, ms*)

Ecteinascidin 594


[184300-09-2]

E-34

C₃₀H₃₂N₂O₁₀S 612.656Isol. from the Caribbean tunicate *Ecteinascidia turbinata*. Shows potent inhibition of DNA and RNA synth. and of RNA polymerase activity. Weakly cytotoxic against several cell lines. Light yellow solid. [α]_D²² -58 (c, 1.1 in MeOH). λ_{max} 207 (ε 60500); 230 (sh) (ε 11000); 287 (ε 2900) (MeOH).Sakai, R. *et al.*, *J.A.C.S.*, 1996, **118**, 9017-9023 (*isol, uv, pmr, cmr, ms, struct*)**Ecteinascidin 597**

[184300-07-0]

E-35



Absolute Configuration

C₃₀H₃₇N₃O₉S 615.703Isol. from the Caribbean tunicate *Ecteinascidia turbinata*. Shows potent inhibition of DNA and RNA synthesis and of RNA polymerase activity. Weakly cytotoxic against several cell lines. Light brown solid. [α]_D²⁵ -49 (c, 0.17 in MeOH). λ_{max} 207 (ε 46000); 230 (sh) (ε 15000); 278 (ε 3500); 285 (ε 3800) (no solvent reported). λ_{max} 207 (ε 46000); 278 (ε 3500); 285 (ε 3800) (MeOH) (Berdy).**N-De-Me: Ecteinascidin 583**

[184300-08-1]

C₂₉H₃₅N₃O₉S 601.676From *Ecteinascidia turbinata*. Shows potent inhibition of DNA and RNA synthesis and of RNA polymerase activity. Weakly cytotoxic against several cell lines. Light yellow solid. [α]_D²² -47 (c, 0.14 in CHCl₃/MeOH, 6:1). λ_{max} 207 (ε 48000); 230 (sh) (ε 9200); 280 (ε 2100); 290 (ε 2300) (no solvent reported). λ_{max} 207 (ε 48000); 280 (ε

2100); 290 (ε 2300) (MeOH) (Berdy).


I'-Deamino, I'-oxo: Ecteinascidin 596

[184300-10-5]

C₃₀H₃₄N₂O₁₀S 614.672Trace constit. of *Ecteinascidia turbinata*. Observed only in a mixt. as a MeOH adduct. Tentative struct.Sakai, R. *et al.*, *J.A.C.S.*, 1996, **118**, 9017-9023 (*isol, uv, ir, cd, pmr, cmr, ms*)Chen, J. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 8028-8032 (*synth*)**Ecteinascidin 722**

[144285-84-7]

E-36


C₃₉H₄₀N₄O₉S 740.832Isol. from the tunicate *Ecteinascidia turbinata*. Cytotoxic agent. Yellow powder. Mp 160-164°. [α]_D -40 (c, 1.64 in CHCl₃).**N¹²-Me: Ecteinascidin 736**

[144285-85-8]

C₄₀H₄₂N₄O₉S 754.859Isol. from *Ecteinascidia turbinata*. Cytotoxic agent. Fine needles (MeCN aq.). Mp 140-150° dec. [α]_D -76 (c, 0.5 in CHCl₃).Pat. Coop. Treaty (WIPO), 1992, 92 09 607; CA, **117**, 205189z (*isol*)Sakai, R. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 11456-11460 (*isol*)**Ecteinascidin 743****Trabectedin, INN, USAN. Yondelis. ET 743. NSC 648766**

[114899-77-3]

E-37

C₃₉H₄₃N₃O₁₁S 761.848Alkaloid from the ascidian *Ecteinascidia*

turbinata. Antineoplastic agent. Investigated for treatment of a variety of human tumours incl. soft tissue sarcoma, osteosarcoma, melanoma and breast cancer. Mechanisms of action include inhibition of minor-groove-interacting transcription factors. Sol. MeOH, CH₂Cl₂, C₆H₆; poorly sol. hexane, H₂O. λ_{\max} 203 (€ 62000); 240 (sh) (€ 9000); 288 (€ 6500) (0.1N HCl) (Derep). λ_{\max} 235 (€ 38000); 252 (sh) (€ 9000); 295 (€ 6500) (0.1N KOH) (Derep). λ_{\max} 202 (€ 81000); 240 (sh) (€ 15000); 287 (€ 6200) (MeOH) (Derep). λ_{\max} 202 (€ 81000); 284 (€ 6600); 289 (€ 6400) (MeOH) (Berdy).

S-Oxide: Ecteinascidin 759B

[114899-29-5]
C₃₉H₄₃N₃O₁₂S 777.848
From *Ecteinascidia turbinata*. Antineoplastic agent. Struct. revised in 2002. Formerly thought to be an N-oxide.

N¹²-Oxide: Ecteinascidin 759C. ET 759C

[146663-68-5]
C₃₉H₄₃N₃O₁₂S 777.848
From *Ecteinascidia turbinata*. Prisms (MeCN). Mp 150° dec. $[\alpha]_{\text{D}}^{25}$ -55 (c, 0.22 in CHCl₃).

N¹²-De-Me: Ecteinascidin 729

[114899-27-3]
C₃₈H₄₁N₃O₁₁S 747.821
From tunicate *Ecteinascidia turbinata*. Antineoplastic agent, immunoregulator. Sol. MeOH, C₆H₆, CH₂Cl₂; poorly sol. hexane, H₂O. λ_{\max} 203 (€ 62000); 240 (sh) (€ 9000); 288 (€ 6500) (0.1N HCl) (Derep). λ_{\max} 235 (€ 38000); 252 (sh) (€ 9000); 295 (€ 6500) (0.1N KOH) (Derep). λ_{\max} 202 (€ 81000); 240 (sh) (€ 15000); 287 (€ 6200) (MeOH) (Derep). λ_{\max} 202 (€ 61000); 283 (€ 5000); 284 (€ 4700) (MeOH) (Berdy). λ_{\max} 204 (€ 61000); 283 (€ 4800); 289 (€ 4300) (MeOH-HCl) (Berdy). λ_{\max} 215 (€ 33800); 258 (€ 8200); 290 (€ 6400) (MeOH/NaOH) (Berdy).

21-Ketone: Ecteinascidin 759A

[114899-30-8]
C₃₉H₄₁N₃O₁₁S 759.832
From *Ecteinascidia turbinata*. Formerly thought to be an N-oxide.

21-Deoxy: Ecteinascidin 745

[114899-28-4]
C₃₉H₄₃N₃O₁₀S 745.849
From *Ecteinascidia turbinata*. Antineoplastic agent, immunoregulator. Sol. MeOH, CH₂Cl₂, C₆H₆; poorly sol. hexane, H₂O. λ_{\max} 202 (€ 52000); 281 (€ 5600); 287 (€ 5400) (MeOH) (Berdy). λ_{\max} 204 (€ 51000); 281 (€ 5200); 287 (€ 5200) (MeOH/HCl) (Berdy). λ_{\max} 215 (€ 36000); 254 (€ 8300); 290 (€ 5900); 298 (€ 5800) (MeOH/NaOH) (Berdy).

21-Deoxy, 21-cyano: Ecteinascidin 770

[114899-80-8]
C₄₀H₄₂N₃O₁₀S 770.859
From *Ecteinascidia turbinata* and *Ecteinascidia thurstoni*. Antineoplastic agent, immunoregulator. Prisms

(MeOH). Sol. MeOH, CH₂Cl₂, C₆H₆; poorly sol. hexane, H₂O. Mp 216-218° dec. $[\alpha]_{\text{D}}^{24}$ -58.5 (c, 1 in CHCl₃). λ_{\max} 216 (€ 66000); 234 (€ 55000); 240 (€ 58000); 263 (€ 25000); 299 (€ 22000); 329 (€ 3900); 342 (€ 3200) (MeOH) (Berdy). λ_{\max} 216 (€ 71000); 234 (€ 57000); 240 (€ 58000); 263 (€ 29000); 299 (€ 24000); 329 (€ 5700); 342 (€ 4900) (MeOH/HCl) (Berdy). λ_{\max} 216 (€ 57000); 234 (€ 57000); 240 (€ 58000); 263 (€ 28000); 299 (€ 22000); 329 (€ 4900); 342 (€ 3700) (MeOH/NaOH) (Berdy).

21-Deoxy, 21-cyano, S-oxide: Ecteinascidin 786

[442851-31-2]
C₄₀H₄₂N₄O₁₁S 786.858
From *Ecteinascidia thurstoni*. Prisms (MeOH). Mp 197-199° dec. $[\alpha]_{\text{D}}^{24}$ -156.9 (c, 0.6 in CHCl₃).

Wright, A.E. *et al.*, *J.O.C.*, 1990, **55**, 4508 (*isol. struct*)

Rinehart, K.L. *et al.*, *J.O.C.*, 1990, **55**, 4512-4515 (*isol. struct*)

Sakai, R. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 11456-11460 (*Ecteinascidin 759C*)
Kerr, R.G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1618 (*biosynth*)

Jimeno, J.M. *et al.*, *Drugs of the Future*, 1996, **21**, 1155-1165 (*rev*)

Sakai, R. *et al.*, *J.A.C.S.*, 1996, **118**, 9017 (*cd. abs config*)

Corey, E.J. *et al.*, *J.A.C.S.*, 1996, **118**, 9202 (*synth*)

Moore, B.M. *et al.*, *J.A.C.S.*, 1997, **119**, 5475 (*drug-DNA adduct*)

Zewail-Foote, M. *et al.*, *J. Med. Chem.*, 1999, **42**, 2493-2497 (*pharmacol*)

Saito, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1549-4557 (*synth*)

Martinez, E.J. *et al.*, *Org. Lett.*, 2000, **2**, 993-996 (*synth*)

Jin, S. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 6775-6779 (*pharmacol*)

Minuzzo, M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2000, **97**, 6780-6784 (*pharmacol*)

Jeedigunta, S. *et al.*, *Tetrahedron*, 2000, **56**, 3303-3307 (*biosynth*)

Ryan, D.P. *et al.*, *Clin. Cancer Res.*, 2001, **7**, 231-242 (*clin trial*)

Erba, E. *et al.*, *Eur. J. Cancer*, 2001, **37**, 97-105 (*pharmacol*)

Zewail-Foote, M. *et al.*, *J.A.C.S.*, 2001, **123**, 6485-6495 (*pharmacol*)

Cvetkovic, R.S. *et al.*, *Drugs*, 2002, **62**, 1185-1192 (*rev*)

Suwanborirux, K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 935-937 (*Ecteinascidins 770, 786*)

Endo, A. *et al.*, *J.A.C.S.*, 2002, **124**, 6552-6554 (*synth*)

Sainz-Diaz, C.I. *et al.*, *Acta Cryst. C*, 2003, **59**, o197-o198 (*cryst struct*)

Nicolaou, K.C. *et al.*, *Classics in Total Synthesis II: More Targets, Strategies, Methods*, Wiley-VCH, 2003, 109 (*rev. synth*)

Menchaca, R. *et al.*, *J.O.C.*, 2003, **68**, 8859-8866 (*synth*)

Jimeno, J. *et al.*, *Mar. Drugs*, 2004, **1**, 14-29 (*rev*)

Zheng, S. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 1754-1759 (*synth*)

Chen, J. *et al.*, *J.A.C.S.*, 2006, **128**, 87-89 (*synth*)


Carter, N.J. *et al.*, *Drugs*, 2007, **67**, 2257-2276 (*rev*)

Beumer, J.-H. *et al.*, *Invest. New Drugs*, 2007, **25**, 1-7 (*pharmacol*)

Fishlock, D. *et al.*, *J.O.C.*, 2008, **73**, 9594-9600 (*synth*)

Edeine

E-38



A₁, R¹ = OH, R² = H
B₁, R¹ = OH, R² = C(NH₂)=NH
D, R¹ = R² = H
F, R¹ = H, R² = C(NH₂)=NH

Large peptide antibiotic. Several components have been isolated. Some strains producing one major component, others a mixt. Prod. from *Bacillus brevis* strains. Active against gram-positive and -negative bacteria, yeasts and fungi. Inhibits DNA and protein biosynth. by combining with nucleic acids.

Edeine A₁ [27656-72-0]

C₃₃H₅₈N₁₀O₁₀ 754.882
H₂O; fairly sol. MeOH; poorly sol. butanol, hexane. λ_{\max} 240 (€ 5000); 290 (€ 4000) (MeOH/NaOH) (Derep). λ_{\max} 270 (€ 3000) (MeOH/HCl) (Derep). λ_{\max} 270 (H₂O) (Berdy). λ_{\max} 270 (HCl) (Berdy). λ_{\max} 240 ; 290 (NaOH) (Berdy).

Edeine B₁ [27656-73-1]

C₃₄H₆₀N₁₂O₁₀ 796.922
Sol. H₂O; poorly sol. butanol, hexane. λ_{\max} 290 (€ 4000) (MeOH/NaOH) (Derep). λ_{\max} 270 (€ 3000) (MeOH) (Derep). λ_{\max} 270 (H₂O) (Berdy). λ_{\max} 270 (HCl) (Berdy). λ_{\max} 240 ; 290 (NaOH) (Berdy).

Edeine C [66524-52-5]

A minor component of undefined struct.

Edeine D [40627-96-1]

C₃₃H₅₈N₁₀O₉ 738.883
Sol. H₂O, MeOH; poorly sol. butanol, hexane. λ_{\max} 240 (€ 5000); 290 (€ 4000) (MeOH/NaOH) (Derep). λ_{\max} 270 (€ 3000) (MeOH/HCl) (Derep).

Decarboxy:

C₃₂H₅₈N₁₀O₇ 694.873
Shows similar activity to Edeine D.

Edeine E [66524-53-6]

A minor component of undefined struct.

Edeine F [71750-64-6]

C₃₄H₆₀N₁₂O₉ 780.923
Also prepd. by amidation of Edeine D. Sol. H₂O; poorly sol. hexane, butanol. λ_{\max} 290 (€ 4000) (MeOH/NaOH) (Derep). λ_{\max} 270 (€ 3000) (MeOH) (Derep).

Borowski, E. *et al.*, *Biochim. Biophys. Acta*, 1966, **130**, 560 (*isol*)

Borowski, E. *et al.*, *Chemotherapy (Basel)*, 1967, **12**, 12 (*isol*)

Hettinger, T.P. *et al.*, *Ann. N.Y. Acad. Sci.*, 1970, **171**, 1002 (*props*)

Hettinger, T.P. *et al.*, *Biochemistry*, 1970, **9**, 1224 (*struct*)


Wojciechowska, H. *et al.*, *Experientia*, 1972, **28**, 1423 (*isol. ms*)

Kurylo-Borowska, Z. *et al.*, *Antibiotics*, (Corcoran, J.W., Ed.), Springer, N.Y., 1975, **3**, 129 (*rev*)

- Polish Pat., 1977, 88 098; CA, **89**, 4528 (isol)
Laland, S.G. et al., *Bioact. Pept. Prod. Microorg.*, (Umezawa, H., Ed.), 1978, 7 (biosynth)
Wojciechowska, H. et al., CA, 1978, **92**, 111285 (synth)
Polish Pat., 1978, 99 496; CA, **91**, 193648 (isol)
Mazerowski, J. et al., *J. Antibiot.*, 1981, **34**, 28 (props)
Wojciechowska, H. et al., *J. Antibiot.*, 1983, **36**, 793 (struct)
Czerwinski, A. et al., *J. Antibiot.*, 1983, **36**, 1001 (synth)
Gumieniak, J. et al., *J. Antibiot.*, 1983, **36**, 1239 (props)
Kurylo-Borowska, Z. et al., *Methods Enzymol.*, 1983, **94**, 441 (isol, rev)
Wojciechowska, H. et al., *Int. J. Pept. Protein Res.*, 1985, **26**, 279 (struct)

Edpetine

[27317-54-0]




Absolute Configuration

- C₂₇H₄₁NO₄ 443.625
Alkaloid from *Petilium eduardii* (Liliaceae). Cryst. (EtOH). Mp 314-315°.
Nuriddinov, R.N. et al., *Khim. Prir. Soedin.*, 1969, **5**, 603; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 523 (ir, ms, pmr)

Eduinine

3-(2,3-Dihydroxy-3-methylbutyl)-4-methoxy-1-methyl-2(1H)-quinolinone [13849-56-4]



- C₁₆H₂₁NO₄ 291.346
Abs. config. revised in 2000.

(S)-form [27495-36-9]

Alkaloid from the bark of *Casimiroa edulis* (Mexican apple) (Rutaceae). Constituent of *Orixa japonica*. Muscle relaxant. Plates (EtOAc). Mp 140° (synthetic). [α]_D²⁰ -32 (CHCl₃) (synthetic). The alkaloid is probably partly racemic. Lower Mps. around 114° have been reported for natural isolates.

2'-Ac: **2'-O-Acetyleduinine**. *Acetoxyeduinine (incorr.)* [207603-52-9]
C₁₈H₂₃NO₅ 333.383
Alkaloid from *Skimmia laureola*. Amorph. light brown solid. [α]_D²⁰ -40 (c, 0.02 in CHCl₃). Synonym is misleading. λ_{max} 229 (log ε 4.55); 231 (log ε 4.57) (MeOH).

3'-Deoxy, 2'-ketone: 4-Methoxy-1-methyl-3-(3-methyl-2-oxobutyl)-2(1H)-quinolinone, 9CI. **Orixiarine**.

Orijanone
[123613-75-2]
C₁₆H₁₉NO₃ 273.331

Alkaloid from *Orixa japonica* and *Skimmia laureola*. Needles (cyclohexane) (synthetic). Mp 80-81° (synthetic). λ_{max} 232 (log ε 4.24); 236 (log ε 4.25) (MeOH).

3'-Deoxy, 3',4'-didehydro: **Ptefoliarine**
C₁₆H₁₉NO₃ 273.331

Alkaloid from *Skimmia laureola*. Light brown gum. [α]_D²⁰ -8 (c, 0.12 in CHCl₃). λ_{max} 232 (log ε 4.39) (MeOH).


3'-Deoxy, 3',4'-didehydro, Ac: **O-Acetylptefoliarine**. *Acetoxyptefoliarine (incorr.)*

C₁₈H₂₁NO₄ 315.368
Alkaloid from *Skimmia laureola*. Amorph. light brown solid. [α]_D²⁰ -8 (c, 0.13 in CHCl₃). λ_{max} 230 (log ε 4.16) (MeOH).

- Iriarte, J. et al., *J.C.S.*, 1956, 4170-4173 (isol, uv, ir, struct)
Toube, T.P. et al., *Tetrahedron*, 1967, **23**, 2061 (pmr, ms)
Boyd, D.R. et al., *J.C.S. (C)*, 1970, 556 (synth, struct)
Naito, T. et al., *Chem. Pharm. Bull.*, 1983, **31**, 366 (synth)
Gaston, J.L. et al., *J.C.S. Perkin 1*, 1989, 905-908 (*Orixiarine*, synth)
Barr, S.A. et al., *Chem. Comm.*, 1994, 153 (synth)
Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1998, **61**, 713-717 (*Orixiarine*, *Acetoxyeduinine*, *Ptefoliarine*, *Acetoxyptefoliarine*)
Boyd, D.R. et al., *J.C.S. Perkin 1*, 2000, 3379-3405 (abs config)
Noshita, T. et al., *Biosci., Biotechnol., Biochem.*, 2001, **65**, 710-713 (*Orixiarine*)
Kumar, R.N. et al., *Heterocycles*, 2002, **57**, 357-360 (*Orixiarine*, synth)
Funayama, S. et al., *CA*, **132**, 262617w (isol, activity)

Edulisone A

E-41



C₃₇H₄₀N₂O₁₀ 672.73
Alkaloid from the bark of *Aglaia edulis*. Needles. Mp 215-218°. [α]_D²⁰ +55 (c, 0.2 in CHCl₃). λ_{max} 208 (log ε 4.2); 278 (log ε 3.47); 329 (log ε 3.06) (EtOH).

19,20-Didehydro: **19,20-Dehydroedulisone A**

C₃₇H₃₈N₂O₁₀ 670.715
Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. [α]_D²² +95 (c, 0.2 in MeOH). λ_{max} 208 (log ε 4.88); 278 (log ε 4.09) (MeOH).


13-Epimer: **Edulisone B**

C₃₇H₄₀N₂O₁₀ 672.73
Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. [α]_D²⁰ +48.5 (c, 0.2 in CHCl₃). λ_{max} 212 (log ε 4.31); 277 (log ε 3.72); 332 (log ε 3.33) (EtOH).

- Kim, S. et al., *Tet. Lett.*, 2005, **46**, 9021-9024 (isol, pmr, cmr, cryst struct)
Kim, S. et al., *J. Nat. Prod.*, 2006, **69**, 1769-1775 (19,20-Dehydroedulisone A)

Efatine

[104330-66-7]




C₃₈H₄₄N₂O₈ 656.774
Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). Amorph. [α]_D +70 (c, 0.86 in CHCl₃).

O¹¹-De-Me, O¹²-Me: **Ambrimine**
[104330-67-8]

- C₃₈H₄₄N₂O₈ 656.774
Alkaloid from the root bark of *Hernandia peltata* (Hernandiaceae). Amorph. [α]_D +128 (c, 0.78 in CHCl₃).
Chalandre, M.C. et al., *C. R. Hebd. Seances Acad. Sci., Ser. II*, 1985, **301**, 1185 (uv, pmr, ms, struct)
Chalandre, M.C. et al., *Can. J. Chem.*, 1986, **64**, 123 (isol)

Egenine

5,8-Dihydro-6-(5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)furo[3,4-e]-1,3-benzodioxol-8-ol, 9CI. **Decumbensine** [6883-44-9]



C₂₀H₁₉NO₆ 369.373
Decumbensine originally descr. (1988) as a benzyloisoquinoline of MF C₁₉H₁₉NO₅. Alkaloid from aerial parts of *Fumaria vaillantii* and from *Corydalis decumbens* (Papaveraceae). Prob. biogenetic precursor of Bicuculline, B-119. Amorph. [α]_D²³ +219 (c, 0.11 in CHCl₃). [α]_D²⁰ +129 (c, 1.1 in CHCl₃). [α]_D²³ +99 (c, 0.12 in MeOH).

7',9-Dipimer: **Corytensine**. *Epi-α-Decumbensine*. *Humosine A* [11014-02-1]
C₂₀H₁₉NO₆ 369.373
Alkaloid from *Corydalis decumbens*, *Corydalis humosa*, *Corydalis ochotensis* and other *Corydalis* spp. (Papaveraceae). Prisms (Me₂CO). Mp 215-

215.5°. $[\alpha]_D^{25} +168$ (c, 0.5 in CHCl_3).

[131614-67-0, 131614-66-9, 131681-56-6, 131681-55-5]

Gözler, B. *et al.*, *Tetrahedron*, 1983, **39**, 577 (isol, uv, pmr, struct)

Wu, T.-S. *et al.*, *Heterocycles*, 1988, **27**, 1565; 1990, **31**, 575 (*Corytensine*)

Zhang, J. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1245 (isol, pmr, cmr)

Rein, K.S. *et al.*, *J.O.C.*, 1991, **56**, 1564 (synth, bibl)


Rozwadowska, M.D. *et al.*, *Annalen*, 1991, 287 (synth, abs config, struct, bibl)

Kessar, S.V. *et al.*, *Tet. Lett.*, 1991, **32**, 3221 (synth)

Egonine

E-44

3-Epitazattadiol
[73060-68-1]



$\text{C}_{18}\text{H}_{23}\text{NO}_5$ 333.383

Alkaloid from bulbs of *Hippeastrum equestre*. Amorph. powder. $[\alpha]_D^{22} +86$ (c, 1 in MeOH).

Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2924-2932 (synth)

Pham, L.H. *et al.*, *Phytochemistry*, 1999, **51**, 327-332 (isol, cd, pmr, cmr, ms)

2,4-Eicosadienoic acid, 9CI

E-45

2,4-Icosadienoic acid
[32839-28-4, 25448-01-5]

$\text{H}_3\text{C}(\text{CH}_2)_{14}\text{CH}=\text{CHCH}=\text{CHCOOH}$

$\text{C}_{20}\text{H}_{36}\text{O}_2$ 308.503

(E,E)-form [38225-16-0]

2-Methylpropylamide: N-(2-Methylpropyl)-2,4-eicosadienamide, 9CI. 2,4-Eicosadienoic acid isobutylamide. 2,4-Icosadienoic acid isobutylamide. N-Iso-butyl-2,4-eicosadienamide

[54794-71-7]

$\text{C}_{24}\text{H}_{45}\text{NO}$ 363.626

Alkaloid from the fruits of *Piper guineense* (Piperaceae). Cryst. (EtOAc). Mp 68° Mp 89-90°.

Piperidine: 1-(1-Oxo-2,4-eicosadienyl)piperidine, 9CI. N-(2,4-Eicosadienyl)-piperidine. 2,4-Eicosadienoic acid piperidine

[74267-84-8]

$\text{C}_{25}\text{H}_{45}\text{NO}$ 375.637

Alkaloid from *Piper retrofractum* (Javanese long pepper). λ_{max} 265 (log ϵ 4.1) (MeOH).

Pyrrolidide: see Trichonine, T-491

[38306-23-9]

Vig, B. *et al.*, *Indian J. Chem.*, 1972, **10**, 564; 1975, **13**, 1358-1359 (synth)

Okogun, J.I. *et al.*, *J.C.S. Perkin I*, 1974, 2195-2198 (2-methylpropylamide, isol)

Addae-Mensah, I. *et al.*, *Phytochemistry*, 1977, **16**, 483-485 (isol, ir, pmr, ms, struct, 2-methylpropylamide)

Vig, B. *et al.*, *J. Indian Chem. Soc.*, 1979, **56**, 935-937 (synth, piperidide)

Kikuzaka, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 1329-1333 (isol, piperidide)

Strunz, G.M. *et al.*, *Can. J. Chem.*, 1996, **74**, 419-432 (synth)

2,14-Eicosadienoic acid

E-46

$\text{H}_3\text{C}(\text{CH}_2)_4\text{CH}=\text{CH}(\text{CH}_2)_{10}\text{CH}=\text{CHCOOH}$

$\text{C}_{20}\text{H}_{36}\text{O}_2$ 308.503

(2E,14Z)-form

Piperidide: N-(2,14-Eicosadienyl)piperidine. 2,14-Eicosadienoic acid piperidide [151391-74-1]

$\text{C}_{25}\text{H}_{45}\text{NO}$ 375.637

Alkaloid from *Piper retrofractum* (Javanese long pepper). Oil. λ_{max} 212 (log ϵ 3.81) (MeOH).

Kikuzaki, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 1329-1333

4,5-Eicosanediamine

E-47

4,5-Icosanediamine. 4,5-Diaminoeicosane.

Necrosamine

$\text{H}_3\text{C}(\text{CH}_2)_{14}\text{CH}(\text{NH}_2)\text{CH}(\text{NH}_2)\text{CH}_2\text{-CH}_2\text{CH}_3$

$\text{C}_{20}\text{H}_{44}\text{N}_2$ 312.581

Prod. in the phospholipid fraction of *Escherichia coli*. Cryst. (EtOH). Mp 225° approx.

Dihydrochloride:

Cryst. Mp 275° (dec.).

Dipicrate:

Cryst. (EtOH). Mp 159-161°.

Dibenzoyl:

Cryst. (EtOH aq.). Mp 72.5-73.5°.

Ikawa, M. *et al.*, *J.A.C.S.*, 1953, **75**, 3439; 6314 (struct, synth)

5,8,11,14-Eicosatetraenoic acid

E-48

5,8,11,14-Icosatetraenoic acid

[7771-44-0]

[27400-91-5]

$\text{H}_3\text{C}(\text{CH}_2)_4\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{COOH}$

$\text{C}_{20}\text{H}_{32}\text{O}_2$ 304.472

► JX3850000

(all-Z)-form

Arachidonic acid

[506-32-1]

[31152-45-1]

Constit. of many animal phospholipids, also of some ferns and mosses. Essential fatty acid. Precursor of Prostaglandin G₂ series prostaglandins. Mp -49.5°. Bp₁ 163°.

► Potentially explosive. LD₅₀ (mus, ivn) 33 mg/kg. CE6675000

Me ester: [2566-89-4]

$\text{C}_{21}\text{H}_{34}\text{O}_2$ 318.498

Bp_{0.7} 194-196°.

Et ester: [1808-26-0]

$\text{C}_{22}\text{H}_{36}\text{O}_2$ 332.525

Liq. d²⁰ 0.9. Bp_{0.04} 121-122°. n_D²⁰ 1.4719.

2-Aminoethyl ester: **Virodhamine**.

O-Arachidonoyl ethanolamine

[287937-12-6]

$\text{C}_{22}\text{H}_{37}\text{NO}_2$ 347.54

Found in rat brain and human hippocampus. Endocannabinoid. Cannabinoid (CB₁) receptor antagonist.

(2-Hydroxyethyl)amide: Arachidonylethanolamide. **Anandamide**

[94421-68-8]

$\text{C}_{22}\text{H}_{37}\text{NO}_2$ 347.54

Isol. from porcine brain. Cannabinoid receptor ligand. Apoptosis inducer.

(4-Hydroxyphenyl)amide: N-(4-Hydroxyphenyl)-5,8,11,14-eicosatetraenamide, 9CI. N-(4-Hydroxyphenyl)arachidonylamide. **AM 404**

[183718-77-6]

$\text{C}_{26}\text{H}_{37}\text{NO}_2$ 395.584

Anandamide transport inhibitor.

20,20,20-Trifluoro: 20-Trifluoroarachidonic acid

[115178-89-7]

$\text{C}_{20}\text{H}_{29}\text{F}_3\text{O}_2$ 358.443

Used in studying physiological conditions of leukotriene B₄ synth. [6610-25-9, 77297-91-7]

Osbond, J.M. *et al.*, *J.C.S.*, 1961, 2779 (synth)

Pyatnova, Yu.B. *et al.*, *Zh. Obshch. Khim.*, 1963, **33**, 1120; *CA*, **59**, 9782b (*Et ester*)

Schlenk, H. *et al.*, *J. Am. Oil Chem. Soc.*, 1965, **42**, 481 (*occur*)

Wagner, H. *et al.*, *Naturwissenschaften*, 1965, **52**, 305 (*occur*)

Frost, D.J. *et al.*, *Chem. Phys. Lipids*, 1975, **15**, 53 (*pmr*)

Fryer, R.I. *et al.*, *J.O.C.*, 1975, **40**, 348 (*synth*)

Gunstone, F.D. *et al.*, *Chem. Phys. Lipids*, 1976, **17**, 1 (*cmr*)

Corey, E.J. *et al.*, *Tet. Lett.*, 1984, **25**, 2729 (*purify*)

Tanaka, Y. *et al.*, *Arch. Biochem. Biophys.*, 1988, **263**, 178 (*synth*)

Devane, W.A. *et al.*, *Science (Washington, D.C.)*, 1992, **258**, 1946 (*Anandamide*)

Khanolkar, A.D. *et al.*, *J. Med. Chem.*, 1996, **39**, 4515-4519 (*AM 404*, *synth*, *pmr*)

Gonzalez, S. *et al.*, *Life Sci.*, 1999, **65**, 327-336 (*AM 404*, *pharmacol*)

Giuffrida, A. *et al.*, *Eur. J. Pharmacol.*, 2000, **408**, 161-168 (*AM 404*, *pharmacol*)

Sarker, K.P. *et al.*, *FEBS Lett.*, 2000, **472**, 39-44 (*Anandamide*, *activity*)

Beltramo, M. *et al.*, *J. Neurosci.*, 2000, **20**, 3401-3407 (*AM 404*, *pharmacol*)

Porter, A.C. *et al.*, *J. Pharmacol. Exp. Ther.*, 2002, **301**, 1020-1024 (*virodhamine*, *pharmacol*)

Anagnostopoulos, D. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2766-2770 (*all-E-form*, *synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AQS750

2,4,8-Eicosatrienoic acid

E-49

2,4,8-Icosatrienoic acid

[27070-56-0]

$\text{H}_3\text{C}(\text{CH}_2)_{10}\text{CH}=\text{CHCH}_2\text{CH}_2(\text{CH}=\text{CH})_2\text{COOH}$

$\text{C}_{20}\text{H}_{34}\text{O}_2$ 306.487

(2E,4E,8Z)-form

2-Methylpropylamide: **2,4,8-Eicosatrienoic acid isobutylamide**. N-Isobutyl-2,4,8-eicosatrienamide

[64543-30-2]
C₂₄H₄₃NO 361.61

Alkaloid from *Piper officinarum* (Javanese long pepper) (Piperaceae). Mp 67-67.5°.

Gupta, O.P. *et al.*, *Phytochemistry*, 1977, **16**, 1436 (uv, ir, pmr, ms, isol, struct)

2,4,14-Eicosatrienoic acid E-50

2,4,14-Icosatrienoic acid
H₃C(CH₂)₄CH=CH(CH₂)₈CH=CHCH=CHCOOH

C₂₀H₃₄O₂ 306.487

(2E,4E,14Z)-form

2-Methylpropylamide: **2,4,14-Eicosatrienoic acid isobutylamide**. N-Isobutyl-2,4,14-eicosatrienamide

[151391-70-7]

C₂₄H₄₃NO 361.61

Alkaloid from *Piper retrofractum* (Javanese long pepper). Oil. λ_{max} 264 (log ε 4.19) (MeOH).

Kikuzaki, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 1329-1333

2,4,16-Eicosatrienoic acid E-51

2,4,16-Icosatrienoic acid
[27070-56-0]

H₃CCH₂CH₂CH=CH(CH₂)₁₀CH=CHCHcCHCOOH

C₂₀H₃₄O₂ 306.487

(2E,4E,16Z)-form

Piperidine: **Pipereicosalidine**

[145237-20-3]

C₂₅H₄₃NO 373.621

Alkaloid from fruits of *Piper retrofractum* (Javanese long pepper) (Piperaceae). Oil.

Ahn, J.W. *et al.*, *Phytochemistry*, 1992, **31**, 3609

8,11,14-Eicosatrienoic acid E-52

8,11,14-Icosatrienoic acid

[7324-41-6]

[27070-56-0]

H₃C(CH₂)₄CH=CHCH₂CH=CHCH₂CH=CH(CH₂)₆COOH

C₂₀H₃₄O₂ 306.487

(all-Z)-form

Dihomogammalinolenic acid. γ-Homolinolenic acid. Dihomo-γ-linolenic acid
[1783-84-2]

Minor component of animal phospholipids. Constit. of *Sargassum pallidum*. Metab. of Linoleic acid. Precursor of PG₁ series of prostaglandins. Intermediate for 8,9-Leukotriene C₃.

N-(2-Hydroxyethyl)amide: **Homo-γ-linolenylethanolamide**
[150314-34-4]

C₂₂H₃₉NO₂ 349.556

Isol. from porcine brain. Endogenous cannabinoid receptor agonist.

Struijk, C.B. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1966, **85**, 1241 (synth)

Hanus, L. *et al.*, *J. Med. Chem.*, 1993, **36**, 3032-3034 (amide deriv, synth, pharmacol)

Pertwee, R. *et al.*, *Eur. J. Pharmacol.*, 1994, **259**, 115-120; 1995, **287**, 145-152 (amide deriv, pharmacol)

Groza, N.V. *et al.*, *Bioorg. Khim.*, 1998, **24**, 458-461 (synth)


Zhukova, N.V. *et al.*, *Phytochemistry*, 1999, **50**, 1209-1211 (isol)

Elacomine

Alkaloid I

[38739-23-0]

[176300-93-9]



C₁₅H₂₀N₂O₂ 260.335

(±)-form [67008-94-0]

Alkaloid from the roots of *Elaeagnus commutata*. Subl. at ca. 150° or dec. at ca. 200° (sealed tube) depending on heating rate.

3-Epimer: Isoelacomine

[176300-94-0]

C₁₅H₂₀N₂O₂ 260.335

Alkaloid from the roots of *Elaeagnus commutata*. Oil. Racemic.

James, M.N.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 2407 (cryst struct)


Pellegrini, C. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 151-168 (synth)

Miyake, F.Y. *et al.*, *Org. Lett.*, 2004, **6**, 711-713 (synth)

Elaeocarpidine

E-54

1,2,3,3a,5,6,11,11b,12,13-Decahydropyridolo[2'',1'':2',3']pyrimido[1',6':1,2]pyrido[3,4-b]indole, 9CI. **Elaeocarpidine**
[20069-07-2]



C₁₇H₂₁N₃ 267.373

Isol. in racemic form. Alkaloid from the leaves of *Elaeocarpus polydactylus*, *Elaeocarpus sphaericus* and *Elaeocarpus densiflorus* (*Elaeocarpus archboldianus*) (Elaeocarpaceae). Needles (EtOH). Mp 229-230°. [α]_D 0 (CHCl₃). λ_{max} 226 (ε 39800); 283 (ε 8000); 290 (sh) (ε 6750) (no solvent reported).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 801-806 (isol, uv, ir, pmr, ms, struct)

Harley-Mason, J. *et al.*, *Chem. Comm.*, 1969, 281 (synth)

Gribble, G.W. *et al.*, *J.O.C.*, 1970, **35**, 1944-1949 (synth, ir, ms, pmr, config)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1971, **24**, 1679-1694 (isol)

Gribble, G.W. *et al.*, *Synth. Commun.*, 1987, **17**, 377-383 (synth)

Chen, C.-K. *et al.*, *J.A.C.S.*, 1988, **110**, 4829-4831 (synth)


Gribble, G.W. *et al.*, *J.O.C.*, 1988, **53**, 3164-3170 (synth)

Diker, K. *et al.*, *J. Nat. Prod.*, 1997, **60**, 791-793 (synth, uv, ir, pmr, cmr)

Elaeocarpiline

E-55

15,16-Dihydroelaeocarpine, 9CI. **Elaeocarpiline**



(7R,8R,9R,16S)-form

C₁₆H₂₁NO₂ 259.347

(7R,8R,9R,16S)-form [21104-44-9]

Alkaloid from the leaves of *Elaeocarpus dolichostylus* and *Elaeocarpus sphaericus* (Elaeocarpaceae). Plates (Me₂CO). Mp 165-166°. [α]_D +395 (c, 0.08 in CHCl₃).

(7S,8S,9S,16S)-form

Epielaeocarpiline

[28230-64-0]

Alkaloid from the leaves of *Elaeocarpus sphaericus* (Elaeocarpaceae). Cryst. (petrol). Mp 72-74°. [α]_D -396 (c, 0.06 in CHCl₃).

(7R,8S,9S,16S)-form

Isoelaeocarpiline

[21104-46-1]

Major alkaloid from the leaves of *Elaeocarpus sphaericus* and *Elaeocarpus dolichostylus* (Elaeocarpaceae). Cryst. (Me₂CO). Mp 146-147°. [α]_D -400 (c, 0.08 in CHCl₃).

(7S,8R,9R,16S)-form

Epiisoelaeocarpiline

[28230-63-9]

Alkaloid from the leaves of *Elaeocarpus sphaericus* (Elaeocarpaceae). Cryst. (petrol). Mp 98-101°. [α]_D +340 (c, 0.08 in CHCl₃).

(7R,8R,9S,16S)-form

Alloelaeocarpiline

[28230-66-2]

Alkaloid from the leaves of *Elaeocarpus sphaericus* (Elaeocarpaceae). Cryst. (Me₂CO). Mp 131-134°. [α]_D -73 (c, 0.03 in CHCl₃) (ca. 90% pure). Not obt. pure, readily isom. to Isoelaeocarpiline. The cryst. material is an approx. 1:1 mixt. of 2 components.

(7S,8S,9R,16S)-form

Epialloelaeocarpiline

[28230-65-1]

Alkaloid from the leaves of *Elaeocarpus sphaericus* (Elaeocarpaceae). Cryst. (Me₂CO). Mp 136-137°. [α]_D +139 (c, 0.11 in CHCl₃). Readily isom. to Epiisoelaeocarpiline.


Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 793-800; 1971, **24**, 1679-1694 (isol, uv, ord, pmr, ms, abs config)

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*Isoelaeocarpiline*, cd, pmr, cmr)

Elaeocarpine

E-56

1,2,3,5,6,6a,12a,12b-Octahydro-11-methyl-12H-[1]benzopyrano[2,3-g]indolizin-12-one, 9CI. Eleaocarpine



(+)-form

$C_{16}H_{19}NO_2$ 257.332

(+)-form [30891-90-8]

Obt. by dehydrogenation of Elaeocarpiline, in E-55. $[\alpha]_D +206$ ($CHCl_3$).

(-)-form [30891-91-9]

Obt. by dehydrogenation of Epieleaocarpiline, in E-55. $[\alpha]_D -210$ ($CHCl_3$).

(±)-form [22964-80-3]

Alkaloid from the leaves of *Elaeocarpus polydactylus*, *Elaeocarpus sphaericus* and *Elaeocarpus ganitrus*. Cryst. (C_6H_6 /hexane, Me_2CO or pentane). Mp 81-82°.

Hydrobromide:

Cryst. (EtOH aq. or H_2O). Mp 303-305° (290-292°).

7-Epimer: Isoelaocarpine. Isoelaocarpine [23092-73-1]

$C_{16}H_{19}NO_2$ 257.332

Alkaloid from the leaves of *Elaeocarpus polydactylus*, *Elaeocarpus sphaericus* and *Elaeocarpus ganitrus*. Cryst. (Me_2CO or pentane). Mp 51-52° Mp 75-76°.

7-Epimer, hydrobromide: Mp 258°.**7-Epimer, picrate:** Mp 245-247°.

Wunderlich, J.A. *et al.*, *Acta Cryst. B*, 1969, **25**, 1436 (cryst struct)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 775; 1971, **24**, 1679 (isol, uv, ir, pmr, ms, synth)

Johns, S.R. *et al.*, *Chem. Comm.*, 1970, 804 (isol, uv, ir, pmr, ms, synth)

Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4395 (synth)

Tanaka, T. *et al.*, *Tetrahedron*, 1973, **29**, 1285 (synth)

Tufariello, J.J. *et al.*, *J.A.C.S.*, 1979, **101**, 7114 (synth)

Ray, A.B. *et al.*, *Phytochemistry*, 1979, **18**, 700 (isol)


Howard, A.S. *et al.*, *Tet. Lett.*, 1980, **21**, 1373 (synth)

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 872-875 (isol, pmr, cmr)

Elaeokanidine A

E-57

Decahydro-3-methylpyrrolo[2,1-f][1,6]naphthyridin-1(2H)-one, 9CI. Eleaokanidine A [33023-06-2]



Relative configuration

$C_{12}H_{20}N_2O$ 208.303

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Needles by

subl. Mp 38-38.5°. $[\alpha]_D +9$ (c, 1.0 in $CHCl_3$).

Dipicrate: Mp 153-155°.

Stereoisomer (1): Eleaokanidine B.**Eleaokanidine B**

[33963-69-8]

$C_{12}H_{20}N_2O$ 208.303

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Cryst. + $2H_2O$ by subl. Mp 93-94°. $[\alpha]_D 0$ ($CHCl_3$).

Stereoisomer (1), dipicrate:

Yellow needles (EtOH). Mp 144-146°.

Stereoisomer (2): Eleaokanidine C.**Eleaokanidine C**

[33963-68-7]

$C_{12}H_{20}N_2O$ 208.303

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Cryst. + $2H_2O$ by subl. Mp 56-58°. $[\alpha]_D +1$ (c, 0.7 in $CHCl_3$).

Stereoisomer (2), dipicrate:


Yellow needles (EtOH). Mp 212-215°.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 817 (isol, ir, pmr, ms, struct)

Elaeokanine A

E-58

1-(1,2,3,5,6,8a-Hexahydro-8-indoliziny)-1-butanone, 9CI. 8-Butyryl-1,2,3,5,6,8a-hexahydroindolizine. Eleaokanine A



(R)-form

$C_{12}H_{19}NO$ 193.288

(R)-form [33023-01-7]

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Oil; yellow needles (EtOH)(as picrate). Mp 163-165° (picrate). $[\alpha]_D +13$ (c, 0.9 in $CHCl_3$). $[\alpha]_D^{26} +63$ (c, 0.93 in $CHCl_3$) (synth).

(S)-form

1'-Alcohol (1'S-?): 1,2,3,5,6,8a-Hexahydro- α -propyl-8-indolizidinemethanol, 9CI. Eleaokanine B. Eleaokanine B

[33023-02-8]

$C_{12}H_{21}NO$ 195.304

Minor alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Gum. $[\alpha]_D -76$ (c, 0.4 in $CHCl_3$). The nat. alkaloid may be a mixture of diastereoisomers.

(±)-form [73971-21-8]

Synthetic. Oil. Mp 139.5-140.5° (134-136°)(as picrate).

1'-Alcohol: Synthetic. Oil.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 817 (isol, uv, ir, pmr, ms, struct)

Tufariello, J.J. *et al.*, *Tet. Lett.*, 1979, 4445 (synth, ir, pmr)

Watanabe, T. *et al.*, *Heterocycles*, 1980, **14**, 1433 (synth, uv, ir, pmr, ms)

Schmitthener, H.F. *et al.*, *J.O.C.*, 1980, **45**, 3372 (synth)

Howard, A.S. *et al.*, *Tet. Lett.*, 1980, **21**, 1373 (synth)

Khatri, N.A. *et al.*, *J.A.C.S.*, 1981, **103**, 6387 (synth)

Wijnberg, B.P. *et al.*, *Tet. Lett.*, 1981, **22**, 5079 (synth)

Otomasu, H. *et al.*, *Tetrahedron*, 1982, **38**, 2627 (synth)

Shono, T. *et al.*, *J.O.C.*, 1984, **49**, 300 (synth)

Chamberlin, A.R. *et al.*, *J.O.C.*, 1984, **49**, 1682 (synth, ir, pmr, cmr, ms)

Flitsch, W. *et al.*, *Annalen*, 1987, 649 (synth, ms)

Flann, C. *et al.*, *J.A.C.S.*, 1987, **109**, 6097 (synth)

Gribble, G.W. *et al.*, *J.O.C.*, 1988, **53**, 3164 (synth, ir, pmr, cmr, ms)

Hua, D.H. *et al.*, *J.O.C.*, 1990, **55**, 2128-2132 (synth, abs config)

Comins, D.L. *et al.*, *J.A.C.S.*, 1991, **113**, 6672

Taber, D.F. *et al.*, *J.O.C.*, 1991, **56**, 1287 (synth)

Arai, Y. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 535 (synth)

Cordero, F.M. *et al.*, *Tetrahedron*, 1993, **49**, 9867 (synth)


Arai, Y. *et al.*, *J.C.S. Perkin 1*, 1994, 15 (synth)

Dieter, R.K. *et al.*, *J.O.C.*, 2006, **71**, 5674-5678 (synth)

Elaeokanine C

E-59

1-(Octahydro-7-hydroxy-8-indoliziny)-1-butanone, 9CI. 8-Butyryl-7-hydroxyindolizidine. Eleaokanine C



(-)-form

$C_{12}H_{21}NO_2$ 211.303

(+)-form

Synthetic. Oil. $[\alpha]_D^{26} +36.9$ (c, 0.58 in $CHCl_3$). $[\alpha]_D^{23} +47$ (c, 0.4 in $CHCl_3$).

(-)-form [33023-03-9]

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Gum. $[\alpha]_D -14$ (c, 1.0 in $CHCl_3$).

Methiodide:

Prisms ($Me_2CO/MeOH$). Mp 203-205°. $[\alpha]_D -11$ (c, 0.06 in $MeOH$).

(±)-form [36451-37-3]

Synthetic. Mp 69-70°.

Methiodide:

Cryst. (Me_2CO + trace of $MeOH$). Mp 203-205°.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 817 (isol, ir, pmr, ms, struct, synth)

Tufariello, J.J. *et al.*, *Tet. Lett.*, 1979, 4445 (synth, ir, pmr)

Watanabe, T. *et al.*, *Heterocycles*, 1980, **14**, 1433 (synth)

Howard, A.S. *et al.*, *Tet. Lett.*, 1980, **21**, 1373 (synth)

Otomasu, H. *et al.*, *Tetrahedron*, 1982, **38**, 2677 (synth)

Shono, T. *et al.*, *J.O.C.*, 1984, **49**, 300 (synth)

Flitsch, W. *et al.*, *Annalen*, 1987, 649 (synth, ms)

Gribble, G.W. *et al.*, *J.O.C.*, 1988, **53**, 3164 (synth, ir, pmr, cmr, ms)


Comins, D.L. *et al.*, *J.A.C.S.*, 1991, **113**, 6672

Arai, Y. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 535 (synth)

Arai, Y. *et al.*, *J.C.S. Perkin 1*, 1994, 15 (synth)

Elaeokanine E E-60

Decahydro-3-methyl-1H-pyrano[2,3-g]indolizin-1-one, 9CI. Eleokanine E



Relative configuration

C₁₂H₁₉NO₂ 209.288

(+)-form [33023-05-1]

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Needles by subl. Mp 57-58.5°. [α]_D²⁵ +35 (c, 0.9 in CHCl₃).

4a-Epimer: Elaeokanine D. Eleokanine D [33023-04-0]

C₁₂H₁₉NO₂ 209.288

Alkaloid from the leaves of *Elaeocarpus kaniensis* (Elaeocarpaceae). Needles by subl. Mp 76-78°. [α]_D²⁵ +51 (c, 0.9 in CHCl₃). The 4a posn. (CA numbering) is adjacent to the pyran ring O.

(±)-form [77699-01-5]

Synthetic. Noncryst.

Hart, N.K. *et al.*, *Chem. Comm.*, 1971, 460 (isol, ir, pmr, ms, struct)


Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 817 (isol, ir, pmr, ms, struct)

Watanabe, T. *et al.*, *Heterocycles*, 1981, **16**, 39 (synth, ir, pmr, ms)

Elaioymin

E-61

4-Methoxy-3-(1-octenyl-ONN-azoxy)-2-butanol, 9CI. Antibiotic 1252 [23315-05-1]



C₁₃H₂₆N₂O₃ 258.36

Azoxy config. (Z- or trans-) is tentative. Isol. from culture filtrates of *Streptomyces hepaticus* and *Streptomyces gelaticus*. Tuberculostatic antibiotic. Neutral yellow oil. Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O. [α]_D²⁵ +38.4 (c, 2.8 in EtOH). λ_{\max} 238 (ε 11000) (MeOH) (Derep). λ_{\max} 256 (NaOH) (Berdy).

► Exp. carcinogen. LD₅₀ (mus, scu) 63 mg/kg. LD₅₀ (mus, ivn) 43.7 mg/kg. EL5075000

Haskell, T. *et al.*, *Antibiot. Chemother.*

(Washington, D.C.), 1954, **4**, 141-144 (isol)

Ehrlich, J. *et al.*, *Antibiot. Chemother.*

(Washington, D.C.), 1954, **4**, 338-342 (activity)

U.K. Pat., 1955, 730 341; *CA*, **49**, 14279 (synth)

Stevens, C.L. *et al.*, *J.A.C.S.*, 1956, **78**, 3229-3230 (struct)

McGahren, W.J. *et al.*, *J.A.C.S.*, 1970, **92**, 1587-1590 (cd)

Moss, R.A. *et al.*, *J.A.C.S.*, 1977, **99**, 1643-1645 (synth)

Parry, R.J. *et al.*, *J.A.C.S.*, 1982, **104**, 339-340;

1984, **106**, 5764-5765 (biosynth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, EAG000

Elatrine

E-62

C₄₀H₅₆N₃O₆ 674.899

Struct. unknown. Impossible mol. formula. Alkaloid from *Thalictrum minus* var. *elatum* (Ranunculaceae). Needles. Mp 180-183° dec. [α]_D²⁶ +229 (EtOH).

Picrate: Mp 174-175° dec.


Picolonate: Mp 178-182° dec.

Nakajima, T. *et al.*, *Yakugaku Zasshi*, 1945, **65**, 422-424; *CA*, **48**, 330h

Elegansamine

E-63

[120090-58-6]



C₂₉H₃₆N₂O₆ 508.613

Minor alkaloid from the branches of *Gelsemium elegans*. Prisms (MeOH). Mp 172-173°.

14R-Hydroxy: 14-Hydroxyelegansamine

C₂₉H₃₆N₂O₇ 524.613

Alkaloid from *Gelsemium elegans*. Amorph. powder. [α]_D²⁰ -90 (c, 0.12 in CHCl₃). λ_{\max} 209 (log ε 4.49); 258 (log ε 3.87) (MeOH).


Ponglux, D. *et al.*, *Tet. Lett.*, 1988, **29**, 5395-5396 (*Elegansamine*, *cryst struct*)

Xu, Y.-K. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1347-1350 (*14-Hydroxyelegansamine*)

Eleuthoside A†

E-64

[180692-76-6]



C₃₆H₄₈N₂O₁₁ 684.782

Constit. of *Eleutherobia aurea*. Inhibits tubulin polym. induction. Shows strong cytotoxic and antitumour effects. Oil. [α]_D²⁵ -9 (c, 0.2 in CHCl₃).

3''-De-Ac: Desmethyleleutherobin. Desmethyleleutherobin [259670-83-2]

C₃₄H₄₆N₂O₁₀ 642.745

Constit. of *Erythropodium caribaeorum*.

4-Me ether, 2''-de-Ac: Isoeleutherobin A [259728-67-1]

C₃₅H₄₈N₂O₁₀ 656.772

Constit. of *Erythropodium caribaeorum*.

4-Me ether, 3''-de-Ac: Eleutherobin

[174545-76-7]

C₃₅H₄₈N₂O₁₀ 656.772

Isol. from the soft coral *Eleutherobia* cf. *albiflora* and *Erythropodium caribaeorum*. Cytotoxin with microtubule stabilising props. Shows potent antitumour activity. Amorph. solid. [α]_D²⁵ -49.3 (c, 3 in MeOH). λ_{\max} 290 (log ε 3.82) (MeOH).

4-Me ether, 2'',3''-dide-Ac: Deacetyeleutherobin

[213824-30-7]

C₃₃H₄₆N₂O₉ 614.734

Constit. of *Erythropodium caribaeorum*.

2'Z-Isomer, 4-Me ether, 3''-de-Ac: Z-Eleutherobin

[259670-84-3]

C₃₅H₄₈N₂O₁₀ 656.772

Constit. of *Erythropodium caribaeorum*.

Schiff, P.B. *et al.*, *Nature (London)*, 1979, **277**, 665-667 (activity)

U.S. Pat., 1995, 5 473 057; *CA*, **124**, 194297

(*Eleutherobin*)

Ketzinel, S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 873-875 (*Eleuthoside A*)

Lindel, T. *et al.*, *J.A.C.S.*, 1997, **119**, 8744-8745 (*Eleutherobin*)

Long, B.H. *et al.*, *Cancer Res.*, 1998, **58**, 1111-1117 (*Eleutherobin*, activity)

Bernardelli, P. *et al.*, *Heterocycles*, 1998, **49**, 531-556 (rev)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 1998, **120**,

8661-8673; 8674-8680 (synth, activity)

Gutteridge, C. E. *et al.*, *J.A.C.S.*, 1999, **121**,

6563-6579 (synth)

Rayl, A.J.S. *et al.*, *The Scientist*, 1999, **13**, 1-3

(clinical)

Roberge, M. *et al.*, *Cancer Res.*, 2000, **60**,

5052-5058 (activity)

Cinel, B. *et al.*, *Org. Lett.*, 2000, **2**, 257-260

(*Deacetyeleutherobin*, *Isoeleutherobin A*,

Desmethyleleutherobin, *Z-Eleutherobin*)

Britton, R. *et al.*, *Tet. Lett.*, 2001, **42**, 2953-

2956 (isol, biosynth)

Castoldi, D. *et al.*, *Angew. Chem., Int. Ed.*,

2005, **44**, 588-591 (*Eleutherobin*, synth)


Castoldi, D. *et al.*, *Chem. Eur. J.*, 2006, **12**, 51-

62 (synth)

Eliamide

E-65

[215714-29-7]



C₂₃H₃₅NO₄ 389.534


Prod. by *Sorangium cellulosum*. Exhibits cytostatic, nematocidal and antifungal activities. Oil. λ_{\max} 223 (ε 19200); 234 (ε

19300) (MeOH).

Ger. Pat., 1998, 19 718 843; CA, 130, 3107d

Elisabethamine**E-66**


[294202-41-8]

C₂₁H₃₃NO₂ 331.497

Alkaloid from *Pseudopterogorgia elisabethae*. Cytotoxic agent. Yellow gum. $[\alpha]_D^{20} +89$. Synth. studies show that the assigned struct. is probably wrong. λ_{\max} 292 (no solvent reported).

Ata, A. et al., *Tet. Lett.*, 2000, 41, 5821-5825 (isol, pmr, cmr)Dai, X. et al., *J.O.C.*, 2007, 72, 1895-1900 (synth)**Ellipticine****E-67**

5,11-Dimethyl-6H-pyrido[4,3-b]carbazole, 9CI. NSC 71795 [519-23-3]

C₁₇H₁₄N₂ 246.311

Numbering systems vary. Systematic CAS numbering (shown) is now more usual. Alkaloid from *Ochrosia elliptica*, *Ochrosia moorei* and several other *Ochrosia* spp., *Senecio* spp., *Strychnos dinklagei*, *Aspidosperma subincanum*, *Bleekeria coccinea* and *Bleekeria vitiensis* (preferred genus name *Ochrosia*). Shows broad-spectrum antineoplastic activity. An intercalating agent, also shows anti-trypanosomal activity. Cytochrome P-450 (CYPIA) inhibitor. Yellow needles (MeOH), orange rods or rosettes (AcOH or CHCl₃). Mp 311-315° dec. Log P 4.37 (calc). λ_{\max} 224 (ε 30200); 237 (sh) (ε 26900); 245 (sh) (ε 21900); 275 (ε 57500); 285 (ε 75900); 293 (ε 70800); 331 (ε 7080); 346 (ε 5010); 380 (ε 3800); 400 (ε 3800) (EtOH) (Derrep). λ_{\max} 239 (ε 17000); 277 (ε 40700); 286 (ε 57500); 294 (ε 55000); 332 (ε 4460); 382 (ε 4080); 400 (ε 3400) (EtOH) (Berdy).

▶ LD₅₀ (mus, orl) 178 mg/kg. LD₅₀ (mus, ivn) 20 mg/kg Exp. adverse cardiovascular effects. Mutagen. UU8825000

N²-Oxide: **Ellipticine N²-oxide**

[37687-33-5]

C₁₇H₁₄N₂O 262.31Alkaloid from *Ochrosia vieillardii*,

Ochrosia moorei and *Strychnos dinklagei* (Apocynaceae, Loganiaceae).

Noncryst.

N²-Me: **N-Methylellipticine**C₁₈H₁₇N₂[⊕] 261.346

Minor quaternary alkaloid from the bark of *Aspidosperma subincanum* (Apocynaceae). Mp 360° dec. (as iodide).

8-Hydroxy: **8-Hydroxy-5,11-dimethylpyrido[4,3-b]carbazole**. 8-Hydroxyellipticine

[72236-82-9]

Minor metab. of ellipticine by *Aspergillus alliaceus*. Yellow prisms (MeOH/EtOAc). Mp 268-270° dec.

12-Hydroxy: **12-Hydroxyellipticine**

[83329-79-7]

C₁₇H₁₄N₂O 262.31

Alkaloid from the stem bark of *Strychnos dinklagei* (Loganiaceae). Noncryst. λ_{\max} 309 (EtOH/HCl). λ_{\max} 278 (sh); 289; 295; 315 (sh) (EtOH).

8-Methoxy: **8-Methoxy-5,11-dimethylpyrido[4,3-b]carbazole**. 8-Methoxyellipticine

[5263-05-8]

C₁₈H₁₆N₂O 276.337

Alkaloid isol. from leaves of *Ochrosia elliptica*. Yellow needles (EtOAc). Mp 280-285° dec.

9-Methoxy: see 9-Methoxyellipticine, M-252

3,4-Dihydro: **3,4-Dihydroellipticine**

[37687-32-4]

C₁₇H₁₆N₂ 248.327

Alkaloid from *Aspidosperma subincanum*, *Ochrosia balansae*, *Ochrosia vieillardii*, *Ochrosia moorei*, *Strychnos dinklagei*, *Pterotaberna inconspicua*, *Hazunta modesta* and other *Hazunta* spp. (Apocynaceae, Loganiaceae). Cubes (MeOH). Mp 293-296° dec.

3,4-Dihydro, N²-Me: **3,4-Dihydro-2-methylellipticine**. N-Methyl-3,4-dihydroellipticine

C₁₈H₁₉N₂[⊕] 263.361

Minor quaternary alkaloid from the bark of *Aspidosperma subincanum* (Apocynaceae). Bright yellow needles (Me₂CO aq.) (as picrate). Mp 296-306° dec. (as iodide) Mp 273-275° dec. (as picrate).

1,2,3,4-Tetrahydro: **1,2,3,4-Tetrahydroellipticine**

[6538-51-8]

C₁₇H₁₈N₂ 250.343

Alkaloid from *Ochrosia balansae*, *Ochrosia vieillardii*, *Strychnos dinklagei* and *Pterotaberna inconspicua* (Apocynaceae, Loganiaceae).

1,2,3,4-Tetrahydro, N²-Me: **Tetrahydro-2-methylellipticine**. u-Alkaloid B. N-Methyltetrahydroellipticine

C₁₈H₂₀N₂ 264.369

Alkaloid from *Aspidosperma ulei* and *Aspidosperma subincanum* (Apocynaceae). Cryst. (MeOH). Mp 215-218° dec.

13-Oxo: **11-Methyl-6H-pyrido[4,3-b]carbazole-5-carboxaldehyde**. **13-Oxoellipticine**

[77251-57-1]

C₁₇H₁₂N₂O 260.295

Alkaloid from the stem bark of *Strychnos dinklagei* (Loganiaceae). Yellow needles (CHCl₃). Mp 275-276°.

13-Oxo, N²-oxide: **13-Oxoellipticine N^b-oxide**

[83329-78-6]

C₁₇H₁₂N₂O₂ 276.294

Alkaloid from the stem bark of *Strychnos dinklagei* (Loganiaceae). Noncryst.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 473A (nmr)

Schmutz, J. et al., *Helv. Chim. Acta*, 1958, 41, 288 (*N-Methyltetrahydroellipticine*)Goodwin, S. et al., *J.A.C.S.*, 1959, 81, 1903-1908 (*8-Methoxyellipticine, isol, uv, ir, pmr*)Woodward, R.B. et al., *J.A.C.S.*, 1959, 81, 4434-4435 (*Ellipticine, synth, struct*)Büchi, G. et al., *Tetrahedron*, 1961, 15, 167

(3,4-Dihydroellipticine, iso, struct, synth)

Budzikiewicz, H. et al., *Structure Elucidation of Natural Products by Mass Spectrometry*, Holden-Day Inc., 1964, 1, 53 (ms)Loder, J.W. et al., *Aust. J. Chem.*, 1966, 19, 1947 (ms)Dalton, L.K. et al., *Aust. J. Chem.*, 1967, 20, 2715 (synth, pmr)Bruneton, J. et al., *Phytochemistry*, 1972, 11,3073-3075 (*Ellipticine N-oxide, 3,4-**Dihydroellipticine, 1,2,3,4-**Tetrahydroellipticine*)Courseille, C. et al., *Acta Cryst. B*, 1974, 30, 2628 (cryst struct)Sainsbury, M. et al., *J.C.S. Perkin I*, 1976,

1155 (synth)

Oikawa, Y. et al., *J.C.S. Perkin I*, 1976, 1479

(synth)

Sainsbury, M. et al., *Synthesis*, 1977, 437

(rev)

Bergman, J. et al., *Tet. Lett.*, 1977, 4663

(synth)

Ahond, A. et al., *Tetrahedron*, 1978, 34, 2385-

2388 (cmr)

Kohn, K.W. et al., *Antibiotics (N.Y.)*, 1979, 5,

195 (rev, pharmacol)

Joule, J.A. et al., *Chem. Comm.*, 1979, 642

(synth, bibl)

Michel, S. et al., *Tet. Lett.*, 1980, 21, 4027 (*13-**Oxoellipticine*)Ahond, A. et al., *J. Nat. Prod.*, 1981, 44, 193-199 (*Ellipticine N-oxide, 3,4-**Dihydroellipticine*)Dolman, D. et al., *Tet. Lett.*, 1981, 22, 2119-2120 (*8-Methoxyellipticine, synth*)Besselièvre, R. et al., *Tetrahedron, Suppl.*,

1981, 241 (synth)

Kano, S. et al., *Heterocycles*, 1982, 19, 1673

(synth)

Michel, S. et al., *J. Nat. Prod.*, 1982, 45, 489-494 (*12-Hydroxyellipticine, Ellipticine N-**oxide, 13-Oxoellipticine, 13-Oxoellipticine N-**oxide, 3,4-Dihydroellipticine, 1,2,3,4-**Tetrahydroellipticine*)Sainsbury, M. et al., *J.C.S. Perkin I*, 1982, 587-590 (*8-Hydroxyellipticine, 8-**Methoxyellipticine*)Saulnier, M.G. et al., *J.O.C.*, 1982, 47, 2810

(synth)

Sainsbury, M. et al., *Org. Magn. Reson.*, 1982,

18, 117 (cmr)

Morfaux, A.M. et al., *Phytochemistry*, 1982,21, 1767-1769 (*3,4-Dihydroellipticine,**1,2,3,4-Tetrahydroellipticine, isol*)Miller, R.B. et al., *J.O.C.*, 1983, 48, 886

(synth)

Murakami, Y. et al., *Tet. Lett.*, 1983, 24, 2189

(synth)

Saulnier, M.G. et al., *Tet. Lett.*, 1983, 24, 3831


(13-Oxoellipticine, synth, pmr)

- Garlich, J.R. *et al.*, *Acta Cryst. C*, 1984, **40**, 1871 (*oxide, cryst struct*)
- Gribble, G.W. *et al.*, *J.O.C.*, 1984, **49**, 4518; 1992, **57**, 5891 (*synth*)
- Weller, D.D. *et al.*, *Tet. Lett.*, 1984, **25**, 2105 (*synth*)
- Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 116 (*rev, pharmacol*)
- Gansser, C. *et al.*, *Farmaco, Ed. Sci.*, 1985, **40**, 459-476 (*8-Methoxyellipticine, synth*)
- Obaza-Nutaitis, J.A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 449 (*13-Oxoellipticine, synth*)
- Kansal, V.K. *et al.*, *Tetrahedron*, 1986, **42**, 2389-2408 (*rev*)
- May, C. *et al.*, *J.C.S. Perkin 1*, 1988, 247 (*synth*)
- Miller, R.B. *et al.*, *Tet. Lett.*, 1989, **30**, 297 (*synth*)
- Hibino, S. *et al.*, *J. Het. Chem.*, 1990, **27**, 1751 (*synth*)
- Yokoyama, Y. *et al.*, *J.C.S. Perkin 1*, 1990, 1319 (*synth*)
- Bäckvall, J.-E. *et al.*, *J.O.C.*, 1990, **55**, 4528 (*synth*)
- Davis, D.A. *et al.*, *Tet. Lett.*, 1990, **31**, 1081 (*synth*)
- Modi, S.P. *et al.*, *Tetrahedron*, 1991, **47**, 6539-6548 (*12-Hydroxyellipticine, synth*)
- Froelich-Ammon, S.J. *et al.*, *J. Biol. Chem.*, 1995, **270**, 14998 (*pharmacol*)
- Mustafin, A.G. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 1999, **48**, 2121-2126 (*synth*)
- Ishikura, M. *et al.*, *Heterocycles*, 2000, **53**, 11-14 (*synth*)
- Diaz, M. *et al.*, *Eur. J. Org. Chem.*, 2001, 4543-4549 (*synth*)
- Miki, Y. *et al.*, *J.C.S. Perkin 1*, 2001, 2213-2216 (*synth*)
- Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 111-116 (*synth, pmr, cmr*)
- Liu, C.-Y. *et al.*, *J.O.C.*, 2007, **72**, 7106-7115 (*synth*)
- Dracinsky, M. *et al.*, *Tet. Lett.*, 2007, **48**, 6893-6895 (*synth*)
- Mal, D. *et al.*, *Tetrahedron*, 2007, **63**, 3768-3781 (*Ellipticine, synth*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EA1850; HKH000

Elliptinine

E-68

[523-16-0]




C₂₀H₂₄N₂O₂ 324.422

Proposed struct.; dubious. Alkaloid from *Ochrosia elliptica* (Apocynaceae). Cryst. (MeOH/C₆H₆). Mp 231-233°. [α]_D²⁴ -255 (c, 0.2 in EtOH).

Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 1903-1908

Elliptinol†

E-69



C₁₈H₂₄N₂O₃ 316.399


Alkaloid from *Aglaia elliptifolia*. Cytotoxic. Powder. Mp 162-163°. [α]_D²² +38.6 (c, 0.05 in CHCl₃). λ_{max} 283 (log ε 4.66) (MeOH).

Wang, S.-K. *et al.*, *J. Nat. Prod.*, 2001, **64**, 92-94

Elmerrillicine

E-70

6,7,7a,8-Tetrahydro-4-methoxy-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-12-ol, 9CI. 11-Hydroxy-3-methoxy-1,2-methylenedioxyynoraporphine



C₁₈H₁₇NO₄ 311.337

(R)-form [61619-35-0]

Alkaloid from the bark of *Elmerrillia papuana* (Magnoliaceae).

N-Ac:

Prisms (Et₂O). Mp 201-203°. [α]_D -474.

N,O-Di-Ac:

Prisms (Et₂O). Mp 210-212°. [α]_D -287.

N-Me: 11-Hydroxy-3-methoxy-1,2-methylenedioxyaporphine. **N-Methylel-merrillicine**

[107882-27-9]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the trunk bark of *Guatteria sagotiana* (Annonaceae). Amorph. [α]_D -73 (c, 0.26 in EtOH).


Cleaver, L. *et al.*, *Aust. J. Chem.*, 1976, **29**, 2003 (*isol, uv, ir, pmr, ms, struct*)

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1078 (*isol, uv, pmr, ms, struct, deriv*)

Emeniveol

E-71

[146001-22-1]



C₂₈H₃₉NO 405.622


Alkaloid from *Emericella nivea*. Pollen growth inhibitor. Needles (MeOH). Mp 179°. [α]_D²⁰ -91 (c, 1 in MeOH). λ_{max} 224 (ε 16600); 277 (sh) (ε 5620); 283 (ε 6170); 291 (ε 5620) (MeOH) (Derep).

Kimura, Y. *et al.*, *Tet. Lett.*, 1992, **33**, 6987 (*isol, pmr, cmr, cryst struct*)

Emetine, BAN

E-72

6',7',10,11-Tetramethoxyemetan, 9CI. *Ipecine. Methylcephaline. NSC 33669* [483-18-1] [65910-27-2 ((±)-form)]



C₂₉H₄₀N₂O₄ 480.646

Alkaloid from *Alangium lamarckii*, *Alangium lonorchii*, *Cephaelis acuminata*, *Psychotria granadensis* and *Uragoga ipe-cachuana* (preferred genus name *Psychotria*) (Alangiaceae, Rubiaceae). The occurrence of Emetine in the common ivy, *Hedera helix* (Araliaceae) and in the roots of *Borreria verticillata* (Rubiaceae) remains unconfirmed. Inhibitor of RNA, DNA and protein synthesis. Orally-active emetic. Amoebicide, general parasiticide, shows antiviral, antineoplastic and antibacterial activity. Also shows neuromuscular and cardiovascular effects. *A. lamarckii* has been extensively used in Indian medicine and the biochemistry of Emetine has been comprehensively studied. Amorph. Mp 74° efferv. (sinters at 70°). [α]_D -46.55 (c, 1.16 in CHCl₃) (-50, -26). [α]_D -22 (EtOH). Log P 4.55 (calc).

▶ Severe skin and eye irritant. Adverse systemic effects by ingestion and intravenous routes. Cumulative poison. Cardiotoxic. Reported fatal dose approx. 2g. LD₅₀ (rat, orl) 68 mg/kg.

Hydrochloride (1:2): Emetine hydrochloride, USAN. Emetinal. Encol. Er- ketin. Hemometina

[316-42-7]

Needles + 7H₂O. Mp 255° dec. (sinters at 235°).

▶ Skin, eye and respiratory tract irritant. Local and cumulative systemic effects when used therapeutically. Cardiotoxic, nephrotoxic, hepatotoxic. LD₅₀ (rat, orl) 0.012 mg/kg. LD₅₀ (rat, ipr) 17 mg/kg. JY5250000

Hydrobromide (1:2):

Needles + 4H₂O. Mp 250-260° (sinters at 245°).

Sulfate:

Needles + 7H₂O. Mp 245° (sinters at 205°).

Camphorsulfonate salt: Canforemetina.

Emetoplisc

[53625-86-8] Antiamoebic agent.

Cryst. (EtOH/Et₂O). Mp 203-204° Mp 134-135°.

N-Ac:

Granules (Et₂O). Mp 97-99°.

O^{6'}-De-Me: Cephaline. Desmethylemetine. Dihydropsychotrine. Alangine B

[483-17-0]

[5884-41-3, 1353-93-1 (Alangine B)]

C₂₈H₃₈N₂O₄ 466.619


- Alkaloid from *Alangium lamarckii*, *Alangium salviifolium*, *Psychotria granadensis*, *Psychotria ipeacacuanha* and *Cephaelis acuminata*. Shows antitumour activity, inhibitor of RNA, DNA and protein biosynth. Emetic and expectorant. Antileishmanial and antimalarial agent. Sol. MeOH, CHCl₃, acids; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 115-116° (104-107°; 110-111°). [α]_D²⁵ -43.4 (CHCl₃). [α]_D -23.4 (c, 0.81 in CHCl₃). Log P 4.27 (calc).
- 0⁶-De-Me, hydrochloride:** Mp 254-257°. [α]_D +21 (H₂O).
- 0⁶-De-Me, 2'-N-(1-deoxy-β-D-fructopyranos-1-yl): 2'-N-(1-Deoxy-β-D-fructopyranos-1-yl) cephaeline** [261515-32-6]
C₃₄H₄₈N₂O₉ 628.761
Alkaloid from *Cephaelis acuminata*. Amorph. powder. [α]_D²⁶ -58 (c, 0.46 in CHCl₃). λ_{max} 209 (log ε 4.5); 223 (sh) (log ε 4.11); 285 (log ε 3.78) (MeOH).
- 0⁶-De-Me, O,N-di-Ac:** Mp 116°.
- 0⁶-De-Me, N-(methylaminocarbonyl): Alangamide** [24529-94-0]
C₃₀H₄₁N₃O₅ 523.671
Alkaloid from the seeds of *Alangium lamarckii*. Needles (EtOH aq. or Et₂O/EtOH). Mp 213°. [α]_D -45 (c, 0.82 in CHCl₃). Possible artifact.
- 0⁶-De-Me, N-Me: Alamarckine** [6793-51-7]
C₂₉H₄₀N₂O₄ 480.646
Semisynthetic; from N-methylation of the total crude alkaloids of *Alangium lamarckii* or of Cephaeline. Cryst. (MeOH). Mp 191-192°. [α]_D -50.5 (c, 1.4 in CHCl₃).
- 0⁶,O⁷-Di-de-Me: 7'-O-Demethylcephaeline** [261515-31-5]
C₂₇H₃₆N₂O₄ 452.592
Alkaloid from *Cephaelis acuminata*. Amorph. powder. [α]_D²⁵ -3.4 (c, 0.56 in MeOH). λ_{max} 208 (log ε 4.54); 229 (sh) (log ε 4.08); 286 (log ε 3.81) (MeOH).
- 0⁶,O⁹-Di-de-Me: 9-O-Demethylcephaeline** [82570-80-7]
Mp 147° (sinters at 124°). [α]_D¹² -55 (c, 0.50 in CHCl₃).
- 0⁶,O¹⁰-Di-de-Me: 10-O-Demethylcephaeline** [29700-91-2]
C₂₇H₃₆N₂O₄ 452.592
Alkaloid from *Alangium longiflorum* and *Cephaelis acuminata*. Cytotoxic. Mp 148° (sinters at 129-130°) (synthetic). [α]_D¹⁷ -53 (c, 0.50 in CHCl₃) (synthetic). λ_{max} 209 (log ε 4.5); 225 (sh) (log ε 4.11); 286 (log ε 3.83) (MeOH).
- 0⁶,O²-Di-de-Me: Demethylcephaeline** [30901-62-3]
C₂₇H₃₆N₂O₄ 452.592
Alkaloid from the stem bark of *Alangium lamarckii*. Amorph. Mp 147-149°. [α]_D -53.5 (CHCl₃). This is probably identical with O¹⁰-Demethylcephaeline above, but this is not yet confirmed (1999). Reported phys. props. identical. λ_{max} 211 (log ε 4.28); 225 (sh) (log ε 3.91); 286 (log ε 3.69) (EtOH). λ_{max} 213 (sh) (log ε 3.48); 227 (log ε 3.96); 247 (log ε 4.05); 301 (log ε 3.81) (EtOH/NaOH).
- 1',2'-Didehydro: O-Methylpsychotrine** [523-01-3]
C₂₉H₃₈N₂O₄ 478.63
Alkaloid from *Psychotria ipeacacuanha*, *Psychotria granadensis* and *Cephaelis ipeacacuanha*. Mp 123-124°. [α]_D +43.2 (c, 1 in EtOH). λ_{max} 241 (ε 18800); 288 (ε 7244); 305 (ε 8320); 354 (ε 8130) (MeOH-HCl) (Berdy). λ_{max} 226 (ε 26900); 278 (ε 9120); 307 (ε 5760) (MeOH-NAOH) (Berdy).
- JY4376000
- 1',2'-Didehydro, oxalate salt:** Mp 161° dec. [α]_D +42 (c, 1.0 in H₂O).
- 1',2'-Didehydro, O⁶-de-Me: Psychotrine** [7633-29-6]
C₂₈H₃₆N₂O₄ 464.603
Alkaloid from the seeds, root bark and stem bark of *Alangium lamarckii*, the roots and branches of *Alangium salviifolium*, and from *Psychotria ipeacacuanha* and *Cephaelis ipeacacuanha*. Shows cytotoxic activity. HIV reverse transcriptase inhibitor. Yellow prisms (Me₂CO aq.). Mp 138° (130-133°, 117-120°). [α]_D +80.2 (+75.50) (MeOH). [α]_D²⁰ +68.4 (c, 1.0 in EtOH). λ_{max} 240 (ε 13900); 288 (ε 3700); 306 (ε 6250); 356 (ε 6800) (MeOH/HCl) (Berdy).
- 1',2'-Didehydro, O⁶-de-Me, oxalate salt:** Mp 131-145°. [α]_D +21.6 (H₂O).
- 1',2'-Didehydro, O⁶,O⁹-di-de-Me: 9-Demethylpsychotrine** [16531-03-6]
[73053-57-3]
C₂₇H₃₄N₂O₄ 450.577
Alkaloid from the root bark and stem bark of *Alangium lamarckii* (Alangiaceae). Dark-yellow granules (EtOH). Mp 166-168°. [α]_D +67.9 (c, 0.50 in MeOH).
- 1',2',3',4'-Tetrahydro: Emetamine** [483-19-2]
C₂₉H₃₆N₂O₄ 476.614
Alkaloid from *Psychotria granadensis* and *Psychotria ipeacacuanha* (Rubiaceae). Needles (EtOAc). Mp 155-156°. [α]_D +12.3 (EtOH).
- 1'-Hydroxy, O⁶,O⁹-di-de-Me: Klugine. 1'-Hydroxy-9-O-demethylcephaeline** C₂₇H₃₆N₂O₅ 468.592
Alkaloid from the stem bark of *Psychotria klugii*. Light brown cryst. (MeCN/CH₂Cl₂). Mp 288-290°. [α]_D -66 (c, 1 in MeOH). λ_{max} 208 (log ε 4.35); 244 (log ε 3.9); 315 (log ε 3.75); 328 (log ε 3.8); 450 (sh) (log ε 3.65) (MeOH).
- 8-Hydroxy, 1',2'-didehydro, O⁶-de-Me: Alangicine** [16531-04-7]
C₂₈H₃₆N₂O₅ 480.603
Alkaloid from the root bark of *Alangium lamarckii* (Alangiaceae). Pale-yellow granules (EtOH). Mp 147-148°. [α]_D +64.1 (c, 0.26 in MeOH).
- 6'-Demethoxy, 8'-hydroxy: Neocephaeline** [261515-30-4]
C₂₈H₃₈N₂O₄ 466.619
Alkaloid from *Cephaelis acuminata*. Amorph. powder. [α]_D²⁵ -87 (c, 1 in CHCl₃). λ_{max} 207 (log ε 4.76); 230 (sh) (log ε 4.12); 282 (log ε 3.71) (MeOH).
- 6'-Demethoxy, 8'-hydroxy, 2'-N-(1-deoxy-β-D-fructopyranos-1-yl): 2'-N-(1-Deoxy-β-D-fructopyranos-1-yl)neocephaeline** [261515-33-7]
C₃₄H₄₈N₂O₉ 628.761
Alkaloid from *Cephaelis acuminata*. Amorph. powder. [α]_D²⁶ -50 (c, 0.18 in CHCl₃). λ_{max} 207 (log ε 4.62); 224 (sh) (log ε 4.05); 283 (log ε 3.55) (MeOH).
- 1'-Epimer, O⁶-de-Me: Isocephaeline** [5884-45-7]
[17132-43-3]
C₂₈H₃₈N₂O₄ 466.619
Alkaloid from the seeds of *Alangium lamarckii* and *Cephaelis acuminata*. Cryst. (Et₂O). Mp 116° (shrinks from 108°). [α]_D²⁰ -69.8 (CHCl₃).
- 1'-Epimer, O⁶,O⁷-di-de-Me: 7'-O-Demethylisocephaeline** C₂₇H₃₆N₂O₄ 452.592
Alkaloid from the stem bark of *Psychotria klugii*. Light brown cryst. (MeCN/CH₂Cl₂). Mp 214-216°. [α]_D -47 (c, 1 in MeOH). λ_{max} 204 (log ε 4.3); 242 (log ε 3.85); 282 (sh) (log ε 3.1); 305 (log ε 3.65); 316 (log ε 3.68); 460 (log ε 3.55) (MeOH).
- 1'-Epimer, O⁶,O¹⁰-di-de-Me: 10-O-Demethylisocephaeline** C₂₇H₃₆N₂O₄ 452.592
Alkaloid from the stem bark of *Alangium longiflorum*. Needles. Mp 114-116°. [α]_D³⁰ -12.8 (c, 0.81 in MeOH).
- Pyman, F.L. *et al.*, *J.C.S.*, 1917, **111**, 419-446 (*Emetine*)
Brossi, A. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 1515-1522 (*Emetine, synth, resohn*)
Van Tamelen, E.E. *et al.*, *J.A.C.S.*, 1959, **81**, 507 (*Emetine, stereochem, bibl*)
Battersby, A.R. *et al.*, *J.C.S.*, 1959, 1744-1748; 1748-1753; 2704-2711; 3512-3521; 1960, 717-725 (*Emetine, Cephaeline, Psychotrine, O-Methylpsychotrine, isol, config, synth*)
Openshaw, H.T. *et al.*, *J.C.S.*, 1963, 1461-1471 (*synth*)
Pakrashi, S.C. *et al.*, *Indian J. Chem.*, 1964, **2**, 379-380 (*Alangine B, Alamarckine*)
Budzikiewicz, H. *et al.*, *Tetrahedron*, 1964, **20**, 399-408 (*Emetine, Psychotrine, Emetamine, isol, ms*)
Teitel, S. *et al.*, *J.A.C.S.*, 1966, **88**, 4068-4071 (*Psychotrine, synth, uv, ir, pmr, ms*)
Szántay, C. *et al.*, *J.O.C.*, 1966, **31**, 1447-1451 (*synth*)
Pakrashi, S.C. *et al.*, *Tet. Lett.*, 1967, 2143-2146 (*Alangicine, 9-Demethylpsychotrine, uv, ir, ms*)
Battersby, A.R. *et al.*, *Chem. Comm.*, 1968, 134-135 (*biosynth*)
Pakrashi, S.C. *et al.*, *Indian J. Chem.*, 1969, **7**, 635-636 (*Alangamide*)
Pakrashi, S.C. *et al.*, *Experientia*, 1970, **26**, 933-934 (*Demethylcephaeline*)
Garg, A.K. *et al.*, *Phytochemistry*, 1972, **11**, 689-695 (*biosynth*)
Koch, M.C. *et al.*, *J.O.C.*, 1975, **40**, 2836-2838 (*Emetine, cmr*)
Mahran, G.H. *et al.*, *Planta Med.*, 1975, **27**, 127-132 (*Emetine, isol, uv, ir, pmr, ms*)

- Nagakura, N. *et al.*, *Chem. Comm.*, 1978, 896-898 (biosynth)
- Takano, S. *et al.*, *Heterocycles*, 1979, **12**, 765-770 (synth)
- Schuij, C. *et al.*, *J.C.S. Perkin 1*, 1979, 970-975 (O-Methylpsychotrine)
- Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1979, 1211-1217 (synth)
- Achari, B. *et al.*, *Planta Med. (Suppl.)*, 1980, 5-7 (Isocephaline)
- Feyns, L.V. *et al.*, *Anal. Profiles Drug Subst.*, 1981, **10**, 289-335 (Emetine, rev. uv, ir, pmr, cmr, ms, anal)
- Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 598-609; 1983, **31**, 2583-2592; 1985, **33**, 144-151; 583 (Psychotrine, Alangicine, uv, ms, pmr, cd, cmr, struct, synth)
- Fujii, T. *et al.*, *Heterocycles*, 1982, **19**, 857-860 (Demethylcephaelines)
- Fujii, T. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 31-50 (bibl. pharmacol)
- Brown, R.T. *et al.*, *Tet. Lett.*, 1984, **25**, 3127-3130 (synth)
- Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 52-57 (antitumour activity)
- Naito, T. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 3530-3533 (synth)
- Hirai, Y. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 1343-1350 (synth)
- Ihara, M. *et al.*, *J.C.S. Perkin 1*, 1990, 1469-1476 (synth)
- Guiles, J.W. *et al.*, *J.O.C.*, 1991, **56**, 6873-6878 (synth)
- Pan, S.J. *et al.*, *Toxicology*, 1995, **97**, 93-104 (tox)
- Ito, A. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1346-1348 (isol, activity)
- Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 582; 1062
- Itoh, A. *et al.*, *Phytochemistry*, 1999, **52**, 1169-1176 (Emetine, Cephaeline, Isocephaline, Neocephaline, O-Demethylcephaelines, 2'-N-(1-deoxyfructopyranosyl))
- Muhammed, I. *et al.*, *J. Nat. Prod.*, 2003, **66**, 962-967 (Klugine, 7'-O-Demethylisocephaline)
- Tietze, L.F. *et al.*, *Chem. Eur. J.*, 2004, **10**, 2722-2731 (synth)
- Itoh, T. *et al.*, *Org. Lett.*, 2006, **8**, 1295-1297 (synth)
- Sakurai, N. *et al.*, *Phytochemistry*, 2006, **67**, 894-897 (10-O-Demethylisocephaline)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, EAL500; EAN000

Emiline

E-73

[36506-99-7]

C₁₉H₂₇NO₆ 365.425

Struct. revised in 1987. Alkaloid from *Emilia flammea* (Asteraceae). Needles (Et₂O). Mp 105-107° (103-105°). [α]_D²⁰ -13.1 (CHCl₃). [α]_D^{22.5} -17.5 (CHCl₃).

►Carcinogenic.

Hydrochloride:

Needles (EtOH). Mp 248-251° (243-245°).

Picrate: Mp 245-247°.


[36506-98-6, 36506-81-7]

- Tomczyk, H. *et al.*, *CA*, 1972, **77**, 19848u (isol)
- Barbour, R.H. *et al.*, *Phytochemistry*, 1987, **26**, 2430 (isol, ir, pmr, cmr, ms, struct)
- Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 640

Emindole DA

E-74

Decahydro-5-(1H-indol-3-ylmethyl)-1,4a-dimethyl-6-methylene-1-(4-methyl-3-pentenyl)-2-naphthalenol, 9CI
[110883-36-8]

C₂₈H₃₉NO 405.622

Indolic diterpenoid antibiotic. Isol. from *Emericella desertorum* and *Emericella striata*. Mycotoxin. Prisms (C₆H₆/hexane). Sol. MeOH, Me₂CO; poorly sol. H₂O. Mp 146-147°. [α]_D -30.7 (c, 2.32 in MeOH). λ_{max} 224 (ε 16600); 277 (sh) (ε 5620); 283 (ε 6170); 291 (ε 5620) (MeOH) (Derep). λ_{max} 224 (ε 16600); 283 (ε 6150); 291 (ε 5610) (MeOH) (Berdy).

Ac:

Needles (hexane). Mp 142.5-143.5°.

5-Epimer: **Emindole SA**

[110883-37-9]

C₂₈H₃₉NO 405.622

From *Emericella desertorum* and *Emericella striata*. Amorph. powder.

Mp 58-60°. [α]_D +32 (c, 0.79 in MeOH). λ_{max} 224 (ε 16600); 277 (sh) (ε 5620); 283 (ε 6170); 291 (ε 5620) (MeOH) (Derep).

5-Epimer, Ac:


Leaflets +½H₂O (MeOH). Mp 102°.

- Nozawa, K. *et al.*, *Chem. Comm.*, 1987, 1157 (isol, struct)
- Nozawa, K. *et al.*, *J.C.S. Perkin 1*, 1988, 1689; 2155 (isol, cmr, pmr, struct)
- Rainier, J.D. *et al.*, *Tet. Lett.*, 2000, **41**, 9419-9423 (synth, Emindole SA)
- Fueki, S. *et al.*, *Org. Lett.*, 2004, **6**, 2697-2700 (biosynth, Emindole DA)

Emindole DB

E-75

Dodecahydro-7-(1H-indol-3-ylmethyl)-α,α,6a,10b-tetramethyl-8-methylene-1H-naphtho[2,1-b]pyran-3-methanol, 9CI
[112900-03-5]

C₂₈H₃₉NO₂ 421.622


Metab. of *Emericella desertorum*. Prisms (C₆H₆/hexane). Mp 165.5-166.5°. [α]_D²² -86.3 (c, 0.48 in MeOH). λ_{max} 224 (ε 38000); 283 (ε 7760); 291 (ε 6920) (MeOH) (Berdy).

Nozawa, K. *et al.*, *J.C.S. Perkin 1*, 1988, 1689 (isol, cmr, struct)

Emindole PA

E-76

[158371-92-7]



Probable Absolute Configuration

R¹ = -C(CH₃)₂CH=CH₂, R² = HC₃₃H₄₇NO₂ 489.74

Struct. revised in 2006. Isol. from *Emericella purpurea* IFO 30849. Mycotoxin; angiotensin II receptor antagonist. Amorph. powder.

21-Epimer: **Emindole PC**C₃₃H₄₇NO₂ 489.74

Isol. from *Emericella purpurea* IFO 30849. Cryst. (petrol). Mp 238-240°.

Hosoe, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 185-187 (isol, pmr, cmr, ms)

Emindole PB

E-77

As Emindole PA, E-76 with R¹ = H, R² = -C(CH₃)₂CH=CH₂

C₃₃H₄₇NO₂ 489.74


Isol. from *Emericella purpurea* IFO 30849. Amorph. powder.

Hosoe, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 185-187 (isol, pmr, cmr, ms)

Emindole SB

E-78

[112900-04-6]

C₂₈H₃₉NO 405.622

Abs. config. assumed to be as shown but not explicitly assigned. Prod. by *Emericella striata* and mangrove fungus No. dz17. Amorph. powder. Mp 58-60°. [α]_D¹⁵ +32 (c, 0.79 in MeOH).

O-Ac:

Leaflets + ½H₂O (MeOH). Mp 102°.


Nozawa, K. *et al.*, *J.C.S. Perkin 1*, 1988, 2607-2610 (isol, ir, pmr, cmr, ms)

Huang, Z. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 655-657 (isol, pmr, cmr)

Endophenazine B

E-79

5-Methyl-9-(3-methyl-2-butenyl)-7-oxo-1(5H)-phenazinecarboxylic acid


C₁₉H₁₈N₂O₃ 322.363

Prod. by various strains of the endosymbiotic *Streptomyces anulatus*. Active against gram-positive bacteria and some fungi. Violet solid. λ_{\max} 236 (log ϵ 4.33); 283 (log ϵ 4.44); 374 (log ϵ 3.86); 516 (log ϵ 3.89); 545 (sh) (MeOH).

Gebhardt, K. *et al.*, *J. Antibiot.*, 2002, **55**, 794-800; 801-806 (*isol, pmr, cmr, ms, activity*)

Enduracididine

E-80

 α ,2-Diamino-4,5-dihydro-1H-imidazole-4-propanoic acid, 9CI*(\alpha,S,4R)*-formC₆H₁₂N₄O₂ 172.186

Contains tautomeric guanidino system. CAS name refers to aminoimidazole tautomer.

***(\alpha,S,4R)*-form** [21209-39-2]

Isol. from seeds of *Lonchocarpus sericeus*. Component of Enduracidin A. Mp 235-236° (as hydrobromide).

N²-(6-Bromo-1H-indol-3-ylcarbonyl): N²-(6-Bromo-1H-indol-3-ylcarbonyl)enduracididine

C₁₅H₁₆BrN₅O₃ 394.227

Isol. from the ascidian *Leptoclinidies dubius*. Sol. MeOH, butanol. $[\alpha]_{\text{D}}^{20}$ +16.5 (c, 0.8 in MeOH). λ_{\max} 222 (log ϵ 2.98); 250 (log ϵ 2.63); 282 (log ϵ 2.39) (MeOH). λ_{\max} 222 (ϵ 1122); 250 (ϵ 1350); 282 (ϵ 1288) (MeOH) (Berdy).

(\alpha,R,4R)*-formAlloenduracididine*

[21380-32-5]

Component of Enduracidin A. Mp 249-252° (as hydrobromide).

Horii, S. *et al.*, *J. Antibiot.*, 1968, **21**, 665-667 (*isol*)

Tsujii, S. *et al.*, *Chem. Lett.*, 1975, 1281-1284 (*synth*)


Fellows, L.E. *et al.*, *Phytochemistry*, 1977, **16**, 1957-1959 (*isol, ms, ir, pmr, cmr*)

Garcia, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 782-785 (6-Bromoindolylcarbonylenduracididine)

Engleromycin

E-81

[77784-06-6]

C₂₈H₃₅NO₆ 481.588


Prod. by *Engleromyces goetzei*. Shows antibiotic and cytotoxic props. Needles (EtOH aq.). Mp 226-228°. $[\alpha]_{\text{D}}^{25}$ +64 (EtOH).

Pedersen, E.J. *et al.*, *Tet. Lett.*, 1980, **21**, 5079 (*isol, pmr, cmr, struct*)

Enicoflavine

E-82

[56050-08-9]


C₁₀H₁₃NO₄ 211.217

Alkaloid from *Enicostema hyssopifolium* (Gentianaceae). Unstable yellow liq. V. reactive, converted into Gentianine, G-59, Genticrucine, G-62, and other compds. on standing at r.t. for three weeks. Genus name given as Enicostema.

Chaudhuri, R.K. *et al.*, *Chem. Ind. (London)*, 1975, 127 (*uv, ir, pmr, ms, isol, struct*)

Enkleia

E-83

5-Hydroxy-4,7-dimethoxybenz[*g*]isoquinolin-1(2H)-one, 9CI [139682-16-9]C₁₅H₁₃NO₄ 271.272

Alkaloid from roots of *Enkleia siamensis* (Thymelaeaceae). Needles. Mp 165-168°.

Boonyaratankornkit, L. *et al.*, *Planta Med.*, 1991, **57**, 582 (*isol, uv, ir, pmr, ms, struct*)

Entadamide A

E-84

N-(2-Hydroxyethyl)-3-(methylthio)-2-propenamide, 9CI
MeSCH=CHCONHCH₂CH₂OH
C₆H₁₁NO₂S 161.224

(E)-form [100477-88-1]

Constit. of *Clinacanthus siamensis* and *Entada phaseoloides*. Syrup.

S-Oxide (R-): **Entadamide C**

[121949-96-0]

C₆H₁₁NO₂S 177.224

Constit. of *Clinacanthus siamensis* and *Entada phaseoloides*. Needles (Me₂CO). Mp 144-145°. $[\alpha]_{\text{D}}^{24}$ +186 (c, 0.13 in MeOH). The synthetic racemate has Mp 122-123° (Me₂CO).

O-β-D-Glucopyranoside: Entadamide A β-D-glucoside

[138916-58-2]

C₁₂H₂₁NO₇S 323.366

Constit. of the seeds of *Entada phaseoloides*. Pale brown syrup. $[\alpha]_{\text{D}}$ -12 (c, 0.05 in MeOH).

Ikegami, F. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 5153 (*isol, ir, cmr, pmr*)

Ikegami, F. *et al.*, *Phytochemistry*, 1989, **28**, 881 (*Entadamide C*)

Dai, J. *et al.*, *Phytochemistry*, 1991, **30**, 3749 (*glycoside*)

Tuntiwachwuttikul, P. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1423-1425 (*isol*)

Entadamide B

E-85

N-(2-Hydroxyethyl)-3,3-bis(methylthio)propanamide, 9CI

[110225-60-0]

(MeS)₂CHCH₂CONHCH₂CH₂OHC₇H₁₅NO₂S₂ 209.333


Isol. from the seeds of *Entada phaseoloides* (Fabaceae).

Ikegami, F. *et al.*, *Phytochemistry*, 1987, **26**, 1525 (*isol, ir, pmr, cmr, ms, struct, synth*)

Enterobactin

E-86

N,N',N''-(2,6,10-Trioxo-1,5,9-trioxacyclododecane-3,7,11-triyl)tris[2,3-dihydroxybenzamide], 9CI. Enterochelin [28384-96-5]

C₃₀H₂₇N₃O₁₅ 669.554

Isol. from *Salmonella typhimurium*, *Escherichia coli*, *Klebsiella pneumoniae*, *Klebsiella aerogenes*, *Enterobacter* sp. and *Shigella* sp. Catecholate siderophore, iron chelator. Sol. MeOH, Me₂CO, bases, DMSO; poorly sol. H₂O. Mp 202-203°. $[\alpha]_{\text{D}}$ +7.4 (EtOH). λ_{\max} 316 (ϵ 4390) (EtOH) (Berdy).

Pollack, J.R. *et al.*, *Biochem. Biophys. Res. Commun.*, 1970, **38**, 989-992 (*isol*)

Rogers, H.A. *et al.*, *Biochim. Biophys. Acta*, 1977, **497**, 548-557 (*isol*)

Corey, E.J. *et al.*, *Tet. Lett.*, 1977, **18**, 3919-3922 (*synth*)

Rastetter, W.H. *et al.*, *J.O.C.*, 1981, **46**, 3579-3590 (*synth*)

Shanzer, A. *et al.*, *Chem. Comm.*, 1983, 846-847 (*synth*)

Pecoraro, V. *et al.*, *J.A.C.S.*, 1983, **105**, 4617-4623; 4623-4633 (*isol, props*)


Tor, Y. *et al.*, *J.A.C.S.*, 1992, **114**, 6661-6671 (*pmr, cd, bibl*)

Rogers, H.J. *et al.*, *J.C.S. Perkin I*, 1995, 3073-3075 (*synth, ir, pmr, cmr*)

Marinez, E.R. *et al.*, *J.O.C.*, 1996, **61**, 3548-3550 (*synth*)

Epibatidine**E-87**

2-(6-Chloro-3-pyridinyl)-7-azabicyclo[2.2.1]heptane, 9CI

**(+)-form**

C₁₁H₁₃ClN₂ 208.69

Unique new class of alkaloids. Log P 1.02 (calc). λ_{max} 217; 270 (sh); 274 (ε 5300) (MeOH) (Derep).

(+)-form [140111-52-0]

Alkaloid from the skin extracts of the Ecuadoran poison frog *Epidobates tricolor*. Neuronal nicotinic acetylcholine receptor agonist (most potent known to date). Also active at neuromuscular nicotinic receptors. Potent non-opioid analgesic (potency reported to be 200-500 times greater than that of morphine). Mp 56-57°. [α]_D²³ +5 (c, 0.35 in CHCl₃). Data refer to synthetic material.

Hydrochloride: Mp 150° dec. [α]_D²⁴ +34.7 (c, 0.36 in MeOH).

(-)-form [152378-30-8]

Needles. Mp 56-57° Mp 64°. [α]_D²³ -5 (c, 0.35 in CHCl₃). Similar pharmacol. profile to (+)-form.

Hydrochloride: Mp 130° dec. [α]_D²⁴ -33.7 (c, 0.16 in MeOH).

(±)-form [148152-66-3]

Synthetic. Mp 50-51°.

4-Epimer: [152377-48-5]

Yellow oil. Pharmacol. inactive.

Spande, T.F. *et al.*, *J.A.C.S.*, 1992, **114**, 3475 (*isol*, *pmr*, *ms*, *struct*)

Corey, E.J. *et al.*, *J.O.C.*, 1993, **58**, 5600 (*synth*)

Broka, C.A. *et al.*, *Tet. Lett.*, 1993, **34**, 3251 (*synth*)

Huang, D.F. *et al.*, *Tet. Lett.*, 1993, **34**, 4477 (*synth*, *resoln*)

Clayton, S.C. *et al.*, *Tet. Lett.*, 1993, **34**, 7493 (*synth*)

Rupniak, N.M.J. *et al.*, *Br. J. Pharmacol.*, 1994, **113**, 1487-1493 (*pharmacol*)

Ko, S.Y. *et al.*, *Chem. Comm.*, 1994, 1775 (*synth*)

Okabe, K. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1432 (*synth*)

Watt, A.P. *et al.*, *J. Liq. Chromatogr.*, 1994, **17**, 1256 (*resoln*)

Fisher, M. *et al.*, *J. Pharmacol. Exp. Ther.*, 1994, **270**, 702-707 (*pharmacol*)

Fletcher, S.R. *et al.*, *J.O.C.*, 1994, **59**, 1771 (*synth*, *pmr*, *abs config*)

Senokuchi, K. *et al.*, *Synlett*, 1994, 343 (*synth*)

Sestanjan, K. *et al.*, *Tet. Lett.*, 1994, **35**, 5417 (*synth*)

Albertini, E. *et al.*, *Tet. Lett.*, 1994, **35**, 9297; 1997, **38**, 681-684 (*synth*)

Szantay, C. *et al.*, *Alkaloids (Academic Press)*, 1995, **46**, 95-125 (*rev*, *synth*)

Bonhaus, D.W. *et al.*, *J. Pharmacol. Exp. Ther.*, 1995, **272**, 1199-1203 (*pharmacol*)

Dehmlow, E.V. *et al.*, *J. Prakt. Chem.*, 1995, **337**, 167 (*rev*)

Harnández, A. *et al.*, *J.O.C.*, 1995, **60**, 2683 (*synth*)

Kotian, P.L. *et al.*, *Synth. Commun.*, 1995, **25**, 63 (*synth*, *resoln*)

Sullivan, J.P. *et al.*, *CNS Drug Rev.*, 1996, **2**, 21 (*rev*)

Chen, Z. *et al.*, *Chem. Rev.*, 1996, **96**, 1179 (*rev*, *bibl*)

Sacaan, A.I. *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, **276**, 509-515 (*pharmacol*)

Bai, D. *et al.*, *J.O.C.*, 1996, **61**, 4600 (*synth*)

Zhang, C. *et al.*, *J.O.C.*, 1996, **61**, 7189 (*synth*)

Müller, C.E. *et al.*, *Pharm. Unserer Zeit*, 1996, **25**, 85 (*rev*)

Xu, R. *et al.*, *Tet. Lett.*, 1996, **37**, 1463 (*synth*)

Trost, B.M. *et al.*, *Tet. Lett.*, 1996, **37**, 7485-7488 (*asym synth*)

Szántay, C. *et al.*, *Tetrahedron*, 1996, **52**, 11053 (*synth*, *pmr*, *cmr*)

Ikeda, M. *et al.*, *J.C.S. Perkin 1*, 1997, 3339-3344 (*synth*, *biol*)

Giblin, G.M.P. *et al.*, *J.C.S. Perkin 1*, 1998, 3689-3697 (*synth*)

Aoyagi, S. *et al.*, *J.O.C.*, 1998, **63**, 8397-8406 (*synth*)

Avenoz, A. *et al.*, *Synthesis*, 1998, 1335-1338 (*synth*)

Jones, C.D. *et al.*, *Tet. Lett.*, 1998, **39**, 1021-1022; 1023-1024 (*synth*)

Aoyagi, S. *et al.*, *Tet. Lett.*, 1998, **39**, 4513-4516 (*synth*)

Zhang, C. *et al.*, *J.C.S. Perkin 1*, 1999, 675-676 (*synth*)

Habermann, J. *et al.*, *J.C.S. Perkin 1*, 1999, 1253-1255 (*synth*)

Barros, M.T. *et al.*, *Tet. Lett.*, 1999, **40**, 557-560 (*synth*)

Hall, A. *et al.*, *J.C.S. Perkin 1*, 2000, 329-343 (*synth*)

Cabanal-Duvillard, I. *et al.*, *Tetrahedron*, 2000, **56**, 3763-3769 (*synth*)

Roy, B. *et al.*, *Heterocycles*, 2001, **55**, 861-871 (*synth*)

Barros, M.T. *et al.*, *J.C.S. Perkin 1*, 2001, 166-173 (*synth*)

Evans, D.A. *et al.*, *Org. Lett.*, 2001, **3**, 3009-3012 (*synth*)

Olivio, H.F. *et al.*, *Org. Prep. Proced. Int.*, 2002, **34**, 1-26 (*rev*)


Pandey, G. *et al.*, *Tetrahedron*, 2002, **58**, 3525-3534 (*synth*)

Lee, C.-L.K. *et al.*, *Org. Lett.*, 2005, **7**, 2965-2967 (*synth*)

Armstrong, A. *et al.*, *J.O.C.*, 2007, **72**, 8019-8024 (*synth*)

Epicoccamide**E-88**

[606139-26-8]



n = 14

C₂₉H₅₁NO₉ 557.723

Tetramic acid deriv. Metab. of the fungus *Epicoccum purpurascens* derived from the jellyfish *Aurelia aurita*, and a terrestrial *Epicoccum* sp. Amorph. powder. [α]_D²⁰ -10.3 (c, 0.1 in EtOH). λ_{max} 282 (ε 7940) (EtOH).

3'-Ac: Epicoccamide B

[959855-22-2]

C₃₁H₅₃NO₁₀ 599.76

Prod. by a terrestrial *Epicoccum* sp. Oil. [α]_D²⁵ -94 (c, 0.2 in MeOH). λ_{max} 224; 283 (MeOH).

6'-Ac: Epicoccamide C

[959855-23-3]

C₃₁H₅₃NO₁₀ 599.76


Prod. by a terrestrial *Epicoccum* sp. Oil. [α]_D²⁵ -52.6 (c, 0.1 in MeOH). λ_{max} 224; 283 (MeOH).

Wright, A.D. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 507-510 (*Epicoccamide*)

Wangun, H.V.K. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1800-1803 (*Epicoccamides B-C*)

Epicoccarine B**E-89**

[944339-71-3]



Relative Configuration

C₂₃H₃₁NO₅ 401.502

Tetramic acid deriv. Prod. by an *Epicoccum* sp. associated with *Pholiota squarrosa*. Red oil. [α]_D²⁵ -179.9 (c, 0.1 in MeOH). λ_{max} 228; 280 (MeOH).

6-Deoxy: Epicoccarine A

[944339-69-9]


C₂₃H₃₁NO₄ 385.502

Prod. by an *Epicoccum* sp. associated with *Pholiota squarrosa*. Red oil. [α]_D²⁵ -45.8 (c, 0.15 in MeOH). λ_{max} 228; 280 (MeOH).

Wangun, H.V.K. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1702-1705 (*isol*, *pmr*, *cmr*, *ms*)

Epicoccin B**E-90**

[952585-65-8]



Absolute Configuration


C₁₈H₁₈N₂O₇S₃ 470.547

Prod. by *Epicoccum nigrum* colonising a *Cordyceps* sp. Powder. [α]_D +111 (c, 0.04 in MeOH). λ_{max} 212 (ε 14600) (MeOH).

Zhang, Y. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1522-1525 (*isol*, *pmr*, *cmr*)

Epicoccin D**E-91**

[952585-67-0]



Absolute Configuration

C₁₈H₁₈N₂O₆S₂ 422.482

Prod. by *Epicoccum nigrum* colonising a *Cordyceps* sp. Powder. [α]_D +130 (c, 0.01 in MeOH). λ_{max} 206 (ε 14500) (MeOH).

2,7-Disulfide analogue: Epicoccin A

[952585-64-7]

C₁₈H₁₈N₂O₆S₃ 454.548
Prod. by *Epicoccum nigrum*. Powder.
[α]_D +365 (c, 0.06 in MeOH). λ_{max} 204
(ε 15100) (MeOH).


2,7:2',7'-Bis(disulfide) analogue: **Epico-**
cin C

[952585-66-9]
C₁₈H₁₈N₂O₆S₄ 486.614
Prod. by *Epicoccum nigrum*. Powder.
[α]_D +218 (c, 0.12 in MeOH). λ_{max} 204
(ε 13100) (MeOH).

Zhang, Y. et al., *J. Nat. Prod.*, 2007, **70**, 1522-
1525 (isol, pmr, cmr, cryst struct)

Epiderstatin E-92

4-[3-(3,5-Dimethyl-6-oxo-2-piperidinylidene)-2-oxopropyl]-2,6-piperidinedione,
9CI
[126602-16-2]



Relative
configuration

C₁₅H₂₀N₂O₄ 292.334

Glutarimide antibiotic. Prod. by *Streptomyces pulveraceus* ssp. *epiderstagenes*. Inhibitor of mitogenic activity induced by epidermal growth factor. Powder. Sol. MeOH, CHCl₃, DMSO, EtOH; poorly sol. H₂O, hexane. Mp 185-187°. [α]_D²¹ +5.3 (c, 0.22 in MeOH). λ_{max} 295 (ε 16700) (MeOH) (Derep). λ_{max} (MeOH-HCl) (Berdy).

Osada, H. et al., *J. Antibiot.*, 1989, **42**, 1599;
1607 (isol, struct, props)

Sonoda, T. et al., *J. Antibiot.*, 1992, **45**, 1963
(abs config)


Ubukata, M. et al., *Nat. Prod. Lett.*, 1992, **1**,
149 (synth)

Dow, R.L. et al., *Tet. Lett.*, 1992, **33**, 309
(synth, config)

Ubukata, M. et al., *J. Antibiot.*, 1995, **48**, 1176
(synth, pmr)

Epilachnene E-93

5-Propyl-1-oxa-4-azacyclopentadec-10-en-15-one, 9CI. 11-Propyl-12-azacyclopentadec-5-en-14-olide
[147363-82-4]



Absolute configuration

C₁₆H₂₉NO₂ 267.411

Major alkaloid from the pupal defensive secretion of the Mexican bean beetle *Epilachna varivestis*. Syrup.

7,8-Didehydro(-): 5-Propyl-1-oxa-4-azacyclopentadeca-7,10-dien-15-one, 9CI. 11-Propyl-12-azacyclopentadeca-5,8-dien-14-olide. **Epilachnadiene**

[147363-83-5]
C₁₆H₂₇NO₂ 265.395

Minor alkaloid from *Epilachna varivestis*.

Lower homologue (side chain): 5-Ethyl-1-oxa-4-azacyclopentadec-10-en-15-one, 9CI. 11-Ethyl-12-azacyclopentadec-5-en-14-olide. **Novepilachnene**

[147363-86-8]
C₁₅H₂₇NO₂ 253.384

Trace alkaloid from *Epilachna varivestis*.

Higher homologue (n = 2): 5-Propyl-1-oxa-4-azacyclohexadec-11-en-16-one, 9CI. 12-Propyl-13-azacyclopentadec-5-en-15-olide. **Homoepilachnene**

[147363-84-6]
C₁₇H₃₁NO₂ 281.437

Trace alkaloid from *Epilachna varivestis*.

Attygalle, A.B. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 5204 (isol, ir, ms, pmr, struct)


Rao, B.V. et al., *Tet. Lett.*, 1996, **37**, 8613
(synth)

Fürstner, A. et al., *Synthesis*, 1997, 792-803
(synth, pmr, cmr, ir, ms)

Farmer, J.J. et al., *Tet. Lett.*, 1997, **38**, 2787
(abs config)

Attygalle, A.B. et al., *Tetrahedron*, 1999, **55**,
955-966 (biosynth, ms)

11-Epileontidane E-94




C₁₄H₂₄N₂ 220.357

Minor alkaloid from leaves and stems of *Maackia amurensis*, identified by ms (Fabaceae).

Kinghorn, A.D. et al., *Phytochemistry*, 1982,
21, 2269

Epilupinine E-95

Isolupinine. Tetralupine
[486-71-5]



(+)-form

C₁₀H₁₉NO 169.266

Diastereoisomeric with Octahydro-2H-quinolizine-1-methanol, O-58. Tetralupine was the racemate.

(+)-form

Alkaloid from leaves and seeds of *Lupinus cosentinii* (prev. designated *Lupinus pilosus*, *Lupinus varius* and *Lupinus digitatus*) (Fabaceae). Needles (petrol). Mp 77-78°. [α]_D +32 (EtOH). [α]_D +35 (H₂O).

Hydrochloride: Mp 172-173°. [α]_D +20.3 (H₂O).

Picrate: Mp 147-148°.

Methiodide: Mp 251-252°.

N-Oxide: **Epilupinine N-oxide**

[38225-02-4]

C₁₀H₁₉NO₂ 185.266

Alkaloid from *Lupinus cosentinii* leaves and seeds (Fabaceae). Hygroscopic needles (MeOH/Me₂CO). Mp 212-215°.

Ac: **Epilupinyl acetate**. Alkaloid LC1
C₁₂H₂₁NO₂ 211.303

Alkaloid from leaves of *Lupinus cosentinii* (Fabaceae). [α]_D²⁵ +33 (c, 1 in MeOH).

Ac, N-oxide: **Epilupinyl acetate N-oxide**. Alkaloid LC8

C₁₂H₂₁NO₃ 227.303

Cryst. + ½ H₂O (Me₂CO/cyclohexane). Mp 192°. [α]_D²⁵ +14 (c, 1 in MeOH).

O-(4-Hydroxy-E-cinnamoyl): **Epilupinyl trans-p-coumarate**. Alkaloid LC5

C₁₉H₂₅NO₃ 315.411

Alkaloid from leaves of *Lupinus cosentinii* (Fabaceae). Off-white cryst. (Me₂CO/cyclohexane). Mp 165°.

O-(4-Hydroxy-Z-cinnamoyl): **Epilupinine cis-p-coumarate**

C₁₉H₂₅NO₃ 315.411

Alkaloid from leaves of *Lupinus cosentinii* (Fabaceae). Off-white cryst. (Me₂CO/cyclohexane). Mp 144°.

O-[α-L-Rhamnopyranosyl-(1→4)-4-hydroxy-E-cinnamoyl]: **Epilupinyl trans-p-rhamnosylcoumarate**

[71657-66-4]

C₂₅H₃₅NO₇ 461.554

Alkaloid from *Lupinus cosentinii* and *Lupinus hirsutus* (Fabaceae). Oil. [α]_D -76 (c, 0.62 in EtOH).

O-[α-L-Rhamnopyranosyl-(1→4)-4-hydroxy-Z-cinnamoyl]: **Epilupinyl cis-p-rhamnosylcoumarate**

[71657-67-5]

C₂₅H₃₅NO₇ 461.554

Alkaloid from *Lupinus cosentinii* and *Lupinus hirsutus* (Fabaceae). Oil. [α]_D -52.8 (c, 0.99 in EtOH).

O-(4-Acetoxy-E-cinnamoyl): **Epilupinyl trans-p-acetoxycinnamate**

[136396-57-1]

C₂₁H₂₇NO₄ 357.449

Alkaloid from *Lupinus hirsutus* (Fabaceae). Oil. [α]_D²³ +20.5 (c, 0.12 in EtOH).

O-(4-Hydroxy-3-methoxy-E-cinnamoyl): **Epilupinyl trans-ferulate**. Alkaloid LC4

C₂₀H₂₇NO₄ 345.438

Alkaloid from leaves of *Lupinus cosentinii* (Fabaceae). Pale yellow cryst. (Me₂CO/cyclohexane). Mp 119°. [α]_D +35 (c, 1 in MeOH). Also present as rhamnoside (component of Alkaloid LC9).

O-[α-L-Rhamnopyranosyl-(1→4)-4-hydroxy-3-methoxy-E-cinnamoyl]: **Epilupinyl rhamnosylferulate**

C₂₆H₃₇NO₈ 491.58

Alkaloid from aerial parts of *Lupinus hirsutus* and from *Lupinus varius* (Fabaceae). Amorph. solid. [α]_D -37.5 (c, 0.056 in EtOH). [α]_D -80 (c, 0.28 in EtOH). Isol. once as a mixt. of E- and Z-isomers (ratio = 1:5), later from *L. varius* as pure E-form. λ_{max} 230 ; 295 (MeOH).

(-)-form

Synthetic. $[\alpha]_D^{22}$ -24.09 (c, 0.22 in EtOH).

(±)-form [486-72-6]

Alkaloid from *Lupinus palmeri* (Fabaceae). Mp 83°.

Picrate:

Yellow needles. Mp 145.5-147°.

Methiodide: Mp 251-253° (248°).

White, E.P. et al., *N.Z. J. Sci. Technol., Sect. B*, 1951, **33**, 50 (isol)

Crow, W.D. et al., *Aust. J. Chem.*, 1955, **8**, 136; 1957, **10**, 77 (isol)

Thomas, A.F. et al., *Can. J. Chem.*, 1955, **33**, 1290

Van Tamelen, E.E. et al., *J.A.C.S.*, 1960, **82**, 502 (synth)

Paszyc, S. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1970, **18**, 15 (synth)

Batra, V. et al., *Indian J. Chem., Sect. B*, 1976, **14**, 636 (isol)

Beck, A.B. et al., *J. Nat. Prod.*, 1979, **42**, 385 (isol, esters, bibl, rhamnosylcoumarate)

Okita, M. et al., *Heterocycles*, 1983, **20**, 401 (synth)

Bremmer, M.L. et al., *J.O.C.*, 1983, **48**, 3661 (synth)

Chamberlin, A.R. et al., *J.O.C.*, 1984, **49**, 1682 (synth)

Ihara, M. et al., *Heterocycles*, 1985, **23**, 1097 (synth)

Hiemstra, H. et al., *J.O.C.*, 1985, **50**, 4014 (synth)

Takahata, H. et al., *Chem. Pharm. Bull.*, 1986, **34**, 4523 (synth)

Grieco, P.A. et al., *J.O.C.*, 1988, **53**, 3325 (synth, ir, pmr)

Nagasaka, T. et al., *Heterocycles*, 1989, **29**, 1209 (synth)

Célérier, J.P. et al., *Tetrahedron*, 1989, **45**, 6161 (synth)

Takamatsu, S. et al., *Phytochemistry*, 1990, **29**, 3923 (rhamnosylcoumarates)

Takamatsu, S. et al., *J. Nat. Prod.*, 1991, **54**, 477 (trans-p-acetylcoumarate, acetoxyacoumarate)

Hua, D.H. et al., *Synthesis*, 1991, 970 (synth)

Edstrom, E.D. et al., *Tet. Lett.*, 1991, **32**, 5709 (synth)

Gesson, J.P. et al., *Tet. Lett.*, 1992, **33**, 3633 (synth)

Pandey, G. et al., *Tet. Lett.*, 1992, **33**, 6533 (synth)

West, F.G. et al., *J.A.C.S.*, 1994, **116**, 8420 (synth)

Suzuki, H. et al., *Phytochemistry*, 1994, **37**, 591 (rhamnosylferulate)

Molander, G.A. et al., *J.O.C.*, 1996, **61**, 6040 (synth)


Mangency, P. et al., *Tetrahedron*, 1998, **54**, 10349-10362 (synth)

Abdel-Halim, O.B. et al., *Phytochemistry*, 1999, **51**, 5-9; **52**, 965 (rhamnosylferulate)

Amorde, S.M. et al., *Org. Lett.*, 2005, **7**, 2031-2033 (synth)

Epinardine A**E-96**

[178822-54-3]




$C_{18}H_{18}N_3O_3^{\oplus}$ 324.358

Alkaloid from an unidentified deep-water green demosponge collected in the South Indian Ocean. Green powder (counterion unspecified). λ_{max} 245 (€ 9800); 367 (€ 6800); 570 (€ 740) (MeOH).

D'Ambrosio, M. et al., *Tetrahedron*, 1996, **52**, 8899 (isol, uv, cd, pmr, cmr, struct)

Epinardine B**E-97**

[178822-55-4]



$C_{18}H_{17}Br_2N_3O_2$ 467.159

Alkaloid from an unidentified deep-water green demosponge collected in the South Indian Ocean. Grey-green powder. λ_{max} 200 (€ 11600); 248 (€ 9300); 340 (€ 4050); 390 (€ 3130) (MeOH). λ_{max} 248 (€ 9300); 340 (€ 4050); 390 (€ 3130) (MeOH) (Berdy).

7,8-Didehydro: Epinardine C

[178822-56-5]

$C_{18}H_{15}Br_2N_3O_2$ 465.143

Isol. from an unidentified deep-water demosponge. Strongly cytotoxic towards L1210 and doxorubicin-resistant L1210/DX murine lymphocytic leukaemia cells *in vitro*. Green powder. λ_{max} 210 (€ 11400); 250 (€ 5900); 290 (€ 4400); 370 (€ 3200); 535 (€ 374) (MeOH).

8 α -Methoxy: Epinardine D

[178822-57-6]


$C_{19}H_{19}Br_2N_3O_3$ 497.185

Isol. from an unidentified deep-water green demosponge. Green powder. Converted to Epinardine C on long standing in Me₂CO soln. λ_{max} 205 (€ 35000); 242 (€ 31000); 335 (€ 24000); 480 (€ 2000) (MeOH).

D'Ambrosio, M. et al., *Tetrahedron*, 1996, **52**, 8899 (isol, uv, cd, pmr, cmr, struct)

Epinine**E-98**

4-[2-(Methylamino)ethyl]-1,2-benzenediol, 9CI. 4-[2-(Methylamino)ethyl]pyrocatechol, 8CI. 4-(β -Methylaminoethyl)-catechol. N-Methyl-2-(3,4-dihydroxyphenyl)ethylamine. Deoxyadrenaline. Desoxyepinephrine. N-Methyldopamine [501-15-5]



$C_9H_{13}NO_2$ 167.207

Alkaloid from *Cytisus scoparius*, *Vicia faba* and *Lophophora williamsii* (Faba-

ceae, Cactaceae). Active metab. of Ibo-pamine. Dopamine D₁- and D₂- agonist and α - and β -adrenoceptor agonist. Positive inotropic agent and vasodilator. Cryst. (EtOH). Mp 188-189°. Log P 0.16 (calc).

Hydrochloride: [62-32-8]

Prisms (H₂O). Mp 179-180°.

▶ LD₅₀ (mus, ipr) 212 mg/kg. UX1925000

O⁴-Phosphate: N-Methyldopamine 4-O-dihydrogen phosphate. Z 2055

$C_9H_{14}NO_5P$ 247.187

Dopaminergic prodrug showing renal selectivity. Mp 175-178° (as hydrochloride).

Pyman, F.L. et al., *J.C.S.*, 1909, 95; 1266; 1610 (synth)

Buck, J.S. et al., *J.A.C.S.*, 1930, **52**, 4119 (synth)

Bretschneider, H. et al., *Monatsh. Chem.*, 1947, **76**, 335 (synth)

Dewhurst, W.G. et al., *Br. J. Pharmacol.*, 1965, **25**, 682 (pharmacol)

Path, P.N. et al., *J. Pharmacol. Exp. Ther.*, 1967, **155**, 1 (pharmacol)

Tocher, R.D. et al., *Phytochemistry*, 1972, **11**, 1161 (isol)

Borgman, R.J. et al., *J. Med. Chem.*, 1973, **16**, 630 (synth)

Giesecke, J. et al., *Acta Cryst. B*, 1976, **32**, 2337 (cryst struct)

Smith, T.A. et al., *Phytochemistry*, 1977, **16**, 9 (rev, occur)

Van Woerkens, L.J. et al., *Br. J. Pharmacol.*, 1992, **107**, 303 (pharmacol)

Boomsma, F. et al., *J. Chromatogr.*, 1992, **574**, 109 (hplc)

Schwinger, R.H.G. et al., *J. Pharmacol. Exp. Ther.*, 1993, **265**, 346 (pharmacol)

Ianelli, S. et al., *Acta Cryst. C*, 1995, **51**, 1338 (cryst struct, phosphate)

Daul, A. et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1995, **352**, 429 (pharmacol)


Martindale, *The Extra Pharmacopoeia*, 31st edn., Pharmaceutical Press, 1996, 1578

Schwinger, R.H.G. et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1996, **354**, 343 (pharmacol)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EAZ000

Epipodopetaline**E-99**

Amazonine. Alkaloid R6. Sweetinine



Relative configuration

$C_{20}H_{33}N_3$ 315.501

Identity of Amazonine with Epipodopetaline is tentative. Sweetinine was the racemate.

(-)-form [62624-21-9]

Alkaloid from *Podopetalum ormondii* (preferred genus name *Ormosia*) (Fabaceae). Mp 118-119°. $[\alpha]_D$ -3.2 (MeOH). Largely racemic. One isolate from *P. ormondii* was wholly racemic (see below).

(±)-form [1361-42-8]

Alkaloid from stem bark of *Sweetia elegans*, bark of *Sweetia panamensis* and *Podopetalum ormondii* (Fabaceae). Mp 174-175°.

Fitzgerald, T.J. *et al.*, *J. Nat. Prod.*, 1964, **27**, 107-110 (*isol*)

Cheng, P.-T. *et al.*, *Tet. Lett.*, 1976, **17**, 4245-4246 (*isol, struct*)


Mackay, M.F. *et al.*, *Cryst. Struct. Commun.*, 1980, **9**, 805-809 (*cryst struct*)

McLean, S. *et al.*, *Can. J. Chem.*, 1981, **59**, 34-37 (*isol, struct*)

Balandrin, M.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 619-622 (*isol, uv, ir, pmr, ms, Sweetinine*)

Epipolasin A**E-100**

4-Isothiocyanatomaaliane

**(+)-form**

X = -NCS

C₁₆H₂₅NS 263.446**(+)-form** [97950-05-5]

Metab. of the sponge *Epipolasis kushimotoensis*. Cryst. [α]_D²⁵ +7.6 (c, 1.0 in CHCl₃).

(-)-form

Metab. of the sponges *Acanthella pulcherrima* and *Axinyssa* sp. and the mollusc *Cadlina luteomarginata*. Oil. [α]_D²⁵ -8 (c, 1.3 in CHCl₃).

Formamide: 4-Formylaminomaaliane

[83631-20-3]

C₁₆H₂₇NO 249.395

Metab. of an *Acanthella* sp. and *Cadlina luteomarginata*. Has -NHCHO replacing -NCS.

[83631-19-0]

Thompson, J.E. *et al.*, *Tetrahedron*, 1982, **38**, 1865-1873 (*isol, pmr, ms*)


Tada, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 1941

Capon, R.J. *et al.*, *Aust. J. Chem.*, 1988, **41**, 979 (*isol*)

Simpson, J.S. *et al.*, *Aust. J. Chem.*, 1997, **50**, 1123-1127 (*isol, pmr, cmr, ms*)

Epipyridone**E-101**

[944339-73-5]




Relative Configuration

C₂₃H₂₉NO₃ 367.487

Related to Leporine A, L-112 and Fusaricide, F-225. Prod. by an *Epicoccum* sp. associated with *Pholiota squarrosa*. Red oil. [α]_D²⁵ +123.3 (c, 0.06 in MeOH). λ_{max} 213 ; 250 (MeOH).

Wangun, H.V.K. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1702-1705 (*isol, pmr, cmr, ms*)

Epistephanine*Aknadine***E-102****(R)-form**C₃₇H₃₈N₂O₆ 606.717

Not the epimer of 1,2,8-Trihydroxyaporphine, T-522.

(R)-form [549-08-6]

Alkaloid from *Stephania capitata*, the roots and terrestrial portion of *Stephania japonica* and from *Stephania hernandifolia* (Menispermaceae). Needles (C₆H₆/petrol). Mp 203-204° (133-134°). [α]_D²⁰ +183.5 (CHCl₃). [α]_D²⁵ +226 (CHCl₃). λ_{max} 233 (ε 34100); 282 (ε 14600) (EtOH).

O⁷-De-Me: 3',4'-Dihydrostephasubine

[115610-45-2]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the stems of *Stephania hernandifolia*. Amorph. [α]_D²⁵ +286 (MeOH). λ_{max} 216 ; 223 ; 283 (EtOH).

O¹²-De-Me: Hypoepistephanine. Pseudoepistephanine

[33116-41-5]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from roots and terrestrial parts of *Stephania japonica* (Menispermaceae). Mp 256-257°. [α]_D^{15.5} +183.8 (CHCl₃).

O¹²-De-Me, N-de-Me: GuattebolineC₃₅H₃₄N₂O₆ 578.663

Alkaloid from the stem bark of *Guatteria boliviana*. Amorph. solid. [α]_D²⁰ +138 (c, 0.8 in CHCl₃). λ_{max} 205 (log ε 4.85); 231 (sh) (log ε 4.56); 283 (log ε 4.21) (EtOH).

O⁷,O¹²-Di-de-Me, N²-de-Me: Pangkorimine

[110360-01-5]

C₃₄H₃₂N₂O₆ 564.637

Alkaloid from *Albertisia* cf. *Albertisia papuana* (Menispermaceae). Exhibits antiparasitic and cytotoxic activities.

[α]_D²⁰ +65 (c, 0.05 in MeOH). λ_{max} 213 (log ε 4.64); 228 (log ε 4.48); 283 (log ε 4.26); 314 (log ε 3.8) (EtOH).

(S)-form [40039-47-2]

Alkaloid from the stems of *Anisocycla grandidieri* (Menispermaceae). Cryst. (MeOH). Mp 198-206°. [α]_D²⁰ -216 (c, 1.94 in CHCl₃).

(ξ)-form**α'-Oxo: Oxoepistephanine**C₃₇H₃₆N₂O₇ 620.701

Alkaloid from *Stephania hernandifolia* (Menispermaceae). Cryst. (MeOH/Et₂O). Mp 222-226° dec. [α]_D²⁷ +272 (CHCl₃). No CAS Reg. no. to 14CI (2001).

Tanaka, K. *et al.*, *Yakugaku Zasshi*, 1944, **64**, 27-35; *CA*, **45**, 5173d (*struct*)

Tomita, M. *et al.*, *Pharm. Bull.*, 1954, **2**, 378-381 (*struct*)

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 996-999; *CA*, **60**, 4201h (*isol*)

Barton, D.H.R. *et al.*, *J.C.S.(C)*, 1966, 2313-2319 (*uv, pmr, ms, biosynth*)

Moza, B.K. *et al.*, *Indian J. Chem.*, 1967, **5**, 281-282 (*Aknadine*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 2061-2064 (*(S)-form, isol, struct*)

Cava, M.P. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 279-280 (*Oxoepistephanine*)

Ray, A.B. *et al.*, *Planta Med.*, 1979, **35**, 167-173 (*Epistephanine*)

Lavault, M. *et al.*, *Can. J. Chem.*, 1987, **65**, 343-347 (*Pangkorimine*)


Patra, A. *et al.*, *Phytochemistry*, 1988, **27**, 653-655 (*3',4'-Dihydrostephasubine*)

Mahiou, V. *et al.*, *Phytochemistry*, 2000, **54**, 709-716 (*Guatteboline, Pangkorimine, activity*)

Chen, C.-K. *et al.*, *Chin. Pharm. J. (Taipei)*, 2003, **55**, 35-47 (*Pangkorimine*)

Epocarbazolin A**E-103**

[146935-39-9]

R = CH₃C₂₂H₂₇NO₄ 369.46

Prod. by *Streptomyces anulatus* T688-8. 5-Lipoxygenase inhibitor. Active against gram-positive bacteria. Pale yellow powder. Sol. MeOH, EtOAc, Me₂CO; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 100°. [α]_D²⁵ +75 (c, 0.5 in MeOH).

Unstable to light. λ_{max} 234 (ε 38000); 258 (ε 32500); 290 (sh) (ε 9000); 304 (ε 15200); 358 (ε 6900); 364 (sh) (ε 6800) (MeOH/NaOH) (Derep). λ_{max} 234 (ε 38600); 250 (sh) (ε 33900); 289 (sh) (ε 10200); 299 (ε 14800); 345 (ε 7100); 359 (ε 7500) (MeOH) (Derep).

Nihei, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 25-33 (*isol, pmr, cmr, struct, props*)

Knöll, J. *et al.*, *Tet. Lett.*, 2006, **47**, 6079-6082 (*synth*)

Epocarbazolin B**E-104**

[146935-40-2]

As Epocarbazolin A, E-103 with

R = -CH₂CH₃C₂₃H₂₉NO₄ 383.486

Prod. by *Streptomyces anulatus* T688-8. 5-Lipoxygenase inhibitor. Active against gram-positive bacteria. Pale yellow powder. Sol. MeOH, EtOAc, Me₂CO; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 100°. [α]_D²⁵ +78 (c, 0.5 in MeOH).


Unstable to light. λ_{max} 234 (ε 38000); 258 (ε 32500); 290 (sh) (ε 9000); 304 (ε 15200); 358 (ε 6900); 364 (sh) (ε 6800) (MeOH/NaOH) (Derep). λ_{max} 234 (ε 38600); 250 (sh) (ε 33900); 289 (sh) (ε 10200); 299 (ε 14800); 345 (ε 7100); 359 (ε 7500) (MeOH) (Derep).

Nihei, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 25-33 (*isol, pmr, cmr, uv, struct, activity*)

Knöll, J. *et al.*, *Tet. Lett.*, 2006, **47**, 6079-6082 (*synth*)

Epothelmin A

E-105



Absolute Configuration

C₁₈H₃₁NO₂ 293.448

Struct. revised in 2005. Prev. assigned as an epoxide. Prod. by fungal strain FKI-0929. Lanosterol synthase inhibitor. Oil. $[\alpha]_D^{25} +22$ (c, 0.14 in MeOH). λ_{\max} 224 (€ 9300) (MeOH).

1-Epimer: Epothelmin B

C₁₈H₃₁NO₂ 293.448

Prod. by fungal strain FKI-0929.

Lanosterol synthase inhibitor. Oil.

$[\alpha]_D^{25} +25$ (c, 0.28 in MeOH). λ_{\max} 224 (€ 11800) (MeOH).

Sakano, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 564-568 (*isol*, *pmr*, *cmr*)

Shibuya, M. *et al.*, *J. Antibiot.*, 2005, **58**, 599-601 (*pmr*, *cmr*, *struct*)


Snider, B.B. *et al.*, *Org. Lett.*, 2005, **7**, 4419-4422 (*synth*, *struct*)

Fürstner, A. *et al.*, *Chem. Asian J.*, 2008, **3**, 310-318 (*Epothelmin B*, *synth*)

Epolactaene

E-106

[167782-17-4]

C₂₁H₂₇NO₆ 389.447

Nat. product is a diastereoisomeric mixture at C-15. Prod. by the marine fungus *Penicillium* sp. BM1689-P. Neuritogenic agent. Amorph. solid. Sol. MeOH, EtOH, DMSO; fairly sol. CHCl₃, EtOAc; poorly sol. H₂O, hexane. $[\alpha]_D^{26} +32$ (c, 0.1 in MeOH). Similar to Fusarin C, F-227. λ_{\max} 232 (€ 21800); 280 (€ 15600) (MeOH) (Berdy).

Takeya, H. *et al.*, *J. Antibiot.*, 1995, **48**, 733-735 (*isol*, *ir*, *uv*, *pmr*, *cmr*)

Kuramochi, K. *et al.*, *Tet. Lett.*, 1999, **40**, 7371-7374 (*synth*)


Marumoto, S. *et al.*, *Tetrahedron*, 1999, **55**, 7129-7144; 7145-7156 (*synth*, *abs config*)

Hayashi, Y. *et al.*, *J.O.C.*, 2002, **67**, 9443-9448 (*synth*)

Epostatin

E-107

[181372-99-6]

C₂₃H₃₃N₃O₅ 431.531

Prod. by *Streptomyces* sp. MJ995-OF5. Inhibitor of dipeptidyl peptidase II. Pale yellow powder. Mp 157-159° dec. $[\alpha]_D^{23} -174$ (c, 0.1 in MeOH). λ_{\max} 218 (sh) (log € 3.54); 305 (log € 4.43) (MeOH).

Akiyama, T. *et al.*, *J. Antibiot.*, 1998, **51**, 253-260; 372-373 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Epothilone A₁

E-108

[252917-29-6]

As Epothilone A, E-110 with

R¹ = R² = H, R³ = R⁴ = CH₃C₂₅H₃₇NO₆S 479.636

C-4 config. not determined. Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22} -69$ (c, 0.1 in MeOH). λ_{\max} 208 (€ 19600); 247 (€ 13600) (MeOH).

4-Epimer: Epothilone A₂

[252917-30-9]

C₂₅H₃₇NO₆S 479.636Prod. by *Sorangium cellulosum*.

Amorph. solid. $[\alpha]_D^{22} -12$ (c, 1 in MeOH). λ_{\max} 210 (€ 15100); 248 (€ 15500) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone A₃

E-109

As Epothilone A, E-110 with

R¹ = R⁴ = H, R² = R³ = CH₃C₂₅H₃₇NO₆S 479.636


Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{25} -76.2$ (c, 1 in MeOH). λ_{\max} 210 (€ 15300); 248 (€ 15500) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone A

E-110

[152044-53-6]

R¹ = H, R² = R³ = R⁴ = CH₃C₂₆H₃₉NO₆S 493.663

Prod. by *Sorangium cellulosum*. Induces cellular microtubuli assembly and stabilisation by a Taxol™-like mechanism.

Competes for same binding site on microtubules as Taxol™. Antiproliferative and potential anticancer agent.

Antifungal and cytotoxic agent. Cryst. (EtOAc/toluene). Mp 95°. $[\alpha]_D^{21} -47.1$ (c, 1 in MeOH). λ_{\max} 211 (€ 17800); 249 (€ 12500) (MeOH).

21-Hydroxy: Epothilone E

[201049-37-8]

C₂₆H₃₉NO₇S 509.663

Prod. by *Sorangium cellulosum*. Antitumour agent. Amorph. solid. $[\alpha]_D^{21} -37.1$ (c, 0.56 in MeOH). λ_{\max} 212 (€ 17800); 247 (€ 13000) (MeOH).

27-Hydroxy: Epothilone A₉

[252917-32-1]

C₂₆H₃₉NO₇S 509.663

Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22} -38$ (c, 0.5 in MeOH). λ_{\max} 211 (€ 15500); 253 (€ 14100) (MeOH).

Bollag, D.M. *et al.*, *Cancer Res.*, 1995, **55**, 2325-2333 (*pharmacol*)

Hoefle, G. *et al.*, *Angew. Chem., Int. Ed.*, 1996, **35**, 1567-1569 (*uv*, *cmr*, *pmr*, *cryst struct*)

Balog, A. *et al.*, *Angew. Chem., Int. Ed.*, 1996, **35**, 2801-2803 (*synth*)

Victory, S.F. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, 893-898 (*conformn*)

Gerth, K. *et al.*, *J. Antibiot.*, 1996, **49**, 560-563 (*isol*, *ir*, *pmr*)

Yang, Z. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 166-168; 525-527 (*synth*)

Schinzer, D. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 523-524 (*synth*)

Wessjohann, L. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 715-718 (*rev*)

Su, D.-S. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 2093-2096 (*activity*)

Finlay, R. *et al.*, *Chem. Ind. (London)*, 1997, 991-996 (*rev*)

Kowalski, R.J. *et al.*, *J. Biol. Chem.*, 1997, **272**, 2534-2541 (*pharmacol*)

Meng, D. *et al.*, *J.A.C.S.*, 1997, **119**, 2733-2734 (*synth*)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 1997, **119**, 7960-7973; 7974-7991 (*synth*)

Meng, D. *et al.*, *J.A.C.S.*, 1997, **119**, 10073-10092 (*synth*)

Nicolaou, K.C. *et al.*, *Nature (London)*, 1997, **387**, 268-272 (*synth*, *bibl*)

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1998, **37**, 2015-2045 (*rev*)

Nicolaou, K.C. *et al.*, *Bioorg. Med. Chem.*, 1999, **7**, 665-697 (*Epothilone E*, *synth*)

Schinzer, D. *et al.*, *Chem. Eur. J.*, 1999, **5**, 2483-2491; 2492-2500 (*synth*)

Kalesse, M. *et al.*, *Eur. J. Org. Chem.*, 1999, 2817-2823 (*synth*)

Taylor, R.E. *et al.*, *J.O.C.*, 1999, **64**, 7224-7228 (*conformn*)

Gerth, K. *et al.*, *J. Antibiot.*, 2000, **53**, 1373-1377 (*biosynth*)

Zhu, B. *et al.*, *Eur. J. Org. Chem.*, 2001, 1701-1714 (*synth*)

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856 (*Epothilone A₆*)

Bode, J.W. *et al.*, *J.A.C.S.*, 2001, **123**, 6410-6424 (*synth*)

Hindupur, R.M. *et al.*, *Tet. Lett.*, 2001, **42**, 7341-7344 (*synth*)

Liu, Z.-Y. *et al.*, *Chem. Eur. J.*, 2002, **8**, 3747-3756 (*synth*)

Altmann, K.-H. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 4086-4110 (*12,13-trans isomer*, *synth*, *activity*)

Epothilone B

E-111

Patupilone, INN. EPO 906

[152044-54-7]

As Epothilone A, E-110 with

R¹ = R² = R³ = R⁴ = CH₃C₂₇H₄₁NO₆S 507.69

Prod. by *Sorangium cellulosum*. Antifungal and cytotoxic agent. Induces cellular microtubuli assembly and stabilisation by a Taxol™-like mechanism. Competes for the same binding site on microtubules as Taxol™. Antiproliferative and potential anticancer agent. Cryst. (EtOAc). Mp 93-94°. $[\alpha]_D^{21} -35$ (c, 0.7 in MeOH). More potent in some cytotoxicity and antifungal assays than Epothilone A, E-110. λ_{\max} 211 (€ 18600); 249 (€ 14100) (MeOH).

21-Hydroxy: Epothilone F

[208518-52-9]

C₂₇H₄₁NO₇S 523.689Prod. by *Sorangium cellulosum*.

Amorph. solid. $[\alpha]_D^{22} -27.4$ (c, 0.46 in MeOH). λ_{\max} 212 (€ 19100); 249 (€

13800) (MeOH).

21-Methyl: Epothilone B₁₀
[252917-33-2]
C₂₈H₄₃NO₆S 521.717
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -27 (c, 0.15 in MeOH). λ_{\max} 212 (ε 15800); 247 (ε 12500) (MeOH).

21-Amino: BMS 310705
[280578-49-6]
C₂₇H₄₂N₂O₆S 522.705
Antineoplastic agent. Solid; characterised by pmr and ms.

Bollag, D.M. *et al.*, *Cancer Res.*, 1995, **55**, 2325-2333 (pharmacol)

Hoeftel, G. *et al.*, *Angew. Chem., Int. Ed.*, 1996, **35**, 1567-1569 (uv, pmr, cmr, cryst struct)

Victory, S.F. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, 893-898 (conformn)

Gerth, K. *et al.*, *J. Antibiot.*, 1996, **49**, 560-563 (isol)

Wessjohann, L. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 715-718 (rev)

Su, D.-S. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 757-759 (synth)

Muehlradt, P.F. *et al.*, *Cancer Res.*, 1997, **57**, 3344-3346 (pharmacol)

Finlay, R. *et al.*, *Chem. Ind. (London)*, 1997, 991-996 (rev)

Kowalski, R.J. *et al.*, *J. Biol. Chem.*, 1997, **272**, 2534-2541 (pharmacol)

Nicolaou, K. C. *et al.*, *J.A.C.S.*, 1997, **119**, 7960-7973; 7974-7991 (synth)

Meng, D. *et al.*, *J.A.C.S.*, 1997, **119**, 10073-10092 (synth)

Nicolaou, K.C. *et al.*, *Nature (London)*, 1997, **387**, 268-272 (synth, bibl)

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1998, **37**, 2015-2045 (rev)

May, S.A. *et al.*, *Chem. Comm.*, 1998, 1597-1598 (synth)

Mulzer, J. *et al.*, *Tet. Lett.*, 1998, **39**, 8633-8636 (synth)

Nicolaou, K.C. *et al.*, *Chem. Comm.*, 1999, 519-520 (synth)

Schinzer, D. *et al.*, *Chem. Eur. J.*, 1999, **5**, 2483-2491; 2492-2500 (synth)

White, J.D. *et al.*, *J.O.C.*, 1999, **64**, 684-685 (synth, bibl)

Taylor, R.E. *et al.*, *J.O.C.*, 1999, **64**, 7224-7228 (conformn)

Nicolaou, K.C. *et al.*, *Chem. Eur. J.*, 2000, **6**, 2783-2800 (Epothilone B₁₀, synth)

Gerth, K. *et al.*, *J. Antibiot.*, 2000, **53**, 1373-1377 (biosynth)

Mulzer, J. *et al.*, *J.O.C.*, 2000, **65**, 7456-7467 (synth)

Mulzer, J. *et al.*, *Monatsh. Chem.*, 2000, **131**, 205-238 (rev, synth)

Pat. Coop. Treaty (WIPO), 2000, ((Bristol-Myers Squibb))00 50 423; CA, **133**, 193030f (BMS 310705, synth, pharmacol)

Klaus, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856 (Epothilone B₁₀)

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856 (Epothilone F, isol, uv, ir, pmr, cmr, ms)

White, J.D. *et al.*, *J.A.C.S.*, 2001, **123**, 5407-5413 (synth)

Bode, J.W. *et al.*, *J.O.C.*, 2001, **66**, 6410-6424 (synth)

Broker, L.E. *et al.*, *Cancer Res.*, 2002, **62**, 4081-4088 (pharmacol)

Altaha, R. *et al.*, *Curr. Pharm. Des.*, 2002, **8**, 1707-1712 (rev)

Uyar, D. *et al.*, *Gynecol. Oncol.*, 2003, **91**, 173-178 (BMS 310705, rev)

Kolman, A. *et al.*, *Curr. Opin. Invest. Drugs*, 2004, **5**, 1292-1297 (BMS 310705, rev)

Jung, J.-C. *et al.*, *J.O.C.*, 2004, **69**, 9269-9284 (synth)

Rubin, E.H. *et al.*, *J. Clin. Oncol.*, 2005, **23**, 9120-9129 (clin trial)

Larkin, J.M.G. *et al.*, *Drugs of the Future*, 2007, **32**, 323 (rev)

Keck, G.E. *et al.*, *J.O.C.*, 2008, **73**, 9675-9691 (synth)

Epothilone C₁

E-112

As Epothilone C, E-115 with
R¹ = R² = H, R³ = R⁴ = R⁵ = CH₃
C₂₅H₃₇NO₅S 463.637
C-4 config. not determined. Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -114 (c, 10 in MeOH). λ_{\max} 211 (ε 16500); 248 (ε 12500) (MeOH).

12E-Isomer: trans-Epothilone C₁
C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -84 (c, 0.2 in MeOH). λ_{\max} 211 (ε 17400); 248 (ε 12900) (MeOH).

4-Epimer: Epothilone C₂
C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*.
Amorph. powder. $[\alpha]_D^{22}$ -11.6 (c, 10 in MeOH). λ_{\max} 212 (ε 15500); 249 (ε 12100) (MeOH).

4-Epimer, 12E-isomer: trans-Epothilone C₂
C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*.
Amorph. powder. $[\alpha]_D^{22}$ -3 (c, 1.5 in MeOH). λ_{\max} 211 (ε 15800); 248 (ε 11900) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone C₃

E-113

As Epothilone C, E-115 with
R¹ = R⁴ = H, R² = R³ = R⁵ = CH₃
C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -62.1 (c, 5 in MeOH). λ_{\max} 212 (ε 16200); 248 (ε 12300) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone C₄

E-114


As Epothilone C, E-115 with
R¹ = R⁵ = H, R² = R³ = R⁴ = CH₃
C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -75.6 (c, 1 in MeOH). λ_{\max} 212 (ε 17200); 248 (ε 12500) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone C

E-115

Deoxyepothilone A
[186692-73-9]



R¹ = H, R² = R³ = R⁴ = R⁵ = CH₃

C₂₆H₃₉NO₅S 477.664
Prod. by *Sorangium cellulosum*. Antitumour agent. Amorph. solid. $[\alpha]_D^{22}$ -74.4 (c, 2 in MeOH). λ_{\max} 213 (ε 16200); 248 (ε 12500) (MeOH).

8,9-Didehydro(E-): Epothilone C₅

C₂₆H₃₇NO₅S 475.648
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -158 (c, 0.5 in MeOH). λ_{\max} 205 (ε 19500); 247 (ε 12700) (MeOH).

10,11-Didehydro(E-): Epothilone C₆

C₂₆H₃₇NO₅S 475.648
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -205.2 (c, 1 in MeOH). λ_{\max} 218 (ε 24600); 237 (ε 28800) (MeOH).

14α-Hydroxy: Epothilone C₇

C₂₆H₃₉NO₆S 493.663
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -11 (c, 1 in MeOH). λ_{\max} 203 (ε 18300); 247 (ε 12500) (MeOH).

27-Hydroxy: Epothilone C₉

C₂₆H₃₉NO₆S 493.663
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -93.4 (c, 1 in MeOH). λ_{\max} 209 (ε 15200); 254 (ε 15700) (MeOH).

4-Demethyl, 9-oxo: 4-Demethyl-9-ketoe-pothilone C

[612057-52-0]
C₂₅H₃₅NO₆S 477.621
Isol. from recombinant *Myxococcus xanthus*. Yellow oil. $[\alpha]_D$ -9.2 (c, 0.05 in MeOH). λ_{\max} 212 (ε 25140); 248 (ε 20600) (MeOH).

16-Demethyl: Epothilone C₈. 27-Norepothilone C

C₂₅H₃₇NO₅S 463.637
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -75.2 (c, 2.5 in MeOH). λ_{\max} 210 (ε 16800); 248 (ε 17800) (MeOH).

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 2097-2103; 1998, **37**, 84-87 (synth, ir, pmr, cmr)

Hardt, I. *et al.*, *J. Nat. Prod.*, 2000, **64**, 847-856 (isol, uv, ir, pmr, cmr, ms)

Storer, R.I. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2521-2525 (synth)

Starks, C.M. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1313-1317 (4-Demethyl-9-ketoe-pothilone C)

Epothilone D₁

E-116

As Epothilone C, E-115 with
R¹ = R³ = R⁴ = R⁵ = CH₃, R² = H
C₂₆H₃₉NO₅S 477.664

C-4 config. not determined. Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -119 (c, 0.5 in MeOH). λ_{\max} 208 (ε 18300); 249 (ε 11900) (MeOH).

4-Epimer: Epothilone D₂

C₂₆H₃₉NO₅S 477.664
Prod. by *Sorangium cellulosum*.
Amorph. solid. $[\alpha]_D^{22}$ -12.5 (c, 1 in MeOH). λ_{\max} 210 (ε 15400); 248 (ε 11200) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone D

E-117

Deoxyepothilone B. NSC 703147. KOS 862

[189453-10-9]

As Epothilone C, E-115 with

 $R^1 = R^2 = R^3 = R^4 = R^5 = CH_3$ C₂₇H₄₁NO₅S 491.691Prod. by *Mycococcus xanthus* and *Sorangium cellulosum*. Antitumour agent.Cryst. (EtOH aq.). Mp 120-121°. $[\alpha]_D^{22}$ -61.3 (c, 2.5 in MeOH). λ_{max} 210 (ε 18400); 248 (ε 13200) (MeOH).**8,9-Didehydro(E-): Epothilone D₅**C₂₇H₃₉NO₅S 489.675Prod. by *Sorangium cellulosum*.Amorph. solid. $[\alpha]_D^{22}$ -150 (c, 0.2 in MeOH). λ_{max} 205 (ε 23300); 248 (ε 13600) (MeOH).**10,11-Didehydro(E-): 10,11-Didehydroepothilone D**C₂₇H₃₉NO₅S 489.675Prod. by genetically modified *Mycococcus xanthus*. Cytotoxic. Oil. $[\alpha]_D$ -74 (c, 0.08 in MeOH).**9-Oxo: 9-Oxoepothilone D**

[612057-56-4]

C₂₇H₃₉NO₆S 505.674Isol. from recombinant *Mycococcus xanthus*.**9-Oxo, 10,11-didehydro(E-): 10,11-Didehydro-9-oxoepothilone D**

[612057-51-9]

C₂₇H₃₇NO₆S 503.658Isol. from recombinant *Mycococcus xanthus*. Yellow oil. $[\alpha]_D$ -84.2 (c, 0.005 in MeOH). λ_{max} 210 (ε 17500); 247 (ε 14900); 286 (ε 10600) (MeOH).**4-Demethyl, 10,11-didehydro(E-): 10,11-Didehydro-4-demethylepothilone D**

[612057-49-5]

C₂₆H₃₇NO₅S 475.648Isol. from recombinant *Mycococcus xanthus*. Yellow oil. $[\alpha]_D$ -36.3 (c, 0.003 in MeOH). λ_{max} 213 (ε 22300); 242 (ε 27600) (MeOH).**4-Demethyl, 9-oxo, stereoisomer 1: 4-Demethyl-9-oxoepothilone D(1)**

[612057-55-3]

C₂₆H₃₇NO₆S 491.647Isol. from recombinant *Mycococcus xanthus*. Yellow oil. $[\alpha]_D$ -79.8 (c, 0.015 in MeOH). Probably C-4 epimer of stereoisomer 2. λ_{max} 210 (ε 30200); 248 (ε 22500) (MeOH).**4-Demethyl, 9-oxo, stereoisomer 2: 4-Demethyl-9-oxoepothilone D(2)**


[612057-55-3]

C₂₆H₃₇NO₆S 491.647Isol. from recombinant *Mycococcus xanthus*. Yellow oil. $[\alpha]_D$ +24.8 (c, 0.05 in MeOH). Probably C-4 epimer of stereoisomer 1. λ_{max} 212 (ε 25580); 248 (ε 19600) (MeOH).**6-Demethyl, 10,11-didehydro(E-): 10,11-Didehydro-6-demethylepothilone D**

[612057-50-8]

C₂₆H₃₇NO₅S 475.648Isol. from recombinant *Mycococcus xanthus*. Yellow oil. $[\alpha]_D$ -53.2 (c, 0.008 in MeOH). λ_{max} 215 (ε 24100); 240 (ε 34000) (MeOH).**A^{11,12}-Isomer, 9-oxo: 11,12-Didehydro-12,13-dihydro-9-oxoepothilone D**C₂₇H₃₉NO₆S 505.674Prod. by genetically engineered *Mycococcus xanthus*.**8-Epimer, 9-oxo: 8-Epi-9-oxoepothilone D**C₂₇H₃₉NO₆S 505.674Prod. by genetically engineered *Mycococcus xanthus*.Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 2097-2103; 1998, **37**, 84-87 (*synth*, *ir*, *pmr*, *cmr*)Chou, T.-C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1998, **95**, 9642-9647 (*pharmacol*)Mulzer, J. *et al.*, *J.O.C.*, 2000, **65**, 7456-7467 (*synth*)Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)White, J.D. *et al.*, *J.A.C.S.*, 2001, **123**, 5407-5413 (*synth*)Arslanian, R.L. *et al.*, *J. Nat. Prod.*, 2002, **65**, 570-572; 1061-1064 (*isol*, *Didehydroepothilone D*)Starks, C.M. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1313-1317 (*Demethylepothilones*)Kolman, A. *et al.*, *Curr. Opin. Invest. Drugs*, 2004, **5**, 657-667 (*rev*)Goodin, S. *et al.*, *J. Clin. Oncol.*, 2004, **22**, 2015-2025 (*pharmacol*)Jung, J.-C. *et al.*, *J.O.C.*, 2004, **69**, 9269-9284 (*synth*)Tang, L. *et al.*, *J. Antibiot.*, 2005, **58**, 178-184 (*9-Oxoepothilones*)Keck, G.E. *et al.*, *J.O.C.*, 2008, **73**, 9675-9691 (*synth*)**Epothilone H₁**

E-118




R = H

C₂₆H₃₉NO₆ 461.597Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -84 (c, 0.2 in MeOH). λ_{max} 203 (ε 19600); 237 (ε 12000) (MeOH).**12α,13α-Epoxyde: Epothilone G₁**C₂₆H₃₉NO₇ 477.597Prod. by *Sorangium cellulosum*.Amorph. solid. $[\alpha]_D^{22}$ -39.7 (c, 1 in MeOH). λ_{max} 203 (ε 15200); 236 (ε 15100) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone H₂**

E-119

As Epothilone H₁, E-118 withR = CH₃C₂₇H₄₁NO₆ 475.624Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -44 (c, 0.25 in MeOH). λ_{max} 203 (ε 14500); 236 (ε 12200) (MeOH).**12α,13α-Epoxyde: Epothilone G₂**C₂₇H₄₁NO₇ 491.623Prod. by *Sorangium cellulosum*.Amorph. solid. $[\alpha]_D^{22}$ -22.6 (c, 1 inMeOH). λ_{max} 202 (ε 21500); 236 (ε 14800) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone I₁**

E-120

R¹ = R⁴ = H, R² = R³ = CH₃C₂₈H₄₃NO₅S 505.717Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -9 (c, 0.02 in MeOH). λ_{max} 203 (ε 19100); 244 (ε 12500) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone I₂**

E-121

As Epothilone I₁, E-120 withR¹ = H, R² = R³ = R⁴ = CH₃C₂₉H₄₅NO₅S 519.744Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -68.5 (c, 1 in MeOH). λ_{max} 210 (ε 12600); 249 (ε 9200) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone I₃**

E-122

As Epothilone I₁, E-120 withR¹ = R² = R³ = R⁴ = CH₃C₃₀H₄₇NO₅S 533.771Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -64 (c, 0.17 in MeOH). λ_{max} 203 (ε 15800); 249 (ε 9000) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone I₄**

E-123


As Epothilone I₁, E-120 withR¹ = R² = H, R³ = R⁴ = CH₃C₂₈H₄₃NO₅S 505.717C-4 config. not determined. Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -51 (c, 0.5 in MeOH). λ_{max} 211 (ε 13400); 250 (ε 9800) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856**Epothilone I₅**

E-124

As Epothilone I₁, E-120 withR¹ = R² = R⁴ = CH₃, R³ = HC₂₉H₄₅NO₅S 519.744C-4 config. not determined. Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -36 (c, 0.5 in MeOH). λ_{max} 208 (ε 14400); 249 (ε 9700) (MeOH).**4-Epimer: Epothilone I₆**C₂₉H₄₅NO₅S 519.744Prod. by *Sorangium cellulosum*.Amorph. solid. $[\alpha]_D^{22}$ -40 (c, 0.35 in MeOH). λ_{max} 204 (ε 14600); 250 (ε 9000) (MeOH).Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone K

E-125

C₁₈H₁₈O₆ 330.337


C₂₂H₃₁NO₄S 405.557
Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{22}$ -7 (c, 0.08 in MeOH). λ_{\max} 212 (ε 16700); 248 (ε 12500) (MeOH).

Hardt, I. et al., *J. Nat. Prod.*, 2001, **64**, 847-856

Epothilone tetrahydrofuran

E-126

7,10-Epoxyepothilone




C₂₇H₃₉NO₆S 505.674
Isol. from recombinant *Myxococcus xanthus*. Pale yellow oil. λ_{\max} 210 (ε 19800); 247 (ε 13500) (MeOH).

Starks, C.M. et al., *J. Nat. Prod.*, 2003, **66**, 1313-1317 (isol, pmr, cmr)

1,4-Epoxy-3-aza-A-homoandrostane-17-ol, 9CI

E-127

3-Aza-A-homoandrostane-17-ol 1,4-epoxide
[25484-32-6]



C₁₉H₃₁NO₂ 305.459
Alkaloid from the salamander *Cryptobranchus maximus*. Needles (Me₂CO/hexane). Mp 186° Mp 191-193°.

Hara, S. et al., *J.A.C.S.*, 1967, **89**, 1041-1042 (synth)

Oka, K. et al., *Tet. Lett.*, 1969, 1987-1990 (synth)


Japan. Pat., 1970, 70 18 663; *CA*, **73**, 56316g
Benn, M. et al., *Can. J. Chem.*, 1974, **52**, 2936-2940 (synth, ms)

Shimizu, Y. et al., *J.O.C.*, 1976, **41**, 1930-1934 (synth)

4,7'-Epoxy-3,8'-bilign-7-ene-3',4',5,9,9'-pentol

E-128

4,7'-Epoxy-3',4',5,9,9'-pentahydroxy-3,8'-bilign-7-ene, 9CI



(7'R,8'R)-form

(7'R*,8'S*)-form

9-Carboxylic acid, 3',5-di-Me ether, 2-(4-hydroxyphenyl)ethylamide: **Grossamide K**
[403647-07-4]

C₂₈H₂₉NO₇ 491.54

Constit. of the bark of *Hibiscus cannabinus*. Yellowish oil.

9,9'-Dicarboxylic acid, 3',5-di-Me ether, bis[2-(4-hydroxyphenyl)ethylamide]: **Tataramide B**
[187655-56-7]

C₃₆H₃₆N₂O₈ 624.689

Isol. from *Acorus tatarinowii*. Possesses Z-config.

[4263-87-0, 150407-42-4, 112835-77-5, 28199-68-0, 86363-02-2]


Wang, M.F. et al., *Chin. Chem. Lett.*, 1997, **8**, 35-36 (Tataramide B)

Seca, A.M.L. et al., *Phytochemistry*, 2001, **56**, 759-767; **58**, 1219-1223 (Boehmenan K, Grossamide K)

Epoxychothalasin H

E-129

[80618-96-8]

C₃₀H₃₉NO₅ 493.642

Prod. by *Phoma* sp. and *Phomopsis sojae*. Inhibits the growth of plants; toxic. Cryst. (Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 134-136° (128°-130°). $[\alpha]_D^{30}$ -85.5 (c, 0.1 in CHCl₃). Not strictly the epoxide of Cytochalasin H in Z-37. λ_{\max} 260 (MeOH) (Berdy).

O-De-Ac: **Deacetylepoxycytochalasin H**

[80618-95-7]

C₂₈H₃₇NO₄ 451.605

Prod. by *Phomopsis sojae*. Plant growth inhibitor, toxic. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 192-194°. $[\alpha]_D^{30}$ -49.1 (c, 0.19 in CHCl₃). λ_{\max} 260 (MeOH) (Berdy).


Cole, R.J. et al., *J. Agric. Food Chem.*, 1982, **30**, 301

Kakeya, H. et al., *J. Nat. Prod.*, 1997, **60**, 669 (isol)

6,17-Epoxy-19,20-dihydro-1-methylsarpagan-17,21-diol, 9CI

E-130

Alkaloid G[†]
[149507-99-3]



Absolute Configuration

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from cell cultures of *Rauwolfia serpentina*. λ_{\max} 224 ; 281 ; 290 (MeOH).


Endreb, S. et al., *Phytochemistry*, 1993, **32**, 725-730 (isol, synth, pmr, cmr, cd)

Li, J. et al., *J.A.C.S.*, 1999, **121**, 6998-7010 (synth, ir, pmr, cmr)

Yu, J. et al., *J.O.C.*, 2003, **68**, 5852-5859 (synth)

12,16-Epoxy-16,17-dihydro-napelline

E-131


C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrum*. Cryst. (MeOH). Mp 232-233°. $[\alpha]_D^{20}$ -50 (c, 0.1 in CHCl₃).

Zhang, F. et al., *Planta Med.*, 2005, **71**, 1073-1076

14,15-Epoxy-3,5-dihydroxy-bufa-20,22-dienolide

E-132

C₂₄H₃₂O₅ 400.514

(3β,5β,14β,15β)-form

Marinobufagin. *Bufo* *Marinobufogenin*
[470-42-8]

Constit. of *Bufo marinus*, *Bufo formosus* and Ch'an Su. Cryst. (Me₂CO/Et₂O). Mp 224-225°. $[\alpha]_D^{16}$ +10 (c, 2.6 in CHCl₃).

▶ EI2963000

O³-Sulfate: **Marinobufogenin O³-sulfate**

[71595-47-6]

[72173-02-5 (NH₄ salt), 72173-03-6 (Na salt)]

C₂₄H₃₂O₈S 480.578

Isol. from skin of *Bufo marinus*. Prisms. Mp 156-160° dec. $[\alpha]_D^{20}$ +7 (c, 0.37 in MeOH) (monohydrate). Isol. as NH₄ salt to which the data refers, also isol. as oily Na salt.

O³-(L-Arginyl)suberoyl): **Marinobufotoxin**

[30685-91-7]

C₃₈H₅₆N₄O₉ 712.882

Component of the toxin of *Bufo ictericus* and *Bufo marinus*. Mp 174-

179°. $[\alpha]_D^{24} +24$ (CHCl₃/MeOH 1:1).
See note under Bufotoxin, B-394.

▶ OP9575000

O³-(L-Glutaminy)suberoyl): Marinobufagin 3-suberoylglutamine ester

[75090-30-1]

C₃₇H₅₂N₂O₁₀ 684.825

Constit. of *Bufo americanus*. Cryst. (MeOH/Et₂O). Mp 166-170°. $[\alpha]_D +15.1$ (c, 0.11 in MeOH).

O³-(L-Argininy)pimeloyl): Marinobufagin 3-pimeloylarginine ester

[72103-58-3]

C₃₇H₅₄N₄O₉ 698.855

Constit. of *Bufo marinus*. Amorph. Mp 186-190°. $[\alpha]_D^{20} +22$ (c, 0.23 in MeOH).

Schröter, H. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 1385-1394 (isol)

Gsell, L. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 551-568 (pmr)

Linde-Tempel, H. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 2188-2196 (*Marinobufotoxin*)


Porto, A.M. *et al.*, *Experientia*, 1971, **27**, 506 (biosynth)

Pettit, G.R. *et al.*, *J.O.C.*, 1974, **39**, 3003-3006 (*Marinobufotoxin, synth*)

Shimada, K. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 1881-1886; 1980, **28**, 1559-1562 (isol, pmr, sulfates, esters)

1,2-Epoxy-1-hydroxymethylpyrrolizidine E-133

Hexahydro-6bH-oxireno[a]pyrrolizine-6b-methanol, 9CI. 1β,2β-Epoxyisoretro-necanol, 8CI. 1-Hydroxymethyl-1,2-epoxy-pyrrolizidine
[15211-03-7]



Relative configuration

C₈H₁₃NO₂ 155.196

Alkaloid from *Crotalaria trifoliolatum*, *Crotalaria aridicola*, *Crotalaria grantiana*, *Heliotropium ternatum*, *Heliotropium molle*, *Heliotropium angiospermum* (major alkaloid) and *Heliotropium subulatum* (major alkaloid) (Fabaceae, Boraginaceae). Large prisms (Me₂CO). Mp 66-67°. $[\alpha]_D^{20} -60.1$ (c, 2.2 in EtOH). The alkaloid isol. from *H. subulatum* by Malik *et al* was originally named Subulacine N-oxide and assigned an erroneous struct.

Picrate:

Yellow needles (EtOH). Mp 173-174° (170-171°). $[\alpha]_D^{20} -8$ (c, 4.97 in Me₂CO).

Me ether: 1,2-Epoxy-1-methoxymethylpyrrolizidine. 1-Methoxymethyl-1,2-epoxypyrrolizidineC₉H₁₅NO₂ 169.223

Alkaloid from *Crotalaria trifoliolatum* and *Crotalaria aridicola* (Fabaceae). Bp_{0.1} 53°. $[\alpha]_D -63$ (c, 1.08 in EtOH). pK_a 8.1.

Me ether, picrate:

Yellow needles (EtOH). Mp 166-168°. $[\alpha]_D^{18} -7.5$ (c, 1.96 in Me₂CO).

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1963,

16, 131; 1967, **20**, 757 (isol, pmr, ms, struct)

Birecka, H. *et al.*, *Phytochemistry*, 1983, **22**, 1167 (isol, ms)

Smith, L.W. *et al.*, *Phytochemistry*, 1984, **23**, 473 (isol)


Malik, A. *et al.*, *Heterocycles*, 1988, **27**, 707

Winkler, T. *et al.*, *Heterocycles*, 1988, **27**, 2331

Stermitz, F.R. *et al.*, *Tet. Lett.*, 1988, **29**, 4943

5-(2,3-Epoxy-3-methylbutyl)-3-[2-hydroxy-1-(hydroxymethyl)-2-methylpropyl]-1H-indole E-134

2-[5-[(3,3-Dimethyloxiranyl)methyl]-1H-indol-3-yl]-3-methyl-1,3-butanediol, 9CI

C₁₈H₂₅NO₃ 303.4**(1'S,2''S)-form**

5'-Hexadecanoyl: [165604-37-5]

C₃₄H₅₅NO₄ 541.813Alkaloid from *Hexalobus crispiflorus*.

Oil. $[\alpha]_D^{21} +30$ (c, 1.4 in CHCl₃). λ_{max} 226 (log ε 4.6); 286 (log ε 3.9); 295 (log ε 3.9) (MeOH).

5'-(9,12-Octadecadienoyl) (Z,Z-):

[93822-73-2]

C₃₆H₅₅NO₄ 565.835Alkaloid from *Hexalobus crispiflorus*.

Oil. $[\alpha]_D^{21} +27$ (c, 1.4 in CHCl₃). λ_{max} 226 (log ε 4.6); 286 (log ε 3.9); 295 (sh) (MeOH).

5'-(9-Octadecenoyl) (Z-): [93822-72-1]


C₃₆H₅₇NO₄ 567.851Alkaloid from *Hexalobus crispiflorus*.

Oil. $[\alpha]_D^{21} +25$ (c, 1.8 in CHCl₃). λ_{max} 226 (log ε 4.6); 286 (log ε 3.9); 295 (sh) (MeOH).

Achenbach, H. *et al.*, *Annalen*, 1995, 1327-1337 (isol, uv, ir, pmr, cmr, ms)


1,2-Epoxy-2-methyl-4-(N-methyltyraminy)-3-butene E-135

2-[3-(3,4-Epoxy-3-methyl-1-butenyl)-4-hydroxyphenyl]-N-methylethylamine
[129743-99-3]

C₁₄H₁₉NO₂ 233.31

Alkaloid from the leaves of *Zanthoxylum coriaceum* (Rutaceae). Oil. Opt. inactive.

Marcos, M. *et al.*, *Phytochemistry*, 1990, **29**, 2315 (isol, uv, ir, pmr, cmr, ms, struct)

13,17-Epoxy-8,10(18)-myrsinadiene-3,5,7,14,15,17-hexol E-136C₂₀H₃₀O₇ 382.453

17-Hydroxymyrsinol is a misleading name since 13,17-Epoxy-3,5,7,15-tetrahydroxy-8,10(18)-myrsinadien-14-one has a C-14 oxo group.

(2β,3β,5α,7β,13α,14α,15β,17S)-form 17-Hydroxydihydromyrsinol. 17-Hydroxy-myrsinol (incorr.)

14-O-(3-Pyridinecarbonyl), 7-benzoyl, 3-propanoyl, 5,15-di-Ac: [213920-14-0]

C₄₀H₄₅NO₁₂ 731.795

Constit. of *Euphorbia seguieriana*. λ_{max} 221 (log ε 1.55); 258 (sh); 261 (log ε 0.72) (EtOH).

3,14-Bis-O-(3-pyridinecarbonyl), 7-O-(2-methylpropanoyl), 5,15-di-Ac: [213920-06-0]

C₄₀H₄₆N₂O₁₂ 746.81Constit. of *Euphorbia seguieriana*.

Cryst. Mp 205-207°. λ_{max} 220 (log ε 1.5); 258 (sh); 260 (log ε 0.7) (EtOH).

3,14-Bis-O-(3-pyridinecarbonyl), 7-O-(2-methylpropanoyl), 5,15,17-tri-Ac: [213920-10-6]

C₄₂H₄₈N₂O₁₃ 788.847

Constit. of *Euphorbia seguieriana*. λ_{max} 222 (log ε 1.55); 260 (log ε 0.73); 267 (sh); 281 (sh) (EtOH).

3,14-Bis-O-(3-pyridinecarbonyl), 7-benzoyl, 5,15-di-Ac: [213920-08-2]

C₄₃H₄₄N₂O₁₂ 780.827

Constit. of *Euphorbia seguieriana*. λ_{max} 221 (log ε 1.5); 251 (sh); 260 (log ε 0.73) (EtOH).

3,7,14-Tris-O-(3-pyridinecarbonyl), 5,15-di-Ac: [213920-19-5]


C₄₂H₄₃N₃O₁₂ 781.815

Constit. of *Euphorbia seguieriana*. λ_{max} 222 (log ε 1.5); 258 (sh); 260 (log ε 0.7); 280 (sh) (EtOH).

Öksüz, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1198-1201 (isol, pmr, cmr)

2,3-Epoxy-6,8-nonadienoic acid E-137

3-(3,5-Hexadiynyl)oxiranecarboxylic acid

C₉H₈O₃ 164.16

2-Phenylethylamide: 3-(3,5-Hexadiynyl)-N-(2-phenylethyl)oxiranecarboxamide.


2,3-Epoxy-6,8-nonadiynoic acid 2-phenylethylamide
[96917-25-8]
C₁₇H₁₇NO₂ 267.327
Isol. from *Acmella ciliata* and *Salmea scandens*. Oil.

(Z)-2-Phenylethylamide: 3-(3,5-Hexadiynyl)-N-(2-phenylethyl)oxirane-carboxamide, 9CI, 1,2-Epoxy-6,8-nonadiynoic acid styrylamide
[75872-78-5]
C₁₇H₁₅NO₂ 265.311

Isol. from *Spilanthes alba*. Oil.

Bohlmann, F. et al., *Phytochemistry*, 1980, **19**, 1535; 1985, **24**, 595 (isol. struct, pmr, ir, ms)
Martin, R. et al., *Phytochemistry*, 1985, **24**, 2295 (isol. struct, pmr, ms)

5,20-Epoxy-2,4,7,10,13-pentahydroxy-11-taxen-9-one E-138



C₂₀H₃₀O₇ 382.453

(2α,4α,5β,7β,10β,13α)-form


2-Benzoyl, 4,10-di-Ac: 1-Dehydroxybaccatin III
[132302-29-5]
C₃₁H₃₈O₁₀ 570.635
Constit. of *Taxus yunnanensis*. Needles (MeOH). Mp 169-171°. [α]_D²³ -45.9 (c, 0.75 in CHCl₃).

13-O-(3S-Benzoylamino-2R-hydroxy-3-phenylpropanoyl), 2-benzoyl, 4,10-di-Ac: 1-Deoxytaxol
[198990-70-4]
C₄₇H₅₁NO₁₃ 837.919
Constit. of *Taxus mairei*. Amorph. powder. [α]_D²² -57 (c, 0.05 in CHCl₃).

Zhang, Z. et al., *Phytochemistry*, 1990, **29**, 3673-3675 (1-Dehydroxybaccatin III)

Shi, Q.-W. et al., *J. Nat. Prod.*, 2006, **69**, 280-283 (1-Deoxytaxol)

16,23-Epoxy-16,28-secosolanidane-3,23,27-triol E-139



C₂₇H₄₅NO₄ 447.657

(3β,5α,22S,23R,25S)-form


Esculeogenin B
[854381-41-2]
Amorph. powder. [α]_D -96.2 (c, 0.05 in Py).

3-O-[[β-D-Glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]]-β-D-glucopyranosyl-(1→4)-β-D-galactopyranoside], 27-O-β-D-glucopyranoside: Esculeoside B
[719294-52-7]
C₅₆H₉₃NO₂₈ 1228.341

Alkaloid from ripe tomatoes (*Lycopersicon esculentum*). Amorph. powder. [α]_D²⁶ -49.2 (c, 0.6 in Py).

Fujiwara, Y. et al., *Tetrahedron*, 2004, **60**, 4915-4920 (Esculeoside B)
Yoshizaki, M. et al., *Chem. Pharm. Bull.*, 2005, **53**, 839-840 (Esculeogenin B)

16,23-Epoxy-16,28-secosolanid-22(28)-ene-3,20,23,27-tetrol E-140



C₂₇H₄₃NO₅ 461.64


(3β,5α,20S,23R,25S)-form

23-Me ether, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]]-β-D-glucopyranosyl-(1→4)-β-D-galactopyranoside], 27-O-β-D-glucopyranoside: Esculeoside C
[886225-73-6]
C₅₇H₉₃NO₂₉ 1256.351

Constit. of the ripe fruit of *Lycopersicon esculentum* var. *cerasiforme* (cherry tomato). Amorph. powder. [α]_D²⁴ -56.7 (c, 0.9 in Py).

Ono, M. et al., *Chem. Pharm. Bull.*, 2006, **54**, 237-239 (Esculeoside C)

5,20-Epoxy-11-taxene-1,2,4,7,9,10,13-heptol E-141



C₂₀H₃₂O₈ 400.468

(1β,2α,5β,7β,9α,10β,13α)-form

10-Benzoyl, 13-(3-dimethylamino-3-phenylpropanoyl) (R-), 2,4,7,9-tetra-Ac: Taxuspine N
[175801-13-5]
C₄₆H₅₇NO₁₄ 847.955

Constit. of *Taxus cuspidata*. Amorph. solid. [α]_D²⁵ -6 (c, 0.26 in CHCl₃). λ_{max} 278 (ε 4900) (MeOH) (Berdy).

9-Ketone, 2-benzoyl, 13-[2-hydroxy-3-(4-methylhexanoylamino)-3-phenylpropanoyl], 4,10-di-Ac: Taxuspianane A
[187740-00-7]
C₄₇H₅₉NO₁₄ 861.981


Constit. of *Taxus cuspidata* var. *nana*.

Powder. [α]_D -40.2 (c, 0.37 in MeOH).

Kobayashi, J. et al., *Tetrahedron*, 1995, **51**, 5971-5978; 1996, **52**, 5391-5396 (*Taxuspines*)
Morita, H. et al., *J. Nat. Prod.*, 1997, **60**, 390-392 (*Taxuspianane A*)
Buchta, M. et al., *Z. Kristallogr.*, 2006, **221**, 97-100 (*Taxuspianane A*, *cryst struct*)

Equisetin E-142

[57749-43-6]



Absolute Configuration

C₂₂H₃₁NO₄ 373.491

Tetramic acid antibiotic. Enolised triketone. Isol. from *Fusarium equiseti* and *Fusarium heterosporum*. Inhibitor of HIV-1 integrase. Active against gram-positive organisms. Amorph. powder. Mp 65-66°. [α]_D²² -278 (c, 0.77 in CHCl₃) ((natural)). [α]_D²² -352 (c, 1 in CHCl₃). λ_{max} 234 (ε 5680); 294 (ε 8860) (EtOH/HCl) (Derep). λ_{max} 252 (ε 8210); 292 (ε 7400) (EtOH/NaOH) (Derep). λ_{max} 232 (ε 6900); 248 (ε 7500); 292 (ε 10000) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 63 mg/kg. KD7986000

N-De-Me: N-Demethylequisetin. *Trichosetin*

C₂₁H₂₉NO₄ 359.464

Isol. from a co-culture of *Catharanthus roseus* and *Trichoderma harzianum*. Phytotoxin. Active against gram-positive bacteria. Pale orange amorph. solid. [α]_D²⁷ -471.6 (c, 0.1 in MeOH). λ_{max} 204 (ε 15000); 252 (ε 5700); 288 (ε 10000) (MeOH).

U.S. Pat., 1974, 467 548; CA, **84**, 41985 (isol)
Vesonder, R.F. et al., *J. Antibiot.*, 1979, **32**, 759 (nmr, ms, ir, uv, struct)

Phillips, N.J. et al., *J.A.C.S.*, 1989, **111**, 8223 (struct)

Turos, E. et al., *J.A.C.S.*, 1989, **111**, 8231 (synth)

Singh, S.B. et al., *Tet. Lett.*, 1998, **39**, 2243-2246 (isol, cmr)

Hazuda, D. et al., *Antiviral Chem. Chemother.*, 1999, **10**, 63-70 (isol, activity)

Yuki, K. et al., *Tet. Lett.*, 2001, **42**, 2517-2519 (synth)

Marfori, E.C. et al., *Tetrahedron*, 2002, **58**, 6655-6658 (*Trichosetin*, biosynth)

Marfori, E.C. et al., *Z. Naturforsch., C*, 2002, **57**, 465-470 (*Trichosetin*, isol, pmr, cmr)

Marfori, E.C. et al., *Phytochemistry*, 2003, **62**, 715-721 (*Trichosetin*, activity)

Burke, L.T. et al., *Org. Biomol. Chem.*, 2005, **3**, 274-280 (synth)

Equisetone E-143

C₁₈H₃₁N₃O₄ 353.461


Struct. unknown. Alkaloid from *Equisetum palustre*. Mp 130-135° Mp 220° dec. (as hydrochloride).

Wöhlbier, W. et al., *Chem. Ber.*, 1950, **83**, 310-314

Eugster, C.H. et al., *Helv. Chim. Acta*, 1953, **36**, 1387-1400

Erbstatin**E-144**

N-[2-(2,5-Dihydroxyphenyl)ethyl]formamide. MH 435A. Antibiotic MH 435A [100827-28-9]

C₉H₉NO₃ 179.175

Isol. from *Streptomyces* sp. MH435-hF3. Epidermal growth factor-receptor; tyrosine kinase inhibitor; exhibits antimicrobial and antineoplastic props; angiogenesis inhibitor; DNA topoisomerase inhibitor; induces cell apoptosis; mitogenic response inhibitor. Yellow cryst. (MeOH/CHCl₃). Sol. MeOH, EtOH, Me₂CO, acids; fairly sol. CHCl₃, EtOAc; poorly sol. H₂O, hexane. Mp 78-82° Mp 150.5-152° (softens at 88-91°) (synthetic). Log P 0.62 (calc). λ_{max} 215 (ε 8200); 278 (ε 14900); 330 (ε 8100) (MeOH/HCl) (Derep). λ_{max} 208 (ε 10700) (MeOH/NaOH) (Derep). λ_{max} 215 (ε 8200); 278 (ε 14900); 286 (sh) ; 330 (ε 8100) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 200 mg/kg. KD9500000

1',2'-Dihydro- N-[2-(2,5-Dihydroxyphenyl)ethyl]formamide. MH 435B. Antibiotic MH 435B

[108997-59-7]

C₉H₁₁NO₃ 181.191From *Streptomyces* sp. MH 435-hF₃.

[108536-24-9]

Umezawa, H. *et al.*, *J. Antibiot.*, 1986, **39**, 170; 314 (isol, struct, props)

Hangauer, D.G. *et al.*, *Tet. Lett.*, 1986, **27**, 5799 (synth, Z-form, pmr)

Isshiki, K. *et al.*, *J. Antibiot.*, 1987, **40**, 1207; 1209 (synth, props)

Dow, R.L. *et al.*, *Tet. Lett.*, 1987, **28**, 2217 (synth)

Deshmukh, M.N. *et al.*, *Synth. Commun.*, 1988, **18**, 1483 (synth)

Kleinschroth, J. *et al.*, *Synthesis*, 1988, 970 (synth)

Ishibashi, H. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2214 (synth)

Takekura, N. *et al.*, *Int. J. Cancer*, 1991, **47**, 938 (activity)

Umezawa, K. *et al.*, *Methods Enzymol.*, 1991, **201**, 379 (activity)

Hsu, C.Y. *et al.*, *Biochem. Pharmacol.*, 1992, **43**, 2471 (enzyme activity)

Oikawa, T. *et al.*, *J. Antibiot.*, 1993, **46**, 785

Markovits, J. *et al.*, *Biochem. Pharmacol.*, 1994, **48**, 549 (pharmacol)

Ercine**E-145**C₂₅H₂₈N₂O₆ 452.506

Struct. unknown. Alkaloid from the upper parts of *Vinca erecta* (Apocynaceae). Cryst. (C₆H₆). Mp 158-159°. [α]_D¹⁸ -121 (c, 1.4 in MeOH).

Hydrochloride:


Cryst. (H₂O). Mp 172-173°. [α]_D¹⁸ -108 (c, 1.4 in MeOH). [α]_D¹⁸ -116.6 (c, 0.2 in EtOH).

Koretskaya, N.I. *et al.*, *Zh. Obshch. Khim.*,

1963, **33**, 2065-2066; *J. Gen. Chem. USSR (Engl. Transl.)*, 1963, **33**, 2010 (isol, uv, ir)

Eremophilene lactam**E-146**

[65012-44-4]


C₁₅H₂₃NO 233.353

Alkaloid from the rhizomes of *Petasites hybridus* (Apocynaceae). Cryst. (C₆H₆ or C₆H₆/Me₂CO). Mp 215-218°. [α]_D²² +199.5 (c, 0.33 in MeOH).

Jizba, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 2438 (isol, uv, ir, pmr, ms, cd, struct)

β,β-Ergoannam**E-147**

[97387-95-6]




Absolute Configuration

R = R' = -CH(CH₃)CH₂CH₃C₃₃H₄₃N₅O₄ 573.734Alkaloid from *Claviceps purpurea*.

Flieger, M. *et al.*, *J. Nat. Prod.*, 1984, **47**, 970 (isol, pmr, cmr, ms, struct)

Ergobalansine**E-148**

[131985-45-0]

C₂₈H₃₅N₅O₅ 521.615

Alkaloid from *Cenchrus echinatus* infected with *Balansia obtecta*. Also prod. from cultures of *Balansia cyperi* and *Balansia obtecta*. Detected in seeds of *Ipomoea piurensis* (Convolvulaceae). Amorph.

8-Epimer: Ergobalansinine

[132074-97-6]

C₂₈H₃₅N₅O₅ 521.615


Alkaloid from *Cenchrus echinatus* infected with *Balansia obtecta*. Also prod. from cultures of *Balansia cyperi* and *Balansia obtecta*. Isol. from seeds of *Ipomoea piurensis* (Convolvulaceae). Amorph.

Powell, R.G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1272 (isol, pmr, cmr, struct)

Jennett-Siems, K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1304 (isol, pmr, ms)

Ergobine**E-149**

[149537-27-9]

R = CH₂CH₃, R' = CH₃C₂₈H₃₃N₅O₅ 519.599

Alkaloid from submerged cultures of *Claviceps purpurea* strain 231 F.I.

Crespi Perellino, N. *et al.*, *J. Nat. Prod.*, 1993, **56**, 489-493 (isol, pmr, ms, struct)

Ergobutine**E-150**

[82564-35-0]

As Ergobine, E-149 with

R = R' = -CH₂CH₃C₂₉H₃₅N₅O₅ 533.626

Isol. from saprophytic cultures of *Claviceps purpurea* strain 231 F.I.

Bianchi, M.L. *et al.*, *J. Nat. Prod.*, 1982, **45**, 191 (isol, ms, struct)

Ergobutyryne**E-151**

[82564-36-1]

As Ergobine, E-149 with

R = -CH₂CH₃, R' = -CH(CH₃)₂C₃₀H₃₇N₅O₅ 547.653

Isol. from saprophytic cultures of *Claviceps purpurea* strain 231 F.I.

Bianchi, M.L. *et al.*, *J. Nat. Prod.*, 1982, **45**, 191 (isol, ms, struct)

Ergocornam**E-152***Lysergylvalylvalylproline lactam*

[77477-94-2]

As β,β-Ergoannam, E-147 with

R = R' = -CH(CH₃)₂C₃₁H₃₉N₅O₄ 545.68


Alkaloid from field ergot. Mp 196-198°. [α]_D²⁰ +57.8 (c, 0.5 in Py).

Flieger, M. *et al.*, *J. Chromatogr.*, 1981, **207**, 139 (isol)

Stuchlik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1982, **47**, 3312 (isol, pmr, cmr, ms, struct)

Ergocornine**E-153**

12'-Hydroxy-2',5'-bis(1-methylethyl)ergotaman-3',6',18-trione, 9CI
[564-36-3]



Absolute Configuration

C₃₁H₃₉N₅O₅ 561.68

Alkaloid from ergot (*Claviceps purpurea*). Vasoconstrictor. Investigational tool for studying ovum implantation in mammals. Pharmacol. effects resemble those of Ergotamine, E-172 but is more toxic and little used clinically. Sol. Me₂CO, CHCl₃, EtOAc; fairly sol. EtOH; poorly sol. H₂O. Mp 182-184° dec. [α]_D²⁰ -175 (c, 1 in CHCl₃). [α]_D -110 (c, 1 in Py). Log P 5.07 (uncertain value) (calc). Component of Ergotoxine. λ_{\max} 311 (ϵ 8130) (MeOH) (Berdy).

► Exp. reprod. effects ; LD₅₀ (rbt, ivn) 1.17 mg/kg. KE7630000

Me ether: 12'-O-Methylergocornine

[112767-18-7]

C₃₂H₄₁N₅O₅ 575.706Alkaloid from *Claviceps purpurea*.

9,10 α -Dihydro: Dihydroergocornine

[25447-65-8]

[5611-87-0]

Mp 185-187° dec. [α]_D²⁰ -48 (c, 0.5 in Py).

8-Epimer: Ergocorninine

[564-37-4]

C₃₁H₃₉N₅O₅ 561.68

Alkaloid from ergot. Prisms (MeOH).

Mp 228° dec. [α]_D²⁰ +409 (c, 1.0 in CHCl₃). [α]_D²⁰ +500 (c, 1.0 in Py).

[8047-28-7, 8006-25-5]

Stoll, A. et al., *Helv. Chim. Acta*, 1943, **26**, 2070-2081; 1951, **34**, 1544 (struct)

Stadler, P.A. et al., *Helv. Chim. Acta*, 1969, **52**, 1549 (synth, config, uv, ir, pmr, epimer)

Voigt, D. et al., *Pharmazie*, 1974, **29**, 697 (ms)

Wurst, M. et al., *J. Chromatogr.*, 1979, **174**, 401 (hplc)

Plattner, R.D. et al., *J. Agric. Food Chem.*, 1983, **31**, 785 (ms)

Crespi-Perellino, N. et al., *J. Nat. Prod.*, 1987, **50**, 1065 (12'-O-Methylergocornine)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1366

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EDA600; EDC565; EDC575

Ergocristam**E-154**

Lysergylvalylphenylalanylproline lactam
[50868-53-6]

As β,β -Ergoannam, E-147 with R = -CH(CH₃)₂, R' = -CH₂Ph

C₃₅H₃₉N₅O₄ 593.724

Alkaloid from *Claviceps purpurea*. Cryst. (C₆H₆ or CH₂Cl₂/Et₂O). Mp 234-236° dec. [α]_D²⁰ +8 (c, 0.5 in CHCl₃). [α]_D +41 (c, 0.5 in Py).

Stütz, P. et al., *Experientia*, 1973, **29**, 936 (isol,


uv, ir, pmr, struct)

Černý, A. et al., *Coll. Czech. Chem. Comm.*, 1976, **41**, 3415 (isol, uv, ir, pmr, struct)

Flieger, M. et al., *J. Chromatogr.*, 1981, **207**, 139 (isol)

Ergocristine**E-155**

12'-Hydroxy-2'-(1-methylethyl)-5'-(phenylmethyl)ergotaman-3',6',18-trione, 9CI
[511-08-0]



Absolute configuration

C₃₅H₃₉N₅O₅ 609.724

Alkaloid from ergot (*Claviceps purpurea*). Vasoconstrictor. Pharmacol. effects resemble those of Ergotamine, E-172 but is more toxic, therefore little used clinically. Mp 165-170°. [α]_D²⁰ -173 (c, 0.7 in CHCl₃). [α]_D -101 (c, 0.8 in Py). Log P 5.56 (uncertain value) (calc). Component of Ergotoxine.

► KE1180000

9,10-Dihydro: Dihydroergocristine, BAN.

Many other names

[17479-19-5]

C₃₅H₄₁N₅O₅ 611.739

Used in treatment of impaired mental function in the elderly. Peripheral and cerebral vasodilator. Log P 5.86 (uncertain value) (calc). Mainly used as a mix. with other dihydroergot alkaloids.

9,10-Dihydro; methanesulfonate: Dihydroergocristine mesilate, BAN

[24730-10-7]

8-Epimer: Ergocristinine

[511-07-9]

C₃₅H₃₉N₅O₅ 609.724

Alkaloid from ergot. Prisms (MeOH).

Mp 226-227° dec. (214° dec.). [α]_D²⁰ +370 (c, 0.4 in CHCl₃). [α]_D²⁰ +471 (c, 0.35 in Py).

2'-Epimer: Aciergocristine

[74081-45-1]

C₃₅H₃₉N₅O₅ 609.724

Minor ergot alkaloid occurring as artifact of isomerisation. Identified by hplc.

2',8-Diepimer: Aciergocristinine

[74081-46-2]

C₃₅H₃₉N₅O₅ 609.724

Minor ergot alkaloid occurring as artifact of epimerisation. Identified by hplc.

[8067-24-1, 8006-25-5]

Stoll, A. et al., *Helv. Chim. Acta*, 1951, **34**, 1544 (struct)

Stadler, P.A. et al., *Helv. Chim. Acta*, 1969, **52**, 1549 (synth, config, uv, ir, epimer)

Voigt, D. et al., *Pharmazie*, 1974, **29**, 697 (ms)

Wurst, M. et al., *J. Chromatogr.*, 1979, **174**, 401 (hplc)

Szepesi, G. et al., *J. Chromatogr.*, 1980, **191**, 101 (Aciergocristine, Aciergocristinine)

Plattner, R.D. et al., *J. Agric. Food Chem.*, 1983, **31**, 785 (ms)

Coppi, G. et al., *Arzneim.-Forsch.*, 1992, **42**, 1381; 1391 (Dihydroergocristine, rev)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1366

Ergocryptam**E-156**

Lysergylvalylleucylproline lactam

[77477-95-3]

As β,β -Ergoannam, E-147 withR = -CH(CH₃)₂, R' = -CH₂CH(CH₃)₂C₃₂H₄₁N₅O₄ 559.707

Alkaloid from field ergot. Mp 108-110°.

[α]_D²⁰ +31.1 (c, 0.5 in Py).


Flieger, M. et al., *J. Chromatogr.*, 1981, **207**, 139 (isol)

Stuchlk, J. et al., *Coll. Czech. Chem. Comm.*, 1982, **47**, 3312 (isol, pmr, cmr, ms, struct)

 α -Ergocryptine**E-157**

12'-Hydroxy-2'-(1-methylethyl)-5'-(2-methylpropyl)ergotaman-3',6',18-trione, 9CI. Ergocryptine. Ergokryptine. Ergomolline

[511-09-1]



Absolute configuration

C₃₂H₄₁N₅O₅ 575.706

Alkaloid from ergot (*Claviceps purpurea*). Pharmacol. effects resemble those of Ergotamine, E-172 but is more toxic and little used clinically. Mp 211-212° dec. (MeOH). [α]_D²⁰ -120 (c, 1 in Py). [α]_D²⁰ -198 (c, 1 in CHCl₃).

► LD₅₀ (rbt, ivn) 0.95 mg/kg. KE1400000

Me ether: 12'-O-Methyl- α -ergocryptine

C₃₃H₄₃N₅O₅ 589.733Alkaloid from *Claviceps purpurea*.

8-Hydroxy: 8-Hydroxy- α -ergocryptine

C₃₂H₄₁N₅O₆ 591.706

Alkaloid from sclerotia of *Claviceps purpurea*. Amorph.

2-Bromo: Bromocriptine, BAN, INN,

USAN. Bromocryptine. Axialit. Bagren. Bromergon. Parilac. Parlodol. Parodel. Umprel. Cabl 154. NSC 169774. Bromergocryptine. Many other names

[25614-03-3]

C₃₂H₄₀BrN₅O₅ 654.602

Dopamine D₂-receptor agonist used for treatment of prolactinomas. Antiparkinsonian agent. Free radical scavenger antioxidant, prolactin inhibitor. Cryst. Mp 215-218° dec. [α]_D²⁰ -195 (c, 1 in CH₂Cl₂). Log P 6.69 (uncertain value) (calc).

► Human and exp. teratogen. Exp. reprod. effects. LD₅₀ (rat, ivn) 72 mg/kg. KE8250000

2-Bromo; methanesulfonate salt: Bromocriptine mesilate, JAN, USAN

[22260-51-1]

Cryst. (2-butanone). Mp 192-196° dec. [α]_D²⁰ +95 (c, 1 in MeOH/CH₂Cl₂).

► Adverse systemic effects reported when

used therapeutically (gastrointestinal, psychiatric, and cardiovascular). Exp. reprod. and teratogenic effects. LD₅₀ (rat, ivn) 10.5 mg/kg. KE1595000

8-Epimer: α-Ergocryptinine

[511-10-4]

Formed when α-Ergocryptine is boiled with MeOH. Cryst. (EtOH). Mp 238° dec. [α]_D²⁰ +405 (c, 0.5 in CHCl₃). [α]_D²⁰ +485 (c, 0.5 in Py).

Schlientz, W. et al., *Pharm. Acta Helv.*, 1968, **43**, 497 (isol, ir, pmr)

Ger. Pat., 1969, 1 926 045; CA, **72**, 43969b (deriv)

Stadler, P.A. et al., *Helv. Chim. Acta*, 1969, **52**, 1549 (synth, ir, uv, pmr)

Voigt, D. et al., *Pharmazie*, 1974, **29**, 697 (ms)

Giron-Forest, D.A. et al., *Anal. Profiles Drug Subst.*, 1979, **8**, 47 (rev, Bromocriptine, anal, synth)

Mehta, A.E. et al., *Drugs*, 1979, **17**, 313 (rev, Bromocriptine, pharmacol)

Wurst, M. et al., *J. Chromatogr.*, 1979, **174**, 401 (hplc)

Thorner, M.O. et al., *Bromocriptine: A Clinical and Pharmacological Review*, Raven Press, N.Y., 1980, (book)

Rucman, R. et al., *Farmaco, Ed. Sci.*, 1983, **38**, 406 (synth, Bromocriptine)

Barbien, R.L. et al., *Fertil. Steril.*, 1983, **39**, 727 (rev, Bromocriptine)

Kanto, J. et al., *Int. J. Clin. Pharmacol., Ther. Toxicol.*, 1983, **21**, 135 (Bromocriptine)

Crespi-Perellino, N. et al., *J. Nat. Prod.*, 1987, **50**, 1065 (12'-O-Methyl-α-ergocryptine)

Yoshikawa, T. et al., *J. Neurochem.*, 1994, **62**, 1034 (pharmacol)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 11291 (synonyms)

Cvak, L. et al., *Phytochemistry*, 1997, **44**, 365 (8-Hydroxy-α-ergocryptine)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, No. 1132

Moldvai, I. et al., *J.O.C.*, 2004, **69**, 5993-6000 (synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, BNB250; EDB100; BNB325

β-Ergocryptine

E-158

β-Ergokryptine

[20315-46-2]

As Ergobine, E-149 with

R = -CH(CH₃)CH₂CH₃ (S-), R' = -CH(CH₃)₂

C₃₂H₄₁N₅O₅ 575.706

Alkaloid from ergot (*Claviceps purpurea*).

Antimigraine agent. Plates (C₆H₆). Mp

173° dec. [α]_D²⁰ -174 (c, 1.5 in CHCl₃).

[α]_D²⁰ -91 (c, 2.0 in Py).

▶ KE1590000

5'-Epimer: 5'-Epiergocryptine

C₃₂H₄₁N₅O₅ 575.706

Alkaloid from *Claviceps purpurea*.

8-Epimer: β-Ergocryptinine

[19467-61-9]

By action of alkali, acid or hot MeOH on β-Ergocryptine. Long needles (MeOH). Mp 220° dec. [α]_D²⁰ +424 (c, 1.0 in CHCl₃). [α]_D²⁰ +492 (c, 1.0 in Py).

Schlientz, W. et al., *Experientia*, 1967, **23**, 991 (isol)

Stadler, P.A. et al., *Helv. Chim. Acta*, 1969, **52**, 1549 (synth, ir, uv, pmr)


Flieger, M. et al., *J. Nat. Prod.*, 1984, **47**, 970 (epimer)

γ-Ergocryptinine

E-159

γ-Ergokryptinine

[881650-40-4]



Absolute Configuration

C₃₂H₄₁N₅O₅ 575.706

Alkaloid from *Claviceps purpurea* CCM 8059.

Cvak, L. et al., *Amino Acids*, 2005, **29**, 145-150 (isol, pmr, cmr, ms, cryst struct)

Ergogaline

E-160

[157135-96-1]

As Ergobine, E-149 with

R = -CH₂CH(CH₃)CH₂CH₃ (R-), R' = -CH(CH₃)₂

C₃₃H₄₃N₅O₅ 589.733

Alkaloid from *Claviceps purpurea*. Cryst. + 1H₂O. Mp 182°.

Cvak, L. et al., *J.C.S. Perkin 2*, 1994, 1861-1865 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Ergoheptine

E-161

[62818-83-1]

As Ergobine, E-149 with

R = -CH₂CH₂CH(CH₃)₂, R' = -CH(CH₃)₂

C₃₃H₄₃N₅O₅ 589.733

Alkaloid from ergot.

Ohmomo, S. et al., *Nippon Nogei Kagaku Kaishi*, 1976, **50**, 543; CA, **86**, 190303x (struct)

Ergohexine

E-162

[62818-82-0]

As Ergobine, E-149 with

R = -CH₂CH₂CH(CH₃)₂, R' = -CH₃

C₃₁H₃₉N₅O₅ 561.68


Alkaloid from ergot.

Ohmomo, S. et al., *Nippon Nogei Kagaku Kaishi*, 1976, **50**, 543; CA, **86**, 190303x (struct)

Ergoladinine

E-163

[176391-56-3]



C₃₁H₃₉N₅O₅ 593.746


The first sulfur-containing ergot alkaloid. Alkaloid from sclerotia of *Claviceps purpurea*. Cryst. (MeOH). Mp not reported.

Cvak, L. et al., *Phytochemistry*, 1996, **42**, 231 (isol, pmr, cmr, ms, cryst struct)

Ergometrine, BAN, INN

E-164

9,10-Didehydro-N-(2-hydroxy-1-methyl-ethyl)-6-methylergoline-8-carboxamide, 9CI. Ergonovine. Ergotrate. Ergobasine. Ergotocin. Ergostetrine



(+)-form

C₁₉H₂₃N₃O₂ 325.41

5HT receptor antagonist. Oxytocic agent used in childbirth. Shows little adrenergic or CNS effect. Log P 1.03 (uncertain value) (calc).

(+)-form [60-79-7]

Alkaloid from ergot (*Claviceps purpurea*) and several spp. in the genera *Argyriaea*, *Stictocardia*, *Rivea* and *Ipomoea* (Convolvulaceae). 5-Hydroxytryptamine antagonist. Plates (EtOAc), needles (Me₂CO). Mp 162-163° dec. Mp 212° dec. (dimorph.). Pharmacol. active isomer.

▶ Adverse systemic effects by intramuscular route. LD₅₀ (mus, ivn) 144 mg/kg. KE5075000

Hydrochloride: Mp 245-246° dec. [α]_D²⁵ +63 (H₂O).

Maleate salt: **Ergonovine maleate, USAN. Ergometrine maleate, JAN. Cornocentin. Ermetrine. Syntometrine. Many other names**

[129-51-1] Used in the prevention and treatment of postpartum haemorrhage. Mp 167° dec. [α]_D²⁵ +48.

▶ Adverse effects reported when used therapeutically. LD₅₀ (mus, ivn) 8.26 mg/kg. KE5250000

Picrate:

Red prisms or hydrated yellow needles. Mp 188-189° dec. (anhyd.) Mp 148° dec. (hydrated).

8-Epimer: Ergometrinine. Ergobasine.

Ergonovinine

[479-00-5]

C₁₉H₂₃N₃O₂ 325.41

Alkaloid from ergot and several *Argyriaea*, *Stictocardia*, *Rivea* and *Ipomoea* spp. (Convolvulaceae). Prisms (Me₂CO). Mp 195-197° dec. [α]_D²⁰ +414 (c, 0.45 in CHCl₃). [α]_D²⁰ +328 (c, 0.7 in MeOH).

▶ KE5067000

8-Epimer; hydrochloride:

Needles + 1H₂O. Mp 175-180° dec.

(-)-form

Obt. by partial synth. Mp 159-162° dec. [α]_D²⁰ -89 (H₂O). Physiologically inactive.

8-Epimer: Synthetic. Prisms (Me₂CO).

Mp 196° dec. $[\alpha]_D^{20}$ -415 (c, 0.4 in CHCl_3).

[129-50-0]

- Jacobs, W.A. *et al.*, *Science (Washington, D.C.)*, 1935, **82**, 16 (struct)
 Smith, S. *et al.*, *J.C.S.*, 1936, 1166 (epimer, isol)
 Stoll, A. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1938, **251**, 155 (epimer, synth)
 Stoll, A. *et al.*, *Helv. Chim. Acta*, 1943, **26**, 922; 944 (synth)
 Kornfeld, E.C. *et al.*, *J.A.C.S.*, 1956, **78**, 3087 (synth)
 Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435 (occur)
 Bach, N.J. *et al.*, *J.O.C.*, 1974, **39**, 1272 (emr)
 Reif, V.D. *et al.*, *Anal. Profiles Drug Subst.*, 1982, **11**, 273 (rev)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1365
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EDB500; LJL000

Ergonine E-165

[29537-61-9]

As Ergobine, E-149 with

R = $-\text{CH}(\text{CH}_3)_2$, R' = $-\text{CH}_2\text{CH}_3$

$\text{C}_{30}\text{H}_{37}\text{N}_5\text{O}_5$ 547.653

Alkaloid from *Claviceps purpurea*. Prisms (EtOH/diisopropyl ether). Mp 207-208°. $[\alpha]_D^{20}$ -182 (c, 1 in CHCl_3). $[\alpha]_D^{20}$ -28.2 (c, 1 in Py).

Stütz, P. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1278 (synth, uv, ir, pmr)

Brunner, R. *et al.*, *Can. J. Chem.*, 1979, **57**, 1638 (isol, uv, ir, pmr)

β -Ergoptine E-166

[65756-55-0]

As Ergobine, E-149 with

R = $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, R' = $-\text{CH}_2\text{CH}_3$

$\text{C}_{31}\text{H}_{39}\text{N}_5\text{O}_5$ 561.68

Isol. from ergot *Claviceps purpurea*. Mp 201-203° dec. $[\alpha]_D^{20}$ -163 (c, 1.0 in CHCl_3).

► KE7100000

8-Epimer: β -Ergoptinine

[65794-73-2]

Synthetic. Mp 205-206° dec. $[\alpha]_D^{20}$ +421 (c, 0.5 in CHCl_3).

Stadler, P.A. *et al.*, *Experientia*, 1977, **33**, 1552 (β -Ergoptinine)

Plattner, R.D. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 785 (ms)

Porter, J.K. *et al.*, *J. Agric. Food Chem.*, 1987, **35**, 359 (isol)

Ergoptine E-167

[29475-05-6]

As Ergobine, E-149 with

R = $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, R' = $-\text{CH}_2\text{CH}_3$

$\text{C}_{31}\text{H}_{39}\text{N}_5\text{O}_5$ 561.68

Alkaloid from *Claviceps purpurea*. Cryst. (Me_2CO aq.). Mp 198-199° (195-197° dec.). $[\alpha]_D^{20}$ -188 (c, 0.8 in CHCl_3).

► KE7050000

8-Epimer: Ergoptinine

[29475-04-5]

$\text{C}_{31}\text{H}_{39}\text{N}_5\text{O}_5$ 561.68


Alkaloid from ergot *Claviceps purpurea*.

Stütz, P. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1278 (synth, uv, ir, pmr)

Brunner, R. *et al.*, *Can. J. Chem.*, 1979, **57**, 1638 (isol, uv, ir, pmr)

Stuchlik, J. *et al.*, *esk. Farm.*, 1981, **30**, 201; *CA*, **95**, 146752e (Ergoptinine)

Ergosecalinine E-168



$\text{C}_{24}\text{H}_{28}\text{N}_4\text{O}_4$ 436.51

Tentative struct. Alkaloid from ergot (*Claviceps purpurea*). Prisms (EtOAc). Mp 217° dec. $[\alpha]_D^{18}$ +298 (c, 0.2 in CHCl_3). $[\alpha]_D^{18}$ +417 (c, 0.22 in Py). No CAS no 1967-2001 (8-14Cl).

Abe, M. *et al.*, *Bull. Agric. Chem. Soc. Jpn.*, 1959, **23**, 246-248; *CA*, **54**, 3482e (isol)

Abe, M. *et al.*, *Nippon Nogei Kagaku Kaishi*, 1960, **34**, 580-584; *CA*, **59**, 3098e (isol)

Ergosine† E-169

[561-94-4]

As Ergobine, E-149 with

R = $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, R' = $-\text{CH}_3$

$\text{C}_{30}\text{H}_{37}\text{N}_5\text{O}_5$ 547.653

Alkaloid from ergot (*Claviceps purpurea*) and from the seeds of *Ipomoea argyrophylla*. Also present in *Argyrea mollis*, *Argyrea luzonensis*, *Argyrea obtusifolia*, *Argyrea ridleyi* and *Argyrea splendens* (Convolvulaceae). Prisms (EtOAc). Mp 228° dec. (220° dec.). $[\alpha]_D^{20}$ -161 (c, 1 in CHCl_3). $[\alpha]_D^{20}$ +16 (c, 1 in Me_2CO). Forms mol. compd. with ergosinine, Mp 200° dec.

► KE7200000

Hydrochloride:

Plates + $1\text{Me}_2\text{CO}$. Mp 235° dec.

Hydrobromide:

Needles + $1\text{Me}_2\text{CO}$. Mp 230° dec.

9,10 α -Dihydro: Dihydroergosine

$\text{C}_{30}\text{H}_{39}\text{N}_5\text{O}_5$ 549.669

Alkaloid from the sclerotia of *Sphacelia sorghi*. Prisms (EtOAc). Mp 212° dec. $[\alpha]_D^{20}$ +10.1 (c, 1.5 in CHCl_3). $[\alpha]_D^{20}$ -52 (c, 0.5 in Py).

8-Epimer: Ergosinine. Ergoclavinine

[596-88-3]

$\text{C}_{30}\text{H}_{37}\text{N}_5\text{O}_5$ 547.653

Alkaloid from ergot (*Claviceps purpurea*) and the seeds of *Ipomoea argyrophylla*. Also present in *Argyrea mollis*, *Argyrea luzonensis*, *Argyrea obtusifolia*, *Argyrea ridleyi* and *Argyrea splendens* (Convolvulaceae). Prisms (EtOH, Me_2CO aq., C_6H_6 or EtOAc), needles + $\frac{1}{2}$ MeOH (MeOH). Mp 228° dec. $[\alpha]_D^{20}$ +420 (c, 1 in CHCl_3). $[\alpha]_D^{20}$ +380 (c, 1 in Me_2CO).

8-Epimer; hydrochloride: Mp 206° dec.

Smith, S. *et al.*, *J.C.S.*, 1937, 396 (isol, epimer)

Stoll, A. *et al.*, *Helv. Chim. Acta*, 1943, **26**, 2070 (Dihydroergosine)

Stadler, P.A. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1911 (synth, epimer)


Stauffer, D. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 1379 (isol, epimer)

Mantle, P.G. *et al.*, *Nature (London)*, 1968, **218**, 581 (Dihydroergosine)

Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435 (occur)

Ergosine† E-170

Tetrahydro-10b-hydroxy-5-(1-methylpropyl)-8H-oxazolo[3,2-a]pyrrolo[2,1-c]pyrazine-3,6(2H,5H)-dione, 9CI [897363-33-6]



Absolute Configuration

$\text{C}_{13}\text{H}_{20}\text{N}_2\text{O}_4$ 268.312

Alkaloid from the leaves of *Ipomoea fistulosa*.

Shah, V.H. *et al.*, *Chemistry (Rajkot, India)*, 2006, **3**, 8-10 (isol)

Ergostine E-171

[2854-38-8]

As Ergobine, E-149 with

R = $-\text{CH}_2\text{Ph}$, R' = $-\text{CH}_2\text{CH}_3$

$\text{C}_{34}\text{H}_{37}\text{N}_5\text{O}_5$ 595.697

Alkaloid from ergot (*Claviceps purpurea*). Prisms (EtOAc). Mp 204-208° dec. $[\alpha]_D^{20}$ -169 (c, 1.0 in CHCl_3).

8-Epimer: Ergostinine

[3268-95-9]


$\text{C}_{34}\text{H}_{37}\text{N}_5\text{O}_5$ 595.697

Alkaloid from ergot (*Claviceps purpurea*). Prisms (MeOH or EtOH). Mp 215-216° dec. $[\alpha]_D^{20}$ +357 (c, 1.0 in CHCl_3). $[\alpha]_D^{20}$ +429 (c, 1.0 in Py).

Schlienz, W. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1921 (isol, ir, struct, synth, epimer)

Ergotamine, BAN, INN E-172

12'-Hydroxy-2'-methyl-5'-(phenylmethyl)ergotaman-3',6',18-trione, 9CI [113-15-5]



Absolute Configuration

$\text{C}_{33}\text{H}_{35}\text{N}_5\text{O}_5$ 581.67


Alkaloid from ergot (*Claviceps purpurea*). Vasoconstrictor used in migraine treatment, powerful uterine contractant. Adrenergic antagonist, inhibitor of vasomotor centres, haemostatic used in obstetrics. Mp 213-214° dec. $[\alpha]_D$ -160 (CHCl_3). Log P 4.63 (uncertain value) (calc).

- LD₅₀ (mus, ipr) 62 mg/kg. KE7700000
Hydrochloride:
 Plates (EtOH aq.). Mp 212°.
 (+)-**Tartrate (2:1): Ergotamine tartrate, JAN, USAN.** Ergomar. Gynergen. Ergostate. Many other names [379-79-3]
 Mp 203° dec. [α]_D²¹ -135 (c, 0.4 in CHCl₃). Component of Cafergot.
- Adverse gastrointestinal and CNS effects when used therapeutically. Abuse can lead to addiction. Exp. reprod. and teratogenic effects. LD₅₀ (rat, ivn) 80 mg/kg. KE8225000
 N¹-Me: **Metergotamine, INN. 1-Methylergotamine, 8CI** [22336-84-1]
 C₃₄H₃₇N₅O₅ 595.697
 Vasoconstrictor. Migraine therapeutic. Cryst. (CH₂Cl₂/MeOH). Mp 167-170° dec. [α]_D²⁰ -163 (c, 0.5 in CHCl₃). Log P 5.1 (uncertain value) (calc).
- Dihydro: Dihydroergotamine, BAN, INN** [511-12-6]
 C₃₃H₃₇N₅O₅ 583.686
 Antiadrenergic agent, vasoconstrictor, antimigraine agent. Diminished oxytocic and vasoconstrictor effects cf. Ergotamine. Prisms (Me₂CO). Mp 239°. [α]_D²⁰ -64 (c, 0.5 in Py). Log P 4.94 (uncertain value) (calc).
- LD₅₀ (mus, scu) 80 mg/kg. II3675000
Dihydro; methanesulfonate: Dihydroergotamine mesylate, JAN, USAN. Migranal. Many other names [6190-39-2]
 Prisms (EtOH). Mp 230-235°.
- KE7920000
8-Hydroxy: 8-Hydroxyergotamine [72170-28-6]
 C₃₃H₃₅N₅O₆ 597.669
 Alkaloid from ergot (*Claviceps purpurea*). Cryst. (EtOAc or CHCl₃). Mp 197°. [α]_D²⁰ +14 (c, 1 in Py).
- 8-Epimer: Ergotaminine** [639-81-6]
 C₃₃H₃₅N₅O₅ 581.67
 Constit. of ergot alkaloids (*Claviceps purpurea*). Vasoconstrictor, uterine contractant. Plates (EtOH). Much less sol. than ergotamine. Mp 252° dec. (241-243°). [α]_D²⁰ +369 (c, 0.5 in CHCl₃). [α]_D²⁰ +397 (c, 0.5 in Py). Dec. at 241-243°.
- 2'-Epimer: Aciergotamine** [52949-35-6]
 C₃₃H₃₅N₅O₅ 581.67
 Minor ergot alkaloid occurring as artifact of epimerisation. Identified by hplc.
- 2',8-Diepimer: Aciergotaminine** [52919-20-7]
 C₃₃H₃₅N₅O₅ 581.67
 Minor ergot alkaloid occurring as artifact of epimerisation. Identified by hplc.
- [56644-29-2, 5989-77-5]
 Smith, S. et al., *J.C.S.*, 1930, 1390 (Ergotaminine)
 Stoll, A. et al., *Helv. Chim. Acta*, 1951, 34, 1544 (struct)
 Hofmann, A. et al., *Experientia*, 1961, 17, 206

- Hofmann, A. et al., *Helv. Chim. Acta*, 1963, 46, 2306 (synth, ir, uv, config)
 Ger. Pat., 1970, 2 017 560; *CA*, 74, 42535 (Metergotamine)
 Bassett, R.A. et al., *Biochem. J.*, 1973, 134, 1 (biosynth)
 Bach, N.J. et al., *J.O.C.*, 1974, 39, 1272 (cmr)
 Kreilgård, B. et al., *Anal. Profiles Drug Subst.*, 1977, 6, 113 (rev)
 Aellig, W.H. et al., *Int. J. Clin. Pharmacol. Biopharm.*, 1977, 15, 106 (Metergotamine)
 Kraljček, A. et al., *Coll. Czech. Chem. Comm.*, 1979, 44, 2255 (8-Hydroxyergotamine)
 Wurst, M. et al., *J. Chromatogr.*, 1979, 174, 401 (hplc)
 Scholten, A. et al., *J. Chromatogr.*, 1979, 176, 349 (Aciergotamine, Aciergotaminine)
 Pierri, L. et al., *J. Med. Chem.*, 1982, 25, 937 (conform)
 Kanto, J. et al., *Int. J. Clin. Pharmacol., Ther. Toxicol.*, 1983, 21, 135 (rev. pharmacol, metab)
 Plattner, R.D. et al., *J. Agric. Food Chem.*, 1983, 31, 785 (ms)
 Clark, B.J. et al., *Discoveries Pharmacol.*, 1984, 2, 3 (rev. pharmacol, tox)
 Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 7985; 7989 (synonyms)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 412; 414
 Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DLK800; EDC000; EDC500

Ergothioneine E-173

α -Carboxy-2,3-dihydro-N,N,N-trimethyl-2-thioxo-1H-imidazole-4-ethanaminium hydroxide inner salt, 9CI. 2-Mercaptohistidine trimethylbetaine. Thiolhistidine trimethylbetaine. Thioneine. Thiasine. Sympectothione [497-30-3]



C₉H₁₅N₃O₂S 229.302
 Present in blood, semen and various mammalian tissues, principally liver and kidneys. Also prod. by a variety of plants and microbes. Cryst. + 2H₂O. Mp 290° dec. [α]_D²⁰ +116.5 (H₂O).

- S-(2-Hydroxyethyl): Sclerothionine** [19833-41-1]
 C₁₁H₁₉N₃O₃S 273.355
 Isol. from *Sclerotinia libertiana*. Rhombic prisms (EtOH/Me₂CO). Mp 218-223° dec. [α]_D²⁴ +7.4 (c, 2 in H₂O).
- S-(2-Amino-2-carboxyethyl), S-oxide:** [53252-98-5]
 C₁₂H₂₀N₄O₅S 332.38
 Isol. from the fungi *Clitocybe acromelalga* and *Neurospora crassa*. Amorph. hygroscopic powder. Mp 188-190° dec. [α]_D²⁴ +74.7 (c, 0.5 in H₂O).
- Heath, H. et al., *J.C.S.*, 1951, 2215 (synth)
 Genghof, D.S. et al., *J. Bacteriol.*, 1964, 87, 852 (biosynth)

- Crossland, J. et al., *Nature (London)*, 1964, 203, 1388
 Matsuo, M. et al., *Agric. Biol. Chem.*, 1967, 31, 353; 1968, 32, 605; 611 (*Sclerothionine*)
 Ishikawa, Y. et al., *J. Biol. Chem.*, 1974, 249, 4420 (deriv)
 Sugihara, A. et al., *Acta Cryst. B*, 1976, 32, 181 (cryst struct)
 Motohashi, N. et al., *Chem. Pharm. Bull.*, 1976, 24, 1737 (nmr)
 Nicholson, J.R. et al., *Biochem. J.*, 1983, 211, 605 (pmr)

Ergovaline E-174

[2873-38-3]
 As Ergobine, E-149 with
 R = -CH(CH₃)₂, R' = -CH₃
 C₂₉H₃₅N₅O₅ 533.626
 Alkaloid from *Claviceps purpurea*. Cryst. (EtOAc). Mp 207-208° dec. [α]_D²⁰ -172 (c, 0.5 in CHCl₃).

Stadler, P.A. et al., *Helv. Chim. Acta*, 1964, 47, 1911 (synth, uv, ir)
 Brunner, R. et al., *Can. J. Chem.*, 1979, 57, 1638 (isol, uv, ir, pmr)

Ericamycin E-175

[11052-01-0]

Absolute Configuration

C₂₈H₂₁NO₈ 499.476
 Prod. by *Streptomyces varius*. Shows antibiotic activity against gram-positive bacteria. Deep red cryst. Mp 260-265°. λ_{\max} 249 (ε 39510); 323 (ε 15800); 345 (ε 13130); 364 (ε 12435); 487 (ε 14450) (MeOH).

Shimura, H. et al., *J. Antibiot., Ser. A*, 1966, 19, 51 (isol)
 U.S. Pat., 1974, 3 769 403; *CA*, 80, 13653s
 Kondo, S. et al., *J. Antibiot.*, 1998, 51, 232-234 (cd, uv, ir, pmr, cmr, struct)

Erichsonine E-176

[102673-54-1]

C₂₀H₂₆N₂O₃ 342.437
 Alkaloid from the stem bark of *Strychnos erichsonii* (Loganiaceae). Yellow prisms (MeOH). Mp 250° dec. [α]_D²⁰ -166 (c, 1.0 in MeOH).

Di-O-Ac:


Cryst. (EtOH). Mp 210°. $[\alpha]_D^{20}$ -115 (c, 0.4 in MeOH).

N,O,O -Tri-Ac: Mp 133-135°. $[\alpha]_D^{20}$ -70 (c, 0.5 in $CHCl_3$).

Forgacs, P. et al., *Phytochemistry*, 1986, **25**, 969 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Erinacean**E-177**

N-(7,8-Dihydro-8-oxo-H-purin-6-yl)- β -alanine, 9CI. N⁶- β -Alanil-8-oxoadenine [161212-38-0]




$C_8H_9N_5O_3$ 223.191

Isol. from the sponge *Isodictya erinacea*. Amorph. powder. λ_{max} 272 (ε 8444) (MeOH).

Moon, B. et al., *J. Nat. Prod.*, 1998, **61**, 116-118 (isol)

Erinacerin A**E-178**


[870193-54-7]



$C_{27}H_{31}NO_4$ 433.546

Related to Stachybotrin A in S-490. Alkaloid from the mushroom *Hericium erinaceum* (lions mane). Oil. Racemic. λ_{max} 215 (log ε 4.6); 244 (log ε 4.2); 292 (log ε 3.4) (MeOH).

Yaoita, Y. et al., *Chem. Pharm. Bull.*, 2005, **53**, 1202-1203 (isol, pmr, cmr)

Erioaustralasine**E-179**

4' α -form
Relative
configuration

$C_{22}H_{25}NO_5$ 383.443

4' α -form**trans-Erioaustralasine**

[152517-75-4]

Alkaloid from aerial parts of *Eriostemon australasius* ssp. *banksii* (Rutaceae). Gum. $[\alpha]_D$ +130 (c, 0.96 in $CHCl_3$).

Deacetoxy: Deacetoxy-trans-erioaustralasine

$C_{20}H_{23}NO_5$ 325.407

Alkaloid from *Halfordia kendack*. Needles (EtOAc). Mp 190-192°. $[\alpha]_D^{30}$ +244 (c, 0.24 in $CHCl_3$). λ_{max} 235 (log ε 3.84); 276 (log ε 3.68); 286 (log ε 3.68); 318 (log ε 3.65); 331 (log ε 3.55) (MeOH).

4' β -form**cis-Erioaustralasine**

[152406-29-6]


Alkaloid from aerial parts of *Eriostemon australasius* ssp. *banksii* (Rutaceae). Cubes (EtOAc). Mp 144-146°. $[\alpha]_D$ +104 (c, 1.1 in $CHCl_3$).

da Cunha, E.V.L. et al., *Aust. J. Chem.*, 1993, **46**, 1507 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Sultana, N. et al., *Phytochemistry*, 2001, **58**, 329-332 (Deacetoxyerioaustralasine)

Erioaustralasine hydrate**E-180**

[379689-36-8]



$C_{22}H_{27}NO_6$ 401.458

Alkaloid from *Halfordia kendack*. Amorph. λ_{max} 234 (log ε 4.17); 275 (log ε 3.98); 285 (log ε 3.99); 317 (log ε 3.9); 324 (log ε 3.77) (MeOH).

Deacetoxy: Deacetoxy-trans-erioaustralasine hydrate

[379689-34-6]

$C_{20}H_{25}NO_4$ 343.422

Minor alkaloid from *Halfordia kendack*. Amorph. $[\alpha]_D^{30}$ +125 (c, 0.03 in $CHCl_3$). Has N-Me replacing N- CH_2OAc . λ_{max} 234 (log ε 3.55); 275 (log ε 3.31); 286 (log ε 3.32); 317 (log ε 3.29); 332 (log ε 3.18) (MeOH).

1'-Epimer, deacetoxy: Deacetoxy-trans-1'-epierioaustralasine hydrate

[379689-37-9]


$C_{20}H_{25}NO_4$ 343.422

Minor alkaloid from *Halfordia kendack*. Amorph. λ_{max} 233 (log ε 3.93); 276 (log ε 3.45); 287 (log ε 3.46); 317 (log ε 3.44); 329 (log ε 3.34) (MeOH).

Sultana, N. et al., *Phytochemistry*, 2001, **58**, 329-332 (isol, pmr, cmr, ms)

Eripinal**E-181**

[69734-95-8]



$C_{23}H_{26}N_2O_5$ 410.469

Prob. Akuammiline-related. Alkaloid from the leaves of *Hunteria congolana* (Apocynaceae). Cryst. (Me₂CO). Mp 165°. $[\alpha]_D$ -154 (c, 0.5 in $CHCl_3$).

N^b-de-Me: Noveripinal

[69734-93-6]


$C_{22}H_{24}N_2O_5$ 396.442

Isol. from leaves of *Hunteria congolana* (Apocynaceae). Cryst. (Me₂CO). Mp 240-245°. $[\alpha]_D$ -98 (c, 0.96 in $CHCl_3$).

Vercauteren, J. et al., *Bull. Soc. Chim. Fr., Part II*, 1982, 291

Eripine**E-182**

Methyl 21,23-deepoxy-23-hydroxyerinin-21-oate, 9CI [20594-89-2]



Absolute
configuration

$C_{23}H_{28}N_2O_5$ 412.485

Prob. akuammiline-related. Alkaloid from *Hunteria umbellata* leaves (Apocynaceae). Needles (MeOH/EtOAc). Mp 186°. $[\alpha]_D^{23}$ -188 (c, 0.44 in $CHCl_3$).

O-Ac:


Needles (MeOH). Mp 76°.

Morita, Y. et al., *Helv. Chim. Acta*, 1968, **51**, 1438 (isol, uv, ir, pmr, ms, struct)

Heatley, F. et al., *J.C.S. Perkin 2*, 1981, 725 (struct)

Erucalexin**E-183**

2'-(Methylthio)spiro[2H-indole-2,5'-(4H)-thiazol]-3(1H)-one



$C_{12}H_{12}N_2O_2S_2$ 280.371

Related to Spirobrassinin, S-426.

(+)-form

Constit. of the leaves of *Erucastrum gallicum*. Phytoalexin. $[\alpha]_D$ +73 (c, 0.09 in MeOH).

(±)-form


Orange solid. Mp 94-97°. λ_{max} 234 (log ε 4.6); 262 (log ε 4.1); 368 (log ε 3.4) (MeCN).

Pedras, M.S.C. et al., *Chem. Comm.*, 2006, 1848-1850 (biosynth)

Pedras, M.S.C. et al., *Org. Biomol. Chem.*, 2006, **4**, 691-701 (isol, synth, pmr, cmr, ms)

Erucifoline

12,13-Epoxy-19-hydroxysenecianan-11,16-dione, 9CI. Alkaloid SC [40158-95-0]



C₁₈H₂₃NO₆ 349.383

Cyclic retronecine diester. Alkaloid from *Senecio erucifolius*, *Senecio aegypticus*, *Senecio erraticus*, *Senecio jacobaea* and *Senecio persoonii* (Asteraceae). Cryst. (EtOAc/EtOH). Mp 195-197°. [α]_D²⁵ -108 (c, 0.99 in CHCl₃).

N-Oxide: Erucifoline N-oxide

C₁₈H₂₃NO₇ 365.382

Alkaloid from whole plants of *Senecio persoonii* (Asteraceae). Glassy solid. [α]_D²⁰ -49 (c, 1 in EtOH).

Ac: O-Acetylerucifoline

[41310-60-5]

C₂₀H₂₅NO₇ 391.42

Constit. of leaves and inflorescences of *Senecio jacobaea*. Also present in inflorescences of *Senecio erucifolius* (Asteraceae). Needles. Mp 127-129°.

[40158-93-8, 67494-13-7]

Schröter, H.-B. *et al.*, *Coll. Czech. Chem. Comm.*, 1960, **25**, 472 (isol)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (cd)


Sedmera, P. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 4112 (isol, pmr, struct)

Witte, L. *et al.*, *Phytochemistry*, 1992, **31**, 559 (O-Acetylerucifoline)

Roeder, E. *et al.*, *Phytochemistry*, 1993, **32**, 1051 (oxide)

Ervafolene

[77784-39-5]



C₄₀H₄₄N₄O₃ 628.813

Aspidosperma-pseudoaspidosperma dimer. Alkaloid from the leaves of *Stenosolen heterophyllus* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. [α]_D²⁰ +236 (c, 1 in MeOH).

14'β,15'β-Epoxy: Ervafoline, 9CI

[70545-44-7]

C₄₀H₄₄N₄O₄ 644.812

Alkaloid from *Ervatamia pandacaqui* and the leaves of *Stenosolen heterophyllus* (preferred genus name *Taber-*

E-184

naemontana) (Apocynaceae). Cryst. (Me₂CO). Mp 258°. [α]_D²⁰ +279 (c, 1 in CHCl₃). λ_{\max} 252 (log ϵ 3.86); 306 (log ϵ 3.95); 326 (log ϵ 4.03) (EtOH).

19'ξ-Hydroxy: 19'-Hydroxyervafolene

[77794-87-7]

C₄₀H₄₄N₄O₄ 644.812

Alkaloid from the leaves of *Stenosolen heterophyllus* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 248°. [α]_D²⁰ +284 (c, 1 in CHCl₃).

19'ξ-Hydroxy, 14'β,15'β-epoxide: 19'-Hydroxyervafoline

[77784-40-8]


C₄₀H₄₄N₄O₅ 660.811

Alkaloid from the leaves of *Stenosolen heterophyllus* (Apocynaceae). Cryst. (MeOH). Mp 258°. [α]_D²⁰ +247 (c, 1 in CHCl₃). λ_{\max} 252 (log ϵ 3.86); 306 (log ϵ 3.95); 326 (log ϵ 4.03) (EtOH).

Henriques, A. *et al.*, *J.O.C.*, 1982, **47**, 803-811 (isol, pmr, ms, cryst struct)

Ervafolidene

[84716-74-5]



Absolute Configuration

C₄₀H₄₆N₄O₄ 646.828

Alkaloid from *Pandaca caducifolia* (Apocynaceae). Cryst. (Me₂CO). Mp 212°. [α]_D²⁰ +56 (c, 1.5 in MeOH).

14'β,15'β-Epoxy: Ervafolidine

[80293-76-1]

C₄₀H₄₆N₄O₅ 662.827

Alkaloid from the leaves of *Stenosolen heterophyllus* (preferred genus name *Tabernaemontana*) (Apocynaceae). Cryst. (Et₂O/hexane). Mp 240°. [α]_D²⁰ +20 (c, 0.05 in CHCl₃).

19'R-Hydroxy, 14'β,15'β-epoxide: 19'R-Hydroxyervafolidine

[80293-77-2]

C₄₀H₄₆N₄O₆ 678.827

Alkaloid from the leaves of *Stenosolen heterophyllus* (Apocynaceae). Cryst. (Me₂CO). [α]_D²⁰ +33 (c, 0.5 in Py).

3-Epimer: Epiervafolidene

[84774-04-9]

C₄₀H₄₆N₄O₄ 646.828

Alkaloid from *Pandaca caducifolia* (Apocynaceae). Amorph. [α]_D²⁰ +35 (c, 2 in MeOH).

3-Epimer, 14'β,15'β-epoxide: 3-Epiervafolidine

[80338-95-0]

C₄₀H₄₆N₄O₅ 662.827

Alkaloid from the leaves of *Stenosolen heterophyllus* (Apocynaceae). Cryst. (MeOH). Mp 260°. [α]_D²⁰ +52 (c, 0.5 in CHCl₃).

3-Epimer, 19'S-hydroxy, 14'β,15'β-epoxide: 19'S-Hydroxy-3-epiervafolidine

[80293-78-3]

C₄₀H₄₆N₄O₆ 678.827


Alkaloid from the leaves of *Stenosolen heterophyllus* (Apocynaceae). Cryst. (Me₂CO/Et₂O). Mp 260°. [α]_D²⁰ +74 (c, 1 in Py).

Zeches, M. *et al.*, *J. Nat. Prod.*, 1982, **45**, 707 (*Ervafolidene*, *Epiervafolidene*)

Henriques, A. *et al.*, *J.O.C.*, 1982, **47**, 803 (*Ervafolidene*, *3-Epiervafolidine*, *Hydroxyervafolidines*)

Ervahainamidine A**E-187**

[126398-90-1]




C₄₄H₅₄N₄O₅ 718.934

Minor alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae).

Feng, X.-Z. *et al.*, *J. Nat. Prod.*, 1989, **52**, 928 (isol, uv, pmr, ms, struct)

Ervahainamidine B**E-188**

[126456-02-8]




C₄₄H₅₄N₄O₅ 718.934

Minor alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae).

Feng, X.-Z. *et al.*, *J. Nat. Prod.*, 1989, **52**, 928 (isol, uv, ir, pmr, ms, struct)

Ervahanine A**E-189**

Methyl 13-(17-methoxy-17-oxovobasan-3α-yl)ibogamine-18-carboxylate, 9CI [80981-93-7]



Absolute Configuration

C₄₂H₅₀N₄O₄ 674.881

Alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae). Amorph. [α]_D²⁵ -97 (c, 0.86 in CHCl₃).

19,20-Dihydro: **19,20-Dihydroervatamine A**

[175478-70-3]

C₄₂H₅₂N₄O₄ 676.897

Alkaloid from the stems of *Ervatamia coronaria*.

3'-Hydroxy, N-de-Me: **3'-Hydroxy-N^d-demethylervatamine A**

[99257-58-6]

C₄₁H₄₈N₄O₅ 676.854

Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Mixt. of 3-epimers.

19'S-Hydroxy: **19S-Hydroxyervatamine A**

C₄₂H₅₀N₄O₅ 690.881

Alkaloid from the leaves and stem bark of *Tabernaemontana corymbosa*. Light yellow oil. [α]_D -105 (c, 0.16 in CHCl₃). λ_{\max} 228 (log ϵ 4.75); 287 (log ϵ 4.22); 294 (log ϵ 4.19) (EtOH).

10'-Methoxy, 3'-oxo: **Conodularine**

C₄₃H₅₀N₄O₆ 718.891

Alkaloid from the stem bark of *Tabernaemontana divaricata*. Light yellow oil. [α]_D +19 (c, 0.26 in CHCl₃). λ_{\max} 223 (log ϵ 4.77); 286 (log ϵ 4.31); 295 (log ϵ 4.3); 312 (log ϵ 4.01) (EtOH).

Feng, X.-Z. *et al.*, *J. Nat. Prod.*, 1981, **44**, 670 (*Ervatamine A*)

Perera, P. *et al.*, *Phytochemistry*, 1985, **24**, 2097 (*3'-Hydroxy-N^d-demethylervatamine A*)

Henriques, A.T. *et al.*, *J. Ethnopharmacol.*, 1996, **50**, 19-25 (*19,20-Dihydroervatamine A*)


Kam, T.-S. *et al.*, *Phytochemistry*, 2003, **63**, 625-629 (*Ervatamine*, *cmr*, *19S-Hydroxyervatamine A*)

Kam, T.S. *et al.*, *Heterocycles*, 2004, **63**, 845-850 (*Conodularine*)

Ervatamine

E-190

Methyl 4-ethyl-2,3,4,4a,5,6,7,12-octahydro-2-methyl-6-oxopyrido[3',4':4,5]cyclohept[1,2-b]indole-12a(1H)-carboxylate, 9CI [33257-13-5]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Ervatamia orientalis* and *Ervatamia lifuana* (Apocynaceae). Na channel blocker in nerve fibres, local anaesthetic blocker. Prisms + 1MeOH (MeOH). Mp 92-98°. [α]_D -3.7 (c, 2.1 in CHCl₃).

Picrate:

Yellow plates (Me₂CO). Mp 236-237° dec.

Methiodide:

Yellow plates (MeOH/Et₂O). Mp 217-220° dec.

19,20-Didehydro: **19,20-Didehydroervatamine**. 19,20-Dehydroervatamine [33228-82-9]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Ervatamia orientalis* (Apocynaceae). Prisms (Et₂O). Mp 198-200° dec. [α]_D +52 (c, 1.0 in CHCl₃).

19,20-Didehydro, N¹-methoxy: **19,20-Didehydro-N¹-methoxyervatamine**. N¹-Methoxy-19,20-dehydroervatamine [130263-09-1]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Ervatamia malaccensis* (Apocynaceae). [α]_D -14 (c, 0.5 in CHCl₃).

5-Oxo, 19,20-didehydro: **19,20-Didehydro-5-oxoervatamine**. 19,20-Dehydro-5-oxoervatamine [147802-44-6]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from leaves of *Tabernaemontana corymbosa* (Apocynaceae).

3-Deoxo, 3,14-didehydro: **Dehydroxyervataminol**

[923286-64-0]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Ervatamia divaricata*. Amorph. powder. [α]_D¹⁶ -61.5 (c, 0.28 in CHCl₃). λ_{\max} 241 (log ϵ 4.26); 315 (log ϵ 4.07) (MeOH).

6 α -Hydroxy, 19,20-didehydro: **19,20-Didehydro-6-hydroxyervatamine**. 19,20-Dehydro-6-hydroxyervatamine [923286-63-9]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Ervatamia divaricata*. Amorph. powder. [α]_D¹⁶ +167.3 (c, 0.15 in CHCl₃). λ_{\max} 235 (log ϵ 4.04); 312 (log ϵ 4.18) (MeOH).

20-Epimer: **20-Epiervatamine**

[33495-49-7]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Ervatamia orientalis* and *Ervatamia lifuana* (Apocynaceae). Prisms (Et₂O/petrol). Mp 185-187° dec. [α]_D -22 (c, 1.1 in CHCl₃).

Husson, A. *et al.*, *Tetrahedron*, 1973, **29**, 3095 (*synth, cryst struct*)

Riche, C. *et al.*, *Acta Cryst. B*, 1974, **30**, 610 (*cryst struct*)

Knox, J.R. *et al.*, *Aust. J. Chem.*, 1975, **28**, 1813; 1825 (*Ervatamine*, *20-Epiervatamine*, *Didehydroervatamine*, *isol, uv, ir, pmr, ms, struct*)

Clivio, P. *et al.*, *Phytochemistry*, 1990, **29**, 2693 (*19,20-Didehydro-N¹-methoxyervatamine*)


Kam, T.-S. *et al.*, *Phytochemistry*, 1993, **32**, 1357 (*19,20-Didehydro-5-oxoervatamine*)

Zhang, H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 54-59 (*19,20-Didehydro-6-hydroxyervatamine*, *Dehydroxyervataminol*)

Ervaticine

Conolidine

[100414-81-1]



Absolute Configuration

C₁₇H₁₈N₂O 266.342

Alkaloid from the leaves of *Ervatamia coronaria* and the stem bark of *Taber-*

naemontana divaricata. Pale yellow amorph. solid or light yellow oil. [α]_D +120 (CHCl₃). λ_{\max} 238 (log ϵ 4.15); 313 (log ϵ 4.22) (EtOH).

19R,20R-Epoxyde: **Conolobine A**

[748788-59-2]

C₁₇H₁₈N₂O₂ 282.341

Alkaloid from the stem bark of *Tabernaemontana divaricata*. Pale yellow oil. [α]_D +24 (c, 0.21 in CHCl₃). λ_{\max} 239 (log ϵ 4.12); 313 (log ϵ 4.26) (EtOH).

19S,20S-Epoxyde: **Conolobine B**

[748788-65-0]

C₁₇H₁₈N₂O₂ 282.341

Alkaloid from the stem bark of *Tabernaemontana divaricata*. Pale yellow oil. [α]_D +159 (c, 0.06 in CHCl₃). λ_{\max} 239 (log ϵ 4.11); 317 (log ϵ 4.14) (EtOH).


Atta-ur-Rahman, *et al.*, *Heterocycles*, 1985, **23**, 2975-2978 (*Ervatamine*)

Kam, T.-S. *et al.*, *Chem. Biodiversity*, 2004, **1**, 646-656 (*Conolidine*, *Conolobines A, B*)

Ervinidine

E-192

7-Ethyl-1,4,5,6,7,8,10,14b-octahydro-1,2-dioxo-2H-3,7-methanoazacycloundecino[5,4-b]indole-9-carboxylic acid methyl ester, 9CI [35989-12-9]



Relative Configuration

C₂₁H₂₄N₂O₄ 368.432


Alkaloid from *Vinca erecta* (Apocynaceae). Needles (MeOH). Mp 283-284° dec. [α]_D -17.3 (CHCl₃). λ_{\max} 232 (log ϵ 3.92); 305 (log ϵ 3.93); 342 (log ϵ 4.1) (EtOH).

Malikov, V.M. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 142; 1971, **7**, 640-643; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 119-120; 1971, **7**, 619-621 (*isol, struct*)

Ervitsine

E-193

4-Ethylidene-1,3,4,5,6,8-hexahydro-2-methyl-13-methylene-1,5-methanoazonino[4,3-b]indol-7(2H)-one, 9CI [65617-99-4]



Absolute Configuration

C₁₉H₂₀N₂O 292.38


Alkaloid from the root bark of *Pandaca boiteau* (Apocynaceae). Cryst. (MeOH). Mp 176-177°. [α]_D²⁰ -160 (c, 1 in CHCl₃).

Andriantsiferana, M. *et al.*, *Tet. Lett.*, 1977, 2587-2590 (*uv, pmr, cryst struct*)

Riche, C. *et al.*, *Acta Cryst. B*, 1979, **35**, 2738-2740 (*cryst struct*)

Bennasar, M.-L. *et al.*, *J.O.C.*, 1997, **62**, 3597-3609 (*synth*)

Erybidine **E-194**
6,7,8,9-Tetrahydro-2,11,12-trimethoxy-7-methyl-5H-dibenzo[*d,f*]azonin-3-ol, 9CI [34083-19-7]




$C_{20}H_{25}NO_4$ 343.422
Alkaloid from the leaves of *Erythrina* x. *bidwillii*, *Erythrina herbacea*, *Erythrina arborescens*, *Erythrina orientalis* and *Erythrina crista-galli* (Fabaceae). Needles (EtOH). Mp 178-180°.

Me ether: Mp 139-140°.

Ito, K. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 1509; 1974, **22**, 2108; 1976, **24**, 52 (*isol, ir, uv, pmr, ms, struct, synth*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1976, **41**, 3210 (*synth*)

Erylusamine TA **E-195**
[178063-82-6]



$R^1 = H, R^2 = Ac, R^3 = -(CH_2)_5NMe_2, n = 6$

$C_{54}H_{100}N_2O_{20}$ 1097.385
Glycolipid. *Isol.* from the marine sponge *Erylus* cf. *lendenfeldi* (tentative identification). Oil. $[\alpha]_D^{25} +2.8$ (c, 2.9 in MeOH).

Goobes, R. *et al.*, *Tetrahedron*, 1996, **52**, 7921-7928 (*isol, ir, pmr, cmr*)

Erylusidine **E-196**
[178063-84-8]

As Erylusamine TA, E-195 with $R^1 = -COCH_2CH(CH_3)_2, R^2 = H, R^3 = -(CH_2)_4NHC(NH_2)=NH, n = 7$

$C_{56}H_{104}N_4O_{20}$ 1153.452
Isol. from the marine sponge *Erylus* cf. *lendenfeldi* (tentative identification). Oil. $[\alpha]_D^{25} -4.1$ (c, 4.7 in MeOH).

Goobes, R. *et al.*, *Tetrahedron*, 1996, **52**, 7921-7928 (*isol, ir, pmr, cmr*)

Erylusine **E-197**

[178063-83-7]

As Erylusamine TA, E-195 with


$R^1 = H, R^2 = Ac, R^3 = -(CH_2)_3N(Me)(CH_2)_4NMe_2, n = 6$

$C_{57}H_{107}N_3O_{20}$ 1154.48

Glycolipid. *Isol.* from the marine sponge *Erylus* cf. *lendenfeldi* (tentative identification). Oil. $[\alpha]_D^{25} +1.9$ (c, 4.3 in MeOH).

Goobes, R. *et al.*, *Tetrahedron*, 1996, **52**, 7921-7928 (*isol, ir, pmr, cmr*)

Erymelanthine **E-198**
[88840-26-0]



$C_{18}H_{20}N_2O_3$ 312.368

Alkaloid from the seeds of *Erythrina melanacantha* (Fabaceae). Mp 160-161°. $[\alpha]_D^{22} +87$ (c, 0.11 in MeOH). First member of a new type of Erythrina alkaloid containing a 16-azaerythrinan skeleton.

8-Oxo: Melanacanthine. 8-Oxoerymelanthine

$C_{18}H_{18}N_2O_4$ 326.351

Alkaloid from *Erythrina melanacantha* (Fabaceae). Minimal structural proof.

De(methoxycarbonyl): Decarbomethoxyerymelanthine


$C_{16}H_{18}N_2O$ 254.331

Alkaloid from seeds of *Erythrina variegata* (Fabaceae). Tentative identification by gc-ms.

Dagne, E. *et al.*, *Tet. Lett.*, 1983, **24**, 5067 (*uv, ir, pmr, cmr, ms, struct*)

Jackson, A.H. *et al.*, *Chem. Biol. Isoquinoline Alkaloids*, (Eds. Phillipson, J.D. *et al.*), Springer, 1985, 62 (*Melanacanthine, Decarbomethoxyerymelanthine*)

Erylodienone **E-199**



$C_{18}H_{19}NO_4$ 313.352

(+)-form
Synthetic. $[\alpha]_D^{21} +25.1$ (c, 0.9 in $CHCl_3$).

(-)-3-Bromocamphor-9-sulphonate: Mp 197-198° dec. $[\alpha]_D^{20} -43.5$ (c, 1.4 in $CHCl_3$).

(-)-form [5531-67-9]
Alkaloid from *Erythrina lithosperma* and *Erythrina variegata* (Fabaceae). Intermediate in the biosynthesis of *Erythrina* alkaloids. $[\alpha]_D^{19} -27$ (c, 0.8 in $CHCl_3$) (for synthetic material). Opt. rotn. of natural

product (with presumed 5*S* chirality) not recorded.

(+)-3-Bromocamphor-9-sulphonate: Mp 203-204° dec. $[\alpha]_D^{20} +44.8$ (c, 0.98 in $CHCl_3$).

2-Alcohol, O¹⁶-Me: Erythritol.

Erythritol†

[50719-80-7]

$C_{19}H_{23}NO_4$ 329.395

Alkaloid from flowers of *Erythrina variegata* (Fabaceae). Brown semisolid.

Deoxy: Coccudienone

[77795-05-2]

$C_{18}H_{19}NO_3$ 297.353

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Amorph. powder.

(±)-form [13168-52-0]

Synthetic. Cryst. (EtOH). Mp 226-229° (224-227°).

Gervay, J.E. *et al.*, *Chem. Comm.*, 1966, 142 (*synth, uv, ir, pmr*)

Mondon, A. *et al.*, *Tet. Lett.*, 1966, 2557

(*synth, uv, ir, ms, pmr*)

Barton, D.H.R. *et al.*, *J.C.S. (C)*, 1968, 1529 (*synth, ir*)

Ghosal, S. *et al.*, *Aust. J. Chem.*, 1971, **24**, 2733

(*isol, ir, uv, ms, pmr*)

Ghosal, S. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 1274

(*isol, ir, uv, ms, pmr*)

Barton, D.H.R. *et al.*, *J.C.S. Perkin I*, 1974, 2278 (*resoln, config, biosynth*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1980, **36**,

3107 (*Coccudienone*)

Hewgill, F.R. *et al.*, *Aust. J. Chem.*, 1985, **38**,

537 (*synth*)

Chawla, H.M. *et al.*, *Fitoterapia*, 1993, **64**, 15;

383 (*Erythritol*)


Rigby, J.H. *et al.*, *Tet. Lett.*, 1999, **40**, 6887-

6890 (*synth, abs config, Erythritol*)

Erysodinophorine **E-200**

[72611-99-5]

[72379-63-6]



$C_{32}H_{38}N_3O_4^{\oplus}$ 528.67

Ester of Hypaphorine, H-764 with Erylodine in E-203. Alkaloid from the pods of *Erythrina arborescens* (Fabaceae). Brown syrupy liq. (as hydroxide).

O¹⁵-De-Me: Erysoinophorine

[73588-32-6]

[73606-99-2]

$C_{31}H_{36}N_3O_3^{\oplus}$ 514.643


Alkaloid from the pod walls of *Erythrina arborescens* (Fabaceae). Brown syrup (as hydroxide).

Tiwari, K.P. *et al.*, *Phytochemistry*, 1979, **18**, 704 (*isol, uv, ir, pmr, ms, struct*)

Eryosphorine

[54773-83-0]

[54615-80-4]

Absolute
configuration $C_{32}H_{38}N_3O_4^{\oplus}$ 528.67

Ester of Hypaphorine, H-764 and Erysovine in E-203. Alkaloid from the seeds of *Erythrina arborescens* (Fabaceae). Brown microcryst. solid (Me₂CO/MeOH)(as chloride). Mp 260° dec. (chloride).

O¹⁶-De-Me: Isoerysopinophorine

[74555-93-4]

[74555-94-5]

 $C_{31}H_{36}N_3O_4^{\oplus}$ 514.643

Alkaloid from the seeds of *Erythrina arborescens* (Fabaceae). Brown syrupy liq. (as hydroxide). Hydrol. to Erysovine and Hypaphorine.


Ghosal, S. *et al.*, *Phytochemistry*, 1974, **13**, 2603 (*isol, uv, pmr, ms, struct*)

Masood, M. *et al.*, *Phytochemistry*, 1980, **19**, 490 (*isol, uv, ir, pmr, ms, struct, Isoerysopinophorine*)

Erystramidine**E-202**

1,2,6,7-Tetrahydro-3,15,16-trimethoxyerythrinan-8-one, 9Cl. 8-Oxoerystrine
[52358-58-4]

[69111-35-9 ((±)-form)]

 $C_{19}H_{21}NO_4$ 327.379

Alkaloid from *Erythrina arborescens* (Fabaceae). $[\alpha]_D^{25} +148.5$ (c, 1.2 in CHCl₃) (synthetic).

O¹⁶-De-Me: 8-Oxoerysodine $C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Erythrina tahitensis* (Fabaceae). Minimal structural proof.

10,11-Didehydro: Erytharbine

[52358-59-5]

 $C_{19}H_{19}NO_4$ 325.363

Alkaloid from *Erythrina arborescens* (Fabaceae). Possibly an artifact.

15-Demethoxy: Isococcolinine

[66835-08-3]

 $C_{18}H_{19}NO_3$ 297.353

Alkaloid from flowers of *Erythrina variegata* (Fabaceae).

16-Demethoxy: Coccolinine. 8-Oxococcolinine. O-Methylcoccolinine

[60888-59-7]

[78463-71-5 ((±)-form)]

 $C_{18}H_{19}NO_3$ 297.353

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Cryst. (MeOH). Mp 174-175°. Poss. an

artifact. C-5 config. not firmly established.

16-Demethoxy, O¹⁵-de-Me: Coccoline. 8-Oxococcolinine

[60229-90-5]

 $C_{17}H_{17}NO_3$ 283.326

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Cryst. (EtOAc). Mp 245-246°. $[\alpha]_D +233$ (c, 1.08 in MeOH). Poss. an artifact. C-5 config. not firmly established.

15- or 16-Demethoxy: Erythromotidienone $C_{18}H_{19}NO_3$ 297.353

Alkaloid from *Erythrina variegata*. Needles (MeOH). Mp 172°. Posn. of aromatic methoxy group not determined. Presumably identical with either Coccolinine or Isococcolinine.

11β-Hydroxy: 11β-Hydroxyerystramidine $C_{19}H_{21}NO_5$ 343.379

Alkaloid from the flowers and pods of *Erythrina lysistemon*. Yellowish solid. $[\alpha]_D +100$ (c, 0.14 in MeOH). λ_{max} 219 (log ε 2.98); 257 (log ε 3.52); 322 (log ε 2.49) (MeOH).

11β-Methoxy: 11β-Methoxyerystramidine $C_{20}H_{23}NO_5$ 357.405

Alkaloid from the flowers of *Erythrina lysistemon*. Brownish solid. $[\alpha]_D +60$ (c, 0.11 in MeOH). λ_{max} 220 (log ε 2.96); 238 (log ε 3.28); 325 (log ε 1.58) (MeOH).

Ito, K. *et al.*, *Yakugaku Zasshi*, 1973, **93**, 1611; 1617; *CA*, **80**, 48212z; 68387p (*isol, struct*)

Pande, H. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 366 (*Coccolinine*)

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1976, **15**, 739 (*Coccoline, isol, uv, ir, pmr, ms, struct*)

Ito, K. *et al.*, *Chem. Comm.*, 1978, 733 (*synth*)

Ju-Ichi, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 563 (*Isococcolinine, synth*)

Chawla, A.S. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 39 (*cmr*)

Jackson, A.H. *et al.*, *Chem. Biol. Isoquinoline Alkaloids, Int. Symp., Phytochem. Soc. Eur., Abstr. Pap.*, 1985, 62 (*8-Oxoerysodine*)

Sano, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 94; 479 (*Erystramidine, Coccolinine, synth, ir, pmr*)

Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1365; 1993, **41**, 965; 2087 (*synth, Erystramidine, Erytharbine*)

Sharma, S.K. *et al.*, *Indian J. Heterocycl. Chem.*, 1992, **2**, 71; *CA*, **119**, 4929h (*Isococcolinine, synth*)

Sharma, S.K. *et al.*, *J. Indian Chem. Soc.*, 1998, **75**, 833-837 (*Erythromotidienone*)

Lee, H.I. *et al.*, *J.O.C.*, 2004, **69**, 8209-8218 (*Erystramidine, synth*)

Juma, B.F. *et al.*, *Phytochemistry*, 2004, **65**, 1397-1404 (*11-Hydroxyerystramidine, 11-Methoxyerystramidine*)


Blake, A.J. *et al.*, *Synthesis*, 2005, 3287-3292 (*Erystramidine, synth, ir, pmr, cmr, ms*)

Gao, S. *et al.*, *Org. Lett.*, 2006, **8**, 2373-2376 (*Erystramidine, synth*)

Erystrine

[27740-43-8]

[70681-68-4 ((-)-form)]

E-203 $C_{19}H_{23}NO_3$ 313.396

Alkaloid from a wide range of *Erythrina* spp. including *Erythrina abyssinica*, *Erythrina arborescens*, *Erythrina attilanensis*, *Erythrina blakei*, *Erythrina caffra*, *Erythrina coralloides*, *Erythrina crista-galli*, *Erythrina flabelliformis*, *Erythrina folkersii*, *Erythrina fusca* (gallito), *Erythrina goldmanii*, *Erythrina guatemalensis*, *Erythrina herbacea*, *Erythrina lithosperma*, *Erythrina livingstoniana*, *Erythrina macrophylla*, *Erythrina mulungu*, *Erythrina oliviae*, *Erythrina poeppigiana*, *Erythrina senegalensis*, *Erythrina steyermarkii*, *Erythrina suberosa*, *Erythrina tajumulcensis*, *Erythrina variegata* and *Erythrina zeheri*. Competitive neuromuscular blocking agent. *E. suberosa* shows antitumour activity and this is ascribed to Erystrine. Yellow cryst. (petrol).

Picrate: Mp 159-161°. $[\alpha]_D +142$ (c, 0.122 in EtOH).

N-Oxide: Erystrine N-oxide

[80153-99-7]

 $C_{19}H_{23}NO_4$ 329.395

Alkaloid from the flowers of *Erythrina mulungu* (Fabaceae). Oil. $[\alpha]_D^{25} +78.3$ (c, 1 in EtOH).

O³-De-Me: Erythravine. 1,2,6,7-Tetrahydro-15,16-dimethoxyerythrinan-3-ol

[19373-79-6]

 $C_{18}H_{21}NO_3$ 299.369

Alkaloid from the seeds of *Erythrina folkersii*. Also detected by glc-ms in the seeds of *Erythrina steyermarkii*, *Erythrina eggersii* and *Erythrina berteroa* (Fabaceae). Faintly yellow oil.

O¹⁵-De-Me: Erysovine

[466-72-8]

 $C_{18}H_{21}NO_3$ 299.369

Alkaloid from a wide range of *Erythrina* spp. (Fabaceae). Cryst. (EtOH). Mp 178-179.5° (167-169°, 175-176°). $[\alpha]_D +252$ (c, 0.123 in EtOH).

O¹⁵-De-Me, O-(carboxymethylsulfonyl): Erysothiovine $C_{20}H_{23}NO_7S$ 421.47

Alkaloid from the seeds of *Erythrina glauca*, *Erythrina pallida* and *Erythrina poeppigiana* (Fabaceae). Shows curarising paralytic activity in frogs. Cryst. + 2H₂O (H₂O). Mp 187°. $[\alpha]_D^{25} +208$ (c, 0.359 in EtOH). Hydrol. to Erysovine and sulfoacetic acid.

O¹⁶-De-Me: Erysovine. 1,2,6,7-Tetrahydro-3,15-dimethoxyerythrinan-16-ol

[7290-03-1]


 $C_{18}H_{21}NO_3$ 299.369

Alkaloid from a wide range of *Erythrina* spp., including *Erythrina abyssinica*, *Erythrina americana*, *Erythrina arborescens*, *Erythrina berter-*

- oana*, *Erythrina blakei*, *Erythrina chiriquensis*, *Erythrina costaricensis*, *Erythrina crista-galli*, *Erythrina cubensis*, *Erythrina decora*, *Erythrina excelsa*, *Erythrina falcata*, *Erythrina flabelliformis*, *Erythrina folkersii*, *Erythrina fusca* (gallito), *Erythrina herbacea*, *Erythrina lithosperma*, *Erythrina macrophylla*, *Erythrina melanacantha*, *Erythrina merrilana*, *Erythrina milbraedii*, *Erythrina oreophila*, *Erythrina pallida*, *Erythrina poeppigiana*, *Erythrina salviflora*, *Erythrina sandwicensis*, *Erythrina senegalensis*, *Erythrina stricta*, *Erythrina subumbrans*, *Erythrina variegata* and *Erythrina velutina* (Fabaceae). Cryst. (EtOH). Mp 204-205°. $[\alpha]_D^{27} +248$ (c, 0.311 in EtOH). $[\alpha]_D^{20} +239$ (c, 0.50 in CHCl_3). Erysocine was a complex of Erysodine with Erysovine.
- O¹⁶-De-Me, O- β -D-glucopyranoside: Glucoerysodine**
[509-49-9]
 $\text{C}_{24}\text{H}_{31}\text{NO}_8$ 461.511
Alkaloid from *Erythrina abyssinica*, *Erythrina senegalensis* and *Erythrina lysistemon* (Fabaceae). Mp 135-137°
Mp 166-168°. $[\alpha]_D +100.92$ (c, 1.56 in CHCl_3). $[\alpha]_D +127$ (H_2O).
- O¹⁶-De-Me, O- α -L-rhamnopyranoside: Rhamnoerysodine**
[133377-63-6]
 $\text{C}_{24}\text{H}_{31}\text{NO}_7$ 445.511
Alkaloid from *Erythrina lysistemon* (Fabaceae). Gum. $[\alpha]_D +83$ (c, 0.13 in CHCl_3).
- O³, O¹⁵-Di-de-Me: Erysoline. 3-Desmethylerysovine**
[52358-61-9]
 $\text{C}_{17}\text{H}_{19}\text{NO}_3$ 285.342
Alkaloid from the seeds of *Erythrina folkersii*. Also detected by glc-ms in the seeds of *Erythrina guatemalensis*, *Erythrina steyermarkii*, *Erythrina berteroa*, *Erythrina subumbrans*, *Erythrina lanata* and *Erythrina acanthocarpa* (Fabaceae). Broad Mp range.
- O³, O¹⁶-Di-de-Me: Erysonine. O³-Desmethylerysodine**
[7290-05-3]
 $\text{C}_{17}\text{H}_{19}\text{NO}_3$ 285.342
Alkaloid from the seeds of *Erythrina costaricensis* and *Erythrina lithosperma*, and from the trunk bark, root bark and seeds of *Erythrina variegata*. Also detected by glc-ms in the seeds of *Erythrina folkersii*, *Erythrina guatemalensis*, *Erythrina steyermarkii*, *Erythrina berteroa* and *Erythrina lanata* (Fabaceae). Cryst. (EtOH). Mp 241-243° dec. $[\alpha]_D^{25} +289$ (0.5% HCl aq.).
- O¹⁵, O¹⁶-Di-de-Me: Erysovine**
[545-68-6]
 $\text{C}_{17}\text{H}_{19}\text{NO}_3$ 285.342
Alkaloid from a wide variety of *Erythrina* spp. including *Erythrina abyssinica*, *Erythrina acanthocarpa*, *Erythrina arborensis*, *Erythrina berteroa*, *Erythrina chiriquensis*, *Erythrina costaricensis*, *Erythrina crista-galli*, *Erythrina cubensis*, *Erythrina decora*, *Erythrina dominguezii*, *Erythrina eggessii*, *Erythrina falcata*, *Erythrina flabelliformis*, *Erythrina folkersii*, *Erythrina fusca* (gallito), *Erythrina herbacea*, *Erythrina lithosperma*, *Erythrina macrophylla*, *Erythrina merrilana*, *Erythrina milbraedii*, *Erythrina oreophila*, *Erythrina pallida*, *Erythrina poeppigiana*, *Erythrina resupinata*, *Erythrina rubrinervia*, *Erythrina salviflora*, *Erythrina sandwicensis*, *Erythrina enegalensis*, *Erythrina subumbrans*, *Erythrina variegata*, and many others (Fabaceae). Cryst. (EtOH). Mp 241-242°. $[\alpha]_D^{25} +265.2$ (EtOH/glycerol). $[\alpha]_D^{25} +225$ (morpholine). $[\alpha]_D^{25} +276$ (HCl aq.).
- O¹⁵, O¹⁶-Di-de-Me, 15-O- β -D-glucopyranoside: 15-Glucoerysopine**
[273409-71-5]
 $\text{C}_{23}\text{H}_{29}\text{NO}_8$ 447.484
Alkaloid from the seeds of *Erythrina latissima*. Dark brown solid. Mp 150-152°. $[\alpha]_D +67.5$ (c, 0.006 in MeOH). λ_{max} 206 (log ϵ 4.38); 222 (log ϵ 4.4); 279 (log ϵ 3.74) (MeOH).
- O¹⁵, O¹⁶-Di-de-Me, 16-O- β -D-glucopyranoside: 16-Glucoerysopine**
[273409-70-4]
 $\text{C}_{23}\text{H}_{29}\text{NO}_8$ 447.484
Alkaloid from the seeds of *Erythrina latissima*. Dark brown solid. Mp 158-160°. $[\alpha]_D +76.5$ (c, 0.006 in MeOH). Authors' numbering scheme differs. λ_{max} 206 (log ϵ 4.38); 222 (log ϵ 4.4); 279 (log ϵ 3.74) (MeOH).
- O¹⁵, O¹⁶-Di-de-Me, O¹⁶-(carboxymethylsulfonfyl): Erysothiopine**
 $\text{C}_{19}\text{H}_{21}\text{NO}_7\text{S}$ 407.443
Alkaloid from the seeds of *Erythrina glauca* (gallito). Shows curarising paralytic activity in frogs. Cryst. (H_2O). Mp 168-169°. $[\alpha]_D^{25} +193$ (c, 0.183 in EtOH). Hydrol. to Erysovine and sulfoacetic acid.
- 10,11-Didehydro, O¹⁵-de-Me: 10,11-Dihydroerysodine**
 $\text{C}_{18}\text{H}_{19}\text{NO}_3$ 297.353
Alkaloid detected by glc-ms in the seeds of *Erythrina latissima*, *Erythrina stricta* and *Erythrina lysistemon* (Fabaceae). Possibly an artifact produced by an elimination reaction from the corresponding 11-methoxy or 11-hydroxy alkaloid.
- 10,11-Didehydro, O¹⁶-de-Me: 10,11-Dihydroerysodine**
 $\text{C}_{18}\text{H}_{19}\text{NO}_3$ 297.353
Alkaloid detected by glc-ms in the seeds of *Erythrina stricta*, *Erythrina lysistemon*, *Erythrina latissima* and *Erythrina caffra* (Fabaceae). Possibly an artifact produced by an elimination reaction from the corresponding 11-methoxy or 11-hydroxy alkaloid.
- 11-Hydroxy: see Erythartine, E-206**
- 8-Oxo: see Erysotramidine, E-202**
- 11-Oxo, 10 β -hydroxy: 10-Hydroxy-11-oxoerysotrine**
 $\text{C}_{19}\text{H}_{21}\text{NO}_5$ 343.379
Alkaloid from the flowers of *Erythrina herbacea*. Amorph. powder. $[\alpha]_D +150$ (c, 0.1 in MeOH). λ_{max} 204 (log ϵ 4.58); 242 (log ϵ 4.11); 283 (log ϵ 3.46) (MeOH).
- 10,11-Dioxo: 10,11-Dioxoerysotrine**
 $\text{C}_{19}\text{H}_{19}\text{NO}_5$ 341.363
Alkaloid from the seed pods of *Erythrina latissima*. Brown solid (Me_2CO). Mp 174-176°. $[\alpha]_D^{25} +167.5$ (c, 0.1 in MeOH). λ_{max} 206 (log ϵ 4.38); 247 (log ϵ 4.17); 292 (sh) (log ϵ 3.62); 351 (log ϵ 3.54) (MeOH).
- Folkers, K. *et al.*, *J.A.C.S.*, 1940, **62**, 1677-1683; 1941, **63**, 15444-1549; 1942, **64**, 1892-1896; 1944, **66**, 1083 (*Erysodine*, *Erysovine*, *Erysovine*, *Erysonine*, *Erysothiopine*, *Erysothiopine*, *isol*)
- Delofeu, V. *et al.*, *J.O.C.*, 1947, **12**, 486-489 (*Erysodine*, *isol*)
- Prelog, V. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 453-461; 1956, **39**, 498-504 (*Erysodine*, *Erysovine*, *Erysovine*, *struct*)
- Lapière, C. *et al.*, *J. Pharm. Belg.*, 1951, **6**, 71; *CA*, **45**, 9806c (*Glucoerysodine*)
- Weiss, U. *et al.*, *Experientia*, 1963, **19**, 108-112 (*Erysodine*, *Ersopine*, *ord*, *config*)
- Mondon, A. *et al.*, *Tet. Lett.*, 1966, 2557-2568 (*Erysonine*, *struct*)
- Barton, D.H.R. *et al.*, *J.C.S.(C)*, 1968, 1529-1537 (*struct*, *stereochem*, *biosynth*)
- Boar, R.B. *et al.*, *J.C.S.(B)*, 1970, 1591-1595 (*Erysodine*, *Erysovine*, *Erysovine*, *ms*)
- Miana, G.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 92 (*isol*, *uv*, *pmr*, *ms*)
- Barton, D.H.R. *et al.*, *J.C.S. Perkin 1*, 1973, 874-880 (*Erysodine*)
- Hargreaves, R.T. *et al.*, *J. Nat. Prod.*, 1974, **37**, 569-580 (*Erysodine*, *Erysovine*, *Erythravine*, *Erysovine*, *Erysonine*, *Erysoline*, *occur*)
- Games, D.E. *et al.*, *J. Nat. Prod.*, 1974, **37**, 581-588 (*Erysodine*, *Erysovine*, *Erythravine*, *Erysovine*, *Erysonine*, *occur*)
- Millington, D.S. *et al.*, *J.A.C.S.*, 1974, **96**, 1909-1917 (*Erythravine*, *Erysoline*, *Erysonine*, *occur*)
- Ito, K. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 52-55 (*isol*, *uv*, *pmr*, *ms*, *Erysodine*)
- Mondon, A. *et al.*, *Chem. Ber.*, 1979, **112**, 1329-1347 (*synth*, *uv*, *ir*, *bibl*)
- Sarragiotto, M.H. *et al.*, *Can. J. Chem.*, 1981, **59**, 2771-2775 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *oxide*)
- Jackson, A.H. *et al.*, *Allertonia*, 1982, **3**, 39; *CA*, **98**, 50334f (*Erysonine*, *Erysovine*, *Erysovine*, *occur*)
- Chawla, A.S. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 39-41 (*cmr*)
- Sano, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 479-500 ((\pm)-form, *synth*, *ir*, *pmr*)
- Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1365-1373; 1993, **41**, 2087-2095 (*synth*)
- Rigby, J.H. *et al.*, *J.A.C.S.*, 1991, **113**, 8975-8976 ((\pm)-form, *synth*)
- Wandji, J. *et al.*, *Phytochemistry*, 1995, **39**, 677-681 (*Erysodine*, *Glucoerysodine*)
- Toda, J. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 906-912 ((\pm)-form, *synth*)
- Wanjala, C.C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 871-873 (*Glucoerysopines*)
- Wanjala, C.C.W. *et al.*, *Planta Med.*, 2002, **68**, 640-642 (*10,11-Dioxoerysotrine*)
- Tanaka, H. *et al.*, *J. Nat. Med. (Tokyo)*, 2008, **62**, 228-231 (*10-Hydroxy-11-oxoerysotrine*)

Erythlaurine**E-204**

Methyl 1,6-didehydro-17-hydroxy-3,16-dimethoxyerythrinan-15-carboxylate, 9CI [77410-42-5]



C₂₀H₂₅NO₅ 359.421

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Oil. [α]_D +232 (EtOH).

Deoxy: Erythroculine

[22150-96-5]

C₂₀H₂₅NO₄ 343.422

Alkaloid from the leaves of *Cocculus laurifolius* and *Hyperbaena columbica* (Menispermaceae). Cryst. (Et₂O/pentane). Mp 79–82°. [α]_D²⁵ +216 (c, 1.05 in CHCl₃).

Deoxy, parent acid, amide: Erythramide

[77410-40-3]

C₁₉H₂₄N₂O₃ 328.41

Alkaloid from *Cocculus laurifolius* (Menispermaceae). Cubes. Mp 87–89°. [α]_D +262 (EtOH). Poss. an artifact arising from Erythroculine, from which it is obt. by NH₄OH treatment. However, it was shown that it was prob. not formed under the conditions used for isol.


Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1951 (*Erythroculine*)

Ju-ichi, M. *et al.*, *Heterocycles*, 1981, **16**, 555 (*Erythlaurine*, *Erythroculine*, *Erythramide*, *ir*, *pmr*, *ms*, *struct*)

Ripperger, H. *et al.*, *Phytochemistry*, 1983, **22**, 2603 (*Erythroculine*, *isol*, *uv*, *ir*, *pmr*, *ms*, *ord*)

Erythraline**E-205**

1,2,6,7-Tetrahydro-3-methoxy-15,16-[methylenebis(oxy)]erythrinan, 9CI



C₁₈H₁₉NO₃ 297.353

(+)-form [466-77-3]

Alkaloid from a wide range of *Erythrina* spp. including *Erythrina abyssinica*, *Erythrina arborescens*, *Erythrina brucei*, *Erythrina barqueroana*, *Erythrina burana*, *Erythrina caffra*, *Erythrina coralloides*, *Erythrina crista-galli*, *Erythrina folkersii*, *Erythrina excelsa*, *Erythrina fusca*, *Erythrina glauca*, *Erythrina globocalyx*, *Erythrina guatemalensis*, *Erythrina huehuetenangensis*, *Erythrina lanceolata*, *Erythrina latissima*, *Erythrina lithosperma*, *Erythrina lysistemon*, *Erythrina macrophylla*, *Erythrina milbraedii*, *Erythrina oliviae*, *Erythrina perrieri*, *Erythrina senegalensis*, *Erythrina sigmoidea*,

Erythrina stricta, *Erythrina steyermarkii*, *Erythrina tahitiensis*, *Erythrina tajumulcensis*, *Erythrina variegata*, *Erythrina velutina*, *Erythrina vespertilio* and *Erythrina zeheri* (Fabaceae). Mp 120° (106–107°). [α]_D^{21.5} +228 (c, 1.35 in EtOH).

Hydrobromide: Mp 243°. [α]_D²⁷ +216.6 (c, 0.500 in H₂O).

Hydroiodide: Mp 252° (245–245.5°). [α]_D +176 (c, 0.229 in H₂O).

O-De-Me: Erythrocarine

[98899-98-0]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the seeds of *Erythrina caribaea* (Fabaceae). Gum.

11α-Hydroxy: 11-Epierythrinine

[160604-69-3]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Erythrina caffra* (Fabaceae). Isol. as an insep. mixt. with Erythrinine.

11β-Hydroxy: Erythrinine†

[29306-29-4]

C₁₈H₁₉NO₄ 313.352

See also Erythrinine, E-211. Alkaloid from the leaves of *Erythrina indica* (now classified as *Erythrina variegata*), *Erythrina x bidwillii*, *Erythrina crista-galli* and *Erythrina crista-galli* cv Manuba Deiko, the trunk bark of *Erythrina lithosperma*, the flowers of *Erythrina brucei*, and from *Erythrina* spp. of Singapore. Also detected by glc-ms in the seeds of *Erythrina stricta*, *Erythrina caffra*, *Erythrina vespertilio*, *Erythrina burana* and *Erythrina perrieri*, and in the leaves of *Erythrina macrophylla* (Fabaceae). Mp 202–204°. [α]_D²⁵ +205 (c, 0.5 in CHCl₃). 11-Config. not certain.

11β-Methoxy: 11-Methoxyerythraline

[31686-06-3]

C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Erythrina lysistemon*. Also detected by glc-ms in the seeds of *Erythrina caffra* and *Erythrina vespertilio* (?) (Fabaceae). Pale yellow gum. [α]_D²⁵ +199 (c, 0.6 in CHCl₃). 11-Config. not certain.

11β-Methoxy, N-oxide: 11β-Methoxyerythraline N-oxide

C₁₉H₂₁NO₅ 343.379

Alkaloid from the flowers of *Erythrina crista-galli* (Fabaceae).

8-Oxo: 8-Oxoerythraline

[58779-40-1]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the leaves of *Erythrina crista-galli* and the seeds of *Erythrina brucei* (Fabaceae). Oil. [α]_D +135 (c, 0.1 in CHCl₃). λ_{max} 205 (log ε 4.57); 248 (log ε 4.12); 294 (sh) (log ε 3.66) (MeOH).

8-Oxo, 1α,2α-epoxide: 8-Oxoerythraline 1,2-epoxide

C₁₈H₁₇NO₅ 327.336

Alkaloid from *Erythrina x bidwillii*. Oil. [α]_D +94 (c, 0.1 in CHCl₃). λ_{max} 205 (log ε 4.59); 289 (log ε 3.78) (EtOH).

8-Oxo, demethoxy: Erythrosotidienone

C₁₇H₁₅NO₃ 281.31

Alkaloid from *Erythrina variegata*. Light brown semisolid.

8-Oxo, 11β-hydroxy: 8-Oxoerythrinine

[90686-27-4]

C₁₈H₁₇NO₅ 327.336

Alkaloid from the flowers of *Erythrina brucei* (Fabaceae). Pale-yellow oil. [α]_D +100 (c, 0.35 in CHCl₃).

8-Oxo, 11β-hydroxy, Ac: Mp 215–217° dec.

8-Oxo, 11β-methoxy: 8-Oxo-11β-methoxyerythraline

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Erythrina lysistemon* (Fabaceae). Minimal structural proof.

11-Oxo: 11-Oxoerythraline

C₁₈H₁₇NO₄ 311.337

Alkaloid detected by glc-ms in the seeds of *Erythrina zeheri* (Fabaceae).

11-Oxo, 10β-hydroxy: Erythbidine B

[214349-99-2]

C₁₈H₁₇NO₅ 327.336

Alkaloid from *Erythrina x bidwillii*. Amorph. solid. [α]_D +148 (c, 0.1 in MeOH). λ_{max} 205 (log ε 4.52); 243 (log ε 4.04); 291 (log ε 3.54) (no solvent reported).

10,11-Dioxo: 10,11-Dioxoerythraline

C₁₈H₁₅NO₅ 325.32

Alkaloid from *Erythrina x bidwillii*. Amorph. solid. [α]_D +254 (c, 0.1 in CHCl₃). λ_{max} 204 (log ε 4.32); 247 (log ε 4.17); 292 (sh) (log ε 3.62); 351 (log ε 3.53) (MeOH).

(±)-form [81750-78-9]

Synthetic. Oil.

Picrate:

Brown prisms (EtOH). Mp 196–199° dec.

8-Oxo: [81750-77-8]

Synthetic. Oil.

3-Epimer, 8-oxo:

Prisms (Et₂O/Me₂CO). Mp 161–165°.

Folkers, K. *et al.*, *J.A.C.S.*, 1940, **62**, 436; 1673 (*isol*, *uv*)

Deulofeu, V. *et al.*, *J.O.C.*, 1947, **12**, 486 (*isol*)

Prelog, V. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 453 (*isol*, *uv*)

Nowacki, W. *et al.*, *Z. Kristallogr.*, 1958, **110**, 89; *CA*, **52**, 15185c (*cryst struct*)

Weiss, U. *et al.*, *Experientia*, 1963, **19**, 108 (*ord*)

Barton, D.H.R. *et al.*, *J.C.S.(C)*, 1970, 1213 (*biosynth*)

Letcher, R.M. *et al.*, *J.C.S.(C)*, 1971, 652 (*11-Methoxyerythraline*)

Hargreaves, R.T. *et al.*, *J. Nat. Prod.*, 1974, **37**, 569 (*occur*)

Games, D.E. *et al.*, *J. Nat. Prod.*, 1974, **37**, 581 (*Erythrinine*, *11-Methoxyerythraline*, *11-Oxoerythraline*, *occur*)

Ito, K. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 52 (*Erythraline*, *Erythrinine*, *isol*, *uv*, *pmr*, *ms*)

Barakat, I. *et al.*, *J. Nat. Prod.*, 1977, **40**, 471 (*occur*)

El-Olemy, M.M. *et al.*, *J. Nat. Prod.*, 1978, **41**, 342 (*abs config*)

Jackson, A.H. *et al.*, *Allertonia*, 1982, **3**, 39; *CA*, **98**, 50334f (*occur*)


Chawla, A.S. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 39 (*cmr*)

Dagne, E. *et al.*, *Phytochemistry*, 1984, **23**, 449 (*Erythrinine*, *8-Oxoerythrinine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*, *synth*)

- Mantle, P.G. *et al.*, *Phytochemistry*, 1984, **23**, 1336 (8-Oxoerythraline)
- Jackson, A.H. *et al.*, *Chem. Biol. Isoquinoline Alkaloids*, (Eds. Phillipson, J.D. *et al.*), Springer, 1985, 62 (8-Oxo-11 β -methoxyerythraline)
- Chawla, A.S. *et al.*, *Phytochemistry*, 1985, **24**, 1821 (Erythrocarine)
- Sano, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 479 (*synth*, Erythraline, 8-Oxoerythraline)
- Chawla, A.S. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1146 (11 β -Methoxyerythraline N-oxide)
- Amer, M.E. *et al.*, *CA*, 1993, **122**, 101566g (11-Epierythrine)
- Sharma, S.K. *et al.*, *J. Indian Chem. Soc.*, 1998, **75**, 833-837 (Erythrosotidienone)
- Tanaka, H. *et al.*, *Phytochemistry*, 1998, **48**, 1461-1463 (Erythridine B)
- Tanaka, H. *et al.*, *Heterocycles*, 1999, **51**, 2759-2764 (10,11-Dioxoerythraline, 8-Oxoerythraline 1,2-epoxide)
- Shimizu, K. *et al.*, *Org. Lett.*, 2003, **5**, 2323-2325 (*synth*, erythrocarine)

Erythartine**E-206**

1,2,6,7-Tetrahydro-3,15,16-trimethoxyerythrinan-11-ol, 9CI. 11-Hydroxyerysotrine
[51666-26-3]



$C_{19}H_{23}NO_4$ 329.395

Alkaloid from the seeds of *Erythrina arborescens*, the leaves of *Erythrina herbacea*, the flowers of *Erythrina variegata* and *Erythrina mulungu*, and from *Erythrina* spp. of Singapore (Fabaceae). $[\alpha]_D^{25} +256$ (c, 1.4 in $CHCl_3$).

N-Oxide: **Erythartine N-oxide**

[80153-97-5]

$C_{19}H_{23}NO_5$ 345.394

Alkaloid from the flowers of *Erythrina lysistemon* and *Erythrina mulungu*. Oil. $[\alpha]_D^{25} +88.57$ (c, 1.4 in EtOH). λ_{max} 219 (log ϵ 2.9); 260 (log ϵ 3.61); 327 (log ϵ 2.31) (MeOH).

Ac: **Erythrascine. 11-Acetoxyerysotrine**

[39027-75-3]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from the seeds of *Erythrina arborescens* (Fabaceae). Cream-coloured needles (EtOH/Me₂CO). Mp 138-140°. $[\alpha]_D^{25} +152$ (c, 0.51 in $CHCl_3$). Doubtful structural assignment. Synthetic compd. was oily, $[\alpha]_D +66^\circ$, and the pmr spectrum differed from that of the alkaloid.

O¹⁵-De-Me: **11-Hydroxyerysotrine**

[54980-18-6]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid detected by glc-ms in the seeds of *Erythrina stricta*, *Erythrina senegalensis*, *Erythrina livingstoniana* and *Erythrina lysistemon* (Fabaceae).

O¹⁶-De-Me: **11-Hydroxyerysodine**

[54980-17-5]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid detected by glc-ms in the seeds of *Erythrina stricta*, *Erythrina lysistemon*, *Erythrina senegalensis*, *Erythrina livingstoniana* and *Erythrina latissima* (Fabaceae).

Me ether: **Erythristemine. 11-Methoxyerysotrine**

[28619-41-2]

$C_{20}H_{25}NO_4$ 343.422

Alkaloid from the leaves of *Erythrina lysistemon* and *Erythrina abyssinica* and from the seeds of *Erythrina arborescens* (Fabaceae). Pale yellow prisms (petrol). Mp 127-129°. $[\alpha]_D^{22} +189$ (c, 0.4 in $CHCl_3$).

Me ether, picrate: Mp 145-150°.

Me ether, N-oxide: **Erythristemine N-oxide**

[80153-98-6]

$C_{20}H_{25}NO_5$ 359.421

Alkaloid from flowers of *Erythrina x bidwillii* (Fabaceae). $[\alpha]_D^{25} -3.8$ (c, 0.01 in $CHCl_3$).

Me ether, O¹⁵-de-Me: **11-Methoxyerysotrine**

[54980-20-0]

$C_{19}H_{23}NO_4$ 329.395

Alkaloid detected by glc-ms in the seeds of *Erythrina abyssinica* and *Erythrina lysistemon* (Fabaceae).

Me ether, O¹⁵-de-Me, O- β -D-glucopyranoside: **11 β -Methoxyglucoerysotrine**

[133377-62-5]

$C_{25}H_{33}NO_9$ 491.537

Alkaloid from *Erythrina lysistemon* (Fabaceae). Gum.

Me ether, O¹⁶-de-Me: **11-Methoxyerysodine**

[54980-19-7]

$C_{19}H_{23}NO_4$ 329.395

Alkaloid detected by glc-ms in the seeds of *Erythrina lysistemon* (Fabaceae).

Me ether, O¹⁶-de-Me, O- β -D-glucopyranoside: **11 β -Methoxyglucoerysodine**

[133377-61-4]

$C_{25}H_{33}NO_9$ 491.537

Alkaloid from *Erythrina lysistemon* (Fabaceae). Gum. $[\alpha]_D +73$ (c, 0.42 in $CHCl_3$).

Me ether, O¹⁵, O¹⁶-di-de-Me: **11-Methoxyerysopine**

[54980-14-2]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid detected by glc-ms in the seeds of *Erythrina caffra* (Fabaceae).

11-Ketone, O¹⁵-de-Me: **11-Oxoerysotrine**

[54980-14-2]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid detected by glc-ms in the seeds of *Erythrina arborescens*, *Erythrina livingstoniana* and *Erythrina tahitensis* (Fabaceae).

11-Ketone, O¹⁶-de-Me: **11-Oxoerysodine**

[54980-13-1]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid detected by glc-ms in the seeds of *Erythrina arborescens*, *Erythrina livingstoniana*, *Erythrina abyssinica* and *Erythrina tahitensis* (Fabaceae).

11-Ketone, O¹⁵, O¹⁶-di-de-Me: **11-Oxoerysopine**

[54980-15-3]

$C_{17}H_{17}NO_4$ 299.326

Alkaloid detected by glc-ms in the seeds of *Erythrina arborescens* and *Erythrina tahitensis* (Fabaceae).

11-Epimer: **11-Epierythartine**

$C_{19}H_{23}NO_4$ 329.395

Alkaloid from the flowers of *Erythrina mulungu*. Amorph. powder. $[\alpha]_D^{25} +8.8$ (c, 1 in MeOH).

11-Epimer, N-oxide: **11-Epierythartine N-oxide**

$C_{19}H_{23}NO_5$ 345.394

Alkaloid from the flowers of *Erythrina lysistemon*. Yellowish oil. $[\alpha]_D +100$ (c, 0.04 in MeOH). λ_{max} 220 (log ϵ 2.82); 240 (log ϵ 3.32); 338 (log ϵ 2.42) (MeOH).

11-Epimer, O³-de-Me: **11 α -Hydroxyerythravine**

$C_{18}H_{21}NO_4$ 315.368

Alkaloid from the flowers of *Erythrina mulungu*. Amorph. powder. $[\alpha]_D^{25} +8.7$ (c, 1 in MeOH). λ_{max} 234 (log ϵ 4.3); 283 (log ϵ 3.45) ($CHCl_3$).

Ghosal, S. *et al.*, *Phytochemistry*, 1972, **11**,

2101-2103 (*Erythrascine*, *Erythristemine*)

Barton, D.H.R. *et al.*, *J.C.S. Perkin 1*, 1973,

874-880 (*Erythristemine*, *cryst struct*)

Ito, K. *et al.*, *Yakugaku Zasshi*, 1973, **93**, 1611-

1616; 1617-1621; 1975, **95**, 358-362; *CA*, **80**,

48212z; 68387p; **82**, 167515b (*isol, struct*)

Hough, E. *et al.*, *Acta Cryst. B*, 1976, **32**, 1154-

1162 (*Erythristemine*, *cryst struct, abs config*)

El-Olemy, M.M. *et al.*, *J. Nat. Prod.*, 1978, **41**,

342-347 (*isol, abs config*)

Ahmad, V.U. *et al.*, *J. Chem. Soc. Pak.*, 1979,

1, 79-80 (*isol*)

Sarragiotto, M.H. *et al.*, *Can. J. Chem.*, 1981,

59, 2771-2775 (*isol, uv, ir, pmr, cmr, ms*,

Erythartine N-oxide)

Amer, M.E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 161-

166 (*Erythrina lysistemon glycosides*)

Chawla, A.S. *et al.*, *Phytochemistry*, 1992, **31**,

372-374 (*Erythristemine N-oxide*)

Isobe, K. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**,

197 (*synth, Ac*)

Juma, B.F. *et al.*, *Phytochemistry*, 2004, **65**,


1397-1404 (*11-Epierythartine N-oxide*)

Flausino, O. *et al.*, *J. Nat. Prod.*, 2007, **70**, 48-

53 (*11-Epierythartine*, *11 α -Hydroxyerythravine*)

Erythratidine**E-207**

1,6-Didehydro-3,15,16-trimethoxyerythrinan-2-ol, 9CI
[41431-22-5]



$C_{19}H_{25}NO_4$ 331.411

Alkaloid from a wide range of *Erythrina* spp. including *Erythrina abyssinica*, *Erythrina arborescens*, *Erythrina barqueroana*, *Erythrina burana*, *Erythrina*

chiriquensis, *Erythrina coralloides*, *Erythrina decora*, *Erythrina eggersii*, *Erythrina excelsa*, *Erythrina falcata*, *Erythrina goldmanii*, *Erythrina guatemalensis*, *Erythrina latissima*, *Erythrina livingstoniana*, *Erythrina macrophylla*, *Erythrina melanacantha*, *Erythrina oliviae*, *Erythrina oreophila*, *Erythrina perrieri*, *Erythrina poeppigiana*, *Erythrina rubrinervia*, *Erythrina salviiflora*, *Erythrina senegalensis*, *Erythrina sigmoidea*, *Erythrina steyermarkii*, *Erythrina tahitensis* and *Erythrina variegata* (Fabaceae). Cryst. (EtOAc/petrol). Mp 120-120.5°. $[\alpha]_D^{25} +258$ (c, 0.581 in CHCl₃). $[\alpha]_D^{25} +273$ (c, 0.109 in EtOH).

Picrate:

Pale yellow plates (Me₂CO). Mp 222-224° (220-222°).

O¹⁵-De-Me: Erytosolvine

[52358-62-0]
C₁₈H₂₃NO₄ 317.384
Alkaloid from seeds of *Erythrina salviiflora*. Also detected in seeds of *Erythrina oliviae*, *Erythrina arborescens*, *Erythrina livingstoniana*, *Erythrina latissima*, *Erythrina tahitensis*, *Erythrina burana* and (possibly) *Erythrina melanacantha* (Fabaceae). Not obt. free of Erysotone. C-2 config. not detd., may not necessarily be as shown.

O¹⁶-De-Me: Erysotone

[26153-74-2]
C₁₈H₂₃NO₄ 317.384
Alkaloid from the seeds of *Erythrina salviiflora*. Also detected in seeds of *Erythrina eggersii*, *Erythrina decora*, *Erythrina oreophila*, *Erythrina variegata*, *Erythrina oliviae* and (possibly) *Erythrina melanacantha* (Fabaceae). Cryst. (CHCl₃/Et₂O). Mp 225-227°. C-2 config. not detd., may not correspond to that in Erysopitine.

O¹⁵, O¹⁶-Di-de-Me: Erysopitine

[38739-04-7]
C₁₇H₂₁NO₄ 303.357
Alkaloid from the trunk bark of *Erythrina variegata*. Mp 168-171°. $[\alpha]_D^{25} +148$ (c, 0.52 in EtOH). 2-Config. uncertain.

2-Ketone: Erythratidinone

[41758-74-1]
C₁₉H₂₃NO₄ 329.395
Alkaloid from the leaves of *Erythrina lithosperma*, also detected by glc-ms in leaves of *Erythrina poeppigiana* (Fabaceae). Cryst. (C₆H₆/petrol). Mp 119-120°. $[\alpha]_D^{25} +358$ (c, 1.121 in CHCl₃).

2-Ketone, picrate: Mp 205-207°.

2-Ketone, O¹⁵-de-Me: Erytosolvine

[52358-63-1]
C₁₈H₂₁NO₄ 315.368
Alkaloid detected in the seeds of *Erythrina salviiflora* (Fabaceae).

2-Ketone, O¹⁶-de-Me: Erysotone

[7236-40-0]
C₁₈H₂₁NO₄ 315.368
Alkaloid from the seeds of *Erythrina salviiflora* (Fabaceae). Needles (Et₂O). Mp 177-179°. $[\alpha]_D^{25} +342$ (c, 0.28 in EtOH).

2-Ketone, O¹⁵, O¹⁶-di-de-Me: Erysoflorinone

[52358-52-8]
C₁₇H₁₉NO₄ 301.341
Alkaloid detected by glc/ms in the seeds of *Erythrina salviiflora* and *Erythrina subumbrans* (Fabaceae).

11-ξ-Hydroxy: 11-Hydroxyerythratidine

[84209-90-5]
C₁₉H₂₅NO₅ 347.41
Alkaloid detected by glc-ms in the leaves of *Erythrina poeppigiana* (Fabaceae).

11-ξ-Hydroxy, O¹⁶-de-Me: 11-Hydroxyerysotone

C₁₈H₂₃NO₅ 333.383
Alkaloid from *Erythrina berteroa* (Fabaceae). Minimal structural proof. The isomeric struct., 11-Hydroxyerysotolvine, cannot be excluded.

11-ξ-Methoxy: 11-Methoxyerythratidine

[84214-98-2]
C₂₀H₂₇NO₅ 361.437
Alkaloid detected by glc-ms in the leaves of *Erythrina macrophylla* (Fabaceae).

11-ξ-Hydroxy, 2-ketone, O¹⁶-de-Me: 11-Hydroxyerysotinine

[84209-91-6]
C₁₈H₂₁NO₅ 331.368
Alkaloid detected by glc-ms in the leaves of *Erythrina macrophylla* (Fabaceae).

10,11-Dioxo, 2-ketone: 10,11-Dioxoerythratidinone

C₁₉H₁₉NO₆ 357.362
Alkaloid from the bark of *Erythrina subumbrans*. Pale yellow solid. $[\alpha]_D^{28} +16.3$ (c, 0.16 in MeOH). λ_{max} 244 (log ε 3.81); 288 (log ε 3.54); 352 (log ε 3.33) (MeOH).

2-Epimer: 2-Epierythratidine

[41431-23-6]
C₁₉H₂₅NO₄ 331.411
Alkaloid from the bark of *Erythrina variegata*. Cryst. (EtOAc/petrol). Mp 67-68°. $[\alpha]_D^{25} +142$ (c, 0.148 in CHCl₃).

2-Epimer, 11-ξ-hydroxy: 11-Hydroxyepierythratidine

[84209-89-2]
C₁₉H₂₅NO₅ 347.41
Alkaloid detected by glc-ms in the leaves of *Erythrina poeppigiana* (Fabaceae).

2-Epimer, 10,11-dioxo: 10,11-Dioxo-2-epierythratidine. 2-Epi-10,11-dioxoerythratidine

C₁₉H₂₁NO₆ 359.378
Alkaloid from the bark of *Erythrina subumbrans*. Pale yellow solid. $[\alpha]_D^{31} +92.8$ (c, 0.17 in MeOH). λ_{max} 246 (log ε 3.77); 291 (log ε 3.76); 352 (log ε 3.72) (MeOH).

Deulofeu, V. et al., *Chem. Ber.*, 1952, **85**, 620 (isol)

Mondon, A. et al., *Tet. Lett.*, 1966, 2557 (Erysotinine)

Ghosal, S. et al., *J. Pharm. Sci.*, 1972, **61**, 1274 (Erysopitine)

Barton, D.H.R. et al., *J.C.S. Perkin 1*, 1973, 874 (Erythratidine, Erythratidinone, 2-Epierythratidine, uv, ir, pmr, ms, struct, synth, abs config)

Hargreaves, R.T. et al., *J. Nat. Prod.*, 1974, **37**, 569 (occur)

Games, D.E. et al., *J. Nat. Prod.*, 1974, **37**, 581 (occur)

Millington, D.S. et al., *J.A.C.S.*, 1974, **96**, 1909 (Erythratidine, Erysotinine, Erysopitine, Erytosolvine, Erytosolvone, Erysotone, isol, uv, pmr, ms)

Barakat, I. et al., *J. Nat. Prod.*, 1977, **40**, 471 (occur)

Jackson, A.H. et al., *Allertonia*, 1982, **3**, 39; 47; *CA*, **98**, 14354d; 50334f (11-Hydroxyerythratidine, 11-Hydroxyepierythratidine, 11-Methoxyerythratidine, 11-Hydroxyerysotinine, Erytosolvone)

Chawla, A.S. et al., *J.C.S. Perkin 1*, 1982, 2903 (11-Hydroxyerysotone)

Chawla, A.S. et al., *Org. Magn. Reson.*, 1983, **21**, 39 (cmr, Erytosolvone)

Chawla, A.S. et al., *Planta Med.*, 1988, **54**, 526 (2-Epierythratidine)


Hosoi, S. et al., *Chem. Pharm. Bull.*, 1996, **44**, 2342 (synth)

Hosoi, S. et al., *J.C.S. Perkin 1*, 2000, 1505-1511 (synth)

Rukachaisirikul, T. et al., *J. Nat. Prod.*, 2008, **71**, 156-158 (Dioxoerythratidinone, Dioxoepierythratidine)

Erythratine**E-208**

1,6-Didehydro-3-methoxy-15,16-[methylenebis(oxy)]erythran-2-ol, 9CI
[5550-20-9]



C₁₈H₂₁NO₄ 315.368

Alkaloid from the seeds of *Erythrina glauca* (gallito), *Erythrina arborescens* and *Erythrina crista-galli*, the pods and trunk bark of *Erythrina lithosperma*, and the leaves of *Erythrina abyssinica*. Also detected by glc-ms in the seeds of *Erythrina subumbrans*. Cryst. + 0.5 H₂O (EtOH). Mp 170°. $[\alpha]_D^{28} +145.5$ (c, 0.371 in EtOH).

Hydrochloride:

Prisms (EtOH). Mp 250°. $[\alpha]_D^{25} +165.1$ (c, 0.333 in H₂O).

Hydrobromide: Mp 241°. $[\alpha]_D^{25} +158.7$ (c, 0.258 in H₂O).

2-Ketone: Erythratinone

[5639-06-5]
C₁₈H₁₉NO₄ 313.352
Alkaloid from *Erythrina crista-galli* and from the trunk bark of *Erythrina lithosperma* (Fabaceae). Mp 136-137°. $[\alpha]_D^{22.5} +409$ (c, 0.35 in EtOH).

11-Hydroxy: 11-Hydroxyerythratine

[84209-92-7]
C₁₈H₂₁NO₅ 331.368
Alkaloid detected by glc-ms in the leaves of *Erythrina macrophylla*. Also tentatively identified as a constit. of *Erythrina subumbrans* seeds (Fabaceae).

10,11-Dioxo: 10,11-Dioxoerythratine

C₁₈H₁₇NO₆ 343.335

Alkaloid from the bark of *Erythrina subumbrans*. Pale yellow solid. $[\alpha]_D^{25} +132.9$ (c, 0.17 in MeOH). λ_{\max} 219 (log ϵ 3.41); 249 (log ϵ 3.38); 289 (log ϵ 3.13); 352 (log ϵ 2.83) (MeOH).

2-Deoxy: Erythramine. Dihydroerythraline

[13268-79-6]
C₁₈H₂₁NO₃ 299.369

Alkaloid from the seeds of *Erythrina sandwicensis*, *Erythrina subumbrans*, *Erythrina crista-galli* and *Erythrina glauca*, and from the pods and trunk bark of *Erythrina lithosperma* (Fabaceae). Cryst. (Me₂CO). Mp 119.5-120.5° (103-104°). $[\alpha]_D^{25} +223$ (c, 1.64 in EtOH).

2-Deoxy, hydrobromide:

Needles (EtOH). Mp 228°. $[\alpha]_D^{26} +203.2$ (c, 0.500 in H₂O).

2-Deoxy, hydroiodide:

Yellow-orange needles (EtOH). Mp 249°. $[\alpha]_D^{25} +220$ (c, 0.500 in H₂O).

11-Methoxy: 11-Methoxyerythratine

C₁₉H₂₃NO₅ 345.394

Alkaloid from the flowers of *Erythrina crista-galli* (Fabaceae). Incorrect struct. shown in paper; subsequently amended but again an erroneous struct. is depicted.

2-Epimer: 2-Epierythratine

[5550-21-0]
C₁₈H₂₁NO₄ 315.368

Trace alkaloid detected by glc-ms in the seeds of *Erythrina subumbrans*. Mp 147-150°. $[\alpha]_D^{24} +280$ (c, 0.345 in EtOH). Physical data for this compd. refer to synthetic material.

2-Epimer, 11-hydroxy: 11-Hydroxy-epierythratine

C₁₈H₂₁NO₅ 331.368

Alkaloid tentatively identified by glc-ms as a constit. of *Erythrina subumbrans* seeds (Fabaceae).

Folkers, K. et al., *J.A.C.S.*, 1939, **61**, 1232; 3053; 1940, **62**, 436; 1673; 1942, **64**, 2146; 1951, **73**, 589 (*Erythratine, Erythramine, isol, uv, struct*)

Deulofeu, V. et al., *J.O.C.*, 1947, **12**, 486 (*Erythratine, Erythramine, isol*)

Prelog, V. et al., *Helv. Chim. Acta*, 1949, **32**, 453 (*Erythramine, synth, uv, struct*)

Barton, D.H.R. et al., *J.C.S. (C)*, 1968, 1529 (*Erythratine, Erythramine, 2-Epierythratine, Erythratinone, struct, stereochem, synth, biosynth*)

Boar, R.B. et al., *J.C.S. (B)*, 1970, 1591 (*ms, Erythratinone*)

Ghosh, D.K. et al., *Curr. Sci.*, 1972, **41**, 578 (*Erythramine, Erythratine, Erythratinone, isol, ms*)

Barton, D.H.R. et al., *J.C.S. Perkin 1*, 1973, 874 (*isol*)

Jackson, A.H. et al., *Allertonia*, 1982, **3**, 39; 47; *CA*, **98**, 14354d; 50334f (*occur, derivs*)

Chawla, A.S. et al., *J. Nat. Prod.*, 1987, **50**, 1146; 1988, **51**, 624 (*11-Methoxyerythratine*)

Rukachaisirikul, T. et al., *J. Nat. Prod.*, 2008, **71**, 156-158 (*10,11-Dioxoerythratine*)

Erythroidine

E-209

C₁₇H₂₂N₂O 270.374

Struct. unknown. Alkaloid from *Lycopodium erythraeum* (Lycopodiaceae).

MacLean, D.B. et al., *Alkaloids (Academic Press)*, 1985, **26**, 241

Erythreine

E-210

C₁₇H₂₂N₂O 270.374

Struct. unknown. Alkaloid from *Lycopodium erythraeum* (Lycopodiaceae).

Complex with Erythroidine, E-209: Isoerythreine

C₃₄H₄₄N₄O₂ 540.747

From *Lycopodium erythraeum*.

MacLean, D.B. et al., *Alkaloids (Academic Press)*, 1985, **26**, 241

Erythrinine[†]

E-211

C₃₀H₃₆N₄O₅ 532.638


Struct. unknown. Alkaloid from the seeds of *Erythrina lithosperma* (Fabaceae). Conts. 4 *O*-Me groups and one amide function. All *N* atoms are present in rings.

Tandon, S.P. et al., *Proc. Natl. Acad. Sci., India, Sect. A*, 1969, **39**, 263-264; *CA*, **73**, 77454r

Erythrocoamide A

E-212

[207226-71-9]



C₂₀H₂₅NO₄ 343.422


Alkaloid from *Dinosperma erythrocoeca*. Insecticidal agent. Needles (hexane/EtOAc). Mp 160-161°. $[\alpha]_D +133$ (c, 0.01 in CHCl₃). λ_{\max} 257; 310 (sh) (MeOH).

Latif, Z. et al., *J. Nat. Prod.*, 1998, **61**, 614-619 (*isol, uv, ir, pmr, cmr, ms*)

Erythrocoamide B

E-213

[207226-73-1]



C₁₆H₂₁NO₄ 291.346


Alkaloid from *Dinosperma erythrocoeca*. Pale yellow solid. Mp 95°. $[\alpha]_D +73$ (c, 0.001 in CHCl₃). λ_{\max} 256 (MeOH).

Latif, Z. et al., *J. Nat. Prod.*, 1998, **61**, 614-619 (*isol, uv, ir, pmr, cmr, ms*)

Erythrocoamide C

E-214

[207226-75-3]



C₂₂H₂₇NO₄ 369.46


Alkaloid from *Dinosperma erythrocoeca*. Pale yellow wax. Mp 42°. $[\alpha]_D +185$ (c, 0.001 in CHCl₃). λ_{\max} 255; 260; 270; 295 (MeOH).

Latif, Z. et al., *J. Nat. Prod.*, 1998, **61**, 614-619 (*isol, uv, ir, pmr, cmr, ms*)

α -Erythroidine, 9CI

E-215

1,2,6,7-Tetrahydro-12,17-dihydro-3-methoxy-16(15H)-oxaerythrinan-15-one, 11CI
[466-80-8]



Absolute configuration

C₁₆H₁₉NO₃ 273.331

Alkaloid from a range of *Erythrina* spp. including *Erythrina americana*, *Erythrina berteroaana*, *Erythrina chiapasana*, *Erythrina coralloides*, *Erythrina globocalyx*, *Erythrina poeppigiana*, *Erythrina salviflora* and *Erythrina standleyana* (Fabaceae). Needles (pentane). Mp 58-60°. $[\alpha]_D^{27} +136$ (c, 0.5 in H₂O). Unstable.

Hydrochloride: Mp 227-228° dec. $[\alpha]_D^{32} +118$ (c, 0.5 in H₂O).

Hydrobromide: Mp 220-222° dec.

Didehydro (?): Dehydro- α -erythroidine

C₁₆H₁₇NO₃ 271.315

Alkaloid detected by glc-ms in the leaves of *Erythrina poeppigiana*. Struct. not clear from publ. data.

8-Oxo: 8-Oxo- α -erythroidine

[84209-93-8]

C₁₆H₁₇NO₄ 287.315

Alkaloid from *Erythrina berteroaana* and *Erythrina poeppigiana*. Cryst. (Me₂CO). Mp 183°. $[\alpha]_D^{23} +164$ (c, 0.1 in MeOH).

8-Oxo, 1 α ,2 α -epoxide: 8-Oxo- α -erythroidine 1,2-epoxide

C₁₆H₁₇NO₅ 303.314

Alkaloid from the wood of *Erythrina poeppigiana*. Oil. $[\alpha]_D^{23} +211$ (c, 0.1 in MeOH). λ_{\max} 216 (log ϵ 4.23); 250 (sh) (log ϵ 3.69) (MeOH).

Δ^{12} -Isomer: β -Erythroidine. 12,13-Didehydro-13,14-dihydro- α -erythroidine, 10CI, 9CI

[466-81-9]

C₁₆H₁₉NO₃ 273.331

Alkaloid from a range of *Erythrina* spp. including *Erythrina americana*, *Erythrina arborescens*, *Erythrina berteroaana*, *Erythrina chiapasana*, *Erythrina coralloides*, *Erythrina globocalyx*, *Erythrina lithosperma*, *Erythrina poeppigiana*, *Erythrina salviflora* and *Erythrina standleyana* (Fabaceae). Neuromuscular and ganglionic blocking agent. Mp 99-100°. $[\alpha]_D^{28} +85$ (c, 0.3 in H₂O).

▶ LD₅₀ (mus, ipr) 24 mg/kg. KF3050000

Δ^{12} -Isomer, hydrochloride: Mp 228-231° dec. $[\alpha]_D^{27} +107$ (c, 0.5 in H₂O).

Δ^{12} -Isomer, hydrobromide: Mp 222.5° dec.

Δ^{12} -Isomer, 8-oxo: 8-Oxo- β -erythroidine
[85198-99-8]

C₁₆H₁₇NO₄ 287.315

Alkaloid from the leaves of *Erythrina berteroaana* (Fabaceae). Glass. $[\alpha]_D +32$ (c, 0.156 in EtOH).

Δ^{12} -Isomer, 1,2,6,7-tetrahydro, 1,6-dide-

hydro: 12,13-Didehydro-2,7,13,14-tetrahydro- α -erythroidine

[23255-54-1]

C₁₆H₂₁NO₃ 275.347

Nicotinic receptor antagonist. Mp 80-82°.

Δ^{12} -Isomer, 1,2,6,7-tetrahydro, 1,6-didehydro, hydrobromide: [29734-68-7]
Cryst. (EtOH). Mp 231-232° dec. [α]_D³⁰ +109 (c, 1.0 in H₂O).

Boekelheide, V. et al., *J.A.C.S.*, 1951, **73**, 2286-2289; 1953, **75**, 2550; 1958, **80**, 3905 (α -Erythroidine, β -Erythroidine, struct)

Weiss, U. et al., *Experientia*, 1963, **19**, 108; 660 (β -Erythroidine, ord)

Hanson, A.W. et al., *Proc. Chem. Soc., London*, 1963, 52 (β -Erythroidine, cryst struct)

Wenzinger, G.R. et al., *Proc. Chem. Soc., London*, 1963, 53 (abs config)

Leete, E. et al., *J.A.C.S.*, 1966, **88**, 4722 (biosynth)

Barton, D.H.R. et al., *J.C.S. Perkin 1*, 1973, 874; 1974, 346; 2278 (isol, biosynth)

Hargreaves, R.T. et al., *J. Nat. Prod.*, 1974, **37**, 569 (occur)

Aguilar, M.I. et al., *Phytochemistry*, 1981, **20**, 2061 (isol, uv, ir, pmr)

Jackson, A.H. et al., *Allertonia*, 1982, **3**, 39; *C.A.*, **98**, 50334f (occur, Dehydro- α -erythroidine)

Chawla, A.S. et al., *J.C.S. Perkin 1*, 1982, 2903 (8-Oxo- α -erythroidine, 8-Oxo- β -erythroidine)

Chawla, A.S. et al., *Org. Magn. Reson.*, 1983, **21**, 39 (cmr)

Damaj, M.I. et al., *Psychopharmacology (Berlin)*, 1995, **117**, 67-73 (Didehydrotetrahydroerythroidine)

Curzon, P. et al., *Brain Res.*, 1996, **714**, 185-191 (Didehydrotetrahydroerythroidine)

Tanaka, H. et al., *Planta Med.*, 2001, **67**, 871-873 (8-Oxo- α -erythroidine 1,2-epoxide)


Fukamoto, H. et al., *Angew. Chem., Int. Ed.*, 2006, **45**, 2731-2734 (β -Erythroidine, synth)

He, Y. et al., *Org. Lett.*, 2006, **8**, 3689-3692 (synth)

Erythropterin E-216

3-(2-Amino-4,5,6,8-tetrahydro-4,6-dioxo-7(3H)-pteridinylidene)-2-oxopropanoic acid

[7449-03-8]



C₉H₇N₅O₅ 265.185

Pigment of wing scales of pierid butterflies. Constit. of human tubercle bacilli, *Mycobacterium lacticola* and *Ephestia kühnellia*. Red cryst. + 1H₂O (dil. HCl). Violet-blue fluor. Forms Lepidopterin, L-109 with NH₃.

Sulfate salt:

Deep-red leaflets (H₂SO₄ aq.).

1',7-Dihydro, 2'-alcohol: 2-Amino-3,4,5,6,7,8-hexahydro- α -hydroxy-4,6-dioxo-7-pteridinepropanoic acid. **Ekap-terin**

[29067-93-4]

C₉H₁₁N₅O₅ 269.216

Constit. of *Ephestia kühnellia*. Yellow microcryst. + 1H₂O. [α]_D²⁰ -144. Fluorescent.

Schöpf, C. et al., *Annalen*, 1936, **524**, 49 (isol)
Pfleiderer, W. et al., *Chem. Ber.*, 1962, **95**, 2195 (struct)


Viscontini, M. et al., *Helv. Chim. Acta*, 1962, **45**, 2479; 1963, **46**, 56 (Ekapterin)

Viscontini, M. et al., *Helv. Chim. Acta*, 1963, **46**, 51 (synth)

Pfleiderer, W. et al., *J. Het. Chem.*, 1992, **29**, 583 (rev)

Erythroskyrin E-217

[4987-27-3]



C₂₆H₃₃NO₆ 455.55

Mycotoxin prod. by *Penicillium islandicum*. Mycotoxin. Orange-red cryst. (EtOH). Sol. MeOH, Py, bases, AcOH; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 130-133°. [α]_D +46.9 (CHCl₃). λ_{\max} 260 (ϵ 13800); 392 (ϵ 60300) (EtOH/NaOH) (Derep). λ_{\max} 260 (ϵ 8910); 409 (ϵ 28200) (EtOH) (Derep).

► LD₅₀ (mus, ipr) 60 mg/kg. UY7600000
Howard, B.H. et al., *Biochem. J.*, 1954, **57**, 212 (isol)

Shoji, J. et al., *Chem. Pharm. Bull.*, 1965, **13**, 1240 (isol, uv, ir, pmr, struct)

Shibata, S. et al., *Chem. Pharm. Bull.*, 1966, **14**, 474 (biosynth)

Ueno, Y. et al., *Jpn. J. Exp. Med.*, 1975, **45**, 525 (tox)


Beutler, J.A. et al., *J. Nat. Prod.*, 1988, **51**, 562 (abs config, pmr, cmr)

Dixon, D.J. et al., *J.C.S. Perkin 1*, 1999, 839-841 (synth)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 688

Erythrosuamine E-218

[22149-20-8]



C₂₅H₃₉NO₆ 449.586

Alkaloid from the bark of *Erythrophleum guineense* (Fabaceae). Noncryst.

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 139-141°.

[α]_D -67 (c, 0.9 in EtOH).

3 β -Acetoxy, N-de-Me: 3-Acetoxynererythroamine

[58189-26-7]

C₂₆H₃₉NO₈ 493.596

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae).

Antineoplastic agent. Log P 2.86

(calc). λ_{\max} 223 (ϵ 11000) (EtOH)

(Derep).

3 β -Acetoxy, N-de-Me; hydrochloride: Mp 173-175°.


3 β -Acetoxy, N-de-Me; N,O-Di-Ac: Mp 201°.

Thorell, A. et al., *Acta Chem. Scand.*, 1968, **22**, 2835 (isol, ir, pmr, struct)

Loder, J.W. et al., *Tet. Lett.*, 1975, 2497 (deriv)

Erythrosuavine E-219

[874460-56-7]




C₂₇H₃₉NO₇ 489.608

Alkaloid from the stem bark of *Erythrophleum suaveolens*. Cryst. (MeOH). Mp 173-174°.

Manfouo, R.N. et al., *Bull. Chem. Soc. Ethiop.*, 2005, **19**, 69-74 (isol, pmr, cmr)

Eschscholtzidine E-220

5,6,11,12-Tetrahydro-8,9-dimethoxy-14-methylbenzo[5,6]cycloocta[1,2-f]-1,3-benzodioxol-5,11-imine, 9CI. 2,3-Dimethoxy-8,9-methylenedioxy-pavinane. O-Methylcaryachine



(+)-form

C₂₀H₂₁NO₄ 339.39

(+)-form

Alkaloid from the bark of *Cryptocarya chinensis* (Lauraceae). Cryst. (H₂O) (as hydrochloride). Mp 178-180° (hydrochloride).

N-Oxide: **Eschscholtzidine N-oxide**

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Cryptocarya chinensis*. Needles (MeOH). Mp 193-194°. [α]_D +145.3 (c, 0.13 in MeOH). λ_{\max} 225 (sh) (log ϵ 3.7); 290 (log ϵ 3.2) (MeOH).

O³-De-Me, N-oxide: (+)-**Isocaryachine N-oxide**

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Cryptocarya chinensis*. Needles (Me₂CO). [α]_D +72.6 (c, 0.07 in MeOH). Mp >280°. λ_{\max} 219 (log ϵ 3.7); 280 (log ϵ 3.2) (MeOH).

(-)-form [6451-67-8]

Alkaloid from *Eschscholtzia californica* and from the fruit and tops of *Thalictrum revolutum* (Papaveraceae, Ranunculaceae). Shows weak antibacterial activity. Amorph. [α]_D²⁴ -194.2 (c, 1.56 in MeOH). λ_{\max} 291 (ϵ 12000) (EtOH) (Berdy).

N-Me: **Eschscholtzidine N-methosalt**

[74111-40-3]

[74111-41-4]
 $C_{21}H_{24}NO_4^{\oplus}$ 354.425
 Quaternary alkaloid from the fruit of *Thalictrum revolutum* (Ranunculaceae). Amorph. solid (as chloride). $[\alpha]_D^{20}$ -170 (c, 0.26 in MeOH) (chloride).

O²-De-Me: 2-Hydroxy-3-methoxy-8,9-methylenedioxy-pavinane. Caryachine
 [37687-27-7]

$C_{19}H_{19}NO_4$ 325.363
 Alkaloid from the leaves of *Cryptocarya chinensis* (Lauraceae). Cryst. (Me₂CO). Mp 174°. $[\alpha]_D^{21}$ -269.6 (c, 1.00 in EtOH).

O²-De-Me, hydrobromide: Mp 295-296° dec.

O²-De-Me, N-oxide: Caryachine N-oxide

$C_{19}H_{19}NO_5$ 341.363
 Alkaloid from *Cryptocarya chinensis*. Needles (Me₂CO). $[\alpha]_D$ -86.9 (c, 0.1 in MeOH). Mp >280°. λ_{max} 223 (sh) (log ϵ 3.7); 284 (log ϵ 3.2) (MeOH).

O²-De-Me, N-Me: Caryachine N-methosalt. N-Methylcaryachinium

[70494-78-9]
 [70518-44-4]

$C_{20}H_{22}NO_4^{\oplus}$ 340.398
 Quaternary alkaloid from the bark of *Cryptocarya chinensis* (Lauraceae). Cryst. (MeOH) (as iodide). Mp 174-175° (iodide). $[\alpha]_D^{24}$ -160 (c, 0.12 in MeOH).

O³-De-Me: Isocaryachine

[65634-33-5]
 $C_{19}H_{19}NO_4$ 325.363

Alkaloid from *Cryptocarya chinensis*.

O³-De-Me, N-oxide: Isocaryachine N-oxide

$C_{19}H_{19}NO_5$ 341.363
 Alkaloid from *Cryptocarya chinensis*. Needles (Me₂CO). $[\alpha]_D$ -245.1 (c, 0.11 in MeOH). Mp >280°. λ_{max} 223 (log ϵ 3.7); 293 (log ϵ 3.2) (MeOH).

O³-De-Me, N-oxide, stereoisomer: Isocaryachine N-oxide B

$C_{19}H_{19}NO_5$ 341.363
 Alkaloid from *Cryptocarya chinensis*. Needles (Me₂CO). $[\alpha]_D$ -26.2 (c, 0.22 in MeOH). Mp >280°. λ_{max} 225 (log ϵ 3.6); 288 (log ϵ 3.3) (MeOH).

12-Hydroxy: 12-Hydroxyeschscholtzidine

$C_{20}H_{21}NO_5$ 355.39
 Alkaloid from *Cryptocarya chinensis*. Needles (MeOH). Mp 201-202°. $[\alpha]_D$ -171.5 (c, 0.13 in MeOH). λ_{max} 225 (sh) (log ϵ 3.72); 288 (log ϵ 3.94) (MeOH).

(±)-form [52437-08-8]

Synthetic. Noncryst.

N-Me:

Cryst. (MeOH) (as iodide). Mp 305° (iodide).

O²-De-Me: (±)-Caryachine

[37686-77-4]
 $C_{19}H_{19}NO_4$ 325.363
 Alkaloid from the leaves of *Cryptocarya chinensis* (Lauraceae). Cryst. (EtOH). Mp 241-242°.

Manske, R.H.F. et al., *Can. J. Chem.*, 1966, **44**, 1259 (uv, struct)

Lu, S.-T. et al., *Yakugaku Zasshi*, 1966, **86**, 177; 296 (Caryachine, Eschscholtzidine, isol, uv, pmr, struct)

Tomita, M. et al., *Yakugaku Zasshi*, 1966, **86**, 414; *CA*, **65**, 5500e (ms)

Chan, R.P.K. et al., *Tetrahedron*, 1967, **23**, 4209 (uv, ord, cd)

Natarajan, S. et al., *Indian J. Chem.*, 1972, **10**, 451; 1974, **12**, 550 (uv, pmr, ms, struct, synth)

Premila, M.S. et al., *Indian J. Chem.*, 1973, **11**, 1084 (synth, uv, pmr)

Wu, J. et al., *J. Nat. Prod.*, 1977, **40**, 294 (isol, pmr)

Dyke, S.F. et al., *Tetrahedron*, 1978, **34**, 241 (synth, uv, pmr, cd, ms)

Chen, C.-H. et al., *J. Nat. Prod.*, 1979, **42**, 163 (Caryachine N-methosalt)

Wu, J. et al., *J. Nat. Prod.*, 1980, **43**, 270 (isol, uv, cd, pmr, struct, deriv)

Chang, W.T. et al., *Phytochemistry*, 1998, **48**, 119-124 (Caryachine N-methosalt)


Lin, F.W. et al., *Chem. Pharm. Bull.*, 2001, **49**, 1292-1294; 2002, **50**, 157-159 (Caryachine oxides, Isocaryachine oxides)

Wu, T.-S. et al., *J. Nat. Prod.*, 2001, **64**, 1404-1407 (N-oxide, 12-hydroxy)

Eschscholtzine

E-221

5,6,12,13-Tetrahydro-15-methylcycloocta[1,2-f:5,6-f']bis[1,3]benzodioxol-5,12-imine, 9CI, 2,3:8,9-Bis(methylenedioxy)-pavinane. Californine. Crychine



(-)-form

$C_{19}H_{17}NO_4$ 323.348

Shows twofold rotational symmetry.

(-)-form [4040-75-9]

Alkaloid from *Eschscholtzia californica*, *Eschscholtzia douglasii*, *Eschscholtzia glauca* and *Cryptocarya chinensis* (Papaveraceae, Lauraceae). Cryst. (MeOH). Mp 128°. $[\alpha]_D^{15}$ -220.2 (c, 1.00 in EtOH) (202). $[\alpha]_D^{19}$ -240 (c, 0.50 in MeOH).

Hydrochloride:

Needles (MeOH/CHCl₃). Mp 244-246°.

Picrate: Mp 177°.

N-Oxide: Eschscholtzine N-oxide

[106449-14-3]

$C_{19}H_{17}NO_5$ 339.347

Alkaloid from the petals of *Eschscholtzia californica* (Papaveraceae). Amorph. $[\alpha]_D^{22}$ -213.75 (c, 0.4 in CHCl₃).

N-Me: Californidine. Eschscholtzine N-methosalt

[18830-99-4]

$C_{20}H_{20}NO_4^{\oplus}$ 338.382

Quaternary alkaloid from *Eschscholtzia californica*, *Eschscholtzia douglasii*, *Eschscholtzia glauca* and *Eschscholtzia oregana* (Papaveraceae).

N-Me, iodide: [17939-30-9]

Cryst. (MeOH). Mp 285-286°. $[\alpha]_D^{18}$ -212 (c, 0.5 in MeOH).

N-Me, perchlorate: [17939-31-0]

Cryst. (MeOH or H₂O). Mp 327-329°. $[\alpha]_D^{23}$ -219 (c, 0.10 in MeOH).

N-De-Me: N-Demethyleschscholtzine. N-Demethylcrychine

$C_{18}H_{15}NO_4$ 309.321

Alkaloid from *Cryptocarya chinensis*. Yellow powder (Me₂CO). $[\alpha]_D$ -74.3 (c, 0.02 in MeOH). λ_{max} 221 (sh) (log ϵ 3.75); 294 (log ϵ 3.92) (MeOH).

12-Hydroxy: 12-Hydroxyeschscholtzine.

12-Hydroxyerychine

Alkaloid from *Cryptocarya chinensis*. Needles (MeOH). Mp 173-174°. $[\alpha]_D$ -143 (c, 0.28 in MeOH). λ_{max} 230 (log ϵ 3.77); 294 (log ϵ 3.91) (MeOH).

5-Oxo: Eschscholtzinone

[119060-95-6]

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D$ -122 (c, 0.18 in MeOH). The first ketonic pavinoid.

(±)-form [65634-34-6]

Synthetic. Mp 223-225° (as picrolonate).

Gertig, H. et al., *Acta Pol. Pharm.*, 1965, **22**,

443; *CA*, **64**, 11547c (isol, uv, ir, deriv)

Manske, R.H.F. et al., *Can. J. Chem.*, 1965, **43**,

2180; 2183 (isol, uv, pmr, ms, struct)

Lu, S.-T. et al., *Yakugaku Zasshi*, 1966, **86**,

177; *CA*, **64**, 17653g (isol, uv, ir, pmr)

Tomita, M. et al., *Yakugaku Zasshi*, 1966, **86**,

414; *CA*, **65**, 5500e (ms)

Slavk, J. et al., *Coll. Czech. Chem. Comm.*,

1967, **32**, 4420; 1975, **40**, 1095 (isol, uv,

struct, deriv)

Barker, A.C. et al., *J.C.S. (C)*, 1967, 1317 (abs

config, synth)

Chan, R.P.K. et al., *Tetrahedron*, 1967, **23**,

4209 (uv, ord, cd)

Urzuá, A. et al., *J. Nat. Prod.*, 1986, **49**, 922

(isol, uv, pmr, ms, cd, struct, oxide)

Gözler, B. et al., *J. Nat. Prod.*, 1988, **51**, 760

(Eschscholtzinone)

Chang, W.-T. et al., *Phytochemistry*, 1998, **48**,

119-124 (Eschscholtzinone)

Marek, R. et al., *Magn. Reson. Chem.*, 1999,

37, 195-202 (N-15 nmr)


Wu, T.-S. et al., *J. Nat. Prod.*, 2001, **64**, 1404-

1407 (12-Hydroxyeschscholtzine, N-

Demethyleschscholtzine)

Esmeraldine A

E-222



R = mixt. of $C_{13}H_{27}$, -
 $(CH_2)_{10}CH(CH_3)CH_2CH_3$, -
 $(CH_2)_nCH(CH_3)_2$ ($n = 12, 13, 14$), $C_{15}H_{31}$,
 $C_{16}H_{33}$, $C_{17}H_{33}$, $C_{17}H_{31}$ plus other un-
 identified components. Mixt. of diaster-
 eoisomers. Isol. from *Streptomyces*
antibioticus strain Tü 2706. Dark-green
 amorph. powder. Mp 121-155° dec. λ_{max}
 201 (ϵ 43700); 257 (ϵ 51300); 345 (ϵ
 11300); 400 (sh); 642 (ϵ 4730) (EtOH/
 HCl). λ_{max} 257 (ϵ 59000); 360 (ϵ 14200);
 400 (sh); 591 (ϵ 7280) (EtOH/NaOH).
 λ_{max} 200 (ϵ 46200); 256 (ϵ 57500); 355 (ϵ
 12700); 400 (sh); 608 (ϵ 3640) (EtOH).

Di-Me ester:

Blue-green powder (CHCl₃/hexane).
Mp 65-73°.

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 2058 (*isol, uv, ir, pmr, struct*)
Van't Land, C.W. *et al.*, *J.O.C.*, 1993, **58**, 6576 (*biosynth*)

Emeraldine B**E-223**

4-[1-[(2-Hydroxy-6-methylbenzoyl)oxy]ethyl]-6-methyl-6H,10H-pyrido[3,2-a:5,6,1-d',e']diphenazine-11,17-dicarboxylic acid, 9CI
[119958-58-6]

As Emeraldine A, E-222 with
R = 2-Hydroxy-6-methylphenyl

C₃₈H₂₈N₄O₇ 652.662

Phenazine antibiotic. *Isol.* from *Streptomyces antibioticus* strain Tü 2706. Dark-green amorph. powder. Sol. MeOH, CHCl₃. Mp 300°. Mixt. of diastereoisomers. λ_{max} 206 (ε 61300); 258 (ε 59100); 293 (ε 9410); 342 (ε 13600); 400 (sh) (ε); 658 (ε 7120) (EtOH/HCl). λ_{max} 216 (sh) (ε); 258 (ε 59100); 306 (ε 13800); 350 (ε 17100); 400 (sh) (ε); 588 (ε 9400) (EtOH/NaOH). λ_{max} 206 (ε 61300); 256 (ε 65400); 301 (ε 9320); 352 (ε 14240); 400 (sh) (ε 10000); 689 (ε 4320) (EtOH).

Di-Me ester: [119936-22-0]

Fine blue-green cryst. (Me₂CO/hexane). Mp 176-180°.

N,O-Di-Ac:


Red powder. Mp 300°.

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 2058 (*isol, uv, ir, pmr, cmr, ms, struct*)

Van't Land, C.W. *et al.*, *J.O.C.*, 1993, **58**, 6576 (*biosynth*)

Estatin A**E-224**

[106455-06-5]



C₁₈H₂₅N₅O₅ 391.426

Peptide-type antibiotic. *Isol.* from *Myceliophthora thermophila*. Specifically inhibits thiol proteases and IgE antibodies. Needles + 1H₂O. Sol. H₂O, DMSO, Py, AcOH; poorly sol. CHCl₃, EtOAc, C₆H₆, hexane. Mp 223-225° dec. [α]_D²⁴ +41.8 (c, 0.6 in H₂O). λ_{max} 247 (ε 160); 252 (ε 180); 259 (ε 199); 264 (ε 160); 269 (ε 86) (H₂O) (Derep).

▶ RR0392000

p-Hydroxybenzyl analogue: Estatin B

[106396-24-1]

C₁₈H₂₅N₅O₆ 407.425

From *Myceliophthora thermophila*. Similar properties to Estatin A. Needles + 1H₂O. Sol. H₂O, Py, AcOH, DMSO; poorly sol. CHCl₃, EtOAc, hexane, C₆H₆. Mp 217-218° dec. [α]_D²⁴ +46.8 (c, 0.2 in 0.1M HCl). λ_{max} 222 (ε 11200); 276 (ε 1560); 283 (ε 1290) (H₂O) (Derep).

▶ RR0391000

Eur. Pat., 1986, 196 189; *CA*, **106**, 67676 (*isol, synth*)

Yaginuma, S. *et al.*, *J. Antibiot.*, 1989, **42**, 1362 (*isol, struct, props*)

1,2-Ethanediamine, 9CI**E-225**

Ethylenediamine, 8CI. 1,2-Diaminoethane [107-15-3]

H₂NCH₂CH₂NH₂

C₂H₈N₂ 60.099

Solvent, emulsifier, rubber latex stabiliser, antifreeze component. Base with synthetic applications. Used in manuf. of chelating agents and fungicides. Li deriv. is alkene isomerisation and dehydrogenating agent. Pharmaceutical aid. Chelating agent; reagent for Cu(I).

Comonomer in polyamides and polyurethanes. Clear, thick, strongly alkaline liq. with ammoniacal odour. Sol. EtOH, H₂O (with hydration); insol. C₆H₆; sl. sol. Et₂O. d₂₀²⁰ 0.9. Mp 11°. Bp 118°. n_D²⁰ 1.4565. pK_a 6.85. Steam-volatile. Forms comds. with metallic salts. Vp 10.7 mmHg (20°). Readily absorbs CO₂ from the air.

▶ Flammable, fl. p. 34°, autoignition temp. 385°. Reacts violently with many materials. Eye, skin and respiratory tract irritant. Conc. solns. corrosive. An allergen and sensitiser. LD₅₀ (rat, orl) 500 mg/kg. Exp. reprod. effects. ACGIH TLV: long-term 10 ppm (Sk). KH8575000

Monohydrate: [6780-13-8]

Oil. Mp 10°. Bp 118°.

Hydrochloride (1:2): [333-18-6]

Monoclinic prisms. Insol. EtOH; sol.

H₂O. Subl. without melting.

▶ LD₅₀ (rat, orl) 500 mg/kg. KV3850000

Sulfate (1:1): [22029-36-3]

Cryst. (EtOH aq.).

Dipicrate: [1593-00-6]

Mp 233-235° dec.

N-Ac: N-(2-Aminoethyl)acetamide, 9CI.

2-Acetamidoethylamine

[1001-53-2]

C₄H₁₀N₂O 102.136

Needles. V. sol. H₂O, EtOH; spar. sol.

Et₂O. Mp 172°. Bp₁₃ 148°. Bp₅ 125-130°.

N,N'-Di-Ac: N,N'-1,2-Ethanediybisacetamide, 9CI

[871-78-3]

C₆H₁₂N₂O₂ 144.173

Mp 51°. Bp₃ 128°.

N,N,N',N'-Tetra-Ac: Tetraacetylene-

ethylenediamine. N,N'-1,2-Ethanediybis[N-

acetylacetamide], 9CI. N,N'-Ethylene-

bis(diacetamide), 8CI

[10543-57-4]

C₁₀H₁₆N₂O₄ 228.247

Comly. important bleach activator. Mp

149-150°.

N-Dodecanoyl: N-(2-Aminoethyl)dode-

canamide, 9CI

[10138-02-0]

C₁₄H₃₀N₂O 242.404

Cryst. (Et₂O). Mp 51-52°.

N,N'-Didodecanoyl: N,N'-1,2-Ethane-

diybisdodecanamide, 9CI

[7003-56-7]

C₂₆H₅₂N₂O₂ 424.709

Cryst. (EtOH). Mp 162-164° (160°).

N,N'-Didodecanoyl, picrate:

Cryst. Mp 61.5°.

N,N'-Dioctadecanoyl: N,N'-Dioctadeca-

noylethanediamine. N,N'-1,2-Ethane-

diybis(octadecanamide), 9CI. N,N'-

Ethylenebis(octadecanamide). N,N'-

Distearoylethylenediamine. Acrawax C.

Acrawax CT. Chemetron 100. Wax C

[110-30-5]

C₃₈H₇₆N₂O₂ 593.031

Antifoaming agent used in polymer

industry. Cryst. (EtOH aq.). Mp 149-

149.5° (140-141°).

N,N'-Di-9Z-octadecenoyl: N,N'-Dioleoy-

lethylenediamine

[110-31-6]

C₃₈H₇₂N₂O₂ 588.999

Cryst. (MeOH). Mp 114-115°.

N-tert-Butyloxycarbonyl: tert-Butyl N-

(2-aminoethyl)carbamate. N-Mono(-

tert-butyloxycarbonyl)ethylenediamine

[57260-73-8]

C₇H₁₆N₂O₂ 160.216

Oil. Bp_{0.3} 79-80°. n_D²⁰ 1.4505.

N-Benzoyl: N-(2-Aminoethyl)benzamide

[1009-17-2]

C₉H₁₂N₂O 164.207

Bp₂ 183-185°.

N,N'-Dibenzoyl: N,N'-1,2-Ethanediybis-

benzamide

[644-33-7]

C₁₆H₁₆N₂O₂ 268.315

Cryst. (EtOH). Mp 246-247°.

N,N'-Bis(4-methylbenzenesulfonyl):

[4403-78-5]

C₁₆H₂₀N₂O₄S₂ 368.477

Cryst. (EtOH). Mp 163-164°.

▶ XT5748000

N-(4-Methoxy-E-cinnamoyl): N-(4-

Methoxycinnamoyl)ethylenediamine

[208465-40-1]

C₁₂H₁₆N₂O₂ 220.271

Alkaloid from *Nicotiana glauca*.

N-Me: N-Methyl-1,2-ethanediamine. 2-

(Methylamino)ethylamine

[109-81-9]

C₃H₁₀N₂ 74.125

Bp 115-116°.

▶ KV5250500

N-Me; hydrochloride: [64670-85-5]

Cryst. + H₂O (EtOH). Mp 130-132°

dec.

N,N,N'-Tri-Me: [142-25-6]

C₃H₁₄N₂ 102.179

Bp 116-118°. n_D²⁰ 1.4190.

N-Et: N-Ethyl-1,2-ethanediamine. 2-

Ethylaminoethylamine

[110-72-5]

C₄H₁₂N₂ 88.152

Refractive oil. V. sol. H₂O. Bp 130°.

N-Et; hydrochloride (1:2): [75776-31-7]

Mp 173°.

N-Et, N'-Me: N-Ethyl-N'-methyl-1,2-

ethanediamine, 9CI. N-Ethyl-N'-

methylthylenediamine, 9CI


[111-37-5]

C₃H₁₄N₂ 102.179

- Liq. which dec. in air. Misc. H₂O, Et₂O. Bp 130-133° (135°).
- N-Et, N'-Me; hydrochloride (1:2):*
Cryst. (EtOH). Mp 217-218°.
- N-Et, N'-Me, dipicrate:* [22509-07-5]
Cryst. (as monohydrate). Mp 175-177°.
- N,N-Di-Et: N,N-Diethyl-1,2-ethanediamine. 2-Diethylaminoethylamine*
[100-36-7]
C₆H₁₆N₂ 116.206
Bp 145°. pK_{a1} 9.88; pK_{a2} 6.82 (H₂O).
- Skin and severe eye irritant. LD₅₀ (rat, orl) 2830 mg/kg. KV3500000
- N,N'-Di-Et: N,N'-Diethyl-1,2-ethanediamine. 1,2-Bis(ethylamino)ethane*
[111-74-0]
C₆H₁₆N₂ 116.206
d 1.43. Bp 151-152°. n_D²⁰ 1.4326.
- N,N'-Di-Et; hydrochloride (1:2):* [52198-62-6]
Leaflets (EtOH). Mp 259-260° dec.
- N,N-Di-Et, N'-Me:* [104-79-0]
C₇H₁₈N₂ 130.233
Bp 157-160°.
- N,N-Di-Et, N',N'-di-Me:* [123-10-4]
C₈H₂₀N₂ 144.259
Mp 207-208° (as hydrobromide). Bp 156-157°.
- N,N'-Di-Et, N,N'-di-Me:* [106-66-1]
C₈H₂₀N₂ 144.259
Comly. available. No phys. props. reported.
- N,N,N'-Tri-Et:* [105-04-4]
C₈H₂₀N₂ 144.259
Bp₁₃ 54-55°.
- N,N,N',N'-Tetra-Et: N,N,N',N'-Tetraethylethanediamine, 9CI*
[150-77-6]
C₁₀H₂₄N₂ 172.313
Bp 189-192° Bp₇ 72°.
- N,N,N',N'-Tetra-Et; hydrochloride (1:2):*
[92422-58-7]
Mp 186-187°.
- N,N,N',N'-Tetra-Et, dipicrate:* Mp 242-243°.
- N-Propyl:* [111-39-7]
C₅H₁₄N₂ 102.179
Bp 147-150°.
- N-Isopropyl:* [19522-67-9]
C₅H₁₄N₂ 102.179
Liq. d 0.82. Bp 137-138°. n_D²⁰ 1.4369.
- N,N-Dipropyl:* [14165-22-1]
C₈H₂₀N₂ 144.259
d₁₅⁵ 0.83. Mp 192-193° (as picrate). Bp₄₁ 96-98°. n_D²³ 1.4384.
- N,N-Diisopropyl: N,N-Diisopropylethylethanediamine*
[121-05-1]
C₈H₂₀N₂ 144.259
Mp 207-209° (as dihydrobromide). Bp 178° Bp_{3,5} 38°.
- N,N'-Diisopropyl:* [4013-94-9]
C₈H₂₀N₂ 144.259
Bp 169-171°. n_D²⁰ 1.4289.
- N,N,N'-Triisopropyl:* [97-13-2]
C₁₁H₂₆N₂ 186.34
Mp 119-120° (as hydrochloride). Bp 201-203°.
- N,N-Bis(2-methylpropyl):* [14156-98-0]
C₁₀H₂₄N₂ 172.313
Bp 210°.
- N,N-Dibutyl:* [3529-09-7]
C₁₀H₂₄N₂ 172.313
Bp₂₄ 114-117°. n_D²⁰ 1.4430.
- N,N'-Dibutyl:* [4013-95-0]
C₁₀H₂₄N₂ 172.313
Bp₄₀ 167°.
- N,N'-Di-tert-butyl:* [4062-60-6]
C₁₀H₂₄N₂ 172.313
Bp 193°.
- N-Benzyl:* [4152-09-4]
C₉H₁₄N₂ 150.223
Liq. Bp₂₀ 162-165° Bp_{0.1} 100°. n_D²⁰ 1.5417. Forms complexes with metal ions with other ligands; ions include Co²⁺, Cu²⁺, Ni²⁺.
- N-Benzyl; hydrochloride (1:2):* [3303-16-0]
[65358-90-9]
Cryst. (EtOH). Mp 257-259°.
- N-Benzyl, dipicrate:* [86013-60-7]
Yellow cryst. (EtOH). Mp 222° dec.
- N-Benzyl, N,N'-di-Me:* [102-11-4]
C₁₁H₁₈N₂ 178.277
Bp₉ 118-120°.
- N,N'-Dibenzyl, N,N'-di-Me:* [102-18-1]
C₁₈H₂₄N₂ 268.401
Oil. d₄⁹ 0.97. Bp₁₀ 198°.
- N-Ph:* [1664-40-0]
C₈H₁₂N₂ 136.196
Oil. Misc. H₂O. Bp 262-264°.
- N-Ph; hydrochloride:* [126214-34-4]
Mp 199-200° (sinters at 190°).
- N,N'-Bis(2-methylphenyl):* [94-92-8]
C₁₆H₂₀N₂ 240.347
Mp 78°.
- N-Nitro:* [58130-90-8]
C₂H₇N₃O₂ 105.096
Isol. from *Agaricus silvaticus* (red staining mushroom). λ_{max} 228 (ε 7940) (no solvent reported) (pH6).
[20829-66-7, 25723-52-8]
- Aldrich Library of FT-IR Spectra, 1st edn.,* 1985, **1**, 289C; 289D; 303A; 303B; 304A; 304B; 304D; 304C; 305B; 1271D (*ir*)
- Aldrich Library of 13C and 1H FT NMR Spectra,* 1992, **1**, 463B; 463C; 488B; 488C; 489A; 490A; 490B; 491A; 491C; 492A; 492C; 1234B; **2**, 573C (*nmr*)
- Aldrich Library of FT-IR Spectra: Vapor Phase,* 1989, **3**, 374D; 387D; 388A; 388B; 389B; 389C; 390A-390C; 390D; 391B; 1165C; 1167C (*ir*)
- Mills, W.S. *et al.*, *J.C.S.*, 1900, **77**, 1021-1024 (*N,N'-ditolyl*)
- Clarke, H.T. *et al.*, *J.C.S.*, 1911, **99**, 1927-1937 (*dibenzyl dimethyl*)
- Johnson, T.B. *et al.*, *J.A.C.S.*, 1916, **38**, 2135-2145 (*N-Me, N,N'-di-Me, synth*)
- v. Braun, J. *et al.*, *Ber.*, 1918, **51**, 737-741 (*N,N'-di-Me, synth*)
- Meyer, K.H. *et al.*, *Ber.*, 1921, **54**, 2274-2282 (*N,N-di-Et N',N'-di-Me*)
- Ing, H.R. *et al.*, *J.C.S.*, 1926, 2348-2351 (*synth*)
- Bailar, J.C. *et al.*, *J.A.C.S.*, 1934, **56**, 955 (*synth*)
- Chitwood, H.C. *et al.*, *J.A.C.S.*, 1935, **57**, 2424-2426 (*N,N'-didodecanoyl*)
- Bloom, M.S. *et al.*, *J.A.C.S.*, 1945, **67**, 539-540 (*N,N'-di-Et, N,N'-dibutyl, N-Ph*)
- Pearson, D.E. *et al.*, *J.A.C.S.*, 1946, **68**, 1225-1229 (*N-isopropyl*)
- Boon, W.R. *et al.*, *J.C.S.*, 1947, 307-318 (*synth*)
- Takase, S. *et al.*, *Nippon Kagaku Kaishi*, 1948, **69**, 154-157; *CA*, **46**, 3951e (*N,N'-dioctadecanoyl*)
- Hall, R.H. *et al.*, *J.A.C.S.*, 1951, **73**, 2213-2218 (*N-nitro*)
- Abe, K. *et al.*, *Yakugaku Zasshi*, 1955, **75**, 153-159 (*N,N'-di-Me, N,N'-di-Et, synth*)
- McKay, A.F. *et al.*, *Can. J. Chem.*, 1956, **34**, 1567-1573 (*N,N-dibutyl, N,N-dipropyl*)
- McKay, A.F. *et al.*, *J.A.C.S.*, 1956, **78**, 486-488 (*N,N-diisopropyl, N,N-dibutyl, N,N-dipropyl*)
- Agre, C.L. *et al.*, *J.O.C.*, 1956, **21**, 561-564 (*N,N'-didodecanoyl*)
- Fukuto, T.R. *et al.*, *J.A.C.S.*, 1957, **79**, 6083-6085 (*tetra-Et*)
- Epstein, P.F. *et al.*, *J.O.C.*, 1959, **24**, 68-72 (*N,N'-dibutyls*)
- Frost, A.E. *et al.*, *J.O.C.*, 1959, **24**, 1581-1582 (*N-benzyl*)
- U.S. Pat.*, 1960, 2 941 005; *CA*, **54**, 17890f (*N,N'-di-Et N,N'-di-Me*)
- Jucker, E. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 2383-2402 (*N-benzyl N,N'-di-Me*)
- Shepherd, R.G. *et al.*, *J. Med. Chem.*, 1962, **5**, 823-835 (*N,N,N'-trisisopropyl*)
- Fieser and Fieser's Reagents for Organic Synthesis*, Wiley, 1969, **2**, 239; 1974, **4**, 231 (*use*)
- Pryde, E.H. *et al.*, *J. Am. Oil Chem. Soc.*, 1969, **46**, 213-218 (*N,N'-di-9-octadecenoyl*)
- Mariella, R.P. *et al.*, *J.O.C.*, 1971, **36**, 735-737 (*synth, tetra-Ac*)
- Jamet-Deleroix, S. *et al.*, *Acta Cryst. B*, 1973, **29**, 977-980 (*cryst struct*)
- Austin, G.R. *et al.*, *Chem. Eng. (N.Y.)*, 29 April, 1974, **81**, 143-150 (*rev, manuf*)
- Chilton, W.S. *et al.*, *Phytochemistry*, 1975, **14**, 2291-2292 (*N-nitro, isol, ir, uv, synth*)
- Butler, R.N. *et al.*, *J.C.S. Perkin I*, 1976, 386-389 (*Acrawax C*)
- Ivin, K.J. *et al.*, *Makromol. Chem.*, 1978, **179**, 591-599 (*synth, pmr*)
- Nilsson, L. *et al.*, *Acta Chem. Scand., Ser. B*, 1983, **37**, 929-933 (*synth*)
- Newkome, G.R. *et al.*, *J.O.C.*, 1983, **48**, 4848-4851 (*pmr, ms*)
- Bjerrum, M.J. *et al.*, *Inorg. Chem.*, 1986, **25**, 816-821 (*cmr*)
- Houssin, R. *et al.*, *Synthesis*, 1988, 259-261 (*N-tert-butylloxycarbonyl, synth, ir, pmr, ms*)
- Bauermeister, S. *et al.*, *Compr. Inorg. Chem.*, 1991, **69**, 811-816 (*N-benzyl*)
- Encyclopedia of Food and Color Additives*, (ed. Burdock, G.A.), CRC Press, 1997, 970
- Morel, A.F. *et al.*, *Planta Med.*, 1998, **64**, 284-285 (*4-methoxycinnamoyl*)
- Martindale, The Extra Pharmacopoeia, 32nd edn.*, Pharmaceutical Press, 1999, 1577
- Thalladi, V.R. *et al.*, *Angew. Chem., Int. Ed.*, 2000, **39**, 918-922 (*cryst struct*)
- Hanaoka, K. *et al.*, *J.C.S. Perkin 2*, 2001, 1840-1843 (*N-tert-butylloxycarbonyl, synth, pmr, cmr*)
- Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn.*, (ed. Snyder, R.), Elsevier, Volume 2, 1990, 102 (*tox, rev*)
- Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 0899
- Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 376
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, DJ1400; EIW000; DPK400; EEA500

2,2'-(1,2-Ethanediy)bis[1-methyl-3-piperidinone], 9CI
1,2-Bis(3-oxo-2-piperidyl)ethane

E-226

C₁₄H₂₄N₂O₂ 252.356**(RS,SR)-form Hyalbidone**


[138596-73-3]

Alkaloid from the hairy roots of *Hyoscyamus albus* (Solanaceae) transformed with *Agrobacterium rhizogenes*. Yellow oil. $[\alpha]_D^{28}$ -0.13 (c, 0.196 in CHCl₃). *meso*-. Small opt. rotn. must be due to impurities.

Sauerwein, M. *et al.*, *Phytochemistry*, 1991, **30**, 2977 (*isol, uv, pmr, cmr, ms, struct*)


6-Ethenyl- E-227**3,4,4b,5,6,7,8,9,10b,12-decahydro-12-hydroxy-5-methyl-1H-pyrido[4',3':4,5]pyrano[3,2-h]isoquinoline-1,10(2H)-dione, 9CI**

[118984-42-2]

C₁₈H₂₂N₂O₄ 330.383

Alkaloid from the whole plant of *Schulnesia guianensis* (Gentianaceae). Mp 192-193°.

Nóbrega, E.M. *et al.*, *J. Nat. Prod.*, 1988, **51**, 962 (*isol, ir, pmr, ms, cryst struct*)

16-Ethenyloxayohimban-17-ol, 9CI E-228C₂₀H₂₄N₂O₂ 324.422

Undefined stereochem.

O-β-D-Glucopyranoside: [62835-74-9]

C₂₆H₃₄N₂O₇ 486.564

Alkaloid from the leaves of *Strychnos decussata* (Loganiaceae). Mp 201°. $[\alpha]_D^{25}$ -134 (c, 0.6 in MeOH).

Petitjean, A. *et al.*, *Phytochemistry*, 1977, **16**, 154-155 (*isol, uv, pmr, ms, struct*)

Ethesdanine


U 20207. Antibiotic U 20207

C₉H₁₆N₂O₅ 232.236

Pyrrolidine antibiotic. Struct. unknown. Prod. by *Streptomyces caelestis* with S-Ethylcysteine. Mainly active against gram-positive bacteria. Sol. H₂O, butanol; poorly sol. CHCl₃, hexane.

▶ LD₅₀ (mus, ipr) 60 mg/kg.*Hydrochloride*: Mp 155-158°.

Meyer, C.E. *et al.*, *Antimicrob. Agents Chemother.*, 1965, **850** (*isol, ir*)

N-Ethoxycarbonylprolina- E-230*Ethyl 2-(aminocarbonyl)-1-pyrrolidine-carboxylate, 9CI*C₈H₁₄N₂O₃ 186.21**(S)-form***L-form*


[62787-66-0]

Alkaloid from the leaves of *Arnica montana* (Asteraceae). Cryst. (Et₂O). Mp 103-103.5°. $[\alpha]_D^{20}$ -55.8 (c, 0.12 in MeOH).

Holub, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1971, **36**, 2189; 1977, **42**, 151 (*isol, struct, ord, cd, ms, pmr, synth*)

4-Ethoxyhexahydro-3a-hydroxy-1H-indole-2,6-dione, 9CI E-231

[116964-15-9]


C₁₀H₁₅NO₄ 213.233

Isol. from *Senecio anonymus* (Asteraceae). Mp 170°. Possibly an artifact.

Zalkow, L.H. *et al.*, *J. Nat. Prod.*, 1988, **51**, 690 (*isol, pmr, ms, cryst struct*)

8-Ethoxy-3-oxo-1,2-dehydropretorsine E-232

[101029-56-5]

C₂₀H₂₇NO₈ 409.435

Alkaloid from the aerial parts of *Senecio grisebachii* (Asteraceae).

Hirschmann, S. *et al.*, *Rev. Latinoam. Quim.*, 1985, **16**, 109; *CA*, **104**, 126516g

Ethylamine, 8CI*Ethanamine, 9CI*

[75-04-7]

H₃CCH₂NH₂C₂H₇N 45.084

Manuf. by catalytic amination of ethanol with ammonia. Prod. by marine algae, *Clostridium* spp., *Candida albicans*, *Brevibacterium linens* and *Streptococcus lactis*. Used in manuf. of resins, rubber, herbicides, etc. Flammable gas with ammoniacal odour. Misc. EtOH, Et₂O, H₂O, salted out by NaOH. Fp -80. Bp 16.6°. pK_a 10.7.

▶ Extremely flammable, fl. p. <-18°, auto-ignition temp. 580°. Skin, respiratory tract and severe eye irritant. High vapour conc. may affect CNS. LD₅₀ (rat, ori) 400 mg/kg. LD₅₀ (rbt, skn) 390 mg/kg. OES: long-term 10 ppm. KH2100000


[16999-99-8]

Lerch, B. *et al.*, *Z. Naturforsch., B*, 1966, **21**, 216 (*occur*)

Steiner, M. *et al.*, *Planta*, 1968, **79**, 113 (*isol*)
Neurath, G.B. *et al.*, *Food Cosmet. Toxicol.*, 1979, **15**, 275 (*isol*)

1-Ethyl-β-carboline E-234*1-Ethyl-9H-pyrido[3,4-b]indole, 9CI*

[20127-61-1]

C₁₃H₁₂N₂ 196.251

Alkaloid from the roots of *Hannoa klaineana* (Simaroubaceae) and the bryozoan *Cribricellina cribraria*. Also from *Costaticella hastata*. Whitish-yellow needles (MeOH/CHCl₃). Mp 202-203° (synthetic) (194-195°).

2-N-Oxide: 1-Ethyl-beta-carboline-2N-oxide [90686-26-3]

C₁₃H₁₂N₂O 212.251

Alkaloid from the roots of *Hannoa klaineana* (Simaroubaceae). Whitish-yellow needles (MeOH/CHCl₃).

Lumonadio, L. *et al.*, *Phytochemistry*, 1984, **23**, 453-455 (*isol, uv, ir, pmr, ms, struct*)


Blackman, A.J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 494-496 (*isol, Costaticella*)

Rocca, P. *et al.*, *Tetrahedron*, 1993, **49**, 3325-3342 (*synth, cmr*)

Ivanov, I. *et al.*, *Heterocycles*, 2005, **65**, 2483-2492 (*synth, pmr*)

Ethylcyclononylprodigiosin E-235*Antibiotic 610FF*


[52589-17-0]

C₂₅H₃₃N₃O 391.555

Pyrrole antibiotic. Prod. by *Actinomadura pelletieri*. Active against gram-positive bacteria. Yellow solid. Red in salt form. λ_{\max} 550 (CHCl₃) (Berdy). λ_{\max} 541 (EtOH-HCl) (Berdy).

Gerber, N.N. et al., *J. Het. Chem.*, 1973, **10**, 925; *Can. J. Microbiol.*, 1976, **22**, 658 (isol, struct)

3-Ethyl-1,3-dihydro-3-methoxy-2H-indol-2-one, 9CI E-236
3-Ethyl-3-methoxyoxindole
[66424-33-7]




C₁₁H₁₃NO₂ 191.229

Isol. from *Streptomyces antibioticus*. Active against gram-positive bacteria. Mp 179°.

Westley, J.W. et al., *J. Antibiot.*, 1979, **32**, 100 (isol)

2-Ethyl-3,5-dimethylpyrazine E-237
FEMA 3150
[13925-07-0]



C₈H₁₂N₂ 136.196

Isol. from an arctic marine bacterium. Isol. from coffee aroma. Also present in raw asparagus, wheat bread, other breads, smoked fatty fish, roast chicken, roast beef, lamb and mutton liver, black tea, hydrolysed soy protein and other foods. Organoleptic agent. Flavouring agent. Bp₁₉ 80-81° Bp₈ 64-66°. n_D^{22} 1.4975.

► UQ3155000

Klein, B. et al., *J.A.C.S.*, 1951, **73**, 2949 (synth)
Goldman, I.M. et al., *Helv. Chim. Acta*, 1967, **50**, 694

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 968-969 (use, occur)

Dickschat, J.S. et al., *Chem. Biodiversity*, 2005, **2**, 318-353 (isol)

3-Ethyl-2,5-dimethylpyrazine E-238
[13360-65-1]

C₈H₁₂N₂ 136.196

Pheromone of various ant spp. Isol. from various marine bacteria and from coffee. Organoleptic agent with nutty roasted odour. Bp 180.5° Bp₁₁ 72-73°. n_D^{24} 1.4945.

Goldman, I.M. et al., *Helv. Chim. Acta*, 1967, **50**, 694

Gelas, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1968, **266**, 625 (synth, uv, ir, pmr)

Ohta, A. et al., *Heterocycles*, 1987, **26**, 2449 (synth)

Buechi, G. et al., *J.O.C.*, 1991, **56**, 2605 (synth)

Dickschat, J.S. et al., *Chem. Biodiversity*, 2005, **2**, 318-353 (isol)

Dickschat, J.S. et al., *Eur. J. Org. Chem.*, 2005, 4141-4153 (isol)

5-Ethyl-2,3-dimethylpyrazine E-239
[15707-34-3]

C₈H₁₂N₂ 136.196

Constit. of Galbanum oil. Urinary signalling pheromone for tree shrew *Tupaia belangeri*. Liq. Bp₁₄ 72°. n_D^{24} 1.4982.

Gelas, J. et al., *C. R. Hebd. Seances Acad. Sci.*, 1968, **266**, 625 (synth)


Rizzi, G.P. et al., *J.O.C.*, 1968, **33**, 1333 (synth)
Netherlands Pat., 1969, 68 121 899; *CA*, **71**, 61421 (synth)

Burrell, J.W.K. et al., *Chem. Ind. (London)*, 1970, 1409 (isol)

von Stralendorff, F. et al., *J. Chem. Ecol.*, 1987, **13**, 655

8-Ethyl-5-(1-ethyl-1-butenyl)-1-octahydro-6-methylindolizine E-240

8-Ethyl-5-(1-ethyl-1-butenyl)-6-methylindolizidine. **Indolizidine 249H**. Dendrobates *Alkaloid 249H*
[220624-30-6]




C₁₇H₃₁N 249.439

Alkaloid from the skin extract of a Panamanian frog *Dendrobates pumilio*.

Tokuyama, T. et al., *Heterocycles*, 1998, **49**, 427-436

N-Ethylhastanecine trachelanthate E-241




C₁₇H₃₂NO₅⁺ 330.444

Quaternary alkaloid from *Lindelofia macrostyla* (Boraginaceae). Mp 149-151° (as chloride). $[\alpha]_D^{20}$ +20.25 (MeOH). Alkaloid not named in the lit.

Tsyru'nikova, L.G. et al., *Zh. Obshch. Khim.*, 1962, **32**, 2705-2709; *J. Gen. Chem. USSR (Engl. Transl.)*, 1962, **32**, 2663-2666 (isol, struct)

Aasen, A.J. et al., *J.O.C.*, 1969, **34**, 4137-4143 (config)

4-Ethyl-2,4-hexadienoic acid, 9CI E-242



C₈H₁₂O₂ 140.182

(E,E)-form [92454-23-4]

Amide: **Antibiotic FR 900411**. FR 900411 [92454-24-5]

C₈H₁₃NO 139.197

Prod. by *Streptomyces griseosporus*.

Cryst. (Et₂O). λ_{\max} 260 (ε 20000) (no solvent reported).

[152873-92-2]


Hino, M. et al., *Chem. Pharm. Bull.*, 1989, **37**, 2864-2866 (amide, synth)

Hino, M. et al., *J. Antibiot.*, 1989, **42**, 1578-1583; 1584-1588; 1589-1592 (amide, isol, synth)

Huh, K.T. et al., *J.O.C.*, 1993, **58**, 6956-6957 (synth)

4-Ethyl-2-(hydroxyimino)-5-nitro-3-hexenamide, 9CI E-243

4-Ethyl-5-nitro-2-oxo-3-hexenamide oxime. **FK409**. FR 900409
[92454-83-6]



C₈H₁₃N₃O₄ 215.208

Prod. by *Streptomyces griseosporus*.

Platelet aggregation inhibitor. Nitric oxide releaser (in soln. pH 7.4). Coronary vasodilator. Hypotensive agent. Prisms (MeOH). Sol. MeOH; fairly sol. EtOAc, H₂O, CHCl₃. Mp 140° dec. $[\alpha]_D$ 0. λ_{\max} 273 (ε 13500) (MeOH/NaOH) (Derep). λ_{\max} 240 (sh) (ε 7000); 273 (ε 13500) (MeOH) (Derep).

[92454-60-9]

Hino, M. et al., *Chem. Pharm. Bull.*, 1989, **37**, 2864 (isol, struct)

Hino, M. et al., *J. Antibiot.*, 1989, **42**, 1578; 1584; 1589 (isol, struct, props)

Japan. Pat., 1991, ((Fujisawa))03 240 763; *CA*, **116**, 143869q


Kita, Y. et al., *Br. J. Pharmacol.*, 1994, **113**, 385 (pharmacol)

Kita, Y. et al., *Eur. J. Pharmacol.*, 1994, **257**, 123 (pharmacol)

Hirasawa, Y. et al., *Eur. J. Pharmacol.*, 1995, **272**, 39 (pharmacol)

6-Ethyl-2-hydroxymethyl-3,4-piperidinediol, 9CI E-244

6-Ethyl-3,4-dihydroxy-2-hydroxymethylpiperidine



(2R,3S,4S,6R)-form

C₈H₁₇NO₃ 175.227

(2R,3S,4S,6R)-form 5-Deoxyadenophorine

Constit. of the roots of *Adenophora* spp. Oil. $[\alpha]_D^{20}$ +52.3 (c, 0.38 in H₂O).

1'-O-β-D-Glucopyranoside: [262616-01-3]

C₁₄H₂₇NO₈ 337.369

Constit. of the roots of *Adenophora* spp. Oil. $[\alpha]_D^{20}$ +17.6 (c, 0.87 in H₂O).

(2R,3R,4R,6R)-form*1 α* -C-Ethylfagomine

Alkaloid from *Adenophora triphylla* var. *japonica*. Inhibits bovine liver β -galactosidase. $[\alpha]_D^{25} +45.7$ (c, 0.71 in H₂O).


Ikeda, K. *et al.*, *Carbohydr. Res.*, 2000, **323**, 73-80 (*isol*, *pmr*, *cmr*)

Asano, N. *et al.*, *Phytochemistry*, 2000, **53**, 379-382 (*1-C-Ethylfagomine*)

Felplin, F.-X. *et al.*, *J.O.C.*, 2004, **69**, 1497-1503 (*synth*, *pmr*, *cmr*, *abs config*)

2-Ethyl-6-hydroxymethyl-3,4,5-piperidinetriol E-245

2-Ethyl-3,4,5-trihydroxy-6-hydroxymethylpiperidine



C₈H₁₇NO₄ 191.227

(2R,3R,4S,5S,6R)-form*Adenophorine*

Alkaloid from the roots of *Adenophora* spp. $[\alpha]_D^{25} +59.7$ (c, 1 in H₂O).

1''-O- β -D-Glucopyranoside:

C₁₄H₂₇NO₉ 353.369

Constit. of the roots of *Adenophora* spp. $[\alpha]_D^{25} +24.9$ (c, 0.92 in H₂O).

(2S,3R,4R,5R,6R)-form [125711-56-0]

Alkaloid from the bulbs of *Scilla socialis*. Powder. $[\alpha]_D^{25} +5$ (c, 0.19 in H₂O).

Ikeda, K. *et al.*, *Carbohydr. Res.*, 2000, **323**, 73-80 (*isol*)

Pearson, M.S.M. *et al.*, *Eur. J. Org. Chem.*, 2007, 4888-4894 (*synth*)

Kato, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 993-997 (*isol*, *pmr*, *cmr*)

2-Ethyl-3-hydroxypyrazine E-246

3-Ethyl-2(1H)-pyrazinone, 9CI. 3-Ethylpyrazinol, 8CI [25680-54-0]



C₆H₈N₂O 124.142

Cryst. (C₆H₆/pentane). Mp 105°.

OH-form

Me ether: 3-Ethyl-2-methoxypyrazine [25680-58-4]

C₇H₁₀N₂O 138.169

Occurs in potato. Bp₂₀ 95-98°. FEMA 3280 is an isomeric mixt. of ethylmethoxypyrazines and methylmethoxypyrazines.

U.S. Pat., 1973, 3 772 039; *CA*, **80**, 58618a (*synth*, *Me ether*, *use*)


Nursten, H.E. *et al.*, *J. Sci. Food Agric.*, 1974, **25**, 643-663 (*Me ether*, *occur*, *ms*)

Sala, C. *et al.*, *J. Chromatogr. A*, 2000, **880**, 93-99 (*Me ether*, *occur*)

Fenaroli's Handbook of Flavor Ingredients, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 563-564 (FEMA 3280)

2-Ethyl-4-hydroxyquinazolinone E-247

2-Ethyl-4(3H)-quinazolinone. 2-Ethyl-4-quinazolinol



C₁₀H₁₀N₂O 174.202

1H-form

N-Me: 2-Ethyl-1-methyl-4(1H)-quinazolinone. *Homoglomerine*

[10553-04-5]

C₁₁H₁₂N₂O 188.229

Alkaloid from the defensive secretion of the arthropod *Glomeris marginata*. Prisms (EtOAc). Mp 148-149°.

3H-form

N-Amino: 3-Amino-2-ethyl-4(3H)-quinazolinone

[50547-51-8]

C₁₀H₁₁N₃O 189.216

Cryst. (EtOH). Mp 122-123° Mp 152-153°.

Chakravarti, D. *et al.*, *Tetrahedron*, 1961, **16**, 224 (*synth*, *uv*, *ir*)

Schildknecht, H. *et al.*, *Naturwissenschaften*, 1967, **54**, 196 (*Homoglomerine*)

Schildknecht, H. *et al.*, *Tet. Lett.*, 1967, 1815 (*biosynth*)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1977, 2347 (*synth*, *pmr*)

Naik, N.R. *et al.*, *J. Indian Chem. Soc.*, 1979, **56**, 708 (*synth*, *uv*, *ir*)


Soliman, R. *et al.*, *Synthesis*, 1979, 803 (*N-amino*)

Bhattacharyya, J. *et al.*, *Heterocycles*, 1980, **14**, 1469 (*cmr*)

Leiby, R.W. *et al.*, *J.O.C.*, 1985, **50**, 2926 (*N-amino*, *pmr*, *ir*)

3-Ethylidene-1,3-dihydro-2H-indol-2-one, 9CI E-248

3-Ethylideneoxindole



C₁₀H₉NO 159.187

(E)-form

CG-SI-1

[58807-92-4]

Prod. by *Colletotrichum fragariae* and *Colletotrichum gloeosporoides*. Spore germination self-inhibitor. Yellow needles (C₆H₆). Mp 140-142°.

(Z)-form

CG-SI-2

[58807-93-5]

Prod. by *Colletotrichum fragariae* and *Colletotrichum gloeosporoides*. Spore germination self-inhibitor. Yellow needles (C₆H₆). Mp 177-178° (173-174°).

Tacconi, G. *et al.*, *J.C.S. Perkin 2*, 1976, 150-154 (*synth*, *ir*, *pmr*, *bibl*)

Nozoye, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2986-2991 (*synth*)


Goehring, R.R. *et al.*, *J.A.C.S.*, 1985, **107**, 435-443 (*synth*, *pmr*)

Ueno, T. *et al.*, *Stud. Nat. Prod. Chem.*, 1991, **9**, 219-248 (*isol*)

Inoue, M. *et al.*, *J. Chem. Ecol.*, 1996, **22**, 2111-2122 (*isol*, *struct*)

3-Ethylidenehexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione, 9CI E-249

1,6-Propano-3-ethylidene-2,5-piperazine-dione



C₉H₁₂N₂O₂ 180.206


(S)-form [160706-82-1]

Isol. from cultured cells of *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Amorph. powder.

Aimi, N. *et al.*, *Heterocycles*, 1994, **38**, 2411 (*isol*, *pmr*, *cmr*, *struct*)

3-Ethylidene-2-pyrrolidinone, 9CI E-250

[27428-45-1]



C₆H₉NO 111.143

(E)-form

Corydactam. Alkaloid P

[930-94-9]

Constit. of *Corydalis pallida* (Papaveraceae). Needles (EtOAc). Mp 176-177.5°.

(Z)-form

Cryst. (petrol). Mp 98-99.5°.

Cummings, W.A. *et al.*, *J.C.S.*, 1964, 4591 (*synth*, *uv*)


Kametani, T. *et al.*, *J.C.S.(C)*, 1970, 1060; 1971, 999 (*isol*, *synth*)

Kaneko, H. *et al.*, *Yakugaku Zasshi*, 1971, **91**, 101 (*isol*)

Warnhoff, H. *et al.*, *Synthesis*, 1976, 331 (*synth*)

2-Ethyl-11-methoxy-3-methyl-3H-[1,6]naphthyridino[6,5,4-*def*]quinoxaline, 9CI E-251

[599173-39-4]




C₁₇H₁₇N₃O 279.341

Alkaloid from a *Xestospongia* sp. Orange gum. λ_{\max} 208 (ϵ 7400); 225 (ϵ 6590); 237 (ϵ 7390); 251 (ϵ 5550); 261 (ϵ 4400); 272 (ϵ 3470); 360 (ϵ 1970) (EtOH).

Calcul, L. *et al.*, *Tetrahedron*, 2003, **59**, 6539-6544 (*isol, pmr, cmr*)

2-Ethyl-3-methyl-2-butenedioic acid, 9CI E-252

2-Pentene-2,3-dioic acid. dibasic-Haematinic acid



$C_7H_{10}O_4$ 158.154
 pK_{a1} 3.2; pK_{a2} 6.1 (20°).

(*E*)-form

2-Ethyl-3-methylfumaric acid
 [28098-80-8]
 Mp 178-183°.

(*Z*)-form

2-Ethyl-3-methylmaleic acid
 Exists in equilib. with the anhydride.
 Constit. of Turkish tobacco.

Di-Me ester: [41654-26-6]
 $C_9H_{14}O_4$ 186.207
 Bp 235°.

Anhydride:
 $C_7H_8O_3$ 140.138
 Bp 230°.

Imide: 3-Ethyl-4-methyl-1H-pyrrole-2,5-dione, 9CI. 2-Ethyl-3-methylmaleimide [20189-42-8]
 $C_7H_9NO_2$ 139.154
 Constit. of tobacco. Isol. from sediments of Tokyo Bay. Degradn. prod. of Chlorophylls. Needles. Mp 72° (68°).

[57959-42-9, 119581-55-4, 84565-24-2]

Küster, W. *et al.*, *Annalen*, 1906, **345**, 10 (*synth*)
 Ebersson, L. *et al.*, *Acta Chem. Scand.*, 1964, **18**, 1276 (*struct*)

Brockmann, H. *et al.*, *Chem. Ber.*, 1973, **106**, 803 (*synth*)

Chuman, T. *et al.*, *Agric. Biol. Chem.*, 1977, **41**, 1021 (*occur*)


Gossauer, A. *et al.*, *Annalen*, 1977, 664 (*synth*)
 Girard, C. *et al.*, *Tet. Lett.*, 1982, **23**, 3683 (*synth*)

Kozono, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 1844-1847 (*Imide, isol, synth, pmr, ms*)

1-Ethyl-3-(2-methylbutyl)peridine, 9CI E-253

Stenusine

[54985-88-5]
 [150447-27-1 (2'R,3S-form), 144301-03-1 (2'RS,3SR-form), 144301-02-0 (2'RS,3RS-form), 150447-26-0 (2'R,3R-form)]



$C_{12}H_{25}N$ 183.336

Natural Stenusine is a mixt. of all four stereoisomers in species-specific ratios.

Alkaloid from the staphylinid beetles *Stenus* spp. Major component of the spreading (flotation) agent of the beetles.

(2'*S*,3*S*)-form [150447-24-8]

Oil. $[\alpha]_{365}^{25}$ +34.4 (c, 0.27 in EtOH).

(2'*S*,3*R*)-form [150447-25-9]

$[\alpha]_{365}^{24}$ +68 (c, 0.52 in EtOH).

Schildknecht, H. *et al.*, *J. Neurochem.*, 1975, **14**, 427 (*isol, ms, pmr, synth*)

Enders, D. *et al.*, *J.O.C.*, 1993, **58**, 4881 (*synth, ir, pmr, cmr, ms*)

Micouin, L. *et al.*, *Tet. Lett.*, 1994, **35**, 2529 (*synth*)


Heilporn, S. *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 193-200 (*synth*)

Gedig, T. *et al.*, *Tetrahedron*, 2007, **63**, 2670-2674 (*synth*)

Lusebrink, I. *et al.*, *J. Nat. Prod.*, 2008, **71**, 743-745 (*biosynth*)

2-Ethyl-6-methyl-3,4,5-piperidinetriol E-254

2-Ethyl-3,4,5-trihydroxy-6-methylpiperidine



$C_8H_{17}NO_3$ 175.227


(2*R*,3*R*,4*R*,5*S*,6*R*)-form 1-Deoxyadenophorine

Constit. of the roots of *Adenophora* spp.
 $[\alpha]_D$ +67.8 (c, 0.83 in H₂O).

Ikeda, K. *et al.*, *Carbohydr. Res.*, 2000, **323**, 73-80

2-Ethyl-5-methylpyrazine, 9CI, 8CI E-255

FEMA 3154
 [13360-64-0]



Found in roast peanuts, roast filberts, cooked beef, cooked lamb, coffee beans, cocoa butter, roast barley and other foodstuffs. Prod. by various marine bacteria. Found in cocoa butter and other foodstuffs. Used in food flavouring. Bp₅₆ 79-80°. n_D^{25} 1.4925. Odour threshold 100 ppb in H₂O.

Goldman, I.M. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 694

U.S. Pat., 1976, 3 952 026; *CA*, **71**, 90131m (*manif, uses*)

Shibamoto, T. *et al.*, *J. Agric. Food Chem.*, 1979, **27**, 1027 (*formn*)

Milic, B.L. *et al.*, *Food Chem.*, 1984, **13**, 165 (*gc*)
 Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*)

5-Ethyl-2-methylpyridine E-256

5-Ethyl- α -picoline. Aldehydecollidine.

FEMA 3546

[104-90-5]

$C_8H_{11}N$ 121.182

Present in dry red beans, cocoa and tea. Solvent, proton acceptor, comly. available. Flavouring agent. Liq. Part misc. H₂O. d^{20} 0.94. Bp 178.3°. n_D^{20} 1.4978. pK_a 6.51 (20°). Steam-volatile.

► Violent reactn. with HNO₃. Severe skin and eye irritant. LD₅₀ (rat, orl) 368 mg/kg; LD₅₀ (rbt, skn) 1000 mg/kg. TJ6825000

Picrate:

Yellow plates or needles (H₂O or EtOH aq.). Mp 164-165°.

N-Oxide: [768-44-5]

C₈H₁₁NO 137.181

► Mod. toxic, skin irritant. TJ7000000

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 742D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 256A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1523A (*ir*)

Org. Synth., Coll. Vol., **4**, 1963, 451 (*bibl*)


Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 253

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1015-1016

Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 2801

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EOS000

5-Ethyl-4-methyl-3-pyridine-carboxylic acid E-257



$C_9H_{11}NO_2$ 165.191

Me ester: Alkaloid LA 2

[4695-97-0]


$C_{10}H_{13}NO_2$ 179.218

Alkaloid artifact generated by NH₃ and secoiridoid glucosides from ripe fruits of the common privet *Ligustrum vulgare* (Oleaceae). Needles (EtOAc/MeOH). Mp 23°.

Willems, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1987, **320**, 1245 (*isol, uv, ir, pmr, cmr, ms, struct*)

5-Ethyl-2-methyl-11H-pyrido[3,4-*a*]carbazolium(1+), 9CI E-258

[76787-84-3]




$C_{18}H_{17}N_2^{\oplus}$ 261.346

Anomalous indole alkaloid. Appears not to be derived from Tryptamine and Secologanin. Alkaloid from the bark of *Aspidosperma gilbertii* (Apocynaceae).

Mp 270° (as hydroxide). CAS no. refers to hydroxide.

Miranda, E.C. *et al.*, *Chem. Ber.*, 1980, **113**, 3245 (*isol, uv, ir, pmr, ms, struct, synth*)

2-Ethyl-4-methylquinazoline E-259
[13535-90-5]



C₁₁H₁₂N₂ 172.229


Metab. of the bacterium *Pseudomonas aeruginosa*. Oil, which cryst. slowly on standing. Bp 259-260°.

Picrate: Mp 100° dec.

Bischler, A. *et al.*, *Ber.*, 1893, **26**, 1384 (*synth*)
Mann, S. *et al.*, *Arch. Mikrobiol.*, 1967, **56**, 324

1-Ethyl-4-methylsulfonyl-β-carboline E-260

1-Ethyl-4-(methylsulfonyl)-9H-pyrido[3,4-b]indole, 9CI
[138683-70-2]




C₁₄H₁₄N₂O₂S 274.343

Alkaloid from the marine bryozoan *Cribricellina cribraria*. Pale green oil. λ_{max} 213 (ε 2690); 250 (ε 1950); 258 (ε 2090); 299 (ε 871); 304 (ε 1070); 350 (ε 417) (MeOH) (Derep).

Prinsep, M.R. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1068 (*isol, uv, ir, pmr, cmr, ms, struct*)

2-Ethyl-4-methylthiazole E-261
FEMA 3680
[15679-12-6]



C₆H₉NS 127.21


Occurs in fennel *Foeniculum vulgare*. Liq. d₄²⁰ 1.03. Bp_{728.5} 160.6-161°. n_D²⁰ 1.5059.

Benbadjarian, A. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 4878 (*synth, pmr, ir, uv*)

Vitzthum, O.G. *et al.*, *J. Food Sci.*, 1974, **39**, 1210 (*isol*)

Haag, A. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 511 (*ms*)

1*b*-Ethyl-1*a*,1*b*,3,4,8*c*,9,10,10*a*-octahydro-2*H*-oxireno[4,5-*b*]pyrido[3,2-*c*]carbazole, 9CI E-262
[80249-75-8]



Relative configuration


C₁₇H₂₀N₂O 268.358

Biosynth. not certain. May be derived from Aspidospermidine, A-1498 by Grob elimination. Minor alkaloid from *Voacanga africana* (Apocynaceae). Cryst. (C₆H₆). Mp 233-235°. [α]_D²⁰ +109 (c, 0.1 in CHCl₃).

Kunesch, N. *et al.*, *Tet. Lett.*, 1981, **22**, 1981 (*uv, ir, ms, pmr, cmr, struct*)

6-Ethyl-8-(2-hydroxyethyl)-5-pentyl-8-indolizineethanol E-263

6-Ethyl-8-(2-hydroxyethyl)-5-pentylindolizidine. *Dendrobates Alkaloid 267J*. *Indolizidine 267J*
[185417-28-1]



C₁₇H₃₃NO 267.454

Tentative struct. and relative stereochem. Alkaloid from skin extracts of a Panamanian population of the frog *Dendrobates pumilio*.

Deoxy: 6,8-Diethyl-8-(2-hydroxyethyl)-5-pentylindolizine. 6,8-Diethyl-5-pentylindolizidine. *Dendrobates Alkaloid 251M*. *Indolizidine 251M*
[185417-27-0]


C₁₇H₃₃N 251.454

Alkaloid from skin extracts of *Dendrobates pumilio*. Tentative struct.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 2 (*isol, ms, ir*)

1-Ethyl-4-(4-pentynyl)quinolizidine E-264

1-Ethyl-4-(4-pentynyl)-2H-quinolizidine. *Dendrobates Alkaloid 233A*. *Quinolizidine 233A*
[151805-15-1]



C₁₆H₂₇N 233.396

Alkaloid from the frog, *Mantella* spp.


Jain, P. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1174-1178 (*isol*)

Michel, P. *et al.*, *J.O.C.*, 2000, **65**, 8908-8918 (*synth*)

Kinderman, S.S. *et al.*, *J.A.C.S.*, 2004, **126**, 4100-4101 (*synth*)

1-Ethyl-4-(2-propenyl)quinolizidine E-265

1-Ethyl-4-(2-propenyl)-2H-quinolizidine, 9CI. *4-Allyl-1-ethylquinolizidine*. *Dendrobates Alkaloid 207I*. *Quinolizidine 207I*
[194344-58-6]
[194351-57-0]



Relative Configuration

C₁₄H₂₅N 207.358

Minor alkaloid from the frog *Dendrobates* spp. Mp 205-207° (as hydrochloride) (synthetic). [α]_D²⁶ -97.9 (c, 0.5 in CHCl₃) (synthetic).


Toyooka, N. *et al.*, *Tetrahedron*, 1997, **53**, 9553-9574 (*synth, ir, ms, pmr, cmr*)

Pearson, W.H. *et al.*, *J.O.C.*, 1998, **63**, 9910-9918 (*synth*)

Michel, P. *et al.*, *Chem. Comm.*, 1999, 2281-2282 (*synth*)

Michel, P. *et al.*, *J.O.C.*, 2000, **65**, 8908-8918 (*synth*)

2-(1-Ethylpropyl)-4-methoxyquinoline, 9CI E-266
[126365-16-0]




C₁₅H₁₉NO 229.321

Alkaloid from roots of *Esenbeckia leiocarpa*. Cryst. (CH₂Cl₂/heptane). Mp 72-73° Mp 77.8-78.5° synthetic.

Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (*isol, uv, pmr, cmr, ms, struct*)

Nakatsu, T. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1508 (*synth*)

2-(1-Ethylpropyl)-1-methyl-4(1*H*)-quinolinone E-267
Leikinine B
[126365-17-1]




C₁₅H₁₉NO 229.321

Alkaloid from the roots and leaves of *Esenbeckia leiocarpa* (Rutaceae). Shows antifeedant activity against *Pectinophora gossypiella*. Pale yellow prisms (Et₂O). Mp 137-140°.

Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (*isol, uv, ir, pmr*)

Nakatsu, T. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1508
(*isol*, *pmr*, *cmr*, *struct*, *synth*)
Coppola, G.M. *et al.*, *J. Het. Chem.*, 1992, **29**,
1873 (*synth*)

3-Ethylpyridine **E-268**
FEMA 3394
[536-78-7]



C₇H₉N 107.155
Found in black tea and raw fish. Fla-
vouring ingredient. Liq. with a tobacco
flavour. Sl. sol. H₂O, sol. EtOH, Et₂O. d²⁰
0.95. Bp 162-165°. n_D²⁰ 1.5020. pK_a 5.56
(25°).

Picrate: Mp 128-130°.

N-Oxide: [14906-62-8]

C₇H₉NO 123.154
Bp₄ 100°.

Aldrich Library of FT-IR Spectra, 1st edn.,
1985, **2**, 735C (*ir*)

Aldrich Library of 13C and 1H FT NMR
Spectra, 1992, **3**, 244C (*nmr*)

Mijajima, G. *et al.*, *Chem. Pharm. Bull.*, 1972,
20, 429 (*cmr*)


Ferles, M. *et al.*, *Coll. Czech. Chem. Comm.*,
1973, **38**, 611 (*synth*)

Fenaroli's Handbook of Flavor Ingredients, 3rd
edn., (ed. Burdock, G.A.), CRC Press, 1995,
2, 266

Encyclopedia of Food and Color Additives, (ed.
Burdock, G.A.), CRC Press, 1997, 1038

Bremner, D.H. *et al.*, *Synth. Commun.*, 1997,
27, 1535-1542 (*N-oxide*)

5-Ethyl-3-pyridinecarboxylic acid **E-269**
acid



C₈H₉NO₂ 151.165

Me ester: **Alkaloid LA 1**
[68686-58-8]


C₉H₁₁NO₂ 165.191

Alkaloid artifact generated by NH₃
and secoiridoid glucosides from ripe
fruits of the common privet *Ligustrum*
vulgare (Oleaceae). Needles (EtOAc/
MeOH). Mp 8°.

Me ester, hydrochloride:
Cryst. (Et₂O). Mp 112°.


Willems, M. *et al.*, *Arch. Pharm. (Weinheim,
Ger.)*, 1987, **320**, 1245 (*isol*, *uv*, *ir*, *pmr*, *cmr*,
ms, *struct*)

3-Ethyl-2-pyrrolidinone, 9CI, 8CI **E-270**
Alkaloid CP-P
[930-92-7]



C₆H₁₁NO 113.159
Found in *Corydalis pallida*. Mp 38-39°
Mp 43-44° Mp 47-48.5°. Hygroscopic.
Cummings, W.A.W. *et al.*, *J.C.S.*, 1964, 4591
(*synth*, *uv*)
Kametani, T. *et al.*, *J.C.S.(C)*, 1970, 1060
(*synth*)
Japan. Pat., 1973, 73 08 484; *CA*, **80**,
41035k
Reddy, P.A. *et al.*, *J. Med. Chem.*, 1996, **39**,
1898-1906 (*synth*, *ir*, *pmr*, *cmr*)

3-Ethyl-1,2,3,4-tetrahydro-1,3-dimethylisoquinoline **E-271**



C₁₃H₁₉N 189.3

(1R*,3R*)-form
Hemiargine D


[710948-57-5]

Constit. of the leaves of *Croton hemi-*
argyreus var. *gymnodiscus*.

Lin, W.-H. *et al.*, *J. Chin. Pharm. Sci.*, 2003,
12, 117-122 (*isol*)

5-[(4-Ethyltetrahydro-5-oxo-3-furanyl)methylidene]-1-methyl-2,4-imidazolidinedione, 9CI **E-272**

4,6-Dehydro-1,2,4,5-tetrahydro-2,5-dioxo-
pilocarpine
[876945-76-5]




Absolute
Configuration

C₁₁H₁₄N₂O₄ 238.243

Related to Pilocarpine, P-424. Alkaloid
from the stems of *Pilocarpus grand-*
iflorus. Yellow oil. [α]_D²⁵ +19.6
(c, 0.005 in CHCl₃). λ_{max} 242 (log ε
3.46); 284 (log ε 3.98)
(MeOH).

De Souza, R.C. *et al.*, *Z. Naturforsch., B.*
2005, **60**, 787-791 (*isol*, *pmr*, *cmr*, *ms*)

Ethyltrimethylpyrazine **E-273**
[17398-16-2]



C₉H₁₄N₂ 150.223


Prod. by various marine bacteria.

Marion, J.P. *et al.*, *Chimia*, 1967, **21**, 510-511;
CA, **68**, 78249m (*synth*)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005,
2, 318-353 (*isol*)

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005,
4141-4153 (*isol*)

Euchrestifoline **E-274**
[31231-34-2]




C₁₈H₁₇NO₂ 279.338

Alkaloid from leaves of *Murraya euchres-*
tifolia. Yellow needles (Et₂O). Mp 187-
189°. λ_{max} 231 (log ε 4.53); 255 (sh) (log ε
3.85); 284 (log ε 4.15); 291 (log ε 4.16); 330
(sh) (log ε 3.52); 386 (log ε 3.8) (MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 785-
789 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Euchrestine A **E-275**
6-Methyl-1-(3-methyl-2-butenyl)-9H-carbazole-2,7-diol, 9CI. 2,7-Dihydroxy-6-methyl-1-prenylcarbazole
[138842-70-3]



C₁₈H₁₉NO₂ 281.354

Alkaloid from the stem bark of *Murraya*
euchrestifolia (Rutaceae). Oil. λ_{max} 213 ;
236 ; 265 ; 295 ; 313 ; 318 ; 329 (MeOH).

O²-*Me*: **7-Methoxy-3-methyl-8-(3-**
methyl-2-butenyl)-9H-carbazole-2-ol. 7-
Hydroxy-2-methoxy-6-methyl-1-prenyl-
carbazole. Isomurrayafoline B
[107903-15-1]


C₁₉H₂₁NO₂ 295.38

Alkaloid from the stem bark of *Mur-*
raya euchrestifolia (Rutaceae). Shows
cytotoxic activity. Prisms (Et₂O). Mp
158-161°. λ_{max} 213 ; 237 ; 264 ; 310 ;
330 (sh) (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**,
450-452; 1991, **39**, 1668-1671

Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**,
893-897 (*Isomurrayafoline B*, *activity*)

Euchrestine C **E-276**
1-Geranyl-2,7-dihydroxy-6-methylcarba-
zole. 1-(3,7-Dimethyl-2,6-octadienyl)-6-
methyl-9H-carbazole-2,7-diol
[138822-95-4]



$C_{23}H_{27}NO_2$ 349.472

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Powder.

2-Me ether: Euchrestine B

[138822-94-3]

$C_{24}H_{29}NO_2$ 363.499

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Pale yellow oil.

7-Me ether: Murrayanol†

[144525-81-5]

$C_{24}H_{29}NO_2$ 363.499

Alkaloid from seeds of *Murraya koenigii* (curry leaf tree) (Rutaceae). Cryst. (CH_2Cl_2 /petrol). Mp 161°.

2,3'-Epoxide: Euchrestine E

[139726-46-8]

$C_{23}H_{27}NO_3$ 365.471

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Pale brown oil. Obt. as a racemate.

1''-Oxo, 2-Me ether: 6-Formyl-1-geranyl-7-hydroxy-2-methoxycarbazole. Murrayaline D

[139726-44-6]

$C_{24}H_{27}NO_3$ 377.482

Alkaloid from the stem bark of *Murraya euchrestifolia*. Pale brown oil. λ_{max} 243 ; 255 (sh) ; 287 ; 330 ; 357 ; 364 ; 368 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1668-1671 (*isol, pmr, cmr, struct*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 2525-2528 (*Murrayaline D*)


Reisch, J. *et al.*, *Phytochemistry*, 1992, **31**, 2877-2879 (*Murrayanol*)

Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893-897 (*activity*)

Euchrestine D

E-277

1-Geranyl-2,7-dihydroxy-3-methylcarbazole. 1-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-9H-carbazole-2,7-diol, 9CI
[138822-96-5]



$C_{23}H_{27}NO_2$ 349.472

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Pale yellow oil. λ_{max} 212 ; 238 ; 265 ; 312 ; 321 (sh) (MeOH).

7-Deoxy (?): Mahanimbilol. Mahanimbinol

[77156-13-9]

$C_{23}H_{27}NO$ 333.472

Alkaloid from *Murraya koenigii* (curry leaf tree). Shows anti-HIV activity. Pale yellow oil. The struct. was first assigned to Mahanimbinol but Reisch *et al* assigned it to Mahanimbilol with

different spectroscopic props. The struct. of Mahanimbinol is therefore currently unresolved. λ_{max} 244 (log ϵ 4.26); 260 (log ϵ 4.14); 296 (log ϵ 3.98); 325 (log ϵ 3.4); 339 (log ϵ 3.38) (EtOH).

Rao, A.V.R. *et al.*, *Chem. Ind. (London)*, 1980, 697-698 (*Mahanimbinol*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1668-1671 (*isol, pmr, cmr, struct*)

Reisch, J. *et al.*, *Phytochemistry*, 1994, **36**, 1073-1076 (*Mahanimbilol*)


Meragelman, K.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 427 (*Mahanimbilol, activity*)

Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893-897 (*activity*)

Eudistalbin A

E-278

7-Bromo- α -(2-methylpropyl)-9H-pyrido[3,4-b]indole-1-methanamine, 9CI
[142755-07-5]



$R^1 = H, R^2 = NH_2$

$C_{16}H_{18}BrN_3$ 332.242

Alkaloid from the marine tunicate *Eudistoma album*. Exhibits cytotoxicity *in vitro* against KB human buccal carcinoma. Amorph. Sol. MeOH. $[\alpha]_D^{25}$ -10 (c, 0.1 in MeOH). λ_{max} 220 (sh) ; 240 (ϵ 45400); 290 (sh) ; 295 (ϵ 23200); 338 (ϵ 5790); 350 (ϵ 5790) (MeOH) (Derep). λ_{max} 240 (ϵ 45386); 295 (ϵ 23157); 338 (ϵ 5789); 350 (ϵ 5789) (MeOH) (Berdy).

Adesanya, S.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 525-527 (*isol, uv, ir, pmr, cmr, ms, struct*)

Eudistalbin B

E-279

1-(7-Bromo-9H-pyrido[3,4-b]indol-1-yl)-3-methyl-1-butanone, 9CI
[142755-08-6]

As Eudistalbin A, E-278 with $R^1R^2 = O$

$C_{16}H_{15}BrN_2O$ 331.211


Alkaloid from the marine tunicate *Eudistoma album*. Amorph. Sol. MeOH. λ_{max} 210 (sh) ; 224 (ϵ 28400); 245 (ϵ 11700); 285 (ϵ 14700); 310 (ϵ 8120); 376 (ϵ 4730) (MeOH) (Derep).

Adesanya, S.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 525-527 (*isol, uv, ir, pmr, ms, struct*)

Eudistomidin A

E-280

6-Bromo-1-(3,4-dihydro-2H-pyrrol-5-yl)-9H-pyrido[3,4-b]indol-8-ol, 9CI
[102673-53-0]



$C_{15}H_{12}BrN_3O$ 330.183

Alkaloid from the Okinawan tunicate *Eudistoma glaucus*. Calmodulin antagonist. Yellow solid. Mp 225-230° dec Mp 260-270° dec Mp 265-280° dec. λ_{max} 222 (ϵ 33000); 254 (ϵ 17000); 371 (ϵ 5500) (MeOH) (Derep).


Kobayashi, J. *et al.*, *Tet. Lett.*, 1986, **27**, 1191-1194 (*isol, uv, ir, pmr, cmr, ms, struct*)

Murakami, Y. *et al.*, *Tet. Lett.*, 1989, **30**, 2099-2100 (*synth*)

Eudistomidin B

E-281

6-Bromo-2,3,4,9-tetrahydro-2-methyl- α -(4-methylphenyl)methyl]-1H-pyrido[3,4-b]indole-1-methanamine, 9CI
[125443-66-5]



$C_{21}H_{24}BrN_3$ 398.345


Alkaloid from the marine tunicate *Eudistoma glaucus*. Cytotoxic. Activates rabbit heart muscle actomyosin ATPase. Yellow foam. Mp 81-83°. $[\alpha]_D^{25}$ -54 (c, 0.2 in MeOH). $[\alpha]_D$ -76.4 (c, 0.3 in $CHCl_3$). λ_{max} 231 (ϵ 26900); 291 (ϵ 6300) (MeOH) (Derep).

Kobayashi, J. *et al.*, *J.O.C.*, 1990, **55**, 3666-3670 (*isol, pmr, cmr, struct*)

Eudistomidin C

E-282

5-Bromo-1-[1-(methylamino)-2-(methylthio)ethyl]-9H-pyrido[3,4-b]indol-6-ol, 9CI
[125422-17-5]



Absolute Configuration

$C_{15}H_{16}BrN_3OS$ 366.281

Alkaloid from the marine tunicate *Eudistoma glaucus*. Cytotoxic. Exhibits calmodulin antagonistic activity. Yellow solid. Mp 120-122°. $[\alpha]_D^{25}$ +15.6 (c, 0.2 in MeOH). λ_{max} 218 (ϵ 21300); 236 (ϵ 23700); 253 (sh); 287 (sh); 298 (ϵ 13100); 368 (ϵ 3800) (MeOH) (Derep).

N¹⁴-Me: N¹⁴-Methyleudistomidin C

[380614-19-7]

$C_{16}H_{18}BrN_3OS$ 380.308


Alkaloid from *Eudistoma gilboverde*. Cytotoxic. Yellowish gum. $[\alpha]_D$ +12.9 (c, 0.07 in MeOH). λ_{max} 216 (log ϵ 4.28); 234 (log ϵ 4.16); 252 (log ϵ 4.06); 287 (sh) (log ϵ 3.77); 298 (log ϵ 3.93); 369 (log ϵ 3.22) (EtOH).

Kobayashi, J. *et al.*, *J.O.C.*, 1990, **55**, 3666-3670 (*isol, pmr, cmr, struct*)

Rashid, M.A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1454-1456 (*N*¹⁴-Methyleudistomidin C)

Eudistomidin F E-283

7-Bromo-2,3-dihydro-2-methyl-3-[(methylthio)methyl]-1H-2,4,10b-triazafuranthen-8-ol, 9CI
[136094-36-5]



C₁₆H₁₆BrN₃OS 378.292
Alkaloid from the marine tunicate *Eudistoma glaucus*. Amorph. [α]_D²⁶ +16 (c, 0.1 in MeOH). Stereochem. not confirmed. λ_{\max} 220 (€ 19000); 237 (€ 18000); 300 (€ 11000) (MeOH) (Derep). λ_{\max} 218 (€ 18000); 237 (€ 17000); 300 (€ 10000) (MeOH) (Berdy).

S-Oxide: Eudistomidin E

[136094-35-4]


C₁₆H₁₆BrN₃O₂S 394.291

Alkaloid from the marine tunicate *Eudistoma glaucus*. Amorph. [α]_D²⁴ +19 (c, 0.1 in MeOH). λ_{\max} 220 (€ 19000); 237 (€ 18000); 300 (€ 11000) (MeOH) (Derep).

Murata, O. *et al.*, *Tet. Lett.*, 1991, **32**, 3539-3542 (*isol, pmr, cmr, struct*)

Eudistomin X E-284

1-(1-Dimethylamino-2-phenylethyl)-9H-pyrido[3,4-b]indol-6-ol



C₂₁H₂₁N₃O 331.416


(ξ)-form [492463-46-4]

Alkaloid from the marine tunicate *Eudistoma* sp. Amorph. brown powder. [α]_D²⁰ -7 (c, 0.49 in MeOH). λ_{\max} 201 ; 234 ; 251 ; 298 ; 373 (MeOH).

Schupp, P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 272-275 (*isol, pmr, cmr, ms*)

Eudistomin A E-285

7-Bromo-1-(1H-pyrrol-2-yl)-9H-pyrido[3,4-b]indol-6-ol, 9CI. 7-Bromo-6-hydroxy-1-(2-pyrrolyl)- β -carboline
[88704-36-3]



C₁₅H₁₀BrN₃O 328.167

Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Mod. active against viruses, gram-positive bacteria and yeasts. Yellow oil. λ_{\max} 210 (€ 17600); 229 (sh) ; 258 (€ 13600); 272 (sh) ; 285 (sh) ; 326 (€ 11600); 373 (€ 6400); 389 (sh) (MeOH) (Derep).

Debromo: 1-(1H-Pyrrol-2-yl)-9H-pyrido[3,4-b]indol-6-ol, 9CI. 6-Hydroxy-1-(2-pyrrolyl)- β -carboline. Eudistomin M
[88704-39-6]

C₁₅H₁₁N₃O 249.271

Alkaloid from the tunicate *Eudistoma olivaceum*. Antiviral agent. ATPase stimulant used as biochem. tool in study of muscular contraction. Yellow prisms (CHCl₃/MeOH). Mp 225-227° (synthetic). λ_{\max} 210 (€ 24000); 231 (€ 16500); 253 (€ 18000); 266 (sh) (€ 12700); 276 (sh) (€ 9900); 289 (sh) (€ 6800); 321 (€ 17100); 371 (€ 9900); 384 (€ 9800) (MeOH) (Derep).

A'-Isomer, 3',4'-dihydro: 7-Bromo-1-(3,4-dihydro-2H-pyrrol-5-yl)-9H-pyrido[3,4-b]indol-6-ol, 9CI. *Eudistomin P*
[88704-48-7]

C₁₅H₁₂BrN₃O 330.183

Found in *Eudistoma olivaceum*. Modestly active against gram-positive bacteria and yeasts. Sol. MeOH, CHCl₃, toluene; poorly sol. H₂O. Mp 128-130°. λ_{\max} 201 (sh) ; 216 (€ 28100); 249 (€ 11800); 300 (€ 13400); 318 (sh) ; 372 (€ 4600); 381 (€ 4800) (MeOH) (Derep). λ_{\max} 216 (€ 28100); 249 (€ 11800); 300 (€ 10700); 372 (€ 4600); 381 (€ 4800) (hexane) (Berdy).

A'-Isomer, 3',4'-dihydro, deoxy: 7-Bromo-1-(3,4-dihydro-2H-pyrrol-5-yl)-9H-pyrido[3,4-b]indole, 9CI. *Eudistomin G*
[88704-43-2]

C₁₅H₁₂BrN₃ 314.184

Found in *Eudistoma olivaceum*. Antiviral agent. Needles (CH₂Cl₂). Sol. MeOH, toluene, CHCl₃; poorly sol. H₂O. Mp 204-206°. λ_{\max} 220 (€ 36900); 238 (€ 17600); 284 (€ 18200); 297 (sh) (€ 10400); 370 (€ 7900) (MeOH) (Derep). λ_{\max} 203 (€ 19000); 219 (€ 21500); 250 (€ 10000); 282 (€ 11700); 299 (€ 8700); 307 (€ 8400); 346 (€ 5100); 362 (€ 7600) (hexane) (Berdy).

A'-Isomer, 3',4'-dihydro, debromo: 6-Hydroxy-1-(1-pyrrolin-2-yl)- β -carboline.

Eudistomin Q

[88704-49-8]

C₁₅H₁₃N₃O 251.287

Found in *Eudistoma olivaceum*. Modestly active against gram-positive bacteria. Sol. MeOH, CHCl₃, toluene; poorly sol. H₂O. Mp 120-125°. λ_{\max} 216 (€ 23500); 250 (€ 8700); 295 (€ 9600); 382 (€ 2900) (MeOH) (Derep). λ_{\max} 214 (€ 19900); 288 (€ 8000); 348 (€ 2500); 362 (€ 2500) (hexane) (Berdy).

Kobayashi, J. *et al.*, *J.A.C.S.*, 1984, **106**, 1526-1528 (*isol, pmr, struct*)

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**, 3378-3387 (*isol, struct, props*)

Hino, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2596-2600 (*Eudistomins M,P, synth*)


Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 7117-7120 (*Eudistomin M, synth*)

Ohizumi, Y. *et al.*, *J. Pharmacol. Exp. Ther.*, 1998, **285**, 381-389 (*Eudistomin M, use*)

Eudistomin B

E-286

[96426-92-5]



C₁₇H₁₆BrN₃O₂ 374.236


Alkaloid antibiotic. Found in the Caribbean tunicate *Eudistoma olivaceum*. Shows activity against viruses, bacteria and yeasts.

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**, 3378-3387 (*isol, struct, props*)

Eudistomin C

E-287

1-Amino-11-bromo-1,2,7,8,13,13b-hexahydro[1,6,2]oxathiazepino[2',3':1,2]pyrido[3,4-b]indol-10-ol, 9CI
[88704-50-1]



C₁₄H₁₆BrN₃O₂S 370.269

Stereochem. of N–O bond revised in 1987. CA numbering shown. Alkaloid from the Caribbean tunicate *Eudistoma olivaceum* and the New Zealand ascidian *Ritterella sigillinoides*. Potent antiviral agent. Pale-yellow oil. [α]_D²⁵ -52 (c, 0.4 in MeOH). Eudistomins C, E, F, K and L are the most potent of the group. λ_{\max} 226 (€ 23400); 287 (€ 8000) (MeOH) (Derep).

N-Methoxycarbonyl: *Eudistomin F*
[96426-93-6]

C₁₆H₁₈BrN₃O₄S 428.306

Found in *Eudistoma olivaceum*. Potent antiviral agent. Oil. Sol. MeOH, CHCl₃, toluene; poorly sol. H₂O, hexane. λ_{\max} 226 (€ 23400); 287 (€ 8000) (MeOH) (Derep). λ_{\max} 225 (€ 10800); 276 ; 287 (€ 4100) (MeOH) (Berdy).

Deoxy: Eudistomin K

[88704-52-3]

C₁₄H₁₆BrN₃OS 354.27

From *Eudistoma olivaceum* and from the New Zealand ascidian *Ritterella sigillinoides*. Potent antiviral agent. Oil. Sol. MeOH, toluene, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -102 (c, 0.2 in MeOH). λ_{\max} 230 (€ 11900); 237 (sh); 284 (€ 3470); 293; 301 (EtOH) (Derep). λ_{\max} 226 ; 229 (€ 17000); 284 (€ 4000); 287 (MeOH) (Berdy).

Deoxy, S-oxide (α -): Eudistomin K sulf-oxide

[115276-12-5]
 $C_{14}H_{16}BrN_3O_2S$ 370.269
 Isol. from *Ritterella sigillinoides*. Shows antiviral props. Light yellow oil. $[\alpha]_D^{25}$ -3.3 (c, 0.09 in MeOH). λ_{max} 230 (ϵ 11900); 237 (sh); 284 (ϵ 3470); 293; 301 (EtOH) (Derep). λ_{max} 229 (ϵ 11920); 284 (ϵ 3470) (MeOH) (Berdy).

Debromo, 9-bromo: Eudistomin E

[88704-51-2]
 $C_{14}H_{16}BrN_3O_2S$ 370.269
 Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Potent antiviral agent. Pale-yellow oil. Sol. MeOH, $CHCl_3$, toluene; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ -18 (c, 0.1 in MeOH). λ_{max} 226 (ϵ 23400); 287 (ϵ 8000) (MeOH) (Derep). λ_{max} 226 (ϵ 18600); 282 (ϵ 6700); 287 (MeOH) (Berdy).

Debromo, 9-bromo, 1-N-Me: N¹⁰-Methyleudistomin E. Eudistomin V†

[230636-55-2]
 $C_{15}H_{18}BrN_3O_2S$ 384.296
 Isol. from *Eudistoma olivaceum*. Not to be confused with Eudistomin V in E-288.

Deoxy, debromo: Debromoeudistomin K

[110597-53-0]
 $C_{14}H_{17}N_3OS$ 275.374
 From *Ritterella sigillinoides*. Sol. MeOH, EtOAc, Me_2CO , C_6H_6 ; poorly sol. H_2O . $[\alpha]_D^{25}$ -58.3 (c, 0.06 in MeOH). λ_{max} 223 (ϵ 11500); 272 (ϵ 2900); 285 (ϵ 17270); 375 (ϵ 15870) (MeOH) (Berdy).

Deoxy, debromo, 10-bromo: Eudistomin L

[88704-55-6]
 $C_{14}H_{16}BrN_3OS$ 354.27
 Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Potent antiviral agent. Sol. MeOH, $CHCl_3$, toluene; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ -77 (c, 0.2 in MeOH). Stereochem. of N–O bond revised in 1987. λ_{max} 230 (ϵ 11900); 237 (sh); 284 (ϵ 3470); 293; 301 (EtOH) (Derep). λ_{max} 226 (ϵ 26600); 288 (ϵ 6100) (MeOH) (Berdy).

Deoxy, debromo, 10-bromo, 1-N-Ac: N-Acyleudistomin L

[123492-40-0]
 $C_{16}H_{18}BrN_3O_2S$ 396.307
 Amorph. solid. Mp 193.5-195.5°.

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1984, **106**, 1524-1526; 1987, **109**, 3378-3387 (*isol, uv, pmr, struct, Eudistomins E,F,K*)

Eur. Pat., 1985, 133 000; *C.A.*, **102**, 226023 (*isol, props*)

Blunt, J.W. *et al.*, *Tet. Lett.*, 1987, **28**, 1825-1826 (*Eudistomins F,K, pmr, cmr, stereochem*)

Lake, R.J. *et al.*, *Tet. Lett.*, 1988, **29**, 2255-2256; 4971-4972 (*Eudistomin K S-oxide, Eudistomin K, cryst struct, abs config*)

Lake, R.J. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1201-1206 (*Debromoeudistomin K*)

Still, I.W.J. *et al.*, *Tet. Lett.*, 1989, **30**, 1041-1044 (*N-Acyleudistomin L, synth, pmr, cmr, ms*)

Liu, J.-J. *et al.*, *Heterocycles*, 1990, **31**, 229-231 (*Eudistomin F, synth*)

Ibanez-Calero, S. *et al.*, *CA*, 1999, **131**, 100027f (*N-Methyleudistomin E*)


Liu, J.-J. *et al.*, *J.C.S. Perkin 1*, 2000, 3477-3486; 3487-3494 (*Eudistomin C,E,F,K,L, synth*)

Yamashita, T. *et al.*, *J.A.C.S.*, 2005, **66**, 15038-15039 (*synth*)

Yamagishi, H. *et al.*, *Org. Lett.*, 2008, **10**, 2369-2372 (*Eudistomin C,E, synth*)

Eudistomin H**E-288**

6-Bromo-1-(3,4-dihydro-2H-pyrrol-5-yl)-9H-pyrido[3,4-b]indole, 9CI. 6-Bromo-1-(1-pyrrolin-2-yl)- β -carboline [88704-44-3]



$C_{15}H_{12}BrN_3$ 314.184

Alkaloid from the Caribbean tunicate *Eudistoma olivaceum*. Shows mod. activity against yeasts and viruses. Powder. Sol. MeOH, $CHCl_3$, toluene; poorly sol. H_2O . Mp 140-142°. λ_{max} 220 (ϵ 36900); 238 (ϵ 17600); 284 (ϵ 18200); 297 (sh) (ϵ 10400); 370 (ϵ 7900) (MeOH) (Derep). λ_{max} 221 (ϵ 18300); 286 (ϵ 8300); 371 (ϵ 3000) (MeOH) (Berdy). λ_{max} 203 (ϵ 24700); 221 (ϵ 34200); 250 (ϵ 12400); 258 (ϵ 10000); 286 (ϵ 17000); 306 (ϵ 9400); 369 (ϵ 8400) (hexane) (Berdy).

Debromo: Eudistomin I. 1-(3,4-Dihydro-2H-pyrrol-5-yl)-9H-pyrido[3,4-b]indole, 9CI. 1-(1-Pyrrolin-2-yl)- β -carboline [88704-45-4]

$C_{15}H_{13}N_3$ 235.288

Found in *Eudistoma olivaceum*. Modestly active against gram-positive bacteria and viruses. Powder. Sol. MeOH, toluene, $CHCl_3$; poorly sol. H_2O . Mp 153-155°. λ_{max} 227 (ϵ 14500); 282 (ϵ 11300); 368 (ϵ 5700) (CH_2Cl_2) (Derep). λ_{max} 216 (ϵ 24400); 247 (ϵ 3500); 277 (ϵ 14100); 298 (ϵ 7800); 366 (ϵ 9200) (MeOH) (Derep). λ_{max} 215 (ϵ 13800); 238 (ϵ 8500); 280 (ϵ 7000); 368 (ϵ 2800) (MeOH) (Berdy). λ_{max} 217 (ϵ 23800); 249 (ϵ 6500); 256 (ϵ 6000); 279 (ϵ 10700); 293 (ϵ 4700); 365 (ϵ 7000) (hexane) (Berdy).

7-Bromo: Eudistomin V†

[211427-62-2]

$C_{15}H_{11}Br_2N_3$ 393.08

Alkaloid from the tunicate *Pseudodistoma aureum*. Yellow gum. λ_{max} 224 (ϵ 30000); 290 (ϵ 16000); 311 (ϵ 11900); 371 (ϵ 6300) (MeOH).

Kobayashi, J. *et al.*, *J.A.C.S.*, 1984, **106**, 1526-1528 (*Eudistomin I, isol, struct*)

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1987, **109**, 3378-3387 (*isol, struct, props, rev*)

Hino, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2596-2600 (*Eudistomins H and I, synth*)

Van Wagenen, B.C. *et al.*, *Tet. Lett.*, 1989, **30**, 3605-3608 (*Eudistomin I, synth*)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 7117-7120 (*Eudistomin I, synth*)


McNulty, J. *et al.*, *J.C.S. Perkin 1*, 1994, 1329-1337 (*synth*)

Shen, G.Q. *et al.*, *Tet. Lett.*, 1994, **35**, 1141-1144; 4923-4926 (*biosynth*)

Davis, R.A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 959-960 (*Eudistomin V*)

Eudistomin T**E-289**

2-Phenyl-1-(9H-pyrido[3,4-b]indol-1-yl)ethanone, 9CI. 1-Phenylacetyl- β -carboline [108335-05-3]



$C_{19}H_{14}N_2O$ 286.332

Alkaloid from the marine tunicate *Eudistoma olivaceum*. Yellow cryst. (MeOH). Mp 160-161°. λ_{max} 227 (ϵ 24100); 280 (ϵ 56100); 380 (ϵ 7000) (no solvent reported) (Derep). λ_{max} 227 (ϵ 39900); 292 (ϵ 21400); 379 (ϵ 7600) (CH_2Cl_2) (Derep).

6-Bromo: Eudistomin S

[108335-04-2]

$C_{19}H_{13}BrN_2O$ 365.228

Alkaloid from *Eudistoma olivaceum*. Yellow cryst. (MeOH). Mp 168°. λ_{max} 227 (ϵ 24100); 280 (ϵ 56100); 380 (ϵ 7000) (no solvent reported) (Derep). λ_{max} 227 (ϵ 39900); 292 (ϵ 21400); 379 (ϵ 7600) (CH_2Cl_2) (Derep).

7-Bromo: Eudistomin R

[108335-03-1]

$C_{19}H_{13}BrN_2O$ 365.228

Alkaloid from *Eudistoma olivaceum*. λ_{max} 227 (ϵ 24100); 280 (ϵ 56100); 380 (ϵ 7000) (no solvent reported) (Derep). λ_{max} 227 (ϵ 39900); 292 (ϵ 21400); 379 (ϵ 7600) (CH_2Cl_2) (Derep).

Kinzer, K.F. *et al.*, *Tet. Lett.*, 1987, **28**, 925-926 (*uv, ir, pmr, ms, struct*)

Still, I.W.J. *et al.*, *Heterocycles*, 1989, **29**, 2057 (*synth, cmr, Eudistomins S and T*)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 7117 (*synth*)

Bracher, F. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1994, **327**, 121 (*synth*)


McNulty, J. *et al.*, *J.C.S. Perkin 1*, 1994, 1329 (*synth*)

Rocca, P. *et al.*, *Synth. Commun.*, 1995, **25**, 3373 (*synth*)

Molina, P. *et al.*, *Tet. Lett.*, 1996, **37**, 9353 (*synth, Eudistomin S, Eudistomin T*)

Eudistomin W**E-290**

1-(1-Hydroxy-2-phenylethyl)-9H-pyrido[3,4-b]indol-6-ol. 6-Hydroxy- α -phenylmethyl-9H-pyrido[3,4-b]indole-1-methanol, 9CI



$C_{19}H_{16}N_2O_2$ 304.348

(S)-form [492463-95-3]


Alkaloid from the marine tunicate *Eudistoma* sp. Amorph. brown powder. $[\alpha]_D^{20}$

-23 (c, 0.24 in MeOH). λ_{\max} 208 ; 234 ; 266 ; 308 ; 405 (MeOH).

Schupp, P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 272-275 (*isol*, *pmr*, *cmr*, *ms*)

Eudistone A

[135340-00-0]




$C_{27}H_{19}N_5O$ 429.48

Alkaloid from the tunicate *Eudistoma* sp. Amorph. yellow powder. λ_{\max} 210 (ϵ 44800); 238 (ϵ 48500); 260 (sh) (ϵ 24700); 323 (ϵ 15300); 338 (ϵ 15500); 359 (sh) (ϵ 13100); 395 (ϵ 10600) (MeOH) (Derep).

He, H.-Y. *et al.*, *J.O.C.*, 1991, **56**, 5369-5371 (*isol*, *uv*, *ir*, *cd*, *pmr*, *cmr*, *struct*)

Eudistone B

[135340-01-1]




$C_{27}H_{17}N_5O$ 427.464

Alkaloid from the tunicate *Eudistoma* sp. Amorph. powder. $[\alpha]_D^{25}$ -177.8 (c, 0.036 in MeOH). λ_{\max} 204 (ϵ 45700); 239 (ϵ 47400); 259 (ϵ 39600); 324 (ϵ 14700) (MeOH) (Derep).

He, H.-Y. *et al.*, *J.O.C.*, 1991, **56**, 5369-5371 (*isol*, *uv*, *ir*, *cd*, *pmr*, *cmr*, *struct*)

Eugenine A**E-293**

2,3,4,7-Tetrahydro-5-methoxy-2,2,7-trimethylpyrano[2,3-f][2,3]benzoxazin-3-ol [913329-95-0]



Absolute Configuration


$C_{15}H_{19}NO_4$ 277.319

Alkaloid from the aerial parts of *Eugenia caryophyllus*.

Abdel-Kader, S.M. *et al.*, *CA*, 2006, **145**, 451680r (*isol*)

Eugenine B**E-294**

3,6-Dihydro-4-methoxy- $\alpha,\alpha,6$ -trimethyl-2H-furo[2,3-f][2,3]benzoxazine-2-methanol [913329-96-1]



Absolute Configuration


$C_{15}H_{19}NO_4$ 277.319

Alkaloid from the aerial parts of *Eugenia caryophyllus*.

Abdel-Kader, S.M. *et al.*, *CA*, 2006, **145**, 451680r (*isol*)

Euphosalicin**E-295**

[330937-97-8]




$C_{42}H_{48}N_2O_{15}$ 820.846

Constit. of *Euphorbia salicifolia*. Amorph. solid. $[\alpha]_D^{25}$ -101 (c, 0.12 in $CHCl_3$). λ_{\max} 219 (log ϵ 3.94); 264 (log ϵ 3.38) (MeOH).

Hohmann, J. *et al.*, *Tetrahedron*, 2001, **57**, 211-215 (*isol*, *pmr*, *cmr*)

Euphosine**E-296**

6,7-Dihydro-7-hydroxy-7-methyl-5H-2-pyridine-4-carboxaldehyde, 9CI [126632-23-3]



Absolute Configuration


$C_{10}H_{11}NO_2$ 177.202

Alkaloid from *Orthocarpus luteus*. Pale yellow solid. $[\alpha]_D^{23}$ +3.3 (c, 0.21 in $CHCl_3$). λ_{\max} 238 (log ϵ 3.4); 266 (log ϵ 2.9); 281 (log ϵ 2.8) (EtOH).

Boros, C.A. *et al.*, *J. Nat. Prod.*, 1990, **53**, 72-80 (*isol*, *uv*, *pmr*, *cmr*, *ms*, *struct*)

Eupolauramine**E-297**

6-Methoxy-5-methylbenzo[h]pyrrolo[4,3,2-de]quinolin-4(5H)-one, 9CI. EL Base 2 [58856-98-7]



$C_{16}H_{12}N_2O_2$ 264.283

Alkaloid from the bark of *Eupomatia laurina* (Eupomatiaceae). Golden-yellow prisms (EtOH). Mp 189-190°.

O-De-Me:

Orange needles ($CHCl_3/Et_2O$). Mp 290° dec.

10-Hydroxy: **Hydroxyeupolauramine. EL Base 3**

[58856-99-8]

$C_{16}H_{12}N_2O_3$ 280.282

Alkaloid from the bark of *Eupomatia laurina* (Eupomatiaceae). Fine orange needles (EtOH). Mp 282-283°.

10-Hydroxy, Me ether:

Orange-yellow prisms (EtOH). Mp 213-215°.

Bowden, B.F. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2659; 1975, **28**, 2681 (*isol*, *uv*, *ir*, *pmr*, *ms*)

Bowden, B.F. *et al.*, *J.C.S. Perkin 2*, 1976, 658 (*cryst struct*)

Levin, J.I. *et al.*, *J.A.C.S.*, 1983, **105**, 1397

(*synth*)

Taylor, W.C. *et al.*, *Aust. J. Chem.*, 1984, **37**,

1095-1104 (*biosynth*,

Hydroxyeupolauramine)

Karuso, P. *et al.*, *Aust. J. Chem.*, 1984, **37**, 1271

(*synth*)

Murakami, Y. *et al.*, *Chem. Pharm. Bull.*, 1988,

36, 3732 (*synth*)

Kikugawa, Y. *et al.*, *Tet. Lett.*, 1988, **29**, 4297

(*synth*)

Wang, X. *et al.*, *Tet. Lett.*, 1991, **32**, 4883

(*synth*)

Makosza, M. *et al.*, *Heterocycles*, 1992, **33**, 585

(*synth*)

Goehring, R.R. *et al.*, *Tet. Lett.*, 1992, **33**,

6045 (*synth*)

Hoarau, C. *et al.*, *J.O.C.*, 2001, **66**, 8064-8069

(*synth*)

Euroline**E-298**


[11085-46-4]

Struct. unknown. Alkaloid from the roots of *Euonymus europaeus* (Celastraceae). Powder. Mp 66-68°.

Bishay, D.W. *et al.*, *Herba Pol.*, 1971, **17**, 97-109; *CA*, **75**, 115894k

Europine**E-299**

[570-19-4]



$C_{16}H_{27}NO_6$ 329.392

Ester of Heliotridine in T-188 with Lasiocarpic acid. Alkaloid from *Heliotropium* spp. (Boraginaceae). Shows antifungal and insect antifeedant activity. Gum or glass. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{25}$ +10.9 (c, 2.5 in EtOH). Abs. config. shown follows from that detd. crystallographically for Lasiocarpine, L-47.

► Highly toxic.

N-Oxide: **Europine N-oxide**

[65582-53-8]

$C_{16}H_{27}NO_7$ 345.392

Alkaloid from *Heliotropium*

maris-mortui and *Heliotropium rotundifolium* (Boraginaceae). Shows anti-tumour activity. Mp 171-172°. $[\alpha]_D^{25} +27$.

O⁷-Ac: 7-Acetyლეუპინე

[123520-99-0]

C₁₈H₂₉NO₇ 371.43

Alkaloid from aerial parts of *Heliotropium bovei* (Boraginaceae). Gum. $[\alpha]_D^{25} +5.9$ (c, 0.67 in EtOH).

5'-O-Ac: 5'-Acetyლეუპინე

[123520-98-9]

C₁₈H₂₉NO₇ 371.43

Alkaloid from *Heliotropium rotundifolium* (Boraginaceae). Gum. $[\alpha]_D^{25} +27.2$ (c, 1.58 in CHCl₃).

O⁷-Angeloyl: see Lasiocarpine, L-47**7-Deoxy: Ilamine**C₁₆H₂₇NO₅ 313.393


Alkaloid from *Heliotropium crassifolium*. Light yellow oil. $[\alpha]_D^{25} +33.3$ (c, 0.2 in EtOH).

7-Deoxy, N-oxide: Ilamine N-oxideC₁₆H₂₇NO₆ 329.392

Alkaloid from *Heliotropium crassifolium*. Light yellow oil. $[\alpha]_D^{25} +20$ (c, 0.1 in EtOH).

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1954, **7**, 287 (*isol, struct*)Crowley, H.C. *et al.*, *CA*, 1957, **51**, 3925 (*isol*)Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1965, **18**, 1625 (*pmr*)Zalkow, L.H. *et al.*, *Phytochemistry*, 1978, **17**, 172 (*oxide*)Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)Asibal, C.F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 726 (*isol, ir, pmr, cmr, ms, struct, 5'-Acetyლეუპინე*)Reina, M. *et al.*, *Phytochemistry*, 1995, **38**, 355 (*7-Acetyლეუპინე*)Farsam, H. *et al.*, *Planta Med.*, 2000, **66**, 389-391 (*Ilamine, Ilamine oxide*)**Eustifoline A****E-300**

3,7-Dihydro-3,3,10-trimethylpyrano[2,3-c]carbazole, 9CI. 5,6-Pyranoglycozoline. Glycomaurin [119157-65-2]

C₁₈H₁₇NO 263.338

Alkaloid from *Glycosmis mauritiana*, *Micromelum falcatum* and *Murraya euchrestifolia* (Rutaceae). Green-yellow needles (CHCl₃/petrol). Mp 195-196° (178°).

N-Me: [125287-16-3]

Pale green-yellow needles (EtOH). Mp 130-131°.

2-Hydroxy: 2-Hydroxyeustifoline A. Glycoborinine

[233279-39-5]

C₁₈H₁₇NO₂ 279.338

Alkaloid from *Glycosmis arborea*.

Cryst. (C₆H₆/MeOH). Mp 220-221°. λ_{max} 262 ; 269 ; 293 ; 335 (MeOH).

2-Methoxy: 2-Methoxyeustifoline A. 2-Methoxyglycomaurin

[220789-48-0]

C₁₉H₁₉NO₂ 293.365

Alkaloid from *Glycosmis rupestris*. Needles. Mp 225-227°. λ_{max} 235 (log ϵ 0.54); 240 (log ϵ 0.48); 256 (log ϵ 5); 293 (log ϵ 0.58); 334 (0.79) (no solvent reported).

Kong, Y.C. *et al.*, *Biochem. Syst. Ecol.*, 1988, **16**, 485-489 (*isol, uv, pmr, ms, struct*)Kumar, V. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1375-1379 (*isol, uv, ir, pmr, ms, synth, struct*)Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1548-1550 (*isol, pmr, cmr, struct*)Rahmani, M. *et al.*, *Planta Med.*, 1998, **64**, 780 (*2-Methoxyeustifoline A*)Chakravarty, A.K. *et al.*, *Phytochemistry*, 1999, **50**, 1263-1266 (*Glycoborinine*)Lebold, T.P. *et al.*, *Org. Lett.*, 2007, **9**, 1883-1886 (*synth*)**Eustifoline B****E-301**

3,7-Dihydro-3,10-dimethyl-3-(4-methyl-3-pentenyl)pyrano[2,3-c]carbazole, 9CI [133740-40-6]


As Eustifoline A, E-300 with

R = -CH₂CH₂CH=C(CH₃)₂C₂₃H₂₅NO 331.457

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1548 (*isol, pmr, cmr, struct*)Lebold, T.P. *et al.*, *Org. Lett.*, 2007, **9**, 1883-1886 (*synth*)**Eustifoline C****E-302**


4-(3,7-Dimethyl-2,6-octadienyl)-6-methyl-9H-carbazol-3-ol, 9CI. 4-Geranyl-3-hydroxy-6-methylcarbazole [133740-41-7]

C₂₃H₂₇NO 333.472

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Amorph. powder.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1548 (*isol, uv, ir, pmr, ms, struct*)Lebold, T.P. *et al.*, *Org. Lett.*, 2007, **9**, 1883-1886 (*synth*)**Eusynstyelamide****E-303**


[157878-21-2]

C₃₂H₄₂Br₂N₁₀O₅ 806.556

Highly modified tryptophan-arginine dipeptide dimer. Metab. from the ascidian *Eusynstyela misakiensis*.

Swersey, J.C. *et al.*, *J. Nat. Prod.*, 1994, **57**, 842 (*isol, uv, ir, pmr, cmr, ms, struct*)**Euthyroideone A****E-304**

[219566-39-9]

C₁₂H₁₃BrN₂O₃S 345.216

Alkaloid from the bryozoan *Euthyroides episcopalis*. Mp 245°. λ_{max} 202 (log ϵ 3.96); 242 (log ϵ 3.72); 383 (log ϵ 3.6); 452 (log ϵ 3.86) (MeOH).

2,3-Didehydro: Euthyroideone B

[219566-38-8]

C₁₂H₁₁BrN₂O₃S 343.201

Alkaloid from *Euthyroides episcopalis*. Mp 244°. λ_{max} 208 (log ϵ 4.5); 310 (log ϵ 4.38); 436 (log ϵ 4.6) (MeOH).

7,8-Didehydro: Euthyroideone C


[219566-40-2]

C₁₂H₁₁BrN₂O₃S 343.201

Alkaloid from *Euthyroides episcopalis*. λ_{max} 203 (log ϵ 3.6); 280 (log ϵ 3.63); 391 (log ϵ 3.32) (MeOH).

Morris, B.D. *et al.*, *J.O.C.*, 1998, **63**, 9545-9547**Euxylophoricine A****E-305**

8,13-Dihydro-2,3-dimethoxyindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI [20999-50-2]

R¹ = R² = Me, R³ = HC₂₀H₁₇N₃O₃ 347.373

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Needles (CHCl₃/MeOH). Mp 296-298°.

7,8-Didehydro: Euxylophoricine B. 2,3-Dimethoxyindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(13H)-one, 9CI

[20999-51-3]

C₂₀H₁₅N₃O₃ 345.357

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Yellow prisms (CHCl₃/MeOH). Mp 310-312°.

- Canonica, L. *et al.*, *Tet. Lett.*, 1968, 4865 (*uv, ir, pmr, struct*)
 Danieli, B. *et al.*, *Phytochemistry*, 1972, **11**, 1833 (*synth*)
 Danieli, B. *et al.*, *Gazz. Chim. Ital.*, 1975, **105**, 45 (*synth*)
 Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 2347 (*synth*)
 Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth, cmr*)
 Hamid, A. *et al.*, *Tet. Lett.*, 2006, **47**, 1777-1781 (*synth*)

Euxylophoricine C E-306

5,15-Dihydro[1,3]dioxolo[4,5-g]indolo[2',3':3,4]pyrido[2,1-b]quinazolin-8(6H)-one, 9CI

[38990-11-3]

As Euxylophoricine A, E-305 with $R^1R^2 = -CH_2-$, $R^3 = H$

$C_{19}H_{13}N_3O_3$ 331.33

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Yellowish cryst. (EtOH). Mp 310-312°.

- Danieli, B. *et al.*, *Phytochemistry*, 1972, **11**, 1833 (*isol, uv, ir, pmr, ms, struct, synth*)
 Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 2347 (*synth*)
 Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth, ir*)

Euxylophoricine D E-307

8,13-Dihydro-2,3,10-trimethoxyindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI

[51059-70-2]

As Euxylophoricine A, E-305 with $R^1 = R^2 = Me$, $R^3 = OMe$

$C_{21}H_{19}N_3O_4$ 377.399

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Plates (CHCl₃). Mp 293-295°.

7,8-Didehydro: **Euxylophoricine E**, 2,3,10-Trimethoxyindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(13H)-one, 9CI

[51059-71-3]

$C_{21}H_{17}N_3O_4$ 375.383

Minor alkaloid from *Euxylophora paraënsis* bark (Rutaceae). Yellowish cryst. (CHCl₃). Mp 290°.

- Danieli, B. *et al.*, *Phytochemistry*, 1973, **12**, 2521 (*isol, uv, ir, pmr, ms, struct, synth*)
 Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth, ir, cmr*)

Euxylophoricine F E-308

8,13-Dihydro-2-hydroxy-3-methoxyindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI

[61017-57-0]

As Euxylophoricine A, E-305 with $R^1 = H$, $R^2 = Me$, $R^3 = H$

$C_{19}H_{15}N_3O_3$ 333.346


Alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Pale-yellow needles (C₆H₆/petrol). Mp 226°.

- Danieli, B. *et al.*, *Phytochemistry*, 1976, **15**, 1095 (*isol, uv, ir, ms, struct, synth*)

Euxylophorine

8,14-Dihydro-2,3-dimethoxy-14-methylindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI

[20999-49-9]



$C_{21}H_{19}N_3O_3$ 361.399

Alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Orange-red needles (anhyd. C₆H₆). Mp 227-230°.

7,8-Didehydro: **Euxylophorine B**, 2,3-Dimethoxy-14-methylindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(14H)-one, 9CI

[38990-12-4]

$C_{21}H_{17}N_3O_3$ 359.384

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Yellow-orange cryst. (CHCl₃). Mp 268-271° dec.

7,8-Didehydro; hydrochloride:

Yellow cryst. (MeOH + HCl). Mp 270-280° dec. Sublimation at 260-270°/0.01 mm gives Euxylophoricine B (q.v.).

10-Methoxy: **Euxylophorine C**, 5,7,8,13-Tetrahydro-2,3,10-trimethoxy-14-methyl-5-oxoindolo[2',3':3,4]pyrido[2,1-b]quinazolinium hydroxide inner salt, 9CI

[50816-62-1]

$C_{22}H_{21}N_3O_4$ 391.426

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Red needles (C₆H₆). Mp 207-209° dec.

10-Methoxy, 7,8-didehydro: **Euxylophorine D**

[51059-69-9]

$C_{22}H_{19}N_3O_4$ 389.41

Minor alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Yellow-orange needles (C₆H₆ or CHCl₃/diisopropyl ether). Mp 256-260° dec.


Canonica, L. *et al.*, *Tet. Lett.*, 1968, 4865 (*uv, ir, pmr, synth*)

Danieli, B. *et al.*, *Phytochemistry*, 1972, **11**, 1833 (*isol, uv, ir, pmr, ms, struct, synth, Euxylophorine B*)

Danieli, B. *et al.*, *Phytochemistry*, 1973, **12**, 2521 (*isol, uv, ir, pmr, ms, struct, synth, Euxylophorine C, Euxylophorine D*)

Evodiainine

8,13,13b,14-Tetrahydro-14-methylindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI. Rhetsine

**E-310**

$C_{19}H_{17}N_3O$ 303.363

Rhetsine was the racemate.

(S)-form [518-17-2]

A major component of the Chinese drug Wou-chou-yu (the dried fruit of *Evodia rutaecarpa*). Also present in *Araliopsis tabouensis* (Rutaceae, Araliaceae). Diuretic and diaphoretic agent. Shows bronchoconstrictive activity. Yellow leaflets (EtOH). Sol. Me₂CO; fairly sol. EtOH, CHCl₃, Et₂O; poorly sol. H₂O, C₆H₆, hexane. Mp 278°. $[\alpha]_D^{25} +352$ (Me₂CO). λ_{max} 272 ; 280 ; 291 ; 335 (MeOH) (Berdy).

7 β -Carboxy: 5,7,8,13,13b,14-Hexahydro-14-methyl-5-oxoindolo[2',3':3,4]pyrido[2,1-b]quinazolin-7-carboxylic acid, 9CI. 7-Carboxyevodiamine

[69754-32-1]

$C_{20}H_{17}N_3O_3$ 347.373

Minor alkaloid from the fruits of *Evodia rutaecarpa* (Rutaceae). Amorph. solid (as Me ester). $[\alpha]_D^{20} +441$ (c, 1 in CHCl₃) (Me ester).

(±)-form [518-18-3]

Alkaloid from the trunk bark of *Zanthoxylum rhetsa* (Rutaceae). Prisms (EtOH/EtOAc). Mp 275-277° (softens at 269°).

Chatterjee, A. *et al.*, *Tetrahedron*, 1959, **7**, 257-261 (*isol*)

Gopinath, K.W. *et al.*, *Tetrahedron*, 1960, **8**, 293-295 (*isol*)

Yamazaki, M. *et al.*, *Tet. Lett.*, 1966, **7**, 3221-3224; 1967, **8**, 3317-3320 (*biosynth*)

Tamás, J. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1976, **89**, 85-89 (*ms*)

Kametani, T. *et al.*, *J.A.C.S.*, 1976, **98**, 6186-6188 (*synth, ir, uv, pmr, ms*)

Danieli, B. *et al.*, *Heterocycles*, 1978, **9**, 803-806 (*synth*)

Danieli, B. *et al.*, *Experientia*, 1979, **35**, 156 (7-Carboxyevodiamine)

Danieli, B. *et al.*, *Chem. Comm.*, 1982, 1092-1093 (*synth, cd, abs config*)

Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246-1255 (*synth*)


Kobayashi, Y. *et al.*, *Planta Med.*, 2000, **66**, 526-530 (*activity*)

Nakayama, A. *et al.*, *Heterocycles*, 2008, **76**, 861-865 (*synth*)

Evodiainine**E-311**

9,14-Dihydro-14-methyl-5H-benzo[2,3][1,7]naphthyridino[8,7-b]quinazolin-5-one

[918476-26-3]



$C_{20}H_{15}N_3O$ 313.358

Alkaloid from *Evodia rutaecarpa*.

Wang, Q. *et al.*, *Zhongguo Yaoke Daxue Xuebao*, 2005, **36**, 520-522; *CA*, **146**, 118069r (*isol*)

Evoevoline E-312

Struct. unknown. Alkaloid from the leaves of *Euonymus europaeus* (Celastraceae). Needles (Et₂O). Mp 261-263°.

Bishay, D.W. *et al.*, *Herba Pol.*, 1971, **17**, 233-241

Evoline E-313


[11085-47-5]

Struct. unknown. Alkaloid from the stems and roots of *Euonymus europaeus* (Celastraceae). Mp 261-263°.

Bishay, D.W. *et al.*, *Herba Pol.*, 1971, **17**, 97-109; *CA*, **75**, 115894k

Evomeliaefolin E-314

3-(1-Hydroxy-3-oxobutyl)-4,7,8-trimethoxy-2(1H)-quinolinone, 9CI [168146-20-1]




C₁₆H₁₉NO₆ 321.329

Alkaloid from heartwood of *Tetradium glabrifolium* (*Evodia meliaefolia*) (Rutaceae). Needles (CHCl₃/MeOH). Mp 226-228°. [α]_D²⁰ -50 (c, 0.015 in MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1995, **40**, 121 (*isol, uv, ir, pmr, ms, struct*)

Evomerrine E-315

7-Hydroxy-4-methoxyfuro[2,3-b]quinoline-8-carboxaldehyde. 8-Formyl-7-hydroxy-4-methoxyfuro[2,3-b]quinoline. 8-Formyl-7-hydroxydictamine. 8-Formylconfusameline



C₁₃H₉NO₄ 243.218

Alkaloid from leaves of *Melicope semecarpifolia* (Rutaceae). Needles (CHCl₃/MeOH). Mp 219-220°. λ_{max} 267 (log ε 4.16); 273 (log ε 4.14); 332 (log ε 3.83) (EtOH).

Tsai, I.L. *et al.*, *Phytochemistry*, 1995, **40**, 1561-1562 (*isol*)

Evomine E-316

[12765-27-4]

Struct. unknown. Alkaloid from *Euonymus europaeus* (Celastraceae). Needles (Et₂O). Mp 232-235°.

Bishay, D.W. *et al.*, *Herba Pol.*, 1971, **17**, 233-241

Evopine E-317


[11085-48-6]

Struct. unknown. Alkaloid from the roots of *Euonymus europaeus* (Celastraceae). Mp 222-225°.

Bishay, D.W. *et al.*, *Herba Pol.*, 1971, **17**, 97-109; *CA*, **75**, 115894k

Excentricine E-318

[155416-29-8]



Absolute Configuration

C₁₈H₂₃NO₅ 333.383

Alkaloid from roots of *Stephania excentrica* (Menispermaceae). Needles (Me₂CO/MeOH). Mp 195-196°. [α]_D²² +261 (c, 0.03 in MeOH). λ_{max} 228 (log ε 4.31); 263 (log ε 4.02); 302 (log ε 3.23) (MeOH).

N-Me: N-Methylexcentricine

[186541-61-7]

C₁₉H₂₅NO₅ 347.41

Alkaloid from roots of *Stephania excentrica*. Needles (CHCl₃). Mp 215-217°. [α]_D¹⁸ +266 (c, 0.02 in MeOH). λ_{max} 228 (log ε 4.42); 264 (log ε 4.02); 305 (log ε 3.51) (MeOH).

14-Deoxy, 14,15-didehydro: **Stephalonganine A**

[908582-38-7]

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Stephania longa*. Amorph. powder. [α]_D²⁰ +157.5 (c, 0.12 in CHCl₃). λ_{max} 227 (log ε 4.38); 265 (log ε 4.03); 301 (log ε 3.46) (MeOH).

14-Deoxy, 14,15-didehydro, N-Me: **Stephalonganine B**

[908582-39-8]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Stephania longa*. Amorph. powder. [α]_D²⁰ +160 (c, 0.4 in CHCl₃).

1-Epimer, 14-deoxy, 14,15-didehydro: **Stephalonganine C**

[908582-40-1]

C₁₈H₂₁NO₄ 315.368


Alkaloid from *Stephania longa*. Amorph. powder. [α]_D²⁰ -155 (c, 0.08 in CHCl₃). λ_{max} 227 (log ε 4.3); 259 (log ε 4.03) (MeOH).

Deng, J.Z. *et al.*, *Nat. Prod. Lett.*, 1993, **2**, 283-268 (*isol, struct*)

Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 1996, **59**, 803-805 (*abs config*)

Deng, J.Z. *et al.*, *J. Nat. Prod.*, 1997, **60**, 294-295 (*N-Methylexcentricine*)

Zhang, H. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1105-1109 (*Stephalonganines A-C*)

Exiguamine A E-319

C₂₅H₂₆N₅O₆[⊕] 492.51

(±)-*form* [921223-01-0]

Constit. of *Neopetrosia exigua*. Deep red cryst. (MeOH) (as hydrochloride). λ_{max} 212 (log ε 3.44); 263 (log ε 3.08); 330 (log ε 2.79) (MeOH).

Brastianos, H.C. *et al.*, *J.A.C.S.*, 2006, **128**, 16046-16047 (*isol, pmr, cmr, cryst struct*)

Eximidine E-320

C₂₀H₂₃NO₄ 341.406


Struct. unknown. Possibly identical with Corydine, C-682. Alkaloid from *Dicentra eximia*. Mp 133°.

Manske, R.H.F. *et al.*, *Can. J. Res.*, 1933, **8**, 592-599; *CA*, **27**, 5148

Exochomine E-321

[142796-25-6]

[142754-96-9]




C₂₆H₃₆N₂O 392.583

Alkaloid from the ladybird *Exochomus quadripustulatus*. Cryst. + 1H₂O (as hydrochloride). Mp 243-248° (hydrochloride). [α]_D²⁰ +168 (c, 0.77 in CHCl₃).

Timmermans, M. *et al.*, *Tet. Lett.*, 1992, **33**, 1281 (*isol, pmr, cmr, cryst struct*)

Fabianine


[6871-51-8]

Relative
configurationC₁₄H₂₁NO 219.326Alkaloid from *Fabiana imbricata* (Solanaceae). Oil. Bp_{0.05} 74°. [α]_D 0.**Picrate:**

Needles (EtOAc). Mp 114-116°.

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1962, **40**, 256 (*isol, uv, pmr, ir, struct*)Sugita, T. *et al.*, *Heterocycles*, 1986, **24**, 29 (*synth, ir, pmr*)Schmed-Hirschmann, G. *et al.*, *Phytochemistry*, 1994, **36**, 1439 (*isol, pmr, cmr*)**Fagarine I**

5,7,8,15-Tetrahydro-3,4-dimethoxy-6-methylbenzo[e][1,3]dioxolo[4,5-k][3]benzazecin-14(6H)-one, 9CI. Thalictrimine†. α-Fagarine. Allocryptopine. γ-Homochelidonine. β-Homochelidonine
[24240-04-8]
[485-91-6]

C₂₁H₂₃NO₅ 369.416

Alkaloid from a wide variety of genera in the Papaveraceae (*Argemone*, *Bocconia*, *Corydalis*, *Dactylicapnos*, *Eschscholtzia*, *Glaucium*, *Hunnemannia*, *Hylomecon*, *Macleaya*, *Meconopsis*, *Papaver*, *Sanguinaria*, *Stylomecon*, *Chelidonium*), Ranunculaceae (*Thalictrum*), Rutaceae (*Fagara*, *Zanthoxylum*) and Sapindaceae (*Pteridophyllum*). Antihypertensive agent, cardiac inhibitor, respiratory stimulant, muscle relaxant. Cryst. (EtOH, Me₂CO, C₆H₆, EtOAc/petrol or CHCl₃/petrol). Poorly sol. hexane. Mp 160-161° (anhyd.). Log P 2.73 (uncertain value) (calc).

Hydrochloride:Cryst. + ½ H₂O. Mp 192-193°.**Picrate:**

Yellow plates (abs. EtOH). Mp 208-209°.

O⁹-De-Me: Hunnemannine

[490-52-8]

C₂₀H₂₁NO₅ 355.39Alkaloid from *Hunnemannia furmariaefolia* and *Argemone fruticosa* (Papaveraceae). Cryst. (CHCl₃/MeOH

F-1

or 2-propanol).

O¹⁰-De-Me: **Thalictrosine**. Alkaloid AA2. Alkaloid ED. Thalictrosine
[22047-92-3]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Thalictrum simplex*, *Thalictrum amurense*, *Argemone albiflora*, *Eschscholtzia californica* and *Eschscholtzia douglasii* (Ranunculaceae, Papaveraceae). Cryst. (MeOH). Mp 261-263°.

O¹⁰-De-Me, hydrochloride:Cryst. (H₂O). Mp 235-238°.

1-Methoxy: **Oreonone**. 1-Methoxyallocryptopine
[56743-52-3]

C₂₂H₂₅NO₆ 399.443Alkaloid from *Papaver curviscapum* (Papaveraceae). Mp 213° (131°, 125-129°).12-Methoxy: **12-Methoxyallocryptopine**C₂₂H₂₅NO₆ 399.443Alkaloid from *Arctomecon* spp. (Papaveraceae).Haworth, R.D. *et al.*, *J.C.S.*, 1926, 445

(synth)

Manske, R.H.F. *et al.*, *J.A.C.S.*, 1942, **64**, 1659 (*Hunnemannine*)Deulofeu, V. *et al.*, *J.O.C.*, 1947, **12**, 217

(struct)

Redemann, C.E. *et al.*, *J.A.C.S.*, 1949, **71**,1030 (*uv, struct*)Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*,1964, **29**, 2479 (*ms*)Ma, J.C.N. *et al.*, *Can. J. Chem.*, 1965, **43**, 1849

(pmr)

Slaviková, L. *et al.*, *Coll. Czech. Chem. Comm.*,1966, **31**, 1355 (*Hunnemannine, isol, uv, ir*)Hruban, L. *et al.*, *Coll. Czech. Chem. Comm.*,1967, **32**, 3414 (*uv*)Giacopello, D. *et al.*, *Tetrahedron*, 1967, **23**,3265 (*Hunnemannine, synth, uv, ir, pmr*)Maturová, M. *et al.*, *Planta Med.*, 1968, **17**,121 (*Oreonone*)Umarov, Kh.S. *et al.*, *Khim. Prir. Soedin.*,1970, **6**, 224; *Chem. Nat. Compd. (Engl. Transl.)*,1970, **6**, 219 (*Thalictrosine*)Haisková, K. *et al.*, *Coll. Czech. Chem. Comm.*,1973, **38**, 3312 (*Hunnemannine, isol, uv, ms*)Teitel, S. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 553


(synth)

Nakashima, T.T. *et al.*, *Org. Magn. Reson.*,1973, **5**, 9 (*cmr*)Vesely, Z. *et al.*, *Coll. Czech. Chem. Comm.*,1975, **40**, 1403 (*Oreonone, synth, uv, ir, ms*)Battersby, A.R. *et al.*, *J.C.S. Perkin I*, 1975,1147 (*biosynth*)Hanaoka, M. *et al.*, *Heterocycles*, 1976, **4**,1685 (*synth*)Iwasa, K. *et al.*, *J.O.C.*, 1982, **47**, 4275 (*pmr, cmr*)Sakai, T. *et al.*, *Acta Cryst. C*, 1988, **44**, 838

(cryst struct)

Sariyar, G. *et al.*, *Planta Med.*, 1989, **55**, 89(Oreonone, *isol, uv, ir, pmr, ms*)Raynie, D.E. *et al.*, *Biochem. Syst. Ecol.*, 1990,**18**, 45 (*12-Methoxyallocryptopine*)Marek, J. *et al.*, *Coll. Czech. Chem. Comm.*,1998, **63**, 416-424 (*cryst struct*)Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999,**37**, 195-202 (*N-15 nmr*)**Fagarine II**


5,7,8,15-Tetrahydro-2,3-dimethoxy-6-methylbenzo[e][1,3]dioxolo[4,5-k][3]benzazecin-14(6H)-one, 9CI
[528-67-6]

C₂₁H₂₃NO₅ 369.416Alkaloid from the leaves and twigs of *Fagara coco* (Rutaceae). Needles (EtOH, EtOAc or C₆H₆). Mp 200-202°.**Hydrochloride:**Cryst. + ½ H₂O (EtOH/Et₂O). Mp 200-202° dec., 215° dec. (rapid heating).**Picrate:**

Yellow needles. Mp 214°.

Redemann, C.E. *et al.*, *J.A.C.S.*, 1949, **71**, 1030 (*isol, uv*)Comin, J. *et al.*, *Tetrahedron*, 1959, **6**, 63 (*ir, struct*)Giacopello, D. *et al.*, *Tetrahedron*, 1964, **20**, 2971 (*synth, uv*)Orito, K. *et al.*, *Heterocycles*, 1980, **14**, 11 (*synth, pmr*)**Fagarine III**C₂₂H₂₆NO₄ 368.452Struct. unknown. Prob. a protopine alkaloid. Trace alkaloid from *Fagara coco* (Rutaceae). Prisms (EtOH). Mp 181-183°. [α]_D²⁷ -300 (CHCl₃).**Hydrochloride:** Mp 232-234° dec.Redemann, C.E. *et al.*, *J.A.C.S.*, 1949, **71**, 1030-1034 (*isol, uv*)**Fagaronine**

2-Hydroxy-3,8,9-trimethoxy-5-methylbenzo[c]phenanthridinium(1+), 9CI
[52259-65-1]

C₂₁H₂₀NO₄⁺ 350.393

Alkaloid from the roots of *Fagara xanthoxyloides* (Rutaceae) and from *Zanthoxylum spinosum*. Antineoplastic agent. Inhibitor of RNA reverse transcriptase activity in oncogenic viruses; very active antileukaemic agent. Shows anti-HIV activity. An intercalating agent. Shows bactericidal props. Yellow needles (EtOAc/MeOH)(as chloride). Mp 202° (198-200°) Mp 260-261° (double Mp)(chloride). Log P -0.06 (uncertain value) (calc). λ_{max} 233 (ε

F-3

F-4

F-5

19500); 272 (ε 35480); 328 (ε 27500) (MeOH) (Berdy). λ_{max} 346 (NaOH) (Berdy).

► DI9861000

Me ether, O-de-Me: **Punctatine**†

[65522-91-0]

C₂₁H₂₀NO₄ 350.393

Alkaloid from the stems and branches of *Zanthoxylum punctatum* (Rutaceae). Exact struct. not known.

Messmer, W.M. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 1858 (*ir, uv, ms, pmr, isol struct*)

Tin-Wa, M. *et al.*, *J. Pharm. Sci.*, 1974, **63**, 1476 (*struct*)

Gillespie, J.P. *et al.*, *J.O.C.*, 1974, **39**, 3239 (*synth*)

Sethi, V.S. *et al.*, *Biochem. Biophys. Res. Commun.*, 1975, **63**, 1070 (*pharmacol*)

Stermitz, F.R. *et al.*, *Phytochemistry*, 1977, **16**, 2003-2006 (*Punctatine*)

Phillips, S.D. *et al.*, *J. Het. Chem.*, 1981, **18**, 223 (*rev*)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2963 (*synth*)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2348 (*synth, uv, ir, pmr, ms*)

Ishii, H. *et al.*, *J.C.S. Perkin 1*, 1987, 671 (*synth, ir, pmr*)

Šmidrkal, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1988, **53**, 3184 (*synth, uv, pmr*)

Tan, G.T. *et al.*, *J. Nat. Prod.*, 1991, **54**, 143 (*anti-HIV activity*)

Barret, Y. *et al.*, *Phytother. Res.*, 1992, **6**, 59 (*rev*)

Lunch, M.A. *et al.*, *Bull. Soc. Chim. Fr.*, 1994, **131**, 718 (*synth*)

Seckárová, P. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 147-152 (*pmr, cmr*)


Fagopyrine

F-6

1,3,4,6,8,13-Hexahydroxy-10,11-dimethyl-2,5-di-2-piperidinylphenanthro[1,10,9,8-opqra]perylene-7,14-dione, 9CI

[72393-03-4]

[72393-47-6, 72393-46-5]



C₄₀H₃₄N₂O₈ 670.717


Photodynamic pigment of *Fagopyrum esculentum* (buckwheat) (Polygonaceae). Violet powder. Sol. Py, to give red solns. Free base not obt. in pure state. tetra-Ac and tetrabenzoyl derivs. obt. chromatographically pure.

Brockmann, H. *et al.*, *Annalen*, 1952, **575**, 53 (*isol*)

Brockmann, H. *et al.*, *Tet. Lett.*, 1979, 1575 (*uv, pmr, cmr, struct*)

Falcatine

[568-23-0]



Absolute configuration

C₁₇H₁₉NO₄ 301.341

Alkaloid from the bulbs of *Nerine falcata* and *Nerine laticoma* (Amaryllidaceae). Prisms (Et₂O). Mp 127-128°. [α]_D²² -197.8 (c, 1.05 in CHCl₃). Unstable to air and light.

Hydrochloride: Mp 220-235° dec., 238-240° dec.

Ac:

Prisms (EtOH). Mp 201-202°.

Demethoxy: **Caranine**

[477-12-3]

C₁₆H₁₇NO₃ 271.315

Alkaloid from *Amaryllis belladonna* and a large number of other spp. in the Amaryllidaceae. Weak analgesic, convulsant and hypotensive agent, acetylcholinesterase inhibitor. Mp 178-180°. [α]_D²⁷ -197 (c, 2 in CHCl₃).

Demethoxy, perchlorate: Mp 260-270° dec.

Demethoxy, Ac: **Belamarine**. *Bellamarine*.

Acetylcaranine

[14383-07-4]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Amaryllis belladonna* and some other spp. in the Amaryllidaceae. Cytotoxic to P388 sarcoma cells *in vitro*, shows no activity *in vivo*. Sol. MeOH, CHCl₃. Mp 184-185° (173-174°). [α]_D -177 (-81) (CHCl₃). λ_{max} 236 (ε 33200); 294 (ε 35100) (MeOH) (Berdy).

2ξ-*Hydroxy*: **Amaryllidine**

C₁₇H₁₉NO₅ 317.341

Alkaloid from the bulbs of *Amaryllis belladonna* var. *purpurea major* (Amaryllidaceae). Cryst. (MeOH/Me₂CO). Mp 204°. [α]_D²⁵ +64 (c, 0.16 in CHCl₃).

2ξ-*Hydroxy, perchlorate*: Mp 134-135°.

2ξ-*Methoxy*: **Parkamine**

C₁₈H₂₁NO₅ 331.368

Alkaloid from bulbs of *Amaryllis parkeri* (Amaryllidaceae). Cryst. (MeOH/Me₂CO or MeOH). Mp 251-253° dec. [α]_D²³ +69 (c, 0.25 in CHCl₃).

2ξ-*Methoxy, perchlorate*: Mp 245° dec.

2ξ-*Methoxy, picrate*: Mp 178° dec.

Wildman, W.C. *et al.*, *J.A.C.S.*, 1955, **77**, 4807 (*isol*)

Boit, H.G. *et al.*, *Chem. Ber.*, 1956, **89**, 2093; 1959, **92**, 2578 (*Parkamine, Amaryllidine*)

Warnhoff, E.W. *et al.*, *J.A.C.S.*, 1957, **79**, 2192 (*Caranine*)

Takeda, K. *et al.*, *J.A.C.S.*, 1958, **80**, 2562 (*Caranine, synth*)

Fales, H.M. *et al.*, *J.A.C.S.*, 1958, **80**, 4395 (*struct*)

Torszell, K. *et al.*, *Acta Chem. Scand.*, 1961, **15**, 947 (*struct*)

Benington, F. *et al.*, *J.O.C.*, 1962, **27**, 142 (*struct*)

F-7

Kotera, K. *et al.*, *Tet. Lett.*, 1966, 2009

(*Caranine, pmr, config*)


Kinstle, T.H. *et al.*, *Tet. Lett.*, 1966, 4659

(*Caranine, ms*)

Faranine

F-8

[182261-47-8]



C₁₇H₁₇NO₇ 347.324


Alkaloid from leaves of *Medicosma fareana*. Gum. [α]_D -2 (c, 0.7 in CHCl₃). Appears to be a degradn. prod. of an acridone precursor. λ_{max} 305 (EtOH).

Habtemariam, S. *et al.*, *Phytochemistry*, 1996, **43**, 291 (*isol, uv, ir, pmr, cmr, ms, struct*)

Fargesine

F-9

3,4,6,6-Tetrahydro-5-methyl-1H-azepino[5,4,3-cd]indol-7-ol N³-oxide



C₁₂H₁₄N₂O₂ 218.255


Alkaloid from *Evodia fargesii*. Amorph. powder. λ_{max} 218 ; 274 (MeOH).

Qu, S.-J. *et al.*, *Planta Med.*, 2006, **72**, 264-266 (*isol, pmr, ms*)

Farinosone A

F-10

3-(8,10-Dimethyl-1-oxo-2,4,6,8-dodecatrienyl)-4-hydroxy-5-(4-hydroxyphenyl)-2(1H)-pyridinone, 9CI



C₂₅H₂₇NO₄ 405.493

Similar to Tenellin, T-72. Prod. by *Paecilomyces farinosus* RCEF 0101. Neurotrophic agent. Bright yellow powder. [α]_D²⁴ -27.2 (c, 0.15 in MeOH). λ_{max} 249 (log ε 4.24); 415 (log ε 4.34) (MeOH).

N-*Hydroxy*: **Farinosone B**

C₂₅H₂₇NO₅ 421.492


Prod. by *Paecilomyces farinosus* RCEF 0101. Bright yellow powder. [α]_D²⁴ -30 (c, 0.07 in MeOH). λ_{max} 257 (log ε 3.64); 416 (log ε 3.88) (MeOH).

Cheng, Y. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1854-1858 (*isol, pmr, cmr*)

Lang, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 810-811 (*isol*)

Farinosone C

F-11



C₁₉H₂₅NO₅ 347.41
 Prod. by *Paecilomyces farinosus* RCEF 0101. Neurotrophic agent. Amorph. powder. [α]_D²⁴ -20 (c, 0.07 in MeOH). λ_{max} 266 (log ε 2.62) (MeOH).

Cheng, Y. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1854-1858 (*isol, pmr, cmr*)


2,6,10-Farnesatrien-1-amine

F-12

3,7,11-Trimethyl-2,6,10-dodecatrien-1-amine, 9CI. 1-Amino-2,6,10-farnesatriene.

Farnesylamine

[6784-46-9]
 [134963-96-5]



C₁₅H₂₇N 221.385

(2E,6E)-form [129121-59-1]

Alkaloid from the ant *Monomorium fieldi*. Oil. Bp_{0.005} 120°.

(2Z,6E)-form [515846-16-9]

Alkaloid from the ant *Monomorium fieldi*.

Coppola, G.M. *et al.*, *Synth. Commun.*, 1993, **23**, 535-541 (*synth, pmr, cmr*)

Patel, D.V. *et al.*, *J. Med. Chem.*, 1995, **38**, 2906-2921 (*synth, pmr*)


Sen, S.E. *et al.*, *Synthesis*, 1995, 756-758 (*synth, pmr, cmr*)

Jones, T.H. *et al.*, *J. Nat. Prod.*, 2003, **66**, 325-326 (*isol, ms*)

Fascaplysin

F-13

12,13-Dihydro-13-oxopyrido[1,2-a:3,4-b']diindol-5-ium(1+), 9CI. NSC 622398 [114719-57-2]



C₁₈H₁₁N₂O⁺ 271.298

Pigment from the marine sponge *Fascaplysinopsis* sp. and from *Hyrtios erecta*, *Smenospongia* sp. and tunicate *Didemnum* sp. Exhibits antimicrobial and cytotoxic props. Inhibits HIV-1-rt and p56^{LCK} tyrosine kinase. Shows antiplasmodial and antiviral activity. Red cryst. (MeOH or CHCl₃) (as chloride). Sol. MeOH, CHCl₃. Mp 232-235° (chloride). CAS no. refers to chloride. λ_{max} 210 (ε 30400); 237 (ε 6480); 274 (ε 5800); 332 (ε 5470); 398 (ε 2300); 427 (ε 2980); 453 (ε 2700) (MeOH/KOH) (Derep). λ_{max} 214 (ε 7000); 262 (ε 6400); 274 (ε 5450); 300 (ε 6800); 333 (ε 3560); 413 (ε 3100) (MeOH) (Derep).

Salt with *Dehydroloffariellolide diacid: Fascaplysin A*. NSC 660649

[132911-49-0]

C₄₃H₄₈N₂O₅ 672.863

Alkaloid-sesterterpene salt from the sponge *Fascaplysinopsis reticulata*. Red oil.

Salt with 16-Oxodehydroloffariellolide

diacid: Fascaplysin B

[135091-13-3]

C₄₃H₄₆N₂O₆ 686.846

Alkaloid-sesterterpene salt from *Fascaplysinopsis reticulata*. No phys. props. reported.

3-Bromo- 3-Bromofascaplysin. NSC

700409

C₁₈H₁₀BrN₂O⁺ 350.194

Isol. from a *Didemnum* sp. and from *Fascaplysinopsis reticulata*. Red solid (as chloride).

10-Bromo- 10-Bromofascaplysin

C₁₈H₁₀BrN₂O⁺ 350.194

Isol. from *Fascaplysinopsis reticulata*. Red solid (as chloride).

3,10-Dibromo- 3,10-Dibromofascaplysin

C₁₈H₉Br₂N₂O⁺ 429.09

Isol. from *Fascaplysinopsis reticulata*. Red solid (as chloride).

Roll, D.M. *et al.*, *J.O.C.*, 1988, **53**, 3276-3278

(*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Jiménez, C. *et al.*, *J.O.C.*, 1991, **56**, 3403-3410

(*Fascaplysin A*)

Jiménez, C. *et al.*, *Tet. Lett.*, 1991, **32**, 1843

(*Fascaplysin B*)

Gribble, G.W. *et al.*, *J.O.C.*, 1992, **57**, 3636-

3642 (*synth*)

Rocca, P. *et al.*, *Tet. Lett.*, 1993, **34**, 7917-7918

(*synth*)

Molina, P. *et al.*, *Tet. Lett.*, 1994, **35**, 8851-

8854 (*synth*)

Radchenko, O.S. *et al.*, *Tet. Lett.*, 1997, **38**,

5339-5342 (*synth*)

Kirsch, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 825-

829 (*isol, pmr, cmr, activity*)

Segraves, N.L. *et al.*, *Tet. Lett.*, 2003, **44**, 3471-

3475 (*3-Bromofascaplysin*)

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**,

783-792 (*occur, 10-Bromofascaplysin, 3,10-*

Dibromofascaplysin)


Zhidkov, M.E. *et al.*, *Tet. Lett.*, 2007, **48**, 7998-

8000 (*Bromofascaplysin, synth*)

Fasciospongine A

F-14

[945975-25-7]



Relative Configuration

C₃₀H₄₇N₃O₅S 561.784

Alkaloid from a *Fasciospongia* sp. Oil.

[α]_D²³ -51.5 (c, 0.26 in MeOH). λ_{max} 205 (log ε 4.28); 230 (sh) (MeOH).

5-Deoxo, 2-oxo: Fasciospongine B

[945975-26-8]

C₃₀H₄₇N₃O₅S 561.784

Isol. from a *Fasciospongia* sp. Oil. [α]_D

-52.4 (c, 0.25 in MeOH). λ_{max} 205 (log

ε 4.27); 230 (sh) (MeOH).


Yao, G. *et al.*, *Org. Lett.*, 2007, **9**, 3037-3040

(*isol, pmr, cmr*)

Fasicularine

F-15

[187618-62-8]



Relative Configuration

C₂₀H₃₄N₂S 334.568

Alkaloid from the Micronesian ascidian *Nephteis fascicularis*. Shows selective activity against a DNA repair-deficient organism. Cytotoxic to Vero cells. Gum. Closely related to Cylindricine B, C-925.

Patil, A.D. *et al.*, *Tet. Lett.*, 1997, **38**, 363 (*isol, ir, pmr, cmr, ms, struct*)

Maeng, J.-H. *et al.*, *Org. Lett.*, 2002, **4**, 331-333 (*synth*)


Fenster, M.D.B. *et al.*, *Chem. Eur. J.*, 2005, **11**, 639-649 (*synth*)

Abe, H. *et al.*, *J.A.C.S.*, 2005, **127**, 1473-1480 (*synth*)

Fasmerianamine B

F-16

[383191-03-5]



Relative Configuration

C₂₄H₂₂N₄O₆S₂ 526.593

Isol. from the ascidian *Hypsistozoa fasmeriana*. Brown gum. Related to 5-Hydroxy-4-(4-hydroxy-3-methoxyphenyl)-4-(2-imidazolyl)-1,2,3-trithiane, H-507.

Me ester: Fasmerianamine A

[383191-02-4]

C₂₅H₂₄N₄O₆S₂ 540.62

Isol. from *Hypsistozoa fasmeriana*.


Brown gum. [α]_D²⁰ -17.5 (c, 0.4 in MeOH). Incorrect MF stated in ref. λ_{max} 206 (log ε 4.13); 283 (log ε 3.35); 311 (log ε 3.18) (MeOH).

Pearce, A.N. *et al.*, *J.O.C.*, 2001, **66**, 8257-8259

Fastigiatine†

F-17

[103805-65-8]




C₁₉H₂₈N₂O 300.443

Minor alkaloid from *Lycopodium fastigiatum* (Lycopodiaceae). Cryst. (Et₂O). Mp 143-146°. [α]_D²³ +289.9 (c, 1.36 in CHCl₃). Novel pentacyclic ring system.

N-De-Me: Des-N-methylfastigiatineC₁₈H₂₆N₂O 286.416From *Lycopodium fastigiatum* (Lycopodiaceae). Oil.Gerard, R.V. *et al.*, *Can. J. Chem.*, 1986, **64**, 943 (*isol, uv, ir, ms, pmr, cmr, cryst struct*)Gerard, R.V. *et al.*, *Phytochemistry*, 1986, **25**, 1143 (*Des-N-methylfastigiatine*)**Fastrine**

[198541-46-7]

C₃₈H₄₂N₂O₈ 654.758Alkaloid from the roots of *Anisocycla jollyana* (Menispermaceae). Powder. [α]_D²⁰ +49 (c, 0.01 in CHCl₃). λ_{max} 278 (log ε 3.6); 285 (log ε 3.7) (MeOH).**5-Deoxy: Jollyanine†**

[198541-45-6]

C₃₈H₄₂N₂O₇ 638.759Alkaloid from the roots of *Anisocycla jollyana* (Menispermaceae). Powder. [α]_D²⁰ -70 (c, 0.01 in CHCl₃). λ_{max} 276 (log ε 3.6); 284 (log ε 3.6) (MeOH).Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1121-1124 (*isol, uv, ir, cd, pmr, cmr, ms*)**Fastudine**

[11050-43-4]

C₁₆H₂₁NO₃ 275.347Tropane alkaloid. Struct. unknown. Alkaloid from *Datura fastuosa* (*Datura metel*) seeds (Solanaceae). Mp 98°. Ester ir 1710 cm⁻¹.Kaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1966, **3**, 212-216; *CA*, **68**, 29909t (*isol*)**Fastunine**

[11050-44-5]

C₁₇H₂₃NO₃ 289.374Tropane alkaloid. Struct. unknown. Alkaloid from *Datura fastuosa* (*Datura metel*) seeds (Solanaceae). Mp 88-90°. Ir ester 1728 cm⁻¹.Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1966, **3**, 212-216; *CA*, **68**, 29909t (*isol*)**Fastusidine**

[58798-69-9]

Tropane alkaloid. Struct. unknown. Alkaloid from *Datura fastuosa* (*Datura metel*) seeds (Solanaceae). Ir ester 1723 cm⁻¹.*Picrate*: Mp 161-162°.Khaleque, M.A. *et al.*, *Sci. Res. (Dacca)*, 1966, **3**, 212-216; *CA*, **68**, 29909t (*isol*)**Fastusine**


[11013-32-4]

C₁₅H₁₉NO₂ 245.321Struct. unknown. Alkaloid from the seeds of *Datura metel* (*Datura fastuosa*) (Solanaceae). Mp 74-76°. [α]_D³⁰ +55 (EtOH).*Picrate*: Mp 209-211°.*Picolonate*: Mp 195-197°.Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1955, **2**, 147-151; *CA*, **65**, 12246h**Fastusinine**


[11013-31-3]

Struct. unknown. Alkaloid from the seeds of *Datura metel* (*Datura fastuosa*) (Solanaceae). Semisolid.*Picrate*: Mp 235°.Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1955, **2**, 147-151; *CA*, **65**, 12246h**Fauridine**

[11013-31-3]

Struct. unknown. Alkaloid from the seeds of *Datura metel* (*Datura fastuosa*) (Solanaceae). Semisolid.*Picrate*: Mp 235°.Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1955, **2**, 147-151; *CA*, **65**, 12246hC₄₀H₄₆N₂O₈ 682.812Alkaloid from *Thalictrum fauriei*. Amorph. solid. [α]_D²⁴ +285.6 (c, 1.04 in MeOH). λ_{max} 222 (sh) (log ε 4.9); 272 (log ε 4.3); 290 (sh) (log ε 4.21); 311 (sh) (log ε 3.83) (MeOH).Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 803-810 (*isol, uv, ir, cd, pmr, cmr, ms*)**Faurine**


[178765-20-3]

C₂₉H₃₃NO₇ 507.582Alkaloid from whole plants of *Thalictrum fauriei*. Amorph. solid. [α]_D²⁵ +155 (c, 0.8 in CHCl₃). λ_{max} 220 (log ε 5.06); 269 (log ε 4.85); 293 (sh) (log ε 4.63); 310 (sh) (log ε 4.43) (MeOH).*Ac*:Amorph. solid. [α]_D²⁴ +251 (c, 0.54 in CHCl₃).**Me ether: O-Methylfaurine**


[178765-21-4]

C₃₀H₃₅NO₇ 521.609From *Thalictrum fauriei*. Amorph. solid. [α]_D²⁵ +244 (c, 1.2 in MeOH). λ_{max} 220 (log ε 4.56); 270 (log ε 4.34); 296 (sh) (log ε 4.12); 310 (sh) (log ε 3.9) (MeOH).Lee, S.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 738 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)**Fauripavine**


F-26

C₃₉H₄₂N₂O₈ 666.769Alkaloid from *Thalictrum fauriei*. Amorph. off-white solid. [α]_D²⁶ +67.8 (c, 0.9 in MeOH). λ_{max} 220 (sh) (log ε 4.96); 271 (log ε 4.44); 292 (sh) (log ε 4.29); 310 (sh) (log ε 3.86) (MeOH).Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 803-810 (*isol, uv, cd, ir, pmr, cmr, ms*)**Faurithaline**

F-27

C₃₉H₄₄N₂O₈ 668.785Alkaloid from *Thalictrum fauriei*. Amorph. solid. [α]_D²⁵ +83.3 (c, 0.9 in MeOH). λ_{max} 220 (log ε 4.82); 268 (sh) (log ε 4.27); 273 (log ε 4.28); 307 (log ε 3.72) (MeOH).**3-Methoxy: 3-Methoxyfaurithaline**C₄₀H₄₆N₂O₉ 698.811Alkaloid from *Thalictrum fauriei*. Amorph. solid. [α]_D²⁵ +66.6 (c, 0.78 in MeOH). λ_{max} 220 (sh) (log ε 4.81); 275 (log ε 4.3); 315 (sh) (log ε 3.54) (MeOH).Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 803-810 (*isol, uv, ir, cd, pmr, cmr, ms*)**Fawcettidine**

[14912-31-3]

C₁₆H₂₃NO 245.364Alkaloid from *Lycopodium fawcettii*, *Lycopodium alopecuroides* and *Lycopodium*

phlegmaria (Lycopodiaceae). Liq. Bp_{0.1} 125°.

Picrate: Mp 222-223°.

Methiodide: Mp 224-226°.

5β-Alcohol: **Epidihydrofawcettidine**

C₁₆H₂₅NO 247.38

Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Mp 156-157°.

8S-Hydroxy: **Anhydroaposevratinine**

[62023-85-2]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium verticillatum* (Lycopodiaceae). Mp 151°. [α]_D²⁵ +211 (CHCl₃).

8S-Hydroxy, Ac: Mp 144°.

8S-Hydroxy, 3,4-didehydro: **Alolycopine**

[23185-50-4]

C₁₆H₂₁NO₂ 259.347

Alkaloid from *Lycopodium alopecuroides* (Lycopodiaceae). Cryst. (Et₂O). Mp 53-56°.

8S-Hydroxy, 3,4-didehydro, Ac:

Cryst. (Et₂O). Mp 180-182°.

11S-Hydroxy: **11S-Hydroxyfawcettidine**

[942502-18-3]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum*. Amorph. solid. [α]_D²⁵ +134.2 (c, 0.28 in CHCl₃).

2S,11S-Dihydroxy: **2S,11S-Dihydroxy-fawcettidine**

[942502-19-4]

C₁₆H₂₃NO₃ 277.363

Alkaloid from *Lycopodium serratum*. Amorph. solid. [α]_D²⁵ +84.7 (c, 0.2 in CHCl₃).

8S,11S-Dihydroxy: **8S,11S-Dihydroxy-fawcettidine**

[942502-20-7]

C₁₆H₂₃NO₃ 277.363

Alkaloid from *Lycopodium serratum*. Amorph. solid.

Burnell, R.H. et al., *J.C.S.*, 1959, 3091-3093 (*isol*)

Ayer, W.A. et al., *Can. J. Chem.*, 1969, **47**, 2449; 2457 (*Alolycopine*)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1970, **18**, 1880 (*struct*)

Nyembo, L. et al., *Bull. Soc. Chim. Belg.*, 1976, **85**, 595; *CA*, **86**, 90108u

(*Anhydroaposevratinine*)

Inubushi, Y. et al., *Yakugaku Zasshi*, 1982, **102**, 434 (*Epidihydrofawcettidine*)

Harayama, T. et al., *Heterocycles*, 1984, **22**, 1327 (*Epidihydrofawcettidine*)

Katakawa, K. et al., *J. Nat. Prod.*, 2007, **70**, 1024-1028 (*11-hydroxy, 2,11-dihydroxy, 8,11-dihydroxy*)

Kozak, J.A. et al., *Angew. Chem., Int. Ed.*, 2008, **47**, 4221-4223 (*synth*)

C₁₈H₂₉NO₃ 307.432

Alkaloid from *Lycopodium annotinum*, *Lycopodium clavatum*, *Lycopodium fawcettii*, *Lycopodium contiguum*, *Lycopodium magellanicum*, *Lycopodium saururus* and *Lycopodium thyooides* (Lycopodiaceae). Cryst. (Et₂O). Mp 166-167°. [α]_D²⁵ -6 (c, 2.0 in EtOH).

Perchlorate:

Prisms (Me₂CO or H₂O). Mp 272-275°.

Methiodide:

Cryst. (MeOH). Mp 296-297° dec. (279-280°).

Ac: Acetyl-fawcettidine

[52998-88-6]

C₂₀H₃₁NO₄ 349.469

Alkaloid from *Lycopodium fawcettii*, *Lycopodium clavatum*, *Lycopodium contiguum*, *Lycopodium magellanicum*, *Lycopodium saururus* and *Lycopodium thyooides* (Lycopodiaceae). Cryst. by subl. Mp 117°.

O-De-Ac: Deacetyl-fawcettidine

[54307-46-9]

C₁₆H₂₇NO₂ 265.395

Alkaloid from *Lycopodium clavatum*, *Lycopodium fawcettii*, *Lycopodium contiguum*, *Lycopodium magellanicum* and *Lycopodium thyooides* (Lycopodiaceae). Needles (Me₂CO). Mp 203-204°.

O-De-Ac, perchlorate: Mp 224-227°.

O-De-Ac, 5-O-(4-hydroxy-3-methoxy-E-cinnamoyl): Lannotinidine C

[851461-38-6]

C₂₆H₃₅NO₅ 441.566

Alkaloid from *Lycopodium annotinum*. Amorph. solid. [α]_D²⁴ -5 (c, 1 in MeOH).

O-De-Ac, 5-ketone: Clavolonine. Alkaloid L34

[466-62-6]

C₁₆H₂₅NO₂ 263.379

Alkaloid from a variety of *Lycopodium* spp. Mp 238°.

8-Ketone: **Acetylannofoline. Obscurumine B**

[132143-35-2]

C₁₈H₂₇NO₃ 305.416

Alkaloid from *Lycopodium obscurum* var. *dendroideum* (Lycopodiaceae). Amorph. solid. [α]_D²⁴ -63 (c, 0.5 in MeOH). Component of Alkaloid L17 which was a mixt. with Acetylacrifoline.

O-De-Ac, 8-ketone: Annofoline

[664-23-3]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Cryst. (MeOH/Et₂O). Mp 156-157°. [α]_D²⁵ -131 (c, 2 in EtOH).

O-De-Ac, 8-ketone, perchlorate:

Cryst. (MeOH/Et₂O). Mp 234-236°.

11,12-Didehydro, *O-de-Ac, 5-O-(4-hydroxy-3-methoxy-E-cinnamoyl): Lannotinidine D*

[851461-39-7]

C₂₆H₃₃NO₅ 439.55

Alkaloid from *Lycopodium annotinum*. Amorph. solid. [α]_D²⁴ -24 (c, 1 in MeOH).

7-Hydroxy, 5-ketone, *O-de-Ac: 7,8β-Dihydroxylycopodine. Sauroine*

[777855-51-3]

C₁₆H₂₅NO₃ 279.378

Alkaloid from the aerial parts of *Huperzia saururus*. [α]_D²⁵ +26.2 (c, 0.08 in MeOH).

8-Epimer (?): **α-Lofoline. Lofoline**

[561-15-9]

C₁₈H₂₉NO₃ 307.432

Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Prisms (MeOH). Mp 211-212°. [α]_D²⁵ -52 (c, 2 in EtOH). C-8 and C-15 configs. not fully clear from published data.

8-Epimer, *methiodide*:

Cryst. (MeOH/Me₂CO). Mp 266-267°.

8-Epimer, *Ac, 1:1 complex with Lycopodine, L-341: Alkaloid L9*

C₃₆H₅₆N₂O₅ 596.849

Isol. from *Lycopodium annotinum* (Lycopodiaceae). Fine needles (Et₂O). Mp 122°.

8-Epimer, *Ac: Acetyllofoline*

[102378-18-7]

C₂₀H₃₁NO₄ 349.469

Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Oil; cryst. (Me₂CO/Et₂O) (as perchlorate). Mp 272-273° dec. (perchlorate). Component of L9 in T-263. Alkaloid L31 was a mixt. of Acetyllofoline and 5,15-Oxidolycopodane, O-163.

Anet, F.A.L. et al., *Can. J. Chem.*, 1959, **37**, 1589-1596 (*Fawcettiine, Lofoline, Annofoline, isol, ir*)

Burnell, R.H. et al., *J.C.S.*, 1959, 3091-3093 (*Fawcettiine, Deacetyl-fawcettidine*)

Burnell, R.H. et al., *Can. J. Chem.*, 1960, **38**, 1927-1932 (*Acetyl-fawcettidine*)

Anet, F.A.L. et al., *Chem. Ind. (London)*,

1960, 1238-1239 (*Annofoline, pmr, struct*)

Burnell, R.H. et al., *Chem. Ind. (London)*, 1960, 1239-1240 (*Fawcettiine, struct, stereochem*)

Burnell, R.H. et al., *Tetrahedron*, 1961, **15**, 173-182 (*Clavolonine*)

Ayer, W.A. et al., *Can. J. Chem.*, 1964, **42**, 949-951 (*Acetyllofoline, Alkaloid L9*)

Ayer, W.A. et al., *Tet. Lett.*, 1964, 2959-2963 (*Annofoline, struct*)

Nakashima, T.T. et al., *Can. J. Chem.*, 1975,

53, 1936-1942 (*Lofoline, Clavolonine, cmr*)

Wenkert, E. et al., *Chem. Comm.*, 1984, 714-715 (*Acetyl-fawcettidine, Deacetyl-fawcettidine, Clavolonine, Annofoline, synth*)

Ayer, W.A. et al., *Can. J. Chem.*, 1990, **68**, 1300-1304 (*Alkaloid L17, Alkaloid L31*)

Ortega, M.G. et al., *Tet. Lett.*, 2004, **45**, 7003-7005 (*Sauroine*)


Evans, D.A. et al., *Angew. Chem., Int. Ed.*, 2005, **44**, 6038-6042 (*Clavolonine, synth*)

Morita, H. et al., *Tetrahedron*, 2005, **61**, 1955-1960 (*Obscurumine B*)

Koyama, K. et al., *Tetrahedron*, 2005, **61**, 3681-3690 (*Lannotinidines C,D*)

Fawcettiine

β-Lofoline
[6899-87-2]




(-)-form

F-29

Fawcettimine

[15228-74-7]

[118892-07-2 ((±)-form), 103498-99-3 ((±)-form hydrobromide)]



Absolute Configuration

F-30

C₁₆H₂₅NO₂ 263.379

Shows ring-chain tautomerism; can exist either as the carbinolamine (shown) or as the aminoketone. Alkaloid from *Lycopodium fawcettii*, *Lycopodium clavatum* and *Lycopodium clavatum* var. *inflexum* (Lycopodiaceae).

Hydrochloride:Cryst. (EtOH/Me₂CO). Mp 235° dec.**Perchlorate:**Cryst. (2-propanol/Me₂CO). Mp 225–226°.**6,7-Didehydro: Lycoposerramine C**

[481048-23-1]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum*. Prisms (EtOAc/hexane). Mp 164–165°. λ_{max} 229 ; 270 (EtOH).

9,10-Didehydro: Lycothunine

[11052-60-1]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum* var. *serratum* f. *intermedium* (Lycopodiaceae).

2R-Hydroxy, 9,10-didehydro: 2R-Hydroxylycothunine

[942502-21-8]

C₁₆H₂₃NO₃ 277.363

Alkaloid from *Lycopodium serratum*. Amorph. powder.

2S-Hydroxy: Lycoposerramine P

[481048-33-3]

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium serratum*. Amorph. powder.

4-Hydroxy: Alopecuridine

[11049-78-8]

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium alopecuroides* and *Lycopodium sieboldii*. Cryst. (Me₂CO). Mp 171–172°.

8S-Hydroxy, 9,10-didehydro: 8S-Hydroxylycothunine

[942502-22-9]

C₁₆H₂₃NO₃ 277.363

Alkaloid from *Lycopodium serratum*. Amorph. solid. [α]_D²¹ +72.2 (c, 0.26 in CHCl₃).

13-Deoxy, N-oxide: Lannotinidine B

[851461-37-5]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium annotinum*. Amorph. solid. [α]_D²⁴ -62 (c, 1 in MeOH).

13-Deoxy, 13,14-didehydro, 5S-alcohol: Lycoposerramine Q

[481048-37-7]

C₁₆H₂₅NO 247.38

Alkaloid from *Lycopodium serratum*. Amorph. solid.

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1961, **39**, 1090–1093 (*isol, ir*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1968, **46**, 15–20 (*Alopecuridine*)

Ayer, W.A. *et al.*, *Tet. Lett.*, 1973, 5045–5048 (*occur*)

Ayer, W.A. *et al.*, *Tetrahedron*, 1974, **30**, 4213–4214 (*Alopecuridine*)

Harayama, T. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2394–2402 (*synth*)

Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 3418–3421 (*Lycothunine, cryst struct*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1989, **54**, 1548–1562 (*synth, ir, pmr, cmr*)

Takayama, H. *et al.*, *Tet. Lett.*, 2002, **43**, 8307–8311 (*cmr, Lycoposerramines*)

Hirasawa, Y. *et al.*, *Org. Lett.*, 2003, **5**, 3991–3993 (*Alopecuridine, cryst struct*)

Koyama, K. *et al.*, *Tetrahedron*, 2005, **61**, 3681–3690 (*Lannotinidine B*)


Linghu, X. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 7671–7673 (*synth*)

Katakawa, K. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1024–1028 (*Lycothunine, Hydroxylycothunines*)

Febrifugine†

F-31

3-[3-(3-Hydroxy-2-piperidinyl)-2-oxopropyl]-4(3H)-quinazolinone, 9CI. Dichroine B. β-Dichroine



Absolute Configuration

C₁₆H₁₉N₃O₃ 301.344

Absolute config. revised in 1999 and again in 2003.

(+)-form [24159-07-7]

Alkaloid from the roots and leaves of the Chinese drug Ch'ang Shan (*Dichroa febrifuga*) and from *Hydrangea* spp. (Hydrangeaceae). Shows high degree of antimalarial activity (*ca.* 100 times as active as Quinine, Q-20) but use limited by toxicity. Also shows antipyretic and emetic activities; an effective coccidiostat. Various synthetic analogues show similar activities and much lower toxicity. Antineoplastic agent. Possesses schizonticide props. Needles (EtOH). Mp 139–140° Mp 154–156° (dimorph.). [α]_D²⁵ +6 (c, 0.5 in CHCl₃) (c, 0.5 in EtOH). Log P -0.33 (calc). Pharmacol. active isomer.

Hydrochloride (1:2): Mp 220–222° dec.

Oxime: Mp 224–225° dec.

Semicarbazone: Mp 187–188° dec.

Bis(benzenesulfonyl): Mp 148–148.5°.

(±)-form [39037-90-6]

Synthetic. Mp 178.5–180.5°.

Hydrochloride (1:2): Mp 201–204° dec.

Hydrobromide:

Cryst. (EtOH). Mp 229–231°.

Koepfli, J.B. *et al.*, *J.A.C.S.*, 1947, **69**, 1837; 1949, **71**, 1048 (*isol, uv*)

Kuehl, F.A. *et al.*, *J.A.C.S.*, 1948, **70**, 2091 (*isol*)

Ablondi, F. *et al.*, *J.O.C.*, 1952, **17**, 14 (*isol*)

Baker, B.R. *et al.*, *J.O.C.*, 1955, **20**, 136 (*synth*)

Barringer, D.F. *et al.*, *J.O.C.*, 1973, **38**, 1933; 1937 (*pmr, abs config*)

Johns, S. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 129 (*rev. pharmacol*)

Burgess, L.E. *et al.*, *Tet. Lett.*, 1996, **37**, 3255 (*synth*)

Uesato, S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1–5 (*pmr, cmr, conform*)

Murata, K. *et al.*, *J. Nat. Prod.*, 1998, **61**, 729–733 (*isol, uv, ir, pmr, cmr, ms*)

Takaya, Y. *et al.*, *J. Med. Chem.*, 1999, **42**, 3163–3166 (*isol*)

Kobayashi, S. *et al.*, *J.O.C.*, 1999, **64**, 6833–6841 (*synth, abs config*)

Takeuchi, Y. *et al.*, *Tetrahedron*, 2003, **59**, 1639–1646 (*synth, struct, bibl*)

Katoh, M. *et al.*, *Heterocycles*, 2006, **67**, 189–204 (*synth*)

Fedamazine

F-32

C₂₀H₂₁N₂O[⊕] 305.399

Struct. unknown. Alkaloid from calabash curare (*Strychnos toxifera*) (Loganiaceae). Mp 233–235° (as picrate). Stable blue col. with Ce(SO₄)₂.


Asmis, H. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1983–1992 (*isol, uv*)

Feldamycin

F-33

N-Methylhistidyl-β-[1-carboxy-2-(1H-imidazol-4-yl)ethyl]amino]-α-aminobutanoic acid, 9CI. BMY 28565. U 48266. Antibiotic BMY 28565. Antibiotic U 48266

[61230-27-1]

C₁₇H₂₅N₇O₅ 407.428

Peptide antibiotic. Prod. by *Streptomyces ficellus* and *Streptomyces calvus*. Active against gram-positive and -negative bacteria. Inhibits melanin synthesis.

Amorph. solid. Sol. H₂O, EtOH, MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁵ -6.6 (c, 1 in H₂O).

▶EK7910000

Argoudelis, A.D. *et al.*, *J. Antibiot.*, 1976, **29**, 1007; 1117 (*isol*)

U.S. Pat., 1976, 3 969 515; CA, **85**, 141320


Reusser, F. *et al.*, *Biochemistry*, 1977, **16**, 3406 (*props*)

Imae, K. *et al.*, *J. Antibiot.*, 1991, **44**, 76 (*synth, abs config*)

Fenfangjine H

F-34

[134227-37-5]

C₃₇H₃₉N₂O₇[⊕] 623.724

Alkaloid from the roots of *Stephania tetrandra*. Antihypertensive agent.

Amorph. orange powder (as hydroxide).

[α]_D²⁷ -89.5 (c, 0.4 in MeOH) (hydroxide). CAS no. refers to hydroxide.

α-Aldehyde: Fenfangjine I

[134242-53-8]

C₃₇H₃₇N₂O₇[⊕] 621.709

Alkaloid from *Stephania tetrandra*.


Antihypertensive agent. Amorph. orange powder (as hydroxide).

[α]_D²⁷ -46.5 (c, 0.3 in MeOH) (hydroxide). CAS no. refers to hydroxide.

Ogino, T. *et al.*, *Heterocycles*, 1998, **48**, 311–317 (*isol, ir, pmr, cmr*)

Ferribactin G173

[535994-96-8]



C₅₀H₇₆N₁₄O₂₀ 1193.232

L-Glu incorr. drawn in ref. diag. Prod. by *Pseudomonas fluorescens* G173. Siderophore. Biosynthetic precursor of Pyoverdine G 173, P-853.

N^{Glu}-Ac: [535994-98-0]

C₅₂H₇₈N₁₄O₂₁ 1235.269

Prod. by *Pseudomonas fluorescens* G173.

N^{Glu}-Di-Ac: [535994-99-1]

C₅₄H₈₀N₁₄O₂₂ 1277.307

Prod. by *Pseudomonas fluorescens* G173. Siderophore.

N^{Glu}-Hydroxy: [535994-97-9]


C₅₀H₇₆N₁₄O₂₁ 1209.232

Prod. by *Pseudomonas fluorescens* G173. Siderophore.

Fernandez, D.U. *et al.*, *Z. Naturforsch.*, C, 2003, **58**, 1-10 (*isol*, *pmr*, *ms*, *N^{Glu}-derivs*)

Ferrirocina, 9CI

[23086-46-6]



C₂₈H₄₄FeN₉O₁₃ 770.555

Secreted by *Aspergillus* spp. Principal siderophore from *Micromonospora gypseum*. Growth factor and iron transport compd. Phytotoxin. Orange-brown cryst. + 7H₂O (EtOH). Sol. MeOH, H₂O. Mp 250° dec. λ_{max} 430 nm (log ε 3.42). λ_{max} 422 (ε 2058); 430 (ε 2630) (H₂O) (Berdy).

[37279-99-5]

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 1907 (*ir*, *struct*)

Llinas, M. *et al.*, *J. Mol. Biol.*, 1972, **68**, 265; 1976, **104**, 853 (*pmr*, *conform*, *struct*, *cmr*)

Fiedler, H.P. *et al.*, *J. Chromatogr.*, 1981, **209**, 103 (*hplc*)

Dell, A. *et al.*, *Biomed. Mass Spectrom.*, 1982, **9**, 158 (*ms*)

Wong, G.B. *et al.*, *J.A.C.S.*, 1983, **105**, 810 (*props*)

Barnes, C.L. *et al.*, *Acta Cryst. C*, 1984, **40**, 922


Bentley, M.D. *et al.*, *Biochemistry*, 1986, **25**, 1455 (*isol*)

Ohra, J. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 113; 1782 (*isol*, *cd*, *props*)

F-35

Ferrimycin A₁, 9CI

[15319-50-3]



C₄₁H₆₅FeN₁₀O₁₄ 977.87

Produced by *Streptomyces griseoflavus* strain ETH 9578. Shows antibiotic props. Sol. H₂O, MeOH, DMF, phenol; fairly sol. AcOH, EtOH, 1-propanol, Py; poorly sol. butanol, hexane. Acyl deriv. of its competitive antagonist Ferrioxamine B, F-39. λ_{max} 229 (E1%/1cm 336); 319 (E1%/1cm 37); 428 (E1%/1cm 27.6) (H₂O) (Berdy).

Hydrochloride (1:2):

Amorph., hygroscopic solid. pK_{a1} 4.11; pK_{a2} 7.92; pK_{a3} 11.4 (2-methoxyethanol).


Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2105 (*isol*, *ir*)

Bickel, H. *et al.*, *Tetrahedron, Suppl.*, No. 8, 1966, 171 (*struct*, *ir*, *uv*, *nmr*)

Ferrioxamine A₂

F-38

[N'-[5-[4-[4-(Acetylhydroxyamino)-butyl]amino]-1,4-dioxobutyl]hydroxyamino]pentyl]-N-(5-aminopentyl)-N-hydroxybutanediamidato(3-)iron(III), 9CI



C₂₄H₄₃FeN₆O₈ 599.486

Component of inhomogeneous Ferrioxamine A complex from *Streptomyces pilosus*. Iron-transport compd. Sol. H₂O, MeOH. pK_a 9.89 (2-methoxyethanol). λ_{max} 430 (ε 2238); 440 (H₂O) (Berdy).

N-Ac:

Amorph. brown-orange powder. Mp 146-147°.

F-37

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 710 (*ir*, *pmr*, *struct*, *synth*)

Ferrioxamine B

F-39

[N'-[5-[4-[4-(Acetylhydroxyamino)-pentyl]amino]-1,4-dioxobutyl]hydroxyamino]pentyl]-N-(5-aminopentyl)-N-hydroxybutanediamidato(3-)iron. Ferrioxamine D

[14836-73-8]

As Ferrioxamine A₂, F-38 with m = n = 5

C₂₅H₄₅FeN₆O₈ 613.513

Siderophore. Metab. of *Actinomyces* spp. Important Sideramine-type growth factor isol. from *Streptomyces pilosus*.

Hygroscopic red-brown solid. Sol. H₂O, MeOH. pK_a 9.74 (2-methoxyethanol). λ_{max} 428 (ε 2800); 430 (ε 2510); 440 (H₂O) (Berdy).

N-Ac: **Ferrioxamine D₁**

C₂₇H₄₇FeN₆O₉ 655.55

Growth factor. Sol. H₂O, MeOH. λ_{max} 430 (ε 2884) (H₂O) (Berdy).

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118; 2129 (*isol*, *ir*)

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 709 (*ir*, *struct*)

Prelog, V. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 631 (*ir*, *synth*)

Bock, J.L. *et al.*, *Biochim. Biophys. Acta*, 1972, **264**, 245 (*Mössbauer*)

Karrer, W. *et al.*, *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 3180

Leong, J. *et al.*, *J.A.C.S.*, 1975, **97**, 293 (*struct*, *synth*)

Tufano, T.P. *et al.*, *J.A.C.S.*, 1981, **103**, 6617 (*metal exch*, *uv*)

Hossain, M.B. *et al.*, *Acta Cryst. C*, 1986, **42**, 1275 (*cryst struct*)

Ferrioxamine C

F-40


Structure unknown but thought to be similar to Ferrioxamine B, F-39. Metab. of *Actinomyces* spp. Growth factor.

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118 (*ir*, *isol*)

Ferrioxamine D₂

F-41

[29825-04-5]



C₂₆H₄₃FeN₆O₉ 639.507

Isol. from *Streptomyces pilosus*. Cryst. (MeOH/Et₂O). Sol. H₂O, MeOH. Mp 220-223°.

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118 (*isol*)

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 710 (*isol, pmr, struct*)

Ferrioxamine E F-42

Nocardamine ferric salt
[20008-20-2]

As Ferrioxamine D₂, F-41 with n = 5

C₂₇H₄₅FeN₆O₉ 653.534

Metab. of *Actinomyces* spp. Growth factor important in iron transport. Amorph. solid. Sol. H₂O, MeOH.

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118 (*isol, ir*)

Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1981 (*ir, uv, struct*)

Van der Helm, D. *et al.*, *J.A.C.S.*, 1976, **98**, 82 (*cryst struct*)

Ferrioxamine F F-43

Structure unknown but thought to be similar to Ferrioxamine B, F-39. Metab. of *Actinomyces* spp. Growth factor. pK_a 9.75 (2-methoxyethanol).

Bickel, H. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 2118 (*ir, isol*)

Ferrioxamine H F-44

[HOOCCH₂CH₂CONH(CH₂)₅N(OH)-COCH₂CH₂CONH(CH₂)₅-N(OH)Ac]Fe

C₂₀H₃₆FeN₄O₈ 516.373

Minor component of the Sideramine complex of actinomycetes. Red amorph. powder.


Adapa, S. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 1818

Ferroverdin F-45

Ferroverdin A. Tris(4-ethenylphenyl 4-hydroxy-3-nitrosobenzoato-N,O)ferate(1-)

[18904-95-5]

[26671-45-4]



C₄₅H₃₀FeN₃O₁₂ 860.592

First formulated as C₃₀H₂₄FeN₂O₈. Later, ligand proved to be *p*-vinylphenyl ester of 3-nitroso-4-hydroxybenzoic acid. X-ray data (1969) show 3 ligands attached to Fe(II) centre. Pigment produced by a *Streptomyces* sp. when cultured in an Fe(III) containing medium. Inhibits cholesteryl ester transfer protein, weakly active against *Acholeplasma laidlawii*. Green powder. Insol. H₂O, C₆H₆, CHCl₃; spar. sol. EtOH, Et₂O, AcOH; Sol. MeOH, MeCN, EtOAc. [α]_D²⁵ -41400 (c, 0.01 in MeOH). Readily reduced by H₂ in presence of Pd catalyst. λ_{max} 285 (ε 44000); 305 (sh) (ε 24000); 440 (ε 6000); 690 (ε 5300) (MeOH). λ_{max} 285 (ε 44000); 440 (ε 6000); 690 (ε 5300) (MeOH) (Berdy).

8-Monohydroxy(E-): Ferroverdin B. Antibiotic WK 5344A. WK 5344A

[259733-69-2]

[290304-15-3]

C₄₅H₃₀FeN₃O₁₃ 876.591

Prod. by *Streptomyces* sp. WK-5344. Inhibitor of cholesteryl ester transfer protein. Green powder. [α]_D²⁵ -3000 (c, 0.01 in MeOH). λ_{max} 285 (ε 44000); 305 (sh) (ε 24000); 440 (ε 6000); 690 (ε 5300) (MeOH).

8-Monocarboxy(Z-): Ferroverdin C.

Antibiotic WK 5344B. WK 5344B

C₄₆H₃₀FeN₃O₁₄ 904.602

Prod. by *Streptomyces* sp. WK-5344. Inhibitor of cholesteryl ester transfer protein. Green powder. [α]_D²⁵ -1000 (c, 0.01 in MeOH). λ_{max} 285 (ε 44000); 305 (sh) (ε 24000); 440 (ε 6000); 690 (ε 5300) (no solvent reported).

[52746-83-5]

Chain, E.B. *et al.*, *Nature (London)*, 1955, **176**, 645 (*isol, uv*)

Ehrenberg, A. *et al.*, *Nature (London)*, 1956, **178**, 379 (*magnetism*)

Ballio, A. *et al.*, *Nature (London)*, 1962, **194**, 769 (*struct, ir, uv*)

Ballio, A. *et al.*, *Proc. R. Soc. London, B*, 1963, **158**, 43 (*struct*)

Candeloro, S. *et al.*, *Nature (London)*, 1969, **224**, 589 (*struct*)

Candeloro De Sanctif, S. *et al.*, *Proc. R. Soc. London, B*, 1973, **184**, 121; 137 (*struct*)


Yang, C.C. *et al.*, *J. Bacteriol.*, 1982, **149**, 381 (*biochem, use*)

Sveshnikova, M.A. *et al.*, *Antibiotiki (Moscow)*, 1983, **28**, 28; *CA*, **99**, 209437m (*biochem, use*)

Tomoda, H. *et al.*, *J. Antibiot.*, 1999, **52**, 1101-1107; 1108-1113 (*isol, uv, pmr, cmr, ms, activity*)

Ferrugine† F-46

(8-Methyl-8-azabicyclo[3.2.1]oct-2-yl)-phenylmethanone, 9CI. 2α-Benzoyltropane [58471-11-7]



Absolute, configuration

C₁₅H₁₉NO 229.321

Alkaloid from the leaves, stems and roots of *Darlingia ferruginea* (Proteaceae). Rosettes (EtOH). Mp 101-102°. [α]_D¹⁹ +55 (CHCl₃).


Picrate: Mp 192-194°.

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 2537 (*isol, uv, ir, pmr, ms, abs config*)

Ahmed, S. *et al.*, *J.O.C.*, 2008, **73**, 8116-8119 (*synth*)

Ferruginine F-47

1-(8-Methyl-8-azabicyclo[3.2.1]oct-2-en-2-yl)ethanone, 9CI. 2-Acetyl-7-methyl-7-azabicyclo[3.2.1]oct-2-ene



(+)-form

C₁₀H₁₅NO 165.235

(+)-form [73069-63-3]

Alkaloid from *Darlingia ferruginea* (Proteaceae). Noncryst. [α]_D¹⁹ +37 (CHCl₃). *Picrate*: Mp 161-163°.

(-)-form [64603-84-5]

Synthetic. Noncryst. [α]_D¹⁹ -37 (CHCl₃).

Hydrochloride: [64626-41-1]

Mp 161-163°. [α]_D¹⁹ -70 (c, 1 in MeOH).

Campbell, H.F. *et al.*, *Can. J. Chem.*, 1977, **55**, 1372 (*synth*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 2537 (*isol, uv, ir, pmr, ms, synth, struct*)

Davies, H.M.L. *et al.*, *J.O.C.*, 1991, **56**, 5696; 1997, **62**, 1095 (*synth, ir, pmr, cmr, ms*)

Rigby, J.H. *et al.*, *J.O.C.*, 1995, **60**, 7392 (*synth*)

Hernández, A.S. *et al.*, *J.O.C.*, 1996, **61**, 314 (*synth*)

Gauthier, I. *et al.*, *J.O.C.*, 1997, **62**, 6704-6705 (*synth*)


Ham, W.-H. *et al.*, *Tet. Lett.*, 1997, **38**, 3247-3248 (*synth*)

Jonsson, S.Y. *et al.*, *J.O.C.*, 2000, **65**, 8458-8464 (*synth*)

Piccardi, R. *et al.*, *Eur. J. Org. Chem.*, 2007, 4752-4757 (*synth*)

Fervenuin F-48

6,8-Dimethylpyrimido[5,4-e]-1,2,4-triazine-5,7(6H,8H)-dione, 9CI. Planomycin. Pulanomycin. Antibiotic 10204-BII [483-57-8]



C₇H₇N₅O₂ 193.165

Isol. from *Streptomyces fervens* and *Streptomyces rubriretili*. Active against Gram-positive and -negative bacteria and possesses antineoplastic props. Yellow cryst. Mp 178-179°. Log P -0.82 (uncertain value) (calc). Known synthetically before isolation. λ_{max} 237 (ε 17000); 275 (ε 1950); 341 (ε 4250) (pH 1) (Derep). λ_{max} 230 (sh) (ε 4850); 275 (ε 1900) (pH 11) (Derep). λ_{max} 239 (ε 17400); 277 (ε 1860); 343 (ε 4570) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 11.2 mg/kg. UW7900000

Pfleiderer, W. *et al.*, *Annalen*, 1958, **615**, 42 (*synth*)

Daves, G.D. *et al.*, *J.O.C.*, 1961, **26**, 5256 (*isol, struct, uv, ir*)

Yoneda, F. *et al.*, *J. Het. Chem.*, 1970, **7**, 1443 (*synth*)

Taylor, E.C. *et al.*, *J.O.C.*, 1975, **40**, 2321 (*synth, nmr*)

Senda, S. *et al.*, *J.A.C.S.*, 1977, **99**, 7358 (*synth*)

Ichiba, M. *et al.*, *J.O.C.*, 1978, **43**, 469 (*synth, ir, uv, ms, nmr*)

CRS Handb. of Antibiotic Compounds, (eds. Berdy, J. *et al.*), CRC Press, Boca Raton, 1984, **5**, 198 (*props*)


Werner-Simon, S. *et al.*, *J. Het. Chem.*, 1996, **33**, 949 (*synth, derivis*)

Koshino, H. *et al.*, *Heterocycles*, 2000, **52**, 811-817 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FBP300

Festuclovine

Dihydroagroclavine
[569-26-6]



Absolute
Configuration

C₁₆H₂₀N₂ 240.347

Alkaloid from *Claviceps purpurea*, *Claviceps gigantea*, *Aspergillus fumigatus*, *Penicillium roquefortii*, *Penicillium crustosum*, *Argyria nervosa* and several other *Argyria* spp., *Ipomoea hildebrandtii*, *Ipomoea violacea* and *Stictocardia tiliaefolia*. Mycotoxin, 5-HT receptor antagonist; active against gram-positive bacteria. Needles (MeOH). Sol. MeOH, CHCl₃. Mp 249-252° (242-243°). [α]_D²⁰ -113 (c, 0.25 in Py). [α]_D¹⁴ -67.4 (c, 0.12 in CHCl₃). λ_{max} 224 (log ε 4.53); 276 (log ε 3.82); 281 (log ε 3.84) (EtOH).

► LD₅₀ (mus, ipr) 45 mg/kg. KE6334000

2-Bromo- **Pibocine A**. 2-Bromofestuclovine [102394-16-1]

C₁₆H₁₉BrN₂ 319.244

Alkaloid from the ascidian *Eudistoma* sp. Moderate cytotoxic agent. Mp 226-228°. [α]_D -36 (c, 0.14 in EtOH). λ_{max} 225 (ε 27400); 281 (ε 7000) (EtOH).

2-Bromo, N¹-methoxy: **Pibocine B**

[388112-41-2]

C₁₇H₂₁BrN₂O 349.27

Alkaloid from a *Eudistoma* sp. Cytotoxic. Thin cryst. (MeOH). Mp >358°. [α]_D -51 (c, 0.19 in EtOH). λ_{max} 231 (ε 4410); 283 (ε 2450) (MeOH).

18-Hydroxy: **α-Dihydrolysergol**. *Dihydrolysergol*. *Dihydroelymoclavine*

[18051-16-6]

C₁₆H₂₀N₂O 256.347

Alkaloid from *Claviceps gigantea*. Also present in the seeds of *Stictocardia campanulata* and *Ipomoea muelleri* (Convolvulaceae). 5-HT receptor antagonist, mycotoxin. Prisms (MeOH). Mp 283°. [α]_D²⁰ -60 (c, 0.13 in EtOH). [α]_D -92 (c, 0.5 in Py). λ_{max} 224 (log ε 4.6); 282 (log ε 3.92); 292 (log ε 3.88) (EtOH).

8-Epimer: **Pyroclavine**

[478-89-7]

C₁₆H₂₀N₂ 240.347

Prod. by *Claviceps purpurea* and *Claviceps gigantea*, also *Penicillium* sp. 5-HT receptor antagonist, mycotoxin. Mp 204°. [α]_D²⁰ -90 (c, 0.2 in Py).

10-Epimer: **Costaclavine**

[436-41-9]

C₁₆H₂₀N₂ 240.347

Alkaloid from *Claviceps purpurea* and *Penicillium chermesinum*. Mp 182-184°. [α]_D +44 (Py). λ_{max} 275; 283; 293 (no solvent reported).

8,10-Diepimer: **Epicostaclavine**

F-49

Mp 132-135°.

Abe, M. *et al.*, *Bull. Agric. Chem. Soc. Jpn.*, 1956, **20**, 59-60; 1959, **33**, 1031-1036; *CA*, **51**, 11365c; **59**, 2878c (*Costaclavine*, *Pyroclavine*, *isol*, *struct*)

Spilsbury, J.F. *et al.*, *J.C.S.*, 1961, 2085-2089 (*isol*, *uv*)

Aguere, S.L. *et al.*, *Experientia*, 1964, **20**, 25-26 (*Costaclavine*, *isol*, *uv*, *ir*)

Aguere, S. *et al.*, *Acta Pharm. Suec.*, 1965, **2**, 231-238 (*Pyroclavine*, *Festuclovine*, *α-Dihydrolysergol*, *isol*)

Semononsky, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1966, **31**, 577-582 (*α-Dihydrolysergol*, *synth*)

Chao, J.M. *et al.*, *Phytochemistry*, 1973, **12**, 2435-2440 (*Festuclovine*, *occur*)

Bach, N.J. *et al.*, *J.O.C.*, 1974, **39**, 1272-1276 (*pmr*, *cmr*)

Voigt, D. *et al.*, *Pharmazie*, 1974, **29**, 697 (*ms*)

Ohmomo, S. *et al.*, *Agric. Biol. Chem.*, 1975, **39**, 1333-1336 (*isol*, *uv*, *ir*)

Ninomiya, I. *et al.*, *Chem. Comm.*, 1976, 624-626 (*Costaclavine*, *Pyroclavine*, *pmr*)

Krepelka, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 1416-1418 (*synth*)

Lee, T.M. *et al.*, *Planta Med.*, 1979, **35**, 247-252 (*α-Dihydrolysergol*, *isol*)

Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 655-657 (*isol*)

Oppolzer, W. *et al.*, *Tetrahedron*, 1983, **39**, 3695 (*Costaclavine*, *synth*, *ir*, *pmr*, *ms*)

Eich, E. *et al.*, *Arzneim.-Forsch.*, 1985, **35**, 1760-1762 (*Festuclovine*, *activity*)

Osanaï, K. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1587-1590 (*Costaclavine*, *synth*)

Makariev, T.N. *et al.*, *Tet. Lett.*, 1999, **40**, 1591-1594 (*Pibocine A*)

Makariev, T.N. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1559-1561 (*Pibocine B*, *Festuclovine*)

Fibraminine

F-50

[1357-05-7]

C₁₈H₁₉NO₈ 377.35

Struct. unknown. Compd. possesses 4 methoxy groups. Alkaloid from *Fibraurea tinctoria* (Menispermaceae). Yellowish prisms. Mp 192-193°. [α]_D¹⁵ +28.33 (c, 1 in CHCl₃).

Hydrobromide:

Yellowish needles. Mp 202-203°.

Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1962, **28**, 89-95; *CA*, **60**, 6887g

Fibranine

F-51

[1357-06-8]

C₂₅H₂₉NO₇ 455.507

Struct. unknown. Compd. possesses 3 methoxy and 2 methylenedioxy groups. Alkaloid from *Fibraurea tinctoria* (Menispermaceae).

Hydrochloride:

Reddish prismatic needles + 1H₂O. Mp 196-198°. [α]_D¹⁵ +273.3 (c, 1 in EtOH).

Hydroiodide:

Yellowish needles. Mp 217-218°.

Perchlorate:

Yellowish needles. Mp 273-274°.

Picrate:

Reddish needles. Mp 212-213°.

Chloroaurate:


Greenish-yellow needles. Mp 185-186°.

Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1962, **28**, 89-95; *CA*, **60**, 6887g

Ficellomycin

F-52

2-[4-[(Aminoiminomethyl)amino]-1-azabicyclo[3.1.0]hex-2-yl]-N-valylglycine, 9CI. U 47929. Antibiotic U 47929 [59458-27-4]



C₁₃H₂₄N₆O₃ 312.371

Dipeptide antibiotic. From *Streptomyces ficellus*. Animal feed additive. Active against gram-positive bacteria. Amorph. solid. Sol. H₂O, EtOH, MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁵ +39 (c, 1 in H₂O).

► LD₅₀ (mus, ivn) 800 mg/kg; LD₅₀ (mus, ipr) 800 mg/kg. LK3660000

Argoudelis, A.D. *et al.*, *J. Antibiot.*, 1976, **29**, 1001 (*isol*)

U.S. Pat., 1976, 542 226; *CA*, **85**, 19047

U.S. Pat., 1976, 3 993 748; *CA*, **86**, 70105


Reusser, F. *et al.*, *Biochemistry*, 1977, **16**, 3406 (*props*)

Kuo, M.-S. *et al.*, *J. Antibiot.*, 1989, **42**, 357 (*pmr*, *cmr*, *struct*)

Ficine

F-53

[2520-36-7]



C₂₀H₁₉NO₄ 337.374

Alkaloid from *Ficus pantoniana* (Moraceae). Proteolytic agent. Shows vermifidal props. Mp 235°. Rearranges to a mixt. with Isoficine, I-233 (in which the pyrrolidiny residue is at C6) in refluxing HCl.

Johns, S.R. *et al.*, *Tet. Lett.*, 1965, 1987 (*uv*, *ir*, *pmr*, *ms*, *struct*)


Ficuseptine

F-54

2,3-Dihydro-6,8-bis(4-methoxyphenyl)-1H-indolizinium(1+), 9CI

[132923-01-4]

[132923-50-3]




C₂₂H₂₂NO₂⁺ 332.421

Alkaloid from the leaves of *Ficus septica* (Moraceae). Shows antimicrobial activity. Cryst. (CH₂Cl₂/2-methoxy-2-methylpropane). Mp 186-187° dec. (as chloride). λ_{max} 244 (ε 7950); 284 (ε 15500); 336 (ε 4000) (MeOH) (Berdy).

Baumgartner, B. *et al.*, *Phytochemistry*, 1990, **29**, 3327 (isol, uv, ir, pmr, cmr, ms, struct)
Bracher, F. *et al.*, *Eur. J. Org. Chem.*, 2002, 2288-2291 (synth)

Ficuseptine C

F-55

C₂₂H₂₁NO₃ 347.413**(R)-form**

Alkaloid from the stems of *Ficus septica*. Gum. [α]_D²⁵ -79.9 (c, 0.05 in MeOH). λ_{max} 257 (log ε 3.46); 282 (log ε 3.42); 340 (log ε 2.68) (MeOH).

7-Methoxy: Ficuseptine BC₂₃H₂₃NO₄ 377.439


Alkaloid from the stems of *Ficus septica*. Gum. [α]_D²⁵ -95.6 (c, 0.1 in MeOH). λ_{max} 255 (log ε 3.8); 261 (log ε 3.91); 281 (log ε 2.59); 340 (log ε 3.24); 358 (log ε 3.18) (MeOH).

Damu, A.G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1071-1075 (isol, cd, pmr, ms)
Fürstner, A. *et al.*, *Chem. Eur. J.*, 2006, **12**, 7398-7410 (synth)

Firefly oxyluciferin

F-56

2-(6-Hydroxy-2-benzothiazolyl)-4(5H)-thiazolone, 9CI. 2-(4-Hydroxy-2-thiazolyl)-6-benzothiazolone, 9CI, 8CI. 4-Hydroxy-2-(6-hydroxy-2-benzothiazolyl)thiazole. 6-Hydroxy-2-(4-hydroxy-2-thiazolyl)benzothiazole. *Luciola oxyluciferin*. *Luciola oxyluciferin* [17002-50-5] [24963-17-5]

C₁₀H₆N₂O₂S₂ 250.302

Tautomeric. Prod. by fireflies, such as the Japanese firefly *Luciola cruciata*. Responsible for firefly luminescence. Orange-yellow powder + H₂O. Mp 169-171° dec. λ_{max} 371 (ε 18900) (MeOH).

6-Ac: [38292-83-0]

Yellow cryst. (MeOH aq.).

Di-Ac: [38292-84-1]

Yellow leaflets (MeOH). Mp 178-181°.

Suzuki, N. *et al.*, *Tetrahedron*, 1972, **28**, 4065-4074; 4075-4082 (synth, ir, pmr, uv, ms, props, 6-Ac, di-Ac)

White, E.H. *et al.*, *J.A.C.S.*, 1980, **102**, 3199-3208 (props)

Suzuki, N. *et al.*, *Heterocycles*, 1983, **20**, 1027-1030 (props)

Kirk-Othmer *Encycl. Chem. Technol.*, 4th edn., Wiley, 1991, **15**, 540 (props)

Gandelman, O.A. *et al.*, *J. Photochem. Photobiol. B*, 1993, **19**, 187-191 (props)


Esteves da Silva, J.C.G. *et al.*, *Tet. Lett.*, 2001, **42**, 8173-8176 (synth, pmr, cmr)

Nakatani, N. *et al.*, *J.A.C.S.*, 2007, **129**, 8756-8765 (tautom, theory)

Fiscalin A

F-57

[149030-11-5]



Absolute Configuration

R = H

C₂₆H₂₇N₅O₄ 473.53


Prod. by the fungus *Neosartorya fischeri*. Substance P inhibitor. Neurokinin binding inhibitor. Cryst. (MeOH/CH₂Cl₂). Sol. MeOH, EtOAc; poorly sol. hexane. Mp 242-243°. [α]_D -169 (c, 0.5 in MeOH). λ_{max} 226 ; 278 ; 306 (prob. MeOH) (Derep).

Wong, S.-M. *et al.*, *J. Antibiot.*, 1993, **46**, 545-553 (isol, pmr, cmr, cryst struct, abs config)

Fiscalin B

F-58

[149008-35-5]



Absolute Configuration

C₂₃H₂₂N₄O₂ 386.452

Prod. by the fungi *Neosartorya fischeri* and *Corynascus setosus*. Substance P inhibitor, neurokinin binding inhibitor. Cryst. (CHCl₃/MeOH). Sol. MeOH, EtOAc; poorly sol. hexane. Mp 164.5-170.5°. [α]_D -124 (c, 0.02 in MeOH). λ_{max} 222 (ε 56000); 268 (ε 15000); 278 (ε 14000); 290 (ε 14500); 310 (ε 5000); 321 (sh) (ε 3500) (no solvent reported) (Derep). λ_{max} 220 (ε 44668); 273 (ε 11480); 290 (ε 8700); 305 (ε 3980); 318 (ε 2950) (MeOH) (Berdy).

Wong, S.-M. *et al.*, *J. Antibiot.*, 1993, **46**, 545-553 (isol, pmr, cmr, abs config)

Fujimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1843-1848 (isol, uv, pmr, cmr)

Wang, H. *et al.*, *J.O.C.*, 2000, **65**, 1022-1030 (synth)

Liu, J.-F. *et al.*, *J.O.C.*, 2005, **70**, 6339-6345 (synth)

Fiscalin C

F-59

[149008-36-6]

As Fiscalin A, F-57 with

R = CH₃C₂₇H₂₉N₅O₄ 487.557


Prod. by the fungus *Neosartorya fischeri*. Substance P inhibitor. Neurokinin binding inhibitor. Amorph. solid. Sol. MeOH, EtOAc; poorly sol. hexane. λ_{max} 226 ; 278 ; 306 (prob. MeOH) (Derep).

Wong, S.-M. *et al.*, *J. Antibiot.*, 1993, **46**, 545-553 (isol, pmr, cmr, props, abs config)

Fischerellin A

F-60

[182227-56-1]



Relative Configuration

C₂₆H₃₆N₂O₂ 408.583

Isol. from the cyanobacterium *Fischerella muscicola*. Potent photosystem-II inhibitor. Exhibits antifungal and herbicidal activity. Powder. Unique struct.; unrelated to all other known cyanobacterial metabolites. λ_{max} 214 (log ε 53000); 229 (log ε 12000); 241 (log ε 16000); 253 (log ε 18000); 268 (log ε 20600); 284 (log ε 24000); 301 (log ε 22000) (no solvent reported).

Hagmann, L. *et al.*, *Tet. Lett.*, 1996, **37**, 6539-6542 (isol, uv, pmr, cmr, struct)

Fischerellin B

F-61

[186901-41-7]



Absolute Configuration

C₂₀H₂₉NO 299.455

Isol. from the cyanobacteria *Fischerella muscicola* and *Fischerella ambigua*. Yellowish solid. λ_{max} 240 ; 252 ; 267 ; 283 (MeOH).


Papke, U. *et al.*, *Tet. Lett.*, 1997, **38**, 379 (isol, uv, pmr, cmr, struct, abs config)

Fischerin

F-62

NFA Toxin

[134822-63-2]



Relative configuration

C₂₃H₂₉NO₇ 431.485

Struct. revised in 1993. Metab. from the fungus *Neosartorya fischeri* var. *fischeri*. Pale yellow amorph. powder. Sol. MeOH, bases; poorly sol. acids. $[\alpha]_D^{26}$ -65 (c, 0.52 in CHCl₃); λ_{\max} 213 (ε 13000); 233 (ε 7700); 290 (ε 4800); 343 (ε 3800) (MeOH) (Berdy).

► Highly toxic, causes lethal peritonitis in mice.

Tri-Ac:


Needles (Me₂CO). Mp 121-123°. $[\alpha]_D^{22}$ -46 (c, 0.44 in CHCl₃).

Fujimoto, H. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1268 (*isol, uv, ir, pmr, cmr, struct*)

Fischerindole L

F-63

[144398-55-0]

C₂₁H₂₃ClN₂ 338.879

The suffix letters for the Fischerindoles are allocated on the basis of analogy with structurally similar Hapalindoles, therefore not all suffix letters are known. Alkaloid from the terrestrial blue-green alga *Fischerella muscicola*. λ_{\max} 220 (ε 38000); 278 (ε 6800); 290 (sh) (ε 5000) (MeOH) (Derep).

10,11-Didehydro-12-Epifischerindole 1 isonitrile

[159189-08-9]

C₂₁H₂₁ClN₂ 336.863

Isol. from *Hapalosiphon welwitschii*.

10-Epimer-12-Epifischerindole G isonitrile

[159249-50-0]

C₂₁H₂₃ClN₂ 338.879

Isol. from *Hapalosiphon welwitschii*.

$[\alpha]_D$ +67 (c, 0.09 in CH₂Cl₂). λ_{\max} 222 (ε 18600); 278 (ε 5880) (MeOH).

10-Epimer, dechloro-12-Epifischerindole U isonitrile

[159189-06-7]

C₂₁H₂₄N₂ 304.434

Isol. from *Hapalosiphon welwitschii*.

λ_{\max} 226 (ε 17500); 278 (ε 5680) (MeOH).

10-Epimer, dechloro, isothiocyanate: 12-Epifischerindole U isothiocyanate

C₂₁H₂₄N₂S 336.5

Isol. from *Hapalosiphon welwitschii*.

$[\alpha]_D$ +231 (c, 0.03 in CH₂Cl₂). Has -NCS replacing -NC. λ_{\max} 206 (sh) (ε 16300); 228 (ε 23000); 278 (ε 5240) (MeOH).

Park, A. *et al.*, *Tet. Lett.*, 1992, **33**, 3257-3260 (*isol, uv, pmr, cmr, ms*)

Stratmann, K. *et al.*, *J.A.C.S.*, 1994, **116**, 9935-9942 (*derivs*)


Richter, J.M. *et al.*, *J.A.C.S.*, 2008, **130**, 17938-17954 (*synth*)

Fissicisine

F-64

3,4,8-Trimethoxy-N,N-dimethyl-1-phenanthreneethanamine, 9CI

[129743-89-1]

C₂₁H₂₅NO₃ 339.433

Alkaloid from the root bark of *Fissistigma glaucescens* (Annonaceae). Amorph. solid.

N-Oxide: Fissicisine N-oxide

[129743-90-4]

C₂₁H₂₅NO₄ 355.433


Alkaloid from the root bark of *Fissistigma glaucescens* (Annonaceae). Yellowish oil.

Wu, Y.-C. *et al.*, *Phytochemistry*, 1990, **29**, 2387 (*isol, uv, pmr, cmr, ms, struct*)

Fissilandione

F-65

[216445-07-7]

C₁₉H₁₅NO₅ 337.331

Alkaloid from *Fissistigma balansae*.

Amorph. violet solid (CHCl₃). Mp 250-252°. λ_{\max} 220 (log ε 4.4); 284 (sh) (log ε 4.07); 306 (log ε 4.15); 333 (log ε 3.96); 590 (log ε 3.3) (EtOH).

N-De-Me: Norfissilandione

[216445-09-9]

C₁₈H₁₃NO₅ 323.304

Alkaloid from *Fissistigma balansae*.

Amorph. violet solid (CHCl₃). Mp 258-261°. λ_{\max} 218 (log ε 4.57); 283 (log ε 4.06); 300 (log ε 4.2); 335 (log ε 3.98); 591 (log ε 3.3) (EtOH).


Chia, Y.-C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1430-1432 (*isol, uv, ir, pmr, cmr, ms*)

Fissistigine B

F-66

2,3,12,12a-Tetrahydro-5,10-dimethoxy-1-methyl-1H-[1]benzoxepino[2,3,4-ij]isoquinolin-8-ol, 9CI

[83685-18-1]

C₁₉H₂₁NO₄ 327.379


Alkaloid from *Fissistigma oldhamii* (Annonaceae).

Xu, C. *et al.*, *Zhongyao Tongbao*, 1982, **7**, 30-31; *CA*, **97**, 212639t

Fissoldhimine

F-67

[163634-04-6]

C₁₄H₂₂N₄O₂ 278.353


Alkaloid from stems of *Fissistigma oldhamii* (Annonaceae). Cryst. (MeOH). Mp 234-236°.

Wu, J.-B. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 2202 (*isol, pmr, cmr, cryst struct*)

Fistularin 1

F-68

[73622-20-5]

C₂₂H₂₁Br₄N₃O₈ 775.039

Metab. of the sponge *Aplysina fistularis* forma *fulva*. Amorph. solid. $[\alpha]_D$ +93.5 (c, 1.2 in MeOH). λ_{\max} 230 (sh) (ε 15200); 284 (sh) (ε 5680) (MeOH) (Derep).


Di-O-Ac: Mp 168-171° dec. $[\alpha]_D$ +122.7 (c, 0.44 in CHCl₃).

Gopichand, Y. *et al.*, *Tet. Lett.*, 1979, 3921 (*isol, uv, ir, pmr, struct*)

Fistularin 2

F-69

[73622-21-6]


C₂₂H₂₁Br₄N₃O₈ 775.039

Metab. of the sponge *Aplysina fistularis* forma *fulva*.

Gopichand, Y. *et al.*, *Tet. Lett.*, 1979, 3921 (*isol, pmr, struct*)

Fistularin 3

[73622-22-7]

C₃₁H₃₀Br₆N₄O₁₁ 1114.022

Complex stereochem. In view of the fact that C-17 configs. remain undetermined, the relationships shown in this entry are tentative. Some samples prev. descr. as Fistularin 3 were mixts. with its 11-epimer, and opt. rotn. is unable to distinguish these. Stereochem. variability occurs between individuals of the same sp. Metab. of the sponge *Aplysina fistularis* forma *fulva*, *Aplysina archeri*, *Agelas oroides*, *Pseudoceratina durissima*, *Verongula* sp., *Verongia aephorba* and *Verongia cavernicola*. Inhibits the growth of feline leukaemia virus. Amorph. solid. $[\alpha]_D^{25} +104.2$ (c, 1.67 in MeOH). λ_{\max} 223 (sh) (€ 26545); 283 (sh) (€ 10387) (MeOH) (Derep).

Tetra-O-Ac: Mp 202-204° dec. $[\alpha]_D^{25} +149.4$ (c, 1.32 in CHCl₃).

11-Ketone: 11-Oxofistularin 3. 11-Ketofistularin 3

[142755-09-7]

C₃₁H₂₈Br₆N₄O₁₁ 1112.006

Metab. of the sponge *Aplysina archeri*. Inhibits the growth of feline leukaemia virus. Pale yellow gum. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{25} +130$ (c, 0.1 in MeOH). λ_{\max} 225 (€ 27000); 242 (€ 14000); 283 (€ 9900) (MeOH) (Berdy).

11-Deoxy: 11-Deoxyfistularin 3

[191112-19-3]

C₃₁H₃₀Br₆N₄O₁₀ 1098.023

Isol. from *Aplysina cavernicola* and *Aplysina fistularis insularis*. Cytotoxic agent. Powder. Mp 128-130°. $[\alpha]_D^{25} +194.2$ (c, 4.12 in MeOH). λ_{\max} 232 (€ 19000); 284 (€ 11000) (MeOH). λ_{\max} 209 (€ 42000); 235 (€ 21600); 283 (€ 11200) (MeOH).

19-Deoxy: 19-Deoxyfistularin 3

[153209-11-1]

C₃₁H₃₀Br₆N₄O₁₀ 1098.023

Metab. from a new sp. of sponge of the family Aplysinellidae, order Verongida, from the Coral Sea. Powder. $[\alpha]_D^{20} +155$ (c, 0.17 in Me₂CO). λ_{\max} 206 (€ 60400); 235 (€ 22000); 283 (€ 10400) (MeOH) (Berdy).

19-Deoxy, 11-ketone: 19-Deoxy-11-oxofistularin 3

[153209-12-2]

C₃₁H₂₈Br₆N₄O₁₀ 1096.007

Metab. from a new sp. of sponge of the family Aplysinellidae, order Verongida, from the Coral Sea. Powder. $[\alpha]_D^{20} +136$ (c, 0.2 in Me₂CO). λ_{\max} 205 (€ 18500);

F-70

280 (€ 3400) (MeOH) (Berdy).

11,19-Dideoxy: 11,19-Dideoxyfistularin 3

[179523-38-7]

C₃₁H₃₀Br₆N₄O₉ 1082.023

Metab. of the verongid sponge *Pseudoceratina durissima* and from *Aplysina lacunosa* and *Verongia cavernicola*. Unstable yellow powder. $[\alpha]_D^{25} +98.5$ (c, 0.1 in MeOH). λ_{\max} 224 (€ 26000); 257 (€ 16000); 284 (€ 10400) (MeOH) (Berdy).

11-Epimer: 11-Epifistularin 3C₃₁H₃₀Br₆N₄O₁₁ 1114.022

Metab. from the marine sponge *Agelas oroides*. Exhibits antibacterial activity and moderate cytotoxicity towards cultured breast cancer cells. Amorph. solid. $[\alpha]_D^{25} +65.2$ (c, 1.04 in Me₂CO). λ_{\max} 233 (€ 13500); 283 (€ 2650) (EtOH) (Berdy).

Stereoisomer: Isofistularin 3

[87099-50-1]

C₃₁H₃₀Br₆N₄O₁₁ 1114.022

Isol. from *Verongia aerophoba* and *Ailochroia crassa*. Cytotoxic agent. Shows antibiotic activity. Sol. Me₂CO, Et₂O. $[\alpha]_D^{25} +108$ (c, 2.75 in MeOH). Isomeric at one or more chiral centres. Genus name often misspelt Ailochroia. λ_{\max} 223 (sh) (€ 26545); 283 (sh) (€ 10387) (MeOH) (Derep).

Stereoisomer (2):C₃₁H₃₀Br₆N₄O₁₁ 1114.022

Isol. from *Verongia aerophoba*. Pale yellow powder. Mp 128°. $[\alpha]_D^{25} +51.6$ (c, 0.6 in MeOH).

Gopichand, Y. et al., *Tet. Lett.*, 1979, **20**, 3921-3924 (*isol, uv, ir, pmr, struct*)

Cimino, G. et al., *Tet. Lett.*, 1983, **24**, 3029-3032 (*isol, pmr, struct, Isofistularin*)

Kernan, M.R. et al., *J. Nat. Prod.*, 1990, **53**,

615-622 (*11,19-Dideoxyfistularin 3*)

Gunasekera, S.P. et al., *J. Nat. Prod.*, 1992, **55**,

509-512 (*11-Oxofistularin 3*)

König, G.M. et al., *Heterocycles*, 1993, **36**,

1351-1358 (*11-Epifistularin 3*)

Mancini, I. et al., *J.C.S. Perkin 1*, 1993, 3121-

3125 (*19-Deoxyfistularin 3, 19-Deoxy-11-*

oxofistularin 3)

Ciminiello, P. et al., *Tetrahedron*, 1997, **53**,

6565-6572 (*11-Deoxyfistularin 3*)

Compagnone, R.S. et al., *J. Nat. Prod.*, 1999,

62, 1443-1444 (*11-Deoxyfistularin 3*)

Gao, H. et al., *Tetrahedron*, 1999, **55**, 9717-

9726 (*isol, cd, pmr, cmr*)

Aydoğmuş, Z. et al., *Turk. J. Chem.*, 1999, **23**,

339-344 (*Isofistularin 3, isol, activity,*

stereoisomer 2)


Rogers, E.W. et al., *J. Nat. Prod.*, 2005, **68**,

891-896 (*stereochem*)

Flabellidine

Alkaloid L5†

F-71

C₁₈H₂₈N₂O 288.432**Natural-form [6900-86-3]**

Alkaloid from *Lycopodium flabelliforme*, *Lycopodium paniculatum*, *Lycopodium thyoides* and *Lycopodium complanatum* (Lycopodiaceae). Cryst. (Me₂CO)(as perchlorate). Mp 281-282° (perchlorate).

Ac:


Cryst. (Me₂CO/Et₂O). Mp 150-152°.**(±)-form**

Ac:

Cryst. (Et₂O/Me₂CO). Mp 140°.Manske, R.H.F. et al., *Can. J. Res., Sect. B*,1942, **20**, 87 (*isol*)Alam, S.N. et al., *Can. J. Chem.*, 1964, **42**,2456 (*uv, ir, pmr, ms, struct*)Braekman, J.C. et al., *Phytochemistry*, 1974,**13**, 2519 (*isol*)Morales, G. et al., *Phytochemistry*, 1979, **18**,1719 (*isol*)Schumann, D. et al., *Annalen*, 1982, 2057(*synth, pmr, cmr, deriv*)**Flabelliformide A**

F-72

[57567-16-5]



Relative Configuration

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from the stems of *Ervatamia flabelliformis*. Needles (MeOH). Mp 264-265°. $[\alpha]_D^{25} -68.8$ (c, 0.1 in EtOH).

15,20-Diepimer: Flabelliformide B

[953423-34-2]

C₂₀H₂₂N₂O₂ 322.406


Alkaloid from the stems of *Ervatamia flabelliformis*. Powder. Mp 252-253°. $[\alpha]_D^{25} -23.3$ (c, 0.1 in EtOH).

Liang, S. et al., *Helv. Chim. Acta*, 2007, **90**, 1467-1470 (*isol, pmr, cmr, ms, cryst struct*)

Flabelliformine†

F-73

Dodecahydro-1-hydroxy-11-methyl-1,9-ethanobenzo[i]quinolizin-14-one, 13CI. 14-Hydroxy-15-methyllycopodan-5-one, 9CI. Clavatine [7096-84-6]



Absolute Configuration

C₁₆H₂₅NO₂ 263.379

Identity of Clavatine and Flabelliformine not definitely establ. Alkaloid from *Lycopodium flabelliforme*, *Lycopodium clavatum* var. *megastachyon* and *Lycopodium lucidulum* (Lycopodiaceae). Mp 210-211° (change in cryst. form at 150°). $[\alpha]_D^{20} -365.7$ (Me₂CO) (Clavatine). Forms a 1:1 mol. complex with Lycopodine. Mp 213-214°, *isol.* from *L. clavatum*.

Hydrobromide: Mp 337° dec.*Methodide*: Mp 335° dec. (317-318°).*1:1 Complex with Lycodoline, L-325*:

*Clavatoxine*C₃₂H₅₀N₂O₄ 526.758Isol. from *Lycopodium clavatum* (Lycopodiaceae). Needles (petrol). Mp 185-186°.6 α -Hydroxy: **4,6 α -Dihydroxylycopodine**6 α -Hydroxyflabelliformine

[791588-39-1]

C₁₆H₂₅N₃O₃ 279.378Alkaloid from *Huperzia serrata*.Amorph. powder. Mp >300° dec. [α]_D²⁵ -56.8 (c, 0.16 in MeOH).

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1938, **18**, 88-95 (*Clavatoxine, isol*)
 Curcumelli-Rodostamo, M. *et al.*, *Can. J. Chem.*, 1962, **40**, 1068-1070 (*isol, uv, ir, pmr, struct*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1962, **40**, 2088-2100; 1969, **47**, 449-455 (*isol*)

Achmatowicz, O. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1964, **12**, 311-313; *CA*, **61**, 13358f (*Clavatine, struct*)

Rodewald, W.J. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1967, **15**, 579-581; *CA*, **68**, 69166e (*Clavatoxine, struct*)


Nakashima, T.T. *et al.*, *Can. J. Chem.*, 1975, **53**, 1936-1942 (*cmr*)

Tan, C.-H. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1963-1967 (*4,6-Dihydroxylycopodine*)

Flabelliformine†

F-74

[1020085-49-7]



Absolute Configuration


C₂₁H₂₆N₂O₃ 354.448Alkaloid from the stems of *Ervatamia flabelliformis*. Powder (MeOH). [α]_D²⁵ -29 (c, 0.2 in EtOH).

Liang, S. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 239-243 (*isol, pmr, cmr, ms*)

Flabelline

F-75

[6899-89-4]



Absolute Configuration

C₁₈H₂₈N₂O 288.432Alkaloid from *Lycopodium flabelliforme* (Lycopodiaceae). Mp 185-187.5°.

Perchlorate: Mp 280-282°.

Methiodide: Mp 281-282°.

N-De-Ac, N-oxalyl: **Huperzine G**

[220205-68-5]

C₁₈H₂₆N₂O₃ 318.415Alkaloid from *Huperzia serrata*. Mp 285°.12-Hydroxy: **Lycoverticine**

[62023-86-3]

C₁₈H₂₈N₂O₂ 304.431Alkaloid from *Lycopodium verticillatum* (Lycopodiaceae). Mp 222-225°.[α]_D -61 (EtOH).

Young, J.C.F. *et al.*, *Can. J. Chem.*, 1963, **41**, 2731 (*isol, ir, pmr, struct*)

Alam, S.N. *et al.*, *Can. J. Chem.*, 1964, **42**, 2456 (*ms*)


Nyembo, L. *et al.*, *Bull. Soc. Chim. Belg.*, 1976, **85**, 595 (*Lycoverticine*)

Wang, B.-D. *et al.*, *Zhiviu Xuebao (Acta Bot. Sin.)*, 1998, **40**, 842-845 (*Huperzine G*)

Flavacol

F-76

3,6-Bis(2-methylpropyl)-2(1H)-pyrazinone, 9CI. 3,6-Diisobutyl-2(1H)-pyrazinone [495-98-7]

C₁₂H₂₀N₂O 208.303Metab. from *Aspergillus flavus* and *Aspergillus ochraceus*. Needles (EtOAc). Mp 149-150° Mp 144.5-147°. λ _{max} 229 (ε 4290); 325 (ε 4660) (EtOH).1-Oxide: **Neoaspergillilic acid**

[2152-59-2]

C₁₂H₂₀N₂O₂ 224.302

Metab. of *Aspergillus sclerotiorum*, *Aspergillus flavus* and *Aspergillus ochraceus*. Shows antibiotic activity. Needles (MeOH aq.). Sol. MeOH, Et₂O, bases; poorly sol. H₂O, hexane. Mp 127.5-128.5°. λ _{max} 234 (ε 10557); 330 (ε 8143) (EtOH). λ _{max} 236 (ε 9150); 328 (ε 10500) (EtOH) (Berdy).
 ▶ LD₅₀ (mus, ipr) 125 mg/kg.

1-Oxide, Fe complex: **Ferrineoaspergillin**

[67269-53-8]

C₃₆H₅₇FeN₆O₆ 725.73

Pigment from *Aspergillus ochraceus*, *Aspergillus parasiticus*, *Aspergillus auricomus*, *Aspergillus melleus*, *Aspergillus sclerotiorum* and *Aspergillus sulphureus*. Red leaflets (Me₂CO aq.). Insol. H₂O. Mp 135-136° (129-133°). λ _{max} 227; 315; 410 (EtOH).

2''-Hydroxy: **Deoxyneo- β -hydroxyaspergillilic acid**. 6-(2-Hydroxy-2-methylpropyl)-3-(2-methylpropyl)-2(1H)-pyrazinone, 9CI

[40034-14-8]

C₁₂H₂₀N₂O₂ 224.302

Metab. from *Aspergillus ochraceus*. Chelating agent. Needles (EtOAc or diisopropyl ether). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 118-119.5°. Mp 122.5-123° (Deoxyneo- β -hydroxyaspergillilic acid). λ _{max} 230 (ε 5016); 326 (ε 5528) (EtOH).

NH-form1,1''- ξ -Dihydroxy: **Neohydroxyaspergillilic acid**

[72598-34-6]

C₁₂H₂₀N₂O₃ 240.302

Metab. of *Aspergillus sclerotiorum* and *Aspergillus flavus*. Exhibits antibiotic and antiphae activity. Light-yellow needles (MeOH aq.). Sol. MeOH, Et₂O, bases; poorly sol. H₂O, hexane.

Mp 170-171° (164-166°). [α]_D²⁵ -57 (c, 0.64 in EtOH). Dimorphic. λ _{max} 236 (ε 6450); 328 (ε 8350) (EtOH) (Berdy). λ _{max} 246 (ε 8000); 360 (ε 7550) (HCl) (Berdy).

▶ LD₅₀ (mus, ipr) 125 mg/kg.1,2''-Dihydroxy: **β -Hydroxyneoaspergillilic acid**

[62309-40-4]

C₁₂H₂₀N₂O₃ 240.302Metab. of *Aspergillus ochraceus*.

Amorph. solid (EtOAc). Mp 143-144°.

OH-form*Me ether*: 3-Methoxy-2,5-bis(2-methylpropyl)pyrazine

[36329-97-2]

C₁₃H₂₂N₂O 222.33Prod. by *Chondromyces crocatus*.

Dunn, G. *et al.*, *J.C.S.*, 1949, 2586 (*synth, struct*)

Weiss, U. *et al.*, *Arch. Biochem. Biophys.*, 1958, **74**, 150 (*Neohydroxyaspergillilic acid*)

Micetich, R.G. *et al.*, *J.C.S.*, 1964, 1507 (*isol, derivs*)

Masaki, M. *et al.*, *Tet. Lett.*, 1965, 4837 (*synth, uv, ir, deriv*)

Yamazaki, M. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 2274-2276 (*Aspergillus ochraceus, isol, uv, ir, pmr*)

Maebayashi, Y. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1320 (*Ferrineoaspergillin*)

Ohta, A. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 1316; 1378 (*deriv, synth*)


Assante, G. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 785 (*isol, uv, ir, ms, derivs*)

Okada, Y. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2259-2262 (*synth*)

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*Me ether*)

Aglaia dasyclada Flavagline compound 1


F-77

C₄₂H₅₂N₂O₁₄ 808.878Alkaloid from *Aglaia dasyclada*.Amorph. [α]_D²⁵ -22.4 (c, 1 in MeOH).Isomer of *Aglaia dasyclada* Flavagline compound 2, F-78.

Chaidir, *et al.*, *J. Nat. Prod.*, 2001, **64**, 1216-1220

Aglaia dasyclada Flavagline compound 2


F-78

C₄₂H₅₂N₂O₁₄ 808.878Alkaloid from *Aglaia dasyclada*.Amorph. [α]_D²⁵ -28.2 (c, 0.04 in MeOH).Isomer of *Aglaia dasyclada* Flavagline compound 1, F-77.

Chaidir, *et al.*, *J. Nat. Prod.*, 2001, **64**, 1216-1220 (*isol, pmr, cmr*)

Flavine adenine dinucleotide F-79

Riboflavin 5'-(trihydrogen diphosphate) 5' → 5'-diester with adenosine, 9CI. Flavinat. Flavinat. Isoalloxazine adenine dinucleotide. FAD. Fademin. Flamitajin. Flavin F
[146-14-5]



$C_{27}H_{33}N_9O_{15}P_2$ 785.556
Isol. from various natural sources incl. the liver, heart, kidney, muscles and yeasts. Vitamin B₂ therapeutic agent. Prosthetic group of various flavoproteins incl. D-amino acid oxidase, fumaric hydrogenase, histaminase glucose oxidase and xanthine oxidase. Yellow microcryst. (H₂O). Light-sensitive.

▶ AU7470000

[1910-41-4, 84366-81-4]

Christie, S.M.H. *et al.*, *J.C.S.*, 1954, 46 (*synth*)
Moffatt, J.G. *et al.*, *J.A.C.S.*, 1958, **80**, 3756 (*synth*)

Yagi, K. *et al.*, *Biochem. Prep.*, 1959, **7**, 51 (*isol*)

Boyer, P.D. *et al.*, *Enzymes*, Ed., Vol. 2, Academic Press, New York, 1960, (*rev*)

Kainosho, M. *et al.*, *Biochemistry*, 1972, **11**, 741 (*pmr*, P-31 *nmr*)

Yagi, K. *et al.*, *Biochemistry*, 1976, **15**, 2877 (*N-15 nmr*)

Yoshimura, J. *et al.*, *Method. Chim.*, 1977, **11**, 76 (*rev*)

Holmgren, A. *et al.*, *Experientia*, Suppl., 1980, **36**, 149 (*rev*)

Shimizu, S. *et al.*, *Appl. Biochem. Biotechnol.*, 1983, **8**, 237 (*synth*)

Ulrich, E.L. *et al.*, *Tet. Lett.*, 1983, **24**, 473 (*pmr*)


Otvos, J.D. *et al.*, *Biochemistry*, 1986, **25**, 7220 (*P-31 nmr*)

Copeland, R.A. *et al.*, *J. Phys. Chem.*, 1986, **90**, 6648 (*uv*, *Raman*)

Nielsen, P. *et al.*, *Methods Enzymol.*, 1986, **122**, 209 (*rev*, *synth*, *props*)

Flavinine F-80

5,6,8,14-Tetrahydro-3-hydroxy-2,6-dimethoxymorphinan-7-one, 9CI
[19777-83-4]



(-)-form

 $C_{18}H_{19}NO_4$ 313.352**(+)-form**N-Me: (+)-**Flavinantine** $C_{19}H_{21}NO_4$ 327.379Alkaloid from *Roemeria refracta*.Amorph. $[\alpha]_D^{25} +13$ (c, 0.13 in MeOH). $[\alpha]_D^{25} +30$ (c, 0.1 in CHCl₃). λ_{max} 240 (log ϵ 4.12); 285 (log ϵ 3.81) (MeOH).

Me ether: N-Nor-2,3,6-trimethoxymorphinandien-7-one

[149155-13-5]

 $C_{19}H_{21}NO_4$ 327.379Alkaloid from stems of *Fissistigma oldhamii* (Annonaceae). Amorph. pale yellow solid. Mp 126-128°. $[\alpha]_D^{25} 0$ (c, 0.02 in CHCl₃).

Me ether, N-Me: O-Methylpallidine. Fissistigine C

[27510-33-4]

[83685-19-2]

 $C_{20}H_{23}NO_4$ 341.406Alkaloid from leaves of *Ocotea acutangula*, *Ceratocarpus palaestinus* and the bark of *Fissistigma oldhamii* (Annonaceae). Needles (Et₂O) or amorph. Mp 118-120°. $[\alpha]_D^{20} +25.2$ (CHCl₃) (+12). Struct. of Fissistigine C revised in 1985. The lower opt. rotn. refers to synthetic material (amorph.). λ_{max} 208 (log ϵ 4.49); 238 (log ϵ 4.19); 280 (log ϵ 3.82) (EtOH).

Me ether, N-Me, N-oxide: O-Methylpallidine N-oxide

[122249-94-9]

 $C_{20}H_{23}NO_5$ 357.405Alkaloid from the whole plant of *Sarcocapnos emeaphylla* (Papaveraceae). Amorph. $[\alpha]_D^{20} +20$ (c, 2.1 in CHCl₃). The first morphinandienone N-oxide.O²-De-Me, O³-Me: Norpallidine

[61774-73-0]

 $C_{18}H_{19}NO_4$ 313.352Alkaloid from *Fumaria vaillantii* (Papaveraceae). Cryst. (CHCl₃). Mp 102°. $[\alpha]_D^{23} -11$ (MeOH).O²-De-Me, O³-Me, N-Me: Pallidine†

[25650-75-3]

 $C_{19}H_{21}NO_4$ 327.379Alkaloid from *Chasmanthera dependens*, *Corydalis pallida*, *Corydalis incisa*, *Corydalis koidzumiana*, *Desmos tiebaghiensis*, *Monodora crispata*, *Ocotea acutangula*, *Ocotea brachybotra*, *Rollinia mucosa*, *Thalictrum dioicum* and *Thalictrum faberi*. Pale yellow syrup. $[\alpha]_D^{18} -32$ (c, 0.16 in EtOH). λ_{max} 210 (log ϵ 4.51); 240 (log ϵ 4.2); 286 (log ϵ 3.91) (EtOH).O²-De-Me, O³-Me, N-Me, N-oxide: Pallidine N-oxide

[937018-74-1]

 $C_{19}H_{21}NO_5$ 343.379Alkaloid from the leaves of *Neolitsea sericea* var. *aurata*.**(-)-form**Alkaloid from *Croton flavens* (Euphorbiaceae). Cryst. + Me₂CO (Me₂CO). Mp 130-132°. $[\alpha]_D^{15} -6$ (EtOH).N-Me: (-)-**Flavinantine**

[19777-82-3]

 $C_{19}H_{21}NO_4$ 327.379Alkaloid from *Croton flavens* (Eu-phorbiaceae). Rods (EtOAc). Mp 130-132°. $[\alpha]_D^{23} -14.5$ (c, 1.1 in EtOH).Me ether, N-Me: **Sebiferine**. O-Methylflavinantine. Probovatine


[23979-25-1]

 $C_{20}H_{23}NO_4$ 341.406From *Litsea sebifera*, *Alseodaphne perakensis*, *Cocculus laurifolius*, *Duguetia obovata*, *Kolobopetalum auriculatum* and *Rhigiocarya racemifera*. Inhibitor of peristaltic reflex in guinea pig ileum. Shows morphine-like centrally-mediated antinociceptive activity. Mp 112-113°. $[\alpha]_D^{24} -13$ (c, 0.1 in CHCl₃) (synthetic). Probovatine, isol. from *D. obovata* (1980) was assigned a different structure, making it the only member of the proposed group of neoproporphine alkaloids. This assignment was erroneous and on reexamination Sebiferine was isol.Me ether, N-Me, N-oxide: **Sebiferine N-oxide**. O-Methylflavinantine N-oxide

[135357-27-6]

 $C_{20}H_{23}NO_5$ 357.405Alkaloid from *Alseodaphne perakensis* (Lauraceae). Pale yellow solid.**(±)-form** [22324-01-2]Cryst. (Me₂CO). Mp 138°.Stuart, K.L. *et al.*, *J.C.S. (C)*, 1969, 1681-1684 (*isol*, *struct*, *cd*)Kotani, E. *et al.*, *Tet. Lett.*, 1973, 4759-4761 (*synth*)Shamma, M. *et al.*, *Phytochemistry*, 1976, **15**, 1802-1803 (*Norpallidine*)Kametani, T. *et al.*, *J.A.C.S.*, 1977, **99**, 3805-3808 (*synth*)Bhakuni, D.S. *et al.*, *Tetrahedron*, 1979, **35**, 2365-2367 (*biosynth*)Vecchiotti, V. *et al.*, *J.C.S. Perkin 1*, 1981, 578-581 (*O-Methylpallidine*)Dubourg, P.A. *et al.*, *Acta Cryst. B*, 1982, **38**, 1657 (*Sebiferine*, *cryst struct*)Xu, C. *et al.*, *Zhongyao Tongbao*, 1982, **7**, 30-31; *CA*, **97**, 212639t (*Fissistigine C*)Chiaroni, A. *et al.*, *Acta Cryst. C*, 1983, **39**, 1311-1312 (*cryst struct*)Blaskó, G. *et al.*, *J.O.C.*, 1984, **49**, 1439-1441 (*synth*)Tojo, E. *et al.*, *J. Nat. Prod.*, 1989, **52**, 415-416 (*O-Methylpallidine N-oxide*)Gözler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 986-988 ((+)-*Flavinantine*)Lajis, N.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 612-614 (*Sebiferine oxide*)Wu, J.B. *et al.*, *Planta Med.*, 1993, **59**, 179-180 (*N-Nor-2,3,6-trimethoxymorphinandienone*)Hara, H. *et al.*, *Tetrahedron: Asymmetry*, 1995, **6**, 1683-1694 (*Sebiferine*, *synth*)Lee, S.-S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 637-642 (*Pallidine N-oxide*)**Flavipucine****F-81**

6-Methyl-2-(3-methyl-1-oxobutyl)-1-oxa-5-azaspiro[2.5]oct-6-ene-4,8-dione, 9CI. Glutamincine. Glutamycin. FRT-B. Fruit rot toxin B



Relative configuration

 $C_{12}H_{15}NO_4$ 237.255

λ_{\max} (dec. slowly) (Derep). λ_{\max} 330 (ε 5400) (95% EtOH) (Derep).

(-)-form [38473-18-6]

Prod. by *Aspergillus flavipes* and *Macrospora* sp. Phytotoxin. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 130-131°. [α]_D²¹ -88 (c, 1 in EtOH). λ_{\max} 326 (ε 7900) (MeOH) (Berdy). λ_{\max} 330 (ε 5400) (EtOH) (Berdy).

(±)-form [61248-17-7]

Synthetic. Mp 153-155°.

Findlay, J.A. et al., *J.C.S. Perkin 1*, 1972, 2071

(ir, uv, ms, pmr, struct)

Girotra, N.N. et al., *Chem. Comm.*, 1976, 566

(synth)

Findlay, J.A. et al., *Synth. Commun.*, 1977, 7, 149 (synth)

White, P.S. et al., *Can. J. Chem.*, 1978, 56, 1904 (cryst struct)

Girotra, N.N. et al., *Heterocycles*, 1978, 9, 417 (synth)


Sassa, T. et al., *Agric. Biol. Chem.*, 1983, 47, 1155; 1417 (isol, uv, ir, pmr)

Wagner, C. et al., *Z. Naturforsch., C*, 1995, 50, 358 (isol, props)

Flavocarpine

F-82

2-Carboxy-3-ethyl-12H-indolo[2,3-a]quinolizin-5-ium inner salt, 9CI [6879-51-2]



C₁₈H₁₄N₂O₂ 290.321

A nor-corynanthe alkaloid. Zwitterionic form predominates for the parent compd. Alkaloid from the stem bark of *Pleio-carpa mutica* (Apocynaceae). Cryst. (AcOH). Mp 307° dec. (evac. tube) Mp 323-325° dec. (synthetic). λ_{\max} 223 (ε 40800); 242 (ε 48600); 250 (ε 51200); 291 (ε 23000); 351 (ε 23500); 389 (ε 22000) (EtOH).

Hydrochloride: Mp 301-302° dec.

Me ester:

Red needles (EtOH). Mp 152-154°.

5,6-Dihydro: **Dihydrovincarpine**

[62926-61-8]

C₁₈H₁₆N₂O₂ 292.337

Minor alkaloid from *Vinca major* var. *elegantissima* (Apocynaceae). Yellow needles (MeOH/CHCl₃). Mp 280° dec. λ_{\max} 222 (sh) (log ε 4.22); 250 (sh) (log ε 3.77); 315 (log ε 3.81); 395 (log ε 3.73) (EtOH).

5,6-Dihydro, 18,19-didehydro: **Vincarpine**

[62926-60-7]

C₁₈H₁₄N₂O₂ 290.321

Minor alkaloid from *Vinca major* var. *elegantissima* (Apocynaceae). Yellow needles (MeOH/CHCl₃). Mp 300° dec. λ_{\max} 222 (sh) (log ε 4.22); 250 (sh) (log ε 3.77); 315 (log ε 3.81); 395 (log ε 3.73) (EtOH).

Büchi, G. et al., *J.A.C.S.*, 1962, 84, 3393-3398 (Flavocarpine, sol, uv, ir, ms, pmr, struct)


Ali, E. et al., *Tet. Lett.*, 1976, 15, 4887-4890

(Vincarpine, Dihydrovincarpine)

Gribble, G.W. et al., *Tet. Lett.*, 1987, 28, 5259-5262 (Flavocarpine, synth)

Flavolipids

F-83




A complex of at least 37 components having different saturated or unsaturated amide groups. Struct. of major component is shown. Related to Aerobactin. Prod. by soil isolate *Flavobacterium* sp. strain MTN 11. Biosurfactants.

Bodour, A.A. et al., *Appl. Environ. Microbiol.*, 2004, 70, 114-120 (isol, pmr, cmr, ms)

Flavopereirine

F-84

3-Ethyl-12H-indolo[2,3-a]quinolizin-5-ium, 9CI. Melinonine G [6784-38-9]



C₁₇H₁₄N₂ 246.311

A bisnor-corynanthe alkaloid. On biogenetic numbering, 6,7- are 5,6-respectively. Alkaloid from the bark of *Geissospermum laeve*, *Geissospermum vellosii*, *Strychnos melinoniana* and *Strychnos longicaudata* (Apocynaceae, Loganiaceae). Inhibits cancer cell DNA synthesis.

Perchlorate: Mp 317-318° dec. (301-305°).

Picrate: Mp 229.5-230.5°.

6,7-Dihydro: **6,7-Dihydroflavopereirine** [42281-83-4]

C₁₇H₁₆N₂ 248.327

Alkaloid from *Strychnos usambarensis* (Loganiaceae). Poorly sol. hexane. Mp 305-307° dec.

6,7-Dihydro; *perchlorate*: Mp 281-282°.

6,7-Dihydro; *hydrobromide*: Mp 317-320° dec.

Bejar, O. et al., *C. R. Hebd. Seances Acad. Sci.*, 1957, 244, 2066 (isol, struct)

Bächli, E. et al., *Helv. Chim. Acta*, 1957, 40, 1167 (isol, uv, ir, struct)

Angenot, L. et al., *Planta Med.*, 1973, 23, 226 (6,7-Dihydroflavopereirine)

Massiot, G. et al., *Tetrahedron*, 1983, 39, 3645 (isol, uv)

Giri, V.S. et al., *Heterocycles*, 1984, 22, 233 (synth, uv, ir, pmr)

Ninomiya, I. et al., *J.C.S. Perkin 1*, 1984, 2035 (synth)

Gribble, G.W. et al., *Tet. Lett.*, 1987, 28, 5259 (synth, Flavopereirine, Dihydroflavopereirine)

Sankar, P.J. et al., *Heterocycles*, 1991, 32, 1109 (synth)

Fürstner, A. et al., *Tetrahedron*, 1995, 51, 773 (synth, Flavopereirine, Dihydroflavopereirine)


Lounasmaa, M. et al., *Heterocycles*, 1996, 43, 1365 (synth, Flavopereirine, Dihydroflavopereirine)

Manna, R.K. et al., *J. Chem. Res., Synop.*, 1999, 350-351 (synth)

Flazine

F-85

1-[5-(Hydroxymethyl)-2-furanyl]-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI [100041-05-2]



C₁₇H₁₂N₂O₄ 308.293

Constit. of the seeds of *Brucea javanica*. Prod. by the marine bacterium strain Bio215. Cryst. (MeOH). Mp 231-233°. Fluorescent. λ_{\max} 217 (log ε 4.87); 275 (log ε 4.85); 365 (log ε 3.02); 379 (log ε 3.11) (MeOH).

Me ester: **Flazine methyl ester**

[104537-94-2]

C₁₈H₁₄N₂O₄ 322.32

Prod. by *Streptomyces* sp. K01-0031. Mp 199-200°.

Me ether: **Flazine methyl ether. O-**

Methylflazine

[159898-11-0]

C₁₈H₁₄N₂O₄ 322.32

Alkaloid from freshly pressed juice of blackcurrant (*Ribes nigrum*). Mp 199-200°.

De(hydroxymethyl): 1-(2-Furanyl)-9H-pyrido[3,4-b]indole-3-carboxylic acid.

Dehydroxymethylflazine

[76135-36-9]

C₁₆H₁₀N₂O₃ 278.267

From blackcurrant (*Ribes nigrum*).

Nakatsuka, S. et al., *Tet. Lett.*, 1986, 27, 3399-3402 (isol, uv, ir, pmr, struct)

Gessner, W.P. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, 321, 95-98 (synth, uv, pmr)

Blech, S. et al., *Z. Naturforsch., C*, 1994, 49, 540-544 (Flazine methyl ether, Dihydroxymethylflazine)


Su, B.-N. et al., *Planta Med.*, 2002, 68, 730-733 (isol, pmr, cmr, ms)

Shabaan, M. et al., *Dissertation*, Univ. of Göttingen, 2004, (marine, isol)

Shiomi, K. et al., *J. Antibiot.*, 2005, 58, 74-78 (Me ester, isol)

Flephilone

[183239-76-1]

C₂₄H₂₇NO₇ 441.48

Prod. by *Trichoderma harzianum*. Rev protein binding inhibitor. Yellow powder. Sol. MeOH, CHCl₃, CH₂Cl₂; poorly sol. H₂O, hexane. λ_{max} 209 (ε 2250); 229 (ε 1280); 286 (ε 710); 389 (ε 570) (MeOH).

Qian-Cutrone, J. *et al.*, *J. Antibiot.*, 1996, **49**, 990 (*isol, uv, ir, pmr, cmr*)

Flexamine

[1357-10-4]

C₁₇H₁₉NO₅ 317.341


Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine flexuosa* (Amaryllidaceae). Mp 226-228° dec. [α]_D 0 (c, 0.2 in CHCl₃).

Methiodide: Mp 245° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109 (*isol*)

Flexicorine

[80765-85-1]



Absolute Configuration

C₄₁H₄₄N₄O₅ 672.822

Alkaloid from the leaves of *Rauwolfia reflexa* (Apocynaceae). Amorph. red solid. Mp 360°. [α]_D²⁵ -519.5 (CHCl₃).

Chatterjee, A. *et al.*, *J.O.C.*, 1982, **47**, 1732-1734 (*cmr, struct*)

FlexineC₁₇H₁₉NO₅ 317.341

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine flexuosa* (Amaryllidaceae). Cryst. (Me₂CO). Mp 212°. [α]_D²⁵ +75 (c, 0.2 in CHCl₃).

Perchlorate: Mp 242° dec.


Methiodide: Mp 221° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 470-471 (*isol*)

F-86

Flindersine

2,6-Dihydro-2,2-dimethyl-5H-pyrano[3,2-c]quinolin-5-one, 9CI
[523-64-8]

C₁₄H₁₃NO₂ 227.262

Alkaloid from *Flindersia australis*, *Fagara holtziana*, *Fagara cholybea*, *Haplophyllum perforatum* and several other spp. (Flindersiaceae, Rutaceae). Phytotoxic. Mp 195-196° dec. λ_{max} 235; 333; 350; 365 (MeOH) (Berdy).

N-Me: N-Methylflindersine

[50333-13-6]

C₁₅H₁₅NO₂ 241.289

Alkaloid from *Fagara chalybea*, *Fagara holtziana*, *Melicope ptefolia*, *Atalantia roxburghiana* and several other spp. (Rutaceae). Phytotoxin, insecticide, larvicide. Mp 83-85°. λ_{max} 226 (ε 83176); 274 (ε 13800); 318 (ε 13490) (MeOH) (Berdy).

N-(Hydroxymethyl): N-Hydroxymethylflindersine

[96400-50-9]

C₁₅H₁₅NO₃ 257.288

Alkaloid from the stems and leaves of *Haplophyllum buxbaumii* (Rutaceae). Amorph. Possibly an artifact derived from Haplophylline during extraction.

N-(Acetoxymethyl): N-Acetoxymethylflindersine

[149998-45-8]

C₁₇H₁₇NO₄ 299.326

Alkaloid from leaves of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil.

N-(3-Methyl-2-butenoyloxymethyl):

Haplophylline

[93710-20-4]

C₂₀H₂₁NO₄ 339.39

Alkaloid from the aerial parts of *Haplophyllum suaveolens* (Rutaceae). Amorph.

1',2'-Dihydro, 1'ξ-hydroxy: 3',4'-Dihydro-4'-hydroxyflindersine. 4'-Hydroxy-3',4'-dihydroflindersine

[100019-40-7]

C₁₄H₁₅NO₃ 245.277

Alkaloid from the leaves of *Geijera balansae* (Rutaceae). Prisms (EtOAc). Mp 208°. [α]_D²⁰ -6 (c, 1 in CHCl₃).

1',2'-Dihydro, 1'ξ-hydroxy, N-Me: 2,3,4,6-Tetrahydro-4-hydroxy-2,2,6-trimethyl-5H-pyrano[3,2-c]quinolin-5-one

[80357-90-0]

C₁₅H₁₇NO₃ 259.304

Alkaloid from the heartwood of *Euxylophora paraënsis* (Rutaceae). Pale yellow needles (MeOH). Mp 159-160°. [α]_D²⁰ +5.45 (c, 0.4 in CHCl₃). Not interconverted with its

F-90

N-de-Me deriv.

1',2'-Dihydro, 2'ξ-hydroxy, N-Me: 2,3,4,6-Tetrahydro-3-hydroxy-2,2,6-trimethyl-5H-pyrano[3,2-c]quinolin-5-one

[928008-45-1]

C₁₅H₁₇NO₃ 259.304

Alkaloid from *Skimmia laureola*. Amorph. solid. [α]_D²⁹ -57 (c, 0.14 in MeOH). λ_{max} 265 (log ε 2.67); 304 (log ε 3.84); 318 (log ε 3.86); 325 (log ε 3.74); 330 (log ε 3.76) (MeOH).

1',2'-Dihydro, 1'R,2'S-dihydroxy: 2,3,4,6-Tetrahydro-3,4-dihydroxy-2,2-dimethyl-5H-pyrano[3,2-c]quinolin-5-one. 3',4'-Dihydroxy-3',4'-dihydroflindersine

[99964-48-4, 100019-42-9, 100019-41-8]

C₁₄H₁₅NO₄ 261.277

Alkaloid from the leaves of *Geijera balansae* (Rutaceae). Prisms (CH₂Cl₂).

Mp 234-235°. [α]_D²⁰ -2 (c, 0.5 in CHCl₃).

1',2'-Dihydro, 1'R,2'S-dihydroxy, N-Me: **Orixalone D**C₁₅H₁₇NO₄ 275.304

Alkaloid from the stems of *Orixa japonica*. Oil. [α]_D²⁴ +3.1 (c, 0.06 in MeOH). λ_{max} 230; 274; 284; 318; 332 (MeOH).

8-Hydroxy, N-(acetoxymethyl): **Desmethylzanthophylline**

[65560-24-9]

C₁₇H₁₇NO₅ 315.325

Alkaloid from stems and branches of *Zanthoxylum monophyllum* (Rutaceae). Noncryst.

8-Hydroxy, 3,4-dihydro, N-Me: **Ravesilone**

[94935-96-3]

C₁₅H₁₇NO₃ 259.304

Alkaloid from the leaves of *Ravenia spectabilis* (Rutaceae). Cryst. (EtOH/petrol). Mp 272°.

5-Methoxy: 5-Methoxyflindersine. **Haplophytine A**

C₁₅H₁₅NO₃ 257.288

Alkaloid from *Haplophyllum acutifolium*. Powder. Mp 209-210.5°. λ_{max} 244 (log ε 7.92) (CHCl₃).

6-Methoxy: **Haplamine**. 6-Methoxyflindersine

[52617-31-9]

C₁₅H₁₅NO₃ 257.288

Alkaloid from above-ground parts of *Haplophyllum perforatum* (Rutaceae). Cryst. (MeOH). Mp 201-202°.

6-Methoxy, N-Me: **N-Methylhaplamine**. 6-Methoxy-N-methylflindersine

[129225-33-8]

C₁₆H₁₇NO₃ 271.315

Alkaloid from the aerial parts of an *Agathosma* sp. (Rutaceae). Yellow plates (Et₂O). Mp 136-138°.

6-Methoxy, 1',2'-dihydro: **Dihydrohaplamine**

[61929-96-2]

C₁₅H₁₇NO₃ 259.304

Alkaloid from *Haplophyllum perforatum*. Cryst. (Me₂CO). Mp 231-232°.

6-Methoxy, 8-hydroxy: **8-Hydroxy-6-methoxyflindersine**. 7-Hydroxy-9-methoxyflindersine

[155474-84-3]

- C₁₅H₁₅NO₄ 273.288
Alkaloid from *Haplophyllum telephoides* (Rutaceae). Amorph.
- 7-Methoxy: **7-Methoxyflindersine**
[177857-27-1]
C₁₅H₁₅NO₃ 257.288
Alkaloid from leaves of *Vepris bilocularis*. Cryst. (Et₂O/petrol). Mp 194-196°.
- 7-Methoxy, N-Me: **7-Methoxy-N-methylflindersine**
C₁₆H₁₇NO₃ 271.315
Alkaloid from the stem bark of *Oricia renieri* (Rutaceae). Oil.
- 8-Methoxy: **8-Methoxyflindersine**
[35989-00-5]
C₁₅H₁₅NO₃ 257.288
Alkaloid from *Myrtopsis macrocarpa* (Rutaceae). Needles (Et₂O). Mp 178°.
- 8-Methoxy, N-Me: **Zanthobungeanine**
[64190-94-9]
C₁₆H₁₇NO₃ 271.315
Alkaloid from roots of *Zanthoxylum bungeanum* (Rutaceae). Cryst. (C₆H₆). Poorly sol. hexane. Mp 77-78°.
- 8-Methoxy, N-(acetoxymethyl): **Zanthophylline**
[65560-23-8]
C₁₈H₁₉NO₅ 329.352
Alkaloid from stems and branches of *Zanthoxylum monophyllum* (Rutaceae). Cryst. (MeOH aq.). Mp 126-127°.
- 8-Methoxy, 1',2'-dihydro, 1'R,2'R-dihydroxy, N-Me: **Zanthodioline**
[198336-59-3]
[193978-31-3]
C₁₆H₁₉NO₅ 305.33
Alkaloid from the bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Amorph. solid. $[\alpha]_D^{25}$ -20 (c. 0.05 in CHCl₃). λ_{\max} 218 (log ϵ 4.14); 234 (log ϵ 4.21); 254 (log ϵ 4.11); 270 (sh) (log ϵ 3.56); 280 (log ϵ 3.59); 327 (log ϵ 3.23) (EtOH).
- 6,7-Dimethoxy, N-Me: **Oricine**, 6,7-Dimethoxy-N-methylflindersine
[33798-66-2]
C₁₇H₁₉NO₄ 301.341
Alkaloid from the timber of *Oricia suaveolens* and the stem bark of *Oricia renieri* (Rutaceae). Cryst. (C₆H₆). Mp 150-155°.
- 7,8-Dimethoxy, N-Me: **Veprisine**, 7,8-Dimethoxy-N-methylflindersine
[76525-26-3]
C₁₇H₁₉NO₄ 301.341
Alkaloid from the stem bark of *Vepris louisii*, *Vepris stolzii* and *Oricia renieri* (Rutaceae). Needles (C₆H₆/hexane). Mp 89-90°.
- 7,8-Dimethoxy, 1',2'-dihydro, 1'R,2'R-dihydroxy, N-Me: **Araliopsinine**
[120481-37-0]
C₁₇H₂₁NO₆ 335.356
Details not publ. The struct. is that given by Grundon (1990). Alkaloid from stem bark of *Araliopsis tabouensis*.
- 7-(3-Methyl-2-butenyloxy): **7-Prenyloxyflindersine**
[177857-28-2]
C₁₉H₂₁NO₃ 311.38
Alkaloid from leaves of *Vepris bilocularis*. Cryst. (Et₂O/petrol). Mp 165-167°.
- 7-(3-Methyl-2-butenyloxy), N-Me: **N-Methyl-7-prenyloxyflindersine**
[177857-29-3]
C₂₀H₂₃NO₃ 325.407
Alkaloid from leaves of *Vepris bilocularis*. Oil.
- 8-(3-Methyl-2-butenyloxy), N-Me: **8-(3,3-Dimethylallyloxy)-N-methylflindersine**, N-Methyl-8-prenyloxyflindersine
[82504-12-9]
C₂₀H₂₃NO₃ 325.407
Alkaloid from the stem bark of *Vepris stolzii* (Rutaceae). Oil.
- 8-(3-Methyl-2-butenyloxy), 7-methoxy, N-Me: **8-(3,3-Dimethylallyloxy)-7-methoxy-N-methylflindersine**, 7-Methoxy-N-methyl-8-prenyloxyflindersine
[82504-13-0]
C₂₁H₂₅NO₄ 355.433
Alkaloid from the stem bark of *Vepris stolzii* (Rutaceae). Oil.
- 8-(2,3-Epoxy-3-methylbutoxy), 7-methoxy, N-Me: **8-(2,3-Epoxy-3,3-dimethylallyloxy)-7-methoxy-N-methylflindersine**
[82504-14-1]
C₂₁H₂₅NO₅ 371.432
Alkaloid from the stem bark of *Vepris stolzii* (Rutaceae). Oil.
- Brown, R.F.C. et al., *Aust. J. Chem.*, 1954, **7**, 348 (*isol, uv, struct*)
Robertson, A.V. et al., *Aust. J. Chem.*, 1963, **16**, 451 (*pmr*)
Lavie, D. et al., *Tetrahedron*, 1968, **24**, 3011 (*ms*)
Abe, M.O. et al., *Phytochemistry*, 1971, **10**, 1167 (*Oricine*)
Akhmedzhanova, V.I. et al., *Khim. Prir. Soedin.*, 1974, **10**, 109; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 121 (*Haplamine*)
Reisch, J. et al., *Phytochemistry*, 1975, **14**, 1678 (*isol*)
Hifnawy, M.S. et al., *Phytochemistry*, 1977, **16**, 1035 (*8-Methoxyflindersine*)
Stermitz, F.R. et al., *Phytochemistry*, 1977, **16**, 2003 (*Zanthophylline, Desmethylzanthophylline*)
Jurd, L. et al., *Aust. J. Chem.*, 1981, **34**, 1625 (*Tetrahydrohydroxytrimethylpyranoquinoline*)
Khalid, S.A. et al., *Phytochemistry*, 1981, **20**, 2761 (*7-Methoxy-N-methylflindersine, Oricine, Veprisine*)
Ren, L. et al., *Yaouxue Xuebao*, 1981, **16**, 672; *CA*, **96**, 48974e (*isol, struct, Zanthobungeanine*)
Khalid, S.A. et al., *J. Nat. Prod.*, 1982, **45**, 343 (*Veprisine, Dimethylallyloxy-N-methylflindersine, Dimethylallyloxy-7-methoxy-N-methylflindersine, Epoxydimethylallyloxy-7-methoxy-N-methylflindersine*)
Ayafor, J.F. et al., *Phytochemistry*, 1982, **21**, 2733 (*Veprisine*)
Bhattacharyya, P. et al., *Phytochemistry*, 1984, **23**, 1825 (*Ravesilone*)
Ulubelen, A. et al., *Phytochemistry*, 1984, **23**, 2123; 1985, **24**, 372; 1994, **35**, 1600 (*Haplophylline, N-Hydroxymethylflindersine, 8-Hydroxy-6-methoxyflindersine*)
Ramesh, M. et al., *Tetrahedron*, 1984, **40**, 4041 (*synth, ir, pmr, Zanthobungeanine*)
- Mitaku, S. et al., *J. Nat. Prod.*, 1985, **48**, 772 (*Hydroxydihydroflindersine, Dihydroxydihydroflindersine*)
Bellino, A. et al., *Heterocycles*, 1986, **24**, 1821 (*Ravesilone, synth*)
Bravo, P. et al., *Gazz. Chim. Ital.*, 1988, **118**, 507 (*Oricine, Zanthobungeanine, 8-Methoxyflindersine, N-Methylflindersine, 7-Methoxy-N-methylflindersine, synth*)
Ngadjui, B.T. et al., *Phytochemistry*, 1988, **27**, 2979 (*Araliopsinine*)
Grundon, M.F. et al., *Nat. Prod. Rep.*, 1990, **7**, 131 (*Araliopsinine*)
Campbell, W.E. et al., *Phytochemistry*, 1990, **29**, 1303 (*pmr, N-Methylhaplamine*)
Jackson, G.E. et al., *Spectrosc. Lett.*, 1990, **23**, 971 (*pmr, cmr, 8-Methoxyflindersine*)
Anand, R.C. et al., *Indian J. Chem., Sect. B*, 1991, **30**, 560; 604 (*synth, Flindersine, N-methylflindersine*)
Subramanian, M. et al., *Z. Naturforsch., B*, 1992, **47**, 1016 (*synth*)
Brader, G. et al., *Annalen*, 1993, 355 (*N-Acetoxyethylflindersine*)
Brader, G. et al., *Phytochemistry*, 1996, **42**, 881 (*7-Methoxyflindersine, 7-Prenyloxyflindersine, N-Methyl-7-prenyloxyflindersine*)
Watters, W.H. et al., *J. Chem. Res., Synop.*, 1997, 184-185 (*Zanthodioline, Veprisine, synth*)
Chen, I.S. et al., *Phytochemistry*, 1997, **46**, 525-529 (*Zanthodioline*)
Akhmedzhanova, V.I. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1999, **35**, 552-553 (*Dihydrohaplamine*)
Ali, M.S. et al., *Phytochemistry*, 2001, **57**, 1277-1280 (*Haplophylline A*)
Lee, Y.R. et al., *Synthesis*, 2001, 1851-1855 (*synth, ir, pmr, N-Methylflindersine*)
Ito, C. et al., *J. Nat. Prod.*, 2004, **67**, 1800-1803 (*Orixalone D*)
Atta-ur-Rahman, et al., *J. Enzyme Inhib. Med. Chem.*, 2006, **21**, 703-710 (*Skimmia laureola constii*)
Thangavel, D. et al., *J. Chem. Res.*, 2007, 124-126 (*synth*)
Wang, X. et al., *Synthesis*, 2007, 3044-3050 (*synth, ir, pmr, N-Methylflindersine, Orixalone D*)
Wang, X. et al., *Tet. Lett.*, 2007, **48**, 6275-6280 (*Orixalone D, synth*)

Floramultinine

F-91

C₂₁H₂₇NO₆ 389.447


Struct. unknown. Compd. possesses 1 NMe and 2 methoxy groups. Alkaloid from the whole plant of *Kreysigia multiflora* (Liliaceae). Sol. H₂O. Mp 165°. $[\alpha]_D^{20}$ +118 (c. 0.17 in EtOH).

Badger, G.M. et al., *J.C.S.*, 1960, 445-447 (*isol, uv, ir*)

Floridimine

F-92

[200067-96-5]

C₁₅H₂₇NO₅ 301.382

Ester of Trachelanthamidine in H-629 with Trachelanthic acid. Alkaloid from *Heliotropium floridum* (Boraginaceae). Oil.

3'-Ac: Floridine

[200067-92-1]

C₁₇H₂₉NO₆ 343.419

Alkaloid from *Heliotropium floridum* (Boraginaceae). Oil. [α]_D -14.2 (c, 0.04 in EtOH).

2'-Epimer: Floridinine

[200067-94-3]

Alkaloid from *Heliotropium floridum* (Boraginaceae). Oil. [α]_D -8.3 (c, 0.1 in EtOH). Ester of Trachelanthamidine in H-629 with Viridifloric acid.

Stereoisomer: Macrotomine. Makrotomine

C₁₅H₂₇NO₅ 301.382

Alkaloid from *Macrotomia echioidea* (Boraginaceae). Large prisms (Me₂CO). Mp 95-97°. [α]_D -6.9 (c, 7.58 in EtOH).

Stereoisomer, picrate:

Fine yellow needles (EtOH). Mp 130-132°.


Men'shikov, G.P. *et al.*, *Zh. Obshch. Khim.*, 1952, **22**, 1457-1461; *J. Gen. Chem. USSR (Engl. Transl.)*, 1952, **22**, 1499-1502 (Macrotomine)

Denisova, S.I. *et al.*, *Zh. Obshch. Khim.*, 1958, **28**, 1882-1885; *J. Gen. Chem. USSR (Engl. Transl.)*, 1958, **28**, 1925-1927 (Macrotomine)

Reina, M. *et al.*, *Phytochemistry*, 1997, **46**, 845-853 (Floridine, Floridinine, Floridine)

Flueggeine A

F-93



Absolute Configuration


C₂₄H₂₆N₂O₄ 406.48

Dimer of Norsescurinine, N-307. Alkaloid from the roots of *Flueggea virosa*. Amorph. powder. [α]_D²⁰ -24 (c, 0.5 in MeOH). λ_{max} 212 (log ε 4.34); 260 (log ε 4.1) (MeOH).

Gan, L.-S. *et al.*, *Org. Lett.*, 2006, **8**, 2285-2288 (isol, pmr, cmr, ms)

Flueggeine B

F-94



Absolute Configuration

C₂₄H₂₈N₂O₅ 424.496

Alkaloid from the roots of *Flueggea virosa*. Amorph. powder. [α]_D²⁰ -70 (c, 0.22 in MeOH). λ_{max} 213 (log ε 4.22) (MeOH).

Gan, L.-S. *et al.*, *Org. Lett.*, 2006, **8**, 2285-2288 (isol, pmr, cmr, ms)

Fluggeine

F-95

Fluggeine

C₁₀H₁₅NO 165.235


Struct. unknown. Alkaloid from *Flueggea virosa*. Needles. Mp 228°.

Paris, R.A. *et al.*, *Ann. Pharm. Fr.*, 1955, **13**, 245-249; *CA*, **49**, 16345f

Fluopsin B

F-96

Tris(N-hydroxy-N-methylmethanethioamidato-O,S)iron, 9CI. Fluopsin F [31323-26-9]



C₆H₁₂FeN₃O₃S₃ 326.224

Fe complex of Thioformin. Isol. from *Pseudomonas fluorescens*. Broad spectrum antibiotic with antifungal and antineoplastic props. Black prisms (MeOH/CHCl₃). Sol. EtOH, CHCl₃; fairly sol. EtOAc, C₆H₆, Et₂O, H₂O. Dec. on heating.

▶ LD₅₀ (mus, ivn) 3 - 9 mg/kg, LD₅₀ (mus, ipr) 30 - 50 mg/kg. NO8780000

Shirahata, K. *et al.*, *J. Antibiot.*, 1970, **23**, 546 (isol, struct)

Mitchell, A.J. *et al.*, *Aust. J. Chem.*, 1977, **30**, 2439 (synth)

Murray, K.S. *et al.*, *Aust. J. Chem.*, 1978, **31**, 983 (cryst struct)


Leong, J. *et al.*, *Inorg. Chem.*, 1978, **17**, 1886 (ir, nmr, synth)

Bell, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1979, **15**, 384 (props)

Fluopsin C

F-97

Bis(N-hydroxy-N-methylmethanethioamidato-O,S)copper, 9CI. Bis(N-methylthioformohydroxamato)copper, 8CI. YC 73. Antibiotic YC 73. MRL 3120. Antibiotic MRL 3120 [31323-25-8]



C₄H₈CuN₂O₂S₂ 243.797

Cu complex of Thioformin. Isol. from *Pseudomonas fluorescens*. Also prod. by *Pseudomonas aeruginosa*. Possesses antibacterial props. Dark-green cryst. (EtOH). Mp 199° dec. λ_{max} 231, 253, 267, 320 and 365 nm (MeOH). λ_{max} 230 (ε 13000); 252 (ε 12500); 266 (ε 14400); 319

(ε 7230); 364 (sh) (ε 3140) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 4.5 mg/kg. GL6490000 [67069-47-0]

Shirahata, K. *et al.*, *J. Antibiot.*, 1970, **23**, 546;

1971, **24**, 140 (isol, struct, synth, ir, nmr)

Egawa, Y. *et al.*, *J. Antibiot.*, 1971, **24**, 124 (struct, synth)

Taylor, D. *et al.*, *Cryst. Struct. Commun.*, 1978, **7**, 237 (cryst struct)


Bell, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1979, **15**, 384 (props)

Ko, H.-R. *et al.*, *CA*, 1994, **121**, 152876 (MRL 3120)

Fluopsin N

F-98

Bis(N-hydroxy-N-methylmethanethioamidato-O,S)nickel, 9CI [31541-90-9]



C₄H₈N₂NiO₂S₂ 238.941

Isol. from *Pseudomonas fluorescens*. Antibiotic props. resemble those of Fluopsin B, F-96. Purple cryst.

Japan. Pat., 1973, 73 10 554; *CA*, **80**, 35824w (synth)


Leong, J. *et al.*, *Inorg. Chem.*, 1978, **17**, 1886 (synth, nmr)

Bell, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1979, **15**, 384 (props)

Fluorocarpamine

F-99

16-De(hydroxymethyl)-4-demethyl-16-(methoxycarbonyl)fluorocurine, 9CI [2254-31-1]



Absolute Configuration

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia plumosa*, *Alstonia angustifolia*, *Alstonia undulata*, *Tabernaemontana citrifolia*, *Hunteria zeylanica*, *Gonioma kamassi* and *Petchia ceylanica*. Amorph. λ_{max} 235 (log ε 4.05); 257 (log ε 3.64); 398 (log ε 3.13) (EtOH).

N^b-Oxide: **Fluorocarpamine N-oxide** [88746-90-1]

C₂₀H₂₂N₂O₄ 354.405

Alkaloid from *Catharanthus roseus* (Apocynaceae).


Kaschnitz, R. *et al.*, *Monatsh. Chem.*, 1965, **96**, 909-921 (isol, ms, uv, struct)

Jacquier, M.J. *et al.*, *Phytochemistry*, 1982, **21**, 2973-2977 (isol)

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1983, **49**, 124-125 (oxide)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1989, **28**, 3221-3225 (isol, ir, uv, pmr, ms)

Takayama, H. *et al.*, *Heterocycles*, 1996, **42**, 87-92 (abs config, cd)

C-FluorocurineFluorocurine
[6879-92-1]Absolute
configuration $C_{20}H_{25}N_2O_2^{\oplus}$ 325.43Alkaloid from *Strychnos panamensis*, *Strychnos variabilis*, *Strychnos scheffleri* and *Strychnos melinoniana* (Loganiaceae).

▶ LN2800000



Iodide:

 $C_{20}H_{25}IN_2O_2$ 452.334Yellow cryst. or powder. Mp 138-140°. $[\alpha]_D^{25} +326$ (c, 0.831 in MeOH). Intense carmine col. with Ce(IV).

Picrate: Mp 179°.

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1947, **30**, 2081; 1952, **35**, 1846; 1964, **47**, 878 (*isol, uv, pmr, struct*)Wieland, T. *et al.*, *Chem. Ber.*, 1952, **85**, 731 (*isol*)Lee, F.G.H. *et al.*, *Diss. Abstr. B*, 1967, **27**, 2297 (*synth*)Angenot, L. *et al.*, *Planta Med.*, 1981, **42**, 364-370; 371-374 (*isol, ir, pmr*)Coune, C. *et al.*, *J. Pharm. Belg.*, 1982, **37**, 189 (*cmr*)**ψ-Fluorocurine**


Pseudofluorocurine

 $C_{20}H_{25}N_2O_2$ 325.43Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 179° (as picrate). Closely similar to C-Fluorocurine, F-100; shows identical colour reacs. and UV spectrum but does not form an insol. iodide or *p*-nitrophenylhydrazone.Meyer, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1208-1213 (*isol, uv*)**C-Fluorocurinine** $C_{21}H_{29}N_2O_2^{\oplus}$ 341.472Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Yellow needles (2-pentanone)(as picrate). Mp 213° (picrate). Carmine col. with Ce(SO₄)₂, becoming wine-red on standing. Schmid, H. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 1864-1879; 1953, **36**, 102-121**5-Fluoro-3,4-dihydro-2,4-dioxo-1(2H)-pyrimidineacetic acid****F-100** $C_6H_5FN_2O_4$ 188.115*Me ester*: 5-Fluoro-1-(methoxycarbonylmethyl)uracil
[56059-28-0] $C_7H_7FN_2O_4$ 202.142Isol. from the sponge *Phakellia fusca*. Needles (MeOH). Mp 178-179° (natural) Mp 185-186° (synthetic). λ_{max} 219 (ε 5725); 268 (ε 12813) (MeOH).*Hydrazide*: [488713-02-6] $C_6H_7FN_4O_3$ 202.145Isol. from the sponge *Phakellia fusca*. Brown solid. Mp > 300°. λ_{max} 219 (ε 5763); 274 (ε 10909) (MeOH).Tada, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 3427-3428 (*synth*)Xu, X.-H. *et al.*, *J. Nat. Prod.*, 2003, **66**, 285-288 (*isol*)**5-Fluoro-2,4-dihydroxypyrimidine**5-Fluoro-2,4(1H,3H)-pyrimidinedione, 9CI. 5-Fluoro-2,4-pyrimidinediol. 5-Fluorouracil. **Fluorouracil**, **BAN**, **INN**, **JAN**, **USAN**. *Adrucil*. *Efudex*. *Fluoroplex*. *Fluracil*. *NSC 19893*. *Ro 2-9757*. 5-FU [51-21-8] $C_4H_3FN_2O_2$ 130.078Dione form predominates. Two intermediate tautomers also possible. Isol. from the sponge *Phakellia fusca*. Reported from Chinese drug Mao Guo Yan Ming (roots of *Rabdosia lasiocarpus*). Antineoplastic agent. Used in combination with Adrenaline, A-152 in the treatment of genital warts (AccuSite). Cryst. (H₂O). Mp 282-283° dec. Log P - 0.97 (calc).▶ Skin and respiratory tract irritant. Gastrointestinal, bone marrow depression and other adverse systemic effects when used therapeutically. Exp. reprod. and teratogenic effects. LD₅₀ (rat, orl) 230 mg/kg. YR0350000Rudy, B.C. *et al.*, *Anal. Profiles Drug Subst.*, 1973, **2**, 221 (*rev*)Wasternack, C. *et al.*, *Pharmazie*, 1987, **42**, 73 (*rev*)Li, T.S.C. *et al.*, *Chinese and Related North American Herbs*, CRC Press, 2002, 124 (*occur*)Xu, X.-H. *et al.*, *J. Nat. Prod.*, 2003, **66**, 285-288 (*Phakellia fusca uracils*)**Fluorosolimoosines**

[1357-19-3 (IV), 1357-17-1 (II), 1357-16-0 (I), 1357-18-2 (III)]

Struct. unknown. Alkaloids from *Strychnos solimoosana* (Loganiaceae). Fluorosolimoosines I, II, III and IV identified, having different colour reacs. Give no colour with Ce(IV).Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148**Flustramine C**

[78127-86-3]

 $C_{16}H_{19}BrN_2$ 319.243Alkaloid from the marine bryozoan *Flustra foliacea*. Oil. λ_{max} 210 (sh) (ε 13000); 230 (ε 23000); 280 (ε 4600) (EtOH/HCl) (Derep). λ_{max} 210 (sh) (ε 10000); 232 (ε 27000); 290 (ε 9400); 307 (ε 4300) (EtOH) (Derep).8,8a-Dihydro: **Dihydroflustramine C**

[93930-02-0]

 $C_{16}H_{21}BrN_2$ 321.259Alkaloid from *Flustra foliacea*. Cryst. (Et₂O/petrol). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 82-84°. $[\alpha]_D^{25} -110$ (c, 1.5 in CH₂Cl₂). λ_{max} 213 (ε 20000); 230 (ε 5600); 309 (ε 3400) (MeOH) (Berdy).8,8a-Dihydro, N¹-Oxide: **Dihydroflustramine C N-oxide**

[104387-11-3]

 $C_{16}H_{21}BrN_2O$ 337.259Minor metab. from *Flustra foliacea*. Oil. $[\alpha]_D^{25} -67.1$ (c, 0.38 in CH₂Cl₂). Somewhat unstable. λ_{max} 226 (ε 2800); 283 (ε 2000) (MeOH) (Berdy).

8,8a-Dihydro, 8-(3-methyl-2-butenyl):

Flustramine A

[71239-64-0]

 $C_{21}H_{29}BrN_2$ 389.377Alkaloid from the marine bryozoan *Flustra foliacea*. Muscle relaxant. Liq. Sol. MeOH, C₆H₆, CHCl₃; fairly sol. hexane. $[\alpha]_D^{20} -76.9$ (c, 6.6 in EtOH). *cis*-3a,8a-Ring junction. λ_{max} 219 (ε 21000); 253 (ε 7700); 305 (ε 3500) (EtOH/HCl) (Derep). λ_{max} 218 (ε 20000); 262 (ε 9200); 317 (ε 3500) (EtOH) (Derep).

8,8a-Dihydro, 2-oxo, 8-(3-methyl-2-butenyl):

Flustramide A

[82534-44-9]


 $C_{21}H_{27}BrN_2O$ 403.361Alkaloid from the marine bryozoan *Flustra foliacea*. Oil. λ_{max} 216 (ε 18200); 260 (ε 7760); 310 (ε 3550) (EtOH) (Derep).Carlé, J.S. *et al.*, *J.A.C.S.*, 1979, **101**, 4012-4013 (*isol*)Carlé, J.S. *et al.*, *J.O.C.*, 1980, **45**, 1586-1589; 1981, **46**, 3440-3443 (*Flustramine A*,*Flustramine C, isol, uv, ir, pmr, ms, struct*)Wulff, P. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1982, **71**, 523-524 (*Flustramide A*)Wright, J.L.C. *et al.*, *J. Nat. Prod.*, 1984, **47**, 893-895 (*Dihydroflustramine C*)Laycock, M.V. *et al.*, *Can. J. Chem.*, 1986, **64**, 1312-1316 (*Dihydroflustramine N¹-oxide*)Kawasaki, T. *et al.*, *Tet. Lett.*, 1996, **37**, 7525-7528 (*synth, cmr*)Morales-Rios, M.S. *et al.*, *J.O.C.*, 1999, **64**, 1086-1087; 2001, **66**, 1186-1192 (*Flustramine***F-106**

- A. Debromoflustramine A, Flustramide A, synth*
 Morales-Rios, M.S. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 677-682 (*Flustramine A, pmr, cmr, conformn*)
 Morales-Rios, M.S. *et al.*, *Tetrahedron*, 2002, **58**, 1479-1484 (*Dihydroflustramine C, synth*)
 Lysek, N. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 1056-1061 (*Flustramine A, isol, pmr, cmr*)
 Fuchs, J.R. *et al.*, *Org. Lett.*, 2005, **7**, 677-680 (*synth*)
 Lindel, T. *et al.*, *Org. Lett.*, 2007, **9**, 283-286 (*synth*)
 Kawasaki, T. *et al.*, *J.O.C.*, 2008, **73**, 5959-5964 (*Flustramine A, Flustramide A, synth*)

Flustramine D

F-107

[104387-12-4]



Relative Configuration

$C_{21}H_{29}BrN_2$ 389.377
 Metab. from the marine bryozoan *Flustra foliacea*. Oil. $[\alpha]_D^{25}$ -86.5 (c, 1.03 in CH_2Cl_2). λ_{max} 231 (ε 7100); 292 (ε 3100) (MeOH) (Berdy).

*N*¹-Oxide: **Flustramine D N-oxide**

[104387-13-5]

 $C_{21}H_{29}BrN_2O$ 405.377


Trace metab. from *Flustra foliacea*. Oil. Unstable. λ_{max} 224 (ε 7100); 285 (ε 3100) (MeOH) (Berdy).

Laycock, M.V. *et al.*, *Can. J. Chem.*, 1986, **64**, 1312-1316 (*isol, uv, ir, pmr, cmr, ms, struct, oxide*)

Flustramine E

F-108

[158642-05-8]



Absolute Configuration

 $C_{16}H_{21}BrN_2$ 321.259

Rel. stereochem. only detd. Alkaloid from the marine bryozoan *Flustra foliacea*. $[\alpha]_D^{20}$ -1136 (c, 0.0088 in EtOH). λ_{max} 210 ; 250 ; 308 (MeOH) (Berdy).

*N*⁸-(3-Methyl-2-butenyl): **Flustramine B**

[71239-65-1]

 $C_{21}H_{29}BrN_2$ 389.377

Alkaloid from the marine bryozoan *Flustra foliacea*. Muscle relaxant. Sol. MeOH, C_6H_6 , $CHCl_3$; fairly sol. hexane. λ_{max} 219 (ε 21000); 253 (ε 7700); 305 (ε 3500) (EtOH/HCl) (Derep). λ_{max} 218 (ε 20000); 262 (ε 9200); 317 (ε 3500) (EtOH) (Derep).

*2-Oxo, N*⁸-(3-methyl-2-butenyl): **Flustramide B**

[105708-75-6]

 $C_{21}H_{27}BrN_2O$ 403.361Minor alkaloid from *Flustra foliacea*.

Oil. $[\alpha]_D^{20}$ -53.79 (EtOH). No stereochem. given. λ_{max} 216 (ε 18200); 260 (ε 7760); 310 (ε 3550) (EtOH) (Derep).

*Debromo, N*⁸-(3-methyl-2-butenyl): **Debromoflustramine B**

[158060-73-2]

 $C_{21}H_{30}N_2$ 310.481

Trace alkaloid from *Flustra foliacea*. $[\alpha]_D^{20}$ -98.2 (c, 0.0153 in $CHCl_3$). (+)-form has been synthesised. λ_{max} 257 ; 308 (MeOH) (Berdy).


4'-Hydroxy, debromo: see Pseudophrynamine 258A, P-720

- Carlé, J.S. *et al.*, *J.A.C.S.*, 1979, **101**, 4012-4013 (*isol, Flustramine B*)
 Carlé, J.S. *et al.*, *J.O.C.*, 1980, **45**, 1586-1589 (*isol, uv, ir, pmr, ms, struct, Flustramine B*)
 Hino, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 1806-1808 (*synth*)
 Keil, P. *et al.*, *Acta Chem. Scand., Ser. B*, 1986, **40**, 555-558 (*Flustramide B*)
 Holst, P.B. *et al.*, *J. Nat. Prod.*, 1994, **57**, 997-1000 (*Flustramine E, Debromoflustramine B*)
 Bruncko, M. *et al.*, *J.O.C.*, 1994, **59**, 5543-5549 (*Debromoflustramine B, synth*)
 Somei, M. *et al.*, *Heterocycles*, 1997, **45**, 2327-2330 (*synth, Debromoflustramine B*)
 Morales-Rios, M.S. *et al.*, *J.O.C.*, 1999, **64**, 1086-1087; 2001, **66**, 1186-1192 (*Flustramide B, Flustramine B, Debromoflustramine B, synth*)
 Morales-Rios, M.S. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 677-682 (*pmr, cmr, conformn*)
 Morales-Rios, M.S. *et al.*, *Tetrahedron*, 2002, **58**, 1479-1484 (*synth*)
 Tan, G.H. *et al.*, *Org. Lett.*, 2003, **5**, 1801-1803 (*synth*)
 Kawasaki, T. *et al.*, *Chem. Comm.*, 2006, 420-422 (*Flustramine B, Flustramide B, synth*)
 Cardoso, A.S.P. *et al.*, *Tetrahedron*, 2007, **63**, 10211-10225 (*Debromoflustramine B, synth*)

Flustraminol A

F-109

[78127-88-5]



Relative Configuration

 $C_{16}H_{21}BrN_2O$ 337.259


Alkaloid from the marine bryozoan *Flustra foliacea*. Brown oil. λ_{max} 215 (ε 18000); 234 (sh) (ε 8600); 246 (ε 5400); 306 (ε 2300) (EtOH/HCl) (Derep). λ_{max} 215 (ε 17000); 256 (ε 6500); 318 (ε 2300) (EtOH) (Derep).

Carlé, J.S. *et al.*, *J.O.C.*, 1981, **46**, 3440-3443 (*isol, uv, ir, pmr, ms, struct*)

Flustraminol B

F-110

[78127-89-6]



Relative Configuration

 $C_{16}H_{21}BrN_2O$ 337.259

Alkaloid from the marine bryozoan *Flustra foliacea*. Brown oil. λ_{max} 220 (ε 17000); 240 (sh) (ε 6300); 298 (ε 2300) (EtOH/HCl) (Derep). λ_{max} 220 (ε 17000); 256 (ε 6300); 310 (ε 2300) (EtOH) (Derep).


Carlé, J.S. *et al.*, *J.O.C.*, 1981, **46**, 3440-3443 (*isol, uv, ir, pmr, ms*)

Suárez-Castillo, O.R. *et al.*, *Heterocycles*, 2007, **71**, 1539-1551 (*synth*)

Flustramine B

F-111

[105708-76-7]



Relative configuration

 $C_{21}H_{29}BrN_2O$ 405.377


Alkaloid from the marine bryozoan *Flustra foliacea*. Also obt. by H_2O_2 oxidn. of Flustramine E, F-108. $[\alpha]_D^{20}$ -180 (EtOH). Prob. struct. Nat. prod may be Flustramine B N-oxide. λ_{max} 216 (ε 18200); 260 (ε 7760); 310 (ε 3550) (EtOH) (Derep).

Keil, P. *et al.*, *Acta Chem. Scand., Ser. B*, 1986, **40**, 555-558 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Flutimide

F-112

1-Hydroxy-5-(2-methylpropyl)-3-(2-methylpropylidene)-2,6(1H,3H)-pyrazin-2(1H)-one
 [162666-34-4]

 $C_{12}H_{18}N_2O_3$ 238.286

Isol. from the fungus *Delitschia confer-taspora*. Endonuclease inhibitor of influenza virus. Antiviral agent. Gum. λ_{max} 270; 360 (sh) (MeOH) (Derep).

Hensens, O.D. *et al.*, *Tet. Lett.*, 1995, **36**, 2005-2008 (*isol, uv, ir, struct*)


Tomassini, J.E. *et al.*, *Antimicrob. Agents Chemother.*, 1996, **40**, 1189-1193 (*activity*)

Singh, S.B. *et al.*, *J.O.C.*, 2001, **64**, 5504-5516 (*synth*)

Fluvibactin

F-113

[103185-30-4]




C₃₁H₃₄N₄O₁₀ 622.63
Isol. from *Vibrio fluvialis*. Siderophore.
 λ_{\max} 248 (E1%/1cm 10.1); 314 (E1%/1cm
5.4) (MeOH) (Berdy).

Yamamoto, S. *et al.*, *J. Biochem. (Tokyo)*,
1993, **113**, 538 (*isol, pmr, ms, struct*)

Fluviol D F-114


[195154-70-2]



C₁₁H₁₀N₁₀O₂ 314.266
Prod. by *Pseudomonas fluorescens*. Un-
stable. λ_{\max} 237 ; 278 ; 374 ; 555 (EtOH).

Smirnov, V.V. *et al.*, *FEMS Microbiol. Lett.*,
1997, **153**, 357-361

Fluivuricin B F-115



Fluivuricin B₁ R₁ = R₂ = H
Fluivuricin B₃ R₁ = CH₃, R₂ = H
Fluivuricin B₅ R₁ = CH₃, R₂ = CONH CH₂CH₂PH

Macrolactam antibiotic. Identity of the
Fluivuricins with the Sch antibiotics does
not seem fully establ. There are some
discrepancies in Mp's and they could be
stereoisomeric. Prod. by an unidentified
actinomycete. Antiviral agent.

Fluivuricin B₁

Sch 38516. *Antibiotic Sch 38516*
[137428-64-9]

C₂₄H₄₆N₂O₅ 442.638
Isol. from *Actinomadura vulgaris* ssp.
lanata. Possesses antifungal props. Need-
les (MeOH). Mp 156-160° dec Mp 262-
263°. $[\alpha]_D^{26}$ -6.7 (c, 0.5 in DMSO). No
stereochem. assigned to Fluivuricin B₁.
Sch 38516 possesses 3R,7S,11S-config.

4'-Epimer: *Antibiotic Sch 38513. Sch*
38513. Fluivuricin B₂

C₂₄H₄₆N₂O₅ 442.638

Prod. by *Actinomadura* spp. Sol.
MeOH, DMSO, butanol; fairly sol.
EtOAc; poorly sol. H₂O.

Fluivuricin B₃

Sch 39185. *Antibiotic Sch 39185*
[137120-29-7]

C₂₅H₄₈N₂O₅ 456.665
Prod. by *Actinomadura vulgaris* ssp.
vulgaris. Antifungal agent. Needles
(MeOH). Mp 263-266° (216-220° dec.).
 $[\alpha]_D^{26}$ -5.8 (c, 0.5 in MeOH).

4'-Epimer: *Antibiotic Sch 38518. Sch*
38518. Fluivuricin B₂. AB 85. Antibiotic
AB 85

[137120-28-6]
C₂₅H₄₈N₂O₅ 456.665

Prod. by *Actinomadura vulgaris* spp.
vulgaris. Antifungal agent. Needles
(MeOH). Mp 261-263° (220° dec.).
 $[\alpha]_D^{26}$ +9.7 (c, 0.5 in MeOH).

N^{3'}-[[2-(2-Phenylethyl)amino]carbo-
nyl]: *Fluivuricin B₅*

[137120-14-0]

C₃₄H₅₇N₃O₆ 603.841

Prod. by *Microtetraspora* sp. Rods.
Sol. MeOH, Py, EtOAc; poorly sol.
H₂O, hexane. Mp 210° dec. λ_{\max} 246 (ε
810); 252 (ε 810); 258 (ε 800); 264 (sh)
(ε 680); 268 (sh) (ε 610) (MeOH)
(Derep).

4'-Epimer, 2'-O-α-D-glucopyranosyl:

Antibiotic Sch 42729. Sch 42729

[151284-51-4]

C₃₁H₅₈N₂O₁₀ 618.807

Prod. by a *Actinomadura* sp. Antifun-
gal agent. Sol. MeOH, butanol,
DMSO; fairly sol. EtOAc; poorly sol.
H₂O.

4'-Epimer, N^{3'}-[[2-(2-phenylethyl)ami-
no]carbonyl]: *Fluivuricin B₄*

[137019-45-5]

C₃₄H₅₇N₃O₆ 603.841

Prod. by *Microtetraspora* sp. Needles.
Sol. MeOH, Py, EtOAc; poorly sol.
H₂O, hexane. Mp 245° dec. λ_{\max} 246 (ε
810); 252 (ε 810); 258 (ε 800); 264 (sh)
(ε 680); 268 (sh) (ε 610) (MeOH)
(Derep).

4'-Epimer, 2'-O-[β-D-glucopyranosyl-
(1→4)-α-D-glucopyranosyl]: *Antibio-
tic Sch 42282. Sch 42282*

C₃₇H₆₈N₂O₁₅ 780.949

Prod. by *Microtetraspora* sp. Antifun-
gal agent. Sol. MeOH, butanol.

[135559-94-3 , 128563-23-5 , 135559-95-4]

Hegde, V.R. *et al.*, *J.A.C.S.*, 1990, **112**, 6403-
6405 (*Sch 38516, cryst struct*)

Naruse, N. *et al.*, *J. Antibiot.*, 1991, **44**, 733-
740; 741-755; 756-761; 940-948 (*Fluivuricins*)

Puar, M.S. *et al.*, *Bioorg. Med. Chem. Lett.*,
1992, **2**, 575 (*biosynth*)

Hegde, V. *et al.*, *J. Antibiot.*, 1992, **45**, 624-632;
1993, **46**, 1109-1115; 1998, **51**, 464-470 (*Sch*
antibiotics)


Cooper, R. *et al.*, *J. Antibiot.*, 1992, **45**, 633-
638 (*isol, cmr, struct*)

Houri, A.F. *et al.*, *J.A.C.S.*, 1995, **117**, 2943-
2944 (*synth*)

Xu, Z. *et al.*, *J.A.C.S.*, 1997, **119**, 10302-10316
(*synth*)

Foetidine I

F-116



C₄₇H₄₉N₅O₉ 827.932

Erroneous mol. formula in paper.

(S)-form [163167-81-5]

Alkaloid from trunk bark of *Mappia*
foetida (Icacinaeae). Shows potent anti-
viral activity. Cryst. (Me₂CO/hexane).
Mp 157-158° dec. $[\alpha]_D^{25}$ -37.9 (c, 0.31 in
MeOH).

9-Methoxy: *Foetidine II*

[173220-98-9]

C₄₈H₅₁N₅O₁₀ 857.958

Alkaloid from *Mappia foetida*. Anti-
tumour and antiviral agent. Cryst. Mp
172-174°. $[\alpha]_D$ -44.6 (c, 0.35 in
MeOH).


Eur. Pat., 1995, 685 481; *CA*, **124**, 141377u
(*isol*)

Pirillo, A. *et al.*, *J.C.S. Perkin 1*, 1995, 583-587
(*isol, pmr, cmr, struct*)

Folicangine

F-117

[32340-00-4]



C₄₂H₄₆N₄O₅ 686.849

Alkaloid from leaves of *Voacanga afri-
cana* (Apocynaceae). Cryst. (MeOH).
Mp 200° dec. $[\alpha]_D$ -271 (CHCl₃).

Kunesch, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970,
4370 (*isol, uv, ir, pmr, ms*)


Rolland, Y. *et al.*, *J.O.C.*, 1976, **41**, 3270
(*cmr*)

Kunesch, N. *et al.*, *Helv. Chim. Acta*, 1977, **60**,
2854 (*struct*)

Folifinine

F-118

8-Hydroxy-4-methoxy-α,α-dimethyl-
furo[2,3-b]quinoline-7-propanol, 9CI. 8-
Hydroxy-7-(3-hydroxy-3-methylbutyl)-4-
methoxyfuro[2,3-b]quinoline
[22329-40-4]



C₁₇H₁₉NO₄ 301.341


Alkaloid from aerial parts of *Haplophyl-
lum foliosum* (Rutaceae). Cryst.
(Me₂CO). Mp 181-182°.

Kurbanov, D. *et al.*, *Khim. Prir. Soedin.*, 1968,
4, 373; *Chem. Nat. Compd. (Engl. Transl.)*,
1968, **4**, 315

Folimidine

F-119

2-(4-Hydroxy-3-methoxyphenyl)-1-methyl-4(1H)-quinolinone, 9CI
[40444-99-3]



C₁₇H₁₅NO₃ 281.31
Alkaloid from *Haplophyllum foliosum*
(Rutaceae). Cryst. (EtOH). Mp 247°.

Me ether:


Cryst. (Et₂O). Mp 186-187°.

Razzakova, D.M. *et al.*, *Khim. Prir. Soedin.*,
1972, **8**, 755; *Chem. Nat. Compd. (Engl.
Transl.)*, 1972, **8**, 737 (*ir, uv, pmr, ms*)

Foliminine

F-120

[52617-28-4]



C₁₇H₁₇NO₃ 283.326
Probable struct. Alkaloid from the
above-ground parts of *Haplophyllum
foliosum* (Rutaceae). Cryst. (MeOH aq.).
Mp 107-108°.

Hydrochloride:


Cryst. (EtOH). Mp 188°.

Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*,
1974, **10**, 52; *Chem. Nat. Compd. (Engl.
Transl.)*, 1974, **10**, 46 (*uv, ir, pmr, ms, isol
struct*)

Folinic acid

F-121

N-[4-(2-Amino-5-formyl-1,4,5,6,7,8-
hexahydro-4-oxo-6-pteridiny)methyl]a-
mino]benzoylglutamic acid, 9CI. 5-Formyl-
5,6,7,8-tetrahydrofolic acid.
Citrovorum factor. Leucovorin. Formylp-
teroylglutamic acid
[58-05-9]



C₂₀H₂₃N₇O₇ 473.444

►WB2650000

(6S,2'S)-form

(-)-L-form
[68538-85-2]

Occurs in yeast, insects etc. Active form
of folic acid in the body. Light yellow
cryst. + 3H₂O. Mp 248-250° dec. [α]_D²⁵
+16.8 (aq. bicarbonate). This is the
biologically active natural stereoisomer.
The “(-)-L” description used in the

biochemical literature is rather mislead-
ing.

Ca salt: **Calcium levofolinat**, INN. *Le-
voleucovorin calcium, USAN. Isovorin.*
CL 307782
[80433-71-2]
[α]_D²⁰ -15.1 (c, 1.82 in H₂O).

(6R,2'S)-form

(+)-L-form
[α]_D²⁰ +28.3 (c, 3.53 in H₂O) (as Ca salt).

(6RS,2'S)-form

dl-L-form
Antidote to folic acid antagonists such as
methotrexate e.g. in cancer chemother-
apy. Cryst. Spar. sol. H₂O to give acid
soln. which slowly decs. Mp 240-250°
dec. pK_{a1} 3.2; pK_{a2} 4.8; pK_{a3} 10.4. This is
the usual coml. form, consisting of a
mixt. of diastereoisomers. Used as a Ca
salt.

Ca salt: **Leucovorin calcium, USAN.**
*Calcium folinat, BAN, INN. Lederfol-
lin. Leucosar. Refolinon. Rescufolin.*
NSC 3590
[41927-89-3]
[1492-18-8]
Yellowish powder + 5H₂O. Sol. H₂O;
insol. EtOH. [α]_D²⁰ +14.26 (c, 3.42 in
H₂O).

[6035-45-6]

Brockman, J.A. *et al.*, *J.A.C.S.*, 1950, **72**, 4325
(*synth*)

Cosulich, D.B. *et al.*, *J.A.C.S.*, 1952, **74**, 3252
Kas, J. *et al.*, *J. Chromatogr.*, 1976, **124**, 53
(*chromatog*)

Pont, O. *et al.*, *Anal. Profiles Drug Subst.*,
1979, **8**, 315 (*bibl*)

Khalifa, E. *et al.*, *Helv. Chim. Acta*, 1980, **63**,
2554 (*synth*)

Feeney, J. *et al.*, *J.C.S. Perkin 2*, 1980, 176
(*pmr, cmr*)

Kerr, D.J. *et al.*, *Br. J. Cancer*, 1989, **60**, 807
(*pharmacol*)


Owens, J. *et al.*, *J.C.S. Perkin 1*, 1993, 871
(*synth*)

Martindale, *The Extra Pharmacopoeia*, 30th
edn., Pharmaceutical Press, 1993, 1040

Foliosidine

F-122

8-(2,3-Dihydroxy-3-methylbutoxy)-4-
methoxy-1-methyl-2(1H)-quinolinone,
9CI. Pholiosidine
[2520-38-9]



C₁₆H₂₁NO₅ 307.346

Alkaloid from *Haplophyllum foliosum*
(Rutaceae). Said to be pharmacologi-
cally active and to be introduced into
USSR medical practice, but no details
given. Mp 140-141°. [α]_D +41.6
(EtOH).

2'-Ac: Foliphorine

[42907-17-5]

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Haplophyllum foliosum*.
Cryst. (hexane/petrol). Mp 83-84°.

2',3'-Isopropylidene: *Foliosidine acetonide*
C₁₉H₂₅NO₅ 347.41

Artifact isol. from *Haplophyllum fo-
liosum*. Cryst. (petrol). Mp 119-120°.

O-De-Me: Mp 235-236°. [α]_D +50.4 (Py).

3'-Deoxy, 2'-ketone: 4-Methoxy-1-
methyl-8-(3-methyl-2-oxobutoxy)-
2(1H)-quinolinone, 9CI. **Folidine**
[102719-91-5]

C₁₆H₁₉NO₄ 289.33

Alkaloid from aerial parts of *Haplo-
phyllum foliosum* (Rutaceae). Cryst.
(Me₂CO/petrol). Mp 148-149°.

Pastukhova, V.I. *et al.*, *Dokl. Akad. Nauk
SSSR*, 1964, **21**, 31; *CA*, **62**, 11864d (*ir*)
Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*,
1968, **4**, 201; *Chem. Nat. Compd. (Engl.
Transl.)*, 1968, **4**, 174 (*pmr*)

Shakirov, T.T. *et al.*, *Khim. Prir. Soedin.*, 1969,
5, 385; *Chem. Nat. Compd. (Engl. Transl.)*,
1969, **5**, 385 (*isol, use*)

Tel'nov, V.A. *et al.*, *Khim. Prir. Soedin.*, 1970,
6, 724; *Chem. Nat. Compd. (Engl. Transl.)*,
1970, **6**, 735 (*acetonide*)


Akhmedzhanova, V.I. *et al.*, *Khim. Prir.
Soedin.*, 1985, **21**, 823; *Chem. Nat. Compd.
(Engl. Transl.)*, 1985, **21**, 782 (*Folidine*)

Reisch, J. *et al.*, *Monatsh. Chem.*, 1988, **119**,
1169 (*synth, pmr, ms, Folidine*)

Akhmedzhanova, V.I. *et al.*, *Chem. Nat.
Compd. (Engl. Transl.)*, 1999, **35**, 552-553
(*Foliphorine*)

Folipidine

F-123



C₂₂H₂₂N₂O₄ 378.427


Alkaloid from the aerial parts of *Haplo-
phyllum pedicellatum*. Cryst. (EtOH). Mp
242-243°. [α]_D -46 (c, 0.67 in Py). λ_{max}
230 ; 248 ; 278 ; 292 (sh) ; 305 (sh) ; 356 ;
370 (EtOH).

Akhmedzhanova, V.I. *et al.*, *Chem. Nat.
Compd. (Engl. Transl.)*, 2005, **41**, 60-64
(*isol, pmr, cmr, ms*)

Fontaineine

F-124

[211690-64-1]




C₂₅H₃₈N₄O₄ 458.6

Alkaloid from *Fontainea pacheri* (Euphorbiaceae). Gum.

Lontsi, D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 953-954 (*isol, ir, pmr, cmr*)

Fontaphilline F-125

Methyl 5-ethenyl-4-[2-[(4-hydroxybenzoyl)oxy]ethyl]-3-pyridinecarboxylate, 9CI
[22667-62-5]




C₁₈H₁₇NO₅ 327.336

Alkaloid from leaves of *Fontanesia phillyroides* (Oleaceae). Needles (MeOH aq.). Mp 80-81° Mp 121-122° (double Mp).

Budzikiewicz, H. *et al.*, *Chem. Ber.*, 1967, **100**, 2798 (*ir, uv, pmr, ms, isol, struct*)

Fontonamide F-126

[109217-15-4]



C₂₀H₂₂ClNO₂ 343.852

Apparently a seco-indole alkaloid which is a singlet oxygen oxidn. prod. of Hapalindole A, H-48. See also Hapalnamide G, H-52. *Isol.* from blue-green alga *Hapalosiphon fontinalis*. Mp 156-157°. [α]_D -141 (c, 0.21 in CHCl₃). λ_{max} 240 (ε 5680); 289 (ε 4770); 345 (ε 1630) (MeOH) (Derep).

Dechloro: Dechlorofontonamide

[123498-02-2]

C₂₀H₂₃NO₂ 309.407


Minor alkaloid from *Hapalosiphon fontinalis*. [α]_D -100 (c, 0.2 in CHCl₃).

Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 3773 (*isol, struct*)

Moore, R.E. *et al.*, *Phytochemistry*, 1989, **28**, 1565 (*Dechlorofontonamide*)

Forbagline A F-127

[177262-35-0]



C₃₇H₄₂N₂O₉ 658.747

Alkaloid from *Aglaia forbesii*. Cryst. (MeOH). Mp 246°. [α]_D²⁰ +58 (c, 1.2 in MeOH). λ_{max} 281 (log ε 3.94) (MeOH).

13-Epimer: Forbagline B


[177468-87-0]

C₃₇H₄₂N₂O₉ 658.747

Alkaloid from *Aglaia forbesii*. Amorph. powder. [α]_D²⁰ +76 (c, 0.2 in MeOH). λ_{max} 277 (log ε 4.26) (MeOH).

Dumontet, V. *et al.*, *Tetrahedron*, 1996, **52**, 6931-6942 (*isol, uv, ir, pmr, cmr*)

Aglaia dasyclada Forbagline-like compound F-128



C₃₅H₃₈N₂O₁₀ 646.693

Related to Forbagline A, F-127. Alkaloid from *Aglaia dasyclada*. Amorph. [α]_D²⁰ -11 (c, 0.1 in MeOH).


4'-Me ether, Me ester, 21-O-β-D-glucopyranoside:

C₄₃H₅₂N₂O₁₅ 836.888

Alkaloid from *Aglaia dasyclada*. Amorph. [α]_D²⁰ -38.5 (c, 0.2 in MeOH).

Chaidir, *et al.*, *J. Nat. Prod.*, 2001, **64**, 1216-1220 (*isol, pmr, cmr*)

11-Formamido-5-eudesmene F-129



C₁₆H₂₇NO 249.395

(4β,7α,10β)-form

Constit. of *Axinella cannabina*. Oil.

Isocyanide: 11-Isocyanato-5-eudesmene

C₁₆H₂₅N 231.38

Constit. of *Axinella cannabina* and *Phakellia* sp. Oil. [α]_D -89.7 (c, 0.8 in CHCl₃). Has -NC replacing -NHCHO.

Isothiocyanate: 11-Isothiocyanato-5-eudesmene

[108639-31-2]

C₁₆H₂₅NS 263.446

Constit. of *Axinella cannabina*, *Acanthella klethra* and *Cadlina luteo-marginata*. Oil. [α]_D -89.7 (c, 0.8 in CHCl₃). Has -NCS replacing -NHCHO.


Ciminiello, P. *et al.*, *Can. J. Chem.*, 1987, **65**, 518

Angerhofer, C.K. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1787 (*isothiocyanate, isol*)

Burgoyne, D.L. *et al.*, *Tetrahedron*, 1993, **49**, 4503

Formobactin F-130

[181417-64-1]




C₃₈H₅₇N₅O₁₀ 743.896

Prod. by a *Nocardia* sp. Siderophore. Free radical scavenger. Powder. Related to Nocobactin NA, N-267.

Murakami, Y. *et al.*, *J. Antibiot.*, 1996, **49**, 839-845 (*isol, uv, ir, pmr, cmr, props*)

Formycin A F-131

1-C-(7-Amino-1H-pyrazolo[4,3-d]pyrimidin-3-yl)-1,4-anhydro-D-ribose, 9CI. Formycin. NSC 102811. Antibiotic 4-215 [6742-12-7]



C₁₀H₁₃N₅O₄ 267.244

Nucleoside-type antibiotic. Prod. by *Nocardia interforma*, *Streptomyces lavendulae* and *Streptomyces gunnaensis*. Shows limited antibacterial activity but possesses antitumour and antiviral props. Insulinotropic agent. Adenosine transport inhibitor. Cryst. Sol. MeOH, H₂O, acids, bases; fairly sol. EtOH; poorly sol. Me₂CO, hexane. Mp 153-155°. [α]_D²⁵ -39.3 (c, 0.5 in 0.1M HCl). pK_{a1} 4.4; pK_{a2} 9.7 (H₂O). λ_{max} 295 (ε 10700) (H₂O) (Derep). λ_{max} 294 (E1%/1cm 390) (H₂O) (Berdy). λ_{max} 234 (E1%/1cm 280); 295 (E1%/1cm 340) (HCl) (Berdy). λ_{max} 235 (E1%/1cm 500); 305 (E1%/1cm 260) (NaOH) (Berdy).

▶ LD₅₀ (mus, orl) 1000 mg/kg; LD₅₀ (mus, ivn) 250-350 mg/kg. UR0719000

4-Me: Mp 268-269°.

6-Me:

Dihydrate. Mp 231-232°.

7-N-Me: Mp 145° dec.

3'-Epimer: Xyloformycin

Synthetic. Shows no antiviral props.

Cryst. Mp 138-140°. [α]_D -97.8 (c, 0.23 in H₂O).

Koyama, G. *et al.*, *J. Antibiot.*, Ser. A, 1965, **18**, 175 (*isol, ir, uv*)

Koyama, G. *et al.*, *Tet. Lett.*, 1966, 597 (*ir, uv, pmr, cryst struct*)

Japan. Pat., 1967, 67 10 928; CA, **68**, 24552 (*isol*)


Japan. Pat., 1968, 68 6 996; CA, **69**, 34691 (*isol*) Prusiner, P. *et al.*, *Biochemistry*, 1973, **12**, 1196 (*cryst struct*)

Ochi, K. *et al.*, *J. Antibiot.*, 1976, **29**, 638 (*biosynth*)

- Chenon, M.-T. *et al.*, *J.A.C.S.*, 1976, **98**, 4736 (*pmr, cmr*)
 Kalvoda, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 1431 (*synth*)
 Lewis, A.F. *et al.*, *J.A.C.S.*, 1980, **102**, 2817
 Spiemulli, E.N. *et al.*, *Cancer Treat. Rep.*, 1983, **67**, 267 (*pharmacol, tox*)
 Buchanan, J.G. *et al.*, *J.C.S. Perkin 1*, 1986, 1267 (*analogs*)
 Cho, B.P. *et al.*, *Nucleosides Nucleotides*, 1994, **13**, 481 (*pmr, cmr, tautom*)

Formycin B F-132

1,4-Dihydro-3-β-D-ribofuranosyl-7H-pyrazolo[4,3-d]pyrimidin-7-one, 9CI. *Laurusin*. *Ohyamycin*. NSC 106486 [13877-76-4]



$C_{10}H_{12}N_4O_5$ 268.229

Nucleoside-type antibiotic. Prod. by *No-cardia interforma*, *Streptomyces lavendulae* and *Streptomyces roseochromogenes-oyaensis*. Shows limited antibacterial activity; possesses antitumour and antiviral props. Nucleoside transporter substrate. Cryst. (H_2O). Mp 254-255°. $[\alpha]_D^{25}$ -52 (c, 0.5 in H_2O). pK_{a1} 8.8; pK_{a2} 10.4. λ_{max} 219 (ϵ 9330); 280 (ϵ 7880) (H_2O) (Derep). \blacktriangleright LD₅₀ (mus, ivn) 1000 mg/kg. UR0800000


Hydrobromide:

Prisms + H_2O (H_2O). Mp 180-180.5°.

- Aizawa, S. *et al.*, *Agric. Biol. Chem.*, 1965, **29**, 375; 377 (*isol*)
 Koyama, G. *et al.*, *J. Antibiot., Ser. A*, 1965, **18**, 175 (*isol, ir, uv*)
Japan. Pat., 1966, 66 13 792; *CA*, **65**, 19272 (*isol*)
 Koyama, G. *et al.*, *Tet. Lett.*, 1966, 597 (*uv, ir, nmr, cryst struct*)
 Acton, E.M. *et al.*, *Chem. Comm.*, 1971, 986 (*synth, nmr*)
 Koyama, G. *et al.*, *Acta Cryst. B*, 1976, **32**, 813 (*cryst struct*)
 Chenon, M.-T. *et al.*, *J.A.C.S.*, 1976, **98**, 4736 (*pmr, cmr*)
 Kalvoda, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 1431 (*synth*)
 Buchanan, J.G. *et al.*, *J.C.S. Perkin 1*, 1984, 2367 (*synth*)
 Plagemann, P.G. *et al.*, *Biochim. Biophys. Acta*, 1990, **1022**, 93-102; 103-109; **1028**, 289-298 (*pharmacol*)

7-Formyldehydrohernangerine F-133

[178765-68-9]




$C_{19}H_{15}NO_5$ 337.331

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Yellowish prisms ($CHCl_3$ /MeOH). Mp 211-213°.

Chen, J.-J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (*isol, uv, ir, pmr, ms, struct*)

7-Formyldehydronornantentine F-134

[169624-28-6]




$C_{20}H_{17}NO_5$ 351.358

Alkaloid from trunk bark of *Hernandia sonora*. Yellowish prisms ($CHCl_3$ /MeOH). Mp 213-214°.

Chen, I.-S. *et al.*, *Phytochemistry*, 1995, **40**, 983 (*isol, uv, ir, pmr, ms, struct*)

7-Formyldehydrovigerine F-135


[169624-27-5]



$C_{19}H_{13}NO_5$ 335.315

Alkaloid from trunk bark of *Hernandia sonora*. Yellowish prisms ($CHCl_3$ /MeOH). Mp 247-249°.

Chen, I.-S. *et al.*, *Phytochemistry*, 1995, **40**, 983 (*isol, uv, ir, pmr, ms, struct*)

7-Formyldehydrothalicsimidine F-136


$C_{23}H_{25}NO_6$ 411.454

Alkaloid from *Annona purpurea* (soncoya). Yellow needles ($CHCl_3$). Mp 160-162°. λ_{max} 217 ; 268 ; 335 (sh) ; 432 (EtOH).

Chang, F.-R. *et al.*, *Phytochemistry*, 1998, **49**, 2015-2018 (*isol, uv, ir, pmr*)

4-(2-Formyl-5-hydroxymethyl-1H-pyrrol-1-yl)butanoic acid F-137

[142629-55-8]



$C_{10}H_{13}NO_4$ 211.217

Alkaloid from the processed roots of *Aconitum* sp. (Kako-bushi-matsu), the seeds of *Allium fistulosum* and the fruit of *Lycium chinense* (Chinese boxthorn). Hepatoprotectant. Amorph. solid. λ_{max} 292 (log ϵ 3.55) (EtOH).

Me ether: 2-Formyl-5-methoxymethyl-1H-pyrrole-1-butanoic acid

$C_{11}H_{15}NO_4$ 225.244

Alkaloid from the fruit of *Lycium chinense* (Chinese boxthorn). Hepatoprotectant. λ_{max} 291 (log ϵ 3.76) (EtOH).

Me ether, Me ester: Methyl 2-formyl-5-methoxymethyl-1H-pyrrole-1-butanoate

$C_{12}H_{17}NO_4$ 239.271

Alkaloid from the fruit of *Lycium chinense* (Chinese boxthorn) and from the bulbs of *Bolbostemma paniculatum*. Yellow oil. λ_{max} 286 (log ϵ 3.39) (EtOH).

Matsui, M. *et al.*, *Nat. Med. (Tokyo)*, 1998, **52**, 232-235 (*isol*)


Sang, S. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 6318-6321 (*isol, pmr, cmr*)

Chin, Y.-W. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 79-81 (*isol, synth, pmr, cmr*)

Liu, W.-Y. *et al.*, *J. Asian Nat. Prod. Res.*, 2003, **5**, 159-163 (*Me ether Me ester*)

2-(2-Formyl-5-hydroxymethyl-1H-pyrrol-1-yl)-3-phenylpropanoic acid F-138

α -Benzyl-2-formyl-5-hydroxymethyl-1H-pyrrole-1-acetic acid



$C_{15}H_{15}NO_4$ 273.288

Me ether, Me ester: Methyl 2-(2-formyl-5-methoxymethyl-1H-pyrrol-1-yl)-3-phenylpropanoate

$C_{17}H_{19}NO_4$ 301.341


Constit. of the bulbs of *Bolbostemma paniculatum*. Yellow oil.

Liu, W.-Y. *et al.*, *J. Asian Nat. Prod. Res.*, 2003, **5**, 159-163 (*isol, pmr, cmr, ms*)

3-(2-Formyl-5-hydroxymethyl-1H-pyrrol-1-yl)-2,5-pyrrolidinedione

F-139

2',3',4',5'-Tetrahydro-5-hydroxymethyl-2',5'-dioxo-1,3'-bi-1H-pyrrole-2-carboxaldehyde

C₁₀H₁₀N₂O₄ 222.2**(-)-form**


Constit. of the dried fruit of *Prunus domestica* (prunes). Powder. $[\alpha]_D^{25}$ -89.2 (c, 0.12 in MeOH). λ_{\max} 259 (log ϵ 3.76); 294 (log ϵ 4.06) (MeOH).

Kayano, S. et al., *Biosci., Biotechnol., Biochem.*, 2004, **68**, 942-944 (isol, pmr, cmr)

3-Formyl-1H-indole-2-sulfonic acid

F-140

[857351-59-8]

C₉H₇NO₄S 225.225


Isol. from *Leptosphaeria maculans*. Inter-termed. in the detoxification pathways of 8H-Isothiazolo[5,4-b]indole, I-336 and its N-methoxy deriv. Sinalexin in I-336. Mp 300-305° dec. λ_{\max} 212 (ε 17400); 244 (ε 9800); 304 (ε 7100) (MeOH).

Pedras, M.S.C. et al., *Org. Biomol. Chem.*, 2005, **3**, 2002-2007 (isol, synth, pmr, cmr)

5-Formyl-1H-pyrrole-2-alkanenitriles

F-141

5-(ω-Cyanoalkyl)-2-pyrrolicarboxaldehydes

**5-Formyl-1H-pyrrole-2-octadecanenitrile**

F-142

5-(17-Cyanoheptadecyl)-1H-pyrrole-2-carboxaldehyde. **Mycalenitrile 1**

[705973-45-1]

C₂₃H₃₈N₂O 358.566

Alkaloid from the sponge *Mycale cecilia*. Oil. n = 17. λ_{\max} 204 (ε 6025); 248 (ε 2540); 300 (ε 10125) (MeOH).

5-Formyl-1H-pyrrole-2-icosanenitrile, 9CI

5-(19-Cyanononadecyl)-1H-pyrrole-2-carboxaldehyde

[233744-62-2]

C₂₅H₄₂N₂O 386.62

Alkaloid from the sponges *Desmampsama anchorata* and *Mycale microsigmata*. Antileishmanial agent. n = 19.

5-Formyl-1H-pyrrole-2-heneicosanenitrile5-(20-Cyanoicosyl)-1H-pyrrole-2-carboxaldehyde. **Mycalenitrile 2**

[705973-46-2]

C₂₆H₄₄N₂O 400.646

Alkaloid from the sponge *Mycale cecilia*. Oil. n = 20. λ_{\max} 204 (ε 6030); 248 (ε 2710); 300 (ε 11790) (MeOH).

5-Formyl-1H-pyrrole-2-docosanenitrile5-(21-Cyanoheicosyl)-1H-pyrrole-2-carboxaldehyde. **Mycalenitrile 3**

[705973-47-3]

C₂₇H₄₆N₂O 414.673

Alkaloid from the sponge *Mycale cecilia*. Oil. n = 21. λ_{\max} 203 (ε 5950); 248 (ε 2510); 300 (ε 10000) (MeOH).

Compagnone, R.S. et al., *Nat. Prod. Lett.*, 1999, **13**, 203-211 (5-Formyl-1H-pyrrole-2-icosanenitrile)

Ortega, M.J. et al., *Tetrahedron*, 2004, **60**, 2517-2524 (Mycalenitriles)

5-Formyl-1H-pyrrole-2-carboxylic acid

F-142

[7126-51-4]

C₆H₅NO₃ 139.11

Prod. in cultures of *Erwinia oxoidiae*. Cryst. (H₂O). Mp 202-203° (212-215° dec.).

Me ester: [1197-13-3]C₇H₇NO₃ 153.137

Cryst. (petrol). Mp 92-93°.

Et ester: [7126-50-3]C₈H₉NO₃ 167.164Cryst. (petrol). Mp 75°. Bp_{0.1} 96-100°.*1-Me, Me ester*:C₈H₉NO₃ 167.164

Cryst. (petrol). Mp 100-102°.

Reichstein, T. et al., *Helv. Chim. Acta*, 1930, **13**, 349 (synth)

Khan, M.K.A. et al., *Tetrahedron*, 1966, **22**, 2095 (synth, ir, pmr)

Muchowski, J.M. et al., *Tet. Lett.*, 1988, **29**, 777 (synth)

Wallace, D.M. et al., *J.O.C.*, 1993, **58**, 7245 (*Et ester*)

Campbell, S.E. et al., *J.C.S. Perkin 1*, 1997, 2195-2202 (*Et ester*, synth, pmr, cmr)


Hong, F. et al., *J.C.S. Perkin 1*, 1997, 2997-3003 (*Me ester*, synth, pmr, cmr)

Fortuneine

F-143

1,2,6,7-Tetrahydro-3,15,16-trimethoxy-C-homoerythrinan, 9CI

[87340-25-8]

C₂₀H₂₅NO₃ 327.422

Alkaloid from the twigs and leaves of *Cephalotaxus fortunei*. Needles (Me₂CO). Mp 110°. $[\alpha]_D^{26}$ -121 (c, 0.2 in MeOH). C-6/C-7 configs. not known for Comosiline. λ_{\max} 233 (log ϵ 4.05); 282 (sh) (log ϵ 3.37) (MeOH).

6β,7β-Epoxyde: Wilsonine

[39024-12-9]

C₂₀H₂₅NO₄ 343.422

Alkaloid from the twigs and leaves of *Cephalotaxus wilsoniana* and *Cephalotaxus fortunei*, the seeds of *Cephalotaxus sinensis* (Cephalotaxaceae). Cryst. (Et₂O). Poorly sol. hexane. Mp 150-151°. $[\alpha]_D$ -51.4 (c, 0.55 in CHCl₃). $[\alpha]_D$ -36 (c, 0.55 in EtOH). λ_{\max} 233 (log ϵ 3.9); 281 (log ϵ 3.41) (EtOH).

3-Epimer: Epifortuneine

[87340-26-9]

Alkaloid from *Cephalotaxus fortunei*. Pale yellow amorph. solid. $[\alpha]_D^{26}$ +134.4 (c, 0.06 in MeOH).

3-Epimer, 6β,7β-epoxyde: 3-Epiwilsonine.

Alkaloid 7. Comosiline

[39024-15-2]

[123163-73-5 (Comosiline)]

C₂₀H₂₅NO₄ 343.422

Alkaloid from the leaves of *Phelline comosa*, twigs and leaves of *Cephalotaxus wilsoniana* and *Cephalotaxus fortunei* and from *Cephalotaxus sinensis*. Also a minor constit. of the seed of *Cephalotaxus harringtonia* var. *drupacea* (Phellinaceae, Cephalotaxaceae). Cryst. (Et₂O). Mp 103-104°. $[\alpha]_D$ +60.7 (c, 0.55 in CHCl₃). $[\alpha]_D$ +75.8 (c, 0.52 in EtOH). λ_{\max} 233 (log ϵ 3.92); 281 (log ϵ 3.44) (EtOH).

3-Epimer, 6β,7β-epoxyde, O¹⁶-de-Me:**Robustiline**

[123050-84-0]

C₁₉H₂₃NO₄ 329.395

Alkaloid from leaves of *Phelline comosa* var. *robusta* (Phellinaceae). Mp 228°. $[\alpha]_D$ +71.2 (c, 0.48 in CHCl₃). λ_{\max} 208 (ε 17900); 282 (ε 2000) (EtOH).

3-Epimer, 6β,7β-epoxyde, O¹⁷-de-Me:**Isorobustiline**

[123011-99-4]

C₁₉H₂₃NO₄ 329.395

From leaves of *Phelline comosa* var. *robusta* (Phellinaceae). Mp 205°. $[\alpha]_D$ +69.7 (c, 0.46 in CHCl₃). λ_{\max} 211 (ε 14250); 282 (ε 3350) (EtOH).

3-Epimer, 8-oxo, 6α,7α-epoxyde: 3-EpiwilsononeC₂₀H₂₃NO₅ 357.405

Alkaloid from *Cephalotaxus wilsoniana*. Cytotoxic. Powder. $[\alpha]_D^{28}$ +11 (c, 0.1 in CHCl₃). λ_{\max} 206 (log ϵ 4.37); 225 (sh) (log ϵ 4.02); 280 (log ϵ 3.56) (MeOH).

Langlois, N. et al., *Bull. Soc. Chim. Fr.*, 1970, 3535-3543 (*Wilsonine*, 3-*Epiwilsonine*)

Powell, R.G. et al., *Phytochemistry*, 1972, **11**, 3317-3320 (*Wilsonine*, 3-*Epiwilsonine*)


Ma, G. et al., *Phytochemistry*, 1983, **22**, 251-253 (*Fortuneine*, *Wilsonine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

- Pusset, J. *et al.*, *Phytochemistry*, 1989, **28**, 1298-1300 (*Robustiline*, *Isorobustiline*)
 Ma, G. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 137-140 (*Epifortuneine*)
 Langlois, N. *et al.*, *Heterocycles*, 1990, **30**, 659-664 (*Robustiline*, *struct*)
 Du, J. *et al.*, *CA*, 1999, **130**, 63637m (*Wilsonine*, *cryst struct*, *abs config*)
 Wang, L.-W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1182-1185 (*3-Epivilsonone*)

Foveoglin A

F-144

[948312-48-9]

C₃₈H₄₀N₂O₈ 652.743

Alkaloid from the leaves of *Aglaia foveolata*. Cytotoxic. Amorph. powder. [α]_D²⁰ -30 (c, 0.16 in CHCl₃). λ_{max} 203 (log ε 4.64); 213 (sh) (log ε 4.57) (MeOH).

10-Epimer: Foveoglin B

[948312-49-0]

C₃₈H₄₀N₂O₈ 652.743


Alkaloid from the leaves of *Aglaia foveolata*. Amorph. powder. [α]_D²⁰ +170 (c, 0.2 in CHCl₃). λ_{max} 203 (log ε 4.76); 214 (sh) (log ε 4.71) (MeOH).

Salim, A.A. *et al.*, *Tetrahedron*, 2007, **63**, 7926-7934 (*isol*)

Fragilamide

F-145

2-[[[4-(4-Hydroxyphenyl)methyl]methylamino]-N-[2-(1H-indol-3-yl)ethenyl]-4-methylhexanamide, 9CI
 [87168-33-0]

C₂₅H₃₁N₃O₂ 405.539


Alkaloid from the marine red alga *Martensia fragilis*. [α]_D -32 (c, 0.62 in MeOH). λ_{max} 229 (ε 15500); 280 (ε 10200) (EtOH) (*Derep*).

Kirkup, M.P. *et al.*, *Tet. Lett.*, 1983, **24**, 2087 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Fragin, 8CI

F-146

N-[2-(Hydroxynitrosamino)-3-methylbutyl]octanamide, 8CI
 [17073-33-5]

C₁₃H₂₇N₃O₃ 273.375

λ_{max} 248 (ε 8000) (EtOH/NaOH) (*Derep*). λ_{max} 230 (ε 7000) (EtOH) (*Derep*).

▶RG7957000

(-)-form

Prod. by *Pseudomonas fragi*. Phytotoxin. Possesses antifungal, antitumour and antiviral props. Plates (hexane). Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O, acids. Mp 80°. [α]_D²⁵ -122 (c, 1.9 in EtOH). pK_a 6.4. λ_{max} 230 (ε 7000) (EtOH) (*Berdy*). λ_{max} 230 (ε 7000) (EtOH-HCl) (*Berdy*). λ_{max} 248 (ε 8000) (EtOH-NaOH) (*Berdy*).
 ▶LD₅₀ (mus, ivn) 80 mg/kg.

(±)-form

Cryst. (hexane). Mp 97°.


Tamura, S. *et al.*, *Agric. Biol. Chem.*, 1967, **31**, 758; 996 (*synth*)

Murayama, A. *et al.*, *Agric. Biol. Chem.*, 1969, **33**, 1599; 1970, **34**, 122; 130 (*isol*, *struct*, *synth*)

Francheline

F-147

[799782-00-6]



Absolute Configuration

C₂₄H₃₇NO₆ 435.559

Unusual 6,12-epoxyaconitane structs. Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrum*. Amorph. powder. Mp 86-88°. [α]_D²⁰ -146.8 (c, 0.5 in CHCl₃).

14-Benzoyl: 13-Hydroxyfranchetine

[874986-10-4]

C₃₁H₄₁NO₇ 539.667

Alkaloid from the roots of *Aconitum nagarum*. Amorph. powder. Mp 97-98°. [α]_D²⁵ -120 (c, 0.1 in CHCl₃).

13-Deoxy, 14-Ac: Vilmorisine

[143109-03-9]

C₂₆H₃₉NO₆ 461.597

Alkaloid from roots of *Aconitum vilmorianum* (Ranunculaceae). Struct. revised in 1997.

13-Deoxy, 14-benzoyl: Franchetine†

[88661-42-1]

C₃₁H₄₁NO₆ 523.668

Alkaloid from roots of *Aconitum franchetii* (Ranunculaceae) and *Aconitum hemsleyanum* var. *pengshiesse*. Struct. revised in 1997.

13-Deoxy, 14-cinnamoyl: Leueandine

[664304-10-3]

C₃₃H₄₃NO₆ 549.706

Alkaloid from the roots of *Aconitum hemsleyanum* var. *leueanthus*. Amorph. powder. Mp 138-140°. [α]_D²⁵ -71.4 (c, 0.5 in CHCl₃). Config. of cinnamoyl group not determined.

13-Deoxy, 3α-hydroxy, 14-benzoyl: 3-Hydroxyfranchetine

[1007889-95-3]

C₃₁H₄₁NO₇ 539.667

Alkaloid from the roots of *Aconitum hemsleyanum* var. *atropurpureum*. Amorph. powder. Mp 116-118°. [α]_D²⁰ -76.7 (c, 0.3 in Me₂CO).

15α-Hydroxy, 14-benzoyl: Beiwudine

[219314-07-5]

C₃₁H₄₁NO₈ 555.667

Alkaloid from the roots of *Aconitum kusnezoffii*. Amorph. powder.

Ding, L.S. *et al.*, *Huaxue Xuebao*, 1992, **50**, 405; *CA*, **117**, 108100f (*Vilmorisine*)

Wang, F.-P. *et al.*, *Phytochemistry*, 1997, **45**, 1539-1542 (*Franchetine*)

Wang, F.-P. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1555-1556 (*Beiwudine*)

Chen, D.-L. *et al.*, *J. Asian Nat. Prod. Res.*, 2003, **5**, 209-213 (*Leueandine*)

Ji, H. *et al.*, *Heterocycles*, 2004, **63**, 2363-2370 (*Francheline*)

Zhang, F. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 1043-1046 (*13-Hydroxyfranchetine*)

Tang, P. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 704-707 (*3-Hydroxyfranchetine*)

Franchetine†

F-148

Pyrrrolizidine alkaloid of unknown struct. Alkaloid from *Senecio franchetii* (Asteraceae). Cryst. (Et₂O). Mp 124-125°. Cooccurs with Sarracine N-Oxide in S-90.


Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 351; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 296-297 (*isol*)

Franganine

F-149

Daechuine S4

[19526-08-0]



Absolute Configuration

C₂₈H₄₄N₄O₄ 500.68

Alkaloid from *Rhamnus frangula*, *Discaria americana*, *Discaria febrifuga*, *Euonymus europaeus*, *Melochia corchorifolia*, the Daechu tree (*Zizyphus jujuba* var. *inermis*) and *Zizyphus spinachristi*. Shows antibacterial, antifungal and sedative props. Needles (petrol). Mp 248°. [α]_D²² -302 (c, 0.1 in CHCl₃).

10',11'-Dihydro, 11'-hydroxy: Discarine H

[95689-32-0]

C₂₈H₄₆N₄O₅ 518.695

Alkaloid from the root bark of *Discaria febrifuga* (Rhamnaceae). Mp 232°. [α]_D²⁰ -266 (MeOH).

Tschesche, R. *et al.*, *Tet. Lett.*, 1968, 2993; 3817 (*isol*, *pmr*, *ms*, *struct*)

Bishay, D.W. *et al.*, *Phytochemistry*, 1973, **12**, 693 (*uv*, *ir*, *ms*)

Herzog, R. *et al.*, *Chem.-Ztg.*, 1984, **108**, 406 (*Discarine H*)


Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443

Da Silva, U.F. *et al.*, *Phytochem. Anal.*, 1996, **7**, 20-23 (*config*)

Tan, N.H. *et al.*, *Chem. Rev.*, 2006, **106**, 840-895 (bibl)

Franguloline**F-150**

Sanjoinine A. Daechuine S1
[19526-09-1]



R = PhCH₂—

C₃₁H₄₂N₄O₄ 534.697

Alkaloid from the leaves of *Rhamnus frangula*, *Euonymus europaeus*, *Ceanothus sanguineus*, *Melochia pyramidata*, *Melochia corchorifolia*, *Discaria febrifuga*, *Discaria longispina*, *Zizyphus mauritiana*, *Zizyphus lotus* and *Zizyphus nummularia*. Also isol. from Sanjoin (seeds of *Zizyphus vulgaris* var. *spinosus*) and from the stem bark of the Daechu tree (*Zizyphus jujuba* var. *inermis*) (Rhamnaceae, Celastraceae, Sterculiaceae). Shows strong sedative activity. Needles (EtOH aq. or CH₂Cl₂/MeOH/Et₂O). Mp 244° (234-236°). [α]_D²² -299 (c, 0.1 in CHCl₃). The *Z. lotus* isolate was stated to be racemic.

N-De-Me: Sanjoinine B. N-Demethylsanjoinine A. N-Demethylfranguloline
[107462-33-9]

C₃₀H₄₀N₄O₄ 520.67

Isol. from the seeds of *Zizyphus vulgaris* var. *spinosus*. Needles (MeOH). Mp 212-214°.

15ξ-Hydroxy: Sanjoinine F

[107462-35-1]

C₃₁H₄₂N₄O₅ 550.697

Isol. from the seeds of *Zizyphus vulgaris* var. *spinosus* and from *Zizyphus lotus*. Needles (EtOAc). Mp 228-229°. [α]_D²⁶ -215 (c, 0.28 in CHCl₃). The *Z. lotus* isolate stated to be racemic.

3'-Epimer: Sanjoinine Ah₁

[107494-19-9]

C₃₁H₄₂N₄O₄ 534.697

Alkaloid from the seeds of *Zizyphus vulgaris spinosus*. Possible artifact.

Tschesche, R. *et al.*, *Tet. Lett.*, 1968, 2993-2998; 3817-3818; 1972, 2609-2612 (isol, pmr, ms, occur, struct)

Mascaretti, O.A. *et al.*, *Phytochemistry*, 1972, **11**, 1133-1137 (isol)

Bishay, D.W. *et al.*, *Phytochemistry*, 1973, **12**, 693-698 (uv, ir, pmr, ms)

Merkuza, V.M. *et al.*, *Phytochemistry*, 1974, **13**, 1279-1282 (ir, ms)

Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (isol)

Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 663-668 (isol, ms)

Medina, E. *et al.*, *Annalen*, 1981, 538-545 (isol, ir, pmr, ms)

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443-448 (*Sanjoinine Ah₁*)


Han, B.H. *et al.*, *Phytochemistry*, 1990, **29**, 3315-3319 (*Sanjoinines B,F*)

Abu-Zarga, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 504-511 ((±)-*Franguloline*, (±)-*Sanjoinine F*)

Xiao, D. *et al.*, *Tet. Lett.*, 1998, **39**, 9631-9632 (synth)

Frangulanine**F-151**

2-(Dimethylamino)-3-methyl-N-[3-(1-methylethyl)-7-(2-methylpropyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]pentanamide, 9CI. Ceanothamine A. Daechuine S2
[25350-22-5]



Absolute Configuration

C₂₈H₄₄N₄O₄ 500.68

Alkaloid from *Rhamnus frangula*, *Zizyphus sativa*, *Ceanothus americanus*, *Hovenia dulcis*, *Hovenia tomentella*, *Zizyphus jujuba* var. *inermis*, *Euonymus europaeus* and *Discaria longispina*. Needles (CHCl₃/petrol, MeOH or CH₂Cl₂/EtOAc). Mp 276-279° (275-276°). [α]_D²⁰ -288 (c, 0.1 in CHCl₃).

N-De-Me: Hovenine A. N-Demethylfrangulanine

[52309-78-1]

C₂₇H₄₂N₄O₄ 486.653

Alkaloid from the root bark of *Hovenia dulcis* (raisin tree) and *Hovenia tomentella* (Rhamnaceae). Mp 215°. Opt. rotn. not reported.

10',11'-Dihydro, 11'-hydroxy: Discarine L
[165561-02-4]

C₂₈H₄₆N₄O₅ 518.695

Alkaloid from root bark of *Discaria febrifuga* (Rhamnaceae). Amorph. powder. [α]_D -30 (c, 0.5 in MeOH). Not interrelated with Frangulanine. Stereochem. may not correspond.

Warnhoff, E.W. *et al.*, *Can. J. Chem.*, 1965, **43**, 2594 (isol, uv, pmr, ms)

Tschesche, R. *et al.*, *Chem. Ber.*, 1967, **100**, 3937 (isol, uv, cd, ir, pmr, ms, struct)

González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915 (pmr, stereochem)

Mascaretti, O.A. *et al.*, *Phytochemistry*, 1972, **11**, 1133 (isol, ir)

Bishay, D.W. *et al.*, *Phytochemistry*, 1973, **12**, 693 (uv, ir, pmr, ms)

Takai, M. *et al.*, *Phytochemistry*, 1973, **12**, 2985 (isol, ir, pmr, ms, deriv)

Chang, C.-J. *et al.*, *Phytochemistry*, 1974, **13**, 1273 (pmr)

Otsuka, H. *et al.*, *Phytochemistry*, 1974, **13**, 2016 (occur)

Takai, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2556; 1976, **24**, 2181 (pmr, conformn, cryst struct)

Haslinger, E. *et al.*, *Tetrahedron*, 1978, **34**, 685 (pmr, cmr, conformn)


Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443

Schmidt, U. *et al.*, *Chem. Comm.*, 1991, 1002 (synth)

Morel, A.F. *et al.*, *Phytochemistry*, 1995, **39**, 431-434 (*Discarine L*)

Fredericamycin A**F-152**

FCRC A48. NSC 305263. Antibiotic FCRC A48
[80455-68-1]



Absolute Configuration

C₃₀H₂₁NO₉ 539.497

Prod. by *Streptomyces griseus*
ATCC49344. Antitumour agent.

Thought to be a generator of free radicals causing DNA strand cleavage. Red powder. Sol. Py, AcOH, DMSO, DMF; fairly sol. MeOH; poorly sol. Me₂CO, H₂O, hexane. Mp 350° dec. Red in acid, blue-green in base. Non-mutagenic in Ames test. λ_{max} 253 (ε 50800); 304 (ε 23500); 317 (ε 20600); 358 (sh) (ε 22000); 392 (ε 20600); 505 (ε 10400) (MeOH/HCOOH) (Derep). λ_{max} 260 (ε 43459); 302 (ε 22991); 316 (ε 25152); 332 (ε 23323); 357 (ε 24044); 373 (ε 19971); 392 (ε 23711); 635 (ε 8033) (MeOH).

Pandey, R.C. *et al.*, *J. Antibiot.*, 1981, **34**, 1389-1401 (isol)

Warnick-Pickle, D.J. *et al.*, *J. Antibiot.*, 1981, **34**, 1402-1407 (props)

Misra, R. *et al.*, *J.A.C.S.*, 1982, **104**, 4478-4479 (cryst struct, pmr)

Parker, K.A. *et al.*, *Tet. Lett.*, 1986, **27**, 3835-3838 (synth, bibl)

Misra, R. *et al.*, *J. Antibiot.*, 1987, **40**, 786-802; 1988, **41**, 976-981 (pmr, cmr, ms, bibl, salts)

Kelly, T.R. *et al.*, *J.A.C.S.*, 1988, **110**, 6471-6480 (synth, bibl)

Saint-Jalmes, L. *et al.*, *Bull. Soc. Chim. Fr.*, 1993, **130**, 447-449 (synth)

Rao, A.V.R. *et al.*, *Heterocycles*, 1994, **37**, 1893-1912 (synth)

Wendt, J.A. *et al.*, *J.A.C.S.*, 1994, **116**, 9921-9926 (synth, bibl)

Clive, D.L.J. *et al.*, *J.A.C.S.*, 1994, **116**, 11275-11286 (synth, bibl)

Boger, D.L. *et al.*, *J.A.C.S.*, 1995, **117**, 11839-11849 (synth, cd, uv, pmr)

Kita, Y. *et al.*, *J.A.C.S.*, 2001, **123**, 3214-3222 (synth, abs config, bibl)


Sontag, B. *et al.*, *J. Antibiot.*, 2004, **57**, 823-828 (isol, uv, pmr, cmr)

Akai, S. *et al.*, *Chem. Eur. J.*, 2005, **11**, 6286-6297 (synth)

Chen, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 431-437 (biosynth)

Fredericamycin B**F-153**

[80450-64-2]



C₃₁H₂₃NO₉ 553.524


Position of methoxy uncertain, either C-11 or C-12. Prod. by *Streptomyces griseus* ATCC 49344. Weak antimicrobial agent. Black powder. Mp 350° dec. λ_{max} 260 ; 328 ; 348 ; 370 ; 387 ; 510 ; 548 ; 586 (MeCN aq. pH 6).

Pandey, R.C. *et al.*, *J. Antibiot.*, 1981, **34**, 1389-1401; 1402-1407 (*isol*, *props*)

Sontag, B. *et al.*, *J. Antibiot.*, 2004, **57**, 823-828 (*isol*, *uv*, *pmr*, *cmr*)

Chen, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 431-437 (*isol*, *biosynth*)

Fredericamycin E F-154
[1011723-58-2]



C₃₁H₂₃NO₁₀ 569.523

OMe position not certain. Prod. by *Streptomyces griseus* ATCC 49344. Cytotoxic. Dark red solid. [α]_D²⁰ +8 (c, 0.05 in CHCl₃). λ_{max} 262 (log ε 3.75); 303 (log ε 3.57); 317 (log ε 3.63); 330 (log ε 3.63); 358 (log ε 3.56); 372 (log ε 3.58); 393 (log ε 3.45); 464 (log ε 3.36); 496 (log ε 3.41) (CHCl₃).

Chen, Y. *et al.*, *J. Nat. Prod.*, 2008, **71**, 431-437 (*isol*, *uv*, *pmr*, *cmr*)

Fritilline A F-155
[1357-22-8]

C₂₇H₄₅NO₃ 431.657

Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria verticillata* (Liliaceae). Dehydrogenation gives Veranthridine.

Chu, T.-C. *et al.*, *CA*, 1960, **54**, 22695

Fritilline B F-156
[1357-23-9]

C₂₇H₄₃NO₃ 429.642

Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria verticillata* (Liliaceae).

Chu, T.-C. *et al.*, *CA*, 1960, **54**, 22695


Fritimine F-157
[664307-32-8]

C₂₃H₃₇NO₂ 359.551

Struct. unknown. Alkaloid from *Fritillaria roylei* (Liliaceae). Smooth muscle relaxant, antihypertensive agent. Much referred to in Chinese herbal formulations. Mp 167°. [α]_D²² -50.

Chou, T.-Q. *et al.*, *CA*, 1933, **27**, 3033

Fromiamycalin F-158
[163597-73-7]



C₄₅H₇₈N₆O₅ 783.148


Alkaloid from the New Caledonian starfish *Fromia monilis*. Cytotoxic. [α]_D -12 (as hydrochloride).

Palagiano, E. *et al.*, *Tetrahedron*, 1995, **51**, 3675-3682 (*isol*, *pmr*, *cmr*, *struct*)

Fructigenine A F-159
Puberuline

[127926-11-8]

[144606-96-2]



R = CH₂Ph

Absolute configuration

C₂₇H₂₉N₃O₃ 443.544

Prod. by *Penicillium fructigenum*, *Penicillium aurantiogriseum*, *Penicillium rugulosum*, *Penicillium puberulum* and *Penicillium piscarium*. Exhibits growth inhibitory activity against oat coleoptiles and L-5178Y cells. Solid. [α]_D -178 (c, 0.24 in CHCl₃). λ_{max} 246 (ε 12600); 276 (ε 2240); 285 (ε 1860) (EtOH) (Derep). λ_{max} 217 ; 250 (ε 9330); 298 (ε 12880) (MeOH) (Berdy).

N-Ac:

Needles (Et₂O/hexane). Mp 196-198°.

N-De-Ac: **Puberuline A**†

[192183-17-8]

C₂₅H₂₇N₃O₂ 401.507

Prod. by *Penicillium rugulosum*.

Stereochem. not determined.

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2937 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Boyes-Korkis, J.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1707 (*isol*, *ms*)

Solov'eva, T.F. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 1997, **33**, 66-69 (*Puberuline A*)

Kozlovskii, A.G. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 2000, **36**, 317-321 (*isol*)

Fructigenine B F-160

Verrucofortine. Verrucosine. Verrucozine

[113706-21-1]

[144525-64-4]

As Fructigenine A, F-159 with

R = -^{1'}CH₂^{2'}CH(CH₃)₂

C₂₄H₃₁N₃O₃ 409.572

Stereochem. of Verrucosine not confirmed. Alkaloid from *Penicillium fructigenum* and *Penicillium verrucosum* var. *cyclopium*. Solid. [α]_D -161.1 (c, 0.47 in CHCl₃). λ_{max} 247 (ε 9950); 277 (ε 1580); 285 (ε 1510) (EtOH) (Derep).

N-De-Ac: **Brevicompanine B**

[215121-47-4]

C₂₂H₂₉N₃O₂ 367.49

Alkaloid from *Penicillium brevicompactum* and *Aspergillus janus* (IBT 22274). Plant growth regulator. Amorph. solid. Mp 79-82°. [α]_D²⁰ -228.3 (c, 0.5 in EtOH). λ_{max} 208 ; 245 ; 300 (EtOH).

1'',2''-Didehydro: **Dehydroverrucosine**

[151298-54-3]

C₂₄H₂₉N₃O₃ 407.511

Prod. by *Penicillium verrucosum* var. *cyclopium*.

3''-Hydroxy: **Verrucosinol. Verrucozanol**

[151756-69-3]

C₂₄H₃₁N₃O₄ 425.527

Prod. by *Penicillium verrucosum* var. *cyclopium*.

Hodge, R.P. *et al.*, *J. Nat. Prod.*, 1988, **51**, 66-73 (*isol*, *pmr*, *cmr*)

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2937-2939 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Solovyeva, T.F. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 1992, **28**, 880-888 (*Verrucosine, Verrucosinol, Dehydroverrucosine*)


Kusano, M. *et al.*, *J.C.S. Perkin 1*, 1998, 2823-2826 (*Brevicompanine B*)

Matsumara, K. *et al.*, *Heterocycles*, 2001, **54**, 727-733 (*Brevicompanine B, synth*)

Sprogoe, K. *et al.*, *Tetrahedron*, 2005, **61**, 8718-8721 (*Brevicompanine B*)

2-(Fructosylamino)benzoic acid, 9CI F-161

1-(o-Carboxyanilino)-1-deoxyfructose



C₁₃H₁₇NO₇ 299.28

D-form [79896-08-5]

Isol. from a *Neurospora crassa* mutant. Patented for analgesic, hypoglycaemic, antitumour, etc. props.

4,6-O-Benzylidene:

Cryst. (Et₂O). Mp 192-194° dec. [α]_D²³ -119.2 (Py).


[79896-09-6]

Lingens, F. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1960, **315**, 58; 1963, **333**, 190 (*isol*, *biosynth*)

Lingens, F. *et al.*, *Annalen*, 1960, **630**, 84 (*synth*)

Fruit rot toxin A **F-162**

6-Methyl-2-(2-methyl-1-oxobutyl)-1-oxa-5-azaspiro[2.5]oct-6-ene-4,8-dione, 9CI.
FRT-A
[87334-35-8]




$C_{12}H_{15}NO_4$ 237.255
Isol. from *Macrophoma* sp. Phytotoxic to apples. Needles (C_6H_6). Mp 134-135.5°. $[\alpha]_D^{25} +13$ (EtOH). Isomer of Flavipucine, F-81.

Stereoisomer: Sapinopyridione

$C_{12}H_{15}NO_4$ 237.255
Prod. by *Sphaeropsis sapinea* strain D-55. Phytotoxin. Needles (CH_2Cl_2 /hexane). Mp 133-135°. $[\alpha]_D^{25} -11.3$ (c, 0.16 in EtOH). Probably the enantiomer of FRT-A. λ_{max} 320 (log ϵ 3.61) (MeCN).
Sassa, T. et al., *Agric. Biol. Chem.*, 1983, **47**, 1155-1157; 1417-1418 (isol, uv, ir, pmr)
Evidente, A. et al., *Phytochemistry*, 2006, **67**, 1019-1025 (Sapinopyridione)

Fruticosine **F-163**

[14058-43-6]



Absolute Configuration

$C_{22}H_{24}N_2O_4$ 380.443
Alkaloid from the leaves of *Kopsia fruticosa* (Apocynaceae). Cryst. (MeOH). Mp 225-226°. $[\alpha]_D^{20} -19$ (c, 2.60 in $CHCl_3$). pK_{a1} 4.62 (80% 2-methoxyethanol aq.). pK_{a1} 4.78 (2-methoxyethanol).

Picrate:

Cryst. + $1C_6H_6$ (C_6H_6). Mp 158-164° dec.

Ac: [14058-53-8]

Cryst. (petrol). Mp 115-118°.

12-Methoxy, N-de(methoxycarbonyl):

Jasminiflorine. 1-Demethoxycarbonyl-12-methoxyfruticosine
[114622-11-6]

$C_{21}H_{24}N_2O_3$ 352.432

Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Mp 230-233°. $[\alpha]_D -55.3$ ($CHCl_3$). λ_{max} 210 (ϵ 31600); 245 (ϵ 6920); 291 (ϵ 2340) (EtOH) (Derep).

16-Epimer: Fruticosamine

[14058-44-7]

$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the leaves of *Kopsia fruticosa* (Apocynaceae). Cryst. (Et₂O/pentane), cryst. + $\frac{1}{2}$ MeOH (MeOH). Mp 161-162° Mp 177-181° (MeOH solvate). $[\alpha]_D^{20} +43$ (c, 2.73 in $CHCl_3$). pK_{a1} 4.04 (80% 2-methoxyethanol aq.).

pK_{a1} 4.19 (2-methoxyethanol). Epimeric at the OH group.

16-Epimer, picrate:

Cryst. (EtOH). Mp 209-213° dec.

Guggisberg, A. et al., *Helv. Chim. Acta*, 1963, **46**, 679; 1966, **49**, 2321 (isol, synth, pmr, ms)

Battersby, A.R. et al., *J.C.S.*, 1963, 22 (isol, uv, ir, pmr, synth)

Battersby, A.R. et al., *J.C.S. (C)*, 1967, 813 (struct, pmr)


Klyne, W. et al., *Helv. Chim. Acta*, 1968, **51**, 1169 (ord, cd)

Ruangrunsi, N. et al., *Tet. Lett.*, 1987, **28**, 3679 (Jasminiflorine)

Glover, R.P. et al., *Magn. Reson. Chem.*, 2005, **43**, 483-485 (pmr, cmr, N-15 nmr)

Fruticosonine **F-164**

[73326-88-2]



Relative Configuration

$C_{20}H_{28}N_2O$ 312.454

Biosynthetically related to the *Aristolotelia* alkaloids. Alkaloid from *Aristolotelia fruticosa* (Elaeocarpaceae). Cryst. (Et₂O). Mp 120-121°. $[\alpha]_D^{20} +45.7$ (c, 0.5 in $CHCl_3$).

3'-ξ-Alcohol: Fruticosoline

[129350-17-0]

$C_{20}H_{30}N_2O$ 314.47


Alkaloid from *Aristolotelia fruticosa* (Elaeocarpaceae). Amorph. $[\alpha]_D^{20} +69.5$ (c, 0.4 in $CHCl_3$). λ_{max} 221; 282; 290 (MeOH).

Chaichit, N. et al., *Chem. Comm.*, 1979, 874-875 (pmr, ms, synth, cryst struct)

Hai, M.A. et al., *Indian J. Chem., Sect. B*, 1990, **29**, 586-587 (Fruticosoline)

Fuligorubin A **F-165**

[108343-55-1]



$C_{20}H_{23}NO_5$ 357.405


Tetramic acid deriv. Isol. from the slime mold *Fuligo septica* (Myxomycetes). Isol. from the sponge *Topsentia roquensis*. Prob. plays a role in slime mold phototaxis. Red cryst. + 2H₂O. Mp 150° dec. λ_{max} 243 (ϵ 14454); 425 (ϵ 26900) (MeOH) (Berdy). λ_{max} 243; 377 (MeOH-NaOH) (Berdy).

Casser, I. et al., *Angew. Chem., Int. Ed.*, 1987, **26**, 586 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Ley, S.V. et al., *Tetrahedron*, 1992, **48**, 1145 (synth)

Fulvanine A **F-166**

[132922-40-8]



Relative Configuration

$C_9H_{13}NO_5$ 215.205

Constit. of *Hemerocallis fulva* var. *kwanso*. Needles. Mp 118-119°. $[\alpha]_D -0.52$ (c, 1 in MeOH).

5-Me ether: Fulvanine D

[155944-19-7]

$C_{10}H_{15}NO_5$ 229.232

Constit. of *Hemerocallis fulva* var. *kwanso*. Powder. Mp 80-83°. $[\alpha]_D -0.3$ (c, 1.4 in MeOH).

6'-Deoxy: Fulvanine B

[132922-41-9]

$C_9H_{13}NO_4$ 199.206

Isol. from *Hemerocallis fulva* var. *kwanso*. Needles. Mp 132-133°. $[\alpha]_D -0.3$ (c, 0.8 in MeOH).

6'-Deoxy, 5-Me ether: Fulvanine C

[132922-42-0]


$C_{10}H_{15}NO_4$ 213.233

Constit. of *Hemerocallis fulva* var. *kwanso*. Amorph. powder.

Inoue, T. et al., *Chem. Pharm. Bull.*, 1990, **38**, 3187-3189; 1994, **42**, 154-155 (isol, pmr, cmr)

Fulvanine E **F-167**

[155944-20-0]



Relative Configuration


$C_9H_{11}NO_4$ 197.19

Constit. of *Hemerocallis fulva* var. *kwanso*. Needles (MeOH). Mp 210-212° dec. $[\alpha]_D +1.3$ (c, 0.2 in MeOH).

Inoue, T. et al., *Chem. Pharm. Bull.*, 1994, **42**, 154-155

Fulvine **F-168**

14,19-Dihydro-13-hydroxy-20-norcrotalaria-11,15-dione, 9CI
[6029-87-4]



Absolute configuration

$C_{16}H_{23}NO_5$ 309.361

Alkaloid from *Crotalaria fulva*, *Crotalaria crispata*, *Crotalaria madurensis* and *Crotalaria paniculata* (Fabaceae). Shows antineoplastic props. Prisms (Me₂CO). Mp 212-213°. $[\alpha]_D^{20} -50.8$ (c, 1 in $CHCl_3$). Log P -0.81 (uncertain value) (calc). Ester of Retronecine in T-188 with Fulvic acid.

▶ Heptatotoxin, pneumotoxin, teratogen. RC1300000

Hydrochloride:

Prisms. Mp 285° dec.

Picrate:

Yellow needles. Mp 250° dec.

N-Oxide: Fulvine N-oxide

$C_{16}H_{23}NO_6$ 325.361

Alkaloid from *Crotalaria fulva* (Fabaceae). Needles (Me₂CO). Mp 198° dec.

N-Oxide, picrate:

Yellow needles. Mp 185° dec.

13-Epimer: **Crispatine**. *Cryspatine*

[6029-88-5]

C₁₆H₂₃NO₅ 309.361

Alkaloid from *Crotalaria crispata* and *Crotalaria madurensis* (Fabaceae). Shows antineoplastic props. Needles (petrol). Mp 137-138°. $[\alpha]_D^{20} +40.7$ (c, 1.94 in EtOH). Log P -0.81 (uncertain value) (calc). Ester of Retronecine in T-188 with Crispatic acid. An alkaloid of *C. leschenaultii*, previously reported as Crispatine has subsequently been shown to be Crotaleschenine, C-773.

► Heptatotoxin, exp. carcinogen.

12-Epimer or 13,14-diepimer (1): [57129-98-3]

C₁₆H₂₃NO₅ 309.361

Alkaloid from *Crotalaria madurensis* (Fabaceae). Mp 242-243°. Ester of Retronecine in T-188 with Chromaduric acid.

12-Epimer or 13,14-diepimer (2): **Isocromadurine**

[57495-69-9]

C₁₆H₂₃NO₅ 309.361

Alkaloid from *Crotalaria madurensis* (Fabaceae). Mp 135-136°. Ester of Retronecine in T-188 with Isochromaduric acid.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1963, **16**, 239 (*isol, struct*)

Chalmers, A.H. *et al.*, *J. Chromatogr.*, 1965, **20**, 270 (*tlc*)

Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (*cd*)

Sussman, J.L. *et al.*, *Acta Cryst. B*, 1973, **29**, 2918 (*cryst struct*)

Rashkes, Ya. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 40 (*ms*)

Rao, P.G. *et al.*, *Experientia*, 1975, **31**, 878 (*Isocromadurine*)

Rao, P.G. *et al.*, *Indian J. Chem.*, 1975, **13**, 870 (*Cromadurine*)

Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417 (*cmr*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Mackay, M.F. *et al.*, *Acta Cryst. C*, 1984, **40**, 470 (*cryst struct, Crispatine*)


Vedejs, E. *et al.*, *J.A.C.S.*, 1984, **106**, 3030 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FOT000

Fumadensine

F-169

[108906-89-4]



C₃₀H₃₄N₂O₆ 518.608

Alkaloid from *Fumaria densiflora* (Papaveraceae). Cryst. (MeOH). Mp 168-170°.


Racemic.

Zarga, M.H.A. *et al.*, *Phytochemistry*, 1987, **26**, 1233 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Fumaflorine

F-170

[180386-83-8]



C₂₀H₁₅NO₇ 381.341

Alkaloid from *Fumaria densiflora* (Papaveraceae). Yellowish needles (MeOH). Mp 218-220°. λ_{max} 236 (log ϵ 4.69); 330 (log ϵ 3.98) (MeOH).

Me ester: [180386-82-7]

C₂₁H₁₇NO₇ 395.368


Alkaloid from aerial parts of *Fumaria demiflora*. Red needles (MeOH). Mp 184-186°. Prob. an artifact. λ_{max} 213 (log ϵ 4.62); 236 (log ϵ 4.54); 286 (sh) (log ϵ 4.02); 334 (log ϵ 4.07) (No solvent reported).

Táborská, E. *et al.*, *Heterocycles*, 1997, **45**, 817-821 (*isol, uv, ir, pmr, cmr, ms, struct*)

Fumaramidine

F-171

6-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]methylene]-6,7-dihydro-8H-1,3-dioxolo[4,5-e]isoindol-8-one, 9CI [76202-52-3]



R = R' = Me

C₂₂H₂₄N₂O₅ 396.442

Alkaloid from *Fumaria parviflora*.

Amorph.

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 169-178 (*isol, pmr, ms*)

Lamblin, M. *et al.*, *Tetrahedron*, 2006, **62**, 2917-2921 (*synth*)

Fumaramine

F-172

6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]-6,7-dihydro-8H-1,3-dioxolo[4,5-e]isoindol-8-one, 9CI. Adumidiceine imide. Alkaloid F46

As Fumaramidine, F-171 with

R, R' = -CH₂-

C₂₁H₂₀N₂O₅ 380.399

(*E*)-form [88728-13-6]

Alkaloid from aerial parts of *Fumaria vaillantii*. Poss. artifact.

(*Z*)-form [30341-99-2]

Alkaloid from *Fumaria parviflora*, Fu-

maria vaillantii and *Corydalis ochroleuca* (Papaveraceae). Yellow cryst. (EtOH).

Mp 220-221°. May be an artifact of work-up.

Shamma, M. *et al.*, *Chem. Comm.*, 1975, 89 (*struct, synth*)

Rodrigo, R.G.A. *et al.*, *Can. J. Chem.*, 1976, **54**, 471 (*ir, pmr, ms*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 169 (*isol, pmr*)

Sener, B. *et al.*, *Phytochemistry*, 1983, **22**, 2073 (*E-form*)


Fumaric acid, 8CI

F-173

(*E*)-2-Butenedioic acid, 9CI. *trans*-Ethylene-1,2-dicarboxylic acid. *Paramaleic acid*. *Glauic acid*. *Boletic acid*. FEMA 2488. E297

[110-17-8]

[7704-73-6]



C₄H₄O₄ 116.073

Geometrical isomer of Maleic acid.

Manuf. by catalytic or thermal isomerisation of maleic anhydride or maleic acid. Occurs in many plants, e.g. *Fumaria officinalis*, *Boletus scaber*, *Fomes ignarius*. Produced by *Rhizopus nigricans*.

Essential respiratory intermed. in plant and animal tissues. Used as di-Na salt for pptn. sepn. of Th from lanthanides. Used in manuf. of unsaturated polyester resins and as acidulant and flavouring agent in foods and pharmaceutical formulations. Used in systemic and topical treatment of psoriasis. Monoclinic, prismatic needles or leaflets (H₂O). Sol. EtOH; spar. sol. H₂O, Et₂O, Me₂CO; prac. insol. C₆H₆, CHCl₃. Mp 300-302° (sealed tube). pK_a 4.54 (25°). At 230° forms maleic anhydride.

► Skin and eye irritant. Nephrotoxic. LD₅₀ (rat, ori) 10700 mg/kg. LS9625000

2-Methylpropylamide: 4-[(2-Methylpropyl)amino]-4-oxo-2-butenic acid, 9CI.

Fumaric acid isobutylamide

[174204-82-1]

C₈H₁₃NO₃ 171.196

Alkaloid from *Piper hancei*.


[5873-57-4, 6915-18-0, 17013-01-3]

Narui, T. *et al.*, *Nat. Med. (Tokyo)*, 1995, **49**, 438-441 (*isobutylamide*)

Fumaricine

F-174

3',4',6,8-Tetrahydro-6',7'-dimethoxy-2'-methylspiro[7H-indeno[4,5-d]-1,3-dioxole-7,1'(2'H)-isoquinolin]-8-ol, 9CI



(-)-form

C₂₁H₂₃NO₅ 369.416

(-)-form**Alkaloid F37**

[24181-77-9]

Alkaloid from *Fumaria officinalis*, *Fumaria schrammii* and *Fumaria parviflora* (Papaveraceae). Prisms (MeOH). Mp 177°. $[\alpha]_D^{25}$ -31 (c, 0.97 in CHCl_3). Prob. identical with *O*-Methylfumarophycinol from Herba Fumariae Officinalis (Mollov *et al.*). λ_{max} 207 (log ϵ 4.74); 235 (log ϵ 3.94); 288 (log ϵ 3.74) (EtOH).

Ac: O-Methylfumarophycine

[36017-45-5]

 $\text{C}_{23}\text{H}_{25}\text{NO}_6$ 411.454

Alkaloid from *Fumaria officinalis* and aerial parts of *Fumaria kralikii* (Papaveraceae). Cryst. ($\text{Et}_2\text{O}/\text{MeOH}$). Mp 124-127°. $[\alpha]_D^{25}$ -51 (CHCl_3). $[\alpha]_D^{25}$ -30 (c, 0.158 in MeOH). λ_{max} 208 (log ϵ 4.79); 232 (sh) (log ϵ 4.18); 285 (log ϵ 3.86) (MeOH).

O²-De-Me: Fumaritine

[24181-78-0]

 $\text{C}_{20}\text{H}_{21}\text{NO}_5$ 355.39

Alkaloid from *Fumaria officinalis*, the roots of *Fumaria schleicheri* and the leaves of *Fumaria muralis* ssp. *boraei* (Papaveraceae). Needles (dry Et_2O). Mp 157°. Opt. rotn. not recorded.

O²-De-Me, hydrochloride: Mp 224°.**O²-De-Me, hydrobromide:** Mp 219°.**O²-De-Me, N-oxide: Fumaritine N-oxide.****Alkaloid F_k-5**

[60604-88-8]

 $\text{C}_{20}\text{H}_{21}\text{NO}_6$ 371.389

Alkaloid from *Fumaria kralikii*, *Fumaria indica* and *Fumaria officinalis* (Papaveraceae). Cryst. ($\text{C}_6\text{H}_6/\text{MeOH}$). Mp 204°. $[\alpha]_D^{25}$ +49.2 (MeOH). λ_{max} 212 (log ϵ 4.29); 287 (log ϵ 3.42) (MeOH).

O²-De-Me, 8-Ac: Fumarophycine

[30833-08-0]

 $\text{C}_{22}\text{H}_{23}\text{NO}_6$ 397.427

Alkaloid from *Fumaria officinalis*, the aerial parts of *Fumaria kralikii* and the leaves of *Fumaria muralis* ssp. *boraei* (Papaveraceae). Cryst. (MeOH). Mp 107-109°. $[\alpha]_D^{19}$ -67.5 (c, 1.0 in CHCl_3).

O²-De-Me, N-Me: Fumaritine N-metho-salt

[82054-20-4]

 $\text{C}_{21}\text{H}_{24}\text{NO}_5^{\oplus}$ 370.424

Tentatively identified as alkaloid in leaves of *Fumaria muralis* ssp. *boraei* (Papaveraceae). No phys. props. reported.

O²-De-Me, N-de-Me: Norfumaritine

[100016-58-8]

 $\text{C}_{19}\text{H}_{19}\text{NO}_5$ 341.363

Alkaloid from leaves of *Fumaria kralikii* (Papaveraceae). Amorph. $[\alpha]_D^{25}$ -16 (c, 0.12 in CHCl_3). λ_{max} 238 (log ϵ 4.65); 240 (sh) (log ϵ 3.89); 287 (log ϵ 3.96) (MeOH).

Ketone: Parfumidine

[31225-67-9]

 $\text{C}_{21}\text{H}_{21}\text{NO}_5$ 367.401

Alkaloid from *Fumaria parviflora*, *Fumaria officinalis* and *Fumaria vaillantii* (Papaveraceae). Cryst. (MeOH, EtOH

or Et_2O). Mp 171-172° (165-168°). $[\alpha]_D^{22}$ +33.3 (c, 0.5 in CHCl_3). λ_{max} 235(log ϵ 4.46); 263 (log ϵ 4.14); 290 (sh);360 (log ϵ 3.4) (no solvent reported).**Ketone, O²-de-Me: Parfumine, Fumarilicine**

[28230-70-8]

[56297-73-5]

 $\text{C}_{20}\text{H}_{19}\text{NO}_5$ 353.374

Alkaloid from *Fumaria kralikii*, *Fumaria parviflora*, *Fumaria rostellata*, *Fumaria vaillantii*, *Fumaria muralis* ssp. *boraei*, *Fumaria judaica*, *Fumaria schleicheri* and *Fumaria schrammii* (Papaveraceae). Cryst. (EtOH or CHCl_3). Mp 118-120° (111-112°). $[\alpha]_D^{23}$ +18 (c, 1.1 in CHCl_3). λ_{max} 202 (log ϵ 4.61); 234 (log ϵ 4.38); 261 (log ϵ 4.01); 355 (log ϵ 3.42) (MeOH).

Ketone, O²-de-Me, O²- β -D-glucopyranoside: Parviflorine[†]

[75969-88-9]

 $\text{C}_{26}\text{H}_{29}\text{NO}_{10}$ 515.516

Alkaloid from *Fumaria parviflora* (Papaveraceae). Yellow cryst. (MeOH). Mp 230-232°. $[\alpha]_D$ +1 (c, 0.0124 in MeOH). λ_{max} 233 (log ϵ 4.5); 260 (log ϵ 4.18); 288 (sh) (log ϵ 3.7); 352 (log ϵ 3.66) (MeOH).

Ketone, O²-de-Me, O²-Ac: Mp 198-199°.**Ketone, O³-de-Me: Isoparfumine**

[107080-29-5]

 $\text{C}_{20}\text{H}_{19}\text{NO}_5$ 353.374

Alkaloid from *Rupicapnos africana* (Papaveraceae). Cryst. (MeOH). Mp 206-208°. $[\alpha]_D^{25}$ +54 (c, 0.792 in CHCl_3). λ_{max} 210 (log ϵ 4.3); 234 (log ϵ 4.35); 260 (log ϵ 4.6); 293 (sh); 350 (log ϵ 3.47) (EtOH).

8-Deoxy, O²-de-Me, N-oxide: Papracinine

[145290-12-6]

 $\text{C}_{20}\text{H}_{21}\text{NO}_5$ 355.39

Alkaloid from aerial parts of *Fumaria indica* (Papaveraceae). Amorph. solid. $[\alpha]_D$ +23 (MeOH). λ_{max} 200 (log ϵ 4.25); 220 (log ϵ 4.13); 291 (log ϵ 3.97); 318 (log ϵ 4.04) (MeOH).

(±)-form [24869-02-1]

Synthetic. Prisms (EtOH/ Et_2O). Mp 147.5-149°.

O²-De-Me: [32420-39-6]

Synthetic. Prisms + 0.5 H_2O (MeOH/ Et_2O). Mp 193-194°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 438-444 (isol)

Platonova, T.F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 181-186; *CA*, **50**, 13960a (Fumaritine, isol)

Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1967, **20**, 557-559; 1969, **189**, 1262; *CA*, **67**, 117013 (Fumarophycine, *O*-Methylfumarophycine, Parfumidine)

Saunders, J.K. *et al.*, *Can. J. Chem.*, 1968, **46**, 2873-2875 (uv, ir, struct)

MacLean, D.B. *et al.*, *Can. J. Chem.*, 1969, **47**, 3593-3599 (Fumaricine, Fumaritine, pmr, ms, struct)

Israilov, I.A. *et al.*, *Dokl. Chem. (Engl. Transl.)*, 1969, **189**, 999-1000 (Parfumine, isol, uv, ir, pmr, ms, struct)

Kishimoto, T. *et al.*, *J.C.S. (C)*, 1969, 2600-2602; 1971, 1644-1647 (synth, ir, pmr, ms)

Israilov, I.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 518 (Parfumidine)

Castillo, M. *et al.*, *Can. J. Chem.*, 1971, **49**,

139-142 (Fumarophycine, *O*-

Methylfumarophycine, pmr, ms, struct)

Yu, C.K. *et al.*, *Can. J. Chem.*, 1971, **49**, 3025-

3037 (ms)

Kiryakov, Kh. *et al.*, *Can. J. Chem.*, 1979, **57**,

53-56 (Fumaritine, Fumaritine N-oxide, ir,

pmr, cmr, ms)

Nasirov, S.M. *et al.*, *Chem. Nat. Compd.*

(Engl. Transl.), 1980, **16**, 55-60 (Parfumine,

cryst struct)

Hussain, S.F. *et al.*, *Tet. Lett.*, 1980, **21**, 1909-

1912 (Fumaritine, Parfumine, Parviflorine,

pmr, cd, abs config)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**,

169-178 (Parfumidine, Parviflorine, isol, pmr,

ms, uv, cd, struct)

Loukis, A. *et al.*, *J. Pharm. Pharmacol., Suppl.*,

1981, **33**, 16P (Fumaritine N-methosalt,

Parfumine)

Alimova, M. *et al.*, *Chem. Nat. Compd. (Engl.*

Transl.), 1982, **18**, 608-609 (isol)

Gozler, B. *et al.*, *J. Nat. Prod.*, 1983, **46**, 433-

435 (*O*-Methylfumarophycine)

Colton, M.D. *et al.*, *J. Nat. Prod.*, 1985, **48**,

846-847 (Norfumaritine)

Castedo, L. *et al.*, *Heterocycles*, 1986, **24**,

2781-2784 (Isoparfumine)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1992,

31, 2869-2872 (Papracinine)

Seger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**,


882-886 (Fumarophycine, Parfumine, *O*-

Methylfumarophycine, pmr, cmr)

Fumariflorine**F-175**

6-[2-(Dimethylamino)ethyl]-1,3-benzodioxole-5-carboxylic acid, 9CI. 2-[2-(Dimethylamino)ethyl]-4,5-methylenedioxybenzoic acid

[76948-81-7]

 $\text{C}_{12}\text{H}_{15}\text{NO}_4$ 237.255

Alkaloid present in *Fumaria parviflora* dried extracts (Papaveraceae). Prob. a final metab. of the secophthalideisoquinoline alkaloids. Isol. as the Et ester.

Et ester: Fumariflorine ethyl ester

[76202-50-1]


 $\text{C}_{14}\text{H}_{19}\text{NO}_4$ 265.308

Alkaloid from *Fumaria parviflora* (Papaveraceae). Wax. Almost certainly an artifact of ethylation.

Hussain, S.F. *et al.*, *Tet. Lett.*, 1980, **21**, 1693 (isol, ms, ir, pmr, struct, synth)

Fumariline**F-176**

7,8-Dihydro-6-methylspiro[1,3-dioxolo[4,5-g]isoquinoline-5(6H),7'-[7H]indeno[4,5-d][1,3]dioxol]-8'(6'H)-one, 9CI

**(S)-form**

C₂₀H₁₇NO₅ 351.358
CAS numbering shown.

(S)-form [20411-03-4]

Alkaloid from *Fumaria officinalis*, *Fumaria parviflora*, *Fumaria schrammii*, *Fumaria vaillantii*, the seeds of *Fumaria indica*, the leaves of *Fumaria muralis* var. *boraei*, and from the flowers and unripe fruit of *Fumaria rostellata* (Papaveraceae). Cryst. (MeOH or CHCl₃/MeOH), also descr. as oil. Mp 74-76° Mp 138° Mp 144°. [α]_D²⁵ +82.5 (c, 0.6 in CHCl₃). [α]_D²⁰ +138 (c, 1.05 in CHCl₃). [α]_D¹⁵ +96 (c, 1.0 in CHCl₃). λ_{max} 203 (log ε 4.6); 237 (log ε 4.31); 263 (log ε 4.05); 294 (log ε 3.66); 355 (log ε 3.51) (EtOH).

▶ WH1306520

(8R)-Alcohol: Dihydrofumariline 1

[87638-96-8]
C₂₀H₁₉NO₅ 353.374

Alkaloid from *Fumaria officinalis* (Papaveraceae). Mp 191-193°. No opt. rotn. reported.

(8S)-Alcohol: Dihydrofumariline 2. Capreoline

[24181-80-4]
C₂₀H₁₉NO₅ 353.374

Synthetic. Alkaloid from *Fumaria capreolata* (Papaveraceae). Mp 135-137°. [α]_D²⁵ +106 (CHCl₃).

8-Alcohol: Dihydrofumariline

C₂₀H₁₉NO₅ 353.374

Alkaloid from *Fumaria schrammii* (Papaveraceae). Mp 184-187°. Config. not definitely detd., but appears to be identical with Dihydrofumariline 2, the discrepancy in Mps being due to dimorphism.

(±)-form [32420-40-9]

Synthetic. Prisms + 0.25H₂O (MeOH). Mp 217-220°.

8-Alcohol: Synthetic. Mp 198-199°.

[56082-99-6]

Saunders, J.K. *et al.*, *Can. J. Chem.*, 1968, **46**, 2873-2875 (*uv, ir, pmr, struct*)

Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1969, **47**, 1103-1105 (*isol*)

MacLean, D.B. *et al.*, *Can. J. Chem.*, 1969, **47**, 3593-3599 (*uv, ir, pmr, ms, struct*)

Kishimoto, T. *et al.*, *J.C.S. (C)*, 1971, 1644-1647 (*synth, ir, pmr*)

Shamma, M. *et al.*, *Chem. Comm.*, 1972, 310-311 (*abs config*)

Pandey, V.B. *et al.*, *Phytochemistry*, 1979, **18**, 695-696 (*isol, uv, ir, pmr, ms*)

Popova, M.E. *et al.*, *Planta Med.*, 1980, **40**, 156-160 (*Dihydrofumariline*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 169-178 (*isol, ir, pmr, ms*)

Mardirossian, Z. *et al.*, *Phytochemistry*, 1983, **22**, 759-761 (*Dihydrofumarilines*)

Hanaoka, M. *et al.*, *Tet. Lett.*, 1985, **26**, 917-920 (*Dihydrofumarilines, synth*)

Fumarinine

F-177

C₁₆H₁₅NO₄ 285.299

Struct. unknown. Alkaloid from roots of *Fumaria schleicheri* (Papaveraceae). Mp 189-190°. Conts. 1NMe gp.


Hydrochloride: Mp 255-257°.

Oxalate: Mp 213°.

Platonova, T.F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 181-186; *CA*, **50**, 13960

Fumaritridine

F-178



Absolute configuration

C₂₁H₂₃NO₅ 369.416

(+)-form [37646-04-1]

Alkaloid from *Fumaria rostellata* (Papaveraceae). Needles (EtOH). Mp 189-193° Mp 198-200°. [α]_D²² +18 (c, 1.0 in CHCl₃). Originally, this alkaloid was erroneously assigned a spirobenzylisoquinoline struct.

Me ether: Fumaritrine

[38907-25-4]

C₂₂H₂₅NO₅ 383.443

Alkaloid from the Bulgarian medicinal herb "Herba Fumariae" (Papaveraceae). Needles (EtOH). Mp 153-155°. [α]_D²⁵ +14.5 (c, 0.055 in CHCl₃).

(±)-form

Me ether: Synthetic. Mp 164.5-166°.

Mollov, N.M. *et al.*, *Phytochemistry*, 1972, **11**, 2331 (*uv, pmr, ms*)

Blaskó, G. *et al.*, *Tet. Lett.*, 1981, **22**, 3143 (*uv, pmr, synth, struct, abs config*)

Hanaoka, M. *et al.*, *Tet. Lett.*, 1983, **24**, 3845 (*synth, pmr, Fumaritrine*)

Fumarizine

F-179

[133740-04-2]

C₂₀H₂₁NO₅ 355.39


Was assigned the same struct. as Marshaline in L-80 but props. of synthetic material differ from those of the natural alkaloid. It is prob. a regiosomer. Alkaloid from *Fumaria indica* (Papaveraceae). Amorph. λ_{max} 240 ; 245 ; 285 (MeOH). Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1989, **60**, 552-553 (*isol, uv, ir, pmr*)
Takaba, K. *et al.*, *Yakugaku Zasshi*, 1997, **117**, 555-559

Fumarofine

F-180

Fumarostelline. Alkaloid F38

[34114-84-6]



C₂₀H₁₉NO₆ 369.373

Alkaloid from *Fumaria officinalis* and *Fumaria microcarpa* (Papaveraceae).

Cryst. (dioxan/MeOH). Mp 256° (shrinking at 240-245°). Originally, this alkaloid was erroneously assigned a spirobenzylisoquinoline struct.

O¹¹-Me:

Cryst. (MeOH/Et₂O). Mp 245-246°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 438 (*isol*)

Yu, C.K. *et al.*, *Can. J. Chem.*, 1971, **49**, 3020 (*uv, pmr, ms*)


Blaskó, G. *et al.*, *Tet. Lett.*, 1981, **22**, 3135 (*struct*)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2172 (*synth, ir, pmr*)

Fumigaclavine B

F-181

6,8-Dimethylergolin-9-ol, 9CI



(-)-form

C₁₆H₂₀N₂O 256.347

(-)-form [6879-93-2]

Alkaloid from *Aspergillus fumigatus* and *Rhizopus arrhizus*. Mycotoxin. Mp 244-245° Mp 265-267° (double Mp). [α]_D²⁵ -6.3 (c, 1.2 in MeOH). λ_{max} 225 (log ε 4.49); 275 (log ε 3.79); 282 (log ε 3.82); 293 (log ε 3.72) (EtOH).

Methodide:

Cryst. (EtOH/Et₂O). Mp 309-310° dec.

O-Ac: Fumigaclavine A

[6879-59-0]

C₁₈H₂₂N₂O₂ 298.384

Alkaloid from *Aspergillus fumigatus* and *Aspergillus tamarit*. Mycotoxin. Needles (MeOH aq.). Mp 84-85°. [α]_D²² -56.7 (c, 1.5 in MeOH) (as hydrochloride).

▶ KE6345400

O-Ac; hydrochloride:

Prisms (EtOH). Mp 304-305° dec.

(inserted at 300°). [α]_{Hg}²² -56.7 (c, 1.5 in MeOH).

8,9-Diepimer: Isofumigaclavine B. Roquefortine B

[58800-20-7]

C₁₆H₂₀N₂O 256.347

Metab. from *Penicillium roquefortii* and *Penicillium crustosum*. Mp 278-282° dec. (222-224°). [α]_D²⁵ -147 (c, 0.50 in Py). λ_{max} 224 (log ε 4.65); 275 (log ε 3.79); 282 (log ε 3.84); 294 (log ε 3.78) (EtOH).

8,9-Diepimer, O-Ac: Isofumigaclavine A. Roquefortine A

[58800-19-4]

C₁₈H₂₂N₂O₂ 298.384

Metab. from *Penicillium roquefortii* and *Penicillium crustosum*. Cryst. (C₆H₆). Mp 190-193° (182° dec.). [α]_D²² -54.1 (c, 0.67 in CHCl₃). [α]_D¹⁵ -130 (c, 0.56 in Py). λ_{max} 226 (log ε 4.42); 271 (sh) (log ε 3.77); 283 (log ε 3.81); 293 (log ε 3.75) (95% EtOH).

(±)-form


Synthetic. Mp 199-200° dec.

8,9-Diepimer: [98570-39-9]

Synthetic. Cryst. (MeOH). Mp 271-273° dec.

Spilsbury, J.F. *et al.*, *J.C.S.*, 1961, 2085-2089 (*Fumigaclavine A*, *isol*, *uv*, *ir*, *struct*)Bach, N.J. *et al.*, *J.O.C.*, 1974, **39**, 1272-1276 (*pmr*, *cmr*, *config*)Ohmono, S. *et al.*, *Agric. Biol. Chem.*, 1975, **39**, 1333-1334 (*Isofumigaclavines*, *isol*, *uv*, *ir*, *pmr*)Scott, P.M. *et al.*, *J. Agric. Food Chem.*, 1976, **24**, 865-868 (*Isofumigaclavines*, *isol*, *uv*, *ms*)Arnoux, B. *et al.*, *J. Chem. Res., Synop.*, 1978, 210-211; 2829-2849 (*Isofumigaclavines*, *pmr*, *cmr*, *cryst struct*, *abs config*)Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 655-657 (*Isofumigaclavines*, *isol*)Kiguchi, T. *et al.*, *Heterocycles*, 1985, **23**, 1925-1928 (*synth*, *pmr*)Ninomiya, I. *et al.*, *J.C.S. Perkin I*, 1985, 941-948 (*Isofumigaclavines*, *synth*, *ir*, *pmr*)Ninomiya, I. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 23-30 (*synth*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 545; 553**Fumigaclavine C****F-182**

[62867-47-4]




Absolute Configuration

C₂₃H₃₀N₂O₂ 366.502Alkaloid prod. by *Aspergillus fumigatus*. Mycotoxin. Needles (MeOH). Mp 194°.


► KE6345500

Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 826 (*isol*, *uv*, *ir*, *ms*, *pmr*, *cmr*, *cryst struct*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 560**Fumimycin****F-183**

[942472-95-9]

C₁₆H₁₅NO₇ 333.297


Related to Sorbicillactone A, S-382.

Prod. by *Aspergillus fumisynnematus* F746. Inhibitor of peptide deformylase. Antibacterial agent. Powder. [α]_D -11.9 (c, 0.26 in MeOH). λ_{max} 212 (log ε 4.38); 239 (log ε 4.26); 257 (log ε 4.13); 266 (log ε 3.98); 316 (log ε 3.46) (MeOH).Kwon, Y.-J. *et al.*, *Org. Lett.*, 2007, **9**, 2449-2451 (*isol*, *pmr*, *cmr*)**Fumiquinazoline I****F-184**

Absolute Configuration

C₂₇H₂₉N₅O₄ 487.557Isol. from an *Acremonium* sp. obt. from the tunicate *Ecteinascidia turbinata*. Solid. Mp 116-120°. [α]_D -138 (c, 0.001 in CHCl₃). λ_{max} 224 (ε 63300); 230 (sh) (ε 57200); 255 (ε 26000); 266 (sh) (ε 21700); 277 (ε 17000); 304 (ε 5100) (MeOH).Belofsky, G.N. *et al.*, *Chem. Eur. J.*, 2000, **6**, 1355-1360 (*isol*, *uv*, *pmr*, *cmr*)Snider, B.B. *et al.*, *J.O.C.*, 2003, **68**, 545-563 (*synth*)**Fumiquinazoline A****F-185**

[140715-85-1]



Absolute Configuration

C₂₄H₂₃N₅O₄ 445.477Prod. by *Aspergillus fumigatus* isol. from the gastrointestinal tract of the fish *Pseudolabrus japonicus*. Cytotoxic agent. Cryst. (CH₂Cl₂). Mp 178-182°. [α]_D³⁵ -214.5 (c, 0.47 in CHCl₃). λ_{max} 208 (ε 37900); 226 (ε 30900); 233 (ε 27400); 256 (ε 14800); 265 (ε 13500); 277 (ε 10230); 305 (ε 3450); 327 (ε 2770) (MeOH) (Berdy).**4-Methoxy: Fumiquinazoline E**


[140715-87-3]

C₂₅H₂₅N₅O₅ 475.503Prod. by *Aspergillus fumigatus*. Cytotoxic agent. Pale yellow powder. Mp 168-172°. [α]_D -143.3 (c, 0.2 in CHCl₃). λ_{max} 210 (log ε 4.54); 226 (log ε 4.48); 233 (log ε 4.44); 256 (log ε 4.16); 278 (log ε 4.06); 304 (log ε 3.62); 317 (log ε 3.52) (EtOH).**4-Epimer: Fumiquinazoline B**

[140852-71-7]

C₂₄H₂₃N₅O₄ 445.477From *Aspergillus fumigatus* in *Pseudolabrus japonicus*. Cytotoxic agent. Cryst. (Me₂CO). Mp 174-176°. [α]_D²¹ -196.7 (c, 0.38 in CHCl₃). λ_{max} 206 (ε 58500); 225 (ε 49200); 232 (ε 44700); 256 (ε 24000); 266 (ε 20900); 277 (ε 17300); 305 (ε 6150); 317 (ε 4800) (MeOH) (Berdy).Numata, A. *et al.*, *Tet. Lett.*, 1992, **33**, 1621-1624 (*isol*, *pmr*, *cmr*, *struct*)Takahashi, C. *et al.*, *J.C.S. Perkin I*, 1995, 2345-2353 (*isol*, *uv*, *ir*, *pmr*, *cmr*)Snider, B.B. *et al.*, *J.O.C.*, 2003, **68**, 545-563 (*synth*)**Fumiquinazoline C****F-186**


[140924-01-2]



Absolute Configuration

C₂₄H₂₁N₅O₄ 443.461Prod. by *Aspergillus fumigatus* isol. from the gastrointestinal tract of the fish *Pseudolabrus japonicus*. Cytotoxic agent. Cryst. + 1Me₂CO (Me₂CO). Mp 244-246°. [α]_D²⁵ -193.7 (c, 0.31 in CHCl₃). λ_{max} 207 (ε 36300); 225 (ε 30200); 260 (ε 11480); 271 (ε 10470); 282 (ε 9550); 304 (ε 4074); 317 (ε 3160) (MeOH) (Berdy).Numata, A. *et al.*, *Tet. Lett.*, 1992, **33**, 1621-1624 (*isol*, *pmr*, *cryst struct*)Takahashi, C. *et al.*, *J.C.S. Perkin I*, 1995, 2345-2353 (*isol*, *uv*, *ir*, *pmr*, *cd*)Snider, B.B. *et al.*, *J.O.C.*, 2003, **68**, 545-563 (*synth*)Afiyatullof, S.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 236-238 (*isol*, *cmr*)**Fumiquinazoline D****F-187**


[140715-86-2]



Absolute Configuration

C₂₄H₂₁N₅O₄ 443.461Prod. by *Aspergillus fumigatus*. Cytotoxic agent. Prisms (Me₂CO). Mp 214-216°. [α]_D²² +86.2 (c, 0.15 in CHCl₃). λ_{max} 205 (log ε 4.3); 225 (log ε 4.26); 232 (log ε 4.24); 254 (log ε 3.91); 265 (log ε 3.84); 276 (log ε 3.77); 304 (log ε 3.43); 316 (log ε 3.54) (EtOH).Takahashi, C. *et al.*, *J.C.S. Perkin I*, 1995, 2345-2353 (*isol*, *uv*, *ir*, *cd*, *pmr*, *cmr*, *cryst struct*)Afiyatullof, S.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 236-238 (*isol*, *cmr*)**Fumiquinazoline F****F-188**

[169626-35-1]



$C_{21}H_{18}N_4O_2$ 358.399
Prod. by a marine-derived *Aspergillus fumigatus*. Cytotoxic agent. Pale yellow powder. Mp 88-90°. $[\alpha]_D^{25}$ -411.2 (c, 1.4 in $CHCl_3$). λ_{max} 207 (log ϵ 4.71); 219 (log ϵ 4.73); 270 (log ϵ 4.13); 277 (log ϵ 4.13); 289 (log ϵ 3.99); 306 (log ϵ 3.78); 320 (log ϵ 3.66) (EtOH).

4-Epimer: Fumiquinazoline G

[169869-87-8]
 $C_{21}H_{18}N_4O_2$ 358.399
Prod. by a marine-derived *Aspergillus fumigatus*. Cytotoxic agent. Pale yellow powder. Mp 119-121°. $[\alpha]_D^{25}$ -462.8 (c, 0.6 in $CHCl_3$). λ_{max} 208 (log ϵ 4.61); 220 (log ϵ 4.67); 273 (log ϵ 4.14); 278 (log ϵ 4.13); 288 (log ϵ 4.01); 307 (log ϵ 3.66); 323 (log ϵ 3.49) (EtOH).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1995, 2345-2353 (*isol, uv, ir, pmr, cmr*)

He, F. *et al.*, *Synlett*, 1997, 483-484 (*synth*)

Wang, H. *et al.*, *J.O.C.*, 2000, **65**, 1022-1030 (*synth*)


Snider, B.B. *et al.*, *Tetrahedron*, 2001, **57**, 3301-3307 (*synth*)

Liu, J.-F. *et al.*, *J.O.C.*, 2005, **70**, 6339-5345 (*synth*)

Fumiquinazoline H

F-189

[278184-55-7]



Absolute Configuration

$C_{27}H_{27}N_5O_4$ 485.541

Isol. from an *Acremonium* sp. obt. from the tunicate *Ecteinascidia turbinata*. Pale yellow solid. Mp 144-147°. $[\alpha]_D^{25}$ -59 (c, 0.001 in $CHCl_3$). λ_{max} 224 (ϵ 27700); 230 (sh) (ϵ 25800); 255 (ϵ 11500); 265 (ϵ 9700); 276 (ϵ 7600); 303 (ϵ 3000) (MeOH).


Belofsky, G.N. *et al.*, *Chem. Eur. J.*, 2000, **6**, 1355-1360 (*isol, uv, pmr, cmr*)

Snider, B.B. *et al.*, *J.O.C.*, 2003, **68**, 545-563 (*synth*)

Fumitremorgin C

F-190

SM-Q
[118974-02-0]



Absolute Configuration

$C_{22}H_{25}N_3O_3$ 379.458

Struct. revised in 1988. Prod. by *Aspergillus fumigatus* and *Neosartorya*

fischeri. Tremorgenic mycotoxin. Reverses of multidrug resistance in cells transfected with the breast cancer resistance protein. Cryst. (EtOAc). Mp 259.5-260.5° (synthetic). $[\alpha]_D^{28}$ -13 (c, 0.53 in MeOH). An early Mp of 125-130° was reported for natural material but this was prob. erroneous or for an impure sample. λ_{max} 224 ; 272 ; 294 (MeOH) (Berdy). λ_{max} 225 ; 273 ; 299 (EtOH) (Berdy).

Demethoxy: Demethoxyfumitremorgin C

[111768-16-2]
[106211-91-0 (12 β -isomer)]
 $C_{21}H_{23}N_3O_2$ 349.432
Prod. by *Aspergillus fumigatus*. Cell cycle progression Inhibitor, mycotoxin. Pale yellow needles. Sol. MeOH. Mp 210-212°. $[\alpha]_D^{30}$ +8 (c, 0.2 in $CHCl_3$). λ_{max} 226 (ϵ 24950); 282 (ϵ 9000); 291 (sh) (ϵ 7450) (MeOH).

12,13-Didehydro, 12 ξ ,13 ξ -epoxide: Epoxyfumitremorgin C

$C_{22}H_{25}N_3O_4$ 395.457
From *Aspergillus fumigatus*. Tremorgenic mycotoxin. Yellow cryst. Mp 197-198°. $[\alpha]_D^{25}$ +18.4.

12-Epimer: [119066-64-7]

Synthetic. Pale yellow prisms (EtOH). Mp 240-247.5° dec. $[\alpha]_D^{25}$ +239 (c, 0.11 in MeOH).

[119066-69-2, 119066-68-1, 118974-03-1, 119066-67-0]

Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 826-830 (*isol*)

Steyn, P.S. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1985, **48**, 60 (*uv, pmr, cmr*)

Plate, R. *et al.*, *J.O.C.*, 1987, **52**, 560-564 (*synth, uv, pmr, ms*)

O'Malley, G.J. *et al.*, *Tet. Lett.*, 1987, **28**, 1131-1154 (*demethoxy, synth*)

Hermkens, P.H.H. *et al.*, *Tetrahedron*, 1988, **44**, 1991-2000 (*synth, uv, pmr, ms*)

Bailey, P.D. *et al.*, *Tet. Lett.*, 1989, **30**, 6421-6422; 2001, **42**, 113-115 (*demethoxy, synth*)

Hino, T. *et al.*, *Tetrahedron*, 1989, **45**, 1941-1944 (*synth, uv, ir, pmr, cmr, ms*)

Abraham, W.R. *et al.*, *Phytochemistry*, 1990, **29**, 1025-1026 (*12,13-*

Dihydroxyfumitremorgin C)
Cui, C.-B. *et al.*, *J. Antibiot.*, 1996, **49**, 527-534; 534-540 (*Demethoxyfumitremorgin C, abs config*)

Hino, T. *et al.*, *Heterocycles*, 1997, **46**, 673-704 (*rev, synth*)

Rabindran, S.K. *et al.*, *Cancer Res.*, 1998, **58**, 5850-5858 (*activity*)

Handbook of Secondary Fungal Metabolites, (ed. Cole, R.J. *et al.*), Academic Press, 2003, **1**, 226


Afiyatullo, S.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 615-617 (*12,13-Dihydroxyfumitremorgin C*)

Wang, F.-W. *et al.*, *Acta Cryst. E*, 2006, **62**, o1375-o1377 (*cryst struct*)

Fumschleicherine

F-191

6-[16-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methyl]-6,7-dihydro-6-hydroxy-8H-1,3-dioxolo[4,5-e]isoindol-8-one, 9CI
[77292-20-7]



$C_{21}H_{22}N_2O_6$ 398.415

Alkaloid from *Fumaria schleicheri* and *Fumaria schrammii* (Papaveraceae). Mp 224-226°. $[\alpha]_D^{20}$ +5.88 (DMSO).

Kiryakov, Kh. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1980, **33**, 1377; *CA*, **94**, 188644f (*isol, struct*)

Fumvaine

F-192

$C_{20}H_{19}NO_6$ 369.373


Struct. unknown. Alkaloid from *Fumaria vaillantii* (Papaveraceae). Cryst. (Me₂CO). Mp 180.5-181.5°. $[\alpha]_D^{25}$ -44. Nonphenolic, conts. 2 OMe groups. *Hydrochloride*: Mp 212°.

Platonova, T.F. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 173-180; *CA*, **50**, 13960

Funebradiol

F-193

3-[2,5-Bis(hydroxymethyl)-1H-pyrrol-1-yl]-4,5-dimethyl-2(5H)-furanone, 9CI
[133086-87-0]



$C_{12}H_{15}NO_4$ 237.255

(R)-form


Alkaloid from the flowers of *Quararibea funebris* (Bombacaceae). Gum. $[\alpha]_D^{25}$ +8.6 (c, 0.3 in MeOH). λ_{max} 219 (ϵ 2190); 266 (ϵ 933) (MeOH) (Derep).

Zennie, T.M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1611 (*isol, pmr, cmr, struct*)

Funebral

F-194

[105708-56-3]



$C_{12}H_{15}NO_4$ 237.255

Alkaloid from the flowers of *Quararibea funebris* (Bombacaceae). $[\alpha]_D^{22}$ -19 (c, 0.05 in MeOH).

Zennie, T.M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 695-698 (isol, uv, ir, pmr, cmr, ms, cd, struct)


Yu, S.-X. *et al.*, *Tet. Lett.*, 1995, **36**, 6205-6208 (synth)

Dong, Y. *et al.*, *J.O.C.*, 1999, **64**, 2657-2666 (synth, conformn)

Tamura, O. *et al.*, *J.O.C.*, 2004, **69**, 1475-1480 (synth)

Funebrine

[90605-98-4]



$C_{18}H_{24}N_2O_5$ 348.398

Alkaloid from the flowers of *Quararibea funebris* (Bombacaceae). Cryst. (cyclohexane/ $CHCl_3$). Mp 232-233°. $[\alpha]_D^{22.5}$ -215 (c, 0.01 in DMSO). λ_{max} 265 (sh) (ϵ 20000); 293 (ϵ 30000) (DMSO) (Derep).

Raffauf, R.F. *et al.*, *J.O.C.*, 1984, **49**, 2714 (isol, uv, ir, pmr, ms, cryst struct)

Zennie, T.M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 695 (pmr)


Dong, Y. *et al.*, *J.O.C.*, 1999, **64**, 2657-2666 (synth, conformn)

Tamura, O. *et al.*, *J.O.C.*, 2004, **69**, 1475-1480 (synth)

Fungerin*Visoltricin*

[185681-81-6]

[139874-44-5]



$C_{13}H_{18}N_2O_2$ 234.297

Alkaloid from *Fusarium tricinctum*, another *Fusarium* sp., *Metarhizium* sp. FKI-1079 and the marine *Microascus* sp. K14. Cytotoxic agent. Shows anticholinesterase and antifungal activities. Needles (hexane/EtOAc). Mp 93-95°. λ_{max} 300 (ϵ 28000) (MeOH).

N¹-Me: N-Methylvisoltricin

[156524-19-5]

[142750-50-3]

 $C_{14}H_{21}N_2O_2^{\oplus}$ 249.332

Quaternary alkaloid prod. by *Fusarium* sp. Acetylcholinesterase inhibitor. Powder (as iodide).

5''-Hydroxy: Hydroxyfungerin A $C_{13}H_{18}N_2O_3$ 250.297

Prod. by *Metarhizium* sp. FKI-1079. Pale yellow powder. λ_{max} 305 (ϵ 27000)

(MeOH).

5''-Hydroxy; N-de-Me, 3-Me (3H-form):**Hydroxyfungerin B** $C_{13}H_{18}N_2O_3$ 250.297Prod. by *Metarhizium* sp. FKI-1079.

Pale yellow powder. λ_{max} 305 (ϵ 27200) (MeOH).

Marta, M. *et al.*, *Acta Med. Rom.*, 1990, **28**, 343-346 (N-Me)

Visconti, A. *et al.*, *J. Agric. Food Chem.*, 1994, **42**, 195-199 (*Visoltricin*)

Kato, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1996, **60**, 2081-2083 (isol, uv, pmr, cmr, N-15 nmr)

Benhida, R. *et al.*, *Tet. Lett.*, 1998, **39**, 5963-5964 (synth)

Martin, G.E. *et al.*, *J. Nat. Prod.*, 2000, **63**, 543-585 (N-15 nmr)


Rieder, J.M. *et al.*, *Tet. Lett.*, 2002, **43**, 2375-2376 (synth, struct)

Koizumi, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 415-420 (activity)

Uchida, R. *et al.*, *J. Antibiot.*, 2005, **58**, 804-809 (*Hydroxyfungerins*)

Funiculosin†*SL 3238. Antibiotic SL 3238*

[11055-06-4]



$C_{27}H_{41}NO_7$ 491.623

Identity with SL 3238 is probable. Related to Oxysporidinone, O-226 and Sambutoxin, S-48. Prod. by *Penicillium funiculosum* and *Penicillium purpurogenum*. Antiviral and antifungal agent. SL 3238 reported as active against gram-positive and -negative bacteria. Fine needles. Sol. MeOH, Me_2CO , butanol, EtOH; fairly sol. EtOAc, $CHCl_3$, Et_2O , C_6H_6 ; poorly sol. hexane, H_2O . Mp 165-166° (160-161°). $[\alpha]_D^{20}$ -102 (c, 0.47 in $CHCl_3$). Opt. rotn. refers to SL 3238. λ_{max} 290 (ϵ 5500) (MeOH) (Derep). λ_{max} 290 (ϵ 5500) (MeOH) (Berdy).

▶ **LD₅₀** (mus, ipr) 4 mg/kg. LT5270000

2''-Deoxy: Antibiotic AF 110. AF 110 [146671-30-9]

 $C_{27}H_{41}NO_6$ 475.624

Prod. by *Penicillium* sp. F-4209. Antifungal and antitumour agent. Yellow powder. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D$ -184 (MeOH). Stereochem. not determined. λ_{max} 213 (ϵ 34000); 290 (ϵ 6700) (MeOH). λ_{max} 213 (ϵ 34000); 290 (ϵ 6700) (MeOH) (Berdy).

Ando, K. *et al.*, *J. Antibiot.*, 1969, **22**, 189-194; 1978, **31**, 533-538 (isol, abs config, cryst struct)

Ger. Pat., 1970, 2005976; *CA*, **73**, 108258m (SL 3238)

Japan. Pat., 1992, 92 321 684; *CA*, **118**, 167161y (AF 110)


Williams, D.R. *et al.*, *Tet. Lett.*, 2000, **41**, 9397-9401 (synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FPD000

Funtudienine

3,5-Conadien-7-one

F-198



Absolute Configuration

$C_{22}H_{31}NO$ 325.493

Alkaloid from bark of *Funtumia latifolia* (Apocynaceae). Amorph. $[\alpha]_D$ -247 (c, 2.5 in $CHCl_3$).

Oxime:

Cryst. (MeOH). Mp 190°. $[\alpha]_D$ -228 (c, 1.0 in $CHCl_3$).


Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 2169 (isol, ir, uv, pmr, ms, struct, synth)

Crabbé, P. *et al.*, *Chem. Ind. (London)*, 1966, 158 (ord)

3-Furanmethanol, 9CI

F-199

3-Furancarbinol. 3-Furfuryl alcohol. 3-(Hydroxymethyl)furan [4412-91-3]



$C_5H_6O_2$ 98.101

Constit. of *Aloe arborescens*, *Eucommia ulmoides*, and *Stellaria aquatica*. Liq. d_4^{20} 1.14. Bp_{17} 79-80°. n_D^{20} 1.4842.

Phenylurethane:

Fine needles. Mp 102.5-105.6°.

O-β-D-Glucopyranoside: [86425-28-7] $C_{11}H_{16}O_7$ 260.243

Isol. from seeds of *Vigna angularis* (azuki bean). Hygroscopic powder. $[\alpha]_D^{26}$ +40 (c, 1.1 in MeOH).

Benzoyl: $C_{12}H_{10}O_3$ 202.209 $Bp_{0.75}$ 95-97°.

O-(2-Pyrrolicarbonyl): 3-Furfuryl 2-pyrrolicarboxylate [119767-00-9]

 $C_{10}H_9NO_3$ 191.186

Alkaloid from the roots of *Pseudostellaria heterophylla*. Oil.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 581D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 18B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1453B (ir)

Sherman, E. *et al.*, *J.A.C.S.*, 1950, **72**, 2195 (synth)

Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 664 (glucoside)

Jurášek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1985, **50**, 2077 (synth)

Reinecke, M.G. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1236; 1989, **52**, 375 (pyrrole-2-carboxylate)


Marstokk, K.M. *et al.*, *Acta Chem. Scand.*, 1993, **47**, 849 (microwave, struct)

Bures, E. *et al.*, *J.O.C.*, 1997, **62**, 8741-8749 (synth, ir, pmr, cmr)

Mace, L.H. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1020-1031 (synth)

1-(2-Furanylmethyl)-1H-pyrrole, 9CI **F-200**

N-(2-Furfuryl)pyrrole. FEMA 3284 [1438-94-4]




C₉H₉NO 147.176

Present in sandalwood oil (*Santalum album*). Flavour ingredient. Bp₁ 77-79°. n_D²¹ 1.5317.

- ▶ LD₅₀ (mus, orl) 380 mg/kg. UX9631000
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 42B (nmr)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1466A (ir)
 Gianturco, M.A. et al., *Tetrahedron*, 1964, 20, 1763; 2951 (isol, ir, pmr, synth)
 Fenaroli's *Handbook of Flavor Ingredients*, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 2, 290
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1149-1150

2-[(2-Furanylmethyl)thio]-6-methylpyrazine, 9CI **F-201**

6-Methyl-2-pyrazinyl 2-furfuryl sulfide. FEMA 3189 [59021-05-5] [59303-07-0, 59303-08-1, 59303-09-2]




C₁₀H₁₀N₂OS 206.268

The phys. props. given are for a mixt. of regioisomers (3-, 5- and 6-methylpyrazines), which are the coml. flavouring material FEMA 3189. Present in aroma of pumpkin seed oil. Present in pumpkin seed oil. Flavouring ingredient. Liq. with roasted coffee-like taste. d₄²⁰ 1.22. Bp₁₀ 153-156°. n_D²⁰ 1.5970. Synthesised as a mixt. of regioisomers (2,3-, 2,5- and 2,6-) to which the phys. props. refer. The FEMA no. refers to this mixt.

U.S. Pat., 1976, 3 989 713; *CA*, 86, 43556d (synth, use)
 Buchbauer, G. et al., *Ernaehrung (Vienna)*, 1998, 22, 246-249; *CA*, 129, 135437a (isol)
 Fenaroli's *Handbook of Flavor Ingredients*, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 1122-1123 (use, props)

5-(3-Furanyl)octahydro-8-methylindolizine, 9CI **F-202**

5-(3-Furanyl)-8-methylindolizidine [60161-29-7]




C₁₃H₁₉NO 205.299

Closely related to the *Nuphar* alkaloids. Trace alkaloid from the scent glands of the Canadian beaver *Castor fiber*. Oil.

Maurer, B. et al., *Helv. Chim. Acta*, 1976, 59, 1169 (isol, ms, struct)
 Tufariello, J.F. et al., *Chem. Comm.*, 1987, 1138 (synth)

Furoclausine A **F-203**

8-Hydroxy-10H-furo[3,2-a]carbazole-4-carboxaldehyde, 9CI [192706-50-6]




C₁₅H₉NO₃ 251.241

Alkaloid from the root bark of *Clausena excavata* (Rutaceae). Yellow oil. λ_{max} 221 ; 239 ; 288 (sh) ; 301 ; 345 (MeOH).

Wu, T.-S. et al., *Heterocycles*, 1997, 45, 969-973 (isol, uv, ir, pmr, cmr)

Furoclausine B **F-204**

[192706-53-9]




C₁₈H₁₇NO₄ 311.337

Alkaloid from the root bark of *Clausena excavata* (Rutaceae). Yellow oil. [α]_D -32.7 (c, 0.02 in MeOH). λ_{max} 202 ; 218 (sh) ; 231 (sh) ; 252 ; 275 (sh) ; 286 (sh) ; 301 ; 339 (MeOH).

Wu, T.-S. et al., *Heterocycles*, 1997, 45, 969-973 (isol, uv, ir, pmr)

Furoeriaustralasine **F-205**

[152406-30-9]



Relative configuration


C₁₉H₂₁NO₃ 311.38

Minor alkaloid from aerial parts of *Eriostemon australasius* (Rutaceae). Oil. [α]_D -117 (c, 1.0 in CHCl₃).

da Cunha, E.V.L. et al., *Aust. J. Chem.*, 1993, 46, 1507 (isol, uv, ir, pmr, cmr, ms, struct)

Furofoline **F-206**

5-Hydroxy-11-methylfuro[2,3-c]acridin-6(11H)-one, 9CI. Furofoline I. Furoacridone [62541-22-4]



R = H

C₁₆H₁₁NO₃ 265.268

Alkaloid from the roots of *Ruta graveolens* (rue) and the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow plates (Me₂CO). Mp 245-246°.

Reisch, J. et al., *Phytochemistry*, 1977, 16, 151 (isol)

Reisch, J. et al., *Annalen*, 1981, 85 (synth)
 Wu, T.-S. et al., *J.C.S. Perkin 1*, 1983, 1681 (isol, uv, ir, pmr, ms, struct)

Furofoline II **F-207**

5-Hydroxy-2-(1-hydroxy-1-methylethyl)-11-methylfuro[2,3-c]acridin-6(11H)-one, 9CI

[82644-80-2]

As Furofoline, F-206 with

R = -C(CH₃)₂OH

C₁₉H₁₇NO₄ 323.348


Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow needles (Me₂CO). Mp 213-215°.

Wu, T.-S. et al., *Heterocycles*, 1982, 19, 1227 (isol, uv, ir, pmr, ms, struct)

Wu, T.-S. et al., *J.C.S. Perkin 1*, 1983, 1681 (isol, uv, ir, pmr, ms, struct)

Furomegistine II **F-208**

[352280-28-5]




C₁₄H₁₃NO₇ 307.259

Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow solid. Racemic. λ_{max} 278 (log ε 2.66); 307 (sh) (MeOH).

Fokialakis, N. et al., *Phytochemistry*, 2001, 57, 593-596

Europaradine **F-209**

[161161-72-4]




C₁₇H₁₃NO₅ 311.293

Alkaloid from roots of *Citrus paradisi* (grapefruit) (Rutaceae). Yellow oil.

Takemura, Y. et al., *Heterocycles*, 1995, **41**, 187
(isol, uv, ir, pmr, cmr, ms, struct)

Furo[2,3-*b*]quinoline-4,5-diol F-210
4,5-Dihydroxyfuro[2,3-*b*]quinoline



C₁₁H₇NO₃ 201.181


Di-Me ether: 4,5-Dimethoxyfuro[2,3-*b*]quinoline. 5-Methoxydictamnine [119708-35-9]

C₁₃H₁₁NO₃ 229.235

Alkaloid from the roots of *Ruta chalepensis* var. *latifolia* (Rutaceae).

Ulubelen, A. et al., *Planta Med.*, 1988, **54**, 551
(isol, uv, ir, pmr, ms, struct)

Furo[2,3-*b*]quinoline-4,6-diol F-211
4,6-Dihydroxyfuro[2,3-*b*]quinoline. 6-Hydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₃ 201.181

OH-form

Di-Me ether: 4,6-Dimethoxyfuro[2,3-*b*]quinoline, 9CI. **Pteleine**. 6-Methoxydictamnine [2221-41-2]

C₁₃H₁₁NO₃ 229.235

Alkaloid from *Platydema campanulata* and *Ptelea trifoliata*; tentatively identified in cells of *Ruta graveolens* (rue) grown in continuous light in liq. medium (Rutaceae). Shows antibacterial activity. Cubes (C₆H₆/petrol). Mp 134-135°. λ_{max} 237 (log ε 4.66); 250 (log ε 4.56); 295 (log ε 3.91); 306 (log ε 3.99); 334 (log ε 3.68); 348 (log ε 3.67) (no solvent reported).

▶ LV2708180

Di-Me ether, picrate:

Yellow needles (EtOH). Mp 195-196° dec.

Di-Me ether, picrolonate: Mp 200-202° dec.

NH-form

N-Me: 6-Hydroxy-9-methylfuro[2,3-*b*]quinolin-4(9H)-one, 9CI. **Dictangustine A** [221457-19-8]

C₁₂H₉NO₃ 215.208

Alkaloid from root bark of *Dictamnus angustifolius*. Yellowish powder (Me₂CO). Mp > 280°. λ_{max} 242; 255; 263; 348; 364 (MeOH).

6-Me ether, N-Me: 6-Methoxy-9-methylfuro[2,3-*b*]quinolin-4(9H)-one, 9CI. **Isopteleine**. 6-Methoxyisodictamnine [2181-84-2]

C₁₃H₁₁NO₃ 229.235

Alkaloid from root and bark of *Platydema campanulata* and *Dictamnus*

caucasicus (Rutaceae). Cryst. (EtOH/Me₂CO). Mp 214-216° (207-209°). Turns pink on standing. λ_{max} 249 (log ε 6.1); 261 (log ε 4.91); 284 (sh) (log ε 4.81); 296 (log ε 4.99); 307 (log ε 5.04); 333 (log ε 4.74); 350 (log ε 4.68) (95%EtOH).

Werny, F. et al., *Tetrahedron*, 1963, **19**, 1293-1305 (6-Methoxydictamnine, isol, uv, ir, struct, synth)

Pai, B.R. et al., *Indian J. Chem.*, 1964, **2**, 491-492 (6-Methoxydictamnine, synth, uv)

Frolova, V.I. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1964, **34**, 3542-3546 (Pteleine, isol, struct)

Kikvidze, I.M. et al., *Khim. Prir. Soedin.*, 1971, **7**, 675; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 659-660 (Isopteleine)

Steck, W. et al., *Phytochemistry*, 1971, **10**, 191-194 (6-Methoxydictamnine, isol, uv, pmr)


Sekiba, T. et al., *Bull. Chem. Soc. Jpn.*, 1973, **46**, 577-580 (Pteleine, synth, uv)

Narasimhan, N.S. et al., *Tetrahedron*, 1974, **30**, 4153-4157 (Pteleine, synth, uv, ir, pmr)

Wu, T.-S. et al., *Phytochemistry*, 1999, **50**, 509-512 (Dictangustine A)

El Sayed, K. et al., *J. Nat. Prod.*, 2000, **63**, 995-997 (Pteleine, activity)

Furo[2,3-*b*]quinoline-4,7-diol F-212
4,7-Dihydroxyfuro[2,3-*b*]quinoline. 7-Hydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₃ 201.181

OH-form

4-Me ether: 4-Methoxyfuro[2,3-*b*]quinolin-7-ol, 9CI. **Confusameline**. 7-O-Demethylevolitrine. 7-Hydroxydictamnine [20643-71-4]

C₁₂H₉NO₃ 215.208

Alkaloid from the leaves of *Evodia elleryana* and *Melicope confusa*, and from the aerial parts of *Melicope lasioneura* (Rutaceae). Pale yellow needles (MeOH). Mp 240-242°.

4-Me ether, 7-Ac:

Needles (EtOH). Mp 156-157°.

Di-Me ether: 4,7-Dimethoxyfuro[2,3-*b*]quinoline, 9CI. **Evolitrine**. 7-Methoxydictamnine [523-66-0]

C₁₃H₁₁NO₃ 229.235

Alkaloid from *Evodia littoralis*, *Evodia lunu-ankenda*, *Phebalium nudum*, *Ruta montana*, *Melicope lasioneura*, *Acronychia laurifolia* and others. Cytotoxic agent. Rosettes of needles (petrol). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 114-115°. λ_{max} 245 (ε 100000); 307 (ε 14150); 318 (ε 13500); 332 (ε 11720) (MeOH) (Berdy). λ_{max} 244 (ε 69700); 308 (ε 10250); 320 (ε 9800); 333 (ε 7980) (EtOH) (Berdy).

▶ LV2708190

Di-Me ether, hydrochloride:

Rods (EtOH/Et₂O). Mp 150-151° dec.

Di-Me ether, picrate:

Pale yellow needles (EtOH). Mp 201-202°.

7-O-(3-Methyl-2-butenyl), 4-Me ether: 4-Methoxy-7-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline, 9CI. 4-Methoxy-7-prenyloxyfuro[2,3-*b*]quinoline.

Melineurine

[85802-28-4]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the aerial parts of *Melicope lasioneura* (Rutaceae). Small prisms (Et₂O). Mp 98-99°. λ_{max} 247 (ε 70800); 290 (sh) (ε 6610); 309 (ε 10500); 321 (ε 9550); 333 (sh) (ε 8320) (EtOH) (Derep).

7-O-(2-ξ-Hydroxy-3-methyl-3-butenyl), 4-Me ether: **Confusadine** [479415-36-6]

C₁₇H₁₇NO₄ 299.326

Alkaloid from the leaves of *Melicope semecarpifolia*. Anti-platelet aggregation agent. Needles (MeOH). Mp 144-146°. [α]_D²⁷ -22.4 (c, 0.11 in MeOH). λ_{max} 246 (log ε 4.83); 309 (log ε 4.03); 320 (log ε 4.05); 333 (sh) (log ε 3.95) (EtOH).

7-O-(2,3-Dihydroxy-3-methylbutyl), 4-Me ether: **Evellerine**

[20547-76-6]

C₁₇H₁₉NO₅ 317.341

Alkaloid from leaves of *Evodia elleryana* (Rutaceae). Prisms (Me₂CO). Mp 149-150°. [α]_D²⁰ +21 (c, 0.30 in CHCl₃). λ_{max} 247 (ε 70800); 290 (sh) (ε 6610); 309 (ε 10500); 321 (ε 9550); 333 (sh) (ε 8320) (EtOH) (Derep).

7-O-(3-Chloro-2-hydroxy-3-methylbutyl), 4-Me ether: 7-(3-Chloro-2-hydroxy-3-methylbutoxy)-4-methoxyfuro[2,3-*b*]quinoline

C₁₇H₁₈ClNO₄ 335.786

Constit. of the leaves of *Melicope bonwickii*. Pale yellow needles (MeOH). Mp 171-173°. λ_{max} 245 (CHCl₃).

7-O-(3,7-Dimethyl-5-oxo-2E,6-octadienyl), 4-Me ether: [1050284-42-8]

C₂₂H₂₃NO₄ 365.428

Alkaloid from the leaves of *Boninia glabra*. Amorph. yellow solid.

7-O-(3,7-Dimethyl-5-oxo-3E,6-octadienyl), 4-Me ether: [1050284-41-7]

C₂₂H₂₃NO₄ 365.428

Alkaloid from the leaves of *Boninia glabra*. Pale yellow cryst. Mp 150-153°.

7-O-(3,7-Dimethyl-5-oxo-3Z,6-octadienyl), 4-Me ether: [1050284-40-6]

C₂₂H₂₃NO₄ 365.428

Alkaloid from the leaves of *Boninia glabra*. Amorph. yellow solid.

NH-form

7-Me ether, N-Et: 9-Ethyl-7-methoxyfuro[2,3-*b*]quinolin-4(9H)-one, 9CI.

Taifine

[79808-96-1]

C₁₄H₁₃NO₃ 243.262

Alkaloid from *Ruta chalepensis* (Rutaceae). Needles (petrol). Mp 110°. Doubtful structural assignment; spectral and physical props. of authentic synthetic material differ from those reported for natural Taifine.

7-Me ether, N-(3-methyl-2-butenyl): 7-Methoxy-9-(3-methyl-2-butenyl)furo[2,3-*b*]quinolin-4(9H)-one, 9CI. 7-

*Methoxy-9-prenylfuro[2,3-*b*]quinolin-4(9H)-one. Acrophylline*

[18904-40-0]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the leaves and bark of *Acronychia haplophylla* (Rutaceae).

Cryst. (MeOH). Mp 119-120°.

7-*Me ether*, N-(3-hydroxy-3-methylbutyl)-9-(3-Hydroxy-3-methylbutyl)-7-methoxyfuro[2,3-*b*]quinolin-4(9H)-one, 9CI. **Acrophyllidine**

[18813-64-4]

C₁₇H₁₉NO₄ 301.341

Alkaloid from the leaves of *Acronychia haplophylla* (Rutaceae). Also obt. by acid-catalysed hydration of Acrophylline. Needles (MeOH). Mp 176-177°.

Cooke, R.G. *et al.*, *Aust. J. Chem.*, 1954, **7**, 273; 1958, **11**, 225 (*Evolitrine*, *isol*, *uv*, *struct*, *synth*)

Sato, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1958, **31**, 161 (*Evolitrine*, *synth*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1968, **21**, 1897 (*Confusameline*, *Evellerine*)

Lahey, F.N. *et al.*, *Aust. J. Chem.*, 1969, **22**, 447 (*Acrophylline*, *Acrophyllidine*)

Prabhakar, S. *et al.*, *Indian J. Chem.*, 1970, **8**, 857 (*Acrophylline*, *synth*, *uv*, *ir*, *pmr*)

Yang, T.H. *et al.*, *Yakugaku Zasshi*, 1971, **91**, 782; *CA*, **75**, 95382m (4-*Me ether*, *isol*, *struct*)

Narasimhan, N.S. *et al.*, *Tetrahedron*, 1974, **30**, 4153 (*Evolitrine*, *synth*, *uv*, *ir*, *pmr*)

El-Tawil, B.A.H. *et al.*, *Z. Naturforsch.*, **B**, 1981, **36**, 1169 (*Taifine*)

Tillequin, F. *et al.*, *J. Nat. Prod.*, 1982, **45**, 486; 1983, **46**, 132 (*Evellerine*, *Evolitrine*, *Melineurine*)

Kang, S.S. *et al.*, *Arch. Pharmacol. Res.*, 1986, **9**, 11; *CA*, **105**, 39390p (4-*Me ether*, *isol*)

Kuo, S.C. *et al.*, *J. Nat. Prod.*, 1986, **49**, 48 (*Taifine*)

Rajamanickam, P. *et al.*, *Indian J. Chem., Sect. B*, 1987, **26**, 910 (*Evolitrine*, *synth*)

Huang, A.-C. *et al.*, *J. Nat. Prod.*, 1995, **58**, 117 (*Acrophylline*, *Acrophyllidine*, *synth*, *uv*, *ir*, *pmr*, *cmr*, *biochem*)

Cui, B. *et al.*, *Phytochemistry*, 1999, **52**, 95-98 (*Evolitrine*, *activity*)

Chen, J.-J. *et al.*, *Planta Med.*, 2002, **68**, 790-793 (*Confusadine*)


Lal, B. *et al.*, *ARKIVOC*, 2005, **ii**, 77-97 (*Evolitrine*, *isol*, *synth*)

Komala, I. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 355-360 (3-chloro-2-hydroxy-3-methylbutyl 4-*Me ether*)

Inada, A. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 727-729 (*Boninia glabra constitis*)

Furo[2,3-*b*]quinoline-4,8-diol F-213

4,8-Dihydroxyfuro[2,3-*b*]quinoline. 8-Hydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₃ 201.181

OH-form

4-*Me ether*: 4-Methoxyfuro[2,3-*b*]quinolin-8-ol, 9CI. 8-Hydroxy-4-methoxyfuro[2,3-*b*]quinoline. 8-Hydroxydictamnine. **Robustine**† [2255-50-7]

C₁₂H₉NO₃ 215.208

Alkaloid from *Thamnosma montana*, *Dictamnus caucasicus*, *Haplophyllum robustum* and *Zanthoxylum* spp. (Rutaceae). Cryst. (EtOH or C₆H₆). Mp 147-148° (142-143°).

4-*Me ether*, hydrochloride:

Cryst. (EtOH). Mp 199-200°.

4-*Me ether*, picrate:

Cryst. (EtOH). Mp 179-180°.

Di-*Me ether*: 4,8-Dimethoxyfuro[2,3-*b*]quinoline. 8-Methoxydictamnine. γ -**Fagarine**

[524-15-2]

C₁₃H₁₁NO₃ 229.235

Alkaloid from *Fagara coco*, *Aegle marmelos* (bael fruit) and *Haplophyllum* spp. (Rutaceae). Active against gram-positive bacteria and fungi. Pale yellow prisms (petrol). Poorly sol. hexane. Mp 142°. λ_{\max} 203; 243; 310; 325 (MeOH) (Berdy). λ_{\max} 244 (ϵ 56210); 290 (ϵ 5250); 311 (ϵ 6610); 327 (ϵ 5750); 340 (ϵ 4900) (EtOH) (Berdy).

▶ LV2708200

Di-*Me ether*, picrate:

Yellow crystal. (EtOH). Mp 177°.

O⁴-*Et*, O⁸-*Me*: 4-Ethoxy-8-methoxyfuro[2,3-*b*]quinoline. O-Ethylnor- γ -fagarine

[105988-99-6]

C₁₄H₁₃NO₃ 243.262

Alkaloid from roots of *Dictamnus dasycarpus* (Rutaceae).

8-O-(3-Methyl-2-butenyl), O⁴-*Me*: 4-Methoxy-8-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline, 9CI. 4-Methoxy-8-prenylxyfuro[2,3-*b*]quinoline. **Haplophydine**. **Haplofidine**. **Haplophidine**

[55727-61-2]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the above-ground parts of *Haplophyllum perforatum* (Rutaceae). Cryst. (EtOH). Mp 111-112°.

NH-form

8-*Me ether*, N-*Me*: 8-Methoxy-9-methylfuro[2,3-*b*]quinolin-4(9H)-one. **Iso- γ -fagarine**

C₁₃H₁₁NO₃ 229.235

Alkaloid from root bark of *Dictamnus angustifolius*. Needles (Me₂CO). Mp 160-161°. λ_{\max} 246; 255; 301; 322; 336; 350 (MeOH).

Fakhrutdinova, I.M. *et al.*, *Chem. Nat. Compd.* (Engl. Transl.), 1965, **1**, 83 (*isol*, *struct*)

Faizutdinova, Z.S. *et al.*, *Chem. Nat. Compd.* (Engl. Transl.), 1967, **3**, 218 (*ms*)

Collins, J.F. *et al.*, *J.C.S. Perkin 1*, 1973, 94-97 (*synth*)

Abdullaeva, K.A. *et al.*, *Chem. Nat. Compd.* (Engl. Transl.), 1974, **10**, 713 (*Haplophydine*)

Narasimhan, N.S. *et al.*, *Tetrahedron*, 1974, **30**, 4153-4157 (*synth*, *uv*, *ir*, *pmr*)

Chang, P.T.O. *et al.*, *J. Nat. Prod.*, 1976, **39**, 134 (*isol*, *ir*, *uv*, *ms*, *pmr*)

Wolters, B. *et al.*, *Planta Med.*, 1981, **43**, 166-174 (γ -*Fagarine*, *activity*)


Lin, T.P. *et al.*, *CA*, 1987, **106**, 30033n (4-Ethoxy-8-methoxyfuroquinoline)

Wu, T.S. *et al.*, *Phytochemistry*, 1999, **50**, 509-512 (*Iso- γ -fagarine*)

Chlouchi, A. *et al.*, *Biochem. Syst. Ecol.*, 2006, **34**, 71-74 (*Robustine*, *pmr*, *cmr*)

Furo[2,3-*b*]quinoline-4,5,6,7,8-pentol F-214

4,5,6,7,8-Pentahydroxyfuro[2,3-*b*]quinoline. 5,6,7,8-Tetrahydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₆ 249.179

OH-form

6,7-Methylene, 4,5,8-tri-*Me ether*: 4,9,10-Trimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline. 4,5,8-Tri-methoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline. 5,8-Dimethoxymaculine. 5-Methoxyflindersiamine

C₁₅H₁₃NO₆ 303.271

Alkaloid from the wood of *Vepris punctata*. Viscous oil. λ_{\max} 242 (ϵ 20800); 258 (ϵ 44300); 330 (ϵ 12400); 350 (ϵ 6400) (MeOH).

Penta-*Me ether*: 4,5,6,7,8-Pentamethoxyfuro[2,3-*b*]quinoline. 5-Methoxyhalfordimine. 6-Methoxyacronycidine. 5,6,7,8-Tetramethoxydictamnine


C₁₆H₁₇NO₆ 319.313

Alkaloid from the wood of *Vepris punctata*. Viscous oil. λ_{\max} 244 (ϵ 24500); 262 (ϵ 46000); 327 (ϵ 9100); 352 (ϵ 4200) (MeOH).

Chaturvedula, V.S.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 532-534 (*isol*, *uv*, *pmr*, *cmr*)

Furo[2,3-*b*]quinoline-4,5,6,7-tetrol F-215

4,5,6,7-Tetrahydroxyfuro[2,3-*b*]quinoline. 5,6,7-Trihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₅ 233.18

OH-form

Tetra-*Me ether*: 4,5,6,7-Tetramethoxyfuro[2,3-*b*]quinoline. 5,6,7-Trimethoxydictamnine

[928849-55-2]

C₁₅H₁₅NO₅ 289.287

Alkaloid from the stem bark of a

Brombya sp.

6,7-Methylene, 4,5-di-*Me ether*: 9,10-Dimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline. 4,5-Dimethoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline. 5-Methoxymaculine

C₁₄H₁₁NO₅ 273.245


Alkaloid from the wood of *Vepris punctata*. Viscous oil. λ_{\max} 248 (log ϵ

21410); 262 (log ϵ 46000); 332 (log ϵ 10000); 355 (log ϵ 4300) (MeOH).

Chaturvedula, V.S.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 532-534 (5-Methoxymaculine)
Halstead, C.W. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 351-356 (5,6,7-Trimethoxydictamnine)

Furo[2,3-*b*]quinoline-4,5,7,8-tetrol, 9CI F-216

4,5,7,8-Tetrahydroxyfuro[2,3-*b*]quinoline



C₁₁H₇NO₅ 233.18

▶ LV2796000

Tetra-Me ether: 4,5,7,8-Tetramethoxyfuro[2,3-*b*]quinoline. 5,7,8-Trimethoxydictamnine. **Acronycidine** [521-43-7]

C₁₅H₁₅NO₅ 289.287

Alkaloid from the bark of *Acronychia baueri* and *Melicope fareana* (Rutaceae). Needles or prisms (MeOH or EtOH). Mp 136-137.5°.

Tetra-Me ether, hydrochloride: Mp 120-121° dec.

Tetra-Me ether, picrate:

Yellow needles (MeOH). Mp 181.5-182.5°.

Lahey, F.N. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1950, **3**, 155 (*uv, struct, bibl*)


Batterman, T.J. *et al.*, *Aust. J. Chem.*, 1965, **18**, 859 (*pmr*)

Prabhakar, S. *et al.*, *Indian J. Chem.*, 1971, **9**, 191 (*synth, uv, ir*)

Sekiba, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1974, **47**, 2895 (*synth, uv, ir, pmr*)

Furo[2,3-*b*]quinoline-4,6,7,8-tetrol F-217

4,6,7,8-Tetrahydroxyfuro[2,3-*b*]quinoline. 6,7,8-Trihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₅ 233.18

▶ JI4642000

OH-form

Tetra-Me ether: 4,6,7,8-Tetramethoxyfuro[2,3-*b*]quinoline, 9CI. **Halfordinine**†. 6,7,8-Trimethoxydictamnine [26949-99-5]

C₁₅H₁₅NO₅ 289.287

Alkaloid from *Halfordia scleroxyla*, *Melicope perspicuineria*, *Oricia suaveolens* and *Teclea verdoorniana* (Rutaceae). Needles (C₆H₆/petrol or MeOH). Mp 150-152°.

6,7-Methylene, 4-Me ether: 9-Methoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-

4-ol, 9CI. 8-Hydroxy-4-methoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline.

Tecleine. 8-Hydroxy-6,7-methylenedioxydictamnine

[82202-82-2]

C₁₃H₉NO₅ 259.218

Alkaloid from the stem bark of *Teclea verdoorniana* (Rutaceae). Needles (EtOH). Mp 256-257°.

6,7-Methylene, 4-Me ether, *Ac*:

Plates (EtOAc). Mp 186-188°.

6,7-Methylene, 4,8-di-Me ether: 4,9-Dimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline, 9CI. 4,8-Dimethoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline. 8-Methoxy-6,7-methylenedioxydictamnine. **Flindersiamine** [522-06-5]

C₁₄H₁₁NO₅ 273.245

Alkaloid from *Flindersia collina*, *Flindersia maculosa*, *Araliopsis soyauxii*, and *Teclea verdoorniana* (Flindersiaceae, Rutaceae). Needles (MeOH). Mp 206-207°.

6,7-Methylene, 4,8-di-Me ether, picrate:

Yellow needles (1-propanol). Mp 200°.

6,7-Methylene, 8-O-(3-methyl-2-butenyl), 4-Me ether: 4-Methoxy-6,7-methylenedioxy-8-prenyloxyfuro[2,3-*b*]quinoline.

Tecleamine

[84658-45-7]

C₁₈H₁₇NO₅ 327.336

Alkaloid from the stem bark of *Teclea ouabanguensis* (Rutaceae). Prisms (Et₂O/petrol). Mp 112-113°.

NH-form

6,7-Methylene, 8-Me ether, N-Me: 4-Methoxy-5-methyl-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-9(5H)-one, 9CI.

Isoflindersiamine

[1357-99-9]

C₁₄H₁₁NO₅ 273.245

Alkaloid from the leaves and twigs of *Helieta parvifolia* (Rutaceae). Also obt. by heating Furo[2,3-*b*]quinoline-4,6,7,8-tetrol, F-217 with MeI. Cryst. (MeOH or MeOH aq.). Mp 212-213° (209-211°).

Anet, F.A. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 412 (*Flindersiamine*, *Isoflindersiamine*)

Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1954, **7**, 181 (*Flindersiamine*)

Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1963, **1**, 348 (*synth*)

Clugston, D.M. *et al.*, *Can. J. Chem.*, 1965, **43**, 2516 (*ms*)

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1968, **21**, 3075 (*Halfordinine*)

Murphy, S.T. *et al.*, *Aust. J. Chem.*, 1974, **27**, 187 (*Halfordinine*)

Chang, P.T.O. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 561 (*Isoflindersiamine, isol, uv, ir, pmr, ms*)

Vaquette, J. *et al.*, *Phytochemistry*, 1976, **15**, 743 (*Flindersiamine, isol, uv, ir, pmr, ms*)

Ayafor, J.F. *et al.*, *J. Nat. Prod.*, 1982, **45**, 714 (*Tecleamine*)


Ayafor, J.F. *et al.*, *J.C.S. Perkin 1*, 1982, 909 (*Tecleine*)

Biavatti, M.W. *et al.*, *J. Braz. Chem. Soc.*, 2002, **13**, 66-70 (*isol, cryst struct*)

Chaturvedula, V.S.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 532-534 (*Flindersiamine, cmr*)

Furo[2,3-*b*]quinoline-4,5,6-triol, 9CI F-218

4,5,6-Trihydroxyfuro[2,3-*b*]quinoline



C₁₁H₉NO₄ 219.196

Tri-Me ether: 4,5,6-Trimethoxyfuro[2,3-*b*]quinoline. 5,6-Dimethoxydictamnine C₁₄H₁₃NO₄ 259.261


Alkaloid from the stems and leaves of *Haplophyllum buxbaumii* (Rutaceae). Amorph.

Ulubelen, A. *et al.*, *Phytochemistry*, 1985, **24**, 372 (*isol, uv, ir, pmr, ms, struct*)

Furo[2,3-*b*]quinoline-4,5,7-triol F-219

4,5,7-Trihydroxyfuro[2,3-*b*]quinoline.

5,7-Dihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₄ 217.181

OH-form

Tri-Me ether: 4,5,7-Trimethoxyfuro[2,3-*b*]quinoline. 5,7-Dimethoxydictamnine [96400-49-6]

Alkaloid from the stems and leaves of *Haplophyllum buxbaumii* (Rutaceae). Amorph.

NH-form

5,7-Di-Me ether, N-Me: 5,7-Dimethoxy-9-methylfuro[2,3-*b*]quinolin-4(9H)-one. **Glycarpine**

[71295-48-2]

C₁₄H₁₃NO₄ 259.261

Minor alkaloid from the leaves of *Glycosmis cyanocarpa* (Rutaceae).

Cryst. (C₆H₆/petrol). Mp 170-171°. The proposed struct. has been questioned.


Sarkar, M. *et al.*, *Phytochemistry*, 1978, **17**, 2145 (*Glycarpine*)

Kuo, S.C. *et al.*, *J. Nat. Prod.*, 1984, **47**, 47

Ulubelen, A. *et al.*, *Phytochemistry*, 1985, **24**, 372 (*tri-Me ether*)

Furo[2,3-*b*]quinoline-4,5,8-triol F-220

4,5,8-Trihydroxyfuro[2,3-*b*]quinoline



C₁₁H₇NO₄ 217.181


4,8-Di-Me ether: 4,8-Dimethoxyfuro[2,3-*b*]quinolin-5-ol, 9CI. 5-Hydroxy-4,8-

dimethoxyfuro[2,3-*b*]quinoline. 5-Hydroxy-8-methoxydictamnine
[128508-21-4]
C₁₃H₁₁NO₄ 245.234
Alkaloid from *Alstonia mairei* (Apocynaceae).

Ye, J. *et al.*, *Huaxue Xuebao*, 1989, **47**, 1012; *CA*, **113**, 74721j

Furo[2,3-*b*]quinoline-4,6,7-triol F-221

4,6,7-Trihydroxyfuro[2,3-*b*]quinoline.
6,7-Dihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₄ 217.181

OH-form

4,6-Di-Me ether: 4,6-Dimethoxyfuro[2,3-*b*]quinolin-7-ol. 7-Hydroxy-4,6-dimethoxyfuro[2,3-*b*]quinoline. **Heliparvifoline**. 7-Hydroxy-6-methoxydictamnine
[59442-99-8]
C₁₃H₁₁NO₄ 245.234

Alkaloid from the leaves and twigs of *Helietta parvifolia* and from the leaves of *Melicope confusa* (Rutaceae). Cryst. (MeOH). Mp 245-247°. The 7-Hydroxy-6-methoxydictamnine (Mp 229°) reported in 1981 from *Monnieria trifolia* appears to be identical with Delbine below. (J. Bhattacharyya *et al.*, *Heterocycles*, 1981, **16**, 371; 1983, **20**, 1063).

4,7-Di-Me ether: 4,7-Dimethoxyfuro[2,3-*b*]quinolin-6-ol. 6-Hydroxy-4,7-dimethoxyfuro[2,3-*b*]quinoline. **Delbine**†
[77145-73-4]
C₁₃H₁₁NO₄ 245.234

Alkaloid from the leaves of *Monnieria trifolia* (Rutaceae). Mp 229-231°. Genus name often given as *Monniera*.

Tri-Me ether: 4,6,7-Trimethoxyfuro[2,3-*b*]quinoline. **Kokusaginine**. 6,7-Dimethoxydictamnine
[484-08-2]
C₁₄H₁₃NO₄ 259.261

Alkaloid present in *Evodia* and *Orixa* spp. (Rutaceae) and in *Flindersia* spp., *Ruta graveolens*, *Acronychia pedunculata* and *Acronychia laurifolia*. Cytotoxic agent. Prisms (EtOH). Mp 171°. λ_{max} 245 (ε 67000); 252 (ε 5900); 308 (ε 12900); 321 (ε 13200); 335 (ε 9500) (EtOH) (Berdy).

▶ LV2798000

Tri-Me ether, hydrochloride: Mp 225° dec.

Tri-Me ether, picrate:

Yellow needles (EtOH). Mp 218°.

7-O-(3-Methyl-2-butenyl), 4,6-di-Me ether: 4,6-Dimethoxy-7-prenyloxyfuro[2,3-*b*]quinoline. **Nobiline**†
[114216-84-1]
C₁₈H₁₉NO₄ 313.352

Alkaloid from the leaves and fruits of *Teclea nobilis* (Rutaceae). Mp 117-119°.

6-O-(3-Methyl-2-butenyl), 4,7-di-Me ether: 4,7-Dimethoxy-6-prenyloxyfuro[2,3-*b*]quinoline. **Tecleanatalensine B**

C₁₈H₁₉NO₄ 313.352

Alkaloid from the leaves of *Teclea natalensis*. Pale yellow gum. λ_{max} 248 (log ε 4.4); 253 (log ε 4.4); 310 (log ε 3.7); 322 (log ε 3.7); 336 (log ε 3.7) (CH₂Cl₂).

7-O-(2ξ,3-Epoxy-3-methylbutyl), 4,6-di-Me ether: **Isotecleoxine**
[651737-87-0]
C₁₈H₁₉NO₅ 329.352

Alkaloid from *Teclea nobilis*. Amorph. solid. [α]_D -13.3 (c, 0.06 in MeOH). λ_{max} 244 (log ε 3.41); 309 (log ε 4.05); 321 (log ε 4.02); 334 (log ε 4.11) (MeOH).

6-O-(2ξ,3-Epoxy-3-methylbutyl), 4,7-di-Me ether: **Tecleoxine**. **Tecleanatalensine A**
[651737-86-9]
C₁₈H₁₉NO₅ 329.352

Alkaloid from *Teclea natalensis* and *Teclea nobilis*. Needles (petrol). Mp 120-121°. [α]_D +11 (c, 0.19 in CH₂Cl₂). λ_{max} 246 (log ε 4.51); 253 (log ε 4.47); 310 (log ε 3.85); 322 (log ε 3.88); 334 (log ε 3.79) (CH₂Cl₂).

4-O-(2S,3-Dihydroxy-3-methylbutyl), 6,7-di-Me ether: **1-[(6,7-Dimethoxyfuro[2,3-*b*]quinolin-4-yl)oxy]-3-methyl-2,3-butanediol**

C₁₈H₂₁NO₆ 347.367

Alkaloid from the aerial parts of *Clausena dumiana*. [α]_D²⁴ -14.4 (c, 0.45 in MeOH). Config. drawn incorrectly in ref. λ_{max} 210; 244; 250; 293; 308; 321; 334 (MeOH).

6-O-(2ξ,3-Dihydroxy-3-methylbutyl), 4,7-di-Me ether: **Montrifoline**. **Nkolbisine**
[77145-74-5]
[95462-87-6]
C₁₈H₂₁NO₆ 347.367

Alkaloid from the stem bark of *Teclea verdoorniana* ((-)-form) and from the aerial parts of *Haplophyllum vulcanicum* ((+)-form). Nkolbisine is also present in the leaves of *Monnieria trifolia* and the stem bark of *Teclea ouabangensis*, but the specific rotms. of the alkaloids from these two sources have not been recorded (Rutaceae). Needles. Mp 190-192°. [α]_D²⁴ -16.5 (c, 2.0 in MeOH) ((-)-form). [α]_D²⁵ +23 (c, 0.5 in MeOH) ((+)-form).

7-O-(2ξ,3-Dihydroxy-3-methylbutyl), 4,6-di-Me ether: **Evolatine**
[524-90-3]
C₁₈H₂₁NO₆ 347.367

Alkaloid from *Evodia alata* (Rutaceae). Cryst. (Me₂CO). Mp 201-202°. [α]_D²⁴ +17.5 (c, 0.3 in EtOH). The alkaloid reported as Evolatine in 1981 from *Monnieria trifolia* appears to be identical with Montrifoline (J. Bhattacharyya *et al.*, *Heterocycles*, 1981, **16**, 371; 1983, **20**, 1063).

6-O-(2ξ-Hydroxy-3-methoxy-3-methylbutyl), 4,7-di-Me ether: **3'-O-Methyl-nkolbisine**. 3'-O-Methylmontrifoline

[651737-88-1]

C₁₉H₂₃NO₆ 361.394

Alkaloid from *Teclea nobilis*. Needles. Mp 168-169°. [α]_D -2.8 (c, 0.04 in MeOH). λ_{max} 245 (log ε 3.26); 251 (log ε 3.25); 308 (log ε 3.98); 320 (log ε 3.97); 334 (log ε 4.11) (MeOH).

6-O-(3-Chloro-2ξ-hydroxy-3-methylbutyl), 4,7-di-Me ether: **Chlorodesnkolbisine**
[651737-89-2]
C₁₈H₂₀ClNO₅ 365.812

Alkaloid from *Monnieria trifolia*. Light yellow powder. Racemic. This struct. was incorrectly assigned to a compd. from *Teclea nobilis* in 2003. λ_{max} 247 (log ε 4.57); 320 (log ε 4.03) (MeOH).

6,7-Methylene, 4-Me ether: 9-Methoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline. 4-Methoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline. 6,7-Methylenedioxydictamnine. **Maculine**
[524-89-0]
C₁₃H₉NO₄ 243.218

Alkaloid from *Flindersia dissosperma* and *Flindersia maculosa* (Flindersiaceae). Needles (MeOH). Mp 196-197°.

▶ JI4644000

6,7-Methylene, 4-Me ether, hydrochloride: Mp 205-210° dec.

6,7-Methylene ether, 4-O-(2ξ,3-dihydroxy-3-methylbutyl): **Maculosine**†
C₁₇H₁₇NO₆ 331.324

Alkaloid from *Flindersia maculosa* bark. Needles (EtOH). Mp 229-230°. [α]_D²⁰ +36 (Py). Originally given the MF C₁₅H₁₅NO₅.

NH-form

6,7-Methylene ether, N-Me: 5-Methyl-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-9(5H)-one. 6,7-Methylenedioxy-9-methylfuro[2,3-*b*]quinolin-9(5H)-one. **Isomaculine**
[482-84-8]
C₁₃H₉NO₄ 243.218

Alkaloid from the root bark of *Esenbeckia pilocarpoides* (Asteraceae), also obt. by heating Maculine with MeI. Needles (CHCl₃). Mp 269° (237-239°).

Anet, F.A.L. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 412 (*Kokusaginine*)

Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1954, **7**, 181 (*Maculine*, *Maculosine*)

Gell, R.J. *et al.*, *Aust. J. Chem.*, 1955, **8**, 114; 422 (*Evolatine*, *Maculine*)

Prager, R.H. *et al.*, *Aust. J. Chem.*, 1960, **13**, 380 (*Maculosine*, *synth*)

Robertson, A.V. *et al.*, *J.C.S.*, 1960, 2458 (*Evolatine*, *ir*)

Robertson, A.V. *et al.*, *Aust. J. Chem.*, 1963, **16**, 450 (*pmr*, *derivs*)

Clugston, D.M. *et al.*, *Can. J. Chem.*, 1965, **43**, 2516 (*ms*, *derivs*)

Mester, I. *et al.*, *Phytochemistry*, 1971, **10**, 2205 (*synth*, *derivs*)

Chang, P.T.O. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 561 (*Heliparvifoline*)

Sekiba, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1978, **51**, 325 (*synth*, *ir*, *pmr*, *derivs*)


Bhattacharyya, J. *et al.*, *Heterocycles*, 1981, **16**, 371 (*Delbine*, *Montrifoline*)

Ranade, A.C. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 528 (*Maculine*, *synth*)

- Ayafor, J.F. *et al.*, *J. Nat. Prod.*, 1982, **45**, 182; 714 (*Nkolbisine*)
 Patra, A. *et al.*, *Heterocycles*, 1984, **22**, 2821 (*Montrifoline*)
 Bevalot, F. *et al.*, *Planta Med.*, 1984, **50**, 522 (*Isomaculine*)
 Kang, S.S. *et al.*, *Arch. Pharmacol. Res.*, 1986, **9**, 11; *CA*, **105**, 39390p (*Heliparvifoline, isol*)
 Yenesew, A. *et al.*, *Phytochemistry*, 1988, **27**, 651 (*Nobiline*)
 Cui, B. *et al.*, *Phytochemistry*, 1999, **52**, 95-98 (*Kokusaginine, activity*)
 He, H.-P. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 3187-3193 (*Clausena dunniana ether*)
 Al-Rehailly, A.J. *et al.*, *Phytochemistry*, 2003, **64**, 1405-1411 (*Tecleoxine, Isotecleoxine, 3'-Methylnkolbisine, Nobiline*)
 Nunes, F.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 864-866 (*Maculine, pmr, cmr*)
 Tarus, P.K. *et al.*, *Phytochemistry*, 2005, **66**, 703-706 (*Tecleanatalensines A,B*)
 Cao, S. *et al.*, *Phytochemistry*, 2008, **69**, 553-557 (*Chlorodesnkolbisine, Nkolbisine*)

Furo[2,3-*b*]quinoline-4,6,8-triol F-222

4,6,8-Trihydroxyfuro[2,3-*b*]quinoline.
 6,8-Dihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₄ 217.181

▶ LV2799000

OH-form

Tri-Me ether: 4,6,8-Trimethoxyfuro[2,3-*b*]quinoline, 9CI. **Maculosidine**. 6,8-Dimethoxydictamnine [522-19-0]
 C₁₄H₁₃NO₄ 259.261
 Alkaloid from *Flindersia maculosa*, *Acronychia laurifolia* and *Eriostemon* spp. (Flindersiaceae, Rutaceae). Cytotoxic. Plates (EtOH). Mp 186°.

8-O-(3-Methyl-2-butenyl), 4,6-di-*Me ether*: 4,6-Dimethoxy-8-(3-methyl-2-butenyloxy)furo[2,3-*b*]quinoline. **Tecleabine** [651737-85-8]
 C₁₈H₁₉NO₄ 313.352
 Alkaloid from *Teclea nobilis*. Needles. Mp 107-108°. λ_{max} 248 (log ε 3.22); 295 (log ε 4.16); 307 (log ε 4.18); 339 (log ε 4.29); 353 (log ε 4.31) (MeOH).

NH-form

6,8-Di-*Me ether*, N-*Me*: 6,8-Dimethoxy-9-methylfuro[2,3-*b*]quinolin-4(9H)-one, 9CI. **Isomaculosidine** [518-96-7]
 C₁₄H₁₃NO₄ 259.261
 Alkaloid from the roots of *Dictamnus albus* and *Dictamnus caucasicus* (Rutaceae). Also obt. by heating Furo[2,3-*b*]quinoline-4,6,8-triol, F-222 with MeI. Needles (EtOH, CHCl₃/hexane or C₆H₆). Mp 170-172° (167-168°).


Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1954, **7**, 181 (*isol, uv, struct*)

Duffield, A.M. *et al.*, *Aust. J. Chem.*, 1962, **15**, 812; 1963, **16**, 123 (*isol*)

- Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1963, **1**, 17 (*synth*)
 Clugston, D.M. *et al.*, *Can. J. Chem.*, 1965, **43**, 2516 (*ms*)
 Kikvidze, I.M. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 659 (*Isomaculosidine*)
 Sekiba, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 577 (*synth, uv*)
 Storer, R. *et al.*, *Tetrahedron*, 1973, **29**, 1217 (*Isomaculosidine*)
 Xu, W.H. *et al.*, *Huaxue Xuebao*, 1984, **42**, 899 (*Maculosidine, isol*)
 Lin, T.P. *et al.*, *J. Nat. Prod.*, 1987, **50**, 631 (*synth, uv, ir, pmr, cmr*)
 Cui, B. *et al.*, *Phytochemistry*, 1999, **52**, 95-98 (*Maculosidine, activity*)
 Al-Rehailly, A.J. *et al.*, *Phytochemistry*, 2003, **64**, 1405-1411 (*Tecleabine*)

Furo[2,3-*b*]quinoline-4,7,8-triol F-223

4,7,8-Trihydroxyfuro[2,3-*b*]quinoline.
 7,8-Dihydroxyfuro[2,3-*b*]quinolin-4(9H)-one



C₁₁H₇NO₄ 217.181

OH-form

4,7-Di-*Me ether*: 4,7-Dimethoxyfuro[2,3-*b*]quinolin-8-ol. 8-Hydroxy-4,7-dimethoxyfuro[2,3-*b*]quinoline. **Isolahopline**. 8-Hydroxy-7-methoxydictamnine [62580-17-0]
 C₁₃H₁₁NO₄ 245.234
 Alkaloid from the leaves of *Zanthoxylum arborescens* and *Teclea simplicifolia* (Rutaceae). Cryst. (hexane/EtOAc), plates (Me₂CO/petrol). Mp 129-130° (121-123°).

4,8-Di-*Me ether*: 7-Hydroxy-4,8-dimethoxyfuro[2,3-*b*]quinoline. 4,8-Dimethoxyfuro[2,3-*b*]quinolin-7-ol. **Haplopinine**. 7-Hydroxy-8-methoxydictamnine [5876-17-5]
 C₁₃H₁₁NO₄ 245.234
 Alkaloid from *Haplophyllum robustum*, *Zanthoxylum cuspidatum*, *Monnieria trifolia* and *Melicope lasioneura* (Rutaceae). Mp 203-204°.

4,8-Di-*Me ether*, 7-O-β-D-glucopyranoside: **Glycohaplopinine**. Glucohaplopinine [74201-15-3]
 C₁₉H₂₁NO₉ 407.376
 Alkaloid from the aerial parts of *Haplophyllum perforatum* (Rutaceae). Mp 217-218°. [α]_D -41 (c, 0.516 in Py).

4,8-Di-*Me ether*, 7-O-α-L-rhamnopyranoside: **Glycoepine** [55740-45-9]
 C₁₉H₂₁NO₈ 391.377
 Alkaloid from *Haplophyllum perforatum* (Rutaceae). Cryst. (MeOH). Mp 224-225°. [α]_D -66.3 (c, 2.32 in Py).

4,8-Di-*Me ether*, 7-O-(O-acetyl-α-L-rhamnopyranoside): **Monoacetylglycoepine**

[67184-79-6]
 C₂₁H₂₃NO₉ 433.414
 Alkaloid from aerial parts of *Haplophyllum perforatum*. Mp 120-121°.

4,8-Di-*Me ether*, 7-O-(di-O-acetyl-α-L-rhamnopyranoside): **Diacetylglycoepine**
 C₂₃H₂₅NO₁₀ 475.451
 Alkaloid from *Haplophyllum perforatum*.

4,8-Di-*Me ether*, 7-O-(tri-O-acetyl-α-L-rhamnopyranoside): **Triacetylglycoepine** [55740-46-0]
 C₂₅H₂₇NO₁₁ 517.488
 Alkaloid from *Haplophyllum perforatum*. Cryst. (C₆H₆/petrol). Mp 181-182°. [α]_D -91 (c, 0.4 in EtOH).

4,8-Di-*Me ether*, 7-O-[β-D-glucopyranosyl-(1→3)-α-L-rhamnopyranoside]: **Haplopinine** [115345-33-0]
 C₂₅H₃₁NO₁₃ 553.519
 Alkaloid from aerial parts of *Haplophyllum perforatum* (Rutaceae). Mp 227-228°. [α]_D -74 (c, 2.83 in Py).

4,8-Di-*Me ether*, 7-O-[β-D-glucopyranosyl-(1→3)-2-O-acetyl-α-L-rhamnopyranoside]: **Haplopidine** [115345-32-9]
 C₂₇H₃₃NO₁₄ 595.556
 Alkaloid from *Haplophyllum perforatum*. Mp 158-160°. [α]_D -77 (c, 2.66 in Py). λ_{max} 249 (log ε 4.87); 275 (sh) (log ε 3.25); 320 (log ε 3.85); 333 (log ε 3.77) (EtOH).

Tri-Me ether: 4,7,8-Trimethoxyfuro[2,3-*b*]quinoline. **Skimmianine**. Chloroxylinine. Pentaphylline. 7,8-Dimethoxydictamnine. β-Fagarine [83-95-4]
 C₁₄H₁₃NO₄ 259.261
 Alkaloid from *Skimmia japonica*, *Haplophyllum tuberculatum*, *Zanthoxylum ovalifolium*, *Zanthoxylum coco*, *Fagara coco*, *Fagara zanthoxyloides*, *Acronychia pedunculata*, *Acronychia laurifolia*, *Stauranthus perforatus* and many other spp., mostly in the Rutaceae. Spasmolytic agent. Shows pharmacol. props. similar to 2-(Methylamino)-1-phenyl-1-propanol, M-386. Mp 176-177°. Log P 2.99 (calc). λ_{max} 204 ; 249 ; 323 (MeOH) (Berdy). λ_{max} 250 (ε 76000); 319 (ε 7200); 332 (ε 16000) (EtOH) (Berdy).

▶ LV2800000

Tri-Me ether, perchlorate: Mp 210°.

Tri-Me ether, picrate: Mp 197°.

4-Et, 7,8-di-*Me ether*: 4-Ethoxy-7,8-dimethoxyfuro[2,3-*b*]quinoline. **O-Ethyl-norskimmianine** [106006-02-4]
 C₁₅H₁₅NO₄ 273.288
 Alkaloid from roots of *Dictamnus dasycarpus* (Rutaceae).

4-O-(2ξ,3-Dihydroxy-3-methylbutyl), 7,8-di-*Me ether*: **Nigdenine** [95378-12-4]
 C₁₈H₂₁NO₆ 347.367
 Alkaloid from the aerial parts of *Haplophyllum vulcanicum* (Rutaceae).


- Cryst. (CHCl₃/hexane). Mp 177-180°. [α]_D²⁵ +5.3 (c, 0.3 in MeOH).
- 7-O-(2,3-Epoxy-3-methylbutyl), 4,8-di-Me ether: **Anhydroevoxine** [24099-25-0] C₁₈H₁₉NO₅ 329.352 Alkaloid from the branch ends of *Evodia xanthoxyloides* and from *Haplophyllum ferganicum* and *Haplophyllum perforatum*. Cryst. (Et₂O/hexane). Mp 141-144°. [α]_D +13 (CHCl₃).
- 7-O-(3-Methyl-2-oxobutyl), 4,8-di-Me ether: **Evoxidine**. 7-(3-Methyl-2-oxobutoxy)-8-methoxydictamnine [572-24-7] C₁₈H₁₉NO₅ 329.352 Alkaloid from *Evodia xanthoxyloides* (Rutaceae). Cryst. (EtOH). Mp 136-137°.
- 7-O-(2R,3-Dihydroxy-3-methylbutyl), 4,8-di-Me ether: 7-(2,3-Dihydroxy-3-methylbutoxy)-8-methoxydictamnine. **Evoxine**. *Haploperine*. *Haplophytine B* [522-11-2] C₁₈H₂₁NO₆ 347.367 Alkaloid from *Evodia xanthoxyloides*, *Monnieria trifolia* and several *Haplophyllum* spp. (Rutaceae). Mp 154-155°. [α]_D²² +14.6 (EtOH). [α]_D +63.6 (c, 0.33 in MeOH) (Haplophytine B).
- 7-O-(2-Acetoxy-3-hydroxy-3-methylbutyl), 4,8-di-Me ether: **Evoxine acetate**. 2'-Acetylevoxine [40817-01-4] [64397-07-5] C₂₀H₂₃NO₇ 389.404 Alkaloid from *Haplophyllum hispanicum*, *Haplophyllum ferganicum* and *Haplophyllum ramosissimum*. Cryst. (C₆H₆). Mp 168.5-169.5°.
- 7-O-(2-Hydroxy-3-methoxy-3-methylbutyl), 4,8-di-Me ether: **Methylevoxine** [56775-80-5] C₁₉H₂₃NO₆ 361.394 Alkaloid from *Haplophyllum glaberrimum*, *Haplophyllum perforatum* and *Haplophyllum obtusifolium*. Needles. Mp 55-56°. λ_{\max} 251 (log ϵ 4.91); 322 (log ϵ 3.78); 334 (log ϵ 3.77) (no solvent reported).
- 7-O-(2R-Hydroxy-3-ethoxy-3-methylbutyl), 4,8-di-Me ether: **Ethylevoxine** [953812-17-4] C₂₀H₂₅NO₆ 375.421 Alkaloid from the leaves of *Choisya ternata*. [α]_D -8.8 (c, 0.2 in CHCl₃) (as 2'-Ac).
- 7-O-(3-Chloro-2R-hydroxy-3-methylbutyl), 4,8-di-Me ether: **Chlorodeoxyevoxine**. *Chlorodesoxyevoxine* [953812-19-6] C₁₈H₂₀ClNO₅ 365.812 Alkaloid from the leaves of *Choisya ternata*. Cryst. (Me₂CO/hexane). Mp 137-138°. [α]_D +14.2 (c, 0.33 in EtOH).
- 7-O-(3-Chloro-2RS-hydroxy-3-methylbutyl), 4,8-di-Me ether: 7-O-(3-Chloro-2-hydroxy-3-methylbutyl)haplopine C₁₈H₂₀ClNO₅ 365.812 Alkaloid from *Monnieria trifolia*. Light yellow powder. Racemic. λ_{\max} 249 (log ϵ 4.68); 320 (log ϵ 3.97) (MeOH).
- 7-O-(3-Methyl-2-butenyl), 4,8-di-Me ether: 4,8-Dimethoxy-7-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline. 7-Isopentenylxyloxy- γ -fagarine. 8-Methoxy-7-prenyloxydictamnine. **Haplopine prenyl ether** [23417-92-7] C₁₈H₁₉NO₄ 313.352 Alkaloid from *Ptelea aptera*, *Melicope lasioneura* and *Choisya ternata* (biosynth.) (Rutaceae). Cryst. (MeOH). Mp 101-103°.
- 7-O-(2-Hydroxy-3-methyl-3-butenyl), 4,8-di-Me ether: **Evodine**[†] [6989-38-4] C₁₈H₁₉NO₅ 329.352 Alkaloid from the leaves of *Evodia xanthoxyloides* (Rutaceae). Mp 153-154°. [α]_D²⁰ -3 (c, 1.0 in CHCl₃). Abs. config. not detd.
- 7-O-(4-Hydroxy-3-methyl-2Z-butenyl), 4,8-di-Me ether: **Haplatine**. *Myrtifoline* [58480-57-2] C₁₈H₁₉NO₅ 329.352 Alkaloid from *Haplophyllum latifolium* (Rutaceae) and *Haplophyllum myrtifolium*. Cryst. (C₆H₆). Mp 139-140°.
- 7-O-(4-Acetoxy-3-methyl-2Z-butenyl), 4,8-di-Me ether: **Acetylhaplatine** [58480-58-3] C₂₀H₂₁NO₆ 371.389 Alkaloid from *Haplophyllum obtusifolium* (Rutaceae). Mp 87-88° (78-80°).
- 7-O-(4-Chloro-3-methyl-2 ζ -butenyl), 4,8-di-Me ether: **Haplobine** [107783-37-9] C₁₈H₁₈ClNO₄ 347.797 Alkaloid from the roots of *Haplophyllum obtusifolium* (Rutaceae). Cryst. (Me₂CO). Mp 151-153°.
- 8-O-(3-Methyl-2-butenyl), 4,7-di-Me ether: 4,7-Dimethoxy-8-prenyloxy-furo[2,3-*b*]quinoline. *Isoplopine prenyl ether*. 7-Methoxy-8-(3-methyl-2-butenyloxy)dictamnine [81536-07-4] C₁₈H₁₉NO₄ 313.352 Alkaloid from *Teclea simplicifolia* and *Zanthoxylum arborescens* (Rutaceae). Cryst. (EtOAc/hexane). Mp 120.5-121.5° (118-119°).
- 7-O-(6,7-Dihydroxy-3,7-dimethyl-2-octenyl), 4,8-di-Me ether: **Haplotubine** C₂₃H₂₉NO₆ 415.485 Alkaloid from *Haplophyllum tuberculatum*. Amorph. yellow powder. [α]_D²² -6 (c, 1.2 in CH₂Cl₂). λ_{\max} 250 (log ϵ 4.87); 320 (log ϵ 3.9); 332 (log ϵ 3.88) (MeOH).
- 7,8-Methylene, 4-Me ether: 6-Methoxy-1,3-dioxolo[4,5-*h*]furo[2,3-*b*]quinoline. 4-Methoxy-7,8-methylenedioxy-furo[2,3-*b*]quinoline. **Kokusagine**. 7,8-Methylenedioxydictamnine [482-32-6] C₁₃H₉NO₄ 243.218 Alkaloid from the roots of *Orixa japonica* (Rutaceae). Needles. Mp 201° (192-193°).
- 7,8-Methylene, 4-Me ether, picrate: Mp 178° (157°).

NH-form

- 7,8-Di-Me ether, N-Et: 9-Ethyl-7,8-dimethoxyfuro[2,3-*b*]quinolin-4(9H)-one. **8-Methoxytaifine** [84323-10-4] C₁₅H₁₅NO₄ 273.288 Alkaloid from *Ruta chalepensis* (Rutaceae). Needles (petrol). Mp 122-123°.
- Deloueu, V. et al., *J.A.C.S.*, 1942, **64**, 2326-2338 (*isol*, *Skimmianine*)
- Eastwood, F.W. et al., *Aust. J. Chem.*, 1954, **7**, 87-98 (*Evoxidine*, *Evoxine*)
- Terasaka, M. et al., *Chem. Pharm. Bull.*, 1954, **2**, 159 (*Kokusagine*)
- Briggs, L.H. et al., *J.C.S.*, 1960, 2458-2460 (*ir. derivs*)
- Prager, R.H. et al., *Aust. J. Chem.*, 1962, **15**, 301-304 (*Evodine*)
- Sidyakin, G.P. et al., *Dokl. Akad. Nauk SSSR. Ser. Khim.*, 1962, **19**, 39-40; *CA*, **57**, 15170 (*Haplopine*)
- Robertson, A.V. et al., *Aust. J. Chem.*, 1963, **16**, 451 (*pmr*, *Skimmianine*, *Evoxine*)
- Pai, B.R. et al., *Indian J. Chem.*, 1964, **2**, 449 (*Kokusagine*, *synth*)
- Clugston, D.M. et al., *Can. J. Chem.*, 1965, **43**, 2516-2521 (*ms*, *Skimmianine*)
- Dreyer, D.L. et al., *Phytochemistry*, 1969, **8**, 1013 (*Haplopine prenyl ether*)
- Dreyer, D.L. et al., *J.O.C.*, 1970, **35**, 2420-2422 (*Anhydroevoxine*)
- Gonzalez, A.G. et al., *An. Quim.*, 1972, **68**, 1133-1138 (*Evoxine acetate*)
- Akhmedzhanova, V.I. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 706-707; 1975, **11**, 291 (*Glycoferine*, *Methylevoxine*)
- Grundon, M.F. et al., *J.C.S. Perkin 1*, 1974, 2181-2184; 1975, 302-304 (*biosynth*, *Skimmianine*, *Evoxine*)
- Nesmelova, E.F. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 706-707 (*Haplatine*)
- Abdullaeva, Kh.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 363; 1978, **14**, 179-182; 1979, **15**, 782-783 (*acetylgycoferines*, *Glycohaplopine*)
- Moulis, C. et al., *Planta Med.*, 1981, **42**, 400-402 (*isol*, *uv*, *pmr*, *ms*, *Evoxine*, *Haplopine*)
- Bessonova, I.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 505-506; **20**, 125 (*Anhydroevoxine*, *Acetylhaplatine*)
- Tillequin, F. et al., *J. Nat. Prod.*, 1982, **45**, 486-488 (*Skimmianine*, *Haplopine*, *Haplopine prenyl ether*)
- Grina, J.A. et al., *J.O.C.*, 1982, **47**, 2648-2651 (*Isoplopine prenyl ether*)
- Mohr, N. et al., *Phytochemistry*, 1982, **21**, 1838 (*8-Methoxytaifine*)
- Patra, A. et al., *Heterocycles*, 1984, **22**, 2821 (*Nigdenine*)
- Razakova, D.M. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 259 (*Anhydroevoxine*)
- Bessonova, I.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 684-686 (*Haplobine*)
- Liu, T.P. et al., *CA*, 1987, **106**, 30033n (*O-Ethylskimmianine*)
- Razulova, Kh.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 731-734; 1988, **24**, 82-84 (*Haplosinine*, *Haplosidine*)
- Wondimu, A. et al., *Phytochemistry*, 1988, **27**, 959-960 (*Isoplopine*)
- Cox, O. et al., *Acta Cryst. C*, 1989, **45**, 1263-1265 (*cryst struct*, *Skimmianine*)
- Jackson, G.E. et al., *Spectrosc. Lett.*, 1990, **23**, 971-982 (*pmr*, *cmr*, *Skimmianine*)
- Chakravarty, A.K. et al., *Phytochemistry*, 1999, **50**, 1263-1266 (*isol*, *pmr*, *cmr*, *ms*, *Skimmianine*)

- Al-Rehaily, A.J. *et al.*, *Phytochemistry*, 2001, **57**, 597-602 (*Haplotubine*)
 Ali, M.S. *et al.*, *Phytochemistry*, 2001, **57**, 1277-1280 (*Haplophytine B*, *pmr*, *cmr*, *ms*)
 Boyd, D.R. *et al.*, *Chem. Comm.*, 2002, 3070-3071 (*Evoxine*, *abs config*)
 Al-Rehaily, A.J. *et al.*, *Phytochemistry*, 2003, **64**, 1405-1411 (*Haplopine prenyl ether*, *Anhydroevoxine*)
 Tarus, P.K. *et al.*, *Phytochemistry*, 2005, **66**, 703-706 (*Isohaplopine prenyl ether*, *cmr*)
 Boyd, D.R. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 2983-2991 (*Chlorodeoxyevoxine*, *Ethylevoxine*)
 Cao, S. *et al.*, *Phytochemistry*, 2008, **69**, 553-557 (7-*O*-(*Chlorohydroxymethylbutyl*)haplopine)

Furo[2,3-*b*]quinolin-4-ol, 9CI **F-224**
 4-Hydroxyfuro[2,3-*b*]quinoline.
*Furo[2,3-*b*]quinolin-4(9H)-one*



$C_{11}H_7NO_2$ 185.182
Picrate: Mp 164-165°.

OH-form

Me ether: 4-Methoxyfuro[2,3-*b*]quinoline. **Dictamnine**
 [484-29-7]

$C_{12}H_9NO_2$ 199.209

Alkaloid from *Dictamnus albus* and widely distributed in the family Rutaceae. Strong smooth muscle contractant. Active against gram-positive bacteria and fungi. Mp 132-133°.

Et ether: 4-Ethoxyfuro[2,3-*b*]quinoline, 9CI. **O-Ethylordictamnine**
 [95874-16-1]

$C_{13}H_{11}NO_2$ 213.235

Alkaloid from roots of *Dictamnus dasycarpus* (Rutaceae).

NH-form

N-Me: 9-Methylfuro[2,3-*b*]quinolin-4(9H)-one. **Isodictamnine**
 [484-74-2]

$C_{12}H_9NO_2$ 199.209

Alkaloid from trunk bark of *Helietta longifolia* and root bark of *Dictamnus albus* (Rutaceae). Mp 186-187°.

Asahina, Y. *et al.*, *Ber.*, 1930, **63**, 2045 (*Dictamnine*, *isol*)

Grundon, M.F. *et al.*, *J.C.S.*, 1957, 2177 (*Dictamnine*, *ir*, *uv*, *synth*)

Robertson, A.V. *et al.*, *Aust. J. Chem.*, 1963, **16**, 451 (*pmr*)

Clugston, D.M. *et al.*, *Can. J. Chem.*, 1965, **43**, 2516 (*ms*)

Mammarella, C.A. *et al.*, *An. Asoc. Quim. Argent.*, 1971, **59**, 239; *CA*, **76**, 43980f (*Isodictamnine*)

Collins, J.F. *et al.*, *J.C.S. Perkin 1*, 1974, 2177 (*biosynth*)


Narasimhan, N.S. *et al.*, *Tetrahedron*, 1974, **30**, 4153-4157 (*synth*)

Wolters, B. *et al.*, *Planta Med.*, 1981, **43**, 166-174 (*activity*)

Lin, T.P. *et al.*, *Hua Hsueh*, 1986, **44**, 96; 1987, **45**, 45; *CA*, **106**, 30033n; **111**, 229011n (*O-Ethylordictamnine*, *Isodictamnine*)

Kuo, S.C. *et al.*, *J. Het. Chem.*, 1991, **28**, 955 (*Isodictamnine*, *synth*)

Fusaricide **F-225**
 [184031-21-8]




Relative Configuration

$C_{17}H_{25}NO_3$ 291.389

Related to Cordopyridone D, C-647. *Isol*. from a *Fusarium* sp. Cytotoxic. Shows antifungal activity. Cryst. Mp 208°. $[\alpha]_D^{20} +194$ (c, 0.12 in $CHCl_3$). λ_{max} 218 (ε 25800); 295 (ε 4200) (MeOH).

McBrien, K.D. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1151 (*isol*, *ir*, *uv*, *pmr*, *cmr*, *cd*, *cryst struct*)
 Snider, B.B. *et al.*, *Synth. Commun.*, 2001, **31**, 2667-2679 (*synth*)

Fusarin A **F-226**
 [100079-50-3]



$C_{23}H_{29}NO_6$ 415.485

Metab. of *Fusarium moniliforme*. λ_{max} 358 (ε 32000) (MeOH) (Derep).

17 α -Hydroxy: **Fusarin D**


$C_{23}H_{29}NO_7$ 431.485

From *Fusarium moniliforme*. λ_{max} 358 (ε 32000) (MeOH) (Derep).

Gelderblom, W.C.A. *et al.*, *Chem. Comm.*, 1984, 122-125 (*isol*)

Steyn, P.S. *et al.*, *Chem. Comm.*, 1985, 1189-1191 (*biosynth*)

Fusarin C **F-227**
 [79748-81-5]



$C_{23}H_{29}NO_7$ 431.485

Prod. by many *Fusarium* spp. in infected corn. Also *isol*. from *Gibberella fujikuroi* and *Nectria coccinea*. λ_{max} 358 (ε 32000) (MeOH) (Derep).

► Mutagenic (comparable to Aflatoxin B₁). YQ2831500

1-Hydroxy: **Fusarin X**

[145569-98-8]

$C_{23}H_{29}NO_8$ 447.484

Prod. by *Fusarium moniliforme*.

8*Z*-Isomer: [90987-21-6]

$C_{23}H_{29}NO_7$ 431.485

From *Gibberella fujikuroi* and *Nectria coccinea*. Cryst. (CH_2Cl_2). Mp 170-172° (synthetic). λ_{max} 350 (ε 25000) (MeOH) (Berdy).

6*Z*-Isomer: [194152-45-9]

$C_{23}H_{29}NO_7$ 431.485

Prod. by *Nectria coccinea*. Pale yellow oil.

Gelderblom, W.C.A. *et al.*, *J. Agric. Food Chem.*, 1984, **32**, 1064 (*isol*, *struct*)

Gaddamidi, V. *et al.*, *J. Agric. Food Chem.*, 1985, **33**, 652 (*cmr*, *conformn*)


Farber, J.M. *et al.*, *J. Agric. Food Chem.*, 1986, **34**, 963 (*isol*)

Barrero, A.F. *et al.*, *Phytochemistry*, 1991, **30**, 2259 (*isol*)

Lu, F.X. *et al.*, *Chem. Res. Toxicol.*, 1993, **6**, 91-96; 97-101 (*Fusarin X*)

Eilbert, F. *et al.*, *J. Antibiot.*, 1997, **50**, 443-445 (6*Z*-isomer, 8*Z*-isomer)

Fusarin F **F-228**
 [139768-10-8]




$C_{23}H_{29}NO_7$ 431.485

Isol. from *Fusarium moniliforme*. λ_{max} 293 ; 370 (ε 40200) (MeOH) (Berdy).

Savard, M.E. *et al.*, *J. Nat. Prod.*, 1992, **55**, 64 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Fuscine **F-229**
 [3-(1,2,3,4-Tetrahydro-6,8-dihydroxy-1-isoquinolinyl)propyl]guanidine, 9CI
 [155210-58-5]




$C_{13}H_{20}N_4O_2$ 264.327

Alkaloid from the body wall of the seastar *Perknaster fuscus antarcticus*. Ichthyotoxin. Oil. CAS no. refers to monohydrochloride. λ_{max} 278 (ε 2510) (MeOH) (Derep).

Harper, M.K. *et al.*, *Nat. Prod. Lett.*, 1992, **1**, 71-74 (*isol*, *pmr*, *cmr*, *uv*, *ir*)


Fuseine **F-230**




$C_{17}H_{13}NO_3$ 279.295

Alkaloid from the trunk wood of *Fusaea longifolia* (Annonaceae). Mp 280° dec. (subl. at 180°). *Abs. config.* not determined, no opt. rotn. reported. λ_{max} 235 (log ε 4.48); 273 (log ε 4.55); 317 (log ε

3.91) (EtOH).

Braz Filho, R. *et al.*, *Phytochemistry*, 1976, **15**, 1187-1188 (*isol, uv, ir, pmr, ms, struct*)**Fusigen, 9CI***Fusarinine C*
[19624-79-4] $C_{33}H_{51}FeN_6O_{12}$ 779.646Isol. from *Fusarium cubense*, *Aspergillus nidulans* and *Penicillium chrysogenum*. Siderophore. Sol. H_2O , $MeOH$. λ_{max} 250 : 439 (ϵ 3162) (H_2O) (Berdy).Dieckmann, H. *et al.*, *Eur. J. Biochem.*, 1967, **3**, 213 (*isol, struct*)Charlang, G. *et al.*, *J. Bacteriol.*, 1982, **150**, 785 (*isol*)**Futoamide**

7-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,6-heptadienamide, 9CI. N-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]isobutylamine [114092-54-5]


**F-232** $C_{18}H_{23}NO_3$ 301.385

Related to 1-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]pyrrolidine, M-452.

(E,E)-form [23477-80-7]Alkaloid from *Piper futokadsura*, *Piper longum* (long pepper) and *Piper hancei*. Cryst. (Me_2CO /hexane). Mp 128-130°.Takahashi, S. *et al.*, *Phytochemistry*, 1969, **8**, 321 (*isol*)Vig, O.P. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 458-460 (*synth*)Blade, R.J. *et al.*, *Tet. Lett.*, 1987, **28**, 3857-3860 (*synth*)Li, S. *et al.*, *Yaoxue Xuebao*, 1987, **22**, 196 (*isol*)Das, B. *et al.*, *Fitoterapia*, 1998, **69**, 548 (*isol, pmr, cmr*)

Gabonine

G-1

C₄₂H₅₆N₄O₈ 744.926

Tentative struct. Prob. artifact arising from autoxidn. of Ibogaline in I-4.

Alkaloid from numerous *Tabernanthe* spp. (Apocynaceae). Mp 223-226°. [α]_D²⁴ +65 (CHCl₃).Dickel, D.F. et al., *J.A.C.S.*, 1958, **80**, 123 (*isol*)
Taylor, W.I. et al., *J.O.C.*, 1965, **30**, 309 (*uv*,
pmr, *struct*)

Galanthamidine


G-2

C₁₈H₂₃NO₅ 333.383Amaryllidaceae alkaloid. Struct. unknown. Isol. from the roots of *Galanthus voronovii* (Amaryllidaceae). Cryst. (MeOH). Mp 211-213°. [α]_D -94.2 (c, 0.53 in MeOH).*Methiodide*: Mp 219° dec.Proskurnina, N.F. et al., *Zh. Obshch. Khim.*, 1956, **26**, 172-173; *J. Gen. Chem. USSR* (*Engl. Transl.*), 1956, **26**, 179-180 (*isol*)

Galanthamine†, 9CI

G-3

4a,5,9,10,11,12-Hexahydro-3-methoxy-11-methyl-6H-benzofuro[3a,3,2-ef][2]-benzazepin-6-ol



(–)-form

C₁₇H₂₁NO₃ 287.358

(+)–form [60384-53-4]

Synthetic. Mp 133-135°. [α]_D¹⁸ +126 (c, 0.083 in EtOH).

1,2-Dihydro, N-de-Me: Demethyldihydrogalanthamine

C₁₆H₂₁NO₃ 275.347Component of the quasi-racemic alkaloid Narcissamine isol. from *Narcissus pseudonarcissus* and *Amaryllis* spp., which was a quasiracemate of (+)-Demethyldihydrogalanthamine and N-Demethylgalanthamine (see below). Mp 77-87° (hydrate) Mp 134° (anhyd.).

1,2-Dihydro, N-de-Me, O,N-di-Ac:

Cryst. (EtOH). Mp 177-179°.

3-Epimer: Synthetic. [α]_D +211 (c, 0.9 in EtOH).

(–)-form

Galantamine, BAN, INN, USAN. Jilkon.*Karantonin*. *Lycoremine*. *Lycorimine*. *Nevalina*. *Nivalin*. *Razadyne*. *Reminyl*. *NIH 7380*

[357-70-0]

Alkaloid from *Galanthus voronovii* and a very large number of other spp. in the Amaryllidaceae. Cholinesterase inhibitor. Analgesic. Investigated for treatment of Alzheimer's disease. Mp 127-129° (133-134°). [α]_D²² -122 (c, 0.60 in EtOH) (-132). pK_a 8.32. Log P 1.11 (uncertain value) (calc). Pharmacol. active isomer.*Hydrochloride*: Mp 254-255° dec.*Hydrobromide*: *Galantamine hydrobromide*, USAN

[1953-04-4]

Mp 246-247° dec.

▶ LD₅₀ (mus, orl) 19 mg/kg. DF8075000*Perchlorate*: Mp 225-227° dec.N-Oxide: *Galanthamine N-oxide*

[134332-50-6]

C₁₇H₂₁NO₄ 303.357Alkaloid from *Lycoris sanguinea* (Amaryllidaceae). Powder. [α]_D²⁵ -122.9 (c, 0.38 in MeOH).Ac: *O-Acetylgalanthamine*

[25650-83-3]

C₁₉H₂₃NO₄ 329.395Trace alkaloid present in bulbs of *Narcissus pseudonarcissus* ssp. *pseudonarcissus* cv. Carlton (Amaryllidaceae). Cryst. (EtOH). Mp 129°. [α]_D²⁵ -61 (c, 1.3 in MeOH).O-(3R-Hydroxybutanoyl): *O-Methylleucotamine*

[82644-83-5]

C₂₁H₂₇NO₅ 373.448Isol. from the leaves and bulbs of *Leucojum aestivum* (Amaryllidaceae). Oil. [α]_D²⁴ -36 (c, 0.75 in CHCl₃). [α]_D²⁵ -49.3 (c, 0.75 in EtOH).N-De-Me: *N-Demethylgalanthamine*

[41303-74-6]

C₁₆H₁₉NO₃ 273.331Alkaloid from the bulbs of *Crinum asiaticum* var. *japonicum* (Amaryllidaceae). Component of the quasi-racemic alkaloid Narcissamine isol. from *Narcissus pseudonarcissus* and *Amaryllis* spp., which was a quasiracemate of Demethylgalanthamine and (+)-Demethyldihydrogalanthamine (see above). Cryst. (EtOAc). Mp 156-158°. [α]_D²³ -62. [α]_D³⁶ -146 (c, 0.28 in CHCl₃).N-De-Me, N-formyl: *N-Formylorgalanthamine*

[109606-37-3]

C₁₇H₁₉NO₄ 301.341Alkaloid from *Narcissus confusus* (Amaryllidaceae). Mp 190-192°. In-corr. called *N-Formylgalanthamine* in the lit.N-De-Me, O,N-di-Ac: *O,N-Diacetyl-N-demethylgalanthamine*

[61948-10-5]

C₂₀H₂₃NO₅ 357.405Alkaloid from the fruit of *Crinum asiaticum* var. *japonicum*. Cryst. (EtOH). Mp 234-235°. *N-Demethylgalanthamine* may be the plant const.

N-De-Me, O-(3ξ-hydroxybutanoyl):

*Leucovernine*C₂₀H₂₅NO₅ 359.421Alkaloid from the bulbs *Leucojum vernum*. Amorph. solid. [α]_D³¹ -42 (c, 1 in CHCl₃). λ_{\max} 209 (log ϵ 3.62); 233 (sh) (log ϵ 3.14); 290 (log ϵ 2.67) (MeOH).

N-De-Me, O-(3ξ-acetoxybutanoyl):

*Acetylleucovernine*C₂₂H₂₇NO₆ 401.458Alkaloid from the bulbs of *Leucojum vernum*. Amorph. solid. λ_{\max} 208 (log ϵ 3.67); 233 (sh) (log ϵ 3.21); 289 (log ϵ 2.69) (MeOH).N-De-Me, N-(2-propenyl): *N-Allylnorgalanthamine*

[112448-56-3]

C₁₉H₂₃NO₃ 313.396Alkaloid from the bulbs of *Lycoris guangxiensis* and leaves of *Leucojum aestivum*. Oil. [α]_D²⁰ -97 (c, 0.31 in MeOH).Me ether: *O-Methylgalanthamine*C₁₈H₂₃NO₃ 301.385Alkaloid from *Chlidanthus fragrans* (Amaryllidaceae). Oil.*Me ether*, N-Me: Mp 272-273° (as iodide). [α]_D -90 (c, 1.0 in MeOH) (as iodide).O-De-Me: *Sanguinine*. *O-Demethylgalanthamine*

[60755-80-8]

C₁₆H₁₉NO₃ 273.331Alkaloid from bulbs of *Lycoris sanguinea* var. *kiushiana* (Amaryllidaceae). Cryst. (EtOH/C₆H₆). Mp 210.5-213° dec. [α]_D²⁷ -133 (c, 0.23 in EtOH).*O-De-Me, perchlorate*: Mp 249-251° dec.*O-De-Me, N-oxide: Sanguinine N-oxide*

[134332-51-7]

C₁₆H₁₉NO₄ 289.33Alkaloid from *Lycoris sanguinea* (Amaryllidaceae). Prisms (MeOH). Mp 191-195°. [α]_D²⁵ -106.1 (c, 0.25 in MeOH).*O-De-Me, O³-Ac: 3-O-Acetylsanguinine*. *3-O-Acetyldemethylchlidanthine*

[125340-13-8]

C₁₈H₂₁NO₄ 315.368Alkaloid from the bulbs of *Crinum kirkii* and *Haemanthus multiflorus*. Needles (MeOH). Mp 215-217°. [α]_D²⁰ -45.8 (c, 0.5 in CHCl₃). Numbering systems differ.*O-De-Me, O⁵-Ac: P11012*

[164723-36-8]

[164723-74-4]

C₁₈H₂₁NO₄ 315.368

Acetylcholinesterase (ACE) inhibitor. CAS no. refers to hydrochloride.

O-De-Me, O³-(3R-hydroxybutanoyl): Leucotamine

[82644-82-4]


C₂₀H₂₅NO₅ 359.421Alkaloid from leaves of *Leucojum aestivum* (Amaryllidaceae). Needles (C₆H₆). Mp 168-171°. [α]_D²⁰ -52.6 (c, 0.74 in CHCl₃).*O-De-Me, O³-(3-hydroxybutanoyl), N-de-Me: Norbutsanguinine*

- [164854-60-8]
 $C_{19}H_{23}NO_5$ 345.394
 From bulbs of *Lycoris sanguinea* (Amaryllidaceae). Oil. $[\alpha]_D^{20}$ -26 (c, 0.1 in $CHCl_3$).
- O-De-Me, O⁶-[(adamantan-1-yl)carboxyl]: P11149**
 [164724-79-2]
 [164724-95-2]
 $C_{27}H_{33}NO_4$ 435.562
 Acetylcholinesterase inhibitor.
 Candidate for the treatment of Alzheimer's disease. CAS no. refers to hydrochloride.
- O-De-Me, O³-Me: Chlidanthine**
 [25650-85-5]
 $C_{17}H_{21}NO_3$ 287.358
 Alkaloid from *Chlidanthus fragrans*, *Haemanthus multiflorus* and *Hippeastrum aulicum* var. *robustum* (Amaryllidaceae). Mp 238-239°. $[\alpha]_D$ -140 (EtOH). The abs. config. is incorrectly given in CAS.
- O-De-Me, O³,N-di-Me:** Mp 263-264° dec.
- O,N-Di-de-Me: Norsanguinine**
 [164854-59-5]
 $C_{15}H_{17}NO_3$ 259.304
 Alkaloid from bulbs of *Lycoris sanguinea* (Amaryllidaceae). Cryst. ($CHCl_3$ /MeOH). Mp 142-144°. $[\alpha]_D^{20}$ -54 (c, 0.1 in MeOH).
- 3-Ketone: Narwedine. Galanthamine. Narvedine**
 [510-77-0]
 $C_{17}H_{19}NO_3$ 285.342
 Alkaloid from *Narcissus cyclamineus*, some other *Narcissus* spp. and from *Galanthus nivalis* (Amaryllidaceae). Biosynth. precursor of Galanthamine. Cryst. (C_6H_6). Mp 189-192°. $[\alpha]_D$ +405 (c, 1.0 in $CHCl_3$).
- 3-Ketone, picrate:** Mp 123°.
- 1,2-Dihydro: Lycoramine. Pseudohomolycorine**
 [21133-52-8]
 $C_{17}H_{23}NO_3$ 289.374
 Alkaloid from *Lycoris radiata*, *Crinum powellii*, and some other spp. in the Amaryllidaceae. Mp 120-121°. $[\alpha]_D^{27}$ -98 (EtOH).
- 1,2-Dihydro; perchlorate:** Mp 138-139°.
- 1,2-Dihydro, N-oxide: Lycoramine N-oxide**
 [134306-21-1]
 $C_{17}H_{23}NO_4$ 305.373
 Alkaloid from *Lycoris sanguinea* (Amaryllidaceae). Powder. $[\alpha]_D$ -91.8 (MeOH).
- 1,2-Dihydro, N-de-Me: N-Demethyllycoramine. N-Demethyldihydrogalanthamine**
 [41432-21-7]
 [41303-53-1]
 $C_{16}H_{21}NO_3$ 275.347
 Alkaloid from the bulbs of *Hymenocallis rotata* (Amaryllidaceae). Prisms + $1H_2O$ (Me_2CO). Mp 123-124°. $[\alpha]_D^{29}$ -39.1 (c, 0.95 in $CHCl_3$).
- 1,2-Dihydro, O-Ac: Acetyllycoramine**
 [53375-68-1]
 $C_{19}H_{25}NO_4$ 331.411
 Alkaloid from *Pancreatum maritimum*. Needles (MeOH). Mp 93-95°. $[\alpha]_D^{22}$ -87 (c, 0.1 in MeOH).
- 1,2-Dihydro, O-(3S-hydroxybutanoyl): Pancratamine**
 [243986-37-0]
 $C_{21}H_{29}NO_5$ 375.464
 Alkaloid from *Pancreatum maritimum*. Oil. $[\alpha]_D^{22}$ -62.9 (c, 0.7 in MeOH).
- 1,2-Dihydro, O-de-Me: O-Demethyllycoramine. O-Demethyldihydrogalanthamine**
 [77754-93-9]
 $C_{16}H_{21}NO_3$ 275.347
 Alkaloid from the bulbs of *Lycoris radiata* (Amaryllidaceae). Prisms (C_6H_6 /EtOH). Mp 204-207°.
- 3-Epimer: Epigalanthamine**
 [1668-85-5]
 $C_{17}H_{21}NO_3$ 287.358
 Alkaloid from the bulbs of *Lycoris squamigera* (Amaryllidaceae). Mp 190°. $[\alpha]_D$ -222 (MeOH).
- 3-Epimer, N-de-Me: Epinorgalanthamine**
 [156040-03-8]
 $C_{16}H_{19}NO_3$ 273.331
 Alkaloid from whole plants of *Narcissus leonensis* (Amaryllidaceae). Mp 150-152°. $[\alpha]_D^{25}$ -62 (c, 0.73 in $CHCl_3$).
- 3-Epimer, 1,2-dihydro, N-de-Me: Epinorlycoramine**
 [156128-69-7]
 $C_{16}H_{21}NO_3$ 275.347
 Alkaloid from whole plants of *Narcissus leonensis* (Amaryllidaceae). $[\alpha]_D^{25}$ +79.1 (c, 0.51 in $CHCl_3$).
- (±)-form [23173-12-8]**
 Synthetic. Mp 121-123° (117-118°).
Hydrochloride:
 Cryst. + $\frac{1}{2}H_2O$ (EtOH). Mp 252-254°.
- N-De-Me, O,N-di-Ac:**
 Cryst. (EtOAc). Mp 204-205°.
- 3-Ketone:** [1668-86-6]
 Synthetic or by racemisation of alkaloid. Also occurs naturally in *Narcissus kristalli* and *Ungernia victoris* (Amaryllidaceae). Mp 187-190°.
- 1,2-Dihydro:** [18747-70-1]
 Mp 98-99°.
- 3-Epimer:** [23173-13-9]
 Synthetic. Needles (EtOH). Mp 199°.
- 3-Epimer, hydrochloride:**
 Cryst. (EtOH). Mp 268-269°.
- Döpke, W. et al., *Chem. Ber.*, 1956, **89**, 1129-1134 (*Chlidanthine, isol*)
 Kobayashi, S. et al., *Chem. Ind. (London)*, 1956, 177-178 (*struct*)
 Boit, H.G. et al., *Chem. Ber.*, 1957, **90**, 2197-2202 (*Narwedine, isol, struct*)
 Barton, D.H.R. et al., *J.C.S.*, 1962, 806-817 (*Narwedine, Epigalanthamine, struct, ir, synth*)
 Laiho, S.M. et al., *J.A.C.S.*, 1964, **86**, 4434-4438 (*N-Demethylgalanthamine, Demethyldihydrogalanthamine, Narcissamine*)
 Williams, D.J. et al., *Proc. Chem. Soc., London*, 1964, 357 (*cryst struct*)
 Hung, S.H. et al., *Yaouxue Xuebao*, 1964, **11**, 1-14; *CA*, **61**, 3154e (*Epigalanthamine, isol*)
 Abdiazimov, K.A. et al., *Khim. Prir. Soedin.*, 1967, **3**, 64-65; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 54-55 (*Narwedine, isol*)
 Kametani, T. et al., *J.C.S. (C)*, 1969, 2602-2605 (*Narwedine, synth, ir, pmr, ms*)
 DeAngelis, G.G. et al., *Tet. Lett.*, 1969, 729-732 (*Epigalanthamine, ord, cd*)
 Bhandarkar, J.G. et al., *J.C.S. (C)*, 1970, 1224-1227 (*Chlidanthine, uv, ir, pmr, ms, struct, biosynth*)
 Kametani, T. et al., *J.O.C.*, 1971, **36**, 1295-1297 (*synth*)
 Nogueiras, C. et al., *Tet. Lett.*, 1971, 3249-3250 (*O-Methylgalanthamine*)
 Kametani, T. et al., *J. Het. Chem.*, 1973, **10**, 35-37 (*N-Demethyldihydrogalanthamine*)
 Zetta, L. et al., *J.C.S. Perkin 2*, 1973, 1180-1184 (*cmr*)
 Fuganti, C. et al., *Tet. Lett.*, 1974, 2261-2264 (*biosynth*)
 Ochi, M. et al., *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3363-3364 (*O,N-Diacetyl-N-demethylgalanthamine*)
 Kobayashi, S. et al., *Chem. Pharm. Bull.*, 1976, **24**, 1537-1543; 2553-2555; 1980, **28**, 3433-3436; 1985, **33**, 5258-5263 (*Sanguinine, N-Demethylgalanthamine, O-Demethyllycoramine, Leucotamine, O-Methylleucotamine*)
 Kametani, T. et al., *Heterocycles*, 1976, **4**, 1111-1114 (*resoln*)
 Shimizu, K. et al., *Heterocycles*, 1977, **8**, 277-282 (*synth*)
 Schultz, A.G. et al., *J.A.C.S.*, 1977, **99**, 8065-8067 (*Lycoramine, synth*)
 Sánchez, I.H. et al., *J.O.C.*, 1984, **49**, 157-163 (*Lycoramine, synth*)
 Westra, P. et al., *Br. J. Anaesth.*, 1986, **58**, 1303 (*pharmacol*)
 Mikhailova, D. et al., *Pharmacology*, 1986, **32**, 301 (*pharmacol*)
 Martin, S.F. et al., *Alkaloids (Academic Press)*, 1987, **30**, 251-376 (*rev*)
 Kihara, M. et al., *Chem. Pharm. Bull.*, 1987, **35**, 1070-1075 (*N-Demethyllycoramine*)
 Teneheva, J. et al., *J. Chromatogr.*, 1987, **421**, 396-400 (*hplc*)
 Ackland, D.J. et al., *J.C.S. Perkin 1*, 1987, 2695-2700 (*Lycoramine, synth*)
 Bastida, J. et al., *Phytochemistry*, 1987, **26**, 1519-1524; 1993, **34**, 1656-1658 (*N-Formylnorgalanthamine, Epinorlycoramine, Epinorgalanthamine*)
 Li, H.-Y. et al., *Planta Med.*, 1987, **53**, 259-261 (*N-Allylnorgalanthamine*)
 Szweczyk, J. et al., *J. Het. Chem.*, 1988, **25**, 1809-1811; 1995, **32**, 195-199 (*synth, ir, pmr, resoln, Epigalanthamine*)
 Vlahov, R. et al., *Tetrahedron*, 1989, **45**, 3329-3345 (*Epigalanthamine, synth*)
 Carroll, P. et al., *Bull. Soc. Chim. Fr.*, 1990, 769-780 (*ir, pmr, cmr, cryst struct*)
 Kobayashi, S. et al., *Phytochemistry*, 1991, **30**, 675-677 (*N-oxides*)
 Parker, K.A. et al., *J.O.C.*, 1992, **57**, 752-755 (*synth, Lycoramine*)
 Ishizaki, M. et al., *J.O.C.*, 1993, **58**, 3877-3885 (*synth, Lycoramine*)
 Shieh, W.-C. et al., *J.O.C.*, 1994, **59**, 5463-5465 (*Narwedine, synth, resoln, bibl*)
Eur. Pat., 1995, ((Hoechst-Roussel))649 846; 653 427; *CA*, **123**, 56362v; 56364x (*P11149*)
 Harvey, A. et al., *Pharmacol. Ther.*, 1995, **68**, 113-128 (*pharmacol, rev*)
 Abdallah, O.M. et al., *Phytochemistry*, 1995, **39**, 477-478 (*Norsanguinine, Norbutanguinine*)

- Kreh, M. *et al.*, *Phytochemistry*, 1995, **40**, 1303-1306 (*O*-Acetylgalanthamine)
- Bores, G.M. *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, **277**, 728-738 (*pharmacol*, P11149, P11012)
- Peeters, O.M. *et al.*, *Acta Cryst. C*, 1997, **53**, 1284-1286 (*cryst struct*, bromide)
- Eichhorn, J. *et al.*, *Phytochemistry*, 1998, **49**, 1037-1047 (*biosynth*, *cmr*)
- Youssef, D.T. *et al.*, *Planta Med.*, 1998, **64**, 669-670 (*isol*, *pmr*, *cmr*)
- Czollner, L. *et al.*, *Tet. Lett.*, 1998, **39**, 2087-2088 (*synth*)
- Martindale, The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1393
- Kuenburg, B. *et al.*, *Org. Process Res. Dev.*, 1999, **3**, 425-431 (*Narwedine*, *synth*, *resoln*)
- Youssef, D.T.A. *et al.*, *Pharmazie*, 1999, **54**, 535-537 (*Pancritamine*, *Acetyllycoramine*)
- Gras, E. *et al.*, *Tet. Lett.*, 1999, **40**, 9243-9244 (*synth*)
- Scott, L.J. *et al.*, *Drugs*, 2000, **60**, 1095-1122 (*rev*)
- Krikorian, D. *et al.*, *Synth. Commun.*, 2000, **30**, 2833-2846 (*synth*, *ir*, *pmr*, *cmr*)
- Node, M. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 3060-3062 (*Narwedine*, *synth*)
- Guillou, C. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 4745-4746 (*synth*)
- Linnemann, E. *et al.*, *J. Labelled Compd. Radiopharm.*, 2001, **44**, 661-669 (*synth*)
- Treu, M. *et al.*, *Molecules*, 2001, **6**, M274 (*Norsanguinine*, *synth*)
- Fan, C.-A. *et al.*, *Org. Lett.*, 2004, **4691**, 4694 (*synth*)
- Machochi, A.K. *et al.*, *Phytochemistry*, 2004, **65**, 3143-3149 (*3-O-Acetylsanguinine*)
- Forgo, P. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1588-1591 (*Leucovernine*, *Acetylleucovernine*)
- Trost, B.M. *et al.*, *J.A.C.S.*, 2005, **127**, 14785-14803 (*synth*)
- Node, M. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 1662-1679 (*synth*)
- Hu, X.-D. *et al.*, *Org. Lett.*, 2006, **8**, 1823-1825 (*synth*)
- Malachowski, W.P. *et al.*, *J.O.C.*, 2007, **72**, 6792-6796 (*Lycoramine*, *synth*)
- Satcharoen, V. *et al.*, *Org. Lett.*, 2007, **9**, 1867-1869 (*synth*)
- Tanimoto, H. *et al.*, *Tet. Lett.*, 2007, **48**, 6267-6270 (*synth*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GBA000

Galanthindole

G-4




$C_{17}H_{15}NO_3$ 281.31
Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. λ_{\max} 220 (log ϵ 4.21); 290 (log ϵ 3.83); 298 (sh) (log ϵ 3.8) (MeOH).

Unver, N. *et al.*, *Planta Med.*, 2003, **69**, 869-871 (*isol*, *pmr*, *cmr*)

Galanthusine

[31278-95-2]




$C_{18}H_{23}NO_5$ 333.383
Alkaloid from *Galanthus caucasicus* (Amaryllidaceae). Cryst. (Me₂CO). Mp 118-119°. $[\alpha]_D^{30}$ -66.6 (c, 0.42 in EtOH).

Tsakadze, D.M. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 773; *Chem. Nat. Compd. (Engl. Transl.)*, 791

Galantini

G-6

D-Ornithinegalantini
[78330-65-1]



$C_{41}H_{84}N_{14}O_{13}$ 981.201
Struct. revised in 1992. Isol. from *Bacillus puvifaciens*. Active against gram-negative and -positive bacteria and acid fast bacteria. Sol. H₂O, MeOH; poorly sol. Me₂CO, hexane.
▶ LD₅₀ (mus, ipr) 50 mg/kg. LW5965000 [55599-74-1]

Shoji, J. *et al.*, *J. Antibiot.*, 1975, **28**, 122
Sakai, N. *et al.*, *J.A.C.S.*, 1992, **114**, 998 (*synth*, *struct*)

Galantini

G-7


D-Lysinegalantini
[78330-64-0]

As Galantini I, G-6 with
R = CH₂CH₂CH₂NH₂
 $C_{42}H_{86}N_{14}O_{13}$ 995.227
Prod. by *Bacillus puvifaciens*. Powder. Sol. H₂O, EtOH, MeOH; poorly sol. Me₂CO, hexane.

Shoji, J. *et al.*, *J. Antibiot.*, 1975, **28**, 122 (*isol*)
Sakai, N. *et al.*, *J.A.C.S.*, 1992, **114**, 998 (*synth*, *struct*)

Galasine

[167568-94-7]



$C_{19}H_{23}NO_7$ 377.393
Alkaloid from whole plants of *Galanthus*

G-5


elwesii (Amaryllidaceae). Prisms (MeOH). Mp 284.9-286.3°. Opt. rotn. not measured due to paucity of material.

Latvala, A. *et al.*, *Phytochemistry*, 1995, **39**, 1229 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cd*, *cryst struct*)

Galipinine

G-9

2-[2-(1,3-Benzodioxol-5-yl)ethyl]-1,2,3,4-tetrahydro-1-methylquinoline. 1,2,3,4-Tetrahydro-2-(3,4-methylenedioxyphenethyl)-1-methylquinoline. Allocuspareine



$C_{19}H_{21}NO_2$ 295.38
Identical structs. assigned to Galipinine and Allocuspareine. Alkaloid from *Galipea officinalis*. Shows activity against *Mycobacterium tuberculosis*. Oil or pale yellow powder. $[\alpha]_D$ -33.4 (c, 0.005 in CHCl₃). λ_{\max} 259 (log ϵ 3.98); 311 (log ϵ 4.41) (CHCl₃) (Galipinine). λ_{\max} 204; 253; 290; 315 (MeOH) (Allocuspareine).
Rakotoson, J.H. *et al.*, *Planta Med.*, 1998, **64**, 762-763 (*isol*, *uv*, *pmr*, *cmr*, *activity*)
Houghton, P.J. *et al.*, *Planta Med.*, 1999, **65**, 250 (*Allocuspareine*)
O'Byrne, A. *et al.*, *Tetrahedron*, 2008, **64**, 8067-8072 (*synth*, *abs config*)

Galipoidine

G-10


[1399-42-4]

$C_{19}H_{15}NO_4$ 321.332
Struct. unknown. Alkaloid from angostura bark (*Cusparia trifoliata*, preferred genus name *Angostura*) (Rutaceae). Needles (EtOH). Mp 233°.
Tröger, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1911, **249**, 174-208

Galipoline

G-11

2-[2-(3,4-Dimethoxyphenyl)ethyl]-4-hydroxyquinoline



$C_{19}H_{19}NO_3$ 309.364
Alkaloid from the bark of angostura (*Galipea officinalis*) (Rutaceae). Cryst. (Et₂O). Mp 193°.

Me ether: Galipine

[525-68-8]
 $C_{20}H_{21}NO_3$ 323.391
Occurs in bark of *Galipea officinalis* (Rutaceae). Antispasmodic agent. Active against *Mycobacterium tuberculosis*. Cryst. (Et₂O). Mp 115-116°. Log P 4.35 (calc).

Me ether, hydrochloride: Mp 165°.


Me ether, picrate: Mp 194°.

Späth, E. *et al.*, *Ber.*, 1924, **57**, 1687 (*isol*, *synth*, *Galipine*)

- Späth, E. *et al.*, *Monatsh. Chem.*, 1929, **52**, 129 (isol, synth, Galipoline)
 Rakotoson, J.H. *et al.*, *Planta Med.*, 1998, **64**, 762-763 (isol, uv, pmr, cmr, Galipine)
 Houghton, P.J. *et al.*, *Planta Med.*, 1999, **65**, 250-254 (isol, pmr, cmr, Galipine, activity)

Gambirine G-13

18,19-Dihydro-9-hydroxycorynantheine
 [29472-77-3]



C₂₂H₂₈N₂O₄ 384.474
 Alkaloid from the leaves of *Uncaria gambier*, *Uncaria callophylla* and *Neonauclea schlechteri* (Rubiaceae). Cryst. (CHCl₃/hexane). Mp 163-165° dec. [α]_D²⁰ +28.6 (c, 0.077 in CHCl₃). λ_{max} 226 (ε 49800); 250 (sh) (ε 14650); 285 (ε 6050); 294 (ε 5550) (95% EtOH).

Me ether: Speciogynine

[4697-67-0]
 C₂₃H₃₀N₂O₄ 398.501
 Alkaloid from *Mitragyna speciosa* (Rubiaceae). Cryst. (Et₂O). Mp 214°. [α]_D^{23.5} +28.4 (c, 0.26 in CHCl₃).

Me ether, picrate:

Orange-red rosettes (EtOH). Mp 224°.

18,19-Didehydro: Gambireine. 9-Hydro-

xcorynantheine
 [142905-21-3]
 C₂₂H₂₆N₂O₄ 382.458
 Alkaloid from leaves of *Uncaria callophylla* (Rubiaceae). λ_{max} 227; 250 (sh); 286; 295 (EtOH).

18,19-Didehydro, Me ether: Paynantheine.

9-Methoxycorynantheine
 [4697-66-9]
 C₂₃H₂₈N₂O₄ 396.485
 Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae). Yellow solid. Mp 98° (softens). [α]_D²³ -28.9 (c, 0.27 in CHCl₃). λ_{max} 227 (log ε 4.63); 272 (sh) (log ε 3.95); 283 (sh) (log ε 3.86); 292 (log ε 3.84) (EtOH).

18,19-Didehydro, Me ether, picrate:

Orange needles (EtOH). Mp 202°.

3-Epimer, 18,19-didehydro, Me ether: 3-Isopaynantheine

[22032-51-5]
 C₂₃H₂₈N₂O₄ 396.485
 Alkaloid from *Mitragyna speciosa* (Rubiaceae). No phys. props. recorded.

3,20-Diepimer, Me ether: Speciociliatine

[14382-79-7]
 C₂₃H₃₀N₂O₄ 398.501
 Alkaloid from *Mitragyna speciosa*, also detected in *Uncaria* spp. (Rubiaceae). Antitussive, analgesic. Amorph. Mp 97-98°. [α]_D²⁴ -21 (c, 0.24 in CHCl₃). Log P 3.93 (calc).

3,20-Diepimer; Me ether, methiodide:
 Pale-cream microcryst. (EtOH). Mp 264°.

Beckett, A.H. *et al.*, *J. Pharm. Pharmacol.*, 1965, **17**, 753-755 (Paynantheine, uv, ir, pmr, struct)

Beckett, A.H. *et al.*, *Planta Med.*, 1966, **14**, 277-288 (Speciociliatine, Speciogynine, isol, uv, ir, pmr, struct)

Merlini, L. *et al.*, *Tet. Lett.*, 1967, 1571-1574 (isol, uv, ir, pmr, ms, cd, ord, struct, abs config)

Lee, C.M. *et al.*, *Tetrahedron*, 1967, **23**, 375-385 (Speciogynine, Speciociliatine, ord, cd, pmr, ir, uv, abs config)

Trager, W.F. *et al.*, *Tetrahedron*, 1967, **23**, 1043-1047 (Paynantheine, uv, ord, cd, abs config)

Shellard, E.J. *et al.*, *Planta Med.*, 1978, **34**, 253-263 (3-Isopaynantheine)

Goh, S.H. *et al.*, *Phytochemistry*, 1985, **24**, 880-881 (isol, uv, ir, pmr, cmr, ms)


Kam, T.S. *et al.*, *Phytochemistry*, 1992, **31**, 2031-2034 (Gambireine)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 916-928 (rev)

Kitajima, M. *et al.*, *J. Nat. Med. (Tokyo)*, 2006, **60**, 28-35 (Speciogynine, Speciociliatine, 3-Isopaynantheine, pmr, cmr)

Gandharamine G-14

1-(4-Hydroxybenzoyl)-6,7-dimethoxy-2-methylisoquinolinium, 9CI
 [81509-33-3]



C₁₉H₁₈NO₄⁺ 324.355

Quaternary alkaloid from *Berberis baluchistanica* and *Thalictrum fendleri* (Berberidaceae, Ranunculaceae). Amorph. solid (as chloride).

O⁷,N-Di-de-Me: 7-Hydroxy-1-(4-hydroxybenzoyl)-6-methoxyisoquinoline
 C₁₇H₁₃NO₄ 295.294


Alkaloid from the roots of *Menispermum dauricum*. Red needles. Mp 230-232°. λ_{max} 233; 252; 276; 297; 342 (no solvent reported).

Zarga, M.H.A. *et al.*, *Heterocycles*, 1982, **18**, 63 (uv, pmr, ms, struct, synth)

Zhang, X. *et al.*, *Phytochemistry*, 2004, **65**, 929-932 (di-de-Me)

Ganoderpurine G-15

4-(6-Amino-9H-purin-9-yl)-4-methyl-2-pentanone, 9CI. N⁹-(1,1-Dimethyl-3-oxobutyl)adenine
 [133086-82-5]




C₁₁H₁₅N₅O 233.272

Alkaloid from the mycelium of *Ganoderma capense*. Mp 151-152°.

Yu, J.G. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 612; *CA*, **114**, 160668z

Garceine G-16

1,2,4-Triheptylpyrazolidine, 9CI. 1,2,4-Triheptyltetrahydropyrazole
 [340829-54-1]




C₂₄H₅₀N₂ 366.672

Alkaloid from *Lotus garcinii*. Powder. Mp 60°.

Ali, M.S. *et al.*, *Turk. J. Chem.*, 2001, **25**, 107-112 (isol, pmr, cmr)

Gardenamide A G-17

[214125-00-5]



Absolute Configuration

C₁₁H₁₃NO₄ 223.228

Constit. of *Gardenia jasminoides* and *Rothmannia urcelliformis*. Amorph. powder. [α]_D +404.4 (c, 0.25 in MeOH). λ_{max} 242 (log ε 4.07) (MeOH).

N-(2-Hydroxyethyl): N-(2-Hydroxyethyl)gardenamide A

[284045-03-0]
 C₁₃H₁₇NO₅ 267.281
 Constit. of *Gardenia jasminoides* and other *Gardenia* spp. Amorph. powder. [α]_D²⁶ +325 (c, 0.16 in MeOH).

N-(3-Carboxypropyl): [936548-28-6]

C₁₅H₁₉NO₆ 309.318
 Constit. of *Gardenia volkensii*. Solid. Mp 84-86°.

Machida, K. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1295-1300 (isol, pmr, cmr)


Machida, K. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 746-748 (2-Hydroxyethylgardenamide A)

Bringmann, G. *et al.*, *Eur. J. Org. Chem.*, 2001, 1983-1987 (isol, synth, cd, pmr, cmr, abs config)

Juma, B.F. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 121-125 (*Gardenia volkensii* constit)

Gardfloramine, 9CI G-18

[56198-74-4]



R¹, R² = -OCH₂O-, R³ = H

Absolute Configuration

C₂₂H₂₄N₂O₅ 396.442
Alkaloid from *Gardneria multiflora* (Loganiaceae). Prisms + ½ H₂O (Me₂CO). Mp 159-160°. [α]_D -248.

18-Demethoxy, 19E-isomer: (19E)-18-Demethoxygardfloramine
[56198-75-5]

C₂₁H₂₂N₂O₄ 366.416
Alkaloid from *Gardneria multiflora* (Loganiaceae). Amorph.


Sakai, S. et al., *Tet. Lett.*, 1975, 715 (uv, ir, pmr, ms)

Sakai, S. et al., *Chem. Pharm. Bull.*, 1987, 35, 453 (cmr, cryst struct)

Gardmultine

G-19

[56197-32-1]



C₄₅H₅₄N₄O₁₀ 810.942
Alkaloid from *Gardneria multiflora* (Loganiaceae). Cryst. + 0.5H₂O. Mp 283-285°.

Sakai, S. et al., *Tet. Lett.*, 1975, 719 (uv, ir, pmr, ms, struct)

Gardneramine

G-20

[34274-91-4]

As Gardfloramine, G-18 with

R¹ = R³ = OMe, R² = H

C₂₃H₂₈N₂O₅ 412.485
Alkaloid from *Gardneria nutans* (Loganiaceae). Prisms (Et₂O). Mp 134-135°. [α]_D -288.

▶ LX3090000

Cyanobromide: Mp 214°.

N^d-Oxide: **Gardneramine N^d-oxide**

[64494-86-6]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Gardneria multiflora* (Loganiaceae).

O¹⁸-De-Me: **O¹⁸-Demethylgardneramine**

[32975-55-6]

C₂₂H₂₆N₂O₅ 398.458

Alkaloid from *Gardneria multiflora*.

18-Demethoxy, 19E-isomer: (19E)-18-

Demethoxygardneramine

[56246-54-9]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Gardneria multiflora* (Loganiaceae). Cryst. (Me₂CO). Mp 160-161.5°. [α]_D -147.

Aimi, N. et al., *Tet. Lett.*, 1971, 2061 (cryst struct)

Sakai, S. et al., *Chem. Pharm. Bull.*, 1975, 23, 2805 (uv, ir, pmr, ms, struct)


Sakai, S. et al., *Tet. Lett.*, 1975, 715 (uv, ir, pmr, config, struct, deriv)

Aimi, N. et al., *Chem. Pharm. Bull.*, 1978, 26, 3444 (cmr)

Gardneramine oxindole, 9CI

G-21

[32975-50-1]



Absolute Configuration

C₂₃H₃₀N₂O₆ 430.5

Obt. by hydrol. of Gardneramine, G-20 with HCOOH aq. Foam.

O¹⁸-De-Me: **O¹⁸-Demethylgardneramine oxindole, 9CI. Alkaloid M**

[64494-82-2]

C₂₂H₂₈N₂O₆ 416.473

Alkaloid from *Gardneria multiflora* (Loganiaceae).

17-Deoxy, 16,17-didehydro: **16,17-Didehydro-17-deoxygardneramine oxindole, 9CI**

[64494-85-5]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Gardneria multiflora* (Loganiaceae).

16-Hydroxy: **16-Hydroxygardneramine oxindole, 9CI. Alkaloid N**

[64530-48-9]

C₂₃H₃₀N₂O₇ 446.499

Alkaloid from *Gardneria multiflora* (Loganiaceae). Mp 115-119°. 19-Config. not detd.

18-Demethoxy, 16-hydroxy: **Chitosenine.**

18-Demethoxy-16-hydroxygardneramine oxindole

[56210-06-1]

C₂₂H₂₈N₂O₆ 416.473

Alkaloid from *Gardneria multiflora* (Loganiaceae). Amorph. Mp 173-174° (as hydrochloride). 19-Config. incorrectly given as Z- in CAS.

16-Epimer: **16R-Gardneramine oxindole, 9CI. Alkaloid J†**

[64550-59-0]

C₂₃H₃₀N₂O₆ 430.5

Alkaloid from *Gardneria multiflora* (Loganiaceae). Mp 124-128°.

16-Epimer, O¹⁸-de-Me: **16R-O¹⁸-Demethylgardneramine oxindole, 9CI. Alkaloid L†**

[64550-58-9]

C₂₂H₂₈N₂O₆ 416.473

Alkaloid from *Gardneria multiflora* (Loganiaceae). Mp 148-152°.

Shinichiro, S. et al., *Chem. Pharm. Bull.*, 1975, 23, 2805-2817 (synth, uv, ir, pmr, struct)

Sakai, S. et al., *Tet. Lett.*, 1975, 715-718 (Chitosenine, ir, uv, pmr, ms, struct, abs config)


Sakai, S. et al., *Yakugaku Zasshi*, 1977, 97, 399-409; *CA*, 88, 23222e (*Gardneria multiflora* constits)

Aimi, N. et al., *Chem. Pharm. Bull.*, 1978, 26, 3444-3449 (Chitosenine, cmr, config)

Gardnerine

G-22

11-Methoxysarpagan-17-ol, 9CI
[23172-92-1]



C₂₀H₂₄N₂O₂ 324.422

Alkaloid from the roots and stems of *Gardneria nutans* (Apocynaceae). Shows ganglion-blocking activity. Mp 243-244°. [α]_D²⁵ -29.4. pK_a 7.36. λ_{max} 229 (log ε 4.56); 269 (log ε 3.7); 298 (log ε 3.77) (MeOH).

▶ VQ5835500

Ac:

Prisms (MeOH aq.). Mp 218-219°.

19,20-ξ-Dihydro, 17-carboxylic acid, N^l-Me, Me ester: **11-Methoxy-N-methyl-dihydropericyclivine**

[135574-53-7]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Tabernaemontana divaricata* (Apocynaceae). λ_{max} 235; 270; 292 (MeOH).

Sakai, S. et al., *Chem. Pharm. Bull.*, 1973, 21, 1783-1798 (uv, ir, pmr, ms, struct, abs config)

Aimi, N. et al., *Chem. Pharm. Bull.*, 1978, 26, 3444-3449 (cmr)


Arambewela, L.S.R. et al., *Phytochemistry*, 1991, 30, 1740-1741 (11-Methoxy-N-methyl-dihydropericyclivine)

Zhou, H. et al., *Tet. Lett.*, 2005, 46, 4219-4224 (synth)

Gardnutine

G-23

6,17-Epoxy-11-methoxysarpagan, 9CI
[23172-98-7]



C₂₀H₂₂N₂O₂ 322.406

Alkaloid from the roots and stems of *Gardneria nutans* (Loganiaceae). Shows ganglion-blocking activity. Mp 319-320°. [α]_D²⁵ +30.3 (solvent not reported). pK_a 5.77.

18-Hydroxy: **Hydroxygardnutine**

[23173-00-4]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the roots and stem of *Gardneria nutans* (Loganiaceae).

Shows ganglion-blocking activity. Mp 311-312°. $[\alpha]_D^{25} +36.2$ (solvent not reported). pK_a 5.64.

Demethoxy, N¹-Me-16-Epidehydroaffinisine. Dehydro-16-epiaffinisine [138989-36-3]

$C_{20}H_{22}N_2O$ 306.407

Alkaloid from leaves and root bark of *Ervatamia hirta* (Apocynaceae). $[\alpha]_D +58.8$ (c, 0.25 in MeOH). λ_{max} 224 (log ϵ 4.37); 281 (log ϵ 3.65); 290 (sh) (log ϵ 3.58) (MeOH).

Sakai, S. et al., *Chem. Pharm. Bull.*, 1973, **21**, 1783-1798 (*Gardmutine, Hydroxygardmutine*)

Clivio, P. et al., *Phytochemistry*, 1991, **30**, 3785-3792 (*16-Epidehydroaffinisine*)


Yu, J. et al., *J.O.C.*, 2003, **68**, 5852-5859 (*16-Epidehydroaffinisine, synth*)

Zhou, H. et al., *Tet. Lett.*, 2005, **46**, 4219-4224 (*synth*)

Gardquinolone

G-24

[156430-98-7]



$C_{20}H_{22}N_2O_3$ 338.405


Prob. a ring-expanded indole alkaloid (sarpagine type). Alkaloid from leaves of *Gardneria nutans* (Loganiaceae). Powder (EtOAc). Mp 265-273° dec.

Takayama, H. et al., *J. Nat. Prod.*, 1994, **57**, 521-523 (*isol, uv, pmr, cmr, ms, cd, struct*)

Garryine, 9CI

G-25

[561-51-3]



Absolute Configuration

$C_{22}H_{33}NO_2$ 343.508

Alkaloid from the bark of *Garrya veatchii* (Garryaceae). Mp 74-82° (hydrate). $[\alpha]_D^{28} -84$ (c, 1.4 in EtOH).

Hydrochloride: Mp 251-252° dec., 263-268°.

16 α ,17-Dihydro, 15-ketone: Isocuauchichicine [467-93-6]

[467-93-6]

Cryst. (MeOH). Mp 134-136°. $[\alpha]_D -84$ (CHCl₃). pK_a 8.1.

15-Epimer: Isogarryfoline. Isolaurifoline [467-92-5]

Obt. by rearr. of Garryfoline in hot

MeOH. Cryst. (Me₂CO or hexane).

Mp 140-144°. $[\alpha]_D -57$ (CHCl₃). pK_a 8.6.

Oneto, J.F. et al., *J. Am. Pharm. Assoc.*, 1946, **35**, 204-207 (*isol*)

Weisner, K. et al., *Can. J. Chem.*, 1952, **30**, 608-626 (*isol, struct, ir*)

Wiesner, K. et al., *J.A.C.S.*, 1954, **76**, 6068-6073 (*isol, struct, ir*)

Djerassi, C. et al., *J.A.C.S.*, 1955, **77**, 4801-4807; 6633 (*Garryine, Isogarryfoline, Isocuauchichicine, synth, ir, nomencl*)

Verbrueggen, H. et al., *J.A.C.S.*, 1962, **84**, 2990-2997 (*abs config*)

Masamune, S. et al., *J.A.C.S.*, 1964, **86**, 290-291 (*synth*)

Nagata, W. et al., *J.A.C.S.*, 1967, **89**, 1499-1504 (*synth*)

Pelletier, S.W. et al., *Tetrahedron*, 1968, **24**, 2019-2038 (*pmr*)

Pelletier, S.W. et al., *J.A.C.S.*, 1977, **99**, 284-286 (*Garryine, Isocuauchichicine, cmr*)


Mody, N.V. et al., *Tetrahedron*, 1978, **34**, 2421-2431 (*cmr*)

Pelletier, S.W. et al., *J. Nat. Prod.*, 1980, **43**, 41-71 (*rev*)

Pelletier, S.W. et al., *J.O.C.*, 1981, **46**, 1840-1846 (*cmr*)

Gasabiimine

G-26



$C_{34}H_{26}N_2O_5$ 542.59


Alkaloid from the roots of *Triclisia sacleuxii*. λ_{max} 205 (log ϵ 4.61); 234 (log ϵ 4.47); 287 (log ϵ 4); 333 (log ϵ 3.59) (MeOH).

Murebwayire, S. et al., *Fitoterapia*, 2006, **77**, 615-617 (*isol, pmr, cmr*)

Gaultherialine A

G-27

[956261-55-5]



$C_{21}H_{19}NO_5$ 365.385


Alkaloid from *Gaultheria nummularioides*. Amorph. red powder. Mp 170-172°. λ_{max} 268 (log ϵ 3.71); 314 (log ϵ 4.03); 380 (log ϵ 3.96); 446 (log ϵ 3.75); 464 (log ϵ 3.76); 514 (log ϵ 3.73) (CHCl₃).

Yang, M.-F. et al., *J. Asian Nat. Prod. Res.*, 2007, **9**, 183-186 (*isol, pmr, cmr*)

GB4 Toxin

G-28

2-(1-Methyl-2-oxopropylidene)phosphorohydrazidothioate oxime [82638-81-1]




$C_{10}H_{22}N_3O_3PS$ 295.342

Isol. from the Florida red tide dinoflagellate *Gymnodinium breve* (*Ptychodiscus brevis*). Ichthyotoxin. Needles (C₆H₆). Mp 82-83°. λ_{max} 230 (solvent not reported) (Derep).

Alam, M. et al., *J.A.C.S.*, 1982, **104**, 5232 (*isol, cryst struct, ms, ir, pmr*)

Geibalansine

G-29



(R)-form

$C_{15}H_{17}NO_3$ 259.304

(R)-form [316172-98-2]

Alkaloid from the stem bark of *Zanthoxylum hyemale*. Needles. Mp 181-182°. $[\alpha]_D^{25} -14$ (c, 0.06 in MeOH).

8,9-Dimethoxy: [366490-63-3]

Alkaloid from *Melicope semecarpifolia*. $[\alpha]_D^{23} -18.6$ (c, 0.06 in CHCl₃).

(S)-form [72855-84-6]

Alkaloid from the leaves of *Geijera balansae* (Rutaceae). Cryst. (MeOH/2-propanol). Mp 181-182°. $[\alpha]_D +12$ (c, 0.86 in MeOH).

Ac: O-Acetylgeibalansine

[72855-85-7]

$C_{17}H_{19}NO_4$ 301.341

Alkaloid from the leaves of *Geijera balansae* (Rutaceae). Amorph. $[\alpha]_D +58$ (c, 1.2 in CHCl₃).

N-Me: N-Methylgeibalansine. Tabouensinium

$C_{16}H_{20}NO_3^{\oplus}$ 274.339

Quaternary alkaloid from the stem bark of *Araliopsis tabouensis*. Powder (as chloride). Mp 136-137° (chloride). $[\alpha]_D -17.2$ (c, 1.06 in DMSO) (chloride). λ_{max} 229 (log ϵ 4.86); 274 (log ϵ 4.19); 283 (log ϵ 4.16); 325 (log ϵ 4.17); 337 (log ϵ 4.03) (MeOH) (chloride).

8,9-Dimethoxy: 8,9-Dimethoxygeibalansine

[142741-29-5]

$C_{17}H_{21}NO_5$ 319.357

Alkaloid from stem bark of *Dutailleya baudouinii*. λ_{max} 243; 315; 327 (EtOH).

(±)-form [89575-83-7]


Synthetic. Cryst. (C₆H₆/EtOH). Mp 175-176°.

Ac: [89495-53-4]


Synthetic. Needles (C₆H₆). Mp 138-139°.Bowman, R.M. *et al.*, *J.C.S. (C)*, 1966, 1504 (synth)Ahond, A. *et al.*, *Phytochemistry*, 1979, **18**, 1415-1416 (uv, ir, pmr, ms, struct, deriv)Ramesh, M. *et al.*, *Heterocycles*, 1984, **22**, 125 (synth, ir, pmr, deriv)Muyard, F. *et al.*, *Phytochemistry*, 1992, **31**, 1087-1089 (8,9-Dimethoxygeibalansine)Barr, S.A. *et al.*, *Chem. Comm.*, 1994, 153 (synth)Boyd, D.R. *et al.*, *J.C.S. Perkin 1*, 2000, 3397-3405 (synth, abs config)Chen, I.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1143-1147 (8,9-Dimethoxygeibalansine)De Moura, N.F. *et al.*, *Planta Med.*, 2002, **68**, 534-538 (R-form, isol, pmr, cmr)Wabo, H.K. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 591-595 (Tabouensintum)**Geijedimerine**

G-30

[99964-51-9]

C₂₈H₂₆N₂O₄ 454.524Alkaloid from the leaves of *Geijera balansae* (Rutaceae). Prisms (MeOH). Mp 205-207°. [α]_D²⁵ 0 (c, 0.5 in EtOH).Mitaku, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 772 (isol, uv, ir, pmr, ms, struct)**Geissolosimine**

G-31


Alkaloid D₂C₃₈H₄₄N₄O 572.792Alkaloid from the bark of *Geissospermum vellosii* (Apocynaceae). Cryst. + 1/2 H₂O (MeOH aq.). Mp 140° (133-135°). [α]_D²⁵ +74 (c, 1.0 in EtOH).19,20E,22,23-Tetrahydro, N^b-oxide: **Di-varicine**

[138683-55-3]

C₃₈H₄₀N₄O₂ 584.76Alkaloid from the roots of *Strychnos divaricans* (Loganiaceae). Prisms (MeOH/EtOAc). Mp 280-282° dec. [α]_D²² +94 (c, 0.1 in EtOH).Rapoport, H. *et al.*, *J.A.C.S.*, 1958, **80**, 1601; *J.O.C.*, 1962, **27**, 2981 (isol, uv, ir, pmr, struct, synth)Mukherjee, R. *et al.*, *Heterocycles*, 1991, **32**, 985 (Divaricine)**Geissoschizine**

Alkaloid R1

[439-66-7]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from *Rhazya stricta*, *Bonafousia tetrastachya* (preferred genus name *Tabernaemontana*) and *Rauwolfia volkensii* (Apocynaceae) and hydrol. prod. from Geissospermine, G-35. Mp 194-196° (187°). [α]_D²¹ +115 (EtOH). [α]_D +72 (CHCl₃).

2,4-Dinitrophenylhydrazone; hydrochloride: Mp 198°.

Me ether: **Geissoschizine methyl ether**

[60314-89-8]

C₂₂H₂₆N₂O₃ 366.459Alkaloid from *Uncaria rhynchophylla* (Rubiaceae). Prisms (Me₂CO/hexane). Mp 190-192°. [α]_D²⁷ +100 (c, 1 in MeOH).N^d,21-Dehydro: **4,21-Dehydrogeissoschizine**

[73385-56-5]

[73385-57-6]

C₂₁H₂₃N₂O₃[⊕] 351.424Main alkaloid from *Guettarda eximia* (Apocynaceae). Intermed. in the biosynth. of heteroyohimbine alkaloids. Hydrate (as chloride). Mp 250° dec. (chloride). Deprotonates to Cathenamine, C-207.

(Z)-Isomer:

Cryst. (MeOH). Mp 125-127°. [α]_D²⁴ -13 (c, 0.247 in Py).


(±)-form

Cryst. (EtOH). Mp 189-190°.

Janot, M.-M. *et al.*, *Tetrahedron*, 1961, **14**, 113 (isol, struct)Yamada, K. *et al.*, *Chem. Comm.*, 1974, 908 (synth)Chatterjee, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 2000 (uv, ir, pmr, ms)Hachmeister, B. *et al.*, *Chem. Ber.*, 1976, **109**, 3825 (synth)Rackur, G. *et al.*, *Chem. Ber.*, 1976, **109**, 3837 (conformn)Damak, M. *et al.*, *Tet. Lett.*, 1976, 4731 (pmr, cmr, config)Aimi, N. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 2067 (deriv)Kan-Fan, C. *et al.*, *Chem. Comm.*, 1979, 1015 (4,21-Dehydrogeissoschizine)Rüffer, M. *et al.*, *Chem. Comm.*, 1979, 1016 (biosynth)Bohlmann, C. *et al.*, *Annalen*, 1985, 1752 (synth)Winterfeldt, E. *et al.*, *Annalen*, 1986, 1262 (Z-isomer)Martin, S.F. *et al.*, *J.A.C.S.*, 1988, **110**, 5925 (synth, bibl)Overman, L.E. *et al.*, *J.A.C.S.*, 1989, **111**, 300 (synth)Wenkert, E. *et al.*, *J.O.C.*, 1989, **54**, 1166 (synth)Takayama, H. *et al.*, *Tet. Lett.*, 1992, **33**, 6831; 1997, **38**, 5307 (pmr, cmr, struct, synth)Tirkkonen, B. *et al.*, *Tetrahedron*, 1994, **50**, 3537 (Z-isomer)Lounasmaa, M. *et al.*, *Tetrahedron*, 1996, **52**, 6803 (synth)Lounasmaa, M. *et al.*, *Heterocycles*, 1999, **51**, 649-670 (rev, conformn)Bennasar, M.-L. *et al.*, *J.O.C.*, 1999, **64**, 9605-9612 (synth)Martin, S.F. *et al.*, *Org. Lett.*, 1999, **1**, 79-81 (synth)Yu, S. *et al.*, *J.A.C.S.*, 2000, **122**, 7827-7828 (synth)Deiters, A. *et al.*, *J.A.C.S.*, 2003, **125**, 4541-4550 (synth)Ariffin, J. *et al.*, *Heterocycles*, 2004, **63**, 663-670 (pmr, cmr, conformn)**Geissoschizol**

G-33

[439-69-0]



Absolute Configuration

C₁₉H₂₄N₂O 296.411Alkaloid from *Rauwolfia vomitoria*, *Aspidosperma oblongum*, *Peschiera laeta*, *Hunteria zeylanica*, *Ervatamia hainanensis*, *Tabernaemontana elegans* and others (Apocynaceae). Cryst. (MeOH/CHCl₃/heptane). Mp 216° (204°). [α]_D²⁰ -70 (Py).O-Ac: **O-Acetylgeissoschizol**

[83378-14-7]

C₂₁H₂₆N₂O₂ 338.449Alkaloid from stem bark of *Rauwolfia vomitoria* (Apocynaceae).Aldehyde: **Geissoschizal**

[41787-67-1]

C₁₉H₂₂N₂O 294.396Alkaloid in *Strychnos nux-vomica* (biosynth.) (Loganiaceae).9-Methoxy: **9-Methoxygeissoschizol**C₂₀H₂₆N₂O₂ 326.438Alkaloid from the bark of *Strychnos guianensis*. λ_{\max} 210 (sh) (log ϵ 4.21); 224 (log ϵ 4.31); 269 (log ϵ 3.74); 292 (log ϵ 3.6) (MeOH).9-Methoxy, N^b-Me: **9-Methoxy-N^b-methylgeissoschizol**C₂₁H₂₉N₂O₂[⊕] 341.472Quaternary alkaloid from *Strychnos guianensis*. Counterion not specified. λ_{\max} 207 (log ϵ 4.25); 220 (log ϵ 4.32); 259 (log ϵ 3.74); 291 (log ϵ 3.52) (MeOH).10-Hydroxy: **10-Hydroxygeissoschizol**

Huntabrine. Huntabrine

[6870-18-4]

C₁₉H₂₄N₂O₂ 312.411Alkaloid from *Amsonia elliptica*, *Rauwolfia vomitoria* and *Ervatamia hainanensis* (Apocynaceae).

10-Hydroxy, N^b-Me: Huntrabrine N-metho saltC₂₀H₂₇N₂O₂[⊕] 327.445

Quaternary alkaloid from *Hunteria eburnea* and *Pleiocarpa mutica* (Apocynaceae). Cryst. (MeOH aq.) (as chloride). Mp 282–286° dec. (chloride). [α]_D²⁶ +56 (c, 0.585 in H₂O).

10-Methoxy: 10-Methoxygeissoschizol.

19,20-Dehydro-10-methoxydihydrocorynantheol. Alkaloid AD-VI

[15266-60-1]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Aspidosperma discolor*, *Aspidosperma oblongum*, *Aspidosperma marcgravianum*, *Rauwolfia vomitoria*, *Rauwolfia obscura* and *Neisosperma kilneri* (Apocynaceae). Cryst. (MeOH/Et₂O or Me₂CO/pentane). Mp 184–185° dec. (181–182.5°). [α]_D²⁴ -64.5 (c, 0.916 in pentane) (-42).

10-Methoxy, N^b-Me: 10-Methoxy-4-methylgeissoschizol

[143257-73-2]

C₂₁H₂₉N₂O₂[⊕] 341.472

Quaternary alkaloid from bark of *Aspidosperma pruinosum* (Apocynaceae). Mp 171–173°. [α]_D²⁵ +36.04 (c, 1.0 in MeOH). Counterion not specified.

Z-Isomer(?): IsogeissoschizolC₁₉H₂₄N₂O 296.411

Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph. [α]_D -26 (c, 0.32 in MeOH). Yellow-green col. with Ce(IV). λ_{max} 227 (log ε 4.67); 273 (log ε 3.72); 281 (log ε 3.81); 290 (log ε 3.57) (EtOH) (unchanged in acid or alkali).

Z-Isomer(?), 10-methoxy: 10-Methoxyiso-geissoschizolC₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph. [α]_D +2 (c, 0.69 in MeOH). Yellow-green col. with Ce(IV). λ_{max} 229 (log ε 4.72); 284 (log ε 3.84); 294 (log ε 3.78); 308 (log ε 3.54) (EtOH) (unchanged in alkali).

Puisieux, F. et al., *C. R. Hebd. Seances Acad. Sci.*, 1959, **249**, 1369 (synth)

Rapoport, H. et al., *J.A.C.S.*, 1960, **82**, 4404 (synth)

Bartlett, M.F. et al., *J.O.C.*, 1963, **28**, 1445 (*Huntrabrine N-methosalt*)

Spieller, G. et al., *Monatsh. Chem.*, 1963, **94**, 779 (ms)

Khan, Z.M. et al., *Helv. Chim. Acta*, 1965, **48**, 1957 (*Huntrabrine N-methosalt*)

Gilbert, B. et al., *Tetrahedron*, 1965, **21**, 1141 (*10-Methoxygeissoschizol*)

Dastoor, N.J. et al., *Helv. Chim. Acta*, 1967, **50**, 213 (*10-Methoxygeissoschizol*)

Heimberger, S.I. et al., *Chem. Comm.*, 1973, 217 (*Geissoschizal*)

Sakai, S. et al., *Yakugaku Zasshi*, 1973, **93**, 483 (*10-Hydroxygeissoschizol*)

Jahodar, L. et al., *Phytochemistry*, 1974, **13**, 2880 (isol)

Iwu, M.M. et al., *Planta Med.*, 1982, **45**, 105 (*Acetylgeissoschizol*)

Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 694-707 (*Iso-geissoschizol*, *10-Methoxyiso-geissoschizol*)

Bohlmann, C. et al., *Annalen*, 1985, 1752 (synth)

Nunes, D.S. et al., *Phytochemistry*, 1992, **31**, 2507 (*10-Methoxy-4-methylgeissoschizol*)

Mavar-Manga, H. et al., *Phytochemistry*, 1996, **43**, 1125 (*9-Methoxygeissoschizol*)

Yu, S. et al., *J.A.C.S.*, 2000, **122**, 7827-7828 (synth)


Penelle, J. et al., *Phytochemistry*, 2000, **53**, 1057-1066 (*9-Methoxy-N-methylgeissoschizol*)

Ma, J. et al., *Org. Lett.*, 2007, **9**, 3491-3494 (synth)

Geissoschizoline

G-34

16α-Curan-17-ol, 9CI. Pereirine [18397-07-4]



Absolute Configuration

C₁₉H₂₆N₂O 298.427

Alkaloid from *Geissospermum vellosii* and *Geissospermum sericeum*, identical with acid hydrol. prod. of Geissospermine, G-35 and therefore possible artifact (Apocynaceae). Mp 84–87° Mp 85–105° Mp 98–143°. [α]_D²⁰ +8 (c, 4 in MeOH). Mp wide and variable owing to solvation.

Picolonate:

Cryst. (EtOH). Mp 209–211°.

N⁴-Oxide: Geissoschizoline N⁴-oxide

[402752-15-2]

C₁₉H₂₆N₂O₂ 314.427

Alkaloid from the bark of *Geissospermum sericeum*. [α]_D²⁰ +31 (c, 0.64 in MeOH). λ_{max} 245 ; 300 (MeOH).

Di-Ac:Cryst. (Me₂CO). Mp 196–197°.**1,2-Didehydro: 1,2-Didehydrogeissoschizoline. 1,2-Dehydrogeissoschizoline**

[402752-17-4]

C₁₉H₂₄N₂O 296.411

Alkaloid from the bark of *Geissospermum sericeum*.

[23365-75-5]

Bertho, A. et al., *Chem. Ber.*, 1958, **91**, 2581; 1961, **94**, 2737 (isol, struct)

Janot, M.M. et al., *Tetrahedron*, 1961, **14**, 113 (isol, ir, uv, struct)


Harley-Mason, J. et al., *Chem. Comm.*, 1969, 665; 1970, 812 (synth)

Steele, J.C.P. et al., *J. Nat. Prod.*, 2002, **65**, 85-88 (isol, uv, pmr, cmr)

Geissospermine

G-35

[427-01-0]



Absolute configuration

C₄₀H₄₈N₄O₃ 632.844

Alkaloid from the bark of *Geissospermum vellosii*, *Geissospermum laeve*, *Geissospermum sericeum* and

Tabernaemontana laevis (Apocynaceae).

Mp 213–214° (sinters at 160°) (anhyd. 217–219° dec.). [α]_D²⁰ -101 (c, 1 in EtOH). Hydrol. gives Geissoschizine, G-32.

▶GN1040000

Hydrochloride: Mp 148°.

Hesse, O. et al., *Ber.*, 1877, **10**, 2162 (isol)

Rapoport, H. et al., *J.A.C.S.*, 1960, **82**, 4404 (uv, ir, struct)

Janot, M.-M. et al., *Tetrahedron*, 1961, **14**, 113 (struct)

Schmid, H. et al., *Biochem. Physiol. Alkaloid. Int. Symp.*, 4th, 1969 (1972), 1972, 337; *CA*, **77**, 140397z (synth)

Chiaroni, A. et al., *Tet. Lett.*, 1976, 4729 (cryst struct)


Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GCG300

Geissovelline

G-36

1-Acetyl-14,19-didehydro-10,11-dimethoxy-4-methyl-3,4-secocondyfolan-3-one, 9CI

[36954-68-4]



Relative Configuration

C₂₃H₃₀N₂O₄ 398.501

Alkaloid from the bark of *Geissospermum vellosii* (Apocynaceae). Cryst.

(CHCl₃/Et₂O or EtOH). Mp 189–190°.

[α]_D²⁵ -125 (c, 1.15 in CHCl₃). pK_a 6.7 (50% EtOH aq.). CAS gives the abs. config. as shown, but it has not been definitely determined. λ_{max} 217 (ε 22600); 262 (ε 17500); 299 (ε 10500) (95% EtOH).

N-De-Ac:


Pale-yellow cryst. Mp 158–159.5°. [α]_D²⁸ -6 (c, 1.07 in CHCl₃). pK_a 7 (50% EtOH aq.). λ_{max} 230 (ε 12500); 305 (ε 6260) (95% EtOH).

Moore, R.E. et al., *J.O.C.*, 1973, **38**, 215-230 (isol, uv, ir, pmr, cmr, ms, struct)

Gelliusine A

G-37

[159903-67-0]

C₃₀H₃₀Br₂N₆O 650.415

Alkaloid from a deep water New Caledonian marine sponge (*Gellius* or *Orina* sp.). Serotonin receptor agonist. Racemic. λ_{max} 226 (ε 52000); 285 (ε 15000) (MeOH).

Diastereoisomer: Gelliusine B

[159992-42-4]

C₃₀H₃₀Br₂N₆O 650.415

From a deep water New Caledonian


marine sponge (*Gellius* or *Orina* sp.). Serotonin receptor agonist. Racemate with different rel. configs. at C-8 and C-8'' from (\pm)-Gelliusine A. λ_{\max} 226 (ϵ 52000); 285 (ϵ 15000) (MeOH) (Berdy).

Bifulco, G. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1294 (*isol, uv, pmr, cmr, struct*)

Gelliusine C

G-38

[171090-79-2]




$C_{30}H_{30}Br_2N_6O$ 650.415

Alkaloid from the deep-water New Caledonian marine sponge *Orina* sp. and from *Gellius* sp. Shows antiserotonin activity. Racemic. λ_{\max} 230 (ϵ 35234); 284 (ϵ 13272) (MeOH) (Berdy).

Bifulco, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1254 (*isol, uv, pmr, cmr, struct*)

Gelliusine D

G-39



$C_{20}H_{21}BrN_4O$ 413.316


(\pm)-*form* [171090-80-5]

Alkaloid from the deep-water New Caledonian marine sponge *Orina* sp. and from *Gellius* sp. Shows antiserotonin activity. λ_{\max} 230 (E1%/1cm 284) (MeOH) (Berdy).

Bifulco, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1254-1260 (*isol, pmr, cmr, struct*)

Gelliusine E

G-40



$C_{20}H_{21}BrN_4O$ 413.316

(\pm)-*form* [171090-81-6]

Alkaloid from the deep-water New

Caledonian marine sponges *Orina* sp. and *Gellius* sp. Shows antiserotonin activity. Neuropeptide Y receptor binding agent. λ_{\max} 230 ; 284 (MeOH) (Berdy).

5-Deoxy, 6-bromo: **Gelliusine F**

[171090-82-7]

$C_{20}H_{20}Br_2N_4$ 476.213


From *Orina* sp. and *Gellius* sp. Antiserotonin agent, neuropeptide Y binding agent. λ_{\max} 230 ; 284 (MeOH) (Berdy).

Bifulco, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1254-1260 (*isol, pmr, cmr, struct*)

Gelsamydine

G-41

[120881-61-0]



$C_{29}H_{36}N_2O_6$ 508.613

Alkaloid from the whole plant of *Gelsemium elegans*. Needles. Mp 194-196°. $[\alpha]_D^{20}$ -126.9 (c, 0.26 in MeOH). λ_{\max} 206 ; 257 (MeOH).

14R-Hydroxy: **14 α -Hydroxygelsamydine**

$C_{29}H_{36}N_2O_7$ 524.613

Alkaloid from *Gelsemium elegans*. Needles. $[\alpha]_D^{20}$ -71.5 (c, 0.07 in $CHCl_3$). λ_{\max} 258 (log ϵ 4.66) (MeOH).

19-Epimer, 19-hydroxy: **19 α -Hydroxygelsamydine**

[181948-91-4]

$C_{29}H_{36}N_2O_7$ 524.613

Alkaloid from whole plant of *Gelsemium elegans*. Yellowish powder. $[\alpha]_D$ -75 (c, 0.01 in MeOH). λ_{\max} 209 (log ϵ 4.29); 257 (log ϵ 3.67) (MeOH).

Lin, L.-Z. *et al.*, *J.O.C.*, 1989, **54**, 3199-3202 (*isol, pmr, cmr, ms, cd, cryst struct*)


Lin, L.-Z. *et al.*, *Phytochemistry*, 1996, **43**, 723-726 (*19 α -Hydroxygelsamydine*)

Xu, Y.-K. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1347-1350 (*14-Hydroxygelsamydine*)

Gelsebanine

G-42

[911847-81-9]



$C_{30}H_{36}N_2O_5$ 504.625


Alkaloid from *Gelsemium elegans*.

Amorph. powder. $[\alpha]_D^{20}$ -8 (c, 0.08 in $CHCl_3$). λ_{\max} 255 (log ϵ 4.06); 287 (sh); 317 (log ϵ 3.08) (MeOH).

Xu, Y.-K. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1347-1350 (*isol, pmr, cmr, ms*)

Gelsedilam

G-43



Absolute Configuration

$C_{17}H_{18}N_2O_4$ 314.34

Related to Gelsedine, G-44. Alkaloid from the leaves of *Gelsemium elegans*. λ_{\max} 209 (log ϵ 4.35); 257 (log ϵ 3.72) (MeOH).

14R-Acetoxy: **14-Acetoxygelsedilam**

$C_{19}H_{20}N_2O_6$ 372.377


Alkaloid from the leaves of *Gelsemium elegans*. λ_{\max} 210 (log ϵ 4.31); 257 (log ϵ 3.65) (MeOH).

Kogure, N. *et al.*, *Org. Lett.*, 2006, **8**, 3085-3088 (*isol, cd, pmr, cmr, ms*)

Gelsedine, 9CI

G-44

[7096-96-0]



Absolute Configuration

$C_{19}H_{24}N_2O_3$ 328.41

Alkaloid from *Gelsemium sempervirens*. Mp 172.6-174°. $[\alpha]_D^{25}$ -158 (c, 1.35 in $CHCl_3$).

Perchlorate: Mp 183°. $[\alpha]_D^{20}$ -96 (c, 0.73 in $CHCl_3$).

4,20-Didehydro: **4,20-Didehydrogelsedine.**

Gelsenicine. Humantenmine

[82354-38-9]

$C_{19}H_{22}N_2O_3$ 326.394

Alkaloid from roots of *Gelsemium elegans*. Exhibits analgesic activity. Mp 166-168°. $[\alpha]_D$ -147. λ_{\max} 210 (log ϵ 4.38); 256 (log ϵ 3.84) (no solvent reported).

19-Oxo, 4,20-didehydro: **19-Oxogelsenicine. 4,20-Didehydro-19-oxogelsedine, 9CI**

[113900-77-9]

$C_{19}H_{20}N_2O_4$ 340.378

Alkaloid from leaves of *Gelsemium elegans*. Plates (Me_2CO). Mp 226-227°. λ_{\max} 209 ; 256 ; 281 (MeOH).

14R-Hydroxy: **14-Hydroxygelsedine**

[100482-87-9]

$C_{19}H_{24}N_2O_4$ 344.41

Alkaloid from stems of *Gelsemium sempervirens*. Needles. Mp 216-218°.

$[\alpha]_D^{25}$ -113 (c, 0.3 in MeOH). λ_{\max} 216 (log ϵ 4.26); 260 (log ϵ 3.68) (MeOH).

14R-Hydroxy, 4,20-didehydro: 4,20-Didehydro-14-hydroxygelsedine, 9CI. 14-Hydroxygelsenicine. Humantenidine [114027-39-3]

[82375-28-8 (Humantenidine)]

$C_{19}H_{22}N_2O_4$ 342.394

Alkaloid from roots of *Gelsemium elegans*. Gum. $[\alpha]_D$ -123. λ_{\max} 210 (log ϵ 4.38); 252 (log ϵ 3.98) (no solvent reported).

14R-Hydroxy, 19-oxo, 4,20-didehydro: 14-Hydroxy-19-oxogelsenicine. 14-Hydroxy-19-oxohumantenmine. 19-Oxohumantenidine

$C_{19}H_{20}N_2O_5$ 356.377

Alkaloid from the leaves of *Gelsemium elegans*. Amorph. $[\alpha]_D^{19}$ -146.1 (c, 0.14 in $CHCl_3$). λ_{\max} 213 (log ϵ 4.28); 256 (log ϵ 3.73) (MeOH).

14R, 15-Dihydroxy, 4,20-didehydro: 14,15-Dihydroxygelsenicine [557113-06-1]

$C_{19}H_{22}N_2O_5$ 358.393

Alkaloid from *Gelsemium elegans*. Amorph. solid. λ_{\max} 212 (log ϵ 4.29); 258 (log ϵ 3.76) (MeOH).

14R-Acetoxy, 4,20-didehydro: 14-Acetoxygelsenicine. 14-O-Acetylhumantenidine

$C_{21}H_{24}N_2O_5$ 384.431

Alkaloid from the leaves of *Gelsemium elegans*. Amorph. $[\alpha]_D^{15}$ -102.6 (c, 1.5 in $CHCl_3$). λ_{\max} 213 (log ϵ 4.27); 258 (log ϵ 3.77) (MeOH).

14R-Acetoxy, 15-hydroxy, 4,20-didehydro: 14-Acetoxy-15-hydroxygelsenicine. 14-O-Acetyl-15-hydroxyhumantenidine

$C_{21}H_{24}N_2O_6$ 400.43

Alkaloid from the leaves of *Gelsemium elegans*. Amorph. $[\alpha]_D^{19}$ -66.3 (c, 0.09 in $CHCl_3$). λ_{\max} 212 (log ϵ 4.26); 258 (log ϵ 3.73) (MeOH).

11-Methoxy: Gelsemicine

[6887-28-1]

$C_{20}H_{26}N_2O_4$ 358.436

Alkaloid from *Gelsemium sempervirens* and *Mostuea brunonis*. Respiratory stimulant. Prisms (Me₂CO). Mp 171-172°. $[\alpha]_D^{25}$ -142 (c, 0.945 in $CHCl_3$).

► Highly toxic. Causing death by respiratory paralysis. Most toxic of the *Gelsemium* alkaloids.

11-Methoxy; hydrochloride: Mp 140-142°.

11-Methoxy, 4,20-didehydro: 4,20-Didehydro-11-methoxygelsedine. 4,20-Didehydrogelsenicine

[76328-63-7]

$C_{20}H_{24}N_2O_4$ 356.421

Alkaloid from *Gelsemium sempervirens* and *Mostuea brunonis*. λ_{\max} 214 ; 285 ; 295 (sh) (MeOH).

11-Methoxy, 14R-hydroxy: 14-Hydroxygelsenicine. Alkaloid C†

[41478-09-5]

$C_{20}H_{26}N_2O_5$ 374.436

Alkaloid from *Gelsemium sempervirens* and *Mostuea brunonis*. Plates (Me₂CO).

11-Methoxy, 14R-hydroxy, 4,20-didehydro: 14-Hydroxy-11-methoxygelsenicine. 4,20-Didehydro-14-hydroxygelsenicine. GS 2

[677761-62-5]

$C_{20}H_{24}N_2O_5$ 372.42

Alkaloid from *Gelsemium sempervirens*. Amorph. λ_{\max} 218 ; 286 ; 296 (sh) (MeOH).

11-Methoxy, 14R-hydroxy, 19-oxo, 4,20-didehydro: 14-Hydroxy-11-methoxy-19-oxogelsenicine. GS 1

[677761-61-4]

$C_{20}H_{22}N_2O_6$ 386.404

Alkaloid from *Gelsemium sempervirens*. Amorph. solid. λ_{\max} 218 ; 287 ; 295 (sh) (MeOH).

Forsyth, W.G.C. *et al.*, *J.C.S.*, 1945, 579-582 (*Gelsemicine, isol*)

Schwarz, H. *et al.*, *J.A.C.S.*, 1953, **75**, 4372 (*Gelsemicine, isol*)

Przybylska, M. *et al.*, *Can. J. Chem.*, 1961, **39**, 2124-2127 (*Gelsemicine, cryst struct, ir*)

Wenkert, E. *et al.*, *Experientia*, 1972, **28**, 377-379 (*cmr, ms*)

Wichtl, M. *et al.*, *Monatsh. Chem.*, 1973, **104**, 87-98; *Ann. Chim. Farm.*, 1973, **104**, 99-104 (*ms, 14-Hydroxygelsenicine*)

Onanga, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1980, **291**, 191-193 (*uv, ir, pmr, cmr, ms, struct, Didehydrogelsenicine*)

Yang, J. *et al.*, *Yaoxue Xuebao*, 1982, **17**, 119-120; 633-634; 1983, **18**, 104-112; 1984, **19**, 437-440; *CA*, **97**, 36099f; 212699n; **99**, 102248y; **103**, 3678h (*Humantenmine, Humantenidine*)

Schun, Y. *et al.*, *J. Nat. Prod.*, 1985, **48**, 788-791 (*14-Hydroxygelsedine*)

Ponglux, D. *et al.*, *Tet. Lett.*, 1988, **29**, 5395-5396 (*cmr*)

Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075-5094 (*19-Oxogelsenicine*)

Kitajima, M. *et al.*, *J.C.S. Perkin 1*, 1994, 1573-1578 (*Gelsemicine, synth*)

Takayama, H. *et al.*, *J.O.C.*, 1994, **59**, 4381 (*Gelsedine, Gelsenicine, synth*)

Beyersbergen, W.G. *et al.*, *J.O.C.*, 2000, **65**, 8317-8325 (*ent-Gelsedine, synth*)


Kitajima, M. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1211-1214 (*GS1, GS2, 4,20-Didehydrogelsenicine*)

Kitajima, M. *et al.*, *Org. Lett.*, 2003, **5**, 2075-2078 (*14,15-Dihydroxygelsenicine*)

Kitajima, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 715-718; 2007, **70**, 142 (*14-Acetoxygelsenicine, 14-Acetoxy-15-hydroxygelsenicine, 14-Hydroxy-19-oxogelsenicine*)

Gelsefuranidine

G-45



Absolute Configuration


$C_{24}H_{24}N_2O_5$ 420.464

Alkaloid from the leaves of *Gelsemium elegans*. λ_{\max} 208 (log ϵ 4.21); 255 (sh) (log ϵ 3.67); 306 (log ϵ 4.14); 362 (sh) (log ϵ 2.77) (MeOH).

Kogure, N. *et al.*, *Org. Lett.*, 2006, **8**, 3085-3088 (*isol, cd, pmr, cmr, ms*)

Gelseiridone

G-46



Absolute Configuration

$C_{29}H_{34}N_2O_8$ 538.596

Highly modified indole alkaloid most closely related to the *Gelsemium* group. Alkaloid from the leaves of *Gelsemium elegans*. λ_{\max} 205 (log ϵ 4.33); 288 (log ϵ 4.28) (MeOH).

Kogure, N. *et al.*, *Org. Lett.*, 2006, **8**, 3085-3088 (*isol, cd, pmr, cmr, ms*)

Gelselegine

G-47

20-(Hydroxymethyl)gelsedine, 9CI

[131653-76-4]

As Gelsedine, G-44 with R = CH₂OH

$C_{20}H_{26}N_2O_4$ 358.436

Alkaloid from *Gelsemium elegans* (Loganiaceae). Prisms (MeOH). Mp 167-168°. $[\alpha]_D$ -30 (c, 0.02 in MeOH).

19R-Hydroxy, 11-methoxy: 19R-Hydroxy-11-methoxygelselegine

[131683-37-9]

$C_{21}H_{28}N_2O_6$ 404.462

Alkaloid from *Gelsemium elegans* (Loganiaceae). Prisms (MeOH). Mp 234-236°. $[\alpha]_D$ -110 (c, 0.02 in MeOH).

14R-Acetoxy: 14-Acetoxygelselegine

$C_{22}H_{28}N_2O_6$ 416.473

Alkaloid from the leaves of *Gelsemium elegans*. Amorph. $[\alpha]_D^{30}$ -62.1 (c, 1 in $CHCl_3$). λ_{\max} 209 (log ϵ 4.23); 258 (log ϵ 3.6) (MeOH).

Lin, L.-Z. *et al.*, *Phytochemistry*, 1990, **29**, 3013 (*isol, pmr, cmr, cryst struct*)

Takayama, H. *et al.*, *J.O.C.*, 1994, **59**, 4381 (*synth*)


Takayama, H. *et al.*, *Tetrahedron*, 1994, **50**, 11813 (*synth, 19-Hydroxy-11-methoxygelselegine*)

Kitajima, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 715-718 (*14-Acetoxygelselegine*)

Gelsemamide

G-48

[122297-34-1]



$C_{20}H_{24}N_2O_3$ 340.421

Novel N(1)-C(2) secoindole alkaloid. Alkaloid from *Gelsemium elegans* (Loganiaceae). Needles (Me₂CO). Mp 183-184°. $[\alpha]_D$ +228.3 (c, 0.046 in MeOH). λ_{\max} 207 (ϵ 30200); 233 (ϵ 6610); 284 (ϵ 1950) (MeOH) (Derep).

11-Methoxy: 11-Methoxygelsemamide

[122297-35-2]


$C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from *Gelsemium elegans* (Loganiaceae). Needles (Me₂CO). Mp 140°. $[\alpha]_D^{25} +215.5$ (c, 0.2 in MeOH). λ_{max} 207 (ϵ 30200); 233 (ϵ 6610); 284 (ϵ 1950) (MeOH) (Derep).

Lin, L.-Z. *et al.*, *Tet. Lett.*, 1989, **30**, 1177 (*isol, uv, ir, pmr, cmr, ms, cd, cryst struct*)
Takayama, H. *et al.*, *Tetrahedron*, 1994, **50**, 8363 (*synth, 11-Methoxygelsemine*)

Gelsemine, 9CI**G-49**

[509-15-9]



Relative configuration

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Gelsemium sempervirens*, *Gelsemium elegans* and *Mostuea stimulans* (Loganiaceae). Analgesic, CNS stimulant. *Gelsemium* Root has been used to treat trigeminal neuralgia and migraine. Also used in homoeopathic medicine. Mp 178°. $[\alpha]_D^{20} +17.8$ (c, 2.03 in CHCl₃). Log P 0.48 (uncertain value) (calc).

► V. toxic. *G. spp.* have been freq. causes of human poisoning. LD₅₀ (mus, ipr) 49 mg/kg. CNS depressant, can cause respiratory arrest. LX9100000

Hydrochloride: Mp 333° dec. (rapid heating). pK_a 9.79 (13°).

Hydrobromide:

Prisms (EtOH aq.). Mp 325° dec.

Picrate:

Yellow plates (MeOH). Mp 203°.

N^b-Oxide (R-): [119363-15-4]

Semisynthetic by oxidation of *Gelsemine*. Amorph. $[\alpha]_D^{25} +20.4$ (c, 0.23 in MeOH).

N^b-Oxide (S-): **Gelsemine N-oxide**

[113900-76-8]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the leaves of *Gelsemium elegans* (Loganiaceae). Amorph. $[\alpha]_D^{25} -16.9$ (c, 0.19 in MeOH).

21-Oxo: 21-OxogelsemineC₂₀H₂₀N₂O₃ 336.36

Alkaloid from *Gelsemium sempervirens* (Loganiaceae). Mp 148-150°.

19R-Hydroxy, 18,19-dihydro: 19-Hydroxydihydrogelsemine. 18,19-Dihydro-19-hydroxygelsemine, 9CI

[134306-20-0]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from *Gelsemium elegans*, *Gelsemium rankinii* and *Gelsemium sempervirens* (Loganiaceae). Powder. Mp 230-232°. $[\alpha]_D -20$ (c, 0.25 in MeOH).

N-Methoxy: Gelsevirine. N-Methoxygelsemine. Alkaloid A†

[38990-03-3]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Gelsemium sempervirens* and *Gelsemium elegans* (Loganiaceae). Oil. Mp 245°. $[\alpha]_D -4.5$.

N-Methoxy, 21-oxo: 21-Oxogelsevirine

[104386-95-0]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from stems of *Gelsemium rankinii* (Loganiaceae). Needles. Mp 226-228°. $[\alpha]_D^{25} -67$ (c, 0.1 in MeOH).

N-Methoxy, 19R-hydroxy, 18,19-dihydro:**19R-Hydroxydihydrogelsevirine**

[114027-49-5]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the roots of *Gelsemium elegans* and other *Gelsemium* spp. (Loganiaceae). Needles. Mp 210-212°. $[\alpha]_D -34$ (c, 0.1 in MeOH).

N-Methoxy, 19S-hydroxy, 18,19-dihydro:**19S-Hydroxydihydrogelsevirine**

[113924-12-2]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Gelsemium elegans*. Amorph. $[\alpha]_D -68$.

19R-Acetoxy, N-methoxy, 18,19-dihydro:**19-Acetoxydihydrogelsevirine**

[134306-19-7]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Gelsemium* spp. (Loganiaceae). Cryst. (Me₂CO). Mp 187-189°. $[\alpha]_D -6.7$ (c, 0.3 in MeOH).

Forsyth, W.G.C. *et al.*, *J.C.S.*, 1945, 579 (*isol*)
Lovell, F.M. *et al.*, *Tet. Lett.*, 1959, **4**, 1 (*cryst struct*)

Conroy, H. *et al.*, *Tet. Lett.*, 1959, **4**, 6 (*ir, pmr, struct*)

Wenkert, E. *et al.*, *Chem. Comm.*, 1970, 961 (*pmr, cmr*)

Wenkert, E. *et al.*, *Experientia*, 1972, **28**, 377 (*pmr, cmr*)

Wichtl, M. *et al.*, *Monatsh. Chem.*, 1973, **104**, 87 (*Gelsevirine*)

Nikiforov, A. *et al.*, *Monatsh. Chem.*, 1974, **105**, 1292 (*21-Oxogelsemine*)

Nagakura, N. *et al.*, *J.C.S. Perkin 1*, 1979, 2308 (*biosynth*)

Stöckigt, J. *et al.*, *Tet. Lett.*, 1979, 2615 (*biosynth*)

Schun, Y. *et al.*, *J. Nat. Prod.*, 1986, **49**, 483 (*21-Oxogelsevirine*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 4668 (*pmr, cmr, struct*)

Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075 (*isol, uv, ir, pmr, struct, oxide*)

Lin, L.-Z. *et al.*, *Phytochemistry*, 1991, **30**, 679 (*19-Hydroxydihydrogelsemine, 19-Acetoxydihydrogelsevirine*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1372

Takayama, H. *et al.*, *Nat. Prod. Lett.*, 1993, **2**, 271 (*Gelsevirine, synth*)

Sheikh, Z. *et al.*, *Chem. Comm.*, 1994, 763 (*synth*)

Dutton, J.K. *et al.*, *Chem. Comm.*, 1994, 765 (*synth*)

Newcombe, N.J. *et al.*, *Chem. Comm.*, 1994, 767 (*synth*)

Kuzmich, D. *et al.*, *J.A.C.S.*, 1994, **116**, 6943 (*synth, 21-Oxogelsemine*)

Speckamp, W.N. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2163 (*synth, Gelsemine, 21-Oxogelsemine*)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1996, **118**, 7426 (*synth*)

Atarashi, S. *et al.*, *J.A.C.S.*, 1997, **119**, 6226 (*synth*)

Madin, A. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 2934-2936 (*synth*)

Ng, F.W. *et al.*, *J.A.C.S.*, 2002, **124**, 9812-9824 (*synth*)

Lin, H. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 36-51 (*rev. synth*)


Han, H.-B. *et al.*, *Acta Cryst. E*, 2007, **63**,

o1369-o1371 (*Gelsevirine, cryst struct*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GCK000

Gelsemoxonine**G-50**

[135626-64-1]



Absolute Configuration

C₁₉H₂₂N₂O₅ 358.393

Structure revised in 2003. Alkaloid from *Gelsemium elegans*. Prisms (C₆H₆). Mp 171-172°. $[\alpha]_D^{25} -41.3$ (c, 1 in MeOH). λ_{max} 209 (log ϵ 4.31); 257 (log ϵ 3.64) (MeOH).

11-Methoxy: 11-Methoxygelsemoxonine.**GS 3**

[677761-63-6]

C₂₀H₂₄N₂O₆ 388.419

Alkaloid from *Gelsemium sempervirens*. Amorph. solid. λ_{max} 219; 286; 295 (sh) (MeOH).

Lin, L.-Z. *et al.*, *Phytochemistry*, 1991, **30**,


1311-1315 (*isol, uv, ir, pmr, cmr, ms*)

Kitajima, M. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1211-1214 (*11-Methoxygelsemoxonine*)

Kitajima, M. *et al.*, *Org. Lett.*, 2003, **5**, 2075-2078 (*cd, pmr, cmr, cryst struct*)

Geneserine**G-51**

2,3,4,4a,9,9a-Hexahydro-2,4a,9-trimethyl-1,2-oxazino[6,5-b]indol-6-ol methylcarbamate (ester), 9CI. Eseridine, INN. Eserine aminoxide. Eserine oxide. Physostigmine oxide

C₁₅H₂₁N₃O₃ 291.349

Anticholinesterase agent used in gastrointestinal disorders and chronic dermatoses. Possesses purgative props.

(-)-form [25573-43-7]

Alkaloid from *Physostigma venenosum* (Fabaceae) also formed from Physostigmine, P-400 by oxidn. Mp 128-129°. $[\alpha]_D -175$ (EtOH). Pharmacol. active isomer.

Picrate: Mp 175°.*Methiodide*: Mp 215°.**(±)-form** [105816-13-5]

Synthetic. Oil.

Bild, N. *et al.*, *Helv. Chim. Acta*, 1967, **50**,

1885-1892 (*ms*)

Hootele, C. *et al.*, *Tet. Lett.*, 1969, 2713-2716

(ir, ms, pmr, struct)

Robinson, B. *et al.*, *J.C.S. (C)*, 1970, 2077-

2078 (*config*)

Shishido, K. *et al.*, *J.C.S. Perkin 1*, 1987, 2491-

2495 (*synth, ir, pmr, bibl*)


Wright, C. *et al.*, *Tet. Lett.*, 1987, **28**, 6389-

6390 (*synth*)

- Yu, Q.-S. *et al.*, *J. Nat. Prod.*, 1989, **52**, 332-336 (*pmr, struct*)
 Bacchi, A. *et al.*, *Acta Cryst. C*, 1994, **50**, 1126-1130 (*cryst struct*)
 Node, M. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 715-719 (*synth, bibl*)

Genipamide G-52

[942914-13-8]




$C_{16}H_{21}NO_6$ 323.345
 Constit. of *Genipa americana*. Syrup.
 $[\alpha]_D^{32} +207.7$ (c, 1.4 in MeOH).

Ono, M. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 632-634 (*Genipamide*)

Gentiactaline G-53

[94444-36-7]




$C_{32}H_{39}N_3O_5$ 545.677
 Alkaloid from the root bark of *Alstonia undulata* (Apocynaceae). $[\alpha]_D -332$ (c, 0.27 in $CHCl_3$).

Guillaume, D. *et al.*, *Phytochemistry*, 1984, **23**, 2407 (*isol, uv, ir, pmr, ms, struct*)

Gentianadine G-54

3,4-Dihydro-1H-pyrano[3,4-c]pyridin-1-one, 9CI
 [6790-32-5]



$C_8H_7NO_2$ 149.149
 Alkaloid from *Gentiana turkistanorum*, *Gentiana olgae*, *Gentiana olivieri* and *Cephalaria* spp. (Gentianaceae, Dipsacaceae). Mp 77-78°.

Hydrochloride: Mp 196-197°.


Samatov, A. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 182; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 150 (*isol, ir*)

Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 14; 66; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 10; 59 (*pmr, ms*)

Dolby, J. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 735; 1972, **26**, 2469 (*synth*)

Gentianal G-55

3,4,6,7-Tetrahydro-1-methyl-6-oxo-1H-pyrano[3,4-c]pyridine-5-carboxaldehyde, 9CI
 [53848-05-8]




$C_{10}H_{11}NO_3$ 193.202
 Alkaloid from gentiana (a Chinese medicinal herb). Mp 208-210°. $[\alpha]_D +0.8$ (approx.).

Xue, Z. *et al.*, *Kexue Tongbao (Chin. edn.)*, 1974, **19**, 378; *CA*, **82**, 13964k (*isol, uv, ir, pmr, struct*)

Gentianamine G-56

[22952-54-1]



$C_{11}H_{11}NO_3$ 205.213
 Alkaloid from *Gentiana olivieri* and *Gentiana turkistanorum* (Gentianaceae). Mp 149-150°.

Picrate: Mp 146-147°.

Ac: Mp 96-97°.


Samatov, A. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 182; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 150 (*isol, uv, ir, ms*)

Akramov, S.F. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 14; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 10 (*pmr*)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (*rev*)

Gentiananine G-57

3,4-Dihydro-6(8)-methoxy-5,8(5,6)-dimethyl-1H-pyrano[3,4-c]pyridin-1-one [11075-48-2]



$R^1, R^2 = CH_3, OMe$


$C_{11}H_{13}NO_3$ 207.229
 Exact struct. not determined. Alkaloid from *Gentiana* spp., *Pedicularis macrochila* and *Swertia* spp. Mp 380-382° (375-380° dec.). λ_{max} 219 (log ϵ 3.89); 265 (log ϵ 3.5) (no solvent reported).

Rakhmatullaev, T.U. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 32-36; 1971, **7**, 128; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 26-28; 1971, **7**, 123 (*isol, ms*)

Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 122; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 116 (*isol, ir, pmr*)

Gentianidine G-58

[2202-12-2]




$C_9H_9NO_2$ 163.176
 Alkaloid from *Gentiana asclepiadea*, *Gentiana macrophylla*, *Erythraea centaureum* (preferred genus name *Centaureum*), *Menyanthes trifoliata* and *Centaureum spicatum* (Gentianaceae, Menyanthaceae). Hypothermic, antihypertensive, antiinflammatory agent and muscle relaxant. Of low toxicity. Mp 131-132°. Log P 1.25 (calc).

Liang, H.-T. *et al.*, *Yaoyue Xuebao*, 1964, **11**, 412; *CA*, **62**, 5309g (*isol, ir, uv, pmr, synth*)
 Popov, S. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1968, **21**, 435; *CA*, **69**, 52368v (*ms*)
 Marekov, N.L. *et al.*, *Tetrahedron*, 1968, **24**, 1323 (*synth*)

Gentianine† G-59

5-Ethenyl-3,4-dihydro-1H-pyrano[3,4-c]pyridin-1-one, 9CI. 4-(2-Hydroxyethyl)-5-vinylnicotinic lactone. Erythricine
 [439-89-4]



$C_{10}H_9NO_2$ 175.187
 Alkaloid from *Gentiana kirilowi* and very many other *Gentiana* spp., from several *Swertia* spp. and several other genera in Gentianaceae. Also found in *Anthocleista procera*, *Strychnos angolensis* and other *Strychnos* spp. CNS stimulant, antihypertensive, antiinflammatory agent and muscle relaxant. Analgesic potentiator of low toxicity. Shows antipsychotic props. Mp 83°. Log P 1.48 (calc). The most widespread alkaloid of its group. λ_{max} 218 (ϵ 17500); 285 (ϵ 990) (no solvent reported).

▶ UQ1374500

Hydrochloride: Mp 171-172° dec.

Picrate: Mp 123-124°.

Proskurnina, N.F. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1944, **14**, 1148-1152; *CA*, **40**, 7213 (*isol, struct*)

Govindachari, T.R. *et al.*, *J.C.S.*, 1957, 551-556; 2725-2726 (*isol, struct, uv, synth*)

Lavie, D. *et al.*, *Chem. Ind. (London)*, 1963, 781-782 (*isol, uv, ir, pmr*)


Bailleul, F. *et al.*, *Phytochemistry*, 1977, **16**, 723-726 (*cmr*)

Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, Strychnos*)

Gentianine lactam

G-60

5-Ethenyl-3,4-dihydro-2,7-naphthyridin-1(2H)-one, 9CI
[149155-04-4]



C₁₀H₁₀N₂O 174.202

CAS numbering shown. Alkaloid from *Strychnos pungenis*.

3,4-Didehydro: **Dehydrogentianine lactam**
[189323-28-2]

C₁₀H₈N₂O 172.186


Alkaloid from *Strychnos xantha*. λ_{max} 205 ; 258 ; 310 (MeOH).

Thépenier, P. *et al.*, *Bull. Soc. R. Sci. Liege*, 1996, **65**, 379-382; 383-386 (*isol, ir, uv, pmr, cmr*)

Gentiatibetine

G-61

Gentiatibetine
[26005-36-7]



C₉H₁₁N₂O₂ 165.191

Alkaloid from *Gentiana asclepiadea*, *Gentiana lutea* (yellow gentian), *Gentiana olivieri*, *Gentiana punctata*, *Gentiana purpurea*, *Gentiana tibetica*, *Menyanthes trifoliata* (Gentianaceae, Menyanthaceae). Mp 161.5°. λ_{max} 263 (ε 270) (EtOH).

Me ether: **Oliveridine**†

[29276-66-2]

C₁₀H₁₃N₂O₂ 179.218

Alkaloid from *Gentiana olivieri* (Gentianaceae). Cryst. (Me₂CO). Mp 260° dec.


Rulko, F. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1967, **41**, 567-571 (*Gentiatibetine*)

Rakhmatullaev, T.U. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 608; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 531-532 (*Oliveridine*)

Gentiocrucine

G-62

3-(Aminomethylene)dihydro-2H-pyran-2,4(3H)-dione, 9CI. *Gentianaine*
[58213-76-6]



C₆H₇N₂O₃ 141.126

Consists of interconverting E- and Z-forms. Alkaloid from *Gentiana cruciata* and *Enicostema hyssopifolium* (Gentianaceae). Cryst. Mp 144-145°. Genus

name given as *Enicostemma*.

[22108-77-6]

Popov, S. *et al.*, *Chem. Ind. (London)*, 1969, 49 (*isol, pmr, ms, ir*)


Ghosal, S. *et al.*, *Tet. Lett.*, 1974, 403 (*isol, struct, uv, ir, pmr, cmr, ms*)

Ganem, B. *et al.*, *J.A.C.S.*, 1976, **98**, 224 (*synth*)

Gentioflavine

G-63

[18058-50-9]



C₁₀H₁₁N₃O 193.202

Alkaloid from *Gentiana asclepiadea* and many other *Gentiana* spp., *Erythraea centaurium* (preferred genus name *Centaurium*), and several *Swertia* spp. (Gentianaceae). Yellow prisms. Mp 218-220° dec.

Semicarbazone: Mp 221-223° dec.

Oxime: Mp 203-205° dec.

Marekov, N.L. *et al.*, *Tetrahedron*, 1968, **24**, 1323 (*uv, ir, pmr, struct*)

Geotrichum alkaloid A

G-64

25822H

Antibiotic A 25822H. A 25822H

[55466-14-3]

C₂₈H₄₃N₃O₂ 425.653

Struct. unknown. Minor metab. of *Geotrichum flavo-brunneum* NRRL 3862. Amorph. Sol. MeOH; poorly sol. H₂O. [α]_D²⁵ +15 (c, 0.147 in MeOH). pK_a 6.9 (66% DMF). Strongly hygroscopic. λ_{max} 277 (ε 13400) (EtOH/HCl) (Derep). λ_{max} 240 (ε 12300); 270 (sh) (EtOH) (Derep). λ_{max} 235 (ε 10000) (EtOH) (Berdy). λ_{max} 270 (ε 8700) (EtOH-HCl) (Berdy).

Chamberlin, J.W. *et al.*, *J. Antibiot.*, 1975, **28**, 102-111 (*isol, uv, ir, pmr*)

Gordee, R.S. *et al.*, *J. Antibiot.*, 1975, **28**, 112-117 (*activity*)


Gephyrotoxin

G-65

Histrionicotoxin D. Dendrobates Alkaloid 287C

[55893-12-4]

[75685-48-2 ((±)-form)]



Absolute Configuration

C₁₉H₂₉NO 287.444

Abs. config. derived chemically, as shown above, is at variance with the crystallographic determination. Minor alkaloid from skin extracts of the neotropical

poison-frog *Dendrobates histrionicus* (Dendrobatidae). Appears to be a weak muscarinic antagonist and acetylcholine antagonist. Mp 231-232° dec. [α]_D -51.5 (c, 1.0 in EtOH).

4',5'-Dihydro: **Dihydrogephyrotoxin**.

Dendrobates Alkaloid 289B

[63983-58-4 ((±)-form)]

C₁₉H₃₁NO 289.46

Minor constit. in *Dendrobates histrionicus* (Dendrobatidae). Terminal triple bond reduced to double.

Tokuyama, T. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 2597-2604 (*isol, pmr, ms, uv, ir*)

Daly, J.W. *et al.*, *Helv. Chim. Acta*, 1977, **60**,

1128-1140 (*isol, cryst struct, pmr, ms, deriv*)

Fujimoto, R. *et al.*, *J.A.C.S.*, 1980, **102**, 7154-7176 (*synth*)

Hart, D.J. *et al.*, *J.O.C.*, 1981, **46**, 3576-3578 (*synth*)

Fujimoto, R. *et al.*, *Tet. Lett.*, 1981, **22**, 4197-4198 (*abs config*)

Hart, D.J. *et al.*, *J.A.C.S.*, 1983, **105**, 1255-1263 (*synth*)

Daly, J.W. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1986, **4**, 110 (*rev, pharmacol*)

Pearson, W.H. *et al.*, *J.O.C.*, 2000, **65**, 7158-7174 (*synth*)

Wei, L.-L. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 1516-1518 (*synth*)

Santarem, M. *et al.*, *J.O.C.*, 2008, **73**, 6466-6469 (*synth*)

Geralbine

G-66

C₂₂H₃₃N₂O₂ 343.508

Steroidal alkaloid. Struct. unknown. Alkaloid from *Veratrum album* (Liliaceae). Plates (EtOAc/Et₂O), prisms (Me₂CO). Mp 221-223°. No measurable opt. rotn. in several solvs. Dec. slowly in soln. Ir shows N-Me and C=O groups.


Hydrochloride: Mp 270°.

Stoll, A. *et al.*, *J.A.C.S.*, 1952, **74**, 4728-4729

Jaspersen-Schib, R. *et al.*, *Pharm. Acta Helv.*, 1961, **36**, 461-471; *CA*, **56**, 1525h

Gerambullin

G-67



C₂₂H₂₉N₂O₅S 419.541

N-Me: **Methylgerambullin**

C₂₃H₃₁N₂O₅S 433.568

Isol. from leaves of *Glycosmis ovoidea*.

1-Hydroxy: **β-Hydroxygerambullin**

C₂₂H₂₉N₂O₆S 435.54

Alkaloid from *Glycosmis chlorosperma*.

Oil. [α]_D²⁰ +21 (c, 0.4 in CHCl₃). λ_{max} 227 ; 275 (sh) (MeOH).

Hofer, O. *et al.*, *Annalen*, 1995, 1789-1794


(*Methylgerambullin*)

Hofer, O. *et al.*, *Phytochemistry*, 2000, **54**, 207-213 (*β-Hydroxygerambullin*)

Gerambullin

G-68

[160896-53-7]



C₂₂H₃₁NO₄S 405.557

Alkaloid from leaves of *Glycosmis angustifolia* (Rutaceae). Mp 139-140°. Authors' numbering system used.

N-Me: Methylgerambullin

[160896-54-8]

C₂₃H₃₃NO₄S 419.584

From leaves of *Glycosmis angustifolia*. Mp 82-85°.

5'-Oxo, N-Me: Methylgerambullone

[160954-12-1]

C₂₃H₃₁NO₅S 433.568

From leaves of *Glycosmis angustifolia*. Mp 128-130°.

5'-Oxo, Δ^{3',4'}-isomer, N-Me: Methylisogerambullone

[160954-13-2]

C₂₃H₃₁NO₅S 433.568

From leaves of *Glycosmis angustifolia*. Oil.

1-Hydroxy: β-Hydroxygerambullin

C₂₂H₃₁NO₅S 421.557

Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 126-128°. [α]_D²⁰ +25 (c, 0.2 in CHCl₃). λ_{max} 224 ; 273 (sh) (MeOH).

8'-Hydroxy: Gerambullol

C₂₂H₃₁NO₅S 421.557

Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 128-129°. λ_{max} 224 (sh) ; 275 (sh) (MeOH).

12-Hydroxy: Sakerine

[173867-24-8]

C₂₂H₃₁NO₅S 421.557

From leaves of *Glycosmis craibii*. Cryst. Mp 89-91°.

12-Hydroxy, 5'-oxo: Sakerone

[173867-27-1]

C₂₂H₂₉NO₆S 435.54

From leaves of *Glycosmis mauritiana*.

1,8'-Dihydroxy: β-Hydroxygerambullol

C₂₂H₃₁NO₆S 437.556

Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 131-133°. [α]_D²⁰ +38 (c, 0.2 in CHCl₃). λ_{max} 224 (sh) ; 273 (sh) (MeOH).

5',12-Dihydroxy: Sakerol

[173867-25-9]

C₂₂H₃₁NO₆S 437.556

From leaves of *Glycosmis mauritiana*. Cryst. Mp 52-54°. [α]_D²⁰ -74 (c, 0.035 in CHCl₃).

5',12-Dihydroxy, 2',3'-dihydro: Dihydroisosakerol

[173906-99-5]

C₂₂H₃₃NO₆S 439.572

From leaves of *Glycosmis craibii*. [α]_D²⁰ -5.9 (c, 0.54 in CHCl₃).

6',7'-Dihydroxy, 6',7'-dihydro: Gerambullindiol

[160896-55-9]

C₂₂H₃₃NO₆S 439.572

From leaves of *Glycosmis angustifolia*. Mp 111-112°. [α]_D⁴³⁶ -6 (c, 0.1 in CDCl₃).

8',12-Dihydroxy: Sakerinol A

C₂₂H₃₁NO₆S 437.556

Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 133-135°. λ_{max} 225 (sh) ; 275 (MeOH).

9',12-Dihydroxy: Sakerinol B

C₂₂H₃₁NO₆S 437.556

Alkaloid from *Glycosmis* ex aff. *pseudoracemosa*. Cryst. (Et₂O). Mp 98-100°. λ_{max} 225 (sh) ; 275 (MeOH).

12-Methoxy, 8'-hydroxy: O-Methylsakerinol A. Methoxygerambullol

C₂₃H₃₃NO₆S 451.583

Alkaloid from *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 125-127°. λ_{max} 227 (sh) ; 275 (MeOH).

Greger, H. et al., *Phytochemistry*, 1994, **37**, 1305-1310 (isol, uv, ir, pmr, cmr, ms, struct)
Hofer, O. et al., *Annalen*, 1995, 1789-1794 (Sakerine, Sakerone, Sakerol, Dihydroisosakerol)


Hinterberger, S. et al., *Tetrahedron*, 1998, **54**, 487-496 (synth)

Hofer, O. et al., *Phytochemistry*, 2000, **54**, 207-213 (*Glycosmis chlorosperma* derivis)

1(10),4-Germacradien-11-amine

G-69

11-Amino-1(10),4-germacradiene



C₁₅H₂₇N 221.385

1(10)Z,4Z)-form [674817-51-7]

Constit. of *Axinyssa* n. sp. Oil. [α]_D²⁵ +28 (c, 0.2 in CHCl₃).


Satitpatipan, V. et al., *J. Nat. Prod.*, 2004, **67**, 503-505 (isol, pmr, cmr)

Germinine

G-70

4,9-Epoxy-3,4,7,14,15,16,20-cevaneheptol, 9CI

[508-65-6]



Absolute configuration

C₂₇H₄₃NO₈ 509.639

Alkaloid from *Veratrum viride* and other *Veratrum* spp. and *Zygadenus venenosus* (Liliaceae). Potent hypotensive agent with characteristic heart action, causing irregularity and prolongation of the beat. Also affects nerve impulses. Derivs. have been used to treat myasthenia gravis. Esters are hypotensive agents, with Germinine being the most active of the nat. alkaloids. Many semisynthetic esters have also been evaluated. They show strong emetic activity with freq. an unacceptably narrow range of therapeutic usefulness. Mp 219-220°. [α]_D²⁵ +5 (EtOH). Forms MeOH solvate which loses MeOH at

163-173°.

▶LD₅₀ (mus, ivn) 139 mg/kg. FL4725000
3,4,7,15,16-Penta-Ac: Mp 257-258° dec. [α]_D²⁴ -88.8 (Py).

3-O-(2R-2-Methylbutanoyl): Protoveratridine

[465-78-1]

C₃₂H₅₁NO₉ 593.756

Alkaloid from *Zygadenus venenosus*, *Veratrum album* and *Veratrum viride* (Liliaceae). Mp 272-273°. [α]_D²² -9 (c, 0.76 in Py).

15-O-(2R-Methylbutanoyl): 2-Methylbutyrylgerminine

[42138-61-4]

C₃₂H₅₁NO₉ 593.756

Alkaloid from *Veratrum lobelianum* (Liliaceae). Mp 224-226°. [α]_D -21.5 (Py).

15-O-(2R-Methylbutanoyl), 3-Ac: Germinidine

[465-77-0]

C₃₄H₅₃NO₁₀ 635.793

Alkaloid from *Veratrum viride*, *Veratrum lobelianum* and *Veratrum nigrum* (Liliaceae). Insecticide against cockroaches. Moderate hypotensive agent. Mp 198-200° Mp 202-203° Mp 242-244° (double Mp). [α]_D +13 (c, 1.67 in CHCl₃).

▶LD₅₀ (mus, ipr) 10 mg/kg. FL4900000

15-O-(2R-Methylbutanoyl), 7-Ac: Neogerminidine. Isogerminidine

[560-49-6]

C₃₄H₅₃NO₁₀ 635.793

Alkaloid from *Zygadenus venenosus* (Liliaceae). Mp 221-223°. [α]_D²² -60 (c, 2 in Py).

15-O-(2R-Methylbutanoyl), 3,7-di-Ac: Neogermitrine

[508-66-7]

C₃₆H₅₅NO₁₁ 677.831

Alkaloid from *Veratrum viride* and *Zygadenus venenosus* (Liliaceae). Strong hypotensive agent. Mp 237-239° (230-231°). [α]_D²³ -78 (c, 0.38 in Py).

▶QP2625000

16-O-(2R-Methylbutanoyl): Verussurinine

[135636-53-2]

C₃₂H₅₁NO₉ 593.756

Alkaloid from the rhizoma and roots of *Veratrum nigrum* var. *ussuriense* (Liliaceae). Amorph. [α]_D²³ -28.4 (c, 0.48 in Py).

3,15-Bis-O-(2-methylbutanoyl): 3,15-Bis(2-methylbutyryl)germinine

[175030-77-0]

C₃₇H₅₉NO₁₀ 677.874

Alkaloid from *Veratrum lobelianum*. Cryst. (C₆H₆). Mp 214-216°. [α]_D -25.15 (c, 0.676 in Py).

3,15-Bis-O-(2-methylbutanoyl), 7-Ac: Germinine

C₃₉H₆₁NO₁₁ 719.911

Alkaloid from *Veratrum lobelianum*. Cryst. (Me₂CO/hexane). Mp 195-197°. Error in CAS.

3-Angeloyl, 15-O-(2R-methylbutanoyl), at

- 7-Ac: Germanitrine**
[639-11-2]
C₃₉H₅₉NO₁₁ 717.895
Alkaloid from *Veratrum fimbriatum* (Liliaceae). Strong hypotensive agent. Mp 228-229°. [α]_D²⁴ -61 (c, 1 in Py).
- 15-Angeloyl: 15-Angeloylgerminine**
[240802-94-2]
C₃₂H₄₉NO₉ 591.74
Alkaloid from *Veratrum taliense*. Needles (Me₂CO/C₆H₆). Mp 219-221°. [α]_D²⁵ +5.5 (c, 0.43 in MeOH).
- 15-Angeloyl, 3-Ac: 3-Acetyl-15-angeloylgerminine**
[942293-83-6]
C₃₄H₅₁NO₁₀ 633.778
Alkaloid from the rhizomes of *Veratrum dahuricum*. Amorph. powder (Me₂CO). Mp 198-200°. [α]_D²⁰ -8 (c, 0.09 in CHCl₃).
- 15-Angeloyl, 3,7-di-Ac: 3,7-Diacetyl-15-angeloylgerminine**
[942293-81-4]
C₃₆H₅₃NO₁₁ 675.815
Alkaloid from the rhizomes of *Veratrum dahuricum*. Amorph. powder (Me₂CO). Mp 231-233°. [α]_D²⁰ +2.8 (c, 0.08 in CHCl₃).
- 3,15-Diangeloyl, 7-Ac: 7-Acetyl-3,15-diangeloylgerminine. Maackininine**
[122332-72-3]
C₃₉H₅₇NO₁₁ 715.879
Minor alkaloid from *Veratrum maackii* (Liliaceae). Prisms. Mp 218-221°. [α]_D +3.85 (c, 0.65 in CHCl₃).
- O-Angeloyl, O-tigloyl, O-Ac: Germinitrine**
[465-76-9]
C₃₇H₅₇NO₁₁ 691.857
Alkaloid from *Veratrum fimbriatum* (Liliaceae). Hypotensive agent. Mp 175°. [α]_D²⁴ -36 (c, 1.12 in Py).
- 3,15-Bis-O-(2-methyl-2-butenoyl): Stenophylline A**
[90541-57-4]
C₃₇H₅₅NO₁₀ 673.842
Alkaloid from *Veratrum stenophyllum* (Liliaceae). Exhibits hypotensive activity.
- 3-O-(2-Hydroxy-2-methylbutanoyl), 15-O-(2-methylbutanoyl): Neogerminine**
[134357-41-8]
C₃₇H₅₉NO₁₁ 693.873
Alkaloid from *Veratrum nigrum*. Same gross struct. as Germinine, stereochemical variation may occur in the acyl groups.
- 3-O-(2R-Hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl): Germinine. Veratensine**
[508-67-8]
C₃₇H₅₉NO₁₁ 693.873
Alkaloid from *Veratrum lobelianum*, *Veratrum oxysepalum*, *Veratrum album*, *Veratrum nigrum* and *Zygadenus* sp. Strong hypotensive agent. Mp 200-203°. [α]_D -7 (Py). λ_{max} 243 (ε 630) (MeOH) (Berdy).
- 3-O-(2R-Hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 7-Ac: Germitrine**
[560-48-5]
C₃₉H₆₁NO₁₂ 735.91
Alkaloid from *Veratrum viride* (Liliaceae). V. strong hypotensive agent. Mp 197-199°. [α]_D²⁵ +11 (CHCl₃).
- ▶ FL4800000
- 3-O-(2R,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl): Germbudine**
[426-34-6]
C₃₇H₅₉NO₁₂ 709.873
Alkaloid from *Veratrum viride* (Liliaceae). Hypotensive agent. Mp 160-164° (157-159°). [α]_D²⁴ -7 (c, 1 in Py). [α]_D +8.02.
- 3-O-(2R,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 7-Ac: Germinalinine**
[58162-51-9]
C₃₉H₆₁NO₁₃ 751.91
Alkaloid from *Veratrum lobelianum* (Liliaceae). Mp 168-170°. [α]_D -52.1 (Py).
- 3-O-(2S,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl): Neogermbudine**
[595-64-2]
C₃₇H₅₉NO₁₂ 709.873
Alkaloid from *Veratrum viride* (Liliaceae). Mp 149-152°. [α]_D²⁵ -12 (c, 1 in Py).
- 3-O-(2S-Acetoxy-3R-hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl): Germinaline**
[23211-84-9]
C₃₉H₆₁NO₁₃ 751.91
Alkaloid from *Veratrum lobelianum* (Liliaceae). Cryst. (C₆H₆), needles (Et₂O). Mp 138-140°, 156-158° (dimorph.). [α]_D +3.7 (c, 1.06 in CHCl₃). Incorrect struct. prev. proposed.
- 3-O-(2S-Acetoxy-3R-hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 7-Ac: Germitetrine. Germitetrine B**
[465-75-8]
C₄₁H₆₃NO₁₄ 793.947
Alkaloid from *Veratrum album* and *Veratrum lobelianum* (Liliaceae). Cryst. (Me₂CO aq.). Mp 233-234°. [α]_D²⁵ -74 (c, 1 in Py). [α]_D -12 (c, 1.0 in CHCl₃).
- 3-O-(3,4-Dimethoxybenzoyl): 3-Veratroylgerminine**
[942293-82-5]
C₃₆H₅₁NO₁₁ 673.799
Alkaloid from the rhizomes of *Veratrum dahuricum*. Amorph. powder (MeOH). Mp 280-283°. [α]_D²⁰ +13 (c, 0.08 in CHCl₃).
- 3-O-(3,4-Dimethoxybenzoyl), 15-O-angeloyl: 15-Angeloyl-3-veratroylgerminine**
[942293-84-7]
C₄₁H₅₇NO₁₂ 755.901
Alkaloid from the rhizomes of *Veratrum dahuricum*. Amorph. powder (Me₂CO). Mp 281-283°. [α]_D²⁰ +18 (c, 0.08 in CHCl₃).
- 15-O-(3,4-Dimethoxybenzoyl): 15-Veratroylgerminine**
[33352-59-9]
C₃₆H₅₁NO₁₁ 673.799
Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Amorph.
- 15-O-(3,4-Dimethoxybenzoyl), 3-Ac: 3-Acetyl-15-veratroylgerminine**
[33352-58-8]
C₃₈H₅₃NO₁₂ 715.836
Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Amorph. [α]_D²⁵ +6.6 (CHCl₃).
- 3-O-(3,4-Dimethoxybenzoyl), 15-O-(2-methylbutanoyl): Verabenzoamine**
[142735-72-6]
C₄₁H₅₉NO₁₂ 757.917
Alkaloid from *Veratrum nigrum* var. *ussuriense*. Amorph. solid. [α]_D²³ +8.7 (c, 0.4 in CHCl₃). Struct. revised in 1998 (prev. assigned struct. now given to Verussurine).
- 3-O-(3,4-Dimethoxybenzoyl), 15-O-(2-methylbutanoyl), 7-Ac: Verussurine**
[214046-03-4]
C₄₃H₆₁NO₁₃ 799.954
Alkaloid from *Veratrum nigrum* var. *ussuriense* (Liliaceae). Amorph. powder. [α]_D²³ +8 (c, 1 in CHCl₃).
- 1α-Acetoxy, 3-O-(2ξ-hydroxy-2-methylbutanoyl), 15-O-(2-methylbutanoyl): Neojerminalanine**
[182693-36-3]
C₃₉H₆₁NO₁₃ 751.91
Alkaloid from rhizomes of *Veratrum album*. Amorph. powder. [α]_D²⁵ -44 (c, 0.05 in Py).
- [36506-68-0]
- Nash, H.A. *et al.*, *J.A.C.S.*, 1953, **75**, 1942-1948 (*Germitetrine B*)
Kupchan, S.M. *et al.*, *J.A.C.S.*, 1953, **75**, 4671-4672 (*Germitetrine B*)
Klohs, M.W. *et al.*, *J.A.C.S.*, 1953, **75**, 4925-4927 (*Germanitrine, Germinitrine*)
Myers, G.S. *et al.*, *J.A.C.S.*, 1955, **77**, 3348-3353; 1956, **78**, 1621-1624 (*Neogerminine, Germbudine*)
Kupchan, S.M. *et al.*, *Chem. Ind. (London)*, 1958, **48**, 1594-1595 (*Germitetrine, struct*)
Kupchan, S.M. *et al.*, *J. Am. Pharm. Assoc.*, 1959, **48**, 440-442 (*Germitetrine, Neogermbudine*)
Kupchan, S.M. *et al.*, *J.A.C.S.*, 1959, **81**, 1913-1921; 1921-1924 (*struct, bibl*)
Tomko, J. *et al.*, *Chem. Zvesti*, 1971, **25**, 69 (*15-Veratroylgerminine, 3-Acetyl-15-veratroylgerminine*)
Shakirov, R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 606-607; 1975, **11**, 566-567; 1983, **19**, 120-121 (*Germinine, Germinaline, Germinalinine*)
Carey, F.A. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 141-144 (*cmr*)
Liang, G. *et al.*, *Yaoxue Xuebao*, 1984, **19**, 190-194; *CA*, **103**, 51156h (*Stenophylline A*)
Zhao, W. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 2920-2928; 1991, **39**, 549-554 (*Maackininine, Verussurinine*)
Han, X. *et al.*, *Magn. Reson. Chem.*, 1991, **29**, 100-112 (*cmr, struct*)
Shakirov, R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 79-81 (*bis(2-methylbutanoyl)*)
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1996, **43**, 907-911 (*Neojerminalanine*)
Shakirov, R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 479-480 (*Germinine*)
Tezuka, Y. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1397-1399 (*Verabenzoamine, Verussurine*)
Zhou, C.X. *et al.*, *Planta Med.*, 1999, **65**, 480-482 (*15-Angeloylgerminine*)
Tang, J. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 769-775 (*Veratrum dahuricum esters*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, EBL500

Gerrardamine

G-71

[11013-56-2]


C₈H₁₅NOS₂ 205.345

Struct. unknown. Alkaloid from *Cassipourea gerrardii* (Rhizophoraceae). Gum. Wright, W.G. *et al.*, *J.C.S. (C)*, 1967, 283-284 (*isol, ir*)

Gerrardine

G-72

3,3'-(1-Methyl-2,5-pyrrolidinediyl)bis-1,2-dithiolan-4-ol, 9CI
[14022-19-6]



Relative Configuration

C₁₁H₁₉NO₂S₄ 325.541

Alkaloid from *Cassipourea gerrardii* and *Cassipourea guianensis* (Rhizophoraceae). Antifungal agent active against *Candida albicans*. Yellow orthorhombic cryst. Sol. MeOH, CHCl₃. Mp 180°. Cryst. from EtOH as solvate, Mp 90° and 178°. Cryst. from C₆H₆ as unstable solvate. λ_{max} 330 (ε 300) (MeOH) (Derp). λ_{max} 336 (MeOH) (Berdy).

Hydrochloride: Mp 207° dec. (180°). [α]_D²³ -172 (c, 1 in H₂O). Sublimes at 195°.

l'-S-Oxide: [97094-26-3]

C₁₁H₁₉NO₃S₄ 341.54

Minor alkaloid from the bark of *Cassipourea guianensis* (Rhizophoraceae). Mp 183-185° dec. [α]_D²⁴ +140.66 (c, 0.39 in CHCl₃). λ_{max} 247 (ε 3630); 330 (ε 148) (CHCl₃) (Derp).

l',l''-S-Dioxide: [97094-27-4]

C₁₁H₁₉NO₄S₄ 357.539

Minor alkaloid from *Cassipourea guianensis* (Rhizophoraceae). Mp 190-193° dec. λ_{max} 247 (ε 6760) (CHCl₃) (Derp).

Wright, W.G. *et al.*, *J.C.S. (C)*, 1967, 283; 284 (*isol, ir, ms, struct*)

Gafner, G. *et al.*, *Acta Cryst. B*, 1971, 27, 565 (*cryst struct*)

Kato, A. *et al.*, *J. Nat. Prod.*, 1984, 47, 706; 1985, 48, 289-292 (*isol, pmr, cmr, oxides*)

Gerrardoline

G-73

[11013-58-4]

C₈H₁₅NO₂S₂ 221.344


Struct. unknown. Alkaloid from *Cassipourea gerrardii* (Rhizophoraceae). Non-cryst. yellow solid. Mp 60°.

Wright, W.E. *et al.*, *J.C.S. (C)*, 1967, 283-284 (*isol, ir*)

Gigantamide A

G-74

[955381-63-2]

C₁₈H₂₂N₂O₃ 314.383


Alkaloid from the leaves of *Aglaia gigantea*. Amorph. solid. [α]_D²⁰ -10 (c, 0.2 in MeOH). λ_{max} 217 (ε 15000); 275 (ε 13510) (MeOH).

Duong, T.N. *et al.*, *J. Nat. Prod.*, 2007, 70, 1640-1643 (*isol, pmr, cmr*)

Gigartinine

G-75

N⁵-[[1-(Aminoiminomethyl)amino]carbonyl]ornithine, 9CI. 5-(3-Amidinoureido)-2-aminovaleric acid

C₇H₁₅N₅O₃ 217.227**(S)-form** [7536-90-5]

[7536-92-7 (nitrate)]

Constit. of many spp. of red algae, e.g. *Gymnogongrus flabelliformis*, *Gelidium amansii*, *Grateloupia livida*, *Grateloupia filicina*, *Polyopes polydeoides*, *Carpopeltis flabellata*, *Hypnea japonica* and *Gracilaria textorii*. Plates (EtOH)(as nitrate). Mp 197°. [α]_D²² +7.5 (c, 2.0 in H₂O).

Ito, K. *et al.*, *Nature (London)*, 1966, 211, 417 (*isol, uv, ir*)

Ito, K. *et al.*, *CA*, 1968, 68, 87527c; 69, 93635y (*isol*)


Laycock, M.V. *et al.*, *Can. J. Biochem.*, 1977, 55, 27 (*isol, pmr, cmr*)

Wakamiya, T. *et al.*, *Tetrahedron*, 1984, 40, 235 (*isol*)

Gilbertine

G-76

1,17-Epoxy-1-methylasycarpidan, 9CI
[85769-42-2]

C₁₈H₂₂N₂O 282.385

Alkaloid from the bark of *Aspidosperma gilbertii* (Apocynaceae). Mp 118-122°. [α]_D²⁵ -149 (CHCl₃).

Miranda, E.C. *et al.*, *Tet. Lett.*, 1982, 23, 5395 (*ir, pmr, cmr, ms, struct*)

Jiricek, J. *et al.*, *J.A.C.S.*, 2004, 126, 3534-3538 (*synth*)

Gindaricine

G-77

Hindaricine

[11075-74-4]

C₁₈H₁₉NO₃ 297.353

Struct. unknown. Hindaricine is a trans-literation error of Gindaricine, via a Russian paper. Alkaloid from tubers of *Stephania glabra* (Menispermaceae). Cryst. (C₆H₆). Mp 193°. [α]_D +137

(EtOH). No further reports to 2007. Not obt. in further investigations of this species.

Hydrochloride: Mp 232°.


Picrate: Mp 150° dec.

Chaudhry, G.R. *et al.*, *J. Sci. Ind. Res., Sect. B*, 1950, 9, 79-83; *CA*, 45, 824a (*isol*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, 33, 2785-2789 (*Stephania glabra constits*)

Ginsenine

G-78



Relative Configuration

C₁₃H₁₄N₂O₂ 230.266


Alkaloid from the fruit of *Panax ginseng*. Powder. [α]_D²⁰ -13.3 (c, 0.36 in MeOH). λ_{max} 221 (log ε 3.42); 272 (log ε 2.75); 278 (log ε 2.74); 289 (log ε 2.63) (MeOH).

Wang, J.-Y. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, 8, 605-608 (*isol, pmr, cmr, ms*)

Giracodazole, INN

G-79

2-Amino-α-(2-amino-1-chloroethyl)-1H-imidazole-4-methanol, 9CI. 4-Amino-1-(2-amino-4-imidazolyl)-2-chloro-1-butanol. **Girolline**. NSC 627434. RP 49532



(2'S,3'S)-form

C₆H₁₁ClN₄O 190.632

Protein synthesis inhibitor, antineoplastic agent. Sol. MeOH, EtOH.

(2'S,3'S)-form [135824-74-7]

Isol. from the sponge *Pseudaxinyssa cantharella*. Powder (as hydrochloride). [α]_D²⁰ +7.9 (c, 0.84 in MeOH). Log P -2.03 (calc). CAS no. refers to hydrochloride.

(2'RS,3'SR)-form [127909-96-0]

Mp 191-193° (hydrochloride).

[132618-95-2, 117678-06-5, 117678-08-7, 110883-46-0]

Ahond, A. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1988, 307, 145 (*isol*)

Bedoya Zurita, M. *et al.*, *Tetrahedron*, 1989, 45, 6713 (*synth, pmr*)

Chiaroni, A. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1991, 312, 49 (*cryst struct, abs config*)

Commercon, A. *et al.*, *Tet. Lett.*, 1991, 32, 1419; 4905 (*synth, config*)

Colson, G. *et al.*, *Biochem. Pharmacol.*, 1992, 43, 1717 (*pharmacol*)


Ahond, A. *et al.*, *Tetrahedron*, 1992, 48, 4327 (*synth*)

Marchais, S. *et al.*, *Tet. Lett.*, 1998, 39, 8085-8088 (*synth*)

Girgensonine

G-80

4-Hydroxy- α -(1-piperidinyl)benzeneacetone nitrile. N-Piperidyl-p-hydroxyphenylacetone nitrile. *Girgensonine* [486-30-6]



$C_{13}H_{16}N_2O$ 216.282

Alkaloid from green parts of *Girgensonia oppositiflora* (Chenopodiaceae).

Cryst. (Et₂O or MeOH). Mp 147-148°.

Hydrochloride: Mp 145-148°.


Picrolonate: Mp 192-194°.

Yurashevskii, N.K. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1946, **16**, 141; *CA*, **40**, 6754 (isol, struct, synth)

Girinimbine

G-81

3,11-Dihydro-3,3,5-trimethylpyrano[3,2-a]carbazole, 9CI [23095-44-5]



$C_{18}H_{17}NO$ 263.338

CAS numbering shown. Alternative numbering frequently used in which the carbazole nucleus is numbered 1-9 and the pyran carbons 1'-5'. Alkaloid from the roots of *Murraya koenigii* (curry leaf tree) and *Clausena heptaphylla*, and from the stem bark of *Murraya exotica* (Rutaceae). Cryst. (cyclohexane, hexane or CH₂Cl₂/hexane). Mp 175°. λ_{max} 237 (log ϵ 4.76); 278 (log ϵ 4.53); 315 (log ϵ 3.88); 328 (log ϵ 4); 340 (log ϵ 4.03); 355 (log ϵ 3.98) (no solvent reported).

N-Me:

Cryst. (EtOH). Mp 150°.

8-Hydroxy: *Koenine*. *Kenine*

[28200-63-7]

$C_{18}H_{17}NO_2$ 279.338

Alkaloid from the leaves of *Murraya koenigii* (curry leaf tree) (Rutaceae).

Cryst. (C₆H₆). Mp 250-252°. λ_{max} 230 (log ϵ 4.65); 236 (log ϵ 4.64); 296 (log ϵ 4.41); 335 (log ϵ 3.84); 356 (log ϵ 3.72) (MeOH).

8-Hydroxy, N-Me: Mp 148°.

8-Methoxy: *Koenimbine*. *Kenimbine*

[21087-98-9]

$C_{19}H_{19}NO_2$ 293.365

Alkaloid from leaves and fruits of *Murraya koenigii* (curry leaf tree) and from stem bark of *Murraya exotica* (Rutaceae). Cryst. (hexane or CHCl₃/hexane). Mp 194-195°. λ_{max} 230 (log ϵ 4.68); 240 (log ϵ 4.69); 300 (log ϵ 4.62); 340 (log ϵ 4.24); 360 (log ϵ 4.2) (MeOH).

9-Hydroxy: *Murrayamine A*. *Mukoeneine C*

[134779-17-2]

$C_{18}H_{17}NO_2$ 279.338

Alkaloid from the leaves of *Murraya euchrestifolia* and roots of *Murraya koenigii* (curry leaf tree) (Rutaceae). Cytotoxic. Needles (Me₂CO). Mp 162-163°. *Mukoeneine C* from *M. koenigii* descr. as an oil. λ_{max} 221 (ϵ 10450); 240 (ϵ 11750); 294 (ϵ 9100); 342 (ϵ 2240); 349 (ϵ 1600); 360 (ϵ 1620) (MeOH) (Berdy).

10-Methoxy: *Mupamine*

[66003-49-4]

$C_{19}H_{19}NO_2$ 293.365

Alkaloid from the root bark of *Clausena anisata* (Rutaceae). Mp 151-152°. λ_{max} 238 (log ϵ 4.8); 263 (log ϵ 4.31); 272 (log ϵ 4.33); 282 (log ϵ 4.43); 315 (sh) (log ϵ 3.75); 330 (log ϵ 3.89); 342 (log ϵ 3.94); 357 (log ϵ 3.9) (MeOH).

1'-Acetoxy: *Murrayamine K*

[175669-19-9]

$C_{20}H_{19}NO_3$ 321.375

Alkaloid from leaves of *Murraya euchrestifolia*. Needles (Me₂CO). Mp 127-128°. λ_{max} 237; 279 (sh); 288; 327; 344; 359 (MeOH).

9-Hydroxy, 1'-acetoxy: *Murrayamine I*

[175669-18-8]

$C_{20}H_{19}NO_4$ 337.374

Alkaloid from leaves of *Murraya euchrestifolia*. Oil. λ_{max} 222; 240; 281 (sh); 295; 324; 360 (MeOH).

8-Methoxy, 9-hydroxy: *Koenigine*. *Kenigine*

[28513-33-9]

$C_{19}H_{19}NO_3$ 309.364

Alkaloid from the leaves of *Murraya koenigii* (curry leaf tree), and from the leaves and stems of *Micromelum zeylanicum* (Rutaceae). Cryst. (EtOH), needles (hexane/EtOH). Mp 183-185°. λ_{max} 243 (log ϵ 4.47); 260 (log ϵ 4.05); 275 (log ϵ 3.94); 290 (log ϵ 4.36); 300 (log ϵ 3.53); 344 (log ϵ 3.99); 357 (log ϵ 3.9) (MeOH).

8,9-Dimethoxy: *Koenigicine*. *Koenidine*.

Koenimbicine. *Kenigicine*. *Kenidine*.

Kenimbicine

[24123-92-0]

$C_{20}H_{21}NO_3$ 323.391

Alkaloid from the leaves of *Murraya koenigii* (curry leaf tree) (Rutaceae). Cryst. (EtOH). Mp 224-225°. Subl. at 210°/0.5 mm. λ_{max} 230 (log ϵ 4.68); 240 (log ϵ 4.69); 300 (log ϵ 4.62); 340 (log ϵ 4.24); 360 (log ϵ 4.2) (MeOH).

8,9-Dimethoxy, N-Me:

Cryst. (EtOH). Mp 190-191°.

8,10-Dimethoxy: *Mukonicine*

[88607-43-6]

$C_{20}H_{21}NO_3$ 323.391

Alkaloid from the leaves of *Murraya koenigii* (curry leaf tree) (Rutaceae). Cryst. (C₆H₆/CHCl₃). Mp 233-234°. λ_{max} 226 (log ϵ 4.7); 240 (log ϵ 4.67); 300 (log ϵ 4.59); 342 (log ϵ 4.26) (EtOH).

Narasimhan, N.S. et al., *Tet. Lett.*, 1968, 5501-5504 (*Koenimbine*, isol, uv, pmr, ms, struct)

Dutta, N.L. et al., *Indian J. Chem.*, 1969, **7**, 307-308 (uv, pmr, struct)

Kureel, S.P. et al., *Chem. Ind. (London)*, 1970, 1262 (*Koenine*, *Koenimbine*, *Girinimbine*, synth)

Joshi, B.S. et al., *Tetrahedron*, 1970, **26**, 1475-1482 (*Girinimbine*, *Koenigine*, *Koenigicine*, *Koenine*, *Koenimbine*, isol, uv, ir, pmr)

Joshi, B.S. et al., *Phytochemistry*, 1972, **11**, 2065-2071 (isol, uv, ir, pmr)

Narasimhan, N.S. et al., *Indian J. Chem., Sect. B*, 1975, **13**, 993-999; 1976, **14**, 329-331 (*Koenine*, *Koenimbine*, *Koenigicine*,

Girinimbine, isol, uv, ir, pmr, ms, struct, synth)

Oikawa, Y. et al., *Heterocycles*, 1976, **5**, 233-238 (synth, ms)

Mester, I. et al., *Annalen*, 1977, 1725-1729; 1980, 241-245 (*Mupamine*, isol, ir, uv, pmr, ms, cmr, struct, synth)

Mester, I. et al., *Z. Naturforsch., B*, 1979, **34**, 650-652 (*Mupamine*, cmr)

Sharma, R.B. et al., *Experientia*, 1980, **36**, 815 (*Koenigicine*, synth)

Sharma, R.B. et al., *Indian J. Chem., Sect. B*, 1981, **20**, 701-703 (*Mupamine*, synth)

Bowen, I.H. et al., *Phytochemistry*, 1982, **21**, 433-437 (*Koenigine*, isol, ms)

Mukherjee, M. et al., *Phytochemistry*, 1983, **22**, 2328-2329 (*Mukonicine*)

Wu, T.-S. et al., *Phytochemistry*, 1991, **30**, 1048-1051; 1996, **41**, 1433-1435 (*Murrayamine A*, *Murrayamine I*, *Murrayamine K*)


Ito, C. et al., *Chem. Pharm. Bull.*, 1993, **41**, 2096-2100 (*Mukoeneine C*)

Knölker, H.J. et al., *Tet. Lett.*, 1996, **37**, 7947-7950 (synth)

Glabratine

G-82

[142750-47-8]



$C_{27}H_{34}N_2O_9$ 530.574

Alkaloid from *Cephalanthus occidentalis* and *Uncaria glabrata*. Needles (EtOH). Mp 265-267°. $[\alpha]_D^{25} +43.9$ (c, 0.1 in MeOH). λ_{max} 225 (log ϵ 4.36); 296 (log ϵ 4.59) (MeOH).


Arbain, D. et al., *J.C.S. Perkin 1*, 1992, 665-666 (isol, pmr, cmr)

Zhang, Z. et al., *Planta Med.*, 2005, **71**, 355-361 (isol, pmr, cmr)

Glaciopyrrole A

G-83

[853885-46-8]




$C_{19}H_{27}NO_4$ 333.427

Metab. of a marine-sediment derived *Streptomyces* sp. Glass. $[\alpha]_D^{25} +16.8$ (c, 0.02 in MeOH). λ_{max} 285 (sh) (ϵ 6700); 333 (ϵ 14400) (MeOH).


Macherla, V.R. et al., *J. Nat. Prod.*, 2005, **68**, 780-783 (*Glaciopyrrole A*)

Glaciapyrrole B G-84


[853885-48-0]

C₁₉H₂₇NO₃ 317.427Metab. of a marine-sediment derived *Streptomyces* sp. Glass. λ_{max} 285 (sh); 335 (MeCN aq.).Macherla, V.R. et al., *J. Nat. Prod.*, 2005, **68**, 780-783 (Glaciapyrrole B)**Glaciapyrrole C** G-85


[853885-50-4]

C₁₉H₂₅NO₂ 299.412Constit. of a marine-sediment derived *Streptomyces* sp. Glass. λ_{max} 285 (sh); 335 (MeCN aq.).Macherla, V.R. et al., *J. Nat. Prod.*, 2005, **68**, 780-783 (Glaciapyrrole C)**Glansreginin A** G-86*Glansresinoside A*

[860267-30-7]

C₂₈H₃₅NO₁₃ 593.583Constit. of the seeds of *Juglans regia*. Pale yellow powder + 1.5 H₂O. [α]_D²³ -37.9 (c, 0.1 in MeOH). λ_{max} 265 (log ε 4.45) (MeOH).Ito, H. et al., *J. Agric. Food Chem.*, 2007, **55**, 672-679 (isol, cd, pmr, cmr, ms)**Glaucamine** G-87

2,3-Dimethoxy-16-methyl-10,11-[methylenebis(oxy)]rheadan-8-ol, 9CI. Alkaloid R-L [2255-44-9]

C₂₁H₂₃NO₆ 385.416Various numbering schemes in use. Alkaloid from *Papaver anomalum*, *Papaver fugax*, *Papaver glaucum*, *Papaver hybridum*, *Papaver nudicaule*, *Papaver rhoeas* (corn poppy) and *Papaver tauricola* (Papaveraceae). Plates (CHCl₃/EtOH). Mp 222-223°. [α]_D²² +298 (c, 0.25 in CHCl₃). λ_{max} 238 (log ε 4); 286 (log ε 3.8) (no solvent reported).

Hydrochloride: Mp 201-203°.

Me ether: **Epiglaucine**. O-Methylglaucamine. *Glaupavine*

[18104-26-2]

C₂₂H₂₅NO₆ 399.443Alkaloid from *Papaver glaucum* (as glaupavine) and *Papaver tauricola* (Papaveraceae). [α]_D²² +293 (c, 0.1 in CHCl₃). λ_{max} 237 (log ε 4.18); 287 (log ε 4.06) (MeOH).*Me ether*, N-*de-Me*: **Papaverrubine H**. 14-Epipapaverrubine B. O-Methylpapaverrubine C

[25651-16-5]

C₂₁H₂₃NO₆ 385.416Alkaloid from *Papaver anomalum* (Papaveraceae). Amorph. Mp 70-71°. [α]_D +308 (CHCl₃).*Et ether*: O-Ethylglaucamine

[77658-93-6]

C₂₃H₂₇NO₆ 413.469Isol. from *Papaver fugax* (Papaveraceae). Prob. an artifact.O²-*De-Me*: **N-Methyl-14-O-demethylepiporphyrroxine**. Alkaloid A4

[18361-67-6]

C₂₀H₂₁NO₆ 371.389Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Prisms (MeOH). Mp 217-218°. [α]_D²⁰ +340 (c, 0.2 in MeOH).O²-*De-Me*, O⁸-*Me*, N-*de-Me*: **Papaverrubine C**. Epiporphyrroxine. 14-Epipapaverrubine D

[22584-46-9]

C₂₀H₂₁NO₆ 371.389Alkaloid from *Meconopsis betonicifolia*, *Papaver albiflorum* and other *Papaver* spp. (Papaveraceae). Prisms (C₆H₆/petrol). Mp 190-191.5°. [α]_D +283 (CHCl₃).8-*Epimer*: **Epiglaucamine**

[35321-05-2]

Semisynthetic (from Glaupavine and Glaucamine). Amorph. [α]_D²¹ +126 (c, 0.27 in CHCl₃).8-*Epimer*, *Me ether*: **Glaudine**. O,N-Dimethylporphyrroxine. O,N-Dimethylpapaverrubine D. N-Methylpapaverrubine B

[5140-40-9]

C₂₂H₂₅NO₆ 399.443Alkaloid from *Papaver armeniacum*, *Papaver fugax*, *Papaver glaucum*, *Papaver rhoeas*, *Papaver somniferum* and *Papaver tauricola* (Papaveraceae). Cryst. (heptane). Mp 103-105°. [α]_D²⁴ +455 (c, 0.5 in CHCl₃). λ_{max} 237 (log ε 4.03); 287 (log ε 3.92) (MeOH).8-*Epimer*, *Me ether*, hydrochloride: Mp 185-187°.8-*Epimer*, *Me ether*, N-*de-Me*: **Papaver-****rubine B**. O-Methylporphyrroxine. O-Methylpapaverrubine D. N-Demethylglaucine

[5140-39-6]

C₂₁H₂₃NO₆ 385.416Alkaloid from *Papaver alpinum*, *Papaver atlanticum*, *Papaver bracteatum*, *Papaver californicum*, *Papaver caucasicum*, *Papaver commutatum*, *Papaver dubium*, *Papaver feddei*, *Papaver fugax*, *Papaver glaucum*, *Papaver heldreichii*, *Papaver latericum*, *Papaver macrostomum*, *Papaver nudicaule*, *Papaver oreophilum*, *Papaver orientale*, *Papaver persicum*, *Papaver pilosum*, *Papaver polychaetum*, *Papaver rhoeas*, *Papaver rupifragum*, *Papaver setigerum*, *Papaver strigosum*, *Papaver trinaefolium* and *Papaver urbanianum* (Papaveraceae). Cryst. (MeOH). Mp 201-203°. [α]_D +398 (CHCl₃). λ_{max} 236 (log ε 4.06); 286 (log ε 3.86) (MeOH).8-*Epimer*, O²-*de-Me*, O⁸-*Me*: **N-Methylporphyrroxine**. N-Methylpapaverrubine D

[18211-29-5]

C₂₁H₂₃NO₆ 385.416May occur in *Papaver somniferum* (Papaveraceae).8-*Epimer*, O²-*de-Me*, O⁸-*Me*, N-*de-Me*: **Papaverrubine D**. Porphyrroxine

[18104-24-0]

C₂₀H₂₁NO₆ 371.389Alkaloid from *Meconopsis betonicifolia* and other *Meconopsis* spp., *Papaver albiflorum* and other *Papaver* spp. (Papaveraceae). Needles (MeOH). Mp 237-239° (234-236°). [α]_D +391 (CHCl₃). λ_{max} 232 (log ε 3.9); 287 (log ε 3.83) (MeOH).8,14-*Diepimer*: **Oreogenine**

[6598-99-8]

C₂₁H₂₃NO₆ 385.416Alkaloid from *Papaver fugax*, *Papaver oreophilum* and *Papaver tauricola* (Papaveraceae). Mp 173-175° (as methiodide). Indefinite Mp.8,14-*Diepimer*, *Me ether*: **Oreodine**. N-Methylpapaverrubine F

[6516-48-9]


C₂₂H₂₅NO₆ 399.443Alkaloid from *Papaver fugax*, *Papaver oreophilum*, *Papaver tauricola* and *Papaver trinaefolium* (Papaveraceae). Cryst. (Et₂O). Mp 184-186°. [α]_D²² +224 (c, 0.1 in CHCl₃). λ_{max} 237 (log ε 3.96); 289 (log ε 3.72) (no solvent reported).8,14-*Diepimer*, *Me ether*, N-*de-Me*: **Papaverrubine F**. N-Demethyloreodine

[19914-48-8]

C₂₁H₂₃NO₆ 385.416Alkaloid from *Papaver commutatum* and *Papaver oreophilum* (Papaveraceae). Needles (Me₂CO). Mp 223-225°. λ_{max} 237 (log ε 4.11); 287 (log ε 3.84) (MeOH).8,14-*Diepimer*, *Et ether*: O-EthyloreogenineC₂₃H₂₇NO₆ 413.469Isol. from *Papaver fugax* and *Papaver tauricola* (Papaveraceae). Prob. an artifact.

- Pfeifer, S. et al., *Pharmazie*, 1964, **19**, 678; 724-786; 1965, **20**, 585-586; 1968, **23**, 82-98; 267-281; 1972, **27**, 48 (*Papaverrubines*, *Glaudine*, *Oreodine*, *Oreogenine*, *Epiglaudine*)
- Pfeifer, S. et al., *Pharmazie*, 1964, **19**, 786; 1965, **20**, 585 (*Oreogenine*)
- Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1965, **30**, 3687-3696 (*Epiglaudine*, *Glaudine*, *Glaupavine*, *isol*, *synth*, *uv*)
- Pfeifer, S. et al., *J. Pharm. Pharmacol.*, 1966, **18**, 133-134 (*Papaverrubine D*, *struct*)
- Cross, A.D. et al., *Pharmazie*, 1966, **21**, 181 (*Oreogenine*, *Oreodine*, *stereochem*)
- Mann, I. et al., *Pharmazie*, 1966, **21**, 700 (*Papaverrubine F*)
- Brochmann-Hanssen, E. et al., *J. Pharm. Sci.*, 1967, **56**, 1658 (*Papaverrubine D*, *Oreodine*, *isol*, *ord*, *cd*, *pmr*, *struct*, *config*)
- Dolejš, L. et al., *Tetrahedron*, 1967, **23**, 2997-3005 (*Oreodine*, *Papaverrubines B,D*, *Glaudine*, *ms*)
- Shamma, M. et al., *Chem. Comm.*, 1968, 212-214 (*Papaverrubines*, *Epiglaudine*, *Glaudine*, *stereochem*)
- Brochmann-Hanssen, E. et al., *J. Pharm. Sci.*, 1968, **57**, 30-35 (*N-Methyl-14-O-demethylepiporphyroxine*, *Epiglaudine*, *Papaverrubine D*, *Glaudine*, *isol*, *uv*, *ir*, *pmr*, *struct*)
- Slaviková, L. et al., *Coll. Czech. Chem. Comm.*, 1980, **45**, 761-763 (*Papaverrubines*)
- Sariyar, G. et al., *Phytochemistry*, 1980, **19**, 2189-2192 (*Oreogenine*, *Oreodine*, *Ethylloregene*, *Epiglaudine*, *Glaudine*, *isol*, *pmr*, *ms*)
- Phillipson, J.D. et al., *Planta Med.*, 1981, **41**, 105-118 (*Glaudine*, *Oreodine*, *Oreogenine*, *isol*)
- Sariyar, G. et al., *Planta Med.*, 1983, **49**, 43-45 (*Oreogenine*, *isol*)

Glaucenamides **G-88**
2-(2,6,6-Trimethyl-4-oxo-2-cyclohexen-1-yl)acetamide
[330832-86-5]



- $C_{11}H_{17}NO_2$ 195.261
Alkaloid from *Fissistigma glaucescens*. Prisms (CHCl₃). $[\alpha]_D^{24}$ -120 (c, 0.15 in CHCl₃). Mp >300°. λ_{max} 237 (log ϵ 3.99) (EtOH).
- Lo, W.-L. et al., *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 1251-1256

Glaucescine **G-89**
[11022-53-0]

- $C_{19}H_{27}NO_3$ 317.427
Struct. unknown. Alkaloid from the bark of *Daphniphyllum glaucescens* (Daphniphyllaceae). Cryst. (Me₂CO). Mp 234-235° (vac. dec.). $[\alpha]_D^{20}$ +68.1 (CHCl₃).
Arthur, H.R. et al., *Phytochemistry*, 1965, **4**, 627-629 (*isol*, *ir*)


Glaucescinine **G-90**

- $C_{19}H_{27}NO_3$ 317.427
Struct. unknown. Alkaloid from the bark

of *Daphniphyllum glaucescens* (Daphniphyllaceae). Cryst. (Me₂CO). Mp 248-250° (vac. dec.).

Arthur, H.R. et al., *Phytochemistry*, 1965, **4**, 627-629 (*isol*)

Glaudelsine **G-91**
[78018-32-3]



$C_{36}H_{48}N_2O_{10}$ 668.783

The abs. config. of the methylsuccinimide residue was shown to be (S)-, as illus., for methyllycaconitine and is prob. the same for all of these alkaloids. Alkaloid from the aerial parts of *Delphinium glaucescens* (Ranunculaceae). Amorph. solid. Mp 80-110°. $[\alpha]_D^{21}$ +36.1 (c, 0.97 in CHCl₃).

Perchlorate:

Amorph. solid (MeOH/Et₂O). Mp 203-213°.

O⁶-Me: **Methyllycaconitine**. *Delartine*. *Delsemidine*. *Mellictine*
[21019-30-7]
 $C_{37}H_{50}N_2O_{10}$ 682.809

Alkaloid from *Delphinium elatum*, *Delphinium dictyocarpum*, *Delphinium tricornis*, *Delphinium grandiflorum*, *Delphinium triste*, *Delphinium crassifolium*, *Delphinium elisabethae*, *Delphinium tamarae*, *Delphinium glaucescens*, *Consolida ambigua* (*Delphinium ajacis*) and others (Ranunculaceae). Nicotinic antagonist. Amorph. Mp 128°. $[\alpha]_D^{22}$ +49.1 (c, 2 in EtOH). λ_{max} 228 (log ϵ 4.13); 276 (log ϵ 3.43) (EtOH).

▶ AR5542000

O⁶-Me, N-de-Et: **N-Deethylmethyllycaconitine**
 $C_{35}H_{46}N_2O_{10}$ 654.756
Alkaloid from *Aconitum cochleare*. $[\alpha]_D^{25}$ +35 (c, 0.25 in CHCl₃).

O⁶-Me, N-de-Et, N-formyl: **Potansine B**
[187655-62-5]
 $C_{36}H_{46}N_2O_{11}$ 682.766
Alkaloid from roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D$ +28.3 (c, 0.92 in CHCl₃).

O⁶, O⁸-Di-Me: **Potansine F**
[664304-11-4]
 $C_{38}H_{52}N_2O_{10}$ 696.836
Alkaloid from the roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D^{10}$ -23.1 (c, 1 in CHCl₃).

O⁸-Et, O¹⁴-de-Me, O⁶-Me, 14-propanoyl: **Alpinine**†
 $C_{41}H_{56}N_2O_{11}$ 752.9

Alkaloid from the leaves of *Delphinium alpinum*.

7,8-Methylene ether, O⁶-Me: **Elatine**
[26000-16-8]

$C_{38}H_{50}N_2O_{10}$ 694.82

Alkaloid from the above-ground parts of *Delphinium elatum* (Ranunculaceae). Cryst. (EtOH). Mp 233-235°. $[\alpha]_D$ +3.4 (CHCl₃).

▶ AR5554000

O¹-De-Me, O⁶-Me: **Grandiflorine**†. *Uraline*
[150044-88-5]

$C_{36}H_{48}N_2O_{10}$ 668.783

Alkaloid from *Delphinium glaucum*, *Delphinium uralense* and *Delphinium grandiflorum*. Amorph. solid. $[\alpha]_D^{23}$ +49.2 (c, 0.6 in CHCl₃).

O¹⁴-De-Me, O⁶-Me: **14-Deacetylnudicauline**. *14-Deacetylrandersoline*
[119347-24-9]

$C_{36}H_{48}N_2O_{10}$ 668.783

Alkaloid from epigeal parts of *Delphinium macrocentrum*, seeds of *Delphinium elatum* and aerial parts of *Delphinium andersonii* (Ranunculaceae). Amorph. off-white solid. $[\alpha]_D^{21}$ +40.7 (c, 0.9 in CHCl₃) (+24).

O¹⁴-De-Me, O⁶-Me, 14-Ac: **Nudicauline**. *Andersoline*
[99815-83-5]

$C_{38}H_{50}N_2O_{11}$ 710.82

Alkaloid from *Delphinium nudicaule*, seeds of *Delphinium elatum* and aerial parts of *Delphinium andersonii* (Ranunculaceae). Amorph.; cryst. (as hydroiodide). Mp 228-230° (hydroiodide). $[\alpha]_D$ +47 (c, 0.42 in CHCl₃).

O¹⁴-De-Me, O⁶-Me, O¹⁴-(2-methylpropanoyl): **14-Deacetyl-14-isobutyrylnudicauline**
[254876-71-6]

$C_{40}H_{54}N_2O_{11}$ 738.873

Alkaloid from the roots of *Delphinium stapeliosum*. Amorph. $[\alpha]_D^{20}$ +26.7 (c, 0.23 in CHCl₃).

O¹⁴-De-Me, O⁶-Me, O¹⁴-(2R-methylbutanoyl): **Elanine**
[129011-70-7]

$C_{41}H_{56}N_2O_{11}$ 752.9

Alkaloid from seeds of *Delphinium elatum* (Ranunculaceae). $[\alpha]_D$ +41.5 (c, 0.71 in CHCl₃).

O¹⁴-De-Me, O⁶-Me, 14-benzoyl: **Ajaculine**
[66731-39-3]

$C_{43}H_{52}N_2O_{11}$ 772.891

Alkaloid from the seeds of *Consolida ambigua*, (formerly *Delphinium ajacis*) (Ranunculaceae). Cryst. (CH₂Cl₂/hexane). Mp 158-161° dec. $[\alpha]_D^{24}$ +65.2 (c, 0.98 in abs. EtOH).

O¹⁶-De-Me, O⁶-Me: **Deacetylgeyerline**
[253328-44-8]

$C_{36}H_{48}N_2O_{10}$ 668.783

Alkaloid from *Delphinium nuttallianum*. Amorph. solid.

O¹⁶-De-Me, O⁶-Me, 16-Ac: **Geyerline**
[213697-78-0]

$C_{38}H_{50}N_2O_{11}$ 710.82

Alkaloid from *Delphinium geyeri*.

Amorph. solid. $[\alpha]_D^{23} +32.3$ (c, 0.3 in CHCl_3); λ_{max} 228 (log ϵ 4.05); 276 (log ϵ 3.41) (EtOH).

O¹⁴, O¹⁶-Di-de-Me, O⁶-Me, 16-Ac: Bearline
 [253337-82-5]
 $\text{C}_{37}\text{H}_{48}\text{N}_2\text{O}_{11}$ 696.793
 Alkaloid from *Delphinium nuttallianum*. Amorph. solid.

O¹⁴, O¹⁶-Di-de-Me, O⁶-Me, 14,16-di-Ac: 14-Acetylbearline
 [253328-43-7]
 $\text{C}_{39}\text{H}_{50}\text{N}_2\text{O}_{12}$ 738.83
 Alkaloid from *Delphinium nuttallianum*. Amorph. solid.

14-Ketone, O¹⁴-de-Me, O⁶-Me: Barbinine
 [123497-99-4]
 $\text{C}_6\text{H}_4\text{N}_2\text{O}_{10}$ 666.767
 Alkaloid from aerial parts of *Delphinium barbeyi* (Ranunculaceae). Amorph. $[\alpha]_D^{26} +35.9$ (c, 0.388 in CHCl_3).

14-Ketone, O¹, O¹⁴-di-de-Me, O⁶-Me: Grandifloricine
 [146471-87-6]
 $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_{10}$ 652.74
 Alkaloid from aerial parts of *Delphinium grandiflorum* (Ranunculaceae).

19-Oxo, O⁶-Me, N-de-Et: Zailine
 [145237-02-1]
 $\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_{11}$ 668.739
 Alkaloid from seeds of *Delphinium zaili* (Ranunculaceae).

10-Hydroxy, O⁶-Me: 10-Hydroxymethyllycaconitine
 [532993-22-9]
 $\text{C}_{37}\text{H}_{50}\text{N}_2\text{O}_{11}$ 698.809
 Alkaloid from *Delphinium excelsum*. Amorph. $[\alpha]_D^{20} +51$ (CHCl_3).

15 β -Hydroxy, O⁶, O⁸-di-Me, N-de-Et, N-formyl: Potanisine E
 [182262-36-8]
 $\text{C}_{37}\text{H}_{48}\text{N}_2\text{O}_{12}$ 712.792
 From roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D +13$ (c, 0.10 in CHCl_3).


16-Demethoxy, O⁶-Me: 16-Demethoxymethyllycaconitine
 $\text{C}_{36}\text{H}_{48}\text{N}_2\text{O}_9$ 652.783
 Alkaloid from the roots of *Delphinium cuneatum*.

Goodson, J.A. *et al.*, *J.C.S.*, 1943, 139-141 (*Methyllycaconitine*)
 Kuzovkov, A.D. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 2746-2749 (*Methyllycaconitine, Elatine*)
 Pelletier, S.W. *et al.*, *J. Nat. Prod.*, 1980, **43**, 395-406 (*Ajacusine*)
 Pelletier, S.W. *et al.*, *J.A.C.S.*, 1981, **103**, 6536-6538 (*config*)
 Pelletier, S.W. *et al.*, *J.O.C.*, 1981, **46**, 3284 (*isol, ir, pmr, cmr, ms, struct*)
 Edwards, O.E. *et al.*, *Can. J. Chem.*, 1982, **60**, 2661-2667 (*config*)
 Kulanthaivel, P. *et al.*, *Heterocycles*, 1985, **23**, 2515-2520 (*Nudicauline*)
 Pelletier, S.W. *et al.*, *Heterocycles*, 1987, **26**, 2835-2840 (*Elatine, synth, pmr, cmr*)
 Pelletier, S.W. *et al.*, *Heterocycles*, 1988, **27**, 2387-2393 (*14-Deacetyludicauline*)
 Benn, M.H. *et al.*, *Phytochemistry*, 1989, **28**, 919-922 (*Deacetyludicauline*)
 Pelletier, S.W. *et al.*, *Phytochemistry*, 1989, **28**, 1521-1523; 1990, **29**, 2381-2383 (*Barbinine, Elatine*)

Pelletier, S.W. *et al.*, *Tetrahedron*, 1989, **45**, 1887-1892 (*Nudicauline, Deacetyludicauline*)
 Li, C. *et al.*, *CA*, 1992, **118**, 143418h (*Grandifloricine*)
 Sun, F. *et al.*, *Phytochemistry*, 1992, **31**, 3247-3250 (*Zailine*)
 Coates, P.A. *et al.*, *Tet. Lett.*, 1994, **35**, 8701-8704 (*abs config, Methyllycaconitine*)
 Pu, H.-Y. *et al.*, *Phytochemistry*, 1996, **43**, 287-290 (*Potanisine E*)
 Pu, H. *et al.*, *Planta Med.*, 1996, **62**, 462-464 (*Potanisine B*)
 Manners, G.D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1086-1089 (*Geyerline, Grandifloricine*)
 Gardner, D.R. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 5049-5058 (*Bearline, 14-Acetylbearline*)
 Shrestha, P.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 2-5; 2004, **67**, 1574-1576 (*Deacetylisobutyryludicauline, Deacetyludicauline, Methyllycaconitine, Geyerline*)
 Gardner, D.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1127-1130 (*Bearline, Acetylbearline, Deacetylgeyerline, isol, pmr, cmr, ms*)
 Shen, X.-L. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1265-1267 (*Elatine, cmr*)
 Chen, D.-L. *et al.*, *J. Asian Nat. Prod. Res.*, 2003, **5**, 209-213 (*Potanisine F*)
 Batbayar, N. *et al.*, *Phytochemistry*, 2003, **62**, 543-550 (*10-Hydroxymethyllycaconitine*)
 Khairitdinova, E.D. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2003, **52**, 2078-2080 (*16-Demethoxymethyllycaconitine*)
 Khairitdinova, E.D. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 575-577 (*Alpinine*)
 Tsyrlina, T.M. *et al.*, *Russ. J. Bioorg. Chem. (Engl. Transl.)*, 2005, **31**, 383-386 (*Uraline*)
 Meriçli, A.H. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 210-217 (*N-Deethylmethyllycaconitine, Methyllycaconitine*)

Glauflavine G-92
 [11024-37-6 (iodide)]
 Struct. unknown. Quaternary alkaloid from *Glaucium flavum* (Papaveraceae). Mp 216-217° (as iodide). $[\alpha]_D +45$ (c, 1.0 in CHCl_3) (iodide).
 Gertik, H. *et al.*, *Diss. Pharm. Pharmacol.*, 1966, **18**, 375-380; *CA*, **66**, 17042

Glaunine G-93
11-Hydroxy-1,2,10-trimethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI
 [57986-73-9]



$\text{C}_{19}\text{H}_{15}\text{NO}_5$ 337.331
 Alkaloid from *Glaucium fimbriigerum* (Papaveraceae).

O¹-De-Me, N-Me, inner salt: 1,11-Dihydroxy-2,10-dimethoxy-6-methyl-7-oxo-7H-dibenzo[de,g]quinolinium. Arosinine
 [73777-68-1]

$\text{C}_{19}\text{H}_{15}\text{NO}_5$ 337.331
 Alkaloid from the roots of *Glaucium flavum* var. *vestitum* (Papaveraceae). Dark-green needles ($\text{CHCl}_3/\text{EtOH}$). Mp 302-305° dec.


O¹-De-Me, O¹¹-Me, N-Me, inner salt: Glaunidine. Arosine
 [72032-70-3]
 $\text{C}_{20}\text{H}_{17}\text{NO}_5$ 351.358
 Alkaloid from *Glaucium flavum* var. *vestitum*, *Glaucium fimbriigerum* and *Aconitum leucostomum* (Papaveraceae, Ranunculaceae). Dark-green prisms ($\text{EtOAc}/\text{CHCl}_3$). Mp 245-248° dec.

O¹⁰-De-Me, O¹¹-Me: 7-Oxohermagine
 [172274-06-5]
 $\text{C}_{19}\text{H}_{15}\text{NO}_5$ 337.331
 Alkaloid from the stem bark of *Hernandia nymphaeifolia*. Shows anti-platelet aggregation activity. Orange prisms (MeOH). Mp 252-254°. λ_{max} 213 (log ϵ 4.52); 274 (log ϵ 4.42); 360 (log ϵ 3.96); 403 (log ϵ 3.92) (EtOH).

Israïlov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 415; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 367 (*Glaunine, Glaunidine, isol, uv, ir, pmr, struct*)
 Castedo, L. *et al.*, *Tet. Lett.*, 1979, 4589 (*Glaunidine, Arosinine*)
 Karimova, S.U. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 224; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 177 (*isol*)
 Zhamierashvili, M.G. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 805; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 576 (*Glaunidine, occur*)
 Chen, J.-J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 156-158 (*7-Oxohermagine*)
 Chen, J.J. *et al.*, *Planta Med.*, 2000, **66**, 251-256 (*activity*)

Glenodinine G-95
 Struct. unknown. Alkaloid isol. from the dinoflagellate *Peridinium polonicum*. Ichthyotoxin. λ_{max} 200 (MeOH).
 Hashimoto, Y. *et al.*, *Nippon Suisan Gakkaishi*, 1968, **34**, 528-534; *CA*, **72**, 76239s

Glidobactamine G-96



Absolute Configuration

$\text{C}_{15}\text{H}_{26}\text{N}_4\text{O}_5$ 342.394
 Parent peptide moiety of the Glidobactins prod. by enzymic deacylation of Glidobactins in a *Pseudomonas* strain. Amorph. solid. $[\alpha]_D^{27} -157$ (c, 0.5 in H_2O).

N^{Thr}-(2E,4E-Decadienoyl): Glidobactin F
 [119259-72-2]
 $\text{C}_{25}\text{H}_{40}\text{N}_4\text{O}_6$ 492.614
 Isol. from *Polyangium brachysporum*

K481-B101 ATCC 53080. Antitumour agent. Powder. Sol. MeOH, DMSO, butanol; fairly sol. CHCl₃, MeCN, EtOAc; poorly sol. H₂O, hexane. Mp 233° dec. λ_{\max} 260 (ε 24200) (95% EtOH) (Derep).

N^{Thr} -(9-Methyl-2E,4E-dodecadienyl): **Cepafungin III**
[130743-09-8]

C₂₆H₄₂N₄O₆ 506.641
Prod. by a *Pseudomonas* sp. Exhibits activities against fungi, yeast and tumours. Powder. Sol. MeOH-H₂O, CHCl₃-MeOH, DMSO; fairly sol. MeOH; poorly sol. EtOAc, hexane. Mp 215-220°. $[\alpha]_D^{25}$ -110 (c, 0.155 in MeOH). λ_{\max} 260 (ε 24200) (95% EtOH).

N^{Thr} -(2E,4E-Dodecadienyl): **Gliodobactin A. Cepafungin II**
[108351-50-4]

C₂₇H₄₄N₄O₆ 520.668
Prod. by a *Pseudomonas* sp. and *Polyangium brachysporum*. Antitumour agent also active against fungi and yeasts. Needles + ½ H₂O (MeOH aq.). Mp 245-250° dec. (259-261°). $[\alpha]_D^{24}$ -127 (c, 0.283 in MeOH). $[\alpha]_D^{24}$ -111 (c, 0.5 in MeOH). λ_{\max} 258 (ε 35500) (MeOH) (Derep).

► JR1673000

N^{Thr} -(7ξ-Hydroxy-2E,4E-dodecadienyl): **Gliodobactin E**
[119259-74-4]

C₂₇H₄₄N₄O₇ 536.667
Isol. from *Polyangium brachysporum* K481-B101. Antitumour agent. Powder. Sol. MeOH, butanol, DMSO; fairly sol. CHCl₃, MeCN, EtOAc; poorly sol. H₂O, hexane. Mp 195° dec. λ_{\max} 258 (ε 35500) (MeOH).

N^{Thr} -(10ξ-Hydroxy-2E,4E-dodecadienyl): **Gliodobactin D**
[119259-73-3]

C₂₇H₄₄N₄O₇ 536.667
Isol. from *Polyangium brachysporum* K481-B101. Antitumour agent. Powder. Sol. MeOH, DMSO, butanol; fairly sol. CHCl₃, MeCN, EtOAc; poorly sol. H₂O, hexane. Mp 204° dec. λ_{\max} 258 (ε 35500) (MeOH).

N^{Thr} -(11-Methyl-2E,4E-dodecadienyl): **Cepafungin I**
[130743-08-7]

C₂₈H₄₆N₄O₆ 534.695
Prod. by a *Pseudomonas* sp. Exhibits activities against fungi, yeast and tumours. Powder + 1H₂O. Sol. MeOH-H₂O, DMSO, CHCl₃-MeOH; fairly sol. MeOH; poorly sol. EtOAc, hexane. Mp 235-240° dec. $[\alpha]_D^{24}$ -124 (c, 0.694 in MeOH). λ_{\max} 260 (ε 32700) (EtOH).

N^{Thr} -(2E,4E-Tetradecadienyl): **Gliodobactin C**
[108351-52-6]

C₂₉H₄₈N₄O₆ 548.721
Isol. from *Polyangium brachysporum* K481-B101. Antitumour agent also active against fungi and yeasts. Needles (MeOH). Sol. MeOH, DMSO, EtOH; fairly sol. EtOAc, MeCN, CHCl₃; poorly sol. H₂O, hexane. Mp 273-275°.

$[\alpha]_D^{24}$ -104 (c, 0.5 in MeOH). λ_{\max} 258 (ε 35500) (MeOH).

► LD₅₀ (mus, ipr) 25 mg/kg. XB7450000

N^{Thr} -(2E,4E,8Z-Tetradecatrienyl): **Gliodobactin B**

[108351-51-5]

C₂₉H₄₆N₄O₆ 546.706

Prod. by *Polyangium brachysporum* K481-B101. Antitumour agent also active against fungi and yeasts. Powder + 1½ H₂O. Sol. MeOH, DMSO, EtOH; fairly sol. EtOAc, MeCN, CHCl₃; poorly sol. H₂O, hexane. Mp 232-234°. $[\alpha]_D^{24}$ -92 (c, 0.5 in MeOH). λ_{\max} 258 (ε 35500) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 8.1 mg/kg.

N^{Thr} -(2E,4E,8Z,11Z-Tetradecatetraenyl): **Gliodobactin PFI**

[138087-92-0]

C₂₉H₄₄N₄O₆ 544.69

Prod. by *Polyangium brachysporum* K481-B101. Cytotoxic. Amorph. powder + 1H₂O. Sol. MeOH, butanol, DMSO; poorly sol. hexane, H₂O. Mp 194-196° (dec.). $[\alpha]_D^{24}$ -114 (c, 0.25 in MeOH). λ_{\max} 261 (ε 34000) (MeOH). λ_{\max} 261 (ε 34000) (MeOH) (Berdy).

N^{Thr} -Acyl (?): **Gliodobactin H**

[119314-46-4]

Prod. by *Pseudomonas* sp. Mp 191° dec. Congener of undetermined struct., isol. only in v. low yield.

13-Hydroxy, N^{Thr} -(2E,4E-dodecadienyl): **Gliodobactin G**

[119259-71-1]

C₂₇H₄₄N₄O₇ 536.667

Isol. from *Polyangium brachysporum* K481-B101. Antitumour agent. Powder. Mp 217° dec. λ_{\max} 258 (ε 35500) (MeOH) (Derep).

Oka, M. *et al.*, *J. Antibiot.*, 1988, **41**, 1331-1337; 1338-1350; 1351-1357; 1358-1365; 1812-1822; 1906-1909 (*Cepafungins, Gliodobactins, isol, struct, synth*)

Numata, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1351-1357 (*Gliodobactamine*)

U.S. Pat., 1988, 4777160 (*Gliodobactins*)

Shoji, J. *et al.*, *J. Antibiot.*, 1990, **43**, 783-787; 788-795 (*Cepafungins*)

Titus, J.A. *et al.*, *J. Ind. Microbiol. Biotechnol.*, 1990, **6**, 215-218 (*manuf*)

Eur. Pat., 1991, 437 244; CA, **116**, 19730y


(*Gliodobactin PFI*)

Schmidt, U. *et al.*, *Chem. Comm.*, 1992, 1687-1689 (*synth*)

Gliocladine A†

Bionectin C

[749216-45-3]



C₂₄H₂₄N₄O₃S₂ 480.611

Not to be confused with Gliocladine A in V-89. Identity with Bionectin C not

G-97

Relative Configuration

confirmed. Prod. by *Gliocladium roseum* OUPS-N132 isol. from *Aplysia kurodai*. Also prod. by *Bionectra byssicola* F120. Cryst. (CH₂Cl₂/MeOH). Mp 174-176°. $[\alpha]_D^{16}$ +263 (c, 0.14 in CHCl₃). λ_{\max} 242 (log ε 4.36); 283 (log ε 4.12); 291 (log ε 4.11) (EtOH).

11-Deoxy: **Gliocladine B†**

[749216-46-4]

C₂₄H₂₄N₄O₂S₂ 464.611


Prod. by *Gliocladium roseum* OUPS-N132 from *Aplysia kurodai*. Cryst. (CH₂Cl₂/MeOH). Mp 145-148°. $[\alpha]_D^{16}$ +200 (c, 0.06 in CHCl₃). λ_{\max} 283 (log ε 3.78); 292 (log ε 3.77) (EtOH).

Usami, Y. *et al.*, *Heterocycles*, 2004, **63**, 1123-1129 (*Gliocladines A,B, isol, cd, pmr, cmr*)

Zheng, C.J. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1816-1819 (*Bionectin C, isol, pmr, cmr*)

Gliocladine C†

[749216-47-5]



C₂₂H₁₆N₄O₃ 384.393


Not to be confused with Gliocladine C, G-99. Prod. by *Gliocladium roseum* OUPS-N132 isol. from *Aplysia kurodai*. Cytotoxic. Pale yellow powder (CH₂Cl₂/MeOH). Mp 180-183°. $[\alpha]_D^{16}$ +131.4 (c, 0.07 in CHCl₃). λ_{\max} 242 (log ε 5.11); 283 (log ε 4.88); 292 (log ε 4.89) (EtOH).

Usami, Y. *et al.*, *Heterocycles*, 2004, **63**, 1123-1129 (*isol, cd, pmr, cmr*)

Overman, L.E. *et al.*, *Org. Lett.*, 2007, **9**, 339-341 (*synth*)

Gliocladine C†

[871335-07-8]



C₂₃H₂₀N₄O₃S₂ 464.568

Not to be confused with Gliocladine C, G-98. Prod. by *Gliocladium roseum* 1A. Amorph. powder. $[\alpha]_D^{19}$ +512.6 (c, 0.33 in Py). λ_{\max} 204 (log ε 0.94); 219 (log ε 0.93); 283 (log ε 0.16); 290 (log ε 0.15) (Py).

G-98

G-99

13-Hydroxy: Antibiotic T 988C. T 988C
 $C_{23}H_{20}N_4O_4S_2$ 480.567
 Prod. by *Tilachlidium* sp. (CANU-T988). Cytotoxic. Amorph. solid. $[\alpha]_D^{20} +277$ (c, 0.0006 in MeOH). λ_{max} 210 (log ϵ 4.21); 234 (log ϵ 4.29); 282 (log ϵ 4.12); 354 (log ϵ 3.63) (MeOH).

Deoxy: Gloclatine
 [929616-13-7]
 $C_{23}H_{20}N_4O_2S_2$ 448.569
 Prod. by *Gliocladium roseum*
 YMF1.00133.

Trisulfide analogue: Glocladine D
 [871335-08-9]
 $C_{23}H_{20}N_4O_3S_3$ 496.634
 Prod. by *Gliocladium roseum* 1A. Amorph. powder. $[\alpha]_D^{25} +604.1$ (c, 0.47 in Py). λ_{max} 204 (log ϵ 0.54); 219 (log ϵ 0.64); 290 (log ϵ 0.11) (Py).

Trisulfide analogue, 13-hydroxy: Antibiotic T 988A. T 988A
 $C_{23}H_{20}N_4O_4S_3$ 512.633
 Prod. by *Tilachlidium* sp. (CANU-T988). Cytotoxic. Amorph. solid. $[\alpha]_D^{20} +366$ (c, 0.002 in MeOH). λ_{max} 210 (log ϵ 4.07); 230 (log ϵ 4.16); 258 (log ϵ 4.19); 292 (log ϵ 4.16) (MeOH).

Tetrasulfide analogue: Glocladine E
 [870480-90-3]
 $C_{23}H_{20}N_4O_3S_4$ 528.7
 Prod. by *Gliocladium roseum* 1A. Amorph. powder. $[\alpha]_D^{19} +562.6$ (c, 0.43 in Py). λ_{max} 206 (log ϵ 0.64); 219 (log ϵ 0.74); 290 (log ϵ 0.17) (Py).


3-Desmethyl: Bionectin A
 $C_{22}H_{18}N_4O_3S_2$ 450.541
 Prod. by *Bionectra byssicola* F120. Antibacterial agent. Active against MRSA. Powder. $[\alpha]_D +584$ (c, 0.2 in MeOH). λ_{max} 220 (log ϵ 4.54); 245 (sh) (log ϵ 4.06); 281 (log ϵ 3.81); 290 (log ϵ 3.79) (MeOH).

3-Desmethyl, 3-(1-hydroxyethyl): Bionectin B
 $C_{24}H_{22}N_4O_4S_2$ 494.594
 Prod. by *Bionectra byssicola* F120. Antibacterial agent. Active against MRSA. Powder. $[\alpha]_D +493$ (c, 0.2 in MeOH). λ_{max} 219 (log ϵ 4.59); 246 (sh) (log ϵ 3.99); 282 (log ϵ 3.81); 291 (log ϵ 3.79) (MeOH).

Feng, Y. et al., *J. Nat. Prod.*, 2004, **67**, 2090-2092 (T 988A, T 988C)
 Dong, J.-Y. et al., *J. Nat. Prod.*, 2005, **68**, 1510-1513 (Glocladines C,D,E)
 Dong, J.-Y. et al., *Chin. Chem. Lett.*, 2006, **17**, 922-924 (Gloclatine)
 Zheng, C.-J. et al., *J. Nat. Prod.*, 2006, **69**, 1816-1819 (Bionectins A,B)

Glochidicine

G-100


 $C_{15}H_{23}N_3O$ 261.366


(±)-form [1386-87-4]

Alkaloid from the leaves of a *Glochidion* sp. (probably *Glochidion philippicum*)

(Euphorbiaceae). Cryst. + $\frac{1}{2}$ H₂O (Me₂CO). Mp 103-105°.
 Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 555 (isol, ir, ms, pmr, struct)
 Lee, Y.S. et al., *Heterocycles*, 1994, **37**, 303 (synth, cmr)

Glochidine

G-101


 $C_{15}H_{23}N_3O$ 261.366

(±)-form [1386-83-0]


Alkaloid from the leaves of a *Glochidion* sp. (probably *Glochidion philippicum*) (Euphorbiaceae). Mp 65-67°.

Picrate:

Yellow prisms (EtOH). Mp 143-144°.
 Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 555 (isol, ir, ms, pmr, struct)
 Lee, Y.S. et al., *Heterocycles*, 1994, **37**, 303 (synth, cmr)

Glucosatisin

G-102


 $C_{21}H_{26}N_2O_{12}S_2$ 562.575

Isol. from woad (*Isatis tinctoria*). Isol. as a mixture with C-3 epimer. λ_{max} 206 (log ϵ 4.29); 228 (log ϵ 3.9); 273 (log ϵ 3.23) (MeOH).

3-Epimer: Epiglucosatisin

$C_{21}H_{26}N_2O_{12}S_2$ 562.575
 Isol. from *Isatis tinctoria*. Isol. as a mixture with C-3 epimer.

3'-ξ-Hydroxy: 3-Hydroxyglucosatisin

$C_{21}H_{26}N_2O_{13}S_2$ 578.574
 Isol. from *Isatis tinctoria*. Isol. as a mixture with C-3 epimer. λ_{max} 210 (log ϵ 3.85); 235 (log ϵ 3.38); 247 (log ϵ 3.17); 289 (log ϵ 2.7) (MeOH).


3-Epimer, 3'-ξ-hydroxy: 3'-Hydroxyepiglucosatisin

$C_{21}H_{26}N_2O_{13}S_2$ 578.574
 Isol. from *Isatis tinctoria*. Isol. as a mixture with C-3 epimer.
 Frechard, A. et al., *Tet. Lett.*, 2001, **42**, 9015-9017

N-[3-(β-D-Glucopyranosyloxy)-2,3-dihydro-2-oxo-1H-indol-3-yl]acetyl]aspartic acid, 9CI

G-103

[99694-85-6]




$C_{20}H_{24}N_2O_{12}$ 484.416
 Constit. of the seedlings of *Vicia faba*.
 Metab. of indolacetic acid. Powder.

Tsurumi, S. et al., *Plant Physiol.*, 1985, **79**, 667 (isol)

Glucose 3-nitropropanoates

G-104



A group of toxic compds. which are esters of 1 mol. of glucose with 1, 2, 3 or 4 mols. of 3-Nitropropanoic acid. The substitution pattern for the Endecaphyllins is unknown and they may be identical with the other compounds of known struct. listed.

6-O-(3-Nitropropanoyl)-α-D-glucopyranose

[59440-99-2]

$C_9H_{15}NO_9$ 281.219
 Constit. of the aerial shoots of *Coronilla varia*.

6-O-(3-Nitropropanoyl)-β-D-glucopyranose

[59441-00-8]

$C_9H_{15}NO_9$ 281.219
 Constit. of the aerial shoots of *Coronilla varia*.

1,6-Bis-O-(3-nitropropanoyl)-β-D-glucopyranose

Cibarian
 [39797-90-5]

$C_{12}H_{18}N_2O_{12}$ 382.28
 Isol. from *Lotus pedunculatus*, *Astragalus* spp. and *Coronilla varia*. Mp 123.5-124°.

2,3-Bis-O-(3-nitropropanoyl)-α-D-glucopyranose

6-O-(2-Propenoyl): 6-O-Acryloyl-2,3-bis-O-(3-nitropropanoyl)-α-D-glucopyranose
 [913739-14-7]

$C_{15}H_{20}N_2O_{13}$ 436.329
 Constit. of the roots of *Indigofera carlesii*.

2,6-Bis-O-(3-nitropropanoyl)-α-D-glucopyranose

Coronarian
 [63505-68-0]

$C_{12}H_{18}N_2O_{12}$ 382.28
 Isol. from *Lotus pedunculatus*, *Corynocarpus laevigatus*, *Coronilla* and *Indigofera* spp. Cryst. (Me₂CO/CCl₄). Mp 151-153°.

3,6-Bis-O-(3-nitropropanoyl)-α-D-glucopyranose

Kirilowin G

Constit. of the roots of *Indigofera kirilowii*.

2-O-(2-Propenoyl): Kirilowin B. 2-O-Acryloyl-3,6-bis-O-(3-nitropropanoyl)-α-D-glucopyranose
 [874363-53-8]

$C_{15}H_{20}N_2O_{13}$ 436.329

Constit. of the roots of *Indigofera carlesii* and *Indigofera kirilowii*. Needles (Me₂CO). Mp 221-222°. [α]_D +60.4 (c, 0.53 in Me₂CO).

Bis-O-(3-nitropropanoyl)-D-glucopyranose (5)

Endecaphyllin C₁

[90114-46-8]

$C_{12}H_{18}N_2O_{12}$ 382.28

Alkaloid from *Indigofera endecaphylla*. Mp 145-146.5°.

Bis-O-(3-nitropropanoyl)-D-glucopyranose (6)

Endecaphyllin D

[90114-46-8]

$C_{12}H_{18}N_2O_{12}$ 382.28

Alkaloid from *Indigofera endecaphylla*. Mp 145-146°.

Bis-O-(3-nitropropanoyl)-D-glucopyranose (7)

Endecaphyllin E

[90114-46-8]

$C_{12}H_{18}N_2O_{12}$ 382.28

Alkaloid from *Indigofera endecaphylla*. Mp 132-134°.

1,2,6-Tris-O-(3-nitropropanoyl)- α -D-glucopyranose

Coronillin

[63368-43-4]

$C_{15}H_{21}N_3O_{15}$ 483.342

Isol. from aerial parts of *Coronilla varia*. Cryst. (Me₂CO/CCl₄). Mp 112.5-114°.

1,2,6-Tris-O-(3-nitropropanoyl)- β -D-glucopyranose

Karakin. Endecaphyllin A

[1400-11-9]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Corynocarpus laevigata*, *Astragalus* spp., *Lotus pedunculatus* and others. Mp 124-125° (122°). [α]_D²⁰ +4.5. Struct. revised in 1975. Identity of Karakin and Endecaphyllin A not certain. Gnanasunderam *et al.* give the 1,4,6-triacyl struct. which is apparently in error (archaic struct. superseded by Harlow's reassignment).

► Toxic.

1,3,6-Tris-O-(3-nitropropanoyl)- β -D-glucopyranose [71454-29-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Astragalus canadensis* and *Corynocarpus laevigatus*.

1,4,6-Tris-O-(3-nitropropanoyl)- β -D-glucopyranose

Corynocarpin

[1391-18-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Astragalus canadensis* and *Corynocarpus laevigatus*. Cryst. (EtOAc/CCl₄). Mp 114.5-115°.

2,3,6-Tris-O-(3-nitropropanoyl)- α -D-glucopyranose

Corollin

[63461-31-4]

$C_{15}H_{21}N_3O_{15}$ 483.342

Constit. of crown vetch *Coronilla varia*. Mp 158.5-160°.

3,4,6-Tris-O-(3-nitropropanoyl)- α -D-glucopyranose [144549-44-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Corynocarpus laevigatus* and *Indigofera linnaei*.

2-O-(2-Propenoyl): Kirilowin A. 2-O-Acryloyl-3,4,6-tris-O-(3-nitropropanoyl)- α -D-glucopyranose

[874363-52-7]

$C_{18}H_{23}N_3O_{16}$ 537.39

Constit. of the roots of *Indigofera kirilowii*. Needles (Me₂CO). Mp 162-163°. [α]_D²⁰ +71.2 (c, 0.17 in Me₂CO).

Tris-O-(3-nitropropanoyl)-D-glucose (7)

Endecaphyllin A₁

[97159-95-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Indigofera endecaphylla*. Mp 81-91°.

Tris-O-(3-nitropropanoyl)-D-glucose (8)

Endecaphyllin A₂

[97159-95-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Indigofera endecaphylla*. Mp 156-158.5°.

Tris-O-(3-nitropropanoyl)-D-glucose (9)

Endecaphyllin B

[97159-95-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Indigofera endecaphylla*. Mp 125-126.5°.

Tris-O-(3-nitropropanoyl)-D-glucose (10)

Endecaphyllin B₁

[97159-95-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Indigofera endecaphylla*. Mp 129-130°.

Tris-O-(3-nitropropanoyl)-D-glucose (11)

Endecaphyllin C

[97159-95-0]

$C_{15}H_{21}N_3O_{15}$ 483.342

Alkaloid from *Indigofera endecaphylla*. Mp 155-156°.

1,2,3,6-Tetrakis-O-(3-nitropropanoyl)- β -D-glucopyranose [155020-57-8]

$C_{18}H_{24}N_4O_{18}$ 584.404

Constit. of *Corynocarpus laevigatus*.

1,2,4,6-Tetrakis-O-(3-nitropropanoyl)- β -D-glucopyranose

Hiptagin. Endecaphyllin X

[19896-10-7]

$C_{18}H_{24}N_4O_{18}$ 584.404

Isol. from *Hiptage madablota*, *Lotus pedunculatus*, *Astragalus flexuosus* and *Indigofera endecaphylla*. Needles (EtOH aq.). Mp 110° (104-104.5°). [α]_D +3.5 (Me₂CO).

1,3,4,6-Tetrakis-O-(3-nitropropanoyl)- β -D-glucopyranose [155020-58-9]

$C_{18}H_{24}N_4O_{18}$ 584.404

Constit. of *Corynocarpus laevigatus*.

2,3,4,6-Tetrakis-O-(3-nitropropanoyl)- α -D-

glucopyranose [122475-42-7]

$C_{18}H_{24}N_4O_{18}$ 584.404

Constit. of *Indigofera suffruticosa* and *Indigofera linnaei*. Cryst. (MeOH). Mp 139-141°.

Skey, W. *et al.*, *Ber.*, 1873, 6, 627 (*isol.*, Karakin) Finnegan, R.A. *et al.*, *J. Pharm. Sci.*, 1965, 54, 1136-1144; 1968, 57, 353-354

(*Endecaphyllins*, Hiptagin)

Sternitz, F.R. *et al.*, *Phytochemistry*, 1972, 11, 3525-3527 (*Coronarian*)

Harlow, M.C. *et al.*, *Phytochemistry*, 1975, 14, 1421-1423 (*Hiptagin*, *Cibarian*, *Karakin*)

Williams, M.C. *et al.*, *Phytochemistry*, 1975, 14, 2306-2308 (*Hiptagin*, *Cibarian*, *Karakin*)

Moyer, B.G. *et al.*, *Phytochemistry*, 1977, 16, 375-377; 1979, 18, 111-113 (*Coronarian*, *Corollin*, *Corynocarpin*, *Cibarian*, *Coronillin*)

Pfeffer, P.E. *et al.*, *Carbohydr. Res.*, 1979, 73, 1-8 (*cmr*)

Gnanasunderam, C. *et al.*, *Phytochemistry*, 1986, 25, 409-410 (*Hiptagin*, *Coronarian*, *Cibarian*, *Karakin*)

Garcez, W.S. *et al.*, *Phytochemistry*, 1989, 28, 1251-1252 (*2,3,4,6-isomer*)

Majak, W. *et al.*, *Phytochemistry*, 1994, 35, 901-903 (*isol.*)


Su, Y.-F. *et al.*, *J. Nat. Prod.*, 2005, 68, 1785-1786 (*Kirilowin A, B*)

Zhang, X.-X. *et al.*, *Fitoterapia*, 2006, 77, 15-18 (*Acryloylnitropropanoylglucoses*)

Su, Y.-F. *et al.*, *Fitoterapia*, 2008, 79, 451-455 (*Kirilowin C-H*)

Glusun I

G-105




$C_{15}H_{11}N_3O_4S$ 329.336

Prod. by the marine-derived *Halomonas* sp. RK377. Yellow solid. Mp 150° dec. λ_{\max} 271 (log ϵ 3.31); 278 (log ϵ 3.32); 356 (log ϵ 3.35) (MeOH).

Liang, L. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*isol.*, *uv*, *pmr*, *cmr*, *ms*)

Glusun II

G-106




$C_{16}H_{13}N_3O_5S$ 359.362

Prod. by the marine-derived *Halomonas* sp. RK377. Orange solid. λ_{\max} 272 (log ϵ 3.72); 365 (log ϵ 3.56) (MeOH).

Liang, L. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*isol.*, *uv*, *pmr*, *cmr*, *ms*)

Glutamic acid **G-107**

2-Aminopentanedioic acid, 9CI. 2-Aminoglutamic acid. Glutaminic acid. Glu. E620. FEMA 3285 [6899-05-4]



C₅H₉NO₄ 147.13

(R)-form

D-form

[6893-26-1]

Component of bacterial cell proteins and peptidolipids. Leaflets (H₂O). Mp 213° dec. (rapid heating). [α]_D²⁶ -12.9 (H₂O). [α]_D +31 (aq. HCl). Tasteless.

Hydrochloride: [617-61-8]

Prisms (HCl aq.). Mp 209° dec. [α]_D²⁰ -24.5 (H₂O).

N-(4-Aminobutyl): *N*-(4-Aminobutyl)-glutamic acid. **Rideopine**

[279688-93-6]

C₉H₁₈N₂O₄ 218.252

Prod. by crown gall tumours.

N-Me: *N*-Methyl-*D*-glutamic acid

[77481-28-8]

C₆H₁₁NO₄ 161.157

Isol. from the mollusc *Scapharca broughtonii*.

(S)-form

L-form

[56-86-0]

Since 1965 the main industrial source has been bacterial fermentation of carbohydrate sources such as molasses and corn starch hydrolysate in the presence of a nitrogen source such as ammonium salts or urea. Annual production approx. 350000t worldwide in 1988. Obt. from acid hydrolyd. of proteins. Seasoning additive in food manuf. (as Na, K and NH₄ salts). Intermed. in 2,5-Diaminopentanoic acid biosynth. Excitatory CNS transmitter. Dietary supplement, nutrient. Rhombic cryst. (EtOH aq.). Spar. sol. EtOH, H₂O (0.84 g/100g at 25°). Mp 224-225° (211-213°, 247-249°) dec. [α]_D²⁵ +17 (c, 2 in H₂O). [α]_D²⁵ +46.8 (c, 2 in 5M HCl). pK_{a1} 2.2; pK_{a2} 4.2; pK_{a3} 9.6 (25°, NH₂). Isoelectric point 3.24. Appetising taste.

► Human systemic effects by ingestion and intravenous routes. LZ9700000

Hydrochloride: [138-15-8] Dietary supplement, nutrient, flavouring agent and enhancer. Used in treatment of achlorhydria or hypochlorhydria. Mp 202° dec Mp 213° (rapid heating).

N-(9Z,12Z-Octadecadienyl): *N*-(9,12-Octadecadienyl)glutamic acid. *N*-Linoleoyl-*L*-glutamic acid

[2601-92-5]

[103188-97-2]

C₂₃H₃₉NO₅ 409.565

Constit. of the oral secretions of the caterpillar *Spodoptera littoralis*. Mp 76-79°.

N-(9Z,12Z,15Z-Octadecatrienyl): *N*-

(9,12,15-Octadecatrienyl)glutamic acid. *N*-Linolenoylglutamic acid

[247150-66-9]

C₂₃H₃₇NO₅ 407.549

Constit. of the oral secretions of the caterpillar *Epirrita autumnata*.

N-(4-Aminobutanoyl): *N*-(4-Aminobutanoyl)glutamic acid

[122910-12-7]

C₉H₁₆N₂O₅ 232.236

Isol. from the toxic mushroom *Clitocybe acromelalga*. Sol. H₂O. [α]_D²⁶ +20.8 (c, 0.16 in H₂O) (synthetic).

► Convulsive poison.

N-(4-Hydroxybenzoyl): *N*-(4-Hydroxybenzoyl)glutamic acid

[119843-02-6]

C₁₂H₁₃NO₆ 267.238

Constit. of *Anthoceros agrestis*.

N-(3,4-Dihydroxybenzoyl): *N*-(3,4-Dihydroxybenzoyl)glutamic acid

[159623-11-7]

C₁₂H₁₃NO₇ 283.237

Constit. of *Anthoceros agrestis*.

N-(4-Hydroxy-Z-cinnamoyl): *N*-cis-(p-Coumaroyl)glutamic acid

[159623-15-1]

C₁₄H₁₅NO₆ 293.276

Constit. of *Anthoceros agrestis*.

N-(3-Hydroxy-4-methoxycinnamoyl) (*E*): *N*-trans-Isoferuloylglutamic acid

[159623-13-9]

C₁₅H₁₇NO₇ 323.302

Constit. of *Anthoceros agrestis*.

N-(3-Hydroxy-4-methoxycinnamoyl) (*Z*): *N*-cis-Isoferuloylglutamic acid

[159623-14-0]

C₁₅H₁₇NO₇ 323.302

Constit. of *Anthoceros agrestis*.

N-Me: *N*-Methyl-*L*-glutamic acid

[6753-62-4]

C₆H₁₁NO₄ 161.157

Isol. from the mollusc *Scapharca broughtonii*. Hygroscopic solid (as hydrochloride). Mp 139-142° (hydrochloride). [α]_D²⁵ +2.7 (c, 7.0 in H₂O).

N-(2-Pyridinylethyl): [146500-72-3]

C₁₂H₁₆N₂O₄ 252.269

Isol. from the toxic mushroom *Clitocybe acromelalga*. Neurotoxic, shows depolarising activity. Mycotoxin. Sol. H₂O, MeOH; poorly sol. EtOAc, hexane. [α]_D²⁵ +129 (c, 0.09 in H₂O). λ_{max} 260 (ε 2140) (H₂O) (Berdy).

[63663-21-8, 7558-63-6, 19285-83-7, 24938-00-9, 25683-11-8, 997-42-2, 11070-68-1, 25513-46-6, 21752-29-4, 5996-22-5]

Jungermann, E. et al., *J.A.C.S.*, 1956, **78**, 172-174 (*N*-dodecanoyl)

Takahira, M. et al., *J. Am. Oil Chem. Soc.*,

1972, **49**, 157-161 (*N*-dodecanoyl)

Hilbig, S. et al., *Angew. Chem., Int. Ed.*, 1985,

24, 1063 (*Xanthodermin*)

Yamano, K. et al., *Tetrahedron*, 1993, **49**, 2427

(*N*-pyridinylethyl)

Trennheuser, F. et al., *Phytochemistry*, 1994,

37, 899-903 (*Anthoceros amides*)

Yamano, K. et al., *Z. Naturforsch., C*, 1994,

49, 157 (*N*-4-Aminobutanoylglutamic acid)


Burger, K. et al., *Eur. J. Org. Chem.*, 2000, 199-

204 (*N*-Me)

Tarui, A. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 2003, **134**, 79-84 (*N*-Me *Scapharca constits*)

Glutamine, 9CI, INN, USAN **G-108**

2,5-Diamino-5-oxopentanoic acid. Glutamic acid 5-amide. γ-Glutamine. Gln. Glu. min. NSC 97925 [6899-04-3]



C₅H₁₀N₂O₃ 146.146

Log P -3.46 (calc).

(R)-form

D-form

[5959-95-5]

Mp 186-188°. [α]_D²⁰ -7.9 (c, 4 in H₂O).

(S)-form

L-form. **Levoglutamide**. *Energlut*. *Glutaven*. *Iperphos*. *Memoril*. *Nutrestore*. *FEMA 3684*. Many other names [56-85-9]

Widely distributed in plants, e.g. beetroot. Flavouring ingredient. Dietary supplement. Needles (EtOH aq.). Spar. sol. H₂O (3.6 g/100 g at 25°), EtOH. Mp 184-185° dec. [α]_D¹⁹ +9.2 (H₂O). [α]_D +46.5 (5M HCl). pK_{a1} 2.17; pK_{a2} 9.01 (25°, 0.1M KNO₃). Isoelectric point 5.65. Sweet taste.

► LD₅₀ (rat, orl) 7500 mg/kg. Exp. reprod. effects (very large dose). MA2275100

Me ester: [32668-14-7]

C₆H₁₂N₂O₃ 160.172

Cryst. (MeOH/Et₂O) (as hydrochloride). Mp 142° (hydrochloride).

p-Nitrobenzyl ester; *hydrobromide*:

[14349-18-9]

Cryst. Mp 175°.

*N*²-(2-Nitrobenzenesulfonyl): *N*²-[(2-Nitrophenyl)thio]-*L*-glutamine, 9CI

[7685-73-6]

C₁₁H₁₃N₃O₅S 299.307

Cryst. Mp 165° (158-160°). [α]_D -74.3 (c, 2 in DMF).

*N*²-Tetradecanoyl: *N*²-Tetradecanoylglutamine. *N*²-Myristoylglutamine

[100652-49-1]

C₁₉H₃₆N₂O₄ 356.504

Constit. of the oral secretions of various caterpillars incl. *Operophtera* sp. and *Spodoptera exigua*.

*N*²-(9Z-Tetradecenyl): *N*²-(9-Tetradecenyl)glutamine. *N*²-Myristoleoylglutamine

[247150-77-2]

C₁₉H₃₄N₂O₄ 354.489

Constit. of the oral secretions of various caterpillars incl. *Heliothis virescens* and *Spodoptera* sp.

*N*²-Hexadecanoyl: *N*²-Hexadecanoylglutamine. *N*²-Palmitoylglutamine

[58725-34-1]

C₂₁H₄₀N₂O₄ 384.558

Constit. of the oral secretions of various caterpillars incl. *Epirrita autumnata* and *Spodoptera* spp.


*N*²-(9Z-Hexadecenyl): *N*²-(9-Hexadecenyl)glutamine. *N*²-Palmitoleoylglu-

- tamine*
[247150-75-0]
C₂₁H₃₈N₂O₄ 382.542
Constit. of the oral secretions of various caterpillars incl. *Chloroclysta truncata* and *Spodoptera* spp.
- N²-(9Z-Octadecenoyl): N²-(9-Octadecenoyl)glutamine. N²-Oleoylglutamine
[247150-73-8]
C₂₃H₄₂N₂O₄ 410.596
Constit. of the oral secretions of various caterpillars incl. *Operophtera* sp. and *Spodoptera* spp.
- N²-(9Z,12Z-Octadecadienoyl): N²-(9,12-Octadecadienoyl)glutamine. N²-Linoleoylglutamine
[247150-70-5]
C₂₃H₄₀N₂O₄ 408.58
Constit. of the oral secretions of various caterpillars incl. *Heliothis virescens* and *Spodoptera* spp.
- N²-(9Z,12Z,15Z-Octadecatrienoyl): N²-(9,12,15-Octadecatrienoyl)glutamine. N²-Linolenoylglutamine
[157598-98-6]
C₂₃H₃₈N₂O₄ 406.564
Constit. of the oral secretions of various caterpillars incl. *Epirrita autumnata* and *Spodoptera* spp.
- N²-(15,16-Epoxy-9Z,12Z-octadecadienoyl): N²-(15,16-Epoxy-9,12-octadecadienoyl)glutamine. N²-(15,16-Epoxylinoleoyl)glutamine
C₂₃H₃₈N₂O₅ 422.564
Constit. of the oral secretions of the caterpillars *Spodoptera exigua* and *Spodoptera frugiperda*.
- N²-(17ξ-Hydroxy-9Z,12Z-octadecadienoyl): N²-(17-Hydroxy-9,12-octadecadienoyl)glutamine. N²-(17-Hydroxylinoleoyl)glutamine
[247150-68-1]
C₂₃H₄₀N₂O₅ 424.579
Constit. of the oral secretions of various caterpillars incl. *Heliothis virescens* and *Spodoptera* spp.
- N²-(15,16-Dihydroxy-9Z,12Z-octadecadienoyl): N²-(15,16-Dihydroxy-9,12-octadecadienoyl)glutamine. N²-(15,16-Dihydroxylinoleoyl)glutamine
C₂₃H₄₀N₂O₆ 440.579
Constit. of the oral secretions of the caterpillars *Spodoptera exigua* and *Spodoptera frugiperda*.
- N-(17S-Hydroxy-9Z,12Z,15Z-octadecatrienoyl): N²-(17-Hydroxylinolenoyl)-glutamine. **Volicitin**
[191548-06-8]
[191547-98-5]
C₂₃H₃₈N₂O₅ 422.564
Isol. from the oral secretions of the caterpillars, *Helicoverpa armigera*, *Mythimna separata* and *Spodoptera exigua*. Elicitor of plant volatiles. [α]_D²¹ +6.6 (c, 0.8 in MeOH).
- N²-Carbamoyl, amide: 2-[(Aminocarbo-nyl)amino]pentanediamide, **9CI**. N²-Carbamoylglutamine amide
[134958-17-1]
C₆H₁₂N₄O₃ 188.186
Isol. from *Ectothiorhodospira merismortui*. [α]_D²⁰ +10.5 (c, 0.6 in DMSO).
- N⁵-Isopropyl: N⁵-Isopropylglutamine
[4311-12-0]
C₈H₁₆N₂O₃ 188.226
Isol. from seeds of *Lunaria annua*. Cryst. (EtOH aq.). [α]_D²² +7.1 (c, 1.7 in H₂O).
- N⁵-(2-Hydroxyethyl): N-(γ-Glutamyl)ethanolamine, **8CI**
[2650-74-0]
C₇H₁₄N₂O₄ 190.199
Constit. of *Agaricus bisporus* (button mushroom), seeds of *Lunaria annua* and defence secretions of the beetle *Chrysolina coerulans*. Cryst. (EtOH aq.). Mp 188.5-191°. [α]_D¹⁹ +5.8 (c, 1.8 in H₂O).
- N⁵-(4-Hydroxyphenyl): γ-Glutaminyl-4-hydroxybenzene
[30382-24-2]
C₁₁H₁₄N₂O₄ 238.243
Isol. from the gill tissue of the edible mushrooms *Agaricus bisporus* (button mushroom) and *Agaricus hortensis*. Neoplasm inhibitor. Sol. H₂O; fairly sol. MeOH. Mp 231.5° (225-226°). [α]_D +30 (1M HCl). λ_{max} 246 (H₂O) (Berdy). λ_{max} 246 (HCl) (Berdy). λ_{max} 260 (NaOH) (Berdy).
- ▶ LD₅₀ (mus, ipr) 5000 mg/kg. MA2276400
- N⁵-(4-Methoxybenzyl): N⁵-(4-Methoxybenzyl)glutamine
C₁₃H₁₈N₂O₄ 266.296
Constit. of the xylem sap of the roots of *Cucurbita maxima* x *Cucurbita moschata*. Amorph. solid. Mp 214-215° dec. [α]_D²² +23 (c, 0.1 in 1M HCl). λ_{max} 224 (log ε 4.1); 273 (log ε 3.2) (H₂O).
- (±)-form [585-21-7]
Prisms (Me₂CO aq.). Mp 186° dec.
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 782D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1278B (nmr)
Archibald, R.M. *et al.*, *Chem. Rev.*, 1945, **37**, 161 (rev, props)
Lepp, A. *et al.*, *Biochem. Prep.*, 1957, **5**, 79 (synth)
Jadot, J. *et al.*, *Biochim. Biophys. Acta*, 1960, **43**, 322 (deriv)
Greenstein, J.P. *et al.*, *Chemistry of the Amino Acids*, (Chapter 25), Wiley, N.Y., 1961, 1929 (isol, synth)
Kline, G.B. *et al.*, *J.O.C.*, 1961, **26**, 1854 (synth)
Zervas, L. *et al.*, *J.A.C.S.*, 1963, **85**, 3660-3666 (2-nitrobenzenesulfonyl)
Larsen, P.O. *et al.*, *Acta Chem. Scand.*, 1965, **19**, 1071 (N⁵-Isopropylglutamine)
Olesen Larsen, P. *et al.*, *Acta Chem. Scand.*, 1967, **21**, 1592-1604 (N⁵-2-hydroxyethyl, isol, synth)
Bak, B. *et al.*, *J. Mol. Spectrosc.*, 1968, **26**, 78 (pmr)
Weaver, R.F. *et al.*, *J. Biol. Chem.*, 1971, **246**, 2010 (γ-Glutaminyl-4-hydroxybenzene)
Koetzle, T.F. *et al.*, *Acta Cryst. B*, 1973, **29**, 2571 (cryst struct)
Voelter, W. *et al.*, *Monatsh. Chem.*, 1974, **105**, 1110 (cmr)
Synth. Prod. Util. Amino Acids, (Kaneko, T. *et al.*, Ed.), Halsted Press, 1974, 109 (synth)
- Lindberg, P. *et al.*, *J.C.S. Perkin 1*, 1977, 684
London, R.E. *et al.*, *J.A.C.S.*, 1978, **100**, 3723 (cmr)
Jaeger, P. *et al.*, *N. Engl. J. Med.*, 1986, **315**, 1120 (rev)
Tachiki, T. *et al.*, *Prog. Ind. Microbiol.*, 1986, **24**, 121 (rev, synth)
Yoshifuji, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2994 (synth)
Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 328
Van Oyccke, S. *et al.*, *Bull. Soc. Chim. Belg.*, 1988, **97**, 297-311 (N⁵-2-hydroxyethyl, isol, synth)
Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, GFO050
Galinski, E.A. *et al.*, *Eur. J. Biochem.*, 1991, **198**, 593 (N-Carbamoylglutamineamide)
Martindale. The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 598
Suresh, S. *et al.*, *Acta Cryst. C*, 1996, **52**, 1313 (cryst struct, DL-form)
Alborn, H.T. *et al.*, *Science (Washington, D.C.)*, 1997, **276**, 945-949 (Volicitin, isol)
Pohnert, G. *et al.*, *Chem. Comm.*, 1999, 1087-1088 (Volicitin, synth)
Inouye, Y. *et al.*, *Phytochemistry*, 1999, **51**, 425-428 (N⁵-4-Methoxybenzylglutamine)
Pohnert, G. *et al.*, *Tetrahedron*, 1999, **55**, 11275-11280 (caterpillar amides)
Spiteller, D. *et al.*, *Tet. Lett.*, 2001, **42**, 1483-1485 (Volicitin, abs config)
Itoh, S. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 1591-1596 (Volicitin, synth)
Mori, N. *et al.*, *Biosci., Biotechnol., Biochem.*, 2003, **67**, 1168-1171 (caterpillar amides)
Spiteller, D. *et al.*, *Tetrahedron*, 2003, **59**, 135-139 (epoxyoctadecadienoyl, dihydroxyoctadecadienoyl derivs)
Sawada, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 2185-2190 (Volicitin, isol, abs config)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GFO050

Glyantrypine

G-109

[142382-42-1]

C₂₀H₁₆N₄O₂ 344.372

Incorrect MF given in the lit. Alkaloid from *Aspergillus clavatus*. Mycotoxin. Amorph. solid. λ_{max} 224 ; 270 (MeOH) (Berdy).


Penn, J. *et al.*, *J.C.S. Perkin 1*, 1992, 1495-1496 (isol, uv, pmr, cmr, ms, struct)

Wang, H. *et al.*, *J.O.C.*, 2000, **65**, 1022-1030 (synth)

Liu, J.-F. *et al.*, *J.O.C.*, 2005, **70**, 6339-6345
(*synth*)

Glybomine C **G-110**

3-Methyl-5-(3-methyl-2-butenyl)-9H-carbazole-2,6-diol. 2,6-Dihydroxy-3-methyl-5-prenyl-9H-carbazole



$C_{18}H_{19}NO_2$ 281.354

Alkaloid from the stems of *Glycosmis arborea*. Oil. λ_{max} 218 ; 237 ; 266 ; 309 ; 338 (MeOH).

6-Me ether: 2-Hydroxy-6-methoxy-3-methyl-5-prenyl-9H-carbazole. **Glybomine B**


$C_{19}H_{21}NO_2$ 295.38

Alkaloid from the stems of *Glycosmis arborea*. Oil. λ_{max} 217 ; 239 ; 263 (sh) ; 307 ; 336 (MeOH).

Ito, C. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1488-1491
(*isol*, *pmr*, *cmr*)

Glycerinopyrin **G-111**

2,3-Dihydroxypropyl 1-hydroxy-4-methyl-1H-pyrrole-2-carboxylate
[129966-44-5]



$C_9H_{13}NO_5$ 215.205


Metab. of *Streptomyces violaceus*. Cryst. powder. Sol. $CHCl_3$, DMSO, Me_2CO , THF, MeOH; fairly sol. H_2O ; poorly sol. hexane. Mp 93-96°. $[\alpha]_D^{20} +7.11$ (c, 1.25 in MeOH). λ_{max} 278 (ϵ 21200) (MeOH) (Berdy).

Schoenewolf, M. *et al.*, *Angew. Chem., Int. Ed.*, 1991, **30**, 183 (*biosynth*)

Schoenewolf, M. *et al.*, *Annalen*, 1991, 77 (*isol*, *pmr*, *cmr*)

Glycobismine A **G-112**

[91147-20-5]



$C_{37}H_{34}N_2O_6$ 602.685


(±)-form

Alkaloid from the bark of *Glycosmis citrifolia* (Rutaceae). Shows *in vitro* antimalarial activity. Yellow needles. Mp 256-258°. The first naturally occurring C-C linked bisacridone alkaloid.

Furukawa, H. *et al.*, *J.C.S. Perkin 1*, 1993, 471
(*uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Glycobismine B **G-113**

[148717-57-1]



$C_{37}H_{34}N_2O_7$ 618.685

Alkaloid from root and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow oil. Racemic.


Diastereoisomer: **Glycobismine C**

[148810-27-9]

$C_{37}H_{34}N_2O_7$ 618.685

Alkaloid from root and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow oil.


Furukawa, H. *et al.*, *J.C.S. Perkin 1*, 1993, 471
(*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Glycobismine D **G-114**

$C_{35}H_{30}N_2O_{10}$ 638.629

Alkaloid from *Glycosmis citrifolia*. Yellow oil. Racemic. λ_{max} 202 (log ϵ 4.61); 225 (log ϵ 4.39); 265 (log ϵ 4.73); 294 (sh) (log ϵ 4.46); 335 (log ϵ 4.25); 405 (log ϵ 3.82) (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 65-70

Glycobismine E**G-115**

$C_{39}H_{34}N_2O_9$ 674.706

Alkaloid from *Glycosmis citrifolia*. Yellow oil. Racemic. λ_{max} 202 ; 211 ; 269 ; 292 (sh) ; 317 ; 340 ; 407 (MeOH).


O-De-Me: **Glycobismine F**

$C_{38}H_{32}N_2O_9$ 660.679

Alkaloid from the roots of *Glycosmis citrifolia*. Yellow oil. Racemic. λ_{max} 270 ; 282 (sh) ; 316 (sh) ; 340 (sh) ; 406 (EtOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 65-70 (*isol*, *pmr*, *cmr*)

Negi, N. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 362-364 (*Glycobismine F*)

Glycobismine G**G-116**


$C_{38}H_{32}N_2O_9$ 660.679

Alkaloid from the roots of *Glycosmis citrifolia*. Yellow oil. Racemic. λ_{max} 270 ; 284 (sh) ; 296 (sh) ; 344 (sh) ; 406 (MeOH).

Negi, N. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 362-364 (*isol*, *pmr*, *cmr*, *ms*)

Glycocinnaspermicidin A **G-117**

LL-BM 123 β_1 . BM 123 β_1 . Antibiotic BM 123 β_1 . Antibiotic LL-BM 123 β_1
[62251-27-8]



C₃₆H₆₁N₁₃O₁₂ 867.958

Glycolipid antibiotic. Related to Cincodine I, C-459. Isol. from *Nocardia* spp. Active against gram-positive and -negative bacteria. Amorph. powder (as hydrochloride). Sol. H₂O; fairly sol. MeOH; poorly sol. Et₂O, hexane, butanol. $[\alpha]_D^{25} +67$ (c, 1 in H₂O). Hydrochloride dec. at ca. 200°. λ_{\max} 286 (E1%/1cm 200) (MeOH) (Berdy). λ_{\max} 286 (E1%/1cm 260) (H₂O) (Berdy). λ_{\max} 286 (HCl) (Berdy).


4'-O-Deglycosyl: Glyocinnaspermicidin**D**

[99260-73-8]

C₃₀H₅₀N₁₀O₉ 694.787

Isol. from *Nocardia* sp. M9615-7F6. Possesses a broad antibacterial spectrum; also active against leukaemia L-1210 cells. Powder. Sol. H₂O; poorly sol. Me₂CO, hexane. Mp 300° dec. $[\alpha]_D^{18} +88$ (c, 0.5 in H₂O). λ_{\max} 218 (€ 13500); 287 (€ 21900) (H₂O) (Derep). λ_{\max} 218 (€ 10750); 287 (€ 21900) (H₂O) (Berdy).

► LD₅₀ (mus, ivn) 25 - 50 mg/kg.
UC6230000

Tresner, H. *et al.*, *J. Antibiot.*, 1978, **31**, 394 (isol)Martin, J.H. *et al.*, *J. Antibiot.*, 1978, **31**, 398 (isol)Ellestad, G.A. *et al.*, *J.A.C.S.*, 1978, **100**, 2515 (struct)Araki, K. *et al.*, *Tet. Lett.*, 1982, **23**, 1705 (synth)Dobashi, K. *et al.*, *J. Antibiot.*, 1985, **38**, 1166 (Glyocinnaspermicidin D)Nishiyama, T. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 4372-4375 (Glyocinnaspermicidin D, synth)**Glycocitlone A****G-118**C₁₆H₁₉NO₃ 273.331

Alkaloid from *Glycosmis citrifolia*. Yellow oil. λ_{\max} 230 (log € 4.46); 286 (sh) (log € 3.93); 296 (log € 4.02); 309 (log € 4) (MeOH).

3'-Deoxy, 3',4'-didehydro: N-Methylschinifoline

[149998-43-6]

C₁₆H₁₇NO₂ 255.316

Alkaloid from roots of *Zanthoxylum schinifolium* and *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil.

3'-Deoxy, 3',4'-didehydro, N-de-Me: 4-Methoxy-3-(3-methyl-1,3-butadienyl)-2(1H)-quinolinone. Schinifoline†

[149998-56-1]

C₁₅H₁₅NO₂ 241.289

Alkaloid from roots of *Zanthoxylum schinifolium*, *Glycosmis cyanocarpa* and *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Cryst. Mp 171-174° dec.

8-Hydroxy: Glycocitlone BC₁₆H₁₉NO₄ 289.33Alkaloid from *Glycosmis citrifolia*.

Amorph. yellow powder. λ_{\max} 202 (log € 4.41); 225 (log € 4.25); 267 (log € 4.21); 296 (log € 4.23); 307 (log € 4.25); 410 (log € 3.3) (MeOH).

8-Methoxy: Glycocitlone CC₁₇H₂₁NO₄ 303.357Alkaloid from *Glycosmis citrifolia*.


Yellow oil. λ_{\max} 230 (log € 4.1); 258 (log € 4.05); 307 (log € 3.88); 321 (sh) (log € 3.81); 350 (log € 3.39) (MeOH).

Brader, G. *et al.*, *Annalen*, 1993, 355


(Schinifoline, N-Methylschinifoline)

Wurz, G. *et al.*, *Nat. Prod. Lett.*, 1993, **3**, 177 (Schinifoline, isol)Ito, C. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 65-70 (Glycocitlones)**Glycocitridine****G-119**

1,2-Dihydro-4,7,8-trimethoxy-2-oxo-3-quinolinecarboxaldehyde, 9CI
[167504-57-6]

C₁₃H₁₃NO₅ 263.249


Alkaloid from leaves of *Glycosmis citrifolia* (Rutaceae). Needles. Mp 174-175°.

Wu, T.-S. *et al.*, *Phytochemistry*, 1995, **39**, 1453 (isol, uv, ir, pmr, ms, struct)**Glycocitrine V****G-120**C₂₂H₂₅NO₈ 431.441


Alkaloid from *Glycosmis citrifolia*. Yellow oil. Racemic. λ_{\max} 202 (log € 4.22); 226 (sh) (log € 4); 263 (log € 4.27); 274 (log € 4.28); 334 (log € 3.84); 399 (log € 3.4) (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 65-70**Glycocitrine III****G-121**

2-(3,7-Dimethyl-2,6-octadienyl)-1,3,5-trihydroxy-10-methyl-9(10H)-acridinone. 2-Geranyl-1,3,5-trihydroxy-1-methylacridone

C₂₄H₂₇NO₄ 393.482Alkaloid from the stems of *Glycosmis*


pentaphylla. Yellow oil. λ_{\max} 214 ; 228 ; 255 ; 267 ; 284 ; 308 (sh) ; 336 (sh) ; 401 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1579-1581**Glycocitrine VI****G-122**C₂₄H₂₇NO₃ 377.482

Alkaloid from *Glycosmis citrifolia*. Yellow oil. λ_{\max} 203 ; 229 ; 286 ; 296 ; 340 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 65-70**Glycoline****G-123**


2,11-Dihydro-5,10-dihydroxy-2,11-dimethyl-2-(4-methyl-3-pentenyl)-6H-pyrano[3,2-b]acridin-6-one

C₂₄H₂₅NO₄ 391.466**(R)-form** [82513-80-2]

Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Orange needles (Et₂O). Mp 216-218°. $[\alpha]_D -15.1$ (c, 1.0 in CHCl₃).

Wu, T.-S. *et al.*, *Heterocycles*, 1982, **19**, 825; *J.C.S. Perkin 1*, 1983, 1681 (isol, uv, ir, pmr, cmr, ms, cd, struct)**Glycolone†****G-124**

3-(2-Butenyl)-4-ethoxy-8-hydroxy-1-methyl-2(1H)-quinolinone

C₁₆H₁₉NO₃ 273.331


Note exceptional opportunities for confusion with Glycolone in T-583. Alkaloid from the root bark of *Glycosmis pentaphylla* (Rutaceae). Fine pale-yellow needles (CCl₄/cyclohexane). Mp 160-162°.

Sinha, S.K.P. *et al.*, *Indian J. Chem., Sect. B*, 1988, **27**, 460 (isol, uv, ir, pmr, struct)

Glycomauroll

G-125

6-Methyl-4-(3-methyl-2-butenyl)-9H-carbazol-3-ol, 9CI. 3-Hydroxy-6-methyl-4-prenylcarbazole
[125287-18-5]



$C_{18}H_{19}NO$ 265.354

Alkaloid from the stem bark of *Glycosmis mauritiana* (Rutaceae). Pale-yellow needles (MeOH), Mp 149-150°.

Kumar, V. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1375


(*isol, uv, ir, pmr, cmr, ms, struct*)

Lebold, T.P. *et al.*, *Org. Lett.*, 2007, **9**, 1883-1886 (*synth*)

Glyothiohexide α

G-126

LL-14E605 α . Antibiotic LL-14E605 α
[158446-31-2]



$C_{58}H_{57}N_{13}O_{15}S_6$ 1368.562

Cyclic depsipeptide antibiotic. Prod. by *Sebekia benihana* LL-14E605. Active against gram-positive bacteria. Sol. CH_2Cl_2 -MeOH; fairly sol. MeOH; poorly sol. Me_2CO , H_2O , hexane. Related to Antibiotic S 54832A, A-1237 and Nosiheptide, N-319. λ_{max} 302 (ϵ 24900); 352 (ϵ 19200); 370 (ϵ 16000) (MeOH/HCl). λ_{max} 295 (ϵ 28800); 400 (ϵ 15300) (MeOH/NaOH). λ_{max} 302 (ϵ 27000); 370 (ϵ 11900); 400 (ϵ 8400) (MeOH).

Stereoisomer: **Glyothiohexide β** . LL-E605 β . Antibiotic LL-E605 β

$C_{58}H_{57}N_{13}O_{15}S_6$ 1368.562

Prod. by *Sebekia benihana* LL-14E605.


Steinberg, D.A. *et al.*, *J. Antibiot.*, 1994, **47**, 887-893; 894-900; 901-908 (*isol, pmr, cmr, struct*)

U.S. Pat., 1995, 5 451 581; *CA*, **124**, 143742h (*Glyothiohexide β*)

Glycozalone A

G-127

2-Benzyl-2,3-dihydro-1-methyl-4(1H)-quinazolinone. Glycozoline A



$C_{16}H_{16}N_2O$ 252.315

(\pm)-**form** [26750-21-0]

Alkaloid from leaves of *Glycosmis cochinchinensis*. Oil (nat.); cryst. ($CHCl_3$ /EtOH)(synthetic). Mp 199-200° (synthetic). λ_{max} 205 (log ϵ 4.25); 226 (log ϵ 4.3); 262 (log ϵ 3.6); 355 (log ϵ 3.22) (MeOH).

4'-Methoxy: 2,3-Dihydro-2-(4-methoxybenzyl)-1-methyl-4(1H)-quinazolinone.

Glycozalone B. Glycozoline B

[252234-33-6]

$C_{17}H_{18}N_2O_2$ 282.341


Alkaloid from leaves of *Glycosmis cochinchinensis*. Oil. λ_{max} 203 (log ϵ 4.4); 226 (log ϵ 4.37); 266 (log ϵ 3.75); 319 (log ϵ 3.41); 367 (log ϵ 3.3) (MeOH).

Chakravarti, D. *et al.*, *J.C.S.*, 1953, 3337-3340 (*synth*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1491-1493 (*isol, ir, pmr, cmr*)

Glymontanine A

G-128



Absolute Configuration

$R^1 = H, R^2 = -CH_2NHCOSMe$

$C_{19}H_{21}NO_8S$ 423.443

Constit. of the aerial parts of *Glycosmis montana*. Powder. $[\alpha]_D^{25}$ -18.4 (c, 0.45 in MeOH).

Wang, J. *et al.*, *Tet. Lett.*, 2005, **46**, 169-172 (*isol, cd, pmr, cmr*)

Glymontanine B

G-129

As Glymontanine A, G-128 with $R^1 = -CH_2NHCOSMe, R^2 = H$

$C_{19}H_{21}NO_8S$ 423.443


Constit. of the aerial parts of *Glycosmis montana*. Powder. $[\alpha]_D^{25}$ -13.3 (c, 0.25 in MeOH).

Wang, J. *et al.*, *Tet. Lett.*, 2005, **46**, 169-172 (*isol, cd, pmr, cmr*)

Glypetelotine

G-130

S-Methyl [2-(1H-indol-3-yl)ethyl]-methylcarbamothioate
[259821-95-9]



$C_{13}H_{16}N_2OS$ 248.348


Alkaloid from the leaves of *Glycosmis petelotii*. Cryst. (petrol/EtOAc). Mp 92.3°. λ_{max} 203 (log ϵ 5.3); 222 (log ϵ 4.66); 282 (log ϵ 3.88); 290 (log ϵ 3.8) (EtOH).

Cuong, N.M. *et al.*, *Phytochemistry*, 1999, **52**, 1711-1714

Glyserin A

G-131

BU 2349A. Antibiotic BU 2349A
[78213-56-6]



$C_{44}H_{75}N_7O_{18}$ 990.113

Glycoside antibiotic. Isol. from *Bacillus cereus*. Active against gram-negative and -positive bacteria. Inhibitor of reverse transcriptase. Sol. H_2O ; fairly sol. MeOH, DMF, DMSO, EtOH; poorly sol. Me_2CO , hexane. λ_{max} 247 (E1%/1cm 131) (H_2O) (Berdy). λ_{max} 247 (HCl) (Berdy). λ_{max} 247 (NaOH aq.) (Berdy).

► LD₅₀ (mus, ivn) 35 mg/kg, LD₅₀ (mus, ims) 285 mg/kg. MD4787000

Kawaguchi, H. *et al.*, *J. Antibiot.*, 1981, **34**, 381 (*isol*)

Tsuno, T. *et al.*, *J. Antibiot.*, 1981, **34**, 390 (*struct*)

Glyserin B

G-132

BU 2349B. Antibiotic BU 2349B
[78213-55-5]

As Glyserin A, G-131 with $R^1 = H, R^2R^3 = CH_2$

$C_{40}H_{66}N_6O_{18}$ 918.991

Glycoside antibiotic. Isol. from *Bacillus cereus*. Active against gram-positive and -negative bacteria. Reverse transcriptase inhibitor. Amorph. powder (as carbonate salt). Sol. H_2O ; fairly sol. MeOH, EtOH; poorly sol. Me_2CO , hexane. Mp 166° dec. (carbonate). $[\alpha]_D^{25}$ +132 (c, 0.5 in H_2O) (carbonate). λ_{max} 247 (E1%/1cm 148) (H_2O) (Berdy). λ_{max} 247 (HCl) (Berdy). λ_{max} 247 (NaOH aq.) (Berdy).

Kawaguchi, H. *et al.*, *J. Antibiot.*, 1981, **34**, 381 (*isol*)

Tsuno, T. *et al.*, *J. Antibiot.*, 1981, **34**, 390 (*struct*)

Glyserin C

G-133

BU 2349C. Antibiotic BU 2349C
[78213-54-4]

As Glyserin A, G-131 with $R^1 = (CH_2)_4NH_2, R^2 = CH_2OH, R^3 = H$
 $C_{44}H_{77}N_7O_{19}$ 1008.128

Glycoside antibiotic. Isol. from *Bacillus cereus*. Reverse transcriptase inhibitor. Amorph. powder (as carbonate salt). Sol. H_2O ; fairly sol. MeOH, EtOH, DMF, DMSO; poorly sol. butanol, hexane. Mp 140-145° (carbonate). $[\alpha]_D^{25}$ +157 (c, 0.5 in H_2O) (carbonate). λ_{max} 247 (E1%/1cm 133) (H_2O) (Berdy). λ_{max} 247 (HCl) (Berdy). λ_{max} 247 (NaOH aq.) (Berdy).

Kawaguchi, H. *et al.*, *J. Antibiot.*, 1981, **34**, 381 (*isol*)

Tsuno, T. *et al.*, *J. Antibiot.*, 1981, **34**, 390 (*struct*)


Gnidine

G-134


$C_{29}H_{51}N_3$ 441.742

Struct. unknown. Alkaloid from *Lycopodium gnidioides* (Lycopodiaceae).


MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Gnidinine**G-135**C₂₉H₅₁N₃O 457.741Struct. unknown. Alkaloid from *Lycopodium gnidioides* (Lycopodiaceae).MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241**Goadsporin****G-136**C₇₂H₉₇N₁₉O₂₀S₂ 1612.806Prod. by *Streptomyces* sp. TP-A0584.Promotes secondary metab. and morphogenesis in streptomycetes. Powder. Mp 172-174°. [α]_D²² -21 (c, 1 in MeOH). λ_{max} 204 (log ε 4.93); 226 (log ε 4.91) (MeOH).Onaka, H. *et al.*, *J. Antibiot.*, 2001, **54**, 1036-1044; 1045-1053 (*isol, uv, pmr, cmr, ms, activity*)Fujii, K. *et al.*, *Tetrahedron*, 2002, **58**, 6873-6879 (*abs config*)**Goebeline****G-137**

13,14-Didehydro-[14,14'-bimatridine]-15,15'-dione, 9CI. Gebeline [25908-93-4]

C₃₀H₄₄N₄O₂ 492.703Dimeric matrine alkaloid; dimer of Sophocarpine, S-371. Artifact formed in thermal treatment of matrine alkaloids. Mp 231° (228-230°). [α]_D -13.2 (EtOH).*Hydroiodide:*Cryst. (H₂O). Mp 361-362°.*Perchlorate:*


Cryst. (MeOH). Mp 315-316°.

Pakanaev, Ya.I. *et al.*, *Zh. Obshch. Khim.*, 1961, **31**, 2428; *J. Gen. Chem. USSR (Engl. Transl.)*, 1961, **31**, 2263 (*isol*)Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 347-350; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 340-343 (*ms, struct*)Sadykov, B. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 606-610; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 635-638 (*synth, ir*)**Golmaenone****G-138**

Absolute Configuration

C₁₉H₂₁N₃O₄ 355.393Alkaloid from the marine fungus *Aspergillus* (culture MFA 212). Radical scavenger and uv protectant. Yellow cryst. (CHCl₃). Mp 160-161°. [α]_D +7.1 (c, 0.4 in CHCl₃). CAS no. not found CA 141. λ_{max} 222 (log ε 1.8); 247 (log ε 2); 327 (log ε 1.9); 368 (log ε 1.7) (CHCl₃).Li, Y. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 375-376 (*isol, cd, pmr, cmr*)**Gomandonine****G-139**

16,17-Epoxy-4,21-dimethyl-7,20-cyclooctadane-1,13,15-triol, 9CI [110382-21-3]

C₂₁H₃₁NO₄ 361.48Alkaloid from the roots of *Aconitum subcuneatum* (Ranunculaceae). Cryst. (MeOH). Mp 248-249°. [α]_D¹² -42.5 (c, 0.12 in MeOH).

13-Ac: 13-Acetylgomandonine

[120122-19-2]

C₂₃H₃₃NO₅ 403.517Minor alkaloid from *Aconitum delphinifolium* (Ranunculaceae). Amorph. solid.1,13-Di-Ac: **Yesoxine**

[112561-70-3]

C₂₅H₃₅NO₆ 445.555Alkaloid from rhizomes of *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Mp 184° dec. [α]_D -37.5 (c, 0.32 in EtOH).Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2615 (*ir, pmr, cmr, ms, cryst struct*)Bando, H. *et al.*, *Heterocycles*, 1987, **26**, 2623 (*Yesoxine*)Kulathaiyel, P. *et al.*, *Phytochemistry*, 1988, **27**, 3998 (*13-Acetylgomandonine*)**Gongrine****G-140**

4-[[[(Aminoiminomethyl)amino]carbonyl]amino]butanoic acid, 9CI. γ-(Guanylureido)butanoic acid. 1-Amidino-3-(3-carboxypropyl)urea. γ-(Amidinoureido)-butyric acid [5998-99-2]

HN=C(NH₂)NHCONH(CH₂)₃COOHC₆H₁₂N₄O₃ 188.186Isol. from red algae *Gymnogongrus flabelliformis* and *Grateloupia filicina*.Prisms (H₂O). Mp 208-209° dec.*Picrate:* [21808-08-2]


Yellow needles. Mp 208-210° dec.

Sinters at ca. 196°.

Ito, K. *et al.*, *Agric. Biol. Chem.*, 1965, **29**, 832; 1969, **33**, 237 (*isol, synth*)Frankel, M. *et al.*, *J.C.S. (C)*, 1967, 2698 (*synth*)Wakamiya, T. *et al.*, *Tetrahedron*, 1984, **40**, 235 (*isol, bibl*)**Goniomine****G-141**

1,2,5,6,6a,7,8,12c-Octahydro-6a-hydroxy-7-methyl-14-methylene-4H-6,12b-ethanoindolizino[8,1-cd][1]benzazepin-13-one, 9CI

[74503-97-2]




Absolute Configuration

C₁₉H₂₂N₂O₂ 310.395A rearranged condylocarpan. Alkaloid from *Gonioma malagasy* (Apocynaceae). Cryst. (MeOH). Mp 223°. [α]_D²⁰ -270 (c, 0.5 in CHCl₃).*Dihydro:* [74503-99-4]


Cryst. (EtOH). Mp 228°.

Chiaroni, A. *et al.*, *J.A.C.S.*, 1980, **102**, 5920 (*ir, uv, cmr, pmr, ms, cryst struct*)**Goniomitine****G-142**

1,2,3,4,4a,5,6,12a-Octahydroindolo[1,2-a][1,8]naphthyridine-7-ethanol, 9CI [109794-95-8]



Absolute configuration


C₁₉H₂₆N₂O 298.427Highly rearranged indole alkaloid related to the *Aspidosperma* group. Alkaloid from the root bark of *Gonioma malagasy*. Cryst. (Et₂O/MeOH). Mp 150°. [α]_D²⁰ -80 (c, 0.9 in CHCl₃).Randriambola, L. *et al.*, *Tet. Lett.*, 1987, **28**, 2123-2126 (*uv, ir, pmr, cmr, ms*)Takano, S. *et al.*, *Chem. Comm.*, 1991, 462-464 (*synth, abs config*)Morales, C.L. *et al.*, *Org. Lett.*, 2008, **10**, 157-159 (*synth*)**Gorchacoine****G-143***Gortschakoine*

C₁₉H₂₃NO₃ 313.396**(R)-form** [16336-17-7]Alkaloid from *Corydalis gortschakovii*. Noncryst. [α]_D²⁰ -40 (c, 0.3 in MeOH).**N-De-Me: Magnococline.** Norgorchacoine [34955-66-3]C₁₈H₂₁NO₃ 299.369Alkaloid from *Magnolia coco* and *Monodora junodii*.**N-Me: Petaline**


[7354-32-7]

C₂₀H₂₆NO₃⁺ 328.43Quaternary alkaloid from *Leontice leontopetalum*. CNS Depressant, shows antiacetylcholine activity. Mp 140-143° (as chloride). [α]_D²⁰ +11.3.**Me ether: Bonzakaline**


[221000-68-6]

C₂₀H₂₅NO₃ 327.422Alkaloid from *Bongardia chrysogonum*. Amorph. light green solid. [α]_D²² -40 (c, 0.3 in MeOH). λ_{\max} 204 (log ϵ 4.26); 227 (log ϵ 3.85); 278 (log ϵ 3.29); 414 (log ϵ 2.05) (MeOH).**(±)-form** [28090-91-7]**N-Me:** [5890-48-2]Synthetic. Mp 134-138° (Me₂CO solvate)(as iodide).McShefferty, J. et al., *J. Pharm. Pharmacol.*, 1956, **8**, 1117 (isol, *Petaline*)Cymerman-Craig, J. et al., *Tetrahedron*, 1966, **22**, 1335 (uv, ord, config, *Petaline*)McCorkindale, N.J. et al., *Tetrahedron*, 1969, **25**, 5475 (synth)Grethe, G. et al., *Helv. Chim. Acta*, 1970, **53**, 874 (synth)Yang, T.-H. et al., *J. Chin. Chem. Soc. (Taipei)*, 1971, **18**, 91-93 (*Magnococline*)Uff, B.C. et al., *J.C.S. Perkin 1*, 1972, 479 (synth)Irgashev, T. et al., *Khim. Prir. Soedin.*, 1977, **13**, 127; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 118 (uv, ir, pmr, ms, struct)Atta-ur-Rahman, et al., *Nat. Prod. Lett.*, 1998, **12**, 161-173 (*Bonzakaline*)Nishiyama, Y. et al., *Nat. Med. (Tokyo)*, 2000, **54**, 338-341 (*Norgorchacoine*)**Goshuyamide II****G-144**3-[2-(1H-Indol-3-yl)ethyl]-1-methyl-2,4-(1H,3H)-quinazolinedione, 9CI. *Goshuyamide II* [95274-42-3]C₁₉H₁₇N₃O₂ 319.362Alkaloid from the fruits of *Evodia rutaecarpa* (Rutaceae). Prisms (EtOH). Mp 216-218°.Bergman, J. et al., *J.O.C.*, 1985, **50**, 1246-1255 (synth, ir)Shoji, N. et al., *J. Nat. Prod.*, 1989, **52**, 1160-1162 (isol, ir, pmr, cmr)**Gossyrubilone****G-145**


6,7-Dihydroxy-2-methyl-5-[[3-methylbutyl]imino]methyl]-8-(1-methylethyl)-1,4-naphthalenedione. 6,7-Dihydroxy-5-(isobutyliminomethyl)-8-isopropyl-2-methyl-1,4-naphthoquinone [69734-70-9]

C₂₀H₂₅NO₄ 343.422Isol. from terminal leaves of *Gossypium hirsutum* (cotton) (Malvaceae). Red cryst. Mp 140-142°.Bell, A.A. et al., *Phytochemistry*, 1978, **17**, 1297 (isol, uv, ir, pmr, ms)**Gostatin****G-146**

5-Amino-2-carboxy-1,4,5,6-tetrahydro-4-oxo-3-pyridineacetic acid, 9CI [78416-84-9]

C₈H₁₀N₂O₅ 214.177Isol. from *Streptomyces sumanensis*. Phytotoxin. Aspartate aminotransferase inhibitor. Cryst. + 1H₂O. Sol. H₂O, DMSO; fairly sol. MeOH; poorly sol. Me₂CO, hexane. [α]_D¹⁶ -117 (c, 1 in 2M NH₄OH). No def. Mp. λ_{\max} 331 (ϵ 11000) (H₂O) (Berdy).Nishino, T. et al., *Agric. Biol. Chem.*, 1983, **47**, 1531; 1961 (isol, props)**Goudotianine****G-147**


5,6-Dihydro-1,2-dimethoxy-6,7-dimethyl-4H-dibenzo[de,g]quinoline-3,9-diol, 9CI. 3,9-Dihydroxy-1,2-dimethoxy-7-methyldehydroaporphine [121985-88-4]

C₂₀H₂₁NO₄ 339.39Revised struct. (1,3-dimethoxy struct. originally assigned). Alkaloid from the leaves and bark of *Guatteria goudotiana* (Annonaceae). Cryst. (CHCl₃) or

amorph. Mp 186-188° (synthetic).

Castedo, L. et al., *J. Het. Chem.*, 1988, **25**, 1561 (synth, ir, uv, pmr, ms, struct)Guinaudeau, H. et al., *J. Nat. Prod.*, 1988, **51**, 389; 438 (uv, pmr, ms)Castedo, L. et al., *Phytochemistry*, 1991, **30**, 2781 (isol)**Gouregine****G-148**


4,5-Dimethoxy-12,12-dimethyl-12H-[1]benzoxepino[2,3,4-ij]isoquinoline-6,10-diol, 9CI [85769-39-7]

C₂₀H₁₉NO₅ 353.374Alkaloid from *Guatteria ouregou* (Annonaceae). Cryst. (MeOH). Mp 112-114°. First member of a new class of curarine-related alkaloids, the α -gem-dimethyltetrahydrocurarines.**Di-Ac:**C₂₄H₂₃NO₇ 437.448

Cryst. (MeOH). Mp 129-131°.

Leboeuf, M. et al., *Tetrahedron*, 1982, **38**, 2889 (pmr, cmr, ms, cryst struct)**Gracicleistanthoside****G-149**


[912851-04-8]



Relative Configuration

C₁₄H₁₉NO₇ 313.307Alkaloid from the roots of *Cleistanthus gracilis*. Viscous brown gum. [α]_D²⁵ -35.7 (c, 0.7 in MeOH).Pinho, P.M. et al., *Phytochemistry*, 2006, **67**, 1789-1792 (isol, pmr, cmr)**Gracilamine****G-150**

[879561-55-4]




Relative Configuration

C₂₅H₃₄N₂O₅ 442.554Alkaloid from *Galanthus gracilis*. Amorph. solid. [α]_D +21.8 (c, 0.13 in MeOH). λ_{\max} 234 (sh) (log ϵ 4.01); 268 (log ϵ 3.37); 295 (log ϵ 3.71) (MeOH).Unver, N. et al., *Turk. J. Chem.*, 2005, **29**, 547-553 (isol, cd, pmr, cmr)

Graciline†

[216444-04-1]

C₁₇H₁₇NO₃ 283.326Alkaloid from *Galanthus gracilis*.Amorph. solid. [α]_D²⁰ +521 (c, 0.17 in MeOH). λ_{max} 248 (sh) (log ε 3.89); 290 (log ε 3.76) (MeOH).

3,4-Dihydro, 3R-hydroxy: 3-Epi-3,4-dihydro-3-hydroxygraciline

C₁₇H₁₉NO₄ 301.341Alkaloid from *Galanthus gracilis*.Amorph. solid. [α]_D²⁰ +110.6 (c, 0.15 in MeOH). λ_{max} 215 (log ε 3.81); 233 (sh) (log ε 3.56); 292 (log ε 3.62) (MeOH).

3,4-Dihydro, 3S-hydroxy: 3,4-Dihydro-3-hydroxygraciline

[216444-08-5]


C₁₇H₁₉NO₄ 301.341Alkaloid from *Galanthus gracilis*.Amorph. solid. [α]_D²⁰ +138 (c, 0.11 in MeOH). λ_{max} 230 (sh) (log ε 4.08); 290 (log ε 3.71) (MeOH).

11S-Acetoxy: 11-Acetoxygraciline

[216444-06-3]

C₁₉H₁₉NO₅ 341.363Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. [α]_D²⁰ +198 (c, 0.17 in MeOH). λ_{max} 252 (sh) (log ε 3.95); 288 (log ε 3.69) (MeOH).Noyan, S. *et al.*, *Heterocycles*, 1998, **48**, 1777-1791 (*isol, uv, cd, ir, pmr, cmr, ms*)Unver, N. *et al.*, *Heterocycles*, 2001, **55**, 641-652 (3-Epidihydroxygraciline)

Graciline†

1,19-Epoxy-20-ethyl-4-methylaconitanine-7,8,14-triol, 9CI
[90475-66-4]C₂₁H₃₁NO₄ 361.48Alkaloid from *Delphinium gracile* (Ranunculaceae). Mp 98-100°.

8-Cinnamoyl: 8-O-Cinnamoylgraciline

[151563-64-3]

C₃₀H₃₇NO₅ 491.626Alkaloid from above-ground parts of *Delphinium cossonianum* (Ranunculaceae). Amorph. [α]_D²⁰ +10.4 (c, 0.41 in CHCl₃).

7-Deoxy: Dehydrocardiopetaline

[151563-63-2]

C₂₁H₃₁NO₃ 345.481Alkaloid from above-ground parts of *Delphinium cossonianum* (Ranunculaceae). Amorph. [α]_D²⁰ +17.8 (c, 0.80 in CHCl₃).

G-151

16β-Methoxy: Pentagydine

[86630-38-8]

C₂₂H₃₃NO₅ 391.506Minor alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 130-131°.

16β-Methoxy, 6β-hydroxy, 7,8-methylene, 14-Me ether: Laxicyminine

[1008106-60-2]

C₂₄H₃₅NO₆ 433.544Alkaloid from *Delphinium laxicymosum* var. *pilostachyum*. Needles. [α]_D²⁰ -1 (c, 0.65 in CHCl₃).

16β-Methoxy, 5,6β-dihydroxy, 7,8-methylene, 14-Me ether: Laxicymine

[1008106-58-8]

C₂₄H₃₅NO₇ 449.543Alkaloid from *Delphinium laxicymosum* var. *pilostachyum*. Cryst. Mp 192-193°. [α]_D²⁰ +4.7 (c, 0.5 in CHCl₃).

16β-Methoxy, 7-deoxy: Karakanine

[50656-90-1]

C₂₂H₃₃NO₄ 375.507Alkaloid from aerial parts of *Aconitum karacolicum* (Ranunculaceae). Cryst. (Me₂CO). Mp 193-195°.16β-Methoxy, 6β-acetoxy, 7-deoxy, O⁸-Me: Dehydrobicoloridine

[167868-58-8]

C₂₅H₃₇NO₆ 447.57Alkaloid from epigeal parts of *Delphinium peregrinum* var. *elongatum* (Ranunculaceae). Amorph. [α]_D²⁰ +29.1 (c, 0.15 in CHCl₃).

6α,16β-Dimethoxy, 7-deoxy: Pentagydine

[84306-91-2]

C₂₃H₃₅NO₅ 405.533Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 198-201°. [α]_D²⁰ +72 (c, 0.12 in EtOH).

6β,16β-Dimethoxy: Gadesine

[70420-60-9]

C₂₃H₃₅NO₆ 421.533Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 174-177°. [α]_D²⁰ +76 (c, 0.27 in EtOH).

6β,16β-Dimethoxy, 14-Ac: 14-Acetylgadesine

[70420-61-0]

C₂₅H₃₇NO₇ 463.57Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Resin.

6β,16β-Dimethoxy, 14-benzoyl: 14-Benzoylgadesine

[103976-33-6]

C₃₀H₃₉NO₇ 525.641Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin.

6β,16β-Dimethoxy, 10β-hydroxy, 14-benzoyl: Gadeline

[103956-40-7]

C₃₀H₃₉NO₈ 541.64Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Resin.

6β,16β-Dimethoxy, 18-hydroxy: 1-O,19-Didehydrotakaosamine

C₂₃H₃₅NO₇ 437.532Alkaloid from *Consolida orientalis*. Amorph. solid. [α]_D²⁵ +55.2 (c, 0.65 in CHCl₃).6β,16β-Dimethoxy, 18-hydroxy, O¹⁴-Me: 18-Hydroxy-14-O-methylgadesine

[81362-18-7]

C₂₄H₃₇NO₇ 451.559Minor alkaloid from *Consolida orientalis* (Ranunculaceae). Mp 110-114°. [α]_D²⁵ +50 (c, 0.8 in CHCl₃).

16β,18-Dimethoxy: Nevadensine

[100447-59-4]

C₂₃H₃₅NO₆ 421.533Minor alkaloid from the aerial parts of *Aconitum nevadense* (Ranunculaceae). Resin.

16β,18-Dimethoxy, 7-deoxy: Nevadenine

[100458-94-4]

C₂₃H₃₅NO₅ 405.533Minor alkaloid from the aerial parts of *Aconitum nevadense* (Ranunculaceae). Resin.

6β,16β,18-Trimethoxy: 18-Methoxygadesine

[85649-35-0]

C₂₄H₃₇NO₇ 451.559Minor alkaloid from *Consolida orientalis* (Ranunculaceae). Mp 180-184°.6β,16β,18-Trimethoxy, O⁸-Me: DehydrodeltatineC₂₅H₃₉NO₇ 465.586Alkaloid from the aerial parts of *Consolida orientalis*. Amorph. solid. [α]_D²⁵ +20 (c, 0.1 in CHCl₃).6β,16β,18-Trimethoxy, O¹⁴-Me: Dehydrodeldelsoline

[58111-41-4]

C₂₅H₃₉NO₇ 465.586Alkaloid from above-ground parts of *Aconitum vulparia* ssp. *neapolitanum* (Ranunculaceae). Cryst. (EtOAc/hexane). Mp 200-203°. [α]_D²⁰ +50.5 (c, 0.77 in CHCl₃).

6β,16β,18-Trimethoxy, 7,8-methylene ether: Campylocine

[960201-79-0]

C₂₅H₃₇NO₇ 463.57Alkaloid from *Delphinium campylocentrum*.

6β,16β,18-Trimethoxy, 14-ketone: Yesoensine

[131652-37-4]

C₂₄H₃₅NO₇ 449.543Alkaloid from *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Amorph. powder. [α]_D²¹ +39.4 (c, 1.67 in CHCl₃).

6β,16β,18-Trimethoxy, 7-deoxy: Delstaphininine

[111537-46-3]

C₂₄H₃₇NO₆ 435.559Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Amorph. [α]_D²⁸ +47.7 (c, 1.1 in CHCl₃).

6β,16β,18-Trimethoxy, 7-deoxy, di-Ac: Delstaphidinine


[112515-34-1]

C₂₈H₄₁NO₈ 519.634Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Mp 192.5-194.5°. [α]_D²⁵ +27.4 (c, 0.27 in CHCl₃).González, A.G. *et al.*, *Tet. Lett.*, 1979, 79; 1981, **22**, 4843; 1983, **24**, 959 (*Gadesine, 18-Hydroxy-14-O-methylgadesine, Pentagydine, ir, pmr, ms, cryst struct*)



- González, A.G. *et al.*, *Phytochemistry*, 1982, **21**, 1781 (*Pentagynine*)
- González, A.G. *et al.*, *Heterocycles*, 1983, **20**, 409; 1984, **22**, 667; 1985, **23**, 2979; 1986, **24**, 1513 (*18-Methoxygadesine*, *14-Benzoylgadesine*, *Gadeline*, *Graciline*, *Nevadensine*, *Nevadenine*, *14-Acetylgadesine*)
- Ross, S.A. *et al.*, *Heterocycles*, 1987, **26**, 2895 (*Delstaphidine*)
- Pelletier, S.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 381 (*Delstaphinine*)
- Wada, K. *et al.*, *Heterocycles*, 1990, **31**, 1081 (*Yesoensine*)
- Sultankhodzhaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 62; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 51 (*Karakanine*)
- De la Fuente, G. *et al.*, *Phytochemistry*, 1993, **34**, 553; 1994, **37**, 271; 1995, **39**, 1459 (*Dehydrocardiopetaline*, *8-Cinnamoylgraciline*, *Dehydrodelsoline*, *Dehydrobicoloridine*)
- Alva, A. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 530-534 (*Dehydrodeltatine*, *18-Hydroxy-14-O-methylgadesine*)
- Alva, A. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 2110-2119 (*Didehydroatakaosamine*)
- Tang, P. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 700-703 (*Laxicymine*, *Laxicyminine*)
- Yan, L.P. *et al.*, *Youji Huaxue*, 2007, **27**, 976-980; *CA*, **27**, 74053r (*Campylocine*)

Grahamine†**G-153**


[125685-27-0]

C₄₈H₆₁N₃O₁₂ 872.023Alkaloid from *Schizanthus grahamii* (Solanales). Oil. $[\alpha]_D^{20} +6.21$ (CHCl₃).Hartmann, R. *et al.*, *Angew. Chem., Int. Ed.*, 1990, **29**, 385 (*isol, pmr, cmr, struct*)**Graminifoline****G-154***Gramminifoline*C₁₈H₂₃NO₅ 333.383Pyrrolizidine alkaloid. Struct. unknown. Minor alkaloid from *Senecio graminifolius* (Asteraceae). Mp 236°.De Waal, H.L. *et al.*, *CA*, 1941, **35**, 5194⁹; 1942, **36**, 6308⁹**Gramodendrine****G-155***1-Acetyl-1,2,3,4-tetrahydro-5-[1-(1H-indol-3-ylmethyl)-2-piperidinyl]pyridine*, 9CI


[83905-67-3]

C₂₁H₂₇N₃O 337.464Alkaloid from the above-ground parts of *Lupinus arbustus* ssp. *calcaratus* (Fabaceae).Keller, W.J. *et al.*, *Phytochemistry*, 1982, **21**, 1415 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)**Grandiamide A****G-156**C₃₈H₄₄N₂O₉ 672.774Constit. of the leaves of *Aglaia grandis*. Amorph. powder. $[\alpha]_D^{25} -108.8$ (c, 0.25 in CHCl₃).Inada, A. *et al.*, *Phytochemistry*, 2000, **53**, 1091-1095**Grandifloritine****G-157**


[146471-88-7]

C₃₅H₄₂N₂O₁₀ 650.724Alkaloid from aerial parts of *Delphinium grandiflorum* (Ranunculaceae). $[\alpha]_D^{30} +38$ (c, 0.07 in CHCl₃).Li, C. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1992, **34**, 466-469; *CA*, **118**, 143418h (*isol, pmr, cmr, ms*)**Grandifoline†****G-158**

[34426-04-5]

C₃₆H₅₃NO₁₂ 691.814Alkaloid from *Malaxis grandifolia* (Orchidaceae). Amorph. $[\alpha]_D^{23} -7$ (c, 0.69 in EtOH). Hydrol. → Laburnine in

H-629.

Lindström, B. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 1900 (*isol, struct*)**Grandirubrine****G-159***10-Hydroxy-4,5,6-trimethoxy-9H-azuleno[1,2,3-ij]isoquinolin-9-one*, 9CI [74631-22-4]C₁₉H₁₅NO₅ 337.331Rearranged troponoid aporphine. CAS numbering shown. Alkaloid from *Abuta grandifolia* and *Cissampelos pareira* (Menispermaceae). Reddish-brown needles (CHCl₃/EtOH). Mp 201-203°. λ_{max} 232 (ε 91200); 254 (ε 61700); 274 (sh) (ε 45700); 296 (ε 38000); 312 (sh) (ε 28800); 343 (sh) (ε 32400); 363 (ε 52500); 384 (ε 25700); 400 (ε 15500); 480 (ε 7940) (95% EtOH) (Derep). λ_{max} 218 (ε 14600); 250 (ε 16600); 262 (ε 15700); 264 (ε 15500); 298 (ε 9800); 300 (ε 9700); 363 (ε 16900); 570 (ε 3400) (EtOH) (Berdy).*O⁹-Me: 4,5,6,9-Tetramethoxy-10H-azuleno[1,2,3-ij]isoquinolin-10-one. Imerubrine*

[58189-33-6]

C₂₀H₁₇NO₅ 351.358Alkaloid from the stems of *Abuta imene* and *Abuta rufescens* (Menispermaceae). Orange-red needles (MeOH/Et₂O). Mp 183-185°.*O¹⁰-Me: Isoimerubrine*

[74684-11-0]

C₂₀H₁₇NO₅ 351.358Alkaloid from *Cissampelos pareira* (Menispermaceae). Reddish-brown needles. Mp 183-185°. λ_{max} 218 (ε 9200); 257 (ε 11200); 304 (ε 3500); 362 (ε 17700); 414 (ε 3400); 480 (ε 1200) (EtOH) (Berdy).*4-Demethoxy: Pareirubrine B*

[152845-78-8]

C₁₈H₁₃NO₄ 307.305Alkaloid from *Cissampelos pareira* (Menispermaceae). Exhibits antileukaemic activity. Reddish-brown needles. Mp 290° dec. λ_{max} 274 (ε 25200); 294 (ε 21000); 364 (ε 20200); 420 (ε 3800); 472 (ε 7700) (EtOH) (Derep). λ_{max} 252 (ε 22900); 300 (ε 13200); 364 (ε 24600); 394 (ε 9300); 416 (ε 6900); 520 (ε 1500) (EtOH) (Berdy).*4-Demethoxy, 10-deoxy: Pareitropone*C₁₈H₁₃NO₃ 291.306Alkaloid from roots of *Cissampelos pareira* (Menispermaceae). Shows potent cytotoxic activity.*11-Methoxy: Pareirubrine A*

[147044-68-6]


C₂₀H₁₇NO₆ 367.357

Alkaloid from *Cissampelos pareira* (Menispermaceae). Exhibits antileukaemic activity. Reddish-brown needles (MeOH). Mp 168-170°. Exists as the 10-oxo tautomer in the cryst. state but as a tautomeric mixt. in soln. λ_{\max} 274 (ϵ 25200); 294 (ϵ 21000); 364 (ϵ 20200); 420 (ϵ 3800); 472 (ϵ 7700) (EtOH) (Derep). λ_{\max} 217 (ϵ 2900); 274 (ϵ 25200); 294 (ϵ 21000); 364 (ϵ 20200); 420 (ϵ 3800); 472 (ϵ 7700) (EtOH) (Berdy).

- Cava, M.P. *et al.*, *Tetrahedron*, 1975, **31**, 1667 (*Imerubrine*, *isol*, *uv*, *ir*, *pmr*)
 Silverton, J.V. *et al.*, *J.A.C.S.*, 1977, **99**, 6708 (*Imerubrine*, *cryst struct*)
 Menachery, M.D. *et al.*, *Heterocycles*, 1980, **14**, 943 (*Grandirubrine*)
 Morita, H. *et al.*, *Chem. Lett.*, 1993, 339 (*Pareirubrine A*, *isol*, *uv*, *pmr*, *cmr*, *cryst struct*)
 Morita, H. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1418; 1478 (*Isoimerubrine*, *Pareirubrine B*)
 Banwell, M.G. *et al.*, *Heterocycles*, 1994, **39**, 205 (*synth*)
 Morita, H. *et al.*, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 597 (*Pareitropone*)
 Boger, D.L. *et al.*, *J.A.C.S.*, 1995, **117**, 12452 (*synth*)
 Lee, J.C. *et al.*, *J.A.C.S.*, 2001, **123**, 3243-3246 (*synth*)
 Feldman, K.S. *et al.*, *J.O.C.*, 2002, **67**, 8528-8537 (*Pareitropone*, *synth*)

Grandisine A


G-160



- $C_{16}H_{23}NO_3$ 277.363
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum. $[\alpha]_D^{23} +38.6$ (c, 0.1 in CH_2Cl_2). λ_{\max} 270 (log ϵ 2.1) (MeOH).
 Carroll, A. *et al.*, *J.O.C.*, 2005, **70**, 1889-1892 (*isol*, *pmr*, *cmr*)
 Maloney, D.J. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 7789-7792 (*synth*)
 Maloney, D.J. *et al.*, *Heterocycles*, 2007, **72**, 167-174 (*synth*)

Grandisine B


G-161



- $C_{16}H_{22}N_2O$ 258.363
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum. $[\alpha]_D^{23} +11$ (c, 0.1 in CH_2Cl_2). λ_{\max} 275 (log ϵ 2.9) (MeOH).
 Carroll, A. *et al.*, *J.O.C.*, 2005, **70**, 1889-1892 (*isol*, *pmr*, *cmr*)

Grandisine C

[911682-66-1]



Absolute Configuration


- $C_{16}H_{23}NO_3$ 277.363
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum. $[\alpha]_D^{22} -59.2$ (c, 0.15 in MeOH). λ_{\max} 204 (log ϵ 3.39); 273 (log ϵ 3.56) (MeOH).

Stereoisomer: Rudrakine

- [72361-62-7]
 $C_{16}H_{23}NO_3$ 277.363
 Alkaloid from the leaves of *Elaeocarpus ganitrus*. Needles (MeOH). Mp 159-160°. Stereochem. not determined. λ_{\max} 272 (ϵ 7300) (MeOH).
 Ray, A.B. *et al.*, *Phytochemistry*, 1979, **18**, 700-701 (*Rudrakine*)
 Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*Grandisine C*)

Grandisine D

[911682-67-2]




Absolute Configuration

- $C_{16}H_{21}NO_2$ 259.347
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum (as trifluoroacetate salt). $[\alpha]_D^{23} +34.6$ (c, 0.09 in MeOH) (trifluoroacetate). λ_{\max} 222 (log ϵ 3.28); 267 (log ϵ 2.8); 325 (log ϵ 2.43) (MeOH) (trifluoroacetate).
 Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*isol*, *pmr*, *cmr*, *ms*)

Grandisine E

[911682-68-3]




- $C_{16}H_{23}NO_3$ 277.363
 Alkaloid from the leaves of *Elaeocarpus grandis*. Gum. λ_{\max} 272 (log ϵ 1.14) (MeOH).

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*isol*, *pmr*, *cmr*, *ms*)

Grandisine F

[911682-69-4]



Absolute Configuration

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*isol*, *pmr*, *cmr*, *ms*)


G-162

$C_{16}H_{24}N_2O_2$ 276.378
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum. $[\alpha]_D^{23} -22.4$ (c, 0.09 in MeOH). λ_{\max} 202 (log ϵ 3.2); 274 (log ϵ 3.5) (MeOH).

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*isol*, *pmr*, *cmr*, *ms*)

Grandisine G

[911682-70-7]




G-166

$C_{17}H_{26}N_2O_2$ 290.405
 Alkaloid from the leaves of *Elaeocarpus grandis*. Yellow gum (as trifluoroacetate salt). λ_{\max} 209 (log ϵ 3.39); 271 (log ϵ 3.08); 308 (log ϵ 3) (MeOH) (trifluoroacetate).

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1295-1299 (*isol*, *pmr*, *cmr*)

Grantaline

[83482-61-5]




G-167

$C_{18}H_{25}NO_6$ 351.399
 Alkaloid from *Crotalaria virgulata* subsp. *grantiana* and from the seeds of *Crotalaria globifera* (Fabaceae). Cryst. (Me₂CO aq.), prisms (EtOH). Mp 219.5-220°. $[\alpha]_D^{18} +33$ (c, 0.065 in $CHCl_3$). $[\alpha]_D^{20} +100.9$ (c, 0.8 in EtOH).

- Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*isol*, *cmr*, *struct*)
 Mackay, M.F. *et al.*, *Acta Cryst. C*, 1983, **39**, 1227 (*cryst struct*)
 Brown, K. *et al.*, *Phytochemistry*, 1984, **23**, 457 (*isol*, *ir*, *ms*, *pmr*)
 Smith, L.W. *et al.*, *Phytochemistry*, 1984, **23**, 473 (*isol*, *ir*, *pmr*, *ms*)

Grantianine

[633-10-3]



G-168

C₁₈H₂₃NO₇ 365.382

Alkaloid from *Crotalaria virgulata* ssp. *grantiana* (*Crotalaria grantiana*) and *Crotalaria globifera* (Fabaceae). Mp 223-224° (204-205°, 209-209.5°). [α]_D²⁷ +50.6 (+44.4) (CHCl₃).

Adams, R. *et al.*, *J.A.C.S.*, 1956, **78**, 4458 (struct)


Brown, K. *et al.*, *Phytochemistry*, 1984, **23**, 457 (isol, pmr, ms, cmr)

Smith, L.W. *et al.*, *Phytochemistry*, 1984, **23**, 473 (isol, pmr)

Stoeckli-Evans, H. *et al.*, *Acta Cryst. C*, 1984, **40**, 1445 (cryst struct)

Granulatimide **G-169**

1*H*-Imidazo[4,5-*a*]pyrrolo[3,4-*c*]carbazole-4,6-(5*H*,11*H*)-dione, 9*CI* [219828-99-6]

C₁₅H₈N₄O₂ 276.254

Alkaloid from the ascidian *Didemnum granulatium*. G2 specific cell cycle checkpoint inhibitor. Yellow solid.

6-Bromo-6-Bromogranulatimide [330846-92-9]

C₁₅H₇BrN₄O₂ 355.15

Alkaloid from *Didemnum granulatium*. Amorph. yellow solid. λ_{\max} 236 ; 282 ; 308 ; 386 (MeOH).


Berlinck, R.G.S. *et al.*, *J.O.C.*, 1998, **63**, 9850-9856 (isol, synth, pmr, cmr, activity)

Britton, R. *et al.*, *J. Nat. Prod.*, 2001, **64**, 254-255 (6-Bromogranulatimide, isol, pmr, cmr)

Yoshida, T. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 872-876; 2003, **51**, 209-214 (synth)

Grassypeptolide

[1010433-25-6]



Absolute Configuration


C₅₆H₇₉N₉O₁₀S₂ 1102.426

Isol. from *Lynghya confervoides*. Cytotoxic. Amorph. solid. [α]_D²⁰ +76 (c, 0.1 in CH₂Cl₂). λ_{\max} 230 (log ϵ 2.2); 260 (log ϵ 1.95); 330 (log ϵ 1.37) (CH₂Cl₂).

Kwan, J.C. *et al.*, *Org. Lett.*, 2008, **10**, 789-792 (isol, pmr, cmr, ms)

Graveoline**G-171**

2-(1,3-Benzodioxol-5-yl)-1-methyl-4(1*H*)-quinolinone, 9*CI*. 1-Methyl-2-(3,4-methylenedioxyphenyl)-4(1*H*)-quinolinone. Rutamine. Foliosine [485-61-0]

C₁₇H₁₃NO₃ 279.295

Alkaloid from *Ruta graveolens* (rue), *Ruta bracteosa* and *Haplophyllum* spp. (Rutaceae). Needles (EtOH). Mp 186-187° Mp 205-205.5° (dimorph.). Rutamine was the metastable lower-melting modification.

N-De-Me: 2-(3,4-Methylenedioxyphenyl)-4(1*H*)-quinolinone. *Norgraveoline* [74054-38-9]

Alkaloid from the aerial parts of *Haplophyllum dubium* (Rutaceae). Mp 288-290° (246-248°).

5'-Hydroxy: 2-(3-Hydroxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone. 3'-Hydroxygraveoline [107316-96-1]

C₁₇H₁₃NO₄ 295.294

Alkaloid from the aerial parts of *Ruta chalepensis* (Rutaceae). Amorph. Incorrect name in CA.

5'-Methoxy: 2-(3-Methoxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone. 3'-Methoxygraveoline [173180-39-7]

C₁₈H₁₅NO₄ 309.321

Alkaloid from trunk bark of *Esenbeckia almawillia* (Rutaceae). Amorph. solid (EtOAc). Mp 193-194°. Numbering systems vary.

5',8-Dimethoxy: 8-Methoxy-2-(3-methoxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone. 3',8-Dimethoxygraveoline [173180-40-0]

C₁₉H₁₇NO₅ 339.347

Alkaloid from trunk bark of *Esenbeckia almawillia* (Rutaceae). Amorph. solid (MeOH). Mp 201-203°. Numbering systems vary.

Arthur, H.R. *et al.*, *Aust. J. Chem.*, 1960, **13**, 510-513 (isol, uv, ir, pmr, struct)

Arthur, H.R. *et al.*, *J.C.S.*, 1961, 4360-4361 (synth)

Schneider, G. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1967, **300**, 953-954 (Rutamine)

Reisch, J. *et al.*, *Naturwissenschaften*, 1967, **54**, 200 (synth)

Razakova, D.M. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 810-812; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 716 (Norgraveoline)

Kasahara, A. *et al.*, *Chem. Ind. (London)*, 1981, 121 (synth, Norgraveoline)


Ulubelen, A. *et al.*, *Phytochemistry*, 1986, **25**, 2692-2693 (3'-Hydroxygraveoline)

Koyama, J. *et al.*, *Chem. Express*, 1992, **7**, 321-324 (synth)

Oliviera, F.M. *et al.*, *Phytochemistry*, 1996, **41**, 647-649 (3'-Methoxygraveoline, 3',8-Dimethoxygraveoline)

Annunziata, R. *et al.*, *Synth. Commun.*, 1996, **26**, 495-501 (synth)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1038-1039 (synth, ir, pmr)

3-Greenwayodendrinol**G-172**3 α -formC₂₃H₃₁NO 337.5043 α -form [85027-87-8]

Alkaloid from the stem bark of *Greenwayodendron suaveolens* (*Polyalthia suaveolens*). Gum.

3 β -form*Polyavolensinol*

[76525-23-0]

Alkaloid from the stem bark of *Greenwayodendron suaveolens* (*Polyalthia suaveolens*). Long needles (petrol/CHCl₃). Mp 167-169°. [α]_D²⁷ +11 (c, 0.56 in CHCl₃) (+6.2).

Ac: *Polyavolensin*

[76525-22-9]

C₂₅H₃₃NO₂ 379.541

Alkaloid from the stem bark of *Greenwayodendron suaveolens* (*Polyalthia suaveolens*). Plates (EtOAc/petrol). Mp 211-214°. [α]_D²⁵ -1.2 (c, 0.5 in CHCl₃) (-3.8).

3-Ketone: *Polyavolensinone*. 3-*Greenwayodendrinone*

[76525-24-1]

C₂₃H₂₉NO 335.488

Alkaloid from the stem bark of *Greenwayodendron suaveolens* (Annonaceae). Needles (EtOAc/petrol). Mp 191-194°. [α]_D²⁵ +46.8 (c, 0.64 in CHCl₃).


Okorie, D.A. *et al.*, *Tetrahedron*, 1980, **36**, 2005 (*Polyavolensin*, *Polyavolensinone*)

Hasan, C.M. *et al.*, *J.C.S. Perkin 1*, 1982, 2807-2812 (*Greenwayodendrinols*, isol, uv, ms, ir, pmr, cmr, cryst struct)

Falshaw, C.P. *et al.*, *Tetrahedron*, 1982, **38**, 2311 (*Polyavolensin*, cryst struct)

Griffinine**G-173**

5-Hydroxy-1,2-dimethoxy-7-methyl-4*H*-dibenzo[de,g]quinolin-4-one


C₁₉H₁₅NO₄ 321.332

Apparently a separately isolable tautomer of Griffithdione, G-175. Alkaloid from the stems of *Goniothalamus griffithii*. Orange needles or powder. Mp >250°. λ_{\max} 215 (log ϵ 1.51); 240 (log ϵ 1.63); 305 (log ϵ 0.68); 317 (log ϵ 0.69); 449 (log ϵ 0.57) (MeOH).

Mu, Q. *et al.*, *Planta Med.*, 2003, **69**, 826-830 (isol, pmr, cmr, ms)

Griffithazanone A

Griffithazanone A
[240122-30-9]



Absolute Configuration


$C_{14}H_{11}NO_4$ 257.245

Alkaloid from the roots of *Goniothalamus griffithii*. Yellow needles (Me₂CO). Mp 208-210°. $[\alpha]_D^{25}$ +146 (c, 0.06 in CHCl₃). λ_{\max} 203 (log ϵ 4.07); 218 (log ϵ 4); 255 (log ϵ 4.29); 286 (log ϵ 3.99); 335 (log ϵ 3.36) (CHCl₃).

Zhang, Y.J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1050-1052

Griffithdione

1,2-Dimethoxy-7-methyl-4H-dibenzo[de,g]quinoline-4,5(6H)-dione
[240122-31-0]




$C_{19}H_{15}NO_4$ 321.332

Alkaloid from the roots of *Goniothalamus griffithii*. Orange needles (CHCl₃). Mp 216-218°. λ_{\max} 244 (log ϵ 4.49); 295 (log ϵ 4.02); 307 (log ϵ 4.14); 330 (log ϵ 4.13); 460 (log ϵ 3.98) (CHCl₃).

Zhang, Y.-J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1050-1052 (isol, uv, ir, pmr, cmr, ms)

Griffithine

[140367-81-3]



Relative Configuration


$C_{24}H_{28}N_4O_4$ 436.51

Alkaloid from leafy shoots of *Sophora griffithii* (Fabaceae). Cream-coloured solid. $[\alpha]_D$ +217 (c, 0.03 in MeOH).

Atta-ur-Rahman, *et al.*, *Magn. Reson. Chem.*, 1991, **29**, 1077-1083 (isol, uv, ir, pmr, cmr, ms, struct)

Grisabine

[62057-36-7]



G-177

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloids covered by this entry are enantiomeric with those covered by Berbaminine, B-98 (1R,1'S-config.) and diastereomeric with those covered by Dauricine, D-88 (R,R-config.) and Thallibrine, T-309 (S,S-config.). Alkaloid from the stems of *Abuta grisebachii* (Menispermaceae) and *Xylomalos monospora*. Mp 148-149°. $[\alpha]_D$ -60.2 (c, 0.5 in CHCl₃).

Di-Me ether: **O,O-Dimethylgrisabine**

[13565-64-5]

$C_{39}H_{46}N_2O_6$ 638.802

Alkaloid from the leaves of *Phaeanthus vietnamensis* (Annonaceae). Exhibits antibacterial activity. Amorph. solid. $[\alpha]_D^{26}$ -26 (c, 0.19 in CHCl₃).

*O*¹²-*De-Me*: **Magnoline. Grisabutine**

[6859-66-1]

$C_{36}H_{40}N_2O_6$ 596.722

Alkaloid from the stems of *Abuta grisebachii* (Menispermaceae) and leaves of *Magnolia fuscata* (Magnoliaceae). Cryst. (MeOH or EtOH). Mp 178-179° Mp 192-193°. $[\alpha]_D$ -50 (c, 0.5 in CHCl₃). Enantiomer of Berbaminine, B-98.

*O*¹²-*De-Me*, *O*⁷-*Me*: **(-)-Temuconine**

[126589-93-3]

$C_{37}H_{42}N_2O_6$ 610.749

Alkaloid from *Aristolochia elegans* (Aristolochiaceae). Amorph. $[\alpha]_D$ -99.8 (c, 2.4 in MeOH). Enantiomer of Temuconine in B-98.

*O*¹²-*De-Me*, *O*⁷, *O*⁷-*di-Me*:

[935751-39-6 (hydrochloride)]

$C_{38}H_{44}N_2O_6$ 624.775

Alkaloid from the stem bark of *Xylomalos monospora*.

Proskurina, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1938, **5**, 1357 (*Magnoline*, isol)

Proskurina, N. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1940, **10**, 707; *CA*, **35**, 2520 (*Magnoline*, struct)

Ahmed, R. *et al.*, *J.O.C.*, 1977, **42**, 2271

(*Grisabine*, *Magnoline*, isol, uv, ir, pmr, ms, struct)

El-Sebakhy, N. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1374 (*(-)-Temuconine*)


Sedmera, P. *et al.*, *Heterocycles*, 1990, **30**, 205 (*O,O-Dimethylgrisabine*)

Nganga, D. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 619-622 (*Xylomalos constiti*)

Griseoviridin

[53216-90-3]

G-178



$C_{22}H_{27}N_3O_7S$ 477.537

Modified cyclic peptide antibiotic. prod. by *Streptomyces griseus*/*Streptomyces griseoviridis*. Antibiotic active against gram-positive bacteria. Cryst. (MeOH) (polymorphic). Sol. MeOH, EtOH, Py; fairly sol. H₂O, butanol; poorly sol. EtOAc, hexane. Mp 161-163° dec., 194-200°, 230-240°. $[\alpha]_D^{25}$ -237 (c, 0.5 in MeOH). λ_{\max} 221 (ϵ 41500); 278 (sh) (ϵ 1500) (MeOH) (Derep). λ_{\max} 221 (E1%/1cm 870) (MeOH) (Berdy). λ_{\max} 220 (ϵ 44000) (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 75 mg/kg, LD₅₀ (mus, ipr) 100 mg/kg, LD₅₀ (mus, scu) 100 mg/kg. ME4200000

Hydrochloride:

Needles (Et₂O/MeOH). Dec. \geq 170°.

Di-Ac:

Cryst. (MeOH). Mp 136-140° dec. $[\alpha]_D^{27}$ -230 (c, 0.44 in MeOH).

Ames, D.E. *et al.*, *J.C.S.*, 1955, 4260 (isol, ir, uv)
Fallona, M.C. *et al.*, *Can. J. Chem.*, 1964, **42**, 371; 394 (ir, uv, nmr)

Birnbaum, G.I. *et al.*, *J.A.C.S.*, 1976, **98**, 1926 (cryst struct)

Bycroft, B.W. *et al.*, *J.C.S. Perkin 1*, 1976, 1996; 1977, 2464 (cmr, pmr, ms, ir, cryst struct, conformn)

Meyers, A.I. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 2537 (synth)

Nikaido, M.M. *et al.*, *Diss. Abstr. Int.*, **B**, 1983, 146 (synth)

Jayaraman, G. *et al.*, *Biochim. Biophys. Acta*, 1994, **1201**, 149 (pmr, struct)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GK000

Grisonomycin

Antibiotic A 10073. A 10073

Related to Ferrimycin A₁, F-37. Prod. by *Streptomyces griseus* sp. A10073. Side-phore. Antibacterial agent. Yellow powder. Sol. H₂O, phenol; poorly sol. butanol, hexane. λ_{\max} 280 (E1%/1cm 82); 425 (E1%/1cm 16.4) (H₂O) (Berdy).


Ger. Pat., 1959, 1 070 782; *CA*, **55**, 21475f

G-179

Grixazone B

SM 43. Antibiotic SM 43. Umycin B
[116511-06-9]

G-180



$C_{18}H_{15}N_3O_7S$ 417.398

Prod. by *Streptomyces* sp. DSM 3813 and by *Streptomyces griseus* under phosphate depletion. Anthelmintic. Lipoygenase inhibitor.

1'-Aldehyde: Grinoxazone A

$C_{18}H_{15}N_3O_6S$ 401.399

Prod. by *Streptomyces griseus* under phosphate depletion. λ_{max} 270 (€ 17100); 414 (€ 14800); 436 (€ 15200) (DMSO).

Eur. Pat., 1988, 260 486; *CA*, **109**, 142586p (isol)

Ohnishi, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 218-223 (isol, pmr, cmr)


Horinouchi, S. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 283-299 (rev, biosynth)

Suzuki, H. *et al.*, *J. Antibiot.*, 2007, **60**, 380-387 (biosynth)

Groenlandicine G-181

11,12-Dihydro-9-hydroxy-8-methoxybenzo[a]-1,3-benzodioxolo[4,5-g]quinolizinium, 9CI. Dehydrochelanthifoline.

Tetradehydrochelanthifoline. Alkaloid B† [38691-95-1]



$C_{19}H_{16}NO_4^{\oplus}$ 322.34

Originally considered to be $C_{21}H_{16}NO_6^{\oplus}$. Alkaloid from *Coptis groenlandica*, other *Coptis* spp., *Corydalis saxicola* and cultures of *Nandina domestica*. Orange needles (as chloride). Mp 270-275° dec.

Me ether: 11,12-Dihydro-8,9-dimethoxybenzo[a]-1,3-benzodioxolo[4,5-g]quinolizinium, 9CI. Epiberberine [6873-09-2]

$C_{20}H_{18}NO_4^{\oplus}$ 336.367

Alkaloid from *Berberis floribunda*, *Coptis chinensis*, *Coptis trifolia* and *Nandina domestica*. Orange cryst. (CHCl₃/MeOH)(as chloride). Mp 260° dec. (chloride). λ_{max} 227 ; 245 ; 268 ; 361 (EtOH).

[41451-54-1]

Chatterjee, D.R. *et al.*, *J. Indian Chem. Soc.*, 1951, **28**, 225-228 (isol, Epiberberine)

Cooper, S.F. *et al.*, *Planta Med.*, 1970, **19**, 23; 1972, **21**, 313 (isol, uv, ir, pmr)

Jewers, K. *et al.*, *J.C.S. Perkin 2*, 1972, 1393 (pmr, struct)

Suguna, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 1219 (struct)

Pai, B.R. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 607-611 (synth, Epiberberine)

Ikuta, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 189-190 (isol, Epiberberine)


Ikuta, A. *et al.*, *Phytochemistry*, 1988, **27**, 2143-2145 (isol, Epiberberine)

Mizuno, M. *et al.*, *Phytochemistry*, 1992, **31**, 717-719 (isol, Epiberberine)

Cheng, X. *et al.*, *Chem. Biodiversity*, 2008, **5**, 1335-1344 (isol, pmr, cmr)

Grossamide G-182

[80510-06-1]



$C_{36}H_{36}N_2O_8$ 624.689

Isol. from *Cannabis sativa* and *Capsicum annuum* var. *grossum* and *Hibiscus cannabinus*. Protease inhibitor. Cryst. (MeOH/CHCl₃/cyclohexane). Mp 174-175° (133-135°).

3'-O-De-Me: Demethylgrossamide

[170171-94-5]

$C_{35}H_{34}N_2O_8$ 610.662

Isol. from seeds of *Xylopia aethiopica*. Amorph. powder.

2-Epimer: Tribulusamide A

[218622-84-5]

$C_{36}H_{36}N_2O_8$ 624.689

Alkaloid from the fruit of *Tribulus terrestris*. Hepatoprotective agent. Amorph. solid. $[\alpha]_D^{25}$ -10.9 (c, 0.1 in MeOH). λ_{max} 220 (log € 4.22); 287 (log € 4.01); 340 (log € 4.07) (MeOH).

2-Epimer, 2''-oxo: Tribulusamide B

[218622-86-7]

$C_{36}H_{34}N_2O_9$ 638.673

Alkaloid from the fruit of *Tribulus terrestris*. Hepatoprotective agent. Amorph. solid. $[\alpha]_D^{25}$ -4.9 (c, 0.2 in MeOH). λ_{max} 220 (log € 4.38); 285 (log € 4.32); 320 (log € 4.18) (MeOH).

Yoshihara, T. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 2593; 1983, **47**, 217 (isol, synth, struct)

Sakakibara, I. *et al.*, *Phytochemistry*, 1991, **30**, 3013-3016 (isol)


Lajide, L. *et al.*, *Phytochemistry*, 1995, **40**, 1105-1112 (Demethylgrossamide)

Li, J.-X. *et al.*, *Planta Med.*, 1998, **64**, 628-631 (Tribulusamides)

Seca, A.M.L. *et al.*, *Phytochemistry*, 2001, **58**, 1219-1223 (isol, pmr, cmr)

Grossularine 1 G-183

[94935-97-4]



$C_{23}H_{18}N_6O$ 394.435

Struct. revised in 1989. First examples of a naturally occurring α -carboline (see also Grossularine 2, G-184). Cmr as-

signments revised in 1996. Isol. from the tunicate *Dendrodoa grossularia*. Exhibits marked cytotoxicity toward murine and human tumour cells. Amorph. yellow powder. Mp 350°. λ_{max} 202 ; 235 ; 264 (sh) ; 340 ; 360 (EtOH) (Derep).

N,N-Di-de-Me: N,N-Didemethylgrossularine 1

[175170-89-5]

$C_{21}H_{14}N_6O$ 366.381

Alkaloid from the ascidian *Polycarpa aurata*. Yellow needles (MeOH/Me₂CO). Mp 333-335°.

Moquin, C. *et al.*, *Tet. Lett.*, 1984, **25**, 5047-5048 (isol, ms)

Moquin-Patthey, C. *et al.*, *Tetrahedron*, 1989, **45**, 3445-3450 (isol, uv, ms, struct)

Abas, S.A. *et al.*, *J.O.C.*, 1996, **61**, 2709-2712 (*N,N*-Didemethylgrossularine 1)


Loukaci, A. *et al.*, *Magn. Reson. Chem.*, 1996, **34**, 143-145 (pmr, cmr)

Molina, P. *et al.*, *Tetrahedron*, 1998, **54**, 9623-9638 (synth)

Miyake, F.Y. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 3280-3282 (synth)

Grossularine 2 G-184

[102488-58-4]



$C_{21}H_{17}N_5O_2$ 371.398

Cmr spectral assignments revised in 1996. Isol. from the tunicate *Dendrodoa grossularia*. Exhibits marked cytotoxicity toward murine and human tumour cells. Cryst. (THF/MeOH). Mp 281-283°. λ_{max} 235; 255; 340; 360 (EtOH) (Derep).


Moquin-Patthey, C. *et al.*, *Tetrahedron*, 1989, **45**, 3445 (isol, pmr, cmr, ms, struct)

Loukaci, A. *et al.*, *Magn. Reson. Chem.*, 1996, **34**, 143 (pmr, cmr)

Molina, P. *et al.*, *Tetrahedron*, 1998, **54**, 9623-9638 (synth)

Guadinomic acid G-185

K01-0509B. Antibiotic K01-0509B [919521-26-9]



$C_{10}H_{18}N_4O_4$ 258.277


Prod. by *Streptomyces* sp. K01-0509. Selective inhibitor of type III secretion system. $[\alpha]_D^{25}$ +31.9 (c, 0.1 in MeOH).

Tsuchiya, S. *et al.*, *Org. Lett.*, 2006, **8**, 5577-5580 (synth, pmr, cmr, abs config)

Iwatsuki, M. *et al.*, *J. Antibiot.*, 2008, **61**, 230-236 (isol, pmr, cmr)

Guadinomine A **G-186**

[948018-04-0]



C₂₀H₃₈N₈O₈ 518.569

Prod. by *Streptomyces* sp. K01-0509.

Bacterial type III secretion system inhibitor.

1'-Deoxy: **Guadinomine B**

[948018-05-1]

C₂₀H₃₈N₈O₇ 502.57


Prod. by *Streptomyces* sp. K01-0509.

Hirose, T. *et al.*, *Chem. Eur. J.*, 2008, **14**, 8220-8238 (*Guadinomine B*, *synth*)

Iwatsuki, M. *et al.*, *J. Antibiot.*, 2008, **61**, 222-229; 230-236 (*isol*, *pmr*, *cmr*)

Guadinomine C₁ **G-187**

[1033827-71-2]



Absolute Configuration

C₂₃H₄₀N₈O₈ 556.618

Prod. by *Streptomyces* sp. K01-0509.

3'-Epimer: **Guadinomine C₂**

[1033827-72-3]

C₂₃H₄₀N₈O₈ 556.618

Prod. by *Streptomyces* sp. K01-0509.


Hirose, T. *et al.*, *Chem. Eur. J.*, 2008, **14**, 8220-8238 (*Guadinomine C₂*, *synth*, *abs config*)

Iwatsuki, M. *et al.*, *J. Antibiot.*, 2008, **61**, 222-229; 230-236 (*isol*, *pmr*, *cmr*)

Guadiscidine **G-188**

6,8-Dihydro-8,8-dimethyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-10-ol, *9CI*

[91174-10-6]



C₁₉H₁₇NO₃ 307.348

Alkaloid from the stem bark of *Guatteria discolor*. Cryst. (MeOH). Mp 269-270°.

Me ether: Guadiscine. 9-O-Methylguadiscidine

[84679-87-8]

C₂₀H₁₉NO₃ 321.375

Alkaloid from the stem bark of *Guatteria discolor*. Amorph.

4,5-Didehydro, Me ether: 4,5-Didehydroguadiscine

C₂₀H₁₇NO₃ 319.359

Alkaloid from the stems and roots of *Hornschurchia obliqua*. Brown amorph. powder.

Deoxy: Demethoxyguadiscine. Deoxyguadiscidine

C₁₉H₁₇NO₂ 291.349

Alkaloid from the stems and roots of *Hornschurchia obliqua*. Brown amorph. powder.

11-Methoxy, Me ether: Guadiscoline

[84679-86-7]

C₂₁H₂₁NO₄ 351.401

Alkaloid from the stem bark of *Guatteria discolor*. Noncryst.

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 353-362 (*isol*, *uv*, *ir*, *pmr*, *ms*)

Fechine, I.M. *et al.*, *Fitoterapia*, 2003, **74**, 29-33 (*Demethoxyguadiscine*, *4,5-Didehydroguadiscine*)

C-Guaianine **G-189**

Struct. unknown

C₂₁H₂₅N₂O⁺ 321.441

MF is tentative. May be C₂₀H₂₅N₂⁺.


Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 305° (as picrate). Blue-viol. col. with Ce(SO₄)₂. Props. similar to those of C-Curarine, C-809.

Giesbrecht, E. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1974-1982

Guaipyridine **G-190**

6,7,8,9-Tetrahydro-2,5-dimethyl-8-(1-methylethenyl)-5H-cyclohepta[b]pyridine, *9CI. Epiguaipyridine*

[41447-48-7]



Absolute configuration

C₁₅H₂₁N 215.338

Originally considered to have *trans*-config. and therefore named Epiguaipyridine. Alkaloid from the essential oil of *Pogostemon patchouly* (Lamiaceae). Liq. [α]_D^{25.5} -34.5 (c, 2.69 in C₆H₆).

5-Epimer, 1',2'-dihydro, 1'-hydroxy: Cananodine

C₁₅H₂₃NO 233.353

Alkaloid from the fruit of *Cananga odorata*. Cytotoxic. Yellow oil. [α]_D²⁵ -76.2 (c, 0.06 in CHCl₃). λ_{max} 204 (log ε 4.15); 222 (sh) (log ε 4.01); 270 (log ε 3.92) (EtOH).

Büchi, G. *et al.*, *J.A.C.S.*, 1966, **88**, 3109 (*isol*, *ir*, *pmr*)


Van der Gen, A. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1972, **91**, 1433 (*synth*, *config*, *pmr*)

Hsieh, T.-J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 616-619 (*Cananodine*)

Craig, D. *et al.*, *Eur. J. Org. Chem.*, 2006, 3558-3561 (*synth*)

Guamamide **G-191**

[657401-19-9]




C₂₅H₄₅NO₈ 487.632

Alkaloid from a *Symploca* sp. Cytotoxic. Amorph. powder. [α]_D²¹ +6 (c, 0.4 in MeOH). λ_{max} 201 (log ε 3.71) (MeOH).

Williams, P.G. *et al.*, *J. Nat. Prod.*, 2004, **67**, 49-53 (*isol*, *pmr*, *cmr*)

Guanacastepene D **G-192**

[378781-31-8]




C₂₂H₂₇NO₃ 353.46

Metab. of fungus CR115.

Brady, S.F. *et al.*, *J.A.C.S.*, 2001, **123**, 9900-9901 (*Guanacastepene D*, *cryst struct*)

Guanacastepene H **G-193**

[378781-63-6]



C₂₂H₂₉NO₅ 387.475

Metab. of fungus CR115.

Brady, S.F. *et al.*, *J.A.C.S.*, 2001, **123**, 9900-9901 (*Guanacastepene H*, *cryst struct*)

Guan-fu base C **G-194**

[1394-51-0]

C₂₂H₃₃NO₂ 343.508

Struct. unknown. Alkaloid from the Chinese herbal drug Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Mp 150°. [α]_D^{16.4} -21.2 (EtOH).

Hydrobromide: Mp 235°.

Nitrate: Mp 222°.

Di-Ac; hydrochloride: Mp 222.5-224°.

Kao, H.-C. *et al.*, *Yaoxue Xuebao*, 1966, **13**, 186-194; *CA*, **65**, 3922g (*isol*)

Guan-fu base D **G-195**

[59113-42-7]

C₂₄H₃₅NO₃ 385.545Struct. unknown. Alkaloid from the Chinese herbal drug Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae).

Nitrate: [1394-52-1]

Mp 210-211°.

Kao, H.-C. *et al.*, *Yaoxue Xuebao*, 1966, **13**, 186-194; *CA*, **65**, 3922g (isol)**Guan-fu base E** **G-196**

[59113-41-6]


C₂₉H₄₃NO₇ 517.661Struct. unknown. Alkaloid from the Chinese herbal drug Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae).

Perchlorate: [1394-53-2]

Mp 272°.

Kao, H.-C. *et al.*, *Yaoxue Xuebao*, 1966, **13**, 186-194; *CA*, **65**, 3922g (isol)**Guan-fu base K** **G-197**


[497831-34-2]



Absolute Configuration

C₂₀H₂₇NO₄ 345.438Alkaloid from the roots of *Aconitum coreanum*.Yang, C. *et al.*, *Zhongcaoyao*, 2002, **33**, 201-203; *CA*, **138**, 166611**Guan-fu base S** **G-198**

[1037700-00-7]

C₂₄H₂₉NO₅ 411.497Alkaloid from the roots of *Aconitum coreanum*. Powder. Mp 182-184°. [α]_D²⁵ +16.5 (c, 1 in CHCl₃).Yang, C.-H. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 759-765 (isol)**Guanidine, 9CI** **G-199**

Carbamidine. Iminourea. Aminoformamidine. Aminomethanamidine. Carbamamidine

[113-00-8]

HN=C(NH₂)₂CH₅N₃ 59.071Shows degenerate imino⇌amino tautomerism; only one set of monosubst. derivs. obtainable. Anion is symmetrical. Occurs widely, e.g. first isol. from vetch seedlings (*Vicia sativa*). Organic base used to form extractable ion-pairs with anionic complexes. Enhances acetylcholine release from nerve endings. Used to treat botulism and other muscle weakness states. Deliquescent cryst. mass. Sol. H₂O, EtOH, acids aq. solns. Mp 50°. p*K*_{a1} 13.59; p*K*_{a2} -11 (25°). Strong base, absorbs CO₂ from air.▶ LD₅₀ (mus, ipr) 350 mg/kg. ME7750000

Hydrochloride: [50-01-1]

Hygroscopic powder. Mp 183°.

▶ MF4300000

Carbonate salt (2:1): [593-85-1]

[100224-74-6, 3425-08-9]

Mp 197°.

▶ FG1750000

Sulfate (2:1): [594-14-9]

[1184-68-5]

Mp 290-293° dec.

Acetate salt: [34771-62-5]

Mp 229-230°.

Nitrate salt: [506-93-4]

Mp 214°.

▶ May explode during prepn. MF4350000

Phosphate salt: [1763-07-1]

[5423-22-3, 38848-02-1, 5423-23-4]

Fertiliser. Fireproofing agent for textiles.

Picrate: Mp 333°.

N-Benzenesulfonyl: Mp 212°.

N-Ac: [5699-40-1]

C₃H₇N₃O 101.108

Mp 145°.

N-Benzoyl:

C₈H₉N₃O 163.179

Cryst. (EtOH). Mp 160°.

N,N'-Di-Ac:

C₅H₉N₃O₂ 143.145

Mp 176-177°.

N,N',N''-Tri-Ac:

C₇H₁₁N₃O₃ 185.182Needles (Me₂CO). Mp 110-112°.

N-tert-Butyloxycarbonyl: [219511-71-4]

C₆H₁₃N₃O₂ 159.188

Cryst. Mp 165° dec.

N-Benzoyloxycarbonyl: [16706-54-0]

C₉H₁₁N₃O₂ 193.205

Mp 140-142°.

N-Isopropyl: [42771-39-1]

C₄H₁₁N₃ 101.151

Gel.

N,N,N',N''-Tetra-Ph: N,N,N',N''-Tetra-phenylguanidine, 9CI

[632-88-2]

C₂₅H₂₁N₃ 363.461

Cryst. (petrol). Mp 130-131°.

N,N,N',N''-Tetra-Ph, N''-benzoyl:

C₃₂H₂₅N₃O 467.569

Cryst. (EtOH). Mp 142-144°.

[5150-56-1, 19227-70-4, 19244-98-5, 10308-84-6, 52470-25-4, 25215-10-5, 5423-23-4, 10199-21-0, 50979-18-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 818A; 818C; 818B (ir)Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1329C (nmr)Weith, W. *et al.*, *Ber.*, 1874, **7**, 843 (tetraphenyl)Johnson, T.B. *et al.*, *J.A.C.S.*, 1912, **34**, 164 (tetraphenyl)Org. Synth., Coll. Vol., **1**, 1932, 302 (synth)Ger. Pat., 1935, 614 818; *CA*, **29**, 8239 (phosphate)Welcher, R.J. *et al.*, *Organic Analytical Reagents*, Van Nostrand, New York, 1947, **2**, 388 (use)Watt, G.W. *et al.*, *Ind. Eng. Chem.*, 1954, **46**, 2599 (synth)Beynon, J.H. *et al.*, *Org. Mass Spectrom.*, 1968, **1**, 169 (ms)Hart, N.K. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1679 (synth, deriv)Corral, R.A. *et al.*, *Chem. Comm.*, 1970, 556 (synth, deriv)Ghosh, P. *et al.*, *Technology*, 1972, **9**, 65-67; *CA*, **79**, 18019d (phosphate)U.S. Pat., 1974, 3 811 992; *CA*, **81**, 93341x (phosphate, use)

Ullmanns Encykl. Tech. Chem., 4. Aufl., 1976, (rev)

Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, 514 (rev)

Smith, R.L. *et al.*, *J.A.C.S.*, 1979, **101**, 191 (cmr)Fritsche-Lang, W. *et al.*, *Chem. Ber.*, 1985, **118**, 2044 (purifn)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1116Katrusiak, A. *et al.*, *Acta Cryst. C*, 1994, **50**, 1161 (cryst struct, nitrate)Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **4**, 2626-2630 (use)Zhang, Q. *et al.*, *J. Med. Chem.*, 2005, **48**, 6472-6481 (N-isopropyl)Goebel, M. *et al.*, *Chem. Comm.*, 2007, 3180-3181 (cryst struct)Schmuck, C. *et al.*, *Eur. J. Org. Chem.*, 2008, 324-329 (N-BOC, N-benzyloxycarbonyl)Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 0483Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 651Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, GKW000; GKY000; GLC000; GLA000; GLB300**Guanidinoacetic acid** **G-200**

N-(Aminoiminomethyl)glycine, 9CI. Glycocyamine. N-Guanylglycine. [(Aminoiminomethyl)amino]acetic acid.

Guanidinoacetic acid. (Carboxymethyl)-guanidine

[352-97-6]

HN=C(NH₂)NHCH₂COOHC₃H₇N₃O₂ 117.107Occurs in the sea anemone *Anthopleura japonica* and other spp. Has cardiotoxic props. Plates (H₂O). Mp 280-284° dec. p*K*_a 2.82 (25°).

Hydrochloride: [14901-20-3]

Plates (conc. HCl). Mp 200° dec.

N-Me: see Creatine, C-731

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 820D (ir)Aldrich Library of NMR Spectra, 2nd edn., 1983, **1**, 680B (nmr)Walter, W. *et al.*, *Angew. Chem.*, 1955, **67**, 275 (synth)Org. Synth., Coll. Vol., **3**, 1955, 440 (synth)

- Fasold, H. *et al.*, *Biochem. Z.*, 1961, **335**, 86 (synth)
 Cramer, F. *et al.*, *Chem. Ber.*, 1962, **95**, 1670 (synth)
 Pant, R. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1964, **335**, 272 (synth)

4-Guanidinobutanoic acid G-201

4-[(Aminoiminomethyl)amino]butanoic acid, 9CI. GGBA

[463-00-3]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}_2\text{CH}_2\text{COOH}$

$\text{C}_5\text{H}_{11}\text{N}_3\text{O}_2$ 145.161

Occurs in plants and animals, e.g. *Pinus* spp., *Lunaria annua*, sea anemone *Anthopleura japonica*, green alga *Enteromorpha intestinalis*. Found in soybean, tea and *Citrus* spp. Depressive neurotransmitter, displays similarities to 4-Aminobutanoic acid, A-715. Cryst. + 2H₂O (H₂O).

▶ Toxic to chick embryos.

Hydrochloride: [13890-14-7]

Mp 184°.

Amide: 4-Guanidinobutyramide. **Tiformin**, INN. Tyformin, BAN

[4210-97-3]

$\text{C}_5\text{H}_{12}\text{N}_4\text{O}$ 144.176

Oral hypoglycaemic agent. Log P -2.9 (calc).

Amide; *hydrochloride*: HL 523.

Augmentin†

[23256-39-5]

Cryst. (MeOH). Mp 108°.

Robin, Y. *et al.*, *Bull. Soc. Chim. Biol.*, 1953, **35**, 285 (biosynth)

Irreverre, F. *et al.*, *Nature (London)*, 1957, **80**, 704 (occur)

Makisumi, S. *et al.*, *J. Biochem. (Tokyo)*, 1961, **49**, 284 (isol)

Pant, R. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1964, **335**, 272 (synth)

Reinbothe, H. *et al.*, *Phytochemistry*, 1964, **3**, 327 (biosynth)

Tomita, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 3628 (cryst struct)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 13387

Okabe, N. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1992 (cryst struct)

N-(4-Guanidinobutyl)-2-(4-hydroxyphenyl)-2-oxoacetamide G-202

$\text{C}_{13}\text{H}_{18}\text{N}_4\text{O}_3$ 278.31

Isol. from the hydroid *Campanularia* sp. Oil. λ_{max} 209 (log ϵ 3.84); 295 (log ϵ 3.89) (MeOH).

Houssen, W.E. *et al.*, *J. Nat. Prod.*, 2005, **68**, 453-455

2-Guanidinoethanesulfonic acid, 8CI G-203

2-[(Aminoiminomethyl)amino]ethanesulfonic acid, 9CI. **Hypotaurocyamine**

[1119-54-6]

[35365-99-2]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$

$\text{C}_3\text{H}_9\text{N}_3\text{O}_2\text{S}$ 151.189

Isol. from *Agelas* spp., *Arenicola marina* and *Phascolosoma* spp. Mp 187-188°.

Phosphate: 2-[[Imino(phosphonoamino)methyl]amino]ethanesulfonic acid, 9CI. 2-(3-Phosphoguanidino)ethanesulfonic acid, 8CI. **Hypotaurocyanamine-phosphoric acid**

[4378-69-2]

$\text{C}_3\text{H}_{10}\text{N}_3\text{O}_5\text{PS}$ 231.169

Isol. from marine annelids.

Robin, Y. *et al.*, *Biochim. Biophys. Acta*, 1962, **63**, 481 (isol)

Desvages, G. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1968, **267**, 1868 (synth)

Berthou, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1972, **274**, 157 (cryst struct)

2-Guanidinoethanesulfonic acid G-204

2-[(Aminoiminomethyl)amino]ethanesulfonic acid, 9CI. N-Amidino-taurine, 8CI. **Taurocyamine**. Guanidinotaurine

[543-18-0]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}_2\text{SO}_3\text{H}$

$\text{C}_3\text{H}_9\text{N}_3\text{O}_3\text{S}$ 167.188

Isol. from urine. Antistaphylococcal, antifibrinolytic. Cryst. (EtOH/Me₂CO). Mp 226-228° Mp 264°.

N¹-Me: **Halichondriasulfonic acid**

$\text{C}_4\text{H}_{11}\text{N}_3\text{O}_3\text{S}$ 181.215

Isol. from *Halichondria rugosa*. Anti-HIV-1 agent. Needles.


Schram, E. *et al.*, *Bull. Soc. Chim. Biol.*, 1957, **39**, 561 (isol)

Fujii, A. *et al.*, *J. Med. Chem.*, 1975, **18**, 502 (synth, pharmacol)

Jin, Y. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 1129-1135 (*Halichondriasulfonic acid*)

1-(2-Guanidinoethyl)-1,2,3,4-tetrahydro-6-hydroxymethyl-β-carboline G-205

1-(2-Guanidinoethyl)-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indole-6-methanol



$\text{C}_{15}\text{H}_{21}\text{N}_5\text{O}$ 287.364

Alkaloid from the web of the spider *Nephila clavipes*.

Marques, M.R. *et al.*, *Chem. Biodiversity*, 2005, **2**, 525-534 (isol, hplc, ms)

4-Guanidino-3-hydroxybutanoic acid G-206

4-[(Aminoiminomethyl)amino]-3-hydroxybutanoic acid, 9CI

[7010-89-1]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}(\text{OH})\text{CH}_2\text{COOH}$

$\text{C}_5\text{H}_{11}\text{N}_3\text{O}_3$ 161.16

Occurs in the sea anemone *Anthopleura japonica*. Cryst. (H₂O). Mp 260° dec.

Robin, Y. *et al.*, *Bull. Soc. Chim. Biol.*, 1953, **35**, 285 (biosynth)

Makisumi, S. *et al.*, *J. Biochem. (Tokyo)*, 1961, **49**, 284 (isol)

Japan. Pat., 1963, 63 2 860; CA, **59**, 11263 (synth)

5-Guanidino-2-oxopentanoic acid G-207

5-[(Aminoiminomethyl)amino]-2-oxopentanoic acid, 9CI. α-Keto-δ-guanidinovaleric acid

[3715-10-4]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}_2\text{CH}_2\text{COCOOH}$

$\text{C}_6\text{H}_{11}\text{N}_3\text{O}_3$ 173.171

Occurs in insects and marine invertebrates. Also found in *Phlox decussata* and *Picea glauca*. Cryst. + 1H₂O (H₂O). Mp 250°. Darkens at 221°.

2,4-Dinitrophenylhydrazone: Mp 250°. Darkens at 218°.

Robin, Y. *et al.*, *Biochim. Biophys. Acta*, 1953, **11**, 403 (isol)

Brandner, G. *et al.*, *Acta Chem. Scand.*, 1964, **18**, 574 (isol)

Biellmann, J.F. *et al.*, *Bioorg. Chem.*, 1977, **6**, 89 (synth)

Cooper, A.J.L. *et al.*, *J. Biol. Chem.*, 1978, **253**, 5407 (synth)

3-Guanidinopropanoic acid G-208

N-(Aminoiminomethyl)-β-alanine, 9CI.

N-Amidino-β-alanine, 8CI. Guanidinepropionic acid

[353-09-3]

$\text{HN}=\text{C}(\text{NH}_2)\text{NHCH}_2\text{CH}_2\text{COOH}$

$\text{C}_4\text{H}_9\text{N}_3\text{O}_2$ 131.134

Occurs in the sea anemone *Anthopleura japonica*. Cryst. (EtOH aq.). Mp 209-211°.

Mourgue, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1948, 181 (synth)


Makisumi, S. *et al.*, *J. Biochem. (Tokyo)*, 1961, **49**, 284 (isol)

Steward, E.G. *et al.*, *Acta Cryst. B*, 1974, **30**, 813 (cryst struct)

3-Guanidinylpyridine G-209

3-Pyridinylguanidine, 9CI, 8CI. **Pyrazinone**

[67087-03-0]




$\text{C}_6\text{H}_8\text{N}_4$ 136.156

Isol. from the New Caledonian marine sponge *Cymbastela cantharella*. Amorph. solid. λ_{max} 203 (ε 2300); 231 (ε 1900); 260 (sh) (ε 1100) (EtOH). λ_{max} 215 (ε 2800); 243 (ε 2000); 277 (ε 900) (EtOH/NaOH aq).

Al Mourabit, A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 290-291 (*isol, synth, uv, ir, pmr, cmr*)

Guanine**G-210**

2-Amino-1,7-dihydro-6H-purin-6-one, 9CI. 2-Aminohypoxanthine. 2-Amino-6-hydroxypurine. 2-Amino-6-purinol. C.I. Natural White 1 [73-40-5]



C₅H₅N₅O 151.127

Occurs widely in animals and plants. Component of nucleic acids. Cryst. Mp 300°.

▶ MF8260000

Hydrochloride: [635-39-2]

[18602-14-7]

Needles + 2H₂O (dil. HCl).

2-N-Ac: [19962-37-9]

C₇H₇N₅O₂ 193.165

Cryst. (EtOH aq.). Mp 350°.

2-N-Benzoyl: [21323-87-5]

C₁₂H₉N₅O₂ 255.235

Cryst. (EtOH aq.). Mp 315-340°.

2-N-Me: N²-Methylguanine. 6-Hydroxy-2-methylaminopurine. 2-(Methylamino)hypoxanthine

[10030-78-1]

C₆H₇N₅O 165.154

Occurs in human urine and as a component of yeast nucleic acid.

Cryst. + 1H₂O (H₂O). Mp 200° dec. Mp 288° (as picrate). [α]_D²⁰ -34.6 (c, 1.01 in DMSO/EtOH 1:1).

2-N,N-Di-Me: N²,N²-Dimethylguanine.

2-Dimethylamino-6-hydroxypurine

[1445-15-4]

C₇H₉N₅O 179.181

Isol. from RNA of various plants. λ_{max} 235 (ε 6900); 258 (ε 14500); 290 (sh) (ε 6300) (H₂O).

7-Oxide: Guanine N⁷-oxide. 7-Hydroxyguanine. Antibiotic 3780

[5227-68-9]

C₅H₅N₅O₂ 167.127

Isol. from culture broths of *Streptomyces* spp. Antitumour, antimicrobial and antiviral agent. Brownish cryst. + 1/3H₂O. Mp 300°. λ_{max} 251 (ε 4510) (pH 1) (Derap). λ_{max} 234 (ε 13760); 291 (ε 3940) (H₂O) (Derap).

▶ LD₅₀ (mus, ipr) 53 mg/kg. MF8585000

1-Amino:

C₅H₆N₆O 166.142

Needles (as hydrochloride). Mp 215-218°. pK_{a1} 3.6; pK_{a2} 10.5.

3-Amino:

C₅H₆N₆O 166.142

Mp 250° (as hydrochloride). pK_{a1} 4.2; pK_{a2} 9.8.

7-Amino:

C₅H₆N₆O 166.142

Monohydrate (as hydrochloride). Mp 200-203° (hydrochloride). pK_{a1} 3.5; pK_{a2} 9.6.

9-Amino:

C₅H₆N₆O 166.142

Needles (1M HCl). Mp 250°. pK_{a1} 2.7; pK_{a2} 9.7.

1,7-Dihydro-form

1-Me: see 1-Methylguanine, M-467

1,7-Di-Me: see 1-Methylguanine, M-467

7-Benzyl: 2-Amino-1,7-dihydro-7-(phenylmethyl)-6H-purin-6-one, 9CI

[182005-51-2 (dihydrochloride)]

C₁₂H₁₁N₅O 241.252

Mp 285° dec. (dihydrochloride).

7-Benzyl, 2-N-Ac: [17495-10-2]

C₁₄H₁₃N₅O₂ 283.289

Mp 241°.

7-Benzyl, 2-N-benzoyl: [182005-36-3]

C₁₉H₁₅N₅O₂ 345.36

Mp 265-267°.

7-β-D-Xylofuranosyl: [99436-41-6]

C₁₀H₁₃N₅O₅ 283.243

Cryst. (H₂O). Mp 262-264°. [α]_D²⁰ -33.3 (c, 0.57 in DMSO).

1,9-Dihydro-form

Cryst. (H₂O).

2-N, 9-Di-Ac: [3056-33-5]

C₉H₉N₅O₃ 235.202

Mp 285°.

9-α-D-Xylofuranosyl: [27462-38-0]

C₁₀H₁₃N₅O₅ 283.243

Mp 186° dec. (260-261°). [α]_D²⁰ -17.1 (c, 0.76 in DMSO).

9-β-D-Xylofuranosyl: [27462-39-1]

C₁₀H₁₃N₅O₅ 283.243

Cryst. (H₂O). Mp 236-240°. [α]_D²⁰ -68.1 (c, 0.72 in DMSO).

▶ MF8610000

1,9-Di-Me: see 1-Methylguanine, M-467

7,9-Dihydro-form

Imidazolium betaine form

7,9-Di-Me: see Herbiopoline, H-143

OH-form

Et ether: 6-Ethoxy-1H-purin-2-amine,

9CI. 2-Amino-6-ethoxypurine. 6-O-

Ethylguanine

[51866-19-4]

C₇H₉N₅O 179.181

Cryst. (H₂O). Mp 293° dec.

Cyclohexylmethyl ether: O⁶-Cyclohexylmethylguanine. NU 2058

[161058-83-9]

C₁₂H₁₇N₅O 247.299

Cyclin-dependent kinase inhibitor. Solid. Mp 202-204°.

Aldrich Library of NMR Spectra, 2nd edn.,

1983, **2**, 593A (*nmr*),

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **2**, 716B (*ir*)

Balsiger, R.W. *et al.*, *J.O.C.*, 1960, **25**, 1573-

1575 (*Et ether*)

Gerster, J.F. *et al.*, *J.O.C.*, 1966, **31**, 3258

(*synth, derivs*)

Rice, J.M. *et al.*, *J.A.C.S.*, 1967, **89**, 2719 (*ms*)

Thewalt, U. *et al.*, *Acta Cryst. B*, 1971, **27**,

2358 (*cryst struct*)

Lister, J.H. *et al.*, *Chem. Heterocycl. Compd.*,

(ed. Weissberger, A. *et al.*), 1971, (*rev*)

Lee, W.W. *et al.*, *J.O.C.*, 1971, **36**, 842 (*synth*)

Sekiyama, M. *et al.*, *Chem. Pharm. Bull.*, 1976,

24, 1331 (*synth*)

Yamazaki, A. *et al.*, *Nucleic Acids Res.*, 1976, **3**, 251 (*synth*)

Dimitrijevič, S.D. *et al.*, *J.O.C.*, 1979, **44**, 400 (*deriv, synth*)

McGee, D.P.C. *et al.*, *Can. J. Chem.*, 1986, **64**, 1885 (*deriv, synth, uv, pmr, cmr*)

Mathlouth, M. *et al.*, *Carbohydr. Res.*, 1986, **146**, 15 (*ir, Raman*)

Gosselin, G. *et al.*, *J. Med. Chem.*, 1986, **29**, 203 (*derivs*)

Nohara, F. *et al.*, *Tet. Lett.*, 1987, **28**, 1287 (*7-oxide, bibl*)

Kohda, K. *et al.*, *Tetrahedron*, 1989, **45**, 6367 (*aminoguanines*)

Maixner, J. *et al.*, *Acta Cryst. C*, 1991, **47**, 2474 (*cryst struct, hydrochloride*)

Yamagata, Y. *et al.*, *Acta Cryst. C*, 1992, **48**, 318 (*cryst struct, 1-amino, 7-amino, 9-amino*)

Ogawa, K. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 343 (*7-oxide*)

Arris, C.E. *et al.*, *J. Med. Chem.*, 2000, **43**, 2797-2804; 2002, **45**, 3381-3393 (*NU 2058, synth, pharmacol*)


Guille, K. *et al.*, *Acta Cryst. C*, 2006, **62**, o515-o517 (*cryst struct*)

Torii, T. *et al.*, *Tetrahedron*, 2006, **62**, 5709-5716 (*7-benzyl derivs*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, GL1000

Guatambuine**G-211**

2,3,4,6-Tetrahydro-1,2,5-trimethyl-1H-pyrido[4,3-b]carbazole, 9CI. u-Alkaloid C



C₁₈H₂₀N₂ 264.369

Antineoplastic agent. Log P 4.72 (calc).

(+)-form [2744-45-8]

Alkaloid from *Aspidosperma australe* and some other *Aspidosperma* spp. (Apocynaceae). Mp 245-248°. [α]_D²⁵ +112 (c, 0.49 in Py). Pharmacol. active isomer. λ_{max} 240 (ε 41300); 250 (ε 31300); 262 (ε 22700); 299 (ε 19300); 330 (ε 4300) (EtOH).

(-)-form

Alkaloid from root bark of *Aspidosperma australe* (Apocynaceae). Mp 247-248°. [α]_D²⁵ -106 (Py). λ_{max} 240 (ε 41000); 250 (ε 31000); 262 (ε 22200); 299 (ε 19200) (EtOH).

(±)-form [11046-16-5]

Alkaloid from aerial bark of *Aspidosperma australe* (Apocynaceae). Mp 224-225°.

Methodide: Mp 299-301°.

(±)-form

N-De-Me: 2,3,4,6-Tetrahydro-1,5-dimethyl-1H-pyrido[4,3-b]carbazole, 9CI. Janetine. Tetrahydroolivacine [69754-02-5]

C₁₇H₁₈N₂ 250.343

Alkaloid from *Ervatamia coronaria* and *Stenosolen heterophyllus* (pre-

ferred genus name *Tabernaemontana* (Apocynaceae). Abs. config. not detd., no opt. rotn. reported. May correspond with (+), (-) or (±)-Guatambuine.

3-Hydroxy, N-de-Me: 3-Hydroxytetrahydrooolivacine

[167696-87-9]

C₁₇H₁₈N₂O 266.342

Alkaloid from stem bark of *Peschiera buchtieni* (Apocyanaceae). [α]_D +8 (c, 0.3 in EtOH).

Schmutz, J. et al., *Helv. Chim. Acta*, 1957, **40**, 1189-1200; 1958, **41**, 288-294 (*isol. uv. ir*)

Ondetti, M.A. et al., *Tetrahedron*, 1961, **15**, 160-166 (*isol. uv. struct*)

Besselièvre, R. et al., *Tet. Lett.*, 1976, 1873-1876 (*synth*)

Gomez Gonzalez, C. et al., *CA*, 1979, **90**, 138069u (*Janetine*)

Kutney, J.P. et al., *Can. J. Chem.*, 1982, **60**, 2426-2430 (*synth*)

Kan, C. et al., *J. Nat. Prod.*, 1984, **47**, 478-481 (*Janetine*)

Azoug, M. et al., *Phytochemistry*, 1995, **39**, 1223-1228 (*3-Hydroxytetrahydrooolivacine*)


Bennasar, M.-L. et al., *J.O.C.*, 2006, **71**, 1746-1749 (*synth*)

De Simone, C.H. et al., *Z. Kristallogr. - New Cryst. Struct.*, 2006, **221**, 233-234 (*(+)-form, cryst struct*)

Guattaguanine

G-212

[116127-41-4]



C₃₈H₄₂N₂O₆ 622.76

Alkaloids in this entry (*S,S*-config.) are epimeric with those under Cordobine, C-643 (*1R,1'S*-) and Rodiasine, R-114 (*1S,1'R*-). Semisynthetic, by methylation of 2'-Norguattaguanine. [α]_D +40 (c, 0.09 in CHCl₃).

N²-De-Me: 2'-Norguattaguanine

[116127-42-5]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). [α]_D +18 (c, 0.12 in CHCl₃).

Di-N-de-Me: 2,2'-Bisnorguattaguanine

[116064-71-2]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). [α]_D +40 (c, 0.8 in CHCl₃).

O¹²-De-Me, O¹²-Me, N-De-Me (?):

Dirosine

[1356-72-5]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the bark of *Ocotea rodiaei* (Lauraceae). Cryst. + 1½ H₂O (as hydrochloride). Mp 303° hydrochloride. [α]_D +97 (c, 1.0 in H₂O). Shown to be a stereoisomer of an *N*-Demethylrodiasine (see Rodiasine, R-114). Assigned here tentatively to the *S,S*-series on the basis of its opt. rotn.

Hearst, P.J. et al., *J.O.C.*, 1964, **29**, 466-470


(*Dirosine*)

Berthou, S. et al., *Tetrahedron*, 1998, **44**, 2193-2201 (*Guattaguanine, Norguattaguanines*)

Guatterine

G-213

[3912-56-9]



C₁₉H₁₉NO₄ 325.363

Alkaloid from *Guatteria psilopus*, *Pachypodanthium confine* and *Polyalthia suaveolens* (Annonaceae). Needles (Me₂CO aq.). Mp 146-148°. [α]_D²⁰ -57.1 (CHCl₃).

N-Oxide: Guatterine N-oxide

[62951-76-2]

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Polyalthia confine* (Annonaceae). Mp 196°. [α]_D -43 (c, 0.8 in CHCl₃).

7-Epimer, N-de-Me: Artabonatine B

[247244-63-9]

C₁₈H₁₇NO₄ 311.337

Alkaloid from fresh unripe fruit of *Artabotrys uncinatus*. Amorph. yellow powder. [α]_D²⁴ -121.5 (c, 0.8 in CHCl₃). Only rel. config. determined. λ_{max} 214 (log ε 4.25); 256 (log ε 4.18); 295 (log ε 3.86); 325 (log ε 3.56) (EtOH).

Harris, W.M. et al., *J.O.C.*, 1965, **30**, 432 (*isol. uv. pmr, ir, struct*)

Bévalot, F. et al., *Ann. Pharm. Fr.*, 1977, **35**, 65; *CA*, **86**, 185953e (*oxide*)


Cavé, A. et al., *Planta Med.*, 1978, **33**, 243 (*isol*)

Hsieh, T.-J. et al., *J. Nat. Prod.*, 1999, **62**, 1192-1193 (*Artabonatine B*)

Guattescidine

G-214

[82404-36-2]



C₁₈H₁₅NO₄ 309.321

Alkaloid from the stem bark of *Guatteria scandens* (Annonaceae). Amorph. [α]_D -165 (c, 0.6 in CHCl₃). Prob. has (*S*)-config.

O⁹-Me: Guattescine. 9-O-Methylguattescidine

[82404-37-3]

C₁₉H₁₇NO₄ 323.348

Alkaloid from the stem bark of *Guatteria scandens* (Annonaceae). Yellow cryst. (MeOH). Mp 160°. [α]_D

+26 (c, 0.87 in CHCl₃). Prob. has (*R*)-config.

O⁹-Me, Ac:

Cryst. (MeOH), Mp 153°. [α]_D +163 (c, 0.5 in CHCl₃).

4,5-Didehydro, O⁹-Me: Dehydroguattescine

[96997-99-8]

C₁₉H₁₅NO₄ 321.332

Alkaloid from the stem bark of *Guatteria schomburgkiana* (Annonaceae).

3-Methoxy: 3-Methoxyguattescidine

[158018-13-4]

C₁₉H₁₇NO₅ 339.347

Alkaloid from stem bark of *Guatteria foliosa* (Annonaceae). Amorph. [α]_D +37 (c, 0.23 in MeOH).

11-Methoxy: Guacolidine

[91174-12-8]

C₁₉H₁₇NO₅ 339.347

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Yellow needles (CH₂Cl₂). Mp 122°. [α]_D +4 (c, 0.78 in MeOH).

11-Methoxy, O⁹-Me: Guacoline. 9-O-Methylguacolidine

[91174-11-7]

C₂₀H₁₉NO₅ 353.374

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Noncryst. [α]_D -37 (c, 1.08 in MeOH).

Hocquemiller, R. et al., *Tetrahedron*, 1982, **38**, 911 (*Guattescidine, Guattescine, isol. uv. pmr, ms, cmr, ir*)

Hocquemiller, R. et al., *J. Nat. Prod.*, 1983, **46**, 335 (*Guacoline, Guacolidine, isol. uv. ir, pmr, ms, cmr, synth*)

Chiaroni, A. et al., *Tetrahedron*, 1983, **39**,

2163 (*pmr, cmr, struct, Guattescidine, Guattescine*)

Hocquemiller, R. et al., *J. Nat. Prod.*, 1984, **47**, 353 (*isol. uv. ir, pmr, cmr, ms, struct, Guacolidine, Guacoline*)

Cortes, D. et al., *J. Nat. Prod.*, 1985, **48**, 254

(*Dehydroguattescine*)

Mahiou, V. et al., *J. Nat. Prod.*, 1994, **57**, 890


(*3-Methoxyguattescidine*)

Atanes, N. et al., *Tetrahedron*, 1994, **50**, 11257 (*Guacoline, synth*)

Guattouregidine

G-215

[84391-85-5]



C₁₉H₁₉NO₅ 341.363

Alkaloid from the stem bark of *Guatteria ouregou* (Annonaceae). Amorph. [α]_D -31 (c, 0.17 in EtOH).

O⁹-Me: Guattouregine. 9-O-Methylguattouregidine

[84391-84-4]

C₂₀H₂₁NO₅ 355.39

Alkaloid from the stem bark of *Guatteria ouregou* (Annonaceae). Noncryst. [α]_D

$[\alpha]_D$ -69 (c, 0.10 in EtOH).

O²-De-Me, O¹-Me: Isoguattouregidine

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Guatteria melosma* (Annonaceae). λ_{\max} 216 (log ϵ 4.3); 235 (sh) (log ϵ 4.12); 268 (log ϵ 4.47); 308 (log ϵ 3.93); 355 (log ϵ 3.89) (no solvent reported).

Leboeuf, M. et al., *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1982, **295**, 191-194 (*Guattouregine, isol, uv, ir, pmr, cmr, ms, struct*)


Leboeuf, M. et al., *Planta Med.*, 1983, **48**, 234-245 (*isol, uv, ir, pmr, cmr, ms, struct*)

Guinaudeau, H. et al., *J. Nat. Prod.*, 1988, **51**, 389-474 (*Isoguattouregidine*)

Guayulamine A

G-216

[447457-74-1]



R = CH₂CH(CH₃)₂

C₁₈H₂₄N₂ 268.401

Alkaloid from the derubberised resin of the hybrid *Parthenium argentatum* x *Parthenium tomentosum*. Oil. Racemate. λ_{\max} 286 (no solvent reported).

Maatooq, G.T. et al., *Z. Naturforsch., C*, 2002, **57**, 211-215 (*isol, pmr, cmr, ms*)

Guayulamine B

G-217

[447457-75-2]

As Guayulamine A, G-216 with

R = CH₂CH₂CH(CH₃)₂

C₁₉H₂₆N₂ 282.428


Alkaloid from the derubberised resin of the hybrid *Parthenium argentatum* x *Parthenium tomentosum*. Oil. Racemate. λ_{\max} 289 (no solvent reported).

Maatooq, G.T. et al., *Z. Naturforsch., C*, 2002, **57**, 211-215 (*isol, pmr, cmr, ms*)

Guettardine

G-218

[91897-59-5]



Absolute Configuration

C₂₀H₃₀N₂O₂ 330.469

Alkaloid from the bark of *Guettarda heterosepala* (Rubiaceae). Amorph. solid. $[\alpha]_D^{20}$ -10 (c, 1 in EtOH). Possible biogenetic intermed. in the formn. of Corynanthe-Cinchona alkaloids. λ_{\max} 226 (log ϵ 4.54); 277 (log ϵ 3.89); 284 (log ϵ 3.91); 292 (log ϵ 3.85) (EtOH).


Brillianceau, M.H. et al., *Tet. Lett.*, 1984, **25**, 2767-2770 (*uv, pmr, cmr, ms, struct*)

Diez, A. et al., *Tetrahedron*, 1990, **46**, 4393-4406 (*synth*)

Guiaflavine

G-219

[229011-55-6]



Absolute Configuration

C₄₀H₄₂N₄O₂²⁺ 610.797

Quaternary alkaloid from the stem bark of *Strychnos guianensis*. Amorph. yellow-brown solid. Counterion not specified. λ_{\max} 208 (log ϵ 4.32); 252 (log ϵ 4.07); 321 (log ϵ 3.93); 433 (log ϵ 3.99) (MeOH).

5',6'-Didehydro: 5',6'-Dehydroguiaflavine

C₄₀H₄₀N₄O₂²⁺ 608.782

Quaternary alkaloid from *Strychnos guianensis*. Orange-brown powder. Counterion not specified. CAS no. not found 14CI. λ_{\max} 207 (log ϵ 4.49); 257 (log ϵ 4.35); 432 (log ϵ 4.11) (MeOH).


Penelle, J. et al., *J. Nat. Prod.*, 1999, **62**, 898-900 (*isol, uv, ir, pmr, cmr, ms*)

Penelle, J. et al., *Phytochemistry*, 2001, **58**, 619-626 (*Dehydroguiaflavine*)

Guianensine

G-220

[171828-69-6]



C₃₉H₃₆N₄O₄ 624.738

Zwitterion. Isol. from the stem bark of *Strychnos guianensis* (Loganiaceae). Orange powder.

Quetin-Leclercq, J. et al., *Phytochemistry*, 1995, **40**, 1557 (*isol, uv, ir, pmr, cmr, ms, struct*)

Guillauminine

G-221

[90599-29-4]

C₄₀H₄₈N₄O₂ 616.845


Struct. unknown. Bisindole alkaloid. Trace alkaloid from the stem bark and aerial parts of *Melodinus guillauminii* (Apocynaceae). $[\alpha]_D$ +186 (c, 1 in CHCl₃).

Zeches, M. et al., *Phytochemistry*, 1984, **23**, 171-174 (*isol, uv, ir, pmr, ms*)

Guineamide A

G-222

[560082-02-2]



C₃₁H₄₃N₅O₆S 613.777


Isol. from *Lyngbya majuscula*. Oil. $[\alpha]_D^{26}$ -8 (c, 0.1 in CHCl₃). λ_{\max} 214 (ϵ 14400) (MeOH).

Tan, L.T. et al., *J. Nat. Prod.*, 2003, **66**, 764-771 (*isol, pmr, cmr*)

Guineamide B

G-223

[560082-04-4]



C₃₂H₄₅N₅O₆S 627.803


Isol. from *Lyngbya majuscula*. Amorph. solid. $[\alpha]_D$ -5 (c, 0.17 in CHCl₃). λ_{\max} 213 (ϵ 24900) (EtOH).

Tan, L.T. et al., *J. Nat. Prod.*, 2003, **66**, 764-771 (*isol, pmr, cmr*)

Guineensine

G-224

13-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4,12-tridecatrienamide, 9CI. 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienic acid isobutylamide



C₂₄H₃₃NO₃ 383.53

(E,E,E)-form [55038-30-7]

Alkaloid from the fruits of *Piper guineense* and *Piper nigrum*, and the roots of *Piper attenuatum* (Piperaceae). ACAT inhibitor. Cryst. (C₆H₆/petrol). Mp 113-115°.

3-Methylbutylamide analogue: 13-(1,3-Benzodioxol-5-yl)-N-(3-methylbutyl)-2,4,12-tridecatrienamide, 9CI. **Pipwaqarine**

$C_{25}H_{35}NO_3$ 397.556

Alkaloid from the fruit of *Piper nigrum* (black pepper). Insecticide. Needles (EtOAc/petrol). Mp 110-112°. λ_{max} 269 (ε 37800); 312 (ε 3800) (MeOH).

Okogun, J.I. *et al.*, *J.C.S. Perkin 1*, 1974, 2195 (isol, uv, ir, pmr, ms, struct)

Dasgupta, S. *et al.*, *Indian J. Chem., Sect. B*, 1979, 17, 538 (isol, uv, ir, pmr, ms)

Okwute, S.K. *et al.*, *Tetrahedron*, 1984, 40, 2541 (synth)

Jacobs, H. *et al.*, *J. Indian Chem. Soc.*, 1999, 76, 713-717 (isol, pmr, cmr)

Park, I.-K. *et al.*, *J. Agric. Food Chem.*, 2002, 50, 1866-1870 (isol, pmr, cmr)


Siddiqui, B.S. *et al.*, *Nat. Prod. Res.*, 2005, 19, 143-150 (Pipwaqarine)

Gusanlung C

G-225

Methyl [2-[4-hydroxy-2-[2-(4-hydroxyphenyl)ethenyl]phenyl]ethyl]carbamate, 9CI

[165134-18-9]



$C_{18}H_{19}NO_4$ 313.352

Alkaloid from stems of *Arcangelisia gusanlung* (Menispermaceae). Yellow needles. Mp 104-105°. Genus name erroneously given as Acangelisia.

Di-Ac: [164920-60-9]

Mp 155-156°.

Di-Me ether: [164920-61-0]


Fine needles (MeOH). Mp 126-127°.

Zhang, J.-S. *et al.*, *Phytochemistry*, 1995, 39, 439 (isol, uv, ir, pmr, cmr, ms, struct)

Gusanlung D

G-226

5,6,13,13a-Tetrahydro-8H-benzo[g]-1,3-benzodioxolo[5,6-a]quinolizin-8-one, 9CI



$C_{18}H_{15}NO_3$ 293.321

(*S*)-form [165306-67-2]

Alkaloid from stems of *Arcangelisia gusanlung*. Needles. Mp 250-251°. $[\alpha]_D^{20}$ -345 (c, 0.018 in $CHCl_3$). Genus name erroneously given as Acangelisia. λ_{max} 222; 273; 294; 320 (MeOH).


Zhang, J.-S. *et al.*, *Phytochemistry*, 1995, 39, 439-442 (isol, pmr, cmr)

Chang, J.-K. *et al.*, *Tetrahedron*, 2008, 64, 3483-3487 (synth)

Gutingimycin

[690998-82-4]

G-227



$C_{47}H_{57}N_5O_{21}$ 1027.988


Closely related to Trioxacarcin A. Prod. by a marine streptomycete (isolate B8652). Yellow solid. $[\alpha]_D^{20}$ -56.5 (c, 0.5 in $CHCl_3$). λ_{max} 269 (log ε 5.53); 399 (log ε 4.96) (MeOH).

Maskey, R.P. *et al.*, *Angew. Chem., Int. Ed.*, 2004, 43, 1281-1283 (isol, pmr, cmr)

Gymnamine

G-228

1,8-Dihydro-1-(1-oxopropyl)lycodin-8-one [38948-15-1]



$C_{19}H_{26}N_2O_3$ 330.426

Alkaloid from leaves of *Gymnema sylvestre* (Asclepiadaceae). Noncryst. solid. Mp 60-70°. First lycopodane-type alkaloid from outside the Lycopodiaceae.

Picrate:


Fine needles. Mp 260°.

Rao, G.S. *et al.*, *Chem. Ind. (London)*, 1972, 537 (isol, ms, struct)

Gymnastatin F

G-229

[229334-19-4]



$C_{24}H_{35}Cl_2NO_5$ 488.45

Prod. by *Gymnascella dankaliensis* isol. from a *Halichondria* sp. Powder. $[\alpha]_D^{26}$ -77.7 (c, 0.16 in $CHCl_3$). λ_{max} 263 (log ε 4.25) (EtOH).

Demethoxy: **Gymnastatin R**

$C_{23}H_{33}Cl_2NO_4$ 458.423

Prod. by *Gymnascella dankaliensis* isol.

from a *Halichondria* sp. Powder. Mp 79-82°. $[\alpha]_D^{24}$ -104.5 (c, 0.48 in EtOH). λ_{max} 264 (log ε 4.47) (EtOH).

6-Dechloro, 5β,6β-epoxide, O-de-Me:

Gymnastatin G

[229334-20-7]

$C_{23}H_{34}ClNO_6$ 455.977

Prod. by *Gymnascella dankaliensis* isol. from a *Halichondria* sp. Powder. $[\alpha]_D$ -53.1 (c, 1.5 in $CHCl_3$). λ_{max} 266 (log ε 4.63) (EtOH).

1-Epimer: **Gymnastatin Q**

$C_{24}H_{35}Cl_2NO_5$ 488.45

Prod. by *Gymnascella dankaliensis* isol. from a *Halichondria* sp. Powder. Mp 105-108°. $[\alpha]_D^{23}$ -34.3 (c, 0.26 in $CHCl_3$). λ_{max} 264 (log ε 4.45) (EtOH).


Amagata, T. *et al.*, *J. Nat. Prod.*, 2006, 69, 1384-1388 (*Gymnastatins F,G*)

Amagata, T. *et al.*, *J. Nat. Prod.*, 2008, 71, 340-345 (*Gymnastatins Q,R*)

Gymnastatin L

G-230

[234757-11-0]



$C_{23}H_{33}NO_3$ 371.519


Metab. of *Gymnascella dankaliensis* isol. from the sponge *Halichondria japonica*.

Amagata, T. *et al.*, *CA*, 1999, 131, 129789s (isol)

Gymnasterone A

G-231

[209169-57-3]



$C_{45}H_{67}NO_5$ 702.028

Constit. of *Gymnascella dankaliensis* isol. from the sponge *Halichondria japonica*. Pale yellow oil. $[\alpha]_D$ -110.7 (c, 1.44 in $CHCl_3$). λ_{max} 270 (log ε 4.22) (EtOH). λ_{max} 270 (ε 16600) (MeOH) (Berdy).


Amagata, T. *et al.*, *Tet. Lett.*, 1998, 39, 3773-3774 (isol, pmr, cmr)

Amagata, T. *et al.*, *J. Nat. Prod.*, 2007, 70, 1731-1740 (stereochem)

Gymnodimine

G-232

[173792-58-0]



C₃₂H₄₅NO₄ 507.712

Isol. from New Zealand oysters (*Tiostrea chilensis*) and the dinoflagellate *Gymnodinium* cf. *mikimotoi*. Shows potent ichthyotoxicity. Amorph. solid. $[\alpha]_D^{25}$ -10.4 (c, 0.13 in MeOH).

$\Delta^{17,29}$ -Isomer, 18R-hydroxy: **Gymnodimine C**

[586959-19-5]

C₃₂H₄₅NO₅ 523.711

Prod. by *Karenia selliformis* (formerly *Gymnodinium selliforme*).

$\Delta^{17,29}$ -Isomer, 18S-hydroxy: **Gymnodimine B**

[266998-21-4]

C₃₂H₄₅NO₅ 523.711

Isol. from *Karenia selliformis* (formerly *Gymnodinium selliforme*).


Seki, T. et al., *Tet. Lett.*, 1995, **36**, 7093-7096 (isol, pmr, cmr, struct)

Stewart, M. et al., *Tet. Lett.*, 1997, **38**, 4889-4890 (cryst struct, abs config)

Miles, C.O. et al., *J. Agric. Food Chem.*, 2000, **48**, 1373-1376; 2003, **51**, 4838-4840 (*Gymnodimines B, C*)

Gynuramine**G-233**

[85611-43-4]

C₁₈H₂₅NO₆ 351.399

Retronecine cyclic diester. Alkaloid from *Gynura scandens* (Asteraceae). Mp 204°. $[\alpha]_D^{20}$ -16 (CHCl₃).

1''-Ac: **Acetylgynuramine**

[85547-34-8]

C₂₀H₂₇NO₇ 393.436


Alkaloid from *Gynura scandens* (Asteraceae). Mp 153-155°. $[\alpha]_D^{20}$ -33 (CHCl₃).

Wiedenfeld, H. et al., *Phytochemistry*, 1982, **21**, 2767 (isol, ir, pmr, cmr, ms, struct, deriv)

Wiedenfeld, H. et al., *Phytochemistry*, 1983, **22**, 2065 (cryst struct, deriv)

Gypsetin**G-234**

[155114-38-8]



Absolute Configuration

C₃₂H₃₆N₄O₄ 540.661

Prod. by *Nannizzia gypsea* var. *incurvata* IFO 9228/IMI 86518. Acyl-CoA:cholesterol acyltransferase inhibitor and cytotoxic agent. Needles (CHCl₃/MeOH). Mp 162°. $[\alpha]_D$ -116.9 (c, 0.14 in CHCl₃). λ_{\max} 205 (ε 58900); 237 (ε 14900); 296 (ε 4430) (MeOH).

2,3-Diepimer: **Fellutanine D**

C₃₂H₃₆N₄O₄ 540.661

Prod. by *Penicillium fellutanum*. Microcryst. (CHCl₃/hexane). Mp 198-201°. $[\alpha]_D^{20}$ -383 (c, 0.1 in MeOH). Struct. subject of erratum in 2001. λ_{\max} 214 (log ε 4.1); 244 (log ε 3.2); 297 (log ε 3.53) (MeOH).

Shinohara, C. et al., *J. Antibiot.*, 1994, **47**, 163-167 (isol, uv, ir, activity)


Nuber, B. et al., *J. Antibiot.*, 1994, **47**, 168-172 (pmr, cmr, cryst struct)

Schkeryantz, J.M. et al., *J.A.C.S.*, 1999, **121**, 11964-11975 (synth)

Kozlovsky, A.G. et al., *J. Nat. Prod.*, 2000, **63**, 698-700; 2001, **64**, 553 (*Fellutanine D*)

Gyrolidine**G-235**

[39020-36-5]

C₃₈H₄₂N₂O₆ 622.76

These alkaloids have (1*S*,1'*R*)-config. (Shamma-Moniot convention) and are enantiomeric with those given under Obaberine, O-1. The older CAS numbering puts the primes on the opposite rings. Alkaloid from *Gyrocarpus amer-*

icanus (Hernandiaceae). Amorph. $[\alpha]_D$ -115 (c, 1.1 in CHCl₃). λ_{\max} 261 (log ε 4.23); 282 (log ε 4.12) (MeOH).

O⁶-De-Me: **Gyrocarpine**

[102487-16-1]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stem bark of *Gyrocarpus americanus* (Hernandiaceae). Cryst. (MeOH/Et₂O). Mp 192°. $[\alpha]_D$ -239 (c, 1 in CHCl₃). λ_{\max} 239 (sh) (log ε 4.34); 284 (log ε 3.87) (MeOH).

O⁷,O¹²-Di-de-Me: **Macolidine**

[66288-77-5]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Abuta grisebachii* (Menispermaceae). Mp 179-181°. $[\alpha]_D^{20}$ -320 (CHCl₃).

O⁷,O¹²-Di-de-Me, N^{2'}-Me: **Macoline**

[66216-59-9]

[66216-60-2]

C₃₇H₄₁N₂O₆[⊕] 609.741

Alkaloid from *Abuta grisebachii* (Menispermaceae). Mp 255-259° (as chloride hydrochloride). $[\alpha]_D^{20}$ -60.6 (MeOH) (chloride hydrochloride).

Galeffi, C. et al., *Farmac. Ed. Sci.*, 1977, **32**, 853; *CA*, **89**, 6465q (*Macoline, Macolidine*)

Chalandre, M.C. et al., *J. Nat. Prod.*, 1986, **49**, 101-105 (*Gyrolidine, Gyrocarpine*)

Gyromitritin**G-236**

Ethylidenemethylhydrazinecarboxaldehyde, 9CI. N'-Ethylidene-N-formyl-N-methylhydrazine. Acetaldehyde N-formyl-N-methylhydrazone

[16568-02-8]

H₃CCH=NNMeCHOC₄H₈N₂O 100.12

Toxin from the fungi *Gyromitra esculenta* and *Helvella gigas*. Freq. cause of mushroom poisoning. Unstable cryst. Mp 19.5° (5°).

► Toxic, carcinogenic. Toxicity varies greatly between individuals. LQ8500000

List, P.H. et al., *Tet. Lett.*, 1967, 1893 (isol)

Zinner, G. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1971, **304**, 706 (synth)

Pyysalo, H. et al., *Acta Chem. Scand., Ser. B*, 1976, **30**, 235; 792 (isol, ms)


Zelenin, K.N. et al., *Zh. Org. Khim.*, 1978, **14**, 490 (pmr, conformn)

Meier-Bratschi, A. et al., *J. Agric. Food Chem.*, 1983, **31**, 1117 (synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, AAH000

Habbemine A

[937023-27-3]

Relative
ConfigurationC₁₆H₂₅NO₃ 279.378

Tautomeric. Alkaloid from the leaves of *Elaeocarpus habbemine*. Yellow gum (as hydrochloride). [α]_D²³ +13.7 (c, 0.13 in MeOH). Isol. as a 1:1 mixt. with Habbemine B to which the data refers. λ_{max} 227 (log ε 3.35); 270 (log ε 2.85); 339 (log ε 2.7) (MeOH).

2-Epimer: Habbemine B

[937023-28-4]


C₁₆H₂₅NO₃ 279.378

Alkaloid from the leaves of *Elaeocarpus habbemine*. Isol. as a 1:1 mixt. with Habbemine A.

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 866-868 (isol, pmr, cmr, ms)

Habranthine

[21502-30-7]

C₁₇H₂₁NO₄ 303.357

Alkaloid from *Habranthus brachyandrus* (*Hippeastrum brachyandrum*) (Amaryllidaceae). Mp 198-199°. [α]_D²⁵ -320.

Wildman, W.C. *et al.*, *Tet. Lett.*, 1968, 4573 (ir, pmr, ms, struct)

HachijodinesHachijodine A R = -(CH₂)₁₂NHOMeB R = -(CH₂)₉CH(CH₃)CH₂CH₂NHOMeC R = -(CH₂)₁₀CH(CH₃)CH₂NHOMeD R = -CH₂CH(CH₃)(CH₂)₁₀NHOMeE R = -(CH₂)₁₃NMeOHF R = -(CH₂)₄C≡C(CH₂)₈NMeOHG R = -(CH₂)₄C≡CCH=CH(CH₂)₈NMeOHCribrocholine B R = -(CH₂)₁₁CH(CH₃)CH₂NHOMe

Alkaloids from either *Amphimedon* sp. or *Xestospongia* sp. Cytotoxic agents.

Hachijodine A [265114-90-7]C₁₈H₃₂N₂O 292.464

From a *Xestospongia* sp. λ_{max} 260 (ε 3300) (MeOH).

Hachijodine B [265114-91-8]C₁₉H₃₄N₂O 306.49

From a *Xestospongia* sp. λ_{max} 260 (ε 3300) (MeOH).

H-1

Hachijodine C**Cribrocholine A**

[265114-92-9]

C₁₉H₃₄N₂O 306.49

Isol. from the sponges *Xestospongia* sp. and *Cribrochalina* sp. [α]_D²⁴ -1 (c, 0.2 in MeOH). Related to Niphatesine C, N-225. λ_{max} 260 (ε 3200) (MeOH). λ_{max} 210 (ε 5630); 262 (ε 4530); 269 (ε 3330) (MeOH).

Hachijodine D [265114-93-0]C₁₉H₃₄N₂O 306.49

From a *Xestospongia* sp. λ_{max} 260 (ε 3200) (MeOH).

N¹²-β-D-Glucopyranosyl: Amphimedoside DC₂₅H₄₄N₂O₆ 468.632

Alkaloid from an *Amphimedon* sp.

Hachijodine E [265114-94-1]C₁₉H₃₄N₂O 306.49

From an *Amphimedon* sp.

Hachijodine F [265114-95-2]C₂₀H₃₂N₂O 316.486

From an *Amphimedon* sp.

Hachijodine G [265114-96-3]C₂₂H₃₄N₂O 342.523

From an *Amphimedon* sp.

Cribrocholine BC₂₀H₃₆N₂O 320.517

Isol. from the sponge *Cribrochalina* sp.

Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 682-684 (isol, pmr, Hachijodines B-G)

Nicholas, G.M. *et al.*, *Tetrahedron*, 2000, **56**, 2921-2927 (*Cribrocholine A, B*)


Goundry, W.R.F. *et al.*, *Tetrahedron*, 2003, **59**, 1719-1729 (*Hachijodines F, G, synth*)

Romeril, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 3273-3277 (*Hachijodine B, synth*)

Takekawa, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1503-1505 (*Amphimedoside D*)

Hackelidine

[122222-01-9]

C₁₅H₂₁NO₆ 311.334

Alkaloid from *Hackelia californica* (Boraginaceae).

7-Ac: Longitubine. 7-Acetylhackelidine

[121817-17-2]

C₁₇H₂₃NO₇ 353.371

Alkaloid from *Hackelia californica* (Boraginaceae).

7-Angeloyl: Latifoline†

[6029-86-3]

C₂₀H₂₇NO₇ 393.436

Diester of retronecine with angelic and latifolic acids. Alkaloid from *Cynoglossum latifolium* and *Hackelia floribunda* (Boraginaceae). Long needles (petrol). Mp 102-103°. [α]_D²⁴ +57 (c, 1.99 in EtOH).

7-Angeloyl, picrate: Mp 173-174°.

7-Angeloyl, N-oxide: Latifoline N-oxide

[98752-06-8]

C₂₀H₂₇NO₈ 409.435

Alkaloid from *Hackelia floribunda* (Boraginaceae). Oil. [α]_D²⁵ +29.1 (c, 1.10 in EtOH).

Stereoisomer, 7-angeloyl: Neolatifoline

[121916-88-9]

C₂₀H₂₇NO₇ 393.436

Alkaloid from *Hackelia longituba*. Gum (impure). [α]_D +48 (EtOH)(impure).

Crowley, H.C. *et al.*, *Aust. J. Chem.*, 1962, **15**, 139 (*Latifoline*)

Matsumoto, T. *et al.*, *Chem. Lett.*, 1973, 773 (*Latifoline, abs config*)

Hagglund, K.M. *et al.*, *J. Nat. Prod.*, 1985, **48**, 638 (*Latifoline, isol, pmr, cmr, Latifoline N-oxide*)

Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1988, **41**, 1827 (*Neolatifoline*)

Stermitz, F.R. *et al.*, *Tet. Lett.*, 1989, **30**, 7153 (*Hackelidine*)

Li, Y. *et al.*, *Huaxue Xuebao*, 1990, **48**, 415; *CA*, **113**, 168971t (*Hackelidine*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1992, **45**, 451 (*Latifoline, cryst struct, abs config*)


Drutu, I. *et al.*, *J.O.C.*, 2001, **66**, 7025-7029 (*Latifoline, synth*)

Haemanthamine

H-5

Natalensine. 3-Epicrinamine. Hemanthamine

[466-75-1]



(+) -form

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Haemanthus puniceus x katherinae* and a v. large number of spp. in the Amaryllidaceae. Mp 203-203.5°. [α]_D²⁵ +19.7 (c, 3.8 in MeOH). [α]_D²⁵ +33 (c, 1.25 in CHCl₃). λ_{max} 240 (ε 3220); 297 (ε 5200) (EtOH).

Picrate: Mp 221°.

Ac: 11-O-AcetylhaemanthamineC₁₉H₂₁NO₅ 343.379

Alkaloid from *Narcissus bujei*. Mp 92-96°. [α]_D²² -9.1 (c, 0.55 in MeOH).

Ac; perchlorate: Mp 209°.

O-De-Me: O-Demethylhaemanthamine.**11-Hydroxyvittatine**

[23610-70-0]

C₁₆H₁₇NO₄ 287.315

Alkaloid from *Galanthus plicatus* ssp. *byzantinus*, *Hippeastrum equestre*, *Hippeastrum puniceum* and *Pantratum maritimum*. Mp 248-250°. [α]_D²⁵ +35 (c, 0.1 in MeOH).

1,2-Dihydro: DihydrohaemanthamineC₁₇H₂₁NO₄ 303.357

Alkaloid from *Crinum powellii* var. *harlemense* (Amaryllidaceae). Cryst. (Me₂CO). Mp 226°. [α]_D²⁴ +78 (c, 0.2 in CHCl₃).

1,2-Dihydro, picrate:

Cryst. (MeOH). Mp 210°.


1,2-Dihydro, O-(2-hydroxybutyl): Narcimarkine

- [77809-91-7]
C₂₁H₂₉NO₅ 375.464
Alkaloid from bulbs of *Narcissus poetrius* (Amaryllidaceae). Fine needles (Me₂CO). Mp 110°. Stereochem. not well defined.
- 4 α -Hydroxy: Albiflomanthine**
[156408-70-7]
C₁₇H₁₉NO₅ 317.341
Alkaloid from bulbs of *Haemanthus albiflos* (Amaryllidaceae). Amorph. solid. $[\alpha]_D^{20} +31$ (c, 0.05 in EtOH).
- 6-Hydroxy:** see Haemanthidine, H-6
- 3-Epimer: Crinamine**
[639-41-8]
C₁₇H₁₉NO₄ 301.341
Alkaloid from *Crinum asiaticum*, many other *Crinum* spp. and several other spp. in the Amaryllidaceae such as *Narcissus tazetta*, *Crinum asiaticum*, *Brunsvigia josephinae*, *Pancreatum foetidum*. Powerful transient hypotensive agent in dogs. Respiratory depressant, antineoplastic agent. Active against gram-positive bacteria, shows antimalarial props. Mp 198-199°. $[\alpha]_D^{28} +157$ (c, 1.65 in CHCl₃). λ_{max} 237 (ϵ 3400); 297 (ϵ 3120) (EtOH) (Berdy).
- LD₅₀ (dog) 10 mg/kg.
- 3-Epimer, perchlorate:** Mp 201-201.5° dec.
- 3-Epimer, picrate:** Mp 273-274° dec.
- 3-Epimer, O-de-Me: Hamayne. Demethylcrinamine. Bulbispermine**
[61948-11-6]
C₁₆H₁₇NO₄ 287.315
12-Config. is R-. Alkaloid from *Crinum asiaticum* var. *japonicum*, *Crinum bulbispermum*, *Crinum latifolium*, *Crinum zeylanicum*, *Crinum jogus*, *Crinum asiaticum-japonicum* and *Brunsvigia radulosa* (Amaryllidaceae). Shows antimalarial props. Plates + 2/3 H₂O (Me₂CO), cryst. (Me₂CO or MeOH/EtOAc). Mp 83-84° Mp 130-132° (dimorph. ?). $[\alpha]_D^{20} +106.7$ (c, 1.02 in MeOH). $[\alpha]_D^{18} +78.2$ (c, 1.1 in EtOH).
- 3-Epimer, O-de-Me, picrate:**
Cryst. (Me₂CO). Mp 205-208°.
- 3-Epimer, O-de-Me, 3-Ac: 3-O-Acetylhamayne**
C₁₈H₁₉NO₅ 329.352
Isol. from the bulbs of *Crinum latifolium* and *Crinum zeylanicum* (Amaryllidaceae). Prisms (C₆H₆). Mp 115-116° (112-114°). $[\alpha]_D^{20} +126$ (c, 0.5 in CHCl₃).
- 3-Epimer, O-de-Me, di-Ac:** Mp 77.5-78° Mp 119-121°.
- 3-Epimer, O-de-Me, 11-O-(3-hydroxybutanoyl): 11-O-(3-Hydroxybutanoyl)-hamayne**
[946570-69-0]
C₂₀H₂₃NO₆ 373.405
Alkaloid from *Galanthus nivalis*.
- 3-Epimer, O-de-Me, 11-O-(3-hydroxybutanoyl), 3-O-(2E-butenoyl): 3-O-(2E-Butenoyl)-11-O-(3-hydroxybutanoyl)-hamayne**
[946570-71-4]
C₂₄H₂₇NO₇ 441.48
Alkaloid from *Galanthus nivalis*.
- 3-Epimer, O-de-Me, 3,11-bis-O-(3-hydroxybutanoyl): 3,11-Bis-O-(3-hydroxybutanoyl)hamayne**
[946570-70-3]
C₂₄H₂₉NO₈ 459.495
Alkaloid from *Galanthus nivalis*.
- 3-Epimer, O-de-Me, 3-O-[3-(3-hydroxybutanoyloxy)butanoyl], 11-O-(3-hydroxybutanoyl): 3,3',11-Tris-O-(3-hydroxybutanoyl)hamayne**
[946572-18-5]
C₂₈H₃₅NO₁₀ 545.585
Alkaloid from *Galanthus nivalis*.
- 3-Epimer, O-de-Me, 3-O-[4-(2-aminoethyl)phenyl]: 3-[4-(2-Aminoethyl)phenoxy]bulbispermine**
C₂₄H₂₆N₂O₄ 406.48
Alkaloid from *Crinum moorei*. Amorph. $[\alpha]_D^{29} +53.3$ (c, 0.04 in CHCl₃).
- 11-Epimer: 11-Epihaemanthamine**
[1472-76-0]
C₁₇H₁₉NO₄ 301.341
Alkaloid from *Haemanthus katherinae*. Prisms (Me₂CO). Mp 205°. $[\alpha]_D -24$ (c, 0.2 in CHCl₃).
- 11-Epimer, O-de-Me: 1,2-Didehydro-3,11-dihydroxycrinane. 3,11-Dihydroxy-1,2-dihydrocrinane**
[151003-81-5]
C₁₆H₁₇NO₄ 287.315
Alkaloid from the bulbs of *Pancreatum maritimum*. Cryst. (CHCl₃). Mp 128°. $[\alpha]_D^{25} -28$ (c, 0.04 in MeOH).
- 3,11-Diepimer, O-de-Me, 3-Ac: Yemenine A**
C₁₈H₁₉NO₅ 329.352
Alkaloid from the bulbs of *Crinum yemense*. Powder. $[\alpha]_D^{25} +61.7$ (c, 0.5 in CHCl₃). λ_{max} 243 (log ϵ 3.56); 292 (log ϵ 3.8) (MeOH).
- Mason, L.H. *et al.*, *J.A.C.S.*, 1955, **77**, 1253 (*Crinamine, isol, uv*)
Wildman, W.C. *et al.*, *J.A.C.S.*, 1958, **80**, 6465 (*Crinamine, struct*)
Fales, H.M. *et al.*, *J.A.C.S.*, 1960, **82**, 197; 3368 (*uv, ir, struct, abs config*)
Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 868; *CA*, **58**, 11415g (*Dihydrohaemanthamine*)
Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902 (*ms*)
Haugwitz, R.D. *et al.*, *J.C.S.*, 1965, 2001 (*pmr*)
Clardy, J. *et al.*, *J.A.C.S.*, 1970, **92**, 6337 (*cryst struct*)
Battersby, A.R. *et al.*, *Chem. Comm.*, 1971, 183 (*biosynth*)
Kirby, G.W. *et al.*, *Chem. Comm.*, 1971, 187; 415 (*biosynth*)
Tsuda, Y. *et al.*, *Chem. Comm.*, 1971, 1555 (*synth*)
Nogueiras, C. *et al.*, *Tet. Lett.*, 1971, 2743 (*11-Epihaemanthamine*)
Ochi, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3363 (*Hamayne, uv, ir, pmr, struct*)
Isobe, K. *et al.*, *Tet. Lett.*, 1976, 2331 (*Crinamine, synth*)
Roques, R. *et al.*, *Acta Cryst. B*, 1977, **33**, 3696 (*Crinamine, cryst struct*)
Onyiriuka, O.S. *et al.*, *Isr. J. Chem.*, 1978, **17**, 185 (*Crinamine, ms*)
- Döpke, W. *et al.*, *Z. Chem.*, 1981, **21**, 71 (*Narcinarkine*)
Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3015 (*Hamayne, 3-Acetylhamayne, isol, ir, pmr, ms, ord*)
Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3023 (*Hamayne, 3-Acetylhamayne, isol, ir, pmr*)
Ali, A.A. *et al.*, *Planta Med.*, 1984, **50**, 424 (*Hamayne, uv, ir, pmr, cmr, ms, cd, struct*)
Amigó, J.M. *et al.*, *Acta Cryst. C*, 1988, **44**, 1497 (*cryst struct*)
Adesanya, S.A. *et al.*, *Int. J. Pharmacogn.*, 1992, **30**, 303-307 (*Crinamine, activity*)
Baudouin, G. *et al.*, *Heterocycles*, 1994, **38**, 965 (*Albiflomanthine*)
Sener, B. *et al.*, *J. Chem. Soc. Pak.*, 1994, **16**, 275-279 (*3,11-Dihydroxy-1,2-dihydrocrinane*)
Viladomat, F. *et al.*, *Phytochemistry*, 1994, **35**, 809-812 (*Hamayne, 3-Acetylhamayne*)
Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591-6600 (*isol, cd*)
Nishimata, T. *et al.*, *J.O.C.*, 1998, **63**, 7586-7587 (*Crinamine, synth*)
Labrana, J. *et al.*, *Phytochemistry*, 1999, **50**, 183-188 (*11-Acetylhaemanthamine*)
Elgorashi, E.E. *et al.*, *Phytochemistry*, 2001, **56**, 637-640 (*Aminoethylphenoxybulbispermine*)
Unver, N. *et al.*, *Planta Med.*, 2003, **69**, 869-871 (*11-Hydroxyvittatine*)
Abdel-Halim, O.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1119-1124 (*Yemenine A*)
Zhang, F.-M. *et al.*, *Tetrahedron*, 2006, **62**, 9446-9455 (*Crinamine, synth*)
Berkov, S. *et al.*, *Phytochemistry*, 2007, **68**, 1791-1798 (*3-Hydroxybutanoylhamaynes*)
Bohno, M. *et al.*, *Tetrahedron*, 2007, **63**, 6977-6989 (*synth*)
Tam, N.T. *et al.*, *J.O.C.*, 2008, **73**, 6258-6264 (*Crinamine, synth*)

Haemanthidine

H-6

Pancreatine. Hemanthidine
[466-73-9]

C₁₇H₁₉NO₅ 317.341

Exists in soln. as a mixt. of C-6 epimers. Alkaloid from *Haemanthus punicese* x *katherinae*, other *Haemanthus* spp. and many other spp. in the Amaryllidaceae. Mp 189-190° (hemihydrate). $[\alpha]_D^{22} -41$ (c, 1 in CHCl₃).

Picrate: Mp 208° dec.

Di-Ac: Mp 200-205° dec.

O⁶-Me: 6-O-Methylhaemanthidine
[120293-53-0]

C₁₈H₂₁NO₅ 331.368
Alkaloid from bulbs of *Pancreatum maritimum* (Amaryllidaceae). Cryst. (Me₂CO). Mp 110-112°. $[\alpha]_D -43$ (c, 0.46 in CHCl₃).

1,2-Dihydro: Hippawine

[639-37-2]

C₁₇H₂₁NO₅ 319.357

Isol. from the *Hippeastrum* hybrid "Queen of the Whites" (Amaryllidaceae). Cryst. (Me₂CO). Mp 203°. [α]_D²⁵ 0 (c, 0.2 in CHCl₃).

1,2-Dihydro, picrate: Mp 152° dec.

1,2-Dihydro, di-Ac: Mp 168° dec.

3-Epimer: **6-Hydroxycrinamine**. 3-Epihaemanthidine

[545-66-4]

C₁₇H₁₉NO₅ 317.341

Alkaloid from *Haemanthus natalensis* and from several *Crinum* spp. (Amaryllidaceae). Antitumour agent mod. active against Rauscher leukaemia virus. Mp 140° (hydrate) Mp 211° (anhyd.). [α]_D²⁵ +46 (c, 1 in CHCl₃). Exists in soln. as a mixt. of C-6 epimers. The 6S-epimer occurs in the crystal. λ_{\max} 240 (ε 3500); 294 (ε 4600) (EtOH) (Berdy).

3-Epimer, methiodide: Mp 174°. [α]_D²³ +57 (c, 1 in EtOH).

3-Epimer, di-Ac: Mp 182-184°.

3-Epimer, O⁶-Me: **6-Methoxycrinamine**

[93452-25-6]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the bulbs of *Crinum zeylanicum* (Amaryllidaceae). Oil; yellow needles (EtOAc)(as picrate). Mp 205-207° (picrate).

3-Epimer, O-de-Me: **Yemenine C**

C₁₆H₁₇NO₅ 303.314

Alkaloid from the bulbs of *Crinum yemense*. Fine needles (MeOH). Mp 152-154°. [α]_D²⁵ -14 (c, 0.2 in MeOH). Exists as a mixt. of C-6 epimers. λ_{\max} 241 (log ε 3.58); 293 (log ε 3.7) (MeOH).

3,11-Diepimer, O-de-Me: **Yemenine B**

C₁₆H₁₇NO₅ 303.314

Alkaloid from the bulbs of *Crinum yemense*. Powder. [α]_D²⁵ +92 (c, 0.2 in MeOH). Exists in soln. as a mixt. of C-6 epimers. λ_{\max} 243 (log ε 3.56); 292 (log ε 3.68) (MeOH).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1954, **87**, 1339-1342 (*isol*)

Uyeo, S. *et al.*, *J.A.C.S.*, 1958, 2590-2591 (*struct*)

Goosen, J. *et al.*, *J.C.S.*, 1960, 1088-1089 (*6-Hydroxycrinamine*)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 470 (*Hippavine*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902-4912 (*ms*)

King, R.W. *et al.*, *J.A.C.S.*, 1965, **87**, 4912-4917 (*pmr, config*)

Haugwitz, R.D. *et al.*, *J.C.S.*, 1965, 2001-2009 (*pmr, bibl*)

Karle, J. *et al.*, *J.A.C.S.*, 1967, **89**, 6510-6514 (*6-Hydroxycrinamine, cryst struct*)

Tsuda, Y. *et al.*, *Tet. Lett.*, 1972, 3153-3156 (*synth*)

Hendrickson, J.B. *et al.*, *J.A.C.S.*, 1974, **96**, 7781-7789 (*synth*)

Isobe, K. *et al.*, *Tet. Lett.*, 1976, 2331-2334 (*6-Hydroxycrinamine, synth*)

Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3023-3027 (*6-Methoxycrinamine*)

Martin, S.F. *et al.*, *J.A.C.S.*, 1984, **106**, 6431-6433 (*synth*)

Vázquez Tato, M.P. *et al.*, *Heterocycles*, 1988, **27**, 2833-2838 (*6-O-Methylhaemanthidine*)

Ishibashi, H. *et al.*, *J.O.C.*, 1993, **58**, 2360-2368 (*synth*)

Viladomat, F. *et al.*, *Phytochemistry*, 1996, **43**, 1379-1384 (*6-Hydroxycrinamine, pmr, cmr*)

Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591-6600 (*cd*)

Nishimata, T. *et al.*, *J.O.C.*, 1998, **63**, 7586-7587 (*synth*)

Baldwin, S.W. *et al.*, *Org. Lett.*, 2000, **2**, 99-102 (*synth*)

Hohmann, J. *et al.*, *Planta Med.*, 2002, **68**, 454-456 (*isol, pmr, cmr*)


Abdel-Halim, O.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1119-1124 (*Yemenines*)

Zhang, F.-M. *et al.*, *Tetrahedron*, 2006, **62**, 9446-9455 (*synth*)

Haematopodin

[151964-21-5]

H-7



C₁₃H₁₂N₂O₃ 244.249

Alkaloid from the mushroom *Mycena haematopus* (blood mycena). Black cryst. +1 H₂O (CH₂Cl₂/MeOH). Sol. MeOH, CHCl₃. [α]_D²³ +571 (c, 0.14 in MeOH). Mp >350°. Artifact formed during workup. λ_{\max} 208 (ε 9900); 242 (ε 18600); 344 (ε 10700); 514 (ε 890) (MeOH).


Baumann, C. *et al.*, *Angew. Chem., Int. Ed.*, 1993, **32**, 1087-1089 (*isol, uv, ir, pmr, cmr, cd, cryst struct*)

Hopmann, C. *et al.*, *Annalen*, 1996, 1117-1120 (*synth*)

Haematopodin B

[1032121-80-4]

H-8



C₁₃H₁₃N₃O₂ 243.265


Alkaloid from *Mycena haematopus* (blood mycena). Red solid. [α]_D²⁵ +490 (c, 0.001 in H₂O). λ_{\max} 242 (log ε 4.32); 357 (log ε 4.06); 507 (log ε 3.37) (H₂O).

Peters, S. *et al.*, *Eur. J. Org. Chem.*, 2008, 319-323 (*isol, cd, pmr, cmr, ms*)

Haemultine

α -Demethoxyhaemanthamine

H-9



C₁₆H₁₇NO₃ 271.315

The struct α -Demethoxyhaemanthamine (illus.) which has been synthesised, was suggested for Haemultine. Since the synthetic and natural alkaloids differ, it has since been proposed that Haemultine was a mixt. with the Δ^2 -isomer (see below). Alkaloid from the bulbs of *Haemanthus multiflorus* (Amaryllidaceae). Prisms (Me₂CO). Mp 174-175° (nat.) Mp 195-195.5° (synthetic). [α]_D²⁵ +10 (c, 0.19 in CHCl₃) (synthetic).

Hydroiodide: Mp 102° (nat.) Mp 140-150° (hydrate) (synthetic).

Picrate:

Prisms (H₂O). Mp 208-210° dec. (nat.) Mp 260-264° (synthetic).

7-Methoxy, stereoisomer: **7-Methoxycrinan-11-ol**

[117091-34-6]

C₁₇H₂₁NO₄ 303.357

Minor alkaloid from rhizomes of *Crinum americanum*. Stereochem. undetermined. May be identical with Fiancine below.

ar-Methoxy: **Fiancine**

C₁₇H₁₉NO₄ 301.341

Isol. from bulbs of *Narcissus tazetta* and *Hippeastrum aulicum* var. *robustum* (Amaryllidaceae). Mp 239-241° dec. [α]_D +75 (c, 0.2 in CHCl₃). Tentative struct.

ar-Methoxy, picrate: Mp 223-225° dec.

Δ^2 -Isomer: β -Demethoxyhaemanthamine

C₁₆H₁₇NO₃ 271.315

Synthetic. Prisms (Me₂CO). Mp 201-202°. [α]_D²⁴ +199 (c, 0.17 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 2462-2465; 1958, **91**, 1965-1967 (*Haemultine, Fiancine, isol*)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1958, **45**, 262-263; 1960, **47**, 109 (*isol*)


Fales, H.M. *et al.*, *J.O.C.*, 1961, **26**, 1617-1621 (*synth, uv, struct*)

Trimino, Z. *et al.*, *Rev. Cubana Quim.*, 1987, **3**, 67; *CA*, **109**, 187251s (*7-Methoxycrinan-11-ol*)

Hainanensine

[79233-02-6]

H-10



C₁₇H₁₇NO₄ 299.326

Alkaloid from *Cephalotaxus hainanensis* (Cephalotaxaceae). Possesses antitumour props. Poorly sol. hexane. Mp 240-244°. [α]_D 0 (MeOH).

Sun, N.-C. *et al.*, *Yaoxue Tongbao*, 1980, **15**, 39; *CA*, **95**, 175614s (*isol, uv, ir, pmr, ms, struct*)


Sun, N.-J. *et al.*, *Yaoxue Xuebao*, 1981, **16**, 24; *CA*, **95**, 175622t (*isol, struct*)

Huang, L. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 157 (*rev*)

Hakurirodine

H-11

16,28-Secosolanida-5,22(28)-diene-3,12,16-triol, 9CI
[56857-49-9]



$C_{27}H_{43}NO_3$ 429.642

Isol. from *Veratrum grandiflorum* (Liliaceae). Key intermed. in biosynth. of Rubijervine, R-144. Needles (Me₂CO). Mp 194-197°. $[\alpha]_D^{19}$ -163 (c, 0.71 in CHCl₃). Undergoes acetylation with ring-opening to give a tetraacetyl deriv.

22R,28-Dihydro: Baikeine

[38636-84-9]

$C_{27}H_{45}NO_3$ 431.657

Alkaloid from *Veratrum grandiflorum* (Liliaceae). Mp 153-153.5°. $[\alpha]_D$ -97.9 (CHCl₃). pK_a 9.65.

22R,28-Dihydro, N-Ac: Mp 141-143°.

$[\alpha]_D$ -63.3.

22R,28-Dihydro, 3-Ac: Baikeidine

[38636-85-0]

$C_{29}H_{47}NO_4$ 473.695

Alkaloid from *Veratrum grandiflorum* (Liliaceae). Not obt. pure.


Itô, S. *et al.*, *Tet. Lett.*, 1972, 2961 (*isol, ms, pmr, Baikeine*)

Kaneko, K. *et al.*, *Phytochemistry*, 1975, **14**, 1295 (*isol, ms, ir, uv, pmr, struct*)

Halfordinol

H-12

4-[2-(3-Pyridinyl-5-oxazolyl)]phenol, 9CI. 5-(4-Hydroxyphenyl)-2-(3-pyridyl)oxazole. Aegelenine
[4210-82-6]



$C_{14}H_{10}N_2O_2$ 238.245

Alkaloid from bark of *Halfordia scleroxyla* (Rutaceae). Fine cream needles. Mp 255-256°.

Ac: Mp 167-180°.

Me ether: 3-[5-(4-Methoxyphenyl)-2-oxazolyl]pyridine, 9CI. **Methylhalfordinol**. 2-(3-Pyridyl)-5-(4-methoxyphenyl)oxazole
[33864-03-8]

$C_{15}H_{12}N_2O_2$ 252.272

Alkaloid from aerial parts of *Triphasia trifolia* (limeberry), ripe fruits of *Aegle marmelos* (bael fruit) and the leaves and stems of *Micromelum zeylanicum* (Rutaceae). Needles (C₆H₆/petrol or EtOH). Mp 100-101° (98-99°).

O-(3-Methyl-2-butenyl): **O-Isopentenyl-halfordinol**. 3-[5-[4-(3-Methyl-2-butenyl)oxy]phenyl]-2-oxazolyl]pyridine, 9CI. O-(3,3-Dimethylallyl)halfordinol.

O-Prenylhalfordinol

[17190-80-6]

$C_{19}H_{18}N_2O_2$ 306.363

Alkaloid from *Aeglopsis chevalieri*, *Aegle marmelos* (bael fruit), *Amyris plumieri* and *Amyris texana* (Rutaceae). Cryst. (EtOAc/hexane). Mp 118-119° (109-111°).

O-(3-Methyl-3-butenyl): O-(3-Methyl-3-butenyl)halfordinol

[17190-81-7]

$C_{19}H_{18}N_2O_2$ 306.363

Alkaloid from *Aeglopsis chevalieri* (Rutaceae). Obt. only in admixture with the isomeric O-Isopentenylhalfordinol. Alkaloid not named in the paper.

O-(2,3-Dihydroxy-3-methylbutyl): Halfordine†

[21059-65-4]

$C_{19}H_{20}N_2O_4$ 340.378

Alkaloid from bark of *Halfordia scleroxyla* (Rutaceae). Cream needles (MeOH). Mp 163-164°.

O-(2,3-Dihydroxy-3-methylbutyl), N-Me: N-Methylhalfordinium

$C_{20}H_{23}N_2O_4^+$ 355.413

Quaternary alkaloid from the bark of *Halfordia scleroxyla* (Rutaceae). Fine yellow needles (EtOAc/MeOH) (as chloride). Mp 235° dec. (chloride). Methylated on the pyridine nitrogen.

O-(3-Methyl-2-oxobutyl): Halfordinone

[21144-32-1]

$C_{19}H_{18}N_2O_3$ 322.363

Alkaloid from bark of *Halfordia scleroxyla* (Rutaceae). Needles (Me₂CO/petrol). Mp 132-133°.

O-(3,7-Dimethyl-2,6-octadienyl) (E-): O-Geranylhalfordinol

[89764-17-0]

$C_{24}H_{26}N_2O_2$ 374.482

Alkaloid from aerial parts of *Amyris plumieri* (Rutaceae).

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1964, **17**, 119; 1965, **18**, 1433 (*ir, ms, pmr, isol, struct*)

Dreyer, D.L. *et al.*, *J.O.C.*, 1968, **33**, 3658 (*derivs*)

Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4393 (*synth*)
Manandhar, M.D. *et al.*, *Phytochemistry*, 1978, **17**, 1814 (*derivs*)

Burke, B.A. *et al.*, *Tet. Lett.*, 1978, 2723 (*deriv*)
Sharma, B.R. *et al.*, *Planta Med.*, 1981, **43**, 102 (*Methylhalfordinol*)

Bowen, I.H. *et al.*, *Phytochemistry*, 1982, **21**, 433 (*Methylhalfordinol*)

Philip, S. *et al.*, *Heterocycles*, 1984, **22**, 9 (*O-Geranylhalfordinol*)

Velez, H. *et al.*, *Rev. Cubana Quim.*, 1986, **2**, 22; *CA*, **109**, 20246g (*Methylhalfordinol*)


Dominguez, X.A. *et al.*, *Heterocycles*, 1988, **27**, 35 (*derivs*)

Molina, P. *et al.*, *Heterocycles*, 1993, **36**, 2255 (*synth, Methylhalfordinol*)

Halichlorine

H-13

[178176-75-5]



Absolute configuration

$C_{23}H_{32}ClNO_3$ 405.963

Alkaloid from the marine sponge *Halichondria okadai*. Shows antiinflammatory, antimetastatic and immunosuppressive props. Mp 183.5-185.5°. $[\alpha]_D$ +240.7 (c, 0.54 in MeOH).

Kuramoto, M. *et al.*, *Tet. Lett.*, 1996, **37**, 3867-3870 (*isol, ir, pmr, cmr, struct*)

Arimoto, H. *et al.*, *Tet. Lett.*, 1998, **39**, 861-862 (*abs config*)

Trauner, D. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 3542-3545 (*synth*)

Matsumura, Y. *et al.*, *Org. Lett.*, 2004, **6**, 965-968 (*synth*)


Zhang, H.-L. *et al.*, *J.O.C.*, 2005, **70**, 4954-4961 (*synth*)

Andrade, R.B. *et al.*, *Org. Lett.*, 2005, **7**, 5733-5735 (*synth*)

Halichonadin A

H-14

[847605-76-9]



$C_{31}H_{52}N_2O$ 468.765


Constit. of a *Halichondria* sp. Cryst. Mp 114-116°. $[\alpha]_D^{24}$ -30 (c, 1 in CHCl₃).

Ishiyama, H. *et al.*, *Tetrahedron*, 2005, **61**, 1101-1105 (*isol, pmr, cmr*)

Halichondramide

H-15

[113275-14-2]



$C_{44}H_{60}N_4O_{12}$ 836.978

Macrolide antibiotic. Related to Kabiramide C in K-2. Prod. by sponge *Halichondria* sp., *Haliclona* sp. and *Polycitorella* sp. Antifungal agent, insecticide, toxic to sea urchins. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 66-68°. $[\alpha]_D$ -100.7 (c, 0.42 in MeOH). λ_{max} 232 (ε 23400) (MeOH) (Derep).

5,6-Dihydro: Dihydrohalichondramide

[116302-29-5]

$C_{44}H_{62}N_4O_{12}$ 838.993

Isol. from *Halichondria* sp. and *Hexabranthus sanguineus*. Glass. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D$ -69.7 (c, 1.68 in MeOH). λ_{max} 247 (ε 32000) (MeOH) (Berdy). λ_{max} 245 (ε 26800) (EtOH) (Berdy).

5,6-Dihydro, 7R-alcohol: Tetrahydrohalichondramide

[116302-33-1]

$C_{44}H_{64}N_4O_{12}$ 841.009

Isol. from *Halichondria* sp. Glass. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D$ -27 (c, 0.54 in MeOH). λ_{max} 247 (ε

34000) (MeOH) (Berdy).

5S-Hydroxy, 5,6-dihydro: Jaspisamide A
[149420-76-8]
C₄₄H₆₂N₄O₁₃ 854.993
Isol. from the Okinawan marine sponge *Jaspis* sp. Cytotoxic. Solid. [α]_D¹⁷ -51 (c, 0.13 in MeOH). λ_{max} 244 (ε 24000) (MeOH) (Berdy).

22S-Hydroxy: Jaspisamide B
[149420-77-9]
C₄₄H₆₀N₄O₁₃ 852.977
Isol. from *Jaspis* sp. Cytotoxic. Solid. [α]_D¹⁹ -112 (c, 0.19 in MeOH). λ_{max} 230 (ε 36000) (MeOH) (Berdy).

5S-Amino, 5,6-dihydro: Halishigamide A
C₄₄H₆₃N₅O₁₂ 854.008
Metab. from the Okinawan marine sponge *Halichondria* sp. Cytotoxic. Antifungal agent. Amorph. solid. [α]_D²⁵ +38 (c, 0.51 in MeOH). λ_{max} 263 (ε 15000) (MeOH).

5Z-Isomer: Isohalichondramide
[116302-30-8]
C₄₄H₆₀N₄O₁₂ 836.978
Minor prod. from *Halichondria* sp. Ichthyotoxin, feeding deterrent and sporicide. Oil. Sol. MeOH, CHCl₃; poorly sol. H₂O.

33R-Methyl: Jaspisamide C
[149420-78-0]
C₄₅H₆₂N₄O₁₂ 851.004
Isol. from *Jaspis* sp. Cytotoxic. Solid. [α]_D¹⁹ -76 (c, 0.37 in MeOH). λ_{max} 230 (ε 32000) (MeOH) (Berdy).

33R-Methyl, 5,6-dihydro: 33-Methyl-dihydrohalichondramide. Dihydrojaspisamide
[119147-13-6]
C₄₅H₆₄N₄O₁₂ 853.02
Isol. from the egg masses of *Hexabranchus* sp. Cytotoxic. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²³ -53 (c, 0.5 in CHCl₃). λ_{max} 242 (ε 24600) (MeOH).


Kernan, M.R. *et al.*, *J.O.C.*, 1988, **53**, 5014-5020 (*isol, pmr, cmr*)

Matsunaga, S. *et al.*, *J.O.C.*, 1989, **54**, 1360-1363 (*33-Methyl-dihydrohalichondramide*)

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 787-791 (*Jaspisamides A,B,C*)

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 150-154 (*Halishigamide*)

Haliclamine B **H-16**
[126622-63-7]




C₃₁H₅₀N₂ 450.749
Alkaloid from an unidentified marine sponge of the genus *Haliclona*. Cytotoxic agent. Oil.

14,15-Dihydro: Haliclamine A
[126622-64-8]
C₃₁H₅₂N₂ 452.765
Alkaloid from *Haliclona* sp. Oil.

Fusetani, N. *et al.*, *Tet. Lett.*, 1989, **30**, 6891-6894 (*isol, ir, pmr, cmr, ms, struct*)

Morimoto, Y. *et al.*, *Tetrahedron*, 1998, **54**, 12197-12214 (*synth*)
Baldwin, J.E. *et al.*, *Tet. Lett.*, 2000, **41**, 733-736 (*synth*)
Michelliza, S. *et al.*, *J.O.C.*, 2002, **67**, 6474-6478 (*Haliclamine A, synth*)

Haliclamine C **H-17**



m = 9, n = 11

C₃₀H₅₄N₂ 442.77
Alkaloid from the Arctic sponge *Haliclona viscosa*. CAS no. not found in CA 142.


Volk, C.A. *et al.*, *Eur. J. Org. Chem.*, 2004, 3154-3158 (*isol, pmr, cmr, ms*)

Haliclamine D **H-18**
As Haliclamine C, H-17 with m = 10, n = 11

C₃₁H₅₆N₂ 456.797
Alkaloid from the Arctic sponge *Haliclona viscosa*. CAS no. not found in CA 142.

Volk, C.A. *et al.*, *Eur. J. Org. Chem.*, 2004, 3154-3158 (*isol, pmr, cmr, ms*)

Haliclونacyclamine A **H-19**
[179733-14-3]



Relative configuration

C₃₂H₅₆N₂ 468.808
Alkaloid from the tropical marine sponge *Haliclona* sp. Exhibits pronounced cytotoxic, antibacterial and antifungal activity. Needles. Mp 149-150°. [α]_D -3.4 (c, 1.21 in CH₂Cl₂).

Methodide (1:2):
Yellow needles (MeOH/EtOAc). Mp 196-197° dec.

15,16-Dihydro: Haliclونacyclamine C
C₃₂H₅₈N₂ 470.824
From *Haliclona* sp. Gum. [α]_D +4.8 (c, 0.3 in CH₂Cl₂).

25,26-Dihydro, 27,28-didehydro: Haliclونacyclamine B
[179733-15-4]
C₃₂H₅₆N₂ 468.808
From *Haliclona* sp. Exhibits pronounced cytotoxic, antibacterial and antifungal activity. Needles. Mp 145-146°. [α]_D +3.4 (c, 0.55 in CH₂Cl₂). Struct. revised in 1998.

25,26-Dihydro, 29,30-didehydro, methio-


dide (1:2):
Yellow needles (MeOH/EtOAc). Mp 170-175° dec.

15,16,25,26-Tetrahydro, 27,28-didehydro: Haliclونacyclamine D
C₃₂H₅₈N₂ 470.824
From *Haliclona* sp. Gum. [α]_D +16.1 (c, 0.3 in CH₂Cl₂).

Charan, R.D. *et al.*, *Tetrahedron*, 1996, **52**, 9111 (*isol, pmr, cmr, cryst struct*)

Clark, R.J. *et al.*, *Tetrahedron*, 1998, **54**, 8811-8826 (*isol, pmr, cmr, cryst struct*)

Haliclونamide B **H-20**
[248270-62-4]



C₄₀H₅₂N₈O₉ 788.899
Isol. from the marine sponge *Haliclona* sp. [α]_D²⁴ -1.01.


O-(3-Methyl-2-butenyl): Haliclونamide A
[350593-22-5]
C₄₅H₆₀N₈O₉ 857.017
Isol. from a *Haliclona* sp. [α]_D²⁴ -0.23.

O-(2-Hydroxy-3-methyl-3-butenyl): Haliclونamide C
[489446-55-1]
C₄₅H₆₀N₈O₁₀ 873.017
Isol. from a *Haliclona* sp.

Guan, L.L. *et al.*, *Biochem. Biophys. Res. Commun.*, 2001, **283**, 976-981 (*isol, ir, pmr, cmr*)

Sera, Y. *et al.*, *Mar. Biotechnol.*, 2002, **4**, 441-446 (*isol*)

Haliclونamide D **H-21**
[489446-56-2]



C₄₀H₅₄N₈O₁₀ 806.914

Isol. from a *Haliclona* sp.

4'-O-(3-Methyl-2-butenyl): *Haliclona*-amide **E**

[489446-57-3]


C₄₅H₆₂N₈O₁₀ 875.032

Isol. from a *Haliclona* sp.

Sera, Y. *et al.*, *Mar. Biotechnol.*, 2002, **4**, 441-446 (*isol*)

Halicyclamine A

[156280-95-4]



C₃₂H₅₀N₂ 462.76

Alkaloid from the marine sponges *Amphimedon* sp. and *Haliclona* sp. Cytotoxic. Yellowish solid. [α]_D²⁵ -7.3 (c, 0.72 in CH₂Cl₂). [α]_D²⁵ -24 (c, 0.1 in MeOH). λ_{max} 236 (MeOH).

6,9,10,N-Dehydro: *Tetradehydrohalicyclamine A*

C₃₂H₄₇N₂⁺ 459.737

Quaternary alkaloid from *Amphimedon* sp. Cytotoxic. Amorph. solid (as TFA salt). [α]_D²⁵ -14.7 (c, 0.1 in MeOH) (TFA salt). λ_{max} 232; 273 (MeOH) (TFA salt).

22ξ-Hydroxy: *22-Hydroxyhalicyclamine A*

C₃₂H₅₀N₂O 478.76

Alkaloid from *Amphimedon* sp. Amorph. solid. [α]_D²⁵ +21 (c, 0.1 in MeOH). λ_{max} 234; 273 (MeOH).


Jaspars, M. *et al.*, *J.O.C.*, 1994, **59**, 3253-3255 (*isol*, *ir*, *pmr*, *cmr*, *ms*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (*rev*)

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1758-1760 (*isol*, *pmr*, *cmr*)

Halicyclamine B

[185559-26-6]



Relative Configuration

C₂₆H₄₂N₂ 382.631

Alkaloid from the marine sponge *Xestospongia* sp. [α]_D -143.5.


Harrison, B. *et al.*, *Tet. Lett.*, 1996, **37**, 9151

(*isol*, *pmr*, *cmr*, *cryst struct*)

Haligramide A

[282096-24-6]

H-24



C₃₇H₄₉N₇O₆S₃ 784.035

Isol. from the sponge *Haliclona nigra*.

Amorph. powder. [α]_D -36.5 (c, 0.1 in MeOH). λ_{max} 206 (log ε 4.35); 242 (log ε 4.11); 254 (log ε 4.15); 260 (log ε 3.99) (EtOH).

36-S-Oxide: *Haligramide B*

[282096-25-7]

C₃₇H₄₉N₇O₇S₃ 800.035

Isol. from *Haliclona nigra*. Amorph. powder. [α]_D -32.5 (c, 0.09 in MeOH). λ_{max} 206 (log ε 4.26); 240 (log ε 3.63) (EtOH).

21,36-S,S-Dioxide: *Waiakeamide*

[179667-64-2]

C₃₇H₄₉N₇O₈S₃ 816.034

Cyclic hexapeptide. Isol. from the sponge *Ircinia dendroides* and a Palauan *Haliclona* sp. Amorph. solid. [α]_D -54 (c, 0.5 in MeOH). λ_{max} 206 (ε 13000); 239 (ε 4400) (MeOH).

21,21,36-S,S,S-Trioxide: *Waiakeamide*

21-sulfone

[518315-43-0]

C₃₇H₄₉N₇O₉S₃ 832.034

Isol. from a Palauan *Haliclona* sp. Amorph. solid. [α]_D²⁶ -108 (c, 0.16 in CHCl₃).

Mau, C.M.S. *et al.*, *J.O.C.*, 1996, **61**, 6302-6304 (*Waiakeamide*)


Rashid, M.A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 956-959 (*Haligramides A,B*)

Fujii, K. *et al.*, *Tetrahedron*, 2002, **58**, 6873-6879 (*Waiakeamide*, *abs config*)

Sera, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 719-721 (*Waiakeamide*, *Waiakeamide 21-sulfone*)

Halimedin

H-25




C₁₀H₁₆N₆O 236.276

First example of a naturally occurring 1,3,5-triazine. Alkaloid from the alga *Halimeda xishaensis*. Cryst. Mp 164-165°.

Su, J.-Y. *et al.*, *Phytochemistry*, 1998, **48**, 583-584 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)

Halipeptin A

H-26



R = -CH₂CH₂OH

Absolute Configuration

C₃₁H₅₄N₄O₇S 626.856

Struct. revised in 2002. MF originally given as C₃₁H₅₄N₄O₉. Isol. from the sponge *Haliclona* sp. Potent antiinflammatory agent. Amorph. solid. [α]_D -16.6 (c, 0.03 in CHCl₃).

O-De-Me: *Halipeptin B*

C₃₀H₅₂N₄O₇S 612.829

Isol. from *Haliclona* sp. Amorph. solid. [α]_D -22.7 (c, 0.002 in CHCl₃).

5'-Deoxy: *Halipeptin D*

[866555-47-7]

C₃₁H₅₄N₄O₆S 610.857

Isol. from the sponge *Leiosella cf. arenifibrosa*. Viscous oil. [α]_D²⁵ -26 (c, 0.2 in CHCl₃).

Randazzo, A. *et al.*, *J.A.C.S.*, 2001, **123**, 10870-10876 (*isol*, *pmr*, *cmr*, *N-15 nmr*)

Monica, C.D. *et al.*, *Tet. Lett.*, 2002, **43**, 5707-5710 (*struct*)

Nicolaou, K.C. *et al.*, *Chem. Eur. J.*, 2005, **11**, 6197-6211 (*Halipeptin D*)

Yu, S. *et al.*, *Chem. Eur. J.*, 2006, **12**, 6572-6584 (*synth*)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 2006, **128**, 4460-4470 (*synth*)

Hara, S. *et al.*, *Tet. Lett.*, 2006, **47**, 1081-1085 (*synth*)

Halipeptin C

H-27

As Halipeptin A, H-26 with

R = CH₃

C₃₀H₅₂N₄O₆S 596.83

Isol. from the sponge *Haliclona* sp. Amorph. solid. [α]_D -30 (c, 0.3 in CHCl₃).


Monica, C.D. *et al.*, *Tet. Lett.*, 2002, **43**, 5707-5710 (*isol*, *pmr*, *cmr*)

Yu, S. *et al.*, *Chem. Eur. J.*, 2006, **12**, 6572-6584 (*synth*)

Halishigamide B

H-28

[185847-12-5]



C₄₃H₆₀N₄O₁₃ 840.966


Metab. from the Okinawan marine sponge

Halichondria sp. Exhibits weak cytotoxicity and modest antifungal activity. Amorph. solid. $[\alpha]_D^{25}$ -72 (c, 0.06 in MeOH). λ_{\max} 231 (ϵ 66000); 260 (sh) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 150-154 (*isol, uv, ir, pmr, cmr, struct*)

Halishigamide C H-29

[185847-13-6]




$C_{44}H_{64}N_4O_{14}$ 873.008

Metab. from the Okinawan marine sponge *Halichondria* sp. Exhibits weak cytotoxicity and modest antifungal activity. Amorph. solid. $[\alpha]_D^{27}$ -70 (c, 0.12 in $CHCl_3$). λ_{\max} 231 (ϵ 56000) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 150-154 (*isol, uv, ir, pmr, cmr, struct*)

Halishigamide D H-30

[185847-14-7]




$C_{44}H_{64}N_4O_{14}$ 873.008

Metab. from the Okinawan marine sponge *Halichondria* sp. Exhibits weak cytotoxicity and modest antifungal activity. Amorph. solid. $[\alpha]_D^{25}$ -88 (c, 0.03 in MeOH). λ_{\max} 235 (ϵ 15000) (MeOH).

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 150-154 (*isol, uv, ir, pmr, cmr, struct*)

Halifulfate 1 H-31

[116302-38-6]



$C_{31}H_{48}O_6S$ 548.783

Constit. of a sponge (*Halichondriidae*). Active against gram-positive bacteria and *Candida albicans*. Phospholipase A2 and serine protease inhibitor. Needles. $[\alpha]_D$ -27.3 (c, 0.01 in MeOH).

1,7,9-Trimethylguaninium salt: [201667-10-9]

$C_{39}H_{59}N_5O_7S$ 741.99


Isol. from the sponge *Coscinoderma mathewsi*. Cryst. Mp 165-170°. λ_{\max} 260 ($\log \epsilon$ 4.04); 287 ($\log \epsilon$ 3.96) (MeOH).

Kernan, M.R. *et al.*, *J.O.C.*, 1988, **53**, 4574-4578 (*isol, pmr, cmr, activity*)

Kimura, J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 248-250; 862 (*trimethylguaninium salt*)

Singh, M. *et al.*, *Planta Med.*, 1999, **65**, 2-8 (*rev*)

Halitoxin H-32




Toxin from the sponges *Haliclona rubens*, *Haliclona viridis*, *Haliclona erina*, *Amphimedon compressa* and *Amphimedon viridis*. Neurotoxin. Haemolytic, ichthyotoxic and antimitotic agent. Neuromuscular blocker. λ_{\max} 267 ; 273 (EtOH).

► LD₅₀ (mus, ipr) 2-5 mg/kg.

[54990-72-6]

Schmitz, F.J. *et al.*, *J.O.C.*, 1978, **43**, 3916-3922 (*isol, uv, pmr, cmr*)

Halitulín H-33



$C_{35}H_{40}N_4O_4$ 580.725

(S)-form [221367-90-4]

Alkaloid from the marine sponge *Haliclona tulearensis*. Cytotoxic agent. Orange foaming oil. $[\alpha]_D$ +7.5 (c, 2.8 in MeOH). λ_{\max} 212 (ϵ 29200); 252 (ϵ 31600); 364 (ϵ 4400) (MeOH). λ_{\max} 214 (ϵ 24700); 264 (ϵ 14800); 350 (ϵ 4650) (MeOH/NaOH) (Berdy).


Kashman, Y. *et al.*, *Tet. Lett.*, 1999, **40**, 997-1000 (*isol, uv, ir, pmr, cmr, ms*)

Banwell, M.G. *et al.*, *J.C.S. Perkin 1*, 2002, 1340-1343 (*synth, struct*)

Heinrich, M.R. *et al.*, *Tetrahedron*, 2003, **59**, 9239-9247 (*synth, abs config*)

Hallacridone H-34

2-Acetyl-5-hydroxy-11-methylfuro[2,3-c]acridin-6(11H)-one, 9CI
[109897-77-0]



$C_{18}H_{13}NO_4$ 307.305

Struct. revised in 1989. Minor alkaloid from *Ruta graveolens* (rue) tissue cultures (Rutaceae). Yellow needles ($CHCl_3$). Mp 295-298°.


Baumert, A. *et al.*, *Pharmazie*, 1987, **42**, 67; *CA*, **107**, 93537w

Baumert, A. *et al.*, *Fitoterapia*, 1988, **59**, 83; *CA*, **109**, 226730u

Reisch, J. *et al.*, *J.C.S. Perkin 1*, 1989, 1047 (*synth, ir, struct*)

Halocyamine A H-35

[122548-03-2]




$C_{27}H_{28}BrN_7O_5$ 610.466

Oligopeptide antibiotic. Isol. from the haemocytes of the ascidian *Halocynthia roretzi*. Active against several bacteria and yeasts. Sol. MeOH, Me₂CO. $[\alpha]_D^{25}$ +5.2 (c, 0.5 in MeOH).

Azumi, K. *et al.*, *Biochemistry*, 1990, **29**, 159-165 (*isol*)

Halocyamine B H-36

[122548-04-3]




$C_{29}H_{32}BrN_7O_6$ 654.519

Oligopeptide antibiotic. Isol. from the haemocytes of the ascidian *Halocynthia roretzi*. Active against several bacteria and yeasts. Sol. MeOH, Me₂CO. $[\alpha]_D^{25}$ +63.1 (c, 0.5 in MeOH). λ_{\max} 203 (E1%/1cm 785); 232 (E1%/1cm 428); 282 (E1%/1cm 276) (MeOH) (Berdy).

Azumi, K. *et al.*, *Biochemistry*, 1990, **29**, 159-165 (*isol*)

Halosaline H-37

α -Propyl-2-piperidineethanol, 9CI. 2-(2-Hydroxypropyl)piperidine
[171019-32-2 (2S,2'S-form)]

(2*R*,2'*R*)-formC₁₀H₂₁NO 171.282**(2*R*,2'*R*)-form** [26648-71-5]Alkaloid from *Haloxylon salicornicum*. [α]_D²⁵ +3.3 (c, 1 in CHCl₃). [α]_D²⁵ -19.5 (c, 0.6 in EtOH).

Ketone: 1-(2-Piperidinyl)-2-pentanone. 2-(2-Oxopentyl)piperidine. **Haloxynine** [157070-30-9] [157070-29-6 (*S*-form)] C₁₀H₁₉NO 169.266 Alkaloid from *Haloxylon salicornicum*. [α]_D²² -21 (c, 1.8 in CHCl₃).

(2*S*,2'*R*)-form [274923-10-3]Alkaloid from *Andrachne aspera*. Mp 32°. [α]_D²⁵ -8 (c, 3.7 in MeOH).**(2*ξ*,2'*ξ*)-form**


N-Me: 1-(*N*-Methyl-2-piperidinyl)-2-pentanol. 2-(2-Hydroxypentyl)-*N*-methylpiperidine [145224-71-1] C₁₁H₂₃NO 185.309 Alkaloid from leaves of *Sedum lancerottense*. Identified by gc-ms.

Ketone: [91016-37-4]C₁₀H₁₉NO 169.266Alkaloid from leaves of *Sedum brissemoreti* and *Sedum lancerottense*.

Ketone, N-Me: 1-(*N*-Methyl-2-piperidinyl)-2-pentanone. *N*-Methyl-2-(2-oxopentyl)piperidine [145224-70-0] C₁₁H₂₁NO 183.293

Alkaloid from leaves of *Sedum fusiforme*, *Sedum lancerottense*, *Sedum nudum*, and from whole plants of *Pratia nummularia*. Identified by gc-ms.Michel, K.H. *et al.*, *Acta Pharm. Suec.*, 1967, **4**, 97-116 (*isol*, *synth*)Michel, K.H. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 3479 (*abs config*)Craig, J.C. *et al.*, *Tetrahedron*, 1978, **34**, 501 (*ord*, *abs config*)Stevens, J.F. *et al.*, *Phytochemistry*, 1992, **31**, 3917-3924 (*derivs*)Louis, C. *et al.*, *Can. J. Chem.*, 1994, **72**, 1347-1350 (*synth*)Ho, L.K. *et al.*, *Planta Med.*, 1995, **61**, 567 (*deriv*)Stragies, R. *et al.*, *Tetrahedron*, 1999, **55**, 8179-8188 (*synth*)Mill, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 762-764 (*isol*, *synth*)Lesma, G. *et al.*, *Tetrahedron*, 2004, **60**, 6437-6442 (*synth*)El-Shazly, A.M. *et al.*, *Pharmazie*, 2005, **60**, 949-952 (*Haloxynine*)**Haloxine****H-38**


Dodecahydro-5-(1-oxobutyl)-6H-pyrido[2,1-f][1,6]naphthyridin-6-one. 9CI [16812-86-5]



Relative Configuration

C₁₆H₂₆N₂O₂ 278.394Alkaloid from *Haloxylon salicornicum*. Cryst. (as hydrochloride). Mp 193.5-196° (hydrochloride).Michel, K.H. *et al.*, *Acta Pharm. Suec.*, 1967, **4**, 97-116 (*isol*, *struct*)Nilsson, B. *et al.*, *Acta Chem. Scand., Ser. B*, 1968, **24**, 252-258 (*cryst struct*)**Haloxyline B****H-39**


3-Hydroxy-1-(22-hydroxy-2,4-docosadienyl)piperidine



Absolute Configuration

C₂₇H₄₉NO₃ 435.689Alkaloid from *Haloxylon salicornicum*. Cryst. Mp 142-143°. [α]_D²⁰ -21.5 (c, 1 in CHCl₃). λ_{max} 262 (log ε 4) (MeOH).

3-Deoxy: 1-(22-Hydroxy-2,4-docosadienyl)piperidine. **Haloxyline A** C₂₇H₄₉NO₂ 419.69

Alkaloid from *Haloxylon salicornicum*. Cryst. Mp 161-162°. λ_{max} 262 (log ε 3.92) (MeOH).Ferheen, S. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 570-572 (*isol*, *pmr*, *cmr*, *ms*)**Hamacanthin A****H-40**

(R)-form

C₂₀H₁₄Br₂N₄O 486.165Abs. configs. not certain. Hamacanthin A itself has now been shown to have *S*-config. but the minor congeners appear to be *R*-.**(R)-form** [354538-21-9]3*R*,4-Dihydro: **cis-3,4-Dihydrohamacanthin A**

[264624-42-2]

C₂₀H₁₆Br₂N₄O 488.181Alkaloid from the sponge *Rhaphisia lacazei*. Abs. config. not confirmed.3*S*,4-Dihydro: **trans-3,4-Dihydrohamacanthin A**


[264624-43-3]

Alkaloid from *Rhaphisia lacazei* and *Spongosorites* sp. [α]_D²³ +16 (c, 1.1 in MeOH).6'-Debromo: **(R)-6'-Debromohamacanthin A**C₁₂H₁₅BrN₄O 311.181Alkaloid from the sponge *Spongosorites* sp. Amorph. yellow powder. [α]_D²³ -

91 (c, 0.2 in MeOH).

6'-Debromo, 3*S*,4-dihydro: **trans-6'-Debromo-3,4-dihydrohamacanthin A** [264624-44-4]C₂₀H₁₇BrN₄O 409.285Alkaloid from *Rhaphisia lacazei* and *Spongosorites* sp. Yellow powder. [α]_D²⁵ +31 (c, 0.38 in MeOH).6''-Debromo: **(R)-6''-Debromohamacanthin A**C₁₂H₁₅BrN₄O 311.181Alkaloid from the sponge *Spongosorites* sp. Amorph. yellow powder. [α]_D²⁵ -76 (c, 0.05 in MeOH).6''-Debromo, 3*S*,4-dihydro: **trans-6''-Debromo-3,4-dihydrohamacanthin A** [264624-45-5]C₂₀H₁₇BrN₄O 409.285Alkaloid from *Rhaphisia lacazei* and *Spongosorites* sp. Yellow powder. [α]_D²⁵ +34 (c, 0.15 in MeOH).**(S)-form** [160098-92-0]Alkaloid from *Hamacantha* sp. and *Spongosorites* sp. Shows antimicrobial activity. Pale yellow powder. [α]_D²⁴ +84 (c, 0.1 in MeOH). λ_{max} 219 (ε 76500); 280 (ε 20600); 325 (ε 13300) (MeOH).

Bis(debromo): (S)-6',6''-Didebromohamacanthin A

C₂₀H₁₆N₄O 328.373Alkaloid from *Spongosorites* sp. Amorph. yellow powder. [α]_D²⁵ +59 (c, 0.72 in MeOH). λ_{max} 221 (log ε 4.17); 268 (log ε 3.92); 330 (log ε 3.73) (MeOH).Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1437-1441 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)Casapullo, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 447-451 (3,4-Dihydrohamacanthins)Jiang, B. *et al.*, *J. O. C.*, 2001, **66**, 4865-4869; 7560 (*synth*, *abs config*)Miyake, F.Y. *et al.*, *Org. Lett.*, 2002, **4**, 941-943 (*synth*)Bao, B. *et al.*, *J. Nat. Prod.*, 2005, **68**, 711-715 (*isol*, *pmr*, *cmr*, *abs config*)Garg, N.K. *et al.*, *Tet. Lett.*, 2005, **46**, 2423-2426 (3,4-Dihydrohamacanthins, *synth*)Kouko, T. *et al.*, *Tetrahedron*, 2005, **61**, 2309-2318 (*synth*)Bao, B. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2-8 (Didebromohamacanthin A)Guinchard, X. *et al.*, *Org. Lett.*, 2007, **9**, 3761-3764 (*synth*)**Hamacanthin B****H-41**

(R)-form

C₂₀H₁₄Br₂N₄O 486.165**(R)-form**3*S*,4-Dihydro: **cis-3,4-Dihydrohamacanthin B**

[264624-39-7]

C₂₀H₁₆Br₂N₄O 488.181Alkaloid from the sponges *Rhaphisia lacazei* and *Spongosorites* sp. Amorph. yellow powder. [α]_D²¹ +101 (c, 0.8 in

MeOH).

6'-Debromo: (R)-6'-Debromohamacanthin BC₂₀H₁₅BrN₄O 407.269Alkaloid from *Spongosorites* sp. Amorph. yellow powder. $[\alpha]_D^{25}$ -288 (c, 0.4 in MeOH). λ_{\max} 227 (log ϵ 4.03); 267 (log ϵ 3.95); 329 (log ϵ 3.84) (MeOH).**6'-Debromo, 3S,4-dihydro: cis-6'-Debromo-3,4-dihydrohamacanthin B**


[264624-40-0]

C₂₀H₁₇BrN₄O 409.285Alkaloid from *Rhaphisia lacazei* and *Spongosorites* sp. Yellow powder. $[\alpha]_D^{25}$ +80 (c, 0.25 in MeOH).**6''-Debromo: (R)-6''-Debromohamacanthin B**C₂₀H₁₅BrN₄O 407.269Alkaloid from *Spongosorites* sp. Amorph. yellow powder. $[\alpha]_D^{25}$ -83 (c, 0.5 in MeOH). λ_{\max} 230 (log ϵ 4.06); 277 (log ϵ 3.98); 322 (log ϵ 3.78) (MeOH).**6''-Debromo, 3S,4-dihydro: cis-6''-Debromo-3,4-dihydrohamacanthin B**


[264624-41-1]

C₂₀H₁₇BrN₄O 409.285Alkaloid from *Rhaphisia lacazei* and *Spongosorites* sp. Yellow powder. $[\alpha]_D^{25}$ +105 (c, 0.2 in MeOH).**Bis(debromo): (R)-6',6''-Didebromohamacanthin B**C₂₀H₁₆N₄O 328.373Alkaloid from *Spongosorites* sp. Amorph. yellow powder. $[\alpha]_D^{25}$ -288 (c, 0.4 in MeOH). λ_{\max} 227 (log ϵ 4.03); 267 (log ϵ 3.95); 329 (log ϵ 3.84) (MeOH).**Bis(debromo), 3S,4-dihydro: 6',6''-Didebromo-3,4-dihydrohamacanthin B**C₂₀H₁₈N₄O 330.388Alkaloid from *Spongosorites* sp.**(S)-form [160098-93-1]**Alkaloid from a deep-water marine sponge *Hamacantha* sp. Shows significant antimicrobial activity. Pale yellow powder. $[\alpha]_D^{24}$ +172 (c, 0.1 in MeOH). λ_{\max} 219 (ϵ 76500); 280 (ϵ 20600); 325 (ϵ 13300) (MeOH) (Berdy).**6'-Debromo: 6'-Debromohamacanthin B**C₂₀H₁₅BrN₄O 407.269Alkaloid from the sponge *Spongosorites* sp. Amorph. yellow solid. Mp 210-212°. $[\alpha]_D^{25}$ -100 (c, 0.2 in MeOH). λ_{\max} 217 (log ϵ 4); 224 (log ϵ 3.94); 274 (log ϵ 3.46); 328 (log ϵ 3.7) (no solvent reported).Gunasekera, S.P. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1437-1441 (*isol, uv, ir, pmr, cmr, ms, struct*)Casapullo, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 447-451 (*3,4-Dihydrohamacanthins*)Jiang, B. *et al.*, *J.O.C.*, 2002, **67**, 1396-1398 (*synth*)Miyake, F.Y. *et al.*, *Org. Lett.*, 2002, **4**, 941-943 (*synth*)Oh, K.-B. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 4927-4931 (*6'-Debromohamacanthin B*)Kouko, T. *et al.*, *Tetrahedron*, 2005, **61**, 2309-2318 (*synth*)Bao, B. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2-8 (*R-form derivs*)Higuchi, K. *et al.*, *Synthesis*, 2007, 669-674 (*3,4-dihydro*)**Hamigeramide**

H-42

C₁₂H₁₁N₃O₂ 229.238Alkaloid from the sponge *Hamigera hamigera*. Pale yellow powder. λ_{\max} 228; 346 (MeOH).Hassan, W. *et al.*, *Mar. Drugs*, 2004, **2**, 88-100 (*isol, pmr, cmr, ms*)**Hamigeran D**

[255882-09-8]




Absolute Configuration

C₂₁H₂₆BrNO₂ 404.346Isol. from the sponge *Hamigera tarangaensis*. Cytotoxic agent. Pale yellow solid. $[\alpha]_D^{25}$ -47.1 (c, 0.21 in CH₂Cl₂).Wellington, K.D. *et al.*, *J. Nat. Prod.*, 2000, **63**, 79-85 (*isol, pmr, cmr*)Cambie, R.C. *et al.*, *Acta Cryst. C*, 2001, **57**, 958-960 (*cryst struct*)**Hamigeroxalamic acid**

H-44

[2-(4-Hydroxyphenyl)vinyl]oxamic acid


C₁₀H₉NO₄ 207.185Alkaloid from the sponge *Hamigera hamigera*. Pale yellow powder. λ_{\max} 248; 320 (MeOH).*Imine: [2-(4-Hydroxyphenyl)vinylamino]iminoacetic acid. Hamigeramine†*C₁₀H₁₀N₂O₃ 206.201Alkaloid from the sponge *Hamigera hamigera*. Pale yellow powder. λ_{\max} 227; 324 (MeOH).Hassan, W. *et al.*, *Mar. Drugs*, 2004, **2**, 88-100 (*isol, pmr, cmr, ms*)**Hanadamine**

H-45

C₂₁H₂₄N₂O₄ 368.432Struct. unknown. Alkaloid from *Uncaria kawakamii* (Rubiaceae). Mp 187°.Kondo, H. *et al.*, *Yakugaku Zasshi*, 1932, **52**, 528-531; *CA*, **27**, 1345**Hanasanagin**


H-46

N-(3,4-Diguanidinobutanoyl)dopa

C₁₅H₂₃N₇O₅ 381.391Constit. of the mushroom *Isaria japonica*. Antioxidant. Pale yellow oil.Sakakura, A. *et al.*, *Tet. Lett.*, 2005, **46**, 9057-9059 (*isol, pmr, cmr*)**Hauouamine A**

H-47

[496962-05-1]




Absolute Configuration

C₃₂H₂₇NO₄ 489.57Alkaloid from the ascidian *Aplidium haouarianum*. Cytotoxic. Solid. $[\alpha]_D^{20}$ -52 (c, 0.4 in MeOH). Dec. at 170°. λ_{\max} 205 (ϵ 62000); 275 (ϵ 11000) (MeOH).**21-Hydroxy: Hauouamine B**

[496962-06-2]

C₃₂H₂₇NO₅ 505.569Alkaloid from *Aplidium haouarianum*. Solid (as penta-Ac). $[\alpha]_D^{20}$ -27.1 (c, 0.14 in CHCl₃) (penta-Ac). λ_{\max} 204 (ϵ 48100); 238 (ϵ 20300) (MeOH) (penta-Ac).Garrido, L. *et al.*, *J.O.C.*, 2003, **68**, 293-299 (*isol, pmr, cmr, cryst struct*)Jeong, J.H. *et al.*, *Org. Lett.*, 2006, **8**, 2309-2312 (*synth*)Gravel, E. *et al.*, *Chem. Comm.*, 2007, 719-721 (*synth*)Burns, N.Z. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 205-208 (*synth, abs config, biosynth*)Belostotskii, A.M. *et al.*, *J.O.C.*, 2008, **73**, 5723-5731 (*conform*)**Hapalindole A**

H-48

8-Chloro-9-ethenyl-2,6,6a,7,8,9,10,10a-octahydro-10-isocyano-6,6,9-trimethylnaphth[1,2,3-cd]indole, 9CI
[92219-95-9]

Absolute Configuration

C₂₁H₂₃ClN₂ 338.879Numbering systems vary. Isol. from the blue-green alga *Hapalosiphon fontinalis*.

Algicide. Antibacterial and antimycotic agent. Yellow plates (CH₂Cl₂/heptane). Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Mp 160-167° dec. [α]_D²⁵ -78 (c, 1.2 in CH₂Cl₂). λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5800) (MeOH) (Derep).

20,21-Epoxyde: Hapalindole N

[106865-65-0]

C₂₁H₂₃ClN₂O 354.878

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -31.5 (c, 1.7 in CHCl₃). Config. at C-20 not known. λ_{\max} 220 (€ 34000); 273 (€ 5700); 280 (€ 5800); 290 (€ 4500) (MeOH) (Derep).

6a,10a-Didehydro: Hapalindole K

[106865-63-8]

C₂₁H₂₁ClN₂ 336.863

From *Hapalosiphon fontinalis* ATCC 39964. Thick yellow plates (CH₂Cl₂/heptane). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 110-240° dec. [α]_D²⁵ -12.5 (c, 1.8 in CHCl₃). λ_{\max} 226 (€ 19300); 301 (€ 9600) (MeOH) (Derep).

10,10a-Didehydro: Hapalindole I

[101968-76-7]

C₂₁H₂₁ClN₂ 336.863

From *Hapalosiphon fontinalis*. Mp 180° dec. [α]_D²⁵ -12 (c, 0.2 in CH₂Cl₂). λ_{\max} 224 (€ 25800); 240 (sh) (€ 13000); 274 (€ 6700); 322 (€ 13400) (MeOH) (Derep). λ_{\max} 222 ; 224 (€ 25800); 274 (€ 6700); 280 ; 291 ; 322 (€ 13900) (MeOH) (Berdy).

Isothiocyanate: Hapalindole B

[92219-96-0]

C₂₁H₂₃ClN₂S 370.945

Isol. from *Hapalosiphon fontinalis*. Algicide. Antibacterial and antimycotic agent. Oil. Sol. MeOH, CHCl₃, butanol; poorly sol. H₂O, hexane. [α]_D²⁵ -194 (c, 5.1 in CH₂Cl₂). Has -NCS replacing -NC. λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep).

10a β -Hydroxy: Hapalindole V. 10-Hydroxyhapalindole G

[106865-68-3]

C₂₁H₂₃ClN₂O 354.878

From *Hapalosiphon fontinalis* ATCC 39964. No phys. props. reported. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{\max} 220 (€ 34000); 273 (€ 5700); 280 (€ 5800); 290 (€ 4500) (MeOH) (Derep).

Dechloro: Hapalindole J

[106928-26-1]

C₂₁H₂₄N₂ 304.434

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ +54.4 (c, 0.9 in CHCl₃). λ_{\max} 222 (€ 38700); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep).

Dechloro, isothiocyanate: Hapalindole M

[106865-64-9]

C₂₁H₂₄N₂S 336.5

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -83.1 (c, 1.8 in

CHCl₃). λ_{\max} 223 (€ 40000); 281 (€ 7300); 291 (€ 5900) (MeOH) (Derep).

Dechloro, β -hydroxy, isothiocyanate:

Hapalindole O

[106865-66-1]

C₂₁H₂₄N₂OS 352.499

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -106 (c, 2.4 in CHCl₃). λ_{\max} 222 (€ 38000); 279 (€ 7000); 290 (€ 5700) (MeOH) (Derep).

6a-Epimer: Hapalindole H

[101968-75-6]

C₂₁H₂₃ClN₂ 338.879

From *Hapalosiphon fontinalis*. Mp 190-193°. [α]_D²⁵ +152 (c, 4.1 in CH₂Cl₂). λ_{\max} 222 (€ 38700); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep).

9-Epimer: Hapalindole L

[106928-27-2]

C₂₁H₂₃ClN₂ 338.879

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -74 (c, 1.1 in CHCl₃). λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5800) (MeOH) (Derep).

9-Epimer, dechloro: 12-Epihapalindole J

[955020-01-6]

C₂₁H₂₄N₂ 304.434

Isol. from *Fischerella* ATCC 43239. Insecticide. Amorph. solid. λ_{\max} 221 ; 281 (MeOH).

10a-Epimer: Hapalindole G

[102045-13-6]

C₂₁H₂₃ClN₂ 338.879

From *Hapalosiphon fontinalis*. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 185° dec. [α]_D²⁵ -43.9 (c, 0.28 in CH₂Cl₂). λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5800) (MeOH) (Derep).

10a-Epimer, dechloro: Hapalindole U

[106928-30-7]

C₂₁H₂₄N₂ 304.434

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ +12 (c, 0.6 in CH₂Cl₂). λ_{\max} 222 (€ 38700); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep). λ_{\max} 222 ; 282 ; 291 (MeOH) (Berdy).

?-Epimer, 20,21-epoxyde: Hapalindole P

[106928-28-3]

C₂₁H₂₃ClN₂O 354.878

From *Hapalosiphon fontinalis* ATCC 39964. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -16.3 (c, 0.8 in CHCl₃). λ_{\max} 220 (€ 34000); 273 (€ 5700); 280 (€ 5800); 290 (€ 4500) (MeOH) (Derep). λ_{\max} 222 ; 280 ; 291 (MeOH) (Berdy).

6a,9-Diepimer: 12-Epihapalindole H

[174063-68-4]

C₂₁H₂₄ClN₂ 339.887

Alkaloid from *Fischerella ambigua* and *Hapalosiphon laingii*. Sol. MeOH. Mp 187-189°. [α]_D²⁵ +217.3 (c, 0.16 in CH₂Cl₂). λ_{\max} 221 (€ 28140); 282 (€ 5150); 291 (sh) (€ 4130) (MeOH).

9,10a-Diepimer: 12-Epihapalindole G

[173241-59-3]

C₂₁H₂₃ClN₂ 338.879

Isol. from *Hapalosiphon laingii*. Sol. MeOH. λ_{\max} 222 (€ 37220); 280 (€ 8200); 291 (sh) (€ 6640) (MeOH).

Moore, R.E. *et al.*, *J.A.C.S.*, 1984, **106**, 6456-6457 (isol, uv, ir, pmr, cmr, Hapalindole B)

Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 1036-1043 (isol, ir, pmr, cmr, cryst struct)

Muratake, H. *et al.*, *Tetrahedron*, 1990, **46**, 6331-6342; 6351-6360 (*Hapalindoles H, J, M, U, synth*)

Sakagami, M. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1393-1398 (*Hapalindole O, synth*)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1994, **116**, 3125-3126 (*Hapalindole G, synth*)


Klein, D. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1781-1785 (*12-Epihapalindoles G, H*)

Becher, P.G. *et al.*, *Phytochemistry*, 2007, **68**, 2493-2497 (*12-Epihapalindole J*)

Hapalindole E

[101968-73-4]

H-49



C₂₁H₂₃ClN₂ 338.879

Isol. from freshwater cyanobacterium *Hapalosiphon fontinalis*. Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Mp 88-90°. [α]_D²⁵ +25.2 (c, 3.1 in CH₂Cl₂). λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5800) (MeOH) (Derep).

Isothiocyanate: Hapalindole F

[101968-74-5]

C₂₁H₂₃ClN₂S 370.945

From *Hapalosiphon fontinalis*. Mp 176-179°. [α]_D²⁵ +93.2 (c, 0.22 in CH₂Cl₂). Has -NCS replacing -NC. λ_{\max} 222 (€ 38000); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep).

Dechloro: Hapalindole C

[101968-71-2]

C₂₁H₂₄N₂ 304.434

Rel. config. illus. Isol. from cyanobacterium *Hapalosiphon fontinalis*. Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Mp 138-143°. [α]_D²⁵ +76.5 (c, 0.41 in CH₂Cl₂). λ_{\max} 222 (€ 38700); 280 (€ 7000); 291 (€ 5700) (MeOH) (Derep).

Dechloro, isothiocyanate: Hapalindole D

[101968-72-3]

C₂₁H₂₄N₂S 336.5

Isol. from *Hapalosiphon fontinalis*. Yellow-orange prisms (CH₂Cl₂). Mp 105-107°. [α]_D²⁵ +239 (c, 3.1 in CH₂Cl₂). λ_{\max} 223 (€ 40000); 281 (€ 7300); 291 (€ 5900) (MeOH) (Derep).

5'-Epimer: 12-Epihapalindole E

[159249-51-1]

C₂₁H₂₃ClN₂ 338.879

Alkaloid from *Hapalosiphon wel-*

witschii, *Hapalosiphon laingii* and *Westiella intricata*. Ichthyotoxic. $[\alpha]_D^{25} +42.9$ (c, 0.3 in CH_2Cl_2). λ_{max} 221 (€ 28400); 282 (€ 5000); 290 (€ 4480) (MeOH) (Berdy).

5'-Epimer, isothiocyanate: 12-Epihapalindole F isothiocyanate

[159249-53-3]
 $\text{C}_{21}\text{H}_{23}\text{ClN}_2\text{S}$ 370.945

Alkaloid from *Hapalosiphon wel-witschii*. $[\alpha]_D^{25} +102$ (c, 0.5 in CH_2Cl_2). Authors' numbering and nomenclature.

5'-Epimer, dechloro: 12-Epihapalindole C

[159249-52-2]
 $\text{C}_{21}\text{H}_{24}\text{N}_2$ 304.434

Alkaloid from *Hapalosiphon wel-witschii*, *Westiella intricata* and *Hapalosiphon laingii*. Ichthyotoxic. $[\alpha]_D^{25} +10.4$ (c, 0.54 in CH_2Cl_2). λ_{max} 219 (€ 18000); 282 (€ 4800) (MeOH) (Berdy). λ_{max} 222 (€ 26900); 283 (€ 4460); 290 (€ 4060) (EtOH) (Berdy).

5'-Epimer, dechloro, isothiocyanate: 12-Epihapalindole D isothiocyanate

[159249-54-4]
 $\text{C}_{21}\text{H}_{24}\text{N}_2\text{S}$ 336.5

Alkaloid from the *Hapalosiphon wel-witschii*.

1',2'-Diepimer, dechloro, isothiocyanate: Hapalindole Q

[106928-29-4]
 $\text{C}_{21}\text{H}_{24}\text{N}_2\text{S}$ 336.5

From *Hapalosiphon fontinalis*. Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. $[\alpha]_D^{25} +24.1$ (c, 1.1 in CH_2Cl_2). λ_{max} 223 (€ 40000); 281 (€ 7300); 291 (€ 5900) (MeOH) (Derep).

1',2',5'-Triepimer, dechloro: 12-Epihapalindole Q

[173241-60-6]
 $\text{C}_{21}\text{H}_{24}\text{N}_2$ 304.434

Alkaloid from *Hapalosiphon laingii*. Ichthyotoxic. Sol. MeOH. λ_{max} 222 (€ 24160); 283 (€ 4850); 291 (sh) (€ 4300) (MeOH).

Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 1036

(*Hapalindoles*, *isol*, *pmr*, *cmr*, *struct*, *Hapalindole D*, *cryst struct*)

Stratmann, K. *et al.*, *J.A.C.S.*, 1994, **116**, 9935

(*12-Epihapalindole isonitriles*, *12-Epihapalindole E isothiocyanates*)


Klein, D. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1781-1785 (*12-Epihapalindole Q isonitrile*)

Kinsman, A.C. *et al.*, *J.A.C.S.*, 2003, **125**, 14120-14125 (*Hapalindole Q*, *synth*)

Richter, J.M. *et al.*, *J.A.C.S.*, 2008, **130**, 17938-17954 (*Hapalindole Q*, *synth*)

Hapalindole T

[106865-67-2]



H-50

$\text{C}_{21}\text{H}_{23}\text{ClN}_2\text{OS}$ 386.944

Isol. from the blue-green alga *Hapalosiphon fontinalis* ATCC39964. Antibacterial and antimycotic agent. Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. $[\alpha]_D^{25} -137$ (c, 1.5 in CH_2Cl_2). λ_{max} 222 (€ 35400); 283 (€ 7100) (MeOH) (Derep).


Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 1036 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Hapalindolinone A

H-51

3-Chloro-2-ethenyl-1-isocyano-2-methyl-5-(1-methylethenyl)spiro[bicyclo[4.1.0]heptane-7,3'-[3H]indol]-2'-(1'H)-one, 9CI

[109151-56-6]



$\text{C}_{21}\text{H}_{21}\text{ClN}_2\text{O}$ 352.862

Isol. from a cultured cyanobacterium and from *Fischerella* sp. and *Hapalosiphon laingii*. Adenylate cyclase inhibitor, vasopressin antagonist and ichthyotoxin. Cryst. Mp 92-96° dec. $[\alpha]_D^{25} -30$. Related to Hapalindole A, H-48. λ_{max} 217 (€ 77000); 253 (€ 3200); 300 (sh) (€ 1500) (no solvent specified) (Derep). λ_{max} 223 (€ 24380); 261 (€ 4620) (MeOH) (Berdy).

Dechloro: Hapalindolinone B

[109151-57-7]

$\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}$ 318.418

Isol. from a cultured cyanobacterium and from *Fischerella* sp. Adenylate cyclase inhibitor. Oil. λ_{max} 217 (€ 77000); 253 (€ 3200); 300 (sh) (€ 1500) (MeOH) (Derep). λ_{max} 223 (€ 24600); 261 (€ 4570) (MeOH) (Berdy).


Schwartz, R.E. *et al.*, *J.O.C.*, 1987, **52**, 3704

(*isol*, *ir*, *pmr*, *cmr*, *cryst struct*)

Hapalonamide G

H-52

[109281-38-1]



$\text{C}_{21}\text{H}_{23}\text{ClN}_2\text{O}_2$ 370.878

Author's numbering shown. See also Fontonamide, F-126. Minor alkaloid from the blue-green alga *Hapalosiphon fontinalis*. Also obt. by oxidn. of Hapalindole A, H-48.

10β-Hydroxy: Hapalonamide V

[123498-01-1]

$\text{C}_{21}\text{H}_{23}\text{ClN}_2\text{O}_3$ 386.877

Minor alkaloid from *Hapalosiphon fontinalis*.

10,15-Diepimer, dechloro: Hapalonamide H

H

[123498-00-0]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_2$ 336.433

Minor alkaloid from *Hapalosiphon fontinalis*. $[\alpha]_D^{25} +13.7$ (c, 0.15 in CHCl_3). Moore, R.E. *et al.*, *J.O.C.*, 1987, **52**, 3773 (*synth*, *pmr*)
 Moore, R.E. *et al.*, *Phytochemistry*, 1989, **28**, 1565 (*isol*)


Haplafine

H-53

4-[(3-Methyl-2-butenyl)oxy]-2(1H)-quinolinone, 9CI. 4-Prenyloxy-2-quinolone.

Haplaphine

[54357-78-7]



$\text{C}_{14}\text{H}_{15}\text{NO}_2$ 229.278

Alkaloid from the aerial parts of flowering *Haplophyllum perforatum* (Rutaceae). Cryst. (Me_2CO). Mp 159-160°.

N-Me: Ravenine

[20105-22-0]

$\text{C}_{15}\text{H}_{17}\text{NO}_2$ 243.305

Alkaloid from *Ravenia spectabilis* (Rutaceae). Mp 120-121°.

N-Me, 2',3'-Dihydroxy: 4-(2,3-Dihydroxy-3-methylbutoxy)-1-methyl-2(1H)-quinolinone

[80357-89-7]

$\text{C}_{15}\text{H}_{19}\text{NO}_4$ 277.319

Alkaloid from the heartwood of *Euxylophora paraensis* (Rutaceae). Needles (C_6H_6). Mp 94°. $[\alpha]_D^{20} +31.8$ (c, 0.9 in CHCl_3).

Paul, B.D. *et al.*, *Indian J. Chem.*, 1969, **7**, 678 (*Ravenine*)


Jurd, L. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1625 (*deriv*)

Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 654; *Chem. Nat. Compd. (Engl. Transl.)*, 619 (*isol*, *ir*, *pmr*, *struct*)

Haplocidiphytine

H-54

[87553-37-5]




$\text{C}_{37}\text{H}_{38}\text{N}_4\text{O}_8$ 666.729

No stereochem. reported for non-aspidosperma part, which presumably shows tautom. with a canthinone struct. (cf. Haplophytine, H-58). Alkaloid from *Haplophyton camicidum* (Apocynaceae). Small needles (EtOH). Mp 334° dec. $[\alpha]_D^{20} +91.5$ (c, 0.55 in CHCl_3). λ_{max} 235 (€ 32900); 265 (€ 20900); 290 (sh) (€ 7000) (EtOH).

Adesomoju, A.A. *et al.*, *Heterocycles*, 1983, **20**, 1511-1517 (*uv, ir, pmr, cmr, ms, struct*)

Haplodimerine H-55
[120931-43-3]




C₂₈H₂₆N₂O₆ 486.523

Dimer of Skimmianine and Flindersine. Alkaloid from the fruit of *Haplophyllum foliosum* (Rutaceae). Prisms (EtOH). Mp 292-293°. Optically inactive.

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1988, **24**, 838; *Chem. Nat. Compd. (Engl. Transl.)*, 714 (*isol, struct, uv, ir, nmr, ms, cryst struct*)

Haplofoline H-56
2,3,4,10-Tetrahydro-2,2-dimethyl-5H-pyrano[2,3-b]quinolin-5-one, 9CI. *Khaplofoline* [6431-83-0]



C₁₄H₁₅NO₂ 229.278

The names Khaplofoline and Haplofoline are equivalent transliterations from the Russian. Alkaloid from *Haplophyllum foliosum* (Rutaceae). Prisms (EtOAc). Mp 272-274°.

Hydrochloride: Mp 78-80°.

N-Ac:

Cryst. (EtOH). Mp 155-156°.

N-Me: *N-Methylkhaplofoline*

[6391-67-9]

C₁₅H₁₇NO₂ 243.305

Alkaloid from the stem bark of *Almeidea guyanensis* (Rutaceae). Needles (Et₂O). Mp 119-120°.

N-(3-Methyl-2-butenyl): **Haplobucharine** *N-Prenylhaplofoline*

[58969-44-1]

C₁₉H₂₃NO₂ 297.396

Alkaloid from *Haplophyllum bucharicum* (Rutaceae). Cryst. (EtOAc). Mp 126°.

Fakhrutdinova, I.M. *et al.*, *Uzb. Khim. Zh.*, 1963, **7**, 41; *CA*, **59**, 15331 (*isol, struct*)

Bowman, R.M. *et al.*, *J.C.S.(C)*, 1966, 1084 (*ir, uv, pmr, synth*)

Nesmelova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 815; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 831 (*Haplobucharine*)

Hammerum, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1977, **31**, 31 (*ms*)

Mouli, C. *et al.*, *Phytochemistry*, 1983, **22**, 2095 (*deriv*)


Bravo, P. *et al.*, *Gazz. Chim. Ital.*, 1988, **118**, 507 (*synth, Khaplofoline, N-Methylkhaplofoline*)

Subramanian, M. *et al.*, *Z. Naturforsch., B*, 1992, **47**, 1016 (*synth*)

Sekar, M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 294-296 (*synth*)

Thangavel, D. *et al.*, *J. Chem. Res.*, 2007, 124-126 (*synth*)

Haplophyllidine H-57
5,6,7,8-Tetrahydro-4,8-dimethoxy-8-(3-methyl-2-butenyl)furo[2,3-b]quinolin-7-ol, 9CI [18063-21-3]



Relative Configuration

C₁₈H₂₃NO₄ 317.384

Alkaloid from the seeds of *Haplophyllum perforatum* and roots of *Haplophyllum glabrinum* (Rutaceae). Strong CNS depressant, hypnotic synergist. Mp 110-111°. [α]_D²⁰ -24 (-16) (1.477 in Me₂CO).

Ac: **Acetylhaplophyllidine**

[23107-57-5]

C₂₀H₂₅NO₅ 359.421

Constit. of *Almeidea coerulea*. Cryst. (MeOH). Mp 147-148°.

5,6-Didehydro: **Perfaminole**

[116987-99-6]

C₁₈H₂₁NO₄ 315.368

Alkaloid from roots of *Haplophyllum glabrinum*. Mod. unstable. Stereochem. not determined but assumed to be the same as Haplophyllidine. λ_{max} 252; 327 (broad) (MeOH).

2',3'-Dihydro, 3'-hydroxy: 5,6,7,8-Tetrahydro-7-hydroxy-4,8-dimethoxy-α,α-dimethylfuro[2,3-b]quinoline-8-propanol.

Perforine

[18063-20-2]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Haplophyllum perforatum* (Rutaceae). Cryst. (MeOH). Mp 182-183°. [α]_D²⁰ +14.56 (c, 1.579 in MeOH). [α]_D +14.31 (c, 0.304 in CHCl₃).

4',2'-Isomer, 3'-hydroxy, 7-Ac:

C₂₀H₂₅NO₆ 375.421

Alkaloid from the leaves of *Almeidea rubra*. Powder. [α]_D²⁵ -8.7 (c, 0.15 in CH₂Cl₂).

7-Epimer: **Dihydroperfaminole**

[117065-23-3]

C₁₈H₂₃NO₄ 317.384

Alkaloid from roots of *Haplophyllum glabrinum*. Trivial name is confusing. λ_{max} 262; 278 (sh); 288 (sh) (MeOH).

Shakirov, T.T. *et al.*, *Dokl. Akad. Nauk SSSR*, No. 9, 1960, 40; *CA*, **56**, 11646a (*isol*)

Faizutdinova, Z.S. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 356; 1968, **4**, 360; 1969, **5**, 273; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 302; 1968, **4**, 304; 1969, **5**, 231 (*Haplophyllidine, Acetylhaplophyllidine, Perforine*)

Grundon, M.F. *et al.*, *Alkaloids (Academic Press)*, 1979, **17**, 105 (*rev*)


Rózsa, Zs. *et al.*, *Phytochemistry*, 1988, **27**, 2369 (*Perfaminole, Dihydroperfaminole*)

Santos, C.S. *et al.*, *J. Braz. Chem. Soc.*, 1998, **9**, 39-42; *CA*, **129**, 79124g

(*Acetylhaplophyllidine, pmr, cmr*)

Ambrozín, A.R.P. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 434-439 (*Almeidea rubra alkaloid*)

Haplophytine H-58
[16625-20-0]



C₃₇H₄₀N₄O₇ 652.746

Alkaloid from *Haplophyton cimidicum* (Apocynaceae). Shows insecticidal props. Mp 300-302° dec. (darkens from 260°, 288-292° dec.). [α]_D²⁵ +109 (c, 1.3 in CHCl₃). λ_{max} 230 (ε 48000); 265 (ε 14300); 326 (ε 5000) (EtOH/NaOH) (Derep). λ_{max} 220 (ε 48500); 265 (ε 14300); 305 (ε 4500) (EtOH) (Derep).

Hydrobromide (1:2): Mp 200-206° dec.

Me ether: Mp 288-291° dec. (280°).

16,17-Dihydro, 14,15-didehydro, N¹-de-Me: **Norisohaplophytine**

[87553-35-3]

C₃₆H₃₈N₄O₇ 638.719

Alkaloid from *Haplophyton cimidicum* (Apocynaceae). Small needles (EtOH). Mp 291° dec. [α]_D²⁰ +111.8 (c, 0.543 in CHCl₃). λ_{max} 230 (ε 15100); 255 (ε 11000); 290 (ε 5900) (EtOH).


Rae, I.D. *et al.*, *J.A.C.S.*, 1967, **89**, 3061-3062 (*uv, ir, pmr, ms, cryst struct*)

Yates, P. *et al.*, *J.A.C.S.*, 1973, **95**, 7842-7850 (*isol, uv, ir, pmr, cmr, struct*)

Cheng, P.-T. *et al.*, *Can. J. Chem.*, 1976, **54**, 726-732 (*cryst struct*)

Adesomoju, A.A. *et al.*, *Heterocycles*, 1983, **20**, 1511-1517 (*Norisohaplophytine*)

Haplosamine H-59
[169306-07-4]




C₁₆H₂₁NO₅ 307.346

Alkaloid from aerial parts of *Haplophyllum perforatum*; also obt. by hydrol. of Dubinidine, D-945. Mp 132-133°. λ_{max} 229 (log ε 5.18); 244 (sh) (log ε 4.72); 265 (log ε 4.39); 274 (log ε 4.49); 284 (log ε 4.4); 312 (log ε 4.32); 326 (log ε 4.42); 340 (log ε 4.29) (no solvent reported).

Rasulova, Kh.A. *et al.*, *Khim. Prir. Soedin.*, 1995, **31**, 586; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 487


Haplosine H-60

3-Ethyl-7-hydroxy-4,8-dimethoxy-2-(1H)-quinolinone, 9CI. 3-Ethyl-2,7-dihydroxy-4,8-dimethoxyquinoline



$C_{13}H_{15}NO_4$ 249.266
Alkaloid from *Haplophyllum perforatum*. Cryst. ($C_6H_6/EtOAc$). Mp 155-156°. λ_{max} 217 (log ϵ 4.57); 252 (log ϵ 4.24); 322 (log ϵ 4.24); 344 (log ϵ 4.22) (EtOH).

Rasulova, K.A. *et al.*, *Khim. Prir. Soedin.*, 1992, **28**, 249; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 214-216 (isol)


Haplotubinsonine H-61

$C_{19}H_{23}NO_4$ 329.395
Alkaloid from *Haplophyllum tuberculatum*. Cryst. (petrol). Mp 177-178°. Racemic. λ_{max} 237 (log ϵ 4.4); 262 (sh) (log ϵ 3.5); 352 (log ϵ 3.7) (MeOH).

Al-Rehaily, A.J. *et al.*, *Phytochemistry*, 2001, **57**, 597-602

Harmalacinine H-62

7,12-Dihydro-10-methoxy-2-methylindolo[2,3-a]quinolizin-4(6H)-one, 9CI [122279-86-1]



$C_{17}H_{16}N_2O_2$ 280.326
A truncated indole alkaloid, original skeleton uncertain. Prob. not secologanin-derived. Alkaloid from the seeds of *Peganum harmala* (Zygophyllaceae). Amorph.

Siddiqui, S. *et al.*, *Heterocycles*, 1989, **29**, 521 (isol, uv, ir, pmr, cmr, ms, struct)

Harmalanine H-63

7,12-Dihydro-10-methoxyindolo[2,3a]-quinolizin-4(6H)-one, 9CI

[116988-01-3]
As Harmalacinine, H-62 with R = H

$C_{16}H_{14}N_2O_2$ 266.299

Truncated indole alkaloid, original skeleton uncertain. Prob. not secologanin-derived. Alkaloid from the seeds of *Peganum harmala* (Zygophyllaceae). Rods ($CHCl_3$). Mp 108-110°.


O-De-Me: Mp 179°.

Siddiqui, S. *et al.*, *Heterocycles*, 1988, **27**, 1401 (isol, uv, ir, pmr, cmr, ms, struct)

Harmalidine H-64

1,2,4,5-Tetrahydro-8-methoxy-4,4-dimethylbenzo[b]pyrrolo[2,3,4-g]pyrrolizine, 9CI

[109794-97-0]



$C_{16}H_{18}N_2O$ 254.331
Alkaloid from the seeds of *Peganum harmala* (Zygophyllaceae). Yellow rods (MeOH). Mp 150-152°. λ_{max} 205 ; 215 ; 260 ; 340 ; 375 (MeOH).


O-De-Me: [109794-99-2]

Mp 210-212°.

Siddiqui, S. *et al.*, *Phytochemistry*, 1987, **26**, 1548 (isol, uv, ir, pmr, cmr, ms, struct)

Harmicine H-65

2,3,5,6,11,11b-Hexahydro-1H-indolizino[8,7-b]indole, 9CI



$C_{14}H_{16}N_2$ 212.294

(R)-form [202258-15-9]

Alkaloid from the leaves of *Kopsia griffithii*. $[\alpha]_D$ +119 (c, 0.09 in $CHCl_3$). Abs. config. revised in 2004. λ_{max} 225 (log ϵ 3.98); 281 (log ϵ 3.3); 292 (log ϵ 3.18) (EtOH).

(±)-form

Mp 174-175°.

Kam, T.-S. *et al.*, *Phytochemistry*, 1998, **47**, 145-147 (isol, uv, pmr, cmr, ms)

Itoh, T. *et al.*, *Heterocycles*, 2004, **63**, 655-661 (synth, abs config)

Knölker, H.J. *et al.*, *Synlett*, 2004, 1767-1768 (synth)

King, F.D. *et al.*, *J. Het. Chem.*, 2007, **44**, 1459-1463 (synth, pmr, cmr)


Lim, S.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1380-1383 (isol, pmr)

Allin, S.M. *et al.*, *Tet. Lett.*, 2007, **48**, 5669-5671 (synth)

Szawalko, J. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 406-413 (synth, abs config)

Harringtonine H-66

Cephalotaxine 4-methyl 2-hydroxy-2-(3-hydroxy-3-methylbutyl)butanedioate (ester), 9CI [26833-85-2]



n = 2

$C_{28}H_{37}NO_9$ 531.602

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus fortunei*, *Cephalotaxus hainanensis*, *Cephalotaxus sinensis* and *Cephalotaxus oliveri* (Cephalotaxaceae). Protein and DNA biosynthesis inhibitor. Shows antineoplastic activity, especially against murine lymphocytic leukaemias. More active than Vincristine against mouse leukaemias and melanomas. Has been used clinically against acute myelocytic leukaemia. Mp 73-75°. $[\alpha]_D$ -104.6 (c, 1.0 in $CHCl_3$). Log P 0.52 (uncertain value) (calc). λ_{max} 261 (ϵ 724); 290 (ϵ 4570) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 4.17 mg/kg. Exp. adverse systemic effects (G.I. tract, heart, haematopoietic organs). FK0250000

Parent acid: 3''-Des-O-methylharringtonine

5'-Des-O-methylharringtonine

$C_{27}H_{35}NO_9$ 517.575

Alkaloid from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Amorph. solid. $[\alpha]_D$ -113 (c, 0.43 in DMSO). λ_{max} 289 (ϵ 3630) (MeOH) (Berdy).

1''S-Hydroxy, parent acid: 1''S-Hydroxy-

3''-des-O-methylharringtonine. 3''S-Hydroxy-5'-des-O-methylharringtonine

[182325-74-2]

$C_{27}H_{35}NO_{10}$ 533.574

From leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Amorph. solid. $[\alpha]_D$ -91 (c, 1.00 in DMSO). λ_{max} 288 (ϵ 3311) (MeOH) (Berdy).

5'-Deoxy: Deoxyharringtonine

[36804-95-2]

$C_{28}H_{37}NO_8$ 515.602

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia* and *Cephalotaxus hainanensis* (Cephalotaxaceae). Shows antineoplastic props. Noncryst.

$[\alpha]_D$ -125.4 (c, 1.76 in $CHCl_3$). Log P 2.73 (uncertain value) (calc). λ_{max} 273 (ϵ 12590) (EtOH) (Derep). λ_{max} 290 (ϵ 14570) (EtOH) (Berdy).

▶ FK0279000

5'-Deoxy, parent acid: Deoxyharringtonic acid

[85623-76-3]

$C_{27}H_{35}NO_8$ 501.575

Alkaloid from *Cephalotaxus hainanensis* (Cephalotaxaceae). Mp 219-220°.

λ_{max} 291 (ϵ 2290) (MeOH) (Berdy).

5'-Deoxy, 11β-hydroxy: 11β-Hydroxy-

deoxyharringtonineC₂₈H₃₇NO₉ 531.602

Isol. from *Cephalotaxus harringtonia* var. *drupacea*. Shows antileukaemic activity against P-388 cells. Oil. $[\alpha]_D^{25}$ -77 (c, 0.085 in MeOH). λ_{\max} 290 (log ϵ 3.3) (MeOH).

Homologue (n = 1), 5'-deoxy: NordeoxyharringtonineC₂₇H₃₅NO₈ 501.575

Alkaloid from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Cytotoxic against P-388 leukaemia. Pale yellowish oil. $[\alpha]_D$ -90 (c, 0.07 in MeOH). λ_{\max} 291 (log ϵ 3.36) (MeOH).

Homologue (n = 3): Homoharringtonine.

Omacetaxine mepesuccinate, INN, USAN. Ceflatonin[®]. CGX 635. NSC 141633

[26833-87-4]

C₂₉H₃₉NO₉ 545.628

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus harringtonia* var. *drupacea*, *Cephalotaxus fortunei*, *Cephalotaxus hainanensis* and *Cephalotaxus sinensis* (Cephalotaxaceae). Shows antitumour activity comparable with that of Harringtonine. Has been used clinically against acute myelocytic leukaemia. Mp 144-146°. $[\alpha]_D$ -119 (c, 0.96 in CHCl₃). Log P 1.05 (uncertain value) (calc). λ_{\max} 261 (ϵ 724); 290 (ϵ 4570) (EtOH) (Derep).

► Adverse systemic effects when used therapeutically. LD₅₀ (mus, orl) 7.456 mg/kg. FK0260000

Homologue (n = 3), parent acid: 3''-Des-O-methylhomoharringtonine. 5'-Des-O-methylhomoharringtonine

[98599-84-9]

C₂₈H₃₇NO₉ 531.602

From leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Amorph. solid. $[\alpha]_D$ -172 (c, 0.50 in MeOH).

Homologue (n = 3), 1''S-hydroxy: Cephalozomine EC₂₉H₃₉NO₁₀ 561.628

Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent. Amorph. solid. $[\alpha]_D$ -131 (c, 2.9 in MeOH). λ_{\max} 291 (ϵ 2100) (MeOH).

Homologue (n = 3), 5'-deoxy: Homodeoxyharringtonine. Deoxyhomoharringtonine

[64543-63-1]

C₂₉H₃₉NO₈ 529.629

Alkaloid isol. from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*, also detected by glc/ms in callus cultures of *Cephalotaxus harringtonia* (Cephalotaxaceae). Cytotoxic against P-388 leukaemia. Pale yellowish oil. $[\alpha]_D$ -122 (c, 1 in MeOH). λ_{\max} 291 (log ϵ 3.55) (MeOH).

Homologue (n = 3), 5'-deoxy, 1''S-hydroxy: Cephalozomine FC₂₉H₃₉NO₉ 545.628

Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent. Amorph. solid. $[\alpha]_D$ -61 (c, 0.2 in MeOH). λ_{\max} 291 (ϵ 1800) (MeOH).

Homologue (n = 3), 5'-deoxy, 11 α -hydroxy: 11 α -HydroxyhomodeoxyharringtonineC₂₉H₃₉NO₉ 545.628

Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*. Shows antileukaemic activity against P-388 cells. Oil. $[\alpha]_D^{25}$ -115 (c, 0.07 in MeOH). λ_{\max} 290 (log ϵ 3.5) (MeOH).

Homologue (n = 3), 5'-deoxy, 11 β -hydroxy: 11 β -HydroxyhomodeoxyharringtonineC₂₉H₃₉NO₉ 545.628

Alkaloid from *Cephalotaxus harringtonia* var. *drupacea*. Shows antileukaemic activity against P-388 cells. Oil. $[\alpha]_D^{25}$ -153 (c, 0.10 in MeOH). λ_{\max} 290 (log ϵ 3.53) (MeOH).

Homologue (n = 3), 5'-epimer, 6'-hydroxy(1): Cephalozomine KC₂₉H₃₉NO₁₀ 561.628

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. $[\alpha]_D$ -99 (c, 0.6 in MeOH). Epimeric with Cephalozomine L at C-6'. λ_{\max} 210 (ϵ 5800); 240 (ϵ 2300); 289 (ϵ 2100) (no solvent reported).

Homologue (n = 3), 5'-epimer, 6'-hydroxy(2): Cephalozomine LC₂₉H₃₉NO₁₀ 561.628

Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. $[\alpha]_D$ -93 (c, 1 in MeOH). Epimeric with Cephalozomine K at C-6'. λ_{\max} 215 (ϵ 4700); 240 (ϵ 2700); 289 (ϵ 2400) (MeOH).

Homologue (n = 4), 5'-deoxy: BishomodeoxyharringtonineC₃₀H₄₁NO₈ 543.656

Alkaloid from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Cytotoxic against P-388 leukaemia cells. Pale yellowish oil. $[\alpha]_D$ -74 (c, 0.27 in CHCl₃). λ_{\max} 293 (log ϵ 3.5) (CHCl₃). λ_{\max} 293 (ϵ 3162) (MeOH) (Berdy).

Powell, R.G. *et al.*, *Tet. Lett.*, 1970, 815 (*pmr, struct*)

Mikolajczak, K.L. *et al.*, *Tetrahedron*, 1972, **28**, 1995 (*uv, ms, pmr, struct*)

Spencer, G.F. *et al.*, *J. Chromatogr.*, 1976, **120**, 335 (*ms, gle*)

Delfel, N.E. *et al.*, *Phytochemistry*, 1977, **16**, 1595 (*Homodeoxyharringtonine*)

Mikolajczak, K.L. *et al.*, *J.O.C.*, 1978, **43**, 4762 (*synth*)

Kelly, T.R. *et al.*, *J.O.C.*, 1979, **44**, 63 (*synth*)

Smith, C.R. *et al.*, *Med. Chem. (Academic Press)*, 1980, **16**, 391 (*rev*)

Weisleder, D. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 114 (*cmr*)

Xue, Z. *et al.*, *Yaouxue Xuebao*, 1981, **16**, 752;

CA, **96**, 82690u (*Deoxyharringtonic acid*)

Hiranuma, S. *et al.*, *J.O.C.*, 1983, **48**, 5321

(*synth, Homoharringtonine*)

Cheng, J. *et al.*, *Yaouxue Xuebao*, 1984, **19**, 178;

CA, **103**, 160738 (*synth*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 63 (*bioactivity*)

Bao, G. *et al.*, *Zhongguo Yixue Kexueyuan Xuebao*, 1988, **10**, 283; CA, **110**, 95588 (*cryst struct, Harringtonine, Homoharringtonine*)

Shen, C. *et al.*, *Huaxue Xuebao*, 1990, **48**, 185;

CA, **113**, 115639 (*cd*)

Martindale, *The Extra Pharmacopoeia*, 30th

edn., Pharmaceutical Press, 1993, 482

Takano, I. *et al.*, *J. Nat. Prod.*, 1996, **59**, 965;

1192 (*Nordeoxy, Homodeoxy,*

Bishomodeoxy, 11-Hydroxydeoxy, 11-

Hydroxyhomodeoxy)

Takano, I. *et al.*, *Phytochemistry*, 1996, **43**, 299 (*isol, derivs*)

Morita, H. *et al.*, *Tetrahedron*, 2000, **56**, 2929-

2934; 2002, **58**, 5489-5495 (*Cephalozomines*)

Marin, D. *et al.*, *Cancer (Philadelphia)*, 2005,

103, 1850-1855 (*homoharringtonine, clin trial*)

Eckelberger, J.D. *et al.*, *J.A.C.S.*, 2006, **128**,

10370-10371 (*Deoxyharringtonine, synth*)

Tang, R. *et al.*, *Mol. Cancer Ther.*, 2006, **5**,

723-731 (*homoharringtonine, pharmacol*)

Quintas-Cardama, A. *et al.*, *Cancer*

(*Philadelphia*), 2007, **109**, 248-255; 2625

(*homoharringtonine, clin trial*)

Lou, Y.J. *et al.*, *Leuk. Lymphoma*, 2007, **48**,

1400-1406 (*homoharringtonine, pharmacol*)

Quintas-Cardama, A. *et al.*, *IDrugs*, 2008, **11**,

356-372 (*homoharringtonine, rev*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties*

of Industrial Materials, 8th edn., Van

Nostrand Reinhold, 1992, HGI575

Harzianic acid**H-67**

α -Hydroxy-4-(1-hydroxy-2,4-octadienyldene)-1-methyl- α -(1-methylethyl)-3,5-dioxo-2-pyrrolidinedipropionic acid, 9CI [157148-06-6]

C₁₉H₂₇NO₆ 365.425

Prod. by *Trichoderma harzianum* SY-307.

Antimicrobial agent. Orange powder.

Sol. MeOH, Me₂CO, MeCN, EtOAc;

poorly sol. toluene, hexane. $[\alpha]_D^{25}$ +19.6

(c, 1 in MeOH). λ_{\max} 263 (ϵ 14100); 287

(ϵ 15500); 337 (ϵ 16600) (MeOH/NaOH)

(Derep). λ_{\max} 244 (ϵ 11000); 299 (sh) (ϵ

10500); 343 (sh) (ϵ 23400); 359 (ϵ 27500);

376 (sh) (ϵ 23400); 398 (sh) (ϵ 10500)

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).


(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

(MeOH) (Derep).

Harzianopyridone**H-68**C₁₄H₁₉NO₅ 281.308**(R)-form**

Metab. of *Trichoderma harzianum*

ATCC64870. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 77-78°. [α]_D²² -12.3 (c, 1 in MeOH). λ_{\max} 237 (ϵ 11500); 271 (ϵ 790); 321 (ϵ 6400) (MeOH) (Berdy). λ_{\max} 243; 267; 331 (EtOH) (Berdy).

(±)-**form** [126637-69-2]

From *Trichoderma harzianum*. Antifungal agent. Needles (EtOAc/petrol). Mp 125°.


Dickinson, J.M. *et al.*, *J.C.S. Perkin 1*, 1989, 1885 (*isol, struct, biosynth*)

Cutler, H.G. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 2629 (*isol, pmr, cmr, props*)

Trecourt, F. *et al.*, *J. Het. Chem.*, 1995, **32**, 1117 (*synth*)

Hastacine

[20361-77-7]



C₁₈H₂₇NO₅ 337.415

Stereoisomeric with Platyphylline, P-516. Alkaloid from *Cacalia hastata* and *Cacalia robusta* (Asteraceae). Shows spasmolytic props. Cryst. (EtOH). Mp 170-171°. [α]_D²⁰ -72.3.

Aasen, A.J. *et al.*, *J.O.C.*, 1969, **34**, 12; 4137-4143 (*struct, pmr*)

Witte, L. *et al.*, *Phytochemistry*, 1993, **32**, 187-196 (*ms, struct*)

Konovalov, V.S. *et al.*, *Zh. Obshch. Khim.*, 1993, **15**, 328; *CA*, **40**, 3760 (*isol*)


Hasubanonine

H-70

7,8-Didehydro-3,4,7,8-tetramethoxy-17-methylhasubanan-6-one, 9CI. O-Methylaknadinine

[1805-85-2]

[30333-82-5 ((±)-form)]



Absolute configuration

C₂₁H₂₇NO₅ 373.448

Alkaloid from *Stephania japonica*, *Stephania japonica* var. *australis* and *Stephania elegans* (Menispermaceae). Mp 116-117°. [α]_D²⁷ -214 (c, 2.0 in MeOH).

Hydrochloride:

Needles (CHCl₃/Et₂O). Mp 210-211°.

[α]_D²⁸ -133 (c, 1.14 in MeOH).

N-Me:

Prisms (MeOH/Et₂O or MeOH aq.) (as iodide). Mp 178° (171-173°) (iodide).

[α]_D²⁷ -61 (c, 0.28 in MeOH) (as iodide).

N-De-Me: Mp 222.5° (as hydrobromide).

[α]_D²⁶ -105 (c, 0.12 in MeOH) (as

hydrobromide).

O³-De-Me: Homostephanoline. O³-Demethylhasubanonine

[2689-15-8]

C₂₀H₂₅NO₅ 359.421

Alkaloid from *Stephania japonica* (Menispermaceae). Mp 233°. [α]_D -247.8 (CHCl₃).

O⁴-De-Me: 7,8-Didehydro-4-hydroxy-3,7,8-trimethoxy-17-methylhasubanan-6-one, 9CI. Aknadinine. *Hernandoline*. 4-Demethylhasubanonine

[24148-86-5]

C₂₀H₂₅NO₅ 359.421

Alkaloid from the roots and rhizomes of *Stephania hernandifolia* and the leaves, stems and roots of *Stephania elegans*, also present in *Stephania sasakii* and *Stephania sutchuenensis* (Menispermaceae). Fine flakes by subl. Mp 70°. [α]_D²⁹ -283 (c, 0.1 in EtOH). Subl. at 120-130°/0.2 mm.

O⁴-De-Me, N-de-Me: Aknadinine. 4-Demethylnorhasubanonine

[24148-89-8]

C₁₉H₂₃NO₅ 345.394

Alkaloid from the roots and rhizomes of *Stephania hernandifolia* (Menispermaceae). Plates (MeOH). Mp 156°. [α]_D²⁷ -200 (c, 0.55 in EtOH).

O⁴-De-Me, 1,1'-dimer: Bisaknadinine

[73461-15-1]

C₄₀H₄₈N₂O₁₀ 716.827

Alkaloid from the stem and root of *Stephania sasakii* (Menispermaceae). Cryst. + 1Me₂CO (Me₂CO). Mp 198-200°. [α]_D -253.3 (CHCl₃). The first known dimeric hasubanan alkaloid.

6 ξ -Alcohol, O⁴-de-Me: 7,8-Didehydro-3,7,8-trimethoxy-17-methylhasubanan-4,6-diol. Hernandolinol

[30452-61-0]

C₂₀H₂₇NO₅ 361.437

Alkaloid from *Stephania hernandifolia* (Menispermaceae). [α]_D -97.9 (EtOH).

16-Oxo: 16-Oxohasubanonine. 7,8-Didehydro-3,4,7,8-tetramethoxy-17-methylhasubanan-6,16-dione, 9CI

[51804-70-7]

[30333-89-2 ((±)-form)]

C₂₁H₂₅NO₆ 387.432

Minor alkaloid from the stems and rhizomes of *Stephania japonica* (Menispermaceae). Prisms (C₆H₆/Et₂O). Mp 161°. [α]_D²³ -105.2 (c, 0.5 in CHCl₃).

16-Oxo, O⁴-de-Me: Aknadilactam. 4-Demethyl-16-oxohasubanonine

[24191-97-7]

C₂₀H₂₃NO₆ 373.405

Alkaloid from *Stephania sasakii* (Menispermaceae). Amorph. [α]_D -212 (CHCl₃).

1-Nitro, O⁴-de-Me: 1-Nitroaknadinine

[154739-06-7]

C₂₀H₂₄N₂O₇ 404.419

Alkaloid from roots of *Stephania sutchuenensis* (Menispermaceae). Yellow needles (Me₂CO). Mp 206° dec. [α]_D -186.7 (c, 0.523 in EtOH).

6-Epimer, 6 ξ -alcohol, O⁴-de-Me: Epihernandolinol

[81027-39-6]

C₂₀H₂₇NO₅ 361.437

Alkaloid from the leaves, stems and roots of *Stephania elegans* (Menispermaceae). Oil. [α]_D²⁷ -16 (c, 1.08 in EtOH).

Kondo, H. *et al.*, *Yakugaku Zasshi*, 1928, **48**, 1141; *CA*, **23**, 2979 (*Homostephanoline, isol*)

Kondo, H. *et al.*, *CA*, 1953, **47**, 5951f (*isol*)
Tomita, M. *et al.*, *Tet. Lett.*, 1964, 2937; 1965, 1019 (*pmr, struct, ms*)

Ibuka, T. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 1809; 1939; 1974, **22**, 782 (*Aknadinine, Homostephanoline, Aknadilactam, 16-Oxohasubanonine, struct, synth*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1968, **33**, 4529 (*synth*)

Kunimoto, J. *et al.*, *Tet. Lett.*, 1969, 3287 (*Aknadilactam, uv, ir, pmr, struct*)

Fadeeva, I.I. *et al.*, *Khim. Pri. Soedin.*, 1970, **6**, 492; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 516 (*Hernandolinol, isol, struct*)

Moza, B.K. *et al.*, *Tetrahedron*, 1970, **26**, 427 (*Aknadinine, Aknadinine, isol, uv, ir, pmr, struct*)

Watanabe, Y. *et al.*, *Phytochemistry*, 1975, **14**, 2675 (*isol, ms, 16-Oxohasubanonine*)

Matsui, M. *et al.*, *Phytochemistry*, 1979, **18**, 1087 (*isol*)

Kunimoto, J. *et al.*, *Heterocycles*, 1980, **14**, 175; 1981, **16**, 351 (*Bisaknadinine, cryst struct, config*)

Kunimoto, J. *et al.*, *Phytochemistry*, 1980, **19**, 2735 (*Bisaknadinine, uv, ord, cd, ir, pmr, ms, synth*)

Singh, R.S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 664 (*Hasubanonine, Aknadinine, Hernandolinol, Epihernandolinol, isol, uv, ir, pmr, ms, struct, synth*)

Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1981, 2016; 2030 (*biosynth*)

Wang, X.-K. *et al.*, *Phytochemistry*, 1994, **35**, 263 (*1-Nitroaknadinine*)

Jones, S.B. *et al.*, *Org. Lett.*, 2006, **8**, 3757-3760 (*synth*)

Hasunohanin

H-71

C₃₁H₃₆N₂O₅ 516.636


Struct. unknown. Alkaloid from *Stephania japonica*. Mp 102-103° Mp 267-268° (as hydrochloride). [α]_D -83 (CHCl₃).

Kondo, H. *et al.*, *Yakugaku Zasshi*, 1928, **48**, 163; 1938, **58**, 46

Haterumaimide I

H-72

[368421-37-8]



C₂₀H₂₈ClNO₄ 381.898


Constit. of a *Lissoclinum* sp. Oil. [α]_D³² +62 (c, 0.77 in MeOH). λ_{\max} 210 (log ϵ 3.56) (MeOH).

Uddin, M.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1169-1173 (*isol, pmr, cmr*)

Haterumaimide C


H-73

[338949-10-3]

 $C_{20}H_{27}Cl_2NO_4$ 416.343Isol. from a *Lissoclinum* sp. Cytotoxic. $[\alpha]_D^{28} +66.6$ (c, 0.06 in MeOH).Uddin, M.J. *et al.*, *Heterocycles*, 2001, **54**, 1039-1048 (isol, pmr, cmr)**Haterumaimide E**

H-74

[338949-12-5]

 $C_{20}H_{29}Cl_2NO_4$ 418.359Isol. from a *Lissoclinum* sp. Cytotoxic. $[\alpha]_D^{29} +29.6$ (c, 0.16 in MeOH).**6-Ketone: Haterumaimide B**

[338949-09-0]

 $C_{20}H_{27}Cl_2NO_4$ 416.343Isol. from a *Lissoclinum* sp. Cytotoxic. $[\alpha]_D^{33} +32.6$ (c, 0.48 in MeOH). **Δ^7 -Isomer, 6-ketone: Haterumaimide D**

[338949-11-4]

 $C_{20}H_{27}Cl_2NO_4$ 416.343Isol. from a *Lissoclinum* sp. Cytotoxic. $[\alpha]_D^{29} -27.7$ (c, 0.16 in MeOH).**2-Dechloro, 6-ketone: Haterumaimide M**

[735278-27-0]

 $C_{20}H_{28}ClNO_4$ 381.898Constit. of *Pleurobranchus alboguttatus*. Amorph. solid. $[\alpha]_D +14.4$ (c, 0.36 in CH_2Cl_2).**2-Dechloro, Δ^7 -isomer, 6-ketone: Haterumaimide L**

[735278-28-1]

 $C_{20}H_{28}ClNO_4$ 381.898Constit. of *Pleurobranchus alboguttatus* and *Pleurobranchus forskalii*. Amorph. solid. $[\alpha]_D -7.2$ (c, 0.43 in CH_2Cl_2).**3-Dechloro, β 3-hydroxy: 3-Hydroxychloro-olissoclimide**

[733753-15-6]

 $C_{20}H_{30}ClNO_5$ 399.913Constit. of *Pleurobranchus alboguttatus*. Amorph. solid.Uddin, M.J. *et al.*, *Heterocycles*, 2001, **54**,

1039-1048 (isol, pmr, cmr)


Fu, X. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1415-1418

(Haterumaimides L-M, 3-Hydroxychloro-olissoclimide)

Haterumaimide F

H-75

[368421-34-5]

 $C_{20}H_{30}ClNO_4$ 383.914Constit. of a *Lissoclinum* sp. Oil. $[\alpha]_D^{29} +53.7$ (c, 0.35 in MeOH). λ_{max} 210 (log ϵ 3.57) (MeOH).**6-Ketone: Haterumaimide G**

[368421-35-6]

 $C_{20}H_{28}ClNO_4$ 381.898Constit. of a *Lissoclinum* sp. Oil. $[\alpha]_D^{29} +63.5$ (c, 0.58 in MeOH). λ_{max} 207 (log ϵ 3.56) (MeOH).**6-Deoxy, 18-hydroxy: Haterumaimide J**

[479249-29-1]

 $C_{20}H_{30}ClNO_4$ 383.914Isol. from a *Lissoclinum* sp. Cytotoxic agent. Oil. $[\alpha]_D^{29} +68$ (c, 0.92 in MeOH). λ_{max} 210 (ϵ 3800) (MeOH).**6-Deoxy, 18-acetoxy: Haterumaimide K**

[479249-30-4]

 $C_{22}H_{32}ClNO_5$ 425.951Isol. from a *Lissoclinum* sp. Cytotoxic. $[\alpha]_D^{33} +59.6$ (c, 0.19 in MeOH). λ_{max} 210 (ϵ 3600) (MeOH). **Δ^7 -Isomer, 6-ketone: Haterumaimide H**

[368421-36-7]

 $C_{20}H_{28}ClNO_4$ 381.898Constit. of a *Lissoclinum* sp. Oil. $[\alpha]_D^{32} +47.6$ (c, 0.31 in MeOH). λ_{max} 205 (log ϵ 3.7); 238 (log ϵ 3.9) (MeOH).Uddin, M.J. *et al.*, *J. Nat. Prod.*, 2001, **64**,

1169-1173 (isol, pmr, cmr)

Uddin, M.J. *et al.*, *Chem. Lett.*, 2002, 1028-


1029 (Haterumaimides J,K)

Haterumalide NE

H-76

Haterumalide E

[245342-52-3]

 $C_{21}H_{29}ClO_7$ 428.909Macrolide antibiotic. Isol. from an *Ircinia* sp. and prod. by *Serratia plymuthica* A153. Cytotoxic agent. Oil.3-Ac: Haterumalide NA. Haterumalide A. *Oocydin A*. FR 177391. Antibiotic FR

177391

[245342-48-7]

[260362-86-5, 288581-04-4]

 $C_{23}H_{31}ClO_8$ 470.946Isol. from an *Ircinia* sp., prod. by *Serratia marcescens*, *Serratia plymuthica* A153. Cytotoxic, antihyperlipidaemic and anti-oomycetic agent. Powder. Mp 106-108°. $[\alpha]_D^{26} -3$ (c, 0.05 in MeOH). $[\alpha]_D +18.2$ (MeOH). $[\alpha]_D +32$ (c, 0.85 in MeOH). Stereochemical identity of the various isolates has not been demonstrated.**3-Ac, butyl ester: Haterumalide NB**

[245342-49-8]

 $C_{27}H_{39}ClO_8$ 527.053Isol. from an *Ircinia* sp. Cytotoxic agent. Oil.**3-Ac, (2-methylene-3-oxobutyl) ester: Haterumalide B**

[245122-93-4]

 $C_{28}H_{37}ClO_9$ 553.048Isol. from a *Lissoclinum* sp., prod. by *Serratia plymuthica* A153. Cytotoxic agent. Oil. $[\alpha]_D 0$ ($CHCl_3$). λ_{max} 206 (ϵ 12000) (MeOH).**3-Ac, (2-sulfoethyl)amide: Biselide D.**

Taurohaterumalide NA

[862167-80-4]

 $C_{25}H_{36}ClNO_{10}S$ 578.079Alkaloid from a *Didemnum* sp. Oil. **δ -Lactone, 3-Ac: Haterumalide X**

[343573-61-5]

 $C_{23}H_{29}ClO_7$ 452.931Prod. by *Serratia plymuthica* A153. Cytotoxic agent.**9R-Hydroxy, 3-Ac: Haterumalide ND**

[245342-51-2]

 $C_{23}H_{31}ClO_9$ 486.945Isol. from an *Ircinia* sp. Cytotoxic agent. Oil.**9R-Hydroxy, 3-Ac, butyl ester: Haterumalide NC**

[245342-50-1]

 $C_{27}H_{39}ClO_9$ 543.052Isol. from an *Ircinia* sp. Cytotoxic agent. Oil.**20-Hydroxy, 3-Ac: Biselide C**

[862167-78-0]

 $C_{23}H_{31}ClO_9$ 486.945Constit. of a *Didemnum* sp. Cytotoxic. Oil.**20-Acetoxy, 3-Ac: Biselide A**

[786712-66-1]

 $C_{25}H_{33}ClO_{10}$ 528.982Constit. of a *Didemnum* sp. Oil.**20-Acetoxy, 3-Ac, (2-methylene-3-oxobutyl) ester: Biselide B**

[786712-67-2]

 $C_{30}H_{39}ClO_{11}$ 611.084Constit. of a *Didemnum* sp. Oil.Strobel, G. *et al.*, *Microbiology (Reading, U.K.)*, 1999, **145**, 3557-3564 (*Oocydin A*)Ueda, K. *et al.*, *Tet. Lett.*, 1999, **40**, 6305-6308;

6309-6312 (isol, pmr, cmr)

Pat. Coop. Treaty (WIPO), 2001, 40442; CA,

135, 32806q (Haterumalide X)Kigoshi, H. *et al.*, *Org. Lett.*, 2003, **5**, 957-960


(synth, abs config)

Gu, Y. *et al.*, *Org. Lett.*, 2003, **5**, 4385-4388

(synth)

Teruya, T. *et al.*, *Chem. Lett.*, 2004, **33**, 1184-1185 (*Biseldide A*)
 Sato, B. *et al.*, *J. Antibiot.*, 2005, **58**, 634-639; 640-647; 648-653; 654-662 (*FR 177391*)
 Hoye, T.R. *et al.*, *J.A.C.S.*, 2005, **127**, 6950-6951 (*Haterumalide NA, synth*)
 Teruya, T. *et al.*, *Tetrahedron*, 2005, **61**, 6561-6567 (*Biselides*)
 Roulland, E. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 3762-3765 (*synth*)
 Schomaker, J.M. *et al.*, *J.A.C.S.*, 2008, **130**, 12228-12229 (*Haterumalide NA, NC, synth*)

Hatomamicin H-77
 YL 0358M-A. Antibiotic YL 0358M-A [116290-93-8]




Relative Configuration

$C_{22}H_{31}NO_5$ 389.491
 Numbering given here corresponds with that of Abikoviromycin, A-14. Alkaloid antibiotic from the culture filtrate of *Saccharopolyspora griseopurpurea*. Exhibits weak antimicrobial activity against gram-positive organisms. Also displays cytotoxic activity against lymphoid leukaemia L1210 and leukaemia P388. Pale yellowish prisms (MeCN). Sol. MeOH, Me₂CO, EtOAc, MeCN, CHCl₃; fairly sol. C₆H₆; poorly sol. H₂O. Mp 93°. $[\alpha]_D^{25} +88$ (c, 1 in MeOH). Labile in soln. but stable in cryst. form. λ_{max} 297 (ε 14200); 320 (sh) (ε 13600) (MeOH) (Derep). λ_{max} 300 (EtOH) (Berdy).


1,7a-Dihydro: Antibiotic YL 0358M-B. YL 0358M-B [118973-61-8]
 $C_{22}H_{33}NO_5$ 391.506
 Prod. by *Saccharomyces griseopurpurea*. Powder. λ_{max} 285 (MeOH).
Japan. Pat., 1988, (*Yamanouchi*)88 122 686; *CA*, **110**, 93563p (*isol*)
 Imai, H. *et al.*, *J. Antibiot.*, 1989, **42**, 1043-1048 (*isol, uv, ir, pmr, cmr, cryst struct*)

Havanine† H-78
 [72755-23-8]




$C_{18}H_{25}NO_5$ 335.399
 Alkaloid from *Hymenocallis arenicola* (Amaryllidaceae). Prisms (Me₂CO). Mp 135-137°.
 Döpke, W. *et al.*, *Z. Chem.*, 1979, **19**, 377 (*uv, ir, pmr, ms, struct*)

Hazuntamine H-79
 [155885-77-1]



$C_{42}H_{54}N_4O_5$ 694.912
 Alkaloid from root bark of *Hazunta modesta* var. *methuenii* subvar. *methuenii* (Apocynaceae). Cryst. (MeOH). Mp 254°. $[\alpha]_D^{20} -88$ (c, 0.7 in Py).
 Bui, A.-M. *et al.*, *Heterocycles*, 1994, **38**, 1025 (*isol, uv, ir, pmr, cmr, ms, struct*)


Hazuntiphyllidine, 9CI H-80
 [137760-56-6]



$C_{40}H_{46}N_4O_3$ 630.828
 Exists as two different forms depending on solvent. Struct. shown is the 2,2'-ring-opened form found in DMSO-d₆. Alkaloid from *Hazunta modesta* var. *modesta* subvar. *divaricata* (Apocynaceae). Cryst. (MeOH). Mp 300°. $[\alpha]_D^{22} -28$ (c, 0.66 in CHCl₃).
2'-Deoxy, 2',16'-didehydro: Anhydrohazuntiphyllidine


[137760-45-3]
 $C_{40}H_{44}N_4O_2$ 612.813
 Alkaloid from *Hazunta modesta* var. *modesta* subvar. *divaricata* (Apocynaceae). Cryst. (hexane). Mp 260°. $[\alpha]_D^{22} -175$ (c, 0.57 in CHCl₃).
 Bui, A.-M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 514 (*isol, uv, ir, pmr, cmr, struct*)

Hazuntiphylline, 9CI H-81
 [103590-56-3]




$C_{40}H_{46}N_4O_3$ 630.828
 Alkaloid from the leaves of *Hazunta modesta* var. *modesta* subvar. *divaricata* (Apocynaceae). Amorph. solid. $[\alpha]_D^{20} +14$ (c, 1 in CHCl₃).
 Bui, A.-M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 321 (*isol, uv, ir, pmr, cmr, ms, struct*)

Hebeirubescensin A H-82
 [887333-23-5]




$C_{26}H_{37}NO_8$ 491.58
 Constit. of *Isodon rubescens*. Amorph. powder. $[\alpha]_D^{19} -23.2$ (c, 0.12 in MeOH).
 Huang, S.-X. *et al.*, *Tetrahedron*, 2006, **62**, 4941-4947 (*Hebeirubescensin A*)

Hebridamine H-83
 [102487-24-1]



$C_{42}H_{50}N_2O_8$ 710.866
 First example of a benzyloquinoline-phenanthrene dimer. Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). Positive opt. rotn.
 Chalandre, M.C. *et al.*, *Can. J. Chem.*, 1986, **64**, 123 (*isol, uv, pmr, ms, struct*)

Hectochlorin H-84




Absolute Configuration

$C_{27}H_{34}Cl_2N_2O_9S_2$ 665.611
 Isol. from *Bursatella leachii* and *Lyngbya majuscula*. Promoter of actin polym. Antifungal and cytotoxic agent. Pale yellow solid. $[\alpha]_D^{25} -8.7$ (c, 1.04 in MeOH).
O-De-Ac: Deacetylhectochlorin
 $C_{25}H_{32}Cl_2N_2O_8S_2$ 623.574
 Isol. from the sea hare *Bursatella leachii*. Potent stimulator of actin assembly. Cytotoxic. Amorph. solid. $[\alpha]_D^{20} -26$ (c, 0.1 in MeOH).

- Marquez, B.L. *et al.*, *J. Nat. Prod.*, 2002, **65**, 866-871 (*isol, ir, pmr, cmr, cryst struct*)
 Cetusic, J.R.P. *et al.*, *Org. Lett.*, 2002, **4**, 1307-1310 (*synth*)
 Suntornchashweij, S. *et al.*, *J. Nat. Prod.*, 2005, **68**, 951-955 (*isol, pmr, cmr*)
 Ramaswamy, A.V. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1977-1986 (*biosynth*)

Hederacine A**H-85**

[616888-47-2]



$C_{16}H_{26}N_2O_3$ 294.393
 Alkaloid from the aerial parts of *Glechoma hederaceae*. Amorph. yellow solid. λ_{max} 232 (log ϵ 4) (MeOH).

 γ -Lactone: **Hederacine B**

[616888-48-3]


 $C_{16}H_{24}N_2O_2$ 276.378

Alkaloid from the aerial parts of *Glechoma hederaceae*. Amorph. yellow solid. λ_{max} 233 (log ϵ 3.98) (MeOH).

Kumarasamy, Y. *et al.*, *Tetrahedron*, 2003, **59**, 6403-6407 (*isol, pmr, cmr, ms*)


Hediamine**H-86**

[183162-31-4]

 $C_{21}H_{19}NO_5$ 365.385

First known C-homoprotoberberine. Alkaloid from roots of *Berberis actinocantha*.

Rahimizadeh, M. *et al.*, *J. Sci., Islamic Repub. Iran*, 1996, **7**, 172; *CA*, **125**, 323013a


Hedycapitelline**H-87** $C_{22}H_{30}N_2O_2$ 354.491

Alkaloid from *Hedyotis capitellata* var. *mollis*. $[\alpha]_D$ -5 (c, 1 in $CHCl_3$). λ_{max} 209 (log ϵ 4.15); 270 (log ϵ 3.53); 316 (log ϵ 4.16) (EtOH).

Phuong, N.M. *et al.*, *Planta Med.*, 1999, **65**, 761-762

Hedycapitine**H-88**

4-(3,3-Dimethyloxiranyl)-2-methylindolo[2,3-a]quinolizine, 9CI

 $C_{20}H_{18}N_2O$ 302.375

Claimed to exist in tautomeric equilib. with the 4-methylene-12H-form.

(ξ)-form [257905-18-3]

Alkaloid from *Hedyotis capitellata* var. *mollis*. $[\alpha]_D$ +4 (c, 1 in $CHCl_3$). λ_{max} 236 (log ϵ 4.12); 271 (log ϵ 3.65); 339 (log ϵ 4) (EtOH).

Phuong, N.M. *et al.*, *Planta Med.*, 1999, **65**, 761-762

Heimine**H-89**

[1357-41-1]

Struct. unknown


 $C_{26}H_{29}NO_5$ 435.519

Prob. a Lythraceae alkaloid (cf. Decaline in V-86). Alkaloid from *Heimia salicifolia* (Lythraceae). Mp 247.5-249°. $[\alpha]_D$ +43 ($CHCl_3$).

Blomster, R.N. *et al.*, *J. Nat. Prod.*, 1964, **27**, 15-24 (*isol, uv, ir*)

Heinsiagenin A**H-90**

[126594-31-8]

 $C_{36}H_{55}NO_4$ 565.835

Sapogenin from *Heinsia crinata*. Cryst. Mp 166-168°. $[\alpha]_D$ +138 (c, 1 in $CHCl_3$).

3-O- β -D-Xylopyranoside: Mussaendoside B

[136133-06-7]

 $C_{41}H_{63}NO_8$ 697.951Constit. of *Mussaenda pubescens*.

Cryst. Mp 194-197°. $[\alpha]_D$ +33.4 (c, 0.26 in MeOH). λ_{max} 265 (ϵ 5648) (MeOH).

3-O- β -D-Glucopyranoside: Mussaendoside A

[136133-05-6]

 $C_{42}H_{65}NO_9$ 727.977Constit. of *Mussaenda pubescens*.

Cryst. Mp 175-177°. $[\alpha]_D$ +60.2 (c, 0.44 in MeOH). λ_{max} 265 (ϵ 18633) (MeOH).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- β -D-xylopyranoside: Mussaendoside C

[136133-07-8]

 $C_{47}H_{73}NO_{13}$ 860.093Constit. of *Mussaenda pubescens*.

Cryst. Mp 186-189°. $[\alpha]_D$ +23 (c, 0.83 in

MeOH). λ_{max} 265 (ϵ 49135) (MeOH).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranoside: Mussaendoside D

[178402-86-3]

 $C_{48}H_{75}NO_{14}$ 890.119Constit. of *Mussaenda pubescens*.

Amorph. powder. $[\alpha]_D^{24}$ +30.2 (c, 0.54 in MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside: [158754-08-6] $C_{54}H_{85}NO_{18}$ 1036.261

Constit. of *Heinsia crinata*. Cryst. (as per-Ac). Mp 105-107° (per-Ac). $[\alpha]_D^{18}$ +21.43 (c, 0.12 in $CHCl_3$) (per-Ac).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside: Mussaendoside E

[178402-87-4]

 $C_{54}H_{85}NO_{19}$ 1052.261Constit. of *Mussaenda pubescens*.

Amorph. powder. $[\alpha]_D^{24}$ -1.4 (c, 0.03 in MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-xylopyranoside: Mussaendoside M

[136864-53-4]

 $C_{59}H_{93}NO_{21}$ 1152.378Constit. of *Mussaenda pubescens*.

Amorph. powder. Mp 178°. $[\alpha]_D$ +20.79 (c, 4.33 in MeOH). λ_{max} 270 (MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside: Mussaendoside O

[157536-44-2]

 $C_{60}H_{95}NO_{22}$ 1182.404Constit. of *Mussaenda pubescens*.

Amorph. powder. $[\alpha]_D^{15}$ +2.4 (c, 0.06 in Py). λ_{max} 265 (MeOH).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside: Mussaendoside F

[176108-51-3]

 $C_{60}H_{95}NO_{23}$ 1198.403Constit. of *Mussaenda pubescens*.

Amorph. powder. $[\alpha]_D^{30}$ -3.2 (c, 0.12 in Py). λ_{max} 265 (MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranosyl-(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-xylopyranoside: Mussaendoside N

[145396-78-7]

 $C_{65}H_{103}NO_{26}$ 1314.52Constit. of *Mussaenda pubescens*.

Amorph. powder. Mp 194°. $[\alpha]_D$ +19.6 (c, 3.5 in MeOH). λ_{max} 268 (MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranosyl-(1 \rightarrow 2)- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside: Mussaendoside G

[178468-01-4]

 $C_{66}H_{105}NO_{27}$ 1344.546Constit. of *Mussaenda pubescens*.

Amorph. powder. $[\alpha]_D^{24}$ +13.6 (c, 0.09 in MeOH).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 6)- $[\alpha$ -L-rhamno-

pyranosyl-(1→2)]-β-D-glucopyranosyl-(1→2)-β-D-glucopyranoside]: [158754-07-5]

C₆₆H₁₀₅NO₂₈ 1360.545

Constit. of *Heinsia crinata*. Cryst. (as per-Ac). Mp 123-124° (per-Ac). [α]_D¹⁸ +14.29 (c, 0.6 in CHCl₃) (per-Ac).

3-O-[β-D-Glucopyranosyl-(1→2)-β-D-glucopyranosyl-(1→6)]-α-L-rhamnopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→6)]-β-D-glucopyranoside]: **Mussaenoside U**

C₇₂H₁₁₅NO₃₂ 1506.688

Constit. of *Mussaenda pubescens*. Amorph. powder. [α]_D²⁵ -14 (c, 0.27 in MeOH). λ_{max} 265 (MeOH).

18-Hydroxy: 18-Hydroxyheinsigenin A
C₃₆H₅₅NO₅ 581.834

Sapogenin from *Mussaenda pubescens*.

18-Hydroxy, 3-O-[α-L-rhamnopyranosyl-(1→2)-β-D-glucopyranosyl-(1→2)]-α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Mussaenoside H**
[178402-88-5]

C₆₀H₉₅NO₂₃ 1198.403

Constit. of *Mussaenda pubescens*. Amorph. powder. [α]_D²⁴ +6.3 (c, 0.2 in Py).

Zhao, W. et al., *Nat. Prod. Sci.*, 1965, **1**, 61 (*Mussaenoside F*)

Babady-Bila, et al., *Tetrahedron*, 1989, **45**, 5907 (*Heinsigenin A*)

Xu, J.-P. et al., *Huaxue Xuebao*, 1991, **49**, 621 (*Mussaenosides A-C*)

Xu, J.-P. et al., *J. Nat. Prod.*, 1992, **55**, 1124 (*Mussaenosides*)

Zhao, W.M. et al., *Chin. Chem. Lett.*, 1994, **5**, 309 (*Mussaenoside O*)

Zhao, W. et al., *J. Nat. Prod.*, 1994, **57**, 1613 (*Mussaenoside O*)


Babady-Bila, et al., *Phytochemistry*, 1994, **36**, 1489 (*saponins*)

Zhao, W. et al., *Phytochemistry*, 1996, **42**, 827; 1131; 1997, **45**, 1073-1078 (*Mussaenosides*)

Helietidine

H-91

[476358-91-5]




C₁₉H₂₁NO₂ 295.38

Alkaloid from the stem bark of *Helietta longifoliata*. Needles. Mp 132-134°.

De Moura, N.F. et al., *Planta Med.*, 2002, **68**, 631-634 (*isol, pmr, cmr*)

Helindicine

H-92



Relative
Configuration

C₁₅H₂₃NO₄ 281.351

Alkaloid from the roots of *Heliotropium indicum*. Resin. [α]_D²⁰ -0.6 (c, 0.05 in MeOH).


Souza, J.S.N. et al., *J. Braz. Chem. Soc.*, 2005, **16**, 1410-1414 (*isol, pmr, cmr, ms*)

Heliospathuline

H-93

[135683-58-8]

[6224-43-7]



Absolute
Configuration

C₁₅H₂₅NO₅ 299.366

Struct. and stereochem. incorr. described in the paper owing to a typographical error and incorrect (*R,S*) assignments. The struct. shown appears to be correct. Alkaloid from *Heliotropium spathulatum*. Yellow oil. [α]_D²⁰ +3.2 (EtOH).

3'-Epimer: **Tessellatine**

C₁₅H₂₅NO₅ 299.366

Alkaloid from *Amsinckia douglasiana* and *Amsinckia tesellata* var. *gloriosa* (Boraginaceae). Oil. [α]_D²⁵ +2.4 (c, 0.45 in EtOH).

2',3'-Diepimer: **Isolycopamine**

C₁₅H₂₅NO₅ 299.366

Alkaloid from *Heliotropium keralense*. [α]_D +19.5 (EtOH).

Ravi, S. et al., *Phytochemistry*, 1990, **29**, 361-364 (*Isolycopamine*)

Roeder, E. et al., *Phytochemistry*, 1991, **30**, 1703-1706 (*isol, pmr, cmr, ms*)

Kelley, R.B. et al., *Phytochemistry*, 1992, **31**, 2513-2518 (*Tessellatine*)

Logie, C.G. et al., *Phytochemistry*, 1994, **37**, 43-109 (*rev, pmr*)


Heliosupine

H-94

Cynoglossophine

[32728-78-2]

[32779-96-7]



C₂₀H₃₁NO₇ 397.467

Alkaloid from *Heliotropium supinum* and *Cynoglossum* spp. (Boraginaceae). Shows antitumour activity. Sol. MeOH, EtOH, CHCl₃; poorly sol. H₂O. Mp 148-149°. [α]_D -4.3 (c, 5.1 in EtOH). CA Index Guide 2002 incorrectly names this compd. Asperumine; Asperumine applies to

Asperumine in T-188.

▶MH6000000

Picrate:

Cryst. + H₂O. Mp 102-106° (97-100°). Dehydrates to a gum on drying in *vacuo*.

N-Oxide: **Heliosupine N-oxide**

[31701-88-9]

C₂₀H₃₁NO₈ 413.467

Alkaloid from *Cynoglossum pictum* and *Paracynoglossum ineretinum* (preferred genus name *Cynoglossum*). Mp 165° dec.

3'-Ac: **Acetylheliosupine**

[31514-30-4]

C₂₂H₃₃NO₈ 439.505

Minor alkaloid from *Cynoglossum officinale* and from *Myosotis sylvatica* (Boraginaceae). [α]_D^{24.5} -1.8 (c, 0.567 in EtOH). Acetylated at the secondary OH group. Originally considered to be the 2'-Ac deriv. λ_{max} 217 (log ε 3.95) (MeOH).

3'-Ac, N-oxide: **3'-Acetylheliosupine N-oxide**

[147802-43-5]

C₂₂H₃₃NO₉ 455.504

Alkaloid from aerial parts of *Heliotropium hirsutissimum* (Boraginaceae). Oil. [α]_D²⁰ +14.9 (c, 0.35 in CHCl₃).

Denisova, S.I. et al., *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1953, **93**, 59-61; *CA*, **49**, 3992h (*isol*)

Culvenor, C.C.J. et al., *Aust. J. Chem.*, 1956, **9**, 512 (*struct*)

Crout, D.H.G. et al., *J.C.S.(C)*, 1966, 1968-1972; 1967, 1233-1234 (*biosynth*)

Pedersen, E. et al., *Dan. Tidsskr. Farm.*, 1970, **44**, 287-291; *CA*, **74**, 72780e (*3'-Ac, isol*)

Pedersen, G. et al., *Org. Mass Spectrom.*, 1970, **4**, 249-256 (*ms*)

Man'ko, I.V. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 523 (*occur, oxide*)

Zalkow, L.H. et al., *J. Nat. Prod.*, 1979, **42**, 612-623 (*isol*)

Smith, L.W. et al., *J. Nat. Prod.*, 1981, **44**, 129-152 (*rev, occur*)

Jones, A.J. et al., *Aust. J. Chem.*, 1982, **35**, 1173-1184 (*cmr*)


Resch, J.F. et al., *Phytochemistry*, 1982, **21**, 2430 (*3'-Ac, uv, pmr, cmr, ms, struct*)

Constantinidis, T. et al., *Phytochemistry*, 1993, **32**, 1335-1337 (*3'-Acetylheliosupine-N-oxide*)

Heliotrine

H-95

[303-33-3]



Absolute
configuration

C₁₆H₂₇NO₅ 313.393

Alkaloid from *Heliotropium europaeum*, *Heliotropium indicum* and *Heliotropium lasiocarpum* (Boraginaceae). Toxin, cause of chronic liver poisoning in animals. Shows ganglion-blocking activity and antineoplastic props. Shows antifungal

activity. Prisms (Me₂CO). Sol. H₂O. Mp 128° (125-126°). [α]_D²⁰ +63.8 (CHCl₃). [α]_D +17.6 (EtOH). Log P -0.39 (calc). The first pyrrolizidine alkaloid to be extensively studied. One of the crystallographic papers shows the wrong enantiomer.

- ▶ Toxic (less so than Lasiocarpine, L-47 or Retrorsine, R-62). Genotoxic agent. MH6125000

Methiodide: Mp 108-110° (softens at 103°).

N-Oxide: Heliotrine N-oxide

[6209-65-0]

C₁₆H₂₇NO₆ 329.392

Alkaloid present in *Heliotropium europaeum* (Boraginaceae). Mp 169.5-170.5° (synthetic). [α]_D +26.6 (EtOH) (synthetic).

- ▶ EK7879100

2'-Ac: 2'-O-Acetylheliotrine

[208662-67-3]


C₁₈H₂₉NO₆ 355.43

Alkaloid from *Heliotropium disciforme*.

- Men'shikov, G.P. *et al.*, *Ber.*, 1932, **65**, 974; 1933, **66**, 875; 1935, **68**, 1051; 1555 (*isol. props. struct*)
- Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1954, **7**, 277; 287; 1965, **18**, 1625 (*isol. struct. pmr*)
- Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*wv*)
- Pedersen, E. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 249 (*ms*)
- Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653 (*cd*)
- Wodak, S.J. *et al.*, *Acta Cryst. B*, 1975, **31**, 569 (*cryst. struct*)
- Zalkow, L.H. *et al.*, *J. Nat. Prod.*, 1979, **42**, 612 (*isol. cryst. struct*)
- Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)
- Marquina, G. *et al.*, *Pharmazie*, 1989, **44**, 870-871 (*Heliotrine, activity*)
- Farsam, H. *et al.*, *Pharm. Pharmacol. Lett.*, 1998, **8**, 79-80 (*2'-Acetylheliotrine*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HAL500

Heliotropamide

[646990-38-7]



Relative Configuration


C₃₆H₃₆N₂O₈ 624.689

Alkaloid from the aerial parts of *Heliotropium ovalifolium*. Amorph. powder. [α]_D²¹ +14 (c, 0.01 in MeOH). λ_{max} 223 (log ε 4.16); 287 (log ε 3.86); 327 (log ε 3.94) (MeOH).

Guntern, A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1550-1553 (*isol. pmr, cmr, ms*)

Hemiargine B

6,6a-Didehydro-11-hydroxy-1,2,10-trimethoxynoraporphine
[710948-55-3]



C₁₉H₁₉NO₄ 325.363


Alkaloid from leaves of *Croton hemiargyreus* var. *gymmodiscus*.

Lin, W.-H. *et al.*, *J. Chin. Pharm. Sci.*, 2003, **12**, 117-122 (*Hemiargine B*)

Hemiasterlin

Milnamide B

[157207-90-4]



R¹ = R² = CH₃

C₃₀H₄₆N₄O₄ 526.718

Peptide antibiotic. Isol. from the marine sponges *Auleta* sp., *Cymbastela* sp. and *Hemiasterella minor*. Cytotoxic. Microtubule formation inhibitor. Cryst. (MeOH/hexane). Mp 120-130°. [α]_D -95 (c, 0.06 in MeOH). λ_{max} 220 (ε 30000); 270 (ε 9000); 295 (ε 5000) (MeOH) (Derep).

N¹-De-Me: Hemiasterlin A

[169181-24-2]

C₂₉H₄₄N₄O₄ 512.691

Isol. from sponges *Cymbastela* sp. and *Siphonochalina* sp. Cytotoxic. Amorph. solid. Sol. MeOH. [α]_D -45 (c, 0.25 in MeOH). λ_{max} 218 (ε 23400); 280 (ε 2800) (MeOH).

Talpir, R. *et al.*, *Tet. Lett.*, 1994, **35**, 4453-4456 (*isol. pmr, cmr, ms*)

Coleman, J.E. *et al.*, *Tetrahedron*, 1995, **51**, 10653-10662 (*Hemiasterlin A*)

Coleman, J.E. *et al.*, *Acta Cryst. C*, 1996, **52**, 1525-1527 (*cryst. struct*)

Gamble, W.R. *et al.*, *Bioorg. Med. Chem.*, 1999, **7**, 1611-1615 (*isol. activity*)

Vedejs, E. *et al.*, *J.O.C.*, 2001, **66**, 7355-7364 (*synth. pmr, cmr*)

Nieman, J.A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 183-199 (*synth. pmr, cmr, activity*)

Chevallier, C. *et al.*, *Org. Lett.*, 2003, **5**, 3737-3739 (*isol. activity*)

Sonnenschein, R. *et al.*, *Org. Lett.*, 2004, **6**, 779-782 (*isol. activity*)

Hemiasterlin C

[246847-61-0]

As Hemiasterlin, H-98 with

R¹ = CH₃, R² = H

C₂₉H₄₄N₄O₄ 512.691


Isol. from the sponge *Siphonochalina* sp. Cytotoxic agent. [α]_D -18.8 (c, 0.11 in MeOH).

H-97

Gamble, W.R. *et al.*, *Bioorg. Med. Chem.*, 1999, **7**, 1611-1615

Hemibastadin 1

[134981-79-6]



C₁₇H₁₆Br₂N₂O₄ 472.132

Numbering systems vary. Minor constituent of the Australian marine sponge *Ianthella basta*. Solid.

4-Sulfate: 4-O-Sulfohemibastadin 1. 1-O-Sulfohemibastadin 1

C₁₇H₁₆Br₂N₂O₇S 552.197

Constit. of *Ianthella basta*. Amorph. solid (as Na salt).

4'-Me ether: 4'-O-Methylhemibastadin 1. 1'-Methoxyhemibastadin 1

[182496-48-6]

C₁₈H₁₈Br₂N₂O₄ 486.159

Constit. of *Ianthella basta*. Oil. λ_{max} 213 (log ε 4.53); 280 (log ε 3.87) (MeOH).

5-Bromo: Hemibastadin 3

[182806-14-0]

C₁₇H₁₅Br₃N₂O₄ 551.029

Constit. of *Ianthella basta*. Isol. as a 3:1 inseparable mixt. with Hemibastadin 2.

5'-Bromo: Hemibastadin 2

[134981-80-9]

C₁₇H₁₅Br₃N₂O₄ 551.029

Constit. of *Ianthella basta*. Solid.

5'-Bromo, 4-sulfate: 4-O-Sulfohemibastadin 2. 1-O-Sulfohemibastadin 2

C₁₇H₁₅Br₃N₂O₇S 631.093

Constit. of *Ianthella basta*. Amorph. solid. λ_{max} 204 ; 282 (MeOH).

5'-Bromo, 4'-Me ether: 4'-O-Methylhemibastadin 2. 1'-Methoxyhemibastadin 2

[182496-49-7]

C₁₈H₁₇Br₃N₂O₄ 565.055

Constit. of *Ianthella basta*. Solid. λ_{max} 213 (log ε 4.5); 281 (log ε 3.87) (MeOH).

Butler, M.S. *et al.*, *Aust. J. Chem.*, 1991, **44**, 287-296 (*isol. pmr, ms, struct*)


Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1996, **59**, 927-934 (*4'-O-Methylhemibastadins, Hemibastadin 3*)

Wasserman, H.H. *et al.*, *J.O.C.*, 1998, **63**, 5581-5586 (*Hemibastadin 2, synth*)

Masuno, M.N. *et al.*, *Mar. Drugs*, 2004, **2**, 176-184 (*4-Sulfohemibastadins*)

Hemibastadinol 1

[182806-09-3]



C₁₇H₁₇Br₂N₂O₄ 459.134

Isol. from the Papua New Guinea marine sponge *Ianthella basta*. Solid. [α]_D²³ -31 (c,

1.83 in MeOH). λ_{\max} 208 (log ϵ 4.43); 281 (log ϵ 3.69) (MeOH).

5'-Bromo: Hemibastadinol 2

[182806-11-7]

$C_{17}H_{16}Br_3NO_4$ 538.03

From *Ianthella basta*. Oil. $[\alpha]_D^{23}$ -24 (c, 0.10 in MeOH). Inseparable mixt. with Hemibastadinol 3, to which data refer.

5-Bromo: Hemibastadinol 3

[182806-12-8]


$C_{17}H_{16}Br_3NO_4$ 538.03

From *Ianthella basta*. Inseparable mixt. with Hemibastadinol 2.

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1996, **59**, 927 (isol, uv, ir, pmr, cmr, ms, struct)

Hemicalyculin A

H-102



$C_{36}H_{55}N_2O_{10}P$ 706.812


Isol. from the sponge *Discodermia calyx*. Amorph. solid. $[\alpha]_D^{20}$ -107 (c, 0.5 in MeOH). λ_{\max} 237 (ϵ 3700); 340 (ϵ 19000) (MeOH).

Wakimoto, T. *et al.*, *Chem. Biol.*, 2002, **9**, 309-319 (isol, pmr, cmr)

Hemifistularin 3

H-103

[153415-35-1]



$C_{18}H_{16}Br_4N_2O_6$ 675.95


Metab. from a new sp. of sponge of the family Aplysinnellidae, order Verongida, from the Coral Sea. Powder (hexane/Et₂O). Mp 73-75°. $[\alpha]_D^{20}$ +110 (c, 0.2 in Me₂CO). λ_{\max} 207 (ϵ 39000); 281 (ϵ 6900) (MeOH) (Berdy).

Mancini, I. *et al.*, *J.C.S. Perkin 1*, 1993, 3121 (isol, uv, ir, pmr, cmr, struct)

Hemsine A

H-104

[530092-42-3]



$C_{32}H_{39}N_5O_4$ 557.691


Alkaloid from the roots of *Paliurus hemsleyanus*. $[\alpha]_D^{26}$ -64.5 (c, 2 in MeOH). λ_{\max} 224 (log ϵ 4.96); 272 (log ϵ 4.3); 280 (log ϵ 4.3); 290 (log ϵ 4.21) (MeOH).

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (isol, pmr, cmr)

Hemsine B

H-105

[530092-43-4]



$C_{36}H_{49}N_5O_5$ 631.814

Alkaloid from the roots of *Paliurus hemsleyanus*. $[\alpha]_D^{26}$ -124 (c, 1 in MeOH). λ_{\max} 217 (log ϵ 4.68); 257 (sh) (log ϵ 4.14); 285 (log ϵ 3.92) (MeOH).

N-De-Me: Ramosine B. N-Demethyl-hemsine B

[530092-40-1]

$C_{35}H_{47}N_5O_5$ 617.787


Alkaloid from the roots of *Paliurus ramosissimus*. $[\alpha]_D^{26}$ -181.5 (c, 2 in MeCN). λ_{\max} 216 (log ϵ 4.6); 233 (log ϵ 4.43); 329 (log ϵ 3.57) (MeCN).

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (isol, pmr, cmr)

Hemsine C

H-106

[530092-44-5]



$C_{41}H_{48}N_6O_5$ 704.867

Alkaloid from the roots of *Paliurus hemsleyanus*. $[\alpha]_D^{26}$ -107 (c, 1 in MeOH).


λ_{\max} 214 (log ϵ 4.28); 280 (log ϵ 4.31) (MeOH).

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (isol, pmr, cmr)

Hemsine D

H-107

[530092-45-6]



$C_{29}H_{38}N_4O_4$ 506.644


Alkaloid from the roots of *Paliurus hemsleyanus*. $[\alpha]_D^{26}$ -573.3 (c, 0.75 in CHCl₃). λ_{\max} 217 (log ϵ 4.7); 230 (log ϵ 3.66); 255 (log ϵ 3.4) (MeOH).

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (isol, pmr, cmr)

Henderine

H-108

[102686-11-3]



$C_{20}H_{15}NO_6$ 365.342


Alkaloid from the whole plants of *Corydalis hendersonii* (*Corydalis nepalensis*) (Papaveraceae).

Lin, M. *et al.*, *Zhivui Xuebao (Acta Bot. Sin.)*, 1986, **28**, 91; *CA*, **105**, 3497q

5-(6-Heneicosenyl)-1H-pyrrole-2-carboxaldehyde, 9CI

H-109

2-Formyl-5-(6-heneicosenyl)-1H-pyrrole



$C_{26}H_{45}NO$ 387.648


(Z)-form [262351-88-2]

Isol. from *Mycale mytilorum*. Pale yellow flakes. Mp 40-43°. λ_{\max} 302 (ϵ 16000) (EtOH).

Reddy, G.B.S. *et al.*, *Bioorg. Med. Chem.*, 2000, **8**, 27-36 (Z-form, ir, ms, pmr, cmr)

Henningsamide

[141544-34-5]

Relative
ConfigurationC₂₁H₂₂N₂O₄ 366.416Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -368 (c, 0.5 in MeOH). λ_{max} 203 ; 242 ; 285 ; 308 (sh) (MeOH).N-De(hydroxyacetyl): **Deshydroxyacetyl-henningsamide**


[141544-35-6]

C₁₉H₂₀N₂O₂ 308.379Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -744 (c, 0.33 in MeOH). λ_{max} 223 ; 287 ; 295 (MeOH).O-Ac: **O-Acetylhenningsamide**

[141565-03-9]

C₂₃H₂₄N₂O₅ 408.453Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -104 (c, 0.5 in CHCl₃). λ_{max} 203 ; 242 ; 284 ; 309 (sh) (MeOH).Massiot, G. et al., *Phytochemistry*, 1991, 30, 3449-3456 (*isol, struct*)**Henningsiine**

[141544-27-6]

Absolute
ConfigurationC₂₁H₂₄N₂O₃ 352.432Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -24 (c, 1 in CHCl₃). CA numbering shown.N^b-Oxide: **Henningsiine N-oxide**

[141544-31-2]

C₂₁H₂₄N₂O₄ 368.432Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -9 (c, 0.35 in CHCl₃).O-Ac: **O-Acetylhenningsiine**

[141544-29-8]

C₂₃H₂₆N₂O₄ 394.469Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -22 (c, 1.3 in CHCl₃).N-De(hydroxyacetyl): **Deshydroxyacetyl-henningsiine**

[141544-28-7]

C₁₉H₂₂N₂O 294.396Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -22 (c, 1.3 in CHCl₃).16-Hydroxy: **16-Hydroxyhenningsiine. 3-Hydroxyhenningsiine**

[141544-30-1]

C₂₁H₂₄N₂O₄ 368.432Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D²⁰ -29 (c, 0.35 in


H-110

CHCl₃).Massiot, G. et al., *Phytochemistry*, 1991, 30, 3449-3456 (*isol, struct*)

Not indexed by CA.

Higa, T. et al., *Pure Appl. Chem.*, 1994, 66, 2227-2230**Hennoxazole A**

[132564-95-5]

C₂₉H₄₂N₂O₆ 514.661Alkaloid from the marine sponge *Polyfibrospongia* sp. Possesses antiviral and peripheral analgesic activities. Light yellow oil. [α]_D²⁵ -47 (c, 3.12 in CHCl₃) (-42.7). λ_{max} 254 (ε 12000) (MeOH) (Derep).4-Ac: **4-O-Acetylhennoxazole A**C₃₁H₄₄N₂O₇ 556.698Alkaloid from a *Polyfibrospongia* sp. CAS no. not found 8-14CI.O²-De-Me: **Hennoxazole E**C₂₈H₄₀N₂O₆ 500.634Alkaloid from *Polyfibrospongia* sp. CAS no. not found 8-14CI.O²-De-Me, O²-Et: **Hennoxazole B**

[132564-96-6]

C₃₀H₄₄N₂O₆ 528.687Alkaloid from *Polyfibrospongia* sp. Analgesic. λ_{max} 254 (ε 12000) (MeOH) (Derep).O²-De-Me, O²-butyl: **Hennoxazole C**


[132564-97-7]

C₃₂H₄₈N₂O₆ 556.741Alkaloid from *Polyfibrospongia* sp. λ_{max} 254 (ε 12000) (MeOH) (Derep).4-Deoxy: **Hennoxazole D**

[132564-98-8]

C₂₉H₄₂N₂O₅ 498.661From *Polyfibrospongia* sp. Light yellow oil. λ_{max} 254 (ε 12000) (MeOH) (Derep).Ichiba, T. et al., *J.A.C.S.*, 1991, 113, 3173-3174 (*isol, struct*)Higa, T. et al., *Pure Appl. Chem.*, 1994, 66, 2227-2230 (4-Acetylhennoxazole A, Hennoxazole E)Williams, D.R. et al., *J.A.C.S.*, 1999, 121, 4924-4925 (*synth*)Yokokawa, F. et al., *Tetrahedron*, 2001, 57, 6311-6327 (*synth, ir, pmr, cmr, ms*)Smith, T.E. et al., *J.O.C.*, 2008, 73, 142-150 (*synth*)**Hennoxazole F**


H-113

C₂₈H₃₈N₂O₅ 482.619Alkaloid from the sponge *Polyfibrospongia* sp. Not indexed by CA.5',6'-Dihydro, 5'-hydroxy: **Hennoxazole G**C₂₈H₄₀N₂O₆ 500.634Alkaloid from *Polyfibrospongia* sp.

H-112

Henrycinol A


[606925-22-8]

Relative
ConfigurationC₂₇H₂₈N₂O₄ 444.529Alkaloid from the roots of *Melodinus henryi*. Amorph. solid. [α]_D²⁰ -43.7 (c, 0.4 in CHCl₃). λ_{max} 232 (log ε 2.6); 278 (log ε 2.4); 280 (log ε 2.8); 290 (log ε 2.3) (EtOH).15-(2-Methylpropanoyl): **Henrycinol B**

[606925-23-9]

C₃₁H₃₄N₂O₅ 514.62Alkaloid from the roots of *Melodinus henryi*. Amorph. powder. [α]_D²⁰ -60.7 (c, 0.7 in CHCl₃). λ_{max} 232 (log ε 2.4); 278 (log ε 2.9); 290 (log ε 2.6) (EtOH).Zhang, Y.W. et al., *Helv. Chim. Acta*, 2003, 86, 415-419 (*isol, pmr, cmr, ms*)**2,3,3',4,4',5,5'-Heptachloro-1'-methyl-1,2'-bi-1H-pyrrole Q1**

H-115


C₉H₃Cl₇N₂ 387.305

Widespread in natural sources. Distribution pattern resembles that of environmental contaminants, e.g. PCBs, but appears to be a genuine nat. prod. Cryst. Mp 154-155.5°. Readily oxidises and polymerises.

Wu, J. et al., *Angew. Chem., Int. Ed.*, 2002, 41, 1740-1743 (*synth, cryst struct*)Vetter, W. et al., *Chemosphere*, 2003, 52, 415-422 (*isol, pmr, cmr*)Vetter, W. et al., *Rapid Commun. Mass Spectrom.*, 2006, 20, 3018-3022 (*cmr, occur*)Vetter, W. et al., *Rev. Environ. Contam. Toxicol.*, 2006, 188, 1-57 (*rev, occur*)**1,9-Heptadecadiene-4,6-diyne-3,8-diol, 9CI**

H-116

[30779-95-4]



(3R,8S,9Z)-form


C₁₇H₂₄O₂ 260.375
Log P 2.49 (calc).**(3R,8S,9Z)-form
Falcavindiol**
[55297-87-5]

Constit. of roots of *Aegopodium podagraria*, *Angelica acutiloba* (Dong Dang Gui), *Angelica japonica*, *Anthriscus sylvestris*, *Aralia cordata* (udo), *Crithmum maritimum* (rock samphire), *Daucus carota* (carrot), *Dendropanax arboreus*, *Falcaria vulgaris*, *Glehnia littoralis*, *Glehnia littoralis leiocarpa*, *Heteromorpha trifoliolata*, *Levisticum officinale* (lovage), *Oenanthe javanica* (water dropwort), *Osmorhiza occidentalis*, *Peucedanum orioselinum*, *Schefflera digitata*, *Apium graveolens* (celery), *Foeniculum vulgare* (fennel), *Petroselinum crispum* (parsley) and *Pastinaca sativa* (parsnip). Constit. of Chinese crude drug Toki. Has antifungal, antibiotic and analgesic activity. Antinociceptive agent. DNA topoisomerase inhibitor. Shows phytotoxic, allelochemical, antimutagenic and antiproliferative props. Pale yellow oil. Sol. MeOH, Et₂O. $[\alpha]_D^{20} +204.8$ (c, 1 in CHCl₃). Log P 2.49 (calc). Pharmacol. active isomer. λ_{\max} 231 ; 244 ; 256 (MeOH) (Berdy). λ_{\max} 233 (€ 1000); 245 (€ 1000); 259 (€ 800) (EtOH) (Berdy). λ_{\max} 232 (€ 400); 244 (€ 400); 258 (€ 200) (Et₂O) (Berdy).

8-O-(1H-Indol-3-ylacetyl): **8-(1H-Indol-3-ylacetyl)falcarindiol** [945857-69-2]
C₂₇H₃₁NO₃ 417.547
Alkaloid from the flower buds of *Hedera rhombea*. Oil. $[\alpha]_D^{26} +43$ (c, 0.4 in CHCl₃). λ_{\max} 221 (log € 4.4); 271 (log € 4); 281 (log € 3.9); 289 (log € 3.9) (Et₂O).

Yamazoe, S. et al., *Phytochemistry*, 2007, **68**, 1706-1711 (*indole-3-acetate*)


14-Heptadecene-4,16-dienoic acid H-117



(Z)-form

2-Phenylethylamide: N-(2-Phenylethyl)-14-heptadecene-4,16-dienamide. **Callyspongamide A** [549494-83-9]
C₂₅H₃₃NO 363.542
Alkaloid from the Red Sea sponge *Callyspongia fistularis*. Cytotoxic. Light yellow oil. λ_{\max} 216 (log € 4.12) (MeOH).
Youssef, D.T.A. et al., *J. Nat. Prod.*, 2003, **66**, 861-862 (*isol, pmr, cmr*)

1,2,3,4,6,9,14-Heptahydroxydihydro-β-agarofuran H-118



(1α,2α,3α,4βOH,6β,9β)-form

C₁₅H₂₆O₈ 334.366

(1α,2α,3α,4βOH,6β,9β)-form

9-(3-Furancarboxyl), 3-(2-methylbutanoyl), 1,2,6,14-tetra-Ac: [828921-28-4]
C₃₃H₄₄O₁₅ 680.702
Constit. of *Euonymus nanoides*. Gum. $[\alpha]_D^{20} +26$ (c, 1.1 in CHCl₃). λ_{\max} 242 (MeOH).

(1α,2α,3β,4βOH,6β,9β)-form Maytol

9-(3-Pyridinecarboxyl), 1,2,6,14-tetra-Ac: **Maytoline** [31146-55-1]
C₂₉H₃₇NO₁₃ 607.61
Alkaloid from the fruit of *Maytenus serrata* (*Maytenus ovatus*) (Celastraceae). Noncryst. $[\alpha]_D^{25} +0.3$ (c, 0.75 in CHCl₃).

9-(3-Pyridinecarboxyl), 1,2,6,14-tetra-Ac: **methiodide**: [31146-59-5]
Cryst. (MeOH/Et₂O). Cryst. unstable in air and rapidly loses solvent of crystallisation to give a powder. Mp 190-193° dec.


9-(3-Pyridinecarboxyl), 6-benzoyl, 1,2,3,14-tetra-Ac: **Maytolidine** [60512-69-8]
C₃₆H₄₁NO₁₄ 711.718
Alkaloid from the fruit of *Maytenus serrata* (Celastraceae). Cryst. (EtOAc/Et₂O). Mp 128-132°.

Bryan, R.F. et al., *J.C.S.(B)*, 1971, 2159 (*cryst struct*)

Kupchan, S.M. et al., *J.O.C.*, 1977, **42**, 115-118 (*Maytoline, Maytolidine*)

Liu, Z.-L. et al., *Planta Med.*, 2004, **70**, 353-358 (*Euonymus nanoides constiti*)

1,2,4,6,8,9,14-Heptahydroxydihydro-β-agarofuran H-119



(1α,2α,4β,6β,8α,9α)-form

C₁₅H₂₆O₈ 334.366

(1α,2α,4β,6β,8α,9α)-form

8-O-(3-Pyridinecarboxyl), 9-benzoyl, 2-O-(2-methylpropanoyl), 1,14-di-Ac: **1,14-Diacetoxy-9-benzoyloxy-4,6-dihydroxy-2-isobutanoyloxy-8-nicotinoyloxydihydro-β-agarofuran** [117714-46-2]
C₃₆H₄₃NO₁₃ 697.735
Alkaloid from *Celastrus angulatus* (Celastraceae). Insect antifeedant. Amorph. $[\alpha]_D^{20} +43.4$ (c, 0.52 in MeOH).

(1α,2α,4β,6β,8α,9β)-form [455260-81-8]

Isol. from *Celastrus angulatus*. Cryst. Mp 177-179°.

1-O-(3-Pyridinecarboxyl), 9-O-(3-furancarboxyl), 14-O-(2-methylpropanoyl), 2,6,8-tri-Ac: [864179-26-0]
C₃₆H₄₃NO₁₅ 729.733
Constit. of *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{25} +9.2$ (c, 1.2 in MeOH). λ_{\max} 216 (log € 3.97); 255 (log € 3.56) (MeOH).

1-O-(3-Pyridinecarboxyl), 9-benzoyl, 14-

O-(2-methylbutanoyl), 8-Ac: [862703-02-4]
C₃₅H₄₃NO₁₂ 669.724
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +3.6$ (c, 0.42 in CHCl₃). λ_{\max} 223 ; 264 (EtOH).

1-O-(3-Pyridinecarboxyl), 9-benzoyl, 14-O-(2-methylbutanoyl), 2,8-di-Ac: [862703-01-3]

C₃₇H₄₅NO₁₃ 711.761
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +9.1$ (c, 0.11 in CHCl₃). λ_{\max} 224 ; 264 (EtOH).

1-O-(3-Pyridinecarboxyl), 9-benzoyl, 14-O-(2-methylbutanoyl), 6,8-di-Ac: [862703-00-2]

C₃₇H₄₅NO₁₃ 711.761
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +9.6$ (c, 0.91 in CHCl₃). λ_{\max} 221 ; 264 (EtOH).

2-O-(3-Pyridinecarboxyl), 9-benzoyl, 14-O-(2-methylbutanoyl), 1,8-di-Ac: [862702-98-5]

C₃₇H₄₅NO₁₃ 711.761
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +42.9$ (c, 0.24 in CHCl₃). λ_{\max} 224 ; 264 (EtOH).

2-O-(3-Pyridinecarboxyl), 9-benzoyl, 14-O-(2-methylbutanoyl), 6,8-di-Ac: [862702-99-6]

C₃₇H₄₅NO₁₃ 711.761
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +43.2$ (c, 0.19 in CHCl₃). λ_{\max} 223 ; 264 (MeOH).

2-O-(3-Pyridinecarboxyl), 1,9-dibenzoyl, 14-O-(2-methylbutanoyl), 8-Ac: [862703-03-5]

C₄₂H₄₇NO₁₃ 773.832
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +59.4$ (c, 0.17 in CHCl₃). λ_{\max} 227 ; 264 (EtOH).

8-O-(3-Pyridinecarboxyl), 9-benzoyl, 1,2,6,14-tetra-Ac:

C₃₆H₄₁NO₁₄ 711.718
Constit. of *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{25} -38.4$ (c, 1 in MeOH). λ_{\max} 229 (log € 4.33); 263 (log € 3.64) (MeOH).

9-O-(3-Pyridinecarboxyl), 2-O-(3-furancarboxyl), 8,14-bis-O-(2-methylpropanoyl), 1,6-di-Ac: [148461-95-4]

C₃₈H₄₇NO₁₅ 757.787
Constit. of *Celastrus angulatus*. Amorph. $[\alpha]_D^{20} +45.8$ (c, 0.52 in MeOH).

1,2-Bis-O-(3-pyridinecarboxyl), 9-benzoyl, 14-O-(2-methylbutanoyl), 8-Ac: [862702-97-4]

C₄₁H₄₆N₂O₁₃ 774.82
Constit. of *Crossopetalum tonduzii*. Glass. $[\alpha]_D^{25} +50.5$ (c, 1.57 in CHCl₃). λ_{\max} 222 ; 263 (EtOH).

1,2-Bis-O-(3-pyridinecarboxyl), 8,9-dibenzoyl, 6,14-di-Ac: **Cangorin E** [152340-58-4]

C₄₅H₄₄N₂O₁₄ 836.848
Constit. of *Maytenus ilicifolia*. Amorph. solid. Mp 103-108°. $[\alpha]_D +26.3$ (c, 0.19 in CHCl₃).

1,8-Bis-O-(3-pyridinecarboxyl), 6,9-dibenzoyl, 2,14-di-Ac: **Cangorin I** [156400-97-4]

C₄₅H₄₄N₂O₁₄ 836.848

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 111-115°. $[\alpha]_D$
+14.8 (c, 0.58 in CHCl_3).

2,8-Bis-O-(3-pyridinecarbonyl), 9-benzoyl, 1,14-di-Ac: **Cangorin F**
[156400-94-1]

$\text{C}_{38}\text{H}_{40}\text{N}_2\text{O}_{13}$ 732.74

Constit. of *Maytenus ilicifolia*. Cryst.
Mp 214-217°. $[\alpha]_D$ +66 (c, 0.59 in
 CHCl_3).

2,8-Bis-O-(3-pyridinecarbonyl), 9-benzoyl, 1,6,14-tri-Ac: **Cangorin D**
[152340-57-3]

$\text{C}_{40}\text{H}_{42}\text{N}_2\text{O}_{14}$ 774.777

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 123-138°. $[\alpha]_D$
+11.2 (c, 0.2 in CHCl_3).

2,8-Bis-O-(3-pyridinecarbonyl), 6,9-dibenzoyl, 1,14-di-Ac: **Cangorin C**
[152340-56-2]

$\text{C}_{45}\text{H}_{44}\text{N}_2\text{O}_{14}$ 836.848

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 122-126°. $[\alpha]_D$ -45
(c, 0.36 in CHCl_3).

1,2,8-Tris-O-(3-pyridinecarbonyl), 9-benzoyl, 14-Ac: **Cangorin G**
[156400-95-2]

$\text{C}_{42}\text{H}_{41}\text{N}_3\text{O}_{13}$ 795.798

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 124-128°. $[\alpha]_D$
+29.4 (c, 1.11 in CHCl_3).

1,2,8-Tris-O-(3-pyridinecarbonyl), 9-benzoyl, 6,14-di-Ac: **Cangorin H**
[156400-96-3]

$\text{C}_{44}\text{H}_{43}\text{N}_3\text{O}_{14}$ 837.835

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 125-130°. $[\alpha]_D$
+43.4 (c, 0.45 in CHCl_3).

1,2,8-Tris-O-(3-pyridinecarbonyl), 6,9-dibenzoyl, 14-Ac: **Cangorin B**
[152340-55-1]

$\text{C}_{49}\text{H}_{45}\text{N}_3\text{O}_{14}$ 899.906

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 137-141°. $[\alpha]_D$
+77.3 (c, 0.33 in CHCl_3).

2,6,8-Tris-O-(3-pyridinecarbonyl), 9-benzoyl, 1,14-di-Ac: **Cangorin A**
[152340-54-0]

$\text{C}_{44}\text{H}_{43}\text{N}_3\text{O}_{14}$ 837.835

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 136-140°. $[\alpha]_D$
+45.7 (c, 0.37 in CHCl_3).

(1 α ,2 α ,4 β ,6 β ,8 β ,9 α)-form [120693-69-8]
Constit. of *Celastrus angulatus*.

8-O-(3-Pyridinecarbonyl), 9-benzoyl, 2-O-(2-methylpropanoyl), 1,14-di-Ac:
[141544-40-3]

$\text{C}_{36}\text{H}_{43}\text{NO}_{13}$ 697.735

Constit. of *Celastrus angulatus*.
Amorph. $[\alpha]_D^{20}$ +43.4 (c, 0.52 in
MeOH).

Ji-Kai, L. et al., *Phytochemistry*, 1991, **30**,
3437-3448 (*Celastrus angulatus* constits)

Itokawa, H. et al., *J. Nat. Prod.*, 1993, **56**,
1479-1485; 1994, **57**, 460-470 (*Cangorins*,
cryst struct)


Tincusi, B.M. et al., *J. Nat. Prod.*, 1998, **61**,
1520-1523 (*Crossopetalum tonduzii* esters)

Jiménez, I.A. et al., *J. Nat. Prod.*, 2003, **66**,
1047-1050 (*Crossopetalum tonduzii* esters)

Mendoza, C.R. et al., *Chem. Biodiversity*,
2005, **2**, 286-294 (*Crossopetalum tonduzii*
esters)

2,3,5,7,8,9,15-Heptahydroxy-6(17),11-jatropha-dien-14-one

H-120



(2 α ,3 β ,5 α ,7 β ,8 α ,9 α ,11E,15 α)-form

$\text{C}_{20}\text{H}_{32}\text{O}_8$ 400.468

(2 α ,3 β ,5 α ,7 β ,8 α ,9 α ,11E,15 β)-form

2-O-(3-Pyridinecarbonyl), 5-O-(2-methylbutanoyl), 7,8,9-tri-Ac: [250293-36-8]

$\text{C}_{37}\text{H}_{49}\text{NO}_{13}$ 715.793

Constit. of *Euphorbia obtusifolia*. Oil.
 $[\alpha]_D$ -6 (c, 0.68 in CHCl_3).

2-O-(3-Pyridinecarbonyl), 5-benzoyl, 3,7,8,9-tetra-Ac: **Guyonianin A**
[936020-20-1]

$\text{C}_{41}\text{H}_{47}\text{NO}_{14}$ 777.821

Constit. of *Euphorbia guyoniana*.

3-O-(3-Pyridinecarbonyl), 7-O-(2-methylpropanoyl), 2,5,8,9-tetra-Ac:
Euphodendroidin G

[578738-22-4]

$\text{C}_{38}\text{H}_{49}\text{NO}_{14}$ 743.803

Constit. of *Euphorbia dendroides*.
Amorph. solid. $[\alpha]_D^{25}$ +3.5 (c, 0.1 in
 CHCl_3).

5-O-(3-Pyridinecarbonyl), 7-benzoyl, 2,8,9-tri-Ac: **Euphodendroidin C**
[578738-18-8]

$\text{C}_{39}\text{H}_{45}\text{NO}_{13}$ 735.783

Constit. of *Euphorbia dendroides*.
Amorph. solid. $[\alpha]_D^{25}$ -5 (c, 0.1 in
 CHCl_3).

2,9-Bis-O-(3-pyridinecarbonyl), 5-O-(2-methylpropanoyl), 3,7,8,15-tetra-Ac:
Euphodendroidin I

[578738-24-6]

$\text{C}_{44}\text{H}_{52}\text{N}_2\text{O}_{15}$ 848.899

Constit. of *Euphorbia dendroides*.
Amorph. solid. $[\alpha]_D^{25}$ -1.3 (c, 0.1 in
 CHCl_3).


Marco, J.A. et al., *Phytochemistry*, 1999, **52**,
479-485 (*Euphorbia obtusifolia* esters)

Corea, G. et al., *J. Med. Chem.*, 2003, **46**,
3395-3402 (*Euphodendroidins*)

Ahmed, A.A. et al., *Nat. Prod. Commun.*,
2006, **1**, 273-279; *CA*, **146**, 478757d
(*Guyonianin A*)

3,5,7,8,9,13,15-Heptahydroxy-6(17),11-jatropha-dien-14-one

H-121



$\text{C}_{20}\text{H}_{32}\text{O}_8$ 400.468

(2 β ,3 β ,5 α ,7 β ,8 α ,9 α ,11E,13 α ,15 β)-form

9-(3-Pyridinecarbonyl), 3,7-diangeloyl, 8-Ac: **Amygdaloidin B**
[853007-72-4]

$\text{C}_{38}\text{H}_{49}\text{NO}_{12}$ 711.805

Constit. of *Euphorbia amygdaloides*.
Amorph. solid. $[\alpha]_D^{25}$ +117.7 (c, 0.1 in
 CHCl_3).

9-(3-Pyridinecarbonyl), 7,8-diangeloyl, 3,5-di-Ac: **Amygdaloidin A**
[853007-71-3]

$\text{C}_{40}\text{H}_{51}\text{NO}_{13}$ 753.842

Constit. of *Euphorbia amygdaloides*.
Amorph. solid. $[\alpha]_D^{25}$ -22.17 (c, 0.1 in
 CHCl_3).

9-(3-Pyridinecarbonyl), 3-(3-hydroperoxy-2-methylenebutanoyl), 7-angeloyl, 8-Ac: **Amygdaloidin C**
[853007-73-5]


$\text{C}_{38}\text{H}_{49}\text{NO}_{14}$ 743.803

Constit. of *Euphorbia amygdaloides*.
Amorph. solid. $[\alpha]_D^{25}$ -16.67 (c, 0.1 in
 CHCl_3).

Corea, G. et al., *Tetrahedron*, 2005, **61**, 4485-
4494 (*Amygdaloidins A-L*)

1,3,6,13,14,16,18-Heptahydroxy-4-methylaconitan-15-one

H-122



(1 α ,3 α ,5 β ,6 α ,14 α ,16 α)-form

$\text{C}_{19}\text{H}_{27}\text{NO}_8$ 397.424

(1 α ,3 α ,5 β ,6 α ,14 α ,16 α)-form

$\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -Tetra-Me, N-Me, 14-benzoyl: **16-Epipyromesaconitine**
[123827-18-9]

$\text{C}_{31}\text{H}_{41}\text{NO}_9$ 571.666

Alkaloid from the the crude drug bushi
obt. from the roots of some *Aconitum*
spp. (Ranunculaceae). Amorph. $[\alpha]_D$ -
79.5 (c, 0.44 in EtOH). Artifact.

▶ AR5574000

$\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -Tetra-Me, N-Et, 14-benzoyl: **16-Epipyroaconitine**
[123834-18-4]

$\text{C}_{32}\text{H}_{43}\text{NO}_9$ 585.693

Alkaloid from the the crude drug bushi
obt. from the roots of some *Aconitum*
spp. (Ranunculaceae). Needles. Mp
166-167° (96-98°). $[\alpha]_D$ -64.3 (c, 0.23 in
EtOH). Artifact.

▶ AR5570000

$\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl): **16-Epipyroresaconitine**
[123930-58-5]

$\text{C}_{33}\text{H}_{45}\text{NO}_{10}$ 615.719

Alkaloid from the the crude drug bushi
obt. from the roots of some *Aconitum*
spp. (Ranunculaceae). Needles. Mp
181-182°. $[\alpha]_D$ -58.9 (c, 0.55 in EtOH).
Artifact.

▶ AR5582000

(1 α ,3 α ,5 β ,6 α ,14 α ,16 β)-form

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Me*, 14-*benzoyl*: **Pyromesaconitine**
[121707-41-3]
C₃₁H₄₁NO₉ 571.666
Alkaloid from the the crude drug bushi obt. from the roots of some *Aconitum* spp. (Ranunculaceae). Amorph. [α]_D +15.4 (c, 0.48 in EtOH). Artifact.

▶ AR5576000


O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-*benzoyl*: **Pyroaconitine**
[561-05-7]
C₃₂H₄₃NO₉ 585.693
Alkaloid from the the crude drug bushi obt. from the roots of some *Aconitum* spp. (Ranunculaceae). Amorph. [α]_D +21.8 (c, 0.22 in EtOH). Artifact formed from Aconitine, A-104 present in the raw aconite roots during the processing procedure. Has not been isol. so far from raw roots of *A.* spp.

▶ AR5572000

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-*methoxybenzoyl*): **Pyrojesaconitine**
[16298-91-2]
C₃₃H₄₅NO₁₀ 615.719
Alkaloid from the the crude drug bushi obt. from the roots of some *Aconitum* spp. (Ranunculaceae). Amorph. [α]_D +20.5 (c, 0.43 in EtOH). Artifact.

▶ AR5584000

Mori, T. et al., *Heterocycles*, 1989, **29**, 873-885 (isol, uv, ir, pmr, cmr, ms)
Ji, H. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 619-624 (16-Epipyroaconitine)

1,6,7,8,14,16,18-Heptahydroxy-4-methylaconitan-19-one H-123C₁₉H₂₇NO₈ 397.424**(1 α ,5 β ,6 β ,14 α ,16 β)-form**

O¹,O⁶,O¹⁴,O¹⁶-*Tetra-Me*, 18-O-(2-*aminobenzoyl*): **Pacifinine**. N-*Desethylpacifiline*. N-*Desethyl*-19-*oxoanthranoyllycoctonine*
[142768-35-2]
C₃₀H₄₀N₂O₉ 572.654
Alkaloid from seeds of *Delphinium elatum* cv. "Pacifig Giant" (Ranunculaceae). Amorph. solid. [α]_D²⁰ +48.5 (c, 0.8 in CHCl₃).

O¹,O⁶,O¹⁴,O¹⁶-*Tetra-Me*, N-*Et*, 18-O-(2-*aminobenzoyl*): **Pacifiline**. 19-*Oxoanthranoyllycoctonine*
[138964-90-6]
C₃₂H₄₄N₂O₉ 600.708
Alkaloid from the stems and leaves of *Delphinium ajacis* and seeds of *Del-*


phinium elatum (Ranunculaceae).

Amorph. [α]_D +58.1 (c, 0.08 in CHCl₃).
O¹,O⁶,O¹⁴,O¹⁶,O¹⁸-*Penta-Me*, N-*Et*: 19-**Oxodelphatine**
[25488-63-5]
C₂₆H₄₁NO₈ 495.612

Alkaloid from the stems and leaves of *Delphinium ajacis* (Ranunculaceae). [α]_D +32.8 (c, 0.31 in CHCl₃).

1-*Deoxy*, 1,2-*didehydro*, O⁸,O¹⁴,O¹⁶-*tri-Me*, N-*Et*, 1 α ,2 α -*epoxide*: **Budelphine**
C₂₄H₃₅NO₈ 465.542
Alkaloid from *Delphinium buschianum*. [α]_D²⁰ +6.9 (c, 0.18 in CHCl₃). Unusual Δ^1 -unsaturated equivalent functionality.

Liang, X. et al., *J. Nat. Prod.*, 1991, **54**, 1283-1287 (19-Oxodelphatine, 19-Oxoanthranoyllycoctonine)
Wada, K. et al., *Phytochemistry*, 1992, **31**, 2135-2138 (Pacifiline, Pacifinine)
Bitis, L. et al., *Helv. Chim. Acta*, 2007, **90**, 2217-2221 (*Budelphine*)


1,6,8,13,14,16,18-Heptahydroxy-4-methylaconitan-19-one H-124C₁₉H₂₇NO₈ 397.424**(1 α ,5 β ,6 α ,14 α ,16 β)-form**

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-*methoxybenzoyl*), 8-*Ac*: **Habaenine A**
[248263-85-6]
C₃₅H₄₇NO₁₁ 657.756
Alkaloid from *Aconitum habaense*. Amorph. solid. [α]_D¹⁵ -44.4 (c, 0.04 in CHCl₃). λ_{\max} 259 (log ϵ 4.47); 313 (log ϵ 3.71); 397 (log ϵ 2.97) (CHCl₃).

Yang, S. et al., *Helv. Chim. Acta*, 2007, **90**, 1160-1164 (isol, pmr, cmr)

Heptaphylline H-125

2-*Hydroxy*-1-(3-*methyl-2-butenyl*)-9H-*carbazole*-3-*carboxaldehyde*, 9CI. 3-*Formyl*-2-*hydroxy*-1-*prenylcarbazole*
[17750-35-5]



C₁₈H₁₇NO₂ 279.338
Alkaloid from the roots of *Clausena heptaphylla* and *Clausena pentaphylla*, the root bark of *Clausena excavata*, and the

leaves of *Clausena lansium* (Rutaceae). Shows antiplasmodial activity. Bright yellow needles (Et₂O or CHCl₃/hexane). Mp 171-172°. λ_{\max} 234 (log ϵ 4.42); 278 (log ϵ 4.53); 298 (log ϵ 4.58); 346 (log ϵ 4.09) (no solvent reported).

2,4-Dinitrophenylhydrazone:

Dark red cryst. (DMF). Mp 315-316°.

Me ether: 3-*Formyl*-2-*methoxy*-1-*prenylcarbazole*. **O-Methylheptaphylline**
[39027-80-0]

Alkaloid from *Clausena suffruticosa*. Mp 139-140°.

7-*Hydroxy*: 2,7-*Dihydroxy*-1-(3-*methyl-2-butenyl*)-9H-*carbazole*-3-*carboxaldehyde*, 9CI. 3-*Formyl*-2,7-*dihydroxy*-1-*prenylcarbazole*. **7-Hydroxyheptaphylline**
[170663-15-7]
C₁₈H₁₇NO₃ 295.337

Alkaloid from root bark of *Clausena lansium* (wampee) (Rutaceae). Yellow needles (CH₂Cl₂/MeOH). Mp 194-196°. λ_{\max} 238 (log ϵ 4.43); 302 (log ϵ 4.71); 340 (log ϵ 4.05) (MeOH).

8-*Hydroxy*: 2,8-*Dihydroxy*-1-(3-*methyl-2-butenyl*)-9H-*carbazole*-3-*carboxaldehyde*, 9CI. 3-*Formyl*-2,8-*dihydroxy*-1-*prenylcarbazole*. **Heptazoline**
[32042-35-6]
C₁₈H₁₇NO₃ 295.337

Alkaloid from *Clausena heptaphylla* and *Micromelum minutum*. Mp 212-214°. λ_{\max} 240 (log ϵ 4.58); 275 (log ϵ 4.62); 298 (log ϵ 4.4) (EtOH).

2',3',8'-*Trihydroxy*, 2',3'-*dihydro*: **Clauszoline D**
[185508-04-7]

C₁₈H₁₉NO₅ 329.352
Alkaloid from stem bark of *Clausena excavata*. Yellow oil. [α]_D²⁵ 0 (c, 0.1 in CHCl₃). λ_{\max} 206; 217; 242; 276; 292; 301 (sh); 356 (MeOH).

2', ξ ,3',7'-*Trihydroxy*, 2',3'-*dihydro*: 1-(2,3-*Dihydroxy*-3-*methylbutyl*)-2,7-*dihydroxy*-9H-*carbazole*-3-*carboxaldehyde*.

Clausine U
C₁₈H₁₉NO₅ 329.352
Alkaloid from *Clausena excavata*. Yellowish powder (Me₂CO). Mp 255-257°. [α]_D -72.8 (c, 0.01 in MeOH). λ_{\max} 201; 221; 240 (sh); 245; 254; 288 (sh); 303; 324 (sh); 339 (MeOH).

6-*Methoxy*: 2-*Hydroxy*-6-*methoxy*-1-(3-*methyl-2-butenyl*)-9H-*carbazole*-3-*carboxaldehyde*, 9CI. 3-*Formyl*-2-*hydroxy*-6-*methoxy*-1-*prenylcarbazole*. **6-Methoxyheptaphylline**
[41679-22-5]
C₁₉H₁₉NO₃ 309.364

Alkaloid from the roots of *Clausena indica* (Rutaceae). Bright yellow cryst. Mp 173-174°. λ_{\max} 207 (log ϵ 4.5); 235 (log ϵ 4.35); 252 (sh) (log ϵ 4.2); 272 (sh) (log ϵ 4.32); 283 (log ϵ 4.47); 311 (log ϵ 4.57); 340 (log ϵ 3.95) (EtOH).

7-*Methoxy*: 2-*Hydroxy*-7-*methoxy*-1-(3-*methyl-2-butenyl*)-9H-*carbazole*-3-*carboxaldehyde*, 9CI. 3-*Formyl*-2-*hydroxy*-7-*methoxy*-1-*prenylcarbazole*. **7-Methoxyheptaphylline**
[119736-84-4]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the root bark of *Clausena harmandiana* (Rutaceae). Cryst. (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 164–166°. λ_{max} 222 (ε 18800); 238 (ε 20000); 244 (ε 20000); 254 (ε 19300); 302 (ε 34700); 340 (ε 7400) (MeOH) (Berdy).

*A*³-Isomer, 2'-ε-Hydroxy: 2-Hydroxy-1-(2-hydroxy-3-methyl-3-butenyl)-9H-carbazole-3-carboxaldehyde. **Clausine S** [250259-38-2]

C₁₈H₁₇NO₃ 295.337

Alkaloid from *Clausena excavata*. Yellowish oil. [α]_D +159.1 (c, 0.002 in MeOH). λ_{max} 203; 236; 249; 279; 291 (sh); 299; 329 (sh); 343 (MeOH).

Chakraborty, D.P. et al., *J. Indian Chem. Soc.*, 1970, **47**, 1197-1198 (*Heptazoline*, uv, ir, pmr, struct)

Joshi, B.S. et al., *Indian J. Chem.*, 1972, **10**, 1123-1124 (6-Methoxyheptaphylline, isol, uv, ir, pmr, ms, struct)

Joshi, B.S. et al., *Phytochemistry*, 1972, **11**, 2065-2071 (isol, uv, ir, pmr, struct, synth)

Anwer, F. et al., *Experientia*, 1977, **33**, 412-413 (isol)

Sharma, R.B. et al., *Chem. Ind. (London)*, 1980, 158; 1982, 268 (synth, 6-Methoxyheptaphylline, *Heptazoline*)

Prakash, D. et al., *Indian J. Chem., Sect. B*, 1980, **19**, 1075-1076 (isol)

Wu, T.-S. et al., *J. Nat. Prod.*, 1982, **45**, 718-720 (isol)

Chaichantipyuth, C. et al., *J. Nat. Prod.*, 1988, **51**, 1285-1288 (7-Methoxyheptaphylline)

Kumar, V. et al., *Phytochemistry*, 1995, **40**, 1563-1565 (7-Hydroxyheptaphylline)

Ito, C. et al., *Chem. Pharm. Bull.*, 1996, **44**, 2231-2235 (*Clauszoline D*)

Wu, T.-S. et al., *Phytochemistry*, 1999, **52**, 523-527 (*Clausine S*, *Clausine U*)


Yenjai, C. et al., *Planta Med.*, 2000, **66**, 277-281 (activity)

Fun, H.-K. et al., *Acta Cryst. E*, 2007, **63**, o3964-o3965 (*Heptazoline*, cryst struct)

Begum, R. et al., *Nat. Prod. Commun.*, 2008, **3**, 815-818 (O-Methylheptaphylline)

2-(2-Heptenyl)-3-methyl-4(1H)-quinolinone H-126

2-(2-Heptenyl)-3-methyl-4-quinolinol. 2-(2-Heptenyl)-4-hydroxy-3-methylquinoline. **PSC-B**, **HMQ** [15436-59-6]

C₁₇H₂₁NO 255.359

λ_{max} 213; 244; 322; 335 (MeOH) (Berdy).

(E)-form [178956-00-8]

Alkaloid from *Pseudomonas cepacia* PC II. Shows antifungal and red pepper growth promoting activity. Cryst. (MeOH aq.). Mp 225-227°.

2',3'-Dihydro: 2-Heptyl-3-methyl-4(1H)-quinolinone. 2-Heptyl-3-methyl-4-qui-

nolinol. 2-Heptyl-4-hydroxy-3-methylquinoline. **PSC-C**

[172484-87-6]

[15436-60-9]

C₁₇H₂₃NO 257.375From *Pseudomonas cepacia* PC II.

Cryst. (MeOH aq.). Mp 227-228°. λ_{max} 215; 239; 322; 335 (MeOH) (Berdy).

Hashimoto, M. et al., *Chem. Pharm. Bull.*,


1967, **15**, 718 (occur, uv, pmr, struct)

Moon, S.-S. et al., *Phytochemistry*, 1996, **42**, 365 (isol, uv, ir, pmr, cmr, ms)

5-(4-Heptenyl)octahydro-8-methylindolizine, 9CI H-127

5-(4-Heptenyl)-8-methylindolizidine.

Dendrobates Alkaloid 235B. **Indolizidine 235B**

C₁₆H₂₉N 235.412

(+)-form

Alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates pumilio*. Trace const. in *Dendrobates histrionicus* (Dendrobatidae). [α]_D +11.3 (c, 1.0 in MeOH).

6',7'-Didehydro: 5-(4,6-Heptadienyl)octahydro-8-methylindolizine, 9CI. Dendrobates Alkaloid 233D. **Indolizidine 233D**

[141643-30-3]

Alkaloid from the skin of the Panamanian frog *Dendrobates pumilio*. [α]_D -3.4 (c, 0.16 in MeOH) (as hydrochloride).

6',7'-Didehydro, 4',5'-dihydro: 5-(6-Heptenyl)-8-methylindolizidine. Dendrobates Alkaloid 235B'. **Indolizidine 235B'**

C₁₆H₂₉N 235.412Alkaloid from *Dendrobates pumilio*.

6'-ε-Hydroxy: 7-(Octahydro-8-methyl-5-indolizinyloxy)-3-hepten-2-ol, 9CI. Dendrobates Alkaloid 251B. **Indolizidine 251B**

[141643-31-4]

C₁₆H₂₉NO 251.411Alkaloid from the skin of the Panamanian frog *Dendrobates pumilio*. [α]_D

+25.9 (c, 0.8 in CHCl₃).

(-)-form

4',5'-Dihydro: 5-Heptyl-8-methylindolizidine. Dendrobates Alkaloid 237D. **Indolizidine 237D**

C₁₆H₃₁N 237.428

Alkaloid from *Dendrobates pumilio* and *Dendrobates speciosus*.

Daly, J.W. et al., *Toxicol.*, 1978, **16**, 163-180 (isol, ms, rev)

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Tokuyama, T. et al., *Tetrahedron*, 1987, **43**, 643; 1991, **47**, 5401 (isol, pmr, cmr, ms, struct, derivs)


Collins, I. et al., *J.C.S. Perkin I*, 1991, 175-182 (synth)

Comins, D.L. et al., *J.O.C.*, 1997, **62**, 8182-8187 (synth)

Toyooka, N. et al., *Chem. Pharm. Bull.*, 2005, **53**, 555-560 (237D, synth, abs config)

Daly, J.W. et al., *J. Nat. Prod.*, 2005, **68**, 1556-1575 (rev)

2-(6-Heptenyl)-6-(4-pentenyl)piperidine H-128

C₁₇H₃₁N 249.439

(2RS,6RS)-form

(±)-trans-form

[127629-08-7]

Alkaloid from the venom of the ant *Monomorium delagoense*. Possesses insecticidal and repellent props.

(2RS,6SR)-form


(±)-cis-form

[127629-06-5]

Alkaloid from *Monomorium delagoense*. Possesses insecticidal and repellent props.

Jones, T.H. et al., *J. Nat. Prod.*, 1990, **53**, 429 (isol, struct, synth)

2-(5-Heptenyl)-6,7,8,8a-tetrahydro-3-methyl-4H-pyrrolo[2,1-b][1,3]oxazin-4-one H-129

C₁₅H₂₃NO₂ 249.352

Isol. from *Penicillium brevicompactum*. Insecticidal agent.

Cantin, A. et al., *Eur. J. Org. Chem.*, 1999, 221-226 (isol, ir, pmr, cmr, ms)

Castillo, M.-A. et al., *J. Agric. Food Chem.*, 1999, **47**, 2120-2124 (props)

1-Heptylamine, 8CI H-130

1-Heptanamine, 9CI. 1-Aminoheptane

[111-68-2]

H₃C(CH₂)₅CH₂NH₂C₇H₁₇N 115.218

Present in teak (*Tectona grandis*) wood. Fp -23. Bp 155.25°. n_D²⁰ 1.4268. pK_a 10.66 (25°).

▶ Flammable, fl. p. 35°. MK0600000

Hydrochloride: [142-93-8]Cryst. (EtOH/Et₂O). Mp 246°.**Picrate**:

Yellow needles. Mp 120-122°.

N-Ac: [14202-55-2]

C₉H₁₉NO 157.255Bp₁ 123° Bp₁₃ 155°.

N-Me: [36343-05-2]

C₈H₁₉N 129.245

Bp 168°.

N-Me, picrate: Mp 97°.

N,N-Di-Me: [5277-11-2]

C₉H₂₁N 143.272

Bp 172°.

N,N-Di-Me, N-oxide: [15290-93-4]

C₉H₂₁NO 159.271

Cryst. Mp 130-134°.

N-Et: [66793-76-8]

C₉H₂₁N 143.272Bp 181° Bp₁₆ 81-83°.

N,N-Di-Et: [26981-81-7]

C₁₁H₂₅N 171.325Bp 198° Bp_{10.5} 82-83°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 451C (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 281D (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 365C (ir)

Gohlke, R.S. et al., Anal. Chem., 1962, 34, 1281 (ms)

Hullot, P. et al., Bull. Soc. Chim. Fr., 1973, 2989 (synth)

Eggert, H. et al., J.A.C.S., 1973, 95, 3710 (cmr)

Kliger, G.A. et al., Neftekimiya, 1977, 17, 602 (synth)

Sax, N.I. et al., Dangerous Properties of Industrial Materials, 5th edn., Van Nostrand Reinhold, 1979, 715

2-Heptyl-1-cyclopropanepropanoic acid H-131

4,5-Methylenedodecanoic acid



(1R,2R)-form

C₁₃H₂₄O₂ 212.331**(1R,2R)-form**

2-Phenylethylamide: Grenadamide

[205521-84-2]

C₂₁H₃₃NO 315.498Isol. from *Lyngbya majuscula*. Toxic to brine shrimp, cannabinoid receptor antagonist. [α]_D -11 (c, 0.1 in CHCl₃). λ_{max} 206 (ε 2600) (MeOH).**(1R,2R)-form**

(5-Acetoxy-4-bromo-1,3-pentadienyl) ester (Z,Z-): Grenadiene

[205521-82-0]

C₂₀H₃₁BrO₄ 415.367Isol. from *Lyngbya majuscula*. Cytotoxic agent. [α]_D -8 (c, 0.1 in CHCl₃). λ_{max} 252 (ε 12200) (MeOH).

5-Acetoxy-1,3-pentadienyl ester (1Z,3E-): Debromogrenadiene

[205521-83-1]

C₂₀H₃₂O₄ 336.47Isol. from *Lyngbya majuscula*. Toxic to brine shrimp. [α]_D +5 (c, 0.1 in CHCl₃). λ_{max} 232 (ε 8000) (hexane).

Sitachitta, N. et al., J. Nat. Prod., 1998, 61, 681-684 (isol, uv, ir, pmr, cmr, ms)

Al Dulayymi, J.R. et al., Tetrahedron, 2004, 60, 341-345 (Grenadamide, synth, abs config)

Green, R. et al., Tet. Lett., 2005, 46, 7931-7934 (Grenadamide, synth)

Avery, T.D. et al., Org. Biomol. Chem., 2006, 4, 323-330 (Grenadamide, synth)


Salim, H. et al., Tet. Lett., 2007, 48, 2059-2062 (Grenadamide, synth)

1-Heptyl-3-hydroxy-2-methyl-9H-carbazole H-132

1-Heptyl-2-methyl-9H-carbazol-3-ol, 9CI.

Carazostatin

[126168-32-9]

C₂₀H₂₅NO 295.424Prod. by *Streptomyces chromofuscus*.

Lipid peroxidation inhibitor, free radical scavenger. Pale yellow powder. Sol.

Me₂CO, CHCl₃, EtOAc; poorly sol. H₂O.Mp 149-152° dec. λ_{max} 218 (ε 153000);

235 (ε 141000); 254 (ε 76000); 266 (ε

60000); 303 (ε 83000); 342 (ε 20000)

(MeOH) (Derep). λ_{max} 218 (ε 153000);

235 (ε 141000); 254 (ε 76000) (MeOH)

(Berdy).

▶ FE6366000

Kato, S. et al., J. Antibiot., 1989, 42, 1879 (isol, struct)

Jackson, P.M. et al., J.C.S. Perkin 1, 1991, 2941 (synth)

Shin, K. et al., Chem. Lett., 1995, 289 (synth)

Choshi, T. et al., J.O.C., 1997, 62, 2535 (synth)

Nonaka, Y. et al., Heterocycles, 2000, 53, 1681-1684 (synth)

Knöflker, H.-J. et al., Tetrahedron, 2002, 58, 8937-8945 (synth, uv, ir, pmr, cmr)


2-Heptyl-4-hydroxyquinoline H-133

2-Heptyl-4(1H)-quinolinone, 9CI. 2-

Heptyl-4-quinolinol. Pyo Ib. Pseudane

VII. MY 12-62a. Antibiotic MY 12-62a

[40522-46-1]

C₁₆H₂₁NO 243.348Alkaloid from a yellow marine pseudomonad. The terrestrial bacterium *Pseudomonas aeruginosa* also yields this compd. Shows antibiotic props. against a number of bacteria. 5-Lipoxygenase inhibitor and Streptomycin antagonist. Cryst. (Me₂CO aq.). Mp 146-147°. Bp_{0.2} 118-122°.

N-Oxide: 2-Heptyl-4-hydroxyquinoline N-oxide. Pyo II. KF 8940. Antibiotic KF 8940

[341-88-8]

[1401-06-5]

C₁₆H₂₁NO₂ 259.347Metab. of *Pseudomonas pyocyanea*, *Pseudomonas aeruginosa*, *Pseudomonas methanica* and *Pseudomonas* sp. KUH-

001. Potent 5-lipoxygenase inhibitor.

Active against *Staphylococcus aureus*.

Synergistic with Vancomycin and

Erythromycin. Cryst. (EtOH). Mp

158-160°.

▶ LD₅₀ (mus, ipr) 40 mg/kg. VC58900001',2'-Didehydro(E-): 2-(1-Heptyl)-4(1H)-quinolinone, 9CI. 2-(1-Heptyl)-4-hydroxyquinoline. A¹-Pseudene VII

[60783-02-0]

C₁₆H₁₉NO 241.332Prod. by *Pseudomonas aeruginosa*.

Antifungal agent.

1',2'-Didehydro(Z-):

C₁₆H₁₉NO 241.332Metab. of *Pseudomonas aeruginosa*.

No CAS no. found to 2007.

4',5'-Didehydro: 2-(4-Heptyl)-4(1H)-quinolinone. Acutine

[36150-05-7]

C₁₆H₁₉NO 241.332

Alkaloid from the above-ground parts

of *Haplophyllum acutifolium* (Ruta-ceae). Prisms (Me₂CO). Mp 122-123°.

Becomes pink on standing in the

light.

NH-form

N-Me: 2-Heptyl-1-methyl-4(1H)-quinolinone. Schinifoline†

[80554-58-1]

C₁₇H₂₃NO 257.375Alkaloid from peel of *Zanthoxylum**schinifolium* (Rutaceae). Cryst. Mp 81-

82°.

8-Methoxy, N-Me: 2-Heptyl-8-methoxy-1-methyl-4(1H)-quinolinone

C₁₈H₂₅NO₂ 287.401Alkaloid from trunk bark of *Esen-**beckia almawillia* (Rutaceae). Oil.**OH-form** [159979-57-4]

Me ether: 2-Heptyl-4-methoxyquinoline, 9CI

[80554-59-2]

C₁₇H₂₃NO 257.375Alkaloid from *Zanthoxylum avicennae*.

Mp 35-38°.

[2503-80-2]

Hays, E.E. et al., J. Biol. Chem., 1945, 159, 725-750 (isol)

Wells, I.C. et al., J. Biol. Chem., 1952, 196, 331-340; 341-345 (synth, uv, struct)

Cornforth, J.W. et al., Biochem. J., 1956, 63,

124-130; 130-137 (isol, props)

Ames, D.E. et al., J.C.S., 1956, 3079-3083

(synth)

Luckner, M. et al., Tet. Lett., 1965, 6, 741-744

(biosynth)

Ritter, C. et al., Eur. J. Biochem., 1971, 18, 391-

400 (biosynth)

Razzakova, D.M. et al., Khim. Prir. Soedin., 1973, 9, 206-210; Chem. Nat. Compd. (Engl. Transl.), 1973, 9, 199-202 (Acutine)

Kozlovskii, A.G. et al., Izv. Akad. Nauk SSSR, Ser. Khim., 1976, 1146-1150 (A¹-Pseudene VII)

Wratten, S.J. et al., Antimicrob. Agents

Chemother., 1977, 11, 411-414 (isol)

Budzikiewicz, H. et al., Monatsh. Chem., 1979, 110, 947-953 (isol, uv)

Somanathan, R. et al., J. Het. Chem., 1981, 18, 1077-1079 (synth, uv, ir, pmr)

Kitamura, S. et al., J. Antibiot., 1986, 39, 1160-1166 (KF8940, isol, struct, props)


Liu, S.L. et al., Yaoxue Xuebao, 1991, 26, 836-840; CA, 117, 86638h (Schinifoline)

Wu, W. et al., Zhongcaoyao, 1992, 23, 115-116;

CA, 117, 76305h (isol, Me ether)


- Guilhon, G.M.S.P. *et al.*, *Phytochemistry*, 1994, **37**, 1193-1195 (2-Heptyl-8-methoxy-1-methyl-4(1H)-quinolinone)
- Debitus, C. *et al.*, *J. Mar. Biotechnol.*, 1998, **6**, 136-141 (*isol*)
- Hwang, S.-Y. *et al.*, *J. Microb. Biotechnol.*, 1998, **8**, 111-118 (*N-oxide, isol, activity*)
- Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur*)
- Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*biosynth*)
- Bredenbruch, F. *et al.*, *J. Bacteriol.*, 2005, **187**, 3630-3635 (*biosynth*)

3-Heptyl-3-hydroxy-2,4(1H,3H)-quinolinedione **H-134**
 MY 12-62c. Antibiotic MY 12-62c [69808-30-6]



- $C_{16}H_{21}NO_3$ 275.347
 Prod. by *Pseudomonas methanica* and *Pseudomonas aeruginosa*. Also prod. by a bacterium *isol.* from the sponge *Suberea creba*. 5-Lipoxygenase inhibitor. Cryst. (EtOAc). Mp 142°.
- Neuenhaus, N. *et al.*, *Z. Naturforsch., B*, 1979, **34**, 313 (*isol, struct, synth*)
- Kitamura, S. *et al.*, *J. Antibiot.*, 1986, **39**, 1160 (*isol, struct, props*)
- Debitus, C. *et al.*, *J. Mar. Biotechnol.*, 1998, **6**, 136-141 (*isol*)

3-Heptyl-5-methylpyrrolizidine **H-135**
 Xenovenine. Pyrrolizidine 223H




$C_{15}H_{29}N$ 223.401

- (3S,5R,8S)-form** [135683-51-1]
 Alkaloid from the venom of the cryptic thief ant *Solenopsis xenoveneum*. $[\alpha]_D^{24}$ +11.7 (c, 0.7 in $CHCl_3$).
- 6'- ζ -Hydroxy-3-(6-Hydroxyheptyl)-5-methylpyrrolizidine. **Pyrrolizidine 239K'** [151805-19-5]
 $C_{15}H_{29}NO$ 239.4
 Alkaloid from the frog *Mantella* sp.
- [74986-28-0 , 105616-96-4 , 88931-18-4 , 161106-53-2]
- Jones, T.H. *et al.*, *J.O.C.*, 1980, **45**, 4778-4780 (*isol, ir, pmr, cmr, ms, struct, synth*)
- Provot, O. *et al.*, *J.O.C.*, 1991, **57**, 2163-2166 (*synth, pmr, abs config*)
- Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*)
- Dhimane, H. *et al.*, *Eur. J. Org. Chem.*, 1998, 1955-1963 (*synth, pmr, cmr*)
- Arredondo, V.M. *et al.*, *J.A.C.S.*, 1999, **121**, 3633-3639 (*synth*)


- Takahata, H. *et al.*, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1293-1295 (*synth*)

2-Heptylpyrrole, 9CI **H-136**
 [878-12-6]



- $C_{11}H_{19}N$ 165.278
 Bp₄₀ 160°.
- N-Sulfo-2-Heptylpyrrole sulfamate** [587875-52-3]
 $C_{11}H_{19}NO_3S$ 245.342
 Alkaloid from the marine annelid *Cirriformia tentaculata*. Glass (as Na salt).
- Garrido, D.O.A. *et al.*, *J.O.C.*, 1984, **49**, 2619-2622 (*synth*)
- Barsby, T. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1110-1112 (*sulfamate*)

Herbaine **H-137**
 Vincaherbine [5308-82-7]




Absolute Configuration

- $C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from *Vinca herbacea* (Apocynaceae). Mp 126-128° dec. $[\alpha]_D^{25}$ -217. λ_{max} 228 (log ϵ 4.63); 274 (log ϵ 3.85); 297 (log ϵ 3.87) (EtOH).

- 10-Methoxy- Herbaceine. Vincaherbinine** [5308-79-2]
 $C_{23}H_{30}N_2O_5$ 414.5
 Alkaloid from *Vinca herbacea* (Apocynaceae). Cryst. (EtOH). Mp 144° dec. $[\alpha]_D^{25}$ -219 (c, 0.544 in Py). λ_{max} 226 (log ϵ 4.53); 280 (sh) (log ϵ 3.83); 300 (log ϵ 4.04) (EtOH).


- 10-Methoxy; methiodide:**
 Orange cryst. (Me₂CO/MeOH). Mp 219-223°.
- Ognyanov, I. *et al.*, *Chem. Ber.*, 1966, **99**, 1008-1014 (*Herbaine, Herbaceine, isol, uv, ir, pmr, ms, struct*)
- Ognyanov, I. *et al.*, *Chem. Comm.*, 1967, 579-581 (*stereochem*)
- Ognyanov, I. *et al.*, *CA*, 1969, **71**, 760p (*isol, struct*)

Herbamide A **H-138**
 [161503-26-0]



- $C_{16}H_{21}Cl_3N_2OS$ 395.779
 Isol. from the marine sponge *Dysidea herbacea*. Oil. $[\alpha]_D^{25}$ +13 (c, 0.013 in $CHCl_3$).
- Clark, W.D. *et al.*, *Tet. Lett.*, 1995, **36**, 1185 (*isol, uv, ir, pmr, cmr, struct*)

Herbamine **H-139**
 Methyl 19,20-didehydro-3,17-dihydroxy-yajmalan-16-carboxylate, 9CI. 2-Epi-3-hydroxyvincamajine [38485-13-1]




Absolute Configuration

- $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from the aerial parts of *Vinca libanotica* (Apocynaceae). Mp 153-155°. $[\alpha]_D^{25}$ -15 (c, 0.66 in $CHCl_3$).

- N-De-Me- Herbaine. 2-Epi-3-hydroxy-quebrachidine** [38485-12-0]
 $C_{21}H_{24}N_2O_4$ 368.432
 Alkaloid from *Vinca herbacea* and *Vinca libanotica* (Apocynaceae). Mp 202-205° dec.

- Vachnadze, V.Yu. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 341; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 334 (*isol, uv, ir, pmr, ms*)
- Aynilian, G.H. *et al.*, *J. Nat. Prod.*, 1974, **37**, 299 (*isol, ir, uv, ms*)
- Aynilian, G.H. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 341 (*pmr, struct*)

Herbindole A **H-140**
 1,6,7,8-Tetrahydro-4,5,6,8-tetramethylcyclopent[g]indole, 9CI [128397-78-4]



R = CH₃

- $C_{15}H_{19}N$ 213.322
 Alkaloid from the sponge *Axinella* sp. Cytotoxic and fish antifeedant. Needles (MeOH). Mp 120-122°. λ_{max} 222 (ϵ 46800); 272 (ϵ 12300) (base in MeOH) (Derep). λ_{max} 210 (ϵ 55000); 226 (sh) (ϵ 47900); 272 (ϵ 8510) (acidic MeOH) (Derep).

- Herb, R. *et al.*, *Tetrahedron*, 1990, **46**, 3089 (*isol, pmr, cmr, struct*)
- Muratake, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 854 (*synth, abs config*)
- Jackson, S.K. *et al.*, *J.O.C.*, 2007, **72**, 1405-1411 (*synth*)

Herbindole B **H-141**

4-Ethyl-1,6,7,8-tetrahydro-5,6,8-trimethylcyclopent[*g*]indole, 9CI
[128397-79-5]

As Herbindole A, H-140 with
R = -CH₂CH₃

C₁₆H₂₁N 227.349

Alkaloid from the sponge *Axinella* sp. Cytotoxic and fish antifeedant. Needles (MeOH). Mp 118-120°. λ_{max} 222 (ε 46800); 272 (ε 12300) (base in MeOH) (Derep). λ_{max} 210 (ε 55000); 226 (sh) (ε 47900); 272 (ε 8510) (acidic MeOH) (Derep). λ_{max} 222 (ε 46773); 272 (ε 12300) (MeOH) (Berdy).

Herb, R. et al., *Tetrahedron*, 1990, **46**, 3089 (isol, pmr, cmr, struct)

Muratake, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 854 (synth, abs config)

Jackson, S.K. et al., *J.O.C.*, 2007, **72**, 1405-1411 (synth)

Herbindole C **H-142**

4-(1-Butenyl)-1,6,7,8-tetrahydro-5,6,8-trimethylcyclopent[*g*]indole, 9CI
[128397-80-8]

As Herbindole A, H-140 with
R = -CH=CHCH₂CH₃(*E*-)

C₁₈H₂₃N 253.386


Alkaloid from the sponge *Axinella* sp. Cytotoxic and fish antifeedant. Oil. Regioisomer of Trikentrin B, T-595. λ_{max} 212 (ε 50100); 226 (ε 42700); 294 (ε 10000) (MeOH) (Derep).

Herb, R. et al., *Tetrahedron*, 1990, **46**, 3089 (isol, pmr, cmr, struct)

Muratake, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 854 (synth, abs config)

Herbipoline **H-143**

2-Amino-6-hydroxy-7,9-dimethylpurinium, 9CI. 7,9-Dimethylguaninium
[524-35-6]



C₇H₁₀N₅O[⊕] 180.189

The +ve charge is delocalised over N-7 and N-9. Isol. as the zwitterionic enolate. Isol. from the giant silicious sponge *Geodia gigas*. Needles (EtOH aq.). Mp 312°.

Picrate: Mp 292-295°.

N²-Me: **Heterimine E**

[196090-68-3]

C₈H₁₂N₅O[⊕] 194.216

Alkaloid from *Heterostemma brownii*. Needles (MeOH) (as chloride). Mp 295° (dec.) (chloride). CAS no. refers to chloride. λ_{max} 236 (log ε 3.9); 253 (sh) (log ε 3.98); 291 (log ε 3.79) (MeOH).

3-Me: **1-Methylherbipoline**

[97174-15-7]

C₈H₁₂N₅O[⊕] 194.216

From the marine sponge *Jaspis* sp. Collagenase inhibitor. Sol. MeOH, Et₂O. Counterion in nat. prod. not characterised. λ_{max} 260 (ε 7600); 282 (ε 4460) (MeOH) (Berdy).

N²,N²-Di-Me: **Heterimine D**

C₉H₁₄N₅O[⊕] 208.243

Alkaloid from the aerial parts of *Heterostemma brownii* (Asclepiadaceae). Needles (MeOH) (as chloride). Mp 196-198° (chloride). CAS no. refers to chloride. λ_{max} 240 (log ε 4); 258 (sh) (log ε 3.85); 300 (log ε 3.69) (MeOH).

Me ether: 2-Amino-6-methoxy-7,9-dimethylpurinium(1+). **Heterimine C**
[176900-99-5]

C₈H₁₂N₅O[⊕] 194.216

Alkaloid from aerial parts of *Heterostemma brownii*. Needles (MeOH) (as chloride). Mp 268-270° (chloride). CAS no. refers to chloride.

Me ether, N²-Me: **Heterimine B**

[176900-98-4]

C₉H₁₄N₅O[⊕] 208.242

From aerial parts of *Heterostemma brownii*. Shows inhibitory activity on K562 and HL-60 cell lines. Needles (MeOH) (as chloride). Mp 229-231° (chloride). CAS no. refers to chloride.

Me ether, N²,N²-di-Me: **Heterimine A**
[176900-97-3]

C₁₀H₁₆N₅O[⊕] 222.269

From aerial parts of *Heterostemma brownii*. Shows inhibitory activity against K562 and HL-60 cell lines. Needles (MeOH) (as chloride). Mp 225-227° (chloride). CAS no. refers to chloride.

Ackermann, D. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1957, **308**, 270; **309**, 286 (Herbipoline, isol, ir)

Bredereck, H. et al., *Chem. Ber.*, 1960, **93**, 1206 (Herbipoline, synth, struct)

Pfleiderer, W. et al., *Annalen*, 1961, **647**, 167-173 (synth)

Yagi, H. et al., *J. Nat. Prod.*, 1994, **57**, 837-838 (1-Methylherbipoline)

Metzger, S. et al., *Chem. Eur. J.*, 1996, **35**, 1228 (Herbipoline, struct, props)

Lin, Y.-L. et al., *Heterocycles*, 1996, **43**, 781 (Heterimines)


Lin, Y.-L. et al., *J. Nat. Prod.*, 1997, **60**, 982-985 (Heterimines D-E)

Peinador, C. et al., *Tetrahedron*, 1997, **53**, 8269 (Heterimine A, synth)

Jakobsen, E. et al., *Heterocycles*, 2000, **53**, 935-940 (Heterimine C, synth)

Herclavine **H-144**

N-[2-(4-Methoxyphenyl)ethyl]-N-methyl-3-phenyl-2-propenamido, 9CI. N-(p-Methoxyphenethyl)-N-methylcinnamamide. N-(2-p-Anisylethyl)-N-methylcinnamamide. Cinnamic acid p-methoxyphenethylamide



C₁₉H₂₁NO₂ 295.38

(*E*)-form [539-18-4]

Alkaloid from the bark of *Zanthoxylum clava-herculis* (Hercules' club) (Rutaceae). Cryst. (Et₂O or petrol). Mp 76°.


LaForge, F.B. et al., *J.O.C.*, 1944, **9**, 250 (isol, struct, synth)

Crombie, L. et al., *J.C.S.*, 1955, 995 (isol, uv, ir)

Chatterjee, A. et al., *J.O.C.*, 1959, **24**, 687 (ir)

Hericerin **H-145**

6-(3,7-Dimethyl-2,6-octadienyl)-2,3-dihydro-7-hydroxy-5-methoxy-2-(2-phenylethyl)-1*H*-isoindol-1-one, 9CI
[140381-53-9]



C₂₇H₃₃NO₃ 419.563

Isol. from the edible lions mane mushroom (*Heridium erinaceum*). Pollen growth inhibitor. Needles + 1/2H₂O (C₆H₆/EtOAc). Mp 138-140°.

5'-Oxo: **Hericenone B**

[126654-53-3]

C₂₇H₃₁NO₄ 433.546


Isol. from the edible lions mane mushroom (*Heridium erinaceum*). Cytotoxic. Cryst. (CHCl₃). Mp 136-138°. λ_{max} 214 (ε 45000); 235 (sh) (ε 16800); 248 (ε 17800) (MeOH) (Derep).

Kawagishi, H. et al., *Tet. Lett.*, 1990, **31**, 373 (Hericenone B)

Kimura, Y. et al., *Agric. Biol. Chem.*, 1991, **55**, 2673 (Hericerin)

Hermidin **H-146**

3,6-Dihydroxy-4-methoxy-1-methyl-2(1*H*)-pyridinone. 3-Hydroxy-4-methoxy-1-methyl-2,6(1*H*,3*H*)-pyridinedione
[92446-30-5]



C₇H₉NO₄ 171.152

Monooxo-form predominates in aq. soln. Chromogenic substance from aqueous extract of *Mercurialis perennis*. Cryst. Bp_{0.01} 110° subl. Readily oxid. by air to give a radical-anion Cyanohermidin followed by a mixt. of dimers Chrysohermidin.

► Powerful sternutator.

Di-Ac:

Needles (CH₂Cl₂). Mp 150-151°.


Forrester, A.R. et al., *Experientia*, 1984, **40**, 688-689 (epr)

Swan, G.A. et al., *J.C.S. Perkin 1*, 1985, 1757-1766 (isol, uv, ir, pmr, struct, synth)

Ostrozhenkova, E. et al., *Phytochemistry*, 2007, **68**, 2816-2824 (biosynth)

Hernandaline

H-147

C₂₉H₃₁NO₇ 505.566**(S)-form** [10210-99-8]

Alkaloid from *Hernandia ovigera* and *Hernandia nymphaeifolia* (Hernandiaceae). Also obt. by KMnO₄ oxidation of Thallicarpine, T-312. Shows anti-platelet aggregation activity. Needles (EtOH). Mp 170-171.5°. [α]_D²⁵ +36.5 (c, 0.10 in CHCl₃). λ_{max} 216 (log ε 4.36); 278 (log ε 4.4); 304 (log ε 4.2) (EtOH).

2'-Alcohol: Hernandalinol

[62874-90-2]

C₂₉H₃₃NO₇ 507.582

Prod. from Thallicarpine, T-312 by *Streptomyces punipalms*. Mp 95-100°. [α]_D²⁶ +35 (c, 1.03 in EtOH). λ_{max} 283 (log ε 4.24); 303 (log ε 4.16) (EtOH).

6a,7-Didehydro: Dehydrohernandaline

[5263-33-2]

C₂₉H₂₉NO₇ 503.551

Alkaloid from trunk bark of *Hernandia sonora* (Hernandiaceae). Yellowish needles (CHCl₃/MeOH). Mp 151-153°.

3-Methoxy: Thaliadine

[67510-96-7]

C₃₀H₃₃NO₈ 535.593

Alkaloid from the roots of *Thalictrum minus* Race B (Ranunculaceae). Also obt. by brief oxidation of Adiantifoline. Shows hypotensive activity. Yellow rosettes (MeOH). Mp 143.5-144.5°. [α]_D²⁶ 0 (c, 0.22 in CHCl₃ or MeOH).

3-Methoxy, 6a,7-didehydro:

Synthetic. Yellow needles (Me₂CO/MeOH). Mp 147-148°.

Cava, M.P. et al., *Tet. Lett.*, 1966, 4279-4282 (isol, uv, ir, pmr, synth, struct)

Kupchan, S.M. et al., *Chem. Comm.*, 1971, 599-600 (synth)

Nabih, T. et al., *J. Med. Chem.*, 1977, **20**, 914-917 (Hernandalinol)

Liao, W.T. et al., *J. Nat. Prod.*, 1978, **41**, 271-276 (Thaliadine)


Chen, I.S. et al., *Phytochemistry*, 1995, **40**, 983-986 (Dehydrohernandaline)

Chen, J.J. et al., *Planta Med.*, 2000, **66**, 251-256 (activity)

Hernandezine

H-148

Thalicsimine. *Thaliximine*
[6681-13-6]

C₃₉H₄₄N₂O₇ 652.786

Alkaloid from *Thalictrum hernandezii*, *Thalictrum fendleri*, *Thalictrum podocarpum*, *Thalictrum rochebrunianum* and other *Thalictrum* spp. (Ranunculaceae). Inhibitor of conditioned avoidance reactions in exptl. animals. Shows strong antiinflammatory activity. Weak anti-neoplastic agent. Antiseptic. Cryst. (hexane). Mp 192-193° (122-124°, 158-159°). [α]_D²⁰ +250 (c, 0.2 in CHCl₃). Log P 7.83 (uncertain value) (calc). Mp dependent on solv. of crystallisation. λ_{max} 282 (ε 7250) (MeOH) (Berdy).

N²-Oxide: Hernandezine 2'-N-oxide

[78414-48-9]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from the aerial parts of *Thalictrum sultanabadense* (Ranunculaceae). Mp 179-180°.

O⁵-De-Me, N²-de-Me: N-Desmethylthaliadine

[65230-06-0]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from the roots of *Thalictrum podocarpum* (Ranunculaceae). Shows antimicrobial activity. Needles (MeOH). Mp 173-174°. [α]_D²⁵ +280 (c, 0.14 in MeOH).

N²-De-Me: Thalissamine. N'-Norhernandezine. 2'-Demethylhernandezine

[26326-54-5]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from *Thalictrum rochebrunianum* roots and the aerial parts of *Thalictrum simplex* (Ranunculaceae). Noncryst. [α]_D²² +143 (c, 0.28 in MeOH). [α]_D¹⁵ +241 (c, 2.6 in CHCl₃). Identity with N'-Norhernandezine established in 1984. Mollov et al quote a negative opt. rotn. but Thalissamine is dextrorotatory since it was related chemically to (+)-Hernandezine.

O⁵-De-Me: Thaliadine. 5-O-Demethylhernandezine

[18251-36-0]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from *Thalictrum fendleri* (whole plant), the roots of *Thalictrum rugosum*, *Thalictrum minus* and *Thalictrum podocarpum*, and from the aerial parts of *Thalictrum simplex* and *Thalictrum sultanabadense* (Ranunculaceae). Antiseptic. Shows antimicrobial activity. Cryst. (Me₂CO). Mp 158-159°. [α]_D²⁵ +235 (CHCl₃). Log P 7.54 (uncertain value) (calc).

1'-Epimer, O⁵-de-Me: Isothaliadine

[64924-28-3]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from the roots of *Thalictrum podocarpum* (Ranunculaceae). Mp 136-138°. [α]_D²⁵ -70 (c, 0.13 in MeOH).

Padilla, J. et al., *Tetrahedron*, 1962, **18**, 427 (isol, uv, ir, pmr, struct)

Battersby, A.R. et al., *J.C.S.*, 1965, 2239 (ord)

Mollov, N.M. et al., *Dokl. Bulg. Akad. Nauk.*, 1967, **20**, 329; 1970, **23**, 181 (*Thaliadine*, *Thalissamine*, isol)

Shamma, M. et al., *Tetrahedron*, 1967, **23**, 2887 (*Hernandezine*, *Thaliadine*, isol, uv, ms, pmr, struct)

Baldas, J. et al., *J.C.S. Perkin 1*, 1972, 592 (ms)

Cava, M.P. et al., *Tet. Lett.*, 1974, 4259 (cd)

Wu, W.-N. et al., *J. Nat. Prod.*, 1977, **40**, 384-394 (*N-Desmethylthaliadine*, *Isothaliadine*, activity)


Wu, J. et al., *J.O.C.*, 1980, **45**, 213 (*N'-Norhernandezine*)

Mukhamedova, S. et al., *Khim. Prir. Soedin.*, 1981, **17**, 250; *CA*, **95**, 58078j (*Hernandezine 2'-N-oxide*)

Guinaudeau, H. et al., *Tetrahedron*, 1984, **40**, 1975 (struct, *Thalissamine*)

Hernandine†

H-149

C₁₈H₁₉NO₅ 329.352


See also Hernandine in L-244 and Hernandine in P-209. Alkaloid from *Hernandia bivalvis* (Hernandiaceae).

Sam, S.K. et al., *Proc. Malays. Biochem. Soc. Conf.*, 1985, 11th; 49; *CA*, **107**, 214804s (struct)

Hernandonine

H-150

8H-Bis[1,3-benzodioxolo[6,5,4-de:4',5'-g]quinolin-8-one, 9CI. 1,2:10,11-Bis(-methylenedioxy)oxoaporphine
[28314-78-5]

C₁₈H₉NO₅ 319.273

Alkaloid from the trunk bark and root bark of *Hernandia ovigera*, and the bark of *Hernandia jamaicensis* and *Hernandia papuana* (Hernandiaceae). Yellow needles (C₆H₆/CHCl₃ or MeOH/CHCl₃). Mp 298-300° dec.

Oxime:

Orange needles (EtOH). Mp 264-265° dec.

3-Methoxy: Oxoduocine

[155944-22-2]

C₁₉H₁₁NO₆ 349.299

Trace alkaloid from roots of *Lindera myrrha* (Lauraceae). Red cryst. (MeOH). Mp 265°.

Ito, K. et al., *Tet. Lett.*, 1970, 3023 (uv, ir, pmr, struct, synth)

Lahey, F.N. et al., *Aust. J. Chem.*, 1971, **24**, 671 (isol, uv, ir, pmr)

Cava, M.P. et al., *Tetrahedron*, 1971, **27**, 2639 (isol, uv, ir, pmr)

Yang, T.H. et al., *J. Chin. Chem. Soc. (Taipei)*, 1976, **23**, 29; *CA*, **85**, 37125a (isol)

Ba Hung Phan, *et al.*, *Phytochemistry*, 1994, **35**, 1363 (*Oxoduocine*)

Herpestine**H-151**

$C_{34}H_{46}N_2O_6$ 578.747

Struct. unknown. Alkaloid from *Monnieria cuneifolia* (Rutaceae). Needles (C_6H_6). Mp 116-118°.


Sulfate: Mp 120° dec.

Tartrate: Mp 209-210°.

Basu, N.K. *et al.*, *Q. J. Pharm. Pharmacol.*, 1947, **20**, 137-139

Herquiline A**H-152**

[71812-08-3]



Relative configuration

$C_{19}H_{26}N_2O_2$ 314.427

Prod. by *Penicillium herquei*. Weakly active against platelet aggregation induced by ADP. Mycotoxin. Cryst. + $\frac{1}{2}H_2O$. Sol. MeOH, C_6H_6 ; fairly sol. hexane; poorly sol. H_2O . Mp 171-174° dec. $[\alpha]_D^{27}$ -388 (c, 1 in MeOH). λ_{max} 288 (MeOH) (Derep). λ_{max} 287 (sh) (ϵ 317) (MeOH/NaOH) (Derep). λ_{max} 288 (sh) (ϵ 283) (MeOH) (Derep). λ_{max} 291 (ϵ 331) (MeOH/HCl) (Berdy).

► LD₅₀ (mus, ipr) 100-300 mg/kg.

Omura, S. *et al.*, *J. Antibiot.*, 1979, **32**, 786

(*isol, uv, ir*)

Furusaki, A. *et al.*, *Chem. Comm.*, 1980, 698


(*cryst struct*)

Enomoto, Y. *et al.*, *J. Antibiot.*, 1996, **49**, 50

(*pmr, cmr*)

Herquiline B**H-153**

15-Methyl-15,17-diazatetracyclo[12.2.2.1^{3,7}.jeicosa-3(20),12(19)-diene-6,9-dione, 9CI
[174423-44-0]



$C_{19}H_{26}N_2O_2$ 314.427


Prod. by *Penicillium herquei*. Active against platelet aggregation induced by ADP. Mycotoxin. Powder. Sol. MeOH, CHCl₃; fairly sol. hexane; poorly sol. H_2O , C_6H_6 , EtOAc. Mp 109-113°. $[\alpha]_D^{25}$ -85.2 (c, 0.3 in MeOH). λ_{max} 203 (ϵ 9100); 222 (sh) (ϵ 1530); 285 (ϵ 780) (MeOH).

Enomoto, Y. *et al.*, *J. Antibiot.*, 1996, **49**, 50-53

(*isol, uv, ir, pmr, cmr, props*)

Herveline C

[160098-81-7]



$C_{40}H_{46}N_2O_7$ 666.813

Alkaloid from stem bark of *Hernandia voyronii* (Hernandiaceae). Cryst. (hexane). Mp 85-86°. $[\alpha]_D^{20}$ -24 (c, 0.3 in CH_2Cl_2).

O⁸-De-Me: **Herveline B**

[160098-80-6]

$C_{39}H_{44}N_2O_7$ 652.786

From stem bark of *Hernandia voyronii* (Hernandiaceae). Cryst. (hexane). Mp 102-104°. $[\alpha]_D^{20}$ +2.4 (c, 0.3 in CH_2Cl_2).

O¹²-De-Me: **Herveline A**

[160098-79-3]

$C_{39}H_{44}N_2O_7$ 652.786

From stem bark of *Hernandia voyronii* (Hernandiaceae). Cryst. (hexane). Mp 114-116°. $[\alpha]_D^{20}$ -2 (c, 0.02 in CH_2Cl_2).

8,12'-Di-O-de-Me: **Herveline D**

$C_{38}H_{42}N_2O_7$ 638.759

From the stem bark of *Hernandia voyronii*. Cryst. (hexane/EtOAc). Mp 120-122°. $[\alpha]_D^{20}$ +1.7 (c, 0.4 in CH_2Cl_2). λ_{max} 225 (log ϵ 4.72); 284 (log ϵ 4.13); 320 (log ϵ 3.4) (MeOH).

Rasoanaivo, P. *et al.*, *Tetrahedron*, 1995, **51**,


1221 (*isol, uv, cd, pmr, cmr, ms, struct*)

Rasoanaivo, P. *et al.*, *Planta Med.*, 1998, **64**,

58-62 (*Herveline D*)

Het₃₈₉

[128941-10-6]



$C_{21}H_{35}N_5O_2$ 389.54

Isol. from venom of the trap-door spider *Hebestatis theveniti* and the tarantula *Harpactirella* sp.

Skinner, W.S. *et al.*, *Toxicon*, 1990, **28**, 541

(*isol*)

Het₄₀₃

[128941-11-7]

As Het₃₈₉, H-155 with

$n = 5$

$C_{22}H_{37}N_5O_2$ 403.567


Isol. from venom of the trap-door spider *Hebestatis theveniti*.

Skinner, W.S. *et al.*, *Toxicon*, 1990, **28**, 541

(*isol*)

H-154**Heteratisine****H-157**

20-Ethyl-6,8-dihydroxy-1-methoxy-4-methylheteratisan-14-one, 9CI. Zeravschanine
[3328-84-5]



$C_{22}H_{33}NO_5$ 391.506

Alkaloid from the roots of *Aconitum heterophyllum*, *Aconitum tanguticum* and *Aconitum zeravschanicum* (Ranunculaceae). Mp 267-269°. $[\alpha]_D^{28}$ +40 (c, 1 in MeOH).

Hydrochloride: Mp 265-270° dec.

6-Ac: **6-Acetylheteratisine**

[52418-02-7]

$C_{24}H_{35}NO_6$ 433.544

Alkaloid from the roots of *Aconitum palmatum* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 156-158°. $[\alpha]_D^{26}$ +18.7 (c, 0.2 in CHCl₃).

6,8-Di-Ac: **6,8-Diacetylheteratisine**

[198422-45-6]

$C_{26}H_{37}NO_7$ 475.581

Constit. of *Aconitum pulchellum*.

6-Benzoyl: **6-Benzoylheteratisine**

[99759-48-5]

$C_{29}H_{37}NO_6$ 495.614

Alkaloid from the roots of *Aconitum heterophyllum* and *Aconitum tanguticum* (Ranunculaceae). Prisms (C_6H_6). Mp 213-214°. $[\alpha]_D^{25}$ +73 (c, 1.0 in EtOH).

O-De-Me: **Heterophyllidine**

[15266-43-0]

$C_{21}H_{31}NO_5$ 377.48

Trace alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Prisms (MeOH). Mp 269-272°. $[\alpha]_D^{26}$ +43.3 (c, 1.26 in MeOH).

6-Deoxy: **Heterophyllisine**

[15266-41-8]

$C_{22}H_{33}NO_4$ 375.507

Trace alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Fine needles (Me₂CO). Mp 178-179°. $[\alpha]_D^{27}$ +15.5 (c, 0.90 in MeOH).

6-Deoxy, 8-Ac: **8-Acetylheterophyllisine**

[187741-57-7]

$C_{24}H_{35}NO_5$ 417.544

Alkaloid from roots of *Delphinium demudatum*. Shows antifungal activity. Brown cryst. (2,3,3-trimethylpentane/ CH_2Cl_2). Mp 152°. $[\alpha]_D$ +38.24 (c, 0.052 in MeOH).

6-Deoxy, O-de-Me: **Heterophylline**[†]

[15266-42-9]

$C_{21}H_{31}NO_4$ 361.48

Trace alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Prisms (MeOH). Mp 221.5-223°. $[\alpha]_D^{30}$ +10.5 (c, 2.0 in MeOH). See also 17-Methylparsonsianidine, M-502 and Aricine, A-1423.

6-Deoxy, O-de-Me, 1-Ac:

Prismatic needles (Et₂O/petrol). Mp 174-176°.

6-Deoxy, 19-oxo, O-de-Me, 1,8-di-Ac:**Souline B**

[244246-52-4]

C₂₅H₃₃NO₇ 459.538

Alkaloid from *Delphinium souliei*. Cryst.

Jacobs, W.A. *et al.*, *J. Biol. Chem.*, 1942, **143**, 605; 1943, **147**, 571 (*Benzoylheteratisine, isol*)

Przybylska, M. *et al.*, *Can. J. Chem.*, 1963, **41**, 2911 (*Heteratisine, cryst struct*)

Pelletier, S.W. *et al.*, *Phytochemistry*, 1968, **7**, 625 (*Heterophylline, Heterophyllidine, Heterophyllisine*)

Aneja, R. *et al.*, *Tetrahedron*, 1973, **29**, 3297 (*Heteratisine, Benzoylheteratisine*)

Vaisov, Z.M. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 800; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 760 (*isol*)


Jiang, Q. *et al.*, *Tet. Lett.*, 1988, **29**, 1875

Desai, H.K. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2193 (*cmr*)

Chen, P. *et al.*, *CA*, 1997, **127**, 356972g (6,8-Diacetylheteratisine)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997, **60**, 472 (*8-Acetylheterophyllisine*)

Pan, X. *et al.*, *Chin. Chem. Lett.*, 1998, **9**, 57-59 (*Souline B*)

Heterocarpine**H-158**

C₁₉H₂₃NO₅ 345.394

Mp 105-106°.


(±)-form

Alkaloid from *Ceratocarpus heterocarpa*. Amorph. powder. Mp 105-106°. λ_{max} 226 (log ε 4.08); 284 (log ε 3.79) (MeOH).

Suau, R. *et al.*, *Phytochemistry*, 1998, **49**, 2551-2555 (*isol, synth, uv, ir, pmr, cmr, ms*)

Heteromine H**H-159**

[116137-80-5]



C₈H₁₃N₅O₂ 211.223

Alkaloid from *Heterostemma brownii* (Asclepiadaceae). Needles (MeOH). Mp 220-222°. λ_{max} 236 (log ε 3.44); 258 (log ε 3.46); 290 (log ε 3.3) (MeOH).

N²-Me: Heteromine G

[196090-70-7]

C₉H₁₅N₅O₂ 225.25

Alkaloid from *Heterostemma brownii*. Needles (MeOH). Mp 105-106°. λ_{max} 242 (log ε 3.61); 266 (log ε 3.62) (MeOH).

N²,N²-Di-Me: Heteromine F

[196090-69-4]

C₁₀H₁₇N₅O₂ 239.277

Alkaloid from *Heterostemma brownii*.

Needles (MeOH). Mp 176-177°. λ_{max}

247 (log ε 4.14); 268 (log ε 3.91)


(MeOH).

Kohda, K. *et al.*, *Tet. Lett.*, 1987, **28**, 6285-6288 (*synth*)

Lin, Y.-L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 982-985 (*isol, uv, ir, pmr, cmr, ms*)

Heteropsine**H-160**

[112543-99-4]



C₃₅H₂₈N₂O₄ 540.617

Alkaloid from the bark of *Unonopsis spectabilis* and roots of *Piptostigma fugax* (Annonaceae). Amorph.

N⁶-Me: N-Methylheteropsine

[112523-85-0]

C₃₆H₃₀N₂O₄ 554.644

Alkaloid from roots of *Piptostigma fugax* (Annonaceae). Powder. Mp 267-269°.


Laprévôt, O. *et al.*, *J. Nat. Prod.*, 1987, **50**, 984 (*isol, uv, ir, pmr, ms, struct*)

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1037 (*N-Methylheteropsine*)

Hetidine**H-161**

2,3-Dihydroxy-21-methyl-6,21-secohetisan-6,13-dione, 9CI

[28154-35-0]



C₂₁H₂₇NO₄ 357.449

Alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Cryst. (C₆H₆/hexane). Mp 218-221°.

Hydroiodide:

Cryst. (MeCN). Mp 305-307°.

3-O-(2-Methylpropanoyl), 2-Ac: **Contorsine** 2-Acetyl-3-isobutyrylheteridine

[144398-40-3]

C₂₇H₃₅NO₆ 469.577

Alkaloid from roots of *Aconitum contortum* (Ranunculaceae). Prisms (Me₂CO). Mp 203-206°. [α]_D -88.1 (c, 0.62 in EtOH).

3-O-(2S-Methylbutanoyl), 2-Ac: **Contortine**

[144398-39-0]

C₂₈H₃₇NO₆ 483.603

Alkaloid from roots of *Aconitum contortum* (Ranunculaceae). Prisms (Me₂CO). Mp 230-233°. [α]_D -82.1 (c,

0.45 in EtOH).

3-Benzoyl, 2-Ac: **Episcopalidine**

[83685-24-9]

C₃₀H₃₃NO₆ 503.594

Alkaloid from the roots of *Aconitum episcopale* (Ranunculaceae).

3-O-(4-Methoxybenzoyl), 2-Ac: **Contortine**

2-Acetyl-3-anisoylheteridine

[144398-38-9]

C₃₁H₃₅NO₇ 533.62

Alkaloid from roots of *Aconitum contortum* (Ranunculaceae). Prisms (Me₂CO). Mp 239° dec. [α]_D -44.9 (c, 0.35 in EtOH).

Δ¹⁵-Isomer: Racemulodine

C₂₁H₂₇NO₄ 357.449

Alkaloid from *Aconitum racemosum* var. *pengzhouense*. Needles (cyclohexane/Me₂CO). Mp 181-183°. [α]_D¹⁷ -24.9 (c, 0.2 in EtOH).

3-Deoxy: **Deacetylheterophyllidine**

[78174-98-8]

C₂₁H₂₇NO₃ 341.449

Alkaloid from aerial parts of *Delphinium albiflorum* (Ranunculaceae). Mp 160-162° (154-158°). [α]_D -59.9 (c, 0.76 in CHCl₃).

3-Deoxy, 2-Ac: **Heterophyllidine**. *Panicutine*

[78174-97-7]

C₂₃H₂₉NO₄ 383.486

Minor alkaloid from *Aconitum heterophyllum*, also *isol.* from *Aconitum paniculatum* (Ranunculaceae). Amorph. or *cryst.* (Et₂O/pentane). Mp 159-161°. [α]_D²⁴ -82 (c, 1.5 in CHCl₃). [α]_D²⁵ -141.1 (c, 0.429 in CHCl₃).

3-Deoxy, 2-ketone: **2-Dehydrodeacetylheterophyllidine**

C₂₁H₂₅NO₃ 339.433

Alkaloid from *Delphinium pentagynum*. Amorph. solid. [α]_D²⁵ -73.3 (c, 0.17 in CHCl₃).

13-Deoxo, 3-deoxy, 2-ketone: **Variogatine**

C₂₁H₂₇NO₂ 325.45

Alkaloid from *Aconitum variegatum*. Amorph. solid. [α]_D²⁰ +20.5 (c, 0.41 in CHCl₃).

[81425-75-4, 31260-17-0]

Pelletier, S.W. *et al.*, *Phytochemistry*, 1968, **7**, 625-635 (*Hetidine*)

Pelletier, S.W. *et al.*, *Chem. Comm.*, 1970, 393 (*Hetidine, cryst struct*)

Pelletier, S.W. *et al.*, *Tet. Lett.*, 1981, **22**, 313 (*Heterophyllidine*)

Katz, A. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 286 (*Panicutine*)

Wang, F. *et al.*, *Yaoxue Xuebao*, 1983, **18**, 514-521; 1985, **20**, 436-445; *CA*, **100**, 64981m; **105**, 24476x (*Episcopalidine, struct, ms*)

Sun, F. *et al.*, *Youji Huaxue*, 1985, 395-398; *CA*, **105**, 24482w (*Episcopalidine, cd*)

Chen, J. *et al.*, *CA*, 1986, **105**, 60804m (*Episcopalidine, ir*)

Pelletier, S.W. *et al.*, *Heterocycles*, 1986, **24**, 1275 (*Heterophyllidine, Panicutine, struct*)

Wang, F. *et al.*, *Youji Huaxue*, 1986, 19-24; *CA*, **105**, 115252n (*Episcopalidine, cmr*)


Niitsu, K. *et al.*, *Heterocycles*, 1992, **34**, 1231 (*Contortine, Contortine, Contorsine*)

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1555-1561 (*Deacetylheterophyllidine*)

Peng, C.S. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 411-414 (*Racemulodine*)

Díaz, J.G. *et al.*, *Phytochemistry*, 2004, **65**, 2123-2127; 2005, **66**, 837-846 (*2-Dehydrodeacetyl-heterophylloidine, Variegatine*)

Hetisan-1,15-diol H-162



C₂₀H₂₇NO₂ 313.439

(1β,15β)-form

Davisine. *Cossonidine*
[173220-99-0]

Alkaloid from aerial parts of *Delphinium davisii*, *Delphinium cossonianum* and *Delphinium cardiopetalum*. Cryst. (Me₂CO/hexane). Mp 130-132° Mp 243-245° (dec.). [α]_D²⁵ +29.9 (c, 1.56 in CHCl₃). Large discrepancy in Mp's noted; one may refer to a salt.


Di-Ac:

Amorph. [α]_D²⁵ +6 (c, 0.75 in CHCl₃).

Reina, M. *et al.*, *J. Nat. Prod.*, 1996, **59**, 145 (*isol, ir, pmr, cmr, ms, struct*)

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 360 (*isol, pmr, cmr, struct*)

Hetisan-2,15-diol, 9CI H-163



C₂₀H₂₇NO₂ 313.439

(2α,15β)-form


Sanyonamine. *2-Hydroxynominine*. *Katsuyama base I*. *Dogo base*
[85352-56-3]

Minor alkaloid from *Aconitum sanyoense*. Mp 276-278°. [α]_D +62.9. λ_{max} 224 (ε 20000); 251 (ε 3600); 285 (ε 3380) (MeOH) (Derop).

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1124 (*struct*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4576; 4579 (*ir, pmr, ms, cryst struct, abs config*)

Hetisan-6,11-diol, 9CI H-164



11α-form

C₂₀H₂₇NO₂ 313.439

11α-form

Venulol

[102517-36-2]

Alkaloid from the aerial parts of *Delphinium venulosum* (Ranunculaceae). [α]_D²² +19.7 (c, 0.15 in MeOH).

11ξ-form

Spiradine B

[19741-51-6]

Alkaloid from *Spiraea japonica*. Mp 259-260°. [α]_D²⁵ -13 (c, 0.25 in CHCl₃). pK_a 9.54 (66% MeOH aq.).

11-Ac: Spiradine C

[19741-52-7]

C₂₂H₂₉NO₃ 355.476

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 248-249°. pK_a 9.02 (66% MeOH).

11-Ketone: 6-Hydroxyhetisan-11-one,

9CI. Spiradine A

[19741-46-9]

C₂₀H₂₅NO₂ 311.423

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 281-282°. pK_a 8.35 (50% MeOH).

11-Ketone; methiodide: Mp 330°.

Goto, G. *et al.*, *Tet. Lett.*, 1968, 1369 (*Spiradines*)

Sasaki, K. *et al.*, *J.C.S.(B)*, 1971, 354


(*Spiradine A, cryst struct*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4579 (*Spiradine A, cd*)

Ulubelen, A. *et al.*, *Phytochemistry*, 1992, **31**, 3239 (*Venulol*)

Fan, L.-M. *et al.*, *J. Integ. Plant Biol.*, 2005, **47**, 120-123 (*Spiradine A,B, pmr, cmr*)

Hetisan-6,13-diol H-165



C₂₀H₂₇NO₂ 313.439

(13R)-form

Spirasine XV

[115610-49-6]

Minor alkaloid from the roots of *Spiraea japonica* var. *fortunei* (Rosaceae). Cryst. (CH₂Cl₂/EtOAc). Mp 156-158°. [α]_D¹⁷ 0 (EtOH).

(13S)-form


Spirasine XIV

[115610-48-5]

Minor alkaloid from the roots of *Spiraea japonica* var. *fortunei* (Rosaceae). Needles (CH₂Cl₂/EtOAc). Mp 244-246°. [α]_D¹⁷ -18.8 (c, 1.6 in EtOH).

Sun, F. *et al.*, *J. Nat. Prod.*, 1988, **51**, 50 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

Hetisan-9,15-diol H-166



C₂₀H₂₇NO₂ 313.439

(15β)-form


2-Desoxyryosenaminol. *9-Hydroxynominine*

[91197-66-9]

Alkaloid from *Aconitum ibukiense* (Ranunculaceae). Mp 287-291° dec. [α]_D²³ +68.5 (c, 0.08 in MeOH).

Sakai, S. *et al.*, *Yakugaku Zasshi*, 1984, **104**, 222; *CA*, **101**, 69329k (*Desoxyryosenaminol*)

Hetisan-11,15-diol H-167



C₂₀H₂₇NO₂ 313.439

▶ MK7000000

(11β,15β)-form

Kobusine⁺

[27530-78-5]

Alkaloid from *Aconitum sachalinense*, *Aconitum yesoense*, *Aconitum lucidusculum* and *Aconitum kamtschaticum*, also *Delphinium davisii* (Ranunculaceae). Prisms (Me₂CO). Mp 267-267.5°. [α]_D¹² +104.4 (MeOH).

Hydrobromide:

Cryst. (EtOH). Mp 285° dec.

Picrate: Mp 282-284° dec.

11-Ketone: 15-Hydroxyhetisan-11-one,

9CI. 11-Dehydrokobusine

[85352-58-5]

C₂₀H₂₅NO₂ 311.423

Alkaloid from aerial parts of *Aconitum talassicum* (Ranunculaceae). Cryst. (Me₂CO). Mp 239-241°.

Suginome, H. *et al.*, *Annalen*, 1940, **545**, 220-228 (*Kobusine, isol*)

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1959, **7**, 44-49; 1962, **10**, 883-886 (*struct*)

Pelletier, S.W. *et al.*, *Chem. Comm.*, 1970, 98-99 (*cryst struct*)

Scott, A.I. *et al.*, *Tetrahedron*, 1971, **27**, 4787-4819 (*cd*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4579-4582; 4583-4584 (*abs config, synth*)


Nishanov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1989, **25**, 857; *Chem. Nat. Compd. (Engl.*

Transl., 1989, **25**, 728 (*11-Dehydrokobusine*)

Wada, K. *et al.*, *Heterocycles*, 1991, **32**, 1297-1300 (*synth*)

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 360-366 (*pmr, cmr*)

Hetisan-11,18-diol H-168



$C_{20}H_{27}NO_2$ 313.439

11 β -form

Davisinol

[175097-22-0]

Alkaloid from aerial parts of *Delphinium davisii*. Amorph. $[\alpha]_D^{25} +27.5$ (c, 0.189 in $CHCl_3$).

18-Benzoyl: 18-Benzoyldavisinol


[174587-13-4]

$C_{27}H_{31}NO_3$ 417.547

Alkaloid from aerial parts of *Delphinium davisii*. Amorph. $[\alpha]_D^{25} +42.3$ (c, 0.26 in $CHCl_3$).

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 360 (*isol, ir, pmr, cmr, struct*)

Hetisan-1,2,3,11,13,14-hexol H-169



(1 β ,2 α ,3 α ,11 α ,13R)-form

$C_{20}H_{27}NO_6$ 377.436

(1 β ,2 α ,3 α ,11 α ,13R)-form

13-Benzoyl, 2-O-(2-methylbutanoyl),

1,3,11-tri-Ac: Cardiodine

[174476-66-5]

$C_{38}H_{45}NO_{11}$ 691.774

Alkaloid from aerial parts of *Delphinium cardiopetalum*. Resin. $[\alpha]_D -26$ (c, 0.05 in EtOH).

(1 β ,2 α ,3 α ,11 α ,13S)-form

2,13-Dibenzoyl, 1,3,11,-tri-Ac: Delphigraciline

[1015757-95-5]


$C_{40}H_{41}NO_{11}$ 711.764

Alkaloid from *Delphinium gracile*. Cryst. (hexane/EtOAc). Mp 208-212°. $[\alpha]_D -95$ (c, 0.02 in $CHCl_3$).

Reina, M. *et al.*, *Phytochemistry*, 1996, **41**, 1235-1250 (*Cardiodine*)

Reina, M. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1048-1055 (*Delphigraciline*)

Hetisan-2,3,9,11,13,14-hexol H-170



$C_{20}H_{27}NO_6$ 377.436

(2 α ,3 α ,11 α ,13R)-form

2-O-(2-Methylbutanoyl), 3-Ac: Glanduline

[188559-93-5]

$C_{27}H_{37}NO_8$ 503.591

Alkaloid from aerial parts of *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 134-137°. $[\alpha]_D +24$ (c, 0.5 in MeOH).

2-O-(2-Methylbutanoyl), 3,9-di-Ac: 9-O-Acetylglanduline

$C_{29}H_{39}NO_9$ 545.628

Alkaloid from *Consolida glandulosa*. Cryst. (hexane/EtOAc/MeOH). Mp 147-150°. $[\alpha]_D^{25} +5.1$ (c, 0.14 in EtOH).

2-O-(2-Methylbutanoyl), 3,13-di-Ac: 13-O-Acetylglanduline

[188560-00-1]

$C_{29}H_{39}NO_9$ 545.628

From aerial parts of *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 110-115°. $[\alpha]_D +15.2$ (c, 0.46 in MeOH).

2-O-(2-Methylbutanoyl), 3,11,13-tri-Ac: 11,13-Di-O-acetylglanduline

[425382-34-9]


$C_{31}H_{41}NO_{10}$ 587.666

Alkaloid from *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 240-241°. $[\alpha]_D^{25} +60.4$ (c, 0.39 in $CHCl_3$).

Almanza, G. *et al.*, *Phytochemistry*, 1997, **44**, 739 (*isol, ir, pmr, cmr, ms, struct*)

Ruiz-Mesia, L. *et al.*, *J. Nat. Prod.*, 2002, **65**, 496-499 (*9-O-Acetylglanduline, 11,13-Diacetylglanduline*)

Hetisan-13-ol, 9CI H-171



$C_{20}H_{27}NO$ 297.439

(13 ξ)-form

Spirasine XI

[102358-20-3]

Alkaloid from the roots of *Spiraea japonica* (Rosaceae). Cryst. Mp 286-288°. $[\alpha]_D^{11} -23.8$ (c, 0.84 in $CHCl_3$). C-13 Config. ambiguously shown.


13-Ketone: Spirasine IV. Hetisan-13-one [102386-48-1]

$C_{20}H_{25}NO$ 295.424

Alkaloid from the roots of *Spiraea japonica* (Rosaceae). Powder. $[\alpha]_D^{17} -95.7$ (c, 1.1 in $CHCl_3$).

Sun, F. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 913-917 (*Spirasines IV,XI, isol, struct*)

Hetisan-15-ol H-172



$C_{20}H_{27}NO$ 297.439

15 β -form

Nominine†. Nomi-base I. 11-Deoxykobusine

[79808-87-0]

Minor alkaloid from *Aconitum sanyoense* (Ranunculaceae). Mp 251-54°. $[\alpha]_D^{24} +53.4$. λ_{max} 224 (ϵ 20000); 251 (ϵ 3600); 285 (ϵ 3380) (MeOH) (Derep).

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4576-4578; 4579-4582 (*ir, pmr, ms, cryst struct, abs config*)


Peese, K.M. *et al.*, *J.A.C.S.*, 2006, **128**, 8734-8735 (*synth*)

Muratake, H. *et al.*, *Tetrahedron*, 2006, **62**, 7093-7112 (*synth*)

Hetisan-11-one H-173

Spirasine IX

[102386-47-0]




$C_{20}H_{25}NO$ 295.424

Alkaloid from the roots of *Spiraea japonica* (Rosaceae). Cryst. Mp 157-158°. $[\alpha]_D^{21} +135.5$ (c, 1 in $CHCl_3$).

Sun, F. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 913-917 (*isol, struct*)

Hetisan-1,2,3,11,13-pentol H-174



$C_{20}H_{27}NO_5$ 361.437

(1 β ,2 α ,3 α ,11 α ,13S)-form

13-Benzoyl, 2-O-(2-methylpropanoyl),

1,11-di-Ac: Cardiopinine

[174423-46-2]
C₃₅H₄₁NO₉ 619.71
Alkaloid from aerial parts of *Delphinium cardiopetalum*. Cryst. (EtOAc/hexane). Mp 218-220°. [α]_D -26.6 (c, 0.08 in EtOH).

13-Benzoyl, 3-O-(2-methylpropanoyl), 1,11-di-Ac: Cardiopinine

[174423-47-3]
C₃₅H₄₁NO₉ 619.71
Alkaloid from *Delphinium cardiopetalum*. Resin. [α]_D -81.3 (c, 0.2 in EtOH).

13-Benzoyl, 2-O-(2-ξ-methylbutanoyl), 1,11-di-Ac: Cardiopine


[174423-45-1]
C₃₆H₄₃NO₉ 633.737
Alkaloid from aerial parts of *Delphinium cardiopetalum*. Cryst. (EtOAc/hexane). Mp 194-197°. [α]_D -26.3 (c, 0.06 in EtOH).

13-Benzoyl, 3-O-(2-ξ-methylbutanoyl), 1,11-di-Ac: Cardiopidine

[174423-48-4]
C₃₆H₄₃NO₉ 633.737
Alkaloid from aerial parts of *Delphinium cardiopetalum*. Resin. [α]_D -22.5 (c, 0.05 in EtOH).

Reina, M. *et al.*, *Phytochemistry*, 1996, **41**, 1235-1250 (*isol, ir, pmr, cmr, ms, struct*)

Hetisan-1,2,11,13,19-pentol H-175




C₂₀H₂₇NO₅ 361.437

(1α,2α,11α,13ξ,19ξ)-form

2-Benzoyl: Delgramine
[152434-56-5]
C₂₇H₃₁NO₆ 465.545
Alkaloid from aerial parts of *Delphinium grandiflorum* (Ranunculaceae).

Li, C. *et al.*, *Huaxue Xuebao*, 1993, **51**, 915; *CA*, **120**, 73352x

Hetisan-2,3,6,11,13-pentol H-176



C₂₀H₂₇NO₅ 361.437


(2α,3ξ,11α,13R)-form

11-O-(2-ξ-Methylbutanoyl), 3-Ac: Geyerinine

[100045-30-5]
C₂₇H₃₇NO₇ 487.592
Minor alkaloid from the flowers, small stems and leaves of *Delphinium geberi* (Ranunculaceae). Gum.

Grina, J.A. *et al.*, *J.O.C.*, 1986, **51**, 390 (*isol, uv, ir, pmr, cmr, ms, struct*)

Hetisan-2,3,7,9,15-pentol H-177




C₂₀H₂₇NO₅ 361.437

(2α,3β,7α,15β)-form

2-Benzoyl: Sadosine
[85753-45-3]
C₂₇H₃₁NO₆ 465.545
Minor alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Needles + 1H₂O (Me₂CO). Mp 222-224°. [α]_D²⁵ +53.1 (c, 0.96 in MeOH).

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 360 (*isol, pmr, cmr, ms, cryst struct*)

Hetisan-2,3,11,13,14-pentol H-178



C₂₀H₂₇NO₅ 361.437

(2α,3α,11α,13R)-form

2-O-(2-Methylpropanoyl), 3,11,13-tri-Ac: Glandulosine†
[425382-33-8]
C₃₀H₃₉NO₉ 557.639
Alkaloid from *Consolida glandulosa*. Amorph. solid. [α]_D²⁵ +120 (c, 0.1 in CHCl₃). Not to be confused with Glandulosine in M-309.

2-O-(2-Methylbutanoyl), 3-Ac: 9-Deoxyglanduline

[425382-32-7]
C₂₇H₃₇NO₇ 487.592
Alkaloid from *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 174-176°. [α]_D²⁵ +6.6 (c, 0.19 in CHCl₃).

2-O-(2-Methylbutanoyl), 3,13-di-Ac: 13-O-Acetyl-9-deoxyglanduline

[188560-03-4]
C₂₉H₃₉NO₈ 529.629
From aerial parts of *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 154-156°. [α]_D +46.6 (c, 0.39 in MeOH).


2-O-(2-Methylbutanoyl), 3,14-di-Ac: 14-O-Acetyl-9-deoxyglanduline

[188560-01-2]
C₂₉H₃₉NO₈ 529.629
From aerial parts of *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 145-148°. [α]_D +20 (c, 0.2 in MeOH).

2-O-(2-Methylbutanoyl), 3,11,13-tri-Ac: 11,13-Di-O-acetyl-9-deoxyglanduline

[188560-04-5]
C₃₁H₄₁NO₉ 571.666
From aerial parts of *Consolida glandulosa*. Cryst. (hexane/EtOAc). Mp 195-198°. [α]_D +36 (c, 0.50 in MeOH).
Almanza, G. *et al.*, *Phytochemistry*, 1997, **44**, 739 (*isol, ir, pmr, cmr, ms, struct*)
Ruiz-Mesia, L. *et al.*, *J. Nat. Prod.*, 2002, **65**, 496-499 (*Glandulosine, 9-Deoxyglanduline*)

Hetisan-1,2,9,15-tetrol, 9CI H-179



C₂₀H₂₇NO₄ 345.438

▶MK730000

(1β,2α,15β)-form

Hypognavinol
[32199-38-5]
Hydrolyt. prod. of Hypognavine. Prisms (MeOH). Mp 307-308°. [α]_D²⁴ +67.7 (MeOH).

Hydrochloride:

Prisms (MeOH/Me₂CO). Mp 310° dec.

2-Benzoyl: Hypognavine

[79808-88-1]
C₂₇H₃₁NO₅ 449.546
Alkaloid from *Aconitum sanyoense* (Ranunculaceae). Needles (Me₂CO). Mp 239-241°. [α]_D²⁴ +127.1 (MeOH).

2-Benzoyl, 1-Ac: 1-O-Acetylhypognavine

[148225-50-7]
C₂₉H₃₃NO₆ 491.583
Alkaloid from roots of *Aconitum sanyoense* var. *tonense* (Ranunculaceae). Prisms (Me₂CO). Mp 127-128° dec. [α]_D³⁰ +116.7 (c, 0.21 in CHCl₃).

2-Benzoyl, 1,15-di-Ac: 1,15-Di-O-acetylhypognavine

[120527-39-1]
C₃₁H₃₅NO₇ 533.62
Alkaloid from roots of *Aconitum sanyoense* var. *tonense* (Ranunculaceae). Amorph. powder; cryst. (EtOH aq., as picrate). Mp 233-239° dec. (picrate). [α]_D³⁰ +83 (c, 0.44 in CHCl₃).

Ochiai, E. *et al.*, *Chem. Pharm. Bull.*, 1953, **1**, 152 (*Hypognavine, isol*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1957, **5**, 1; 1958, **6**, 448; 1959, **7**, 50,55 (*Hypognavine, struct*)


Pelletier, S.W. *et al.*, *Tet. Lett.*, 1971, 795 (*cryst struct, deriv*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4573 (*Hypognavine, struct, abs config*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 2927 (*1-Acetylhyppognavine, 1,15-Diacetylhyppognavine*)

Hetisan-1,2,9,19-tetrol

H-180



(1β,2α,19R)-form

C₂₀H₂₇NO₄ 345.438

(1β,2α,19R)-form

Septentriosine

[115569-73-8]

Minor alkaloid from the roots of *Aconitum septentrionale* (Ranunculaceae). Cryst. (MeOH). Mp 260-262°. [α]_D²⁸ +20.78 (c, 0.55 in MeOH).

Hydrochloride:

Cryst. (MeOH). Mp 299-300°. [α]_D -7.5 (c, 1.0 in MeOH).

2-Ac: 2-Acetylseptentriosine

[147677-10-9]

C₂₂H₂₉NO₅ 387.475

Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Cryst. + 1H₂O (Et₂O/hexane), cryst. (Et₂O/MeOH). Mp 182-184° (with swelling and darkening at 150-156°). [α]_D²⁵ +6.4 (c, 1 in EtOH).

1,2,19-Tri-Ac: [147677-11-0]

Cryst. (Me₂CO). Mp 210.5-212.5°.

(1β,2α,19S)-form

2-Ac: Septenine

[169626-10-2]

C₂₂H₂₉NO₅ 387.475

Alkaloid from the roots of *Aconitum septentrionale*. Cryst. (CHCl₃/MeOH). Mp 190-192°.


Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 265 (*Septentriosine, isol, ir, pmr, cmr, ms, cryst struct*)

Usmanova, S.K. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 349-351 (*Septenine*)

Ross, S.A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 424 (*2-Acetylseptentriosine*)

Hetisan-1,2,11,13-tetrol

H-181



(1β,2α,11α,13R)-form

C₂₀H₂₇NO₄ 345.438

(1β,2α,11α,13R)-form

2-Benzoyl, 1,11-di-Ac: Tadzhaconine. Anthoroidine

[11087-94-8]

C₃₁H₃₅NO₇ 533.62

Alkaloid from *Aconitum zeravschani-*

cum and Aconitum anthoroidine (Ranunculaceae). Cryst. (EtOH). Mp 236-237° (dec.).

(1β,2β,11α,13R)-form

11-Benzoyl, 1,2-di-Ac: Paniculatin†

[1400-72-2]

C₃₁H₃₅NO₇ 533.62

Alkaloid from *Aconitum paniculatum* (Ranunculaceae). Rhombic prisms (MeOH). Mp 263°. See also Lycopaniculatin, L-334. Struct. revised in 1989.

Brunner, G.E. *et al.*, *Schweiz. Apoth.-Ztg.*, 1922, **60**, 357; *CA*, **16**, 2962 (*Paniculatin, isol*)

Staehelin, E. *et al.*, *Pharm. Acta Helv.*, 1980, **55**, 221; *CA*, **93**, 201040g (*Paniculatin, isol*)


Katz, A. *et al.*, *J. Nat. Prod.*, 1989, **52**, 430 (*Paniculatin, uv, ir, pmr, cmr, ms, struct*)

Yusupova, I.M. *et al.*, *Khim. Prir. Soedin.*, 1992, **28**, 382-388; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 335-339 (*Tadzhaconine*)

Vaisov, Z.M. *et al.*, *Khim. Prir. Soedin.*, 1993, 86-87 (*Anthoroidine*)

Hetisan-2,3,6,13-tetrol

H-182



C₂₀H₂₇NO₄ 345.438

(2α,3α,13S)-form

N-Me: Vakhmadine

[135250-81-6 (hydroxide)]


C₂₁H₃₀NO₄[⊕] 360.472

Quaternary alkaloid from *Aconitum palmatum* (Ranunculaceae). Cryst. (EtOH) (as hydroxide). Mp 263-273° (hydroxide). [α]_D²⁴ -37.8 (c, 0.1 in MeOH).

Jiang, Q. *et al.*, *J. Nat. Prod.*, 1991, **54**, 525 (*isol, ir, pmr, cmr, ms, struct*)

Hetisan-2,3,7,13-tetrol

H-183



C₂₀H₂₇NO₄ 345.438

(2α,3β,7β,13S)-form

2-Benzoyl, 13-Ac: Jaluene

C₂₉H₃₃NO₆ 491.583


Alkaloid from the roots of *Aconitum jaluense*. Amorph. solid (MeOH). Mp

114-116°. [α]_D²³ +19.3 (c, 0.27 in CHCl₃).

Shim, S.H. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 451-455 (*isol, pmr, cmr, ms*)

Hetisan-2,3,9,15-tetrol

H-184



(2α,3α,15β)-form

C₂₀H₂₇NO₄ 345.438

(2α,3α,15β)-form

3-Epiignavinol

[119240-18-5]

Alkaloid from the roots of *Aconitum japonicum*, var. *montanum* (Ranunculaceae). Prisms. Mp 292-293.5° dec. [α]_D²³ +49.1 (c, 0.11 in MeOH).

(2α,3β,15β)-form

Anhydroignavinol

[30373-79-6]

Alkaline hydrolyt. prod. of Ignavine. Mp 302-304°.

2-Benzoyl: Ignavine

[1357-76-2]

C₂₇H₃₁NO₅ 449.546

Alkaloid from *Aconitum sanyoense*, *Aconitum tasiro-montanum* and the roots of *Aconitum japonicum* and *Aconitum carmichaeli* (Ranunculaceae). Mp 172-174°. [α]_D +85.3 (EtOH).

▶ MK7350000

Ochiai, E. *et al.*, *Yakugaku Zasshi*, 1956, **76**, 550-553; *CA*, **50**, 13372f (*Ignavine, isol*)

Ochiai, E. *et al.*, *Chem. Pharm. Bull.*, 1959, **7**, 556-559 (*Ignavine, struct*)

Pelletier, S.W. *et al.*, *Tet. Lett.*, 1970, 4825-4827 (*cryst struct*)

Saito, H. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1844-1850 (*Ignavine, pharmacol*)

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4600-4601 (*abs config, cryst struct*)


Takayama, H. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 525-532; *CA*, **97**, 141690q (*Ignavine, isol*)

Hikino, H. *et al.*, *J. Nat. Prod.*, 1983, **46**, 178-182 (*Ignavine, isol*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3210-3212 (*3-Epiignavinol, cryst struct*)

Hetisan-2,3,11,13-tetrol

H-185



C₂₀H₂₇NO₄ 345.438

(2β,3α,11α,13S)-form


2-Benzoyl, 3,13-di-Ac: Cossonine

[152343-38-9]
 $C_{31}H_{35}NO_7$ 533.62
 Alkaloid from above-ground parts of *Delphinium cossonianum* (Ranunculaceae). Amorph. $[\alpha]_D^{25} +45$ (c, 0.16 in $CHCl_3$).

De la Fuente, G. *et al.*, *Phytochemistry*, 1993, **34**, 553 (*isol, ir, pmr, cmr, ms, struct*)

Hetisan-2,6,7,13-tetrol

H-186



$C_{20}H_{27}NO_4$ 345.438

(2 α ,7 α ,13R)-form

13-Ketone, N-Me, 2-Ac: **2-O-Acetyl-7-hydroxyorochrine**

[959853-60-2]


$C_{23}H_{30}NO_5^{\oplus}$ 400.494

Alkaloid from *Aconitum orochryseum*. Pale green solid (as hydroxide). Mp 110-112° (hydroxide). $[\alpha]_D^{25} -31.8$ (c, 0.34 in MeOH) (hydroxide).

Wangchuk, P. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1808-1811 (*isol, pmr, cmr, ms*)

Hetisan-2,7,9,15-tetrol

H-187



$C_{20}H_{27}NO_4$ 345.438

(2 α ,7 α ,15 β)-form

2-Benzoyl: **Torokonine. 7-Hydroxyryosamine**

[110397-61-0]


$C_{27}H_{31}NO_5$ 449.546

Alkaloid from the roots of *Aconitum subcuneatum* (Ranunculaceae). Cryst. (Me_2CO). Mp 198.5-199°. $[\alpha]_D^{25} +71.7$ (c, 0.14 in MeOH).

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2615 (*Torokonine*)

Hetisan-2,7,13,15-tetrol

H-188



$C_{20}H_{27}NO_4$ 345.438

(2 α ,7 β ,13R,15 β)-form


Tanwusine A†

Alkaloid from *Aconitum tanguticum* (Ranunculaceae). Not the same as Tanwusine in H-203 CAS No. not found to 2006.

Liang, X.-T. *et al.*, *Pure Appl. Chem.*, 1986, **58**, 711 (*rev*)

Hetisan-2,11,13,14-tetrol

H-189



$C_{20}H_{27}NO_4$ 345.438

13-Configs. in these alkaloids are difficult to interpret from the accessible lit. and may not be as indicated here.

(2 α ,11 α ,13R)-form

Tangutisine. 14-Hydroxyhetisine

[139219-95-7]

Alkaloid from *Aconitum tanguticum* (Ranunculaceae). Cryst. (H_2O) (as hydrochloride). Mp 310-315° dec. (hydrochloride).

2-Ac: **Acorine. Guan-fu base I. Guan-fu base Y**

[110225-59-7]

$C_{22}H_{29}NO_5$ 387.475

Alkaloid from the roots of *Aconitum koreanum*. Cryst. (Me_2CO). Mp 218-219°.

Ac: **Guan-fu base B**

[1394-49-6]

$C_{22}H_{29}NO_5$ 387.475

Alkaloid from the Chinese herbal drug Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Mp 204°. $[\alpha]_D^{26} +16$ ($CHCl_3$). Site of acylation (2, 11 or 13) not known.

2,13-Di-Ac: **Guan-fu base A**

[1394-48-5]

$C_{24}H_{31}NO_6$ 429.512

Alkaloid from Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Shows antiarrhythmic props. Mp 198°. $[\alpha]_D^{23.8} +49$ ($CHCl_3$).

► LD₅₀ (mus, ipr) 421 mg/kg. LD₅₀ (mus, ivn) 134 mg/kg. MK7900000

2,11,13-Tri-Ac: **Guan-fu base G**

[78969-72-9]

$C_{26}H_{33}NO_7$ 471.549

Alkaloid from Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Mp 178°.

Tetra-Ac: Mp 154.5-155°.

2-Propanoyl: **Acoridine**

[144425-27-4]

$C_{23}H_{31}NO_5$ 401.502

Alkaloid from epigeal parts of *Aconitum koreanum* (Ranunculaceae). Needles ($Me_2CO/CHCl_3$). Mp 204-206°. $[\alpha]_D^{25} +16$ (c, 0.663 in MeOH).

2-Propanoyl, 13-Ac: **13-Acetylacoridine. Guan-fu base O**

[150901-10-3]

[252565-78-9]

$C_{25}H_{33}NO_6$ 443.539

Alkaloid from *Aconitum koreanum*.

2-Propanoyl, 11,13-di-Ac: **Guan-fu base R**

[1004550-87-1]

$C_{27}H_{35}NO_7$ 485.576

Alkaloid from the roots of *Aconitum koreanum*.

2-O-(2-Methylpropanoyl): **2-Isobutyryl-14-hydroxyhetisine. Guan-fu base Z**

[103847-13-8]

$C_{24}H_{33}NO_5$ 415.528

Alkaloid from Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Needles. Mp 230-231°.

2-O-(2-Methylpropanoyl), N-oxide: **Guan-fu base Z N-oxide**

[131916-92-2]

$C_{24}H_{33}NO_6$ 431.528

Alkaloid from epigeal parts of *Aconitum koreanum* (Ranunculaceae). Needles. Mp 317-319°. $[\alpha]_D^{20} +25$ (c, 0.34 in EtOH).

2-O-(2-Methylpropanoyl), 13-Ac: **Guan-fu base F**

[79030-10-7]

$C_{26}H_{35}NO_6$ 457.566

Alkaloid from Guan-Bai-Fu-Tzu (*Aconitum koreanum*) (Ranunculaceae). Mp 184°.

2-O-(2-Methylpropanoyl), 13-Ac, N-oxide: **Guan-fu base F N-oxide**

[131966-22-8]

$C_{26}H_{35}NO_7$ 473.565

Alkaloid from epigeal parts of *Aconitum koreanum* (Ranunculaceae). Mp 240-242°.

2-O-(2-Methylpropanoyl), 11,13-di-Ac: **Guan-fu base P**

[1004550-88-2]

$C_{28}H_{37}NO_7$ 499.603

Alkaloid from the roots of *Aconitum koreanum*.

13-Benzoyl, 2-Ac: **Zeravschanisine. Zeravschanizine**

[192705-33-2]

$C_{29}H_{33}NO_6$ 491.583

Alkaloid from *Aconitum zeravschanicum* (Ranunculaceae). Exhibits antiarrhythmic and anaesthetic activities. Cryst. (EtOH) (as perchlorate). Mp 287-289° dec. (perchlorate).

(2 α ,11 α ,13S)-form

2-Ketone, N-oxide: **14-Hydroxyhetisinone N-oxide**

[1015699-80-5]

$C_{20}H_{25}NO_5$ 359.421

Alkaloid from *Delphinium gracile*. Gum. $[\alpha]_D -125$ (c, 0.2 in EtOH).

Kao, H.-C. *et al.*, *Yaoxue Xuebao*, 1966, **13**, 186-194; *CA*, **65**, 3922g (*Guan-fu bases A, B, isol*)


Liu, J.-H. *et al.*, *Zhongcaoyao*, 1981, **12**, 1-2; *CA*, **95**, 138450q (*Guan-fu bases, struct*)

Reinecke, M.G. *et al.*, *Heterocycles*, 1986, **24**, 49-61 (*Guan-fu base Z, pmr, cmr, struct*)

Reinecke, M.G. *et al.*, *Tetrahedron*, 1986, **42**, 6621-6626 (*Guan-fu bases A, Y*)

- Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 690-692; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 573-575 (*Acorine*)
 Yusipova, I.M. *et al.*, *Khim. Prir. Soedin.*, 1990, **26**, 378-383; 383-386; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 314-318; 318-320 (*Guan-fu base F Z N-oxide*)
 Joshi, B.S. *et al.*, *Heterocycles*, 1991, **32**, 1793-1804 (*Tangutisine*)
 Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*, 1991, **27**, 91-93; *Chem. Nat. Compd. (Engl. Transl.)*, 1991, **27**, 79-81 (*Acoridine*)
 Salimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1992, **28**, 375-382; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 329-334 (*Zeravschanisine*)
 Wang, Y.P. *et al.*, *Zhongguo Yaoli Xuebao (Acta Pharmacol. Sin.)*, 1992, **13**, 231-234 (*Guan-fu base A, activity*)
 Liu, J. *et al.*, *CA*, 1993, **119**, 221622g (*13-Acetylacoridine*)
 Mericli, A.H. *et al.*, *Pharmazie*, 2000, **55**, 696-698 (*Guan-fu base Y, isol, pmr, cmr*)
 Yang, C. *et al.*, *Zhongcaoyao*, 2004, **35**, 1328-1330; *CA*, **144**, 11244 (*Guan-fu base P*)
 Jiang, K. *et al.*, *Yaoxue Xuebao*, 2006, **41**, 128-131 (*Guan-fu base R*)
 Reina, M. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1048-1055 (*14-Hydroxyhetisinone N-oxide*)

Hetisan-2,11,13,15-tetrol H-190



C₂₀H₂₇NO₄ 345.438

(11α,13R,15β)-form

2-Ketone: 11,13,15-Trihydroxyhetisan-2-one

2-Ketone, 11-benzoyl: **Cardiopetamine**

[88841-41-2]
 C₂₇H₂₉NO₅ 447.53

Alkaloid from *Delphinium cardiopetalum* and aerial parts of *Aconitum napellus* (Ranunculaceae). Mp 302-305° dec. [α]_D +65 (c, 1.4 in EtOH).

2-Ketone, 11-benzoyl, 15-Ac: **15-Acetylcardiopetamine**

[88840-05-5]
 C₂₉H₃₁NO₆ 489.567

Alkaloid from *Delphinium cardiopetalum* and aerial parts of *Aconitum napellus* (Ranunculaceae). Mp 236-237°. [α]_D +12 (c, 0.51 in EtOH).

2,13-Diketone: 11,15-Dihydroxyhetisan-2,13-dione

C₂₀H₂₃NO₄ 341.406

2,13-Diketone, 11-benzoyl, 15-Ac: **15-Acetyl-13-dehydrocardiopetamine**

[88840-09-9]
 C₂₉H₂₉NO₆ 487.551


Alkaloid from aerial parts of *Aconitum napellus* (Ranunculaceae). Mp 253-255°. [α]_D -46 (c, 0.03 in EtOH).

González, A.G. *et al.*, *Tet. Lett.*, 1983, **24**, 3765 (*Cardiopetamine, 15-*

Acetylcardiopetamine, uv, ir, pmr, cryst struct)

De la Fuente, G. *et al.*, *Heterocycles*, 1989, **29**, 1577 (*15-Acetyl-13-dehydrocardiopetamine*)

Hetisan-2,11,13,19-tetrol H-191



C₂₀H₂₇NO₄ 345.438

(2α,11α,13S,19S)-form

Vakhmatine

[135273-17-5]

Alkaloid from *Aconitum palmatum* (Ranunculaceae). Also isol. from seeds of *Consolida ambigua* (Ranunculaceae). Cryst. (MeOH). Mp 170.5-174.5°. [α]_D²⁴ +12.6 (c, 0.2 in MeOH).

13-Ac: **13-O-Acetylvakhmatine**

[172045-40-8]


C₂₂H₂₉NO₅ 387.475

Alkaloid from seeds of *Consolida ambigua* (Ranunculaceae). [α]_D -20 (c, 3.11 in CHCl₃).

Jiang, Q. *et al.*, *J. Nat. Prod.*, 1991, **54**, 525 (*isol, ir, pmr, cmr, ms, struct*)

Venkateswarlu, V. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1527 (*13-O-Acetylvakhmatine*)

Hetisan-6,11,12,15-tetrol H-192



C₂₀H₂₇NO₄ 345.438

(11β,15α)-form

15-O-(2-Methylpropanoyl): **Cardionine**

[123151-94-0]

C₂₄H₃₃NO₅ 415.528

Alkaloid from above-ground parts of *Delphinium cardiopetalum* (Ranunculaceae). Cryst. (EtOAc). Mp 235° dec. [α]_D +4.68 (c, 0.13 in EtOH).

15-O-(2-Methylpropanoyl), hydroiodide: Mp 265-270° dec.

15-O-(2-Methylpropanoyl), picrate: Mp 158-160° dec.

15-O-(2-Methylpropanoyl), 11-Ac: **11-Acetylcardionine**

[123151-95-1]


C₂₆H₃₅NO₆ 457.566

Alkaloid from aerial parts of *Delphi-*

nium gracile (Ranunculaceae). Gum. [α]_D -5.71 (c, 0.14 in CHCl₃).

De la Fuente, G. *et al.*, *J.O.C.*, 1990, **55**, 342 (*isol, ir, pmr, cmr, ms, struct*)

Hetisan-7,11,15,19-tetrol H-193



C₂₀H₂₇NO₄ 345.438

(7α,11α,15β,19R)-form

Ternatidine

[193901-49-4]

7-O-(2-Methylpropanoyl): **Ternatine**†


[193901-34-7]

C₂₄H₃₃NO₅ 415.528

Alkaloid from the aerial parts of *Delphinium ternatum*. Cryst. (Me₂CO). Mp 236-238°.

Narzullaev, A.S. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 1997, **46**, 184-185 (*isol, pmr, cmr*)

Hetisan-1,2,13-triol H-194



C₂₀H₂₇NO₃ 329.438

(1β,2α,13R)-form

Tri-Ac: **Venudelphine**


[154887-96-4]

C₂₆H₃₃NO₆ 455.55

Alkaloid from aerial parts of *Delphinium venulosum* (Ranunculaceae). [α]_D²² 0 (c, 0.1 in CHCl₃).

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 780 (*isol, ir, pmr, cmr, ms, struct*)

Hetisan-1,2,15-triol H-195



C₂₀H₂₇NO₃ 329.438

(1β,2α,15β)-form

2-Benzoyl: **Deacetylhanamisine**. *Hanamiyama base*. 9-Deoxyhypognavine [148245-81-2]
 $C_{27}H_{31}NO_4$ 433.546
 Alkaloid from *Aconitum sanyoense* (Ranunculaceae). Plates (Me₂CO/MeOH). Mp 246-247° (243-244.5°). $[\alpha]_D^{28} +120.6$. $[\alpha]_D^{16} +130$ (c, 0.11 in MeOH).

2-Benzoyl, 1-Ac: **Hanamisine**

[87201-23-8]
 $C_{29}H_{33}NO_5$ 475.583
 Alkaloid from *Aconitum sanyoense* and *Aconitum sanyoense* var. *tonense* (Ranunculaceae). Prisms (Me₂CO). Mp 124-127°. $[\alpha]_D^{25} +122.6$ (c, 1.06 in MeOH).


Ochiai, E. et al., *Yakugaku Zasshi*, 1956, **76**, 1436; *CA*, **51**, 6662d (*Deacetylhanamisine, isol*)

Okamoto, T. et al., *Chem. Pharm. Bull.*, 1983, **31**, 1431 (*Hanamisine, pmr, cmr, ms, abs config, cryst struct*)

Takayama, H. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2927 (*Deacetylhanamisine, struct*)

Hetisan-1,7,13-triol

H-196



$C_{20}H_{27}NO_3$ 329.438


(1β,7α,13S)-form

13-Benzoyl: **Crassicauline B**. *Crassicauline II* [79620-39-6]
 $C_{27}H_{31}NO_4$ 433.546
 Minor alkaloid from the roots of *Aconitum crassicaule* (Ranunculaceae). Powder. Mp 311-315° dec. (sealed tube).

Wang, F.-P. et al., *Planta Med.*, 1981, **42**, 375; 1985, 443 (*isol, uv, ir, pmr, ms, struct*)

Hetisan-1,7,15-triol

H-197



$C_{20}H_{27}NO_3$ 329.438

(1β,7α,15β)-form


7-Hydroxycossonidine. 7-Hydroxydavisine
 Alkaloid from *Consolida oliveriana*. Amorph. solid. $[\alpha]_D^{25} +25$ (c, 0.1 in CHCl₃).

Grandez, M. et al., *J. Nat. Prod.*, 2002, **65**, 513-516 (*isol, pmr, cmr, ms*)

Hetisan-2,6,13-triol, 9CI

H-198

$C_{20}H_{27}NO_3$ 329.438



$C_{20}H_{27}NO_3$ 329.438

(2α,13S)-form

Tatsirine

[132922-83-9]
 Alkaloid from the roots of *Delphinium tatsienense*. Mp 260-263°.

2-Ketone: 6,13-Dihydroxyhetisan-2-one.

Delnuttidine

[163597-05-5]
 $C_{20}H_{25}NO_3$ 327.422
 Alkaloid from *Delphinium nuttallianum*.

13-Ketone: 2,6-Dihydroxyhetisan-13-one.

Panicudine

[178451-93-9]
 $C_{20}H_{25}NO_3$ 327.422
 Alkaloid from *Aconitum paniculatum*. Cryst. Mp 249-250°.

13-Ketone, N-Me: **Orochrine**

[959906-92-4]
 $C_{21}H_{28}NO_3$ 342.457
 Alkaloid from *Aconitum orochryseum*. Needles (as hydroxide). Mp 210-214° (hydroxide). $[\alpha]_D^{23} -65.2$ (c, 0.62 in MeOH) (hydroxide).

13-Ketone, N-Me, 2-Ac: **2-O-Acetylorochrine**

[959853-59-9]
 $C_{23}H_{30}NO_4$ 384.494
 Alkaloid from *Aconitum orochryseum*. Amorph. solid (as hydroxide). Mp 148-150° (hydroxide). $[\alpha]_D^{22} -82.5$ (c, 0.82 in CHCl₃) (hydroxide).

2,13-Diketone: 6-Hydroxyhetisan-2,13-dione. **Paniculadine**

[188904-81-6]
 $C_{20}H_{23}NO_3$ 325.407
 Alkaloid from *Aconitum paniculatum*. Cryst. (Me₂CO). Mp 276-278°. λ_{max} 301 (log ε 5.11) (EtOH).

Zhang, X. et al., *Heterocycles*, 1990, **31**, 1879-1888 (*Tatsirine*)


Bai, Y. et al., *Phytochemistry*, 1994, **37**, 1717-1724 (*Delnuttidine*)

Bessonova, I.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 705-707; 1996, **32**, 561-563 (*Panicudine, Paniculadine*)

Wangchuk, P. et al., *J. Nat. Prod.*, 2007, **70**, 1808-1811 (*Orochrine, 2-Acetylorochrine*)

Hetisan-2,7,11-triol

H-199



(2β,7β,11β)-form

Souline F


[244616-34-0]
 Alkaloid from *Delphinium souliei*. Cryst. (MeOH). Mp > 350. $[\alpha]_D^{24} -1.2$ (c, 0.85 in MeOH).

He, L. et al., *Chin. Chem. Lett.*, 1999, **10**, 395-396 (*isol, pmr, cmr*)

He, L. et al., *Indian J. Chem., Sect. B*, 2001, **40**, 1285-1286 (*isol, pmr, cmr*)

Hetisan-2,9,15-triol

H-200



$C_{20}H_{27}NO_3$ 329.438

(2α,15β)-form

Ryosenaminol. 1-Deoxyhypognavinol

[88142-54-5]
 Alkaloid from *Aconitum ibukiense* (Ranunculaceae). Mp 287-290° dec. $[\alpha]_D^{29} +66.8$ (c, 0.38 in MeOH).

2-Benzoyl: **Ryosenamine**. 1-Deoxyhypognavine

[88142-55-6]
 $C_{27}H_{31}NO_4$ 433.546


Alkaloid from *Aconitum ibukiense* (Ranunculaceae). Also isol. from the processed aconite kako-bushi-matsu. Mp 213-215°. $[\alpha]_D^{12} +96.8$ (c, 0.20 in MeOH).

Sakai, S. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3338 (*isol, ir, uv, pmr, cmr, ms, cryst struct*)

Mori, T. et al., *Heterocycles*, 1989, **29**, 873 (*Ryosenamine*)

Hetisan-2,9,19-triol

H-201



$C_{20}H_{27}NO_3$ 329.438

(2α,19S)-form


2-Ac: **Acsinatine**

[1353-76-0]
 $C_{22}H_{29}NO_4$ 371.475
 Alkaloid from roots of *Aconitum leucostomum*. Cryst. (Et₂O/MeOH). Mp 251-253°. Incorrect C-19 stereochem. given in CA.

Telnov, V.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 346-348 (*Acsinatine*)

Hetisan-2,11,13-triol, 9CI

H-202



C₂₀H₂₇NO₃ 329.438

(2α,11α,13R)-form

Hetisine. *Delatine*

[10089-23-3]

Alkaloid from the roots of *Aconitum heterophyllum* and *Delphinium tatsienense*, the seeds of *Delphinium elatum*, and the above-ground parts of *Delphinium cardinale* (Ranunculaceae). Cryst. (MeOH/Me₂CO). Mp 261-264° (256.5-259°). [α]_D²⁵ +10.9 (c, 1.5 in CHCl₃). [α]_D²⁵ +13.7 (c, 1.02 in EtOH).

Hydrochloride:

Cryst. (H₂O). Mp 274-275°. [α]_D¹⁸ +13.4 (c, 2 in H₂O).

11-Ac: 11-Acetylhetisine

[116107-39-2]

C₂₂H₂₉NO₄ 371.475

Alkaloid from *Delphinium nuttallianum*. Cryst. (Me₂CO/hexane). Mp 264-266°.

13-Ac: 13-Acetylhetisine

[103956-38-3]

C₂₂H₂₉NO₄ 371.475

Alkaloid from *Delphinium macrocentrum* and *Delphinium nuttallianum* (Ranunculaceae). Amorph. solid. Mp 241-243°. Mp refers to a semisynthetic sample.

O¹¹,O¹³-Di-Ac: 11,13-Diacetylhetisine

[105742-72-1]

C₂₄H₃₁NO₅ 413.513

Alkaloid from *Delphinium nuttallianum*. Mp 225-227°. [α]_D +26.1 (CHCl₃).

Tri-Ac:

Amorph.

13-Cinnamoyl: Palmasine

[116085-15-5]

C₂₉H₃₃NO₄ 459.584

Minor alkaloid from roots of *Aconitum palmatum* (Ranunculaceae). Cryst. (Me₂CO). Mp 252-254°.

13-Cinnamoyl, 11-Ac: Palmadine

[116085-14-4]

C₃₁H₃₅NO₅ 501.621

Alkaloid from roots of *Aconitum palmatum* (Ranunculaceae). Cryst. (Me₂CO). Mp 269-271°. [α]_D²⁰ +11.2 (c, 0.2 in CHCl₃).

2-Ketone: Hetisinone. *11,13-Dihydroxyhetisan-2-one*. *Dehydrohetisine*

[4829-55-4]

C₂₀H₂₅NO₃ 327.422

Alkaloid from *Delphinium cardinale*, *Delphinium cardiopetalum* and from the roots of *Delphinium tatsienense* and *Aconitum heterophyllum* (Ranunculaceae). Rhombs (C₆H₆). Mp 273-275° (268-270°). [α]_D +18. A semisynthetic sample showed Mp 195-197° (Gliński);

discrepancy unexplained.

2-Ketone, 13-Ac: 13-Acetylhetisinone

[82209-94-7]

C₂₂H₂₇NO₄ 369.46

Alkaloid from *Delphinium cardiopetalum*.

Jacobs, W.A. *et al.*, *J. Biol. Chem.*, 1942, **143**, 605; 1947, **170**, 189 (*isol, uv*)

Goodson, J. *et al.*, *J.C.S.*, 1943, 139 (*isol*)

Solo, A.J. *et al.*, *J.A.C.S.*, 1959, **81**, 4439;

J.O.C., 1962, **27**, 2702 (*ir, pmr, struct*)

Przybylska, M. *et al.*, *Acta Cryst. B*, 1962, **16**, 871 (*cryst struct*)

Benn, M.H. *et al.*, *Can. J. Chem.*, 1966, **44**, 1 (*isol, uv, ord, ir, pmr, struct*)

Aplin, R.T. *et al.*, *Can. J. Chem.*, 1968, **46**, 2635 (*Hetisinone*)

Pelletier, S.W. *et al.*, *Phytochemistry*, 1968, **7**, 625 (*Hetisinone*)

González, A.G. *et al.*, *An. Quím., Ser. C*, 1981, **77**, 171; *CA*, **97**, 20730u (*Hetisinone*, *Acetylhetisinone*)

Pelletier, S.W. *et al.*, *Heterocycles*, 1983, **20**, 1347 (*isol, occur*)

Varughese, K.I. *et al.*, *J. Nat. Prod.*, 1984, **47**, 470 (*cryst struct*)

Benn, M. *et al.*, *Heterocycles*, 1986, **24**, 1605 (*13-Acetylhetisine*)

Gliński, J.A. *et al.*, *Heterocycles*, 1988, **27**, 185


Jiang, Q. *et al.*, *Tet. Lett.*, 1988, **29**, 1875 (*Palmadine*, *Palmasine*)

Bai, Y. *et al.*, *Heterocycles*, 1990, **31**, 1233-1236 (*11,13-Diacetylhetisine*)

Bai, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1717-1724 (*11-Acetylhetisine*)

Hetisan-2,11,15-triol

H-203



C₂₀H₂₇NO₃ 329.438

(2α,11β,15β)-form

Isohypognavinol. *Tanwusine*†

[34692-47-2]

Alkaloid from *Aconitum tanguticum* (Ranunculaceae).

2-Benzoyl: Isohypognavine

[3368-91-0]

C₂₇H₃₁NO₄ 433.546

Alkaloid from *Aconitum majimai* and *Aconitum japonicum* (Ranunculaceae). Mp 135°.

▶ MK8400000

2-Benzoyl, hydrochloride: Mp 190-192° dec.

2-Benzoyl, 11-Ac: 11-Acetylisoypognavine

[71239-52-6]

C₂₉H₃₃NO₅ 475.583

Alkaloid from *Aconitum japonicum* (Ranunculaceae). Mp 187.5-188.5°.

[α]_D¹⁹ +74.1 (c, 0.8 in CHCl₃).

2-Benzoyl, 11,15-di-Ac: Diacetylisoypognavine

[71239-53-7]

C₃₁H₃₅NO₆ 517.621

Alkaloid from *Aconitum japonicum* (Ranunculaceae). Mp 181-183°. [α]_D¹⁸

+55.3 (c, 0.2 in CHCl₃).

Ochiai, E. *et al.*, *Yakugaku Zasshi*, 1956, **76**, 550; *CA*, **50**, 13372f (*Isohypognavine, isol*)

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1124-1128 (*Isohypognavine, struct*)


Sakai, S. *et al.*, *Yakugaku Zasshi*, 1979, **99**, 647-656; *CA*, **91**, 105185w (*11-Acetylisoypognavine*, *Diacetylisoypognavine*)

Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4576; 4579 (*ir, pmr, ms, cryst struct, abs config*)

Chen, D. *et al.*, *Zhongcaoyao*, 1985, **16**, 338; *CA*, **104**, 95320 (*Tanwusine*)

Hetisan-3,15,19-triol

H-204



C₂₀H₂₇NO₃ 329.438

(3α,15β,19S)-form

15-Ac: Andersosine

[159982-64-6]


C₂₂H₂₉NO₄ 371.475

Alkaloid from aerial parts of *Delphinium andersonii* (Ranunculaceae). Plates (MeOH). Mp 310°.

Joshi, B.S. *et al.*, *Tetrahedron*, 1994, **50**, 12283 (*isol, ir, cd, pmr, cmr, ms, struct*)

Hetisan-6,11,15-triol

H-205



C₂₀H₂₇NO₃ 329.438

(11β,15β)-form

Pseudokobusine

[27901-01-5]

Alkaloid from *Aconitum yesoense* and *Aconitum ludiscusculum* (Ranunculaceae). Mp 271-272° dec.

▶ MK8200000

15-O-(2S-Methylbutanoyl): Yesodine

[131652-36-3]

C₂₅H₃₅NO₄ 413.556

Alkaloid from *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Amorph. powder. [α]_D²² -9.4 (c, 0.34 in CHCl₃).

15-Benzoyl: 15-Benzoylpseudokobusine

[112574-95-5]

C₂₇H₃₁NO₄ 433.546

Alkaloid from rhizomes of *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Amorph. [α]_D -6.9 (c, 0.13 in EtOH).

15-O-(3,4-Dimethoxybenzoyl): 15-(3,4-

Dimethoxybenzoyl)pseudokobusine. 15-Veratrolypseudokobusine

[112561-71-4]

C₂₉H₃₅NO₆ 493.599

Alkaloid from the rhizomes of *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Amorph. [α]_D -6.7 (c, 0.18 in EtOH).

Natsume, M. *et al.*, *Chem. Pharm. Bull.*, 1962, **10**, 879 (*Pseudokobusine*)


Bando, H. *et al.*, *Heterocycles*, 1987, **26**, 2623 (*15-Benzoylpseudokobusine, 15-Veratrolypseudokobusine*)

Wada, K. *et al.*, *Heterocycles*, 1990, **31**, 1081 (*Yesodine*)

Merikli, A.H. *et al.*, *Heterocycles*, 2000, **53**, 1987-1996 (*Pseudokobusine*)

Hetisan-6,13,15-triol

H-206



C₂₀H₂₇NO₃ 329.438

(13S,15β)-form

Acorientine


[174232-21-4]

Alkaloid from aerial parts of *Aconitum orientale*. [α]_D +13.5 (c, 0.1 in CHCl₃).

Ulubelen, A. *et al.*, *Phytochemistry*, 1996, **41**, 957-961 (*isol, ir, pmr, cmr, ms*)

Hetisan-7,11,13-triol

H-207



C₂₀H₂₇NO₃ 329.438

(7β,11α,13S)-form

Delfissinol


[152606-55-8]

Alkaloid from aerial parts of *Delphinium fissum* ssp. *anatolicum* (Ranunculaceae). Amorph. [α]_D -39.1 (c, 0.2 in MeOH). Struct. incorrectly drawn in ref.

Ulubelen, A. *et al.*, *Phytochemistry*, 1993, **34**, 1165-1167 (*isol, ir, pmr, cmr, ms*)

Hetisan-7,11,15-triol

H-208



(7α,11α,15α)-form

C₂₀H₂₇NO₃ 329.438

(7α,11α,15α)-form

7-Benzoyl, 11-Ac: Tiantaishandine

[959843-33-5]

C₂₉H₃₃NO₅ 475.583

Alkaloid from the roots of *Delphinium tiantaishanense*. Amorph. powder. Mp 248-249°. [α]_D²⁰ +35.6 (c, 0.85 in CHCl₃).

(7α,11α,15β)-form

11-Ac: Delnutrine

[163564-88-3]

C₂₂H₂₉NO₄ 371.475


Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae).

Bai, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1717-1724 (*Delnutrine*)

Li, J. *et al.*, *Molecules*, 2007, **12**, 353-360 (*Tiantaishandine*)

Hetisan-11,15,19-triol

H-209



C₂₀H₂₇NO₃ 329.438

(11β,15β,19S)-form

Talatisine


[80248-73-3]

Alkaloid from *Aconitum talassicum* (Ranunculaceae). Cryst. (EtOH). Mp 246-246.5°. [α]_D +38 (EtOH).

Karimov, Z. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 335; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 259 (*Talatisine*)

Hetisan-11,15,20-triol

H-210



C₂₀H₂₇NO₃ 329.438

(11β,15β)-form

Orgetine

Alkaloid from *Aconitum orientale*. Cryst. (Me₂CO). Mp 280-282°. [α]_D²² +40 (c, 0.36 in EtOH).


Beshitaishvili, L.V. *et al.*, *Khim. Prir. Soedin.*, 1992, **28**, 240; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 206-208 (*Orgetine*)

2,2',3,4',5,5'-Hexabromo-1,3'-bi-1H-indole

H-211

Rivularin A

[81387-87-3]



C₁₆H₆Br₆N₂ 705.661

Isol. from the marine blue-green alga *Rivularia firma*. Prisms (MeCN). Mp 263-264°. [α]_D²⁰ +18.7 (c, 1 in MeCN). Exhibits opt. activity owing to restricted rotn. λ _{max} 226 (ε 63000); 287 (ε 17000) (MeCN).

Norton, R.S. *et al.*, *J.A.C.S.*, 1982, **104**, 3628-3635 (*isol, uv, ir, pmr, cmr*)


Maehr, H. *et al.*, *J.O.C.*, 1984, **49**, 1549-1553 (*nomenclature*)

3,3',4,4',5,5'-Hexabromo-2,2'-bi-1H-pyrrole

H-212

Bistribromopyrrole

[54705-15-6]



C₈H₂Br₆N₂ 605.541

Pyrrole antibiotic. Isol. from marine *Chromobacterium* sp. Active against *Chromobacterium* sp. Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O. λ _{max} 258 (MeOH) (Berdy). λ _{max} 272 (MeOH/NaOH) (Berdy).

N,N'-Di-Me: 3,3',4,4',5,5'-Hexabromo-1,1'-dimethyl-2,2'-bi-1H-pyrrole [253798-63-9]

C₁₀H₆Br₆N₂ 633.595

Isol. from the eggs of various seabirds. Mp 247-248°.

Andersen, R.J. *et al.*, *Mar. Biol. (Berlin)*, 1974, **27**, 281 (*isol*)

Gribble, G.W. *et al.*, *Chem. Comm.*, 1999, 2195-2196 (*synth, pmr, ms*)

1,1,1,15,15,15-Hexachloro-3,12-pentadecadiyn-8-amine

H-213

8-Amino-1,1,1,15,15,15-hexachloro-3,12-pentadecadiyne

Cl₃CCH₂C≡

C(CH₂)₃CH(NH₂)(CH₂)₃C≡

CCH₂CCl₃

C₁₅H₁₉Cl₆N 426.038

(+)-form

N-Ac: 8-Acetamido-1,1,1,15,15,15-hexachloro-3,12-pentadecadiyne [345261-10-1]

C₁₇H₂₁Cl₆NO 468.075

Isol. from the cyanobacterium *Microcoleus lyngbyaceus*. Oil.

Dechloro, N-Ac: 8-Acetamido-1,1,1,15,15-pentachloro-3,12-pentadecadiyne
[345261-09-8]
C₁₇H₂₂Cl₅NO 433.63
Isol. from the cyanobacterium *Microcoleus lyngbyaceus*. Pale yellow oil. [α]_D +36 (c, 0.08 in CHCl₃).

1,15-Didechloro, 3,3,4,4-tetrahydro, N-Ac: 8-Acetamido-1,1,15,15-pentachloro-3-pentadecyne
[345261-19-0]
C₁₇H₂₆Cl₅NO 437.662
Isol. from the cyanobacterium *Microcoleus lyngbyaceus*. Oil. [α]_D +30 (c, 0.07 in CHCl₃).

1,15-Didechloro, 3,3,4,4-tetrahydro, N-Ac: 8-Acetamido-1,1,15,15-tetrachloro-3-pentadecyne
[345261-11-2]
C₁₇H₂₇Cl₄NO 403.217
Isol. from the cyanobacterium *Microcoleus lyngbyaceus*. Oil. [α]_D +24 (0.08 in CHCl₃).

Orsini, M.A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 572-577 (isol, ir, pmr, cmr)

Hexacosanoic acid, 9CI H-214

Cerinic acid. Cerotic acid. Ceric acid. Ceratimic acid. EB 82B
[506-46-7]
H₃C(CH₂)₂₄COOH
C₂₆H₅₂O₂ 396.696
Constit. of various oils, waxes and bacterial lipids. Mp 88-89°.

Me ester: [5802-82-4]
C₂₇H₅₄O₂ 410.722
Mp 63-68°. Bp₁₅ 286° Bp_{3,5} 248°.

Et ester: [29030-81-7]
C₂₈H₅₆O₂ 424.749
Mp 59.5-59.8°.

Nonyl ester: Nonyl hexacosanoate
C₃₅H₇₀O₂ 522.937
Constit. of *Artabotrys odoratissimus*. Cryst. (C₆H₆). Mp 83°.

Amide: [22715-86-2]
C₂₆H₅₃NO 395.711
Mp 105-107°.

2-Methylpropylamide: N-(2-Methylpropyl)hexacosanamide, 9CI. Hexacosanoic acid isobutylamide. Longamide†
[119736-87-7]
C₃₀H₆₁NO 451.818
Minor constit. of fruits of *Piper longum* (long pepper) (Piperaceae). Amorph. powder. Mp 72°.

Levene, P.A. *et al.*, *J. Biol. Chem.*, 1924, **59**, 905 (synth)

Helz, G.E. *et al.*, *J. Biol. Chem.*, 1936, **116**, 203

Bernhard, K. *et al.*, *Helv. Chim. Acta*, 1948, **31**, 977

Asselineau, J. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1953, **237**, 1804

Tember, G.A. *et al.*, *CA*, 1974, **80**, 47410a (synth)

Watanabe, S. *et al.*, *Z. Naturforsch., C*, 1975, **30**, 825 (ms)

Koul, S.K. *et al.*, *Phytochemistry*, 1988, **27**, 3523 (Longamide)

Mehta, B.K. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 1304-1306 (Nonyl hexacosanoate)

2,9-Hexadecadiene-12,14-diyneic acid, 9CI H-215

H₃CC≡CC≡CCH₂CH=CH(CH₂)₅CH=CHCOOH
C₁₆H₂₀O₂ 244.333

(2E,9Z)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2,9-hexadecadiene-12,14-diyneic acid isobutylamide
[120727-32-4]
Isol. from the roots of *Echinacea angustifolia*. Oil. λ_{max} 199, 211sh nm.
Bauer, R. *et al.*, *Phytochemistry*, 1989, **28**, 505 (isol, struct, ir, uv, pmr, ms)

2,4-Hexadecadienoic acid, 9CI H-216

[25377-52-0]
H₃C(CH₂)₁₀CH=CHCH=CHCOOH
C₁₆H₂₈O₂ 252.396

(all-E)-form [59404-48-7]

2-Methylpropylamide: N-Isobutyl-2,4-hexadecadienamide
C₂₀H₃₇NO 307.518
Constit. of the fruits of *Piper guineense* and *Evodia huiphephensis* (Piperaceae, Rutaceae). Noncryst. Mp 78.5-84° (refers to synthetic material).

(2E,4Z)-form

Me ester: [54977-89-8]
C₁₇H₃₀O₂ 266.423
Used as aroma compd. in the food industry.

Et ester:
C₁₈H₃₂O₂ 280.45
Aroma compd.

(2ξ,4ξ)-form

Piperidide: 2,4-Hexadecadienoic acid piperidide
[486447-33-0]
C₂₁H₃₇NO 319.529
Constit. of *Piper guineense*.

[28933-88-2]

Okogun, J.I. *et al.*, *J.C.S. Perkin 1*, 1974, 2195 (pmr, ms)

Swiss Pat., 1974, 557 645; *CA*, **82**, 138023 (synth)

Vig, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 1358 (synth)

Reisch, J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 862 (isol, uv, ir, pmr, ms, struct, isobutylamide)

Strunz, G.M. *et al.*, *Can. J. Chem.*, 1996, **74**, 419 (synth, isobutylamide)

Adesina, S.K. *et al.*, *Pharmazie*, 2003, **58**, 425 (piperidide)

2,7-Hexadecadienoic acid H-217

[25377-52-0]
H₃C(CH₂)₇CH=CH(CH₂)₃CH=CHCOOH
C₁₆H₂₈O₂ 252.396

(2E,7Z)-form

Pyrrolide: 2,7-Hexadecadienoic acid pyrrolide. 1-(1-Oxo-2,7-hexadienyl)-1H-pyrrole, 9CI
[111509-31-0]
C₂₀H₃₁NO 301.471

Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

[28933-88-2]

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (isol, uv, ir, pmr, ms, struct)

2,7-Hexadecadien-10-ynoic acid H-218

H₃C(CH₂)₄C≡CCH₂CH=CH(CH₂)₃CH=CHCOOH
C₁₆H₂₄O₂ 248.364

(2E,7Z)-form

Pyrrolide: 2,7-Hexadeca-10-ynoic acid pyrrolide. 1-(1-Oxo-2,7-hexadecadien-10-ynyl)-1H-pyrrole, 9CI
[111509-28-5]
C₂₀H₂₇NO 297.439
Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Pyrrolidide: 2,7-Hexadeca-10-ynoic acid pyrrolidide. 1-(1-Oxo-2,7-hexadecadien-10-ynyl)pyrrolidine, 9CI
[111509-34-3]
C₂₀H₃₁NO 301.471
From underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (isol, uv, ir, pmr, cmr, ms, struct)

6,8-Hexadecadien-10-ynoic acid H-219

H₃C(CH₂)₄C≡CCH=CHCH=CH(CH₂)₄COOH
C₁₆H₂₄O₂ 248.364

(6E,8E)-form

Pyrrolidide: 6,8-Hexadecadien-10-ynoic acid pyrrolidide. 1-(1-Oxo-6,8-hexadecadien-10-ynyl)pyrrolidine, 9CI
[111509-35-4]
C₂₀H₃₁NO 301.471
Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (isol, uv, ir, pmr, ms, struct)

2,4,6,8,10,12,14-Hexadecaheptaenedioic acid H-220

HOOCCH=CHCH=CHCH=CHCH=CHCH=CHCOOH
C₁₆H₁₆O₄ 272.3

(all-E)-form

Red tablets. Mp 300°. Probable config.

Di-Me ester:

C₁₈H₂₀O₄ 300.354
Orange cryst. (AcOH). Mp 236°.

Di-Et ester: [5941-46-8]

C₂₀H₂₄O₄ 328.407
Orange cryst. (AcOH). Mp 217°.

1-L-Aspartic acid, 16-L-isoleucine diamide: Boletocrocin A
[220376-33-0]

C₂₆H₃₂N₂O₈ 500.547

Isol. from the fruit bodies of *Boletus laetissimus*. Orange solid. Mp 268°

(dec.). $[\alpha]_D^{25} +25$ (c. 2.1 in MeOH). λ_{\max} 246 (log ϵ 4); 312 (log ϵ 3.83); 389 (log ϵ 4.65); 410 (log ϵ 4.81); 434 (log ϵ 4.77) (MeOH).

1-L-Asparagine, 16-L-isoleucine diamide: Boletocrocin B

[220376-47-6]
C₂₆H₃₃N₃O₇ 499.563
Isol. from the fruit bodies of *Boletus aetissimus*. Orange solid. Mp 260° (dec.). $[\alpha]_D^{25} +3.4$ (c. 1.8 in MeOH). λ_{\max} 245 (log ϵ 3.92); 310 (log ϵ 3.69); 389 (log ϵ 4.51); 409 (log ϵ 4.68); 433 (log ϵ 4.65) (MeOH).

1-L-Aspartic acid, 16-L-valine diamide: Boletocrocin E

[220379-17-9]
C₂₅H₃₀N₂O₈ 486.521
Isol. from the fruit bodies of *Boletus laetissimus*.

1-L-Glutamic acid, 16-L-isoleucine diamide: Boletocrocin F

[220379-21-5]
C₂₇H₃₄N₂O₈ 514.574
Isol. from the fruit bodies of *Boletus laetissimus*.

1-L-Glutamine, 16-L-isoleucine diamide: Boletocrocin G

[220379-25-9]
C₂₇H₃₅N₃O₇ 513.589
Isol. from the fruit bodies of *Boletus laetissimus*.

[1441-62-9 (di-Et ester, ns), 77186-01-7 (di-Me ester, ns)]

Kuhn, R. *et al.*, *Ber.*, 1937, **70**, 1318-1330 (synth, di-Me ester, di-Et ester)

Kahner, L. *et al.*, *Phytochemistry*, 1998, **49**, 1693-1697 (*Boletocrocins*)

Hexadecanoic acid **H-221**

Palmitic acid, USAN. Aethalic acid. FEMA 2832

[57-10-3]
H₃C(CH₂)₁₄COOH

C₁₆H₃₂O₂ 256.428

Occurs in the form of esters (glycerides) in oils and fats of vegetable and animal origin, e.g. present in lipids of *Physalis physalis* (Portuguese-man-of-war). Usually obt. from palm oil. Widely distributed in plants. Used in detn. of water hardness. Active ingredient of LevovistTM, used in echo enhancement in sonographic Doppler B-mode imaging. Ultrasound contrast medium. Cryst. Mp 63-64°. Bp 390° Bp₁₀₀ 268.5° Bp₁₅ 215°.

► Skin irritant. LD₅₀ (mus, ivn) 57 mg/kg. RT4550000

Amide: Hexadecanamide, 9CI. Palmitamide, 8CI

[629-54-9]
C₁₆H₃₃NO 255.443

Isol. from seeds of Mexican apple (*Casimiroa edulis*). Cryst. (EtOH). Mp 106-107°. Bp₁₂ 235-236°.

Benzylamide: N-Benzylhexadecanamide. Macamide B

[74058-71-2]
C₂₃H₃₉NO 345.567

Alkaloid from the tubers of *Lepidium meyenii* (maca). Amorph. solid. λ_{\max} 208 (log ϵ 4.03) (MeOH).

3-Methoxybenzylamide: N-(3-Methoxybenzyl)hexadecanamide

C₂₄H₄₁NO₂ 375.593
Alkaloid from the tubers of *Lepidium meyenii* (maca). Powder. λ_{\max} 216 (log ϵ 3.94); 274 (log ϵ 3.41) (MeOH).

[4991-47-3, 408-35-5, 542-42-7, 2624-31-9]


Kincl, F.A. *et al.*, *J.C.S.*, 1956, 4163 (*isol, amide*)

Muhammad, I. *et al.*, *Phytochemistry*, 2002, **59**, 105-110 (*Macamide B*)

Zhao, J.-P. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 690-693 (*3-methoxybenzylamide*)

3,5,10,13-Hexadecatetraen-2-amine **H-222**

2-Amino-3,5,10,13-hexadecatetraene



C₁₆H₂₇N 233.396

(2R,3E,5E,10Z,13Z)-form [356073-88-6]

Isol. from a *Pseudodistoma* sp. Oil. $[\alpha]_D -6$ (c. 0.05 in MeOH). λ_{\max} 229 (log ϵ 3.38) (MeOH).

Rashid, M.A. *et al.*, *Tetrahedron*, 2001, **57**, 5751-5755 (*Pseudodistoma constit, isol, ir, pmr, cmr, uv*)

2,6,8,12-Hexadecatetraen-10-ynoic acid **H-223**

H₃CCH₂CH₂CH=CHC≡CCH=CHCH=CHCH₂CH₂CH=CHCOOH

C₁₆H₂₀O₂ 244.333

(2E,6E,8E,12Z)-form

2-Methylpropylamide: 2,6,8,12-Hexadecatetraen-10-ynoic acid isobutylamide. N-(2-Methylpropyl)-2,6,8,12-hexadecatetraenamide, 9CI

[81427-12-5]
C₂₀H₂₉NO 299.455

Alkaloid from *Achillea tomentosa* (Asteraceae). Gum.

2,3-Didehydropyrrolidide: 2,6,8,12-Hexadecatetraen-10-ynoic acid dehydropyrrolidide, 2,3-Dihydro-1-(1-oxo-4,6,8,12-hexadecatetraen-10-ynyl)-1H-pyrrole, 9CI

[81427-13-6]
C₂₀H₂₅NO 295.424

Alkaloid from *Achillea tomentosa* (Asteraceae). Gum.

Greger, H. *et al.*, *Phytochemistry*, 1981, **20**, 2579 (*isol, uv, ir, ms, struct*)

2,6,8-Hexadecatrien-10-ynoic acid **H-224**

H₃C(CH₂)₄C≡CCH=CHCH=CHCH₂CH₂CH=CHCOOH

C₁₆H₂₂O₂ 246.349

(2E,6E,8E)-form

Pyrrolide: 2,6,8-Hexadecatrien-10-ynoic acid pyrrolide

C₂₀H₂₅NO 295.424

Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Mp 34°.

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (*isol, uv, ir, pmr, ms, struct*)

7-Hexadecen-10-ynoic acid **H-225**

H₃C(CH₂)₄C≡CCH₂CH=CH(CH₂)₅COOH

C₁₆H₂₆O₂ 250.38

(Z)-form


Pyrrolide: 7-Hexadecen-10-ynoic acid pyrrolide, 1-(1-Oxo-7-hexadecen-10-ynyl)-1H-pyrrole, 9CI

[111509-30-9]
C₂₀H₂₉NO 299.455

Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (*isol, uv, ir, pmr, ms, struct*)

6,7,8,9,10,14-Hexadehydro-4,5-epoxy-3,6-dimethoxy-17-methylthebinan **H-226**



C₁₉H₁₉NO₃ 309.364

Unusual type of morphine alkaloid. Alkaloid from capsules of *Papaver bracteatum* cv. "Arya I" (Papaveraceae). Mp 174-175°.


Methodide:

Needles. Mp 172-173°.

Theuns, H.G. *et al.*, *J.C.S. Perkin 1*, 1984, 1701 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Hexadellin B **H-227**

[121135-01-1]



C₂₁H₂₃Br₄N₃O₅ 717.046

Alkaloid from the deep water sponge *Hexadella* sp.

*N*²⁰-Me: *Purealidin S*

C₂₂H₂₅Br₄N₃O₅ 731.073

Alkaloid from the Fijian sponge *Druinella* sp. Cytotoxic. Oil. λ_{\max} 280 (log ϵ 3.28) (MeOH).

N²⁰,N²⁰-Di-Me: Purealidin Q

[167394-82-3]

C₂₃H₂₇Br₄N₃O₅ 745.099

Alkaloid from the Okinawan sponge *Psammaphysilla porea*. Cytotoxic agent. Enzyme inhibitor. Oil (as trifluoroacetate). Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D¹⁹ +9.1 (c, 0.39 in MeOH). λ_{max} 277 (ε 1700); 284 (ε 1380) (MeOH) (Berdy).

N²⁰-(13-Methyltetradecanoyl): Purpurealidin C

[799246-88-1]

C₃₆H₅₁Br₄N₃O₆ 941.431

Isol. from *Psammaphysilla purpurea*. Oil. [α]_D²⁸ +158.5 (c, 0.2 in CHCl₃). λ_{max} 218 (ε 2500); 282 (ε 10000) (MeOH).

N²⁰-Hexadecanoyl: Purpurealidin D

[799246-89-2]

C₃₇H₅₃Br₄N₃O₆ 955.458

Isol. from *Psammaphysilla purpurea*. Oil. λ_{max} 218 (ε 2500); 282 (ε 10000) (MeOH).

3'ξ,4'ξ-Dihydro, N⁹,N²⁰,N²⁰-tri-Me:**Purpurealidin A**

[799246-86-9]

C₂₄H₃₁Br₄N₃O₅ 761.142

Isol. from the sponge *Psammaphysilla purpurea*. Oil. [α]_D²⁸ +9.5 (c, 0.2 in MeOH). λ_{max} 277 (ε 1700); 284 (ε 1400) (MeOH).

Morris, S.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 677 (isol, ir, pmr, cmr, ms, struct)


Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403 (Purealidin Q)

Tabudravu, J.N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1798-1801 (Purealidin S)

Tilvi, S. *et al.*, *Tetrahedron*, 2004, **60**, 10207-10215 (Purpurealidins A, C, D, Purealidin Q)

Hexadellin C**H-228**

[245436-91-3]


C₂₅H₂₉Br₄N₃O₇ 803.136

Zwitterionic. Isol. from the sponge *Aiolochroia crassa*. Amorph. solid. [α]_D +102 (c, 0.07 in MeOH). λ_{max} 207 (ε 8300); 284 (ε 1200) (EtOH).

Gao, H. *et al.*, *Tetrahedron*, 1999, **55**, 9717-9726

3,4,5,6,7,8-Hexahydro-1,7-benzodiazecin-2(1H)-one**H-229**

[98155-99-8]

C₁₂H₁₆N₂O 204.271


Minor alkaloid from *Mackinlaya macro-*

sciadia (Araliaceae). Cryst. (CCl₄). Mp 92.5-93.5°. Artifact. Formed in low yield from dec. of 6,8,9,11-Tetrahydro-7H-pyrido[2,1-*b*]quinazoline, T-227, the major base, on storage, and is prepd. by hydrol. of the major base in H₂O.

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1985, **38**, 1007-1008 (occur, synth, pmr, struct)


1,4,5,6,7,8-Hexahydro-1,3-diazocine-2,4-dicarboxylic acid, 9CI**H-230**

[203579-33-3]

C₈H₁₂N₂O₄ 200.194

Zwitterionic with charge delocalisation over the two N's. Isol. from the sponge *Axinyssa terpnis*. Needles (MeOH). Mp 210-212°. [α]_D -66.3 (c, 0.9 in MeOH).

Li, C.-J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 387-389 (isol, ir, pmr, cmr, ms)

Hexahydro-2-hydroxy-2-methyl-1H-pyrrolizine-1-carboxylic acid, 9CI**H-231**

(1R,2S,7aS)-form

C₉H₁₅NO₃ 185.222**(1R,2S,7aS)-form Neotussilaginic acid**

[147634-91-1]

Genuine alkaloid from *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*.

Me ester: Neotussilagine. Isotussilaginine [147730-89-0]

C₁₀H₁₇NO₃ 199.249

Alkaloid artifact from *Tussilago farfara* (coltsfoot), *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*, resulting from the use of MeOH during isol.

(1R,2R,7aS)-form**Neoisotussilaginic acid**

[147634-92-2]

Genuine alkaloid from *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*.

Me ester: Neoisotussilagine. Tussilaginine [147730-90-3]

C₁₀H₁₇NO₃ 199.249

Alkaloid artifact from *Tussilago farfara* (coltsfoot), *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*, resulting from the use of MeOH during isol.

(1S,2S,7aS)-form**Tussilagic acid**

[147634-89-7]

Genuine alkaloid from *Echinacea purpurea*, *Echinacea angustifolia*, *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*.

Me ester: Tussilagine

[80151-77-5]

Alkaloid artifact from *Echinacea purpurea*, *Echinacea angustifolia*, *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*, resulting from the use of MeOH during isol. Also isol. from *Tussilago farfara* (coltsfoot). Mp 193° (dec.). [α]_D²⁵ -2.7 (c, 0.15 in EtOH).

(1S,2R,7aS)-form**Isotussilaginic acid**

[147634-90-0]

Genuine alkaloid from *Echinacea purpurea*, *Echinacea angustifolia*, *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*.

Me ester: Isotussilagine

[91108-32-6]

C₁₀H₁₇NO₃ 199.249

Alkaloid artifact from *Tussilago farfara* (coltsfoot), *Echinacea angustifolia*, *Arnica montana*, *Arnica chamissonis* ssp. *foliosa*, *Arnica amplexicaulis* and *Arnica sachalinensis*, resulting from the use of MeOH during isol. [α]_D²⁵ +129 (c, 0.17 in CHCl₃).

Röder, E. *et al.*, *Planta Med.*, 1981, **43**, 99 (ir, pmr, cmr, ms, struct)

Weidenfeld, H. *et al.*, *Arch. Pharm.*

(Weinheim, Ger.), 1983, **316**, 367 (cryst struct, abs config)


Röder, E. *et al.*, *Arch. Pharm.* (Weinheim, Ger.), 1984, **317**, 403 (synth)

Passreiter, C.M. *et al.*, *Phytochemistry*, 1992, **31**, 4135 (stereoisomers)

Passreiter, C.M. *et al.*, *Planta Med.*, 1992, **58**, 556-557 (isol)


Ma, D. *et al.*, *Tet. Lett.*, 1998, **39**, 9067-9068 (synth, pmr)

Ma, D. *et al.*, *J.C.S. Perkin 1*, 1999, 1703-1707 (synth)

1,2,3,4,7,8-Hexahydro-6-(4-hydroxyphenyl)-8-(2,3,4,5-tetrahydroxypentyl)-2,4,7-pteridinetrione**H-232**C₁₇H₁₈N₄O₈ 406.351**D-ribo-form**

6-(4-Hydroxyphenyl)-8-D-ribosyl-2,4,7-(1H,3H,8H)-pteridinetrione, 9CI. 6-(p-Hydroxyphenyl)-8-(1-D-ribityl)-2,4,7-trioxohexahydropteridine

[32507-81-6]

Isol. from *Pseudomonas ovalis*. Yellow cryst. (H₂O). Mp 272-273° dec.Suzuki, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1971, **44**, 1869 (*synth*)**Hexahydro-7-hydroxy-1H-pyrrolizine-1-methanol, 9CI***1-Hydroxy-7-hydroxymethylpyrrolizidine*(1*R*,7*R*,7*aR*)-formC₈H₁₅NO₂ 157.212**(1*R*,7*R*,7*aR*)-form***(1α,7β,7αα)-form. Turneforicidine. Racemonecine*

[21850-67-9]

Hydrol. prod. from Turneforicine and Retusine, Needles (Me₂CO). Mp 118-120°. [α]_D²⁶ -13.9 (c, 0.36 in MeOH). Racemonecine (Mp 138-140°, [α]_D -11.4 (CHCl₃)) the necine base from Racemocine and Racemoneine, was assigned the (1*S*,7*S*,7*aS*)-stereochem. but apparently with no supporting evidence. On grounds of opt. rotn. it appears to be identical with Turneforicidine. Racemocine [α]_D -16 (CHCl₃) appears to be the enantiomer of Farfugine which supports this assignment.

*Hydrochloride:*S. hygroscopic long needles (EtOH/Et₂O). Mp 120° (116°).*1'-Angeloyl: 9-Angeloylturneforicidine*

[1007096-45-8]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Pittocaulon bombycop-hole* and *Pittocaulon hintonii*.*7-Angeloyl: 7-Angeloylturneforicidine*

[135637-69-3]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Pittocaulon* spp. and *Senecio integrifolius* var. *fauriri*. [α]_D +5 (c, 1 in EtOH).*Angeloyl: Turneforicine. Angeloylturneforicidine*C₁₃H₂₁NO₃ 239.314Alkaloid from *Tournefortia sibirica* (Boraginaceae). Oil; cryst. (as hydrochloride). Mp 169-170° (hydrochloride). [α]_D -58. Posn. of esterification not. detd.*1'-Tigloyl: Racemocine. Racemozine*

[139490-40-7]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Senecio racemosus*. Gum. [α]_D -16 (c, 0.25 in CHCl₃).*1'-O-(2-Ethoxycarbonyl-3-methyl-2-butenoyl): Racemoneine*C₁₆H₂₅NO₅ 311.377Alkaloid from *Senecio racemosus* (Asteraceae). Gum. [α]_D²⁰ -24.3 (c, 0.2 in CHCl₃). Struct. revised in 1993.**(1*S*,7*R*,7*aR*)-form***(1β,7β,7αα)-form. Platynecine. Mikane-cine*

[520-62-7]

Necine component of various pyrrolizidine alkaloids, e.g. Platyphylline, P-516, Sarracine, S-90. Mp 148-148.5°. [α]_D -56.8 (CHCl₃). p*K*_a 10.2 (24°).*Picrate:* Mp 184°.*1'-Angeloyl: 9-Angeloylplatynecine*

[94132-25-9]

Alkaloid from stems and leaves of *Castilleja rhexifolia* aff. *miniata* (Scrophulariaceae).*1'-Angeloyl, N-oxide: 9-Angeloylplatynecine N-oxide*C₁₃H₂₁NO₄ 255.313Isol. from the stems and leaves of *Castilleja rhexifolia* aff. *miniata* (Scrophulariaceae).*7-Angeloyl: 7-Angeloylplatynecine*

[94054-32-7]

Alkaloid from the stems and leaves of *Castilleja rhexifolia* aff. *miniata* (Scrophulariaceae). [α]_D^{23.5} -98 (c, 0.004 in EtOH).*Diangeloyl: Diangeloylplatynecine*

[47928-73-4]

C₁₈H₂₇NO₄ 321.416Alkaloid from the roots of *Senecio macedonicus*. Oil.*1'-O-(3-Methyl-2-butenoyl): Fuchsisenecionine. O-Senecioylplatynecine*C₁₃H₂₁NO₃ 239.314Alkaloid from *Senecio fuchsii*. [α]_D²² -120.2.*1'-O-(2,3-Dihydroxy-2-methylbutanoyl):*see Ipanguline D₃, I-166*1'-O-[2*S*-Hydroxy-2-(1*S*-hydroxyethyl)-4-methylpentanoyl], N-oxide:*C₁₆H₂₉NO₆ 331.408Alkaloid from the leaves of *Anchusa strigosa*. Orange oil. [α]_D²⁵ +3 (c, 0.1 in MeOH). λ_{max} 235 (sh); 282 (MeOH).*1'-O-(Phenylacetyl): Ipanguline D₇*

[210698-85-4]

C₁₆H₂₁NO₃ 275.347Alkaloid from *Ipomoea hederifolia*.*7-O-(Phenylacetyl): Ipanguline D₆*

[63503-40-2]

C₁₆H₂₁NO₃ 275.347Alkaloid from *Ipomoea hederifolia*.**(1*S*,7*S*,7*aR*)-form***Dihydroheliotridine*

[21824-59-9]

Mp 76-77°. [α]_D -34 (EtOH).*1'-Angeloyl: 9-Angeloyldihydroheliotridine*Alkaloid from *Pittocaulon praecox*.[α]_D -8.2 (MeOH).*7-Angeloyl: 7-Angeloyldihydroheliotridine*Alkaloid from *Pittocaulon praecox*.[α]_D +7.5 (MeOH).**(1*S*,7*R*,7*aS*)-form***Hastancine*

[480-84-2]

Necine base from Hastacine, H-69. Mp

113-114°. [α]_D -9.1 (MeOH).*1'-Angeloyl: 9-Angeloylhastancine*

[173010-19-0]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Senecio aquaticus* ssp. *barbareifolius* and *Senecio chrysocoma*. Yellow oil.**(1*R*,7*S*,7*aR*)-form**Synthetic. Prisms (Me₂CO). Mp 113-114°. [α]_D¹⁸ +8.5 (c, 2.2 in EtOH).**(1*S*,7*S*,7*aS*)-form***(+)-Turneforicidine*

Oil. Rare enantiomer. Isol. for the first time in 1983 as component of Farfugine.


1'-Angeloyl: Farfugine

[86194-08-3]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Farfugium japonicum* (Asteraceae). Oil. Mp 156.5-157.5° (as picrate). [α]_D²³ +23 (c, 0.54 in EtOH).Orechov, A. *et al.*, *Ber.*, 1935, **68**, 1886 (*Platynecine*)Kononov, V.S. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1945, **15**, 328; *CA*, **40**, 3760 (*Hastancine*)Men'shikov, G.P. *et al.*, *J. Gen. Chem.**USSR (Engl. Transl.)*, 1952, **22**, 1465-1467; *CA*, **47**, 7512 (*Turneforicine, Turneforicidine*)Aasen, A.J. *et al.*, *Aust. J. Chem.*, 1969, **22**,2657 (*struct, synth*)Aasen, A.J. *et al.*, *J.O.C.*, 1969, **34**, 4137*(struct, synth, ms, pmr)*Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*,1972, **37**, 3918 (*cd*)Viscontini, M. *et al.*, *Helv. Chim. Acta*, 1972,**55**, 73 (*synth, ms, pmr*)Gupta, V.P. *et al.*, *Curr. Sci.*, 1975, **44**, 451 (*ir*)Danishefsky, S. *et al.*, *J.A.C.S.*, 1977, **99**, 7711*(synth)*Röder, E. *et al.*, *Phytochemistry*, 1977, **16**, 1462*(Fuchsisenecionine)*Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417*(cmr)*Hart, D.J. *et al.*, *Chem. Comm.*, 1983, 135*(synth)*Niwa, H. *et al.*, *Chem. Lett.*, 1983, 789*(Farfugine)*Rüeger, H. *et al.*, *Heterocycles*, 1983, **20**, 1331*(synth)*Ohsawa, T. *et al.*, *J.O.C.*, 1983, **48**, 3644 (*synth, Turneforicidine*)Roby, M.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 846*(Angeloylplatynecines)*Freer, A.A. *et al.*, *Acta Cryst. C*, 1987, **43**,2020 (*cryst struct, Platynecine*)Ahmed, W. *et al.*, *Heterocycles*, 1991, **32**, 1729-1736 (*Racemocine*)Roeder, E. *et al.*, *Phytochemistry*, 1991, **30**,1735-1737 (*7-Angeloylturneforicidine*)Ahmed, W. *et al.*, *Fitoterapia*, 1993, **64**, 361*(struct, Racemoneine)*Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**,43-109 (*pmr*)Knight, D.W. *et al.*, *J.C.S. Perkin 1*, 1997, 2087*(synth)*Jenett-Siems, K. *et al.*, *Phytochemistry*, 1998,**47**, 1551-1560 (*Ipangulines*)Goti, A. *et al.*, *Eur. J. Org. Chem.*, 2000, 3633-3645 (*synth*)Wee, A.G.H. *et al.*, *J.O.C.*, 2001, **66**, 8513-8517*(synth, pmr, cmr)*An, D.K. *et al.*, *Org. Lett.*, 2001, **3**, 2961-2963*(synth)*

- Christov, V.S. *et al.*, *Fitoterapia*, 2002, **73**, 171-173 (9-Angeloylthastaneceine)
 Christov, V. *et al.*, *Z. Naturforsch. C*, 2002, **57**, 780-784 (*Diangeloylplatynecine*)
 Zhou, C.-Y. *et al.*, *J.O.C.*, 2004, **69**, 7072-7082 (*synth*)
 Vanecko, J.A. *et al.*, *Org. Lett.*, 2005, **7**, 2949-2952 (*synth*)
 Siciliano, T. *et al.*, *Phytochemistry*, 2005, **66**, 1593-1600 (*Anchusa strigosa ester oxide*)
 Jiang, Y. *et al.*, *Asian J. Pharmacodyn. Pharmacokinet.*, 2006, **6**, 187-192 (*tox. derivs*)
 Marin-Loaiza, J.C. *et al.*, *Phytochemistry*, 2008, **69**, 154-167 (*isol. pmr, cmr, ms*)


Hexahydro-7a-hydroxy-3H-pyrrolizin-3-one, 9CI H-234
 5-Hydroxy-1-azabicyclo[3.3.0]octan-2-one. **Pyrrolam C**
 [104885-96-3]



C₇H₁₁NO₂ 141.169
 Sol. MeOH, CHCl₃; poorly sol. H₂O.

- (±)-*form* [119100-90-2]
 Prod. by *Streptomyces olivaceus*. Oil.
Me ether: Hexahydro-7a-methoxy-3H-pyrrolizin-3-one. **Pyrrolam B**
 [126424-77-9]
 C₈H₁₃NO₂ 155.196
 From *Streptomyces olivaceus*. Sol. MeOH, CHCl₃; poorly sol. H₂O.
 (1-Ethoxyethyl) ether: Hexahydro-7a-(1-ethoxyethoxy)-3H-pyrrolizin-3-one. **Pyrrolam D**
 [126424-78-0]
 C₁₁H₁₉NO₃ 213.276
 From *Streptomyces olivaceus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Isol. as a mix of diastereoisomers.
 Yoshifuji, S. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 5042 (*synth*)
 Ikeda, M. *et al.*, *Heterocycles*, 1988, **27**, 943 (*synth*)
 Grote, R. *et al.*, *Annalen*, 1990, 525 (*isol. pmr, cmr, struct*)

Hexahydro-3-imino-1,2,4-oxadiazepine-3-carboxylic acid, 9CI H-235
Desaminocanavanine. Deaminocanavanine




C₅H₉N₃O₃ 159.144
 Cyclic anhydride of Canavanine, C-74; the name Desaminocanavanine is rather misleading.

- (S)-*form*
L-form
 [21539-44-6]
 May occur with Canavanine, C-74 in


- Jack bean meal (*Canavalia ensiformis*) and other plants. Cryst. Mp 257-258° dec. [α]_D +25. Unreactive to ninhydrin.
 Rosenthal, G.A. *et al.*, *Phytochemistry*, 1972, **11**, 2827 (*synth. bibl*)

1,2,3,10,11,11a-Hexahydro-11-[4-(1H-indol-3-yl)-2-oxo-3-but-nylidene]-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one H-236



- C₂₄H₂₁N₃O₂ 383.449
 (11Z,11aS,14E)-*form*
 Isol. from the myxomycete *Fuligo candida*. Yellow solid. [α]_D²⁶ +149 (c, 0.6 in MeOH). λ_{max} 415 (ε 8700) (MeOH).
 Nakatani, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 368-370 (*isol. cd, pmr, cmr*)

2,3,5,6,11,11b-Hexahydro-3-oxo-1H-indolizino[8,7-b]indole-5-carboxylic acid, 9CI H-237




- C₁₅H₁₄N₂O₃ 270.287
 (5S,11bS)-*form*
Cuscutamine
 [122170-93-8]
 Alkaloid from *Clerodendrum trichotomum* (Verbenaceae) and from the dried fruits of *Cuscuta chinensis* (Convolvulaceae). Amorph. pale yellow powder. [α]_D²⁷ +120.6 (c, 0.73 in MeOH).
Me ester: [70004-59-0]
 Solid (CHCl₃). Mp 109-111°. [α]_D -110 (c, 0.218 in MeOH).

- (5S,11bR)-*form* [122170-94-9]
 Alkaloid from *Clerodendrum trichotomum* (Verbenaceae).
Me ester: [82535-75-9]
 Solid (CHCl₃/hexane). Mp 202-203° (185°). [α]_D +187 (c, 0.214 in MeOH).
 Toyoda, Y. *et al.*, *Chem. Lett.*, 1982, 903 (*ir, pmr, ms, struct, synth*)
 Irikawa, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 880 (*isol. ir, pmr, cmr, ms, struct*)
 Yahara, S. *et al.*, *Phytochemistry*, 1994, **37**, 1755 (*Cuscutamine, isol. pmr, cmr, struct*)

- Fontaine, H. *et al.*, *Synth. Commun.*, 1997, **27**, 2817-2824 (*synth*)


1,2,3,10,11,11a-Hexahydro-11-(2-oxopropylidene)-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one H-238



C₁₅H₁₆N₂O₂ 256.304

- (S)-(Z)-*form*
 Isol. from the myxomycete *Fuligo candida*. Cytotoxic. Plates. Mp 140-144°. [α]_D²⁶ +657 (c, 1.4 in MeOH). Possible artifact. λ_{max} 338 (ε 60000) (MeOH).
 Nakatani, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 368-370 (*isol. cd, pmr, cmr*)


4-(1,2,3,10,11,11a-Hexahydro-5-oxo-5H-pyrrolo[2,1-c][1,4]benzodiazepin-11-ylidene)-3-oxobutanoic acid H-239



C₁₆H₁₆N₂O₄ 300.313

- (S)-(Z)-*form*
 Isol. from the myxomycete *Fuligo candida*. Amorph. solid. [α]_D²⁶ +466 (c, 1 in MeOH). Unstable. λ_{max} 340 (ε 19000) (MeOH).
 Nakatani, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 368-370 (*isol. cd, pmr, cmr*)

3,4,6,7,8,9-Hexahydrophenazine Polycartine A H-240
 [112448-71-2]




C₁₂H₁₄N₂ 186.256
 Isol. from the fruits of *Idesia polycarpa* (Flacourtiaceae). Pale-yellow cryst. (MeOH aq.). Mp 69-71°. λ_{max} 260 (ε); 323 (ε) (MeOH) (Derep).

- Moritake, M. *et al.*, *Tet. Lett.*, 1987, **28**, 1425 (*isol. uv, ir, pmr, cmr, ms, struct*)

3,4,7,8,10,10a-Hexahydro-2H,5H-pyrrolo[2,1-c:3,4-b']bis[1,4]thiazin-5-one


H-241

Aminoethylcysteine ketimine decarboxylated dimer
[83923-12-0]C₉H₁₂N₂OS₂ 228.339**(ξ)-form**


Detected in human plasma and urine and bovine cerebellum, possibly arising from dietary sources. Constit. of fresh vegetables, e.g. garlic, spinach, tomato, asparagus, aubergine, onion, pepper and courgette. Antioxidant. Mp 137-138°.

Hermann, P. et al., *Chem. Ber.*, 1961, **94**, 442-445 (*synth*)Antonucci, A. et al., *Amino Acids*, 1994, **7**, 83-88 (*synth*)Macone, A. et al., *J. Agric. Food Chem.*, 2002, **50**, 2169-2172 (*occur, bibl*)**1,2,3,4,5,6-Hexahydroxyacridone**


H-242

C₁₃H₉NO₇ 291.2172,3,4,5-Tetra-Me ether, N-Me: 1,6-Dihydroxy-2,3,4,5-tetramethoxy-10-methylacridone. **Glyfoline**
[82354-35-6]C₁₈H₁₉NO₇ 361.351Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Orange plates (Me₂CO). Mp 215-217°. λ_{max} 272 (ε 37300); 333 (ε 13200); 409 (ε 3400) (MeOH) (Berdy).Wu, T.-S. et al., *J.C.S. Perkin I*, 1983, 1681 (*uv, ir, pmr, ms, struct*)Su, T.-L. et al., *J. Med. Chem.*, 1992, **35**, 2703 (*synth*)**1,2,3,8,9,10-Hexahydroxyporphine**

H-243

C₁₇H₁₇NO₆ 331.324**(S)-form**1,2:8,9-Bis(methylene), 3,10-di-Me ether: 4a,5,6,7-Tetrahydro-8,13-dimethoxy-5-methyl-4H-bis[1,3]benzodioxolo[6,5,4-de:5',4'-g]quinoline, 9CI. 3,10-Dimethoxy-1,2:8,9-bis(methylenedioxy)porphine. **Ocotominarine**
[72170-07-1]C₂₁H₂₁NO₆ 383.4Alkaloid from the leaves of *Ocotea minarum* (Lauraceae). Mp 148-149°. [α]_D²⁵ +40 (c, 1.0 in CHCl₃).Vecchiotti, V. et al., *Farmaco, Ed. Sci.*, 1979, **34**, 829 (*isol, uv, pmr, ms, struct*)**2,3,14,20,24,25-Hexahydro-ycholest-7-en-6-one**

H-244

C₂₇H₄₄O₇ 480.64**(2β,3β,5β,14α,20S,24ξ)-form Pinnasterone**


[146959-88-8]

Constit. of *Vitex pinnata*. Amorph. solid (MeOH/EtOAc). Mp 198-200°. [α]_D²⁵ +41.2 (c, 0.2 in MeOH).24-O-(2-Pyrrolicarbonyl): **Canescensterone**C₃₂H₄₇NO₈ 573.725Constit. of *Vitex canescens*. Cryst. (MeOH/EtOAc). Mp 152-153°. [α]_D²⁶ +18.8 (c, 0.09 in MeOH).24-Epimer: **24-Epipinnasterone**

[474524-74-8]

C₂₇H₄₄O₇ 480.64Constit. of *Vitex scabra* stem bark.Suksamrarn, A. et al., *Phytochemistry*, 1993, **32**, 303 (*Pinnasterone*)Suksamrarn, A. et al., *Phytochemistry*, 1995, **38**, 473 (*Canescensterone*)Suksamrarn, A. et al., *J. Nat. Prod.*, 2002, **65**, 1690-1692 (*24-Epipinnasterone*)**3,5,7,13,15,18-Hexahydroxy-6,12-cyclo-14-lathyrone**


H-245

C₂₀H₃₂O₇ 384.469

3,18-Bis(3-pyridinecarbonyl), 5-benzoyl, 7-Ac: [213920-27-5]

C₄₁H₄₄N₂O₁₁ 740.805Constit. of *Euphorbia seguieriana*.Cryst. Mp 178°. λ_{max} 220 (log ε 1.5); 253 (log ε 0.63); 260 (log ε 0.7) (EtOH).Öksüz, S. et al., *J. Nat. Prod.*, 1998, **61**, 1198-1201 (*isol, pmr, cmr*)**1,2,4,6,8,9-Hexahydroxydihydro-β-agarofuran**


H-246



(1α,2α,4β,6β,8α,9β)-form

C₁₅H₂₆O₇ 318.366**(1α,2α,4β,6β,8β,9β)-form**6-O-(1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 2,9-dibenzoyl, 1,8-di-Ac: **Reissantin B**
[628290-36-8]C₄₀H₄₃NO₁₃ 745.779Constit. of *Reissantia buchananii*.Powder. Mp 168-172°. [α]_D²⁵ +45.7 (c, 0.32 in CHCl₃).Chang, F.-R. et al., *J. Nat. Prod.*, 2003, **66**, 1416-1420 (*Reissantin B*)**1,2,4,6,9,14-Hexahydroxydihydro-β-agarofuran**

H-247



(1α,2α,4β,6α,9β)-form

C₁₅H₂₆O₇ 318.366**(1α,2α,4β,6β,9β)-form** [31146-58-4]Constit. of *Celastrus angulatus*.1-(3-Pyridinecarbonyl), 9-benzoyl, 2,6,14-tri-Ac: **2,6,14-Triacetoxo-9-(benzoyloxy)dihydro-1-nicotinoyloxy-β-agarofuran**
[130774-23-1]C₃₄H₃₉NO₁₂ 653.682Alkaloid from the root bark of *Celastrus angulatus* (Celastraceae). Insect antifeedant. Amorph. solid. [α]_D²⁰ +43.9 (c, 0.5 in MeOH).1-(3-Pyridinecarbonyl), 9-(3-furancarboxonyl), 2,6,14-tri-Ac: **2,6,14-Triacetoxo-9-furoyloxydihydro-1-nicotinoyloxy-β-agarofuran**
[130774-22-0]C₃₂H₃₇NO₁₃ 643.643Alkaloid from the root bark of *Celastrus angulatus* (Celastraceae). Insect antifeedant. Amorph. solid. [α]_D²⁰ +23.9 (c, 0.5 in MeOH).1-(3-Pyridinecarbonyl), 9-(3-furancarboxonyl), 14-(2-methylpropanoyl), 2,6-di-Ac: **2,6-Diacetoxo-9-furoyloxydihydro-14-(2-methylpropanoyloxy)-1-nicotinoyloxy-β-agarofuran**

[130774-20-8]

C₃₄H₄₁NO₁₃ 671.697

Alkaloid from the root bark of *Celastrus angulatus* (Celastraceae). Insect antifeedant. Mp 127-128°. [α]_D²⁰ +34.5 (c, 0.5 in MeOH).

1-(3-Pyridinecarbonyl), 9-(3-furancarbo-
nyl), 14-(2-methylbutanoyl), 2,6-di-Ac:
2,6-Diacetoxy-9-furoxyloxydihydro-14-
(2-methylbutanoyloxy)-1-nicotinoy-
loxy- β -agarofuran

[130774-21-9]

C₃₅H₄₃NO₁₃ 685.724

Alkaloid from the root bark of *Celastrus angulatus* (Celastraceae). Insect antifeedant. Amorph. solid. [α]_D²⁰ +30.1 (c, 0.5 in MeOH).

2-(3-Pyridinecarbonyl), 6,9-dibenzoyl,
1,14-di-Ac: [389865-22-9]

C₃₉H₄₁NO₁₂ 715.752

Constit. of *Maytenus magellanica* and
Maytenus chubutensis. Amorph. solid.
[α]_D²⁵ +15.4 (c, 0.22 in CHCl₃). λ _{max} 229
; 273 (EtOH).

9-(3-Pyridinecarbonyl), 1,2,6,14-tetra-
Ac: **Maytine**

[31146-56-2]

C₂₉H₃₇NO₁₂ 591.611

Alkaloid from the fruits of *Maytenus serrata* (Celastraceae). Noncryst.

9-(3-Pyridinecarbonyl), 1-benzoyl,
2,6,14-tri-Ac: **2,6,14-Triacetoxy-1-ben-**
zoyloxydihydro-9-nicotinoyloxy- β -agar-
ofuran

[126061-43-6]

C₃₄H₃₉NO₁₂ 653.682

Alkaloid from *Celastrus angulatus*
(Celastraceae).

9-(3-Pyridinecarbonyl), 1-(3-furancarbo-
nyl), 2,6,14-tri-Ac: **2,6,14-Triacetoxy-**
1-furoxyloxydihydro-9-nicotinoyloxy- β -
agarofuran

[126006-73-3]

C₃₂H₃₇NO₁₃ 643.643

Alkaloid from *Celastrus angulatus*
(Celastraceae).

9-(3-Pyridinecarbonyl), 1-(3-furancarbo-
nyl), 6-(2-methylpropanoyl), 2,14-di-
Ac: **2,14-Diacetoxy-1-furoxyloxydihy-**
dro-6-(2-methylpropanoyloxy)-9-nico-
tinoyloxy- β -agarofuran

[126006-71-1]

C₃₄H₄₁NO₁₃ 671.697

Alkaloid from *Celastrus angulatus*
(Celastraceae).

9-(3-Pyridinecarbonyl), 1-(3-furancarbo-
nyl), 6-(2-methylbutanoyl), 2,14-di-Ac:
2,14-Diacetoxy-1-furoxyloxydihydro-6-
(2-methylbutanoyloxy)-9-nicotinoy-
loxy- β -agarofuran

[126006-72-2]

C₃₅H₄₃NO₁₃ 685.724

Alkaloid from *Celastrus angulatus*
(Celastraceae).


Kupchan, S.M. et al., *J.O.C.*, 1977, **42**, 115
(*Maytine*)

Liu, J. et al., *Phytochemistry*, 1990, **29**, 2503-
2506 (*Celastrus angulatus* constits)

Kennedy, M.L. et al., *J. Med. Chem.*, 2001,
44, 4668-4676 (*Maytenus magellanica*
constit)

1,2,6,8,9,14-Hexahydroxydi- hydro- β -agarofuran

H-248

(1 α ,2 α ,6 β ,8 α ,9 α)-formC₁₅H₂₆O₇ 318.366

(1 α ,2 α ,6 β ,8 α ,9 α)-form Alatof†

[57461-75-3]

Constit. of *Euonymus alatus*. Amorph.

8-(3-Pyridinecarbonyl), 9-benzoyl,

2,6,14-tri-Ac: [185414-42-0]

C₃₄H₃₉NO₁₂ 653.682

Constit. of *Celastrus angulatus*. Needles
(Me₂CO). Mp 200-202°. [α]_D -43.1 (c,
0.51 in MeOH). λ _{max} 201 (log ϵ 3.16);
219 (log ϵ 3.14); 263 (log ϵ 2.49) (MeOH).

3-Pyridinecarbonyl, benzoyl, tetra-Ac:

Ever 6

[105286-90-6]

C₃₆H₄₁NO₁₃ 695.719

Isol. from *Euonymus verrucosus*. Mp
89-96°.

8-(3-Pyridinecarbonyl), 1,9-dibenzoyl,

2,6,14-tri-Ac: [185414-41-9]

C₄₁H₄₃NO₁₃ 757.79

Constit. of *Celastrus angulatus*. Needles
(Me₂CO). Mp 206-208°. [α]_D -51.3 (c,
0.54 in MeOH). λ _{max} 201 (log ϵ 3.29);
226 (log ϵ 3.26); 263 (log ϵ 2.53) (MeOH).

3-Pyridinecarbonyl, dibenzoyl, tri-Ac:

Ever 7

[105286-93-9]

C₄₁H₄₃NO₁₃ 757.79

Isol. from *Euonymus verrucosus*. Cryst.
(MeOH). Mp 83-95°.

(1 α ,2 α ,6 β ,8 α ,9 β)-form

2,8-Bis-(3-pyridinecarbonyl), 9-benzoyl,

1,6,14-tri-Ac: **Cangorin J**

[156400-98-5]

C₄₀H₄₂N₂O₁₃ 758.777

Constit. of *Maytenus ilicifolia*.
Amorph. solid. Mp 115-120°. [α]_D +20
(c, 0.26 in CHCl₃).

Begley, M.J. et al., *J.C.S. Perkin 1*, 1986, 535-
539 (*Ever series*)


Itokawa, H. et al., *J. Nat. Prod.*, 1994, **57**, 460
(*Cangorin J*)

Wang, Y. et al., *J. Nat. Prod.*, 1997, **60**, 178-
179 (*alkaloids*)

1,4,6,8,9,14-Hexahydroxydi- hydro- β -agarofuran

H-249

5,11-Epoxy-1,4,6,8,9,14-eudesmanhexol

(1 α ,4 β ,6 β ,8 α ,9 α)-formC₁₅H₂₆O₇ 318.366

(1 α ,4 β ,6 β ,8 α ,9 α)-form

IsoalatoI

1-O-(3-Pyridinecarbonyl): **Tripterrege-**
line AC₂₁H₂₉NO₈ 423.462

Alkaloid from the stem bark of *Tripter-*
terygium regelii (Celastraceae).

8-O-(3-Pyridinecarbonyl), 1-O-(E-cinna-
moyl), 9-benzoyl, 6,14-di-Ac: **Tripto-**
fordinine A1

[112494-33-4]

C₄₁H₄₃NO₁₂ 741.79

Alkaloid from leaves of *Tripter-*
terygium wilfordii var. *regelii* (Celastraceae). Mp
193-194°. [α]_D -81.4 (MeOH).

8-O-(3-Pyridinecarbonyl), 1-O-(Z-cinna-
moyl), 9-benzoyl, 6,14-di-Ac: **Tripto-**
fordinine A2

[112571-89-8]

C₄₁H₄₃NO₁₂ 741.79

Alkaloid from *Tripter-*
terygium wilfordii
var. *regelii*. Mp 93-94°. [α]_D -101
(MeOH).

9-O-(3-Pyridinecarbonyl), 1-benzoyl:

Tripterregeline BC₂₈H₃₃NO₉ 527.57

Alkaloid from the stem bark of *Tripter-*
terygium regelii (Celastraceae).

9-O-(3-Pyridinecarbonyl), 1-benzoyl,

8,14-di-Ac: **Tripterregeline C**C₃₂H₃₇NO₁₁ 611.644

Alkaloid from the stem bark of *Tripter-*
terygium regelii (Celastraceae).

9-O-(3-Pyridinecarbonyl), 8-benzoyl,

1,6,14-tri-Ac: [339151-91-6]

C₃₄H₃₉NO₁₂ 653.682

Constit. of *Tripter-*
terygium wilfordii.
Amorph. powder. [α]_D²⁵ -52.4 (c, 0.4 in
MeOH). λ _{max} 216 (log ϵ 4.31); 253 (log
 ϵ 3.64) (MeOH).

(1 α ,4 β ,6 β ,8 β ,9 α)-form

8-O-(N-Methyl-6(1H)-pyridone-3-carbo-
nyl), 1-O-(3-furancarbo-nyl), 9-benzoyl,6-Ac: **Reissantin F**

[910651-20-6]

C₃₈H₄₁NO₁₄ 735.74Constit. of *Reissantia buchananii*.

Powder. Mp 138-139°. [α]_D²⁴ +35.7 (c,
0.23 in CHCl₃). λ _{max} 219 (log ϵ 4.35);
270 (log ϵ 4.07) (MeOH).

[128385-39-7, 128197-56-8]

Takaishi, Y. et al., *Chem. Pharm. Bull.*, 1987,
35, 3534 (*Triptofordinins*)


Han, B.H. et al., *Arch. Pharmacol Res.*, 1989,
12, 310 (*Tripterregelines*)

Duan, H.-Q. et al., *Phytochemistry*, 2001, **56**,
341-346 (*Tripter-*
terygium wilfordii ester)

Chang, F.-R. et al., *Planta Med.*, 2006, **72**, 92-
96 (*Reissantin F*)


2,3,5,7,8,15-Hexahydroxy- 6(17),11-jatrophadiene-9,14-dione

H-250


(2 α ,3 β ,5 α ,7 β ,8 α ,11E,13 α ,15 β)-form

C₂₀H₃₀O₈ 398.452**(2 α ,3 β ,5 α ,7 β ,8 α ,11E,13 α ,15 β)-form**

2-(3-Pyridinecarbonyl), 3,5,7,8,15-penta-Ac:

C₃₆H₄₃NO₁₄ 713.734Constit. of *Euphorbia semiperfoliata*. Gum.Appendino, G. et al., *J. Nat. Prod.*, 1998, **61**, 749-756 (*Euphorbia semiperfoliata* esters)**2,3,5,7,9,15-Hexahydroxy-6(17),11-jatrophadien-14-one** H-251C₂₀H₃₂O₇ 384.469**(2 α ,3 β ,5 α ,7 β ,9 α ,11E,13 α H,15 β)-form**

9-(3-Pyridinecarbonyl), 2,3,5,7,15-penta-Ac: [210108-85-3]

C₃₆H₄₅NO₁₃ 699.75Constit. of *Euphorbia pepul*.Jakupovic, J. et al., *Phytochemistry*, 1998, **47**, 1601-1609 (*isol*, *pmr*, *cmr*)**2,3,5,8,9,15-Hexahydroxy-6(17),11-jatrophadien-14-one** H-252C₂₀H₃₂O₇ 384.469**(2 α ,3 β ,5 α ,8 α ,9 α ,11E,13 β ,15 β)-form**9-(3-Pyridinecarbonyl), 3-benzoyl, 5,8-di-Ac: **Euphocharacin B**

[777896-13-6]

C₃₇H₄₃NO₁₁ 677.747Constit. of *Euphorbia characias*.Amorph. solid. [α]_D²⁵ +117.7 (c, 0.1 in CHCl₃).9-(3-Pyridinecarbonyl), 3-benzoyl, 5,8,15-tri-Ac: **Euphocharacin A**

[777896-12-5]

C₃₉H₄₅NO₁₂ 719.784Constit. of *Euphorbia characias*.Amorph. solid. [α]_D²⁵ -22.17 (c, 0.1 in CHCl₃).3,9-Dibenzoyl, 5,8-di-Ac: **Euphocharacin C**


[777896-15-8]

C₃₈H₄₄O₁₁ 676.759Constit. of *Euphorbia characias*.Amorph. solid. [α]_D²⁵ +16.67 (c, 0.1 in CHCl₃).9-(3-Pyridinecarbonyl), 3-(2-methylbutanoyl), 5,8,15-tri-Ac: **Euphocharacin D**

[777896-17-0]

C₃₇H₄₉NO₁₂ 699.794Constit. of *Euphorbia characias*.Amorph. solid. [α]_D²⁵ +8 (c, 0.1 inCHCl₃).9-(3-Pyridinecarbonyl), 3-benzoyl, 8,15-di-Ac: **Euphocharacin L**


[777896-33-0]

C₃₇H₄₃NO₁₁ 677.747Constit. of *Euphorbia characias*.Amorph. solid. [α]_D²⁵ -40 (c, 0.1 in CHCl₃).Corea, G. et al., *Planta Med.*, 2004, **70**, 657-665 (*isol*, *pmr*, *cmr*)**1,6,7,8,14,16-Hexahydroxy-4-methylnaconitan-19-one** H-253C₁₉H₂₇NO₇ 381.425**(1 α ,6 β ,14 α ,16 β)-form**7,8-Methylene ether, O¹,O⁶,O¹⁶-tri-Me:**Tongolenine D**

[198754-26-6]

C₂₃H₃₃NO₇ 435.516Alkaloid from *Delphinium tongolense* (Ranunculaceae). Needles. Mp 310-311°. [α]_D²⁴ +5.7 (c, 0.27 in CHCl₃).

Possible artifact.

He, L. et al., *Chin. Chem. Lett.*, 1997, **8**, 791-792 (*isol*, *pmr*, *cmr*, *struct*)He, L. et al., *Indian J. Chem., Sect. B*, 1998, **37**, 612-614**1,6,13,14,16,18-Hexahydroxy-4-methylnaconitan-15-one** H-254C₁₉H₂₇NO₇ 381.425**(1 α ,5 β ,6 α ,14 α ,16 α)-form**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl: **16-Epipyrophyaconitine**

[123842-96-6]

C₃₁H₄₁NO₈ 555.667Alkaloid from the processed aconite kako-bushi-matsu from the roots of some *Aconitum* spp. (Ranunculaceae).Amorph. [α]_D -65 (c, 0.18 in EtOH).

Artifact.


▶ AR5578000

(1 α ,5 β ,6 α ,14 α ,16 β)-formO¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl: **Pyrophyaconitine**

[123842-97-7]

C₃₁H₄₁NO₈ 555.667Alkaloid from kako-bushi-matsu from the roots of some *Aconitum* spp.(Ranunculaceae). Amorph. [α]_D +15.6 (c, 0.18 in EtOH). Artifact formed from Hypaconitine in M-343 present in the raw aconite roots during the processing procedure. Has not been isol. so far from the raw roots of *A. spp.*

▶ AR5580000

Mori, T. et al., *Heterocycles*, 1989, **29**, 873 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)**3,5,7,13,15,17-Hexahydroxy-8,10(18)-myrsinadien-14-one** H-255C₂₀H₃₀O₇ 382.453


Various structs. revised in 2001.

(2 β ,3 β ,5 α ,7 β ,13 β ,15 β)-form5-O-(3-Pyridinecarbonyl), 3-butanoyl, 7,15-di-Ac: **Decipinol ester A**C₃₄H₄₃NO₁₁ 641.714Constit. of *Euphorbia decipiens*. [α]_D²³ -25.3 (c, 0.786 in CHCl₃). λ _{max} 204 ; 217 ; 264 (MeOH).17-O-(3-Pyridinecarbonyl), 3,7,15-tri-Ac: **Decipinone A**

[220705-93-1]

C₃₆H₄₅NO₁₂ 683.751Constit. of *Euphorbia decipiens*. Powder. [α]_D²⁷ -9.1 (c, 0.22 in CHCl₃).

17→14 Acetal, 5-O-(3-pyridinecarbonyl), 3,7,15-tri-Ac: [616217-03-9]

C₃₂H₃₉NO₁₁ 613.66Constit. of *Euphorbia decipiens*. Oil. λ _{max} 199 ; 270 (MeOH).Ahmad, V.U. et al., *Planta Med.*, 1998, **64**, 728-731 (*Decipinones A-C*)Jassbi, A.R. et al., *Helv. Chim. Acta*, 2002, **85**, 1706-1713 (*abs config*)Ahmad, V.U. et al., *Chem. Pharm. Bull.*, 2003, **51**, 719-723 (*acetal derivs*)**3,8,14,15,16,18-Hexahydroxyypregnan-20-one** H-256C₂₁H₃₄O₇ 398.495**(3 β ,5 α ,8 β ,14 β ,15 β ,16 α)-form****Shitakigenin**

[68457-66-9]

O¹⁸-(2-Methylaminobenzoyl): **Stephan-****thraniline C**

[68564-11-4]


C₂₉H₄₁NO₈ 531.645Alkaloid from stems of *Stephanotis japonica* (Asclepiadaceae). Amorph.

$[\alpha]_D +4.2$ (c, 0.58 in CHCl_3).

Terada, S. *et al.*, *Tet. Lett.*, 1978, **23**, 1995

2,3-Hexalobine E H-257

3-(2,3-Epoxy-3-methylbutyl)-2-(3-hydroxy-3-methyl-1-butenyl)-1H-indole
[103772-00-5]



$\text{C}_{18}\text{H}_{23}\text{NO}_2$ 285.385

Alkaloid from stem bark of *Hexalobus crispiflorus*. Oil. $[\alpha]_D^{21} +12$ (c, 1.0 in CHCl_3). λ_{max} 231 (log ϵ 4.3); 241 (sh); 306 (log ϵ 4.4); 315 (log ϵ 4.4) (MeOH).

Achenbach, H. *et al.*, *Annalen*, 1995, 1327-1337 (isol, uv, ir, pmr, cmr, ms, struct)

Hexanedioic acid, 9CI H-258

Adipic acid. Butane-1,4-dicarboxylic acid. E355. FEMA 2011

[124-04-9]

$\text{HOOCCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{COOH}$

$\text{C}_6\text{H}_{10}\text{O}_4$ 146.143

Obt. industrially by oxidn. of Cyclohexane by air/catalyst followed by HNO_3 oxidn. of Cyclohexanol.

Cyclohexanone intermed. Constit. of leaves and stems of *Anthyllis sericea*. Also isol. from guava fruit (*Psidium guajava*), papaya (*Carica papaya*) and raspberry (*Rubus idaeus*). Used in manuf. of nylons, esp. nylon-6,6; also plasticisers, resins and as food acidulant. Used in pptn. of Al, Zr, U(VI) and standardisation of base solns. Important industrial chemical, 46th in order of volume for USA in 1994 (production 0.90 million tons/year). Monoclinic cryst. (HNO_3). Sol. EtOH; mod. sol. H_2O ; spar. sol. Et_2O . Mp 153° (149-150°). Bp 337.5° Bp₁₀₀ 265° Bp₁₅ 216°. $\text{p}K_{\text{a}1}$ 4.41; $\text{p}K_{\text{a}2}$ 5.41 (20°). $\text{p}K_{\text{a}1}$ 4.44; $\text{p}K_{\text{a}2}$ 5.44 (25°). Sublimes. Does not readily form a monomeric anhydride; forms a polymeric anhydride which gives the unstable monomer on dist.

- Fl. p. 196°, autoignition temp. 420°. Severe eye and mucous membrane irritant. LD₅₀ (mus, orl) 1900 mg/kg. AU8400000

Diamide: Hexanediamide. Adipamide

[628-94-4]

$\text{C}_6\text{H}_{12}\text{N}_2\text{O}_2$ 144.173

Isol. from *Rhizoclonium hieroglyphicum*. Prisms. Mp 220°. Subl. 0.01 150-160.


- AU7800000

[23311-84-4, 19090-60-9, 27178-16-1, 3385-41-9, 110-29-2, 29254-17-9]

Dembitsky, V.M. *et al.*, *Phytochemistry*, 2000, **54**, 965-967 (isol, diamide)

2-(5-Hexenyl)-3,4-dihydro-5-nonyl-2H-pyrrole, 9CI H-259

[107342-26-7]



$\text{C}_{19}\text{H}_{35}\text{N}$ 277.492


Alkaloid from the venom glands of the ant *Monomorium minutum*.

Clément, J.-L. *et al.*, *C. R. Hebd. Seances Acad. Sci., Ser. III*, 1986, **303**, 669 (isol, ms, struct)

3-(5-Hexenyl)octahydro-5-(4-pentenyl)indolizine, 9CI H-260

3-(5-Hexenyl)-5-(4-pentenyl)indolizidine. Mantella Alkaloid 275C. Indolizidine 275C

[151805-31-1]



$\text{C}_{19}\text{H}_{33}\text{N}$ 275.476

Provisional struct. Minor or trace alkaloid from skin extracts of 2 populations of the Madagascan frog *Mantella madagascariensis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (isol, ir, ms)

1-Hexylamine H-261

1-Hexanamine, 9CI. 1-Aminohexane

[111-26-2]

$\text{H}_3\text{C}(\text{CH}_2)_4\text{CH}_2\text{NH}_2$

$\text{C}_6\text{H}_{15}\text{N}$ 101.191

Metab. of *Claviceps purpurea* and *Claviceps microcephala*. Fp -19. Bp₇₄₂ 129-130°. $\text{p}K_{\text{a}}$ 10.56 (25°).

- Flammable, fl. p. 29° (oc). Severe skin and eye irritant. LD₅₀ (rat, orl) 670 mg/kg. LD₅₀ (rbt, skn) 420 mg/kg. MQ4540000

Hydrochloride: [142-81-4]

Leaflets. Mp 219°.

N-Ac:

$\text{C}_8\text{H}_{17}\text{NO}$ 143.228

Bp₁₄ 152°.

N-Me: [35161-70-7]

$\text{C}_7\text{H}_{17}\text{N}$ 115.218

d 0.76. Bp 140-142° (137°). n_D^{20} 1.4160.

N,N-Di-Me: [4385-04-0]

$\text{C}_8\text{H}_{19}\text{N}$ 129.245

Bp 146-150°. n_D^{20} 1.4136.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 281C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 451B; 485A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 365B; 385B (ir)

v. Braun, J. *et al.*, *Ber.*, 1910, **43**, 3596 (synth)

Steiner, M. *et al.*, *Naturwissenschaften*, 1955, **42**, 345 (anal)

Gohlke, R.S. *et al.*, *Anal. Chem.*, 1962, **34**,

1281 (ms)

Hartmann, T. *et al.*, *Planta Med.*, 1964, **12**, 340 (isol)

Eggert, H. *et al.*, *J.A.C.S.*, 1973, **95**, 3710 (cmr) *Fieser and Fieser's Reagents for Organic Synthesis*, Wiley, 1979, **7**, 170 (use)

Nikishin, G.I. *et al.*, *Tetrahedron*, 1985, **41**,

4279 (deriv synth, ir, pmr, ms)


Erra-Balsells, R. *et al.*, *Aust. J. Chem.*, 1988, **41**, 103 (pmr, cmr, ms)

Kabalka, G.W. *et al.*, *Organometallics*, 1989, **8**, 1093-1095 (N-Me, synth, cmr)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HFK000

2-Hexyl-3,5-dimethylpyrazine H-262

[70303-40-1]



$\text{C}_{12}\text{H}_{20}\text{N}_2$ 192.303

Constit. of the glands of various ants incl. *Odontomachus* sp. and *Streblognathus* sp.


Longhurst, C. *et al.*, *J. Insect Physiol.*, 1978, **24**, 833-837 (isol)

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1998, **24**, 125-134 (isol, ms)

2-Hexyl-4-hydroxyquinoline H-263

2-Hexyl-4(1H)-quinolinone. 2-Hexyl-4-quinolinol, 9CI

[678172-85-5]



$\text{C}_{15}\text{H}_{19}\text{NO}$ 229.321

Metab. of *Pseudomonas aeruginosa*.

N-Oxide: 2-Hexyl-4(1H)-quinolinone N-oxide. 2-Hexyl-4-hydroxyquinoline N-oxide

[678172-92-4]

$\text{C}_{15}\text{H}_{19}\text{NO}_2$ 245.321

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(E-): 2-(1-Hexenyl)-4(1H)-quinolinone. 2-(1-Hexenyl)-4-hydroxyquinoline. 2-(1-Hexenyl)-4-quinolinol

$\text{C}_{15}\text{H}_{17}\text{NO}$ 227.305

Metab. of *Pseudomonas aeruginosa*.

No CAS no. found to 2007.

1',2'-Didehydro(Z-):

$\text{C}_{15}\text{H}_{17}\text{NO}$ 227.305

Metab. of *Pseudomonas aeruginosa*.


No CAS no. found to 2007.

Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (occur, oxide, 1',2'-didehydro)

Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (biosynth)

5-Hexylindolizidine H-264

5-Hexyloctahydroindolizidine, 9CI. Indolizidine 209D. Dendrobates Alkaloid 209D [129784-76-5, 153922-74-8]

C₁₄H₂₇N 209.374**(5R,9R)-form** [129709-01-9]

Isol. from skin extracts of an undescribed *Dendrobates* sp. Oil. $[\alpha]_D^{26}$ -80.4 (c, 1 in CH₂Cl₂).

Polniaszek, R.P. *et al.*, *J.O.C.*, 1990, **55**, 4688-4693 (*synth, pmr, cmr*)

Lee, E. *et al.*, *Tet. Lett.*, 1996, **37**, 1445-1446 (*synth*)

Kim, G. *et al.*, *Tetrahedron: Asymmetry*, 2001, **12**, 2073-2076 (*synth*)

Reddy, P.G. *et al.*, *J.O.C.*, 2004, **69**, 3093-3101 (*synth, pmr, cmr*)

Patil, N.T. *et al.*, *Tet. Lett.*, 2005, **46**, 2101-2103 (*synth*)


Takahata, H. *et al.*, *Heterocycles*, 2006, **67**, 407-411 (*synth*)

Alegret, C. *et al.*, *J.O.C.*, 2008, **73**, 8661-8664 (*synth*)

Pohmakotr, M. *et al.*, *Tetrahedron*, 2008, **64**, 2339-2347 (*synth*)

2-Hexyl-8-methoxy-1-methyl-4(1H)-quinolinone, 9CI

[159979-56-3]


C₁₇H₂₃NO₂ 273.374

Alkaloid from trunk bark of *Esenbeckia almayillia* (Rutaceae). Oil.

Guilhon, G.M.S.P. *et al.*, *Phytochemistry*, 1994, **37**, 1193-1195 (*isol, ir, pmr, cmr, ms, struct*)

3-Hexyloctahydro-5-methylindolizine, 9CI

3-Hexyl-5-methylindolizidine



(3R*,5R*,8aR*)-form

C₁₅H₂₉N 223.401**(3R,5S,8aS)-form** [153481-99-3]

Synthetic. Oil. $[\alpha]_D^{24}$ +27.9 (c, 1.35 in hexane).

(3S,5S,8aS)-form [153482-03-2]

Synthetic. Oil. $[\alpha]_D^{24}$ +90.1 (c, 0.28 in MeOH).

(3R,5R,8aR)-form [208849-62-1]

Isol. from the venom of thief ants, *Solenopsis* sp.

(3R,5S,8aR)-form [92512-78-2]

Isol. from the venom of *Solenopsis* sp.

(3R,5S,8aS)-form [181378-97-2]

Isol. from the venom of *Solenopsis* sp.


Takahata, H. *et al.*, *Tetrahedron*, 1993, **49**, 11205-11212 (*synth, ir, pmr, cmr*)

Jones, T.H. *et al.*, *J. Chem. Ecol.*, 1996, **22**, 1221-1236 (*isol*)

Gorman, J.S.T. *et al.*, *J. Chem. Ecol.*, 1998, **24**, 933-943 (*isol*)

3-Hexylpyridine, 9CI

[6311-92-8]

C₁₁H₁₇N 163.262

Found in orange oil. Perfumery and flavouring ingredient. Liq. with sweet fragrance. Bp_{0.05} 68-71°.

Eur. Pat., 1992, 470 391; *CA*, **116**, 194165 (*synth, use*)


Thomas, A.F. *et al.*, *J. Agric. Food Chem.*, 1992, **40**, 2236-2243 (*synth, isol, ms, pmr*)

Tereshko, A.B. *et al.*, *Zh. Org. Khim.*, 1995, **31**, 289-290; *J. Org. Chem. USSR (Engl. Transl.)*, 1995, **31**, 258-259 (*synth*)

Schwab, P.F.H. *et al.*, *J.O.C.*, 2002, **67**, 443-449 (*synth, pmr*)

2-Hexylpyrrole, 9CI

[1551-14-0]

C₁₀H₁₇N 151.251Bp_{1.5} 92°.

N-Sulfo: 2-Hexylpyrrole sulfamate

[587875-53-4]

C₁₀H₁₇NO₃S 231.315

Alkaloid from the marine annelid *Cirriformia tentaculata*. Glass (as Na salt).

Utimoto, K. *et al.*, *Tet. Lett.*, 1981, **22**, 4277-4288 (*synth*)


Garrido, D.O.A. *et al.*, *J.O.C.*, 1984, **49**, 2619-2622 (*synth*)

Barsby, T. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1110-1112 (*sulfamate*)

1-[(5-Hexyltetrahydro-2-furanyl)acetyl]-1H-pyrrole, 9CI

3,6-Epoxydodecanoic acid pyrrolide

[111509-36-5]


C₁₆H₂₅NO₂ 263.379

Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (*isol, uv, ir, pmr, cmr, ms, struct*)

Heyneatine

[76129-65-2]



Absolute Configuration

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the wood and stem bark of *Ervatamia heyneana* (Apocynaceae). Cytotoxic agent. Powder (90% EtOH aq.). $[\alpha]_D^{26}$ -6 (c, 0.14 in CHCl₃). λ_{max} 220 (ε 31600); 280 (ε 10480); 300 (ε 8520) (MeOH) (Berdy).

19-Epimer, demethoxy: **3,19-Oxidocoronaridine**

C₂₁H₂₄N₂O₃ 352.432


Alkaloid from the stem bark of *Ta-bernaemontana dichotoma* (Apocynaceae). Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Exists in equilib. with its open form, a mixt. of 3R- and 3S-Hydroxy-19R-heyneanine. λ_{max} 226 (ε 16600); 286 (ε 3800); 293 (ε 3650) (EtOH) (Berdy).

Gunasekera, S.P. *et al.*, *Phytochemistry*, 1980, **19**, 1213 (*Heyneatine*)

Perera, P. *et al.*, *Phytochemistry*, 1985, **24**, 2097 (*Oxidocoronaridine*)

Hibiscuwanin A

N-(4'-O-Feruloylferuloyl)tyramine

C₂₈H₂₇NO₇ 489.524

Constit. of the stems of *Hibiscus taiwanensis*. Pale yellow powder. λ_{max} 290 (log ε 4.23); 318 (log ε 4.3) (MeOH).


7'Z-Isomer: **Hibiscuwanin B**C₂₈H₂₇NO₇ 489.524

Constit. of the stems of *Hibiscus taiwanensis*. Pale yellow powder. λ_{max} 288 (log ε 4.19); 315 (log ε 4.22) (MeOH).

Wu, P.-L. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 2193-2197 (*isol, pmr, cmr, ms*)

Hibispeptin A

[204439-16-7]

R = -CH₂PhC₃₉H₅₀N₆O₈ 730.859

Cyclic peptide. Constit. of the root bark of *Hibiscus syriacus*. Shows antimicrobial, cytotoxic, antioxidative and tyrosinase inhibitory activities. Powder. $[\alpha]_D$ -38 (c,

0.2 in CHCl₃/MeOH). λ_{\max} 206 (ϵ 31200); 224 (sh) (ϵ 11300); 276 (ϵ 8600) (MeOH).
 Yoo, I.-D. *et al.*, *Phytochemistry*, 1997, **47**, 799-802 (*isol. activity*)
 Yun, B.-S. *et al.*, *Tet. Lett.*, 1998, **39**, 993-996 (*isol. uv, ir, pmr, cmr*)
 Yun, B.-S. *et al.*, *Tetrahedron*, 1998, **54**, 15155-15160 (*activity*)

Hibispeptin B**H-273**

[219825-49-7]

As Hibispeptin A, H-272 with


R = CH₂CH(CH₃)₂C₃₆H₅₂N₆O₈ 696.842

Cyclic peptide. Constit. of the root bark of *Hibiscus syriacus*. Powder. $[\alpha]_D^{25}$ -42.7 (c, 0.75 in CHCl₃/MeOH). λ_{\max} 209 (ϵ 37500); 221 (sh) (ϵ 24400); 272 (ϵ 23300); 325 (ϵ 9800); 360 (ϵ 9000) (MeOH).

Yun, B.-S. *et al.*, *Tetrahedron*, 1998, **54**, 15155-15160

Hicksoane A**H-274**

[1033618-61-9]



Absolute Configuration

C₂₀H₂₇IN₄O₂ 482.363

Alkaloid from *Subergorgia hicksoni*. Pale yellow powder. $[\alpha]_D^{20}$ +31 (c, 0.01 in MeOH). λ_{\max} 227 (log ϵ 4.21); 286 (log ϵ 3.75) (MeOH).

6'-Deiodo, 4'-iodo: Hicksoane B

[1033618-62-0]

C₂₀H₂₇IN₄O₂ 482.363Alkaloid from *Subergorgia hicksoni*.

Pale yellow powder. $[\alpha]_D^{20}$ +37 (c, 0.009 in MeOH). λ_{\max} 228 (log ϵ 4.17); 285 (log ϵ 3.65) (MeOH).

4'-Iodo: Hicksoane C

[1033618-63-1]


C₂₀H₂₆I₂N₄O₂ 608.26Alkaloid from *Subergorgia hicksoni*.

Yellow powder. $[\alpha]_D^{20}$ +41 (c, 0.007 in MeOH). λ_{\max} 225 (log ϵ 4.37); 284 (log ϵ 3.86) (MeOH).

Rezanka, T. *et al.*, *Eur. J. Org. Chem.*, 2008, 1265-1270 (*isol. pmr, cmr*)

Higenamine**H-275**

1,2,3,4-Tetrahydro-1-[(4-hydroxyphenyl)methyl]-6,7-isoquinolinediol, 9CI. 1,2,3,4-Tetrahydro-6,7-dihydroxy-1-(4-hydroxybenzyl)isoquinoline. Demethylcoclaurine [17072-47-8]

**(R)-form**C₁₆H₁₇NO₃ 271.315

Historically, Higenamine was the racemate and Demethylcoclaurine the opt. active form, probably the (S)-form.

(R)-form [106032-53-5]

Alkaloid from the seed embryo of *Nelumbo nucifera* (East India lotus) (Nelumbonaceae). Prisms (EtOH). Mp 242-244° dec. $[\alpha]_D^{26.5}$ +16 (c, 1 in MeOH). Abs. config. not proven.

N-Me, N-oxide (R-): N-Methylhigenamine N-oxide

[240430-84-6]

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Gnetum parvifolium*. Needles (H₂O). Mp 156-159°. $[\alpha]_D$ -9.6 (c, 0.12 in DMSO). λ_{\max} 225 (sh) (log ϵ 4.35); 284 (log ϵ 3.64) (H₂O).

O⁶-Me: Coclaurine. Sanjoinine K

[2196-60-3]

[486-39-5]

C₁₇H₁₉NO₃ 285.342

Alkaloid from *Xylopia papuana* and *Alseodaphne archboldiana* (Annonaceae, Lauraceae). Also isol. from Sanjoin (seeds of *Zizyphus vulgaris* var. *spinosus*) (Rhamnaceae). Anti-HIV and hypotensive agent. Mp 217-218°. $[\alpha]_D$ +47 (EtOH). The *A. archboldiana* isolate, $[\alpha]_D$ +22°, was a partial racemate (1:1 mixt. of (+)- and (±)-forms).

O⁶-Me, N,O,O-tri-Ac: Mp 199°. $[\alpha]_D$ -76 (EtOH).

O⁶,N-Di-Me: N-Methylcoclaurine

[5096-70-8]

C₁₈H₂₁NO₃ 299.369

Alkaloid from *Phyllica rogersii* (Rhamnaceae). Weak neuromuscular inhibitor. Mp 182-183°. $[\alpha]_D$ -80 (c, 0.17 in EtOH). $[\alpha]_D$ -62 (c, 0.13 in CHCl₃). Remarkably (+)-Coclaurine forms a (-)-N-methyl deriv. λ_{\max} 226 (ϵ 20600); 286 (ϵ 7100) (EtOH).

O⁷-Me: Isococlaurine

[574-75-4]

C₁₇H₁₉NO₃ 285.342

Alkaloid occurring to small extent in commercial *Radix pareirae* bravae. Isol. from roots of *Cissampelos mucronata*. Plates (MeOH). Mp 216-217°. $[\alpha]_D$ +12 (MeOH).

O⁷-Me, hydrochloride:Plates + 1H₂O (H₂O). Mp 175-176°. $[\alpha]_D$ +28 (H₂O). $[\alpha]_D$ +16 (MeOH).**O⁶,O⁷-Di-Me:** see Armepavine, A-1444**O⁶,N,N-Tri-Me: (-)-Magnocurarine**

[6801-40-7]

C₁₉H₂₄NO₃[⊕] 314.404

Alkaloid from *Magnolia obovata* and several other *Magnolia* spp. and from *Michelia fuscata* (Magnoliaceae). Curarising and ganglionic blocking agent. Hypotensive. Mp 199-200° (as chloride). $[\alpha]_D$ -91 (H₂O).

O⁷,N,N-Tri-Me: Lotusine

[6871-67-6]

C₁₉H₂₄NO₃[⊕] 314.404

Quaternary alkaloid from the embryo of *Nelumbo nucifera* (East India lotus) and from *Tiliacora racemosa* (Nelum-

bonaceae). Mp 213-215° (as chloride). $[\alpha]_D$ -15 (MeOH) (chloride). Another natural product (cyanogenic glycoside) reported in 1901 and named Lotusin was erroneous. λ_{\max} 282 (EtOH).

O⁶,N,N-Tri-Me: Luxandrine

[107584-81-6]

C₁₉H₂₄NO₃[⊕] 314.404

Quaternary alkaloid from the trunk bark of *Pseudoxandra sclerocarpa*. Pale brown amorph. solid (as chloride). $[\alpha]_D$ +27 (c, 0.24 in CHCl₃/MeOH). Dec. in soln.

O⁴,O⁶,N-Tri-Me: N-Demethylcolletine.**4'-Methyl-N-methylcoclaurine**

[19879-50-6]

C₁₉H₂₃NO₃ 313.396

Obt. synthetically as a prod. from cleavage of Neferine, N-87 by Na/liq. NH₃. Alkaloid from *Aconitum leucostomum*. Oil. $[\alpha]_D^{20}$ -74 (c, 0.7 in MeOH).

O⁴,O⁶,N,N-Tetra-Me: Colletine

[15500-51-3]

C₂₀H₂₆NO₃[⊕] 328.43

Quaternary alkaloid from the aerial parts of *Colletia spinosissima* (Rhamnaceae). Cryst. + 1H₂O (EtOAc/EtOH) (as chloride). Mp 130-132° (chloride). $[\alpha]_D^{20}$ -132.8 (c, 1.07 in EtOH).

(S)-form [22672-77-1]**O⁷-Me:** [486-39-5]

Obt. by resoln. of (±)-O,O-Dibenzylisococlaurine and subsequent acid hydrolyd. Cryst. (MeOH). Mp 195-197°. $[\alpha]_D$ -14 (c, 0.5 in MeOH).

O⁷-Me, hydrochloride: [3422-42-2]

Needles (MeOH). Mp 242-243°. $[\alpha]_D$ -18 (c, 0.5 in MeOH). $[\alpha]_D$ -27.5 (c, 1.0 in H₂O).

O⁶,N-Di-Me: [3423-07-2]

Obt. by reductive fission of Tetrandrine, T-286 or Isotetrandrine, I-328. Mp 134-136° Mp 176-177°. $[\alpha]_D^{22}$ +82 (c, 0.496 in EtOH).

O⁶,N-Di-Me, O⁷- α -D-xylopyranoside:**Latericine**

[93631-49-3]

C₂₃H₂₉NO₇ 431.485

Alkaloid from *Papaver latericine*, *Papaver pilosum* and *Papaver monanthum* (Papaveraceae). Needles + 1MeOH (MeOH). Mp 216-223°. $[\alpha]_D$ +93 (c, 0.555 in EtOH). $[\alpha]_D$ +40 (c, 0.521 in Py). Stereochem. incorrectly assigned in CAS.

O⁶,N,N-Tri-Me: (+)-Magnocurarine

[63428-12-6]

C₁₉H₂₄NO₃[⊕] 314.404

Alkaloid from *Gyrocarpus americanus* and *Lindera oldhamii* (Hernandiaceae, Lauraceae). Mp 199-200° dec.(as chloride). $[\alpha]_D^{25}$ +106 (c, 1 in H₂O).

O⁴,O⁷,N-Tri-Me: Roefractine

[131724-50-0]

C₁₉H₂₃NO₃ 313.396

Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D$ +73 (c, 0.14 in MeOH).

O,O,O,N-Tetra-Me, N-oxide: O-Methyl-roefractine N-oxide

[240122-94-5]

C₂₀H₂₅NO₄ 343.422Alkaloid from *Delphinium fangshanense*. [α]_D -2.3 (c, 1.7 in H₂O). λ_{max} 230 (log ε 4.4); 285 (log ε 3.61) (no solvent reported).

O,O,O,N,N-Penta-Me: [770645-80-2]

Mp 115° dec. (hydrate) (as iodide) Mp 139° dec (anhyd.) (iodide).

(±)-form [5843-65-2]

Alkaloid from *Aconitum japonicum*, *Asiasarum heterotropoides* and *Gnetum parvifolium* (Ranunculaceae, Aristolochiaceae, Gnetaceae).

Hydrochloride: Mp 265° (258° dec.).

Hydrobromide: Mp 269-270°.

N-Me: N-Methylhigenamine

[66277-20-1]

C₁₇H₁₉NO₃ 285.342Alkaloid from *Gnetum parvifolium*. Cryst. (EtOH). Mp 148-150°. λ_{max} 225 (sh) (log ε 4.6); 285 (log ε 4.08) (EtOH).O⁶-Me: [19442-69-4]Alkaloid from *Cocculus laurifolius*. Mp 220-221°.O⁷-Me: [2033-08-1]

Synthetic. Mp 135-136°.

O⁷-Me, hydrochloride: [17934-80-4]

Mp 215°.

O⁶,N-Di-Me: [1472-62-4]

Mp 161-162° (anhyd.).

(ξ)-form

4'-O-β-D-Glucopyranoside: [192324-89-3]

C₂₂H₂₇NO₈ 433.457Alkaloid from *Phoebe chekiangensis* (Lauraceae). [α]_D -2.3 (c, 0.4 in MeOH). λ_{max} 200 ; 222 ; 292 (no solvent reported).N-Me, O⁷-β-D-glucopyranoside: N-Methylhigenamine 7-O-β-D-glucopyranoside

[154418-17-4]

C₂₃H₂₉NO₈ 447.484Alkaloid from bark of *Phellodendron amurense*. Pale brownish prisms + 2½H₂O (Py aq.). Mp 158-159°. [α]_D -35.8 (c, 1.0 in Py).


[94997-19-0]

King, H. et al., *J.C.S.*, 1940, 737-746 (*Isococlaurine*)Finkelstein, J. et al., *J.A.C.S.*, 1951, 73, 550-553 (*Coclaurine, synth*)Tomita, M. et al., *Yakugaku Zasshi*, 1951, 71, 1069-1075; *CA*, 46, 5059 (*Magnocurarine*)McKenzie, A.K. et al., *Aust. J. Chem.*, 1953, 6, 180-185 (*Magnocurarine*)Arndt, R.R. et al., *J.C.S.*, 1963, 2547-2549 (*N-Methylcoclaurine*)Furukawa, H. et al., *Yakugaku Zasshi*, 1965, 85, 335-338; 472-475; *C.A.*, 63, 4351d; 5692d (*4'-Methyl-N-methylcoclaurine, Lotusine*)Preininger, V. et al., *Coll. Czech. Chem. Comm.*, 1966, 31, 3345-3352 (*Latericine*)Albonico, S.M. et al., *J.C.S.(C)*, 1966, 1340-1342 (*Colletine, Magnocurarine, uv, ord, abs config*)Johns, S.R. et al., *Aust. J. Chem.*, 1967, 20, 1729-1735 (*Coclaurine, isol, pmr, ms*)Sanchez, E. et al., *Tetrahedron*, 1967, 23, 1139-1143 (*Colletine*)Fridrichsons, J. et al., *Tetrahedron*, 1968, 24, 5785-5789 (*Coclaurine, cryst struct*)Koshiyama, H. et al., *Chem. Pharm. Bull.*, 1970, 18, 2564-2568 (*Demethylcoclaurine, isol, uv, ir, pmr, ms, struct, synth*)Bhakuni, D.S. et al., *Tetrahedron*, 1972, 28, 1093-1095 (*Isococlaurine, abs config*)Masaki, N. et al., *J.C.S. Perkin I*, 1977, 717-719 (*cryst struct, Demethylcoclaurine*)Torres, R. et al., *J. Nat. Prod.*, 1979, 42, 430-431 (*N-Demethylcolletine*)Saxena, N.K. et al., *J. Indian Chem. Soc.*, 1980, 57, 773-774 (*Lotusine, uv, ir, pmr*)Nishibe, S. et al., *J. Nat. Prod.*, 1986, 49, 547-548 (*4'-Methyl-N-methylcoclaurine*)Cortes, D. et al., *Phytochemistry*, 1986, 25, 2693-2695 (*Luxandrine*)Han, B.H. et al., *Pure Appl. Chem.*, 1989, 61, 443-448 (*isol, Coclaurine*)Gozler, B. et al., *J. Nat. Prod.*, 1990, 53, 666-668 (*Roefractine*)Ida, Y. et al., *Phytochemistry*, 1994, 35, 209-215 (*N-Methylhigenamine 7-glucoside*)Hegde, V.R. et al., *Biorg. Med. Chem. Lett.*, 1997, 7, 1207-1212 (*glucoside*)Xu, Q. et al., *J. Nat. Prod.*, 1999, 62, 1025-1027 (*N-Methylhigenamine, N-Methylhigenamine oxide*)Zhang, S. et al., *Phytochemistry*, 1999, 51, 333-336 (*O-Methylroefractine N-oxide*)Tshibangu, J.N. et al., *Phytochem. Anal.*, 2003, 14, 13-22 (*Isococlaurine, isol, hplc*)Yang, J. et al., *Magn. Reson. Chem.*, 2005, 43, 184-185 (*Lotusine, pmr, cmr*)

Himalayamine

H-276

[86702-42-3]


C₂₀H₁₇NO₆ 367.357Alkaloid from *Meconopsis villosa* (Papaveraceae). Cryst. (CHCl₃/petrol). Mp 218°. [α]_D²⁵ +137 (c, 0.185 in MeOH).Allais, D.P. et al., *Tet. Lett.*, 1983, 24, 2445 (*uv, cd, pmr, ms, struct*)

Himandridine

H-277

Himandreline

[15521-76-3]



Absolute Configuration

C₃₀H₃₇NO₇ 523.625Himandreline (originally assigned formula C₃₂H₄₁NO₇) was shown to be the polymorph, Mp 189-190°, of Himandridine. Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (preferred genus name *Galbulimima*) (Himantandraceae). Prisms(C₆H₆/hexane, MeOH aq. or EtOAc/petrol). Mp 189-190° Mp 204-205°. [α]_D¹⁸ -22 (c, 2.0 in CHCl₃). pK_a 7.7 (50% EtOH).

Hydroiodide:

Needles (EtOH/Et₂O). Mp 194-195°.

13-Ac: Alkaloid GB6

[15401-76-0]

C₃₂H₃₉NO₈ 565.662Alkaloid from the bark of *Himantandra baccata* (Himantandraceae).Prisms (Me₂CO), needles (Et₂O/hexane). Mp 154° Mp 170°. [α]_D¹⁹ -12 (c, 1.2 in CHCl₃).

20-Ac: Alkaloid GB7

[15549-97-0]

C₃₂H₃₉NO₈ 565.662Alkaloid from the bark of *Himantandra baccata* (Himantandraceae).Prisms (heptane). Mp 183°. [α]_D +35 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, 20-Ac: Alkaloid GB5

[15342-51-5]

C₂₄H₃₃NO₇ 447.527Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms(EtOAc). Mp 234°. [α]_D +60 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, 13,20-di-Ac: Alkaloid GB3

[15488-89-8]

C₂₆H₃₅NO₈ 489.564Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms(EtOAc). Mp 237°. [α]_D +97 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, tetra-Ac: Alkaloid GB2

[15373-50-9]

C₃₀H₃₉NO₁₀ 573.639Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms(EtOAc/cyclohexane). Mp 238°. [α]_D +38 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, 13-benzoyl, 20-Ac: Alkaloid GB4

[15342-50-4]

C₃₁H₃₇NO₈ 551.635Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms(MeOH/CHCl₃). Mp 238°. [α]_D +63 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, 14-benzoyl, 13,16-di-Ac: Alkaloid GB1. GB 1

[15342-91-3]

C₃₃H₃₉NO₉ 593.672Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms(MeOH/CHCl₃). Mp 242°. [α]_D +8 (c, ca. 1 in CHCl₃).

O-De-Me, O-debenzoyl, 14-benzoyl, tri-Ac: Himbosine

[15448-14-3]

C₃₅H₄₁NO₁₀ 635.71Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms

(Himantandraceae). Prisms

(MeOH/CHCl₃). Mp 262°. [α]_D¹⁵ +55 (c, 0.89 in CHCl₃). [α]_D²⁵ +28 (c, 1.1 in CHCl₃).

O-De-Me, O-debenzoyl, 14-benzoyl, tri-Ac, hydrobromide:
Needles (EtOH). Mp 228°.

13-Deoxy: Himandrine

[15610-97-6]

C₃₀H₃₇NO₆ 507.625

Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Prisms (EtOH) or glistening leaflets (EtOH aq.). Mp 185-186°. [α]_D²⁰ -38 (c, 1.22 in CHCl₃).

►MR7000000

13-Deoxy, hydrochloride: Mp 222° dec.

13-Deoxy, O-debenzoyl: Alkaloid GB8.

Debenzoylhimandrine

[15401-61-3]

C₂₃H₃₃NO₅ 403.517

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms (cyclohexane or Et₂O/hexane). Mp 188° (183-185°). [α]_D²⁵ +65 (c, 2.1 in CHCl₃).

13-Deoxy, O-debenzoyl, 16-Ac: Alkaloid GB9

[15342-52-6]

C₂₅H₃₅NO₆ 445.555

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms (heptane). Mp 168°. [α]_D +23 (c, ca. 1 in CHCl₃).

13-Deoxy, O-debenzoyl, 16,20-di-Ac: Alkaloid GB10

[15423-42-4]

C₂₇H₃₇NO₇ 487.592

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms (EtOAc/cyclohexane). Mp 162°. [α]_D +19 (c, ca. 1 in CHCl₃).

13-Deoxy, O-de-Me, O-debenzoyl, 20-Ac: Alkaloid GB11

[15342-53-7]

C₂₄H₃₃NO₆ 431.528

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms (EtOAc). Mp 239°. [α]_D +64 (c, ca. 1 in CHCl₃).

13-Deoxy, O-de-Me, O-debenzoyl, tri-Ac: Alkaloid GB12

[15342-54-8]

C₂₈H₃₇NO₈ 515.602

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Prisms (EtOAc/cyclohexane). Mp 222°. [α]_D +23 (c, ca. 1 in CHCl₃).

Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 283-287 (*isol*)

Pinhey, J.T. *et al.*, *Aust. J. Chem.*, 1961, **14**, 106-134 (*nomencl*)

Lovell, F.M. *et al.*, *Proc. Chem. Soc., London*, 1964, 58 (*Himbosine, cryst struct*)

Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*GB alkaloids, isol*)

Mander, L.N. *et al.*, *Aust. J. Chem.*, 1967, **20**, 981-1019; 1021-1027 (*Himandridine, Himbosine, uv, ir, pmr, struct*)


Guise, G.B. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1029-1035 (*uv, ir, pmr, struct, synth*)

Willis, A.C. *et al.*, *Aust. J. Chem.*, 2006, **59**, 629-632 (*abs config, Himbosine, cryst struct*)

Himanimide A

[464895-13-4]

H-278



C₂₂H₂₁NO₃ 347.413

Prod. by the basidiomycete *Serpula himantoides*. Yellow oil. λ_{\max} 230 (log ϵ 4.1); 356 (log ϵ 3.6) (MeOH).

N-Hydroxy: Himanimide C

[464189-92-2]

C₂₂H₂₁NO₄ 363.412

Prod. by *Serpula himantoides*. Antifungal and cytotoxic agent. Yellow oil. λ_{\max} 232 (log ϵ 4.3); 290 (log ϵ 3.7); 374 (log ϵ 3.6) (MeOH).

3,4-Dihydro, N-hydroxy: Himanimide D

[464194-69-2]

C₂₂H₂₃NO₄ 365.428

Prod. by *Serpula himantoides*. Oil. Possesses *cis*-config. Racemate. λ_{\max} 225 (sh) (log ϵ 4.1); 277 (log ϵ 3.4) (MeOH).

2'',3''-Dihydro, 2'' ζ ,3''-dihydroxy: Himanimide B

[464194-68-1]

C₂₂H₂₃NO₅ 381.427

Prod. by *Serpula himantoides*. Yellow oil. λ_{\max} 229 (log ϵ 4.3); 356 (log ϵ 3.8) (MeOH).

Aqueveque, P. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 257-262 (*isol, pmr, cmr, activity*)

Sellés, P. *et al.*, *Org. Lett.*, 2005, **7**, 605-608 (*Himanimide C, synth*)

Cheng, C.-F. *et al.*, *Tetrahedron*, 2008, **64**, 4347-4353 (*Himanimide D, synth*)

Himanthine

H-279

C₃₇H₄₀N₂O₆ 608.733


Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the stem bark of *Berberis himalaica* (Berberidaceae). Needles (MeOH). Mp 206-207°. [α]_D³³ -198.7.

Chatterjee, R. *et al.*, *J. Indian Chem. Soc.*, 1952, **29**, 921-924 (*isol, uv*)

Himbacine

H-280

4-[2-(1,6-Dimethyl-2-piperidinyl)ethynyl]decahydro-3-methylnaphtho[2,3-c]furan-1(3H)-one, 9CI
[6879-74-9]



Absolute Configuration

C₂₂H₃₅NO₂ 345.524

Alkaloid from the bark of *Himantandra*

baccata and *Himantandra belgraveana* (preferred genus name *Galbulimima*) (Himantandraceae). Muscarine M₄-receptor antagonist, research tool for differentiating between muscarinic receptor sites. Antispasmodic agent. Needles (heptane). Mp 132°. [α]_D¹⁴ +63 (c, 1.04 in CHCl₃). pK_a 9.3 (50% EtOH). Log P 4.36 (uncertain value) (calc).

►QL0843500

Hydrochloride:

Cryst. + 1H₂O (Me₂CO/Et₂O). Mp 267°.

N-Oxide:

Needles (EtOAc). Mp 192-194° dec. Hygroscopic. Mp depends on rate of heating.

N-De-Me: Himbeline

[518-99-0]

C₂₁H₃₃NO₂ 331.497

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Needles (petrol). Mp 100°. [α]_D¹⁸ +19 (c, 2.4 in CHCl₃).

N-De-Me; hydrochloride:

Needles + 1H₂O (EtOH). Mp 265-266°.

N-De-Me, N-Ac:

Needles (petrol). Mp 159°.

10,10a-Didehydro: Himgravine

[6879-75-0]

C₂₂H₃₃NO₂ 343.508

Alkaloid from the bark of *Himantandra baccata* and *Himantandra belgraveana* (Himantandraceae). Needles (petrol). Mp 120°. [α]_D¹⁸ +47 (c, 1.5 in CHCl₃). pK_a 9.3 (50% EtOH).

2'-Epimer, N-de-Me: Himandravine

[6879-73-8]

C₂₁H₃₃NO₂ 331.497

Alkaloid from the bark of *Himantandra belgraveana* (Himantandraceae). Needles (hexane). Mp 119°. [α]_D¹⁸ +23 (c, 1.89 in CHCl₃).

2'-Epimer, N-de-Me, hydrochloride:

Prisms (Me₂CO/EtOH). Mp 238-240°.

Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 283-287 (*Himbeline, Himandravine, Himgravine*)

Abraham, R.J. *et al.*, *Aust. J. Chem.*, 1961, **14**, 64-68 (*Himgravine, struct*)

Pinhey, J.T. *et al.*, *Aust. J. Chem.*, 1961, **14**, 106-134 (*Himbeline, Himandravine, Himgravine, ir, struct*)

Fridrichsons, J. *et al.*, *Acta Cryst.*, 1962, **15**, 119-128 (*cryst struct*)

Galbraith, M.N. *et al.*, *Aust. J. Chem.*, 1963, **16**, 112-122 (*synth, derivis*)

Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*Himbeline, Himandravine, Himgravine, isol*)

Hart, D.J. *et al.*, *J.O.C.*, 1997, **62**, 5023-5033 (*Himbeline, synth, pmr, cmr*)

Chackalamannil, S. *et al.*, *J.O.C.*, 1999, **64**, 1932-1940 (*Himbeline, synth*)

Chackalamannil, S. *et al.*, *Org. Lett.*, 2001, **3**, 1427-1429 (*Himandravine, synth*)

Takadoi, M. *et al.*, *Tetrahedron*, 2002, **58**, 9903-9923 (*synth*)


Takadoi, M. *et al.*, *Bioorg. Med. Chem.*, 2003, **11**, 1169-1186 (*bibl, pharmacol*)

Tchabanenko, K. *et al.*, *Tetrahedron*, 2005, **61**, 11649-11656 (*synth*)

Willis, A.C. *et al.*, *Aust. J. Chem.*, 2006, **59**, 629-632 (*abs config*)

Himbadine

[24932-15-8]

Absolute
ConfigurationC₂₁H₃₁NO₂ 329.481

Alkaloid from the bark of *Himantandra baccata* (preferred genus name *Galbulimima*) (Himantandraceae). Cryst. + ½ H₂O (MeOH aq.). Mp 114-115°. [α]_D²⁵ -42 (c, 1.43 in CHCl₃). pK_a 8 (50% MeOH). Anhyd. material is a gum.

Hydrochloride:Prisms (EtOH/Et₂O). Mp 233-235°.**Oxime:**Buff-coloured needles (Me₂CO). Mp 200-204°.**2,4-Dinitrophenylhydrazone:**Orange needles (C₆H₆/petrol). Mp 148°.**Ac:**Cryst. (MeOH). Mp 120°. [α]_D²⁰ -27 (c, 1.51 in CHCl₃). Subl. at 0.1mm.**N-De-Me: Alkaloid GB13. GB 13**


[16625-34-6]

C₂₀H₂₉NO₂ 315.455

Alkaloid from the bark of *Himantandra baccata* (Himantandraceae). Prisms (heptane). Mp 185°. [α]_D -84 (c, ca. 1 in CHCl₃).

N-De-Me; hydrochloride:Prisms (EtOH/Et₂O). Mp 275-277°.Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 283-287 (*isol*)Pinhey, J.T. *et al.*, *Aust. J. Chem.*, 1961, **14**, 106-134 (*nomencl*)Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*GB13, isol*)Mander, L.N. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1473-1491 (*Himbadine, GB13, uv, ir, pmr, struct*)Mander, L.N. *et al.*, *J.A.C.S.*, 2003, **125**, 2400-2401 (*GB13, synth*)Willis, A.C. *et al.*, *Aust. J. Chem.*, 2006, **59**, 629-632 (*abs config*)Movassaghi, M. *et al.*, *J.A.C.S.*, 2006, **128**, 8126-8127 (*GB13, synth, stereochem*)Evans, D.A. *et al.*, *J.A.C.S.*, 2007, **129**, 1048-1049 (*GB 13, synth*)**Himeic acid A**

[838847-67-9]



X = O

C₂₂H₂₉NO₈ 435.473**H-281**

Prod. by a marine-derived *Aspergillus* sp. Ubiquitin-activating enzyme inhibitor. [α]_D²⁶ -15 (c, 0.14 in MeOH). λ_{max} 234 (log ε 4.5); 254 (log ε 4.4) (MeOH).

N-Deacyl: 11-[5-(Aminocarbonyl)-4-oxo-4H-pyran-2-yl]-10-undecenoic acid. Himeic acid B

[838847-69-1]

C₁₇H₂₃NO₅ 321.372

Isol. from a marine-derived *Aspergillus* sp. λ_{max} 227 (log ε 4.3); 266 (log ε 4.1) (MeOH).

Tsukamoto, S. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 191-194 (*isol, pmr, cmr*)**Himeic acid C**

[838847-71-5]

As Himeic acid A, H-282 with


X = NH

C₂₂H₃₀N₂O₇ 434.488


Prod. by a marine-derived *Aspergillus* sp. [α]_D²⁶ -9.8 (c, 0.32 in MeOH). λ_{max} 252 (log ε 4.4) (MeOH).

Tsukamoto, S. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 191-194 (*isol, pmr, cmr*)**H-283****Himeradine A**

[552831-75-1]

C₂₉H₄₅N₃OAlkaloid from *Lycopodium chinense*.Morita, H. *et al.*, *J.O.C.*, 2003, **68**, 4563-4566 (*isol, pmr, cmr*)**H-284****Himgaline**

[16585-08-3]

Absolute
ConfigurationC₂₀H₃₁NO₂ 317.47

Alkaloid from bark of *Himantandra baccata* and *Himantandra belgraveana* (preferred genus name *Galbulimima*) (Himantandraceae). Needles (C₆H₆). Mp 226°. [α]_D -76 (c, 1 in CHCl₃). pK_a 11 (50% MeOH aq.).

Hydrochloride:Needles (MeOH/Et₂O). Mp 311°.**Picrate:**

Yellow needles (MeOH aq.). Mp 136-137°.

Di-Ac:**H-285**

Prisms (MeCN). Mp 143-145°. [α]_D²⁰ -95 (c, 3.1 in CHCl₃).

Binns, S.V. *et al.*, *Aust. J. Chem.*, 1965, **18**, 569-573 (*isol*)Mander, L.N. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1705-1718 (*ir, pmr, struct*)Willis, A.C. *et al.*, *Aust. J. Chem.*, 2006, **59**, 629-632 (*cryst struct, abs config*)Shah, U. *et al.*, *J.A.C.S.*, 2006, **128**, 12654-12655 (*synth*)Evans, D.A. *et al.*, *J.A.C.S.*, 2007, **129**, 1048-1049 (*synth*)**Himgrine****H-286**

Struct. unknown

C₂₂H₃₃NO₃ 359.508


Alkaloid from the bark of *Himantandra belgraveana* (preferred genus name *Galbulimima*) (Himantandraceae). Noncryst.

Perchlorate:Needles + 1H₂O (EtOH aq.). Mp 143-144°.**Methodide:**

Needles (EtOH). Mp 222-223°.

Brown, R.F.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 283-287 (*isol*)**Hinckdentine A****H-287**

[112663-91-9]


Absolute
ConfigurationC₁₉H₁₄Br₃N₃O 540.051

Alkaloid from the marine bryozoan *Hincksinoflustra denticulata*. Mp 196-197°. [α]_D²⁵ +274 (solvent unspecified).

Blackman, A.J. *et al.*, *Tet. Lett.*, 1987, **28**, 5561-5562 (*isol, uv, ir, pmr, ms, struct, abs config*)**Hinnuliquinone****H-288**

2,5-Bis[2-(1,1-dimethyl-2-propenyl)-1H-indol-3-yl]-3,6-dihydroxy-2,5-cyclohexadiene-1,4-dione

[78860-37-4]

C₃₂H₃₀N₂O₄ 506.6

Isol. from *Nodulisporium hinnuleum*. Inhibitor of HIV-1 protease. Dark red cryst. Poorly sol. hexane. Mp 243°. Isomeric with Cochliodinol, C-547. λ_{max} 280 (ε 26500); 288 (ε 17400); 506 (ε 950) (MeOH). λ_{max} 280 (ε 18600); 288 (ε 19500); 322 (ε 20400); 516 (ε 1470) (EtOH/NaOH).

Di-Ac:

Purple rods (Ac₂O). Mp 240°.


O'Leary, M.A. *et al.*, *J.C.S. Perkin I*, 1984, 567-569 (isol, uv, ir, pmr, ms, biosynth)
 Singh, S.B. *et al.*, *Biochem. Biophys. Res. Commun.*, 2004, **324**, 108-113 (isol, activity)

Hippacine† **H-289**C₁₇H₂₁NO₅ 319.357Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Hippeastrum vittatum* (Amaryllidaceae). Mp 245-246°.

Döpke, W. *et al.*, *Pharmazie*, 1966, **21**, 323-324;
CA, **65**, 10632e (isol)

Hippadine **H-290**

7H-[1,3]Dioxolo[4,5-j]pyrrolo[3,2,1-de]phenanthridin-7-one, 9CI. Pratorine. Alkaloid N3
 [52886-06-3]

C₁₆H₉NO₃ 263.252

Identity of Alkaloid N3 with Hippadine is not certain. Alkaloid from *Hippeastrum vittatum*, *Crinum pratense*, *Crinum bulbispermum*, *Crinum asiaticum* var. *japonicum*, *Crinum latifolium* and *Hae-manthus kalbreyeri* and (Alkaloid N3) from *Lycoris sanguinea* (Amaryllidaceae). Mp 209-210°. λ_{max} 230 (log ε 4.25); 238 (log ε 4.29); 248 (log ε 4.42); 255 (log ε 4.26); 300 (log ε 4.21); 308 (sh) (log ε 4.03); 335 (log ε 3.97); 350 (log ε 3.99); 368 (log ε 3.87) (MeOH).

▶ JI5025000

4,5-Dihydro: Anhydrolycorin-7-one

[40360-71-2]

C₁₆H₁₁NO₃ 265.268

Alkaloid from the bulbs of *Crinum pratense* (Amaryllidaceae). Pale yellow prisms (CHCl₃/MeOH). Mp 232-234° (228-230°). λ_{max} 242 (ε 30000); 274 (ε 13000); 342 (ε 4000) (95% EtOH).

Deoxo: 7H-[1,3]Dioxolo[4,5-j]pyrrolo[3,2,1-de]phenanthridine, 8,9-Methylenedioxy-11H-pyrrolo[3,2,1-de]phenanthridine, **4,5-Dehydroanhydrolycorine**

[212-38-4]

C₁₆H₁₁NO₂ 249.268

Alkaloid from the bulbs of *Pancreatium biflorum* (Amaryllidaceae) infected with *Imperata cylindrica*. Mp 154-156°.

Deoxo, picrate:

Reddish-brown needles. Mp 143-145° dec.

4-Hydroxy, deoxo, 4,5-dihydro: 4-Hydroxyanhydrolycorine

[136850-09-4]

C₁₆H₁₃NO₃ 267.284

Alkaloid from leaves of *Hymenocallis caymanensis* (Amaryllidaceae).

8-Hydroxy: Kalbretorine

[98899-99-1]

C₁₆H₉NO₄ 279.251

Alkaloid from the roots of *Hae-manthus kalbreyeri* (Amaryllidaceae). Mp 245-248° dec. λ_{max} 225 (log ε 4.32); 234 (log ε 4.02); 248 (log ε 4.31); 255 (log ε 4.11); 292 (log ε 3.78); 335 (log ε 3.24); 355 (log ε 3.48) (MeOH). λ_{max} 222 (log ε 4.27); 250 (log ε 4.29); 260 (log ε 4.2); 298 (log ε 3.66); 348 (log ε 3.33); 375 (log ε 3.4) (MeOH/NaOMe).

8-Methoxy:

Brown microcryst. (petrol/Me₂CO). Mp 258-260°.

Cook, J.W. *et al.*, *J.C.S.*, 1954, 4176 (deoxo, synth)

Humber, L.G. *et al.*, *J.C.S.*, 1954, 4622-4630 (synth)

Hara, H. *et al.*, *Tet. Lett.*, 1972, 5031-5033 (synth)

Takagi, S. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 617-622; *CA*, **81**, 74924y (isol)

El-Moghazi, A.M. *et al.*, *Planta Med.*, 1976, **29**, 156-159 (isol)

Ghosal, S. *et al.*, *Phytochemistry*, 1981, **20**, 2003-2007; 1983, **22**, 2305-2309; 1985, **24**, 1825-1828; 1986, **25**, 1097 (*Hippadine*, *Kalbretorine*, *Anhydrolycorin-7-one*, *4,5-Dehydroanhydrolycorine*)

Ali, A.A. *et al.*, *Planta Med.*, 1981, **43**, 407 (pmr, cmr, ms, struct)

Hayakawa, K. *et al.*, *Tet. Lett.*, 1987, **28**, 5895-5898 (synth)

Trimino, Z. *et al.*, *Rev. Cubana Quim.*, 1989, **5**, 55-57; *CA*, **115**, 203303s (*4-Hydroxyanhydrolycorine*)

Siddiqui, M.A. *et al.*, *Tet. Lett.*, 1990, **31**, 1523-1526 (synth)

Lauk, U. *et al.*, *Tet. Lett.*, 1991, **32**, 65-68 (synth)

Grigg, R. *et al.*, *Tet. Lett.*, 1991, **32**, 3859-3862 (synth)

Sakamoto, T. *et al.*, *Heterocycles*, 1993, **36**, 2597-2600 (synth)

Iwao, M. *et al.*, *Heterocycles*, 1994, **38**, 1717-1720 (*Hippadine*, *Anhydrolycorinone*, *Kalbretorine*, synth)

Banwell, M.G. *et al.*, *Chem. Comm.*, 1995, 2551-2553 (synth)

Shao, H.W. *et al.*, *Chin. Chem. Lett.*, 1996, **7**, 13-14; *CA*, **124**, 289955b (synth)

Hutchings, R.H. *et al.*, *J.O.C.*, 1996, **61**, 1004-1013 (*Hippadine*, *Kalbretorine*, synth)

Padwa, A. *et al.*, *J.O.C.*, 1998, **63**, 3986-3997; 1999, **64**, 3595-3607 (synth)

Harrowen, D.C. *et al.*, *Synthesis*, 1999, 1300-1302 (synth, ir, uv, pmr, cmr, ms)

Miki, Y. *et al.*, *Tet. Lett.*, 1999, **40**, 4347-4348 (synth)

Boger, D.L. *et al.*, *J.O.C.*, 2000, **65**, 9120-9124 (synth)

Campbell, W.E. *et al.*, *Phytochemistry*, 2000, **53**, 587-591 (cmr)

Harayama, T. *et al.*, *Tetrahedron*, 2004, **60**, 1611-1616 (*Anhydrolycorinone*, synth)

Ganton, M.D. *et al.*, *Org. Lett.*, 2005, **7**, 4777-4779 (synth)

Mentzel, U.V. *et al.*, *J.O.C.*, 2006, **71**, 5807-5810 (synth)

Hippandrine **H-291**C₁₇H₁₉NO₄ 301.341

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Hippeastrum brachyandrum* (Amaryllidaceae). Prisms (Me₂CO). Mp 194°. [α]_D²⁵ -110 (c, 0.2 in CHCl₃).

Picrate:

Cryst. (MeOH). Mp 179° dec.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1959, **92**, 2582-2584 (isol)

Hippauline **H-292**

[1357-53-5]

C₁₇H₁₉NO₄ 301.341


Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Hippeastrum aulicum* var. *robustum* (Amaryllidaceae). Mp 163-164°. [α]_D²⁵ +10 (c, 0.2 in CHCl₃).

Perchlorate: Mp 175° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109 (isol)

Hippeastidine **H-293**

[66276-51-5]

C₁₈H₂₅NO₄ 319.4

Alkaloid from the bulbs of *Hippeastrum ananuca* (Amaryllidaceae).

▶ KM5830000

O³-De-Me: O³-Demethylhippeastidine
[81904-08-7]C₁₇H₂₃NO₄ 305.373

Alkaloid from *Hippeastrum ananuca* bulbs (Amaryllidaceae). Mp 175°. Alkaloid not named in the abstr.


Watson, W.H. *et al.*, *Cryst. Struct. Commun.*, 1977, **6**, 797 (cryst struct)

Pacheco, P. *et al.*, *Rev. Latinoam. Quim.*, 1978, **9**, 28; *CA*, **89**, 103746q

Pacheco, P. *et al.*, *Bol. Soc. Chil. Quim.*, 1982, **27**, 289; *CA*, **96**, 214311h (deriv)

Hippeastrine **H-294**

Trisphaerine. Trispherine
[477-17-8]

C₁₇H₁₇NO₅ 315.325

Alkaloid from *Amaryllis belladonna*, *Hippeastrum vittatum* and many other spp. in the Amaryllidaceae. Mp 214-215°. [α]_D²⁵ +160 (c, 0.3 in CHCl₃). λ_{max} 236 (log ε 3.4); 268 (log ε 2.7); 308 (log ε 2.6) (MeOH).

Picrate: Mp 256°.**Methodide:** Mp 238°.**N-Oxide: Hippeastrine N-oxide**

[137810-41-4]

C₁₇H₁₇NO₆ 331.324

Alkaloid from the flowers of *Lycoris radiata* (Amaryllidaceae). Cubes (EtOAc). Mp 172-173°. [α]_D²⁵ +81.7 (c,

0.23 in MeOH). Data refers to synthetic compd.

Me ether: Ungerine. Nivaline†

[477-13-4]

C₁₈H₁₉NO₅ 329.352

Alkaloid from *Ungernia severtzovii*, *Ungernia trisphaera* and *Leucocjum aestivum* (Amaryllidaceae). Mp 135-136°. [α]_D +103.

Me ether, hydrochloride: Mp 270-271°.

Yunusov, S. *et al.*, *Dokl. Akad. Nauk SSSR*, 1953, **6**, 44; *CA*, **49**, 1281b (*Ungerine*)

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129; 2093 (*isol, ir*)

Briggs, C.K. *et al.*, *J.A.C.S.*, 1956, **78**, 2899 (*Ungerine*)

Kitagawa, T. *et al.*, *J.C.S.*, 1959, 3741 (*struct, config*)

Proskurnina, N.F. *et al.*, *CA*, 1963, **59**, 14035h (*deriv*)

Abduazimov, K.A. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1963, **153**, 1315; *CA*, **60**, 9324e (*Ungerine*)

Hawthorn, W.A. *et al.*, *J.C.S.*, 1965, 1991 (*pmr*)

Katakawa, J. *et al.*, *Heterocycles*, 1984, **22**, 2213 (*synth*)

Kihara, M. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1849 (*Hippeastrine N-oxide*)


Astley, S.T. *et al.*, *Tet. Lett.*, 1993, **34**, 2035 (*synth*)

Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591 (*cd*)

Evidente, A. *et al.*, *Phytochemistry*, 2004, **65**, 2113-2118 (*isol, cd, pmr, cmr, ms*)

Hippocasin H-295

1,2,3,3a,4,5,6,6a,7,9a-Decahydro-8-methylpyrido[2,1,6-de]quinolizine, 9CI



Absolute Configuration

C₁₃H₂₁N 191.316

Natural-form [60022-25-5]

Alkaloid from the defence secretion of the ladybug *Hippodamia caseyi*. Oil.

N-Oxide: Hippocasin N-oxide

[60022-26-6]

C₁₃H₂₁NO 207.315

Constit. of *Hippodamia caseyi*. Oil.

N-Oxide; hydrochloride: [60047-14-5]

Cryst. (EtOAc/MeOH). [α]_D²⁵ +14.8 (c, 0.73 in MeOH). Dec. >220° without melting.

3a-Epimer, N-oxide: 2-Dehydrococcelline

C₁₃H₂₁NO 207.315

Constit. of the defence secretion of *Anatis ocellata*. Oil. [α]_D²⁵ +8 (c, 0.16 in CH₂Cl₂).

(±)-form [75556-13-7]

Synthetic. Oil; yellow needles (EtOH)(as picrate). Mp 137.5-139° (picrate).

N-Oxide: [75556-12-6]

Synthetic. Oil; feathery cryst. (EtOAc/MeOH)(as hydrochloride). Mp 210° dec. (hydrochloride).


Ayer, W.A. *et al.*, *Can. J. Chem.*, 1976, **54**, 1807 (*isol, ir, ms, synth, oxide, ord, pmr, cryst struct*)

Mueller, R.M. *et al.*, *J.O.C.*, 1984, **49**, 2217 (*synth, ir, pmr, ms*)

Lebrun, B. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1148-1149 (*Dehydrococcelline*)

Hippocrateine III H-296

[169626-41-9]




C₄₆H₅₆N₂O₁₉ 940.95

Alkaloid from root and stem barks of *Hippocratea excelsa* (Celastraceae). Gum. Note differing regiochemistry of nicotinoyl group compared with Per(-deacyl)emarginatine, P-246.

Calzada, F. *et al.*, *Phytochemistry*, 1995, **40**, 583 (*isol, uv, ir, pmr, cmr, struct*)

Hippospongine C H-297

[184301-92-6]




C₂₇H₃₄N₂O₄ 450.577

Constit. of *Hippospongia* sp. Pale yellow oil. λ_{max} 206 (log ε 4.19); 276 (log ε 4.39); 287 (log ε 4.39); 303 (log ε 4.34) (EtOH). λ_{max} 206 (ε 13800); 276 (ε 24000); 287 (ε 24000); 303 (ε 21900) (MeOH) (Berdy).

Rochfort, S.J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1024-1028 (*isol, pmr, cmr, ms*)

Hippospongine E H-298

[184301-94-8]



C₂₄H₃₀N₂O₄ 410.512

Constit. of a *Hippospongia* sp. Pale yellow oil. λ_{max} 279 (log ε 3.95) (EtOH). λ_{max} 279 (ε 8900) (MeOH) (Berdy).

ω-N-Ac: Hippospongine F

[184301-95-9]

C₂₆H₃₂N₂O₅ 452.549


Constit. of a *Hippospongia* sp. Pale yellow oil. λ_{max} 265 (log ε 4.02) (EtOH). λ_{max} 265 (ε 10450) (MeOH) (Berdy).

Rochfort, S.J. *et al.*, *J. Nat. Prod.*, 1996, **59**,

1024 (*isol, pmr, cmr, ms*)

Hirsutellone A H-299

[857672-62-9]



Relative Configuration

R = H


C₂₈H₃₁NO₄ 445.557

Related to Pyrrocidine B, P-940 and Antibiotic GKK 1032B, A-1183. Alkaloid from the insect pathogenic fungus *Hirsutella nivea* BCC 2594. Cryst. Mp 155-157°. [α]_D²⁰ +168 (c, 0.2 in MeOH). λ_{max} 203 (log ε 4.42); 228 (sh) (log ε 3.72); 281 (sh) (log ε 2.99) (MeOH).

Isaka, M. *et al.*, *Tetrahedron*, 2005, **61**, 5577-5583 (*isol, pmr, cmr, cryst struct*)

Hirsutellone B H-300

[857672-63-0]



Relative Configuration

R = H

C₂₈H₃₃NO₄ 447.573

Related to Pyrrocidine B, P-940 and Antibiotic GKK 1032A₂, A-1182. Alkaloid from the insect pathogenic fungus *Hirsutella nivea* BCC 2594. Amorph. solid. Mp 261-263°. [α]_D²⁷ +256 (c, 0.2 in MeOH). λ_{max} 203 (log ε 4.37); 227 (sh) (log ε 3.78); 276 (sh) (log ε 2.99) (MeOH).

3'β,4'β-Epoxy: Hirsutellone C

[857672-64-1]

C₂₈H₃₁NO₅ 461.557

Alkaloid from *Hirsutella nivea* BCC 2594. Amorph. solid. Mp 234-235° dec. [α]_D²⁰ +129 (c, 0.2 in MeOH). λ_{max} 203 (log ε 4.42); 230 (log ε 3.93); 277 (log ε 3.12) (MeOH).

Isaka, M. *et al.*, *Tetrahedron*, 2005, **61**, 5577-5583 (*isol, pmr, cmr*)

Hirsutellone D H-301

[857672-65-2]

As Hirsutellone A, H-299 with

R = CH₃

C₂₉H₃₃NO₄ 459.584

Related to Pyrrocidine B, P-940 and Antibiotic GKK 1032B, A-1183. Alkaloid from *Hirsutella nivea* BCC 2594.

Amorph. solid. Mp 106-109°. $[\alpha]_D^{29} +214$ (c, 0.07 in MeOH). λ_{\max} 204 (log ϵ 4.46); 229 (sh) (log ϵ 3.78); 283 (sh) (log ϵ 3.07) (MeOH).

Isaka, M. *et al.*, *Tetrahedron*, 2005, **61**, 5577-5583 (*isol*, *pmr*, *cmr*)

Hirsutellone E**H-302**

[857672-66-3]

As Hirsutellone B, H-300 with


R = CH₃C₂₉H₃₅NO₄ 461.6

Related to Pyrrocidine B, P-940 and Antibiotic GKK 1032A₂, A-1182. Alkaloid from *Hirsutella nivea* BCC 2594. Amorph. solid. λ_{\max} 227 (sh) ; 276 (MeOH).

Isaka, M. *et al.*, *Tetrahedron*, 2005, **61**, 5577-5583 (*isol*, *pmr*, *cmr*)

Hirsutellone F**H-303**

[901129-86-0]


C₅₆H₆₂N₂O₈ 891.114Prod. by *Trichoderma* sp. BCC 7579.

Antitubercular agent. Solid. Mp 235-237°. $[\alpha]_D^{25} +169$ (c, 0.25 in CHCl₃). λ_{\max} 203 (log ϵ 4.6); 228 (sh) (log ϵ 4.03); 285 (sh) (log ϵ 3.41) (MeOH).

Isaka, M. *et al.*, *Org. Lett.*, 2006, **8**, 2815-2817 (*isol*, *pmr*, *cmr*)

Hirudinoidine A**H-304**

[1021944-70-6]

C₁₁H₁₀N₄O₃S₂ 310.357

Alkaloid from the leech *Hirudo nipponica*. Yellow prisms (CHCl₃/MeOH). λ_{\max} 227 (log ϵ 1.03); 250 (log ϵ 1.68); 254 (log ϵ 1.68); 365 (log ϵ 0.55) (MeOH).

N³-De-Me: Hirudinoidine B

[1021944-72-8]

C₁₀H₈N₄O₃S₂ 296.33

Alkaloid from *Hirudo nipponica*. Orange-brown powder. λ_{\max} 248 (log ϵ 1.2); 363 (log ϵ 0.45) (MeOH).

N¹,N³-Di-de-Me: Hirudinoidine C

[1021944-73-9]

C₉H₆N₄O₃S₂ 282.303

Alkaloid from *Hirudo nipponica*. Orange-brown powder. λ_{\max} 244 (log ϵ 1.12); 358 (log ϵ 0.39); 362 (log ϵ 0.4) (MeOH).

Li, Y.-B. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 303-307 (*isol*, *pmr*, *cmr*, *ms*, *cryst struct*)

Hirudonine**H-305**

3-[[[4-[(Aminoiminomethyl)amino]butyl]amino]propyl]guanidine, 9CI. 4-[[[3-Guanidinopropyl]amino]butyl]guanidine, 8CI. 4-Aza-1,8-diguaminooctane. Diaminospermidine. Hyrudonine [2465-97-6]

HN=C(NH₂)NH(CH₂)₄NH(CH₂)₃NHC(NH₂)=NHC₉H₂₃N₇ 229.328

Alkaloid from *Hirudo medicinalis* (medical leech). Mp 169° (as hydrated sulfate salt).

Picrate:

Yellow platelets. Mp 226-228°.

Robin, Y. *et al.*, *Biochim. Biophys. Acta*, 1957, **24**, 381 (*isol*)


Robin, Y. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1961, **252**, 1224 (*isol*, *struct*, *synth*)

Robin, Y. *et al.*, *Comp. Biochem. Physiol.*, 1967, **22**, 787; *C.A.*, **67**, 98070d (*biosynth*)

Mitchinson, A. *et al.*, *Chem. Comm.*, 1994, 2613 (*synth*)

Hispidospermidine**H-306**

[154648-89-2]



Absolute Configuration

C₂₅H₄₇N₃O 405.666Prod. by *Chaetosphaeronomia hispidulum*.

Phospholipase C inhibitor. Oil. Sol. DMSO, acids, MeOH, CH₂Cl₂; poorly sol. H₂O. $[\alpha]_D^{24} -60$ (c, 1.45 in CHCl₃). λ_{\max} (MeOH) (Berdy).

Sakai, A. *et al.*, *J. Antibiot.*, 1994, **47**, 1-5; 6-15 (*isol*, *ir*, *pmr*, *cmr*)

Frontier, A.J. *et al.*, *J.A.C.S.*, 1997, **119**, 6686-6687 (*synth*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1998, **120**, 4039-4040 (*synth*)

Frontier, A.J. *et al.*, *J.A.C.S.*, 2000, **122**, 6151-6159 (*synth*)


Tamiya, J. *et al.*, *Tetrahedron*, 2003, **59**, 6921-6932 (*synth*)

Histamine**H-307**

1H-Imidazole-4-ethanamine, 9CI. 4-(2-Aminoethyl)-1H-imidazole. Ergamine.

Scombrotosin

[51-45-6]

C₅H₉N₃ 111.146

1H-form predominates ca. 4:1 in aq. soln. At physiological pH exists as a mixt. of base: monocation: dication in ratio 1:96:3 with both base and monocation capable of 1H \rightleftharpoons 3H tautom. Found in *Claviceps purpurea*, *Veronica fistularis*, *Furcellaria lumbicalis*, *Anemonia sulcata*, *Coprinus comatus*, *Geodia baretii*, *Octopus* sp., *Metridium senile*, *Actinia* sp. and many

other plants and animals. Cardiotoxic, vasodilator, gastric secretion stimulant. Substance responsible for localised oedema response in mammalian tissues. Diagnostic aid. Sol. H₂O, MeOH. Mp 86°. Bp_{0.8} 167°. pK_{a1} 6.02; pK_{a2} 9.7 (25°, 0.1M KCl).

► Adverse human effects by subcutaneous (and other non-oral routes of) administration. Potent capillary dilator. Allergic skin effects. Exp. reprod. effects. LD₅₀ (rat, ivn) 630 mg/kg; LD₅₀ (mus, ims) 12 mg/kg. MS1050000

Hydrochloride (1:2): Histamine dihydrochloride, USAN. Amin-Glaukosan. Histamyl. Imido. Peremin. Many other names

[56-92-8] Allergic therapy, diagnostic aid. Mp 244-246°.

► Exp. reprod. effects. LD₅₀ (mus, ivn) 370 mg/kg. MS1575000

Phosphate (1:2): Histamine phosphate, USAN. Histapon

[6890-40-0]

[51-74-1] Gastric secretion stimulant. Mp 140°.

Dipicrate: Mp 239°.**N⁰-Ac: N-Acetylhistamine**

[673-49-4]

C₇H₁₁N₃O 153.183

May be considered a regular const. of amphibian skin whenever histamine is present in large amounts, e.g. *Litoria caerulea*, *Litoria aurea*, *Litoria moorei*. Isol. from leaves of *Spinacia oleracea* (spinach). Mammalian metab. of histamine. Prisms (EtOH/Et₂O or EtOH/EtOAc). Mp 147-148°.

N⁰-(4-Oxodecanoyl): N⁰-(4-Oxodecanoyl)histamine

[7222-82-4]

C₁₅H₂₅N₃O₂ 279.381

Alkaloid from the leaves of a *Glochidion* sp. (probably *Glochidion philippicum*). Cryst. (CHCl₃ or CHCl₃/EtOH). Mp 115-117°. Readily cyclises to Glochidine, G-101 and Glochidicine, G-100.

N⁰-Cinnamoyl: see **N⁰-Cinnamoylhistamine**, C-455

N⁰-Me: N-Methylhistamine

[673-50-7]

C₆H₁₁N₃ 125.173

Constit. of the skin of amphibians, e.g. *Nyctimystes disrupta*, *Litoria glandulosa* and *Leptodactylus pentadactylus labyrinthicus*.

► NI7355000

N⁰-Me, dihydrochloride: Mp 176-177°.

N⁰-Me, dipicrate: Mp 188°.

N⁰-Me, 1-N-β-D-glucopyranosyl: Casimidine

[490-41-5]

Hydrol. prod. of Casimiroedine, C-176. Cryst. (MeOH). Mp 207-209°. $[\alpha]_D +11$ (EtOH).

N⁰,N⁰-Di-Me: see 4-[2-(Dimethylamino)ethyl]imidazole, D-731

N⁰-(3-Methylbutanoyl): N-[2-(1H-Imidazol-4-yl)ethyl]-3-methylbutanamide. 4-[2-[(2-Methylbutanoyl)amino]ethyl]

imidazole. *N*^ω-Isopentanoylhistamine.

Dolichotheleine

[23100-08-5]

C₁₀H₁₇N₃O 195.264

Alkaloid from the cactus *Dolichothele sphaerica* (Cactaceae). Mp 130-131°.

N^ω-(3-Methylbutanoyl), picrate: Mp 150-152°.

N^ω-(2E,4Z-Decadienoyl): N-[2-(1H-Imidazol-4-yl)ethyl]-2,4-decadienamide. ***N*^ω-2,4-Decadienoylhistamine** [55582-43-9]

Alkaloid from *Acacia longifolia* (Fabaceae). Cryst. (MeOH aq.). Mp 145-147° (143-145°).

N^ω-(2-Hydroxy-4-oxodecanoyl): 2-Hydroxy-N-[2-(1H-imidazol-4-yl)ethyl]-4-oxodecanamide. **ω-(2-Hydroxy-4-oxodecanoyl)histidine** [139339-31-4]

C₁₅H₂₅N₃O₃ 295.381

Alkaloid from the exudate of *Jatropha gossypifolia* (Euphorbiaceae). Amorph. solid. Mp 138-140°.

1H-form

*N*¹-Me: tele-Methylhistamine

[501-75-7]

C₆H₁₁N₃ 125.173

Metab. of histamine. Cryst. (EtOH aq.). Mp 218-219° (as picrate).

*N*¹-Me, dihydrochloride: [6481-48-7]

Cryst. (EtOH). Mp 205-207°.

3H-form

N^ω-Me, *N*^ω-(4-oxodecanoyl), *N*³-Ac: N-[2-(1-Acetyl-1H-imidazol-5-yl)ethyl]-N-methyl-4-oxodecanamide. **3-Acetyl-ω-methyl-ω-(4-oxodecanoyl)histidine** [139339-32-5]

C₁₈H₂₉N₃O₃ 335.445

Alkaloid from the exudate of *Jatropha gossypifolia* (Euphorbiaceae). Mp 196-198°.

Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 489B; 492C (pmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 617D; 623B (ir)

Koessler, K.K. et al., *J.A.C.S.*, 1918, 40, 1716 (synth)

Tabor, H. et al., *J. Biol. Chem.*, 1949, 180, 703 (isol, synth, N-Acetylhistamine)

Appel, W. et al., *Arzneim.-Forsch.*, 1959, 9, 22 (isol, derivs)

Johns, S.R. et al., *Aust. J. Chem.*, 1967, 20, 555 (ω-N-(4-Oxodecanoyl)histamine)

Reisch, J. et al., *J. Pharm. Pharmacol.*, 1968, 20, 81 (ms)

Rosenberg, H. et al., *Phytochemistry*, 1970, 9, 655; 1976, 15, 501 (Dolichotheleine, isol, ir, pmr, ms, struct, synth, biosynth)

Fales, H.M. et al., *Arch. Mass Spectral Data*, 1971, 2, 650 (ms)

Rosenberg, H. et al., *J. Nat. Prod.*, 1971, 34, 372; 1974, 37, 313 (biosynth)

Horan, H. et al., *J.C.S.(C)*, 1971, 2083 (Dolichotheleine, biosynth)

Milne, G.W.A. et al., *Anal. Chem.*, 1973, 45, 1952 (cims)

Ganellin, C.R. et al., *J. Med. Chem.*, 1973, 16, 610 (pmr)

Bonnet, J.J. et al., *J.A.C.S.*, 1973, 95, 4829 (cryst struct)

Panzcia, R.P. et al., *J.A.C.S.*, 1973, 95, 8737 (Casimidine)

Repke, D.B. et al., *J. Nat. Prod.*, 1975, 38, 101 (*Acacia longifolia* alkaloid)

Roseghini, M. et al., *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1976, 54, 31 (occur, derivs)

Durant, G.J. et al., *J. Med. Chem.*, 1976, 19, 923 (*N*¹-Me)

Bryn, S.R. et al., *J.O.C.*, 1976, 41, 2283 (conform)

Roseghini, M. et al., *Z. Naturforsch., C*, 1976, 31, 118 (occur, derivs)

Reynolds, W.F. et al., *Can. J. Biochem.*, 1977, 55, 576 (cmr, tautom)

Bertaccini, G. et al., *Handb. Exp. Pharmacol.*, 1982, 59, 201 (rev)

Ferrigni, N.R. et al., *Rev. Latinoam. Quim.*, 1984, 14, 131 (Dolichotheleine, pmr, cmr)

Green, J.P. et al., *Agents Actions*, 1987, 22, 1 (rev, metab)

Potvin, P.G. et al., *Chem. Comm.*, 1987, 672 (*N*¹-Me)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 321

Ahmad, M.U. et al., *Indian J. Chem., Sect. B*, 1992, 31, 67 (*Jatropha gossypifolia* alkaloids)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 775

Worth, G.A. et al., *J.A.C.S.*, 1994, 116, 239 (tautom)

Al-Badr, A.A. et al., *Anal. Profiles Drug Subst.*, 2001, 27, 159-264 (rev)

Ramifrez, F.J. et al., *J.A.C.S.*, 2003, 125, 2328-2340 (tautom, conform, ir)

Galmarini, C.M. et al., *Curr. Opin. Invest. Drugs*, 2004, 5, 1298-1310 (rev)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HGD000; HGE000; HGD500


Histidinol

H-308

β-Amino-1H-imidazole-4-propanol, 9CI. 2-Amino-3-(4-imidazolyl)propanol. **Histidinol**

[501-28-0]

[70142-15-3]



C₆H₁₁N₃O 141.172

(S)-form [4836-52-6]

Hydrochloride (1:2): [1596-64-1]

Mp 193-195°. [α]_D¹⁹ -3.98 (H₂O).

► NI8260000

O-Phosphate:

C₆H₁₂N₃O₄P 221.152

Intermed. in histidine biosynth. Plates + 4H₂O.

N-Ac: N-Acetylhistidinol

[20462-55-9]

C₈H₁₃N₃O₂ 183.21

Isol. from a histidine-deficient *Streptomyces coelicolor* strain.

N-Benzoyl:

C₁₃H₁₅N₃O₂ 245.28

Mp 207-208°. [α]_D¹⁹ -47.6 (EtOH).

N-Benzoyl; hydrochloride: Mp 178°.

*N*¹-Me: [205885-22-9]

C₇H₁₃N₃O 155.199

Needles (EtOAc). Mp 80-81°. [α]_D²⁰ -7.2 (c, 1.4 in MeOH).

*N*³-Me: [65096-23-3]

[145104-39-8]

C₇H₁₃N₃O 155.199

Cryst. (EtOH/Et₂O) (as dihydrochloride).

Mp 195-198° (dihydrochloride). [α]_D²⁰ -2.5 (c, 1.2 in 1N HCl).

Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 492B (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 622B (ir)

Karrer, P. et al., *Helv. Chim. Acta*, 1953, 36, 570 (synth)

Bauer, H. et al., *Biochem. Prep.*, 1955, 4, 46-50 (synth)

Ballio, A. et al., *CA*, 1966, 66, 101841 (N-Ac, isol)

Lintermans, J. et al., *Biochim. Biophys. Acta*, 1972, 273, 18-20 (synth)

Bentley, K.W. et al., *Org. Prep. Proced. Int.*, 1973, 5, 5-7 (synth)


Beyerman, H.C. et al., *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1977, 96, 222-224 (3-Me, synth, pmr)

Suga, T. et al., *Acta Cryst. C*, 1997, 53, 134-136 (cryst struct, phosphate)

Fahrni, C.J. et al., *Helv. Chim. Acta*, 1998, 81, 491-506 (1-Me, synth, ir, pmr, cmr)

Histrionicotoxins

H-309



Histrionicotoxin

	R ¹	R ²
235A	-CH=CH ₂	-CH ₂ CH=CH ₂
259	-CH=CHC=CH	-CH ₂ CH=CH ₂
283A	-CH=CHC=CH	-CH ₂ CH=CHC=CH
285A	-CH=CHC=CH	-CH ₂ CH ₂ CH=C=CH ₂
285B	-CH=CHCH=CH ₂	-CH ₂ CH=CHC=CH
285C	-CH=CHC=CH	-(CH ₂) ₃ C=CH
285E	-CH=CHC=CH	-CH ₂ CH=CHCH=CH ₂
287A	-CH=CHCH=CH ₂	-CH ₂ CH ₂ CH=C=CH ₂
287B	-CH=CHCH=CH ₂	-CH ₂ CH=CHCH=CH ₂
287D	-CH=CHCH=CH ₂	-(CH ₂) ₃ C=CH
291A	-CH ₂ CH ₂ CH=CH ₂	-(CH ₂) ₃ CH=CH ₂

All ethylene bonds (Z)-

Histrionicotoxin 235A

7-Ethenyl-2-(2-propenyl)-1-azaspiro[5.5]undecan-8-ol, 9CI. HTX-235A.

Dendrobates Alkaloid 235A

C₁₅H₂₅NO 235.369

Minor or trace alkaloid from skin extracts of the poison-frogs *Dendrobates auratus*, *Dendrobates granuliferus*, *Dendrobates histrionicus*, *Dendrobates parvulus*, *Dendrobates truncatus* and *Dendrobates trivittatus* (Dendrobatidae).

Histrionicotoxin 259A

HTX-259A. Dendrobates Alkaloid 259A

C₁₇H₂₅NO 259.391

A major alkaloid of *Dendrobates tinctorius* and *Dendrobates trivittatus*; minor constit. of *Dendrobates auratus*, *Dendrobates azureus*, *Dendrobates granuliferus*, *Dendrobates histrionicus* (several populations), *Dendrobates occaltator* and *Dendrobates truncatus* (Dendrobatidae).

Histrionicotoxin 283A

HTX-283A. Dendrobates Alkaloid 283A [34272-51-0]

C₁₉H₂₅NO 283.413

A major alkaloid from *Dendrobates*

azureus, *Dendrobates granuliferus*, *Dendrobates histrionicus* (all populations), *Dendrobates occulator*, *Dendrobates parvulus*, *Dendrobates pictus*, *Dendrobates tinctorius*, *Dendrobates trivittatus* and *Dendrobates truncatus* (Dendrobatidae). Also isol. from skin extracts of the Madagascan frog *Mantella madagascariensis* (Ranidae, subfamily Mantellinae). Affects nerve channel membranes. Mp 225–228° (as hydrochloride). $[\alpha]_D^{25}$ -96.3 (c, 1.0 in EtOH).

► Rel. nontoxic to mammals.

(17E)-isomer: Δ^{17} -**trans-Histrionicotoxin 283A**. Δ^{17} -trans-HTX-283A [85718-94-1] C₁₉H₂₅NO 283.413 Trace alkaloid from *Dendrobates histrionicus* (Dendrobatidae). Opt. rotn. not recorded. May be an artifact of isoln.

Histrionicotoxin 285A

Isodihydrohistrionicotoxin. HTX-285A. *Dendrobates Alkaloid 285A* [34272-52-1]

C₁₉H₂₇NO 285.428 A major alkaloid of *Dendrobates azureus*, *Dendrobates granuliferus*, *Dendrobates histrionicus* (all populations), *Dendrobates occulator*, *Dendrobates parvulus*, *Dendrobates pictus* and *Dendrobates tinctorius* (Dendrobatidae). Also isol. from skin extracts of *Mantella madagascariensis* (Ranidae, subfamily Mantellinae). Mp 240–243° (as hydrochloride). $[\alpha]_D^{25}$ -35.3 (c, 0.5 in EtOH).

Histrionicotoxin 285B

Neodihydrohistrionicotoxin. HTX-285B. *Dendrobates Alkaloid 285B* [55475-52-0]

C₁₉H₂₇NO 285.428 Minor alkaloid from *Dendrobates histrionicus* (several populations), *Dendrobates pictus*, *Dendrobates trivittatus* and *Dendrobates truncatus* (Dendrobatidae). Mp 195–200° (as hydrochloride). $[\alpha]_D^{25}$ -125.9 (c, 1.1 in EtOH).

Histrionicotoxin 285C

Allodihydrohistrionicotoxin. HTX-285C. *Dendrobates Alkaloid 285C* [63983-63-1]

C₁₉H₂₇NO 285.428 A major alkaloid of *Dendrobates histrionicus* (all populations), *Dendrobates pictus*, *Dendrobates trivittatus*, *Dendrobates truncatus*; minor constit. in *Dendrobates granuliferus* and *Dendrobates tinctorius* (Dendrobatidae). Also isol. from skin extracts of *Mantella madagascariensis* (Ranidae, subfamily Mantellinae). Mp 247–250° (as hydrochloride). $[\alpha]_D^{25}$ -43.4 (c, 1.2 in EtOH).

Histrionicotoxin 285E

Dihydrohistrionicotoxin. *Syndihydrohistrionicotoxin*. HTX-285E. *Dendrobates Alkaloid 285E* [55785-62-1]

C₁₉H₂₇NO 285.428 Minor alkaloid of *Dendrobates histrioni-*

cus (Dendrobatidae). $[\alpha]_D^{25}$ -122 (c, 1.0 in EtOH) (hydrochloride).

Histrionicotoxin 287A

Isotetrahydrohistrionicotoxin. HTX-287A. *Dendrobates Alkaloid 287A* [55522-87-7]

C₁₉H₂₉NO 287.444 Minor alkaloid of *Dendrobates azureus*, *Dendrobates histrionicus* (all Colombian populations) and *Dendrobates pictus* (Dendrobatidae).

Histrionicotoxin 287B

Tetrahydrohistrionicotoxin. HTX-287B. *Dendrobates Alkaloid 287B* [55475-49-5]

C₁₉H₂₉NO 287.444 Minor alkaloid from *Dendrobates histrionicus* (several populations) and *Dendrobates parvulus* (Dendrobatidae).

Histrionicotoxin 287D

Allotetrahydrohistrionicotoxin. HTX-287D. *Dendrobates Alkaloid 287D* [63983-64-2]

C₁₉H₂₉NO 287.444 Trace alkaloid from the Guayacana population of *Dendrobates histrionicus* (Dendrobatidae).

Histrionicotoxin 291A

Octahydrohistrionicotoxin. HTX-291A. *Dendrobates Alkaloid 291A* [55475-50-8]

C₁₉H₃₃NO 291.476 A major alkaloid from one population of *Dendrobates histrionicus*, minor constit. in other populations, also in *Dendrobates parvulus*, *Dendrobates tinctorius* and *Dendrobates truncatus* (Dendrobatidae). Mp 180–181° (as hydrobromide).

Daly, J.W. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1971, **68**, 1870 (isol. uv, ir, pmr, ms, *cryst struct*, HTX-283A, HTX-285E)

Karle, I.L. et al., *J.A.C.S.*, 1973, **95**, 4036 (*cryst struct*, HTX-285E)

Tokuyama, T. et al., *Helv. Chim. Acta*, 1974, **57**, 2597 (HTX-285A, HTX-285B, HTX-287A, HTX-287B, HTX-291A)

Fukuyama, T. et al., *J.O.C.*, 1975, **40**, 2011 (*synth*, HTX-291A)

Daly, J.W. et al., *Helv. Chim. Acta*, 1977, **60**, 1128 (HTX-285C, HTX-287D) Tokuyama, T. et al., *Tetrahedron*, 1983, **39**, 49 (HTX-259A, HTX-285E, trans-HTX-283E)

Daly, J.W. et al., *Toxicol.*, 1984, **22**, 905 (isol. ms, HTX-283A, HTX-285A, HTX-285C) Tanner, D. et al., *Tetrahedron*, 1998, **54**, 7907–7918 (*synth*, perhydroHTX-283A)

Daly, J.W. et al., *Alkaloids: Chem. Biol. Perspect.*, 1999, **13**, 1–161 (rev. pharmacol)

Williams, G.M. et al., *J.A.C.S.*, 1999, **121**, 4900–4901 (*synth*)

Stockman, R.A. et al., *Tet. Lett.*, 2000, **41**, 9163–9165 (*synth*)

Davison, E.C. et al., *J.C.S. Perkin 1*, 2002, 1494–1514 (*synth*)

Daly, J.W. et al., *J. Nat. Prod.*, 2005, **68**, 1556–1575 (rev)


Karatholuvhu, M.S. et al., *J.A.C.S.*, 2006, **128**, 12656–12657 (HTX-238A, *synth*)

Sinclair, A. et al., *Nat. Prod. Rep.*, 2007, **24**, 298–326 (rev. *synth*)

Macdonald, J.M. et al., *Org. Lett.*, 2008, **10**, 4227–4229 (HM-X-285A, *synth*)

Hobartine

4-(1H-Indol-3-ylmethyl)-2,2,6-trimethyl-3-azabicyclo[3.3.1]non-6-ene, 9CI



C₂₀H₂₆N₂ 294.439

Various numbering systems have been used.

(-)-form [73004-61-2]

Alkaloid from *Aristotelia peduncularis* (Elaeocarpaceae). Mp 149–150.5°. $[\alpha]_D^{22}$ -20 (c, 1.66 in CHCl₃). $[\alpha]_D^{28}$ -27 (c, 1.69 in CHCl₃) (synthetic). λ_{max} 223 (log ϵ 4.6); 283 (log ϵ 3.84); 291 (log ϵ 3.78) (EtOH).

11,16-Dihydro, 11 β ,16 β -dihydroxy: **Hobartidiol** [131669-90-4]

C₂₀H₂₈N₂O₂ 328.453 Alkaloid from *Aristotelia australasica*. Cryst. (CHCl₃). Mp 270°. $[\alpha]_D^{20}$ +120 (c, 0.8 in MeOH). λ_{max} 222 (log ϵ 4.07); 282 (log ϵ 3.76); 290 (log ϵ 3.69) (EtOH).

17-Hydroxy: **17-Hydroxyhobartine**

[131669-91-5] C₂₀H₂₆N₂O 310.438 Alkaloid from *Aristotelia australasica*. Amorph. $[\alpha]_D^{20}$ +113 (c, 0.4 in CHCl₃). Numbering systems vary. Doubtful structural assignment: synthetic material not identical with the nat. prod. λ_{max} 220 (log ϵ 4.18); 282 (log ϵ 3.63); 290 (log ϵ 3.56) (EtOH).

15-Oxo: **Serratenone**

[85954-13-8] C₂₀H₂₄N₂O 308.422 Alkaloid from *Aristotelia serrata*. Amorph. $[\alpha]_D^{19}$ -45.3 (c, 1.0 in CHCl₃). Abs. config. revised in 2002 to that of other *Aristotelia* alkaloids.

8-Oxo, 9,N-didehydro: **8-Oxo-9-dehydrohobartine**

[128508-04-3] C₂₀H₂₂N₂O 306.407 Alkaloid from aerial parts of *Aristotelia chilensis*. Cryst. (CCl₄). Mp 225°.

$\Delta^{11,17}$ -Isomer: **Makomakine**

[79559-56-1] C₂₀H₂₆N₂ 294.439 Alkaloid from *Aristotelia serrata*. Cryst. (CHCl₃). Mp 106–106.5° (99–100°). $[\alpha]_D^{19}$ +131.2 (c, 0.5 in CHCl₃). $[\alpha]_D^{25}$ +142.5 (c, 2.69 in CHCl₃).

$\Delta^{11,17}$ -Isomer, 8-oxo, 9,N-didehydro: **8-Oxo-9-dehydromakomakine**. 11,12-Didehydromakomakin-10-one

[126267-21-8] C₂₀H₂₂N₂O 306.407 Alkaloid from aerial parts of *Aristotelia chilensis*. Cryst. (CCl₄). Mp 257°. Numbering systems vary.

12,14-Diepimer: **Isohobartine**

C₂₀H₂₆N₂ 294.439 Alkaloid from *Aristotelia serrata* and *Aristotelia fruticosa*. Cryst. (CHCl₃).

Mp 134-135°. $[\alpha]_D^{19}$ -30 (CHCl₃).

(±)-**form** [82769-16-2]

Synthetic. Mp 166-167°.

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 2539-2546 (*isol, uv, ir, pmr, cmr, ms*)

Bick, I.R.C. *et al.*, *Heterocycles*, 1981, **16**, 1301-1303 (*Makomakine*)

Mirand, C. *et al.*, *J.O.C.*, 1982, **47**, 4169-4170 (*Hobartine, Makomakine, synth*)

Stevens, R.V. *et al.*, *Chem. Comm.*, 1983, 384-386 (*Hobartine, Makomakine, synth, cmr*)

Bick, I.R.C. *et al.*, *Heterocycles*, 1983, **20**, 667 (*Serratenone*)

Darbre, T. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 1040-1052 (*synth, abs config*)

Bick, I.R.C. *et al.*, *Alkaloids (Academic Press)*, 1985, **24**, 113-151 (*rev, Isohobartine*)

Gribble, G.W. *et al.*, *J.O.C.*, 1985, **50**, 5900-5902 (*synth*)

Watson, W.H. *et al.*, *Acta Cryst. C*, 1989, **45**, 1322-1324 (*8-Oxo-9-dehydromakomakine, cryst struct*)

Quirion, J.C. *et al.*, *J. Nat. Prod.*, 1990, **53**, 713-716 (*17-Hydroxyhobartine, Hobartidiol*)

Céspedes, C. *et al.*, *Phytochemistry*, 1990, **29**, 1354-1356 (*8-Oxo-9-dehydrohobartine, 8-Oxo-9-dehydromakomakine*)

Beerli, R. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 110-116 (*synth*)

Burkard, S. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 275-289 (*Serratenone, synth, ir, pmr, cmr, ms*)

Dobler, M. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 1411-1420 (*17-Hydroxyhobartine, synth*)


Stoermer, D. *et al.*, *J.O.C.*, 1993, **58**, 564-568 (*Makomakine, synth*)

Galli, R. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 3400-3413 (*Serratenone, 8-Oxo-9-dehydromakomakine, synth, pmr*)

Hodgkinsine

[18210-71-4]

H-311



Absolute Configuration

C₃₃H₃₈N₆ 518.703

Alkaloid from *Hodgkinsonia frutescens* (Rubiaceae), *Psychotria lyciflora* and *Psychotria oleosides*. Platelet aggregation inhibitor, analgesic, sedative. Cryst. (C₆H₆). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 128°. λ_{max} 232 (ε 39000); 252 (ε 30200); 310 (ε 10400); 326 (ε 8600) (MeOH) (Berdy).

3a,3a',8a,8a'-Tetraepimer: Hodgkinsine B

C₃₃H₃₈N₆ 518.703

Alkaloid from *Psychotria* sp. $[\alpha]_D^{27}$ -77 (c, 1 in CHCl₃).

Anet, F.A.L. *et al.*, *Aust. J. Chem.*, 1961, **14**, 173-174 (*isol, uv*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1974, **27**, 639-646 (*ms, pmr*)

Fridrichsons, J. *et al.*, *Tetrahedron*, 1974, **30**, 85-92 (*cryst struct*)

Verotta, L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 392-396 (*isol, pmr, cmr, cd*)

Jannic, V. *et al.*, *J. Nat. Prod.*, 1999, **62**, 838-843 (*isol, cd, ms*)

Kodanko, J.J. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2528-2531 (*synth, abs config, Hodgkinsine B*)

Hodorine

C₁₉H₃₁NO₅ 353.458

Struct. unknown. Alkaloid from the roots of *Stemona sessilifolia* (Stemoneaceae). Amorph.

Hydrochloride: Mp 244-247° dec.

Hydrobromide: Mp 258-259°.

Furuya, T. *et al.*, *Chem. Zentralbl.*, 1913, 1823 (*isol*)

Suzuki, K. *et al.*, *Yakugaku Zasshi*, 1929, **49**, 457-464; *CA*, **23**, 36031

H-312

Holacimine

[84991-64-0]

Struct. unknown. Alkaloid from the bark of *Holarrhena antidysenterica* (Apocynaceae). Mp 335-336°.

Siddiqui, S. *et al.*, *Pak. J. Sci. Ind. Res.*, 1982, **25**, 201-203


H-313

Holadienine

Cona-1,4-dienin-3-one

[2841-61-4]

H-314



C₂₂H₃₁NO 325.493

Alkaloid from the bark of *Holarrhena floribunda* and from *Didymeles madagascariensis* (Apocynaceae, Didymelaceae). Mp 109° subl. $[\alpha]_D$ +80 (c, 1 in CHCl₃).

N-De-Me: Norholadienine. Norholadiene

[86171-05-3]

C₂₁H₂₉NO 311.466

Alkaloid from bark of *Holarrhena pubescens* (Apocynaceae). Fine needles (Me₂CO). Mp 216-218°. Called Norholadiene in the paper.

N-De-Me, N-(N-methylcarbamoyl): Holamide

[162857-95-6]

C₂₃H₃₂N₂O₂ 368.518

Alkaloid from bark of *Holarrhena pubescens* (*Holarrhena antidysenterica*) (Apocynaceae). Shows hypotensive activity in rats. Plates (Me₂CO). Mp 248-250°. $[\alpha]_D$ +126 (c, 0.331 in CHCl₃).

14,15-Didehydro: 14,15-Dehydroholadienine. Cona-1,4,14-trienin-3-one

[91147-30-7]

C₂₂H₂₉NO 323.477

Alkaloid from the stem bark of *Didymeles madagascariensis* (Didymelaceae). $[\alpha]_D$ -8 (c, 0.64 in CHCl₃).

18-Oxo: 3,18-Dioxo-1,4-conadienine. 18-Oxoholadienine

[113846-39-2]

C₂₂H₂₉NO₂ 339.477

Alkaloid from leaves of *Didymeles madagascariensis* and *Didymeles perrieri* (Didymelaceae). $[\alpha]_D$ +32 (c, 0.8 in EtOH).

18-Oxo, 14,15-didehydro: 14-Dehydro-18-oxoholadienine

[113846-38-1]

C₂₂H₂₇NO₂ 337.461

Alkaloid from leaves of *Didymeles madagascariensis* and *Didymeles perrieri* (Didymelaceae). Cryst. (Et₂O). Mp 185-187°. $[\alpha]_D$ +64 (c, 1.1 in EtOH).

16,18-Dioxo, 14,15-didehydro: 3,16,18-Trioxo-1,4,14-conatrienine

C₂₂H₂₅NO₃ 351.444

Trace alkaloid from leaves of *Didymeles madagascariensis* (Didymelaceae). CAS No. not found to 2006.

11α-Hydroxy: 11α-Hydroxycona-1,4-dienin-3-one, 9CI. Regholarrhenine A

[114687-90-0]

C₂₂H₃₁NO₂ 341.492

Alkaloid from the stem bark of *Holarrhena antidysenterica* (Apocynaceae). Light yellow prisms (petrol). Mp 197-198°. $[\alpha]_D^{33}$ -105 (c, 0.2 in MeOH).

11α-Hydroxy, N-de-Me: Regholarrhenine B

[114687-91-1]

C₂₁H₂₉NO₂ 327.466

Alkaloid from the stem bark of *Holarrhena antidysenterica*. Needles (MeOH). Mp 245-247°. $[\alpha]_D^{33}$ -83.5 (c, 0.3 in MeOH).

11α-Acetoxy: Mp 101-103°.

11α-Hydroxy, 18-oxo, N-de-Me: 11-Hydroxycona-1,4-dienine-3,18-dione. Pubamide

[403647-88-1]

C₂₁H₂₇NO₃ 341.449

Alkaloid from the bark of *Holarrhena pubescens*. Orange rods. Mp 130-132°. λ_{max} 201; 245 (MeOH).

11α-Acetoxy, 17,20-didehydro, N-de-Me: Pubescinine

[162857-94-5]

C₂₃H₂₉NO₃ 367.487

Alkaloid from bark of *Holarrhena pubescens* (*Holarrhena antidysenterica*) (Apocynaceae). Shows hypotensive activity in rats. Fine needles (Me₂CO). Mp 138-140°. $[\alpha]_D$ +51 (c, 0.276 in CHCl₃).

15α-Hydroxy, N-de-Me: Kurchinidine

[151271-81-7]

C₂₁H₂₉NO₂ 327.466

Alkaloid from bark of *Holarrhena antidysenterica*. Plates (CHCl₃/MeOH). Mp 114-116°.

16α-Hydroxy, 18-oxo, 14,15-didehydro: 16-Hydroxy-3,18-dioxo-1,4,14-conatrienine

[113846-40-5]

C₂₂H₂₇NO₃ 353.46

Alkaloid from leaves of *Didymeles perrieri* (Didymelaceae). $[\alpha]_D$ -30 (c, 0.46 in EtOH).

18α-Hydroxy, 14,15-didehydro: 18-Hydroxy-3-oxo-1,4,14-conatrienine

[113846-37-0]

C₂₂H₂₉NO₂ 339.477

Alkaloid from leaves of *Didymeles madagascariensis* (Didymelaceae). Cryst. (CHCl₃/Et₂O). Mp 218-220°. [α]_D²⁰ +64 (c, 0.4 in EtOH).

11 α ,18 α -Dihydroxy, N-de-Me, N-hydroxymethyl: Pubatriol [609343-73-9]

C₂₂H₃₁NO₄ 373.491

Alkaloid from the bark of *Holarrhena pubescens*. Orange rods (MeOH). Mp 110-112°. λ_{\max} 210 ; 246 (MeOH).

Janot, M.M. *et al.*, *Ann. Pharm. Fr.*, 1967, **25**, 733 (*isol. struct. ir. uv. pmr. ms. cd*)

Hoyer, G.A. *et al.*, *Planta Med.*, 1978, **34**, 47-52 (*isol*)

Sánchez, V. *et al.*, *Bull. Soc. Chim. Fr.*, 1984, 71-76; 1987, 877-884 (*Didymeles constits*)

Bhutani, K.K. *et al.*, *Phytochemistry*, 1988, **27**, 925-928 (*Regholarrhenines*)

Siddiqui, B.S. *et al.*, *Phytochemistry*, 1993, **33**, 925-928 (*Kurchinidine*)

Begum, S. *et al.*, *Phytochemistry*, 1994, **36**, 1537-1541 (*Norholadiene*)


Siddiqui, B.S. *et al.*, *Heterocycles*, 1995, **41**, 267-276 (*Holamide, Pubescimine*)

Siddiqui, B.S. *et al.*, *Phytochemistry*, 2001, **58**, 1199-1204 (*Pubamide*)

Siddiqui, B.S. *et al.*, *Heterocycles*, 2003, **60**, 909-915 (*Pubatriol*)

Holaline**H-315**

3-Dimethylaminoconan-5-ol [3559-32-8]




C₂₄H₄₂N₂O 374.609

Minor alkaloid from the bark of *Holarrhena floribunda* (Apocynaceae). Cryst. (CH₂Cl₂/MeOH). Mp 267°. [α]_D²⁰ +36 (CHCl₃/MeOH 1:1).

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 4315 (*isol. ir. pmr. ms. struct*)

Holantogenin**H-316**

18,20-Epoxyypregnane-3,14,20-triol [28708-12-5]



C₂₁H₃₄O₄ 350.497

Cryst. (dioxan aq.). Mp 125°. [α]_D²⁰ -31 (CHCl₃).

3-O-(4-Amino-2,4,6-trideoxy-3-O-methyl- β -D-ribo-hexopyranoside): Holantosine A [28719-38-2]

C₂₈H₄₇NO₆ 493.682

Constit. of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (Me₂CO)(as *N*-acetyl). Mp 260-261° (*N*-acetyl). [α]_D²⁰ -28 (CHCl₃) (*N*-acetyl).

3-O-(4-Amino-2,4,6-trideoxy-3-O-methyl- α -L-arabino-hexopyranoside): Holantosine C

[34312-24-8]

C₂₈H₄₇NO₆ 493.682

Constit. of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (as *N*-acetyl). Mp 245° (*N*-acetyl). [α]_D²⁰ -73 (CHCl₃/MeOH, 7:3) (*N*-acetyl).

3-O-(4-Amino-2,4,6-trideoxy-3-O-methyl- β -D-xylo-hexopyranoside): Holantosine E

[40738-39-4]

C₂₈H₄₇NO₆ 493.682

Alkaloid from leaves of *Holarrhena antidysenterica*. Cryst. (Me₂CO/hexane)(as *N*-acetyl). Mp 152° (*N*-acetyl). [α]_D²⁰ -47 (c, 1 in CHCl₃) (*N*-acetyl).

Janot, M.M. *et al.*, *Tetrahedron*, 1970, **26**, 1695 (*isol*)


Khuong-Hui, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1971, 864 (*isol*)

Choay, P. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **272**, 782 (*synth*)

Goutarel, R. *et al.*, *Carbohydr. Res.*, 1972, **24**, 297 (*Holantosine E*)

Holarmine**H-317**

4-Methyl-19-norconan-1,3,5(10)-trienine, 9CI [17905-60-1]




C₂₂H₃₁N 309.494

Alkaloid from the bark of *Holarrhena floribunda* (Apocynaceae). Mp 198°. [α]_D²⁰ +90 (c, 1 in CHCl₃).

Janot, M.M. *et al.*, *Ann. Pharm. Fr.*, 1967, **25**, 733 (*isol. struct. ir. uv. pmr. ms. synth*)

Holarrhenine**H-318**

3-Dimethylaminocon-5-enin-12-ol. 12-Hydroxyconessine [561-22-8]



C₂₄H₄₀N₂O 372.593

Alkaloid from *Holarrhena congolensis* and *Holarrhena mitis* (Apocynaceae). Needles (EtOAc). Mp 197-198°. [α]_D²⁰ -7.1 (CHCl₃).

Absolute configuration

Dihydrobromide: Mp 265-268° (anhyd.). [α]_D²⁰ +11 (H₂O).

O-(4-Methyl-3-pentenoyl): Holarrhetine

C₃₀H₄₈N₂O₂ 468.721

Alkaloid from *Holarrhena africana* (Apocynaceae). Mp 74-75°. [α]_D²⁰ -4.6 (c, 1.12 in EtOH). [α]_D²⁰ -14.9 (c, 1.12 in CHCl₃). Rare occurrence of 4-methyl-3-pentenoic acid residue in natural product: (See also Holafrine below).

N³-De-Me: Holarrheline. 12-Hydroxyconessimine

[2842-93-5]

C₂₃H₃₈N₂O 358.566

Alkaloid from the bark of *Holarrhena floribunda* (Apocynaceae). Mp 190°. [α]_D²⁰ +18 (c, 1 in EtOH).

N³-De-Me, N-Ac: [17905-57-6]

Mp 207°. [α]_D²⁰ +10 (c, 1 in CHCl₃).

N³-De-Me, O-(4-methyl-3-pentenoyl): Holarrhesine. Holafrine

[70866-29-4]

C₂₉H₄₆N₂O₂ 454.695

Alkaloid from the bark of *Holarrhena africana* (Apocynaceae). Platelets (Me₂CO). Mp 116-117°. [α]_D²⁰ -19.1 (c, 0.93 in CHCl₃).

5 α ,6-Dihydro, N³,N³-di-de-Me: 3-Aminoconan-12-ol. Funtessine

C₂₂H₃₈N₂O 346.555

Alkaloid from bark of *Funtumia latifolia* (Apocynaceae). Cryst. (MeOH). Mp 195°. [α]_D²⁰ +49 (c, 1 in CHCl₃).

5 α ,6-Dihydro, N³,N³-di-de-Me, O,N-di-Ac: Mp 300° dec. [α]_D²⁰ +25 (c, 0.9 in CHCl₃).

5 α ,6-Dihydro, N³,N³-di-de-Me, N³-isopropylidene:

Cryst. (Me₂CO). Mp 215-216°. [α]_D²⁰ +45 (c, 1 in CHCl₃).

Uffer, A. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1834 (*isol. ir. struct*)

Rostock, H. *et al.*, *Helv. Chim. Acta*, 1958, **41**, 11 (*Holafrine*)

v. Hove, L. *et al.*, *Tetrahedron*, 1959, **7**, 104 (*struct*)

Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 1831 (*Funtessine*)

Vandevoude, G. *et al.*, *Tet. Lett.*, 1966, 2657 (*struct*)

Janot, M.M. *et al.*, *Ann. Pharm. Fr.*, 1967, **25**, 733 (*Holarrheline*)

Hoyer, G.A. *et al.*, *Planta Med.*, 1978, **34**, 47 (*Holarrhesine*)

Zirihhi, G.N. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2637-2640 (*Holarrhetine, Holarrhesine*)

Holarrhessimine**H-319**

C₂₂H₃₆N₂O 344.539

Steroidal alkaloid. Struct. unknown. Alkaloid from the bark of *Holarrhena antidysenterica* (Apocynaceae). Cryst. (EtOAc). Mp 160-164°. [α]_D¹⁸ -31 (c, 0.775 in CHCl₃).

Hydrobromide: Mp 293-294°.

Methodide: Mp 279-280°.

Picrate: Mp 250-255° dec.

Tschesche, R. *et al.*, *Chem. Ber.*, 1954, **87**, 1719-1725

Holarrhine**H-320**

[61089-72-3]

C₂₀H₃₈N₂O₃ 354.532

Steroidal alkaloid, prob. of conanine type. Struct. unknown. Alkaloid from *Holarrhena antidysenterica* and *Wrightia tomentosa* (Apocynaceae). Needles (MeOH/EtOAc). V. sol. MeOH, EtOH, spar. sol. CHCl₃, almost insol. EtOAc. Mp 240°. [α]_D²⁵ -17.01 (c, 4.99 in MeOH). The anal. figures quoted do not correspond to the mol. formula claimed and are a better fit for C₂₄H₄₄N₂O₄.

Hexachloroplatinate: [63448-28-2]

Yellowish amorph. powder. Darkens >270°; chars >300°.


Picrate: [63408-15-1]

Pale-yellow semicryst. powder. Mp 320°.

Siddiqui, S. et al., *J. Indian Chem. Soc.*, 1932, 9, 562 (isol)

Holidine**H-321**

[95066-35-6]

C₁₉H₂₄N₂O₃ 328.41

Alkaloid from the leaves of *Phelline* sp. aff. *Phelline lucida* (Phellinaceae). [α]_D +175 (c, 1.0 in CHCl₃). Possibly an artifact.

Parent acid, amide: **Phellinamide**

[95066-34-5]

C₁₈H₂₃N₃O₂ 313.399


Alkaloid from *Phelline billiardieri* and from the leaves of *Phelline* sp. aff. *Phelline lucida* (Phellinaceae). Cryst. (Me₂CO). Mp 206°. [α]_D +180 (c, 1.0 in CHCl₃). Artifact.

Langlois, N. et al., *Heterocycles*, 1984, 22, 2453 (uv, cd, ir, pmr, cmr, ms, struct)

Langlois, N. et al., *J. Nat. Prod.*, 1988, 51, 499

Holonamine**H-322**

11-Hydroxy-23-norcona-1,4,18(22)-trienin-3-one



Absolute Configuration

C₂₁H₂₇NO₂ 325.45

Minor alkaloid from bark of *Holarrhena antidysenterica* (Apocynaceae). Mp 257-259°. [α]_D²⁵ -14.8 (c, 1.1 in MeOH). "Di-Ac" and "Tri-Ac" derivs. are descr.; the struct. of these is not clear.

A^{20(N)}-Isomer: **Kurchinine**

[155023-57-7]

C₂₁H₂₇NO₂ 325.45

Alkaloid from bark of *Holarrhena pubescens* (Apocynaceae). Plates (MeOH). Mp 202-204°.

11-Deoxy: **Maingayine**

[27317-25-5]

C₂₁H₂₇NO 309.45

Alkaloid from the bark of *Paravallis maingayi* (Apocynaceae). Mp 150-152°. [α]_D +2.1 (CHCl₃).

11-Deoxy, 1,2-dihydro: **Malouetafrine**

[50980-95-5]

C₂₁H₂₉NO 311.466

Alkaloid from leaves of *Malouetia heudelotii* (Apocynaceae). Cryst. (Me₂CO). Mp 205°. [α]_D +87.

12α-Hydroxy: **11,12-Dihydroxynorcona-N(18),1,4-trienin-3-one**C₂₁H₂₇NO₃ 341.449

Alkaloid from the stem bark of *Funtumia africana* (Apocynaceae). Mp 240-245°.

12α-Hydroxy, 11-deoxy: **12-Hydroxynorcona-N(18),1,4-trienin-3-one**C₂₁H₂₇NO₂ 325.45

Alkaloid from the stem bark of *Funtumia africana* (Apocynaceae). Mp 110°. [α]_D²² +50 (c, 0.104 in CHCl₃).

Tschesche, R. et al., *Chem. Ber.*, 1964, 97, 2316; 2326 (*Holonamine*)

Davis, J.B. et al., *Chem. Ind. (London)*, 1970, 627 (*Maingayine*)


Khuong-Huu, M.F. et al., *Phytochemistry*, 1973, 12, 1813 (*Malouetafrine*)

Wagner, H. et al., *Planta Med.*, 1987, 444 (*Funtumia africana* alkaloids)

Siddiqui, B.S. et al., *J. Nat. Prod.*, 1994, 57, 27 (*Kurchinine*)

Holstiine, 9CI**H-323**

[11053-97-7]



Absolute Configuration

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Strychnos henningsii* (*Strychnos holstii*) (Loganiaceae). Mp 247-248° dec Mp 262-264° Mp 290-292°.

O-Ac:

Cryst. (EtOAc). Mp 194-196°. [α]_D²⁰ +222.3 (c, 1 in CHCl₃).

Me ether: **Holstiline**. ar-Demethoxyrindline

[57103-48-7]

C₂₃H₂₈N₂O₄ 396.485

Alkaloid from *Strychnos henningsii* (Loganiaceae). Mp 219-220°. [α]_D²⁰ +202 (c, 0.25 in CHCl₃).

10-Methoxy, Me ether: **Rindline**

[18361-65-4]

C₂₄H₃₀N₂O₅ 426.511

Alkaloid from the bark of *Strychnos henningsii* (Loganiaceae). Prisms (Et₂O). Mp 214-216°. [α]_D¹⁹ +194 (c, 1 in CHCl₃). Struct. revised in 1975.

Janot, M.-M. et al., *C. R. Hebd. Seances Acad. Sci.*, 1951, 232, 853-855 (isol, uv, ir)

Grossert, J.S. et al., *J.C.S.*, 1965, 2812-2814 (isol, Rindline)

Spiteller-Friedmann, M. et al., *Annalen*, 1968, 711, 205-206 (ms, Rindline)

Bisset, N.G. et al., *Phytochemistry*, 1975, 14, 1411-1414 (*Holstiine*, *Holstiline*, ir, pmr, ms, struct)

Chapya, W.A. et al., *Gazz. Chim. Ital.*, 1983, 113, 773-775 (isol)


Cherif, A. et al., *J. Nat. Prod.*, 1990, 53, 793-802 (pmr, cmr, config)

Martin, G.E. et al., *J. Nat. Prod.*, 2000, 63, 543-585 (N-15 nmr)

Homaline**H-324**

1,1'-(1,4-Butanediy)bis[hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1H)-one], 9CI

[20410-93-9]



Absolute configuration

C₃₀H₄₂N₄O₂ 490.687

Alkaloid from the leaves of *Homalium pronyense* and from other *Homalium* spp. (Flacourtiaceae). Cryst. (Me₂CO). Mp 134°. [α]_D -34 (CHCl₃).

Dihydrochloride:

Cryst. + 1.5H₂O (MeOH). Mp 250°. [α]_D +7 (MeOH).

Pais, M. et al., *Tetrahedron*, 1973, 29, 1001 (isol, ir, pmr, ms, struct)

Lefebvre-Soubeyran, O. et al., *Acta Cryst. B*, 1976, 32, 1305 (cryst struct)

Crombie, L. et al., *Chem. Comm.*, 1983, 960 (synth)

Wasserman, H.H. et al., *Tetrahedron*, 1983, 39, 2459 (synth)


Crombie, L. et al., *J.C.S. Perkin I*, 1993, 2047 (synth)

Ensch, C. et al., *Helv. Chim. Acta*, 2002, 85, 1659-1677 (synth)

Homoanatoxin a**H-325**

1-(9-Azabicyclo[4.2.1]non-2-en-2-yl)-1-propanone, 9CI. *HomoAnTx*

[142926-84-9]

C₁₁H₁₇NO 179.261

Isol. from *Oscillatoria formosa* (*Phormidium formosum*) and *Raphidiopsis mediterranea*. Potent nicotinic acetylcholine agonist. Glass (as hydrochloride). Sol. H₂O, MeOH, EtOH. (±)-Form shows similar potency to (+)-enantiomer of Anatoxin a, A-974. λ_{max} 227 (no solvent reported).

▶ LD₅₀ (mus, ipr) 0.25 mg/kg.

4S-Hydroxy: 1-(4-Hydroxy-9-azabicyclo[4.2.1]non-2-en-2-yl)-1-propanone.

4-Hydroxyhomoanatoxin a

[666829-18-1]

C₁₁H₁₇NO₂ 195.261

Isol. from cyanobacterium *Raphidiopsis mediterranea*.


[142926-85-0, 142926-86-1, 142861-06-1]

Wonnacott, S. et al., *Biochem. Pharmacol.*,

1992, **43**, 419 (*synth, ir, pmr, activity*)
 Skulberg, O.M. *et al.*, *Environ. Toxicol. Chem.*,
 1992, **11**, 321 (*isol, pmr, cmr, activity*)
 Zotov, A. *et al.*, *Analyst (London)*, 1993, **118**,
 753 (*hplc*)
 Thomas, P. *et al.*, *Drug Dev. Res.*, 1994, **31**, 147
 Furey, A. *et al.*, *Rapid Commun. Mass*
Spectrom., 2003, **17**, 583-588 (*ms*)
 Namikoshi, M. *et al.*, *Toxicol.*, 2003, **42**, 533-
 538 (*4-Hydroxyhomoanatoxin a*)

Homoarginine **H-326**

N⁶-(Aminoiminomethyl)lysine, 9CI. N⁶-
 Amidinolylysine, 8CI. 2-Amino-6-guanidi-
 nohexanoic acid



C₇H₁₆N₄O₂ 188.229

(S)-form

L-form
 [156-86-5]

Present in the seeds of *Lotus helleri* and
 some members of genus *Lathyrus*.

▶ OL5550000

Hydrochloride: [1483-01-8]

Cryst. (EtOH aq.). Mp 207-209°. [α]_D²²
 +13 (c, 0.8 in 1M HCl).

Picrate:

Cryst. (EtOH aq.). Mp 202-203° dec.

(±)-form

Cryst. (EtOH aq.) (as hydrochloride). Mp
 224° (hydrochloride).

N^z-Benzoyl:

Cryst. (H₂O). Mp 273°.

Aldrich Library of 13C and 1H FT NMR
Spectra, 1992, **1**, 1281C (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn.,
 1985, **1**, 786C (*ir*)

Stevens, C.M. *et al.*, *J. Biol. Chem.*, 1950, **183**,
 139 (*synth*)


Rao, S.L.N. *et al.*, *Biochemistry*, 1963, **2**, 298
 (*isol*)

Bell, E.A. *et al.*, *Phytochemistry*, 1966, **5**, 1211
 (*occur*)

Eisele, K. *et al.*, *Annalen*, 1975, 2033 (*synth*)

Homobuxaquamarine **H-327**

[109305-89-7]



C₂₇H₄₁NO₂ 411.626

The C-4 config. is prob. the opposite of
 that shown; see note under Buxaquama-
 rine, B-462. Alkaloid from the leaves
 of *Buxus papillosa* (Buxaceae). Amorph.
 [α]_D +21.9 (c, 0.64 in CHCl₃).

Choudhary, M.I. *et al.*, *J. Nat. Prod.*, 1987, **50**,

84-88 (*isol, uv, ir, pmr, ms, struct*)

Homocinchonidine **H-328**

[1399-77-5]

C₁₉H₂₂N₂O 294.396

Struct. unknown. Alkaloid from the bark
 of *Cinchona* spp. (Rubiaceae). Prisms
 (EtOH). Mp 207.5°. [α]_D -107.3 (EtOH).

Hydrochloride:

Cryst. + 2H₂O. [α]_D -138 (EtOH).


Ac: [α]_D -34 (EtOH).

Hesse, O. *et al.*, *Ber.*, 1877, **10**, 2152; 1881, **14**,
 1890

Hesse, O. *et al.*, *Annalen*, 1880, **205**, 194

Homocryptolepinone **H-329**

[171439-52-4]



C₁₇H₁₄N₂O 262.31


Alkaloid from roots of *Cryptolepis san-*
guinolenta. Fine greenish-yellow needles
 (MeOH).

Sharaf, M.H.M. *et al.*, *J. Het. Chem.*, 1995, **32**,
 1631 (*isol, uv, ir, pmr, cmr, ms*)

Homodaphniphylic acid **H-330**

Daphnan-23-oic acid, 9CI

[25455-54-3]



Absolute
 Configuration

C₂₂H₃₅NO₂ 345.524

Me ester: **Methyl homodaphniphyllate**

[23496-03-9]

[104115-43-7]

C₂₃H₃₇NO₂ 359.551

Alkaloid from the fruits of
Daphniphyllum macropodum and
Daphniphyllum teijsmanni. Cryst.
 (MeOH/Et₂O) (as hydrochloride). Mp
 233-234° (sealed tube) (hydrochloride).

2,3-Didehydro, *Me ester*: **Methyl 2,3-**
didehydrohomodaphniphyllate. Caly-
phylline L

C₂₃H₃₅NO₂ 357.535

Alkaloid from *Daphniphyllum calyci-*
num. Amorph. solid. [α]_D²⁵ -3.5 (c, 0.5 in
 CHCl₃).

17*R*-Hydroxy: **17-Hydroxyhomodaphni-**
phylic acid

C₂₂H₃₅NO₃ 361.523

Alkaloid from the seeds of *Daphni-*
phyllum calycinum. Microcryst.
 (MeOH). Mp 97-99°. [α]_D -17 (c, 0.2 in
 MeOH).

17*R*-Hydroxy, *Me ester*: **Methyl 17-hy-**

droxyhomodaphniphyllate

C₂₃H₃₇NO₃ 375.55

Alkaloid from the fruit of *Daphni-*
phyllum macropodum. Yellowish oil.
 [α]_D²⁰ -47.5 (c, 0.1 in MeOH).

Irikawa, H. *et al.*, *Tet. Lett.*, 1969, 1821-1824
 (*Me ester, synth, ir, pmr, ms*)

Toda, M. *et al.*, *Tetrahedron*, 1974, **30**, 2683-
 2688 (*Me ester, isol*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1992, **57**, 2531-
 2544; 2585-2594 (*Me ester, synth*)

El Bitar, H. *et al.*, *J. Nat. Prod.*, 2004, **67**,
 1094-1099 (*17-Hydroxyhomodaphniphylic*
acid)


Wang, X.N. *et al.*, *Helv. Chim. Acta*, 2007, **90**,
 2156-2162 (*Methyl 17-*
hydroxyhomodaphniphyllate)

Saito, S. *et al.*, *Tetrahedron*, 2008, **64**, 1901-
 1908 (*Calyciphylline L*)

Homoerysotine **H-331**

2-Hydroxyisotaxodine

[93888-49-4]



C₁₉H₂₅NO₄ 331.411

2-Hydroxyisotaxodine and Homoeryso-
 tine have not been compared. No phys.
 props. were reported for the latter.

Alkaloid from the twigs and foliage of
Athrotaxis cupressoides, *Athrotaxis sela-*
ginoides and *Athrotaxis laxifolia* (Taxo-
 diaceae) and from *Phelline* sp. aff.

Phelline lucida (Phellinaceae). Noncryst.
 [α]_D¹⁹ +60.69 (c, 1.0 in CHCl₃).

2-*Ac*: **2-Acetoxyisotaxodine**

[93888-50-7]

C₂₁H₂₇NO₅ 373.448

Trace alkaloid from the twigs and
 foliage of *Athrotaxis cupressoides* and
Athrotaxis selaginoides (Taxodiaceae).
 Inseparable mixt. with 2-acetoxytaxo-
 dine.

O¹⁶-*De-Me*, O¹⁷-*Me*: **Hydroxytaxodine**

[93888-53-0]

C₁₉H₂₅NO₄ 331.411

Alkaloid from twigs and foliage of
Athrotaxis cupressoides, *Athrotaxis sel-*
aginoides and *Athrotaxis laxifolia*
 (Taxodiaceae). Cryst. (Me₂CO). Mp
 192-193°. [α]_D¹⁹ +51.47 (c, 1.03 in
 CHCl₃).

O¹⁶-*De-Me*, O¹⁷-*Me*, 2-*Ac*: **2-Acetoxy-**
taxodine

[93888-54-1]

C₂₁H₂₇NO₅ 373.448

Alkaloid from twigs and leaves of
Athrotaxis cupressoides (Taxodiaceae).
 Inseparable mixt. with 2-acetoxyiso-
 taxodine.

2-*Epimer*: **2-Epihydroxyisotaxodine**

[93888-48-3]

C₁₉H₂₅NO₄ 331.411

Alkaloid from the twigs and foliage of
Athrotaxis cupressoides, *Athrotaxis sel-*
aginoides and *Athrotaxis laxifolia*


(Taxodiaceae). Inseparable mixt. with 2-hydroxytaxodine.

2-Epimer, O¹⁷-Me: 7-Deoxycephalofortuneine

[128508-20-3]
C₂₀H₂₇NO₄ 345.438
Alkaloid from *Cephalotaxus fortunei* (Cephalotaxaceae).

Panichanun, S. *et al.*, *Tetrahedron*, 1984, **40**, 2677; 2685 (*isol, uv, ir, pmr, ms, struct, derivs*)
Razafimbelo, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1985, **300**, 441 (*isol, uv, ir, pmr, ms, cd, struct*)
Ma, G. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 137-140; *CA*, **113**, 74722k (7-Deoxycephalofortuneine)

Homofascaplysate A H-332
[694436-60-7 ((+)-form, chloride)]




C₂₀H₁₅N₂O₃[⊕] 331.35
Quaternary alkaloid *isol.* from the sponge *Fascaplysinopsis reticulata*. Yellow solid (as chloride). [α]_D²⁵ +14 (c, 0.25 in MeOH) (chloride).

10-Bromo, 6,7-dihydro: 10-Bromo-6,7-dihydrohomofascaplysate A

[1001396-30-0]
C₂₀H₁₆BrN₂O₃[⊕] 412.262
Alkaloid from an undescribed thorectid sponge.
Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 783-792 (*Homofascaplysate A*)
Ankietty, S. *et al.*, *Nat. Prod. Commun.*, 2007, **2**, 1145-1148 (*10-Bromodihydrohomofascaplysate A*)

Homofascaplysin A H-333
[132911-50-3]




C₂₁H₁₇N₂O₂[⊕] 329.377
Isol. from the sponge *Hyrtios erecta* and from *Fascaplysinopsis reticulata*. Inhibits HIV reverse transcriptase (HIV-rt). Shows cytotoxic, antibacterial and antiplasmodial activities. Inhibits p56^{lck} tyrosine kinase. Brown oil. λ_{max} 202 (ε 11300); 220 (ε 11200); 264 (ε 8100); 334 (ε 5250) (MeOH) (Derep). λ_{max} 256 (ε 2010); 314 (ε 838) (EtOH) (Berdy).

Salt with Dehydrofuffariellolide diacid:
C₄₆H₅₄N₂O₆ 730.942
Alkaloid-sesterterpene salt from the sponge *Fascaplysinopsis reticulata*. Red viscous oil. [α]_D²⁰ -9.36 (c, 0.0064 in MeOH).

Jiménez, C. *et al.*, *J.O.C.*, 1991, **56**, 3403 (*isol*)

uv, ir, pmr, cmr, ms, struct, activity)
Kirsch, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 825-829 (*isol, activity*)

Homofascaplysin B H-334
[132938-12-6]



R = COCOOMe

C₂₁H₁₄N₂O₃ 342.353
Alkaloid from the sponge *Fascaplysinopsis reticulata*. Red oil. λ_{max} 268 (ε 12100); 294 (ε 5980); 330 (ε 7690) (MeOH) (Derep).

3-Bromo: 3-Bromohomofascaplysin B
[693790-73-7]
C₂₁H₁₃BrN₂O₃ 421.249
Isol. from a *Didemnum* sp. Yellow solid.

Jimenez, C. *et al.*, *J.O.C.*, 1991, **56**, 3403-3410 (*isol, uv, ir, pmr, cmr, struct*)
Gribble, G.W. *et al.*, *J.O.C.*, 1992, **57**, 3636-3642 (*synth*)
Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 783-792 (*3-Bromohomofascaplysin B*)

Homofascaplysin B1 H-335
[694436-61-8]

As Homofascaplysin B, H-334 with R = -COCOEt
C₂₂H₁₆N₂O₃ 356.38
Isol. from a *Didemnum* sp. Red oil.

3-Bromo: 3-Bromohomofascaplysin B1
[693790-74-8]
C₂₂H₁₅BrN₂O₃ 435.276
Isol. from a *Didemnum* sp. Yellow solid.

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 783-792 (*isol, pmr, cmr*)

Homofascaplysin C H-336
12H-Pyrido[1,2-a:3,4-b']diindole-13-carboxaldehyde, 9CI


[132911-52-5]
As Homofascaplysin B, H-334 with R = CHO

C₁₉H₁₂N₂O 284.317
Alkaloid from *Didemnum* sp. and *Fascaplysinopsis reticulata*. Yellow oil. λ_{max} 266 (ε 7810); 292 (ε 2980) (MeOH) (Derep).

3-Bromo: 3-Bromohomofascaplysin C
[193790-75-9]
C₁₉H₁₁BrN₂O 363.213
Isol. from a *Didemnum* sp. Yellow solid.


Jimenez, C. *et al.*, *J.O.C.*, 1991, **56**, 3403-3410 (*isol, uv, ir, pmr, cmr, struct*)
Gribble, G.W. *et al.*, *J.O.C.*, 1992, **57**, 3636-3642 (*synth*)
Carter, D.S. *et al.*, *J.O.C.*, 1999, **64**, 8537-8545 (*synth*)
Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 783-792 (*3-Bromohomofascaplysin C*)

Homofluorodaturatin H-337
[80757-46-6]



C₁₅H₁₄N₂O₂ 254.288
Struct. revised in 1988. Alkaloid from seeds of *Datura stramonium* (Solana-ceae). Amorph. λ_{max} 268 (ε 2800); 368 (ε 8600); 455 (ε 5200) (MeOH/NaOH) (Derep). λ_{max} 217 (ε 22200); 261 (ε 6780); 387 (ε 20700) (MeOH) (Derep).
Maier, I. *et al.*, *Monatsh. Chem.*, 1981, **112**, 1425 (*isol, uv, ir, pmr, ms*)
Robien, W. *et al.*, *Sci. Pharm.*, 1988, **56**, 133; *CA*, **110**, 193165d (*struct*)

Homolaudanosine H-338
1-[2-(3,4-Dimethoxyphenyl)ethyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, 9CI




C₂₂H₂₉NO₄ 371.475

(R)-form
Synthetic. [α]₇₈ -13.5 (c, 0.89 in EtOH).

(S)-form [85611-40-1]
Alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae). [α]_D²⁵ +11 (c, 0.21 in EtOH).
Picrate: Mp 78-79°.
Aladesanmi, A.J. *et al.*, *J. Nat. Prod.*, 1983, **46**, 127 (*isol, uv, pmr, ms, cmr, cd, struct*)
Wanner, K.T. *et al.*, *Heterocycles*, 1989, **29**, 29 (*synth*)
Gottlieb, L. *et al.*, *J.O.C.*, 1990, **55**, 5659 (*synth*)

Homomoenjodaramine H-339
[220749-24-6]



C₂₉H₄₈N₂O 440.711
Alkaloid from *Buxus hyrcana*. Acetylcholinesterase inhibitor. Amorph. pow-

der. $[\alpha]_D^{25} +40$ (c, 1.12 in CHCl_3). λ_{max} 237 (log ϵ 3.82); 244 (log ϵ 3.85) (MeOH).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1998, **49**, 481-488 (*isol*, *pmr*, *cmr*, *uv*)

Homophleine H-340


$\text{C}_{56}\text{H}_{90}\text{N}_2\text{O}_9$ 935.335

Erythrophleum alkaloid. Struct. unknown. Alkaloid from the bark of *Erythrophleum guineense* (Fabaceae). Cardiac stimulant, hypotensive agent, local anaesthetic. Amorph. No further data to 2007.

Dalma, G. *et al.*, *Ann. Chim. Appl.*, 1935, **25**, 569-571; *CA*, **30**, 6378 (*isol*)

Homopiptanthine H-341

[38965-96-7]



$\text{C}_{21}\text{H}_{35}\text{N}_3$ 329.528

Stereoisomeric with Jamine, J-16. Note quasi-enantiomeric relationship. Alkaloid from the stem bark of *Bowdichia virgiloides* (Fabaceae). Mp 184-185°. $[\alpha]_D^{21} -40$ (c, 0.32 in CHCl_3).

16,17-Didehydro: Homopodopetaline

[114420-75-6]

$\text{C}_{21}\text{H}_{33}\text{N}_3$ 327.512

Alkaloid from *Ormosia coutinhoi*, *Ormosia macrophylla* and *Ormosia semicastrata*. Also produced synthetically by the action of HCHO on Podopetaline, P-529. Plates (Me₂CO). Mp 117-118.5°. $[\alpha]_D -147.7$ (c, 0.2 in MeOH).

6-Epimer, 16,17-didehydro: Homo-6-epi-podopetaline. Homosweetinine

[114370-18-2]

$\text{C}_{21}\text{H}_{33}\text{N}_3$ 327.512

5Alkaloid from *Ormosia coutinhoi* and *Ormosia nobilis* and *Sweetia elegans*.

McLean, S. *et al.*, *Can. J. Chem.*, 1981, **59**, 34 (*Homopodopetaline*, *synth*, *ms*)

Balandrin, M.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 619-622 (*Homosweetinine*)


Kinghorn, A.D. *et al.*, *Phytochemistry*, 1988, **27**, 439-444 (*Homopodopetaline*, *occur*)

Torrenegra, R. *et al.*, *Phytochemistry*, 1989, **28**, 2219-2221 (*isol*, *ir*, *ms*, *struct*)

Le, P.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 283-293 (*Homopodopetaline*)

Homoplatensimide H-342

[1036714-38-1]



$\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_6$ 458.553

Prod. by *Streptomyces platensis* MA7327. Gum.

Me ester: [1036714-39-2]


$\text{C}_{26}\text{H}_{36}\text{N}_2\text{O}_6$ 472.58

Prod. by *Streptomyces platensis* MA7327. $[\alpha]_D^{23} -16$ (c, 0.5 in MeOH). λ_{max} 236 (MeOH).

Jayasuriya, H. *et al.*, *Tet. Lett.*, 2008, **49**, 3648-3651 (*isol*, *pmr*, *cmr*)

Homopseudopteroxazole H-343

[549520-74-3]



$\text{C}_{26}\text{H}_{37}\text{NO}$ 379.584


Homologue of Pseudopteroxazole, P-724. Alkaloid from *Pseudopterozorgia elisabethae*. Active against *Mycobacterium tuberculosis*. Yellowish oil. $[\alpha]_D^{25} +103.2$ (c, 0.9 in CHCl_3). λ_{max} 204 (ϵ 25500); 244 (ϵ 9000); 279 (ϵ 3500) (MeOH).

Rodriguez, I.I. *et al.*, *J. Nat. Prod.*, 2003, **66**, 855-857 (*isol*, *pmr*, *cmr*)

Homopumiliotoxin 321B H-344

Octahydro-3-(7-hydroxy-2,5-dimethyl-4-octenylidene)-1-methyl-2H-quinolizin-1-ol, 9CI

[150044-81-8]



$\text{C}_{20}\text{H}_{35}\text{NO}_2$ 321.502

Isol. from toads of the genus *Melanophryniscus*.

7'-Ketone: Homopumiliotoxin 319A

[150044-79-4]

$\text{C}_{20}\text{H}_{33}\text{NO}_2$ 319.486

Isol. from the genus *Melanophryniscus*.

***A*^{5,6'}-Isomer, 7'-ketone: Homopumiliotoxin 319B**

[150044-80-7]

$\text{C}_{20}\text{H}_{33}\text{NO}_2$ 319.486

Isol. from the genus *Melanophryniscus*.


Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 357-373 (*isol*)

Kibayashi, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1157-1159 (*struct*)

Homopumiliotoxin 223G H-345

Dendrobates Alkaloid 223G

[109175-44-2]



Absolute Configuration

$\text{C}_{14}\text{H}_{25}\text{NO}$ 223.358

Alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates pumilio* (Dendrobatidae). $[\alpha]_D^{28} +1.7$ (c, 1 in CHCl_3) (synthetic).

Hydrochloride: Mp 183-184° (synthetic).

$[\alpha]_D^{25} +48.1$ (c, 0.5 in MeOH) (synthetic).

Tokuyama, T. *et al.*, *Tetrahedron*, 1987, **43**, 643 (*isol*, *pmr*, *ms*, *struct*)

Edwards, M.W. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1188 (*isol*)

Kibayashi, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1157-1159 (*synth*, *abs config*)

Santos, L.S. *et al.*, *Tet. Lett.*, 2001, **42**, 6999-7001 (*synth*)


Chen, B.-F. *et al.*, *Tetrahedron*, 2004, **60**, 10223-10231 (*synth*)

Huang, P.-Q. *et al.*, *Org. Lett.*, 2006, **8**, 1435-1438 (*synth*)

Voituriez, A. *et al.*, *Org. Lett.*, 2007, **9**, 4705-4708 (*synth*)

Homosecodaphniphylic acid H-346

12,16-Cyclo-1,12-secodaphnan-23-oic acid



Absolute Configuration

$\text{C}_{22}\text{H}_{35}\text{NO}_2$ 345.524

Me ester: **Methyl homosecodaphniphyllate**

[28519-09-7]

[118099-25-5 (\pm)-form]]

$\text{C}_{23}\text{H}_{37}\text{NO}_2$ 359.551

Alkaloid from *Daphniphyllum*

macropodium and the fruits of

Daphniphyllum teijsmanni

(Daphniphyllaceae). Cryst. (hexane).

Mp 102.5-103°.

N-Ac, Me ester:

Plates (hexane). Mp 105-106°.

Irikawa, H. *et al.*, *Tet. Lett.*, 1969, 1821 (*ir*, *ms*)

Sasaki, K. *et al.*, *J.C.S.(B)*, 1971, 1565 (*cryst struct*, *abs config*)

Toda, M. *et al.*, *Tetrahedron*, 1972, **28**, 1477;

1974, **30**, 2683 (*isol*, *ir*, *pmr*, *ms*, *struct*)

Ruggeri, R.B. *et al.*, *J.A.C.S.*, 1988, **110**, 8734

(*synth*)


Heathcock, C.H. *et al.*, *J.O.C.*, 1992, **57**, 2544

(*synth*)

Nicolaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, 465 (*bibl. synth*)

Homosedinone H-347

1-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidiny]-2-butanone, 9CI. 2-(2-Hydroxy-2-phenylethyl)-1-methyl-6-(2-oxobutyl)piperidine [145774-74-9]



$C_{18}H_{27}NO_2$ 289.417
Trace alkaloid from *Sedum acre*. Cryst. (hexane). Mp 62-63°. $[\alpha]_D^{20}$ -40 (c, 0.2 in $CHCl_3$).

6-Epimer: 6-Epimomosedinone

[145307-61-5]

$C_{18}H_{27}NO_2$ 289.417

Alkaloid from *Sedum acre*.

Stereoisomer: Lelobanololine

[128508-05-4]

$C_{18}H_{27}NO_2$ 289.417


Alkaloid from whole plants of *Lobelia davidii*. Red-brown oil. $[\alpha]_D^{24}$ -28.4 ($CHCl_3$).

Zhang, M.-Z. *et al.*, *Phytochemistry*, 1990, **29**, 1353 (*Lelobanololine*)

Durant, A. *et al.*, *Can. J. Chem.*, 1992, **70**, 2722 (*occur, pmr, cmr, ms, synth, struct*)

Homostachydrine H-348

2-Carboxy-1,1-dimethylpiperidinium hydroxide inner salt, 9CI. Pipecolic acid methylbetaine [1195-94-4]



$C_8H_{15}NO_2$ 157.212

(S)-form [472-22-0]

Isol. from seeds of *Medicago sativa* and *Achillea millefolium*. Fine needles. Mp 300° dec. $[\alpha]_D$ -9 (H_2O).

Hydrochloride: Mp 216-217° dec. $[\alpha]_D$ -13.3 (c, 1.43 in EtOH).

(±)-form

V. hygroscopic cryst. Mp 207-208°.

Hydrochloride: Mp 210-211° dec.

Wiehler, G. *et al.*, *Can. J. Chem.*, 1958, **36**, 339 (*isol, struct, synth*)

Robertson, A.V. *et al.*, *Can. J. Chem.*, 1959, **37**, 829 (*abs config*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1959, **78**, 134 (*abs config*)

Pailer, M. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1960, **293**, 646 (*isol*)

Buechel, K.H. *et al.*, *Chem. Ber.*, 1962, **95**, 2453 (*synth*)

Homothermopsine H-349

[11024-41-2]


$C_{17}H_{24}N_2O$ 272.389

Prob. quinolizidine alkaloid. Struct. unknown. Alkaloid from *Thermopsis lanceolata* (Fabaceae). Small, fine needles (EtOH). Mp 224-225°. $[\alpha]_D$ +86.9 (c, 1.4 in MeOH). Cooccurs with Anagryne, A-970, Sparteine, S-385, Thermopsine, T-363 and Caulophylline in C-938.

Orechhoff, A. *et al.*, *Ber.*, 1934, **67**, 1394-1398

Homotyramine H-350

4-(3-Aminopropyl)phenol, 9CI. 3-(p-Hydroxyphenyl)propylamine [57400-89-2]



$C_9H_{13}NO$ 151.208

Mp 68.5-69.5°. Bp₃ 153.5-155.5°.

Hydrochloride: [40829-38-7]

Mp 160-161°.

N-Me: **Methylhomotyramine**. 4-[(3-Methylamino)propyl]phenol [32180-92-0]

$C_{10}H_{15}NO$ 165.235

Alkaloid from *Croton humilis* (Euphorbiaceae). Cryst. Mp 105-108°.

Me ether, N-(3-hydroxy-4-methoxy-E-cinnamoyl): **Mirabliamide**

[220360-38-3]

$C_{20}H_{23}NO_4$ 341.406

Alkaloid from *Mirabilis himalaica*.

Goldschmidt, G. *et al.*, *Monatsh. Chem.*, 1914, **35**, 388 (*synth*)

Mazitova, F.N. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1963, 2063; *CA*, **60**, 5371c (*synth*)

Stuart, K.L. *et al.*, *Phytochemistry*, 1971, **10**, 460 (*isol, struct, pmr, ms, deriv*)

Zhang, G. *et al.*, *CA*, 1999, **130**, 165486d (*Mirabliamide*)

N-Homoveratroylhomoveratrylamine H-351

N-[2-(3,4-Dimethoxyphenyl)ethyl]-3,4-dimethoxybenzeneacetamide, 9CI [139-76-4]



$C_{20}H_{25}NO_5$ 359.421

Alkaloid from *Pleiospermium alatum* leaves (Rutaceae). Needles (C_6H_6 /petrol). Mp 122.5-123° (119-121°).

Kundu, A.B. *et al.*, *Chem. Ind. (London)*, 1975, 433-434 (*isol, uv, ir, ms, struct, synth*)

Kozikowski, A.P. *et al.*, *J.O.C.*, 1980, **45**, 2548-2550 (*synth*)


Patra, A. *et al.*, *Org. Magn. Reson.*, 1981, **16**, 65-67 (*cmr*)

Martin, N.H. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 762-774 (*synth*)

Konyukov, V.N. *et al.*, *CA*, 1983, **99**, 212765x (*synth*)

Hopeanine H-352

1,5,6,7,7a,8,9,10,11,12a-Decahydro-2-(2-pyrrolidinylmethyl)-2H-pyrano[2,3,4-ij]-pyrido[1,2-b][2,7]naphthyridine, 9CI [68156-57-0]



$C_{19}H_{29}N_3O$ 315.458


Alkaloid from *Brunfelsia hopeana* (Solanaaceae).

Iyer, R. *et al.*, *CA*, 1978, **89**, 191130n (*isol*)

Gellert, E. *et al.*, *CA*, 1980, **92**, 42210y (*isol*)

Hopromalinol H-353

1-[4-[Hexahydro-4-(2-hydroxyheptyl)-5-methyl-2-oxo-1,5-diazocin-1(2H)-yl]butyl]hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1H)-one, 9CI [49620-05-5]



R = Ph, R' = $CH_2CH(OH)(CH_2)_4CH_3$

$C_{31}H_{52}N_4O_3$ 528.777

Alkaloid from the leaves of *Homalium pronyense* (Flacourtiaceae). Noncryst. $[\alpha]_D$ -17 (c, 2.5 in $CHCl_3$).

Pais, M. *et al.*, *Tetrahedron*, 1973, **29**, 1001 (*isol, ir, pmr, ms, struct*)

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1993, 2055 (*synth*)

Hopromine H-354

1-[4-[4-Heptylhexahydro-5-methyl-2-oxo-1,5-diazocin-1(2H)-yl]butyl]hexahydro-5-methyl-4-pentyl-1,5-diazocin-2(1H)-one, 9CI [49620-03-3]

As Hopromalinol, H-353 with R = $-(CH_2)_4CH_3$, R' = $-(CH_2)_6CH_3$

$C_{30}H_{58}N_4O_2$ 506.814

Alkaloid from the leaves of *Homalium pronyense* (Flacourtiaceae). Noncryst. $[\alpha]_D$ -10 (c, 3 in $CHCl_3$).

Pais, M. *et al.*, *Tetrahedron*, 1973, **29**, 1001 (*isol, ir, pmr, ms, struct*)

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1993, 2055 (*synth*)

Ensch, C. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1659-1677 (*synth, abs config*)

Hoprominol**H-355**

1-[4-[Hexahydro-4-(2-hydroxyheptyl)-5-methyl-2-oxo-1,5-diazocin-1(2H)-yl]butyl]hexahydro-5-methyl-4-pentyl-1,5-diazocin-2(1H)-one, 9CI
[49620-04-4]

As Hopromalinol, H-353 with

R = $-(\text{CH}_2)_4\text{CH}_3$, -m = $-\text{CH}_2\text{CH}(\text{OH})(\text{CH}_2)_4\text{CH}_3$

$\text{C}_{30}\text{H}_{58}\text{N}_4\text{O}_3$ 522.813


Alkaloid from the leaves of *Homalium pronyense* (Flacourtiaceae). Noncryst. $[\alpha]_{\text{D}}^{20}$ -19 (c, 2 in CHCl_3).

Pais, M. *et al.*, *Tetrahedron*, 1973, **29**, 1001 (*isol, ir, pmr, ms, struct*)

Crombie, L. *et al.*, *J.C.S. Perkin 1*, 1993, 2055 (*synth*)

Hordatine B, 8CI**H-356**

[10502-21-3]



$\text{C}_{29}\text{H}_{40}\text{N}_8\text{O}_5$ 580.686

Alkaloid from *Hordeum vulgare* (barley), *Hordeum bulbosum*, *Hordeum jubatum*, *Hordeum murinum* and *Hordeum spontaneum* (Poaceae). Antifungal antibiotic, phytotoxin. Sol. H_2O . Mp 132-135° (as dipicrate). $[\alpha]_{\text{D}}^{23}$ +54 (c, 1 in H_2O). λ_{max} 223 (ε 22000); 301 (ε 15500); 316 (ε 16200) (EtOH) (Berdy).

4'-O- α -D-Glucopyranoside: *Hordatine B* glucoside

$\text{C}_{35}\text{H}_{50}\text{N}_8\text{O}_{10}$ 742.828

Present in barley (*Hordeum vulgare*) as an inseparable mixt. with Hordatine A glucoside, known as Hordatine M (Poaceae). $[\alpha]_{\text{D}}^{23}$ +60 (c, 1 in H_2O).

O-De-Me: *Hordatine A*

[7073-64-5]

$\text{C}_{28}\text{H}_{38}\text{N}_8\text{O}_5$ 566.659

Alkaloid from *Hordeum vulgare*, *Hordeum bulbosum*, *Hordeum distichon*, *Hordeum jubatum* and *Hordeum murinum* (Poaceae). Antifungal agent. Sol. H_2O . Mp 132-135° (as dipicrate). $[\alpha]_{\text{D}}^{23}$ +69 (c, 1 in H_2O). λ_{max} 229 (ε 20700); 298 (ε 18400); 307 (ε 18400) (MeOH) (Berdy). λ_{max} 235 (ε 17000); 297 (ε 19400); 305 (ε 18700) (MeOH/NaOH) (Berdy).

O-De-Me, 4'-O- α -D-glucopyranoside: *Hordatine A* glucoside

$\text{C}_{34}\text{H}_{48}\text{N}_8\text{O}_{10}$ 728.801

Component of Hordatine M from barley (see above).


[16995-91-8]

Stoessl, A. *et al.*, *Can. J. Chem.*, 1967, **45**, 1745 (*isol, uv, ir, nmr, struct, synth*)

Smith, T.A. *et al.*, *Phytochemistry*, 1978, **17**, 1093 (*occur*)

Horde-nine**H-357**

4-(2-Dimethylaminoethyl)phenol, 9CI. 4-Hydroxy-N,N-dimethylphenethylamine. N,N-Dimethyltyramine. Peyocactine. Anhaline. Cactine. Eremursine
[539-15-1]



$\text{C}_{10}\text{H}_{15}\text{NO}$ 165.235

Alkaloid from *Anhalonium fissuratum*, *Hordeum vulgare* (barley), a very wide range of plant spp. esp. in the Cactaceae. Also present in the Amaryllidaceae, Poaceae, Fabaceae and a few algae and fungi including marine alga *Phyllophora nervosa*. Diuretic, disinfectant, antihypertensive (in large doses) agent. Used for treatment of dysentery. Feeding repellent for grasshoppers. Shows similar actions to 2-(Methylamino)-1-phenyl-1-propanol, M-386. Mp 118°. Bp₁₁ 173-174°. Log P 1.17 (calc).

Hydrochloride: Mp 176-177°.

Sulfate (2:1): [622-64-0]

Mp 207-208°.

O- β -D-Glucopyranoside: *Horde-nine* O-glucoside

[93710-65-7]

$\text{C}_{16}\text{H}_{25}\text{NO}_6$ 327.377

Alkaloid from *Pancreatium biflorum*. Solid. Mp 100-108° dec. $[\alpha]_{\text{D}}^{20}$ -55.4 (c, 0.5 in EtOH). λ_{max} 225 (log ε 3.92); 266 (sh) (log ε 3.61); 275 (log ε 3.52); 305 (sh) (log ε 2.98) (MeOH).

O- α -L-Rhamnopyranoside: *Horde-nine* O-rhamnoside

$\text{C}_{16}\text{H}_{25}\text{NO}_5$ 311.377

Alkaloid from *Selaginella doederleinii* (Selaginellaceae). Amorph. solid. $[\alpha]_{\text{D}}^{20}$ -96 (c, 1 in MeOH).

O-[6-O-E-Cinnamoyl- β -D-glucopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranoside]: *Horde-nine* O-(6''-O-trans-cinnamoyl-3'-O- β -D-glucopyranosyl- α -L-rhamnopyranoside)

[111537-51-0]

$\text{C}_{31}\text{H}_{41}\text{NO}_{11}$ 603.665

Alkaloid from *Selaginella doederleinii* (Selaginellaceae). Amorph. solid. $[\alpha]_{\text{D}}^{20}$ -84 (c, 1 in MeOH).

O-[4-Hydroxycinnamoyl-(\rightarrow 6)- β -D-glucopyranosyl-(1 \rightarrow 3)- α -L-rhamnopyranoside]: *Horde-nine* O-(6''-O-p-coumaroyl-3'-O- β -D-glucopyranosyl- α -L-rhamnopyranoside)

[130748-25-3]

$\text{C}_{31}\text{H}_{41}\text{NO}_{12}$ 619.664

Alkaloid from *Selaginella doederleinii* (Selaginellaceae). Amorph. $[\alpha]_{\text{D}}^{20}$ -81 (c, 0.2 in MeOH).

N-Ac: N-Acetylhorde-nine

$\text{C}_{12}\text{H}_{18}\text{NO}_2$ 208.28

Alkaloid from *Stapelia hirsuta*. Buff powder. Mp 120-123°. Counterion not specified.

N-Me: *Candicine*. Maltoxin

[6656-13-9]

$\text{C}_{11}\text{H}_{18}\text{NO}^{\oplus}$ 180.269

Quaternary alkaloid from *Trichocereus candicans*, *Trichocereus lamprochlorus*, *Trichocereus spachianus*, several *Desmodium* spp. (Cactaceae, Fabaceae) and several other spp. in different families. Also from mescal (*Lophophora williamsii*). Ganglionic blocker and vasoconstrictor with nicotine-like action. Curarising in large doses.

► Toxic, LD₅₀ (rat) 50 mg/kg. BR1574700

N-Me, chloride: Mp 285° dec.

N-Me, iodide: [1976-98-3]

Mp 230-231°.

Voswinkel, H. *et al.*, *Ber.*, 1912, **45**, 1004 (*synth*)

Buck, J.S. *et al.*, *J.A.C.S.*, 1938, **60**, 1789 (*synth, deriv*)

Badger, G.M. *et al.*, *Aust. J. Chem.*, 1963, **16**, 734 (*isol, pmr*)

Rao, G.S. *et al.*, *J. Pharm. Pharmacol.*, 1970, **22**, 545

Güven, K.C. *et al.*, *Phytochemistry*, 1970, **9**, 1893 (*isol, Phyllophora*)

Cherayil, G.D. *et al.*, *J. Pharm. Sci.*, 1973, **62**, 2054 (*synth*)

Rao, K.V. *et al.*, *Planta Med.*, 1975, **27**, 31 (*isol, deriv*)

Srinivasan, P.R. *et al.*, *Org. Magn. Reson.*, 1976, **8**, 198 (*cmr*)

Kruger, T.L. *et al.*, *J.O.C.*, 1977, **42**, 4161 (*ms*)

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*occur*)

Meyer, B.N. *et al.*, *J. Nat. Prod.*, 1983, **46**, 688 (*O-Methylcandicine*)

Ghosal, S. *et al.*, *Phytochemistry*, 1984, **23**, 1167 (*glucoside*)

Lin, R.C. *et al.*, *J. Nat. Prod.*, 1990, **53**, 882 (*glycosides*)


Parvez, M. *et al.*, *Acta Cryst. C*, 1991, **47**, 1450 (*cryst struct*)

Shabana, M. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 710-714 (*Candicine*, N-Acetylhorde-nine, glucoside)

Hormaomycin**H-358***Takaokamycin*

[121548-21-8]

[92092-69-8]




Absolute Configuration

$\text{C}_{55}\text{H}_{69}\text{ClN}_{10}\text{O}_{14}$ 1129.661

Depsipeptide antibiotic. Prod. by *Streptomyces griseoflavus* W384 and *Streptomyces* sp. AC-1978. Morphogenic agent, antibiotic production stimulator, cyto-differentiation Inducer, mycelium formation inducer. Powder. Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. Mp 166-168°. $[\alpha]_{\text{D}}^{20}$ +20.8 (c, 0.5 in MeOH). λ_{max} 206 (ε 43000); 278 (ε 9150) (MeOH).

Omura, S. *et al.*, *J. Antibiot.*, 1984, **37**, 700-705 (*Takaokamycin*)

- Andres, N. *et al.*, *Helv. Chim. Acta*, 1989, **72**, 426-437 (*isol, pmr, cmr, ms*)
 Roessner, E. *et al.*, *Angew. Chem., Int. Ed.*, 1990, **29**, 64-65 (*struct*)
 Otaguru, K. *et al.*, *J. Antibiot.*, 2003, **56**, 322-323 (*isol, struct*)
 Zlatopolskiy, B.D. *et al.*, *Chem. Eur. J.*, 2004, **10**, 4708-4717; 4718-4727; 5568; 2005, **11**, 2929-2945 (*synth, pmr, conformm, abs config*)

Horsfiline**H-359***(R)*-formC₁₃H₁₆N₂O₂ 232.282**(R)-form** [136247-72-8]

Alkaloid from the leaves of *Horsfieldia superba* (Myristicaceae). Cryst. (Me₂CO). Mp 125-126°. [α]_D²⁰ -7.2 (c, 1 in MeOH).

Hydrochloride: [α]_D -28 (c, 0.6 in MeOH).

(S)-form

Synthetic. [α]_D +5.2 (c, 0.5 in MeOH). Mp not detd. due to paucity of material for recrystallisation.

Hydrochloride: [α]_D +28 (c, 0.6 in MeOH).

(±)-form [136316-07-9]

Cryst. (Me₂CO). Mp 156-157°.

(ξ)-form**Demethoxy: Coerulescine**

[66859-18-5]

C₁₂H₁₄N₂O 202.255

Alkaloid from the leaves of *Phalaris coerulescens*. Pale yellow gum. [α]_D²⁰ -0.8 (c, 0.02 in MeOH).

- Jossang, A. *et al.*, *J.O.C.*, 1991, **56**, 6527 (*isol, uv, ir, pmr, cmr, ms, synth, struct*)
 Jones, K. *et al.*, *Chem. Comm.*, 1992, 1767 (*synth*)
 Bascop, S.-I. *et al.*, *Heterocycles*, 1994, **38**, 725 (*synth*)
 Pellegrini, C. *et al.*, *Tetrahedron: Asymmetry*, 1994, **5**, 1979 (*synth, abs config*)
 Palmisano, G. *et al.*, *Tetrahedron: Asymmetry*, 1996, **7**, 1 (*synth*)
 Anderton, N. *et al.*, *Phytochemistry*, 1998, **48**, 437-439 (*isol, pmr, cmr, ms, Coerulescine*)
 Lakshmaiah, G. *et al.*, *J.O.C.*, 1999, **64**, 1699-1704 (*synth*)
 Fischer, C. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 1175-1181 (*synth*)
 Somei, M. *et al.*, *Heterocycles*, 2000, **53**, 7-10 (*Coerulescine, synth*)
 Cravotto, G. *et al.*, *J.O.C.*, 2001, **66**, 8447-8453 (*synth*)
 Lizo, D.E. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 117-122 (*Horsfiline, Coerulescine, synth*)
 Murphy, J.A. *et al.*, *Org. Lett.*, 2005, **7**, 3287-3289 (*synth*)
 Chang, M.-Y. *et al.*, *Tet. Lett.*, 2005, **46**, 8463-8465 (*synth*)


Hortesine**H-360**

Deriv. of cytosine containing a phosphate group. Metab. of *Streptomyces mutabilis* NRRL 2528. Antifungal agent. Sol. H₂O; fairly sol. MeOH; poorly sol. EtOH,

hexane. λ_{max} 267 (E_{1%}/1cm 100) (pH 7 buffer) (Berdy). λ_{max} 270 (H₂O) (Berdy).
 U.S. Pat., 1961, 2 972 569 (*isol, activity*)
 U.S. Pat., 1965, 3 169 902; CA, **62**, 11115 (*isol, props*)

Hortiacine**H-361**

8,13-Dihydro-10-methoxyindo-*lo*[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI. 10-Methoxyrutaecarpine [522-55-4]


C₁₉H₁₅N₃O₂ 317.346

Alkaloid from the bark of *Hortia arborea* and *Hortia badinii* and from the trunk wood of *Hortia longifolia* (Rutaceae). Pale-yellow needles (EtOAc). Mp 252-252.5°. Also obt. by subliming Hortiacine hydrochloride at 230-270°/0.5 mm.

- Pachter, I.J. *et al.*, *J.A.C.S.*, 1960, **82**, 5187 (*isol, struct, synth*)
 Danieli, B. *et al.*, *Gazz. Chim. Ital.*, 1975, **105**, 45 (*synth*)
 de Barros Corrêa, D. *et al.*, *Phytochemistry*, 1975, **14**, 2059 (*isol*)
 Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246 (*synth, ir, ms*)

Hostasine**H-362**

[955092-43-0]



Relative Configuration

C₁₈H₂₁NO₇ 363.366

Alkaloid from *Hosta plantaginea*. Cryst. (EtOH). Mp 202-205°. [α]_D²⁶ -34 (c, 0.21 in MeOH). λ_{max} 223 (log ε 4.34); 263 (log ε 3.87); 344 (log ε 1.79); 360 (log ε 1.01); 397 (log ε 1.32) (MeOH).

10-Me ether: 10-O-Methylhostasine

[955089-62-0]

C₁₉H₂₃NO₇ 377.393

Alkaloid from *Hosta plantaginea*. Amorph. solid. [α]_D²⁶ -65.7 (c, 0.38 in MeOH). λ_{max} 223 (log ε 4.38); 261 (log ε 3.88); 291 (log ε 3.3); 369 (log ε 1.98); 392 (log ε 2.13) (MeOH).

8-Demethoxy: 8-Demethoxyhostasine

[955089-53-9]

C₁₇H₁₉NO₆ 333.34

Alkaloid from *Hosta plantaginea*. Amorph. solid. [α]_D²⁶ -29.5 (c, 0.36 in MeOH). λ_{max} 223 (log ε 4.28); 260 (log ε 3.71); 297 (log ε 3.62); 392 (log ε 2.02) (MeOH).

8-Demethoxy, 10-Me ether: 8-Demethoxy-10-O-methylhostasine

[955089-57-3]

C₁₈H₂₁NO₆ 347.367


Alkaloid from *Hosta plantaginea*. Acetylcholinesterase inhibitor.

Amorph. solid. [α]_D²⁶ -76.4 (c, 0.53 in MeOH). λ_{max} 224 (log ε 4.37); 258 (log ε 3.76); 295 (log ε 3.65); 343 (log ε 1.57); 380 (log ε 1.9) (MeOH).

Wang, Y.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1458-1461 (*isol, pmr, cmr, cryst struct*)

Hostasinine A**H-363**

[960010-06-4]

C₁₇H₁₇NO₆ 331.324


Alkaloid from *Hosta plantaginea*. Needles (MeOH). Mp 229-231°. [α]_D¹⁶ +124.4 (c, 0.45 in MeOH). λ_{max} 244 (log ε 3.87); 297 (log ε 3.48) (MeOH).

Wang, Y.-H. *et al.*, *Org. Lett.*, 2007, **9**, 5279-5281 (*isol, synth, cd, pmr, cmr, ms, cryst struct*)

Hosukinidine**H-364**

Veratraman-3-ol, 9CI. Veratra-5,12-dien-3-ol


[72765-23-2]

C₂₇H₄₃NO 397.643

Alkaloid from the rhizomes of illuminated *Veratrum grandiflorum* (Liliaceae). Mp 176.5-177.5°. [α]_D -56.5 (c, 0.27 in MeOH). Unusual configs. at C-17 and C-20.

N,O-Di-Ac: Mp 195-197°. [α]_D -55.6 (c, 0.23 in CHCl₃).

Kaneko, K. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 2534 (*isol, cryst struct*)

Hoveine**H-365**C₃₈H₄₈N₄O₄ 624.822

Alkaloid from the whole plant of *Hovea longipes* (Fabaceae). Cryst. + $\frac{1}{2}$ CHCl₃. Mp 190-193°. [α]_D²⁰ -11 (c, 0.27 in CHCl₃/MeOH, 1:1).

Hydrochloride (1:2):

Needles + 4H₂O. Mp ca.° 280 (dec.). [α]_D²⁰ -9 (c, 0.30 in EtOH/H₂O, 4:1).

Hydrobromide (1:2):

Cryst. + 2H₂O (EtOH). Mp 280° dec. [α]_D²⁰ -8 (c, 0.26 in EtOH/H₂O, 4:1).

Tetra-Ac:


Noncryst. powder. Mp 153-155°.

Fitzgerald, J.S. et al., *An. Quim.*, 1972, **68**, 737 (isol, ir, struct, abs config)

Huangshanine

H-366

O⁷-Methylfaberonine
[88313-35-3]



C₄₂H₅₀N₂O₉ 726.865

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Yellowish powder. Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 70-75°. [α]_D²⁵ +120.6 (c, 0.43 in MeOH). λ_{max} 281 (ε 25500); 302 (ε 15900) (MeOH) (Berdy).

O⁷-De-Me: **Faberonine**. 7'-Desmethyl-huangshanine

[91926-00-0]
C₄₁H₄₈N₂O₉ 712.838

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Yellow solid. [α]_D²⁴ +82.8 (c, 0.498 in MeOH).

O⁷, O¹¹-Di-de-Me: **Iznikine**

[82958-14-3]
C₄₀H₄₆N₂O₉ 698.811

Minor alkaloid from the roots and rhizomes of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). [α]_D²⁵ +76 (c, 0.068 in MeOH).

6a,7-Didehydro: **Dehydrohuangshanine**

[91948-29-7]
C₄₂H₄₈N₂O₉ 724.849

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Yellow solid.

Guinaudeau, H. et al., *Tet. Lett.*, 1982, **23**, 2523 (Iznikine)


Lin, L.-Z. et al., *Planta Med.*, 1983, **49**, 55 (Huangshanine)

Wagner, H. et al., *Tetrahedron*, 1984, **40**, 2133 (Huangshanine, Faberonine, Dehydrohuangshanine)

Humantenirine

H-367

4-Demethyl-11-methoxyhumantenine, 9CI
[82375-30-2]



C₂₁H₂₆N₂O₄ 370.447

Alkaloid from Hu-Man-Teng (root of *Gelsemium elegans*) and from the stems of *Gelsemium rankinii* (Loganiaceae). Needles (Me₂CO). Mp 167-170°. [α]_D²⁰ -135 (c, 0.56 in MeOH).

N⁴-Me: **11-Methoxyhumantenine**

[93914-74-0]
C₂₂H₂₈N₂O₄ 384.474

Alkaloid from the whole plant of *Gelsemium elegans* (Loganiaceae). Yellow powder. [α]_D²² -146.5 (c, 0.06 in MeOH).

O¹¹-De-Me: **11-Hydroxyrankinidine**. 11-O-Demethylhumantenirine

[122590-03-8]
C₂₀H₂₄N₂O₄ 356.421

Alkaloid from the whole plant of *Gelsemium elegans* (Loganiaceae). Needles. Mp 212-214°. [α]_D²² -135 (c, 0.06 in MeOH).

O¹¹-De-Me, N⁴-Me: **11-Hydroxyhumantenine**, 9CI

[122590-04-9]
C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the whole plant of *Gelsemium elegans* (Loganiaceae). Needles. Mp 176-177°. [α]_D²⁰ -130 (c, 0.05 in MeOH).

11-Demethoxy: **Rankinidine**. 4-Demethylhumantenine, 9CI

[106466-66-4]
C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the stems of *Gelsemium rankinii* (Loganiaceae). Needles (Me₂CO). Mp 175-178°. [α]_D²⁰ -126 (c, 0.07 in MeOH). Dec. rapidly to gum, even at 5°.

11-Demethoxy, N⁴-Me: **Humantenine**, 9CI

[82375-29-9]
C₂₁H₂₆N₂O₃ 354.448

Alkaloid from roots of *Gelsemium elegans* (Loganiaceae). Gum. [α]_D -142.

11-Demethoxy, 19,20-dihydro, 20R-hydroxy: **20-Hydroxydihydrorankinidine**

[135626-62-9]
C₂₀H₂₆N₂O₄ 358.436

Alkaloid from *Gelsemium elegans* (Loganiaceae). Needles (Me₂CO). Mp 173-174°. [α]_D -165 (c, 0.02 in MeOH).

Bis(demethoxy): **N-Desmethoxyrankinidine**

[122590-02-7]
C₁₉H₂₂N₂O₂ 310.395

Alkaloid from the whole plant of *Gelsemium elegans* (Loganiaceae). Needles. Mp 258-260°. [α]_D²² -169.2 (c, 0.052 in MeOH).

Bis(demethoxy), N⁴-Me: **1-Demethoxyhumantenine**

[123629-91-4]
C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Gelsemium elegans* (Loganiaceae). Needles. Mp 238-240°. [α]_D -188.5 (c, 0.052 in MeOH).

15-Hydroxy, 11-demethoxy, N⁴-Me: **15-Hydroxyhumantenine**, 9CI

[135626-63-0]
C₂₁H₂₆N₂O₄ 370.447

Alkaloid from whole plants of *Gelsemium elegans*. Needles. Mp 213-215°.

[α]_D -82.2 (c, 0.028 in MeOH).

Yang, J. et al., *Yaoxue Tongbao*, 1982, **17**, 119; *CA*, **97**, 36099f (Humantenine)

Yang, J. et al., *Yaoxue Xuebao*, 1983, **18**, 104; 1984, **19**, 686; *CA*, **99**, 102248y; **102**, 21236w (isol, cryst struct, Humantenine)

Schun, Y. et al., *J. Nat. Prod.*, 1986, **49**, 806 (Rankinidine)

Xu, C. et al., *Zhongguo Yixue Kexueyuan Xuebao*, 1987, **9**, 95; *CA*, **108**, 187030y (cryst struct, abs config, Humantenine)

Lin, L.-Z. et al., *J. Nat. Prod.*, 1989, **52**, 588 (derivs)

Kitajima, M. et al., *J.C.S. Perkin 1*, 1991, 1773 (synth, N-Desmethoxyrankinidine)

Lin, L.-Z. et al., *Phytochemistry*, 1991, **30**, 1311 (15-Hydroxyhumantenine)

Takayama, H. et al., *Nat. Prod. Lett.*, 1993, **2**, 271 (synth)


Phisalaphong, C. et al., *Tet. Lett.*, 1993, **34**, 4035 (synth, 20-Hydroxydihydrorankinidine)

Hunteracine

H-368

[28999-86-2]

[53643-60-0 (chloride)]



Absolute configuration

C₁₈H₂₃N₂O⁺ 283.392

Indole alkaloid, poss. Corynanthe-related. Alkaloid from *Hunteria eburnea* (Apocynaceae). Mp 343-344° dec. (as chloride). [α]_D -90 (c, 0.1 in MeOH aq.) (chloride).

Bartlett, M.F. et al., *J.O.C.*, 1963, **28**, 1445 (isol, uv)

Burnell, R.H. et al., *Chem. Comm.*, 1970, 772; *Can. J. Chem.*, 1974, **52**, 2327 (uv, ir, pmr, ms, cryst struct)

Hunteriamine

H-369

[1357-71-7]

[C₃₈-₃₉H₄₆-₄₈N₄O₂]

Struct. unknown. Dimeric indole alkaloid. Alkaloid from *Hunteria eburnea* (Apocynaceae). Shows hypotensive props.

Perchlorate:

Cryst. Mp 279-281° dec. [α]_D +29.4 (c, 1 in MeOH).


Hydrochloride: Mp 310-315° dec. [α]_D²³ +27.5 (c, 1 in MeOH).

Renner, U. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1963, **331**, 105-108 (isol)

Hupehenidine

H-370

3,6-Dihydroxy-11,12-secocevan-18-one
[123857-37-4]




$C_{27}H_{45}NO_3$ 431.657

Alkaloid from the Hubei-beimu (bulbs of *Fritillaria hupehensis*) (Liliaceae).

Wu, J. et al., *Zhongcaoyao*, 1989, **20**, 194; *CA*, **111**, 228955m

Hupehensine H-371

17,23-Epoxy-5,6-dihydro-3-hydroxyveratraman-11-one, 9CI
[105814-56-0]



$C_{27}H_{41}NO_3$ 427.626

The config. at C-17 is poss. erroneous. Alkaloid from the bulbs of *Fritillaria hupehensis* (Liliaceae).

17,20,23-Triepimer: *Songbeisine*

[129214-36-4]

$C_{27}H_{41}NO_3$ 427.626

Alkaloid from bulbs of *Fritillaria unibracteata* (Liliaceae).

Wu, J. et al., *Yaoyue Xuebao*, 1986, **21**, 546;

CA, **106**, 15699r (isol, uv, ir, pmr, ms, struct)


Yu, S. et al., *Zhongcaoyao*, 1990, **21**, 2; *CA*,

113, 112481r (*Songbeisine*)

Wang, I.P. et al., *Yaoyue Xuebao*, 1992, **27**, 273

Hupeol H-372

[209168-34-3]



$C_{11}H_{13}NO_3$ 207.229


Alkaloid from *Maaackia hupehensis*. Regarded as an intermediate in the metab. of lupine alkaloids. Needles ($CH_2Cl_2/MeOH$). $[\alpha]_D^{23} +32.3$ (c, 0.26 in EtOH).

Wang, Y.-H. et al., *J. Chem. Res., Synop.*, 1998, 196-197 (isol, pmr, cmr, ms)

Wang, Y.-H. et al., *Heterocycles*, 2000, **53**, 545-548 (synth)

Huperzine I H-373

[291764-11-9]




$C_{16}H_{23}NO_2$ 261.363

Alkaloid from *Huperzia serrata*. Needles

($CHCl_3/Me_2CO$). Mp 154-156°. $[\alpha]_D +141.1$ (c, 0.05 in MeOH).

Gao, W.-Y. et al., *Chin. J. Chem.*, 2000, **18**, 614-616

Huperzine V H-374



Absolute Configuration


$C_{28}H_{47}N_3O$ 441.699

Alkaloid from *Huperzia serrata*. Prisms. CAS No. not found to 2008.

Liu, H.Q. et al., *Chin. Chem. Lett.*, 2004, **15**, 303-304 (isol, pmr, cmr)

Huperzine A H-375

Selagine. Fordine. Isoselagine
[102518-79-6]



$C_{15}H_{18}N_2O$ 242.32

Identity of Huperzine A with Selagine established in 1989 and with Isoselagine in 1993. Alkaloid from *Lycopodium serratum* (= *Huperzia serrata*), *Lycopodium selago*, *Lycopodium saururus*, *Lycopodium erythraeum* and *Lycopodium gnidioides* (Lycopodiaceae). Shows very strong anticholinesterase activity and markedly increases efficiency for learning and memory in animals. Currently under clinical investigation in China for treatment of myasthenia gravis and Alzheimer's disease. Mp 230° (224-226°, 214-215°). $[\alpha]_D^{24.5} -150.4$ (c, 0.498 in MeOH). $[\alpha]_D -99$. λ_{max} 231 (ε 10200); 313 (ε 7760) (EtOH) (Derep).

►PB9185700

6β-Hydroxy: 6β-Hydroxyhuperzine A

[125295-13-8]

$C_{15}H_{18}N_2O_2$ 258.319

Minor alkaloid from *Lycopodium selago* (Lycopodiaceae). Cryst. ($Me_2CO/MeOH$). Mp 207-210°. $[\alpha]_D -206$ (c, 0.37 in MeOH).

10-Hydroxy: 10-Hydroxyhuperzine A.

Carinatamine A

[929286-89-5]

$C_{15}H_{18}N_2O_2$ 258.319

Alkaloid from *Lycopodium carinatum*. Amorph. solid. $[\alpha]_D^{25} -30$ (c, 0.3 in MeOH).

6,6-Dimethyl: Dimethylhuperzine A

[316792-28-6]

$C_{17}H_{22}N_2O$ 270.374

Potent acetylcholinesterase inhibitor. $[\alpha]_D -64.2$ (c, 0.61 in $CHCl_3$).

[132435-40-6, 103735-86-0, 130791-77-4, 174876-78-9, 120786-18-7]

Yoshimura, H. et al., *Tet. Lett.*, 1960, **No. 12**, 14 (*Selagine*)

Valenta, Z. et al., *Tet. Lett.*, 1960, **No. 10**, 26 (isol, uv, ir, pmr)

Chen, C.H. et al., *T'ai-wan Yao Hsueh Tsa Chih*, 1984, **36**, 1; *CA*, **101**, 143940w (*Isoselagine*)

Liu, J.-S. et al., *Can. J. Chem.*, 1986, **64**, 837 (uv, ir, pmr, cmr, struct)

Liu, J. et al., *Huaxue Xuebao*, 1986, **44**, 1035; *CA*, **107**, 115821p (struct)

Ayer, W.A. et al., *Can. J. Chem.*, 1989, **67**, 1538 (6β-Hydroxyhuperzine A, *Isoselagine*)

Xia, Y. et al., *J.A.C.S.*, 1989, **111**, 4116 (synth)

Qian, L. et al., *Tet. Lett.*, 1989, **30**, 2089 (synth)

Geib, S.J. et al., *Acta Cryst. C*, 1991, **47**, 824 (cryst struct)

Yamada, F. et al., *J.A.C.S.*, 1991, **113**, 4695 (synth)

Kozikowski, A.P. et al., *J.O.C.*, 1991, **56**, 4636 (synth)

Kozikowski, A.P. et al., *Adv. Med.*, 1992, **1**, 175 (rev, synth, props)

Kozikowski, A.P. et al., *Chem. Comm.*, 1993, 860 (synth)

Campiani, G. et al., *J.O.C.*, 1993, **58**, 7660 (synth)

Zhou, B.N. et al., *Phytochemistry*, 1993, **34**, 1425 (pmr, cmr)

Sun, C.M. et al., *Planta Med.*, 1993, **59**, 467 (*Isoselagine*, struct)

Kozikowski, A.P. et al., *Bioorg. Med. Chem. Lett.*, 1996, **6**, 259-262; 2000, **10**, 2467-2469 (dimethylhuperzine A, isomers, pharmacol)

Kaneko, S. et al., *Heterocycles*, 1997, **46**, 27-30 (synth)

Kaneko, S. et al., *Tetrahedron*, 1998, **54**, 5471-5484 (synth, isomers, pmr)

Chassaing, C. et al., *Tet. Lett.*, 1999, **40**, 8805-8809 (synth)

Bai, D.L. et al., *Curr. Med. Chem.*, 2000, **7**, 355-374 (rev, chem, pharmacol)

Haudrechy, A. et al., *Tetrahedron*, 2000, **56**, 3181-3187 (synth)

Camps, P. et al., *Tetrahedron*, 2000, **56**, 4541-4553 (synth)

Wang, R. et al., *Acta Pharmacol. Sin.*, 2006, **27**, 1-26 (rev)

Choo, C.Y. et al., *Bioorg. Med. Chem.*, 2007, **15**, 1703-1707 (*Carinatamine A*)

Ma, X. et al., *J. Ethnopharmacol.*, 2007, **113**, 15-34 (rev)

Niemitz, M. et al., *Magn. Reson. Chem.*, 2007, **45**, 878-882 (pmr)

Lucey, C. et al., *Org. Biomol. Chem.*, 2007, **5**, 301-306 (synth)

Little, J.T. et al., *Expert Opin. Invest. Drugs*, 2008, **17**, 209-215 (pharmacol)


Ma, X. et al., *Phytochemistry*, 2008, **69**, 2022-2028 (*Huperzine A*, isol)

Huperzine B H-376

8,15-Didehydro-1(18H)-lycodinone, 9CI

Fordimine

[103548-82-9]



Absolute Configuration

C₁₆H₂₀N₂O 256.347

Unclear whether Fordimine, assigned the same struct. as Huperzine B but claimed to be a new alkaloid, is a stereoisomer of Huperzine B. Alkaloid from *Lycopodium serratum*, *Huperzia serrata* and *Phlegmariurus fordii* (preferred genus name *Huperzia*). Exhibits marked anticholinesterase activity. Mp 270-271°. [α]_D²⁵ -54.2 (c, 0.203 in MeOH). λ_{\max} 231 (ϵ 10200); 313 (ϵ 7760) (EtOH) (Derep).

▶OL2780000

N^z-Me: N-Methylhuperzine B

[110037-64-4]

C₁₇H₂₂N₂O 270.374

Alkaloid from *Lycopodium serratum* (*Huperzia serrata*) (Lycopodiaceae). This N atom is normally tertiary in Lycopodium alkaloids and does not have a locant number in the usual numbering scheme.

11R-Hydroxy: 11-Hydroxyhuperzine B.

Carinatamine B

[929103-66-2]

C₁₆H₂₀N₂O₂ 272.346


Alkaloid from *Lycopodium carinatum*. Amorph. solid. [α]_D²⁵ -16 (c, 0.3 in MeOH).

Liu, J.-S. et al., *Can. J. Chem.*, 1986, **64**, 837-839 (*uv, ir, pmr, cmr, struct*)Liu, J. et al., *Huaxue Xuebao*, 1986, **44**, 1035-1040; *CA*, **107**, 115821p (*struct*)Li, J. et al., *Zhongcaoyao*, 1987, **18**, 50-51; *CA*, **107**, 93545x (*N-Methylhuperzine B*)Chu, B.M. et al., *Yaoxue Xuebao*, 1988, **23**, 115-121; *CA*, **109**, 70347 (*Fordimine*)Wu, B. et al., *J.O.C.*, 1997, **62**, 5978-5981(*synth, pmr, cmr*)Lee, I.Y.C. et al., *Tet. Lett.*, 2004, **45**, 285-287 (*synth*)Choo, C.Y. et al., *Bioorg. Med. Chem.*, 2007, **15**, 1703-1707 (*Carinatamine B*)

Huperzine F'

H-377

[927704-42-5]



Absolute Configuration

C₁₆H₁₉NO₃ 273.331Alkaloid from *Huperzia serrata*.

Amorph. purple powder. Mp 117-119° dec. [α]_D²⁵ -3182 (c, 0.004 in MeOH). λ_{\max} 204 (log ϵ 1.01); 287 (log ϵ 0.4); 556 (log ϵ 1.23) (MeOH).

12-Deoxy: Huperzine E'

[927704-41-4]

C₁₆H₁₉NO₂ 257.332Alkaloid from *Huperzia serrata*.


Amorph. purple powder. [α]_D²⁵ -4750 (c, 0.004 in MeOH). λ_{\max} 205 (log ϵ 1.48); 286 (log ϵ 0.78); 547 (log ϵ 2.62) (MeOH).

Wang, H.-B. et al., *Helv. Chim. Acta*, 2007, **90**, 153-157 (*isol, pmr, cmr*)

Huperzine F

H-378

[371918-18-2]



Absolute Configuration

C₁₆H₂₁NO₃ 275.347Alkaloid from *Huperzia serrata*.

2,3-Dihydro: Huperzine O

[314285-43-3]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*.

12-Deoxy: Huperzine E

[371918-17-1]

C₁₆H₂₁NO₂ 259.347Alkaloid from *Huperzia serrata*.

12-Deoxy, 2-chloro: 2-Chlorohuperzine E

[927704-40-3]

C₁₆H₂₀ClNO₂ 293.792


Alkaloid from *Huperzia serrata*. Yellowish needles. Mp 149-150° dec. [α]_D²⁵ -63.4 (c, 0.46 in CHCl₃). λ_{\max} 204 (log ϵ 0.72); 324 (log ϵ 2.53) (MeOH).

Wang, B.-D. et al., *Youji Huaxue*, 2000, **20**, 812-814; 2001, **21**, 606-610; *CA*, **134**, 68763; **135**, 344620t (*isol, struct*)Wang, H.-B. et al., *Helv. Chim. Acta*, 2007, **90**, 153-157 (*2-Chlorohuperzine E*)

Huperzine H

H-379

[245432-14-8]

C₁₆H₂₁NO₂ 259.347


Alkaloid from *Huperzia serrata*. Needles. Mp 168-170°. [α]_D -216 (c, 0.02 in CHCl₃).

Gao, W.Y. et al., *Chin. Chem. Lett.*, 1999, **10**, 463-466

Huperzine J

H-380

[316821-09-7]



Absolute Configuration

C₁₇H₃₀N₂O 278.437

Alkaloid from *Huperzia serrata* and *Lycopodium serratum*. Prisms (EtOAc/hexane). Mp 120-123°. [α]_D²⁰ +32.2 (c, 0.05 in CHCl₃).

N^b-Oxide: Huperzine L

[316821-11-1]

C₁₇H₃₀N₂O₂ 294.436

Alkaloid from *Huperzia serrata*. Yellowish oil. [α]_D²⁰ +69.4 (c, 0.08 in CHCl₃).

N-De-Me: Lycoposerramine X. Carinatamine C

[927813-32-9]

C₁₆H₂₈N₂O 264.41

Alkaloid from *Lycopodium carinatum* and *Lycopodium serratum*. Amorph. powder. [α]_D²⁵ +35 (c, 0.3 in MeOH) (*Carinatamine C*).

5-Epimer: Lycoposerramine Y

[927813-33-0]

C₁₇H₃₀N₂O 278.437

Alkaloid from *Lycopodium serratum*. Amorph. powder.

5-Epimer, N-de-Me: Huperzine K

[316821-10-0]

C₁₆H₂₈N₂O 264.41

Alkaloid from *Huperzia serrata*. Yellowish oil. [α]_D²⁰ -16.8 (c, 0.06 in CHCl₃).

8a-Epimer, N-de-Me: Lycoposerramine Z

[927813-34-1]

C₁₆H₂₈N₂O 264.41


Alkaloid from *Lycopodium serratum*. Amorph. powder.

Gao, W. et al., *Planta Med.*, 2000, **66**, 664-667 (*Huperzines J-L*)Katakawa, K. et al., *Heterocycles*, 2006, **69**, 223-229 (*Lycoposerramines X, Y and Z, cryst struct*)Choo, C.Y. et al., *Bioorg. Med. Chem.*, 2007, **15**, 1703-1707 (*Carinatamine C*)

Huperzine P

H-381

[292601-58-2]



Absolute Configuration

C₁₆H₂₁NO₃ 275.347


Alkaloid from *Huperzia serrata*. Prisms. Mp 320.5-321.5°. [α]_D²⁵ +2.5 (c, 1 in CHCl₃).

Tan, C.-H. et al., *Tet. Lett.*, 2000, **41**, 5733-5736

Huperzine Q

H-382

[441756-04-3]

C₁₆H₂₅NO₂ 263.379Alkaloid from the whole plant of *Hu-*

perzia serrata. Prisms (Me₂CO). Mp 192-194°. [α]_D²⁵ -0.28 (c, 0.44 in CHCl₃).


N-Oxide: Huperzine Q N-oxide. Oxyhuperzine Q
[441756-05-4]
C₁₆H₂₅NO₃ 279.378
Alkaloid from the whole plant of *Huperzia serrata*. Amorph. powder. [α]_D²⁵ +0.06 (c, 0.12 in CHCl₃).

Tan, C.-H. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1058-1061 (*isol, pmr, cmr, ms, cryst struct*)

Huperzine R

H-383

[443776-29-2]



Relative Configuration


C₁₅H₂₁NO₃ 263.336
Alkaloid from the whole plant of *Huperzia serrata*. Prisms (petrol/Me₂CO). Mp 189-191°. [α]_D²⁵ -0.1 (c, 0.42 in CHCl₃).

Tan, C.-H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1021-1022 (*isol, pmr, cmr, cryst struct*)

Huperzine S

H-384

[618457-04-8]




C₁₆H₂₃NO₃ 277.363
Alkaloid from *Huperzia serrata*. Needles (petrol/Me₂CO). Mp 146-148°. [α]_D²⁵ +156.6 (c, 1.34 in CHCl₃).

Tan, C.-H. *et al.*, *Can. J. Chem.*, 2003, **81**, 315-318 (*isol, pmr, cmr*)

Huperzine T

H-385

[618457-05-9]




C₁₆H₂₃NO₃ 277.363
Alkaloid from *Huperzia serrata*. Amorph. powder. [α]_D²⁵ -1.5 (c, 0.07 in CHCl₃).

Tan, C.-H. *et al.*, *Can. J. Chem.*, 2003, **81**, 315-318 (*isol, cd, pmr, cmr*)

Huperzine U

H-386

[618457-06-0]




C₁₆H₂₂N₂O₂
Alkaloid from *Huperzia serrata*. Amorph. powder. Mp >300° dec. Racial.

Tan, C.-H. *et al.*, *Can. J. Chem.*, 2003, **81**, 315-318 (*isol, cd, pmr, cmr*)

Huperzine W

H-387

[439694-98-1]




C₁₄H₂₁NO₂ 235.325
Alkaloid from the whole plant *Huperzia serrata*. Yellowish oil.

Tan, C.H. *et al.*, *Chin. Chem. Lett.*, 2002, **13**, 331-332 (*isol, pmr, cmr*)

Huperzine

H-388

[119188-49-7]



Absolute Configuration

C₁₇H₂₂N₂O 270.374
Alkaloid from *Lycopodium serratum* (also given as *Huperzia serrata*; preferred genus name *Lycopodium*) and *Lycopodium casuarinoides* (*Lycopodium casuarinoides*) (Lycopodiaceae). Shows modest antiacetylcholinesterase activity. Needles (CH₂Cl₂/Me₂CO). Mp 251-253° (249-250°). [α]_D²⁵ -26.9 (c, 0.2056 in CHCl₃) (-25.3).

N¹³-Oxide: Huperzine N¹³-oxide

C₁₇H₂₂N₂O₂ 286.373
Alkaloid from *Lycopodium casuarinoides*. Powder. [α]_D²⁰ -8.1 (c, 0.73 in MeOH). λ_{\max} 234 (log ϵ 4); 306 (log ϵ 3.84) (MeOH).

N-De-Me: N-Demethylhuperzine

C₁₆H₂₀N₂O 256.347
Alkaloid from aerial parts of *Lycopodium casuarinoides* (Lycopodiaceae). Exhibits anticholinesterase activity. Amorph. powder.

N,N-Di-de-Me: Huperzine C

[163089-71-2]
C₁₅H₁₈N₂O 242.32

Alkaloid from *Lycopodium casuarinoides* (*Lycopodium casuarinoides*) (Lycopodiaceae). Shows antiacetylcholinesterase activity. Cryst. (Me₂CO). Mp 233-235°.


16-Hydroxy: Huperzine D

[163047-27-6]
C₁₇H₂₂N₂O₂ 286.373
Alkaloid from *Lycopodium casuarinoides* (*Lycopodium casuarinoides*) (Lycopodiaceae). Cryst. (CH₂Cl₂/MeOH). Mp 269-271°. No antiacetylcholinesterase activity.

Shen, Y.-C. *et al.*, *J. Nat. Prod.*, 1994, **57**, 824-826 (*isol, pmr, cmr, N-Demethylhuperzine*)
Liu, J.-S. *et al.*, *Phytochemistry*, 1994, **37**, 1759-1761 (*Huperzine C, Huperzine D*)
Yin, S. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 138-143 (*isol, pmr, cmr, N-oxide*)

Huperzine C

H-389




C₁₆H₂₄N₂ 244.379
Alkaloid from *Huperzia serrata*. Mp 42-44°. λ_{\max} 268 (log ϵ 3.6) (no solvent reported).

Yuan, S.-Q. *et al.*, *Yaoxue Xuebao*, 2004, **39**, 116-118 (*isol, pmr, cmr*)

Hyatelactam

H-390

[894786-47-1]




C₂₇H₃₉NO₃ 425.61
Constit. of *Hyatella intestinalis*. Amorph. solid. [α]_D²⁵ +56 (c, 0.1 in CHCl₃). λ_{\max} 217 (ϵ 7900); 278 (ϵ 2500) (MeOH).

Hernández-Guerrero, C.J. *et al.*, *Tetrahedron*, 2006, **62**, 5392-5400 (*Hyatelactam*)

Hyderabadine

H-391


[88607-18-5]



C₂₁H₂₈N₂O₂ 340.464
Alkaloid from the leaves of *Ervatamia coronaria* (Apocynaceae). Amorph.


Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 1983, **38**, 1310 (*isol, uv, ir, pmr, cmr, ms*)

struct)

Hydnellin B H-392C₃₁H₃₂N₂O₁₁ 608.601


Closely related to Sarcodonin, S-76. Isol. from *Hydnellum suaveolens*. [α]_D²⁰ +12.6 (c, 0.7 in CHCl₃) (as tetra-Ac). Not obt. in pure state.

Hashimoto, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 912-914 (isol, pmr, cmr)

Hydrakouminol H-393
[105418-81-3]C₂₀H₂₄N₂O₂ 324.422

Isol. from a long-stored extract of *Gelsemium elegans* in hydrochloric acid. Prob. artifact.

Liu, Z. *et al.*, *Huaxue Xuebao*, 1986, **44**, 239; *CA*, **106**, 18909v

Hydrastine H-394C₂₁H₂₁NO₆ 383.4

All 4 stereoisomers as well as both racemates have been isol. from plants. Other spp. from which hydrastine(s) of unspecified config. have been isol. include *Corydalis fimbriifera*, *Stylomecon heterophylla*, *Berberis laurina* and *Hydrastis canadensis* (Papaveraceae, Berberidaceae). Antihypertensive agent. Antibacterial, sedative agent. Has been used clinically to treat uterine haemorrhagia. Claimed use in ophthalmology for pupil dilation and topical anaesthesia. Log P 2.76 (uncertain value) (calc). Some stereoisomers may be artifacts.

▶ MU6030000

(1R,9R)-form

(-)-α-Hydrastine. *Stylophylline* [4370-85-8]

Alkaloid from *Stylomecon heterophylla*

and *Hydrastis canadensis* (Papaveraceae). Mp 162-163.5°. [α]_D -141 (c, 1.0 in CHCl₃).

(1R,9S)-form

(-)-β-Hydrastine

[118-08-1]

Alkaloid from *Hydrastis canadensis* and *Berberis laurina* (Papaveraceae, Berberidaceae). Cryst. (EtOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 133-135°. [α]_D -68 (CHCl₃).

▶ Adverse human effects by ingestion.

Causes convulsions and gastro-intestinal disturbance. LD₅₀ (rat, ipr) 104 mg/kg. MU6030000

O^d-De-Me: Isohydrastidine

[76851-95-1]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Hydrastis canadensis* and *Corydalis pseudo-adunca* (Papaveraceae). Cryst. (EtOAc/hexane). Mp 163-166°. [α]_D²⁰ -77.7 (c, 0.6 in CHCl₃).

O^s-De-Me: Hydrastidine

[76851-94-0]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Hydrastis canadensis* (Papaveraceae). Cryst. (EtOAc/hexane). Mp 172-174°. [α]_D²⁰ -60.3 (c, 0.7 in CHCl₃).

(1S,9S)-form

(+)-α-Hydrastine

Alkaloid from *Fumaria schleicheri*, *Fumaria parviflora* and *Fumaria vaillantii* (Papaveraceae). Cryst. (EtOH). Mp 159-161°. [α]_D +127.7 (c, 1.1 in CHCl₃).

(1S,9R)-form

(+)-β-Hydrastine

Alkaloid from *Corydalis gortschakovii*, *Corydalis pseudo-adunca* and *Corydalis stricta* (Papaveraceae). Cryst. (MeOH). Mp 131-132°. [α]_D¹⁸ +63 (CHCl₃).

O^d-De-Me: Corfitaline. Corphthaline

[77492-95-6]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Corydalis pseudo-adunca* (Papaveraceae).

(1RS,9RS)-form

(±)-α-Hydrastine. *Hydrastine b*

Alkaloid from *Fumaria schleicheri* (Papaveraceae), also synthetic. Mp 137° Mp 149°. pK_a 6.63 (15°).

(1RS,9SR)-form

(±)-β-Hydrastine. *Hydrastine a*

Alkaloid from *Fumaria schleicheri* (Papaveraceae). Cryst. (EtOH). Mp 153° (151°).

Haworth, R.D. *et al.*, *J.C.S.*, 1950, 1776 (synth)

Ohta, M. *et al.*, *Tet. Lett.*, 1963, 859 (abs config)

Safe, S. *et al.*, *Can. J. Chem.*, 1964, **42**, 160 (pmr, config)

Snatzke, G. *et al.*, *Tetrahedron*, 1969, **25**, 5059 (ord)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (cmr)

Moniot, J.L. *et al.*, *J.A.C.S.*, 1976, **98**, 6714 (synth)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1976, 1221 (synth)

Moniot, J.L. *et al.*, *Heterocycles*, 1978, **9**, 145 (pmr)

Holland, H.L. *et al.*, *Can. J. Chem.*, 1979, **57**, 1588-1597 (biosynth)

Messina, I. *et al.*, *Gazz. Chim. Ital.*, 1980, **110**, 539 (*Hydrastidine, Isohydrastidine*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1980, 851 (*Corphthaline*)

Blaskó, G. *et al.*, *J. Nat. Prod.*, 1982, **45**, 105 (occur)

Shono, T. *et al.*, *J.O.C.*, 1983, **48**, 1621 (synth)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 869 (synth)

Rajnikant, *et al.*, *Acta Cryst. C*, 1993, **49**, 504 (cryst struct)

Martindale. The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 1377

Seger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**, 882-886 (pmr, cmr)

1,2-Hydrazinedicarboxylic acid, 9CI H-395

Bicarbamie acid, 8CI. *Hydrazodicarboxylic acid*. *Hydrazoformic acid*. *Hydrazodiformic acid*. *Hydrazobisformic acid* [5814-86-8]

HOOCNHNHCOOH

C₂H₄N₂O₄ 120.065

Not isol. in free state. Ester derivs. used as dispersants in lubricating oils and as blowing agents for polymer foams.

Monoamide: 2-(Aminocarbonyl)hydrazinecarboxylic acid, 9CI

C₂H₅N₃O₃ 119.08


Constit. of the seeds of *Butea monosperma*.

Sharma, S. *et al.*, *Indian J. Chem., Sect. B*, 1991, **30**, 715-716 (isol, amide)

Hydrocotarnine H-396

5,6,7,8-Tetrahydro-4-methoxy-6-methyl-1,3-dioxolo[4,5-g]isoquinoline, 9CI.

1,2,3,4-Tetrahydro-8-methoxy-2-methyl-6,7-methylenedioxyisoquinoline [550-10-7]

C₁₂H₁₅NO₃ 221.255

Alkaloid from opium (*Papaver somniferum*). (Papaveraceae). Reduction prod. of Cotarnine in N-292. Plates (petrol). Mp 55.5-56.5°. Bp₁₅ 182-183°.

▶ JI4825000

Hydrochloride:

Cryst. (EtOH). Mp 215-217° Mp 247-249° dec.

Hydrobromide: Mp 237-238° (229°).

Picrate: Mp 173°.

Hesse, O. *et al.*, *Annalen*, (Suppl.), 1872, **8**, 261 (isol)

Beckett, G.H. *et al.*, *J.C.S.*, 1875, **28**, 573 (synth)

Bandow, E. *et al.*, *Ber.*, 1898, **31**, 1577 (synth)

Pyman, F.L. *et al.*, *J.C.S.*, 1912, **101**, 1595 (synth)

Dey, B.B. *et al.*, *J. Indian Chem. Soc.*, 1935, **12**, 421 (synth)



Clayson, D.B. *et al.*, *J.C.S.*, 1949, 2016 (synth)

Schneider, W. *et al.*, *Annalen*, 1958, **615**, 34 (synth)

- Habermehl, G. *et al.*, *Annalen*, 1970, **742**, 138 (ms)
 Charnock, G.A. *et al.*, *J.C.S. Perkin 2*, 1972, 856 (synth, pmr, ms)
 Göber, B. *et al.*, *Pharmazie*, 1973, **28**, 221 (synth)
 Harsányi, K. *et al.*, *Annalen*, 1975, 1201 (synth)
 Minamikawa, J. *et al.*, *Can. J. Chem.*, 1979, **57**, 1720 (synth)

Hydrocotyline H-397C₂₂H₃₃NO₈ 439.505Struct. unknown. Alkaloid from *Hydrocotyle asiatica*.Basu, N.K. *et al.*, *Q. J. Pharm. Pharmacol.*, 1947, **20**, 135-136; *CA*, **42**, 1025g (isol)**Hydrohydrastinine** H-398


5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinoline, 9CI. 1,2,3,4-Tetrahydro-2-methyl-6,7-methylenedioxyisoquinoline [494-55-3]

C₁₁H₁₃NO₂ 191.229Alkaloid from *Corydalis cava* and *Corydalis tuberosa* (Papaveraceae). Degradn. prod. of hydrastine and cotarnine. Needles (petrol). Mp 66° (61-62°).*Hydrochloride:*Cryst. (H₂O or EtOH). Mp 276-278° (263-269°, 274°).*Hydrobromide:* Mp 276-278°.*Picrate:* Mp 175-176°.Pyman, F.L. *et al.*, *J.C.S.*, 1912, **101**, 1595 (synth)Späth, E. *et al.*, *Ber.*, 1931, **64**, 1131 (isol)Clayson, D.B. *et al.*, *J.C.S.*, 1949, 2016 (synth)Schneider, W. *et al.*, *Annalen*, 1958, **615**, 34 (synth)Bobbitt, J.M. *et al.*, *J.O.C.*, 1967, **32**, 2225 (synth)Habermehl, G. *et al.*, *Annalen*, 1970, **742**, 138 (ms)Göber, B. *et al.*, *Pharmazie*, 1973, **28**, 221 (synth)Moniot, J.L. *et al.*, *Heterocycles*, 1978, **9**, 145 (pmr)Moniot, J.L. *et al.*, *J.O.C.*, 1979, **44**, 4337 (synth)Ruchirawat, S. *et al.*, *Synth. Commun.*, 1984, **14**, 1221 (synth, pmr, ms)Takano, S. *et al.*, *Heterocycles*, 1993, **35**, 47 (synth)**7-Hydroperoxy-17-hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-5,13,19-triene-1,21-dione** H-399

Relative configuration

C₂₈H₃₅NO₅ 465.588Metab. from an unidentified *Daldinia* sp. Amorph. solid. [α]_D +16.9 (c, 0.09 in MeOH).Buchanan, M.S. *et al.*, *Phytochemistry*, 1996, **41**, 821 (isol, pmr, cmr, ms, struct)**3-(Hydroxyacetyl)-1H-indole** H-400

2-Hydroxy-1-(1H-indol-3-yl)ethanone, 9CI [2400-51-3]

C₁₀H₉NO₂ 175.187Alkaloid from the red marine alga *Prionitis lanceolata*. Also isol. from the sponge *Tedania ignis* and from liquid cultures of the fungus *Lactarius deliciosus*. Plant growth regulator. CNS stimulant. Needles (EtOAc); cryst. (H₂O or C₆H₆). Mp 162-163° Mp 173-174°. λ_{max} 212 (ε 10232); 240 (ε 5500); 294 (ε 4466) (MeOH) (Berdy).

▶ KM5779000

*Oxime:*C₁₀H₁₀N₂O₂ 190.201

Mp 116-118°.

O-Ac: [27536-15-8]C₁₂H₁₁NO₃ 217.224Cryst. (H₂O). Mp 140°.*O-Benzoyl:* [2400-52-4]C₁₇H₁₃NO₃ 279.295

Cryst. (MeOH). Mp 187-189°.

*N,O-Di-Ac:*C₁₄H₁₃NO₄ 259.261


Oil. Mp 0° approx.

5-Hydroxy-5-Hydroxy-3-(hydroxyacetyl)-1H-indole. Hyrtiosin A†

[132922-98-6]

C₁₀H₉NO₃ 191.186Alkaloid from the marine sponge *Hyrtios erecta*. Needles. Mp 196-197°. λ_{max} 215 (ε 59200); 252 (ε 35000); 270 (ε 22400); 303 (ε 23300) (MeOH) (Berdy).

[34951-71-8, 34951-79-6]

Suvorov, N.N. *et al.*, *Khim. Geterotsikl. Soedin.*, 1965, 265; 1971, 778; *Chem. Heterocycl. Compd.*, 1965, 173; 1971, 725 (synth, oxime, ir)*U.K. Pat.*, 1970, 1 183 919; *CA*, **72**, 132517v (synth, O-Ac)Vereshchagin, A.L. *et al.*, *Khim. Geterotsikl. Soedin.*, 1983, 46; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1983, 40 (O-Ac)Tsuji, S. *et al.*, *J.O.C.*, 1988, **53**, 5446 (synth)Bernart, M. *et al.*, *Phytochemistry*, 1990, **29**, 3697 (isol)Kobayashi, J. *et al.*, *Tetrahedron*, 1990, **46**, 7699 (Hyrtiosin A)Dillman, R.L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1056 (isol, uv, pmr, cmr, ms)Ayer, W.A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 839 (isol, pmr, ms, struct)Böhlendorf, B. *et al.*, *Annalen*, 1996, 49 (isol, ir, pmr, cmr, ms)**2-(Hydroxyacetyl)-1H-pyrrole** H-401C₆H₇NO₂ 125.127


N-(2-Acetoxyethyl), O-Ac: 2-(Acetoxyacetyl)-1-(2-acetoxyethyl)-1H-pyrrole [136396-66-2]

C₁₂H₁₅NO₅ 253.254Constit. of the red alga *Gracilariopsis lemaneiformis*. Oil.

N-(2-Acetoxypropyl), O-Ac: 2-(Acetoxyacetyl)-1-(2-acetoxypropyl)-1H-pyrrole [136396-64-0]

C₁₃H₁₇NO₅ 267.281Constit. of red alga *Gracilariopsis lemaneiformis*. Oil. [α]_D²³ -47.2 (c, 0.29 in MeOH).Jiang, Z.D. *et al.*, *J. Nat. Prod.*, 1991, **54**, 403-407 (isol, uv, ir, pmr, cmr, ms, struct, derivs)**1-Hydroxyacridone** H-402

1-Hydroxy-9(10H)-acridinone, 9CI [65582-54-9]

C₁₃H₉NO₂ 211.22Constit. of the roots of *Boenninghausenia albiflora* (Rutaceae). Yellow needles (Me₂CO). Mp 280° (subl., dec.).

N-Me: 1-Hydroxy-10-methylacridone

[16584-54-6]

C₁₄H₁₁NO₂ 225.246Alkaloid from the roots of *Boenninghausenia albiflora* and *Ruta graveolens* (rue). Also isol. from the callus cultures obt. from the meristematic cells of *Ruta graveolens* (Rutaceae). Yellow needles (Me₂CO). Mp 192-194°.

Me ether, N-Me: 1-Methoxy-10-methylacridone


[16584-58-0]

C₁₅H₁₃NO₂ 239.273Alkaloid from a new Australian species, assigned as aff. *Samadera bidwillii*. Yellow amorph. solid. Mp 164°.Hughes, G.K. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 206 (synth)Reisch, J. *et al.*, *Experientia*, 1971, **27**, 1005 (deriv, isol, pmr, struct)Scharlemann, W. *et al.*, *Z. Naturforsch., B*, 1972, **27**, 806 (deriv, isol, uv)Rózsa, Zs. *et al.*, *Pharmazie*, 1975, **30**, 753 (isol, deriv)Adams, J.H. *et al.*, *J.C.S. Perkin 1*, 1977, 2173 (deriv, synth, uv, ir, pmr)Rózsa, Zs. *et al.*, *Phytochemistry*, 1978, **17**, 169 (uv, ir, pmr, ms, deriv, isol, synth)Gibbons, S. *et al.*, *Phytochemistry*, 1997, **44**, 1109 (1-Methoxy-10-methylacridone)

Coppola, G.M. *et al.*, *Org. Prep. Proced. Int.*, 1999, **31**, 225-227 (*synth, ir, pmr*)

22-Hydroxyacuminatine H-403

7-(Hydroxymethyl)benz[6,7]indolizino[1,2-b]quinolin-11(13H)-one, 9CI [123086-77-1]



C₂₀H₁₄N₂O₂ 314.343

Prob. a rearranged indole alkaloid (camptothecin type). Alkaloid from the seeds of *Camptotheca acuminata*. Cytotoxic agent. Yellow cryst. (Me₂CO). Mp 258-260° dec. The parent Acuminatine appears to be unknown. λ_{max} 222 (log ε 4.45); 251 (log ε 4.51); 274 (sh) (log ε 4); 285 (log ε 4); 313 (sh) (log ε 3.91); 365 (sh) (log ε 4.21); 380 (log ε 4.23) (MeOH) (Berdy).

Aldehyde: **22-Oxoacuminatine**

[850415-08-6]

C₂₀H₁₂N₂O₂ 312.327

Alkaloid from the root bark of *Camptotheca acuminata*. Amorph. yellow powder. λ_{max} 221 (log ε 4.43); 251 (log ε 4.38); 285 (log ε 4.01); 381 (log ε 4.26) (MeOH).

Lin, L.-Z. *et al.*, *Phytochemistry*, 1989, **28**, 1295-1297 (*isol, uv, pmr*)

Zhang, Z. *et al.*, *Planta Med.*, 2004, **70**, 1216-1221 (**22-Oxoacuminatine**)

Ma, Z. *et al.*, *Tet. Lett.*, 2004, **45**, 6721-6723 (*synth*)


Xiao, X. *et al.*, *J. Med. Chem.*, 2006, **49**, 1408-1412 (*synth, pharmacol*)

Babjak, M. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 407-409 (*synth*)

Zhou, H.-B. *et al.*, *J.O.C.*, 2007, **72**, 6270-6272 (*synth*)

16-Hydroxyalloibogamine H-404

17-Hydroxyalloibogamine



C₁₉H₂₄ N₂O 296.411

(±)-form


Minor alkaloid from the stem and root of *Strychnos ngouniensis* (Loganiaceae). Amorph. First natural *allo*-iboga alkaloid.

Massiot, G. *et al.*, *Chem. Comm.*, 1983, 1018-1019 (*uv, ir, pmr, cmr, ms, cryst struct*)

Hydroxyaphylline

Oxaphylline

H-405



C₁₅H₂₄N₂O₂ 264.367

Probable struct. Alkaloid from *Anabasis aphylla* (Chenopodiaceae). Needles (Me₂CO). Mp 165-167°. [α]_D +39.2. Dehydrates to Aphyllidine salts on treatment with acids.


Methiodide: Mp 223-225°.

Sadykov, A.S. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 1736; *J. Gen. Chem. USSR (Engl. Transl.)*, 1723 (*isol, struct*)

Sadykov, A.S. *et al.*, *Abh. Dtsch. Akad. Wiss. Berlin. Kl. Chem., Geol. Biol.*, 1966, 279; *CA*, **67**, 32865k (*isol*)

16-Hydroxyaspidofractinine H-406

Aspidofractinin-16-ol



Absolute Configuration

C₁₉H₂₄N₂O 296.411


Alkaloid from leaves of *Kopsia officinalis*. Amorph. powder. [α]_D²⁶ -15.7 (c, 0.33 in CHCl₃). λ_{max} 242 ; 290 ; 391 (CHCl₃).

Zhou, H. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 515-519 (*isol, pmr, cmr*)

7-Hydroxy-2H-1,4-benzothiazin-3(4H)-one H-407

9CI

[91375-75-6]



C₈H₇NO₂S 181.215

Prod. by the marine-derived *Halomonas* sp. RK377. Cryst. (Me₂CO). Mp 217°.

Jones, G.H. *et al.*, *J. Med. Chem.*, 1987, **30**, 295-303 (*synth*)

Crescenzi, O. *et al.*, *Gazz. Chim. Ital.*, 1990, **120**, 21-24 (*synth*)


Liang, L. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*isol, pmr, cmr, ms*)

6-Hydroxy-5-benzothiazoleacetic acid H-408

C 304A. M 4582. Antibiotic C 304A.

Antibiotic M 4582

[115585-81-4]



C₉H₇NO₃S 209.225

Prod. by an *Actinosynema* sp. and *Paecilomyces lilacinus*. Aldose reductase inhibitor. λ_{max} 226 ; 247 ; 274 ; 291 ; 305 (MeOH) (Berdy).

Japan. Pat., 1988, 88 30 493; *CA*, **109**, 91257 (*isol*)


Japan. Pat., 1989, 89 181 793; *CA*, **112**, 96995 (*isol*)

2-Hydroxy-2H-1,4-benzoxazin-3(4H)-one H-409

9CI

Blepharigenin. HBOA

[27779-92-6]



(R)-form

C₈H₇NO₃ 165.148

(R)-form

Constit. of juvenile wheat (*Triticum aestivum*). Biosynth. intermed. of DIBOA. Yellow needles. Mp 207°.

O-β-D-Glucopyranoside: **Blepharin**.

HBOA-Glc

[27625-86-1]

C₁₄H₁₇NO₈ 327.29

Isol. from *Blepharis edulis* (Acanthaceae) and seedlings of rye and sweet corn (*Zea mays*). Mp 226-227°. [α]_D +127.3 (H₂O).

O-Ac: [23520-35-6]

C₁₀H₉NO₄ 207.185

Cryst. (MeOH). Mp 177°.

O-Benzoyl:

C₁₅H₁₁NO₄ 269.256

Cryst. (MeOH/EtOAc). Mp 207°.

Me ether: 2-Methoxy-2H-1,4-benzoxazin-3(4H)-one, 9CI

[27392-93-4]

C₉H₉NO₃ 179.175

Cryst. (MeOH aq.). Mp 168-171°.

Me ether, N-Me: 2-Methoxy-4-methyl-2H-1,4-benzoxazin-3(4H)-one

C₁₀H₁₁NO₃ 193.202

Cryst. (MeOH aq.). Mp 71°. Bp₁ 180-185°.

N-Hydroxy: 2,4-Dihydroxy-2H-1,4-benzoxazin-3(4H)-one. DIBOA

[17359-54-5]

C₈H₇NO₄ 181.148

Isol. from seedlings of rye (*Secale cereale*) and sweet corn (*Zea mays*). Antialgal agent. Cryst. (Et₂O/cyclohexane). Mp 152°.

▶DM3850000

N-Hydroxy, 2-Me ether: 4-Hydroxy-2-methoxy-2H-1,4-benzoxazin-3(4H)-one, 9CI

[55574-10-2]

C₉H₉NO₄ 195.174

Cryst. Mp 136-138°.

6-Hydroxy, 2-O-β-D-glucopyranoside: 6-Hydroxyblepharin

[651728-01-7]

C₁₄H₁₇NO₉ 343.29

Constit. of *Lamium galeobdolon*. Amorph. solid.

(S)-form

N-Hydroxy, 2-O-β-D-glucopyranoside:

DIBOA-Glc. GDIMBOA

[155835-54-4]

[157241-60-6, 22260-47-5]

C₁₄H₁₇NO₉ 343.29

Isol. from seedlings of rye (*Secale cereale*), sweet corn (*Zea mays*) and seeds of *Acanthus mollis*. Cryst. Mp 186.5-187°. [α]_D²⁵ +106 (c, 1 in H₂O). λ_{max} 254 (ε 8300); 281 (ε 6380) (prob. MeOH) (Derep).


7-Chloro, O-β-D-glucopyranoside:

C₁₄H₁₆ClNO₈ 361.735Constit. of *Acanthus ilicifolius*. Amorph. powder. [α]_D²⁶ +198 (c, 0.33 in DMSO).

7-Chloro, N-hydroxy, O-β-D-glucopyranoside: 7-Cl-DIBOA-Glc

C₁₄H₁₆ClNO₉ 377.734Constit. of *Acanthus ebracteatus*.Amorph. powder. [α]_D²² +18.6 (c, 0.86 in DMSO).Lal, J.B. *et al.*, *J. Indian Chem. Soc.*, 1940, **17**, 269 (isol, deriv)Virtanen, A.L. *et al.*, *Acta Chem. Scand.*, 1960, **14**, 499; 502; 504 (synth, uv)Chatterjee, A. *et al.*, *Chem. Ind. (London)*, 1969, 328 (synth)Hofman, J. *et al.*, *Eur. J. Biochem.*, 1969, **8**, 109-112 (HBOA-Glc)Tipton, C.L. *et al.*, *Phytochemistry*, 1973, **12**, 347 (biosynth)Wilson, A.S. *et al.*, *Cryst. Struct. Commun.*, 1982, **11**, 809-813 (DIBOA, cryst struct)Wolf, R.B. *et al.*, *J. Nat. Prod.*, 1985, **48**, 59-63 (DIBOA-Glc)Atkinson, J. *et al.*, *J.O.C.*, 1991, **56**, 1788 (synth, pmr, ms, uv, ir)Tietze, L.F. *et al.*, *Synthesis*, 1991, 1118 (synth, pmr, abs config)Sicker, D. *et al.*, *Synthesis*, 1993, 771 (synth)Hartenstein, H. *et al.*, *Phytochemistry*, 1994, **35**, 827-828 (DIBOA-Glc, isol, abs config)Bravo, H.R. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 1569 (DIBOA, activity)Kamperdick, C. *et al.*, *Pharmazie*, 1997, **52**, 965-966 (isol, pmr, cmr)Tays, K. *et al.*, *Synth. Commun.*, 1998, **28**, 903-912 (N-hydroxy)Tanabe, J. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1614-1617 (isol, biosynth)Baumeler, A. *et al.*, *Phytochemistry*, 2000, **53**, 213-222 (isol, pmr)Kanchanapoom, T. *et al.*, *Phytochemistry*, 2001, **58**, 637-640; 811-817 (7-chloro derivs)Alipieva, K.I. *et al.*, *Phytochemistry*, 2003, **64**, 1413-1417 (6-Hydroxyblepharin)Macias, F. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 991-1000 (synth, DIBOA-Glc, isol, uv, pmr, cmr)Schullehner, K. *et al.*, *Phytochemistry*, 2008, **69**, 2668-2677 (biosynth)**N-[4-(4-Hydroxybenzoylamino)butyl]-1,3-dimethylumazine-6-carboxamide** H-410

6-[[[4-(4-Hydroxybenzoyl)amino]butyl]amino]carbonyl]-1,3-dimethyl-2,4(1H,3H)-pteridinedione




n = 4

C₂₀H₂₂N₆O₅ 426.431Isol. from the leech *Limnatis nilotica*.Cryst. (CHCl₃/MeOH aq.). Mp 256-257°.λ_{max} 194; 249; 337 (MeOH).

Homologue (n = 5): N-[5-(4-Hydroxybenzoylamino)pentyl]-1,3-dimethylumazine-6-carboxamide

C₂₁H₂₄N₆O₅ 440.458Isol. from *Limnatis nilotica*. Beige powder. Mp 268-270° dec. λ_{max} 194; 249; 337 (MeOH).Voerman, G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 938-941 (isol, synth, pmr, cmr, ms)**1-(4-Hydroxybenzoyl)-β-carboline** H-4111-(4-Hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₁**

[1007893-74-4]

C₁₈H₁₂N₂O₂ 288.305Alkaloid from a *Eudistoma* sp. Amorph. yellow solid. λ_{max} 223 (log ε 3.46); 301 (log ε 3.25); 382 (log ε 2.87) (MeOH).6-Bromo: 6-Bromo-1-(4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₂**

[1007893-75-5]

C₁₈H₁₁BrN₂O₂ 367.201Alkaloid from a *Eudistoma* sp. Amorph. yellow solid. λ_{max} 233 (log ε 3.74); 307 (log ε 3.75); 387 (log ε 3.27) (MeOH).3'-Bromo: 1-(3-Bromo-4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₃**

[1007893-76-6]

C₁₈H₁₁BrN₂O₂ 367.201Alkaloid from a *Eudistoma* sp. Amorph. yellow solid. λ_{max} 240 (log ε 3.88); 306 (log ε 3.91); 389 (log ε 3.57) (MeOH).3',6-Dibromo: 6-Bromo-1-(3-bromo-4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₄**

[1007893-77-7]

C₁₈H₁₀Br₂N₂O₂ 446.097Alkaloid from a *Eudistoma* sp. Amorph. yellow solid. λ_{max} 240 (log ε 3.88); 306 (log ε 3.91); 389 (log ε 3.57) (MeOH).

3',5'-Dibromo: 1-(3,5-Dibromo-4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole.

Eudistomin Y₅

[1007893-78-8]

C₁₈H₁₀Br₂N₂O₂ 446.097Alkaloid from a *Eudistoma* sp. Amorph. yellow solid. λ_{max} 224 (log ε 3.79); 296 (log ε 3.61); 388 (log ε 3.5) (MeOH).3',5',6-Tribromo: 6-Bromo-1-(3,5-dibromo-4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₆**


[1007893-79-9]

C₁₈H₉Br₃N₂O₂ 524.993Alkaloid from a *Eudistoma* sp.Amorph. yellow solid. λ_{max} 239 (log ε 3.97); 304 (log ε 3.95); 392 (log ε 3.75) (MeOH).3',5',7-Tribromo: 7-Bromo-1-(3,5-dibromo-4-hydroxybenzoyl)-9H-pyrido[3,4-b]indole. **Eudistomin Y₇**


[1007893-80-2]

C₁₈H₉Br₃N₂O₂ 524.993Alkaloid from a *Eudistoma* sp.Amorph. yellow solid. λ_{max} 225 (log ε 3.68); 299 (log ε 3.42); 392 (log ε 3.28) (MeOH).Wang, W. *et al.*, *J. Nat. Prod.*, 2008, **71**, 163-166 (isol, pmr, cmr)**2-(4-Hydroxybenzoyl)-4(5)-(4-hydroxyphenyl)-1H-imidazole** H-412

[251907-19-4]



1H-form

C₁₆H₁₂N₂O₃ 280.282Alkaloid from the red ascidian *Botryllus leachi*. Amorph. yellow solid. λ_{max} 248 (ε 7600); 305 (ε 6000); 360 (ε 13700) (MeOH).Duran, R. *et al.*, *Tetrahedron*, 1999, **55**, 13225-13232 (isol)Mahboobi, S. *et al.*, *Monatsh. Chem.*, 2004, **135**, 333-342 (synth, pmr, tautom)***Agelenopsis aperta* Hydroxybenzoylpentamine toxins** H-413

Agel 379 m = n = p = 3, q = 4
 Agel 379a m = 4, n = p = q = 3
 Agel 379b m = n = q = 3, p = 4
 Agel 379c m = p = q = 3, n = 4

See also Agel 452, A-176 for related toxins. Acylpolyamine alkaloid complex isol. from the venom of the spider *Agelenopsis aperta*.

Agel 379

AG 379

C₂₀H₃₇N₅O₂ 379.545N²-Hydroxy: **Agel 395a**. AG 395aC₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta*.4-Deoxy, 2,5-dihydroxy: **Agel 395b**. AG 395b. HO 395C₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta* and *Hololena curta*.**Agel 379a**

AG 379a

C₂₀H₃₇N₅O₂ 379.5454-Deoxy, 2,5-dihydroxy: *Agel 395c*. *AG 395c*C₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta*.**Agel 379b***AG 379b*C₂₀H₃₇N₅O₂ 379.545N²-Hydroxy: *Agel 395*. *AG 395*C₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta*.


4-Deoxy, 2,5-dihydroxy:

C₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta*.**Agel 379c***AG 379c*

4-Deoxy, 2,5-dihydroxy:

C₂₀H₃₇N₅O₃ 395.544From *Agelenopsis aperta*.Chesnov, S. et al., *Helv. Chim. Acta*, 2001, **84**, 2178-2197; 2002, **85**, 2827-2846 (*isol, synth, ms*)**4-Hydroxybenzyl alcohol, 8CI H-414**4-Hydroxybenzenemethanol, 9CI. α -4-Dihydroxytoluene. *p*-Hydroxymethylphenol. *Gastrodigenin*

[623-05-2]

C₇H₈O₂ 124.139Constit. of muskmelon (*Cucurbita moschata*) seedlings and from infected bulbs of *Orchis militaris* and *Loroglossum hircinum* (preferred genus name *Himantoglossum*). Cofactor for indoleacetic acid oxidase. Prisms or needles (H₂O). Mp 124.5-125.5°. p*K*_{a1} 9.73 (25°).4-O-[3-Pyridinecarbonyl-(\rightarrow 5)- β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: *Cucurbitoside L*C₂₄H₂₉NO₁₂ 523.493Constit. of the seeds of *Cucurbita pepo*. Amorph. solid. [α]_D²⁵ -80.6 (c, 0.4 in MeOH). λ_{\max} 220 (log ϵ 4.19); 264 (log ϵ 3.5) (MeOH).Li, W. et al., *J. Nat. Prod.*, 2005, **68**, 1754-1757 (*Cucurbitosides K-M*)**2-Hydroxybenzylamine H-415**2-(Aminomethyl)phenol, 9CI. α -Amino-*o*-cresol, 8CI. *Salicylamine* [932-30-9]C₇H₉NO 123.154Found in flowers of *Reseda odorata*. Cryst. (EtOH/petrol) or solid. Mp 97-99° Mp 129° (prob. refers to a hydrate). Sublimes.Sorensen, H. et al., *Phytochemistry*, 1970, **9**, 865-870 (*isol*)Olsen, O. et al., *Phytochemistry*, 1980, **19**, 1783-1787 (*isol, pmr, cmr*)**3-Hydroxybenzylamine H-416**3-(Aminomethyl)phenol, 9CI. α -Amino-*m*-cresol, 8CI

[73604-31-6]

C₇H₉NO 123.154Isol. from *Reseda media*. Cryst. (MeOH). Mp 173-174°.*Me ether*: 3-Methoxybenzylamine

[5071-96-5]

C₈H₁₁NO 137.181Bp₆ 103-104°.

▶ DP5550000

Me ether, N,N-di-*Me*: 3-Methoxy-N,N-dimethylbenzenemethanamine, 9CI

[15184-99-3]

C₁₀H₁₅NO 165.235Pale yellow liq. Bp₁₃ 105°. *n*_D²⁰ 1.5140.N,N-Di-*Me*: 3-(Dimethylaminomethyl)-phenol

[60760-04-5]

C₉H₁₃NO 151.208Prisms (C₆H₆). Mp 108°.Stedman, E. et al., *J.C.S.*, 1927, 1902-1906(N,N-di-*Me*, *Me ether* N,N-di-*Me*)Shoppee, C.W. et al., *J.C.S.*, 1932, 696-711

(Me ether)

Ger. Pat., 1953, 865 455; *CA*, **52**, 17182 (*synth*)Olsen, O. et al., *Phytochemistry*, 1980, **19**, 1783

(isol, pmr, cmr)

Yamagiwa, Y. et al., *Tetrahedron*, 1987, **43**,3387 (Me ether N,N-di-*Me*)Vallée, C. et al., *Adv. Synth. Catal.*, 2005, **347**,1827-1834 (N,N-di-*Me*)**4-Hydroxybenzylamine H-417**4-(Aminomethyl)phenol, 9CI. α -Amino-*p*-cresol

[696-60-6]

C₇H₉NO 123.154Isol. from seeds of *Sinapis alba* (white mustard) (Brassicaceae) and *Sinapis arvensis*. Plates + 1H₂O (H₂O). Mp 114-115° dec.*Hydrobromide*: [90430-14-1]

Solid. Mp 175-180°.

N-Ac: N-Acetyl-4-hydroxybenzylamine.

Antibiotic BMY 40660. BMY 40660



[34185-04-1]

C₉H₁₁NO₂ 165.191Isol. from *Actinomadura verrucosopora*. Active against gram-positive bacteria. Sol. MeOH; poorly sol.CHCl₃. Mp 113-117°. λ_{\max} 292 (ϵ 5620); 367 (ϵ 1150) (MeOH/NaOH)(Derep). λ_{\max} 277 (ϵ 4790); 310 (sh) (ϵ)(MeOH) (Derep). λ_{\max} 367 (ϵ 1150)

(MeOH) (Berdy).

Olsen, O. et al., *Phytochemistry*, 1980, **19**, 1783

(isol, pmr, cmr)

Larsen, P.O. et al., *Phytochemistry*, 1984, **23**,895 (*isol*)Chichak, K.S. et al., *J.O.C.*, 2005, **70**, 7956-7962 (*synth*)**8-(4-Hydroxybenzyl)-2,3,9,10-berbintetrol H-418**(8*R**,13*aR**)-formC₂₄H₂₃NO₅ 405.449**(8*R*,13*aR*)-form**O²-*Me*, O⁹- α -D-glucopyranoside: [145134-72-1]C₃₁H₃₅NO₁₀ 581.618Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (MeOH) (as hepta-Ac). Mp 122.5-125° (hepta-Ac). CAS No. refers to hepta-Ac. λ_{\max} 230 (log ϵ 4.27); 276 (log ϵ 3.69) (MeOH) (as hepta-Ac).**(8*R*,13*aS*)-form**O²-*Me*, O¹⁰- β -xylopyranoside: [145134-69-6]C₃₀H₃₃NO₉ 551.592Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (MeOH). Mp 163-166°. [α]₅₇₈²² +106.4 (c, 0.52 in MeOH). λ_{\max} 231 (log ϵ 3.98); 233 (log ϵ 3.89); 282 (log ϵ 3.52) (MeOH).O²-*Me*, O¹⁰- β -glucopyranoside: [145134-70-9]C₃₁H₃₅NO₁₀ 581.618Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. red-brownish solid (MeOH). Mp 150-152°. [α]₅₇₈²² +231.6 (c, 0.73 in MeOH). λ_{\max} 230 (log ϵ 3.96); 281 (log ϵ 3.58) (MeOH).O²-*Me*, O^{4'}- β -glucopyranoside: [145134-71-0]C₃₁H₃₅NO₁₀ 581.618Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (as hepta-Ac). Mp 150-153° (hepta-Ac). Reg. No. refers to hepta-Ac. λ_{\max} 230 (log ϵ 4.04); 277 (log ϵ 3.7) (MeOH) (as hepta-Ac).O³-*Me*, O¹⁰- β -glucopyranoside, N-oxide: [190953-12-9]C₃₁H₃₅NO₁₁ 597.618Alkaloid from leaves of *Aristolochia gigantea*. Amorph. yellow solid (MeOH). Mp 220-224° (dec.). [α]₅₇₈²⁵ -24.6 (c, 0.12 in MeOH). λ_{\max} 215 (log ϵ 3.22); 230 (log ϵ 3.31); 250 (log ϵ 3.3); 265 (log ϵ 3.29); 275 (log ϵ 3.31); 295 (log ϵ 3.31); 325 (sh) (log ϵ 3.08); 385 (log ϵ 2.88) (MeOH).Lopes, L.M.X. et al., *Phytochemistry*, 1992, **31**, 4005-4009; 1997, **45**, 431-435 (*isol, uv, ir, pmr, cmr, ms, struct*)**8-(4-Hydroxybenzyl)-2,3,10,11-berbintetrol H-419**(8*R**,13*aR**)-formC₂₄H₂₃NO₅ 405.449**(8*R*,13*aR*)-form**O²-*Me*: 8-(4-Hydroxybenzyl)-2-methoxy-3,10,11-berbintetrol [145134-66-3]

C₂₅H₂₅NO₅ 419.476

Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid. Mp 178-180°. [α]_D²² -65.3 (c, 2.17 in MeOH). λ_{\max} 203 (log ϵ 4.36); 290 (log ϵ 3.58) (MeOH).

O²-Me, O¹⁰- β -xylopyranoside: [145134-65-2]

C₃₀H₃₃NO₉ 551.592

Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (MeOH). Mp 165-168°. [α]_D²² +14 (c, 1.83 in MeOH). λ_{\max} 201 (log ϵ 4.73); 203 (log ϵ 4.74); 286 (log ϵ 3.49) (MeOH).

O²-Me, O¹⁰- β -glucopyranoside: [145134-64-1]

C₃₁H₃₅NO₁₀ 581.618

Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (MeOH). Mp 166-170°. [α]_D²² +52.2 (c, 1.11 in MeOH). λ_{\max} 203 (log ϵ 4.79); 285 (log ϵ 4.14) (MeOH).

(8R,13aS)-form [240430-85-7]

Alkaloid from *Gnetum parvifolium*. Needles (MeOH aq.) (as penta-Ac). Mp 166-168° (penta-Ac). λ_{\max} 275 (log ϵ 3.92) (EtOH) (penta-Ac).

O²-Me, O¹⁰- β -glucopyranoside: [145134-67-4]

C₃₁H₃₅NO₁₀ 581.618

Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (as hepta-Ac). Mp 124.5-127° (hepta-Ac). [α]_D²² -101.5 (c, 0.31 in MeOH) (hepta-Ac). CAS No. refers to hepta-Ac. λ_{\max} 203 (log ϵ 5.18); 281 (log ϵ 4.24) (MeOH) (hepta-Ac).

O³-Me, O¹⁰- β -glucopyranoside: [190952-98-8]

C₃₁H₃₅NO₁₀ 581.618

Alkaloid from leaves of *Aristolochia gigantea*. Amorph. yellow solid (MeOH). Mp 201-204°. [α]_D²⁵ -50.8 (c, 0.04 in MeOH). λ_{\max} 210 (log ϵ 3.65); 230 (log ϵ 3.71); 250 (log ϵ 3.69); 270 (log ϵ 3.71); 290 (log ϵ 3.71); 325 (log ϵ 3.63); 385 (log ϵ 3.22) (MeOH).


O²,O⁴-Di-Me: [145134-68-5]

C₂₆H₂₇NO₅ 433.503

Alkaloid from leaves of *Aristolochia gigantea* (Aristolochiaceae). Amorph. yellow solid (MeOH). Mp 191-193°. [α]_D²² -34.4 (c, 0.25 in MeOH). λ_{\max} 201 (log ϵ 4.35); 293 (log ϵ 3.77) (MeOH).

Lopes, L.M.X. *et al.*, *Phytochemistry*, 1992, **31**, 4005-4009; 1997, **45**, 431-435 (*Me ether derivs, isol, uv, ir, pmr, cmr, ms, struct*)

Xu, Q. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1025-1027 (*8R*,13aS*-form*)

8-(4-Hydroxybenzyl)-2,3,10-berbintrioI H-420C₂₄H₂₃NO₄ 389.45**(8R,13aR)-form**

O²,O³-Di-Me: **8-(4-Hydroxybenzyl)-2,3-dimethoxy-10-berbinol**

[195971-32-5]


C₂₆H₂₇NO₄ 417.504

Alkaloid from *Aristolochia constricta* (Aristolochiaceae). Amorph. yellow solid. Mp 150-162°. [α]_D²⁵ -65.3 (c, 1 in MeOH). λ_{\max} 290 (log ϵ 4.36) (MeOH).

Rastrelli, L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1065-1069 (*isol, uv, pmr, cmr*)

(4-Hydroxybenzyl)carbamic acid H-421

[(4-Hydroxyphenyl)methyl]carbamic acid. (4-Hydroxybenzyl)aminoformic acid

C₈H₉NO₃ 167.164

Benzyl ester: [75383-60-7]

C₁₅H₁₅NO₃ 257.288

Cryst. (C₆H₆). Mp 90.5-91° dec.

O- α -L-Rhamnopyranoside, *Et ester*:

[208346-80-9]

C₁₆H₂₃NO₇ 341.36

Isol. from seed pods of *Moringa oleifera* (horseradish tree). λ_{\max} 202; 222; 252 (MeOH).

O-(4-O-Acetyl- α -L-rhamnopyranoside),

Me ester: Niazicinin A

[163812-19-9]

C₁₇H₂₃NO₈ 369.371

Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). λ_{\max} 200; 223; 271 (MeOH).

O-(Tri-O-acetyl- α -L-rhamnopyranoside),

Me ester: [163812-18-8]C₂₁H₂₇NO₁₀ 453.445

Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). Isol. as two conformational (rotational) isomers. λ_{\max} 201; 222; 247 (MeOH).

O-(4-O-Acetyl- α -L-rhamnopyranoside),

Et ester (1): Niazimin A

[159768-73-7]

C₁₈H₂₅NO₈ 383.397

Constit. of *Moringa oleifera* (horseradish tree). Plates (CHCl₃/MeOH). Mp 168-170°.

O-(4-O-Acetyl- α -L-rhamnopyranoside),

*Et ester (2): Niazimin B*C₁₈H₂₅NO₈ 383.397

Constit. of *Moringa oleifera* (horseradish tree). Needles (CHCl₃/MeOH). Mp 182-184°. Conformational isomer of Niazimin A, differing in the orientation of the NH proton with respect to the carbonyl group.

O-(Tri-O-acetyl- α -L-rhamnopyranoside),

Et ester: [159733-93-4]C₂₂H₂₉NO₁₀ 467.472

Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). λ_{\max} 203; 222; 246 (MeOH).

Nakano, M. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2212-2216 (*synth, benzyl ester*)

Faizi, S. *et al.*, *J. C. S. Perkin I*, 1994, 3035-3040 (*Niazimin A, Niazimin B*)


Faizi, S. *et al.*, *Phytochemistry*, 1995, **38**, 957-963 (*isol, uv, pmr, cmr, derivs*)

Leuck, M. *et al.*, *Carbohydr. Res.*, 1998, **312**, 33-44 (*synth*)

Faizi, S. *et al.*, *Planta Med.*, 1998, **64**, 225-228 (*Et ester rhamnoside*)

2-(4-Hydroxybenzyl)imidazole H-422

4-(1H-Imidazol-2-ylmethyl)phenol. Semilepidine A

C₁₀H₁₀N₂O 174.202

Powder (CH₂Cl₂/MeOH). Mp 164-166°. Artifact.

O- β -D-Galactopyranoside: **Semilepidinose A**

C₁₆H₂₀N₂O₆ 336.344

Alkaloid from the seeds of *Lepidium sativum* (garden cress). Amorph. powder. Mp 188-190° dec.

3'-Methoxy: **2-(4-Hydroxy-3-methoxybenzyl)imidazole. Semilepidine B**

C₁₁H₁₂N₂O₂ 204.228

Powder (CH₂Cl₂/MeOH). Mp 187-190°. Artifact.

3'-Methoxy, O- β -D-galactopyranoside: **Semilepidinose B**

C₁₇H₂₂N₂O₇ 366.37

Alkaloid from the seeds of *Lepidium sativum* (garden cress). Amorph. powder. Mp 212-215° dec.


Maier, U.H. *et al.*, *Phytochemistry*, 1998, **49**, 1791-1795 (*isol, pmr, cmr, ms*)

5-(4-Hydroxybenzyl)-2,4-imidazolidinedione H-423

5-(p-Hydroxybenzyl)hydantoin

[58942-04-4]

[98819-07-9]



(S)-form

C₁₀H₁₀N₂O₃ 206.201**(S)-form** [40856-79-9][α]_D -165 (EtOH). [α]_D -292 (1M NaOH).**(±)-form** [67337-72-8]

Cryst. (EtOH). Mp 258-259°.


(ξ)-form

Isol. from a marine endophytic fungus No. 1893. No details are given for its isol. from the fungus.

Suzuki, T. *et al.*, *Agric. Biol. Chem.*, 1973, **37**,

411-416; 1976, **40**, 225-226 (*synth ord pmr*)
 Marton, J. et al., *J. Agric. Food Chem.*, 1993, **41**, 148-152 (*synth*)
 Chen, G. et al., *Tetrahedron*, 2003, **59**, 4907-4909 (*isol*)

1-(4-Hydroxybenzyl)-6,7-isoquinolinediol H-424
 6,7-Dihydroxy-1-(4-hydroxybenzyl)isoquinoline



C₁₆H₁₃NO₃ 267.284

6-Me ether: 7-Hydroxy-1-(4-hydroxybenzyl)-6-methoxyisoquinoline. **Juzirine** [64069-53-0]
 C₁₇H₁₅NO₃ 281.31
 Alkaloid from the leaves of *Zizyphus jujuba* (Chinese date). Cryst. (Me₂CO). Mp 203-205°.

6,7-Di-Me ether: 4-[(6,7-Dimethoxy-1-isoquinolinyl)methyl]phenol, 9CI. 1-(4-Hydroxybenzyl)-6,7-dimethoxyisoquinoline. **Crykonisine** [5544-58-1]
 C₁₈H₁₇NO₃ 295.337

Alkaloid from *Cryptocarya konishii* and *Machilus acuminatissima* (Lauraceae). Needles. Mp 235-236° dec.

Tri-Me ether: 6,7-Dimethoxy-1-(4-methoxybenzyl)isoquinoline [41498-25-3]
 [41498-26-4]

C₁₉H₁₉NO₃ 309.364

Alkaloid from *Ocotea macrophylla*. Cryst. (Et₂O). Mp 125°. λ_{max} 238 (log ε 4.77); 269 (log ε 3.78); 278 (log ε 3.76); 314 (log ε 3.57); 326 (log ε 3.63) (EtOH).

6,7-Methylene ether: 1-(4-Hydroxybenzyl)-6,7-methylenedioxyisoquinoline. **Neolitacuminine**

C₁₇H₁₃NO₃ 279.295

Alkaloid from the stem bark of *Neolitsea acuminatissima*. Amorph. powder. λ_{max} 261 (log ε 4.56); 298 (log ε 4.45); 330 (log ε 4.38) (MeOH).

6,7-Methylene, 4'-Me ether: 1-(4-Methoxybenzyl)-6,7-methylenedioxyisoquinoline [55857-56-2]
 C₁₈H₁₅NO₃ 293.321

Alkaloid from *Ocotea macrophylla*. Cryst. (CHCl₃). Mp 122°. λ_{max} 237 (log ε 4.77); 267 (log ε 3.84); 276 (log ε 3.85); 315 (log ε 3.65); 329 (log ε 3.74) (EtOH).

6,7-Methylene, 4'-Me ether, N-Me: 4-Methoxy-2-methyl-6,7-methylenedioxyisoquinolinium

C₁₉H₁₈NO₃⁺ 308.356

Quaternary alkaloid from *Doryphora sassafras*. Brown gum (as chloride).

λ_{max} 228 (log ε 4.18); 253 (log ε 4.43); 283 (log ε 3.6); 313 (log ε 3.78); 333 (sh) (log ε 3.7); 348 (sh) (log ε 3.6) (MeOH) (chloride).

Lu, S.-T. et al., *Yakugaku Zasshi*, 1967, **87**, 1278 (*Crykonisine*)

Coomes, R.M. et al., *J.O.C.*, 1973, **38**, 3701-3704 (*synth*)

Franca, N.C. et al., *Phytochemistry*, 1975, **14**, 1671-1672 (*Ocotea isolates*)


Ziyaev, R. et al., *Khim. Prir. Soedin.*, 1977, **13**, 239; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 204 (*Juzirine*)

Marsaioli, A.J. et al., *Phytochemistry*, 1978, **17**, 1655-1658 (*cmr*)

Carroll, A.R. et al., *J. Nat. Prod.*, 2001, **64**, 1572-1573 (*Doryphora isolate*)

Chang, F.R. et al., *J. Nat. Prod.*, 2002, **65**, 255-258 (*Neolitacuminone*)

2-(4-Hydroxybenzyl)-4-oxo-2-azetidinecarboxylic acid H-425
 2-[(4-Hydroxyphenyl)methyl]-4-oxo-2-azetidinecarboxylic acid. **Streptovercillone**



C₁₁H₁₁NO₄ 221.212


(S)-form [941687-54-3]

Prod. by *Streptomyces morookaense* strain SC1169. Antifungal agent.

Amorph. powder. [α]_D²⁰ +62.8 (c, 0.32 in MeOH). λ_{max} 225 (log ε 3.89); 278 (log ε 3.17) (MeOH).

Feng, N. et al., *J. Antibiot.*, 2007, **60**, 179-183 (*isol, pmr, cmr, ms*)

5-(4-Hydroxybenzyl)-2-pyridinecarboxylic acid H-426
 5-[(4-Hydroxyphenyl)methyl]-2-pyridinecarboxylic acid, 9CI. **Phenopicolinic acid** [56153-30-1]




C₁₃H₁₁NO₃ 229.235

Isol. from culture filtrates of the fungus *Paecilomyces*. Dopamine β-hydroxylase inhibitor; strong hypotensive action. Sol. Py, bases, DMSO; fairly sol. MeOH, EtOAc, H₂O; poorly sol. C₆H₆, hexane. Mp 222-226°. Log P 2.2 (calc). λ_{max} 271 (ε 11100) (HCl) (Berdy). λ_{max} 271 (ε 8000); 277 (ε 7300); 300 (ε 3700) (NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 350 mg/kg. US5770000 U.K. Pat., 1975, 1 452 728; CA, **86**, 104446s Nakamura, T. et al., *J. Antibiot.*, 1975, **28**, 477

(4-Hydroxybenzyl)thiocarbamic acid H-427

[(4-Hydroxyphenyl)methyl]carbamothioic acid. (4-Hydroxybenzyl)aminothioformic acid



C₈H₉NO₂S 183.231

Known as naturally occurring O-alkyl esters, the E- and Z-isomers of which are separately isolable.

4'-O-β-L-Rhamnopyranoside, O-α-D-glucopyranosyl ester:

C₂₀H₂₉NO₁₁S 491.515

Constit. of the fruit of *Moringa oleifera* (horseradish tree). Amorph. powder.

4'-O-α-L-Rhamnopyranoside, O-Me ester (E-): **Niazinin A**

[148152-07-2]

C₁₅H₂₁NO₆S 343.4

Constit. of the horseradish tree (*Moringa oleifera*, Moringaceae). Hypotensive agent. λ_{max} 200 ; 224 ; 245 (MeOH).

4'-O-α-L-Rhamnopyranoside, O-Me ester (Z-): **Niazinin B**

[147821-47-4]

C₁₅H₂₁NO₆S 343.4

Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). Hypotensive agent. λ_{max} 200 ; 224 ; 245 (MeOH).

4'-O-α-L-Rhamnopyranoside, O-Et ester (E-): **Niazimicin A**

[147821-49-6]

C₁₆H₂₃NO₆S 357.427

Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). Hypotensive agent. Antitumour promoter. Chemopreventive agent. [α]_D²⁵ -96.8 (c, 0.5 in CHCl₃). λ_{max} 200 ; 223 ; 246 (MeOH).

4'-O-(4-O-Acetyl-α-L-rhamnopyranoside), O-Me ester (E-): **Niazicin A** [159768-74-8]

C₁₇H₂₃NO₇S 385.437

Constit. of *Moringa oleifera* (horseradish tree). λ_{max} 201 ; 223 ; 245 (MeOH).

4'-O-(4-O-Acetyl-α-L-rhamnopyranoside), O-Me ester (Z-): **Niazicin B** C₁₇H₂₃NO₇S 385.437

Constit. of *Moringa oleifera* (horseradish tree). λ_{max} 201 ; 223 ; 245 (MeOH).

4'-O-(4-O-Acetyl-α-L-rhamnopyranoside), O-Et ester (E-): **Niaziminin A** [147821-51-0]

C₁₈H₂₅NO₇S 399.464


Constit. of *Moringa oleifera* (horseradish tree) (Moringaceae). Hypotensive agent. Cryst. (CHCl₃/MeOH). Mp 168-170°. λ_{max} 195 ; 201 ; 222 ; 246 (MeOH).

4'-O-(4-O-Acetyl-α-L-rhamnopyranoside), O-Et ester (Z-): **Niaziminin B**

[147921-21-9]

C₁₈H₂₅NO₃S 399.464Constit. of *Moringa oleifera* (horse-radish tree) (Moringaceae). Hypotensive agent.4'-O-(Tri-O-acetyl- α -L-rhamnopyranoside), O-Me ester (E-): [147821-48-5]C₂₁H₂₇NO₆S 469.512Constit. of *Moringa oleifera* (horse-radish tree) (Moringaceae). λ_{\max} 201 ; 222 ; 247 (MeOH).4'-O-(Tri-O-acetyl- α -L-rhamnopyranoside), O-Et ester (Z-):C₂₁H₂₇NO₆S 469.512Constit. of *Moringa oleifera* (horse-radish tree) (Moringaceae). λ_{\max} 200 ; 221 ; 248 (MeOH).4'-O-(Tri-O-acetyl- α -L-rhamnopyranoside), O-Et ester: [147821-50-9]C₂₂H₂₉NO₆S 483.538Constit. of *Moringa oleifera* (horse-radish tree) (Moringaceae). λ_{\max} 203 ; 221 ; 246 (MeOH).O- α -L-Rhamnopyranoside, Et ester (Z-): **Niazimicin B**C₁₆H₂₃NO₆S 357.427Constit. of *Moringa oleifera* (horse-radish tree).Faizi, S. et al., *J.C.S. Perkin 1*, 1992, 3237-3241; 1994, 3035-3040 (Niazinins, Niaziminins, Niazimicin, Niazicins)Faizi, S. et al., *Phytochemistry*, 1995, **38**, 957-963 (isol, uv, pmr, cmr)Leuck, M. et al., *Carbohydr. Res.*, 1998, **312**, 33-44 (synth, bibl)Murakami, A. et al., *Planta Med.*, 1998, **64**, 319-323 (isol, activity)Guevara, A.P. et al., *Mutat. Res.*, 1999, **440**, 181-188 (activity)Francis, J.A. et al., *Helv. Chim. Acta*, 2004, **87**, 317-326 (4'-rhamnoside glucosyl ester)**2-(α -Hydroxybenzyl)-3,6-tropanediol H-428**

2-(Hydroxyphenylmethyl)-8-methyl-8-azabicyclo[3.2.1]octane-3,6-diol, 9CI

C₁₅H₂₁NO₃ 263.336

Stereochem. of alkaloids not fully known; they may not all correspond stereochemically.

O³-Benzoyl: **Knightolamine**

[75638-70-9]

C₂₂H₂₅NO₄ 367.444Alkaloid from the leaves of *Knightia strobilina* (Proteaceae). Cryst. (Me₂CO). Mp 231-232°. [α]_D²⁰ +66 (CHCl₃).O⁶-Ac: **Knightalbinol**


[75638-71-0]

C₁₇H₂₃NO₄ 305.373Alkaloid from the leaves of *Knightia strobilina* (Proteaceae). Amorph. [α]_D²⁰ +54 (CHCl₃).O⁶-Benzoyl: **Knightia Alkaloid F**

[55249-52-0]

C₂₂H₂₅NO₄ 367.444Minor alkaloid from *Knightia deplanchei* (Proteaceae). Cryst. (Me₂CO). Mp 164-167°. [α]_D 0 (CHCl₃). Tentative plane struct. only detd.; appears to be racemic.O⁶-Benzoyl, O³-cinnamoyl: **Knightia Alkaloid E**

[55249-51-9]

C₃₁H₃₁NO₅ 497.59Minor alkaloid from *Knightia deplanchei* (Proteaceae). Viscous oil. [α]_D 0 (CHCl₃). Tentative plane struct. only detd.; prob. racemic.Lounasmaa, M. et al., *J.O.C.*, 1975, **40**, 3694-3697 (cmr, struct, config)Lounasmaa, M. et al., *Planta Med.*, 1975, **27**, 83-88 (O⁶-benzoyl, O³-benzoyl O³-cinnamoyl)Lounasmaa, M. et al., *Phytochemistry*, 1980, **19**, 953-955 (ir, pmr, ms, struct, Knightolamine, Knightalbinol)**2-(α -Hydroxybenzyl)-3-tropanol H-429**3-Hydroxy-8-methyl- α -phenyl-8-azabicyclo[3.2.1]octane-2-methanol, 9CIC₁₅H₂₁NO₂ 247.336**(1R,1'R,2S,3R)-form**O³-Ac: **Knightinol**


[77053-06-6]

C₁₇H₂₃NO₃ 289.374Alkaloid from *Knightia strobilina* (Proteaceae). Mp 153-154°. [α]_D²⁰ +24 (CHCl₃). Has 2R-config. (change of Cahn-Ingold-Prelog priorities).**(1R,1'R,2S,3R)-form**Di-Ac: **Acetylknightinol**

[77053-07-7]

C₁₉H₂₅NO₄ 331.411Alkaloid from *Knightia strobilina* (Proteaceae). Mp 118-120°. [α]_D²⁰ -18 (CHCl₃).O³-Benzoyl: **Darlingia Alkaloid DF**

[73045-48-4]

C₂₂H₂₅NO₃ 351.444Minor alkaloid from leaves, stems and root of *Darlingia ferruginea* (Proteaceae). Needles (petrol). Mp 136-138°. [α]_D¹⁹ +28 (CHCl₃). Has 2R-config. (change of Cahn-Ingold-Prelog priorities).Bick, I.R.C. et al., *Aust. J. Chem.*, 1979, **32**, 2537-2543 (Alkaloid DF)Lounasmaa, M. et al., *Phytochemistry*, 1980, **19**, 949-952 (Knightinol, Acetylknightinol)Lounasmaa, M. et al., *Planta Med.*, 1983, **48**, 56-58 (synth)Majewski, M. et al., *J.O.C.*, 1995, **60**, 5825-5830 (Knightinol, synth, pmr, cmr, abs config)**1-Hydroxy-3,6-bis[2-(methylthio)ethyl]-2(1H)-pyrazinone H-430**
[37167-10-5]C₁₀H₁₆N₂O₂S₂ 260.381Metab. from *Aspergillus flavus*. Mp 106-107°.MacDonald, J.C. et al., *Can. J. Biochem.*, 1972, **50**, 543 (isol, pmr, ms, struct)**4-Hydroxy-2-butenic acid, H-431**
9CI

4-Hydroxycrotonic acid, 8CI

[4013-24-5]

HOCH₂CH=CHCOOHC₄H₆O₃ 102.09**(E)-form** [24587-49-3]

Cryst. (EtOAc). Mp 109°. Readily undergoes polycondensation.

Et ester: [10080-68-9]

C₆H₁₀O₃ 130.143d₄¹⁷ 1.08. Bp₁₃ 119-120°. n_D²³ 1.4610.

Ac:

C₆H₈O₄ 144.127Cryst. (Et₂O/petrol).**(Z)-form**Nitrile, O- β -D-glucopyranoside: **Alliarioside**C₁₀H₁₅NO₆ 245.232Constit. of *Alliaria petiolata*. Feeding deterrent for larvae of *Pieris napi oleracea*. Prisms (as tetra-Ac). Mp 84-86° (tetra-Ac). [α]_D -4 (c, 0.4 in CHCl₃) (tetra-Ac). λ_{\max} 218 (MeOH).

[29576-13-4, 82744-23-8, 4508-99-0]

Laporte, J.F. et al., *Bull. Soc. Chim. Fr.*, 1969, 1340 (synth, ir, pmr)Boulanger, T. et al., *J. Crystallogr. Spectrosc. Res.*, 1987, **17**, 71 (cryst struct)Dardoize, F. et al., *Tetrahedron*, 1989, **45**, 7783 (synth, pmr)Haribal, M. et al., *J. Nat. Prod.*, 2001, **64**, 440-443 (Alliarioside)**(4-Hydroxybutyl)guanidine, H-432**
9CI

4-Guanidino-1-butanol

[17581-95-2]

HN=C(NH₂)NH(CH₂)₃CH₂OHC₅H₁₃N₃O 131.177Alkaloid from *Leonurus sibiricus*.

Monohydrochloride: [27091-78-7]

Cryst. (2-propanol). Mp 107-109°.

Sulfate (2:1): [98600-31-8]

Cryst. (EtOH). Mp 168-169°.

O-(4-Hydroxy-3,5-dimethoxybenzoyl):

Leonurine. 4-Guanidinobutyl syringate [24697-74-3]C₁₄H₂₁N₃O₅ 311.337Alkaloid from *Leonurus artemisia* (a Chinese herbal medicine used in ob-

stetrics) and *Leonurus sibiricus* (Lamiaceae). Uterotonic agent. Mp 229-230°. p*K*_a 7.9 (H₂O).

O-(4-Hydroxy-3,5-dimethoxybenzoyl), hydrochloride: Mp 195-196°.

[106936-57-6]

Fishbein, L. et al., *J.A.C.S.*, 1954, **76**, 3217-3219 (synth)

Goto, T. et al., *Tet. Lett.*, 1962, **3**, 545-548 (Leonurine, isol, struct)

Sugiyama, S. et al., *Tetrahedron*, 1969, **25**, 5155-5161 (Leonurine, synth, struct)


Ludwig, B.J. et al., *J. Med. Chem.*, 1970, **13**, 60-63 (synth)

Reuter, G. et al., *Pharmazie*, 1971, **26**, 777 (isol)

Yeung, H.W. et al., *Planta Med.*, 1977, **31**, 51 (Leonurine, isol, ir, pmr, ms, struct)

Cheng, K.F. et al., *Experientia*, 1979, **35**, 571-572 (Leonurine, synth)

2-(2-Hydroxybutyl)-6-(2-hydroxypropyl)-1-methylpiperidine H-433 8-Ethyl-10-propyllobelidol



(2*R**,2'*R**,2''*R**,6*S*)-form

C₁₅H₃₁NO₂ 257.415

(2*R**,2'*R**,2''*R**,6*S*)-form [212006-68-3]
Alkaloid from *Siphocampylus verticillatus*. Amorph. Mp 55° (as hydrochloride).

(2*R**,2'*S**,2''*R**,6*S*)-form [212515-05-4]
Alkaloid from *Dialypetalum floribundum*.

(2*S*,2'*R*,2''*R*,6*S*)-form

3,4-Didehydro: α²-Ethyl-1,2,3,6-tetrahydro-1-methyl-α²-propyl-2,6-pyridinediethanol, 9CI. 1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxypropyl)-1-methylpyridine. *Andrachcine* [109028-29-7]

C₁₅H₂₉NO₂ 255.4

Alkaloid from *Andrachne aspera* (Euphorbiaceae). Gum. [α]_D²⁵ -87 (c, 1.2 in MeOH). Has 2*R*-config. owing to introduction of double bond.

3,4-Didehydro, 2'-ketone: *Andrachcinine* [274925-35-8]

C₁₅H₂₇NO₂ 253.384

Alkaloid from *Andrachne aspera*. Oil. [α]_D²² -108 (c, 2.5 in CHCl₃).

(2*S*,2'*S*,2''*S*,6*S*)-form

N-De-Me: 2-(2-Hydroxybutyl)-6-(2-hydroxypropyl)piperidine. *Aspertine D* [442155-63-7]

C₁₄H₂₉NO₂ 243.389

Alkaloid from the aerial parts of *Andrachne aspera*. Gum. [α]_D -97 (c, 0.8 in CHCl₃).

3,4-Didehydro, N-de-Me: 1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxypropyl)pyridine. *Aspertine A* [442155-60-4]

C₁₄H₂₇NO₂ 241.373

Alkaloid from the aerial parts of *Andrachne aspera*. Gum. [α]_D -103 (c, 0.5 in CHCl₃).

Biaivatti, M.W. et al., *Phytochemistry*, 1998, **48**, 747-749 (isol, ir, pmr, ms)


Krebs, H.C. et al., *Phytochemistry*, 1998, **48**, 911-913 (isol, pmr, cmr, ms)

Mill, S. et al., *J. Nat. Prod.*, 2000, **63**, 762-764 (isol, pmr, cmr, abs config)

Ahmad, V.U. et al., *Turk. J. Chem.*, 2002, **26**, 245-250 (*Aspertine A, D*)

Miguel, O.G. et al., *Z. Naturforsch.*, C, 2002, **57**, 81-84 (isol, pmr, cmr)

2-(2-Hydroxybutyl)-6-(2-hydroxypropyl)-1-methylpiperidine H-434 10-Ethyl-8-methyllobelidol. 8-Methyl-10-ethyllobelidol [96777-12-7]



C₁₃H₂₇NO₂ 229.362

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Oil. Opt. inactive. Hydrochloride, hydrobromide, hydroiodide and perchlorate were noncryst.

Hexachloroplatinate: Mp 217-218° dec.

Picrate: [110555-04-9]

Cryst. (2-propanol). Mp 133-134°.

4,5-Didehydro: α²-Ethyl-1,2,3,6-tetrahydro-α²-1,1-dimethyl-2,6-pyridinediethanol, 8CI. 6-(2-Hydroxybutyl)-2-(2-hydroxypropyl)-1-methyl-Δ³-piperidine. *Dehydro-8-methyl-10-ethyllobelidol*. Alkaloid D3 [105694-51-7]

C₁₃H₂₅NO₂ 227.346


Alkaloid from *Lobelia syphilitica* and *Lobelia berlandieri* (Campanulaceae). Extremely hygroscopic cryst. (as hydrochloride). Mp 120° (sealed tube)(hydrochloride). [α]_D²⁵ -110 (c, 1.0 in EtOH). Props. refer to the *L. syphilitica* isolate, a 2,6-*trans*-isomer which gives on hydrogenation 10-ethyl-8-methyllobelidol identical to the (opt. inactive) natural alkaloid. The *L. berlandieri* isolate may not necessarily be identical.

Schöpf, C. et al., *Annalen*, 1957, **608**, 88-127 (isol, struct)

Tschesche, R. et al., *Chem. Ber.*, 1961, **94**, 3327-3336 (didehydro)

Williams, H.J. et al., *J. Agric. Food Chem.*, 1987, **35**, 19-22 (didehydro)

2-(2-Hydroxybutyl)piperidine H-435 α-Ethyl-2-piperidineethanol, 9CI. 1-(2-Piperidinyl)-2-butanol. 8-Ethylnorlobelol I



(2*R*,2'*R*)-form

C₉H₁₉NO 157.255

(2*R*,2'*R*)-form

Synthetic. Mp 86-87°. [α]_D²² -26 (c, 2.1 in EtOH).

(2*S*,2'*R*)-form

Alkaloid from *Andrachne aspera*. [α]_D²² -15.7 (c, 2.1 in MeOH).

(2*S*,2'*S*)-form

Alkaloid from *Lobelia inflata* (Campanulaceae). Cryst. (Et₂O). Mp 87°. [α]_D^{21.5} +22.3 (c, 1.56 in EtOH). Readily subl.

Hydrochloride: Mp 135°.

Picrate: Mp 116-117°.

(2ξ,2'ξ)-form

Alkaloid from leaves of *Sedum brissemoretti* and *Sedum fusiforme*. Identified by gc-ms.

N-Me: 1-(N-Methyl-2-piperidinyl)-2-butanol. 2-(2-Hydroxybutyl)-N-methylpiperidine

C₁₀H₂₁NO 171.282

Alkaloid from leaves of *Sedum brissemoretti*, *Sedum nudum*, *Sedum lancertottense* and *Sedum fusiforme*. Identified by gc-ms.

8-Ketone: 1-(2-Piperidinyl)-2-butanone.

2-(2-Oxobutyl)piperidine

C₉H₁₇NO 155.239

Alkaloid from leaves of *Sedum brissemoretti*, *Sedum nudum*, *Sedum lancertottense* and *Sedum fusiforme*. Identified by gc-ms.

8-Ketone, N-Me: 1-(N-Methyl-2-piperidinyl)-2-butanone. N-Methyl-2-(2-oxobutyl)piperidine

C₁₀H₁₉NO 169.266

Alkaloid from leaves of *Sedum brissemoretti*, *Sedum nudum*, *Sedum lancertottense* and *Sedum fusiforme*. Identified by gc-ms.

(2*RS*,2'*RS*)-form

Synthetic. Cryst. (hexane). Mp 65-66°.

8-Ketone, N-Me: Alkaloid from whole plants of *Pratia nummularia*. Pale yellow oil.

[81738-59-2]

Wieland, H. et al., *Annalen*, 1931, **491**, 14 (isol)
Schöpf, C. et al., *Annalen*, 1957, **608**, 88 (isol, struct)

Stevens, J.F. et al., *Phytochemistry*, 1992, **31**, 3917-3924 (derivs)


Ho, L.K. et al., *Planta Med.*, 1995, **61**, 567 (deriv)

Mill, S. et al., *Annalen*, 1996, 2083-2086 (synth, abs config)

Mill, S. et al., *J. Nat. Prod.*, 2000, **63**, 762-764 (isol, synth)

Chen, L.-J. et al., *Tetrahedron: Asymmetry*, 2008, **19**, 715-720 (synth)

5-(3-Hydroxybutyl)-2-pyridinecarboxylic acid, 9CI H-436 5-(3-Hydroxybutyl)picolinic acid. *Fusarinoic acid* [26108-30-5]



(*S*)-form

C₁₀H₁₃NO₃ 195.218

(*S*)-form [33130-74-4]

Isol. from *Gibberella fujikuroi*. Cryst.

(CH₂Cl₂/MeOH)(synthetic); oil (nat.). Mp 108–109°. $[\alpha]_D^{25} +20.5$ (c, 1.05 in MeOH). The oily natural product appears to have been largely racemic.

Me ester:

Oil. $[\alpha]_D^{25} +17.9$ (c, 1.13 in CHCl₃).

(±)-**form** [33130-77-7]

Mp 122–123°.

Steiner, K. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 845 (synth)

Frederiks, J.C. *et al.*, *Verh. Schweiz. Naturforsch. Ges.*, 1971, **151**, 84


Buyuk, G. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 682 (synth)

Renslo, A.R. *et al.*, *J.O.C.*, 1998, **63**, 7840-7850 (*S*-form, synth)

Song, J.J. *et al.*, *J.O.C.*, 2001, **66**, 605-608 (synth)

2-(4-Hydroxybutyl)-4(3H)-quinazolinone

2-(4-Hydroxybutyl)-4-quinazolinol
[60915-16-4]



C₁₂H₁₄N₂O₂ 218.255


Alkaloid from the leaves of *Dichroa febrifuga*. Cryst. (EtOAc). Mp 175–177°.

Deng, Y. *et al.*, *J. Chin. Pharm. Sci.*, 2000, **9**, 116-118 (isol)

Mhaske, S.B. *et al.*, *Tetrahedron*, 2004, **60**, 3417-3420 (synth, pmr, cmr)

1-Hydroxycanthin-6-one

1-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI
[80787-59-3]



C₁₄H₈N₂O₂ 236.229

Alkaloid from the heartwood of *Ailanthus giraldii* and root bark of *Ailanthus altissima* (Simaroubaceae). Yellow cryst. (MeOH/Me₂CO). Mp 220°.

1-O-[β-D-Xylopyranosyl-(1→6)-β-D-glucopyranoside]: **Ailantcanthinoside B**
[960002-01-1]

C₂₅H₂₆N₂O₁₁ 530.487

Alkaloid from the root bark of *Ailanthus altissima*. Powder. Mp 186–188°. λ_{max} 225 ; 358 ; 376 (Py).

Ac:

C₁₆H₁₀N₂O₃ 278.267

Cryst. (MeOH). Mp 205°.

Me ether: 1-Methoxycanthin-6-one. 1-Methoxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI
[60755-86-4]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from the wood and root bark

of *Ailanthus altissima*, and from the root bark of *Ailanthus excelsa* (Simaroubaceae). Cytotoxic to guinea pig keratocytes. Shows significant anti-HIV activity. Pale yellow needles (Me₂CO or MeOH). Sol. MeOH, Me₂CO; poorly sol. H₂O. Mp 250–250.5° (246°). λ_{max} 259 ; 270 ; 280 ; 346 ; 360 ; 383 (MeOH) (Berdy). λ_{max} 279 (ε 12300); 330 (ε 7500); 361 (ε 17400); 370 (ε 16300); 378 (ε 19000) (EtOH) (Berdy).

Me ether, N-oxide: 1-Methoxycanthin-6-one N(3)-oxide

[77370-00-4]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the root bark of *Ailanthus altissima* (Simaroubaceae).

Yellow needles (CHCl₃). Mp 256–257°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1532 (isol, uv, ir, pmr, ms, struct)

Cordell, G.A. *et al.*, *J. Nat. Prod.*, 1978, **41**, 166 (isol)

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 390 (isol, uv, ir, pmr, ms, derivs)

Varga, E. *et al.*, *Fitoterapia*, 1981, **52**, 183 (isol)

Khan, S.A. *et al.*, *Phytochemistry*, 1981, **20**,

2062 (isol, uv, ir, pmr, ms, struct)

Hagen, T.J. *et al.*, *J.O.C.*, 1989, **54**, 2170 (synth, pmr, 1-Me ether)

Xu, Z. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1712-1715 (activity)

Suzuki, H. *et al.*, *Synthesis*, 2005, 28-31 (1-Me ether, synth)

Zhang, L.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2007, **9**, 253-259 (*Ailantcanthinoside B*)

2-Hydroxycanthin-6-one

2-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI

C₁₄H₈N₂O₂ 236.229

Alkaloid from cell cultures of *Ailanthus altissima* (Simaroubaceae). Amorph. solid.

Me ether: 2-Methoxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI. 2-Methoxycanthin-6-one
[116353-93-6]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from stem wood of *Quassia amara* (Surinam quassia) (Simaroubaceae). Pale yellow powder (MeOH/CHCl₃). Mp 262–264°.

Crespi-Perellino, N. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1010 (isol, uv, pmr, ms, struct)

Njar, V.C.O. *et al.*, *Planta Med.*, 1993, **59**, 259 (2-Methoxycanthin-6-one)

4-Hydroxycanthin-6-one

4-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI

C₁₄H₈N₂O₂ 236.229

Alkaloid from cell cultures of *Ailanthus altissima* (Simaroubaceae). Yellow amorph. solid (MeOH).

Me ether: 4-Methoxycanthin-6-one. 4-Methoxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI. Cordatanine

[5023-08-5]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from the stem and root bark of *Charpentiera obovata* and from *Drymaria cordata* (Aramanthaceae, Caryophyllaceae). Needles (C₆H₆). Mp 220–221°.

Me ether, picrate:

Yellow needles (EtOH). Mp 221.5–222.5° dec.

Me ether, methiodide:

Cryst. +1 H₂O (H₂O + trace HI). Mp 218° dec.


Nelson, E.R. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 768 (synth)

Scheuer, P.J. *et al.*, *J. Nat. Prod.*, 1965, **28**, 95 (isol, uv, synth)

Chen, W. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1986, **28**, 450; *CA*, **106**, 15732w (deriv)

5-Hydroxycanthin-4-one

5-Hydroxy-4H-indolo[3,2-de][1,5]-naphthyridin-4-one, 9CI



C₁₄H₈N₂O₂ 236.229

Me ether: 5-Methoxycanthin-4-one. **Drymaritine**

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from *Drymaria diandra*. Anti-HIV agent. Pale yellow solid. Mp 181–183°. λ_{max} 222 (log ε 3.75); 237 (sh) (log ε 3.64); 264 (log ε 3.6); 288 (log ε 3.41); 297 (log ε 3.41); 333 (sh) (log ε 3.18); 349 (log ε 3.43); 366 (log ε 3.41) (MeOH).

Hsieh, P.-W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1175-1177 (isol, uv, pmr, cmr, ms)

5-Hydroxycanthin-6-one

5-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI
[64118-73-6]

C₁₄H₈N₂O₂ 236.229

Alkaloid from the root bark of *Simarouba amara* and from *Ailanthus altissima* (Simaroubaceae). Orange-yellow platelets (MeOH). Sol. MeOH, CHCl₃. Mp 259–261° dec.

O-[β-D-Xylopyranosyl-(1→6)-β-D-glucopyranoside]: **Ailantcanthinoside A**
[960002-00-0]

C₂₅H₂₆N₂O₁₁ 530.487

Alkaloid from the root bark of *Ailanthus altissima*. Powder (MeOH). Mp 200–202°. λ_{max} 225 ; 304 ; 343 ; 358 ; 376 (Py).

O-[β-D-Glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Bruceolline B**. **Bruceacanthinoside**
[159194-91-9]

C₂₆H₂₈N₂O₁₂ 560.513

Alkaloid from stems of *Brucea javanica* and root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Pale yellow amorph. powder. Mp 192–194°. $[\alpha]_D^{25} -41$ (c, 1 in Py). $[\alpha]_D^{25} -64.4$ (c, 0.45 in DMSO).

Ac:

Needles (C₆H₆). Mp 231.5–232°.

Me ether: 5-Methoxycanthin-6-one
[15071-56-4]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from the bark of *Zanthoxylum caribaeum*, *Zanthoxylum elephantiasis*, *Pentaceras australis* and *Odyendea gabonensis*, the wood of *Picrasma excelsa* (Jamaican quassia wood), and the root bark of *Ailanthus excelsa* (Rutaceae, Simaroubaceae). Antineoplastic agent. Cytotoxic to guinea pig keratocytes. Pale yellow needles (MeOH, CHCl₃/MeOH or EtOH/CH₂Cl₂). Mp 243-244° (237-238°) dec. Log P 2.35 (uncertain value) (calc).

Me ether, hydrochloride:

Needles (MeOH). Mp 205-208°.

Me ether, picrate:

Yellow needles (MeOH). Mp 244-246°.

Haynes, H.F. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 387 (isol)

Nelson, E.R. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 563 (uv, struct, synth)

Bartlett, M.F. *et al.*, *J.A.C.S.*, 1960, **82**, 5941 (synth)

Awad, A.T. *et al.*, *J. Pharm. Sci.*, 1967, **56**, 279 (isol)

Della Casa, D. *et al.*, *J.C.S. (C)*, 1967, 2155 (isol, uv, ir, pmr)

Lassak, E.V. *et al.*, *Phytochemistry*, 1977, **16**, 1126 (isol, synth, uv, ms, struct)

Cordell, G.A. *et al.*, *J. Nat. Prod.*, 1978, **41**, 166 (isol, uv, ir, pmr, ms)

Wagner, H. *et al.*, *Planta Med.*, 1979, **36**, 113 (isol)

Forgacs, P. *et al.*, *Planta Med.*, 1982, **46**, 187 (isol)

Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1416 (*Bruceacanthinoid*)

Ouyang, Y. *et al.*, *Phytochemistry*, 1994, **36**, 1543 (*Bruceolline B*)

Zhang, L.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2007, **9**, 253-259 (*Ailantcanthinoside A*)

8-Hydroxycanthin-6-one H-443

8-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI
[66762-19-4]

C₁₄H₈N₂O₂ 236.229

Alkaloid from the root bark of *Ailanthus excelsa* and the trunk bark of *Odyendea gabonensis* (Simaroubaceae). Cryst. (CHCl₃). Mp 340° (270-275° dec.).

Cordell, G.A. *et al.*, *J. Nat. Prod.*, 1978, **41**, 166 (isol, uv, ir, pmr, ms, struct)

Forgacs, P. *et al.*, *Planta Med.*, 1982, **46**, 187 (isol)

9-Hydroxycanthin-6-one H-444

9-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI
[138544-91-9]

C₁₄H₈N₂O₂ 236.229

Alkaloid from the wood of *Simaba multiflora*, from the roots of *Eurycoma longifolia* and from stems of *Picrolemma granatensis* (Simaroubaceae). Yellow needles. Mp 288-293° (285-286°). The nat. prod. from *S. multiflora* has been reassigned this struct. by Kardono, *et al.* λ_{max} 238 (sh) (ε 15850); 274 (ε 15850); 310 (ε 10960); 352 (ε 14130) (MeOH).

N³-Oxide: 9-Hydroxycanthin-6-one N³-oxide

[137739-75-4]

C₁₄H₈N₂O₃ 252.229

Alkaloid from the roots of *Eurycoma*

longifolia (Simaroubaceae). Red needles. Mp 248-250° dec. λ_{max} 230 (ε 33110); 289 (ε 3200); 360 (ε 19500); 417 (ε 1200) (MeOH).

O-β-D-Glucopyranoside: [339077-98-4]

C₂₀H₁₈N₂O₇ 398.371

Alkaloid from the roots of *Eurycoma harmandiana*. Amorph. yellow powder. λ_{max} 285 (log ε 3.16); 321 (log ε 3.14); 367 (log ε 3.22) (MeOH/HCl).

Me ether: 9-Methoxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI. 9-Methoxycanthin-6-one. 8-Methoxycanthin-6-one (incorr.)

[74991-91-6]

C₁₅H₁₀N₂O₂ 250.256

Yellow cryst. (EtOH). Mp 178-180°. λ_{max} 272 (ε 14750); 310 (ε 4500); 353 (ε 7000) (MeOH). λ_{max} 278 (ε 25000); 320 (ε 12000); 380 (ε 18700) (MeOH/HCl) (Berdy).

Me ether, N³-oxide: 9-Methoxycanthin-6-one N³-oxide

[137739-74-3]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the roots of *Eurycoma longifolia* and stems of *Picrolemma granatensis* (Simaroubaceae). Orange-yellow powder. Mp 238-240° (193° dec.). λ_{max} 242 (ε 1000); 286 (ε 15500); 293 (ε 18600); 360 (ε 6300); 422 (ε 5600) (CHCl₃).

1,2-Dihydro: 1,2-Dihydro-9-hydroxy-6-canthinone. 4,5-Dehydrofluorodaturatin. 1,2-Dehydrofluorodaturatin (incorr.). 1,2-Dehydrofluorodaturine (incorr.)

[95855-89-3]

C₁₄H₁₀N₂O₂ 238.245

Alkaloid from the seeds of *Datura stramonium* var. *stramonium* (Solana-ceae). Brown cryst. Mp 360° (blackens >73°). Also obt. by air oxidn. of Fluorodaturatin, therefore prob. artifact. Incorr. called 1,2-Dehydrofluorodaturine in *CA*. The struct. was assigned (and the name 1,2-Dehydrofluorodaturatin allocated) before the correct struct. of Fluorodaturatin was assigned.

1,2,4,5-Tetrahydro: 1,2,4,5-Tetrahydro-9-hydroxy-6-canthinone. Fluorodaturatin.

Alkaloid GF

[80757-43-3]

C₁₄H₁₂N₂O₂ 240.261

Alkaloid from the seeds of *Datura stramonium* var. *stramonium* (Solana-ceae). Yellow cryst. Dec. >125°, turns black above 140°; does not melt < 360°. Largely enolised. Struct. revised in 1988. λ_{max} 268 (ε 2800); 368 (ε 8600); 455 (ε 5200) (MeOH/NaOH) (Derep). λ_{max} 217 (ε 22200); 261 (ε 6780); 387 (ε 20700) (MeOH) (Derep).

Giesbrecht, A.M. *et al.*, *Phytochemistry*, 1980, **19**, 313-315 (*Me ether*, isol, uv, ir, pmr, ms, struct)

Maier, I. *et al.*, *Monatsh. Chem.*, 1981, **112**, 1425-1439 (*Fluorodaturatin*)

Polonsky, J. *et al.*, *Tet. Lett.*, 1982, **23**, 869-872 (isol, uv, ir, pmr, struct)

Jurenitsch, J. *et al.*, *Sci. Pharm.*, 1984, **52**, 301-306 (*Dehydrofluorodaturin*)

Robien, W. *et al.*, *Sci. Pharm.*, 1988, **56**, 133 (*Fluorodaturatin*, struct)

Kardono, L.B.S. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1360-1367 (*N-oxide*, *Me ether*, *Me ether N-oxide*, *isol*, *uv*, *ir*, *ms*, *pmr*, *cmr*)

Rodrigues Fo, E. *et al.*, *Phytochemistry*, 1992, **31**, 2499-2501 (*Me ether N-oxide*)

Kanchanapoom, T. *et al.*, *Phytochemistry*, 2001, **56**, 383-386 (*glucoside*)

10-Hydroxycanthin-6-one H-445

10-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI. *Aervine*. *Ervine*†
[86293-41-6]

C₁₄H₈N₂O₂ 236.229

Alkaloid from aerial parts of *Aerva lanata* (Amaranthaceae) and from *Simaba multiflora*. Yellow cryst. (MeOH or CHCl₃/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 310-313° dec. λ_{max} 238 (sh) (ε 15800); 283 (sh) (ε 13800); 297 (ε 15500); 320 (sh) (ε 12900); 356 (ε 11000); 428 (ε 8510) (MeOH/NaOH) (Derep). λ_{max} 238 (sh) (ε 15500); 269 (sh) (ε 16600); 275 (ε 17800); 304 (sh) (ε 11000); 312 (ε 11700); 352 (ε 14800) (MeOH) (Derep).

O-β-D-Glucopyranoside: *Aervoside*. *Ervo-side*

[139767-01-4]

C₂₀H₁₈N₂O₇ 398.371

From aerial parts of *Aerva lanata* (Amaranthaceae). Bright yellow cryst. (MeOH or EtOH). Mp 215-218°.

Me ether: 10-Methoxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI. 10-Methoxycanthin-6-one. *Methylaervin*
[86293-40-5]

C₁₅H₁₀N₂O₂ 250.256

From aerial parts of *Aerva lanata* (Amaranthaceae) Alkaloid from aerial parts of *Aerva lanata* (Amaranthaceae) and from *Simaba multiflora*. Bright yellow needles (MeOH or CHCl₃/EtOH). Mp 194-196°. λ_{max} 212 (ε 15800); 232 (sh) (ε 6460); 266 (sh) (ε 8910); 274 (ε 11500); 310 (ε 4900); 352 (ε 6760) (MeOH) (Derep).

Zapesochayna, G. *et al.*, *Planta Med.*, 1992, **58**, 192 (isol, uv, ir, pmr, cmr, ms, struct)

11-Hydroxycanthin-6-one H-446

11-Hydroxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI. *Amarorine*
[75969-83-4]

C₁₄H₈N₂O₂ 236.229

Alkaloid from the bark of *Amaroria soulameoides* and from *Soulamea soulameoides*, *Brucea antidysenterica* and *Quassia kerstingii* (Simaroubaceae). Red-brown cryst. + 1H₂O (MeOH aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 323-325°. λ_{max} 326 (ε 33200); 383 (ε 40900) (MeOH) (Berdy). λ_{max} 352; 445 (MeOH-NaOH) (Berdy). λ_{max} 325 (ε 9100); 385 (ε 17800) (EtOH) (Berdy).

N-Oxide: 11-Hydroxycanthin-6-one N-oxide

[158784-71-5]

C₁₄H₈N₂O₃ 252.229

Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Yellow needles. Mp 300°. Intense yellow-green fluor. in MeOH.

O-β-D-Glucopyranoside: *Bruceolline G*.

11-O- β -D-Glucopyranosyloxycanthin-6-one

[165967-63-5]

C₂₀H₁₈N₂O₇ 398.371

Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Amorph. pale yellow powder.

O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Bruceolline A**

[159194-90-8]

C₂₆H₂₈N₂O₁₂ 560.513

Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Pale yellow amorph. powder (MeOH). Mp 241-243° dec. $[\alpha]_D^{25}$ -62.2 (c, 1 in Py).

Me ether: **11-Methoxycanthin-6-one**. 11-Methoxy-6H-indolo[3,2,1-de][1,5]-naphthyridin-6-one, 9CI. *Amaroridine* [75969-82-3]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from bark of *Amaroria sou-lameoides* (Simaroubaceae). Mp 237-238°.


Clarke, P.J. *et al.*, *J.C.S. Perkin 1*, 1980, 1614 (isol, uv, ir, pmr, ms, cryst struct)

Ouyang, Y. *et al.*, *Phytochemistry*, 1994, 36, 1543 (oxide, Bruceolline A)

Ouyang, Y. *et al.*, *Phytochemistry*, 1995, 39, 911 (Bruceolline G)

1-Hydroxy-9H-carbazole-3-carboxaldehyde H-447

3-Formyl-1-hydroxycarbazole. **Demethyl-murrayanine** [123497-84-7]



C₁₃H₉NO₂ 211.22

Alkaloid from *Clausena heptaphylla* (Rutaceae). Plates (CHCl₃). Mp 237-239°. λ_{\max} 226 (log ϵ 4.4); 244 (log ϵ 4.51); 255 (log ϵ 4.39); 278 (log ϵ 4.59); 291 (log ϵ 4.45); 336 (log ϵ 4.22); 346 (log ϵ 4.22) (EtOH).

Me ether: 1-Methoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-1-methoxycarbazole. **Murrayanine**† [723-97-7]

C₁₄H₁₁NO₂ 225.246

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) and the roots of *Clausena heptaphylla* (Rutaceae). Active against gram-positive bacteria. Needles (C₆H₆/petrol). Sol. MeOH, C₆H₆; poorly sol. hexane. Mp 168-170° Mp 219° dec. (synthetic). Bp_{0.05} 110-115° subl. λ_{\max} 239 (ϵ 29510); 247 (ϵ 19950); 274 (ϵ 36310); 289 (ϵ 36300) (MeOH) (Berdy).

Me ether, picrate:

Fine orange needles (C₆H₆/petrol). Mp 198-200° dec.

Me ether, oxime:

C₁₄H₁₂N₂O₂ 240.261

Needles (C₆H₆/petrol). Mp 155-156°.

Me ether, N-Me:

C₁₃H₁₃NO₂ 239.273

Cream-coloured needles (C₆H₆/petrol). Mp 150.5-151.5°.

Das, K.C. *et al.*, *Experientia*, 1965, 21, 340

(Murrayanine, activity)

Chakraborty, D.P. *et al.*, *Tetrahedron*, 1965, 21, 681-685 (Murrayanine, isol, uv, ir, pmr, struct)

Crum, J.D. *et al.*, *Chem. Comm.*, 1966, 417-419 (Murrayanine, synth, uv, ir, pmr)

Chakraborty, D.P. *et al.*, *J.O.C.*, 1968, 33,

1265-1268 (Murrayanine, synth, uv, ir)

Bhattacharyya, P. *et al.*, *Phytochemistry*, 1973, 12, 1831-1832 (Murrayanine, isol, uv, ir)

Ngadjui, B.T. *et al.*, *Phytochemistry*, 1989, 28,

1517-1519 (isol, uv, ir, pmr, cmr, ms, struct)

Knölker, H.-J. *et al.*, *Tetrahedron*, 1993, 49,

11221-11234 (Murrayanine, synth)

Bringmann, G. *et al.*, *Planta Med.*, 1998, 64,

54-57 (synth, pmr, cmr)

Bringmann, G. *et al.*, *Synthesis*, 1998, 1501-

1505 (synth, Me ether, cmr)

Benavides, A. *et al.*, *Synthesis*, 2004, 2499-2504

(Murrayanine, synth)

Bernal, P. *et al.*, *Helv. Chim. Acta*, 2007, 90,

1449-1454 (Murrayanine, synth)

2-Hydroxy-9H-carbazole-3-carboxaldehyde H-448

3-Formyl-2-hydroxycarbazole. **Mukonal** [203223-67-5]

C₁₃H₉NO₂ 211.22

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae).

Cryst. (C₆H₆/CHCl₃). Mp 238°. λ_{\max} 229 (ϵ 26360); 247 (ϵ 16220); 278 (ϵ 34990); 297 (ϵ 36140); 345 (ϵ 11560) (EtOH).

O-Ac:

Cryst. (C₆H₆). Mp 210°.

Me ether: 2-Methoxy-9H-carbazole-3-carboxaldehyde. 3-Formyl-2-methoxycarbazole. **Glycosinine**. O-Methylmukonal [51971-08-5]

C₁₄H₁₁NO₂ 225.246

Alkaloid from *Murraya siamensis* and roots of *Glycosmia pentaphylla* (Rutaceae). Mp 189-189.5° (185°). λ_{\max} 296 (log ϵ 4.64); 350 (log ϵ 4.19) (EtOH).

Joshi, B.S. *et al.*, *Chem. Ind. (London)*, 1968, 685 (uv, ir, pmr, synth)

Bhagwanth, M.R.R. *et al.*, *Indian J. Chem.*,

1969, 7, 1065-1069 (synth)

Bhattacharyya, P. *et al.*, *Phytochemistry*, 1984,

23, 471-472 (isol, uv, ir, pmr, cmr, ms)

Ruangrunsi, N. *et al.*, *J. Nat. Prod.*, 1990, 53,

946-952 (O-Methylmukonal)

Wu, T.S. *et al.*, *Phytochemistry*, 1993, 32, 449-

451 (Glycosinine)

Knölker, H.J. *et al.*, *Synlett*, 1996, 737-740

(Mukonal, O-Methylmukonal, synth)

6-Hydroxy-9H-carbazole-3-carboxaldehyde H-449

3-Formyl-6-hydroxycarbazole

C₁₃H₉NO₂ 211.22

Me ether: 6-Methoxy-9H-carbazole-3-carboxaldehyde. 8-Methoxy-2-carbazolecarboxaldehyde (obsol.). 3-Formyl-6-methoxycarbazole [107816-72-8]

C₁₄H₁₁NO₂ 225.246

Alkaloid from the roots of *Clausena lansium* (wampee) (Rutaceae). Prisms (Me₂CO/hexane). Mp 135-136°.

Hewlins, M.J.E. *et al.*, *J. Chem. Res., Synop.*, 1986, 292

7-Hydroxy-9H-carbazole-3-carboxaldehyde, 9CI H-450

6-Formyl-2-hydroxycarbazole [142750-13-8]

C₁₃H₉NO₂ 211.22

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Powder (Me₂CO).

Me ether: 7-Methoxy-9H-carbazole-3-carboxaldehyde, 9CI. 6-Formyl-2-methoxycarbazole. **Clausoline K** [187110-71-0]

C₁₄H₁₁NO₂ 225.246

Alkaloid from *Clausena excavata*.

Powder. λ_{\max} 226 (sh); 232; 252 (sh); 293; 324 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1992, 40, 230; 1997, 45, 48-52 (isol, uv, ir, pmr, ms, struct)

Krahl, M.P. *et al.*, *Org. Biomol. Chem.*, 2006, 4, 3215-3219 (synth)

8-Hydroxy-9H-carbazole-3-carboxaldehyde H-451

6-Formyl-1-hydroxycarbazole

C₁₃H₉NO₂ 211.22


Me ether: 8-Methoxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-8-methoxycarbazole. 6-Formyl-1-methoxycarbazole. **Mukolidine** [87264-45-7]

C₁₄H₁₁NO₂ 225.246

Minor alkaloid from roots of *Murraya koenigii* (curryleaf tree). Cryst. (C₆H₆). Mp 152-155°.

Roy, S. *et al.*, *J. Indian Chem. Soc.*, 1982, 59, 1369 (Mukolidine)

1-Hydroxy-9H-carbazole-3-carboxylic acid, 9CI H-452



C₁₃H₉NO₃ 227.219

Me ester: **Clausine E**. **Clausoline I**

[182261-83-2]

C₁₄H₁₁NO₃ 241.246

Alkaloid from stem bark of *Clausena excavata*. Powder (Me₂CO). Mp 218-220°. λ_{\max} 222 (sh) (log ϵ 4.01); 239 (log ϵ 4.14); 248 (sh) (log ϵ 4.08); 269 (sh) (log ϵ 4.22); 276 (log ϵ 4.23); 311 (log ϵ 3.65); 324 (log ϵ 3.64); 337 (log ϵ 3.5) (MeOH).

Me ether: 1-Methoxy-9H-carbazole-3-carboxylic acid. **Mukoic acid**. **Mukeic acid**

[3889-89-2]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (C₆H₆ or EtOAc). Mp 242°. λ_{\max} 210 (log ϵ 4.58); 230 (log ϵ 3.92); 235 (log ϵ 4.51) (no solvent)

reported).

Me ether, picrate: Mp 225-228°.

Me ether, Me ester: Mukonine

[23523-94-6]

C₁₃H₁₃NO₃ 255.273

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (CHCl₃/C₆H₆). Mp 198-200° (195°). λ_{max} 236 (log ε 4.6); 245 (log ε 4.57); 266 (log ε 4.64); 274 (log ε 4.73); 306 (log ε 3.88); 320 (log ε 3.78) (EtOH).

Choudhury, B.K. *et al.*, *Phytochemistry*, 1971, **10**, 1967-1970 (*Mukoic acid, isol, uv, ir, pmr, struct*)

Chakraborty, D.P. *et al.*, *Phytochemistry*, 1978, **17**, 834-855 (*Mukonine, synth*)

Knölker, H.-J. *et al.*, *Tet. Lett.*, 1993, **49**, 11221-11236 (*synth*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 133-140 (*Clausine E*)

Ho, C. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 48-52 (*Clauszoline I*)

Knölker, H.J. *et al.*, *Tet. Lett.*, 1997, **38**, 533-536 (*synth*)

Bringmann, G. *et al.*, *Synthesis*, 1998, 1501-1505 (*Mukoic acid, Mukonine, Clausine E, synth*)

Brenna, E. *et al.*, *Tetrahedron*, 1998, **54**, 1585-1588 (*synth*)

Zempoalteca, A. *et al.*, *Heterocycles*, 2002, **57**, 259-267 (*synth*)

Bergman, J. *et al.*, *Org. Prep. Proced. Int.*, 2006, **38**, 593-599 (*Clausine E, synth*)

2-Hydroxy-9H-carbazole-3-carboxylic acid, 9CI

[14501-64-5]

C₁₃H₉NO₃ 227.219

Mp 273-274°.

Me ester: Mukonidine. 2-Hydroxy-3-carbomethoxycarbazole

[24949-00-6]

C₁₄H₁₁NO₃ 241.246

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (C₆H₆/CHCl₃). Mp 168-170° Mp 188° Mp 190° Mp 245°.

Doubts have been expressed as to the correctness of this struct.; see below. λ_{max} (EtOH). λ_{max} 230 (log ε 4.8); 248 (log ε 4.57); 266 (log ε 4.64); 272 (log ε 4.73); 308 (log ε 3.88); 316 (log ε 3.78); 323 (log ε 3.18) (EtOH).

Me ether, Me ester: 2-Methoxy-3-carbomethoxycarbazole. Clausine L. O-Methylmukonidine

[14501-65-6]

C₁₅H₁₃NO₃ 255.273

Alkaloid from stem bark of *Murraya koenigii* (curryleaf tree) and leaves of *Clausena excavata* (Rutaceae). Cryst. (C₆H₆/CHCl₃). Plates (Me₂CO). Mp 133-135° Mp 205°. Methylmukonidine and Clausine L not compared. Phys. data are not identical. λ_{max} 227 (log ε 4.5); 250 (log ε 4.28); 255 (log ε 4.1); 285 (log ε 4.13) (EtOH).

Chakraborty, D.P. *et al.*, *Indian J. Chem.*, 1966, **4**, 416 (*synth*)

Joshi, B.S. *et al.*, *J.C.S. (C)*, 1969, 1518 (*synth*)

Chakraborty, D.P. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 1114-1115 (*Mukonidine, isol, uv, ir, ms, struct, synth*)

Wu, T.S. *et al.*, *Phytochemistry*, 1993, **32**, 449-451 (*Clausine L*)

Bhattacharyya, P. *et al.*, *Phytochemistry*, 1994, **35**, 1085-1086 (*O-Methylmukonidine*)

Forke, R. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 2481-2483 (*Clausine L, synth*)

6-Hydroxy-9H-carbazole-3-carboxylic acid

H-454

8-Hydroxy-2-carbazolecarboxylic acid (obsol.)

C₁₃H₉NO₃ 227.219

Me ether, Me ester: Methyl 6-methoxy-9H-carbazole-3-carboxylate. Methyl 8-methoxy-2-carbazolecarboxylate (obsol.)

[132922-58-8]

C₁₅H₁₃NO₃ 255.273

Alkaloid from the roots of *Clausena lansium* (wampee) (Rutaceae). Prisms (Et₂O/hexane). Mp 147-149°.

Hewlin, M.J.E. *et al.*, *J. Chem. Res., Synop.*, 1986, 292 (*synth*)

Li, W.-S. *et al.*, *Phytochemistry*, 1991, **30**, 343 (*isol*)

7-Hydroxy-9H-carbazole-3-carboxylic acid, 9CI

H-455

C₁₃H₉NO₃ 227.219

Me ester: Clausine M

[250259-33-7]

C₁₄H₁₁NO₃ 241.246

Alkaloid from *Clausena excavata*. Yellowish needles (Me₂CO). Mp 200-203°.

Me ether: 7-Methoxy-9H-carbazole-3-carboxylic acid. Clausine N

[250259-34-8]

C₁₄H₁₁NO₃ 241.246

Alkaloid from *Clausena excavata*. Yellow powder (Me₂CO). Mp 215-218°.

Me ether, Me ester: 3-Carbomethoxy-7-methoxycarbazole. Clausine C. Clauszoline L

[186002-61-9]

C₁₅H₁₃NO₃ 255.273

Alkaloid from stem bark of *Clausena excavata*. Yellowish needles (Me₂CO). Mp 195-197°. λ_{max} 220; 239 (sh); 249; 282; 308 (sh); 320 (sh) (MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 1427-1429; 1999, **52**, 523-527 (*Clausines*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 48-52 (*Clauszoline L*)


Krahl, M.P. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3215-3219 (*synth*)

Watanabe, T. *et al.*, *Chem. Comm.*, 2007, 4516-4518 (*Clausine C, synth*)

4-Hydroxy-β-carboline-1-carboxaldehyde

H-456

4-Hydroxy-9H-pyrido[3,4-b]indole-1-carboxaldehyde, 9CI. 1-Formyl-4-hydroxy-β-carboline [74690-70-3]



C₁₂H₈N₂O₂ 212.207

Isol. from *Picrasma quassioides* (Simaroubaceae). Inhibitor of xanthine oxidase. Cryst. (MeOH aq). Mp 296-298°. λ_{max} 210; 240; 283; 370 (EtOH).

Me ether: 4-Methoxy-9H-pyrido[3,4-b]indole-1-carboxaldehyde. 1-Formyl-4-methoxy-β-carboline. Kumujancine [92631-69-1]

C₁₃H₁₀N₂O₂ 226.234

Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Yellow needles. Mp 209-210°.

Ger. Pat., 1980, 2 941 449; CA, **93**, 114495 (*isol, synth, uv, ir*)


Yang, J. *et al.*, *Huaxue Xuebao*, 1984, **42**, 679-683; CA, **101**, 207605a

4-Hydroxy-β-carboline-1-carboxylic acid

H-457

4-Hydroxy-9H-pyrido[3,4-b]indole-1-carboxylic acid, 9CI

[74690-79-2]



C₁₂H₈N₂O₃ 228.207

Needles (MeOH/THF). Mp 158-159° dec.

Me ester: [74690-72-5]

C₁₃H₁₀N₂O₃ 242.234

Isol. from *Picrasma quassioides*. Prisms (dioxan). Mp 241-242°. λ_{max} 280; 304; 350; 365 (EtOH).

Me ether, Me ester: Methyl 4-methoxy-β-carboline-1-carboxylate. 4-Methoxy-1-methoxycarbonyl-β-carboline [60807-25-2]

C₁₄H₁₂N₂O₃ 256.26

Alkaloid from root bark and cell cultures of *Ailanthus altissima*. Pale yellow needles (C₆H₆). Mp 191-192°. λ_{max} 236 (log ε 4.32); 247 (log ε 4.29); 267 (log ε 4.41); 278 (log ε 4.31); 303 (log ε 3.94); 363 (log ε 3.87) (EtOH).

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1532-1536 (*Me ether Me ester, synth, uv, ir, pmr, ms*)

Ger. Pat., 1980, 2 941 449; CA, **93**, 114495 (*Me ester, isol*)

Varga, E. *et al.*, *Planta Med.*, 1980, **40**, 337-339 (*Me ether Me ester, isol*)

Crespi-Pellerino, N. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1010-1014 (*Me ether Me ester, isol*)

Suzuki, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 2170-2172 (*Me ether Me ester, synth*)

Suzuki, H. *et al.*, *Tetrahedron*, 1997, **53**, 1593 (*Me ether Me ester, synth*)

6-Hydroxy-β-carboline-1-carboxylic acid

H-458

6-Hydroxy-9H-pyrido[3,4-b]indole-1-carboxylic acid, 9CI

C₁₂H₈N₂O₃ 228.207

Me ester: Begonanline

C₁₃H₁₀N₂O₃ 242.234

Alkaloid from the rhizomes of *Begonia nantoensis*. Yellow syrup. λ_{\max} 206 (log ϵ 3.82); 267 (log ϵ 3.04); 316 (log ϵ 2.56); 392 (log ϵ 2.04) (MeOH).

Wu, P.-L. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 345-349 (*isol*, *pmr*, *cmr*)

7-Hydroxy- β -carboline-1-carboxylic acid H-459

7-Hydroxy-9H-pyrido[3,4-b]indole-1-carboxylic acid, 9CI

C₁₂H₈N₂O₃ 228.207

Me ether, *Me ester*: **Harmic acid methyl ester**. *Methyl harmate* [57498-79-0]

C₁₄H₁₂N₂O₃ 256.26

Alkaloid from *Banisteriopsis caapi* (Malpighiaceae). Light-yellow needles (CHCl₃). Mp 118° dec. λ_{\max} 256 (log ϵ 4.29); 282 (log ϵ 4.25); 323 (log ϵ 4.12) (CHCl₃) (Solvent reported may be incorr).

Me ether, *amide*: 7-Methoxy- β -carboline-1-carboxamide. **Harmic amide**

[62230-09-5]

C₁₃H₁₁N₃O₂ 241.249

Alkaloid from *Banisteriopsis caapi* (Malpighiaceae). Needles (CHCl₃/MeOH). Mp 226-227° dec. λ_{\max} 255 (log ϵ 4.54); 279 (log ϵ 4.44); 321 (log ϵ 4.35) (CHCl₃) (Solvent reported may be incorr).

3,4-Dihydro, *Me ether*: 3,4-Dihydro-7-hydroxy- β -carboline-1-carboxylic acid.

Harmalinic acid


[57498-77-8]

C₁₃H₁₂N₂O₃ 244.249

Alkaloid from *Banisteriopsis caapi* (Malpighiaceae). Yellow plates (MeOH). Mp 224-225°. λ_{\max} 250 (log ϵ 3.9); 372 (log ϵ 4.36) (MeOH).

Hashimoto, Y. *et al.*, *Phytochemistry*, 1975, **14**, 1633-1635; 1976, **15**, 1559-1560 (*Methyl harmate*, *Harmic amide*, *Harmalinic acid*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)

7-Hydroxyceve-4-ene-3,18-dione H-460



C₂₇H₃₉NO₃ 425.61

(7 β ,25 α)-form

Veralodine

[41787-59-1]

Alkaloid from *Veratrum lobelianum* (Liliaceae). Cryst. (Me₂CO). Mp 254-257°. [α]_D +96.4 (c, 0.89 in CHCl₃).


Ac: Mp 249-250°. [α]_D +76.4 (c, 0.102 in CHCl₃).

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 770; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 751-766 (*isol*, *ir*, *pmr*, *struct*)

Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 630 (*cd*, *config*)

20-Hydroxyceve-8-en-3-one H-461

8,9-Didehydro-20-hydroxy-3-cevanone, 9CI



C₂₇H₄₁NO₂ 411.626

(5 α ,20 β ,25 β)-form

Edpetisine

[61966-02-7]


Alkaloid from *Petilium eduardii* (aerial parts) (Liliaceae). Cryst. (Me₂CO). Mp 169-171°. [α]_D +5.31 (c, 1.6 in CHCl₃).

Nabiev, A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 403-404; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 357-358 (*isol*, *ir*, *ms*, *pmr*, *struct*)

14-Hydroxycodeine H-462

7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6,14-diol, 9CI

[4829-46-3]



Absolute Configuration

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Papaver bracteatum*. Metab. of Thebaine (Papaveraceae). Yellow cryst.

6-Ketone: **14-Hydroxycodeinone**

[508-54-3]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Papaver bracteatum* (Papaveraceae). Pale-yellow oil.

► G8225000

Misra, A.L. *et al.*, *J. Chromatogr.*, 1972, **71**, 554 (*tlc*)


Theuns, H.G. *et al.*, *Phytochemistry*, 1977, **16**, 753 (*isol*, *struct*)

Coop, A. *et al.*, *Tetrahedron*, 1999, **55**, 11429-11436 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HJX500

3-(4-Hydroxy-3,5-dimethoxybenzylidene)-2-indolinone H-463

[193621-08-8]



C₁₇H₁₅NO₄ 297.31


(*E*)-form [189098-75-7]

Alkaloid from the roots of *Isatis indigotica* (Brassicaceae). Yellow prisms. Mp 204-206°. [α]_D +90.7 (c, 0.1 in EtOH). λ_{\max} 211 (log ϵ 4.44); 256 (log ϵ 4.22); 379 (log ϵ 4.22) (MeOH).

Wu, X. *et al.*, *Planta Med.*, 1997, **63**, 55-57 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

2-Hydroxy-3,8-dimethoxy-pavinane H-464

5,6,11,12-Tetrahydro-2,9-dimethoxy-13-methylidibenzo[a,e]cycloocten-5,11-imino-3-ol, 9CI. 2,9-Dimethoxy-3-hydroxypavinane (*incorr.*). 3,8-Dimethoxy-9-hydroxypavinane (*incorr.*)



C₁₉H₂₁NO₃ 311.38

(-)-form [41498-94-6]

Alkaloid from the above-ground parts of *Argemone munita* var. *rotundata* (Papaveraceae). Cryst. (MeOH). Mp 197-198°. [α]_D²⁷ -254 (c, 1.59 in MeOH).


(\pm)-form [41498-95-7]

Synthetic. Mp 162°.

Coomes, R.M. *et al.*, *J.O.C.*, 1973, **38**, 3701 (*isol*, *ir*, *pmr*, *ms*, *struct*, *synth*)

3-Hydroxy-4,5-dimethoxyphenethylamine H-465

5-(2-Aminoethyl)-2,3-dimethoxyphenol, 9CI, 8CI. 3-Demethylmescaline [16046-07-4]



C₁₀H₁₅NO₃ 197.233

Alkaloid from mescal (*Lophophora williamsii*), *Trichocereus pachanoi* (Cactaceae) and *Acacia rigidula*. Prob. intermed. in mescaline biosynth.; precursor of Pelotine and Anhalidine in *L.*

williamsii. Plant growth inhibitor. Identified by ms.

Hydrochloride: [13062-71-0]

Flat needles (EtOH/C₆H₆). Mp 178-179°.

N-Formyl: N-Formyl-3-demethylmescaline

C₁₁H₁₅NO₄ 225.244

Alkaloid from *Lophophora williamsii* (Cactaceae).

N-Ac: N-Acetyl-3-demethylmescaline. N-[2-(3-Hydroxy-4,5-dimethoxyphenylethyl)acetamide, 9CI

[4958-20-7]

C₁₂H₁₇NO₄ 239.271

Alkaloid from *Lophophora williamsii* (Cactaceae).

N-Me: 3-Hydroxy-4,5-dimethoxymethylphenethylamine

[35144-96-8]

C₁₁H₁₇NO₃ 211.26

Present in *Lophophora williamsii* as immediate precursor of Pelletine and Anhalidine (Cactaceae). Mp 159-161° (as hydrochloride, synthetic).

N-Di-Me: 3-Hydroxy-4,5-dimethoxydimethylphenethylamine. 5-[2-(Dimethylamino)ethyl]-2,3-dimethoxyphenol, 9CI. N,N-Dimethyl-3-hydroxy-4,5-dimethoxyphenethylamine. 3-Demethyltrichocereine

[38184-65-5]

C₁₂H₁₉NO₃ 225.287

Alkaloid from *Pelecyphora aselliformis*, also detected in *Lophophora williamsii* (Cactaceae). Mp 155-156° (as oxalate).

Späth, E. *et al.*, *Monatsh. Chem.*, 1922, **43**, 93 (synth)

Agurell, S. *et al.*, *Chem. Comm.*, 1968, 1638

Kapadia, G.J. *et al.*, *Chem. Comm.*, 1968, 1688 (isol, ms, derivs)

Lee, F.G.H. *et al.*, *J. Med. Chem.*, 1969, **12**, 959 (synth, deriv)

Agurell, S. *et al.*, *J. Nat. Prod.*, 1969, **32**, 206 (isol)


Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1969, **58**, 1157 (isol, ms, synth)

Lundstrom, J. *et al.*, *Acta Pharm. Suec.*, 1971, **8**, 485; *CA*, **76**, 56708y (deriv)

Neal, J.M. *et al.*, *Science (Washington, D.C.)*, 1972, **176**, 1131 (deriv)

4-[(2-Hydroxy-4,5-dimethoxyphenyl)amino]-3-hydroxy-4-oxobutanoic acid H-466

N-(2-Hydroxy-4,5-dimethoxyphenyl)- α -malamic acid




C₁₂H₁₅NO₇ 285.253

Constit. of the leaves of *Justicia ghiesbreghtiana*. Pink cryst. Mp 175-176°. [α]_D²⁵ -56 (c, 1 in MeOH). λ_{\max} 252 (ε 7250); 302 (ε 7010) (MeCN).

Ismail, L.D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1174-1176 (isol, synth, uv, ir, pmr, cmr, ms)

5-Hydroxy-3,4'-dimethyl-[1,2'-bipyrrole]-2,5'(2'H,5'H)-dione, 9CI H-467

(1,5-Dihydro-3-methyl-2-oxo-2H-pyrrol-5-yl)-1,5-dihydro-5-hydroxy-3-methyl-2H-pyrrol-2-one. 5-Hydroxy-3-methyl-1-(3-methyl-2-oxo-2H-pyrrolin-5-yl)-3-pyrrolin-2-one [79641-96-6]



C₁₀H₁₂N₂O₃ 208.216

Alkaloid from the bulbs of *Lilium candidum* (Liliaceae). Cryst. (CHCl₃/MeOH). Mp 194-198°.

3',4'-Dihydro: 1,5-Dihydro-5-hydroxy-3-methyl-1-(4-methyl-5-oxo-2-pyrrolidinyl)-2H-pyrrol-2-one, 9CI. 5-Hydroxy-3-methyl-1-(3-methyl-2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one [114054-73-8]

C₁₀H₁₄N₂O₃ 210.232

Isol. from petals of *Lilium candidum* (Liliaceae). Mp 169-171°. [α]_{Hg}²⁰ +206 (c, 0.15 in MeOH).

5-Deoxy, 3',4'-dihydro: 1,5-Dihydro-3-methyl-1-(4-methyl-5-oxo-2-pyrrolidinyl)-2H-pyrrol-2-one, 9CI. 3-Methyl-1-(3-methyl-2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one [114020-01-8]

C₁₀H₁₄N₂O₂ 194.233


Isol. from the petals of *Lilium candidum* (Liliaceae). Mp 172-174°. [α]_{Hg}²⁰ +248 (c, 0.25 in MeOH).

Yakushijin, K. *et al.*, *Heterocycles*, 1981, **16**, 1157 (synth)

Haladová, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1988, **53**, 157; 1991, **56**, 436 (isol, struct)

2-(14-Hydroxy-14,15-dimethylhexadecyl)-4(1H)-quinolinone H-468

4-Hydroxy-2-(14-hydroxy-14,15-dimethylhexadecyl)quinoline



C₂₇H₄₃NO₂ 413.642

(+)-form

Alkaloid from the leaves of *Dictyoloma vandellianum*. Yellow gum. [α]_D +3 (c, 0.2 in CHCl₃). λ_{\max} 236; 314; 325 (MeOH).

3-Methoxy: 2-(14-Hydroxy-14,15-dimethylhexadecyl)-3-methoxy-4(1H)-quinolinone


C₂₈H₄₅NO₃ 443.668

Alkaloid from the leaves of *Dictyoloma vandellianum*. Yellow gum. [α]_D +4 (c, 0.3 in CHCl₃). λ_{\max} 237; 288; 335 (MeOH).

Sartor, C.F.P. *et al.*, *Phytochemistry*, 2003, **63**,

185-192 (isol, pmr, cmr, ms)

11-Hydroxy-4,12-dimethyl-13-(2-methyl-4-thiazolyl)-8,12-tridecadien-3-one H-469



R = H

C₁₉H₂₉NO₂S 335.51

Prod. by *Sorangium cellulosum*. Precursor of Epothilones. Amorph. solid. [α]_D²² -25 (c, 0.5 in MeOH). λ_{\max} 212 (ε 17700); 247 (ε 13400) (MeOH).

Homologue (R = CH₃): 11-Hydroxy-4,8,12-trimethyl-13-(2-methyl-4-thiazolyl)-8,12-tridecadien-3-one


C₂₀H₃₁NO₂S 349.536

Prod. by *Sorangium cellulosum*. Precursor of Epothilones. Amorph. solid. [α]_D²² -9 (c, 0.02 in MeOH). λ_{\max} 203 (ε 19100); 244 (ε 12500) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

9-Hydroxy-6,10-dimethyl-11-(2-methyl-4-thiazolyl)-6,10-undecadien-3-one H-470

4-(3-Hydroxy-2,6-dimethyl-9-oxo-1,5-undecadienyl)-2-methylthiazole




C₁₇H₂₅NO₂S 307.456

Epothilone fragment. Isol. from recombinant *Myxococcus xanthus*. Yellow oil. [α]_D +1.8 (c, 0.027 in MeOH). λ_{\max} 211 (ε 17600); 247 (ε 12800) (MeOH).

Starks, C.M. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1313-1317 (isol)

6-Hydroxy-2,6-dimethyl-2,7-octadienoic acid H-471

Linalolic acid. Linalool-1-oic acid. Menthaifolic acid [28420-25-9]



C₁₀H₁₆O₃ 184.235

(S,E)-form [75979-26-9]

Constit. of *Juniperus thurifera*, *Monilinia mali* and *Penstemon ambiguus*. [α]_D +16.8 (c, 1.1 in CHCl₃).


Amide: Concinnamide. Acacialactam (in-corr.) [126262-46-2]

C₁₀H₁₇NO₂ 183.25

Constit. of *Acacia concinna*. Oil. [α]_D +4.3 (c, 0.16 in MeOH). Struct. revised


in 1994. λ_{\max} 210 (ϵ 15700) (MeOH).
[67711-07-3 , 121843-29-6 , 34935-27-8]
Sekine, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**,
3164-3165; 1997, **45**, 148 (*Concinnamide*)

3-Hydroxy-2,3-dimethyl-5-oxo-2-pyrrolidinedicarboxylic acid H-472
3-Hydroxy-2,3-dimethyl-5-oxoproline



$C_7H_{11}NO_4$ 173.168
Residue present in the lipopolysaccharide of *Pseudomonas fluorescens* 361.
Naberezhnykh, G.A. *et al.*, *Bioorg. Khim.*, 1987, **13**, 1426

18-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13,19-triene-1,21-dione H-473



Relative Configuration

$C_{28}H_{35}NO_3$ 433.589
Metab. of an unidentified *Daldinia* fungus. Amorph. solid. $[\alpha]_D$ -98.4 (c, 0.34 in $CHCl_3$).

6 β ,7 β -Epoxyde: 6,7-Epoxy-18-hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-13,19-diene-1,21-dione
 $C_{28}H_{35}NO_4$ 449.589
From *Daldinia* sp. Amorph. solid. $[\alpha]_D$ -57.7 (c, 0.23 in $CHCl_3$).

12-Hydroxy: 12,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13,19-triene-1,21-dione
 $C_{28}H_{35}NO_4$ 449.589
From *Daldinia* sp. Amorph. solid. $[\alpha]_D$ -70.4 (c, 0.31 in $CHCl_3$).

19,20-Dihydro, 19 β -hydroxy, 6 β ,7 β -epoxyde: 6,7-Epoxy-18,19-dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-13-ene-1,21-dione
 $C_{28}H_{37}NO_5$ 467.604
From *Daldinia* sp. Amorph. solid. $[\alpha]_D$ -49 (c, 0.35 in $CHCl_3$).

19,20-Dihydro, 19 β -methoxy, 6 β ,7 β -epoxyde: 6,7-Epoxy-18-hydroxy-19-methoxy-16,18-dimethyl-10-phenyl[11]cytochalasa-13-ene-1,21-dione
 $C_{29}H_{39}NO_5$ 481.631
Isol. from *Microporellus subsessilis*. Amorph. solid (MeOH). $[\alpha]_D^{20}$ -115 (c, 0.0007 in $CHCl_3$).

19,20-Dihydro, 19 β -methoxy, 12-hydroxy: 12,18-Dihydroxy-19-methoxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13-


diene-1,21-dione
 $C_{29}H_{39}NO_5$ 481.631
Isol. from *Microporellus subsessilis*. Needles (MeOH). Mp 136-138°. $[\alpha]_D^{28}$ -102 (c, 0.0003 in $CHCl_3$).

18-Deoxy, 17 β -hydroxy: 17-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13,19-triene-1,21-dione
 $C_{28}H_{35}NO_3$ 433.589
From *Daldinia* sp. Amorph. solid. $[\alpha]_D$ -126.6 (c, 0.54 in $CHCl_3$).

Buchanan, M.S. *et al.*, *Phytochemistry*, 1996, **41**, 821 (*isol, pmr, cmr, ms, struct*)
Kurnia, D. *et al.*, *Phytochemistry*, 2007, **68**, 697-702 (*Microporellus constits*)

2-Hydroxy-3,5-dimethylpyrazine, 9CI H-474

3,5-Dimethyl-2(1H)-pyrazinone, 9CI.
3,5-Dimethyl-2-pyrazinol
[60187-00-0]



$C_6H_8N_2O$ 124.142
NH-Form predominates. Cryst. (EtOH). Mp 149.5-151°.

OH-form

Me ether: 2-Methoxy-3,5-dimethylpyrazine, 9CI
[92508-08-2]
 $C_7H_{10}N_2O$ 138.169
Isol. from *Crotalaria ochroleuca* seeds and prod. by microorganisms, e.g. *Chondromyces crocatus*. Causes objectionable odour in foods. Liq. with unpleasant musty odour.

Klein, B. *et al.*, *J.O.C.*, 1964, **29**, 2623-2626 (*synth*)
Macdonald, J.C. *et al.*, *Tetrahedron*, 1976, **32**, 655-660 (*pmr, cmr*)
Ohta, A. *et al.*, *J. Het. Chem.*, 1983, **20**, 311-320 (*synth*)
Mottram, D.S. *et al.*, *Chem. Ind. (London)*, 1984, 448-449 (*Me ether, isol, synth, ms*)
Leitner, H. *et al.*, *Chem. Mikrobiol. Technol. Lebensm.*, 1990, **12**, 151-158; *CA*, **113**, 3275f (*Me ether, isol*)
Simpson, R.F. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 5425-5430 (*Me ether, isol, synth, pmr, cmr, ms*)
Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*Me ether, isol, synth, pmr, cmr*)

3-Hydroxy-2,5-dimethylpyrazine H-475

3,6-Dimethyl-2(1H)-pyrazinone, 9CI.
3,6-Dimethyl-2-pyrazinol
[16289-18-2]

$C_6H_8N_2O$ 124.142
Cryst. (C_6H_6). Mp 210-211°.

OH-form

Me ether: 3-Methoxy-2,5-dimethylpyrazine, 9CI
[19846-22-1]
 $C_7H_{10}N_2O$ 138.169
Prod. by *Chondromyces crocatus*. Liq. with musty odour. Bp₂ 41°.
Klein, B. *et al.*, *J.O.C.*, 1964, **29**, 2623 (*synth*)

Macdonald, J.C. *et al.*, *Tetrahedron*, 1976, **32**, 655 (*cmr, pmr*)
Ohta, A. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 2027 (*synth*)
Mottram, D.S. *et al.*, *Chem. Ind. (London)*, 1984, 448 (*synth, ms, deriv*)
Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*)

4-Hydroxy-1,1-dimethylpyrrolidinium-2-carboxylate H-476

2-Carboxy-4-hydroxy-1,1-dimethylpyrrolidinium hydroxide inner salt, 9CI. 4-Hydroxyproline betaine
[105869-44-1]



$C_7H_{13}NO_3$ 159.185

(2R,4R)-form

Turicine. Combretine A
[515-24-2]
[12785-71-6 (Combretine A)]
Alkaloid from *Betonica wiesin* (preferred genus name *Stachys*), *Canavalia ensiformis* (jack bean), *Combretum micranthum* and *Stachys sylvatica*. Needles or prisms+1H₂O (EtOH aq.). V. sol. H₂O. Mp 260° dec. $[\alpha]_D^{20}$ +37.5 (c, 2.4 in H₂O). Identity of Combretine A not certain.

O-(4-Bromo-1H-pyrrole-2-carbonyl):

Damituricine
[949091-37-6]
 $C_{12}H_{15}BrN_2O_4$ 331.166
Alkaloid from *Axinella damicornis*. Serotonin receptor modulator. $[\alpha]_D^{20}$ +10.7 (c, 0.001 in H₂O).

(2S,4R)-form

Betonicine. Achillein
[515-25-3]
Constit. of *Achillea millefolium* (yarrow), *Betonica officinalis* (preferred genus name *Stachys*), *Marrubium vulgare* (horehound) and *Stachys sylvatica*. Shows antiinflammatory props. Prisms (EtOH). Mp 252° dec. $[\alpha]_D^{15}$ -36.6 (H₂O).
4-O-(4-Hydroxy-E-cinnamoyl): 4-O-p-Coumaroylbetonicine
 $C_{16}H_{19}NO_5$ 305.33
Constit. of *Phlomis bruneogaleata*. Plates (MeCN/MeOH). Mp 239-241°. $[\alpha]_D^{20}$ +3.9 (c, 0.1 in MeOH). λ_{\max} 229; 304 (MeOH).

(2RS,4RS)-form

Combretine B
[12785-70-5]
Alkaloid from the leaves of *Combretum micranthum* (Combretaceae). Mp 248-249° dec. $[\alpha]_D^{20}$ 0 (c, 2.7 in H₂O). Identity not certain.


[119303-01-4 , 101198-81-6]

Schulze, E. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1911, **76**, 258; 1912, **79**, 236 (*isol*)
Goodson, J.A. *et al.*, *J.C.S.*, 1919, **115**, 923 (*isol*)

- Paudler, W.W. *et al.*, *Chem. Ind. (London)*, 1963, 1693-1694 (*isol. struct*)
 Mandava, N. *et al.*, *Annalen*, 1970, **741**, 167 (*pmr, config*)
 Ogan, A.U. *et al.*, *Planta Med.*, 1972, **21**, 210-217 (*Combretines A,B*)
 Musich, J.A. *et al.*, *J.O.C.*, 1977, **42**, 139 (*synth*)
 Blunden, G. *et al.*, *Magn. Reson. Chem.*, 1986, **24**, 965-971 (*pmr, cmr*)
 Jones, G.P. *et al.*, *Acta Cryst. C*, 1988, **44**, 2208-2211 (*cryst struct*)
 Blunden, G. *et al.*, *Biochem. Syst. Ecol.*, 1996, **24**, 71-81 (*occur*)
 Kirmizibekmez, H. *et al.*, *Planta Med.*, 2004, **70**, 711-717 (*4-Coumaroylbetonicine*)
 Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 5877-5887 (*Damituricine*)

4-Hydroxy-2,3-dimethylquinoline H-477

2,3-Dimethyl-4(1H)-quinolinone, 9CI.
 2,3-Dimethyl-4-quinolinol, 4-Hydroxy-3-methylquinaldine
 [58596-45-5]



- $C_{11}H_{11}NO$ 173.214
 Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae). Prisms + H_2O (H_2O). Spar. sol. EtOH. Mp 315°. Subl. at ca. 300°.
 N-Acetoxyethyl: 1-Acetoxyethyl-2,3-dimethyl-4(1H)-quinolinone
 [151077-57-5]
 $C_{14}H_{15}NO_3$ 245.277
 Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae). Needles ($CHCl_3/MeOH$). Mp 175-178°.
 Conrad, M. *et al.*, *Ber.*, 1891, **24**, 2990-2992 (*synth*)
 Mander-Jones, B. *et al.*, *CA*, 1933, **27**, 1350 (*synth*)
 Royer, R. *et al.*, *J.C.S.*, 1948, 106-110 (*synth*)
 Scheuer, P.J. *et al.*, *J.C.S.*, 1963, 5569-5571 (*synth, ir, uv, pmr*)
 Ahsan, M. *et al.*, *Phytochemistry*, 1993, **33**, 1507 (*isol, uv, ir, pmr, cmr, ms, struct*)

6-Hydroxy-4,8-dimethylquinoline H-478


- 4,8-Dimethyl-6-quinolinol, 9CI
 [194143-58-3]
 $C_{11}H_{11}NO$ 173.214
 Alkaloid from the marine cyanobacterium *Lyngbya majuscula*. Amorph. powder. λ_{max} 234 (log ϵ 4.47); 283 (log ϵ 3.51); 322 (log ϵ 3.28); 333 (log ϵ 3.28) (MeOH).
 O-(2,4-Di-O-methyl- β -D-xylopyranoside): [194143-57-2]
 $C_{18}H_{23}NO_5$ 333.383
 From *Lyngbya majuscula*. Amorph. powder. $[\alpha]_D^{25}$ -56.7 (c, 0.15 in $CHCl_3$). λ_{max} 236 (log ϵ 4.65); 286 (log ϵ 3.57); 316 (log ϵ 3.44); 328 (log ϵ 3.44) (MeOH).
 Orjala, J. *et al.*, *Phytochemistry*, 1997, **45**, 1087-1090 (*isol, uv, ir, pmr, cmr, ms, struct*)

6-Hydroxy-5,7-dimethylquinoline H-479

- 5,7-Dimethyl-6-quinolinol, 9CI. *Phlegmariurine N*
 [126552-19-0]
 $C_{11}H_{11}NO$ 173.214
 Alkaloid *isol.* from *Phlegmariurus fordii* (preferred genus name *Huperzia*). Mp 179-181°. λ_{max} 328 (log ϵ 3.71) (EtOH).
 Miao, Z.C. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 114-117; *CA*, **112**, 179528j (*isol, pmr, cmr*)

4-Hydroxy-N,N-dimethyltryptamine H-480

3-(2-Dimethylaminoethyl)-1H-indol-4-ol, 9CI. 3-(2-Dimethylaminoethyl)-4-hydroxyindole. *Psilocine*. CX 59
 [520-53-6]




- $C_{12}H_{16}N_2O$ 204.271
 Alkaloid from fungi *Psilocybe* and *Conocybe* spp., *Copelandia chlorocystis* and from *Stropharia* spp. Euphoric, hallucinogenic. Sol. MeOH, butanol; fairly sol. H_2O . Mp 173-176° dec. Log P 1.16 (calc). λ_{max} 222 (ϵ 39800); 260 (ϵ 5000); 267 (ϵ 6300); 283 (ϵ 5000); 293 (ϵ 4000) (MeOH) (Berdy).
 ▶ LD₅₀ (rat, ivn) 75 mg/kg. NM2625000
 O-Phosphate: *Psilocybine*, BAN, INN. *Indocybin*. CY 39
 [520-52-5]
 $C_{12}H_{17}N_2O_4P$ 284.251
 Active principle of hallucinogenic mushrooms *Psilocybe mexicana* and other *Psilocybe* spp. Also from *Stropharia* spp. *Inocybe aeruginascens* and *Conocybe cyanopus*. Psychotomimetic agent, hallucinogen. Cryst. (MeOH or H_2O). Sol. MeOH, H_2O , butanol. Mp 220-228° (185-195°). λ_{max} 221 (ϵ 18620); 268 (ϵ 6920); 280 (ϵ 5500); 290 (ϵ 4360) (MeOH) (Berdy). λ_{max} 220 (ϵ 39800); 267 (ϵ 6300); 290 (ϵ 4000) (EtOH) (Berdy).
 ▶ Human adverse systemic effects by ingestion and intraperitoneal routes incl. toxic psychosis. LD₅₀ (rat, ivn) 280 mg/kg; LD₅₀ (mus, ipr) 420 mg/kg. NM3150000
 N^b-Me, O-phosphate: N^b-Methylpsilocybine. *Aeruginascine*
 [114264-95-8]
 $C_{13}H_{20}N_2O_4P^{\oplus}$ 299.286
 Quaternary alkaloid from *Inocybe aeruginascens*. Glassy solid. Counterion not specified. λ_{max} 219 (log ϵ 3.92); 267 (log ϵ 3.2); 282 (sh) (log ϵ 3.1); 288 (sh) (log ϵ 3) (H_2O).
 N-De-Me, O-phosphate: *Baeocystine*
 [21420-58-6]
 $C_{11}H_{15}N_2O_4P$ 270.224
 Alkaloid from *Psilocybe semilanceolata* and *Psilocybe baeocystis*. Mp 254-258° dec.
 Di-N-de-Me, O-phosphate: *Norbaeocys-*

tine
 [21420-59-7]
 $C_{10}H_{13}N_2O_4P$ 256.197
 Alkaloid from *Psilocybe baeocystis*. Mp 188-192° dec.

- Julia, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1959, **269**, 51 (*synth*)
 Hofmann, A. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 1557-1572 (*isol, synth, ir, uv*)
 Leung, A.Y. *et al.*, *J. Pharm. Sci.*, 1967, **56**, 146; 1968, **57**, 1667-1671 (*Baeocystine, Norbaeocystine*)
 Agurell, S. *et al.*, *Acta Chem. Scand.*, 1968, **22**, 1210-1218 (*biosynth*)
 Hofmann, A. *et al.*, *Bull. Narc.*, 1971, **23**, 3-14 (*rev*)
 Aboul-Enein, H.Y. *et al.*, *Am. J. Pharm.*, 1974, **146**, 91-95 (*rev, Psilocybine*)
 Weber, H.P. *et al.*, *J.C.S. Perkin 2*, 1974, 942-946; 946-948 (*cryst struct*)
 Repke, D.B. *et al.*, *J. Pharm. Sci.*, 1977, **66**, 113; 743 (*isol, glc, ms, derivs*)
 Unger, S.E. *et al.*, *Anal. Lett.*, 1979, **12**, 1157-1167 (*ms*)
 White, P.C. *et al.*, *J. Chromatogr.*, 1979, **169**, 453-456 (*hplc*)
 Weeks, R.A. *et al.*, *J. Nat. Prod.*, 1979, **49**, 469-474 (*isol, uv, ms*)
 Chilton, W.S. *et al.*, *J. Psychedelic Drugs*, 1979, **11**, 61-69 (*rev*)
 Repke, D.B. *et al.*, *J. Het. Chem.*, 1981, **18**, 175 (*synth, bibl*)
 Seeger, R. *et al.*, *Dtsch. Apoth. -Ztg.*, 1985, **125**, 65-66 (*rev, Psilocybine*)
 Brenneisen, R. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 487-489 (*synth, uv, ir, pmr, cmr, P-31 nmr, ms, bibl, Baeocystine*)
 Sakagami, H. *et al.*, *Heterocycles*, 1999, **51**, 1131-1135 (*synth*)
 Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1622
 Nichols, D.E. *et al.*, *Synthesis*, 1999, 935-938 (*Psilocine, Psilocybine*)
 Shirota, O. *et al.*, *J. Nat. Prod.*, 2003, **66**, 885-887 (*synth, pmr, cmr, ms*)
 Gathergood, N. *et al.*, *Org. Lett.*, 2003, **5**, 921-923 (*synth*)
 Jensen, N. *et al.*, *Planta Med.*, 2006, **72**, 665-666 (*Aeruginascine*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, HKE000; Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PHU500

2-(4-Hydroxy-3,5-dinitrophenyl)ethylamine H-481


4-(2-Aminoethyl)-2,6-dinitrophenol, 9CI.
 4-Hydroxy-2,6-dinitrophenethylamine.
 3,5-Dinitrotyramine



- $C_8H_9N_3O_5$ 227.176
 N-Ac: N-2-[(4-Hydroxy-3,5-dinitrophenyl)ethyl]acetamide. N-Acetyl-3,5-dinitrotyramine
 $C_{10}H_{11}N_3O_6$ 269.213
 Isol. from the marine-derived *Flavobacterium* sp. T436.
 Schuhmann, I. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*isol*)

4-Hydroxyepilupinine H-482

Octahydro-2-hydroxy-2H-quinolizine-1-methanol, 9CI
[135531-65-6]



C₁₀H₁₉NO₂ 185.266

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

11-O-Benzoyl: 11-O-Benzoyl-4-hydroxyepilupinine

[135531-70-3]

C₁₇H₂₃NO₃ 289.374

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

4-O-(2-Pyrrolicarbonyl): 4-(2-Pyrrolicarbonyloxy)epilupinine

[135531-69-0]

C₁₅H₂₂N₂O₃ 278.35

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

11-O-(2-Pyrrolicarbonyl): 4-Hydroxy-11-O-(2-pyrrolicarbonyl)epilupinine

[135531-71-4]


C₁₅H₂₂N₂O₃ 278.35

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides*.

Veen, G. et al., *Phytochemistry*, 1991, **30**, 1891

3-Hydroxy-22,26-epiminocholest-22(N)-en-6-one H-483

3-Hydroxy-16,28-secosolanid-22(28)-en-6-one, 9CI



(3β,5α,25R)-form

C₂₇H₄₃NO₂ 413.642

(3β,5α,25R)-form

Obt. by acid hydrolysis of the glucoside. Amorph. powder. [α]_D²⁸ +35.5 (c, 0.22 in CHCl₃).

O-β-D-Glucopyranoside: [145867-04-5]

C₃₃H₅₃NO₇ 575.784

Alkaloid from bulbs of *Fritillaria persica* (Liliaceae). Amorph. powder. [α]_D²⁰ -44 (c, 0.50 in MeOH).

(3β,5α,20ξ,25S)-form**Petiline**

[26989-18-4]

Alkaloid from *Petilium raddeanum* and *Petilium eduardii* (Liliaceae). Antiinflammatory agent. Mp 205-206°. [α]_D -51.1 (c, 1.2 in EtOH).

Hydrochloride: Mp 288-289°.

(3β,5α,20ξ,25ξ)-form**Korsiline**

[62929-45-7]

Alkaloid from *Korolkowia sewerzowii*

(Liliaceae). Mp 194-196°. [α]_D 0 (c, 0.8 in EtOH).

Nuriddinov, R.N. et al., *Khim. Prir. Soedin.*, 1968, 168-174; 1969, 604-605; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, 145; 1969, 525 (*Petiline, isol, ir, uv, pmr, ms, struct*)

Samikov, K. et al., *Khim. Prir. Soedin.*, 1976, **12**, 827-828; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 748 (*Korsiline*)


Nakhatov, I. et al., *Khim. Prir. Soedin.*, 1981, 616-619; *CA*, **96**, 65672x (*isol*)

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1986, 345-347; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 319-321 (*Petiline, ord*)

Ori, K. et al., *Phytochemistry*, 1992, **31**, 4337-4341 (*3β,5α,25R-form, glucoside*)

β-Hydroxyergothioneine H-484

α-Carboxy-2,3-dihydro-β-hydroxy-N,N,N-trimethyl-2-thioxo-1H-imidazole-4-ethaniminium hydroxide inner salt. β-Hydroxy-2-mercaptohistidine trimethylbetaine



Absolute Configuration

C₉H₁₅N₃O₃S 245.302

Isol. from the mushroom *Lyophyllum conatum*. Hepatoprotectant. Prisms (EtOH aq.). Mp 195-200° dec. [α]_D +35 (c, 0.27 in H₂O). λ_{max} 250 (ε 9200) (H₂O). λ_{max} 235 (NaOH aq.).

Kimura, C. et al., *Biosci., Biotechnol., Biochem.*, 2005, **69**, 357-363 (*isol, uv, pmr, cmr, ms, cryst struct*)

N-(1-Hydroxyethyl)benzamide H-485

N-(1-Hydroxyethyl)-N-phenylbenzamide. N-Benzoyl-N-phenylaminomethylcarbinol PhCONPhCH(OH)CH₃

C₁₅H₁₅NO₂ 241.289

Alkaloid from *Oxytropis muricata*. Cryst. (CHCl₃). Mp 147-149°.


Hydrochloride:

Cryst. Mp 194-197°.

Duboshina, Z.N. et al., *Zh. Obshch. Khim.*, 1963, **33**, 2071; *CA*, **59**, 12851h (*isol*)

1-(1-Hydroxyethyl)-β-carboline H-486

α-Methyl-9H-pyrido[3,4-b]indole-1-methanol, 9CI



C₁₃H₁₂N₂O 212.251

λ_{max} 211 (ε 6030); 234 (ε 9770); 287 (sh) (ε 6170); 337 (ε 1020) (EtOH) (Derep).

(S)-form [110282-66-1]


Alkaloid from the marine bryozoan *Costaticella hastata*. Mp 163-169° dec. [α]_D²⁵ -11.4 (MeOH).

Blackman, A.J. et al., *J. Nat. Prod.*, 1987, **50**, 494 (*isol, uv, ir, pmr, ms, struct*)

Bracher, F. et al., *Annalen*, 1993, 837 (*synth*)

1-(2-Hydroxyethyl)-β-carboline H-487

9H-Pyrido[3,4-b]indole-1-ethanol, 9CI
[102828-03-5]



C₁₃H₁₂N₂O 212.251

Alkaloid from the stem bark of *Soulamea fraxinifolia* (Simaroubaceae). Cryst. (MeOH). Mp 192°. λ_{max} 214 (ε 20900); 235 (ε 37200); 240 (sh) (ε 38000); 250 (ε 25100); 282 (sh) (ε 10000); 288 (ε 16200); 336 (ε 4680); 350 (ε 4570) (EtOH) (Derep).

Charles, B. et al., *J. Nat. Prod.*, 1986, **49**, 303 (*isol, uv, pmr, ms, struct*)

Torisawa, Y. et al., *Tetrahedron*, 1991, **47**, 8067 (*synth*)

N-(2-Hydroxyethyl)cinnamide H-488

N-(2-Hydroxyethyl)-3-phenyl-2-propenamide, 9CI. **Idrocilamide**, INN. Brolitene. Srilane. Talval. LCB 29

[6961-46-2]
PhCH=CHCONHCH₂CH₂OH

C₁₁H₁₃NO₂ 191.229

Alkaloid from leaves of *Erythrophleum chlorostachys* (Fabaceae). Muscle relaxant, antiinflammatory agent. Cryst. (CHCl₃/petrol or EtOH/petrol). Mp 101° (97°). Log P 1.12 (calc).

▶ LD₅₀ (mus, orl) 2950 mg/kg. Adverse effects incl. gastrointestinal disturbances and drowsiness. GD7360000

N-Me: N-(2-Hydroxyethyl)-N-methylcinnamide. N-(2-Hydroxyethyl)-N-methyl-3-phenyl-2-propenamide, 9CI
[30687-16-2]

C₁₂H₁₅NO₂ 205.256

Alkaloid from leaves of *Erythrophleum chlorostachys* (Fabaceae). Cryst. (EtOH/petrol or C₆H₆). Mp 78°.

▶ LD₅₀ (mus, orl) 3000 mg/kg. GD7380000

N-Me, 4-hydroxy: N-(2-Hydroxyethyl)-N-methyl-p-hydroxycinnamide. N-(2-Hydroxyethyl)-3-(4-hydroxyphenyl)-N-methyl-2-propenamide, 9CI
[35241-60-2]

C₁₂H₁₅NO₃ 221.255

Alkaloid from leaves of *Erythrophleum chlorostachys* (Fabaceae). Cryst. (EtOH). Mp 176-177°.

Behrens, O.K. et al., *J. Biol. Chem.*, 1948, **175**, 771 (*synth*)

Griffin, W.J. et al., *Phytochemistry*, 1971, **10**, 2793 (*isol, ir, struct, synth*)


Grand, M. et al., *Eur. J. Med. Chem. (Chim. Ther.)*, 1974, **9**, 205 (*pharmacol*)

Xu, J. et al., *Zhongguo Yiyao Gongye Zazhi*, 1990, **21**, 70; *CA*, **113**, 70738x (*pharmacol, tox*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1205

3-(1-Hydroxyethyl)coronaridine H-489

[82464-33-3]



Absolute Configuration


C₂₃H₃₀N₂O₃ 382.502

Incorrectly called β-hydroxyethyl in the ref. Alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae). Amorph. [α]_D -25.4 (c, 0.4 in CHCl₃). λ_{max} 228 (log ε 4.3); 286 (log ε 3.58); 294 (log ε 3.52) (EtOH).

Feng, X.Z. et al., *Planta Med.*, 1982, **44**, 212-214 (isol, uv, pmr)

8-(1-Hydroxyethyl)dihydrochelerythrine H-490

12,13-Dihydro-1,2-dimethoxy-α,12-dimethyl[1,3]benzodioxolo[5,6-c]phenanthridine-13-methanol, 9CI




C₂₃H₂₃NO₅ 393.438

(R)-form [928006-52-4]

Alkaloid from the roots of *Zanthoxylum nitidum*. Yellow solid. Mp 152-173°. [α]_D²² +8 (c, 0.2 in CHCl₃).

Hu, J. et al., *Chem. Biodiversity*, 2006, **3**, 990-995 (isol, pmr, cmr, ms)

8-(2-Hydroxyethyl)dihydro-sanguinarine H-491



C₂₂H₁₉NO₅ 377.396

λ_{max} 232 (log ε 4.2); 284 (log ε 4.6); 322 (log ε 4.8) (MeOH).

(S)-form

Alkaloid from the roots of *Corydalis flabellata*. Cryst. (MeOH/C₆H₆). Mp 212-213°. [α]_D²⁰ +29.2 (c, 0.05 in CHCl₃).

Ac:

Cryst. (CHCl₃/petrol). Mp 180-181°.

2'-Carboxylic acid: **Spallidamine**

[129388-65-4]

C₂₂H₁₇NO₆ 391.379


Alkaloid from aerial parts of *Corydalis bulbosa* and the roots of *Corydalis flabellata* (Papaveraceae). Amorph. solid. Mp 140-141°. [α]_D +35 (c, 0.04 in CHCl₃). Abs. config. not determined but prob. S-. λ_{max} 206; 231; 286; 322 (MeOH).

Ito, C. et al., *Phytochemistry*, 1990, **29**, 2044-2045 (*Spallidamine*)

Koul, S. et al., *Planta Med.*, 2002, **68**, 262-265 (isol, pmr, cmr, ms)

2-(2-Hydroxyethyl)-5-(hydroxymethyl)-3,4-pyrrolidinediol, 9CI H-492

3,4-Dihydroxy-5-(hydroxymethyl)-1-pyrrolidinemethanol, 2,5-Imino-2,5,6-trideoxy-heptitol, 3,4-Dihydroxy-2-(2-hydroxyethyl)-5-(hydroxymethyl)pyrrolidine



(2R,3R,4R,5R)-form

C₇H₁₅NO₄ 177.2

(2R,3R,4R,5R)-form

D-manno-form. 2,5,6-Trideoxy-2,5-imino-D-manno-heptitol

[205762-20-5]

Alkaloid from the bulbs of *Hyacinthus orientalis* and *Scilla sibirica*. Inhibitor of α-glucosidase and chitin synthase. [α]_D +98.5 (c, 1.13 in H₂O). [α]_D +46 (c, 1.15 in H₂O). Significantly lower α_D recorded for the synthetic product.

3-O-β-D-Mannopyranoside:

C₁₃H₂₅NO₉ 339.342

Alkaloid from the bulbs of *Scilla sibirica*. Inhibitor of α-fucosidase. Powder. [α]_D +2 (c, 0.32 in H₂O).

3-O-[β-D-Mannopyranosyl-(1→4)-β-D-mannopyranoside]:

C₁₉H₃₅NO₁₄ 501.484

Alkaloid from the bulbs of *Scilla sibirica*. Inhibitor of α-fucosidase. Powder. [α]_D +10.4 (c, 0.69 in H₂O).

(2R,3S,4S,5S)-form

D-gulo-form. 2,5,6-Trideoxy-2,5-imino-D-gulo-heptitol

[205762-22-7]

Alkaloid from the bulbs of *Hyacinthus orientalis* and *Scilla sibirica*. Inhibitor of α-fucosidase. [α]_D +41.4 (c, 0.56 in H₂O).


Asano, N. et al., *J. Nat. Prod.*, 1998, **61**, 625-628 (isol, pmr, cmr)

Yamashita, T. et al., *J. Nat. Prod.*, 2002, **65**, 1875-1881 (*Scilla sibirica* alkaloids)

Behr, J.-B. et al., *Tetrahedron: Asymmetry*, 2002, **13**, 111-113 (D-manno-form, D-gulo-form, synth, pmr, cmr)

1-(2-Hydroxyethyl)-4-methoxy-β-carboline H-493

4-Methoxy-9H-pyrido[3,4-b]indole-1-ethanol, 9CI [77369-98-3]



C₁₄H₁₄N₂O₂ 242.277

Alkaloid from the root bark of *Ailanthus altissima* (Simaroubaceae). Pale-yellow needles. Mp 253-254° dec.

O-Ac:

Pale-yellow needles (Me₂CO). Mp 175-177°.

1'ξ-Hydroxy: 1-(4-Methoxy-9H-pyrido[3,4-b]indol-1-yl)-1,2-ethanediol, 9CI. 1-(1,2-Dihydroxyethyl)-4-methoxy-β-carboline


[77369-99-4]

C₁₄H₁₄N₂O₃ 258.276

Alkaloid from root bark of *Ailanthus altissima* (Simaroubaceae). Pale-yellow needles (CHCl₃). Mp 189-190°. [α]_D²⁰ -2.56 (c, 0.78 in EtOH).

Ohmoto, T. et al., *Chem. Pharm. Bull.*, 1981, **29**, 390-395 (isol, uv, ir, pmr, cmr, ms, struct)

5-(1-Hydroxyethyl)-4-methyl-3-pyridinecarboxylic acid H-494



C₉H₁₁NO₃ 181.191

(ξ)-form

Me ester: 3-(1-Hydroxyethyl)-4-methyl-5-methoxycarbonylpyridine

[149155-03-3]


C₁₀H₁₃NO₃ 195.218

Alkaloid from *Strychnos cocculoides* and *Strychnos spinosa*. [α]_D -27 (c, 0.58 in CHCl₃).

Delaude, C. et al., *Bull. Soc. R. Sci. Liege*, 1992, **61**, 429-440 (Me ester, isol, pmr)

5-(2-Hydroxyethyl)-4-methylthiazole H-495

4-Methyl-5-thiazoleethanol, 9CI. 2-(4-Methyl-5-thiazolyl)ethanol. FEMA 3204 [137-00-8]



C₆H₉NOS 143.209

Present in porcini (*Boletus edulis*). Oil with nut-like odour on extreme dilution. Bp₁₀ 142-145° Bp₃ 127-129°. n_D²⁰ 1.5470.

N-Me:

[16311-69-6 (iodide)]

C₇H₁₂NOS⁺ 158.244

Mp 88-89° (as iodide). Light-sensitive.

Ac: FEMA 3205

[656-53-1]

C₈H₁₁NO₂S 185.246

Flavour ingredient. Mp 131°. Bp₄ 123°.

Benzoyl: [31299-77-1]

C₁₃H₁₃NO₂S 247.317


Mp 192°. Bp₄ 185°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 643C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 111B; 116B (nmr)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1493D; 1495B (ir)
 Buchman, E.R. et al., *J.A.C.S.*, 1936, **58**, 1803 (deriv)
U.K. Pat., 1951, 641 426; *CA*, **45**, 2509
 Takeda, Y. et al., *Yakugaku Zasshi*, 1951, **71**, 84; *CA*, **45**, 9532
 Dornow, A. et al., *Chem. Ber.*, 1953, **86**, 1404
U.S. Pat., 1954, 2 654 760; *CA*, **48**, 12810
 Miyatake, K. et al., *Yakugaku Zasshi*, 1955, **75**, 1054; *CA*, **50**, 5633
 Thiel, M. et al., *Annalen*, 1958, **619**, 161
 Rubtsov, I.A. et al., *CA*, 1970, **73**, 56015
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1862; 1863 (occur, props)

6-(1-Hydroxyethyl)-1-phenazinecarboxylic acid H-496

Saphenic acid. DC 86Y. Antibiotic DC 86Y
 [94448-14-3]




$C_{15}H_{12}N_2O_3$ 268.271
 Phenazine antibiotic. Prod. by *Streptomyces luteogriseus* and *Streptomyces antibioticus*. Biol. inactive. Yellow needles. Sol. MeOH, C_6H_6 ; poorly sol. H_2O . Mp 223-225°. $[\alpha]_D^{20}$ +55.8 (c, 1.0 in DMSO). λ_{max} 252 (€ 79000); 363 (€ 14000) (MeOH) (Derep). λ_{max} 253 (€ 84000); 367 (€ 16000) (MeOH) (Berdy). λ_{max} 255 (€ 58100); 365 (€ 13200) (EtOH) (Berdy).
 ▶ LD₅₀ (mus, ipr) 300 mg/kg.
Me ester: [73634-72-7]
 $C_{16}H_{14}N_2O_3$ 282.298
 Prod. by *Streptomyces* sp. Phosphodiesterase inhibitor.
 6-Deoxy- α -L-glucopyranos-2-yl ester: 2'-L-Quinoyosyl saphenate
 [137570-43-5]
 $C_{21}H_{22}N_2O_7$ 414.414
 Isol. from a *Streptomyces* sp. Amorph. yellow solid. $[\alpha]_D$ -35 (c, 0.49 in MeOH). λ_{max} 252 (€ 74100); 365 (€ 14200) (EtOH) (Derep).
 6-Deoxy- α -L-glucopyranos-3-yl ester: 3'-L-Quinoyosyl saphenate
 [137570-42-4]
 $C_{21}H_{22}N_2O_7$ 414.414
 Isol. from a *Streptomyces* sp. Amorph. yellow solid. $[\alpha]_D$ -40 (c, 0.73 in MeOH). λ_{max} 252 (€ 74100); 365 (€ 14200) (EtOH) (Derep).
 O-(Hydroxyacetyl): **Antibiotic DC 86M**. DC 86M
 [94448-15-4]
 $C_{17}H_{14}N_2O_5$ 326.308
 Prod. by *Streptomyces luteogriseus*. Active against gram-positive and -negative bacteria and murine sarcoma. Yellow needles. Sol. MeOH, C_6H_6 ; poorly sol. H_2O . Mp 185-187°. $[\alpha]_D^{20}$ -

43.8 (c, 0.5 in $CHCl_3$). λ_{max} 252 (€ 79000); 363 (€ 14000) (MeOH) (Derep).
 ▶ LD₅₀ (mus, ipr) 20 mg/kg.
 O-(Chloroacetyl): **Antibiotic DC 86C**. DC 86C
 $C_{17}H_{13}ClN_2O_4$ 344.753
 Semisynth., prod. from DC 86Y. Shows strong antibacterial activity. Yellow solid. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 154-156°. $[\alpha]_D^{25}$ -54 (c, 0.6 in $CHCl_3$). λ_{max} 254 ; 364 (MeOH) (Berdy).
 ▶ LD₅₀ (mus, ipr) 40 mg/kg.
 O-(tert-Butoxyacetyl): **Antibiotic DC 86R**. DC 86R
 $C_{21}H_{22}N_2O_5$ 382.415
 Semisynthetic. Biol. inactive. Yellow needles. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 189-192°. $[\alpha]_D^{25}$ -34.6 (c, 0.5 in $CHCl_3$). λ_{max} 252 ; 364 (MeOH) (Berdy).
 ▶ LD₅₀ (mus, ipr) 250 mg/kg.
 O-Tetradecanoyl: [120482-16-8]
 $C_{29}H_{38}N_2O_4$ 478.63
 From *Streptomyces antibioticus*.
 O-(12-Methyltridecanoyl): [120464-89-3]
 $C_{29}H_{38}N_2O_4$ 478.63
 From *Streptomyces antibioticus*.
 O-(12-Methyltetradecanoyl): [120464-90-6]
 $C_{30}H_{40}N_2O_4$ 492.657
 From *Streptomyces antibioticus*.
 O-(14-Methylpentadecanoyl): [120464-91-7]
 $C_{31}H_{42}N_2O_4$ 506.684
 From *Streptomyces antibioticus*.
 O-Hexadecanoyl: [120464-92-8]
 $C_{31}H_{42}N_2O_4$ 506.684
 From *Streptomyces antibioticus*.
 O-(14-Methylhexadecanoyl):
 $C_{32}H_{44}N_2O_4$ 520.711
 From *Streptomyces antibioticus*.
 O-(16-Methylheptadecanoyl): [120464-94-0]
 $C_{33}H_{46}N_2O_4$ 534.737
 From *Streptomyces antibioticus*.
 O-Benzoyl: O-Benzoylsaphenic acid. **Shisen 1**
 [221300-00-1]
 $C_{22}H_{16}N_2O_4$ 372.379
 Prod. by *Streptoplanospora viridis*. Active against MRSA. Yellow powder. λ_{max} 254 ; 364 (no solvent reported).
 O-(2-Hydroxy-6-methylbenzoyl): **Saphenamycin**. A 32256. Antibiotic A 32256
 [83198-27-0]
 $C_{23}H_{18}N_2O_5$ 402.406
 From *Streptomyces canarius* and other *Streptomyces* spp. Active against gram-positive and -negative bacteria. Weak antitumour props. Mosquito larvicide. Yellow prisms ($CHCl_3$ /hexane). Mp 200-202°. λ_{max} 254 (€ 95000); 300 (€ 15000); 348 (€ 17000); 364 (€ 22000) (EtOH/NaOH) (Derep). λ_{max} 207 (€ 51000); 255 (€ 86700); 365 (€ 13700) (EtOH) (Derep).
 ▶ SG1576540
Me ether: 6-(1-Methoxyethyl)-1-phenazinecarboxylic acid. **Saphenic acid methyl ether**


[120464-87-1]
 $C_{16}H_{14}N_2O_3$ 282.298
 From *Streptomyces antibioticus*. Yellow needles (Me_2CO /hexane). Mp 158-159° dec. $[\alpha]_D^{24}$ +9.1 (c, 0.54 in $CHCl_3$).
Me ether, Me ester: [73634-71-6]
 $C_{17}H_{16}N_2O_3$ 296.325
 From *Streptomyces* sp. Phosphodiesterase inhibitor.
 I'-Ketone: 6-Acetyl-1-phenazinecarboxylic acid
 [120464-88-2]
 $C_{15}H_{10}N_2O_3$ 266.256
 From *Streptomyces antibioticus*. Yellow needles (Me_2CO). Mp 219-221°. Turns brown in light.
 [137570-45-7, 137570-44-6]
Japan. Pat., 1980, 80 3 737; *CA*, **92**, 213541 (isol, struct, props)
 Kitahara, M. et al., *J. Antibiot.*, 1982, **35**, 1412 (*Saphenamycin*)
 Asano, K. et al., *J. Antibiot.*, 1986, **39**, 619 (isol)
 Takahashi, K. et al., *J. Antibiot.*, 1986, **39**, 624 (struct, uv, pmr, ms, props)
 Geiger, A. et al., *J. Antibiot.*, 1988, **41**, 1542 (derivs)
 Pathirana, C. et al., *J.O.C.*, 1992, **57**, 740 (glycosyl ester)
 Van't Land, C.W. et al., *J.O.C.*, 1993, **58**, 6576 (*Saphenamycin*, biosynth)
 Yue, H. et al., *CA*, 1999, **130**, 234411q (*Shisen I*)
 Petersen, L. et al., *Synthesis*, 1999, 1763-1766 (synth, cmr)
 Laursen, J.B. et al., *Bioorg. Med. Chem.*, 2003, **11**, 723-731 (*Saphenamycin*, synth)

3-(2-Hydroxyethyl)-6-prenylindole H-497


6-(3-Methyl-2-butenyl)-1H-indole-3-ethanol, 9CI. 3-(2-Hydroxyethyl)-6-(3-methyl-2-butenyl)indole. 6-Prenyltryptophol
 [583060-24-6]




$C_{15}H_{19}NO$ 229.321
 Alkaloid from a marine *Streptomyces* sp. (BL-49-58-005). Cytotoxic. λ_{max} 226 ; 282 (MeOH).
Carboxylic acid, nitrile: 6-(3-Methyl-2-butenyl)-1H-indole-3-acetonitrile. 3-Cyanomethyl-6-prenylindole
 [583060-26-8]
 $C_{15}H_{16}N_2$ 224.305
 Prod. by a marine *Streptomyces* sp. (BL-49-58-005). λ_{max} 226 ; 278 (MeOH).
Aldehyde, oxime: [583060-25-7]
 $C_{15}H_{18}N_2O$ 242.32
 Prod. by a marine *Streptomyces* sp. (BL-49-58-005). Cytotoxic. λ_{max} 226 ; 280 (MeOH).
 Sánchez López, J.M. et al., *J. Nat. Prod.*, 2003, **66**, 863-864 (isol, pmr, cmr)

5-(1-Hydroxyethyl)-3-pyridinecarboxylic acid, 9CI H-498C₈H₉NO₃ 167.164*Me ester: 3-(1-Hydroxyethyl)-5-methoxycarbonylpyridine*
[38940-64-6]C₉H₁₁NO₃ 181.191Alkaloid from *Nauclea diderrichii* (Rubiaceae), *Isertia haenkeana*, *Strychnos cocculoides* and *Strychnos spinosa*. Cryst. (Me₂CO). Mp 53-56° (52°). [α]_D²⁰ +23 (c, 2.5 in CHCl₃). Spec. rotn. refers to the *Isertia* isolate.*Me ether, Me ester: 3-(1-Methoxyethyl)-5-methoxycarbonylpyridine*


[38940-66-8]

C₁₀H₁₃NO₃ 195.218Alkaloid from *Nauclea diderrichii* (Rubiaceae). Cryst. (hexane). Mp 47-48°. [α]_D²⁶ +50 (c, 5 in MeOH).McLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**, 1478-1485 (*synth*, *ir*, *pmr*, *ms*)Bruix, M. *et al.*, *Phytochemistry*, 1993, **33**, 1257-1261 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)Din Belle, D. *et al.*, *Tetrahedron*, 1996, **52**, 11361-11378 (*synth*)**2-(1-Hydroxyethyl)-4(3H)-quinazolinone, 8CI** H-499*Chrysogine*
[18326-30-2]



(S)-form

C₁₀H₁₀N₂O₂ 190.201**(S)-form** [42599-89-3]Metab. of the fungi *Alternaria citri*, *Fusarium culmorum*, *Fusarium sambucinum* and *Penicillium chrysogenum*. Cryst. (EtOAc). Mp 190-192° (179-182°). [α]_D -25 (c, 0.39 in MeOH). [α]_D -41 (c, 2.5 in EtOH). λ_{max} 226 ; 304 (MeOH) (Berdy). λ_{max} 226 (ε 18620); 230 (ε 20892); 238 (ε 12882); 265 (ε 7244); 273 (ε 6660); 292 (ε 2884); 305 (ε 3890); 316 (ε 3235) (EtOH) (Berdy).*l'-Ketone: see 2-Acetyl-4(3H)-quinazolinone, A-75***(±)-form** [14422-59-4]Synthetic. Needles (MeOH/EtOAc or Me₂CO). Mp 190-191°.Uskokovic, M. *et al.*, *J.O.C.*, 1964, **29**, 582 (*synth*, *uv*, *ir*)Suter, P.J. *et al.*, *J.C.S. (C)*, 1967, 2240 (*isol*, *synth*)Hikino, H. *et al.*, *Yakugaku Zasshi*, 1973, **93**, 619; *CA*, **79**, 40922g (*isol*, *struct*)Blight, M.M. *et al.*, *J.C.S. Perkin 1*, 1974, 1691 (*isol*)Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 2347 (*synth*)Chadwick, D.J. *et al.*, *Acta Cryst. C*, 1983, **39**, 454 (*isol*, *cryst struct*)Bergman, J. *et al.*, *Tetrahedron*, 1990, **46**, 1295 (*synth*)Niederer, D. *et al.*, *Tet. Lett.*, 1992, **33**, 3997 (*isol*, *ir*, *pmr*, *cmr*, *abs config*)Maiti, D.K. *et al.*, *J. Chem. Res., Synop.*, 1996, 306 (*synth*)Bergman, J. *et al.*, *J. Chem. Res., Synop.*, 1997, 224 (*synth*)**3-(6-Hydroxyheptyl)-5-propylpyrrolizidine** H-500*Hexahydro-α-methyl-5-propyl-1H-pyrrolizine-3-hexanol, 9CI. Pyrrolizidine 267H'*C₁₇H₃₃NO 267.454**(3S,5R,6'ξ,8S)-form** [151805-21-9]Alkaloid from the Madagascan frog *Mantella* sp.*Ketone: 7-(Hexahydro-5-propyl-1H-pyrrolizine-3-yl)-2-heptanone, 9CI. 3-(6-Oxoheptyl)-5-propylpyrrolizidine. Pyrrolizidine 265H'*

[151805-20-8]

C₁₇H₃₁NO 265.438Alkaloid from *Mantella* sp.Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016-1038 (*isol*)Takahata, H. *et al.*, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1293-1295 (*synth*)**16-Hydroxy-4-hexadecenoic acid** H-501HOCH₂(CH₂)₁₀CH=CHCH₂CH₂COOHC₁₆H₃₀O₃ 270.411*Methylamide: 16-Hydroxy-N-methyl-4-hexadecenamide*C₁₇H₃₃NO₂ 283.453Alkaloid from *Michelia champaka*. Cryst. (MeOH). Mp 52°.Sharma, S. *et al.*, *Indian J. Chem., Sect. B*, 1998, **37**, 1219-1220**4-Hydroxy-2-hexenoic acid** H-502C₆H₁₀O₃ 130.143**(2E,4S)-form***2-Methylpropylamide: 4-Hydroxy-N-(2-methylpropyl)-2-hexenamide. 4-Hydroxy-N-isobutyl-2-hexenamide. Piptahsine*

[622405-50-9]

C₁₀H₁₉NO₂ 185.266Alkaloid from the dried seeds of *Piper nigrum*. Needles (petrol/EtOAc). Mp115-116°. Quoted as *S*-isomer in text but drawn as *R*-isomer. λ_{max} 213 (ε 6965) (MeOH).Siddiqui, B.S. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2760-2767 (*Piptahsine*)**Hydroxyhomodolabellin** H-503
HomohydroxydolabellinC₂₅H₃₄Cl₂N₂O₉S₂ 641.589Isol. from a *Lyngbya* sp. Cytotoxic. Amorph. solid. [α]_D²⁵ -10 (c, 0.05 in CHCl₃). CAS no. not found to CA 138. λ_{max} 203 (log ε 4.47); 238 (log ε 4.06) (MeOH).Luesch, H. *et al.*, *Tetrahedron*, 2002, **58**, 7959-7966 (*isol*, *pmr*, *cmr*)**5-Hydroxyhomolycorine** H-504
[13255-05-5]C₁₈H₂₁NO₅ 331.368Alkaloid from the bulbs of *Crinum defixum* (Amaryllidaceae). Cryst. (Me₂CO/MeOH). Mp 168-170°. λ_{max} 228 (ε 22100); 265 (ε 8030); 302 (ε 4070) (EtOH).*O⁹-De-Me, O⁵-Me: 5-Methoxy-9-O-demethylhomolycorine*

[167416-33-3]

C₁₈H₂₁NO₅ 331.368Alkaloid from whole plants of *Galanthus elwesii* (Amaryllidaceae). Prisms (MeOH/CHCl₃). Mp 232-236°. [α]_D²¹ +75.2 (c, 0.067 in MeOH). λ_{max} 200 (log ε 4.45); 227 (log ε 4.51); 265 (log ε 4.01); 305 (log ε 3.8) (MeOH).*O¹⁰-De-Me: 5-Hydroxy-10-O-demethylhomolycorine*

[119309-00-1]

C₁₇H₁₉NO₅ 317.341Alkaloid from *Narcissus tortifolius* and *Narcissus dubius* (Amaryllidaceae). Cryst. (MeOH/CH₂Cl₂). Mp 272-275°. [α]_D +98.8 (c, 0.52 in EtOH).*O¹⁰-De-Me, O¹⁰-(3ξ-hydroxybutanoyl), O⁵-Ac: Dubiusine*

[119308-98-4]

C₂₃H₂₇NO₈ 445.468

Alkaloid from the aerial parts of *Narcissus dubius* (Amaryllidaceae). Cryst. + ½H₂O (CHCl₃). Mp 226-228°. λ_{max} 210 (log ε 4.15); 232 (log ε 4.24); 276 (log ε 3.8); 308 (log ε 3.66) (EtOH).

4β,12β-Epoxyde, O⁵-Me: Galwesine

[167568-93-6]

C₁₉H₂₃NO₆ 361.394

Alkaloid from whole plants of *Galanthus elwesii* (Amaryllidaceae). Yellow amorph. powder. [α]_D²¹ +13.4 (c, 0.3 in MeOH). λ_{max} 225 (log ε 4.34); 267 (log ε 3.85); 303 (log ε 3.64) (MeOH).

4β,12β-Epoxyde, O⁹-de-Me, O⁵-Me: 9-O-Demethylgalwesine

[167416-36-6]

C₁₈H₂₁NO₆ 347.367

From whole plants of *Galanthus elwesii* (Amaryllidaceae). Prisms. Mp 241-242°. [α]_D²¹ +25.7 (c, 0.14 in MeOH). λ_{max} 227 (log ε 4.42); 265 (log ε 3.92); 306 (log ε 3.72) (MeOH).

16-Hydroxy, 4β,12β-epoxyde, O⁵-Me: 16-Hydroxygalwesine

[167416-34-4]

C₁₉H₂₃NO₇ 377.393

From whole plants of *Galanthus elwesii* (Amaryllidaceae). Prisms (EtOAc). Mp 154-156°. [α]_D²¹ +1.8 (c, 0.22 in MeOH). λ_{max} 202 (log ε 4.36); 226 (log ε 4.46); 269 (log ε 3.99); 303 (log ε 3.76) (MeOH).

16-Hydroxy, 4β,12β-epoxyde, O⁹-de-Me, O⁵-Me: 16-Hydroxy-9-O-demethylgalwesine

[167416-35-5]

C₁₈H₂₁NO₇ 363.366

From whole plants of *Galanthus elwesii* (Amaryllidaceae). Amorph. powder. Mp 107-110°. [α]_D²¹ +5.2 (c, 0.155 in MeOH). [α]_D²¹ +23 (c, 0.48 in CHCl₃). λ_{max} 200 (log ε 4.27); 226 (log ε 4.31); 266 (log ε 3.81); 305 (log ε 3.6) (MeOH). λ_{max} 243 ; 276 ; 341 (MeOH/NaOH).


Jeffs, P.W. *et al.*, *J.O.C.*, 1985, **50**, 1732-1737 (5-Hydroxyhomolycorine, *isol, uv, cd, pmr*)

Bastida, J. *et al.*, *Phytochemistry*, 1988, **27**, 3657-3660; 1990, **29**, 2683-2684 (*Dubiusine, 5-Hydroxy-10-O-demethylhomolycorine*)

Latvala, A. *et al.*, *Phytochemistry*, 1995, **39**, 1229-1240 (*Galwesine, 9-O-Methylgalwesine, 16-Hydroxygalwesine, 16-Hydroxy-9-O-demethylgalwesine, 5-Methoxy-9-O-demethylhomolycorine, cryst struct*)

5-Hydroxy-3-(2-hydroxyethyl)-1H-indole H-505

5-Hydroxy-1H-indole-3-ethanol, 9CI. 5-Hydroxytryptophol [154-02-9]

C₁₀H₁₁NO₂ 177.202

Serotonin metab. in rat. Present in bovine pineal tissue and *Bufo alvarius*. Isol. from the sponge *Hyrtios erectus*. Prisms. Mp 105-107°. λ_{max} 210 (ε 14400); 276 (ε 4940); 300 (sh) (ε 3450) (MeOH).

► Exp. reprod. effects. NL8512500

Picrate: Mp 150-152°.

5-O-β-D-Glucopyranoside:

C₁₆H₂₁NO₇ 339.344

Constit. of the stem bark of *Tetractron sinense*. Cytotoxic. Pale brown oil. λ_{max} 224 ; 278 ; 296 (MeOH).

5-Me ether: 3-(2-Hydroxyethyl)-5-methoxyindole. 5-Methoxytryptophol [712-09-4]

C₁₁H₁₃NO₂ 191.229

Present in bovine pineal tissue and *Bufo alvarius*.

► Human and exp. reprod. effects. NL8513000

Kveder, S. *et al.*, *Biochem. J.*, 1962, **85**, 447 (*synth*)

McIsaac, W.M. *et al.*, *Science (Washington, D.C.)*, 1965, **148**, 102


Salmoun, M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1173-1176 (*isol, pmr, cmr*)

Wang, Y.-F. *et al.*, *Chem. Biodiversity*, 2006, **3**, 1023-1030 (5-glucoside)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MFT300

2-Hydroxy-3-(5-hydroxy-1H-indol-3-yl)propanoic acid H-506

α,5-Dihydroxy-1H-indole-3-propanoic acid, 9CI. 3-(5-Hydroxy-3-indolyl)lactic acid. *Hyrtioerectine C*

C₁₁H₁₁NO₄ 221.212


(S)-form [151764-92-0]

Alkaloid from the sponge *Hyrtios erectus*. Amorph. solid. [α]_D²⁵ -31.3 (c, 0.13 in MeOH). λ_{max} 229 (log ε 3.76); 276 (log ε 3.56); 301 (log ε 3.44) (MeOH).

Youssef, D.T.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1416-1419 (*isol, pmr, cmr*)

5-Hydroxy-4-(4-hydroxy-3-methoxyphenyl)-4-(2-imidazolyl)-1,2,3-trithiane H-507

4-(4-Hydroxy-3-methoxyphenyl)-4-(1H-imidazol-2-yl)-1,2,3-trithian-5-ol, 9CI



Relative Configuration

C₁₃H₁₄N₂O₃S₃ 342.463

(+)-form [383191-04-6]

Isol. from the ascidian *Aplidium* sp. D. Cytotoxic. Yellow gum. [α]_D²⁰ +26 (c, 0.1 in MeOH). λ_{max} 216 (ε 13750); 255 (ε 8100); 290 (ε 5800) (MeOH/KOH) (Derep). λ_{max} 283 (ε 3900) (MeOH) (Derep).

(-)-form [383191-01-3]

Isol. from the ascidian *Hypsistozoa fasmariana*. Cytotoxic. [α]_D²⁰ -26 (c, 0.1 in MeOH).


[123060-46-8, 123060-45-7]

Copp, B.R. *et al.*, *Tet. Lett.*, 1989, **30**, 3703-3706 (*isol, struct*)

Pearce, A.N. *et al.*, *J.O.C.*, 2001, **66**, 8257-8259 (*isol, cd, pmr, cmr*)

4-Hydroxy-7-[1-hydroxy-2-(methylamino)ethyl]-2(3H)-benzothiazolone S 1319


[220752-37-4]

C₁₀H₁₂N₂O₃S 240.282

Isol. from the marine sponge *Dysidea* sp. β₂-Adrenoceptor agonist.

Suzuki, H. *et al.*, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 1361-1364 (*isol, pmr, cmr, activity*)

Fairhurst, R.A. *et al.*, *Org. Lett.*, 2005, **7**, 4697-4700 (*synth*)

4-Hydroxy-2-(hydroxymethyl)-2-butenic acid, 9CI H-509C₅H₈O₄ 132.116

(E)-form

5-O-E-Cinnamoyl, Me ester: [54118-68-2]

C₁₅H₁₆O₅ 276.288

Constit. of *Enkianthus perulatus*. Oil. Sol. MeOH, Et₂O; poorly sol. H₂O. λ_{max} 217 (ε 27800); 222 (ε 22800); 278 (ε 22900) (EtOH).

(Z)-form [111833-91-1]

Nitrile, 4-O-β-D-glucopyranoside: 4-(β-D-Glucopyranosyloxy)-2-(hydroxymethyl)-2-butenenitrile, 9CI.

Sarmentosin†. *Nigrum*

[71933-54-5]

C₁₁H₁₇NO₇ 275.258

Isol. from *Ribes nigrum* (blackcurrant), *Sedum sarmentosum* and from the moth *Abroxas grossulariata*. Antithe-

patotoxic compd. $[\alpha]_D^{24}$ -18 (c, 0.3 in MeOH). Has (*E*)-config.

5-O-(4-Hydroxybenzoyl), nitrile, 4-O- β -D-glucopyranoside: [930275-35-7]
C₁₈H₂₁NO₉ 395.365

Constit. of redcurrants (*Ribes rubrum*). Amorph. powder. Has (*E*)-config. (change of Cahn-Ingold priorities). λ_{\max} 211 ; 255 (MeCN).

5-O-(4-Hydroxy-3-methoxybenzoyl), nitrile, 4-O- β -D-glucopyranoside: [930275-36-8]
C₁₉H₂₃NO₁₀ 425.391

Constit. of redcurrants (*Ribes rubrum*). Amorph. powder. Has (*E*)-config. (change of Cahn-Ingold priorities). λ_{\max} 211 ; 263 (MeCN).

5-O-(3,4,5-Trihydroxybenzoyl), nitrile, 4-O- β -D-glucopyranoside: **Rhodiocyanoside B**
[168433-87-2]

C₁₈H₂₁NO₁₁ 427.364
Constit. of *Rhodiola quadrifida*. Powder. $[\alpha]_D$ -12.2 (MeOH). Has (*E*)-config. (change of Cahn-Ingold priorities). λ_{\max} 217 (log ϵ 4.48); 279 (log ϵ 3.99) (MeOH).

5-O-(4-Hydroxy-E-cinnamoyl), nitrile, 4-O- β -D-glucopyranoside: **5-O-p-Coumaroylnigrumin**
C₂₀H₂₃NO₉ 421.403

Constit. of the seeds of *Ribes nigrum* (blackcurrant). Powder. Has (*E*)-config. (change of Cahn-Ingold priorities).

5-O-(4-Hydroxy-3-methoxy-E-cinnamoyl), nitrile, 4-O- β -D-glucopyranoside: **5-O-Feruloylnigrumin**
C₂₁H₂₅NO₁₀ 451.429

Constit. of the seeds of *Ribes nigrum* (blackcurrant). Has (*E*)-config. (change of Cahn-Ingold priorities).

2,3-Epoxyde, nitrile, 4-O- β -D-glucopyranoside: **Epoxy-sarmentosin**. Sarmentosin epoxyde
[81907-02-0]

C₁₁H₁₇NO₈ 291.257
Isol. from *Sedum cepaea*. Hygroscopic powder.

[77459-99-5 ; 111833-90-0]

Kobayashi, A. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 1997-2000 (5-cinnamoyl Me ester)

Gu, Y.-X. *et al.*, *CA*, 1981, **94**, 18382j (*cryst struct*, Sarmentosin)

Nahrstedt, A. *et al.*, *Phytochemistry*, 1982, **21**, 107 (epoxyde)

Alen, R. *et al.*, *Acta Chem. Scand., Ser. B*, 1987, **41**, 76 (*synth*, ms)

Chu, G. *et al.*, *Bioorg. Med. Chem. Lett.*, 1993, **3**, 1343 (*synth*, Sarmentosin)

Nishida, R. *et al.*, *Phytochemistry*, 1994, **36**, 37 (*isol*)

Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1245-1247 (*Rhodiocyanoside B*)

Lu, Y. *et al.*, *Phytochemistry*, 2002, **59**, 465-468 (*Nigrumins*)

Xu., R. *et al.*, *Synth. Commun.*, 2003, **33**, 3355-3363 (*synth*, Sarmentosin)

Schwarz, B. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 1405-1410 (*redcurrant constits*)

4-Hydroxy-3-hydroxymethyl-2-butenic acid

H-510

(HOCH₂)₂C=CHCOOH

C₅H₈O₄ 132.116

Nitrile: 4-Hydroxy-3-hydroxymethyl-2-butenenitrile. 3-Cyano-2-hydroxymethyl-2-propen-1-ol. 2-(Cyanomethylene)-1,3-propanediol
[29768-68-1]

C₅H₇NO₂ 113.116
Constit. of *Cardiospermum halicacabum*, *Heliotropium* spp. and other Sapindaceae. Component of Jia Ku Gua.

Nitrile(*E*-), O- β -D-glucopyranoside: [84976-19-2]
C₁₁H₁₇NO₇ 275.258
Constit. of the secretions of *Leptocoris isolata*.

Nitrile(*Z*-), O- β -D-glucopyranoside: 4-(β -D-Glucopyranosyloxy)-3-(hydroxymethyl)-2-butenenitrile, 9CI. **Sutherlandin**
[110115-57-6]

C₁₁H₁₇NO₇ 275.258
Isol. from leaves of *Acacia sutherlandii*. Powder.

Nitrile, O'-octadecanoyl, O'-(14Z-eicosenoyl):
C₄₃H₇₇NO₄ 672.086

Constit. of the seeds of *Koelreuteria paniculata*.

Nitrile(*E*-), O-(4-hydroxy-E-cinnamoyl), O- β -D-glucopyranoside: **Sutherlandin trans-p-coumarate**
C₂₀H₂₃NO₉ 421.403

Constit. of *Sorbaria sorbifolia* var. *stellipila*. Needles. Mp 88-89°. $[\alpha]_D^{18}$ +27.8 (c, 0.09 in MeOH). λ_{\max} 211 ; 315 (MeOH).

Nitrile, 2S,3R-epoxyde, O- β -D-glucopyranoside: **Sutherlandin epoxyde**
[173792-47-7]

C₁₁H₁₇NO₈ 291.257
Constit. of *Exochorda giraldii*, *Exochorda serratifolia* and *Rhodotypos scandens*.

Braekman, J.C. *et al.*, *Biochem. Syst. Ecol.*, 1982, **10**, 355-364 (*isol*)

Swenson, W.K. *et al.*, *Phytochemistry*, 1987, **26**, 1835-1836 (*Sutherlandin*)

Lechtenberg, M. *et al.*, *Phytochemistry*, 1994, **37**, 1039-1043 (*isol*, *pmr*, *cmr*)

Lechtenberg, M. *et al.*, *Phytochemistry*, 1996, **41**, 779-785 (*Sutherlandin epoxyde*)

Spitzer, V. *et al.*, *Phytochemistry*, 1996, **42**, 1357-1360


Akita, H. *et al.*, *Tetrahedron: Asymmetry*, 1999, **10**, 2429-2439 (*Sutherlandin*, *synth*)

Kim, D.K. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1766-1767 (*Sutherlandin trans-p-coumarate*)

Yang, X. *et al.*, *Yaoxue Xuebao*, 2000, **35**, 279-283 (*isol*, *nitrile*, *ester*)

6-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1H)-quinolinone

H-511



C₁₄H₁₇NO₃ 247.293

Alkaloid from the leaves of *Spathelia excelsa*. Yellow powder. λ_{\max} 240 ; 332 ; 346 (MeOH).

Da Paz Lima, M. *et al.*, *Phytochemistry*, 2005, **66**, 1560-1566 (*isol*, *pmr*, *cmr*)

7-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1H)-quinolinone

H-512


C₁₄H₁₇NO₃ 247.293

Alkaloid from the leaves of *Spathelia excelsa*. Yellow powder. λ_{\max} 240 ; 331 ; 347 (MeOH).

Da Paz Lima, M. *et al.*, *Phytochemistry*, 2005, **66**, 1560-1566 (*isol*, *pmr*, *cmr*)

7-Hydroxy-4-(5-hydroxy-methyl-2-furanyl)-2(1H)-quinolinone

H-513



C₁₄H₁₁NO₄ 257.245


Alkaloid from the whole plant of *Aquilegia ecalcarata*. Cytotoxic. Amorph. orange powder. Mp 278-280° dec. λ_{\max} 238 ; 262 ; 360 (MeOH).

Chen, S.-B. *et al.*, *Planta Med.*, 2002, **68**, 554-556 (*isol*, *pmr*, *cmr*, *ms*)

9-Hydroxy-6-hydroxymethyl-1-phenazinecarboxylic acid, 9CI

H-514

Antibiotic T 41348. T 41348
[60160-03-4]



C₁₄H₁₀N₂O₄ 270.244

Phenazine antibiotic. Isol. from *Streptomyces recifensis*. Active against gram-positive and -negative bacteria. Orange-red cryst. Sol. MeOH. λ_{\max} 267 (ϵ 42000); 368 (ϵ 8700) (MeOH) (Berdy).

9-Me ether: 6-Hydroxymethyl-9-methoxy-1-phenazinecarboxylic acid. **Griseoluteic acid**. B 4607B. Antibiotic B 4607B
[489-76-9]

C₁₅H₁₂N₂O₄ 284.271

From *Streptomyces griseoluteus* and also *Vibrio* and *Bacillus* spp. Degradation prod. of Griseoluteins. Active against HeLa cells and Ehrlich carcinoma. Orange cryst. Sol. H₂O, MeOH. λ_{\max} 263 ; 350 ; 365 (MeOH) (Berdy).

9-Me ether, Me ester:

Orange-yellow needles. Mp 189°.

9-*Me ether, 1'-O-(aminoacetyl): Pelagiomicin C*. 2088C. Antibiotic 2088C. B 4607A. Antibiotic B 4607A. O-Glycylgriseoluteic acid [173485-82-0] C₁₇H₁₅N₃O₅ 341.323
Prod. by the marine bacterium *Pelagiomonas variabilis* and by a *Vibrio* sp. Active against gram-positive and -negative bacteria. Cytotoxic agent. Red-orange solid. λ_{max} 210 (log ε 4.67); 267 (log ε 4.65); 369 (log ε 4.09) (EtOH).

9-*Me ether, 1'-O-(hydroxyacetyl): Griseolutein A*

[573-84-2]

C₁₇H₁₄N₂O₆ 342.307

Produced by *Streptomyces griseoluteus*. Shows antibiotic activity against gram-positive and -negative bacteria. Orange-yellow needles. Sol. bases; fairly sol. MeOH, EtOAc; poorly sol. C₆H₆, H₂O, hexane. Mp 193° dec. λ_{max} 265 (ε 67700); 362 (ε 11300) (MeOH) (Derep). λ_{max} 267; 368 (EtOH) (Berdy).

► LD₅₀ (mus, scu) 500 mg/kg. SG1576550

9-*Me ether, 1'-O-(2-aminopropanoyl): O-Alanylgriseoluteic acid*. B 4607C. Antibiotic B 4607C

C₁₈H₁₇N₃O₅ 355.349

Prod. by *Vibrio* sp. Yellow powder. Sol. H₂O, MeOH. [α]_D²⁰ -3.7 (MeOH). λ_{max} 263; 365 (MeOH) (Berdy).

9-*Me ether, 1'-O-(2-amino-3-methylbutanoyl) (S-): Pelagiomicin B*. 2088B. Antibiotic 2088B. O-Valylgriseoluteic acid

[173485-81-9]

C₂₀H₂₁N₃O₅ 383.403

Prod. by *Pelagiomonas variabilis*. Red-orange needles. Sol. MeOH, butanol, CHCl₃. λ_{max} 210 (log ε 4.7); 267 (log ε 4.66); 369 (log ε 4.1) (EtOH).

9-*Me ether, 1'-O-(2-amino-3-hydroxy-3-methylbutanoyl) (S-): Pelagiomicin A*. 2088A. Antibiotic 2088A. O-(3-Hydroxyvalyl)griseoluteic acid

[173485-80-8]

C₂₀H₂₁N₃O₆ 399.402

Prod. by *Pelagiomonas variabilis*. Antitumour agent. Red-orange needles. Mp 130° dec. [α]_D²⁰ +19.8 (c, 1 in CHCl₃). λ_{max} 209 (log ε 4.59); 265 (log ε 4.62); 368 (log ε 3.96) (EtOH).

5,10-Dihydro, 9-*Me ether, N⁵-(hydroxyacetyl): Griseolutein B*

C₁₇H₁₆N₂O₆ 344.323

Isol. from *Streptomyces griseoluteus*. Shows antibiotic activity similar to that of Griseolutein A but weaker. Yellow needles (EtOAc). Sol. Py, bases; fairly sol. dioxan, MeOH; poorly sol. butyl acetate, toluene, Et₂O. Mp 220° dec. λ_{max} 282 (ε 10200); 342 (ε 5850) (MeOH) (Derep).

► LD₅₀ (mus, ivn) 200 mg/kg, LD₅₀ (mus, scu) 400 mg/kg.

Nakamura, S. *et al.*, *J. Antibiot., Ser. A*, 1957, **10**, 265; 1959, **12**, 26; 55; 133 (*synth, props*)

Nakamura, S. *et al.*, *Chem. Pharm. Bull.*, 1958, **6**, 539; 543; 547 (*synth, props*)

Yagishita, K. *et al.*, *J. Antibiot., Ser. A*, 1960, **13**, 83 (*isol*)

Holliman, F.G. *et al.*, *Chem. Comm.*, 1970, 1423 (*synth, bibl*)

Challand, S.R. *et al.*, *Chem. Comm.*, 1970, 1423 (*Griseoluteins A-B, synth, ms, ir, pmr, struct*)

Ueda, K. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 1931 (*synth*)

Japan. Pat., 1976, 76 32 790; *CA*, **85**, 107461 (*isol, props*)

Japan. Pat., 1995, 95 330 741; *CA*, **124**, 315159s (*Antibiotic 2088A*)

Imamura, N. *et al.*, *CA*, 1996, **124**, 170245e (*Pelagiomicins*)


Japan. Pat., 1996, 96 217 760; *CA*, **125**, 273725m

Imamura, N. *et al.*, *J. Antibiot.*, 1997, **50**, 8-12 (*Pelagiomicins A-C*)

5-Hydroxy-2-(hydroxymethyl)pyridine

H-515

5-Hydroxy-2-pyridinemethanol, 9CI
[40222-77-3]



C₆H₇NO₂ 125.127

Acid dehydration product of 5-Amino-5-deoxygalactose, A-726. Alkaloid from *Codonopsis pilosula* and *Papaver nudicaule*. Cryst. (MeCN). Mp 122-124°. Subl. 0.05 12.5.

Picrate: Mp 182-183°.

N-Oxide: [73737-02-7]

C₆H₇NO₃ 141.126

Mp 183-184°.

1'-Ac: [89098-99-7]

C₈H₉NO₃ 167.164

Cryst. (C₆H₆). Mp 121-123°.

Di-Ac: [31181-78-9]

C₁₀H₁₁NO₄ 209.201

Liq. Bp_{0.8} 122-128°.

Di-Ac, N-oxide:

C₁₀H₁₁NO₅ 225.201

Cryst. (EtOAc). Mp 121-122°.

[74386-51-9]

Elming, N. *et al.*, *Acta Chem. Scand.*, 1956, **10**, 1603-1605 (*synth*)

Inouye, S. *et al.*, *Tetrahedron*, 1968, **20**, 2125-2144 (*synth*)

French, F.A. *et al.*, *J. Med. Chem.*, 1970, **13**, 1124-1130; 1974, **17**, 172

Loh, W. *et al.*, *J. Med. Chem.*, 1980, **23**, 631-634 (*N-oxide, synth*)

Hass, W. *et al.*, *Annalen*, 1982, 1615-1622 (*di-Ac*)

Deady, L.W. *et al.*, *Aust. J. Chem.*, 1983, **36**, 2565-2568 (*synth, pmr, Ac*)

Hachisu, M. *et al.*, *J. Pharmacobio-Dyn.*, 1983, **6**, 922-931 (*synth*)

Miyake, Y. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 661-666 (*formn*)

Wang, Z.T. *et al.*, *CA*, 1989, **111**, 12386d (*isol*)


Zhang, Y. *et al.*, *CA*, 1998, **128**, 306192c (*isol*)

Müller, C. *et al.*, *Tetrahedron*, 1998, **54**, 10703-10712 (*synth, pmr, cmr*)

4-Hydroxy-5-hydroxymethyl-2-pyrrolidinecarboxylic acid

H-516

4-Hydroxy-5-hydroxymethylproline



C₆H₁₁NO₄ 161.157

(2R,4R,5S)-form

Synthetic isomer. [α]_D²⁰ +16.7 (c, 0.42 in H₂O).

(2R,4S,5R)-form

[921201-49-2 (hydrochloride)]

[α]_D²⁵ +29.8 (c, 0.6 in 1M HCl) (hydrochloride).

(2S,4S,5R)-form

Bulgecinine

[95863-87-9]

Constit. of bulgecins, glycopeptides prod. by *Pseudomonas acidophila* and *Pseudomonas mesoacidophila*. [α]_D²⁰ -13.1 (c, 0.95 in H₂O).

Shinagawa, S. *et al.*, *Tetrahedron*, 1984, **40**, 3465

Wakamiya, T. *et al.*, *Tet. Lett.*, 1985, **26**, 4759 (*synth*)

Bashyal, B.P. *et al.*, *Tet. Lett.*, 1986, **27**, 3205 (*synth*)

Yuasa, Y. *et al.*, *Chem. Comm.*, 1994, 1383 (*synth, bibl*)

Maeda, M. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 962 (*synth, bibl*)

Panday, S.K. *et al.*, *Synth. Commun.*, 1997, **27**, 1373-1384 (*Bulgecinine, synth*)

Fehn, S. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 2001-2005 (*synth*)

Krasinski, A. *et al.*, *Tet. Lett.*, 2001, **42**, 2019-2021 (*synth*)

Chavan, S.P. *et al.*, *Tet. Lett.*, 2005, **46**, 439-441 (*Bulgecinine, synth*)

Trost, B.M. *et al.*, *Chem. Eur. J.*, 2006, **12**, 6607-6620 (*Bulgecinine, synth*)


Chandrasekhar, S. *et al.*, *Tetrahedron: Asymmetry*, 2006, **17**, 2864-2869 (*Bulgecinine, 2R,4S,5R-form, synth*)

Toumi, M. *et al.*, *Tet. Lett.*, 2008, **49**, 1175-1179 (*Bulgecinine, synth*)

1-Hydroxy-1-hydroxymethylpyrrolizidine

H-517

Hexahydro-1-hydroxy-1H-pyrrolizine-1-methanol, 9CI



C₈H₁₅NO₂ 157.212

(1R,8S)-form

Yellow oil.

(1S,8S)-form

Curassanecine


[82842-00-0]

Necine base from *Heliotropium curassavicum* (Boraginaceae).

[138752-57-5, 138752-56-4]

Subramanian, P.S. *et al.*, *Aust. J. Chem.*, 1980, **33**, 1357 (*isol, ms*)
 Mohanraj, S. *et al.*, *Phytochemistry*, 1982, **21**, 1775 (*struct*)
 Gramain, J.-C. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1062 (*synth, abs config, cryst struct*)
 Dieter, R.K. *et al.*, *Tet. Lett.*, 2002, **43**, 7725-7728 (*synth*)

2-Hydroxy-1-hydroxymethylpyrrolizidine H-518
Hexahydro-2-hydroxy-1H-pyrrolizine-1-methanol, 9CI



(1*S*,2*R*,7*aR*)-form

C₈H₁₅NO₂ 157.212

(1*S*,2*R*,7*aR*)-form
 (1*β*,2*β*,7*aβ*)-form. *Macronecine*

[21824-61-3]
 [21823-73-4 ((±)-form)]
 Obt. by hydrol. of Macrophylline. Long needles (Me₂CO). Mp 126-128° (129°). [α]_D²⁰ +49.3 (c, 1.6 in EtOH).

Hydrochloride:

Cryst. (EtOH), Mp 152-153°. [α]_D¹⁸ +49.4 (c, 1.37 in EtOH).

O^{1'}-Angeloyl: *Macrophylline*†
 [27841-97-0]

C₁₃H₂₁NO₃ 239.314
 Alkaloid from *Senecio macrophyllus* (Asteraceae). Mp 42-44°. [α]_D²⁰ +34.5 (c, 1.68 in EtOH). See also Macrophylline, M-21.

*O*²-(3-Methyl-2-butenoyl): *O*²-*Senecioid-macronecine*

C₁₃H₂₁NO₃ 239.314
 Alkaloid from *Senecio caudatus* (Asteraceae). Alkaloid not named in the lit.

O^{1'}-(3-Methyl-2-butenoyl): *O*⁹-*Senecioid-macronecine*
 C₁₃H₂₁NO₃ 239.314
 Alkaloid from *Senecio caudatus* (Asteraceae). Not named in the lit.

(1*R*,2*R*,7*aS*)-form [21823-66-5]
 Fine needles (Me₂CO). Mp 128-128.5°. [α]_D¹⁸ -42.1 (c, 1 in EtOH).

O^{1'}-Angeloyl, *N*-oxide: **2*α*-Hydroxy-9-angeloyloxy-(*-*)-trachelanthamidine *N*-oxide**

[119060-86-5]
 C₁₃H₂₁NO₄ 255.313
 Alkaloid from aerial parts of *Senecio deferens* (Asteraceae). The *N*-oxide of the enantiomer of Macrophylline.

(1*S*,2*R*,7*aS*)-form
 (1*β*,2*β*,7*aα*)-form. *Petasinecine*

[70494-70-1]
 Obt. by hydrol. of Petasinine. Mp 132-134°.

O^{1'}-Angeloyl: *Hectorine*
 C₁₃H₂₁NO₃ 239.314
 Alkaloid from *Brachyglottis hectori*. Glass.

*O*²-Angeloyl: *Petasinine*
 [70474-33-8]

C₁₃H₂₁NO₃ 239.314
 Alkaloid from *Petasites japonicus* (sweet coltsfoot) (Asteraceae). Amorph. powder. [α]_D²⁵ +16 (c, 2.5 in EtOH).

*O*²-Angeloyl, *O*^{1'}-[*α*-*L*-rhamnopyranosyl-(→4)-4-hydroxycinnamoyl]: *Petasinose*
 [70474-34-9]
 C₂₈H₃₇NO₉ 531.602
 Alkaloid from flower stalks of *Petasites japonicus* (sweet coltsfoot) (Asteraceae). Amorph. powder. [α]_D²⁵ -38 (c, 1.1 in EtOH).

(1*R*,2*S*,7*aS*)-form

Synthetic. Mp 124-127°. [α]_D -49.4 (c, 0.96 in EtOH).

Danilova, A.V. *et al.*, *Zh. Obshch. Khim.*, 1955, **25**, 831; *CA*, **50**, 2626 (*isol, struct, Macrophylline*)

Aasen, A.J. *et al.*, *J.O.C.*, 1969, **34**, 4143 (*struct, synth, pmr*)

Yamada, K. *et al.*, *Tet. Lett.*, 1978, 4543 (*isol, pmr, struct, Petasinine, Petasinose*)
 Rüeger, H. *et al.*, *Heterocycles*, 1983, **20**, 235 (*synth*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151 (*isol, ir, ms, struct, derivs*)
 Hirschmann, G.S. *et al.*, *Planta Med.*, 1988, **54**, 360 (*isol, pmr, ms, struct, oxide*)

Ito, H. *et al.*, *J.A.C.S.*, 1994, **116**, 5469 (*Macronecine, synth*)


Ma, D. *et al.*, *J.C.S. Perkin 1*, 1999, 1703-1707 (*synth*)

Sarkar, T.K. *et al.*, *Tetrahedron*, 2005, **61**, 1155-1165 (*Macronecine, synth*)

Bai, Y. *et al.*, *ARKIVOC*, 2006, iii, 34-42 (*Hectorine, Petasinine*)

2-Hydroxy-2-(3-hydroxyphenyl)acetic acid H-519

α,3-Dihydroxybenzeneacetic acid, 9CI. *m*-Hydroxymandelic acid, 8CI. *α*,3-Dihydroxyphenylacetic acid. 3-Hydroxyphenylglycollic acid. *α*,3-Dihydroxy-*α*-toluic acid
 [17119-15-2]



(*R*)-form

C₈H₈O₄ 168.149

(*R*)-form [17513-99-4]

Nitrile, 2-*O*-(6-*O*-acetyl-*β*-*D*-glucopyranoside): **6''-*O*-Acetylholocalin**

C₁₆H₁₉NO₈ 353.328
 Constit. of *Sambucus nigra* (elderberry). [α]_D -47 (c, 0.2 in EtOH). Possible artifact. λ_{max} 278 (EtOH).

(*S*)-form [12514-00-0]

Nitrile, 2-*O*-*β*-*D*-glucopyranoside: *Zierin*
 [645-02-3]
 C₁₄H₁₇NO₇ 311.291
 Isol. from above ground parts of *Zieria laevigata*. Needles (EtOAc/CHCl₃). Mp 156° (softens at 153°). [α]_D²⁰⁻³ -29.5.

Nitrile, 2-*O*-[*β*-*D*-xylopyranosyl-(1→6)-*β*-*D*-glucopyranoside]: *Zierinxylamide*
 [82083-98-5]

C₁₉H₂₅NO₁₁ 443.407
 Constit. of the fruit of *Xeranthemum cylindraceum*.

Nitrile, 2-*O*-[*β*-*D*-glucopyranosyl-(1→4)-3,4-dihydroxycinnamoyl-(→5)-*β*-*D*-apiofuranosyl-(1→4)-*β*-*D*-xylopyranosyl-(1→6)-*β*-*D*-glucopyranoside]: *Xeranthin*
 [129761-12-2]
 C₃₉H₄₉NO₂₃ 899.809
 Constit. of the fruit of *Xeranthemum cylindraceum*. Powder. [α]_D²⁰ -88.1 (c, 1.29 in MeOH).

[19988-46-6]

Finnemore, H. *et al.*, *J. Proc. R. Soc. N.S.W.*, 1936, **70**, 135 (*Zierin*)

Gmelin, R. *et al.*, *Phytochemistry*, 1973, **12**, 457-461 (*Holocalin*)

Schwarzmaier, U. *et al.*, *Chem. Ber.*, 1976, **109**, 3250-3251 (*pmr, Zierin*)

Moehrl, H. *et al.*, *Pharmazie*, 1980, **35**, 756-761 (*synth, Zierin*)

Huebel, W. *et al.*, *Planta Med.*, 1982, **44**, 178-180 (*Zierinxylamide*)

Schwind, P. *et al.*, *Phytochemistry*, 1990, **29**, 1903-1911 (*Xeranthin, Zierinxylamide*)

Greca, M.D. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 175-182 (*Holocalin, 6''-Acetylholocalin*)

2-Hydroxy-2-(4-hydroxyphenyl)acetic acid H-520

α,4-Dihydroxybenzeneacetic acid, 9CI. *p*-Hydroxymandelic acid, 8CI. *α*,4-Dihydroxyphenylacetic acid. 4-Hydroxyphenylglycollic acid. *α*,4-Dihydroxy-*α*-toluic acid. ***Pisolithin B***

[1198-84-1]
 C₈H₈O₄ 168.149

(*R*)-form [13244-78-5]

Metab. of the fungus *Pisolithus tinctorius*. Cryst. + ½H₂O (H₂O). Mp 102-103° (anhyd.). [α]_D -144.4 (H₂O). Readily isomerises during isoln. λ_{max} 228; 274 (HCl) (Berdy). λ_{max} 248 (NaOH) (Berdy).

Nitrile, 2-*O*-*β*-*D*-glucopyranoside: ***Taxiphyllin***. *Phyllanthin*†. *Phyllanthoside*†
 [21401-21-8]

Cyanogenic glucoside of *Taxus* spp., *Bambusa vulgaris*, *Bambusa guadua*, *Triglochin maritima*, *Stenocarpus sinuatus*, *Sorghum* sp. *Girgensohnia oppositiflora* and *Macadamia ternifolia*. Algicide, phytotoxic. Cryst. (EtOH/C₆H₆). Mp 168-169° Mp 176°. [α]_D²⁰ -66.7 (c, 0.37 in EtOH). Diastereoisomeric, not enantiomeric, with Dhurrin (see below). λ_{max} 216 (ε 17780); 222 (ε 15490); 277 (ε 21880) (MeOH) (De-rep).

(*S*)-form [13244-75-2]

Plates + 1H₂O (H₂O). Mp 114-115°. [α]_D +154.7 (c, 1.36 in H₂O).

Nitrile, 2-*O*-*β*-*D*-glucopyranoside: ***Dhurrin***
 [499-20-7]

Cyanogenic glucoside isol. from *Sorghum vulgare* (sorghum), *Sorghum halapense* and *Suckleya suckleyana*. Algicide, phytotoxin. Cryst. + 1H₂O (H₂O); cryst. + 1EtOH (EtOH). Sol. MeOH, H₂O. Mp 165°. λ_{max} 228 (ε 13000) (MeOH) (Berdy).

Nitrile, 2-*O*-[3,4,5-trihydroxybenzoyl-(→

2)- β -*D*-glucopyranoside]: **Glochidacuminoside D**

C₂₁H₂₁NO₁₁ 463.397

Constit. of the leaves of *Glochidion acuminatum*. Amorph. powder. $[\alpha]_D^{25}$ -72.8 (c, 0.67 in MeOH). λ_{\max} 223 (log ϵ 4.19); 275 (log ϵ 3.98) (MeOH).

Nitrile, 2-O-[β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranoside]: **Dhurrin 6'-glucoside**

C₂₀H₂₇NO₁₂ 473.433

Constit. of *Sorghum bicolor* (sorghum).

(ξ)-form

Nitrile, 4'-O- β -*D*-glucopyranoside: *p*-Hydroxymandelonitrile glucoside

C₁₄H₁₇NO₇ 311.291

Constit. of *Goodia latifolia*, *Nandina domestica* and *Thalictrum aquilegifolium*. Unstable. Config. of mandelonitrile residue undetd.

Nitrile, di-O- β -*D*-glucopyranoside:

C₂₀H₂₇NO₁₂ 473.433

Constit. of *Thalictrum aquilegifolium*.

Nitrile, 4'-O-[3,4-dihydroxycinnamoyl-(\rightarrow 4)- β -*D*-glucopyranoside]:

Nandinin[†]

[91919-94-7]

C₂₃H₂₃NO₁₀ 473.435

Constit. of *Nandina domestica*. Pale yellow powder. Mp 101-104° dec. λ_{\max} 329 (MeOH).

[70013-61-5, 78689-25-5, 59845-69-1, 68758-69-0, 82492-24-8, 82492-25-9, 69322-02-7, 69322-01-6, 79694-14-7, 82503-24-0]

Plouvier, V. et al., *C. R. Hebd. Seances Acad. Sci.*, 1964, **259**, 665 (*Taxiphyllin*)

Towers, G.H.N. et al., *Tetrahedron*, 1964, **20**, 71-77 (*Taxiphyllin*, isol, uv, ir, pmr)

Mao, C.H. et al., *Phytochemistry*, 1965, **4**, 297 (*Taxiphyllin*, struct)

Schwartzmaier, U. et al., *Chem. Ber.*, 1976, **109**, 3250; 3379 (*abs config*)

Olechno, J.D. et al., *Phytochemistry*, 1984, **23**, 1784-1785 (*Nandinin*)

Selmar, D. et al., *Phytochemistry*, 1996, **43**, 569 (*Dhurrin 6'-glycoside*)

Nielsen, J.S. et al., *Arch. Biochem. Biophys.*, 1999, **368**, 121-130 (*Dhurrin*, biosynth)

Calderón, A.I. et al., *Acta Cryst. C*, 2003, **59**, o174-o176 (*Taxiphyllin*, *cryst struct*)

Otsuka, H. et al., *Chem. Pharm. Bull.*, 2004, **52**, 591-596 (*Glochidacuminoside D*)


Seigler, D.S. et al., *Phytochemistry*, 2005, **66**, 1567-1580 (*Dhurrin*, *Taxiphyllin*)

6-Hydroxyhyoscyamine

H-521

6-Hydroxy-8-methyl-8-azabicyclo[3.2.1]oct-3-yl α -(hydroxymethyl)benzeneacetate, 9*CI*. Anisodamine. Alkaloid V[†]

[17659-49-3]



(2'S,3R,6R)-form

C₁₇H₂₃NO₄ 305.373

Anticholinergic, antispasmodic agent. Reduces effects of acute myocardial infarction in rabbits. Shows atropine-like activity. Log P 0.11 (calc).

(2'S,3R,6R)-form [126371-43-5]

Alkaloid from *Duboisia* sp. and *Hyoscyamus albus*. Needles (MeOH). Mp 67°. $[\alpha]_D^{25}$ +1.2 (c, 1.9 in Me₂CO).

(2'S,3S,6S)-form [55869-99-3]

Alkaloid from *Datura ferox*, *Duboisia* sp., *Przewalskia tangutica*, *Scopoliatanguticus* and *Physochlaina dubia*. Cryst. (C₆H₆). Mp 61-62°. $[\alpha]_D^{20}$ -13.5 (c, 1.9 in MeOH).

►CY1634520

Hydrobromide: Mp 162° (156-157°). $[\alpha]_D^{23}$ -6.8.

N-Oxide: **6-Hydroxyhyoscyamine N-oxide**

[54519-13-0]

C₁₇H₂₃NO₅ 321.372

Alkaloid from *Physochlaina alaiica* (Solanaceae). Cryst. (EtOH). Mp 105-106°. Config. not detd.

Di-Ac:

Cryst. (Et₂O). Mp 74.5-75.5°.

Stereoisomer: [855784-79-1]

C₁₇H₂₃NO₄ 305.373

Alkaloid from *Datura stramonium*.

Relative config. only determined.

Complex stereochem. Appears from CAS to be the 3-epimer of previous isolates but with undetermined side-chain config.

(\pm)-form

6-Hydroxyatropine

Alkaloid from *Physochlaina alaiica* (Solanaceae). Mp 68-69°.

[52646-91-0]

Fodor, G. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 91 (*synth*)

Romeike, A. et al., *Naturwissenschaften*, 1962, **49**, 281 (*isol*)

Mirzamatov, P.T. et al., *Khim. Prir. Soedin.*, 1972, 493; 1974, 416; 540 (*isol*, pmr, ms, oxide)

Hsiao, P.K. et al., *CA*, 1981, **95**, 138447u (*isol*)

Shi, J. et al., *CA*, 1984, **100**, 114755p

(*pharmacol*)

Ishimaru, K. et al., *Phytochemistry*, 1989, **28**, 3507-3509 (*isol*, pmr, cmr)

Doncheva, T. et al., *CA*, 2005, **143**, 74802f (*Datura stramonium* alkaloid)


Muñoz, M.A. et al., *J. Nat. Prod.*, 2006, **69**, 1335-1340 (*vcd*, *abs config*)

5-Hydroxy-2,4-imidazolidin-2-one

H-522

5-Hydroxyhydantoin

[29410-13-7]



C₃H₄N₂O₃ 116.076

(\pm)-form

Cryst. (AcOH). Mp 140-142°. Compd. descr. in early ref. had different props.

(ξ)-form

Isol. from the tunicate *Botryllus schlosseri*.

Biltz, H. et al., *Ber.*, 1921, **54**, 1802-1828 (*synth*)


Abblard, J. et al., *Bull. Soc. Chim. Fr.*, 1971, 942-946 (*synth*, pmr)

Usov, A.I. et al., *Russ. J. Bioorg. Chem. (Engl. Transl.)*, 2002, **28**, 147-151 (ξ -form, *isol*)

2-Hydroxy-3-(4-imidazolyl)propanoic acid

H-523

α -Hydroxy-1*H*-imidazole-4-propanoic acid, 9*CI*. Hydroxydesaminohistidine. 3-(4-Imidazolyl)lactic acid [876-19-7]



C₆H₈N₂O₃ 156.141

Intermed. in histidine biosynth. by *Escherichia coli*.

(*S*)-form [14403-45-3]

Urinary metab. of histidine. Cryst. + 1H₂O (H₂O). Mp 204°. $[\alpha]_D^{25}$ -43 (c, 2.8 in H₂O).

Me ester: [80941-71-5]

C₇H₁₀N₂O₃ 170.168

Cryst. (MeOH/Et₂O) (as hydrochloride). Mp 140-142° (hydrochloride). $[\alpha]_D^{25}$ -22 (c, 1.9 in MeOH) (hydrochloride).

(\pm)-form

Cryst. + 1H₂O. Mp 222°.

Et ester:

C₈H₁₂N₂O₃ 184.194

Cryst. (CHCl₃). Mp 118-119°.

[72710-85-1, 86093-33-6]

Pyman, F.L. et al., *J.C.S.*, 1911, **99**, 1386-1401 (*synth*)

Fargher, R.G. et al., *J.C.S.*, 1919, **115**, 1015-1020 (*Et ester*, *synth*)


Hedegaard, J. et al., *Biochem. Biophys. Res. Commun.*, 1966, **25**, 335-339 (*occur*)

Noordam, A. et al., *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1981, **100**, 441-446 (*synth*, *ord*)

Okabe, N. et al., *Acta Cryst. C*, 1999, **55**, 217-218 (*S-form*, *cryst struct*)

4-Hydroxy-1*H*-indole-3-acetic acid

H-524



C₁₀H₉NO₃ 191.186

Amide, O- β -*D*-glucopyranoside:

C₁₆H₂₀N₂O₇ 352.343

Alkaloid from the roots of *Capparis tenera*. Mp 257-260°. $[\alpha]_D^{25}$ -59.3 (c, 0.09 in MeOH aq.). λ_{\max} 203 (sh) (log ϵ 1.13); 221 (log ϵ 1.45); 272 (log ϵ 0.28); 290 (log ϵ 0.2) (MeOH aq.).

Nitrile: 4-Hydroxy-1*H*-indole-3-acetonitrile. 3-(Cyanomethyl)-4-hydroxyindole

[118573-52-7]

C₁₀H₈N₂O 172.186

Present in rapeseed. Identified by ms.

Nitrile, O-β-D-glucopyranoside: Cappari-
loside A

[229483-41-4]

C₁₆H₁₈N₂O₆ 334.328Constit. of the fruit of *Capparis spinosa* (caper). Amorph. $[\alpha]_D^{20}$ -58.8 (c, 0.4 in MeOH). λ_{\max} 267; 278; 289 (MeOH).**Nitrile, O-[β-D-glucopyranosyl-(1→6)-β-**
D-glucopyranoside]: Cappari-
loside B

[229483-42-5]

C₂₂H₂₈N₂O₁₁ 496.47Constit. of the fruit of *Capparis spinosa* (caper). Amorph. $[\alpha]_D^{20}$ -23.7 (c, 0.3 in MeOH). λ_{\max} 272; 279; 289 (MeOH).**Me ether, nitrile: 4-Methoxy-1H-indole-3-**
acetonitrile. 3-(Cyanomethyl)-4-meth-
oxyindole. Arvelexin

[4837-74-5]

C₁₁H₁₀N₂O 186.213Constit. of *Brassica pekinensis* infected with *Plasmodiophora brassicae*. Also from *Rorippa sylvestris* and *Thalaspia arvensis*. Phytoalexin. Prisms (CHCl₃/hexane). Mp 145-146°.Nomoto, M. *et al.*, *Agric. Biol. Chem.*, 1970, **34**, 1590-1592 (*isol*)Slominski, B.A. *et al.*, *J. Chromatogr.*, 1988, **454**, 285-291 (*occur*)Yamada, F. *et al.*, *Heterocycles*, 1993, **36**, 2783-2804 (*synth*)Calis, I. *et al.*, *Phytochemistry*, 1999, **50**, 1205-1208 (*Cappari-*
losides)Somei, M. *et al.*, *Heterocycles*, 2000, **53**, 1573-1578 (*Cappari-*
loside A, synth)Pedras, M.S.C. *et al.*, *Phytochemistry*, 2003, **64**, 949-956 (*Arvelexin*)Su, D.-M. *et al.*, *Chem. Biodiversity*, 2008, **4**, 2852-2862 (*amide glucoside*)**5-Hydroxy-1H-indole-3-**
acetic acid, 9CI

H-525

[54-16-0]

C₁₀H₉NO₃ 191.186

Present in brain and cerebrospinal fluid. Serotonin metabolite. Mp 165°.

▶ LD₅₀ (mus, ipr) 1125 mg/kg. NL3650000**N-Me, Me ether, nitrile:** [176688-98-5]C₁₂H₁₂N₂O 200.24

Cryst. Mp 105-106°.

Me ether: 5-Methoxy-1H-indole-3-acetic
acid

[3471-31-6]

C₁₁H₁₁NO₃ 205.213

Present in bovine pineal tissue. Mp 150°.

▶ LD₅₀ (mus, ipr) 98 mg/kg. Exp. neoplas-**Me ether, Me ester:** [23304-48-5]C₁₂H₁₃NO₃ 219.24Cryst. (C₆H₆/petrol). Mp 73-74°.**Me ether, Et ester:** [57000-49-4]C₁₃H₁₅NO₃ 233.266Cryst. (C₆H₆/petrol). Mp 97-98°.**Me ether, nitrile:** [2436-17-1]C₁₁H₁₀N₂O 186.213

Oil.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 670B; 670C (*ir*)*Aldrich Library of 13C and 1H FT NMR**Spectra*, 1992, **3**, 142C (*nmr*)Hoshino, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1936, **11**, 221; *CA*, **30**, 5982Asero, B. *et al.*, *Farmaco, Ed. Sci.*, 1956, **11**, 219; *CA*, **50**, 13870Lerner, A.B. *et al.*, *J. Biol. Chem.*, 1960, **235**, 1992McIsaac, W.M. *et al.*, *Science (Washington, D.C.)*, 1965, **148**, 102Olivieri, G. *et al.*, *Med. Sci. Res.*, 1990, **18**, 99 (*metab*)Tsotinis, A. *et al.*, *J. Med. Chem.*, 2006, **49**, 3509-3519 (*Me ether nitrile, N-Me Me ether nitrile*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, HLJ000; MES850**6-Hydroxy-1H-indole-3-**
acetic acid

H-526

[31031-05-7]

C₁₀H₉NO₃ 191.186

No phys. props. reported.

Amide: 6-Hydroxy-1H-indole-3-aceta-
amide

[192184-73-9]

C₁₀H₁₀N₂O₂ 190.201Alkaloid from the edible mushroom *Agrocybe cylindracea*. Lipid peroxidation inhibitor. Powder. λ_{\max} 224 (ε 9870); 274 (ε 1690); 295 (ε 2360) (MeOH).**L-Valine amide: N-(6-Hydroxy-1H-indol-**
3-ylacetyl)valine. 6-Hydroxyindole-3-
acetylvaline

[945774-95-8]

C₁₅H₁₈N₂O₄ 290.318Metab. of 1H-Indole-3-acetic acid, I-81 in *Arabidopsis* sp.**L-Phenylalanine amide: N-(6-Hydroxy-**
1H-indol-3-ylacetyl)phenylalanine. 6-
Hydroxyindole-3-acetylphenylalanine

[945774-96-9]

C₁₉H₁₈N₂O₄ 338.362Metab. of 1H-Indole-3-acetic acid, I-81 in *Arabidopsis* sp.**Nitrile, O-β-D-glucopyranoside:**C₁₆H₁₈N₂O₆ 334.328Alkaloid from *Isatis indigotica*. Mp 240-242°.**Me ether: 6-Methoxy-1H-indole-3-acetic**
acid

[103986-22-7]


C₁₁H₁₁NO₃ 205.213Cryst. (H₂O). Mp 163-164° (dec.).**Me ether, nitrile: 6-Methoxy-1H-indole-3-**
acetonitrile. 3-(Cyanomethyl)-6-meth-
oxyindole

[23084-35-7]

C₁₁H₁₀N₂O 186.213

Needles (EtOH). Mp 113-114°.

[116621-16-0]

Findlay, S.P. *et al.*, *J.O.C.*, 1948, **13**, 560 (*deriv, synth*)Toyota, M. *et al.*, *J.C.S. Perkin I*, 1992, 547-552 (*synth*)Shinada, T. *et al.*, *Tet. Lett.*, 1996, **37**, 7099-7102 (*synth*)Kim, W.-G. *et al.*, *J. Nat. Prod.*, 1997, **60**, 721 (*amide, isol, uv, ir, pmr, cmr, ms*)Li, B. *et al.*, *Yaoxue Xuebao*, 2000, **35**, 508-510 (*nitrile, glucoside*)Kai, K. *et al.*, *Phytochemistry*, 2007, **68**, 1651-1663 (*aminoacid amides*)**4-Hydroxy-1H-indole-3-car-**
boxaldehyde, 9CI H-527
3-Formyl-4-hydroxyindole
[81779-27-3]C₉H₇NO₂ 161.16**O-β-D-Glucopyranoside:**C₁₅H₁₇NO₇ 323.302Alkaloid from the roots of *Capparis tenera*. Amorph. solid. λ_{\max} 216 (log ε 2.46); 243 (log ε 2.43); 306 (log ε 1.1) (MeOH).**Me ether: 4-Methoxy-1H-indole-3-car-**
boxaldehyde

[90734-97-7]

C₁₀H₉NO₂ 175.187

Cryst. (MeOH). Mp 162-163°.

Somei, M. *et al.*, *Heterocycles*, 1986, **24**, 3065-3069 (*synth*)Pindur, U. *et al.*, *Monatsh. Chem.*, 1990, **121**, 77-80 (*Me ether, synth, pmr*)Yamada, F. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 92-99 (*synth*)Su, D.-M. *et al.*, *Chem. Biodiversity*, 2008, **4**, 2852-2862 (*glucoside*)**5-Hydroxy-1H-indole-3-car-**
boxaldehyde, 9CI H-528
3-Formyl-5-hydroxyindole

[3414-19-5]

C₉H₇NO₂ 161.16Alkaloid from the marine sponge *Hyrtios erecta*. Needles. Mp 220-221°. λ_{\max} 215 (ε 12500); 252 (ε 11300); 271 (ε 9000); 298 (ε 1500) (MeOH) (Berdy).**Me ether: 5-Methoxy-1H-indole-3-car-**
boxaldehyde

[10601-19-1]

C₁₀H₉NO₂ 175.187Cryst. (CH₂Cl₂/petrol). Mp 174-175°.**Me ether, N-Me: 5-Methoxy-1-methyl-**
1H-indole-3-carboxaldehyde

[39974-94-2]

C₁₁H₁₁NO₂ 189.213

Needles (EtOH). Mp 134°.

Reimann, E. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 823-826 (*Me ether N-Me, synth, ms, pmr, anal*)Kobayashi, J. *et al.*, *Tetrahedron*, 1990, **46**, 7699-7702 (*isol*)Moody, C.J. *et al.*, *J.C.S. Perkin I*, 1993, 2561 (*Me ether, synth, ir, pmr, cmr, ms*)Naylor, M.A. *et al.*, *J. Med. Chem.*, 1997, **40**, 2335-2346 (*Me ether N-Me, synth, pmr*)Sauleau, P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1676-1679 (*isol, pmr, cmr*)**6-Hydroxy-1H-indole-3-car-**
boxaldehyde H-529
3-Formyl-6-hydroxyindole

[192184-71-7]


C₉H₇NO₂ 161.16

Alkaloid from the edible mushroom

Agroclype cylindracea. Lipid peroxidation inhibitor. Powder. λ_{\max} 220 (ϵ 3950); 235 (ϵ 3500); 276 (ϵ 2730); 300 (ϵ 1820) (MeOH).

Kim, W.G. *et al.*, *J. Nat. Prod.*, 1997, **60**, 721-723 (*isol, uv, pmr, cmr, ms*)

5-Hydroxy-1*H*-indole-3-carboxylic acid, 9CI H-530
[3705-21-3]



$C_9H_7NO_3$ 177.159
Cryst. (C_6H_6 /EtOH). Mp 187°.

Me ester: [112332-96-4]
 $C_{10}H_9NO_3$ 191.186
Isol. from the sponge *Hyrtios erectus*. Yellow needles.

Et ester: [24370-69-2]
 $C_{11}H_{11}NO_3$ 205.213
Mp 185-186°.

Benzyl ether: [24370-73-8]
 $C_{16}H_{13}NO_3$ 267.284
Mp 203-204° dec.

Marchelli, R. *et al.*, *Can. J. Chem.*, 1969, **47**, 4375-4381; 1971, **49**, 1296-1300 (*synth, ms, uv*)
Nakatsuka, S. *et al.*, *Heterocycles*, 1987, **26**, 1471-1474 (*Me ester, synth, pmr*)
Sauléau, P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1676-1679 (*Me ester, isol, pmr, cmr*)

6-Hydroxy-1*H*-indole-3-carboxylic acid, 9CI H-531
[24370-78-3]

$C_9H_7NO_3$ 177.159
Cryst. (EtOH aq.). Mp 229°.

Me ester: [112332-97-5]
 $C_{10}H_9NO_3$ 191.186
Alkaloid from a *Spongosorites* sp. Oil.

Me ester, O-[α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]:
 $C_{22}H_{29}NO_{12}$ 499.471
Constit. of the roots of *Clematis manshurica*. Light brown gum. λ_{\max} 192 ; 214 ; 279 ; 322 (MeOH).

Et ester: [24370-70-5]
 $C_{11}H_{11}NO_3$ 205.213
Mp 174-176° dec.

Me ether, Me ester:
 $C_{11}H_{11}NO_3$ 205.213
Constit. of the roots of *Clematis manshurica*. Powder. Mp 163-165°. λ_{\max} 218 ; 228 ; 275 ; 322 (MeOH).


Me ether, Et ester: [88612-61-7]
 $C_{12}H_{13}NO_3$ 219.24
Mp 141-142°.

Et ether, Me ester:
 $C_{12}H_{13}NO_3$ 219.24
Constit. of the roots of *Clematis manshurica*. Amorph. powder. Mp 172-174°. λ_{\max} 218 ; 228 ; 275 ; 322 (MeOH).

Benzyl ether: [24370-74-9]
 $C_{16}H_{13}NO_3$ 267.284
Mp 185-187° dec.

Marchelli, R. *et al.*, *Can. J. Chem.*, 1969, **47**, 4375-4381; 1971, **49**, 1296-1300 (*synth, ms, uv*)
Kasahara, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 927-928 (*Me ether Et ester*)
Nakatsuka, S. *et al.*, *Heterocycles*, 1987, **26**, 1471-1474 (*Me ester, synth, pmr*)
Shi, S.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 73-78 (*Clematis manshurica* constits)
Bao, B. *et al.*, *Mar. Drugs*, 2007, **5**, 31-39 (*Me ester, isol, pmr, cmr, ms*)

4-Hydroxy-1*H*-indole-2,3-dione H-532
4-Hydroxyisatin



$C_8H_5NO_3$ 163.132
Orange prisms. Mp 260° dec.

*Me ether: 4-Methoxy-1*H*-indole-2,3-dione. 4-Methoxyisatin. Isalexin* [108937-87-7]
 $C_9H_7NO_3$ 177.159
Alkaloid from *Brassica napus* ssp. *rapifera* (rutabaga) and *Brassica oleracea* var. *botrytis* (cauliflower). Phytoalexin. Mp 197-198°. λ_{\max} 199 (log ϵ 4.2); 234 (log ϵ 3.9); 335 (log ϵ 3.4) (no solvent reported).

Giovannini, E. *et al.*, *Helv. Chim. Acta*, 1957, **31**, 249-255 (*synth*)
Hewawasam, P. *et al.*, *Tet. Lett.*, 1994, **35**, 7303-7306 (*Isalexin, synth*)
Pedras, M.S.C. *et al.*, *J.O.C.*, 2004, **69**, 4471-4476 (*Isalexin*)

6-Hydroxy-1*H*-indole-2,3-dione, 9CI H-533

6-Hydroxyisatin
[116569-08-5]


$C_8H_5NO_3$ 163.132
Prod. by the marine-derived *Streptomyces* sp. B1848. Orange cryst. (MeOH). Dec. >325° without melting. λ_{\max} 263 ; 280 (sh) ; 349 (sh) ; 420 (sh) (MeOH).

*Me ether: 6-Methoxy-1*H*-indole-2,3-dione. 6-Methoxyisatin* [52351-75-4]
 $C_9H_7NO_3$ 177.159
Yellow needles (H_2O). Mp 233-235°.

*O,N-Di-Me: 6-Methoxy-1-methyl-1*H*-indole-2,3-dione. 6-Methoxy-N-methylisatin* [35162-27-7]
 $C_{10}H_9NO_3$ 191.186
Alkaloid from the aerial parts of *Boronella koniambiensis*. Bright orange needles (cyclohexane/EtOAc). Mp 196°.

Giovannini, E. *et al.*, *Helv. Chim. Acta*, 1948, **31**, 1381-1391 (*synth, Me ether*)
Pavlidis, V.H. *et al.*, *Synth. Commun.*, 1994, **24**, 533 (*synth*)
Shabaan, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (*isol, pmr, cmr, ms*)
Grougnet, R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1083-1086 (*6-Methoxy-N-methylisatin*)
Kaila, N. *et al.*, *J. Med. Chem.*, 2007, **50**, 21-39 (*Me ether*)

3-Hydroxy-1*H*-indole-2-thiol H-534
*1,3-Dihydro-3-hydroxy-2*H*-indole-2-thione. 3-Hydroxy-2-mercaptindole*



C_8H_7NOS 165.215

SH-form

O,S-Di- β -D-glucopyranoside: Calanthoside

$C_{20}H_{27}NO_{11}S$ 489.499
Alkaloid from *Calanthe discolor*. Powder. $[\alpha]_D^{25}$ -12 (c, 0.2 in MeOH). λ_{\max} 222 (log ϵ 3.5); 290 (log ϵ 2.7) (MeOH).

*S-Me: 3-Hydroxy-2-(methylthio)-1*H*-indole*
 C_9H_9NOS 179.242
Constit. of *Murex trunculus*. Biosynthetic precursor of Indigotin, I-72.

S-Me, O-sulfate:
 $C_9H_9NO_4S_2$ 259.306
Isol. from *Murex trunculus*.

*S-Me, S,S-dioxide: 3-Hydroxy-2-(methylsulfonyl)-1*H*-indole*
 $C_9H_9NO_3S$ 211.241
Constit. of *Purpura haemastoma*. Prob. biosynthetic precursor of Indigotin, I-72.

Baker, J.T. *et al.*, *Tet. Lett.*, 1968, 43 (*S-Me sulfate*)

Fouquet, H. *et al.*, *Angew. Chem., Int. Ed.*, 1971, **10**, 816-817 (*S-Me sulfate, isol*)


Baker, J.T. *et al.*, *Tet. Lett.*, 1976, 1233-1234 (*S-Me sulfate, occur*)

Murakami, T. *et al.*, *Heterocycles*, 2001, **54**, 957-966 (*Calanthoside*)

Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-769 (*S-Me sulfate, rev*)

2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-hexanone H-535

*3-(2-Hydroxy-4-methyl-3-oxohexyl)-1*H*-indole*
[79338-81-1]



$C_{15}H_{19}NO_2$ 245.321

Metab. from the bacterium *Xenorhabdus bovienii* A2. Possesses antifungal and antibacterial props. Oil. Sol. MeOH, $CHCl_3$; poorly sol. hexane. $[\alpha]_D^{25}$ +87 (c, 1.0 in $CHCl_3$).


*O-Ac: 3-(2-Acetoxy-4-methyl-3-oxohexyl)-1*H*-indole*

[79338-82-2]
 $C_{17}H_{21}NO_3$ 287.358
From *Xenorhabdus bovienii* A2. Possesses antifungal and antibacterial props. Oil. Sol. MeOH, $CHCl_3$; poorly sol. hexane. $[\alpha]_D^{25}$ +57 (c, 1.1 in $CHCl_3$). λ_{\max} 273 (ϵ 6560); 280 (ϵ 6990); 289 (ϵ 6200) (MeOH) (Berdy).

Li, J. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1081 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-pentanone H-536

3-(2-Hydroxy-4-methyl-3-oxopentyl)-1*H*-indole
[161889-68-5]



C₁₄H₁₇NO₂ 231.294

Metab. from the bacterium *Xenorhabdus bovienii* A2. Possesses antifungal and antibacterial props. Oil. $[\alpha]_D^{25}$ +86 (c, 1.1 in CHCl₃). λ_{\max} 273 ; 279 ; 289 (MeOH) (Berdy).

O-Ac: 3-(2-Acetoxy-4-methyl-3-oxopentyl)-1*H*-indole
[161889-67-4]


C₁₆H₁₉NO₃ 273.331

From *Xenorhabdus bovienii* A2. Possesses antifungal and antibacterial props. Oil. Sol. MeOH, CHCl₃. λ_{\max} 273 (€ 6700); 279 (€ 7050); 289 (€ 5900) (MeOH) (Berdy).

Li, J. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1081 (*isol*, *pmr*, *ms*, *struct*)

2-Hydroxy-1-(1*H*-indol-3-yl)-1,4-pentanedione, 9CI H-537

3-(2-Hydroxy-1,4-dioxopentyl)indole. 4-Hydroxy-5-(3-indolyl)-5-oxo-2-pentanone
[107749-00-8]



C₁₃H₁₃NO₃ 231.251


(±)-form

Alkaloid from the sponges *Dysidea etheria* and *Ulosa ruetzleri*. Possesses plant growth regulatory activity. Off-white solid. Sol. Me₂CO, MeOH.

Cardellina, J.H. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1065-1067 (*isol*, *ir*, *pmr*, *ms*, *struct*)

3-Hydroxy-4-(1*H*-indol-3-yl)-1-phenyl-2-butanone H-538

Kurasoin B
[183008-45-9]



C₁₈H₁₇NO₂ 279.338

(S)-form [182232-66-2]

Prod. by *Paecilomyces* sp. FO-3684.

Protein farnesyltransferase inhibitor. Powder. λ_{\max} 205 (sh) (€ 14790); 218 (€ 17020); 273 (€ 3490); 283 (€ 3630); 289 (€ 3210) (MeOH).

4''-Hydroxy: 3-Hydroxy-1-(4-hydroxyphenyl)-4-(1*H*-indol-3-yl)-2-butanone.

Soraphinol A

[1006373-58-5]

C₁₈H₁₇NO₃ 295.337

Prod. by *Sorangium cellulosum* JW1059. Oil. $[\alpha]_D^{25}$ +28.1 (c, 0.3 in MeOH). λ_{\max} 202 (log € 4.46); 221 (log € 4.53); 280 (log € 3.82) (MeOH).

Uchida, R. *et al.*, *J. Antibiot.*, 1996, **49**, 886;

932 (*isol*, *synth*, *uv*, *ir*, *pmr*, *cmr*)

Sunazuka, T. *et al.*, *J. Antibiot.*, 1997, **50**, 453 (*synth*, *abs config*)

Li, X.-M. *et al.*, *Bull. Korean Chem. Soc.*, 2007, **28**, 835-836 (*Soraphinol A*)

Tsuchiya, S. *et al.*, *Heterocycles*, 2007, **72**, 91-94 (*synth*)


Fernandes, R.A. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 15-18 (*synth*)

2-Hydroxy-3-(3-indolyl)propanoic acid H-539

α -Hydroxy-1*H*-indole-3-propanoic acid,

9CI. 3-(3-Indolyl)lactic acid

[1821-52-9]



C₁₁H₁₁NO₃ 205.213

(R)-form

D-form

[101312-07-6]

Isol. from the yeast *Malassezia furfur*. Mp 100-103°. $[\alpha]_D^{25}$ -6 (c, 0.28 in H₂O). λ_{\max} 221 (log € 4.56); 281 (log € 3.8); 290 (log € 3.74) (MeOH).

(S)-form

L-form

[7417-65-4]

Cryst. (Et₂O/petrol). Mp 100-101° (99°). $[\alpha]_D^{20}$ -5.36 (H₂O).

N-Me: α -Hydroxy-1-methyl-1*H*-indole-3-propanoic acid. 2-Hydroxy-3-(1-methylindolyl)propanoic acid

C₁₂H₁₃NO₃ 219.24

Constit. of the skins of peanuts (*Arachis hypogaea*). Needles (MeOH). Mp 216° dec. $[\alpha]_D^{25}$ +32.2 (c, 0.13 in MeOH).

Me ether: α -Methoxy-1*H*-indole-3-propanoic acid. 2-Methoxy-3-(3-indolyl)propanoic acid

C₁₂H₁₃NO₃ 219.24

Constit. of the skins of peanuts (*Arachis hypogaea*). Needles (MeOH). Mp 252° dec. $[\alpha]_D^{25}$ +22.2 (c, 0.1 in MeOH).

(±)-form [832-97-3]

Mp 144-145°.

(ξ)-form

Metab. of various bacteria *Agrobacterium* spp., *Acetobacter* spp., *Aspergillus* spp. and *Streptomyces* spp. incl. the marine *Streptomyces* sp. B2150.

Et ester, O-β-*D*-glucopyranoside: [143884-93-9]

C₁₉H₂₅NO₈ 395.408

Constit. of grapes (Riesling wine).

Me ether, dimethylamide: α -Methoxy-N,N-dimethyl-1*H*-indole-3-propanamide, 9CI. 3-(1*H*-Indol-3-yl)-2-methyl-N,N-dimethylpropanamide
[201284-05-1]

C₁₄H₁₈N₂O₂ 246.308

Alkaloid from the seeds of *Erythrina brucei*.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 669B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 140C (*nmr*)

Ehrlich, F. *et al.*, *Ber.*, 1911, **44**, 888

Bauguess, L.C. *et al.*, *J. Biol. Chem.*, 1934, **104**, 679

Ratusky, J. *et al.*, *Chem. Listy*, 1957, **51**, 109; *CA*, **51**, 13843 (*synth*)

Marinos, V.A. *et al.*, *Phytochemistry*, 1992, **31**, 2755-2759; 4399 (*isol*)


Bugge, J. *et al.*, *Pharm. Pharmacol. Lett.*, 1997, **7**, 155-156 (*Me ether dimethylamide*)

Lou, H. *et al.*, *Planta Med.*, 2001, **67**, 345-349 (*N-Me, Me ether*)

Shabaan, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (*marine, isol*)

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 1472-1485 (*isol*, *pmr*, *cmr*, *ms*)

19-Hydroxyingol H-540



C₂₀H₃₀O₇ 382.453

8-Tigloyl, 12,19-di-Ac:

C₂₉H₄₀O₁₀ 548.629

Constit. of *Euphorbia acruensis*. Oil. $[\alpha]_D$ +2 (c, 4 in CHCl₃).

8-Tigloyl, 3,12,19-tri-Ac:

C₃₁H₄₂O₁₁ 590.666

Constit. of *Euphorbia acruensis*. Oil. $[\alpha]_D$ -15 (c, 1 in CHCl₃).

7,8-Ditigloyl, 3,12-di-Ac:

C₃₄H₄₆O₁₁ 630.731

Constit. of *Euphorbia acruensis*. Oil. $[\alpha]_D$ -29 (c, 4.8 in CHCl₃).

8-Benzoyl, 3,7,12-tri-Ac:

C₃₃H₄₀O₁₁ 612.672

Constit. of *Euphorbia hermentiana*. Gum. $[\alpha]_D^{25}$ -45.1 (c, 0.2 in CHCl₃).

7-Phenylacetyl, 3,19-di-Ac:

C₃₂H₄₀O₁₀ 584.662

Constit. of *Euphorbia poisonii*. $[\alpha]_D^{23}$ +11 (c, 0.93 in CHCl₃). λ_{\max} 201 (€ 10230) (MeOH) (Berdy).

7-Phenylacetyl, 3,12,19-tri-Ac:


C₃₄H₄₂O₁₁ 626.699

Constit. of *Euphorbia poisonii*. Flakes. Mp 180-182°. $[\alpha]_D^{23}$ -4.3 (c, 1.03 in CHCl₃). λ_{\max} 201 (€ 5623); 212 (€ 6000) (MeOH) (Berdy).

8-(3-Pyridinecarbonyl), 7-(phenylacet-
tyl), 3,12,19-tri-Ac:
C₄₀H₄₅NO₁₂ 731.795
Constit. of *Euphorbia poissonii*. Cryst.
Mp 198-199°. [α]_D²³ -30 (c, 0.52 in
CHCl₃). λ _{max} 201 (ε 23442); 247 (ε
2570); 263 (ε 3020) (MeOH) (Berdy).
19-(3-Pyridinecarbonyl), 8-benzoyl,
3,7,12-tri-Ac: [225640-78-8]
C₃₉H₄₃NO₁₂ 717.768
Constit. of *Euphorbia marginata*.
Lin, L.-J. et al., *Phytochemistry*, 1983, **22**,
2795-2799 (benzoyl ester)
Branch, S.K. et al., *Magn. Reson. Chem.*, 1992,
30, 632 (isol, pmr, cmr)
Fatope, M.O. et al., *Bioorg. Med. Chem.*, 1996,
4, 1679 (isol, pmr, cmr)
Marco, J.A. et al., *Phytochemistry*, 1998, **49**,
1095-1099 (*Euphorbia acruens* esters)

20-Hydroxyisoaflavinine H-541

24,25-Didehydro-10,11-dihydro-20-hydro-
xyflavinone
[116865-09-9]



C₂₈H₃₉NO₂ 421.622
Metab. of *Aspergillus flavus*. Prisms
(MeOH). Mp 146-148° Mp 259-262°.
[α]_D²⁴ +56 (c, 0.1 in MeOH). [α]_D +0.9 (c,
0.34 in MeOH).


20-Epimer: **20-epi-Hydroxyisoaflavinine**
[124815-91-4]

C₂₈H₃₉NO₂ 421.622
From *Aspergillus tubingensis*. Mp 79-
82°. [α]_D -5.6 (c, 1.06 in CHCl₃).

Gloer, J.B. et al., *J.O.C.*, 1988, **53**, 5457 (isol)
Nozawa, K. et al., *Chem. Pharm. Bull.*, 1989,
37, 626 (isol, cryst struct)
TePaske, M.R. et al., *Tetrahedron*, 1989, **45**,
4961 (isol)

9-Hydroxyisoascididemnin H-542

9-Hydroxy-8H-benzo[b]pyrido[4,3,2-
de][1,10]phenanthrolin-8-one, 9CI
[215590-37-7]



C₁₈H₉N₃O₂ 299.288
Alkaloid from the sponge *Biemna fortis*.
Neuronal differentiation inducer.
Amorph. yellow solid. Mp 293-295° dec.
(as Me ether). Possible artifact. λ _{max} 227
(ε 15000); 277 (ε 10600); 365 (ε 5270)

(MeOH).


9-Deoxy, 9-amino: 9-Amino-8H-ben-
zo[b]pyrido[4,3,2-de][1,10]phenan-
throlin-8-one, 9CI. **9-
Aminoisoascididemnin**
[191849-14-6]

C₁₈H₁₀N₄O 298.303
Alkaloid from the sponge *Biemna
fortis*. Neuronal differentiation indu-
cer. Amorph. yellow solid (natural).
Red powder (CHCl₃/petrol) (syn-
thetic). Mp > 250° (synthetic). λ _{max}
231 (ε 20000); 282 (ε 14100); 373 (ε
7020) (MeOH).

Kitahara, Y. et al., *Tetrahedron*, 1998, **54**,
8421-8432 (synth, pmr, cmr)
De la Fuente, J.A. et al., *Bioorg. Med. Chem.*,
2001, **9**, 1807-1814 (synth, pmr, cmr)
Aoki, S. et al., *Bioorg. Med. Chem.*, 2003, **11**,
1969-1973 (isol, pmr, cmr)

1-Hydroxy-6-isopropyl-3-(1-methylpropyl)-2(1H)-pyrazinone H-543

1-Hydroxy-6-(1-methylethyl)-3-(1-
methylpropyl)-2(1H)-pyrazinone
[63538-07-8]




C₁₁H₁₈N₂O₂ 210.275
Metab. from *Aspergillus oryzae*. Oil.

Cu salt:
Grass-green cryst. (Me₂CO). Mp 205-
207°.

Ueno, T. et al., *Agric. Biol. Chem.*, 1977, **41**,
901 (isol, uv, pmr, ms, struct)

1-Hydroxy-6-isopropyl-3-[2-(methylthio)ethyl]-2(1H)-pyrazinone H-544

1-Hydroxy-6-(1-methylethyl)-3-[2-
(methylthio)ethyl]-2(1H)-pyrazinone
[37167-11-6]




C₁₀H₁₆N₂O₂S 228.315
Metab. from *Aspergillus flavus*. Mp 69-
75°.

MacDonald, J.C. et al., *Can. J. Biochem.*,
1972, **50**, 543 (isol, pmr, ms, struct)

2-Hydroxy-3-isopropylpyra- zine H-545

3-Isopropyl-2-pyrazinol. 2-Hydroxy-3-(1-
methylethyl)pyrazine, 9CI. 3-(1-Meth-
ylethyl)-2(1H)-pyrazinone



C₇H₁₀N₂O 138.169

OH-form

Me ether: 2-Isopropyl-3-methoxy-pyrazine
[25773-40-4]

C₈H₁₂N₂O 152.196
Occurs in petitgrain and galbanum oil.
Prod. by *Penicillium* spp., *Pseudomonas*
spp. and *Streptomyces* spp. Perfumery
and flavouring ingredient.

1'-Hydroxy, Me ether: 2-(1-Hydroxy-1-
methylethyl)-3-methoxy-pyrazine. 3-
Methoxy- α,α -dimethylpyrazinemetha-
nol, 9CI

[38346-80-4]
C₈H₁₂N₂O₂ 168.195
Prod. by *Chondromyces crocatus*. Oil.

Aldrich Library of FT-IR Spectra, 1st edn.,
1985, **2**, 842B (ir)

Aldrich Library of 13C and 1H FT NMR
Spectra, 1992, **3**, 404B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase,
1989, **3**, 1560C (ir)

Parliament, T.H. et al., *J. Agric. Food Chem.*,
1972, **20**, 682-684 (struct)

Bramwell, A.F. et al., *J.C.S. Perkin 1*, 1972,
2004-2007 (synth)

Bramwell, A.F. et al., *Tetrahedron*, 1973, **29**,
3939-3942 (pmr)

Calabretta, P.J. et al., *Cosmet. Perfum.*, 1975,
90, 74; 76; 79-80; *CA*, **83**, 183292c (synth,
props)

Murray, K.E. et al., *J. Sci. Food Agric.*, 1975,
26, 973-986; *CA*, **84**, 29266u (occur)

Gerber, N.N. et al., *Dev. Ind. Microbiol.*,
1978, **20**, 225-238; *CA*, **91**, 206932t (isol,
rev)

Karahadian, C. et al., *J. Agric. Food Chem.*,
1985, **33**, 339-343 (isol)

McInver, R.C. et al., *ACS Symp. Ser.*, 1986,
317, 266 (isol)

Maga, J.A. et al., *Food Res. Int.*, 1987, **3**, 269-
284 (occur, rev)

Arnoldi, A. et al., *Heterocycles*, 1988, **27**,
2875-2882 (ms)

Leete, E. et al., *Spec. Publ. - R. Soc. Chem.*,
1992, **95**, 75 (rev, biosynth)

Fenaroli's Handbook of Flavor Ingredients, 4th
edn., (ed. Burdock, G.A.), CRC Press, 2001,
1052-1053 (FEMA 3358)

Schulz, S. et al., *Tetrahedron*, 2004, **60**, 3863-
3872 (*Chondromyces crocatus* constiti)

2-Hydroxy-5-isopropylpyra- zine H-546

5-(1-Methylethyl)-2(1H)-pyrazinone. 5-
Isopropyl-2-pyrazinol
C₇H₁₀N₂O 138.169

OH-form

Me ether: 2-Isopropyl-5-methoxy-pyrazine
[68039-46-3, 56891-99-7]

C₈H₁₂N₂O 152.196
Present in various vegetable sources,
e.g. carrot, cucumber. Component of
FEMA 3358 together with its
regioisomers. See also 2-Hydroxy-3-
isopropylpyrazine, H-545. Not
characterised except as isomeric
mixture.

Stofberg, J. et al., *Perfum. Flavor.*, 1984, **9**, 53-
56; 58-59; 62; 65-72; 76-83 (use)
Fenaroli's Handbook of Flavor Ingredients, 4th
edn., (ed. Burdock, G.A.), CRC Press, 2001,
1052-1053 (use, occur)

2-Hydroxy-6-isopropylpyrazine H-547

6-(1-Methylethyl)-2(1H)-pyrazinone. 6-Isopropyl-2-pyrazinol
 $C_7H_{10}N_2O$ 138.169

OH-form


Me ether: 2-Isopropyl-6-methoxy-pyrazine
 $C_8H_{12}N_2O$ 152.196
 Present in various vegetable sources, e.g. carrot, cucumber. Component of FEMA 3358 together with its regioisomers. Not characterised except as isomeric mixt.

Stofberg, J. *et al.*, *Perfum. Flavor.*, 1984, **9**, 53-56; 58-59; 62; 65-72; 76-83 (*use*)

Fenaroli's Handbook of Flavor Ingredients, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 1052-1053 (*occur, use*)

5-Hydroxy-3-isothiocyanato-4-methoxyindole H-548

3-Isothiocyanato-4-methoxy-1H-indol-5-ol, 9CI. **Rapalexin B**
 [929083-73-8]




$C_{10}H_8N_2O_2S$ 220.251

Phytoalexin isol. from white rust infected *Brassica rapa*. No phys. props. reported.

Pedras, M.S.C. *et al.*, *Chem. Comm.*, 2007, 368-370 (*isol, synth, pmr, cmr*)

Hydroxylupanine H-549

7,10-Dihydro-8-(1-hydroxy-1-methylethyl)-10-methyl-1,3-dioxolo[4,5-h]furo[2,3-b]quinolin-6(8H)-one, 9CI. 3,9-Dihydro-2-(1-hydroxy-1-methylethyl)-9-methyl-7,8-methylenedioxyfuro[2,3-b]quinolin-4(2H)-one



$C_{16}H_{17}NO_5$ 303.314

(+)-form

Alkaloid from the leaves of *Ptelea trifoliata* (Rutaceae). Cryst. (MeOH). Mp 224-227°. $[\alpha]_D^{25} +6$ (c, 0.59 in EtOH).

(-)-form

Alkaloid from the leaves of *Lunasia amara* (Rutaceae). Cryst. (C_6H_6). Mp 228-230°. $[\alpha]_D^{25} -5.9$ (c, 0.276 in EtOH).

(±)-form


Synthetic.

Needles (MeOH/Et₂O). Mp 222-225°.

Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 6209 (*isol, uv, ir, struct*)

Reisch, J. *et al.*, *Tet. Lett.*, 1969, 3803 (*occur, struct*)

Neville, C.F. *et al.*, *J.C.S. Perkin 1*, 1991, 259 (*synth*)

3-Hydroxylupanine H-550

$C_{15}H_{24}N_2O_2$ 264.367

3α-form [129242-22-4]

Alkaloid from leaves of *Ammopiptanthus mongolicus* (Fabaceae) and from *Leontice leontopetalum* (Berberidaceae). Cryst. (toluene), needles (CHCl₃/petrol). Mp 143-145° (130-132°). $[\alpha]_D^{20} +73.8$ (c, 1 in CHCl₃). $[\alpha]_D +54.2$ (c, 0.15 in MeOH).

3β-form [129443-39-6]

Alkaloid from *Pearsonia* sp. (Fabaceae). Mp 94-95°. Incorr. descr. as 4β-Hydroxylupanine (see 4-Hydroxylupanine, H-551). Racemic.


Proksa, B. *et al.*, *Coll. Czech. Chem. Comm.*, 1990, **55**, 1257

Verdoorn, G.H. *et al.*, *Phytochemistry*, 1990, **29**, 1297

Al-Tel, T.H. *et al.*, *Phytochemistry*, 1991, **30**, 2393

4-Hydroxylupanine H-551

Dodecahydro-2-hydroxy-7,14-methano-4H,6H-dipyrido[1,2-a:1',2'-e]diazocin-4-one, 9CI. 4-Hydroxy-2-oxosparteine



$C_{15}H_{24}N_2O_2$ 264.367

4α-form**Chamaetin**

[31190-54-2]

Alkaloid from *Chamaecytisus frivaldskyanus*, *Chamaecytisus albus* ssp. *microphyllus*, *Chamaecytisus janae*, *Chamaecytisus supinus*, *Chamaecytisus calcareus* and *Chamaecytisus polytrichus* (Fabaceae). Cryst. Mp 207-209°. $[\alpha]_D^{25} +59.2$ (c, 0.38 in EtOH).

Perchlorate: [23360-88-5]

Mp 170-172°.

4β-form**Nuttalline†**

[23360-87-4]

Alkaloid from *Lupinus nuttallii* and *Lupinus hartwegii* (Fabaceae). Mp 108-109°. $[\alpha]_D^{26} +25.3$ (c, 2 in EtOH).

Ac. Nuttalline acetate. 4β-Acetoxy-lupanine

[68385-76-2]

$C_{17}H_{26}N_2O_3$ 306.404

Alkaloid detected in *Lupinus polyphyllus*, *Lupinus angustifolius*, *Lupinus mutabilis* and *Lupinus albus*.

4ξ-form [81149-32-8]

Alkaloid from *Lupinus mutabilis* seeds (Fabaceae).

O-Angeloyl: 4-(*Angeloyloxy*)lupanine [86632-28-2]

$C_{20}H_{30}N_2O_3$ 346.469

Minor alkaloid detected in *Lupinus mutabilis* seeds (Fabaceae).

Goldberg, S.I. *et al.*, *Chem. Comm.*, 1969, 660 (*isol, struct, uv, Nuttalline*)

Cho, Y.D. *et al.*, *Arch. Mass Spectral Data*, 1972, **2**, 732 (*ms*)


Anderson, J.N. *et al.*, *J.O.C.*, 1976, **41**, 3441 (*isol, Nuttalline*)

Daily, A. *et al.*, *Tet. Lett.*, 1978, 1453 (*isol, pmr, abs config, Chamaetin*)

Hatzold, T. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 934 (*isol, glc*)

Rastogi, R. *et al.*, *J. Indian Chem. Soc.*, 1984, **61**, 918 (*isol, Nuttalline*)

Mühlbauer, P. *et al.*, *Planta Med.*, 1988, **54**, 237 (*Nuttalline acetate*)

6-Hydroxylupanine H-552

$C_{15}H_{24}N_2O_2$ 264.367

6α-form [278617-91-7]

Alkaloid from *Lygos raetam* var. *sarcocarpa* (preferred genus name *Genista*) and *Maackia amurensis*. Cryst. Mp 116-117°. $[\alpha]_D^{23} -73.5$ (c, 0.17 in EtOH).

Me ether: 6α-Methoxylupanine

[278617-92-8]

$C_{16}H_{26}N_2O_2$ 278.394

Alkaloid from *Maackia amurensis*. Oil. $[\alpha]_D^{23} -16.5$ (c, 0.42 in EtOH).


6β-form [103847-11-6]

Alkaloid from *Bolusanthus speciosus*.

Asres, K. *et al.*, *Phytochemistry*, 1986, **25**, 1449-1452 (*isol, ir, pmr, ms, cd, struct*)

Abdel-Halim, O.B. *et al.*, *Phytochemistry*, 1995, **40**, 1323-1325 (*isol, pmr, cmr, ms*)

Wang, Y.-H. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 641-645 (*isol, pmr, cmr, ms*)

12-Hydroxylupanine H-553

$C_{15}H_{24}N_2O_2$ 264.367


12α-form [145204-92-8]

Alkaloid from aerial parts of *Lygos raetam* var. *sarcocarpa* (preferred genus name *Genista*) (Fabaceae). Cryst. Mp 173-175°. $[\alpha]_D^{24} +20$ (c, 0.15 in EtOH).

Abdel-Halim, O.B. *et al.*, *Phytochemistry*, 1992, **31**, 3251 (*isol, ir, pmr, cmr, ms, struct*)

13-Hydroxylupanine **H-554**

13-Hydroxy-2-oxosparteine. Octalupine.
Hydroxylupanine
[15358-48-2]



(+)-form

$C_{15}H_{24}N_2O_2$ 264.367
Alkaloid from *Cytisus scoparius*, *Chamaecytisus absinthoides*, *Calpurnia aurea* ssp. *aurea*, *Lupinus angustifolius*, *Lupinus albus*, *Lupinus hilarianus*, *Lupinus perennis*, *Lupinus polyphyllus*, *Lupinus wyethii*, *Cadia* spp., *Genista cinerea*, *Sarothamnus catalaunicus* and *Thermopsis cinerea*. Mp 178-180° (172-174°) Mp 76° (hydrate). $[\alpha]_D^{20} +74.1$ (c, 3.6 in H_2O). $[\alpha]_D +51.8$.

►OK5746000

Hydroiodide: Mp 234-238°.

Ac: **13-Acetoxyupanine**

[72822-06-1]
 $C_{17}H_{26}N_2O_3$ 306.404

Alkaloid from *Lupinus polyphyllus*, *Lupinus mutabilis* and *Lupinus albus*. Mp 103-105°.

Propanoyl: **13-Propanoxyupanine**. 13-Propoxyupanine (incorr.)

[118636-92-3]

$C_{18}H_{28}N_2O_3$ 320.431

Alkaloid detected in *Lupinus polyphyllus*, *Lupinus mutabilis*, *Lupinus angustifolius* and *Lupinus albus*.

Butanoyl: **13-Butyroxylupanine**

[118636-94-5]

$C_{19}H_{30}N_2O_3$ 334.458

Alkaloid detected in *Lupinus polyphyllus*, *Lupinus mutabilis*, *Lupinus angustifolius* and *Lupinus albus*.

O-(2-Methylpropanoyl): **13-Isobutyryloxyupanine**

[118636-93-4]

$C_{19}H_{30}N_2O_3$ 334.458

Alkaloid detected in *Lupinus polyphyllus*, *Lupinus mutabilis*, *Lupinus angustifolius* and *Lupinus albus*.

Pentanoyl: **13-Valeroyloxyupanine**

[118636-96-7]

$C_{20}H_{32}N_2O_3$ 348.484

Alkaloid detected in *Lupinus albus*.

O-(2-Methylbutanoyl): **13-(2-Methylbutyryloxy)upanine**

[81362-35-8]

$C_{20}H_{32}N_2O_3$ 348.484

Alkaloid from cell cultures of *Lupinus polyphyllus*, detected by glc/ms (Fabaceae). Also from *Rothia trifoliata*.

O-(3-Methylbutanoyl): **13-Isovaleroyloxyupanine**

[118636-95-6]

$C_{20}H_{32}N_2O_3$ 348.484

Alkaloid detected in *Lupinus polyphyllus*, *Lupinus mutabilis*, *Lupinus angustifolius* and *Lupinus albus*.

Hexanoyl: **13-Caproyloxyupanine**

[134958-48-8]

$C_{21}H_{34}N_2O_3$ 362.511

Alkaloid from *Lupinus angustifolius*.

Angeloyl: **13-Angeloyloxyupanine**

[57943-35-8]

$C_{20}H_{30}N_2O_3$ 346.469

Alkaloid from *Calpurnia aurea*, seeds of *Lupinus mutabilis*, and from cell cultures of *Lupinus polyphyllus*, detected by glc/ms (Fabaceae).

Tigloyl: **13-Tigloyloxyupanine**

[57943-34-7]

$C_{20}H_{30}N_2O_3$ 346.469

Alkaloid from *Lupinus polyphyllus* and seeds of *Lupinus mutabilis* (Fabaceae). Oil. Mp 260° dec. (as perchlorate). Bp_{0.001} 180-200° (synthetic).

O-(4-Hydroxy-2-methyl-2E-butenoyl):

13-(4-Hydroxytigloyloxy)upanine

[149155-69-1]

$C_{20}H_{30}N_2O_4$ 362.468

Alkaloid from seeds of *Ormosia krugii* (Fabaceae). Yellow oil. $[\alpha]_D^{20} +42$ (c, 0.1 in MeOH).

O-(3-Hydroxy-5-octenoyl) (Z-): **Cineroctine**

[155454-00-5]

$C_{23}H_{36}N_2O_4$ 404.548

Alkaloid from aerial parts of *Genista cinerea* subsp. *cinerea*, *Genista cinerea* ssp. *speciosa* and *Genista ramosissima* (Fabaceae). Yellow oil. $[\alpha]_D^{30} +6.4$ (c, 1 in MeOH).

Benzoyl: **13-Benzoyloxyupanine**

[34226-97-6]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from *Lupinus angustifolius*, *Lupinus albus*, *Lupinus polyphyllus*, seeds of *Lupinus mutabilis* and *Genista cinerea* (Fabaceae). Mp 205°. $[\alpha]_D^{20} +35$.

O-(3-Hydroxy-4-methoxybenzoyl): **Isocinevanine**

[36101-56-1]

$C_{23}H_{30}N_2O_5$ 414.5

Alkaloid from *Sarothamnus patens* (Fabaceae). Mp 210-211°.

O-(4-Hydroxy-3-methoxybenzoyl): **Cinevanine**. 13-Vanilloxyloxyupanine

[34221-21-1]

$C_{23}H_{30}N_2O_5$ 414.5

Alkaloid from *Genista cinerea*, also detected in cell cultures of *Lupinus polyphyllus* (Fabaceae). Mp 207-208°. $[\alpha]_D +69$ (MeOH).

O-(3,4-Dimethoxybenzoyl): **Cineverine**

[10547-06-5]

$C_{24}H_{32}N_2O_5$ 428.527

Alkaloid from *Genista cinerea* (Fabaceae). Mp 155-157° (148-149°). $[\alpha]_D^{20} +65$ (c, 1 in EtOH). $[\alpha]_D +71$ (MeOH).

O-(3,5-Dihydroxy-4-methoxybenzoyl):

Cinegalline

[20772-38-7]

$C_{23}H_{30}N_2O_6$ 430.5

Alkaloid from *Genista cinerea* (Fabaceae). Mp 223-225°. $[\alpha]_D +46.7$ (EtOH).

O-(3-Hydroxy-4,5-dimethoxybenzoyl):

Cinegalleine

[27570-28-1]

$C_{24}H_{32}N_2O_6$ 444.527

Alkaloid from *Genista cinerea* (Fabaceae). Mp 182°. $[\alpha]_D^{20} +48$ (c, 0.25 in MeOH).

O-(3-Formyloxy-4,5-dimethoxybenzoyl):

Formylcinegalleine

[33192-90-4]

$C_{25}H_{32}N_2O_7$ 472.537

Alkaloid from *Genista cinerea* (Fabaceae). Needles (MeOH). Mp 188-189°.

O-(3,4,5-Trimethoxybenzoyl): **Sarodesmine**

[30430-48-9]

$C_{25}H_{34}N_2O_6$ 458.553

Constit. of branches of *Sarothamnus catalaunicus* (Fabaceae). Needles (Et₂O or MeOH). Mp 174°. $[\alpha]_D^{20} +73.6$ (c, 1 in MeOH).

E-Cinnamoyl: **trans-13-Cinnamoyloxyupanine**

[5835-04-1]

$C_{24}H_{30}N_2O_3$ 394.513

Alkaloid from *Lupinus angustifolius*, *Lupinus polyphyllus* and seeds of *Lupinus mutabilis* (Fabaceae). Mp 166°. $[\alpha]_D^{20} +42$.

Z-Cinnamoyl: **cis-13-Cinnamoyloxyupanine**

[6068-29-7]

$C_{24}H_{30}N_2O_3$ 394.513

Alkaloid from *Lupinus* spp. and seeds of *Lupinus mutabilis*. Mp 145° (as hydrochloride). $[\alpha]_D^{20} +26.1$.

O-Phenylacetyl: **13-Hydroxylupanine phenylacetate**

[57943-36-9]

$C_{23}H_{30}N_2O_3$ 382.502

Alkaloid in leaves and twigs of *Cadia purpurea* (Fabaceae). Identified only by tlc comparison with synthetic sample.

O-(4-Hydroxyphenylacetyl): **Cadiaine**.

13-Hydroxylupanine p-hydroxyphenylacetate

[36445-14-4]

$C_{23}H_{30}N_2O_4$ 398.501

Alkaloid in leaves and twigs of *Cadia purpurea* (Fabaceae). Identified only by tlc comparison with synthetic sample.

O-(2-Pyrrolicarbonyl): **Calpurnine**. **Oroboidine**. **Hoe 933**

[6874-80-2]

$C_{20}H_{27}N_3O_3$ 357.452

Alkaloid from *Calpurnia lasiogyne*, *Calpurnia aurea*, *Virgilia capensis* and *Cadia ellisiana* (Fabaceae). Shows hypotensive effects. Prisms (EtOAc). Mp 152-154°. $[\alpha]_D^{22} +59$ (c, 1 in CHCl₃). $[\alpha]_D^{20} +47$ (c, 1.16 in EtOH).

►Highly toxic (fish, mice). LD₅₀ (mus, orl) 32 mg/kg. LD₅₀ (mus, ivn) 3.1 mg/kg. UX9372400

O-(2-Pyrrolicarbonyl), perchlorate:

Prisms (MeOH). Mp 279° (255-260°).

Me ether: **13-Methoxyupanine**

[4697-77-2]

$C_{16}H_{26}N_2O_2$ 278.394

Alkaloid from combined leaf/hypocotyl extracts of *Lupinus polyphyllus* (Fabaceae).

Et ether: **13-Ethoxyupanine**

[58165-60-9]

C₁₇H₂₈N₂O₂ 292.42

Alkaloid from *Cadia purpurea*, identified by ms (Fabaceae).

13-Epimer: **Jamaidine**. Epihydroxylupanine

[6870-60-6]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from seeds of *Ormosia jamaicensis* and from *Lupinus polyphyllus* (Fabaceae), also obt. by reacen. of Angustifoline, A-1013 with CH₂O. Mp 194.5-195° Mp 170° (synthetic) Mp 190° (polymorph, synthetic). [α]_D²⁰ +63.7 (c, 0.66 in EtOH). [α]_D²⁰ +47 (c, 1.5 in MeOH) (synthetic).

13-Epimer, perchlorate: Mp 175-177°.

13-Epimer, Ac: [3382-87-4]

Mp 170.5-171°.

13-Epimer, Me ether: **13-Epimethoxylupanine**. Alkaloid W95. 13β-Methoxylupanine

[3382-85-2]

C₁₆H₂₆N₂O₂ 278.394

Alkaloid from *Lupinus angustifolius* (Fabaceae) and *Acosmium panamense*. Mp 97°. [α]_D²⁴ +64.5 (c, 1.24 in EtOH) (synthetic).

Marion, L. et al., *Can. J. Chem.*, 1951, **29**, 721-724 (*isol, ir*)

Bohlmann, F. et al., *Chem. Ber.*, 1958, **91**, 2194-2205 (*ir*)

Lloyd, H.A. et al., *J.O.C.*, 1960, **25**, 1959-1962; 1961, **26**, 2143-2145 (*Jamaidine*)

Bratek, M.D. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1961, **9**, 705 (*Jamaidine, 13-Epimethoxylupanine*)

Bratek-Wiewiorowska, M.D. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1963, **11**, 629 (*isol, esters*)

Bohlmann, F. et al., *Chem. Ber.*, 1963, **96**, 2254-2255 (*tiglate*)

Goosen, A. et al., *J.C.S.*, 1963, 3067 (*Calpurnine*)

Gerrans, G.C. et al., *J.C.S.*, 1964, 2202 (*Calpurnine*)

Bohlmann, F. et al., *Chem. Ber.*, 1965, **98**, 3133-3141 (*synth*)

Vazquez, M.D. et al., *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1966, **62**, 837; *CA*, **66**, 76216h (*Jamaidine, synth*)

Faugeras, G. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1966, **263**, 436; 1968, **267**, 538; 1970, **270**, 203; 1971, **271**, 611; 1219 (*isol, esters*)

Fales, H.M. et al., *J.A.C.S.*, 1970, **92**, 1590-1597 (*ms*)

Faugeras, G. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 1372 (*isol, esters*)

Wysocka, W. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1974, **22**, 831-842 (*13-epimer, synth*)

Klyne, W. et al., *J.C.S. Perkin 1*, 1974, 2565-2570 (*cd*)

Bohlmann, F. et al., *Chem. Ber.*, 1975, **108**, 1043-1051 (*struct, cmr*)

Radema, M.H. et al., *Planta Med.*, 1975, **28**, 143 (*13-Ethoxylupanine*)

Lindner, G. et al., *Arzneim.-Forsch.*, 1976, **26**, 1651 (*Calpurnine, pharmacol*)

Kaluski, Z. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1977, **25**, 347; *CA*, **88**, 23224g (*cryst struct*)

Van Eijk, J.L. et al., *Planta Med.*, 1977, **32**, 275 (*Capurnine, isol*)

Wink, M. et al., *Planta Med.*, 1982, **44**, 15-20 (*Lupinus polyphyllus constits*)

Hatzold, T. et al., *J. Agric. Food Chem.*, 1983, **31**, 934-938 (*Lupinus mutabilis constits*)

Mühlbauer, P. et al., *Planta Med.*, 1988, **54**, 237-239 (*esters, synth, ms*)

Strack, D. et al., *Prog. Hemostasis Thromb.*, 1991, **31**, 1493-1498 (*13-Caproyloxylupanine*)

Veen, G. et al., *Phytochemistry*, 1992, **31**, 4343-4345 (*13-epimer tigloyl deriv*)

Nasution, M.P. et al., *Phytochemistry*, 1993, **32**, 1603-1605 (*13α-(4-Hydroxytigloyloxy)lupanine*)

Van Rensen, I. et al., *Phytochemistry*, 1994, **35**, 421-424 (*Cineractine, Cineverine*)

Veitch, N.C. et al., *Phytochemistry*, 1997, **45**, 847-850 (*13-Epimethoxylupanine*)


Borowiak, T. et al., *J. Mol. Struct.*, 2005, **753**, 27-34 (*cryst struct, 13-epimer*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CAZ125 (*Calpurnine*)

α-Hydroxymagnocurarine

H-555

1a-Hydroxymagnocurarine



C₁₉H₂₄NO₄⁺ 330.403

(1*R,αR*)-form [152154-55-7]

[152154-56-8]


Alkaloid from stems of *Cryptocarya konishii* (Lauraceae). Amorph. solid (as perchlorate). Mp 142° (perchlorate). [α]_D^{24.5} +35 (c, 0.50 in MeOH) (perchlorate).

Lee, S.-S. et al., *J. Nat. Prod.*, 1993, **56**, 1971 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

2-Hydroxy-4-mercaptopyrimidine

H-556

3,4-Dihydro-4-thioxo-2(1*H*)-pyrimidinone, 9CI. 4-Mercapto-2-pyrimidinol. 4-Thiouracil [591-28-6]



C₄H₄N₂OS 128.154

Di-NH form predominates. Intermediate mono-NH tautomers also possible. Isol. from *Streptomyces libani soldani*. Broad antibacterial agent with low specific activity. Shows antineoplastic activity. Yellow cryst. (EtOH). Mp 317-321°. λ_{max} 241 (ε 5380); 253 (sh) (ε 5100); 342 (ε 20900) (MeOH/KOH) (Derep). λ_{max} 246 (ε 3960); 327 (ε 17000) (MeOH) (Derep).

N¹-Benzoyl: [215544-47-1]
C₁₁H₈N₂O₂S 232.262
Yellow needles (MeCN). Mp 181° dec.

N¹,N³-Di-Me: [49785-67-3]
C₆H₈N₂O₂S 156.208

Yellow solid. Mp 183°.

Fox, J.J. et al., *J.A.C.S.*, 1960, **82**, 486-489 (*synth*)

Mizuno, Y. et al., *Chem. Pharm. Bull.*, 1962, **10**, 647-652 (*synth*)

Dolak, L. et al., *Antimicrob. Agents Chemother.*, 1977, **11**, 569-570 (*isol, cmr, ms, props*)

Still, I.W.J. et al., *Can. J. Chem.*, 1978, **56**, 725-729 (*cmr*)

Brown, D.J. et al., *Aust. J. Chem.*, 1980, **33**, 1147-1152 (*synth*)

Tajiri, A. et al., *Biochim. Biophys. Acta*, 1981, **656**, 1-15 (*uv*)

Lapucha, A.R. et al., *Synthesis*, 1987, 256-258 (*synth, ir, pmr, cmr, ms*)

Isono, K. et al., *J. Antibiot.*, 1988, **41**, 1711-1739 (*rev*)

Kaneko, K. et al., *Synthesis*, 1988, 152-154 (*synth*)

Frieden, M. et al., *J.C.S. Perkin 1*, 1998, 2827-2832 (*synth, pmr, cmr, N-benzoyl*)


Friden, M. et al., *J.C.S. Perkin 1*, 1998, 2827-2832 (*synth, pmr*)

Rodriguez, J. et al., *Chem. Eur. J.*, 1999, **5**, 3549-3561 (*1,3-di-Me*)

β-Hydroxymescaline

H-557

α-(Aminomethyl)-3,4,5-trimethoxybenzenemethanol, 9CI. 2-Amino-1-(3,4,5-trimethoxyphenyl)ethanol. β-Hydroxy-3,4,5-trimethoxyphenethylamine [13079-18-0]



C₁₁H₁₇NO₄ 227.26

(±)-form [92536-08-8]

Impurity in illegally synthesised Mescaline, M-219. Needles (MeOH/Et₂O). Mp 196-199°.

(ξ)-form

Alkaloid detected in *Pereskia grandiflora* (Cactaceae).


Friedman, O.M. et al., *J. Med. Chem.*, 1963, **6**, 227 (*synth*)

Doetsch, P.W. et al., *J. Chromatogr.*, 1980, **189**, 79 (*occur*)

2-(4-Hydroxy-3-methoxybenzoyl)imidazole

H-558

2-Vanilloylimidazole [114703-13-8]



C₁₁H₁₀N₂O₃ 218.212


Metab. from the Australian ascidian *Aplidium pliciferum*. Also obt. from aged extracts of an undescr. New Zealand *Aplidium* sp. (not detected in freshly-prepared ascidian extracts). Mp 226°. Prob. not a naturally occurring metab. Genus name given as *Aplydium*. λ_{max} 369 (ε 4300) (MeOH/KOH) (Derep). λ_{max} 240

(ϵ 2300); 291 (ϵ 2400); 324 (ϵ 2600)
(MeOH) (Derep).

Arabshahi, L. *et al.*, *Tet. Lett.*, 1988, **29**, 1099-1102 (*isol*, *pmr*, *struct*)
Copp, B.R. *et al.*, *Tet. Lett.*, 1989, **30**, 3703-3706 (*uv*, *ir*, *cmr*, *cryst struct*)

11-Hydroxy-12-methoxycoptisine H-559


[30426-64-3]
[32585-02-7]



$C_{20}H_{16}NO_6^{\oplus}$ 366.349
Alkaloid from *Coptis groenlandica* (Ranunculaceae). Mp 270-275° (as chloride).
[30426-69-8, 30426-70-1, 30426-68-7]
Cooper, S.F. *et al.*, *Planta Med.*, 1971, **19**, 23

1-(1-Hydroxy-2-methoxyethyl)-4-methoxy- β -carboline H-560

β ,4-Dimethoxy-9H-pyrido[3,4-b]indole-1-ethanol, 9CI
[89915-40-2]



$C_{15}H_{16}N_2O_3$ 272.303
Alkaloid from the root bark of *Ailanthus altissima* (Simaroubaceae). Needles (Me₂CO). Mp 223°.


Ac:

Needles (Me₂CO). Mp 125°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **32**, 170 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

4-Hydroxy-3-methoxy-2-methyl-9H-carbazole-1-carboxaldehyde, 9CI H-561

Carbazomycin E. *Carbazomycin*al
[103744-20-3]



$C_{15}H_{13}NO_3$ 255.273
Isol. from *Streptovorticillium ehimensense*. Pale-yellow needles (EtOAc/hexane). Mp 224°. λ_{max} 214 (ϵ 24400); 227 (ϵ 23200); 263 (ϵ 12200); 295 (ϵ 16300); 320 (ϵ 5100); 372 (ϵ 8400) (MeOH) (Derep).


6-Methoxy-4-Hydroxy-3,6-dimethoxy-2-methyl-9H-carbazole-1-carboxaldehyde, 9CI. *Carbazomycin F*. 6-Methoxycarbazomycin
[103744-21-4]
 $C_{16}H_{15}NO_4$ 285.299
From *Streptovorticillium ehimensense*. Pale-yellow needles (EtOAc/hexane). λ_{max} 215 (ϵ 27000); 227 (ϵ 24500); 245 (ϵ 13000); 268 (ϵ 13000); 310 (ϵ 17000); 382 (ϵ 8500) (MeOH) (Derep).

Kondo, S. *et al.*, *J. Antibiot.*, 1986, **39**, 727 (*isol*, *struct*)

Naid, T. *et al.*, *J. Antibiot.*, 1987, **40**, 157 (*isol*)
Knöflker, H.-J. *et al.*, *Heterocycles*, 1991, **32**, 2443 (*synth*, *Carbazomycin E*)

6-Hydroxy-5-methoxy-1-methyl-2,3,4(1H)-pyridinetrione, 9CI H-562

SEN 34. *Antibiotic SEN 34*
[70158-18-8]




$C_7H_7NO_5$ 185.136
Isol. from *Streptomyces aburaviensis*. Weak antimicrobial activity against gram-positive and -negative bacteria. Yellow prisms. Sol. MeOH, CHCl₃; fairly sol. H₂O; poorly sol. hexane. Mp 161-162°. λ_{max} 215 (ϵ 8732); 275 (ϵ 9309); 385 (ϵ 1924) (MeOH) (Berdy). λ_{max} 276; 385 (EtOH) (Berdy). λ_{max} 216; 276; 385 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 160 mg/kg.

Ozaki, M. *et al.*, *Heterocycles*, 1978, **11**, 191 (*isol*)

1-Hydroxy-5-methoxy-6-methyl-2(1H)-pyridinone, 9CI H-563

Cepabactin. *BN 227*. *G 1549*. *Antibiotic BN 227*. *Antibiotic G 1549*
[72731-33-0]



$C_7H_9NO_3$ 155.153
Prod. by *Pseudomonas* sp. Active against gram-positive and gram-negative bacteria. Siderophore. Cryst. (EtOAc). Mp 115°. λ_{max} 229 (ϵ 8680); 330 (ϵ 7150) (MeOH) (Derep).

► UU7786100

Fe complex (3:1): *Antibiotic BN 227F*. *BN 227F*
[73349-00-5]

$C_{21}H_{24}FeN_3O_9$ 518.282
Prod. by *Pseudomonas* sp. Active against gram-positive and -negative bacteria. Red crust. (CHCl₃). Mp 156°. λ_{max} 224 (ϵ 63000); 318 (ϵ 15100) (MeOH) (Derep).

Cu complex (2:1): [73349-01-6] Active against gram-positive bacteria and fungi. Green cryst. Mp 308-310° dec.

Al complex (3:1): *Antibiotic BN 229A*. *BN 229A*

[73298-51-8]

$C_{21}H_{24}AlN_3O_9$ 489.417

From *Pseudomonas cepacia*. Active against *Staphylococcus aureus*.

Itoh, J. *et al.*, *J. Antibiot.*, 1979, **32**, 1089; 1980, **33**, 377 (*isol*, *struct*)

Barker, N.W. *et al.*, *J. Antibiot.*, 1979, **32**, 1096 (*isol*, *struct*)


Japan. Pat., 1979, 79 157 501; *CA*, **92**, 162178 (*deriv*)

Japan. Pat., 1979, 79 132 577; *CA*, **93**, 6121 (*isol*)

Meyer, J.M. *et al.*, *J. Gen. Microbiol.*, 1989, **135**, 1479 (*isol*, *uv*, *pmr*, *props*)

Ohta, A. *et al.*, *Heterocycles*, 1990, **30**, 875 (*synth*, *ms*, *ir*, *pmr*)

5-Hydroxy-4-methoxy-methyl-1H-pyrrole-3-carboxylic acid H-564



$C_7H_9NO_4$ 171.152

Me ester: *Methyl 5-hydroxy-4-methoxy-methyl-1H-pyrrole-3-carboxylate*
[128922-72-5]

$C_8H_{11}NO_4$ 185.179

Alkaloid from stems of *Gnetum montanum* (Gnetaceae). Mp 194-197°.


Zhou, J. *et al.*, *Zhivwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 878-882; *CA*, **113**, 94757d (*isol*, *deriv*)

3-Hydroxy-4-methoxyphenethylamine H-565

5-(2-Aminoethyl)-2-methoxyphenol. 4-O-Methyl-dopamine

[3213-30-7]

[645-33-0 (hydrochloride)]



$C_9H_{13}NO_2$ 167.207

Alkaloid from *Acacia rigidula*, *Pachycereus pecten-aboriginum*, the soft coral *Nephthea* sp., and human urine. Shows cardiovascular and CNS activity. Needles (MeOH/EtOAc) (as hydrochloride). Mp 206.5-207° (hydrochloride). Identified chromatographically as a nat. prod.

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): *N-trans-Feruloyl-4-O-methyl-dopamine*
[78510-20-0]

$C_{19}H_{21}NO_5$ 343.379

Alkaloid from roots of the pseudocereal *Chenopodium album* (lambsquar-

ters) (Chenopodiaceae). Oil.
 N-(3,4-Dimethoxy-E-cinnamoyl): **N-(4-O-Methyl-trans-feruloyl)-4-O-methyl-dopamine**
 [651768-68-2]
 $C_{20}H_{23}NO_5$ 357.405
 Alkaloid from *Chenopodium album* (lambsquarters). Oil.
 N-Me: 3-Hydroxy-4-methoxy-N-methyl-phenethylamine
 [35266-68-3]
 $C_{10}H_{15}NO_2$ 181.234
 Alkaloid from *Acacia rigidula*.
 N,N,N-Tri-Me: 3-Hydroxy-4-methoxy-N,N,N-trimethylbenzeneethanaminium. **Salicifoline**
 [6882-07-1]
 $C_{12}H_{20}NO_2^{\oplus}$ 210.295
 Alkaloid from *Magnolia grandiflora* roots, *Magnolia kobus*, *Magnolia denudata*, *Magnolia stellata* and *Michelia alba* (Magnoliaceae). Mp 260-261° dec. (as chloride).
 Ramirez, F.A. et al., *J.A.C.S.*, 1950, **72**, 2781-2782 (synth)
 Tomita, M. et al., *Yakugaku Zasshi*, 1952, **72**, 197-203; 727-731; 766-767; 1256-1260; *CA*, **47**, 1627; 12288; 12409; **48**, 2639 (Salicifoline)
 Nakano, T. et al., *Pharm. Bull.*, 1954, **2**, 321; 1956, **4**, 409 (Salicifoline)
 Kakimoto, Y. et al., *J. Biol. Chem.*, 1962, **237**, 208 (isol)
 Yang, T.H. et al., *Yakugaku Zasshi*, 1962, **82**, 816-820; *CA*, **58**, 7991 (Salicifoline)
 Agurell, S. et al., *J. Nat. Prod.*, 1971, **34**, 183-187 (isol, struct)
 Gregson, R.P. et al., *Experientia*, 1981, **37**, 493-494 (isol)
 Kohno, M. et al., *Bull. Chem. Soc. Jpn.*, 1990, **63**, 1252-1254 (synth, ir, pmr)
 Horio, T. et al., *Phytochemistry*, 1993, **33**, 807-808 (N-Feruloyl-4-methyl-dopamine)
 Clement, B.A. et al., *Phytochemistry*, 1998, **49**, 1377-1380 (isol)
 Cutillo, F. et al., *Phytochemistry*, 2003, **64**, 1381-1387 (4-O-Methylferuloyl-4-O-methyl-dopamine)

4-Hydroxy-3-methoxyphenethylamine H-566


4-(2-Aminoethyl)-2-methoxyphenol, 9CI, 8CI. 3-O-Methyl-dopamine. 3-Methoxytyramine. Homovanillylamine. 4-(2-Aminoethyl)guaiacol
 [554-52-9]
 $C_9H_{13}NO_2$ 167.207
 Alkaloid from *Lophophora williamsii*, *Carnegiea gigantea*, *Trichocereus* spp. and other cacti, also detected in *Echinocereus merkeri*. Constit. of human brain tissue and urine. Intermed. in mescaline biosynth. hypotensive agent. Prisms (EtOH). Mp 159° (156-157°).
 Hydrochloride: [1477-68-5]
 Prisms (EtOH). Mp 210-211°.
 N-(2-Hydroxy-5-methoxybenzoyl): **Aduncamide**
 [154992-23-1]
 $C_{17}H_{19}NO_5$ 317.341
 Constit. of *Piper aduncum*. λ_{max} 280 (ε 25118); 323 (ε 36300) (MeOH) (Berdy).
 N-(3-Hydroxy-4-methoxy-E-cinnamoyl): **N-(4-O-Methyl-trans-caffeoyl)-3-O-methyl-dopamine**. N-trans-Isoferuloyl-3-

O-methyl-dopamine
 [125002-88-2]
 $C_{19}H_{21}NO_5$ 343.379
 Alkaloid from *Chenopodium album* (lambsquarters). Oil.
 N-(4-Hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyl-3-methoxytyramine**. N-trans-Feruloyl-3-O-methyl-dopamine
 [78510-19-7]
 $C_{19}H_{21}NO_5$ 343.379
 Alkaloid from *Annona cherimola* (cherimoya), immature green onions and spinach and from the wood of *Actinodaphne longifolia*. Needles. Mp 111-113°. λ_{max} 220 (ε 3.42); 290 (ε 2.67); 319 (ε 3.15) (EtOH).
 N-(4-β-D-Galactopyranosyloxy-3-methoxy-E-cinnamoyl): **Cimicifugamide**
 [155159-01-6]
 $C_{25}H_{31}NO_{10}$ 505.521
 Alkaloid from the rhizomes of *Cimicifuga dahurica*.
 N-(4-β-D-Glucopyranosyloxy-3-methoxy-E-cinnamoyl), 4-O-β-D-glucopyranoside: **Aristomanoside**
 $C_{31}H_{41}NO_{15}$ 667.663
 Alkaloid from the stems of *Aristolochia manshuriensis*. Amorph. yellow powder (CHCl₃/MeOH). Mp 195-197°. λ_{max} 219 (log ε 4.3); 228 (log ε 4.3); 284 (log ε 4.29); 313 (log ε 4.24) (MeOH).
 N-(4-Hydroxy-3-methoxy-Z-cinnamoyl): **N-cis-Feruloyl-3-methoxytyramine**
 $C_{19}H_{21}NO_5$ 343.379
 Alkaloid from *Annona cherimola* (cherimoya) and *Aristolochia gehrtii*. Needles. Mp 107-110° (248-251). λ_{max} 222 (ε 3.47); 290 (ε 2.64); 319 (ε 3.2) (EtOH).
 N-(4-β-D-Galactopyranosyloxy-3-methoxy-Z-cinnamoyl): **Isocimicifugamide**
 [156318-64-8]
 $C_{25}H_{31}NO_{10}$ 505.521
 Constit. of *Chenopodium dahurica*. Amorph. powder. Mp 97-100°. $[\alpha]_D$ -46.2 (c, 0.13 in MeOH).
 N-(4-β-D-Glucopyranosyloxy-3-methoxy-Z-cinnamoyl): **Drodrinin**
 [102062-99-7]
 $C_{25}H_{31}NO_{10}$ 505.521
 Constit. of *Scopolia japonica*. Antibacterial agent. Sol. MeOH. λ_{max} 210; 230; 287; 312 (MeOH).
 N-Me: **4-Hydroxy-3-methoxymethylphenethylamine**
 $C_{10}H_{15}NO_2$ 181.234
 Alkaloid in *Trichocereus courantii* and *Lophophora williamsii* (Cactaceae).
 N,N-Di-Me: 4-[2-(Dimethylamino)ethyl]-2-methoxyphenol, 9CI. **4-Hydroxy-3-methoxydimethylphenethylamine**. N,N-Dimethyl-3-methoxytyramine. *Magnosprengerine*
 [35266-63-8]
 $C_{11}H_{17}NO_2$ 195.261
 Alkaloid from *Ariocarpus agavoides* and *Magnolia sprengeri* (Cactaceae, Magnoliaceae). Mp 258-260°.
 N,N,N-Tri-Me: 4-Hydroxy-3-methoxy-N,N,N-trimethylbenzeneethanaminium, 9CI. **4-Hydroxy-3-methoxytrimethylphenethylammonium**

$C_{12}H_{20}NO_2^{\oplus}$ 210.295
 Quaternary alkaloid from *Alhagi pseudalhagi* (Fabaceae).
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1294C (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 623A (nmr)
 Kratzl, K. et al., *Monatsh. Chem.*, 1952, **83**, 1038 (synth)
 Agurell, S. et al., *J. Nat. Prod.*, 1969, **32**, 40 (isol, synth)
 Lundstrom, J. et al., *Acta Chem. Scand.*, 1971, **25**, 3489
 Milne, G.W.A. et al., *Anal. Chem.*, 1973, **45**, 1952 (ms)
 Crosby, D.M. et al., *J. Nat. Prod.*, 1973, **36**, 418 (isol)
 Cherayil, G.D. et al., *J. Pharm. Sci.*, 1973, **62**, 2054 (synth)
 Bruhn, J.G. et al., *Econ. Bot.*, 1974, **27**, 241; *CA*, **80**, 93142n (isol, deriv)
 Ghosal, S. et al., *Planta Med.*, 1974, **26**, 318 (isol, deriv)
 Pardanani, J.H. et al., *J. Nat. Prod.*, 1977, **40**, 585 (isol)
 Strombom, J. et al., *Acta Pharm. Suec.*, 1978, **15**, 127; *CA*, **89**, 126107w (isol)
 Doetsch, P.W. et al., *J. Chromatogr.*, 1980, **189**, 79 (occur)
 Meyer, B.N. et al., *Phytochemistry*, 1980, **19**, 719 (occur)
 Pummangura, S. et al., *J. Nat. Prod.*, 1981, **44**, 498 (isol)
Japan. Pat., 1985, 85 199 395; *CA*, **104**, 205524m (Drodrinin)
 Cao, Z. et al., *Zhongcaoyao*, 1985, **16**, 386; *CA*, **104**, 17645c (Magnosprengerine)
 Tanaka, H. et al., *Phytochemistry*, 1989, **28**, 2516 (N-Feruloyl-3-methyl-dopamine)
 Okabe, N. et al., *Acta Cryst. C*, 1992, **48**, 1698 (cryst struct)
 Li, C.J. et al., *Chin. Chem. Lett.*, 1993, **4**, 891 (Cimicifugamide)
 Orjala, J. et al., *Nat. Prod. Lett.*, 1993, **2**, 231-236 (Aduncamide)
 Li, C.J. et al., *Yaoxue Xuebao*, 1994, **29**, 129 (Isocimicifugamide)
 Ding, P.Y. et al., *Chin. Chem. Lett.*, 1996, **7**, 545 (Cimicifugamide, synth)
 Chen, C.-Y. et al., *Phytochemistry*, 1998, **49**, 1443-1447 (N-feruloyl derivs)
 Navickienė, H.M.D. et al., *J. Braz. Chem. Soc.*, 2001, **12**, 467-472 (N-cis-feruloyl)
 Wu, P.-L. et al., *J. Nat. Prod.*, 2003, **66**, 996-998 (Aristomanoside)
 Cutillo, F. et al., *Phytochemistry*, 2003, **64**, 1381-1387 (4-O-Methylcaffeoyl-3-O-methyl-dopamine)
 Xiao, H. et al., *J. Agric. Food Chem.*, 2006, **54**, 8417-8424 (4-Hydroxy-3-methoxy-E-cinnamoyl, isol)

2-(2-Hydroxy-4-methoxyphenyl)-5,8-dimethoxy-3-propyl-4(1H)-quinolinone H-567

4-Hydroxy-2-(2-hydroxy-4-methoxyphenyl)-5,8-dimethoxy-3-propylquinoline
 [1010070-86-6]




$C_{21}H_{23}NO_5$ 369.416

Alkaloid from the fruit of *Casimiroa edulis*. Cryst. (CHCl₃). Mp 115-116°. λ_{\max} 280 (ϵ 14294); 300 (ϵ 8140); 320 (ϵ 6590) (EtOH).

Awaad, A.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 576-580 (isol, pmr, cmr)

4-Hydroxy-4-(4-methoxyphenyl)-1-methyl-5-thioxo-2-imidazolidinone, 9CI H-568



C₁₁H₁₂N₂O₃S 252.293

(±)-form

Metab. from the Indian ocean ascidian *Polycarpa clavata*. Amorph. solid. Prob. an artifact. CAS no. not found 8-14CI. λ_{\max} 202 (log ϵ 3.99); 220 (log ϵ 3.91); 278 (log ϵ 3.88) (MeOH).

Me ether: 4-Methoxy-4-(4-methoxyphenyl)-1-methyl-5-thioxo-2-imidazolidinone

C₁₂H₁₄N₂O₃S 266.32


Metab. from the ascidians *Polycarpa clavata* and *Polycarpa aurata*. Inosine monophosphate dehydrogenase inhibitor. Needles. Mp 127-129°. Prob. an artifact. CAS no. not found 8-14CI. λ_{\max} 203 (log ϵ 4.14); 222 (log ϵ 4.17); 279 (log ϵ 4.1) (MeOH). λ_{\max} 224 (ϵ 8822); 282 (ϵ 7640) (MeOH) (Berdy).

Abas, S.A. *et al.*, *J.O.C.*, 1996, **61**, 2709-2712 (*Me ether*, isol, uv, pmr, cmr, ms, struct)

Kang, H. *et al.*, *Tet. Lett.*, 1996, **37**, 2369-2372 (*Me ether*, isol, ir, uv, pmr, cmr, struct)

α-(4-Hydroxy-3-methoxyphenyl)-2-thiazolemethanol, 9CI H-569

[110281-34-0]



C₁₁H₁₁NO₃S 237.279

Minor metab. from the Australian ascidian *Aplydium pliciferum*. Mp 154-155°.

Ketone: (4-Hydroxy-3-methoxyphenyl)-2-thiazolylmethanone, 9CI. 2-(4-Hydroxy-3-methoxybenzoyl)thiazole [110281-33-9]

C₁₁H₉NO₃S 235.263


Metab. from *Aplydium pliciferum*. Mp 104°. λ_{\max} 233 (ϵ 4900); 300 (ϵ 4800); 341 (ϵ 5600) (EtOH) (Derep).

Arabshahi, L. *et al.*, *Tet. Lett.*, 1988, **29**, 1099-

1102 (isol, uv, ir, pmr, cmr, ms, struct)

7-Hydroxy-4-methoxy-8-prenylfuro[2,3-b]quinoline H-570

4-Methoxy-8-(3-methyl-2-butenyl)-furo[2,3-b]quinolin-7-ol, 9CI [107882-26-8]



C₁₇H₁₇NO₃ 283.326

Alkaloid from the leaves of *Sarcomelicope glauca* (Rutaceae) and aerial parts of *Haplophyllum tuberculatum*. Yellowish cryst. (C₆H₆). Mp 106-108°. λ_{\max} 252 (ϵ 34700); 306 (sh) (ϵ 5890); 318 (ϵ 7080); 330 (ϵ 6920); 342 (ϵ 6170) (MeOH) (Derep).

Me ether: 4,7-Dimethoxy-8-(3-methyl-2-butenyl)furo[2,3-b]quinoline, 9CI. 7-Methoxy-8-prenyldictamine. 8-(Dimethylallyl)-7-methoxydictamine [208170-82-5]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Almeidea coerulea*.


Cryst. Mp 86-88°.

Matiku, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1091 (isol, uv, ir, pmr, ms, struct)

Al-Yahya, M.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 899 (isol, cmr)

Santos, C.S. *et al.*, *J. Braz. Chem. Soc.*, 1998, **9**, 39-42; *C.A.*, **129**, 79124g (*Me ether*)

12-Hydroxy-16-methoxy-11,12,13,14-tetrahydrocamaoensine H-571



C₁₅H₁₆N₂O₃ 272.303

Alkaloid from the seeds of *Camoensia brevicalyx* (Fabaceae). Brown amorph. solid.

Waterman, P.G. *et al.*, *Phytochemistry*, 1982, **21**, 215 (isol, uv, ir, pmr, ms, struct)

2-Hydroxy-2-methylbutanoic acid, 9CI H-572

Ethylmethylglycollic acid. 2-Ethylactic acid [3739-30-8]



C₅H₁₀O₃ 118.132

(R)-form [37505-02-5]

Mp 73.5-74.5°. $[\alpha]_{\text{D}}^{24}$ -8.9 (c, 3.0 in CHCl₃).

Me ester: [55568-78-0]

C₆H₁₂O₃ 132.159

Oil. Bp 153°. $[\alpha]_{\text{D}}^{25}$ -6.7 (neat).

4-Phenylphenacyl ester: Mp 122°. $[\alpha]_{\text{D}}^{22}$ -2.1 (c, 3 in CHCl₃).

Nitrile, O-β-D-glucopyranoside: **Lotaustralin**

[534-67-8]

C₁₁H₁₉NO₆ 261.274

Glycoside from *Lotus australis*, other *Lotus* spp., *Trifolium repens* (white clover) and other plants. Needles (EtOAc). Mp 125-126°. $[\alpha]_{\text{D}}$ -19.1 (c, 1 in H₂O).

Nitrile, O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Neolinustatin**

[72229-42-6]

C₁₇H₂₉NO₁₁ 423.416

Isol. from *Passiflora pendens* and flaxseed meal. Cryst. (EtOH). Mp 190-192°. $[\alpha]_{\text{D}}^{25}$ -37 (c, 0.37 in H₂O).

Nitrile, O-[β-D-Apiofuranosyl-(1→6)-β-D-glucopyranoside]: **6'-Apiosyllotaustralin**

[170033-25-7]

C₁₆H₂₇NO₁₀ 393.39

Constit. of the roots of *Manihot esculenta* (cassava). Probable abs. config.

(S)-form [37505-07-0]

Mp 74-75°. $[\alpha]_{\text{D}}^{25}$ +9.1 (c, 1.6 in CHCl₃).

Benzyl ester: [587832-08-4]

C₁₂H₁₆O₃ 208.257

Oil. $[\alpha]_{\text{D}}^{19}$ -4.9 (c, 1.35 in CHCl₃).

4-Phenylphenacyl ester: Mp 117.5-118°. $[\alpha]_{\text{D}}^{25}$ +2.5 (c, 1.5 in CHCl₃).

Nitrile, O-β-D-glucopyranoside: **Epilotaustralin**

[55758-42-4]

C₁₁H₁₉NO₆ 261.274

Isol. from *Triticum monococcum* (wheat) and *Passiflora warmingii*.

(±)-form [37505-03-6]

Needles by subl. V. sol. H₂O, Et₂O. Mp 72-73°. Bp₁₆ 133-134°. pK_{a1} 3.73 (25°).

Me ester: [32793-34-3]

Oil. Bp 151.6-152°.

Amide: [1112-12-5]

C₅H₁₁NO₂ 117.147

Mp 160°.

Nitrile: 2-Cyano-2-butanol

[73683-34-8]

C₅H₉NO 99.132

Oil. Bp 180° Bp_{20.5} 91°.

Ac, nitrile:

C₇H₁₁NO₂ 141.169

Oil. Mp 195°.

Me ether:

C₆H₁₂O₃ 132.159

Bp₂₁ 112-113°.


Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 519C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 808A (nmr)

- Schmidt, R.R. *et al.*, *Synthesis*, 1994, 255 (synth, Ac, pmr)
 Jordis, U. *et al.*, *Org. Prep. Proced. Int.*, 1997, 29, 549-560 (nitrile)
 Akita, H. *et al.*, *Tetrahedron: Asymmetry*, 1999, 10, 2429-2439 (Osmaronin, synth)
 Yang, X. *et al.*, *Yaoxue Xuebao*, 2000, 35, 279-283 (isol, nitrile, esters)

6-Hydroxy-5-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde H-577

6-Formyl-3-hydroxy-4-prenylcarbazole. *Micromeline*[†]




C₁₈H₁₇NO₂ 279.338
 Alkaloid from the stem bark of *Micromelum hirsutum*. Yellow powder. λ_{max} 316 (log ε 3.3); 329 (log ε 3.7) (Me₂CO).

Ma, C. *et al.*, *Planta Med.*, 2005, 71, 261-267 (isol, pmr, cmr)

5-(3-Hydroxy-3-methyl-1-butenyl)-4,6-dimethoxyfuro[2,3-b]quinoline H-578

[953812-16-3]



C₁₈H₁₉NO₄ 313.352

(E)-form

Alkaloid from the leaves of *Choisya ternata*. Cryst. (EtOAc/hexane). Mp 144-145°.

Me ether: 4,6-Dimethoxy-5-(3-methoxy-3-methyl-1-butenyl)furo[2,3-b]quinoline [953812-15-2]

C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Choisya ternata*. Cryst. (EtOAc/hexane). Mp 94°.

7-Methoxy, 3'-hydroperoxide: 5-(3-Hydroperoxy-3-methyl-1-butenyl)-4,6,7-trimethoxyfuro[2,3-b]quinoline [953812-12-9]


C₁₉H₂₁NO₆ 359.378

Alkaloid from the leaves of *Choisya ternata*. Cryst. (EtOAc/hexane). Mp 138-139°.

Boyd, D.R. *et al.*, *Org. Biomol. Chem.*, 2007, 5, 2983-2991 (isol, pmr, cmr)

(4-Hydroxy-3-methyl-2-butenyl)guanidine H-579

4-Hydroxygalegine [1557-67-1]



C₆H₁₃N₃O 143.188

(E)-form [26806-04-2]
 Mp 196° (as flavianate).


(Z)-form [26806-06-4]
 Constit. of *Galega officinalis*. Mp 149-151° (as picrate).

Pufahl, K. *et al.*, *Experientia*, 1961, 17, 302 (isol)

Olomucki, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1965, 260, 4519 (synth)

Leonard, N.J. *et al.*, *Chem. Comm.*, 1972, 133 (synth)

5-(2-Hydroxy-3-methyl-3-butenyl)-4,6,7-trimethoxyfuro[2,3-b]quinoline H-580



C₁₉H₂₁NO₅ 343.379

(±)-form [953812-14-1]

Alkaloid from the leaves of *Choisya ternata*. Cryst. (CHCl₃). Mp 95°.

2'-Hydroperoxide: 5-(2-Hydroperoxy-3-methyl-3-butenyl)-4,6,7-trimethoxyfuro[2,3-b]quinoline [953812-10-7]


C₁₉H₂₁NO₆ 359.378

Alkaloid from the leaves of *Choisya ternata*. Cryst. (CH₂Cl₂). Mp 169-170°.

Boyd, D.R. *et al.*, *Org. Biomol. Chem.*, 2007, 5, 2983-2991 (isol, pmr, cmr)

4-(2-Hydroxy-3-methyl-3-butenyl)yukocitrine H-581

[176391-57-4]




C₂₄H₂₅NO₅ 407.465

Alkaloid from stem bark of *Bosistoa transversa*. Yellow oil. [α]_D +16.4 (c, 0.0005 in CHCl₃).

Auzi, A.A. *et al.*, *Phytochemistry*, 1996, 42, 235 (isol, uv, ir, pmr, ms, struct)

5-Hydroxymethylcanthin-6-one H-582

5-(Hydroxymethyl)-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI [89915-37-7]



C₁₅H₁₀N₂O₂ 250.256

Alkaloid from the root bark of *Ailanthus altissima* (Simaroubaceae). Needles (MeOH). Mp 246-247° dec.

Ac:

Needles (Me₂CO). Mp 154°.

9-Methoxy: 5-(Hydroxymethyl)-9-methoxy-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI. 5-Hydroxymethyl-9-methoxycanthin-6-one

C₁₆H₁₂N₂O₃ 280.282


Alkaloid from the roots of *Eurycoma longifolia*. Yellow powder (MeOH). Mp 235° dec. λ_{max} 263 (log ε 3.24); 273 (log ε 3.35); 299 (log ε 2.99); 309 (log ε 3.03); 346 (log ε 3.02); 376 (log ε 2.83) (MeOH).

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984, 32, 170 (isol, uv, ir, pmr, ms, struct)

Kuo, P.-C. *et al.*, *J. Nat. Prod.*, 2003, 66, 1324-1327 (9-methoxy)

1-Hydroxy-3-methyl-9H-carbazole H-583

3-Methyl-9H-carbazol-1-ol [14960-81-7]



C₁₃H₁₁NO 197.236

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (EtOH aq.). Mp 157-158°.

Me ether: 1-Methoxy-3-methyl-9H-carbazole, 9CI. *Murrayafoline A* [4532-33-6]

C₁₄H₁₃NO 211.263

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Cytotoxic agent. Plates (hexane). Mp 52-54°.

Me ether, picrate:

Brown needles (C₆H₆). Mp 188-190°.

Benzyl ether: [220985-64-8]

C₂₀H₁₇NO 287.36

Oil.

- Chakraborty, D.P. *et al.*, *J.O.C.*, 1968, **33**, 1265 (synth)
 Wu, T.-S. *et al.*, *Heterocycles*, 1983, **20**, 1267 (occur, pmr)
 Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4132 (isol, uv, ir, pmr, cmr, ms)
 Martin, T. *et al.*, *J.C.S. Perkin 1*, 1988, 235 (synth, uv, ir, pmr, ms)
 Knölker, H.-J. *et al.*, *Tetrahedron*, 1993, **49**, 11221 (synth)
 Bhattacharyya, P. *et al.*, *Phytochemistry*, 1994, **35**, 1085 (isol, uv, ir, pmr, cmr, struct)
 Murakami, Y. *et al.*, *Heterocycles*, 1998, **49**, 127-132 (*Murrayafoline A*, synth)
 Bringmann, G. *et al.*, *Synthesis*, 1998, 1501-1505 (synth, Me ether, ir, pmr, cmr, ms)
 Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893-897 (*Murrayafoline A*, activity)
 Lin, G. *et al.*, *Tetrahedron*, 2000, **56**, 7163-7171 (synth, pmr, ms, benzyl ether)
 Benavides, A. *et al.*, *Synthesis*, 2004, 2499-2504 (*Murrayafoline A*, synth)
 Mal, D. *et al.*, *Tet. Lett.*, 2006, **47**, 1071-1075 (*Murrayafoline A*, synth)

1-Hydroxy-6-methyl-9H-carbazole H-584

6-Methyl-9H-carbazol-1-ol, 9CI
 [73910-76-6]

C₁₃H₁₁NO 197.236
 Cryst. (C₆H₆). Mp 194-195°.

Me ether: 1-Methoxy-6-methyl-9H-carbazole. *Glycozolicine*
 [73910-79-9]
 [143438-96-4]

C₁₄H₁₃NO 211.263
 Alkaloid from the roots of *Glycosmis pentaphylla*. Cryst. (C₆H₆) or fine needles (hexane). Mp 150° (synthetic) Mp 137-138° (natural). Struct. of nat. prod. revised in 2001 but not fully confirmed.

- Shah, G.D. *et al.*, *Indian J. Chem., Sect. B*, 1979, **18**, 451 (synth, Me ether)
 Patel, B.P.J. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 20-23 (synth)
 Roy, S. *et al.*, *J. Indian Chem. Soc.*, 1982, **59**, 1369-1371 (synth, Me ether)
 Jash, S.S. *et al.*, *Phytochemistry*, 1992, **31**, 2503-2505 (*Glycozolicine*, isol, cmr)
 Chakravarty, A.K. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 484-489 (*Glycozolicine*, synth, struct)

2-Hydroxy-3-methyl-9H-carbazole H-585

3-Methyl-9H-carbazol-2-ol, 9CI
 [24224-30-4]

C₁₃H₁₁NO 197.236
 Alkaloid from the roots of *Murraya koenigii* (curryleaf tree) (Rutaceae) and stem bark of (*Clausena excavata*). Mp 245-246°.

Me ether: 2-Methoxy-3-methyl-9H-carbazole
 [24224-28-0]

C₁₄H₁₃NO 211.263
 Alkaloid from the seeds of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (C₆H₆ or C₆H₆/petrol). Mp 225° (218°).

- Chakraborty, D.P. *et al.*, *Chem. Ind. (London)*, 1969, 1662 (synth, uv, ir)
 Bhattacharyya, P. *et al.*, *Chem. Ind. (London)*, 1986, 246 (isol, uv, ir, pmr, cmr)

- Knölker, H. *et al.*, *J. Indian Chem. Soc.*, 1994, **71**, 345 (synth, 2-Methoxy-3-methylcarbazole)
 Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 133-140 (isol)
 Knölker, H.J. *et al.*, *Synlett*, 1996, 737 (synth)

2-Hydroxy-7-methyl-9H-carbazole H-586

7-Methyl-9H-carbazol-2-ol, 9CI
 [151985-61-4]

C₁₃H₁₁NO 197.236
 Alkaloid from aerial parts of *Cimicifuga simplex* (Ranunculaceae). Pale yellow amorph. powder (CHCl₃/MeOH). Mp 269-271°.

Ac:
 C₁₅H₁₃NO₂ 239.273
 Needles (EtOAc). Mp 245-247°.

Kusano, G. *et al.*, *Heterocycles*, 1993, **36**, 2367 (isol, uv, ir, pmr, cmr, ms, struct)

3-Hydroxy-6-methyl-9H-carbazole H-587

6-Methyl-9H-carbazol-3-ol, 9CI. *Glycozolinine*. *Glycozolinol*
 [5257-08-9]

C₁₃H₁₁NO 197.236
 Alkaloid from the seeds of *Glycosmis pentaphylla* (Rutaceae). Cryst. (C₆H₆). Mp 231-232°.

O-Ac:
 C₁₅H₁₃NO₂ 239.273
 Cryst. (C₆H₆). Mp 210°.

Me ether: 3-Methoxy-6-methyl-9H-carbazole, 9CI. *Glycozoline*
 [5234-30-0]
 C₁₄H₁₃NO 211.263

Alkaloid from the roots of *Glycosmis mauritiana* and the root bark of *Glycosmis pentaphylla* (Rutaceae). Also from *Murraya koenigii*. Mp 181-182°. λ_{max} 227 (ε 33000); 252 (ε 14500); 264 (ε 40000); 304 (ε 14700) (EtOH) (Berdy).

- Me ether, picrate: Mp 182°.
 Carruthers, W. *et al.*, *Chem. Comm.*, 1966, 272 (synth, uv, pmr)
 Chakraborty, D.P. *et al.*, *Tet. Lett.*, 1966, 661 (uv, ir, pmr, struct)
 Bhattacharyya, P. *et al.*, *J. Indian Chem. Soc.*, 1976, **53**, 321 (synth)
 Rastogi, K. *et al.*, *Phytochemistry*, 1980, **19**, 945 (isol)
 Mukherjee, S. *et al.*, *Phytochemistry*, 1983, **22**, 1064 (isol, uv, ir, pmr, struct, *Glycozolinine*)
 Bhattacharyya, P. *et al.*, *Indian J. Chem., Sect. B*, 1984, **23**, 49 (isol, uv, ir, pmr, synth, *Glycozolinine*)
 Iwao, M. *et al.*, *Heterocycles*, 1993, **36**, 1483 (synth, *Glycozolinine*)

4-Hydroxy-6-methyl-9H-carbazole H-588

6-Methyl-9H-carbazol-4-ol
 C₁₃H₁₁NO 197.236


Me ether: 5-Methoxy-3-methyl-9H-carbazole, 9CI. *Glycoborine*
 [359865-27-3]
 C₁₄H₁₃NO 211.263

Alkaloid from roots of *Glycosmis arborea*. Fine needles (petrol/hexane). Mp 155-156°. Struct. formerly as-

- signed to *Glycozolicine*.
 Chakravarty, A.K. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 484-489 (isol, synth, pmr, cmr)

3-Hydroxy-6-methyl-9H-carbazole-1-carboxaldehyde H-589

1-Formyl-3-hydroxy-6-methylcarbazole




- C₁₄H₁₁NO₂ 225.246
 Me ether: 3-Methoxy-6-methyl-9H-carbazole-1-carboxaldehyde. 1-Formyl-3-methoxy-6-methylcarbazole
 [356549-34-3]
 C₁₅H₁₃NO₂ 239.273
 Alkaloid from the leaves of *Murraya koenigii*. Light yellow cryst. Mp 181-182°. λ_{max} 227 (log ε 4.78); 262 (log ε 4.1); 303 (log ε 4.41); 398 (log ε 3.87) (MeOH).

Chowdhury, B.K. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 490-494 (isol, synth, pmr)

1-Hydroxymethyl-β-carboline H-590

9H-Pyrido[3,4-b]indole-1-methanol, 9CI
 [17337-22-3]




- C₁₂H₁₀N₂O 198.224
 Alkaloid from the stems of *Picrasma ailanthoides* and the wood of *Picrasma quassioides* (Simaroubaceae). Yellow needles (MeOH). Mp 228-230° dec.

Hydrochloride:
 Yellow needles (MeOH). Mp 240° dec.

- O-Ac:
 Pale-yellow needles (Et₂O). Mp 113-113.5°.
 Kondo, Y. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 837 (isol, uv, pmr, struct, synth)
 Koike, K. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 471 (cmr)
 Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3579 (isol, uv, ir, synth)

6-Hydroxy-1-methyl-β-carboline H-591

1-Methyl-9H-pyrido[3,4-b]indol-6-ol, 9CI. 6-Hydroxyharman
 [67767-19-5]



- C₁₂H₁₀N₂O 198.224
 Alkaloid from the roots of *Grewia mollis*

and from *Ophiorrhiza japonica*. Also from the bryozoan *Cribricellina cribraria*. Pale-yellow prisms. Mp 276-278°.

Hydrochloride:

Cryst. (MeOH). Mp 320° dec.

Me ether: 6-Methoxy-1-methyl-β-carboline. 6-Methoxyharman. **Isoharminine.**

Coharminine

[3589-72-8]

C₁₃H₁₂N₂O 212.251

Alkaloid from *Mucuna pruriens* and leaves and stems of *Virola cuspidata* (Myristicaceae, Fabaceae). Also isol. from *Peganum harmala* (Zygophyllaceae). Prisms (MeOH). Mp 275-276° dec. λ_{max} 232 (ε 33800); 246 (sh) (ε 29400); 257 (sh) (ε 21900); 285 (sh) (ε 15000); 290 (ε 15600); 296 (ε 25000); 354 (ε 7500) (EtOH).

▶ UV0168000

3,4-Dihydro, Me ether: 3,4-Dihydro-6-methoxy-1-methyl-β-carboline. 6-Methoxyharmalan

[3589-73-9]

C₁₃H₁₄N₂O 214.266

Alkaloid from the leaves and stems of *Virola cuspidata* (Myristicaceae).

▶ UU9802000

Cook, J.W. et al., *J.C.S.*, 1951, 1203-1207 (synth)

Cassady, J.M. et al., *J. Nat. Prod.*, 1971, **34**, 161-162 (6-Methoxyharman, 6-Methoxyharmalan)

Ghosal, S. et al., *Planta Med.*, 1972, **21**, 200-203 (6-Methoxyharman, occur)

Rosler, H. et al., *J. Nat. Prod.*, 1978, **41**, 383-384 (isol, uv, ir, pmr, ms, struct)

Aimi, N. et al., *Chem. Pharm. Bull.*, 1986, **34**, 3064-3066 (isol)

Ayoub, M.T. et al., *Phytochemistry*, 1991, **30**, 1046-1047 (Isoharminine)

Rocca, P. et al., *Tet. Lett.*, 1994, **35**, 2003-2004 (synth)

Harwood, D.T. et al., *Nat. Prod. Res.*, 2003, **17**, 15-19 (isol, pmr, cmr)

7-Hydroxy-1-methyl-β-carboline H-592

1-Methyl-9H-pyrido[3,4-b]indol-7-ol, 9CI. **Harmol.** 7-Hydroxyharman

[487-03-6]

C₁₂H₁₀N₂O 198.224

Alkaloid from *Carex brevicollis*, *Elaeagnus angustifolia* (Russian olive), *Hippophae rhamnoides*, *Banisteriopsis caapi*, *Banisteriopsis inebrians*, *Peganum harmala*, *Tribulus terrestris*, *Zygophyllum fabago* and *Passiflora incarnata* (maypops) (Cyperaceae, Eleagnaceae, Malphigiaceae, Zygophyllaceae, Passifloraceae) and the watermoss *Fontinalis squamosa*. Shows antiparasitic activity. Active against gram-positive bacteria and fungi. Sol. CHCl₃, Me₂CO, bases; poorly sol. H₂O. Mp 304-307°. λ_{max} 210; 240; 300; 325 (MeOH) (Berdy).

O-Ac: [257938-78-6]

C₁₄H₁₂N₂O₂ 240.261

Needles (EtOH). Mp 155°.

Me ether: 7-Methoxy-1-methyl-9H-pyrido[3,4-b]indole. 7-Methoxy-1-methyl-β-carboline. **Harmine.** Yageine. **Banisterine.** Telepathine

[442-51-3]

C₁₃H₁₂N₂O 212.251

Alkaloid from *Peganum harmala*, several *Banisteriopsis* spp., *Passiflora edulis* (passionfruit) and several other spp. (Zygophyllaceae, Malphigiaceae, Passifloraceae). Acid-base fluorescence indicator (pH 7.2-8.9, colour change blue → yellow). Antiparkinsonian agent. CNS stimulant. Component of a South American hallucinogenic drink. Active against gram-positive bacteria and fungi. Cryst. Mp 264-265° (257-259°). pK_a 7.61 (20°). Log P 3.12 (calc). λ_{max} 210; 240; 300; 338 (MeOH) (Berdy). λ_{max} 298 (log ε 4.21); 324 (log ε 3.72); 332 (log ε 3.68) (MeCN). λ_{max} 242 (log ε 4.62); 301 (log ε 4.2) (EtOH).

▶ Adverse gastrointestinal and CNS effects in humans by intramuscular route. LD₅₀ (mus, scu) 243 mg/kg; LD₅₀ (mus, ivn) 38 mg/kg. UV0175000

Me ether, hydrochloride: [343-27-1]

Dihydrate. Mp 268-270° dec Mp 321° (anhyd.).

▶ MG9450000

Me ether, picrate: Mp 249-250° dec.

Me ether, N²-oxide: Harmine N-oxide

[57498-78-9]

C₁₃H₁₂N₂O₂ 228.25

Alkaloid from *Banisteriopsis caapi*

(Malphigiaceae). Needles (MeOH).

Mp 226-227° dec. λ_{max} 213 (log ε 4.45);

248 (log ε 4.72); 330 (log ε 4.62)

(MeOH).

3,4-Dihydro: 4,9-Dihydro-1-methyl-3H-pyrido[3,4-b]indol-7-ol. 3,4-Dihydro-7-hydroxy-1-methyl-β-carboline. **Harmalol**

[525-57-5]

[6028-07-5]

C₁₂H₁₂N₂O 200.24

Alkaloid from *Peganum harmala*, *Passiflora incarnata* (maypops), *Amsonia tabernaemontana*, *Apocynum cannabinum* and *Hippophae rhamnoides* (Zygophyllaceae, Passifloraceae, Apocynaceae, Eleagnaceae). Traditional dye for wool. CNS stimulant. Active against gram-positive bacteria and fungi. Fine orange-yellow or red needles + 3H₂O (EtOH aq.). Sol. Me₂CO, CHCl₃; poorly sol. H₂O. Mp 100-105°. Log P 1.31 (uncertain value) (calc). Readily oxid. in air. λ_{max} 218; 260; 376 (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 175 mg/kg.

3,4-Dihydro, hydrochloride: [6028-00-8]

Green cryst. (EtOH), red needles

(EtOH). Mp 264-267° dec.

3,4-Dihydro, Me ether: 4,9-Dihydro-7-methoxy-1-methyl-3H-pyrido[3,4-b]indole. 3,4-Dihydro-7-methoxy-1-methyl-β-carboline. **Harmaline.** Dihydroharmine. **Harmidine**

[304-21-2]

C₁₃H₁₄N₂O 214.266

Alkaloid from *Peganum harmala*, *Passiflora incarnata*, *Banisteria caapi* (preferred genus name *Heteropterys*) and some *Banisteriopsis* spp. (Zygophyllaceae, Passifloraceae, Malphigia-

ceae). CNS stimulant. Possesses antiparkinsonian props. Component of a S. American hallucinogenic drink. Active against gram-positive bacteria and fungi. Sol. Et₂O. Mp 250-251° (227-229°). Log P 1.91 (uncertain value) (calc). λ_{max} 218 (ε 17700); 260 (ε 7950); 376 (ε 10050) (MeOH) (Berdy). λ_{max} 260 (log ε 4.07); 380 (log ε 4.49) (0.1M HCl). λ_{max} 300 (log ε 4.24); 324 (log ε 3.8); 336 (log ε 3.73) (MeOH). λ_{max} 216; 228; 261; 341; 352; 380 (MeOH).

▶ LD₅₀ (rat, scu) 120 mg/kg. UU9800000

3,4-Dihydro, Me ether, hydrochloride: [6027-98-1]

Mp 212° Mp 239-240° (dihydrate).

▶ MG8750000

3,4-Dihydro, Me ether, N-Ac: Mp 204-205°.

1,2,3,4-Tetrahydro: 2,3,4,9-Tetrahydro-1-methyl-1H-pyrido[3,4-b]indol-7-ol, 8CI. 1,2,3,4-Tetrahydro-7-hydroxy-1-methyl-β-carboline. **Tetrahydroharmol**

[17952-75-9]

C₁₂H₁₄N₂O 202.255

Alkaloid from *Elaeagnus angustifolia* (Russian olive), *Shepherdia argentea* and *Shepherdia canadensis* (Eleagnaceae). Mp 254-255°.

1,2,3,4-Tetrahydro, N²,O-di-Ac: [28090-84-8]

C₁₆H₁₈N₂O₃ 286.33

Cryst. (EtOAc). Mp 202°.

1,2,3,4-Tetrahydro, N²-Me: N^b-Methyltetrahydroharmol

[57116-40-2]

C₁₃H₁₆N₂O 216.282

Alkaloid from *Elaeagnus angustifolia* (Russian olive) (Eleagnaceae). Cryst. (EtOH). Mp 268-270°. Racemic.

[28090-85-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 682C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 166B; 166C (nmr)

Späth, E. et al., *Ber.*, 1930, **63**, 120-125 (*Harmine, Harmaline, synth*)

Doig, G.G. et al., *J.C.S.*, 1952, 3912-3916 (*Harmaline, uv*)

Spenser, I.D. et al., *Can. J. Chem.*, 1959, **37**, 1851-1858 (*Harmaline, synth, uv, ir*)

Lutowski, J. et al., *CA*, 1961, **55**, 21479a (*Harmol, isol*)

Robinson, B. et al., *Chem. Ind. (London)*, 1965, 605 (*Harmaline, pmr*)

Ayer, W.A. et al., *Can. J. Chem.*, 1970, **48**, 1980-1984 (*Tetrahydroharmol*)

Ribas, I. et al., *CA*, 1972, **77**, 123811n (*isol*)

Bishop, E. et al., *Indicators*, 1972, 695 (*Harmine, use*)

Phillipson, J.D. et al., *J. Chromatogr.*, 1975, **405**, 163 (*Harmaline, uv, tlc, glc*)

Hashimoto, Y. et al., *Phytochemistry*, 1975, **14**, 1633-1635 (*Harmine N-oxide*)

Shoemaker, D.W. et al., *J. Chromatogr.*, 1979, **174**, 159-164 (*Harmine, Harmol, glc, ms*)

Ross, S.A. et al., *Fitoterapia*, 1980, **6**, 309-312 (*Harmalol, Harmaline, activity*)

Allen, J.R.F. et al., *Phytochemistry*, 1980, **19**, 1573-1582 (*rev, bibl*)

Coune, C.A. et al., *Phytochemistry*, 1980, **19**, 2009-2011 (*Harmalol, Harmaline, cmr*)

Al-Shamma, A. et al., *J. Nat. Prod.*, 1981, **44**, 745-747 (*Harmine, activity*)

- Verpoorte, R. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 328-335 (*Harmaline, cmr*)
 Ahmad, A. *et al.*, *J. Ethnopharmacol.*, 1992, **35**, 289-294 (*Harmine, activity*)
 Biondic, M.C. *et al.*, *J.C.S. Perkin 2*, 1992, 1049-1058 (*Harmaline, uv*)
 Salm, R.F. *et al.*, *Phytochemistry*, 1998, **49**, 887-892 (*isol*)
 Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1587 (*Harmine, Harmaline*)
 Ponce, M.A. *et al.*, *J. Het. Chem.*, 2001, **38**, 1071-1082 (*O-Ac*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, HA1500 (*Harmine*)


8-Hydroxy-1-methyl-β-carboline H-593

1-Methyl-9H-pyrido[3,4-b]indol-8-ol, 9CI. **8-Hydroxyharman**
 [521305-60-2]

C₁₂H₁₀N₂O 198.224
 Alkaloid from the bryozoan *Cribricellina cribraria*. Yellow oil. λ_{max} 216 (ε 4000); 242 (ε 6500); 277 (ε 1300); 286 (ε 1500); 340 (ε 800) (EtOH).

Harwood, D.T. *et al.*, *Nat. Prod. Res.*, 2003, **17**, 15-19 (*isol, pmr, cmr, ms*)

9-Hydroxy-3-methyl-2-decenoic acid H-594



C₁₁H₂₀O₃ 200.277

(2Z,9R)-form

O-(15*Z*-Docosenoyl), 2-sulfoethyl ester: **Irciniasulfonic acid A₁**
 C₃₅H₆₄O₇S 628.952
 Isol. from the sponge *Ircinia* sp. Multidrug resistance modulator. Isol. as a mixt. with other Irciniasulfonic acids A, to which data refers. λ_{max} 220 (ε 6100) (MeOH).

O-(15*Z*-Docosenoyl), 2-sulfoethylamide: **Irciniasulfonic acid B₁**
 [901764-82-7 (Irciniasulfonic acid B)]
 C₃₅H₆₅NO₆S 627.968
 Isol. from *Ircinia* sp. Isol. as a mixt. with B₂.

O-Tricosanoyl, 2-sulfoethyl ester: **Irciniasulfonic acid A₃**
 C₃₆H₆₈O₇S 644.995
 Isol. from *Ircinia* sp.

O-(5*Z*,9*Z*-Tetracosadienoyl), 2-sulfoethyl ester: **Irciniasulfonic acid A₂**
 C₃₇H₆₆O₇S 654.99
 Isol. from *Ircinia* sp.

O-(5*Z*,9*Z*-Tetracosadienoyl), 2-sulfoethylamide: **Irciniasulfonic acid B₂**
 [901764-82-7 (Irciniasulfonic acid B)]
 C₃₇H₆₇NO₆S 654.005
 Isol. from *Ircinia* sp. Isol. as a mixt. with B₁.


Kawakami, A. *et al.*, *Tet. Lett.*, 2001, **42**, 3335-3337 (*isol, pmr, cmr*)

Dobbs, A.P. *et al.*, *Synlett*, 2005, 652-654 (*synth*)

Emura, C. *et al.*, *Tetrahedron*, 2006, **62**, 5682-5685 (*Irciniasulfonic acid B*)

8-Hydroxymethyl-dihydroinditidine H-595

6-Hydroxymethyl-dihydroinditidine
 [96627-11-1]




C₂₂H₂₁NO₅ 379.412

Minor alkaloid from the stem bark of *Fagaropsis angolensis* (Rutaceae). Yellow gum.

Khalid, S.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 118 (*isol, uv, ir, pmr, ms, struct*)

8-Hydroxymethyl-dihydroguanine H-596

13,14-Dihydro-13-methyl[1,3]benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridine-14-methanol, 9CI
 [67951-20-6]
 [88588-10-7]




C₂₁H₁₇NO₅ 363.369

Possibly an artifact. Alkaloid from the roots of *Chelidonium japonicum* (Papaveraceae) and aerial parts of *Fumaria vaillantii*. Cryst. (CHCl₃/EtOH). Mp 249-251.5°. λ_{max} 212 (log ε 4.24); 235 (log ε 4.39); 284 (log ε 4.42); 322 (log ε 4.02); 350 (sh) (log ε 3.54) (MeOH).

Itokawa, H. *et al.*, *Phytochemistry*, 1978, **17**, 839-840 (*isol, uv, ir, pmr, ms, struct*)

2-(10-Hydroxy-10-methyldecyl)-3-methoxy-4(1H)-quinolinone H-597



C₂₃H₃₅NO₃ 373.534

(+)-form

Alkaloid from the leaves of *Spathelia excelsa*. Oil. [α]_D +5.4 (c, 0.02 in CHCl₃). λ_{max} 241; 248; 325; 334 (MeOH).

Da Paz Lima, M. *et al.*, *Phytochemistry*, 2005, **66**, 1560-1566 (*isol, pmr, cmr*)

2-(11-Hydroxy-11-methyldecyl)-3-methoxy-4(1H)-quinolinone H-598


C₂₃H₃₅NO₃ 373.534

Alkaloid from the leaves of *Spathelia excelsa*. Oil. λ_{max} 241; 248; 326; 336 (MeOH).

Da Paz Lima, M. *et al.*, *Phytochemistry*, 2005, **66**, 1560-1566 (*isol, pmr, cmr*)

3-Hydroxy-8,9-methylenedioxybenzo[h]quinoline H-599

[1,3]Benzodioxolo[5,6-h]quinolin-3-ol, 9CI. **Toddaquinoline**
 [152406-73-0]



C₁₄H₉NO₃ 239.23

Alkaloid from the root bark of *Toddalia asiatica* (Rutaceae). Prisms (MeOH/Et₂O). Mp 235-237°.

Ac:

Needles (CHCl₃/MeOH). Mp 174-176°.

Me ether:


Prisms (MeOH). Mp 145-148°.

Chen, I.-S. *et al.*, *Phytochemistry*, 1993, **34**, 1449 (*isol, uv, ir, pmr, struct*)

Harrowven, D.C. *et al.*, *Tetrahedron*, 2001, **57**, 4447-4454 (*synth*)

6-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid H-600

10-Hydroxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. **Aristolochic acid C**. **Aristolochic acid IIIa**
 [4849-90-5]



C₁₆H₉NO₇ 327.25

Alkaloid from *Aristolochia debilis* and from the tubers of *Aristolochia rotunda*. An active principle of "Tong Cheng Hu Geng" (roots of *Aristolochia tagala*) and of the Chinese drug Fang-chi. Also found in the butterfly *Zerynthia polyxena*. Cytotoxic. Needles + 5H₂O. Mp 280° dec.

Me ester: Mp 254°.

6-O-β-D-Glucopyranoside:

C₂₂H₁₉NO₁₂ 489.392

Alkaloid from roots of *Aristolochia cinnabarina* (Aristolochiaceae). Yellow amorph. powder. [α]_D²⁵ -46.6 (c, 0.09 in MeOH).

Me ether: **Aristolochic acid III**. *10-Methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid*, 9CI. *6-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid*
 [7267-92-7]

C₁₇H₁₁NO₇ 341.276

Alkaloid from the roots of *Aristolochia argentina*, *Aristolochia esperanzae*, *Aristolochia clematitidis* and *Aristolochia longa* (Aristolochiaceae).

Me ether, Me ester: Aristolochic acid III methyl ester

C₁₈H₁₃NO₇ 355.303

Alkaloid from *Aristolochia cucurbitifolia*. Yellowish needles. Mp 272-274° (265-266°). λ_{max} 216 (log ε 4.06); 256 (log ε 4.33); 278 (sh) (log ε 3.91); 300 (log ε 3.81); 350 (log ε 3.63); 384 (log ε 3.59) (MeOH).

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 973; 1470; *CA*, **53**, 21841e; **54**, 6688i (*isol. ir. uv*)

Sasagawa, S. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 921; *CA*, **59**, 1552g (*struct*)

Pailer, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 863 (*isol. deriv*)

Carboni, S. *et al.*, *Gazz. Chim. Ital.*, 1966, **96**, 641 (*isol. ir*)

Priestap, H.A. *et al.*, *An. Asoc. Quim. Argent.*, 1971, **59**, 245; *CA*, **76**, 43943w (*isol. deriv*)

Rothschild, M. *et al.*, *Insect Biochem.*, 1972, **2**, 334; *CA*, **77**, 137607n (*isol. ms*)

Chen, Z.-L. *et al.*, *Huaxue Xuebao*, 1981, **39**, 237; *CA*, **95**, 156428t (*isol*)

Ding, L. *et al.*, *Zhongcaoyao*, 1981, **12**, 436; *CA*, **96**, 214267y (*isol*)

De Pascual Teresa, J. *et al.*, *Phytochemistry*, 1983, **22**, 2745 (*ir, pmr, ms*)

Priestap, H.A. *et al.*, *Magn. Reson. Chem.*, 1989, **27**, 460 (*cmr*)


Hong, L. *et al.*, *Phytochemistry*, 1994, **37**, 237 (*glucoside*)

Leu, Y.-L. *et al.*, *Phytochemistry*, 1998, **48**, 743-745 (*isol. uv, pmr, ms*)

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1006-1009 (*Aristolochic acid III Me ester, activity*)

7-Hydroxy-1-methylenepyrrolizidine H-601

2,3,5,6,7,7a-Hexahydro-7-methylene-1H-pyrrolizin-1-ol, 8CI



(7R,7aR)-form

C₈H₁₃NO 139.197

The 7a posn. acc. to CAS numbering is often numbered 8.

(7R,7aR)-form

Minor alkaloid from *Crotalaria goreensis* (Fabaceae). Mp 35-36°. Bp_{0.1} 41°. [α]_D¹⁹ -150 (c, 1.2 in EtOH). pK_a 9.5.

Picrate:

Needles. Mp 203.5-204°. [α]_D -47.5 (c, 0.68 in Me₂CO).

Angeloyl: 7-Angeloyl-1-methylenepyrrolizidine

C₁₃H₁₉NO₂ 221.299

Alkaloid from *Senecio chrysocoma* (Asteraceae). Gum.

N-Oxide: 7-Hydroxy-1-methylenepyrrolizidine N-oxide

C₈H₁₃NO₂ 155.196

Alkaloid from epigeal parts of *Senecio schweinfurthii* (Asteraceae). Oil.

(7R,7aS)-form [2520-33-4]

Major alkaloid from *Crotalaria goreensis*, also *isol.* from *Crotalaria podocarpa*, *Crotalaria maypurensis* and *Crotalaria aegyptiaca* (Fabaceae). Needles (petrol). Mp 33-35°. Bp_{0.03} 62°. [α]_D¹⁸ +36.1 (c, 1.39

in EtOH). pK_a 9.3.

Picrate:

Yellow needles (EtOH). Mp 173.5-174.5°. [α]_D¹⁸ -24.2 (c, 1.01 in Me₂CO).

(7S,7aR)-form

Angeloyl:

C₁₃H₁₉NO₂ 221.299

Alkaloid from *Senecio chrysocoma* (Asteraceae). Gum.

Angeloyl, N-oxide: 7-Angeloyl-1-methylenepyrrolizidine N-oxide

C₁₃H₁₉NO₃ 237.298

Alkaloid from *Senecio chrysocoma* (Asteraceae). Gum.

[6257-42-7]

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1961, **14**, 284 (*isol. struct, abs config*)

Robins, D.J. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 115 (*occur*)


Liddell, J.R. *et al.*, *Phytochemistry*, 1993, **34**, 1198 (*derivs*)

Benn, M.H. *et al.*, *Phytochemistry*, 1995, **40**, 1327 (*oxide*)

Logie, C.G. *et al.*, *S. Afr. J. Chem.*, 1997, **50**, 72-74 (*synth, abs config*)

2-Hydroxymethyl-5-(1-hydroxypentyl)-3,4-pyrrolidinediol H-602

3,4-Dihydroxy-2-hydroxymethyl-5-(1-hydroxypentyl)pyrrolidine



Relative Configuration

C₁₀H₂₁NO₄ 219.28

(2R,3R,4R,5R,6S)-form

Alkaloid from *Adenophora triphylla* var. *japonica*. Inhibitor of almond β-glucosidase. [α]_D +174.3 (c, 0.32 in H₂O).


Asano, N. *et al.*, *Phytochemistry*, 2000, **53**, 379-382

3-Hydroxymethyl-1-methoxycarbazole H-603

1-Methoxy-9H-carbazole-3-methanol,

9CI. *Koehnline*

[3909-78-2]



C₁₄H₁₃NO₂ 227.262

Alkaloid from the root bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Exhibits cytotoxic activity vs. KB cells.

Cryst. (C₆H₆/petrol). Mp 142° (130°). Partially converted to Murrayanine (see 1-Hydroxy-9H-carbazole-3-carboxaldehyde, H-447) on standing at r.t.

O-Ac:

Cryst. (Py aq.). Mp 110°.

Chakraborty, D.P. *et al.*, *Tetrahedron*, 1965, **21**, 681 (*synth*)

Fiebig, M. *et al.*, *Phytochemistry*, 1985, **24**, 3041 (*isol. uv, ir, pmr, cmr, ms*)

Knölker, H.-J. *et al.*, *Tetrahedron*, 1993, **49**, 11221 (*synth*)

Bringmann, G. *et al.*, *Synthesis*, 1998, 1501-1505 (*synth, ms*)


Knölker, H.-J. *et al.*, *Chem. Rev.*, 2002, **102**, 4303-4427 (*rev*)

6-Hydroxymethyl-1-methoxycarbazole H-604

8-Methoxy-9H-carbazole-3-methanol,

9CI. *Mukoline*

[87264-41-3]



C₁₄H₁₃NO₂ 227.262

Minor alkaloid from roots of *Murraya koenigii* (curryleaf tree). Cryst. (C₆H₆). Mp 118°.

Picrate:

Cryst. (C₆H₆). Mp 230°.

O-Ac:


Cryst. (C₆H₆/petrol). Mp 145°.

N-Me:

Cryst. (C₆H₆/petrol). Mp 140°.

Roy, S. *et al.*, *J. Indian Chem. Soc.*, 1982, **59**, 1369 (*isol. uv, ir, pmr, ms, struct*)

N-(1-Hydroxymethyl-2-methoxyethyl)-7-methoxy-4-eicosanamide H-605



C₂₅H₄₉NO₄ 427.666

(1'R,4E,7S)-form [253443-61-7]

Isol. from an Australian cyanobacterium. Plates. Mp 32-34°. [α]_D²⁰ -3 (c, 0.33 in CHCl₃).

2'-Ac: N-(1-Acetoxyethyl-2-methoxyethyl)-7-methoxy-4-eicosanamide [253443-58-2]


C₂₇H₅₁NO₅ 469.704

Isol. from an Australian cyanobacterium. Needles. Mp 39-40.5°. [α]_D²⁰ -6.1 (c, 0.58 in CHCl₃).

Wan, F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1696-1699 (*isol. ir, pmr, cmr*)

3-(Hydroxymethyl)-4-(1-methyl-1H-imidazol-5-yl)-1-phenyl-1-butanone H-606

[38993-91-8]



C₁₅H₁₈N₂O₂ 258.319

Alkaloid from *Pilocarpus microphyllus*.

11-Deoxy, 7,11-didehydro: 3-Methylene-4-(1-methyl-1H-imidazol-5-yl)-1-phenyl-1-butanone


[942060-97-1]

C₁₅H₁₆N₂O 240.304Alkaloid from *Pilocarpus microphyllus*.


Abreu, I.N. *et al.*, *Rapid Commun. Mass Spectrom.*, 2007, **21**, 1205-1213 (isol, ms)
 Sawaya, A.C.H.F. *et al.*, *Molecules*, 2008, **13**, 1518-1529 (ms, chromatog)

2-(Hydroxymethyl)-6-methyl-3,4,5-piperidinetriol H-607

1,2,6-Trideoxy-2,6-iminoheptitol

(2*R*,3*R*,4*R*,5*R*,6*R*)-formC₇H₁₅NO₄ 177.2**(2*R*,3*R*,4*R*,5*R*,6*R*)-form**7-Deoxy- α -homomannonojirimycinAlkaloid from the bulbs of *Scilla sibirica*. Syrup. [α]_D -11.1 (c, 0.44 in H₂O).**(2*R*,3*R*,4*R*,5*R*,6*S*)-form**7-Deoxy- β -homomannonojirimycin. β -HomofuconojirimycinAlkaloid from *Angylocalyx pynaertii*, *Scilla sibirica* and *Scilla socialis*. Inhibitor of α -fucosidases. [α]_D -21.5 (c, 0.73 in H₂O).**(2*R*,3*R*,4*R*,5*S*,6*R*)-form**7-Deoxy- α -homonojirimycinAlkaloid from the bulbs of *Scilla sibirica* and *Scilla socialis*. Inhibitor of α -glucosidase. Powder. [α]_D +78.9 (c, 0.46 in H₂O).Asano, N. *et al.*, *Eur. J. Biochem.*, 2001, **268**, 35-41 (isol, pmr, cmr)Yamashita, T. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1875-1881 (isol, pmr, cmr)**4-(Hydroxymethyl)-6-methyl-2,3,5-pyridinetriol** H-608


2,3,5-Trihydroxy-4-(hydroxymethyl)-6-methylpyridine [95508-58-0]

C₇H₉NO₄ 171.152


Tentative struct. assigned. Prod. by *Streptovorticillum hiroshimensis*. Dopamine β -hydroxylase inhibitor, antihypertensive agent. Cryst. (H₂O). Sol. H₂O; fairly sol. MeOH; poorly sol. hexane. λ_{\max} 276 (HCl) (Berdy). λ_{\max} 227; 304 (NaOH) (Berdy).

U.S. Pat., 1984, 4 487 761; *CA*, **102**, 130458r**2-Hydroxymethyl-4-methyl-quinazoline** H-609

4-Methyl-2-quinazolinemethanol, 9CI [13535-91-6]

C₁₀H₁₀N₂O 174.202Metab. of the bacterium *Pseudomonas aeruginosa*.Mann, S. *et al.*, *Arch. Mikrobiol.*, 1967, **56**, 324**3-Hydroxy-5-methyl-1-naphthalenecarboxylic acid, 9CI** H-610


3-Hydroxy-5-methyl-1-naphthoic acid, 8CI [34581-87-8]

C₁₂H₁₀O₃ 202.209Obt. by alkaline hydrolysis of Carzinoiphilin. Needles (C₆H₆/EtOAc). Mp 246°.*Me ether*, [1-(aminocarbonyl)-2,3-epoxy-2-methylpropyl] ester (1*S*,2*S*-):

[110450-52-7]

C₁₈H₁₉NO₅ 329.352Prod. by *Streptomyces griseofuscus*.Needles (CHCl₃/hexane). Mp 153-154°. [α]_D²⁵ +48 (c, 0.33 in MeOH). A fragment of Azinomycin A. Biol. inactive. λ_{\max} 217 (ϵ 46600); 245 (sh); 303 (ϵ 3300); 343 (ϵ 4900) (MeOH).Yokoi, K. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4554-4561 (*Streptomyces constits*)**4-Hydroxy-2-methyl-6-nonylpiperidine** H-611

2-Methyl-6-nonyl-4-piperidinol, 9CI

C₁₅H₃₁NO 241.416**(2*R**,4*S**,6*S**)-form****Dendrobates Alkaloid 241D**


[120030-06-0]

Alkaloid from skin extracts of the Panamanian poison frog *Dendrobates speciosus*. Also detected as a trace constit. in certain populations of *Dendrobates pumilio*. [α]_D²⁵ +39 (c, 0.2 in MeOH).*Oxo*: **Dendrobates Alkaloid 255**


[120041-46-5]

C₁₅H₂₉NO 239.4Alkaloid from skin extracts of *Dendrobates speciosus*. Tentative struct. assignment. Prob. a side chain oxo deriv. of 241D.Edwards, M.W. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1188 (isol, pmr, ms, struct)Edwards, M.W. *et al.*, *Synthesis*, 1994, 1167 (synth)Ciblat, S. *et al.*, *J.C.S. Perkin 1*, 2000, 353-357 (synth)Girard, N. *et al.*, *Tet. Lett.*, 2007, **48**, 4097-4099 (synth)**3-Hydroxy-6-methyl-2-(11-oxododecyl)pyridine** H-612

12-(3-Hydroxy-6-methyl-2-pyridinyl)-2-dodecanone

C₁₈H₂₉NO₂ 291.433Alkaloid from the flowers of *Senna spectabilis*. Cytotoxic. Oil. λ_{\max} 224 (log ϵ 3.9); 288 (log ϵ 3.8) (MeOH).*N-Oxide*:C₁₈H₂₉NO₃ 307.432Alkaloid from the flowers of *Senna spectabilis*. Cytotoxic. Cryst. Mp 71-72°. λ_{\max} 222 (log ϵ 4); 260 (log ϵ 3.6); 303 (log ϵ 3.2) (MeOH).Sriphong, L. *et al.*, *Planta Med.*, 2003, **69**, 1054-1056 (isol, pmr, cmr, ms)**5-Hydroxy-6-methyl-2-piperidinecarboxylic acid, 9CI** H-613

5-Hydroxy-6-methylpipercolic acid. 2-Carboxy-5-hydroxy-6-methylpiperidine

(2*S*,5*S*,6*S*)-formC₇H₁₃NO₃ 159.185**(2*S*,5*S*,6*S*)-form** [54957-04-9]


Found in seeds of the Fagaceae esp.

Fagus sylvatica. Cryst. (H₂O/EtOH/Et₂O). [α]_D²³ -11 (c, 1.3 in H₂O). [α]_D²³ +1.1 (c, 2.2 in 2*M* HCl).**(2*S*,5*R*,6*S*)-form** [54836-98-5]Found in seeds of *Fagus sylvatica*. Cryst. [α]_D²³ -46.4 (c, 2.8 in H₂O). [α]_D²³ -33.8 (c, 2.8 in 2*M* HCl).*Hydrochloride*:Cryst. (H₂O/EtOH/Et₂O). [α]_D²² -35.6 (c, 1 in H₂O).Kristensen, I. *et al.*, *Tetrahedron*, 1976, **32**,

2799 (isol, struct, ir, nmr, cd)

Kasai, T. *et al.*, *Phytochemistry*, 1978, **17**, 1911 (isol)**2-(Hydroxymethyl)-3,4-piperidinediol, 9CI** H-614

3,4-Dihydroxy-2-(hydroxymethyl)piperidine. 3,4-Dihydroxy-2-piperidinemethanol. 1,2,5-Trideoxy-1,5-iminoheptitol

(2*R*,3*R*,4*R*)-form

C₆H₁₃NO₃ 147.174

Care needed with numbering; if named as an iminosugar the numbering is different.

(2R,3R,4R)-form**Fagomine**

[53185-12-9]

Alkaloid from buckwheat seeds (*Fagopyrum esculentum*) (Polygonaceae). Also from seeds of *Castanospermum australe* and *Xanthocercis zambeziaca* (Fabaceae). Needles (MeOH/EtOH). Mp 186-188°. [α]_D²⁵ +23 (c, 1 in H₂O). [α]_D¹ +37 (c, 1 in 0.1M HCl).

Hydrochloride: Mp 176-177°.

3-O- β -D-Glucopyranosyl: **3-O- β -D-Glucopyranosylfagomine**

[104958-79-4]

[96602-64-1]

C₁₂H₂₃NO₈ 309.316

Alkaloid from the seeds of *Xanthocercis zambeziaca* (Fabaceae). Cryst. (MeOH/EtOH/Me₂CO). Mp 232-233° dec. [α]_D²⁰ -3.1 (c, 1.2 in H₂O). Can also be named as 4-glucopyranosylfagomine (confusion with 4-isomer below).

4-O- β -D-Glucopyranosyl: **4-O- β -D-Glucopyranosylfagomine**C₁₂H₂₃NO₈ 309.316

Alkaloid from leaves and roots of *Xanthocercis zambeziaca*. [α]_D -18.2 (c, 0.48 in H₂O). Can also be numbered as the 3-isomer (see above).

(2R,3R,4S)-form**4-Epifagomine**

[156639-77-9]

Alkaloid from roots of *Morus alba* (white mulberry) (Moraceae) and from leaves and roots of *Xanthocercis zambeziaca* (Fabaceae). Solid. Mp 220-222° synthetic. [α]_D²⁵ +77.15 (c, 0.68 in H₂O). Numbering systems vary.


(2R,3S,4S)-form**3,4-Diepifagomine**

Alkaloid from leaves of *Xanthocercis zambeziaca*. [α]_D -8.7 (c, 0.3 in H₂O) (natural). [α]_D²⁵ +13.4 (c, 0.32 in H₂O) (synthetic). No CAS no. found.

Koyama, M. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 1111 (*ir, pmr, ms, struct*)Evans, S.V. *et al.*, *Tet. Lett.*, 1985, **26**, 1465 (3-O- β -D-glucopyranosyl, *pmr, struct*)Fleet, G.W.J. *et al.*, *Tet. Lett.*, 1985, **26**, 1469 (*abs config, synth*)Molyneux, R.J. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1198-1206 (*isol*)Asano, N. *et al.*, *Carbohydr. Res.*, 1994, **259**, 243 (3-Epifagomine)Kato, A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 312-314 (*glucoside, diepimer*)Schneider, C. *et al.*, *Eur. J. Org. Chem.*, 1998, 1155-1159 (*synth*)Banba, Y. *et al.*, *Tetrahedron: Asymmetry*, 2001, **12**, 817-819 (*synth*)Takahata, H. *et al.*, *J.O.C.*, 2003, **68**, 3603-3607 (*synth*)Takahata, H. *et al.*, *Tetrahedron*, 2004, **60**, 8199-8205 (*Fagomine, 4-Epifagomine, synth*)Yokoyama, H. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 852-856 (*Fagomine, synth*)**6-(Hydroxymethyl)-3,4-piperidinediol, 9CI**

H-615

1,4,5-Trideoxy-1,5-iminoheptitol




(2S,4R,5R)-form

C₆H₁₃NO₃ 147.174**(2S,4R,5R)-form****1,4-Dideoxyaltronojirimycin**Alkaloid from *Conarus ferrugineus*.Amorph. solid. [α]_D -9 (c, 1.2 in H₂O).**(2S,4R,5S)-form****1,4-Dideoxyallonojirimycin**Alkaloid from *Conarus ferrugineus*.Amorph. solid. Mp 166-170°. [α]_D +62 (c, 0.92 in H₂O).**(2S,4S,5R)-form****1,4-Dideoxymannonojirimycin**Alkaloid from *Conarus ferrugineus*.Syrup. [α]_D -37.5 (c, 0.14 in H₂O).Asano, N. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1238-1242 (*isol, pmr, cmr*)**2-(Hydroxymethyl)-3,4,5-piperidinetriol, 12CI**

H-616

3,4,5-Trihydroxy-2-piperidinemethanol.

3,4,5-Trihydroxy-2-hydroxymethylpiperidine. 1,5-Dideoxy-1,5-iminohexitol



(2R,3R,4R,5R)-form

C₆H₁₃NO₄ 163.173

CAS name was changed for 12CI period. Different numbering systems depending on whether named as a piperidine or as an iminosugar.

(2R,3R,4R,5R)-form

1,5-Dideoxy-1,5-imino-D-mannitol, 9CI.

1-Deoxymannonojirimycin. LU 1. Anti-

biotic LU 1

[84444-90-6]

Alkaloid from the seeds of *Lonchocarpus sericeus*, *Lonchocarpus costaricensis* (Fabaceae), *Commelina communis* and *Streptomyces lavendulae*. Inhibitor of mannosidase and α -glucosidase. Cryst. (MeOH/Et₂O). Mp 187-188°. [α]_D²⁰ -26.7 (c, 0.12 in MeOH). [α]_D²⁴ -45.5 (c, 1.02 in H₂O).

Hydrochloride: [73465-43-7]

Mp 175-180°. [α]_D -10.9 (c, 0.3 in H₂O).1'-O- α -L-Rhamnopyranoside: **6-O- α -L-Rhamnopyranosyl-1-deoxymannonojirimycin**

[330594-47-3]

C₁₂H₂₃NO₈ 309.316Alkaloid from the bark of *Angylocalyx pynaertii*. [α]_D -42.6 (c, 0.33 in H₂O).5-O- α -D-Galactopyranoside: 2-O- α -D-Galactopyranosyl-1-deoxymannonojirimycinC₁₂H₂₃NO₉ 325.315Constit. of *Conarus ferrugineus*. Powder. [α]_D +51.1 (c, 0.96 in H₂O).3-O- β -D-Glucopyranoside: 4-O- β -D-Glucopyranosyl-1-deoxymannonojirimycinC₁₂H₂₃NO₉ 325.315Constit. of *Albizia myriophylla*. Powder. [α]_D -26.3 (c, 0.41 in H₂O).4-O- β -D-Glucopyranoside: 3-O- β -D-Glucopyranosyl-1-deoxymannonojirimycinC₁₂H₂₃NO₉ 325.315Constit. of *Conarus ferrugineus*. Powder. [α]_D -27.9 (c, 0.35 in H₂O).5-O- β -D-Glucopyranoside: 2-O- β -D-Glucopyranosyl-1-deoxymannonojirimycinC₁₂H₂₃NO₉ 325.315Constit. of *Albizia myriophylla*. Powder. [α]_D -28.3 (c, 0.86 in H₂O).**(2R,3R,4R,5S)-form**1,5-Dideoxy-1,5-imino-D-glucitol, 9CI. **1-Deoxynojirimycin**. Moranoline. Antibiotic S-GI. S-GI

[19130-96-2]

Alkaloid from *Morus* spp. (Moraceae), *Jacobinia suberecta* and *Commelina communis*. Also prod. by various *Bacillus* spp. and *Streptomyces lavendulae* ssp. *trehalostaticus*. α -Glucosidase inhibitor. Animal feedstuff additive. Shows anti-HIV activity. Antihyperglycaemic activity *in vivo*. Cryst. (MeOH). Mp 206° (195-197°). [α]_D²² +44.7 (c, 1 in H₂O). [α]_D²⁰ -34.9 (c, 5 in H₂O). pK_a 6.6. Log P -2.85 (calc).

Hydrochloride: [73285-50-4]

Amorph. powder. Mp 196-198°. [α]_D²² +38 (c, 1 in H₂O).1'-Phosphate: **1-Deoxynojirimycin 6-phosphate**

[134462-64-9]

C₆H₁₄NO₇P 243.153Isol. from the sponge *Lendenfeldia chondrodes*. Amorph. solid. [α]_D²⁴ +27 (c, 0.19 in H₂O).1'-O- α -D-Galactopyranoside: **6-O- α -D-Galactopyranosyl-1-deoxynojirimycin**

[156639-78-0]

C₁₂H₂₃NO₉ 325.315Alkaloid from *Morus alba* (white mulberry). Powder. [α]_D +107 (c, 0.1 in H₂O).5-O- α -D-Galactopyranoside: **2-O- α -D-Galactopyranosyl-1-deoxynojirimycin**

[155168-05-1]

C₁₂H₂₃NO₉ 325.315From leaves of *Morus bombycis* and from *Morus alba* (white mulberry). Powder. [α]_D +118.8 (c, 0.1 in H₂O).1'-O- β -D-Glucopyranoside: **6-O- β -D-Glucopyranosyl-1-deoxynojirimycin**

[156714-96-4]

C₁₂H₂₃NO₉ 325.315Alkaloid from *Morus alba* (white mulberry). Powder.3-O- α -D-Glucopyranoside: **4-O- α -D-Glu-**

- copyranosyl-1-deoxyojirimycin**. 4-O- α -D-Glucopyranosylmoranoline
[80312-32-9]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry). Needles (H₂O). Mp 189-192°. [α]_D²⁴ +128.6 (c, 1 in H₂O).
- 3-O- β -D-Glucopyranoside: **4-O- β -D-Glucopyranosyl-1-deoxyojirimycin**
[152375-46-7]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry). [α]_D +25 (c, 0.42 in H₂O).
- 4-O- α -D-Glucopyranoside: **3-O- α -D-Glucopyranosyl-1-deoxyojirimycin**
[156714-94-2]
[150447-77-1]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry) and *Scilla sibirica*. [α]_D +137.5 (c, 0.52 in H₂O).
- 4-O- β -D-Glucopyranoside: **3-O- β -D-Glucopyranosyl-1-deoxyojirimycin**
[143900-42-9]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry). Powder. [α]_D +18.1 (c, 0.74 in H₂O).
- 5-O- α -D-Glucopyranoside: **2-O- α -D-Glucopyranosyl-1-deoxyojirimycin**
[156714-93-1]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry). [α]_D +127.6 (c, 0.61 in H₂O).
- 5-O- β -D-Glucopyranoside: **2-O- β -D-Glucopyranosyl-1-deoxyojirimycin**
[156714-95-3]
C₁₂H₂₃NO₉ 325.315
Alkaloid from *Morus alba* (white mulberry). [α]_D -0.8 (c, 0.51 in H₂O).
- N-Me: **1,5-Dideoxy-1,5-(methylimino)-D-glucitol**, 9CI. **N-Methyl-1-deoxyojirimycin**. N-Methylmoranoline
[69567-10-8]
C₇H₁₅NO₄ 177.2
Alkaloid from *Morus alba* (white mulberry) and *Morus bombycis*. Isol. from a *Streptomyces* sp. Cryst. (EtOH). Mp 141-142°. [α]_D²⁴ +15.5 (H₂O).
- N-Me, 1'-phosphate: **N-Methyl-1-deoxyojirimycin 6-phosphate**
[915158-93-9]
C₇H₁₆NO₇P 257.18
Isol. from the sponge *Lendenfeldia chondrodes*. Amorph. solid. [α]_D²⁴ +9.7 (c, 0.1 in H₂O).
- N-Butyl: **N-Butyldeoxyojirimycin**. **Miglustat**, **BAN**, **INN**, **USAN**. **Butyl-DNJ**. **OXAIDS**. **Vevesca**. **Zavesca**. **OGT 918**. **SC48334**
[72599-27-0]
C₁₀H₂₁NO₄ 219.28
Ceramide-specific glucosyltransferase inhibitor. Inhibits biosynthesis of glucosphingolipids. Antiviral activity, of potential use against HIV infections. A substrate balance (deprivation) therapy for the treatment of Gaucher's disease and other lysosomal storage disorders. Log P -0.4 (calc).
- (**2R,3R,4S,5R**)-form
1,5-Dideoxy-1,5-imino-D-altritol. **1-Deoxy-D-altronojirimycin**
[135395-59-4]
Alkaloid from the bark of *Angylocalyx pynaertii* and the bulbs of *Scilla sibirica*. Mp 175°. [α]_D²⁰ -30.7 (c, 0.13 in MeOH) (synthetic). [α]_D +19.1 (c, 0.74 in H₂O) (natural).
- (**2R,3R,4S,5S**)-form
1-Deoxy-1,5-imino-D-allitol. **1-Deoxy-D-allonojirimycin**
[158341-94-7]
[α]_D²⁵ +30.5 (c, 1 in H₂O).
- (**2R,3S,4R,5R**)-form
1,5-Dideoxy-1,5-imino-D-talitol. **1-Deoxy-D-talonojirimycin**
[143406-74-0]
Mp 150-151°. [α]_D²⁰ +30.4 (c, 0.3 in EtOH). [α]_D²³ -22.4 (c, 1.6 in MeOH).
- (**2R,3S,4R,5S**)-form
1,5-Dideoxy-1,5-imino-D-galactitol. **1-Deoxy-D-galactonojirimycin**. **1-Deoxy-D-galactostatin**. **Migalstat**, **INN**. **AT 1001**
[108147-54-2] α -Galactosidase A enzyme inhibitor. Irregular prisms (EtOH aq.) (as hydrochloride). Mp 240-241.5° dec. (hydrochloride). [α]_D²³ +52.8 (c, 1 in H₂O). [α]_D +61.3 (c, 0.4 in H₂O).
Hydrochloride: Migalstat hydrochloride, **USAN**. **Amigal**
[75172-81-5] Used in the treatment of Fabry disease.
- (**2R,3S,4S,5R**)-form
1,5-Dideoxy-1,5-imino-D-iditol. **1-Deoxy-D-idonojirimycin**
Cryst. (MeOH/Et₂O). Mp 137-139°. [α]_D²⁵ +28 (c, 0.5 in H₂O).
- (**2R,3S,4S,5S**)-form
1-Deoxy-D-gulonojirimycin. **1-Deoxy-1,5-imino-D-gulitol**
[254104-05-7]
Mp 148-148.6°. [α]_D²⁰ -16.2 (c, 0.3 in EtOH).
- (**2S,3R,4R,5R**)-form
1,5-Dideoxy-1,5-imino-L-gulitol. **1-Deoxy-L-gulonojirimycin**
[84518-54-7]
Alkaloid from the bark of *Angylocalyx pynaertii*. Oil. [α]_D²¹ +8.8 (c, 0.25 in H₂O) (synthetic). [α]_D +14 (c, 0.56 in H₂O) (natural). Abs. config. of natural product was revised in 2005.
- (**2S,3R,4R,5S**)-form
1,5-Dideoxy-1,5-imino-L-iditol. **1-Deoxy-L-idonojirimycin**
[16647-80-6]
Cryst. (MeOH/Et₂O). Mp 137-139°. [α]_D²⁵ -27 (c, 0.33 in H₂O).
Hydrochloride: [210223-32-8]
Cryst. (MeOH aq.). [α]_D²⁰ +32.3 (c, 1.0 in H₂O).
N-Pentyl: **OGT 2378**
[441061-33-2]
C₁₁H₂₃NO₄ 233.307
Antineoplastic agent. Inhibits glucosylceramide synth. Powder.
- (**2S,3R,4S,5R**)-form
1,5-Dideoxy-1,5-imino-L-galactitol
[126663-71-6]
[α]_D²⁰ -48.3 (c, 1.06 in H₂O).
Hydrochloride: [126663-84-1]
Mp 225-230°. [α]_D²⁰ -54.1 (c, 1.0 in H₂O).
- (**2S,3S,4R,5S**)-form
1,5-Dideoxy-1,5-imino-L-altritol. **1-Deoxy-L-altronojirimycin**
[188779-13-7]
Oil. [α]_D -7 (c, 0.4 in MeOH). [α]_D²⁰ -34.6 (c, 0.3 in H₂O).
- (**2S,3S,4S,5R**)-form
1,5-Dideoxy-1,5-imino-L-glucitol
[146747-37-7]
Mp 193-195°. [α]_D²⁵ -46 (c, 0.3 in H₂O).
- (**2S,3S,4S,5S**)-form
1,5-Dideoxy-1,5-imino-L-mannitol
[117821-07-5]
[α]_D²² +26.6 (H₂O).
Hydrochloride: [118464-54-3]
Mp 184-186°. [α]_D²⁰ +12.6 (H₂O).
Yagi, M. et al., *CA*, 1977, **86**, 167851r (isol)
Fellows, L.E. et al., *Chem. Comm.*, 1979, 977 (isol, pmr, cmr, ms, cd, struct, *Deoxymannojirimycin*)
Schmidt, D.D. et al., *Naturwissenschaften*, 1979, **66**, 584 (isol)
Murao, S. et al., *Agric. Biol. Chem.*, 1980, **44**, 219 (isol)
Legler, G. et al., *Carbohydr. Res.*, 1984, **61**, 128 (*Deoxymannojirimycin*, synth)
Bernotas, R.C. et al., *Tet. Lett.*, 1984, **25**, 165; 1985, **26**, 1123 (*Deoxymannojirimycin*)
Ezure, Y. et al., *Agric. Biol. Chem.*, 1985, **49**, 2159-2165 (*4-Glucosyl-1-deoxyojirimycin*, synth)
Fleet, G.W.J. et al., *Tet. Lett.*, 1985, **26**, 1469; 1988, **29**, 2817 (*D-manno, L-manno*, synth)
Daigo, K. et al., *Chem. Pharm. Bull.*, 1986, **34**, 2243 (isol)
Yoshikuni, Y. et al., *Agric. Biol. Chem.*, 1988, **52**, 121 (*Moranoline*, pharmacol)
Ziegler, T. et al., *Angew. Chem., Int. Ed.*, 1988, **27**, 716 (*Deoxymannojirimycin*)
Paulsen, H. et al., *Annalen*, 1988, 1121 (synth, *L-galacto*)
Fleet, G.W.J. et al., *FEBS Lett.*, 1988, **237**, 128 (*N-butyl*, synth)
Ezure, Y. et al., *J. Antibiot.*, 1988, **41**, 1142 (*Deoxymannojirimycin*, isol)
Cole, M.D. et al., *J. Chromatogr.*, 1988, **445**, 295 (hplc)
Karpas, A. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 9229 (pharmacol, *N-butyl*)
Furumoto, T. et al., *J. Antibiot.*, 1989, **42**, 1302 (*Deoxymannojirimycin*)
Von der Ostern, C.H. et al., *J.A.C.S.*, 1989, **111**, 3924 (*enzymatic synth*)
Paulsen, H. et al., *Annalen*, 1990, 953 (synth, *L-galacto*)
Ermer, P. et al., *Helv. Chim. Acta*, 1991, **74**, 2043 (*Moranoline*, synth)
Behling, J. et al., *Synth. Commun.*, 1991, **21**, 1383 (*Moranoline*, synth)
Baxter, E.W. et al., *Bioorg. Med. Chem. Lett.*, 1992, **2**, 1419 (*N-butyl*, synth)
Chida, N. et al., *Carbohydr. Res.*, 1992, **237**, 185 (*Deoxyojirimycin*, synth)
Faber, E.D. et al., *Pharm. Res.*, 1992, **9**, 1442 (*Deoxyojirimycin*, metab)
Hardick, D.J. et al., *Tetrahedron*, 1992, **48**, 6285 (*biosynth*)
Oshiro, Y. et al., *Toxicol. Lett.*, 1992, **60**, 275 (*N-butyl*, tox)

Ratner, L. *et al.*, *AIDS Res. Hum. Retroviruses*, 1993, **9**, 291 (*N*-butyl, HIV-1)

Fowler, P.A. *et al.*, *Carbohydr. Res.*, 1993, **246**, 377 (*L*-ido)

Hempel, A. *et al.*, *J. Med. Chem.*, 1993, **36**, 4082 (*Deoxymannojirimycin, cryst struct*)

Asano, N. *et al.*, *Carbohydr. Res.*, 1994, **253**, 235-245; **259**, 243-255 (*N*-Methyl-*I*-deoxymannojirimycin, glycosides, 2-*O*- α -*D*-Galactopyranosyl-*I*-deoxymannojirimycin)

Zou, W. *et al.*, *Carbohydr. Res.*, 1994, **254**, 25 (*Deoxymannojirimycin, synth*)

Platt, F.M. *et al.*, *J. Biol. Chem.*, 1994, **269**, 8362 (*N*-butyl, pharmacol)

Park, K.H. *et al.*, *J.C.S. Perkin 1*, 1994, 2871 (*synth, D*-manno)

Hughes, A.B. *et al.*, *Nat. Prod. Rep.*, 1994, **11**, 135 (*Moranoline, rev*)

Block, T.M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1994, **91**, 2235 (*N*-butyl, antihepatitis B activity)

Johnson, C.R. *et al.*, *Synlett*, 1995, 313-314 (*D*-talo-*Deoxyltronojirimycin, Deoxymannojirimycin, synth, pmr, cmr*)

Ikota, N. *et al.*, *Heterocycles*, 1997, **46**, 637-643 (*Moraniline, synth*)

Xu, J.-M. *et al.*, *J.C.S. Perkin 1*, 1997, 741-746

Schaller, C. *et al.*, *Carbohydr. Res.*, 1998, **314**, 25-35 (*D*-gluco, *L*-gluco, *D*-ido, *L*-ido)

Kazmaier, U. *et al.*, *Eur. J. Org. Chem.*, 1998, 1833-1840 (*Deoxyltronojirimycin*)

Davis, B.G. *et al.*, *Tetrahedron: Asymmetry*, 1998, **9**, 2947-2960 (*L*-gulo)

Kim, H.-S. *et al.*, *Planta Med.*, 1999, **65**, 437-439 (*isol, activity*)

Ruiz, M. *et al.*, *Synlett*, 1999, 204-206, (*D*-talo)

Uriel, C. *et al.*, *Synlett*, 1999, 593-595 (*synth, D*-galacto, *L*-altro)

Matos, C.R.R. *et al.*, *Synthesis*, 1999, 571-573 (*synth, Deoxymannojirimycin*)

Meyers, A.I. *et al.*, *Tetrahedron*, 1999, **55**, 8931-8952 (*L*-manno, *synth*)

Cox, T. *et al.*, *Lancet*, 2000, **355**, 1481-1485 (*N*-butyl, pharmacol)

Lee, B.W. *et al.*, *Synthesis*, 2000, 1305-1309 (*L*-ido)

Ranes, M.K. *et al.*, *Br. J. Cancer*, 2001, **84**, 1107-1114 (*N*-butyl, pharmacol)

Asano, N. *et al.*, *Eur. J. Biochem.*, 2001, **268**, 35-41 (*Angylocalyx pynaertii* isolates)

Watson, A.A. *et al.*, *Phytochemistry*, 2001, **56**, 265-295

Spreitz, J. *et al.*, *Carbohydr. Res.*, 2002, **337**, 183-186 (*Deoxymannojirimycin, synth*)

Joseph, C.C. *et al.*, *Carbohydr. Res.*, 2002, **337**, 1083-1087 (*L*-gulo form, *D*-talo-form, *synth*)

Schiffmann, R. *et al.*, *Drugs*, 2002, **62**, 733-742 (*N*-butyl, rev)

Ruiz, M. *et al.*, *Tetrahedron: Asymmetry*, 2002, **13**, 795-799 (*D*-allo, *D*-gulo, *synth*)

Weiss, M. *et al.*, *Cancer Res.*, 2003, **63**, 3654-3658 (*OGT 2378*)

Lachmann, R.H. *et al.*, *Curr. Opin. Invest. Drugs*, 2003, **4**, 472-479 (*miglustat, rev*)

McCormack, P.L. *et al.*, *Drugs*, 2003, **63**, 2427-2434 (*miglustat, rev*)

Sorbera, L.A. *et al.*, *Drugs of the Future*, 2003, **28**, 229-236 (*N*-butyl, rev)

Banwell, M.G. *et al.*, *Org. Biomol. Chem.*, 2003, 2035-2037 (*Deoxymannojirimycin 6-rhamnoside, synth*)

Zimran, A. *et al.*, *Philos. Trans. R. Soc. London, B*, 2003, **358**, 961-966 (*N*-butyl, pharmacol)

Somfai, P. *et al.*, *Tetrahedron*, 2003, **59**, 1293-1299 (*Deoxymannojirimycin, synth*)

Pyun, S.-J. *et al.*, *Heterocycles*, 2004, **62**, 333-341 (*Deoxygalactonojirimycin, synth*)

Kimura, T. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 1415-1418 (*Deoxymannojirimycin, detn, hplc*)

Amat, M. *et al.*, *Tet. Lett.*, 2004, **45**, 5355-5358 (*D*-gulo-form, *synth*)

Morwenna, M.S.M. *et al.*, *Eur. J. Org. Chem.*, 2005, 2159-2191 (*rev, synth*)

Asano, N. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1238-1242 (*5*-galactoside, *3*-*4*-*5*-glucosides)

Boglio, C. *et al.*, *Org. Lett.*, 2005, **7**, 4851-4854 (*Deoxymannojirimycin, synth*)

Pyun, S.-J. *et al.*, *Tetrahedron*, 2005, **61**, 1413-1416 (*Deoxygulonojirimycin, synth, abs config*)

Sakai, R. *et al.*, *J. Antibiot.*, 2006, **59**, 507-511 (*I*-*Deoxymannojirimycin 6*-phosphates)

Ghosh, S. *et al.*, *Tet. Lett.*, 2006, **47**, 6041-6044 (*synth*)

Song, X. *et al.*, *Tet. Lett.*, 2007, **48**, 3115-3118 (*synth*)

2-Hydroxy-2-methylpropanoic acid, 9CI H-617

2-Methylactic acid, 8CI. *2*-Hydroxyisobutyric acid. Acetonic acid

[594-61-6]

(H₃C)₂C(OH)COOH

C₄H₈O₃ 104.105

Hygroscopic prisms (Et₂O). Sol. H₂O; spar. sol. C₆H₆. Mp 79°. Bp 212° Bp₁₂ 114°.

Nitrile: *2*-Hydroxy-2-methylpropanenitrile. *2*-Cyano-2-propanol. *2*-Hydroxyisobutyronitrile. Acetone cyanohydrin [75-86-5]

Used in place of HCN in Gattermann reaction for formylation of arenes. V. sol. H₂O; spar. sol. petrol. Mp -19°. Bp₂₃ 82°.

►Hydrolyses readily to HCN and Me₂CO. Fl. p. 64/74°, autoignition temp. 688°. LD₅₀ (rat, orl) 17.8 mg/kg. LD₅₀ (rbt, skn) 17 mg/kg. Human fatalities reported from skin contact or oral ingestion. OD9275000

Nitrile, O- β -*D*-glucopyranoside: *2*-(β -*D*-Glucopyranosyloxy)-2-methylpropanenitrile, 9CI. **Linamarin**. Phaseolunatin. *Manihot*toxin

[554-35-8]

C₁₀H₁₇NO₆ 247.247

Occurs in flax (*Linum usitatissimum*), manioc (*Manihot utilisissimum*), *Phaseolus lunatus*, *Trifolium repens* (white clover) and other plants. First isol. in 1830. Needles. Mp 143-144°. [α]_D -28.5 (c, 3.86 in H₂O).

►Exp. teratogen. TZ4850000

Nitrile, O-[β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranoside]: **Linustatin**. *Linamarin gentiobioside*

[72229-40-4]

C₁₆H₂₇NO₁₁ 409.389

Isol. from *Passiflora pendens* and flax seed meal. Cryst. (EtOH). Mp 123-123.5°. [α]_D²⁵ -37 (c, 0.31 in H₂O).

Butler, G.W. *et al.*, *Phytochemistry*, 1965, **4**, 127 (*Linamarin*)

Malik, B.K. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 76 (*Linamarin*)


Smith, C.R. *et al.*, *J.O.C.*, 1980, **45**, 507 (*Linustatin*)

Hirata, K. *et al.*, *CA*, 1990, **114**, 162500n (*Linamarin, anal*)

Olafsdottir, E.S. *et al.*, *Phytochemistry*, 1992, **31**, 4129 (*biosynth*)

Nakajima, N. *et al.*, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 453-458 (*Linamarin, synth, pmr, cmr*)

2-(1-Hydroxy-1-methylpropyl)-3-methoxypyrazine H-618



C₉H₁₄N₂O₂ 182.222

Prod. by *Chondromyces crocatus*. Oil.

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol, synth, pmr, cmr*)

2-(1-Hydroxy-2-methylpropyl)-3-methoxypyrazine H-619


C₉H₁₄N₂O₂ 182.222

Prod. by *Chondromyces crocatus*.

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol, synth, pmr, cmr*)

1-Hydroxy-3-(2-methylpropyl)-6-[2-(methylthio)ethyl]-2(1H)-pyrazinone H-620

1-Hydroxy-3-isobutyl-6-[2-(methylthio)ethyl]-2(1H)-pyrazinone [37175-51-2]



C₁₁H₁₈N₂O₂S 242.341

Metab. from *Aspergillus flavus*. Mp 114-116°.

MacDonald, J.C. *et al.*, *Can. J. Biochem.*, 1972, **50**, 543 (*isol, pmr, ms, struct*)

1-Hydroxy-6-(2-methylpropyl)-3-[2-(methylthio)ethyl]-2(1H)-pyrazinone H-621

1-Hydroxy-6-isobutyl-3-[2-(methylthio)ethyl]-2(1H)-pyrazinone [37175-52-3]


C₁₁H₁₈N₂O₂S 242.341

Metab. from *Aspergillus flavus*. Mp 114-116°.

MacDonald, J.C. *et al.*, *Can. J. Biochem.*, 1972, **50**, 543 (*isol, pmr, ms, struct*)

4-Hydroxy-2-methyl-6-propylpiperidine H-622

2-Methyl-6-propyl-4-piperidinol, 9CI



C₉H₁₉NO 157.255

(2*R*,4*S*,6*S*)-form

Trace alkaloid from *Picea pungens*. Cryst. (EtOAc). Mp 88-89° (*synthetic*).


2'-Oxo: *1*-(4-Hydroxy-6-methyl-2-piperi-

dinyl)-2-propanone. 4-Hydroxy-2-methyl-6-(2-oxopropyl)piperidine
C₉H₁₇NO₂ 171.239

Trace alkaloid from *Picea pungens*. Oil (synthetic).

Tawara, J.N. et al., *J. Nat. Prod.*, 1999, **62**, 321-323 (isol, synth, pmr, cmr, ms)

6-(Hydroxymethyl)-2,4(1H,3H)-pteridinedione H-623
6-Hydroxymethylumazine, 8CI
[10129-99-4]




C₇H₆N₄O₃ 194.149

Isol. from leaves of *Spinacia oleracea*. Mp 260-262° dec.

Sugiura, K. et al., *CA*, 1966, **65**, 15375; 20508 (isol, uv, synth, struct)

2-Hydroxy-3-methylpyrazine H-624
3-Methyl-2(1H)-pyrazinone, 9CI. 3-Methylpyrazinol, 8CI
[19838-07-4]



C₅H₆N₂O 110.115

Cryst. (EtOAc). Mp 151-152°.

OH-form

Me ether: 2-Methoxy-3-methylpyrazine
[2847-30-5]

C₆H₈N₂O 124.142

Prod. by *Chondromyces crocatus*. Flavouring ingredient. Liq. Bp₆₀ 87°.

Et ether: 2-Ethoxy-3-methylpyrazine
[32737-14-7]

C₇H₁₀N₂O 138.169

Flavouring ingredient. Bp₄₈ 88-90°.

Et ether, 4-oxide:

C₇H₁₀N₂O₂ 154.168

Cryst. (EtOH), Mp 121-122.5°.

Karmus, G. et al., *J.A.C.S.*, 1952, **74**, 1580 (synth)

Hirschberg, A. et al., *J.O.C.*, 1961, **26**, 2356-2360 (*Et ether*, synth)

Klein, B. et al., *J.O.C.*, 1964, **29**, 2623-2626 (*Et ether*, *Et ether N-oxide*)

Fr. Pat., 1965, 1 391 212; *CA*, **62**, 16270g (*FEMA 3183*)

Kolor, M.G. et al., *Org. Mass Spectrom.*, 1971, **5**, 959-966 (*Me ether*, ms)

Nakel, G.M. et al., *J. Agric. Food Chem.*, 1972, **20**, 682-684 (*Me ether*, struct)

Bramwell, A.F. et al., *Tetrahedron*, 1972, **28**, 4155-4170; 1973, **29**, 3939 (synth, pmr)

Parliment, T.H. et al., *J. Agric. Food Chem.*, 1973, **21**, 714-716 (synth)


Calabretta, P.J. et al., *Cosmet. Perfum.*, 1975, **90**, 74; 76; 79-80 (*Me ether*, synth, use)

Macdonald, J.C. et al., *Tetrahedron*, 1976, **32**, 655 (pmr, cmr)

Fenaroli's Handbook of Flavor Ingredients, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 1057-1058; *Fenaroli's Handbook of Flavor Ingredients*, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 1118 (*FEMA 3569*, *FEMA 3183*)

Schulz, S. et al., *Tetrahedron*, 2004, **60**, 3863-3872 (isol)

4-Hydroxy-2-methylpyridine H-625
2-Methyl-4(1H)-pyridinone. 2-Methyl-4-pyridinol
[18615-86-6]



NH-form

OH-form

C₆H₇NO 109.127

NH-form predominates. Isol. from the fungus *Physozorinus sanguinolentus*.

Cryst. (Me₂CO). Mp 174-176°. Bp_{0.003} 180-189°.

Hydrochloride: Mp 116°.

N-Oxide:

C₆H₇NO₂ 125.127

Mp 192-193°.

OH-form

Me ether, N-oxide:

C₇H₉NO₂ 139.154

Mp 78-80°.

Et ether: 4-Ethoxy-2-methylpyridine

C₈H₁₁NO 137.181

Bp ca. ° 220.

Ph ether, N-oxide:

C₁₂H₁₁NO₂ 201.224

Mp 161-162° (as hydrochloride).

Benzyl ether, N-oxide: [63071-08-9]

C₁₃H₁₃NO₂ 215.251

Fluffy cryst. (Me₂CO). Mp 154-156°.

Collie, J.N. et al., *J.C.S.*, 1925, **127**, 962

Suzuki, et al., *Yakugaku Zasshi*, 1948, **68**, 126

Kato, T. et al., *Yakugaku Zasshi*, 1955, **75**,

1239; *CA*, **50**, 8666

Nantka-Namirski, P. et al., *Acta Pol. Pharm.*,

1967, **24**, 231; *CA*, **69**, 2827y


Hom, R.K. et al., *J.O.C.*, 1996, **61**, 2624

(*benzyl ether oxide*)

Svensson, M. et al., *Phytochemistry*, 2001, **56**,

747-751 (isol, pmr, cmr, ms)

5-(Hydroxymethyl)-1H-pyrrole-2-carboxaldehyde H-626
2-Formyl-5-(hydroxymethyl)pyrrole
[67350-50-9]



C₆H₇NO₂ 125.127

Cryst. (C₆H₆). Mp 94-95°.

O-(4-Hydroxybutanoyl): (5-Formyl-1H-pyrrol-2-yl)methyl 4-hydroxybutanoate
C₁₀H₁₃NO₄ 211.217

Alkaloid from the seeds of *Castanea*

sativa. Oil. λ_{max} 255 (sh); 291 (MeOH).

N-Me: 5-Hydroxymethyl-1-methylpyrrolecarboxaldehyde

[29813-44-3]

C₇H₉NO₂ 139.154

Oil or cryst. Mp 23-24°. Subl. 0.01 80.

N-Me, O-Ac:

C₉H₁₁NO₃ 181.191

Cryst. Mp 20-20.2°.

N-(3-Methylbutyl): **Ganoine**

[133086-80-3]

C₁₁H₁₇NO₂ 195.261

Alkaloid from the mycelium of *Ganoderma capense*.

N-(2-Phenylethyl): **Ganodine**

[133086-81-4]

C₁₄H₁₅NO₂ 229.278

Alkaloid from the mycelium of *Ganoderma capense*.

N-[2-(4-Hydroxyphenyl)ethyl]: **Pyrrrolezanthine**

C₁₄H₁₅NO₃ 245.277

Alkaloid from the stems of *Zanthoxylum simulans* (Szechuan pepper).

Needles (C₆H₆). Mp 106-108°. λ_{max} 206 (log ε 3.9); 224 (log ε 3.8); 296 (log ε 3.89) (EtOH).

Me ether: 5-(Methoxymethyl)-1H-pyrrole-2-carboxaldehyde. 2-Formyl-5-(methoxymethyl)-1H-pyrrole

C₇H₉NO₂ 139.154

Alkaloid from the roots of *Salvia miltiorrhiza*. Brown oil.

Kato, H. et al., *Agric. Biol. Chem.*, 1970, **34**,

1071-1077 (synth, uv, pmr)

Jurch, G.R. et al., *Carbohydr. Res.*, 1970, **15**,

233-239 (synth, derivs, ir, pmr, ms)

Severin, T. et al., *Chem. Ber.*, 1975, **108**, 1768-

1775 (synth, derivs, ir, pmr, ms)

Olsson, K. et al., *Acta Chem. Scand., Ser. B*,

1977, **31**, 469-474; 1978, **32**, 249-256 (synth,

derivs, ir, pmr, ms)

Yang, J.J. et al., *Yaoxue Xuebao*, 1990, **25**, 555;

612; *CA*, **114**, 160668z; 164574p (*Ganoine*,

Ganodine)

Hiermann, A. et al., *Fitoterapia*, 2002, **73**, 22-

27 (4-hydroxybutanoyl)


Yang, Y.-P. et al., *Phytochemistry*, 2002, **61**,

567-572 (*Pyrrrolezanthine*)

Don, M.-J. et al., *J. Nat. Prod.*, 2005, **68**, 1066-

1070 (*Me ether*)

3-Hydroxy-4-methyl-2-pyrrolidinecarboxylic acid H-627
3-Hydroxy-4-methylproline, 9CI



(2S,3S,4R)-form

C₆H₁₁NO₃ 145.158

(2S,3S,4R)-form

Hydrochloride: [208709-19-7]

[α]_D²⁴ -4 (c, 0.48 in MeOH).

(2S,3S,4S)-form [54615-51-9]


Component of the antibiotic complex Antibiotic A 30912. Mp 265° dec. [α]_D -21 (c, 0.425 in H₂O).

Keller-Schierlein, W. et al., *Helv. Chim. Acta*,

- 1974, **57**, 2459-2477 (*isol*)
 Koyama, G. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 2477-2483 (*cryst struct*)
 Keller-Jusler, C. *et al.*, *Tet. Lett.*, 1976, 4147 (*isol*)
 Kurokawa, N. *et al.*, *J.A.C.S.*, 1986, **108**, 6041-6043 (*synth*)
 Mulzer, J. *et al.*, *J.A.C.S.*, 1989, **111**, 7500-7504 (*synth, pmr, cmr, ir*)
 Langlois, N. *et al.*, *Tetrahedron*, 2000, **56**, 2437-2448 (2*S*,3*S*,4*S*-form, 2*S*,3*S*,4*R*-form, *synth, pmr, cmr*)
 Raghavan, S. *et al.*, *Tet. Lett.*, 2003, **44**, 7459-7462 (2*R*,3*R*,4*R*-form, *synth*)
 Inaba, H. *et al.*, *Heterocycles*, 2005, **65**, 2657-2665 (2*R*,3*R*,4*R*-form, *synth*)

3-Hydroxy-5-methyl-2-pyrrolidinecarboxylic acid H-628

3-Hydroxy-5-methylproline, 9CI



C₆H₁₁NO₃ 145.158

(2*R**,3*S**,5*R**)-form

(2*α*,3*α*,5*α*)-form

[55137-75-2]

Isol. from hydrolysates of Actinomycin Z.

(2*RS*,3*SR*,4*RS*)-form

(±)-(2*α*,3*α*,5*α*)-form

[61248-03-1]

Needles (Me₂CO aq.). Mp 267-268°.

(2*RS*,3*RS*,5*RS*)-form

(±)-(2*α*,3*β*,5*α*)-form

[61248-05-3]

Plates (Me₂CO aq.). Mp 204-205° dec.

(2*RS*,3*RS*,5*SR*)-form

(±)-(2*α*,3*β*,5*β*)-form

[61248-07-5]

Plates (Me₂CO aq.). Mp 259-260° dec.

(2*RS*,3*SR*,5*SR*)-form

(±)-(2*α*,3*α*,5*β*)-form

[61248-04-2]

Plates (Me₂CO aq.). Mp 242-243° dec.


Katz, E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1975, **63**, 502 (*isol, struct, ms*)

Mauger, A.B. *et al.*, *J.O.C.*, 1977, **42**, 1000 (*struct, synth, ms, pmr*)

Fushiyama, S. *et al.*, *Chem. Lett.*, 1987, 2229 (*synth*)

1-Hydroxymethylpyrrolizidine H-629

Hexahydro-1H-pyrrolizine-1-methanol, 9CI



C₈H₁₅NO 141.213

Only simple esters are included here. Many others have individual entries. p*K*_a 10.83 (26°).

(1*R*,7*aR*)-form

Lindelofidine. (+)-*Isoretronecanol*

[488-06-2]

Necine base from pyrrolizidine alkaloids, also isol. from *Thesium minkwitzianum*, *Vandopsis lissochiloides* and *Vandopsis gigantea* (Santalaceae, Orchidaceae). Oil or *cryst*. Mp 40-41°. Bp₂ 116-117°. [α]_D²³ +72 (c, 1 in EtOH).

Picrate: Mp 193-194°.

Ac: *Lindelofidine acetate*

[26673-87-0]

C₁₀H₁₇NO₂ 183.25

Alkaloid from *Vandopsis lissochiloides* and *Vandopsis gigantea* (Orchidaceae). Oil. [α]_D²⁴ +69 (c, 0.6 in CHCl₃).

4-Hydroxy-E-cinnamoyl: *Thesinine*

[488-02-8]

C₁₇H₂₁NO₃ 287.358

Alkaloid from *Thesium minkwitzianum* (Santalaceae). Mp 38-40°.

O-[α-L-Rhamnopyranosyl-(→4)-4-hydroxy-E-cinnamoyl]: *Thesinine 4'-rhamnoside*

C₂₃H₃₁NO₇ 433.5

Alkaloid from *Tephrosia kirilowii*. Amorph. solid. [α]_D²³ -72.7 (c, 0.55 in MeOH). λ_{max} 220 (log ε 4.22); 298 (log ε 4.22) (MeOH).

O-[β-D-Glucopyranosyl-(→4)-4-hydroxy-E-cinnamoyl]: *Thesinine 4'-glucoside*

C₂₃H₃₁NO₈ 449.5

Alkaloid from the seeds of *Borago officinalis* (borago). Amorph. solid. [α]_D²⁰ -22.8 (c, 2 in DMSO). λ_{max} 204 (log ε 4.15); 225 (log ε 4.06); 297 (log ε 4.28); 305 (log ε 4.29) (MeOH).

(1*R*,7*aS*)-form

Trachelanthamidine

[526-64-7]

Alkaloid present in *Eupatorium maculatum* and *Phalaenopsis equestris* (Asteraceae, Orchidaceae); also obt. by *hydrol.* of *Trachelanthamine* and *Viridiflorine*. Oil. Bp₁₂ 134°. [α]_D²⁰ -14.91 (c, 1 in EtOH).

O-[2*S*-Hydroxy-2-(1*S*-hydroxyethyl)-4-methylpentanoyl]:

C₁₆H₂₉NO₄ 299.409

Alkaloid from the roots of *Anchusa strigosa*. Yellow oil. [α]_D²⁵ +2 (c, 0.1 in MeOH). λ_{max} 233 (sh); 275 (MeOH).

O-(4-Hydroxy-3,5-dimethoxybenzoyl): *Alafine*. *Syringyltrachelanthamidine*

C₁₇H₂₃NO₅ 321.372

Mp 167-168°. [α]_D +5 (CHCl₃).

(1*S*,7*aR*)-form

Laburnine

[3348-73-0]

Alkaloid from the seeds of *Cytisus laburnum* and from *Vandopsis* spp. (Fabaceae, Orchidaceae). Oil. Bp₁₂ 134°. [α]_D²⁰ +17.01 (c, 1 in EtOH).

Ac: *Laburnine acetate*. *Acetylalaburnine*

C₁₀H₁₇NO₂ 183.25

Alkaloid from *Vandopsis lissochiloides* and *Vandopsis gigantea* (Orchidaceae). [α]_D²⁴ +12 (c, 5.1 in EtOH).

Tigloyl: *Tigloylallaburnine*. *Laburnine tiglate*

C₁₃H₂₁NO₂ 223.314

Alkaloid in *Planchonella thyrsoidea*

(preferred genus name *Pouteria*) and *Planchonella anteridifera* (Sapotaceae). Identified after *hydrol.* Prob. occurs as the angelate in the plant.

O-Benzoyl: *Benzoylallaburnine*. *Laburnine benzoate*

C₁₅H₁₉NO₂ 245.321

Alkaloid detected in *Planchonella thyrsoidea* and *Planchonella anteridifera* (Sapotaceae). Identified after *hydrol.* to *Laburnine* and *benzoic acid*.

(1*S*,7*aS*)-form

Isoretronecanol

[526-63-6]

Alkaloid present in *Planchonella equestris*, also obt. by *hydrol.* of other pyrrolizidine alkaloids (Sapotaceae). Oil. Bp₂ 116-117°. [α]_D²⁰ -77.5 (c, 2.5 in MeOH). p*K*_a 10.83 (26°).

Tigloyl: *Isoretronecyl tiglate*

C₁₃H₂₁NO₂ 223.314

Minor alkaloid from a *Planchonella* sp. (Sapotaceae).

O-(2*R*-Hydroxy-3*R*-methylpentanoyl):

Cremastrine

C₁₄H₂₅NO₃ 255.356

Alkaloid from the bulbs of *Cremastra appendiculata*. Inhibitor of the muscarinic M3 receptor. Syrup. [α]_D²⁵ -26.8 (c, 1 in EtOH).

(1*RS*,7*aRS*)-form

(±)-*Isoretronecanol*

[18929-90-3]

Yellow oil. Bp₃ 118-119°.

Picrate:

Yellow needles (EtOH). Mp 192-193° (189.5-191°).

(1*RS*,7*aSR*)-form

(±)-*Trachelanthamidine*

[18929-91-4]

Yellow oil.

Picrate: Mp 174-175.5°.

Leonard, N.J. *et al.*, *J.A.C.S.*, 1950, **72**, 2537 (*synth, struct*)

Arendaruk, A.P. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 670; *CA*, **54**, 24835d (*isol, Lindelofidine, Thesinine*)

Tsuda, Y. *et al.*, *Can. J. Chem.*, 1963, **41**, 1919 (*isol*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1259; 1968, **21**, 1393 (*Benzoylallaburnine, Tigloylallaburnine, isol, esters*)

Lindström, B. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 3352 (*Acetylalaburnine*)

Pais, M. *et al.*, *Ann. Pharm. Fr.*, 1971, **29**, 57 (*Alafine*)

Abdullaev, U.A. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 634; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 602 (ms)

Brandänge, S. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 1096 (*isol, Laburnine, Lindelofidine, acetates*)

Danishesky, S. *et al.*, *J.A.C.S.*, 1977, **99**, 4783 (*synth*)

Borch, R.F. *et al.*, *J.O.C.*, 1977, **42**, 1225 (*synth*)
 Pinnick, H.W. *et al.*, *J.O.C.*, 1978, **43**, 4662-4663 (*Isoretronecanol, Trachelanthamidine, synth*)

Muchowski, J.M. *et al.*, *Tet. Lett.*, 1980, **21**, 4585 (*synth*)


Robins, D.J. *et al.*, *J.C.S. Perkin I*, 1981, 909-913 (*synth*)

Robins, D.J. *et al.*, *Chem. Comm.*, 1982, 1289-1290 (*Trachelanthamidine, synth*)

- Terao, Y. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 3167 (synth)
- Iwashita, T. *et al.*, *J.O.C.*, 1982, **47**, 230 (synth)
- Flitsch, W. *et al.*, *Annalen*, 1983, 521 (synth)
- Tatsuta, K. *et al.*, *J.A.C.S.*, 1983, **105**, 4096 (synth)
- Blum, Z. *et al.*, *Acta Chem. Scand., Ser. B*, 1984, **38**, 297 (synth)
- Kametani, T. *et al.*, *Heterocycles*, 1984, **22**, 729 (synth)
- Chamberlin, A.R. *et al.*, *J.O.C.*, 1984, **49**, 1682 (synth)
- Gramain, J.-C. *et al.*, *J.O.C.*, 1985, **50**, 710 (synth)
- Ortiz, C. *et al.*, *Tet. Lett.*, 1985, **26**, 2831 (synth)
- Celerier, J.P. *et al.*, *Tet. Lett.*, 1987, **28**, 6597 (Isoretronecanol, synth, pmr, cmr)
- Keusenkothen, P.F. *et al.*, *J.C.S. Perkin 1*, 1994, 2485 (synth)
- Knight, D.W. *et al.*, *J.C.S. Perkin 1*, 1997, 2089 (synth)
- David, O. *et al.*, *J.O.C.*, 1999, **64**, 3122-3131 (synth, pmr, cmr)
- Bertrand, S. *et al.*, *Tet. Lett.*, 1999, **40**, 3173-3174 (synth)
- Herrmann, M. *et al.*, *Phytochemistry*, 2002, **60**, 399-402 (Thesinine, Thesinine 4'-glucoside)
- Braca, A. *et al.*, *Planta Med.*, 2003, **69**, 835-841 (*Anchusa strigosa* ester)
- Lemaire, S. *et al.*, *Eur. J. Org. Chem.*, 2004, 2840-2847 (Isoretronecanol, synth)
- Chang, M.-Y. *et al.*, *Heterocycles*, 2005, **65**, 2965-2971 (*Trachelanthamidine*, synth)
- Ikeda, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 572-573 (*Cremastrine*)
- Pereira, E. *et al.*, *Tet. Lett.*, 2005, **46**, 2691-2693 (Isoretronecanol, synth)
- Wang, Y.-H. *et al.*, *J. Asian Nat. Prod. Res.*, 2008, **10**, 25-31 (*Thesinine 4'-rhamnoside*)
- Ishibashi, H. *et al.*, *Tetrahedron*, 2008, **64**, 7771-7773 (*Trachelanthamidine*, synth)

1-Hydroxy-7-methylpyrrolizidine H-630

Hexahydro-7-methyl-1H-pyrrolizin-1-ol, 9CI. 7-Methyl-1-pyrrolizidinol



C₈H₁₅NO 141.213

(1R,7S,7aR)-form

Retronecanol

[567-39-5]

Necine base obt. by redn. of retronecine or by reductive hydrolysis of pyrrolizidine alkaloids, e.g. Retrorsine. Cryst. (petrol). Mp 95-96°. Bp₃₀ 140°. [α]_D²⁵ -91.1 (EtOH). pK_a 10.88 (25°).

Picrate: Mp 208°.

Ac:

C₁₀H₁₇NO₂ 183.25
Oil. Bp₁₂ 107°.

Ac, picrate:

Long prisms (EtOH). Mp 176°.

O-(4-Methoxybenzoyl): **Ehretinine**

[76231-29-3]

C₁₆H₂₁NO₃ 275.347

Alkaloid from the leaves of *Ehretia aspera* (Ehretiaceae). Mp 187-188°.

[α]_D²⁵ -108 (c, 1 in MeOH). First report of a retronecanol ester alkaloid. May be an artifact arising from hydrogenol-

ysis of retronecine diester alkaloids during isol.

Barger, G. *et al.*, *J.C.S.*, 1935, 11

Warren, F.L. *et al.*, *J.C.S.*, 1958, 4574 (abs config)

Neuner-Jehle, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 321 (ms)

Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (cd)


Suri, O.P. *et al.*, *Phytochemistry*, 1980, **19**, 1273 (isol, ir, pmr, ms, struct, synth, Ehretinine)

Robins, D.J. *et al.*, *Alkaloids (London)*, 1982, **12**, 61 (Ehretinine)

Pandey, G. *et al.*, *Tet. Lett.*, 1996, **37**, 2285-2288; 1998, **39**, 8371 (synth)

1-Hydroxymethylpyrrolizidine methyl 2-O-acetyl-2-isopropylmalate H-631

[77156-31-1]



C₁₈H₂₉NO₆ 355.43


Ester of (S)-methyl 2-isopropylmalate with 1-hydroxymethylpyrrolizidine, H-629 of undetd. config. Alkaloid from *Parsonsia heterophylla* and *Parsonsia spiralis* (Apocynaceae). Prob. a mixt. of diastereoisomers with stereochem. corresp. to Laburnine and Lindelofidine.

Cole, F.E. *et al.*, *Biochemistry*, 1973, **12**, 3346 (abs config)

Edgar, J.A. *et al.*, *Tet. Lett.*, 1980, **21**, 2657 (isol, pmr, ms, struct)

4-Hydroxy-2-methylquinazolinone H-632

2-Methyl-4(1H)-quinazolinone, 9CI. 2-Methyl-4-quinazolinol [1769-24-0]



C₉H₈N₂O 160.175

1H-Form predominates. Prod. by *Bacillus cereus*. Inhibitor of poly(ADP-ribose) synthetase. Needles (EtOH). Sol. MeOH, C₆H₆; poorly sol. H₂O, hexane. Mp 240-242° (232-234°). λ_{max} 232 (ε 38900); 269 (ε 10200); 291 (ε 7240); 302 (ε 5750) (MeOH/HCl) (Derep). λ_{max} 229 (ε 38000); 280 (ε 14500); 309 (ε 9120) (MeOH/NaOH) (Derep). λ_{max} 225 (ε 40700); 263 (ε 12900); 303 (ε 7080); 314 (ε 5890) (MeOH) (Derep). λ_{max} 229 (ε 38000); 280 (ε 14400); 309 (ε 9200) (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, orl) 859 mg/kg. VA3677000

Hydrochloride: [29378-39-0]

Cryst. (MeOH). Dec. >320°.

1H-form

N-Me: 1,2-Dimethyl-4(1H)-quinazolinone. **Glomerine**

[7471-65-0]

C₁₀H₁₀N₂O 174.202

Alkaloid from the defensive secretion of the arthropod *Glomeris marginata*. Yellow needles (EtOAc). Mp 209-211°.

3H-form

N-(2-Methoxycarbonylphenyl):

C₁₇H₁₄N₂O₃ 294.309

Alkaloid from the roots of *Aconitum pseudo-laeve* var. *erectum*. Amorph. powder (MeOH). λ_{max} 227 (log ε 4.23); 231 (log ε 4.24); 266 (log ε 3.87); 273 (log ε 3.85); 306 (log ε 3.5); 316 (log ε 3.42) (EtOH).

[110287-94-0, 92442-54-1, 132305-21-6, 86030-13-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 882A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 467A (nmr)

Bogert, M.T. *et al.*, *J.A.C.S.*, 1900, **22**, 523; 1902, **24**, 1031 (synth)

Schildknecht, H. *et al.*, *Z. Naturforsch., B*, 1966, **21**, 121; 552 (*Glomerine*, isol, uv, ir, pmr, ms, struct, synth)

Schildknecht, H. *et al.*, *Tet. Lett.*, 1967, 1815 (biosynth)

Goncalves, H. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 2599 (synth)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 2347 (synth)

Kirmani, M.Z. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 526 (ms)

Naik, N.R. *et al.*, *J. Indian Chem. Soc.*, 1979, **56**, 708 (*Glomerine*, synth)

Bhattacharyya, J. *et al.*, *Heterocycles*, 1980, **14**, 1469 (cmr)

Etter, M.C. *et al.*, *J.C.S. Perkin 2*, 1983, 115 (synth, cryst struct)

Gupton, J.T. *et al.*, *Synth. Commun.*, 1984, **14**, 1013 (synth, ir, pmr)

Spasov, S. *et al.*, *Magn. Reson. Chem.*, 1985, **23**, 795 (pmr, cmr)

Bajardi, M.L. *et al.*, *Heterocycles*, 1986, **24**, 1367 (synth)

Singh, H. *et al.*, *Tetrahedron*, 1986, **42**, 1449 (synth)

Shishoo, C.J. *et al.*, *Indian J. Chem., Sect. B*, 1989, **28**, 1039 (synth)

Yoshida, S. *et al.*, *J. Antibiot.*, 1991, **44**, 111 (isol, props)


Smith, K. *et al.*, *Zh. Org. Khim.*, 2003, **39**, 452-462; *Russ. J. Org. Chem. (Engl. Transl.)*, 2003, **39**, 430-435 (synth, ir, pmr, cmr, ms)

Rocco, S.A. *et al.*, *Synthesis*, 2004, 429-435 (synth, ir, pmr)

Shim, S.H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 400-402 (*N-methoxycarbonylphenyl*)

4-Hydroxy-2-methylquinolinone H-633

2-Methyl-4(1H)-quinolinone, 9CI. 2-Methyl-4(1H)-quinolinone, 8CI. 4-Hydroxyquinaldine. 2-Methyl-4-quinolinol. γ-Quinaldone. Pseudane I [5660-24-2]



C₁₀H₉NO 159.187Tautomeric. Cryst. (H₂O). Mp 234-235°.**Hydrochloride:**Prisms + 1H₂O (H₂O). Mp 217° (anhyd.).**Methiodide:**Cryst. + 1H₂O, losing H₂O at 100°. Mp 201° (anhyd.).**N-Oxide:**C₁₀H₉NO₂ 175.187

Mp 246-248°.

NH-form**N-Me:** 1,2-Dimethyl-4(1H)-quinolinone [6760-40-3]C₁₁H₁₁NO 173.214Alkaloid from *Platydesma campanulata* and *Acronychia baueri* (Rutaceae). Needles (C₆H₆). Sol. H₂O, EtOH, hot C₆H₆. Mp 176°.**N-Me, picrate:** Mp 234-238°.**N-Acetoxyethyl:** 1-Acetoxyethyl-2-methyl-4(1H)-quinolinoneC₁₃H₁₃NO₃ 231.251Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae). Needles (CHCl₃/MeOH). Mp 160-164°.**OH-form****O-Ac:** [117498-01-8]C₁₂H₁₁NO₂ 201.224Liq. Bp_{0.2} 100-102°.**Me ether:** 4-Methoxy-2-methylquinoline.

4-Methoxyquinaldine

[31835-53-7]

C₁₁H₁₁NO 173.214Alkaloid from aerial parts of *Ruta montana*. Cryst. (H₂O). Mp 82° (63-65°). Bp 294-298°.**Et ether:** 4-Ethoxy-2-methylquinoline. 4-Ethoxyquinaldine

[46272-56-4]

C₁₂H₁₃NO 187.241


Mp 48°.

[607-67-0]

*Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 737C (nmr)**Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 860C (ir)*Knorr, L. *et al.*, *Ber.*, 1897, **30**, 922-926 (methiodide)Limpach, L. *et al.*, *Ber.*, 1931, **64**, 969-970 (synth)Backeberg, O.E. *et al.*, *J.C.S.*, 1931, 2814-2817 (synth)Werny, F. *et al.*, *Tetrahedron*, 1963, **19**, 1293 (isol, uv, pmr, synth, deriv)Lamberton, J.A. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1995 (isol, ms, pmr, deriv)Werner, W. *et al.*, *Tetrahedron*, 1969, **25**, 255 (tautom)Schweizer, E.E. *et al.*, *J.O.C.*, 1977, **42**, 200-205 (Me ether, Et ether)Claret, P.A. *et al.*, *Spectrosc. Lett.*, 1977, **10**, 25 (pmr, cmr)Muscio, O.J. *et al.*, *J.O.C.*, 1989, **54**, 166 (deriv, synth)Ulubelen, A. *et al.*, *Fitoterapia*, 1991, **62**, 279 (4-Methoxyquinaldine)Ahsan, M. *et al.*, *Phytochemistry*, 1993, **33**, 1507 (N-acetoxyethyl)**10-Hydroxymethylsparteine**


H-634

[139112-20-2]

C₁₆H₂₈N₂O 264.41Constit. of *Genista sessilifolia* (Fabaceae). Mp 168-170° (as picrate).Nasution, M.P. *et al.*, *Tet. Lett.*, 1991, **32**, 5915**1-Hydroxy-4-(methylthio)-canthin-6-one**


H-635

1-Hydroxy-4-methylthio-6H-indolo[3,2,1-de][1,5]naphthyridin-6-one, 9CI


C₁₅H₁₀N₂O₂S 282.322**S-Oxide(S-): 9-Deoxycurtisin**C₁₅H₁₀N₂O₃S 298.322Alkaloid from the mushroom *Boletus curtisii*. Pale yellow solid. [α]_D²⁰ -70 (c, 0.05 in MeOH). λ_{max} 210 (log ε 1); 242 (sh) (log ε 0.3); 276 (sh) (log ε 0.21); 286 (log ε 0.26); 308 (log ε 0.19); 352 (log ε 0.18); 366 (log ε 0.3); 384 (log ε 0.35); 414 (log ε 0.2) (MeOH).Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, cd, ir, pmr, cmr, ms)**2-(12-Hydroxy-12-methyltridecyl)-4(1H)-quinolinone**

H-636

4-Hydroxy-2-(12-hydroxy-12-methyltridecyl)quinoline


C₂₃H₃₅NO₂ 357.535Alkaloid from the leaves of *Dictyoloma vandellianum*. Yellow gum. λ_{max} 238 ; 325 ; 335 (MeOH).**3-Methoxy:** 2-(12-Hydroxy-12-methyltridecyl)-3-methoxy-4(1H)-quinolinoneC₂₄H₃₇NO₃ 387.561Alkaloid from the leaves of *Dictyoloma vandellianum*. Yellow gum. λ_{max} 239 ; 330 ; 348 (MeOH).Sartor, C.F.P. *et al.*, *Phytochemistry*, 2003, **63**, 185-192 (isol, pmr, cmr, ms)**10-Hydroxy-18-(N-naphthalenyl-N-phenylamino)betaenone C**

H-637

C₃₇H₄₅NO₅ 583.766Prod. by a *Microsphaeropsis* sp. isol. from the sponge *Aplysina aerophoba*. Powder. [α]_D -44.4 (c, 1 in EtOH). λ_{max} 221 ; 350 (MeOH).Brauers, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 739-745**3-Hydroxy-1,4-naphthoquinone-2-carboxylic acid**

H-638


1,4-Dihydro-3-hydroxy-1,4-dioxo-2-naphthalenecarboxylic acid, 9CI

C₁₁H₆O₅ 218.165**Amide:** [103646-20-4]C₁₁H₇NO₄ 217.181Constit. of *Rubia cordifolia*. Yellow cryst. (MeOH). Mp 228-230° dec. (subl. at 200°).**Me ether, amide:** 2-Carbamoyl-3-methoxy-1,4-naphthoquinone

[144258-80-0]

C₁₂H₉NO₄ 231.207Constit. of *Rubia cordifolia*. Yellow cryst. (MeOH). Mp 150-151°.Bretschneider, H. *et al.*, *Monatsh. Chem.*, 1957, **88**, 652 (synth)Koyama, J. *et al.*, *Phytochemistry*, 1992, **31**, 2907 (isol, pmr, cmr, struct)**17-Hydroxy-10-nitroestr-4-en-3-one**

H-639

C₁₈H₂₅NO₄ 319.4**17β-form****Nitrohedon**


[61573-76-0]

Constit. of the male of fossil plant *Palaeogelos doleros*.

Roedder, W. *et al.*, *Pharm. Ztg.*, 1976, **121**, 479-484 (*isol*)

2-(4-Hydroxy-3-nitrophenyl)ethylamine H-640

4-(2-Aminoethyl)-2-nitrophenol, 9CI. 4-Hydroxy-2-nitrophenethylamine. 3-Nitrotyramine [49607-15-0]



C₈H₁₀N₂O₃ 182.179

Isol. from the cactus *Cereus validus*. Yellow orange prisms. Mp 217° dec.

Hydrochloride: [63195-80-2]

Yellow-brown plates (EtOH). Mp 214-215°.

Nitrate: [63195-79-9]

Mp 214-216° dec.

N-Ac: N-(4-Hydroxy-3-nitrophenylethyl)acetamide

C₁₀H₁₂N₂O₄ 224.216

Metab. of *Pyricularia oryzae*. Isol. from the marine-derived *Flavobacterium* sp. T436. Yellow needles (C₆H₆/hexane). Mp 95-98°.

Me ether: 2-(4-Methoxy-3-nitrophenylethyl)amine

C₉H₁₂N₂O₃ 196.205

Oil.

Me ether; hydrochloride:

Yellow needles (MeOH). Mp 231-232°.

N-(2-Methylpropanoyl): N-(2-Methylpropionyl)-3-nitrotyramine

[173791-66-7]

C₁₂H₁₆N₂O₄ 252.269

Isol. from a facultatively anaerobic, halophilic bacterium sediment from the Great Salt Plains, Oklahoma. Shows cytotoxicity against murine leukaemia P-388. Fine needles (CH₂Cl₂). Mp 99-100°. λ_{max} 274 (ε 8860); 357 (ε 3350) (MeOH) (Berdy).

N-(3-Methylbutanoyl): N-(3-Methylbutanoyl)-3-nitrotyramine

[173791-67-8]

C₁₃H₁₈N₂O₄ 266.296

Isol. from a facultatively anaerobic halophilic bacterium from sediment from the Great Salt Plains, Oklahoma. Fine needles (CH₂Cl₂). Mp 88-89°.

Waser, E. *et al.*, *Helv. Chim. Acta*, 1923, **6**, 55 (*synth*)

Callow, R.K. *et al.*, *J.C.S.*, 1929, 1453 (*Me ether*)

Neme, G. *et al.*, *Phytochemistry*, 1977, **16**, 277 (*isol*)

Sviridov, S.I. *et al.*, *Khim. Prir. Soedin.*, 1990, **26**, 811; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 691 (*isol, deriv*)

Fu, X. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1950 (*N*-2-Methylpropionyl, *N*-3-Methylbutanoyl)

Schuhmann, I. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*N-Ac, marine, isol*)

4-Hydroxy-3-nitrosobenzoic acid H-641

[32749-97-6]

C₇H₅NO₄ 167.121

Fe complex (2:1): Actinoveridin A

C₁₄H₈FeN₂O₈ 388.073

Isol. from *Streptomyces griseus*. Dark-green solid. Mp 300°. λ_{max} 268 (ε 26100); 295 (ε 22700); 433 (ε 7200); 686 (ε 6700) (MeOH) (Berdy).

Amide: [162152-90-1]

C₇H₆N₂O₃ 166.136

Prod. by *Streptomyces murayamensis*. Pale yellow cryst. (EtOH/hexane). Mp 145° dec. λ_{max} 233 (ε 11000); 280 (sh) (ε 5300); 373 (ε 2950) (MeOH). λ_{max} 233 (ε 11000); 373 (ε 2950) (MeOH) (Berdy). λ_{max} 232 (ε 11300); 288 (ε 4600); 372 (ε 3060) (MeOH-HCl) (Berdy). λ_{max} 206 (ε 10800); 267 (ε 13000); 449 (ε 4450) (MeOH-NAOH) (Berdy).


Cronheim, G. *et al.*, *J.O.C.*, 1947, **12**, 7 (*synth*)
Lazar, J. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1978, **98**, 327 (*synth*)

Kurobane, I. *et al.*, *J. Antibiot.*, 1987, **40**, 1131 (*isol, struct, bibl*)

Cone, M.C. *et al.*, *Tetrahedron*, 1995, **51**, 3095-3102 (*amide*)

4-Hydroxy-2-nonylquinoline H-642

2-Nonyl-4(1H)-quinolinone. 2-Nonyl-4-quinolinol. *Pyo Ic. Pseudane IX* [55396-45-7]



C₁₈H₂₅NO 271.402

Isol. from *Pseudomonas aeruginosa*. Also prod. by a bacterium isol. from the sponge *Suberea creba*. Antibacterial agent. Cryst. Sol. MeOH, Et₂O; poorly sol. H₂O, acids. Mp 138.8-139.2°. λ_{max} 213 (ε 23500); 236 (ε 26900); 316 (ε 10400); 327 (ε 10000) (MeOH) (Berdy). λ_{max} 236 (E1%/1cm 1050); 316 (E1%/1cm 431); 328 (E1%/1cm 415) (EtOH) (Berdy).

Oxalate: Mp 158-159°.

N-Oxide: [316-66-5]

C₁₈H₂₅NO₂ 287.401

Metab. of *Pseudomonas aeruginosa*. Leaflets (EtOH). Mp 148-149°.

1',2'-Didehydro(E)-: 2-(1-Nonenyl)-4(1H)-quinolinone. 4-Hydroxy-2-(1-nonenyl)quinoline. 2-(1-Nonenyl)-4-quinolinol. Δ¹-Pseudene IX. *Pyo III* [60783-01-9]

[1033-24-5]

C₁₈H₂₃NO 269.386

Isol. from *Pseudomonas aeruginosa*. Also prod. by a bacterium isol. from the sponge *Suberea creba*. Moderately active against gram-positive bacteria. Cryst. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 152.8-153.5°. λ_{max} 258 (ε 39000); 267 (ε 38000); 308 (ε 8700); 330 (ε 11000) (MeOH) (Berdy). λ_{max} 258 (E1%/1cm 1460); 266 (E1%/1cm 1376); 307 (E1%/1cm 340); 338 (E1%/1cm 434) (EtOH) (Berdy).

1',2'-Didehydro(Z)-:

C₁₈H₂₃NO 269.386

Metab. of *Pseudomonas aeruginosa*.

3',4',6',7'-Tetradehydro: 2-(3,6-Nonadienyl)-4(1H)-quinolinone. 4-Hydroxy-2-(3,6-nonadienyl)quinoline. 2-(3,6-Nonadienyl)-4-quinolinol [29479-51-4]

C₁₈H₂₁NO 267.37

Alkaloid from *Vepris ampody* (Rutaceae). Mp 103°. Config. of double bonds unknown.

9'-Hydroxy: 2-(9-Hydroxynonyl)-4(1H)-quinolinone

[29479-52-5]

C₁₈H₂₅NO₂ 287.401

Alkaloid from leaves of *Vepris ampody* (Rutaceae). Cryst. (Me₂CO). Mp 50°.

8'-Oxo: 2-(8-Oxononyl)-4(1H)-quinolinone

[263026-21-7]

C₁₈H₂₃NO₂ 285.385

Alkaloid from *Ruta montana*. λ_{max} 216 (log ε 4.5); 236 (log ε 4.2); 320 (log ε 3.8); 332 (log ε 3.8) (MeOH).

8'-Oxo, N-Me: 1-Methyl-2-(8-oxononyl)-4(1H)-quinolinone

[263026-23-9]

C₁₉H₂₅NO₂ 299.412

Alkaloid from *Ruta montana*. λ_{max} 220 (log ε 4.2); 235 (log ε 4.5); 320 (log ε 3.8); 332 (log ε 3.8) (MeOH).

NH-form

N-Me: 1-Methyl-2-nonyl-4(1H)-quinolinone

[68353-24-2]

C₁₉H₂₇NO 285.428

Alkaloid from *Ruta graveolens* (rue) and the ripe fruit of *Evodia rutaecarpa* (Rutaceae). Plates (C₆H₆/cyclohexane). Mp 71-75°.

N-Acetoxymethyl: 1-Acetoxymethyl-2-nonyl-4(1H)-quinolinone

[156547-73-8]

C₂₁H₂₉NO₃ 343.465

Alkaloid from aerial parts of *Boronia bowmanii* (Rutaceae). Amorph. solid. Mp 79°.

OH-form

8'-Oxo, *Me ether*: 4-Methoxy-2-(8-oxononyl)quinoline. 9-(4-Methoxy-2-quinolinyl)-2-nonanone

[263026-22-8]

C₁₉H₂₅NO₂ 299.412

Alkaloid from *Ruta montana*. λ_{max} 228 (log ε 4.5); 300 (log ε 2.9); 314 (log ε 2.8) (MeOH).

Hays, E.E. *et al.*, *J. Biol. Chem.*, 1945, **159**, 725-750 (*isol, uv*)

Wells, I.C. *et al.*, *J. Biol. Chem.*, 1952, **196**, 331-340 (*isol, struct, synth*)

Kan-Fan, C. *et al.*, *Phytochemistry*, 1970, **9**, 1283-1291 (9'-hydroxy, 3',4',6',7'-tetradehydro)

Ritter, C. *et al.*, *Eur. J. Biochem.*, 1971, **18**, 391-400 (*biosynth*)

Wratten, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1977, **11**, 411-414 (*isol*)


Kamikado, T. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 1515-1519 (1-Methyl-2-nonyl-4(1H)-quinolinone)

Budzikiewicz, H. *et al.*, *Monatsh. Chem.*, 1979, **110**, 947-953 (*isol, uv*)

Grundon, M.F. *et al.*, *Phytochemistry*, 1979, **18**, 1768-1769 (1-Methyl-2-nonyl-4(1H)-quinolinone)


- Somanathan, R. *et al.*, *J. Het. Chem.*, 1981, **18**, 1077-1079 (*synth, uv, ir, pmr*)
- Ahsan, M. *et al.*, *J. Nat. Prod.*, 1994, **57**, 670-672 (*1-Acetoxyethyl-2-nonyl-4(1H)-quinolinone*)
- Debitus, C. *et al.*, *J. Mar. Biotechnol.*, 1998, **6**, 136-141 (*isol*)
- Touati, D. *et al.*, *Phytochemistry*, 2000, **53**, 277-279 (*8'-oxo derivs*)
- Biaivatti, M.W. *et al.*, *J. Braz. Chem. Soc.*, 2002, **13**, 66-70 (*isol, pmr, cmr*)
- Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur, 1',2'-didehydro, oxide*)
- Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*biosynth*)
- Bredenbruch, F. *et al.*, *J. Bacteriol.*, 2005, **187**, 3630-3635 (*biosynth*)

3-Hydroxy-3-nonyl-2,4(1H,3H)-quinolinedione, 9CI H-643
[69808-31-7]




- $C_{18}H_{25}NO_3$ 303.4
Metab. from *Pseudomonas aeruginosa*.
Mp 146°.
- Neuhaus, W. *et al.*, *Z. Naturforsch., B*, 1979, **34**, 313 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

3-Hydroxy-11-norcytisine H-644
2,3,4,5-Tetrahydro-8-hydroxy-1,4-methanopyrido[1,2-a][1,4]diazepin-7(1H)-one, 9CI
[121063-35-2]



- $C_{10}H_{12}N_2O_2$ 192.217
Alkaloid from the green pods of *Laburnum anagyroides*.
- Hayman, A.R. *et al.*, *Phytochemistry*, 1989, **28**, 673-675 (*isol, pmr, ms*)
- Yohannes, D. *et al.*, *Org. Lett.*, 2008, **10**, 5353-5356 (*synth, pmr, cmr*)

N²-(17-Hydroxy-9,12,15-octadecatrienyl)glutamine H-645



$C_{23}H_{38}N_2O_5$ 422.564

(2S,9'Z,12'Z,15'Z,17'ξ)-form

- 17'-Phosphate: N²-(17-Phosphonoxylinolenoyl)glutamine*
 $C_{23}H_{39}N_2O_8P$ 502.544
Isol. from the oral secretions of *Spodoptera exigua*.
- 17'-(9Z,12Z-Octadecadienyl): N²-(17-*

Linoleoxylinolenoyl) glutamine

- $C_{41}H_{68}N_2O_6$ 684.998
Constit. of the oral secretions of the caterpillars *Spodoptera* spp. and *Heliothis virescens*.

17'-(9Z,12Z,15Z-Octadecatrienyl): N²-(17-Linoleoxylinolenoyl) glutamine

- $C_{41}H_{66}N_2O_6$ 682.982
Constit. of the oral secretions of the caterpillars *Spodoptera* spp. and *Heliothis virescens*.

15',16'-Dihydro, 17'-phosphate: N²-(17-Phosphonoxylinolenoyl) glutamine

- $C_{23}H_{41}N_2O_8P$ 504.559
Isol. from the oral secretions of *Spodoptera exigua*.

15',16'-Dihydro, 17'-(9Z,12Z-octadecadienyl): N²-(17-Linoleoxylinolenoyl) glutamine


- $C_{41}H_{70}N_2O_6$ 687.014
Constit. of the oral secretions of the caterpillars *Spodoptera* spp. and *Heliothis virescens*.

15',16'-Dihydro, 17'-(9Z,12Z,15Z-octadecatrienyl): N²-(17-Linolenoxylinolenoyl) glutamine

- $C_{41}H_{68}N_2O_6$ 684.998
Constit. of the oral secretions of the caterpillars *Spodoptera* spp. and *Heliothis virescens*.

- Spiteller, D. *et al.*, *J.O.C.*, 2003, **68**, 8743-8749; 2004, **69**, 1104-1108

3-Hydroxy-2-octadecylindole H-646
1,2-Dihydro-2-octadecyl-3H-indol-3-one. 2-Octadecyl-1H-indol-3-ol. Fistulosin



- $C_{26}H_{43}NO$ 385.632
Tautomeric with oxo form. Alkaloid from the roots of *Allium fistulosum* (Welsh onion). Antifungal agent. Cryst. Mp 80-83°. λ_{max} 220 ; 237 ; 252 ; 277 ; 302 (EtOH).

- Phay, N. *et al.*, *Phytochemistry*, 1999, **52**, 271-274 (*isol, uv, ir, pmr, cmr*)

8-Hydroxy-2,4,6-octatrienoic acid H-647

- $HOCH_2C\equiv CC\equiv CC\equiv CCOOH$
 $C_8H_4O_3$ 148.118

Amide: 8-Hydroxy-2,4,6-octatrienamide.


- Agrocybin†**
[544-44-5]
 $C_8H_5NO_2$ 147.133

- From *Agrocybe dura* and *Marasmius oreades* (fairy ring mushroom). Antifungal antibiotic. Phytotoxin. Cryst. (EtOH aq.). Sol. MeOH, EtOH, Me₂CO, Et₂O; fairly sol. H₂O; poorly sol. hexane. Mp 130-140° dec. λ_{max} 216 (ε 67600); 224 (ε 87000); 269 (ε 1740); 286 (ε 2400); 304 (ε 3020); 325 (ε 1950) (MeOH) (Berdy).

- Causes dermatitis. LD₅₀ (mus, ivn) 6 mg/kg. RH2100000

- Jones, E.R.H. *et al.*, *J.C.S.*, 1953, 3719
- Bu'Lock, J.D. *et al.*, *Chem. Ind. (London)*, 1954, 990 (*struct*)
- Ashworth, P.J. *et al.*, *J.C.S.*, 1958, 950 (*synth, ir*)
- Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1371 (*isol*)

4-Hydroxy-2-octylquinoline H-648
2-Octyl-4(1H)-quinolinone. 2-Octyl-4-quinolinol, 9CI
[678172-88-8]



- $C_{17}H_{23}NO$ 257.375
Metab. of *Pseudomonas aeruginosa*.

N-Oxide: 2-Octyl-4(1H)-quinolinone N-oxide. 4-Hydroxy-2-octylquinoline N-oxide. 2-Octyl-4-quinolinol N-oxide [2503-84-6]

- $C_{17}H_{23}NO_2$ 273.374
Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(E)-: 2-(1-Octenyl)-4(1H)-quinolinone. 4-Hydroxy-2-(1-octenyl)quinoline


- $C_{17}H_{21}NO$ 255.359
Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(Z)-:
 $C_{17}H_{21}NO$ 255.359
Metab. of *Pseudomonas aeruginosa*.
No CAS no. found to 2007.

- Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur, oxide, 1',2'-didehydro, ms*)

- Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*biosynth*)

6-(2-Hydroxy-2-oxido-4H-1,3,2-dioxaphosphorin-6-yl)-2,4(1H,3H)-pteridinedione, 9CI H-649
[149230-97-7]



- $C_9H_7N_4O_6P$ 298.151
Isol. from the marine polychaete *Odontosyllis undecimdonga*. Pale yellow cryst. Mp 225-235° dec.


N³-Me: [149230-96-6]
 $C_{10}H_9N_4O_6P$ 312.178
Isol. from *Odontosyllis undecimdonga*. Pale yellow needles (MeOH aq.). Mp 230-235° dec.

N⁷,N³-Di-Me: [149230-95-5]
 $C_{11}H_{11}N_4O_6P$ 326.205
Isol. from *Odontosyllis undecimdonga*. Pale yellow prisms (MeOH aq.). Mp 220-227° dec.

- Inou, S. *et al.*, *Heterocycles*, 1993, **35**, 147-150 (*isol, pmr, cmr, P-31 nmr*)

1-Hydroxy-4-oxo-2,5-cyclohexadiene-1-acetic acid

[55604-87-0]

C₈H₈O₄ 168.149

Constit. of *Ajuga parviflora*, *Pseudogynoxys cunninghamii*, *Jacaranda caucana* and the marine alga *Delesseria sanguinea*. Larvicide, leishmanicidal agent. Cryst. (EtOAc). Sol. Et₂O, MeOH; poorly sol. H₂O. Mp 103-104°. Log P -0.69 (calc). λ_{max} 224 (log ε 5.8) (MeOH).

Amide: *1-Hydroxy-4-oxo-2,5-cyclohexadiene-1-acetamide*. *4-(Carbamoylmethyl)-4-hydroxy-2,5-cyclohexadiene-1-one*. *Verongiaquinol*

[86254-99-1]

C₈H₉NO₃ 167.164

Parent compd. unknown.

Di-Me acetal, amide: *1-Hydroxy-4,4-dimethoxy-2,5-cyclohexadiene-1-acetamide*

C₁₀H₁₅NO₄ 213.233Isol. from *Verongia aerophoba*. Cryst. Mp 187-189°.

Di-Et acetal, amide: *4,4-Diethoxy-1-hydroxy-2,5-cyclohexadiene-1-acetamide*

[253308-66-6]

C₁₂H₁₉NO₄ 241.286Isol. from the sponge *Verongia aerophoba*. Cryst. Mp 183-187°.

Et-Me acetal, amide: *4-Ethoxy-1-hydroxy-4-methoxy-2,5-cyclohexadiene-1-acetamide*

C₁₁H₁₇NO₄ 227.26Isol. from *Verongia aerophoba*. Cryst. Mp 183-184°.

3-Bromo, amide: *3-Bromoverongiaquinol*

[137006-40-7]

C₈H₈BrNO₃ 246.06

Isol. from the marine sponge *Verongia cavernicola* and from *Suberea creba*. Sticky semi-solid. Sol. MeOH, C₆H₆; poorly sol. H₂O. Racemic. λ_{max} 235 (ε 5500) (MeOH) (Berdy).

3-Bromo, 5-chloro, amide: *3-Bromo-5-chloroverongiaquinol*

[98349-14-5]

C₈H₇BrClNO₃ 280.505Isol. from *Verongia cavernicola*. Cryst. Mp 175-177°. Racemic.

3,5-Dibromo, amide: *3,5-Dibromoverongiaquinol*

[16628-93-6]

C₈H₇Br₂NO₃ 324.956

Isol. from *Verongia cauliformis* (*Aplysina cauliformis*). Shows antibiotic props. Needles (Et₂O/Me₂CO). Mp 195-196°. λ_{max} 200 (sh) (ε) (MeOH/NaOH) (Derep). λ_{max} 257 (ε 8300) (MeOH) (Derep).

3,5-Dichloro, amide: *Dichloroverongiaquinol*

[88290-16-8]

H-650

C₈H₇Cl₂NO₃ 236.054

Isol. from the Mediterranean sponges *Aplysina cavernicola* and *Aplysina fistularis*. Inhibitor of gram-positive and gram-negative bacteria. Cryst. Mp 162-163°.

Evans, D.A. *et al.*, *J.O.C.*, 1977, **42**, 350-352 (*3,5-dibromo amide*)

D'Ambrosio, M. *et al.*, *Experientia*, 1983, **39**, 1091-1092 (*Dichloroverongiaquinol, isol, uv, ir, pmr, ms, synth*)

Fischer, A. *et al.*, *Tet. Lett.*, 1983, **24**, 131-134 (*3,5-dibromo amide, synth*)

D'Ambrosio, M. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 1484-1492 (*3-Bromoverongiaquinol*)

Goo, Y.M. *et al.*, *Arch. Pharmacol. Res.*, 1985, **8**, 21-30 (*Dichloroverongiaquinol*)


Bicchierini, N. *et al.*, *Tet. Lett.*, 1991, **32**, 4039-4040 (*bromoamide, dibromoamide, chlorobromoamide*)

Aydogmuş, Z. *et al.*, *Turk. J. Chem.*, 1999, **23**, 339-344 (*acetal amides*)

N-(3-Hydroxy-1-oxocyclopent-2-en-2-yl)-3-(4-hydroxy-3-methoxyphenyl)propenamide

2880-II

[117820-36-7]

C₁₅H₁₅NO₅ 289.287

Metab. of *Streptomyces griseoflavus*. Yellow powder. Sol. DMSO, DMF; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 272°. λ_{max} 268 (ε 30500); 338 (ε 28500) (MeOH/HCl) (Derep). λ_{max} 252 (ε 46100); 364 (ε 28800) (MeOH/NaOH) (Derep). λ_{max} 250 (ε 51800); 322 (ε 20900) (MeOH) (Derep). λ_{max} 250 (ε 51800); 322 (ε 20900) (MeOH) (Berdy). λ_{max} 268 (ε 51800); 338 (ε 28500) (MeOH-HCl) (Berdy). λ_{max} 252 (ε 46100); 364 (ε 28800) (MeOH-NaOH) (Berdy).


Grote, R. *et al.*, *J. Antibiot.*, 1988, **41**, 1275 (*isol, pmr, cmr, ir*)

Ebenezer, W.J. *et al.*, *Synth. Commun.*, 1991, **21**, 351 (*synth*)


Macdonald, G. *et al.*, *Tetrahedron*, 1998, **54**, 9823-9836 (*synth*)

8-Hydroxy-3-oxo-1,7(11)-eremophiladien-12,8-lactam

H-652

C₁₅H₁₉NO₃ 261.32**8αOH-form***Me ether*: [261164-36-7]C₁₆H₂₁NO₃ 275.347Constit. of *Senecio flavus*. Oil.Torres, P. *et al.*, *Phytochemistry*, 1999, **52**,1507-1513 (*isol, pmr, cmr*)**8-Hydroxy-1-oxo-7(11)-eremophilene-12,8-lactam**

H-653

C₁₅H₂₁NO₃ 263.336**(8αOH)-form***Me ether: 8-Methoxy-1-oxo-7(11)-eremophilene-12,8-lactam*

[849700-46-5]


C₁₆H₂₃NO₃ 277.363

Constit. of *Senecio aegyptius* var. *discoideus*. Oil. [α]_D²⁵ -58.7 (c, 2.74 in CHCl₃).

Mohamed, A.E.H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 439-442 (*Senecio aegyptius* constit)

6-Hydroxy-α-oxo-1H-indole-3-acetic acid

H-654

2-(6-Hydroxy-3-indolyl)-2-oxoacetic acidC₁₀H₇NO₄ 205.17*Me ester*: [956611-98-6]C₁₁H₉NO₄ 219.196


Alkaloid from a *Spongosorites* sp. Amorph. yellow powder.

Bao, B. *et al.*, *Mar. Drugs*, 2007, **5**, 31-39 (*isol, pmr, cmr, ms*)

25-Hydroxy-24-oxokibdelone C

H-655

[934464-85-4]



Relative Configuration

C₂₉H₂₉NO₁₂ 583.548

Poorly named. Prod. by *Kibdelosporangium* sp. (MST-108465). Possible artifact.

3-Me ether: 25-Methoxy-24-oxokibdelone C

[934464-84-3]


C₃₀H₃₁NO₁₂ 597.574

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. yellow solid. [α]_D²⁵ +164 (c, 0.01 in CHCl₃). Possible artifact. λ_{max} 208 (ε 21200); 221 (sh); 254 (ε 23100); 317 (sh); 388 (sh); 429

(ε 12100) (EtOH).

Ratnayake, R. *et al.*, *Chem. Eur. J.*, 2007, **13**, 1610-1619 (*isol*, *pmr*, *cmr*)**7-Hydroxy-3-oxo-3,7-secorhynchophylline** H-656


[128638-15-3]



Absolute Configuration

C₂₂H₂₈N₂O₆ 416.473Alkaloid from the leaves of *Uncaria salaccensis* (*Uncaria attenuata*) (Rubiaceae). Amorph. The nat. prod. is a mixt. of C-7 epimers.Ponglux, D. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 573-575 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *synth*, *struct*)Takayama, H. *et al.*, *Tetrahedron*, 1999, **55**, 6841-6846 (*config*)**4-Hydroxy-2-pentadecylquinoline** H-657

2-Pentadecyl-4(1H)-quinolinone. 2-Pentadecyl-4-quinolinol, 9CI

C₂₄H₃₇NO 355.562λ_{max} 215 (log ε 4.18); 239 (log ε 4.46); 321 (log ε 4.15); 334 (log ε 4.16) (MeOH).

N-Me: 1-Methyl-2-pentadecyl-4(1H)-quinolinone, 9CI

[59443-03-7]

C₂₅H₃₉NO 369.589Alkaloid from the leaves and fruits of *Evodia rutaecarpa* (Rutaceae). Leaflets (EtOAc/Et₂O). Mp 80°.

6',7'-Didehydro(Z-), N-Me: 1-Methyl-2-(6-pentadecenyl)-4(1H)-quinolinone [120693-51-8]

C₂₅H₃₇NO 367.573Alkaloid from fruits of *Evodia rutaecarpa* (Rutaceae). Oil. Isol. as an inseparable mixt. with the Δ^{10'}-isomer.

9',10'-Didehydro(Z-), N-Me: 1-Methyl-2-(9-pentadecenyl)-4(1H)-quinolinone [170661-70-8]

C₂₅H₃₇NO 367.573Alkaloid from fruits of *Evodia rutaecarpa*. Oil. Isol. as an inseparable mixt. with the Δ^{10'}-isomer.

10',11'-Didehydro(Z-), N-Me: 1-Methyl-2-(10-pentadecenyl)-4(1H)-quinolinone [120693-50-7]

C₂₅H₃₇NO 367.573Alkaloid from fruits of *Evodia rutaecarpa* (Rutaceae). Oil. Isol. as an inseparable mixt. with the Δ^{6'}-isomer.

6',7',9',10'-Tetradehydro(Z,Z-), N-Me: 1-Methyl-2-(6,9-pentadecadienyl)-4(1H)-quinolinone


[120693-52-9]

C₂₅H₃₅NO 365.558Alkaloid from fruits of *Evodia rutaecarpa* (Rutaceae). Oil.

6',7',9',10',12',13'-Hexadehydro(all-E)-: 2-(6,9,12-Pentadecatrienyl)-4(1H)-quinolinone, 9CI. Hapovine, 9CI [80981-97-1]

C₂₄H₃₁NO 349.515Alkaloid from *Haplophyllum popovii* (Rutaceae).Kamikado, T. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 605-609 (*N-Me*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)Razakova, D.M. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 528-529; *CA*, **96**, 100871k (*Hapovine*)Sugimoto, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 4453-4461 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)Tang, Y.-Q. *et al.*, *Phytochemistry*, 1996, **43**, 719-722 (*Evodia rutaecarpa constits*)Deziel, F. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*biosynth*)**4-Hydroxy-2-(1,3-pentadienyl)piperidine** H-658

2-(1,3-Pentadienyl)-4-piperidinol, 9CI



(1'E,2R,3'E,4R)-form

C₁₀H₁₇NO 167.25

(1'E,2R,3'E,4R)-form [261364-07-2]

Prod. by *Streptomyces* sp. GT 41006.Yellow cryst. [α]_D²⁰ +10.8 (c, 1.1 in MeOH). λ_{max} 230 (ε 17400) (MeOH).

(1'E,2S,3'E,4R)-form [137121-80-3]

Prod. by a *Streptomyces* sp. Cryst.(Et₂O). Mp 67-68°. [α]_D²⁰ -13 (c, 1 in CHCl₃) (natural). [α]_D²⁵ -37 (c, 1 in CHCl₃) (synthetic).

(1'E,2S,3'E,4S)-form

Antibiotic SS 20846A. SS 20846A


[104259-19-0]

From a *Streptomyces* sp. Inhibits intestine motility. Oil. Sol. MeOH, CHCl₃.[α]_D²⁰ -15 (c, 1 in CHCl₃) (natural). [α]_D²⁵ -20 (c, 1.4 in CHCl₃) (synthetic). λ_{max} 228 (MeOH) (Berdy). λ_{max} 228 (E1%/1cm 3100); 300 (E1%/1cm 50) (EtOH) (Berdy).

4-Deoxy: 2-(1,3-Pentadienyl)piperidine [137692-89-8]

C₁₀H₁₇N 151.251From a *Streptomyces* sp. [α]_D²⁰ +27 (c, 1 in CHCl₃).Japan. Pat., 1986, 86 35 788; *CA*, **105**, 132137 (2S,4S-form, *isol*, *struct*, *props*)Grabley, S. *et al.*, *J. Antibiot.*, 1991, **44**, 797-800 (2S,4R-form, *isol*)Takemoto, Y. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1906-1909 (*synth*)Ripoche, I. *et al.*, *J.C.S. Perkin I*, 1998, 3485-3492 (*synth*, *pmr*, *cmr*)Yokoyama, H. *et al.*, *Tet. Lett.*, 1998, **39**, 5971-5974 (*synth*)Maul, C. *et al.*, *J. Antibiot.*, 1999, **52**, 1124-1134 (2R,4R-form, *isol*, *uv*, *pmr*, *cmr*)Davis, F.A. *et al.*, *Org. Lett.*, 2000, **2**, 1041-1043 (*synth*)Sabat, M. *et al.*, *Tet. Lett.*, 2001, **42**, 1209-1212 (*synth*)**3-Hydroxy-4-pentenoic acid, 9CI** H-659

[81357-28-0]

C₅H₈O₃ 116.116

(R)-form [38996-04-2]

Oil or cryst. Mp 41.5-42.5°. Bp_{0.04} 69°. [α]_D¹⁷ -26 (c, 1 in 95% EtOH) (>99.8% op).

Nitrile: 3-Hydroxy-4-pentenenitrile, 9CI.

1-Cyano-3-buten-2-ol

[7451-85-6]

C₅H₇NO 97.116Produced by enzymatic hydrol. of Glucorapiferin (Progoitrin). Isol. from *Crambe abyssinica* seed meal. Shows phytotoxic activity. Oil.

Nitrile, 1-naphthylurethane:

Cryst. Mp 89-92°.

(S)-form [38996-05-3]

Et ester: [349649-09-8]

C₇H₁₂O₃ 144.17Bp_{0.125} 42°. [α]_D²⁰ -5 (c, 0.98 in CHCl₃) (>99% ee).Nitrile: **Crambene**

[6071-81-4]

Found in cruciferous vegetables incl. *Crambe abyssinica*. Prod. by enzymatic hydrol. of Epiglucoarapiferin (Epiprogoitrin). Oil. n_D²⁰ 1.4544.▶ LD₅₀ (rat, scu) 200 mg/kg. Exp. teratogen. SB4500000

Nitrile, 1-naphthylurethane:

Cryst. (C₆H₆/heptane). Mp 93°.

(±)-form [38996-03-1]

Bp_{0.07} 74-76°. n_D³⁰ 1.4633.

Et ester: [38996-01-9]

[134896-00-7]

C₇H₁₂O₃ 144.17Liq. Bp_{0.6} 57°.

tert-Butyl ester: [122763-67-1]

[159787-33-4]

C₉H₁₆O₃ 172.224Bp_{0.04} 35°. n_D²⁰ 1.4374.

Nitrile: [27451-36-1]

Bp₃₀ 132-133°.

▶ SB2270000

Nitrile, 1-naphthylurethane: Mp 112.5-113.5°.


[27451-36-1]

Bissinger, W.E. *et al.*, *J.A.C.S.*, 1947, **69**, 2955-2961 (*nitrile*, *synth*)Daxenbichler, M.E. *et al.*, *Biochemistry*, 1966, **5**, 692-697 (*isol*, *pmr*, *ord*, *nitrile*)Spencer, G.F. *et al.*, *J. Sci. Food Agric.*, 1980, **31**, 359-367 (*ms*, *nitrile*)Das, N.B. *et al.*, *Tetrahedron*, 1983, **39**, 2247-2253 (*synth*, *nitrile*)

- Graf, S. *et al.*, *Annalen*, 1993, 1091-1098 (synth, pmr, cmr)
 Vaughn, S.F. *et al.*, *J. Chem. Ecol.*, 1998, **24**, 1117-1126 (nitrile, isol, activity)
Org. Synth., Coll. Vol., 9, 1998, 423-435 (Et ester, synth, ir, pmr)
 Jeffery, E.H. *et al.*, *Handbook of Nutraceuticals and Functional Foods*, (ed. Wildman, R.E.C.), CRC Press, 2001, 169-191
 Niedoborski, T.E. *et al.*, *J. Agric. Food Chem.*, 2001, **49**, 3394-3399 (Crambene, isol, purifn)
 Song, J. *et al.*, *Tetrahedron: Asymmetry*, 2001, **12**, 387-391 (S-form, synth, Et ester)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, COP400

4-Hydroxy-2-pentylquinoline H-660

2-Pentyl-4(1H)-quinolinone. 2-Pentyl-4-quinolinol
 [62869-70-9]



C₁₄H₁₇NO 215.294
 Alkaloid from *Boronia ternata* var. *elongata*, *Pseudomonas aeruginosa* and a yellow marine pseudomonad. Antibacterial agent. Cryst. Sol. MeOH, CHCl₃. Mp 146° (141-142°). λ_{max} 235 (log ε 5.13); 282 (log ε 4.12); 294 (log ε 4.28); 310 (log ε 4.63); 315 (log ε 4.74); 327 (log ε 4.71) (MeOH).

N-Oxide: 2-Pentyl-4(1H)-quinolinone N-oxide. 4-Hydroxy-2-pentylquinoline N-oxide
 [91292-96-5]

C₁₄H₁₇NO₂ 231.294

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(E-): 2-(1E-Pentenyl)-4(1H)-quinolinone. 4-Hydroxy-2-(1E-pentenyl)quinoline
 C₁₄H₁₅NO 213.279

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(Z-): 2-(1Z-Pentenyl)-4(1H)-quinolinone. 4-Hydroxy-2-(1Z-pentenyl)quinoline
 C₁₄H₁₅NO 213.279

Metab. of *Pseudomonas aeruginosa*.

NH-form

N-Me: 1-Methyl-2-pentyl-4(1H)-quinolinone
 [22048-98-2]

C₁₅H₁₉NO 229.321

Alkaloid from *Boronia ternata* var. *elongata* and fruit of *Evodia rutaecarpa*. Powder (EtOAc/Et₂O) or brownish oil. Mp 65-66°. λ_{max} 213 (log ε 4.4); 239 (log ε 4.46); 321 (log ε 4.13); 333 (log ε 4.15) (MeOH). λ_{max} 213 (log ε 4.62); 235 (log ε 4.67); 316 (log ε 4.14); 325 (log ε 4.13) (MeOH).

8-Methoxy, N-Me: 8-Methoxy-1-methyl-2-pentyl-4(1H)-quinolinone
 [159979-55-2]

C₁₆H₂₁NO₂ 259.347

Alkaloid from trunk bark of *Esenbeckia almarillia*. Oil.

OH-form

Me ether: 4-Methoxy-2-pentylquinoline
 [22048-99-3]

C₁₅H₁₉NO 229.321

Alkaloid from *Galipea officinalis* and *Galipea longiflora*. Oil. Bp_{0.14} 135°.

1',2'-Didehydro, Me ether: 4-Methoxy-2-(1-pentenyl)quinoline
 [124902-97-2]

C₁₅H₁₇NO 227.305

Alkaloid from stem bark of *Galipea bracteata*. Amorph.

Buu-Hoi, N.G. *et al.*, *J.O.C.*, 1953, **18**, 1209-1224 (synth)

Werner, W. *et al.*, *Tetrahedron*, 1969, **25**, 255-261 (synth)

Kimimado, T. *et al.*, *Agric. Biol. Chem.*, 1976, **40**, 605-609 (N-Me)

Wratten, S.J. *et al.*, *Antimicrob. Agents Chemother.*, 1977, **11**, 411-414 (isol, struct)

Thomsen, I. *et al.*, *Acta Chem. Scand., Ser. B*, 1988, **42**, 309-313 (synth, pmr)

Sugimoto, T. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 4453-4461 (N-Me)

Fournet, A. *et al.*, *Can. J. Chem.*, 1989, **67**, 2116-2118 (4-Methoxy-2-pentylquinoline, 4-Methoxy-2-(1-pentenyl)quinoline)

Guilhon, G.M.S.P. *et al.*, *Phytochemistry*, 1994, **37**, 1193-1195 (8-Methoxy-1-methyl-2-pentyl-4(1H)-quinolinone)


Lepine, F. *et al.*, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (occur, oxide, 1',2'-didehydro, ms)

Deziel, E. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (biosynth)

Agier, C. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 698-703 (isol, *Boronia*, pmr, cmr, ms)

2-Hydroxy-1-phenazinecarboxylic acid, 9CI H-661

[4075-25-6]



C₁₃H₈N₂O₃ 240.218

Pigment produced by *Pseudomonas aerofaciens*. Shows antimicrobial activity. Fine orange needles (C₆H₆). Sol. bases. Mp 225-226° dec. (darkens >220°). pK_a 7.6 (66.8% dioxan aq.). λ_{max} 252 ; 363 (MeOH) (Berdy). λ_{max} 287 ; 363 ; 490 (MeOH/NaOH) (Berdy).

Me ester: [13389-20-3]

C₁₄H₁₀N₂O₃ 254.245

Cryst. (MeOH). Mp 140-141°.

Me ether: 2-Methoxy-1-phenazinecarboxylic acid

C₁₄H₁₀N₂O₃ 254.245

Cryst. (C₆H₆). Mp 204-205°.

Me ether, Me ester: [13392-00-2]

C₁₅H₁₂N₂O₃ 268.271

Yellow cryst. (CCl₄). Mp 129° subl.

Olson, E.S. *et al.*, *J.O.C.*, 1967, **32**, 2887 (isol, uv, ir, struct)

Herbert, R.B. *et al.*, *Tet. Lett.*, 1968, 1907 (synth)

Brooke, P.K. *et al.*, *J.C.S. Perkin 1*, 1976, 2248 (synth)

Herbert, R.P. *et al.*, *J.C.S. Perkin 1*, 1979, 2411 (biosynth)

4-Hydroxy-1-phenazinecarboxylic acid H-662

[79417-75-7]

C₁₃H₈N₂O₃ 240.218

Minor pigment from *Pseudomonas aerofaciens*.

Römer, A. *et al.*, *Z. Naturforsch., B*, 1981, **36**, 1037 (isol, uv, ir, pmr, ms, struct)

6-Hydroxy-1-phenazinecarboxylic acid H-663

C₁₃H₈N₂O₃ 240.218

Me ether, Me ester: 6-Methoxy-1-phenazinecarboxylic acid methyl ester, 9CI.

Mycomethoxin B

[39011-76-2]

C₁₅H₁₂N₂O₃ 268.271

Isol. from *Streptomyces luteoreticuli*.

Active against mycobacteria. Yellow needles. Mp 150-151° (139-141°).

Yamanaka, S. *et al.*, *CA*, 1972, **77**, 162986 (isol)

Brooke, P.K. *et al.*, *J.C.S. Perkin 1*, 1976, 2248 (synth, uv, pmr)

Buckland, P.R. *et al.*, *J. Chem. Res., Synop.*, 1981, 363 (biosynth)

Roemer, A. *et al.*, *Org. Magn. Reson.*, 1982, **19**, 66 (pmr)

9-Hydroxy-1-phenazinecarboxylic acid H-664

[23462-26-2]

C₁₃H₈N₂O₃ 240.218

Formed by *Pseudomonas phenazinium*. Mp 270°.

Me ether: 9-Methoxy-1-phenazinecarboxylic acid

[23531-25-1]

C₁₄H₁₀N₂O₃ 254.245

Mp 258-261°.

Me ether, Me ester:

C₁₅H₁₂N₂O₃ 268.271

Mp 115-120°.

Ph ether: 9-Phenoxy-1-phenazinecarboxylic acid

[103942-79-6]

C₁₉H₁₂N₂O₃ 316.315

Solid. Mp 232-234°.


Yoshida, K. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 703

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1969, **6**, 297

Byng, G.S. *et al.*, *Biochem. Soc. Trans.*, 1975, **3**, 742

Brooke, P.K. *et al.*, *J.C.S. Perkin 1*, 1976, 2248 (synth)

Spicer, J.A. *et al.*, *J. Med. Chem.*, 2000, **43**, 1350-1358 (Ph ether, synth, pmr)

2-Hydroxy-1,6-phenazinedicarboxylic acid H-665

C₁₄H₈N₂O₅ 284.228

1'-Me ester: **Antibiotic SB 212021**. SB 212021

[170591-55-6]

C₁₅H₁₀N₂O₅ 298.254

Prod. by a *Streptomyces* sp. Inhibitor of metalloenzymes. Sol. MeOH, bases, butanol; poorly sol. H₂O. λ_{\max} 210 (€ 23563); 247 (€ 11680); 294 (€ 17672); 370 (€ 2840); 520 (€ 1178) (MeOH) (Berdy). λ_{\max} 203 (€ 20145); 248 (€ 16421); 268 (€ 17914); 369 (€ 5938); 431 (€ 2582) (MeOH/HCl) (Berdy). λ_{\max} 210 (€ 73271); 246 (€ 17554); 288 (€ 19778); 370 (€ 4000); 520 (€ 2528) (MeOH/NaOH) (Berdy).

[83297-76-1]

Roemer, A. et al., *Org. Magn. Reson.*, 1982, **19**, 66 (pmr, deriv)Gilpin, M.L. et al., *J. Antibiot.*, 1995, **48**, 1081 (isol, deriv)**4-Hydroxy-1,6-phenazinedicarboxylic acid** H-666C₁₄H₈N₂O₅ 284.228

Di-Me ester: [79417-76-8]

C₁₆H₁₂N₂O₅ 312.281Minor pigment from *Pseudomonas cepacia*.


Me ether, di-Me ester:

C₁₇H₁₄N₂O₅ 326.308

Mp 147° (142-144°).

Römer, A. et al., *Z. Naturforsch., B*, 1981, **36**, 1037 (isol, uv, ir, pmr, ms, struct, synth)**4-Hydroxyphenethylamine** H-6674-(2-Aminoethyl)phenol, 9CI. 2-(p-Hydroxyphenyl)ethylamine. **Tyramine**. *Tyrosamine*

[51-67-2]

C₈H₁₁NO 137.181

Widespread biogenic amine, found in plant spp. such as *Magnolia*, *Desmodium* spp., *Pisum sativum*, *Hordeum vulgare*, *Actinodaphne* sp., *Cannabis sativa*, *Piper nigrum* (pepper) mescal (*Lophophora williamsii*) and other cacti and in putrefied animal tissues. Found in urine of patients with Parkinson's disease. Diagnostic aid. Vasopressor. Mp 164-164.5° (161°). pK_{a1} 9.3; pK_{a2} 10.9 (25°). Log P 0.62 (calc).

▶ LD₅₀ (mus, ivn) 229 mg/kg. SJ5950000

Hydrochloride: [60-19-5]

Mp 269°.

▶ SJ6050000

N-Ac: N-[2-(4-Hydroxyphenyl)ethyl]acetamide. N-(p-Hydroxyphenethyl)acetamide, 8CI. **N-Acetyltyramine**

[1202-66-0]

C₁₀H₁₃NO₂ 179.218

Metab. of pathogenic fungi, mycobacteria, enterobacteria and from *Bombyx mori* at the chrysalis stage. Mp 134-135° (sinters at 128°).

▶ AC3852300

N-Ac, O-[4-methoxy-E-cinnamoyl-(→6)-β-D-glucopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→3)]-α-L-rhamnopyranoside]: [952340-41-9]

C₃₈H₅₁NO₁₈ 809.817Constit. of the aerial parts of *Schnabelia tetradonta*.

N-Ac, O-[4-methoxy-Z-cinnamoyl-(→6)-β-D-glucopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→3)]-α-L-rhamnopyranoside]:

C₃₈H₅₁NO₁₈ 809.817

Constit. of the aerial parts of *Schnabelia tetradonta*. Needles (MeOH). Mp 198-199°. [α]_D²⁵ -14.1 (c, 0.75 in MeOH).

N-Ac, O-[4-methoxy-E-cinnamoyl-(→6)-β-D-glucopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→3)]-4-O-acetyl-α-L-rhamnopyranoside]: [952340-42-0]

C₄₀H₅₃NO₁₉ 851.854Constit. of the aerial parts of *Schnabelia tetradonta*. CA name incorrect.

N-Ac, O-[4-methoxy-Z-cinnamoyl-(→6)-β-D-glucopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→3)]-4-O-acetyl-α-L-rhamnopyranoside]:

C₄₀H₅₃NO₁₉ 851.854

Constit. of the dried roots of *Schnabelia tetradonta*. Needles (MeOH). Mp 156-158°. [α]_D²⁵ -16.4 (c, 0.19 in MeOH).

O,N-Di-Ac: [14383-56-3]

C₁₂H₁₅NO₃ 221.255

Mp 100.5°.

N-(2-Methylbutanoyl): N-[2-(4-Hydroxyphenyl)ethyl]-2-methylbutanamide. N-(2-Methylbutanoyl)tyramine. **Antibiotic YUA 001**. YUA 001

[218153-66-3]

C₁₃H₁₉NO₂ 221.299

Prod. by *Corynebacterium* sp. YUA 25. Aldose reductase inhibitor. Powder. Mp 103°. λ_{\max} 224; 276 (MeOH).

N-Tigloyl: N-[2-(4-Hydroxyphenyl)ethyl]tiglamide. N-Tigloyltyramine

C₁₃H₁₇NO₂ 219.283Constit. of the wood of *Hortia colombiana*.

N-(5Z,8Z,11Z-Tetradecatrienoyl): N-(5,8,11-Tetradecatrienoyl)tyramine

C₂₂H₃₁NO₂ 341.492Isol. from *Muricea austera*. Oil.N-Tetracosanoyl: **Violyedoamide**. N-[2-(4-Hydroxyphenyl)ethyl]tetracosanamide, 9CI. N-Tetracosanoyltyramine

[113122-70-6]

C₃₂H₅₇NO₂ 487.808Constit. of *Viola yedoensis*.N-Benzoyl: N-[2-(4-Hydroxyphenyl)ethyl]benzamide, 9CI. **N-Benzoyltyramine**

[41859-54-5]

C₁₅H₁₅NO₂ 241.289

Alkaloid from seeds of the Mexican apple (*Casimiroa edulis*) and from *Aniba riparia*. Plates (Et₂O). Mp 161-162°.

N-(2-Hydroxybenzoyl): N-(2-Hydroxybenzoyl)tyramine

[112356-52-2]

C₁₅H₁₅NO₃ 257.288Constit. of the fruit of *Aniba riparia*.

Cryst. (Et₂O). Mp 135-137°. λ_{\max} 225 (€ 5650); 240 (€ 4000); 305 (€ 1650) (MeOH).

O-(3,4,5-Trihydroxybenzoyl): **O-Gallolyltyramine**C₁₅H₁₅NO₅ 289.287

Alkaloid from the leaves of *Cupaniopsis macropetala*. Pim2 enzyme inhibitor. Brown oil (as trifluoroacetate salt). λ_{\max} 212 (log € 4.68); 272 (log € 4.3) (MeOH) (trifluoroacetate).

N-Cinnamoyl: **N-Cinnamoyltyramine**. N-[2-(4-Hydroxyphenyl)ethyl]cinnamide

[103188-44-9]

[20384-14-9]

C₁₇H₁₇NO₂ 267.327Alkaloid from *Piper steerni*. Cryst.

(EtOH). Mp 194-195° (synthetic).

N-(4-Hydroxy-E-cinnamoyl): **N-p-trans-Coumaroyltyramine**. N-(p-Hydroxyphenyl)ethyl p-hydroxycinnamide. *Parprazine*

[36417-86-4]

C₁₇H₁₇NO₃ 283.326

Constit. of *Actinodaphne* spp., *Allium chinense*, *Cannabis sativa*, *Evodia lahae*, *Fumaria indica* and *Vicia faba*. Cryst. (MeOH). Mp 260-261°.

▶ LD₅₀ (mus, ipr) 780 mg/kg.N-(4-Hydroxy-Z-cinnamoyl): **N-p-cis-Coumaroyltyramine**

[111129-11-4]

C₁₇H₁₇NO₃ 283.326

Alkaloid from *Allium chinense* and *Aristolochia kankauensis*. Needles (CHCl₃/MeOH). Mp 234-235°.

N-(3,4-Dihydroxy-E-cinnamoyl): **N-trans-Caffeoyltyramine**

[103188-48-3]

C₁₇H₁₇NO₄ 299.326

Constit. of *Annona crassiflora*, *Annona montana*, *Mangifera indica* and *Cannabis sativa*. Powder. Mp 219.5-222.5° (214-215°). λ_{\max} 221 (€ 14125); 294 (€ 10960); 321 (€ 12020) (MeOH) (Berdy). λ_{\max} 208 (€ 12500); 225 (€ 13182); 242 (€ 10300); 308 (€ 7080); 349 (€ 13800) (MeOH/NaOH) (Berdy).

N-(3,4-Dihydroxy-Z-cinnamoyl): **N-cis-Caffeoyltyramine**

[219773-48-5]

C₁₇H₁₇NO₄ 299.326

Alkaloid from stems of *cherimoya* (*Annona cherimola*). Yellow oil. λ_{\max} 203 (log € 2.43); 220 (sh) (log € 3.36); 280 (log € 2.53); 310 (log € 3.21) (EtOH).

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyltyramine**. *Moupinamide*

[66648-43-9]

[65646-26-6]


C₁₈H₁₉NO₄ 313.352

Alkaloid from *Aristolochia moupinensis*, *Cannabis sativa*, *Asimina triloba*, *Annona montana*, *Tribulus terrestris*, *Piper nigrum*, *Canganga odorata* and other plant spp. Platelet aggregation inhibitor, hepatoprotective. Cryst. (Me₂CO/CHCl₃). Mp 144.5-145°.

- N-(4-Hydroxy-3-methoxy-Z-cinnamoyl):
N-cis-Feruloyltyramine
[80510-09-4]
C₁₈H₁₉NO₄ 313.352
Alkaloid from *Aristolochia kankauensis*, *Capsicum annuum* var. *grossum* and *Tinospora tuberculata*. Pale yellow oil.
- N-(4-Hydroxy-3,5-dimethoxy-E-cinnamoyl): **N-trans-Sinapoyltyramine**
[200125-11-7]
C₁₉H₂₁NO₅ 343.379
Alkaloid from *Porcelia macrocarpa*. Amorph. powder. λ_{max} 205 (log ε 4.21); 225 (log ε 4.14); 320 (log ε 3.96) (MeOH).
- N-(4-Hydroxy-3,5-dimethoxy-Z-cinnamoyl): **N-cis-Sinapoyltyramine**
C₁₉H₂₁NO₅ 343.379
Alkaloid from *Lindera glauca*. Amorph. powder. λ_{max} 225; 320 (EtOH).
- N-[3-(3,4-Dihydroxyphenyl)propanoyl]: **N-Dihydrocaffeoyltyramine**
C₁₇H₁₉NO₄ 301.341
Alkaloid from the root bark of *Lycium chinense* (Chinese boxthorn).
- N-[3-(4-Hydroxy-3-methoxyphenyl)propanoyl]: **N-Dihydroferuloyltyramine**
[184877-32-5]
C₁₈H₂₁NO₄ 315.368
Alkaloid from stems of cherimoya (*Annona cherimola*). Amorph. powder. Mp > 300°. λ_{max} 206 (log ε 2.73); 255 (sh) (log ε 3.65); 280 (log ε 2.34) (EtOH).
- N-Jasmonoyl: **N-Jasmonoyltyramine**
[202661-57-2]
C₂₀H₂₇NO₃ 329.438
Constit. of the pollen of *Petunia hybrida*.
- N-Me: 4-[2-(Methylamino)ethyl]phenol, **9CI. N-Methyltyramine**
[370-98-9]
C₉H₁₃NO 151.208
Alkaloid from *Hordeum vulgare*, *Panicum miliaceum* (proso millet), some *Coryphantha* spp. and several other spp. Also found in mescal (*Lophophora williamsii*), *Trichocereus* spp. and other cacti (Poaceae, Cactaceae). Mp 130-131° (127-128°).
- ▶ SL8300000
- N-Me, O-α-L-rhamnopyranoside: **N-Methyltyramine O-α-L-rhamnopyranoside**
[111537-50-9]
C₁₅H₂₃NO₅ 297.35
Alkaloid from *Selaginella doederleinii* (Selaginellaceae). Amorph. solid. [α]_D²⁰ -131 (c, 0.3 in MeOH).
- N-Me, N-formyl: **N-Formyl-N-methyltyramine**
C₁₀H₁₃NO₂ 179.218
Alkaloid from *Cyatobasis fruticulosa*. Needles (MeOH). Mp 112-114°. λ_{max} 226 (log ε 2.8); 270 (log ε 2.8) (MeOH).
- N,N-Di-Me: see Hordenine, H-357
- Me ether: see 2-(4-Methoxyphenyl)ethylamine, M-280
- O-(3-Methyl-2-butenyl), N-tigloyl: N-[2-(4-Prenyloxyphenyl)ethyl]tiglamide
[172837-74-0]
C₁₈H₂₅NO₂ 287.401
Constit. of *Boronia megastigma* (brown boronia). Cryst. (Et₂O). Mp 69-71°.
- O-(3-Methyl-2-butenyl), N-benzoyl: **Hortiamide**
[106055-13-4]
C₂₀H₂₃NO₂ 309.407
Alkaloid from roots of *Hortia regia* (Rutaceae). Cryst. Mp 109-111°.
- O-(3-Methyl-2-butenyl), N-E-cinnamoyl: N-[2-[4-[(3-Methyl-2-butenyl)oxy]phenyl]ethyl]-3-phenyl-2-propenamido. **N-2-[4-(3,3-Dimethylallyloxy)phenyl]ethylcinnamide**. N-Cinnamoyl-p-prenyloxyphenethylamine
[87596-51-8]
C₂₂H₂₅NO₂ 335.445
Alkaloid from the leaves of *Aegle marmelos* (bael fruit) (Rutaceae). Cryst. (MeOH). Mp 125-126°.
- [70185-64-7, 20375-37-5]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1289C; 1289D (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 612A; 612B (*nmr*)
Barger, G. *et al.*, *J.C.S.*, 1909, **95**, 1127; 1722 (*synth*)
Kincl, F.A. *et al.*, *J.C.S.*, 1956, 4163 (*N-Benzoyltyramine*)
Butenandt, A. *et al.*, *Arch. Biochem. Biophys.*, 1959, **83**, 76 (*N-Acetyltyramine*)
Mann, J.D. *et al.*, *J. Biol. Chem.*, 1963, **238**, 676-681 (*N-Me, isol*)
Kappe, T. *et al.*, *J. Med. Chem.*, 1965, **8**, 368 (*uv*)
Boulton, A.A. *et al.*, *Nature (London)*, 1967, **215**, 132 (*occur*)
Rondest, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 2411-2414 (*N-4-hydroxycinnamoyl*)
Takahashi, T. *et al.*, *Agric. Biol. Chem.*, 1970, **34**, 1256-1258 (*isol, sawa millet*)
Stuart, K.L. *et al.*, *Phytochemistry*, 1971, **10**, 460 (*isol, derivs*)
Milne, G.W. *et al.*, *Anal. Chem.*, 1973, **45**, 1952 (*ms*)
Tamura, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1974, **47**, 2682 (*cryst struct*)
Lambert, F. *et al.*, *Org. Magn. Reson.*, 1975, **72**, 66 (*pmr*)
Kruger, T.L. *et al.*, *J.O.C.*, 1977, **42**, 4161 (*derivs, ms*)
Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*rev, occur, derivs*)
Doetsch, P.W. *et al.*, *J. Chromatogr.*, 1980, **189**, 79 (*occur*)
Yoshihara, T. *et al.*, *Agric. Biol. Chem.*, 1981, **45**, 2593-2598 (*N-Feruloyltyramines*)
Fukuda, N. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 156-161 (*N-Feruloyltyramines*)
Govindachari, T.R. *et al.*, *Phytochemistry*, 1983, **22**, 755 (*N-cinnamoyl-p-prenyl*)
Xu, L. *et al.*, *Yaoxue Xuebao*, 1984, **19**, 48; *CA*, **101**, 107360y (*Moupinamide*)
Chao, L.R. *et al.*, *J. Nat. Prod.*, 1987, **50**, 477 (*N-Methyltyramine rhamnopyranoside*)
Barbosa-Filho, J.M. *et al.*, *Phytochemistry*, 1987, **26**, 2615-2617 (*N-2-hydroxybenzoyl*)
Xiao, Y. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1987, **29**, 532; *CA*, **108**, 109524y (*Violydoenamido*)
Tseng, C.-F. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 396-400 (*synth, cinnamic amides*)
Tinto, W.F. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1676 (*Hortiamide*)
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1992, **31**, 2869-2872 (*Paprazine*)
Martindale, *The Extra Pharmacopoeia, 30th edn.*, Pharmaceutical Press, 1993, 780
Weyerstahl, P. *et al.*, *Amalgen*, 1994, 1043 (*Prenyloxyphenylethyltiglamide*)
Wu, T.-S. *et al.*, *Phytochemistry*, 1994, **36**, 1063 (*N-p-cis-Coumaroyltyramine*)
- Posso, O. *et al.*, *Rev. Colomb. Quim.*, 1994, **23**, 53-62; *CA*, **122**, 286624r (*N-Cinnamoyltyramine*)
Wu, Y.-C. *et al.*, *Planta Med.*, 1995, **61**, 146 (*N-Caffeoyltyramine*)
Nishioka, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 1138-1141 (*N-p-Coumaroyltyramine, isol, synth*)
Chaves, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 879-881 (*N-Sinapoyltyramine*)
Bahn, Y.-S. *et al.*, *J. Antibiot.*, 1998, **51**, 902-907 (*YUA 001*)
Miersch, O. *et al.*, *Phytochemistry*, 1998, **47**, 327-329 (*N-Jasmonoyltyramine*)
Chen, C.-Y. *et al.*, *Phytochemistry*, 1998, **49**, 1443-1447 (*Annona cherimola amides*)
Kalač, P. *et al.*, *Biologically-active Phytochemicals in Food*, (eds. Pfannhauser, W. *et al.*), Royal Society of Chemistry, 2001, 217-220 (*sauerkraut*)
Chang, Y.-C. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 811-815 (*N-Sinapoyltyramines, N-Feruloyltyramines*)
Lee, H.H. *et al.*, *Arch. Pharmacol. Res.*, 2002, **25**, 428-432 (*N-Dihydrocaffeoyltyramine*)
Suarez, L.E.C. *et al.*, *J. Braz. Chem. Soc.*, 2002, **13**, 339-344 (*Hydroxyphenylethyltiglamide*)
Dou, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1777-1781 (*Schnabelia tetradonta C38 glycoside*)
Dou, H. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2797-2804 (*Schnabelia tetradonta C40 glycoside*)
Bahçeevli, A.K. *et al.*, *J. Nat. Prod.*, 2005, **68**, 956-958 (*N-Formyl-N-methyltyramine*)
Gutiérrez, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1379-1383 (*Muricea austera constit*)
Li, B. *et al.*, *Tianran Chanwu Yanjiu Yu Kaifa*, 2006, **18**, 61-64; *CA*, **147**, 443770w (*Schnabelia trans-cinnamoyl glycosides*)
Zhao, P.-J. *et al.*, *Chem. Biodiversity*, 2007, **4**, 899-904 (*N-Ac, isol, pmr, cmr*)
Davis, R.A. *et al.*, *J. Nat. Prod.*, 2008, **71**, 451-452 (*Gallolytyramine*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, TOF750; TOG250

N-(4-Hydroxyphenethyl)-2,4- H-668 decadienamide

2,4-Decadienoic acid p-hydroxyphenethylamide



C₁₈H₂₅NO₂ 287.401

(E,E)-form [24738-52-1]

Constit. of the roots of *Anacyclus pyrethrum* (Asteraceae). Mp 132-133°.

Burden, R.S. *et al.*, *J.C.S. (C)*, 1969, 2477 (*isol, uv, ir, pmr, struct, synth*)

N-(4-Hydroxyphenethyl)-2,4- H-669 dodecadienamide

2,4-Dodecadienoic acid p-hydroxyphenethylamide

As N-(4-Hydroxyphenethyl)-2,4-decadienamide, H-668 with n = 6

C₂₀H₂₉NO₂ 315.455

(E,E)-form [24738-53-2]

Constit. of the roots of *Anacyclus pyrethrum* (Asteraceae). Mp 140-141°.

Burden, R.S. *et al.*, *J.C.S.(C)*, 1969, 2477 (*isol. synth*)

N-(4-Hydroxyphenethyl)-2,4-tetradecadienamide H-670

2,4-Tetradecadienoic acid p-hydroxyphenethylamide

As N-(4-Hydroxyphenethyl)-2,4-decadienamide, H-668 with n = 8

C₂₂H₃₃NO₂ 343.508

(E,E)-form [24738-54-3]


Constit. of the roots of *Anacyclus pyrethrum* (Asteraceae). Mp 141.5-142.5°.

Burden, R.S. *et al.*, *J.C.S.(C)*, 1969, 2477 (*isol. synth*)

2-Hydroxy-2-phenylacetamide H-671

α-Hydroxybenzeneacetamide, 9CI. Mandelamide, 8CI

[4410-31-5]



(R)-form

C₈H₉NO₂ 151.165

(R)- and (S)-forms confused in early lit.

(R)-form [24008-62-6]

Plates (C₆H₆). Mp 123-124°. [α]_D²⁰ -73 (c, 1.60 in Me₂CO).

O-β-D-Glucopyranoside: [151649-54-6]

C₁₄H₁₉NO₇ 313.307

Constit. of *Olinia emarginata* and *Olinia ventosa*. [α]_D²⁰ -135 (c, 0.4 in MeOH).

O-(6-O-Benzoyl-β-D-glucopyranoside):

Persicaside

[1007096-61-8]

C₂₁H₂₃NO₈ 417.415

Constit. of the seeds of *Prunus persica* (peach). Antiinflammatory agent. Cryst. Config. not confirmed.

O-[β-D-Xylopyranosyl-(1→6)-β-D-glucopyranoside]: **Lucuminamide**

C₁₉H₂₇NO₁₁ 445.422

Constit. of *Calocarpum sapota* (preferred genus name *Pouteria*). Powder.

N-Me:

C₉H₁₁NO₂ 165.191

Mp 94-95°.

Me ether: 2-Methoxy-2-phenylacetamide

C₉H₁₁NO₂ 165.191

Plates (petrol). Mp 108-109°. [α]_D¹⁸ -103.6 (Me₂CO).

(S)-form [24008-63-7]

Mp 123°. [α]_D²⁰ +73 (c, 1.60 in Me₂CO).

N-Benzyl:

C₁₅H₁₅NO₂ 241.289

Catalyst for enantioselective addn. of dimethylzinc to *α*-keto esters. Solid (MeOH). Mp 134-135°. [α]_D²⁵ +83.2 (c, 0.54 in CHCl₃).

(±)-form

Plates (C₆H₆ or EtOH). Spar. sol. Et₂O. Mp 134-135°.

O-Ac: [6635-16-1]

C₁₀H₁₁NO₃ 193.202

Needles (H₂O). Mp 112-113°.

N,N-Di-Me: [2019-71-8]

C₁₀H₁₃NO₂ 179.218

Mp 158°.

Me ether: [56598-63-1]

Plates (H₂O or Et₂O). Sol. EtOH,

C₆H₆. Mp 112-114°.

Me ether, N,N-di-Me:

C₁₁H₁₅NO₂ 193.245

Mp 41°.

[65645-88-7]

Freudenberg, K. *et al.*, *Annalen*, 1933, **501**, 199

(*synth. abs config*)

Org. Synth., *Coll. Vol.*, 3, 1955, 536 (*synth*)

Iriuchijima, S. *et al.*, *J.A.C.S.*, 1975, **97**, 596

(*synth. ir*)

Schwanghart, A.-D. *et al.*, *Chem. Ber.*, 1977, **110**, 778 (*resoln*)

Nährstedt, A. *et al.*, *Phytochemistry*, 1993, **34**, 433-436 (*isol. glucoside*)

Takeda, T. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 697 (*Lucuminamide*)

Papakyprianou, A. *et al.*, *Org. Prep. Proced. Int.*, 2002, **34**, 436-440 (*synth*)

Tietze, L.F. *et al.*, *Synthesis*, 2004, 2236-2239 (*S-form, synth*)


Rho, J.-R. *et al.*, *Bull. Korean Chem. Soc.*, 2007, **28**, 1289-1293 (*Persicaside*)

Blay, G. *et al.*, *Synthesis*, 2007, 3754-3757 (*S-form, N-benzyl*)

4-Hydroxyphenylacetic acid H-672

4-Hydroxybenzeneacetic acid, 9CI. p-Hydroxy-*α*-toluic acid

[156-38-7]



C₈H₈O₃ 152.149

Constit. of sweet clover (*Melilotus officinalis*) and also from seeds of *Taraxacum officinale* yeast, wool and other biol. sources incl. an unidentified marine-derived fungus. Needles (H₂O). Sol.

EtOH, Et₂O, hot H₂O. Mp 148-150°. pK_a 4.59 (10% EtOH aq.).

▶ AI2680000

Amide: 4-Hydroxybenzeneacetamide,

9CI. p-Hydroxyphenylacetamide

[17194-82-0]

C₈H₉NO₂ 151.165

Isol. from the sponge *Suberites tylob-tusa*. Prod. by *Cytophaga marinoflava* sp. AM13. and an unidentified marine-derived fungus. Antitumour agent.

Leaflets (H₂O). Mp 175°.

▶ AC3851500

Amide, N-formyl: N-Formyl-4-hydroxy-

benzeneacetamide, 9CI. N-[2-(4-Hy-

droxyphenyl)acetyl]formamide

[913691-27-7]

C₉H₉NO₃ 179.175

Isol. from an unidentified marine-derived fungus.

Nitrile: 4-Hydroxyphenylacetoneitrile. 4-

Hydroxybenzyl cyanide. 4-(Cyanomethyl)phenol

[14191-95-8]

C₈H₇NO 133.149

Isol. from *Brassica alba* as a dec. prod. of 4-Hydroxybenzyl glucosinolate. Needles (H₂O). Mp 72°. Bp 330°.

▶ AM0530000

Nitrile, O-*α*-L-rhamnopyranoside: **Niazirin**

[122001-32-5]

C₁₄H₁₇NO₅ 279.292

Constit. of the leaves of the horseradish tree (*Moringa oleifera*, Morinaceae). Hypotensive agent.

▶ Mutagen.

Nitrile, O-(4-O-acetyl-*α*-L-rhamnopyra-

noside): **Niazirin**

[159397-73-6]

C₁₆H₁₉NO₆ 321.329

Constit. of horseradish tree (*Moringa oleifera*, Moringaceae) leaves. Hypotensive agent. Needles (CHCl₃/MeOH). Mp 183-184°. Struct. originally given as the 3-Ac isomer but revised without comment. 4-Ac is more likely (cf. other *M. oleifera* isolates). λ_{max} 200 ; 222 (MeOH).

Nitrile, O-(tri-O-acetyl-*α*-L-rhamnopyra-

noside): [219501-37-8]

C₂₀H₂₃NO₈ 405.404

Constit. of horseradish tree (*Moringa oleifera*, Moringaceae) leaves. Cryst. (EtOAc/petrol). Mp 103° (synthetic). λ_{max} 200 ; 220 ; 278 (MeOH).

Nitrile, O-[β-D-apiofuranosyl-(1→6)-β-

D-glucopyranoside]: [845531-45-5]

C₁₉H₂₅NO₁₀ 427.407

Constit. of the roots of *Semiaquilegia adoxoides*. [α]_D²⁰ -68.5 (c, 1 in MeOH).

Nitrile, O-[β-D-glucopyranosyl-(1→3)-*α*-

L-rhamnopyranoside]: [764656-60-2]

C₂₀H₂₇NO₁₀ 441.434

Constit. of the fruit of *Moringa oleifera* (horseradish tree). Amorph. powder.

O-(3-Methyl-2-butenyl), nitrile: 4-(3-

Methyl-2-butenyloxy)phenylacetoni-

trile. **Zanthonitrile**

[230308-50-6]

C₁₃H₁₅NO 201.268

Alkaloid from the fruit of *Zanthoxylum integrifolium*. Yellowish oil.

Li, S. *et al.*, *CA*, 1984, **100**, 117826y (*amide, nitrile*)

Faizi, S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1256-

1261 (*Niazirin, Niazirin*)

Strauss, C.R. *et al.*, *Org. Prep. Proced. Int.*,

1995, **27**, 552-555 (*amides*)

Kumar, A. *et al.*, *Synth. Commun.*, 1997, **27**,

1133-1141 (*synth, amide*)

Leuck, M. *et al.*, *Carbohydr. Res.*, 1998, **312**,

33-44 (*Niazirin derivs, synth, struct*)

Chen, I.-S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 833-

837 (*Zanthonitrile*)

Zhang, H. *et al.*, *Chin. J. Chem.*, 2004, **22**,

1200-1203 (*nitrile 6-apiosylglucoside*)

Shabaan, M. *et al.*, *Dissertation*, Univ. of

Göttingen, 2004, (*amide, isol, Cytophaga*)

Francis, J.A. *et al.*, *Helv. Chim. Acta*, 2004, **87**,

317-326 (*nitrile 3-glucosylrhamnoside, Me ester rhamnoside*)

2-Hydroxy-2-phenylacetone nitrile H-673

α -Hydroxybenzeneacetone nitrile, 9CI. Benzaldehyde cyanohydrin. Amygdonitrile. Mandelonitrile
[532-28-5]



C₈H₇NO 133.149

Glycosides widespread in (mostly higher) plants as cyanogenic glycosides. Glycosides of both enantiomers and of the racemate occur.

(R)-form [10020-96-9]

Needles. Mp 28.5-29.5°. [α]_D²⁵ +46.9 (C₆H₆).

▶ Eye irritant. LD₅₀ (rat, orl) 116 mg/kg. OO8400000

O-Ac: [119718-89-7]

C₁₀H₉NO₂ 175.187
[α]_D²⁰ +8.8 (c, 1 in CHCl₃).

O- β -D-Allopyranoside: **Passiedulin**

C₁₄H₁₇NO₆ 295.291

Constit. of the leaves and stems of *Passiflora edulis* (passion fruit). [α]_D²⁵ -69 (c, 0.24 in MeOH).

O- β -D-Glucopyranoside: **Prunasin**

[99-18-3]

C₁₄H₁₇NO₆ 295.291

Isol. from kernels of *Prunus* spp., immature fruits of *Passiflora* spp. and leaves of perilla (*Perilla frutescens* var. *acuta*). Needles (CHCl₃). Mp 147-148°. [α]_D -29.6 (H₂O).

▶ UL3420000

O-(6-O-Malonyl- β -D-glucopyranoside):**6'-O-Malonylprunasin**

[121748-23-0]

C₁₇H₁₉NO₉ 381.338

Constit. of *Lotononis fruticoides* and *Lotononis* aff. *falcata*.

O-[3,4,5-Trihydroxybenzoyl-(\rightarrow 6)- β -D-glucopyranoside]: **6'-O-Galloylprunasin**

C₂₁H₂₁NO₁₀ 447.398

Constit. of *Monochaetum multiflorum* and *Phyllagathis rotundifolia*. Pinkish amorph. powder. [α]_D²⁷ -81.2 (c, 0.22 in MeOH) (-17). λ _{max} 276 (log ϵ 3.98) (MeOH). λ _{max} 269 (log ϵ 2.8); 351 (log ϵ 2.4) (MeOH).

O-[2,6-Bis(3,4,5-trihydroxybenzoyl)- β -D-glucopyranoside]: **2',6'-Di-O-galloylprunasin**

C₂₈H₂₅NO₁₄ 599.504

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ -77.2 (c, 0.12 in MeOH). λ _{max} 278 (log ϵ 4.31) (MeOH).

O-[3,6-Bis(3,4,5-trihydroxybenzoyl)- β -D-glucopyranoside]: **3',6'-Di-O-galloylprunasin**

C₂₈H₂₅NO₁₄ 599.504

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ -8.2 (c, 0.27 in MeOH). λ _{max} 276 (log ϵ 4.28) (MeOH).

O-[4,6-Bis(3,4,5-trihydroxybenzoyl)- β -*D*-glucopyranoside]: **4',6'-Di-O-galloylprunasin**

C₂₈H₂₅NO₁₄ 599.504

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ -73.1 (c, 0.12 in MeOH). λ _{max} 277 (log ϵ 4.21) (MeOH).

O-[2,3,6-Tris(3,4,5-trihydroxybenzoyl)- β -D-glucopyranoside]: **2',3',6'-Tri-O-galloylprunasin**

C₃₅H₂₉NO₁₈ 751.61

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ +57 (c, 0.16 in MeOH). λ _{max} 278 (log ϵ 4.39) (MeOH).

O-[3,4,6-Tris(3,4,5-trihydroxybenzoyl)- β -D-glucopyranoside]: **3',4',6'-Tri-O-galloylprunasin**

C₃₅H₂₉NO₁₈ 751.61

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ -65 (c, 0.23 in MeOH). λ _{max} 278 (log ϵ 4.49) (MeOH).

O-[2,3,4,6-Tetrakis(3,4,5-trihydroxybenzoyl)- β -D-glucopyranoside]: **2',3',4',6'-Tetra-O-galloylprunasin**

C₄₂H₃₃NO₂₂ 903.716

Constit. of *Phyllagathis rotundifolia*. Tan amorph. powder. [α]_D²⁷ +14.2 (c, 0.25 in MeOH). λ _{max} 279 (log ϵ 4.58) (MeOH).

O-[4-Hydroxy-E-cinnamoyl-(\rightarrow 4)- β -D-glucopyranoside]: **4'-O-p-Coumaroylprunasin**

[190581-72-7]

C₂₃H₂₃NO₈ 441.437

Constit. of *Microlepis strigosa*.

O-[3,4-Dihydroxy-E-cinnamoyl-(\rightarrow 4)- β -D-glucopyranoside]: **4'-O-Caffeoylprunasin**

[190581-73-8]

C₂₃H₂₃NO₉ 457.436

Constit. of *Microlepis strigosa*.

O-[3,4-Dihydroxy-E-cinnamoyl-(\rightarrow 6)- β -D-glucopyranoside]: **Grayanin**

[110978-97-7]

C₂₃H₂₃NO₉ 457.436

Constit. of *Prunus grayana* bark. Pale yellow powder. [α]_D²⁵ -39.1 (c, 0.7 in Me₂CO).

O-[β -D-Apiofuranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Oxyanthin**

[140681-88-5]

C₁₉H₂₅NO₁₀ 427.407

Constit. of *Oxyanthus pyriformis* and *Psydrax livida*. Mp 158° (as per-Ac).

[α]_D²⁰ -81.1 (c, 0.95 in MeOH).

O-[5-O-Benzoyl- β -D-apiofuranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **5'-O-Benzoyloxyanthin**

[140701-86-6]

C₂₆H₂₉NO₁₁ 531.515

Constit. of *Oxyanthus pyriformis* and *Psydrax livida*. Mp 183° (as per-Ac).

[α]_D²⁸ -59.4 (c, 3.3 in MeOH).

O-[α -L-Arabinopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Vicianin**

[155-57-7]

C₁₉H₂₅NO₁₀ 427.407

Isol. from seeds of vetch (*Vicia angustifolia*). Also from *Gerbera jamesonii*

and ferns (*Davallia* spp.). Needles (H₂O). Mp 160°. [α]_D -20.7.

▶ Toxic.

O-[β -D-Xylopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Lucumin**. **Lucuminoside**

[1392-28-5]

C₁₉H₂₅NO₁₀ 427.407

Isol. from marmalade plum (*Lucuma mammosa*, preferred genus name *Pouteria*) seeds. Mp 184-185°. [α]_D²⁰ -50.5 (c, 0.4 in H₂O).

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Mandelonitrile rutinoid**

[184002-37-7]

C₂₀H₂₇NO₁₀ 441.434

Constit. of the fruit of purple passion fruit (*Passiflora edulis*).

O-[β -D-Glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]: **Mandelonitrile sophoroside**

[99520-78-2]

C₂₀H₂₇NO₁₁ 457.433

Isol. from leaves of perilla (*Perilla frutescens* var. *acuta*). Needles (H₂O). Mp 193-195°. [α]_D²² -22.6 (c, 0.14 in MeOH).

O-[β -D-Glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Mandelonitrile gentiobioside**. **Amygdalin**. **Amygdaloid**. **Glucoprunasin**. **Laetrile**. **Vitamin B₁₇**. **NSC 15780**

[29883-15-6]

[672-72-0]

C₂₀H₂₇NO₁₁ 457.433

Bitter glycoside of the Rosaceae, found esp. in kernels of cherries, peaches and apricots. Present in cold pressed bitter almond oil from the above sources prior to enzymic hydrolysis and steam distillation for food use.

Antiinflammatory agent. Trihydrate. Poorly sol. Et₂O. Mp 214°. [α]_D²⁵ -40.6 (H₂O). Log P -4.18 (uncertain value) (calc).

▶ LD₅₀ (rat, orl) 522 mg/kg. Exp. teratogen. OO8450000**(S)-form**O- β -D-Allopyranoside: Constit. of the leaves and stems of passion fruit (*Passiflora edulis*).O- β -D-Glucopyranoside: **Sambunigrin**

[99-19-4]

C₁₄H₁₇NO₆ 295.291

Isol. from leaves of elderberry (*Sambucus nigra*) and from other plants. Needles (EtOAc). Mp 151-152°. [α]_D -76.3 (EtOAc).

O-[β -D-Apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside]:

C₁₉H₂₅NO₁₀ 427.407

Constit. of *Sambucus nigra* (elderberry). [α]_D -73 (c, 0.7 in EtOH). λ _{max} 278 (EtOH).

O-[3,4,5-Trihydroxybenzoyl-(\rightarrow 6)- β -D-glucopyranoside]: **6'-O-Galoylsambunigrin**

C₂₁H₂₁NO₁₀ 447.398

Constit. of *Elaeocarpus sericopetalus*. Oil. [α]_D²⁵ -41 (c, 0.4 in MeOH). λ _{max} 216 (log ϵ 9.1); 278 (log ϵ 10.2) (MeOH).

(±)-form [613-88-7]

Isol. from peach kernels (*Prunus persica*).
Prisms or oil. Mp 21.5-22°.

O-β-D-Glucopyranoside: Prulaurasin

[138-53-4]

C₁₄H₁₇NO₆ 295.291

Isol. from cherry laurel leaves (*Prunus laurocerasus*), *Cotoneaster* spp. and other plants. Needles (EtOAc). Mp 122-123°. [α]_D -53 (H₂O).

▶ UL3200000

O-[β-D-Xylopyranosyl-(1→6)-β-D-glucopyranoside]: Epilucumin

[89460-01-5]

C₁₉H₂₅NO₁₀ 427.407Isol. from fruits of *Anthemis cairica*.**Ac:** [5762-35-6]C₁₀H₉NO₂ 175.187Bp₂₅ 152° Bp₁₁ 137-138°.**Benzoyl:** [4242-46-0]C₁₅H₁₁NO₂ 237.257

Found in defensive secretions of millipedes. Needles (EtOH). Mp 63-64°. Releases HCN and acetophenone.

Me ether: [13031-13-5]C₉H₉NO 147.176Bp₁₄ 116-118°.**Et ether:** [33224-69-0]C₁₀H₁₁NO 161.203Bp₁₆ 122-124°.

[51371-34-7, 60981-47-7, 61091-09-6, 61091-10-9, 29883-16-7]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 426B; 426D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1483B; 1484A (nmr)

Caldwell, R.J. et al., *J.C.S.*, 1907, **91**, 671

(*Sambunigrin*, *Prulaurasin*)

Power, F.B. et al., *J.C.S.*, 1909, **95**, 243; 1910,

97, 1099 (*Prunasin*)

Hawarth, W.N. et al., *J.C.S.*, 1923, 3120; 1924,

1337 (*Amygdalin*)

Chaudhury, D.N. et al., *J.C.S.*, 1949, 2054

(*Vicianin*)

Jones, M.B. et al., *Science (Washington, D.C.)*,

1961, **134**, 284 (isol. (±)-form)

Towers, G.H.N. et al., *Tetrahedron*, 1964, **20**,

71 (pmr, glucosides)

Kofod, H. et al., *Phytochemistry*, 1968, **8**, 1509

(*Vicianin*)

Schwartzmaier, U. et al., *Chem. Ber.*, 1976,

109, 3250 (pmr, *Amygdalin*)

Duffey, S.S. et al., *J. Chem. Ecol.*, 1977, **3**, 101

(*benzoyl*)

Turczan, J.W. et al., *J. Assoc. Off. Anal. Chem.*,

1978, **61**, 192; 1979, **62**, 190 (pmr, derivs)

Dreifuss, P.A. et al., *Biomed. Mass Spectrom.*,

1980, **7**, 201 (ms, *Amygdalin*)

Huebel, W. et al., *Arch. Pharm. (Weinheim, Ger.)*,

1981, **314**, 609 (cmr)

Spencer, K.C. et al., *J. Agric. Food Chem.*,

1983, **31**, 794-796 (*Prunasin*)

Nahrstedt, A. et al., *Planta Med.*, 1983, **49**,

143 (*Epilucumin*)

Aritomi, M. et al., *Phytochemistry*, 1985, **24**,

2438 (*sophoroside*)

Shimomura, H. et al., *Phytochemistry*, 1987,

26, 2363 (*Grayanin*)

Rockenbach, J. et al., *Phytochemistry*, 1992, **31**,

567 (*Oxyanthin*)

Cardona, L. et al., *Phytochemistry*, 1992, **31**,

3507 (*Prunasin*, cmr)

Martindale, *The Extra Pharmacopoeia*, 30th

edn., *Pharmaceutical Press*, 1993, 1428

Takeda, T. et al., *Chem. Pharm. Bull.*, 1997, **45**,

697 (*Lucumin*)

Wada, H. et al., *Nat. Med. (Tokyo)*, 1997, **51**,

69-70 (*Cinnamoylprunasin*)

Nakajima, N. et al., *Biosci., Biotechnol.,*

Biochem., 1998, **62**, 453-458 (*Prunasin,*

synth, pmr, cmr)

Chassagne, D. et al., *Phytochemistry*, 1998, **49**,

757-759 (*rutinoside*)

Lechtenberg, M. et al., *Biochem. Syst. Ecol.*,

1999, **27**, 607-612 (*6'-Malonylprunasin*)

Greca, M.D. et al., *Nat. Prod. Lett.*, 2000, **14**,

175-182 (*2-apisosylglucoside*)

Christensen, J. et al., *Org. Lett.*, 2001, **3**, 2193-

2195 (*Passiedulin*)

Isaza, J.H. et al., *Phytochemistry*, 2001, **58**,

321-327 (*6'-Galloylprunasin*)

Ling, S.-K. et al., *J. Nat. Prod.*, 2002, **65**, 131-

135 (*Phyllagathis galloylprunasin*)

Seigler, D.S. et al., *Phytochemistry*, 2002, **60**,

873-882 (*alloyprunasin*)

Belokon, Yu.N. et al., *Eur. J. Org. Chem.*,

2006, 4609-4617 (*R-form*, *O-Ac*)

Miller, R.E. et al., *Phytochemistry*, 2006, **67**,

1365-1371 (*6'-O-Galloylsambunigrin*)

Lewis, R.J. et al., *Sax's Dangerous Properties*

of Industrial Materials, 8th edn., *Van*


Nostrand Reinhold, 1992, AOD500;

MAP250; IHO700

1-(4-Hydroxyphenyl)-3H-benzo[a]pyrrolo[2,3-c]phenazine H-674

4-(3H-Benzo[a]pyrrolo[2,3-c]phenazin-1-yl)phenol, 9CI

[156753-18-3]

C₂₄H₁₅N₃O 361.402


Prod. by *Nostoc commune*. Antibacterial agent and spore formn. inhibitor. Cell division inhibitor. Phytotoxin.

Japan. Pat., 1994, 94 116 159; *CA*, **121**,

151281k

7-Hydroxy-2-phenyl-4H-3,1-benzoxazin-4-one H-675**Dianthalexine**

[85915-62-4]

C₁₄H₉NO₃ 239.23

Alkaloid from *Dianthus caryophyllus* (Caryophyllaceae). Cryst. Sol. MeOH, Me₂CO; poorly sol. hexane. Mp 229-231°. λ_{max} 261 (ε 44900); 292 (ε 14200); 302 (ε 13500) (MeOH) (Berdy).

Bouillant, M.-L. et al., *Tet. Lett.*, 1983, **24**, 51

(uv, pmr, ms, struct)

Hauteville, M. et al., *J. Het. Chem.*, 1988, **25**,


715 (*synth*)

Crombie, L. et al., *Tet. Lett.*, 1990, **31**, 2647

(*synth*)

2-(4-Hydroxyphenyl)ethanol H-676

4-Hydroxybenzeneethanol, 9CI. p-Hydroxyphenethyl alcohol. **Tyrosol**. *Salidrosl* [501-94-0]

C₈H₁₀O₂ 138.166

Prod. by several *Ceratocystis* spp. Isol. from leaves of *Ligustrum ovalifolium* and bark of *Fraxinus excelsior*. Prod. by *Gibberella fujikuroi* and by the marine strain *Bacillus subtilis* KMM3427 associated with sea sponge *Verongia* sp. Also isol. from peanuts. Needles (CHCl₃). Sol. H₂O, EtOH, Et₂O, Me₂CO; spar. sol. petrol. Mp 85-86°.

1-O-Carbamoyl: 2-(4-Hydroxyphenyl)ethyl carbamate. *Tyrosol carbamate* [1001207-09-5]


C₉H₁₁NO₃ 181.191

Prod. by a marine-derived *Arthrinium* sp.

Gautschi, J.T. et al., *Nat. Prod. Commun.*, 2007, **2**, 541-546 (*carbamate*)

2-(4-Hydroxyphenyl)ethenamine H-677

4-Hydroxystyrylamine. 4-(2-Aminoethenyl)phenol

C₈H₉NO 135.165

N-Ac: [214217-93-3]

C₁₀H₁₁NO₂ 177.202

Prod. by *Streptomyces* sp. T-950510.


Antitumour agent.

Japan. Pat., 1998, 98 259 174; *CA*, **129**,

289264d (*N-Ac*)

2-(3-Hydroxyphenyl)ethylamine H-678

3-(2-Aminoethyl)phenol, 9CI. 3-Hydroxyphenethylamine. m-Tyramine [588-05-6]

C₈H₁₁NO 137.181

Found in human brain. Tan needles (C₆H₆). Mp 103-104° (72-75°).

Hydrochloride: [3458-98-8]

Cryst. (EtOH/Et₂O). Mp 145°.

▶ SJ5975000


[38449-59-1]

Corrodi, H. et al., *Helv. Chim. Acta*, 1966, **49**, 798 (*synth*)

Anderson, W.K. *et al.*, *J. Med. Chem.*, 1984, 27, 1321 (*synth*)

N-[2-(4-Hydroxyphenyl)ethyl]-3-methyl-2-dodecenamide, 9CI **H-679**

3-Methyl-2-dodecenoic acid p-hydroxyphenethylamide



C₂₁H₃₃NO₂ 331.497

(Z)-form [75513-94-9]

Isol. from *Muricea austera* and *Simularia flexibilis*. Shows antiinflammatory, cytotoxic and cardiotoxic activities. Sol.

MeOH, CH₂Cl₂; poorly sol. H₂O. Mp 70-72°. λ_{max} 278 (ε 2860) (MeOH) (Derep).

2'',3''-ξ-Dihydro: **N-[2-(4-Hydroxyphenyl)ethyl]-3-methyl-2-dodecenamide**

C₂₁H₃₅NO₂ 333.513

Isol. from *Muricea austera*. Oil. [α]_D²⁵ -26.8 (c, 0.5 in CHCl₃).

3-Hydroxy, Me ether: **N-[2-(3-Hydroxy-4-methoxyphenyl)ethyl]-3-methyl-2-dodecenamide, 9CI**

[75513-96-1]

C₂₂H₃₅NO₃ 361.523

Isol. from *Simularia flexibilis*. Shows antiinflammatory and cardiotoxic props. Oil which slowly solidifies. Sol. MeOH, CH₂Cl₂; poorly sol. H₂O. λ_{max} 280 (ε 3400); 287 (sh) (MeOH) (Derep).

3-Methoxy: **N-[2-(4-Hydroxy-3-methoxyphenyl)ethyl]-3-methyl-2-dodecenamide, 9CI**

[75513-95-0]

C₂₂H₃₅NO₃ 361.523

Isol. from *Simularia flexibilis*. Shows antiinflammatory and cardiotoxic props. Oil which slowly solidifies. Sol. MeOH, CH₂Cl₂; poorly sol. H₂O. λ_{max} 280 (ε 3400); 287 (sh) (MeOH) (Derep).

[75513-93-8 E-form]

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1980, 33,


1799 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Sheu, J.-H. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1999, 46, 253-257 (*isol, activity*)

Gutiérrez, M. *et al.*, *J. Nat. Prod.*, 2006, 69, 1379-1383 (*Muricea austera constiti*)

1-(4-Hydroxyphenyl)-2-isocynoethene **H-680**

4-(2-Isocyanoethenyl)phenol



C₉H₇NO 145.16

(E)-form [298711-81-6]

Prod. by *Erwinia carotovora*.


O-(6-Deoxyhexopyranoside): [957062-49-6]

C₁₅H₁₇NO₅ 291.303

Prod. by *Erwinia carotovora*.

Brady, S.F. *et al.*, *J.A.C.S.*, 2007, 129, 12102-12103 (*isol, pmr, cmr, ms*)

2'-O-[3-(4-Hydroxyphenyl)-2-methoxypropenoyl]-3'-O-[3-(1H-imidazol-4-yl)propenoyl]-S-methyl-5'-thioadenosine **H-681**



C₂₇H₂₇N₇O₇S 593.619

Isol. from the ascidian *Atrialium robustum*. Amorph. solid. [α]_D²⁴ -115 (c, 0.18 in MeOH). λ_{max} 293 (ε 44800) (MeOH).

S-Oxide:

C₂₇H₂₇N₇O₈S 609.618


Isol. from *Atrialium robustum*.

Amorph. solid. [α]_D²⁴ -124.1 (c, 0.08 in MeOH). λ_{max} 288 (ε 44000) (MeOH).

Kehraus, S. *et al.*, *J. Med. Chem.*, 2004, 47, 2243-2255 (*isol, pmr, cmr, ms*)

3-(4-Hydroxyphenyl)-N-[4-(3-methyl-2,5-dioxo-1-pyrrolidinyl)-butyl]-2-propenamide, 9CI **H-682**

N-(4-Methylsuccinimidobutyl)-p-coumaramide



C₁₈H₂₂N₂O₄ 330.383

(±)-(E)-form [128866-07-9]


Alkaloid from bulbs of *Lilium regale* (Liliaceae). Needles (MeOH). Mp 184-185.5° (natural) Mp 192-193° (synthetic).

Mimaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1990, 38, 541 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kuehne, P. *et al.*, *Tetrahedron*, 1993, 49, 4575 (*synth*)

Linden, A. *et al.*, *Acta Cryst. C*, 1997, 53, 1472-1475 (*cryst struct*)

5-[(4-Hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone **H-683**



C₁₀H₈N₂O₂S 220.251


(Z)-form [280137-26-0]

Prod. by *Burkholderia mallei*. Cryst.

Brady, S.F. *et al.*, *J.A.C.S.*, 2007, 129, 12102-12103 (*isol, pmr, cryst struct*)

3-(4-Hydroxyphenyl)-4-(2-methylpropyl)-1H-pyrrole-2,5-dione 2-(4-Hydroxyphenyl)-3-isobutylmaleimide **H-684**

[1040394-24-8]



C₁₄H₁₅NO₃ 245.277

Prod. by *Antrodia camphorata*. Yellow needles. Mp 199-201°. λ_{max} 231 (log ε 4.4); 269 (log ε 4.4); 360 (log ε 3.8) (MeOH).

O-(3-Methyl-2-butenyl): 3-[4-(3-Methyl-2-butenyloxy)phenyl]-4-(2-methylpropyl)-1H-pyrrole-2,5-dione. 2-Isobutyl-3-(4-prenyloxyphenyl)maleimide. **Antrodin B**. *Camphorataimide B*. *Antro-pyrroledione*

[656830-25-0]

C₁₉H₂₃NO₃ 313.396

Prod. by *Antrodia camphorata* and *Antrodia cinnamomea*. Cytotoxic. Yellow needles (hexane/EtOAc). Mp 110-111°. λ_{max} 230 (log ε 4.3); 272 (log ε 3.5); 355 (log ε 3.7) (MeOH).

O-(3-Methyl-2-butenyl), N-hydroxy: **Antrodin C**. *Camphorataimide C*

[656830-26-1]

C₁₉H₂₃NO₄ 329.395

Prod. by *Antrodia camphorata* and *Antrodia cinnamomea*. Cytotoxic. Yellow oil. λ_{max} 232 (log ε 4.3); 296 (log ε 3.7); 374 (log ε 3.7) (MeOH).

2-Deoxo, 2ξ, N-dihydroxy, O-(3-methyl-2-butenyl):

C₁₉H₂₅NO₄ 331.411

Isol. from *Sparassis crispa* (cauliflower mushroom). Inhibitor of melanin synth. Oil.

3R*,4R*-Dihydro, O-(3-methyl-2-butenyl), N-hydroxy: **Antrodin E**. *Camphorataimide D*

[656832-05-2]

C₁₉H₂₅NO₄ 331.411

Prod. by *Antrodia camphorata* and *Antrodia cinnamomea*. Oil. [α]_D²³ +3 (c, 0.2 in MeOH). λ_{max} 227 (log ε 4.3); 275 (log ε 3.4); 283 (log ε 3.3) (MeOH).

3R,4S-Dihydro, N-hydroxy: [1040394-21-5]

C₁₄H₁₇NO₄ 263.293

Prod. by *Antrodia camphorata*. Mp 189-193°. [α]_D²⁵ +9.6 (c, 0.11 in MeOH). λ_{max} 221 (log ε 4.16); 275 (log ε 3.46); 283 (log ε 3.38) (MeOH).

3R,4S-Dihydro, O-(3-methyl-2-butenyl): [1030374-53-8]

C₁₉H₂₅NO₃ 315.411


Prod. by *Antrodia camphorata*. Gum. [α]_D²⁵ +24 (c, 0.2 in MeOH).

3R*,4S*-Dihydro, O-(3-methyl-2-butenyl), N-hydroxy: **Antrodin D**. *Camphorataimide E*

[656832-04-1]
C₁₉H₂₅NO₄ 331.411
Prod. by *Antrodia camphorata*,
Antrodia cinnamomea and *Sparassiss
crispa* (cauliflower mushroom). Oil.
[α]_D²⁵ +2.5 (c, 0.2 in MeOH). λ_{max} 225
(log ε 4.3); 275 (log ε 3.3); 283 (log ε
3.2) (MeOH).

Nakamura, N. *et al.*, *J. Nat. Prod.*, 2004, **67**,
46-48 (*O*-prenyl, *isol*, *pmr*, *cmr*, *cryst struct*)
Baag, M.M. *et al.*, *Synthesis*, 2006, 1005-1008
(*synth*, *ir*, *pmr*, *cmr*)
Kawagishi, H. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 2007, **71**, 1804-1806 (*Sparassiss
crispa constit*)
Stewart, S.G. *et al.*, *Tet. Lett.*, 2007, **48**, 2241-
2244 (*synth*)
Chien, S.-C. *et al.*, *J. Agric. Food Chem.*, 2008,
56, 7017-7022 (*Antrodia camphorata constits*)
Cheng, C.-F. *et al.*, *Tetrahedron*, 2008, **64**,
4347-4353 (*Antrodins D,E, synth*)

2-(4-Hydroxyphenyl)-1-methyl-4(1H)-quinolinone, 9CI H-685
Reevesianine A
[109030-96-8]



C₁₆H₁₃NO₂ 251.284
Alkaloid from the root and stem bark of
Skimmia reevesiana (Rutaceae). Plates
(MeOH). Mp 322-326°.

Ac: [109030-97-9]
Plates (Me₂CO). Mp 124-126°.

6-Methoxy: Reevesianine B

[109030-98-0]
C₁₇H₁₅NO₃ 281.31
Alkaloid from the root and stem bark of
Skimmia reevesiana (Rutaceae).
Plates (MeOH). Mp 304-306°.


6-Methoxy, Ac: [109030-99-1]
Mp 200-202°.

Wu, T.S. *et al.*, *Phytochemistry*, 1987, **26**, 873-
875 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *deriv*)

Kuo, S.C. *et al.*, *J. Med. Chem.*, 1993, **36**,
1146-1156 (*synth*, *ir*, *pmr*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1999,
47, 1038-1039 (*synth*, *ir*, *pmr*)

1-Hydroxy-4-phenyl-2,7-naphthyridine H-686
4-Phenyl-2,7-naphthyridin-1(2H)-one,
9CI. Lophocladine A




C₁₄H₁₀N₂O 222.246
Alkaloid from *Lophocladia* sp. Displays
affinity for NMDA receptors. δ-Opioid
receptor antagonist. Needles (DMSO).

Mp 211-213°. λ_{max} 224 (log ε 4.06);
250 (log ε 4.03); 314 (log ε 3.88)
(MeOH).

Gross, H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 640-
644 (*isol*, *pmr*, *cmr*, *ms*)

2-(4-Hydroxyphenyl)-2-oxoa-cetic acid H-687

4-Hydroxy-α-oxobenzeneacetic acid, 9CI.
p-Hydroxyphenylglyoxylic acid, 8CI. 4-
Hydroxybenzoylformic acid. Pisolithin A
[15573-67-8]



C₈H₆O₄ 166.133
Metab. of the fungus *Pisolithus tinctorius*.
Needles (Et₂O/C₆H₆/petrol). Mp 177.5-
178°. λ_{max} 206 ; 226 ; 296 (HCl) (Berdy).
λ_{max} 240 ; 334 (NaOH) (Berdy).

*Me ether, methylamide: 2-(4-Methoxy-
phenyl)-N-methyl-2-oxoacetamide*
[244264-57-1]
C₁₀H₁₁NO₃ 193.202
Isol. from the ascidian *Polycarpa
aurata*. Oil. λ_{max} 225 (ε 8738); 293 (ε
1675) (EtOH).

[54537-30-3]

Wessels, M. *et al.*, *J. Nat. Prod.*, 2001, **64**,
1556-1558 (*Me ether derivs*)

Hydroxyphenylpropanedioic acid H-688

*Hydroxyphenylmalonic acid. Phenyltar-
tronic acid*

PhC(OH)(COOH)₂

C₉H₈O₅ 196.159

Parent acid not well known.

Di-Me ester: [92607-06-2]

C₁₁H₁₂O₅ 224.213

Needles. Mp 67°. Bp₁₁ 165°.

Di-Et ester:

C₁₃H₁₆O₅ 252.266

Cryst. mass. Mp 28°. Bp₁₀ 170°.

Me ether, di-Me ester: [124604-00-8]

C₁₂H₁₄O₅ 238.24

Oil.

*Diamide: 2-Hydroxy-2-phenylpropanedia-
mide. Armillarisin B*

[53696-75-6]

C₉H₁₀N₂O₃ 194.19

Isol. from *Armillariella tabescens*.

Shows insecticidal activity. Solid

(EtOH). Mp 159-161°.

Guyot, A. *et al.*, *C. R. Hebd. Seances Acad.*

Sci., 1909, **148**, 564 (*synth*)

Weishengwu Xuebao, 1974, **14**, 9; *CA*, **82**,
40463c (*isol*, *diamide*)


Ger. Pat., 1983, 3 140 275; *CA*, **99**, 22142u
(*synth*)

Toth, G. *et al.*, *CA*, 1988, **109**, 62247u (*props*,
diamide)

Kawabata, J. *et al.*, *Annalen*, 1990, 181 (*synth*,
pmr)

3-(4-Hydroxyphenyl)propa- noic acid H-689

4-Hydroxybenzenepropanoic acid, 9CI. 3-
(p-Hydroxyphenyl)propionic acid, 8CI.
Dihydro-p-coumaric acid. p-Hydroxyhy-
drocinnamic acid. Phloretic acid. Phlor-
etic acid. HPPA
[501-97-3]



C₉H₁₀O₃ 166.176

Constit. of urine. Isol. from sulfite
liquors, prod. by *Clostridium butyricum*.
Isol. from heartwood of *Haplormisia
monophylla* (Fabaceae). A prod. of
tyrosine metab.; conc. in urine increases
in patients with gastrointestinal diseases.
Fluorescent substrate for the detn. of
peroxidase. Prisms (Et₂O). Mp 129-130°.
pK_a 4.76 (25°).

▶MW5342000

Methylamide: 3-(4-Hydroxyphenyl)-N-
methylpropanamide

C₁₀H₁₃NO₂ 179.218

Prod. by *Micromonospora* sp. P1068.

Pale yellow solid. Mp 86°.


Gutierrez-Lugo, M.-T. *et al.*, *Nat. Prod. Res.*,
2005, **19**, 645-652 (*methylamide*)

3-(3-Hydroxyphenyl)-2-pro- penoic acid, 9CI H-690

*m-Hydroxycinnamic acid. m-Coumaric
acid. m-Cumaric acid*

[588-30-7]

[25429-38-3]



C₉H₈O₃ 164.16

(E)-form [14755-02-3]

Present in roots of *Citrus sinensis* (or-
ange), *Citrus limon* (lemon) and *Citrus
paradisi* (grapefruit). Prisms (H₂O). Sol.
EtOH, Et₂O, C₆H₆, hot H₂O. Mp 191°
(194°). pK_a 4.44.

Me ester: [66417-46-7]

C₁₀H₁₀O₃ 178.187

Leaflets. Mp 85°.

*Me ether, pyrrolidide: 1-(m-Methoxycin-
namoyl)pyrrolidine, 8CI. m-Methoxy-*
cinnamic acid pyrrolidide

[29647-01-6]

C₁₄H₁₇NO₂ 231.294

Alkaloid from the roots of *Piper
methisticum* (kava) (Piperaceae). Need-
les (petrol). Mp 90-93°.

[6099-04-3]

Achenbach, H. *et al.*, *Chem. Ber.*, 1970, **103**,
2535 (*Me ether pyrrolidide*)

Chen, I.-S. *et al.*, *Fitoterapia*, 2007, **78**, 414-419 (*Me ether pyrrolidide*)

3-(4-Hydroxyphenyl)-2-propenoic acid, 9CI H-691

p-Hydroxycinnamic acid, 8CI. *p*-Hydroxyphenylacrylic acid. ***p*-Coumaric acid**. *Naringenic acid*. *Naringeninic acid*. *p*-*Coumaric acid*
[7400-08-0]

C₉H₈O₃ 164.16

▶ LD₅₀ (mus, ipr) 657 mg/kg. Exp. reprod. effects. GD9094000

(*E*)-form [501-98-4]

Widespread in plants, e.g. peel of black cherry (*Prunus serotina*), lentil seeds and from red clover (*Trifolium pratense*) and *Daviesia latifolia*. Occurs as many glycosides. Found by Bate-Smith in 48% of investigated dicotyledonous and 55% of monocotyledonous spp. Shows cytostatic activity. Immunoactive agent, inhibitor of stilbene oxidase. Cryst. + 1H₂O (cold H₂O), anhyd. cryst. (hot H₂O). Sol. Et₂O, hot EtOH; spar. sol. C₆H₆; insol. petrol. Mp 210-213°. p*K*_{a1} 4.64; p*K*_{a2} 9.45 (25°). SOCl₂-activated polycondensation gives the homopolyester with an all-*trans* struct. Polyesters with other hydroxy acids are also known. λ_{max} 225; 290; 310 (MeOH) (Berdy). λ_{max} 223 (ε 14450); 286 (ε 19000) (EtOH) (Berdy).

▶ GD9095000

Me ether, pyrrolidide: *N*-(4-Methoxycinnamoyl)pyrrolidine. *p*-Methoxycinnamic acid pyrrolidide. **Piperlotine A** [389572-70-7]
C₁₄H₁₇NO₂ 231.294
Alkaloid from the leaves of *Piper lolot*. Syrup. λ_{max} 225; 300 (MeOH).

(*Z*)-form [4501-31-9]

Found in *Larix sibirica* bark and *Miscanthus floridulus*. Cryst. (C₆H₆ or H₂O). Mp 134-134.5° dec. (subl. from 124°).

▶ GD9095100

Me ether, pyrrolidide: **Piperlotine B** [958296-12-3]
C₁₄H₁₇NO₂ 231.294
Alkaloid from the leaves of *Piper lolot*. Syrup. λ_{max} 215; 273 (MeOH).


[155339-67-6, 155339-66-5, 830-09-1, 1929-30-2, 2979-06-8, 3943-97-3, 7139-64-2, 13080-39-2]

Wang, M.F. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 35-36 (*Tataramide A*)

Li, C.-Y. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 9436-9442 (*Piperlotines A, B*)

3-(4-Hydroxyphenyl)quinoline H-692

4-(3-Quinoliny)phenol, 9CI
[107915-36-6]



C₁₅H₁₁NO 221.258

Alkaloid from *Peganum nigellastrum*. Yellow needles (C₆H₆). Mp 224.5-226° (221-222°).

Me ether: 3-(4-Methoxyphenyl)quinoline [57479-32-0]

C₁₆H₁₃NO 235.285

Cryst. Mp 85-86°.

Kaslow, C.E. *et al.*, *J.O.C.*, 1958, **23**, 271-276 (*synth*)

Ishikura, M. *et al.*, *Heterocycles*, 1985, **23**, 2375-2386 (*Me ether*)


Cacchi, S. *et al.*, *Tetrahedron*, 1996, **52**, 10225-10240 (*synth, ir, pmr, cmr*)

Giroux, A. *et al.*, *Tet. Lett.*, 1997, **38**, 3841-3844 (*Me ether*)

Ma, Z.-Z. *et al.*, *Phytochemistry*, 2000, **53**, 1075-1078 (*isol*)

4-Hydroxy-2-phenylquinoline H-693

2-Phenyl-4(1*H*)-quinolinone, 9CI. 2-Phenyl-4-quinolinol
[14802-18-7]



C₁₅H₁₁NO 221.258

NH-form predominates. Plates (EtOH). Mp 255-257°.

NH-form

N-Me: 1-Methyl-2-phenyl-4(1*H*)-quinolinone

[17182-60-4]

C₁₆H₁₃NO 235.285

Alkaloid from *Balfourodendron riedelianum* (Rutaceae). Needles (MeOH). Mp 143-144°.

OH-form [1144-20-3]

Benzoyl: [880250-95-3]

C₂₂H₁₅NO₂ 325.366

Needles (petrol) or rhombs (dimorph.). Mp 96° (needles) Mp 110-111° (rhombs). Dimorphic, melts and resolidifies to give higher melting form.

Methanesulfonyl: [147197-91-9]

C₁₆H₁₃NO₃S 299.35

Cryst. (C₆H₆/petrol). Mp 132-134°.

4-Methylbenzenesulfonyl: [147197-92-0]

C₂₂H₁₇NO₃S 375.447

Cryst. (C₆H₆/petrol). Mp 126-128°.

Me ether: 4-Methoxy-2-phenylquinoline

[22680-62-2]

C₁₆H₁₃NO 235.285

Alkaloid from the leaves of *Lumasia amara* (Rutaceae). Needles (pentane). Mp 69-70° (66-67°).

Me ether, perchlorate:

Rods (EtOH). Mp 215-218°.

Me ether, methiodide:

Bright yellow needles (MeOH). Mp 148-151°.

Et ether: [22680-63-3]

C₁₇H₁₅NO 249.312

Mp 74-75°.

Goodwin, S. *et al.*, *J.A.C.S.*, 1957, **79**, 2239-2241 (*Me ether, isol, uv*)

Price, J.R. *et al.*, *Aust. J. Chem.*, 1959, **12**, 589-600 (*ir*)

Rapoport, H. *et al.*, *J.A.C.S.*, 1960, **82**, 4395-4404 (*N-Me, isol*)

Ogata, Y. *et al.*, *Tetrahedron*, 1971, **27**, 2765-2770 (*synth, uv*)

Koyama, J. *et al.*, *Chem. Express*, 1992, **7**, 321-324 (*synth, N-Me*)

Kuo, S.-C. *et al.*, *J. Med. Chem.*, 1993, **36**, 1146-1156 (*synth, Me ether, ir, pmr*)

Singh, O.V. *et al.*, *Synth. Commun.*, 1993, **23**, 277-283 (*synth*)

Jensen, S. *et al.*, *Acta Chem. Scand.*, 1994, **49**, 53-56 (*synth*)

Fürstner, A. *et al.*, *J.O.C.*, 1994, **59**, 5215-5229 (*synth, pmr, cmr, ms*)

Prakash, O. *et al.*, *Synth. Commun.*, 1994, **24**, 2169-2172 (*synth*)

Varma, R.S. *et al.*, *Tet. Lett.*, 1998, **39**, 9113-9116 (*synth*)

Koyama, T. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1038-1039 (*N-Me, synth, ir, pmr*)


Sato, S. *et al.*, *J. Het. Chem.*, 1999, **36**, 1189-1193 (*synth*)

George, L. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 558-564 (*O-benzoyl*)

Hemanth, K. *et al.*, *Tetrahedron*, 2007, **63**, 9531-9535 (*Me ether, Et ether*)

2-(2-Hydroxyphenyl)thiazole H-694

2-(2-Thiazolyl)phenol, 9CI
[83053-38-7]




C₉H₇NOS 177.226

Isol. from *Pseudomonas cepacia*. Mp 51°.

Bukovits, G.J. *et al.*, *Z. Naturforsch., B*, 1982, **37**, 877 (*isol*)

2-(2-Hydroxyphenyl)-4-thiazolecarboxylic acid, 9CI H-695

Aeruginic acid. *Eruginic acid*
[27501-91-3]



C₁₀H₇NO₃S 221.236

Isol. from cultures of *Pseudomonas aeruginosa* and *Pseudomonas cepacia*. Antibiotic showing antimicrobial, antiinflammatory and hypotensive activity. Sol. CHCl₃, bases, EtOH, Me₂CO, MeOH; fairly sol. EtOAc, Et₂O; poorly sol. H₂O, hexane. Mp 271-272°. λ_{max} 276 (ε 3000); 320 (ε 2250) (0.1*N* HCl) (Derep). λ_{max} 240 (sh) (ε 3750); 282 (ε 1400); 362 (ε 1800) (0.1*N* NaOH) (Derep). λ_{max} 280 (ε 2400); 324 (ε 1850) (EtOH) (Derep). λ_{max} 220; 279; 322 (ε 10500) (MeOH) (Berdy). λ_{max} 212; 282; 362 (EtOH/NaOH) (Berdy).

Me ester: [27501-92-4]

C₁₁H₉NO₃S 235.263

Isol. from cultures of *Pseudomonas aeruginosa*. Mp 138°.

Ac, Me ester: [27501-93-5]
Mp 99°.

Aldehyde: 2-(2-Hydroxyphenyl)-4-thiazolecarboxaldehyde, 9CI

[83053-39-8]
C₁₀H₇NO₂S 205.237
From *Pseudomonas cepacia*. Mp 126-127°.

Alcohol: 2-(2-Hydroxyphenyl)-4-(hydroxymethyl)thiazole, Aeruginol

C₁₀H₉NO₂S 207.253
Metab. from *Pseudomonas aeruginosa*. Off white amorph. solid. λ_{max} 324 (MeOH) (Berdy). λ_{max} 365 (MeOH/NaOH) (Berdy).

4S,5-Dihydro: Dihydroaeruginic acid

C₁₀H₉NO₃S 223.252
Antibiotic from *Pseudomonas fluorescens* strain PFM2. Exhibits antimicrobial activity. [α]_D²⁷ +47 (c, 0.10 in CHCl₃). λ_{max} 214 (ε 31750); 252 (ε 15800); 315 (ε 6670) (MeCN) (Berdy).

4S,5-Dihydro, Me ester: [α]_D²⁵ +11.3 (c, 0.15 in MeOH).

4ξ,5-Dihydro, alcohol: 4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolemethanol. 4-Hydroxymethyl-2-(2-hydroxyphenyl)-2-thiazoline, Aerugine

[112515-23-8]
C₁₀H₁₁NO₂S 209.268
Prod. by *Pseudomonas aeruginosa* 590, *Pseudomonas fluorescens* MM-B16 and *Streptomyces fradiae* SU-1. Cryst. (hexane). Mp 85-88°. [α]_D +28 (c, 1.0 in CHCl₃). λ_{max} 213 (ε 12300); 251 (ε 5500); 317 (ε 2090) (MeOH) (Berdy). λ_{max} 213 (ε 6760); 276 (ε 5250); 298 (ε 1950) (MeOH/HCl) (Berdy). λ_{max} 353 (ε 4466) (MeOH/NaOH) (Berdy).

Ger. Pat., 1969, 2 045 819; *CA*, 75, 18612j
Yamada, Y. *et al.*, *Agric. Biol. Chem.*, 1970, 34, 780 (isol, uv)

Bukovits, G.J. *et al.*, *Z. Naturforsch., B*, 1982, 37, 877 (isol)

Zunnundzhanov, A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1987, 23, 461-465 (*Aerugine, isol*)

Yang, W. *et al.*, *J. Nat. Prod.*, 1993, 56, 1993 (*Aeruginol, isol, uv, pmr, struct*)


Carmi, R. *et al.*, *J. Nat. Prod.*, 1994, 57, 1200 (*Dihydroaeruginic acid*)

Loughlin, W.A. *et al.*, *Aust. J. Chem.*, 2000, 53, 457-462 (*Dihydroaeruginic acid, synth*)

Lee, J.Y. *et al.*, *Appl. Environ. Microbiol.*, 2003, 69, 2023-2031 (*Aerugine, isol, pmr, cmr, ms, activity*)

3-Hydroxy-2-piperidinecarboxylic acid H-696

3-Hydroxypiperidic acid, 8CI. 2-Carboxy-3-hydroxypiperidine
[56879-47-1]



(2R,3R)-form

C₆H₁₁NO₃ 145.158

(2R,3R)-form
(-)-trans-form

[176019-04-8]
Mp 232° dec. [α]_D²² -14 (c, 0.4 in 10% HCl aq.).

(2R,3S)-form

(+)-cis-form
[119593-44-1]
No phys. props. reported.

(2S,3R)-form

(-)-cis-form
[112241-70-0]
Cryst. (MeOH). Mp 300°. [α]_D²⁰ -93.3 (c, 0.3 in H₂O).

(2S,3S)-form

(+)-trans-form
[176019-05-9]
Mp 232-236°. [α]_D²⁰ +13.5 (c, 0.2 in 10% HCl aq.).

Amide: [908040-24-4]

C₆H₁₂N₂O₂ 144.173
Cryst. (MeOH/Et₂O). Mp 149-150°. [α]_D²⁰ +43.8 (c, 0.4 in 10% HCl).

(2R,S,3SR)-form

(±)-cis-form
[122088-79-3]
Isol. from halophyte leaves, flowers and fruits. Cryst. (EtOH aq.). Mp 284-285° dec.

Me ester

C₇H₁₃NO₃ 159.185
Mp 191-192° dec. (hydrochloride).

[122019-48-1]

Fowden, L. *et al.*, *Biochem. J.*, 1958, 70, 629 (*isol, struct, synth*)

Pleiningner, H. *et al.*, *Chem. Ber.*, 1959, 92, 1579 (*synth*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1987, 35, 3482 (*synth, pmr*)

Drummond, J. *et al.*, *J. Med. Chem.*, 1989, 32, 2116 (*synth*)

Roemmele, R.C. *et al.*, *J.O.C.*, 1989, 54, 1866 (*synth*)

Battistini, L. *et al.*, *Tetrahedron: Asymmetry*, 1997, 8, 2975-2987 (*synth*)

Haddad, M. *et al.*, *Tet. Lett.*, 2001, 42, 5223-5225 (*R,R-form, synth*)

Bodas, M.S. *et al.*, *Tet. Lett.*, 2004, 45, 8461-8463 (*S,S-form, synth*)

Kumar, P. *et al.*, *J.O.C.*, 2005, 70, 360-363 (*R,R, S,S-forms*)

Liang, N. *et al.*, *J.O.C.*, 2005, 70, 10182-10185 (*synth*)

Kim, I.S. *et al.*, *Tet. Lett.*, 2006, 47, 7289-7293 (*S,S-form, synth*)

Feng, C.-G. *et al.*, *Tetrahedron*, 2006, 62, 7459-7465 (*2S,3S-form, amide*)


Phansavath, P. *et al.*, *J. Chem. Res.*, 2007, 313-316 (*synth*)

Pham, V.-T. *et al.*, *Tetrahedron: Asymmetry*, 2008, 19, 318-321 (*2S,3R-form, synth*)

Liu, L.-X. *et al.*, *Tetrahedron: Asymmetry*, 2008, 19, 1200-1203 (*2S,3S-form, synth*)

4-Hydroxy-2-piperidinecarboxylic acid, 9CI H-697

4-Hydroxypiperidic acid, Pegaline
[14228-16-1]



(2R,4R)-form

C₆H₁₁NO₃ 145.158

(2R,4R)-form

d-trans-form
[189385-64-6]
Gum. [α]_D²⁰ +12.6 (c, 0.8 in H₂O).

O-(4-Bromo-1H-pyrrole-2-carbonyl):

Damipipicoline

[949091-36-5]
C₁₁H₁₃BrN₂O₄ 317.139
Isol. from *Axinella damicornis*. Serotonin receptor modulator. [α]_D²⁰ +5.4 (c, 0.001 in H₂O).

(2R,4S)-form

d-cis-form
[175671-49-5]
Gum. [α]_D²⁰ +19.3 (c, 0.7 in H₂O).

(2S,4R)-form

L-cis-form
Isol. from leaves of *Calliandra pittieri* and *Strophanthus scandens*. Cryst. + 1 or 2H₂O (EtOH aq.). Mp 265° dec. [α]_D²⁵ -23.5 (c, 1 in H₂O) (99.8% ee).

Me ester

C₇H₁₃NO₃ 159.185
Beige solid (as hydrochloride). Mp 172.5-174.5° dec. (hydrochloride). [α]_D²⁵ +9.9 (c, 1.01 in MeOH) (95.8% ee).

tert-Butylamide

C₁₀H₂₀N₂O₂ 200.28
Solid. Mp 165.5-168°. [α]_D²⁵ -18.5 (c, 1.02 in MeOH).

N-Benzoyl

C₁₃H₁₅NO₄ 249.266
Mp 104° Mp 191° (dimorph.).

N-(2,4-Dinitrophenyl)

Yellow prisms (EtOH aq.). Mp 134°.

(2S,4S)-form

L-trans-form
[4382-31-4]
Isol. from heartwood, sapwood and leaves of *Acacia* spp. and *Armeria* spp. (Fabaceae, Plumbaginaceae). Prisms (EtOH aq.). Mp 294° dec. [α]_D²⁰ -13 (c, 1 in H₂O). [α]_D²⁰ -2.7 (c, 1 in 5M HCl).

Hydrochloride

Cryst. (EtOH aq.). Mp 161-163°.

4-O-Sulfate: [99694-77-6]

C₆H₁₁NO₆S 225.222
Isol. from seeds of *Peltophorum africanum*, *Peltophorum adnatum*, *Peltophorum dubium*, *Peltophorum ferrugineum*, *Peltophorum inerme*, *Peltophorum pterocarpum* and *Peltophorum tonkinense*. Also isol. from various sponges. Glutamate receptor agonist. Cryst. + 1H₂O (H₂O). Mp 248° dec. [α]_D²⁰ +6.5 (H₂O).

N-Benzoyl

C₁₃H₁₅NO₄ 249.266
Needles (EtOH aq.). Mp 174°. [α]_D¹⁵ -54 (c, 1 in EtOH).

N-Me: Ovalin

[73710-91-5]
C₇H₁₃NO₃ 159.185
Obt. from seeds of *Milletia ovalifolia* (Fabaceae). Mp 280-281°.

N-(2,4-Dinitrophenyl)

Orange prisms (EtOH aq.). Mp 183°.

Me ether: 4-Methoxy-2-piperidinecar-

boxylic acid. **4-Methoxypipelic acid**
[135607-86-2]
C₇H₁₃NO₃ 159.185
Isol. from *Inga paterno* and other *Inga*
spp. (Fabaceae).

(2R,5S)-form

(±)-trans-form
[143616-98-2]

Mp 161-163° (as hydrochloride). CAS
no. refers to hydrochloride.

Clark-Lewis, J.W. *et al.*, *J.C.S.*, 1961, 189 (*isol. struct*)

Shooley, J.N. *et al.*, *Acta Chem. Scand.*, 1962, **16**, 2547 (*pmr, conformm*)

Witkop, B. *et al.*, *J.A.C.S.*, 1964, **86**, 1844 (*synth*)

Murakoshi, I. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1723 (*isol, struct*)

Gupta, R.K. *et al.*, *Phytochemistry*, 1979, **18**, 2021-2022 (*Ovalin*)

Callens, R.E.A. *et al.*, *Bull. Soc. Chim. Belg.*, 1982, **91**, 713 (*synth*)

Evans, S.V. *et al.*, *Phytochemistry*, 1985, **24**, 2593-2596 (*4-sulfate*)

Morton, T.C. *et al.*, *Phytochemistry*, 1991, **30**, 2397 (*4-methoxypipelic acid*)

Pellicciari, R. *et al.*, *Med. Chem. Res.*, 1992, **2**, 491 (*synth, sulfate*)

Gillard, J. *et al.*, *J.O.C.*, 1996, **61**, 2226 (*2S,4R-form, synth, ir, pmr, cmr, bibl*)

Di Nardo, C. *et al.*, *J.O.C.*, 1999, **64**, 6119-6125 (*synth*)

Haddad, M. *et al.*, *Tetrahedron: Asymmetry*, 1999, 4231-4237 (*2S,4R-form, synth, pmr, cmr*)

Davis, F.A. *et al.*, *Synthesis*, 2000, 2106-2112 (*isomers, synth*)

Brooks, C.A. *et al.*, *Tet. Lett.*, 2000, **41**, 3551-3553 (*synth*)

Agami, C. *et al.*, *Eur. J. Org. Chem.*, 2001, 2385-2389 (*2S,4R-form, synth, pmr, cmr*)

Lloyd, R.C. *et al.*, *Org. Process Res. Dev.*, 2002, **66**, 762-766 (*isomers, synth, pmr, cmr*)

Sakai, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 784-787 (*4-sulfate, isol, pmr, cmr, activity*)

Cordero, F.M. *et al.*, *Eur. J. Org. Chem.*, 2006, 3235-3241 (*2R,4R-form, 2S,4S-form, synth, pmr*)


Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 5877-5887 (*Damipicolin*)

Occhiato, E.G. *et al.*, *Eur. J. Org. Chem.*, 2008, 524-531 (*2S,4R-form, synth, pmr, cmr*)

Alegret, C. *et al.*, *Eur. J. Org. Chem.*, 2008, 1789-1796 (*synth*)

5-Hydroxy-2-piperidinecarboxylic acid H-698

5-Hydroxypipelic acid
[13096-31-6]



(2S,5R)-form

C₆H₁₁NO₃ 145.158**(2S,5R)-form**

L-trans-form
[50439-45-7]

Present in dates (*Phoenix dactylifera*), Rhodesian teak (*Rhapix excelsa*), Acacia, leaves of *Morus alba* (white mulberry) and the seeds of *Caesalpinia* spp. and *Gleditsia* spp. (Arecaceae, Fabaceae, Moraceae, Caesalpinaceae). Mp 235°

dec. [α]_D²⁰ -23.1 (c, 1 in H₂O).

Hydrochloride:

Cryst. (EtOH aq.). Mp 210-215° dec.

[α]_D²⁰ -10.9 (c, 0.92 in H₂O).

N-Benzoyloxycarbonyl: [84409-89-2]

C₁₄H₁₇NO₅ 279.292

Cryst. (Me₂CO/petrol). Mp 150-152°.

[α]_D²⁰ -17.9 (c, 1 in Me₂CO).

(2S,5S)-form

L-cis-form

[63088-78-8]

Present in the leaves of *Morus alba* (white mulberry), *Gleditsia* spp. and *Lathyrus japonicus* (Fabaceae). Cryst. (EtOH aq.). Mp 230-235° dec. [α]_D²² -29.5 (c, 2 in H₂O). [α]_D²² -17.3 (6M HCl).

Hydrochloride:

Cryst. (EtOH aq./Et₂O). Mp 183-185° dec. [α]_D²⁴ -18.5 (c, 1 in H₂O).

Hydrobromide: Mp 200-201°.

Lactone: 2-Oxa-5-azabicyclo[2.2.2]octan-3-one

C₆H₉NO₂ 127.143

Platelets (as hydrobromide). Mp 228-231° (hydrobromide).

N-Benzoyloxycarbonyl: [917507-80-3]

C₁₄H₁₇NO₅ 279.292

Mp 131-132°. [α]_D²⁴ +76.5 (c, 0.17 in MeOH).

(2RS,5SR)-form

(±)-trans-form

Hydrochloride:

Cryst. (EtOH). Mp 192-194°.

Virtanen, A.I. *et al.*, *Acta Chem. Scand.*, 1954, **8**, 1290 (*isol*)

Witkop, B. *et al.*, *J.A.C.S.*, 1957, **79**, 192-197; 1964, **86**, 1844 (*isol, synth*)

Kristensen, I. *et al.*, *Tetrahedron*, 1976, **32**, 2799 (*nmr*)

Hatanaka, S.-I. *et al.*, *Phytochemistry*, 1977, **16**, 1041 (*isol*)

Evans, C.S. *et al.*, *Phytochemistry*, 1978, **17**, 1127 (*isol, occur*)

Callen, R.E.A. *et al.*, *Bull. Soc. Chim. Belg.*, 1982, **91**, 713 (*synth*)


Adams, D.R. *et al.*, *Chem. Comm.*, 1996, 349-350 (*synth*)

Botman, P.N.M. *et al.*, *Org. Lett.*, 2004, **6**, 4941-4944 (*synth*)

Jung, J.-C. *et al.*, *Tetrahedron: Asymmetry*, 2005, **17**, 2479-2486 (*2S,5R-form, 2S,5S-form, N-benzoyloxycarbonyl*)

5-Hydroxy-2-piperidine-methanol, 9CI H-699

6-(Hydroxymethyl)-3-piperidinol. 5-Hydroxy-2-(hydroxymethyl)piperidine

C₆H₁₃NO₂ 131.174**(2S,5R)-form**


1,3,4-Trideoxynojirimycin

Alkaloid from the pods of *Angylocalyx pynaertii*. [α]_D +36.6 (c, 0.23 in H₂O).

Yasuda, K. *et al.*, *J. Nat. Prod.*, 2002, **65**, 198-202 (*isol, pmr, cmr*)

7-Hydroxypleurocorine H-700

[155210-50-7]



Relative configuration


C₂₅H₃₁N₃O₄ 437.538

The parent Pleurocorine is not known. Alkaloid from leaves of *Pleurostyliya opposita* (Celastraceae). Amorph. [α]_D 0 (c, 1 in CHCl₃).

Séguineau, C. *et al.*, *Heterocycles*, 1994, **38**, 181 (*isol, ir, pmr, cmr, struct*)

2-Hydroxy-6-prenylindole H-701

6-(3-Methyl-2-butenyl)-1H-indol-2-ol. 1,3-Dihydro-6-(3-methyl-2-butenyl)-2H-indol-2-one. 6-Prenyl-2-indolinone. 6-Prenyloxindole


C₁₃H₁₅NO 201.268

Tautomeric with oxindole and 3H-forms. Alkaloid from *Monodora angolensis*. Orange solid.

Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (*isol, pmr, cmr*)

6-(1-Hydroxypropyl)lumazine H-702

6-(1-Hydroxypropyl)-2,4(1H,3H)-pteridinedione, 9CI



(S)-form

C₉H₁₀N₄O₃ 222.203**(S)-form**

1-Me: 6-(1-Hydroxypropyl)-1-methyllumazine. Leucettidine

[79121-29-2]

C₁₀H₁₂N₄O₃ 236.23

Isol. from the calcareous sponge

Leucetta microraphis. Sol. MeOH.

[α]_D²¹ -35.9 (c, 1.26 in MeOH).

Struct. revised in 1988. λ _{max} 237

(ϵ 13200); 248 (sh) (ϵ 12000); 334

(ϵ 7080) (MeOH) (Derep). λ _{max} 218

(ϵ 10400); 245 (ϵ 19500); 288 (ϵ 2900);

343 (ϵ 8320) (MeOH/NaOH)

(Derep).

3-Me: 6-(1-Hydroxypropyl)-3-methylumazine

[155758-91-1]

C₁₀H₁₂N₄O₃ 236.23

Isol. from the marine polychaete *Odontosyllis undecimdongata*. $[\alpha]_D^{25}$ -76 (c, 0.938 in MeOH) (synthetic).

1,3-Di-Me: 6-(1-Hydroxypropyl)-1,3-dimethylumazine

[155758-92-2]
C₁₁H₁₄N₄O₃ 250.257
From *Odontosyllis undecimdongata*.
Mp 173-174° (synthetic). $[\alpha]_D$ -59.6 (c, 1.079 in MeOH) (synthetic).

1'-Ketone: 6-(1-Oxopropyl)-2,4(1H,3H)-pteridinedione. 6-Propionyllumazine

[126857-77-0]
C₉H₈N₄O₃ 220.187
Isol. from the marine polychaete *Odontosyllis undecimdongata*. Mp 278-280° dec.

1'-Ketone, 3-Me: 3-Methyl-6-(1-oxopropyl)-2,4(1H,3H)-pteridinedione. 3-Methyl-6-propionyllumazine

[71014-16-9]
C₁₀H₁₀N₄O₃ 234.214
Isol. from *Odontosyllis undecimdongata*. Cryst. (CHCl₃/hexane). Mp 234° (227-228°).

1'-Ketone, 1,3-di-Me: 1,3-Dimethyl-6-(1-oxopropyl)-2,4(1H,3H)-pteridinedione. 1,3-Dimethyl-6-propionyllumazine

[71014-17-0]
C₁₁H₁₂N₄O₃ 248.241
Isol. from *Odontosyllis undecimdongata*. Cryst. (C₆H₆/petrol). Mp 146-147° (141-142°).

3'-Hydroxy, 1'-ketone, 3-Me: 6-(β-Hydroxypropionyl)-3-methylumazine

[135048-63-4]
C₁₀H₁₀N₄O₄ 250.213
From *Odontosyllis undecimdongata*. Mp 200° dec.

3'-Methoxy, 1'-ketone, 3-Me: 6-(β-Methoxypropionyl)-3-methylumazine

[135048-61-2]
C₁₁H₁₂N₄O₄ 264.24
From *Odontosyllis undecimdongata*. Mp 193-196° dec.

3'-Hydroxy, 1'-ketone, 1,3-di-Me: 6-(β-Hydroxypropionyl)-1,3-dimethylumazine

[163132-70-5]
C₁₁H₁₂N₄O₄ 264.24
From *Odontosyllis undecimdongata*. Mp 168-169° dec.

3'-Methoxy, 1'-ketone, 1,3-di-Me: 6-(β-Methoxypropionyl)-1,3-dimethylumazine

[135048-62-3]
C₁₂H₁₄N₄O₄ 278.267
From *Odontosyllis undecimdongata*. Mp 129-130°.

(±)-form

Cryst.

1-Me: Mp 195°.

Cardellina, J.H. *et al.*, *J.O.C.*, 1981, **46**, 4782-4784 (*Leucettidine*, isol, uv, ir, pmr, ms)

Baur, R. *et al.*, *Annalen*, 1984, 1798-1814 (*1'-ketone 1,3-di-Me*, synth)

Pfleiderer, W. *et al.*, *Tetrahedron*, 1988, **44**, 3373-3378 (*Leucettidine*, struct, synth, uv, pmr)


Inoue, S. *et al.*, *Chem. Lett.*, 1990, 367-368 (6-Propionyllumazine, 3-Methyl-6-propionyllumazine, 1,3-Dimethyl-6-propionyllumazine, isol)

Tanino, H. *et al.*, *Heterocycles*, 1994, **38**, 971-974 (*Odontosyllis undecimdongata* constits)

Kakoi, H. *et al.*, *Heterocycles*, 1995, **41**, 789-797 (*Odontosyllis undecimdongata* constits)

Hydroxypropyllycodine H-703

α-Methyllycodine-1-ethanol, 9CI
[123118-09-2]




C₁₉H₂₈N₂O 300.443
Alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Opaque solid. $[\alpha]_D$ +11 (c, 0.39 in MeOH).

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077 (isol, uv, ir, pmr, cmr, struct)

2-(2-Hydroxypropyl)-1-methyl-5-(2-oxopropyl)pyrrolidine H-704

1-[5-(2-Hydroxypropyl)-1-methyl-2-pyrrolidinyl]-2-propanone, 9CI. 2-Acetyl-5-(2-hydroxypropyl)-1-methylpyrrolidine
[202057-29-2]



C₁₁H₂₁NO₂ 199.292
Alkaloid from *Erythroxylum lucidum* (Erythroxylaceae) and *Merremia* sp. (Convolvulaceae).

2'-Ketone: 1,1'-(1-Methyl-2,5-pyrrolidinediyl)bis-2-propanone. 2,5-Diacetyl-1-methylpyrrolidine
[36295-25-7]
[36295-52-0]


C₁₁H₁₉NO₂ 197.277
Alkaloid from *Erythroxylum lucidum* (Erythroxylaceae) and *Merremia* sp. (Convolvulaceae). Mp 176-177° (as picrate).

Hess, J. *et al.*, *Chem. Ber.*, 1972, **105**, 441-453 (synth)

Brachet, A. *et al.*, *Phytochemistry*, 1997, **46**, 1439-1442 (isol, struct)

2-(2-Hydroxypropyl)-6-methylpiperidine H-705

α,6-Dimethyl-2-piperidineethanol, 9CI. Pinidinol



C₉H₁₉NO 157.255

(2R,2'R,6R)-form [129785-16-6]

Alkaloid from *Picea engelmannii* and its root parasite *Pedicularis bracteosa*. Also found in *Picea abies*, *Picea pungens* and *Picea sitchensis*. Shows antifeedant activity on the Spruce budworm. $[\alpha]_D^{26}$ -15 (c, 0.55 in CHCl₃).

Stermitz, F.R. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1019-1020 (*cryst struct, abs config*)

Schneider, M.J. *et al.*, *J. Nat. Prod.*, 1991, **54**, 905-909 (isol)


Molander, G.A. *et al.*, *J.O.C.*, 2001, **66**, 4344-4347 (synth)

Gebauer, J. *et al.*, *Heterocycles*, 2006, **68**, 2129-2132 (synth)

Davis, F.A. *et al.*, *J.O.C.*, 2008, **73**, 9619-9626 (synth)

2-(3-Hydroxypropyl)-4(3H)-quinazolinone H-706

Pegamine
[31431-93-3]



C₁₁H₁₂N₂O₂ 204.228
Alkaloid from *Peganum harmala* (Zygophyllaceae). Mp 160-161°.


O-Ac: Mp 173-174°.

Khachimov, Kh.N. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 453; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 462 (isol, uv, ir, pmr, ms, struct)

Mhaske, S.B. *et al.*, *J.O.C.*, 2002, **66**, 9038-9040 (synth)

4-Hydroxy-2-propylquinoline H-707

2-Propyl-4(1H)-quinolinone. 2-Propyl-4-quinolinol



C₁₂H₁₃NO 187.241

Alkaloid from the leaves and terminal branches of *Boronia ternata*. Needles (C₆H₆/petrol). Mp 214-216° (as hydrochloride). Artifact of the isol. procedure.

NH-form

N-Me: 1-Methyl-2-propyl-4(1H)-quinolinone. Leptomerine
[22048-97-1]

C₁₃H₁₅NO 201.268
Alkaloid from the aerial parts of *Haplophyllum leptomerum* (Rutaceae). Cryst. (EtOAc). Mp 149-150°.

N-(Acetoxymethyl): 1-Acetoxymethyl-2-propyl-4(1H)-quinolinone
C₁₅H₁₇NO₃ 259.304

Alkaloid from leaves and terminal branches of *Boronia ternata* (Rutaceae). Cryst. (C₆H₆/petrol). Mp 112°.

OH-form

Me ether: 4-Methoxy-2-propylquinoline.

Chimarine A

[22048-96-0]

C₁₃H₁₅NO 201.268

Alkaloid from the trunk bark of
Galipea longiflora (Rutaceae). Oil.

Duffield, A.M. *et al.*, *Aust. J. Chem.*, 1963, **16**, 292 (*isol. uv. struct. deriv*)


Werner, W. *et al.*, *Tetrahedron*, 1969, **25**, 255-261 (*synth*)

Akhmedzhanova, V.I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 78-79 (*Leptomerine*)

Fournet, A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1547-1552 (*Chimarine A*)

6-Hydroxy-2-pyridineacetic acid H-708 acid

1,6-Dihydro-6-oxo-2-pyridineacetic acid.
2-Pyridone-6-acetic acid



C₇H₇NO₃ 153.137

Et ester:

C₉H₁₁NO₃ 181.191

Cryst. (C₆H₆). Mp 170°.

Amide:

C₇H₈N₂O₂ 152.152

Cryst. (EtOH). Mp 254°.

NH-form

N-β-D-Glucopyranosyl: [117803-24-4]

C₁₃H₁₇NO₈ 315.279

Isol. from the Algerian newt, *Pleurodeles walthii*. Cryst. λ_{max} 316 (H₂O).

N-Me, Et ester: [103096-12-4]

C₁₀H₁₃NO₃ 195.218

Mp 100-101°.

N-Benzyl: [107922-40-7]

C₁₄H₁₃NO₃ 243.262

Cryst. (H₂O). Mp 169° dec.

OH-form

Et ether, Et ester: [99986-53-5]

C₁₁H₁₅NO₃ 209.244

Oil. Bp_{0.3} 80°.

Et ether, nitrile: 2-(Cyanomethyl)-6-ethoxypyridine

[53676-89-4]

C₉H₁₀N₂O 162.191

Mp 54°. Bp_{0.05} 95°.


Adams, R. *et al.*, *J.A.C.S.*, 1954, **76**, 3168;

1959, **81**, 2537 (*derivs. synth*)

Yoshida, T. *et al.*, *Eur. J. Biochem.*, 1988, **175**, 41-44 (*l-glucosyl*)

4-Hydroxy-3-pyridinecarboxylic acid H-709

1,4-Dihydro-4-oxo-3-pyridinecarboxylic acid. 4(1H)-Pyridone-3-carboxylic acid. 4-Hydroxynicotinic acid [609-70-1]



C₆H₅NO₃ 139.11

Needles (H₂O). Mp 250°.

Me ester: [67367-24-2]

C₇H₇NO₃ 153.137

Needles (hexane). Mp 221-222°.

Et ester: [57905-31-4]

C₈H₉NO₃ 167.164

Cryst. (EtOH). Mp 219-220°.

Amide: 4-Hydroxynicotinamide

[7418-63-5]

C₆H₆N₂O₂ 138.126

Isol. from ferns *Arachniodes festina* and *Arachniodes nigrospinoso*. Cryst. (H₂O). Mp 276-278° (263-265°).

NH-form

1-α-D-Ribofuranosyl, amide: [171284-19-8]

C₁₁H₁₄N₂O₆ 270.241

Isol. from human urine.

1-β-D-Ribofuranosyl, amide: 1,4-Dihydro-4-oxo-1-β-D-ribofuranosyl-3-pyridinecarboxamide, 9CI. 1-β-D-Ribofuranosyl-4-pyridone-3-carboxamide

[78686-01-8]

C₁₁H₁₄N₂O₆ 270.241

Isol. from human urine. Constit. of *Rothmannia longiflora*. Cryst. (EtOH). Mp 211-212°. [α]_D²⁵ -97.1 (c. 0.25 in MeOH).

N-Me, nitrile: 1,4-Dihydro-1-methyl-4-oxo-3-pyridinecarbonitrile. 3-Cyano-1-methyl-4(1H)-pyridinone. **Mallorepine** [767-98-6]

C₇H₈N₂O 134.137

Alkaloid from *Mallotus repandus*.

Prisms (MeOH). Mp 185-186° (179-181°). λ_{max} 258 (log ε 4.21); 263 (log ε 4.2); 289 (log ε 3.76) (MeOH).

OH-form

O-β-D-Glucopyranoside:

C₁₂H₁₅NO₈ 301.252

Alkaloid from *Conyza bonariensis*.

Needles. Mp 203-205°. [α]_D²⁵ -250.4 (c. 1.35 in MeOH).

Me ether: 4-Methoxy-3-pyridinecarboxylic acid

[10177-31-8]

C₇H₇NO₃ 153.137

Plates (EtOH). Mp 178-179° dec.

Me ether, Me ester: [10177-32-9]

C₈H₉NO₃ 167.164

Cryst. (cyclohexane). Mp 82-84°.

Me ether, nitrile: 3-Cyano-4-methoxypyridine

[74133-20-3]

C₇H₈N₂O 134.137

Alkaloid from *Hernandia nymphaefolia* (Hernandiaceae). Needles (Et₂O). Mp 124°.

Me ether, nitrile, N-oxide: 3-Cyano-4-methoxypyridine 1-oxide. **Malloapeltine** [220698-57-7]

C₇H₈N₂O₂ 150.137

Alkaloid from the roots of *Mallotus apelta*. Amorph. powder. Mp 204-206°. λ_{max} 220 (log ε 1.93); 280 (log ε 1.53) (MeOH). λ_{max} 220 (ε 85); 280 (ε 32) (MeOH) (Berdy).

Et ether: 4-Ethoxy-3-pyridinecarboxylic acid

C₈H₉NO₃ 167.164

Prisms (EtOH). Mp 178° dec.

[10177-31-8]

Wieland, T. *et al.*, *Annalen*, 1961, **642**, 163-173 (*synth. nitrile, amide, Me ether amide*)

Robinson, T. *et al.*, *Phytochemistry*, 1965, **4**, 75-79 (*synth. Malloapeltine*)

Ross, W.C.J. *et al.*, *J.C.S. (C)*, 1966, 1816-1821 (*synth*)

Kitagawa, T. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1403-1414 (*uv. struct*)

Hikino, H. *et al.*, *Planta Med.*, 1978, **33**, 385-388 (*Mallorepine*)

Yukuskijin, K. *et al.*, *Phytochemistry*, 1980, **19**, 161-162 (*isol. Me ether nitrile, ir. uv. pmr*)

Tanaka, N. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 1355-1358 (*amide*)

Dutta, S.P. *et al.*, *Nucleic Acid Chem.*, 1991, **4**, 152-154 (*synth. riboside*)

Chheda, G.B. *et al.*, *Nucleosides Nucleotides*, 1995, **14**, 1519-1537 (*isol. riboside*)

Leroy, F. *et al.*, *Synth. Commun.*, 1996, **26**, 2257 (*Me ether Me ester*)

Cheng, X.-F. *et al.*, *Phytochemistry*, 1998, **49**, 2193-2194 (*Malloapeltine*)


Bringmann, G. *et al.*, *Phytochemistry*, 1999, **51**, 271-276 (*isol. riboside*)

Kong, L.D. *et al.*, *Phytochemistry*, 2001, **58**, 645-651 (*glucoside*)

5-Hydroxy-2-pyridinecarboxylic acid, 9CI H-710

5-Hydroxypicolinic acid

[15069-92-8]



C₆H₅NO₃ 139.11

Detected in culture filtrate of *Nocardia* sp. Isol. from marine macrophytes. Cryst. + 1H₂O (H₂O) or powder. Mp 269-271° (258°). λ_{max} 250 ; 283 (EtOH) (Berdy). λ_{max} 252 (EtOH-HCl) (Berdy). λ_{max} 273 ; 303 (EtOH-NaOH) (Berdy).

Diesel, B.F. *et al.*, *J.A.C.S.*, 1949, **71**, 1866

(*synth*)

Paulsen, H. *et al.*, *Chem. Ber.*, 1973, **106**, 1525 (*synth*)

Entsch, B. *et al.*, *J. Biol. Chem.*, 1976, **251**, 2550 (*biochem*)

Grachev, V.T. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1977, 2273 (*uv. ir*)


Lezina, V.P. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1980, 98; 1981, 2218 (*pmr*)

Oehlke, J. *et al.*, *Pharmazie*, 1983, **38**, 591; 624 (*synth. derivs. ms. uv. pmr*)

Makar'eva, T.N. *et al.*, *Khim. Prir. Soedin.*, 1989, 140-141; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 125-126 (*isol*)

6-Hydroxy-3-pyridinecarboxylic acid H-711

1,6-Dihydro-6-oxo-3-pyridinecarboxylic acid, 9CI. 6-Hydroxynicotinic acid, 8CI. 6(1H)-Pyridone-3-carboxylic acid. α-Pyridone-5-carboxylic acid [5006-66-6]



C₆H₅NO₃ 139.11

Pyridone (*NH*)-form predominates. Needles (EtOH). Mp 304° dec. pK_{a1} 3.82 (20°).

Me ester: [66171-50-4]

$C_7H_7NO_3$ 153.137

Leaflets (Me₂CO). Mp 164°. pK_a 9.92 (20°).

Et ester: [18617-50-0]

$C_8H_9NO_3$ 167.164

Cryst. (Me₂CO). Mp 149-150°.

NH-form

N-Me: 1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarboxylic acid, 9CI. *Nudifloric acid*

[3719-45-7]

$C_7H_7NO_3$ 153.137

Mp 238°.

N-Me, Me ester: 1-Methyl-5-carbomethoxy- α -pyridone

[6375-89-9]

$C_8H_9NO_3$ 167.164

Alkaloid from the leaves of *Ampelocera ruizii* (Ulmaceae). Cryst. (Me₂CO). Mp 132°.

N-Me, amide: [701-44-0]

$C_7H_8N_2O_2$ 152.152

Cryst. Mp 214-215°.

N-Me, nitrile: 5-Cyano-1-methyl-2(1H)-pyridinone. *Nudiflorine*

[768-45-6]

$C_7H_6N_2O$ 134.137

Alkaloid from the leaves of *Trewia nudiflora* (Euphorbiaceae). Needles (CHCl₃/petrol). Mp 161°.

OH-form

Me ether: 6-Methoxy-3-pyridinecarboxylic acid

[66572-55-2]

$C_7H_7NO_3$ 153.137

Needles (H₂O). Mp 237-238° Mp 171.5-172.5°.

Me ether, Et ester: [74925-37-4]

$C_9H_{11}NO_3$ 181.191

Prisms (EtOH). Mp 71°. Bp_{0.25} 135°.

Me ether, nitrile: 3-Cyano-6-methoxypyridine

[15871-85-9]

$C_7H_6N_2O$ 134.137

Solid. Mp 97°.

Et ether: 6-Ethoxy-3-pyridinecarboxylic acid

$C_8H_9NO_3$ 167.164

Cryst. (EtOH). Mp 183°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 787D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 322A (nmr)

Reissert, A. *et al.*, *Ber.*, 1895, 28, 119 (*Et ether*)

Meyer, H. *et al.*, *Monatsh. Chem.*, 1907, 28, 59 (*derivs*)

Org. Synth., Coll. Vol., 4, 1963, 532 (*synth*)

Mukherjee, R. *et al.*, *Tetrahedron*, 1966, 22, 1461 (*isol, uv, pmr, synth, Nudiflorine*)

Gault, H. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1968, 266, 131 (*synth*)

Weber, H. *et al.*, *Chem. Ber.*, 1971, 104, 1478; 1985, 118, 4086 (*synth*)

Burnell, R.H. *et al.*, *J. Nat. Prod.*, 1975, 38, 444 (*isol, uv, ir, pmr, ms, struct, synth, deriv*)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1976, 24, 1870 (*synth*)

Bernofsky, C. *et al.*, *Anal. Biochem.*, 1979, 96, 189 (*synth*)

Buurman, D.J. *et al.*, *J. Het. Chem.*, 1986, 23, 1015-1018 (*N-Me amide, N-Me nitrile*)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 4, 2769-2771 (*use*)


Ghosez, L. *et al.*, *Tetrahedron*, 1999, 55, 3387-3400 (*Me ester*)

Khanna, I.K. *et al.*, *J. Med. Chem.*, 2000, 43, 3168-3185 (*Me ether nitrile, synth, pmr*)

Sobczak, A. *et al.*, *Synth. Commun.*, 2005, 35, 2993-3001 (*Et ester*)

Gupta, S. *et al.*, *Acta Cryst. E*, 2007, 63, o2784 (*cryst struct*)

4-Hydroxy-2-[4-(2-pyridinyl)-1,3-butadienyl]benzoic acid H-712



$C_{16}H_{13}NO_3$ 267.284

Me ester: [125508-38-5]


$C_{17}H_{15}NO_3$ 281.31

Alkaloid from the aerial parts of *Dehaasia kurzii* (Lauraceae).

Rahman, M.A. *et al.*, *CA*, 1990, 112, 95569k (*isol, uv, ir, pmr, ms, struct*)

N-[(5-Hydroxy-2-pyridinyl)-methyl]adenosine, 9CI H-713

6-(5-Hydroxy-2-pyridinylmethylamino)-9 β -ribofuranosylpurine. *AMG 1* [123369-41-5]




$C_{16}H_{18}N_6O_5$ 374.355

Isol. from *Armillaria mellea* (honey mushroom). Cytokinin. Powder. Sol. H₂O, MeOH; poorly sol. EtOAc, hexane. λ_{max} 268 (MeOH) (Berdy). λ_{max} 265 (MeOH-HCl) (Berdy). λ_{max} 264 (MeOH-NaOH) (Berdy).

Watanabe, N. *et al.*, *Planta Med.*, 1990, 56, 48 (*isol, pmr, cmr, struct*)

4-Hydroxy-1H-pyrrole-2-carboxylic acid H-714

Hydroxyminaline



989

$C_5H_5NO_3$ 127.099

Component of the molecule pectase found in *Penicillium* and *Aspergillus*. Grey-brown amorph. powder which cannot be recryst. or subl. Dec. readily in soln.

Me ether: 4-Methoxy-2-pyrrolecarboxylic acid

$C_6H_7NO_3$ 141.126

Cryst. by subl. Mp 179-180°.

Me ether, Me ester:

$C_7H_9NO_3$ 155.153

Cryst. (MeOH). Mp 85-86°.

Me ether, Et ester:

$C_8H_{11}NO_3$ 169.18

Cryst. (C₆H₆). Mp 55-58°.

Minagawa, T. *et al.*, *Proc. Imp. Acad. (Tokyo)*,


1945, 21, 33; 37; *CA*, 47, 151i; 152b (*isol*)

Kuhn, R. *et al.*, *Chem. Ber.*, 1956, 89, 1423 (*synth, uv*)

Rapoport, H. *et al.*, *J.A.C.S.*, 1962, 84, 630 (*derivs, synth, uv, ir*)

4-Hydroxy-2-pyrrolidinecarboxylic acid H-715

4-Hydroxyproline. *Oxyproline*



$C_5H_9NO_3$ 131.131

(2R,4R)-form

D-allo-form. *D*-cis-form

[2584-71-6]

Mp 237-241°. $[\alpha]_D^{25}$ +58.6 (c, 0.65 in H₂O). Inispid taste.

(2R,4S)-form

D-trans-form

[3398-22-9]

Mp 274°. $[\alpha]_D^{21}$ +75.2 (H₂O). Inispid taste.

(2S,4R)-form

L-trans-form

[51-35-4]

Constit. of proteins. Chiral building block. Mp 274°. $[\alpha]_D^{26}$ -74.6. pK_{a1} 1.92; pK_{a2} 9.73 (25°). Sweet taste.

β -*D*-Galactopyranosyl ester: [79284-73-4]

$C_{11}H_{19}NO_8$ 293.273

Isol. from hydrolysed wheat endosperm.

O- β -*D*-Galactopyranoside: [80185-44-0]

$C_{11}H_{19}NO_8$ 293.273

Isol. from hydrolysed wheat endosperm.

Me ester: [1499-56-5]

$C_6H_{11}NO_3$ 145.158

Needles (MeOH/EtOAc) (as hydrochloride). Mp 162-164°.

Et ester, hydrochloride: [33996-30-4]

[61478-25-9]

Cryst. Mp 153-153.5°.

N-Ac: *Oxaceprol*, *INN*. *Jonctum*. *CO 61*

[33996-33-7]

$C_7H_{11}NO_4$ 173.168

Antirheumatic, dermatological agent.

Mp 135°. $[\alpha]_D$ -118.5 (H₂O). Log P - 1.11 (calc).

O-Benzoyl: [54665-21-3]
C₁₂H₁₃NO₄ 235.239
Mp 218-219°. $[\alpha]_D^{25}$ -6.66 (c, 0.21 in MeOH).

N-Benzoyl, Me ester: [31560-20-0]
C₁₃H₁₅NO₄ 249.266
Cryst. (EtOAc). Mp 145-146° (141°). $[\alpha]_D^{25}$ -142.4 (c, 1.6 in CHCl₃).

N-Benzoyl, tert-butyl ester: [158177-75-4]
C₁₆H₂₁NO₄ 291.346
Cryst. (Et₂O). Mp 90°. $[\alpha]_D^{21}$ -137.2 (c, 1 in CHCl₃).

N-Benzoyloxycarbonyl: [13504-85-3]
C₁₃H₁₅NO₅ 265.265
Cryst. Mp 106-107°. $[\alpha]_D^{24}$ -75.5 (c, 1.03 in CHCl₃).

N-Me: N-Methyl-trans-4-hydroxy-L-proline. **4-Hydroxyhygrinic acid.** *Aceprolinum*. Joint. Problaston. Thioprol [4252-82-8]
C₆H₁₁NO₃ 145.158
Constit. of *Capsicum annuum* (sweet pepper), *Copaifera* spp., *Croton gubougia*, *Dalbergia sympathetica*, *Erythroxylum argentinum*, *Toddalia aculeata* and *Afromosia elata* (preferred genus name *Pericopsis*). Also from the red alga *Chondria coerulescens* and other red algae.
Cryst. (MeOH). Mp 238-240° dec. $[\alpha]_D$ -86.6 (c, 1.5 in H₂O). Isolate from *Dalbergia sympathetica* originally thought to be a 2-piperidone but struct. was corrected by Winkler, *et al* in 2006.

N-Me, hydrochloride: Mp 181-183°. $[\alpha]_D$ -55 (c, 2.3 in MeOH).

N-Me, O-Ac: [90088-50-9]
C₈H₁₃NO₄ 187.195
Cryst. (EtOAc). $[\alpha]_D$ +67 (c, 2 in MeOH).

N-Heptyl: [76666-35-8]
C₁₂H₂₃NO₃ 229.319
Hplc stationary phase for resoln. of amino acids.

N-Hexadecyl: [76652-69-2] Hplc stationary phase for resoln. of amino acids.

N-Benzyl, Et ester: [154342-88-8]
C₁₄H₁₉NO₃ 249.309
Oil. $[\alpha]_D^{20}$ -56.3 (c, 1 in CHCl₃).

N-Benzyl, amide: [500733-22-2]
C₁₂H₁₆N₂O₂ 220.271
Yellowish cryst. Mp 39°. $[\alpha]_D^{20}$ -79.6 (c, 1 in CHCl₃).

Me ether: 4-Methoxy-2-pyrrolidinecarboxylic acid
C₆H₁₁NO₃ 145.158
Mp 202°. $[\alpha]_D^{20}$ -56 (c, 1 in H₂O).

Me ether, N-Me: trans-4-Methoxy-N-methyl-L-proline [131559-49-4]
C₇H₁₃NO₃ 159.185
Alkaloid from *Petiveria alliacea*. Mp 251-253°. $[\alpha]_D^{20}$ -32.9 (c, 0.7 in MeOH).

N-Nitroso: [30310-80-6]
C₅H₈N₂O₄ 160.129
Cryst. (MeOH/butanol). Mp 138-140° dec.

► Potential carcinogen.

(2S,4S)-form

L-allo-form. *L*-cis-form

[618-27-9]

Occurs in toxic peptides of the death cap mushroom (*Amanita phalloides*). Mp 238-241°. $[\alpha]_D^{25}$ -57.7 (c, 0.65 in H₂O). Sweet taste.

Me ester

Needles + 1H₂O (EtOAc/hexane) (as hydrochloride). Mp 79-80° (hydrochloride). $[\alpha]_D$ -49.2 (c, 0.5 in MeOH).

N-Ac: [66267-44-5]

Cryst. (EtOH/Et₂O). Mp 144-145°. $[\alpha]_D$ -91.5 (c, 2 in H₂O).

N-Ac, Me ester: [78804-96-3]

C₈H₁₃NO₄ 187.195
Cryst. (Et₂O/EtOH). Mp 78°.

N-tert-Butyloxycarbonyl: [13726-69-7]

C₁₀H₁₇NO₅ 231.248
Oil.

N-tert-Butyloxycarbonyl, dicyclohexylamine salt (1:1): [21157-12-0]

Cryst. (EtOAc/petrol). Mp 194-195°. $[\alpha]_D^{25}$ -25.3 (DMF).

N-Benzoyloxycarbonyl:

Cryst. (EtOAc/petrol). Mp 106-107°. $[\alpha]_D^{20}$ -72 (c, 1 in CHCl₃).

(2RS,4RS)-form

D,L-allo-form. (\pm)-cis-form

[49761-17-3]

Mp 238°.

Et ester:

[51814-20-1 (hydrochloride)]

C₇H₁₃NO₃ 159.185

Mp 133-136° (as hydrochloride).

N-Ac:

Cryst. (EtOH/Et₂O). Mp 143-144°.

N-Ac, Me ester:

Cryst. (Et₂O/EtOH). Mp 79-80°.

(2RS,4SR)-form

(\pm)-trans-form

[618-28-0]

Cryst. (EtOH aq.). Mp 247°.

Et ester:

Cryst. (EtOH). Mp 142°.

[67463-44-9, 89771-43-7]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 583D; 584B; 584A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 887B; 887C (*nmr*)

Goodson, J.A. *et al.*, *J.C.S.*, 1919, 923 (*isol. struct. deriv*)

Patchett, A.A. *et al.*, *J.A.C.S.*, 1957, **79**, 185-192 (*N*-benzyloxycarbonyl)

Morgan, J.W.W. *et al.*, *Chem. Ind. (London)*, 1964, 542 (*deriv*)

Abraham, R.J. *et al.*, *J.C.S.*, 1964, 3739 (*pmr*)

Schnabel, E. *et al.*, *Annalen*, 1967, **702**, 188-196 (*N*-tert-butylloxycarbonyl)

U.K. Pat., 1971, 1 246 141; *CA*, **75**, 140679b (*Oxaceprol*)

Thomas, W.A. *et al.*, *Chem. Comm.*, 1972, 788 (*pmr, cryst struct, Oxaceprol*)

Koetzle, T.F. *et al.*, *Acta Cryst. B*, 1973, **29**, 231 (*cryst struct*)

Kaneko, T. *et al.*, *Synth. Prod. Util. Amino Acids*, 1974, 123 (*rev*)

Shamala, N. *et al.*, *Acta Cryst. B*, 1976, **32**, 3267 (*cryst struct*)

Roques, B.P. *et al.*, *Tetrahedron*, 1976, **32**, 1517 (*cmr, Oxaceprol*)

Ramaswamy, S.G. *et al.*, *J.O.C.*, 1977, **42**, 3440 (*synth*)

Hospital, M. *et al.*, *Biopolymers*, 1979, **18**, 1141 (*cryst struct, Oxaceprol*)

Davankov, V.A. *et al.*, *Chromatographia*, 1980, **13**, 677 (*N*-heptyl, *N*-hexadecyl)

Hara, J. *et al.*, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 3871 (*synth*)

Baker, G.L. *et al.*, *J.O.C.*, 1981, **46**, 2954-2960 (*Et ester, synth, ir, pmr, cmr*)

Field, L. *et al.*, *J.O.C.*, 1981, **46**, 17114182 (*synth, derivs*)

Strahm, A. *et al.*, *Phytochemistry*, 1981, **20**, 1061 (*galactosyl ester*)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 13051

Sciuto, S. *et al.*, *Phytochemistry*, 1983, **22**, 2311 (*isol, pmr, cmr, deriv*)

Inigo, R.P.A. *et al.*, *An. Asoc. Quim. Argent.*, 1984, **72**, 1428 (*isol, deriv*)

Thaning, M. *et al.*, *Helv. Chim. Acta*, 1986, **69**, 1711 (*derivs*)

Figliuolo, R. *et al.*, *Phytochemistry*, 1987, **26**, 3255 (*isol, ir, pmr, cmr, ms, deriv*)

Jones, G.P. *et al.*, *Phytochemistry*, 1987, **26**, 3343 (*isol, deriv*)

Jones, G.P. *et al.*, *Acta Cryst. C*, 1988, **44**, 2208 (*cryst struct, deriv*)

Papaioannon, D. *et al.*, *Acta Chem. Scand.*, 1990, **44**, 243 (*synth, pmr*)

De Martino, G. *et al.*, *J. Het. Chem.*, 1990, **27**, 507 (*synth*)

De Sousa, J.R. *et al.*, *Phytochemistry*, 1990, **29**, 3653 (*Me ether N-Me*)

Bridges, R.J. *et al.*, *J. Med. Chem.*, 1991, **34**, 715-725 (*N*-benzyloxycarbonyl)

Remuzon, P. *et al.*, *J. Med. Chem.*, 1992, **35**, 2898 (*Ac*)

Burger, K. *et al.*, *Angew. Chem., Int. Ed.*, 1993, **32**, 285 (*synth*)

Matsuoka, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1994, **58**, 1747 (*synth*)

Mehlfuehrer, M. *et al.*, *Chem. Comm.*, 1994, 1291 (*synth*)

Seki, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 1161 (*synth, pmr, cmr*)

Blanco, M.-J. *et al.*, *J.O.C.*, 1996, **61**, 4748 (*Me ester, synth, pmr, cmr*)

Remuzon, P. *et al.*, *Tetrahedron*, 1996, **52**, 13803-13835 (*2S,4R*-form, *rev. use in chiral synth*)

Lowe, G. *et al.*, *J.C.S. Perkin 1*, 1997, 539-546 (*isom, pmr*)

Baldwin, J.E. *et al.*, *Tetrahedron*, 1997, **53**, 5233-5254 (*2S,4R*-form, *N*-benzoyl esters, *synth*)

Lewis, A. *et al.*, *J.C.S. Perkin 1*, 1998, 3767-3776; 3777-3794 (*Me ester, synth, ir, pmr, cmr, ms*)

Ohtake, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1999, **72**, 2737-2754 (*O*-benzoyl)

Puripattanavong, J. *et al.*, *Planta Med.*, 2000, **66**, 740-745 (*isol, N-Me*)

Kimura, R. *et al.*, *Bull. Chem. Soc. Jpn.*, 2002, **75**, 2517-2525 (*2S,4S*-form, *synth, ir, pmr*)

Croce, P.D. *et al.*, *Tetrahedron: Asymmetry*, 2002, **13**, 197-201 (*2R,4R*-form, *synth, pmr, cmr*)

Schuch, C.M. *et al.*, *Tetrahedron: Asymmetry*, 2002, **13**, 1973-1980 (*2R,4S*-form, *synth*)

El-Ashry, E.H. *et al.*, *Carbohydr. Res.*, 2003, **338**, 2265-2290 (*rev, synth*)


Heindl, C. *et al.*, *Tetrahedron: Asymmetry*, 2003, **14**, 3141-3152; 3153-3172 (*2S,4R*-form, *Et ester N*-benzyl, *amide N*-benzyl)

Winkler, T. *et al.*, *Magn. Reson. Chem.*, 2006, **44**, 571-572 (*N-Me, pmr, cmr*)

Jain, S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1005-1010 (*N-Me, pmr, cmr, cryst struct*)

Fonari, M.S. *et al.*, *Tetrahedron*, 2006, **62**, 8779-8786 (2*S*,4*R*-form, *N*-nitroso, synth, cryst struct, pmr, cmr)
Dekebo, A. *et al.*, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 421-426 (*Capsicum*, isol)

4-Hydroxy-3-pyrrolidinecarboxylic acid H-716



C₅H₉NO₃ 131.131


(3*R**,4*S**)-form (+)-trans-form

N,N-Di-Me, betaine: 4-Hydroxy-*N,N*-dimethyl-3-pyrrolidinecarboxylate [117845-15-5]
C₇H₁₃NO₃ 159.185
Isol. from the Mediterranean red alga *Grateloupia proteus*. Off-white hygroscopic powder. [α]_D²⁵ +14.42 (c, 0.34 in H₂O).

Sciuto, S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1017 (isol, pmr, cmr, ms, struct)

3-Hydroxy-2-pyrrolidine-methanol, 9CI H-717

3-Hydroxy-2-(hydroxymethyl)pyrrolidine. 1,2,4-Trideoxy-1,4-iminopentitol. CYB 3



C₅H₁₁NO₂ 117.147

(2*R*,3*S*)-form

D-erythro-form
Alkaloid from the seeds of *Castanospermum australe*. Mp 108-112° (as hydrochloride). [α]_D²¹ +46.5 (H₂O) (hydrochloride).

N-(2-Hydroxyethyl): 3-Hydroxy-*N*-(2-hydroxyethyl)-2-(hydroxymethyl)pyrrolidine. 3-Hydroxy-2-(hydroxymethyl)-1-pyrrolidineethanol, 9CI [136600-18-5]
C₇H₁₅NO₃ 161.2

Alkaloid from the seeds of *Castanospermum australe*. Oil.

Nash, R.J. *et al.*, *Chem. Comm.*, 1985, 738 (pmr, ms, cryst struct)

Ikota, N. *et al.*, *Heterocycles*, 1988, **27**, 2535 (synth)

Molyneux, R.J. *et al.*, *Biochemistry*, 1991, **30**, 9981 (2-hydroxyethyl)

Griffart-Brunet, D. *et al.*, *Tet. Lett.*, 1994, **35**, 119 (synth)

Huwe, C.M. *et al.*, *Synthesis*, 1997, 61 (synth, pmr, cmr)

Malle, B.M. *et al.*, *J. Carbohydr. Chem.*, 2000, **19**, 573-583 (synth, pmr, cmr)


Hulme, A.N. *et al.*, *J.C.S. Perkin 1*, 2002, 1083-1091 (synth)

El-Ashry, H.E. *et al.*, *Carbohydr. Res.*, 2003, **338**, 2265-2290 (rev, synth)

Merino, P. *et al.*, *Eur. J. Org. Chem.*, 2008, 2929-2947 (synth)

4-Hydroxy-2-pyrrolidinone, 9CI H-718

4-Hydroxy-2-pyrrolidone, 8CI [25747-41-5]



(*R*)-form

C₄H₇NO₂ 101.105

(*R*)-form [40759-90-8]
Cryst. Mp 157-158°. [α]_D²⁵ +57.3 (c, 1.40 in H₂O).

(*S*)-form [22677-21-0]
Isol. from *Amanita muscaria*. Cryst. Mp 155-157°. [α]_D²⁵ -55.5 (c, 1.04 in H₂O).

(±)-form [62624-29-7]
Plates (Me₂CO). Mp 123-125°.

► UY5775500

Ac: 4-Acetoxy-2-pyrrolidone [64097-45-6]
C₆H₉NO₃ 143.142
Needles (C₆H₆/hexane). Mp 91-92°.

N-tert-Butyloxycarbonyl: [409341-03-3]
C₉H₁₅NO₄ 201.222
Solid. Mp 150.4-151.4° dec.

N-Benzyl: [125370-52-7]
C₁₁H₁₃NO₂ 191.229
Oil.

Matsumoto, T. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 716 (isol, ir, pmr)

Seiler, N. *et al.*, *Z. Anorg. Allg. Chem.*, 1970, **252**, 127 (ms)

Piffner, G. *et al.*, *Farmaco, Ed. Sci.*, 1977, **32**, 602 (synth)

Pellegata, R. *et al.*, *Synthesis*, 1978, 614 (synth)
Baker, J.T. *et al.*, *J.O.C.*, 1979, **44**, 2798 (synth, nmr)

Bladé-Font, A. *et al.*, *Tet. Lett.*, 1980, 2443 (abs config, synth)

Konda, Y. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 1083 (synth, ir, pmr)

Pellegata, R. *et al.*, *Tetrahedron*, 1985, **41**, 5607 (synth, pmr)


Seki, M. *et al.*, *Synthesis*, 1999, 745-747 (*S*-form, synth, ir, pmr, cmr)

Daoust, B. *et al.*, *Tetrahedron*, 1999, **55**, 3495-3514 (*N*-benzyl)

Tian, Z. *et al.*, *Org. Process Res. Dev.*, 2002, **6**, 416-418 (*N*-tert-butyloxycarbonyl, synth, pmr, cmr, ms)

5-Hydroxy-2-pyrrolidinone, 9CI H-719

5-Hydroxy-2-pyrrolidone, 8CI [62312-55-4]



(*R*)-form

C₄H₇NO₂ 101.105

Constit. of *Jatropha curcas*.

(*R*)-form [168111-92-0]

Constit. of *Hyptis verticillata*. Isol. from the mushroom *Macrolepiota neomastoides*. Amorph. powder. [α]_D²⁵ +5 (c, 0.05 in MeOH). λ_{max} 198 (log ε 1.55); 224 (sh) (log ε 0.61); 241 (sh) (log ε 0.55) (H₂O). λ_{max} 203 (log ε 5.33); 229 (log ε 5.04) (MeOH).

(±)-form

Cryst. Mp 98-99°.

Me ether: 5-Methoxy-2-pyrrolidinone

[63853-74-7]

C₅H₉NO₂ 115.132

Cryst. (EtOAc/hexane). Mp 59-60.5°.

(±)-form

Me ether: *Pterolactam*

[38072-88-7]

[63853-74-7]

C₅H₉NO₂ 115.132

Constit. of *Pteridium aquilinum* (bracken fern). Cryst. (petrol). Mp 56-67°. [α]_D²⁵ +2 (CHCl₃).

De Mayo, P. *et al.*, *Chem. Ind. (London)*, 1962, 1576 (synth)

Takatori, K. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 1087 (isol, *Me ether*)

Iwasaki, T. *et al.*, *J.O.C.*, 1979, **44**, 1552 (synth, *Me ether*)

Cue, B.W. *et al.*, *Org. Prep. Proced. Int.*, 1979, **11**, 285 (synth)

Barco, A. *et al.*, *Synthesis*, 1979, 68 (synth, ir, pmr)

Lundgren, D.W. *et al.*, *J. Biol. Chem.*, 1980, **255**, 4481 (ms)

Phan, X.T. *et al.*, *J.O.C.*, 1983, **48**, 5164 (synth, *Me ether*)

Farina, F. *et al.*, *Heterocycles*, 1984, **22**, 1733-1739 (synth)

Shono, T. *et al.*, *J.O.C.*, 1985, **50**, 3243 (synth, *Me ether*)


Kuhnt, M. *et al.*, *Planta Med.*, 1995, **61**, 227-232 (isol, uv, pmr, cmr, ms)

Staubmann, R. *et al.*, *Phytochemistry*, 1999, **50**, 337-338 (isol, pmr, cmr)

Kim, K.H. *et al.*, *Bull. Korean Chem. Soc.*, 2008, **29**, 1591-1593 (isol, cd, pmr, cmr)

6-Hydroxy-1-pyrrolizidine-carboxylic acid H-720

Hexahydro-6-hydroxy-1*H*-pyrrolizine-1-carboxylic acid



(1*R*,6*R*,7*aR*)-form

C₈H₁₃NO₃ 171.196

(1*R*,6*R*,7*aR*)-form

Cryst. (Me₂CO/MeOH) (as hydrochloride). Mp 147-148° (hydrochloride). [α]_D +32 (c, 1 in MeOH).

Me ester: *Amphorogynine D*

C₉H₁₅NO₃ 185.222

Alkaloid from *Amphorogynine spicata*. Amorph. gum. [α]_D +17 (c, 1 in CHCl₃).

Et ester: [73354-03-7]

C₁₀H₁₇NO₃ 199.249

Needles (EtOAc). Mp 110°. [α]_D¹⁸ +73.4 (c, 4 in CHCl₃).

6-*O*-[3-(4-Hydroxy-3-methoxyphenyl)propanol], *Me ester*: *Amphorogynine A*
C₁₉H₂₅NO₆ 363.41

Alkaloid from *Amphorogyne spicata*. Cryst. (MeOH/heptane). Mp 108°. [α]_D +53 (c, 1 in CHCl₃).

(1R,6S,7aR)-form

6-O-[3-(4-Hydroxy-3-methoxyphenyl)propanoyl], Me ester: **Amphorogynine C**

C₁₉H₂₅NO₆ 363.41

Alkaloid from *Amphorogyne spicata*. Cryst. (MeOH). Mp 130°. [α]_D -2 (c, 1 in CHCl₃).

(1S,6S,7aR)-form

6-O-[3-(4-Hydroxy-3-methoxyphenyl)propanoyl], Me ester: **Amphorogynine B**

Alkaloid from *Amphorogyne spicata*. Gum. [α]_D -7 (c, 1 in CHCl₃).

(1S,6S,7aS)-form

Et ester: [78175-02-7]

Needles (EtOAc). Mp 110°. [α]_D¹⁸ -72.1 (c, 4 in CHCl₃).

Robins, D.J. *et al.*, *J.C.S. Perkin 1*, 1989, 1339-1342 (*synth*, *ester*)


Huong, D.T.T. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1444-1446 (*isol*, *ir*, *pmr*, *cmr*)

Yoda, H. *et al.*, *Tet. Lett.*, 2003, **44**, 1643-1646 (*Amphorogynine A*, *synth*)

Roche, C. *et al.*, *J.O.C.*, 2005, **70**, 8352-8363 (*Amphorogynine A, D*, *synth*)

2-Hydroxy-4-quinolinecarboxaldehyde H-721

1,2-Dihydro-2-oxo-4-quinolinecarboxaldehyde. 4-Formyl-2-hydroxyquinoline. 4-Formylcarbostyryl



C₁₀H₇NO₂ 173.171

NH-form

N-Mercapto: 4-Formyl-1-mercapto-2(1H)-quinolinone. 4-Formyl-N-mercaptocarbostyryl

C₁₀H₇NO₂S 205.237

Prod. by *Pseudomonas fluorescens* strain G308. Antifungal agent. λ_{\max} 222 ; 280 ; 318 (no solvent reported).

Fakhouri, W. *et al.*, *Phytochemistry*, 2001, **58**, 1297-1303 (*N-mercapto*, *isol*, *pmr*, *cmr*, *ms*)

8-Hydroxy-4-quinolinecarboxaldehyde, 9CI H-722

8-Hydroxycinchonaldehyde, 8CI. 4-Formyl-8-hydroxyquinoline

[14510-07-7]

C₁₀H₇NO₂ 173.171

Alkaloid from *Broussonetia zeylanica*. Orange cryst. (EtOH aq.). Mp 144-145°. λ_{\max} 244 ; 263 ; 325 (no solvent reported).

Oxime: [71294-68-3]

C₁₀H₈N₂O₂ 188.185

Alkaloid from *Broussonetia zeylanica*. Cryst. Subl. 222. Natural product originally assigned a bipyridine struct.

Me ether: 8-Methoxy-4-quinolinecarboxaldehyde. 4-Formyl-8-methoxyquinoline

C₁₁H₉NO₂ 187.198

Mp 109-110°.

Buechi, J. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1676-1683 (*synth*)

Fujikawa, F. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 844-849 (*synth*)


Gunatilaka, A.A.L. *et al.*, *J. Chem. Res., Synop.*, 1979, 61 (*isol*)

Dehmlow, E.V. *et al.*, *Phytochemistry*, 1990, **29**, 3993-3995 (*oxime*)

2-Hydroxy-4-quinolinecarboxylic acid H-723

1,2-Dihydro-2-oxo-4-quinolinecarboxylic acid. 2(1H)-Quinolinone-4-carboxylic acid. 2-Hydroxycinchoninic acid. Carbostyryl-4-carboxylic acid

[84906-81-0]



C₁₀H₇NO₃ 189.17

NH-form is major tautomer. Isol. from poppy straw (*Papaver somniferum*). Needles (H₂O). Mp 343° (anhyd.) Mp 310° (monohydrate).

Me ester:

C₁₁H₉NO₃ 203.197

Prod. by a marine bacterium strain He159b. Needles (H₂O). Mp 248-251°.

Et ester: 4-Ethoxycarbonyl-2(1H)-quinolinone

[5466-27-3]

C₁₂H₁₁NO₃ 217.224

Alkaloid from the fruit of *Brucea javanica* and *Parthenocissus tricuspidata*. Needles (EtOH aq.). Mp 207-210°.

NH-form [15733-89-8]

N-Me:

C₁₁H₉NO₃ 203.197

Mp 244°.

OH-form

Me ether: 2-Methoxy-4-quinolinecarboxylic acid

[10222-62-5]

C₁₁H₉NO₃ 203.197

Cryst. (MeOH). Mp 175°.

Et ether: 2-Ethoxy-4-quinolinecarboxylic acid

C₁₂H₁₁NO₃ 217.224

Needles (H₂O). Mp 145-146°.

Et ether, Et ester:

C₁₄H₁₅NO₃ 245.277

Needles. Mp 86°.

Wojahn, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1931, **269**, 422 (*synth*)

Sugasawa, S. *et al.*, *Yakugaku Zasshi*, 1937, **57**, 296 (*synth*)

Schmid, H. *et al.*, *Helv. Chim. Acta*, 1945, **28**, 722 (*isol*)

Org. Synth., Coll. Vol., 3, 1955, 456 (*synth*)

Kadaba, P.K. *et al.*, *Synthesis*, 1972, 628 (*deriv*)

Sawayama, T. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 2305 (*synth*)

Wang, Y.Z. *et al.*, *Yaoxue Xuebao*, 1982, **17**, 466; *CA*, **97**, 107056 (*isol*, *deriv*)

Mirek, J. *et al.*, *Z. Naturforsch., A*, 1982, **37**, 1276 (*tautom*, *uv*)


Yu, Y.N. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 382; *CA*, **113**, 227957 (*isol*, *deriv*)

Shabaan, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (*Me ester*, *marine*, *isol*)

3-Hydroxy-2-quinolinecarboxylic acid, 9CI H-724

3-Hydroxyquinaldic acid, 8CI

[15462-45-0]



C₁₀H₇NO₃ 189.17

Isol. from *Streptomyces griseoflavus*.

Yellow needles (Me₂CO). Mp 200° dec.

(196-198°). λ_{\max} 237 (€ 11000); 328 (€ 2400); 369 (€ 2200) (MeOH/HCl) (Derep).

λ_{\max} 228 (€ 14500); 286 (€ 1800); 353 (€ 2100) (MeOH/NaOH)

(Derep). λ_{\max} 217 (€ 16300); 228 (€ 16100); 291 (€ 1500); 352 (€ 2100) (MeOH) (Derep).

Ca salt: [110429-27-1]

Sol. MeOH; poorly sol. H₂O. Mp 170°. λ_{\max} 217 (€ 16300); 228 (€ 16100); 292 (€ 1500); 352 (€ 2100)

(MeOH) (Berdy). λ_{\max} 237 (€ 11000); 328 (€ 2400); 369 (€ 2200) (MeOH-HCl) (Berdy). λ_{\max} 228 (€ 14500); 286 (€ 1800); 353 (€ 2100) (MeOH-NAOH) (Berdy).

Me ester: [171917-61-6]

C₁₁H₉NO₃ 203.197

Solid. Mp 122-124°.

Amide: [15462-44-9]

C₁₀H₈N₂O₂ 188.185

Pale-yellow needles (Me₂CO). Mp 215-216°.

Nitrile: 2-Cyano-3-hydroxyquinoline

[15462-43-8]

C₁₀H₆N₂O 170.17

Needles (MeOH). Mp 183° Mp 300° dec. No explanation given for widely differing Mp's.

Me ether, Me ester: [110429-26-0]

C₁₂H₁₁NO₃ 217.224

Mp 78-80°.

Benzyl ether: [110429-24-8]

C₁₇H₁₃NO₃ 279.295

Needles (MeOH). Mp 150-151° dec.

Benzyl ether, Me ester:

C₁₈H₁₅NO₃ 293.321

Oil.

Kaneko, C. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 663 (*synth*)

Breiding-Mack, S. *et al.*, *J. Antibiot.*, 1987, **40**, 953 (*isol*, *props*)

Boger, D.L. *et al.*, *J.O.C.*, 1995, **60**, 7369

(*derivs*)

Bergeron, R.J. *et al.*, *J. Med. Chem.*, 1999, **42**, 95-108 (*nitrile*)

4-Hydroxy-2-quinolinecarboxylic acid H-725

1,4-Dihydro-4-oxo-2-quinolinecarboxylic acid, 9CI. 4-(1H)-Quinolinone-2-carboxylic acid. 4-Hydroxyquinaldinic acid. *Kynurenic acid*. *Transitorine*. *Transitorine* [492-27-3]

C₁₀H₇NO₃ 189.17

The oxo form is the major tautomer. A normal prod. of mammalian tryptophan metab. Isol. from urine of dogs (Liebig, 1953). Alkaloid from *Ephedra transitoria* (Ephedraceae). Shows anticonvulsant props. Needles. Ppt. from soln. of Na salt by HCl. Insol. H₂O, Et₂O; sol. hot EtOH. Mp 123° (nat. prod.) Mp 287-288° dec.

Me ester:

C₁₁H₉NO₃ 203.197
Mp 224°.

Et ester: [13720-90-6]

C₁₂H₁₁NO₃ 217.224
Mp 215° (213°).

OH-form

Me ether: 4-Methoxy-2-quinolinecarboxylic acid [15733-83-2]

C₁₁H₉NO₃ 203.197
Needles (H₂O). Mp 196-197° dec.

Me ether, Me ester: [67976-94-7]

C₁₂H₁₁NO₃ 217.224
Mp 148-149°.

Me ether, Et ester:

C₁₃H₁₃NO₃ 231.251
Prisms (C₆H₆/petrol). Mp 131-133°.

Oxo-form

N-Me, Et ester: [60442-66-2]

C₁₃H₁₃NO₃ 231.251
Mp 110-112°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 448C (nmr)

Besthorn, E. et al., *Ber.*, 1921, **54**, 1330 (synth)

Riegel, B. et al., *J.A.C.S.*, 1946, **68**, 2685 (ester)

Wald, D.K. et al., *J.O.C.*, 1966, **31**, 3369 (synth, ir)

Narumiya, S. et al., *J. Biol. Chem.*, 1979, **254**, 7007 (metab)

Okabe, N. et al., *Acta Cryst. C*, 1996, **52**, 1827 (cryst struct)

Al-Khalil, S. et al., *J. Nat. Prod.*, 1998, **61**, 262-263 (isol, *Ephedra*)

Michael, J.P. et al., *Nat. Prod. Rep.*, 2000, **17**, 603-620 (*Transitorine*)

6-Hydroxy-4-quinolinecarboxylic acid H-726

6-Hydroxycinchoninic acid. *Xanthoquininic acid*. *Xanthoquinic acid* [4312-44-1]

C₁₀H₇NO₃ 189.17

Plates (H₂O). Mp 325° dec.

Methiodide:

Orange-yellow cryst. (EtOH). Mp 302°.

Et ester: [31610-09-0]

C₁₂H₁₁NO₃ 217.224

Mp 185.5°.

Amide: 6-Hydroxy-4-quinolinecarboxamide. *Xanthoquininamide*

[51094-27-0]

C₁₀H₈N₂O₂ 188.185

Alkaloid from the roots of *Microdesmis keayana*. Pale yellow powder. λ_{max} 205 (log ε 4.17); 232 (log ε 3.94); 270 (log ε 3.91); 308 (log ε 3.23) (MeOH).

Me ether: 6-Methoxy-4-quinolinecarboxylic acid. *Quininic acid*

[86-68-0]

C₁₁H₉NO₃ 203.197

Used as a satd. aq. soln. as acid-base fluorescent indicator (pH range: 4.0-5.0; colour change: yellow → blue). Yellow prisms (dil. HCl). Spar. sol. H₂O; sol. acids, alkalis. Mp 285° dec. pK_a 3.05. Yellow colour in acid soln. Blue fluor. in EtOH destroyed by H₂O or acids.

Me ether, Me ester: [19834-77-6]

C₁₂H₁₁NO₃ 217.224
Mp 84°.

Me ether, Et ester: [5345-57-3]

C₁₃H₁₃NO₃ 231.251
Insol. H₂O. Mp 69°.

Me ether, nitrile: 4-Cyano-6-methoxyquinoline

[6443-89-6]

C₁₁H₈N₂O 184.197

Yellow needles (C₆H₆). Mp 157°.

Et ether: 6-Ethoxy-4-quinolinecarboxylic acid. *Optoquinic acid*

[525-39-3]

C₁₂H₁₁NO₃ 217.224

Yellow needles (propanol). Mp 278°.

John, H. et al., *Ber.*, 1930, **63**, 2657-2661 (*Me ether*)

John, H. et al., *J. Prakt. Chem.*, 1930, **128**, 194 (synth)

Rabe, P. et al., *Ber.*, 1931, **64**, 2487-2500 (*Me ether*)

Thielpape, E. et al., *Ber.*, 1939, **72B**, 1432-1443 (*Me ether*)

King, H. et al., *J.C.S.*, 1942, 401-404 (*Me ether*)

Campbell, K.N. et al., *J.O.C.*, 1946, **11**, 803-811 (*Me ether Et ester*)

Benassi, C.A. et al., *Gazz. Chim. Ital.*, 1967, **97**, 3 (synth, ir)

Schultz, O.-E. et al., *Annalen*, 1970, **740**, 192-195 (*Me ether*)

Bishop, E. et al., *Indicators*, Pergamon, Oxford, 1972, (use, *Me ether*)

Zamble, A. et al., *Chem. Pharm. Bull.*, 2007, **55**, 643-645 (*Xanthoquininamide*)

8-Hydroxy-2-quinolinecarboxylic acid H-727

[1571-30-8]

C₁₀H₇NO₃ 189.17

Cryst. Mp 210°.

► UZ9450000

Diethylamide:

C₁₄H₁₆N₂O₂ 244.293
Solid. Mp 104°.

Me ether: 8-Methoxy-2-quinolinecarboxylic acid

[21638-90-4]

C₁₁H₉NO₃ 203.197

Isol. from scent gland of beetle *Ilybius fenestratus*. Yellow needles. Mp 75-75.5°.

Benzyl ether: [176696-00-7]

C₁₇H₁₃NO₃ 279.295
Mp 109°.

Tamada, T. et al., *CA*, 1969, **70**, 113301n (tox) Schildknecht, H. et al., *Z. Naturforsch., B*, 1969, **24**, 38 (isol, synth)


Caris, C. et al., *Tetrahedron*, 1996, **52**, 4659-4672 (*benzyl ether, synth, pmr, cmr*)

Albrecht, M. et al., *Eur. J. Org. Chem.*, 2007, 4902-4908 (*diethylamide*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HOE000

4-Hydroxy-2,3-quinolinedicarboxylic acid H-728

1,4-Dihydro-4-oxo-2,3-quinolinedicarboxylic acid, 9CI



C₁₁H₇NO₅ 233.18

NH-form

N-Me, di-Me ester: 2,3-Dicarbomethoxy-1-methyl-4(1H)-quinolinone

[181289-63-4]

C₁₄H₁₃NO₅ 275.26

Alkaloid from leaves of *Sarcomelicope dogniensis*. Amorph. λ_{max} 223 (log ε 3.97); 247 (log ε 3.87); 321 (log ε 3.74); 337 (log ε 3.79) (MeOH).

N-Me, di-Et ester: [73281-87-5]

Mp 96-98°.

N-Me, 3-amide: 3-(Aminocarbonyl)-1,4-dihydro-1-methyl-4-oxo-2-quinolinecarboxylic acid. *Quinoloninic acid*

[888728-71-0]

C₁₂H₁₀N₂O₄ 246.222

Prod. by *Penicillium citrinum* (MST-F10130). solid. λ_{max} 221 (ε 9300); 250 (ε 8600); 258 (ε 8500); 304 (sh) (ε 3500); 316 (ε 4000) (MeOH).

N-Me, imide: 4-Methyl-1H-pyrrolo[3,4-b]quinoline-1,3,9(2H,4H)-trione. *Quinolonimide*

[888728-70-9]

C₁₂H₈N₂O₃ 228.207

Prod. by *Penicillium citrinum* (MST-F10130). Pale yellow solid. λ_{max} 224 (ε 12500); 251 (sh) (ε 8700); 269 (sh) (ε 5600); 313 (ε 3800); 325 (ε 4300) (MeOH).

Coppola, G.M. et al., *J. Het. Chem.*, 1979, **16**, 1605 (synth, di-Et ester)


Coppola, G.M. et al., *Org. Magn. Reson.*, 1981, **17**, 242 (cmr, di-Et ester)

Mitaku, S. et al., *Nat. Prod. Lett.*, 1995, **7**, 219 (isol, di-Me ester)

Clark, B. et al., *Org. Biomol. Chem.*, 2006, **4**, 1512-1519 (*Quinoloninic acid, Quinolonimide*)

5-Hydroxy-1-β-D-ribofuranosyl-1*H*-imidazole-4-carboxaldehyde, 9CI

H-729

F 2787. Antibiotic F 2787
[143577-73-5]C₉H₁₂N₂O₆ 244.204

Nucleoside-type antibiotic. Prod. by *Streptomyces filipinensis-artispiralis*. Plant growth regulator and herbicide. Powder. Sol. H₂O; fairly sol. MeOH; poorly sol. butanol, Me₂CO, hexane. [α]_D -43 (H₂O). λ_{max} 215 (E1%/1cm 320); 240 (E1%/1cm 160); 305 (ε 695) (H₂O) (Berdy). λ_{max} 210 (E1%/1cm 210); 245 (E1%/1cm 150) (HCl) (Berdy).


▶ LD₅₀ (mus, ipr) 500 mg/kg.

Japan. Pat., 1992, 92 112 896; CA, 117, 145326e

N-(6-Hydroxy-9-β-D-ribofuranosyl-9*H*-purin-2-yl)alanine

H-730

Guanosinepropionic acid

C₁₃H₁₇N₅O₇ 355.307


Isol. from a *Fusarium* sp. Cryst. (MeOH/Me₂CO). [α]_D²⁵ +9.9 (c, 1.0 in H₂O).

Gerster, J.F. et al., J.A.C.S., 1965, 87, 3752 (synth, uv)

Ballio, A. et al., Gazz. Chim. Ital., 1966, 96, 337 (isol)

11-Hydroxysaxitoxin

H-731



11α-form

C₁₀H₁₇N₇O₅ 315.288**11α-form [78780-57-1]**Isol. from *Gonyaulax tamarensis*.11-O-Sulfate: **Gonyautoxin II. GTX2**

[60508-89-6]

C₁₀H₁₇N₇O₈S 395.352

From *Gonyaulax* and *Protogonyaulax* spp. and present in other marine organisms. Neurotoxin. Sol. H₂O.

▶ Toxic.

11-O-Sulfate, N¹⁴-sulfonic acid: **Protogonyautoxin I. Toxin C₁. Toxin PX₁**

[80173-30-4]

C₁₀H₁₇N₇O₁₁S₂ 475.417

From *Gonyaulax* and *Protogonyaulax* spp. Neurotoxin. Prisms + H₂O (MeOH aq.).

▶ Toxic.

Decarbamoyl: **Decarbamoyl-11α-hydroxysaxitoxin**C₉H₁₆N₆O₄ 272.263

Isol. from shellfish. No CAS no. 8-14 CI.

Decarbamoyl, 11-O-sulfate: **Decarbamoylgonyautoxin II**

[86996-87-4]

C₉H₁₆N₆O₇S 352.327

Isol. from various shellfish.

Decarbamoyl, O¹³-Ac, 11-O-sulfate: **13-Acetyldecarbamoylgonyautoxin II. LWTX3**

[200816-97-3]

C₁₁H₁₈N₆O₈S 394.365

Isol. from *Lyngbya wollei*.

Decarbamoyl, 13-O-(4-hydroxybenzoyl), 11-O-sulfate: **GC 1**

[603125-80-0]

C₁₆H₂₀N₆O₉S 472.435

Isol. from *Gymnodinium catenatum*.

De(carbamoyloxy), 11-O-sulfate: **Decarbamoyloxygonyautoxin II**

[186249-38-7]

C₉H₁₆N₆O₆S 336.328

Isol. from *Alexandrium* sp.

N¹-Hydroxy: **11α-Hydroxyneosaxitoxin**C₁₀H₁₇N₇O₆ 331.288

Isol. from shellfish. No CAS no found 8-14CI.

N¹-Hydroxy, 11-O-sulfate: **Gonyautoxin I. GTX1**

[60748-39-2]

C₁₀H₁₇N₇O₉S 411.352

Prod. by *Gonyaulax* and *Protogonyaulax* spp. and isol. from shellfish and other marine organisms. Neurotoxin, causal agent, with Saxitoxin, of shellfish poisoning. Sol. H₂O. The Gonyautoxins, esp. the N-sulfates, show reduced toxicity compared with Saxitoxin.

▶ Toxic.

N¹-Hydroxy, 11-O-sulfate, N¹⁴-sulfonic acid: **Protogonyautoxin 3. Toxin C₃. Toxin PX₃**

[89614-45-9]

C₁₀H₁₇N₇O₁₂S₂ 491.416

From *Gonyaulax* and *Protogonyaulax* spp. Neurotoxin.

▶ Toxic.

N¹-Hydroxy, decarbamoyl: **Decarbamoyl-11α-hydroxyneosaxitoxin**C₉H₁₆N₆O₅ 288.263

No CAS no. found 8-14CI.

N¹-Hydroxy, decarbamoyl, 11-O-sulfate:**Decarbamoylgonyautoxin I**

[122075-86-9]

C₉H₁₆N₆O₈S 368.327

Isol. from *Gymnodinium catenatum* and *Alexandrium* sp.

11β-form [99685-70-8]

Isol. from mussels (*Mytilus* sp.) during an intense bloom of *Alexandrium tamarensis*.

11-O-Sulfate: **Gonyautoxin III. GTX3**

[60537-65-7]

C₁₀H₁₇N₇O₈S 395.352

From *Gonyaulax* and *Protogonyaulax* spp. and other marine organisms. Neurotoxin. Sol. H₂O.

▶ Toxic.

11-O-Sulfate, N¹⁴-sulfonic acid: **Gonyautoxin VIII. GTX8. Protogonyautoxin 2. Toxin C₂. Toxin PX₂**

[80226-62-6]

C₁₀H₁₇N₇O₁₁S₂ 475.417

From *Gonyaulax* and *Protogonyaulax* spp. Neurotoxin. Large cryst. + H₂O (H₂O).

▶ Toxic.

UU6858700

Decarbamoyl: **Decarbamoyl-11β-hydroxysaxitoxin**C₉H₁₆N₆O₄ 272.263

Isol. from shellfish. No CAS no found 8-14CI.

Decarbamoyl, 11-O-sulfate: **Decarbamoylgonyautoxin III**

[87038-53-7]

C₉H₁₆N₆O₇S 352.327

Isol. from various shellfish.

Decarbamoyl, O¹³-Ac, 11-O-sulfate: **13-Acetyldecarbamoylgonyautoxin III. LWTX2**

[200816-96-2]

C₁₁H₁₈N₆O₈S 394.365

Isol. from *Lyngbya wollei*.

Decarbamoyl, 13-O-(4-hydroxybenzoyl), 11-O-sulfate: **GC 2**

[603125-81-1]

C₁₆H₂₀N₆O₉S 472.435

Isol. from *Gymnodinium catenatum*.

De(carbamoyloxy), 11-O-sulfate: **Decarbamoyloxygonyautoxin III**

[155666-11-8]

C₉H₁₆N₆O₆S 336.328

Isol. from *Gymnodinium catenatum* and *Alexandrium* sp.

N¹-Hydroxy: **11β-Hydroxyneosaxitoxin**C₁₀H₁₇N₇O₆ 331.288

Isol. from shellfish. No CAS no found 8-14CI.

N¹-Hydroxy, 11-O-sulfate: **Gonyautoxin IV. GTX4**

[64296-26-0]

C₁₀H₁₇N₇O₉S 411.352

From *Gonyaulax* and *Protogonyaulax* spp. and other marine organisms. Neurotoxin. Sol. H₂O.

▶ Toxic.

N¹-Hydroxy, 11-O-sulfate, N¹⁴-sulfonic acid: **Protogonyautoxin 4. Toxin C₄. Toxin PX₄**

[89674-98-6]

C₁₀H₁₇N₇O₁₂S₂ 491.416

From *Gonyaulax* and *Protogonyaulax* spp. Neurotoxin.

▶ Toxic.

N¹-Hydroxy, decarbamoyl: Decarbamoyl-11β-hydroxyneosaxitoxinC₉H₁₆N₆O₅ 288.263

Isol. from shellfish.

N¹-Hydroxy, decarbamoyl, 11-O-sulfate: Decarbamoylgonyautoxin IV


[122169-51-1]

C₉H₁₆N₆O₈S 368.327Isol. from *Gymnodinium catenatum* and *Alexandrium* sp.**12-Deoxy, decarbamoyl, O¹³-Ac, 11-O-sulfate: LWTX1**

[200816-95-1]


C₁₁H₁₈N₆O₇S 378.365Isol. from *Lyngbya wollei*.Onoue, Y. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 420-430 (props)Oshima, Y. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 1707-1711 (chromatog, bibl)Maruyama, J. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 2783-2788 (ms)Hall, S. *et al.*, *Tet. Lett.*, 1984, **25**, 3537-3538 (Toxins C₃, C₄)Shimizu, Y. *et al.*, *Tetrahedron*, 1984, **40**, 539-544 (props)Anderson, D.M. *et al.*, *Toxicol.*, 1989, **27**, 665-674 (Decarbamoylgonyautoxins)Hall, S. *et al.*, *ACS Symp. Ser.*, 1990, 418 (book)Gallacher, S. *et al.*, *Appl. Environ. Microbiol.*, 1997, **63**, 239-245 (Decarbamoyloxygonyautoxins)Onodera, H. *et al.*, *Nat. Toxins*, 1997, **5**, 146-151 (LWTX toxins)Food Sci. Technol., *Seafood and Freshwater Toxins*, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (revs)Negri, A. *et al.*, *Chem. Res. Toxicol.*, 2003, **16**, 1029-1033 (GCI, GC2)Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2004, 2533-2551 (rev)Llewellyn, L.E. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 200-222 (rev)Dell'Aversano, C. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1518-1523 (isol, pmr, cmr, ms)Mulcahy, J.V. *et al.*, *J.A.C.S.*, 2008, **130**, 12630-12631 (Gonyautoxin III, synth)**3-Hydroxy-3,4-secocoronaridine H-732**

[131653-78-6]

C₂₁H₂₈N₂O₃ 356.464Alkaloid from the leaves and stem bark of *Ervatamia polyneura* (Apocynaceae). [α]_D²⁵ -46 (c, 0.5 in CHCl₃).Clivio, P. *et al.*, *Phytochemistry*, 1990, **29**, 3007 (isol, uv, ir, pmr, cmr, ms struct)**4a-Hydroxyskytanthine H-733****Octahydro-2,4,7-trimethyl-1H-cyclopenta[c]pyridin-4a-ol, 9CI. Isooxyskytanthine**

[22324-99-8]

[947688-44-0 (monohydrochloride)]




Absolute configuration

C₁₁H₂₁NO 183.293Alkaloid from *Tecoma stans*. Mp 91-92°.

Picrate: Mp 170-170.5°.

Dickinson, E.M. *et al.*, *Tetrahedron*, 1969, **25**, 1523 (ir, pmr, isol)Jones, G. *et al.*, *Chem. Comm.*, 1971, 994 (cryst struct)Ferguson, G. *et al.*, *J.C.S. Perkin 2*, 1975, 1124 (cryst struct)Berg, W. *et al.*, *Pharmazie*, 1977, **32**, 41 (ms)Cossy, J. *et al.*, *J.O.C.*, 1993, **58**, 2351 (synth, pmr, cmr, ms)Marzouk, M. *et al.*, *Z. Naturforsch.*, C, 2006, **61**, 783-791 (isol, pmr, cmr)**7a-Hydroxyskytanthine H-734****Octahydro-2,4,7-trimethyl-1H-cyclopenta[c]pyridin-7a-ol, 9CI**

[22325-03-7]



(4R,4aS*,7R*,7aR*)-form


C₁₁H₂₁NO 183.293**(4R,4aS,7R,7aR)-form**N-Oxide: **Kinabalarine G**C₁₁H₂₁NO₂ 199.292Alkaloid from *Kopsia dasyrachis*.**(4ξ,4αξ,7ξ,7αξ)-form**Alkaloid from *Tecoma stans*. Mp 87-89°.

Methiodide:



Cryst. (EtOH). Mp 310-312° dec.

Dickinson, E.M. *et al.*, *Tetrahedron*, 1969, **25**, 1523-1529 (isol, ir, pmr, ms)Kam, T.S. *et al.*, *Phytochemistry*, 1999, **52**, 959-963 (Kinabalarine G)**4-Hydroxyskytanthine H-735****Hydroxyskytanthine I**


[16625-64-2]

C₁₁H₂₁NO 183.293Minor alkaloid from the leaves of *Skytanthus acutus* (Apocynaceae). Cryst. (cyclohexane or by subl.). Mp 94-95°.[α]_D¹⁹ +38.5 (c, 1.50 in cyclohexane). [α]_D¹⁹ +35.8 (c, 1.80 in MeOH).Appel, H.H. *et al.*, *Scientia (Valparaiso)*, 1961, **28**, 5; *CA*, **57**, 2332g (isol)Casinovi, C.G. *et al.*, *Chem. Ind. (London)*, 1963, 984 (ir, pmr)Adolphsen, G. *et al.*, *Tetrahedron*, 1967, **23**, 3147 (isol, pmr, ms, struct)**7-Hydroxyskytanthine H-736****Hydroxyskytanthine II**

[16625-65-3]

C₁₁H₂₁NO 183.293Minor alkaloid from the leaves of *Skytanthus acutus* (Apocynaceae).Cryst. (cyclohexane or by subl.). Mp 119-120°. [α]_D¹⁹ -38.5 (c, 1.00 in MeOH).Adolphsen, G. *et al.*, *Tetrahedron*, 1967, **23**, 3147 (isol, pmr, ms, struct)**4-Hydroxysparteine H-737**C₁₅H₂₆N₂O 250.383**4α-form**Alkaloid from bark of *Acosmium panamense* (Fabaceae). Oil. [α]_D²³ -31. [α]₃₆₅²³ -103 (c, 0.1 in EtOH).Balandrin, M.F. *et al.*, *Heterocycles*, 1982, **19**, 1931 (isol, struct, ms, synth)**7-Hydroxysparteine H-738**

[36565-37-4]



Relative configuration

C₁₅H₂₆N₂O 250.383Alkaloid in *Cytisus hirsutus* and *Chamaecytisus ciliatus* (Fabaceae).**6-Epimer: 7-Hydroxy-β-isosparteine**C₁₅H₂₆N₂O 250.383Alkaloid from *Lupinus sericeus* (Fabaceae). Cryst. solid by subl. Mp 103.5-104.5°. [α]_D²⁵ -8 (c, 3.0 in EtOH). pK_a 10.7 (50% EtOH aq.). Crystallises with difficulty.


6-Epimer, perchlorate: Mp 207.5-208.5°.

6-Epimer, Ac:Noncryst. Mp 231-236.5° (as perchlorate). [α]_D²⁵ +23.5 (c, 0.66 in EtOH).Pinkerton, J.M.H. *et al.*, *J.O.C.*, 1967, **32**, 1828 (cryst struct, epimer)Carmack, M. *et al.*, *J.O.C.*, 1967, **32**, 3045 (isol, struct, epimer)Neuner-Jehle, N. *et al.*, *Monatsh. Chem.*, 1967, **98**, 836 (ms)

Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1971, **24**, 1657; *CA*, **76**, 138214y (isol)
Daily, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 889 (occur)

14-Hydroxysparteine

H-739

C₁₅H₂₆N₂O 250.383**14 α -form**

Alkaloid from *Laburnum watereri*. Characterised by ms only.


Bohlmann, F. *et al.*, *Chem. Ber.*, 1965, **98**, 653 (synth)

Greinwald, R. *et al.*, *Biochem. Physiol. Pflanz.*, 1990, **186**, 1 (occur)

3-Hydroxystachydrine

H-740

2-Carboxy-3-hydroxy-1,1-dimethylpyrrolidinium hydroxide inner salt, 9CI. 3-Hydroxyproline dimethylbetaine [51795-35-8]



(2R,3S)-form

C₇H₁₃NO₃ 159.185

Alkaloid in *Boscia*, *Cadaba*, *Capparis*, *Maerua*, *Morisonia*, *Neothorelia*, *Poitanedora*, *Steriphoma*, *Thilachium* and *Tirania* spp. (no stereochem. detd.). Characteristic of the Cappariaceae.

(2R,3S)-form

D-cis-form. 3-Hydroxystachydrine *b* [32213-41-5]

Alkaloid from *Courbonia virgata* (preferred genus name *Maerua*). Cryst. + 1H₂O (EtOH aq.). Mp 209-210° dec. [α]_D²² +53 (c, 2.5 in H₂O).

Hydrochloride:

Prisms (EtOH). Mp 201-202°.

(2S,3S)-form

L-trans-form. 3-Hydroxystachydrine *a*

Alkaloid from *Courbonia virgata*. Prisms. Mp 250° (efferv.) (245°). [α]_D³⁰ +10 (c, 29 in H₂O).

Hydrochloride:

Needles (EtOH). Mp 196-197° dec.

Picrate:

Yellow cryst. (EtOH). Mp 160°.

Cornforth, J.W. *et al.*, *J.C.S.*, 1952, 597 (isol, struct)

Sakiyama, F. *et al.*, *J.A.C.S.*, 1964, **86**, 1842 (synth, config)

Mandava, N. *et al.*, *Annalen*, 1970, **741**, 167 (pmr, config)


Delaveau, P. *et al.*, *Phytochemistry*, 1973, **12**, 2893 (occur)

Yousif, G. *et al.*, *Fitoterapia*, 1984, **55**, 117; *CA*, **102**, 21169b (occur)

8-Hydroxystreptazalone

H-741

4-Acetyl-2a,3,5,6,7,7b-hexahydro-3,5-dihydroxy-1H-2-oxa-7a-azacyclopent[*c*-*d*]inden-1-one [479581-78-7]



Relative Configuration

C₁₁H₁₃NO₅ 239.227

Prod. by *Streptomyces* sp.


Nomura, I. *et al.*, *Org. Lett.*, 2002, **4**, 4301-4304 (synth)

Nomura, I. *et al.*, *J.O.C.*, 2004, **69**, 1803-1812 (synth)

(4-Hydroxystyryl)trimethylammonium(1+)

H-742

2-(4-Hydroxyphenyl)-N,N,N-trimethylethenaminium, 9CI

C₁₁H₁₆NO⁺ 178.253**(E)-form** [778550-97-3]

[162827-15-8 (chloride)]


Metab. of the Palauan sponge *Axinyssa aplysinoides*. Highly hygroscopic solid (as chloride).

Compagnone, R.S. *et al.*, *J. Nat. Prod.*, 1995, **58**, 145-148 (isol, uv, ir, pmr, cmr)

Sumaryono, W. *et al.*, *CA*, 2006, **145**, 502464h (isol)

19-Hydroxytabersonine

H-743



(19R)-form

C₂₁H₂₄N₂O₃ 352.432**(19R)-form** [59086-85-0]

Alkaloid from *Catharanthus ovalis* and *Melodinus celastroides* (Apocynaceae).

[α]_D²² -365 (c, 0.53 in CHCl₃). λ _{max} 220 (ε 10500); 300 (ε 10500); 330 (ε 13500) (EtOH).

14 β ,15 β -Epoxide: **(19R)-Epimisiline**. 19-Hydroxypachysiphine [109063-93-6]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the leaves of *Petchia ceylanica* (Apocynaceae). Mp 252° dec.

[α]_D²⁷ -382 (c, 0.3 in CHCl₃).

14,15-Dihydro: see Minovincine, M-631

11-Hydroxy: **11,19R-Dihydroxytabersonine**

[138615-17-5]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Melodinus fusiformis* (Apocynaceae). Shows significant spermatocidal activity.

11-Methoxy: **Vandrikidine**. 20-Hydroxy-16-methoxytabersonine

[50656-92-3]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Craspidospermum verticillatum* (Apocynaceae). No phys. props. reported. λ _{max} 240 ; 328 (MeOH).

(19S)-form

14,15-Dehydro-19-epiminovincinine. 6,7-Dehydro-20-epiminovincinine

[59086-86-1]

Alkaloid from *Catharanthus ovalis*, *Catharanthus lanceus* and *Melodinus celastroides* (Apocynaceae). Cryst. (MeOH). Mp 163°. [α]_D²² -375 (c, 0.49 in CHCl₃). λ _{max} 220 (ε 13000); 300 (ε 10000); 330 (ε 14000) (EtOH).

14 α ,15 α -Epoxide: **Cathovalinine**

[61825-77-2]

C₂₁H₂₄N₂O₄ 368.432

Minor alkaloid from *Catharanthus ovalis* (Apocynaceae). Cryst. (C₆H₆). Mp 223°. [α]_D -492 (c, 0.30 in CHCl₃).

14 β ,15 β -Epoxide: **(19S)-Epimisiline**

[109063-94-7]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Petchia ceylanica* (Apocynaceae). Mp 198° dec. [α]_D²⁷ -399 (c, 0.2 in CHCl₃).

(19 ξ)-form [80286-66-4]

14 β ,15 β -Epoxide: **Demethoxyhoerhammerinine**. **Hoerhammericine**. **Hoerhammericine**

[19459-04-2]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Catharanthus trichophyllus* (Apocynaceae). Shows antitumour activity. Needles (C₆H₆). Mp 140-144° (138-140°). [α]_D²⁵ -40.3 (c, 2.04 in CHCl₃). [α]_D²⁵ -395 (c, 0.1 in CHCl₃). λ _{max} 299 (log ε 4.14); 328 (log ε 4.32) (MeOH) (Berdy).

11-Methoxy, 14 β ,15 β -epoxide: **Hoerhammerinine**. **Hoerhammerinine**

[19459-05-3]

Alkaloid from *Catharanthus lanceus* (Apocynaceae). Needles (EtOH). Mp 209.5-211° dec. [α]_D²⁵ -381 (c, 1 in CHCl₃). λ _{max} 242 ; 300 ; 325 (MeOH).

Blomster, R.N. *et al.*, *Naturwissenschaften*, 1968, **55**, 298 (Hoerhammericine)

Farnsworth, N.R. *et al.*, *Z. Naturforsch., B.*, 1968, **23**, 1061-1063 (Hoerhammerinine)

Abraham, D.J. *et al.*, *J.O.C.*, 1969, **34**, 1575 (Hoerhammericine)


Gabetta, B. *et al.*, *Fitoterapia*, 1976, **47**, 6-7; *CA*, **85**, 177738c (19S-Hydroxytabersonine)

Cordell, G.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 366-369 (Hoerhammericine)

Chiaroni, A. *et al.*, *Tetrahedron*, 1976, **32**, 1899 (Cathovalinine, isol, uv, ir, pmr, ms, cryst struct)

- Gabetta, B. *et al.*, *Fitoterapia*, 1979, **35**, 1151 (14,15-Dehydro-19-epiminovicinine)
- Langlois, N. *et al.*, *J.O.C.*, 1979, **44**, 2468-2471 (19R-Hydroxytabersonine, 19S-Hydroxytabersonine)
- Langlois, N. *et al.*, *Phytochemistry*, 1979, **18**, 467-471 (19R-Hydroxytabersonine, 19S-Hydroxytabersonine)
- Lounasmaa, M. *et al.*, *Acta Chem. Scand., Ser. B*, 1980, **34**, 379-381 (Vandrikidine, *pmr*, *config*)
- Kurz, W.G.W. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 1891-1896 (Vandrikidine, *biosynth*)
- Kutney, J.P. *et al.*, *Phytochemistry*, 1980, **19**, 2589-2595 (Vandrikidine, *biosynth*, *uv*, *pmr*, *ms*)
- Caron, C. *et al.*, *Heterocycles*, 1981, **16**, 645-646 (19R-Hydroxytabersonine)
- Chen, W. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 906-912; 1986, **21**, 187-190; *CA*, **105**, 3520s; 3558k (19-Hydroxy-11-methoxytabersonine)
- Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1987, **26**, 543 (Epimisilines)
- He, X. *et al.*, *Huaxue Xuebao*, 1992, **50**, 96-101; *CA*, **116**, 191092c (11,19R-Dihydroxytabersonine)

12-Hydroxy-2,4,8,10-tetradecatetraenoic acid H-744




(2E,4E,8Z,10E,12E)-form

2-Methylpropylamide: 12-Hydroxy-N-isobutyl-2,4,8,10-tetradecatetraenamide. **Lanyuamide VI**
 $\text{C}_{18}\text{H}_{29}\text{NO}_2$ 291.433
 Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. $[\alpha]_{\text{D}}^{25}$ -20.9 (c, 0.13 in CHCl_3). λ_{max} 237 (log ϵ 4.23); 259 (log ϵ 4.27) (EtOH).

Tsai, I.-L. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 830-833

11-Hydroxy-2,4,8-tetradecatrienoic acid H-745




(2E,4E,8Z,11E)-form

2-Methylpropylamide: 11-Hydroxy-N-isobutyl-2,4,8-tetradecatrienamide. **Lanyuamide IV**
 $\text{C}_{18}\text{H}_{31}\text{NO}_2$ 293.448
 Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. $[\alpha]_{\text{D}}^{25}$ -42.9 (c, 0.11 in CHCl_3). λ_{max} 260 (log ϵ 3.85) (EtOH).

Tsai, I.-L. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 830-833

4-Hydroxy-2-tetradecylquinoline H-746

2-Tetradecyl-4(1H)-quinolinone



$\text{C}_{23}\text{H}_{35}\text{NO}$ 341.536

NH-form

N-Me: 1-Methyl-2-tetradecyl-4(1H)-quinolinone


$\text{C}_{24}\text{H}_{37}\text{NO}$ 355.562
 Alkaloid from the fruit of *Evodia rutaecarpa*. Powder. λ_{max} 212 ; 238 ; 321 ; 333 (MeOH).

Ko, J.S. *et al.*, *Planta Med.*, 2002, **68**, 1131-1133 (*isol*, *pmr*, *cmr*, *ms*)

10-Hydroxy-1,4,6,9-tetramethylpyrido[3,2-g]quinoline-2,8(1H,9H)-dione H-747

BE 12233. Antibiotic BE 12233

[136158-10-6]




$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_3$ 284.314

Prod. by *Streptomyces* BA-12233. Antineoplastic agent. Mp 300°. λ_{max} 225 ; 283 ; 355 ; 374 (MeOH) (Berdy).

Japan. Pat., 1991, 91 30 688; *CA*, **115**, 157136 (*isol*, *props*)

7-Hydroxy-2,8-tridecadiene-10,12-dienoic acid H-748




(2E,7E,8E)-form

2-Methylpropylamide: 7-Hydroxy-N-(2-methylpropyl)-2,8-tridecadiene-10,12-dienamide
 [1017577-34-2]
 $\text{C}_{17}\text{H}_{23}\text{NO}_2$ 273.374
 Alkaloid from *Spilanthes callimorpha*.

Li, G.-P. *et al.*, *J. Integ. Plant Biol.*, 2007, **49**, 1608-1610 (*isol*)

2-(12-Hydroxytridecyl)-3-methoxy-4(1H)-quinolinone H-749



$\text{C}_{23}\text{H}_{35}\text{NO}_3$ 373.534

(+)-form

Alkaloid from the leaves of *Spathelia excelsa*. Oil. $[\alpha]_{\text{D}}^{25}$ +20.7 (c, 0.02 in CHCl_3). λ_{max} 240 ; 248 ; 326 ; 335 (MeOH).

12'-Ketone: 3-Methoxy-2-(12-oxotridecyl)-4(1H)-quinolinone
 $\text{C}_{23}\text{H}_{33}\text{NO}_3$ 371.519


Alkaloid from the leaves of *Spathelia excelsa*. Gum. λ_{max} 242 ; 247 ; 324 ; 335 (MeOH).

Da Paz Lima, M. *et al.*, *Phytochemistry*, 2005,

66, 1560-1566 (*isol*, *pmr*, *cmr*)

4-Hydroxy-2-tridecylquinoline H-750

2-Tridecyl-4(1H)-quinolinone. 2-Tridecyl-4-quinolinol, 9CI
 [14427-54-4]



$\text{C}_{22}\text{H}_{33}\text{NO}$ 327.509

Alkaloid from fruits of *Evodia rutaecarpa*. Metab. of *Pseudomonas aeruginosa*. Amorph. powder. Mp 132-134°. λ_{max} 212 (log ϵ 4.44); 235 (log ϵ 4.54); 315 (log ϵ 4.12); 327 (log ϵ 4.09) (MeOH).

N-Oxide: 2-Tridecyl-4(1H)-quinolinone N-oxide
 [185855-03-2]

$\text{C}_{22}\text{H}_{33}\text{NO}_2$ 343.508

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(E)-: 2-(1E-Trideceny)-4(1H)-quinolinone. 4-Hydroxy-2-(1E-trideceny)quinoline
 $\text{C}_{22}\text{H}_{31}\text{NO}$ 325.493

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(Z)-: 2-(1Z-Trideceny)-4(1H)-quinolinone. 4-Hydroxy-2-(1Z-trideceny)quinoline
 $\text{C}_{22}\text{H}_{31}\text{NO}$ 325.493

Metab. of *Pseudomonas aeruginosa*.

NH-form

N-Me: 1-Methyl-2-tridecyl-4(1H)-quinolinone, 9CI. **Dihydroevocarpine**
 [15266-35-0]

$\text{C}_{23}\text{H}_{35}\text{NO}$ 341.536

Alkaloid from the leaves and fruits of *Evodia rutaecarpa* (Rutaceae). Active against *Helicobacter pylori*. Needles (EtOAc/Et₂O). Mp 74-75° (57°).

7',8'-Didehydro(Z-), N-Me: 1-Methyl-2-(7-trideceny)-4(1H)-quinolinone
 [182056-19-5]

$\text{C}_{23}\text{H}_{33}\text{NO}$ 339.52

Alkaloid from fruits of *Evodia rutaecarpa*. Oil. *Isol.* as an inseparable mixt. with Evocarpine.

8',9'-Didehydro, N-Me: 1-Methyl-2-(8-trideceny)-4(1H)-quinolinone. **Evo-carpine**
 [15266-38-3]


$\text{C}_{23}\text{H}_{33}\text{NO}$ 339.52

Alkaloid from the fruit of *Evodia rutaecarpa* (Rutaceae). Active against *Helicobacter pylori*. Mp 34-38°.

Geometry of side-chain double bond not definitely known, inferred to be probably (Z-). λ_{max} 215 (log ϵ 4.25); 240 (log ϵ 4.34); 322 (log ϵ 4.04); 335 (log ϵ 4.06) (MeOH).

8',9'-Didehydro, N-Me, picrate: Mp 96-96.5°.

4',5',7',8'-Tetrahydro(Z,Z-), N-Me: 1-Methyl-2-(4,7-tridecadienyl)-4(1H)-quinolinone
 [120693-53-0]

C₂₃H₃₁NO 337.504Alkaloid from fruits of *Evodia rutaecarpa* (Rutaceae). Oil.**8-Methoxy, N-Me: 8-Methoxy-1-methyl-2-tridecyl-4(1H)-quinolinone, 9CI**
[132113-93-0]C₂₄H₃₇NO₂ 371.562Alkaloid from roots of *Esenbeckia leiocarpa* (Rutaceae).Tschesche, R. et al., *Tetrahedron*, 1967, **23**, 1873-1881 (*Evocarpine*, uv, ir, ms, pmr, isol, struct)Kamikado, T. et al., *Agric. Biol. Chem.*, 1976, **40**, 605-609 (*N-Me*)Sugimoto, T. et al., *Chem. Pharm. Bull.*, 1988, **36**, 4453-4461 (*N-Me*)Delle Monache, F. et al., *Gazz. Chim. Ital.*, 1990, **120**, 387-389 (*8-methoxy-N-Me*)Tang, Y.-Q. et al., *Phytochemistry*, 1996, **43**, 719-722 (*Evodia rutaecarpa* constits)Rho, T.C. et al., *Biol. Pharm. Bull.*, 1999, **22**, 1141-1143 (*Evocarpin*, *Dihydroevocarpine*, isol, activity)Lepine, F. et al., *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (*occur, 1',2'-didehydro*)Deziel, E. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (*biosynth*)**1-Hydroxy-2,9,10-trimethoxyaporphine H-751**5,6,6a,7-Tetrahydro-2,9,10-trimethoxy-6-methyl-4H-dibenzo[de,g]quinolin-1-ol, 9CI. O⁹-Methylisoboldine. *Thalicmidine*

(S)-form

C₂₀H₂₃NO₄ 341.406 λ_{\max} 220 (ϵ 33110); 280 (ϵ 13180); 305 (ϵ 13180) (EtOH) (Berdy).**(S)-form****Thaliporphine**

[5083-88-5]

Alkaloid from *Thalictrum fendleri* and other spp. (Ranunculaceae, Papaveraceae, Lauraceae, Rutaceae, Annonaceae, Berberidaceae, Magnoliaceae, Euphorbiaceae, Lauraceae). Soporific and narcotic enhancer. Hypotensive agent, produces motility disturbances and convulsions. Mp 170-172°. $[\alpha]_D$ +44 (c, 0.97 in EtOH). λ_{\max} 220 (log ϵ 4.52); 280 (log ϵ 4.12); 305 (log ϵ 4.12) (EtOH).► Toxic, LD₅₀ (mus/rbt, ivn) 10 mg/kg.**N-Oxide: Thalimidine N-oxide**

[41607-13-0]

C₂₀H₂₃NO₅ 357.405Alkaloid from *Thalictrum minus* and *Berberis integerrima* (Ranunculaceae, Berberidaceae). Mp 192-193° dec. λ_{\max} 227 (log ϵ 4.42); 282 (log ϵ 4.1); 308 (log ϵ 4.07) (no solvent reported).**N-Me: Fagara base. Aporphinium DVT**
[7224-60-4]

[18482-48-9, 68889-50-9, 47476-23-3]

C₂₁H₂₆NO₄[⊕] 356.441Quaternary alkaloid from *Fagara tinguassoiba* and *Fagara rhoifolia* (Rutaceae). Mp 215-219° (as chloride). $[\alpha]_D^{25}$ +30.2 (c, 2.16 in H₂O).**N-De-Me: 1-Hydroxy-2,9,10-trimethoxy-noraporphine. Wilsonirine. Aducaine. Northaliporphine**

[17807-64-6]

C₁₉H₂₁NO₄ 327.379Alkaloid from *Croton wilsonii* and *Monodora angolensis* (Euphorbiaceae, Annonaceae). Mp 211-213° dec. (108-110°). $[\alpha]_D$ +47 (c, 0.13 in MeOH).**(±)-form [2755-00-2]**

Synthetic. Mp 192-194°.

N-Me: [15358-35-7]

Mp 243-245° (as iodide). CAS no. refers to iodide.

N-De-Me: [39945-43-2]


Synthetic. Mp 214-216° dec. (210-213°).

Ac: [35587-63-4]

Mp 156-158°.

Tschesche, R. et al., *Tet. Lett.*, 1965, 445-449 (*struct*)Shamma, M. et al., *Tetrahedron*, 1967, **23**, 2887-2892 (*isol, uv, pmr*)Ismailov, Z.F. et al., *Khim. Prir. Soedin.*, 1968, **4**, 196; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 169-170 (*ms*)Johns, S.R. et al., *Aust. J. Chem.*, 1970, **23**, 363-368 (*Wilsonirine*)Hoshino, O. et al., *Chem. Comm.*, 1971, 1533 (*synth*)Khozhdaev, V.G. et al., *Khim. Prir. Soedin.*, 1972, **8**, 631-633; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 599-601 (*oxide*)Cava, M.P. et al., *J.O.C.*, 1973, **38**, 2394-2397 (*synth, ir, pmr, ms, Wilsonirine*)Karimov, A. et al., *Khim. Prir. Soedin.*, 1978, **14**, 419; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 360-361 (*oxide*)Jackmann, L.M. et al., *J. Nat. Prod.*, 1979, **42**, 437-449 (*cmr*)Castedo, L. et al., *Heterocycles*, 1980, **14**, 1135-1138 (*synth*)Hoshino, O. et al., *Heterocycles*, 1987, **25**, 151-153 (*synth, pmr, Wilsonirine*)Hara, H. et al., *Tetrahedron: Asymmetry*, 1995, **6**, 1683-1692 (*synth, Wilsonirine*)**10-Hydroxy-1,2,9-trimethoxyaporphine H-752**

5,6,6a,7-Tetrahydro-1,2,9-trimethoxy-6-methyl-4H-dibenzo[de,g]quinolin-10-ol, 9CI



(S)-form

C₂₀H₂₃NO₄ 341.406**(S)-form****Lirioferine**

[6883-42-7]

Alkaloid from the discoloured sapwood of *Liriodendron tulipifera* (Magnoliaceae). Needles (EtOAc). Mp 172-174°. $[\alpha]_D$ +128.6 (CHCl₃).**N-Me: Coccarmine**

[7096-88-0]

C₂₁H₂₆NO₄ 356.441Quaternary alkaloid from the roots of *Cocculus sarmentosus* (Menispermaceae). Cryst. + 3H₂O (as iodide). Mp 205-207° (sinters at 185°) (iodide) Mp 257-259° dec. (iodide). $[\alpha]_D^{22}$ +27.9 (c, 0.93 in EtOH).**N-De-Me: 10-Hydroxy-1,2,9-trimethoxy-noraporphine. Norlirioferine**


[86941-27-7]

C₁₉H₂₁NO₄ 327.379Alkaloid from the bark of *Phoebe pittieri* (Lauraceae). Mp 94-96°. No opt. rotn. recorded, abs. config. not detd. λ_{\max} 220 (log ϵ 4.36); 281 (log ϵ 3.83); 302 (log ϵ 3.95); 312 (sh) (log ϵ 3.89) (MeOH). λ_{\max} 280 (sh) (log ϵ 3.8); 310 (sh) (log ϵ 3.95); 315 (log ϵ 3.98); 330 (sh) (log ϵ 3.91) (MeOH/KOH).**(±)-form**

Synthetic. Mp 140-145°.

Hydroiodide:


Cryst. (EtOH). Mp 227-228°.

Tomita, M. et al., *Yakugaku Zasshi*, 1963, **83**, 190-194 (*Coccarmine*)Baarschers, W.H. et al., *Tetrahedron*, 1965, **21**, 2155-2158 (*synth, pmr, uv*)Chen, C.-L. et al., *Phytochemistry*, 1976, **15**, 1161-1167 (*isol, uv, pmr, ms, struct*)Castedo, L. et al., *Heterocycles*, 1980, **14**, 1135-1138 (*synth*)Castedo, G. et al., *An. Quim., Ser. C*, 1982, **78**, 103-107; *CA*, **97**, 56087d (*cmr*)Castro, C.O. et al., *Phytochemistry*, 1985, **24**, 203-204 (*Norlirioferine*)**4-Hydroxy-1',3,4'-tris(4-hydroxyphenyl)-3'-[2-(4-hydroxyphenyl)-ethyl]-7'-(sulfooxy)spiro[furan-2(5H),2'(3'H)-pyrrolo[2,3-c]carbazole]-5,5'(6'H)-dione, 9CI**
[149444-91-7]C₄₃H₃₀N₂O₁₂S 798.782Isol. from the marine sponge *Dictyodendrilla* sp. Potent aldose reductase inhibitor. Purple solid (as Na salt). Mp 300° (Na salt). Opt. inactive (racemic). λ_{\max} 229 (ϵ 40800); 289 (ϵ 20900); 324 (ϵ 19800); 391 (ϵ 9700); 475 (ϵ 1200) (EtOH) (Derep).

[149444-90-6]

Sato, A. et al., *J.O.C.*, 1993, **58**, 7632 (*isol, uv, ir, pmr, cmr, struct*)**2-Hydroxytryptamine** **H-754**


3-(2-Dimethylaminoethyl)oxindole. 3-(2-Dimethylaminoethyl)-2H-indol-2-one

C₁₀H₁₂N₂O 176.218

Oxo-form likely to predominate.

N²-Me: 3-[2-(Methylamino)ethyl]-1H-indol-2-ol, 9CI. **2-Hydroxy-N-methyltryptamine**. 3-(2-Methylaminoethyl)oxindole

[106987-89-7]

C₁₁H₁₄N₂O 190.244Alkaloid from the roots of *Desmanthus illinoensis* (Fabaceae).Thompson, A.C. et al., *J. Agric. Food Chem.*, 1987, **35**, 361-365 (*isol, ir, pmr, ms, struct*)**5-Hydroxytryptamine** **H-755**3-(2-Aminoethyl)-1H-indol-5-ol, 9CI. 3-(2-Aminoethyl)-5-hydroxyindole. **Serotonin**. Enteramine. Hippophaine. Anthemovister. Thrombocytin. Thrombotonin. 5 HT [50-67-9]C₁₀H₁₂N₂O 176.218

Found in blood and other tissues.

Widespread alkaloid from terrestrial and marine organisms, e.g. *Mucuna pruriens*, bananas (*Musa* sp.) and other fruits, basidiomycete *Panaeolus campanulatus*, cotton (*Gossypium hirsutum*) and other plants. Also from the sea anemone *Calliactis parasitica*, *Hippophae rhamnoides*, *Panaeolus* sp., *Pentacta crassa*, *Thelenoia ananas*, *Octopus vulgaris*, *Eledone moschata*, *Corallistes undulatus*, *Physalia* sp., *Hydra littoralis*, *Sycon ciliatum* and arthropod venoms and constit. of toad venom (*Bufo* spp.). Vasoconstrictor, hypotensive agent, neurotransmitter. Plays a major role in regulating behaviour such as aggression, depression, psychosis. pK_{a1} 1; pK_{a2} 4.52; pK_{a3} 15 (20°). Log P 0.61 (calc). Echinoderm name given apparently incorrectly as *Pentacta crasa*.

► LD₅₀ (mus, orl) 60 mg/kg. Exp. reprod. and teratogenic effects. NM2450000

Hydrochloride: [153-98-0][21591-86-6] Sol. H₂O. Mp 167-168°.

Light-sensitive. Aq. solns. stable at acid pH.

Creatinine sulfate complex: *Antemovis* [971-74-4]

[61-47-2]

Found in mammalian sera. Potent vasoconstrictor. Plates +1H₂O. Mp 214-216° dec. (monohydrate).

► NM2550000

O-Sulfate: [16310-20-6]C₁₀H₁₂N₂O₄S 256.282Metab. of 5-hydroxytryptamine excreted in mammalian urine. Cryst. (H₂O). Mp 192-194°.N^b-Ac: [1210-83-9]C₁₂H₁₄N₂O₂ 218.255

Metab. of 5-hydroxytryptamine. Enzyme inhibitor. Melanophore which produces night coloration in fish. Mp 120-122°.

N^b-Eicosanoyl: N^b-Eicosanoylserotonin [21249-34-3]C₃₀H₅₀N₂O₂ 470.737Constit. of coffee (*Coffea arabica*). Mp 122.5-123°. λ_{max} 278 (log ε 3.77) (no solvent reported).N^b-(11Z-Eicosenoyl): N^b-(11Z-Eicosenoyl)serotonin. **Scorodocarpine C**

[381719-44-4]

C₃₀H₄₈N₂O₂ 468.721Alkaloid from the fruit of *Scorodocarpus borneensis*.N^b-Docosanoyl: N^b-Docosanoylserotonin.**Scorodocarpine A**

[21249-35-4]

C₃₂H₅₄N₂O₂ 498.791Alkaloid from the fruit of *Scorodocarpus borneensis*. Constit. of green coffee beans.N^b-(13Z-Docosenoyl): N^b-(13Z-Docosenoyl)serotonin. **Scorodocarpine B**

[381719-43-3]

C₃₂H₅₂N₂O₂ 496.775Alkaloid from the fruit of *Scorodocarpus borneensis*.N^b-Tetracosanoyl: N^b-Tetracosanoylserotonin

[21249-36-5]

C₃₄H₅₈N₂O₂ 526.844Constit. of coffee (*Coffea arabica*). No phys. props. reported.N^b-Carbamoyl: N-Carbamoylserotonin.**Bufoserotonin A**

[1002722-85-1]

C₁₁H₁₃N₃O₂ 219.243Isol. from the venom of *Bufo* spp. (ChanSu). Amorph. solid (MeOH). Mp 156-158°. λ_{max} 221; 277; 300 (MeOH).N^b-(3-Carboxypropanoyl): **Bufobutanoic acid**

[74010-65-4]

C₁₄H₁₆N₂O₄ 276.291Alkaloid from the Chinese traditional drug Ch'an Su (prepared skin secretions of *Bufo* spp.). Amorph. solid. λ_{max} 225; 274; 300 (MeOH).N^b-(7-Carboxyheptanoyl): **Bufoserotonin B**

[1002722-86-2]

C₁₈H₂₄N₂O₄ 332.399Isol. from the venom of *Bufo* spp. (ChanSu). Amorph. powder. λ_{max} 236; 309 (MeOH).N^b-(15ξ-Hydroxy-9Z-hexadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide G**

[702702-58-7]

C₃₈H₆₀N₂O₁₃ 752.898Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. λ_{max} 277 (log ε 3.34); 301 (sh) (log ε 3.23) (MeOH).N^b-(14ξ-Hydroxyhexadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide C**

[702702-42-9]

C₃₈H₆₂N₂O₁₃ 754.913Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. [α]_D²⁰ -34 (c, 0.01 in MeOH). λ_{max} 276 (log ε 3.53); 300 (sh) (log ε 3.42) (MeOH).N^b-(15ξ-Hydroxyhexadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide B**

[702702-41-8]

C₃₈H₆₂N₂O₁₃ 754.913Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. [α]_D²⁰ -34 (c, 0.01 in MeOH). λ_{max} 277 (log ε 3.32); 300 (sh) (log ε 3.22) (MeOH).N^b-(16ξ-Hydroxyoctadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide E**

[702702-55-4]

C₄₀H₆₆N₂O₁₃ 782.967Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Pale brown solid. λ_{max} 275 (log ε 3.3); 300 (sh) (log ε 3.2) (MeOH).N^b-(16ξ-Hydroxy-9Z-octadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide F**

[702702-56-5]

C₄₀H₆₄N₂O₁₃ 780.951Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. λ_{max} 276 (log ε 3.31); 301 (sh) (log ε 3.21) (MeOH).N^b-(17ξ-Hydroxyoctadecenoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide D**

[702702-54-3]

C₄₀H₆₆N₂O₁₃ 782.967Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. λ_{max} 277 (log ε 3.43); 301 (sh) (log ε 3.3) (MeOH).N^b-(17R-Hydroxy-6Z,9Z-octadecadienoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide A**

[702702-22-5]

C₄₀H₆₂N₂O₁₃ 778.935Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Pale brown powder. [α]_D²⁰ -35 (c, 0.01 in MeOH). λ_{max} 276 (log ε 3.5); 300 (sh) (log ε 3.4) (MeOH).N^b-(17ξ-Hydroxy-6Z,9Z-octadecadienoyl), O-[[β-D-glucopyranosyl-(1→6)-β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Withanamide I**

[702702-72-5]

C₄₆H₇₂N₂O₁₈ 941.077

- Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. λ_{\max} 278 (log ϵ 3.43); 302 (sh) (log ϵ 3.3) (MeOH).
- N^b -(17 ξ -Hydroxy-6Z,9Z,12Z,15Z-octadecatetraenoyl), O-[β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **Withanamide H** [702702-59-8]
C₄₀H₅₈N₂O₁₃ 774.904
Alkaloid from the fruit of *Withania somnifera*. Lipid peroxidation inhibitor. Amorph. powder. λ_{\max} 276 (log ϵ 3.53); 301 (sh) (log ϵ 3.38); 330 (sh) (log ϵ 3.2) (MeOH).
- N^b -(4-Hydroxy-E-cinnamoyl): N^b -p-Coumaroylserotonin. **Ipobscurine A** [201301-83-9]
[68573-24-0]
C₁₉H₁₈N₂O₃ 322.363
Alkaloid from seeds of *Ipomoea obscura*, *Carthamus tinctorius* (safflower) and dried *Amorphophallus konjac* (Devil's tongue). Needles. Mp 195-196°.
- N^b -(4-Hydroxy-E-cinnamoyl), O- β -D-glucopyranoside: [76423-56-8]
C₂₅H₂₈N₂O₈ 484.505
Alkaloid from *Carthamus tinctorius* (safflower). Needles. Mp 240-242°.
[α]_D²⁵ -14.9 (c, 0.7 in MeOH).
- N^b -(4-Hydroxy-Z-cinnamoyl): N^b -cis-p-Coumaroylserotonin [314298-49-2]
C₁₉H₁₈N₂O₃ 322.363
Constit. of dried *Amorphophallus konjac* (Devil's tongue). Amorph. powder. Mp 205-206°. λ_{\max} 277 (log ϵ 4.4) (MeOH aq.).
- N^b -(4-Hydroxy-3-methoxy-E-cinnamoyl): N^b -(E)-Feruloylserotonin. **Moschamine** [193224-22-5]
[68573-23-9]
C₂₀H₂₀N₂O₄ 352.389
Alkaloid from *Carthamus tinctorius* (safflower) and *Centaurea moschata* (Asteraceae). Needles. Mp 115-117°. Isol. with Z-isomer as a mixt.
- N^b -(4-Hydroxy-3-methoxy-E-cinnamoyl), O- β -D-glucopyranoside: [201301-85-1]
C₂₆H₃₀N₂O₉ 514.531
Alkaloid from *Carthamus tinctorius* (safflower). Needles. Mp 143-145°.
[α]_D²⁵ -8.8 (c, 1.2 in MeOH).
- N^b -(4-Hydroxy-3-methoxy-Z-cinnamoyl): N^b -(Z)-Feruloylserotonin. **cis-Moschamine** [193224-24-7]
C₂₀H₂₀N₂O₄ 352.389
Alkaloid from the seeds of *Centaurea moschata* (Asteraceae). Isol. with E-isomer as a mixt.
- N^b -Me: 5-Hydroxy-N^o-methyltryptamine. N^b -Methylserotonin, 9CI [1134-01-6]
[15558-50-6]
C₁₁H₁₄N₂O 190.244
Isol. from *Citrus unshiu* (satsuma mandarin). Component of oviposition-

stimulating complex for the butterfly *Papilio xuthus*. Constit. of toad venom (*Bufo* spp.). Serotonin metab. in schizophrenic patients treated with MAO inhibitors. Mp 153-156° (as oxalate).

N^b , N^b -Di-Me: see Bufotenine, B-393
Me ether: see 5-Methoxytryptamine, M-288

Benzyl ether: [20776-45-8]
[52055-23-9]
Cryst. (MeOH) (as hydrochloride). Mp 265° (248-250°) (hydrochloride).

Hamlin, K.E. et al., *J.A.C.S.*, 1951, **73**, 5007-5008 (synth, uv)

Speeter, H.E. et al., *J.A.C.S.*, 1951, **73**, 5514-5515 (synth)

Asero, B. et al., *Annalen*, 1952, **576**, 69-74 (synth, uv)

Bowden, K. et al., *Nature (London)*, 1954, **174**, 925-926 (isol)

Abramovitch, R.A. et al., *Chem. Ind. (London)*, 1955, 1255 (synth)

Stoll, A. et al., *Helv. Chim. Acta*, 1955, **38**, 1452-1472 (N-Me)

Mathias, A.P. et al., *Nature (London)*, 1957, **180**, 658-659 (isol, occur)

Harms, U. et al., *Z. Lebensm.-Unters.-Forsch.*, 1968, **138**, 75-80 (coffee amides)

Thewalt, U. et al., *Acta Cryst. B*, 1972, **28**, 82-92 (cryst struct)

Couch, M.W. et al., *Anal. Biochem.*, 1972, **50**, 612-622 (ms)

Folstar, P. et al., *J. Agric. Food Chem.*, 1979, **27**, 12-15 (coffee amides, hplc)

Biagini, S. et al., *J.C.S. Faraday 2*, 1983, **79**, 491-496 (pmr, cmr)

Borgulya, J. et al., *Synthesis*, 1983, 29-30 (O-sulfate)

Falcon, J. et al., *Ann. Endocrinol.*, 1986, **47**, 65-66 (N^b -Ac, activity)

Eich, E. et al., *Planta Med.*, 1986, **52**, 523 (*Ipobscurine A*)

Nishida, R. et al., *Experientia*, 1987, **43**, 342-344 (5-Hydroxy-N^o-methyltryptamine)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1411

Zhang, H.L. et al., *Chem. Pharm. Bull.*, 1997, **45**, 1910-1914 (isol, cinnamides)

Somei, M. et al., *Heterocycles*, 1997, **46**, 91-94 (synth)

Sarker, S.D. et al., *Nat. Prod. Lett.*, 1997, **9**, 189-199 (*Moschamines*)

Zaunreiter, M. et al., *NeuroReport*, 1998, **9**, 1475-1479; *CA*, **129**, 173288 (N^b -Ac, activity)

Kamano, Y. et al., *Heterocycles*, 1999, **50**, 499-503 (*Bufobutanoic acid*)

Neuropsychopharmacology, Special Supplement Issue, Serotonin 50th Anniversary, 1999, **21**, 2S-115S (rev)

Niwa, T. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2269-2271 (*N*-coumaroylserotonin)

Kurauchi, T. et al., *Heterocycles*, 2000, **53**, 1017-1019 (synth, *Bufobutanoic acid*)

Somei, M. et al., *Chem. Pharm. Bull.*, 2001, **49**, 87-96 (synth)

Wiert, C. et al., *Phytochemistry*, 2001, **58**, 653-656 (*Scorodocarpines*)


Jayaprakasam, B. et al., *Tetrahedron*, 2004, **60**, 3109-3121 (*Withanamides*)

Zhang, P. et al., *Chem. Pharm. Bull.*, 2005, **53**, 1582-1586 (toad venom constits)

Liu, R.-H. et al., *Helv. Chim. Acta*, 2007, **90**, 2427-2431 (*Bufoserotonins A,B*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, (AJX500, AJX750)

10-Hydroxy-2,4,6,8-undecatetraenoic acid, 9CI



C₁₁H₆O₃ 186.167

Amide: [83475-37-0]


C₁₁H₇NO₂ 185.182

Major metabolite isol. from *Mycena viridimarginata*. Light yellow cryst. turning brown immediately. Poorly sol. hexane. λ_{\max} 236; 247; 299; 319; 341; 367 (Et₂O) (Berdy).

Jente, R. et al., *Phytochemistry*, 1985, **24**, 553 (isol, struct)

4-Hydroxy-2-undecylquinoline

2-Undecyl-4(1H)-quinolinone, 9CI. 2-Undecyl-4-quinolinol, 9CI [56183-46-1]



C₂₀H₂₉NO 299.455

Alkaloid from *Evodia rutaecarpa*, *Ptelea trifoliata* and roots of *Ruta graveolens* (rue) (as the main component of an inseparable mixt. of 2-alkylquinolones contg. the 2-dodecyl, 2-tridecyl and 2-tetradecyl homologues). Metab. of *Pseudomonas aeruginosa*. Antagonist of Dihydrostreptomycin. Cryst. (Me₂CO). Mp 130-132°. λ_{\max} 213 (log ϵ 4.43); 235 (log ϵ 4.44); 315 (log ϵ 4.03); 327 (log ϵ 4.01) (MeOH).

N-Oxide: 2-Undecyl-4(1H)-quinolinone N-oxide

[2503-86-8]

C₂₀H₂₉NO₂ 315.455

Metab. of *Pseudomonas pyocyanea* and *Pseudomonas aeruginosa*. Isol. from flowers of *Ptelea trifoliata* and roots of *Ruta graveolens* (rue) (Rutaceae). Leaflets. Mp 148.5-149.5°.

N-Me: 1-Methyl-2-undecyl-4(1H)-quinolinone, 9CI

[59443-02-6]

C₂₁H₃₁NO 313.482

Alkaloid from the leaves and fruits of *Evodia rutaecarpa* (Rutaceae). Active against *Helicobacter pylori*. Needles (EtOAc/Et₂O). Mp 68.5-70°.

1',2'-Didehydro(E-): 2-(1-Undecenyl)-4(1H)-quinolinone. 4-Hydroxy-2-(1-undecenyl)quinoline [678176-79-9]

C₂₀H₂₇NO 297.439

Metab. of *Pseudomonas aeruginosa*.

1',2'-Didehydro(Z-):

C₂₀H₂₇NO 297.439

Metab. of *Pseudomonas aeruginosa*.

3',4'-Didehydro: 2-(3-Undecenyl)-4(1H)-quinolinone. **Pyo V** [71932-11-1]

C₂₀H₂₇NO 297.439

From *Pseudomonas aeruginosa*. Mp

114°. λ_{\max} 213 (ε 37300); 236 (ε 37000); 316 (ε 12200); 327 (ε 12900) (MeOH) (Berdy).

5,6'-Didehydro(Z-), N-Me: 1-Methyl-2-(5-undecenyl)-4(1H)-quinolinone

$C_{21}H_{29}NO$ 311.466

Alkaloid from fruits of *Evodia rutaecarpa*. Oil. Isol. as an inseparable mixt. with the Δ^6 -isomer.

6,7'-Didehydro(Z-), N-Me: 1-Methyl-2-(6-undecenyl)-4(1H)-quinolinone

$C_{21}H_{29}NO$ 311.466

Alkaloid from fruits of *Evodia rutaecarpa* (Rutaceae). Oil.

10'-Acetoxy, N-(acetoxyethyl): 1-Acetoxyethyl-2-(10-acetoxyundecyl)-4(1H)-quinolinone

$C_{25}H_{35}NO_5$ 429.555

Alkaloid from a new Australian species tentatively aff. *Samadera bidwillii*. Dark yellow oil. $[\alpha]_D^{25}$ -1 (c, 0.45 in MeOH).

10'-Oxo: 2-(10-Oxoundecyl)-4(1H)-quinolinone

[29479-53-6]

$C_{20}H_{27}NO_2$ 313.439

Alkaloid from *Evodia rutaecarpa* and *Vepris ampody*. Cryst. (Me₂CO). Mp 126°.

10'-Oxo, N-Me: 1-Methyl-2-(10-oxoundecyl)-4(1H)-quinolinone

$C_{21}H_{29}NO_2$ 327.466

Alkaloid from the fruit of *Evodia rutaecarpa* var. *bodinaieri*. Amorph. powder. λ_{\max} 216; 239; 321; 334 (MeOH).

Cornforth, J.W. et al., *Biochem. J.*, 1956, **63**, 124-130 (isol)

Kan-Fan, C. et al., *Phytochemistry*, 1970, **9**, 1283-1291 (10'-oxo)

Reisch, T. et al., *Phytochemistry*, 1975, **14**, 840-841 (isol, ir, pmr)

Kamikado, T. et al., *Agric. Biol. Chem.*, 1976, **40**, 605-609 (N-Me)

Budzikiewicz, H. et al., *Monatsh. Chem.*, 1979, **110**, 947-953 (N-oxide)

Sugimoto, T. et al., *Chem. Pharm. Bull.*, 1988, **36**, 4453-4461 (*Evodia rutaecarpa* constits)

Tang, Y.-Q. et al., *Phytochemistry*, 1996, **43**, 719-722 (*Evodia rutaecarpa* constits)

Gibbons, S. et al., *Phytochemistry*, 1997, **44**, 1109-1114 (10'-acetoxy-N-acetoxyethyl)

Tae, C.R. et al., *Biol. Pharm. Bull.*, 1999, **22**, 1141 (N-Me, activity)

Lepine, F. et al., *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 862-869 (occur, 1',2'-didehydro)


Deziel, E. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 1339-1344 (biosynth)

Yang, X.-W. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 697-703 (isol, pmr, cmr, ms)

8-Hydroxy-1-vinyl-β-carboline H-758

1-Ethenyl-9H-pyrido[3,4-b]indol-8-ol, 9CI

[138683-69-9]



$C_{13}H_{10}N_2O$ 210.235

Alkaloid from the marine bryozoans

Cribricellina cribraria and *Catenicella cribraria*. Exhibits cytotoxicity. Yellow oil. Pmr data revised in 1993. λ_{\max} 240 (ε 11700); 262 (ε 11500); 287 (ε 10800); 386 (ε 6610) (MeOH/NaOH) (Derep). λ_{\max} 230 (ε 11500); 244 (ε 12300); 296 (ε 9330); 369 (ε 8510) (MeOH) (Derep). λ_{\max} 240 (ε 11750); 262 (ε 11450); 287 (ε 10715); 386 (ε 6600) (EtOH) (Berdy).


Prinsep, M.R. et al., *J. Nat. Prod.*, 1991, **54**, 1068 (isol, uv, ir, pmr, cmr, struct)

Beutler, J.A. et al., *J. Nat. Prod.*, 1993, **56**, 1825 (isol, pmr)

Hygrine

H-759

1-(1-Methyl-2-pyrrolidinyl)-2-propanone, 9CI. 2-Acetyl-1-methylpyrrolidine



(S)-form

$C_8H_{15}NO$ 141.213

Isol. samples of Hygrine are racemic due to ready racemisation esp. in alkaline soln. In the plant, enantiomers are present and both can be precursors of tropane alkaloids, depending on spp. Alkaloid from *Erythroxyllum truxillense*, *Erythroxyllum coca*, *Nicandra physaloides*, *Convolvulus hamadana*, *Datura*, *Atropa*, *Hyoscyamus*, *Physalis*, *Dendrobium* and *Cochlearia* spp. (Erythroxyllaceae, Solanaceae, Convolvulaceae, Orchidaceae, Brassicaceae). Intermed. in biosynth. of tropane alkaloids.

(R)-form [496-49-1]

(+)-Tartrate: Mp 129-130°. $[\alpha]_D^{21}$ +30.1 (c, 6.1 in H₂O).

(S)-form [65941-22-2]

Liq. d_4^{25} 0.94. Bp 193-195° Bp₅₀ 111-113°.

(+)-Tartrate: Mp 67-68°. $[\alpha]_D^{21}$ +4.1 (c, 6.5 in H₂O) (-1.8).

N-De-Me, N-E-cinnamoyl: **trans-Dendrochrysanine**

[851439-76-4]

$C_{16}H_{19}NO_2$ 257.332

Alkaloid from the stems of *Dendrobium chrysanthum*. Viscous oil. $[\alpha]_D^{20}$ -19.2 (c, 3.4 in CHCl₃).

N-De-Me, N-Z-cinnamoyl: **cis-Dendrochrysanine**

[851439-77-5]

$C_{16}H_{19}NO_2$ 257.332

Alkaloid from the stems of *Dendrobium chrysanthum*. Viscous oil. $[\alpha]_D^{20}$ -17.7 (c, 3.5 in CHCl₃).

(±)-form [45771-52-6]

Bp₁₆ 88°.

Picrate:

Yellow needles. Mp 158° (155°).

Oxime: Mp 125°.

(ξ)-form

N-De-Me: **Norhygrine**. N-Demethylhygrine

$C_7H_{13}NO$ 127.186

Alkaloid from whole plants of *Nierembergia hippomanica* (Solanaceae).

CAS erroneously assigns the stereochem. as R- and CAS no. 130325-43-8.

1',2'-Didehydro(E-): 1-(1-Methyl-2-pyrrolidinylidene)-2-propanone. 1',2-Dehydrohygrine

[126412-05-3]

[39178-30-8]

$C_8H_{13}NO$ 139.197

Alkaloid from the stem bark of *Erythroxyllum lucidum*. λ_{\max} 304 (ε 26000) (EtOH aq.).

Liebermann, C. et al., *Ber.*, 1889, **22**, 675-679; 1895, **28**, 578-585 (isol, struct)

Willstätter, R. et al., *Ber.*, 1900, **33**, 1160-1166 (synth)

Šorm, F. et al., *Coll. Czech. Chem. Comm.*, 1947, **12**, 245-250 (synth)

Galinovsky, F. et al., *Monatsh. Chem.*, 1953, **84**, 798-808 (synth, resoln)

Lukeš, R. et al., *Coll. Czech. Chem. Comm.*, 1960, **25**, 483-491 (abs config)

Lüning, B. et al., *Acta Chem. Scand.*, 1965, **19**, 1607-1611 (isol, ms, pmr)

Romeike, A. et al., *Naturwissenschaften*, 1965, **52**, 619-620 (isol)

McGaw, B.A. et al., *Phytochemistry*, 1978, **17**, 257-259 (occur, props)

McGaw, B.A. et al., *Tet. Lett.*, 1979, 3135-3138 (biosynth)

Langenskiöld, T. et al., *Heterocycles*, 1983, **20**, 671-675 (synth)

Ghirlando, R. et al., *Tetrahedron*, 1984, **40**, 2879-2884 (synth)

Leete, E. et al., *Phytochemistry*, 1985, **24**, 953-955 (biosynth)

Nagasaka, T. et al., *Heterocycles*, 1989, **29**, 155-164 (synth, ir, pmr)

Pomilio, A.B. et al., *Phytochemistry*, 1996, **41**, 1393-1398 (Norhygrine)

Brachet, A. et al., *Phytochemistry*, 1997, **46**, 1439-1442 (Dehydrohygrine)

Yang, L. et al., *Heterocycles*, 2005, **65**, 633-636 (Dendrochrysanines)

Konno, H. et al., *Heterocycles*, 2006, **68**, 2579-2585 (Dendrochrysanines, synth)

Lee, J.-H. et al., *J.O.C.*, 2006, **71**, 6690-6692 (synth)

Arévalo-García, E.B. et al., *Tet. Lett.*, 2008, **49**, 3995-3996 (synth)


Hygriline

H-760

α,1-Dimethyl-2-pyrrolidineethanol, 9CI.

1-(1-Methyl-2-pyrrolidinyl)-2-propanol.

2-(2-Hydroxypropyl)-1-methylpyrrolidine



(2R,2'R)-form

$C_8H_{17}NO$ 143.228

(2R,2'R)-form [496-47-9]

Alkaloid from *Erythroxyllum coca* (minor congener of cocaine) and from *Cochlearia arctica* (Erythroxyllaceae, Brassicaceae). Needles (petrol). Mp 33-34°. $[\alpha]_D$ -63.2 (c, 11.4 in H₂O).

3,5-Dinitrobenzoyl: Mp 68°.

(+)-Tartrate:

Cryst. + 2H₂O. Mp 69°. $[\alpha]_D^{21}$ +4.1 (c, 6.5 in H₂O). $[\alpha]_D^{18}$ -1.8.

(2S,2'S)-form [1617-83-0]

Major alkaloid from leaves of *Carallia*

brachiata and from *Gynotroches axillaris* (Rhizophoraceae). Mp 29-31°. $[\alpha]_D^{25}$ +87 (c, 0.90 in H₂O) (+84). $[\alpha]_D$ +50 (c, 0.77 in EtOH). $[\alpha]_D$ +39.3 (EtOH) (synthetic).
Methiodide: Mp 124°. $[\alpha]_D^{20}$ +11 (c, 2.69 in H₂O).

Ac; hydrochloride:

Cryst. (Me₂CO). Mp 177°.

Ac, picrate: Mp 128-129° (120-123°).

(2R,2'S)-form

Pseudohygroline

[496-48-0]

Alkaloid from roots of *Schizanthus hoo-keri* (Solanaceae). Oil. $[\alpha]_D^{25}$ +73.13 (c, 1.68 in EtOH).

Methiodide:

Cryst. (MeOH). Mp 118°.

Ac, picrate: Mp 165° (163°).

(2RS,2'RS)-form

(±)-Hygroline

Cryst. or oil. Mp 24-25°. Bp₁₂ 84°. n_D^{20} 1.4597.

(+)-Tartrate: Mp 132°.

Ac, picrate: Mp 127-128°.

(2RS,2'SR)-form

(±)-Pseudohygroline

Bp₁₃ 90-94°. n_D^{20} 1.4653.

Picrate: Mp 95°.

Methiodide:

Cryst. (EtOH/Et₂O). Mp 117-119°.

Ac, picrate:

Cryst. (EtOH). Mp 158°.

N-De-Me: Norpseudohygroline

Bp_{1,2} 85-87°. n_D^{20} 1.4792.

Späth, E. *et al.*, *Ber.*, 1943, **76**, 942

Galinovsky, F. *et al.*, *Monatsh. Chem.*, 1953, **84**, 798 (*synth, resoln*)

Lukeš, R. *et al.*, *Coll. Czech. Chem. Comm.*, 1960, **25**, 483 (*abs config*)

Platonova, T.F. *et al.*, *CA*, 1964, **60**, 8350a (*isol*)

Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1965, **18**, 589 (*isol, ms, pmr*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1303 (*isol*)

San Martin, A. *et al.*, *Phytochemistry*, 1980, **19**, 2007 (*isol, Pseudohygroline*)

Shono, T. *et al.*, *J.O.C.*, 1986, **51**, 2590 (*synth*)

Takahata, H. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 2801-2810 (*synth*)


Enierga, G. *et al.*, *Tet. Lett.*, 1998, **39**, 2813-2814 (*synth*)

Knight, D.W. *et al.*, *Tet. Lett.*, 1999, **40**, 5915-5918 (*synth*)

Hymeniacidine

H-761

5-(Formylamino)-6-(methylamino)-2,4-pyrimidinedione. 5-(Formylamino)-6-(methylamino)uracil. N-Methyl-N-[1,2,3,4-tetrahydro-1,3-dimethyl-6-(methylamino)-2,4-dioxo-5-pyrimidinyl]-formamide, 9CI
 [104509-28-6]



C₈H₁₂N₄O₃ 212.208


Seco-xanthine formamide. Alkaloid from the sponge *Hymeniacidon* sp. Viscous pale yellow oil (natural). Cryst. (EtOAc) (synthetic). Mp 158-159° (synthetic). Exists as two formamide rotamers. λ_{max} 271 (ε 11400) (MeOH).

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2002, **65**, 368-370 (*isol, synth, pmr, cmr, ms*)

Hymenialdisine

H-762

4-(2-Amino-4-oxo-2-imidazolidin-5-ylidene)-2-bromo-4,5,6,7-tetrahydropyrrolo[2,3-c]azepin-8-one
 [82005-12-7]



C₁₁H₁₀BrN₅O₂ 324.136

Isol. from the Okinawan marine sponge *Hymeniacidon aldii*, the Red Sea sponge *Acanthella aurantiaca* and the sponges *Axinella verrucosa* and *Axinella carteri*. Exhibits potent cytotoxicity against murine P388 lymphocytic leukaemia cells. Shows insecticidal activity. Inhibits interleukin-1 stimulated rheumatoid synovial fibroblasts. Antiinflammatory agent. Yellow needles + 1MeOH (MeOH aq.) or yellow amorph. solid. Sol. DMSO, insol. most solvs. Mp 160-164° dec. λ_{max} 229 (ε 11900); 270 (ε 11700); 346 (ε 15900) (MeOH) (Derep).

(E)-Isomer: (10E)-Hymenialdisine

C₁₁H₁₀BrN₅O₂ 324.136

Isol. from the marine sponge *Stylotella aurantium*.

Debromo: Debromohymenialdisine

[75593-17-8]

C₁₁H₁₁N₅O₂ 245.24

Isol. from the Great Barrier Reef sponge *Phakellia flabellata*, *Hymeniacidon aldii*, and from an unidentified sponge from Korolevu, Fiji. Exhibits potent cytotoxicity against murine P388 lymphocytic leukaemia cells. Also shows insecticidal activity. Cryst. + 2H₂O (H₂O). Mp 220-225° dec. λ_{max} 224 (ε 8600); 252 (ε 6300); 338 (ε 15900) (HCl salt) (Derep). λ_{max} 228 (ε); 272 (ε); 365 (ε) (5% NaOH) (Derep). λ_{max} 228 (ε 11700); 262 (ε 10000); 346 (ε 18000) (MeOH) (Derep).

Debromo; hydrochloride: Mp 230-235° dec.

Debromo, (E)-isomer: (10E)-Debromohymenialdisine

C₁₁H₁₁N₅O₂ 245.24

Isol. from the marine sponge *Stylotella aurantium*.

3-Bromo: 3-Bromohymenialdisine

[175421-15-5]

C₁₁H₉Br₂N₅O₂ 403.032

From *Axinella carteri*. Cytotoxic agent. λ_{max} 269; 332 (MeOH) (Berdy).

3-Bromo, (E)-isomer: Spongiacidin A

[206433-68-3]

C₁₁H₉Br₂N₅O₂ 403.032

Isol. from a *Hymeniacidon* sp. and

Stylissa carteri. Enzyme inhibitor.

Amorph. solid. λ_{max} 206 (ε 10000); 270 (ε 7000); 332 (ε 7000) (MeOH).

3-Bromo, 2-debromo, (E)-isomer: Spongiacidin B

[206443-95-0]

C₁₁H₁₀BrN₅O₂ 324.136

Isol. from a *Hymeniacidon* sp. Enzyme inhibitor. Amorph. solid. λ_{max} 204 (ε 10000); 272 (ε 10000); 331 (ε 9000) (MeOH).

Sharma, G.M. *et al.*, *Chem. Comm.*, 1980, 435 (*deriv*)

Mattia, C.A. *et al.*, *Acta Cryst. B*, 1982, **38**, 2513 (*cryst struct*)

Cimino, G. *et al.*, *Tet. Lett.*, 1982, **23**, 767 (*isol, uv, pmr, cmr, cryst struct*)

Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2321 (*isol, uv, ir, cmr, ms, cryst struct, deriv*)

Schmitz, F.J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 47 (*isol, uv, ir, pmr, ms*)

Utkina, N.K. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 578; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 547 (*isol, deriv*)

Annoura, H. *et al.*, *Tet. Lett.*, 1995, **36**, 413 (*synth, Hymenialdisine, Debromohymenialdisine*)

Supriyono, A. *et al.*, *Z. Naturforsch., C*, 1995, **50**, 669 (*3-Bromohymenialdisine*)

Williams, D.H. *et al.*, *Nat. Prod. Lett.*, 1996, **9**, 57 (*10E-Hymenialdisine, 10E-Debromohymenialdisine*)

Roshak, A. *et al.*, *J. Pharmacol. Exp. Ther.*, 1997, **283**, 955-961 (*pharmacol*)

Xu, Y. *et al.*, *J.O.C.*, 1997, **62**, 456 (*synth, Hymenialdisine, Debromohymenialdisine*)

Inaba, K. *et al.*, *J. Nat. Prod.*, 1998, **61**, 693-695 (*Spongiacidins*)

Barrios Sosa, A.C. *et al.*, *J.O.C.*, 2000, **65**, 610-611 (*Debromohymenialdisine, synth*)


Hoffmann, H. *et al.*, *Synthesis*, 2003, 1753-1783 (*rev*)

Papeo, G. *et al.*, *Org. Lett.*, 2005, **7**, 5641-5644 (*synth*)

Hymenocardine

H-763

N,N-Dimethyl-L-isoleucyl-N-[7-(1H-indol-3-ylmethyl)-3,3-dimethyl-5,8,11-trioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-12,14,15-trien-4-yl]-L-valinamide, 9CI
 [19771-56-3]



Absolute Configuration

C₃₇H₅₀N₆O₆ 674.839

Alkaloid from *Hymenocardia acida* (Hymenocardiaceae). Cryst. + 1H₂O


(Me₂CO). Mp 261°. [α]_D -124 (c, 1 in CHCl₃).

Pais, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 2979 (isol, uv, ir, pmr, ms, struct)

Pais, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869 (cmr)

Hypaphorine**H-764**

α -Carboxy-N,N,N-trimethyl-1H-indole-3-ethaniminium hydroxide inner salt, 9CI. Trimethyltryptophan betaine. 1-Trimethylammonio-3-(3-indolyl)propionate. Tryptophan betaine. Glyyunnanenine. Lenticine



C₁₄H₁₈N₂O₂ 246.308

(R)-form*D*-form**6-Bromo- 6-Bromohypaphorine**

[158250-30-7]

C₁₄H₁₇BrN₂O₂ 325.205

Isol. from the Okinawan marine sponge *Aplysina* sp. Yellow needles. Mp 275° dec. [α]_D¹⁹ -27 (c, 0.80 in MeOH/CF₃COOH 8:1). λ _{max} 227 (ε 36000); 285 (ε 9300); 293 (ε 8000) (MeOH) (Berdy).

(S)-form*L*-form

[487-58-1]

Present in seeds of the genus *Erythrina* (Fabaceae). Isol. from *Abrus precatorius*, *Astragalus lusitanicus* and seedlings of lentil (*Lens culinaris*). Feeding deterrent for a seed-eating rodent (*Lionys salvini*). Neurotoxin. Cryst. (H₂O or dil. acid). V. sol. H₂O, EtOH, insol. most other solvs. Mp 255°. [α]_D²⁰ +140 (c, 1 in H₂O).

► Convulsive poison.

5-Bromo- 5-BromohypaphorineC₁₄H₁₇BrN₂O₂ 325.205

Isol. from the sponge *Thorectandra* sp. Yellow solid. [α]_D²⁵ +46.3 (c, 0.5 in MeOH).

6-Bromo- [64364-14-3]C₁₄H₁₇BrN₂O₂ 325.204

Metab. of the sponge *Pachymatisma johnstoni*. Rods. Mp 275-280° dec. [α]_D¹⁵ +58 (MeOH/CF₃COOH 8:1).

5-Iodo, 7-chloro- Plakohypaphorine FC₁₄H₁₆ClIN₂O₂ 406.65

Isol. from *Plakortis simplex*. Pale yellow solid. [α]_D²⁵ +26.7 (c, 1.5 in MeOH/TFA). λ _{max} 226 (ε 17200); 286 (ε 3740) (MeOH).

7-Iodo- Plakohypaphorine A

[502686-54-6]

C₁₄H₁₇IN₂O₂ 372.205

Isol. from the sponge *Plakortis simplex*. Pale yellow solid. [α]_D²⁵ +17.3 (c,

1.5 in MeOH/CF₃COOH). λ _{max} 224 (ε 17210); 286 (ε 3740) (MeOH).

5,6-Diiodo- Plakohypaphorine DC₁₄H₁₆I₂N₂O₂ 498.102

Isol. from *Plakortis simplex*. Pale yellow solid. [α]_D²⁵ +27.1 (c, 2 in MeOH/TFA). λ _{max} 224 (ε 17200); 288 (ε 3700) (MeOH).

5,7-Diiodo- Plakohypaphorine C

[502686-56-8]

C₁₄H₁₆I₂N₂O₂ 498.102

Isol. from the sponge *Plakortis simplex*. Yellow solid. [α]_D²⁵ +29.1 (c, 1 in MeOH/CF₃COOH). λ _{max} 231 (ε 8190); 284 (ε 1210) (MeOH).

6,7-Diiodo- Plakohypaphorine B

[502686-55-7]

C₁₄H₁₆I₂N₂O₂ 498.102

Isol. from the sponge *Plakortis simplex*. Yellow solid. [α]_D²⁵ +30.4 (c, 1.2 in MeOH/CF₃COOH). λ _{max} 236 (ε 18600); 290 (ε 3690) (MeOH).

5,6,7-Triiodo- Plakohypaphorine EC₁₄H₁₅I₃N₂O₂ 623.998

Isol. from *Plakortis simplex*. Pale yellow solid. [α]_D²⁵ +31.2 (c, 4 in MeOH/TFA). λ _{max} 224 (ε 17210); 288 (ε 3740) (MeOH).

(±)-form [64363-86-6]

Mp 253-255°.

Van Romburgh, P. *et al.*, *J.C.S.*, 1911, **99**,

2068-2071 (struct, synth)

Cahill, W.M. *et al.*, *J. Biol. Chem.*, 1938, **126**,

29-36 (synth)

Folkers, K. *et al.*, *J.A.C.S.*, 1940, **62**, 436-441

(isol)

Hofinger, M. *et al.*, *Phytochemistry*, 1975, **14**,

475-477 (isol, biosynth)

Raverty, W.D. *et al.*, *J.C.S. Perkin 1*, 1977,

1204-1211 (6-Bromohypaphorine, synth,

isol)

Kondo, K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1008-

1011 (isol, uv, ir, pmr, cmr, struct, 6-

Bromohypaphorine)

Hu, J.-F. *et al.*, *CA*, 1996, **124**, 50632g

(Glyyunnanenine)


Campagnuolo, C. *et al.*, *Eur. J. Org. Chem.*,

2003, 284-287; 2004, 3227-3232

(Plakohypaphorines)

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2005, **68**,

1484-1488 (5-Bromohypaphorine)


Bel-Kassaoui, H. *et al.*, *Nat. Prod. Res.*, 2008,**22**, 461-465 (isol, synth, pmr, cmr)**Hypecoleptopine****H-765**C₂₄H₁₉N₃O₆ 445.431**(±)-form [401947-62-4]**Alkaloid from *Hypecoum leptocarpum*.

Amorph. powder (MeOH). Mp 206.5-209.5°. λ _{max} 260 (log ε 4.56); 295 (sh); 350 (log ε 3.44) (MeOH).

Li, B.-G. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 1215-1218 (isol, uv, pmr, cmr, ms)

Hypecorine**H-766**

[41787-56-8]

C₂₀H₁₉NO₅ 353.374**(±)-form [88271-32-3]**

Alkaloid from flowering *Hypecoum erectum* (Hypecoaceae). Cryst. (Me₂CO). Mp 160-162° (154-156°). The nat. alkaloid is presumably racemic. Phys. props. refer to synthetic material. λ _{max} 208 (log ε 4.55); 248 (log ε 4.15); 294 (log ε 3.85); 369 (log ε 3.89) (MeOH).

8-Oxo- Leptopidine

[221347-16-6]

C₂₀H₁₇NO₆ 367.357

Alkaloid from *Hypecoum leptocarpum*. Yellow needles (MeOH/CHCl₃). Mp 345-350°. λ _{max} 240 (log ε 3.96); 307 (log ε 3.66); 330 (log ε 3.65); 410 (log ε 3.88) (MeOH).

13-Oxo- Hypecorinine. Corydalispirone

[41787-57-9]

[56142-90-6]

C₂₀H₁₇NO₆ 367.357

Alkaloid from *Hypecoum erectum*, *Pteridophyllum racemosum* and *Corydalis incisa* (Hypecoaceae, Papaveraceae). Pteridophyllaceae, Papaveraceae). Prisms (CHCl₃/MeOH). Mp 196-198°. Racemic. λ _{max} 240 (log ε 4.36); 292 (log ε 4.08); 320 (log ε 3.93) (MeOH).

13-Oxo, perchlorate:

Pale brown needles (MeOH). Mp 248-250° dec.

8,13-Dioxo- 8-Oxohypecorinine

[163165-34-2]

C₂₀H₁₅NO₇ 381.341

Alkaloid from whole plants of *Hypecoum procumbens* var. *glaucescens* (Hypecoaceae). Amorph. orange powder. Racemic. λ _{max} 224 (log ε 4.46); 245 (sh) (log ε 4.11); 308 (log ε 3.82); 362 (log ε 3.8) (MeOH).

8,13-Dioxo, N-oxide- 8-Oxohypecorinine N-oxide

[401892-68-0]


C₂₀H₁₅NO₈ 397.34

Alkaloid from *Hypecoum leptocarpum*. Amorph. yellow powder (MeOH). Mp 200-205°. Racemic. λ _{max} 248 (log ε 4.06); 309 (log ε 3.87); 374 (log ε 3.77) (MeOH).

- Yakhontova, L.D. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 624-628; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 592-595 (*Hypecorine, Hypecorinine, isol, struct*)
- Nonaka, G. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 294-298 (*Hypecorinine, uv, ir, pmr, ms, struct*)
- Ikuta, A. *et al.*, *Phytochemistry*, 1976, **15**, 577-578 (*Hypecorinine, isol, uv, ir, pmr, ms*)
- Nalliah, B.C. *et al.*, *Can. J. Chem.*, 1978, **56**, 1378-1380 (*Hypecorinine, synth, ir, pmr*)
- Gözler, B. *et al.*, *J.C.S. Perkin 1*, 1983, 2431-2434 (*synth, uv, ir, pmr, ms*)
- Iwasa, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 998-1008 (*synth, pmr, cmr, ms*)
- Rozwadowska, M.D. *et al.*, *Tetrahedron*, 1988, **44**, 1221-1226 (*Hypecorinine, synth, ir, pmr, ms*)
- Saad, H.E.A. *et al.*, *Phytochemistry*, 1995, **38**, 1049-1051 (*8-Oxohypecorinine*)
- Zhou, Y. *et al.*, *Phytochemistry*, 1999, **50**, 339-343 (*Leptopidine*)
- Li, B.-G. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 1215-1218 (*8-Oxohypecorinine N-oxide*)

Hypecoumine

Decumbenine C
[100163-16-4]




$C_{19}H_{11}NO_6$ 349.299

Same struct. given for Hypecoumine and Decumbenine C: they appear to be identical. Alkaloid from *Hypecoum leptocarpum* (Hypecoaceae) and *Corydalis decumbens* (Papaveraceae). Cryst. Mp 202-204° (*Hypecoumine*) Mp 211-213° (*Decumbenine C*). $[\alpha]_D^{25} +45.1$ (c, 0.07 in $CHCl_3$).

- Chen, B. *et al.*, *Yaouxue Xuebao*, 1985, **20**, 658-661; *CA*, **104**, 65907m
- Zhang, J.-S. *et al.*, *Phytochemistry*, 1995, **39**, 435-437 (*Decumbenine C*)

Hypecumine

[102719-88-0]




$C_{23}H_{22}N_2O_6$ 422.437

Alkaloid from *Hypecoum procumbens* (Hypecoaceae).

- Önür, M.A. *et al.*, *Planta Med.*, 1986, **70** (*isol, uv, pmr, ms, struct*)
- Fitzgerald, J.J. *et al.*, *Tet. Lett.*, 1994, **35**, 9191 (*synth*)

Hyperaspine

[356547-86-9]



Absolute Configuration

$C_{19}H_{30}N_2O_3$ 334.458


Alkaloid from the ladybird beetle *Hyperaspis campestris*. λ_{max} 263 (ε 3650) (no solvent reported).

- Lebrun, B. *et al.*, *Tet. Lett.*, 2001, **42**, 4621-4623 (*isol, pmr, cmr*)
- Zhu, W. *et al.*, *Org. Lett.*, 2003, **5**, 5063-5066 (*synth*)
- Dooms, C. *et al.*, *Eur. J. Org. Chem.*, 2005, 1378-1383 (*synth, abs config*)
- Comins, D.L. *et al.*, *Org. Lett.*, 2005, **7**, 5227-5228 (*synth*)
- Krishna, P.R. *et al.*, *Tet. Lett.*, 2007, **48**, 6924-6927 (*synth*)

Hypercratine

H-770

1-[2,3-Dihydro-2-(4-hydroxy-3-methoxyphenyl)-3-(2-pyrrolidinyl)-1,4-benzodioxin-6-yl]-2-(1-pyrrolidinyl)ethanone, 9CI [88607-22-1]




$C_{25}H_{30}N_2O_5$ 438.522

Alkaloid from the roots of *Ruspolia hypercrateriformis* (Acanthaceae). Opt. inactive.

- Neukomm, G. *et al.*, *Planta Med.*, 1983, **48**, 246 (*isol, uv, ir, pmr, cmr, ms, struct*)

Hyperectine

[94656-46-9]



Relative Configuration

$C_{24}H_{21}N_3O_6$ 447.446

Alkaloid from *Hypecoum erectum* and *Hypecoum leptocarpum* (Hypecoaceae). Mp 237-238°.

1-Epimer: Isohyperectine

[170384-75-5]

$C_{24}H_{21}N_3O_6$ 447.446


Alkaloid from *Hypecoum erectum* and *Hypecoum leptocarpum*. Cryst. ($CHCl_3/MeOH$). Mp 239-240°.

- Perel'son, M.E. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 628-635; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 592-598
- Yakhontova, L.D. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 835-839; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 744-747 (*Isohyperectine*)
- Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 1813-1816 (*isol, uv, pmr, cmr*)

Hypodemapyrazine

H-772

2,3-Dihydro-4-oxa-1,3a-diazacyclohepta[def]fluoren-10(1H)-one, 9CI [151936-22-0]



$C_{13}H_{10}N_2O_2$ 226.234


Alkaloid from the fern *Hypodematium sinense* (Woodsiaceae).

- Lu, Y. *et al.*, *Chin. Chem. Lett.*, 1993, **4**, 609-610; *CA*, **120**, 73332r (*isol, ir, uv, pmr, ms*)

Hypoestestatin 2

H-773

[97415-40-2]



$C_{24}H_{27}NO_4$ 393.482

Alkaloid from *Hypoestes verticillaris* (Acanthaceae). Inhibits cell growth. Highly cytotoxic. Sol. $MeOH, CHCl_3$; poorly sol. H_2O . Mp 206-212°. $[\alpha]_D^{25} -80$ (c, 0.525 in CH_2Cl_2). λ_{max} 259 (ε 41600); 285 (ε 25700); 340 (ε 1600) ($MeOH$) (Berdy).

Deoxy: Hypoestestatin 1

[97387-93-4]

$C_{24}H_{27}NO_3$ 377.482

Alkaloid from *Hypoestes verticillaris* (Acanthaceae). Cell growth inhibitor. Highly cytotoxic. Pale yellow solid. Sol. $MeOH, CHCl_3$; poorly sol. H_2O . Mp 191-197°. $[\alpha]_D^{25} -36.4$ (c, 0.55 in CH_2Cl_2). Extremely sensitive to air and acid. λ_{max} 258 (ε 28100); 286 (ε 15500); 342 (ε 870); 360 (ε 510) ($MeOH$) (Berdy).

- Pettit, G.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 913-919 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

Hypotuberostemonine


H-774

Struct. unknown

Minor alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Mp 183°. $[\alpha]_D^{19} +63.22$ (MeOH).Kondo, H. *et al.*, *CA*, 1957, **51**, 1540f; *Chem. Zentralbl.*, 1958, **129**, 14057 (*isol*)Alkaloid from the leaves of *Buxus hyrcana*. Amorph. solid. $[\alpha]_D^{24} +15.8$ (c, 0.13 in CHCl₃). λ_{\max} 225 (log ϵ 4.2) (MeOH).Choudhary, M.I. *et al.*, *Chem. Biodiversity*, 2006, **3**, 1039-1052 (*isol*)**Hyrtiomanzamine**


H-781

[177988-05-5]

C₁₈H₁₇N₄O₂S[⊕] 353.424Related to Dragmacidonamine A, D-933. Counterion not specified. Charge is delocalised. Alkaloid from the marine sponge *Hyrtios erecta*. Shows immunosuppressive activity. Orange glassy solid. λ_{\max} 394 (MeOH) (Berdy).**Deoxo: Gesashidine A**C₁₈H₁₉N₄OS[⊕] 339.44Alkaloid from an unidentified sponge of the family Thorectidae. Amorph. yellow solid. λ_{\max} 212 (ε 25000); 246 (ε 7300); 289 (ε 4400); 297 (ε 5600); 364 (ε 1400) (MeOH).Bourguet-Kondracki, M.L. *et al.*, *Tet. Lett.*, 1996, **37**, 3457-3460 (*isol*, *pmr*, *cmr*)Iinuma, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1109-1110 (*Gesashidine A*)**Hyperpine**


H-775

[312487-95-9]

C₁₉H₂₅NO₇ 379.409Alkaloid from the bark of *Hyperpa neocaledonica*.Montagnac, A. *et al.*, *CA*, 2001, **134**, 27525x (*isol*)**Hyrtiazepine**


H-778

3,4-Dihydro-7-hydroxy-6-(5-hydroxy-1H-indol-3-yl)-1H-azepino[5,4,3-cd]indole-4-carboxylic acid [922495-16-7]

C₂₀H₁₅N₃O₄ 361.356Alkaloid from *Hyrtios erectus*. Yellow solid. $[\alpha]_D^{25} -16.9$ (c, 0.13 in MeOH). λ_{\max} 227 (log ϵ 4.22); 351 (log ϵ 3.73); 435 (log ϵ 3.72) (EtOH).Sauleau, P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1676-1679 (*isol*, *pmr*, *cmr*)**Hyrtinadine A**


H-779

[925253-33-4]

C₂₀H₁₄N₄O₂ 342.356Isol. from a *Hyrtios* sp. (SS-1127). Cytotoxic. Amorph. solid. λ_{\max} 277 (ε 7600); 339 (ε 3100) (MeOH).Endo, T. *et al.*, *J. Nat. Prod.*, 2007, **70**, 423-424 (*isol*, *pmr*, *cmr*)**Hyrtiosin B[†]**


H-782

Bis(5-hydroxy-1H-indol-3-yl)ethanedione, 9CI [132922-99-7]


C₁₈H₁₂N₂O₄ 320.304Alkaloid from the marine sponge *Hyrtios erecta*. Pale green solid. Mp > 400. λ_{\max} 212 (ε 30000); 255 (ε 12800); 290 (ε 12100); 332 (ε 11000) (MeOH) (Berdy).**Dideoxy: Di-1H-indol-3-ylethanedione, 9CI. Di-3-indolylglyoxal** [65610-87-9]C₁₈H₁₂N₂O₂ 288.305Alkaloid from the sponge *Smenospongia* sp. Yellow cryst. (Me₂CO aq.). Mp 279-280°. λ_{\max} 247 (log ϵ 4.28); 266 (log ϵ 4.27); 326 (log ϵ 4.27) (no solvent reported).Bergman, J. *et al.*, *Tetrahedron*, 1990, **46**, 6061-6066 (*Bisindolyethanedione, synth*)Kobayashi, J. *et al.*, *Tetrahedron*, 1990, **46**, 7699-7702 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)Bergman, J. *et al.*, *Synthesis*, 1999, 580-585 (*synth*, *ir*, *pmr*, *cmr*)McKay, M.J. *et al.*, *J. Nat. Prod.*, 2002, **65**, 595-597 (*Bisindolyethanedione*)Krasnyushkin, M.M. *et al.*, *Zh. Org. Khim.*, 2005, **41**, 895-902; *Russ. J. Org. Chem. (Engl. Transl.)*, 2005, **41**, 875-883 (*dideoxy, synth*)**Hyrcanine**

H-776

[220749-23-5]


C₂₆H₃₉NO 381.6Alkaloid from *Buxus hyrcana*. Needles (Me₂CO). Mp 118°. $[\alpha]_D^{25} -33$ (c, 0.09 in CHCl₃). λ_{\max} 236 (log ϵ 4); 243 (log ϵ 4.01) (MeOH).Atta-ur-Rahman, *et al.*, *Heterocycles*, 1998, **49**, 481-488 (*isol*, *pmr*, *cmr*, *uv*)**Hyrtioerectine A**

H-780

C₂₁H₁₃N₃O₅ 387.351Alkaloid from the sponge *Hyrtios erectus*. Amorph. yellow solid. λ_{\max} 387 (log ϵ 4.05) (MeOH).Youssef, D.T.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1416-1419 (*isol*, *pmr*, *cmr*)**Hyrcanone**


H-777

[934386-81-9]

C₃₃H₄₈N₂O₂ 504.754

Hysodricanine A
[64309-17-7]**H-783**


Tschesche, R. *et al.*, *Phytochemistry*, 1977, **16**,
1025-1028 (*isol, uv, ir, pmr, ms, struct*)
Tschesche, R. *et al.*, *Pharmazie*, 1981, **36**, 511
(*isol, uv, ir, pmr, ms*)

C₃₅H₄₅N₅O₅ 615.771

Alkaloid from the bark of *Zizyphus hysodrica* and *Zizyphus hutchinsonii* (Rhamnaceae). Needles (MeOH/petrol). Mp 93-96°. [α]_D²⁰ -215 (c, 0.05 in CHCl₃).

Dihydro:


Cryst. (MeOH aq.). Mp 130-132°. [α]_D²⁰ -186 (c, 0.03 in MeOH).

Hysodricanine B**H-784**C₃₀H₄₃N₅O₆ 569.7

Cyclic peptide alkaloid. Constit. of *Zizyphus hysodrica*. Cryst. (CHCl₃/cyclohexane). Mp 35°. λ_{\max} 360 ; 395 (CHCl₃).

Hysodricanine C**H-785**

[220376-45-4]


C₃₀H₄₃N₅O₆ 569.7

Cyclic peptide alkaloid. Constit. of *Zizyphus hysodrica*. Cryst. (CHCl₃/cyclohexane). Mp 35°.

Khokhar, I. *et al.*, *J. Nat. Sci. Math.*, 1996, **36**, 149-154; *CA*, **130**, 165491b (*isol*)

Ianthelline

[105596-36-9]


C₁₅H₁₇Br₂N₃O₃ 475.139

Amino acid-derived antibiotic. Isol. from the sponge *Ianthella ardis* and *Pseudoceratina crassa*. Shows antibacterial and antifungal props. Pale yellow powder. Mp 113-115°.

Litaudon, M. *et al.*, *Tet. Lett.*, 1986, **27**, 4455 (isol, ir, pmr, cmr, ms, struct)

Ianthesine B

I-2

C₂₂H₂₃Br₄N₃O₇ 761.056

Alkaloid from the marine sponge *Ianthella* sp. Cryst. (MeOH aq.). Mp 154-157°. [α]_D²² -97 (c, 0.58 in MeOH). λ_{max} 207 (ε 45400); 283 (ε 7300) (MeOH).

20-Epimer, N²⁰-di-Me: **Ianthesine A**

C₂₄H₂₇Br₄N₃O₇ 789.109

Alkaloid from *Ianthella* sp. Cryst. (MeOH aq.). Mp 154-156°. [α]_D²² -118 (c, 1 in MeOH). λ_{max} 210 (ε 29500); 283 (ε 7390) (MeOH).

N²⁰-Sulfo: **Ianthesine D**


C₂₂H₂₃Br₄N₃O₁₀S 841.12

Alkaloid from *Ianthella* sp. Powder (as Na salt). Mp 190° dec. (Na salt). [α]_D²⁵ -69 (c, 0.19 in MeOH) (Na salt). C-20 config. not determined. λ_{max} 206 (ε 53200); 220 (sh) (ε 25000); 282 (ε 6760) (MeOH) (Na salt).

Okamoto, Y. *et al.*, *Tetrahedron*, 2000, **56**, 5813-5818

Ianthesine C

I-3

C₄₄H₄₄Br₈N₆O₁₆S 1584.16


Alkaloid from the marine sponge *Ianthella* sp. Yellow powder (DMSO aq.) (as Na salt). Mp 200° dec. (Na salt). [α]_D²⁶ -93 (c, 0.86 in DMSO) (Na salt). λ_{max} 231 (ε 26900); 283 (ε 11900) (MeOH/DMSO) (Na salt).

Okamoto, Y. *et al.*, *Tetrahedron*, 2000, **56**, 5813-5818

Ibogamine, 9CI

[481-87-8]

[2288-55-3 (+/-)-form]



Absolute Configuration

C₁₉H₂₄N₂ 280.412

Alkaloid from *Tabernanthe iboga* and several other *Tabernanthe* spp., *Pandaca retusa*, *Stemmadenia galeottiana* and *Stemmadenia glabra* (Apocynaceae). Shows moderate to weak cytotoxic activity and weak antibacterial activity. Also shows some CNS, bradycardic and hypotensive activity. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 162-164°. [α]_D¹⁸ -54.7 (c, 1 in EtOH). Log P 4.03 (uncertain value) (calc). λ_{max} 225 (log ε 4.58); 285 (log ε 3.96); 293 (log ε 3.94) (MeOH).

Hydrochloride: Mp 247-250°.

19R-Hydroxy: **19R-Hydroxyibogamine**

[73543-09-6]

C₁₉H₂₄N₂O 296.411

Alkaloid from the roots of *Tabernanthe quadrangularis* and *Pandaca mocquerysii* var. *pendula* (Apocynaceae). Needles (MeOH). Mp 172-174°. [α]_D²⁰ -28 (c, 1 in CHCl₃). λ_{max} 227 (log ε 4.49); 276 (sh) (log ε 3.73); 283 (log ε 3.89); 291 (log ε 3.81) (MeOH).

19S-Hydroxy: **19S-Hydroxyibogamine**

[437764-31-3]

C₁₉H₂₄N₂O 296.411

Alkaloid from *Tabernaemontana corymbosa* and *Pandaca mocquerysii* var. *pendula*. Pale yellow oil. [α]_D -21 (c, 0.16 in CHCl₃). λ_{max} 226 (log ε 4.28); 283 (log ε 3.66); 292 (log ε 3.62) (EtOH).

10-Methoxy: **Ibogaine**. Endabuse

[83-74-9]

[6516-52-5 (+/-)-form]

C₂₀H₂₆N₂O 310.438

Major alkaloid from *Tabernanthe iboga* (5-6% in root bark), also from *Tabernanthe pubescens*, *Tabernaemontana crassa*, *Voacanga thouarsii* and several other spp. in the Apocynaceae. Strong CNS stimulant, hallucinogen, anticonvulsant agent. Possesses antihypertensive props. *T. iboga* is an African folk remedy against fatigue. Sigma 2-receptor ligand and serotonergic agent. Under investigation for treatment of withdrawal symptoms from drugs of abuse. Shows antimycobacterial activity. Sol. org. solvs. Mp 152-153°. [α]_D -53 (EtOH). Log P 4.05 (uncertain value) (calc). Dec. by heat and light. λ_{max} 228 (log ε 4.4); 300 (log ε 3.92) (MeOH).

10-Methoxy, hydrochloride: [5934-55-4]

Mp 299-300° dec. [α]_D -63 (EtOH). [α]_D -49 (H₂O).

▶ LD₅₀ (rat, orl) 482 mg/kg. NH6730000

10-Methoxy, 19R-hydroxy: **19-Epiibogaine**

[17378-49-3]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from leaves of *Tabernaemontana dichotoma*. λ_{max} 226 (log ε 4.32); 288 (log ε 3.79) (EtOH).

10-Methoxy, 19S-hydroxy: **Iboxygaine**.

Kimvuline. Decarbomethoxyvoacangarine

[82-55-3]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Tabernanthe iboga*, *Daturicarpa elliptica* (preferred genus name *Tabernanthe*), *Pandaca speciosa* and several other spp. in the Apocynaceae. Shows weak antibacterial and some CNS, bradycardial and hypotensive activity. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 235-236°. [α]_D²⁰ -10 (EtOH). [α]_D²⁰ -15 (c, 0.41 in CHCl₃). Log P 1.84 (uncertain value) (calc). λ_{max} 228; 288 (MeOH) (Berdy).

11-Methoxy: **Tabernanthine**. 11-Methoxyibogamine. 13-Methoxyibogamine

[83-94-3]

C₂₀H₂₆N₂O 310.438

Alkaloid from *Tabernanthe iboga*, *Tabernanthe laurifolia* and *Stemmadenia donnell-smithii* (Apocynaceae). Shows some CNS, bradycardial and hypotensive activity. Active against gram-positive bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 211-212° (209°). [α]_D²⁵ -35 (CHCl₃). [α]_D -40 (Me₂CO). Log P 4.05 (uncertain value) (calc). λ_{max} 229 (ε 35970); 271 (ε 4410); 299 (ε 5810) (MeOH) (Berdy).

11-Methoxy, 3RS-hydroxy: **3-Hydroxytabernanthine**

[524729-87-1]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from *Tabernaemontana calcaria*. Oil. [α]_D -17 (c, 0.24 in CHCl₃). λ_{max} 222 (log ε 4.02); 286 (log ε 3.68); 294 (log ε 3.26) (MeOH).

10,11-Dimethoxy: **Ibogaline**. Descarbomethoxyconopharyngine

[482-18-8]

C₂₁H₂₈N₂O₂ 340.464

Alkaloid from *Tabernanthe iboga* bark (Apocynaceae). Hypotensive and bradycardic agent. CNS stimulant. Cryst. (MeOH aq.). Mp 141-143°. Log P 3.5 (uncertain value) (calc).

10,11-Dimethoxy; hydrochloride: Mp 264-266° dec.

Delourme-Houdé, J. *et al.*, *Ann. Pharm. Fr.*, 1946, **4**, 30-36; *CA*, **41**, 1390d (Tabernanthine)

Burckhardt, C.A. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 642-643 (isol)

Goutarel, M. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 742-748 (uv, ir)

Goutarel, R. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1958, **246**, 279-281 (Iboxygaine, isol, uv, ir)

Bartlett, M.F. *et al.*, *J.A.C.S.*, 1958, **80**, 126-136 (Ibogamine, Tabernanthine, struct)

Walls, F. *et al.*, *Tetrahedron*, 1958, **2**, 173-182 (Tabernanthine)


Neuss, N. *et al.*, *J.O.C.*, 1959, **24**, 2047-2048 (Ibogaline)

- Biemann, K. *et al.*, *J.A.C.S.*, 1961, **83**, 4805 (*Ibogaine, Iboxygaine, ms, struct*)
- Binst, G.V. *et al.*, *Tet. Lett.*, 1964, 973 (*pmr*)
- Büchi, G. *et al.*, *J.A.C.S.*, 1966, **88**, 3099-3109 (*Ibogaine, synth*)
- Hirai, S. *et al.*, *Chem. Comm.*, 1968, 1016-1017 (*synth*)
- Thomas, D.W. *et al.*, *J. Nat. Prod.*, 1968, **31**, 1-8 (*ms*)
- Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 157-176 (*Tabernanthe, activity*)
- Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 2258-2266 (*Iboxygaine, abs config*)
- Rosenmund, P. *et al.*, *Chem. Ber.*, 1975, **108**, 1871-1895 (*Ibogamine, Ibogaine, synth, ms*)
- DeBellefont, M. *et al.*, *Phytochemistry*, 1975, **14**, 1649-1652 (*19-Hydroxyibogamines*)
- Harmouche, A. *et al.*, *Ann. Pharm. Fr.*, 1976, **34**, 31-35 (*pmr*)
- Wenkert, E. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2437 (*cmr*)
- Damak, M. *et al.*, *Tet. Lett.*, 1976, 3531-3534 (*cmr*)
- Trost, B.M. *et al.*, *J.A.C.S.*, 1978, **100**, 3930-3931 (*synth*)
- Achenbach, H. *et al.*, *Z. Naturforsch., B*, 1980, **35**, 219-225 (*19-Hydroxyibogamine*)
- Perera, P. *et al.*, *Planta Med.*, 1983, **47**, 148-150 (*19-Epiiboxygaine*)
- Imanishi, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4202-4211 (*synth, ir, pmr, ms*)
- Van Beek, T.A. *et al.*, *J. Ethnopharmacol.*, 1985, **14**, 315-318; *CA*, **104**, 106242 (*Ibogamine, activity*)
- Huffmann, J.W. *et al.*, *J.O.C.*, 1985, **50**, 1460-1464 (*synth*)
- Kuehne, M.E. *et al.*, *J.O.C.*, 1985, **50**, 1464-1467 (*synth*)
- Danieli, B. *et al.*, *Alkaloids (Academic Press)*, 1986, **127**, 1-130 (*rev, pharmacol*)
- Herdeis, C. *et al.*, *Annalen*, 1991, 99-104 (*synth*)
- Soriano-García, M. *et al.*, *Acta Cryst. C*, 1992, **48**, 2055-2057 (*cryst struct*)
- Bornmann, W.G. *et al.*, *J.O.C.*, 1992, **57**, 1752-1760 (*Ibogamine, synth*)
- Sershen, H. *et al.*, *Neurochem. Res.*, 1994, **19**, 1463-1466 (*Ibogaine, pharmacol*)
- Gallagher, C.A. *et al.*, *Biochem. Pharmacol.*, 1995, **49**, 73-79 (*Ibogaine, gc-ms*)
- Mach, R.H. *et al.*, *Life Sci.*, 1995, **57**, PL57-PL62 (*Ibogaine, pharmacol*)
- Henry, K.J. *et al.*, *Tet. Lett.*, 1996, **37**, 8289-8292 (*Ibogamine, synth*)
- Rastogi, N. *et al.*, *FEMS Immunol. Med. Microbiol.*, 1998, **20**, 267-273 (*Ibogaine, activity*)
- Alper, K.R. *et al.*, *Alkaloids (Academic Press)*, 2001, **56**, 1-38 (*Ibogaine, !rev*)
- White, J.D. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 4306-4327 (*Ibogamine, synth*)
- Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 669-672 (*19S-Hydroxyibogamine*)
- Chaturvedula, V.S.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 528-531 (*3-Hydroxytabernanthe*)
- Höck, S. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 542-557 (*19R-Hydroxyibogamine, synth*)

Iboluteine

I-5

[468-11-1]



Absolute Configuration

C₂₀H₂₆N₂O₂ 326.438Alkaloid from *Tabernanthe iboga*, *Tabernanthe montana* spp., *Pandaca ochracea*, *Pandaca speciosa* and *Rejoua aurantiaca*

(preferred genus name *Tabernaemontana*) (Apocynaceae). Mp 142°. [α]_D²⁵ -114 (CHCl₃). Also an autoxidn. prod. of Ibogaine in I-4 therefore poss. artifact. Wenkert shows the incorr. abs. config.

Oxime: Mp 293-294°.

Demethoxy: Desmethoxyiboluteine. Ibo-gamine pseudoindoxyl
[73777-64-7]

C₁₉H₂₄N₂O 296.411

Alkaloid from *Tabernaemontana iboga* and from the roots of *Tabernaemontana quadrangularis* (Apocynaceae). Cryst. (C₆H₆ or MeOH). Mp 141° (137-138°). [α]_D²⁰ -62 (c, 0.2 in CHCl₃). Prob. artifact of autoxidn.

Demethoxy, 19R-hydroxy: 19R-Hydroxyibogamine pseudoindoxyl. 20R-Hydroxyibogamine pseudoindoxyl
[73459-04-8]


C₁₉H₂₄N₂O₂ 312.411

Alkaloid from the roots of *Tabernaemontana quadrangularis* (Apocynaceae). Yellow oil. [α]_D²⁰ -16 (c, 0.1 in CHCl₃).

Goutarel, R. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 742 (*ir, uv, synth*)Dickel, D.F. *et al.*, *J.A.C.S.*, 1958, **80**, 123 (*struct*)Thomas, D.W. *et al.*, *Tetrahedron*, 1968, **24**, 4223 (*ms, uv, ir*)Wenkert, E. *et al.*, *Heterocycles*, 1977, **7**, 753 (*cmr, config*)Achenbach, H. *et al.*, *Z. Naturforsch., B*, 1980, **35**, 219 (*Ibogamine pseudoindoxyl, 19-Hydroxyibogamine pseudoindoxyl*)**Ibophyllidine**

I-6

Methyl 2,3-didehydro-7-ethyl-E,20,21-trinoraspidospermidine-3-carboxylate, 9CI
[59871-84-0]
[78019-50-8 (±)-form]



Absolute Configuration

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from the leaves of *Tabernanthe iboga* and *Tabernanthe subsessilis*, the stems of *Tabernaemontana flavicans* and the bark of *Tabernaemontana albiflora* and *Daturicarpa elliptica* (preferred genus name *Tabernanthe*) (Apocynaceae). Noncryst. [α]_D +134 (c, 1 in CHCl₃).

N^b-Oxide: Ibophyllidine N^b-oxide

[76190-14-2]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the stems of *Tabernaemontana flavicans* (Apocynaceae). Oil. [α]_D²⁰ +187 (CHCl₃).

20-Epimer: 20-Epiibophyllidine

[74170-68-6]

[78019-52-0 (±)-form]


C₂₀H₂₄N₂O₂ 324.422

Alkaloid from the bark of *Tabernaemontana albiflora* (Apocynaceae). Amorph. [α]_D²⁰ +518 (c,

0.3 in CHCl₃) (natural). [α]_D²⁰ +322 (c, 1.3 in CHCl₃) (natural). [α]_D²⁰ +268 (c, 1 in CHCl₃) (synthetic).

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1976, **32**,2539 (*isol, uv, ir, pmr, cmr, ms, struct*)Kan, C. *et al.*, *Tet. Lett.*, 1980, **21**, 55 (*uv, ir, pmr, ms, struct, epimer*)Achenbach, H. *et al.*, *Z. Naturforsch., B*, 1980,**35**, 1465 (*isol, uv, ir, pmr, ms, cd, oxide, synth*)Kuehne, M.E. *et al.*, *J.O.C.*, 1981, **46**, 3443(*synth, uv, ir, pmr, ms, epimer*)Barsi, M.-C. *et al.*, *Chem. Comm.*, 1985, 88(*synth*)Jegham, S. *et al.*, *Tet. Lett.*, 1989, **30**, 1959(*synth*)Bornmann, W.G. *et al.*, *J.O.C.*, 1992, **57**, 1752(*synth*)**Iboquine**

I-7


C₂₀H₂₄N₂O₂ 324.422

Alkaloid from roots of *Tabernanthe iboga* (Apocynaceae). Mp 284°. Prob. artifact of autoxidn. of Ibogaine in I-4.

Goutarel, R. *et al.*, *Ann. Pharm. Fr.*, 1953, **11**, 272 (*isol*)Dickel, D.F. *et al.*, *J.A.C.S.*, 1958, **80**, 123**Ibotenic acid**

I-8

α-Amino-2,3-dihydro-3-oxo-5-isoxazoleacetic acid, 9CI. Premuscimol
[2552-55-8]

C₅H₆N₂O₄ 158.113

(±)-form [60573-88-8]

Prod. by *Amanita muscaria*, *Amanita pantherina*, *Amanita strobiliformis* and *Tricholoma muscarium*. Narcotizing agent, CNS depressant, psychomimetic agent, pesticide. Aspartic acid receptor agonist. Cryst. + 1H₂O. Sol. H₂O. Mp 151-152° dec. pK_{a1} 2; pK_{a2} 5.1; pK_{a3} 8.2. ▶ LD₅₀ (mus, ivn) 15 mg/kg. NY2100000

Et ester: [127020-25-1]C₇H₁₀N₂O₄ 186.167

Solid (EtOH/Et₂O) (as hydrochloride). Mp 184-186° dec. (hydrochloride).

Et ether, Et ester:C₉H₁₄N₂O₄ 214.221

Solid (EtOH/Et₂O) (as hydrochloride). Mp 172-174° dec. (hydrochloride).

[25690-45-3, 31758-99-3]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 636D (*ir*)


Gagneux, A.R. *et al.*, *Tet. Lett.*, 1965, 2081(*synth*)Yokobe, T. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1236 (*ms*)Nakamura, N. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 46 (*synth*)

- Cilton, W.S. *et al.*, *J. Nat. Prod.*, 1976, **39**, 150 (occur)
 Leslie, D.T. *et al.*, *J. Pharm. Sci.*, 1978, **67**, 485 (glc, ms)
 Madsen, U. *et al.*, *Acta Chem. Scand.*, 1990, **44**, 96 (deriv)
 Ohta, T. *et al.*, *Nat. Med. (Tokyo)*, 1995, **49**, 354 (isol, uv, ir, pmr, cmr)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AKG250

Iboxygaine hydroxyindolenine

I-9

[23627-68-1]



Absolute Configuration


$C_{20}H_{26}N_2O_3$ 342.437
 Alkaloid from the leaves, stems and bark of *Peschiera lundii* (Apocynaceae). Mp 223°. $[\alpha]_D^{25}$ +111 (c, 0.026 in $CHCl_3$).

Hwang, B. *et al.*, *J.O.C.*, 1969, **34**, 412 (isol, uv, ir, struct, synth)

Iboxyphylline

I-10

Methyl 2,3-didehydro-7-hydroxy-8-methyl-E-homo-20,21-dinoraspidospermidine-3-carboxylate, 9CI
 [62861-08-9]



Absolute Configuration

$C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the leaves of *Tabernanthe iboga* and *Tabernanthe subsessilis*. Cryst. + $1H_2O$ (Et_2O or $MeCN$). Mp 245°. $[\alpha]_D$ +444 (c, 1 in $CHCl_3$).

O-Ac:

Cryst. (MeOH). Mp 167°. $[\alpha]_D$ +466 (c, 1 in $CHCl_3$).

Khuong-Huu, F. *et al.*, *Tetrahedron*, 1976, **32**, 2539-2543 (isol, uv, ir, pmr, cmr, ms, cryst struct)


Bornmann, W.G. *et al.*, *J.O.C.*, 1992, **57**, 1752-1760 (synth)

Tóth, F. *et al.*, *Heterocycles*, 2008, **75**, 65-76 (synth)

Icaceine

I-11

17-(Dimethylamino)-3,6-dihydroxy-4-methyl-15-oxaandrost-7-ene-4-carboxylic acid, 9CI
 [74991-71-2]

 $C_{22}H_{33}NO_4$ 375.507

First known pimarane-derived diterpene alkaloid. Stereochem. not fully determined but prob. the same as that of Icacine. Alkaloid from the leaves and roots of *Icacina guesfeldtii* (Icacinaceae). Needles ($MeOH/CHCl_3$). Mp 220-250° dec., subl. $[\alpha]_D^{20}$ -196.7 (c, 0.995 in $CHCl_3$).

Ac:

Needles. Mp 190-215° dec. subl. $[\alpha]_D$ -180.3 (c, 0.295 in $CHCl_3$).

N-De-Me: **De-N-methylicaceine**

[75008-99-0]

 $C_{21}H_{31}NO_4$ 361.48


Alkaloid from *Icacina guesfeldtii* (Icacinaceae). Mp 210-235° dec., subl. $[\alpha]_D^{20}$ -165.8 (c, 0.535 in $CHCl_3$).

On'okoko, P. *et al.*, *Phytochemistry*, 1980, **19**, 303-305 (isol, ir, pmr)

Icajine

I-12

19-Methyl-16,19-secostrychnidine-10,16-dione, 9CI. N-Methyl-sec-pseudostrychnine
 [5525-31-5]



Absolute Configuration

 $C_{22}H_{24}N_2O_3$ 364.443

Several numbering schemes in the lit. Only derivative synonyms using the biogenetic scheme favoured in DNP are given here. Alkaloid from the leaves of *Strychnos icaja* and *Strychnos wallichiana* (Loganiaceae). Weak convulsive agent. Mp 271-272° dec. $[\alpha]_D$ -10 (c, 1 in $CHCl_3$). λ_{max} 253 (log ϵ 4.23); 291 (log ϵ 3.52) (EtOH).

N-Oxide: **Icajine N-oxide**

[52080-27-0]

 $C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from leaves of *Strychnos wallichiana* (Loganiaceae). Needles (MeOH). λ_{max} 210 (log ϵ 4.66); 255 (log ϵ 4.32); 283 (log ϵ 3.91); 289 (sh) (log ϵ 3.8) (EtOH).

N-De-Me, N-cyano: **N-Cyano-sec-pseudostrychnine**

[52080-24-7]

 $C_{22}H_{21}N_3O_3$ 375.426

Alkaloid from leaves of *Strychnos wallichiana*, also detected in *Strychnos ignatii* (Loganiaceae). Needles (Me_2CO). Mp 235° dec. λ_{max} 222 (log ϵ 4.62); 257 (log ϵ 4.76); 283 (log ϵ 4.25) (EtOH).

12-Hydroxy: **Vomicine**. *Strychnicine*.*Struxine*

[125-15-5]

 $C_{22}H_{24}N_2O_4$ 380.443Alkaloid from *Strychnos nux-vomica*

(leaves and seeds) and *Strychnos icaja* (Loganiaceae). Weak convulsive agent. Mp 250° dec. Mp 281-282° dec. $[\alpha]_D$ +100 (c, 1 in $CHCl_3$).

12-Hydroxy, hydrochloride: Mp 245° dec. (browns from 210°).

12-Hydroxy, 19 α ,20 α -epoxide: **19,20-Epoxyvomicine**. 19,20-Epoxy-12-hydroxy-N-methyl-sec-pseudostrychnine
 [22029-94-3]

 $C_{22}H_{24}N_2O_5$ 396.442

Alkaloid from leaves of *Strychnos icaja* (Loganiaceae). Small needles (MeOH). Mp 252-255° dec. $[\alpha]_D$ +112 (c, 1 in $CHCl_3$).

▶ ZB2505000

15-Hydroxy: **15-Hydroxyicajine**. 15-Hydroxy-N-methyl-sec-pseudostrychnine
 [22029-96-5]

 $C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from leaves of *Strychnos icaja* and *Strychnos wallichiana* (Loganiaceae). Needles (MeOH). Mp 262-265° dec. $[\alpha]_D$ -60 (c, 1.1 in $CHCl_3$).

▶ VS3615000

15-Hydroxy, 19 α ,20 α -epoxide: **19,20-Epoxy-15-hydroxyicajine**. 19,20-Epoxy-15-hydroxy-N-methyl-sec-pseudostrychnine
 [22029-99-8]

 $C_{22}H_{24}N_2O_5$ 396.442

Alkaloid from the leaves of *Strychnos icaja* (Loganiaceae). Rods (MeOH). Mp 266-268° dec. $[\alpha]_D$ +3 (c, 1 in $CHCl_3$).

12,15-Dihydroxy, 19 α ,20 α -epoxide: **19,20 α -Epoxy-15-hydroxyvomicine**.

19,20-Epoxy-12,15-dihydroxy-N-methyl-sec-pseudostrychnine
 [22029-97-6]

 $C_{22}H_{24}N_2O_6$ 412.441

Alkaloid from leaves of *Strychnos icaja* (Loganiaceae). Rods (Me_2CO). Mp 239-242°. $[\alpha]_D$ +86 (c, 1.01 in $CHCl_3$).

10-Methoxy: **N-Methyl-sec-pseudo- β -colubrine**

[52080-22-5]

 $C_{23}H_{26}N_2O_4$ 394.469

Alkaloid from fruits of *Strychnos nux-vomica* and leaves of *Strychnos wallichiana* (Loganiaceae). Needles (Me_2CO). A 10- or 11-methoxy (see below) icajine was isol. from leaves of *S. icaja*. λ_{max} 226 (log ϵ 4.38); 261 (log ϵ 3.95); 300 (log ϵ 3.92) (EtOH).

10-Methoxy, 19 α ,20 α -epoxide: **19,20-Epoxy-10-methoxyicajine**. 19,20-Epoxy-10-methoxy-N-methyl-sec-pseudostrychnine
 [27239-52-7]

 $C_{23}H_{26}N_2O_5$ 410.469

Alkaloid from leaves of *Strychnos icaja* (Loganiaceae). Flat prisms (MeOH). Mp 265-267° dec.

11-Methoxy: **11-Methoxyicajine**

[73710-81-3]

 $C_{23}H_{26}N_2O_4$ 394.469

Alkaloid from *Strychnos nux-vomica* (Loganiaceae). Mp 240-242°.

11-Methoxy, 12-hydroxy: **12-Hydroxy-11-methoxy-N-methyl-sec-pseudostrychnine**


- [50837-33-7]
 $C_{23}H_{26}N_2O_5$ 410.469
 Alkaloid from seeds of *Strychnos wallichiana* (*Strychnos cinnamomifolia*) (Loganiaceae). $[\alpha]_D^{20} +33$ (c, 0.24 in $CHCl_3$); λ_{max} 232; 270; 301 (EtOH).
- 11-Methoxy, 12-hydroxy, 19 α ,20 α -epoxide: 19,20-Epoxy-11-methoxyvomicine.** 19,20-Epoxy-12-hydroxy-11-methoxy-N-methyl-sec-pseudostrychnine
 [27281-86-3]
 $C_{23}H_{26}N_2O_6$ 426.468
 Alkaloid from leaves of *Strychnos icaja* (Loganiaceae). Noncryst. $[\alpha]_D^{20} +78$ (c, 1.05 in $CHCl_3$).
- 11-Methoxy, 12,15-dihydroxy, 19 α ,20 α -epoxide: 19,20-Epoxy-15-hydroxy-11-methoxyvomicine.** 19,20-Epoxy-12,15-dihydroxy-11-methoxy-N-methyl-sec-pseudostrychnine
 [62421-66-3]
 $C_{23}H_{26}N_2O_7$ 442.468
 Alkaloid from *Strychnos icaja* leaves (Loganiaceae). Prisms (EtOAc). Mp 255-260°. λ_{max} 237 (log ϵ 4.21); 270 (log ϵ 3.8) (EtOH).
- 12-Methoxy, 19 α ,20 α -epoxide: 19,20-Epoxy-12-methoxyicajine.** 19,20-Epoxy-12-methoxy-N-methyl-sec-pseudostrychnine
 [62421-68-5]
 $C_{23}H_{26}N_2O_5$ 410.469
 Minor alkaloid from dried fruits of m'boundou (*Strychnos* sp.) from Gabon (Loganiaceae). Noncryst. λ_{max} 217 (log ϵ 4.29); 256 (log ϵ 4.11); 295 (log ϵ 3.85) (EtOH).
- 12-Methoxy, 15-hydroxy, 19 α ,20 α -epoxide: 19,20-Epoxy-15-hydroxy-12-methoxyicajine.** 19,20-Epoxy-15-hydroxy-12-methoxy-N-methyl-sec-pseudostrychnine
 [62421-69-6]
 $C_{23}H_{26}N_2O_6$ 426.468
 Major alkaloid from dried fruits of m'boundou (*Strychnos* sp.) (Loganiaceae). Noncryst. λ_{max} 217 (log ϵ 4.29); 256 (log ϵ 4.11); 295 (log ϵ 3.58) (EtOH).
- 10,11-Dimethoxy: Novacine.** N-Methyl-sec-pseudobrucine
 [466-64-8]
 [30864-62-1]
 $C_{24}H_{28}N_2O_5$ 424.496
 Alkaloid from *Strychnos nux-vomica* seeds, *Strychnos icaja* and *Strychnos wallichiana* leaves (Loganiaceae). Prisms (EtOAc). Mp 231-232°. $[\alpha]_D^{20} -17.7$ (c, 5 in $CHCl_3$). λ_{max} 215 (log ϵ 4.36); 266 (log ϵ 4.04); 300 (log ϵ 3.88) (EtOH).
- 10,11-Dimethoxy, hydrochloride:**
 Prismatic needles + H_2O (EtOH). Mp 229-230°.
- 10,11-Dimethoxy, N-de-Me, N-cyano: N-Cyano-sec-pseudobrucine**
 [52080-25-8]
 $C_{24}H_{25}N_3O_5$ 435.479
 Alkaloid from leaves of *Strychnos wallichiana* (Loganiaceae). Needles (Me_2CO).

- 10,11-Dimethoxy, 19,20 α -epoxide: 19,20-Epoxyvomicine.** 19,20-Epoxy-N-methyl-sec-pseudobrucine. Alkaloid C†
 [19879-54-0]
 $C_{24}H_{28}N_2O_6$ 440.495
 Alkaloid from the leaves of *Strychnos icaja* (Loganiaceae). Hexagonal plates (MeOH). Mp 239-241°. $[\alpha]_D^{20} +12$ (c, 1.09 in $CHCl_3$).
- 10,11-Dimethoxy, 15-hydroxy: 15-Hydroxyvomicine**
 [52080-26-9]
 $C_{24}H_{28}N_2O_6$ 440.495
 Alkaloid from leaves of *Strychnos wallichiana* (Loganiaceae). Fine needles (MeOH). Mp 200° dec. λ_{max} 217 (log ϵ 4.64); 265 (log ϵ 4.39); 301 (log ϵ 4.27) (EtOH).
- 10,11-Dimethoxy, 15-hydroxy, 19 α ,20 α -epoxide: 19,20-Epoxy-15-hydroxyvomicine.** 19,20-Epoxy-14-hydroxy-10,11-dimethoxy-N-methyl-sec-pseudostrychnine. Alkaloid B'
 [22029-98-7]
 $C_{24}H_{28}N_2O_7$ 456.494
 Alkaloid from leaves of *Strychnos icaja* (Loganiaceae). Rods (Me_2CO). Mp 251-253° dec. $[\alpha]_D^{20} +3$ (c, 0.96 in $CHCl_3$).
- 11,12-Dimethoxy, 19 α ,20 α -epoxide: 19,20-Epoxy-11,12-dimethoxyicajine.** 19,20-Epoxy-11,12-dimethoxy-N-methyl-sec-pseudostrychnine
 [62421-67-4]
 $C_{24}H_{28}N_2O_6$ 440.495
 Alkaloid from *Strychnos icaja* leaves (Loganiaceae). Noncryst. λ_{max} 226 (log ϵ 4.3); 256 (log ϵ 4.11); 295 (log ϵ 3.58) (EtOH).
- [19774-65-3]
 Wieland, H. et al., *Annalen*, 1929, **469**, 193-215 (*Vomicine, sol*)
 Leuchs, H. et al., *Ber.*, 1939, **72**, 965-972; 1940, **73**, 885-892 (*synth*)
 Martin, W.F. et al., *J.C.S.*, 1952, 3603-3604 (*Novacine*)
 Smith, G.F. et al., *Alkaloids (Academic Press)*, 1965, **8**, 591-671 (*rev, bibl*)
 Bisset, N.G. et al., *C. R. Hebd. Seances Acad. Sci.*, 1965, **261**, 5237-5238 (*isol, uv, ms*)
 Bisset, N.G. et al., *Tet. Lett.*, 1968, 3107-3110 (*Epoxyvomicine*)
 Bisset, N.G. et al., *J. Pharm. Pharmacol.*, 1973, **25**, 563-569 (*12-Hydroxy-11-methoxy-N-methyl-sec-pseudostrychnine*)
 Bisset, N.G. et al., *Tetrahedron*, 1973, **29**, 4137-4148 (*19,20-Epoxy-10-methoxyicajine, Strychnos icaja constits*)
 Bisset, N.G. et al., *Phytochemistry*, 1974, **13**, 255-258; 259-263; 265-269; 1976, **15**, 1973-1976 (*Epoxydes, 15-Hydroxyvomicine, N-Cyano-sec-pseudobrucine, N-Cyano-sec-pseudostrychnine, Icajine N-oxide, N-methyl-sec-pseudo- β -colubrine*)
 Verpoorte, R. et al., *Org. Magn. Reson.*, 1977, **9**, 567-571; 1984, **22**, 345-348 (*cmr*)
 Rodriguez, F. et al., *Phytochemistry*, 1979, **18**, 2065 (*11-Methoxyicajine*)
 Mostad, A. et al., *Acta Chem. Scand., Ser. B*, 1984, **38**, 821-827 (*cryst struct*)
 Mostad, A. et al., *Acta Chem. Scand., Ser. B*, 1985, **39**, 141-148 (*19,20-Epoxy-15-hydroxyicajine, cryst struct*)

Ichthyopterin

I-13

2-Amino-6-(1,2-dihydroxypropyl)-4,7(1H,8H)-pteridinedione, 9CI. 6-(1,2-Dihydroxypropyl)isoxanthopterin. 7-Hydroxybiopterin. 2-Amino-4,7-dihydroxy-6-(1,2-dihydroxypropyl)pteridine
 [490-58-4]
 [18503-60-1 (N,O,O-tri-Ac), 18503-57-6 (mono-Ac)]



$C_9H_{11}N_5O_4$ 253.217
 Pigment present in the scales of various fish (e.g. *Cyprinus carpio*). Also isol. from scorpion flies (*Panorpa japonica*). Cryst. (EtOH)(as tri-Ac). Mp 188-196° dec. (tri-Ac). Stereochem. not determined.


2'-Deoxy, N⁸-Me: 2-Amino-6-(1-hydroxypropyl)-8-methyl-4,7(1H,8H)-pteridinedione. 6-(1-Hydroxypropyl)-8-methylisoxanthopterin
 [78944-83-9]
 $C_{10}H_{13}N_5O_3$ 251.244
 Pterin from the firefly *Photinus pyralis*. Mp 350°. $[\alpha]_D^{20} +113.1$ (c, 0.011 in MeOH).

Hüttel, R. et al., *Annalen*, 1943, **554**, 69-82 (*isol, fish*)
 Tschesche, R. et al., *Chem. Ber.*, 1958, **91**, 2081-2089 (*synth*)
 Hüttel, R. et al., *Chem. Ber.*, 1960, **93**, 2439-2441 (*struct*)
 Ohta, K. et al., *J. Biochem. (Tokyo)*, 1968, **63**, 127
 Goetz, M.A. et al., *Experientia*, 1981, **37**, 679-680 (*2'-deoxy N-Me*)
 Nakagoshi, M. et al., *Experientia*, 1983, **39**, 742-744 (*Panorpa japonica constit, isol, bibl*)

Icumazole A

I-14

[159311-91-8]




$C_{33}H_{45}NO_7$ 567.721
 Prod. by *Sorangium cellulosum*. Antifungal agent. Yellow oil. $[\alpha]_D^{20} +165$ (MeOH). λ_{max} 214 (ϵ 29510); 302 (ϵ 46770); 316 (ϵ 38900) (MeOH). λ_{max} 214 (ϵ 29512); 302 (ϵ 46773); 316 (ϵ 38900) (MeOH) (Berdy).

2'''-O-Pentofuranoside: Icumazole B

[159311-92-9]
 $C_{38}H_{53}NO_{11}$ 699.837
 Prod. by *Sorangium cellulosum*. Antifungal agent. Yellow oil. λ_{max} 214; 302; 316 (MeOH). λ_{max} 214; 302; 316 (MeOH) (Berdy).
 Ger. Pat., 1994, 4 305 486; CA, **121**, 300667v (*isol*)

Icumazole F

[159311-93-0]


C₂₇H₃₇NO₆ 471.592

Prod. by *Sorangium cellulosum*. Antifungal agent. Yellow oil. λ_{\max} 214 ; 302 ; 316 (MeOH). λ_{\max} 214 ; 302 ; 316 (MeOH) (Berdy).

Ger. Pat., 1994, 4 305 486; CA, 121, 300667v

Ideamine A

[137821-13-7]



C₁₄H₂₃NO₅ 285.339
 $[\alpha]_D^{31} +4$ (c, 1.25 in EtOH).

N-Oxide: **Ideamine A N-oxide**

[137760-62-4]


C₁₄H₂₃NO₆ 301.339

Isol. from adult bodies of the Apocynaceae-feeding danaine butterfly *Idea leuconoe*. $[\alpha]_D^{25} +5$ (c, 2.0 in MeOH).

Nishida, R. et al., *Agric. Biol. Chem.*, 1991, 55, 1787 (isol, pmr, cmr, ms, struct)

Idiospermuline

[169397-89-1]



Absolute
Configuration

C₃₅H₄₂N₆ 546.757

Closely related to Hodgkinsine, H-311. Alkaloid from seeds of *Idiospermum australiense* (Idiospermaceae). Powder. Mp 95-97°. $[\alpha]_D^{25} -2.5$ (c, 1.0 in CHCl₃).


Methiodide (1:3):

Needles (MeOH). Mp 230-231° dec. (subl. at 220-225°). $[\alpha]_D^{25} -2.3$ (c, 0.1 in H₂O).

Duke, R.K. et al., *J. Nat. Prod.*, 1995, 58, 1200 (isol, uv, cd, ir, pmr, cmr, ms, cryst struct)
 Overman, L.E. et al., *Tetrahedron*, 2003, 59, 6905-6919 (synth)

IFB Lactam 1

2,15-Dihydroxy-7-methyl-6-azabicyclo[11.3.0]hexadec-3-en-5-one


C₁₆H₂₇NO₃ 281.394

Prod. by *Trichoderma* sp. obt. from cabbage. Active against *Gloesporium kaki* Hori. Cryst. (MeOH).

Hu, S.-C. et al., *Acta Cryst. E*, 2006, 62, o5777-o5778 (isol, cryst struct)

Ifflaiamine

3,9-Dihydro-2,3,3,9-tetramethylfuro[2,3-b]quinolin-4(2H)-one, 9CI

C₁₅H₁₇NO₂ 243.305

(-)-**form** [31520-95-3]

Alkaloid from *Flindersia iffliana* (Flindersiaceae). Mp 122-125°. $[\alpha]_D^{25} -0.6$ (-6.20) (MeOH). Forms monohydrate, Mp 62-3°, and dihydrate, Mp 56-8°.

Picrate: Mp 207-209° (189-192°).

(±)-**form** [18194-15-5]

Mp 128-129°.

Picrate: Mp 179-182°.


Bosson, J.A. et al., *Aust. J. Chem.*, 1963, 16, 480 (isol, ir, uv, pmr, struct)

Chamberlain, T.R. et al., *J.C.S.(C)*, 1971, 910 (isol, synth, pmr)

Hammerum, S. et al., *Acta Chem. Scand., Ser. B*, 1977, 31, 31 (ms)

Iforrestine

2-Amino-8,9-dihydrooxazolo[4,5-g]pyrido[2,1-c][1,4]benzodiazepine-7,13(6H,7aH)-dione, 9CI
 [125287-08-3]

C₁₄H₁₂N₄O₃ 284.274

Alkaloid from *Isotropis forrestii* (Fabaceae). Nephrotoxin. Needles (MeOH/

I-18


CHCl₃). Mp 330-331° (dec. from 275°).

$[\alpha]_D +362.6$ (c, 1.15 in EtOH).

Colegate, S.M. et al., *Aust. J. Chem.*, 1989, 42, 1249 (isol, ir, pmr, cmr, ms, cryst struct)

Igzamide

[149355-80-6]


C₁₂H₁₀BrN₃O₂ 308.134

Alkaloid from the northeastern Pacific sponge *Plocamissa igzo*. Yellow solid. Genus name given as Plocamissma, which appears to be an error. λ_{\max} 228 (ε 12800); 294 (ε 4900) (MeOH) (Berdy).

Dumdei, E. et al., *J. Nat. Prod.*, 1993, 56, 792-794 (isol, uv, ir, pmr, cmr, ms, struct)

Iheyamine A

[221160-80-1]


C₁₉H₁₃N₃O 299.331

Alkaloid from an ascidian *Polycitrella* sp. Cytotoxic agent. Amorph. purple solid (as trifluoroacetate). λ_{\max} 284 (ε 16000); 325 (sh) (ε 33000); 341 (ε 36000); 360 (sh) (ε 30000); 388 (ε 19000); 499 (ε 5200) (MeOH) (as trifluoroacetate salt). λ_{\max} 287 ; 346 ; 395 ; 515 (MeOH) (Berdy). λ_{\max} 348 (ε 42000); 480 (ε 4400) (MeOH/NaOH) (Berdy).

Sasaki, T. et al., *Tet. Lett.*, 1999, 40, 303-306 (isol, uv, ir, pmr, cmr)

Iheyamine B

[221160-84-5]



Relative
Configuration


C₂₅H₂₃N₃O₃ 413.475

Alkaloid from an ascidian *Polycitrella* sp. Cytotoxic agent. Amorph. purple solid. $[\alpha]_D^{25} -16$ (c, 0.0002 in CHCl₃). λ_{\max} 276 (sh) (ε 18000); 300 (ε 36000); 307 (ε 36000); 330 (sh) (ε 14000); 526 (ε 7400) (CHCl₃). λ_{\max} 284 (ε 16000); 341 (ε 36000); 388 (ε 19000); 499 (ε 5200) (MeOH) (Berdy).

Sasaki, T. et al., *Tet. Lett.*, 1999, 40, 303-306 (isol, uv, ir, pmr, cmr)

Ikimine A

α -Methyl-3-pyridinedodecanal O-methylloxime, 9CI
[131479-30-6]



$C_{19}H_{32}N_2O$ 304.475
Alkaloid from the Okinawan sponge *Niphates* sp. and another sponge. Cytotoxic agent. Shows antibacterial and antifungal activity. Oil. Occurs as a partial racemate of the *E*-form with the *S*-enantiomer, illus., predominating. The *Z*-isomer, also isol., is prob. an artifact. λ_{max} 260 (ϵ 2090); 265 (ϵ 7760); 272 (ϵ 1660) (MeOH) (Derep). λ_{max} 260 (ϵ 2090); 263 (ϵ 3300); 265 (ϵ 8912); 272 (ϵ 1660) (MeOH) (Berdy).

[131564-74-4]

Carroll, A.R. *et al.*, *Tetrahedron*, 1990, **46**, 6637 (*isol, uv, ir, pmr, cmr, ms, struct, activity*)


Kobayashi, J. *et al.*, *J.C.S. Perkin I*, 1992, 1291 (*isol, uv, ir, pmr, cmr, ms*)

Bracher, F. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 223 (*synth*)

Nicholas, G.M. *et al.*, *Tetrahedron*, 2000, **56**, 2921-2927 (*isol, abs config*)

Ikimine B

β -Methyl-3-pyridinedodecanal O-methylloxime, 9CI
[131479-31-7]



$C_{19}H_{32}N_2O$ 304.475
The struct. illus. has been shown to be in doubt (2004). Alkaloid from an unidentified sponge. Cytotoxic. Oil. The *syn*-isomer, also isol., is prob. an artifact. λ_{max} 260 (ϵ 2090); 265 (ϵ 7760); 272 (ϵ 1660) (MeOH) (Derep). λ_{max} 260 (ϵ 1995); 264 (ϵ 10964); 274 (ϵ 1348) (MeOH) (Berdy).


[131564-75-5]

Carroll, A.R. *et al.*, *Tetrahedron*, 1990, **46**, 6637-6644 (*isol, uv, ir, pmr, cmr, ms, struct*)

Romeril, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 3273-3277 (*struct*)

Ikimine D

N-Methoxy-14-(3-pyridinyl)-3-tetradecyn-1-amine, 9CI
[131479-33-9]




$C_{20}H_{32}N_2O$ 316.486
Alkaloid from an unidentified sponge. Cytotoxic. Oil. λ_{max} 260 (ϵ 2000); 265 (ϵ 6610); 270 (ϵ 2880) (MeOH) (Derep).

Carroll, A.R. *et al.*, *Tetrahedron*, 1990, **46**, 6637 (*isol, uv, ir, pmr, cmr, ms, struct*)

I-24**Ileabethoxazole**

[890022-30-7]




Relative Configuration

$C_{21}H_{27}NO_2$ 325.45
Alkaloid from *Pseudopterogorgia elisabethae*. Antimycobacterial agent. Pale yellow oil. $[\alpha]_D^{20} +6.8$ (c, 1 in $CHCl_3$). λ_{max} 225 (ϵ 22200); 277 (ϵ 2300); 288 (ϵ 1800) (MeOH).

Rodriguez, I.I. *et al.*, *Tet. Lett.*, 2006, **47**, 3229-3232 (*isol, pmr, cmr*)

Ilicicolin H, 9CI

[12689-26-8]



$C_{27}H_{31}NO_4$ 433.546
Isol. from *Cylindrocladium ilicicola*. Antifungal antibiotic. Yellow needles (C_6H_6). Mp 144-150°. $[\alpha]_D^{23} -17.4$. λ_{max} 248 (ϵ 23200); 349 (ϵ 5300) (EtOH) (Derep).


Hayakawa, S. *et al.*, *J. Antibiot.*, 1971, **24**, 653 (*isol*)

Matsumoto, M. *et al.*, *Tet. Lett.*, 1976, 3827 (*ir, uv, ms, nmr, struct, stereochem*)

Tanabe, M. *et al.*, *Tetrahedron*, 1983, **39**, 3569 (*biosynth*)

Illudinine

[18500-63-5]



$C_{16}H_{17}NO_3$ 271.315
Alkaloid from the microfungus *Clitocybe illudens*. Mp 228-229° dec.

Me ester: Mp 83-84°.

Nair, M.S.R. *et al.*, *J.O.C.*, 1969, **34**, 240 (*ir, uv, pmr, struct*)

Wenkert, E. *et al.*, *J.O.C.*, 1973, **38**, 4305 (*synth*)

Woodward, R.B. *et al.*, *J.A.C.S.*, 1977, **99**, 8007 (*synth*)

Girija, T. *et al.*, *J.C.S. Perkin I*, 1991, 1467 (*synth*)

Shanker, P.S. *et al.*, *Indian J. Chem., Sect. B*, 1993, **32**, 1209 (*synth*)

Teske, J.A. *et al.*, *J.O.C.*, 2008, **73**, 342-345 (*synth*)

I-27**Illukumbin**

3-(Methylthio)-N-(2-phenylethenyl)-2-propenamide, 9CI
 $MeSCH=C=CHCONH^+CH=CHPh$
 $C_{12}H_{13}NOS$ 219.307

(E,E)-form**Illukumbin A**

[152135-69-8] Not yet known as a nat. prod.

N-Me: Methyllukumbin A

[152175-17-2]

$C_{13}H_{15}NOS$ 233.334

From leaves of *Glycosmis mauritiana* (Rutaceae). Shows antifungal activity. Mp 83-84°. Originally reported as a cinnamic acid deriv. λ_{max} 222 ; 311 (Et₂O) (Berdy).

(1'Z,2E)-form**Illukumbin B**

[152135-60-9]

Isol. from leaves of *Glycosmis mauritiana* (Rutaceae). Exhibits antifungal activity. Mp 120-122°. Originally reported with an isomeric struct. λ_{max} 223 ; 297 (Et₂O) (Berdy).

N-Me: Methyllukumbin B

[152175-16-1]

$C_{13}H_{15}NOS$ 233.334

From leaves of *Glycosmis mauritiana* (Rutaceae). Shows antifungal activity. Mp 71-72°. Originally reported as a cinnamic acid deriv. λ_{max} 283 (Et₂O) (Berdy).


Greger, H. *et al.*, *Tetrahedron*, 1992, **48**, 1209-1218 (*activity*)

Greger, H. *et al.*, *Phytochemistry*, 1993, **34**, 175-179 (*isol, uv, ir, pmr, cmr, ms*)

Hinterberger, S. *et al.*, *Tetrahedron*, 1994, **50**, 6279-6286; 1998, **54**, 487-496 (*struct, synth*)

Imbiline 1

[139220-13-6]



$C_{17}H_{12}N_2O_3$ 292.293

Alkaloid from aerial parts of *Eupomatia bennettii* and bark of *Eupomatia laurina* (Eupomatiaceae). Red prisms (EtOH). Mp 212-214°.

N-De-Me: Hadranthine B

$C_{16}H_{10}N_2O_3$ 278.267

Alkaloid from the stem bark of *Duguetia hadrantha*. Cytotoxic agent. Orange plates (MeCN/CH₂Cl₂). Mp 214-216°. λ_{max} 204 (log ϵ 4.3); 242 (log ϵ 3.85); 282 (sh) (log ϵ 3.1); 305 (log ϵ 3.65); 316 (log ϵ 3.68); 460 (log ϵ 3.55) (MeOH).

11-Hydroxy: Imbiline 2

[139220-14-7]

$C_{17}H_{12}N_2O_4$ 308.293

Alkaloid from aerial parts of *Eupo-*

matia bennettii and bark of *Eupomatia laurina* (Eupomatiaceae). Dark red prisms (EtOH). Mp 267-270°.

9-Methoxy: **Imbiline 3**

[139253-59-1]

C₁₈H₁₄N₂O₄ 322.32

Alkaloid from aerial parts of *Eupomatia bennettii* and bark of *Eupomatia laurina* (Eupomatiaceae). Fine orange-red needles (petrol). Mp 240-244°.

10-Methoxy: **Hadranthine A**

C₁₈H₁₄N₂O₄ 322.32

Alkaloid from the stem bark of *Duguetia hadrantha*. Antimalarial agent. Light purple plates (CH₂Cl₂). Mp 288-290°. λ_{max} 208 (log ε 4.35); 244 (log ε 3.9); 315 (log ε 3.75); 328 (log ε 3.8); 450 (sh) (log ε 3.65) (MeOH).

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1991, **44**, 1615 (*isol, uv, ir, pmr, cmr, ms, struct*)


Muhammad, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 559-562 (*Hadranthines*)

Kitahara, Y. *et al.*, *Tetrahedron*, 2003, **59**, 2885-2891 (*synth*)

Imbricatin†

I-32

[105372-70-1]



Absolute configuration

C₂₄H₂₆N₄O₇S 514.558

Alkaloid from the starfish *Dermasterias imbricata*. Amorph. solid. First example of a benzyltetrahydroisoquinoline alkaloid from a marine organism, and apparently the first example from a non-plant source.

Pathirana, C. *et al.*, *J.A.C.S.*, 1986, **108**, 8288-8289 (*pmr, cmr, struct*)


Burgoyne, D.L. *et al.*, *Can. J. Chem.*, 1991, **69**, 20-27 (*abs config*)

Ohba, M. *et al.*, *Tetrahedron*, 1999, **55**, 4999-5016 (*deriv, synth*)

Imeluteine

I-33

4,5,6,9,10-Pentamethoxyindeno[1,2,3-ij]isoquinoline, 9CI
[38366-03-9]



C₂₀H₁₉NO₅ 353.374

Alkaloid from the stems of *Abuta imene* and *Abuta rufescens* (Menispermaceae).

Yellow prisms (2-propanol/MeOH). Mp 146-147°.

O^o-De-Me: **Norimeluteine**

[152606-56-9]

C₁₉H₁₇NO₅ 339.347

Alkaloid from the roots and wood of *Cissampelos pareira* (Menispermaceae). Cytotoxic to P388 cells. Yellow powder. λ_{max} 210 (ε 25400); 240 (ε 22800); 256 (ε 23200); 292 (ε 18200) (MeOH) (Berdy).

Cava, M.P. *et al.*, *Tetrahedron*, 1975, **31**, 1667 (*isol, uv, ir, pmr, struct*)

Boger, D.L. *et al.*, *J.O.C.*, 1984, **49**, 4050 (*synth, ir, pmr, ms*)

Zhao, B. *et al.*, *Tet. Lett.*, 1991, **32**, 5277 (*synth*)


Morita, H. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1307 (*Norimeluteine*)

Fu, J.-M. *et al.*, *Can. J. Chem.*, 1994, **72**, 227 (*synth, pmr, cmr*)

Imenine

I-34

1,2,3,4-Tetramethoxy-7*H*-dibenzo[de,g]-quinolin-7-one, 9CI
[24268-94-8]



C₂₀H₁₇NO₅ 351.358

Alkaloid from the stems of *Abuta imene* and *Abuta rufescens* (*Abuta splendida*) (Menispermaceae). Yellow needles (MeOH). Mp 206-207°.

Glick, M.D. *et al.*, *Chem. Comm.*, 1969, 1217 (*uv, ir, pmr, struct*)

Cava, M.P. *et al.*, *J.O.C.*, 1973, **38**, 60 (*synth, uv, pmr, ms*)

Cava, M.P. *et al.*, *Tetrahedron*, 1975, **31**, 1667 (*isol, uv, ir, pmr*)

Skiles, J.W. *et al.*, *Can. J. Chem.*, 1979, **57**, 1642 (*isol*)

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1328-1330 (*isol, cmr*)

Imidazole, 9CI, 8CI

I-35

1,3-Diazole. Glyoxaline. Iminazole
[288-32-4]



C₃H₄N₂ 68.078

In general, imidazoles show 1*H* ⇌ 3*H* tautomerism. 4-Substituted imidazoles are indistinguishable from 5-substituted unless blocked by *N*-substitution. Isol. from the seeds of *Lens culinaris* (lentil). Also present in the seeds of other legumes: *Adenantha pavonina*, *Amphimas pterocarpoides*, *Cathormion altissimum* (preferred genus name *Albizia*), *Erythrophleum* sp., *Lathyrus rotundifolius*, *Lathyrus sylvestris*, *Macrotyloma uniflor-*


um (horse gram), *Parkia bicolor*, *Pso-phocarpus tetragonolobus* (winged bean), *Tephrosia platycarpa*, *Vigna radiata* (mung bean). Used as aq. soln. to accelerate the colour reaction in photometric detn. of Co with porphine. Epoxy resin curing agent. Sol. H₂O, EtOH, Et₂O, Me₂CO. Mp 88-90°. Bp 255°. p*K*_{a1} 7; p*K*_{a2} 14.52 (25°, acidic). Sublimes. ▶ LD₅₀ (mus, ori) 880 mg/kg. NI3325000 [17009-90-4, 36877-68-6]

Hayman, A.R. *et al.*, *Phytochemistry*, 1987, **26**, 3247-3248 (*isol, pmr, ms*)

1*H*-Imidazole-1-acetic acid, 9CI

I-36

[22884-10-2]



C₅H₆N₂O₂ 126.115

Isol. from *Coprinus atramentarius* (common ink cap). Cryst. (MeOH aq.). Mp 268-269° dec.

Et ester: [17450-34-9]

C₇H₁₀N₂O₂ 154.168

Mp 50°. Bp₁₇ 160°.

Easson, A.P.T. *et al.*, *J.C.S.*, 1932, 1806

List, P.H. *et al.*, *Arzneim.-Forsch.*, 1960, **10**, 34-40 (*isol*)


Birkofer, L. *et al.*, *Chem. Ber.*, 1960, **93**, 2804 (*synth*)

Sunjic, V. *et al.*, *Croat. Chem. Acta*, 1969, **41**, 107-110 (*struct*)

1*H*-Imidazole-4(5)-acetic acid, 9CI

I-37

[645-65-8]



C₅H₆N₂O₂ 126.115

Isol. from various fungi, e.g. *Polyporus sulphureus*, *Inocybe patouillardii*, *Coprinus atramentarius* (common ink cap) and *Phallus impudicus* (common stinkhorn). Cryst. + 1H₂O. Mp 222° dec.

[51718-81-1]

Tabor, H. *et al.*, *J.A.C.S.*, 1955, **77**, 505-506 (*biochem*)

List, P.H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1959, **292**, 260-271; 777-787; 1962, **295**, 564-571 (*isol*)

List, P.H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1960, **319**, 17-21; *CA*, **55**, 18901g (*isol*)

Okabe, N. *et al.*, *Acta Cryst. C*, 1999, **55**, 1142-1144 (*cryst struct*)

1*H*-Imidazole-4(5)-ethanol, 9CI

I-38

2-(4-Imidazolyl)ethanol. *Histaminol*
[872-82-2]

C₅H₈N₂O 112.131

Metab. of *Saccharomyces rouxii*; found in urine.

Hydrochloride:

Cryst. (Me₂CO/petrol). Mp 120°.

O-Ac:


Cryst. (EtOH) (as oxalate salt). Mp 149-152° (oxalate).

Me ether: 4-(2-Methoxyethyl)imidazole

Cryst. (MeOH/Et₂O) (as oxalate salt). Mp 158-159° (oxalate).

LaRue, T. et al., *Experientia*, 1966, **22**, 729
Schunack, W. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1974, **307**, 517 (synth)

1H-Imidazole-4(5)-propanoic acid, 9CI I-39
3-(4-Imidazolyl)propionic acid
[1074-59-5]



C₆H₈N₂O₂ 140.141

Isol. from the fungus *Coprinus atramentarius* (common ink cap). Also in *Ustilago maydis*. Cryst. (1-butanol aq.). Mp 206-208°.

Me ester: [31434-93-2]

C₇H₁₀N₂O₂ 154.168

Cryst. (Me₂CO) (as hydrochloride). Mp 106-107° (hydrochloride).

Et ester: [52237-38-4]

C₈H₁₂N₂O₂ 168.195

Bp_{0.05} 143-152°.

Windaus, A. et al., *Ber.*, 1910, **43**, 499


Akabori, S. et al., *Ber.*, 1933, **66**, 156

List, P.H. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1960, **319**, 17 (isol)

Schunack, W. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1974, **307**, 517

2,4-Imidazolidinedione, 9CI I-40

Hydantoin. Diketotetrahydroglyoxaline. Glycollylurea. 2,4-Dioximidazolidine. 2,4(5)-Dihydroxyimidazole
[461-72-3]



C₃H₄N₂O₂ 100.077

Present in tobacco smoke. Needles (MeOH). Spar. sol. H₂O, sol. alkalis. Mp 217-220°. pK_{a1} 8.93 (25°).

► Exp. teratogen. MT8210000

3-Me: 3-Methyl-2,4-imidazolidinedione.

3-Methylhydantoin


[6843-45-4]

C₄H₆N₂O₂ 114.104

Isol. from an endophytic mangrove fungus No. 1839. Prisms (H₂O). Sol. H₂O. Mp 155-157°. Sublimes.

Chen, G. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2006, **42**, 138-141 (3-Me, isol)

5-[[3-(1H-Imidazol-4-yl)propenyl]amino]pentanoic acid I-41



C₁₁H₁₅N₃O₃ 237.258


(E)-form

Isol. from the ascidian *Atrium robustum*. Amorph. solid. λ_{max} 277 (ε 39500) (H₂O).

Kehraus, S. et al., *J. Med. Chem.*, 2004, **47**, 2243-2255 (isol, pmr, cmr)

N⁶-[3-(1H-Imidazol-4-yl)propenyl]lysine I-42

2-Amino-6-[[3-(1H-imidazol-4-yl)-2-propenyl]amino]hexanoic acid



C₁₂H₁₈N₄O₃ 266.299


(S,E)-form

Isol. from the ascidian *Atrium robustum*. Amorph. solid. [α]_D²⁵ -9 (c, 1.7 in H₂O). λ_{max} 280 (ε 25500) (H₂O).

Kehraus, S. et al., *J. Med. Chem.*, 2004, **47**, 2243-2255 (isol, pmr, cmr)

N²-[3-(1H-Imidazol-4-yl)propenyl]histamine I-43

3-(1H-Imidazol-4-yl)-N-[2-(1H-imidazol-4-yl)ethyl]propanamide
[50767-80-1]



C₁₁H₁₅N₅O 233.272

Isol. from the Philippine gastropod *Drupa concatenata*. Cryst. (MeOH). Mp 163-164°.


Roseghini, M. et al., *Experientia*, 1973, **29**, 940-941 (isol)

Baures, P.W. et al., *Molecules*, 2002, **7**, 813-816 (synth, pmr, cmr)

2-(1H-Imidazol-4-yl)quinoline, 9CI I-44

Normacrorine

[2054-67-3]



C₁₂H₉N₃ 195.223

Alkaloid from the aerial parts of *Macrorungia longistrobis* (preferred genus name *Anisotes*) (Acanthaceae). Plates (Me₂CO/hexane). Mp 156-157°. pK_a 5.36 (50% MeOH).

Picrate:

Yellow needles (MeOH). Mp 211-212° dec.

1'-Me: 2-(1-Methyl-1H-imidazol-4-yl)-

quinoline, 9CI. Macrorine

[2552-96-7]

C₁₃H₁₁N₃ 209.25

Alkaloid from the aerial parts of *Macrorungia longistrobis* (Acanthaceae). Needles (Me₂CO). Mp 160°. pK_a 4.87 (50% MeOH).

3'-Me: 2-(1-Methyl-1H-imidazol-5-yl)-

quinoline. Isomacrorine

[2552-97-8]

C₁₃H₁₁N₃ 209.25

Alkaloid from the aerial parts of *Macrorungia longistrobis* (Acanthaceae). Needles (hexane/Me₂CO). Mp 110° (105-107°). pK_a 5.09 (50% EtOH).

Arndt, R.R. et al., *J.C.S.*, 1964, 5969

(*Macrorine, Isomacrorine*)

Jordaan, A. et al., *J.C.S.*, 1965, 3001


(*Normacrorine*)

Joynt, V.P. et al., *J.C.S. (B)*, 1966, 980 (ms)

Wuonola, M.A. et al., *Tetrahedron*, 1976, **32**, 1085 (synth, uv, ir, pmr, ms)

2,2'-Iminobis[4-hydroxybenzoic acid], 9CI I-45

5,5'-Dihydroxydiphenylamine-2,2'-dicarboxylic acid. Dianthramine
[136945-65-8]



C₁₄H₁₁NO₆ 289.244


Alkaloid from *Dianthra caryophyllus* infected with *Fusarium oxysporum* f. sp. *dianthi*. λ_{max} 221 (sh); 255; 283 (sh) (EtOH).

Niemann, G.J. et al., *Planta Med.*, 1990, **56**, 564 (isol)

Niemann, G.J. et al., *Physiol. Mol. Plant Pathol.*, 1991, **38**, 417-432 (isol)

3,3'-[Iminobis(methylene)]bis-2(3H)furanone, 9CI I-46


Di(α-methylene-γ-butyrolactonyl)amine
[96562-86-6]





C₁₀H₁₅NO₄ 213.233

Isol. from *Garnieria spathulaefolia* (Proteaceae). Oil which partially crystallizes on standing. Artifact.

Lounasmaa, M. et al., *Heterocycles*, 1985, **23**, 939 (isol, ir, pmr, cmr, ms, struct)


23,29-Imino-B(9a)-homo-19-norstigmasta-1(10),7,9(11),23(N)-tetraen-3-amine I-47C₂₉H₄₄N₂ 420.68**(3α,5α,24ξ)-form** [215245-26-4]Alkaloid from the marine sponge *Corticium* sp. [α]_D²⁵ -50 (c, 0.01 in CHCl₃). λ_{max} 207 (log ε 3.62); 248 (log ε 3.72) (MeOH).**9,11-Dihydro-23,29-Imino-B(9a)-homo-19-norstigmasta-1(10),7,23(N)-trien-3-amine**

[215245-28-6]


C₂₉H₄₆N₂ 422.696Alkaloid from *Corticium* sp. [α]_D²⁵ -8.1 (c, 0.01 in CHCl₃). λ_{max} 210 (log ε 3.74); 240 (log ε 3.5) (MeOH).De Marino, S. *et al.*, *Tet. Lett.*, 1998, **39**, 7611-7614 (*isol, uv, pmr, cmr*)**2-Iminomethyl-3-methyl-6-methylamino-9H-purine** I-48**3,7-Dihydro-N,3-dimethyl-2-(methylimino)-2H-purin-6-amine, 9CI** [112058-06-7]C₈H₁₂N₆ 192.223Isol. from the sea-anemone *Sagartia troglodytes*. Phytotoxic agent. Cryst. (MeOH). Mp 226-227°. λ_{max} 288 (ε 14000) (MeOH) (Berdy).De Rosa, S. *et al.*, *J. Nat. Prod.*, 1987, **50**, 876 (*isol, uv, pmr, ms, cryst struct*)**2-Iminopiperidine** I-49**2-Piperidinimine, 9CI**C₅H₁₀N₂ 98.147Prod. by *Streptomyces atroolivaceus* 70757. NO formation inhibitor. Mp 208-209° dec. Bp 121-135°. Sample not pure.*Hydrochloride*: [16011-96-4]

Mp 159-161°.

▶ TN0404000

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 825C (*ir*)Renault, J. *et al.*, *CA*, 1956, **50**, 9409g (*synth*)Moriconi, E.J. *et al.*, *J.O.C.*, 1968, **33**, 2109-2111 (*synth, ir*)
Japan. Pat., 1996, 2 08 475; *CA*, **125**, 238709n (*isol*)**5-Imino-2-pyrrolidinecarboxylic acid** I-50**5-Iminoproline, 9CI**C₅H₈N₂O₂ 128.13**(S)-form****N-Me: 5-Imino-1-methyl-2-pyrrolidinecarboxylic acid. 5-Imino-1-methylproline**

[915127-35-4]

C₆H₁₀N₂O₂ 142.157Isol. from *Cliona tenuis*. [α]_D²⁵ -10.1 (c, 0.1 in H₂O) (as Me ester). Originally assigned as struct. of Pyrostatin B.Castellanos, L. *et al.*, *Org. Lett.*, 2006, **8**, 4967-4970 (*isol, synth, pmr, cmr, abs config*)**5-Imino-2H-pyrrol-2-ol, 9CI** I-51**2-Hydroxy-5-iminoazacyclopent-3-ene** [71765-74-7]C₄H₆N₂O 98.104Pyrroline antibiotic. Isol. from *Streptovorticillium parvisporogenes*. Weakly active against gram-positive and -negative bacteria. Sol. H₂O. λ_{max} 205 (ε 11300); 240 (ε 1580) (hydrochloride salt in H₂O) (Derep). λ_{max} 205 (ε 11300); 240 (ε 1580) (H₂O) (Berdy).▶ LD₅₀ (mus, ivn) 100 - 200 mg/kg.*Hydrochloride*: [71765-75-8]Needles (propanol). Mp 108° dec. [α]_D²⁴ 0 (c, 0.8 in H₂O).


▶ UY8691300

Okuyama, A. *et al.*, *J. Antibiot.*, 1979, **32**, 768 (*isol, uv, ir, pmr*)**Imperoline** I-52C₂₇H₄₅NO₃ 431.657Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria imperialis* var. *lutea* (Liliaceae). Mp 201-204° (as di-Ac). Redn. prod. of Imperonine, I-53.Paul, L. *et al.*, *Chem. Ber.*, 1958, **91**, 1968**Imperonine** I-53C₂₇H₄₃NO₃ 429.642

Steroidal alkaloid. Prob. the ketone corresponding to Imperoline, I-52.

Struct. unknown. Alkaloid from *Fritillaria imperialis* (Liliaceae). Mp 237-238°. [α]_D²³ -8 (CHCl₃).*Ac*: Mp 208-209°.Paul, L. *et al.*, *Chem. Ber.*, 1958, **91**, 1968**Impranin** I-54


[571167-90-3]

C₂₈H₄₅NO₂ 427.669Alkaloid from the bulbs of *Fritillaria imperialis*. Amorph. powder. [α]_D²⁵ +28 (c, 0.05 in MeOH). λ_{max} 289 (log ε 2.47) (MeOH).**6β-Alcohol: Dihydroimpranin**

[571167-91-4]


C₂₈H₄₇NO₂ 429.685Alkaloid from the bulbs of *Fritillaria imperialis*. Amorph. powder. [α]_D²⁵ -32 (c, 0.08 in MeOH).Akhtar, M.N. *et al.*, *Phytochemistry*, 2003, **63**, 115-122 (*isol, pmr, ms*)**Inaequidenine, 9CI** I-55

[64215-80-1]

C₂₁H₂₅NO₇ 403.431Alkaloid from *Senecio inaequidens* roots (Asteraceae). Oil.Bohlmann, F. *et al.*, *Phytochemistry*, 1977, **16**, 965 (*isol, ms*)**Inandenin-10-one** I-56

[122890-12-4]

[122890-15-7]

C₂₃H₄₅N₃O₂ 395.627Alkaloid from the bark of *Oncinotis nitida*. Oil.**10-Alcohol: Inandenin-10-ol**

[122890-13-5, 122890-14-6]

C₂₃H₄₇N₃O₂ 397.643From *Oncinotis nigra*. Inseparable mixt. with other Inandeninols.

10-Alcohol, 11-hydroxy: Inandenin-10,11-diolC₂₃H₄₇N₃O₃ 413.643From bark of *Oncinotis nitida*. Oil.Badawi, M.M. *et al.*, *Pure Appl. Chem.*, 1973, **33**, 81 (occur, *Inandenin-10-one*, *Inandenin-10,11-diol*)Guggisberg, A. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 85 (occur, *Inandenin-10-ol*)Bienz, S. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 1708 (*synth*, *Inandenin-10-one*, *Inandenin-10-ol*)**Inandenin-11-one****I-57**C₂₃H₄₅N₃O₂ 395.627Alkaloid from leaves of *Oncinotis tenuiloba*. Tentative identification.**11-Alcohol: Inandenin-11-ol**C₂₃H₄₇N₃O₂ 397.643Alkaloid from *Oncinotis nigra*. Inseparable mixt. with other Inandeninols.Guggisberg, A. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 85 (occur, *Inandenin-11-ol*)Doll, M.K.-H. *et al.*, *Phytochemistry*, 1995, **39**, 689 (occur)**Inandenin-12-one****I-58***Inandenin A*

[29579-65-5]

[29579-67-7]

C₂₃H₄₅N₃O₂ 395.627Alkaloid from leaves of *Oncinotis inandensis*, *Oncinotis nitida* and *Oncinotis tenuiloba*. Noncryst. Usually isol. as an inseparable mixt. with Inandenin-13-one, I-59. First successful separation of the mixt. achieved in 1995 via the N,N'-diacetate derivs.**12-Alcohol: Inandenin-12-ol**C₂₃H₄₇N₃O₂ 397.643Alkaloid from *Oncinotis nigra* and leaves of *Oncinotis tenuiloba*. Isol. and characterised as the N,N'-diacetate.Veith, H.J. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1355 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 3026 (*struct*)Trost, B.M. *et al.*, *J.A.C.S.*, 1982, **104**, 6881 (*synth*)Guggisberg, A. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 85 (occur, *Inandenin-12-ol*)Doll, M.K.-H. *et al.*, *Phytochemistry*, 1995, **39**, 689 (*isol*, *Inandenin-12-one*, *Inandenin-12-ol*)**Inandenin-13-one****I-59***Inandenin B*


[29579-66-6]

C₂₃H₄₅N₃O₂ 395.627Alkaloid from leaves of *Oncinotis inandensis*, *Oncinotis nitida* and *Oncinotis tenuiloba*. See note under Inandenin-12-one, I-58.**13-Alcohol: Inandenin-13-ol**

[29579-86-0]

C₂₃H₄₇N₃O₂ 397.643Alkaloid from *Oncinotis nigra* and leaves of *Oncinotis tenuiloba*. Isol. and characterised as the N,N'-diacetate.Veith, H.J. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1355 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 3026 (*struct*)Guggisberg, A. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 85 (occur, *Inandenin-13-ol*)Doll, M.K.-H. *et al.*, *Phytochemistry*, 1995, **39**, 689 (*isol*, *Inandenin-13-one*, *Inandenin-13-ol*)**Incarvillateine****I-60**

[129748-10-3]

C₄₂H₅₈N₂O₈ 718.929Alkaloid from the aerial parts of *Incarvillea sinensis*. Antinociceptive agent. Cryst. (MeOH). Mp 217.2-217.7°. [α]_D -10.8 (MeOH).**N-Oxide: Incarvillateine N-oxide**

[185054-71-1]

C₄₂H₅₈N₂O₉ 734.928Alkaloid from *Incarvillea sinensis*.Powder. [α]_D¹⁸ -19.2 (c, 0.5 in MeOH).

Obt. as an inseparable mixt. of the two possible N-oxides.

5'-Methoxy: Methoxyincarvillateine


[185054-70-0]

C₄₃H₆₀N₂O₉ 748.955Alkaloid from *Incarvillea sinensis*.Powder. [α]_D²⁰ -4 (c, 0.6 in CHCl₃).**Mono(demethoxy): Incarvillateine D**


[244259-23-2]

C₄₁H₅₆N₂O₇ 688.903Alkaloid from *Incarvillea sinensis*.Powder. [α]_D¹⁶ -5.1 (c, 0.3 in CHCl₃).**Bis(demethoxy): Incarvillateine C**

[244259-22-1]

C₄₀H₅₄N₂O₆ 658.876Alkaloid from *Incarvillea sinensis*.Powder. [α]_D¹⁷ -3.2 (c, 0.3 in CHCl₃).Chi, Y.-M. *et al.*, *Phytochemistry*, 1990, **29**,2376-2378 (*isol*, *cryst struct*)Chi, Y.-M. *et al.*, *Phytochemistry*, 1997, **46**,763-769 (*N-oxide*, *Methoxyincarvillateine*)Nakamura, M. *et al.*, *J. Nat. Prod.*, 1999, **62**,1293-1294 (*activity*)Nakamura, M. *et al.*, *Phytochemistry*, 1999,**51**, 595-597 (*Incarvillateines C,D*)Ichikawa, M. *et al.*, *J.A.C.S.*, 2004, **126**, 16553-16558 (*synth*)Chi, Y.-M. *et al.*, *Chem. Pharm. Bull.*, 2005,**53**, 1472-1474 (*isol*, *pmr*, *cmr*, *cryst struct*)Tsai, A.S. *et al.*, *J.A.C.S.*, 2008, **130**, 6316-7317(*synth*)**Incarvillateine E****I-61**C₅₃H₈₁N₃O₈ 888.238Alkaloid from the aerial parts of *Incarvillea sinensis*. Powder. [α]_D²³ -6.4 (c, 0.32 in CHCl₃).Chi, Y.-M. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 1178-1179 (*isol*, *pmr*, *cmr*, *ms*)**Incarvilleine****I-62***6-Hydroxyskytanthine*

[145307-22-8]



Absolute Configuration

C₁₁H₂₁NO 183.293Numbering systems vary. Standardised here on that shown. Alkaloid from aerial parts of *Incarvillea sinensis* (Bignoniaceae). Mp 93.4-93.8°. [α]_D²⁴ -8 (c, 1.24 in CHCl₃).**O-(8-Hydroxy-2,6-dimethyl-2-octenoyl) (E-): Incarvine D**

[185123-29-9]

C₂₁H₃₇NO₃ 351.528From aerial parts of *Incarvillea sinensis* (Bignoniaceae). Powder. [α]_D¹⁶ -4.2 (c, 0.6 in CHCl₃).**O-(8-Carboxy-3,7-dimethyl-2,6-octadienoyl): Incarvine B**

[168009-88-9]

C₂₁H₃₃NO₄ 363.496From aerial parts of *Incarvillea sinensis* (Bignoniaceae). Powder. [α]_D +14 (c, 0.45 in CHCl₃). Ester of Incarvilleine with 2,6-Dimethyl-2,6-octadienedioic acid.**O-(4-Hydroxy-3-methoxy-E-cinnamoyl): Incarvine C. Incarvilleine 7-O-ferulate**

[168009-89-0]

C₂₁H₂₉NO₄ 359.464From aerial parts of *Incarvillea sinensis* (Bignoniaceae). Powder. [α]_D -20.8 (c, 0.46 in CHCl₃).**4a,5-Didehydro: 6-Hydroxydehydroxy-****tanthine**C₁₁H₁₉NO 181.277Alkaloid from fruits of *Tecoma stans* (Bignoniaceae). Oil. Stereochem. uncertain. Called 7-hydroxy in the lit.**4a-Hydroxy: 4a-Hydroxyincarvilleine**

[185054-68-6]



C₁₁H₂₁NO₂ 199.292Alkaloid from *Incarvillea sinensis*.Powder. [α]_D -6.1 (c, 1.5 in CHCl₃).**4a,7a-Diepimer: Kinabalarine F**

[190962-58-4]

C₁₁H₂₁NO 183.293Alkaloid from *Kopsia pauciflora*. Light yellow oil. [α]_D +56 (c, 0.02 in CHCl₃).

7,7a-Diepimer: Kinabalarine A[183815-45-4]
C₁₁H₂₁NO 183.293Alkaloid from the leaves of *Kopsia pauciflora* (Apocynaceae). Plates. Mp 92-93°. [α]_D +26 (c, 0.35 in CHCl₃).**7,7a-Diepimer, 6-ketone: Kinabalarine B**[190961-66-1]
C₁₁H₁₉NO 181.277Alkaloid from the leaves of *Kopsia pauciflora* (Apocynaceae). Light yellow oil. [α]_D -100 (c, 0.02 in CHCl₃).**7,7a-Diepimer, 6-ketone, N-de-Me: Kinabalarine C**[190961-75-2]
C₁₀H₁₇NO 167.25Alkaloid from the leaves of *Kopsia pauciflora*. Light yellow oil. [α]_D +25 (c, 0.09 in CHCl₃).**4,4a,7-Triepimer: Kinabalarine D**[190962-01-7]
C₁₁H₂₁NO 183.293Alkaloid from *Kopsia pauciflora*. Light yellow oil. [α]_D -13 (c, 0.3 in CHCl₃).**4,4a,7-Triepimer, 6-ketone: Kinabalarine E**[190962-45-9]
C₁₁H₁₉NO 181.277Alkaloid from the leaves of *Kopsia pauciflora*. Light yellow oil. [α]_D +134 (c, 0.06 in CHCl₃).Chi, Y.-M. et al., *Phytochemistry*, 1992, **31**, 2930; 1995, **39**, 1485 (isol, ir, pmr, cmr, ms)Lins, A.P. et al., *Phytochemistry*, 1993, **34**, 876 (6-Hydroxydehydrokytanthine)Kam, T.-S. et al., *Nat. Prod. Lett.*, 1996, **8**, 231-235 (pmr, cmr, cryst struct, Kinabalarine A)Chi, Y.-M. et al., *Chem. Pharm. Bull.*, 1997, **45**, 495-498 (cryst struct, abs config)Kam, T.-S. et al., *J. Nat. Prod.*, 1997, **60**, 673 (Kinabalarines, isol, pmr, cmr, ms)Chi, Y.-M. et al., *Phytochemistry*, 1997, **46**, 763-769 (Incarvine D)Ichikawa, M. et al., *J.A.C.S.*, 2004, **126**, 16553-16558 (synth)Honda, T. et al., *J.O.C.*, 2007, **72**, 6541-6547 (synth)**Incarvine A****I-63**

[168434-17-1]

C₃₂H₅₂N₂O₄ 528.774Alkaloid from aerial parts of *Incarvillea sinensis* (Bignoniaceae). Powder. [α]_D -0.9 (CHCl₃).**N-Oxide: Incarvine A N-oxide**C₃₂H₅₂N₂O₅ 544.773Alkaloid from *Incarvillea sinensis* (Bignoniaceae). Powder. [α]_D²⁷ -14.8 (c, 0.4 in CHCl₃). Obt. as an inseparable mixture of the two possible N-oxides.Chi, Y.-M. et al., *Phytochemistry*, 1995, **40**, 353 (isol, pmr, cmr, ms, struct)Chi, Y.-M. et al., *Phytochemistry*, 1997, **46**, 763-769 (N-oxide)**Indanomycin****I-64**X 14547A. Antibiotic X 14547A
[66513-28-8]R = CH₃C₃₁H₄₃NO₄ 493.685Polyether-type antibiotic. Prod. by *Streptomyces antibioticus*, *Streptomyces galbus* and *Streptomyces griseofuscus*. Ionophore active against gram-positive bacteria, shows antihypertensive props. Growth promoter for ruminants. Insecticide. Sol. MeOH, DMF, DMSO, Et₂O; poorly sol. H₂O. Mp 138-141°. [α]_D -328 (c, 1 in CHCl₃). λ_{max} 244 (ε 32000); 291 (ε 16100) (EtOH) (Derep). λ_{max} 244 (ε 32000); 291 (ε 16100) (EtOH) (Berdy). λ_{max} 245 (24000); 292 (12500) (no solvent reported).▶ LD₅₀ (mus, orl) 129 mg/kg. UP6100000

Me ester: [76567-01-6]

Glassy solid. [α]_D²¹ -236 (c, 2.55 in CHCl₃).**8,9-Dihydro: 8,9-Dihydroindanomycin**C₃₁H₄₅NO₄ 495.701Prod. by *Streptomyces galbus*.**16-Deethyl: 16-Deethylindanomycin. A 83094A. Antibiotic A 83094A. Omomycin**

[117615-33-5]


C₂₉H₃₉NO₄ 465.631From *Streptomyces setonii*. Active against gram-positive bacteria and coccidia. Pale yellow powder. Sol. MeOH, C₆H₆; poorly sol. H₂O, hexane. λ_{max} 244 (ε 32000); 291 (ε 16100) (EtOH) (Derep).▶ LD₅₀ (mus, ipr) 196.4 mg/kg, LD₅₀ (mus, orl) 630 mg/kg. UP6120000**Homologue (R = CH₂CH₃): Homoindanomycin**

[129258-44-2]

C₃₂H₄₅NO₄ 507.712Prod. by *Streptomyces galbus*. λ_{max} 244 (E1%/1cm 608); 290 (E1%/1cm 325) (MeOH) (Berdy).**Homologue (R = CH₂CH₃), 4-methyl: 4-Methylhomoindanomycin**C₃₃H₄₇NO₄ 521.739Prod. by *Streptomyces galbus*. Cryst. λ_{max} 246; 291 (MeOH). λ_{max} 246; 291 (MeOH) (Berdy).Ger. Pat., 1978, 2 735 456; CA, **89**, 40863c (isol) Westley, J.W. et al., *J.A.C.S.*, 1978, **100**, 6784 (struct)Liu, C.-M. et al., *J. Antibiot.*, 1979, **32**, 95 (isol)Westley, J.W. et al., *J. Antibiot.*, 1979, **32**, 100 (isol)Edwards, M.P. et al., *J.C.S. Perkin 1*, 1984, 1761 (synth)Nicolaou, K.C. et al., *J.O.C.*, 1985, **50**, 1440 (synth)Burke, S.D. et al., *Tet. Lett.*, 1985, **26**, 1163 (synth, bibl)Larsen, S.H. et al., *J. Antibiot.*, 1988, **41**, 1170 (deriv)Hungarian Pat., 1989, 49 909; CA, **113**, 130720x (Homoindanomycin, Dihydroindanomycin, Methylhomoindanomycin)Burke, S.D. et al., *J.O.C.*, 1994, **59**, 332 (synth)**Indeno[1,2,3-ij][2,7]naphthyridine, 9CI****I-65****1,6-Diazafluoranthene. Eupolauridine.**

Canangine. EL Base 1

[58786-39-3]

C₁₄H₈N₂ 204.231Alkaloid from the bark of *Eupomatia laurina*, *Cleistopholis patens* and *Cananga odorata* (ylang ylang) (Eupomatiaceae, Annonaceae). Shows antiyeast activity. Mp 156-157°. λ_{max} 228 (log ε 4.33); 233 (log ε 4.34); 278 (log ε 4.23); 288 (log ε 4.2); 296 (sh) (log ε 3.98); 335 (log ε 3.57); 350 (log ε 3.81); 367 (log ε 3.8) (EtOH).**Hydrochloride:**Orange cryst. (MeOH/Et₂O). Mp 248-249° dec.**N-Oxide: Eupolauridine N-oxide**

[96889-95-1]

C₁₄H₈N₂O 220.23Alkaloid from the root bark of *Cleistopholis patens* (Annonaceae). Yellow needles (CHCl₃). Mp 186-188°.**Di-N-Oxide: Eupolauridine-di-N-oxide**

[96889-96-2]

C₁₄H₈N₂O₂ 236.229Alkaloid from *Cleistopholis patens*. Orange plates (CHCl₃). Mp 190-195°. λ_{max} 230 (log ε 3.96); 250 (log ε 4.24); 295 (log ε 4.3); 405 (log ε 3.62); 440 (log ε 3.64) (MeOH).**N-Me: 1-Methylindeno[1,2,3-ij][2,7]naphthyridinium**

[58786-40-6]

C₁₅H₁₁N₂⁺ 219.265

Red prismatic needles (MeOH) (as iodide). Mp 267-269° dec. (iodide).

3-Methoxy: 3-Methoxyeupolauridine

[139220-17-0]

C₁₅H₁₀N₂O 234.257Alkaloid from bark of *Eupomatia laurina* (Eupomatiaceae). Yellow needles (petrol). Mp 166-168°. λ_{max} 216 (ε 21880); 226 (ε 21380); 237 (ε 23440); 285 (ε 9120); 298 (ε 12020); 306 (ε 12880); 320 (ε 6760); 341 (ε 3800); 358 (ε 6030); 376 (ε 6310) (EtOH).Bowden, B.F. et al., *Aust. J. Chem.*, 1972, **25**, 2659-2669; 1975, **28**, 2681-2701 (isol, uv, pmr, ms, struct, synth)Leboeuf, M. et al., *Ann. Pharm. Fr.*, 1975, **33**, 43-47 (isol)Leboeuf, M. et al., *J. Nat. Prod.*, 1976, **39**, 459-460 (uv, ir, pmr, ms, struct)Taylor, W.C. et al., *Aust. J. Chem.*, 1984, **37**, 1097-1104 (biosynth)

- Waterman, P.G. *et al.*, *Phytochemistry*, 1985, **24**, 523-527 (oxide, dioxide)
 Hufford, C.D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 961-964 (activity)
 Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1991, **44**, 1615-1626 (3-Methoxyeupolauridine)
 Tong, T.H. *et al.*, *Synth. Commun.*, 1992, **22**, 1773-1782 (synth)

Indicamine†**I-66**C₁₄H₂₃NO 221.342

Monoterpene alkaloid. Struct. unknown. Alkaloid from *Pedicularis dolichorrhiza*, *Plantago albicans*, *Plantago indica*, *Plantago notata*, *Plantago ovata* and *Plantago psyllium* (African plantain) (Scrophulariaceae, Plantaginaceae). Mp 127-128° (as picrate).

Daniilova, A.V. *et al.*, *Zh. Obshch. Khim.*, 1952, **22**, 2237-2239; *CA*, **48**, 691c (isol)


Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 195; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 168

Karawya, M.S. *et al.*, *U. A. R. J. Pharm. Sci.*, 1971, **12**, 53-61; *CA*, **78**, 13729e (isol)

Indicaxanthin**I-67**

4-[2-(2-Carboxy-1-pyrrolidinyl)ethenyl]-2,3-dihydro-2,6-pyridinedicarboxylic acid, 9CI

[2181-75-1]

C₁₄H₁₆N₂O₆ 308.29

Constit. of the fruits of *Opuntia ficus-indica* (Indian fig), flowers of *Mirabilis jalapa* and petals of *Portulaca grandiflora* (Cactaceae, Nyctaginaceae, Portulacaceae). Pigment of the small betaxanthin group which replace anthocyanins in some plants of the Centrospermae. Orange cryst. (H₂O). Mp 160-162° dec. [α]_D²⁰ +394 (c, 1 in 0.1M phosphate buffer; pH 7). pK_a 3.3. Undergoes rapid E/Z-isomerism in soln.

Piattelli, M. *et al.*, *Tetrahedron*, 1964, **20**, 2325 (isol, struct, abs config)

Piattelli, M. *et al.*, *Phytochemistry*, 1965, **4**, 817 (isol)


Piattelli, M. *et al.*, *Phytochemistry*, 1972, **11**, 2499 (biosynth)

Wyler, H. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 1793 (pmr)

Stintzing, F.C. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1008-1016 (pmr, cmr)

Indicine†**I-68**

[480-82-0]

C₁₅H₂₅NO₅ 299.366

Ester of Retronecine, in T-188 with Trachelanthic acid. Diastereoisomeric with Intermedine, I-160 and Echinatine, E-12. Alkaloid from *Heliotropium indicum* (Boraginaceae) and twigs of *Messerschmidia argentea*. Shows strong antitumour activity. Prisms (petrol). Mp 97-98°. [α]_D +22.3 (c, 1.65 in EtOH). V. difficult to crystallise.

▶EK7789900

Picrate:

Leaflets (C₆H₆). Mp 81-86° Mp 90° (dimorph.).

Methiodide:

Blades (EtOH/Et₂O). Mp 159-160°. [α]_D +12.5 (c, 4.32 in EtOH).

N-Oxide: Indicine N-oxide. NSC 132319 [41708-76-3]

C₁₅H₂₅NO₆ 315.366Constit. of *Heliotropium indicum*(Boraginaceae) and twigs of *Messerschmidia argentea*. Exhibits strong

antitumour activity vs. Walker 256 carcinosarcoma, poss. by a mechanism of purine antagonism; clinically tested antineoplastic agent (1985). Deliquescent cryst. (MeOH). Mp 130-131° dec. [α]_D²⁰ +34 (c, 7 in EtOH). Log P -2.8 (uncertain value) (calc).

▶Haematopoietic and hepatotoxic effects reported when used therapeutically. LD₅₀ (mus, ipr) 1917 mg/kg. EK7791200

3'-Ac: Acetylcindicine

[11014-09-8]

C₁₇H₂₇NO₆ 341.403Alkaloid from *Heliotropium indicum*

(Boraginaceae) and twigs of *Messerschmidia argentea*. Gum. [α]_D²⁵ -13.8 (c, 2.2 in EtOH).

3'-Ac, N-oxide: 3'-Acetylcindicine N-oxideC₁₇H₂₇NO₇ 357.403

Isol. from twigs of *Messerschmidia argentea*. Gum. [α]_D²⁵ -3.7 (c, 5.4 in EtOH).

Tri-Ac: [39870-08-1]Viscous oil. [α]_D -9.25 (c, 4.4 in EtOH).**Tri-Ac, picrate:**

Leaflets (EtOH). Mp 131-132°.

3'-Epimer: Morifoline

[95462-13-8]

C₁₅H₂₅NO₅ 299.366

Alkaloid from *Critonia morifolia*. Ester of Retronecine, in T-188 with (+)-Viridifloric acid.

2'- or 3'-Epimer, 3'-Ac: 9-(3'-Acetylviridifloryl)retronecine

[115793-79-8]

C₁₇H₂₇NO₆ 341.403

Alkaloid from roots of *Heliotropium curassavicum* var. *curassavicum* (Boraginaceae). Config. of side-chain centres not detd. May be the Ac deriv. of Morifoline.

2' or 3'-Epimer, O³-Me: SincamidineC₁₆H₂₇NO₅ 313.393

Alkaloid from *Amsinckia intermedia* (Boraginaceae). Noncryst. Hydrol. gives an acid Mp 138-139°, [α]_D²⁰ +5.8°, isomeric with Heliotrinic acid. May be the Me ether of Morifoline.

Mattocks, A.R. *et al.*, *J.C.S.*, 1961, 5400 (isol, struct)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1955 (*Sincamidine*)

Mattocks, A.R. *et al.*, *J.C.S.(C)*, 1967, 329 (*Acetylcindicine*)

Kugelman, M. *et al.*, *J. Nat. Prod.*, 1976, **39**, 125 (*pharmacol, oxide*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (cmr)

Letendre, L. *et al.*, *Cancer (Philadelphia)*, 1984, **54**, 1256 (*N-oxide, activity, tox*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 32 (rev. *pharmacol, oxide*)

Zalkow, L.H. *et al.*, *J. Med. Chem.*, 1985, **28**, 687 (synth)

Vedejs, E. *et al.*, *J.O.C.*, 1985, **50**, 2170 (synth)

Mattocks, A.R. *et al.*, *Chemistry and Toxicology of Pyrrolizidine Alkaloids*, Academic Press, London, 1986, 312 (*oxide, rev*)

Nishimura, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 4107 (synth, enantiomers)

Davicino, J.G. *et al.*, *Phytochemistry*, 1988, **27**, 960 (9-(3'-Acetylviridifloryl)retronecine)

Moore, D.J. *et al.*, *Toxicol. Appl. Pharmacol.*, 1989, **101**, 271 (*oxide, tox*)

Miser, J.S. *et al.*, *Invest. New Drugs*, 1991, **9**, 339 (*clin trial, N-oxide*)

Miser, J.S. *et al.*, *Am. J. Clin. Oncol.*, 1992, **15**, 135 (*clin trial, N-oxide*)

Ogawa, T. *et al.*, *Tetrahedron*, 1993, **49**, 1571 (synth, *Indicine, oxide*)

Nambu, M. *et al.*, *Chem. Comm.*, 1996, 1619 (synth)

Ogihara, K. *et al.*, *Phytochemistry*, 1997, **44**, 545 (3'-Acetylcindicine *N-oxide*)

Wiedenfeld, H. *et al.*, *Phytochemistry*, 1998, **49**, 1463-1465 (*Morifoline*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ICD100

Indicine**I-69**

[11014-14-5]


Pyrrolizidine alkaloid, struct. unknown. Alkaloid from *Heliotropium indicum* (Boraginaceae). Gum. Hydrol. gives retronecine in T-188 and a noncryst. acid resembling trachelanthic acid.

Mattocks, A.R. *et al.*, *J.C.S.(C)*, 1967, 329-331

Indigoidin**I-70**

5-Amino-3-[5-amino-1,6-dihydro-2,6-dioxo-3(2H)-pyridinylidene]-2,6(1H,3H)-pyridinedione, 9CI, 5,5'-Diamino-4,4'-dihydroxy-3,3'-diaz-2,2'-diphenoquinone [2435-59-8]

[2435-61-2 (di-Ac)]

C₁₀H₈N₄O₄ 248.198


Blue pigment from *Corynebacterium insidiosum*, *Pseudomonas indigofera*, *Arthrobacter atrocyaneus* and *Arthrobacter polychromogenes*. Blue solid. Extremely insol. all org. solvs., sol. hot conc. acids. λ_{max} 605 (log ε 4.33) (*N*-Methylpyrrolidine).

Kuhn, R. *et al.*, *Arch. Mikrobiol.*, 1965, **51**, 71-84 (isol)

- Kuhn, R. *et al.*, *Chem. Ber.*, 1965, **98**, 2139-2153 (*synth, struct, ir, pmr, uv*)
 Dieris, C.-G. *et al.*, *Synthesis*, 1979, 948-950 (*synth*)
 Takahashi, H. *et al.*, *J. Biol. Chem.*, 2007, **282**, 9073-9081 (*isol, ms*)

Indigotisocoumarin A I-71

6-Hydroxy-7-(hydroxymethyl)-3-(1H-pyrrol-3-yl)-1H-2-benzopyran-1-one
 [924890-68-6]




C₁₄H₁₁NO₄ 257.245
 Constit. of *Isatis indigotica*.

Wang, F. *et al.*, *Shenyang Yaoke Daxue Xuebao*, 2005, **22**, 187-188; *CA*, **146**, 224723j (*isol*)

Indigotin† I-72

Δ^{2,2'}-Biindoline-3,3'-dione, 8CI. Δ^{2,2'}-Bipseudoindoxyl. Diindogen. Indigo. Indigo blue. C.I. Natural Blue 1. C.I. Vat Blue 1. C.I. 75780
 [482-89-3]



C₁₆H₁₀N₂O₂ 262.267
 Known since ancient times; obt. from various plants, e.g. *Indigofera tinctoria*, where it occurs as its precursor Indican. Constit. of Tyrian purple obt. from *Murex trunculus*. Blue powder with copper lustre. Insol. H₂O. Mp 390-392°. Sublimes.

▶ DU2988400

I,I'-Di-Ac: [33934-64-4]
 C₂₀H₁₄N₂O₄ 346.342
 Red cryst. Mp 200-220°.

I,I'-N-Dibenzoyl:
 C₃₀H₁₈N₂O₄ 470.483
 Violet cryst. Mp 254° Mp 257°.

N-Me: [23831-49-4]
 C₁₇H₁₂N₂O₂ 276.294
 Mp 254°.

Di-N-Me: [18996-72-0]
 C₁₈H₁₄N₂O₂ 290.321
 Sol. nonpolar solvs. Mp 182-183°. Green in soln. Shows bathochromicity compared with indigotin owing to nonpolarity. Has no affinity for cotton fibres.

N-Et: [97207-48-2]
 C₁₈H₁₄N₂O₂ 290.321
 Mp 245°.

Di-N-Et: [109670-05-5]
 C₂₀H₁₈N₂O₂ 318.374
 Mp 161°.

N-Benzyl: [449761-23-3]

C₂₃H₁₆N₂O₂ 352.392
 Mp 210°.

I,I'-N-Dibenzyl: [858229-81-9]

C₃₀H₂₂N₂O₂ 442.516
 Mp 165°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 394B (*ir*)

Harley-Mason, J. *et al.*, *J.C.S.*, 1950, 2907 (*synth*)

Pummerer, R. *et al.*, *Annalen*, 1954, **590**, 173-194 (*N,N-di-Me*)

Yamanaka, T. *et al.*, *CA*, 1956, **50**, 304 (*synth*)
 Gosteli, J. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1980-1983 (*synth*)

Christophersen, C. *et al.*, *Tet. Lett.*, 1977, 1747-1748

Miehe, G. *et al.*, *Angew. Chem., Int. Ed.*, 1991, **30**, 964-967 (*cryst struct, bibl, N,N-di-Me*)

Wouters, J. *et al.*, *J. Soc. Dyers Colour.*, 1991, **107**, 226-269 (*isol, hplc*)

Kirk-Othmer Encycl. Chem. Technol., 4th edn., Wiley, 1991, **8**, 796-800 (*rev*)

Rademacher, P. *et al.*, *Chem. Ber.*, 1992, **125**, 1773-1775 (*pe, struct*)

Grimme, G. *et al.*, *Chem. Ber.*, 1993, **126**, 1015-1021 (*struct*)


Maugard, T. *et al.*, *Phytochemistry*, 2001, **58**, 897-904 (*isol, uv*)

Tanoue, Y. *et al.*, *The Fischer Indole Synthesis*, 2001, **67**, 726-729 (*synth*)

Sigma-Aldrich Library of Stains, Dyes and Indicators, 403

Indirubin I-73

3-(1,3-Dihydro-3-oxo-2H-indol-2-ylidene)-1,3-dihydro-2H-indol-2-one, 9CI. Δ^{2,3'}-Biindoline-2',3'-dione. Courouppitine B. Indigo red. C.I. 75790
 [479-41-4]



C₁₆H₁₀N₂O₂ 262.267
 Alkaloid from *Indigofera* spp. as minor constit. of indigo. Isol. from the molluscs *Hexaplex trunculus* and *Murex brandaris* (as component of Tyrian purple). Constit. of *Courouppitine*, root of *Isatis indigotica*, *Isatis tinctoria*, *Polygonum tinctorium* and *Baphicacanthus cusia*. Isol. from the fungal pigment of a *Schizophyllum commune* mutant. Inhibitor of Lewis lung carcinoma and Walker 256 carcinosarcoma in mice. Red cryst. Sol. EtOH, AcOH; poorly sol. H₂O, hexane. Mp > 400° subl. λ_{max} 522; 561 (xylene). λ_{max} 207; 239; 290; 360; 540 (MeOH) (Berdy).

▶ DU2995000

I,I'-Di-Ac: Mp 192-193°.

Oxime: [160807-49-8]
 C₁₆H₁₁N₃O₂ 277.282
 Glycogen synthase kinase (GSK-3) inhibitor. Deep red cryst. Mp 246°.

6-Bromo: 6-Bromindirubin
 [200273-66-1]

C₁₆H₉BrN₂O₂ 341.163
 Isol. from the mollusc *Hexaplex trunculus*. Selective inhibitor of GSK-3.

6'-Bromo: [200273-67-2]

C₁₆H₉BrN₂O₂ 341.163

Isol. from the mollusc *Hexaplex trunculus*.

6,6'-Dibromo: 6,6'-Dibromindirubin

[171565-08-5]

C₁₆H₈Br₂N₂O₂ 420.059

Isol. from the molluscs *Hexaplex trunculus* and *Nucella lapillus*. λ_{max} 305 (log ε 4.43); 350 (sh); 367 (log ε 3.94); 387 (sh); 525 (sh); 552 (log ε 4.07) (CHCl₃). λ_{max} 305 (log ε 4.28); 371 (log ε 3.73); 552 (log ε 3.89) (CCl₄).

De Diesbach, H. *et al.*, *Helv. Chim. Acta*, 1945, **28**, 690-700 (*synth*)

v. Eller-Pandraud, H. *et al.*, *Bull. Soc. Chim. Fr.*, 1958, 712-714 (*cryst struct*)

Epstein, E. *et al.*, *CA*, 1966, **66**, 102443 (*isol*)

Plieninger, D.W. *et al.*, *Chem. Ber.*, 1966, **99**, 3063-3069 (*synth*)

Sen, A.K. *et al.*, *Tet. Lett.*, 1974, 609-610 (*isol*)
 Bergmann, J. *et al.*, *Tet. Lett.*, 1977, 2625-2626 (*struct*)

Bergmann, J. *et al.*, *Tetrahedron*, 1985, **41**, 2879-2891 (*isol, struct, props*)

Shen, Y. *et al.*, *Yaoxue Tongbao*, 1985, **19**, 601-602; *CA*, **102**, 10071 (*isol*)

Ji, X. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 137-139; *CA*, **103**, 98313 (*activity*)

Zhang, S. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 301-305; *CA*, **103**, 92918 (*detn*)

Koren, Z.C. *et al.*, *Isr. J. Chem.*, 1995, **35**, 117-124 (*hplc*)

Clark, R.J.H. *et al.*, *J. Soc. Dyers Colour.*, 1997, **113**, 316-321 (*Bromindirubins*)

Leclerc, S. *et al.*, *J. Biol. Chem.*, 2001, **276**, 251-260 (*oxime, pharmacol, synth, pmr, cmr*)

Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-769 (*6,6'-Dibromindirubin, rev*)

Maugard, T. *et al.*, *Phytochemistry*, 2001, **58**, 897-904 (*isol, uv*)

Meijer, L. *et al.*, *Chem. Biol.*, 2003, **10**, 1255-1266 (*Bromindirubins*)

Mak, N.K. *et al.*, *Biochem. Pharmacol.*, 2004, **67**, 167-174 (*oxime, pharmacol*)


Polychronopoulos, P. *et al.*, *J. Med. Chem.*, 2004, **47**, 935-946 (*synth, oxime, acetoxime, pmr, pharmacol*)

Sato, N. *et al.*, *Nat. Med. (N.Y.)*, 2004, **10**, 55-63 (*pharmacol*)

Xie, Y. *et al.*, *Neurosci. Lett.*, 2004, **367**, 355-359 (*oxime, pharmacol*)

Indisocin I-74

3-(Acetyloxy)-3-(2-chloro-2-isocyanoethenyl)-1,3-dihydro-2H-indol-2-one, 9CI. 3-(Acetyloxy)-3-(2-chloro-2-isocyanoethenyl)-2-oxoindole. 3-Acetoxy-3-(2-chloro-2-isocyanovinyl)-2-oxoindole
 [90632-49-8]



C₁₃H₉ClN₂O₃ 276.678

Indole antibiotic. Isol. from *Nocardia blackwellii*. Active against gram-positive and -negative bacteria and fungi. Sol. MeOH, C₆H₆; poorly sol. H₂O. [α]_D²⁵ +20 (c, 0.03 in MeOH). Stable in org. solvs. Related to 3-(2-Isocyanoethenyl)-1H-indole, 1-216. λ_{max} 208 (ε 1120); 225 (sh) (ε 730); 255 (sh) (ε 340); 290 (sh) (ε 80)

(MeOH) (Derep). λ_{\max} 208 (E1%/1cm 1120) (MeOH) (Berdy).

N-Me: N-Methylindisocin

[90632-48-7]

C₁₄H₁₁ClN₂O₃ 290.705

Isol. from *Nocardia blackwellii*. Active against gram-positive and -negative bacteria and fungi. $[\alpha]_{\text{D}}^{25}$ +22 (c, 0.04 in MeOH). λ_{\max} 208 (ε 1120); 225 (sh) (ε 730); 255 (sh) (ε 340); 290 (sh) (ε 80) (MeOH) (Derep).


[114114-51-1, 114114-64-6, 114114-52-2, 114114-63-5]

Isshiki, K. *et al.*, *J. Antibiot.*, 1987, **40**, 1195; 1199; 1202 (isol, uv, ir, pmr, ms, struct, synth, props)

Indizoline

I-75

1-Methoxy-2-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, 9CI
[54313-18-7]



C₁₉H₁₉NO₂ 293.365


Alkaloid from the roots of *Clausena indica* (Rutaceae). Plates (hexane/CH₂Cl₂). Mp 170-171°. Isomeric with Atanisatine, A-1530.

Joshi, B.S. *et al.*, *Indian J. Chem.*, 1974, **12**, 437 (isol, uv, ir, pmr, ms, struct)

Indocarbazostatin

I-76

[253450-08-7]



C₂₈H₂₁N₃O₇ 511.49

Prod. by *Streptomyces* sp. TA-0403. Inhibitor of NGF-induced neurite outgrowth. Pale yellow cryst. (Me₂CO/C₆H₆). Mp 256° dec. $[\alpha]_{\text{D}}^{26}$ +51.3 (c, 0.05 in MeOH). λ_{\max} 236 (log ε 4.61); 283 (log ε 3.36); 290 (log ε 4.43); 326 (log ε 4.6) (MeOH).

Me ester analogue: Indocarbazostatin C

C₂₇H₁₉N₃O₇ 497.463

Prod. by a mutant strain, *Streptomyces* sp. MUV-6-83. Yellow powder. Mp 259-262°. λ_{\max} 236 (log ε 4.44); 286 (log ε 4.1); 293 (log ε 4.2); 325 (log ε 4.4) (MeOH).

8-Amino: Indocarbazostatin B

[334831-87-7]

C₂₈H₂₂N₄O₇ 526.504

Prod. by *Streptomyces* sp. TA-0403. Inhibitor of NGF-induced neurite growth. Yellow powder. Mp 255° dec. $[\alpha]_{\text{D}}^{26}$ -48.7 (c, 0.05 in MeOH). λ_{\max} 236

(log ε 4.36); 270 (log ε 4.11); 292 (log ε 4.04); 327 (log ε 4.04) (MeOH).

8-Amino, Me ester analogue: Indocarbazostatin D

C₂₇H₂₀N₄O₇ 512.478

Prod. by a mutant strain, *Streptomyces* sp. MUV-6-83. Orange powder. Mp > 300°. $[\alpha]_{\text{D}}$ -44 (c, 0.05 in MeOH). λ_{\max} 237 (log ε 4.06); 269 (log ε 3.9); 291 (log ε 3.85); 325 (log ε 3.64) (MeOH).

Ubukata, M. *et al.*, *J. Antibiot.*, 1999, **52**, 921-924 (Indocarbazostatin)


Matsuura, N. *et al.*, *J. Antibiot.*, 2002, **55**, 355-362; 363-370 (Indocarbazostatin B)

Feng, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 627-633 (Indocarbazostatin C,D)

Indolactam I

I-77

[132738-83-1]




C₁₈H₂₅N₃O₂ 315.414

Lactam antibiotic. Isol. from *Streptovercillium blastmyceticum*. Tumour promotor. Amorph. λ_{\max} 228 (ε 27600); 289 (ε 7500); 300 (ε 7500) (MeOH) (Derep). Irie, K. *et al.*, *Tet. Lett.*, 1990, **31**, 7337 (isol, struct, pmr, cd)

Indolactam V

I-78

[90365-57-4]



C₁₇H₂₃N₃O₂ 301.388

Prod. by *Streptomyces blastmyceticum* sp. NA34-17. Induces the early antigen of Epstein-Barr virus. CSF inducer. Needles (EtOH aq.) or amorph. solid. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 130-165°. $[\alpha]_{\text{D}}^{27}$ -170 (c, 0.499 in EtOH). λ_{\max} 228 (ε 27600); 289 (ε 7500); 300 (ε 7500) (MeOH) (Derep).

O-Ac: O-Acetylindolactam V

[91403-61-1]

C₁₉H₂₅N₃O₃ 343.425

Prod. by *Streptomyces blastmyceticum* sp. NA34-17. Induces the early antigen of Epstein-Barr virus. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 100-110°. $[\alpha]_{\text{D}}^{26}$ -57 (c, 0.315 in EtOH). λ_{\max} 227 (ε 27800); 295 (ε 7200) (EtOH) (Berdy).

O-Malonoyl: O-Malonylindolactam V

C₂₀H₂₅N₃O₅ 387.435

Prod. by *Streptomyces blastmyceticum*

sp. NA34-17. Amorph. solid. $[\alpha]_{\text{D}}^{23}$ -47.4 (c, 0.3 in MeOH). λ_{\max} 226 (ε 24300); 298 (ε 6800) (MeOH).

N-De-Me: N-Desmethylindolactam V

[90365-52-9]

C₁₆H₂₁N₃O₂ 287.361

Isol. from *Streptomyces blastmyceticum*. Needles (EtOAc/hexane). Mp 199-201°. $[\alpha]_{\text{D}}^{25}$ -81 (c, 0.25 in EtOH).

9-Epimer: Epiindolactam V

[90365-55-2]

Needles (CHCl₃). Mp 134-136°. $[\alpha]_{\text{D}}^{20}$ -84.9 (c, 0.75 in MeOH).

7-(3,7-Dimethyl-2,6-octadienyl): 7-Geranylindolactam V

C₂₇H₃₉N₃O₂ 437.624

Metab. of *Streptomyces blastmyceticum*. Tumour promotor. Amorph. powder. $[\alpha]_{\text{D}}^{28}$ -119 (c, 0.47 in MeOH). λ_{\max} 229 (ε 30200); 286 (ε 8900) (MeOH) (Berdy).

[90365-56-3, 84622-58-2, 132341-58-3, 90365-54-1]

Endo, Y. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 3457; 1984, **32**, 358 (synth, abs config)

Irie, K. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 1269 (isol, uv, ir, pmr, cd)

Endo, Y. *et al.*, *Tetrahedron*, 1986, **42**, 5905-5924; 1987, **43**, 3695-3704 (synth, stereochem, props)

Hagiwara, N. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 641-648 (N-Desmethylindolactam V)

Mascal, M. *et al.*, *Chem. Comm.*, 1988, 589 (synth)

Masuda, T. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 2257-2260 (synth)

Irie, K. *et al.*, *Tet. Lett.*, 1990, **31**, 101 (biosynth)

Kogan, T.P. *et al.*, *Tetrahedron*, 1990, **46**, 6623 (synth, bibl)

Mascal, M. *et al.*, *J.C.S. Perkin 1*, 1992, 823 (synth)

Kawai, T. *et al.*, *J.O.C.*, 1992, **57**, 6150 (epi-Indolactam V)

Semmelhack, M.F. *et al.*, *Tet. Lett.*, 1993, **34**, 1395 (synth)

Irie, K. *et al.*, *J. Nat. Prod.*, 1994, **57**, 363 (7-Geranylindolactam V)


Quick, J. *et al.*, *Tet. Lett.*, 1994, **35**, 8549 (synth)

Irie, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1669-1670 (Malonylindolactam V, biosynth)

Indole, 9CI, 8CI

I-79

1-Benzazole, 2,3-Benzopyrrole, 1-Azaindene. Ketole (obsol.). FEMA 2593



1H-form

C₈H₇N 117.15


Constit. of coal tar. Constit. of several flower oils, esp. of *Jasminum* and *Citrus* spp. (Oleaceae, Rutaceae). A prod. of bacterial dec. of proteins, ubiquitous in faeces. Constit. of anal gland secretion of aardwolf *Proteles cristatus*. Prod. by marine bacterium strain Ta-6 from *Tapes philippinarum* and by a range of microorganisms isol. from fish and shellfish. Perfumery and flavouring ingredient. Used as 0.5% EtOH soln. for

photometric detn. of NO_2^- . λ_{max} 530 nm (ϵ 9500, 2.5-5M HCl). Cryst. (H_2O). Sol. EtOH, MeOH, Et_2O ; poorly sol. H_2O . Mp 52°. Bp 253-254° Bp₅ 122.5-124°. $\text{p}K_{\text{a}1}$ -2.4; $\text{p}K_{\text{a}2}$ 16.97 (25°, NH, KOH aq.). λ_{max} 216; 270; 278; 281; 295 (MeOH) (Berdy).

- LD₅₀ (rat, orl) 1000 mg/kg. NL2450000
Stowe, B.B. et al., *Prog. Chem. Org. Nat. Prod.*, 1959, **17**, 250-297 (occur)
Apps, P.J. et al., *J. Chem. Ecol.*, 1989, **15**, 1681-1688 (occur)
Kirk-Othmer *Encycl. Chem. Technol.*, 4th edn., Wiley, 1991, **14**, 161-173 (rev)
Takao, T. et al., *Biosci., Biotechnol., Biochem.*, 1994, **58**, 1780-1783 (isol, occur)

1H-Indole-3-acetaldehyde, I-80 9CI

β -Indolylacetaldehyde. 3-(Formylmethyl)indole. Skototenin [2591-98-2]



$\text{C}_{10}\text{H}_9\text{NO}$ 159.187

Isol. from various plants incl. *Brassica* spp. Intermed. in biosynth. of 1H-Indole-3-acetic acid, I-81 from Tryptophan, T-640. Syrup. Bp_{0.00005} 120°.

Oxime: [2776-06-9]
[95394-24-4, 95394-25-5]

$\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}$ 174.202

Metab. of *Archangium gephyra* and various plant spp. incl. *Brassica* spp. Cryst. (CHCl_3). Mp 140-141°. Isol. as a mixt. of *E/Z*-isomers.

Semicarbazone: Mp 142-155°.

2,4-Dinitrophenylhydrazone: Mp 196-202°.

Gordon, S.A. et al., *Arch. Biochem.*, 1949, **43**, 7096

Brown, J.B. et al., *Nature (London)*, 1952, **169**, 335 (synth, uv, ir)

Ahmad, A. et al., *Can. J. Chem.*, 1960, **38**, 2523 (synth, oxime)

Ihara, M. et al., *Tetrahedron*, 1985, **41**, 2109 (synth, ir, pmr)


Schlecht, M.F. et al., *J. Med. Chem.*, 1990, **33**, 386 (synth)

Bohlendorf, B. et al., *Annalen*, 1996, 49-53 (isol, oxime)

1H-Indole-3-acetic acid, 9CI I-81

3-Indolylacetic acid. Heteroauxin. Rhizopin. IAA. Rhizipon A

[87-51-4]



$\text{C}_{10}\text{H}_9\text{NO}_2$ 175.187

Isol. from the marine alga *Undaria pinnatifida*. Widely distributed in higher

plants, also in bacteria, algae, yeasts and fungi. Plant growth hormone (auxin).

Involved in root development. Phytotoxic agent. Cryst. (CHCl_3). Sol. EtOH, Me_2CO ; fairly sol. H_2O , CHCl_3 , Mp 164-165°. $\text{p}K_{\text{a}1}$ 4.36; $\text{p}K_{\text{a}2}$ 16.9 (NH). λ_{max} 229 (ϵ 10400); 273 (ϵ 9900); 278 (ϵ 10000); 289 (ϵ 8700) (MeOH) (Berdy).

- Exp. reprod. and teratogenic effects. NL3150000

β -D-Glucopyranosyl ester: [19817-95-9]

$\text{C}_{16}\text{H}_{19}\text{NO}_7$ 337.329

Metab. of 1H-Indole-3-acetic acid, I-81 in *Arabidopsis* sp. and other plants. Isol. from rice (*Oryza sativa*) and other plants.

Me ester: [1912-33-0]

$\text{C}_{11}\text{H}_{11}\text{NO}_2$ 189.213

Isol. from immature seeds of beach pea (*Lathyrus maritimus*), *Vicia amurensis*, wild soybean (*Glycine soja*), lobiya (*Vigna catiang* var. *sinensis*) and hyacinth bean (*Dolichos lablab*). Mp 125° (as picrate).

Et ester: [778-82-5]

Mp 44-45°. Bp₁ 164-166°.

- NL3528000

Chloride: [50720-05-3]

$\text{C}_{10}\text{H}_8\text{ClNO}$ 193.632

Flakes (Et_2O). Mp 63-65°.

Amide: 1H-Indole-3-acetamide

[879-37-8]

$\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}$ 174.202

Isol. from etiolated seedlings of the black gram (*Phaseolus mungo*). Also from the sponges *Dysidea etheria* and *Ulosa ruetzleri*. Plant growth regulator. Mp 150-151°.

Hydrazide: [5448-47-5]

$\text{C}_{10}\text{H}_{11}\text{N}_3\text{O}$ 189.216

Mp 138-139°.

- NL3600000

Nitrile: 1H-Indole-3-acetonitrile. 3-Cyanomethyl-1H-indole

[771-51-7]

$\text{C}_{10}\text{H}_8\text{N}_2$ 156.187

Myrosinase-induced hydrol. prod. of indole glucosinolates, found in cabbage and other crucifers. Antineoplastic agent. Plant growth hormone. Mp 36.5-37°. Bp_{0.2} 157°. Log P 1.55 (calc). Functions by enzymic hydrol. to the acid.

- LD₅₀ (rat, scu) 255 mg/kg. AM0700000

N-Benzenesulfonyl: [945401-04-7]

$\text{C}_{16}\text{H}_{13}\text{NO}_4\text{S}$ 315.349

Solid. Mp 171-173°.

N-Me: [1912-48-7]

$\text{C}_{11}\text{H}_{11}\text{NO}_2$ 189.213

Mp 128°. Bp_{0.1} 130-150°.

N-Me, amide:

$\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}$ 188.229

Solid.

N-Me, nitrile: [51584-17-9]

$\text{C}_{11}\text{H}_{10}\text{N}_2$ 170.213

Cryst. Mp 58-60°.

N-β-D-Glucopyranosyl: [351346-16-2]

$\text{C}_{16}\text{H}_{19}\text{NO}_7$ 337.329

Constit. of redcurrants (*Ribes rubrum*), rice (*Oryza sativa*) and maize (*Zea mays*). Isol. from *Cortinarius brunneus*.

Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +18.1 (c, 0.23 in MeOH). λ_{max} 215; 267 (MeCN).

λ_{max} 222 (log ϵ 4.35); 273 (log ϵ 3.72) (MeOH).

N-β-D-Glucopyranosyl, Me ester: [23099-32-3]

$\text{C}_{17}\text{H}_{21}\text{NO}_7$ 351.355

Constit. of redcurrants (*Ribes rubrum*). Isol. from *Cortinarius brunneus*.

Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +4.7 (c, 0.06 in MeOH). λ_{max} 215; 267 (MeCN). λ_{max} 221 (log ϵ 4.12); 273 (log ϵ 3.45) (MeOH).

N-Methoxy, nitrile: 1-Methoxy-1H-indole-3-acetonitrile, 9CI. **Caulilexin C** [30536-48-2]

$\text{C}_{11}\text{H}_{10}\text{N}_2\text{O}$ 186.213

Alkaloid from the Japanese radish Daikon (*Raphanus sativus* var. *hortensis*) inoculated with *Pseudomonas cichorii* and from clubroots of Chinese cabbage (*Brassica pekinensis*) infected with *Plasmodiophora brassicae*. Also from *Brassica oleracea* var. *botrytis* (cauliflower). Phytoalexin.

N-tert-Butoxycarbonyl:

$\text{C}_{15}\text{H}_{17}\text{NO}_4$ 275.304

Solid. Mp 117-120°.

N-tert-Butoxycarbonyl, Me ester:

$\text{C}_{16}\text{H}_{19}\text{NO}_4$ 289.33

Solid. Mp 53°.

[2338-19-4]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 667C; 675D; 676D; 678C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 138B; 149A; 154A; 159A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1503C (ir)

Henbest, H.B. et al., *J.C.S.*, 1953, 3796-3801 (isol, nitrile)

Stowe, B.B. et al., *Prog. Chem. Org. Nat. Prod.*, 1959, **17**, 248-297 (rev, bibl)

Isogai, Y. et al., *Chem. Pharm. Bull.*, 1963, **11**, 1217-1218 (isol, amide)

Johnson, H.E. et al., *J.O.C.*, 1963, **28**, 1246-1248 (synth)

Stefanescu, P. et al., *Rev. Chim. (Bucharest)*, 1969, **20**, 65; *CA*, **69**, 30311 (synth)

Nomoto, M. et al., *Agric. Biol. Chem.*, 1970, **34**, 1590-1592 (*Caulilexin C*, isol)

Abe, H. et al., *Agric. Biol. Chem.*, 1972, **36**, 2259-2260 (isol, uv, ms)

Karrer, W. et al., *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, nos. 2514; 2515 (bibl)

Suvorov, N.N. et al., *CA*, 1973, **79**, 146320g (synth)

Chandrasekhar, K. et al., *Acta Cryst. B*, 1982, **38**, 2534-2535 (cryst struct)

Vekrel, J. et al., *Spectrochim. Acta A*, 1983, **39**, 887-894 (ir, pmr, ms)

Acheson, R.M. et al., *J. Chem. Res., Synop.*, 1984, 101 (*Caulilexin C*, synth)

Somei, M. et al., *Heterocycles*, 1985, **23**, 1101-1106 (*Caulilexin C*, synth)

Cardellina, J.H. et al., *J. Nat. Prod.*, 1986, **49**, 1065-1067 (isol, amide)

Pfeiffer, D. et al., *Cryst. Res. Technol.*, 1987, **22**, KI (cryst struct)


Raucher, S. et al., *J.A.C.S.*, 1987, **109**, 442-446 (chloride)

Morales-Rios, M.S. et al., *Magn. Reson. Chem.*, 1987, **25**, 377-395 (cmr)

Katayama, N. et al., *Plant Cell Physiol.*, 1987, **28**, 383-386 (isol, Me ester)

- Wall, M.E. *et al.*, *J. Nat. Prod.*, 1988, **51**, 129-135 (isol, nitrile)
Agrochemicals Handbook, 3rd edn., Royal Society of Chemistry, 1992, A232
 Nagarathnam, D. *et al.*, *J. Het. Chem.*, 1992, **29**, 953-958 (synth)
 Fukuyama, T. *et al.*, *J.A.C.S.*, 1994, **116**, 3127-3128 (synth)
 Faul, M. *et al.*, *J.O.C.*, 1998, **63**, 6053-6058 (*N-Me*, amide, synth, pmr, cmr, uv, ir)
 Shimada, A. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 187-189 (isol, activity)
 Tsotinis, A. *et al.*, *J. Med. Chem.*, 2006, **49**, 3509-3519 (*N-Me* nitrile)
Pesticide Manual, 14th edn., 2006, No. 476
 Pedras, M.S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1503-1509 (*Caulilexin C*)
 Schwarz, B. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 1405-1410 (*N-glucosyl*)
 Knör, S. *et al.*, *J. Med. Chem.*, 2007, **50**, 4329-4339 (*N-Boc*)
 Kai, K. *et al.*, *Phytochemistry*, 2007, **68**, 2512-2522 (*N-glucosyl*, *glucosyl ester*)
 Roy, S. *et al.*, *Synth. Commun.*, 2007, **37**, 1879-1886 (*N-benzenesulfonyl*)
 Teichert, A. *et al.*, *Chem. Biodiversity*, 2008, **5**, 664-669 (*N-glucosyl*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 11th edn., J. Wiley, 2004, ICN000; ICW000

Agelenopsis aperta Indoleacetylptentamine toxins **I-82**



Agel 416 m = n = q = 3, p = 4
 Agel 416a m = n = p = 3, q = 4
 Agel 416b m = 4, n = p = q = 3
 Agel 416c m = p = q = 3, n = 4

C₂₃H₄₀N₆O 416.609
 Acylpolyamine alkaloid complex isol. from the venom of the spider *Agelenopsis aperta*.

Agel 416
AG 416

- [133805-31-9]
 C₂₃H₄₀N₆O 416.609
 Isol. from the venom of the spider *Agelenopsis aperta*.
*N*²-Hydroxy: **Agel 432f**. *AG 432f*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*.
Hydroxy(2): **Agel 432**. *AG 432*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not determined.
Hydroxy(3): **Agel 432b**. *AG 432b*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not determined.
*N*²,*4*-Dihydroxy: **Agel 448**. *AG 448*. *PB 448*
 [128550-00-5]
 C₂₃H₄₀N₆O₃ 448.607
 From *Agelenopsis aperta* and *Paracaelotes birulae*.
*N*²,*N*³,*4*-Trihydroxy: **Agel 464a**. *AG 464a*
 C₂₃H₄₀N₆O₄ 464.607
 From *Agelenopsis aperta*.

Agel 416a
AG 416a

- C₂₃H₄₀N₆O 416.609
 Isol. from the venom of the spider *Agelenopsis aperta*.
*N*²-Hydroxy: **Agel 432g**. *AG 432g*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*.
Hydroxy(2): **Agel 432a**. *AG 432a*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not determined.
Hydroxy(3): **Agel 432c**. *AG 432c*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not determined.
*N*²,*4*-Dihydroxy: **Agel 448a**. *AG 448a*
 C₂₃H₄₀N₆O₃ 448.607
 From *Agelenopsis aperta*.
*N*²-(*3*-Aminopropyl), *N*²-hydroxy: **Agel 489**. *AG 489*
 [128549-96-2]
 C₂₆H₄₇N₇O₂ 489.703
 Alkaloid from the venom of the spider *Agelenopsis aperta*.
*N*⁵-(*3*-Aminopropyl), *N*⁴-hydroxy: **Agel 489b**. *AG 489b*
 C₂₆H₄₇N₇O₂ 489.703
 Alkaloid from the venom of *Agelenopsis aperta*.
*N*⁵-(*3*-Aminopropyl), *N*²,*N*³-dihydroxy: **Agel 505b**. *AG 505b*
 C₂₆H₄₇N₇O₃ 505.702
 Alkaloid from the venom of *Agelenopsis aperta*.
*N*²-(*3*-Aminopropyl), *N*²,*4*-dihydroxy: **Agel 505**. *AG 505*
 [128549-97-3]
 [128550-22-1]
 C₂₆H₄₇N₇O₃ 505.702
 Alkaloid from the venom of the spider *Agelenopsis aperta*.
*N*⁵-(*3*-Aminopropyl), *N*²,*N*²,*4*-trihydroxy: **Agel 521**. *AG 521*
 C₂₆H₄₇N₇O₄ 521.702
 Alkaloid from the venom of *Agelenopsis aperta*.
*N*⁵-(Aminoiminomethyl), *N*²,*4*-dihydroxy: **PB 490**
 C₂₄H₄₂N₈O₃ 490.648
 Isol. from the venom of the spider *Paracaelotes birulai*.

Agel 416b
AG 416b

- HO 416b*
 [133823-88-8]
 C₂₃H₄₀N₆O 416.609
 Isol. from the venom of the spiders *Agelenopsis aperta* and *Hololena curta*.
*N*²-Hydroxy: **Agel 432h**. *AG 432h*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*.
*N*³-Hydroxy: **Agel 432i**. *AG 432i*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*.
Hydroxy(3): **Agel 432d**. *AG 432d*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not


- determined.
Hydroxy(4): **Agel 432e**. *AG 432e*
 C₂₃H₄₀N₆O₂ 432.608
 From *Agelenopsis aperta*. OH group attached to indole ring but posn. not determined.
*N*²,*4*-Dihydroxy: **Agel 448b**. *AG 448b*
 C₂₃H₄₀N₆O₃ 448.607
 From *Agelenopsis aperta*.
*N*²,*N*³-Dihydroxy: **Agel 448c**. *AG 448c*
 C₂₃H₄₀N₆O₃ 448.607
 From *Agelenopsis aperta*.
*N*²,*N*³,*4*-Trihydroxy: **Agel 464**. *AG 464*
 C₂₃H₄₀N₆O₄ 464.607
 From *Agelenopsis aperta*.

Agel 416c
AG 416c

- C₂₃H₄₀N₆O 416.609
 Isol. from the venom of the spider *Agelenopsis aperta*.
 Jasys, V.J. *et al.*, *J.A.C.S.*, 1990, **112**, 6696-6704 (*AG505*, *AG489*)
 Schulz, S. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 314 (rev)
 Hidai, Y. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1570-1576 (*AG 416*, *AG 489*, synth, pmr, cmr, ms)
 Chesnov, S. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 3295-3305; 2001, **84**, 2178-2197; 2002, **85**, 2827-2846 (isol, synth, ms, bibl)

1H-Indole-3-butanoic acid, **I-83**
9CI

4-Indol-3-ylbutyric acid, *ISO*. *Hormex*. *Hormodin*. *IBA*. *Seradix*
 [133-32-4]




- C₁₂H₁₃NO₂ 203.24
 Widespread in plants. Plant growth regulator: "Rooting powder". Cryst. (C₆H₆/petrol). Insol. H₂O; sol. Me₂CO, Et₂O, EtOH. Mp 124°.
 ▶ LD₅₀ (mus, ori) 100 mg/kg. NL5250000
Me ester: [15591-70-5]
 C₁₃H₁₅NO₂ 217.267
 Cryst. (MeOH aq.). Mp 73-74°. Bp₆ 230°.
Et ester: [49850-32-0]
 C₁₄H₁₇NO₂ 231.294
 Cryst. (cyclohexane). Mp 39-40°.
Hydrazide: [27086-07-3]
 C₁₂H₁₅N₃O 217.27
 Cryst. (EtOH). Mp 112°.
Amide: [6245-91-6]
 C₁₂H₁₄N₂O 202.255
 Solid. Mp 86-87°.
 [110627-85-5]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 668A (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 139A (nmr)
 Jackson, R.W. *et al.*, *J.A.C.S.*, 1930, **52**, 5029-5035 (synth, *Me ester*, hydrazide)
 Suvarov, N.N. *et al.*, *CA*, 1954, **48**, 12078
 Bullock, M.W. *et al.*, *J.A.C.S.*, 1956, **78**, 5854-5857 (*Me ester*, *Et ester*)

- Avramenko, V.G. *et al.*, *CA*, 1970, **73**, 14615 (synth)
- Bowman, R.E. *et al.*, *Chem. Ind. (London)*, 1971, 154 (synth)
- Sharma, J.P. *et al.*, *Talanta*, 1976, **23**, 841 (detn)
- Harris, R.L.N. *et al.*, *Aust. J. Plant Physiol.*, 1977, **4**, 235 (activity)
- Chandrasekhar, K. *et al.*, *Acta Cryst. B*, 1980, **36**, 1165 (cryst struct)
- Wurst, M. *et al.*, *J. Chromatogr.*, 1984, **286**, 237 (hplc)
- Soriano-Garcia, M. *et al.*, *Rev. Latinoam. Quim.*, 1984, **15**, 64 (cryst struct)
- Soriano-Garcia, M. *et al.*, *J. Crystalllogr. Spectrosc. Res.*, 1987, **17**, 207 (cryst struct, picrate)
- Morales-Rios, M.S. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 377 (cmr)
- Pesticide Manual*, 9th edn., 1991, No. 7270
- Agrochemicals Handbook*, 3rd edn., Royal Society of Chemistry, 1992, A231
- Epstein, E. *et al.*, *Physiol. Plant.*, 1993, **88**, 382 (rev, occur, synth, metab)
- Mewshaw, R.E. *et al.*, *J. Med. Chem.*, 2004, **47**, 3823-3842 (amide)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ICP000


N-(1H-Indole-3-carbonyl)urea I-84

N-(Aminocarbonyl)-1H-indole-3-carboxamide, 9CI
[159308-53-9]



C₁₀H₉N₃O₂ 203.2
Metab. from the sponge *Zyssa massalis*.
Mancini, I. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1886-1894 (isol, uv, pmr, cmr, ms, struct)

1H-Indole-3-carbothioic acid, 9CI I-85



C₉H₇NOS 177.226

OH-form

N-Me, amide: 1-Methyl-1H-indole-3-carbothioamide
[24662-21-3]
C₁₀H₁₀N₂S 190.268
Yellow solid. Mp 125-126°.


SH-form

Me ester: S-Methyl 1H-indole-3-carbothioate. Methyl 1H-indolethiolcarboxylate
[54584-28-0]
C₁₀H₉NOS 191.253
Isol. from the marine-derived *Oceani-bulbus indolifex* Hel 45. Cryst. (C₆H₆/petrol). Mp 164-166°.

- Barbero, M. *et al.*, *Synthesis*, 1988, 300-302 (synth, pmr, cmr)
- Moody, C.J. *et al.*, *Anti-Cancer Drugs*, 1997, **8**, 489-499 (N-Me amide)
- Wagner-Döbler, I. *et al.*, *Int. J. Syst. Evolut. Microbiol.*, 2004, **54**, 1177-1184 (isol)

1H-Indole-3-carboxaldehyde, I-86

9CI
Indole-3-aldehyde. 3-Formylindole
[487-89-8]



C₉H₇NO 145.16
Alkaloid from the stem bark of *Murraya exotica* and from the red alga *Botryocladia leptopoda*. Also found in barley and tomato seedlings, cotton, *Pseudomonas syringae*, *Rhizobium* sp. and the coral *Tubastrea coccinea*. Mp 180° Mp 197-199°. pK_{a2} 12.36 (25°, NH).

▶ NL5993600

N-Methoxy: 1-Methoxy-1H-indole-3-carboxaldehyde, 9CI. 1-Methoxy-3-formylindole
[67282-55-7]
C₁₀H₉NO₂ 175.187
Stress metab. from the Japanese radish Daikon (*Raphanus sativus* var. *hortensis*) inoculated with *Pseudomonas cichorii*. Prisms + ½H₂O (Et₂O/hexane). Mp 50-51°. Bp_{0.06} 109-110°.
Schneider, E.A. *et al.*, *J. Exp. Bot.*, 1972, **23**, 152-170 (occur)
Acheson, R.M. *et al.*, *J.C.S. Perkin I*, 1978, 1117-1125 (N-methoxy, synth)
Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 677-681 (N-methoxy, synth)
Monde, K. *et al.*, *Phytochemistry*, 1995, **39**, 581-586 (1-Methoxy-3-indolecarboxaldehyde, isol)

1H-Indole-5-carboxaldehyde I-87

5-Formylindole
[1196-69-6]
C₉H₇NO 145.16
Alkaloid from *Monodora angolensis*. Mp 99-101°. n_D²⁰ 1.4196.

Ethylene acetal: 5-(1,3-Dioxolan-2-yl)-1H-indole, 9CI
[152879-74-8]
C₁₁H₁₁NO₂ 189.213
Light yellow cryst. (toluene). Mp 127-128°.

- Troxler, F. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1616 (synth)
- Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4116 (synth, ir, pmr, bibl)
- Moyer, M.P. *et al.*, *J.O.C.*, 1986, **51**, 5106 (synth, pmr)
- Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (isol, pmr, cmr)
- Cody, W.L. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 59-68 (synth)
- Fetter, J. *et al.*, *J. Het. Chem.*, 2005, **42**, 137-139 (synth, ethylene acetal)

1H-Indole-7-carboxaldehyde I-88

7-Formylindole
[1074-88-0]
C₉H₇NO 145.16
Isol. from roots of *Esenbeckia leiocarpa*. Needles (hexane). Mp 87-89°.
Troxler, F. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1616-1628 (synth)

Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4116-4125 (synth, pmr)

Moyer, M.P. *et al.*, *J.O.C.*, 1986, **51**, 5106-5110 (synth, pmr)

Delle Monache, *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435-439 (isol, uv, ir, pmr, cmr, ms)

Uchil, V.R. *et al.*, *Synth. Commun.*, 2006, **36**, 1051-1056 (synth, pmr)

Kochnev, A.I. *et al.*, *Russ. J. Org. Chem. (Engl. Transl.)*, 2007, **43**, 571-575 (synth, pmr)

Mohanakrishnan, A.K. *et al.*, *Synth. Commun.*, 2007, **37**, 4343-4352 (synth)

1H-Indole-2-carboxylic acid I-89

[1477-50-5]

C₉H₇NO₂ 161.16
Alkaloid from *Strychnos cathayensis*. Cryst. (C₆H₆ or Et₂O/petrol). Mp 203°. pK_{a2} 17.13 (25°, NH, KOH aq.).

Cheng, M.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 235-239 (isol)

1H-Indole-3-carboxylic acid I-90

Indole-β-carboxylic acid
[771-50-6]
C₉H₇NO₂ 161.16
Present in plants, e.g. apple (*Pyrus malus*), garden pea (*Pisum sativum*), *Brassica* spp. and the marine algae *Undaria pinnatifida* and *Botryocladia leptopoda*. Mp 210-218° (198-200°). pK_{a1} 3.87; pK_{a2} 15.59 (25°, NH). λ_{max} 270; 280; 288 (EtOH).

α-L-Rhamnopyranosyl ester: [309297-78-7]
C₁₅H₁₇NO₆ 307.302
Prod. by *Streptomyces* sp. GT 61150. Light yellow powder (CHCl₃/MeOH). Mp 132-133°. [α]_D²⁵ -36.8 (c, 0.1 in MeOH). λ_{max} 211 (log ε 3.45); 227 (sh) (log ε 1.56); 284 (log ε 0.97) (MeOH).

[α-L-Rhamnopyranosyl-(1→6)-β-D-glucopyranosyl] ester:
C₂₁H₂₇NO₁₁ 469.444
Constit. of the roots of *Clematis manshurica*. Amorph. powder. Mp 130-132°. [α]_D²⁰ -121 (c, 2 in MeOH). λ_{max} 210; 228; 285; 322 (MeOH).

Me ester: [942-24-5]
C₁₀H₉NO₂ 175.187
Isol. from the red alga *Botryocladia leptopoda* and a *Spongosorites* sp. Phytotoxin. Mp 147-148° (140°). Probably an artifact.

Et ester: [776-41-0]
C₁₁H₁₁NO₂ 189.213
Mp 122-124°.

Amide: 1H-Indole-3-carboxamide, 9CI
[1670-85-5]
C₉H₈N₂O 160.175
Isol. from the sponge *Zyssa massalis*.
Solid (EtOAc/hexane). Mp 201-203°.

Nitrile: 1H-Indole-3-carbonitrile, 9CI. 3-Cyanoindole
[5457-28-3]
C₉H₆N₂ 142.16
Alkaloid from a facultatively anaerobic, halophilic bacterium. Mp 178°.

N-Ac, nitrile:
C₁₁H₈N₂O 184.197
Mp 202°.

N-Ethoxycarbonyl, Et ester: Diethyl 1H-indole-1,3-dicarboxylate
[13328-43-3]
C₁₄H₁₅NO₄ 261.277
Cryst. (EtOH). Mp 102-104°.

N-Me: 1-Methyl-1H-indole-3-carboxylic acid
[32387-21-6]
C₁₀H₉NO₂ 175.187
Cryst. (2-propanol) or plates (Me₂CO aq.). Mp 212° (200-201° dec.). λ_{max} 215 ; 243 (sh) ; 286 ; 298 (sh) (MeOH).

N-Me, Me ester: [108438-43-3]
C₁₁H₁₁NO₂ 189.213
Solid. Mp 85-87°.

N-Me, Et ester: [56559-60-5]
C₁₂H₁₃NO₂ 203.24
Mp 69.7-70.2°.

N-Me, amide: [118959-44-7]
C₁₀H₁₀N₂O 174.202
Solid. Mp 176-177°.

N-Me, nitrile: 3-Cyano-1-methyl-1H-indole
[24662-37-1]
C₁₀H₈N₂ 156.187
Cryst. (Et₂O/hexane). Mp 60-61°.

N-(1,1-Dimethyl-2-propenyl), Me ester:
[306776-02-3]
C₁₅H₁₇NO₂ 243.305
Alkaloid from the fungus *Aporpium caryae*. Antifungal agent. Oil. λ_{max} 240 (log ε 3.05); 244 (log ε 3.05); 288 (log ε 3.03) (CH₂Cl₂).

N-(2,3-Dihydroxy-1,1-dimethylpropyl)(S-), Me ester: [306776-03-4]
C₁₅H₁₉NO₄ 277.319
Alkaloid from the fungus *Aporpium caryae*. Oil. [α]_D -11.8 (c, 0.21 in CH₂Cl₂). λ_{max} 242 (log ε 3.05); 248 (log ε 3.05); 284 (log ε 3.03) (CH₂Cl₂).

N-Benzyl: [27018-76-4]
C₁₆H₁₃NO₂ 251.284
Cryst. (EtOAc/hexane). Mp 194-196°
Mp 198-204°.

N-Hydroxy: 1-Hydroxy-1H-indole-3-carboxylic acid, 9CI
[18377-48-5]
C₉H₇NO₃ 177.159
Pale orange solid. Mp 135-137° dec.

N-Hydroxy, amide: [69111-90-6]
C₉H₈N₂O₂ 176.174

Green-brown rods (H₂O). Mp 181-182° dec.


N-Hydroxy, nitrile: 1H-Indole-3-carbonitrile N-oxide
[69111-88-2]
C₉H₆N₂O 158.159
Cryst. (C₆H₆/hexane). Mp 127-128°.

N-Methoxy, Me ester: Methyl 1-methoxy-1H-indole-3-carboxylate
[18377-50-9]
C₁₁H₁₁NO₃ 205.213
Prod. by Japanese horseradish (*Wasabia japonica*). Phytoalexin. Antifungal agent. Prisms (CHCl₃). Mp 45-46° (40-41°) Mp 40-41°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 138A; 158B (nmr)
Whalley, W.B. *et al.*, *J.C.S.*, 1954, 1651-1653 (*N-Me, synth*)
Millich, F. *et al.*, *J.O.C.*, 1958, 23, 1096-1102 (*N-Me, synth*)
Hart, G. *et al.*, *J.O.C.*, 1962, 27, 2940-2942 (*N-Me, synth*)
Kasperek, S. *et al.*, *Can. J. Chem.*, 1966, 40, 2805-2811 (*synth*)
Glombitza, K.W. *et al.*, *Planta*, 1966, 69, 135-149 (*biosynth*)
Katner, A.S. *et al.*, *Org. Prep. Proced. Int.*, 1970, 2, 297-303 (*synth*)
Aldridge, D.C. *et al.*, *J.C.S.(C)*, 1971, 1623-1627 (*isol*)
Abe, H. *et al.*, *Agric. Biol. Chem.*, 1972, 36, 2259-2260 (*isol, uv, ms*)
Acheson, R.M. *et al.*, *J.C.S. Perkin I*, 1978, 1117-1125 (*N-hydroxy*)
Mehta, G. *et al.*, *Synthesis*, 1978, 374-376 (*synth, amide, nitrile*)
Acheson, R.M. *et al.*, *J. Chem. Res., Synop.*, 1984, 101; *J. Chem. Res., Miniprint*, 1984, 1301-1319 (*N-hydroxy, N-methoxy Me ester, synth*)

Plate, R. *et al.*, *Tetrahedron*, 1986, 42, 6511-6518 (*N-Me, uv*)
Bano, S. *et al.*, *Planta Med.*, 1987, 53, 117-118 (*isol, uv, pmr, cmr, ms, esters*)
Baiocchi, L. *et al.*, *J. Het. Chem.*, 1988, 25, 1905-1909 (*synth, N-Me, N-benzyl*)
Swain, C.J. *et al.*, *J. Med. Chem.*, 1992, 35, 1019-1031 (*N-Me, nitrile*)
Buttery, C.D. *et al.*, *J.C.S. Perkin I*, 1993, 1425-1431 (*N-Me, synth, ir, pmr*)
Mancini, I. *et al.*, *Helv. Chim. Acta*, 1994, 77, 1886-1894 (*amide, isol*)
Vorbrüggen, H. *et al.*, *Tetrahedron*, 1994, 50, 6549-6558 (*nitrile*)
Fu, X. *et al.*, *J. Nat. Prod.*, 1995, 58, 1950-1954 (*isol, nitrile*)
Prashad, M. *et al.*, *Synth. Commun.*, 1995, 25, 95-100 (*Me ester, synth*)
Moody, C.J. *et al.*, *Anti-Cancer Drugs*, 1997, 8, 489-499 (*N-Me Me ester, N-Me amide*)
Pedras, M.S.C. *et al.*, *Phytochemistry*, 1998, 49, 1959-1965 (*1-methoxy Me ester*)
Hu, J.-F. *et al.*, *J. Antibiot.*, 2000, 53, 944-953 (*rhamnosyl ester*)
Levy, L.M. *et al.*, *Phytochemistry*, 2000, 54, 941-943 (*N-1,1-dimethylpropyl derivs, synth*)
Somei, M. *et al.*, *Heterocycles*, 2001, 54, 425-432 (*1-methoxy Me ester, synth*)
Everett, S.A. *et al.*, *J.C.S. Perkin 2*, 2001, 1989-1997 (*N-Me amide*)
Sala, G.D. *et al.*, *Tet. Lett.*, 2002, 43, 8839-8841 (*N-1,1-dimethylpropyl derivs, synth*)
Hu, S.-C. *et al.*, *Acta Cryst. E*, 2005, 61, 01654-01656 (*Me ester, cryst struct*)
Shi, S.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, 8, 73-78 (*6-rhamnosylglucosyl ester*)
Bao, B. *et al.*, *Mar. Drugs*, 2007, 5, 31-39 (*Me ester, isol*)

1H-Indole-2,3-diol I-91
1,3-Dihydro-3-hydroxy-2H-indol-2-one, 9CI. Dioxindole. o-Aminomandelic lactam. 3-Hydroxyoxindole. 2,3-Dihydroxy-yindole
[61-71-2]
[5638-85-7]



C₈H₇NO₂ 149.149

Oxindole form predominates. Prod. by a marine bacterial strain isol. from gills of *Sphyræna japonica* and prod. by a range of microorganisms present in fish and shellfish. Prod. of metab. of Indole, I-79 by microorganisms. Antioxidant. Cryst. (H₂O or EtOH). Mp 167-168°.

▶ NM3289460

Hydrochloride: Mp 156°.

Acyl deriv.: Isatan C

[397243-11-7]
C₂₀H₁₃NO₈ 395.325
Isol. from the leaves of *Isatis tinctoria*. Precursor of Indigotin, I-72. Full struct. not determined. Ester with a (C₁₁H₇O₅)-CO acyl group. Not indexed by CA. λ_{max} 218 ; 280 (no solvent reported).

Fujioka, M. *et al.*, *Biochim. Biophys. Acta*, 1965, 158, 70-78 (*isol*)

Maugard, T. *et al.*, *Phytochemistry*, 2001, 58, 897-904 (*Isatan C*)

1H-Indole-2,5-diol I-92
1,3-Dihydro-5-hydroxy-2H-indol-2-one, 9CI. 5-Hydroxyoxindole. 2,5-Dihydroxy-1H-indole
[3416-18-0]

C₈H₇NO₂ 149.149

Constit. of the leaves of *Isatis tinctoria*. Metab. of 1H-Indol-2-ol, I-103, excreted in urine. Cryst. (H₂O or by subl.), needles (EtOAc/petrol). Mp 270° (265-266°) dec.

Me ether: 1,3-Dihydro-5-methoxy-2H-indol-2-one. 5-Methoxyoxindole

[7699-18-5]
C₉H₉NO₂ 163.176
Cryst. (EtOH). Mp 156-157° (153-154°).

N,O-Di-Me: 1,3-Dihydro-5-methoxy-1-methyl-2H-indol-2-one
C₁₀H₁₁NO₂ 177.202
Cryst. Mp 97-98.5° (91-92.5°).

Beer, R.J. *et al.*, *J.C.S.*, 1953, 1262 (*synth*)

Becket, A.H. *et al.*, *Tetrahedron*, 1968, 24, 6093 (*synth, ir, uv, pmr, deriv*)

Reio, L. *et al.*, *J. Chromatogr.*, 1974, 88, 119 (*chromatog*)

Bocelli, G. *et al.*, *J. Mol. Struct.*, 1982, 96, 127 (*cryst struct*)

Sakamoto, T. *et al.*, *J. Het. Chem.*, 1988, 25, 1279 (*synth, use*)

Andreani, A. *et al.*, *J. Het. Chem.*, 1988, 25, 1519 (*deriv*)


Crestini, C. *et al.*, *Synth. Commun.*, 1994, 24, 2835 (*Me ether*)

Li, L. *et al.*, *CA*, 1996, 125, 270492x (*isol*)

Smith, K. *et al.*, *J.C.S. Perkin I*, 1996, 2793 (*N,O-di-Me*)

Li, B. *et al.*, *Yaouxue Xuebao*, 2000, **35**, 508-510 (isol)

1H-Indole-5,6-diol, 9CI I-93
5,6-Dihydroxyindole. Dopamine lutine [3131-52-0]



C₈H₇NO₂ 149.149

Alkaloid from the leaves of *Rhaphidophora korthalsii* and *Epipremnum pinnatum*. Cytotoxic agent. Intermed. in the tyrosinase-catalysed biosynth. pathway from tyrosine to melanin. Needles (C₆H₆/petrol). Sol. hot H₂O; insol. petrol. Mp 140° dec. Unstable on storage, undergoes spont. oxidn. and condensation to melanin.

► Toxic.

N,6-O-Disulfo: Ancorinolate B
[473740-08-8]

C₈H₇NO₈S₂ 309.277

Alkaloid from the sponge *Ancorina* sp. Powder (as di-Na salt). λ_{max} 222 (log ε 4.08); 265 (log ε 3.61); 296 (log ε 3.39) (MeOH) (di-Na salt).

Wong, K.T. *et al.*, *Nat. Prod. Lett.*, 1996, **9**, 137-140 (isol, ir, pmr, cmr, ms)

Meragelman, K.M. *et al.*, *J.O.C.*, 2002, **67**, 6671-6677 (*Ancorinolate B*)


1H-Indole-6,7-diol, 9CI I-94
6,7-Dihydroxyindole [151980-99-3]

C₈H₇NO₂ 149.149

Poss. marine alkaloid from several *Agelas* spp. Identified on the basis of colour reacs. only; struct. of the alkaloid requires reinvestigation. Could be the isomer 4,6-Dihydroxyindole.

Stempien, M.F. *et al.*, *Am. Zool.*, 1966, **6**, 363 (isol)

1H-Indole-2,3-dione, 9CI I-95
2,3-Indolinedione. *Isatin* [91-56-5]



C₈H₅NO₂ 147.133

Isol. from the fungal pigment of a mutant of *Schizophyllum commune*. Prod. by *Alteromonas* sp.12 and a marine *Halo- monas* sp. RK377. Also prod. by an *Alteromonas* sp. commensal in shrimp embryos *Palaemon macrodactylus* and in egg masses of the muricid mollusc *Trunculariopsis trunculus*. Intermed. for indigoid dyestuffs. Used as 0.2% MeOH soln. for photometric detn. of Thiophene, Proline, hydroxyproline. Chromatog. spray for amino acids. Shows antifungal activity in the marine environment.

Orange cryst. Spar. sol. H₂O. Mp 203.5° dec. Sublimes.

► Exp. reprod. effects. NL7873000

[73859-66-2, 73859-64-0]


Epstein, E. *et al.*, *CA*, 1966, **66**, 102443 (isol) Popp, F.D. *et al.*, *Adv. Heterocycl. Chem.*, 1975, **18**, 1-58 (rev)

Liang, L. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (marine, isol, pmr, cryst struct)

Smith, K. *et al.*, *Synthesis*, 2003, 2047-2051 (synth, ir, pmr, cmr, ms)

Pelosi, G. *et al.*, *Acta Cryst. C*, 2005, **61**, o589-o592 (cryst struct)

1H-Indole-3-ethanol, 9CI I-96
3-(2-Hydroxyethyl)indole. 2-(3-Indoly-1)ethanol. **Tryptophol** [526-55-6]



C₁₀H₁₁NO 161.203

Isol. from Tryptophan fermentations, plant seedlings and *Candida* spp., *Aspergillus niger*, *Agrobacterium tumefaciens*, *Ceratocystis* spp. and *Rhizobium* spp. Also isol. from the sponge *Ircinia spinulosa*. Plant auxin. Active against gram-positive bacteria and *Candida albicans*. Associated with fungal diseases of trees. Prisms (C₆H₆/petrol); plates (Et₂O/petrol). Sol. most org. solvs.; spar. sol. H₂O, petrol. Mp 59°. Bp₂ 174°. Red. soln. in warm conc. H₂SO₄.

► LD₅₀ (mus, ipr) 351 mg/kg. KL3685000

Picrate:

Red needles (H₂O). Mp 100-101°.

O-β-D-Galactopyranoside: Tryptophol galactoside [95088-52-1]

C₁₆H₂₁NO₆ 323.345

Isol. from *Euglena gracilis*.

O-β-D-Glucopyranoside: Tryptophol glucoside [40883-44-1]

C₁₆H₂₁NO₆ 323.345

Isol. from numerous plant and algae. Plant growth promotor.

O-[β-D-Xylopyranosyl-(1→6)-β-D-glucopyranoside]: [149817-65-2]

C₂₁H₂₉NO₁₀ 455.461

Isol. from *Lycium chinense* (Chinese boxthorn). Pale yellow powder. [α]_D²⁰ -32.7 (c, 0.4 in MeOH).

O-Ac: [13137-14-9]

C₁₂H₁₃NO₂ 203.24

Oil.

N-Benzoyl:

C₁₇H₁₅NO₂ 265.311

Yellow prisms and plates (petrol). Mp 76°.

O-Phenylacetyl: 2-(3-Indolyl)ethyl phenylacetate. Monaspiloinole

C₁₈H₁₇NO₂ 279.338

Isol. from *Monascus pilosus*. Oil. λ_{max} 280 (log ε 3.74) (MeOH).

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **2**, 659B (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **3**, 130A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1502A (ir)

Jackson, R.W. *et al.*, *J. Biol. Chem.*, 1930, **88**, 659

Snyder, H.R. *et al.*, *J.A.C.S.*, 1948, **70**, 3770-3771 (synth, bibl)

Nogrady, T. *et al.*, *Can. J. Chem.*, 1964, **42**, 485-486 (synth)

Narayanan, T.K. *et al.*, *Antimicrob. Agents Chemother.*, 1976, **9**, 375 (isol, props)

Magnus, V. *et al.*, *Carbohydr. Res.*, 1979, **76**, 261 (synth, glucoside)

Lacan, G. *et al.*, *Plant Physiol.*, 1984, **76**, 889; 1985, **78**, 447 (glucoside, galactoside)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1986, **64**, 904 (isol)

Yahara, S. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 703 (xylosylglucoside)

Novák, L. *et al.*, *Annalen*, 1995, 1877 (Ac)

Böhlendorf, B. *et al.*, *Annalen*, 1996, 49 (isol, ir, pmr, cmr, ms)


Erdogan, I. *et al.*, *Biochem. Syst. Ecol.*, 2000, **28**, 793-794 (isol, sponge)

Garcez, W.S. *et al.*, *J. Braz. Chem. Soc.*, 2005, **16**, 1382-1386 (glucoside, isol, pmr, cmr)

Cheng, M.-J. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 394-397 (*Monaspiloinole*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, ICS000

1H-Indole-3-methanamine I-97
3-Indolylmethanamine. 3-(Aminomethyl)indole. **N-Dinorgramine**. **N-Bisnorgramine** [22259-53-6]



C₉H₁₀N₂ 146.191

Detected in *Hordeum vulgare* (barley).

Cryst. (C₆H₆). Mp 103-104° (101-102°).

1'-N-Ac:

C₁₁H₁₂N₂O 188.229

Cryst. (C₆H₆). Mp 133-134°.

1'-N-Benzoyl:

C₁₆H₁₄N₂O 250.299

Mp 157-159°.

N^d-Me: 1-Methyl-1H-indole-3-methanamine. 3-(Aminomethyl)-1-methylindole. Donaxamine

[19293-60-8]

C₁₀H₁₂N₂ 160.218

Alkaloid from the aerial parts of

Arundo donax. Mp 178-179°.

1'-N-Me: 3-(Methylaminomethyl)indole. N-Norgramine

C₁₀H₁₂N₂ 160.218

Detected in *Hordeum vulgare* (barley).

Cryst. (EtOH)(as picrate). Mp 176-

176.5° (picrate).

1',1'-N-Di-Me: see 3-[(Dimethylamino)-methyl]indole, D-736

1-Methoxy, 1'-N-formyl: 3-(Formylaminomethyl)-1-methoxyindole. Caulilexin B


C₁₁H₁₂N₂O₂ 204.228

Alkaloid from *Brassica oleracea* var.

botrytis (cauliflower). Phytoalexin. Cryst. (CH₂Cl₂/MeOH). Mp 71-73°. λ_{max} 221 (log ε 4.4); 290 (log ε 3.7) (MeOH).

- Putochin, N. *et al.*, *Ber.*, 1926, **59**, 1987
 Mudd, S.H. *et al.*, *Nature (London)*, 1961, **189**, 489 (*N-Norgramine*)
 Gower, B.E. *et al.*, *J.A.C.S.*, 1963, **85**, 3683 (*synth*)
 Zhalolov, I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 528-530 (*Donaxamine*)
 Csomós, P. *et al.*, *Tetrahedron*, 2005, **61**, 9257-9262 (*synth, N-benzoyl*)
 Gaspari, P. *et al.*, *J. Med. Chem.*, 2006, **49**, 684-692 (*synth*)
 Pedras, M.S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1503-1509 (*Caulilexin B*)
 Kutschy, P. *et al.*, *Tetrahedron*, **54**, 3549-3566 (*synth, 1-Me*)

1H-Indole-3-methanol **I-98**
 3-Hydroxymethylindole. Indole-3-carbinol [700-06-1]




C₉H₉NO 147.176
 Isol. from cruciferous vegetables. Shown clinically to have anti-cancer props. Potential nutraceutical. Platelets (C₆H₆). Mp 89-91° Mp 158°. pK_{a1} 16.5 (25°, NH, KOH aq.). Mp dependent on rate of heating.

- Exp. teratogenic effects. NL9483000
 N-Ac:
 C₁₁H₁₁NO₂ 189.213
 Cryst. (C₆H₆/EtOAc). Mp 137-139°.
 O,N-Di-Ac:
 C₁₃H₁₃NO₃ 231.251
 Cryst. (EtOH). Mp 89-90°.
 N-Me: [6965-44-2]
 C₁₀H₁₁NO 161.203
 Needles. Mp 30°. Bp₅ 150-160°.
 ► NL9486900
 N-Me, trinitrobenzene complex:
 Cryst. (MeOH). Mp 139-141°.
 Mingoia, G. *et al.*, *Gazz. Chim. Ital.*, 1932, **62**, 844 (*synth*)
 Ames, D.E. *et al.*, *J.C.S.*, 1956, 1984 (*synth*)
 Leete, E. *et al.*, *J.A.C.S.*, 1959, **81**, 6023 (*N-methyl*)
 Mattocks, A.R. *et al.*, *J.C.S. Perkin 1*, 1978, 896 (*N-methyl*)
 Broadbent, T.A. *et al.*, *Curr. Med. Chem.*, 1998, **5**, 337-352; 469-491 (*rev. pharmacol*)
 Jeffery, E.H. *et al.*, *Handbook of Nutraceuticals and Functional Foods*, (ed. Wildman, R.E.C.), CRC Press., 2001, 169-191
 Goyal, R.N. *et al.*, *J.C.S. Perkin 2*, 2001, 618-623 (*props. bibl*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ICP100

1H-Indole-7-methanol, 9CI **I-99**
 7-Hydroxymethylindole [1074-87-9]
 C₉H₉NO 147.176
 Alkaloid from roots of *Esenbeckia leiocarpa*. Needles (CH₂Cl₂/hexane). Mp 56-56.5°.

- Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4116 (*synth*)
 Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (*isol, pmr, cmr, ms*)
 Uchil, V.R. *et al.*, *Synth. Commun.*, 2006, 1051-1056 (*synth, ir, pmr*)

1H-Indole-3-propanoic acid **I-100**
 3-(3-Indolyl)propionic acid. Antibiotic KG 431B. IPA. KG 431B [830-96-6]



C₁₁H₁₁NO₂ 189.213
 Isol. from *Claviceps* (ergot) cultures. Mp 134° (111-112°).


- NM1329000
 Me ester: [5548-09-4]
 C₁₂H₁₃NO₂ 203.24
 Mp 79-80°.
 ► NM1480000
 Cyclohexyl ester: **Indobinine** [109897-76-9]
 C₁₇H₂₁NO₂ 271.358
 Alkaloid from roots of *Rauwolfia serpentina* (Apocynaceae). Elongated rods (MeOH). Mp 180-181°.
 Benzyl ester: **Indobine** [85612-35-7]
 C₁₈H₁₇NO₂ 279.338
 Alkaloid from the roots of *Rauwolfia serpentina* (Apocynaceae). Needles (MeOH). Mp 163-164°.
 Amide: **1H-Indole-3-propanamide** [5814-93-7]
 C₁₁H₁₂N₂O 188.229
 Cryst. (MeOH aq.). Mp 134-136°.
 Methylamide: **N-Methyl-1H-Indole-3-propanamide**. 3-(3-Indolyl)-N-methylpropanamide [69397-85-9]
 C₁₂H₁₄N₂O 202.255
 Prod. by *Omphalotus olearius* (common chanterelle). Cryst. (CH₂Cl₂/petrol). Mp 97.5-99° (synthetic).
 Hydrazide: [20401-90-5]
 C₁₁H₁₃N₃O 203.243
 Mp 129-130°.

Nitrile: 3-(2-Cyanoethyl)indole [4414-76-0]
 C₁₁H₁₀N₂ 170.213
 Mp 67-68°.

- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 667D (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 138C (*nmr*)
 Yamano, T. *et al.*, *CA*, 1963, **58**, 763 (*isol*)
 Ghosal, S. *et al.*, *J. Indian Chem. Soc.*, 1964, **41**, 496 (*synth*)
 Bowman, R.E. *et al.*, *Chem. Ind. (London)*, 1971, 154 (*synth*)
 Siddiqui, S. *et al.*, *Indian J. Chem., Sect. B*, 1987, **26**, 279 (*Indobinine*)
 Siddiqui, S. *et al.*, *Z. Naturforsch., B*, 1987, **42**, 783 (*Indobine*)
 Thompson, A.M. *et al.*, *J. Med. Chem.*, 1994, **37**, 598-609 (*synth, amide*)
 Mayer, A. *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 25-32 (*methylamide*)

- Okabe, N. *et al.*, *Acta Cryst. C*, 1998, **54**, 386-387 (*cryst struct*)
 Mewshaw, R.E. *et al.*, *J. Med. Chem.*, 2004, **47**, 3823-3842 (*amide*)
 Black, P.J. *et al.*, *Eur. J. Org. Chem.*, 2006, 4367-4378 (*nitrile*)

Indolizomycin **I-101**
 Octahydro-5-(5-methyl-1,3,5-heptatrienyl)-6bH-cycloprop[a]oxireno[g]indolizine-6b-ol, 9CI [94935-24-7]




C₁₇H₂₃NO₂ 273.374
 Triene antibiotic. Isol. from *Streptomyces griseus*. Weakly active against gram-positive and gram-negative bacteria. Pale-yellow thick oil. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁰ -28.6 (c, 0.5 in MeOH). LD₅₀ 12.5 mg/kg (i.p. mice). Related to Cyclizidine. λ_{max} 260 (sh) (ε 29100); 268 (ε 33400); 277 (sh) (ε 27300) (MeOH) (Derep). λ_{max} 268 (ε 33400) (MeOH) (Berdy).
 ► LD₅₀ (mus, ipr) 12.5 - 25 mg/kg. GZ2247500

6b-Deoxy: Deoxyindolizomycin [94935-27-0]
 [α]_D²⁷ -12.6 (c, 1 in MeOH).

- Hotta, K. *et al.*, *J. Antibiot.*, 1984, **37**, 1491 (*isol*)
 Yamashita, F. *et al.*, *J. Antibiot.*, 1985, **38**, 58; 64 (*isol*)
 Kim, G. *et al.*, *J.A.C.S.*, 1993, **115**, 30 (*synth*)
 Nicolaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, 471 (*bibl, synth*)
 Groaning, M.D. *et al.*, *Tet. Lett.*, 1999, **40**, 4639-4642 (*synth*)

Indolmycin **I-102**
 5-[1-(1H-Indol-3-yl)ethyl]-2-(methylamino)-4(5H)-oxazolone, 9CI. PA 155A. Antibiotic PA 155A [21200-24-8]
 [20893-33-8 (Isoindolmycin), 21200-23-7]



Absolute Configuration

C₁₄H₁₅N₃O₂ 257.291
 Isol. from *Streptomyces albus* BA 3972 and *Streptomyces griseus* ATCC 12648. Shows activity against gram-positive bacteria including mycobacteria. L-Tryptophan antagonist. Prisms (MeOH or EtOAc). Sol. MeOH, Me₂CO, butanol,

EtOH; fairly sol. H₂O, C₆H₆, Et₂O; poorly sol. hexane. Mp 209-210°. [α]_D²⁵ -214 (c, 2 in MeOH).

► LD₅₀ (mus, scu) 800mg/kg.

Dipicrate:

Orange-red plates (EtOH/Et₂O). Mp 148-149°.

[13786-41-9 ±-form]

Schach von Wittenau, M. *et al.*, *J.A.C.S.*, 1963, **85**, 3425-3431 (*synth, struct*)

Preobrazhenskaya, M.N. *et al.*, *Tetrahedron*, 1968, **24**, 6131-6143 (*synth*)

Chan, T.H. *et al.*, *J.O.C.*, 1970, **35**, 3519-3521 (*abs config*)

Hornemann, V. *et al.*, *J.A.C.S.*, 1971, **93**, 3028-3035 (*biosynth*)

Hirota, A. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 147-151 (*isol, uv, ir, ms*)

Dirlam, J.P. *et al.*, *J.O.C.*, 1986, **51**, 4920-4924 (*synth*)

Akita, H. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 323-328 (*synth*)

Bando, T. *et al.*, *Heterocycles*, 1997, **46**, 111-114 (*synth*)


Hasuoka, A. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1604-1608 (*synth*)

Sutou, N. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 1833-1838 (*synth*)

1H-Indol-2-ol

I-103

1,3-Dihydro-2H-indol-2-one, 9CI. 2-Indolinone, 8CI. 2-Hydroxyindole. Oxindole [59-48-3]



C₈H₇NO 133.149

In solid state exists solely as oxo tautomer which also predominates in soln. Prod. by *Chromobacterium violaceum* and *Penicillium* spp. Also found in the oil of *Narcissus geranium*. Shows phospholipase A₂ inhibitory activity. Shows weak cytotoxic effect. Needles (H₂O). Mp 127°. Bp₂₃ 227°. Reduces Tollen's reagent.


► LD₅₀ (mus, orl) 400 mg/kg. NM2080500

Haun, M. *et al.*, *Biol. Res.*, 1992, **25**, 21 (*isol*)
Witter, L. *et al.*, *Z. Naturforsch., C*, 1998, **53**, 60-66 (*isol, activity*)

1H-Indol-3-ol, 9CI

I-104

1,2-Dihydro-3H-indol-3-one, 9CI. Indoxyl. Dihydro-3-oxoindole. 3-Hydroxyindole. Pseudoindoxyl



oxo-form

OH-form

C₈H₇NO 133.149

OH-form prob. favoured for parent.

Unstable yellow solid. Mp 85°. Readily oxidised in air, especially in alkaline soln. to Indigotin, I-72.

N-Ac: [33025-60-4]

C₁₀H₉NO₂ 175.187

Needles (H₂O). Mp 139° (135°).

Shown by ir studies to exist in soln. as oxo-form.

OH-form [480-93-3]

O-Sulfate:

C₈H₇NO₄S 213.214

Isol. from *Murex trunculus*.

O-β-D-Glucopyranoside: **Indican**. Plant indican. Uroanthin

[487-60-5]

C₁₄H₁₇NO₆ 295.291

Occurs in *Polygonum tinctoria*, *Isatis tinctoria*, *Celosia argentea* and *Indigofera* spp. Biosynth. precursor to Indigotin, I-72. Shows growth inhibitory activity on the germination of lettuce. Cryst. + 3H₂O (H₂O). Mp 57-58° (hydrate) Mp 180° (anhyd.). [α]_D¹⁵ -66.2 (H₂O).

O-β-D-ribo-Hexopyranos-3-uloside: **Isatan B**

[20307-14-6]

C₁₄H₁₅NO₆ 293.276

Constit. of *Isatis tinctoria*. Precursor of Indigotin, I-72. Beige amorph. powder. [α]_D²⁰ -14 (c, 1 in MeOH). Readily undergoes hydrol. Occurs as hydrate in aq. solns. λ_{\max} 223 (ε 32200); 281 (ε 5700) (MeOH).

O-(6-O-Malonyl-β-D-ribo-hexopyranos-3-uloside): **Isatan A**

C₁₇H₁₇NO₉ 379.323

Constit. of *Isatis tinctoria*. Precursor of Indigotin, I-72. Beige amorph. powder. [α]_D²⁰ -10.5 (c, 1 in MeOH). Readily undergoes hydrol. Occurs as hydrate in aq. solns. λ_{\max} 212 (log ε 2.03); 251 (log ε 3.51); 280 (log ε 4.61) (MeOH).

O-[β-D-Glucopyranosyl-(1→6)-β-D-glucopyranoside]: 3-Indolyl gentiobioside.

Glucoindican

C₂₀H₂₇NO₁₁ 457.433

Constit. of *Calanthe discolor* and *Calanthe liukiensis*. Powder. [α]_D²⁵ +164 (MeOH). λ_{\max} 224 (log ε 5.4); 282 (log ε 4.7) (no solvent reported).

O-Ac: 3-Acetoxy-1H-indole

[608-08-2]

C₁₀H₉NO₂ 175.187

Alkaloid from *Strychnos cathayensis*. Cryst. (H₂O). Mp 127.5°.

► LD₅₀ (mus, ipr) 600 mg/kg. AI3325000

1,3-Di-Ac: [16800-67-2]

C₁₂H₁₁NO₃ 217.224

Cryst. (H₂O). Mp 82°.

1-Benzoyl, 3-Ac: [90539-82-5]

C₁₇H₁₃NO₃ 279.295

Cryst. (EtOH). Mp 87°.

3-Benzoyl, 1-Ac: [110912-16-8]

C₁₇H₁₃NO₃ 279.295

Cryst. (2-propanol). Mp 158-160°.

1-Me, 3-Ac: [3260-63-7]

C₁₁H₁₁NO₂ 189.213

Cryst. (EtOH aq.). Mp 56.5°.

Et ether: 3-Ethoxy-1H-indole

[916756-92-8]

C₁₀H₁₁NO 161.203

Purple oil.

[3260-61-5, 487-94-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 675B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 148B (*nmr*)

Spencer, G. *et al.*, *J. Soc. Chem. Ind., London*, 1931, **50**, 63T (*N-Ac, 3-O-Ac, synth*)

Robertson, A. *et al.*, *J.C.S.*, 1933, 30 (*Indican*)

Holt, S.J. *et al.*, *J.C.S.*, 1958, 1217 (*N-Ac, conformn, ir*)

Kirby, G.W. *et al.*, *Chem. Comm.*, 1965, 381 (*synth*)

Raileanu, D. *et al.*, *Rev. Roum. Chim.*, 1967, **12**, 105; *CA*, **68**, 21775 (*N-Ac, di-Ac, synth*)

Baker, J.T. *et al.*, *Tet. Lett.*, 1968, 43; 1976, 1233-1234 (*sulfate*)

Fouquet, H. *et al.*, *Angew. Chem., Int. Ed.*, 1971, **10**, 816-817 (*sulfate, isol*)

Rodd's Chem. Carbon Compd. (2nd edn.), 1973, **4A**, 454 (*rev*)

Strobel, J. *et al.*, *Biochem. Physiol. Pflanz.*, 1989, **184**, 321 (*occur*)

Traldi, P. *et al.*, *Rapid Commun. Mass Spectrom.*, 1990, **4**, 335 (*ms*)

Xia, Z.-Q. *et al.*, *Phytochemistry*, 1992, **31**, 2695 (*Indican, Isatan B, biosynth, cmr*)

Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 886-888 (*Glucoindican*)

Sawabe, A. *et al.*, *Stud. Plant Sci.*, 1999, **6**, 290-296 (*Indican, isol, activity*)

Cheng, M.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2001, **48**, 235-239 (*3-Acetoxyindole, isol*)

Cooksey, C.J. *et al.*, *Molecules*, 2001, **6**, 736-769 (*sulfate, rev*)

Oberthür, C. *et al.*, *Chem. Biodiversity*, 2004, **1**, 174-182 (*Isatans A,B*)


Clawson, R.W. *et al.*, *Tetrahedron*, 2006, **62**, 10829-10834 (*Et ether*)

3H-Indolo[3,2,1-de][1,5]-naphthyridine-2,6-dione, 9CI

I-105

Canthin-2,6-dione

[86293-43-8]



C₁₄H₈N₂O₂ 236.229

Alkaloid from the wood of *Simaba multiflora* (Simaroubaceae). Yellow micro-needles. Mp 290-305°. λ_{\max} 227 (ε 27500); 247 (ε 21900); 253 (ε 20400); 291 (ε 9770); 300 (ε 10200); 400 (sh) (ε 8130); 421 (ε 12900); 445 (ε 13800) (MeOH) (Derep).

N³-Me: 3-Methylcanthin-2,6-dione

[82652-21-9]

C₁₅H₁₀N₂O₂ 250.256

Alkaloid from the roots of *Picrasma quassioides* (Simaroubaceae). Orange-red needles (CHCl₃/MeOH). Mp 330°.

Strong yellow-green fluor. in soln. λ_{\max} 227 (log ε 4.25); 242 (log ε 4.33); 250 (log ε 4.3); 287 (log ε 3.87); 345 (log ε 3.55); 444 (log ε 4.1); 462 (log ε 4.1) (EtOH).

N³-Methoxy: 3-Methoxycanthin-2,6-dione

[74991-92-7]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the wood of *Simaba multiflora* and the trunk bark of *Simaba cuspidata* var. *typica* (Simaroubaceae). Red cryst. (AcOH). Mp 330°. Strong green fluor. in soln. λ_{\max} 227 (€ 17400); 232 (sh) (€ 17400); 248 (€ 16600); 253 (sh) (€ 16200); 292 (€ 10200); 302 (€ 10200); 325 (€ 6460); 403 (sh) (€ 9330); 420 (€ 13500) (MeOH) (Derep). λ_{\max} 226 (€ 14500); 247 (€ 13500); 290 (€ 4300); 302 (€ 4500); 325 (€ 2700); 422 (€ 7100); 446 (€ 8000) (MeOH) (Berdy).

4-Methoxy, 5-hydroxy, N³-Me: 5-Hydroxy-4-methoxy-3-methylcanthin-2,6-dione

[129724-30-7]

C₁₆H₁₂N₂O₄ 296.282

Alkaloid from the wood of *Quassia amara* (Surinam quassia) (Simaroubaceae). Orange-red needles (MeOH). Mp 250-254°.

10-Hydroxy, N³-methoxy: 10-Hydroxy-3-methoxycanthin-2,6-dione

[86293-42-7]

C₁₅H₁₀N₂O₄ 282.255

Alkaloid from the wood of *Simaba multiflora* (Simaroubaceae). Yellow microneedles. Mp 280-290° dec.

5-Methoxy: 5-Methoxy-3H-indolo[3,2,1-de][1,5]naphthyridine-2,6-dione. 5-Methoxycanthin-2,6-dione. Indicanthione

[35817-57-3]

C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the wood of *Samadera indica* (Simaroubaceae). Stout red needles (CHCl₃/MeOH). Mp 258° dec. Displays intense yellow-green fluor. in organic solvents.

Iyer, V.S. *et al.*, *Curr. Sci.*, 1972, **41**, 140-141 (Indacanthinone)

Giesbrecht, A.M. *et al.*, *Phytochemistry*, 1980, **19**, 313-315 (3-Methoxycanthin-2,6-dione)


Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1204-1209 (3-Methylcanthin-2,6-dione)

Arisawa, M. *et al.*, *J. Nat. Prod.*, 1983, **46**, 222-225 (Canthin-2,6-dione, 10-Hydroxy-3-methoxy-2,6-canthindione)

Barbetti, P. *et al.*, *Planta Med.*, 1990, **56**, 216-217 (5-Hydroxy-4-methoxy-3-methyl-2,6-canthindione)

3H-Indol-3-one, 9CI, 8CI**I-106**

3-Oxoindole. Dehydroindoxyl. 3-Pseudoindolone
[67285-12-5]

C₈H₅NO 131.134

Pigment from the edible fungus *Pleurotus salmoneostramineus*. Dark orange-red

prisms. Mp 110-112° dec. λ_{\max} 208 (€ 1318); 220 (€ 1288); 282 (€ 501); 290 (€ 489); 456 (€ 692); 480 (MeOH) (Berdy).

Oxime: 3-Hydroxyiminoindole. 3-Nitrosoindole

[76983-82-9]

C₈H₆N₂O 146.148

Brown needles (C₆H₆). Mp 180-182° dec. Tautomeric.

l-Oxide: Isatogen

[5814-98-2]

C₈H₅NO₂ 147.133*l*-Oxide, oxime: [69111-91-7]C₈H₆N₂O₂ 162.148

Yellow-green needles (H₂O). Mp 167-169°. Subl. ca. 150.

Neunhoeffer, O. *et al.*, *Chem. Ber.*, 1961, **94**, 2965-2967

Stamm, H. *et al.*, *Method. Chim.*, 1975, **6**, 329 (rev)

Hiremath, S.P. *et al.*, *Adv. Heterocycl. Chem.*, 1978, **22**, 123-181 (rev)

Acheson, R.M. *et al.*, *J.C.S. Perkin 1*, 1978, 1117-1125 (*l*-oxide, oxime)


Hiremath, S.P. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 767-769 (oxime)

Takehuma, S. *et al.*, *J.A.C.S.*, 1994, **116**, 8849-8850 (isol)

Portela-Cubillo, F. *et al.*, *Chem. Comm.*, 2007, 4041-4043 (synth)

7H-Indolo[3,2-*j*]phenanthridine-7,13(12H)-dione, 9CI**I-107****Calothrixin B**

[254114-34-6]

C₁₉H₁₀N₂O₂ 298.3

Alkaloid from the cyanobacterium *Calothrix* sp. obtained from soil. Exhibits antimalarial and cytotoxic activities. Amorph. orange-red solid. λ_{\max} 283 (€ 15000); 352 (€ 2900); 405 (€ 1710) (EtOH).

*N*⁵-Oxide: **Calothrixin A**

[254114-33-5]

C₁₉H₁₀N₂O₃ 314.3

Alkaloid from *Calothrix* sp. obtained from soil. Exhibits antimalarial and cytotoxic activities. Long wine-red needles (DMSO). Dec. at 280°. λ_{\max} 292 (€ 19000); 362 (€ 4260); 413 (€ 3100) (EtOH).

Rickards, R.W. *et al.*, *Tetrahedron*, 1999, **55**, 13513-13520 (isol)

Bernardo, P.H. *et al.*, *J.O.C.*, 2003, **68**, 8906-8909 (synth)

Tohyama, S. *et al.*, *Tet. Lett.*, 2005, **46**, 5263-5264; 2006, **47**, 5859-5861 (synth)


Sissouma, D. *et al.*, *J.O.C.*, 2006, **71**, 8384-8389 (synth)

Bennasar, M.-L. *et al.*, *Org. Lett.*, 2006, **8**, 561-564 (synth)

McErlean, C.S.P. *et al.*, *Tetrahedron*, 2007, **63**, 10963-10970 (synth)

Indolo[2,1-*b*]quinazoline-6,12-dione, 9CI**I-108**

6,12-Dihydro-6,12-dioxoindolo[2,1-*b*]quinazoline. **Tryptanthrine**. **Courouputine A** [13220-57-0]

C₁₅H₈N₂O₂ 248.24

Isol. from the yeast *Candida lipolytica* grown in the presence of L-Tryptophan. Also from *Courouputa guianensis* (Lecythidaceae), *Isatis indigotica* and *Polygonum tiuiforum*. Prod. by *Cytophaga marinoflava* sp. AM13.1. Antifungal and antimicrobial agent showing specific activity against dermatophytes. Active against *Helicobacter pylori*. Yellow needles. Sol. MeOH, CHCl₃; poorly sol. H₂O, bases, acids. Mp 266-267°. Subl. *in vacuo*. Courouputine A was initially assigned an incorr. struct. λ_{\max} 248 ; 252 ; 277 ; 311 ; 328 ; 387 (MeOH) (Berdy). λ_{\max} 254 (€ 34800); 280 (€ 7000); 340 (€ 7200); 405 (€ 6600) (EtOH) (Berdy).

Brufani, M. *et al.*, *Experientia*, 1971, **27**, 1249 (cryst struct, pmr, ms)

Fedeli, W. *et al.*, *J.C.S. Perkin 2*, 1974, 1621 (cryst struct)

Bergman, J. *et al.*, *Tet. Lett.*, 1977, 2625 (struct, synth)

Karpf, H. *et al.*, *Tet. Lett.*, 1978, 3007 (synth, ir, uv, ms)

Honda, G. *et al.*, *Planta Med.*, 1979, **37**, 172 (props)

Mitscher, L.A. *et al.*, *Heterocycles*, 1981, **15**, 1017 (synth)

Capuano, L. *et al.*, *Chem. Ber.*, 1983, **116**, 741 (synth)

Eguchi, S. *et al.*, *Heterocycles*, 1992, **33**, 153 (synth)

Cornforth, J. *et al.*, *J.C.S. Perkin 1*, 1996, 2787 (synth)

Hashimoto, T. *et al.*, *Nat. Med. (Tokyo)*, 1999, **53**, 27 (isol, activity)


Shaaban, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (Cytophaga, isol, pmr, cmr)

Batanero, B. *et al.*, *Tet. Lett.*, 2006, **47**, 8201-8203 (synth)

Bowman, W.R. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 103-113 (synth)

Indolo[3,2-*b*]quinoline**I-109****Quindoline**

[243-58-3]

**10H-form**C₁₅H₁₀N₂ 218.257

10H-Form appears to predominate for parent compd. Alkaloid from the roots of *Cryptolepis sanguinolenta* (Asclepiadaceae). Active against gram-positive bacteria. Anticomplementary agent. Needles

(EtOH) (isol. from the plant as amorph. solid). Mp 187° Mp 247-248°. Blue fluor. in soln. Subl. with part. dec. λ_{\max} 228 ; 276 ; 344 (MeOH) (Berdy). λ_{\max} 226 ; 272 ; 286 ; 368 (MeOH/HCl) (Berdy).

Hydrochloride:

Yellow cryst. H₂O, EtOH.

5-Oxide: [13220-52-5]

C₁₅H₁₀N₂O 234.257

Yellow prisms (EtOH). Mp 294-296°.

5H-form

5-Me: 5-Methyl-5H-quinodoline, 9CI.

Cryptolepine

[480-26-2]

C₁₆H₁₂N₂ 232.284

Alkaloid from *Cryptolepis sanguinolenta* (Asclepiadaceae). Antibacterial, antihyperthermic, antimalarial and antihyperglycaemic agent. Platelet aggregation inhibitor. Active against gram-positive bacteria. Deep purple needles. Mp 175-178°. Forms a monohydrate, Mp 166-169°.

5-Me: hydrochloride: [72782-09-3]

Mp 263-265°.

[42375-21-3]

Fichter, F. *et al.*, *Ber.*, 1906, **39**, 3932-3942;

1910, **43**, 3489-3499 (*synth*)

Armit, J.N. *et al.*, *J.C.S.*, 1922, **121**, 827-839

(*synth*)

Holt, S.J. *et al.*, *J.C.S.*, 1947, 607-611 (*synth*)

Gellert, E. *et al.*, *Helv. Chim. Acta*, 1951, **34**,

642-651 (*isol, uv, struct, Cryptolepine*)

Dwuma-Badu, D. *et al.*, *J. Pharm. Sci.*, 1978,

67, 433-434 (*isol, uv, ir, pmr, ms, Cryptolepine*)

Oyekan, A.O. *et al.*, *Gen. Pharmacol.*, 1988,

19, 233-237; 1993, **24**, 461-469; 1285-1290

(*Cryptolepine, pharmacol*)

Tackie, A.N. *et al.*, *J. Het. Chem.*, 1991, **28**,

1429-1435; 1996, **33**, 789-797 (*pmr, cmr, Cryptolepine*)

Spitzer, T.D. *et al.*, *J. Het. Chem.*, 1991, **28**,

2065-2070 (*pmr, cmr*)

Oyekan, A.O. *et al.*, *J. Cardiovasc. Pharmacol.*,

1994, **23**, 601-611 (*pharmacol*)

Paulo, A. *et al.*, *J. Ethnopharmacol.*, 1994, **44**,

127-130 (*activity*)

Sawer, I.K. *et al.*, *J. Appl. Bacteriol.*, 1995, **79**,

314-321 (*Cryptolepine, activity*)

Cimanga, K. *et al.*, *Planta Med.*, 1996, **62**, 22-

27 (*isol, pmr, cmr, ir, uv, activity, Cryptolepine*)

Cooper, M.M. *et al.*, *Tet. Lett.*, 1996, **37**, 4283-

4286 (*synth*)

Cimanga, K. *et al.*, *J. Nat. Prod.*, 1997, **60**,

688-691 (*Cryptolepine, activity*)

Bierer, D.E. *et al.*, *J. Med. Chem.*, 1998, **41**,

894-901; 2754-2764 (*Cryptolepine, isol, synth*)

Hadden, C.E. *et al.*, *J. Het. Chem.*, 1999, **36**,

525-531 (*pmr, cmr*)

Yang, S.-W. *et al.*, *J. Nat. Prod.*, 1999, **62**, 976-

983 (*isol, synth, pmr, cmr, Cryptolepine*)

Csanyi, D. *et al.*, *Synth. Commun.*, 1999, **29**,

3959-3969 (*synth*)

Radl, S. *et al.*, *J. Het. Chem.*, 2000, **37**, 855-862

(*synth, uv, pmr*)

Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2002, **85**,

3823-3827 (*synth, ir, pmr, cmr*)

Dhanabal, T. *et al.*, *Tetrahedron*, 2006, **62**,

6258-6263 (*Cryptolepine, synth*)


Tousek, J. *et al.*, *Magn. Reson. Chem.*, 2008,

46, 42-51 (*pmr, cmr, N-15 nmr*)

Indolo[3,2-c]quinoline

3,4-Benzo- γ -carboline

[239-09-8]



11H-form

C₁₅H₁₀N₂ 218.257

Parent compd. should exist as 11H-form.

Prisms (EtOH aq.). Mp 320° (310°) subl.

Picrate:

Orange-yellow needles (EtOH). Mp

268°.

5H-form

N-Me: 5-Methyl-5H-indolo[3,2-c]quino-

line. *Cryptosanguinolentine. Isocryptole-*

lepine

[165467-65-2]

C₁₆H₁₂N₂ 232.284

Alkaloid from the roots of *Cryptolepis*

sanguinolenta (Asclepiadaceae). Cryst.

Mp 132-133°.

[76401-99-5]

Russel, O. *et al.*, *J.C.S.*, 1965, 5458 (*synth*)

Dave, V. *et al.*, *Tetrahedron*, 1975, **31**, 1255

(*synth*)

Stadlbauer, W. *et al.*, *Monatsh. Chem.*, 1987,

118, 81 (*synth*)

Pousset, J.-L. *et al.*, *Phytochemistry*, 1995, **39**,

735 (*isol, uv, ir, pmr, cmr, ms, struct, deriv*)

Dubovitskii, S.V. *et al.*, *Izv. Akad. Nauk, Ser.*

Khim., 1996, 2797; *CA*, **126**, 157678v (*synth*)

Sharaf, M.H.M. *et al.*, *J. Het. Chem.*, 1996, **33**,

239 (*isol, pmr, cmr, struct, deriv*)

Molina, A. *et al.*, *J.O.C.*, 1996, **61**, 5587 (*synth*)

Timari, G. *et al.*, *Synlett*, 1997, 1067-1068

(*synth*)

Murray, P.E. *et al.*, *J. Chem. Res., Synop.*,

1998, 376-377 (*synth*)

Fresneda, P.M. *et al.*, *Tetrahedron*, 2001, **57**,

6197-6202 (*Isocryptolepine, synth*)

Kumar, R.N. *et al.*, *Tet. Lett.*, 2002, **43**, 3327-

3328 (*Isocryptolepine, synth*)

Dhanabal, T. *et al.*, *Tet. Lett.*, 2005, **46**, 4509-

4510 (*Cryptosanguinolentine, synth*)

Dhanabal, T. *et al.*, *Tetrahedron*, 2006, **62**,

6258-6263 (*Isocryptolepine, synth*)

Miki, Y. *et al.*, *Tet. Lett.*, 2007, **48**, 9093-9095

(*Isocryptolepine, synth*)

Tousek, J. *et al.*, *Magn. Reson. Chem.*, 2008,

46, 42-51 (*pmr, cmr, N-15 nmr*)

10H-Indolo[3,2-b]quinoline-

11-carboxylic acid

Quindoline-11-carboxylic acid. Flavindin

C₁₆H₁₀N₂O₂ 262.267

Yellow ppt. Mp 336-337° dec.

Me ester:

C₁₇H₁₂N₂O₂ 276.294

Yellow needles (MeOH aq.). Mp 154-

155°.

Amide:

C₁₆H₁₁N₃O 261.282

Mp 334-336°.

5H-form

N-Me: 5-Methyl-5H-indolo[3,2-b]quino-

line-11-carboxylic acid. *Cryptolepinoic*

acid

C₁₇H₁₂N₂O₂ 276.294

I-110

Alkaloid from *Cryptolepis sanguinolenta*. Amorph. orange powder. λ_{\max} 223 (log ϵ 3.55); 244 (log ϵ 3.3); 280 (log ϵ 3.69); 352 (sh) (log ϵ 3.24); 366 (log ϵ 3.52) (MeOH).

N-Me. Me ester: *Methyl cryptolepinoate*

C₁₈H₁₄N₂O₂ 290.321

Alkaloid from *Cryptolepis sanguinolenta*. Dark reddish powder.

Fichter, R. *et al.*, *Ber.*, 1910, **43**, 3489

Holt, S.J. *et al.*, *J.C.S.*, 1947, 607

Paulo, A. *et al.*, *Planta Med.*, 2000, **66**, 30-34


(*Cryptolepinoic acid*)

10H-Indolo[3,2-b]quinolin-9-

ol

9-Hydroxy-10H-quinodoline

I-112



C₁₅H₁₀N₂O 234.257

O- β -D-Glucopyranoside: *Jusbetonin*

C₂₁H₂₀N₂O₆ 396.399

Alkaloid from the leaves of *Justicia*

betonica. Amorph. yellow powder. [α]_D²⁵

-48.2 (c, 0.54 in MeOH). Mp >300°.

Subbaraju, G.V. *et al.*, *J. Nat. Prod.*, 2004, **67**,


461-462 (*isol, pmr, cmr*)

Indolo[2,3-a]quinolizine, 9CI

Indolopyridocoline

[239-15-6]

I-113



C₁₅H₁₀N₂ 218.257

Truncated indole alkaloid, original ske-

leton uncertain. Prob. not scologanin-

derived. Alkaloid from *Gonioma kamassi*

(Apocynaceae). Orange-yellow, noncryst.

solid.

Perchlorate:

Yellow-green cryst. Mp 282-285° dec.

Picrate: Mp 252-255°.

1,2,3,4,6,7-Hexahydro: 1,2,3,4,6,7-Hexa-

hydroindolo[2,3-a]quinolizine. Tetra-

methylenedihydro- β -carboline

C₁₅H₁₆N₂ 224.305

Alkaloid from *Nitraria schoberi*. Mp

81-83°.

1,2,3,4,6,7-Hexahydro, hydrochloride: Mp

252°.

Kaschnitz, R. *et al.*, *Monatsh. Chem.*, 1965,

96, 909 (*isol, uv, ms, struct*)

Gribble, G.W. *et al.*, *Tet. Lett.*, 1987, **28**, 5259

(*synth*)

Fürstner, A. *et al.*, *Tetrahedron*, 1995, **51**, 773

(*synth*)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996,


32, 937-1035; *Chem. Nat. Compd. (Engl.*

Transl.), 1996, **32**, 932-1028 (*hexahydro*)

Kraus, G.A. *et al.*, *Synlett*, 1997, 107 (*synth*)


Indolo[3,2,1-ij]quino[4,5-bc][1,5]naphthyridine, 9CI
Komarovidinine
[85412-78-8]

I-114


Porter, J.K. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 88-93 (*isol, uv, ir, pmr, ms*)C₂₀H₁₁N₃ 293.327Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (MeOH/CHCl₃). Mp 254-255°.N³-Oxide: **Komarovidinine N³-oxide**
[342880-44-8]C₂₀H₁₁N₃O 309.326Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (EtOH/Me₂CO). Mp 263-264°.Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 635-638; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 601-603 (*isol, struct*)Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 396-398 (*isol, Komarovidinine N-oxide*)**1H-Indol-3-ylacetyl-myoinositol**

I-115


[73925-84-5]

C₁₆H₁₉NO₇ 337.329The inositol residue has the *myo*-config. but the point of attachment is variable (i.e. exact config. is unknown). Present in *Oryza sativa* (rice) and *Zea mays* (corn). Hall, P.J. *et al.*, *Phytochemistry*, 1980, **19**, 2121**4-(1H-Indol-3-yl)-1,2,3-butanetriol, 9CI**

I-116

3-(2,3,4-Trihydroxybutyl)indole
[61183-25-3]C₁₂H₁₅NO₃ 221.255Alkaloid from the clavicipitaceous fungus *Balansia epichloë* which parasitizes pasture grasses. λ_{max} 221 (log ε 4.65); 272 (log ε 3.78); 279 (log ε 3.8); 288 (log ε 3.73) (MeOH).**4-(1H-Indol-3-yl)-3-buten-2-one, 9CI**

I-117

3-(3-Oxo-1-butenyl)-1H-indole
[31951-75-4]C₁₂H₁₁NO 185.225**(E)-form** [57598-80-8]Alkaloid from the sponge *Tedania ignis*. Cryst. (petrol or C₆H₆). Mp 145-146°. Possible artifact. λ_{max} 223 (ε 19000); 276 (ε 9400); 353 (ε 17800) (no solvent reported).**6-Bromo-4-(6-Bromo-1H-indol-3-yl)-3-buten-2-one**


[654649-67-9]

C₁₂H₁₀BrNO 264.121Alkaloid from the sponge *Iotrochota birotulata*. Powder. Mp 145-148°. λ_{max} 270 ; 290 ; 340 (MeOH).Nonnenmacher, A. *et al.*, *Annalen*, 1983, 2135-2140 (*synth, pmr*)Dillman, R.L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1056-1061 (*isol*)Li, L. *et al.*, *Pharmazie*, 2003, **58**, 680-681 (6-bromo)**4-(1H-Indol-6-yl)-3-buten-2-one, 9CI**


I-118


6-(3-Oxo-1-butenyl)-1H-indole
[56774-96-0]C₁₂H₁₁NO 185.225**(E)-form** [40863-48-7]Alkaloid from *Monodora angolensis*. Pale yellow cryst. (cyclohexane/C₆H₆). Mp 135-137°.Ishii, H. *et al.*, *Tetrahedron*, 1975, **31**, 933-938 (*synth, pmr*)Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (*isol, pmr, cmr, ms*)**1-(3-Indolyl)-β-carboline**

I-119

1-(1H-Indol-3-yl)-9H-pyrido[3,4-b]indole, 9CI. Eudistomin U
[155885-64-6]C₁₉H₁₃N₃ 283.332Alkaloid from the Caribbean ascidian *Lissoclinum fragile*. Exhibits antibacterialactivity. DNA binder. Yellow foam. Sol. MeOH. λ_{max} 220 (ε 8000); 240 (ε 3500); 252 (ε 10500); 270 (ε 7700); 340 (ε 9020); 380 (ε 13000); 470 (ε 5120) (MeOH) (Berdy).**3,4-Dihydro-3,4-Dihydroeudistomin U. Isoeudistomin U. Eudisin B**
[155885-65-7]C₁₉H₁₅N₃ 285.348From *Lissoclinum fragile* and a *Eudistoma* sp. Exhibits antibacterial activity. Yellow foam. Originally assigned an α-carboline struct. by Badre *et al* (1994). The name Isoeudistomin U is now misleading. λ_{max} 208 (ε 27500); 248 (ε 11200); 274 (ε 6100); 282 (ε 6100); 330 (ε 9560); 397 (ε 9530) (MeOH). λ_{max} 210 (ε 60000); 243 (sh); 314 (ε 9340) (MeOH).**3,4-Dihydro, 6'-bromo: 19-Bromo-3,4-dihydroeudistomin U. 19-Bromoisoeudistomin U (incorr.). Eudisin A**
[186820-55-3]C₁₉H₁₄BrN₃ 364.244Alkaloid from a *Eudistoma* sp. tunicate. Antitumour agent. Yellow-green powder. λ_{max} 220 (ε 56600); 290 (ε 17500); 325 (ε 20700); 396 (ε 10400) (MeOH). λ_{max} 206 (ε 70000); 245 (sh); 292 (ε 7040); 315 (ε 9020) (MeOH). λ_{max} 206 (ε 70000); 292 (ε 7040); 315 (ε 9020) (MeOH-NaOH) (Berdy).Badre, A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 528-533 (*isol, uv, ir, pmr, cmr, ms, struct*)Massiot, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1636-1639 (*Isoeudistomin U*)Molina, P. *et al.*, *Tet. Lett.*, 1995, **36**, 3581-3582 (*synth*)Rocca, P. *et al.*, *Tet. Lett.*, 1995, **36**, 7085-7088 (*synth*)Kang, H. *et al.*, *Nat. Prod. Lett.*, 1996, **9**, 7-12 (*Isoeudistomin U, 19-Bromoisoeudistomin U*)**N^α-(1H-Indol-3-ylcarbonyl)arginine**

I-120

*(R)-form*C₁₅H₁₉N₅O₃ 317.347**(R)-form** [178495-40-4]Isol. from the ascidian *Leptoclinides dubius*. Sol. MeOH, butanol. [α]_D²⁰ -133.3 (c, 0.1 in MeOH). λ_{max} 214 (log ε 3.65); 246 (log ε 3.13); 282 (log ε 3.11); 286 (log ε 3.1) (MeOH).**(S)-form****6'-Bromo: N^α-(6-Bromo-1H-indol-3-ylcarbonyl)arginine**
[178495-41-5]C₁₅H₁₈BrN₅O₃ 396.243Isol. from *Leptoclinides dubius*. Sol. MeOH, butanol. [α]_D²⁰ +3.6 (c, 0.5 in MeOH). λ_{max} 222 (log ε 3.78); 250 (log ε 3.43); 280 (log ε 3.18) (MeOH).García, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 782-785 (*isol*)

N^z-(1H-Indol-3-ylcarbonyl)-histidine I-121C₁₅H₁₄N₄O₃ 298.301**(S)-form**6'-Bromo: N^z-(6-Bromo-1H-indol-3-yl-carbonyl)histidine


[178495-42-6]

C₁₅H₁₃BrN₄O₃ 377.197

Isol. from the ascidian *Leptoclinides dubius*. Sol. MeOH, butanol. [α]_D²⁰ +26.6 (c, 0.5 in MeOH). λ_{max} 222 (log ε 3.53); 250 (log ε 3.17); 282 (log ε 2.92) (MeOH).

Garcia, A. et al., *J. Nat. Prod.*, 1996, **59**, 782-785 (isol)

3-Indolylmethyl glucosinolate I-122

Glucobrassicin
[4356-52-9]C₁₆H₂₀N₂O₉S₂ 448.474

Constit. of *Brassica* and *Raphanus* spp., e.g. rape (*Brassica napus* var. *napus*) and Brussels sprouts (*Brassica oleracea* var. *gemmifera*). Minor constit. of *Moricandia arvensis* (Brassicaceae). Mp 149-150° (as Me₄N salt). [α]_D²² -13.3 (Me₄N salt).

N-Ac: (1-Acetyl-1H-indol-3-yl)methyl glucosinolate. N-Acetylglucobrassicin [103951-50-4]

C₁₈H₂₂N₂O₁₀S₂ 490.511Present in *Tovaria pendulata* seedlings.

N-Sulfonic acid: (1-Sulfo-1H-indol-3-yl)-methyl glucosinolate, 9CI. Sulfoglucobrassicin [29702-28-1]

C₁₆H₂₀N₂O₁₂S₃ 528.538Isol. from woad (*Isatis tinctoria*).

Cryst. (EtOH aq.) (as Me₄N salt). Mp 156° dec. (Me₄N salt).

N-Methoxy: Neoglucobrassicin. N-Methoxyglucobrassicin [5187-84-8]

C₁₇H₂₂N₂O₁₀S₂ 478.5

Widespread in *Brassica* spp. and found in the Brassicaceae, Tovariaceae, Caparidaceae and Resedaceae. Mp 175° dec. (as brucine salt).

4-Hydroxy: 4-Hydroxyindol-3-ylmethyl glucosinolate. 4-Hydroxyglucobrassicin [83327-20-2]

[87592-99-2, 96828-27-2, 96828-26-1, 87593-00-8]

C₁₆H₂₀N₂O₁₀S₂ 464.473

Isol. from *Brassica* sp. and other crucifers. Formulated as the 5-hydroxy isomer by Goetz et al. Readily oxidises in the presence of air and light. λ_{max} 267; 284; 293 (MeOH).

4-Methoxy: 4-Methoxyindol-3-ylmethyl glucosinolate. 4-Methoxyglucobrassicin [83327-21-3]

C₁₇H₂₂N₂O₁₀S₂ 478.5

Isol. from *Brassica* sp. and other crucifers. Gummy solid. λ_{max} 267; 282; 292 (MeOH).

N,4-Dimethoxy: 1,4-Dimethoxyglucobrassicin [334832-34-7]

C₁₈H₂₄N₂O₁₁S₂ 508.526

Isol. from the roots of *Barbarea vulgaris* ssp. *arcuata*.

Gmelin, R. et al., *Suom. Kemistil. B*, 1961, **34**, 15-18 (isol)

Gmelin, R. et al., *Acta Chem. Scand.*, 1962, **16**, 1378-1384 (*Neoglucobrassicin*)

Gmelin, R. et al., *CA*, 1965, **63**, 5584e (isol, struct)

Schraudolf, H. et al., *Experientia*, 1965, **21**, 520-522 (occur)

Elliott, M.C. et al., *Phytochemistry*, 1970, **9**, 1629-1632 (*N-sulfonic acid*)

Fenwick, G.R. et al., *Biomed. Mass Spectrom.*, 1980, **7**, 410-412 (ms)

Truscott, R.J.W. et al., *Biochem. Biophys. Res. Commun.*, 1982, **107**, 1258-1264; 1368-1375 (*4-methoxy, 4-hydroxy, isol, struct*)

Truscott, R.J.W. et al., *J. Sci. Food Agric.*, 1983, **34**, 247-254 (isol)

Hanley, A.B. et al., *J. Sci. Food Agric.*, 1983, **34**, 867-869 (isol)

Goetz, J.K. et al., *Phytochemistry*, 1983, **22**, 905-907 (*N-methoxy, 4-hydroxy, isol*)

Cox, I.J. et al., *Carbohydr. Res.*, 1984, **132**, 323-329 (cmr, pmr)

Hanley, A.P. et al., *Phytochemistry*, 1985, **24**, 598-600 (isol, struct)

Schraudolf, H. et al., *Z. Naturforsch., C*, 1986, **41**, 526-528 (acetyl)

Belkhir, A. et al., *Phytochemistry*, 1990, **29**, 1315-1316 (isol, uv, pmr)


Viaud, M.C. et al., *Tet. Lett.*, 1990, **31**, 1417-1418 (synth)

Agerbirk, N. et al., *J. Agric. Food Chem.*, 2001, **49**, 1502-1507 (*1,4-Dimethoxyglucobrassicin*)

Fahey, J.W. et al., *Phytochemistry*, 2001, **56**, 5-51 (rev, occur)

1-(1H-Indol-7-yl)-3-methyl-1,2,3-butanetriol, 9CI I-123

7-(1,2,3-Trihydroxy-3-methylbutyl)-1H-indole



(1'R*,2'R*)-form

C₁₃H₁₇NO₃ 235.282**(1'R*,2'R*)-form**threo-form. *Leiocarpatriol A*

[132200-50-1]

Isol. from the roots of *Esenbeckia leio-carpa*. Amorph. Sensitive to light and air.


(1'R*,2'S*)-formerythro-form. *Leiocarpatriol B*

[132113-91-8]

Isol. from the roots of *Esenbeckia leio-carpa*. Amorph. Sensitive to light and air.

Delle Monache, F. et al., *Gazz. Chim. Ital.*, 1990, **120**, 387 (isol, pmr, cmr)

4-(3-Indolyl)-2-methyl-1-butanol I-124

β-Methyl-1H-indole-3-butanol, 9CI. *Paniculidine C. Paniculol*C₁₃H₁₇NO 203.283**(R)-form** [97399-95-6]

Alkaloid from the root bark of *Murraya paniculata* (Rutaceae). Oil. [α]_D²⁰ +45 (c, 0.035 in CHCl₃).

N-Methoxy: 1-Methoxy-β-methyl-1H-indole-3-butanol, 9CI. *Paniculidine B* [97399-94-5]

C₁₄H₁₉NO₂ 233.31

Alkaloid from root bark of *Murraya paniculata* (Rutaceae). Oil. [α]_D²⁰ +21 (c, 0.025 in CHCl₃).

Carboxylic acid, Me ester: β-Methyl-1H-indole-3-butanoic acid, 9CI. 4-(3-Indolyl)-2-methylbutanoic acid. *Paniculidine A*

[97399-93-4]

C₁₄H₁₇NO₂ 231.294

Alkaloid from root bark of *Murraya paniculata* (Rutaceae). Oil. [α]_D²⁴ -31.9 (c, 0.1 in CHCl₃).

[120786-74-5, 129172-71-0]

Kinoshita, T. et al., *Chem. Pharm. Bull.*, 1985, **33**, 1770 (uv, ir, pmr, cmr, ms, struct, synth)

Somei, M. et al., *Chem. Pharm. Bull.*, 1985, **33**, 5147 (synth)

Cheskis, B.A. et al., *J.C.S. Perkin 1*, 1989, 1353

Kinoshita, T. et al., *Phytochemistry*, 1989, **28**, 147 (isol, uv, ir, pmr, cmr, ms, struct)

Boumendjel, A. et al., *Bull. Soc. Chim. Fr.*, 1990, 645 (synth, *Paniculidine A*)


Ito, C. et al., *J.C.S. Perkin 1*, 1990, 2047 (isol, pmr)


Cheskis, B.A. et al., *Zh. Org. Khim.*, 1990, **26**, 425 (synth)

Selvakumar, N. et al., *J.O.C.*, 2004, **69**, 4429-4432 (synth)



1-(1H-Indol-3-yl)-3-methyl-2-buten-1-one, 9CI I-125

3-(3-Methyl-1-oxo-2-butenyl)-1H-indole [50615-00-4]



C₁₃H₁₃NO 199.252Alkaloid from root bark of *Pamburus missionis* (Rutaceae). Needles (CH₂Cl₂/hexane). Mp 136.5-138° (130-131°).Somei, M. *et al.*, *Tet. Lett.*, 1973, 2451 (*synth*)
Wenkert, E. *et al.*, *J.O.C.*, 1986, 51, 2343 (*synth*)Kumar, V. *et al.*, *Phytochemistry*, 1994, 36, 879 (*isol, uv, ir, pmr, cmr, ms*)**6-(1*H*-Indol-3-yl)-5-methyl-3,5-heptadien-2-one, 9CI** I-126C₁₆H₁₇NO 239.316**(3*E*,5*E*)-form** [136685-36-4]Isol. from the sponge *Tedania ignis*. Phytotoxic agent. λ_{max} 224 (ε 12300); 290 (ε 8300) (EtOH) (Berdy).Dillman, R.L. *et al.*, *J. Nat. Prod.*, 1991, 54, 1056 (*isol, uv, ir, pmr, ms, cmr, struct*)**2-(1*H*-Indol-3-ylmethyl)-1*H*-indole-3-acetic acid, 9CI** I-127


[68232-56-4]

C₁₉H₁₆N₂O₂ 304.348Prod. by *Sulfolobus acidocaldarius*. Pink cryst. (CHCl₃/hexane). Undefined Mp. λ_{max} 224 (ε 31000); 282 (ε 15200); 290 (ε 13200) (MeOH). λ_{max} 225; 283; 291 (EtOH) (Berdy).Suzuki, Y. *et al.*, *Agric. Biol. Chem.*, 1978, 42, 1315-1321 (*synth*)Amat-Guerri, F. *et al.*, *J. Chem. Res., Synop.*, 1984, 160-161 (*synth*)White, R.H. *et al.*, *J. Bacteriol.*, 1987, 169, 5859-5860 (*isol*)**N^b-(3-Indolylmethyl)-5-methoxy-N^b-methyltryptamine** I-128N-(1*H*-Indol-3-ylmethyl)-5-methoxy-N-methyl-1*H*-indole-3-ethanamine [173180-36-4]C₂₁H₂₃N₃O 333.432Alkaloid from the roots of *Antirhea lucida* (Rubiaceae). Amorph. powder. λ_{max} 221 (log ε 4.1); 282 (log ε 3.5)




(EtOH).

Weniger, B.R. *et al.*, *Planta Med.*, 1995, 61, 569-570 (*isol, pmr, cmr, uv, struct*)**4-(3-Indolylmethyl)-7-(2-methylpropyl)-1,2,3,5,6-pentathiepane** I-129


3-[(7-Isobutyl-1,2,3,5,6-pentathiepan-4-yl)methyl]indole. 4-(3-Indolemethyl)-7-isobutyl-1,2,3,5,6-pentathiepane [151261-55-1]

C₁₅H₁₉NS₅ 373.652Prod. by the sulfur-metabolising hyperthermophilic archaeobacteria *Thermococcus acidaminovorans* and *Thermococcus tadjuricus*. Solid. Mp 123°.Ritzau, M. *et al.*, *Annalen*, 1993, 871**3-(3-Indolylmethyl)-5-(2-methylpropyl)-1,2,4-trithiolane** I-130

3-[(5-Isobutyl-1,2,4-trithiolan-3-yl)-methyl]indole. 3-(3-Indolemethyl)-5-isobutyl-1,2,4-trithiolane [151261-46-0]

C₁₅H₁₉NS₃ 309.52Prod. by the sulfur-metabolising archaeobacteria *Thermococcus acidaminovorans* and *Thermococcus tadjuricus*. Solid. Mp 98°.Ritzau, M. *et al.*, *Annalen*, 1993, 871**3-(1*H*-Indol-3-yl)-2-methylpropanoic acid** I-131α-Methyl-1*H*-indole-3-propanoic acid, 9CI. *Rhizophorine* [15142-91-3] [69088-73-9]C₁₂H₁₃NO₂ 203.24**(ξ)-form** [61755-28-0]Constit. of *Rhizophora mucronata*. Plant growth inhibitor. Off-white needles (H₂O). Mp 129-130°.*Et ester*: [57901-11-8]C₁₄H₁₇NO₂ 231.294Bp_{0.75} 165°.Johnson, H.E. *et al.*, *J.O.C.*, 1963, 28, 2030-2032 (*synth*)Saha, P.K. *et al.*, *Plant Biochem. J.*, 1978, 5, 65-68 (*isol*)**N-[2-(1*H*-Indol-3-yl)-2-oxoethyl]acetamide, 9CI** I-132N-Acetyl-β-oxotryptamine. 3-(Acetamidoacetyl)indole. 34*N* [73053-91-5]C₁₂H₁₂N₂O₂ 216.239Prod. by *Streptomyces ramulosus*. Shows no antimicrobial activity. Cryst. (EtOH). Mp 216-218° (203-204°). λ_{max} 241 (ε 11000); 261 (ε 8200); 300 (ε 11000) (MeOH) (Derep).*l'*-Hydroxy: N-[1-Hydroxy-2-(1*H*-indol-3-yl)-2-oxoethyl]acetamide. 34*M*

[87084-38-6]

C₁₂H₁₂N₂O₃ 232.238Prod. by *Streptomyces ramulosus*. Shows no antibacterial activity. Needles (MeOH). Mp 146-147°. [α]_D²⁰ -92 (c, 1.0 in MeOH). λ_{max} 241 (ε 11000); 261 (ε 8200); 300 (ε 11000) (MeOH) (Derep).Oikawa, Y. *et al.*, *Heterocycles*, 1979, 12, 1457-1462 (*synth*)Yoshioka, T. *et al.*, *J. Chem. Res., Synop.*, 1981, 194-195; *J. Chem. Res., Miniprint*, 1981, 2252-2281 (*synth*)Chen, Y. *et al.*, *J. Antibiot.*, 1983, 36, 913-915 (*isol, uv, ir, pmr, cmr, ms, struct*)**3-(1*H*-Indol-3-yl)-2-oxopropanoic acid** I-133α-Oxo-1*H*-indole-3-propanoic acid, 9CI. 2-Hydroxy-3-(1*H*-indol-3-yl)-2-propenoic acid. 3-(3-Indolyl)pyruvic acid [392-12-1] [144334-60-1]C₁₁H₉NO₃ 203.197

Tautomer ratio solvent dependent in soln. Crystallises as the (Z)-enol tautomer. Prod. from tryptophan by microorganisms; poss. present in higher plants. Grey cryst. + 1AcOH (AcOH). Mp 211°.

▶ NM1880000

Oxime:C₁₁H₁₀N₂O₃ 218.212


Mp 175°.

4-Nitrophenylhydrazone: Mp 153-154°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 669A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 140B (nmr)
 Gränacher, C. et al., *Helv. Chim. Acta*, 1924, **7**, 575 (synth)
 Kaper, J.M. et al., *Arch. Biochem. Biophys.*, 1963, **103**, 475
 Srivastava, B.I.S. et al., *Plant Physiol.*, 1964, **39**, 381; *CA*, **61**, 16442
 Matchett, W.H. et al., *Biochim. Biophys. Acta*, 1965, **107**, 222
 Havaaldsen, R. et al., *Acta Chem. Scand.*, 1967, **21**, 1095 (synth)
 Nazario, J. et al., *Arch. Biochem. Biophys.*, 1968, **123**, 457-461 (tautom, ir)
 Okabe, N. et al., *Acta Cryst. C*, 1998, **54**, 1330-1331 (cryst struct)

5-(1*H*-Indol-3-yl)-3-penten-2-one, 9CI I-134




C₁₃H₁₃NO 199.252

(E)-form [136685-35-3]
 Isol. from the sponge *Tedania ignis*. Possibly an artifact.

Dillman, R.L. et al., *J. Nat. Prod.*, 1991, **54**, 1056 (isol, uv, ir, pmr, ms, cmr, struct)

1-(1*H*-Indol-3-yl)-1,2,3-propanetriol, 9CI I-135

3-(1,2,3-Trihydroxypropyl)indole. 3-(3-Indolyl)glycerol
 [13615-41-3]



C₁₁H₁₃NO₃ 207.229


Alkaloid from the clavicipitaceous fungus *Balansia epichloë* which parasitizes pasture grasses. The nat. prod. was a 3:2 mixt. of *erythro* and *threo* isomers. The racemates have been synthesised and characterised. λ_{max} 220 (log ε 4.95); 273 (log ε 4); 280 (log ε 4.02); 289 (log ε 3.95) (MeOH).

[66701-28-8, 66701-27-7]

Porter, J.K. et al., *J. Agric. Food Chem.*, 1977, **25**, 88-93 (isol, synth, uv, ir, pmr, ms)

3-(1*H*-Indol-3-yl)-2-propenoic acid I-136

1*H*-Indole-3-propenoic acid. 3-(3-Indolyl)acrylic acid
 [1204-06-4]



(*E*)-form

C₁₁H₉NO₂ 187.198

Major auxin from roots of *Lens culinaris* (lentil) (Fabaceae).

►Exp. carcinogenic data. NL3680000

(E)-form [29953-71-7]
 Alkaloid from the red alga *Chondria* sp. Red-brown cryst. (H₂O or AcOH). Mp 195-196°.

Me ester: [19626-92-7]
 [2756-97-0 (non-stereospecific)]
 C₁₂H₁₁NO₂ 201.224
 Cryst. (C₆H₆). Mp 153-154°.

Me ester, N-Ac: [19626-93-8]
 C₁₄H₁₃NO₃ 243.262
 Cryst. (C₆H₆). Mp 179-180°.

Amide: 3-(1*H*-Indol-3-yl)-2-propenamido. 3-(3-Indolyl)acrylamide
 [218591-77-6]

C₁₁H₁₀N₂O 186.213
 Alkaloid from *Chondria atropurpurea*. Anthelmintic agent. Amber cryst. (Me₂CO). Mp 212.8-214.2°. λ_{max} 226 (log ε 1.49); 275 (log ε 1.25); 322 (log ε 1.55) (MeOH). λ_{max} 226; 275; 322 (MeOH) (Berdy).

Amide, N^b-formyl: N-Formyl-3-(1*H*-indol-3-yl)-2-propenamido, 9CI. N-Formyl-3-indoleacrylamide
 [142677-08-5]
 C₁₂H₁₀N₂O₂ 214.223
 From *Chondria* sp. Oil.

Nitrile: [85452-78-4]

C₁₁H₈N₂ 168.198
 Beige solid. Mp 143-144°.

7'-Hydroxy-3-(7-Hydroxy-1*H*-indol-3-yl)-2-propenoic acid, 9CI. 7-Hydroxy-3-indoleacrylic acid
 [142677-11-0]
 C₁₁H₉NO₃ 203.197
 From *Chondria* sp.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 668B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 139B (nmr)

Furst, A. et al., *Arch. Biochem. Biophys.*, 1951, **31**, 190 (synth)

Inhoffen, H.H. et al., *Annalen*, 1963, **668**, 104-121 (*Me ester, Me ester N-Ac, synth*)

Rappe, C. et al., *Acta Chem. Scand.*, 1964, **18**, 818 (config)

Hofinger, M. et al., *Phytochemistry*, 1970, **9**, 1757 (isol)


Palermo, J.A. et al., *Tet. Lett.*, 1992, **33**, 3097 (*Chondria derivis, isol*)

Davyt, D. et al., *J. Nat. Prod.*, 1998, **61**, 1560-1563 (isol, amide)

Slätt, J. et al., *J. Het. Chem.*, 2005, **42**, 141-145 (nitrile)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, ICO000

N^a-[3-(1*H*-Indol-3-yl)propenoyl]tryptophan I-137



C₂₂H₁₉N₃O₃ 373.41

Deriv. of 3-(1*H*-Indol-3-yl)-2-propenoic acid, I-136.

(S,E)-form

L-trans-form

Et ester: [142677-12-1]


C₂₄H₂₃N₃O₃ 401.464

Minor constit. of a red alga *Chondria* sp. Prob. artifact.

Palermo, J.A. et al., *Tet. Lett.*, 1992, **33**, 3097-3100 (*Et ester, iso*)

3-(1*H*-Indol-3-yl)quinoline, 9CI I-138

3-(3-Quinoliny)-1*H*-indole
 [146535-51-5]



C₁₇H₁₂N₂ 244.295


Alkaloid from *Peganum nigellastrum*. Cryst. (hexane/EtOH). Mp 173-175°.

Maguire, M.P. et al., *J. Med. Chem.*, 1994, **37**, 2129-2137 (synth, pmr)

Ma, Z.-Z. et al., *Phytochemistry*, 2000, **53**, 1075-1078 (isol)

5-(1*H*-Indol-3-yl)-2(3*H*)-thiazolethione I-139

3-(2-Mercaptothiazol-5-yl)-1*H*-indole
 [929282-55-3]




C₁₁H₈N₂S₂ 232.33

Prod. by *Escherichia coli* eDNA clone CSLG18.

Brady, S.F. et al., *J.A.C.S.*, 2007, **129**, 12102-12103 (isol, pmr, cryst struct)

Infracropicrine I-140

1,2,3,12-Tetrahydro-12-oxoindolo[3,2,1-de]pyrido[3,2,1-ij][1,5]naphthyridin-4-ium(1+), 9CI
 [91147-09-0]




C₁₇H₁₃N₂O⁺ 261.302

Alkaloid from the fruit bodies of the gilled agaric *Cortinarius infractus*. Pale yellow powder (as chloride). Mp 350° (darkens above 200°). Responsible for the bitter taste of the toadstool.

Steglich, W. *et al.*, *Tet. Lett.*, 1984, **25**, 2341
(*isol, uv, ir, pmr, cmr, ms, struct*)

Ingamine A

[156310-17-7]



$C_{30}H_{44}N_2O$ 448.69

Alkaloid from the marine sponge *Xestospongia ingens*. Exhibits *in vitro* cytotoxicity against murine leukaemia P388 cells. Glass. Sol. MeOH, EtOAc. $[\alpha]_D +131$ (c, 0.18 in MeOH).

Deoxy: Ingamine B

[156310-18-8]


$C_{30}H_{44}N_2$ 432.691

From *Xestospongia ingens*. Exhibits *in vitro* cytotoxicity against murine leukaemia P388 cells. Glass. Sol. MeOH, EtOAc. $[\alpha]_D +108$ (c, 0.5 in MeOH).

Kong, F. *et al.*, *Tetrahedron*, 1994, **50**, 6137; 1995, **51**, 2895 (*isol, ir, pmr, cmr, ms, struct, abs config*)

Ingamine

[155210-52-9]



$C_{26}H_{40}N_2O$ 396.615

Alkaloid from the Papua New Guinea *Xestospongia ingens*. Cytotoxic. Amorph. solid. Sol. MeOH, EtOAc. $[\alpha]_D +62$ (c, 0.14 in MeOH).

Deoxy: Keramaphidin B

[157536-35-1]

$C_{26}H_{40}N_2$ 380.615

Alkaloid from *Xestospongia ingens* and from the Okinawan marine sponge *Amphimedon* sp. Exhibits cytotoxicity against P388 murine leukaemia and KB human epidermoid carcinoma cells. Poss. biogenetic precursor of manzamine alkaloids (e.g. Manzamine A, M-91). Amorph. solid. $[\alpha]_D +29.8$ (c, 1.1 in MeOH). The isolate (Mp 131-132°) from *Amphimedon* sp. is racemic.

Kong, F. *et al.*, *Tet. Lett.*, 1994, **35**, 1643 (*isol, pmr, cmr, struct*)

Kobayashi, J. *et al.*, *Tet. Lett.*, 1994, **35**, 4383 (*Keramaphidin B*)


Kong, F. *et al.*, *Tetrahedron*, 1995, **51**, 2895 (*abs config*)

Tsuda, M. *et al.*, *Heterocycles*, 1997, **46**, 765-795 (rev)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (rev, *Keramaphidin B*)

Ingamine B

[164081-00-9]



$C_{27}H_{42}N_2O$ 410.642

Minor alkaloid from the Papua New Guinea sponge *Xestospongia ingens*. Powder. $[\alpha]_D +22.4$ (c, 0.25 in MeOH).

27,28-Dihydro, 24,25-didehydro(Z-), Ac:

Ingamine C acetate

[164081-02-1]


$C_{29}H_{44}N_2O_2$ 452.679

Minor alkaloid from *Xestospongia ingens*. Glass. $[\alpha]_D +41.6$ (c, 0.09 in MeOH).

Kong, F. *et al.*, *Tetrahedron*, 1995, **51**, 2895 (*isol, pmr, cmr, ms, struct*)

Ingamine D

[164081-03-2]




$C_{28}H_{42}N_2O$ 422.653

Minor alkaloid from the Papua New Guinea sponge *Xestospongia ingens*. Glass.

Kong, F. *et al.*, *Tetrahedron*, 1995, **51**, 2895 (*isol, pmr, cmr, ms, struct*)

Ingamine E

[164081-04-3]



$C_{30}H_{44}N_2O$ 448.69

Minor alkaloid from the Papua New Guinea sponge *Xestospongia ingens*. Glass. $[\alpha]_D -23.8$ (c, 0.062 in MeOH).

Deoxy: Ingamine F

[164122-97-8]

$C_{30}H_{44}N_2$ 432.691

Minor alkaloid from *Xestospongia*


I-143

ingens. Glass. $[\alpha]_D -64.3$ (c, 0.062 in MeOH).

Kong, F. *et al.*, *Tetrahedron*, 1995, **51**, 2895 (*isol, pmr, cmr, ms, struct*)

Ingamine G

[787551-03-5]



Probable Absolute Configuration


$C_{32}H_{50}N_2O$ 478.76

Alkaloid from the Brazilian sponge *Pachychalina* sp. Cytotoxic. Glassy solid. $[\alpha]_D^{20} -59.2$ (c, 0.05 in MeOH). λ_{max} 230 (ε 2500); 283 (ε 625) (MeOH).

De Oliveira, J.H.H.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1685-1689 (*isol, pmr, cmr*)

Ingol

[51847-86-0]



$C_{20}H_{30}O_6$ 366.453

Hydrol. prod. from *Euphorbia ingens* and *Euphorbia kamerunica*. Antineoplastic agent.

Tetra-Ac: [51906-02-6]

$C_{28}H_{38}O_{10}$ 534.602

Constit. of *Euphorbia kamerunica*.

Cryst. (isopropyl ether). Poorly sol.

hexane. Mp 172-175°. λ_{max} 212 (ε 2570); 226 (ε 1410) (MeOH) (Berdy).

3-O-(2-Methylbutanoyl), 7,8,12-tri-Ac:

[928298-74-2]

$C_{31}H_{44}O_{10}$ 576.683

Constit. of *Euphorbia cornigera*. Resin.

$[\alpha]_D^{25} +95.8$ (c, 0.17 in $CHCl_3$). λ_{max} 218 (log ε 4.32); 228 (log ε 4.93); 278 (log ε 2.93); 286 (log ε 2.82) (MeOH).

7-O-(2-Methylbutanoyl), 3,8,12-tri-Ac:

[928298-75-3]

$C_{31}H_{44}O_{10}$ 576.683

Constit. of *Euphorbia cornigera*.

8-(2-Methylbutanoyl), 3,7,12-tri-Ac:

[928298-76-4]

$C_{31}H_{44}O_{10}$ 576.683

Constit. of *Euphorbia tirucalli* and

Euphorbia cornigera. Oil or resin. $[\alpha]_D$

+92 ($CHCl_3$). $[\alpha]_D^{25} +94.8$ (c, 0.17 in

$CHCl_3$). λ_{max} 218 (log ε 4.32); 228 (log

ε 4.93); 278 (log ε 2.93); 286 (log ε 2.82)

(MeOH).

8-(3-Methylbutanoyl), 3,7,12-tri-Ac:

$C_{31}H_{44}O_{10}$ 576.683

Constit. of *Euphorbia acruensis*. Oil.

12-O-(2-Methylbutanoyl), 3,7,8-tri-Ac:

[928298-77-5]

$C_{31}H_{44}O_{10}$ 576.683

Constit. of *Euphorbia cornigera*. Resin.


$[\alpha]_D^{25} +97.8$ (c, 0.17 in $CHCl_3$). λ_{max} 218

(log ε 4.32); 228 (log ε 4.93); 278 (log ε

- 2.93); 286 (log ϵ 2.82) (MeOH).
- 7-Angeloyl, 3,12-di-Ac: [627512-36-1]
C₂₉H₄₀O₉ 532.63
Constit. of *Euphorbia nivulia*. [α]_D²⁵ -3.6 (c, 0.125 in CHCl₃).
- 7-Angeloyl, 3,8,12-tri-Ac: [92998-77-1]
C₃₁H₄₂O₁₀ 574.667
Constit. of *Euphorbia kamerunica*.
- 8-Angeloyl, 3,7,12-tri-Ac:
C₃₁H₄₂O₁₀ 574.667
Constit. of *Euphorbia kamerunica*.
- 7-Tigloyl, 3,12-di-Ac:
C₂₉H₄₀O₉ 532.63
Constit. of *Euphorbia kamerunica*.
- 7-Tigloyl, 3,8,12-tri-Ac: [92910-93-5]
C₃₁H₄₂O₁₀ 574.667
Constit. of *Euphorbia kamerunica*.
- 8-Tigloyl:
C₂₅H₃₆O₇ 448.555
Constit. of *Euphorbia antiquorum*. [α]_D +21.8 (c, 0.12 in CHCl₃).
- 8-Tigloyl, 12-Ac:
C₂₇H₃₈O₈ 490.592
Constit. of *Euphorbia antiquorum*. [α]_D +10 (c, 0.16 in CHCl₃).
- 8-Tigloyl, 3,12-di-Ac:
C₂₉H₄₀O₉ 532.63
Constit. of *Euphorbia antiquorum* and *Euphorbia lactea*. [α]_D -16.2 (c, 0.13 in CHCl₃).
- 8-Tigloyl, 3,7,12-tri-Ac:
C₃₁H₄₂O₁₀ 574.667
Constit. of *Euphorbia kamerunica*.
- 12-Tigloyl, 3,7-di-Ac: [81980-37-2]
C₂₉H₄₀O₉ 532.63
Constit. of *Euphorbia kamerunica*. Sol. MeOH, Et₂O; poorly sol. H₂O, hexane. λ_{\max} 209 (ϵ 30200); 224 (ϵ 15850) (MeOH) (Berdy).
- 12-Tigloyl, 3,7,8-tri-Ac: [81427-04-5]
C₃₁H₄₂O₁₀ 574.667
Constit. of *Euphorbia kamerunica*. Sol. MeOH, Et₂O; poorly sol. hexane. λ_{\max} 208 (ϵ 17000); 226 (ϵ 14120) (MeOH) (Berdy).
- 7,8-Ditigloyl, 3,12-di-Ac:
C₃₄H₄₆O₁₀ 614.731
Constit. of *Euphorbia acurensis*. Oil.
- 8-Benzoyl, 3,12-di-Ac: [124657-64-3]
C₃₁H₃₈O₉ 554.636
Constit. of *Euphorbia antiquorum* and *Euphorbia nivulia*. Powder. Mp 95-97°. [α]_D -21.8 (c, 0.16 in CHCl₃). [α]_D²⁵ -27 (c, 0.5 in CHCl₃).
- 8-Benzoyl, 3,7,12-tri-Ac:
C₃₃H₄₀O₁₀ 596.673
Constit. of *Euphorbia kamerunica*.
- 7-(Phenylacetyl), 3,8,12-tri-Ac:
C₃₄H₄₂O₁₀ 610.7
Constit. of *Euphorbia* spp.
- 8-(Phenylacetyl), 3,7,12-tri-Ac: [403740-67-0]
C₃₄H₄₂O₁₀ 610.7
Constit. of *Euphorbia resinifera*. Amorph. solid. [α]_D²⁵ +15 (c, 0.002 in CHCl₃).
- 8-(3-Pyridinecarbonyl), 3,7,12-tri-Ac:
C₃₂H₃₉NO₁₀ 597.661
Constit. of *Euphorbia ingens*.
- 8-(3-Pyridinecarbonyl), 7-benzoyl, 3,12-di-Ac: [462119-30-8]
C₃₇H₄₁NO₁₀ 659.732
Constit. of *Euphorbia nivulia*. Powder. Mp 90-92°. [α]_D²⁵ -61 (c, 0.5 in CHCl₃).
- 8-Me ether, 3,12-di-Ac: [625830-44-6]
C₂₅H₃₆O₈ 464.555
Constit. of *Euphorbia nivulia*. [α]_D²⁵ +35 (c, 0.25 in CHCl₃).
- 8-Me ether, 3,7,12-tri-Ac: **Tirucalicine** [77573-15-0]
C₂₇H₃₈O₉ 506.592
Constit. of *Euphorbia tirucalli*. Needles (EtOH). Mp 148-150°. [α]_D +16.13 (c, 3.41 in CHCl₃).
- 8-Me ether, 3-O-(2-methylbutanoyl), 7,12-di-Ac: [928298-78-6]
C₃₀H₄₄O₉ 548.672
Constit. of *Euphorbia cornigera*. Resin. [α]_D²⁵ -9.8 (c, 0.17 in CHCl₃). λ_{\max} 208 (log ϵ 4.32); 225 (log ϵ 4.93); 273 (log ϵ 2.93); 284 (log ϵ 2.82) (MeOH).
- 8-Me ether, 7-O-(2-Methylbutanoyl), 3,12-di-Ac: [928298-79-7]
C₃₀H₄₄O₉ 548.672
Constit. of *Euphorbia cornigera*. Resin. [α]_D²⁵ +4.8 (c, 0.17 in CHCl₃). λ_{\max} 208 (log ϵ 4.32); 225 (log ϵ 4.93); 273 (log ϵ 2.93); 284 (log ϵ 2.82) (MeOH).
- 8-Me ether, 12-O-(2-methylbutanoyl), 3,7-di-Ac: [928298-80-0]
C₃₀H₄₄O₉ 548.672
Constit. of *Euphorbia cornigera*. Resin. [α]_D²⁵ +9.8 (c, 0.17 in CHCl₃). λ_{\max} 208 (log ϵ 4.32); 225 (log ϵ 4.93); 273 (log ϵ 2.93); 284 (log ϵ 2.82) (MeOH).
- 8-Me ether, 7-angeloyl, 3-Ac: [625830-43-5]
C₂₈H₄₀O₈ 504.619
Constit. of *Euphorbia nivulia*. [α]_D -1 (c, 0.25 in CHCl₃).
- 8-Me ether, 7-angeloyl, 3,12-di-Ac: [90027-10-4]
C₃₀H₄₂O₉ 546.656
Constit. of *Euphorbia kamerunica*.
- 8-Me ether, 7-tigloyl, 3,12-di-Ac: [89984-06-5]
C₃₀H₄₂O₉ 546.656
Constit. of latex of *Euphorbia kamerunica*.
- 8-Me ether, 7-benzoyl, 3,12-di-Ac: [89984-05-4]
C₃₂H₄₀O₉ 568.663
Constit. of *Euphorbia kamerunica* and *Euphorbia hermentiana*. Gum. [α]_D -7.2 (c, 0.08 in CHCl₃).
- 2-Epimer, tetra-Ac:
C₂₈H₃₈O₁₀ 534.602
Constit. of *Euphorbia portulacoides*.
- 2-Epimer, 7-(2-methylpropanoyl), 3,8,12-tri-Ac:
C₃₀H₄₂O₁₀ 562.656
Constit. of *Euphorbia portulacoides*.
- 2-Epimer, 7-(2-methylbutanoyl), 3,8,12-tri-Ac:
C₃₁H₄₄O₁₀ 576.683
Constit. of *Euphorbia portulacoides*.
- 2-Epimer, 7-benzoyl, 3,8,12-tri-Ac:
C₃₃H₄₀O₁₀ 596.673
Constit. of *Euphorbia portulacoides*.
- 2-Epimer, 8-benzoyl, 3,7,12-tri-Ac:
C₃₃H₄₀O₁₀ 596.673
- Constit. of *Euphorbia canariensis*. Oil. [α]_D -41 (c, 1 in CHCl₃).
- 2,3-Diepimer, 8-benzoyl, 7,12-di-Ac:
C₃₁H₃₈O₉ 554.636
Constit. of *Euphorbia canariensis*. Oil. [α]_D -25 (c, 3.4 in CHCl₃).
- 2,3-Diepimer, 8-(2-methylpropanoyl), 7,12-di-Ac:
C₂₈H₄₀O₉ 520.619
Constit. of *Euphorbia canariensis*. Oil. [α]_D -8.5 (c, 1.4 in CHCl₃).
- Schroeder, G. et al., *Planta Med.*, 1979, **35**, 235 (derivs)
- Lotter, H. et al., *Tet. Lett.*, 1979, 77 (cryst struct)
- Abo, K. et al., *J. Nat. Prod.*, 1981, **45**, 365 (derivs)
- Abo, K. et al., *Phytochemistry*, 1981, **20**, 2535-2537 (derivs)
- Abo, K. et al., *Planta Med.*, 1981, **43**, 392 (derivs)
- Lin, L.-J. et al., *Phytochemistry*, 1983, **22**, 2795-2799 (isol, esters)
- Evans, F.J. et al., *Prog. Chem. Org. Nat. Prod.*, 1983, **44**, 1 (rev)
- Connolly, J.D. et al., *Tet. Lett.*, 1984, **25**, 3773 (struct, deriv)
- Khan, A.Q. et al., *Heterocycles*, 1988, **27**, 2851 (Tirucalicine)
- Gewali, M.B. et al., *Chem. Pharm. Bull.*, 1989, **37**, 1547 (derivs)
- Khan, A.Q. et al., *J. Nat. Prod.*, 1990, **53**, 728 (isol, deriv)
- Morgenstern, T. et al., *Phytochemistry*, 1996, **41**, 1149-1153 (*Euphorbia portulacoides* isolates)
- Marco, J.A. et al., *Phytochemistry*, 1997, **45**, 563-570; 1998, **49**, 1095-1099 (*Euphorbia canariensis* esters, *Euphorbia acurensis* esters)
- Ahmed, A.A. et al., *Fitoterapia*, 1999, **70**, 140-143 (8-tigloyl 3,12-di-Ac)
- Fattorusso, E. et al., *Eur. J. Org. Chem.*, 2002, 71-78 (8-phenylacetyl 3,7,12-tri-Ac)
- Ravikanth, V. et al., *Phytochemistry*, 2002, **59**, 331-335 (*Euphorbia nivulia* constits)
- Ravikanth, V. et al., *Chem. Pharm. Bull.*, 2003, **51**, 431-434 (*Euphorbia nivulia* constits)
- Baloch, I.B. et al., *Planta Med.*, 2006, **72**, 830-834 (*Euphorbia cornigera* constits)

Inosine, 9CI, INN, JAN I-148

1,9-Dihydro-9- β -D-ribofuranosyl-6H-purine-6-one, 9CI. 9- β -D-Ribofuranosylhypoxanthine, 8CI. Hypoxanthine riboside. Hypoxanthosine. Aminosin[†]. Carnine. Delimmin. Inosie. Oxiamine. Trophicardyl [58-63-9]



C₁₀H₁₂N₄O₅ 268.229
Present in meat extracts and sugar beet. Also prod. by microorganisms, e.g. *Bacillus subtilis*, *Escherichia coli*, *Saccharomyces cerevisiae*, *Fusarium* spp. A minor constit. of t-RNAs. Activates cellular functions. Cardiotonic. Suggested to be

capable of forming base pairs with Adenine, A-137, Cytosine, C-961 or Uracil, U-42 thus contributing to genetic code degeneracy by causing stable mis-pairings. Used to treat cardiac disorders. Egg-release pheromone of *Nereis succinea*. Mp 215° dec. $[\alpha]_D^{18}$ -49.2 (c, 0.9 in H₂O). p*K*_{a1} 1.5; p*K*_{a2} 8.85; p*K*_{a3} 12.5 (25°). Log P -3.76 (calc). λ_{\max} 249 (ε 12200) (H₂O) (pH 6).

► Probable mutagen. Exp. reprod. effects (very high dose) LD₅₀ (mus, ivn) 3000 mg/kg. NM7460000

3'-O- α -D-Glucopyranosyl: 3'-O- α -D-Glucopyranosylinosine

C₁₆H₂₂N₄O₁₀ 430.371

Isol. from the crustacean *Ligia exotica*. Amorph. solid. $[\alpha]_D^{20}$ +38 (c, 0.34 in H₂O). λ_{\max} 248 (ε 8400) (H₂O).

2'-Me: 2'-O-Methylinosine

[3881-21-8]

C₁₁H₁₄N₄O₅ 282.255

Isol. from the starfish *Asterias rolllestoni*. Mp 153-156°.

OH-form

6-O-[2-Hydroxy-1-(hydroxymethyl)ethyl]: 6-O-(1,3-Dihydroxyisopropyl)inosine

[193889-60-0]

C₁₃H₁₈N₄O₇ 342.308

Constit. of *Pedicularis longiflora*.

[35908-31-7]

Hall, R.H. et al., *Biochem. Biophys. Res. Commun.*, 1963, **13**, 394 (isol, deriv)

Munns, A.R.I. et al., *Acta Cryst. B*, 1970, **26**, 1114 (cryst struct)

Chenon, M.T. et al., *J.A.C.S.*, 1975, **97**, 4627 (cmr)

Westhof, E. et al., *Z. Naturforsch., C*, 1975, **30**, 131 (pmr)

Hawkes, G.E. et al., *J.C.S. Perkin 2*, 1977, 1268 (N-15 nmr)

Wang, C. et al., *CA*, 1997, **127**, 173867g (Dihydroxyisopropylinosine)

Zeeck, E. et al., *Chemoecology*, 1998, **8**, 77-84 (biochem, *Nereis*)

Kim, S.H. et al., *J. Nat. Prod.*, 2000, **63**, 1188-1191 (3'- α -D-glucosyl)

Li, G.-Q. et al., *J. Chin. Pharm. Sci.*, 2004, **13**, 81-86 (2'-O-Methylinosine, isol)

Insulamine

I-149

C₁₆H₁₇NO₃ 271.315


Amariyllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Narcissus* sp. (Amariyllidaceae). Cryst. (Me₂CO). Mp 177-178°. $[\alpha]_D^{25}$ -95 (c, 0.1 in CHCl₃). Boit, H.-G. et al., *Chem. Ber.*, 1957, **90**, 2197-2202 (isol)

Insularoline

I-150

Insulanoline

[478-62-6]



C₃₇H₃₈N₂O₆ 606.717

Alkaloid from the rhizomes of *Cyclea insularis* and roots of *Cyclea sutchuenensis* (Menispermaceae). Shows significant inhibitory activity against human colon adenocarcinoma. Plates + 1H₂O. Mp 195° dec. $[\alpha]_D^{14}$ +48.6 (MeOH). λ_{\max} 211 (ε 54954); 280 (ε 4467) (EtOH) (Berdy). λ_{\max} 287; 299 (EtOH-NaOH) (Berdy).

Me ether: Insularine. O-Methylinsularoline

[549-07-5]

C₃₈H₄₀N₂O₆ 620.744

Alkaloid from the rhizomes of *Cyclea insularis*, the roots of *Cyclea sutchuenensis* and *Cissampelos pareira*, the roots and terrestrial portion of *Stephania japonica*, and from the stems, roots and rhizomes of *Paracycloa ochiaiana* (Menispermaceae). Cryst. (Et₂O). Mp 157°. $[\alpha]_D^{22}$ +11.36 (c, 0.44 in EtOH).

Me ether, dipicrate: Mp 230-231° dec.

Me ether, 2 β -N-oxide: Insularine 2 β -N-oxide

C₃₈H₄₀N₂O₇ 636.743

Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae).

Me ether, 2' β -N-oxide: Insularine 2' β -N-oxide

C₃₈H₄₀N₂O₇ 636.743

Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae).

Tomita, M. et al., *Nippon Kagaku Kaishi*, 1943, **64**, 64; 70; 77; 142; 147; *CA*, **41**, 3803 (isol, struct)

Satomi, M. et al., *Itsuo Kenkyusho Nempo*, 1955, **6**, 31; *CA*, **50**, 10112f (isol, *Insularine*)

Tomita, M. et al., *Yakugaku Zasshi*, 1957, **77**, 997; 1967, **87**, 1285; *CA*, **52**, 3832e; **58**, 36765w (*Insularine*, *Insularoline*, isol, struct)

Kikuchi, T. et al., *Yakugaku Zasshi*, 1958, **78**, 1408; 1413; *CA*, **53**, 7219f; 7220a (isol, struct)

Bick, I.R.C. et al., *J.C.S.*, 1961, 1896 (pmr) Kunitomo, J. et al., *Yakugaku Zasshi*, 1962, **82**, 1152; *CA*, **58**, 4614a (abs config)


Baldas, J. et al., *J.C.S. Perkin 1*, 1972, 599 (ms) Dwuma-Badu, D. et al., *Phytochemistry*, 1975, **14**, 2520 (isol, uv, ir, ms)

Lai, S. et al., *Huaxue Xuebao*, 1993, **51**, 1133; *CA*, **120**, 129507w (*Insularine oxides*)

Insulopinine

I-151

[102965-05-9]



C₄₀H₄₆N₄O₂ 614.829


Alkaloid from *Melodinus insulae-pinorum* (Apocynaceae).

Batchily, F. et al., *Ann. Pharm. Fr.*, 1985, **43**, 359; *CA*, **105**, 75870d

Intebrine

I-152

[169238-47-5]



C₂₀H₁₉NO₇ 385.373


Alkaloid from leaves of *Berberis integririma*. Prisms (MeOH). Mp 193-194°. λ_{\max} 223 (log ε 4.98); 250 (log ε 4.62); 304 (log ε 4.45) (EtOH).

Karimov, A. et al., *Khim. Prir. Soedin.*, 1993, 65-70; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 53-57 (isol, uv, cryst struct)

Intebrinine

I-153

[169238-46-4]



C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Berberis integerrima*. Cryst. (MeOH)(as hydrochloride). Mp 223-224° (hydrochloride).


Karimov, A. et al., *Khim. Prir. Soedin.*, 1993, 70-73; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 57-60 (isol, uv, pmr, cryst struct)

Integerrenine

I-154

2-(Dimethylamino)-3-methyl-N-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]pentanamide, 9CI

[18067-39-5]



Absolute Configuration

C₃₁H₄₂N₄O₄ 534.697

Alkaloid from the roots of *Ceanothus integerrimus*, root bark of *Zizyphus nummularia*, bark of *Heisteria nitida* and leaves of *Melochia pyramidata*. Needles (CHCl₃/petrol). Mp 278° dec. $[\alpha]_D^{20}$ -228 (c, 0.2 in CHCl₃).

N-De-Me: Nummularine D. N-Demethylintegerrenine

[58775-93-2]

C₃₀H₄₀N₄O₄ 520.67

Alkaloid from the root bark of *Zizyphus nummularia* (Rhamnaceae). Cryst. (MeOH). Mp 265-268°. $[\alpha]_D^{20}$ -186 (c, 0.2 in CHCl₃).

Dihydro:

Needles (CHCl₃/petrol). Mp 318° (314-316°). [α]_D²⁰ -40.2 (c, 0.13 in CHCl₃).

Tschesche, R. *et al.*, *Chem. Ber.*, 1967, **100**, 3924-3936 (*isol, uv, cd, ir, pmr, ms, struct*)


Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (*Nummularine D*)

Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 220-227 (*isol*)

Medina, E. *et al.*, *Annalen*, 1981, 538-545 (*isol, uv, ir, pmr*)

Intergerressine I-155

2-(Dimethylamino)-N-[5,8-dioxo-3-phenyl-7-(phenylmethyl)-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]-3-methylbutanamide, 9CI [17948-40-2]



Absolute Configuration

C₃₃H₃₈N₄O₄ 554.688

Major alkaloid from the roots of *Ceanothus integerrimus*. Also isol. from *Retanilla ephedra* (Rhamnaceae). Mp 285° dec. [α]_D²⁰ -164 (c, 0.2 in CHCl₃).

Dihydro:


Needles (CHCl₃/petrol). Mp 321°. [α]_D²⁰ -74.6 (c, 0.13 in CHCl₃).

Tschesche, R. *et al.*, *Chem. Ber.*, 1967, **100**, 3924 (*isol, uv, cd, ir, pmr, ms, struct*)

Bhakuni, D.S. *et al.*, *Rev. Latinoam. Quim.*, 1974, **5**, 158; *CA*, **82**, 108803w (*isol*)

Integerrine I-156

2-(Dimethylamino)-N-[7-(1H-indol-3-ylmethyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]-3-methylbutanamide, 9CI [18397-13-2]



C₃₅H₃₉N₅O₄ 593.724

Minor alkaloid from roots of *Ceanothus integerrimus* (Rhamnaceae). Needles (CHCl₃/petrol). Mp 258°. Opt. rotn. not reported.

N-De-Me: **N-Demethylintegerrine**

[70440-98-1]

C₃₄H₃₇N₅O₄ 579.697

Alkaloid from the root bark of *Ceanothus integerrimus* var. *integerrimus* (Rhamnaceae). Mp 350°. No stereochem. shown.


Tschesche, R. *et al.*, *Tet. Lett.*, 1968, 1311 (*isol*),

ms, struct)

Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 220-227 (*N-de-Me*)

Integramine I-157

[289042-21-3]




C₂₈H₃₅N₂O₂ 417.59

Alkaloid from *Zanthoxylum integrifolium*. Prisms (Me₂CO). Mp 208-210°. λ_{max} 232 (log ε 4.08); 265 (log ε 4.26); 290 (sh) (log ε 3.81) (MeOH).

Liu, S.-L. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 571-574 (*isol, ir, uv, pmr, cmr*)

Integramycin I-158




C₃₆H₄₉NO₉ 639.784

Prod. by *Actinoplanes* sp. ATCC202188. HIV-1 integrase inhibitor. Powder. [α]_D²³ +30.6 (c, 2.35 in MeOH).

Singh, S.B. *et al.*, *Org. Lett.*, 2002, **4**, 1123-1126 (*isol, pmr, cmr*)

Integramidine I-159

N-[6-(6-Hydroxy-1,3-benzodioxol-5-yl)-naphtho[2,3-d]-1,3-dioxol-5-yl]-N-methylformamide, 9CI [72459-16-6]



C₂₀H₁₅NO₆ 365.342


A secobenzo[c]phenanthridine alkaloid. Minor alkaloid from the root-xylem of *Zanthoxylum integrifolium* (Rutaceae). Prisms (CHCl₃/MeOH). Mp 302-304°.

Ishii, H. *et al.*, *Heterocycles*, 1979, **12**, 1037 (*ir, pmr, synth*)

Intermedine†

[10285-06-0]

I-160



C₁₅H₂₅NO₅ 299.366

Ester of Retronecine, in T-188 with Trachelanthic acid. Diastereoisomeric with Indicine, I-68 and Echinatine, E-12. Main alkaloid from *Amsinckia lycopsoides* and *Amsinckia hispida* (Boraginaceae). Cryst. (usually obt. as a gum). Mp 140-142°. [α]_D²⁰ +7.8 (c, 1.49 in EtOH) (+4.4). ▶ Hepatotoxin.

O⁷-Ac: **7-Acetylintermedine**

[74243-01-9]

C₁₇H₂₇NO₆ 341.403

Alkaloid from leaves of *Symphytum uplandicum* (Boraginaceae). Identified hydrolytically.

O³,O⁷-Di-Ac: **3',7-Diacetylintermedine**

C₁₉H₂₉NO₇ 383.441

Alkaloid from *Amsinckia menziesii* var. *intermedia* (Boraginaceae). Oil. [α]_D²⁵ +2.5 (c, 0.57 in CHCl₃).

7-O-Propanoyl: **7-Propionylintermedine**

C₁₈H₂₉NO₆ 355.43

Alkaloid from *Pulmonaria obscura*.

7-O-(2-Methylpropanoyl): **7-Isobutyrylintermedine**

C₁₉H₃₁NO₆ 369.457

Alkaloid from *Pulmonaria obscura*.

7-O-(2-Methylbutanoyl): **7-(2-Methylbutyryl)intermedine**

C₂₀H₃₃NO₆ 383.484

Alkaloid from *Pulmonaria obscura*.

7-O-(3-Methylbutanoyl): **7-(3-Methylbutyryl)intermedine**. 7-Isovaleroylintermedine

C₂₀H₃₃NO₆ 383.484

Alkaloid from *Pulmonaria obscura*.

7-O-(3-Methyl-2-butenoyl): see Echiupnine, E-32

7-O-Angeloyl: see Echiumine, E-31

3'-Epimer: **Lycopsamine**

[10285-07-1]

C₁₅H₂₅NO₅ 299.366

Alkaloid from *Amsinckia lycopsoides*, *Amsinckia hispida*, *Eupatorium coelestinum*, *Anchusa officinalis*, *Echium plantagineum*, *Heliotropium spathulatum* etc. (Boraginaceae). Cryst. (usually obt. as gum). Mp 132-134°.

[α]_D +3.3. [α]_D²⁰ +1.4 (EtOH). Cryst. with difficulty. Ester of Retronecine, in T-188 with Viridifloric acid. The struct. is incorrectly given in the 1991 paper, which substitutes the struct. for Heliospathuline (error in struct. diagrams).

▶ Hepatotoxin.

3'-Epimer, N-oxide: **Lycopsamine N-oxide**

[95462-15-0]

C₁₅H₂₅NO₆ 315.366

Isol. from adult bodies of the Apocynaceae-feeding danaine butterfly *Idea leuconoe*. [α]_D³² +6 (c, 2.3 in MeOH).

3'-Epimer, O⁷-Ac: 7-Acetyllycopsamine
[73544-48-6]

C₁₇H₂₇NO₆ 341.403

Alkaloid from leaves of *Symphytum uplandicum*, *Anchusa officinalis* and *Anchusa menziesii* (Boraginaceae). Identified hydrolytically.

3'-Epimer, O³-Ac: 3'-Acetyllycopsamine
[86702-00-3]

C₁₇H₂₇NO₆ 341.403

Alkaloid from *Amsinckia menziesii* (Boraginaceae). Gum. [α]_D²⁵ +9.3 (c, 1.3 in CHCl₃).

3'-Epimer, O⁷, O³-di-Ac: 3',7-Diacetyllycopsamine
[86702-01-4]

C₁₉H₂₉NO₇ 383.441

Alkaloid from *Amsinckia menziesii* (Boraginaceae). Faint-amber oil. [α]_D²⁵ -2.6 (c, 1.7 in CHCl₃).

3'-Epimer, 7-O-propanoyl: 7-Propionyllycopsamine

C₁₈H₂₉NO₆ 355.43

Alkaloid from *Pulmonaria obscura*.

3'-Epimer, 7-O-(2-methylpropanoyl): 7-Isobutyryllycopsamine

C₁₉H₃₁NO₆ 369.457

Alkaloid from *Pulmonaria obscura*.

3'-Epimer, 7-O-(2-methylbutanoyl): 7-(2-Methylbutyryl)lycopsamine

C₂₀H₃₃NO₆ 383.484

Alkaloid from *Pulmonaria obscura*.

3'-Epimer, O³-(3-methylbutanoyl): 9-(3'-Isovaleryl)viridiflorylretronecine
[114926-21-5]

C₂₀H₃₃NO₆ 383.484

Alkaloid from roots of *Heliotropium curassavicum* (2 varieties) (Boraginaceae).

3'-Epimer, 7-O-(3-methylbutanoyl): 7-(3-Methylbutyryl)lycopsamine. 7-Isovaleryllycopsamine

C₂₀H₃₃NO₆ 383.484

Alkaloid from *Pulmonaria obscura*.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1955; 1980, **33**, 1105 (*isol. struct., synth. Acetylintermediate, Acetyllycopsamine*)

Broch-Due, A.I. *et al.*, *Acta Chem. Scand., Ser. B*, 1980, **34**, 75 (*isol., pmr*)

Herz, W. *et al.*, *Experientia*, 1981, **37**, 683 (*isol. props*)

Culvenor, C.C.J. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 958 (*isol*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Mackay, M.F. *et al.*, *Acta Cryst. C*, 1983, **39**, 785 (*cryst. struct., Lycopsamine, Intermediate*)

Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1983, **36**, 769 (*isol., pmr, cmr, ms, struct., Acetyllycopsamine, Diacetyllycopsamine*)

Zalkow, L.H. *et al.*, *J. Med. Chem.*, 1985, **28**, 687 (*synth*)

Davicino, J.G. *et al.*, *Phytochemistry*, 1988, **27**, 960 (*Isovalerylviridiflorylretronecine*)

Nishida, R. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 1787 (*Lycopsamine N-oxide*)

Wiedefeld, H. *et al.*, *Planta Med.*, 1991, **57**, 57B (*isol. from A. conyzoides*)

Kelley, R.B. *et al.*, *Phytochemistry*, 1992, **31**, 2513 (*3',7-Diacetylintermediate*)

Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*rev. pmr*)


Nambu, M. *et al.*, *Chem. Comm.*, 1997, 1619 (*synth*)

Haberer, W. *et al.*, *Planta Med.*, 2002, **68**, 480-482 (*isol., 7-acyl esters*)

Inthomycin

I-161

3-Hydroxy-2,2,4-trimethyl-10-(5-oxazolyl)-4,6,8-decatrienamide, 9CI



(4E,6E,8E)-form

C₁₆H₂₂N₂O₃ 290.361

(4E,6E,8E)-form

Inthomycin C

[133004-93-0]

Isol. from *Streptomyces* sp. Possesses antifungal and herbicidal props.

(4Z,6E,8E)-form

Inthomycin B

[133004-92-9]

From *Streptomyces* sp. Shows antifungal and herbicidal props.

10ξ-Hydroxy: Phthoxazolin D

C₁₆H₂₂N₂O₄ 306.361

From *Streptomyces* sp. KO-7888. Antifungal agent, herbicide (cellulose biosynthesis inhibitor). Powder. Sol. MeOH, EtOH, Me₂CO; poorly sol. H₂O, CHCl₃. Mp 63-69°. [α]_D²⁵ +51 (c, 0.2 in MeOH).

(4Z,6Z,8E)-form

Inthomycin A. Phthoxazolin A. OM 5714.

Antibiotic OM 5714

[130288-22-1]

Prod. by *Streptomyces* sp. OM-5714. Specific inhibitor of cellulose biosynth. Mp 58-62°. [α]_D²⁰ +81.8 (c, 1.52 in Me₂CO). [α]_D¹⁸ +37.4 (c, 1 in CH₂Cl₂). Substruct. element of Oxazolomycin. O-155. λ_{max} 203 (ε 8900); 215 (sh) (ε 8200); 243 (sh) (ε 7600); 253 (sh) (ε 8200); 265 (sh) (ε 10300); 275 (ε 12000); 285 (sh) (ε 10000); 330 (sh) (ε 1000) (MeOH) (Derrep).

►HE1790000

10ξ-Hydroxy (1): Phthoxazolin B

C₁₆H₂₂N₂O₄ 306.361

From *Streptomyces* sp. KO-7888. Antifungal agent, herbicide (cellulose biosynth. inhibitor). Powder. Sol. MeOH, EtOH, Me₂CO; poorly sol. H₂O, CHCl₃. Mp 63-65°. [α]_D²⁵ +141 (c, 0.6 in MeOH). λ_{max} 278 (MeOH) (Berdy).

10ξ-Hydroxy (2): Phthoxazolin C

C₁₆H₂₂N₂O₄ 306.361

From *Streptomyces* sp. KO-7888. Antifungal agent, herbicide (cellulose biosynth. inhibitor). Powder. Sol. MeOH, Me₂CO, EtOH; poorly sol. H₂O, CHCl₃. Mp 64-67°. [α]_D²⁵ +30 (c, 0.2 in MeOH). 10-Epimer of Phthoxazolin B. λ_{max} 267; 288 (MeOH) (Berdy).

Henkel, T. *et al.*, *Annalen*, 1991, 367 (*isol., pmr, cmr, struct*)

Tanaka, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 1208; 1214 (*Phthoxazolin*)

Shiomi, K. *et al.*, *J. Antibiot.*, 1995, **48**, 714 (*Phthoxazolins*)


Henaff, N. *et al.*, *Tetrahedron*, 2000, **56**, 5193-5204 (*synth*)

Webb, M.R. *et al.*, *Tetrahedron*, 2008, **64**, 4778-4791 (*synth*)

Involucratine

I-162

1-[[Dodecahydro-6,9-bis(methylene)-2-oxoazuleno[4,5-b]furan-3-yl]methyl]proline, 9CI. Saussureamine B [126209-82-3]



C₂₀H₂₇NO₄ 345.438

Alkaloid from *Saussurea involucrata* and *Saussurea lappa* (Asteraceae). Powder. [α]_D²⁵ -25.9 (c, 0.8 in MeOH). Involucratine and Saussureamine B not compared.

Li, Y. *et al.*, *Phytochemistry*, 1989, **28**, 3395 (*Involucratine*)

Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 214-216 (*Saussureamine B*)

Matsuda, H. *et al.*, *Tetrahedron*, 2000, **56**, 7763-7777 (*synth, abs config*)

Iodoacetic acid, 9CI, 8CI

I-163

[64-69-7]

ICH₂COOH

C₂H₃IO₂ 185.949

Constit. of the red alga *Asparagopsis taxiformis*. Used as a chromatographic derivatisation reagent for the S-carboxymethylation of cystine and cysteine in peptides. Plates (petrol). V. sol. Et₂O; sol. H₂O, EtOH. Mp 82-83°. pK_a 3.18 (25°, H₂O).

►Corrosive and irritating to eyes, skin and mucous membranes. LD₅₀ (mus, orl) 83 mg/kg. AI3500000

Me ester: Methyl iodoacetate

[5199-50-8]

C₃H₅IO₂ 199.976

Liq. Bp 169-171°.

Et ester: Ethyl iodoacetate. KSK

[623-48-3]

C₄H₇IO₂ 214.003

War gas, tear gas. Oil. Bp 178-180°

Bp₁₆ 73°.

►Highly lachrymatory and irritant. AI3575000

Chloride: Iodoacetyl chloride

[38020-81-4]

C₂H₂ClIO 204.394

Yellow-brown oil. d₂₅²⁵ 2.25. Bp₁₅ 49-52°.

Amide: Iodoacetamide, 9CI

[144-48-9]

C₂H₄INO 184.964

Constit. of *Asparagopsis taxiformis*. Reagent for cleavage of methionine containing peptides. Flaky cryst. (H₂O). Mp 95°.

►LD₅₀ (mus, orl) 74 mg/kg. AC4200000
Nitrile: Iodoacetoneitrile. Cyanoiodo-

methane

[624-75-9]

C₂H₂IN 166.949Oil. Bp₁₂ 76-77°.

Anhydride: [54907-61-8]

C₄H₄L₂O₃ 353.883Cryst. (EtOH aq.). V. sol. CHCl₃, EtOAc, Et₂O; less sol. H₂O, EtOH. Mp 46°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 509C; 651C; 751B; 848D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 791B; 1010A; 1172A; 1366B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 701A; 807C (ir)

v. Braun, J. et al., Ber., 1908, 41, 2130 (synth, nitrile, amide)

Abderhalden, E. et al., Ber., 1908, 41, 2852 (synth)

Jacobs, W.A. et al., J.A.C.S., 1919, 41, 2090 (amide)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, 1, 504 (amide, use)

Paraskewas, S. et al., Synthesis, 1974, 574 (amide)

Moodie, L.M. et al., J. Chromatogr., 1976, 124, 315 (use)

Woolard, F.X. et al., Tetrahedron, 1976, 32, 2843-2846 (amide, isol)

Karpitskaya, L.G. et al., Zh. Org. Khim., 1976, 12, 462 (synth)

Woolard, F.X. et al., Phytochemistry, 1979, 18, 617-620 (isol)

Dillon, K.B. et al., J. Magn. Reson., 1980, 39, 499 (pmr)

Lorenzi-Riatsch, A. et al., Helv. Chim. Acta, 1981, 64, 1563 (ms)

Cooper, J.D.H. et al., J. Chromatogr., 1982, 227, 158 (use)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 4, 2822-2824 (nitrile, use)

Luxon, S.G. et al., Hazards in the Chemical Laboratory, 5th edn., Royal Society of Chemistry, 1992, 730

Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold, 1992, ELQ000; IDW000; IDZ000

(Iodomethyl)trimethylammonium(1+) I-164

1-Iodo-N,N,N-trimethylmethanaminium, 9CI

[39895-69-7]

Me₃N⁺CH₂IC₄H₁₁IN⁺ 200.042Isol. from the Japanese gastropod *Turbo marmorata*.

Chloride: [65603-17-0]

C₄H₁₁ClIN 235.495

Counterion for natural product.

Iodide: [39741-91-8]

C₄H₁₁I₂N 326.947

Cryst. (1,2-ethanediol). Mp 192°

dec.(approx.) Mp 240° dec.(approx.)

(double Mp).

Schreiber, J. et al., Angew. Chem., Int. Ed., 1971, 10, 330-331 (synth)

Laber, R.A. et al., Spectrochim. Acta A, 1977, 33, 1079-1081 (ir, Raman)

Kotaki, Y. et al., CA, 1978, 88, 100056h (isol)


Wittmann, H. et al., Monatsh. Chem., 1982, 113, 1451-1458 (synth, pmr)

Almarzoqi, B. et al., Tetrahedron, 1986, 42, 601-607 (synth)

Ipalbidine

I-165

4-(1,2,3,5,8,8a-Hexahydro-7-methyl-6-indoliziny)phenol, 9CI. 6-(4-Hydroxyphenyl)-7-methyl-1,2,3,5,8,8a-hexahydroindolizine

C₁₅H₁₉NO 229.321

(+) -form [31470-51-6]

Obt. by resoln. of racemate. Cryst. (C₆H₆/cyclohexane). Mp 72-82°. [α]_D²⁵ +233.5 (c, 1 in CHCl₃). Contains some solvent of crystallisation.

Picrate:

Cryst. (MeOH). Mp 183-185°. [α]_D²⁵ +112.6.Ac: [α]_D²⁵ +149.6 (c, 1 in MeOH).

(-) -form [31470-52-7]

Obt. by resoln. of racemate. Glass. Mp 82-84°. [α]_D²⁵ -237 (c, 1 in CHCl₃). [α]_D -190.5 (c, 1 in MeOH).

Hydrochloride:

Cryst. + 1H₂O (MeCN). Mp 103.5-105°. [α]_D²⁵ -170 (c, 0.1 in MeOH).

Picrate:

Cryst. (MeOH). Mp 184-186°. [α]_D²⁵ -114.4 (c, 1 in MeOH).

(±) -form [26294-41-7]

Alkaloid from the seeds of *Ipomoea alba* and *Ipomoea muricata* (Convolvulaceae). Cryst. (EtOAc or EtOH). Mp 149-150° (144-146°).

Hydrochloride:

Cryst. + 1H₂O (MeOH/Et₂O or MeCN). Mp 107-109°.

Picrate:

Cryst. (MeOH). Mp 163-165° Mp 178°.

O-β-D-Glucopyranoside: *Ipalbine*

[23544-46-9]

C₂₁H₂₉NO₆ 391.463Alkaloid from seeds of *Ipomoea alba* (Convolvulaceae). Cryst. (MeOH/MeCN). Mp 118°. [α]_D²³ +32.5 (c, 0.03 in EtOH). [α]_D²⁵ +65.8 (c, 0.7 in EtOH).O-[4-Hydroxy-E-cinnamoyl-(→6)-β-D-glucopyranoside]: *Ipomine*

[65370-71-0]

C₃₀H₃₅NO₈ 537.608Alkaloid from seeds of *Ipomoea muricata* (Convolvulaceae). Amorph. solid + 2H₂O (Et₂O). Mp 139-143°. [α]_D +46.4. Ikhiri et al isolated Ipalbidine and Ipomine with opt. rotns. completely different from those previously reported. They suggest that their Ipomine, [α]_D -46°(EtOH) is the enantiomer of that previously isol., but since the chirality of Ipomine is entirely due to a D-Glucose residue, this is extremely improbable. It could be the glycoside of an enantiomer of Ipalbidine. They also suggest photoisomerism of the p-Hydroxycinnamoyl residue as a possibility.O-[4-Hydroxy-Z-cinnamoyl-(→6)-β-D-glucopyranoside]: *Isoipomine*

[110044-49-0]

C₃₀H₃₅NO₈ 537.608Alkaloid from the seeds of *Ipomoea alba* (Convolvulaceae). Obt. in admixture with Ipomine.O-[4-Hydroxy-3-methoxycinnamoyl-(→6)-β-D-glucopyranoside]: *Methoxyipomine*

[110011-73-9]

C₃₁H₃₇NO₉ 567.635Alkaloid from the seeds of *Ipomoea alba* (Convolvulaceae). Amorph.O-[4-Hydroxy-3,5-dimethoxycinnamoyl-(→6)-β-D-glucopyranoside]: *Di-methoxyipomine*

[110011-74-0]

C₃₂H₃₉NO₁₀ 597.661Alkaloid from the seeds of *Ipomoea alba* (Convolvulaceae). Amorph.

Ac: Mp 67-69°.

Gourley, J.M. et al., Chem. Comm., 1969, 709 (isol, uv, ir, pmr, ms, struct, Ipalbine, Ipalbidine)

Govindachari, T.R. et al., Tetrahedron, 1970, 26, 3829 (synth)

Wick, A.E. et al., Helv. Chim. Acta, 1971, 54, 513 (synth, uv, ir, pmr, ms, resoln)

Stevens, R.V. et al., Tet. Lett., 1977, 979 (synth)

Dawidar, A.M. et al., Tetrahedron, 1977, 33, 1733 (isol, uv, ir, ms)

Chari, V.M. et al., Planta Med., 1978, 34, 93 (cmr, struct)

Hedges, S.H. et al., J. Chem. Res., Synop., 1979, 1 (synth)

Howard, A.S. et al., J.O.C., 1980, 45, 1713 (synth)

Cragg, J.E. et al., Tet. Lett., 1981, 2127 (synth)

Iida, H. et al., J.C.S. Perkin I, 1985, 261 (synth)

Jefford, C.W. et al., Helv. Chim. Acta, 1986, 69, 2048 (synth, uv, ir, pmr, ms)

Danishefsky, S.J. et al., J.O.C., 1986, 51, 3915 (synth)

Ikhiri, K. et al., J. Nat. Prod., 1987, 50, 152 (isol, pmr, cmr, ms, derivs)

Sheehan, S.M. et al., J.O.C., 1997, 62, 438 (synth)


Ikeda, M. et al., Heterocycles, 1999, 50, 31-34 (synth)

Honda, T. et al., Tet. Lett., 2003, 44, 3035-3038 (synth)

Ipanguline D₃

I-166

[210779-90-1]

C₁₃H₂₃NO₅ 273.328Alkaloid from *Ipomoea hederifolia* (Convolvulaceae). Other Ipangulines have been isol. but were not fully characterised.3''-Ac: *Ipanguline D₅*


[210779-91-2]

C₁₅H₂₅NO₆ 315.366

- Alkaloid from *Ipomoea hederifolia* (Convolvulaceae).
- 2'',3''-Di-Ac: **I panguline D₉**
[210698-86-5]
C₁₇H₂₇NO₇ 357.403
Alkaloid *Ipomoea hederifolia* (Convolvulaceae). Stereochem. of side-chain unclear.
- 7-O-(2-Methylbutanoyl): **I panguline C₁**
[210779-92-3]
C₁₈H₃₁NO₆ 357.446
Alkaloid from *Ipomoea hederifolia*. Abs. config. of 2-methylbutanoyl residue not determined.
- 7-O-(2-Methylbutanoyl), 3''-Ac: **I panguline C₂**
[210779-93-4]
C₂₀H₃₃NO₇ 399.483
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(2-Methylbutanoyl), 2'',3''-di-Ac: **I panguline C₃**
[210698-91-2]
C₂₂H₃₅NO₈ 441.52
Alkaloid from *Ipomoea hederifolia*.
- 3''-(2-Methylbutanoyl): **I panguline D₁₀**
[210698-88-7]
C₁₈H₃₁NO₆ 357.446
Alkaloid from *Ipomoea cristulata* (Convolvulaceae).
- 2'' or 3''-Tigloyl: **I panguline D₁₄**
C₁₈H₂₉NO₆ 355.43
Alkaloid from *Ipomoea x peregrina* (Convolvulaceae). Stereochem. of side-chain uncertain.
- 2'' or 3''-Tigloyl, stereoisomer: **I panguline D₁₅**
C₁₈H₂₉NO₆ 355.43
Alkaloid from seeds of *Ipomoea x peregrina* (Convolvulaceae). Stereochem. of side-chain uncertain.
- 3'',7-Bis(2-methylbutanoyl): **I panguline C₄**
[210698-92-3]
C₂₃H₃₉NO₇ 441.564
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(2-Hydroxybenzoyl): **I panguline B₁**
[152273-76-2]
C₂₀H₂₇NO₇ 393.436
Alkaloid from above-ground parts (without fruits) of *Ipomoea hederifolia*.
- 7-O-(2-Hydroxybenzoyl), 3''-Ac: **I panguline B₂**
[210779-99-0]
C₂₂H₂₉NO₈ 435.473
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(2-Hydroxybenzoyl), 2'',3''-di-Ac: **I panguline B₃**
[210698-96-7]
C₂₄H₃₁NO₉ 477.51
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(Phenylacetyl): **I panguline A₁**
[152273-75-1]
C₂₁H₂₉NO₆ 391.463
Alkaloid from seeds of *Ipomoea hederifolia* (Convolvulaceae). Oil. [α]_D²¹ -69 (c, 0.98 in MeOH).
- 7-O-(Phenylacetyl), 3''-Ac: **I panguline A₂**
[210779-98-9]
C₂₃H₃₁NO₇ 433.5
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(Phenylacetyl), 2'',3''-di-Ac: **I panguline A₃**
[210780-00-0]
C₂₅H₃₃NO₈ 475.538
Alkaloid from *Ipomoea hederifolia*.
- 7-O-(Phenylacetyl), 3''-(2-methylbutanoyl): **I panguline A₄**
[210698-97-8]
C₂₆H₃₇NO₇ 475.581
Alkaloid from *Ipomoea hederifolia*.
- 2''-Epimer: **Isoipanguline D₃**
C₁₃H₂₃NO₅ 273.328
Alkaloid from *Ipomoea hederifolia* (Convolvulaceae).
- 2''-Epimer, 3''-Ac: **Isoipanguline D₅**
C₁₅H₂₅NO₆ 315.366
Alkaloid from *Ipomoea hederifolia* (Convolvulaceae).
- 2''-Epimer, 3''-(2-methylbutanoyl): **Isoipanguline D₁₀**
C₁₈H₃₁NO₆ 357.446
Alkaloid from *Ipomoea hederifolia* and *Ipomoea quamoclit*.
- 2''-Epimer, 7-O-(2-methylbutanoyl): **Isoipanguline C₁**
C₁₈H₃₁NO₆ 357.446
Alkaloid from *Ipomoea hederifolia*.
- 2''-Epimer, 7-O-(2-methylbutanoyl), 3''-Ac: **Isoipanguline C₂**
C₂₀H₃₃NO₇ 399.483
Alkaloid from *Ipomoea hederifolia*.
- 2''-Epimer, 7-O-(2-hydroxybenzoyl): **Isoipanguline B₁**
[152375-56-9]
C₂₀H₂₇NO₇ 393.436
Alkaloid from above-ground parts (without fruits) of *Ipomoea hederifolia*. Oil. [α]_D²¹ -108 (c, 0.90 in MeOH).
- 2''-Epimer, 7-O-(2-hydroxybenzoyl), 3''-Ac: **Isoipanguline B₂**
C₂₂H₂₉NO₈ 435.473
Alkaloid from *Ipomoea hederifolia*.
- 2''-Epimer, 7-O-(phenylacetyl): **Isoipanguline A₁**
[152375-55-8]
C₂₁H₂₉NO₆ 391.463
Alkaloid from seeds of *Ipomoea hederifolia* (Convolvulaceae). Oil. [α]_D²¹ -61 (c, 1.47 in MeOH).
- 2''-Epimer, 7-O-(phenylacetyl), 3''-Ac: **Isoipanguline A₂**
C₂₃H₃₁NO₇ 433.5
Alkaloid from *Ipomoea hederifolia*.
- 2''-Epimer, 7-O-(phenylacetyl), 2'',3''-di-Ac: **Isoipanguline A₃**
C₂₅H₃₃NO₈ 475.538
Alkaloid from *Ipomoea hederifolia*.
- Jenett-Siems, K. *et al.*, *Phytochemistry*, 1993, **34**, 437; 1998, **47**, 1551-1560; 2005, **66**, 223-231 (*isol, pmr, cmr, ms, struct*)

Ipecosidic acid

[137318-69-5]



Absolute Configuration

C₂₆H₃₃NO₁₂ 551.546

Most refs. refer to Ipecoside. Alkaloid from *Cephaelis ipecacuanha* (Rubiaceae). Cryst. (MeOH). Mp 248-250°. [α]_D²² -166 (c, 0.2 in MeOH). λ_{max} 207 (log ε 4.52); 226 (log ε 4.19); 233 (sh) (log ε 4.15); 286 (log ε 3.6) (MeOH).

Me ester: **Ipecoside**

[15401-60-2]

C₂₇H₃₅NO₁₂ 565.573

Alkaloid from the roots of *Cephaelis acuminata* and *Cephaelis ipecacuanha* (Rubiaceae). Mp 174-175° (monohydrate) Mp 176-178° (dihydrate).

N-De-Ac, Me ester: **Deacetylpecoside**

[21618-62-2]

C₂₅H₃₃NO₁₁ 523.536

Alkaloid from *Alangium lamarckii*. [α]_D -104 (c, 0.8 in MeOH) (as hydrochloride).

N-De-Ac, N-(4-hydroxy-3-methoxy-E-cinnamoyl), Me ester: **trans-Cephaeloside**

[148717-73-1]

C₃₅H₄₁NO₁₄ 699.707

Alkaloid from roots of *Cephaelis ipecacuanha* (Rubiaceae). Needles (MeOH). Mp 170-172°. [α]_D²⁷ -193 (c, 1.0 in MeOH).

N-De-Ac, N-(4-hydroxy-3-methoxy-Z-cinnamoyl), Me ester: **cis-Cephaeloside**

[148717-74-2]

C₃₅H₄₁NO₁₄ 699.707

Alkaloid from roots of *Cephaelis ipecacuanha* (Rubiaceae). Needles + 1½H₂O (MeOH). Mp 164-165°. [α]_D²⁸ -183 (c, 1.0 in MeOH).

Hexa-Ac, Me ester:

Amorph. powder + 2H₂O. [α]_D²⁵ -168.8 (c, 0.77 in CHCl₃).

6-Me ether, Me ester: **6-O-Methylpecoside**

[137318-68-4]

C₂₈H₃₇NO₁₂ 579.6

Alkaloid from *Cephaelis ipecacuanha* (Rubiaceae). Needles (MeOH/Et₂O). Mp 154-155°. [α]_D²² -184 (c, 1 in MeOH).

6-Me ether, N-de-Ac: **N-Deacetyl-6-O-methylpecosidic acid**

[265096-92-2]

C₂₅H₃₃NO₁₁ 523.536

Alkaloid from *Alangium kurzii*. Cryst. (MeOH). Mp 203-205°. [α]_D -144 (c, 0.2 in Py). [α]_D -124 (c, 0.34 in MeOH). λ_{max} 207 (log ε 4.51); 228 (log ε 4.18); 286 (log ε 3.56); 291 (sh) (log ε 3.52) (MeOH).

6-Me ether, N-de-Ac, N-(4-hydroxy-3-methoxy-E-cinnamoyl), Me ester: **6-O-Methyl-trans-cephaeloside**

[147677-04-1]

C₃₆H₄₃NO₁₄ 713.734

Alkaloid from roots of *Cephaelis ipecacuanha* (Rubiaceae). Needles (MeOH). Mp 162-164°. [α]_D²² -189 (c, 1.0 in MeOH).

6-Me ether, N-de-Ac, N-(4-hydroxy-3-methoxy-Z-cinnamoyl), Me ester: 6-O-Methyl-cis-cephaeloside

[147731-71-3]

C₃₆H₄₃NO₁₄ 713.734

Alkaloid from roots of *Cephaelis ipecacuanha* (Rubiaceae). Needles + 1½H₂O (MeOH). Mp 154-155°. [α]_D²⁸ -157 (c, 1.0 in MeOH).

7-Me ether, Me ester: 7-O-Methylpecoside

[137318-76-4]

C₂₈H₃₇NO₁₂ 579.6

Alkaloid from roots of *Cephaelis ipecacuanha* (Rubiaceae). Amorph. powder. [α]_D³¹ -194 (c, 1.0 in MeOH).

6,7-Di-Me ether, Me ester: [73611-00-4]

[34461-36-4]

Cryst. + 1.5 H₂O (MeOH/Et₂O). Mp 136-138°. [α]_D²¹ -168 (c, 0.54 in MeOH). [α]_D²⁸ -190 (c, 1 in MeOH).

4R-Hydroxy, Me ester: 4R-Hydroxypecoside

C₂₇H₃₅NO₁₃ 581.572

Alkaloid from the roots of *Cephaelis acuminata*. Amorph. powder. [α]_D²⁴ -140 (c, 0.37 in MeOH). λ_{max} 208 (log ε 4.5); 227 (log ε 4.17); 237 (sh) (log ε 4.09); 286 (log ε 3.53) (MeOH).

6-Deoxy, 8-hydroxy, Me ester: Neoipecoside

[122587-76-2]

C₂₇H₃₅NO₁₂ 565.573

Alkaloid from *Cephaelis ipecacuanha* (Rubiaceae). Needles (MeOH). Mp 184-185°. [α]_D²⁸ -161 (MeOH).

6-Deoxy, 8-hydroxy, O⁷-Me, Me ester: 7-Methylneoipecoside

[122587-77-3]

C₂₈H₃₇NO₁₂ 579.6

Alkaloid from *Cephaelis ipecacuanha* (Rubiaceae). Powder. [α]_D³⁰ -150 (EtOH).

6-Deoxy, 8-hydroxy, 3,4-didehydro, Me ester: 3,4-Dehydroneopecoside

[137318-70-8]

C₂₇H₃₃NO₁₂ 563.557

Alkaloid from *Cephaelis ipecacuanha* (Rubiaceae). Cryst. (H₂O). Mp 154-156°. [α]_D²³ -206 (c, 1.4 in MeOH).

6''-O-α-D-Glucopyranosyl, Me ester: 6''-O-α-D-Glucopyranosylpecoside

C₃₃H₄₅NO₁₇ 727.715

Alkaloid from the roots of *Cephaelis acuminata*. Amorph. powder. [α]_D²⁷ -108 (c, 0.24 in MeOH). λ_{max} 207 (log ε 4.53); 225 (log ε 4.18); 234 (sh) (log ε 4.11); 287 (log ε 3.61) (MeOH).

6''-O-β-D-Glucopyranosyl, Me ester: 6''-O-β-D-Glucopyranosylpecoside

C₃₃H₄₅NO₁₇ 727.715

Alkaloid from the roots of *Cephaelis acuminata*. Amorph. powder. [α]_D²⁸ -149 (c, 1 in MeOH).

1-Epimer, N-de-Ac, Me ester: Deacetylisoipecoside

[69400-51-7]

C₂₅H₃₃NO₁₁ 523.536

Alkaloid from *Alangium lamarckii*. [α]_D²⁸ -170 (c, 1 in MeOH) (as hydrochloride).

1-Epimer, 6-Me ether, N-de-Ac: N-Deacetyl-6-O-methylisoipecosidic acid

[342036-85-5]

C₂₅H₃₃NO₁₁ 523.536

Alkaloid from the dried fruit of *Alangium lamarckii*. Cryst. (MeOH). Mp 192-194°. [α]_D²² -203 (c, 1.1 in H₂O). λ_{max} 230 (log ε 4.14); 285 (log ε 3.58); 292 (sh) (log ε 3.52) (MeOH).

1-Epimer, 7-Me ether, N-de-Ac: N-Deacetyl-7-O-methylisoipecosidic acid

[342036-86-6]

C₂₅H₃₃NO₁₁ 523.536

Alkaloid from the dried fruit of *Alangium lamarckii*. Cryst. (MeOH). Mp 250-252°. [α]_D²⁴ -201 (c, 0.2 in H₂O). λ_{max} 229 (log ε 4.12); 285 (log ε 3.54); 292 (sh) (log ε 3.54) (MeOH).

1-Epimer, 6,7-di-Me ether, N-de-Ac: N-Deacetyl-6,7-di-O-methylisoipecosidic acid

[342036-87-7]

C₂₆H₃₅NO₁₁ 537.563

Alkaloid from the dried fruit of *Alangium lamarckii*. Cryst. (MeOH). Mp 198-201°. [α]_D²⁶ -205 (c, 0.5 in H₂O). λ_{max} 233 (log ε 4.2); 279 (sh) (log ε 3.51); 284 (log ε 3.54); 287 (sh) (log ε 3.52) (MeOH).

1-Epimer, 6''-O-α-D-glucopyranosyl, 16-Me ether, N-de-Ac: [342036-88-8]

C₃₁H₄₃NO₁₆ 685.678

Alkaloid from the dried fruit of *Alangium lamarckii*. Powder. [α]_D³⁰ -177 (c, 0.3 in H₂O). λ_{max} 228 (log ε 4.12); 285 (log ε 3.54); 292 (sh) (log ε 3.46) (MeOH).

Bellet, P. et al., *Ann. Pharm. Fr.*, 1954, **12**, 466-470; *CA*, **49**, 4234a (6,7-di-Me ether Me ester, synth)

Battersby, A.R. et al., *Chem. Comm.*, 1967, 219-221 (*Ipecoside*, uv, ir, pmr, ms, struct)

Battersby, A.R. et al., *J.C.S. (C)*, 1969, 1187-1192 (synth, uv, ir, pmr, ms)

Battersby, A.R. et al., *Chem. Comm.*, 1971, 901-902 (*biosynth*)

Garg, A.K. et al., *Phytochemistry*, 1972, **11**, 689-695 (*biosynth*)

Roberts, P.J. et al., *Acta Cryst. B*, 1974, **30**, 133-139 (6,7-di-Me ether Me ester, *cryst struct, abs config*)

Nagakura, N. et al., *Chem. Comm.*, 1978, 896 (*biosynth, cd*)

Hoefle, G. et al., *Chem. Ber.*, 1980, **113**, 566-576 (synth, uv, ir, pmr, ms)

Bhakuni, D.S. et al., *J.C.S. Perkin 1*, 1983, 1949-1952 (*Deacetylisoipecoside, Deacetylisoipecoside*)

Itoh, A. et al., *Chem. Pharm. Bull.*, 1989, **37**, 1137-1139 (*Neoipecoside, 7-O-Methylneoipecoside*)

Itoh, A. et al., *Phytochemistry*, 1991, **30**, 3117-3123 (*Ipecosidic acid, 6-O-Methylpecoside, 3,4-Dehydroneopecoside*)

Nagakura, N. et al., *Phytochemistry*, 1993, **32**, 761-765 (*Cephaelosides*)

Itoh, A. et al., *Phytochemistry*, 1994, **36**, 383-387 (*7-O-Methylpecoside*)


Tanahashi, T. et al., *Chem. Pharm. Bull.*, 2000, **48**, 415-419 (*N-Deacetyl-6-O-methylpecosidic acid*)

Itoh, A. et al., *Phytochemistry*, 2001, **56**, 623-630 (*Isoipecosidic acid derivs*)

Itoh, A. et al., *Phytochemistry*, 2002, **59**, 91-97 (*6''-Glucopyranosylpecosides, 4R-Hydroxypecoside*)

Ipobscurine B

[571144-95-1]

C₂₉H₃₀N₂O₇ 518.565


Alkaloid from the seeds of *Ipomoea obscura*. Yellow solid. [α]_D²⁰ -35 (c, 0.28 in MeOH).

Eich, E. et al., *Planta Med.*, 1989, **55**, 607 (*isol, struct*)

Jenett-Siems, K. et al., *Phytochemistry*, 2003, **62**, 1257-1263 (*isol, pmr, cmr*)

Ipobscurine C

[571144-96-2]

C₂₉H₂₈N₂O₇ 516.549

Alkaloid from seeds of *Ipomoea obscura*. Cryst. Mp 193-195°. [α]_D²⁰ -44 (c, 0.16 in CHCl₃).

Z-Isomer: Ipobscurine D

[571144-97-3]


C₂₉H₂₈N₂O₇ 516.549

Alkaloid from the seeds of *Ipomoea obscura*. Yellow solid. Mp 245-247°.

Jenett-Siems, K. et al., *Phytochemistry*, 2003, **62**, 1257-1263 (*isol, synth, pmr, cmr, ms*)

Ipohardine

2,3-Dihydro-6-(4-hydroxyphenyl)-7-methyl-1H-indolizinium, 9CI. *Ipalbidinium*
[108937-65-1]

C₁₅H₁₆NO⁺ 226.297

Alkaloid from *Ipomoea hardwickii* and *Ipomoea alba* (Convolvulaceae).

[108937-64-0, 108937-66-2]


Yu, Z. et al., *Jiegou Huaxue*, 1985, **4**, 152; 156; *CA*, **107**, 40161n; 40162p (*cryst struct*)

Ikhiri, K. et al., *J. Nat. Prod.*, 1987, **50**, 152 (*isol, uv, pmr, cmr*)

Daab, J.C. et al., *Monatsh. Chem.*, 2003, **134**, 573-583 (*synth*)

Ircinamine**I-171**

1-[(4,5-Dihydro-3-methoxy-4-methylene-1H-pyrrol-2-yl)thio]-1-tridecanone
[437980-03-5]




C₁₉H₃₃N₂O₂ 339.541
Alkaloid from the sponge *Ircinia* sp.
Cytotoxic. Amorph. solid. λ_{max} 242 (ε
1270) (MeOH).

Kuramoto, M. *et al.*, *Chem. Lett.*, 2002, 464-
465 (isol, pmr, cmr, ms)


Ircinamine B**I-172**

2-(Eicosylthio)-4,5-dihydro-3-methoxy-4-
methylene-1H-pyrrole
[915403-62-2]



C₂₆H₄₉NOS 423.745
Alkaloid from the sponge *Dactylia* sp.
Cytotoxic. Oil.

Sato, S. *et al.*, *Tet. Lett.*, 2006, 47, 7871-7873
(isol, pmr, cmr, ms)

Irciniastatin A**I-173***Psymberin*Absolute
Configuration

C₃₁H₄₇NO₁₁ 609.712
Isol. from the sponges *Ircinia* cf. *ramosa*
and *Psammocinia* aff. *bulbosa*. Cytotoxic.
Amorph. powder. [α]_D +29 (c, 0.02 in
MeOH). λ_{max} 215 ; 230 (sh) ; 270 ; 310
(MeOH aq.).

11-Ketone: Irciniastatin B

C₃₁H₄₅NO₁₁ 607.697
Isol. from *Ircinia* cf. *ramosa*. Cytotoxic.
Amorph. powder. [α]_D -4.7 (c, 0.15 in
MeOH). λ_{max} 215 ; 230 (sh) ; 270 ; 310
(MeOH).

Pettit, G.R. *et al.*, *J. Med. Chem.*, 2004, 47,
1149-1152 (isol, pmr, cmr)

Cichewitz, R.H. *et al.*, *Org. Lett.*, 2004, 6,
1951-1954 (*Psymberin*)

Jiang, X. *et al.*, *J.A.C.S.*, 2005, 127, 11254-
11255 (synth, abs config)

Kiren, S. *et al.*, *Org. Lett.*, 2005, 7, 2905-2908
(pmr, cmr, config)

Robinson, S.J. *et al.*, *J. Nat. Prod.*, 2007, 70,
1002-1009 (isol, activity)


Shangguan, N. *et al.*, *Org. Lett.*, 2007, 9, 1093-
1096 (synth)

Huang, X. *et al.*, *Org. Lett.*, 2007, 9, 2597-2600
(synth)

Rubio, B.K. *et al.*, *J.O.C.*, 2008, 71, 1475-1478
(isol)

Smith, A.B. *et al.*, *Org. Lett.*, 2008, 10, 5625-
5628 (synth)

Huang, X. *et al.*, *Tet. Lett.*, 2008, 49, 3592-
3595 (synth)

Ircinol A**I-174**

(+)-form

C₂₆H₄₀N₂O₂ 412.614

The enantiomer of Ircinol A itself so far
characterised has the opposite abs. con-
fig. to other Manzamine alkaloids, but
Ircinol A belonged to the 'normal' series.

(+)-form**Aldehyde: Ircinal A**

[139975-55-6]

C₂₆H₃₈N₂O₂ 410.598

Alkaloid from the Okinawan marine
sponge *Ircinia* sp. and *Amphimedon* sp.
Exhibits cytotoxicity against L1210
murine leukaemia cells and KB human
epidermoid carcinoma cells. Plausible
biogenetic precursor of manzamine
alkaloids. Mp 70°. [α]_D²⁵ +48 (c, 2.9 in
CHCl₃). λ_{max} 231 (ε 8500) (MeOH)
(Derep).

12-Deoxy, 12,28-epoxide, 1-aldehyde:**12,28-Oxaircinal A**C₂₆H₃₆N₂O₂ 408.583

Alkaloid from an *Acanthostrongylo-*
phora sp. Yellow powder. [α]_D²⁵ +56.6 (c,
0.06 in CHCl₃). λ_{max} 208 (log ε 3.63);
268 (log ε 3.78) (MeOH).

(-)-form [157000-77-6]

Alkaloid from the Okinawan marine
sponge *Amphimedon* sp. Cytotoxic. En-
dothelin converting enzyme inhibitor.
Amorph. solid. Mp 83-85°. [α]_D¹⁸ -19 (c,
0.54 in MeOH).

Kondo, K. *et al.*, *J.O.C.*, 1992, 57, 2480 (*Ircinal*
A)

Tsuda, M. *et al.*, *Tetrahedron*, 1994, 50, 7957
(isol, ir, pmr, cmr, ms, struct)

Tsuda, M. *et al.*, *Heterocycles*, 1997, 46, 765-
794 (rev)


Winkler, J.D. *et al.*, *J.A.C.S.*, 1998, 120, 6425-
6426 (synth)

Martin, S.F. *et al.*, *J.A.C.S.*, 1999, 121, 866-
867 (*Ircinal A*, synth)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 2006, 69, 1034-
1040 (12,28-Oxaircinal A)

Ircinol B**I-175**

[157000-78-7]

C₂₆H₄₂N₂O₂ 414.63

Alkaloid from the Okinawan marine
sponge *Amphimedon* sp. Cytotoxic, en-
dothelin converting enzyme inhibitor.
Amorph. solid. Mp 78-79°. [α]_D¹⁸ -2.8 (c,
0.12 in MeOH). Possesses opposite con-
fig. to Manzamine alkaloids.

Enantiomer, 1-aldehyde: Ircinal B

[139975-56-7]

C₂₆H₄₀N₂O₂ 412.614

Alkaloid from the Okinawan marine
sponge *Ircinia* sp. and *Amphimedon* sp.
Exhibits cytotoxicity against L1210
murine leukaemia cells and KB human
epidermoid carcinoma cells. Plausible
biogenetic precursor of manzamine
alkaloids. Mp 95°. [α]_D²⁵ +18 (c, 1.1 in
CHCl₃). λ_{max} 224 (ε 12000) (MeOH)
(Derep).

Kondo, K. *et al.*, *J.O.C.*, 1992, 57, 2480 (*Ircinal*
B)

Tsuda, M. *et al.*, *Tetrahedron*, 1994, 50, 7957
(isol, ir, pmr, cmr, ms struct)

Tsuda, M. *et al.*, *Heterocycles*, 1997, 46, 765-
794 (rev)

Irenine**I-176**C₁₇H₂₃NO₃ 289.374


Amaryllidaceae alkaloid. Struct. un-
known. Originally, Irenine was thought
to be Dihydroepigalanthamine, but this
was disproved by Barton *et al.* Isol. from
the bulbs of *Narcissus* sp. (Amaryllida-
ceae). Prisms (EtOAc). Mp 128°. [α]_D²⁵
+120 (c, 0.2 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, 90, 2197-
2202 (isol, ir)

Barton, D.H.R. *et al.*, *J.C.S.*, 1962, 806-817
(struct)

Irgigaine**I-177**

2-Methyl-6-(9-phenylnonyl)-3-piperidi-
nol. 3-Hydroxy-2-methyl-6-(9-phenylno-
nyl)piperidine
[180779-56-0]



Absolute Configuration

C₂₁H₃₅NO 317.514

Revised abs. config. Alkaloid from tubers
of *Arisarum vulgare* (Araceae). Toxic to
brine shrimp larvae. Oil. [α]_D²⁵ -14 (c, 0.3
in CHCl₃).

N-Me: N-Methylirgigaine

[180779-57-1]

C₂₂H₃₇NO 331.54

From tubers of *Arisarum vulgare*. Toxic
to brine shrimp larvae. Oil. [α]_D²⁵ -8 (c, 5
in CHCl₃).

Hydrochloride: [α]_D²⁰ +2.8 (c, 0.9 in
CHCl₃).


Melhaoui, A. *et al.*, *Nat. Prod. Lett.*, 1995, 7,
101 (isol, ir, pmr, cmr, struct)

Pahl, A. *et al.*, *Tet. Lett.*, 1998, 39, 2095-2096
(synth, ir, pmr, cmr, ms, cryst struct)

Ma, N. *et al.*, *Chin. J. Chem.*, 2003, 21, 1356-
1359 (synth)

Irniine

1-Methyl-2-(9-phenylnonyl)pyrrolidine,
9CI



C₂₀H₃₃N 287.487

(R)-form [144425-08-1]

Alkaloid from the tubers of *Arisarum vulgare* (Araceae). Cytotoxic. Displays anti gram-positive bacterial and antifungal activity. Oil. $[\alpha]_D^{21}$ -35 (c, 0.2 in CH₂Cl₂). $[\alpha]_D^{20}$ -55 (c, 1 in MeOH) (synthetic).

▶ Toxic.

2'-Methoxy-2-[9-(2-methoxyphenyl)-nonyl]-1-methylpyrrolidine. **Irnidine**
C₂₁H₃₅NO 317.514

Alkaloid from tubers of *Arisarum vulgare*. Oil. $[\alpha]_D^{25}$ -20 (c, 0.3 in CHCl₃).

▶ Toxic.

(S)-form

Synthetic. Displays anti gram-positive bacterial and antifungal activity. $[\alpha]_D^{20}$ +50.7 (c, 1.8 in MeOH).

Melhaoui, A. et al., *J. Nat. Prod.*, 1992, **55**, 950 (isol, ir, pmr, cmr, ms, struct)

Jossang, A. et al., *Heterocycles*, 1996, **43**, 755 (synth)

Takahata, H. et al., *Heterocycles*, 1997, **46**, 349-356 (synth)


Melhaoui, A. et al., *Planta Med.*, 1998, **64**, 476-477 (Irnidine)

Lamkadem, M. et al., *Nat. Prod. Res.*, 2004, **18**, 311-318; 2005, **19**, 573-580 (pmr, cmr, ms, activity)

Iromycin A

I-179

6-(3,7-Dimethyl-2,5-octadienyl)-4-hydroxy-3-methyl-5-propyl-2(1H)-pyridinone, 9CI. 2-(3,7-Dimethyl-2,5-octadienyl)-4,6-dihydroxy-5-methyl-3-propylpyridine. NK 26588. Antibiotic NK 26588 [213137-53-2]



Tautomeric. Prod. by *Streptomyces bottropensis* Gö Dra 17 and *Streptomyces* sp. NK 26588. Inhibitor of nitric oxide synthase. λ_{\max} 206 (ε 32120); 291 (ε 7065) (MeOH).

7'-Hydroxy: **Iromycin B**

[913690-66-1]

C₁₉H₂₉NO₃ 319.443

Prod. by *Streptomyces bottropensis* Gö Dra 17. λ_{\max} 207 (ε 41595); 290 (ε 9540) (MeOH).

Japan. Pat., 1998, 98 237 044; CA, **129**, 244203s (NK 26588)

Surup, F. et al., *J.O.C.*, 2007, **72**, 5085-5090; 5091-5097 (Iromycins, isol, synth, biosynth, pmr, cmr, cryst struct)

Iromycin C

I-180

[945634-84-4]

As Iromycin A, I-179 with

R = H

C₁₈H₂₇NO₂ 289.417

Prod. by *Streptomyces bottropensis* Gö Dra 17. Inhibitor of nitric oxide synthase. λ_{\max} 207 (ε 8430); 293 (ε 1990) (MeOH).

7'-ξ-Hydroxy: **Iromycin D**

[945634-85-5]

C₁₈H₂₇NO₃ 305.416


Prod. by *Streptomyces bottropensis* Gö Dra 17. $[\alpha]_D^{20}$ -150 (c, 0.1 in MeOH). λ_{\max} 206 (ε 3770); 289 (ε 910) (MeOH).

Surup, F. et al., *J.O.C.*, 2007, **72**, 5085-5090 (isol, pmr, cmr)

Isaindigotidione

I-181

[197720-99-3]



C₂₃H₂₂N₂O₅ 406.437


Alkaloid from the roots of *Isatis indigotica* (Brassicaceae). Cryst. Mp 240-242°. $[\alpha]_D^{20}$ +90.9 (c, 0.05 in DMSO). λ_{\max} 209 (log ε 4.92); 235 (sh) (log ε 4.53); 265 (log ε 4.02); 278 (sh) (log ε 3.95); 355 (log ε 3.89) (EtOH).

Wu, X. et al., *Tetrahedron*, 1997, **53**, 13323-13328 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Isaindigotone

I-182

[189316-00-5]



C₂₀H₁₈N₂O₄ 350.373

Alkaloid from the roots of *Isatis indigotica*. Yellow prisms. Mp 247-248°. $[\alpha]_D^{20}$ -5 (c, 0.02 in EtOH). λ_{\max} 210 (log ε 3.3); 228 (sh) (log ε 3); 365 (log ε 3.22) (EtOH).

Wu, X. et al., *Tetrahedron*, 1997, **53**, 13323-13328 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Molina, P. et al., *Synthesis*, 2000, 1523-1525 (synth)


Liu, J.-F. et al., *Org. Lett.*, 2005, **7**, 3363-3366 (synth, pmr, cmr)

Isatinone A

I-183

3-[Methoxy(phenyl)methylene]-2-indolinone

[936632-48-3]



C₁₆H₁₃NO₂ 251.284


Alkaloid from *Isatis costata*. Antifungal agent. Pale yellow solid. Mp 178-179°. λ_{\max} 208 ; 231 ; 270 (MeOH).

Fatima, I. et al., *Molecules*, 2007, **12**, 155-162 (isol, pmr, cmr, ms)

Isatinone B

I-184

[936632-49-4]



C₃₁H₃₃NO₄ 483.606


Alkaloid from *Isatis costata*. Antifungal agent. Pale yellow solid. Mp 189-191°. $[\alpha]_D^{18}$ +89.7 (c, 0.02 in MeOH). λ_{\max} 205 ; 232 ; 275 (MeOH).

Fatima, I. et al., *Molecules*, 2007, **12**, 155-162 (isol, pmr, cmr, ms)

Isatisine A

I-185

[953786-75-9]



C₂₂H₁₈N₂O₆ 406.394


Alkaloid from the leaves of *Isatis indigotifera*. Yellow needles (as 9,13-isopropylidene). $[\alpha]_D^{14}$ -283.1 (c, 0.46 in MeOH) (9,13-isopropylidene). Isol. as the 9,13 isopropylidene deriv., thought to be an artifact.

Liu, J.-F. et al., *Org. Lett.*, 2007, **9**, 4127-4129 (isol, pmr, cmr, cryst struct)

Ismine

I-186


6-[2-(Methylamino)phenyl]-1,3-benzodioxole-5-methanol, 9CI. 2-Hydroxy-methyl-2'-methylamino-4,5-methylene-dioxybiphenyl
[1805-78-3]



C₁₅H₁₅NO₃ 257.288


Alkaloid from the bulbs of an *Ismene* sp. (preferred genus name *Hymenocallis*), *Crinum powellii*, *Sprekelia formosissima*, *Narcissus bicolor*, *Narcissus pallidiflorus* and *Narcissus obesus* (Amaryllidaceae). Cryst. (C₆H₆/hexane). Mp 99.5-100.5°.

Picrate: Mp 158-159°.

N-Formyl: N-FormylisimineC₁₆H₁₅NO₄ 285.299Alkaloid from *Galanthus plicatus* ssp. byzantinus. Amorph. solid. λ_{\max} 208 (log ϵ 4.64); 292 (log ϵ 3.71) (MeOH).**Me ether: O-Methylisimine**C₁₆H₁₇NO₃ 271.315Alkaloid from the flowers of *Hippeastrum vittatum*. Oil. λ_{\max} 245 (log ϵ 4.16); 290 (log ϵ 4.65) (MeOH).Highet, R.J. et al., *J.O.C.*, 1961, **26**, 4767 (*isol, uv, ir, struct, bibl*)Hill, R.K. et al., *J.O.C.*, 1965, **30**, 1571 (*synth*)Fuganti, C. et al., *Tet. Lett.*, 1973, 1785*(biosynth)*Prabhakar, S. et al., *J. Chem. Res., Synop.*,1985, 394 (*synth*)Siddiqui, M.A. et al., *Tet. Lett.*, 1988, **29**, 5463*(synth, pmr)*Viladomat, F. et al., *Phytochemistry*, 1990, **29**,1307 (*isol*)Codina, C. et al., *Phytochemistry*, 1990, **29**,2685 (*isol*)Viladomat, F. et al., *J. Nat. Prod.*, 1992, **55**,804 (*isol*)Cowden, C.J. et al., *J. Nat. Prod.*, 1994, **57**,1746 (*pmr, cmr*)Viladomat, F. et al., *Acta Cryst. C*, 1998, **54**,81-82 (*cryst struct*)Unver, N. et al., *Heterocycles*, 2001, **55**, 641-652 (*N-Formylisimine*)Youssef, D.T.A. et al., *J. Nat. Prod.*, 2001, **64**,839-841 (*O-Methylisimine*)**Isoalchorneine****I-187**2,3,5,6-Tetrahydro-1-methoxy-2,6-bis(1-methylethenyl)-1H-imidazo[1,2-a]imidazole, 9CI
[41758-40-1]C₁₂H₁₉N₃O 221.302Alkaloid from the root bark and leaves of *Alchornea floribunda* (Euphorbiaceae).Liq. $[\alpha]_D$ 0 (c, 1 in CHCl₃).Khuong-Huu, F. et al., *Tetrahedron*, 1972, **28**,5207 (*isol, ir, pmr, ms, struct*)Büchi, G. et al., *J.O.C.*, 1989, **54**, 4494 (*synth*)**Isoalfileramine****I-188**

[73326-86-0]


[73365-21-6]

C₃₀H₄₂N₂O₂ 462.674Isol. from leaves *Zanthoxylum chiriquinum* (Rutaceae). Also prod. by treatment of Alfileramine, A-260 with acid. Tan cryst. (EtOAc/MeOH) (as hydrobromide). Mp 284-287° (hydrobromide). Opt. inactive. Poss. an artifact of isol. procedure.**N^a, N^b-Di-de-Me: N^a, N^b-Didemethylisofaliferamine**



[129743-96-0]

C₂₈H₃₈N₂O₂ 434.62Alkaloid from the leaves of *Zanthox-**ylum coriaceum* (Rutaceae). Opt. inactive.Caolo, M.A. et al., *Tetrahedron*, 1979, **35**, 1487; 1492 (*synth*)Marcos, M. et al., *J. Nat. Prod.*, 1990, **53**, 459 (*isol*)Marcos, M. et al., *Phytochemistry*, 1990, **29**, 2315 (*Didemethylisofaliferamine*)**Isoanantine****I-189**

4-(1-Methyl-1H-imidazol-5-yl)-3-(phenylmethylene)-2-pyrrolidinone, 9CI


C₁₅H₁₅N₃O 253.303 λ_{\max} 218 (ϵ 15800); 277 (ϵ 25000) (prob. MeOH) (Derep).**(-)-form** [50656-85-4]Alkaloid from the root bark of *Cynometra lujae* (Fabaceae). Cryst. (Me₂CO). Mp 190°. $[\alpha]_D^{20}$ -347 (c, 1 in CHCl₃).**(±)-form** [68108-27-0]

Synthetic. Mp 201°.

Tchissambou, L. et al., *Tetrahedron*, 1982, **38**, 2687 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)Naito, T. et al., *Chem. Pharm. Bull.*, 1997, **45**, 1932-1939 (*synth*)**Isoaquileidine****I-190**C₂₀H₂₀N₂O₅ 368.388Alkaloid from *Aquilegia ecalcarata*.Amorph. powder (MeOH). Mp 232-233°. $[\alpha]_D$ +19 (c, 0.43 in MeOH). λ_{\max} 286; 322 (sh) (MeOH).Chen, S.-B. et al., *J. Nat. Prod.*, 2001, **64**, 85-87**Isoasterriquinone****I-191***Isoasterriquinone* SU 5220. *Asterriquinone* SU 5220
[78860-47-6]C₃₂H₃₀N₂O₄ 506.6Prod. by *Aspergillus candidus*, *Aspergillus terreus* and *Aspergillus coradidus*. Also obt. by treatment of Asterriquinone, A-1520 with HCl/AcOH. Inhibits binding of proteins to the EGF receptor tyrosine kinase. Dark purple needles (hexane/CH₂Cl₂). Mp 150-151° dec. λ_{\max} 225; 283; 289; 450 (MeOH). λ_{\max} 225 (ϵ 47500); 282 (ϵ 31600); 288 (ϵ 31000); 477 (ϵ 2700) (EtOH) (Berdy).Arai, K. et al., *Chem. Pharm. Bull.*, 1981, **29**, 991 (*synth*)Kaji, A. et al., *Chem. Pharm. Bull.*, 1994, **42**, 1682 (*isol, uv, ir, pmr*)Alvi, K.A. et al., *J. Antibiot.*, 1999, **52**, 215-223 (*isol, uv, ir, pmr, cmr, activity*)**Isoatisine, 9CI****I-192**

[510-38-3]

[111001-28-6]

C₂₂H₃₃NO₂ 343.508Alkaloid from the roots of *Aconitum heterophyllum* (Ranunculaceae). Also obt. from Atisine, A-1534 in hot MeOH. Mp 149.5-152°. $[\alpha]_D$ -22.4 (c, 1.45 in 95% EtOH). pK_a 10.3 (50% MeOH).**Hydrochloride:**Needles (Me₂CO). Mp 304-307°. $[\alpha]_D^{30}$ -6.9 (c, 1.17 in 95% EtOH). Exists as a quaternary ammonium salt rather than a protonated salt.**7 α -Hydroxy: 7-Hydroxyisoatisine**

[70022-66-1]

C₂₂H₃₃NO₃ 359.508Alkaloid from *Consolida glandulosa*. Amorph.**19-Epimer: 19-Epiisoatisine**


[111001-27-5]

C₂₂H₃₃NO₂ 343.508Trace alkaloid from the Chinese herb Guan-Bai-Fu (*Aconitum koreanum*) (Ranunculaceae). Inseparable mixt. with Isoatisine.Pelletier, S.W. et al., *J.A.C.S.*, 1965, **87**, 777-798; 1977, **99**, 284-286; 1978, **100**, 7976-7987 (*ir, cmr, cryst struct, abs config*)Pelletier, S.W. et al., *Phytochemistry*, 1968, **7**, 625-635 (*isol, synth, ir*)Pelletier, S.W. et al., *Tetrahedron*, 1968, **24**, 2019-2038 (*pmr*)Mody, N.V. et al., *Tetrahedron*, 1978, **34**, 2421-2431 (*cmr*)Reinecke, M.G. et al., *J.O.C.*, 1987, **52**, 5051-5053 (*19-Epiisoatisine*)González-Coloma, A. et al., *Chem. Biodiversity*, 2004, **1**, 1327-1335 (*7-Hydroxyisoatisine*)

Isobatzelline A

I-193

7-Amino-6-chloro-3,4-dihydro-1-methyl-2-(methylthio)pyrrolo[4,3,2-de]quinolin-8(1H)-one, 9CI
[133401-01-1]



$C_{12}H_{12}ClN_3OS$ 281.765

Alkaloid from the marine sponge *Batzella* sp. Exhibits cytotoxic and moderate antifungal activities. Brown solid. λ_{max} 262 (ϵ 15500); 342 (ϵ 9000); 430 (ϵ 3100) (MeOH) (Derep).

3,4-Didehydro: Isobatzelline D

[133401-04-4]

 $C_{12}H_{10}ClN_3OS$ 279.749

From *Batzella* sp. Exhibits cytotoxic and moderate antifungal activities. Red-brown solid. λ_{max} 239 (ϵ 33900); 263 (ϵ 25000); 439 (ϵ 25200) (MeOH) (Derep).

Dechloro: Isobatzelline B

[133401-02-2]

 $C_{12}H_{13}N_3OS$ 247.32

From *Batzella* sp. Exhibits cytotoxic and moderate antifungal activities. Red-brown solid. λ_{max} 264 (ϵ 14200); 362 (ϵ 7000); 402 (ϵ 5000) (MeOH) (Derep).

De(methylthio): Isobatzelline C

[133401-03-3]

 $C_{11}H_{10}ClN_3O$ 235.672

From *Batzella* sp., *Zyzzya massalis* and *Zyzzya fuliginosa*. Exhibits cytotoxic and moderate antifungal activities. Green-brown or red solid. λ_{max} 244 (ϵ 9700); 344 (ϵ 5900); 394 (ϵ 2600) (MeOH) (Derep). λ_{max} 224 ($\log \epsilon$ 3.72); 341 ($\log \epsilon$ 2.26); 422 ($\log \epsilon$ 3.43) (MeOH).

De(methylthio), 3,4-didehydro: Isobatzelline E

[437980-21-7]

 $C_{11}H_8ClN_3O$ 233.656

Isol. from *Zyzzya fuliginosa* and *Zyzzya massalis*. Orange solid. λ_{max} 224 (ϵ 21500); 290 (ϵ 4800); 422 (ϵ 13400) (MeOH).

Sun, H.H. *et al.*, *J.O.C.*, 1990, **55**, 4964-4966 (isol, uv, ir, pmr, ms, struct, cmr)

Tao, X.L. *et al.*, *Tetrahedron*, 1994, **50**, 2017 (Isobatzelline C, synth)

Yamada, F. *et al.*, *Heterocycles*, 1995, **41**, 1905-1908 (Isobatzelline C, synth)

Roberts, D. *et al.*, *J.O.C.*, 1997, **62**, 568 (Isobatzelline C, synth)

Alvarez, M. *et al.*, *Tet. Lett.*, 1998, **39**, 679-680 (Isobatzelline B, synth)

Alvarez, M. *et al.*, *Eur. J. Org. Chem.*, 1999, 1173-1183 (synth)


Chang, L.C. *et al.*, *J. Nat. Prod.*, 2002, **65**, 775-778 (Isobatzellines C,E)

Dijoux, M.-G. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 6035-6044 (Isobatzellines C,E)

Isobellendine

I-194

6,7,8,9-Tetrahydro-2,10-dimethylcyclohepta[b]pyran-5,8-imin-4(5H)-one, 9CI



Absolute Configuration

$C_{12}H_{15}NO_2$ 205.256

(+)-form [72362-45-9]

Alkaloid from upper leaves, flowers and flowering stems of *Bellendena montana* (Proteaceae). Fine needles (Et₂O). Mp 114-116°. $[\alpha]_D^{19}$ +143 (CHCl₃).

5 ζ ,11 ζ -Dihydro: 5,11-Dihydroisobellendine

[72362-46-0]

[72401-73-1]

 $C_{12}H_{17}NO_2$ 207.272

Alkaloid from *Bellendena montana* (Proteaceae). Oil. Mp 172-174° (as picrate). $[\alpha]_D^{19}$ -53 (CHCl₃).

(±)-form

Mp 88-90° subl.

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 1827-1840 (isol, uv, pmr, ms, struct, synth, deriv)


Lounasmaa, M. *et al.*, *Tet. Lett.*, 1981, 5179-5180 (synth)

Majewski, M. *et al.*, *J.O.C.*, 1995, **60**, 5825-5830 (synth, abs config)

Isobengamide E

I-195

[118477-05-7]



Absolute configuration

$C_{17}H_{30}N_2O_6$ 358.434

Metab. of an undescribed Fijian marine sponge (Jaspidae). Anthelmintic. Nematocidal. Oil. $[\alpha]_D^{20}$ +17.1 (c, 0.052 in MeOH).


Adamczeski, M. *et al.*, *J.A.C.S.*, 1989, **111**, 647 (isol, ir, pmr, cmr, ms, struct)

Adamczeski, M. *et al.*, *J.O.C.*, 1990, **55**, 240 (abs config)

Isoboldine

I-196

1,9-Dihydroxy-2,10-dimethoxyaporphine. N-Methyllaurelliptine



(R)-form

 $C_{19}H_{21}NO_4$ 327.379**(R)-form** β -N-Oxide: **Isoboldine N-oxide** $C_{19}H_{21}NO_5$ 343.379

Alkaloid from *Cryptocarya chinensis*. Brown powder. Mp 177-179°. $[\alpha]_D$ -90.3 (c, 0.14 in MeOH). λ_{max} 219 ($\log \epsilon$ 3.08); 271 (sh) ($\log \epsilon$ 2.68); 279 ($\log \epsilon$ 2.7); 304 ($\log \epsilon$ 2.7) (MeOH).

(S)-form [3019-51-0]

Alkaloid from a wide variety of genera in the Annonaceae (*Annona*, *Enantia*, *Uvaria*, *Monodora*, *Schefferomitra*, *Xylopia*, *Desmos*, *Guatteria*, *Polyalthia*), Berberidaceae (*Berberis*, *Mahonia*), Lauraceae (*Beilschmiedia*, *Cassytha*, *Litsea*, *Neolitsea*, *Nectandra*, *Phoebe*, *Ocotea*, *Sassafras*, *Machilus*), Fabaceae (*Erythrina*), Menispermaceae (*Cocculus*, *Stephania*), Monimiaceae (*Peumus*), Papaveraceae (*Corydalis*, *Fumaria*, *Glaucium*), Symlocaceae (*Symplocos*), Ranunculaceae (*Delphinium*, *Aconitum*, *Thalictrum*), Rhamnaceae (*Zizyphus*), Hernandiaceae (*Hernandia*), Nandinaceae (*Nandina*) and Monimiaceae (*Laurelia*). Antifeedant, insecticide. Mp 178-180° dec. $[\alpha]_D^{13}$ +54 (c, 0.20 in EtOH). λ_{max} 219 (ϵ 38020); 280 (ϵ 14454); 304 (ϵ 16220) (MeOH) (Berdy).

N-Me: Laurifoline⁺

[7224-61-5]

 $C_{20}H_{24}NO_4^{\oplus}$ 342.414

Quaternary alkaloid from *Cocculus laurifolius* and *Zanthoxylum elephantiasis* (Menispermaceae, Rutaceae). Mp 253° dec. (as chloride). $[\alpha]_D$ +14 (c, 6.22 in MeOH).

N-De-Me: 1,9-Dihydroxy-2,10-dimethoxy-ynoraporphine. Laurelliptine. Norisoboldine

[23599-69-1]

 $C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Beilschmiedia elliptica*, *Beilschmiedia podagrica*, *Cassytha pubescens*, *Monodora tenuifolia*, *Litsea xylanica* and some other *Litsea* spp., *Zizyphus jujuba*, *Nectandra rigida* and (possibly) *Monanthes cauliflora* (Lauraceae, Annonaceae, Rhamnaceae). Shows antifungal activity. Cryst. (C_6H_6 /EtOH). Mp 190-192°. $[\alpha]_D^{21}$ +47.3 (c, 1.0 in EtOH).

N-De-Me, hydrochloride:

Cryst. (MeOH/Et₂O or EtOH/Et₂O). Mp 270-280°. $[\alpha]_D^{27}$ +22.2 (c, 0.5 in EtOH).

N-De-Me, picrate:

Deep orange needles + ½ H₂O (EtOH aq.). Mp 194-195° dec.

N-De-Me, N-Ac:

Prisms (EtOH). Mp 298-300° dec. $[\alpha]_D$ +340 (c, 0.19 in CHCl₃/MeOH, 1:1).

N-De-Me, tri-Ac:

Needles (C_6H_6 /petrol). Mp 187-190°. $[\alpha]_D^{17}$ +76 (c, 1.4 in EtOH).

N-De-Me, N-hexadecanoyl: Laurelliptin-1-hexadecanone

[862587-53-9]

 $C_{34}H_{49}NO_5$ 551.765

Alkaloid from the stems of *Cocculus orbiculatus*. Obt. as a mixt. of *E/Z*-enolates with *N*-octadecanoyl homologue, to which data refers. Brown powder. $[\alpha]_D^{25} +154.4$ (c. 1 in MeOH). λ_{\max} 221 ; 281 ; 304 (MeOH).

N-De-Me, N-octadecanoyl: Laurelliptin-1-octadecanone

[862587-54-0]

$C_{36}H_{53}NO_5$ 579.818

Alkaloid from the stems of *Cocculus orbiculatus*. Obt. as a mixt. of *E/Z*-enolates with *N*-hexadecanoyl analogue above.

(±)-**form** [5164-93-2]

Alkaloid from *Glaucium flavum* (Papaveraceae). Mp 130-131° (123-135°).

Picrate: Mp 163-165°.

Tomita, M. *et al.*, *Pharm. Bull.*, 1953, 1, 1 (*Laurifoline*)

Clezy, P.S. *et al.*, *Aust. J. Chem.*, 1966, 19, 135 (*isol, uv, pmr, Isoboldine, Laurelliptine*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, 19, 2331; 1969, 22, 1277 (*isol, uv, pmr, Isoboldine, Laurelliptine*)

Chan, W.W.C. *et al.*, *J.C.S.(C)*, 1966, 753 (*synth, ir, uv, pmr*)

Albonico, S.M. *et al.*, *J.C.S.(C)*, 1966, 1340 (*ord, abs config, Laurifoline*)

Jackson, A.H. *et al.*, *J.C.S.(C)*, 1966, 2061 (*synth, uv, pmr*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, 33, 323 (*racemate, isol*)

Wada, K. *et al.*, *J. Agric. Food Chem.*, 1968, 16, 471

Dominguez, X.A. *et al.*, *Phytochemistry*, 1974, 13, 680 (*isol, uv, pmr, ms, Laurifoline*)

Brown, G.M. *et al.*, *Acta Cryst. B*, 1977, 33, 2051 (*cryst struct, abs config*)

Kametani, T. *et al.*, *J.A.C.S.*, 1977, 99, 3805 (*synth*)

Roques, R. *et al.*, *Acta Cryst. B*, 1978, 34, 837 (*cryst struct, Laurelliptine*)

Ricca, G.S. *et al.*, *Gazz. Chim. Ital.*, 1979, 109, 1 (*cmr, pmr*)

Marsaioli, A.J. *et al.*, *Phytochemistry*, 1979, 18, 165 (*cmr*)

Paulo, M. de Q. *et al.*, *J. Ethnopharmacol.*, 1992, 36, 39-41 (*Laurelliptine, activity*)


Lin, F.-W. *et al.*, *Chem. Pharm. Bull.*, 2001, 49, 1292-1294 (*N-oxide*)

Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2005, 68, 1056-1060 (*Laurelliptin-1-hexadecanone, Laurelliptin-1-octadecanone*)

Isononafousine

I-197

[76202-19-2]



$C_{35}H_{40}N_4O_3$ 564.726


Alkaloid from the leaves of *Bonafousia tetrastrachya* (preferred genus name *Tabernaemontana*) (Apocynaceae).

Amorph. $[\alpha]_D +19$ (c. 0.36 in $CHCl_3$).

Damak, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1980, 490 (*isol, uv, ir, pmr, cmr, ms, struct*)

Isoborreverine

[69461-07-0]



$C_{32}H_{40}N_4$ 480.695

Alkaloid from *Borreria verticillata* and *Flindersia fournieri* (Rubiaceae, Flindersiaceae). Noncryst. Opt. inactive. λ_{\max} 226 (ε 19000); 287 (ε 12500); 294 (ε 10000) (EtOH) (Berdy).

N^d-Me: 4-Methylisoborreverine

[73706-29-3]

$C_{33}H_{42}N_4$ 494.722

Isol. from *Flindersia fournieri* (Flindersiaceae). Cryst. (Me₂CO). Mp 209°. $[\alpha]_D^{20}$ 0 (CHCl₃).

N^d,N^d-Di-Me: 4,4'-Dimethylisoborreverine

[73706-30-6]

$C_{34}H_{44}N_4$ 508.748

Isol. from *Flindersia fournieri* (Flindersiaceae). Amorph. $[\alpha]_D^0$ (CHCl₃).

14',15'-Dihydro, 15'-hydroxy: 15'-Hydroxy-14',15'-dihydroisoborreverine

[73706-31-7]

$C_{32}H_{42}N_4O$ 498.71


Isol. from *Flindersia fournieri* (Flindersiaceae). Amorph. $[\alpha]_D^{20}$ 0 (CHCl₃).

Tillequin, F. *et al.*, *J. Nat. Prod.*, 1979, 42, 92 (*uv, ir, pmr, ms, struct, synth*)

Tillequin, F. *et al.*, *Phytochemistry*, 1979, 18, 1559; 2066 (*derivs*)

Isobrafouedine

[104021-42-3]



Relative Configuration

$C_{18}H_{22}N_2O_2$ 298.384


Alkaloid from the stem bark of *Strychnos dinklagei* (Loganiaceae). Amorph. $[\alpha]_D^{20} -13$ (c. 0.05 in MeOH). λ_{\max} 227 (log ε 4.18); 275 (sh) (log ε 3.5); 285 (log ε 3.52); 292 (log ε 3.51) (EtOH).

Michel, S. *et al.*, *J. Nat. Prod.*, 1986, 49, 452-455 (*uv, ir, pmr, ms, struct*)

Isobutyropyrrrothine†

I-200

6-Isobutyrylamino-4,5-dihydro-4-methyl-1,2-dithiolo[4,3-b]pyrrole



NHCOCH(CH₃)₂

Me

$C_{10}H_{14}N_2OS_2$ 242.365


Isol. from *Streptomyces pimprina*. Active against gram-positive bacteria. Orange-red plates. Mp 228-229°. Related to Holothin and Thiolutin.

Bhate, D.S. *et al.*, *Experientia*, 1960, 16, 504; *Hind. Antibiot. Bull.*, 1964, 6, 153 (*isol, props*)

Isocephalotaxine

I-201

[73208-54-5]



$C_{18}H_{21}NO_4$ 315.368

Alkaloid from the fruit of *Cephalotaxus fortunei*. Amorph. solid. $[\alpha]_D^{21} -47$ (c. 0.5 in CHCl₃).

Ketone: Isocephalotaxinone

[50908-91-3]

[56144-23-1]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Cephalotaxus fortunei*.

Ketone, O-de-Me: Demethylcephalotaxinone. Desmethylcephalotaxinone

[51020-45-2]

$C_{17}H_{17}NO_4$ 299.326

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus fortunei*, *Cephalotaxus hainanensis* and *Cephalotaxus sinensis* (Cephalotaxaceae). Pale-yellow solid (MeOH). Mp 102-107°. $[\alpha]_D +40$ (semisynthetic). $[\alpha]_D^{26} +2.3$ (c. 0.52 in MeOH). The nat. alkaloid is nearly racemic.

Ketone, O-de-Me, stereoisomer:

$C_{17}H_{17}NO_4$ 299.326

Alkaloid from the heartwood of *Cephalotaxus wilsonii*. Brownish powder. $[\alpha]_D^{28} +52$ (c. 0.08 in MeOH). Incorrectly indexed by CAS. λ_{\max} 212 (log ε 4.64); 288 (log ε 4.22) (MeOH).

Powell, R.G. *et al.*, *Phytochemistry*, 1973, 12, 2987-2991 (*Demethylcephalotaxinone*)

Weinreb, S.M. *et al.*, *J.A.C.S.*, 1975, 97, 2503-2516 (*Demethylcephalotaxinone, synth, uv, ir, pmr*)

Lin, W. *et al.*, *Yaoxue Xuebao*, 1985, 20, 283-287; *CA*, 103, 85049r (*Isocephalotaxinone*)

Bocar, M. *et al.*, *J. Nat. Prod.*, 2003, 66, 152-154 (*Isocephalotaxine*)

Wang, L.W. *et al.*, *J. Nat. Prod.*, 2004, 67, 1182-1185 (*Demethylcephalotaxinone stereoisomer*)

Isochaksine

I-202

[11029-95-1]

$C_{11}H_{20}N_3O_2^{\oplus}$ 226.298

Struct. unknown. Alkaloid from the seeds, roots and leaves of *Cassia absus* (Fabaceae).

Chloride:


$C_{11}H_{20}ClN_3O_2$ 261.751

Mp 250-252° (248°).

Picrate: Mp 182-184°.

Siddiqui, S. *et al.*, *Proc. - Indian Acad. Sci., Sect. A*, 1935, **2**, 421-425; *CA*, **30**, 1799 (*isol*)
 Cheema, M. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1965, **158**, 307-313 (*pharmacol*)
 Krishna Rao, R.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 299-300 (*isol*)

Isochelidonine I-203
 [142741-30-8]



Absolute Configuration

C₂₀H₁₉NO₅ 353.374
 Alkaloid from aerial parts of *Chelidonium majus* (Papaveraceae). Cryst. (EtOH). Mp 124-125°. [α]_D²⁰ +72 (c, 0.23 in EtOH).


De Rosa, S. *et al.*, *Phytochemistry*, 1992, **31**, 1085 (*isol, uv, ir, pmr, struct*)

Isochondourarine I-204

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from Curare. Cryst. (MeOH/Me₂CO) (as dihydrochloride). Mp 278° dec. (dihydrochloride). [α]_D²⁰ -150 (H₂O) (dihydrochloride).

Bodendorf, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1954, **287**, 555; *CA*, **52**, 10119h (*isol*)

Isochromophilone VI I-205
 [167173-91-3]



C₂₃H₂₈ClNO₅ 433.931
 Prod. by *Penicillium multicolor* FO-3216. Acyl-CoA: cholesterol transferase (ACAT) inhibitor. Red powder. Mp 122-125°. [α]_D²⁰ +2170 (c, 0.1 in MeOH). Related to Sclerotiorin. λ_{max} 235 (ε 15600); 370 (ε 23400); 485 (ε 3900) (MeOH) (Berdy).


N-De(hydroxyethyl), N-(3-carboxypropyl): **Isochromophilone IX**

C₂₅H₃₀ClNO₆ 475.968

Prod. by a *Penicillium* sp. (MIN-AP9902). [α]_D²⁰ +430 (c, 0.1 in MeOH).
 Arai, N. *et al.*, *J. Antibiot.*, 1995, **48**, 696 (*isol, uv, ir, pmr, cmr*)
 Michael, A.P. *et al.*, *Aust. J. Chem.*, 2003, **56**, 13-15 (*Isochromophilone IX*)

Isochrysohermidin I-206
 [104006-84-0]

[128726-53-4 (*R*,S**)-form]




C₁₆H₂₀N₂O₁₀ 400.341
 Alkaloid from the whole plant of *Mercurialis leiocarpa*. DNA cross-linking

agent. Needles (MeOH). Mp 266-268°. Racemic.

Masui, Y. *et al.*, *Phytochemistry*, 1986, **25**, 1470-1471 (*isol, pmr, cmr, ms, cryst struct*)
 Abe, K. *et al.*, *Phytochemistry*, 1989, **28**, 960 (*synth*)
 Wasserman, H.H. *et al.*, *Tetrahedron*, 1997, **53**, 8731-8738 (*synth*)
 Nicolaou, K.C. *et al.*, *Classics in Total Synthesis II: More Targets, Strategies, Methods*, 2003, 15-30 (*rev, synth, bibl*)

Isococculidine I-207
 1,2-Didehydro-3,15-dimethoxyerythrinan, 9CI. O-Methylisococculine



C₁₈H₂₃NO₂ 285.385
 Rel. config. only detd.

(+)-form [60229-91-6]
 Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Weak cholinergic receptor blocking agent. Cryst. (C₆H₆/hexane). Mp 95-96°. [α]_D²⁰ +124 (c, 1.2 in MeOH).

►KF1950000

Methiodide:

Cryst. (Et₂O/MeOH). Mp 198-199°.

O¹⁵-De-Me: **Isococculine**

[67657-98-1]

C₁₇H₂₁NO₂ 271.358

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Mp 182-184°. [α]_D²⁰ +164 (c, 1.0 in CHCl₃).

11ξ-Methoxy, O¹⁵-de-Me: **Cocculimine**

[77795-08-5]

C₁₈H₂₃NO₃ 301.385

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Amorph. powder. λ_{max} 233 ; 282 (EtOH). λ_{max} 305 (EtOH/NaOH).


(±)-form [78463-72-6]

Synthetic. Oil.

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1976, **15**, 739-741 (*isol, uv, ir, pmr, ms, struct*)
 Bhakuni, D.S. *et al.*, *Tetrahedron*, 1980, **36**, 2153-2156; 3107-3114 (*Isococculine, Cocculimine, isol, pmr, ms, synth, biosynth*)
 Ju-ichi, M. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 396-401 (*synth, ir, pmr, ms*)
 Madhusudanan, K.P. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 907-908 (*ms*)
 Prakash, O. *et al.*, *J. Nat. Prod.*, 1988, **51**, 603-605 (*pmr, cmr*)

Isocorydine I-208

11-Hydroxy-1,2,10-trimethoxyaporphine. Luteanine†



(*R*)-form

C₂₀H₂₃NO₄ 341.406

(*R*)-form [55056-91-2]

Alkaloid from leaves of *Annona senegalensis* (Annonaceae). Amorph. powder. Mp 78-80°. [α]_D²⁵ -190 (c, 0.02 in CHCl₃).

α-N-Oxide: **Isocorydine N-oxide**

[25405-80-5]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Milusa velutina*, *Glaucium* sp. and *Berberis integerrima* (Berberidaceae). Amorph. solid or gum. Mp 228-229° (as hydrochloride). The *M. velutina* isolate (confusingly called (-)-Isocorydine N-oxide by the authors), to which the opt. rotn. refers, is the only isolate with determined stereochem. It appears to have different stereochem from the *B. integerrima* isolate (no opt. rotn. reported) to which the Mp refers. λ_{max} 221 (log ε 4.31); 268 (log ε 3.87); 306 (log ε 3.46) (MeOH).

N-De-Me: [194923-28-9]

Alkaloid from *Annona cherimola* (cherimoya) (Annonaceae). Brown powder. Mp 200-203°. [α]_D²⁴ -46 (c, 0.5 in CHCl₃). λ_{max} 223 ; 265 ; 309 (no solvent reported).

N-De-Me, N-(methoxycarbonyl): N-Methoxycarbonylnorisocorydine. **Romucosine H**

[344928-15-0]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Annona cherimola* (cherimoya). Amorph. brown powder. Mp 230-233°. [α]_D²⁴ -43 (c, 0.01 in CHCl₃). λ_{max} 222 (log ε 4.39); 270 (log ε 4.19); 310 (log ε 3.8) (EtOH).

(*S*)-form [475-67-2]

Alkaloid from a wide variety of genera in the Annonaceae (*Annona*, *Enantia*, *Asimina*, *Artabotrys*), Berberidaceae (*Berberis*, *Mahonia*), Lauraceae (*Cryptocarya*, *Dehaasia*, *Beilschmiedia*, *Litsea*, *Phoebe*, *Ocotea*), Menispermaceae (*Stephania*), Monimiaceae (*Peumus*), Papaveraceae (*Corydalis*, *Dicentra*, *Glaucium*, *Papaver*, *Dicranostigma*), Ranunculaceae (*Thalictrum*), Rhamnaceae (*Phyllaea*), Hernandiaceae (*Hernandia*), Monimiaceae (*Atherosperma*, *Doryphora*) and Pteridophyllaceae (*Pteridophyllum*). Adrenocytic, sedative and cholinergic agent. Shows virtually no antitussive activity. Poorly sol. hexane. Mp 185°. [α]_D²³ +215 (c, 1 in CHCl₃). λ_{max} 210 ; 267 ; 302 (MeOH) (Berdy).

►LD₅₀ (rat, ipr) 10.9 mg/kg. CE1057950

Hydrochloride: [13552-72-2]

Mp 215-218° dec. [α]_D²⁰ +170 (c, 0.5 in CHCl₃).

N-Me: **Menisperine**. N-Methylisocorydine. *Chakranine*

[25342-82-9]

C₂₁H₂₆NO₄[⊕] 356.441

Quaternary alkaloid from *Bragantia wallichii* (preferred genus name *Thottea*), *Cryptocarya angulata*, *Cryptocarya triplinervis*, *Fagara coco*, *Legnephora moorei* and many other spp. (Aristo-

lochiaceae, Lauraceae, Rutaceae, Menispermaceae). Prisms (EtOH) (as chloride). Mp 217-218° dec. (slow heating) (chloride) Mp 235° (rapid heating) (chloride). $[\alpha]_D^{15} +146.3$ (c, 0.17 in CHCl_3).

N-De-Me: Norisocorydine. Sanjoinine Ib [475-70-7]

$\text{C}_{19}\text{H}_{21}\text{NO}_4$ 327.379

Alkaloid from a variety of genera in the Annonaceae (*Annona*, *Xylopia*), Hernandiaceae (*Hernandia*), Monimiaceae (*Nemuaron*, *Peumus*), Papaveraceae (*Glaucium*, *Corydalis*), Lauraceae (*Cryptocarya*), and Rhamnaceae (*Zizyphus*). Mp 203-205° (as hydrobromide). $[\alpha]_D^{20} +158.5$ (c, 0.1 in EtOH) (hydrobromide). Free base unstable.

4R-Hydroxy: Rhopalotine. Crabbine [151271-85-1]

$\text{C}_{20}\text{H}_{23}\text{NO}_5$ 357.405

Alkaloid from *Corydalis lutea* and *Papaver rhopalothece* (Papaveraceae). Cryst. (MeOH). Mp 191°. $[\alpha]_D +162$ (c, 0.2 in MeOH).


(±)-form [36284-37-4]

Mp 151-152°.

- Gadamer, J. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1911, **249**, 641 (*isol*)
 Comin, J. et al., *J.O.C.*, 1954, **19**, 1774 (*Isocorydine, Menisperine, isol, uv, struct*)
 Ruediger, A. et al., *Helv. Chim. Acta*, 1959, **42**, 754 (*Norisocorydine*)
 Katritzky, A.R. et al., *J.C.S.*, 1960, 1950 (*Menisperine, ir, pmr, struct*)
 Kuck, A.M. et al., *J.O.C.*, 1961, **26**, 5253 (*synth, uv*)
 Albonico, S.M. et al., *J.C.S. (C)*, 1966, 1340 (*ord, derivs*)
 Jackson, A.H. et al., *J.C.S. (C)*, 1966, 2181 (*ms*)
 Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1277 (*isol, pmr*)
 Groebel, A. et al., *Planta Med.*, 1970, **18**, 66 (*Menisperine, isol, uv, pmr, ms*)
 Bhakuni, D.S. et al., *Phytochemistry*, 1972, **11**, 1819 (*Norisocorydine, isol, uv, pmr, ms*)
 Karimov, A. et al., *Khim. Prir. Soedin.*, 1978, **14**, 419; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 360 (*Isocorydine N-oxide*)
 Marsaioli, A.J. et al., *Phytochemistry*, 1979, **18**, 165 (*cmr, derivs*)
 Ringdahl, B. et al., *J. Nat. Prod.*, 1981, **44**, 80 (*cd*)
 Toure, S. et al., *Acta Cryst. C*, 1985, **41**, 1827 (*cryst struct*)
 Sariyar, G. et al., *Planta Med.*, 1990, **56**, 232 (*Rhopalotine*)
 Ribár, B. et al., *Acta Cryst. C*, 1992, **48**, 945 (*cryst struct*)
 Yang, M.H. et al., *Phytochemistry*, 1993, **33**, 943 (*Crabbine*)
 You, M. et al., *J. Nat. Prod.*, 1995, **58**, 598 (*isol, (R)-Isocorydine*)
 Lee, S.-S. et al., *J. Nat. Prod.*, 1996, **59**, 55 (*cd*)
 Chen, C.-Y. et al., *J. Chin. Chem. Soc. (Taipei)*, 1997, **44**, 313 (*R-Norisocorydine*)
 Rasoanaivo, P. et al., *Planta Med.*, 1998, **64**, 58 (*Norisocorydine, cmr*)
 Marek, R. et al., *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)
 Hasan, C.M. et al., *Nat. Prod. Lett.*, 2000, **14**, 393-397 (*α-N-oxide*)
 Chen, C.-Y. et al., *Phytochemistry*, 2001, **56**, 753-757 (*Romucosine H*)

Isocorymine

[16843-68-8]



$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Hunteria umbellata* seeds and from *Hunteria zeylanica* (Apocynaceae). Cryst. (propanol). Mp 183-185°. $[\alpha]_D -243$.

Ac:

Cryst. (propanol). Mp 160°.

N⁴-De-Me: Norisocorymine

[69734-94-7]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Isol. from the leaves of *Hunteria congolana* and leaves and stem bark of *Hunteria zeylanica* (Apocynaceae). Amorph. solid + 1/2 H_2O . $[\alpha]_D -10$ (c, 1.0 in CHCl_3).

Deoxy, 21-oxo (lactone): Erinine

[4825-07-4]

$\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_4$ 380.443

Alkaloid from the leaves of *Hunteria umbellata* and *Hunteria eburnea* (Apocynaceae). Cryst. ($\text{CHCl}_3/\text{MeOH}$). Mp 267-269°. $[\alpha]_D -186$ (CHCl_3).

Deoxy, 21-oxo, picrate:

Cryst. (MeOH). Mp 195-197° dec.

Deoxy, 21-oxo, 19,20ξ-dihydro: Erinicine

[4684-41-7]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Hunteria eburnea* leaves and *Hunteria umbellata* (Apocynaceae). Mp 210-212°.

Bycroft, B.W. et al., *Helv. Chim. Acta*, 1965, **48**, 1598 (*Erinine, Erinicine*)

Bevan, C.W.L. et al., *Tetrahedron*, 1967, **23**, 3809-3821 (*ms, pmr*)

Morfaux, A.M. et al., *Ann. Pharm. Fr.*, 1969, **27**, 679 (*Erinine, Erinicine, isol*)

Heatley, F. et al., *J.C.S. Perkin 2*, 1981, 725-729 (*struct*)

Arambewala, L.S.R. et al., *Phytochemistry*, 1981, **20**, 349-350 (*Isocorymine, isol*)

Vercauteren, J. et al., *Bull. Soc. Chim. Fr.*, Part II, 1982, 291-296 (*Norisocorymine*)


Isocorypalmine

I-210

5,8,13,13a-Tetrahydro-3,9,10-trimethoxy-6H-dibenzo[a,g]quinolizin-2-ol, 9CI.

3,9,10-Trimethoxyberbin-2-ol, 8CI. 2-Hydroxy-3,9,10-trimethoxytetrahydroprotoberberine. Tetrahydrocolumbamine.

Casualtine



(R)-form

I-209

$\text{C}_{20}\text{H}_{23}\text{NO}_4$ 341.406

(R)-form [53447-14-6]

Alkaloid from *Corydalis cava*, in partially racemised form (Papaveraceae). Cryst. (MeOH). Mp 239-241° (220-222°). $[\alpha]_D +303$ (CHCl_3).

(S)-form [483-34-1]

Alkaloid from the leaves of *Bocconia frutescens*, *Corydalis lutea*, *Corydalis ophiocarpa*, other *Corydalis* spp., *Dicranostigma* spp., *Glaucium fimbriigerum*, *Hydrastis canadensis*, *Thalictrum* spp., *Pachypodanthium confine* and *Tinomisium petiolare* (Papaveraceae, Ranunculaceae, Annonaceae). Mp 241-242°.

▶ NQ7685000

α-N-Oxide: Isocorypalmine α-N-oxide

$\text{C}_{20}\text{H}_{23}\text{NO}_5$ 357.405

Alkaloid from *Corydalis tashiroi*. Needles ($\text{CHCl}_3/\text{MeOH}$). Mp 247-249°. $[\alpha]_D^{25} -24.7$ (c, 0.25 in CHCl_3). λ_{max} 209 (log ε 4.55); 225 (sh) (log ε 4.19); 283 (log ε 3.76) (EtOH).

β-N-Oxide: Isocorypalmine β-N-oxide

$\text{C}_{20}\text{H}_{23}\text{NO}_5$ 357.405

Alkaloid from *Corydalis tashiroi*. Needles (CHCl_3). Mp 224-226°. $[\alpha]_D^{25} -38.6$ (c, 0.23 in CHCl_3). λ_{max} 210 (log ε 4.53); 225 (sh) (log ε 4.18); 284 (log ε 3.73) (EtOH).

β-N-Me: β-N-Methylisocorypalminium.

N-Methyltetrahydrocolumbaminium

[92758-34-4]

[95462-85-4, 96442-66-9]

$\text{C}_{21}\text{H}_{26}\text{NO}_4^{\oplus}$ 356.441

Quaternary alkaloid from *Glaucium squamigerum*, *Cymbopetalum brasiliense* and *Tinospora heinannensis*. Prisms (MeOH) (as iodide). Mp 175-176° (iodide). $[\alpha]_D^{24} -127$ (c, 0.13 in MeOH). CAS no. refers to iodide.

Me ether: see Tetrahydrocolumbamine, T-212

8-Oxo: 8-Oxoisocorypalmine

[142808-34-2]

$\text{C}_{20}\text{H}_{21}\text{NO}_5$ 355.39

Alkaloid from stems of *Coscinium fenestratum* (Menispermaceae). Gum. $[\alpha]_D -222$ (c, 0.072 in CHCl_3).

(±)-form [6487-33-8]

Alkaloid from *Corydalis cava* (Papaveraceae). Prisms ($\text{CHCl}_3/\text{MeOH}$). Mp 221-222° (215-216°).

Corrodi, H. et al., *Helv. Chim. Acta*, 1956, **39**, 889

Tomita, M. et al., *Yakugaku Zasshi*, 1967, **87**, 881 (*isol*)

Naruto, S. et al., *Phytochemistry*, 1972, **11**, 2462 (*isol*)

Slavik, J. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 3206; 1979, **44**, 2261; 1984, **49**, 1318 (*isol, N-Methylisocorypalminium*)

Battersby, A.R. et al., *J.C.S. Perkin 1*, 1975, 1147 (*biosynth*)

Casnow, J. et al., *Anal. Biochem.*, 1976, **74**, 343 (*glc, ms*)


Ribár, B. et al., *Acta Cryst. C*, 1992, **48**, 944 (*cryst struct*)

Pinho, P.M.M. et al., *Phytochemistry*, 1992, **31**, 1403 (*8-Oxoisocorypalmine*)

Guo, Y. et al., *Chem. Pharm. Bull.*, 1999, **47**, 287-289 (*N-Methylisocorypalminium*)

Chen, J.-J. *et al.*, *Planta Med.*, 1999, **65**, 643-647; 2001, **67**, 423-427 (*N-oxides*)

Isocryprochine I-211




$C_{19}H_{25}NO_3$ 315.411
Alkaloid from the wood of *Cryptocarya chinensis*. Needles (Me_2CO). Mp 186-187°. $[\alpha]_D^{25}$ -22.2 (c, 0.03 in MeOH). λ_{max} 227 (log ϵ 3.83); 285 (log ϵ 3.27) (MeOH).

Wu, T.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1404-1407

7-Isocyanato-2,10-bisaboladiene I-212

7-Isocyanato-7,8-dihydro- α -bisabolene



$C_{16}H_{25}NO$ 247.38


(6R,7R)-form [105281-35-4]

Constit. of a *Ciocalypta* sp. Oil. $[\alpha]_D^{25}$ -24.3 (c, 0.094 in hexane). λ_{max} 205 (ϵ 5000); 248 (ϵ 500) (hexane) (Derep).

Gulavita, N.K. *et al.*, *J.O.C.*, 1986, **51**, 5136 (*isol, cryst struct*)

3-Isocyano-7,9-bisaboladiene I-213

4-(1,5-Dimethyl-1,3-hexadienyl)-1-isocyano-1-methylcyclohexane, 9CI. 3-Isocyanotheonellin. *Theonelline isocyanide* [105281-40-1]



$C_{16}H_{25}N$ 231.38

Constit. of *Phyllidia pustulosa* and *Halichondria* cf. *lendenfeldi*. Oil. λ_{max} 203 (ϵ 6400); 232 (sh) (ϵ 1100); 239 (ϵ 11600) (MeOH) (Derep).

Isothiocyante: *Theonelline isothiocyante*

[94663-94-2]

$C_{16}H_{25}NS$ 263.446

Constit. of the sponge *Theonella* cf. *swinhoi*. Shows antibacterial props. Oil. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Has -NCS relacing -NC. λ_{max} 238 (ϵ 30200) (MeOH) (Derep). λ_{max} 230 (MeOH) (Berdy).

Formamide: *Theonelline formamide*. 3-Formamidotheonellin

[94663-93-1]

$C_{16}H_{27}NO$ 249.395

Constit. of the sponge *Theonella* cf. *swinhoi* and *Axinyssa* sp. Oil. Sol. MeOH, $CHCl_3$, Me_2CO ; poorly sol. H_2O . Has -NHCHO replacing -NC. λ_{max} 238 (ϵ 24000) (MeOH) (Derep).

Formamide, 11-ethoxy: 11-Ethoxy-3-formamidotheonellin

[937800-57-2]

$C_{18}H_{31}NO_2$ 293.448

Constit. of *Axinyssa* aff. *variabilis*. Oil. λ_{max} 238 (log ϵ 4.18) (MeOH).

Nakamura, H. *et al.*, *Tet. Lett.*, 1984, **25**, 5401-5404 (*isothiocyante, formamide*)

Gulavita, N.K. *et al.*, *J.O.C.*, 1986, **51**, 5136

Kassühlke, K.E. *et al.*, *J.O.C.*, 1991, **56**, 3747-3750 (*isol*)


Fusetani, N. *et al.*, *Tet. Lett.*, 1991, **32**, 7291

Kitano, Y. *et al.*, *J.C.S. Perkin 1*, 2002, 2251-2255 (*synth*)

Mao, S.-C. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 588-593 (11-Ethoxy-3-formamidotheonellin)

7-Isocyano-2,10-bisaboladiene I-214

7-Isocyano-7,8-dihydro- α -bisabolene



(6R,7R)-form

$C_{16}H_{25}N$ 231.38

λ_{max} 195 (ϵ 14500) (hexane) (Derep).

(6R,7R)-form [105281-36-5]

Constit. of a *Ciocalypta* sp. Oil. $[\alpha]_D^{22}$ -49.9 (c, 0.033 in hexane).

(6R,7S)-form [957138-58-8]

Constit. of *Acanthella cavernosa*. Oil. $[\alpha]_D$ +60.8 (c, 0.049 in $CHCl_3$).

Isothiocyante: 7-*Isothiocyante*-2,10-bisaboladiene

[105281-46-7]

$C_{16}H_{25}NS$ 263.446

Isol. from a *Halichondria* sp. and *Phyllidia pustulosa*. Oil. $[\alpha]_D^{20}$ +60.5 (c, 6.8 in $CHCl_3$). Has -NCS in place of -NC.

Sullivan, B.W. *et al.*, *J.O.C.*, 1986, **51**, 5134-5136 (*isothiocyante, isol, pmr, cmr*)


Gulavita, N.K. *et al.*, *J.O.C.*, 1986, **51**, 5136 (*isocyanide*)

Kassühlke, K.E. *et al.*, *J.O.C.*, 1991, **56**, 3747-3750 (*isothiocyante, isol*)

Jumaryatno, P. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1725-1730 (*Acanthella cavernosa* constit)

8-Isocyano-1(12)-cycloamphilectene I-215

[108695-82-5]



$C_{21}H_{31}N$ 297.483


Struct. revised in 1987. Constit. of *Adocia* spp. and a *Halichondria* sp. Gum. Poorly sol. hexane. $[\alpha]_D^{20}$ +39.6 (c, 0.6 in $CHCl_3$).

Kaslauskas, R. *et al.*, *Tet. Lett.*, 1980, 315 (*isol*)

Molinski, T.F. *et al.*, *J.O.C.*, 1987, **52**, 3334 (*isol, cryst struct*)

3-(2-Isocyanoethenyl)-1H-indole I-216

2-(3-Indolyl)vinyl isocyanide. *Indoleacrylonitrile*. B 371. *Antibiotic B 371*. SF 2636. *Antibiotic SF 2636*



$C_{11}H_8N_2$ 168.198

The struct. given in the 1990 CAS abstract is prob. incorrect.

(Z)-form [61168-06-7]

Isol. from an unidentified *Pseudomonas* sp. and from the marine *Streptomyces* sp. SF2636. Shows antibiotic activity against *Staphylococcus* and other bacteria and yeasts. Probable biosynth. precursor to various cyanobacterial indole alkaloids. Extremely unstable, storeable only in soln. for a short period. λ_{max} 273 (sh); 279; 288 (MeOH/HCl) (Derep). λ_{max} 275 (sh); 313 (MeOH) (Derep).


Evans, J.R. *et al.*, *J. Antibiot.*, 1976, **29**, 850-852 (*isol*)

Hoppe, I. *et al.*, *Annalen*, 1984, 600-607 (*synth, ir, pmr, ms*)

Hatsu, M. *et al.*, *CA*, 1990, **113**, 112056f (*isol*)

Stratmann, K. *et al.*, *J.A.C.S.*, 1994, **116**, 9935-9942 (*biochem*)

5-Isocyano-6-eudesmene I-217




$C_{16}H_{25}N$ 231.38

(4β,5β,10α)-form [786703-09-1]

Constit. of *Phyllidiella pustulosa*. Oil. $[\alpha]_D$ +23 (c, 2 in $CHCl_3$).

Manzo, E. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1701-1704 (*isol, pmr, cmr*)

6-Isocyano-4(15)-eudesmene I-218



(5α,6α,7α,10β)-form

$C_{16}H_{25}N$ 231.38

(5α,6α,7α,10β)-form [91466-62-5]

Isol. from *Axinella cannabina*. Oil. $[\alpha]_D$ +39.6 (c, 0.3 in CHCl_3).

Isothiocyante: [91466-63-6]

$\text{C}_{16}\text{H}_{25}\text{NS}$ 263.446

Isol. from *Axinella cannabina*. Oil. $[\alpha]_D$ +41 (c, 0.35 in CHCl_3).

Formamide: [91466-64-7]

$\text{C}_{16}\text{H}_{27}\text{NO}$ 249.395

Isol. from *Axinella cannabina*. Oil. $[\alpha]_D$ +48.5 (c, 0.3 in CHCl_3).

(5 α ,6 α ,7 β ,10 α)-form

Constit. of *Axinella cannabina* and *Acanthella acuta*. Cryst. or oil. Mp 78-79°. $[\alpha]_D$ +92.9 (c, 1.8 in CHCl_3). Incorr. config. assigned in CAS.

Isothiocyante: 6-Isothiocyanto-4(15)-eudesmene

$\text{C}_{16}\text{H}_{25}\text{NS}$ 263.446

Constit. of *Acanthella acuta* and *Axinella cannabina*. Mp 52-53°. $[\alpha]_D$ +88.4 (c, 1.0 in CHCl_3). Has -NCS replacing -NC. Incorr. config. assigned in CAS.

Formamide: 6-Formamido-4(15)-eudesmene

$\text{C}_{16}\text{H}_{27}\text{NO}$ 249.395

Constit. of *Axinella cannabina*. Oil. Has -NHCHO replacing -NC. Incorr. config. assigned in CAS.

(5 α ,6 α ,7 β ,10 β)-form

Halichonadin C

[847605-78-1]

Constit. of a *Halichondria* sp. Amorph. solid. $[\alpha]_D^{19}$ -130 (c, 1 in CHCl_3).

Isothiocyante: Acanthene B

$\text{C}_{16}\text{H}_{25}\text{NS}$ 263.446

Constit. of an *Acanthella* sp. Solid. $[\alpha]_D$ -34 (c, 0.18 in CHCl_3). Incorr. config. assigned in CAS.

Formamide: Acanthene C

$\text{C}_{16}\text{H}_{27}\text{NO}$ 249.395

Constit. of *Cadlina luteomarginata* which feed on *Acanthella* sp. Amorph. solid. Incorr. config. assigned in CAS.

Ciminiello, P. *et al.*, *J.O.C.*, 1984, **49**, 3949-3951 (5 α ,6 α ,7 α ,10 β -form *Axinella cannabina* constits)

Ciminiello, P. *et al.*, *J. Nat. Prod.*, 1987, **50**, 217-220 (5 α ,6 α ,7 β ,10 α -form *Axinella* and *Acanthella* constits)


Burgoyne, D.L. *et al.*, *Tetrahedron*, 1993, **49**, 4503-4510 (*Acanthene* B,C)

Ishiyama, H. *et al.*, *Tetrahedron*, 2005, **61**, 1101-1105 (*Halichonadin* C)

4-Isocyanomaaliane

I-219

[133109-45-2]



$\text{C}_{16}\text{H}_{25}\text{N}$ 231.38

Isol. from the mollusc *Cadlina luteomarginata*. $[\alpha]_D$ +36 (c, 0.2 in CHCl_3).


Thompson, J.E. *et al.*, *Tetrahedron*, 1982, **38**, 1865 (*isol*, *pmr*, *cmr*, *ms*)

Burgoyne, D.L. *et al.*, *Tetrahedron*, 1993, **49**, 4503-4510 (*isol*)

9-Isocyanoneopupekanane

I-220

[119323-93-2]



$\text{C}_{16}\text{H}_{25}\text{N}$ 231.38

Constit. of a *Ciocalypta* sp. Oil. $[\alpha]_D$ +33 (c, 1 in CHCl_3).

Karuso, P. *et al.*, *J.O.C.*, 1989, **54**, 2092-2095; 2095-2097 (*isol*, *pmr*, *cmr*, *biosynth*)

Ho, T.-L. *et al.*, *J.O.C.*, 1999, **64**, 8965-8967 (*synth*)

Krishna, A. *et al.*, *Tetrahedron*, 2005, **61**, 8855-8859 (*synth*)


3-Isocyano-3,7,11,15-tetra-

I-221

methyl-1,6,10,14-hexadecatetraene, 9CI

[55784-75-3]

[154614-35-4]



$\text{C}_{21}\text{H}_{33}\text{N}$ 299.498

(6E,10E)-(+)-form

Geranyllinaloyl isocyanide

[57766-63-9]

Isol. from *Halichondria* sp. Oil. $[\alpha]_D$ +15 (c, 2.8 in CCl_4).

Isothiocyante: Geranyllinaloyl isothiocyante

$\text{C}_{21}\text{H}_{33}\text{NS}$ 331.564

Present in a *Halichondria* sp. sponge. Has -NCS in place of -NC.

Formamide: Geranyllinaloyl formamide

$\text{C}_{21}\text{H}_{35}\text{NO}$ 317.514

Present in a *Halichondria* sp. sponge. Has -NHCHO replacing -NC.

Burreson, B.J. *et al.*, *Tetrahedron*, 1975, **31**,

2015-2018 (*isol*, *ir*, *pmr*, *cmr*)


Ichikawa, Y. *et al.*, *J.C.S. Perkin 1*, 1993, 2429-2432 (*synth*, *ir*, *pmr*, *cmr*)

Isocyclolabenzine

I-222

3,4,5,6,8,9,10,11,12,13-Decahydro-9-phenyl-2,13-methano-2H-2,6,10-benzotriazacyclopentadecine-1,7-dione, 9CI

[70503-72-9]



Absolute Configuration

$\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}_2$ 377.485

CAS numbering shown. Alkaloid from

Maytenus mossambicensis var. *mossambicensis* (Celastraceae).

N^{10} -Ac: Mp 255°. $[\alpha]_D^{25}$ +221 (c, 0.356 in CHCl_3).

13-Hydroxy: 13-Hydroxyisocyclolabenzine

[82497-77-6]

$\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}_3$ 393.485

Alkaloid from *Maytenus mossambicensis* var. *mossambicensis* (Celastraceae). Mp 135°. $[\alpha]_D^{24}$ +113.1 (c, 0.603 in CHCl_3).

13-Hydroxy, N^{10} -Ac: Mp 315-320°.

[104595-98-4 ((±)-form)]

Wagner, H. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 739 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Iida, H. *et al.*, *J.O.C.*, 1986, **51**, 4701 (*synth*, *ir*, *pmr*)


Schultz, K. *et al.*, *Tetrahedron*, 1996, **52**, 14189 (*synth*, *abs config*, *cryst struct*)

Linden, Y.L.A. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 579-591 (*13-Hydroxyisocyclolabenzine*, *synth*)

Isocycloneosamandaridine

I-223

[63653-59-8]



$\text{C}_{21}\text{H}_{31}\text{NO}_3$ 345.481

Probable struct. shown. The originally proposed struct. (referred to as Cyclo-neosamandaridine) was synthesised and found to be incorr. with consequent name change. Alkaloid from the skin of *Salamandra maculosa maculosa*. Mp 270-272° (synthetic) Mp 281-283° (natural).

Hydrochloride: Mp 250° dec.


Habermehl, G. *et al.*, *Chem. Ber.*, 1965, **98**, 3001-3005 (*isol*, *ir*, *ms*)

Oka, K. *et al.*, *J.O.C.*, 1978, **43**, 4408-4408 (*synth*, *struct*, *bibl*)

Isocynometrine

I-224

[79659-62-4]



(-)-form

$\text{C}_{16}\text{H}_{19}\text{N}_3\text{O}_2$ 285.345

Alkaloid from the trunk bark of *Cynometra lujae* (Fabaceae). Cryst. (Me_2CO /pentane). Mp 181°. $[\alpha]_D^{20}$ -66 (c, 1 in CHCl_3).

Benzoyl: Isocynodine

[85644-19-5]


$\text{C}_{23}\text{H}_{23}\text{N}_3\text{O}_3$ 389.453

Alkaloid from the root bark, trunk bark and leaves of *Cynometra lujae*

(Fabaceae). Cryst. (Me₂CO/hexane). Mp 167°. [α]_D²⁰ +139 (c, 1 in CHCl₃).

- Chiaroni, A. *et al.*, *J. Chem. Res., Synop.*, 1981, 182; *J. Chem. Res., Miniprint*, 1981, 2116-2145 (*cryst struct, abs config*)
Tchissambou, L. *et al.*, *Tetrahedron*, 1982, **38**, 2687-2695 (*isol, ir, pmr, cmr, struct, synth, deriv*)
Xie, X. *et al.*, *J.O.C.*, 2001, **66**, 6545-6550 (*synth*)
Xu, W. *et al.*, *J.O.C.*, 2006, **71**, 3854-3858 (*synth*)

6-Isodaucene 14-isonitrile I-225
[112767-02-9]




C₁₆H₂₅N 231.38
Metab. of marine sponge *Acanthella acuta*. Oil. [α]_D +44 (c, 1.0 in CHCl₃).

Isothiocyanate: 6-Isodaucene 14-isothiocyanate
[112757-35-4]
C₁₆H₂₅NS 263.446
Metab. of *Acanthella acuta*. Oil. [α]_D +36.2 (c, 0.2 in CHCl₂). Has -NCS replacing -NC.

Mayol, L. *et al.*, *Tetrahedron*, 1987, **43**, 5381

Isodihydrocadambine I-226
[55624-02-7]



C₂₇H₃₄N₂O₁₀ 546.573
Alkaloid from the heartwood of *Anthocephalus cadamba* (Rubiaceae). [α]_D²⁵ -66 (MeOH) (as penta-Ac).

3-Epimer: 3β-Isodihydrocadambine
[62014-69-1]

C₂₇H₃₄N₂O₁₀ 546.573
Alkaloid from the leaves of *Anthocephalus cadamba*, *Nauclea cadamba* and *Uncaria* spp. Exhibits hypotensive activity. Amorph. solid; needles (MeOH) (as penta-Ac). Mp 170-171° (penta-Ac). [α]_D -3.3 (c, 0.368 in MeOH).

- Brown, R.T. *et al.*, *Tet. Lett.*, 1974, 3335; 1976, 2723 (*uv, ir, pmr, ms, cd, struct, epimer*)
Endo, K. *et al.*, *Planta Med.*, 1983, **49**, 188 (*epimer*)
Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 232-233 (*3β-Isodihydrocadambine*)


C-Isodihydrotoxiferine I I-227
C₂₀H₂₃N₂[⊕] 291.415
Minimum formula. Struct. unknown.

Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 242° dec. (sintering from 200°) (as picrate). [α]_D²⁰ -566 (H₂O) (as chloride).

- Wieland, H. *et al.*, *Annalen*, 1941, **547**, 156-179; 1947, **558**, 144-156 (*isol, uv*)
Wieland, T. *et al.*, *Chem. Ber.*, 1962, **85**, 731

Isodomoic acid A I-228

2-Carboxy-4-(5-carboxy-1-methyl-1,4-hexadienyl)-3-pyrrolidineacetic acid, 9CI
[101899-44-9]



C₁₅H₂₁NO₆ 311.334
Isol. from the red alga *Chondria armata* and the diatom *Nitzschia navis-varingica*. Anthelmintic; exhibits insecticidal activity. Sol. MeOH. Mp 185-187° dec. [α]_D²⁵ -70 (c, 0.1 in H₂O).

1'E-Isomer: Isodomoic acid B

[101977-25-7]
C₁₅H₂₁NO₆ 311.334
Isol. from *Chondria armata* and *Nitzschia navis-varingica*. Anthelmintic; insecticide; vermifuge. Sol. H₂O, MeOH. Mp 182-183° dec. [α]_D²⁵ -8.1 (c, 0.14 in H₂O). λ_{max} 214 (ε 9850) (H₂O) (Berdy).


Δ1',1''-Isomer: Isodomoic acid C

[101899-45-0]
C₁₅H₂₁NO₆ 311.334
Isol. from *Chondria armata* and *Pseudo-nitzschia australis*. Insecticide. Sol. MeOH, H₂O. Mp 257-260° dec. [α]_D²⁵ -30 (c, 0.015 in MeOH). λ_{max} 213 (ε 5030) (MeOH) (Berdy).

- Maeda, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4892 (*isol, uv, ir, pmr, cmr, ms*)
Holland, P.T. *et al.*, *Chem. Res. Toxicol.*, 2005, **18**, 814-816 (*Isodomoic acid C, isol*)
Clayden, J. *et al.*, *J.A.C.S.*, 2005, **127**, 2412-2413 (*Isodomoic acid C, synth*)
Clayden, J. *et al.*, *Tetrahedron*, 2005, **61**, 5713-5724 (*rev*)
Kotaki, Y. *et al.*, *Toxicol.*, 2005, **46**, 946-953 (*isol*)

Isodomoic acid G I-229

2-Carboxy-4-(5-carboxy-1-methyl-2-hexenyldiene)-3-pyrrolidineacetic acid
[188346-81-8]



Absolute Configuration

C₁₅H₂₁NO₆ 311.334
Isol. from the red alga *Chondria armata*.


4Z-Isomer: Isodomoic acid H

[188346-82-9]
C₁₅H₂₁NO₆ 311.334
Isol. from *Chondria armata*.

- Zaman, L. *et al.*, *Toxicol.*, 1997, **35**, 205-212 (*isol, pmr*)
Ni, Y. *et al.*, *Org. Lett.*, 2003, **5**, 3771-3773 (*synth, abs config*)

Isoechinulin A I-230

3-[[2-(1,1-Dimethyl-2-propenyl)-5-(3-methyl-2-butenyl)-1H-indol-3-yl]methylene]-6-methyl-2,5-piperazinedione, 9CI
[60422-87-9]



C₂₄H₂₉N₃O₂ 391.512
Isol. from *Aspergillus ruber*. Mycotoxin.

3e,3h-Didehydro: Isoechinulin B

[60422-88-0]
C₂₄H₂₇N₃O₂ 389.496
Isol. from *Aspergillus ruber*. Mycotoxin.

3e,3h-Didehydro, 5b,5c-epoxide: Isoechinulin C

[60422-89-1]
C₂₄H₂₇N₃O₃ 405.496
Isol. from *Aspergillus ruber*.

5b,5c-Dihydro, 5bR-chloro, 5c-hydroxy: Variicolorin B

[957780-66-4]
C₂₄H₃₀ClN₃O₃ 443.972
Isol. from *Aspergillus varicolor* B-17. Amorph. powder. [α]_D²⁵ -29 (c, 0.1 in MeOH). λ_{max} 208 (log ε 3.5); 228 (log ε 3.6); 287 (log ε 3); 342 (log ε 3.2) (MeOH).

5b,5c-Dihydro, 5c-chloro, 5bR-hydroxy: Variicolorin A

[957780-65-3]
C₂₄H₃₀ClN₃O₃ 443.972
Isol. from *Aspergillus varicolor* B-17. Amorph. powder. [α]_D²⁵ -39 (c, 0.1 in MeOH). λ_{max} 210 (log ε 3.6); 228 (log ε 3.6); 285 (log ε 3.1); 340 (log ε 3.2) (MeOH).

5b,5c-Dihydro, 5bR,5c-dihydroxy: 19,20-Dihydroxyisoechinulin A

[744253-48-3]
C₂₄H₃₁N₃O₄ 425.527
Prod. by a marine *Aspergillus* sp. Oil. λ_{max} 209 (log ε 3.9); 226 (log ε 3.9); 289 (log ε 3.4); 340 (log ε 3.5) (MeOH).

5b,5c-Dihydro, 5bR,5c-dihydroxy, isopropylidene: Variicolorin D

[957780-70-0]
C₂₇H₃₅N₃O₄ 465.591
Isol. from *Aspergillus varicolor* B-17. Amorph. powder. [α]_D²⁵ -51 (c, 0.1 in MeOH). λ_{max} 210 (log ε 3.6); 229 (log ε 3.7); 288 (log ε 3.2); 340 (log ε 3.3)

(MeOH).

5b,5c-Dihydro, 5b-oxo: **Variecolorin E**
[957780-71-1]
C₂₄H₂₉N₃O₃ 407.511
Isol. from *Aspergillus variecolor* B-17.
Amorph. powder. [α]_D²⁵ -20 (c, 0.1 in
MeOH); λ_{max} 203 (log ε 3.4); 230 (log ε
3.4); 285 (log ε 3.1); 345 (log ε 3.2)
(MeOH).

3eξ-Methoxy: **Variecolorin I**
[957780-80-2]

C₂₅H₃₁N₃O₃ 421.538
Isol. from *Aspergillus variecolor* B-17.
Amorph. powder. Racemic. λ_{max} 208
(log ε 3.8); 228 (log ε 3.7); 285 (log ε
3.2); 350 (log ε 3.3) (MeOH).


Δ^{5c}-Isomer, 5bR-hydroxy: **Variecolorin C**
[957780-67-5]

C₂₄H₂₉N₃O₃ 407.511
Isol. from *Aspergillus variecolor* B-17.
Amorph. powder. [α]_D²⁵ -44 (c, 0.1 in
MeOH); λ_{max} 209 (log ε 3.4); 228 (log ε
3.4); 290 (log ε 3.1); 345 (log ε 3.2)
(MeOH).

Nagasawa, H. *et al.*, *Tet. Lett.*, 1976, 1601-
1604 (*isol, uv, ir, pmr, ms*)
Nagasawa, H. *et al.*, *Agric. Biol. Chem.*, 1979,
43, 1759-1763 (*cmr, struct*)
Li, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 337-340
(*Dihydroxyisochininulin A*)
Wang, W.-L. *et al.*, *J. Nat. Prod.*, 2007, **70**,
1558-1564 (*Variecolorins A,B,C,E,I*)
Cole, R.J. *et al.*, *Handbook of Toxic Fungal*
Metabolites, Academic Press, 1981, 475; 476;
477

Isoelaecarpicine I-231

(2-Hydroxy-6-methylphenyl) (octahydro-
7-hydroxy-8-indoliziny) methanone, 9CI.
7-Hydroxy-8-(2-hydroxy-6-methylbenzoyl)
indolizidine



(+) -form

C₁₆H₂₁NO₃ 275.347**(+) -form** [21104-52-9]

Alkaloid from the leaves of *Elaeocarpus*
fuscoides and *Elaeocarpus polydactylus*.
Prisms (EtOH). Mp 164-166°. [α]_D +29
(c, 0.30 in CHCl₃).

(±) -form [72521-90-5]

Synthetic. Mp 143-145°.


7-Deoxy, 7,8-didehydro: **Elaeocarpenine**
[942293-11-0]

C₁₆H₁₉NO₂ 257.332
Alkaloid from the leaves of *Elaeocar-*
pus fuscoides. Shows binding affinity
for the human δ-opioid receptor. Yel-
low gum (as trifluoroacetate). λ_{max} 223
(log ε 3.29) (trifluoroacetate).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**,
775-792 (*isol, uv, ir, pmr, ms*)
Tufariello, J.J. *et al.*, *J.A.C.S.*, 1979, **101**, 7114-
7116 (*synth*)
Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**,
872-875 (*isol, pmr, cmr, Elaeocarpenine*)

Isofebrifugine I-232

3-[(Octahydro-2-hydroxyfuro[3,2-b]pyri-
din-2-yl)methyl]-4(3H)-quinazolinone,
9CI. α-Dichroine. Dichroine A
[32434-44-9]




C₁₆H₁₉N₃O₃ 301.344
Stereochem. revised in 1999. Alkaloid
from the roots of the Chinese drug
Ch'ang Shan (*Dichroa febrifuga*) and
Hydrangea umbellata (Hydrangeaceae).
Cryst. (MeOH or CHCl₃/EtOH). Mp
138-139° (129-130°). [α]_D²³ +131 (c, 0.35 in
CHCl₃).

Bis(benzenesulfonyl): Mp 182.5-183.5°.
Koepfli, J.B. *et al.*, *J.A.C.S.*, 1947, **69**, 1837;
1949, **71**, 1048 (*isol, uv, struct*)
Ablondi, F. *et al.*, *J.O.C.*, 1952, **17**, 14-18 (*isol*)
Uesato, S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**,
1-5 (*pmr, cmr, conformm*)
Murata, K. *et al.*, *J. Nat. Prod.*, 1998, **61**, 729-
733 (*isol, uv, ir, pmr, cmr, ms*)
Takeuchi, Y. *et al.*, *Chem. Pharm. Bull.*, 1999,
47, 905-906 (*synth*)
Takaya, Y. *et al.*, *J. Med. Chem.*, 1999, **42**,
3163-3166 (*isol*)
Kobayashi, S. *et al.*, *J.O.C.*, 1999, **64**, 6833-
6841 (*synth, config*)
Takeuchi, Y. *et al.*, *Synthesis*, 1999, 1814-1818
(*synth*)
Ooi, H. *et al.*, *Org. Lett.*, 2001, **3**, 953-955
(*synth*)
Takeuchi, Y. *et al.*, *Tetrahedron*, 2001, **57**,
1213-1218 (*synth*)
Katoh, M. *et al.*, *Heterocycles*, 2006, **67**, 189-
204 (*synth*)
Wee, A.G.H. *et al.*, *Org. Lett.*, 2008, **10**, 3869-
3872 (*synth*)

Isoficine I-233

5,7-Dihydroxy-6-(1-methyl-2-pyrrolidi-
nyl)-2-phenyl-4H-1-benzopyran-4-one,
9CI. 5,7-Dihydroxy-6-(1-methylpyrrolidi-
nyl)flavone
[2255-62-1]




C₂₀H₁₉NO₄ 337.374
Minor alkaloid from *Ficus pantoniana*
(Moraceae). Mp 168°.

Johns, S.R. *et al.*, *Tet. Lett.*, 1965, 1987 (*pmr,*
struct)

Isoflavipucine I-234

[61277-39-2]




C₁₂H₁₅NO₄ 237.255
Isol. from *Aspergillus flavipes*. Also obt.
by heating Flavipucine, F-81 at 138-140°.
Cryst. (Et₂O). Mp 166-171°. [α]_D²¹ 0 (c, 1
in EtOH). λ_{max} 221 (ε 26100); 307 (ε
10300) (EtOH) (Derep).

Findlay, J.A. *et al.*, *Can. J. Chem.*, 1977, **55**,
600 (*isol, uv, ir, pmr, cmr, ms, struct*)

Isoflustramine D I-235

[104387-14-6]


Relative
ConfigurationC₂₁H₂₉BrN₂ 389.377

Isol. as a 65:35 inseparable mixt. with
Flustramine D, F-107. Metab. from the
marine bryozoan *Flustra foliacea*. [α]_D²⁵ -
14.6 (c, 0.07 in CH₂Cl₂). λ_{max} 240 ; 290
(MeOH) (Berdy).

Laycock, M.V. *et al.*, *Can. J. Chem.*, 1986, **64**,
1312-1316 (*isol, uv, ir, pmr, ms, struct*)

Isogentialutine I-236

[55399-77-4]




C₉H₁₁NO 149.192
Alkaloid from the roots of *Gentiana*
tibetica (Gentianaceae). Mp 131°.

Rulko, F. *et al.*, *Pol. J. Pharmacol. Pharm.*,
1974, **26**, 561; *CA*, **82**, 57984w (*isol, uv, ir,*
pmr, ms, struct)

Isogranulatimide I-237

[219829-00-2]




C₁₅H₈N₄O₂ 276.254
Alkaloid from the ascidian *Didemnum*
granulatum. G2 specific cell cycle check-
point inhibitor. Deep purple crystalline.
(MeCN aq.) or amorph. red solid. λ_{max}
210 (ε 10230); 231 (ε 10650); 280 (ε 6550);
470 (ε 1880) (MeOH).

Berlinck, R.G.S. *et al.*, *J.O.C.*, 1998, **63**, 9850-
9856 (*isol, uv, ir, pmr, cmr, synth*)
Vervoort, H.C. *et al.*, *J. Nat. Prod.*, 1999, **62**,
389-391 (*isol, uv, ir, pmr, cmr, ms*)
Piers, E. *et al.*, *J.O.C.*, 2000, **65**, 530-535
(*synth*)

Isoharringtonine

[26833-86-3]

Absolute
Configuration $C_{28}H_{37}NO_9$ 531.602

Alkaloid from *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus harringtonia* var. *drupacea*, *Cephalotaxus hainanensis*, *Cephalotaxus wilsoniana* and *Cephalotaxus sinensis* (Cephalotaxaceae). Shows significant antineoplastic activity. Mp 69-72.5°. $[\alpha]_D$ -99.6 (c, 1.06 in $CHCl_3$). Log P 1.04 (uncertain value) (calc). λ_{max} 261 (ϵ 724); 290 (ϵ 4570) (EtOH) (Derep).

▶ NR2935500

Parent acid: **Isoharringtonic acid**. 5'-Des-O-methylisoharringtonine
[55825-71-3]

 $C_{27}H_{35}NO_9$ 517.575

Alkaloid from *Cephalotaxus hainanensis* (Cephalotaxaceae) and *Cephalotaxus harringtonia* var. *drupacea*. Amorph. solid. Mp 224-228°. $[\alpha]_D$ -113 (c, 0.33 in DMSO).

3''-Hydroxy: **Cephalzomine C**

[288300-74-3]

 $C_{28}H_{37}NO_{10}$ 547.601

Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent. Amorph. solid. $[\alpha]_D$ -122 (c, 2.3 in MeOH). λ_{max} 291 (ϵ 2100) (MeOH).

3''-Epimer, 3''-hydroxy: **Cephalzomine D**

[182325-75-3]


 $C_{28}H_{37}NO_{10}$ 547.601

Alkaloid from *Cephalotaxus harringtonia* var. *nana*. Cytotoxic agent. Amorph. solid. $[\alpha]_D$ -97 (c, 13.7 in MeOH). λ_{max} 291 (ϵ 2100) (MeOH).

Powell, R.G. *et al.*, *Tet. Lett.*, 1970, 815 (struct)Brandänge, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1974, **28**, 1237 (abs config)Weisleder, D. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 114 (cmr)Xue, Z. *et al.*, *Yaoxue Xuebao*, 1981, **16**, 752; *CA*, **96**, 82690u (Isoharringtonic acid)Huang, L. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 157 (rev)Takano, I. *et al.*, *Phytochemistry*, 1996, **43**, 299 (Isoharringtonic acid)Morita, H. *et al.*, *Tetrahedron*, 2000, **56**, 2929-2934 (Cephalzomines)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, IK5500**Isohematinic acid**

I-239

4-Methylene-2,5-dioxo-3-pyrrolidinepropanoic acid, 9CI. Ishaematinic acid



I-238

 $C_8H_9NO_4$ 183.163

Sol. MeOH, Me_2CO , bases; fairly sol. EtOAc; poorly sol. H_2O , C_6H_6 , hexane. λ_{max} 225 (ϵ 8000) (MeOH) (Derep). λ_{max} 225 ($E1\%/1cm$ 450) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) 300 mg/kg.


(±)-form [86408-37-9]

Alkaloid from *Actinoplanes philippinensis*. Cryst. (MeOH). Mp 138°.

▶ UY1881500

Takeuchi, M. *et al.*, *J. Antibiot.*, 1983, **36**, 493 (isol)Itoh, Y. *et al.*, *J. Antibiot.*, 1983, **36**, 497 (uv, ir, pmr, cmr, struct)**Isokibdelone A**

[916660-08-7]

Absolute
Configuration $C_{29}H_{24}ClNO_{10}$ 581.962

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. orange solid. $[\alpha]_D$ +133 (c, 0.01 in $CHCl_3$). λ_{max} 239 (ϵ 27600); 271 (ϵ 26600); 319 (ϵ 16300); 404 (ϵ 11700); 442 (sh) (EtOH).

11-O- α -Rhamnopyranoside: [916660-07-6] $C_{35}H_{34}ClNO_{14}$ 728.105

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. orange solid. $[\alpha]_D$ +60 (c, 0.01 in $CHCl_3$). λ_{max} 210 (ϵ 25000); 229 (ϵ 23200); 271 (ϵ 17400); 410 (ϵ 6700); 442 (sh) (EtOH).

6,7-Dihydro: **Isokibdelone B**

[916660-09-8]

 $C_{29}H_{26}ClNO_{10}$ 583.978

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. orange solid. $[\alpha]_D$ +82 (c, 0.02 in $CHCl_3$). λ_{max} 203 (ϵ 13700); 244 (ϵ 9500); 266 (sh); 326 (sh); 447 (ϵ 3700); 583 (sh) (EtOH).


5,16-Hydroquinone, 6,7-dihydro: **Isokibdelone C**

[916660-10-1] Artifact prod. after storage.

Ratnayake, R. *et al.*, *Org. Lett.*, 2006, **8**, 5267-5270 (isol, pmr, cmr)**Isoleontalbine**

I-241

5,17-Didehydromatridin-15-one, 9CI [40444-77-7]

 $C_{15}H_{22}N_2O$ 246.352


Stereoisomer of Darvasine, D-80 and

Leontalbine, L-93. Alkaloid from *Leontice smirnowii* (Leonticaceae). Oil. Mp 207° (as perchlorate). $[\alpha]_D$ -147 (EtOH). λ_{max} 240 (log ϵ 4.2) (no solvent reported).

Bohlmann, F. *et al.*, *Chem. Ber.*, 1958, **91**, 2176-2189 (config)Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 410-512; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 386-512 (rev) **α -Isolupanine**

I-242

11-Isolupanine. Tetrahydrothermopsine. Alkaloid P7



(±)-form

 $C_{15}H_{24}N_2O$ 248.367

(±)-form [486-87-3]

Alkaloid from *Lupinus sericeus*, *Lupinus mutabilis*, *Lupinus leontopetalum* tubers and *Lupinus caudatus* (Fabaceae). Rectangular plates. Mp 75-76° Mp 83-84° (dimorph.). $[\alpha]_D^{23}$ +65.9 (c, 1.495 in EtOH).

Perchlorate:

Cryst. (MeOH), Mp 248-249°.

Picrate:

Long yellow needles (EtOH). Mp 197-198°.


(-)-form

By hydrogenation of (-)-Thermopsine in T-363. Mp 75-76°.

Marion, L. *et al.*, *Can. J. Chem.*, 1951, **29**, 22-29; 1953, **31**, 181-186 (isol, ir, struct)Baranowski, P. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1966, **40**, 73-80 (synth)Wiewirowski, M. *et al.*, *Can. J. Chem.*, 1967, **45**, 1447-1457 (ir, pmr, conformn)Panov, P.P. *et al.*, *CA*, 1972, **77**, 72554w (isol)Vulfson, N.S. *et al.*, *Chem. Heterocycl. Compd.*, 1974, **10**, 221-229 (ms)Bohlmann, F. *et al.*, *Chem. Ber.*, 1975, **108**, 1043-1051 (cmr)Anderson, J.N. *et al.*, *J.O.C.*, 1976, **41**, 3441-3444 (isol)Kennelly, E.J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1385-1389 (isol, pmr, cmr)**Isomalindine-16-carboxylate**

I-243

[203255-40-7]


 $C_{21}H_{21}N_3O_2$ 347.416

Zwitterionic. Alkaloid from *Ophiorrhiza* cf. *communis* (Rubiaceae). Needles (MeOH/EtOAc). Mp 193° (dec.). $[\alpha]_D$ +667 (c, 0.03 in MeOH).

Arbain, D. *et al.*, *Aust. J. Chem.*, 1997, **50**, 1109-1110 (isol, pmr, cmr, struct)

Isomatrine

[17801-36-4]

Absolute
configurationC₁₅H₂₄N₂O 248.367


Stereoisomer of Matrine, M-121, Alломатrine, A-628, Sophoridine, S-380 and Darvasamine, D-79. Alkaloid from roots of *Sophora flavescens* (Fabaceae). Mp 132-134°. [α]_D²⁵ +44 (CHCl₃).

Hydrobromide:

V. deliquescent cryst. Mp 166.5-168°.

Methiodide: Mp 213° dec.Ueno, A. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2560 (*isol, ir, pmr, cryst struct*)Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 71; *Chem. Nat. Compd. (Engl. Transl.)*, 66 (*rev, stereochem*)**Isomigrastatin**

[415952-70-4]

Absolute
ConfigurationC₂₇H₃₉NO₇ 489.608

Glutarimide-lactone antibiotic. Closely related to Lactimidomycin, L-7. Prod. by *Streptomyces platensis* NRRL18993. Antitumour agent. Oil. [α]_D²⁵ +170 (c, 0.18 in CHCl₃).

O-De-Me: 8-O-DemethylisomigrastatinC₂₆H₃₇NO₇ 475.581

Prod. by *Streptomyces platensis* NRRL18993. Oil. [α]_D²³ +83 (c, 0.1 in CHCl₃).

16,17-Didehydro: 16,17-DidehydroisomigrastatinC₂₇H₃₇NO₇ 487.592

Prod. by *Streptomyces platensis* NRRL18993. Oil. [α]_D +123.9 (c, 0.18 in CHCl₃).

16,17-Didehydro, O-de-Me: 16,17-Didehydro-8-O-demethylisomigrastatinC₂₆H₃₅NO₇ 473.565

Prod. by *Streptomyces platensis* NRRL18993. Oil.

17R-Hydroxy: 17-HydroxyisomigrastatinC₂₇H₃₉NO₈ 505.607

Prod. by *Streptomyces platensis* NRRL18993. Oil. [α]_D²⁵ +140 (c, 0.1 in CHCl₃).

17R-Hydroxy, O-de-Me: 17-Hydroxy-8-O-demethylisomigrastatinC₂₆H₃₇NO₈ 491.58

Prod. by *Streptomyces platensis* NRRL18993.

Demethoxy: Demethoxyisomigrastatin**I-244**C₂₆H₃₇NO₆ 459.581

Prod. by *Streptomyces platensis* NRRL18993. Oil. [α]_D²⁵ +77 (c, 0.1 in CHCl₃).

Demethoxy, 16,17-didehydro: 16,17-DidehydrodemethoxyisomigrastatinC₂₆H₃₅NO₆ 457.566

Prod. by *Streptomyces platensis* NRRL18993. Oil. [α]_D²⁵ +7.3 (c, 0.05 in CHCl₃).

Demethoxy, 17R-hydroxy: 17-Hydroxydemethoxyisomigrastatin. 8,9-Dihydro-9-hydroxylactimidomycin

[866629-46-1]


C₂₆H₃₇NO₇ 475.581

Prod. by *Streptomyces amphibiosporus* ATCC53964 and *Streptomyces platensis* NRRL18993. Oil. [α]_D²⁵ +80 (c, 0.1 in CHCl₃).

Woo, E.J. *et al.*, *J. Antibiot.*, 2002, **55**, 141-146 (*Isomigrastatin*)Ju, J. *et al.*, *J.A.C.S.*, 2005, **127**, 1622-1623; 11930-11931 (*isol, biosynth, derivs*)Krauss, I.J. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 5576-5579 (*synth*)Ju, J. *et al.*, *Org. Lett.*, 2007, **9**, 5183-5186 (*isol, derivs*)**Isomitomycin A**

AX 2. Antibiotic AX 2


[91917-64-5]

Absolute
ConfigurationC₁₆H₁₉N₃O₆ 349.343


Quinone antibiotic. Metab. of *Streptomyces caespitosus* NRRL12508. Prod. by intramolecular rearrangement of Mitomycin A, M-653. Shows antitumour props. Yellow-orange needles (CHCl₃). Sol. MeOH, CHCl₃; fairly sol. H₂O; poorly sol. hexane. Mp 78-80°. [α]_D²³ -273 (c, 0.17 in CHCl₃). Isomerises to Mitomycin A, M-653. λ_{max} 205 (ε 10800); 294 (ε 5200); 410 (ε 340) (MeOH) (Berdy). ▶ LD₅₀ (mus, ia) 27 mg/kg, LD₅₀ (mus, ipr) 7.8 mg/kg.

Kono, M. *et al.*, *J.A.C.S.*, 1987, **109**, 7224-7225 (*isol, struct*)Kasai, M. *et al.*, *J.O.C.*, 1992, **57**, 7296-7299 (*synth, pmr*)**Isomitraphyllinol**

[949081-83-8]

Absolute
ConfigurationC₂₁H₂₆N₂O₅ 386.447

Alkaloid from the leaves of *Mitragyna hirsuta*.

Kitajima, M. *et al.*, *J. Nat. Med. (Tokyo)*, 2007, **61**, 192-195 (*isol, pmr, cmr*)**I-246**C₁₅H₁₆O₄ 260.289**Ac: Isomurralonginol acetate**

[112667-51-3]

C₁₇H₁₈O₅ 302.326

Constit. of *Murraya exotica*. Oil. [α]_D +18.8 (c, 0.2 in CHCl₃).

O-(3-Methylbutanoyl): Isomurralonginol isoaltrate

[131652-27-2]

C₂₀H₂₄O₅ 344.407

Constit. of *Murraya paniculata*. Pale yellow syrup. [α]_D +14.5 (c, 0.13 in CHCl₃).

O-(3-Pyridinecarbonyl): Isomurralonginol nicotinate


[112606-77-6]

C₂₁H₁₉NO₅ 365.385


Constit. of *Murraya paniculata*. Yellow oil. [α]_D +31.8 (c, 0.145 in CHCl₃).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 4277 (*isol, uv, ir, pmr, ms*)Ito, C. *et al.*, *Heterocycles*, 1987, **26**, 2959Ito, C. *et al.*, *J.C.S. Perkin 1*, 1990, 2047 (*isol, uv, ir, pmr, cmr, ms*)**Isomurrayazoline**

[85547-20-2]

C₂₃H₂₅NO 331.457**I-248****Isomonodoroinole**

6,6'-[1,2-Bis(2-methyl-1-propenyl)-1,2-ethanediy]bis-1H-indole, 9CI. 4,5-Di-6-indolyl-2,7-dimethyl-2,6-octadiene [110325-62-7]

C₂₆H₂₈N₂ 368.521

Alkaloid from the seeds of *Monodora myristica* (Annonaceae).

Muhammad, I. *et al.*, *J. Bangladesh Acad. Sci.*, 1987, **11**, 1-7; *CA*, **107**, 172452t**Isomurralonginol****I-249**


8-[1-(Hydroxymethyl)-2-methyl-2-propenyl]-7-methoxy-2H-1-benzopyran-2-one, 9CI

[112606-75-4]

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Cryst. (C₆H₆/CHCl₃). Mp 269-270°. [α]_D -7.33 (CHCl₃).

Bhattacharya, L. *et al.*, *Phytochemistry*, 1982, **21**, 2432 (*isol, uv, ir, pmr, ms, struct*)

Isonaamidine A I-251
[110189-04-3]



C₂₁H₁₉N₅O₄ 405.412

Two tautomers of the -NH(6)C=N(3'') group are possible. Alkaloid from the Red Sea sponge *Leucetta chagosensis* and from mollusc *Notodoris citrina*. λ_{\max} 227 (€ 20000); 264 (€ 540); 358 (€ 6000); 366 (€ 6800); 380 (€ 590) (dioxan) (Derep).

O¹¹-Me: Isonaamidine B

[121819-70-3]

C₂₂H₂₁N₅O₄ 419.439

Alkaloid from *Leucetta chagosensis*. Amorph. yellow solid (as Zn complex). Also *isol.* as the bimolecular Zn complex with Isonaamidine D. λ_{\max} 226 (€ 19400); 276 (€ 4300); 374 (€ 8000) (MeOH) (Berdy). λ_{\max} 228 (€ 22400); 278 (€ 3900); 368 (€ 8700); 380 (€ 8500) (MeOH) (as Zn complex).

O¹¹-Me, N¹¹-de-Me: Isonaamidine D

C₂₁H₁₉N₅O₄ 405.412

Alkaloid from *Leucetta cf. chagoensis*. Amorph. yellow solid. λ_{\max} 226 (€ 18900); 278 (€ 4000); 370 (€ 6600) (MeOH).

Di-Me ether: Isonaamidine C

[146845-43-4]

C₂₃H₂₃N₅O₄ 433.466

Alkaloid from a *Leucetta* sp. and from *Notodoris gardineri*. Yellow solid (as Zn complex). *Isol.* as the Zn complex, to which the CAS no. refers. λ_{\max} 263; 275; 367; 580 (MeOH) (Zn complex). λ_{\max} 241 (€ 13100); 275 (€ 7400); 380 (€ 19100) (CHCl₃).

10-Methoxy, di-Me ether: Isonaamidine E

C₂₄H₂₅N₅O₅ 463.492

Alkaloid from *Leucetta chagosensis*. Cytotoxic. Amorph. yellow solid. λ_{\max} 237 (€ 23840); 279 (€ 5550); 383 (€ 11340) (CHCl₃).

Carmely, S. *et al.*, *Tetrahedron*, 1989, **45**, 2193-2200 (*isol, uv, ir, pmr, cmr, ms, struct*)

Alvi, K.A. *et al.*, *Tetrahedron*, 1993, **49**, 329 (*Isonaamidine C*)


Copp, B.R. *et al.*, *J. Med. Chem.*, 1998, **41**, 3909-3911 (*Isonaamidine C*)

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 384-386 (*Isonaamidine D* complexes)

Gross, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1190-1193 (*Isonaamidine E*)

Nakamura, S. *et al.*, *Heterocycles*, 2003, **60**, 583-598 (*synth*)

Isonaamine A I-252
4,4'-[(2-Amino-1H-imidazole-1,4-diyl)-bis(methylene)]bisphenol, 9CI. 2-Amino-1,4-bis(p-hydroxybenzyl)imidazole [110189-02-1]



C₁₇H₁₇N₃O₂ 295.34

Alkaloid from the Red Sea sponge *Leucetta chagosensis* and the Indo-Pacific nudibranchs *Notodoris gardineri* and *Notodoris citrina*. Yellow powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 227; 265; 275; 294; 300 (MeOH) (Derep).

3''-Methoxy, 4',4''-di-Me ether: Isonaamine C

C₂₀H₂₃N₃O₃ 353.42

Alkaloid from the sponge *Leucetta chagosensis*. Cytotoxic. Amorph. yellow solid. λ_{\max} 251 (€ 6080); 276 (€ 6450); 379 (€ 1210) (CHCl₃).


Carmely, S. *et al.*, *Tetrahedron*, 1989, **45**, 2193 (*isol, ir, pmr, cmr, ms, struct*)

Alvi, K.A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1509 (*isol, uv, pmr*)

Molina, P. *et al.*, *J.O.C.*, 1999, **64**, 2540-2544 (*synth, pmr, cmr*)

Gross, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1190-1193 (*Isonaamine C*)

Isonaamine B I-253
[203116-58-9]




C₁₉H₂₁N₃O₂ 323.394

Alkaloid from the sponge *Leucetta cf. chagosensis*. Yellow glass. λ_{\max} 226 (€ 19300); 276 (€ 5800); 308 (€ 3700) (MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 384-386 (*isol, uv, ir, pmr, cmr*)

Isonargenicin A1 I-254
[74666-93-6]




C₂₈H₃₇NO₈ 515.602

Prod. by *Nocardia argentinensis*. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Co-metabolite and possible rearrangement product of Nargenicin A₁ in N-276.

Cane, D.E. *et al.*, *J.A.C.S.*, 1984, **106**, 784-787 (*isol, struct*)

Isonitrin A I-255
4-Isocyano-3'-methylspiro[6-oxabicyclo[3.1.0]hex-3-ene-2,2'-oxirane], 9CI [83016-48-2]




C₈H₇NO₂ 149.149

Prod. by *Trichoderma hamatum*. Active against gram-positive and gram-negative bacteria, yeasts and fungi. Needles. Sol. MeOH. Mp 91-91.5°. [α]_D +9 (c, 0.2 in MeOH). λ_{\max} 230 (€ 13900) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 160 mg/kg. WH1424250
Fujiwara, A. *et al.*, *Agric. Biol. Chem.*, 1982, **46**, 1803; 1811 (*isol, props*)

Isonitrinic acid E I-256
3-(4-Isocyano-6-oxabicyclo[3.1.0]hex-1-yl)-2-propenoic acid, 9CI [83016-51-7]




C₉H₇NO₃ 177.159

Prod. by *Trichoderma hamatum*. Weakly active antibacterial agent. Needles. Sol. MeOH, Et₂O, CCl₄, CHCl₃; poorly sol. H₂O, hexane. Mp 114-117° dec. [α]_D +155.5 (c, 0.2 in MeOH). λ_{\max} 223 (€ 14160) (MeOH) (Berdy). λ_{\max} 222 (€ 15500) (Et₂O) (Berdy).

▶ LD₅₀ (mus, ipr) 240 mg/kg. UD3390800
Fujiwara, A. *et al.*, *Agric. Biol. Chem.*, 1982, **46**, 1803; 1811 (*isol, struct*)

Isonoyaine I-257
[132185-18-3]




C₂₁H₂₃NO₇ 401.415

Alkaloid from *Sarcocapnos saetabensis*. Powder. λ_{\max} 241; 292; 313 (EtOH).

Blanco, O. *et al.*, *Phytochemistry*, 1991, **30**, 2071-2074 (*isol, pmr, cmr*)

Isoobscurinine

I-258


C₁₇H₂₄N₂O 272.389

Alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Off-white plates (hexane/Me₂CO). Mp 106-107°. [α]_D +107 (c, 1.1 in MeOH). Possibly an artifact. Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077 (*isol, ir, pmr, cmr, ms, struct*)

Isoocinotine

I-259

Docosahydro-1H-pyrido[1,2-*j*][1,5,10]triazacyclodocosin-10(11H)-one, 9CI

C₂₃H₄₅N₃O 379.628**(R)-form** [21008-80-0]

Alkaloid from the stem bark of *Oncinotis nitida*. Cryst. (pentane). Mp 66-71° (block). [α]_D -37 (c, 0.067 in MeOH).

(±)-form [62107-31-7]


Synthetic. Cryst. (Et₂O/pentane). Mp 52-68°.

Badawi, M.M. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1813-1817 (*struct*)
Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 414-434; 1976, **59**, 3013-3025 (*isol, ir, ord, cd, ms, abs config, synth*)
Scheiper, B. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 11960-11965 (*synth, pmr, cmr, ms*)
Cheng, H.-Y. *et al.*, *Tetrahedron*, 2007, **63**, 3000-3005 (*synth*)

Isooxodesmosine

I-260

[147044-50-6]

C₂₃H₃₇N₅O₇ 495.575


Cross-linking amino acid. Isol. from bovine aorta elastin.

Suyama, K. *et al.*, *Bioorg. Med. Chem. Lett.*, 1992, **2**, 1767-1770 (*isol, pmr*)

Isopalythazine

I-261

4,6,7,9-Tetrahydro-1H,3H-dipyran[3,4-b:4',3'-e]pyrazine-3,7-dimethanol, 9CI [72681-95-9]



Relative Configuration

C₁₂H₁₆N₂O₄ 252.269


Isol. from the zoanthid *Palythoa tuberculosa*. Mp 216-219°.

Uemura, D. *et al.*, *Chem. Lett.*, 1979, 1481-1482 (*isol, cmr, struct, synth*)
Jarglis, P. *et al.*, *Angew. Chem., Int. Ed.*, 1982, **21**, 141-142 (*synth*)

Isoperlolyrine

I-262

5-(5H-Pyrido[4,3-*b*]indol-1-yl)-2-furanmethanol, 9CI. 1-(5-Hydroxymethyl-2-furanyl)-γ-carboline [91897-55-1]

C₁₆H₁₂N₂O₂ 264.283


Alkaloid from the seeds of *Gloriosa superba* (Liliaceae). Mp 186° dec.

Dvořáčková, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1984, **49**, 1536 (*isol, uv, pmr, ms, struct*)

Isophellibilidine

I-263

1,6-Didehydro-12-(hydroxymethyl)-3-methoxy-c-homo-D-nor-16-oxaerythrinan-15(12H)-one, 9CI [80249-90-7]



Absolute Configuration

C₁₇H₂₃NO₄ 305.373


Alkaloid from *Phelline billiardieri* (Phellinaceae). Mp 132°. [α]_D +204 (c, 1.2 in CHCl₃).

Langlois, N. *et al.*, *Tet. Lett.*, 1981, **22**, 2263 (*uv, ir, cd, ms, pmr, cmr, synth, struct*)

Isophomazarin

I-264

7-Butyl-5,10-dihydro-4,6,9-trihydroxy-8-methoxy-5,10-dioxobenzo[g]quinoline-2-carboxylic acid, 9CI [71295-01-7]

C₁₉H₁₇NO₈ 387.345

Isol. from cultures of *Pyrenochaeta terrestris*. Red needles. Mp 215-216°.

Effenberger, R. *et al.*, *J.C.S. Perkin 1*, 1979, 823

Isophysostigmine

I-265


Struct. unknown. Tentative occur. Not detected by Salway. Alkaloid from seeds of *Physostigma venenosum* (Fabaceae). Insol. H₂O. Mp 202° (H₂SO₄ salt). Shows similar chem. and physiol. activity to Physostigmine (no details given of chemistry).

Ogiu, *et al.*, *Apoth. Ztg.*, 1904, **19**, 891
Salway, A.H. *et al.*, *J.C.S.*, 1911, **99**, 2148-2159 (*Physostigma venenosum constits*)
Robinson, B. *et al.*, *J.C.S.*, 1964, 1503-1506 (*Physostigma venenosum constits*)

Isopiperolein B

I-266

1-[10-(1,3-Benzodioxol-5-yl)-1-oxo-9-decyl]pyrrolidine. 10-(3,4-Methylene-dioxyphenyl)-9-decenoic acid pyrrolidide

C₂₁H₂₉NO₃ 343.465


Alkaloid from the berries of *Piper nigrum* (pepper). Yellow oil. λ_{max} 260 (no solvent reported).

Srinivas, P.V. *et al.*, *Phytochemistry*, 1999, **52**, 957-958

Isopiscerythramine

I-267

4'-Amino-3',5,7-trihydroxy-5'-methoxy-2',8-diprenylisoflavone [151590-42-0]

C₂₆H₂₉NO₆ 451.518


Constit. of the root bark of *Piscidia erythrina*. Pale yellow powder. Mp 155-156°. λ_{max} 211; 260 (sh); 266; 300 (sh); 330 (sh) (MeOH).

Moriyama, M. *et al.*, *Phytochemistry*, 1993, **32**, 1317-1325 (*isol, uv, pmr, cmr, ms*)

Isoplatydesmine

I-268

3,9-Dihydro-2-(1-hydroxy-1-methyl-ethyl)-9-methylfuro[2,3-*b*]quinolin-4(2H)-one, 9CI



(R)-form

C₁₅H₁₇NO₃ 259.304**(R)-form** [27495-37-0]

Alkaloid from *Pelea barbiger* (preferred genus name *Melicope*) and *Araliopsis*

soyauuxii (Rutaceae). Cryst. (CHCl₃). Mp 191°. [α]_D²⁰ +60 (c, 1 in CHCl₃).

Me ether: O-Methylisoplatydesmine

[220961-34-2]

C₁₆H₁₉NO₃ 273.331

Constit. of *Skimmia laureola*. Cryst. [α]_D²⁰ +40 (c, 0.1 in CHCl₃). λ_{\max} 211 (log ϵ 2.08); 237 (log ϵ 2.33) (MeOH).

(ξ)-form

2'-Hydroxy: **Folisine**

[36069-06-4]

C₁₅H₁₇NO₄ 275.304

Alkaloid from *Haplophyllum foliosum* (Rutaceae). Cryst. (MeOH). Mp 236-237°. [α]_D²⁵ -123 (c, 1.22 in MeOH). λ_{\max} 215 (log ϵ 4.5); 237 (log ϵ 4.43); 251 (sh) (log ϵ 4.24); 299 (sh) (log ϵ 4.01); 310 (log ϵ 4.12); 321 (log ϵ 4.07) (MeOH).

2'-Hydroxy, hydrochloride: Mp 230°.

(\pm)-form [41234-42-8]

Synthetic. Mp 213-215° (186-187°, 201-203°).

Bowman, R.M. *et al.*, *J.C.S. (C)*, 1966, 1504 (synth)

Bessonova, I.A. *et al.*, *Khim. Prir. Soedin.*, 1971, 7, 629; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, 7, 608 (Folisine)

Higa, T. *et al.*, *Phytochemistry*, 1974, 13, 1269 (isol, uv, ir, ms, struct)

Vaquette, J. *et al.*, *Phytochemistry*, 1976, 15, 743 (isol, uv, ir, pmr, ms, config)


Coppola, G.M. *et al.*, *J. Het. Chem.*, 1983, 20, 1589 (synth, uv, ir, pmr, cmr)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, 12, 223-229 (Methylisoplatydesmine, isol)

Isopolyalthenol

I-269

[99615-96-0]



C₂₃H₃₁NO 337.504

Alkaloid from *Polyalthia suaveolens* (Annonaceae). Cryst. (petrol). Mp 129°. [α]_D²⁰ -49 (c, 0.8 in EtOH).


Kunesch, N. *et al.*, *Tet. Lett.*, 1985, 26, 4937 (uv, ir, pmr, cmr, ms, struct)

2-Isopropenyl-5-isopropylpyrazine

I-270

2-(1-Methylethenyl)-5-(1-methylethyl)pyrazine

[704909-07-9]



C₁₀H₁₄N₂ 162.234


Prod. by *Chondromyces crocatus*. Liq.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (isol, synth, pmr, cmr, ms)

2-Isopropenyl-6-isopropylpyrazine

I-271

2-(1-Methylethenyl)-6-(1-methylethyl)pyrazine
[704909-06-8]



C₁₀H₁₄N₂ 162.234


Prod. by *Chondromyces crocatus*.

Schulz, S. *et al.*, *Tetrahedron*, 2004, 60, 3863-3872 (isol)

2-Isopropyl-3,5-dimethoxy-6-methylpyrazine

I-272

2,6-Dimethoxy-3-methyl-5-(1-methylethyl)pyrazine, 9CI
[32021-42-4]



C₁₀H₁₆N₂O₂ 196.249

Constit. of Galbanum oil.


Burrell, J.W.K. *et al.*, *Chem. Ind. (London)*, 1970, 1409 (synth)

Takken, H.J. *et al.*, *J. Agric. Food Chem.*, 1975, 23, 638 (synth)

2-Isopropyl-10-methoxybenzimidazo[6,7,1-def][1,6]naphthyridine

I-273

[597553-93-0]



C₁₆H₁₅N₃O 265.314


Alkaloid from a *Xestospongia* sp. Brown gum.

Calcul, L. *et al.*, *Tetrahedron*, 2003, 59, 6539-6544 (isol, pmr, cmr)

6-Isopropyl-3-(4-methoxybenzyl)-4-methyl-2,5-morpholine-dione

I-274

3-(4-Methoxyphenylmethyl)-4-methyl-6-(1-methylethyl)-2,5-morpholinedione, 9CI



C₁₆H₂₁NO₄ 291.346

(3*R**,6*R**)-form


Isol. from the sea hare *Bursatella leachii*. Amorph. solid. [α]_D²⁰ -58.2 (c, 0.05 in MeOH). λ_{\max} 223 (log ϵ 2.93); 250 (log ϵ 2.2); 273 (log ϵ 2.08); 281 (log ϵ 1.99) (MeOH).

Suntornchashweij, S. *et al.*, *J. Nat. Prod.*, 2005, 68, 951-955 (isol, pmr, cmr)

2-Isopropyl-3-methoxy-5-(1-methylpropyl)pyrazine

I-275

3-Methoxy-2-(1-methylethyl)-5-(1-methylpropyl)pyrazine
[870543-97-8]



C₁₂H₂₀N₂O 208.303


Prod. by *Chondromyces crocatus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (isol, synth, pmr, cmr, ms)

2-Isopropyl-3-methoxy-5-(2-methylpropyl)pyrazine

I-276

3-Methoxy-2-(1-methylethyl)-5-(2-methylpropyl)pyrazine
[68290-69-7]



C₁₂H₂₀N₂O 208.303


Prod. by *Chondromyces crocatus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (isol, synth, pmr, cmr, ms)

3-Isopropyl-2-methoxy-5-methylpyrazine

I-277

2-Methoxy-5-methyl-3-(1-methylethyl)pyrazine, 9CI
[32021-41-3]



C₉H₁₄N₂O 166.222

Constit. of Galbanum oil.


Burrell, J.W.K. *et al.*, *Chem. Ind. (London)*, 1970, 1409 (synth)








Takken, H.J. *et al.*, *J. Agric. Food Chem.*, 1975, 23, 638 (synth)

3-Isopropyl-2-methoxy-pyrido[2,3-b]pyrazine

I-278

2-Methoxy-3-(1-methylethyl)pyrido[2,3-b]pyrazine, 9CI
[103518-16-7]



C₁₁H₁₃N₃O 203.243Isol. from the lipid fraction extract of the springtail *Tetodontophora bielensis*. Noncryst.Buděšínský, M. *et al.*, *Coll. Czech. Chem. Comm.*, 1986, **51**, 956 (*isol, ms, pmr, struct*)**11-Isopropyl-5-methyl-5H-indolo[3,2-b]quinoline** I-279*11-Isopropyl-5-methyl-5H-quinoline. 11-Isopropylcryptolepine* [219843-82-0]C₁₉H₁₈N₂ 274.365Alkaloid from the dried roots of *Cryptolepis sanguinolenta*. Deep purple solid.Hadden, C.E. *et al.*, *J. Nat. Prod.*, 1999, **62**, 238-240 (*isol, pmr, cmr, N-15 nmr*)**5-Isopropyl-2-(2-methylpropyl)-1H-imidazole** I-280*5-(1-Methylethyl)-2-(2-methylpropyl)-1H-imidazole. 2-Isobutyl-5-isopropyl-1H-imidazole. Catharsitoxin F*C₁₀H₁₈N₂ 166.266Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.Suenaga, K. *et al.*, *Tet. Lett.*, 2001, **42**, 7079-7081 (*isol, pmr*)**2-Isopropyl-5-(1-methylpropyl)pyrazine** I-281*2-(1-Methylethyl)-5-(1-methylpropyl)pyrazine* [595605-19-9]C₁₁H₁₈N₂ 178.277Prod. by *Chondromyces crocatus* and *Paenibacillus polymyxa*.Beck, H.C. *et al.*, *FEMS Microbiol. Lett.*, 2003, **220**, 67-73 (*isol*)Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*)**2-Isopropyl-5-(2-methylpropyl)pyrazine** I-282*2-(1-Methylethyl)-5-(2-methylpropyl)pyrazine. 2-Isobutyl-5-isopropylpyrazine* [68290-70-0]C₁₁H₁₈N₂ 178.277Prod. by *Chondromyces crocatus* and *Paenibacillus polymyxa*.Beck, H.C. *et al.*, *FEMS Microbiol. Lett.*, 2003, **220**, 67-73 (*isol*)Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol*)**2-Isopropyl-5-methylpyrazine, 8CI** I-283*2-Methyl-5-(1-methylethyl)pyrazine, 9CI. FEMA 3554* [13925-05-8]C₈H₁₂N₂ 136.196Present in various cooked foods. Prod. by *Chondromyces crocatus*. Liq. with peanut-like odour. Bp₁₃ 68-70° (as 1:1 mixt. with the 2,6-isomer). n_D²³ 1.4865.Goldman, I.M. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 694-705 (*synth, ms, isol*)Peer, H.G. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1969, **88**, 1335-1336 (*synth*)Oser, B.L. *et al.*, *Food Technol. (Chicago)*, 1978, **32**, 60-62; **32(2)**, 64-66; 68-70 (*use*)Cantalego, M.J. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 1853-1860 (*glc, detn, earth-almond*)Ulrich, D. *et al.*, *Am. J. Potato Res.*, 2000, **77**, 111-117 (*glc, detn*)Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*)**2-Isopropyl-6-methylpyrazine** I-284*2-Methyl-6-(1-methylethyl)pyrazine, 9CI* [24541-74-0]C₈H₁₂N₂ 136.196Prod. by *Chondromyces crocatus*. No phys. props. reported.Peer, H.G. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1969, **88**, 1335-1336 (*synth*)Bramwell, A.F. *et al.*, *J.C.S.(C)*, 1971, 1627-1632 (*synth*)Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*)**Isopropylpyrazine** I-285*(1-Methylethyl)pyrazine, 9CI. FEMA 3940* [29460-90-0]C₇H₁₀N₂ 122.169Prod. by *Chondromyces crocatus*. Flavouring agent. Oil.Oerteh, R.P. *et al.*, *Anal. Chem.*, 1972, **44**, 1589 (*Raman*)Rizzi, G.P. *et al.*, *J.O.C.*, 1974, **39**, 3598 (*pmr, synth*)Konakahara, T. *et al.*, *Heterocycles*, 1979, **12**, 373Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (*isol*)**Isoprosopinine A** I-286*12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-6-dodecanone, 9CI. 3-Hydroxy-2-hydroxymethyl-6-(7-oxododecyl)piperidine*C₁₈H₃₅NO₃ 313.479

Alkaloid obt. only as a mixt. with Isoprosopinine B, I-287.

(+)-form [38764-79-3]Alkaloid from roots and stem bark of *Prosopis africana* (Fabaceae). Cryst. (EtOAc). Mp 95° (isomeric mixt.). [α]_D +16 (CHCl₃).**(±)-form**

Synthetic. Mp 86-87°.

Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Belg.*, 1972, **81**, 425; 443 (*isol, struct, ms, abs config*)Khuong-Huu, Q. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 267 (*cmr*)Birkinshaw, T.N. *et al.*, *Tet. Lett.*, 1987, **28**, 813 (*synth, ir, pmr, cmr*)**Isoprosopinine B** I-287*12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-5-dodecanone, 9CI. 3-Hydroxy-2-hydroxymethyl-6-(8-oxododecyl)piperidine*

As Isoprosopinine A, I-286 with x = 7, y = 3

C₁₈H₃₅NO₃ 313.479

Alkaloid obt. only as a mixt. with Isoprosopinine A, I-286.

Natural-form [38764-80-6]Alkaloid from roots and stem bark of *Prosopis africana* (Fabaceae).**(±)-form**


Synthetic. Mp 90-91°.

[96894-88-1]

Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Belg.*, 1972, **81**, 425; 443 (*isol, ms, struct, abs config*)Khuong-Huu, Q. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 267 (*cmr*)Holmes, A.B. *et al.*, *Chem. Comm.*, 1985, 37 (*synth*)Birkinshaw, T.N. *et al.*, *Tet. Lett.*, 1987, **28**, 813 (*synth, ir, pmr, cmr*)**Isopsychotridine A** I-288

[112295-92-8]

[98985-58-1]



C₅₅H₆₂N₁₀ 863.162

Alkaloid from *Psychotria oleoides* (Rubiaceae). Amorph. [α]_D +66 (c, 1 in CHCl₃).

Stereoisomer (1): **Isopsychotridine B**

[112295-91-7]

C₅₅H₆₂N₁₀ 863.162

Alkaloid from *Psychotria oleoides* (Rubiaceae). Amorph. [α]_D +70 (c, 1 in CHCl₃). Tentative stereochem. assignments have been made.

Stereoisomer (2): **Isopsychotridine C**

Alkaloid D[†]

[129651-29-2]

C₅₅H₆₂N₁₀ 863.162

Alkaloid from leaves of *Psychotria forsteriana* (Rubiaceae). Exhibits potent cytotoxicity. Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D +183.3 (c, 0.05 in EtOH). λ_{max} 243 (ε 17780); 303 (ε 10230) (EtOH) (Berdy).

Roth, A. *et al.*, *Planta Med.*, 1985, 289 (*isol, uv, ir, cmr, ms, struct*)


Libot, F. *et al.*, *J. Nat. Prod.*, 1987, 50, 468-473 (*isol, uv, ir, pmr, ms, struct*)

Jannic, V. *et al.*, *J. Nat. Prod.*, 1999, 62, 838-843 (*stereochem, Isopsychotridine B*)

Isopsychotridine E

I-289

[168611-80-1]



C₅₅H₆₂N₁₀ 863.162


Alkaloid from *Calycodendron milnei*.

Shows analgesic props.

Saad, H.-E.A. *et al.*, *Phytochemistry*, 1995, 61, 313-316

Isopteleflorine

I-290



(S)-form

C₁₆H₁₇NO₅ 303.314

(S)-form [174513-97-4]

Alkaloid from *Orixa japonica*. Oil. [α]_D²⁰ -18.3 (c, 0.2 in CHCl₃). λ_{max} 255 (13000); 307 (1300); 325 (1100) (MeCN).

Ac: **O-Acetylisopteleflorine**

[174231-90-4]

C₁₈H₁₉NO₆ 345.351

Alkaloid from stems of *Orixa japonica*.

Oil. [α]_D²⁵ +6.4 (c, 0.7 in CHCl₃).

N-Me: **Hydroxyluninium(1+)**, O⁴-

Methylhydroxyluninium(1+)

[60593-04-6]

[70474-30-5 (perchlorate), 55812-57-2

(chloride)]

C₁₇H₂₀NO₅[⊕] 318.349

Alkaloid from the leaves of *Ptelea*

trifoliata (Rutaceae). Mp 225° (133-

135°) (as chloride) Mp 176-178° (as

perchlorate). [α]_D²⁰ -15.6 (c, 0.01 in

MeOH) (chloride).

(±)-form [134353-95-0]

Synthetic. Solid (Et₂O/diisopropyl ether).

Mp 163-166°.

N-Me:

Synthetic. Cryst. (MeOH/Et₂O) (as

iodide). Mp 159-161° (iodide).

(ξ)-form

Deoxy, N-Me: **O-Methyluninium**

[26605-52-7]

C₁₇H₂₀NO₄[⊕] 302.349

Alkaloid from the leaves of *Lunasia*

quercifolia (Rutaceae). Microcryst.

(MeOH/Et₂O) (as perchlorate). Mp

208-209° (perchlorate). [α]_D²⁴ -23.8 (c,

1.2 in MeOH) (perchlorate).

Deoxy, 1',2'-didehydro, N-Me: **Ptelefoli-**

donium(1+)

[59719-27-6]

C₁₇H₁₈NO₄[⊕] 300.333

Alkaloid from the leaves of *Ptelea*

trifoliata (Rutaceae).

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1966, 19,

2185-2187 (*O-Methyluninium*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1975, 38,

109-116 (*O-Methylhydroxyluninium, struct*)

Rideau, M. *et al.*, *Phytochemistry*, 1979, 18,

155-159 (*Hydroxyluninium, Ptelefolidonium*)

Petit-Paly, G. *et al.*, *Planta Med.*, 1989, 55,

209-210 (*Ptelefolidonium*)

Neville, C.F. *et al.*, *J.C.S. Perkin 1*, 1991, 259-

262 (*O-Methylhydroxyluninium, synth*)

Funayama, S. *et al.*, *Phytochemistry*, 1996, 41,

1231-1233 (*O-Acetylisopteleflorine*)

Noshita, T. *et al.*, *Biosci., Biotechnol., Biochem.*,

2001, 65, 710-713 (*isol, uv, pmr, cmr, ms*)


Boyd, D.R. *et al.*, *Chem. Comm.*, 2002, 3070-

3071 (*abs config*)

Isoptilocaluin

I-291

[78777-03-4]



Absolute Configuration

C₁₅H₂₅N₃ 247.383

Constit. of the Caribbean sponge *Ptilocaluis* aff. *Ptilocaluis spiculifer*. Shows potent antitumour props. and activity against gram-positive bacteria. Poorly sol. hexane. λ_{max} 225 (ε 10000) (MeOH) (Derep).


Harbour, G.C. *et al.*, *J.A.C.S.*, 1981, 103, 5604-5606 (*isol, pmr, cmr, ms*)

Yu, M. *et al.*, *J.O.C.*, 2008, 73, 9065-9074 (*synth, pmr, cmr*)

Isopyoverdin Pp BTP 1

I-292

[159325-01-6]



Absolute Configuration

C₄₄H₆₀N₁₂O₂₀ 1077.026

Isol. from *Pseudomonas putida* BTP 1.

Siderophore.


Jacques, P. *et al.*, *Z. Naturforsch., C*, 1995, 50, 622-629 (*isol, struct*)

Michalke, R. *et al.*, *Z. Naturforsch., C*, 1997, 52, 549-550 (*config*)

Isopyoverdin Pp CFML 90-33

I-293

[334009-11-9]



Absolute Configuration

C₄₉H₇₁N₁₃O₂₂ 1194.174

Isol. from *Pseudomonas putida* strain


CFML 90-33. Siderophore.

Sultana, R. *et al.*, *Tetrahedron*, 2001, 57, 1019-1023 (*isol, pmr, cmr, ms*)

Isopyoverdin Pp CFML 90-44

I-294

[349112-28-3]



Absolute Configuration

C₅₈H₈₇N₁₇O₂₆ 1438.424

Conts. 2 D-Ser and 2 L-Ser residues, posns. unknown. Isol. from *Pseudomonas putida* CFML 90-44. Siderophore.

Sultana, R. *et al.*, *Z. Naturforsch., C*, 2001, 56, 303-307

Isopyroine

I-295


[1399-97-9]

Struct. unknown. MF uncertain. Alka-

loid from *Isopyrum biternatum*. Cryst. (EtOH). Mp 160° Mp 255-257° (as hydrochloride).

Frankforter, G.B. *et al.*, *J.A.C.S.*, 1903, **25**, 99-102 (*isol*)

Isoapyruthaldine I-296
[307342-48-9]




C₄₄H₅₂N₂O₁₀ 768.902

Alkaloid from *Isopyrum thalictroides*. Amorph. yellow solid. λ_{max} 215 (log ε 6.72); 242 (log ε 6.66); 250 (sh) (log ε 6.59); 260 (sh) (log ε 6.44); 282 (sh) (log ε 6.56); 292 (log ε 6.65) (EtOH).

Istatkova, R.S. *et al.*, *Phytochemistry*, 2000, **54**, 959-964 (*isol, struct*)

Isoapyruthaline I-297
[189316-07-2]




C₄₂H₅₀N₂O₁₀ 742.864

Alkaloid from roots and rhizomes of *Isopyrum thalictroides*. Amorph. solid. [α]_D²⁵ +21 (c, 0.45 in CHCl₃). λ_{max} 208 (log ε 6.02); 232 (sh) (log ε 5.65); 283 (log ε 5.14) (EtOH).

Philipov, S.A. *et al.*, *Phytochemistry*, 1997, **44**, 1591 (*isol, uv, ir, pmr, ms, cd, struct*)

Isoapythaldine I-298
[307342-50-3]




C₄₃H₄₈N₂O₁₀ 752.86

Alkaloid from *Isopyrum thalictroides*. Amorph. yellow solid. λ_{max} 215 (log ε 6.64); 242 (log ε 6.61); 250 (sh) (log ε 6.54); 260 (sh) (log ε 6.4); 282 (sh) (log ε 6.48); 292 (log ε 6.56) (EtOH).

Istatkova, R.S. *et al.*, *Phytochemistry*, 2000, **54**, 959-964 (*isol, struct*)

Isoapythaline I-299
[189316-08-3]




C₄₀H₄₆N₂O₈ 682.812

From roots and rhizomes of *Isopyrum thalictroides*. Amorph. solid. [α]_D²⁵ -6 (c, 0.35 in CHCl₃). λ_{max} 207 (log ε 6.43); 236 (log ε 6.1); 260 (log ε 5.79); 283 (log ε 5.74) (EtOH).

Philipov, S.A. *et al.*, *Phytochemistry*, 1997, **44**, 1591

Isoquinocycline A I-300
PA 371γ. Antibiotic PA 371γ
[37332-17-5]



C₃₃H₃₄N₂O₁₀ 618.639

Anthracycline antibiotic. *Isol.* from *Streptomyces aureofaciens*. Active against gram-positive bacteria and mycobacteria. Orange powder (as hydrochloride). [α]_D²⁵ +112 (c, 1 in AcOH) (hydrochloride). Furumai *et al* gave orig. sugar moiety as L-lyxo. This was later corrected to L-xylo. λ_{max} 231 (ε 55000); 260 (ε 19100); 292 (ε 10700); 430 (ε 12000) (0.1N HCl) (Derep). λ_{max} 245 (ε 52500); 280 (sh) (ε 15500); 500 (ε 15100) (0.1N NaOH) (Derep). λ_{max} 228 (ε 71800); 258 (ε 45000); 288 (sh) (ε 18300); 425 (ε 18900) (MeOH) (Derep).

1''-Ketone: **Isoquinocycline B**. γ-Activity X. PA 371ε. Antibiotic PA 371ε [37332-18-6]

C₃₃H₃₂N₂O₁₀ 616.623
From *Streptomyces aureofaciens*. Active against gram-positive bacteria and mycobacteria. Orange powder (as hydrochloride). [α]_D²⁵ +24 (MeOH) (hydrochloride).

2'-Epimer, 1''-ketone: **Quinocycline B**.

Kosinostat
[37231-76-8]
C₃₃H₃₂N₂O₁₀ 616.623
Prod. by *Micromonospora* sp. TP-A0468 and *Streptomyces aureofaciens*.

Dark yellow powder. Mp >° 155 dec. [α]_D²⁵ +116.7 (c, 0.11 in MeOH/HCl). Identity with *Kosinostat* most probable. Direct comparison not possible. λ_{max} 228 (log ε 4.76); 258 (log ε 4.09); 291 (log ε 4.09); 423 (log ε 4.2) (MeOH).

Celmer, C.D. *et al.*, *Antibiot. Annu.*, 1958, 484; 493 (*isol*)

Cosulich, D.B. *et al.*, *Tet. Lett.*, 1963, 453; 1964, 750 (*struct*)

Tulinsky, A. *et al.*, *J.A.C.S.*, 1964, **86**, 5368-5369 (*cryst struct*)


Matern, U. *et al.*, *Eur. J. Biochem.*, 1972, **29**, 1-4; 5 (*struct*)

Paulsen, H. *et al.*, *Chem. Ber.*, 1978, **111**, 869-878 (*synth*)

Fukushima, K. *et al.*, *CA*, 1979, **93**, 114889x (*ms*)

Furumai, T. *et al.*, *J. Antibiot.*, 2002, **55**, 128-133; 134-140; 2003, **56**, C1 (*isol, uv, pmr, cmr, ms, revised struct*)

Isoquinoline, 9CI, 8CI I-301
2-Benzazine. Benzopyridine. 2-Azaphthalene. FEMA 2978
[119-65-3]




C₉H₇N 129.161

Isol. from the upper parts of *Spigelia anthelmia* (Loganiaceae). Used for photometric detn. of Ni (λ_{max} 410 nm, ε 7900). Flavouring agent. Hygroscopic cryst. or oil. Sol. EtOH, Et₂O, Me₂CO, C₆H₆, d₄²⁰ 1.1. Mp 25.5-26°. Bp 242° Bp₄₀ 142°. pK_a 5.4 (25°).

▶ Fl. p. >107°. Severe eye and skin irritant. LD₅₀ (rat, orl) 360 mg/kg. LD₅₀ (rbt, skn) 590 mg/kg. NW6825000

Wagner, H. *et al.*, *Planta Med.*, 1986, 378 (*isol, uv, pmr, ms*)

6,7-Isoquinolinediol I-302
6,7-Dihydroxyisoquinoline
[48125-47-9]



C₉H₇NO₂ 161.16

7-Me ether: 6-Hydroxy-7-methoxyisoquinoline. 7-Methoxy-6-isoquinolinol C₁₀H₉NO₂ 175.187
Light brown solid. Mp 198-200°.

Di-Me ether: 6,7-Dimethoxyisoquinoline. **Backebergine**
[15248-39-2]

C₁₁H₁₁NO₂ 189.213
Alkaloid from *Backebergia militaris* (preferred genus name *Pachycereus*) (Cactaceae). Cryst. (Et₂O). Mp 93-94°.

Di-Me ether, hydrochloride: Cryst. + 3H₂O (dil. HCl). Mp 219° (208-210° dec.).

Di-Me ether, N-Me: 6,7-Dimethoxy-2-methylisoquinolinium(1+)

[23901-08-8]
[74046-23-4]
C₁₂H₁₄NO₂[⊕] 204.248
Quaternary alkaloid from the fruits of *Thalictrum revolutum* (Ranunculaceae). Fine needles (EtOH/hexane) (as chloride). Mp 185.5-186.5° (chloride).

Methylene ether, N-Me: see N-Me, in 6,7-Methylenedioxyisoquinoline, M-446

Birch, A.J. *et al.*, *J.C.S. Perkin I*, 1974, 2185 (synth)

Wehrli, P.A. *et al.*, *Synthesis*, 1974, 288 (synth)
Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252 (cmr)

Wu, J. *et al.*, *J. Nat. Prod.*, 1980, **43**, 270 (di-Me ether N-Me, isol, uv, pmr, ms, struct)

Boger, D.L. *et al.*, *Tetrahedron*, 1981, **37**, 3977 (synth)

Ferrigni, N.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 839 (isol, ms, synth)

Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1989, **28**, 2833 (pmr, cmr)

Kuczniarz, R. *et al.*, *Synth. Commun.*, 1999, **29**, 1617-1625 (7-Me ether)

7,8-Isoquinolinediol, 9CI I-303

7,8-Dihydroxyisoquinoline
[142913-22-2]
[142913-21-1 (hydrobromide)]

C₉H₇NO₂ 161.16
Solid; yellow needles + ½H₂O (as hydrobromide). Mp 243-245° (hydrobromide).

Di-Me ether: 7,8-Dimethoxyisoquinoline.

Isobackebergine

[16503-95-0]
C₁₁H₁₁NO₂ 189.213

Alkaloid from *Backebergia militaris* (preferred genus name *Pachycereus*) (Cactaceae). Oil. Bp₁ 140° Bp_{0.15} 107-109°.

Di-Me ether, picrate:

Wispy needles + 1EtOH (EtOH). Mp 204°.

1,2,3,4-Tetrahydro: see 1,2,3,4-Tetrahydro-7,8-dihydroxyisoquinoline, T-159

Djerassi, C. *et al.*, *J.O.C.*, 1956, **21**, 975 (synth)

Bevis, M.J. *et al.*, *Tetrahedron*, 1969, **25**, 1585; 1971, **27**, 1253 (synth)


Ferrigni, N.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 839 (isol, ms, synth)

Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1989, **28**, 2833 (pmr, cmr)

U.S. Pat., 1993, 5 194 433; *CA*, **117**, 111380c (synth, ir, pmr)

1,5,6,7-Isoquinolinetetrol I-304

5,6,7-Trihydroxy-1(2H)-isoquinolinone.
1,5,6,7-Tetrahydroxyquinoline



C₉H₇NO₄ 193.159

NH-form

Tri-Me ether, N-Me: 5,6,7-Trimethoxy-2-methyl-1(2H)-isoquinolinone. **Thalactamine**
[23434-97-1]

C₁₃H₁₅NO₄ 249.266

Alkaloid from the above-ground parts of *Thalictrum minus* and the leaves of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Needles (petrol), plates (MeOH). Mp 112-114°.

3,4-Dihydro, 6,7-di-Me ether, N-Me: 3,4-Dihydro-5-hydroxy-2-methyl-1(2H)-isoquinolinone. **5-Hydroxy-N-methylcorydaline**

C₁₂H₁₅NO₄ 237.255

Alkaloid from the leaves of *Thalictrum delavayi*. Plates (CHCl₃). Mp 230-231°.

3,4-Dihydro, tri-Me ether, N-Me: 3,4-Dihydro-5,6,7-trimethoxy-2-methyl-1(2H)-isoquinolinone. **N-Methylthalidaldine**

[23434-99-3]

C₁₃H₁₇NO₄ 251.282

Trace alkaloid from *Thalictrum fendleri* (whole plant) (Ranunculaceae). Mp 104-106°. Thalidaldine appears to be unknown as an alkaloid.

Mollov, N.M. *et al.*, *Tet. Lett.*, 1969, 1951

(*Thalactamine*, uv, ir, pmr, ms, struct)

Shamma, M. *et al.*, *Tetrahedron*, 1971, **27**, 727 (*N-Methylthalidaldine*)

Duchevska, Kh. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1975, **8**, 134; *CA*, **84**, 150809a

(*Thalactamine*, synth)

Belgaonkar, V.H. *et al.*, *J. Het. Chem.*, 1978, **15**, 257 (*Thalactamine*, synth, uv, ir, pmr)


Mashekar, U.C. *et al.*, *Indian J. Chem., Sect. B*, 1979, **18**, 301 (*Thalactamine*, synth)

Başer, K.H.C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 704 (*Thalactamine*, isol, uv, ir, pmr, ms)

Wang, Y. *et al.*, *Zhivwu Xuebao (Acta Bot. Sin.)*, 2003, **45**, 500-502 (5-Hydroxy-N-methylcorydaline)

1,6,7-Isoquinolinetriol I-305

6,7-Dihydroxy-1(2H)-isoquinolinone,
9CI. 1,6,7-Trihydroxyisoquinoline



C₉H₇NO₃ 177.159

6-Me ether: 7-Hydroxy-6-methoxy-1(2H)-isoquinolinone. 1,7-Dihydroxy-6-methoxyisoquinoline. **N-Demethyl-doryphornine**

[711006-89-2]

C₁₀H₉NO₃ 191.186

Alkaloid from the roots of *Menispermum dauricum*. Rods (MeOH). Mp 257-259°. λ_{max} 243 (log ε 3.24); 279 (log ε 1.14); 290 (log ε 1.17); 325 (log ε 0.83) (MeOH).

6,7-Di-Me ether: 1-Hydroxy-6,7-dimethoxyisoquinoline. 6,7-Dimethoxy-1(2H)-isoquinolinone

[16101-63-6]

C₁₁H₁₁NO₃ 205.213

Cryst. (CHCl₃/C₆H₆). Mp 244-245° (237°). λ_{max} 246 (log ε 4.26); 253 (sh) (log ε 4); 268 (log ε 3.45); 278 (log ε 3.48); 290 (log ε 3.51); 310 (log ε 3.25); 322 (log ε 3.35); 335 (log ε 3.24) (EtOH).

Methylene ether: 1,3-Dioxolo[4,5-g]isoquinolin-5(6H)-one, 9CI. 1-Hydroxy-6,7-methylenedioxyisoquinoline. 3,4-Dehydronoroxyhydrastinine. 6,7-Methylenedioxyisocarbostyryl. 6,7-Methylenedioxy-1(2H)-isoquinolinone

[24188-76-9]

C₁₀H₇NO₃ 189.17

Alkaloid from *Corydalis bungeana* and *Thalictrum rugosum*. Cryst. (CHCl₃ or EtOH); needles (C₆H₆/Me₂CO). Mp 278° (268-270°). NH-form predominates.

3,4-Dihydro: 3,4-Dihydro-1,6,7-isoquinolinetriol. 3,4-Dihydro-1,6,7-trihydroxyisoquinoline. 3,4-Dihydro-6,7-dihydroxy-1(2H)-isoquinolinone. **Pericampylinone A**

Iseluxine

[330847-76-2]

C₉H₉NO₃ 179.175

Alkaloid from *Aristolochia elegans* and *Iseia luxurians*. Pale yellow needles (CHCl₃). Mp 229-230°. λ_{max} 220 (log ε 4.23); 267 (log ε 3.75); 304 (log ε 3.69) (MeOH).

3,4-Dihydro, 6-Me ether: 3,4-Dihydro-7-hydroxy-6-methoxy-1(2H)-isoquinolinone. 3,4-Dihydro-6-methoxy-1,7-isoquinolinediol. **Northalfoline**

[157669-72-2]

C₁₀H₁₁NO₃ 193.202

Alkaloid from pedicels of *Lindera megaphylla* (Lauraceae). Rods (MeOH/Me₂CO). Mp 222-224°.

3,4-Dihydro, 6,7-di-Me ether: 3,4-Dihydro-6,7-dimethoxy-1(2H)-isoquinolinone. 3,4-Dihydro-6,7-dimethoxy-1-isoquinolinol. 3,4-Dihydro-6,7-dimethoxyisocarbostyryl. **Corydaline**

[493-49-2]

C₁₁H₁₃NO₃ 207.229

Alkaloid from *Berberis baluchistanica* and *Enantia polycarpa* (Berberidaceae, Annonaceae). Needles (butanone), prisms (C₆H₆). Mp 174-175°.

3,4-Dihydro, methylene ether: 7,8-Dihydro-1,3-dioxolo[4,5-g]isoquinolin-5(6H)-one. **Noroxyhydrastinine**. 3,4-Dihydro-6,7-methylenedioxy-1(2H)-isoquinolinone

[21796-14-5]

C₁₀H₉NO₃ 191.186

Alkaloid from *Fumaria parviflora*, *Corydalis ophiocarpa*, and from the roots of *Thalictrum alpinum*, *Thalictrum minus* var. *adanthifolium*, *Thalictrum rugosum* and *Thalictrum foliolosum* (Papaveraceae, Ranunculaceae). Cryst. (MeOH or C₆H₆). Mp 182-183° Mp 187-187.5°.

NH-form

O⁶,N-Di-Me: 7-Hydroxy-6-methoxy-2-methyl-1(2H)-isoquinolinone. **Doryphornine**

[54302-47-5]

C₁₁H₁₁NO₃ 205.213

Alkaloid from the bark of *Doryphora sassafras* (Monimiaceae). Needles (CHCl₃). Mp 215-217°.

Di-Me ether, N-Me: 6,7-Dimethoxy-2-methyl-1(2H)-isoquinolinone. 6,7-Dimethoxy-2-methylisocarbostyryl. N-

Methyl-6,7-dimethoxyisoquinolone
[20323-75-5]

C₁₂H₁₃NO₃ 219.24

Alkaloid from *Hernandia ovigera*, *Stephania sasakii*, and from the roots of *Thalictrum alpinum* and *Thalictrum isopyroides* (Hernandiaceae, Menispermaceae, Ranunculaceae). Needles (MeOH/CHCl₃). Mp 112-113° (98-100°). λ_{max} 245 (log ε 3.8); 270 (log ε 3.11); 280 (log ε 3.18); 290 (log ε 3.18); 335 (log ε 2.9) (MeOH).

Methylene ether, N-Me: Doryanine. 6-Methyl-1,3-dioxolo[4,5-g]isoquinolin-5(6H)-one. 2-Methyl-6,7-methylenedioxy-1(2H)-isoquinolinone
[24880-42-0]

C₁₁H₉NO₃ 203.197

Alkaloid from leaves and bark of *Doryphora sassafras* (Monimiaceae). Needles (Me₂CO/petrol or C₆H₆/hexane). Mp 162-163°. λ_{max} 230 (log ε 4.45); 250 (log ε 4.51); 255 (log ε 3.85); 295 (log ε 3.91); 325 (log ε 3.67) (MeOH).

Methylene ether, N-(2-acetyl-5,6-methylenedioxybenzyl): N-(2-Acetyl-5,6-methylenedioxybenzyl)-6,7-methylenedioxy-1(2H)-isoquinolinone
[813466-13-6]

C₂₀H₁₅NO₆ 365.342

Alkaloid from *Corydalis bungeana*. Cryst. (MeOH).

3,4-Dihydro, 6-Me ether, N-Me: Thalifoline
[21796-15-6]

C₁₁H₁₃NO₃ 207.229

Alkaloid from the roots of *Thalictrum minus* var. *adiantifolium*, and from the leaves, stems, roots and fruit of *Cryptocarya longifolia* (Ranunculaceae, Lauraceae). Prisms (MeOH). Mp 210-211° Mp 220°.

3,4-Dihydro, di-Me ether, N-Me: 3,4-Dihydro-6,7-dimethoxy-2-methyl-1(2H)-isoquinolinone. N-Methylcorydaldine
[6514-05-2]

C₁₂H₁₅NO₃ 221.255

Alkaloid from *Hernandia ovigera*, *Papaver bracteatum*, *Papaver urbanianum* and *Thalictrum fendleri* (Hernandiaceae, Papaveraceae, Ranunculaceae). Also obt. by degradn. of Warifteine, W-9, Thaliracebine, T-332 and Adiantifoline, A-139. Cryst. (MeOH, EtOH or petrol). Mp 125-126° (120.5°).

3,4-Dihydro, methylene ether, N-Me: 7,8-Dihydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5(6H)-one. 3,4-Dihydro-2-methyl-6,7-methylenedioxy-2(1H)-isoquinolinone. Oxyhydrastinine
[552-29-4]

C₁₁H₁₁NO₃ 205.213

Alkaloid from *Fumaria schleicheri*, *Argemone mexicana* and *Papaver dubium* var. *glabrum* (Papaveraceae). Cryst. (petrol). Mp 98°. λ_{max} 207

(log ε 4.39); 217 (log ε 4.47); 222 (log ε 4.46); 262 (log ε 3.84); 303 (log ε 3.9) (MeOH).

Gharbo, S.A. et al., *J. Nat. Prod.*, 1965, **28**, 237-244 (*Doryanine*)

Cava, M.P. et al., *Tet. Lett.*, 1966, 4279-4282 (*N-Methylcorydaldine*)

Eloy, F. et al., *Helv. Chim. Acta*, 1969, **52**, 1755-1762 (*methylene ether, synth*)

Doskotch, R.W. et al., *J. Nat. Prod.*, 1969, **32**, 29-35 (*isol, uv*)

Mollov, N.M. et al., *Tet. Lett.*, 1969, 1951-1952 (*tri-Me, uv, ir, pmr*)

Doskotch, R.W. et al., *Tetrahedron*, 1969, **25**, 469-475 (*Thalifoline, isol, uv, ir, pmr, ms, struct, synth*)

Shamma, M. et al., *Tetrahedron*, 1971, **27**, 727-733 (*N-Methylcorydaldine, isol, ir, pmr, ms, struct*)

Chen, C.R. et al., *J. Nat. Prod.*, 1974, **37**, 493 (*Doryphorine, Doryanine*)

Shamma, M. et al., *J.A.C.S.*, 1974, **96**, 7809-7811 (*Corydaldine, occur*)

Jackson, A.H. et al., *J.C.S. Perkin I*, 1974, 1911-1920 (*Corydaldine, N-Methylcorydaldine, synth, pmr*)

Markosyan, S.S. et al., *Arm. Khim. Zh.*, 1976, **29**, 1053-1057; *CA*, **86**, 136375j (*Oxyhydrastinine*)

Passannanti, S. et al., *J. Het. Chem.*, 1977, **14**, 103-106 (*methylene ether, synth, ir, pmr*)

Belgaonkar, V.H. et al., *J.C.S. Perkin I*, 1977, 702-706 (*Doryanine, synth*)

Theuns, H.G. et al., *Phytochemistry*, 1977, **16**, 753-755 (*N-Methylcorydaldine, ms*)

Belgaonkar, V.H. et al., *J. Het. Chem.*, 1978, **15**, 257-261 (*tri-Me, synth, uv, ir, pmr*)

Abduzhabbarova, S. et al., *Khim. Pri. Soedin.*, 1978, **14**, 472-474; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 400-402 (*tri-Me, isol, uv, ir, pmr, ms*)

Shamma, M. et al., *Carbon-13 NMR Shift Assignments of Amines and Alkaloids*, Plenum Press, N.Y., 1979, 116 (*cmr*)

Wu, W.-N. et al., *J. Nat. Prod.*, 1980, **43**, 372-381 (*tri-Me, Noroxyhydrastinine, isol, uv, ir, pmr, ms*)

Kondo, Y. et al., *J.C.S. Perkin I*, 1980, 911-918 (*6,7-di-Me ether, synth, ir, pmr, uv*)

Bick, I.R.C. et al., *Aust. J. Chem.*, 1981, **34**, 195-207 (*Thalifoline, isol, cryst struct*)

Modi, A.R. et al., *Indian J. Chem., Sect. B*, 1981, **20**, 813-815 (*tri-Me, synth*)

Chattopadhyay, S.K. et al., *J. Nat. Prod.*, 1981, **44**, 45-49 (*Noroxyhydrastinine*)

Castedo, L. et al., *Tet. Lett.*, 1981, **22**, 2233-2236 (*Doryanine, synth*)

Kunitomo, J. et al., *Yakugaku Zasshi*, 1981, **101**, 431-436; *CA*, **95**, 204236c (*tri-Me, isol*)

Hanaoka, M. et al., *Chem. Pharm. Bull.*, 1982, **30**, 2793-2796 (*Noroxyhydrastinine, synth, ir, pmr, ms*)

Joshi, V. et al., *Indian J. Chem., Sect. B*, 1983, **22**, 65-66 (*Corydaldine, N-Methylcorydaldine, synth*)

Hussain, S.F. et al., *Phytochemistry*, 1983, **22**, 319-320 (*Oxyhydrastinine*)

Thakur, P. et al., *J. Indian Chem. Soc.*, 1991, **68**, 526-528 (*Doryanine, synth*)

Lee, A.W.M. et al., *J.C.S. Perkin I*, 1992, 309-310 (*N-Methylcorydaldine, Oxyhydrastinine, Doryanine, synth*)

Chou, C.-J. et al., *J. Nat. Prod.*, 1994, **57**, 689-694 (*Northalifoline, synth, isol, pmr, cmr, ir, uv*)

Schimming, T. et al., *Z. Naturforsch., C*, 2000, **55**, 1023-1025 (*Isehixine*)

Wu, T.-S. et al., *J. Nat. Prod.*, 2002, **65**, 1522-1525 (*Pericampylinone A*)

Wang, Y.-C. et al., *Synthesis*, 2002, 2187-2190 (*Thalifoline, synth, pmr, cmr, ms*)

Zhang, X. et al., *Phytochemistry*, 2004, **65**, 929-932 (*N-Demethyldoryphorine*)

Xie, C. et al., *Phytochemistry*, 2004, **65**, 3041-3047 (*N-acetylmethylenedioxybenzyl deriv*)

1,6,8-Isoquinolinetriol I-306

1,6,8-Trihydroxyisoquinoline. 6,8-Dihydroxy-1(2H)-isoquinolinone, 9CI. Siamine B
[92446-27-0]

C₉H₇NO₃ 177.159

Alkaloid from the leaves of *Cassia siamea* (Fabaceae). Mp 235° (as hydrochloride).

El-Sayyad, S.M. et al., *J. Nat. Prod.*, 1984, **47**, 708 (*isol, uv, ir, pmr, ms, struct*)

5,6,7-Isoquinolinetriol I-307

5,6,7-Trihydroxyisoquinoline
[120092-56-0]

C₉H₇NO₃ 177.159

CAS no. refers to the iodide.

Tri-Me ether, 5,6,7-Trimethoxyisoquinoline. Isonortehaunine
[36982-71-5]

C₁₂H₁₃NO₃ 219.24

Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae). Oil. Mp 180-181° (as picrate).

Tri-Me ether, N-Me: N-Methyl-5,6,7-trimethoxyisoquinolinium(1+)
C₁₃H₁₆NO₃⁺ 234.274

Quaternary alkaloid from *Isopyrum thalictroides* (Ranunculaceae).

3,4-Dihydro, tri-Me ether: 3,4-Dihydro-5,6,7-trimethoxyisoquinoline. Dehydronortehaunine
[93474-25-0]

C₁₂H₁₅NO₃ 221.255

Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).

1,2,3,4-Tetrahydro, tri-Me ether, N-Me: see 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-2-methylisoquinoline, T-238

Birch, A.J. et al., *J.C.S. Perkin I*, 1974, 2185 (*5,6,7-tri-Me ether*)


Boger, D.L. et al., *Tetrahedron*, 1981, **37**, 3977 (*5,6,7-tri-Me ether*)

Roush, R.A. et al., *Anal. Chem.*, 1985, **57**, 109

Kostalova, D. et al., *Chem. Pap.*, 1988, **42**, 841; *CA*, **110**, 170219k

1-(8-Isoquinolinylmethyl)pyridinium I-308

Sibiridine
[709612-22-6]




C₁₅H₁₃N₂⁺ 221.281

Quaternary alkaloid from *Nitraria schoberi* and *Nitraria sibirica*. Cryst. (EtOH) (as hydroxide). Mp 192-193° (hydroxide).

λ_{\max} 220 (log ϵ 4.1); 287 (log ϵ 3.51 broad); 307 (log ϵ 3.46); 320 (log ϵ 3.54) (EtOH) (hydroxide).

Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2003, **39**, 292-293 (*isol, uv, ir, ms*)

Isorhazicine**I-309**


$C_{21}H_{24}N_2O_4$ 368.432

A 2,3-secoajmaline. This struct. may be incorrect. The struct. of Rhazicine has been revised. See under Lanceomigine, L-28. Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Hygroscopic needles. $[\alpha]_D^{25} +61.4$ (CHCl₃).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1987, **26**, 405 (*isol, uv, ir, pmr, cmr, ms, struct*)

Isorubijervine**I-310**

Solanid-5-ene-3,18-diol
[468-45-1]



Absolute configuration

$C_{27}H_{43}NO_2$ 413.642

Alkaloid from *Veratrum album*, *Veratrum californicum*, *Veratrum eschscholtzii* and *Veratrum grandiflorum* (Liliaceae). Shows antifungal activity. Mp 241-244°. $[\alpha]_D^{25} +9.2$ (EtOH).

O^3 - β -D-Glucopyranoside: **Isorubijervosine**
[468-46-2]

$C_{33}H_{53}NO_7$ 575.784

Alkaloid from *Veratrum viride* and *Veratrum eschscholtzii* (Liliaceae). Fine needles (MeOH). Mp 279-280°. $[\alpha]_D^{25} -20$ (c, 1.45 in Py).

18-Ac: Mp 219°.

Di-Ac: Mp 202°.

12 β -Hydroxy: **Rubivirine**. 12-Hydroxyisorubijervine

[164178-46-5]

$C_{27}H_{43}NO_3$ 429.642

Alkaloid from roots and rhizomes of *Veratrum viride* (Liliaceae). Needles (MeOH). Mp 239-241°. $[\alpha]_D^{25} +28.5$ (c, 0.1 in CHCl₃).

Klohs, M.W. *et al.*, *J.A.C.S.*, 1953, **75**, 2133 (*isol, ir*)

Höhne, E. *et al.*, *Tetrahedron*, 1966, **22**, 673 (*cryst struct*)

Wolters, B. *et al.*, *Planta Med.*, 1970, **19**, 189-193 (*activity*)


Keeler, R.F. *et al.*, *Phytochemistry*, 1974, **13**, 2336 (*isol*)

Kaneko, K. *et al.*, *Tet. Lett.*, 1978, 4801 (*isol*)

El Sayed, K.A. *et al.*, *Phytochemistry*, 1995, **38**, 1547 (*Rubivirine*)

Isosalviamine A

[878475-29-7]

I-311

R = H

$C_{19}H_{13}NO_2$ 287.317

Alkaloid from the roots of *Salvia trijuga*. Amorph. yellow powder (Me₂CO). Mp 178-180°. λ_{\max} 239 (log ϵ 3.61); 258 (sh) (log ϵ 3.74); 267 (log ϵ 3.93); 276 (log ϵ 4); 282 (sh) (log ϵ 3.8); 290 (sh) (log ϵ 3.53); 303 (log ϵ 3.47); 311 (sh) (log ϵ 3.11); 326 (log ϵ 2.55); 342 (log ϵ 2.64); 359 (log ϵ 2.57) (MeOH).

Lin, F.-W. *et al.*, *Heterocycles*, 2006, **68**, 159-165 (*isol, uv, pmr, cmr, ms*)

Isosalviamine B**I-312**

[878475-30-0]

As Isosalviamine A, I-311 with

R = CH₃


$C_{20}H_{15}NO_2$ 301.344

Alkaloid from the roots of *Salvia trijuga*. Pale yellow needles (Me₂CO). Mp 175-177°. λ_{\max} 259 (log ϵ 3.47); 267 (sh) (log ϵ 3.63); 276 (log ϵ 3.66); 283 (sh) (log ϵ 3.49); 289 (sh) (log ϵ 3.35); 303 (log ϵ 3.2); 310 (sh) (log ϵ 2.91); 361 (log ϵ 2.15) (MeOH).

Lin, F.-W. *et al.*, *Heterocycles*, 2006, **68**, 159-165 (*isol, uv, pmr, cmr, ms*)

Isoschizogamine**I-313**

[2779-07-9]



Absolute Configuration

$C_{21}H_{24}N_2O_3$ 352.432

Struct. revised in 1998. Prob. related biogenetically to Schizozygine, S-137. Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Cryst. (EtOAc or petrol). Mp 184-185° (181.5-184.5°). $[\alpha]_D^{25} -241$ (c, 0.8 in CHCl₃).

10-Demethoxy: **Isoschizogaline**

[2671-29-6]

$C_{20}H_{22}N_2O_2$ 322.406

Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Cryst. (Et₂O). Mp 110-112°. $[\alpha]_D^{25} -262.3$ (c, 1 in CHCl₃). Struct. revised in 2002.

Hajicek, J. *et al.*, *Tet. Lett.*, 1998, **39**, 505-508 (*pmr, cmr, struct*)

Hubbs, J.L. *et al.*, *Org. Lett.*, 1999, **1**, 1315-1317 (*synth*)


Kariba, R.M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 566-569 (*Isoschizogaline*)

Zhou, J. *et al.*, *J.O.C.*, 2007, **72**, 3808-3815 (*synth, abs config*)

Stephens, P.J. *et al.*, *Chirality*, 2008, **20**, 454-470 (*cd, abs config*)

Isoschumanniphytine**I-314**

[96889-79-1]



$C_{16}H_9NO_5$ 295.251

Struct. revised in 1987. Alkaloid from the root bark of *Schumanniphyton magnificum* (Rubiaceae). Anti-HIV agent. Yellow cryst. (CHCl₃). λ_{\max} 226 (ϵ 12880); 245 (ϵ 9120); 260 (ϵ 9120); 272 (ϵ 3630); 314 (ϵ 954) (MeOH) (Berdy).

N-Me: **N-Methylisoschumanniphytine**

$C_{17}H_{12}NO_5^{\oplus}$ 310.285


Quaternary alkaloid from *Schumanniphyton magnificum* (Rubiaceae). Anti-HIV agent. λ_{\max} 204 ; 225 ; 236 ; 355 (MeOH) (Berdy).

Houghton, P.J. *et al.*, *Planta Med.*, 1985, 23-27; 1987, **53**, 262-263; 264-266 (*isol, uv, ir, pmr, ms, struct, N-Me*)

Kelly, T.R. *et al.*, *J.O.C.*, 1992, **57**, 1593-1597 (*synth, pmr, struct*)

Isoscoulerine**I-315**

5,8,13,13a-Tetrahydro-2,10-dimethoxy-6H-dibenzo[a,g]quinolizine-3,9-diol, 9CI. 3,9-Dihydroxy-2,10-dimethoxytetrahydroprotoberberine



(S)-form

$C_{19}H_{21}NO_4$ 327.379

(S)-form [62249-73-4]

Alkaloid from *Stephania hainanensis*.

N-Me(cis-): **Steponine**

[17132-69-3]

[17132-70-6]

$C_{20}H_{24}NO_4^{\oplus}$ 342.414

Quaternary alkaloid from dried rhizomes of *Stephania japonica*, *Stephania sasakii* and *Stephania cepharantha* (Menispermaceae). Also isol. from roots of *Cissampelos mucronata*. Prisms (as chloride). Mp 235° dec. (chloride). $[\alpha]_D^{25} -129.9$ (chloride).


Di-Me ether: see Tetrahydropalmatine, T-212

(±)-form [62057-90-3]

Synthetic. Prisms (MeOH). Mp 218-221°. Tomita, M. *et al.*, *Yakugaku Zasshi*, 1957, **77**, 274-277; *CA*, **51**, 11361g (*Steponine*)
 Kunimoto, J. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 1010-1011; *CA*, **68**, 3051g (*Steponine*)
 Brochmann-Hanssen, E. *et al.*, *J.O.C.*, 1977, **42**, 3588-3591 (*synth, ir, pmr, ms*)
 Ohiri, F.C. *et al.*, *Planta Med.*, 1983, **49**, 162-164 (*pmr*)
 Fang, S. *et al.*, *Zhongcaoyao*, 1987, **18**, 146-149; *CA*, **107**, 93551w (*isol*)
 Tshibangu, J.N. *et al.*, *Phytochem. Anal.*, 2003, **14**, 13-22 (*Steponine, isol, hplc*)

Isosogoline A**I-316**

[117694-97-0]




Relative Configuration

$C_{23}H_{19}N_3O_3$ 385.421
 Alkaloid from the Red Sea tunicate *Eudistoma* sp. $[\alpha]_D^{24}$ -660 (c, 0.001 in $CHCl_3$). λ_{max} 248 (ϵ 13400); 297 (ϵ 25000); 368 (ϵ 6100); 388 (ϵ 7700); 550 (ϵ 7800) (MeOH/HCl) (*Derep*). λ_{max} 242 (ϵ 13900); 274 (ϵ 19100); 330 (ϵ 9400); 360 (ϵ 2300); 386 (ϵ 3100); 470 (ϵ 3500) (MeOH) (*Derep*).

Rudi, A. *et al.*, *J.O.C.*, 1989, **54**, 5331 (*isol, uv, ir, pmr, cmr, cd, struct*)

Isosovanine**I-317**

1-(3-Hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline
 [70400-40-7]



$C_{18}H_{15}NO_4$ 309.321
 Trace alkaloid from *Hedycarya angustifolia* (Monimiaceae). Pale yellow needles (EtOH). Mp 148° (natural) Mp 182-183° (synthetic).

O-De-Me, O^{3'}, N-di-Me: **Escholamidine**
 [55861-73-9]

$C_{19}H_{18}NO_4^{\oplus}$ 324.356
 Quaternary alkaloid from *Eschscholtzia oregana*. Small prisms (H_2O) (as iodide). Mp 214-215° (iodide). Cryst. dec. and turn red-brown in air.


Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1975, **40**, 1095 (*Escholamidine*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1985, **38**, 1571 (*cryst struct*)

Chantimakorn, V. *et al.*, *Aust. J. Chem.*, 1989, **42**, 209 (*synth, uv, ir, pmr, ms*)

Isosinomenine**I-318**

5,6-Didehydro-4-hydroxy-3,6-dimethoxy-17-methylmorphinan-7-one, 9CI, 8,14-Dihydrosalutaridine

**(+)-form** $C_{19}H_{23}NO_4$ 329.395**(+)-form** [510-42-9]

Poss. minor alkaloid from *Sinomenium acutum*. Cryst. (Et_2O). Mp 198-202°. $[\alpha]_D$ +73 (c, 1.2 in EtOH).

14-Epimer: Ocobotrine

[60761-55-9]

 $C_{19}H_{23}NO_4$ 329.395

Alkaloid from *Ocotea brachybotra* (Lauraceae). Cryst. + $\frac{1}{2}$ EtOAc (EtOAc/ Et_2O). Mp 97-99°. $[\alpha]_D^{20}$ -93 (c, 1 in $CHCl_3$). λ_{max} 210 ($\log \epsilon$ 4.52); 264 ($\log \epsilon$ 4.01) (MeOH).

14-Epimer, hydrochloride: Cryst. (MeOH). Mp 296-297°.

(-)-form

Alkaloid from *Croton linearis*, *Croton balsamifera* and *Croton discolor* (Euphorbiaceae). Mp 198-203°. $[\alpha]_D^{15}$ -76.1 (c, 0.59 in MeOH). λ_{max} 206 (ϵ 32730); 230 (ϵ 6960); 265 (ϵ 7550) (EtOH).

Ac: O-Acetyldihydrosalutaridine $C_{21}H_{25}NO_5$ 371.432

From *Croton linearis* (Euphorbiaceae). Mp 210°. $[\alpha]_D^{15}$ -22.1 (MeOH).

N-De-Me: 8,14-Dihydronorsalutaridine $C_{18}H_{21}NO_4$ 315.368

Alkaloid from *Croton linearis* (Euphorbiaceae). Cryst. + EtOAc. Mp 208-212°. $[\alpha]_D^{15}$ -69.1 (MeOH). λ_{max} 206 (ϵ 35900); 235 (ϵ 8611); 261 (ϵ 8950) (EtOH).

14-Epimer: Milonine

[171485-61-3]

 $C_{19}H_{23}NO_4$ 329.395

Alkaloid from leaves of *Cissampelos sympodialis* (Menispermaceae). Reddish amorph. powder. Mp 74-76°. $[\alpha]_D^{25}$ +50 (c, ca. 0.6 in MeOH). It appears that Milonine should be the enantiomer of Ocobotrine, but the opt. rotns. do not support this assignment. λ_{max} 224; 262; 300 (MeOH).

Barton, D.H.R. *et al.*, *J.C.S.(C)*, 1968, 929-936 (**(+)-form, occur, synth**)

Haynes, L.J. *et al.*, *J.C.S.(C)*, 1968, 951-957 (**(-)-form, isol, struct, ms**)


Vecchiotti, V. *et al.*, *Tet. Lett.*, 1976, 1631-1634 (*Ocobotrine, uv, ir, pmr, ms, struct*)

De Freitas, M.R. *et al.*, *Phytochemistry*, 1995, **40**, 1553-1555 (*Milonine*)

Rasoanaivo, P. *et al.*, *Planta Med.*, 1998, **64**, 58-62 (*Ocobotrine, cmr*)

Isositsirikine**I-319**

[6519-27-3]



Absolute Configuration

 $C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from *Catharanthus roseus*, *Aspidosperma marcgravianum*, *Aspidosperma oblongum*, a number of *Strychnos* spp. and others. $[\alpha]_D$ -20 ($CHCl_3$). λ_{max} 224 ($\log \epsilon$ 4.55); 283 ($\log \epsilon$ 3.92); 291 ($\log \epsilon$ 3.84) (MeOH).

Sulfate (1:2): Mp 263.5°.

Picrate:

Yellow needles (MeOH). Mp 216°.

 β -N⁴-Oxide: (16R)-Isositsirikine (4S)-N-oxide

[88607-61-8]

 $C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from the stem bark of *Aspidosperma marcgravianum* (Apocynaceae). Amorph. $[\alpha]_D$ -56 (c, 0.54 in MeOH). Named 16-Epiisositsirikine 3S,4S-N-oxide in the original literature (confusing).

N⁴- α -Me: Diploceline

[69306-89-4]

 $C_{22}H_{29}N_2O_3^{\oplus}$ 369.483

Alkaloid from *Strychnos gossweileri* roots (Loganiaceae). Shows antiparasitic and weak antibacterial activity. 16-Config. not determined, may be the oxide of 16-Epiisositsirikine. λ_{max} 220 ($\log \epsilon$ 4.54); 270 ($\log \epsilon$ 3.86); 280 ($\log \epsilon$ 3.82); 289 ($\log \epsilon$ 3.68) (MeOH).

O-Ac: O-Acetylisositsirikine

[6519-40-0]

 $C_{23}H_{28}N_2O_4$ 396.485

Alkaloid from the roots and leaves of *Rhazya stricta* (Apocynaceae). $[\alpha]_D$ -2.5 (MeOH). λ_{max} 202 ($\log \epsilon$ 4.22); 282 ($\log \epsilon$ 3.27); 289 ($\log \epsilon$ 3.15) (MeOH).

9-Methoxy: Strychnorubigine. 9-Methoxy-(16R)-E-isositsirikine

[72994-76-4]

 $C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the root bark of *Strychnos rubiginosa* (Loganiaceae) and from *Strychnos lucens* and *Strychnos madagascariensis*. Cryst. (EtOAc/hexane). Mp 126-127° dec. $[\alpha]_D$ +53 (c, 0.17 in $CHCl_3$). λ_{max} 227 ($\log \epsilon$ 4.68); 274 (sh) ($\log \epsilon$ 3.91); 284 (sh) ($\log \epsilon$ 3.84); 293 ($\log \epsilon$ 3.84) (EtOH).

9-Methoxy, O-Ac:

Cryst. (EtOAc/hexane). Mp 96-97°.

11-Methoxy: Hervine

[16049-27-7]

 $C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Vinca herbacea* (Apocynaceae). Cryst. (EtOH). Mp 173-175°. $[\alpha]_D$ -93 (c, 0.20 in EtOH). 16-

Config. not determined, may be 11-methoxy-16-epiisositsirikine. λ_{\max} 228 (log ϵ 4.45); 270 (log ϵ 3.6); 290 (log ϵ 3.7) (EtOH).

19Z-Isomer: 19Z-Isositsirikine

[80287-17-8]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the leaves of *Alstonia sphaerocapitata* and from *Catharanthus roseus* cell suspension cultures (Apocynaceae). Mp 183°. $[\alpha]_D^{25}$ -32 (c, 1 in $CHCl_3$). λ_{\max} 227; 283; 290 (MeOH).

3-Epimer: Rhazimanine

[119618-47-2]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the fruits of *Rhazya stricta* (Apocynaceae). Amorph. $[\alpha]_D^{26}$ 0 ($CHCl_3$). May be identical with Bhimberine below. Unstable to air and light. λ_{\max} 222; 268 (sh); 273 (sh); 282; 290 (MeOH).

3-Epimer, 10-hydroxy, 19,20 β -dihydro: 19,20-Dihydro-10-hydroxyisositsirikine

[916334-46-8]
 $C_{21}H_{28}N_2O_4$ 372.463
 Alkaloid from the stems of *Ervatamia yunnanensis*.

16-Epimer: 16-Epiisositsirikine. (16S)-Isositsirikine

[60031-91-6]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from *Alstonia cuspa*, *Alstonia marcgravianum*, *Alstonia oblongum*, *Peschiera echinata*, *Pterotaberna inconspicua*, *Tabernaemontana psorocarpa*, *Tabernaemontana pachysiphon*, *Tabernaemontana chippii*, *Tabernaemontana ventricosa* and a number of *Strychnos* spp. Cryst. (EtOAc). Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 186-187° (182-184° dec.). $[\alpha]_D^{28}$ -205 (c, 0.2 in $CHCl_3$) (-145). λ_{\max} 226 (ϵ 31620); 285 (ϵ 7940); 293 (ϵ 6310) (EtOH).

16-Epimer, β -N⁴-oxide: (16S)-Isositsirikine (4S)-N-oxide

[88607-62-9]
 $C_{21}H_{26}N_2O_4$ 370.447
 Alkaloid from the root bark of *Alstonia marcgravianum* (Apocynaceae). Amorph. $[\alpha]_D$ -56 (c, 0.54 in MeOH).

16-Epimer, N⁴- α -Me: 16-Epidiploceline

[75364-53-3]
 $C_{22}H_{29}N_2O_3$ 369.483
 Alkaloid from *Strychnos gossweileri* roots. 16-Config. not certain, may be as in Isositsirikine.

16-Epimer, N¹-hydroxymethyl: N¹-Hydroxymethyl-(16S)-isositsirikine

$C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from the roots of *Rauwolfia yunnanensis*. Amorph. yellow solid. $[\alpha]_D^{24}$ +48 (c, 0.23 in Py). λ_{\max} 225 (log ϵ 4.46); 275 (log ϵ 3.81); 280 (log ϵ 3.81); 291 (log ϵ 3.7); 347 (log ϵ 2.75); 363 (log ϵ 2.78) (MeOH).

16-Epimer, 9-methoxy: 9-Methoxy-

(16S)-E-isositsirikine

[149250-44-2]
 $C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from the root bark of *Strychnos lucens* and *Strychnos mada-gascariensis*. $[\alpha]_D$ -108 (c, 0.11 in MeOH). λ_{\max} 225; 254; 351 (MeOH).

16-Epimer, 19Z-isomer: 16-Epi-Z-isositsirikine

[80287-16-7]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the leaves of *Catharanthus roseus* and *Rhazya stricta* (Apocynaceae). Shows *in vivo* antineoplastic activity vs. human nasopharynx KP and mouse P388 tumour cells. Mp 181°. $[\alpha]_D^{25}$ -41 (c, 0.1 in MeOH).

3,16-Diepimer: Bhimberine

[102043-86-7]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the fruits and leaves of *Rhazya stricta* (Apocynaceae). Amorph. Sensitive to light and air. Evidence has been presented (Lounasmaa *et al*) that Bhimberine has the (16R)-config. and is identical with Rhazimanine. λ_{\max} 222 (log ϵ 4.2); 268 (sh) (log ϵ 3.61); 273 (sh) (log ϵ 3.61); 282 (log ϵ 3.61); 290 (log ϵ 3.55) (MeOH).

Kutney, J.P. *et al.*, *Tetrahedron*, 1966, **22**, 321-336 (struct, uv, ir, ms, pmr)

Ogyanov, I. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 754-758 (Hervine)

Simões, J.C. *et al.*, *Phytochemistry*, 1976, **15**, 543-544 (16-Epiisositsirikine)

Coune, C.A. *et al.*, *Phytochemistry*, 1978, **17**, 1447-1448 (Diploceline)

Hirata, T. *et al.*, *Chem. Comm.*, 1979, 1081-1083 (abs config)

Coune, C.A. *et al.*, *Herba Hung.*, 1980, **19**, 189-193; *CA*, **93**, 21790e (16-Epidiploceline)

Marini-Bettolo, G.B. *et al.*, *Phytochemistry*, 1980, **19**, 992-994 (Strychnorubigine)

Mukhopadhyay, S. *et al.*, *J. Nat. Prod.*, 1983, **46**, 409-413 (16-Epi-Z-isositsirikine)

Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694-707 (oxides)

Caron, C. *et al.*, *Phytochemistry*, 1984, **23**, 2355-2357 (16R-Z-Isositsirikine)

Kohl, W. *et al.*, *Planta Med.*, 1984, **50**, 242-244 (16R-Z-Isositsirikine)

Winterfeldt, E. *et al.*, *Annalen*, 1986, 1262-1267 (synth)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1986, **24**, 703-708 (Bhimberine)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1986, **25**, 1731-1733 (Rhazimanine)

Freund, R. *et al.*, *Annalen*, 1988, 1007-1008 (Rhazimanine, Bhimberine)

Naito, T. *et al.*, *Heterocycles*, 1988, **27**, 1603-1606 (Isositsirikine, 16-Epiisositsirikine, Rhazimanine, Bhimberine, synth)

Naito, T. *et al.*, *Tet. Lett.*, 1989, **30**, 2941-2944 (Z-Isositsirikine, 16-Epi-Z-isositsirikine, synth)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1991, **30**, 1285-1293 (O-Acetylisositsirikine)

Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1992, **61**, 429-440 (9-Methoxyisositsirikines)

Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 131-133 (Rhazimanine, Bhimberine)

Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (occur, *Strychnos*)


Liang, S. *et al.*, *Chin. Chem. Lett.*, 2006, **17**, 783-786 (19,20-Dihydro-10-hydroxyisositsirikine)

Hu, X.-J. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1344-1350 (N-Hydroxymethylisositsirikine)

Isosorphidine

(5 β ,6 β)-Matridin-15-one, 9CI

I-320



$C_{15}H_{24}N_2O$ 248.367

Stereoisomer of Allomatrine, A-628, Darvasamine, D-79, Isomatrine, I-244, Matrine, M-121 and Sophoridine, S-380.

(+)-form [6838-36-4]

Alkaloid from *Sophora alopecuroides* (Fabaceae). Cryst. (petrol). Mp 111-112°. $[\alpha]_D$ +101 (EtOH).

Rulko, F. *et al.*, *Zh. Obshch. Khim.*, 1962, **32**, 1690-1695 (struct)

Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 331-332; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 282 (ms)


Kushmuradov, Y.K. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 244-247; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 207-210 (biosynth)

Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 460-465; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 340-344 (cryst struct)

α -Isosparteine

I-321

11 β -Sparteine. Genistein†



(-)-form

$C_{15}H_{26}N_2$ 234.384

Log P 2.15 (calc).

(+)-form

From (+) Sparteine, S-385 by epimerisation. Mp 110-115° (hydrate). $[\alpha]_D^{25}$ +48.4 (MeOH). pK_{a1} 2.5; pK_{a2} 12.1 (66% DMF aq.).

(-)-form [446-95-7]

Alkaloid from *Cytisus scoparius*, *Lupinus caudatus* and *Sarothamnus scoparius* (Fabaceae). Mp 96-117° (hydrate). $[\alpha]_D$ -56 (MeOH).

Sulfate: Mp 267°.

Dipicrate: Mp 221°.

(±)-form

Mp 98-105° (hydrate) Mp 78-80° (anhyd.).

Perchlorate: Mp 160-162°.

Leonard, N.J. *et al.*, *J.A.C.S.*, 1950, **72**, 1316 (synth, ir)

Marion, L. *et al.*, *Can. J. Chem.*, 1951, **29**, 22; 297 (isol, struct)

Kettelhack, D. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1954, **287**, 1 (isol)


Fales, H.M. *et al.*, *J.A.C.S.*, 1970, **92**, 1590 (ms)

Schöpf, C. *et al.*, *Annalen*, 1972, **755**, 86 (synth)

Przybylska, M. *et al.*, *Acta Cryst. B*, 1974, **30**, 2455 (cryst struct)

Klyne, W. *et al.*, *J.C.S. Perkin I*, 1974, 2565 (cd)
 Bohlmann, F. *et al.*, *Chem. Ber.*, 1975, **108**, 1043 (cmr)
 Oinuma, H. *et al.*, *J.C.S. Perkin I*, 1990, 2593 (synth)
 Norcross, N.R. *et al.*, *J.O.C.*, 2008, **73**, 7939-7951 (synth)

β-Isosparteine I-322
Spartalupine. Pusilline. Spathulatine. Nonalupine



(-)-form

C₁₅H₂₆N₂ 234.384

Care needed with numbering of derivs owing to twofold rotational symmetry. The term 'β-Isosparteine' was first used by Winterfeld *et al* (1934) for a different compd. (semisynthetic). For the naturally occurring alkaloid the name Spartalupine has historical precedence, but β-Isosparteine is preferred as being more descriptive. Nonalupine was actually a hydrate of β-Isosparteine and Spathulatine was its hydrochloride.

(+)-form

From (-)-Sparteine by epimerisation. Oil. [α]_D²⁵ +15.8 (c, 1.25 in EtOH). Air-sensitive.

(-)-form [492-06-8]

Alkaloid from *Lupinus argenteus stenophyllus*, *Lupinus solosericeus*, *Sophora secundiflora* (Fabaceae). Forms metal complexes. Mp 32-32.5°. Bp_{0.05} 110°. [α]_D²⁵ -15.4 (EtOH).

Perchlorate: Mp 216° (211.5-212°).

Perchlorate (1:2):

Small stout prisms (MeOH). Mp 247-249° dec.

Dipicrate: Mp 133-133.5°.

(±)-form

Dipicrate: Mp 220.5-221° dec.

(ξ)-form

10-Oxo- 10-Oxo-β-isosparteine

C₁₅H₂₄N₂O 248.367

Minor alkaloid from *Lupinus sericeus* (Fabaceae). Oil. [α]_D²⁵ -4.8 (c, 0.15 in EtOH).

17-Oxo, 2ξ-hydroxy: Lupanoline. 2-Hydroxy-17-oxo-β-isosparteine

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Lupinus sericeus* (tentative identification) (Fabaceae). Stout prisms (Me₂CO). Mp 174-176°. [α]_D²³ +31 (c, 0.726 in EtOH). [α]_D²³ +64.4 (c, 0.88 in H₂O).

17-Oxo, 2ξ-hydroxy, hydrochloride:

Prisms (Me₂CO/MeOH). Mp 275-276°.

10,17-Dioxo: 10,17-Dioxo-β-isosparteine

C₁₅H₂₂N₂O₂ 262.351

Minor alkaloid from *Lupinus sericeus* (Fabaceae).

[24915-04-6]

Couch, J.F. *et al.*, *J.A.C.S.*, 1940, **62**, 554 (*Lupanoline, isol*)
 Marion, L. *et al.*, *Can. J. Chem.*, 1953, **31**, 181; 187 (*Lupanoline, isol, struct*)
 Carmack, M. *et al.*, *J.A.C.S.*, 1955, **77**, 4435 (*isol, struct, synth*)
 Greenhalgh, R. *et al.*, *Can. J. Chem.*, 1956, **34**, 456
 Winterfeld, K. *et al.*, *Annalen*, 1966, **698**, 230 (*synth*)
 Carmack, M. *et al.*, *J.O.C.*, 1967, **32**, 3045 (*isol, ir, nomencl*)
 Skolik, J. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1969, **17**, 201 (*ir, conformn*)
 Kim, I.C. *et al.*, *J. Agric. Food Chem.*, 1982, **30**, 796 (*10-Oxo-β-isosparteine, 10,17-Dioxo-β-isosparteine*)
 Norcross, N.R. *et al.*, *J.O.C.*, 2008, **73**, 7939-7951 (*synth*)

Isostemonidine I-323

C₁₉H₃₁NO₅ 353.458

Struct. unknown. Alkaloid from the roots of *Stemona ovata* (Stemonaceae). Mp 137°. [α]_D -48.9 (Me₂CO). Can be written C₁₇H₂₈(NH)(CO)(OH)₂(COO). Suzuki, K. *et al.*, *Yakugaku Zasshi*, 1934, **54**, 567-572; *CA*, **31**, 106 (*isol*)

Isostemonine I-324

[24945-83-3]

C₂₂H₃₃NO₄ 375.507


Isomeric with Stemoninine, S-521 and Tuberostemonine, T-664. Struct. unknown. Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Mp 212-216° dec.

Pfeifer, S. *et al.*, *Pharmazie*, 1968, **23**, 342 (*isol, uv*)

Isostephodeline I-325

Stephaphylline

[56648-85-2]



Relative configuration

C₂₁H₂₇NO₅ 373.448

Struct. revised in 1989. The struct. originally proposed for Isostephodeline has now been assigned to Tannagine, T-28. Alkaloid from roots of *Stephania delavayi*, tuberous roots of *Stephania suberosa* and aerial parts of *Stephania zippeliana* (Menispermaceae). Mp 197° (184-185°). [α]_D +160 (c, 2 in EtOH). [α]_D +93 (c, 1.0 in CHCl₃). [α]_D²⁵ +77.3 (c, 0.35 in CHCl₃). λ_{max} 211; 225; 273 (EtOH).

14-Epimer: Erromangine

[123688-33-5]


C₂₁H₂₇NO₅ 373.448

Alkaloid from aerial parts of *Stephania zippeliana* (Menispermaceae). [α]_D +15 (c, 0.85 in CHCl₃). λ_{max} 212; 224; 273 (EtOH).

Perel'son, M.E. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 188; *Chem. Nat. Compd. (Engl.*

Transl.), 1975, **11**, 197 (*uv, pmr, isol*)
 Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 2391 (*Stephaphylline*)
 Charles, B. *et al.*, *Can. J. Chem.*, 1989, **67**, 1257-1260 (*Isostephodeline, Erromangine, isol, uv, ir, pmr, cmr, ms, struct*)
 Guinadeau, H. *et al.*, *Phytochemistry*, 1998, **49**, 2561-2563 (*cd, pmr, cmr*)

Isotrychnopentamine B I-326



C₃₅H₄₃N₅O 549.758


Proposed struct. Alkaloid from *Strychnos usambarensis* leaves (Loganiaceae).

Angenot, L. *et al.*, *J. Pharm. Belg.*, 1978, **33**, 11 (*isol*)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 54 (*rev*)

Isotaifine I-327

[84323-09-1]



C₁₄H₁₃NO₃ 243.262

Struct. unknown. The proposed struct. illus. is incorrect. Props. of authentic synthetic material differ from those of the alkaloid. Alkaloid from *Ruta chalepensis* (Rutaceae). Needles (petrol). Mp 123-125°.


Mohr, N. *et al.*, *Phytochemistry*, 1982, **21**, 1838-1839 (*isol, uv, ir, pmr*)

Shich, B. *et al.*, *Hua Hsueh*, 1990, **48**, 309-316; *CA*, **116**, 174486s

Isotetrandrine I-328

O-Methylberbamine

[477-57-6]



C₃₈H₄₂N₂O₆ 622.76


Alkaloids covered by this entry (1R,1'S-series) are enantiomeric with those included under Peinamine, P-171. Alkaloid from *Atherosperma moschatum*, *Berberis empetrifolia*, *Berberis kawakamii*, *Berberis mingeisensis*, *Berberis morrisonensis*, *Berberis poirerii*, *Berberis thunbergii*, *Cy- clea barbata*, *Doryphora aromatica*, *Iso-*

- pyrum thalictroides*, *Laurelia sempervirens* (Peruvian nutmeg), *Limaciopsis loagensis*, *Mahonia japonica*, *Mahonia lomarii-folia*, *Mahonia morrisonensis*, *Mahonia philippinensis*, *Pycnarrhena australiana*, *Pycnarrhena manillensis*, *Stephania cepharantha*, *Stephania elegans*, *Thalictrum foetidum*, *Tiliacora funifera*, and *Triclisia gillettii* (Monimiaceae, Berberidaceae, Menispermaceae, Ranunculaceae). Shows *in vitro* antineoplastic activity against HeLa cells and Ehrlich ascites. Antiinflammatory, antihypertensive agent. Active against gram-positive bacteria. Prisms (MeOH). Mp 182-183°. $[\alpha]_D^{20} +150.7$ (c, 0.85 in CHCl_3). Log P 8.3 (uncertain value) (calc). λ_{max} 218 (ε 31600); 240 (sh) (ε 15800); 282 (ε 12600) (MeOH) (Derep).
- ▶ LD₅₀ (mus, orl) 6400 mg/kg; LD₅₀ (mus, ipr) 160 mg/kg. NX7285000
- N²-Oxide: Isotetrandrine N-2'-oxide**
[70191-83-2]
C₃₈H₄₂N₂O₇ 638.759
Alkaloid from the roots of *Limaciopsis loagensis* (Menispermaceae). Cryst. (MeOH). Mp 191-192°. $[\alpha]_D +94$ (CHCl_3).
- N²-De-Me: 2-Norisotetrandrine**
[123930-96-1]
C₃₇H₄₀N₂O₆ 608.733
Alkaloid from the tubers of *Stephania pierrii* (Menispermaceae). $[\alpha]_D +100$ (c, 0.16 in CHCl_3). λ_{max} 208 (log ε 4.82); 239 (sh) (log ε 4.34); 282 (log ε 3.86) (MeOH).
- N²-De-Me: 2'-Norisotetrandrine**
[70191-82-1]
C₃₇H₄₀N₂O₆ 608.733
Alkaloid from the roots of *Limaciopsis loagensis* (Menispermaceae). Non-cryst. $[\alpha]_D +26$ (CHCl_3).
- N²-Me: 2'-N-Methylisotetrandrine**
[68331-87-3]
C₃₉H₄₅N₂O₆[⊕] 637.794
Quaternary alkaloid from shoots of *Berberis oblonga* (Berberidaceae). Cryst. (THF) (as iodide). Mp 221-222° (iodide). $[\alpha]_D +29.5$ (c, 0.1 in CHCl_3).
- O⁶-De-Me: Cycleabarbaine. 6-O-De-methylisotetrandrine**
[152697-36-4]
C₃₇H₄₀N₂O₆ 608.733
Alkaloid from roots of *Cyclea barbata* (Menispermaceae). $[\alpha]_D +20$ (c, 0.1 in CHCl_3).
- O⁷-De-Me: Thalrugosine. Thaligine. Isotetrandrine**
[33889-68-8]
C₃₇H₄₀N₂O₆ 608.733
Alkaloid from *Cyclea barbata*, *Thalictrum polygamum*, *Thalictrum lucidum*, *Thalictrum rugosum*, *Thalictrum minus* Race B, *Thalictrum sachalinense*, *Limaciopsis loagensis*, *Pycnarrhena novoguineensis*, *Tiliacora funifera* and *Stephania japonica* var. *australis* (Menispermaceae, Ranunculaceae). Shows antibacterial activity. Cryst. + 0.5H₂O (Et₂O). Mp 212-214°. $[\alpha]_D^{30} +128$ (MeOH). λ_{max} 282 (ε 8710) (EtOH).
- O⁷-De-Me, N²-de-Me: 2-Northalrugosine**
[65995-42-8]
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from stems of *Pycnarrhena ozantha* (Menispermaceae). $[\alpha]_D +209$ (c, 0.16 in CHCl_3).
- O⁷-De-Me, N²,N²-di-de-Me: Bisnorthalrugosine**
[111537-45-2]
C₃₅H₃₆N₂O₆ 580.679
Alkaloid from the stems of *Pycnarrhena ozantha* (Menispermaceae). $[\alpha]_D +142$ (c, 0.13 in CHCl_3).
- O¹²-De-Me: Berbamine. Berbenine**
[478-61-5]
C₃₇H₄₀N₂O₆ 608.733
Alkaloid from *Atherosperma moschatum*, many *Berberis* spp., *Cyclea*, *Limaciopsis*, *Mahonia* and other genera (Monimiaceae, Berberidaceae, Menispermaceae, Ranunculaceae). Antihypertensive, leucocytotic agent. Strong curarising agent. Shows antibacterial activity. Cryst. (C₆H₆). Mp 155-157°. $[\alpha]_D +109$ (c, 0.24 in CHCl_3). Log P 7.83 (uncertain value) (calc). Isom. to Pycnamine in P-301 when treated with MeOH (40°/24h then 2 days at r.t.). λ_{max} 284 (log ε 3.79) (EtOH).
- ▶ Highly toxic; LD₅₀ (mus, ipr) 75 mg/kg. DR9450000
- O¹²-De-Me, hydrochloride (1:2):**
Needles + 7H₂O. Mp 257-258° dec. (2% HCl). $[\alpha]_D^{20} +64.2$ (c, 1.0 in H₂O).
- O¹²-De-Me, N²-β-oxide: Berbamine 2'-β-N-oxide**
[103839-20-9]
C₃₇H₄₀N₂O₇ 624.732
Alkaloid from aerial parts of *Berberis brandisiana* (Berberidaceae).
- O¹²-De-Me, N²-de-Me: 2-N-Norberbamine**
[39028-61-0]
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from *Pycnarrhena australiana* whole plant (Menispermaceae). Cryst. + 0.66 C₆H₆ (C₆H₆). Mp 166-168°. $[\alpha]_D +117$ (c, 0.23 in CHCl_3).
- O¹²-De-Me, N²-Me: 2'-N-Methylberbamine**
[68231-29-8]
C₃₈H₄₃N₂O₆[⊕] 623.767
Quaternary alkaloid from *Berberis oblonga* (Berberidaceae). Amorph. solid (as iodide). CAS no. refers to iodide.
- O¹²-De-Me, N²,N²-di-Me:**
Needles + 1H₂O (Me₂CO aq.). Mp 287-289° dec.
- O⁶,O¹²-Di-de-Me: Aquifoline. 6-O-De-methylberbamine**
[107900-77-6]
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from roots of *Mahonia aquifolium* (also present in stem and seeds) (Berberidaceae). Cryst. (C₆H₆/CHCl₃). Mp 168°. $[\alpha]_D^{25} +80$ (c, 0.1 in MeOH). λ_{max} 229 (sh) (log ε 4.6); 282 (log ε 2.39) (MeOH).
- O⁷,O¹²-Di-de-Me: Obamegine. Stepholine**
[479-37-8]
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from *Berberis tschonoskyana*, *Stephania japonica*, *Thalictrum lucidum*, *Thalictrum rugosum*, *Xanthorhiza simplicissima*, *Mahonia repens* and *Triclisia gillettii* (Berberidaceae, Menispermaceae, Ranunculaceae). Shows antibacterial activity. Antineoplastic agent. Needles (C₆H₆). Mp 171-173°. $[\alpha]_D^{19} +273$ (CHCl_3). Log P 7.54 (uncertain value) (calc). λ_{max} 280 (log ε 3.19) (MeOH) (Berdy).
- O⁷,O¹²-Di-de-Me, N²-de-Me: 2-N-Norobamegine**
[38962-94-6]
Alkaloid from *Pycnarrhena australiana* and *Pycnarrhena ozantha* (Menispermaceae). Cryst. + 2CHCl₃ (CHCl_3). Sol. CHCl₃-MeOH; fairly sol. Me₂CO. Mp 188-190° dec. $[\alpha]_D +290$ (c, 0.25 in CHCl_3). λ_{max} 283 (ε 7530) (MeOH) (Berdy).
- O⁷,O¹²-Di-de-Me, N²,N²-di-de-Me: Bisnorobamegine**
[111537-44-1]
C₃₄H₃₄N₂O₆ 566.652
Alkaloid from the stems of *Pycnarrhena ozantha* (Menispermaceae). $[\alpha]_D +260$ (c, 0.65 in CHCl_3).
- Bick, I.R.C. et al., *Aust. J. Chem.*, 1956, **9**, 111-119; 1980, **33**, 225-228 (*isol, synth, Berbamine*)
Tomita, M. et al., *Yakugaku Zasshi*, 1959, **79**, 317-321; 1963, **83**, 940-944; *CA*, **53**, 17161e; **60**, 4202a (*Obamegine, isol, struct*)
Kugo, T. et al., *Yakugaku Zasshi*, 1959, **79**, 322-324; *CA*, **53**, 17161f (*Obamegine*)
Yang, T.-H. et al., *Yakugaku Zasshi*, 1960, **80**, 847-849; *CA*, **54**, 23187i (*isol*)
Battersby, A.R. et al., *J.C.S.*, 1965, 2239-2247 (*uv, ord*)
Tomimatsu, T. et al., *J. Pharm. Sci.*, 1966, **55**, 208-209 (*Obamegine, uv, ir, pmr*)
Falco, M.R. et al., *Tet. Lett.*, 1968, 1953-1959 (*pmr*)
Mitscher, L.A. et al., *Dtsch. Apoth. -Ztg.*, 1971, **111**, 1704-176 (*activity*)
Loder, J.W. et al., *Aust. J. Chem.*, 1972, **25**, 2193-2197 (*2-N-Norobamegine, 2-N-Norberbamine, isol, uv, pmr, ms, struct*)
Sioumis, A.A. et al., *Aust. J. Chem.*, 1972, **25**, 2251-2254 (*2-N-Norobamegine, 2-N-Norberbamine, isol, uv, pmr, ms*)
Mitscher, L.A. et al., *J. Nat. Prod.*, 1972, **35**, 167-176 (*Thalrugosine, Obamegine, isol, uv, ir, pmr, ms, cd, struct*)
Baldas, J. et al., *J.C.S. Perkin 1*, 1972, 592-596 (*ms*)
Shamma, M. et al., *Experientia*, 1973, **29**, 517-518 (*Thaligine*)
Tackie, A.N. et al., *J. Nat. Prod.*, 1974, **37**, 1-5 (*isol, uv, pmr, ms*)
Kuroda, H. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2413-2420 (*activity*)
Akasu, M. et al., *Phytochemistry*, 1976, **15**, 471-473 (*Berberamine, isol, uv, ir, pmr, ms*)
Ayim, J.S.K. et al., *J. Nat. Prod.*, 1977, **40**, 561-565 (*Thalrugosine, isol*)
Karimov, A. et al., *Khim. Prir. Soedin.*, 1978, **14**, 227-231; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 186-189 (*2'-N-Methylberbamine*)
Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1979, **15**, 818-823; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 723-727 (*cd, abs config*)
Cavé, A. et al., *Planta Med.*, 1979, **35**, 31-41 (*2'-Norisotetrandrine, Isotetrandrine 2'-N-oxide*)

Koike, L. *et al.*, *Tet. Lett.*, 1979, 3765-3768 (cmr)
 Bhakuni, D.S. *et al.*, *Tetrahedron*, 1980, **36**, 2149-2151 (biosynth)
 Owusu, P.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 61-66 (*Obamegine, isol, uv, ir, pmr*)
 Moulis, C. *et al.*, *J. Nat. Prod.*, 1981, **44**, 101-103 (*isol, uv, ir, pmr, ms*)
 Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 274-278; 1986, **49**, 538-539 (*Berberamine 2'-β-N-oxide, isol*)
 Nakova, E. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 86-91; 91-95; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 83-87; 88-91 (*synth*)
 Buck, K.T. *et al.*, *Alkaloids (Academic Press)*, 1987, **30**, 147 (*pharmacol*)
 Kostálová, D. *et al.*, *Coll. Czech. Chem. Comm.*, 1987, **52**, 242-246 (*Aquifoline*)
 Abouchacra, M.L. *et al.*, *J. Nat. Prod.*, 1987, **50**, 375-380 (*2-Northalrugosine, Bisorthalrugosine, Bisorobamegine*)
 Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1989, **52**, 846-851 (*2-Norisorotetrandrine*)
 Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1989-1992 (*Cycleabarbatine*)
 Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*Thalrugosine, pmr, cmr*)
 Tanahashi, T. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 370-373 (*2'-N-Methylisotetrandrine*)
 Chen, C.-K. *et al.*, *Chin. Pharm. J. (Taipei)*, 2003, **55**, 35-47 (*Norobamegine, Obamegine, Thalrugosine*)

Isotalicherberine I-329

1-Isotalicherberine
 [25514-42-5]



$C_{37}H_{40}N_2O_6$ 608.733

Stereoisomer of Thalicherberine, T-313. Alkaloid from the aerial parts of *Berberis chilensis* (Berberidaceae). Cryst. (CHCl₃/cyclohexane). Mp 153-155°. $[\alpha]_D^{20}$ -205 (c, 0.4 in CHCl₃).

Me ether: O-Methylisotalicherberine. 7-O-Methylberlarine
 [19879-44-8]

$C_{38}H_{42}N_2O_6$ 622.76
 Alkaloid from the trunk bark and roots of *Berberis laurina* and from the aerial parts of *Berberis chilensis* (Berberidaceae). Mp 208-209°. $[\alpha]_D$ -195 (c, 0.5 in CHCl₃).

O⁷-De-Me: 7-O-Demethylisotalicherberine
 [73711-14-5]

$C_{36}H_{38}N_2O_6$ 594.706
 Alkaloid from the aerial parts of *Berberis chilensis* (Berberidaceae). Cryst. (CHCl₃/cyclohexane). Mp 245-247°. $[\alpha]_D^{20}$ +230 (c, 0.2 in CHCl₃).


Me ether, O⁷-De-Me: Belarine
 [35471-06-8]

$C_{37}H_{40}N_2O_6$ 608.733
 Alkaloid from the root bark of *Berberis laurina* (Berberidaceae). Mp 158-160°. $[\alpha]_D$ -222 (c, 0.2 in CHCl₃)
 [73711-13-4]

Falco, M.R. *et al.*, *Tet. Lett.*, 1968, 1953; *Chem. Comm.*, 1971, 1056 (*O-Methylisotalicherberine, Belarine*)
 Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 597 (*ms, Belarine*)
 Torres, R. *et al.*, *Gazz. Chim. Ital.*, 1979, **109**, 567 (*isol, uv, ir, pmr, ms, ord, struct, Isotalicherberine, O-Methylisotalicherberine*)

Isotalicrine I-330

[246852-81-3]




$C_{42}H_{52}N_2O_{10}$ 744.88

Alkaloid from *Isopyrum thalictroides*. Amorph. solid. $[\alpha]_D$ +33 (c, 0.31 in CHCl₃). λ_{max} 208 (log ε 6.62); 236 (sh) (log ε 6.06); 285 (log ε 5.74) (EtOH).

Philipov, S.A. *et al.*, *Phytochemistry*, 1999, **51**, 1161-1165 (*isol, uv, ir, pmr, cmr, ms*)

Isotalictrine I-331

[241482-19-9]




$C_{41}H_{44}N_2O_{10}$ 724.806

Alkaloid from *Isopyrum thalictroides*. Amorph. solid. Racemic. λ_{max} 208 (log ε 6.61); 236 (sh) (log ε 6.32); 259 (log ε 6.29); 285 (log ε 5.84) (EtOH).

Philipov, S.A. *et al.*, *Phytochemistry*, 1999, **51**, 1161-1165 (*isol, uv, ir, pmr, cmr, ms*)

Isotalirine I-332

[246852-82-4]




$C_{42}H_{52}N_2O_9$ 728.881

Alkaloid from *Isopyrum thalictroides*. Amorph. solid. Racemic. λ_{max} 208 (log ε 7.2); 236 (sh) (log ε 6.82); 285 (log ε 6.39) (EtOH).

Philipov, S.A. *et al.*, *Phytochemistry*, 1999, **51**, 1161-1165 (*isol, uv, ir, pmr, cmr, ms*)

Isotalisopavine I-333

10,11-Dihydro-3,7,8-trimethoxy-12-methyl-10,5-(iminomethano)-5H-dibenzo[a,d]cyclohepten-4-ol, 9CI




$C_{20}H_{23}NO_4$ 341.406

Alkaloid from *Thalictrum minus*. Amorph. solid. $[\alpha]_D^{25}$ -169 (c, 3.4 in CHCl₃).

Sidjimov, A.K. *et al.*, *Phytochemistry*, 1998, **48**, 403-405 (*isol, pmr, cmr, ms*)

Isotalmidine I-334




$C_{37}H_{36}N_2O_7$ 620.701

Alkaloid from *Isopyrum thalictroides*. Pale yellow solid. λ_{max} 210 (log ε 6.52); 240 (sh) (log ε 6.17); 290 (log ε 5.77); 312 (log ε 5.77) (EtOH).

Istatkova, R.S. *et al.*, *Phytochemistry*, 2000, **54**, 959-964

Isothapsakone A I-335

[302598-27-2]




$C_{35}H_{36}N_2O_9$ 628.677

Alkaloid from *Aglaia oligophylla*. Insecticidal agent. Amorph. solid. $[\alpha]_D^{20}$ +12.2 (c, 1 in EtOH).

Dreyer, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 415-420 (*isol, cd, pmr, cmr, ms*)

8*H*-Isothiazolo[5,4-*b*]indole, I-336

Brassilexin
 [119752-76-0]



$C_9H_6N_2S$ 174.226

Isol. from leaves of brown mustard (*Brassica juncea*) (Brassicaceae). Phytoalexin. Microcryst. Mp 172-173° (164-167°). λ_{\max} 218 (ϵ 50000); 245 (ϵ 14000); 264 (ϵ 12000) (MeOH) (Derep).

N-Ac:

$C_{11}H_8N_2OS$ 216.263
Mp 172-178°.

N-Methoxy: 8-Methoxy-8H-isothiazolo[5,4-b]indole. **Sinalexin**

[200192-82-1]

$C_{10}H_8N_2OS$ 204.252

Isol. from leaves of white mustard (*Sinapis alba*) (Brassicaceae) under biotic or abiotic elicitation. Phytoalexin.

Devys, M. et al., *Tet. Lett.*, 1988, **29**, 6447-6448 (isol, uv, ir, pmr, cmr, ms)

Devys, M. et al., *Synthesis*, 1990, 214-215 (synth, ir, uv, pmr, ms)

Pedras, M.S.C. et al., *Phytochemistry*, 1997, **46**, 833-837 (*Sinalexin*)


Pedras, M.S.C. et al., *Phytochemistry*, 2000, **53**, 161-176 (rev)

Pedras, M.S.C. et al., *J.O.C.*, 2005, **70**, 1828-1834 (synth)

Pedras, M.S.C. et al., *Org. Biomol. Chem.*, 2005, **3**, 2002-2007 (metab)

3-Isothiocyanato-4-methoxy-1H-indole, 9CI I-337

4-Methoxyindole 3-isothiocyanate. *Rapalexin A*
[929083-72-7]




$C_{10}H_8N_2OS$ 204.252

Phytoalexin isol. from white rust infected *Brassica rapa*. No phys. props. reported.

Pedras, M.S.C. et al., *Chem. Comm.*, 2007, 368-370 (isol, synth, pmr, cmr)

Isotrikentrin B I-338

4-(1-Butenyl)-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, 9CI
[107368-94-5]
[152784-95-7]



Absolute Configuration

$C_{17}H_{21}N$ 239.36

Inseparable mixt. with *cis*-Trikenrin B (ratio 2:3) (see Trikenrin B, T-595). Isol. from the marine sponge *Trikentrion flabelliforme*. Possesses antimicrobial activity. Unstable oil which darkens on storage. Sol. MeOH, $CHCl_3$, λ_{\max} 240 (ϵ 44900); 298 (ϵ 12300) (MeOH).

Capon, R.J. et al., *Tetrahedron*, 1986, **42**, 6545-6550 (isol, uv, ir, pmr, cmr, ms, struct)


Muratake, H. et al., *Chem. Pharm. Bull.*, 1994, **42**, 854-864 (abs config, synth)

Isotrilobine I-339

Homotrilobine. N,O-Dimethylcoccoline.

O-Methylcoccoline

[26195-62-0]



Absolute Configuration

$C_{36}H_{36}N_2O_5$ 576.691

Alkaloids covered by this entry are enantiomeric with those included in Micranthine, M-591 and diastereomeric with those in Apateline, A-1331. The name Isotrilobine is misleading. Homotrilobine is more accurate. Alkaloid from the stems of *Anisocycla grandidieri*, *Albertisia papuana* and *Cocculus pendulus*, the stems and roots of *Cocculus hirsutus*, the roots of *Cocculus sarmentosus* and *Stephania hermandifolia*, the roots and tops of *Pachygone pubescens*, the bark and trunk of *Cocculus laurifolius* and from *Cocculus trilobus* (all parts) (Menispermaceae). Shows antitumor activity against HeLa-S₃ cells and Ehrlich ascites. Shows antibacterial activity. Needles (Me₂CO). Mp 217-218°. $[\alpha]_D^{20} +325$ (c, 0.41 in $CHCl_3$).

► LD₅₀ (mus, ipr) 115 mg/kg. NY0890000

2-N-Oxide: **Isotrilobine 2-N-oxide**

[139953-39-2]

$C_{36}H_{36}N_2O_6$ 592.69

Alkaloid from roots of *Cocculus trilobus* (Menispermaceae). Light yellow cryst. (MeOH). Mp 178-179°. $[\alpha]_D^{20} +150.9$ (c, 0.91 in $CHCl_3$).

N²-De-Me: **O-Methylcoccoline**

[54352-71-5]

$C_{35}H_{34}N_2O_5$ 562.664

Alkaloid from the stems of *Albertisia papuana* (Menispermaceae). Cryst. (EtOAc). Mp 235-237°. $[\alpha]_D +236$ (c, 1.2 in $CHCl_3$). Phys. constants are for synthetic material.

N²-De-Me, N²-β-oxide: **O-Methylcoccoline 2'-β-N-oxide**

[149492-48-8]

$C_{35}H_{34}N_2O_6$ 578.663

Alkaloid from roots of *Anisocycla cymosa* (Menispermaceae). Amorph. powder.

N²-De-Me: **Trilobine**

[6138-73-4]

$C_{35}H_{34}N_2O_5$ 562.664

Alkaloid from *Anisocycla grandidieri*, *Pachygone ovata*, *Cocculus hirsutus*, *Cocculus sarmentosus*, *Cocculus laurifolius*, *Cocculus trilobus* and *Stephania japonica*. Convulsive agent. Shows *in vitro* antitumor activity against HeLa-S₃ cells. Cryst. (Me₂CO). Mp 235-239°. $[\alpha]_D^{20} +305$ (c, 1.26 in $CHCl_3$).

► LD₅₀ 500 mg/kg.

N²,N²-Di-de-Me: **Nortrilobine**

[91897-39-1]

$C_{34}H_{32}N_2O_5$ 548.637

Alkaloid from the roots of *Pachygone*

ovata (Menispermaceae). Powder (MeOH). Mp 177-180°. $[\alpha]_D^{25} +216$ (c, 0.44 in $CHCl_3$).

O⁶-De-Me: **12-O-Methyltricordatine**

[187086-17-5]

$C_{35}H_{34}N_2O_5$ 562.664

Alkaloid from stem bark of *Pachygone dasycarpa*. Shows antiplasmodial activity. $[\alpha]_D +55$ (c, 0.196 in MeOH). λ_{\max} 206 (log ϵ 4.82); 236 (sh) (log ϵ 4.29); 286 (log ϵ 3.73) (MeOH).

O¹²-De-Me: **Coccoline**. *Efirine*. *Trigillentine*

[26279-88-9]

$C_{35}H_{34}N_2O_5$ 562.664

Alkaloid from *Cocculus pendulus*, *Cocculus laeabe*, *Triclisia gillettii*, *Triclisia dictyophylla*, *Triclisia patens*, *Synclisia scabrada* and *Anisocycla papuana* (Menispermaceae). Minute needles (abs. EtOH). Mp 272-274°. $[\alpha]_D^{23} +292$ (c, 0.76 in $CHCl_3$). $[\alpha]_D^{23} +337$ (c, 0.86 in Py).

O¹²-De-Me, N²-oxide: **Coccoline 2-N-oxide**

[91106-34-2]

$C_{35}H_{34}N_2O_6$ 578.663

Alkaloid from the stems and roots of *Cocculus hirsutus* (Menispermaceae). Amorph. (MeOH). Mp 182-187°. $[\alpha]_D^{25} +125$ (c, 0.5 in MeOH).

O¹²-De-Me, Ac:

Rosettes of prisms (EtOAc/hexane). Mp 166-168° dec.

O¹²-De-Me, N²-de-Me: **Coccoline**

[54352-70-4]

$C_{34}H_{32}N_2O_5$ 548.637

Alkaloid from the leaves and stems of *Cocculus pendulus*, the roots of *Cocculus laeabe*, and the stems of *Synclisia scabrada* and *Albertisia papuana* (Menispermaceae). Amorph. powder. Mp 197-199°. $[\alpha]_D +204$ (c, 1.3 in $CHCl_3$).

O¹²-De-Me, N²-de-Me, N²-β-oxide:

Coccoline 2'-β-N-oxide

[149492-49-9]

$C_{34}H_{32}N_2O_6$ 564.637

Alkaloid from roots of *Anisocycla cymosa* (Menispermaceae). Amorph. powder.

O¹²-De-Me, N²,N²-di-de-Me: **2'-Norcoccoline**. *12-O-Demethyltrilobine*

[39986-72-6]

$C_{33}H_{30}N_2O_5$ 534.61

Alkaloid from roots of *Anisocycla cymosa*, *Anisocycla grandidieri* and *Albertisia* sp. (Menispermaceae). Cryst. Mp 256-258°. $[\alpha]_D +238$ (c, 0.15 in $CHCl_3$). $[\alpha]_D +332$ (c, 1.52 in Py).

O⁶,O¹²-Di-de-Me: **Tricordatine**

[51076-20-1]

$C_{34}H_{32}N_2O_5$ 548.637

Alkaloid from the roots of *Triclisia subcordata* and the stems of *Cocculus pendulus* (Menispermaceae). Cryst. ($CHCl_3$ /MeOH). Mp 280° dec. $[\alpha]_D^{22} +247.9$ (c, 1.17 in Py). Insolubility in common organic solvents precluded the recording of pmr spectra.

10-Hydroxy: **10-Hydroxyisotrilobine**

$C_{36}H_{36}N_2O_6$ 592.69

Alkaloid from the stems of *Cocculus*

orbiculatus. Light yellow powder. $[\alpha]_D^{25} +96.1$ (c, 1 in MeOH). λ_{\max} 220 ; 233 ; 288 (MeOH).

Kondo, H. *et al.*, *Yakugaku Zasshi*, 1926, **532**, 461-465; 1927, **542**, 265-278 (*Trilobine, isol. struct*)

Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1953, **1**, 1-5 (*Trilobine, isol*)

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 977-979; 1963, **83**, 190-194; *CA*, **53**, 22042i; **59**, 3971h (*Trilobine, isol*)

Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 282-288; 288-292 (*Trilobine, struct*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2289-2290 (*isol, pmr*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 2061-2064 (*12-O-Demethyltrilobine*)

Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592-596 (*ms*)

Tackie, A.N. *et al.*, *Phytochemistry*, 1973, **12**, 2509-2511 (*Tricordatine*)

Joshi, P.P. *et al.*, *Indian J. Chem.*, 1974, **12**, 649-650 (*Coccoline*)

Weber, N. *et al.*, *Phytochemistry*, 1974, **13**, 2326 (*Coccoline*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1975, **31**, 2575-2579 (*Coccoline, Coccoline*)

Kuroda, H. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 2413-2420 (*Isotrilobine, activity*)

Tripathi, V.J. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 62-63 (*isol, uv, pmr, ms*)

Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 1636-1644 (*Trilobine, synth*)

Bhakuni, D.S. *et al.*, *J.C.S. Perkin 1*, 1978, 121-125 (*biosynth*)

Dasgupta, S. *et al.*, *J. Nat. Prod.*, 1979, **42**, 399-406 (*Trilobine, isol, uv, ir, pmr, ms*)

Bick, I.R.C. *et al.*, *Heterocycles*, 1981, **16**, 2105-2108 (*biosynth*)

Spiff, A.I. *et al.*, *J. Nat. Prod.*, 1981, **44**, 160-165 (*Coccoline*)

Leboeuf, M. *et al.*, *Plant. Med. Phytother.*, 1982, **16**, 280-291 (*O-Methylcoccoline*)

Ohiri, F.C. *et al.*, *Planta Med.*, 1983, **47**, 87-89 (*Coccoline, Coccoline*)

El-Shabrawy, A.O. *et al.*, *Heterocycles*, 1984, **22**, 993-995 (*Coccoline 2-N-oxide*)

El-Kawi, M.A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 459-465 (*Nortrilobine*)

Hussain, S.F. *et al.*, *Tetrahedron*, 1984, **40**, 2513-2517 (*Tricordatine, Coccoline, isol, cd, uv, pmr, ms, struct*)

Lavault, M. *et al.*, *Can. J. Chem.*, 1987, **65**, 343-347 (*2'-Norcoccoline*)

Chen, H.S. *et al.*, *Yaoxue Xuebao*, 1991, **26**, 755-758; *CA*, **116**, 170119b (*Isotrilobine 2-N-oxide*)

Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1993, **56**, 957-960 (*Coccoline 2'-β-N-oxide, 12-O-Methylcoccoline 2'-β-N-oxide, 2'-Norcoccoline*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1997, **60**, 258-260 (*12-O-Methyltricordatine*)


Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*Coccoline, pmr*)

Atta-ur-Rahman, *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 802-806 (*Coccoline, Coccoline*)

Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1056-1060 (*10-Hydroxyisotrilobine*)

Isotriperideine I-340

Tetradecahydro-2H,11H-triptyrido[1,2-a:1',2'-c:3'',2''-e]pyrimidine, 9CI [482-59-7]



$C_{15}H_{27}N_3$ 249.398

A large number of stereoisomers are theoretically possible but it appears that the nat. alkaloid is identical with the *all-trans*-isomer (illus.) prepared synthetically and characterised by nmr. Alkaloid from *Coelidium fourcadei* dried branches and leaves (Fabaceae). Cryst. (hexane). Mp 97-98° (94-96°). Prob. an artifact of thermal dec.

Schöpf, C. *et al.*, *Chem. Ber.*, 1951, **84**, 690; 1952, **85**, 937 (*synth*)

Haglid, F. *et al.*, *Acta Chem. Scand.*, 1966, **20**, 2896 (*synth*)

Arndt, R.R. *et al.*, *J. S. Afr. Chem. Inst.*, 1968, **21**, 54; *CA*, **70**, 4342f (*ir, uv, ms, pmr, isol, struct*)

Kessler, H. *et al.*, *J.O.C.*, 1977, **42**, 66 (*cmr, config*)

Warning, K. *et al.*, *Tet. Lett.*, 1979, 1565 (*synth*)

Isotuberostemonine I-341

Struct. unknown. Minor alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Mp 118-120°. $[\alpha]_D^{25} -86.15$ (MeOH). Mol. formula $C_{22}H_{33}NO_4$ or $C_{22}H_{35}NO_4$.

Hydroiodide (1:3):

Reddish-brown needles. Mp 214-216° dec.

Perchlorate:


Cryst. + $\frac{1}{2}$ H₂O (MeOH). Mp 211-213°.

Kondo, H. *et al.*, *CA*, 1957, **51**, 1540f (*isol*)

Kondo, H. *et al.*, *Chem. Zentralbl.*, 1958, **129**, 14057 (*isol*)

Isotylocrebrine I-342

9,11,12,13,13a,14-Hexahydro-3,4,6,7-tetramethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinoline, 9CI, 3,4,6,7-Tetramethoxyphenanthroindolizidine



(*S*)-form

$C_{24}H_{27}NO_4$ 393.482

Abs. config. of natural Isotylocrebrine has been reversed from (*R*)- to (*S*)-.

(*R*)-form

β-N-Oxide: (10S,13aR)-Isotylocrebrine N-oxide

$C_{24}H_{27}NO_5$ 409.481

Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 230-235° dec. $[\alpha]_D^{25} -66.6$ (c, 0.02 in MeOH). λ_{\max} 245 (log ε 3.24); 263 (log ε 5.39); 283 (log ε 3.12); 287 (log ε 3.08); 309 (log ε 2.41); 344 (log ε 2.24); 359 (log ε 2.19) (MeOH).

7-Demethoxy: 3,4,6-Trimethoxyphenanthroindolizidine. Ficuseptine D

$C_{23}H_{25}NO_3$ 363.455

Alkaloid from the stems of *Ficus septica*. Gum. $[\alpha]_D^{25} -75.6$ (c, 0.1 in

MeOH). λ_{\max} 252 (sh) (log ε 3.58); 260 (log ε 3.62); 280 (log ε 2.68); 342 (log ε 2.61); 351 (log ε 3.21) (MeOH).

(*S*)-form [50657-02-8]

Minor alkaloid from the leaves of *Tylophora asthmatica* and from *Tylophora tanakae* (Asclepiadaceae). Cryst. (CH₂Cl₂/MeOH). Mp 212-214° (198-203° dec.). $[\alpha]_D^{25} +22.43$ (c, 1.1 in CHCl₃).

N-Oxide: Isotylocrebrine N-oxide

[165606-50-8]

$C_{24}H_{27}NO_5$ 409.481

From leaves of *Tylophora tanakae* (Asclepiadaceae). Solid. $[\alpha]_D^{20} +28.9$ (c, 0.30 in MeOH).

β-N-Oxide: (10S)-Isotylocrebrine N-oxide

$C_{24}H_{27}NO_5$ 409.481

Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 205-210° dec. $[\alpha]_D^{25} +6.5$ (c, 0.02 in MeOH). λ_{\max} 245 (log ε 3.21); 264 (log ε 5.41); 285 (log ε 3.17); 305 (log ε 2.89); 344 (log ε 2.21) (MeOH).

O³-De-Me: 3-De-O-methylisotylocrebrine

[165606-48-4]

$C_{23}H_{25}NO_4$ 379.455

From leaves of *Tylophora tanakae* (Asclepiadaceae). Mp 193-203° dec. $[\alpha]_D^{28} +34.5$ (c, 0.80 in CHCl₃/MeOH, 1:1).

O³,O⁶-Di-de-Me: 3,6-Dide-O-methylisotylocrebrine

[209684-49-1]

$C_{22}H_{23}NO_4$ 365.428

Alkaloid from *Tylophora tanakae*.

Brownish needles. Mp 233-238° dec.

$[\alpha]_D^{32} +102.7$ (c, 0.2 in CHCl₃/MeOH).

λ_{\max} 261 (log ε 4.44); 280 (log ε 4.15);

286 (log ε 4.14); 305 (sh) (log ε 3.75);

316 (log ε 3.75) (MeOH).

14α-Hydroxy, N-oxide: 14α-Hydroxyisotylocrebrine N-oxide

[165606-51-9]

$C_{24}H_{27}NO_6$ 425.48

From leaves of *Tylophora tanakae*

(Asclepiadaceae). Mp 212-215° dec.

$[\alpha]_D^{26} +8.3$ (c, 0.12 in CHCl₃/MeOH,

1:1).

14α-Hydroxy, O³-de-Me: 3-De-O-methyl-14α-hydroxyisotylocrebrine

[165606-49-5]

$C_{23}H_{25}NO_5$ 395.454

From leaves of *Tylophora tanakae*

(Asclepiadaceae). Mp 210-213° dec.

$[\alpha]_D^{27} +91.2$ (c, 0.35 in CHCl₃/MeOH,

1:1).

14α-Hydroxy, O³-de-Me, N-oxide: 3-De-O-methyl-14α-hydroxyisotylocrebrine N-oxide

[165606-52-0]

$C_{23}H_{25}NO_6$ 411.454

From leaves of *Tylophora tanakae*

(Asclepiadaceae). Mp 215-223° dec.

$[\alpha]_D^{30} +5.5$ (c, 0.12 in MeOH).

14α-Hydroxy, O³,O⁴-di-de-Me, N-oxide: 3,4-Dide-O-methyl-14α-hydroxyisotylocrebrine N-oxide

$C_{22}H_{23}NO_6$ 397.427

Isol. from the butterfly *Ideopsis similis* reared on the host plant, *Tylophora tanakae*.

14 α -Hydroxy, O³,O⁶-di-de-Me: **3,6-Dide-O-methyl-14-hydroxyisotylocrebrine** [209802-55-1]
 C₂₂H₂₃NO₅ 381.427
 Alkaloid from *Tylophora tanakae*.
 Amorph. powder. $[\alpha]_D^{29} +17.4$ (c, 0.3 in MeOH). λ_{\max} 241 (log ϵ 4.3); 260 (log ϵ 4.38); 280 (log ϵ 4.21); 285 (sh) (log ϵ 4.18); 304 (log ϵ 3.89); 316 (log ϵ 3.87) (MeOH).

(±)-form

Synthetic. Cryst. (CHCl₃/MeOH). Mp 219°.

(ξ)-form

O⁴-De-Me: **4-Hydroxy-3,6,7-trimethoxyphenanthroindolizidine**. Alkaloid B[†] [30062-23-8]
 C₂₃H₂₅NO₄ 379.455
 Alkaloid from *Tylophora crebriflora* (Asclepiadaceae). Cryst. (as Ac). Mp 226-228° (Ac).

14ξ-Hydroxy: **14-Hydroxy-3,4,6,7-tetramethoxyphenanthroindolizidine**. Alkaloid A[†] [30062-22-7]
 C₂₄H₂₇NO₅ 409.481
 Alkaloid from *Tylophora crebriflora* (Asclepiadaceae). Cryst. (as Ac). Mp 197-198° (Ac).

14ξ-Hydroxy, O⁴-de-Me: **4,14-Dihydroxy-3,6,7-trimethoxyphenanthroindolizidine**. Alkaloid C[†] [30062-24-9]
 C₂₃H₂₅NO₅ 395.454
 Alkaloid from *Tylophora crebriflora* (Asclepiadaceae). Cryst. (as di-Ac). Mp 216-218° (di-Ac). Stereochem. not determined.

[50657-02-8]

Gellert, E. *et al.*, *J.C.S.*, 1962, 1008-1014 (synth)

Rao, K.V. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1608-1611 (*Tylophora crebriflora constits*)

Govindachari, T.R. *et al.*, *Tetrahedron*, 1973, **29**, 891-897 (isol, uv, pmr, ms, struct)

Govindachari, T.R. *et al.*, *J.C.S. Perkin 1*, 1974, 1161-1165 (ord)

Abe, F. *et al.*, *Phytochemistry*, 1995, **39**, 695-699; 2001, **56**, 697-701 (isol, pmr, *Tylophora tanakae constits*)


Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 767-769 (3,6-Didemethylisotylocrebrines)

Damu, A.G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1071-1075 (*Ficus septica constits*)

Isovariopinsine

I-343

10-Methoxy-N,N-dimethylphenanthro[3,4-d]J-1,3-dioxole-5-ethanamine. 1-(2-Dimethylaminoethyl)-6-methoxy-3,4-methylenedioxyphenanthrene [22137-49-1]



C₂₀H₂₁NO₃ 323.391

Alkaloid from the leaves, twigs and bark


of *Hedycarya angustifolia* (Monimiaceae). Yellow solid. Mp 155-157°.

Geewanda, Y.A. *et al.*, *Heterocycles*, 1987, **26**, 447 (isol, uv, ir, pmr, ms, struct)

Isovoacangine

I-344

[596-54-3]



Absolute Configuration

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Conopharyngia durissima*, *Stemmadenia donnell-smithii* and *Tabernaemontana laurifolia* (Apocynaceae). Shows weak antibacterial and some CNS, brachycardial and hypotensive activity. Cryst. (Et₂O/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 156-157°. $[\alpha]_D^{25} -52$ (CHCl₃). λ_{\max} 227 (ϵ 37150); 270 (sh) (ϵ 6170); 278 (ϵ 4580); 300 (ϵ 7840) (MeOH).

Hydrochloride:

Cryst. (MeOH). Mp 225-228°.

O-De-Me: **11-Hydroxycoronaridine**

[76202-23-8]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the leaves of *Anartia cf. meyeri* (Apocynaceae) (preferred genus name *Tabernaemontana*). Amorph. $[\alpha]_D -34$ (c, 0.6 in CHCl₃).

3-Hydroxy: **3-Hydroxyisovoacangine**

[56867-70-0, 56867-71-1]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from the root bark of *Gabunia eglandulosa* (preferred genus name *Tabernaemontana*) and from *Voacanga* sp. (Apocynaceae). Cryst. (C₆H₆). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 159-160°. $[\alpha]_D -80$ (c, 0.021 in CHCl₃). Mixt. of C-3 epimers. λ_{\max} 226 (log ϵ 4.58); 266 (sh) (log ϵ 3.73); 275 (log ϵ 3.77); 298 (log ϵ 3.93) (EtOH).

3-Oxo: **3-Oxoisovoacangine**

[102488-57-3]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from a callus culture of *Tabernaemontana elegans* (Apocynaceae). λ_{\max} 275 (sh); 288 (sh); 295 (MeOH).

19R-Hydroxy: **19-Epiisovoacristine**

[438201-17-3]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from *Tabernaemontana corymbosa*. Oil. $[\alpha]_D -31$ (c, 0.05 in CHCl₃). λ_{\max} 226 (log ϵ 4.31); 283 (log ϵ 3.64); 294 (log ϵ 3.61) (EtOH).

19S-Hydroxy: **Isovoacristine**

[5525-37-1]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from the bark of *Tabernaemontana laurifolia* (Apocynaceae). Amorph. Mp 104-107° (DMSO solvate) Mp 211-213° (picrate). $[\alpha]_D^{28} -19.6$ (CHCl₃). Forms a DMSO solvate, Mp 104-107°. λ_{\max} 227 (ϵ 32115); 297 (ϵ 5192) (EtOH).

19S-Hydroxy, O-de-Me: **11-Hydroxy-**

heyneanine

[79235-41-9]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from the leaves of *Anartia cf. meyeri* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. $[\alpha]_D -22$ (c, 0.37 in CHCl₃).

19-Oxo: **Isovoacryptine**

[438201-18-4]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Tabernaemontana corymbosa*. Pale yellow oil. $[\alpha]_D +17$ (c, 0.85 in CHCl₃). λ_{\max} 225 (log ϵ 4.25); 280 (log ϵ 3.88); 298 (log ϵ 3.97) (EtOH).

Walls, F. *et al.*, *Tetrahedron*, 1958, **2**, 173-182 (isol, uv, struct)

Renner, U. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 1572-1581 (isol, uv, ir)

Cava, M.P. *et al.*, *Chem. Ind. (London)*, 1965, 2064 (*Isovoacristine*)

Agwada, V.C. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1001-1016 (*3-Hydroxyisovoacangine*)

Ladhar, F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 459-465 (*11-Hydroxyheyneanine*, *11-Hydroxycoronaridine*)

van der Heijden, R. *et al.*, *Phytochemistry*, 1986, **25**, 843-846 (*3-Oxoisovoacangine*)


Soriano-Garcia, M. *et al.*, *J. Crystallogr. Spectrosc. Res.*, 1991, **21**, 681-685 (cryst struct)

Kam, T.-S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 669-672 (*Isovoacryptine*, *19-Epiisovoacristine*)

Isowillardiine

I-345

α -Amino-3,6-dihydro-2,6-dioxo-1(2H)-pyrimidinepropanoic acid. β -(2,4-Dihydroxy-3-pyrimidinyl)alanine



C₇H₉N₃O₄ 199.166

(S)-form [21381-33-9]

From seeds of *Pisum sativum* (peas) (Fabaceae).

Lambein, F. *et al.*, *Biochem. Biophys. Res. Commun.*, 1968, **32**, 474 (isol, uv, struct)

Ashworth, T.S. *et al.*, *Biochem. J.*, 1972, **129**, 897 (biosynth)

Janzen, D.H. *et al.*, *Phytochemistry*, 1977, **16**, 223 (biosynth, tox)


Al-Baldaw, N.F. *et al.*, *CA*, 1980, **93**, 180253m (metab)

Tsuchida, K. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2748 (synth)

De Bruyn, A. *et al.*, *Org. Magn. Reson.*, 1991, **29**, 641 (pmr, cmr)

4-Isoxazolecarboxylic acid

I-346



C₄H₃NO₃ 113.073

Metab. of a *Streptomyces* sp. Phytotoxic. Plates (toluene), cryst. (CHCl₃ or by subl.). Mp 123-124° (121-123°). Dec. at 140° or on keeping. λ_{\max} 212 (E1%/1cm

390) (MeOH) (Berdy). λ_{\max} 215 (E1%/1cm 277) (H₂O) (Berdy).

Et ester: [80370-40-7]

C₆H₇NO₃ 141.126

Liq. Bp₃ 30° Bp_{13.5} 93° (lit. gives a pressure range).

Nitrile: 4-Cyanoisoxazole. 4-Isoxazole-carbonitrile

[68776-58-9]

C₄H₂N₂O 94.073

Sticky pale yellow solid. Mp 63-65°.

Panizzi, L. *et al.*, *Gazz. Chim. Ital.*, 1947, **77**, 206 (*synth, ester*)

Maggioni, P. *et al.*, *Gazz. Chim. Ital.*, 1966, **96**, 443 (*synth, ir*)

Angus, R.O. *et al.*, *Synthesis*, 1988, 746 (*nitrile, synth*)

Kobinata, K. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 1415 (*isol, ir, props*)

3-Isoxazolidinone, 9CI I-347

4,5-Dihydro-3(2H)-isoxazolone. 3-Isoxazolidone

[1192-07-0]



C₃H₅NO₂ 87.078

Isol. from seedlings of *Canavalia ensiformis* (jack bean). Cryst. Mp 70°.

Shunk, C.H. *et al.*, *J.O.C.*, 1957, **22**, 76 (*synth*)
Olive, J.L. *et al.*, *Bull. Soc. Chim. Fr.*, 1976, 1589 (*synth, pmr*)


Sugii, M. *et al.*, *Phytochemistry*, 1981, **20**, 451 (*isol*)

Wolfe, S. *et al.*, *Can. J. Chem.*, 2003, **81**, 937-960 (*synth, pmr*)

4-Isoxazolol I-348

4-Hydroxyisoxazole. *Triumferol*. 4(5H)-Isoxazolone

[80348-66-9]



C₃H₃NO₂ 85.062

OH-form predominates in aq. soln., *4H*-oxo-form in gas phase. Isol. from *Triumfetta rhomboidea*. Antigermination


principle. Cryst. (Et₂O/hexane). Mp 67-68°. λ_{\max} 270 (ε) (MeOH/NaOH) (Derep). λ_{\max} 240 (ε 1100) (MeOH) (Derep).

Kusumi, T. *et al.*, *Tet. Lett.*, 1981, 3451 (*isol, synth*)

Karelson, M.M. *et al.*, *J.C.S. Perkin 2*, 1990, 195 (*tautom*)

5-Isoxazolol I-349

5(4H)-Isoxazolone, 9CI. 2-Isoxazolin-5-one, 8CI. 5-Hydroxyisoxazole. 5(2H)-Isoxazolone [1072-48-6]



C₃H₃NO₂ 85.062

Exists in several tautomeric forms. *2H*-Oxo-form predominates in aq. soln., *4H*-Oxo-form in gas phase. Cryst. Mp 67-69°.

2H-Oxo-form [43228-53-1]

N-β-D-Glucopyranosyl: 2-β-D-Glucopyranosyl-3-isoxazolin-5-one

[51581-00-1]

C₉H₁₃NO₇ 247.204

Constit. of the seedlings of *Lathyrus odoratus* and from the defence secretion of *Chrysomela tremulae*. Amorph. solid.

N-[3-Nitropropanoyl-(→6)-β-D-glucopyranosyl]: 2-[6-(3-Nitropropanoyl)-β-D-glucopyranosyl]-3-isoxazolin-5-one [83566-20-5]

C₁₂H₁₆N₂O₁₀ 348.266

Constit. of the defence secretion of *Chrysomela tremulae*. Amorph. solid.

N-Benzoyl: [174902-45-5]

C₁₀H₇NO₃ 189.17

Cryst. Mp 87-89°.

N-(2-Aminoethyl): 2-(2-Aminoethyl)-3-isoxazolin-5-one

[54019-50-0]

C₅H₈N₂O₂ 128.13

Constit. of the seedlings of *Lathyrus odoratus*.

N-(β-Glutaminylaminoethyl): 2-(β-Glutaminylaminoethyl)-3-isoxazolin-5-one [53987-20-5]

C₁₀H₁₅N₃O₅ 257.246

Constit. of the seedlings of *Lathyrus odoratus*.

OH-form

Me ether: 5-Methoxyisoxazole [31681-57-9]

C₄H₅NO₂ 99.089

Mp 7°. Bp_{0.3} 24-25°.

De Sarlo, F. *et al.*, *J.C.S. (C)*, 1971, 86-89 (*synth, Me ether*)

Teyseyre, J. *et al.*, *J. Mol. Struct.*, 1972, **12**, 191 (*tautom*)

Van Rompuy, L. *et al.*, *Biochem. Biophys. Res. Commun.*, 1974, **56**, 199

Lambien, F. *et al.*, *Biochem. Biophys. Res. Commun.*, 1974, **61**, 155

Kuo, Y.-H. *et al.*, *Plant Physiol.*, 1982, **70**, 1283

Pasteels, J.M. *et al.*, *Tetrahedron*, 1982, **38**, 1891

Ikegami, F. *et al.*, *Phytochemistry*, 1984, **23**, 1567 (*derivs*)


Karelson, M.M. *et al.*, *J.C.S. Perkin 2*, 1990, 195 (*tautom*)

Gould, I.R. *et al.*, *J.C.S. Perkin 2*, 1993, 1771 (*tautom*)

Prager, R.H. *et al.*, *J.C.S. Perkin 1*, 1997, 2659-2664 (*N-Benzoyl, synth, pmr, cmr, ir, ms*)

Itampolin A

I-350



C₃₁H₃₃Br₄N₃O₆ 863.234

Constit. of *Istrochota purpurea*. Pale green oil. [α]_D²¹ +6 (c, 0.66 in MeOH).

27ξ-Hydroxy: *Itampolin B*

C₃₁H₃₃Br₄N₃O₇ 879.234

Constit. of *Istrochota purpurea*. Pale green oil. [α]_D²¹ -2.5 (c, 0.25 in MeOH).

Sorek, H. *et al.*, *Tet. Lett.*, 2006, **47**, 7237-7239 (*isol, pmr, cmr*)

Ivonine

I-351

C₂₉H₃₄N₇O₉S 656.695

Struct. unknown. Alkaloid from the mushroom *Phallus impudicus* (common stinkhorn). Mp 262°.

Stefănescu, P. *et al.*, *Med. Promst. SSSR*, 1960, **14**, 15-17; *CA*, **55**, 11763i

Jaborandine


J-1

$C_{18}H_{28}N_2O_2$ 304.431
Struct. unknown. Alkaloid from *Piper jaborandi* (Piperaceae). Shows local anaesthetic props. pK_a 8.14 (2-Ethoxyethanol). The *p*-Aminobenzoate ester of an aminoalcohol $C_{11}H_{23}NO$, prob. a piperidine.
Hydrochloride: Mp 134°. Sometimes resolidifies and remelts at 154-156°.
Picrolonate: Mp 148-149°.
Wiesner, K. *et al.*, *Can. J. Chem.*, 1951, **29**, 352-354

Jacobine

J-2

15,20-Epoxy-15,20-dihydro-12-hydroxy-senecionan-11,16-dione, 9CI
[6870-67-3]



$C_{18}H_{25}NO_6$ 351.399
Retronecine cyclic diester. Alkaloid of *Senecio jacobaea*, *Senecio brasiliensis* and *Senecio cineraria* (Asteraceae). Plates (EtOH). Mp 228° (219°). $[\alpha]_D^{17}$ -40 (CHCl₃). Samples of higher rotation have been reported including one of -139° (CHCl₃); this may be due to stereomutation.
▶ **Hepatotoxin, genotoxin.** NY5775000
Hydrochloride: Mp 220°. $[\alpha]_D^{19}$ -14.3 (H₂O).


13,19-Didehydro: Jacozine
[5532-23-0]

$C_{18}H_{23}NO_6$ 349.383
Alkaloid from *Senecio jacobaea*, *Senecio alpinus* and *Senecio incanus* (Asteraceae). Tabular prisms (EtOH). Mp 228° sl. dec. (evac. tube). $[\alpha]_D$ -140 (c, 1.90 in CHCl₃).
Bradbury, R.B. *et al.*, *Aust. J. Chem.*, 1954, **7**, 378; 1956, **9**, 258 (*isol, bibl*)
Geissman, T.A. *et al.*, *Aust. J. Chem.*, 1959, **12**, 247 (*struct*)
Bradbury, R.B. *et al.*, *J.A.C.S.*, 1959, **81**, 5201 (*struct*)
Masamune, S. *et al.*, *J.A.C.S.*, 1960, **82**, 5253 (*struct, abs config*)
Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1964, **17**, 233; 1965, **18**, 1625 (*struct, pmr, Jacozine*)
Klásek, A. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 1088 (*isol, Jacozine*)
Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)
Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (*cd*)
Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)
Rohrer, D.C. *et al.*, *Acta Cryst. C*, 1984, **40**, 1449 (*cryst struct*)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, JAK000

Jacoline

J-3

15,20-Dihydro-12,15,20-trihydroxy-senecionan-11,16-dione, 9CI
[480-76-2]



R¹ = OH, R² = H

$C_{18}H_{27}NO_7$ 369.414
Retronecine cyclic diester. Alkaloid from *Senecio jacobaea* (Asteraceae). Cryst. (EtOH). Mp 221.5°. $[\alpha]_D$ +48 (c, 1.155 in CHCl₃).

O¹²-Ac: O-Acetyljacoline
[123403-03-2]
 $C_{20}H_{29}NO_8$ 411.451
Alkaloid from the roots of *Cirsium wallichii* (Asteraceae). Noncryst. $[\alpha]_D^{20}$ +47 (c, 2.1 in EtOH).

O-Ac, picrate: Mp 148-150°.
Bradbury, R.B. *et al.*, *Aust. J. Chem.*, 1954, **7**, 378 (*isol*)
Geissman, T.A. *et al.*, *Aust. J. Chem.*, 1959, **12**, 247 (*struct*)
Bradbury, R.B. *et al.*, *J.A.C.S.*, 1959, **81**, 5201 (*struct, bibl*)
Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)
Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653-3664 (*cd*)
Gable, R.W. *et al.*, *Acta Cryst. C*, 1988, **44**, 1942 (*cryst struct*)
Negi, R.K.S. *et al.*, *Indian J. Chem., Sect. B*, 1989, **28**, 524 (*O-Acetyljacoline*)

Jaconine

J-4

20-Chloro-15,20-dihydro-12,15-dihydroxy-senecionan-11,16-dione, 9CI
[480-75-1]

As Jacoline, J-3 with R¹ = Cl, R² = H
 $C_{18}H_{26}ClNO_6$ 387.859
Alkaloid from *Senecio jacobaea* (Asteraceae). Prisms (EtOH). Mp 147°. $[\alpha]_D$ +52.5 (c, 1.144 in EtOH). MF originally given as $C_{30}H_{32}ClNO_7$.

▶ VT5704000


18-Hydroxy: 18-Hydroxyjacoline

$C_{18}H_{26}ClNO_7$ 403.859
Alkaloid from aerial parts of *Senecio selloi*. No phys. props. reported.
Barger, G. *et al.*, *J.C.S.*, 1937, 584 (*isol*)
Bradbury, R.B. *et al.*, *Aust. J. Chem.*, 1954, **7**, 378 (*isol*)
Geissman, T.A. *et al.*, *Aust. J. Chem.*, 1959, **12**, 247 (*struct*)
Bradbury, R.B. *et al.*, *J.A.C.S.*, 1959, **81**, 5201 (*struct, bibl*)
Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)
Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653-3664 (*cd*)
Gable, R.W. *et al.*, *Acta Cryst. C*, 1988, **44**, 1942 (*cryst struct*)
Krebs, H.C. *et al.*, *Phytochemistry*, 1996, **43**, 1227 (*18-Hydroxyjacoline*)

Jacpaniculine

J-5

[162857-96-7]




$C_{30}H_{33}NO_9$ 551.592
Lignanamide alkaloid from fruits of *Jacquemontia paniculata* var. *paniculata* (Convolvulaceae). Cryst. Mp not reported.

7'-Epimer: Isojacpaniculine
[162990-76-3]
 $C_{30}H_{33}NO_9$ 551.592
Alkaloid from fruits of *Jacquemontia paniculata* var. *paniculata* (Convolvulaceae). Cryst. Mp not reported.
Henrici, A. *et al.*, *Phytochemistry*, 1994, **37**, 1637 (*isol, pmr, cmr, ms, struct*)

Jaculadine

J-6

[27317-49-3]




$C_{17}H_{21}NO_3$ 287.358
Alkaloid from *Croton linearis* (Euphorbiaceae). Noncryst. Mp 201-204° (as tri-Ac). $[\alpha]_D$ -164.7 (c, 0.59 in MeOH) (tri-Ac).
Stuart, K.L. *et al.*, *J.C.S.(C)*, 1970, 1228-1230 (*isol, uv, ir, pmr, ms, struct*)

Jadiffine

J-7

[120374-32-5]



$C_{20}H_{24}N_2O_2$ 324.422
Appears to be a structural variant related to Macroline, M-17. Alkaloid from *Vinca difformis* (Apocynaceae).
Garnier, J. *et al.*, *Phytochemistry*, 1989, **28**, 308 (*pmr, cmr, struct*)

Jadomycin V

J-8

[689281-75-2 (3aR), 689281-68-3 (3aS)]
As Jadomycin Ala, J-10 with


R = -CH(CH₃)₂
 $C_{29}H_{29}NO_6$ 535.549
Metab. of *Streptomyces venezuelae* ISP 5230. Mixt. of diastereoisomers, 67% 3aS, 33% 3aR.

Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (isol, pmr, cmr)
 Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (isol, biosynth)

Jadomycin A

J-9

[138863-47-5]



Absolute Configuration

C₂₄H₂₁NO₆ 419.433

Naphthoquinone antibiotic. Prod. by *Streptomyces venezuelae* ISP5230 (ATCC10712). Green cryst. solid. Mp 167-168°. λ_{max} 214 (ε 37200); 239 (ε 18600); 289 (sh); 314 (ε 21400); 429 (ε 6030) (MeOH) (Derep).

6-O-(2,6-Dideoxy-α-L-ribo-hexopyranoside): **Jadomycin B**

[149633-99-8]

[689281-67-2 (3aS), 689281-74-1 (3aR)]

C₃₀H₃₁NO₉ 549.576

Prod. by *Streptomyces venezuelae* ISP5230 (ATCC10712) after heat shock. Obt. as a mixt. of diastereoisomers at C-3a, R:S 33:67%. λ_{max} 313 (MeOH) (Berdy).

6-O-(6-Deoxy-α-L-altro-hexopyranoside): **ILEVS 1080**

[913188-61-1]

C₃₀H₃₁NO₁₀ 565.576

Isol. from *Streptomyces venezuelae* VS1080. Exists as a mixt. of 3aS and 3aR diastereoisomers.

Ayer, S.W. *et al.*, *Tet. Lett.*, 1991, **32**, 6301-6304 (*Jadomycin A*)

Ayer, S.W. *et al.*, *J. Antibiot.*, 1993, **46**, 869-871 (*Jadomycin B*)

Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (biosynth)

Zheng, J.-T. *et al.*, *J. Antibiot.*, 2005, **58**, 405-408 (activity)


Jakeman, D.L. *et al.*, *Chem. Comm.*, 2006, 3738-3740 (biosynth, ILEVS 1080, isol, pmr)

Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (analogues)

Jadomycin Ala

J-10

[689281-70-7 (3aS), 689281-77-4 (3aR)]

R = CH₃C₂₇H₂₅NO₉ 507.496

Metab. of *Streptomyces venezuelae* ISP 5230. Mixt. of diastereoisomers, 60%

3aS, 40% 3aR.

Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (isol, pmr, cmr)

Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (isol, biosynth)

Jadomycin F

J-11

[689281-69-4 (3aS), 689281-76-3 (3aR)]

As *Jadomycin Ala*, J-10 withR = -CH₂PhC₃₃H₂₉NO₉ 583.593

Metab. of *Streptomyces venezuelae* ISP 5230. Mixt. of diastereoisomers at 3a, 66% 3aS 34% 3aR.

Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (isol, pmr, cmr)

Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (isol, biosynth)

Jadomycin S

J-12

[689281-80-9 (3aR), 689281-73-0 (3aS)]

As *Jadomycin Ala*, J-10 withR = CH₂OHC₂₇H₂₅NO₁₀ 523.495

Metab. of *Streptomyces venezuelae* ISP 5230. Mixt. of diastereoisomers 95% 3aR, 5% 3aS.

Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (isol, pmr, cmr)

Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (isol, biosynth)

Jadomycin T

J-13

[689281-78-5 (3aR), 689281-72-9 (3aS)]

As *Jadomycin Ala*, J-10 withR = -CH(OH)CH₃ (R-)C₂₈H₂₇NO₁₀ 537.522

Metab. of *Streptomyces venezuelae* ISP 5230. Mixt. of diastereoisomers at C-3a 95% 3aR, 5% 3aS.


Rix, U. *et al.*, *J.A.C.S.*, 2004, **126**, 4496-4497 (isol, pmr, cmr)

Borrisow, C.N. *et al.*, *ChemBioChem*, 2007, **8**, 1198-1203 (isol, biosynth)

Jafrine

J-14

[479672-79-2]



Absolute Configuration


C₂₁H₂₈N₂O₃ 356.464

Alkaloid from the flowers of *Tagetes patula*. Yellow gum. [α]_D²⁶ +19.2 (c, 0.1 in CHCl₃).

Faizi, S. *et al.*, *Tetrahedron*, 2002, **58**, 6185-6197 (isol, synth, cd, pmr, cmr, ms)

Jamaicamide A

J-15

C₂₇H₃₆BrClN₂O₄ 567.949

Isol. from *Lyngbya majuscula*. Neurotoxin. Pale yellow oil. [α]_D²⁵ +44 (c, 1.5 in MeOH). λ_{max} 272 (log ε 3.9) (MeOH).

Debromo: Jamaicamide BC₂₇H₃₇ClN₂O₄ 489.053

Isol. from *Lyngbya majuscula*. Pale yellow oil. [α]_D²⁵ +53 (c, 0.61 in MeOH). λ_{max} 272 (log ε 3.9) (MeOH).


Debromo, 19,20-dihydro: Jamaicamide CC₂₇H₃₉ClN₂O₄ 491.069

Isol. from *Lyngbya majuscula*. Pale yellow oil. [α]_D²⁵ +49 (c, 0.39 in MeOH). λ_{max} 273 (log ε 3.8) (MeOH).

Edwards, D.J. *et al.*, *Chem. Biol.*, 2004, **11**, 817-833 (isol, biosynth, pmr, cmr)

Jamine

J-16

Homoomosanine, 9CI

(-)-form

C₂₁H₃₅N₃ 329.528

Stereoisomeric with Homopiptanthine, H-341.

(-)-form [10550-80-8]

Alkaloid from *Ormosia costulata* (Fabaceae) and *Bowdichia virgiloides*, formed by the action of CH₂O on Ormosanine, O-115. Mp 126-127° (116-117°). [α]_D²⁵ -10.4 (c, 1.2 in CHCl₃) (-1.6, -3.8).

18-Epimer: 18-Epihomoomosanine

[123123-43-3]

C₂₁H₃₅N₃ 329.528

Alkaloid from the stem bark of *Bowdichia virgiloides* (Fabaceae). Cryst. Mp 93-95°. [α]_D²¹ +6 (c, 1.03 in CHCl₃).

11,18-Diepimer: Homodasyrcarpine

[67670-84-2]

C₂₁H₃₅N₃ 329.528

Alkaloid from *Ormosia costulata* (Fabaceae). Cryst. (CH₂Cl₂/hexane). Mp 130-140°. [α]_D²⁴ +16 (CHCl₃). Relative config. only known.

(±)-form [4673-41-0]

Alkaloid from seeds of *Ormosia panamensis* and *Ormosia jamaicensis* (Fabaceae). Mp 153-154°.

Clarke, R.T. *et al.*, *J.C.S.*, 1963, 535-539 (*Homodasyrcarpine*)

Karle, I.L. *et al.*, *Tet. Lett.*, 1963, 2065-2067 (cryst struct)

Naegeli, P. *et al.*, *Tet. Lett.*, 1963, 2069-2073 (isol, struct, resoln, synth)

Frank, J.K. *et al.*, *Acta Cryst. B*, 1978, **34**, 2316-2319 (isol, cryst struct)

Wang, A.H.J. *et al.*, *Acta Cryst. B*, 1978, **34**, 2319-2321 (*Homodasyrcarpine*)


Torrengra, G.R. *et al.*, *Planta Med.*, 1985, 276-277 (isol)

Torrengra, R. *et al.*, *Phytochemistry*, 1989, **28**, 2219-2221 (*18-Epihomoomosanine*)

Le, P.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 283-293 (pmr, cmr)

Jamtine

[111261-78-0]

C₂₀H₂₅NO₄ 343.422

Synthetic studies (2003) raise serious doubts about the structural assignments in this series. λ_{\max} 220 ; 239 ; 300 (MeOH).

N-Oxide: Jamtine N-oxide

[111261-79-1]

C₂₀H₂₅NO₅ 359.421

Alkaloid from the leaves of *Cocculus hirsutus* (Menispermaceae). Viscous oil.

O³-De-Me: Hirsutine†

[135250-40-7]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Cocculus hirsutus* (Menispermaceae). Gum. $[\alpha]_D^{25}$ +178 (CHCl₃).

O²,O³-Di-de-Me: Haiderine. Jamtinine†

[154722-71-1]

C₁₈H₂₁NO₄ 315.368

Alkaloid from aerial parts (Haiderine) or whole plants (Jamtinine) of *Cocculus hirsutus* (Menispermaceae). Gum. $[\alpha]_D^{25}$ +167 (CHCl₃). Haiderine and Jamtinine not compared, but same struct. assigned. The name Jamtinine is misleading as this name was previously used for 6-Oxojamtine.

6-Oxo: Jamtinine†

[151079-01-5]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Cocculus hirsutus* (Menispermaceae). Gum. $[\alpha]_D^{25}$ +177 (CHCl₃).

Ahmad, V.U. *et al.*, *Heterocycles*, 1987, **26**, 1251-1255 (*Jamtine N-oxide*)

Rasheed, T. *et al.*, *J. Nat. Prod.*, 1991, **54**, 582-584 (*Hirsutine*)

Ahmad, V.U. *et al.*, *Nat. Prod. Lett.*, 1993, **2**, 105-109 (*Haiderine*)

Ahmad, V.U. *et al.*, *Phytochemistry*, 1993, **33**, 735-736 (*Jamtinine*)


Durrani, S. *et al.*, *Fitoterapia*, 1995, **66**, 172 (*Jamtinine*)

Padwa, A. *et al.*, *J.O.C.*, 2003, **68**, 929-941 (*synth, struct*)

Gill, C.D. *et al.*, *Tetrahedron*, 2003, **59**, 9213-9230 (*synth, struct*)

Janolusimide

[103612-45-9]



Absolute Configuration


C₁₉H₃₃N₃O₅ 383.487**J-17**

Peptide toxin. Isol. from *Janolus crinitatus*. Neurotoxin, atropine antagonist. Amorph. $[\alpha]_D^{25}$ -10.3 (c, 2.5 in CHCl₃).

▶ LD₅₀ (mus, ipr) 5 mg/kg.

Sodano, G. *et al.*, *Tet. Lett.*, 1986, **27**, 2505-2508 (*isol, struct, nmr*)

Giordano, A. *et al.*, *Tet. Lett.*, 2000, **41**, 3979-3982 (*synth, abs config*)

Janoxepin**J-19**

Absolute Configuration

C₁₉H₂₃N₃O₃ 341.409

Homologue of Cinereain, C-449. Prod. by *Aspergillus janus* (IBT 22274). Antiplasmodial agent. Orange-red cryst. Mp 88-89°. $[\alpha]_D^{25}$ -3.4 (c, 1.5 in MeOH).

Sprogog, K. *et al.*, *Tetrahedron*, 2005, **61**, 8718-8721 (*isol, pmr, cmr*)

Janthitrem A**J-20**

[73561-89-4]

C₃₇H₄₇NO₆ 601.781

Struct. unknown. Prod. by *Penicillium janthinellum*. Tremorgenic mycotoxin. Fluorescent.


Gallagher, R.T. *et al.*, *Appl. Environ. Microbiol.*, 1980, **39**, 272-273 (*isol*)

Lauren, D.R. *et al.*, *J. Chromatogr.*, 1982, **248**, 150-154

Penn, J. *et al.*, *Phytochemistry*, 1993, **32**, 1431-1434 (*isol, ms*)

Janthitrem E**J-21**

[90986-50-8]

C₃₇H₄₉NO₆ 603.797

Rel. stereochem. at C-22 and C-23 depicted. Prod. by *Penicillium janthinellum*. Tremorgenic mycotoxin. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 228 (ε 17700); 265 (ε 30000); 330 (ε 17000) (MeOH) (Berdy).

10-Ac: Janthitrem F

[90986-52-0]

C₃₉H₅₁NO₇ 645.834

From *Penicillium janthinellum*. Tremorgenic mycotoxin. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 228 (ε 24200); 265 (ε 22400); 330 (ε 12200) (MeOH) (Berdy).

22-Deoxy, 10-Ac: Janthitrem G

[90986-51-9]

C₃₉H₅₁NO₆ 629.835

From *Penicillium janthinellum*. Tremorgenic mycotoxin. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 228 (ε 23000); 263 (ε 15700); 331 (ε 10800) (MeOH) (Berdy).

22-Deoxy, 10-ketone: Shearinine B

[163136-25-2]

C₃₇H₄₇NO₅ 585.782

Isol. from the ascostromata of *Eupeenicillium shearii*. Mycotoxin. Shows antiinsectant activity. Mp 165-170° dec. $[\alpha]_D$ -76 (c, 0.20 in CHCl₃). C-23 config. not defined.

34-Deoxy, 34,35-didehydro: Janthitrem B

[73561-90-7]

C₃₇H₄₇NO₅ 585.782

Isol. from *Penicillium janthinellum*. Tremorgenic mycotoxin.

22,34-Dideoxy, 34,35-didehydro: Janthitrem C

[73561-91-8]

C₃₇H₄₇NO₄ 569.783

Isol. from *Penicillium janthinellum*. Tremorgenic mycotoxin. Amorph. solid.

De Jesus, A.E. *et al.*, *J.C.S. Perkin 1*, 1984, 697 (*isol, uv, ir, pmr, cmr, struct*)


Wilkins, A.L. *et al.*, *J. Agric. Food Chem.*, 1992, **40**, 1307 (*struct, config*)

Penn, J. *et al.*, *Phytochemistry*, 1993, **32**, 1431 (*struct*)

Belofsky, G.N. *et al.*, *Tetrahedron*, 1995, **51**, 3959 (*Shearinine B*)

Janussine A**J-22**

[97466-24-5]

C₃₀H₃₂N₄O 464.609

Alkaloid from the root bark and stem bark of *Strychnos johnsonii* (Loganiaceae). $[\alpha]_D$ +492 (c, 0.5 in MeOH).

Stereoisomer: Janussine B

[97400-83-4]

C₃₀H₃₂N₄O 464.609

Alkaloid from the root bark and stem bark of *Strychnos johnsonii* (Loganiaceae). $[\alpha]_D$ +153 (c, 0.2 in MeOH).


Massiot, G. *et al.*, *Tet. Lett.*, 1985, **26**, 2441-2444 (*uv, ir, pmr, cmr, ms, struct*)

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (*isol, uv, ir, pmr, ms*)

Japindine

J-23

N,N'-Bis(20-dimethylaminopregnan-3-yl)-*N*-methylthiourea, 9CI
[52483-07-5]



$C_{48}H_{84}N_4S$ 749.284


Alkaloid from root bark of *Chonemorpha macrophylla* (Apocynaceae). Mp 275–280° dec. subl. $[\alpha]_D^{25} +5.4$ (MeOH/CHCl₃ 1:4). Yields Chonemorphine and *N*-Acetylchonemorphine on acetylation.

Banerji, J. *et al.*, *Indian J. Chem., Sect. B*, 1978, **16**, 346

Jasminidine

J-24

1,4-Dihydro-1-methyl-2,7-naphthyridin-3(2H)-one



(S)-form

$C_9H_{10}N_2O$ 162.191

(S)-form [68711-43-3]

Alkaloid from the leaves of *Syringa vulgaris* (Oleaceae). Cryst. (CHCl₃/Et₂O). Mp 190–194° dec. $[\alpha]_D^{25} -3.2$ (c, 0.35 in CHCl₃).

(ξ)-form

5-Methoxy: *Acanthicifoline*

[76193-60-7]

 $C_{10}H_{12}N_2O_2$ 192.217

Alkaloid from *Acanthus ilicifolius* (Acanthaceae). Mp 168–170°. $[\alpha]_D^{24} -31.6$ (CHCl₃).


Ripperger, H. *et al.*, *Phytochemistry*, 1978, **17**, 1069–1070 (*Jasminidine*)

Tiwara, K.P. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1980, **54**, 857–858 (*Acanthicifoline*)

Jasminine†

J-25

Methyl 5,6,7,8-tetrahydro-8-methyl-6-oxo-2,7-naphthyridine-4-carboxylate, 9CI



$C_{11}H_{12}N_2O_3$ 220.227

(S)-form [19634-30-1]

Alkaloid from *Jasminum gracile* and several other *Jasminum* spp., *Ligustrum novoguineense*, *Olea paniculata* and *Syringa vulgaris* (Oleaceae). Mp 175–176°. $[\alpha]_D^{24} -29.8$ (c, 1 in CHCl₃).

1,4-Dihydro: Dihydrojasminine

[206440-44-0]

 $C_{11}H_{14}N_2O_3$ 222.243

Alkaloid from *Osmanthus austrocaledonica*. $[\alpha]_D^{20} +300$ (c, 0.01 in CHCl₃). Rel. config. only determined. 4*H*-*cis*-to Me group. λ_{max} 206 (log ϵ 4.27); 344 (log ϵ 3.83) (MeOH).

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1968, **21**,

1321–1326 (*uv, ir, pmr, ms, isol, struct*)

Ripperger, H. *et al.*, *Phytochemistry*, 1978, **17**,

1069–1070 (*ir, pmr, ms, config*)

Benkrief, R. *et al.*, *Phytochemistry*, 1998, **47**,

825–832 (*Dihydrojasminine, isol, uv, ir, pmr, cmr*)


Bennasar, M.-L. *et al.*, *Tetrahedron*, 2004, **60**,

6785–6789 (*synth*)**Jaspamide**

J-26

Jaspakinolide

[102396-24-7]



Absolute Configuration

$C_{36}H_{45}BrN_4O_6$ 709.679

Cyclodepsipeptide antibiotic. Isol. from the marine sponge *Jaspis* sp. F-actin inhibitor. Active against yeasts, nematodes and carcinoma. Possesses antifungal, anthelmintic and insecticidal props. Cytotoxic. $[\alpha]_D +35$ (c, 3.62 in MeOH). λ_{max} 281 (ϵ 5400); 290 (ϵ 4100) (MeOH) (Derep).

31-Hydroxy: Jaspamide E

[942155-33-1]

 $C_{36}H_{45}BrN_4O_7$ 725.678

Isol. from *Jaspis splendans*. Amorph. solid. $[\alpha]_D^{25} +42.2$ (c, 0.05 in CHCl₃).

4Z-Isomer, 35-oxo: Jaspamide G

[942155-35-3]

 $C_{36}H_{43}BrN_4O_7$ 723.662

Isol. from *Jaspis splendans*. Amorph. solid. $[\alpha]_D^{25} -6.7$ (c, 0.06 in CHCl₃). Has 35*E*-config.

Δ^{4,35}-Isomer, 5ξ-hydroxy: Jaspamide C

[219774-75-1]

 $C_{36}H_{45}BrN_4O_7$ 725.678

Isol. from *Jaspis splendans*. Glass. $[\alpha]_D^{25} +25.4$ (c, 0.0013 in CHCl₃). λ_{max} 220 (log ϵ 3.37); 276 (log ϵ 2.71) (MeOH).

Δ^{4,35}-Isomer, 5-oxo: Jaspamide B

[219774-74-0]

 $C_{36}H_{43}BrN_4O_7$ 723.662

Isol. from *Jaspis splendans*. Glass. $[\alpha]_D^{25} +11.4$ (c, 0.0014 in CHCl₃). λ_{max} 232 (log ϵ 3.75); 278 (log ϵ 3.66) (MeOH).

31-Methyl, 15-epimer: Jaspamide D

[942155-32-0]

 $C_{37}H_{47}BrN_4O_6$ 723.706

Isol. from *Jaspis splendans*. Amorph. solid. $[\alpha]_D^{25} +20.1$ (c, 0.05 in CHCl₃).

4-Demethyl: Jaspamide F

[942155-34-2]

 $C_{35}H_{43}BrN_4O_6$ 695.652

Isol. from *Jaspis splendans*. Amorph. solid. $[\alpha]_D^{25} -15.7$ (c, 0.07 in CHCl₃).

Zabriskie, T.M. *et al.*, *J.A.C.S.*, 1986, **108**,

3123 (*isol, ir, pmr, cmr, cryst struct*)

Crews, P. *et al.*, *Tet. Lett.*, 1986, **27**, 2797 (*isol, uv, ir, pmr, cmr, ms, struct*)

Braekman, J.C. *et al.*, *J. Nat. Prod.*, 1987, **50**,

994 (*isol, pmr, cmr, ir, uv, ms*)

Grieco, P.A. *et al.*, *J.A.C.S.*, 1988, **110**, 1630

(synth)

Kato, S. *et al.*, *Tet. Lett.*, 1988, **29**, 6465 (*synth*)

Inman, W. *et al.*, *J.O.C.*, 1989, **54**, 2523 (*props, bibl*)

Konopelski, J.P. *et al.*, *J.O.C.*, 1991, **56**, 1355

(synth)

Kang, S.K. *et al.*, *Synlett*, 1991, 175 (*synth*)

Rao, A.V.R. *et al.*, *Tet. Lett.*, 1993, **34**, 7085

(synth)

Hirai, Y. *et al.*, *Heterocycles*, 1994, **39**, 603

(synth)

Imaeda, T. *et al.*, *Tet. Lett.*, 1994, **35**, 591

(synth)

Ashworth, P. *et al.*, *Synthesis*, 1995, 199 (*synth*)

Zampella, A. *et al.*, *J. Nat. Prod.*, 1999, **62**,

332–334 (*Jaspamides B-C*)

Tabudravu, J.N. *et al.*, *Org. Biomol. Chem.*,

2005, **3**, 745–749 (*conformn*)


Gosh, A.K. *et al.*, *Org. Lett.*, 2007, **9**, 2425–

2427 (*synth*)

Gala, F. *et al.*, *Tetrahedron*, 2007, **63**, 5212–

5219 (*Jaspamides D-G*)**6(17),11-Jatrofadiene-2,3,5,7,8,9,14,15-octol**

J-27



$C_{20}H_{34}O_8$ 402.484

(2α,3β,5α,7β,8α,9α,11E,14β,15β)-form*9-(3-Pyridinecarbonyl), 3-benzoyl,**2,5,7,14-tetra-Ac*: [210108-89-7] $C_{41}H_{49}NO_{14}$ 779.836Constit. of *Euphorbia peplus*.*9-(3-Pyridinecarbonyl), 3-benzoyl, 7-(2-**methylpropanoyl), 2,5,14-tri-Ac*:


[210108-87-5]

 $C_{43}H_{53}NO_{14}$ 807.89Constit. of *Euphorbia peplus*.

Jakupovic, J. *et al.*, *Phytochemistry*, 1998, **47**,

1601–1609 (*isol, pmr, cmr*)**Javacarboline**

J-28



C₂₀H₂₀N₂O₂ 320.39

Truncated indole alkaloid, original skeleton uncertain. Prob. not secologanin-derived.


(±)-**form** [157232-63-8]

Alkaloid from the stem of *Picrasma javanica* (Simaroubaceae). Exhibits *in vitro* cytotoxicity against human tumour PC-6 cells and murine lymphocytic leukaemia P-388 cells. Pale yellow prisms (MeOH aq.). Mp 220-222°. λ_{max} 235 (ε 13500); 308 (ε 8710); 380 (ε 10471) (MeOH) (Berdy).

Koike, K. *et al.*, *Heterocycles*, 1994, **38**, 1413 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Javanicunine A

[915137-81-4]



Absolute Configuration

C₂₄H₃₀N₂O₄ 410.512

Similar to Mollenine A, M-670. Prod. by *Eupenicillium javanicum* IFM 54704. Amorph. solid. [α]_D²² -152.3 (c, 1 in CH₂Cl₂). λ_{max} 247 (log ε 3.95); 278 (log ε 3.14); 285 (log ε 3.09) (CH₂Cl₂).

10a-Epimer, 10a-hydroxy: **Javanicunine B** [915137-82-5]

C₂₄H₃₀N₂O₅ 426.511

Prod. by *Eupenicillium javanicum* IFM 54704. Amorph. solid. [α]_D²² -118.5 (c, 0.2 in CH₂Cl₂). λ_{max} 246 (log ε 3.87); 278 (log ε 3.13); 285 (log ε 3.09) (MeOH).

Nakadate, S. *et al.*, *Heterocycles*, 2006, **68**, 1969-1972 (*isol, pmr, cmr*)

Javanine†


J-30

Struct. unknown. Alkaloid from the bark of *Cinchona calisaya* var. *javanica* (Rubiaceae). Rhombic leaflets (H₂O).

Hesse, O. *et al.*, *Ber.*, 1877, **10**, 2152

Javaniside

J-31



Absolute Configuration


C₂₆H₃₀N₂O₉ 514.531

Alkaloid from the leaves of *Alangium javanicum*. DNA cleavage agent. Amorph. powder. [α]_D²¹ -50.2 (c, 0.2 in MeOH).

Pham, V.C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1147-1152 (*isol, pmr, cmr, ms*)

Javaphylline

[17138-12-4]



Absolute Configuration

C₂₂H₂₆N₂O₅ 398.458

Alkaloid from *Mitragyna javanica* (Rubiaceae). Mp 180°.

Shellard, E.J. *et al.*, *Planta Med.*, 1967, **15**, 245
Shellard, E.J. *et al.*, *J. Chromatogr.*, 1968, **32**, 472; 1975, **105**, 163

Jaxartinine

J-33


C₁₀H₁₅NO 165.235

Struct. unknown. Alkaloid from *Anabasis jaxartica* (Chenopodiaceae). Cryst. (Me₂CO). Mp 166-169°.

Platonova, T.F. *et al.*, *Zh. Obshch. Khim.*, 1958, **28**, 3128-3131; *CA*, **53**, 7506e

Jenamidine B

[677756-96-6]

C₁₃H₁₈N₂O₄ 266.296

Structs. revised in 2004. Formerly assigned as pyridopyrimidines. Prod. by *Streptomyces* sp. strain HKI0297. Light yellowish oil. [α]_D²⁵ +8.4 (c, 0.6 in MeOH). Isol. as a 1:1 mixt. of stereoisomers B₁/B₂. λ_{max} 228 (log ε 4.12); 299 (log ε 3.08); 333 (log ε 3.26) (MeOH).

4'-Deoxy, 6'-hydroxy: **Jenamidine C**

[677756-98-8]

C₁₃H₁₈N₂O₄ 266.296

Prod. by *Streptomyces* sp. strain HKI0297. Light yellowish gum. [α]_D²⁵ +1.8 (c, 0.39 in MeOH). Isol. as a 1:1 mixt. of stereoisomers C₁/C₂. λ_{max} 228 (log ε 4.23); 248 (log ε 3.53); 299 (log ε 3.13); 332 (log ε 3.48) (MeOH).

7a-Deoxy: **Jenamidine A**

[677756-94-4]

C₁₃H₁₈N₂O₃ 250.297

Prod. by *Streptomyces* sp. strain HKI0297. Antiproliferative agent against chronic myeloid leukaemic cells. Light yellowish oil. [α]_D²⁵ +6.8 (c, 0.7 in MeOH). Isol. as a 1:1 mixt. of stereoisomers A₁/A₂. λ_{max} 240 (log ε 4.01); 280 (log ε 3.11); 326 (log ε 3.32) (MeOH).

Hu, J.-F. *et al.*, *J. Antibiot.*, 2003, **56**, 747-754 (*isol, pmr, cmr, ms*)


Snider, B.B. *et al.*, *Tet. Lett.*, 2004, **45**, 6725-6727 (*pmr, cmr, struct*)

Duvall, J.R. *et al.*, *J.O.C.*, 2006, **71**, 8579-8590 (*synth, struct*)

J-32

Jerusalemine

J-35

C₁₉H₂₃NO₅ 345.394

The 9CI name is defective.

(S)-**form** [138704-11-7]

Alkaloid from *Colchicum decaisnei* (Liliaceae). Amorph. solid. [α]_D -35 (c, 0.17 in MeOH).


Abu Zarga, M.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 936 (*isol, uv, ir, pmr, cmr, ms, struct*)

Banwell, M.G. *et al.*, *Chem. Comm.*, 1994, 2647 (*synth*)

Jervine

J-36

17,23-Epoxy-3-hydroxyveratraman-11-one. *Jervin-11-one* [469-59-0]



Absolute Configuration

C₂₇H₃₉NO₃ 425.61

Alkaloid from *Veratrum album* var. *grandiflorum*, *Veratrum californicum*, several other *Veratrum* spp. and *Amianthium muscaetoxicum* (Liliaceae). Starting material for synth. of c-nor-D-homosteroids. Shows antibacterial and antifungal activity. Severe teratogen responsible for cyclopic malformation in sheep grazing on *V.* spp. Antihypertensive agent. Mp 243-244°. [α]_D -147 (EtOH). λ_{max} 250 (ε 16400); 252 (ε 14000); 300 (ε 80); 360 (ε 70) (EtOH) (Berdy).

▶ Reprod. and teratogenic effects; LD₅₀ (mus, scu) 29 mg/kg; LD₅₀ (mus, ivn) 9.3 mg/kg. WG9700000

Hydrochloride: Mp 308°.

O-β-D-Glucopyranoside: **Pseudojervine** [36069-05-3]

C₃₃H₄₉NO₈ 587.752

Alkaloid from *Veratrum album* and some other *Veratrum* spp. (Liliaceae). Shows antifungal activity. Sol. CHCl₃, C₆H₆; fairly sol. EtOH; poorly sol. Et₂O. Mp 300-301°. [α]_D -131 (CHCl₃/EtOH 1:1). [α]_D²⁵ -97 (c, 0.12 in CHCl₃/EtOH).

▶ LD₅₀ (mus, ivn) 50-150 mg/kg. NY9062000

O-Ac: **O-Acetyljervine**

[14788-78-4]

C₂₉H₄₁NO₄ 467.647

Alkaloid from the leaves of *Veratrum album* (Liliaceae). Powder. Mp 240°. [α]_D -120 (MeOH).

▶ Exp. reprod. and teratogenic effects. NY8900000

N-Methoxycarbonyl: **Verapatuline**

C₂₉H₄₁NO₅ 483.647
Alkaloid from *Veratrum patulum*.
Amorph. solid. [α]_D²³ -49.9 (c, 0.3 in EtOH). λ_{max} 252 (log ε 4.25) (EtOH).

N-(2-Methoxycarbonylethyl): **Methyl jervine-N-3'-propanoate** [132943-48-7]
Alkaloid from the leaves of *Veratrum album* (Liliaceae). Powder. Mp 234°. [α]_D -76 (MeOH).

3-Ketone: **Jervinone**
C₂₇H₃₇NO₃ 423.594
Alkaloid from rhizomes of *Veratrum album* (Liliaceae). Amorph. powder. Mp 246-248°. [α]_D²⁵ +38.5 (c, 1.00 in CHCl₃).

11α-Alcohol, N-methoxycarbonyl: **Neoverapatuline** [944455-07-6]
C₂₉H₄₃NO₅ 485.662
Alkaloid from *Veratrum nigrum*. Powder (CHCl₃). Mp 155-156°. [α]_D²⁰ -6 (c, 0.01 in CHCl₃).

11β-Alcohol: **Veratrobazine**. *Jervin-11β-ol* [20226-97-5]
C₂₇H₄₁NO₃ 427.626
Alkaloid from *Veratrum album* and *Veratrum grandiflorum* (Liliaceae). Shows antifungal activity. Large prisms (C₆H₆/MeOH). Mp 285-288° (dec. from 270°). [α]_D -76.6 (c, 0.58 in 95% EtOH). [α]_D²⁰ -126 (c, 0.65 in Py).

► Teratogenic.

11β-Alcohol, 3,11-di-Ac: Mp 208-210°. [α]_D²⁰ -77 (c, 0.46 in Py).

Deoxo: **Cyclopamine**. *Alkaloid V†. 11-Deoxojervine* [4449-51-8]
C₂₇H₄₁NO₂ 411.626
Isol. from *Veratrum californicum* and *Veratrum album* (Liliaceae). Severe teratogen responsible for cyclopic malformation in sheep grazing on *V. californicum*. Needles (EtOH). Mp 237-238°. [α]_D²⁵ -48 (c, 1 in MeOH/CHCl₃ 2:1).

► Exp. reprod. and teratogenic effects. GY0750000

Deoxo, 3-O-β-D-glucopyranoside: **Cycloposine** [23185-94-6]
C₃₃H₅₁NO₇ 573.768
Alkaloid from *Veratrum californicum* (Liliaceae). Teratogen causing cyclopic malformation in sheep grazing on *V. californicum*. Mp 267-269°. [α]_D²⁵ -51 (c, 1 in EtOH). Acid hydrol. produces Veratramine, V-71 by aromatisation.

► Exp. teratogen.

Deoxo, O,N-di-Ac: Mp 195-197°. [α]_D²⁵ -5 (EtOH). A lower addnl. Mp at 163-164° also reported.

1α-Hydroxy, 5α,6-dihydro: **5α,6-Dihydro-1α-hydroxyjervine**
C₂₇H₄₁NO₄ 443.625
Alkaloid from rhizomes of *Veratrum album* (Liliaceae). Amorph. powder. Mp 245°. [α]_D -70 (CHCl₃).

1β-Hydroxy, 5β,6-dihydro: **5β,6-Dihydro-1β-hydroxyjervine**

[944358-35-4]
C₂₇H₄₁NO₄ 443.625
Alkaloid from *Veratrum nigrum*. Powder (CHCl₃). Mp 239-240°. [α]_D²⁰ -57 (c, 0.01 in MeOH/CHCl₃).

23α-Methoxy, deoxo: **23-Methoxy-11-deoxojervine. 23-Methoxycyclopamine**
C₂₈H₄₃NO₃ 441.653
Alkaloid from *Veratrum nigrum*. Amorph. yellow powder. [α]_D²⁵ +44 (c, 0.05 in CHCl₃).

Jacobs, W.A. et al., *J. Biol. Chem.*, 1947, **170**, 635 (*uv*)

Stoll, A. et al., *Helv. Chim. Acta*, 1955, **38**, 1964 (*Veratrobazine*)

Budzikiewicz, H. et al., *Tetrahedron*, 1964, **20**, 2267 (*ms*)

Masamune, T. et al., *J.A.C.S.*, 1967, **89**, 4521 (*synth*)

Reeke, G.N. et al., *J.A.C.S.*, 1968, **90**, 1663 (*Veratrobazine, cryst struct*)

Kupchan, S.M. et al., *J.A.C.S.*, 1968, **90**, 2730 (*abs config*)

Keeler, R.F. et al., *Phytochemistry*, 1969, **8**, 223 (*Cyclopamine*)

Keeler, R.F. et al., *Steroids*, 1969, **13**, 579 (*Cycloposine*)

Wolters, B. et al., *Planta Med.*, 1970, **19**, 189-193 (*activity*)

Sprague, P.W. et al., *Tetrahedron*, 1971, **27**, 4857 (*cmr*)

Bryden, M.M. et al., *CA*, 1973, **79**, 143407e (*tox*)

Han, Y.B. et al., *Yakhhak Hoeji*, 1973, **17**, 13-16; 137-140 (*activity*)

Kutney, J.P. et al., *Can. J. Chem.*, 1975, **53**, 1796 (*synth*)

Keeler, R.F. et al., *Alkaloids: Chem. Biol. Perspect.*, 1984, **4**, 389 (*rev, tox*)

Atta-ur-Rahman, et al., *Phytochemistry*, 1991, **30**, 368 (*O-Acetyljervine, Methyl jervine-N-3'-propanoate*)

Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1992, **55**, 565 (*Jervinone*)

Atta-ur-Rahman, et al., *Planta Med.*, 1993, **59**, 569 (*5α,6-Dihydro-1α-hydroxyjervine*)

Atta-ur-Rahman, et al., *Phytochemistry*, 1996, **43**, 907 (*Pseudojervine, bibl*)

Tezuka, Y. et al., *J. Nat. Prod.*, 1998, **61**, 1078-1081 (*Verapatuline*)


Cong, Y. et al., *Helv. Chim. Acta*, 2007, **90**, 1038-1042 (*5β,6-Dihydro-1β-hydroxyjervine, Neoverapatuline*)

Wang, B. et al., *Helv. Chim. Acta*, 2008, **91**, 244-248 (*23-Methoxycyclopamine*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, CPT750; JDA000; JCS000

Jhelumine J-37

[85588-85-8]



C₃₆H₃₈N₂O₇ 610.705
Alkaloid from *Berberis lycium* (Berberidaceae). [α]_D²⁵ +28 (c, 0.6 in MeOH).

O⁷-Me: **Chenabine**

[85588-86-9]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from *Berberis lycium*. [α]_D²⁵

+40 (c, 0.18 in MeOH).

α'-Alcohol: **Chenabinol**

[122148-81-6]

C₃₇H₄₂N₂O₇ 626.748

Obt. by NaBH₄ reduct. of Chenabine (see Jhelumine, J-37).

α'-Alcohol, Me ether: **Chenabinol methyl ether**

[122127-75-7]

C₃₈H₄₄N₂O₇ 640.775


Alkaloid from *Berberis brandisiana*. Amorph. [α]_D +21 (c, 0.11 in MeOH).

Leet, J.E. et al., *Heterocycles*, 1983, **20**, 425-429 (*Jhelumine, Chenabine*)

Hussain, S.F. et al., *J. Nat. Prod.*, 1989, **52**, 317-319 (*Chenabinol, Chenabinol methyl ether*)

Jietacin A J-38

[109766-61-2]



C₁₈H₃₄N₂O₂ 310.479

Prod. by *Streptomyces* sp. (KP197; P8889). Exhibits potent nematocidal props. Possesses weak activity against some fungi. Sol. DMSO, hexane, EtOAc, Me₂CO; poorly sol. H₂O, MeOH. λ_{max} 228 (ε 3650); 250 (sh) (ε 2400) (cyclohexane) (Derep). λ_{max} 218 (ε 3650) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 200 - 600 mg/kg. RZ2020000

Omura, S. et al., *J. Antibiot.*, 1987, **40**, 623 (*isol, props*)

Imamura, N. et al., *J. Antibiot.*, 1989, **42**, 156 (*struct*)

Giencke, W. et al., *Annalen*, 1989, 671 (*synth*)

Jietacin B J-39

[109766-62-3]

As Jietacin A, J-38 with

n = 6

C₁₉H₃₆N₂O₂ 324.506

From *Streptomyces* sp. (KP197; P8889). Similar biol. props. as Jietacin A, J-38. Sol. DMSO, hexane, EtOAc, Me₂CO; poorly sol. H₂O, MeOH. λ_{max} 228 (ε 3650); 250 (sh) (ε 2400) (cyclohexane) (Derep). λ_{max} 228 (ε 3650) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 200 - 600 mg/kg.

Omura, S. et al., *J. Antibiot.*, 1987, **40**, 623 (*isol, props*)

Imamura, N. et al., *J. Antibiot.*, 1989, **42**, 156 (*struct*)

Jingsimycin J-40

[75718-53-5]


Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptomyces hygrosopicus*. Yellow powder. Sol. MeOH, H₂O, DMF. [α]_D +42 (H₂O). λ_{max} 222; 270 (MeOH). λ_{max} 222; 270 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 1000 - 3000 mg/kg.

Lu, W.-Z. et al., *CA*, 1981, **94**, 2894j

Jiufengsine

Jiufengsine
[737792-40-4]




$C_{38}H_{54}N_2O_{11}$ 714.851

Deriv. of 4-Methylaconitane-1,6,7,8,14,16,18-heptol, M-309. Alkaloid from the roots of *Delphinium potaninii* var. *jiufengshanense*. Mp 85-87°. $[\alpha]_D^{29} +22.8$ (c, 1 in $CHCl_3$).

Shen, X.-L. et al., *Zhongguo Tianran Yaowu*, 2004, **2**, 152-154; *CA*, **141**, 187848 (*Jiufengsine*)

Joalin

[150999-04-5]




$C_{21}H_{31}NO_3$ 345.481

Constit. of a *Dictyota* sp. Semicryst. $[\alpha]_D^{29} -59$ (c, 0.01 in $CHCl_3$). λ_{max} 205 (ε 12300); 245 (MeOH) (Berdy).

Guella, G. et al., *J.C.S. Perkin I*, 1993, 1545 (*isol*, *pmr*, *cmr*)

Jolantamine

Alkaloid K8. Alkaloid MJ1. *Iolantamine*
[37376-03-7]



$C_{19}H_{23}NO_3$ 313.396

Alkaloid from *Colchicum kesselringii* and *Merendera jolantae* (Liliaceae). Mp 215-216° (193-197°). $[\alpha]_D +112$ ($CHCl_3$). Same plain struct. as Bulbocodine, B-398. Jolantamine is said to have the 6aR config. (same as Bulbocodine) and therefore if a different alkaloid must be the 8a-epimer. Evidence for its non-identity with Bulbocodine is however weak and they have not been directly compared.

Hydrochloride: Mp 248-250°.

N-Me: *Jolantamine*. *Iolantamine*
[62249-76-7]

$C_{20}H_{26}NO_3^{\oplus}$ 328.43

Quaternary alkaloid from *Merendera jolantae* (Liliaceae). Cryst. (Me_2CO)

J-41

MeOH)(as hydroxide). Mp 269-270° (hydroxide).

Me ether: *Crociflorinone*

[55437-98-4]

$C_{20}H_{25}NO_3$ 327.422

Alkaloid from *Colchicum kesselringii* (Liliaceae). Cryst. (Me_2CO/Et_2O). Mp 246-248°.

Yusupov, M.K. et al., *J. Gen. Chem. USSR*


(*Engl. Transl.*), 1964, **34**, 1672 (*isol*)

Yusupov, M.K. et al., *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1973, **208**, 1123; *CA*, **79**, 79010r (*uv*, *pmr*, *struct*)

Turdikulov, K. et al., *Khim. Prir. Soedin.*, 1974, **10**, 810; 1976, **12**, 555; *Chem. Nat. Compd.* (*Engl. Transl.*), 1974, **10**, 844; 1976, **12**, 501 (*Crociflorinone*, *Jolantamine*)

Jolantimine

Iolantimine
[60142-20-3]



$C_{18}H_{21}NO_4$ 315.368


Alkaloid from *Colchicum kesselringii* and *Merendera jolantae* (Liliaceae). Cryst. ($Me_2CO/MeOH$). Mp 272-273°. $[\alpha]_D +98$ (c, 0.5 in MeOH).

Abdullaeva, D.A. et al., *Khim. Prir. Soedin.*, 1976, 121; *CA*, **85**, 106633d

J-44

Jolantimine

Iolantimine
[64986-29-4]



$C_{38}H_{44}N_2O_5$ 608.776


Probable struct. Dimeric phenethylisoquinoline alkaloid. Alkaloid from *Merendera jolantae* (Liliaceae).

Usmanov, A.M. et al., *Khim. Prir. Soedin.*, 1977, **13**, 422; *Chem. Nat. Compd.* (*Engl. Transl.*), 1977, **13**, 360 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

J-45

Joramine

[161897-98-9]



$C_{23}H_{38}N_6O_5$ 478.59

Alkaloid from the spider *Nephila clavata*. λ_{max} 276 (ε 854) (MeOH).

J-46

2'-Hydroxy: *Spidamine*

[161897-97-8]

$C_{23}H_{38}N_6O_6$ 494.59

Alkaloid from the spider *Nephila clavata*. $[\alpha]_D^{25} -4.96$ (c, 0.6 in H_2O). λ_{max} 279 (ε 1690) (MeOH).


Chiba, T. et al., *Chem. Pharm. Bull.*, 1994, **42**, 1864-1869; 1995, **43**, 2177-2181; 1996, **44**, 972-979 (*isol*, *pmr*, *cmr*, *ms*, *synth*)
Nihei, K. et al., *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)

Jordanine

J-47

4-[(3-Methylbutyl)amino]benzeneacetamide, 9CI

[123690-75-5]



$C_{13}H_{20}N_2O$ 220.314


Alkaloid from *Bongardia chrysogonum*. Cryst. (MeOH). Mp 167-170°. λ_{max} 253 (log ε 3.51); 368 (log ε 1.92); 402 (log ε 1.97); 424 (log ε 1.92) (MeOH).

Alfatafa, A.A. et al., *J. Nat. Prod.*, 1989, **52**, 818 (*isol*, *synth*, *uv*, *ir*, *pmr*, *ms*)

Jorumycin

J-48

[304852-37-7]



$C_{27}H_{30}N_2O_9$ 526.542

Related to Renieramycin A, R-41. *Isol.* from the marine nudibranch *Jorunna funebris*. Antitumour agent. Unstable pale yellow powder. $[\alpha]_D -57$ (c, 0.05 in $CHCl_3$). λ_{max} 266 (ε 15000) (MeOH).

Fontana, A. et al., *Tetrahedron*, 2000, **56**, 7305-7308 (*isol*)


Saito, N. et al., *Tetrahedron*, 2004, **60**, 3873-3881 (*synth*, *pmr*, *cmr*)

Lane, J.W. et al., *J.A.C.S.*, 2005, **127**, 12684-12690 (*synth*)

Joubertiamine

J-49

4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)-2-cyclohexen-1-one, 9CI



C₁₆H₂₁NO₂ 259.347**(R)-form***Me ether: O-Methyljoubertinamine*

[77844-21-4]

C₁₇H₂₃NO₂ 273.374Alkaloid from *Sceletium subvelutium* (Aizoaceae). Syrup. Bp_{0.01} 110-120° (bath). [α]_D²⁵ -51 (c, 1.45 in MeOH). λ_{max} 227 (log ε 4.19); 276 (log ε 3.25); 282 (log ε 3.21) (MeOH).**3'-Methoxy, Me ether: 4-(3,4-Dimethoxyphenyl)-4-[2-(dimethylamino)ethyl]-2-cyclohexen-1-one. 3'-Methoxy-4'-O-methyljoubertinamine**

[59096-18-3]

C₁₈H₂₅NO₃ 303.4Minor alkaloid from *Sceletium namaquense*. Oil. λ_{max} 230 (ε 14100); 280 (ε 2000); 346 (ε 600) (95% EtOH).**(S)-form** [28379-30-8]Alkaloid from the aerial parts of *Sceletium joubertii* (Aizoaceae). Liq. Opt. rotn. not reported. λ_{max} 226 (ε 13610); 278 (ε 1960) (96% EtOH).**(±)-form** [34603-53-7]Synthetic. Yellow needles (Et₂O), cryst. (MeCN). Mp 126-130° Mp 162.5-164°.*Me ether*: [34603-52-6]

Synthetic. Oil.

2,3-Dihydro: Dihydrojoubertinamine. 4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)cyclohexanone

[28513-25-9]

C₁₆H₂₃NO₂ 261.363Alkaloid from the aerial parts of *Sceletium joubertii* (Aizoaceae). Achiral. λ_{max} 226 (ε 7905); 278 (ε 1857) (96% EtOH).**2,3-Dihydro, Me ether: 4-[2-(Dimethylamino)ethyl]-4-(4-methoxyphenyl)cyclohexanone. O-Methylidihydrojoubertinamine**

[77784-04-4]

C₁₇H₂₅NO₂ 275.39Alkaloid from *Sceletium subvelutium* (Aizoaceae). Syrup. Achiral.**2,3-Dihydro, (N,N-dimethylamino)ethyl ether: Amisine. 4-[4-[2-(Dimethylamino)ethoxy]phenyl]-4-[2-(dimethylamino)ethyl]cyclohexanone, 9CI**


[76129-90-3]

C₂₀H₃₂N₂O₂ 332.485Alkaloid from the above-ground parts of *Hymenocallis arenicola* (Amaryllidaceae). Needles (Me₂CO). Mp 74-80°.**5,6-Didehydro: 4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)-2,5-cyclohexadien-1-one. Dehydrojoubertinamine. Didehydrojoubertinamine**

[28564-22-9]

C₁₆H₁₉NO₂ 257.332Trace alkaloid from the aerial parts of *Sceletium joubertii* (Aizoaceae). Pale yellow oil. Achiral.**5,6-Didehydro, Me ether: O-Methyldehydrojoubertinamine**

[121983-83-3]

C₁₇H₂₁NO₂ 271.358Alkaloid from whole plants of *Sceletium subvelutium* (Aizoaceae). Achiral. λ_{max} 226 (log ε 3.24); 276 (log ε 2.51); 283 (log ε 2.45) (MeOH).Arndt, R.R. *et al.*, *Tet. Lett.*, 1970, 3237-3240 (*Joubertinamine, Dihydrojoubertinamine, Dehydrojoubertinamine, isol, uv, ir, pmr, ms, struct*)Stevens, R.V. *et al.*, *J.O.C.*, 1972, **37**, 2138-2140 (*synth, ir, pmr*)Martin, N.H. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 1-19 (*ms*)Capps, T.M. *et al.*, *J.C.S. Perkin 2*, 1977, 1098-1104 (*3'-Methoxy-4'-O-methyljoubertinamine*)Strauss, H.F. *et al.*, *Tetrahedron*, 1978, **34**, 127-130 (*O-Methyljoubertinamine, synth*)Döpke, W. *et al.*, *Z. Chem.*, 1980, **20**, 298-299 (*Amisine*)Nieuwenhuis, J.J. *et al.*, *J.C.S. Perkin 1*, 1981, 284-286 (*O-Methyljoubertinamine, O-Methylidihydrojoubertinamine, isol, uv, cd, ir, pmr, struct*)Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3373-3380 (*synth, ir, pmr, ms*)Herbert, R.B. *et al.*, *Tetrahedron*, 1990, **46**, 7105-7118 (*O-Methyldehydrojoubertinamine*)Bauermeister, S. *et al.*, *J.C.S. Perkin 1*, 1991, 561-565 (*synth, O-Methyljoubertinamine*)Parkinson, C.J. *et al.*, *J.C.S. Perkin 1*, 1991, 1053-1057 (*synth, O-Methyljoubertinamine*)Stephenson, G.R. *et al.*, *Tetrahedron*, 1993, **49**, 5649-5662 (*synth, O-Methyljoubertinamine*)Matsumura, Y. *et al.*, *Tetrahedron*, 1993, **49**, 8503-8512 (*synth*)**Joubertinamine****J-50****4-(3,4-Dimethoxyphenyl)-4-[2-(methylamino)ethyl]-2-cyclohexen-1-ol, 9CI**C₁₇H₂₅NO₃ 291.389**(-)-form** [71294-61-6]Alkaloid from *Sceletium joubertii* (Aizoaceae). Noncryst. [α]_D²⁰ -18 (CHCl₃).**O,N-Di-Ac:**Noncryst. [α]_D²⁰ +21 (CHCl₃).**1-Epimer, N-Me: 4-(3,4-Dimethoxyphenyl)-4-[2-(dimethylamino)ethyl]-2-cyclohexen-1-ol. 3'-Methoxy-4'-O-methyljoubertinaminol**

[59096-21-8]

C₁₈H₂₇NO₃ 305.416Alkaloid from *Sceletium namaquense* (Aizoaceae). Solid. Mp 100.5°. [α]_D²⁵ -12.4 (c, 2.6 in MeOH).**(±)-form**


Synthetic. Oil.

1-Epimer: Epijoubertinamine

Synthetic. Oil.

Psotta, K. *et al.*, *J.C.S. Perkin 1*, 1979, 1063 (*isol, uv, ir, pmr, ms, struct*)Jeffs, P.W. *et al.*, *J.O.C.*, 1982, **47**, 3611 (*3'-Methoxy-4'-O-methyljoubertinaminol*)Sánchez, I.H. *et al.*, *Tet. Lett.*, 1983, **24**, 551 (*synth*)Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3373 (*synth, ir, pmr*)Tam, N.T. *et al.*, *Org. Lett.*, 2007, **9**, 3391-3392 (*synth*)**Jubanine A****J-51**

[60375-07-7]

R¹ = -CH(CH₃)CH₂CH₃
R² = -CH₂PhC₄₀H₄₉N₅O₆ 695.857Alkaloid from the root or stem bark of *Zizyphus jujuba*, *Zizyphus nummularia* and *Zizyphus spina-christi* (Rhamnaceae). Amorph. [α]_D²⁰ -326 (c, 0.12 in MeOH). Stereochemistry unspecified.**N-De-Me: Nummularine H. N-Desmethyljubanine A**

[64309-23-5]

C₃₉H₄₇N₅O₆ 681.83Alkaloid from the bark of *Zizyphus nummularia* and stems of *Paliurus ramosissimus* (Rhamnaceae). Needles (MeOH). Mp 194-196°. [α]_D²⁰ -343 (c, 0.27 in MeOH).**Dihydro:** Mp 247-249°. [α]_D²⁰ -148 (c, 0.06 in MeOH).Tschesche, R. *et al.*, *Phytochemistry*, 1976, **15**, 541-542 (*Jubanine A*)Tschesche, R. *et al.*, *Chem. Ber.*, 1977, **110**, 2649 (*Nummularine H*)**Jubanine B****J-52**

[60375-08-8]

As Jubanine A, J-51 with

R¹ = R² = CH₂PhC₄₃H₄₇N₅O₆ 729.874Alkaloid from the stem bark of *Zizyphus jujuba* and from *Zizyphus nummularia* (Rhamnaceae). Amorph. Mp 97-100°. [α]_D²⁰ -218 (c, 0.28 in MeOH). Stereochemistry unspecified.**N-De-Me: Nummularine O. N-Desmethyljubanine B**

[107602-86-8]

[103197-36-0 (*N-Desmethyljubanine B*)]C₄₂H₄₅N₅O₆ 715.847Alkaloid from the root bark of *Zizyphus nummularia* (Rhamnaceae). Powder (MeOH). Mp 159-161°. [α]_D²⁰ -239 (c, 0.2 in MeOH). Identity of the two isolates not certain. *N-Desmethyljubanine B* supported only by ms evidence.**N²-Deacyl, N²-di-Me: Xylopyrine B**

[1010795-96-6]


C₃₄H₃₈N₄O₅ 582.698Alkaloid from the root bark of *Zizyphus xylopyra*. Cryst. (MeOH). Mp 206-208°. [α]_D²⁵ -190 (c, 0.15 in CHCl₃). λ_{max} 270 (log ε 2.8); 320 (log ε 2.4) (MeOH).Tschesche, R. *et al.*, *Phytochemistry*, 1976, **15**, 541-542 (*Jubanine B*)Miana, G.A. *et al.*, *Fitoterapia*, 1985, **56**, 363-364; *CA*, **105**, 75886p (*N-Desmethyljubanine B*)

- Pandey, V.B. *et al.*, *Phytochemistry*, 1986, **25**, 2690-2691 (*Nummularine O*)
 Singh, A.K. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1114-1120 (*Xylopyrine B*)

Jubanine C

J-53

[159226-00-3]




C₃₉H₄₇N₅O₅ 665.831
 Alkaloid from the stem bark of *Zizyphus jujuba* (Chinese date). Cryst. (MeOH).
 Mp 233-235° dec. λ_{max} 214 (MeOH).

Tripathi, M. *et al.*, *Fitoterapia*, 2001, **72**, 507-510

Julandine

J-54

8-(3,4-Dimethoxyphenyl)-1,3,4,6,9,9a-hexahydro-7-(4-methoxyphenyl)-2H-quinolizine, 9CI



(R)-form

C₂₄H₂₉NO₃ 379.498

(R)-form

Synthetic. Needles (EtOAc/hexane). Mp 138.5-140°. [α]_D²⁵ -71.6 (c, 0.33 in CHCl₃).

(S)-form [20772-34-3]

Constit. of *Boehmeria cylindrica* and *Boehmeria platyphylla* (Urticaceae). Shows antifungal activity. Needles (Me₂CO). Poorly sol. hexane. Mp 134.5-135.5°. [α]_D +4.6 (c, 0.5 in CHCl₃). Although natural Jalandine is weakly optically active, it is largely racemic.

4',4''-Di-O-de-Me: 4',4''-Dide-O-methyl-julandine

C₂₂H₂₅NO₃ 351.444

Alkaloid from *Boehmeria siamensis*. Cryst. Mp 123-125°.

(±)-form [23365-39-1]

Synthetic. Long needles (EtOAc/hexane). Mp 139-140° (134-136°).

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1968, **21**, 2579 (*isol, uv, pmr, ms, struct*)

Paton, J.M. *et al.*, *J.C.S. (C)*, 1969, 1309 (*synth, ir, pmr*)

Herbert, R.B. *et al.*, *Chem. Comm.*, 1978, 794 (*synth*)

Trigo, G.G. *et al.*, *J. Het. Chem.*, 1979, **16**, 1625 (*synth, ir, pmr, ms*)

Cragg, J.E. *et al.*, *J.C.S. Perkin 1*, 1982, 2487 (*synth, pmr, ms*)

Al-Shamma, A. *et al.*, *Phytochemistry*, 1982, **21**, 485 (*biochem*)

Iida, H. *et al.*, *J.O.C.*, 1984, **49**, 2412 (*synth, pmr, ms*)

Grieco, P.A. *et al.*, *J.O.C.*, 1988, **53**, 3325 (*synth, ir, pmr*)

Suzuki, H. *et al.*, *J.O.C.*, 1995, **60**, 6114 (*synth, abs config*)


Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1996, **118**, 12082 (*synth*)

Luo, Y.G. *et al.*, *Chin. Chem. Lett.*, 2001, **12**, 337-338 (4',4''-di-O-de-Me)

Julifloridine

J-55

5-Hydroxy-6-methyl-2-piperidinedecanol, 9CI



C₁₈H₃₇NO₂ 299.496

(2R,5R,6R)-form [66731-40-6]

Minor alkaloid from *Prosopis juliflora* leaves (Fabaceae). Mp 82-83°. [α]_D +18 (c, 0.84 in MeOH) (synthetic).

(2R,5S,6S)-form

Alkaloid from *Prosopis alba*. Resin. [α]_D²⁰ -2.8 (c, 1.24 in CHCl₃).

(±)-form [87304-12-9]

Cryst. (EtOH). Mp 87-90°.

N-Me: 5-Hydroxy-1,6-dimethyl-2-piperidinedodecanol. **N-Methyljulifloridine** [132972-82-8]

C₁₉H₃₉NO₂ 313.523

Alkaloid from *Prosopis juliflora* (Fabaceae). Struct. requires revision (1998).

Ahmad, V.U. *et al.*, *Z. Naturforsch., B*, 1978, **33**, 347 (*isol, ms, struct*)

Paterne, M. *et al.*, *C. R. Hebd. Seances Acad. Sci., Ser. II*, 1983, **296**, 433 (*synth*)

Ahmad, V.U. *et al.*, *Sci. Pharm.*, 1990, **58**, 409 (*N-Methyljulifloridine*)

Kiguchi, T. *et al.*, *Tetrahedron*, 1998, **54**, 15589-15606 (*synth, struct*)

Astudillo, S.L. *et al.*, *Planta Med.*, 1999, **65**, 161-162 (*isol, pmr, cmr, ms*)


Lemire, A. *et al.*, *Org. Lett.*, 2005, **7**, 2747-2750 (*synth*)

Zhai, H. *et al.*, *J.O.C.*, 2007, **72**, 3853-3858 (*synth*)

Juliflorine

J-56

Juliprosopine [76202-00-1]



Relative Configuration

C₄₀H₇₅N₃O₂ 630.051

The two piperidine systems need not have the same abs. config. Alkaloid from *Prosopis juliflora*, *Prosopis fraxta* and *Prosopis glandulosa* (Fabaceae). Shows antibacterial and antifungal props. Amoebicide. Noncryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D +0.7 (c, 1.05 in EtOH). [α]_D +10 (c, 2.15 in CHCl₃). Several O,N-acetyl and N-Me derivs. were all noncryst.

▶TN7637000

3''-Oxo: 3''-Oxojuliflorine. 3''-Oxojuliprosopine

C₄₀H₇₃N₃O₃ 644.035

Alkaloid from *Prosopis juliflora*. Gum. [α]_D²³ +4 (c, 1 in MeOH).

Stereoisomer: **Julifloricine**

[66771-80-0]

C₄₀H₇₅N₃O₂ 630.051

Alkaloid from leaves of *Prosopis juliflora* (Fabaceae). Shows antibacterial props. Gum. [α]_D +4.3 (c, 1.4 in CHCl₃). Two structs. stereoisomeric with Juliflorine have been proposed but current spectroscopic data cannot distinguish between them.

Stereoisomer (2): **Juliflorinine**

[122441-92-3]

C₄₀H₇₅N₃O₂ 630.051

Alkaloid from leaves of *Prosopis juliflora* (Fabaceae). Sol. MeOH, C₆H₆. [α]_D +3.9 (c, 0.03 in CHCl₃). Has (2α,5β,6β) configs. of the piperidine systems. λ_{max} 208 ; 285 (MeOH) (Berdy).

Ahmad, V.U. *et al.*, *Z. Naturforsch., B*, 1978, **33**, 347 (*isol*)

Ahmad, V.U. *et al.*, *J. Chem. Soc. Pak.*, 1979, **1**, 137; 1985, **7**, 347 (*struct, ms, pmr, cmr, deriv*)

Ott-Longoni, R. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 2119 (*isol, ir, pmr, cmr, ms, struct*)

Ahmad, A. *et al.*, *Planta Med.*, 1986, 285 (*props*)

Aqeel, A. *et al.*, *Arzneim.-Forsch.*, 1989, **39**, 652 (*props, Julifloricine*)

Ahmad, V.U. *et al.*, *J. Nat. Prod.*, 1989, **52**, 497 (*Juliflorinine*)

Nakano, H. *et al.*, *Phytochemistry*, 2004, **65**, 587-591 (3''-Oxojuliflorine)


Snyder, B.B. *et al.*, *Org. Lett.*, 2005, **7**, 2715-2718 (*synth*)

Juliprosine

J-57

2,3-Dihydro-6,8-bis[10-(5-hydroxy-6-methyl-2-piperidinyl)decyl]-1H-indolizinium, 9CI

[80233-19-8]



Absolute Configuration

C₄₀H₇₂N₃O₂[⊕] 627.028

Alkaloid from *Prosopis juliflora* (Fabaceae). Oil (as chloride). Sol. MeOH, CHCl₃. [α]_D²⁵ +11 (c, 0.50 in CHCl₃) (chloride). λ_{max} 279 (ε 6200) (EtOH) (Berdy).

3'-Ketone: 3'-Oxojuloprosine

C₄₀H₇₀N₃O₂[⊕] 625.012

Alkaloid from *Prosopis juliflora*. Plant growth inhibitor. Isol. as a mixt. with 3-Oxojuloprosine.

3''-Ketone: 3-Oxojuliprosine

C₄₀H₇₀N₃O₂[⊕] 625.012

Alkaloid from *Prosopis juliflora*. Plant growth inhibitor. Gum. [α]_D²⁸ +3 (c, 1 in MeOH). Isol. as a mixt. with 3'-Oxojuliprosine to which data refers.

1,2-Didehydro: 6,8-Bis[10-(5-hydroxy-6-methyl-2-piperidinyl)decyl]-1H-indolizinium. Juliprosinene

[123061-99-4]

C₄₀H₇₀N₃O₂[⊕] 625.012

Alkaloid from the leaves of *Prosopis juliflora* (Fabaceae). Gum (as chloride). Sol. MeOH, C₆H₆. [α]_D²⁸ +9.5 (c, 0.04 in CHCl₃) (chloride). λ_{max} 208; 285 (MeOH) (Berdy).

5',5'',6',6''-Tetraepimer: Isojuliprosine

[123805-40-3]

[123750-53-8]

C₄₀H₇₂N₃O₂[⊕] 627.028

Alkaloid from *Prosopis juliflora* (Fabaceae). Yellow oil (as chloride). Sol. MeOH, CHCl₃. λ_{max} 280 (EtOH) (Berdy).

Dätwyler, P. et al., *Helv. Chim. Acta*, 1981, **64**, 1959 (isol, spectra)

Ahmad, A. et al., *Fitoterapia*, 1989, **60**, 86-89 (Isojuliprosine)

Ahmed, V.U. et al., *J. Nat. Prod.*, 1989, **52**, 497-501 (Juliprosinene)


Nakano, H. et al., *Phytochemistry*, 2004, **65**, 587-591 (Oxojuliprosines)

Snyder, B.B. et al., *Org. Lett.*, 2005, **7**, 2715-2718 (synth)

Julocrotine

J-58

N-[2,6-Dioxo-1-(2-phenylethyl)-3-piperidinyl]-2-methylbutanamide, 9CI [492-87-5]



Absolute Configuration

C₁₈H₂₄N₂O₃ 316.399

Originally considered to be C₁₉H₂₆N₂O₃. Alkaloid from *Julocrotion montevidensis*, *Julocrotion subpannosus*, *Julocrotion camporum* and *Croton membranaceus*. Mp 108-109°. Bp_{0.25} 160-165° subl. [α]_D²⁵ -9 (c, 1.24 in CHCl₃).

2'R-Hydroxy: Julocrotol

C₁₈H₂₄N₂O₄ 332.399

Alkaloid from the aerial parts of *Croton cuneatus*. Orange oil. [α]_D²⁵ -56 (c, 1.2 in CHCl₃).

2'S-Hydroxy: Isojulocrotol

C₁₈H₂₄N₂O₄ 332.399

Alkaloid from *Croton cuneatus*. Green oil. [α]_D²⁵ +63 (c, 1.3 in CHCl₃).

2'-Oxo: Julocrotone

C₁₈H₂₂N₂O₄ 330.383

Alkaloid from *Croton cuneatus*.

Amorph. yellow solid. Mp 89-92°. [α]_D²⁵ -10 (c, 1 in CHCl₃).

Nakano, T. et al., *J.O.C.*, 1961, **26**, 1184-1191 (Julocrotone)

Aboagye, F.A. et al., *Fitoterapia*, 2000, **71**, 461-462 (Julocrotone)


Suárez, A.I. et al., *Nat. Prod. Res.*, 2004, **18**, 421-426 (Julocrotol, Isojulocrotol, Julocrotone)

Moreira, R.Y.O. et al., *Int. J. Quantum Chem.*, 2008, **108**, 513-520 (cryst struct)

Junosidine

J-59

2,11-Dihydro-5-hydroxy-10-methoxy-2,2,11-trimethyl-6H-pyrano[3,2-b]acridin-6-one, 9CI [110883-39-1]



C₂₀H₁₉NO₄ 337.374

Alkaloid from the root bark of *Citrus junos* (yuzu) (Rutaceae). Orange needles. Mp 188-189°.

O-De-Me: Yukocitrine

[145940-32-5]

C₁₉H₁₇NO₄ 323.348

Alkaloid from roots of *Citrus yuko* (Rutaceae). Yellow oil.

9-Hydroxy: Honyumine

[100595-86-6]

C₂₀H₁₉NO₅ 353.374

Alkaloid from the root bark of *Citrus grandis* (pummelo) (Rutaceae). Yellow granules (Me₂CO). Mp 175-176°.

Wu, T.S. et al., *Heterocycles*, 1986, **24**, 41-43 (Honyumine)


Ju-ichi, M. et al., *Heterocycles*, 1987, **26**, 2077 (Junosidine)

Takemura, Y. et al., *Heterocycles*, 1992, **34**, 2123-2130 (Yukocitrine)

Jurubine

J-60

[14256-61-2]



Absolute conuguration

C₃₃H₅₇NO₈ 595.815

Alkaloid from roots of *Solanum paniculatum* and *Solanum torvum* (pea eggplant) (Solanaceae) and from *Solanum cha-coense*. Mp 212-214°. [α]_D¹⁸ -30.9 (c, 1.04 in Py). Paniculine from *Solanum paniculatum* was a mixt. of Jurubine and its C-25 epimer.

N-Salicylidene: Mp 176-177°. [α]_D²⁵ -34.9 (Py).


Penta-Ac: Mp 165-168°. [α]_D²⁵ -41.1 (Py).

Ripperger, H. et al., *Chem. Ber.*, 1967, **100**, 1725 (isol, ir, ms, pmr, struct)

Schreiber, K. et al., *CA*, 1968, **69**, 27721 (isol)

Jussiaeine A

J-61



C₁₃H₂₀N₂O₂ 236.313

Alkaloid from *Ulex jussiaei*. Oil. [α]_D²⁵ +3.3 (c, 0.26 in CHCl₃). λ_{max} 219 (log ε 3.34); 267 (log ε 3.56); 274 (sh) (MeOH).


Maximo, P. et al., *J. Nat. Prod.*, 2000, **63**, 201-204 (isol, pmr, cmr)

Honda, T. et al., *J.O.C.*, 2005, **70**, 499-504 (synth)

Justiciamide

J-62

4-Hydroxy-N^d-(2-hydroxy-4,5-dimethoxyphenyl) glutamine



C₁₃H₁₈N₂O₇ 314.294


Alkaloid from *Justicia ghiesbreghtiana*. Mp 185°. [α]_D²⁵ -2.7 (c, 0.07 in H₂O). λ_{max} 244 (ε 6220); 296 (ε 5775) (H₂O).

Lorenz, P. et al., *Phytochemistry*, 1999, **52**, 63-66 (isol, uv, ir, pmr, cmr)

Juziphine

J-63

Yuziphine. Yuzifine



(R)-form

C₁₈H₂₁NO₃ 299.369

(R)-form [64091-05-0]

Alkaloid from the aerial parts of *Corydalis gortschakovii* and the leaves of *Zizyphus jujuba* (Chinese date) (Papaveraceae, Rhamnaceae).

N-Oxide: Juziphine N-oxide

[65954-43-0]

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Corydalis gortschakovii* (Papaveraceae). Cryst. (MeOH). Mp 197-198°. [α]_D²⁵ -29 (MeOH).

N-De-Me: **Norjuziphine**

[74119-87-2]

C₁₇H₁₉NO₃ 285.342Alkaloid from *Corydalis bulleyana* and *Fumaria vaillantii* (Papaveraceae). Plates (MeOH). Mp 199-201°. [α]_D²¹ +17.3 (MeOH).

N-Me: 1,2,3,4-Tetrahydro-8-hydroxy-1-[(4-hydroxyphenyl)methyl]-7-methoxy-2,2-dimethylisoquinolinium(1+).

Oblongine†

[152230-57-4]

[152230-59-6]

C₁₉H₂₄NO₃[⊕] 314.403Quaternary alkaloid from stems of *Litsea cubeba* (mountain pepper) (Lauraceae). Amorph. solid (as perchlorate). Mp 109.5°(perchlorate). [α]_D²⁴ -11 (c, 1.0 in MeOH).**(S)-form**

N-Me: [60008-01-7]

[60134-32-9]

Quaternary alkaloid from *Berberis oblonga*. Mp 160° (as iodide). [α]_D 9.**(±)-form**

N-Me: [97185-37-0]


Quaternary alkaloid from the roots of *Tiliacora dinklagei* (Menispermaceae).Cryst. (Me₂CO/petrol) (as iodide). Mp 161° (153-155°) (iodide).

[152321-76-1]

Karimov, A. et al., *Khim. Prir. Soedin.*, 1976,12, 117-118 (*Oblongine, isol*)Ziyaev, R. et al., *Khim. Prir. Soedin.*, 1977, 13, 239; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, 13, 204Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1977, 13, 834; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, 13, 702Alimova, M. et al., *Khim. Prir. Soedin.*, 1979, 15, 874; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, 15, 783 (*Norjuziphine*)Dwuma-Badu, D. et al., *J. Nat. Prod.*, 1983, 46, 342 (*Oblongine, isol, uv, ir, pmr, ms, struct, synth*)Hong, H. et al., *Planta Med.*, 1986, 193 (*Norjuziphine*)Lee, S.S. et al., *J. Nat. Prod.*, 1993, 56, 1971 (*Oblongine*)Chen, J.-J. et al., *Planta Med.*, 2001, 67, 423-427 (*Norjuziphine, isol, pmr, cmr*)

Kabiramide I

[927836-97-3]

C₄₇H₇₁N₅O₁₅ 946.102Isol. from the sponge *Pachastrissa nux*.Petchprayoon, C. *et al.*, *Heterocycles*, 2006, **69**, 447-456 (isol, pmr, cmr)


K-1

Cytotoxic. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D²³ +6 (c, 0.1 in CHCl₃). λ_{max} 245 (ε 26000) (MeOH) (Derep). λ_{max} 245 ; 345 (ε 25000) (MeOH) (Berdy).

32-Demethoxy, 31,32-didehydro(E-), 3-O-carbamoyl: **Kabiramide G**

C₄₇H₆₇N₅O₁₃ 910.072Isol. from *Pachastrissa nux*. [α]_D +38 (c, 0.4 in CHCl₃).Matsunaga, S. *et al.*, *J.A.C.S.*, 1986, **108**, 847-849 (*Kabiramide C*)Matsunaga, S. *et al.*, *J.O.C.*, 1989, **54**, 1360-1363 (*Kabiramides A,B,D,E*)Petchprayoon, C. *et al.*, *Heterocycles*, 2006, **69**, 447-456 (*Pachastrissa nux* constits)**Kabiramide D**

[101550-95-2]

C₄₇H₇₀N₄O₁₃ 899.089

Isol. from the eggs of *Hexabranchnus* sp. and from *Pachastrissa nux*. Cytotoxic agent. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D²³ -5 (c, 0.1 in CHCl₃). λ_{max} 245 (ε 26000) (MeOH) (Derep).

3-Ac: **Kabiramide E**

[101550-96-3]

C₄₉H₇₂N₄O₁₄ 941.126

Isol. from the eggs of *Hexabranchnus* sp. Cytotoxic agent. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D²³ -20 (c, 0.1 in CHCl₃). λ_{max} 245 (ε 26000) (MeOH) (Derep).

3-O-Carbamoyl: **Kabiramide C**

[100045-78-1]

C₄₈H₇₁N₅O₁₄ 942.114

Isol. from the eggs of *Hexabranchnus* sp. and the sponge *Pachastrissa nux*. Shows antifungal props. Solid. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D²³ +20 (c, 0.1 in CHCl₃). Macrolide antibiotic. Related to Halichondramide, H-15. λ_{max} 245 (ε 26000) (MeOH) (Derep).

22-O-De-Me: **Kabiramide F**C₄₆H₆₈N₄O₁₃ 885.062Isol. from *Pachastrissa nux*. [α]_D -26 (c, 0.46 in CHCl₃).22-O-De-Me, 3-O-carbamoyl: **Kabiramide B**

[101550-94-1]

C₄₇H₆₉N₅O₁₄ 928.087


Isol. from the eggs of *Hexabranchnus* sp. and from *Pachastrissa nux*. Cytotoxic. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D²³ +8 (c, 0.1 in CHCl₃). λ_{max} 245 (ε 26000) (MeOH) (Derep).

47-Hydroxy, 3-O-carbamoyl: **Kabiramide A**

[101550-93-0]

C₄₈H₇₁N₅O₁₅ 958.113Isol. from the eggs of *Hexabranchnus* sp.**Kabiramide H**

[927836-94-0]


C₄₈H₇₅N₅O₁₆ 978.144Isol. from the sponge *Pachastrissa nux*. [α]_D -35 (c, 0.3 in CHCl₃).Petchprayoon, C. *et al.*, *Heterocycles*, 2006, **69**, 447-456 (isol, pmr, cmr)

K-3

Kainic acid, INN, JAN

K-4

2-Carboxy-4-(1-methylethenyl)-3-pyrrolidinediacetic acid, 9CI. 3-Carboxymethyl-4-isopropenylproline. α-Kaininic acid. Digenic acid. Digenin. α-Kainic acid [487-79-6]

C₁₀H₁₅NO₄ 213.233

Constit. of red algae *Alsidium helminthochorton*, *Centroceras clavulatum* and *Digenea simplex*. Glutamate receptor agonist. Neurotoxin, formerly used as an anthelmintic agent. Cryst. + 1H₂O (EtOH aq.). Sol. H₂O. Mp 253-254° dec. [α]_D²⁴ -14.8 (c, 1 in H₂O). Log P -1.21 (calc).

► V. neurotoxic. UX9665250

Di-Me ester: [4071-37-8]Bp₄ 145°. [α]_D²⁰ +23.*N-Ac*: [59845-92-0]Mp 161-162°. [α]_D -53 (H₂O).

9,10-Dihydro, 9-hydroxy: 2-Carboxy-4-(1-hydroxy-1-methylethyl)-3-pyrrolidinediacetic acid, 9CI. 1'-Hydroxykainic acid

[159834-87-4]

C₁₀H₁₇NO₅ 231.248Constit. of the red alga *Palmaria palmata*.

4-Epimer: α-Allokainic acid. α-Allokainic acid

[4071-39-0]

C₁₀H₁₅NO₄ 213.233

Constit. of *Digenea simplex*. Mp 238-242° dec. [α]_D²⁰ +7.7 (c, 1.3 in H₂O).

[92180-28-4, 58002-62-3, 92180-29-5]

Watase, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1957, **30**, 889 (*cryst struct*)Nitta, I. *et al.*, *Nature (London)*, 1958, **181**, 761 (*struct*)Impellizzeri, G. *et al.*, *Phytochemistry*, 1975, **14**, 1549-1557 (*algae, isol*)

Kainic Acid Tool Neurobiol., (eds. McGeer, E.G. *et al.*), Raven Press, 1978, (*book*)

Oppolzer, W. *et al.*, *Tet. Lett.*, 1978, 3397-3400 (α-Allokainic acid, synth)Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 2282 (*synth, bibl*)Nadler, J.V. *et al.*, *Life Sci.*, 1979, **24**, 289 (*rev, props*)Kraus, G.A. *et al.*, *Tet. Lett.*, 1983, **24**, 3427-3430 (α-Allokainic acid, synth)DeShong, P. *et al.*, *Tet. Lett.*, 1986, **27**, 3979-3982 (α-Allokainic acid, synth)


Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 785

Yoo, S.-E. *et al.*, *J.O.C.*, 1994, **59**, 6968-6972 (*synth*)Agami, C. *et al.*, *J.O.C.*, 1994, **59**, 7937-7940 (*synth*)Ramsey, U.P. *et al.*, *Nat. Toxins*, 1994, **2**, 286-292 (1'-Hydroxykainic acid)Parsons, A.F. *et al.*, *Tetrahedron*, 1996, **52**, 4149-4174 (*rev*)Todeschi, N. *et al.*, *Bioform. Med. Chem.*, 1997, **5**, 1943-1957 (*pmr, conformn*)Bachi, M.D. *et al.*, *Pure Appl. Chem.*, 1998, **70**, 259-262 (*rev, synth*)Rubio, A. *et al.*, *Tet. Lett.*, 1998, **39**, 2171-2174 (*synth*)Chevliakov, M.V. *et al.*, *J.A.C.S.*, 1999, **121**, 11139-11143 (*synth*)Campbell, A.D. *et al.*, *J.C.S. Perkin 1*, 2000, 3194-3204 (*synth*)Miyata, O. *et al.*, *Tetrahedron*, 2000, **56**, 6199-6207 (*synth, bibl*)Anderson, J.C. *et al.*, *J.O.C.*, 2003, **68**, 6160-6163 (*synth*)Trost, B.M. *et al.*, *Org. Lett.*, 2003, **5**, 1467-1470 (*synth*)Cook, G.R. *et al.*, *Org. Lett.*, 2004, **6**, 2481-2484 (*synth*)Martinez, M.M. *et al.*, *Eur. J. Org. Chem.*, 2005, 1427-1443 (*synth*)Scott, M.E. *et al.*, *Org. Lett.*, 2005, **7**, 3045-3047 (*synth*)Morita, Y. *et al.*, *Org. Lett.*, 2005, **7**, 4337-4340 (*synth*)Pandey, S.K. *et al.*, *Org. Lett.*, 2006, **8**, 5665-5668 (*synth*)Sakaguchi, H. *et al.*, *Org. Lett.*, 2007, **9**, 1635-1638 (*synth*)Chalker, J.M. *et al.*, *Org. Lett.*, 2007, **9**, 3825-3828 (*synth*)Tomooka, K. *et al.*, *Tet. Lett.*, 2008, **49**, 6327-6329 (*synth*)**Kaitocephalin**

K-5

Antibiotic PF 1191. PF 1191

[198710-92-8]



Absolute Configuration

C₁₈H₂₁Cl₂N₃O₉ 494.284

Revision of abs. config. originally assigned in 2001. Prod. by *Eupenicillium shearii* PF-1191. Glutamate receptor antagonist and nerve cell protectant. Powder. Mp 235-238° dec. $[\alpha]_D^{21}$ -31 (c, 0.7 in H₂O). λ_{\max} 217 (ϵ 18200); 297 (ϵ 8200) (H₂O).

Shin-ya, K. *et al.*, *Tet. Lett.*, 1997, **38**, 7079-7082 (*isol, uv, ir, pmr, cmr, N-15 nmr*)

Pat. Coop. Treaty (WIPO), 1998, 41 503; *CA*, **129**, 274808k (*isol*)

Okue, M. *et al.*, *Tet. Lett.*, 2002, **43**, 857-860; 861-864 (*synth, abs config*)


Kawasaki, M. *et al.*, *Org. Lett.*, 2005, **7**, 4165-4167 (*synth*)

Vaswani, R.G. *et al.*, *J.O.C.*, 2008, **73**, 1661-1681 (*synth*)

Kalasinamide

K-6

5,10-Dimethoxy-4-methylbenzo[*g*]quinolin-2(1*H*)-one, 9*CI*. *Atemoine*



C₁₆H₁₅NO₃ 269.299

Alkaloid from *Annona atemoya* (custard apple) and *Polyalthia suberosa*. Orange needles (EtOH/CH₂Cl₂). Mp 233.8-235.5°. λ_{\max} 216 (log ϵ 4.52); 247 (log ϵ 4.81); 264 (sh) (log ϵ 4.78); 274 (log ϵ 4.91); 284 (log ϵ 4.87); 310 (log ϵ 3.95); 323 (log ϵ 4.13); 338 (log ϵ 4.22); 392 (log ϵ 3.66) (EtOH).

6- or 9-Methoxy: *Geovanine*

[11971-82-5]

C₁₇H₁₇NO₄ 299.326

Alkaloid from the trunkwood of *Annona ambotay* (Annonaceae). Yellow powder. Mp 190-192°.

De Oliveira, A.B. *et al.*, *Phytochemistry*, 1987, **26**, 2650-2651 (*Geovanine*)

Tuchinda, P. *et al.*, *Phytochemistry*, 2000, **53**, 1079-1082 (*isol, uv, pmr, cmr, ms*)


Wu, Y.-C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 406-408 (*Atemoine*)

Gandy, M.N. *et al.*, *J. Nat. Prod.*, 2008, **71**, 866-868 (*synth*)

Kalkitoxin

K-7

[247184-89-0]



Absolute Configuration

C₂₁H₃₈N₂O₅ 366.61

Isol. from *Lyngbya majuscula*. Neurotoxin and ichthyotoxin. Cytotoxic.

Berman, F.W. *et al.*, *Toxicol.*, 1999, **37**, 1645-1648 (*isol, activity*)

Wu, M. *et al.*, *J.A.C.S.*, 2000, **122**, 12041-12042 (*synth, struct*)


White, J.D. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 2092-2012 (*synth*)

Yokokawa, F. *et al.*, *Tetrahedron*, 2004, **60**, 6859-6880 (*synth, abs config*)

Kanagawamicin

K-8

AB 116. Antibiotic *AB 116*
[84873-16-5]



C₁₃H₁₇N₅O₆ 339.307

Nucleoside antibiotic. Prod. by *Actinoplanes kanagawaensis*. Shows antitumour and weak antibacterial props. Amorph. powder +1 H₂O. Sol. H₂O, DMSO, Py, MeOH; poorly sol. Me₂CO, hexane. Mp 225-228°. $[\alpha]_D^{25}$ -14 (c, 1 in DMSO). λ_{\max} 231 (ϵ 14100); 272 (ϵ 6310); 297 (ϵ 6920) (0.1*M* HCl) (*Derep*). λ_{\max} 225 (sh) (ϵ 9175); 268 (ϵ 9175); 282 (sh) (H₂O at pH 11) (*Derep*). λ_{\max} 232 (ϵ 19680); 272 (ϵ 6881); 298 (ϵ 7610) (H₂O) (*Derep*).

N,O,O-Tri-Ac:

Amorph. powder. Mp 170-171°.

[86330-89-4]

Naruto, S. *et al.*, *Heterocycles*, 1983, **20**, 27 (*isol, struct*)


Japan. Pat., 1983, 83 32 893; *CA*, **99**, 37106 (*isol*)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (*rev*)

Kapakahine A

K-9

[181862-46-4]



C₅₈H₇₂N₁₀O₉ 1053.268


Cyclic peptide. Isol. from the sponge *Cribrachalina olemda*. Protein phosphatase inhibitor. Amorph. solid. $[\alpha]_D^{20}$ -131 (c, 1 in MeOH). λ_{\max} 206 (ϵ 39000); 232 (ϵ 19000); 274 (ϵ 9500) (MeOH).

Yeung, B.K.S. *et al.*, *J.O.C.*, 1996, **61**, 7168 (*isol, uv, pmr, cmr*)

Kapakahine B

K-10

[165406-94-0]



Absolute Configuration

C₄₉H₅₂N₈O₆ 849

Isol. from the sponge *Cribrachalina olemda*. Cytotoxic agent. Amorph. solid. Sol. MeOH, EtOAc; poorly sol. hexane. $[\alpha]_D^{20}$ -70 (c, 0.3 in MeOH). λ_{\max} 214 (ϵ 21000); 250 (ϵ 8400); 280 (ϵ 5600); 294 (ϵ 4000) (MeOH).

N-De(2-amino-3-phenylpropanoyl): *Kapakahine F*

[530097-82-6]

C₄₀H₄₃N₇O₅ 701.824

Isol. from *Cribrachalina olemda*.

Nakao, Y. *et al.*, *J.A.C.S.*, 1995, **117**, 8271-8272 (*isol, uv, ir, pmr, cmr*)


Yeung, B.K.S. *et al.*, *J.O.C.*, 1996, **61**, 7168-7173 (*abs config*)

Nakao, Y. *et al.*, *Org. Lett.*, 2003, **5**, 1387-1390 (*Kapakahine F*)

Kapakahine C

K-11

[181862-47-5]



C₅₈H₇₂N₁₀O₁₀ 1069.268

Cyclic peptide. Isol. from the sponge *Cribrachalina olemda*. Amorph. solid. $[\alpha]_D^{20}$ -120 (c, 0.5 in MeOH). λ_{\max} 208 (ϵ 44000); 253 (ϵ 12000); 279 (ϵ 41000); 305 (ϵ 1200) (MeOH).

4,5-Diepimer: *Kapakahine D*

[182074-03-9]

C₅₈H₇₂N₁₀O₁₀ 1069.268


Isol. from *Cribrachalina olemda*.

Amorph. solid. $[\alpha]_D^{20}$ -30.7 (c, 0.9 in MeOH). λ_{\max} 206 (ϵ 35000); 247 (ϵ 9000); 279 (ϵ 2800); 304 (ϵ 690) (MeOH).


Yeung, B.K.S. *et al.*, *J.O.C.*, 1996, **61**, 7168 (*isol, uv, ir, pmr, cmr*)

Kapakahine E


[530097-81-5]

C₅₇H₅₇N₉O₈ 996.133Isol. from the sponge *Cribrochalina olemda*. Cytotoxic.Nakao, Y. *et al.*, *Org. Lett.*, 2003, **5**, 1387-1390 (isol, pmr, ms)**Kapakahine G**


[530097-83-7]

C₅₅H₆₆N₁₀O₁₀S 1059.253Isol. from the sponge *Cribrochalina olemda*.Nakao, Y. *et al.*, *Org. Lett.*, 2003, **5**, 1387-1390 (isol, pmr, cmr, ms)**Karacicine**



[91926-06-6]

C₂₇H₄₄N₂O 412.657Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. λ_{max} 210 ; 241 ; 248 ; 257 (no solvent reported).Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B.*, 1984, **39**, 839 (isol, pmr, ir, uv, ms)**Karachine**

[80908-02-7]

**K-12**C₂₆H₂₇NO₅ 433.503Minor alkaloid from *Berberis aristata* (Berberidaceae). Cryst. (EtOAc). Mp 146-148°.Blaskó, G. *et al.*, *J.A.C.S.*, 1982, **104**, 2039 (uv, ir, pmr, ms, struct)Stevens, R.V. *et al.*, *Chem. Comm.*, 1983, 1425 (synth)**Karakoramine**

[85588-84-7]

C₂₅H₂₇NO₅ 421.492Alkaloid from *Berberis lycium* (Berberidaceae). Amorph. [α]_D²⁵ +71 (c, 0.17 in MeOH). Poss. derived by *in vivo* oxidn. of Berbaminine, B-98.Leet, J.E. *et al.*, *Heterocycles*, 1983, **20**, 425 (isol, uv, pmr, ms, cd, struct)**Karnamicin A₁**BU 3557A₁. Antibiotic BU 3557A₁ [122535-48-2]R = -CH(OH)CH₂CH₂CH(OH)CH₃C₁₆H₂₁N₃O₆S 383.424Prod. by *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, CHCl₃, EtOAc, DMSO, C₆H₆, EtOH; poorly sol. hexane, H₂O. Mp 56.5-58.5°. [α]_D²⁵ -11 (c, 0.5 in CHCl₃). λ_{max} 220 (ε 25700); 243 (ε 25700); 318 (ε 7500) (MeOH/HCl). λ_{max} 220 (ε 23400); 251 (ε 21600); 281 (ε 17200); 342 (ε 9000) (MeOH/NaOH). λ_{max} 219 (ε 25000); 242 (ε 22700); 317 (ε 6700) (MeOH).1''-Deoxy: **Karnamicin B₂**. BU 3557B₂. Antibiotic BU 3557B₂ [122535-52-8]C₁₆H₂₁N₃O₅S 367.425Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Rods (CHCl₃). Sol. MeOH, EtOH, CHCl₃, EtOAc, C₆H₆, DMSO; poorly sol. hexane, H₂O. Mp 171-172.5°. λ_{max} 220 (sh) (ε 23500); 243 (ε 26900); 318 (ε 8100) (MeOH/HCl). λ_{max} 221 (ε 23300); 252 (ε 21100); 293 (ε 16600); 342 (ε 9000) (MeOH/NaOH). λ_{max} 219 (ε 26100); 242 (ε 23500); 317 (ε 6800) (MeOH).1''-Deoxy, 4''-ketone: **Karnamicin B₁**. BU 3557B₁. Antibiotic BU 3557B₁ [122535-51-7]**K-16**C₁₆H₁₉N₃O₅S 365.409Prod. by *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Needles (CHCl₃/hexane). Sol. MeOH, C₆H₆, EtOAc, DMSO, CHCl₃, EtOH; poorly sol. hexane, H₂O. Mp 151-153°. [α]_D²⁵ 0 (c, 0.5 in CHCl₃). λ_{max} 220 (sh) (ε 23900); 243 (ε 26400); 318 (ε 8000) (MeOH/HCl). λ_{max} 220 (ε 23300); 251 (ε 21300); 292 (ε 16600); 341 (ε 9100) (MeOH/NaOH). λ_{max} 219 (ε 25900); 242 (ε 23600); 316 (ε 6900) (MeOH).4''-Deoxy: **Karnamicin C₅**. BU 3557C₅.Antibiotic BU 3557C₅ [122535-58-4]C₁₆H₂₁N₃O₅S 367.425Prod. by *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, EtOH, CHCl₃, EtOAc, DMSO, C₆H₆; poorly sol. hexane, H₂O. Mp 138-140°. [α]_D²⁵ +19 (c, 1 in CHCl₃). λ_{max} 220 (ε 25700); 243 (ε 23900); 317 (ε 7100) (MeOH). λ_{max} 220 (ε 24600); 243 (ε 24700); 317 (ε 7400) (MeOH/HCl). λ_{max} 217 (ε 22200); 252 (ε 20700); 283 (ε 16700); 343 (ε 8900) (MeOH/NaOH).1'',4''-Dideoxy, 3''-hydroxy: **Karnamicin C₁**. BU 3557C₁. Antibiotic BU 3557C₁ [122535-54-0]C₁₆H₂₁N₃O₅S 367.425Prod. by *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, C₆H₆, EtOAc, DMSO, EtOH, CHCl₃; poorly sol. hexane, H₂O. Mp 152.5-154°. [α]_D²⁵ +14 (c, 0.25 in CHCl₃). λ_{max} 220 (ε 25000); 243 (ε 22200); 317 (ε 6600) (MeOH). λ_{max} 222 (sh) (ε 23100); 242 (ε 23500); 317 (ε 7500) (MeOH/HCl). λ_{max} 220 (ε 22100); 252 (ε 19800); 294 (ε 15700); 342 (ε 8400) (MeOH/NaOH).Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (*Karnamicins*, isol, struct, props)Umamura, K. *et al.*, *Tet. Lett.*, 1997, **38**, 4811-4814 (*Karnamicin B₁*, synth)**Karnamicin A₂**BU 3557A₂. Antibiotic BU 3557A₂ [122535-49-3]As *Karnamicin A₁*, K-17 with R = -''CH(OH)CH₂CH₂C(OH)(CH₃)₂C₁₇H₂₃N₃O₆S 397.451Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, CHCl₃, EtOAc, DMSO, C₆H₆, EtOH; poorly sol. hexane, H₂O. Mp 60-63°. [α]_D²⁵ -14 (c, 0.5 in CHCl₃). λ_{max} 220 (ε 23900); 243 (ε 23900); 318 (ε 6900) (MeOH/HCl). λ_{max} 219 (ε 25500); 242 (ε 23500); 316 (ε 6800) (MeOH). λ_{max} 220 (ε 23400); 251 (ε 21500); 282 (ε 17000); 342 (ε 9000) (MeOH/NaOH).1''-Deoxy: **Karnamicin B₃**. BU 3557B₃.Antibiotic BU 3557B₃ [122535-53-9]C₁₇H₂₃N₃O₅S 381.452Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Rods (CHCl₃). Sol. MeOH, EtOH, CHCl₃, C₆H₆, DMSO, EtOAc; poorly sol. hexane, H₂O. Mp 187.5-189.5°. [α]_D²⁵ 0 (c, 0.5 in CHCl₃). λ_{max} 220 (ε**K-18**

25400); 242 (ε 23200); 317 (ε 6800) (MeOH). λ_{max} 220 (sh) (ε 22100); 243 (ε 26500); 318 (ε 7900) (MeOH/HCl). λ_{max} 218 (ε 21400); 252 (ε 20200); 294 (ε 16200); 342 (ε 8700) (MeOH/NaOH).

Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (Karnamicins, *isol, struct, props*)

Karnamicin C₂**K-19**

BU 3557C₂. Antibiotic BU 3557C₂ [122535-55-1]

As Karnamicin A₁, K-17 with R = -CH₂(CH₂)₂CH(CH₃)CH₂OH

C₁₇H₂₃N₃O₅S 381.452

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Mp 130.5-133°. [α]_D²⁵ -8 (c, 0.25 in CHCl₃). λ_{max} 220 (ε 26000); 243 (ε 24000); 318 (ε 7100) (MeOH). λ_{max} 222 (sh) (ε 22500); 243 (ε 26600); 318 (ε 8200) (MeOH/HCl). λ_{max} 217 (ε 22100); 253 (ε 20600); 294 (ε 16800); 342 (ε 9100) (MeOH/NaOH).

Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (*isol, struct, props*)

Karnamicin C₃**K-20**

BU 3557C₃. Antibiotic BU 3557C₃ [122535-56-2]

As Karnamicin A₁, K-17 with R = -CH₂CH₂³CH(CH₃)⁴CH(OH)CH₃

C₁₇H₂₃N₃O₅S 381.452

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Mp 165.5-167.5°. [α]_D²⁵ +20 (c, 1 in CHCl₃). λ_{max} 220 (ε 25800); 242 (ε 23200); 317 (ε 6800) (MeOH). λ_{max} 222 (sh) (ε 25000); 244 (ε 28300); 318 (ε 8400) (MeOH/HCl). λ_{max} 220 (ε 17300); 251 (ε 21000); 294 (ε 16600); 341 (ε 8800) (MeOH/NaOH).

4'-Ketone: **Karnamicin C₄**. BU 3557C₄. Antibiotic BU 3557C₄ [122535-57-3]

C₁₇H₂₁N₃O₅S 379.436

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, EtOH, CHCl₃, EtOAc, DMSO, C₆H₆; poorly sol. hexane, H₂O. Mp 124.5-126.5°. [α]_D²⁵ 0 (c, 1 in CHCl₃). λ_{max} 220 (ε 25900); 243 (ε 23900); 317 (ε 7100) (MeOH). λ_{max} 223 (sh) (ε 23200); 243 (ε 26100); 318 (ε 8000) (MeOH/HCl). λ_{max} 217 (ε 22300); 252 (ε 20600); 294 (ε 16700); 342 (ε 9100) (MeOH/NaOH).

4'-Deoxy, 3'-hydroxy: **Karnamicin D₃**. BU 3557D₃. Antibiotic BU 3557D₃ [122535-61-9]

C₁₇H₂₃N₃O₅S 381.452

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, EtOH, CHCl₃, EtOAc, DMSO, C₆H₆; poorly sol. hexane, H₂O. Mp 153-154°. [α]_D²⁵ +4 (c, 0.18 in CHCl₃). λ_{max} 220 (ε 25200); 244 (ε 23200); 316 (ε 7300) (MeOH). λ_{max} 220 (sh) (ε 21000); 244 (ε 25700); 318 (ε 7800) (MeOH/HCl). λ_{max} 221 (ε 23300); 254 (ε 22200); 294 (ε 17900); 344 (ε 9800) (MeOH/NaOH).

Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (Karnamicins, *isol, struct, props*)

Karnamicin D₂**K-21**

BU 3557D₂. Antibiotic BU 3557D₂ [122535-60-8]

As Karnamicin A₁, K-17 with -CH₂(CH₂)₂^{4'}CH₂⁵COCH₃

C₁₇H₂₁N₃O₅S 379.436

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, EtOAc, DMSO, C₆H₆, EtOH, CHCl₃; poorly sol. hexane, H₂O. Mp 139-140°. [α]_D²⁵ 0 (c, 0.2 in CHCl₃). λ_{max} 220 (ε 25800); 244 (ε 23900); 315 (ε 7900) (MeOH). λ_{max} 219 (sh) (ε 21400); 244 (ε 25400); 319 (ε 7900) (MeOH/HCl). λ_{max} 220 (ε 21500); 254 (ε 20900); 294 (ε 16800); 342 (ε 9400) (MeOH/NaOH).

5'-Deoxy, 4'-oxo: **Karnamicin D₄**. BU 3557D₄. Antibiotic BU 3557D₄ [122535-62-0]


C₁₇H₂₁N₃O₅S 379.436

Isol. from *Saccharothrix aerocolonigenes*. Active against fungi and yeasts. Powder. Sol. MeOH, C₆H₆, DMSO, EtOAc, CHCl₃, EtOH; poorly sol. hexane, H₂O. Mp 164-165.5°. [α]_D²⁵ 0 (c, 0.2 in CHCl₃). λ_{max} 220 (ε 25400); 244 (ε 24000); 315 (ε 7500) (MeOH). λ_{max} 219 (sh) (ε 22000); 244 (ε 26200); 319 (ε 7900) (MeOH/HCl). λ_{max} 221 (ε 22400); 254 (ε 21600); 293 (ε 17200); 344 (ε 9700) (MeOH/NaOH).

Nishio, M. *et al.*, *J. Antibiot.*, 1989, **42**, 852-868 (*isol, struct, props*)

Kauluamine**K-22**

[170474-99-4]



Absolute Configuration


C₇₂H₉₄N₈O₃ 1119.586

Alkaloid from an Indonesian marine sponge, *Prianos* sp. Shows mod. immunosuppressive activity. Unstable pale yellow solid. Sol. MeOH, CHCl₃, CH₂Cl₂. [α]_D²⁵ +0.7 (c, 0.18 in CHCl₃).

Ohtani, I.I. *et al.*, *J.A.C.S.*, 1995, **117**, 10743-10744 (*isol, ir, pmr, cmr, struct*)

Kayawongine**K-23**

2-(3,4-Dimethoxyphenyl)-3-(4-methoxyphenyl)quinolizidine [86688-02-0]



C₂₄H₃₁NO₃ 381.514


Major alkaloid from the leaves of *Cissus rheifolia* (Vitidaceae). Mp 124-124.5°.

[α]_D²⁵ -116 (c, 0.85 in CHCl₃).

Saifah, E. *et al.*, *J. Nat. Prod.*, 1983, **46**, 353 (*isol, uv, pmr, cmr, ms, cd, struct*)

Kealiinine A**K-24**

[700813-15-6]



C₂₀H₁₉N₃O₃ 349.388

Alkaloid from the sponge *Leucetta chagosensis*. Cytotoxic. Yellow-brown powder. λ_{max} 250 ; 314 ; 338 (MeOH).

6-Me ether: **Kealiinine B**

[700813-16-7]

C₂₁H₂₁N₃O₃ 363.415

Alkaloid from *Leucetta chagosensis*.

Dark brown solid. λ_{max} 250 ; 314 ; 338 (MeOH).

5-Methoxy, 6-Me ether: **Kealiinine C**

[700813-17-8]

C₂₂H₂₃N₃O₄ 393.441


Alkaloid from *Leucetta chagosensis*.

Dark brown solid. λ_{max} 250 ; 314 ; 338 (MeOH).

Hassan, W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 817-822 (*isol, pmr, cmr, ms*)

Kealiiquinone**K-25**

6,7-Dimethoxy-4-(4-methoxyphenyl)-1-methyl-1H-naphth[2,3-d]imidazole-2,5,8(3H)-trione, 9CI [124535-78-0]



C₂₁H₁₈N₂O₆ 394.383

CAS registry no. refers to the trioxo tautomer. Alkaloid from the sponge *Leucetta* sp. Long red needles (MeOH/CH₂Cl₂/hexane). Mp 300° dec. λ_{max} 230 (ε 15100); 296 (ε 22400); 388 (ε 1380) (MeOH) (Derep).

2-Deoxy, 2-amino: **2-Amino-2-deoxykealiiquinone**. 2-Deoxy-2-aminokealiiquinone [189748-83-2]

C₂₁H₁₉N₃O₅ 393.398

Alkaloid from the sponge *Leucetta chagosensis*. Red needles (MeOH/CH₂Cl₂). Mp > 250° dec. λ_{max} 220 (log

ϵ 4.49); 294 (log ϵ 4.57); 382 (log ϵ 3.73) (MeOH).


Akee, R.K. *et al.*, *J.O.C.*, 1990, **55**, 1944 (*isol, uv, ir, pmr, cmr, cryst struct*)

Kawasaki, I. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1393-1398 (*synth, uv, ir, pmr, cmr, cryst struct*)

Fu, X. *et al.*, *J. Nat. Prod.*, 1997, **60**, 497 (2-Deoxy-2-aminoketiquinone)

Kedarcidin**K-26**

[128512-39-0]



$C_{53}H_{60}ClN_3O_{16}$ 1030.52

Chromoprotein enediyne antibiotic. Struct. of chromophore shown. Apoptein moiety is a linear polypeptide consisting of 114 amino acid residues of known struct. Struct. of chromophore revised in 1997 and stereochem. again in 2007. Prod. by *Streptoalloteichus* sp. LS85-6. Antineoplastic agent. Active against gram-positive bacteria. Buff powder. Sol. EtOAc. λ_{max} 258 (ϵ 43800); 306 (ϵ 15500); 313 (ϵ 15200) (H₂O/protein) (Derep). λ_{max} 256 (ϵ 60300); 316 (ϵ 14500) (MeOH) (Derep).

Lam, K.S. *et al.*, *J. Antibiot.*, 1991, **44**, 472 (*props*)

Hofstead, S.J. *et al.*, *J. Antibiot.*, 1992, **45**, 1250 (*isol*)

Leet, J.E. *et al.*, *J.A.C.S.*, 1992, **114**, 7946; 1993, **115**, 8432 (*struct, pmr, cmr, ms*)

Zein, N. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 8009-8012 (*pharmacol*)

Constantine, K.L. *et al.*, *Biochemistry*, 1994, **33**, 11438 (*bibl*)

Lam, K.S. *et al.*, *J. Ind. Microbiol. Biotechnol.*, 1994, **13**, 356-360 (*manuf*)


Zein, N. *et al.*, *NATO ASI Ser. Ser. C*, 1996, **479**, 53 (*rev*)

Kawata, S. *et al.*, *J.A.C.S.*, 1997, **119**, 12012-12013 (*synth, struct*)

Ren, F. *et al.*, *J.A.C.S.*, 2007, **129**, 5381-5383 (*synth*)

Keenamide A**K-27**

[177742-52-8]




$C_{30}H_{48}N_6O_6S$ 620.812

Isol. from the marine mollusc *Pleurobranchus forskalii*. Cytotoxic agent. Off-white powder. $[\alpha]_D^{+24}$ (c, 0.3 in MeOH).

Wesson, K.J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 629-631 (*isol, ir, pmr, cmr, ms*)

Keitine**K-28**

[40766-51-6]



Absolute configuration

$C_{21}H_{29}NO_4$ 359.464

Alkaloid from *Liparis keitaoensis* (Orchidaceae). Noncryst. $[\alpha]_D^{23}$ -7 (c, 0.6 in EtOH). Ester of Laburnine.

O^{4'}- β -D-Glucopyranoside: **Keitaoine**

[40766-50-5]

$C_{27}H_{39}NO_9$ 521.606

Alkaloid from *Liparis keitaoensis* (Orchidaceae). Noncryst. $[\alpha]_D^{23}$ -13 (c, 0.8 in EtOH).

3'-Demethoxy, *O*^{4'}- β -D-glucopyranoside:

Malaxine

[19128-95-1]

$C_{26}H_{37}NO_8$ 491.58

Alkaloid from *Liparis bicallosa*, *Liparis hachijoensis* and *Malaxis conges-ta* (Orchidaceae). Mp 151-159°. $[\alpha]_D^{22}$ -31 (c, 2.7 in EtOH).

1-Epimer, *O*^{4'}- β -D-glucopyranoside:

Hammarbine

[40766-49-2]

$C_{27}H_{39}NO_9$ 521.606

Minor alkaloid of *Hammarbya paludosa* (Orchidaceae). Amorph. $[\alpha]_D^{23}$ +9 (c, 0.31 in EtOH). Hydrol. \rightarrow Lindolofidine.

Leander, K. *et al.*, *Tet. Lett.*, 1967, 3477 (*isol, Malaxine*)


Nishikawa, K. *et al.*, *Tetrahedron*, 1969, **25**, 2723 (*struct, Malaxine*)

Tanino, H. *et al.*, *Tetrahedron*, 1969, **25**, 3033 (*synth, Malaxine*)

Linström, B. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 2963 (*Keitine, Keitaoine, Hammarbine*)

Keramidine**K-29**

[88839-97-8]



$C_{12}H_{14}BrN_5O$ 324.18

Isol. from the Okinawan sea sponge *Agelas* sp. Antagonist of serotonergic receptors. Powder. Sol. MeOH, CHCl₃. Mp 183-187°. λ_{max} 269 (ϵ 21400) (MeOH) (Derep).

9,10-Dihydro: **9,10-Dihydrokeramidine**

$C_{12}H_{16}BrN_5O$ 326.195

Isol. from an *Agelas* sp. Amorph. solid.

λ_{max} 202 (ϵ 21200); 280 (ϵ 19700) (MeOH).

Nakamura, H. *et al.*, *Tet. Lett.*, 1984, **25**, 2475-2478 (*isol, uv, ir, pmr, cmr, ms*)

Daninos-Zeghal, S. *et al.*, *Tetrahedron*, 1997, **53**, 7605-7614 (*synth*)


Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 122-125 (*isol, pmr*)

Lindel, T. *et al.*, *Tet. Lett.*, 1998, **39**, 2541-2544 (*synth, pmr*)

Endo, T. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1262-1267 (9,10-Dihydrokeramidine)

Keramamide B**K-30**

[137041-25-9]

R¹ = R² = -CH₂CH₃

$C_{54}H_{77}BrN_{10}O_{12}$ 1138.166

Cyclic peptide. Isol. from the marine sponge *Theonella* sp. $[\alpha]_D^{23}$ -50 (c, 0.7 in MeOH). λ_{max} 267 (ϵ 23500); 312 (ϵ 4300) (MeOH) (Derep). λ_{max} 267 (ϵ 23500); 312 (ϵ 4300) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *J.A.C.S.*, 1991, **113**, 7812-7813 (*isol, pmr, cmr*)

Shioiri, T. *et al.*, *Heterocycles*, 2003, **61**, 23-29 (*synth*)

Keramamide D**K-31**

[137041-27-1]

As Keramide B, K-30 with

R¹ = R² = -CH₃

$C_{52}H_{73}BrN_{10}O_{12}$ 1110.112

Cyclic peptide. Isol. from the marine sponge *Theonella* sp. λ_{max} 267 (ϵ 23500); 312 (ϵ 4300) (MeOH) (Derep).

9''-O-Sulfate: **Keramamide M**

$C_{52}H_{73}BrN_{10}O_{15}S$ 1190.177

Isol. from a *Theonella* sp. Amorph. solid. λ_{max} 269 (ϵ 20000) (MeOH).

Kobayashi, J. *et al.*, *J.A.C.S.*, 1991, **113**, 7812-7813 (*isol*)

Tsuda, M. *et al.*, *Tetrahedron*, 1999, **55**, 12543-12548 (*Keramamide M*)

Keramamide E**K-32**

[161995-32-0]

As Keramide B, K-30 with

R¹ = CH₃, R² = CH₂CH₃

$C_{53}H_{75}BrN_{10}O_{12}$ 1124.139

Cyclic peptide. Isol. from the sponge *Theonella* sp. Solid. $[\alpha]_D^{22}$ -39 (c, 0.1 in MeOH). λ_{max} 269 (ϵ 30900); 310 (sh) (MeOH). λ_{max} 269 (ϵ 30900); 310 (MeOH) (Berdy).

9''-O-Sulfate: **Keramamide N**

$C_{53}H_{75}BrN_{10}O_{15}S$ 1204.204


Isol. from a *Theonella* sp. Amorph. solid. λ_{max} 270 (ϵ 18000) (MeOH).

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**, 2525-2532 (*isol, uv, ir, pmr, cmr, ms*)

Tsuda, M. *et al.*, *Tetrahedron*, 1999, **55**, 12543-12548 (*Keramamide N*)

Keramamide F**K-33**

[143330-97-6]



$C_{43}H_{56}N_{10}O_{11}S$ 921.042

Cyclic thiopeptide antibiotic. Isol. from a *Theonella* sp. Amorph. Mp 187° dec. $[\alpha]_D^{21}$ -25 (c, 0.86 in MeOH). λ_{max} 209 (ε 30800); 220 (ε 30800); 275 (ε 23200); 339 (ε 9800) (MeOH) (Berdy).

13-Epimer: Keramamide G

[162063-07-2]

$C_{43}H_{56}N_{10}O_{11}S$ 921.042


Isol. from a *Theonella* sp. Solid. $[\alpha]_D^{21}$ +10 (c, 0.1 in MeOH). λ_{max} 224 (ε 26500); 279 (ε 21200); 340 (ε 11700) (MeOH) (Berdy).

Itagaki, F. *et al.*, *J.O.C.*, 1992, **57**, 5540 (*isol*, *pmr*, *cmr*, *struct*)

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**, 2525 (*Keramamide G*)

Keramamide H**K-34**

[161995-33-1]



$C_{43}H_{57}BrN_{10}O_{12}S$ 1017.953

Cyclic peptide. Isol. from the sponge *Theonella* sp. Solid. $[\alpha]_D^{20}$ -42 (c, 0.05 in MeOH). λ_{max} 233; 277 (ε 23100) (MeOH). λ_{max} 277 (ε 23100) (MeOH) (Berdy).

2'-Debromo, 5'-deoxy: Keramamide J

[161995-34-2]

$C_{43}H_{58}N_{10}O_{11}S$ 923.058

Isol. from *Theonella* sp. Solid. $[\alpha]_D^{18}$ +8.4 (c, 0.1 in MeOH). λ_{max} 222 (ε 45200); 278 (ε 27200) (MeOH). λ_{max} 228 (ε 45200); 278 (ε 27200) (MeOH) (Berdy).

2'-Debromo, 5'-deoxy, N¹-Me: Keramide K

$C_{44}H_{60}N_{10}O_{11}S$ 937.084

Isol. from the sponge *Theonella* sp. Cytotoxic agent. Amorph. solid. $[\alpha]_D^{28}$ -25 (c, 0.1 in MeOH). λ_{max} 269 (ε 20000) (MeOH). λ_{max} 269 (ε 20000) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**,


2525-2532 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Uemoto, H. *et al.*, *Tetrahedron*, 1998, **54**, 6719-6724 (*Keramamide K*)

Sowinski, J.A. *et al.*, *Chem. Comm.*, 1999, 981-982 (*synth*, *Keramamide J*)

Keronopsin B₁**K-35**

[158182-29-7]



$C_{18}H_{16}BrNO_3$ 374.233

Alkaloid from the marine ciliate *Pseudokeronopsis rubra*. Chemical defence substance. Black cryst. (Me₂CO). Mp 135-139°. λ_{max} 229 (ε 14454); 266 (ε 18620); 421 (ε 70794) (MeOH) (Berdy).

O-Sulfate: Keronopsin A₁

[158182-31-1]

$C_{18}H_{16}BrNO_6S$ 454.297

Alkaloid from *Pseudokeronopsis rubra*. Brick-red amorph. powder (as Na salt). Unstable in the crude extract. Converted into Keronopsin B₁ with traces of acid. In dry state may polymerise spontaneously. CAS no. refers to Na salt. λ_{max} 221; 267; 325; 339; 439 (MeOH) (Berdy).

17-Bromo: Keronopsin B₂

[158182-30-0]

$C_{18}H_{15}Br_2NO_3$ 453.129

Alkaloid from *Pseudokeronopsis rubra*. Brown-red amorph. powder. λ_{max} 229 (ε 15850); 439 (ε 61659) (MeOH) (Berdy).

17-Bromo, O-sulfate: Keronopsin A₂

[158182-32-2]


$C_{18}H_{15}Br_2NO_6S$ 533.194

Alkaloid from *Pseudokeronopsis rubra*. Unstable in the crude extract. Converted into Keronopsin B₂ with traces of acid. CAS no. refers to Na salt. λ_{max} 222; 269; 326; 434 (MeOH) (Berdy).

Höfle, G. *et al.*, *Angew. Chem., Int. Ed.*, 1994, **33**, 1495-1497 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Kesselridine**K-36****O¹²-Demethylkesselringine**

[56775-79-2]



Absolute Configuration

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from *Colchicum kesselringii* (Liliaceae). Mp 232-234°. $[\alpha]_D$ -50 (Py).

O⁶-Me: Regelamine. Norregeline. Regalamine

[58111-42-5]

$C_{19}H_{25}NO_4$ 331.411

Alkaloid from *Colchicum kesselringii* (Liliaceae). Mp 226°. $[\alpha]_D$ +33

(MeOH). Not correlated with Kesselridine. λ_{max} 218 (log ε 4.34); 290 (log ε 3.59) (MeOH).

O⁶-Me, N-de-Me: Jolantidine. Iolantidine

[89759-22-8]

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from above-ground parts of *Merendera jolantae* (Liliaceae). Cryst. (Me₂CO). Mp 275-277°. $[\alpha]_D$ +102 (c, 0.4 in MeOH). λ_{max} 216; 290 (no solvent reported).

O¹²-Me: Kesselringine

[54692-48-7]

$C_{19}H_{25}NO_4$ 331.411

Alkaloid from *Colchicum kesselringii* (Liliaceae). Mp 194-196°. $[\alpha]_D^{18}$ +93.2 (c, 1.02 in CHCl₃). $[\alpha]_D^{25}$ +53.3 (c, 3.0 in MeOH). λ_{max} 231 (log ε 3.85); 293 (log ε 3.44) (EtOH). λ_{max} 224 (log ε 4.3); 255 (log ε 3.76); 305 (log ε 3.56) (EtOH/NaOH).

O¹²-Me; hydrochloride: Mp 252-253°.

O⁶, O¹²-Di-Me: Regeline

[62653-08-1]

$C_{20}H_{27}NO_4$ 345.438

Alkaloid from *Colchicum kesselringii* (Liliaceae). Cryst. (Me₂CO). Mp 198-200°. $[\alpha]_D$ +93 (c, 1.5 in MeOH). λ_{max} 216 (log ε 4.04); 225 (log ε 3.96); 290 (log ε 3.37) (no solvent reported).

O⁶, O¹²-Di-Me; hydrochloride:

Cryst. (Me₂CO). Mp 249-250°.

11-Epimer, O¹²-Me: Luteine†

[62624-08-2]

$C_{19}H_{25}NO_4$ 331.411

Alkaloid from *Colchicum luteum* (Liliaceae). Mp 228-230°. $[\alpha]_D$ +93 (MeOH). Proposed struct. λ_{max} 210 (log ε 4.7); 285 (log ε 3.7) (MeOH).

11-Epimer, O⁶, O¹²-di-Me: Regelinine.*Regilinine*

[97806-68-3]

$C_{20}H_{27}NO_4$ 345.438

Alkaloid from *Colchicum kesselringii* aerial parts (Liliaceae). Mp 253-254°. Unstable, readily oxidised. Methylation gives Luteine. λ_{max} 243; 293 (no solvent reported).

Kasimov, A.K. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 194-197; 383; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 202-205; 395

(Regelamine, *uv*, *ir*, *pmr*, *ms*, *struct*)

Yusupov, M.K. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 350-354; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 305-308 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Abdullaeva, D.A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 783-787; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 702-705 (*Regeline*)

Mukhamedyarova, N.L. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 801-804; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 718-721 (*Luteine*)


Yusupov, M.K. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 1581-1587 (*pmr*, *struct*, *synth*, *Kesselridine*, *Regelamine*)

Chommadov, B. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 790-791; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 755 (*Jolantidine*)

Yusupov, M.K. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 419-420; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 396-397 (*Regelinine*)

Ketoanhydrokinamycin K-37

1a,2,4,9,10,10a-Hexahydro-5,10-dihydroxy-1a-methyl-2,4,9-trioxo-3H-benz[*b*]oxireno[*h*]carbazole-3-carbonitrile, 9CI. Antibiotic Y1†
[120796-25-0]




C₁₈H₁₀N₂O₆ 350.287
Metab. of *Streptomyces murayamaensis*.
Mp 300° dec. λ_{max} 236 (ε 43300); 255 (ε 28100); 306 (ε 18600); 410 (ε 9010); 453 (ε 9790) (CHCl₃) (Derep).

Seaton, P.J. et al., *J. Antibiot.*, 1989, **42**, 179; 189 (isol, pmr, cmr, struct)

31-Keto-12,34-oxa-32,33-dihydroircinal A K-38

[721429-61-4]



Absolute Configuration


C₂₆H₃₆N₂O₃ 424.582

Related to Ircinol A, I-174. Isol. from an *Acanthostrongylophora* sp. Powder (MeOH). Mp 164° dec. [α]_D²⁵ +44 (c, 0.1 in CHCl₃). λ_{max} 235 (log ε 3.93) (MeOH).

Yousaf, M. et al., *J. Med. Chem.*, 2004, **47**, 3512-3517 (isol, pmr, cmr)

Khyberine K-39

[77795-10-9]



C₃₆H₃₈N₂O₆ 594.706

Trace alkaloid from the roots of *Berberis calliobotrys*. Also obt. by the acid-catalysed dienone-phenol rearr. of Epivaldiberine (Berberidaceae). Cryst. (CHCl₃/MeOH). Mp 145-147°.

7-Me ether: **Kalashine**

[76372-24-2]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the roots of *Berberis calliobotrys* and *Berberis orthobotrys* (Berberidaceae). Obt. in too low a yield for crystallisation.


Hussain, S.F. et al., *Tet. Lett.*, 1980, **21**, 3315; 4573 (isol, uv, cd, pmr, ms, struct, abs config)

Hussain, S.F. et al., *J. Nat. Prod.*, 1981, **44**, 274 (isol, uv, Kalashine)

Guinaudeau, H. et al., *Chem. Comm.*, 1982, 1122 (synth)

Kibdelone A K-40

[934464-77-4]



Relative Configuration

C₂₉H₂₄ClNO₁₀ 581.962

Prod. by *Kibdelosporangium* sp. (MST-108465). Cytotoxic. Orange solid. [α]_D²⁵ +72 (c, 0.01 in CHCl₃). λ_{max} 214 (ε 24500); 254 (ε 24900); 311 (ε 20000); 420 (ε 11000); 447 (sh) (EtOH).

12-O-*α*-L-Rhamnopyranoside: **Kibdelone A rhamnoside**

[934464-80-9]

C₃₅H₃₄ClNO₁₄ 728.105

Prod. by *Kibdelosporangium* sp. (MST-108465).

10-Ketone: **10-Oxokibdelone A**

[934464-83-2]

C₂₉H₂₂ClNO₁₀ 579.946

Prod. by *Kibdelosporangium* sp. (MST-108465).

6,7-Dihydro: **Kibdelone B**

[934464-78-5]

C₂₉H₂₆ClNO₁₀ 583.978

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. orange solid. [α]_D²⁵ +157 (c, 0.01 in CHCl₃). λ_{max} 208 (ε 22000); 258 (ε 23800); 308 (sh); 402 (ε 9670); 444 (sh) (EtOH).

6,7-Dihydro, 12-O-*α*-L-rhamnopyranoside:

Kibdelone B rhamnoside

[934464-81-0]

C₃₅H₃₆ClNO₁₄ 730.12

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. orange solid. [α]_D²⁵ +150 (c, 0.1 in CHCl₃). λ_{max} 205 (ε 18600); 258 (ε 16300); 308 (sh); 435 (ε 5700) (EtOH).

5,16-Hydroquinone, 6,7-dihydro: **Kibdelone C**

[934464-79-6]

C₂₉H₂₈ClNO₁₀ 585.994

Prod. by *Kibdelosporangium* sp. (MST-108465). Amorph. yellow solid. [α]_D²⁵ +49 (c, 0.01 in CHCl₃). λ_{max} 217 (ε 22800); 258 (ε 27500); 272 (sh); 308 (sh); 339 (sh); 396 (ε 20100) (EtOH).

5,16-Hydroquinone, 6,7-dihydro, 12-O-*α*-L-rhamnopyranoside: **Kibdelone C rhamnoside**

[934464-82-1]


C₃₅H₃₈ClNO₁₄ 732.136

Prod. by *Kibdelosporangium* sp. (MST-108465).

Ratnayake, R. et al., *Chem. Eur. J.*, 2007, **13**, 1610-1619 (isol, pmr, cmr)

Kifunensine K-41

Hexahydro-6,7,8-trihydroxy-5-(hydroxymethyl)imidazo[1,2-*a*]pyridine-2,3-dione, 9CI. FR 900494. Antibiotic FR 900494
[109944-15-2]



C₈H₁₂N₂O₆ 232.193

Prod. by *Kitasatospora kifunense*. Immunomodulator, α-mannosidase inhibitor. Prisms (H₂O). Sol. H₂O; fairly sol. MeOH, EtOH; poorly sol. Me₂CO, CHCl₃. Mp 120-136° dec Mp 280°. [α]_D²⁵ +58 (c, 0.1 in H₂O). Genus name given as Kitasatospora, but Kitasatospora is preferred. λ_{max} 222 (sh) (ε 5600) (H₂O/HCl) (Derep). λ_{max} 224 (ε 5600) (H₂O/NaOH) (Derep). λ_{max} 226 (ε 5340) (H₂O) (Derep). λ_{max} 227 (ε 5000) (MeOH) (Berdy).


Iwami, M. et al., *J. Antibiot.*, 1987, **40**, 612 (isol, props)

Kayakiri, H. et al., *J.O.C.*, 1989, **54**, 4015 (pmr, cmr, struct)

Kayakiri, H. et al., *Chem. Pharm. Bull.*, 1991, **39**, 1378; 1392; 1397 (pmr, cmr, cryst struct, synth)

Kigamicinone K-42

[864183-17-5]



Absolute Configuration

C₂₈H₂₅NO₁₁ 551.506

Kigamicins are active against gram-positive bacteria incl. MRSA and are antitumour agents.

11-O-(2,3,6-Trideoxy-β-D-erythro-hexopyranoside): **Kigamicin A**

[680571-49-7]

C₃₄H₃₅NO₁₃ 665.649

Prod. by *Amycolatopsis* sp. ML630-mF1. Yellow powder. Mp 225-227°. [α]_D²⁴ -153 (c, 1 in MeOH). λ_{max} 217 (log ε 4.67); 236 (log ε 4.64); 254 (log ε 4.64); 280 (log ε 4.54); 341 (log ε 4.28) (MeOH).

11-O-[2,3,6-Trideoxy-β-D-erythro-hexopyranosyl-(1→4)-2,3,6-trideoxy-β-D-erythro-hexopyranoside]: **Kigamicin B**

[680571-50-0]

C₄₀H₄₅NO₁₅ 779.793

Prod. by *Amycolatopsis* sp. ML630-mF1. Yellow powder. Mp 225-227°.

λ_{\max} 217 (log ϵ 4.73); 236 (log ϵ 4.71); 254 (log ϵ 4.71); 280 (log ϵ 4.61); 341 (log ϵ 4.35) (MeOH).

11-O-[2,6-Dideoxy-3-O-methyl- β -D-arabino-hexopyranosyl-(1 \rightarrow 4)-2,3,6-trideoxy- β -D-erythro-hexopyranoside]:

Kigamicin C

[680571-51-1]

C₄₁H₄₇NO₁₆ 809.819

Prod. by *Amycolatopsis* sp. ML630-mF1. Yellow powder. Mp 210-212°. $[\alpha]_D^{24}$ -154 (c, 1 in MeOH). λ_{\max} 217 (log ϵ 4.37); 236 (log ϵ 4.34); 254 (log ϵ 4.32); 280 (log ϵ 4.22); 341 (log ϵ 4.02) (MeOH).

11-O-[2,6-Dideoxy-3-O-methyl- β -D-arabino-hexopyranosyl-(1 \rightarrow 4)-2,6-dideoxy-3-O-methyl- β -D-arabino-hexopyranosyl-(1 \rightarrow 4)-2,3,6-trideoxy- β -D-erythro-hexopyranoside]:

Kigamicin D

[680571-52-2]

C₄₈H₅₉NO₁₉ 953.989

Prod. by *Amycolatopsis* sp. ML630-mF1. Yellow powder. Mp 210-212°. $[\alpha]_D^{24}$ -190.6 (c, 1 in MeOH). λ_{\max} 217 (log ϵ 4.56); 236 (log ϵ 4.52); 254 (log ϵ 4.5); 280 (log ϵ 4.39); 341 (log ϵ 4.12) (MeOH).

11-O-[2,6-Dideoxy-3-O-methyl- β -D-arabino-hexopyranosyl-(1 \rightarrow 4)-2,6-dideoxy-3-O-methyl- β -D-arabino-hexopyranosyl-(1 \rightarrow 4)-2,3,6-trideoxy- β -D-erythro-hexopyranoside]:

Kigamicin E

[680571-53-3]

C₅₅H₇₁NO₂₂ 1098.159

Prod. by *Amycolatopsis* sp. ML630-mF1. Yellow powder. Mp 210-212°. $[\alpha]_D^{24}$ -175.2 (c, 1 in MeOH). λ_{\max} 217 (log ϵ 4.59); 236 (log ϵ 4.55); 254 (log ϵ 4.52); 280 (log ϵ 4.42); 341 (log ϵ 4.16) (MeOH).

Kunimoto, S. *et al.*, *J. Antibiot.*, 2003, **56**, 1004-1011; 1012-1017 (*isol, uv, pmr, cmr, activity*)


Someno, T. *et al.*, *J. Antibiot.*, 2005, **58**, 56-60 (*abs config*)

Kikumycin A, 8CI

K-43

TAN 868B. Antibiotic TAN 868B

[37913-77-2]

C₁₃H₁₇N₇O₂ 303.323

Oligopeptide-type antibiotic. Isol. from *Streptomyces phaeochromogenes* and *Streptomyces idiomorphus*. Shows antiviral, antitumour and antibacterial properties. Fine needles (as sulfate salt). Mp 285° dec. (sulfate). $[\alpha]_D^{25}$ +12.8 (c, 1 in 1M HCl) (sulfate). λ_{\max} 230 (ϵ 17200); 321 (ϵ 33300) (H₂O) (as hydrochloride) (Derep). λ_{\max} 240 (sh) (ϵ); 356 (ϵ 40000) (0.1M NaOH) (Derep).

► UX9353000

Pyrrolo-N-Me: **Kikumycin B**

[37913-78-3]

C₁₄H₁₉N₇O₂ 317.35

From *Streptomyces phaeochromogenes*. Antiviral, antibacterial and antitumour agent. Sol. H₂O; fairly sol. MeOH; poorly sol. butanol, hexane. Mp 220-223° (as hydrochloride). $[\alpha]_D^{25}$ +14 (c, 1 in H₂O). λ_{\max} 230 (ϵ 17200); 321 (ϵ 33300) (H₂O) (as hydrochloride) (Derep). λ_{\max} 240 (sh) (ϵ); 356 (ϵ 40000) (0.1M NaOH) (Derep). λ_{\max} 239 (E1%/1cm 430); 328 (E1%/1cm 790) (H₂O) (Berdy). λ_{\max} 242 (ϵ 10500); 320 (ϵ 19500) (HCl) (Berdy).

► LD₅₀ (mus, ipr) 200 mg/kg. UX9354000
4R-Hydroxy: **Antibiotic TAN 868A. TAN 868A**

[111631-13-1]

C₁₃H₁₇N₇O₃ 319.322

From *Streptomyces idiomorphus*. Active against bacteria, fungi and a protozoan; and possesses cytotoxic props. Powder + 2H₂O (as hydrochloride). Sol. MeOH, DMSO, H₂O; fairly sol. EtOAc; poorly sol. Et₂O, hexane. $[\alpha]_D$ +61.6 (c, 1.05 in H₂O). λ_{\max} 230 (ϵ 17200); 321 (ϵ 33300) (H₂O) (as hydrochloride) (Derep). λ_{\max} 240 (sh); 356 (ϵ 40000) (0.1M NaOH) (Derep).

► LD₅₀ (mus, ipr) 25 mg/kg, LD₅₀ (mus, scu) 50 mg/kg.

Kikuchi, M. *et al.*, *J. Antibiot., Ser. A*, 1965, **18**, 243 (*isol*)

Takaishi, T. *et al.*, *Tet. Lett.*, 1972, 1873

(*struct, ir, uv, ms, nmr*)


Takaishi, T. *et al.*, *Org. Mass Spectrom.*, 1974, **9**, 635 (*ms, struct*)

Takizawa, M. *et al.*, *J. Antibiot.*, 1987, **40**, 1220 (*isol*)

Kimbasine A

K-44

[135626-55-0]

C₂₉H₄₅NO₅ 487.678

Norsesterterpene alkaloid from the sponge *Igernella notabilis*. Cytotoxic. $[\alpha]_D$ -69.1 (c, 1.1 in CHCl₃).

O-Debutanoyl, O-Ac: **Kimbasine B**

[135626-56-1]

C₂₇H₄₁NO₅ 459.625

Alkaloid from *Igernella notabilis*. Cytotoxic agent. $[\alpha]_D$ -64.1 (c, 0.1 in CHCl₃).

Cardellina, J.H. *et al.*, *Tet. Lett.*, 1991, **32**, 2347 (*isol, pmr, cmr, struct*)

Kimorexin A

K-45

[163663-02-3]

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Kitasatospora kimorex-ae*. Antifungal agent. Genus name given


as *Kitasatospora* but *Kitasatospora* is preferred.

Yeo, W.-H. *et al.*, *J. Microb. Biotechnol.*, 1994, **4**, 349-353

Kiridine

K-46

[161068-73-1]

C₂₂H₃₁NO₆ 405.49

Alkaloid from *Aconitum kirinense*. Possible artifact.

Feng, F. *et al.*, *CA*, 1995, **122**, 128605f

Kirigamycin C

K-47

Antibiotic 2908C

Nucleoside antibiotic. Struct. unknown. Prod. by *Streptomyces* sp. 2908 and *Streptomyces microflavus*. Powder. Sol. H₂O; fairly sol. MeOH, DMSO, DMF; poorly sol. EtOH, hexane. $[\alpha]_D$ +12.8 (H₂O).


Proc. Int. Ferment. Symp., Berlin, 5th, 1976, 234

Kirinine B

K-48

1,19-Epoxydenudatine

[161068-72-0]

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Aconitum kirinense*. Cubes (CHCl₃/MeOH). Mp 157-158°.

11-Ac: 11-Acetyl-1,19-epoxydenudatine

[132215-96-4]

C₂₄H₃₃NO₄ 399.529

Alkaloid from aerial parts of *Aconitum barbatum* (Ranunculaceae).

Cryst. (CHCl₃/heptane). Mp 202-203°. $[\alpha]_D^{20}$ +99.7 (c, 1 in CHCl₃).

O¹¹-Me: **Vilmorinianine. 1,19-Epoxy-11-methoxydenudatine**

[203513-11-5]

C₂₃H₃₃NO₃ 371.519

Alkaloid from *Aconitum vilmorinianum* var. *altifidum*. $[\alpha]_D$ +60 (c, 0.4 in CHCl₃).


Proksa, B. *et al.*, *Planta Med.*, 1990, **56**, 461 (11-Ac)

Zhang, J. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1997, **39**, 582-584; *CA*, **128**, 190400n (*Vilmorinianine*)

Feng, F. *et al.*, *Phytochemistry*, 1998, **49**, 2557-2559 (*isol, ir, pmr, cmr, ms*)

Kirondrine

[198541-68-3]


C₃₀H₂₈N₄O₄ 508.576

Dimer of 4,7-Dimethoxy-1-vinyl-β-carboline, D-723. Alkaloid from the root bark of *Perriera madagascariensis* (Simaroubaceae). Mp 267°. [α]_D -15.6 (c, 0.5 in MeOH) (hydrochloride). It is not clear whether the Mp given refers to the free base or the hydrochloride. λ_{max} 210 (log ε 3.77); 247 (log ε 4.16); 300 (log ε 3.55); 336 (log ε 3.76); 367 (log ε 3.65) (MeOH).

Krebs, H.C. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1183-1185 (*isol, uv, ir, pmr, cmr*)

Kisantine†

[1673-92-3]

C₂₁H₂₈N₂O₃ 356.464


Alkaloid from the roots of *Tabernanthe iboga* (Apocynaceae). Cryst. (EtOH). Mp 236-238°. [α]_D²⁵ -15 (CHCl₃). No further work reported since 1965. λ_{max} 213 (ε 31700); 269 (ε 6270); 296 (sh) (ε 4710) (no solvent reported).

Dickel, D.F. *et al.*, *J.A.C.S.*, 1958, **80**, 123-125 (*isol, uv, ir*)

Taylor, W.I. *et al.*, *J.O.C.*, 1965, **30**, 309-310 (*uv, ir, pmr, ms, struct*)

Kisantine†

K-51

C₄₄H₅₀N₄O₁₁ 810.899

Alkaloid from the leaves of *Pterotaberna inconspicua* (Apocynaceae). [α]_D -21 (c, 0.33 in MeOH). Unstable.

Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 1085 (*isol, uv, ir, pmr, cmr, ms, struct*)

K-49


KitagineC₇H₇NO₃ 153.137

Struct. unknown. Constit. of *Canavalia ensiformis* (jack bean) seeds (Fabaceae). Mp 240-242°. Opt. inactive.

Ackermann, D. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1939, **262**, 103-110 (*isol*)

Kobutimycin A

[145458-91-9]

R = CH₃C₁₉H₂₅NO₅ 347.41

Prod. by a *Streptomyces* sp. Herbicide. Oil. Sol. MeOH, hexane; poorly sol. H₂O. [α]_D²⁵ +88.2 (c, 1 in CHCl₃). λ_{max} 230 (ε 11700); 322 (ε 11500) (MeOH/HCl) (Derep). λ_{max} 220 (sh) (ε 6800); 258 (ε 16300) (MeOH/NaOH) (Derep). λ_{max} 210 (ε 5600); 257 (ε 14500) (MeOH) (Derep).

Kanbe, K. *et al.*, *J. Antibiot.*, 1992, **45**, 1700 (*isol, pmr, cmr, struct*)

Kobutimycin B

[145458-92-0]

As Kobutimycin A, K-53 with


R = CH₂CH₃C₂₀H₂₇NO₅ 361.437

Prod. by a *Streptomyces* sp. Herbicide. Oil. Sol. MeOH, hexane; poorly sol. H₂O. [α]_D²² +107.2 (c, 1 in CHCl₃). λ_{max} 230 (ε 11700); 322 (ε 11500) (MeOH/HCl) (Derep). λ_{max} 220 (sh) (ε 6800); 258 (ε 16300) (MeOH/NaOH) (Derep). λ_{max} 210 (ε 5600); 257 (ε 14500) (MeOH) (Derep). λ_{max} 207 (ε 7180); 256 (ε 16100) (MeOH) (Berdy). λ_{max} 232 (ε 12900); 322 (ε 12800) (MeOH/HCl) (Berdy). λ_{max} 211 (ε 7000); 257 (ε 15400) (MeOH/NaOH) (Berdy).

Kanbe, K. *et al.*, *J. Antibiot.*, 1992, **45**, 1700 (*isol, pmr, cmr, struct*)

Koeniginequinone A

7-Methoxy-3-methyl-1H-carbazole-1,4(9H)-dione. 7-Methoxy-3-methyl-1,4-carbazolequinone [110519-58-9]

C₁₄H₁₁NO₃ 241.246

K-52

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree). Red needles (C₆H₆/petrol). Mp 240° (dec.). λ_{max} 227 (log ε 4.65); 260 (log ε 4.51); 286 (sh) (log ε 3.8); 387 (log ε 3.72) (MeOH).

6-Methoxy-6,7-Dimethoxy-3-methyl-1H-carbazole-1,4(9H)-dione. **Koeniginequinone B**

[211183-78-7]

C₁₅H₁₃NO₄ 271.272


Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree). Reddish-brown needles (C₆H₆/petrol). Mp 256° (dec.). λ_{max} 225 (log ε 4.61); 263 (log ε 4.39); 293 (sh) (log ε 3.21); 464 (log ε 3.09) (MeOH).

Saha, C. *et al.*, *Phytochemistry*, 1998, **48**, 363-366 (*isol, synth, uv, ir, pmr*)

Knölker, H.-J. *et al.*, *Heterocycles*, 2003, **60**, 1049-1052 (*synth*)

Kohatine

[92664-90-9]

C₃₄H₃₂N₂O₆ 564.637

Redrawn here to correspond with other related alkaloids, e.g. Isotrilobine, I-339, with interchange of the primed and unprimed locants. The names for derivs. are as given in the lit. using the alternative scheme. Alkaloid from the stems of *Cocculus pendulus* (Menispermaceae). [α]_D²⁵ +183 (c, 0.2 in MeOH).

O¹²-Me: **12'-O-Methylkohatine**. *Kohatamine*

[109269-68-3]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +99.4 (CHCl₃).

1,2-Didehydro: **Dehydrokohatine**

[109269-72-9]

C₃₄H₃₀N₂O₆ 562.621

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +53 (CHCl₃).

1,2-Didehydro, N^{2'}-β-oxide: **1',2'-Dehydrokohatine 2β-N-oxide**

C₃₄H₃₀N₂O₇ 578.62

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +81.2 (MeOH).

1,2-Didehydro, O¹²-Me: **12'-O-Methyldehydrokohatine**. *1',2'-Didehydrokohatamine*

[109269-71-8]

C₃₅H₃₂N₂O₆ 576.648

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +97 (CHCl₃).

1,2-Didehydro, di-Me ether, N^{2'}-de-Me: **O,O-Dimethyl-1',2'-dehydrokohatine**

C₃₅H₃₂N₂O₆ 576.648

Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D +56.6 (MeOH).

1,2,3,4-Tetrahydro: Siddiquine


[109269-70-7]

C₃₄H₂₈N₂O₆ 560.605Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D²¹ +172 (CHCl₃).**1,2,3,4-Tetrahydro, O¹²-Me: Siddiquamine**


[109269-69-4]

C₃₅H₃₀N₂O₆ 574.632Alkaloid from *Cocculus pendulus* (Menispermaceae). [α]_D²¹ +113 (CHCl₃).Hussain, S.F. *et al.*, *Tetrahedron*, 1984, **40**, 2513 (*isol, uv, pmr, ms, cd, struct*)Atta-ur-Rahman, *et al.*, *Pure Appl. Chem.*, 1986, **58**, 663 (*derivs*)Guinaudeau, H. *et al.*, *Phytochemistry*, 1987, **26**, 829 (*Kohatamine, Dehydrokohatamine, Dehydrokohatine, Siddiquine, Siddiquamine*)**Kokusaginoline****K-57**C₁₇H₁₃NO₅ 311.293Struct. unknown. Alkaloid from the root bark of *Orixa japonica* (Rutaceae). Cryst. + 1½H₂O. Mp 283°.*Mono-Ac*: Mp 172.5°.Terasaka, M. *et al.*, *Yakugaku Zasshi*, 1933, **53**, 1046-1056; *CA*, **29**, 7337**Komarine****K-58**


[335373-40-5]




Absolute Configuration

C₂₀H₁₇N₃ 299.374Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (Me₂CO/EtOH). Mp 224-225°. λ_{max} 210 (log ε 4.41); 252 (log ε 4.16); 306 (log ε 4.08); 370 (log ε 3.8) (EtOH).Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 76-78 (*isol, pmr*)**Komaroine****K-59***1-(3-Propylphenyl)-9H-pyrido[3,4-b]indole, 9CI. 1-(3-Propylphenyl)-β-carboline* [94898-73-4]C₂₀H₁₈N₂ 286.376Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (CH₂Cl₂/hexane). Mp 144-145°. λ_{max} 215 (log ε 4.44); 235 (log ε 4.42); 280 (log ε 4.12); 290 (log ε 4.12); 350 (log ε 3.81) (EtOH).Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 378-379 (*isol, uv, ir, pmr, ms, struct*)Bracher, F. *et al.*, *Annalen*, 1992, 1315-1319 (*synth, ir, pmr, cmr, ms*)**Komavicine****K-60**


[169626-15-7]

C₂₀H₁₇N₃ 299.374Alkaloid from aerial parts of *Nitraria komarovii*. Amorph.Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin., 1993*, **29**, 33; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 26**Komavine****K-61**



[6716-66-1]

C₁₆H₂₀N₂ 240.347Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (hexane). Mp 131-132° Mp 279-281° (as hydrochloride). λ_{max} 225 (log ε 4.69); 282 (log ε 4.14); 291 (log ε 4.03) (EtOH).*N*²-*Ac*: *N*²-*Acetylkomavine*


[80616-05-3]

C₁₈H₂₂N₂O 282.385Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (2-propanol). Mp 162-163°. λ_{max} 225 (log ε 4.62); 282 (log ε 4.03); 291 (sh) (log ε 3.96) (EtOH).Carrasco, N. *et al.*, *J.O.C.*, 1973, **38**, 4342-4343 (*synth*)Bobowski, G. *et al.*, *J. Het. Chem.*, 1981, **18**, 1179-1187 (*synth, uv, pmr*)Rodriguez, J.G. *et al.*, *J. Appl. Chem.*, 1983, **30**, 373-378 (*synth, pmr*)Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 61-64 (*isol, pmr, cmr, synth*)**Konbamidin****K-62***N-(Carboxymethyl)tryptophan, 9CI**(R)*-formC₁₃H₁₄N₂O₄ 262.265*(R)*-form*d*-form

[160955-40-8]

Alkaloid from the Okinawan marine sponge *Ircinia* sp. Exhibits cytotoxicity against HeLa cells *in vitro*. Powder. [α]_D²¹ +15 (c, 0.27 in MeOH). λ_{max} 228 (ε 18200); 281 (ε 3310); 290 (ε 2690) (MeOH) (Berdy).*(S)*-form*L*-formSynthetic. Not significantly cytotoxic. [α]_D²¹ -12.6 (c, 1.9 in MeOH).Shinonaga, H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1603 (*isol, uv, ir, pmr, cmr, synth, struct*)**Kongboentine A****K-63**C₂₄H₄₀N₂O₄ 420.591Unusual aminoconitane. Alkaloid from the roots of *Aconitum kongboense*. Amorph. powder. Mp 94-96°.Ping, A. *et al.*, *J. Asian Nat. Prod. Res.*, 2004, **6**, 151-154 (*isol, pmr, cmr*)**Koniamborine****K-64***7-Methoxy-5-methyl-4H-pyrano[3,2-b]indol-4(5H)-one*C₁₃H₁₁NO₃ 229.235Alkaloid from the aerial parts of *Boronella koniambiensis*. Needles (cyclohexane/EtOAc). Mp 142°. λ_{max} 242 (log ε 3.83); 264 (log ε 3.96); 279 (log ε 3.83); 322 (log ε 4.21) (MeOH).*2,3-Dihydro: 2,3-Dihydrokoniamborine*C₁₃H₁₃NO₃ 231.251Alkaloid from *Boronella koniambiensis*. Amorph. solid.Grougnet, R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1083-1086 (*isol, pmr, cmr, ms*)Clawson, R.W. *et al.*, *Tet. Lett.*, 2007, **48**, 6019-6021 (*synth*)**Kopsanol****K-65***Kopsan-22-ol, 9CI*

[6582-68-9]



Absolute Configuration

C₂₀H₂₄N₂O 308.422Alkaloid from *Aspidosperma duckei*, *Aspidosperma cuspa*, *Aspidosperma macrocarpon* and *Aspidosperma verbascifolium* (Apocynaceae). Needles (CHCl₃ or MeOH). Mp 239-245° dec. [α]_D²⁷ +16.7 (c, 0.6 in CHCl₃). pK_a 8.2 (66% DMF).*N*³-*Formyl*: *N*³-*Formylkopsanol*

[24382-01-2]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from *Aspidosperma verbascifolium* (Apocynaceae).

O-Ac:

Cryst. (Me₂CO/hexane or MeOH). Mp 202-205°. [α]_D²⁶ +37 (c, 2.97 in CHCl₃).

22-Ketone: **Kopsanone**. *Kopsan-22-one*, 9CI

[6662-83-5]

C₂₀H₂₂N₂O 306.407

Alkaloid from *Aspidosperma duckei*, *Aspidosperma cuspa*, *Aspidosperma macrocarpon*, *Aspidosperma verbascifolium* and *Vinca erecta* (Apocynaceae). Cryst. (MeOH). Mp 161-163°. [α]_D²⁵ -21.3 (c, 3.47 in CHCl₃). pK_a 6.1 (67% DMF aq.).

22-Ketone; methiodide:

Cryst. (MeOH aq.). Mp 287-289° dec.

22-Ketone, N-Ac:

Cryst. (EtOAc). Mp 184-185°.

22-Epimer: **22-Epikopsanol**

[24992-91-4]

C₂₀H₂₄N₂O 308.422

Alkaloid from *Aspidosperma duckei*, *Aspidosperma cuspa*, *Aspidosperma macrocarpon* and *Aspidosperma verbascifolium* (Apocynaceae). Cryst. (MeOH). Mp 196-197°. [α]_D²⁴ -75.5 (c, 0.98 in CHCl₃). pK_a 7.45 (66% DMF).

22-Epimer, N^a,O-di-Ac:

Cryst. (EtOAc). Mp 231-235° dec. [α]_D²² +20.1 (c, 2.08 in CHCl₃).

Ferreira Filho, J.M. et al., *J.C.S. (C)*, 1966, 1260-1266 (*isol*, *Kopsanone*, *Epikopsanol*, *uv. ord*, *ir*, *pmr*, *ms*, *struct*)

Craven, B.M. et al., *Chem. Comm.*, 1968, 955-956 (*Kopsanone*, *abs config*, *cryst struct*)

Brackman, J.C. et al., *Bull. Soc. Chim. Belg.*, 1969, **78**, 63-68 (*Aspidosperma verbascifolium* *constits*)

Simões, J.C. et al., *Phytochemistry*, 1976, **15**, 543-544 (*Aspidosperma cuspa* *constits*)


Magnus, P. et al., *J.A.C.S.*, 1984, **106**, 2105-2114 (*Kopsanone*, *synth*, *pmr*)

Pihko, A.J. et al., *Tetrahedron*, 2005, **61**, 8769-8807 (*synth*, *rev*)

Kopsaporine

K-66

Demethoxykopsingine
[18555-88-9]



Absolute
Configuration

C₂₃H₂₆N₂O₆ 426.468

CAS sometimes gives the opposite incorrect abs. configs. Alkaloid from the leaves of *Kopsia singapurensis* and stem bark of *Kopsia teoi*. Needles (EtOH). Mp 234° dec. [α]_D²⁷ +48. pK_a 5.63. λ_{max} 207 (log ε 4.53); 246 (log ε 4.15); 279 (log ε 3.43); 287 (log ε 3.39) (MeOH).

N-*De*(methoxycarbonyl): **Kopsinol**. N-*Decarbomethoxykopsaporine*
[149355-62-4]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from stem bark of *Kopsia*

teoi. [α]_D +82 (c, 0.022 in CHCl₃).

12-Hydroxy: **Kopsingarine**. 17-Hydroxy-kopsaporine

[1358-67-4]

C₂₃H₂₆N₂O₇ 442.468

Alkaloid from the leaves of *Kopsia singapurensis*. Needles (EtOH aq.). Mp 230° dec. [α]_D +14. pK_a 5.83. Struct. proposed on comparative IR and UV data with Kopsingine and Kopsaporine. λ_{max} 247 (log ε 4.14); 280 (log ε 3.36); 288 (log ε 3.32) (no solvent reported).

12-Hydroxy, 14α,15α-epoxide: **Kopsimaline B**

C₂₃H₂₆N₂O₈ 458.467

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D -5 (c, 0.53 in CHCl₃). λ_{max} 217 (log ε 4.38); 244 (log ε 3.97); 284 (log ε 3.59) (EtOH).

11-Methoxy, 12-hydroxy, 14α,15α-epoxide: **Kopsimaline C**

C₂₄H₂₈N₂O₉ 488.493

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D -27 (c, 0.09 in CHCl₃). λ_{max} 220 (log ε 4.53); 252 (log ε 3.89); 283 (log ε 3.42) (EtOH).

12-Methoxy: **Kopsingine**

[1358-68-5]

C₂₄H₂₈N₂O₇ 456.494

Alkaloid from *Kopsia singapurensis* leaves and *Kopsia teoi*. Prisms (CHCl₃/EtOH). Mp 270-274° dec. [α]_D²⁹ +75. pK_a 5.27. λ_{max} 217 (log ε 4.56); 253 (log ε 4.04); 282 (log ε 3.38); 288 (log ε 3.36) (MeOH).

12-Methoxy, 14α,15α-epoxide: **Kopsimaline E**. 14,15-Epoxykopsingine

[199981-81-2]

C₂₄H₂₈N₂O₈ 472.494

Alkaloid from *Kopsia teoi* and *Kopsia singapurensis*. [α]_D +15 (c, 0.3 in CHCl₃). Stereochem. revised in 2008. λ_{max} 217 (log ε 4.34); 254 (log ε 3.83); 283 (log ε 3.15); 289 (log ε 3.14) (EtOH).

12-Methoxy, 14,15-dihydro:

Prisms (CHCl₃/MeOH). Mp 268-270° dec.

12-Methoxy, 14,15-dihydro, 15α-hydroxy: **Kopsinganol**

[149355-63-5]

C₂₄H₃₀N₂O₈ 474.51

Trace alkaloid from stem bark of *Kopsia teoi*. [α]_D +39 (c, 1.80 in CHCl₃). CAS gives the incorrect opposite abs. config. λ_{max} 217 (log ε 4.56); 254 (log ε 4.07); 282 (log ε 3.38); 290 (log ε 3.35) (EtOH).

12-Methoxy, 17-deoxy: **Kopsinginine**. 17-*Dehydroxykopsingine*

[149355-59-9]

C₂₄H₂₈N₂O₆ 440.495

Alkaloid from stem bark of *Kopsia teoi*. λ_{max} 217 (log ε 4.44); 254 (log ε 3.97); 280 (log ε 3.34); 289 (log ε 3.32) (EtOH).

12-Methoxy, 11-hydroxy: **11-Hydroxykopsingine**

[199981-78-7]

C₂₄H₂₈N₂O₈ 472.494

Alkaloid from *Kopsia teoi*. [α]_D +28 (c,

0.3 in CHCl₃). λ_{max} 220 (log ε 4.43); 252 (log ε 3.87); 289 (log ε 3.27) (EtOH).

11,12-Dimethoxy: **11-Methoxykopsingine**

[199981-79-8]

C₂₅H₃₀N₂O₈ 486.521

Alkaloid from *Kopsia teoi*. [α]_D -24 (c, 0.2 in CHCl₃). λ_{max} 220 (log ε 4.37); 251 (log ε 3.77); 280 (log ε 3.08); 286 (log ε 3.05) (EtOH).

11,12-Methylenedioxy: **11,12-Methylenedioxykopsaporine**

[152243-73-7]

C₂₄H₂₆N₂O₈ 470.478

Alkaloid from *Kopsia teoi* and *Kopsia singapurensis*. Amorph. powder. [α]_D²⁰ +21 (c, 0.8 in CHCl₃).

11,12-Methylenedioxy, N-*de*(methoxycarbonyl): 11,12-Methylenedioxykopsinol. **Kopsincine**

C₂₂H₂₄N₂O₆ 412.441

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D +22 (c, 0.38 in CHCl₃). λ_{max} 219 (log ε 4.27); 246 (log ε 3.69); 279 (log ε 3.07) (EtOH).

11,12-Methylenedioxy, 14α,15α-epoxide: **Kopsimaline A**

C₂₄H₂₆N₂O₉ 486.477

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D +159 (c, 0.13 in CHCl₃). λ_{max} 225 (log ε 4.43); 255 (log ε 3.97); 286 (log ε 3.24) (EtOH).

11,12-Methylenedioxy, 14,15-dihydro, 17-ketone, N-*de*(methoxycarbonyl): **Kopsosfinone**

C₂₂H₂₄N₂O₆ 412.441

Alkaloid from *Kopsia singapurensis*. Amorph. solid. [α]_D +37 (c, 0.27 in CHCl₃). λ_{max} 218 (log ε 4.23); 244 (log ε 3.73); 284 (log ε 3.01) (EtOH).

11,12-Methylenedioxy, 19β-hydroxy, 14α,15α-epoxide: **Kopsimaline D**

C₂₄H₂₆N₂O₁₀ 502.477

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D +83 (c, 0.36 in CHCl₃). λ_{max} 225 (log ε 4.15); 245 (log ε 3.72); 256 (log ε 3.62); 286 (log ε 3.07) (EtOH).

17-Epimer: **Kopsilosine A**. 17-*Epikopsaporine*

[958247-15-9]

C₂₃H₂₆N₂O₆ 426.468

Alkaloid from *Kopsia singapurensis*. Pale yellow oil. [α]_D²⁵ +16 (c, 0.25 in CHCl₃). λ_{max} 206 (log ε 4.31); 244 (log ε 3.96); 278 (log ε 3.28); 285 (log ε 3.23) (EtOH).

17-Epimer, N-*de*(methoxycarbonyl): 17-*Epikopsinol*. **Kopsilosine H**

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from *Kopsia singapurensis*. Yellowish oil. [α]_D -13 (c, 0.17 in CHCl₃). λ_{max} 203 (log ε 3.59); 243 (log ε 3.01); 293 (log ε 2.62) (EtOH).

17-Epimer, 14,15-dihydro: **Kopsilosine B**

[958247-16-0]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from *Kopsia singapurensis*. Oil. [α]_D²⁵ +50 (c, 0.23 in CHCl₃). λ_{max} 204 (log ε 4.32); 244 (log ε 3.96); 281 (log ε 3.27); 287 (log ε 3.23) (EtOH).

17-Epimer, 14,15-dihydro, N-*de*(methoxycarbonyl): **Kopsilosine I**

$C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from *Kopsia singapurensis*.
Oil. $[\alpha]_D^{25}$ -29 (c, 0.09 in $CHCl_3$). λ_{max} 205 (log ϵ 3.9); 242 (log ϵ 3.34); 293 (log ϵ 2.96) (EtOH).

17-Epimer, 14,15-dihydro, 15 α -hydroxy:

Kopsilosine C

[364781-91-9]

$C_{23}H_{28}N_2O_7$ 444.483

Alkaloid from *Kopsia singapurensis*.
Pale yellow cryst. (Et₂O). Mp 172-174°. $[\alpha]_D^{25}$ -40 (c, 0.28 in $CHCl_3$). λ_{max} 206 (log ϵ 3.28); 243 (log ϵ 2.98); 279 (log ϵ 2.02); 288 (log ϵ 1.76) (EtOH).

17-Epimer, 10-methoxy, 14,15-dihydro:

Kopsilosine D

[958247-17-1]

$C_{24}H_{30}N_2O_7$ 458.51

Alkaloid from *Kopsia singapurensis*.
Oil. $[\alpha]_D^{25}$ -63 (c, 0.1 in $CHCl_3$). λ_{max} 201 (log ϵ 4.59); 249 (log ϵ 4.27); 297 (log ϵ 3.67) (EtOH).

17-Epimer, 10-methoxy, 15 α -hydroxy,

14,15-dihydro: **Kopsilosine E**

[958247-18-2]

$C_{24}H_{30}N_2O_8$ 474.51

Alkaloid from *Kopsia singapurensis*.
Oil. $[\alpha]_D^{25}$ -25 (c, 0.15 in $CHCl_3$). λ_{max} 202 (log ϵ 4.46); 249 (log ϵ 4.14); 298 (log ϵ 3.51) (EtOH).

17-Epimer, 12-methoxy, 15 α -hydroxy,

14,15-dihydro: **Kopsilosine F**

[958247-19-3]

$C_{24}H_{30}N_2O_8$ 474.51

Alkaloid from *Kopsia singapurensis*.
Pale yellow oil. $[\alpha]_D^{25}$ +9 (c, 0.28 in $CHCl_3$). λ_{max} 216 (log ϵ 3.57); 252 (log ϵ 3.1); 283 (log ϵ 2.55) (EtOH).

Kiang, A.K. *et al.*, *CA*, 1959, **53**, 14131 (*isol. Kopsingine, Kopsingarine*)

Thomas, D.W. *et al.*, *J.A.C.S.*, 1967, **89**, 3235-3242 (*Kopsaporine, Kopsingine, uv, ir, pmr, ms, struct, Kopsingarine*)

Kam, T.-S. *et al.*, *Phytochemistry*, 1993, **32**, 1343-1345; 1996, **42**, 539-541 (*Kopsingine, cryst struct, Kopsinginine, Kopsinol, Kopsingamol*)


Kam, T.-S. *et al.*, *Phytochemistry*, 1997, **46**, 789-792 (*Hydroxykopsingine, Methoxykopsingine, Epoxykopsingine*)

Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1783-1789 (*Kopsilosines A-F*)

Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 53-57 (*Kopsimalines A-E, Kopsifoline, Kopsinicine, Kopsilosines H,I*)

Kopsidasine

[85559-71-3]



Absolute Configuration

$C_{24}H_{28}N_2O_6$ 440.495

Alkaloid from the leaves of *Kopsia dasyrachis* (Apocynaceae). Gum. $[\alpha]_D^{25}$ -133 (c, 3.58 in $CHCl_3$).

N⁴-Oxide: Kopsidasine N⁴-oxide

[85559-72-4]

$C_{24}H_{28}N_2O_7$ 456.494


Alkaloid from leaves of *Kopsia dasyrachis* (Apocynaceae). Cryst. (MeOH/Et₂O). $[\alpha]_D^{23}$ -159.5 (c, 1.013 in $CHCl_3$).

Hombberger, K. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2548-2557 (*isol, uv, ir, pmr, cmr, ms, struct*)

Magnus, P. *et al.*, *Tetrahedron*, 2002, **58**, 3423-3443 (*synth*)

Kopsidasinine

[85559-78-0]



Absolute Configuration

$C_{24}H_{28}N_2O_6$ 440.495

Alkaloid from the leaves of *Kopsia dasyrachis* (Apocynaceae). Gum. $[\alpha]_D^{23}$ -139.1 (c, 1.307 in $CHCl_3$).

Demethoxy: 10-Demethoxykopsidasinine

[117259-13-9]

$C_{23}H_{26}N_2O_5$ 410.469

Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Prisms. Mp 194-196°. $[\alpha]_D^{20}$ -159 (c, 1.00 in MeOH).

Demethoxy, 12-methoxy: 10-Demethoxy-12-methoxykopsidasinine

[184870-16-4]

$C_{24}H_{28}N_2O_6$ 440.495

Alkaloid from stems of *Kopsia pauciflora*. $[\alpha]_D$ -115 (c, 0.114 in $CHCl_3$). λ_{max} 216 (log ϵ 4.3); 249 (log ϵ 3.82); 280 (log ϵ 2.98); 287 (log ϵ 3.04) (EtOH).


Hombberger, K. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2548 (*isol, uv, ir, pmr, cmr, ms, struct*)

Hamburger, M.O. *et al.*, *Phytochemistry*, 1988, **27**, 2719 (*10-Demethoxykopsidasinine*)

Kam, T.-S. *et al.*, *Phytochemistry*, 1996, **43**, 1385 (*10-Demethoxy-12-methoxykopsidasinine*)

Kopsifine

[204514-78-3]



Absolute Configuration

R, R' = -CH₂-

$C_{23}H_{22}N_2O_7$ 438.436

Alkaloid from *Kopsia dasyrachis*. $[\alpha]_D$ +97 (c, 0.04 in $CHCl_3$). λ_{max} 223 (log ϵ 4.41); 250 (log ϵ 3.94); 285 (log ϵ 3.16); 295 (log ϵ 3.1) (EtOH).

De(methoxycarbonyl): Decarbomethoxy-

kopsifine. Demethoxycarbonylkopsifine

[237078-12-5]

$C_{21}H_{20}N_2O_5$ 380.399

Alkaloid from *Kopsia dasyrachis*. $[\alpha]_D$ +52 (c, 0.07 in $CHCl_3$). λ_{max} 220 (log ϵ 4.72); 243 (log ϵ 4.23); 288 (log ϵ 3.51) (EtOH).

5-Deoxo: 11,12-Methylenedioxykopsine

[237078-14-7]


$C_{23}H_{24}N_2O_6$ 424.452

Alkaloid from *Kopsia dasyrachis*. $[\alpha]_D$ -13 (c, 0.18 in $CHCl_3$). λ_{max} 225 (log ϵ 4.61); 244 (log ϵ 4.2); 287 (log ϵ 3.41); 294 (log ϵ 3.39) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 159-169 (*isol, uv, pmr, cmr, ms*)

Kopsifoline C

[603131-70-0]



Relative Configuration

$C_{22}H_{28}N_2O_6$ 416.473

Alkaloid from *Kopsia fruticosa*. Oil. $[\alpha]_D$ +87 (c, 0.08 in $CHCl_3$). λ_{max} 214 (log ϵ 4.35); 248 (log ϵ 3.72); 294 (log ϵ 3.33) (EtOH).

14-Deoxy: Kopsifoline B

[603131-69-7]

$C_{22}H_{28}N_2O_5$ 400.474

Alkaloid from *Kopsia fruticosa*. Oil. $[\alpha]_D$ -46 (c, 0.66 in $CHCl_3$). λ_{max} 213 (log ϵ 4); 248 (log ϵ 3.88); 295 (log ϵ 3.44) (EtOH).

14,15-Dideoxy, 14,15-didehydro: Kopsifoline A

[603131-68-6]

$C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from *Kopsia fruticosa*. Oil. $[\alpha]_D$ -11 (c, 0.43 in $CHCl_3$). λ_{max} 213 (log ϵ 4.26); 248 (log ϵ 3.93); 296 (log ϵ 3.51) (EtOH).

2,14,15-Trideoxy, 1,2,14,15-tetradehydro: Kopsifoline E

[603131-67-5]

$C_{22}H_{24}N_2O_3$ 364.443

Alkaloid from *Kopsia fruticosa*. Light yellow oil. $[\alpha]_D$ +84 (c, 0.15 in $CHCl_3$). λ_{max} 230 (log ϵ 4.05); 253 (log ϵ 3.63); 309 (log ϵ 3.56) (EtOH).

Demethoxy, 2,14,15-trideoxy, 1,2,14,15-tetradehydro: Kopsifoline D

[603131-66-4]

$C_{21}H_{22}N_2O_2$ 334.417

Alkaloid from *Kopsia fruticosa*. Light yellow oil. $[\alpha]_D$ -27 (c, 0.09 in $CHCl_3$). λ_{max} 224 (log ϵ 4.15); 249 (log ϵ 3.68); 280 (log ϵ 3.56) (EtOH).

Demethoxy, 11-methoxy, 2,14,15-trideoxy, 14,15-didehydro: Kopsifoline F

[603131-65-3]

$C_{22}H_{26}N_2O_3$ 366.459


Alkaloid from *Kopsia fruticosa*. Oil. $[\alpha]_D$ -112 (c, 0.07 in $CHCl_3$). λ_{max} 212 (log ϵ 4.56); 248 (log ϵ 3.99); 295 (log ϵ 3.57) (EtOH).

Kam, T.-S. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 991-998 (*Kopsifolines, isol, pmr, cmr, ms*)

Kopsijasmine

K-71

*N*¹-Methoxycarbonyl-16,17-didehydrokopsinine. 10-Demethoxy-21-deoxykopsidasine
[114639-86-0]



$C_{23}H_{26}N_2O_4$ 394.469
Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Mp 199-202°. $[\alpha]_D$ -202 (CHCl₃). λ_{max} 204 (ε 16700); 239 (ε 7400); 277 (ε 1600); 286 (ε 1300) (EtOH) (Derep).

N-De(methoxycarbonyl): **Kopsijasminine**. *Δ*^{16,17}-Kopsinine. 16,17-Didehydrokopsinine
[947251-86-7]
 $C_{21}H_{24}N_2O_2$ 336.433
Alkaloid from the stem bark of *Kopsia teoi*. Light yellow oil. $[\alpha]_D$ -60 (c, 0.05 in CHCl₃). λ_{max} 206 (log ε 4.36); 243 (log ε 3.77); 292 (log ε 3.3) (EtOH).

12-Hydroxy: **12-Hydroxy-N-methoxycarbonyl-*Δ*^{16,17}-kopsinine**
[164661-14-7]
 $C_{23}H_{26}N_2O_5$ 410.469
Alkaloid from leaves and stems of *Kopsia profunda* (Apocynaceae). $[\alpha]_D$ -121 (c, 0.04 in CHCl₃).

12-Methoxy: **12-Methoxy-N-methoxycarbonyl-*Δ*^{16,17}-kopsinine**
[129743-93-7]
 $C_{24}H_{28}N_2O_5$ 424.496
Alkaloid from the stems and leaves of *Kopsia profunda* (Apocynaceae). Needles (EtOH). Mp 150-152°. $[\alpha]_D$ -75 (c, 0.123 in CHCl₃).

12-Methoxy, N⁴-oxide: **12-Methoxy-N-methoxycarbonyl-*Δ*^{16,17}-kopsinine N⁴-oxide**
[164661-16-9]
 $C_{24}H_{28}N_2O_6$ 440.495
Alkaloid from stems and leaves of *Kopsia profunda* (Apocynaceae). Mp 244-245°. $[\alpha]_D$ -163 (c, 0.15 in CHCl₃).

11,12-Methylenedioxy: **N-Methoxycarbonyl-11,12-methylenedioxy-*Δ*^{16,17}-kopsinine**
[129743-92-6]
 $C_{24}H_{26}N_2O_6$ 438.479
Alkaloid from the stems and leaves of *Kopsia profunda* (Apocynaceae). Prisms (MeOH). Mp 198-200°. $[\alpha]_D$ -180 (c, 0.17 in CHCl₃).


11,12-Methylenedioxy, N⁴-oxide: **N-Methoxycarbonyl-11,12-methylenedioxy-*Δ*^{16,17}-kopsinine N-oxide**
[164661-15-8]
 $C_{24}H_{26}N_2O_7$ 454.479
Alkaloid from leaves and stems of *Kopsia profunda* (Apocynaceae). $[\alpha]_D$

-125 (c, 0.06 in CHCl₃).
Ruangrunsi, N. *et al.*, *Tet. Lett.*, 1987, **28**, 3679 (pmr, cmr, ms, struct)
Magnus, P. *et al.*, *J.O.C.*, 1988, **53**, 5772 (synth, uv, ir, pmr, ms)
Kam, T.-S. *et al.*, *Phytochemistry*, 1990, **29**, 2321; 1995, **39**, 469 (derivs)
Lim, S.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1380-1383 (Kopsijasminine)

Kopsijasminilam

K-72

[114639-87-1]



$C_{23}H_{26}N_2O_6$ 426.468
New skeletal group of *Kopsia* alkaloids. Can be regarded as 20,21-secoaspido-fractines. Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Mp 245-246°. $[\alpha]_D$ -220 (CHCl₃). λ_{max} 208 (ε 41700); 245 (ε 17800); 280 (ε 2950); 290 (ε 2690) (MeOH) (Derep).

20-Deoxy: **Deoxykopsijasminilam**
[114622-12-7]
 $C_{23}H_{26}N_2O_5$ 410.469
Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Mp 192-195°. $[\alpha]_D$ -210 (CHCl₃). λ_{max} 208 (ε 41700); 245 (ε 17800); 280 (ε 2950); 290 (ε 2690) (MeOH) (Derep).


14,15-Didehydro: ***Δ*¹⁴-Kopsijasminilam**
[114622-13-8]
 $C_{23}H_{24}N_2O_6$ 424.452
Alkaloid from the leaves of *Kopsia jasminiflora* (Apocynaceae). Amorph. powder.

Ruangrunsi, N. *et al.*, *Tet. Lett.*, 1987, **28**, 3679 (uv, ir, pmr, cmr, ms, cryst struct, derivs)
Kuehne, M.E. *et al.*, *J.O.C.*, 2000, **65**, 6434-6440 (synth)
Magnus, P. *et al.*, *Tetrahedron*, 2002, **58**, 3423-3443 (synth)

Kopsilactone

K-73

[180841-82-1]




$C_{13}H_{21}NO_2$ 223.314
Minor alkaloid from leaves of *Kopsia macrophylla*. Amorph. $[\alpha]_D^{25}$ +43 (c, 0.8 in CHCl₃).

Kan-Fan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 283 (isol, ir, pmr, cmr, ms, struct)

Relative
Configuration**Kopsinaline**

K-74

Methyl 3-hydroxyaspidofractinine-3-carboxylate, 9CI
[88607-44-7]

Absolute
Configuration

$C_{21}H_{26}N_2O_3$ 354.448
Kopsinaline itself is currently unknown.

N-Methoxycarbonyl: **Kopsiflorine**. N-Methoxycarbonylkopsinaline
[1358-64-1]
 $C_{23}H_{28}N_2O_5$ 412.485
Alkaloid from the leaves of *Kopsia longiflora* and *Kopsia dasyrachis* (Apocynaceae). Shows multidrug resistance (MDR) reversing effect. Needles (petrol). Mp 144-145°. $[\alpha]_D^{23}$ -66.9 (c, 1.41 in CHCl₃). pK_a 6.38 (70% MeOH).

N-Methoxycarbonyl, N⁴-oxide: **Kopsiflorine N⁴-oxide**
[237078-10-3]
 $C_{23}H_{28}N_2O_6$ 428.484
Alkaloid from *Kopsia dasyrachis*. $[\alpha]_D$ -57 (c, 0.32 in CHCl₃). λ_{max} 206 (log ε 4.38); 242 (log ε 4.04); 278 (log ε 3.23); 284 (log ε 3.2) (EtOH).

11-Methoxy, 12-hydroxy, N-methoxycarbonyl: **N-Methoxycarbonyl-12-hydroxy-11-methoxykopsinaline**
[114495-07-7]
 $C_{24}H_{30}N_2O_7$ 458.51
Alkaloid from the fruits of *Kopsia officinalis*. Yellow cryst. Mp 184°. $[\alpha]_D^{20}$ -104.6 (c, 0.0015M in CHCl₃). λ_{max} 223 (log ε 4.54); 255 (log ε 3.98); 280 (log ε 3.22); 290 (log ε 3.22) (MeOH).

12-Methoxy: **12-Methoxykopsinaline**
[88607-45-8]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from the roots of *Kopsia officinalis* (Apocynaceae). Amorph. $[\alpha]_D$ -7 (c, 0.17 in CHCl₃).

12-Methoxy, N-methoxycarbonyl: **Kopsilongine**. 12-Methoxy-N-methoxycarbonylkopsinaline. N-Methoxycarbonyl-12-methoxykopsinaline
[1358-65-2]
 $C_{24}H_{30}N_2O_6$ 442.511
Alkaloid from the roots of *Kopsia longiflora* and (Kopsilongine) from bark and leaves of *Kopsia officinalis* (Apocynaceae). Cryst. (MeOH or Me₂CO). Mp 210-211° (206°). $[\alpha]_D$ -5 (c, 1.31 in CHCl₃). pK_a 6.6 (70% MeOH). The identity of the two isolates is not certain. Mps are similar. Most data refers to the later isolate, N-Methoxycarbonyl-12-methoxykopsinaline.

12-Methoxy, N-methoxycarbonyl, perchlorate: Plates (H₂O). Mp 246-248° dec. Deriv. of Kopsilongine.

12-Methoxy, N-methoxycarbonyl, N⁴-

oxide: Kopsilongine N⁴-oxide

[958247-20-6]

C₂₄H₃₀N₂O₇ 458.51Alkaloid from *Kopsia singapurensis*.Oil. [α]_D²⁵ -29 (c, 0.21 in CHCl₃). λ_{max} 216 (log ε 4.15); 252 (log ε 3.58); 287 (log ε 2.51) (EtOH).**12-Methoxy, 11-hydroxy, N-methoxycarbonyl: N-Methoxycarbonyl-11-hydroxy-12-methoxykopsinaline**

[127608-79-1]

C₂₄H₃₀N₂O₇ 458.51Alkaloid from the fruits of *Kopsia officinalis*. Yellow cryst. Mp 207-209°. λ_{max} 203 (log ε 4.45); 253 (log ε 3.98); 293 (log ε 3.43) (MeOH).**11,12-Dimethoxy, N-methoxycarbonyl: 11,12-Dimethoxy-N-methoxycarbonylkopsinaline. N-Methoxycarbonyl-11,12-dimethoxykopsinaline. 11-Methoxykopsilongine**

[88607-47-0]

C₂₅H₃₂N₂O₇ 472.537Alkaloid from the roots of *Kopsia officinalis* (Apocynaceae). Amorph.[α]_D -4 (c, 0.68 in CHCl₃).**11,12-Dimethoxy, N-methoxycarbonyl, N⁴-oxide: 11-Methoxykopsilongine N⁴-oxide**

[237078-11-4]

C₂₅H₃₂N₂O₈ 488.536Alkaloid from *Kopsia dasyrachis*. [α]_D -11 (c, 0.13 in CHCl₃). λ_{max} 223 (log ε 4.44); 252 (log ε 3.86); 282 (log ε 3.12); 289 (log ε 3.12) (EtOH).**11,12-Methylenedioxy: 11,12-Methylenedioxykopsinaline**

[88607-48-1]

C₂₂H₂₆N₂O₅ 398.458Alkaloid from the roots of *Kopsia officinalis* and from *Kopsia dasyrachis* (Apocynaceae). Amorph. [α]_D -12 (c, 0.70 in CHCl₃).**11,12-Methylenedioxy, N⁴-oxide: 11,12-Methylenedioxykopsinaline N⁴-oxide**

[221149-28-6]

C₂₂H₂₆N₂O₆ 414.457Alkaloid from *Kopsia griffithii*. [α]_D -16 (c, 0.19 in CHCl₃). λ_{max} 221 (log ε 4.35); 246 (log ε 3.77); 282 (log ε 2.97) (EtOH).**11,12-Methylenedioxy, N-methoxycarbonyl: Kopsamine. N-Methoxycarbonyl-11,12-methylenedioxykopsinaline**

[1358-62-9]

C₂₄H₂₈N₂O₇ 456.494Alkaloid from *Kopsia officinalis*, *Kopsia longiflora* and *Kopsia flavida* (prob. the same as *Kopsia pruniformis*) (Apocynaceae). Prisms (MeOH or Me₂CO). Mp 208°. [α]_D -48.2 (c, 1.0 in CHCl₃) (-42).**11,12-Methylenedioxy, N-methoxycarbonyl, N⁴-oxide: Kopsamine N⁴-oxide**

[125295-15-0]

C₂₄H₂₈N₂O₈ 472.494Alkaloid from *Kopsia officinalis*. Cryst. Mp 251-252°. [α]_D²⁰ -40.7 (CHCl₃). λ_{max} 226 (log ε 4.5); 247 (log ε 4.06); 255 (log ε 3.96); 285 (log ε 3.21); 295 (log ε 3.21) (MeOH).**11,12-Methylenedioxy, 14,15-didehydro: 14,15-Didehydro-11,12-methylenedioxykopsinaline. Kopsamidine A**

[949932-19-8]

C₂₂H₂₄N₂O₅ 396.442Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D +97 (c, 0.15 in CHCl₃). λ_{max} 220 (log ε 4.46); 245 (log ε 3.88); 281 (log ε 3.11) (EtOH).**11,12-Methylenedioxy, 21-hydroxy, N-methoxycarbonyl: Paucifinine**

[199981-75-4]

C₂₄H₂₈N₂O₈ 472.494Alkaloid from the leaves of *Kopsia pauciflora* (Apocynaceae). [α]_D -91 (c, 0.08 in CHCl₃). λ_{max} 248 (log ε 4.08); 255 (log ε 4.03); 285 (log ε 3.13); 290 (log ε 3.09) (EtOH).**11,12-Methylenedioxy, 21-hydroxy, N-methoxycarbonyl, N⁴-oxide: Paucifinine N-oxide**

[200053-20-9]

C₂₄H₂₈N₂O₉ 488.493Alkaloid from the leaves of *Kopsia pauciflora* (Apocynaceae). [α]_D -36 (c, 0.3 in CHCl₃). λ_{max} 227 (log ε 4.5); 248 (log ε 4.06); 255 (log ε 3.97); 285 (log ε 3.21); 290 (log ε 3.17) (EtOH).Crow, W.D. et al., *Aust. J. Chem.*, 1955, **8**, 129-135; 1962, **15**, 130-138 (*Kopsilongine, Kopsiflorine*)Feng, X.Z. et al., *Planta Med.*, 1983, **48**, 280-282 (*Kopsia officinalis constits*)Wu, J. et al., *Jiegou Huaxue*, 1987, **6**, 47-52 (*11-methoxy-12-hydroxy-N-methoxycarbonyl, cryst struct*)Wu, J. et al., *C.A.*, 1988, **108**, 204851m(*Kopsamine, cryst struct*)Zheng, J.-J. et al., *Acta Chim. Sin. (Engl. edn.)*, 1989, 168-175 (*Kopsamine, Kopsamine N-oxide, 12-methoxy-11-hydroxy-N-methoxycarbonyl, 11-methoxy-12-hydroxy-N-methoxycarbonyl*)Kam, T.S. et al., *Phytochemistry*, 1997, **46**, 785-787; 1999, **50**, 75-79; **51**, 159-169(*Paucifinine, Paucifinine N-oxide, 11,12-Methylenedioxykopsinaline N-oxide, Kopsiflorine N-oxide, 11-Methoxykopsilongine N-oxide*)Kam, T.-S. et al., *Bioorg. Med. Chem. Lett.*, 1998, **8**, 2769-2772 (*Kopsiflorine, activity*)Rho, M.-C. et al., *Planta Med.*, 1999, **65**, 307-310 (*Kopsiflorine, activity*)Magnus, P. et al., *Tetrahedron*, 2002, **58**, 3423-3443 (*11-Methoxykopsilongine, synth*)Lim, K.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*Kopsamidine A*)Subramaniam, G. et al., *J. Nat. Prod.*, 2007, **70**, 1783-1789 (*Kopsilongine N⁴-oxide*)**Kopsinarine**


[237078-13-6]

As Kopsifine, K-69 with

R = R' = Me

C₂₄H₂₆N₂O₇ 454.479Alkaloid from *Kopsia dasyrachis*. [α]_D +97 (c, 0.25 in CHCl₃). λ_{max} 219 (log ε 4.66); 240 (log ε 4.15); 280 (log ε 3.32); 288 (log ε 3.3) (EtOH).Kam, T.-S. et al., *Phytochemistry*, 1999, **51**, 159-169 (*isol, uv, pmr, cmr, ms*)**K-75****Kopsine***Methyl 3-hydroxy-22-oxokopsan-1-carboxylate, 9CI*

[559-48-8]



Absolute Configuration

C₂₂H₂₄N₂O₄ 380.443Alkaloid from the leaves of *Kopsia fruticosa* (Apocynaceae). Cryst. (MeOH). Mp 217-218° (210-213°). [α]_D -17.5 (c, 2.15 in CHCl₃). pK_{a1} 4.28 (2-Methoxyethanol aq.). pK_{a1} 4.45 (2-Methoxyethanol).

Perchlorate: Mp 284° dec.


Picrate: Mp 230° dec.

N-De(methoxycarbonyl): DecarbomethoxykopsineC₂₀H₂₂N₂O₂ 322.406Alkaloid from the leaves of *Kopsia fruticosa* (Apocynaceae). Cryst. (Me₂CO aq. or MeOH aq.). Mp 101-127°.**11,12-Dimethoxy: 11,12-Dimethoxykopsine. Kopsinidine B**

[949932-22-3]

C₂₄H₂₈N₂O₆ 440.495Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D +113 (c, 0.18 in CHCl₃). λ_{max} 219 (log ε 4.28); 239 (log ε 3.79); 285 (log ε 3.01) (EtOH).Govindachari, T.R. et al., *Helv. Chim. Acta*, 1962, **45**, 1146; 1963, **46**, 572 (*uv, ir, pmr, ms*)Guggisberg, A. et al., *Helv. Chim. Acta*, 1963, **46**, 679; 1969, **52**, 76(*Decarbomethoxykopsine*)Battersby, A.R. et al., *J.C.S.*, 1963, 22 (*isol, pmr, ms*)Magnus, P. et al., *J.A.C.S.*, 1989, **111**, 6707(*synth, ir, pmr*)Watt, W. et al., *Acta Cryst. C*, 1993, **49**, 171(*cryst struct*)Glover, R.P. et al., *Magn. Reson. Chem.*, 2005, **43**, 483-485 (*pmr, cmr, N-15 nmr*)Lim, K.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*Kopsinidine B*)**Kopsinidine A**


[949932-21-2]

**K-77**C₂₁H₂₂N₂O₄ 366.416Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D -15 (c, 0.17 in CHCl₃). λ_{max} 221 (log ε 4.48); 242 (log ε 3.99); 277 (log ε 3.23) (EtOH).Lim, K.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*isol, pmr, cmr*)

Kopsinine

K-78

Methyl aspidofractinine-3-carboxylate,
9CI. *Ervinine*. *Copsinine*. *Erectine*
[559-51-3]



Absolute
Configuration

$C_{21}H_{26}N_2O_2$ 338.449

Alkaloid from *Kopsia longiflora*, *Kopsia hainanensis*, *Pleiocarpa mutica*, *Pleiocarpa tubicina*, *Pleiocarpa pycnantha* var. *tubicina*, *Aspidosperma multiflorum*, *Aspidosperma populifolium*, *Alstonia venenata*, *Vinca erecta*, *Vinca herbacea*, *Hunteria elliptica* and *Hunteria zeylanica* (Apocynaceae). CNS analeptic agent, antinarcotic. Prisms (petrol or Et₂O/hexane). Mp 104-105°. $[\alpha]_D^{25}$ -76.9 (c, 2.09 in CHCl₃). p*K*_a 7.5 (70% MeOH). λ_{max} 204 (log ϵ 4.35); 246 (log ϵ 3.83); 295 (log ϵ 3.45) (EtOH).

Hydrochloride:

Needles + 2H₂O (MeOH/Et₂O). Mp 244-245°.

Hydrochloride (1:2):

Needles (EtOH). Mp 253-254°.

Picrate:

Yellow needles (EtOH). Mp 222-223° (213-214° dec.).

N⁴-Oxide: Kopsinoline

[7290-10-0]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from bark of *Pleiocarpa tubicina* and *Kopsia hainanensis*; and the roots of *Pleiocarpa mutica* (Apocynaceae). Mp 158-160° dec. $[\alpha]_D^{25}$ -70.1 (c, 0.48 in CHCl₃). λ_{max} 205 (log ϵ 4.43); 244 (log ϵ 3.87); 295 (log ϵ 3.47) (EtOH).

N¹-Formyl: Aspidofractine

[2348-67-6]

$C_{22}H_{26}N_2O_3$ 366.459

Alkaloid from *Alstonia refractum* and *Alstonia populifolium* (Apocynaceae). Mp 195-196°. $[\alpha]_D^{24}$ -128 (c, 1.3 in CHCl₃).

N-Methoxycarbonyl: Pleiocarpine

[559-52-4]

$C_{23}H_{28}N_2O_4$ 396.485

Major alkaloid from the roots of *Pleiocarpa mutica*, also isol. from *Pleiocarpa pycnantha*, *Hunteria eburnea* and *Comularia camerunensis* (preferred genus name *Hunteria*) (Apocynaceae). Cryst. (MeOH or pentane/MeOH). Mp 142.5°. $[\alpha]_D^{25}$ -145 (c, 1.1725 in CHCl₃). p*K*_a 6.19 (MCS).

N-Methoxycarbonyl, hydrochloride:

Cryst. (EtOH or EtOH/Et₂O). Mp 234-235° dec. (229-230°).

N-Methoxycarbonyl, N⁴-oxide: Pleiocarpoline

$C_{23}H_{28}N_2O_5$ 412.485

Alkaloid from *Pleiocarpa mutica* and

the bark of *Pleiocarpa tubicina*. Also isol. from *Hunteria eburnea* (Apocynaceae). Prisms (MeOH/Et₂O). Mp 234-235° dec Mp 248-250° dec. $[\alpha]_D^{24}$ -131.6 (c, 0.94 in CHCl₃). CAS no. not found 8-14CI. λ_{max} 208 (log ϵ 4.41); 245 (log ϵ 4.13); 281 (log ϵ 3.33); 287 (log ϵ 3.32) (EtOH).

N¹-Me: Pleiocarpine

[559-50-2]

$C_{22}H_{28}N_2O_2$ 352.475

Minor alkaloid from the roots of *Pleiocarpa mutica*. Also isol. from *Hunteria eburnea* and *Comularia camerunensis* (preferred genus name *Hunteria*) (Apocynaceae). Cryst. (pentane). Mp 135-136°. $[\alpha]_D^{22}$ -124 (c, 1.461 in CHCl₃). p*K*_a 6.94 (2-methoxyethanol).

N¹-Me, picrate:

Cryst. (MeOH). Mp 222-223° dec.

N¹-Me, N⁴-oxide: Pleiocarpoline

[18559-10-9]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from the bark of *Pleiocarpa tubicina*, the roots of *Pleiocarpa mutica* and from *Hunteria eburnea* (Apocynaceae). Prisms (MeOH/Et₂O). Mp 210-211° dec Mp 239-260° dec. $[\alpha]_D^{26}$ -111 (c, 2.575 in CHCl₃). λ_{max} 207 (log ϵ 4.42); 251 (log ϵ 4); 300 (log ϵ 3.53) (EtOH).

Parent acid, N⁴-Me: Kopsinic acid methosalt

[1348-45-4]

$C_{21}H_{27}N_2O_2^{\oplus}$ 339.456

Alkaloid from root bark of *Pleiocarpa tubicina* and roots of *Pleiocarpa mutica* (Apocynaceae). Cryst. (MeOH) (as chloride). Mp 282-295° dec. (chloride). $[\alpha]_D^{25}$ -107 (c, 0.42 in 50% MeOH aq.). λ_{max} 248 (log ϵ 3.85); 295 (log ϵ 3.45) (EtOH).

15 β -Hydroxy: Hydroxykopsinine I. Ld 90

[26619-90-9]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from the stem bark of *Melodinus australis* (Apocynaceae). Prisms (CHCl₃/Et₂O). Mp 190-197°. $[\alpha]_D^{25}$ -62 (c, 1.142 in CHCl₃).

17 β -Hydroxy, N-methoxycarbonyl: N-Carbomethoxy-17-hydroxykopsinine

[116988-03-5]

$C_{23}H_{28}N_2O_5$ 412.485

Alkaloid from the stem bark of *Kopsia deverrei* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ -132 (c, 0.18 in CHCl₃). λ_{max} 245 (log ϵ 4.21); 281 (log ϵ 3.58); 288 (log ϵ 3.55) (EtOH).

19 α -Hydroxy: Hydroxykopsinine II. Ld 91

[26619-91-0]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from *Melodinus australis* (Apocynaceae). Prisms (CHCl₃/Et₂O). Mp 187-192°. $[\alpha]_D^{23}$ -63 (c, 1.037 in CHCl₃).

3-Oxo, hydroxy: 3-Oxohydroxykopsinine

[90341-31-4]

$C_{21}H_{24}N_2O_4$ 368.432

Trace alkaloid from the stem bark and aerial parts of *Melodinus guillauminii*.

$[\alpha]_D$ -14.5 (c, 0.15 in CHCl₃). Full struct. not known. OH group possibly on C-18 or C-19.

5-Oxo: Kopsinilam

[464-68-6]

$C_{21}H_{24}N_2O_3$ 352.432

Alkaloid from *Pleiocarpa mutica*, *Pleiocarpa tubicina*, *Hunteria eburnea* and *Vinca erecta* (Apocynaceae). Prisms (Me₂CO/pentane). Mp 254-254.5°. $[\alpha]_D^{25}$ -13 (c, 2.27 in CHCl₃).

5-Oxo, N¹-Me: Pleiocarpinilam

$C_{22}H_{26}N_2O_3$ 366.459

Minor alkaloid from *Pleiocarpa mutica*, *Pleiocarpa tubicina* and *Hunteria eburnea* (Apocynaceae). Cryst. (Et₂O or Me₂CO/pentane). Mp 249-250°. $[\alpha]_D^{25}$ -53.4 (c, 1.385 in CHCl₃). CAS no. not found 8-14CI. λ_{max} 253 (log ϵ 4.01); 300 (log ϵ 3.49) (EtOH).

5-Oxo, 3,N-dehydro: Kopsinilamine

[36101-53-8]

$C_{21}H_{23}N_2O_3^{\oplus}$ 351.424

Alkaloid from *Vinca erecta* (Apocynaceae). Mp 243-244°. $[\alpha]_D$ -65.5 (0.76 in Me₂CO). Reduction gave Kopsinilam. Stereochemistry not explicit. λ_{max} 246 (log ϵ 3.75); 296 (log ϵ 3.34) (EtOH).

5,6-Dioxo, N-methoxycarbonyl: 5,6-Dioxopleiocarpine. 10,11-Dioxopleiocarpine

$C_{23}H_{24}N_2O_6$ 424.452

Alkaloid from the stem bark of *Pleiocarpa mutica* (Apocynaceae). Yellow cryst. Mp 285-287°. $[\alpha]_D^{17}$ -265 (CHCl₃). CAS no. not found 8-14CI.

12-Methoxy, N-formyl: Refractine. Vello-sine

[2517-52-4]

$C_{23}H_{28}N_2O_4$ 396.485

Alkaloid from *Aspidosperma refractum* and *Aspidosperma populifolium* (Apocynaceae). Mp 157.5-159° Mp 191-192° (dimorph.). λ_{max} 216 (log ϵ 4.43); 260 (log ϵ 4.12); 287 (sh) (log ϵ 3.44) (EtOH).

12-Methoxy, N-Ac: 6-Demethoxyppyrrifoline

[88153-40-6]

$C_{24}H_{30}N_2O_4$ 410.512

Alkaloid from leaves of *Aspidosperma pyriformis* (Apocynaceae). Cryst. (petrol). Mp 121-123°. λ_{max} 216 (log ϵ 4.34); 259 (log ϵ 3.97); 290 (sh) (log ϵ 3.6) (MeOH).

12-Methoxy, N-methoxycarbonyl: 12-Methoxypleiocarpine

[202195-91-3]

$C_{24}H_{30}N_2O_5$ 426.511

Alkaloid from the leaves of *Kopsia griffithii* (Apocynaceae). $[\alpha]_D$ -79 (c, 0.1 in CHCl₃). λ_{max} 216 (log ϵ 4.23); 253 (log ϵ 3.78); 282 (log ϵ 3.17); 288 (log ϵ 3.2) (EtOH).

15 α -Methoxy: Kopsamidine B. 15-Methoxykopsinine

[949932-20-1]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D$ -46 (c, 0.22 in CHCl₃). λ_{max} 212 (log ϵ 4.05); 245 (log ϵ 3.73); 295 (log ϵ 3.36) (EtOH).

15 α -Methoxy, N-formyl: Refractidine

[6883-09-6]

C₂₃H₂₈N₂O₄ 396.485

Alkaloid from *Aspidosperma refractum* (Apocynaceae). Mp 158-160°. [α]_D -140 (CHCl₃). pK_a 6.5. Several errors in the lit.; in two refs. the -COOMe group is omitted, and the abs. config. is wrongly shown. λ_{\max} 205 (log ϵ 4.36); 253 (log ϵ 4.16); 278 (log ϵ 3.72); 288 (log ϵ 3.67) (EtOH).

12,15 α -Dimethoxy, N-Ac: Pyrifoline

[6878-72-4]

C₂₅H₃₂N₂O₅ 440.538

Alkaloid from *Aspidosperma pyriformium* trunk bark and cambium (Apocynaceae). Mp 141.5-143.5°. [α]_D +102 (CHCl₃). pK_a 6.4 (66% DMF). λ_{\max} 217 (log ϵ 4.41); 261 (log ϵ 4.04); 287 (sh) (log ϵ 3.63) (EtOH).

16-Epimer: 16-Epikopsinine

[28161-78-6]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Kopsia singapurensis*. Yellowish oil. [α]_D²⁵ +7 (c, 0.22 in CHCl₃). λ_{\max} 201 (log ϵ 4.23); 241 (log ϵ 3.45); 291 (log ϵ 2.89) (EtOH).

Crow, W.D. et al., *Aust. J. Chem.*, 1955, **8**, 129-135 (*isol, uv*)

Gilbert, B. et al., *Experientia*, 1960, **16**, 61-62 (*Refractine, Pyrifoline*)

Kump, W.G. et al., *Helv. Chim. Acta*, 1961, **44**, 1503-1516; 1962, **45**, 854-858; 1090-1095; 1965, **48**, 1002-1012 (*Kopsinine, Kopsinine N-oxide, Kopsinilam, Kopsinoline, Pleiocarpine, Pleiocarpoline, Pleiocarpinine, Pleiocarpolinine, isol, uv, ir, struct*)

Djerassi, C. et al., *J.A.C.S.*, 1962, **84**, 1499-1501 (*Aspidofractine, Refractine*)

Battersby, A.R. et al., *J.C.S.*, 1962, 3245-3248 (*Pleioicarpine*)

Gilbert, B. et al., *Tet. Lett.*, 1962, 59-67 (*Pyrifoline, Refractidine*)

Djerassi, C. et al., *Helv. Chim. Acta*, 1963, **46**, 742-751 (*ms*)

Linde, H.H.A. et al., *Helv. Chim. Acta*, 1965, **48**, 1822-1842 (*Melodinus australis constits*)

Khan, Z.M. et al., *Helv. Chim. Acta*, 1965, **48**, 1957-1965; 1967, **50**, 625-627 (*Kopsinic acid methosalt*)

Achenbach, H. et al., *Tet. Lett.*, 1965, 3239-3242 (*10,11-Dioxopleioicarpine*)

Gilbert, B. et al., *Tetrahedron*, 1965, **21**, 1141-1166 (*Kopsinine, Aspidofractine, Refractine*)

Govindachari, T.R. et al., *Tetrahedron*, 1965, **21**, 2951-2956 (*isol, uv*)

Thomas, D.W. et al., *J.A.C.S.*, 1966, **88**, 3423-3426 (*Kopsinoline, Pleiocarpoline, Pleiocarpolinine*)

Raymond-Hamet, et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1967, **265**, 71-74 (*Refractine, Velloisine*)

Rakhimov, D.A. et al., *Khim. Prir. Soedin.*, 1967, **3**, 354-355; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 300-301 (*Kopsinilam*)

Gilbert, B. et al., *Alkaloids (Academic Press)*, 1968, **11**, 205-306 (*abs config*)

Klyne, W. et al., *Helv. Chim. Acta*, 1968, **51**, 1169-1184 (*abs config*)

Guggisberg, A. et al., *Helv. Chim. Acta*, 1969, **52**, 76-89 (*abs config*)

Linde, H.H.A. et al., *Pharm. Acta Helv.*, 1970, **45**, 248-253 (*Refractidine, Pyrifoline*)

Malikov, V.M. et al., *Khim. Prir. Soedin.*, 1971, **7**, 793; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 765-766 (*Kopsinilamine*)

Bruneton, J. et al., *Planta Med.*, 1982, **46**, 58-59 (*Pleioicarpine, Pleiocarpinine*)

Craveiro, A.A. et al., *Phytochemistry*, 1983, **22**, 1526-1528 (*Pyrifoline, 6-Demethoxy-pyrifoline*)

Zeche, M. et al., *Phytochemistry*, 1984, **23**, 171-174 (*3-Oxohydroxykopsinine*)

Magnus, P. et al., *Chem. Comm.*, 1985, 184-188 (*Kopsinine, Kopsinilam, synth*)

Ogawa, M. et al., *Tet. Lett.*, 1987, **28**, 3985-3986 (*synth*)

Kan-Fan, C. et al., *J. Nat. Prod.*, 1988, **51**, 703-707 (*N-Carbomethoxy-17-hydroxykopsinine*)

Wenkert, E. et al., *J.O.C.*, 1988, **53**, 4875-4877 (*synth*)

Lapr evote, O. et al., *Org. Mass Spectrom.*, 1990, **25**, 563-565 (*Kopsinoline, ms*)

Varea, T. et al., *J. Nat. Prod.*, 1993, **56**, 2166-2169 (*16-Epi-17-hydroxy- $\Delta^{14,15}$ -kopsinine*)


Kam, T.-S. et al., *Phytochemistry*, 1998, **47**, 145-147; 1999, **50**, 171-175 (*12-Methoxypleioicarpine, 17-Hydroxy- $\Delta^{14,15}$ -kopsinine*)

Lim, K.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*Kopsamidine B*)

Subramaniam, G. et al., *J. Nat. Prod.*, 2007, **70**, 1783-1789 (*16-Epikopsinine*)

Kopsinitarine A

[157207-85-7]



Absolute Configuration

C₂₃H₂₂N₂O₆ 422.437

Novel cage struct.; see also Mersingine A, M-216. Trace alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D +23 (c, 0.093 in CHCl₃). λ_{\max} 215 (log ϵ 4.41); 245 (log ϵ 3.85); 289 (log ϵ 3.06) (EtOH).

N-De(methoxycarbonyl): **Kopsinitarine B** [157207-86-8]

C₂₁H₂₀N₂O₄ 364.4

Trace alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D +188 (c, 0.051 in CHCl₃). λ_{\max} 212 (log ϵ 4.6); 246 (log ϵ 3.9); 290 (log ϵ 3.33) (EtOH).

14,15-Dihydro, 15 α -hydroxy: **Kopsinitarine D**

[184245-98-5]

C₂₃H₂₄N₂O₇ 440.452

Alkaloid from leaves of *Kopsia teoi*. Cryst. (CH₂Cl₂/Et₂O). Mp 258-260°. [α]_D -46 (c, 0.079 in CHCl₃). λ_{\max} 217 (log ϵ 4.43); 246 (log ϵ 3.86); 288 (log ϵ 3.03) (EtOH).

14,15-Dihydro, 15 α -hydroxy, N-de(methoxycarbonyl): **Kopsinitarine C**

[157207-87-9]

C₂₁H₂₂N₂O₅ 382.415

Trace alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D -96 (c, 0.026 in CHCl₃). λ_{\max} 213 (log ϵ 4.49); 246 (log ϵ 3.8); 290 (log ϵ 3.15) (EtOH).

Demethoxy, 14,15-dihydro: **Kopsinitarine E**

[947251-85-6]

C₂₂H₂₂N₂O₅ 394.426

Alkaloid from the stem bark of *Kopsia teoi*. Pale yellow oil. [α]_D -21 (c, 0.04 in


CHCl₃). λ_{\max} 207 (log ϵ 4.13); 238 (log ϵ 3.82); 279 (log ϵ 3.14); 287 (log ϵ 3.06) (MeOH).

Kam, T.-S. et al., *J. Nat. Prod.*, 1996, **59**, 1109-1112 (*isol, uv, pmr, cmr, ms, cryst struct*)

Lim, S.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1380-1383 (*Kopsinitarine E*)

Kopsinone

[116988-05-7]



Absolute Configuration

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from the stem bark of *Kopsia deverrei* (Apocynaceae). Amorph. [α]_D²⁰ +87 (c, 1.2 in CHCl₃). λ_{\max} 241 (log ϵ 4.17); 279 (log ϵ 3.55); 288 (log ϵ 3.51) (EtOH).

17 β -Alcohol, N-de(methoxycarbonyl):

Kopsinginol

[149355-61-3]

C₁₉H₂₂N₂O 294.396

Trace alkaloid from the stem bark of *Kopsia teoi*. [α]_D +288 (c, 0.016 in CHCl₃).

14,15-Dihydro, N-de(methoxycarbonyl):

Kopsonoline

[19634-39-0]

C₁₉H₂₂N₂O 294.396

Alkaloid from the stem bark of *Kopsia teoi*. Light yellow oil. [α]_D +27 (c, 0.41 in CHCl₃). λ_{\max} 209 (log ϵ 4.06); 241 (log ϵ 3.77); 291 (log ϵ 3.32) (EtOH).

10-Methoxy: **10-Methoxykopsinone**

[116979-55-6]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from the leaves of *Kopsia deverrei* (Apocynaceae). Cryst. (Me₂CO/Et₂O). Mp 178°. [α]_D²⁰ -11 (c, 1.4 in CHCl₃). λ_{\max} 219 (log ϵ 4.27); 246 (log ϵ 4.03); 297 (log ϵ 3.38) (EtOH).

10-Methoxy, 14,15-dihydro: **14,15-Dihydro-10-methoxykopsinone**

[116979-53-4]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the leaves of *Kopsia deverrei* (Apocynaceae). Cryst. (Me₂CO/Et₂O). Mp 214°. [α]_D²⁰ -11 (c, 1.2 in CHCl₃). λ_{\max} 219 (log ϵ 4.29); 246 (log ϵ 4.06); 296 (log ϵ 3.43) (EtOH).

12-Methoxy: **12-Methoxykopsinone**

[116979-54-5]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from the leaves of *Kopsia deverrei* (Apocynaceae). Cryst. (Me₂CO/Et₂O). Mp 176°. [α]_D²⁰ +69 (c, 2.1 in CHCl₃). λ_{\max} 219 (log ϵ 4.23); 246 (log ϵ 3.94); 279 (log ϵ 3.26) (EtOH).

Kan-Fan, C. et al., *J. Nat. Prod.*, 1988, **51**, 703-707 (*Kopsinone*)


Do Carmo Carreiras, M. et al., *J. Nat. Prod.*, 1988, **51**, 806-808 (*Kopsia deverrei constits*)

Kam, T.S. *et al.*, *Phytochemistry*, 1996, **42**, 539-541 (*Kopsingol*)

Lim, S.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1380-1383 (*Kopsonoline*)

Kopsirachin K-81

2-(3,4-Dihydroxyphenyl)-3,4-dihydro-6,8-bis(octahydro-2,4,7-trimethyl-1H-2-pyridin-1-yl)-2H-1-benzopyran-3,5,7-triol, 9CI
[89759-08-0]




$C_{37}H_{52}N_2O_6$ 620.828

Alkaloid from the leaves of *Kopsia dasyrachis* (Apocynaceae). Amorphous powder. $[\alpha]_D^{25} +65.8$ (c, 0.987 in $CHCl_3$).

Homburger, K. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 237 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kopsone K-82

[180841-83-2]



Relative Configuration


$C_{11}H_{19}NO$ 181.277

Alkaloid from leaves of *Kopsia macrophylla*. Amorphous. $[\alpha]_D^{25} +132$ (c, 1.25 in $CHCl_3$).

Kan-Fan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 283 (*isol, ir, pmr, cmr, ms, struct*)

Kororamide K-83

[261373-26-6]




$C_{45}H_{64}N_{10}O_{10}S_2$ 969.194

Isol. from *Lyngbya majuscula*. Amorphous solid. λ_{max} 238 (ϵ 6400) (MeOH).

Mitchell, S.S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 279-282 (*isol, pmr, cmr, cd, uv*)

Korormicin A K-84

[200433-05-2]



$C_{25}H_{39}NO_5$ 433.587

Prod. by the marine bacterium *Pseudoalteromonas* sp. F-420 isol. from an alga. Active against marine gram-negative bacteria. Oil. $[\alpha]_D^{26} -24.4$ (c, 0.3 in EtOH). λ_{max} 233 ($\log \epsilon$ 4.8) (EtOH).

Yoshikawa, K. *et al.*, *J. Antibiot.*, 1997, **50**, 949-953; 1999, **52**, 182-185 (*isol, uv, ir, pmr, cmr, activity*)

Uehara, H. *et al.*, *Tet. Lett.*, 1999, **40**, 8641-8645 (*synth, abs config*)

Kobayashi, Y. *et al.*, *Eur. J. Org. Chem.*, 2001, 1873-1881 (*synth, abs config*)

Korormicin B K-85

As Korormicin A, K-84 with n = 5

$C_{23}H_{35}NO_5$ 405.533

Stereochem. not confirmed. Prod. by *Pseudoalteromonas* sp. F-420. Oil. λ_{max} 232 ($\log \epsilon$ 4.55) (EtOH).

Yoshikawa, K. *et al.*, *J. Antibiot.*, 2003, **56**, 866-870

Korormicin C K-86

As Korormicin A, K-84 with n = 6

$C_{24}H_{37}NO_5$ 419.56

Prod. by *Pseudoalteromonas* sp. F-420. Oil. Stereochem. not confirmed. λ_{max} 232 ($\log \epsilon$ 4.59) (EtOH).

Yoshikawa, K. *et al.*, *J. Antibiot.*, 2003, **56**, 866-870

Korormicin D K-87

As Korormicin A, K-84 with n = 8


$C_{26}H_{41}NO_5$ 447.614

Prod. by *Pseudoalteromonas* sp. F-420. Oil. Stereochem. not confirmed. λ_{max} 233 ($\log \epsilon$ 4.43) (EtOH).

Yoshikawa, K. *et al.*, *J. Antibiot.*, 2003, **56**, 866-870

Korsevine K-88

Corsevine
[18209-21-7]



$C_{28}H_{45}NO_2$ 427.669

Alkaloid from *Fritillaria imperialis* and *Korolkowia sewerzowii*. Needles (Me_2CO). Mp 170-172°. $[\alpha]_D^{25} -87$ (c, 0.5 in MeOH).

Hydrobromide:

Cryst. (H_2O). Mp 244-245°.

Oxime: Mp 200-201°.

3-O- β -D-Glucopyranoside: *Korsemine*

[74119-90-7]

$C_{34}H_{55}NO_7$ 589.811

Alkaloid from *Korolkowia sewerzowii*. $[\alpha]_D -46.9$ (c, 0.81 in MeOH).


Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 398; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 333 (*ir, uv, ms, pmr*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 823; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 728 (*Korsemine*)

Akhtar, M.N. *et al.*, *Phytochemistry*, 2003, **63**, 115-122 (*isol, pmr*)

Korsevine K-89

23,26-Epoxy-3-hydroxy-16,28-secosolanid-22(28)-en-6-one. *Corsevine*
[27336-00-1]



Absolute configuration probable structure

$C_{27}H_{41}NO_3$ 427.626


Alkaloid from *Korolkowia sewerzowii* (Liliaceae). Mp 224-225°. $[\alpha]_D -16$ (sol. not specified).

Ac: Mp 154-155°.

Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 600; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 519 (*ir, pmr, ms, struct*)

Korundamine A K-90

[212957-33-0]



Absolute Configuration


$C_{47}H_{48}N_2O_8$ 768.905

Alkaloid from *Ancistrocladus korupensis*. Exhibits antimalarial and cytotoxic properties. Tan solid. λ_{max} 206 (ϵ 56234); 232 (ϵ 50120); 261 (ϵ 33880); 331 (ϵ 18620); 347 (ϵ 21880); 381 (ϵ 22390) (MeOH) (Berdy).

Hallock, Y.F. *et al.*, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 1729-1734 (*isol, pmr, cmr*)

Korupensamine A

[158182-18-4]



Absolute configuration

 $C_{23}H_{25}NO_4$ 379.455

Alkaloid from leaves and twigs of *Ancistrocladus korupensis* (Ancistrocladaceae). Shows antimalarial activity. Light brown solid. $[\alpha]_D^{25}$ -75.5 (c, 1.84 in MeOH).

5'-Me ether: Korupensamine C

[158182-19-5]

 $C_{24}H_{27}NO_4$ 393.482

From leaves and twigs of *Ancistrocladus korupensis* (Ancistrocladaceae). Light brown solid. $[\alpha]_D^{25}$ -62 (c, 0.54 in MeOH).

6-Me ether, N-Me: Ancistrolikokine A

[297749-36-1]

 $C_{25}H_{29}NO_4$ 407.508

Alkaloid from *Ancistrocladus likoko*. Solid. Mp 230-231°. $[\alpha]_D^{25}$ +79.6 (c, 0.25 in MeOH).

5',6-Di-Me ether, N-Me: Ancistrocongoline B

[455255-18-2]

 $C_{26}H_{31}NO_4$ 421.535

Alkaloid from the root bark of *Ancistrocladus congolensis*. Solid (MeOH). Mp 161-162°. $[\alpha]_D^{23}$ -12 (c, 0.22 in $CHCl_3$).

4'-O-De-Me, 5'-Me ether, N-Me: Ancistrocongoline A

[455255-21-7]

 $C_{24}H_{27}NO_4$ 393.482

Alkaloid from the stem and root bark of *Ancistrocladus congolensis*. Light brown powder. $[\alpha]_D^{23}$ +6.9 (c, 0.05 in $CHCl_3$).

3-Epimer, N-Me: Korupensamine D

[158182-20-8]

 $C_{24}H_{27}NO_4$ 393.482

From leaves and twigs of *Ancistrocladus korupensis* (Ancistrocladaceae). Solid. $[\alpha]_D^{25}$ +6 (c, 0.3 in MeOH).

Atropisomer: Korupensamine B

[158252-04-1]

 $C_{23}H_{25}NO_4$ 379.455

From leaves and twigs of *Ancistrocladus korupensis* (Ancistrocladaceae). Shows antimalarial activity. Light brown solid. $[\alpha]_D^{25}$ +65 (c, 0.76 in MeOH).

Atropisomer, 5',8-di-Me ether, N-Me: Ancistrocongoline C

[455255-19-3]

 $C_{26}H_{31}NO_4$ 421.535

Alkaloid from the root bark of *Ancistrocladus congolensis*. Brownish powder. Mp 140-141°. $[\alpha]_D^{24}$ +128.1 (c, 0.12 in $CHCl_3$).

Atropisomer, 4'-O-de-Me, 5',8-di-Me**ether: Korupensamine E** $C_{24}H_{27}NO_4$ 393.482

From the leaves of *Ancistrocladus korupensis* (Ancistrocladaceae). Antimalarial agent. Pale yellow solid. $[\alpha]_D^{25}$ +15.4 (c, 0.26 in MeOH). λ_{max} 232 (log ϵ 4.46); 289 (log ϵ 3.68); 308 (log ϵ 3.75); 322 (log ϵ 3.67); 337 (log ϵ 3.58) (MeOH).

Atropisomer, 4'-O-de-Me, 5',8-di-Me**ether, N-Me: Ancistrolikokine C. N-Methylkorupensamine E**

[297749-38-3]

 $C_{25}H_{29}NO_4$ 407.508

Alkaloid from *Ancistrocladus likoko*. Amorph. solid. $[\alpha]_D^{20}$ -54.9 (c, 0.11 in EtOH).

Atropisomer, 1-epimer, 4'-O-de-Me, 5',8-di-Me ether: Ancistrolikokine B

[297749-37-2]

 $C_{24}H_{27}NO_4$ 393.482

Alkaloid from *Ancistrocladus likoko*. Amorph. solid. $[\alpha]_D^{25}$ -166.4 (c, 0.11 in EtOH).

Bringmann, G. *et al.*, *Heterocycles*, 1994, **39**, 503 (synth)

Hallock, Y.F. *et al.*, *J.O.C.*, 1994, **59**, 6349

(isol, uv, ir, pmr, cmr, cd, cryst struct)

Hoye, T.R. *et al.*, *Tet. Lett.*, 1996, **37**, 3097;

3099 (synth)

Hallock, Y.F. *et al.*, *J. Nat. Prod.*, 1997, **60**, 677

(Korupensamine E)

Hoye, T.R. *et al.*, *J.O.C.*, 1999, **64**, 7184-7201

(synth)

Bringmann, G. *et al.*, *J. Nat. Prod.*, 2000, **63**,

1333-1337 (Ancistrolikokines)

Bringmann, G. *et al.*, *J.O.C.*, 2000, **65**, 2069-

2077 (synth)

Bringmann, G. *et al.*, *Synthesis*, 2000, 1843-

1847 (synth)

Kamikawa, K. *et al.*, *Tetrahedron*, 2000, **56**,

2325-2337 (synth)

Bringmann, G. *et al.*, *J. Nat. Prod.*, 2002, **65**,


1096-1101 (Ancistrocongolines)

Watanabe, T. *et al.*, *J.O.C.*, 2004, **69**, 4152-

4158 (synth)

Kottamide A

[446862-69-7]

 $R^1 = -CH(CH_3)CH_2CH_3$, $R^2 = -CH(CH_3)_2$ $C_{21}H_{24}Br_2N_4O_2$ 524.254

Isol. from the New Zealand ascidian *Pycnoclavella kottae*. Cytotoxic agent. Amorph. solid. $[\alpha]_D^{20}$ +160 (c, 0.2 in MeOH). λ_{max} 236 (log ϵ 4.46); 284 (log ϵ 4.14) (MeOH).

5-Debromo: Kottamide B

[446862-70-0]

 $C_{21}H_{25}BrN_4O_2$ 445.358

Isol. from the New Zealand ascidian *Pycnoclavella kottae*. Amorph. solid. $[\alpha]_D^{20}$ +245 (c, 0.2 in MeOH). Obt. as a mixt. with Kottamide C to which data

refers. λ_{max} 232 (log ϵ 4.44); 286 (log ϵ 4.13) (MeOH).

6-Debromo: Kottamide C

[446862-71-1]

 $C_{21}H_{25}BrN_4O_2$ 445.358

Isol. from *Pycnoclavella kottae*. Obt. as a mixt. with Kottamide B.

Appleton, D.R. *et al.*, *J.O.C.*, 2002, **67**, 5402-5404 (isol, pmr, cmr, N-15 nmr, ms)

Kottamide D**K-93**

[446862-72-2]

As Kottamide A, K-92 with


 $R^1 = CH_3$, $R^2 = -CH(CH_3)CH_2CH_3$ $C_{19}H_{20}Br_2N_4O_2$ 496.201

Isol. from the New Zealand ascidian *Pycnoclavella kottae*. Antiinflammatory and cytotoxic agent. Amorph. solid. $[\alpha]_D^{20}$ +150 (c, 0.2 in MeOH). λ_{max} 237 (log ϵ 4.39); 286 (log ϵ 4.11) (MeOH).

Appleton, D.R. *et al.*, *J.O.C.*, 2002, **67**, 5402-5404 (isol, pmr, cmr, N-15 nmr, ms)

Kottamide E**K-94**

[642492-69-1]

 $C_{16}H_{14}Br_2N_4O_3S_2$ 534.252

Alkaloid from the ascidian *Pycnoclavella kottae*. Amorph. solid. λ_{max} 202 (log ϵ 4.36); 239 (log ϵ 4.41); 305 (log ϵ 3.92) (MeOH).

Appleton, D.R. *et al.*, *Tet. Lett.*, 2003, **44**, 8963-8965 (isol, pmr, cmr)

Koumicine**K-95**

[1358-74-3]


 $C_{26}H_{30}N_2O_4$ 434.534

Indole alkaloid. Struct. unknown. Alkaloid from *Gelsemium elegans* (Loganiaceae). Mp 232-234°.

Liu, C.-T. *et al.*, *Huaxue Xuebao*, 1961, **27**, 47-58; *CA*, **59**, 1404ib

Koumine**K-96**

1,2,18,19-Tetrahydro-3,17-epoxy-7,20(2H,19H)-cyclovobasan, 9CI
[1358-76-5]



(-)-form

 $C_{20}H_{22}N_2O$ 306.407

Main alkaloid from *Gelsemium elegans* (Loganiaceae). Cryst. (Me_2CO). Mp

172°. $[\alpha]_D$ -272 (EtOH). λ_{\max} 220 (log ϵ 4.32); 262 (log ϵ 3.84) (EtOH).

N^b-Oxide(R-): Semisynthetic by oxidation of Koumine. Cryst. + 2H₂O (Me₂CO). Mp 214-216°. $[\alpha]_D^{19}$ -157 (c, 0.14 in MeOH).

N^b-Oxide(S-): **Koumine N-oxide** [113900-75-7]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from the leaves of *Gelsemium elegans* (Loganiaceae). Cryst. + ¼ H₂O (Me₂CO). Mp 111-113°. $[\alpha]_D^{19}$ -237 (c, 0.14 in MeOH).

1,2-Dihydro: **Dihydrokoumine**

[104654-06-0]
C₂₀H₂₄N₂O 308.422
Alkaloid from roots of *Gelsemium elegans* (Loganiaceae).

18,19-Dihydro, 19R-hydroxy:

19R-Kouminol
[126398-76-3]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from the roots or whole plant of *Gelsemium elegans*. Needles or amorph. solid. Mp 198-200°. $[\alpha]_D$ -232.7 (c, 0.52 in MeOH) (-153.8).

18,19-Dihydro, 19S-hydroxy:

19S-Kouminol
[126398-77-4]
Alkaloid from roots of *Gelsemium elegans*. Amorph. $[\alpha]_D^{18}$ -209.9 (c, 1.0 in EtOH).

[83705-29-7, 119241-69-9]

Liu, C.-T. *et al.*, *J.A.C.S.*, 1981, **103**, 4634-4635 (*pmr, cmr, cryst struct*)
Khuong-Huu, F. *et al.*, *Tet. Lett.*, 1981, 733-734 (*uv, ir, pmr, ms, cryst struct*)
Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075-5094 (*isol, uv, ir, pmr, ms, struct, oxide*)
Sun, F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1180-1182 (*Kouminols*)
Takayama, H. *et al.*, *Heterocycles*, 1990, **30**, 325-327 (*synth*)
Magnus, P. *et al.*, *J.A.C.S.*, 1990, **112**, 5220-5230 (*synth*)
Lin, L.Z. *et al.*, *Phytochemistry*, 1990, **29**, 965-968 (*Kouminols*)
Zhang, Z. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 365-368 (*Dihydrokoumine*)
Bailey, P.D. *et al.*, *J.C.S. Perkin 1*, 1993, 441-449 (*synth*)

Kouminidine

K-97

[1358-77-6]

C₂₀H₂₆N₂O₄ 358.436

Struct. unknown. Alkaloid from *Gelsemium elegans* (Loganiaceae). Prisms (MeOH/Et₂O). Mp 299° dec.

Chi, Y.F. *et al.*, *J.A.C.S.*, 1938, **60**, 1723 (*isol*)
Liu, C.-T. *et al.*, *Huaxue Xuebao*, 1961, **27**, 47-58; *CA*, **59**, 14041b

Krelagine

K-98

[1358-78-7]

C₁₇H₁₉NO₄ 301.341

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Crinum powellii* var. *krelagei* (Amaryllidaceae). Cryst. (Me₂CO). Mp 202° dec. $[\alpha]_D^{25}$ +290 (c, 0.2 in CHCl₃).

Picrate: Mp 268° dec.

Methodide: Mp 260° dec.

Methyl perchlorate: Mp 233° dec.


Methopicrate: Mp 256-257° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 498 (*isol*)

Kreysiginine

K-99

Alkaloid CC21



(+)-form

C₂₁H₂₇NO₅ 373.448

Alkaloid CC21 was the (-)-enantiomer.

(+)-form [19741-86-7]

Alkaloid from *Kreysigia multiflora* (Liliaceae) and seeds of *Iphigenia stellata*. Prisms (EtOH or Et₂O). Mp 149°. $[\alpha]_D^{24}$ +89 (c, 1.98 in EtOH). λ_{\max} 218 (ϵ 36220); 274 (ϵ 2240) (no solvent reported).

Hydrobromide:

Cryst. + 1H₂O (EtOH/Et₂O). Mp 142-143°.

Methodide:

Prisms + 1Me₂CO (Me₂CO). Mp 150° (softens).

(-)-form

Alkaloid from *Colchicum cornigerum* (Liliaceae). Mp 151-154°. $[\alpha]_D^{22}$ -100 (c, 0.55 in EtOH). Opt. rotn. decreased to -42° on recryst. due to racemisation. λ_{\max} 212 (log ϵ 4.6); 231 (sh) (log ϵ 4.02); 275 (log ϵ 3.07); 282 (log ϵ 3.07) (EtOH).

Badger, G.M. *et al.*, *J.C.S.*, 1960, 445-447 (*isol, uv*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1968, 695-697 (*pmr, struct*)

Hart, N.K. *et al.*, *Tet. Lett.*, 1968, 2891-2894 (*pmr, struct*)

Potěšilová, H. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540-3552 (*Alkaloid CC-21*)


Fridrichsons, J. *et al.*, *Tetrahedron*, 1970, **26**, 1869-1877 (*cryst struct*)

Kreysiginone

K-100

Dienone I

[17441-87-1]



Relative configuration

C₂₀H₂₃NO₄ 341.406

Natural-form

Alkaloid from *Kreysigia multiflora* (Liliaceae). Mp 193-194°. Opt. act. not given.

9,10-Dihydro: **Dihydrokreysiginone**

[17750-51-5]

C₂₀H₂₅NO₄ 343.422

Minor alkaloid from *Kreysigia multiflora*. Mp 217-222° dec. Probably corresponds in config. at the spiro-centre to Kreysiginone.

(±)-form

Synthetic. Cryst. (C₆H₆). Mp 145-149° (155°, resolid., 193-194°).

8-Epimer: **Dienone II**

Synthetic. Mp 202° dec.

Battersby, A.R. *et al.*, *Chem. Comm.*, 1967, 934 (*isol, ms, uv, ir, pmr, deriv*)

Harmon, R.E. *et al.*, *J. Het. Chem.*, 1970, **7**, 1077 (*synth, ir, uv, ms, pmr*)


Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1974, 1394 (*isol*)

Kametani, T. *et al.*, *J.A.C.S.*, 1977, **99**, 3805 (*synth*)

Kribine

K-101

[69241-15-2]



C₂₀H₂₄N₂O₂ 324.422

Mixt. of C-17 epimers. Alkaloid from stem bark of *Strychnos dale*, *Strychnos elaeocarpa*, *Strychnos nigriflora*, *Strychnos spinosa*, *Strychnos usambarensis* (Loganiaceae). Rearr. to Akagerine, A-222 on acid treatment.

Me ether (17β-): **17-O-Methylkribine**.

21-O-Methylkribine

[69307-96-6]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Strychnos dale* and *Strychnos elaeocarpa* (Loganiaceae). $[\alpha]_D^{22}$ -52 (c, 0.02 in CHCl₃). Prob. artifact. The struct. was assigned (and the name 21-O-methylkribine allocated) before the correct struct. of Kribine was assigned. The 17-configs. of this isomer and its 17-epimer (see below) are therefore not well established and may be reversed.

Me ether (17α-): **Epi-17-O-methylkribine**.

Epi-21-O-methylkribine

[69241-16-3]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Strychnos dale* and *Strychnos elaeocarpa* (Loganiaceae). $[\alpha]_D^{22}$ -212 (c, 0.02 in CHCl₃).


10-Hydroxy, *Me ether* (17α-): **10-Hydroxy-17-O-methylkribine**.

10-Hydroxy-21-O-methylkribine

[73360-06-2]

C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Strychnos decussata* (Loganiaceae).**10-Hydroxy, Me ether (17β -): 10-Hydroxyepi-17-O-methylkrigine. 10-Hydroxyepi-21-O-methylkrigine**
[73367-21-2]C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Strychnos decussata*. Amorph. powder. Mp 176-180° dec.Rolfen, W. *et al.*, *Planta Med.*, 1978, **34**, 264-273 (*isol, uv, ir, pmr, cmr, ms*)
Rolfen, W.N.A. *et al.*, *J. Nat. Prod.*, 1980, **43**, 97-102 (*10-Hydroxy-17-O-methylkrigine, 10-Hydroxyepi-17-O-methylkrigine*)
Verpoorte, R. *et al.*, *J.C.S. Perkin I*, 1984, 1455-1457 (*cmr, struct*)
Delaude, C *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, Strychnos*)**Krigeine**

[905-37-3]

K-102C₁₈H₂₁NO₆ 347.367Alkaloid from the bulbs of *Nerine krigiei* (Amaryllidaceae). Cryst. (Me₂CO aq.). Mp 209-210° dec. [α]_D²⁵ +234 (c, 0.15 in CHCl₃).**7-Ketone (lactone): Neronine**

[1167-58-4]

C₁₈H₁₉NO₆ 345.351Alkaloid from bulbs of *Nerine krigiei* (Amaryllidaceae). Hygroscopic prisms + 1H₂O (EtOAc or C₆H₆/cyclohexane). Mp 196-197°. [α]_D²⁴ +161.6 (c, 0.57 in CHCl₃).**7-Ketone, Ac:**

Cryst. (EtOH). Mp 201-202°.

7-Ketone (lactone), O-de-Me: NorneronineC₁₇H₁₇NO₆ 331.324Alkaloid from the bulbs of *Pancreatium longiflorum* (Amaryllidaceae). Needles (CHCl₃/MeOH or EtOH). Mp 238-240° dec. [α]_D³⁰ +90 (MeOH).**7-Ketone, O-de-Me, di-Ac:**Needles (Me₂CO). Mp 248-250° dec. [α]_D³⁰ +20 (CHCl₃).**5-Deoxy: Krigenamine**



[1165-00-0]

C₁₈H₂₁NO₅ 331.368Alkaloid from the bulbs of *Nerine krigiei* (Amaryllidaceae). Needles (Me₂CO, EtOH or EtOAc). Mp 210-211°. [α]_D +210 (c, 0.43 in CHCl₃).**5-Deoxy; methiodide:** Mp 235-237° dec. [α]_D +150 (c, 1.2 in H₂O).**5-Deoxy, 7-ketone (lactone): Didehydrokrigenamine. Oxokrigenamine**
[1165-01-1]C₁₈H₁₉NO₅ 329.352Alkaloid from the bulbs of *Nerine krigiei* (Amaryllidaceae). Prisms (EtOAc). Mp 147-148°. [α]_D²⁰ +117 (c, 1.0 in CHCl₃). Apparently an artifact.**5-Deoxy, 7-ketone; methiodide:**


Prisms (MeOH). Mp 254-255° dec.

Briggs, C.K. *et al.*, *J.A.C.S.*, 1956, **78**, 2899 (*isol, uv, Krigeine, Neronine*)
Garbutt, D.F.C. *et al.*, *J.C.S.*, 1962, 5010 (*isol, ir, uv, Krigenamine*)
Jeffs, P.W. *et al.*, *Tet. Lett.*, 1963, 217 (*struct, Neronine*)
Hawksworth, W.A. *et al.*, *J.C.S.*, 1965, 1991 (*pmr, struct*)
Rangaswami, S. *et al.*, *Tet. Lett.*, 1966, 4481 (*Norneronine*)
Clardy, J. *et al.*, *J.O.C.*, 1972, **37**, 49 (*config*)**Kryptocurine***Cryptocurine*Struct. unknown. Alkaloid from calabash curare (from bark of an unknown plant) (Loganiaceae). No col. with Ce(SO₄)₂. Forms a cryst. tetraphenylborate. Chloride, picrate and reineckate were noncryst.


► Mod. toxic.

Meyer, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 1214-1218 (*isol, uv*)**K-103****Kuafumine****4,9,10-Trimethoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI**
[107208-76-4]**K-104**C₂₀H₁₅NO₆ 365.342Alkaloid from *Fissistigma glaucescens* (Annonaceae). Cytotoxic. Reddish needles (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 230-232°. λ_{max} 214 (ε 20890); 245 (ε 13800); 283 (ε 23990); 375 (ε 2400) (MeOH) (Berdy). λ_{max} 214; 238; 245; 375 (MeOH-NaOH) (Berdy).Wu, Y.-C. *et al.*, *Heterocycles*, 1987, **26**, 9 (*uv, ir, pmr, struct*)**Kuchinoenamine****K-105**C₂₃H₂₅Br₃N₂O₅ 649.173Alkaloid from the sponge *Hexadella* sp. Antibacterial agent. Amorph. solid. [α]_D²¹ +21 (c, 0.05 in MeOH). λ_{max} 244 (ε 8620); 310 (ε 12730) (MeOH).Matsunaga, S. *et al.*, *J.O.C.*, 2005, **70**, 1893-1896 (*isol, pmr, cmr*)**Kumujansine**


[116064-72-3]

K-106C₂₈H₂₅N₄O₂[⊕] 449.531Alkaloid from *Picrasma quassioides* (Simaroubaceae). Light yellow granular cryst. (as chloride). Mp 249-250° dec. (chloride).**8',11-Dimethoxy: Kumujansine B. Kumujantine**
[116064-73-4]C₃₀H₂₉N₄O₄[⊕] 509.583Alkaloid from *Picrasma quassioides* (Simaroubaceae). Light yellow granular cryst. (as chloride). Mp 270° dec. (chloride).Yang, J.S. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 267-272; *CA*, **109**, 107726n**Kuramerine**

[21284-19-5]

K-107C₂₈H₄₄NO₈[⊕] 522.657Alkaloid from *Liparis kurameri* and *Liparis kumokiri* (Orchidaceae). [α]_D²⁰ -19.7 (MeOH) (as chloride).**Tetra-Ac:**

Yellow needles (EtOH)(as picrate). Mp 150-151° (picrate).

Nishikawa, K. *et al.*, *Tet. Lett.*, 1967, 2579 (*isol, struct, ir, uv*)Tanino, H. *et al.*, *Tetrahedron*, 1969, **25**, 3033 (*synth*)**Kuraramine****6-[5-(Hydroxymethyl)-1-methyl-3-piperidinyl]-2(1H)-pyridinone, 9CI**
[79659-61-3]**K-108**C₁₂H₁₈N₂O₂ 222.286Alkaloid from the fresh flowers of *Sophora flavescens* (Fabaceae). Amorph. solid. [α]_D²⁰ +8.4 (c, 0.52 in EtOH). λ_{max} 226 (log ε 3.59); 304 (log ε 3.64) (EtOH).**Epimer: Isokuraramine**

[85799-36-6]

C₁₂H₁₈N₂O₂ 222.286

Alkaloid from *Sophora flavescens* (Fabaceae). Amorph.


Murakoshi, I. *et al.*, *Phytochemistry*, 1981, **20**, 1407-1409; 1982, **21**, 2379-2384 (*isol. pmr. cmr, ms*)

Honda, T. *et al.*, *J.O.C.*, 2005, **70**, 499-504 (*synth*)

Kurchamide

K-109

[148717-72-0]



$C_{21}H_{27}NO_3$ 341.449

Alkaloid from bark of *Holarrhena antidysenterica*. Irregular plates (MeOH). Mp 84-86°. λ_{max} 238 (MeOH).

N-De-Me: **Norkurchamide**

[609343-72-8]

$C_{20}H_{25}NO_3$ 327.422

Alkaloid from the bark of *Holarrhena pubescens*. Orange rods (MeOH). Mp 130-132°. λ_{max} 206 ; 246 (MeOH).

Begum, S. *et al.*, *Heterocycles*, 1993, **36**, 717-723 (*Kurchamide. struct*)

Siddiqui, B.S. *et al.*, *Heterocycles*, 2003, **60**, 909-915 (*Norkurchamide*)

Kurchenine

K-110

$C_{21}H_{32}N_2O_2$ 344.496

Struct. unknown. Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Platelets (MeOH). Mp 335-336°. $[\alpha]_D$ -92 (2M HCl). Conts. no N-Me gps. Prob. a steroidal alkaloid.

Sulfate:

Needles (EtOH). $[\alpha]_D$ -78.3 (H₂O).

Bertho, A. *et al.*, *Ber.*, 1933, **66**, 786-790

Kurchicine

K-111

[1400-17-5]

Steroidal alkaloid. Struct. unknown. Alkaloid from *Holarrhena floribunda* and *Wrightia tomentosa* (Apocynaceae). Poorly documented. No struct. reported to 2007.

▶ OC7849100

Jayaswal, S.B. *et al.*, *CA*, 1977, **86**, 12205j (*isol*)

Jayaswal, S.B. *et al.*, *Indian J. Pharm.*, 1977, **39**, 37-39 (*isol*)


Zaki, A.Y. *et al.*, *CA*, 1984, **100**, 117842a (*isol*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, KHU136

Kurchilidine

K-112

[148717-71-9]



$C_{22}H_{31}NO$ 325.493

Alkaloid from bark of *Holarrhena antidysenterica*. Irregular plates (MeOH).


Mp 118-120°. λ_{max} 238 (MeOH).

Begum, S. *et al.*, *Heterocycles*, 1993, **36**, 717 (*isol. uv. ir. pmr. ms. struct*)

Kurcholesine

K-113

[6869-47-2]



$C_{25}H_{44}N_2O_2$ 404.635

Produced from extract of the rind of *Holarrhena antidysenterica* by methylation with CH₂O/HCOOH (Apocynaceae). Mp 218.5-221.5°. $[\alpha]_D$ -4 (CHCl₃).

Hydrochloride: Mp 255-265° dec.

Stereoisomer, 17,20-didehydro: Holacine

[84953-46-8]

$C_{26}H_{44}N_2O_2$ 416.646

Alkaloid from the bark of *Holarrhena antidysenterica* (Apocynaceae). Needles (EtOAc/MeOH). Mp 270-271°. $[\alpha]_D^{27}$ -49 (EtOH). Undetermined stereochem.

7,20-Diepimer: Regholarrenine E

[128718-02-5]

$C_{25}H_{44}N_2O_2$ 404.635

Alkaloid from stem bark of *Holarrhena antidysenterica* (Apocynaceae). Brown solid (MeOH). Mp 285-287°.

Tschesche, R. *et al.*, *Chem. Ber.*, 1962, **95**, 1114 (*isol. uv. ir*)

Tschesche, R. *et al.*, *Tet. Lett.*, 1964, 1659 (*struct. ms. ir. pmr*)

Siddiqui, S. *et al.*, *Pak. J. Sci. Ind. Res.*, 1982, **25**, 201 (*Holacine*)

Bhutani, K.K. *et al.*, *Phytochemistry*, 1990, **29**, 969 (*Regholarrenine E*)


Kuroyuridine

K-114

17,23-Epoxy-5,6-dihydroveratraman-

2,3,6-triol, 9CI

[123085-17-6]



$C_{27}H_{43}NO_4$ 445.641

Alkaloid from the fresh bulbs of *Fritillaria camtschatcensis* and *Fritillaria maximowiczii*. Amorph. powder. $[\alpha]_D^{25}$ -10.1 (c, 0.4 in MeOH).

6-Ketone: Pengbeisine B

$C_{27}H_{41}NO_4$ 443.625

Alkaloid from the fresh bulbs of

Fritillaria monatha. Needles (Me₂CO).

23-Epimer: 23-Epikuroyuridine. 23-Iso-kuroyuridine

[169786-65-6]

$C_{27}H_{43}NO_4$ 445.641

Alkaloid from fresh bulbs of *Fritillaria maximowiczii*. Amorph. $[\alpha]_D^{25}$ -14.2 (c, 0.40 in MeOH).

2,3-Diepimer, 6-ketone: Pengbeisine A

$C_{27}H_{41}NO_4$ 443.625

Alkaloid from the fresh bulbs of *Fritillaria monatha*. Needles (Me₂CO).


Mimaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1090-1092 (*Fritillaria camtschatcensis constii*)

Qian, Z.-Z. *et al.*, *Phytochemistry*, 1995, **40**, 979-981 (*Kuroyuridine, 23-Epikuroyuridine*)

Liu, H.N. *et al.*, *Chin. Chem. Lett.*, 2008, **19**, 544-546 (*Pengbeisines A,B*)

Kurramine

K-115



$C_{33}H_{28}N_2O_5$ 532.595

(R)-form

O⁶-Me: 1,2-Dehydromicranthine

[58207-93-5]

$C_{34}H_{30}N_2O_5$ 546.621

Alkaloid from the leaves and terminal twigs of an unnamed *Daphandra* sp. (Monimiaceae). Mp 188-194° dec. $[\alpha]_D^{20}$ -150 (CHCl₃).

(S)-form [92664-89-6]

Alkaloid from the stems of *Cocculus pendulus* (Menispermaceae). $[\alpha]_D$ +83 (c, 0.13 in MeOH).

2'-α-N-Oxide: Kurramine 2'-α-N-oxide

$C_{33}H_{28}N_2O_6$ 548.594

Alkaloid from *Cocculus pendulus*. Amorph. yellow powder. $[\alpha]_D^{25}$ +50 (c, 0.01 in MeOH). λ_{max} 201 (log ε 4.58); 228 (log ε 4.45); 264 (log ε 4.1); 290 (log ε 3.8); 333 (log ε 3.66) (MeOH).

2'-β-N-Oxide: Kurramine 2'-β-N-oxide

$C_{33}H_{28}N_2O_6$ 548.594

Alkaloid from *Cocculus pendulus*. Amorph. yellow powder. $[\alpha]_D^{25}$ +60 (c, 0.01 in MeOH). λ_{max} 203 (log ε 4.61); 235 (log ε 4.41); 263 (log ε 4.02); 291 (log ε 3.81); 341 (log ε 3.52) (MeOH).

O⁶-Me: 1,2-Dehydroapateline

[68711-77-3]

$C_{34}H_{30}N_2O_5$ 546.621

Alkaloid from the bark of *Daphandra apatela* and *Doryphora aromatica*, and from the stems of *Cocculus pendulus*. Mp 196-198° dec. (192-198° dec.). $[\alpha]_D^{20}$ +137 (CHCl₃).

O⁶-Me, dipicrate:

Cryst. + 1.5H₂O. Mp 198-200° dec.

Di-Me ether: 1,2-Dehydrotelobine

$C_{35}H_{32}N_2O_5$ 560.648

Alkaloid from the bark of *Doryphora*

apatela and the stems of *Albertisia papuana* (Monimiaceae, Menispermaceae). Mp 168-172° dec. $[\alpha]_D^{19} +172$ (CHCl₃).

Di-Me ether, dipicrate:

Cryst. + 1.5H₂O. Mp 187-192° dec.

Di-Me ether, N-de-Me: 1,2-Dehydro-2'-nortelobine

C₃₄H₃₀N₂O₅ 546.621

Alkaloid from *Cocculus pendulus*. $[\alpha]_D +100$ (CHCl₃).

Bick, I.R.C. *et al.*, *J.C.S. Perkin 1*, 1972, 2884 (*Dehydrotelobine*)

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1975, 2219 (*1,2-Dehydromicranthine*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1978, **31**, 2077 (*Dehydroapateline*)

Leboeuf, M. *et al.*, *Plant. Med. Phytother.*, 1982, **16**, 280 (*Dehydrotelobine*)

Hussain, S.F. *et al.*, *Tetrahedron*, 1984, **40**, 2513 (*Kurramine, Dehydroapateline*)


Atta-ur-Rahman, *et al.*, *Pure Appl. Chem.*, 1986, **58**, 663 (*1,2-Dehydro-2'-nortelobine*)

Atta-ur-Rahman, *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 802-806 (*Kurramine N-oxides, Dehydroapateline*)

Kutzneride 1

K-116

[878997-78-5]



C₃₇H₄₉Cl₂N₇O₁₂ 854.739

Prod. by the actinomycete *Kutzneria* sp. 744. Moderate antifungal agent. Powder. $[\alpha]_D^{20} -16.7$ (c, 0.33 in MeOH).

2''',3'''-Didehydro: Kutzneride 9

[918664-82-1]

C₃₇H₄₇Cl₂N₇O₁₂ 852.723

Prod. by *Kutzneria* sp. 744. Powder.

4''R-Chloro: Kutzneride 2

[878997-79-6]

C₃₇H₄₈Cl₃N₇O₁₂ 889.184

Prod. by *Kutzneria* sp. 744. Powder.

$[\alpha]_D^{20} -19.4$ (c, 0.29 in MeOH).

3'-Epimer: Kutzneride 3

[878997-80-9]

C₃₇H₄₉Cl₂N₇O₁₂ 854.739

Prod. by *Kutzneria* sp. 744. Powder.

$[\alpha]_D^{20} -30.9$ (c, 0.16 in MeOH).

3'-Epimer, 2''',3'''-didehydro: Kutzneride 4

[878997-81-0]

C₃₇H₄₇Cl₂N₇O₁₂ 852.723

Prod. by *Kutzneria* sp. 744. Powder.

$[\alpha]_D^{20} -17.9$ (c, 0.07 in MeOH).

3'-Epimer, 4''R-chloro: Kutzneride 8

[918664-81-0]

C₃₇H₄₈Cl₃N₇O₁₂ 889.184

Prod. by *Kutzneria* sp. 744. Powder.

3'-Epimer, 4''R-hydroxy, 2''',3'''-didehydro: Kutzneride 6

[918664-79-6]

C₃₇H₄₇Cl₂N₇O₁₃ 868.723

Prod. by *Kutzneria* sp. 744. Powder.

3''''-Demethyl: Kutzneride 7

[918664-80-9]

C₃₆H₄₇Cl₂N₇O₁₂ 840.712

Prod. by *Kutzneria* sp. 744. Powder.

3''''-Demethyl, 3'-epimer: Kutzneride 5

[918664-78-5]

C₃₆H₄₇Cl₂N₇O₁₂ 840.712

Prod. by *Kutzneria* sp. 744. Powder.


Broberg, A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 97-102 (*isol, pmr, cmr*)

Pohanka, A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1776-1781 (*isol, pmr, cmr*)

Kwangsine

K-117

5,5'-Bis(6-hydroxygirinimbine A)



C₃₆H₃₂N₂O₄ 556.66


Alkaloid from *Murraya kwangsiensis*.

Mp 263-273°. $[\alpha]_D^{19} -3.3$ (c, 0.1 in CHCl₃). λ_{max} 222 (log ϵ 4.8); 236 (log ϵ 4.77); 294 (log ϵ 4.77); 335 (log ϵ 4.2) (MeOH).

Xie, F.-Z. *et al.*, *Yaoxue Xuebao*, 2000, **35**, 826-828

Labiataamide A



[174285-78-0]

C₂₉H₄₅NO₈ 535.676Constit. of *Eunicella labiata*. Oil. [α]_D²⁰ +12.8 (c, 0.6 in CHCl₃).**3-Deacetoxy: Labiataamide B**

[174285-79-1]

C₂₇H₄₃NO₆ 477.64Constit. of *Eunicella labiata*. Oil. [α]_D²⁰ -6 (c, 0.2 in CHCl₃).Roussis, V. *et al.*, *Tetrahedron*, 1996, **52**, 2735 (isol, pmr, cmr)**Labuanine A**

10,11-Dihydro-8-hydroxy-9H-benzo[b]pyrido[4,3,2-de][1,10]phenanthroline-9-one

C₁₈H₁₁N₃O₂ 301.304Isol. from the marine sponge *Biemna fortis*. Neuronal differentiation inducer. Amorph. yellow solid. λ_{max} 227 (ε 18000); 288 (ε 7720); 317 (ε 5190); 357 (ε 6070) (MeOH).Aoki, S. *et al.*, *Bioorg. Med. Chem.*, 2003, **11**, 1969-1973 (isol, pmr, cmr)**Laccaic acid A₁**7-[5-[2-(Acetylamino)ethyl]-2-hydroxyphenyl]-9,10-dihydro-3,5,6,8-tetrahydroxy-9,10-dioxo-1,2-anthracenedicarboxylic acid, 9CI. *Laccaic acid A* [15979-35-8]C₂₆H₁₉NO₁₂ 537.436Pigment from stick lac (secretion of the insect *Laccifer lacca*). Food colourant. Red, strongly dichroic platelets (MeOH) or bright red needles (H₂O or MeOH). Does not melt but chars from 230°.*Penta-Me ether, di-Me ester*: Mp 115-116°.**N-De-Ac: Laccaic acid E**

[14597-16-1]

C₂₄H₁₇NO₁₁ 495.398

L-1


Pigment from stick lac. Amorph. red solid. Mp >360°. λ_{max} 291 (log ε 4.5); 360 (sh) (log ε 4.03); 470 (sh) (log ε 3.91); 494 (log ε 3.97); 515 (sh) (log ε 3.88) (MeOH).

[60687-93-6]

Pandhare, E.D. *et al.*, *Tetrahedron, Suppl.*, No. 8, 1966, 229Burwood, R. *et al.*, *J.C.S.(C)*, 1967, 842-851 (isol, uv, pmr, struct)Pandhare, E.D. *et al.*, *Tet. Lett.*, 1967, 2437 (isol, struct)Oka, H. *et al.*, *J. Chromatogr. A*, 1998, **813**, 71-77 (chromatog)Quye, A. *et al.*, *Chem. Rev. (Deddington, U.K.)*, 1999, **8**, 27-31 (rev)**Laccarin**

L-4

3-Acetyl-1,4,5,6,7,7a-hexahydro-7-methyl-2H-pyrrolo[3,2-b]pyridin-2-one [175669-28-0]




Relative Configuration

C₁₀H₁₄N₂O₂ 194.233Alkaloid from fruit bodies of the mushrooms *Lactarius subplinthogalus* and *Laccaria vinaceoavellanea*. Shows moderate phosphodiesterase inhibitory activity. Powder. [α]_D²⁵ +188 (c, 0.47 in CHCl₃). λ_{max} 235 (ε 11220); 293 (ε 11750) (MeOH) (Berdy).**N⁴-(4-Aminobutanoyl): Laccarin A** [766509-22-2]C₁₄H₂₁N₃O₃ 279.338Alkaloid from *Lactarius subplinthogalus*. Powder. [α]_D²⁰ -51.8 (c, 1.05 in MeOH). λ_{max} 235 (log ε 2.5); 239 (log ε 2.5) (MeOH).Matsuda, M. *et al.*, *Heterocycles*, 1996, **43**, 685-690 (isol, uv, ir, pmr, cmr, struct)Wang, Y. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 159-162 (*Laccarin A*)**Lachnanthopyridone**


L-5

5-Hydroxy-7-phenyl-1H-benz[de]isoquinoline-1,6(2H)-dione, 9CI [37011-65-7]

C₁₈H₁₁NO₃ 289.29Poor cryst. (Me₂CO). Mp 338-342°.**N-(1-Carboxy-2-methylbutyl):** [60924-82-5]C₂₄H₂₁NO₅ 403.434Alkaloid from blooms of *Lachnanthes tinctoria*. Rosettes of orange cryst. Mp 218-219°.**N-(2-Hydroxyethyl): N-(2-Hydro-****xyethyl)lachnanthopyridone** [37009-33-9]C₂₀H₁₅NO₄ 333.343Alkaloid from *Lachnanthes tinctoria*. Orange cryst. (EtOAc). Mp 155°.**Me ether, N-butyl: N-Butyl-O-methyl-lachnanthopyridone**C₂₃H₂₁NO₃ 359.424Alkaloid from *Xiphidium caeruleum*.Edwards, J.M. *et al.*, *Tet. Lett.*, 1972, 1631 (isol, struct, deriv)Bazan, A.C. *et al.*, *Phytochemistry*, 1976, **15**, 1413 (isol, synth)Opitz, S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1122-1130 (*N-Butyl-O-methyllachnanthopyridone*)**Lactacystin**

L-6

[133343-34-7]



Absolute Configuration

C₁₅H₂₄N₂O₇S 376.43Prod. by a *Streptomyces* sp. Induces differentiation of neuroblastoma cells. Needles. Sol. H₂O, Py, DMSO, MeOH; poorly sol. EtOAc, hexane, C₆H₆, CHCl₃. Mp 237-238° dec. [α]_D²⁵ +71.3 (c, 0.5 in MeOH). λ_{max} 239 (E1%/1cm 170) (MeOH) (Berdy).


[133398-98-8]

Omura, S. *et al.*, *J. Antibiot.*, 1991, **44**, 113; 117 (isol, pmr, cmr, struct, props)Corey, E.J. *et al.*, *J.A.C.S.*, 1992, **114**, 10677 (synth)Sunazuka, T. *et al.*, *J.A.C.S.*, 1993, **115**, 5302 (synth)Corey, E.J. *et al.*, *Tet. Lett.*, 1993, **34**, 6969; 6973; 6977 (synth)Uno, H. *et al.*, *J.A.C.S.*, 1994, **116**, 2139 (synth)Nakagawa, A. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2411 (biosynth)Nakagawa, A. *et al.*, *Tet. Lett.*, 1994, **35**, 5009 (biosynth)Chida, N. *et al.*, *Chem. Comm.*, 1995, 793 (synth)Takahashi, S. *et al.*, *J. Antibiot.*, 1995, **48**, 1015 (biosynth)Chida, N. *et al.*, *Tetrahedron*, 1997, **53**, 16287-16298 (synth)Panek, J.S. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 1093-1095 (synth)Corey, E.J. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1-10 (rev, synth)Masse, C.E. *et al.*, *Eur. J. Org. Chem.*, 2000, 2513-2528 (rev, synth)Green, M.P. *et al.*, *Tet. Lett.*, 2002, **43**, 6609-6611 (synth)Ooi, H. *et al.*, *J.O.C.*, 2004, **69**, 7765-7768 (synth)Wardrop, D.J. *et al.*, *Chem. Comm.*, 2005, 5106-5108 (synth)Balskus, E.P. *et al.*, *J.A.C.S.*, 2006, **128**, 6810-6812 (synth)Fukuda, N. *et al.*, *J.O.C.*, 2006, **71**, 1220-1225 (synth)Shibasaki, M. *et al.*, *Chem. Asian J.*, 2007, **2**, 20-38 (rev, synth)

Hayes, C.J. *et al.*, *J.O.C.*, 2008, **73**, 2041-2051 (synth)
 Pattenden, G. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 3428-3438 (synth)

Lactimidomycin**L-7**

BM Y 28886. BU 4146T. Antibiotic BM Y 28886. Antibiotic BU 4146T [134869-15-1]



Absolute Configuration

C₂₆H₃₅NO₆ 457.566

Glutarimide-lactone antibiotic. Closely related to Isomigrastatin, I-245. Prod. by *Streptomyces amphibiosporus* ATCC53964. Active against fungi and tumours. Pale yellow solid + 1/4H₂O. Mp 121-125°. [α]_D²⁵ -20 (c, 0.5 in DMSO).

8,9-Dihydro: **8,9-Dihydro-lactimidomycin** [959984-76-0]

C₂₆H₃₇NO₆ 459.581

Prod. by *Streptomyces amphibiosporus* ATCC53964. Oil. [α]_D²⁵ +36.3 (c, 0.12 in CHCl₃).

8,9-Dihydro, 8S-hydroxy: **8,9-Dihydro-8-hydroxylactimidomycin** [910217-36-6]

C₂₆H₃₇NO₇ 475.581


Prod. by *Streptomyces amphibiosporus* ATCC53964. Powder. [α]_D²⁵ +176 (c, 0.3 in CHCl₃).

Sugawara, K. *et al.*, *J. Antibiot.*, 1992, **45**, 1433-1441 (isol, pmr, cmr, props)

Ju, J. *et al.*, *Org. Lett.*, 2007, **9**, 5183-5186 (isol, pmr, cmr)

Lactivicin**L-8**

2-[4-(Acetylamino)-3-oxo-2-isoxazolidinyl]tetrahydro-5-oxo-2-furancarboxylic acid, 9CI. Antibiotic TAN 588. TAN 588 [107167-31-7] [113194-68-6, 113194-69-7]

C₁₀H₁₂N₂O₇ 272.214

Cyclic peptide antibiotic. Prod. by *Empedobacter lactamgensis* and *Lysobacter albus*. Active against gram-positive and -negative bacteria. β-Lactamase inhibitor. [α]_D²⁵ -24.1 (H₂O) (as Na salt). Epimerises readily. λ_{max} 216 (sh) (ε 4050) (H₂O) (Derep).

[106944-73-4, 106944-74-5, 159150-63-7, 105629-02-5, 106962-13-4, 106962-14-5]

Natsugari, H. *et al.*, *Chem. Comm.*, 1987, **62** (synth)


Harada, S. *et al.*, *Tetrahedron*, 1988, **44**, 6589 (isol, pmr, cmr, ir, props)

Nozaki, Y. *et al.*, *J. Antibiot.*, 1989, **42**, 84 (props)

Nakao, Y. *et al.*, *Spec. Publ. - R. Soc. Chem.*, 1989, **70**, 119 (rev)

Lactonomycin

[182234-02-2]



Absolute Configuration

C₂₈H₂₇NO₁₂ 569.521

Prod. by *Streptomyces rishiriensis* MJ773-88K4. Active against gram-positive bacteria. Yellowish powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 168-171°. [α]_D²⁷ +34 (c, 0.27 in MeOH). λ_{max} 228 (ε 11300); 257 (ε 13200); 300 (ε 20000); 395 (ε 6700); 412 (ε 7500) (MeOH). λ_{max} 232 (ε 11600); 250 (ε 12900); 287 (ε 12200); 323 (ε 11300); 367 (ε 10700); 398; 448 (MeOH-NaOH) (Berdy).

4'-Epimer, 3'β-hydroxy: **Lactonomycin Z**

C₂₈H₂₇NO₁₃ 585.52


Prod. by *Streptomyces sanglieri* strain AK 623. Antitumour agent. Yellow powder. Abs. config. not determined. λ_{max} 210 (sh); 229 (sh); 258 (sh); 300; 398 (sh); 415 (sh) (MeOH).

Matsumoto, N. *et al.*, *J. Antibiot.*, 1999, **52**, 269-275; 276-280 (isol, uv, pmr, cmr, abs config)

Höltzel, A. *et al.*, *J. Antibiot.*, 2003, **56**, 1058-1061 (*Lactonomycin Z*)

Lactine**L-10**

6a,7,8,9-Tetrahydro-12-methoxy-6H-benzo[de]-1,3-benzodioxolo[4,5-g]quinolin-11-ol, 9CI. 2-Hydroxy-1-methoxy-10,11-methylenedioxy-noraporphine



(R)-form

C₁₈H₁₇NO₄ 311.337

(R)-form [75638-75-4]

Alkaloid from the bark of *Litsea laeta* (Lauraceae). Mp 296° dec. [α]_D²⁵ -23.7.

Me ether: 1,2-Dimethoxy-10,11-methylenedioxy-noraporphine [α]_D²⁵ -14.8 (c, 0.355 in EtOH).

(S)-form [385366-60-9]

Alkaloid from the trunk bark of *Hernandia nymphaeifolia*. Amorph. powder. [α]_D²⁴ +177 (c, 0.11 in CHCl₃). λ_{max} 224 (log ε 4.37); 271 (log ε 4.07); 307 (log ε 3.71) (EtOH).

(ξ)-form

3-Methoxy, Me ether: 1,2,3-Trimethoxy-10,11-methylenedioxy-noraporphine.

Polygospermine

[69477-64-1]

C₂₀H₂₁NO₅ 355.39**L-9**

Alkaloid from the trunk bark of *Polyalthia oligosperma* (Annonaceae). λ_{max} 222; 276; 306 (sh) (EtOH).


Guineaudeau, H. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 166-172 (*Polygospermine*)

Rastogi, R.C. *et al.*, *Phytochemistry*, 1980, **19**, 998-999 (*R-Laetine*)

Chen, J.-J. *et al.*, *Planta Med.*, 2001, **67**, 593-598 (*S-Laetine*)

Lagerine**L-11**

4'-Demethoxy-6'-hydroxydecaline, 9CI [34532-74-6]

C₂₅H₂₉NO₅ 423.508

Alkaloid from *Lagerstroemia indica* (Lythraceae). Mp 210°. [α]_D -184 (CHCl₃). pK_a 7.3 (EtOH).

Ac:

Cryst. (Et₂O). Mp 217°.

Methiodide:

Cryst. (EtOH). Mp 266°.

Me ether: **Methylagerine**

[33033-88-4]

C₂₆H₃₁NO₅ 437.535

Alkaloid from *Lagerstroemia indica* (Lythraceae). Cryst. by subl. Mp 240°. [α]_D -178 (CHCl₃).

Me ether, N-oxide:


Cryst. (EtOAc). Mp 204-206°.

Ferris, J.P. *et al.*, *J.A.C.S.*, 1971, **93**, 2958 (isol, uv, ir)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2191 (struct, synth, ir, pmr, ms)

Lagunamycin**L-12**

6-Diazo-3-methyl-4-(1,3,5-trimethyl-1-hexenyl)-2,5,7,8(1H,6H)-quinolinetrone, 9CI [150693-65-5]



Absolute Configuration

C₁₉H₂₁N₃O₄ 355.393


Prod. by *Streptomyces* sp. AA0310. 5-Lipoxygenase inhibitor. Yellow needles (MeOH aq.). Mp 134-136° dec. [α]_D²⁶ -33 (c, 0.2 in MeOH). λ_{max} 233 (ε 16300); 264 (sh) (ε 9500); 284 (ε 9100); 352 (ε 25200) (MeOH/NaOH) (Derep). λ_{max} 258 (ε 18300); 302 (ε 15800); 353 (sh) (ε 3400); 420 (ε 1200) (MeOH) (Derep).

▶ VC3215000

Nihei, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 900-907; 1031-1033 (isol, pmr, cmr, struct)

Hosokawa, S. *et al.*, *Tet. Lett.*, 2006, **47**, 6183-6186 (*synth, abs config*)

Lahadinine A L-13
[200053-18-5]



Absolute Configuration

R,R' = -CH₂-

C₂₅H₂₇N₃O₇ 481.504
Alkaloid from *Kopsia pauciflora* (Apocynaceae). Oil. [α]_D -184 (c, 0.1 in CHCl₃). λ_{max} 227 (log ε 4.36); 248 (log ε 3.89); 255 (log ε 3.84); 285 (log ε 2.97); 290 (log ε 2.92) (EtOH).


Kam, T.-S. *et al.*, *Phytochemistry*, 1997, **46**, 785-787 (*isol, uv, ir, pmr, cmr, ms*)

Lahadinine B L-14
[200053-19-6]
As Lahadinine A, L-13 with
R = R' = Me

C₂₆H₃₁N₃O₇ 497.547
Alkaloid from *Kopsia pauciflora* (Apocynaceae). Oil. [α]_D -123 (c, 0.03 in CHCl₃). λ_{max} 222 (log ε 4.39); 253 (log ε 3.79); 284 (log ε 3.06); 289 (log ε 3.03) (EtOH).

Kam, T.-S. *et al.*, *Phytochemistry*, 1997, **46**, 785-787 (*isol, uv, ir, pmr, cmr, ms*)
Magnus, P. *et al.*, *Tetrahedron*, 2002, **58**, 3423-3443 (*synth*)


Lahoramine L-15
8,9-Dimethoxy-13-methyl-1,3-dioxolo[6,7]indeno[2,1-a][3]benzazepinium(1+)
[81189-54-0]
[80787-70-8]



C₂₁H₁₈NO₄[⊕] 348.377
Quaternary alkaloid from *Fumaria parviflora* (Papaveraceae). Obt. as chloride, in amts. too small for crystallisation.


Blaskó, G. *et al.*, *Tet. Lett.*, 1981, **22**, 3127 (*isol, uv, ms, struct*)

Lahoricine L-16
17,19-Epoxy-19,20-dihydroakuamillan, 9CI
[93961-19-4]



C₁₉H₂₂N₂O 294.396
Stereochem. undefined. Alkaloid from the leaves of *Ervatamia coronaria* (Apocynaceae). Amorph. [α]_D +99 (CHCl₃). Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B.*, 1984, **39**, 1289 (*uv, ir, pmr, ms, struct*)


Lahorine L-17
4-Methyl[1,3]dioxolo[4,5-h]-1,3-dioxolo[6,7]indeno[2,1-a][3]benzazepinium
[81189-53-9]
[80787-72-0]



C₂₀H₁₄NO₄[⊕] 332.335
Quaternary alkaloid from *Fumaria parviflora* (Papaveraceae). Yellow cryst. (CHCl₃) (as chloride). Mp 253-255° dec. (chloride). First representative of a new class of isoquinoline alkaloids, the indenobenzazepines.

Blaskó, G. *et al.*, *Tet. Lett.*, 1981, **22**, 3127 (*isol, uv, pmr, ms, struct*)


Lajollamycin L-18



C₃₆H₅₃N₃O₁₀ 687.829
Closely related to Oxazolomycin, O-155. Prod. by the marine-derived *Streptomyces nodosus* (NPS007994). Antibacterial. Bright yellow solid. Mp 92-95°. [α]_D²⁵ +75 (c, 0.00002 in MeOH). λ_{max} 229 (ε 63000); 306 (ε 40000); 375 (ε 22000) (MeOH).

Manam, R.R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 240-243 (*isol, pmr, cmr*)

Lakshminine L-19
6-Amino-5-methoxy-7H-dibenzo[de,h]quinolin-7-one
[479669-27-7]



C₁₇H₁₂N₂O₂ 276.294
Alkaloid from the woody vines of *Sciadotenia toxifera*. Orange-yellow powder. λ_{max} 250 (log ε 3.46); 264 (sh) (log ε 3.13); 350 (log ε 1.96); 400 (sh) (log ε 2.05) (EtOH).

Killmer, L. *et al.*, *J. Nat. Prod.*, 2003, **66**, 115-118 (*isol, pmr, cmr*)

Lallemancine L-20

C₂₃H₂₉N₃O₂ 379.501
Struct. unknown. Alkaloid from *Lallemantia peltata* (Lamiaceae). Mp 191-193°. *Perchlorate*: Mp 221-222°. *N-Ac*: Mp 210-211°.

Platanova, T.F. *et al.*, *Med. Promst. SSSR*, 1962, **16**, 14; *CA*, **57**, 4759h (*isol, ir*)

Lamarckine L-21

Lamarckine
C₁₃H₁₂N₂O₆ 292.248
Struct. unknown. Alkaloid from *Alangium lamarckii* (Alangiaceae). Yellow cryst. Mp 60-62°.

Picrate: Mp 148-150°. *Oxalate*: Mp 73-75°. *Aurichloride*: Mp 255-258°. *Chloroplatinate*: Mp 229-230°.

Basu, N.K. *et al.*, *Indian J. Pharm.*, 1950, **12**, 98-99; *CA*, **44**, 8601b


Lamarckinine[†] L-22

C₂₅H₃₃N₃O₄ 439.553
Struct. unknown. Alkaloid from the bark of *Alangium lamarckii* (Alangiaceae). Mp 265-268° dec.

Khan, N.H. *et al.*, *Pak. J. Biol. Agric. Sci.*, 1965, **8**, 211-213; *CA*, **64**, 9782f

Lambertine L-23

Dihydroberberine
[120834-89-1]
[119420-08-5]



C₂₀H₁₉NO₄ 337.374
Alkaloid from *Berberis lambertii*, *Berberis chitria* and *Berberis vulgaris* (barberry) (Berberidaceae). Yellow needles (MeOH or Me₂CO). Mp 163-164°. λ_{max} 280 (log ε 4.21); 368 (log ε 4.3) (no solvent reported).

N-Me: *N-Methylambertine*. *N-Methyl-dihydroberberine*, 7,8-Dihydro-*N*-methylberberine

[47474-50-0]
[26668-90-6]
C₂₁H₂₂NO₄[⊕] 352.409

Alkaloid from *Berberis heteropoda* (Berberidaceae). Cryst. (as chloride). Mp 210-212° (chloride).

Chatterjee, R. *et al.*, *J. Indian Chem. Soc.*, 1955, **32**, 609 (*isol, uv, struct, synth*)

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 2365-2371 (*deriv, synth*)

Dobhal, M.P. *et al.*, *Pharmazie*, 1988, **43**, 659-660 (*isol, pmr, cmr*)


Yusupov, M.M. *et al.*, *Khim. Prir. Soedin.*, 1990, **26**, 128-129; *Chem. Nat. Compd.* (*Engl. Transl.*), **105**, 106 (*isol*)

Yusupov, M.M. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 53-59; *Chem. Nat. Compd.* (*Engl.*

Transl., 1993, **29**, 43-48
(*Didhydromethylberberine, isol, deriv*)
Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999,
37, 195-202 (*N-15 nmr*)

Lamellarin H**L-24**

[115982-22-4]

C₂₅H₁₅NO₈ 457.395

Authors' numbering scheme shown. Alkaloid from the marine ascidian *Didemnum chartaceum*. Amorph. solid. λ_{\max} 208 (ϵ 31000); 282 (ϵ 24000); 303 (sh); 324 (sh); 340 (ϵ 6000); 370 (ϵ 9000); 389 (ϵ 12000) (MeOH) (Derep). λ_{\max} 218 (ϵ 30000); 244 (sh); 297 (ϵ 20000); 316; 402 (ϵ 13000) (MeOH/NaOH) (Derep). λ_{\max} 205 (ϵ 44000); 298 (ϵ 20000); 405 (ϵ 13000) (MeOH/NaOH) (Berdy).

9,12,21-Tri-Me ether: Lamellarin D

[97614-65-8]

C₂₈H₂₁NO₈ 499.476

Isol. from *Lamellaria* sp. Inhibitor of cell division in fertilised sea urchin eggs. Pale yellow powder. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 218 (ϵ 30000); 244 (sh); 297 (ϵ 20000); 316; 402 (ϵ 13000) (MeOH/NaOH) (Derep). λ_{\max} 208 (ϵ 31000); 282 (ϵ 24000); 303 (sh); 324 (sh); 340 (ϵ 6000); 370 (ϵ 9000); 389 (ϵ 12000) (MeOH) (Derep). λ_{\max} 212; 280; 368; 387 (MeOH) (Berdy).

9,13,21-Tri-Me ether: Lamellarin N

[149379-26-0]

C₂₈H₂₁NO₈ 499.476

Alkaloid from the unidentified *Didemnum* sp. and a *Lamellaria* sp. Needles (MeOH) (as tri-Ac). Mp 283-285° (tri-Ac). λ_{\max} 204 (ϵ 29500); 279 (ϵ 22500); 365 (ϵ 6000); 386 (ϵ 4000) (MeOH) (Berdy). λ_{\max} 204 (ϵ 50000); 295 (ϵ 16800); 402 (ϵ 11300) (MeOH/NaOH) (Berdy).

8,9,12,21-Tetra-Me ether, 20-O-sulfate:**Lamellarin α 20-sulfate**

[229956-47-2]

C₂₉H₂₃NO₁₁S 593.567

Isol. from *Lamellaria* sp. HIV-1 integrase inhibitor. Solid (as Na salt). Mp 145-148° (Na salt). λ_{\max} 204 (ϵ 31500); 276 (ϵ 19500); 303 (ϵ 16000); 324 (sh); 338 (sh); 364 (ϵ 5500); 386 (ϵ 7000) (MeOH).

8,9,13,21-Tetra-Me ether: Lamellarin α

[475232-29-2]

C₂₉H₂₃NO₈ 513.503

Alkaloid from *Didemnum obscurum*. Amorph. solid. Mp 228-230°. λ_{\max} 206 (log ϵ 1.99); 281 (log ϵ 1.91); 302 (log ϵ 1.79); 323 (log ϵ 1.54) (MeOH).

8,9,12,13,21-Penta-Me ether: Lamellarin η η C₃₀H₂₅NO₈ 527.529

Alkaloid from *Didemnum obscurum*. Amorph. solid. Mp 265-269°. λ_{\max} 236 (log ϵ 0.76); 308 (log ϵ 0.55) (MeOH).

5,6-Dihydro, 9-Me ether: Lamellarin S

[181423-71-2]

C₂₆H₁₉NO₈ 473.438

Metab. from an Australian tunicate, *Didemnum* sp. Oil. $[\alpha]_D^{25} +66.5$ (c, 0.84 in MeOH). CAS name is defective. First example of this structural class to show atropisomerism. Abs config. not determined. Slowly racemises.

5,6-Dihydro, 13-Me ether: Lamellarin β

[425381-27-7]

C₂₆H₁₉NO₈ 473.438

Isol. from a *Didemnum* sp. Cytotoxic. Amorph. solid. λ_{\max} 205 (ϵ 43000); 268 (sh); 278 (ϵ 23000); 315 (ϵ 17000); 336 (ϵ 17000) (MeOH).

5,6-Dihydro, 9,20-di-Me ether: Lamellarin Z

[221169-77-3]

C₂₇H₂₁NO₈ 487.465

Alkaloid from *Didemnum chartaceum*. Amorph. solid. λ_{\max} 207 (ϵ 31000); 265 (sh) (ϵ 16000); 276 (ϵ 18000); 314 (ϵ 16000); 333 (ϵ 15000); 348 (sh) (ϵ 14000); 352 (sh) (ϵ 11000) (MeOH).

5,6-Dihydro, 9,12,21-tri-Me ether,**8,13,20-tri-Ac: Lamellarin χ** C₃₄H₂₉NO₁₁ 627.603

Isol. from *Didemnum obscurum*. Amorph. solid. Mp 164-166°. λ_{\max} 208 (log ϵ 0.55); 274 (log ϵ 0.32) (MeOH).

5,6-Dihydro, 8,13,21-tri-Me ether, 20-sulfate:**Lamellarin Y 20-sulfate**

[189084-06-8]

C₂₈H₂₃NO₁₁S 581.556

Alkaloid from an unidentified ascidian. Powder (as Na salt). CAS no. refers to Na salt. λ_{\max} 204 (ϵ 35000); 265 (sh); 273 (ϵ 23000); 309 (ϵ 21000); 328 (sh) (MeOH) (Na salt).

5,6-Dihydro, 9,13,20-tri-Me ether:**Lamellarin G**

[115982-21-3]

C₂₈H₂₃NO₈ 501.492

Alkaloid from *Didemnum chartaceum*. Prisms (DMSO). Mp 263-265°. λ_{\max} 204 (ϵ 55000); 278 (ϵ 12000); 324 (ϵ 12000); 346 (ϵ 8000); 368 (sh) (MeOH/NaOH) (Derep). λ_{\max} 206 (ϵ 46000); 267 (sh); 276 (ϵ 25000); 314 (ϵ 21000); 334 (ϵ 21000) (MeOH) (Derep).

5,6-Dihydro, 9,13,20-tri-Me ether, 8-sulfate:**Lamellarin G 8-sulfate**

[221169-76-2]

C₂₈H₂₃NO₁₁S 581.556

Alkaloid from *Didemnum chartaceum*. Amorph. solid. λ_{\max} 205 (ϵ 46000); 225 (sh) (ϵ 35000); 266 (sh) (ϵ 24000); 275 (ϵ 28000); 301 (sh) (ϵ 20000); 314 (ϵ 27000); 336 (ϵ 22000) (MeOH).

5,6-Dihydro, 9,13,21-tri-Me ether:**Lamellarin L**

[149378-57-4]

C₂₈H₂₃NO₈ 501.492

Alkaloid from an unidentified *Didemnum* sp. Immunomodulator. Amorph.

powder. Mp 285-287°.

5,6-Dihydro, 9,13,21-tri-Me ether, 20-sulfate:**Lamellarin L 20-sulfate**

[221169-75-1]

C₂₈H₂₃NO₁₁S 581.556

Alkaloid from *Didemnum chartaceum*. Amorph. solid. λ_{\max} 203 (ϵ 32000); 221 (sh) (ϵ 21000); 267 (sh) (ϵ 16000); 276 (ϵ 19000); 314 (ϵ 16000); 338 (sh) (ϵ 14000) (MeOH).

5,6-Dihydro, 8,9,13,21-tetra-Me ether:**Lamellarin U**

[189083-79-2]

C₂₉H₂₅NO₈ 515.518

Alkaloid from an unidentified ascidian. Mp 200-204°.

5,6-Dihydro, 8,9,13,21-tetra-Me ether, 20-sulfate:**Lamellarin U 20-sulfate**

[189084-01-3]

C₂₉H₂₅NO₁₁S 595.583

Alkaloid from an unidentified ascidian. Mp 222-226°. λ_{\max} 202 (ϵ 40000); 265 (sh); 273 (ϵ 23500); 309 (ϵ 20000); 327 (sh) (MeOH).

5,6-Dihydro, 9,12,13,21-tetra-Me ether:**Lamellarin J**

[149355-76-0]

C₂₉H₂₅NO₈ 515.518

Alkaloid from the unidentified *Didemnum* sp. and a *Lamellaria* sp. Immunomodulator. Amorph. powder. Mp 216-220°.

5,6-Dihydro, 8,9,12,13,21-penta-Me ether:**Lamellarin F**

[115982-20-2]

C₃₀H₂₇NO₈ 529.545

Alkaloid from *Didemnum chartaceum*. Prisms (DMSO). Mp 273-276° dec. λ_{\max} 205 (ϵ 64000); 265 (sh); 288 (ϵ 28000); 323 (ϵ 23000) (MeOH/NaOH) (Derep). λ_{\max} 208 (ϵ 48000); 268 (sh); 278 (ϵ 38000); 310 (ϵ 23000); 330 (sh) (MeOH) (Derep). λ_{\max} 210 (ϵ 47000); 278 (ϵ 21000); 312 (ϵ 28000) (MeOH) (Berdy). λ_{\max} 207 (ϵ 52000); 290 (ϵ 28000); 322 (ϵ 31000) (MeOH/NaOH) (Berdy).

5,6-Dihydro, 5 ζ ,7-dihydroxy, 7,8,9,12,20-penta-Me ether:**Lamellarin A**

[97614-62-5]

C₃₀H₂₇NO₁₀ 561.544

Metab. of *Lamellaria* sp. MDR inhibitor, immunomodulator. Pale yellow prisms (MeOH). Mp 168-172° dec. Consists of a 1:1 mixt. of geom. isomers generated by restricted rotn. around the pyrrole-aryl bond. Opt. inactive, prob. due to rapid interconversion at C-5 through an open-chain tautomer. λ_{\max} 218 (ϵ 46000); 287 (ϵ 33000); 319 (ϵ 36000); 378 (ϵ 9000) (MeOH/NaOH) (Derep). λ_{\max} 215 (ϵ 41000); 275 (ϵ 33000); 309 (ϵ 28000); 325 (sh); 326 (ϵ 25000) (MeOH) (Derep).

7-Hydroxy, 8,9,12,21-tetra-Me ether:**Lamellarin M**

[149378-58-5]


C₂₉H₂₃NO₉ 529.502

Alkaloid from the unidentified *Didemnum* sp. and a *Lamellaria* sp. Immunomodulator. Amorph. powder.

- 7-Hydroxy, 8,9,13,21-tetra-Me ether:
Lamellarin X
[189083-94-1]
C₂₉H₂₃NO₉ 529.502
Alkaloid from an unidentified ascidian. Powder. λ_{\max} 204 (€ 30000); 276 (€ 15000); 303 (€ 14000); 324 (sh); 338 (sh); 365 (sh) (MeOH).
- 7-Hydroxy, 7,8,9,12,21-penta-Me ether:
Lamellarin B
[97614-63-6]
C₃₀H₂₅NO₉ 543.529
Isol. from the marine probranch mollusc *Lamellaria* sp. Dehydration prod. of Lamellarin A. Immunomodulator. Pale yellow needles (MeOH). Mp 258-259°. λ_{\max} 218 (€ 30000); 244 (sh); 297 (€ 20000); 316; 402 (€ 13000) (MeOH/NaOH) (Derep). λ_{\max} 208 (€ 31000); 282 (€ 24000); 303 (sh); 324 (sh); 340 (€ 6000); 370 (€ 9000); 389 (€ 12000) (MeOH) (Derep).
- 7-Hydroxy, 7,8,9,12,21-penta-Me ether, 20-sulfate: **Lamellarin B 20-sulfate**
[221169-73-9]
C₃₀H₂₅NO₁₂S 623.593
Alkaloid from *Didemnum chartaceum*. Amorph. solid. λ_{\max} 202 (€ 31000); 224 (sh) (€ 18000); 283 (€ 27000); 301 (sh) (€ 21000); 321 (sh) (€ 11000); 365 (€ 8000); 386 (€ 10000) (MeOH).
- 7-Hydroxy, 8,9,12,13,21-penta-Me ether:
Lamellarin ε
C₃₀H₂₅NO₉ 543.529
Alkaloid from *Didemnum obscurum*. Amorph. solid. Mp 271-275°. λ_{\max} 209 (log € 2.61); 277 (log € 2.37); 304 (log € 2.33); 325 (log € 2.1); 367 (log € 1.69) (MeOH).
- 7-Hydroxy, 5,6-dihydro, 8,9,12,21-tetra-Me ether: **Lamellarin K**
[149378-56-3]
C₂₉H₂₅NO₉ 531.518
Alkaloid from an unidentified *Didemnum* sp. Immunomodulator. Amorph. powder. Mp 196-198°. λ_{\max} 203 (€ 40000); 278 (€ 23000); 312 (€ 20000); 337 (€ 17000) (MeOH) (Berdly).
- 7-Hydroxy, 5,6-dihydro, 8,9,13,21-tetra-Me ether: **Lamellarin E**
[115982-19-9]
C₂₉H₂₅NO₉ 531.518
Alkaloid from *Didemnum chartaceum*. Prisms (MeOH). Mp 228-232°. λ_{\max} 205 (€ 64000); 265 (sh); 288 (€ 28000); 323 (€ 23000) (MeOH/NaOH) (Derep). λ_{\max} 208 (€ 48000); 268 (sh); 278 (€ 38000); 310 (€ 23000); 330 (sh) (MeOH) (Derep).
- 7-Hydroxy, 5,6-dihydro, 7,8,9,12,21-penta-Me ether: **Lamellarin C. Dihydrolamellarin B**
[97614-64-7]
C₃₀H₂₇NO₉ 545.545
Metab. of a *Lamellaria* sp. Needles (MeOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 225-230°. λ_{\max} 218 (€ 46000); 287 (€ 33000); 319 (€ 36000); 378 (€ 9000) (MeOH/NaOH) (Derep). λ_{\max} 215 (€ 41000); 275 (€ 33000); 309 (€ 28000); 325 (sh); 326 (€ 25000) (MeOH) (Derep).
- 7-Hydroxy, 5,6-dihydro, 7,8,9,12,21-penta-Me ether, 20-sulfate: **Lamellarin C 20-sulfate**
[221169-74-0]
C₃₀H₂₇NO₁₂S 625.609
Alkaloid from *Didemnum chartaceum*. Amorph. solid. λ_{\max} 205 (€ 45000); 225 (sh) (€ 31000); 265 (sh) (€ 21000); 276 (€ 25000); 310 (€ 22000); 337 (sh) (17000) (MeOH).
- 7-Hydroxy, 5,6-dihydro, 7,8,9,12,13,21-hexa-Me ether: **Lamellarin I**
[149355-75-9]
C₃₁H₂₉NO₉ 559.571
Alkaloid from an unidentified *Didemnum* sp. MDR inhibitor. Off-white irregular prisms (MeOH). Mp 218-220°.
- 7-Methoxy, 8,12,21-tri-Me ether, 9,13,20-tri-Ac: **Lamellarin φ**
C₃₅H₂₉NO₁₂ 655.614
Isol. from *Didemnum obscurum*. Amorph. solid. Mp 276-279°. λ_{\max} 224 (log € 0.44); 298 (log € 0.32) (MeOH).
- 7-Methoxy, 8,9,13,21-tetra-Me ether: **Lamellarin W**
[189083-85-0]
C₃₀H₂₅NO₉ 543.529
Alkaloid from an unidentified ascidian. Pale yellow solid. Mp 224-228°. λ_{\max} 205 (€ 30000); 279 (€ 15000); 302 (€ 13000); 336 (sh); 364 (€ 5000); 385 (€ 6000) (MeOH).
- 7-Methoxy, 8,9,12,13,21-penta-Me ether: **Lamellarin ζ**
C₃₁H₂₇NO₉ 557.556
Alkaloid from *Didemnum obscurum*. Amorph. solid. Mp 268-272°. λ_{\max} 230 (log € 0.36); 404 (log € 0.23) (MeOH).
- 7-Methoxy, 5,6-dihydro, 8,9,13,21-tetra-Me ether: **Lamellarin T**
[189083-78-1]
C₃₀H₂₇NO₉ 545.545
Alkaloid from an unidentified ascidian. Powder. Mp 214-218°.
- 7-Methoxy, 5,6-dihydro, 8,9,13,21-tetra-Me ether, 20-sulfate: **Lamellarin T 20-sulfate**
[189084-00-2]
C₃₀H₂₇NO₁₂S 625.609
Alkaloid from an unidentified ascidian. Powder (as Na salt). CAS no. refers to Na salt. λ_{\max} 204 (€ 47000); 265 (sh) (€ 19350); 274 (€ 24000); 309 (€ 21000); 330 (sh) (€ 16600) (MeOH) (Na salt).
- 7-Methoxy, 5,6-dihydro, 5ξ-hydroxy, 8,9,13,21-tetra-Me ether: **Lamellarin V**
[189083-80-5]
C₃₀H₂₇NO₁₀ 561.544
Alkaloid from an unidentified ascidian. Powder. Mp 162-166°.
- 7-Methoxy, 5,6-dihydro, 5ξ-hydroxy, 8,9,13,21-tetra-Me ether, 20-sulfate: **Lamellarin V 20-sulfate**
[189084-02-4]
C₃₀H₂₇NO₁₃S 641.608
Alkaloid from an unidentified ascidian. Mp 174-178°. λ_{\max} 202 (€ 29000); 264 (sh); 273 (€ 17000); 304 (€ 14000); 328 (sh) (MeOH).
- 14-Methoxy, 13-deoxy, 7-hydroxy, 5,6-dihydro, 8,9,12,21-tetra-Me ether: **Lamellarin γ**
C₃₀H₂₇NO₉ 545.545
Alkaloid from *Didemnum obscurum*. Amorph. solid. Mp 163-165°. λ_{\max} 206 (log € 1.88); 278 (log € 1.65); 310 (log € 1.63); 331 (log € 1.55); 365 (log € 0.87) (MeOH).
- Andersen, R.J. *et al.*, *J.A.C.S.*, 1985, **107**, 5492-5495 (*Lamellarins A-D*)
Lindquist, N. *et al.*, *J.O.C.*, 1988, **53**, 4570-4574 (*Lamellarins E-H*)
Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1993, **46**, 489-501 (*Lamellarins I-N*)
Urban, S. *et al.*, *Aust. J. Chem.*, 1996, **49**, 711-713 (*Lamellarin S*)
Heim, A. *et al.*, *Angew. Chem., Int. Ed.*, 1997, **36**, 155-156 (*Lamellarin G, synth*)
Banwell, M. *et al.*, *Chem. Comm.*, 1997, 2259-2260 (*Lamellarin K, synth*)
Pat. Coop. Treaty (WIPO), 1997, ((*Pharma Mar*))97 01336; *CA*, **1327**, 166474 (*activity*)
Reddy, M.V.R. *et al.*, *Tetrahedron*, 1997, **53**, 3457-3466 (*Lamellarins T-Y*)
Ishibashi, F. *et al.*, *Tetrahedron*, 1997, **53**, 5951-5962 (*synth*)
Reddy, M.V.R. *et al.*, *J. Med. Chem.*, 1999, **42**, 1901-1907 (*Lamellarin α 20-sulfate*)
Davis, R.A. *et al.*, *J. Nat. Prod.*, 1999, 419-424 (*Didemnum chartaceum constiis*)
Steglich, W. *et al.*, *Chem. Eur. J.*, 2000, **6**, 1147-1152 (*Lamellarin L, synth*)
Ruchirawat, S. *et al.*, *Tet. Lett.*, 2001, **42**, 1205-1208 (*Lamellarin G, synth*)
Ham, J. *et al.*, *Bull. Korean Chem. Soc.*, 2002, **23**, 163-166 (*Lamellarin β*)
Iwao, M. *et al.*, *Tet. Lett.*, 2003, **44**, 4443-4446 (*Lamellarin G, synth*)
Bailly, C. *et al.*, *Curr. Med. Chem. Anticancer Agents*, 2004, **4**, 363-378 (*rev*)
Krishnaiah, P. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1168-1171 (*Lamellarins α, γ, ε*)
Handy, S.T. *et al.*, *J.O.C.*, 2004, **69**, 2362-2366 (*Lamellarin G, synth*)
Reddy, S.M. *et al.*, *Tetrahedron*, 2005, **61**, 9242-9247 (*Lamellarins ζ, η, φ, χ*)
Ploypradith, P. *et al.*, *J.O.C.*, 2006, **71**, 9440-9448 (*synth*)
Peschko, C. *et al.*, *Synthesis*, 2006, 3048-3057 (*Lamellarin G, Lamellarin K, synth*)
Yamaguchi, T. *et al.*, *Tet. Lett.*, 2006, **47**, 3755-3757 (*Lamellarin α 20-sulfate, synth*)
Fujikawa, N. *et al.*, *Tetrahedron*, 2006, **62**, 594-604 (*Lamellarins D,L,N, synth*)
Fan, H. *et al.*, *Chem. Rev.*, 2008, **108**, 264-287 (*rev*)
Liermann, J.C. *et al.*, *J.O.C.*, 2008, **73**, 4526-4531 (*synth*)
Baunbaeck, D. *et al.*, *Mar. Drugs*, 2008, **6**, 5134-527 (*sar, props*)

Lamellarin Q L-25

Methyl 3,4-bis(4-hydroxyphenyl)-1H-pyrrole-2-carboxylate
[168010-02-4]



C₁₈H₁₅NO₄ 309.321

Isol. from the sponge *Dendrilla cactos*. Pale yellow oil. λ_{\max} 240 (ϵ 14000); 287 (ϵ 12000) (EtOH).

N-(4-Hydroxyphenyl): **Lamellarin R**
[168010-03-5]

$C_{24}H_{19}NO_5$ 401.418

Isol. from the sponge *Dendrilla cactos*. Green oil. λ_{\max} 229 (ϵ 13500); 283 (ϵ 6000); 336 (ϵ 2200); 418 (ϵ 1200) (EtOH).

N-[2-(4-Methoxyphenyl)-2-oxoethyl]:

Lamellarin O

[158402-61-0]

$C_{27}H_{23}NO_6$ 457.482

Isol. from the Australian marine sponge *Dendrilla cactos*. Unstable pale yellow oil; pale yellow cryst. (as di-Me ether). Mp 55-58° (di-Me ether). λ_{\max} 277 (ϵ 5200) (EtOH) (Berdy).

N-[2-(2-Hydroxy-4-methoxyphenyl)-2-oxoethyl]: **Lamellarin P**

[158402-60-9]

$C_{27}H_{23}NO_7$ 473.481

From *Dendrilla cactos*. Moderately stable pale yellow oil. λ_{\max} 231 (ϵ 5800); 276 (ϵ 6200); 305 (ϵ 3000) (EtOH) (Berdy).

Urban, S. et al., *Aust. J. Chem.*, 1994, **47**, 1919-1924; 1995, **48**, 1491-1494 (*isol, pmr, cmr, ms, struct, uv, ir*)

Fürstner, A. et al., *J.O.C.*, 1995, **60**, 6637

(*synth*)

Banwell, M.G. et al., *Chem. Comm.*, 1997, 207

(*synth*)

Boger, D.L. et al., *J.A.C.S.*, 1999, **121**, 54-62

(*synth*)


Marfil, M. et al., *Tetrahedron*, 2004, **60**, 8659-8668 (*synth*)

Fukuda, T. et al., *Tetrahedron*, 2008, **64**, 328-338 (*synth*)

Lamprolobine

L-26

1-[(Octahydro-2H-quinolizin-1-yl)-methyl]-2,6-piperidinedione, 9CI. 5-(Glutarimidomethyl)quinolizidine [18688-40-9]



Absolute Configuration

$C_{15}H_{24}N_2O_2$ 264.367

(+)-form

Alkaloid from *Lamprolobium fruticosum* and *Lupinus holosericeus* (Fabaceae). Oil, cryst. at 0°. $[\alpha]_D^{20}$ +29 (c, 1.5 in EtOH).

Picrate: Mp 153-154°.

9 β -Hydroxy: **9 β -Hydroxylamprolobine**

[80324-07-8]

$C_{15}H_{24}N_2O_3$ 280.366

Alkaloid from the leaves of *Sophora velutina* var. *zimbabwensis* (Fabaceae). $[\alpha]_D^{20}$ +24 (EtOH).

6-Epimer: **Epilamprolobine**

[80374-24-9]

$C_{15}H_{24}N_2O_2$ 264.367

Alkaloid from leaves of *Sophora tomentosa* (Fabaceae). Needles (C_6H_6 /hexane). Mp 101.5°. $[\alpha]_{365}^{20}$ -13.8 (EtOH).

6-Epimer, 1 β -oxide: **Epilamprolobine N-oxide**

$C_{15}H_{24}N_2O_3$ 280.366

Alkaloid from leaves, stems, seeds and pods of *Sophora tomentosa* (Fabaceae). Amorph. solid. $[\alpha]_D^{20}$ +14.9 (c, 0.77 in EtOH).

Deoxo, 6-epimer: **17-Deoxyepilamprolobine**. 17-Desoxy-cis-lamprolobine

[220908-80-5]

$C_{15}H_{26}N_2O$ 250.383

Constit. of *Bongardia chrysogonum*. Amorph. solid. $[\alpha]_D^{22}$ -70 (c, 0.9 in MeOH). λ_{\max} 203 (log ϵ 2.67) (MeOH).

(±)-form

Noncryst. Mp 192-193° (as picrate).

6-Epimer:

Cryst. (hexane/EtOAc). Mp 68-68.5°.

Hart, N.K. et al., *Aust. J. Chem.*, 1968, **21**,

1619 (*isol, ir, ms, struct*)

Yamada, Y. et al., *Agric. Biol. Chem.*, 1970, **34**, 1536 (*synth*)

Goldberg, S.I. et al., *J.O.C.*, 1970, **35**, 242

(*synth*)

Wenkert, E. et al., *J.O.C.*, 1970, **35**, 515 (*synth*)

Keller, W.J. et al., *Phytochemistry*, 1980, **19**,

2233 (*isol*)

Keller, W.J. et al., *J. Nat. Prod.*, 1981, **44**, 357

(*isol*)

Murakoshi, I. et al., *Phytochemistry*, 1981, **20**,

1725 (*Epilamprolobine, Epilamprolobine*

oxide)

Asres, K. et al., *J. Nat. Prod.*, 1986, **49**, 117

(*isol, ir, pmr, cmr, ms, struct, 9-*

Hydroxylamprolobine)

Michael, J.P. et al., *Tetrahedron*, 1992, **48**,

10211 (*synth, Lamprolobine,*

Epilamprolobine)

Atta-ur-Rahman, et al., *Nat. Prod. Lett.*, 1998,

12, 161-173 (*Deoxyepilamprolobine*)

Lanceine

L-27

[1358-85-6]

Struct. unknown. MF = $C_{20-24}H_{26-36}N_2O_{3-4}$. Alkaloid from *Catharanthus lancea* (Apocynaceae). Mp 198°. $[\alpha]_D^{20}$ +64 (c, 0.5 in EtOH).

Janot, M.M. et al., *Ann. Pharm. Fr.*, 1957, **15**,

474-478 (*isol, ir, uv*)


Blomster, R.N. et al., *J. Pharm. Sci.*, 1967, **56**,

284-286 (*isol, ir*)

Lanceomigine

L-28

Methyl 2,17-epoxy-2-hydroxy-1-methyl-1,17-cyclo-1,2(1H,2H)-secoakuammilan-16-carboxylate, 9CI [79659-63-5]



Absolute Configuration

$C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from *Hunteria congolana*, *Hunteria zeylanica* and *Alstonia lanceolata* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ +32 (c, 1 in $CHCl_3$).

N⁴-Oxide: **Lanceomigine N⁴-oxide**

[79659-64-6]

$C_{22}H_{26}N_2O_5$ 398.458

Alkaloid from the stem bark of *Alstonia lanceolata* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ +79 (c, 0.63 in $CHCl_3$).

N-De-Me: **Rhazicine**

[93772-10-2]

$C_{21}H_{24}N_2O_4$ 368.432

Alkaloid from fresh leaves of *Rhazya stricta* and from leaves of *Melodinus acutiflorus* (Apocynaceae). Prisms (MeOH) or hygroscopic needles. Mp 122.6-124.4°. $[\alpha]_D^{20}$ +69.8 (c, 0.76 in $CHCl_3$) (+65). Struct. revised in 1987.

N-De-Me, N⁴-oxide: **Rhazicine N-oxide**

[117138-43-9]

$C_{21}H_{24}N_2O_5$ 384.431

Alkaloid from leaves of *Melodinus acutiflorus* (Apocynaceae). Needles (C_6H_6 /EtOH). Mp 188-188.1°. $[\alpha]_D^{22}$ +120.4 (c, 0.467 in $CHCl_3$).

Vercauteren, J. et al., *Phytochemistry*, 1981, **20**,

1411 (*isol, uv, ir, pmr, cmr, ms, struct, oxide*)

Atta-ur-Rahman, et al., *Heterocycles*, 1984,

22, 2183 (*Rhazicine*)

Wen-Lan, H. et al., *Phytochemistry*, 1987, **26**,

2625 (*Rhazicine, isol, uv, ir, pmr, cmr, cryst*

struct)


Hu, W.-L. et al., *Planta Med.*, 1988, **54**, 235

(*Rhazicine N-oxide*)

Lanciferine

L-29

[56440-64-3]



Absolute Configuration

$C_{31}H_{32}N_2O_7$ 544.603

Alkaloid from the aerial parts of *Alstonia lanceolifera* and *Alstonia boullindaensis* (Apocynaceae). Cryst. (EtOH). Mp 259-262°. $[\alpha]_D^{20}$ -60 (c, 1 in $CHCl_3$).

10-Hydroxy: **10-Hydroxylanciferine**

[56486-42-1]

$C_{31}H_{32}N_2O_8$ 560.602

Alkaloid from the aerial parts of *Alstonia lanceolifera* and *Alstonia boullindaensis* (Apocynaceae). Cryst. (E_2O /MeOH). Mp 228-231°. $[\alpha]_D^{20}$ -48 (c, 1 in $CHCl_3$).

10-Methoxy: **10-Methoxylanciferine**

[56440-65-4]

$C_{32}H_{34}N_2O_8$ 574.629

Alkaloid from the aerial parts of *Alstonia lanceolifera* and *Alstonia boullindaensis* (Apocynaceae). Cryst. (MeOH). Mp 251-254°. $[\alpha]_D^{20}$ -64 (c, 1

in CHCl_3).


Lewin, G. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1975, **280**, 987 (isol, struct, derivs)

Lewin, G. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 1096 (isol, uv, ir, pmr, ms, derivs)

Lewin, G. *et al.*, *Bull. Soc. Chim. Fr.*, Part 2, 1980, 400 (struct)

Lanitine L-30

3-(Dimethylamino)-2-hydroxypregn-5-en-18,20-olide
[30452-63-2]



$\text{C}_{23}\text{H}_{35}\text{NO}_3$ 373.534

Alkaloid from stem bark of *Kibatalia gitingensis* (Apocynaceae). Mp 184-186°.


2,3-Diepimer: **Lanitinine**
[30452-64-3]

Alkaloid from the stem bark of *Kibatalia gitingensis* (Apocynaceae). Mp 223-225°.

Bernal-Santos, R.M. *et al.*, *Philipp. J. Sci.*, 1967, **96**, 411; *CA*, **76**, 10364p

Lankanescine L-31

12-Ethyl-1,2,7b,8,12b,12c-hexahydro-2-methoxy-2-methylpyrido[4,3-b]pyrrolo[1,2,3-lm]carbazole, 9CI
[367938-59-8]




$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}$ 320.433

Poss. related to the Ellipticine and Olivacine groups; biosynth. not yet known. Alkaloid from *Rauwolfia canescens*. λ_{max} 260 ; 279 (MeOH).

Arambewela, L.S.R. *et al.*, *Pharm. Biol.*, 2001, **39**, 239-240 (isol, pmr, cmr, ms)

Lankongensisine A L-32




$\text{C}_{18}\text{H}_{27}\text{NO}_5$ 337.415

Alkaloid from the roots of *Ligularia lankongensis*. Yellow oil. $[\alpha]_{\text{D}}^{24}$ +48.4 (c, 4.8 in CHCl_3).

Tan, A.M. *et al.*, *Chin. Chem. Lett.*, 2004, **15**, 68-70 (isol, pmr, cmr)

Lankongensisine B L-33




$\text{C}_{18}\text{H}_{27}\text{NO}_5$ 337.415

Alkaloid from the roots of *Ligularia lankongensis*. Yellow oil. $[\alpha]_{\text{D}}^{20}$ +76.7 (c, 1.2 in CHCl_3).

Tan, A.M. *et al.*, *Chin. Chem. Lett.*, 2004, **15**, 68-70 (isol, pmr, cmr)

Lannotinidine A L-34

[851461-36-4]



Relative Configuration


$\text{C}_{17}\text{H}_{25}\text{NO}_3$ 291.389

Alkaloid from *Lycopodium annotinum*. Amorph. solid. $[\alpha]_{\text{D}}^{23}$ +47 (c, 1 in MeOH).

Koyama, K. *et al.*, *Tetrahedron*, 2005, **61**, 3681-3690 (isol, pmr, cmr)

Lannotinidine E L-35

[851461-40-0]



$\text{C}_{16}\text{H}_{23}\text{NO}_4$ 293.362

Alkaloid from *Lycopodium annotinum* var. *acrifolium*. Amorph. solid. $[\alpha]_{\text{D}}^{18}$ -7 (c, 0.2 in MeOH).

10-Deoxy, 9,10-didehydro, 11-ketone:

Lannotinidine F

[851461-41-1]


$\text{C}_{16}\text{H}_{19}\text{NO}_3$ 273.331

Alkaloid from *Lycopodium annotinum*. Amorph. solid. $[\alpha]_{\text{D}}^{24}$ -22 (c, 1 in MeOH).

Koyama, K. *et al.*, *Tetrahedron*, 2005, **61**, 3681-3690 (isol, pmr, cmr)

Lannotinidine G L-36

[851461-42-2]



$\text{C}_{16}\text{H}_{21}\text{NO}_2$ 259.347

Alkaloid from *Lycopodium annotinum*. Amorph. solid. $[\alpha]_{\text{D}}^{18}$ -2 (c, 1 in MeOH).

Koyama, K. *et al.*, *Tetrahedron*, 2005, **61**, 3681-3690 (isol, pmr, cmr)

Lanopylins L-37



Lanopylin	R
A ₁	R = $-(\text{CH}_2)_{13}\text{CH}(\text{CH}_3)_2$
B ₁	R = $-(\text{CH}_2)_{15}\text{CH}_3$
C ₁	R = $-(\text{CH}_2)_{11}\text{CH}(\text{CH}_3)_2$
D ₁	R = $-(\text{CH}_2)_{13}\text{CH}_3$
E ₁	R = $-(\text{CH}_2)_{12}\text{CH}(\text{CH}_3)_2$
F ₁	R = $-(\text{CH}_2)_{14}\text{CH}_3$
G ₁	R = $-(\text{CH}_2)_{14}\text{CH}(\text{CH}_3)_2$
G ₂	R = $-(\text{CH}_2)_{10}\text{CH}=\text{CH}(\text{CH}_2)_4\text{CH}(\text{CH}_3)_2$
H ₁	R = $-(\text{CH}_2)_{16}\text{CH}_3$
H ₂	R = $-(\text{CH}_2)_{10}\text{CH}=\text{CH}(\text{CH}_2)_6\text{CH}_3$
I ₁	R = $-(\text{CH}_2)_{13}\text{CH}(\text{CH}_3)_2$
I ₂	R = $-(\text{CH}_2)_{10}\text{CH}=\text{CH}(\text{CH}_2)_5\text{CH}(\text{CH}_3)_2$
J ₁	R = $-(\text{CH}_2)_{17}\text{CH}_3$
J ₂	R = $-(\text{CH}_2)_{10}\text{CH}=\text{CH}(\text{CH}_2)_7\text{CH}_3$

Structs. given are tentative except for Lanopylins A₁ and B₁. Prod. by *Streptomyces* sp. K99-5041. Lanosterol synthase inhibitors.

Lanopylin A₁

2-Methyl-3-(15-methylhexadecylidene)-1-pyrroline

$\text{C}_{22}\text{H}_{41}\text{N}$ 319.573

Oil. λ_{max} 235 (ε 14500) (MeOH).

12',13'-Didehydro(Z)-: 2-Methyl-3-(15-methyl-12Z-hexadecenylidene)-1-pyrroline. **Lanopylin C₂**

$\text{C}_{22}\text{H}_{39}\text{N}$ 317.557

Lanopylin B₁

3-Heptadecylidene-2-methyl-1-pyrroline

$\text{C}_{22}\text{H}_{41}\text{N}$ 319.573

Oil. λ_{max} 235 (ε 13900) (MeOH).

12',13'-Didehydro(Z)-: 3-(12Z-Heptadecenylidene)-2-methyl-1-pyrroline. **Lanopylin D₂**

$\text{C}_{22}\text{H}_{39}\text{N}$ 317.557

Lanopylin C₁

2-Methyl-3-(13-methyltetradecylidene)-1-pyrroline

$\text{C}_{20}\text{H}_{37}\text{N}$ 291.519

Lanopylin D₁

2-Methyl-3-pentadecylidene-1-pyrroline

$\text{C}_{20}\text{H}_{37}\text{N}$ 291.519

Lanopylin E₁

2-Methyl-3-(14-methylpentadecylidene)-1-pyrroline

$\text{C}_{21}\text{H}_{39}\text{N}$ 305.546

Lanopylin F₁

3-Hexadecylidene-2-methyl-1-pyrroline

$\text{C}_{21}\text{H}_{39}\text{N}$ 305.546

Lanopylin G₁

2-Methyl-3-(16-methylheptadecylidene)-1-pyrroline

$\text{C}_{23}\text{H}_{43}\text{N}$ 333.599

12',13'-Didehydro(E-): 2-Methyl-3-(16-methyl-12Z-heptadecenylidene)-1-pyrroline. **Lanopylin E₂**
C₂₃H₄₁N 331.584

Lanopylin G₂

2-Methyl-3-(18-methyl-12Z-nonadecenylidene)-1-pyrroline
C₂₅H₄₅N 359.637

Lanopylin H₁

2-Methyl-3-octadecylidene-1-pyrroline
C₂₃H₄₃N 333.599

12',13'-Didehydro(Z-): 2-Methyl-3-(12Z-octadecenylidene)-1-pyrroline. **Lanopylin F₂**
C₂₃H₄₁N 331.584

Lanopylin H₂

3-(12Z-Eicosenylidene)-2-methyl-1-pyrroline
C₂₅H₄₅N 359.637

Lanopylin I₁

2-Methyl-3-(17-methyloctadecylidene)-1-pyrroline
C₂₄H₄₅N 347.626

12',13'-Didehydro(Z-): 2-Methyl-3-(17-methyl-12Z-octadecenylidene)-1-pyrroline. **Lanopylin A₂**
C₂₄H₄₃N 345.61
Oil.

Lanopylin I₂

2-Methyl-3-(19-methyl-12Z-eicosenylidene)-1-pyrroline
C₂₆H₄₇N 373.664

Lanopylin J₁

2-Methyl-3-nonadecylidene-1-pyrroline
C₂₄H₄₅N 347.626

12',13'-Didehydro(Z-): 2-Methyl-3-(12Z-nonadecenylidene)-1-pyrroline. **Lanopylin B₂**
C₂₄H₄₃N 345.61
Oil.

Lanopylin J₂

3-(12Z-Heneicosenylidene)-2-methyl-1-pyrroline
C₂₆H₄₇N 373.664

Sakano, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 817-826 (isol, pmr, cmr, ms)
Snider, B.B. *et al.*, *J.O.C.*, 2005, **70**, 1087-1088 (Lanopylin B₁, synth)

Lansamide 1**L-38**

N-Methyl-3-phenyl-N-(2-phenylethenyl)-2-propenamide, 9CI. N-Methyl-N-styrylcinnamamide. *Lansamide 1*
PhCH=CHNMeCOCH=CHPh
C₁₈H₁₇NO 263.338

(E,E)-form [77527-97-0]

Alkaloid from the leaves and seeds of *Clausena lansium* (wampee). Mp 124-125° (119-120°). λ_{max} 218 ; 290 ; 306 ; 327 (MeOH).

(Z,E)-form

Lansiamamide B. N-Methyl-N-cis-styrylcinnamamide
[121817-37-6]

Alkaloid from the seeds of *Clausena lansium* (wampee). Yellowish plates (Et₂O). Mp 72-73°. λ_{max} 218 ; 226 ; 285 (MeOH).

N-De-Me: Lansiamamide A

[121817-36-5]
C₁₇H₁₅NO 249.312

Constit. of the seeds of *Clausena lansium* (wampee). Pale yellow needles (cyclohexane/Et₂O). Mp 121-123°. λ_{max} 218 ; 265 ; 303 (MeOH).

Prakash, D. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 1075 (isol, uv, ir, pmr, ms, struct, synth)


Lin, J.-H. *et al.*, *Phytochemistry*, 1989, **28**, 621-622 (isol, pmr, ms)

Stefanutti, I. *et al.*, *Tet. Lett.*, 2000, **41**, 3735-3738 (synth)

Huang, X.-S. *et al.*, *Acta Cryst. E*, 2006, **62**, 1987-1988 (Lansiamamide B, cryst struct)

Lansamide 2

Lansimide 2
[119574-06-0]



Relative Configurations

C₁₈H₃₆N₂O₅ 360.493


A 1:1 complex of cyclic amides Lansamide 3 in D-575 and ζ-Clausenamide, C-501. Constit. of *Clausena lansium* (wampee). Spasmodic agent. Cryst. (MeOH/C₆H₆). Mp 173-174°. Racemic. λ_{max} 229 ; 254 (MeOH).

Ji, X. *et al.*, *Acta Cryst. C*, 1992, **48**, 1082-1085 (conform, cryst struct)

Lakshmi, V. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 355-358 (isol, pmr, cmr, cryst struct)

Lanuginosine**L-40**

10-Methoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. 9-Methoxy-1,2-methylenedioxy-7-oxodibenzo[de,g]quinoline. *Oxoxylidine*. 9-Methoxylyridodine
[23740-25-2]



C₁₈H₁₁NO₄ 305.289

Alkaloid from a variety of genera in the Annonaceae (*Annona*, *Enantia*, *Guatteria*, *Polyalthia*, *Xylopia*), Magnoliaceae (*Magnolia*, *Michelia*, *Liriodendron*, *Talauma*) and Menispermaceae (*Stephania*). Shows antimicrobial activity. Mp

319-321°.

▶RB6035500**O-De-Me: Oxanolobine**

[76788-85-7]

C₁₇H₉NO₄ 291.262

Alkaloid from stem bark of *Guatteria melosma* (Annonaceae). Amorph. orange solid (MeOH). Mp 270-275° dec.

Talapatra, S.K. *et al.*, *Chem. Ind. (London)*, 1969, 1056 (isol, uv, ir)

Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1970, **8**, 475 (synth)

Kupchan, S.M. *et al.*, *J.O.C.*, 1970, **35**, 1682 (isol, pmr, ms, uv, ir)

Talapatra, S.K. *et al.*, *Tetrahedron*, 1975, **31**, 1105 (isol, uv, ir, ms, pmr)


Phoebe, C.H. *et al.*, *Heterocycles*, 1980, **14**, 1977 (Oxanolobine)

Villar, A. *et al.*, *Planta Med.*, 1986, **52**, 556-557 (activity)

Zhang, Z. *et al.*, *J. Nat. Prod.*, 2002, **65**, 856-859 (cmr)

Lapatin A**L-41**

[859152-45-7]



Absolute Configuration


C₂₃H₁₉N₅O₃ 413.435

Prod. by *Penicillium lapatayae* (IBT 10870). [α]_D²² +22 (c, 1.1 in EtOH). λ_{max} 206 (log ε 4.43); 237 (log ε 4.05); 255 (log ε 3.96); 257 (log ε 3.97); 306 (log ε 3.5) (EtOH).

Larsen, T.O. *et al.*, *J. Nat. Prod.*, 2005, **68**, 871-874 (isol, uv, cd, pmr, cmr)

Lapatin B**L-42**

[859152-47-9]



Absolute Configuration

C₂₀H₁₄N₄O₃ 358.356

Prod. by *Penicillium lapatayae* (IBT 10870). [α]_D²² -20 (c, 1.6 in EtOH). λ_{max} 255 (log ε 3.36); 265 (log ε 3.38); 277 (log ε 3.31); 302 (log ε 3.02); 314 (log ε 2.9) (EtOH).


Larsen, T.O. *et al.*, *J. Nat. Prod.*, 2005, **68**, 871-874 (isol, uv, cd, pmr, cmr)

Leca, D. *et al.*, *J.O.C.*, 2007, **72**, 4284-4287 (synth)

Walker, S.J. *et al.*, *Tet. Lett.*, 2007, **48**, 6214-6216 (synth)

Lapidilectine A

[143070-42-2]

C₂₄H₂₈N₂O₆ 440.495

N⁴-C21:C-20-C21 Bissecoaspidofractines. Alkaloid from bark and leaves of *Kopsia lapidilecta* (Apocynaceae). Amorph. [α]_D -33 (c, 0.8 in CHCl₃).

14,15-Dihydro, 15α-hydroxy: Epilapidi-lectinol

[150881-26-8]

C₂₄H₃₀N₂O₇ 458.51

Alkaloid from bark and leaves of *Kopsia lapidilecta* (Apocynaceae). Amorph. [α]_D +19 (c, 0.8 in CHCl₃).

14,15-Dihydro, 15β-hydroxy: Lapidilectinol

[151003-94-0]

C₂₄H₃₀N₂O₇ 458.51

Alkaloid from bark and leaves of *Kopsia lapidilecta* (Apocynaceae). Amorph. [α]_D -5 (c, 0.5 in CHCl₃).

3-Oxo: Lapidilectam

[150881-25-7]

C₂₄H₂₆N₂O₇ 454.479

Alkaloid from bark and leaves of *Kopsia lapidilecta* (Apocynaceae). [α]_D +77 (c, 0.55 in CHCl₃).

16-Epimer: Isolapidilectine A

[151003-93-9]

C₂₄H₂₈N₂O₆ 440.495


Alkaloid from bark and leaves of *Kopsia lapidilecta* (Apocynaceae). Amorph. [α]_D +54 (c, 0.72 in CHCl₃).

Awang, K. *et al.*, *Tet. Lett.*, 1992, **33**, 2493 (isol, uv, pmr, cmr, ms, struct)

Awang, K. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1134 (config)

Lapidilectine B

[143052-16-8]

C₂₁H₂₂N₂O₄ 366.416

Aspidofractine alkaloids showing N⁴-C-21:C-20-C21 cleavage and elimination of C-21. Alkaloid from bark and leaves of *Kopsia lapidilecta*. Amorph. [α]_D +7.6 (c, 0.9 in CHCl₃).

10-Methoxy: Tenuisine A

[185741-73-5]

C₂₂H₂₄N₂O₅ 396.442

Alkaloid from the leaves of *Kopsia tenuis*. [α]_D +77 (c, 0.8 in CHCl₃). Struct. revised in 2004. λ_{max} 206 (log ε 4.46); 247 (log ε 3.99); 304 (log ε 3.64)

L-43

(EtOH).

10-Methoxy, 3-oxo: Tenuisine C

[185742-51-2]

C₂₂H₂₂N₂O₆ 410.426

Alkaloid from the leaves of *Kopsia tenuis*. [α]_D +87 (c, 0.06 in CHCl₃). Struct. revised in 2004. λ_{max} 203 (log ε 4.41); 247 (log ε 4.1); 304 (log ε 3.55) (EtOH).

10,11-Dimethoxy: Tenuisine B

[185742-22-7]

C₂₃H₂₆N₂O₆ 426.468

Alkaloid from the leaves of *Kopsia tenuis*. Struct. revised in 2004. [α]_D +51 (c, 0.09 in CHCl₃). λ_{max} 214 (log ε 4.48); 252 (log ε 4.08); 300 (log ε 3.7) (EtOH).

Awang, K. *et al.*, *Tet. Lett.*, 1992, **33**, 2493-2496 (isol, uv, pmr, cmr, struct)


Pearson, W.H. *et al.*, *J.O.C.*, 2004, **69**, 9109-9122 (synth)

Kam, T.-S. *et al.*, *Tetrahedron*, 2004, **60**, 10739-10745 (Tenuisines)

Pihko, A.J. *et al.*, *Tetrahedron*, 2005, **61**, 8769-8807 (synth, rev)

Largazole

[1009815-87-5]

C₂₉H₄₂N₄O₅S₃ 622.873

Isol. from a *Symplocos* sp. Histone deacetylase inhibitor. Cytotoxic. Potent antiproliferative agent. Amorph. solid. [α]_D²⁰ +22 (c, 0.1 in MeOH).

Taori, K. *et al.*, *J.A.C.S.*, 2008, **130**, 1806-1807; 13506 (isol, pmr, cmr, ms)

Ying, Y. *et al.*, *J.A.C.S.*, 2008, **130**, 8455-8459 (synth)

Bowers, A. *et al.*, *J.A.C.S.*, 2008, **130**, 11219-11222 (synth)


Nasveschuk, C.G. *et al.*, *Org. Lett.*, 2008, **10**, 3595-3598 (synth)

Ghosh, A.K. *et al.*, *Org. Lett.*, 2008, **10**, 3907-3909 (synth)

Larutensine

Larutensine

[138615-19-7]

C₁₉H₂₂N₂O 294.396

Alkaloid from the bark and stems of *Kopsia larutensis* and bark of *Kopsia terenganensis* (Apocynaceae). Amorph. [α]_D +22 (c, 0.17 in CHCl₃).

L-45

Awang, K. *et al.*, *Phytochemistry*, 1991, **30**, 3164 (isol, uv, pmr, cmr, struct)

Kam, T.-S. *et al.*, *Phytochemistry*, 1992, **31**, 2936 (isol, uv, pmr, cmr, ms, struct)


Lounasmaa, M. *et al.*, *Heterocycles*, 1993, **36**, 751 (synth)

Uzir, S. *et al.*, *Tet. Lett.*, 1997, **38**, 1571 (isol)

Lasiocarpine

[303-34-4]

L-47

C₂₁H₃₃NO₇ 411.494

Diester of heliotridine with angelic and lasiocarpic acids. Major alkaloid from *Heliotropium europaeum*, *Heliotropium lasiocarpum*, *Heliotropium arbainense*, *Heliotropium curassavicum*, *Heliotropium eichwaldii*, *Heliotropium indicum* and other *Heliotropium* spp. (Boraginaceae). Antineoplastic agent. Hepatotoxin, causes liver damage in grazing animals. Shows activity against hepatoma and antifungal activity. Plates (petrol). Mp 95.5-97°. [α]_D¹⁶ -3.5 (c, 2.0 in EtOH). [α]_D¹⁵ +0.9 (c, 6.3 in CHCl₃). Log P 0.46 (calc).

► Highly toxic. Possible human carcinogen (IARC 2B). OE7875000

N-Oxide: Lasiocarpine N-oxide

[127-30-0]

C₂₁H₃₃NO₈ 427.494

From *Heliotropium europaeum* (Boraginaceae). Shows antifungal and antibacterial activity. Prisms or needles. Mp 134-135° dec. (variable). [α]_D¹⁷ +13.1 (c, 4.97 in EtOH).

► OE7900000

O^{3'}-Ac: Acetylasiocarpine

[57538-10-0]

C₂₃H₃₅NO₈ 453.531

From *Heliotropium europaeum* (Boraginaceae). Noncryst. [α]_D¹⁹ -0.9 (c, 2 in EtOH).

O^{3'}-Ac, N-oxide: 3'-Acetylasiocarpine N-oxideC₂₃H₃₅NO₉ 469.531

Alkaloid from aerial parts of *Heliotropium hirsutissimum* (Boraginaceae). Gummy solid. [α]_D²⁰ +10 (c, 0.27 in CHCl₃). Named 4'-Acetylasiocarpine N-oxide in the lit.

3'-Deoxy: 7-Angelyheliotrine

[56317-17-0]

C₂₁H₃₃NO₆ 395.495

Minor alkaloid from *Heliotropium eichwaldii* (Boraginaceae). [α]_D¹⁶ -13.77 (c, 0.8 in EtOH). Ester of heliotridine with angelic and heliotric acids.


3'-Deoxy, picrate: Mp 191-193°.

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1954, **7**, 277; 287; 1975, **28**, 2319 (struct, Acetylasiocarpine)

- Mattocks, A.R. *et al.*, *Nature (London)*, 1968, **217**, 723 (*tox*)
 Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*wv*)
 Pedersen, E. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 249 (*ms*)
 Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (*cd*)
 Suri, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 505 (*7-Angelylheliotrine*)
 Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417 (*cmr*)
 Hay, D.G. *et al.*, *Acta Cryst. B*, 1982, **38**, 155 (*cryst struct*)
 Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)
 Marquina, G. *et al.*, *Pharmazie*, 1989, **44**, 870-871 (*activity*)
 Constantinidis, T. *et al.*, *Phytochemistry*, 1993, **32**, 1335 (*Acetyllassiocarpine N-oxide*)
IARC Monog. (Web),
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LBG000

Lasiodine A†

[22326-90-5]


L-48 $C_{39}H_{49}N_5O_7$ 699.845

A seco-ansapeptide alkaloid. Alkaloid from the leaves of *Lasiodiscus marmoratus* (Rhamnaceae). Cryst. (Me₂CO). Mp 195°. $[\alpha]_D^{25} +38$ (CHCl₃). Contains a D-phenylalanine residue.

- Marchand, J. *et al.*, *Tetrahedron*, 1969, **25**, 937 (*isol, uv, ir, pmr, ms, struct*)
 País, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869 (*cmr*)

Lasiodine B

[22400-56-2]

L-49

Absolute Configuration


 $C_{35}H_{47}N_5O_5$ 617.787

Alkaloid from the leaves of *Lasiodiscus marmoratus* (Rhamnaceae). Cryst. + 0.5H₂O (Me₂CO). Mp 221°. $[\alpha]_D^{25} -301$ (c, 1 in CHCl₃/MeOH 1:1).

- Marchand, J. *et al.*, *Tetrahedron*, 1969, **25**, 937 (*isol, uv, ir, pmr, ms, struct*)
 Marchand, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1971, 3742; 1972, 4699 (*stereochem*)
 González Sierra, M. *et al.*, *Chem. Comm.*, 1972, 915 (*stereochem*)
 País, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869 (*cmr*)

Lassiocarpine

[125287-01-6]

L-51 $C_{29}H_{39}NO_6$ 497.63

Alkaloid from roots of *Aconitum kojimae* var. *lassiocarpium* and *Aconitum fukuto-mei* (Ranunculaceae). Cryst. (Me₂CO). Mp 141-143°. $[\alpha]_D^{25} -17.4$ (c, 0.49 in MeOH).


11-Epimer, O-debenzoyl: 11-Epi-16,17-dihydroxylepenine. 16,17-Dihydroxy-11-epilepenine $C_{22}H_{35}NO_5$ 393.522

Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrium*. Powder. Mp 205-207°. $[\alpha]_D^{20} -62$ (c, 0.1 in MeOH). The semitrivial names are not strictly correct.

- Takayama, H. *et al.*, *Tet. Lett.*, 1989, **30**, 3441 (*isol, ir, pmr, cmr, struct*)
 Takayama, H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 936 (*isol*)
 Zhang, F. *et al.*, *Planta Med.*, 2005, **71**, 1073-1076 (*11-Epi-16,17-dihydroxylepenine*)

Lastourvilline**L-52**

5,6,6a,7-Tetrahydro-9,10-dimethoxy-6-methyl-4H-dibenzo[de,g]quinoline-1,2-diol, 9CI. 1,2-Dihydroxy-9,10-dimethoxyaporphine


 $C_{19}H_{21}NO_4$ 327.379**(S)-form** [70518-70-6]

Alkaloid from the bark of *Artabotrys lastourvillensis* (Annonaceae). Large pale-yellow cryst. (CH₂Cl₂/MeOH). Mp 128-136° dec. $[\alpha]_D^{25} +120$ (c, 0.5 in EtOH). Crystals turn violet when exposed to air.

- Eloumi-Ropivia, J. *et al.*, *J. Nat. Prod.*, 1985, **48**, 460 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lasubine I**L-53**

4-(3,4-Dimethoxyphenyl)octahydro-2H-quinolizin-2-ol, 9CI. 2-Hydroxy-4-(3,4-dimethoxyphenyl)quinolizidine
 [68622-77-5]
 [88931-04-8 ((±)-form)]



Absolute Configuration

 $C_{17}H_{25}NO_3$ 291.389

The abs. configs. of all the alkaloids do not appear to have been firmly established, but are prob. as given here. Alkaloid from *Lagerstroemia subcostata* (Lythraceae), also obt. by basic hydrolysis of Subcosine I. Mp 120.5-122°. $[\alpha]_D^{25} -8.8$ (c, 0.34 in MeOH).

O-(3,4-Dimethoxycinnamoyl): Subcosine I

[68622-78-6]

[88931-06-0 ((±)-form)]

 $C_{28}H_{35}NO_6$ 481.588

Minor alkaloid from leaves of *Lagerstroemia subcostata* (Lythraceae). Oil. $[\alpha]_D^{25} +68$ (c, 0.20 in MeOH).

O^{3'}-De-Me: Demethylasubine I

[59373-46-5]

 $C_{16}H_{23}NO_3$ 277.363

Minor alkaloid from young seedlings of *Heimia salicifolia* (Lythraceae), absent from older plants. Cryst. (MeOH/Et₂O). Mp 193-194°. Equilib. mixt. of conformers in soln.

O^{3'}-De-Me, 2-O-(4-hydroxycinnamoyl):**5-Epidemethoxyabresoline**

[66512-90-1]

 $C_{25}H_{29}NO_5$ 423.508

Minor alkaloid from young seedlings of *Heimia salicifolia* (Lythraceae). Equilib. mixt. of conformers in DMSO soln.

2-Ketone, O^{3'}-de-Me: 1,3,4,6,7,8,9,9a-Octahydro-4-(3-hydroxy-4-methoxyphenyl)-2-quinolizidinone $C_{16}H_{21}NO_3$ 275.347

Trace alkaloid from young seedlings of *Heimia salicifolia* (Lythraceae). Stereochem. not determined.

9a-Epimer: Lasubine II

[68681-73-2]

[89771-49-3 ((±)-form)]

 $C_{17}H_{25}NO_3$ 291.389

Alkaloid from *Lagerstroemia subcostata* (Lythraceae), also obt. by basic hydrolysis of Subcosine II. Oil. $[\alpha]_D^{25} -34.7$ (c, 0.32 in MeOH).

9a-Epimer, O-(3,4-dimethoxycinnamoyl):**Subcosine II**

[68681-74-3]

 $C_{28}H_{35}NO_6$ 481.588

Minor alkaloid from leaves of *Lagerstroemia subcostata* (Lythraceae). Oil. $[\alpha]_D^{25} +85.3$ (c, 0.64 in MeOH) (natur-

al). $[\alpha]_D^{25} +121$ (c, 3.5 in MeOH) (synthetic).

9a-Epimer, O³-de-Me: **Demethylasubine II**

[52656-92-5]
C₁₆H₂₃NO₃ 277.363

Minor alkaloid from young seedlings of *Heimia salicifolia* (Lythraceae), absent from older plants. Cryst. (EtOAc). Mp 94-95°.

9a-Epimer, O³-de-Me, 2-O-(4-hydroxycinnamoyl): **Demethoxyabresoline**

[66408-51-3]
C₂₅H₂₉NO₅ 423.508

Minor alkaloid from young seedlings of *Heimia salicifolia* (Lythraceae). Noncryst. solid.

9a-Epimer, O³-de-Me, 2-O-(4-hydroxy-3-methoxycinnamoyl): **Abresoline**

[53778-14-6]
C₂₆H₃₁NO₆ 453.534

Rel. config. only known. Minor alkaloid from *Heimia salicifolia* (Lythraceae). Amorph. solid.

Rother, A. *et al.*, *J. Nat. Prod.*, 1975, **38**, 477 (*Demethylasubines, ketone O-de-Me*)

Hörhammer, R.B. *et al.*, *J.O.C.*, 1975, **40**, 656 (*Abresoline*)

Fuji, K. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 2515 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Rother, A. *et al.*, *Phytochemistry*, 1978, **17**, 305 (*Demethoxyabresolines*)

Quick, J. *et al.*, *Org. Prep. Proced. Int.*, 1979, **11**, 111 (*synth, deriv*)

Takano, S. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3892 (*synth, ir, pmr*)

Iida, H. *et al.*, *J.O.C.*, 1984, **49**, 1909 (*synth, ir, pmr, cmr, ms*)

Hoffmann, R.W. *et al.*, *Annalen*, 1986, 1823 (*synth, pmr, cmr, Lasubine II*)

Narasaka, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 525 (*synth*)

Ent, H. *et al.*, *Heterocycles*, 1988, **27**, 237 (*synth, ir, pmr, ms*)

Brown, J.D. *et al.*, *J.A.C.S.*, 1988, **110**, 7445 (*synth, ir, pmr, cmr, Lasubine II*)

Comins, D.L. *et al.*, *J.O.C.*, 1992, **57**, 5807 (*synth*)

Pilli, R.A. *et al.*, *J.O.C.*, 1995, **60**, 717 (*Lasubine II, synth*)

Bardot, V. *et al.*, *Heterocycles*, 1998, **48**, 507-518 (*synth, pmr, cmr*)

Chalard, P. *et al.*, *Tetrahedron: Asymmetry*, 1998, **9**, 4361-4368 (*synth*)

Ratni, H. *et al.*, *Org. Lett.*, 1999, **1**, 1997-1999 (*synth*)

Ukaji, Y. *et al.*, *Heterocycles*, 2000, **52**, 563-566 (*Lasubine II, synth*)

Davis, F.A. *et al.*, *Org. Lett.*, 2003, **5**, 3855-3857 (*synth*)

Gracias, V. *et al.*, *Org. Lett.*, 2003, **5**, 4999-5001 (*Lasubine II, synth*)

Zaja, M. *et al.*, *Tetrahedron*, 2004, **60**, 9629-9634 (*Lasubine II, synth*)

Back, T.G. *et al.*, *J.O.C.*, 2005, **70**, 967-972 (*Lasubine II, synth*)

Atobe, M. *et al.*, *Tet. Lett.*, 2005, **46**, 2669-2673 (*Abresoline, synth*)

Yu, R.T. *et al.*, *J.A.C.S.*, 2006, **128**, 12370-12371 (*Lasubine II, synth*)


Liu, S. *et al.*, *Tet. Lett.*, 2006, **47**, 7681-7684 (*synth*)

Mancheño, O.G. *et al.*, *J.O.C.*, 2007, **72**, 10294-10297 (*synth*)

Lim, J. *et al.*, *Tet. Lett.*, 2008, **49**, 88-89 (*synth*)

Latifine

1,2,3,4-Tetrahydro-4-(4-hydroxyphenyl)-6-methoxy-2-methyl-5-isoquinolinol, 9CI. 1,2,3,4-Tetrahydro-5-hydroxy-4-(4-hydroxyphenyl)-6-methoxy-2-methylisoquinoline



C₁₇H₁₉NO₃ 285.342

(R)-form

Synthetic. Mp 213-217° dec. $[\alpha]_D +9.94$ (c, 0.644 in MeOH). The observed specific rotn. of the synthetic material is unexpectedly lower than that of the natural product even after repeated recrystallization.

(S)-form [93915-33-4]

Alkaloid from the leaves and bulbs of *Crinum latifolium* (Liliaceae). Cubes (MeOH). Mp 215-217°. $[\alpha]_D^{20} -27.9$ (c, 0.32 in MeOH).

Di-Me ether:

Oil. $[\alpha]_D^{25} +2.7$ (c, 0.37 in MeOH).

(±)-form

Synthetic. Cryst. (EtOH). Mp 212-215° dec.

Di-Me ether:

Red cryst. (CH₂Cl₂/hexane). Mp 87-88°.

Takano, S. *et al.*, *J.C.S. Perkin 1*, 1985, 2447 (*synth*)

Kobayashi, S. *et al.*, *J. Chem. Res., Synop.*, 1986, 280 (*isol, ir, pmr, cmr, ms, ord, cryst struct, synth*)


Katakawa, J. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3928 (*synth*)

Gore, V.G. *et al.*, *J.C.S. Perkin 1*, 1988, 481 (*synth, ir, pmr, deriv*)

Couture, A. *et al.*, *J.C.S. Perkin 1*, 1999, 789-794 (*synth*)

Latifolian A

L-55



Relative Configuration

C₂₄H₂₂NO₆⁺ 420.441

Quaternary alkaloid from the stem bark of *Gnetum latifolium*. JNK3 kinase inhibitor. Solid (as TFA salt). $[\alpha]_D^{26} -33$ (c, 0.21 in MeOH) (TFA salt). λ_{max} 206 (ε 37200); 290 (ε 6400) (MeOH) (TFA salt).

4'-Me ether: Latifolian B

C₂₅H₂₄NO₆⁺ 434.468

Quaternary alkaloid from the stem

L-54


bark of *Gnetum latifolium*. JNK3 kinase inhibitor. Solid (as TFA salt). $[\alpha]_D^{26} -12$ (c, 0.14 in MeOH) (TFA salt). λ_{max} 206 (ε 35480); 289 (ε 5260) (MeOH) (TFA salt).

Rochfort, S.J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1080-1082 (*isol, pmr, cmr*)

Latifoline†

L-56

Con-5-enin-3-ol, 9CI. *Conan-5-en-3-ol* [4438-06-6]



C₂₂H₃₅NO 329.525

Minor alkaloid from the bark of *Funtumia latifolia* (Apocynaceae). Mp 129°. $[\alpha]_D -4$ (c, 0.8 in CHCl₃).

▶ GK7621300

Ac: Mp 160°. $[\alpha]_D -15$ (c, 1.2 in CHCl₃).

N-De-Me: Norlatifoline

[6704-71-8]

C₂₁H₃₃NO 315.498

Alkaloid from the bark of *Funtumia latifolia* (Apocynaceae). Cryst. (Me₂CO or by subl.). Mp 192°. $[\alpha]_D -26$ (c, 1 in CHCl₃).

N-De-Me, O,N-di-Ac: Mp 124°. $[\alpha]_D +12$ (c, 2 in CHCl₃).

12β-Hydroxy: Funtuline

C₂₂H₃₅NO₂ 345.524

Alkaloid from the bark of *Funtumia latifolia* (Apocynaceae). Cryst. (MeOH). Mp 235°. $[\alpha]_D -6$ (CHCl₃/MeOH 4:1).

12β-Hydroxy, O,O-di-Ac:

C₂₆H₃₉NO₄ 429.598

Mp 237°. $[\alpha]_D -15$ (c, 1 in CHCl₃).

Janot, M.M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1962, **254**, 1326 (*isol, struct, synth*)


Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2486 (*Norlatifoline*)

Nagata, W. *et al.*, *Tet. Lett.*, 1963, 869 (*synth*)

Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 787 (*Funtuline*)

Latifolinine

L-57



C₂₂H₃₃NO 327.509

Alkaloid from *Funtumia latifolia* (Apocynaceae). Oil.


Oxime:

C₂₂H₃₄N₂O 342.523


Cryst. (MeOH). Mp 225°. $[\alpha]_D +190$ (MeOH).

18-Oxo: Malouetamide

[50906-87-1]

C₂₂H₃₁NO₂ 341.492Alkaloid from *Malouetia heudelotii* (Apocynaceae). Mp 182°. [α]_D²⁵ +144 (CHCl₃).Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2486 (Latifolinine)Khuong-Huu, F. *et al.*, *Phytochemistry*, 1973, **12**, 1813-1816 (*Malouetamide*)**Latindine****L-58**C₄₁H₃₉N₃O₁₀ 733.773Not indexed by CA. Alkaloid from *Crinum latifolium* (Amaryllidaceae). Mp >300° dec.Ghosal, S. *et al.*, *Phytochemistry*, 1985, **24**, 2141-2156 (rev)**Latonduine A****L-59**

[572873-52-0]



R = H

C₁₀H₇Br₂N₅O 373.006Alkaloid from *Stylissa carteri*. Pale yellow solid. Dec. at 290°. λ_{max} 245 (ε 19830); 284 (ε 14220) (MeOH).Linington, R.G. *et al.*, *Org. Lett.*, 2003, **5**, 2735-2738 (*isol, synth, pmr, cmr*)**Latonduine B****L-60**


[572873-51-9]

As Latonduine A, L-59 with

R = -COOH

C₁₁H₇Br₂N₅O₃ 417.016Alkaloid from *Stylissa carteri*. Pale yellow cryst. (as Et ester). Mp 228-231° (Et ester). λ_{max} 203 (ε 24130); 252 (ε 25920); 284 (sh); 348 (ε 5890) (MeOH).Linington, R.G. *et al.*, *Org. Lett.*, 2003, **5**, 2735-2738 (*isol, pmr, cmr*)**Latrunculin A†****L-61**

[76343-93-6]



Absolute Configuration

C₂₂H₃₁NO₅S 421.557Toxin from the Red Sea sponge *Latrunculia magnifica* and *Latrunculia corticata*. Also found in three Pacific sponges, *Spongia mycofijiensis*, *Hyatella* sp., an unidentified specimen, and in *Chromodoris hamiltoni*, *Chromodoris lochii* and *Chromodoris elisabethina*. Ichthyotoxin, inhibitor of protein kinase C, shows cytochalasin-like activity, potential anti-glaucoma agent. Oil. Poorly sol. H₂O. [α]_D²⁴ +152 (c, 1.2 in CHCl₃). λ_{max} 218 (ε 23500); 268 (sh) (ε) (MeOH) (Derep).*Me ether:*Cryst. (C₆H₆). Mp 164-165°. [α]_D²⁴ +315 (c, 0.33 in CHCl₃).*N-(Hydroxymethyl): Latrunculin G*


[122876-69-1]

C₂₃H₃₃NO₆S 451.583Isol. from *Latrunculia magnifica* plus formaldehyde. Cell division inhibitor. Oil. [α]_D²⁵ +70 (c, 2.2 in CHCl₃).*6,7-Epoxyde: 6,7-Epoxylatrunculin A*

[122876-48-6]

C₂₂H₃₁NO₆S 437.556Constit. of *Latrunculia magnifica* and *Spongia* sp. Oil. Sol. MeOH. [α]_D²⁵ +37 (c, 1.2 in CHCl₃). λ_{max} 218 (ε 23500); 268 (sh) (ε) (MeOH) (Derep).Kashman, Y. *et al.*, *Tet. Lett.*, 1980, **21**, 3629-3632 (*isol, uv, pmr, cmr, ms, cryst struct*)Groweiss, A. *et al.*, *J.O.C.*, 1983, **48**, 3512-3516 (*Latrunculin A, cryst struct*)Spector, I. *et al.*, *Science (Washington, D.C.)*, 1983, **219**, 493 (rev)Okuda, R.K. *et al.*, *Experientia*, 1985, **41**, 1355-1356Kashman, Y. *et al.*, *Tetrahedron*, 1985, **41**, 1905 (*abs config*)Blasberger, D. *et al.*, *Annalen*, 1989, 1171 (*epoxyde*)U.S. Pat., 1989, 4 857 538; CA, **112**, 30647q (*Latrunculin G*)White, J.D. *et al.*, *J.O.C.*, 1992, **57**, 5292 (*synth, bibl*)Jefford, C.W. *et al.*, *Tet. Lett.*, 1996, **37**, 159 (*abs config, cryst struct, bibl*)Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 115-134; 135-149 (*synth, bibl*)**Latrunculin B†****L-62**

[76343-94-7]



Absolute Configuration

C₂₀H₂₉NO₅S 395.519Isol. from the Red Sea sponge *Latrunculia magnifica*; also from *Latrunculia corticata*, *Spongia* sp., *Glossodoris quadricolor* and *Chromodoris hamiltonii*. Strongly ichthyotoxic; actin polym. inhibitor; shows insecticidal props. [α]_D²⁴ +112 (c, 0.48 in CHCl₃). λ_{max} 212 (ε

17300); 269 (sh) (ε) (MeOH) (Derep).

17300); 269 (sh) (MeOH) (Derep).

Me ether: 15-O-Methylatrunculin B


[105917-28-0]

C₂₁H₃₁NO₅S 409.546Isol. from *Negombata magnifica*. [α]_D²⁵ +108 (c, 1.5 in CHCl₃).*N-Ac: N-Acetylpatrunculin B*C₂₂H₃₁NO₆S 437.556Isol. from *Negombata magnifica*. Oil. [α]_D²⁵ +48 (c, 0.05 in EtOH).*N-(Hydroxymethyl): Latrunculin H*

[122876-74-8]

C₂₁H₃₁NO₆S 425.545Isol. from *Latrunculia magnifica* plus formaldehyde. Cell division inhibitor. Oil.*16-Epimer: 16-Epilatrunculin B*C₂₀H₂₉NO₅S 395.519Isol. from the sponge *Negombata magnifica*.Kashman, Y. *et al.*, *Tet. Lett.*, 1980, **21**, 3629 (*isol, uv, ir, pmr, cmr, ms, struct*)Groweiss, A. *et al.*, *J.O.C.*, 1983, **48**, 3512 (*isol, uv, ir, pmr, cmr, ms, struct*)Spector, I. *et al.*, *Science (Washington, D.C.)*, 1983, **219**, 493 (rev)Mebs, D. *et al.*, *J. Chem. Ecol.*, 1985, **11**, 713 (*isol*)Kashman, Y. *et al.*, *Tetrahedron*, 1985, **41**, 1905 (*pmr, cmr*)U.S. Pat., 1989, 4 857 538; CA, **112**, 30647q (*Latrunculin H*)Hoye, T.R. *et al.*, *J.A.C.S.*, 2002, **124**, 7405-7410 (*16-Epilatrunculin B*)Vilozny, B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1055-1057 (*15-O-Methylatrunculin B*)El Sayed, K.A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 219-223 (*N-Acetylpatrunculin B*)Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 115-134 (*synth, bibl*)**Latrunculin C****L-63**

[76376-32-4]

C₂₀H₃₁NO₅S 397.535Macrolide. Structs. originally reported for Latrunculins C and D were reversed. Isol. from the Red Sea Sponge *Latrunculia magnifica*. Ichthyotoxin. Oil. λ_{max} 212 (ε 17300); 269 (sh) (ε) (MeOH) (Derep).*15-Ketone, 11-Me ether: Latrunculin D*


[98155-17-0]

C₂₁H₃₁NO₅S 409.546From *Latrunculia magnifica*. Ichthyotoxin. Oil. Sol. MeOH, hexane; poorly sol. H₂O. λ_{max} 212 (ε 17300); 269 (sh) (ε) (MeOH) (Derep).Kashman, Y. *et al.*, *Tetrahedron*, 1985, **41**, 1905-1914 (*isol, uv, ir, pmr, cmr, ms, struct*)

Smith, A.B. *et al.*, *J.A.C.S.*, 1992, **114**, 2995-3007 (*synth*)
 Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 115-134 (*synth*)

Latrunculin M

[122876-49-7]


C₂₁H₃₃NO₅S 411.561

Minor constit. of the Red Sea sponge *Latrunculia magnifica*. Oil. Sol. MeOH. [α]_D²⁵ +11 (c, 0.2 in CHCl₃).

Blasberger, D. *et al.*, *Annalen*, 1989, 1171 (*isol*, *ir*, *pmr*, *cmr*, *ms*)
 Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 115-134 (*synth*)

Latrunculin S

[175992-99-1]


C₂₂H₃₃NO₅S 423.572

Macrolide antibiotic. *Isol.* from the sponge *Fasciospongia rimosa*. Cytotoxic agent. [α]_D²⁶ +110 (c, 0.2 in CHCl₃).

Tanaka, J. *et al.*, *Chem. Lett.*, 1996, 255 (*isol*, *ir*, *pmr*, *cmr*)
 Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 115-134 (*synth*)

Latrunculin T

[122876-49-7]

C₂₀H₂₇NO₅S 393.503


Isol. from *Negombata magnifica*. Antimicrobial agent. Oil. [α]_D²⁵ +109.5 (c, 0.11 in EtOH).

El Sayed, K.A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 219-223 (*isol*, *pmr*, *cmr*)

Absolute Configuration

Lauberine

[19879-48-2]

C₃₇H₄₀N₂O₆ 608.733

The alkaloids covered by this entry (2*R*,2'*S*-) belong to the enantiomeric series to those included in Dryadophnine, D-944 (2*S*,2'*R*-) and are diastereomeric with those in Thalmine, T-341 (2*S*,2'-). Alkaloid from the trunk bark and roots of *Berberis laurina* (Berberidaceae). [α]_D -481 (c, 0.2 in CHCl₃). Oxidises rapidly in air.

Hydrobromide (1:2):

Cryst. (EtOH). Mp 250-255°. [α]_D -335 (c, 0.6 in MeOH).

Me ether: [α]_D -369 (c, 0.2 in CHCl₃).*O*⁶-*De-Me*: **Berbilaurine**

[120416-86-6]

C₃₆H₃₈N₂O₆ 594.706

Minor alkaloid from roots of *Berberis laurina* (Berberidaceae).

*O*¹²-*De-Me*: **12-O-Demethylauberine**

[123880-31-9]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis chilensis* (Berberidaceae).

Falco, M.R. *et al.*, *Tet. Lett.*, 1968, 1953 (*pmr*, *ms*, *struct*)


Baldas, J. *et al.*, *Tet. Lett.*, 1968, 6315 (*pmr*)

Baldas, J. *et al.*, *J.C.S. Perkin I*, 1972, 597 (*ms*)

Torres, R. *et al.*, *Bol. Soc. Chil. Quim.*, 1989, **34**, 11; *CA*, **111**, 233323h (12-*O*-

Demethylauberine)

Weber, J.F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 81 (*Berbilaurine*)

Laudanidine*Tritopine*C₂₀H₂₅NO₄ 343.422

The name Laudanidine was originally applied to the racemate.

(R)-form [301-21-3]

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Mp 184-185°. [α]_D²⁸ -101 (EtOH).

Picrate: Mp 174-175°.*Ac*: Mp 98°.**(S)-form** [3122-95-0]

Alkaloid from *Machilus obovatifolia*, *Machilus arisaensis*, *Thalictrum dasycarpum* and *Thalictrum revolutum* (Lauraceae, Ranunculaceae). Mp 181-182°. [α]_D²⁸

L-67

+134 (c, 0.51 in MeOH). [α]_D +87 (c, 0.56 in CHCl₃).

N-Me: **N-Methylaudanidinium**C₂₁H₂₈NO₄⁺ 358.456

Quaternary alkaloid from tubers of

Corydalis solidia (Papaveraceae).

Amorph. solid (as iodide). [α]_D²³ +72 (c, 0.04 in MeOH).

Me ether: see Laudanosine, L-69**(±)-form****Laudanine**

[85-64-3]

Alkaloid from *Papaver somniferum* (opium poppy) and *Xylopija pancheri* (Papaveraceae, Annonaceae). Convulsive agent producing strychnine-like tetany. Small doses are hypertensive. Mp 166-167°. Log P 2.91 (calc).

Picrate: Mp 176-177°.*Benzoyl*: Mp 156-157°.*6-O-De-Me, N-Me*: **Xylopinidine**C₂₀H₂₆NO₄⁺ 344.43

Alkaloid from the bark and roots of *Xylopija parviflora*. Amorph. powder (as perchlorate). λ_{max} 282 (log ε 3.68) (MeOH) (perchlorate).

Späth, E. *et al.*, *Monatsh. Chem.*, 1926, **47**, 733-740 (*synth*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 593-601 (*isol*, *uv*, *pmr*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1985, **50**, 2299-2309 (*N*-

Methylaudanidinium)


Blanchfield, J.T. *et al.*, *Phytochemistry*, 2003, **63**, 711-720 (*isol*, *pmr*, *cmr*)

Nishiyama, Y. *et al.*, *Phytochemistry*, 2004, **65**, 939-944 (*Xylopinidine*)

Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 1073

Laudanosine**L-69**

N-Methyltetrahydropapaverine. Laudanine methyl ether

**(S)-form**C₂₁H₂₇NO₄ 357.449**(R)-form** [85-63-2]

Synthetic. Mp 83-85°. [α]_D²⁰ -85 (c, 0.45 in EtOH).

(S)-form [2688-77-9]

Alkaloid from *Papaver somniferum* (opium poppy) and *Argemone grandiflora* (Papaveraceae). Major metabolite of the neuromuscular blocker Atracurium(2+). Convulsive agent acting on the extrapyramidal system and mesencephalon. Mp 89°. [α]_D²⁵ +103 (EtOH). [α]_D +52 (CHCl₃).

Methiodide: Mp 225-227° (218-221°).[α]_D +120.*N-De-Me*: **Norlaudanosine**

[4747-98-2]

Light yellow oil. [α]_D²⁵ -21.9 (c, 1 in CHCl₃).

(±)-form [1699-51-0]


Synthetic. Mp 114-115° (67-70°).

▶ NX5070000

Hydrochloride: Mp 123°.*Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, **1**, 1318A (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **2**, 667C (*nmr*)Cymmerman-Craig, J. *et al.*, *Tetrahedron*, 1966, **22**, 1335-1339 (*uv, ord, config*)Elliott, I.W. *et al.*, *J. Het. Chem.*, 1970, **7**, 1229-1230 (*synth*)Preininger, V. *et al.*, *Alkaloids (Academic Press)*, 1975, **15**, 207 (*pharmacol*)Konda, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 1025-1031; 1977, **25**, 69-74 (*synth, ir, pmr, ms*)Singh, S.P. *et al.*, *J. Het. Chem.*, 1978, **15**, 541-544 (*cmr*)Takano, S. *et al.*, *Chem. Comm.*, 1982, 769-770 (*synth*)Czarnocki, Z. *et al.*, *Can. J. Chem.*, 1986, **64**, 2205-2210 (*synth*)Gawley, R.E. *et al.*, *Tet. Lett.*, 1988, **29**, 301-302 (*synth*)Gottlieb, L. *et al.*, *J.O.C.*, 1990, **55**, 5659-5662 (*synth*)Coppola, G.M. *et al.*, *J. Het. Chem.*, 1991, **28**, 1769-1772 (*synth*)Comins, D.L. *et al.*, *Tet. Lett.*, 1991, **32**, 2995-2996 (*synth*)Takano, S. *et al.*, *Heterocycles*, 1993, **35**, 47-52 (*synth*)*Martindale, The Extra Pharmacopoeia, 30th edn.*, Pharmaceutical Press, 1993, 1200Kitamura, M. *et al.*, *J.O.C.*, 1994, **59**, 297-310 (*synth*)Comins, D.L. *et al.*, *Tetrahedron*, 1997, **48**, 16327-16340 (*synth*)Mujahidin, D. *et al.*, *Eur. J. Org. Chem.*, 2005, 2689-2693 (*synth*)Werner, F. *et al.*, *Eur. J. Org. Chem.*, 2007, 3911-3915 (*S-form, N-de-Me*)**Laudanosoline****L-70**

[485-33-6]

[3184-36-9]

C₁₇H₁₉NO₄ 301.341

Synthetic or semisynthetic parent of a group of benzylisoquinoline alkaloids.

(R)-form [57231-32-0]**Hydrobromide:** [38221-29-3]

[84030-13-7]

Mp 124-125°. [α]_D -57.3 (c, 1 in EtOH).**Hydrochloride:** [163455-60-5]

[3184-36-9]

O⁷,N-Di-Me: Semiaquilegine A

[935266-65-2]



C₁₉H₂₄NO₃⁺ 330.403Quaternary alkaloid from the roots of *Semiaquilegia adoxoides*. Amorph. powder. Mp 220-222°. [α]_D²² -113 (c, 0.5 in H₂O). λ_{max} 206; 254; 282 (no solvent reported).**(S)-form** [57231-31-9]**Hydrobromide:**

[38221-28-2]

Cryst. Mp 124-125°. [α]_D +57.8 (c, 1 in EtOH).**3',6-Di-Me ether:** see Orientaline, O-111**4',6-Di-Me ether:** see Reticuline, R-59**4',7-Di-Me ether:** see Protosinomenine, P-668**3',4',6-Tri-Me ether:** see Codamine, C-554**3',4',7-Tri-Me ether:** see Pseudolaudanine, P-715**4',6,7-Tri-Me ether:** see Laudanidine, L-68**Tetra-Me ether:** see Laudanosine, L-69**(±)-form** [19481-42-6]

Mp 114-116°.

[84030-13-7, 163455-61-6]

Teitel, S. *et al.*, *J. Med. Chem.*, 1972, **15**, 845-846 (*synth, N-de-Me*)Gibson, E.P. *et al.*, *J.C.S. Perkin 2*, 1980, 1696-1700 (*uv*)Desbourges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664-693; 852-859 (*codamine oxide*)Goralski, C.T. *et al.*, *Ind. Eng. Chem. Res.*, 1989, **28**, 221-224 (*synth*)Stammel, W. *et al.*, *Chirality*, 1995, **7**, 10-19 (*chiral hplc*)Niu, F. *et al.*, *Chin. J. Chem.*, 2006, **24**, 1788-1791 (*Semiaquilegine A*)**Laughine****L-71**C₁₁H₁₈BrN₅O 316.2Isol. from the sponge *Eurypon laughlini*. Pale yellow oil. λ_{max} 206 (ε 3800); 265 (ε 4020) (MeOH).Williams, D.E. *et al.*, *J. Nat. Prod.*, 2005, **68**, 327-330 (*isol, pmr, cmr*)**Laureline****L-72****10-Methoxy-1,2-methylenedioxyaporphine**C₁₉H₁₉NO₃ 309.364**(R)-form** [81-38-9]Alkaloid from the bark of *Laurelia novae-zealandiae* and from the leaves, twigs and bark of *Hedycarya angustifolia* (Monimiaceae). Cryst. (petrol). Mp 114°. [α]_D -97.7 (EtOH).**Hydrochloride:** Mp 280°.**Methiodide:**

Cryst. (EtOH). Mp 223°.

N-De-Me: 10-Methoxy-1,2-methylene-**dioxynoraporphine. Norlaureline**

[65012-41-1]

C₁₈H₁₇NO₃ 295.337Alkaloid from the stem bark of *Guatteria elata* (Annonaceae).

Rapidly darkens on exp. to air and light.

N-De-Me, N-Ac:Cryst. (Me₂CO). Mp 208-209°. [α]_D²⁵ -336 (c, 0.25 in MeOH).**O-De-Me: 10-Hydroxy-1,2-methylene-****dioxyporphine. Mecambroline. Isofi-****gapavine. Phoebe base 2**C₁₈H₁₇NO₃ 295.337Alkaloid from *Phoebe clemensii* (Lauraceae). Mp 220-222° (darkens at 210°). [α]_D -76.5 (c, 0.366 in CHCl₃).**O-De-Me, hydrochloride:** Mp 330°. [α]_D -28.3 (c, 0.315 in H₂O).**O-De-Me, N-Me: Michepressine**

[25454-85-7]


C₁₉H₂₀NO₃⁺ 310.372Quaternary alkaloid from *Michelia compressa* (Magnoliaceae). Mp 235-236° dec. (as iodide). [α]_D¹⁵ -131 (c, 1 in MeOH).**6,6a-Didehydro, N-de-Me: 6,6a-Dehydro-****norlaureline**C₁₈H₁₅NO₃ 293.321Alkaloid from the leaves, twigs and bark of *Hedycarya angustifolia* (Monimiaceae). Brown gum.**(S)-form**Synthetic. Mp 114°. [α]_D +97.6 (EtOH).**O-De-Me:** [3466-56-6]C₁₈H₁₇NO₃ 295.337Alkaloid from *Meconopsis cambrica*, also from the genera *Laurelia*, *Phoebe* and *Papaver* (Papaveraceae, Monimiaceae, Lauraceae). Hypotensive in mice and rabbits. Mp 254-255°. [α]_D²³ +76 (c, 0.48 in CHCl₃).**O-De-Me, hydrochloride:** Mp 264-266°.**O-De-Me, picrate:** Mp 179-180°.**(±)-form**

Synthetic.

Hydrochloride:Cryst. + 1H₂O (0.01M HCl aq.). Mp 230° dec.Aston, B.C. *et al.*, *J.C.S.*, 1910, **97**, 1381 (*isol*)
Barger, G. *et al.*, *Helv. Chim. Acta*, 1931, **14**, 481 (*struct*)Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1932, **15**, 394 (*synth*)Faltis, F. *et al.*, *Ber.*, 1944, **77**, 686 (*synth*)Ito, K. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 703; *CA*, **55**, 23571h (*Mecambroline, isol*)Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 1720 (*Mecambroline, isol, uv, ir*)Barton, D.H.R. *et al.*, *Chem. Comm.*, 1966, 259 (*biosynth*)Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1277 (*pmr*)Gibson, M.S. *et al.*, *J.C.S. (C)*, 1970, 2234 (*synth, uv, ir, ms*)Hsu, C.C. *et al.*, *J. Nat. Prod.*, 1975, **38**, 544; 1977, **40**, 505 (*Norlaureline*)Minamikawa, J. *et al.*, *Can. J. Chem.*, 1979, **57**, 1720 (*synth, uv*)Geewanda, Y.A. *et al.*, *Heterocycles*, 1987, **26**, 447 (*6,6a-Dehydronorlaureline*)

Laurifonine

6,7,8,9-Tetrahydro-2,3,12-trimethoxy-7-methyl-5H-dibenzo[d,f]azonine, 9CI [56261-25-7]



C₂₀H₂₅NO₃ 327.422

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae).

Amorph.

Perchlorate:

Cryst. (MeOH/Et₂O). Mp 191° (182-185°).

Methodide: Mp 165-170°.

N-De-Me: Laurifine

[56261-28-0]

C₁₉H₂₃NO₃ 313.396

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae).

Amorph.

O³-De-Me: Laurifinine

[56261-29-1]

C₁₉H₂₃NO₃ 313.396

Alkaloid from the leaves of *Cocculus laurifolius* (Menispermaceae). Mp 179-181°.

O³-De-Me; perchlorate:

Cryst. (MeOH/Et₂O). Mp 243-245°.

Kupchan, S.M. *et al.*, *Heterocycles*, 1976, **4**, 235 (synth)

Pande, H. *et al.*, *J.C.S. Perkin 1*, 1976, 2197 (isol, uv, ir, pmr, ms, struct, Laurifonine, Laurifine, Laurifinine)


Ito, K. *et al.*, *Heterocycles*, 1978, **9**, 485 (synth, pmr, struct)

Bhakuni, D.S. *et al.*, *Indian J. Chem., Sect. B*, 1981, **20**, 531 (synth, ms)

Bremner, J.B. *et al.*, *Heterocycles*, 1985, **23**, 1451 (synth, Laurifonine, Laurifine)

Lauroidionine

[191654-78-1]




C₂₀H₁₅NO₆ 365.342

Modified aporphine. Alkaloid from the stems of *Phoebe formosana* (Lauraceae). Black prisms (MeOH). Mp > 300°. λ_{max} 204 (log ε 4.2); 266 (log ε 3.99); 344 (log ε 3.99); 430 (log ε 3.73); 537 (log ε 3.5) (MeOH).

Chen, C.-C. *et al.*, *J. Nat. Prod.*, 1997, **60**, 826-827 (isol, uv, ir, pmr, cmr)

L-73**Lauroschooltzie**

5,6,6a,7-Tetrahydro-1,2,10-trimethoxy-6-methyl-4H-dibenzo[d,e]quinolin-9-ol, 9CI. 9-Hydroxy-1,2,10-trimethoxyaporphine. N-Methyllaurotetanine. Rogersine



C₂₀H₂₃NO₄ 341.406

(S)-form [2169-44-0]

Alkaloid from *Eschscholtzia californica* (Papaveraceae) and *Phyllica rogersii* (Rhamnaceae), also from the Hernandiaceae, Lauraceae, Magnoliaceae, Annonaceae, Monimiaceae, Ranunculaceae and Menispermaceae. Cardiac depressant, hypotensive agent, also shows antibacterial props. Mp 100-105°. [α]_D²⁵ +111 (c, 0.86 in MeOH).

▶CE1057900

N-Oxide (R-): N-Methyllaurotetanine N^β-oxide

[100010-00-2]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Glossocalyx brevipes* and *Neolitsea sericea* var. *aurata*.

Amorph. [α]_D²⁵ +49 (c, 0.15 in MeOH).

N-Oxide (S-): N-Methyllaurotetanine N^α-oxide

[937018-79-6]

C₂₀H₂₃NO₅ 357.405

Alkaloid from the leaves of *Neolitsea sericea* var. *aurata*.

N-Me: Xanthoplanine

[6872-88-4]

C₂₁H₂₆NO₄[⊕] 356.441

Quaternary alkaloid from *Zanthoxylum planispium*, *Fagara hyemalis*, *Fagara nigrescens*, *Hernandia ovigera* (Rutaceae, Hernandiaceae) and *Litsea cubeba* (mountain pepper) (Lauraceae). Mp 207-209° (as iodide). [α]_D²¹ +62 (c, 0.83 in EtOH).

N-De-Me: 9-Hydroxy-1,2,10-trimethoxy-noraporphine. Laurotetanine. Litsoene

[128-76-7]

C₁₉H₂₁NO₄ 327.379

Alkaloid from a wide variety of genera in the Annonaceae (*Xylopi*, *Desmos*, *Guatteria*), Hernandiaceae (*Hernandia*, *Illigeria*), Lauraceae (*Actinodaphne*, *Cassytha*, *Cryptocarya*, *Lindera*, *Litsea*, *Neolitsea*, *Machilus*), Monimiaceae (*Dryadodaphne*, *Palmeria*, *Peumus*, *Hedycarya*, *Monimia*, *Nemuaron*, *Laurelia*). Shows antiplatelet aggregation activity. Mp 125°. [α]_D²⁵ +95 (c, 0.50 in EtOH). λ_{max} 219 ; 282 ; 304 (MeOH) (Berdy).

▶RB6025000

N-De-Me, hydrochloride: Mp 245°.

N-De-Me, N-ethoxycarbonyl: N-Ethoxycarbonyllaurotetanine

[947263-10-7]

C₂₂H₂₅NO₆ 399.443

Alkaloid from the roots of *Lindera angustifolia*.

(±)-form [38849-65-9]

Mp 199-202° dec. (143-144°).

N-De-Me: [65877-55-6]

C₁₉H₂₁NO₄ 327.379

Synthetic.

Ruegger, A. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 754-762 (isol, uv)

Ishii, H. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 238 (Xanthoplanine)

Pachler, K.G.R. *et al.*, *Tetrahedron*, 1965, **21**,

2159 (Laurotetanine, struct, pmr)

Albonico, S.M. *et al.*, *J.C.S. (C)*, 1966, 1340

(Xanthoplanine, uv, ord)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1972, 1435

(synth, spectra, derivs)

Govindachari, T.R. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 873 (synth, Laurotetanine)

Marsaioli, A.J. *et al.*, *Phytochemistry*, 1979, **18**, 165 (cmr, Xanthoplanine)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (cd, Laurotetanine, N-Methyllaurotetanine)

Montgomery, C.T. *et al.*, *J. Nat. Prod.*, 1985,

48, 833 (N-Methyllaurotetanine β-N-oxide)

Hara, H. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**,

1946 (synth)

Lee, S.S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1971-

1976 (Xanthoplanine, isol, uv, ir, pmr)

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 55 (cd)

Chen, J.-J. *et al.*, *Planta Med.*, 2000, **66**, 251-

256 (activity)

Zhao, O. *et al.*, *Yaoxue Xuebao*, 2005, **40**, 931-934 (N-Ethoxycarbonyllaurotetanine)

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 637-

642 (N-oxides)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties*

of Industrial Materials, 8th edn., Van

Nostrand Reinhold, 1992, LBO200; TKX700

Lavanducyanin**L-76**

5-[(2,4,4-Trimethyl-1-cyclohexen-1-yl)-methyl]-1-(5H)-phenazinone, 9CI. 1,9-Dihydro-5-[(2,4,4-trimethyl-1-cyclohexen-1-yl)methyl]phenazin-1-one. WS 9659A. Antibiotic YP 02978L-C. Antibiotic WS 9659A. Antibiotic YP 0298L-C [122228-60-8]



C₂₂H₂₄N₂O 332.444


Phenazine antibiotic. Prod. by *Streptomyces* spp. Antitumour agent and testosterone 5α-reductase inhibitor. Deep blue prisms (MeOH/Et₂O). Mp 161-162° (135-136°). Related to Phenazinomycin, P-329. λ_{max} 274 (ε 9700); 386 (ε 5000); 508 (ε 1400) (MeOH/HCl) (Derep). λ_{max} 237 (ε 12400); 319 (ε 9700); 705 (ε) (MeOH/KOH) (Derep). λ_{max} 237 (ε 12400); 319 (ε 9700); 705 (ε 1700) (MeOH) (Derep). λ_{max} 238 (ε 21600); 320 (ε 23200) (MeOH) (Derep).

▶SG2000000

2-Chloro: *Antibiotic WS 9659B*. WS 9659B
[123313-61-1]
C₂₂H₂₃ClN₂O 366.889
From a *Streptomyces* sp. Testosterone 5 α -reductase inhibitor. Deep blue prisms (MeOH/Et₂O). Sol. MeOH, CHCl₃; fairly sol. Et₂O, C₆H₆; poorly sol. H₂O, hexane. Mp 152-153°. λ_{\max} 241 (ε 15400); 328 (ε 20500) (MeOH) (Derep). λ_{\max} 238 (E1%/1cm 650); 320 (E1%/1cm 700); 730 (MeOH) (Berdy).
Imai, H. *et al.*, *J. Antibiot.*, 1989, **42**, 1196 (*isol. struct*)
Nakayama, O. *et al.*, *J. Antibiot.*, 1989, **42**, 1221; 1230; 1235 (*isol. struct. props*)
Imai, S. *et al.*, *J. Antibiot.*, 1993, **46**, 1232 (*props*)
Miyake, M. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2083 (*synth*)
Kinoshita, Y. *et al.*, *Synlett*, 1995, 186 (*synth*)

Lavanduquinocin

L-77

C₂₆H₃₁NO₃ 405.536**(R)-form** [167114-85-4]

Prod. by *Streptomyces viridochromogenes* 2942-SVS3. Neuronal cell protecting agent. Red-brown powder. Mp 157-158°. λ_{\max} 232 (ε 17500); 269 (ε 15900); 429 (ε 3100) (MeOH). λ_{\max} 244 (ε 16400); 288 (ε 16300); 472 (ε 4300) (MeOH/NaOH).

(±)-form


Black cryst. Mp 221°. λ_{\max} 193 (ε 20800); 211 (ε 21500); 231 (ε 25100); 268 (ε 23100); 427 (ε 4900) (MeOH).

Shin-ya, K. *et al.*, *J. Antibiot.*, 1995, **48**, 574-578 (*isol. pmr. cmr*)
Knölker, H.-J. *et al.*, *Tet. Lett.*, 1998, **39**, 2537-2540 (*synth. uv. ir. pmr. cmr. ms*)
Knölker, H.-J. *et al.*, *Chirality*, 2000, **12**, 526-528 (*synth*)

Lavendamycin

L-78

1-(7-Amino-5,8-dihydro-5,8-dioxo-2-quinoliny)-4-methyl-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI. K 82A. Antibiotic K 82A [81645-09-2]

C₂₂H₁₄N₄O₄ 398.377

Quinone antibiotic. Isol. from *Streptomyces lavendulae*. Antineoplastic antibiotic. Mp 300° dec. Log P 3.2 (uncertain value) (calc). λ_{\max} 234 (ε 19600); 246 (ε 19800); 391 (ε 8400) (MeOH) (Derep).


[81677-52-3]

Doyle, T.W. *et al.*, *Tet. Lett.*, 1981, **22**, 4595 (*struct*)
Balitz, A.W. *et al.*, *J. Antibiot.*, 1982, **35**, 259
Kende, A.S. *et al.*, *Heterocycles*, 1984, **21**, 91 (*rev. synth*)
Boger, D.L. *et al.*, *J.O.C.*, 1985, **50**, 5790 (*synth. bibl*)
Boger, D.L. *et al.*, *J. Med. Chem.*, 1987, **30**, 1918 (*pharmacophore. synth*)
Behforouz, M. *et al.*, *J.O.C.*, 1993, **58**, 7089; 1996, **61**, 6552 (*synth. bibl*)
Molina, P. *et al.*, *Tet. Lett.*, 1994, **35**, 1453 (*synth*)

Lavendustin A

L-79

[125697-92-9]

C₂₁H₁₉NO₆ 381.384

Prod. by *Streptomyces griseolavendus*. Tyrosine kinase inhibitor. Angiogenesis inhibitor. Powder + 1½ H₂O. Sol. MeOH, Me₂CO, EtOH, DMSO; poorly sol. CHCl₃, H₂O, Et₂O. Mp 158-162°. λ_{\max} 216 (ε 24000); 280 (ε 4900); 297 (ε 5130) (MeOH) (Derep).

5'-Deoxy: Lavendustin B

[125697-91-8]

C₂₁H₁₉NO₅ 365.385

From *Streptomyces griseolavendus*. Tyrosine kinase inhibitor. Powder. Sol. MeOH, Me₂CO, DMSO, EtOH; poorly sol. CHCl₃, H₂O, Et₂O. Mp 140-144°. λ_{\max} 216 (ε 24000); 275 (ε 5010); 310 (ε 2090) (MeOH) (Derep).

Onoda, T. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1252-1257 (*isol. struct*)

Hu, D.E. *et al.*, *Br. J. Pharmacol.*, 1995, **114**, 262-268 (*activity*)


Green, J. *et al.*, *J.O.C.*, 1995, **60**, 4287-4290 (*synth*)

Fan, T.D. *et al.*, *Trends Biochem. Sci.*, 1995, **16**, 57-66 (*activity*)

Ledecorine

L-80

5-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)methyl]-1,3-benzodioxol-4-ol, 9CI. 1,2,3,4-Tetrahydro-1-(2-hydroxy-3,4-methylenedioxybenzyl)-6,7-methylenedioxyisoquinoline

**(R)-form**C₁₉H₁₉NO₅ 341.363**(R)-form** [68676-56-2]

Alkaloid from *Corydalis ledebouriana* and *Fumaria vaillantii* (Papaveraceae). Mp 199-200°. $[\alpha]_D^{25}$ -112 (c, 0.42 in MeOH). Abs. config. shown incorrectly as S- in 1980 paper and in CAS.

Me ether: Marshalline

[68676-57-3]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Corydalis marshalliana* (Papaveraceae). Mp 134-135°. $[\alpha]_D^{25}$ -105 (c, 0.2 in CHCl₃) (nat.). $[\alpha]_D^{25}$ -63.2 (c, 0.44 in CHCl₃) (synthetic). This struct. was also assigned independently to Fumarizine, F-179. Takaba *et al.* showed that their synthetic product was not identical with Fumarizine but did not compare their synthetic material with Marshalline. The pmr spectra appear v. similar although the opt. rotns. differ.

(±)-form [74924-41-7]

Synthetic. Gum.

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 537-538; 1984, **20**, 672-673; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 465-466; 1984, **20**, 644 (*Ledecorine, Marshalline, isol. ir. pmr. ms, ord. struct*)

Murugesan, N. *et al.*, *Heterocycles*, 1980, **14**, 585-587 (*synth. uv. pmr. ms*)

Takaba, K. *et al.*, *Heterocycles*, 1996, **43**, 1777-1786 (*Marshalline, synth*)


Lehmannine†

L-81

12,13-Didehydromatridin-15-one, 9CI.

12,13-Dehydromatrine

[58480-54-9]

C₁₅H₂₂N₂O 246.352

Alkaloid from epigeal parts of *Ammothamnus lehmannii* (preferred genus name *Sophora*) (Fabaceae). Mp 93-94°. $[\alpha]_D^{24}$ +37.03 (c, 0.54 in EtOH).

Picrate: [58480-55-0]

Mp 144-145°.

N'β-Oxide: Lehmannine N-oxide

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Ammothamnus lehmannii* (Fabaceae). Mp 136°.

Kushmadurov, Ya.K. *et al.*, *Khim. Prir. Soedin.*, 1975, 377; *Chem. Nat. Compd. (Engl. Transl.)*, 389 (*isol. ir. uv. ms. pmr. struct*)


Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1981, 757; *Chem. Nat. Compd.*

(Engl. Transl.), 552 (cryst struct, oxide)

Leinamycin

L-82

DC 107. Antibiotic DC 107
[120500-15-4]



$C_{22}H_{26}N_2O_6S_3$ 510.655

Prod. by a *Streptomyces* sp. S-140.


Antitumour agent. Active against gram-positive and negative bacteria. Appears to function by thiol-activated DNA cleavage. Needles + $\frac{3}{4}$ EtOAc (CHCl₃). Sol. MeOH, DMSO, CHCl₃; poorly sol. H₂O, hexane. Mp 155° dec. $[\alpha]_D^{25}$ +140 (c, 0.1 in MeOH). λ_{max} 211 (ε 14000); 322 (ε 12000) (MeOH) (Derep). λ_{max} 211 (ε 14000); 322 (ε 12000) (MeOH) (Berdy).

- ▶ LD₅₀ (mus, ivn) 2.8 mg/kg. WH1306540 Hara, M. *et al.*, *J. Antibiot.*, 1989, **42**, 333-335; 1768-1774 (isol)
- Pattenden, G. *et al.*, *J. Het. Chem.*, 1992, **29**, 607 (synth)
- Hirayama, N. *et al.*, *Chem. Lett.*, 1993, 1957 (abs config)
- Kanda, Y. *et al.*, *J.A.C.S.*, 1993, **115**, 8451 (synth)
- Mitra, K. *et al.*, *J.A.C.S.*, 1997, **119**, 11691-11692 (biochem)
- Tang, G.-L. *et al.*, *Chem. Biol.*, 2004, **11**, 33-45 (biosynth)
- Tang, G.-L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 387-393 (biosynth)

Leiocarpadiol

L-83

[126365-19-3]




$C_{18}H_{25}NO_2$ 287.401

Alkaloid from roots of *Esenbeckia leiocarpa* (Rutaceae). Amorph. solid. $[\alpha]_D^{20}$ +2.5 (c, 0.6 in CHCl₃). λ_{max} 264 (sh) (log ε 3.74); 271 (log ε 3.76); 280 (log ε 3.7) (no solvent reported).

Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (isol, uv, ir, pmr, cmr, ms, struct)

Leiodelide A

L-84




$C_{31}H_{45}NO_9$ 575.698

Isol. from the sponge *Leiodermatium* sp. Cytotoxic. Pale yellow oil. λ_{max} 224 (log ε 3.2) (MeOH).

Sandler, J.S. *et al.*, *J.O.C.*, 2006, **71**, 7245-7251; 8684 (isol, cd, pmr, cmr, ms)

Leiodelide B

L-85



$C_{31}H_{44}BrNO_9$ 654.594


Isol. from the sponge *Leiodermatium* sp. Cytotoxic. Pale yellow oil. λ_{max} 221 (log ε 2.8) (MeOH).

Sandler, J.S. *et al.*, *J.O.C.*, 2006, **71**, 7245-7251; 8684 (isol, cd, pmr, cmr, ms)

Lelobanidine

L-86

α -Ethyl-1-methyl- α' -phenyl-2,6-piperidine-diethanol, 9CI. 8-Ethyl-10-phenyllobelidol
[492-48-8]



$C_{18}H_{29}NO_2$ 291.433

Three (-)-forms and one racemate have been isol. All are 2,6-cis but side-chain configs. and abs. configs. are unknown.

(-)(1)-form**(-)-Lelobanidine I**

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Mp 86° (as hydrochloride). $[\alpha]_D$ -41.5 (EtOH).

(-)(2)-form**(-)-Lelobanidine II**

Minor alkaloid from *Lobelia inflata* and *Lobelia nicotianaefolia* (Campanulaceae).

Cryst. + H_2O (as hydrochloride). Mp 102-105° (hydrochloride). $[\alpha]_D$ -41.2 (EtOH). On oxidn. gives the same diketone as (-)-Lelobanidine I.

(-)(3)-form**(-)-Lelobanidine III. α -Lelobanidine**

Alkaloid from *Lobelia nicotianaefolia* (Campanulaceae). $[\alpha]_D^{25}$ -37.1 (c, 0.46 in EtOH). Identity of the natural Lelobanidine III with semisynthetic (-)- α -Lelobanidine (obt. from Lobinandinine, L-221) not certain. Opt. rotns. were identical, but α -Lelobanidine did not form a cryst. hydrochloride.

Hydrochloride:

Cryst. + $1\frac{1}{2}$ H₂O. Mp 111°.

(+)-form**N-De-Me: Norlelobanidine**

$C_{17}H_{27}NO_2$ 277.406

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Rosettes (petrol). Mp 90°. $[\alpha]_D$ +62.8. This is the N-demethyl deriv. of the enantiomer of (-)-Lelobanidine I.

N-De-Me; hydrochloride: Mp 193°.

(±)-form**(±)-Lelobanidine I**

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Mp 68°. Has been resolved to give (+)-Lelobanidine I.

Hydrochloride: Mp 79°.

(±)-form**Hydroxy: Hydroxylelobanidine**

$C_{18}H_{29}NO_3$ 307.432

Alkaloid from *Sedum acre*. OH-Subst. in the piperidine ring, posn. undetd. Stereochem. unknown.

Wieland, H. *et al.*, *Annalen*, 1939, **540**, 103 (isol, struct)

Gedeon, J. *et al.*, *Pharm. Acta Helv.*, 1955, **30**, 185 (isol)


Spiteller-Friedmann, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 104 (ms)

Francis, L.P.S. *et al.*, *Planta Med.*, 1977, **32**, 268 (isol, Lelobanidine, Hydroxylelobanidine)

Lemairamine

L-87

N-(3,4-Dimethoxyphenylethyl) cinnamide



$C_{19}H_{21}NO_3$ 311.38

(E)-form [29946-61-0]

Alkaloid from *Zanthoxylum lemairie*. Needles. Mp 119-120°. λ_{max} 206; 216 (sh); 224 (sh); 277 (MeOH).

N-Me: Lemairamide

[193901-52-9]


$C_{20}H_{23}NO_3$ 325.407

Alkaloid from *Zanthoxylum lemairie*. Powder.

Adesina, S.K. *et al.*, *Planta Med.*, 1997, **63**, 286-287 (isol, synth, uv, pmr, cmr, ms)

Lemobiline**L-88**

3,9-Dihydro-2,2,3,9-tetramethylfuro[2,3-b]quinolin-4(2H)-one, 9CI. Alkaloid Fl-A. Spectabiline†



C₁₅H₁₇NO₂ 243.305

Shows antineoplastic activity. Log P 2.42 (clc).

(-)-form [26126-76-1]

Alkaloid from *Ravenia spectabilis* and *Flindersia iffilaiana* (Rutaceae, Flindersiaceae). Mp 53-54°. [α]_D -5.9 (-0.9) (CHCl₃). Forms monohydrate, Mp 98-9°. Pharmacol. active isomer.

Picrate: Mp 198° dec. (188-191°).

(±)-form

Mp 106-108°.

Picrate: Mp 186-190°.


Talapatra, S.K. *et al.*, *Tet. Lett.*, 1969, 4789 (*ir, uv, pmr, ms, isol, struct*)

Chamberlain, T.R. *et al.*, *J.C.S. (C)*, 1971, 910 (*synth, isol*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Lemonnierin**L-89**

[75775-38-1]



C₂₆H₃₇N₅O₆ 515.608


Prod. by *Pseudomonas lemmonieri*. Blue pigment. λ_{\max} 625 (log ϵ 4.75) (Py).

Ferguson, G. *et al.*, *J.C.S. Perkin 1*, 1980, 1782-1787

Jain, K.C. *et al.*, *J.C.S. Perkin 1*, 1980, 1788-1794

Lemonomycin**L-90**

LL-AP-191 γ . Antibiotic LL-AP-191 γ



C₂₇H₄₁N₅O₉ 551.636

Isol. from *Streptomyces candidus*. Active against gram-positive and -negative bacteria. Antitumour agent. Sol. MeOH, CHCl₃, acids, DMSO; poorly sol. H₂O. Mp > 200° dec. [α]_D -116 (CHCl₃). λ_{\max} 272 ; 362 (MeOH) (Berdy). λ_{\max} 271

(E1%/1cm 214); 325 (E1%/1cm 17) (HCl) (Berdy).


Whaley, H.A. *et al.*, *Antimicrob. Agents Chemother.*, 1964, 83-86 (*isol, uv, props*)

He, H. *et al.*, *Tet. Lett.*, 2000, **41**, 2067-2071 (*pmr, cmr, struct*)

Ashley, E.R. *et al.*, *J.A.C.S.*, 2003, **125**, 15000-15001 (*synth*)

Lennoxamine**L-91**

5,6,12b,13-Tetrahydro-9,10-dimethoxy-8H-1,3-dioxolo[4,5-h]isoindolo[1,2-b][3]benzazepin-8-one, 9CI [95530-38-4]



C₂₀H₁₉NO₅ 353.374

(±)-form

Minor alkaloid from *Berberis darwinii* (Berberidaceae). Cryst. (MeOH). Mp 228-229° (225°).

Deoxo: **Chilenamine**. Schopf's base VI [95585-66-3]

C₂₀H₂₁NO₄ 339.39

Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Cryst. or amorph. Mp 180-181° (177-179°).

Teitel, S. *et al.*, *Can. J. Chem.*, 1972, **50**, 2022 (*synth, uv, ir, pmr*)

Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957 (*occur, uv, ir, pmr, ms*)

Napolitano, E. *et al.*, *J.C.S. Perkin 1*, 1986, 785 (*synth*)

Yasuda, S. *et al.*, *Heterocycles*, 1990, **30**, 335 (*synth*)

Moody, C.J. *et al.*, *J.C.S. Perkin 1*, 1990, 2929 (*synth*)

Koseki, Y. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1604 (*synth*)

Rodríguez, G. *et al.*, *J.O.C.*, 1996, **61**, 2780 (*synth*)

Ruchirawat, S. *et al.*, *Tet. Lett.*, 2000, **41**, 8007-8010 (*synth, pmr, cmr, ms*)

Couture, A. *et al.*, *Tetrahedron*, 2000, **56**, 1491-1499 (*synth*)

Koseki, Y. *et al.*, *Heterocycles*, 2003, **59**, 527-540 (*synth*)

Sahakitpichan, P. *et al.*, *Tetrahedron*, 2004, **60**, 4169-4172 (*synth*)

Comins, D.L. *et al.*, *Org. Lett.*, 2005, **7**, 95-98 (*synth*)

Taniguchi, T. *et al.*, *Org. Lett.*, 2005, **7**, 4389-4390 (*synth*)


Honda, T. *et al.*, *Tet. Lett.*, 2005, **46**, 6823-6825 (*synth*)

Couty, S. *et al.*, *Tet. Lett.*, 2006, **47**, 767-769 (*synth*)

Fuwa, H. *et al.*, *Heterocycles*, 2008, **76**, 521-539 (*synth*)

Lenticellarine**L-92**

[117590-94-0]



C₁₉H₂₅NO₅ 347.41

Appears to be a homoseco-erythrina alkaloid. Alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae) and *Lagarostrobos colensoi* (Dacrydium *colensoi*). Has moderate molluscicidal activity. Gum which darkens upon standing. [α]_D²⁸ +16 (c, 0.165 in EtOH).

Picrate: Mp 78-80°.

2 α -Hydroxy: **2-Hydroxylenticellarine** [155272-49-4]

C₁₉H₂₅NO₆ 363.41

Alkaloid from stems of *Dysoxylum lenticellare* (Meliaceae).

2 α -Methoxy: **2-Methoxylenticellarine** [155272-48-3]

C₂₀H₂₇NO₆ 377.436

Alkaloid from stems of *Dysoxylum lenticellare* (Meliaceae).

Aladesanmi, A.J. *et al.*, *Planta Med.*, 1986, 522 (*isol, uv, ir, pmr, cmr, ms, struct*)

Aladesanmi, A.J. *et al.*, *Phytochemistry*, 1988, **27**, 3789-3792 (*isol, uv, ir, pmr, cmr, ms, struct*)

Aladesanmi, A.J. *et al.*, *Phytochemistry*, 1994, **35**, 1361 (2 α -Hydroxylenticellarine, 2 α -Methoxylenticellarine, *isol, pmr, ir*)


Bloor, S.J. *et al.*, *Phytochemistry*, 1996, **41**, 801-802 (*isol*)

Leontalbine**L-93**

5,17-Didehydromatridin-15-one, 9CI†.

5,17-Dehydromatrine. 5,17-Didehydromatrine

[6475-07-6]



(-)-form

C₁₅H₂₂N₂O 246.352

Stereoisomeric with Isoleontalbine, I-241 and Darvasine, D-80.

▶ OQ1753000

(-)-form

Alkaloid from epigeal parts of *Leontice albertii* (Leonticaceae). Oil. Bp₅ 180-185°. [α]_D²⁶ -167 (c, 1.15 in EtOH). Abs. config. corresponds to (-)-Matrine.

Hydrochloride: Mp 255-257°.

Picrate:

Cryst. (EtOH). Mp 214-216°.

(+)-form

IS-Oxide: **5,17-Dehydromatrine N-oxide** [70509-82-9]

C₁₅H₂₂N₂O₂ 262.351

Minor alkaloid from *Euchresta japonica* (Fabaceae). Hygroscopic cryst. (Me₂CO). [α]_D +209.3 (EtOH).

Bohlmann, F. *et al.*, *Chem. Ber.*, 1958, **91**, 2176 (*synth*)

Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1966, **2**, 67; 1967, **3**, 26; 1968, **4**, 106; *Chem. Nat. Compd. (Engl. Transl.)*, 1966,

2, 55; 1967, 3, 21; 1968, 4, 90 (*isol, struct, ms*)

Ohmiya, S. *et al.*, *Phytochemistry*, 1978, 17, 2021 (*oxide*)


Leontalbinine L-94

7,11-Didehydromatridin-15-one, 9CI.

7,11-Dehydromatrine

[16665-57-9]

[46862-63-9]



Relative configuration

C₁₅H₂₂N₂O 246.352

Alkaloid from *Leontice albertii* and flowers of *Sophora flavescens* (Leonticaceae, Fabaceae). Prisms (Et₂O). Mp 107-108°. [α]_D²⁴ -137.2 (EtOH).

N¹-Oxide: **Leontalbinine N-oxide**

[147731-96-2]

C₁₅H₂₂N₂O₂ 262.351

Minor alkaloid from seeds of *Sophora flavescens* var. *angustifolia* (Fabaceae). [α]_D -161.7 (EtOH).

13-Hydroxy: **Albertine**. 7,11-Didehydro-13-hydroxymatridin-15-one, 9CI. 13-Hydroxy-7,11-dehydromatrine [20078-85-7]

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Leontice albertii* (Leonticaceae). Mp 161°. [α]_D -101.

Iskandarov, S. *et al.*, *Khim. Prir. Soedin.*, 1967, 3, 26; 1968, 4, 137; 1972, 8, 628; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, 3, 21; 1968, 4, 118; 1972, 8, 596 (*isol, struct*)

Murakoski, I. *et al.*, *Phytochemistry*, 1982, 21, 2379 (*isol, ms, uv, pmr, cmr struct*)

Sequine, T. *et al.*, *Yakugaku Zasshi*, 1993, 113, 53; *CA*, 118, 251444x (*oxide*)

Leontamine L-95

C₁₄H₂₆N₂ 222.373

Struct. unknown. Alkaloid from *Leontice eversmannii* (Leonticaceae). Oil. d₄²⁰ 0.99. Bp_{0.4} 118-119°. [α]_D¹⁵ +2.53 (neat). n_D 1.5113.

Methiodide (1:2): Mp 265-268°.


Dipicrate: Mp 194-195°.

Orechoff, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1932, 270, 329-334

Platonova, T.F. *et al.*, *Zh. Obshch. Khim.*, 1953, 23, 880-886

Leonticine L-96

3-[2-(Dimethylamino)ethyl]-6-methoxy-2-[2-(4-methoxyphenyl)ethenyl]phenol, 9CI. *Petaline methine*



C₂₀H₂₅NO₃ 327.422

(*E*)-form [2609-29-2]

Alkaloid from *Leontice leontopetalum*, *Corydalis claviculata* and *Alphonsea sclerocarpa*. Needles (Me₂CO). Mp 121-123° (118.5-119.5°). Poss. artifact.

McShefferty, J. *et al.*, *J. Pharm. Pharmacol.*, 1956, 8, 1117 (*isol*)


McCorkindale, N.J. *et al.*, *Tetrahedron*, 1969, 25, 5475 (*uv, ir, pmr*)

Tadić, D. *et al.*, *J. Nat. Prod.*, 1987, 50, 518 (*isol*)

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1990, 53, 1280 (*isol*)

Leontiformine L-97

2-(Octahydro-2H-quinolizin-3-yl)-1-piperidinecarboxaldehyde, 9CI. 3-(N-Formyl-2-piperidyl)quinolizidine [29073-26-5]



Relative configuration

C₁₅H₂₆N₂O 250.383

Alkaloid from *Leontice leontopetalum* (Leonticaceae). Mp 61-63°. [α]_D²² +51.9 (c, 0.26 in EtOH).

Hydrobromide: Mp 275-276°. [α]_D²² +57.7.

Deformyl: Leontiformidine. Octahydro-3-(2-piperidinyl)-2H-quinolizine, 9CI [29073-28-7]

C₁₄H₂₆N₂ 222.373

Alkaloid from the roots of *Leontice leontopetalum* (Leonticaceae). Mp 44-46°. [α]_D²² +2.7.

Mollov, N.M. *et al.*, *Tetrahedron*, 1970, 26,


3805 (*isol, ir, pmr, struct, synth*)

Schöpf, C. *et al.*, *Annalen*, 1971, 753, 50 (*synth, deriv*)

Panov, P.P. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1972, 25, 55 (*isol, deriv*)

Takatsu, N. *et al.*, *Chem. Pharm. Bull.*, 1987, 35, 891 (*synth*)

Leontismidine L-98



C₁₅H₂₄N₂O₂ 264.367

Config. not known. Stereoisomer of Albertamine in D-79 and 13α-Hydroxymatrine in M-121. Alkaloid from *Leontice smirnowii*. Cryst. (EtOH). Mp 110°. λ_{max} 220 (no solvent reported).

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*,


1996, 32, 615-681; *Chem. Nat.*

Compd. (Engl. Transl.), 1996, 32, 596-675 (*rev*)

Leontismine L-99

11-Hydroxymatridin-15-one, 9Cf. 11-Hydroxyleontine

[41787-66-0]



C₁₅H₂₄N₂O₂ 264.367

Stereoisomer of Darvasoline, D-81. Alkaloid from *Leontice smirnowii* (Leonticaceae). Mp 168-169°. [α]_D +70.7.

Tkeshelashvili, E.G. *et al.*, *CA*, 1973, 78, 156646m (*isol, struct*)

Leonuridine L-100

C₆H₁₂N₂O₃ 160.172

Struct. unknown. Alkaloid from the seeds of *Leonurus sibiricus*. Mp 222°.

Tang, T.-H. *et al.*, *J. Chin. Chem. Soc. (Peking)*, 1940, 7, 105-110; *CA*, 35, 4913³ (*isol*)

Leonurinine L-101

[849664-80-8]


Struct. unknown. Alkaloid from the seeds of *Leonurus sibiricus*. Mp 262-263°.

Hsu, C.-F. *et al.*, *J. Chin. Chem. Soc. (Peking)*, 1934, 2, 337-339; *CA*, 29, 2659⁹ (*isol*)

Ruan, J. *et al.*, *Zhongcaoyao*, 2003, 34, S15-S19

Lepadiformine B L-102

[883942-41-4]



Absolute Configuration

C₁₇H₃₁NO 265.438

Alkaloid from *Clavelina moluccensis*. Pale yellow oil. [α]_D +3 (c, 1 in CHCl₃). λ_{max} 215 (log ε 2.98); 296 (log ε 2.55) (EtOH).

Deoxy: Lepadiformine C

[883942-42-5]

C₁₇H₃₁N 249.439

Alkaloid from *Clavelina moluccensis*.

Pale yellow oil. [α]_D +11 (c, 1 in CHCl₃). λ_{max} 205 (log ε 2.57); 275 (log ε 1.63) (EtOH).

7β-Acetoxy, 1'-deoxy, 1'-thiocyanato:

Cylindricine H

[164740-25-4]

C₂₀H₃₂N₂O₂S 364.551

Alkaloid from the ascidian *Clavelina cylindrica*. Oil. CA name is incorrect.

7α-Acetoxy, 1'-deoxy, 1'-isothiocyanato:

Cylindricine I

[164740-26-5]

C₂₀H₃₂N₂O₂S 364.551

Alkaloid from *Clavelina cylindrica*. Oil.


Has -NCS replacing -SCN. CA name is incorrect.

- Li, C. *et al.*, *Aust. J. Chem.*, 1994, **47**, 1355-1361; 1995, **48**, 955-965 (*Cylindricines*)
Sauviat, M.-P. *et al.*, *J. Nat. Prod.*, 2006, **69**, 558-562 (*Lepadiformines*)

Lepadin B

L-103

[168434-12-6]



Absolute Configuration

C₁₈H₃₁NO 277.449

Alkaloid from the flatworm *Prostheceraeus villatus* and its tunicate prey *Clavelina lepadiformis*. Exhibits significant *in vitro* cytotoxicity against human cancer cell lines. Oil. [α]_D -96 (MeOH). *Prostheceraeus villatus* appears to be an authors' misspelling of *P. vittatus*.

O-(Hydroxyacetyl): **Lepadin A**

[141544-70-9]

C₂₀H₃₃NO₃ 335.486

Alkaloid from *Prostheceraeus villatus* and *Clavelina lepadiformis*. Exhibits significant *in vitro* cytotoxicity against human cancer cell lines. Oil. [α]_D -8.5 (c, 0.002 in CHCl₃).

7-Oxo, O-(hydroxyacetyl): **Lepadin C**

[168434-13-7]


C₂₀H₃₁NO₄ 349.469

Alkaloid from *Prostheceraeus villatus* and *Clavelina lepadiformis*. Oil. [α]_D -25 (MeOH).

Steffan, B. *et al.*, *Tetrahedron*, 1991, **47**, 8729-8732 (*Lepadin A*, *isol*, *pmr*, *cmr*, *struct*)Kubaneck, J. *et al.*, *Tet. Lett.*, 1995, **36**, 6189-6192 (*Lepadin B*, *Lepadin C*)Toyooka, N. *et al.*, *Tetrahedron*, 1999, **55**, 10673-10684 (*synth*)Ozawa, T. *et al.*, *J.O.C.*, 2001, **66**, 3338-3347 (*synth*)Pu, X.-T. *et al.*, *J.O.C.*, 2006, **71**, 6562-6572 (*synth*, *abs config*)Barbe, G. *et al.*, *J.A.C.S.*, 2008, **130**, 13873-13875 (*ent-form*, *synth*)**Lepadin D**

L-104

Decahydro-3-hydroxy-5-(5-hydroxyocetyl)-2-methylquinoline
[444914-17-4]
[444914-18-5 (quaternary salt)]



Absolute Configuration

C₁₈H₃₅NO₂ 297.48

Alkaloid from a *Didemnum* sp. Oil. [α]_D²² +3 (c, 0.2 in MeOH). [α]_D²² -14.4 (c, 0.27 in MeOH) (quaternary salt). *Isol.* also in oily cationic form (counterion not specified).

3-O-(2E-Octenoyl): **Lepadin E**

[444914-21-0]

C₂₆H₄₇NO₃ 421.662Alkaloid from a *Didemnum* sp. Oil.[α]_D²² -2 (c, 0.1 in MeOH). λ_{max} 263

(ε 2110) (MeOH).

3-O-(2E,4E-Octadienoyl): **Lepadin H**

[412328-25-7]

C₂₆H₄₅NO₃ 419.646Alkaloid from *Aplidium tabascum*. Red gum.[α]_D +8.2 (c, 0.75 in CH₂Cl₂).Unstable. λ_{max} 261 (ε 16000) (MeOH).2-Epimer, 3-O-(2E-octenoyl): **Lepadin F**

[412328-23-5]

[444914-19-6]

C₂₆H₄₇NO₃ 421.662Alkaloid from *Aplidium tabascum* and*Didemnum* sp. Antiplasmodial and

antitrypanosomal agent. Red gum or

clear oil. [α]_D +5.5 (c, 0.12 in CH₂Cl₂).[α]_D -1.5 (c, 0.1 in CHCl₃). Unstable.λ_{max} 262 (ε 3000) (MeOH). λ_{max} 261

(ε 16000) (MeOH).

2-Epimer, 3-O-(2E,4E-octadienoyl):

Lepadin G

[412328-24-6]


C₂₆H₄₅NO₃ 419.646Alkaloid from the ascidian *Aplidium**tabascum*. Red gum. [α]_D +12.5 (c, 0.31in CH₂Cl₂). Unstable. λ_{max} 263 (ε

24000) (MeOH).

Wright, A.D. *et al.*, *J. Med. Chem.*, 2002, **45**, 3067-3072 (*isol*, *pmr*, *cmr*)Davis, R.A. *et al.*, *J. Nat. Prod.*, 2002, **65**, 454-457 (*isol*, *pmr*, *cmr*)Pu, X.-T. *et al.*, *J.O.C.*, 2006, **71**, 6562-6572 (*synth*, *abs config*)Niethe, A. *et al.*, *J.O.C.*, 2008, **73**, 3088-3093 (*ent-Lepadin F,G*, *synth*)Li, G. *et al.*, *Org. Lett.*, 2008, **10**, 4991-4994 (*Lepadin F*, *synth*)**Lepinine**

L-105

[111524-32-4]

C₂₂H₃₃NO₃ 359.508Alkaloid from *Aconitum pseudohuiliense*and *Aconitum leucostomum* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 120-122°.N-Oxide: **Lepinine N-oxide**

[169134-44-5]

C₂₂H₃₃NO₄ 375.507Alkaloid from aerial parts of *Aconitum**kirinense* (Ranunculaceae).11-Ac: **Lepetine**

[111509-08-1]

C₂₄H₃₅NO₄ 401.545Alkaloid from *Aconitum pseudohuiliense*

(Ranunculaceae). Cryst.

(Me₂CO/hexane). Mp 130-131°.λ_{max} 247 (no solvent reported).15-Ac: **Kirinine A**

[182128-35-4]

C₂₄H₃₅NO₄ 401.545Alkaloid from *Aconitum kirinense*.O¹-Me: **Lepidine**

[111509-09-2]

C₂₃H₃₅NO₃ 373.534Alkaloid from *Aconitum pseudohuiliense*

(Ranunculaceae).

N-De-Et, N-Me: **Stenocarpine**C₂₁H₃₁NO₃ 345.481Alkaloid from *Aconitella stenocarpa*

(Ranunculaceae). Cryst. (hexane/EtOAc). Mp 179-182°.

[α]_D -44.3 (c, 0.2 in CHCl₃).19,21(N)-Didehydro, N-Et, 1-Ac: **Kirinine C**

[161068-71-9]

C₂₂H₂₉NO₄ 371.475Alkaloid from roots of *Aconitum**kirinense*. Needles (CHCl₃/MeOH).

Mp 218-220°.


11-Epimer: **11-Epilepenine**

[260365-65-9]

C₂₂H₃₃NO₃ 359.508Alkaloid from *Aconitum barbatum* var.*hispidum*.Liang, X.-T. *et al.*, *Pure Appl. Chem.*, 1986, **58**, 711 (*rev*)Song, W. *et al.*, *CA*, 1987, **107**, 233050jNishanov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 734-735; *Chem. Nat. Compd.*(Engl. Transl.), 1993, **29**, 651-653 (*Lepinine N-oxide*)Yue, J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 277-279 (*Lepinine*, *Lepetine*)Feng, F. *et al.*, *J. Chin. Pharm. Sci.*, 1997, **6**, 14-17 (*Kirinine A*)Martin, G.F. *et al.*, *Phytochemistry*, 1997, **46**, 1087-1090 (*cmr*, *Stenocarpine*)Batsuren, D. *et al.*, *Heterocycles*, 1998, **49**, 327-341 (*isol*, *pmr*, *cmr*)Feng, F. *et al.*, *Phytochemistry*, 1998, **49**, 2557-2559 (*Kirinine C*)Miao, Z. *et al.*, *Bopuxue Zazhi*, 1999, **16**, 547-552 (*11-Epilepenine*)**Lepidine**

L-106

[100108-65-4]

C₂₁H₂₀N₄O₂ 360.415Alkaloid from the seeds of *Lepidium**sativum* (garden cress) (Brassicaceae). Cryst. (CH₂Cl₂/MeOH).O-De-Me: **Lepidine B**

[220433-99-8]

C₂₀H₁₈N₄O₂ 346.388Alkaloid from *Lepidium sativum* (gar-den cress). Yellow cryst. (CH₂Cl₂/

MeOH). Mp 212-214°.

8-Demethoxy, 10-hydroxy: **Lepidine D**
[220434-01-5]

C₂₀H₁₈N₄O₂ 346.388

Alkaloid from *Lepidium sativum* (garden cress). Yellow cryst. (CH₂Cl₂/MeOH). Mp 202-204°.


8-Demethoxy, 10-methoxy: **Lepidine C**
[220434-00-4]

C₂₁H₂₀N₄O₂ 360.415

Alkaloid from *Lepidium sativum* (garden cress). Yellow cryst. (CH₂Cl₂/MeOH). Mp 194-195°.

Bahroun, A. *et al.*, *J. Soc. Chim. Tunis.*, 1985, **2**, 15-24; *CA*, **104**, 65910g


Maier, U.H. *et al.*, *Phytochemistry*, 1998, **49**, 1791-1795 (*Lepidines B-D, isol, pmr, cmr, ms*)

Lepidine E**L-107**

C₂₀H₁₈N₄O₂ 346.388

Alkaloid from the seeds of *Lepidium sativum* (garden cress). Yellow cryst. (CH₂Cl₂/MeOH). Mp 196-198°.

Maier, U.H. *et al.*, *Phytochemistry*, 1998, **49**, 1791-1795 (*isol, pmr, cmr, ms*)

Lepidine F**L-108**


C₂₀H₁₈N₄O₂ 346.388

Alkaloid from the seeds of *Lepidium sativum* (garden cress). Pale yellow cryst. (CH₂Cl₂/MeOH). Mp 245-247°.

Maier, U.H. *et al.*, *Phytochemistry*, 1998, **49**, 1791-1795 (*isol, pmr, cmr, ms*)

Lepidopterin**L-109**

3-[2-Amino-4-hydroxy-7(3H)-pteridinylidene]-2-iminopropanoic acid, **9CI**. 2-Amino-4-hydroxy- α -imino- $\Delta^7(3H)\beta$ -pteridinepropionic acid, **8CI**
[29067-92-3]




C₉H₈N₆O₃ 248.201

Constit. of flour moth *Ephestia kühnellia*.

Viscontini, M. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1783; 1962, **45**, 2479; 1963, **46**, 51 (*isol, synth, uv*)

Lepiotin A**L-110**

5-Hydroxy-1-(4-hydroxyphenyl)-2-pyrrolidinone
[207446-07-9]



Absolute Configuration

C₁₀H₁₁NO₃ 193.202

Alkaloid from the mushroom *Macrolepiota neomastoidea*. Oil. Sol. MeOH, butanol. $[\alpha]_D^{25} +7.7$ (c, 0.07 in MeOH). λ_{max} 203 (log ϵ 4.02); 230 (log ϵ 4.32); 282 (log ϵ 3.28) (2-propanol).

5-Me ether: **Lepiotin B**

[207446-08-0]

C₁₁H₁₃NO₃ 207.229


Alkaloid from the mushrooms *Chlorophyllum molybdites* and *Macrolepiota neomastoidea*. Oil. Sol. MeOH, butanol. $[\alpha]_D^{25} +8.3$ (c, 0.2 in MeOH). λ_{max} 202 (log ϵ 4.09); 232 (log ϵ 3.77); 279 (log ϵ 3.13) (MeOH).

Ohta, T. *et al.*, *Heterocycles*, 1998, **47**, 883-891 (*isol, synth, uv, ir, pmr, cmr*)

Kim, K.H. *et al.*, *Bull. Korean Chem. Soc.*, 2008, **29**, 1591-1593 (*isol, cd, pmr, cmr, abs config*)

Lepistine**L-111**

4-Acetyloctahydro-2,5-methano-5H-pyrrodo[3,2-b]-1,4-oxazine, **9CI**
[55623-00-2]



Absolute configuration


C₁₀H₁₆N₂O₂ 196.249

Fungal alkaloid from *Clitocybe fasciculata*. Hygroscopic liq. Mp 242° (as hydrochloride). Bp_{0.1} 140-150°. n_D^{20} 1.5270.

Laing, M. *et al.*, *Tet. Lett.*, 1975, 269 (*cryst struct*)

Leporine A**L-112**

2,6,6a,7,8,9,10,10a-Octahydro-2-methoxy-10-methyl-4-phenyl-6-(1-propenyl)-1H-[2]benzopyrano[4,3-c]pyridin-1-one, **9CI**
[140381-54-0]



Relative Configuration

C₂₃H₂₇NO₃ 365.471

Related to Fusaricide, F-225. Isol. from the sclerotia of *Aspergillus leporis* NRRL 3216. Insecticide. Oil. $[\alpha]_D -21.9$ (c, 0.8 in CH₂Cl₂). λ_{max} 240 (ϵ 9500); 283 (ϵ 1750); 344 (ϵ 650); 370 (ϵ 570) (MeOH) (Derep). λ_{max} 201 (ϵ 26900); 242 (ϵ 23900); 287 (ϵ 6200) (MeOH) (Berdy). λ_{max} 200 (ϵ 27000); 242 (ϵ 23000); 295 (ϵ 3000) (MeOH/HCl) (Berdy). λ_{max} 213 (ϵ 62700); 242 (ϵ 23000); 295 (ϵ 3000) (MeOH/NaOH) (Berdy).

O-De-Me: **Leporine B**

C₂₂H₂₅NO₃ 351.444

Isol. from fungal strain OSI 54042.

Hexokinase II gene inducing agent.

TePaske, M.R. *et al.*, *Tet. Lett.*, 1991, **32**, 5687-5690 (*isol, pmr, cmr*)

Snider, B.B. *et al.*, *J.O.C.*, 1996, **61**, 2839-2844 (*synth, uv, pmr, cmr*)

Zhang, C. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 1433-1435 (*Leporin B*)

Leptactinine**L-113**

Struct. unknown. Alkaloid from the root bark of *Leptactina senemgambica*. Long needles. Mp 264-266° Mp 258° dec. (as picrate).

Paris, R. *et al.*, *Ann. Pharm. Fr.*, 1946, **4**, 233-236; *CA*, **41**, 7057c (*isol*)

Leptagline**L-114**

N-[3-(Methylthio)-2-propenyl]-N'-(phenylacetyl)putrescine. *Agelaiothioduline*

[261158-69-4]

MeSCH=CHCONH(CH₂)₄NH-COCH₂Ph

C₁₆H₂₂N₂O₂S 306.428

(E)-form

Alkaloid from *Aglaia leptantha* and *Aglaia edulis*. Needles (MeOH). Mp 140-141°. λ_{max} 271 (log ϵ 4.4) (MeOH).


Saifah, E. *et al.*, *Phytochemistry*, 1999, **52**, 1085-1088 (*Agelaiothioduline*)

Greger, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 616-620 (*Leptagine*)

Detterbeck, R. *et al.*, *Tetrahedron*, 2002, **58**, 6887-6893 (*synth*)

Leptanthine†**L-115**

[271586-49-3]



C₁₂H₁₈N₂O₂S₂ 286.418

Alkaloid from *Aglaia leptantha*. $[\alpha]_D^{20} +9$ (c, 0.4 in MeOH). λ_{max} 226 (log ϵ 3.81); 273 (log ϵ 4.26) (MeOH).

1-Deacyl, 1-(phenylacetyl): **Isoagleptine**

[271586-48-2]

C₁₆H₂₀N₂O₂S 304.412

Alkaloid from *Aglaia leptantha*. Mp 149-151°. $[\alpha]_D^{20} -2$ (c, 0.2 in MeOH). λ_{max} 272 (log ϵ 4.26) (MeOH).

1-Deacyl, 1-E-cinnamoyl: **Aglamide A**

[922149-07-3]

C₁₇H₂₀N₂O₂S 316.423

Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. Racemic. λ_{\max} 219 (log ϵ 4.32); 273 (log ϵ 4.6) (MeOH).

1-Deacyl, 1-E-cinnamoyl, S-oxide: **Aglaamide B**

[922149-08-4]

C₁₇H₂₀N₂O₃S 332.423

Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. Racemic. λ_{\max} 206 (log ϵ 4.49); 275 (log ϵ 4.43) (MeOH).

N²-Deacyl, N²-(phenylacetyl): **Agleptine**

[271586-47-1]

C₁₆H₂₀N₂O₂S 304.412

Alkaloid from *Aglaia leptantha*. Mp 121-123°. $[\alpha]_{\text{D}}^{20}$ -3 (c, 0.2 in MeOH). λ_{\max} 282 (log ϵ 4.08) (MeOH).

N²-Deacyl, N²-E-cinnamoyl: **Pyrrrolotenin**

C₁₇H₂₀N₂O₂S 316.423

Alkaloid from the leaves of *Aglaia tenuicaulis*. Cryst. Mp 167-170°. $[\alpha]_{\text{D}}^{20}$ -24 (c, 0.5 in MeOH). λ_{\max} 217 (log ϵ 4.2); 223 (sh) (log ϵ 4.16); 277 (log ϵ 4.48) (MeOH aq.).

Greger, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 616-620 (*Aglaia leptantha* consists)


Kim, S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1769-1775 (*Aglamides A, B*)

Greger, H. *et al.*, *Phytochemistry*, 2008, **69**, 928-938 (*Pyrrrolotenin*)

Leptinidine

L-116

Solanid-5-ene-3,23-diol, 9CI. 23-Hydroxysolanthrene
[24884-17-1]



C₂₇H₄₃NO₂ 413.642

Aglycone obt. by hydrol. of Leptines. Alkaloid from *Solanum chacoense* and other *Solanum* spp. Mp 247-248°. $[\alpha]_{\text{D}}$ -24 (-19) (CHCl₃).

3-O- β -D-Glucopyranoside: **Leptinidine 3-glycoside**

C₃₃H₅₃NO₇ 575.784

Alkaloid from *Solanum orbignianum*. Amorph. powder.

3-O-[α -L-Rhamnopyranosyl-(1 \rightarrow 2)-[α -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside]: **Leptinine I**

C₄₅H₇₃NO₁₅ 868.069

Alkaloid from *Solanum chacoense* and *Solanum orbignianum* (Solanaceae). Mp 230°. $[\alpha]_{\text{D}}$ -90 (Py).

3-O-[β -D-Glucopyranosyl-(1 \rightarrow 3)-[α -L-rhamnopyranosyl-(1 \rightarrow 2)]- β -D-galactopyranoside]: **Leptinine II**

C₄₅H₇₃NO₁₆ 884.069

Alkaloid from *Solanum chacoense* and *Solanum orbignianum* (Solanaceae). Prisms (MeOH aq.). Mp ca.° 255. $[\alpha]_{\text{D}}$ -62 (Py).

3-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-(1 \rightarrow 4)]- β -D-galactopyranoside]: **Solanum Alkaloid SL-c**

C₄₅H₇₃NO₁₇ 900.068

Alkaloid from the stems of *Solanum lyratum* (Solanaceae). Needles (MeOH). Sol. MeOH, H₂O; poorly sol. hexane. Mp 222-225°. $[\alpha]_{\text{D}}^{22}$ -51.5 (c, 0.54 in Py). Actually an inseparable mixt. of the 3-O- β -D-glucopyranosyl (1 \rightarrow 2)- β -D-glucopyranosyl (1 \rightarrow 4)- β -D-galactopyranosides of Leptinidine and 5,6-Dihydroleptinidine below.

3-O-[β -D-Glucopyranosyl-(1 \rightarrow 2)-[β -D-xylopyranosyl-(1 \rightarrow 3)]- β -D-glucopyranosyl-(1 \rightarrow 4)]- β -D-galactopyranoside]: **Solanum Alkaloid SL-d**

C₅₀H₈₁NO₂₁ 1032.184

Alkaloid from *Solanum lyratum* (Solanaceae). Needles (MeOH aq.). Sol. MeOH, H₂O; poorly sol. hexane. Mp 242-246° dec. $[\alpha]_{\text{D}}^{23}$ -62.3 (c, 0.72 in Py). Actually an inseparable mixt. of the Lycotetraosides of Leptinidine and 5,6-Dihydroleptinidine.

Glycoside (Unknown struct. 1): **Leptinine III**

Alkaloid from *Solanum chacoense* (Solanaceae). Not well characterised, sugar residue not identified.

Glycoside (Unknown struct. 2): **Leptinine IV**

Alkaloid from *Solanum chacoense* (Solanaceae). Not well characterised, sugar residue not identified.

23-Ac: [100994-54-5]

Alkaloid from *Solanum chacoense*. Obt. by hydrol. of Leptines. Small needles (MeOH aq.). Mp 192-196°.

23-O-Ac, glycoside (Unknown struct. 1): **Leptine I**

C₄₇H₇₅NO₁₆ 910.107

Alkaloid from *Solanum chacoense* (Solanaceae). Cholinesterase inhibitor. Needles (H₂O). $[\alpha]_{\text{D}}$ -85 (Py). Gives 2 L-rhamnose and 1 D-glucose on hydrol.

23-O-Ac, glycoside (Unknown struct. 2): **Leptine III**

Alkaloid from *Solanum chacoense* (Solanaceae). Not well characterised, sugar residue not identified.

23-O-Ac, glycoside (Unknown struct. 3): **Leptine IV**

Alkaloid from *Solanum chacoense* (Solanaceae). Not well characterised, sugar residue not identified.

Di-O-Ac: **Di-O-acetylleptinidine**

C₃₁H₄₇NO₄ 497.717

Alkaloid from *Solanum chacoense* (Solanaceae). Needles (MeOH). Mp 194-196° (196-200°). $[\alpha]_{\text{D}}$ -36 (CHCl₃).

5,6-Dihydro: 5,6-Dihydrolepidine

C₂₇H₄₅NO₂ 415.658

Identified as glycosides as a component of alkaloids SL-c and SL-d (see above).

5,6-Dihydro, 3-O-[β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-(1 \rightarrow 4)]- β -D-galactopyranoside]:

C₄₅H₇₃NO₁₇ 902.084

A component of Alkaloid SL-c (see

above).

5,6-Dihydro, 3-O-[β -D-glucopyranosyl-(1 \rightarrow 2)]-[β -D-xylopyranosyl-(1 \rightarrow 3)]- β -D-glucopyranosyl-(1 \rightarrow 4)]- β -D-galactopyranoside]:

C₅₀H₈₃NO₂₁ 1034.2

A component of Alkaloid SL-d (see above).

Kuhn, R. *et al.*, *Chem. Ber.*, 1961, **94**, 1088; 1096; 1962, **95**, 1748 (*isol, struct*)

Ripperger, H. *et al.*, *Chem. Ber.*, 1969, **102**, 4080 (*synth*)


Murakami, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 67 (*Alkaloids SL-c, SL-d*)

Lawson, D.R. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 4122-4126 (*pmr, cmr, ms, isol, di-Ac*)

Coelho, R.M. *et al.*, *Phytochemistry*, 1998, **49**, 893-897 (*isol, pmr, cmr, ms*)

Leptocarpine†

L-117



Relative configuration

C₂₁H₂₁NO₆ 383.4


Alkaloid from whole plants of *Hypecoum leptocarpum* (Papaveraceae). Powder (CHCl₃/MeOH). Mp 166-168°. $[\alpha]_{\text{D}}^{25}$ -94.3 (c, 0.6 in MeOH).

Zhang, G.-L. *et al.*, *Phytochemistry*, 1995, **40**, 1813 (*isol, uv, ir, pmr, cmr, ms, struct*)

Leptocarpinine

L-118

[221347-12-2 (chloride)]



C₂₂H₂₀NO₆⁺ 394.403


Alkaloid from *Hypecoum leptocarpum*. Amorph. powder (as chloride). Mp > 350° (chloride). λ_{\max} 259 (log ϵ 4.23); 303 (log ϵ 4.01); 376 (log ϵ 3.94) (MeOH) (chloride).

Yi, Z. *et al.*, *Phytochemistry*, 1999, **50**, 339-343 (*isol, uv, ir, pmr, cmr, ms*)

Leptophyllin A

L-119

12-(5-Hydroxy-6-methyl-2-piperidinyl)-1,2-dodecanediol, 9CI
[165689-18-9]



C₁₈H₃₇NO₃ 315.495

Alkaloid from leaves of *Cassia leptophylla* (Fabaceae). Oil. $[\alpha]_D +2.5$ (c, 0.02 in CHCl_3).

O³-Ac: 3-Acetylleptophyllin A

[165404-55-7]


$\text{C}_{20}\text{H}_{39}\text{NO}_4$ 357.532

From leaves of *Cassia leptophylla* (Fabaceae). Oil. $[\alpha]_D +3$ (c, 0.03 in CHCl_3).

Bolzani, V. da.S. *et al.*, *Tetrahedron*, 1995, **51**, 5929 (isol, pmr, cmr, ms, struct)

Leptophyllin B L-120

5-Hydroxy-6-methyl-2-piperidineundecanoic acid, 9CI
[165689-19-0]




$\text{C}_{17}\text{H}_{33}\text{NO}_3$ 299.453

Alkaloid from leaves of *Cassia leptophylla* (Fabaceae). Oil. $[\alpha]_D -3.6$ (c, 0.03 in MeOH).

Bolzani, V. da.S. *et al.*, *Tetrahedron*, 1995, **51**, 5929 (isol, pmr, cmr, ms, struct)

Leptopidine L-121

[221347-15-5]



$\text{R}^1, \text{R}^2 = -\text{CH}_2-$, $\text{R}^3 = \text{H}$, $\text{R}^4 = \text{CH}_3$

$\text{C}_{20}\text{H}_{20}\text{NO}_6^{\oplus}$ 370.381

Alkaloid from *Hypocoum leptocarpum*. Yellow powder (MeOH) (as chloride). Mp 235-239° (chloride). λ_{max} 243 (log ϵ 5.41); 305 (log ϵ 4.05); 352 (log ϵ 4.05) (MeOH) (chloride).

Zhou, Y. *et al.*, *Phytochemistry*, 1999, **50**, 339-343 (isol, uv, ir, pmr, cmr, ms)

Leptopine L-122

[221347-13-3]

As Leptopidine, L-121 with

$\text{R}^1, \text{R}^2 = \text{R}^3, \text{R}^4 = -\text{CH}_2-$

$\text{C}_{20}\text{H}_{18}\text{NO}_6^{\oplus}$ 368.365

Alkaloid from *Hypocoum leptocarpum*. Yellow cubes (MeOH) (as chloride). Mp 158-161° (chloride). λ_{max} 248 (log ϵ 4.25); 300 (log ϵ 3.84); 365 (log ϵ 3.94) (MeOH) (chloride).

1-Hydro: Dihydroleptopine

[214537-74-3]

$\text{C}_{20}\text{H}_{19}\text{NO}_6$ 369.373

Alkaloid from *Hypocoum leptocarpum*. Amorph. powder (MeOH). Mp 185-189°. Racemic.

Zhou, Y. *et al.*, *Phytochemistry*, 1999, **50**, 339-

343 (isol, uv, ir, pmr, cmr, ms)

Li, B.-G. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 1215-1218 (*Dihydroleptopine*)

Leptopinine L-123

[221347-14-4]

As Leptopidine, L-121 with

$\text{R}^1 = \text{R}^2 = \text{H}$, $\text{R}^3 = \text{R}^4 = \text{CH}_3$


$\text{C}_{20}\text{H}_{22}\text{NO}_6^{\oplus}$ 372.397

Alkaloid from *Hypocoum leptocarpum*. Yellow powder (MeOH) (as chloride). Mp 207-210° (chloride). λ_{max} 245 (log ϵ 4.22); 310 (log ϵ 3.97); 353 (log ϵ 3.97) (MeOH) (chloride).

Zhou, Y. *et al.*, *Phytochemistry*, 1999, **50**, 339-343 (isol, uv, ir, pmr, cmr, ms)

Leptosin A L-124

[159518-74-8]



x = 2

Absolute Configuration

$\text{C}_{32}\text{H}_{32}\text{N}_6\text{O}_7\text{S}_6$ 805.037
Metab. from a strain of *Leptosphaeria* sp. originally isol. from the marine alga *Sargassum tortile*. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells and significant antitumour activity against Sarcoma 180 ascites. Pale yellow powder. Sol. MeOH, CHCl_3 ; poorly sol. hexane. Mp 216-218°. $[\alpha]_D +237$ (c, 0.49 in CHCl_3). λ_{max} 209 (ϵ 40700); 242 (ϵ 15500); 298 (ϵ 6750) (EtOH) (Berdy).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1994, 1859-1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin B L-125

[159518-75-9]

As Leptosin A, L-124 with

x = 3, y = 2

$\text{C}_{32}\text{H}_{32}\text{N}_6\text{O}_7\text{S}_5$ 772.971

Metab. from a strain of *Leptosphaeria* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Sol. MeOH, CHCl_3 ; poorly sol. hexane. Mp 210-213°. $[\alpha]_D +392$ (c, 0.50 in CHCl_3). λ_{max} 208 (ϵ 28200); 244 (ϵ 10000); 299 (ϵ 4800) (EtOH) (Berdy).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1994, 1859-1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin C L-126

[159518-76-0]

As Leptosin A, L-124 with

x = 2, y = 2

$\text{C}_{32}\text{H}_{32}\text{N}_6\text{O}_7\text{S}_4$ 740.905


Metab. from a strain of *Leptosphaeria* sp.

and a *Sesquicillium* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells and significant antitumour activity against Sarcoma 180 ascites. Pale yellow powder. Sol. MeOH, CHCl_3 ; poorly sol. hexane. Mp 208-210°. $[\alpha]_D +237$ (c, 0.36 in CHCl_3). λ_{max} 206 (ϵ 60400); 283 (ϵ 17000); 301 (ϵ 5150) (EtOH) (Berdy).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1994, 1859-1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin D L-127

[159518-77-1]



x = 4, y = 2

Absolute Configuration

$\text{C}_{25}\text{H}_{24}\text{N}_4\text{O}_3\text{S}_2$ 492.622

Metab. from the mycelium of a strain of *Leptosphaeria* sp. attached to the marine alga *Sargassum tortile*. Also from *Sesquicillium* sp. Shows potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Sol. MeOH, CHCl_3 ; poorly sol. hexane. Mp 190-192°. $[\alpha]_D +436$ (c, 0.51 in CHCl_3). λ_{max} 206 (ϵ 39800); 219 (ϵ 41600); 240 (ϵ 11450); 272 (ϵ 5900); 282 (ϵ 6750); 290 (ϵ 6750) (EtOH) (Berdy).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1994, 1859-1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin E L-128

[159518-78-2]

As Leptosin D, L-127 with

x = 3

$\text{C}_{25}\text{H}_{24}\text{N}_4\text{O}_3\text{S}_3$ 524.688

Metab. from a strain of *Leptosphaeria* sp. and by *Sesquicillium* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Sol. MeOH, CHCl_3 ; poorly sol. hexane. Mp 229-231°. $[\alpha]_D +563$ (c, 0.32 in CHCl_3). λ_{max} 206 (ϵ 42600); 218 (ϵ 44700); 240 (ϵ 13150); 273 (ϵ 6700); 282 (ϵ 6920); 291 (ϵ 6490) (EtOH) (Berdy).

Takahashi, C. *et al.*, *J.C.S. Perkin 1*, 1994, 1859-1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin F L-129

[159518-79-3]

As Leptosin D, L-127 with

x = 4

$\text{C}_{25}\text{H}_{24}\text{N}_4\text{O}_3\text{S}_4$ 556.754

Metab. from a strain of *Leptosphaeria* sp. and *Sesquicillium* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 219–221°. [α]_D²⁵ +452 (c, 0.39 in CHCl₃). λ_{max} 206 (ε 45700); 216 (ε 49000); 240 (ε 15500); 272 (ε 7550); 281 (ε 7950); 290 (ε 7950) (EtOH) (Berdy).

Takahashi, C. et al., *J.C.S. Perkin 1*, 1994, 1859–1864 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin G₁ L-130

[162232-33-9]
As Leptosin A, L-124 with
x = 3, y = 3

C₃₂H₃₂N₆O₇S₆ 805.037
Metab. from a strain of *Leptosphaeria* sp. and *Sargassum tortile*. Exhibits cytotoxicity against P388 lymphocytic leukaemia cells. Powder. Mp 210–212°. [α]_D²⁴ +558 (c, 0.45 in CHCl₃). λ_{max} 210 (ε 52480); 240 (ε 13800); 296 (ε 6166) (EtOH) (Berdy).

Takahashi, C. et al., *Phytochemistry*, 1995, 38, 155–158 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin G₂ L-131

[159334-39-1]
As Leptosin A, L-125 with
x = 2, y = 3

C₃₂H₃₂N₆O₇S₅ 772.971
Metab. from a strain of *Leptosphaeria* sp. and *Sargassum tortile*. Exhibits cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Mp 210–215°. [α]_D²⁴ +303 (c, 0.58 in CHCl₃). λ_{max} 206 (ε 58000); 240 (ε 20417); 296 (ε 6310) (EtOH) (Berdy).

Takahashi, C. et al., *Phytochemistry*, 1995, 38, 155–158 (isol, uv, ir, pmr, ms, cd, struct)

Leptosin G L-132

[159334-43-7]
As Leptosin A, L-124 with
x = 4, y = 3

C₃₂H₃₂N₆O₇S₇ 837.103
Metab. of a strain of *Leptosphaeria* sp. and *Sargassum tortile*. Exhibits cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Mp 205–210°. [α]_D²⁴ +481 (c, 0.40 in CHCl₃). λ_{max} 206 (ε 81283); 240 (ε 26300); 296 (ε 8130) (EtOH) (Berdy).

Takahashi, C. et al., *Phytochemistry*, 1995, 38, 155 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin H L-133


[159334-40-4]
As Leptosin A, L-124 with
x = 2, y = 4

C₃₂H₃₂N₆O₇S₆ 805.037
Metab. from a strain of *Leptosphaeria* sp. and *Sargassum tortile*. Exhibits cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Mp 214–215°. [α]_D²⁴ +298 (c, 0.47 in CHCl₃). λ_{max} 207 (ε 64565); 240 (ε 23442); 302 (ε 7080) (EtOH) (Berdy).

Takahashi, C. et al., *Phytochemistry*, 1995, 38, 155–158 (isol, uv, ir, pmr, cmr, ms, cd, struct)

Leptosin I L-134

[160472-96-8]



Relative Configuration

C₃₂H₃₂N₆O₇S₄ 740.905
Metab. from a strain of *Leptosphaeria* sp. originally isol. from the marine alga *Sargassum tortile*. Cytotoxic agent. Pale yellow powder. Sol. DMSO, CHCl₃, MeOH; poorly sol. H₂O, hexane. Mp 218–220°. [α]_D²⁴ +212 (c, 0.13 in CHCl₃). λ_{max} 209 (log ε 4.45); 240 (log ε 3.91); 305 (log ε 3.45) (EtOH).


3'-Epimer: Leptosin J

[160550-15-2]
C₃₂H₃₂N₆O₇S₄ 740.905
From *Leptosphaeria* sp. Cytotoxic agent. Pale yellow powder. Sol. DMSO, MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 215–216°. [α]_D²⁴ +188 (c, 0.21 in CHCl₃). λ_{max} 209 (log ε 4.45); 240 (log ε 3.94); 298 (log ε 3.44) (EtOH).

Takahashi, C. et al., *J. Antibiot.*, 1994, 47, 1242–1249 (isol, uv, ir, pmr, cmr, cd, struct)

Leptosin K₁ L-135

[159334-36-8]



Absolute Configuration

C₃₄H₃₆N₆O₆S₅ 785.025
Metab. from a strain of *Leptosphaeria* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Mp 209–212°. [α]_D +88.9 (c, 0.32 in CHCl₃). λ_{max} 207 (ε 51286); 237 (ε 19500); 300 (ε 5495) (EtOH) (Berdy).

Takahashi, C. et al., *Tetrahedron*, 1995, 51, 3483–3498 (isol, uv, ir, pmr, ms, cd, cryst struct)

Leptosin K₂ L-136

[159334-37-9]
As Leptosin K₁, L-135 with
x = 4

C₃₄H₃₆N₆O₆S₆ 817.091
Metab. from a strain of *Leptosphaeria* sp. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Pale yellow powder. Mp 214–216°. [α]_D +482.8 (c, 0.44 in CHCl₃). λ_{max} 205 (ε 34950); 242 (ε 14125); 296 (ε 4365) (EtOH) (Berdy).

Takahashi, C. et al., *Tetrahedron*, 1995, 51, 3483–3498 (isol, uv, ir, pmr, cmr, ms, cd, cryst struct)

Leptosin K L-137


[159334-35-7]
As Leptosin K₁, L-135 with
x = 2

C₃₄H₃₆N₆O₆S₄ 752.959
Metab. from a strain of *Leptosphaeria* sp. OUPS-4 originally isolated from the marine alga *Sargassum tortile*. Exhibits potent cytotoxicity against P388 lymphocytic leukaemia cells. Prisms (EtOAc). Mp 222–224°. [α]_D +76.7 (c, 0.37 in CHCl₃). λ_{max} 206 (ε 60256); 240 (ε 19050); 302 (ε 6025) (EtOH).

Takahashi, C. et al., *Tetrahedron*, 1995, 51, 3483–3498 (isol, uv, ir, pmr, cmr, ms, cd, cryst struct)

Leptosin M₁ L-138

[406213-88-5]



Absolute Configuration

C₃₃H₃₆N₆O₈S₂ 708.815
Metab. of *Leptosphaeria* sp. OUPS-4. Pale yellow powder. Mp 219–222°. [α]_D +140 (c, 0.18 in CHCl₃). λ_{max} 212 (log ε 4.73); 238 (log ε 4.3); 300 (log ε 3.83) (EtOH).

Yamada, T. et al., *Tetrahedron*, 2002, 58, 479–487 (isol, pmr, cmr, uv, ir, cd, struct)

Leptosin M L-139

[406213-87-4]
As Leptosin M₁, L-138 with
n = 4

C₃₃H₃₆N₆O₈S₄ 772.947
Metab. of *Leptosphaeria* sp. OUPS-4 isol. from the marine alga *Sargassum tortile*. Pale yellow powder. Mp 223–226°. [α]_D +478 (c, 0.1 in CHCl₃). λ_{max} 206 (log ε 4.64); 243 (log ε 4.08); 304 (log ε 3.62) (EtOH).

11,11'-Diepimer: Leptosin N

[406213-89-6]

C₃₃H₃₆N₆O₈S₄ 772.947

Metab. of *Leptosphaeria* sp. OUPS-4. Pale yellow powder. Mp 226-228°. [α]_D +276 (c, 0.16 in CHCl₃). λ_{max} 212 (log ε 4.75); 240 (log ε 4.36); 302 (log ε 3.86) (EtOH).

Yamada, T. et al., *Tetrahedron*, 2002, **58**, 479-487 (isol, uv, pmr, cmr, cd, struct)

Leptosin N₁**L-140**

[406213-90-9]


As Leptosin M₁, L-138 with n = 3C₃₃H₃₆N₆O₈S₃ 740.881

Metab. of *Leptosphaeria* sp. OUPS-4. Pale yellow powder. Mp 227-229°. [α]_D +347 (c, 0.14 in CHCl₃). λ_{max} 212 (log ε 4.74); 240 (log ε 4.34); 302 (log ε 3.82) (EtOH).

Yamada, T. et al., *Tetrahedron*, 2002, **58**, 479-487 (isol, uv, ir, pmr, cmr, cd, struct)

Leptosin O**L-141**

[721960-15-2]



R = -S-S-Me

C₃₃H₃₆N₆O₇S₂ 692.815

Metab. of *Leptosphaeria* sp. OUPS-N80 isol. from *Sargassum tortile*. Cytotoxic. Pale yellow powder. Mp 220-222°. [α]_D -99 (c, 0.08 in CHCl₃). λ_{max} 211 (log ε 4.76); 238 (log ε 4.58); 265 (log ε 4.3); 298 (log ε 4.08) (EtOH).

3-Epimer: Leptosin P

[721960-16-3]

C₃₃H₃₆N₆O₇S₂ 692.815

Metab. of *Leptosphaeria* sp. OUPS-N80 from *Sargassum tortile*. Cytotoxic. Pale yellow powder. Mp 233-235°. [α]_D +35 (c, 0.17 in CHCl₃). λ_{max} 209 (log ε 4.49); 238 (log ε 4.3); 270 (log ε 3.99); 300 (log ε 3.78) (EtOH).

Yamada, T. et al., *Heterocycles*, 2004, **63**, 641-652 (isol, cd, pmr, cmr, ms)

Leptosin Q**L-142**

[721960-17-4]

As Leptosin O, L-141 with

R = -SMe

C₃₃H₃₆N₆O₇S 660.749

Metab. of *Leptosphaeria* sp. OUPS-N80 isol. from *Sargassum tortile*. Pale yellow

powder. Mp 231-233°. [α]_D -92 (c, 0.07 in CHCl₃). λ_{max} 212 (log ε 4.76); 238 (log ε 4.32); 268 (log ε 4.02); 299 (log ε 3.75) (EtOH).

3-Epimer: Leptosin R

[721960-18-5]

C₃₃H₃₆N₆O₇S 660.749

Metab. of *Leptosphaeria* sp. OUPS-N80 isol. from *Sargassum tortile*. Pale yellow powder. Mp 221-223°. [α]_D -24 (c, 0.1 in CHCl₃). λ_{max} 212 (log ε 4.95); 238 (log ε 4.72); 267 (log ε 4.35); 298 (log ε 4.14) (EtOH).

Yamada, T. et al., *Heterocycles*, 2004, **63**, 641-652 (isol, pmr, cmr, ms)

Leptosin S**L-143**

[721960-19-6]

As Leptosin O, L-141 with

R = NH₂C₃₂H₃₅N₇O₇ 629.671

Metab. of *Leptosphaeria* sp. OUPS-N80 isol. from *Sargassum tortile*. Pale yellow powder. Mp 231-233°. [α]_D -56 (c, 0.07 in CHCl₃). λ_{max} 212 (log ε 4.77); 238 (log ε 4.61); 270 (log ε 4.1); 300 (log ε 4.04) (EtOH).

Yamada, T. et al., *Heterocycles*, 2004, **63**, 641-652 (isol, pmr, cmr, ms)

Lettocine**L-144**C₁₇H₂₅NO₂ 275.39

Struct. unknown. Alkaloid from *Holarhena antidysenterica* bark (Apocynaceae). Light-brown cryst. powder (CHCl₃/petrol). Mp 350-352°.

Hydroiodide:

Yellowish-brown cryst. powder (EtOH aq.). Mp 256° dec.


Methiodide:

Cryst. (MeOH). Mp 235°.

Picrate:

Cryst. (EtOH). Mp 198°.

Peacock, D.H. et al., *J.C.S.*, 1935, 734-735

Lettowianthine**L-145***Annobraine*C₁₉H₁₁NO₄ 317.3

Alkaloid from the root bark of *Letto-wianthus stellatus* and from *Annona glabra* (pond apple) (Annonaceae). Dark red solid (CHCl₃); red needles (CHCl₃). Mp 314-317° dec. (265-267°). λ_{max} 254 (log ε 4.02); 293 (log ε 3.42); 323 (sh) (log ε 3.5); 333 (log ε 3.55); 365 (log ε 3.23); 509 (log ε 3.07) (MeOH) (Lettowianthine). λ_{max} 232 (sh) (log ε 3.97); 253 (log ε 4.37); 280 (log ε 3.8); 291 (log ε 3.84); 333 (log ε 3.64); 355 (sh) (log ε 3.43); 373 (log ε 3.2) (EtOH) (Annobraine).

11-Hydroxy: **11-Hydroxylettowianthine**

C₁₉H₁₁NO₅ 333.3


Alkaloid from the root bark of *Letto-wianthus stellatus* (Annonaceae). Dark red amorph. solid. λ_{max} 260 (log ε 4.09); 292 (log ε 3.48); 311 (sh) (log ε 3.28); 355 (log ε 3.25); 505 (log ε 3.02) (MeOH).

Chang, F.R. et al., *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 913-920 (*Annobraine*)

Nkunya, M.H.H. et al., *Phytochemistry*, 2000, **53**, 1067-1073 (*Lettowianthine*, *11-Hydroxylettowianthine*)

Leucamide A**L-146**

[447461-16-7]

C₂₉H₃₇N₇O₆S 611.721


Isol. from the Australian sponge *Leucetta microraphis*. Cytotoxic agent. Amorph. solid. [α]_D²³ -69.3 (c, 0.6 in CHCl₃). λ_{max} 246 (ε 9270) (CHCl₃).

Kehraus, S. et al., *J.O.C.*, 2002, **67**, 4989-4992 (isol, pmr, cmr)

Wang, W. et al., *J.O.C.*, 2003, **68**, 1636-1639 (*synth*)

Leucascandrolide A**L-147**

[175448-18-7]



Absolute Configuration

C₃₈H₅₆N₂O₁₀ 700.868

Isol. from the marine sponge *Leucascandra caveolata*. Shows potent antifungal activity against *Candida albicans*; shows strong cytotoxicity. Amorph. solid. Sol. MeOH, CHCl₃, CH₂Cl₂, EtOH; poorly sol. H₂O. [α]_D²⁰ +41 (EtOH). λ_{max} 205 (ε 15000); 262 (ε 11000) (EtOH).

D'Ambrosio, M. et al., *Helv. Chim. Acta*, 1996, **79**, 51-60 (isol, uv, ir, pmr, cmr, ms, *cryst struct*, *activity*)

Hornberger, K.R. et al., *J.A.C.S.*, 2000, **122**, 12894-12895 (*synth*, *activity*)

Wang, Y. et al., *J.A.C.S.*, 2002, **124**, 13670-13671 (*synth*)

Williams, D.R. et al., *Angew. Chem., Int. Ed.*, 2003, **42**, 3934-3938 (*synth*)


Fettes, A. et al., *J.O.C.*, 2003, **68**, 9274-9283 (*synth*)

Crimmins, M.T. et al., *Org. Lett.*, 2003, **5**, 4641-4644 (*synth*)

- Williams, D.R. *et al.*, *Org. Lett.*, 2003, **5**, 5035-5038 (*synth*)
 Paterson, I. *et al.*, *Tetrahedron*, 2003, **59**, 6833-6849 (*synth*)
 Su, Q. *et al.*, *J.O.C.*, 2007, **72**, 2-24 (*synth*)
 Van Orden, L.J. *et al.*, *J.O.C.*, 2007, **72**, 5784-5793 (*synth*)
 Ferrié, L. *et al.*, *Org. Lett.*, 2007, **9**, 2461-2464 (*synth*)

Leucenine L-148

α -Amino-3-hydroxy-1-(4H)-pyridinepropionic acid, 9CI. Leucenol. Leucaenine. Leucaenol. Mimosine



C₈H₁₀N₂O₄ 198.178
 Possesses depilatory props.

(S)-form [500-44-7]

Alkaloid from *Leucaena glauca* and *Mimosa pudica*. Cryst. (H₂O). Sol. H₂O. Mp 235-236°. [α]_D²² -20 (H₂O). p*K*_{a1} 2.1; p*K*_{a2} 7.2; p*K*_{a3} 9.2 (phenolic OH). λ _{max} 282 (ε 16980) (MeOH). λ _{max} 284 (ε 15000) (H₂O) (Berdy). λ _{max} 277 (HCl).

O-β-D-Glucopyranoside: **Mimoside** [36518-12-4]

C₁₄H₂₀N₂O₉ 360.32

Alkaloid from *Mimosa pudica* and *Leucaena leucocephala*. Mp 178-179°. [α]_D²⁶ -60.9 (c, 0.46 in H₂O).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 597D (*ir*)

Aldrich Library of NMR Spectra, 2nd edn., 1983, **1**, 506D (*nmr*)

Wibaut, J.P. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1946, **65**, 65; 1950, **69**, 37 (*struct*)

Bickel, A.F. *et al.*, *J.A.C.S.*, 1948, **70**, 326 (*struct*)

Adams, R. *et al.*, *J.A.C.S.*, 1949, **71**, 705 (*synth*)

Spenser, I.D. *et al.*, *Can. J. Chem.*, 1962, **40**, 1374 (*synth*)

Hylin, J.W. *et al.*, *Phytochemistry*, 1964, **3**, 161 (*biosynth*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1964, **83**, 1078 (*abs config*)

Murakoshi, T. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2655


Mostad, A. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 164 (*cryst struct*)

Mostad, A. *et al.*, *Acta Chem. Scand., Ser. B*, 1974, **28**, 249 (*cryst struct*)

Murakoshi, I. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1473 (*synth*)

Leucettamidine L-149

[147395-97-9]




C₂₅H₂₄N₆O₅ 488.502

Alkaloid from the marine sponge *Leucetta microraphis*. Exhibits significant leukotriene B₄ receptor binding activity. Antiinflammatory agent. λ _{max} 205 (ε 4200); 285 (ε 800); 386 (ε 6000) (CHCl₃/MeOH) (Derep). λ _{max} 205 ; 237 ; 363 (MeOH) (Berdy).

Chan, G.W. *et al.*, *J. Nat. Prod.*, 1993, **56**, 116 (*isol, pmr, struct*)

Leucettamine A L-150

4,5-Bis(1,3-benzodioxol-5-ylmethyl)-1-methyl-1H-imidazol-2-amine, 9CI [147395-95-7]



C₂₀H₁₉N₃O₄ 365.388


Alkaloid from the marine sponge *Leucetta microraphis*. Exhibits potent leukotriene B₄ receptor binding activity. Antiinflammatory agent. Yellowish amorph. solid. λ _{max} 205 (ε 4280); 285 (ε 796) (MeOH) (Derep).

Boehm, J.C. *et al.*, *J. Med. Chem.*, 1993, **36**, 3333 (*synth*)

Chan, G.W. *et al.*, *J. Nat. Prod.*, 1993, **56**, 116 (*isol, uv, ir, pmr, cmr, ms, struct, activity*)

Leucettamine B L-151

[147395-96-8]



C₁₂H₁₁N₃O₃ 245.237

Alkaloid from the marine sponge *Leucetta microraphis*. Cream solid. λ _{max} 205 (ε 2320); 237 (ε 1830); 363 (ε 3710) (MeOH) (Derep).


Chan, G.W. *et al.*, *J. Nat. Prod.*, 1993, **56**, 116 (*isol, uv, ir, pmr, cmr, ms, struct*)

Molina, P. *et al.*, *Tet. Lett.*, 1994, **35**, 2235 (*synth*)

Roue, N. *et al.*, *Tetrahedron*, 1999, **55**, 14729-14738 (*synth*)

Leucettamine C L-152

2-Amino-3,5-dihydro-5-[(4-methoxyphenyl)methylidene]-3-methyl-4H-imidazol-4-one, 9CI [497085-85-5]



C₁₂H₁₃N₃O₂ 231.254

Alkaloid from the sponge *Leucetta* sp. Amorph. yellow powder. λ _{max} 207 (log ε 4.26); 230 (log ε 4.1); 368 (log ε 4.4) (MeOH).

Crews, P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 177-182 (*isol, pmr, cmr*)

Leucine, 9CI, USAN L-153

2-Amino-4-methylpentanoic acid. 2-Aminoisocaproic acid. 4-Methylnorvaline. *Leu. Chenopodine* [7005-03-0]



C₆H₁₃NO₂ 131.174

[α]_D¹⁶ +14 (c, 1.3 in H₂O). Log P -1.67 (calc). Derivs. useful in peptide synth. are listed elsewhere.

(R)-form

D-form

[328-38-1]

Isol. from aerial parts of *Coronilla varia* and seeds of *Coronilla scorpioides*. Plates (EtOH). Mp 293° (sealed tube). [α]_D²⁰ +11.1 (H₂O). [α]_D²⁰ -17 (20% HCl).

▶ OH2840000

Me ester: [23032-21-5]

C₇H₁₅NO₂ 145.201

Mp 149-150° (as hydrochloride).

Amide: [15893-47-7]

C₆H₁₄N₂O 130.189

Solid (as hydrochloride). [α]_D -11 (c, 0.5 in MeOH).

N-Ac: [19764-30-8]

C₈H₁₅NO₃ 173.211

Mp 112-114°. [α]_D²³ +23.2 (EtOH).

N-Benzoyl: [57357-55-8]

C₁₃H₁₇NO₃ 235.282

Cryst. + 1/2 H₂O (Et₂O/petrol). Spar. sol. hot H₂O. Mp 60° Mp 105-107° (anhyd.). [α]_D²⁰ -6.39 (KOH).

Homopolymer: *D-Leucine homopolymer*.

Poly-D-leucine

[26335-59-1] Catalyst for the asymmetric epoxidation of unsaturated ketones.

Solid.


N-(2-Nitrobenzenesulfonyl): N-[(2-

- Nitrophenylthio*]-L-leucine
[50571-61-4]
[7685-67-8, 7675-50-5]
C₁₂H₁₆N₂O₄S 284.335
Solid. Mp 108-109° (102-106°). [α]_D²⁰ -112.7 (c, 2 in DMF). Also obt. as dicyclohexylamine salt.
- (S)-form**
L-form. FEMA 3297
[61-90-5]
Obt. by hydrol. of most proteins. Flavouring ingredient. Dietary supplement; nutrient. Plates (EtOH aq.). Spar. sol. H₂O (2.3 g/100 cm³ at 25°). Mp 337° (293-295°) dec. [α]_D²⁰ -10.42 (H₂O). [α]_D¹⁵ +17.3 (20% HCl). pK_{a1} 2.33; pK_{a2} 9.74 (25°). Log P -1.67 (calc). Isoelectric point 5.98. Bitter taste.
► Exp. reprod. and teratogenic effects (v. large doses). OH2850000
Me ester: [2666-93-5]
Bp₁₂ 79-79.5°. [α]_D¹⁷ +16.52.
Me ester, hydrochloride: [7517-19-3]
Mp 118° (α-form) Mp 150-151° (β-form). [α]_D¹⁶ +14 (c, 1.3 in H₂O).
Et ester: [2743-60-4]
C₈H₁₇NO₂ 159.228
Bp₇₆₁ 196° Bp₁₈ 88°.
Et ester, hydrochloride: [2743-40-0]
Prisms (EtOAc/petrol). Mp 134°. [α]_D +18.4.
tert-Butyl ester: [21691-53-2]
C₁₀H₂₁NO₂ 187.281
Bp_{0.15} 45°. [α]_D²⁵ +21.6 (c, 2.5 in EtOH).
tert-Butyl ester, hydrochloride: [42031-13-0]
Cryst. (EtOH/Et₂O). Mp 173°. [α]_D²⁰ +19.8 (c, 2 in EtOH).
Benzyl ester: [1738-69-8]
C₁₃H₁₉NO₂ 221.299
Oil.
Benzyl ester, hydrochloride: [2462-35-3]
Mp 128°. [α]_D³⁰ -8 (c, 2 in HCl).
Amide: 2-Amino-4-methylpentanamide, 9CI. Leucine amide. Leucinamide
[687-51-4]
C₆H₁₄N₂O 130.189
Constit. of the seeds of *Trifolium repens* var. *giganteum*. Mp 254-256° (as hydrochloride). [α]_D²⁰ +10 (c, 5 in H₂O).
(2-Naphthyl)amide: (S)-2-Amino-4-methyl-N-2-naphthalenylpentanamide, 9CI
[732-85-4]
C₁₆H₂₀N₂O 256.347
Mp 105-108°.
Anilide:
C₁₂H₁₈N₂O 206.287
Cryst. (as hydrochloride). Mp 218-222°.
N-Formyl, Me ester: [54259-27-7]
C₈H₁₅NO₃ 173.211
Oil. [α]_D -43.2 (c, 1 in MeOH).
N-Ac: [1188-21-2]
Mp 189-190°. [α]_D²⁴ -40.2 (Py). [α]_D²⁴ -23.1 (EtOH).
N-(Chloroacetyl): [688-12-0]
C₈H₁₄ClNO₃ 207.656
Cryst. Mp 136°. [α]_D -15.8 (EtOH).
- N-Benzoyl: [1466-83-7]
Mp 60° (hydrated) Mp 105-107° (anhyd.).
N-Benzoyl, Me ester: [3005-60-5]
C₁₄H₁₉NO₃ 249.309
Mp 80-82°. [α]_D +38 (c, 1 in CHCl₃).
N-(3,5-Dinitrobenzoyl): [7495-01-4]
C₁₃H₁₅N₃O₇ 325.277
Used in chromatographic resolu. of enantiomers. Mp 186-188°. [α]_D²⁰ -14.3 (c, 3 in EtOH).
4-Nitrobenzoyloxycarbonyl: [122411-88-5]
C₁₄H₁₈N₂O₆ 310.306
Solid + H₂O. Mp 60-61° (monohydrate) Mp 81-85.5° (anhydr.). [α]_D²⁷ -15.8 (c, 1 in 1N NaOH).
N-Me: N-Methylleucine, 9CI. 4-Methyl-2-(methylamino)pentanoic acid. 2-Methylaminoisocaproic acid
[3060-46-6]
C₇H₁₅NO₂ 145.201
Residue present in antibiotics, e.g. Ilamycins, Sporidesmolides. Cryst. (Me₂CO aq.). Mp 200° subl. [α]_D +21.4 (H₂O). [α]_D +31.3 (5M HCl).
N-Me, Me ester: [35026-08-5]
C₈H₁₇NO₂ 159.228
Mp 126.5-127° (as hydrochloride). [α]_D²⁵ +31.4 (c, 1 in EtOH).
N-Me, N-tert-butylloxycarbonyl: [53363-89-6]
C₁₂H₂₃NO₄ 245.318
Mp 56-57°. [α]_D²⁶ -24.6 (c, 0.5 in EtOH).
N-Me, N-benzylloxycarbonyl: [33099-08-0]
C₁₅H₂₁NO₄ 279.335
Mp 74-75°.
N-(2,4-Dinitrophenyl): [1655-57-8]
Mp 94-95°.
N-(2,4,6-Trinitrophenyl):
C₁₂H₁₄N₄O₈ 342.265
Cryst. (MeOH aq.). Mp 98-99°.
Polymer with 5-methylglutamate: **Leuciglumer, INN**
[41385-14-2]
Dermatological agent. [α]_D²⁰ -44 (c, 0.01 in HCl aq.).
Homopolymer: L-Leucine homopolymer.
Poly-L-leucine
[25248-98-0]
Catalyst for the asymmetric epoxidation of unsaturated ketones.
Solid. [α]_D²⁰ -103.4 (c, 0.32 in CF₃CO₂H).
N-Jasmonoyl: N-Jasmonoylleucine
[120330-91-8]
C₁₈H₂₉NO₄ 323.431
Prod. by rice (*Oryza sativa*) leaves after wounding. Cryst. (CHCl₃/hexane). Mp 145-147° (synthetic). [α]_D²⁵ -64.9 (c, 0.23 in MeOH) (synthetic).
- (±)-form** [328-39-2]
Dietary supplement, nutrient. Plates (H₂O). V. spar. sol. EtOH. Mp 293-295° (sealed tube). pK_{a1} 2.33; pK_{a2} 9.74 (25°).
Et ester: [2899-43-6] Misc. EtOH, Et₂O, C₆H₆, petrol. Bp₇₆₁ 196° Bp₁₈ 88°.
- Amide: [13079-20-4]
Prisms (C₆H₆). Sol. H₂O; v. sol. EtOH, Me₂CO; spar. sol. C₆H₆.
N-Ac: **Acetylleucine, INN. Tanganil. RP 7452**
[99-15-0] Drug used to treat dizziness, antiemetic. Mp 160-162° (157-158°). Log P 0.84 (calc).
N-Ac, amide:
C₈H₁₆N₂O₂ 172.227
Mp 157-158°.
N-(Chloroacetyl): [7154-83-8]
Cryst. Mp 142°.
N-Benzoyl: [17966-67-5]
Needles or plates. Sol. EtOH, Et₂O, CHCl₃, AcOH; spar. sol. C₆H₆; insol. petrol.
N-Benzoyl, Me ester: [38449-08-0]
C₁₄H₁₉NO₃ 249.309
Cryst. (petrol). Mp 95-96°.
N-Benzoyl, amide:
C₁₃H₁₈N₂O₂ 234.297
Plates (petrol). Mp 171°.
Nitrile: [65707-62-2]
C₆H₁₂N₂ 112.174
Mp 170-172° (as hydrochloride) Mp 203-208° (as hydrochloride).
N-Me: [2566-33-8]
Needles (EtOH aq.). Sublimes.
N-(2,4-Dinitrophenyl): [10484-03-4]
Yellow cryst. Mp 203°.
[13366-40-0, 5845-53-4, 10466-61-2]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 575A; 575C; 575B (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 875B (nmr)
Fischer, E. et al., Ber., 1900, 33, 2370; 2377; 1901, 34, 445; 1905, 38, 615; 3997; 4002
Ronwin, E. et al., J.O.C., 1953, 18, 127 (N-chloroacetyl)
Roeske, R.W. et al., Chem. Ind. (London), 1959, 1121-1127 (tert-butyl ester)
Russell, D.W. et al., Biochim. Biophys. Acta, 1960, 38, 382; 45, 411 (N-Me)
Taschner, E. et al., Annalen, 1961, 646, 134-136 (tert-butyl ester)
Greenstein, J.P. et al., Chemistry of the Amino Acids, (Chapter 31), Wiley, N.Y., 1961, 2075; 2750; 2758 (rev)
Heyns, K. et al., Annalen, 1963, 667, 194 (ms)
Quitt, P. et al., Helv. Chim. Acta, 1963, 46, 327 (N-Me)
Zervas, L. et al., J.A.C.S., 1963, 85, 3660-3666 (2-nitrobenzenesulfonyl)
Mizoguchi, T. et al., Chem. Pharm. Bull., 1970, 18, 1465-1474 (Me ester)
Benoiton, N.L. et al., Prog. Pept. Res., [Proc. Am. Pept. Symp.], 1970 (1972), 1972, 145 (N-Me)
Kricheldorf, H.R. et al., Synthesis, 1974, 422-424 (2-nitrobenzenesulfonyl)
Di Blasio, B. et al., Acta Cryst. B, 1975, 31, 601; 2941 (cryst struct)
Oguri, T. et al., Chem. Pharm. Bull., 1975, 23, 167 (synth)
Harding, M.M. et al., Acta Cryst. B, 1976, 32, 633 (cryst struct)
Kasai, T. et al., Agric. Biol. Chem., 1976, 40, 2489 (S-form, amide)
U.K. Pat., 1978, 1 562 373; C.A., 90, 110004v (synth, use, Leuciglumer)
Martinez, J. et al., Int. J. Pept. Protein Res., 1979, 13, 22-27 (2-naphthylamide)

- Banfi, S. *et al.*, *Tetrahedron*, 1984, **40**, 5207-5211 (*homopolymer, synth, ir*)
- Kovalev, V.N. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 724; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 687 (*isol*)
- Romani, S. *et al.*, *Synthesis*, 1985, 512-513 (2-nitrobenzenesulfenyl DCHA salt)
- Paventi, M. *et al.*, *Can. J. Chem.*, 1987, **65**, 282 (*nitrile*)
- Tamogami, S. *et al.*, *J. Chromatogr. A*, 1988, **822**, 310-315 (*N-Jasmonoylleucine*)
- Kramell, R. *et al.*, *Tetrahedron*, 1988, **44**, 5791-5807 (*N-Jasmonoylleucine, synth, pmr, ms*)
- Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, LER000; LES000
- Fr. Pat.*, 1989, 2 625 507; *CA*, **113**, 29326y (*synth, Leuciglumer*)
- Eur. Pat.*, 1990, ((SKB))403 252; *CA*, **115**, 71370z (*D-form, homopolymer, synth*)
- Itsuno, S. *et al.*, *J.O.C.*, 1990, **55**, 6047-6049 (*homopolymer, S-form, synth*)
- Flisak, J.R. *et al.*, *J.O.C.*, 1993, **58**, 6247-6254 (*homopolymer, S-form, ir, pmr*)
- Martindale, The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 597
- Fenaroli's Handbook of Flavor Ingredients*, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 429
- Allen, J.V. *et al.*, *Adv. Biochem. Eng./Biotechnol.*, 1996, **63**, 125-144 (*homopolymer, rev*)
- Lasterra-Sanchez, M.E. *et al.*, *J.C.S. Perkin 1*, 1996, 343-348 (*homopolymer, synth*)
- Encyclopedia of Food and Color Additives*, (ed. Burdock, G.A.), CRC Press, 1997, 1567-1569
- Wang, Y. *et al.*, *J.C.S. Perkin 2*, 1997, 899-904 (*amide, ir, cmr, cryst struct*)
- Allen, J.V. *et al.*, *J.C.S. Perkin 1*, 1998, 3171-3179 (*L-form, homopolymer*)
- Giard, T. *et al.*, *Synthesis*, 1998, 297-300 (*N-formyl Me ester*)
- Porter, M.J. *et al.*, *Bioorg. Med. Chem.*, 1999, **7**, 2145-2156 (*homopolymer, rev*)
- Miyazawa, T. *et al.*, *J.C.S. Perkin 1*, 2001, 82-86 (*anilide*)
- Onofrio, A.B. *et al.*, *J.C.S. Perkin 2*, 2001, 1863-1868 (*Me ester, synth*)
- Dutton, F.E. *et al.*, *J. Med. Chem.*, 2003, **46**, 2057-2073 (*benzyl ester*)
- Nadia, K. *et al.*, *J. Het. Chem.*, 2004, **41**, 57-60 (*N-benzoyl Me ester*)
- Isidro-Llobet, A. *et al.*, *Eur. J. Org. Chem.*, 2005, 3031-3039 (*4-nitrobenzyloxycarbonyl*)
- Rossi, J.-C. *et al.*, *Eur. J. Org. Chem.*, 2007, 662-668 (*R-form, amide*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LER000; LES000

Leucoagaricone**L-154**


[99280-75-8]

C₁₀H₁₃N₃O 191.232

Assumed to be a const. of the toadstool *Agaricus xanthoderma*. Colourless precursor of Agaricone, A-175. Cryst. (as hydrochloride). Mp 224° (hydrochloride).

Hilbig, S. *et al.*, *Angew. Chem., Int. Ed.*, 1985,
24, 1063-1065 (*isol*)**Leuconolam**

[93710-27-1]



Relative Configuration

C₁₉H₂₂N₂O₃ 326.394

Alkaloidal artifact from *Leuconotis griffithii* and *Leuconotis eugenifolia* (Apocynaceae). Cryst. (EtOH aq.). Mp 263-264°. [α]_D²⁸ -515.8 (c, 0.25 in MeOH). [α]_D -28.3 (c, 0.7 in CHCl₃).

O-Ac:

Cryst. (MeOH). Mp 235-236°.

N-Me: N-Methylleuconolam

[160985-67-1]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from roots of *Rhazya stricta* (Apocynaceae).

Me ether: 21-O-Methylleuconolam

[109305-79-5]

C₂₀H₂₄N₂O₃ 340.421


Alkaloid from leaves and stems of *Leuconotis eugenifolia* (Apocynaceae). Cryst. (MeOH). Mp 155-156° (140-150°). [α]_D²⁹ -597.8 (c, 0.28 in MeOH).

21-Epimer: EpileuconolamC₁₉H₂₂N₂O₃ 326.394

Alkaloid from *Leuconotis eugenifolia* (Apocynaceae). An artifact.

Goh, S.H. *et al.*, *Tet. Lett.*, 1984, **25**, 3483 (*uv, ir, pmr, cmr, ms, struct*)Wei, C. *et al.*, *Acta Cryst. C*, 1986, **42**, 349 (*cryst struct*)Goh, S.H. *et al.*, *Tet. Lett.*, 1986, **27**, 2501 (*Epileuconolam*)Hiller, W. *et al.*, *Acta Cryst. C*, 1989, **45**, 1258 (*cryst struct*)Goh, S.H. *et al.*, *Tetrahedron*, 1989, **45**, 7899(*uv, ir, pmr, cmr, ms, 21-O-Methylleuconolam*)Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 169 (*21-O-Methylleuconolam*)Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1995, **5**, 245 (*N-Methylleuconolam*)Banwell, M.G. *et al.*, *ARKIVOC*, 2006, **iii**, 163-174 (*synth*)**Leuconoxine****L-156**

[155416-24-3]



Relative Configuration

C₁₉H₂₂N₂O₂ 310.395

Highly rearranged indole alkaloid prob. most closely related to the Aspidosperma group. Alkaloid from leaves and

stems of *Leuconotis eugenifolia* (Apocynaceae), stem bark and leaves of *Kopsia griffithii* and stems of *Kopsia dasyrachis*. Prisms (CHCl₃/MeOH). Mp 238-242°. [α]_D²⁵ -88 (c, 1.2 in MeOH).

6-Oxo: 6-Oxoleuconoxine


[947251-84-5]

C₁₉H₂₀N₂O₃ 324.379

Alkaloid from the leaves of *Kopsia griffithii*. Oil. [α]_D +75 (c, 0.03 in CHCl₃). λ_{max} 202 (log ε 4.42); 234 (log ε 4.12); 251 (log ε 4.02); 349 (log ε 3.1) (EtOH).

Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 169-171 (*isol, pmr, cmr, struct*)Kam, T.-S. *et al.*, *Phytochemistry*, 1998, **47**, 145-146; 1999, **50**, 75-79; **51**, 159-169 (*occur*)Lim, S.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1380-1383 (*6-Oxoleuconoxine*)**Leucosolenamine A****L-157**

[926319-60-0]

C₁₇H₁₇N₇O₄ 383.366


Alkaloid from the calcareous sponge *Leucosolenia* sp. Brown solid.

4'-(N-Methylimine): Leucosolenamine B

[926319-67-7]

C₁₈H₂₀N₈O₃ 396.408

Alkaloid from a *Leucosolenia* sp. Brown solid.

Ralifo, P. *et al.*, *J. Nat. Prod.*, 2007, **70**, 33-38 (*isol, pmr, cmr, ms*)**Leucoxine****L-158****8-Hydroxy-9,10-dimethoxy-1,2-methylenedioxyaporphine****(S)-form**C₂₀H₂₁NO₅ 355.39**(S)-form** [1358-95-8]

Alkaloid from the leaves and stems of *Ocotea leucoxylon* and from the leaves of *Ocotea brachybotra* and *Ocotea minarum* (Lauraceae). Mp 213-217° dec. [α]_D²³ +81 (c, 0.518 in EtOH).

Me ether: 8,9,10-Trimethoxy-1,2-methylenedioxyaporphine. Ocopodine

[19893-95-9]
 $C_{21}H_{23}NO_5$ 369.416
 Alkaloid from *Ocotea macropoda* and from leaves of *Ocotea minarum* and *Ocotea brachybotra* (Lauraceae). Needles (EtOH). Mp 116-117°. $[\alpha]_D^{25} +87$ (c, 0.82 in EtOH).

6a,7-Didehydro, Me ether: **Dehydroocopodine**

[33117-76-9]
 $C_{21}H_{21}NO_5$ 367.401
 Alkaloid from *Ocotea macropoda* (Lauraceae). Golden-yellow plates (EtOH). Mp 113°.

(±)-form

Me ether: Synthetic. Syrup.

Me ether, hydrochloride:
 Prisms. Mp 230-234° dec.

Me ether, styphmate:
 Prisms. Mp 117-122°.

Goodwin, S. et al., *Chem. Ind. (London)*, 1960, 691 (isol)

Hirai, K. et al., *Yakugaku Zasshi*, 1960, **80**, 608; *CA*, **54**, 22696d (synth, *Ocopodine*)

Cava, M.P. et al., *Tet. Lett.*, 1968, 2437 (uv, pmr, struct, *Ocopodine*)

Cava, M.P. et al., *Tetrahedron*, 1971, **27**, 2639 (*Dehydroocopodine*)


Cava, M.P. et al., *J.O.C.*, 1975, **40**, 3601 (*Dehydroocopodine*)

Vecchietti, V. et al., *Farmaco, Ed. Sci.*, 1977, **32**, 767; 1979, **34**, 829; *CA*, **88**, **92**; 133240s (isol, uv, pmr, struct)

Ringdahl, B. et al., *J. Nat. Prod.*, 1981, **44**, 80 (cd)

Leucoxylophine L-159

6,7,7a,8-Tetrahydro-4,9,10,11-tetramethoxy-7-methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline, 9CI. 3,9,10,11-Tetramethoxy-1,2-methylene-dioxyaporphine



$C_{22}H_{25}NO_6$ 399.443

(S)-form [1358-96-9]

Alkaloid from the leaves and stems of *Ocotea leucoxylophine* and from the leaves of *Ocotea minarum* (Lauraceae). Also isol. from aerial parts of *Thalictrum simplex* (Ranunculaceae). Mp 54-58°. $[\alpha]_D^{24} +52$ (c, 0.925 in EtOH). Prob. a solvate.

Methodide: Mp 227-230° dec. $[\alpha]_D^{24} +33$ (c, 0.17 in EtOH).

N-Oxide: **Leucoxylophine N-oxide**

[166266-75-7]
 $C_{22}H_{25}NO_7$ 415.442

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae).

N-De-Me: **Norleucoxylophine**

[72176-43-3]
 $C_{21}H_{23}NO_6$ 385.416
 Alkaloid from the leaves of *Ocotea minarum* (Lauraceae). Mp 255-260° (as hydrochloride). $[\alpha]_D +41$ (c, 1.0 in MeOH).

O⁸-De-Me: **Ocoxylophine**

[66113-87-9]
 $C_{21}H_{23}NO_6$ 385.416
 Alkaloid from the leaves of *Ocotea leucoxylophine* (Lauraceae). Needles (MeOH). Mp 161-162°. $[\alpha]_D +45$ (CHCl₃).

Goodwin, S. et al., *Chem. Ind. (London)*, 1960, 691 (occur, uv)


Ahmad, R. et al., *Heterocycles*, 1977, **7**, 927 (*Leucoxylophine*, *Ocoxylophine*, uv, pmr, ms, struct)

Vecchietti, V. et al., *Farmaco, Ed. Sci.*, 1979, **34**, 829 (isol)

Valcheva, M.P. et al., *Phytochemistry*, 1995, **39**, 683 (*Leucoxylophine N-oxide*)

Leuhistin L-160

α -(1-Amino-3-methylbutyl)- α -hydroxy-1H-imidazole-4-propanoic acid, 9CI. 3-Amino-2-hydroxy-2-(1H-imidazol-4-ylmethyl)-5-methylhexanoic acid [129085-76-3]



Absolute Configuration

$C_{11}H_{19}N_3O_3$ 241.289


Prod. by *Bacillus laterosporus*. Aminopeptidase M inhibitor, immunostimulant. Needles (H₂O)(as hydrochloride). Sol. H₂O, EtOH, DMSO, MeOH; poorly sol. Me₂CO, hexane, EtOAc, CHCl₃. Mp 180-183° (hydrochloride). $[\alpha]_D^{25} -51.4$ (c, 1 in MeOH) (hydrochloride).

Aoyagi, T. et al., *J. Antibiot.*, 1991, **44**, 573; 579; 683 (isol, pmr, cmr, struct, biosynth)

Hecker, S.J. et al., *J.O.C.*, 1993, **58**, 1762 (synth)

Leupyrrin C L-161

[618911-34-5]



$C_{40}H_{56}N_2O_{10}$ 724.89

Partial stereochem. only determined. Prod. by *Sorangium cellulosum* strains So ce705 and So ce690. Amorph. solid.

21-Me ether: **Leupyrrin A₁**

[618911-30-1]
 $C_{41}H_{58}N_2O_{10}$ 738.917
 Prod. by *Sorangium cellulosum*. Active against fungi and eukaryotic cells. Amorph. solid. Mp 95°. $[\alpha]_D^{20} +12$ (c, 4.1 in MeOH). λ_{max} 260 (log ϵ 4.2); 286 (log ϵ 4.46) (MeOH).

2',3'-Didehydro: **Leupyrrin D**

[618911-35-6]
 $C_{40}H_{54}N_2O_{10}$ 722.874
 Prod. by *Sorangium cellulosum*. Amorph. solid.

2',3'-Didehydro, 21-Me ether: **Leupyrrin B₁**

[618911-32-3]
 $C_{41}H_{56}N_2O_{10}$ 736.901
 Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{20} +11$ (c, 4.6 in MeOH). λ_{max} 228 (log ϵ 4.33); 261 (log ϵ 4.26); 286 (log ϵ 4.45) (MeOH).

19,20-Didehydro, 21-Me ether: **Leupyrrin A₂**

[618911-31-2]
 $C_{41}H_{56}N_2O_{10}$ 736.901
 Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{20} -36$ (c, 5.4 in MeOH). λ_{max} 282 (log ϵ 4.43); 298 (log ϵ 4.57) (MeOH).

2',3',19,20-Tetradehydro, 21-Me ether:


Leupyrrin B₂
 [618911-33-4]
 $C_{41}H_{54}N_2O_{10}$ 734.885
 Prod. by *Sorangium cellulosum*. Amorph. solid. $[\alpha]_D^{20} +7$ (c, 4.6 in MeOH). λ_{max} 286 (log ϵ 4.32); 310 (log ϵ 4.16) (MeOH).

Bode, H.B. et al., *J. Nat. Prod.*, 2003, **66**, 1203-1206 (isol, uv, pmr, cmr)

Bode, H.B. et al., *Angew. Chem., Int. Ed.*, 2004, **43**, 4163-4167 (biosynth)

Leurosine L-162

4'-Deoxy-3',4'-epoxyvincalculoblastine, 9CI. **Vinleurosine**, INN. NSC 528004. Lilly 32645 [23360-92-1]



Absolute Configuration

$C_{46}H_{56}N_4O_9$ 808.97

Alkaloid from *Catharanthus roseus*, *Catharanthus lanceus*, *Catharanthus ovalis* and *Catharanthus longifolius* (Apocynaceae). Antineoplastic agent. Possesses antihyperglycaemic activity. Mp 202-205° dec. (anhyd.). Log P 2.85 (uncertain value) (calc). λ_{max} 214 (E1%/1cm 575); 259 (E1%/1cm 175) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 80 mg/kg. OH6370000

Sulfate salt: **Vinleurosine sulfate**, USAN

[1404-95-1]

Mp 238-242° dec. $[\alpha]_D$ -8.3 (MeOH).**N^{4'}-Oxide: Pleurosine.** Leurosine N^{6'}-oxide

[39608-80-5]

C₄₆H₅₆N₄O₁₀ 824.969Alkaloid from *Catharanthus roseus* (Apocynaceae). Noncryst. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{\max} 217 (ε 55000); 260 (ε 16700); 291 (ε 14200) (EtOH) (Berdy).**N-De-Me, N-formyl: 4'-Deoxy-3',4'-epoxy-22-oxovincaleukoblastine.** N-De-methyl-N-formylleurosine. Vinformide, INN. F-Leurosine. Neuroformine [54022-49-0]C₄₆H₅₄N₄O₁₀ 822.953Antineoplastic agent. Cryst. (MeOH). Mp 209-211° (212-216°). $[\alpha]_D^{20}$ +80.3 (c, 1 in CHCl₃). Log P 2.3 (uncertain value) (calc).

▶ Mutagenic props. LR2962000

N-De-Me, N-formyl, sulfate salt: Vinformide sulfate. Alzegir [58115-81-4]Mp 248-252°. $[\alpha]_D^{20}$ +37 (c, 1.0 in H₂O).**Deacetoxy: Deacetoxyleurosine**

[82267-12-7]


C₄₄H₅₄N₄O₇ 750.933Alkaloid from *Catharanthus roseus*. Cryst. (CHCl₃). Mp 202°.**21'-ζ-Hydroxy: 21'-Hydroxyleurosine**C₄₆H₅₆N₄O₁₀ 824.969Alkaloid from the aerial parts of *Catharanthus ovalis* (Apocynaceae). Mp 246°.**21'-Oxo: 21'-Oxoleurosine**C₄₆H₅₄N₄O₁₀ 822.953Alkaloid from *Catharanthus roseus* (Apocynaceae). Needles (MeOH). Mp 212-215°. $[\alpha]_D^{25}$ +55 (c, 0.10 in CHCl₃). λ_{\max} 220 (ε 61500); 270 (ε 18200); 290 (ε 16600) (EtOH) (Berdy).**16'-De(methoxycarbonyl): Vincovalinine**

[61927-01-3]

C₄₄H₅₄N₄O₇ 750.933Alkaloid from the aerial parts of *Catharanthus ovalis* (Apocynaceae). $[\alpha]_D$ +19 (c, 0.3 in CHCl₃).Neuss, N. et al., *J.A.C.S.*, 1959, **81**, 4754 (ir, uv, pmr, ms, struct, oxide)Neuss, N. et al., *Tet. Lett.*, 1968, 783 (ir, uv, pmr, ms, struct, oxide)Ger. Pat., 1974, ((*Vegyeszeti Gyár*))2 404 120; CA, **81**, 169686m (Vinformide)Taylor, W.I. et al., *Catharanthus Alkaloids*, Marcel, Dekker, 1975, (rev)Wenkert, E. et al., *Helv. Chim. Acta*, 1975, **58**, 1560 (cmr, struct)Gorog, S. et al., *J. Chromatogr.*, 1977, **139**, 203 (Vinformide, hplc)Kutney, J.P. et al., *Can. J. Chem.*, 1978, **56**, 62 (synth)Kutney, J.P. et al., *Heterocycles*, 1978, **9**, 201 (Vinformide, synth)Langlois, N. et al., *Helv. Chim. Acta*, 1980, **63**, 793 (Vincovalinine, 21'-Hydroxyleurosine)El-Sayed, A. et al., *J. Nat. Prod.*, 1980, **43**, 157 (21'-Oxoleurosine)Csuka, O. et al., *Oncology*, (Suppl. 1), 1980, 37, 83 (Vinformide, pharmacol)De Bruyn, A. et al., *Bull. Soc. Chim. Belg.*, 1982, **91**, 75-85 (Deacetoxyleurosine)Ronai-Lukacs, S. et al., *Eur. J. Drug Metab. Pharmacokinet.*, 1982, **7**, 47 (Vinformide, metab. pharmacol)Kutney, J.P. et al., *Helv. Chim. Acta*, 1982, **65**, 2088 (biosynth)**Leurosine**

[117590-95-1]

L-163

C₄₉H₆₀N₄O₁₀ 865.034Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Amorph. $[\alpha]_D^{24}$ +86 (c, 0.02 in CHCl₃).Atta-ur-Rahman, et al., *J.C.S. Perkin I*, 1988, 2175 (isol, uv, ir, pmr, cmr, ms, struct)**Leurosivine**

[1358-99-2]

L-164


C₄₁H₅₄N₃O₉ 732.892Bisindole alkaloid (*Aspidosperma*-cleavamine type). Struct. unknown. Alkaloid from the roots of *Catharanthus roseus* (Apocynaceae). Noncryst.; cryst. (EtOH)(as sulfate). Mp 335° dec. (sulfate).**Sulfate:**

Cryst. (EtOH). Mp 335° dec.

Svoboda, G.H. et al., *J. Nat. Prod.*, 1963, **26**, 141-153 (isol, uv, ir)Svoboda, G.H. et al., *J. Pharm. Sci.*, 1963, **52**, 407-408 (isol, uv, ir)**Liberine**

[56119-16-5]


L-165

7,9-Dihydro-2-methoxy-1,9-dimethyl-1H-purine-6,8-dione, 9CI. O²,1,9-Trimethyluric acidC₈H₁₀N₄O₃ 210.192Constit. of *Coffea liberica* and other *Coffea* spp. Also isol. from a Chinese tea, *Camellia assamica* var. *kucha*. Needles (EtOH aq.). Mp 269-273°. Melts and resolidifies due to thermalrearrangement to 8-Oxocaffeine in U-45. λ_{\max} 212 (ε 13520); 245 (ε 7790); 282 (ε 8155) (MeOH).**7-Me: 7,9-Dihydro-2-methoxy-1,7,9-trimethyl-1H-purine-6,8-dione, 9CI.****Methyliberine.** O²,1,7,9-Tetramethyluric acid

[51168-26-4]


C₉H₁₂N₄O₃ 224.219Isol. from *Coffea* spp. and *Camellia assamica* var. *kucha*. Cryst. (MeOH). Mp 207.3-208.4°. λ_{\max} 213 (ε 17080); 247 (ε 8430); 284 (ε 8145) (MeOH).Wanner, H. et al., *Phytochemistry*, 1975, **14**, 747-750 (isol, synth, uv, ms)Zheng, X.-Q. et al., *Phytochemistry*, 2002, **60**, 129-134 (isol, bibl)**Lidimycin, INN**

L-166

5-(Hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl)-2-pentenoic acid, 9CI. **8CI.** **Lydimycin, USAN.** α-Dehydrobiotin. U 15965. Antibiotic U 15965 [10118-85-1]C₁₀H₁₄N₂O₃S 242.298Isol. from culture filtrates of *Streptomyces lydicus* NRRL2433. Biotin anti-metab. Antifungal agent. Needles (MeOH). Sol. MeOH, Et₂O; fairly sol. H₂O. Mp 238-240° Mp 256-257.5°. $[\alpha]_D^{25}$ +105.7 (c, 1.2 in 0.1M NaOH). pK_a 4.32. Log P -0.12 (calc). λ_{\max} 203 (ε 15000) (EtOH) (Berdy).Hanka, L.J. et al., *Science (Washington, D.C.)*, 1966, **154**, 1667 (isol)Field, G.F. et al., *J.A.C.S.*, 1970, **92**, 3520 (synth)Martin, D.G. et al., *Tet. Lett.*, 1971, 3791 (synth)Field, G.F. et al., *J.O.C.*, 1976, **41**, 3853 (synth, ir, nmr)Tanaka, M. et al., *J. Biotechnol.*, 1987, **5**, 209 (synth)Pearson, B.M. et al., *Lett. Appl. Microbiol.*, 1990, **10**, 89 (synth)**Lienkonine**

[83151-88-6]

L-167



Absolute Configuration

C₂₁H₂₅N₄O₄ 355.433Alkaloid from *Corydalis ochotensis* (Papaveraceae). Cryst. (EtOH). Mp 166-167°. $[\alpha]_D^{29}$ -180 (c, 1.0 in EtOH).Lu, S.-T. et al., *Phytochemistry*, 1982, **21**, 809 (isol, uv, ir, pmr, ms, struct)

Ligundentine


L-168

Pyrrrolizidine alkaloid. Struct. unknown. Alkaloid from *Ligularia dentata* and *Ligularia brachyphylla* (Asteraceae). Cooccurs with Ligularine and Clivorine. Klásek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1971, **36**, 2205-2215 (*isol, pmr*)

Ligularine

L-169

[34429-54-4]



Absolute Configuration


C₂₃H₃₁NO₉ 465.499

Cyclic otonecine diester. Alkaloid from *Ligularia elegans*, *Ligularia hodgsonii* and *Ligularia dentata* (Asteraceae). Noncryst. [α]_D²⁴ -34 (c, 0.82 in CHCl₃). Klásek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1971, **36**, 2205-2215 (*isol, ir, pmr*)
Lin, G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 857-860 (*isol, pmr, cmr, struct*)

Ligularizine

L-170

[90364-92-4]

C₂₁H₂₉NO₈ 423.462

Cyclic diester of Otonecine, O-136. Alkaloid from the aerial parts and roots of *Ligularia dentata* (Asteraceae). Yellow needles (EtOH)(as picrate). Mp 210-211° (picrate). [α]_D²⁵ -24.5 (c, 1.08 in CHCl₃) (picrate).


▶ VS3595400

Asada, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 475 (*isol, ir, pmr, cmr, ms, struct*)

Lihouidine

L-171

[714965-76-1]


C₃₆H₂₇N₅O₆ 625.639

Alkaloid from the marine sponge *Suberea* sp. nov. Purple-red prisms (MeCN). Mp > 330°. Racemic. λ_{\max} 210 (ε 54400); 246 (ε 74200); 287 (sh) (ε 24000); 314 (ε 19000); 340 (ε 14700); 406 (ε 5100); 430 (ε 5600); 444 (ε 5600); 514 (ε 10500); 543 (sh) (ε 10200) (EtOH).

Bowden, B.F. *et al.*, *J.O.C.*, 2004, **69**, 7791-7793 (*isol, uv, pmr, cmr, cryst struct*)

Lilacinone

L-172

C₂₂H₁₅N₃O₁₁ 497.374


Isol. from *Lactarius lilacinus*. Red solid. Mp > 300° dec. Racemic. λ_{\max} 210 (log ε 4.06); 250 (log ε 4.11); 309 (sh) (log ε 3.7); 354 (log ε 3.82); 519 (log ε 3.24) (MeOH).

Spiteller, P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1402-1403 (*isol, uv, pmr, cmr, ms*)

Lilaline

L-173

3-Methyl-5-[3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4-oxo-4H-1-benzopyran-8-yl]-2-pyrrolidinone, 9CI. 3,4',5,7-Tetrahydroxy-8-(4-methyl-5-oxo-2-pyrrolidinyl)-flavone [110011-49-9]

C₂₀H₁₇NO₇ 383.357


Alkaloid from the aerial parts of *Lilium candidum* (Liliaceae). Yellow prisms (Me₂CO). Mp 247°. [α]_D²⁷ +65 (c, 0.2 in MeOH).

Mašterová, I. *et al.*, *Phytochemistry*, 1987, **26**, 1844 (*isol, uv, ir, pmr, cmr, ms, struct*)

Limaciamine

L-174

N,N'-Carbonimidoylbis[2-ethylhexanamide], 9CI. N,N'-Bis(2-ethylhexanoyl)-guanidine [202747-40-8]

C₁₇H₃₃N₃O₂ 311.467


Isol. from the skin extracts of nudibranch *Limacia clavigera*. Glass. Related to Triophamine, T-617.

Graziani, E.I. *et al.*, *J. Nat. Prod.*, 1998, **61**, 285-286 (*isol, pmr, cmr, ms*)

Limacusine

L-175

6,6',12'-Trimethoxy-2,2'-dimethoxyxanthan-7-ol, 9CI [10172-03-9]

C₃₇H₄₀N₂O₆ 608.733

Alkaloids covered by this entry (*R,R*-config.) are enantiomeric with those covered by Repandine, R-46 and diastereomeric with those under Oxycanthine with those under Oxycanthine from *Limacia oblonga* and *Limacia cuspidata* (Menispermaceae). Antineoplastic agent. Cryst. (MeOH/Me₂CO). Mp 235-237°. [α]_D +110 (CHCl₃). Log P 8.08 (uncertain value) (calc). λ_{\max} 283 (no solvent reported).

N²-β-Oxide: **Limacusine 2'-β-N-oxide** [172045-41-9]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from leaves of *Anisocycla jollyana* (Menispermaceae). Mp 215°. [α]_D²⁰ +157 (c, 1.1 in CHCl₃). λ_{\max} 218 (log ε 4.72); 283 (log ε 3.9) (MeOH).

N²-De-Me: **2-Norlimacusine**

[96744-71-7]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Sciadotenia eichleriana* (Menispermaceae). Cryst. (MeOH). Mp 172°. [α]_D +167 (c, 0.7 in CHCl₃).

Me ether: **O-Methylimacusine**

[13017-15-7]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from the stem bark of *Gyrocarpus americanus* (Hernandiaceae). Cryst. (MeOH). Mp 210-212°.

O¹²-De-Me: **Candicusine. 12-O-De-methylimacusine**

[99945-41-2]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the roots of *Curarea candicans* (Menispermaceae). [α]_D +75 (c, 0.07 in MeOH).

O¹²-De-Me, N²,N²-di-de-Me: **Pangkoramine**

[110416-09-6]

C₃₄H₃₄N₂O₆ 566.652

Alkaloid from *Albertisia* cf. *Albertisia papuana* (Menispermaceae). Amorph. [α]_D +126 (c, 0.05 in MeOH). λ_{\max} 214 (log ε 4.72); 230 (log ε 4.39); 280 (log ε 4.58) (EtOH).

O^{6'}-De-Me, O^{7'}-Me: **Gyrocarpusine**

[102518-66-1]


Alkaloid from the stem bark of *Gyrocarpus americanus* (Hernandiaceae). Amorph. [α]_D +66 (c, 1 in CHCl₃). λ_{\max} 234 (log ε 4.42); 283 (log ε 3.85) (MeOH).

- Tomita, M. *et al.*, *Tet. Lett.*, 1966, 4293-4296 (*Limacusine*, *pmr*, *ms*, *struct*)
 Tomita, M. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 793-796; 1560-1561; *CA*, **68**, 3048m; 84980j (*isol*, *uv*, *pmr*, *ms*, *struct*, *Limacusine*, *O-Methylimacusine*)
 Lavault, M. *et al.*, *J. Chem. Res., Synop.*, 1985, 248-249; *J. Chem. Res., Miniprint*, 1985, 2786-2793 (*Candicusine*)
 Damas, P. *et al.*, *J. Nat. Prod.*, 1985, **48**, 69-71 (*Limacusine*, *Norlimacusine*)
 Chalandre, M.-C. *et al.*, *J. Nat. Prod.*, 1986, **49**, 101-105 (*O-Methylimacusine*, *Gyrocarpusine*)
 Lavault, M. *et al.*, *Can. J. Chem.*, 1987, **65**, 343-347 (*Pangkoramine*)
 Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1587-1589 (*Limacusine 2'-β-N-oxide*)

Limalongine

L-176

[121255-00-3]

C₁₈H₂₃NO₅ 333.383

Modified hasubanan-type alkaloid. Alkaloid from the stem bark of *Limacia oblonga* (Menispermaceae). [α]_D²⁰ +290 (c, 0.15 in CHCl₃).

10ξ-Chloro: Clolimalongine

[121254-99-7]

C₁₈H₂₂ClNO₅ 367.828


Alkaloid from the stem bark of *Limacia oblonga* (Menispermaceae). [α]_D²⁰ +300 (c, 0.1 in CHCl₃).

Berthou, S. *et al.*, *J.O.C.*, 1989, **54**, 3491 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Limaspermidine

L-177

[72821-58-0]



Absolute Configuration

C₁₉H₂₆N₂O 298.427

Alkaloid from *Aspidosperma rhombeo-signatum* (Apocynaceae). Mp 227-228°.

12-Hydroxy, N-Ac: Limapodine

[2636-98-8]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from the bark extracts of *Aspidosperma limae* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 177°. [α]_D²³ +110 (c, 0.84 in CHCl₃).

12-Hydroxy, N-propanoyl: Limaspermine.*Dihydrohaplocine (incorr.)*

[5516-64-3]

C₂₂H₃₀N₂O₃ 370.491

Alkaloid from *Aspidosperma limae* (Apocynaceae). Cryst. (MeOH or Me₂CO/hexane). Mp 175-175.5°. [α]_D²¹ +108 (c, 0.78 in CHCl₃).

12-Methoxy: Cyllindrocarpinol

[28189-98-2]

Mp 145-147°. [α]_D²⁰ +2 (MeOH).**12-Methoxy, N-formyl: N-Formylcyllindrocarpinol**C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Aspidosperma cyllindrocarpon* (Apocynaceae). Mp 101-105°. [α]_D²⁰ -150 (c, 0.0016 in MeOH).

12-Methoxy, N-Ac: N-AcetylcyllindrocarpinolC₂₂H₃₀N₂O₃ 370.491

Alkaloid from *Aspidosperma cyllindrocarpon* and *Aspidosperma dispernum* (Apocynaceae). Mp 210-215°.

12-Methoxy, N-propanoyl:C₂₃H₃₂N₂O₃ 384.517Mp 170°. [α]_D²⁹ +89 (c, 0.823 in CHCl₃).**11-Methoxy, 12-hydroxy, N-Ac: Aspido-limidinol. 11-Methoxyylimapodine**C₂₂H₃₀N₂O₄ 386.49

Alkaloid from *Aspidosperma album* and *Aspidosperma limae* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 219-220.5°. [α]_D²⁰ +131 (c, 0.83 in CHCl₃). Props. given for free base refer to Methoxyylimapodine from *A. limae*.

11-Methoxy, 12-hydroxy, N-Ac, hydrobromide:

Cryst. (MeOH aq.). Mp 197-199°. [α]_D²⁰ +70 (c, 0.66 in H₂O). Props. for hydrobromide refer to Aspidolimidinol from *A. album*.

11-Methoxy, 12-hydroxy, N-propanoyl:**11-Methoxyylimaspermine**C₂₃H₃₂N₂O₄ 400.517

Alkaloid from *Aspidosperma limae* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 174-175°. [α]_D²³ +118 (c, 0.79 in CHCl₃).

Pinar, M. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 2260 (*isol*, *uv*, *ms*, *pmr*, *struct*)
 Djerassi, C. *et al.*, *Tet. Lett.*, 1962, 1001-1009 (*Cyllindrocarpinol*, *synth*)

Pinar, M. *et al.*, *Annalen*, 1963, **668**, 97 (*Limapodine*, *11-Methoxyylimaspermine*)

Ferrari, C. *et al.*, *Can. J. Chem.*, 1964, **42**, 2705-2709 (*Aspidolimidinol*)

Cava, M.P. *et al.*, *Tetrahedron*, 1964, **20**, 581 (*stereochem*)

Klyne, W. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 833 (*ord*, *abs config*, *synth*)

Milborrow, B.V. *et al.*, *J.C.S. (C)*, 1969, 417 (*N-Formylcyllindrocarpinol*, *N-Acetylcyllindrocarpinol*)

Honma, Y. *et al.*, *Heterocycles*, 1976, **5**, 47 (*Limaspermidine*)

Lawson, G. *et al.*, *Tetrahedron*, 1977, **33**, 1641 (*synth*)

Wegmann, A. *et al.*, *Anal. Chem.*, 1978, **50**, 830 (*ms*)

Medina, J.D. *et al.*, *Planta Med.*, 1979, **37**, 165 (*Limaspermidine*)

Pearson, A.J. *et al.*, *J.C.S. Perkin 1*, 1982, 2467 (*Limaspermidine*, *synth*)


Fukuda, Y. *et al.*, *Heterocycles*, 2004, **62**, 787-792 (*Limaspermine*, *synth*)

Limocrocin

L-178

N,N'-Bis(2-hydroxy-5-oxo-1-cyclopenten-1-yl)-2,4,6,8,10,12,14-hexadecaheptaenediamide, 9CI

[24112-57-0]

C₂₆H₂₆N₂O₆ 462.501

Prod. by *Streptomyces limosus*. Antiviral agent. Active against gram-positive bacteria. Dark-red cryst. (AcOH). Sol. MeOH, CHCl₃. Mp 316° dec. λ_{max} 258 (ε 45000); 422 (ε 100000); 445 (ε 88000) (H₂O) (Derep). λ_{max} 258 (ε 45000); 422 (ε 100000); 445 (ε 88000) (H₂O) (Derep). λ_{max} 258 ; 420 (Na₂CO₃) (Berdy).

Brockmann, H. *et al.*, *Chem. Ber.*, 1953, **86**, 1110; 1955, **88**, 419 (*isol*, *struct*)

Macdonald, G. *et al.*, *Tetrahedron*, 1998, **54**, 9823-9836 (*synth*)

Linanthenine

L-179

Limnanthenine

Struct. unknown. Alkaloid from *Limnanthemum humboldtianum* (preferred genus name *Nymphoides*). Mp 285° dec. (darkens at 260°). [α]_D²⁰ 0. Genus *Limnanthemum* assigned by the authors to the Gentianaceae, but normally assigned to the Menyanthaceae.

Hydrochloride:

Needles (EtOH). Mp 194-195°.


Sulfate: Mp 287°.**Picrate:**Yellow prisms (Me₂CO aq.). Mp 182°.

Ribeiro, O. *et al.*, *CA*, 1952, **46**, 3219

Linarinic acid

L-180

1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazoline-1-carboxylic acid

C₁₂H₁₂N₂O₂ 216.239**(S)-form** [481710-98-9]

Alkaloid from *Linararia vulgaris*. Needles (MeOH). Mp 218° dec. [α]_D¹⁸ -217 (0.01 in MeOH) (natural). [α]_D¹⁸ -290 (c, 0.01 in MeOH) (synthetic). λ_{max} 211 ; 281 (MeOH).


Hua, H. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1393-1394 (*isol*, *pmr*, *cmr*)

Cheng, M. *et al.*, *Tetrahedron: Asymmetry*, 2006, **17**, 179-183 (*synth*, *abs config*)

Linavuline

L-181

[215664-11-2]


C₁₀H₁₃NO₃ 195.218

Alkaloid from *Linararia vulgaris*.

Hua, H. *et al.*, *CA*, 1999, **130**, 2166k

Lindenialine**L-182**

2,4a,5,6,7,7a-Hexahydro-7-methyl-1H-2-pyridine-4-carboxaldehyde, 9CI
[114542-52-8]




C₁₀H₁₅NO 165.235

Alkaloid from the leaves of *Lindenia austro-caledonica* (Rubiaceae). Fine needles (MeOH). Mp 162°. [α]_D²⁰ +20 (c, 1.6 in EtOH). Artifact.

Saad, H.E.A. *et al.*, *Tet. Lett.*, 1988, **29**, 615 (isol, uv, ir, pmr, cmr, ms, struct)

Lindeniamine**L-183**

[114558-81-5]



Relative Configuration


C₂₀H₂₆N₂O₂ 326.438

Alkaloid artifact from the leaves of *Lindenia austro-caledonica* (Rubiaceae). Cryst. (MeOH). Mp 264° dec. [α]_D²⁰ -26 (c, 0.4 in MeOH).

Saad, H.E.A. *et al.*, *Tet. Lett.*, 1988, **29**, 615 (isol, uv, ir, pmr, cmr, ms, struct)

Lindheimerine**L-184**

[68831-67-4]



C₂₂H₃₁NO₂ 341.492


Minor alkaloid from the bark and leaves of *Garrya ovata* var. *lindheimeri* (Garryaceae). Amorph. [α]_D²⁴ -113.8 (c, 1.0 in CHCl₃).

Pelletier, S.W. *et al.*, *Heterocycles*, 1978, **9**, 1409-1412 (ir, pmr, cmr, struct, synth)

Pelletier, S.W. *et al.*, *J.O.C.*, 1981, **46**, 1840-1846 (isol, cmr, pmr)

Lipogrammistin A**L-185**

[151756-67-1]



Absolute configuration

C₃₅H₆₆N₄O₃ 590.931

Isol. from the mucus secreted by the soapfish *Diploprion bifasciatum*. Shows haemolytic activity and exhibits erythrocyte transformation. Ichthyotoxic. [α]_D²⁰ +18.5 (c, 0.86 in MeOH). Exists in soln. as a mixt. of 4 conformers.


Onuki, H. *et al.*, *J.O.C.*, 1998, **63**, 3925-3932 (synth, abs config)

Kobayashi, Y. *et al.*, *Bioorg. Med. Chem.*, 1999, **7**, 2073-2081 (activity)

Kan, T. *et al.*, *Tetrahedron*, 2002, **58**, 6267-6276 (synth)

α-Lipomycin**L-186**

[51053-40-8]



C₃₂H₄₅NO₉ 587.709

Pentaene antibiotic. Prod. by *Streptomyces aureofaciens*. Active against gram-positive organisms. Orange-red amorph. powder. Sol. MeOH, Et₂O, bases; poorly sol. H₂O, hexane. Mp 105°. [α]_D²⁰ -229 (c, 0.1 in MeOH). λ_{max} 260 (ε 16400); 267 (ε 15800); 301 (ε 10600); 455 (ε 55800) (MeOH/HCl) (Derep). λ_{max} 260 (ε 15800); 270 (ε 15300); 287 (ε 10600); 403 (ε 47000) (MeOH) (Derep). λ_{max} 259 (ε 14700); 269 (ε 13500); 285 (ε 10600); 393 (ε 55900) (MeOH/NaOH) (Derep).

Aglycone: β-Lipomycin

[51053-41-9]

C₂₆H₃₅NO₆ 457.566

Prod. by *Streptomyces aureofaciens*.

Active against gram-positive bacteria.

Sol. bases, Et₂O, MeOH; poorly sol.

H₂O, hexane. [α]_D²⁰ -176 (c, 0.1 in

MeOH). λ_{max} 261 (ε 15800); 269 (ε

15100); 301 (ε 9830); 453 (ε 57600)

(MeOH/HCl) (Derep). λ_{max} 260 (ε

14400); 269 (ε 13700); 287 (ε 10100);

393 (ε 57600) (MeOH/NaOH) (Derep).

λ_{max} 260 (ε 14600); 269 (ε 14600); 287

(ε 10100); 399 (ε 9830); 460 (ε 47500)


(MeOH) (Derep).

Kunze, B. *et al.*, *Arch. Microbiol.*, 1972, **86**, 147 (isol)

Schabacher, K. *et al.*, *Tet. Lett.*, 1973, 2691 (struct)

Zeeck, A. *et al.*, *Annalen*, 1975, 2079

Nolte, M.J. *et al.*, *J.C.S. Perkin 1*, 1980, 1057 (cryst struct)

Lipopurealins**L-187**

Lipopurealin A; R = (CH₂)₂CH₃
B; R = (CH₂)₃CH(CH₃)₂
C; R = (CH₂)₄CH₃
D; R = (CH₂)₅CH=CH(CH₂)₄CH(CH₃)₂ (Z-)
E; R = (CH₂)₆CH₃

Metabs. of the Okinawan marine sponge

Psammaphysilla pura.

Lipopurealin A [101541-48-4]

C₃₁H₄₈Br₂N₆O₄ 728.566

Na⁺ and K⁺-ATPase inhibitor.

Amorph. solid (as hydrochloride). Sol.

MeOH, CHCl₃; poorly sol. H₂O. Mp 94-

95° (hydrochloride). CAS no. refers to

hydrochloride. λ_{max} 284 (ε 970) (MeOH) (Berdy).

Lipopurealin B [101541-49-5]

C₃₂H₅₀Br₂N₆O₄ 742.593

Na⁺ and K⁺-ATPase inhibitor.

Amorph. solid (as hydrochloride). Sol.

MeOH, CHCl₃; poorly sol. H₂O. Mp 93-

95° (hydrochloride). CAS no. refers to

hydrochloride. λ_{max} 284 (ε 930) (MeOH) (Berdy).

Lipopurealin C [101541-50-8]

C₃₃H₅₂Br₂N₆O₄ 756.62

Na⁺ and K⁺-ATPase inhibitor.

Amorph. solid (as hydrochloride). Sol.

MeOH, CHCl₃; poorly sol. H₂O. Mp

108-110° (hydrochloride). CAS no. refers

to hydrochloride. λ_{max} 284 (ε 910)

(MeOH) (Berdy).

Lipopurealin D [164301-28-4]

C₃₄H₅₂Br₂N₆O₄ 768.631

Amorph. solid. Oxime has *E*-config.

Lipopurealin E [164301-29-5]

C₃₆H₅₈Br₂N₆O₄ 798.7

Amorph. solid. Oxime has *E*-config.


Wu, H. *et al.*, *Experientia*, 1986, **42**, 855

(Lipopurealins A-C)

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1995, **58**, 467 (Lipopurealins D-E)

Liposidomycin**L-188**

[118817-12-2]



Liposidomycin A-I R = -(CH₂)₇CH=CHCH₂CH=CHCH₂¹³CH₂CH₂CH₂CH₃

B-I R = -(CH₂)₈CH(CH₃)₂

H-I R = -(CH₂)₁₀CH₃

K-I R = -(CH₂)₈CH=CHCH₂¹²CH=CH(CH₂)₄CH₃

L-I R = -(CH₂)₈CH(CH₃)₂

X-I R = -(CH₂)₆CH₃

Y-I R = -CH₂⁷CH=CHCH₂⁸CH=CH(CH₂)₃CH₃

Nucleoside antibiotic complex. Compounds

mostly not abstracted by CAS. Isol. from

Streptomyces griseosporus, *Streptomyces*

sp. RK-1061 and *Streptomyces* sp. SN-1061M.

Inhibits bacterial peptidoglycan synthetase.

Sol. H₂O, DMSO; fairly sol. MeOH, EtOH, AcOH; poorly

sol. Me₂CO, hexane. Mp 190° dec. [α]_D²⁴

+18.7 (c, 0.3 in H₂O). Liposidomycins

have been divided into 4 structural types.

Not all of the members of each homologous

series have been detected. λ_{max} 260

(ε 5150) (0.1N HCl) (Derep). λ_{max} 261

(ε 7160) (0.1N NaOH) (Derep). λ_{max} 262

(ε 7260) (H₂O) (Derep). λ_{max} 260

(E1%/1cm 60) (H₂O) (Berdy). λ_{max} 258

(E1%/1cm 44) (HCl) (Berdy). λ_{max} 259

(E1%/1cm 58) (NaOH) (Berdy).

Liposidomycin A-I [99751-55-0]

C₄₄H₆₇N₅O₂₁S 1034.1
Powder.

Desulfo: Liposidomycin A-III
[213201-84-4]

C₄₄H₆₇N₅O₁₈ 954.036
From *Streptomyces* sp. SN-1061M.
Powder.

3-O-Deacetyl: Liposidomycin A-II

C₃₈H₅₉N₅O₁₈S 905.973
From *Streptomyces* sp. SN-1061M.
Powder.

3-O-Deacetyl, desulfo: Liposidomycin A-IV

C₃₈H₅₉N₅O₁₅ 825.909
From *Streptomyces* sp. SN-1061M.
Powder.

10,11-Dihydro: Liposidomycin G-I. Antibiotic RK 1061G. RK 1061G

C₄₄H₆₉N₅O₂₁S 1036.116
Prod. by *Streptomyces* sp. RK-1061.

10,11-Dihydro, desulfo: Liposidomycin G-III

C₄₄H₆₉N₅O₁₈ 956.052
Prod. by *Streptomyces* sp. SN-1061M.

13,14-Didehydro(Z-): Liposidomycin V-I

C₄₄H₆₅N₅O₂₁S 1032.084
Not detected.

13,14-Didehydro, desulfo: Liposidomycin V-III

C₄₄H₆₅N₅O₁₈ 952.02
From *Streptomyces* sp. SN-1061M.

7,8,10,11-Tetrahydro: Liposidomycin M-I

C₄₄H₇₁N₅O₂₁S 1038.132
Prod. by *Streptomyces* sp. SN-1061M.

7,8,10,11-Tetrahydro, desulfo: Liposidomycin M-III

C₄₄H₇₁N₅O₁₈ 958.068
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin B-I [99751-53-8]

C₄₂H₆₇N₅O₂₁S 1010.078
Inhibitor of bacterial peptidoglycan
synthetase. Powder. Mp 190° dec. [α]_D²⁴
+17.3 (c, 0.4 in H₂O).

Desulfo: Liposidomycin B-III

C₄₂H₆₇N₅O₁₈ 930.014
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin H-I

Antibiotic RK 1061H. RK 1061H

C₄₃H₆₉N₅O₂₁S 1024.105

Desulfo: Liposidomycin H-III

C₄₃H₆₉N₅O₁₈ 944.041
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin K-I

C₄₆H₇₁N₅O₂₁S 1062.154

Desulfo: Liposidomycin K-III

C₄₆H₇₁N₅O₁₈ 982.09
Prod. by *Streptomyces* sp. SN-1061M.

12,13-Dihydro: Liposidomycin N-I

C₄₆H₇₃N₅O₂₁S 1064.17
Prod. by *Streptomyces* sp. SN-1061M.

12,13-Dihydro, desulfo: Liposidomycin N-III

C₄₆H₇₃N₅O₁₈ 984.105
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin L-I

C₄₄H₇₁N₅O₂₁S 1038.132

Desulfo: Liposidomycin L-III

C₄₄H₇₁N₅O₁₈ 958.068
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin X-I

C₄₁H₆₅N₅O₂₁S 996.051
Not detected.

Desulfo: Liposidomycin X-II

C₄₁H₆₅N₅O₁₈ 915.987
Prod. by *Streptomyces* sp. SN-1061M.

Liposidomycin Y-I

C₄₂H₆₃N₅O₂₁S 1006.047
Not detected.

Desulfo: Liposidomycin Y-III

C₄₂H₆₃N₅O₁₈ 925.982
Prod. by *Streptomyces* sp. SN-1061M.

8,9-Dihydro: Liposidomycin Z-I

C₄₂H₆₅N₅O₂₁S 1008.062
Prod. by *Streptomyces* sp. SN-1061M.

8,9-Dihydro, desulfo: Liposidomycin Z-III

C₄₂H₆₅N₅O₁₈ 927.998
Prod. by *Streptomyces* sp. SN-1061M.

5,6,8,9-Tetrahydro: Liposidomycin C-I

[99751-54-9]
C₄₂H₆₇N₅O₂₁S 1010.078
Prod. by *Streptomyces* sp. RK-1061.
Powder. Mp 190° dec. [α]_D²⁴ +18.9 (c,
0.9 in H₂O).

5,6,8,9-Tetrahydro, desulfo: Liposidomycin C-III

C₄₂H₆₇N₅O₁₈ 930.014
Prod. by *Streptomyces* sp. SN-1061M.

5,6,8,9-Tetrahydro, 3-deacetyl: Liposidomycin C-II

C₃₆H₅₉N₅O₁₈S 881.951
Prod. by *Streptomyces* sp. SN-1061M.

5,6,8,9-Tetrahydro, 3-deacetyl, desulfo: Liposidomycin C-IV

C₃₆H₅₉N₅O₁₅ 801.887
Prod. by *Streptomyces* sp. SN-1061M.

Isono, K. *et al.*, *J. Antibiot.*, 1985, **38**, 1617-1621 (*isol, uv, ir, pmr*)

Kimura, K. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 1811-1815 (*isol, biochem*)

Ubukata, M. *et al.*, *J.O.C.*, 1992, **57**, 6392-6403 (*pmr, ms, uv, struct*)

Kimura, K. *et al.*, *J. Antibiot.*, 1998, **51**, 640-646; 647-654 (*isol, struct, props*)

Esumi, Y. *et al.*, *J. Antibiot.*, 1999, **52**, 281-287

Knapp, S. *et al.*, *J.O.C.*, 2001, **66**, 5822-5831 (*synth*)


Lipoxamycin**L-189**

2-Amino-N,3-dihydroxy-N-(14-methyl-3,10-dioxopentadecyl)propanamide, 9CI.

2-Amino-1,4-dihydroxy-18-methyl-4-aza-3,7,14-nonadecanetrione. N-Hydroxy-N-

(15-methyl-3,10-dioxopentadecyl)seramide. Enactin IVa. Neoenactin M₁

[32886-15-0]



C₁₉H₃₆N₂O₅ 372.504

Isol. from *Streptomyces virginiae*, *Streptomyces olivoreticuli* ssp. *neoenactus* and *Streptomyces roseoviridis*. Shows antifungal activity. Mp 68-70°. Related to Neoenactins. λ_{max} 240 (ε 5500) (MeOH/NaOH) (Derep). λ_{max} 210 (ε 5900) (MeOH) (Derep).

▶ MU5230000

Sulfate: Mp 155°.

10'-Alcohol: *Enactin Va*. Hydroxylipoxamycin [130640-31-2]

C₁₉H₃₈N₂O₅ 374.52
Isol. from *Streptomyces roseoviridis* and another *Streptomyces* sp. Mp 125.5-128.5° (as sulfate). λ_{max} 208 (ε 3334) (MeOH).


Whaley, M.A. *et al.*, *J.A.C.S.*, 1971, **93**, 3767 (*isol, struct*)

Roy, S.K. *et al.*, *J. Antibiot.*, 1987, **40**, 266 (*isol*)
Yamamoto, K. *et al.*, *J. Antibiot.*, 1990, **43**, 1012 (*Enactins*)

Mandala, S.M. *et al.*, *J. Antibiot.*, 1994, **47**, 376-379 (*Hydroxylipoxamycin*)

Lipoxazolidinone A**L-190**

[908138-32-9]



R¹ = -(CH₂)₆CH₃, R² = -(CH₂)₃CH₃

C₁₉H₃₁NO₃ 321.459

Prod. by a *Marinispora* sp. (strain NPS008920). Antibacterial agent. Oil. [α]_D -31 (c, 0.02 in MeOH). λ_{max} 253 (log ε 3.96); 310 (log ε 4.36) (MeOH).

Macherla, V.R. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1454-1457 (*isol, pmr, cmr*)

Lipoxazolidinone B**L-191**

[908138-34-1]

As Lipoxazolidinone A, L-190 with R¹ = -(CH₂)₆CH₃, R² = -(CH₂)₄CH₃

C₂₀H₃₃NO₃ 335.486
Prod. by a *Marinispora* sp. (strain NPS008920). Antibacterial agent. Oil. λ_{max} 255 ; 310 (MeCN aq.).

Macherla, V.R. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1454-1457 (*isol, pmr, cmr*)

Lipoxazolidinone C**L-192**

[908138-36-3]


As Lipoxazolidinone A, L-190 with R¹ = -(CH₂)₅CH₃, R² = -(CH₂)₃CH₃

C₁₈H₂₉NO₃ 307.432
Prod. by a *Marinispora* sp. (strain NPS008920). Antibacterial agent. Oil. λ_{max} 255 ; 310 (MeCN aq.).

Macherla, V.R. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1454-1457 (*isol, pmr, cmr*)

Liriodendronine**L-193**

1,2-Dihydroxy-7H-dibenzo[de,g]quinolin-7-one, 9CI
[65400-36-4]



C₁₆H₉NO₃ 263.252

Alkaloid from the discoloured sapwood of *Liriodendron tulipifera* (Magnoliaceae). Shows anti-yeast activity. Violet cryst. (Et₂O). Mp 265-270° dec.

N-Me: N-Methyliriodendronine

[132872-61-8]

C₁₇H₁₁NO₃ 277.279

Alkaloid from *Stephania dinklagei*. Zwitterionic.

O¹-Me: 2-Hydroxy-1-methoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI. 7-Oxodehydroasimilobine

[138690-42-3]

C₁₇H₁₁NO₃ 277.279

Alkaloid from *Monocyclanthus vignei* (Annonaceae). Brown powder.

O²-Me, N-Me: O,N-Dimethyliriodendronine

[59626-93-6]

C₁₈H₁₃NO₃ 291.306

Alkaloid from *Gutteria chrysopetala* (Annonaceae). Dark green needles (MeOH). Mp 275-278° dec. Zwitterionic oxoaporphine.

Di-Me ether: 1,2-Dimethoxydibenz[de,g]quinolin-7-one. Lysicamine. Dae-chualkaloid C. Oxonuciferine

[15444-20-9]

C₁₈H₁₃NO₃ 291.306

Alkaloid from a variety of genera in the Annonaceae (*Annona*, *Gutteria*, *Enantia*, *Polyalthia*), Araceae (*Lysichiton*), Magnoliaceae (*Liriodendron*), Menispermaceae (*Abuta*, *Chasmanthera*, *Stephania*, *Telitoxicum*) and Rhamnaceae (*Colubrina*, *Zizyphus*). Shows antimicrobial and nematocidal activity. Yellow needles. Mp 185-187° Mp 211-212° dec. λ_{max} 255 ; 283 ; 335 (no solvent reported). λ_{max} 235 (ε 29500); 270 (ε 25700); 307 (ε 5750); 400 (ε 8700) (EtOH) (Berdy). λ_{max} 249 (ε 21400); 276 (ε 27700); 306 (ε 6600); 453 (ε 3800) (EtOH/HCl) (Berdy).

▶RB6030000

Methylene ether: see Liriodenine, L-194

Katsui, N. *et al.*, *Tet. Lett.*, 1966, 6257

(*Lysicamine*, *isol*, *synth*, *uv*, *ir*, *pmr*)

Cava, M.P. *et al.*, *Tetrahedron*, 1972, **28**, 4299 (*synth*)

Senter, P.D. *et al.*, *Phytochemistry*, 1977, **16**, 2015 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Hufford, C.-D. *et al.*, *J. Polym. Sci.*, 1980, **69**, 1180-1187 (*Lysicamine*, *activity*)

Hocquemiller, R. *et al.*, *Plant. Med. Phytother.*, 1984, **18**, 165 (*O,N-Dimethyliriodendronine*)

Saá, C. *et al.*, *Tet. Lett.*, 1985, **26**, 4559

(*synth*)

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443 (*Lysicamine*, *isol*)

Pabuccuoglu, V. *et al.*, *Arch. Pharm.*

(*Weinheim*, *Ger.*), 1991, **324**, 29-33 (*synth*, *activity*)

Achenbach, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1331 (*7-Oxodehydroasimilobine*)


Atanes, N. *et al.*, *J.O.C.*, 1991, **56**, 2984 (*synth*)

Hsieh, J.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1999, **46**, 607-612 (*Lysicamine*, *isol*, *uv*, *ir*, *pmr*)

Camacho, M.D.R. *et al.*, *Planta Med.*, 2000, **66**, 478-480 (*N-Methyliriodendronine*)

Liriodenine**L-194**

8H-Benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. 1,2-Methylene-dioxyoaporphine. *Spermatheridine*. *Micheline B. Oxoushinsunine*
[475-75-2]



C₁₇H₉NO₃ 275.263

Alkaloid from a wide variety of genera in the Annonaceae (*Annona*, *Asimina*, *Cananga*, *Cleistopholis*, *Enantia*, *Fusaea*, *Isolona*, *Gutteria*, *Melodorum*, *Pseuduvaria*, *Hexalobus*, *Mitrella*, *Monodora*, *Pachypodanthium*, *Polyalthia*, *Uvariopsis*, *Schefferomitra*, *Xylopia*), Araceae (*Lysichiton*), Eupomatiaceae (*Eupomatia*), Lauraceae (*Litsea*, *Neolitsea*), Magnoliaceae (*Liriodendron*, *Magnolia*, *Michelia*, *Elmerrillia*, *Talauma*), Menispermaceae (*Pycnarrhena*, *Stephania*, *Chasmanthera*, *Pachygone*, *Rhigiocarya*), Monimiaceae (*Dryadodaphne*), Nelumbonaceae (*Nelumbo*), Papaveraceae (*Roemeria*, *Papaver*), Rhamnaceae (*Colubrina*), Rutaceae (*Zanthoxylum*), *Atherosperma*, *Doryphora*, *Laurelia* and Monimiaceae (*Atherosperma*, *Doryphora*, *Laurelia*, *Siparuna*). Shows leishmanicidal activity. Antineoplastic agent. Shows some cytotoxicity against human nasopharyngeal carcinoma cells. Shows good activity against *Trichophyton mentagrophytes* and *Syncephalestrum racemosum*. Also effective against barley net blotch, rice blast and broad bean chocolate spot disease. Muscarinic receptor antagonist, topoisomerase II inhibitor. Yellow needles (CHCl₃). Mp 275-276° dec. Log P 2.94 (uncertain value) (calc). Rel. nontoxic but shows some mutagenic activity. λ_{max} 257 (ε 21400); 277 (ε 18200); 329 (ε 4680); 392 (ε 4900); 455 (ε 3800) (EtOH/HCl) (Derep). λ_{max} 248 (ε 16600); 268 (ε 13500); 309 (ε 4170); 413 (ε 6610) (EtOH) (Derep). λ_{max} 247 (ε 10960); 267 (ε 9550); 310 (ε 2884); 413 (ε 4680) (MeOH) (Berdy). λ_{max} 204 ; 248 ; 268 ; 311 ; 415 (EtOH) (Berdy).

▶RB6036000

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1964, 1629 (*isol*, *ir*, *uv*, *pmr*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1403 (*ms*)

Kessar, S.V. *et al.*, *Tet. Lett.*, 1980, **21**, 3307 (*synth*, *uv*)

Nimgirawath, S. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1061 (*synth*)


Gupta, Y.P. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 429 (*synth*)

Fevrier, A. *et al.*, *Planta Med.*, 1999, **65**, 46-49 (*activity*)

Zhang, Z. *et al.*, *J. Nat. Prod.*, 2002, **65**, 856-859 (*cmr*)

Liriotulipiferine**L-195**

5,6,6a,7-Tetrahydro-1,9-dimethoxy-6-methyl-4H-dibenzo[de,g]quinoline-2,10-diol, 9CI. 2,10-Dihydroxy-1,9-dimethoxyaporphine



C₁₉H₂₁NO₄ 327.379

(S)-form [60755-81-9]

Alkaloid from the discoloured sapwood of *Liriodendron tulipifera* (Magnoliaceae). Needles (EtOAc). Mp 184-186°. [α]_D +174.4 (CCl₄).

Hydrochloride:

Needles (MeOH). Mp 218-220° dec.

N-Me:

C₂₀H₂₄NO₄[⊕] 342.414

Mp 214-216° dec. (as iodide).

N-De-Me: 2,10-Dihydroxy-1,9-dimethoxyaporphine. **Laetanine**
[72361-67-2]


C₁₈H₁₉NO₄ 313.352

Alkaloid from the bark of *Litsea laeta* (Lauraceae). Light-amber cryst. (EtOAc/MeOH). Mp 226-228°. [α]_D +105 (c, 0.4 in MeOH).

Chen, C.-L. *et al.*, *Phytochemistry*, 1976, **15**, 1161 (*isol*, *uv*, *pmr*, *ms*, *struct*)

Borthakur, N. *et al.*, *Phytochemistry*, 1979, **18**, 910 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *Laetanine*)

Rasoanaivo, P. *et al.*, *Planta Med.*, 1998, **64**, 58-62 (*cmr*, *Laetanine*)


Lissoclibadin 1**L-196**

$C_{39}H_{57}N_3O_6S_7$ 888.358
Isol. from *Lissoclinum* cf. *badium*. Active against *Ruegeria atlantica*. $[\alpha]_D$ -3.6 (c, 0.1 in $CHCl_3$) (as tris(TFA) salt). λ_{max} 279 (ϵ 134300); 318 (sh) (ϵ 26500) (tris(TFA) salt).

Liu, H. *et al.*, *Tetrahedron*, 2005, **61**, 8611-8615 (isol, pmr, cmr)

Lissoclibadin 2**L-197**

[865369-78-4]



$C_{26}H_{38}N_2O_4S_5$ 602.927

Disulfide analogue of Lissoclibadin 3, L-198. Isol. from *Lissoclinum* cf. *badium*. λ_{max} 270 (ϵ 14300); 316 (ϵ 6500) (as bis(trifluoroacetate)).

5-*De(methylthio)*, 3-*O-de-Me: Lissoclibadin 5*
[930086-79-6]
 $C_{24}H_{34}N_2O_4S_4$ 542.808
Isol. from *Lissoclinum* cf. *badium*. λ_{max} 255 (log ϵ 4.1); 302 (log ϵ 3.86) (MeOH) (as bis(trifluoroacetate)).


Bis(*demethylthio*), 3,6'-*di-O-de-Me: Lissoclibadin 4*
[930086-78-5]
 $C_{22}H_{30}N_2O_4S_3$ 482.688
Isol. from *Lissoclinum* cf. *badium*. λ_{max} 258 (log ϵ 4.06); 311 (log ϵ 3.81) (MeOH) (as bis(trifluoroacetate)).

Liu, H. *et al.*, *Tetrahedron*, 2005, **61**, 8611-8615 (*Lissoclibadin 2*)

Nakazawa, T. *et al.*, *J. Nat. Prod.*, 2007, **70**, 439-442 (*Lissoclibadins 4,5*)

Lissoclibadin 3**L-198**

[865376-21-2]



$C_{26}H_{38}N_2O_4S_4$ 570.861

Isomeric with Lissoclinotoxin E, L-209. Isol. from *Lissoclinum* cf. *badium*. Cytotoxic. Amorph. (as bis-TFA salt). λ_{max}

274 (ϵ 13300); 318 (ϵ 4500) (MeOH) (bis-TFA salt).

5-*De(methylthio)*, *O*³-*de-Me: Lissoclibadin 6*

[930086-80-9]

$C_{24}H_{34}N_2O_4S_3$ 510.742


Isol. from *Lissoclinum* cf. *badium*. λ_{max} 260 (log ϵ 4.25); 310 (log ϵ 3.93) (MeOH) (as bis-TFA salt).

Liu, H. *et al.*, *Tetrahedron*, 2005, **61**, 8611-8615 (*Lissoclibadin 3*)

Nakazawa, T. *et al.*, *J. Nat. Prod.*, 2007, **70**, 439-442 (*Lissoclibadin 6*)

Lissoclin C**L-199**

7-*Bromo-2,3,4,9-tetrahydro-1-(1H-imidazol-4-ylmethyl)-1H-pyrido[3,4-b]indole, 9CI*
[158761-14-9]



$C_{15}H_{15}BrN_4$ 331.214

Alkaloid from the tropical ascidian *Lissoclinum* sp. Isol. as a virtual racemate, prob. by racemisation during isol. λ_{max} 231 (ϵ 21900); 285 (ϵ 4170) (MeOH) (TFA salt).

Debromo: 2,3,4,9-Tetrahydro-1-(1H-imidazol-4-ylmethyl)-1H-pyrido[3,4-b]indole. Haploscleridamine

[426258-83-5]

$C_{15}H_{16}N_4$ 252.318


Alkaloid from a sponge of the order Haploscleridae. Cathepsin K inhibitor. Gum. $[\alpha]_D$ -3.4 (c, 0.78 in MeOH). Racemate or virtual racemate. λ_{max} 235 (log ϵ 3.72); 270 (log ϵ 3.41); 278 (log ϵ 3.55); 288 (log ϵ 3.06) (MeOH).

Searle, P.A. *et al.*, *J.O.C.*, 1994, **59**, 6600-6605 (isol, uv, ir, pmr, cmr, struct)

Patil, A.D. *et al.*, *J. Nat. Prod.*, 2002, **65**, 628-629 (*Haploscleridamine*)

Lissoclinamide 1**L-200**

[87411-84-5]




$C_{35}H_{43}N_7O_5S_2$ 705.901

Isol. from the marine tunicate *Lissoclinum patella*. Related to Ulicyclamide, U-9.

Wasylyk, J.M. *et al.*, *J.O.C.*, 1983, **48**, 4445-4449 (isol, ir, pmr, ms, struct)

Lissoclinamide 4**L-201**

[120853-16-9]



$C_{38}H_{43}N_7O_5S_2$ 741.934

Cyclopeptide from the ascidian *Lissoclinum patella*. Exhibits marked cytotoxicity. Powder (Et_2O). Mp 152-154°. $[\alpha]_D$ +45 (c, 0.7 in $CHCl_3$).

18 α ,19-*Dihydro: Lissoclinamide 7*

[126297-39-0]

$C_{38}H_{45}N_7O_5S_2$ 743.949

Isol. from *Lissoclinum patella*. Shows selective metal binding props. Cytotoxic.

28,29-*Didehydro: Lissoclinamide 5*

[120853-17-0]

$C_{38}H_{41}N_7O_5S_2$ 739.918

From *Lissoclinum patella*. Cytotoxic (less than Lissoclinamide 4 by two orders of magnitude).

21-*Epimer: Lissoclinamide 6*

[121209-53-8]

$C_{38}H_{43}N_7O_5S_2$ 741.934

Trace const. of *Lissoclinum patella*. Cytotoxic. Possibly an artifact.

Stereoisomer: Lissoclinamide 8

[126452-98-0]

$C_{38}H_{43}N_7O_5S_2$ 741.934

Isol. from *Lissoclinum patella*. Cytotoxic. Full config. not detd.

Degnan, B.M. *et al.*, *J. Med. Chem.*, 1989, **32**, 1349-1354 (isol, pmr, cmr, struct)

Schmitz, F.J. *et al.*, *J.O.C.*, 1989, **54**, 3463-3472 (isol, ir, pmr, cmr, struct)

Hawkins, C.J. *et al.*, *J. Med. Chem.*, 1990, **33**, 1634-1638 (*Lissoclinamides 7 and 8*)


Wipf, P. *et al.*, *J.A.C.S.*, 1996, **118**, 12358-12367; 1998, **120**, 4105-4112 (*Lissoclinamide 7, synth, abs config, conformn*)

Boden, C.D.J. *et al.*, *J.C.S. Perkin 1*, 2000, 875-882 (*synth, abs config*)


Morris, L.A. *et al.*, *Tetrahedron*, 2001, **57**, 3199-3207 (activity)

Lissoclinamide 9



[309963-07-3]

Absolute
ConfigurationC₃₅H₄₅N₇O₅S₂ 707.916Isol. from *Lissoclinum patella*. Pale yellow oil. λ_{max} 308 (ε 3370) (CHCl₃).Morris, L.A. *et al.*, *Tetrahedron*, 2000, **56**, 8345-8353 (*isol, pmr, cmr, cd*)**Lissoclinamide 10**




[309963-08-4]

C₃₆H₄₉N₇O₅S₂ 723.959Isol. from *Lissoclinum patella*. Pale yellow oil. λ_{max} 308 (ε 2900) (CHCl₃).Morris, L.A. *et al.*, *Tetrahedron*, 2000, **56**, 8345-8353 (*isol, pmr, cmr, cd*)**Lissoclinidine**

[496909-77-4]


C₂₀H₁₇N₃O₂S 363.439Alkaloid from the ascidian *Lissoclinum notti*. Hygroscopic purple solid (as TFA salt). λ_{max} 218 (log ε 4.41); 281 (log ε 4.26); 295 (log ε 4.21); 310 (log ε 4.1); 384 (log ε 3.61); 548 (log ε 3.4) (MeOH/TFA).Appleton, D.R. *et al.*, *Tetrahedron*, 2002, **58**, 9779-9783 (*isol*)**L-202****Lissoclinotoxin A**9-(2-Aminoethyl)-6-methoxy-7-benzopentathiepinol
[133883-05-3]C₉H₁₁NO₂S₅ 325.521Originally reported with an incorrect trithiane struct. Chiral molecule exhibiting unusual stereoisomerism due to restricted inversion, although isol. samples are racemic. Isol. from the tunicate *Lissoclinum perforatum*. Exhibits potent antimicrobial and antifungal activity. Amorph. yellow solid. Fairly sol. MeOH, DMSO; poorly sol. butanol, hexane. Mp 245-250°. λ_{max} 215 (ε 10400); 246 (ε 8870) (EtOH) (Derep). λ_{max} 213 (ε 14100) (MeOH) (Berdy).*Me ether*: 8,9-Dimethoxy-6-benzopentathiepinethanamine, 9CI. **Varacin** [134029-48-4]C₁₀H₁₃NO₂S₅ 339.548Isol. from the ascidians *Lissoclinum vareau* and *Polycitor* sp. Exhibits potent antifungal and antimicrobial activity. Cytotoxic towards the human colon cancer HCT 116. Light yellow powder (also descr. as a glass). Mp 258-260° dec. λ_{max} 214 ; 244 (MeOH) (Derep). λ_{max} 212 ; 244 (EtOH) (Berdy).*Me ether, N-Ac*:

Light yellow cryst. (MeOH). Mp 136-137°.


O-De-Me: 9-(2-Aminoethyl)-6,7-benzopentathiepinol, 9CI. **Dide-O-methylvaracin**. *De-O-methyl*lissoclinotoxin A [159645-89-3]C₈H₉NO₂S₅ 311.495Isol. from a *Eudistoma* sp. Yellow powder.8-(*Methylthio*), *Me ether*: 9-(2-Aminoethyl)-6,7-dimethoxy-8-(*methylthio*)benzopentathiepin. 5-(*Methylthio*)**varacin** [159645-87-1]C₁₁H₁₅NO₂S₆ 385.641Isol. from a *Lissoclinum* sp. CAS no. refers to trifluoroacetate.8-(*Methylthio*), N,N,O-*tri-Me*: 6-Amino-8,9-dimethoxy-N,N-dimethyl-7-(*methylthio*)benzopentathiepin. N,N-Dimethyl-5-(*methylthio*)**varacin** [159645-83-7]C₁₃H₁₉NO₂S₆ 413.695Isol. from *Lissoclinum japonicum*. Pale yellow oil (as trifluoroacetate). CAS no. refers to trifluoroacetate. λ_{max} 211 (ε 16500) (MeOH).Davidson, B.S. *et al.*, *J.A.C.S.*, 1991, **113**,4709-4710 (*Varacin*)
Litaudon, M. *et al.*, *Tet. Lett.*, 1991, **32**, 911-914 (*isol, pmr, cmr*)Ford, P.W. *et al.*, *J.O.C.*, 1994, **59**, 5955-5960 (*synth, struct*)Searle, P.A. *et al.*, *J.O.C.*, 1994, **59**, 6600-6605 (*isol, uv, ir, pmr, cmr*)**L-205**Litaudon, M. *et al.*, *Tetrahedron*, 1994, **50**, 5323-5334 (*uv, pmr, ms, struct*)Compagnone, R.S. *et al.*, *Tetrahedron*, 1994, **50**, 12785-12792 (*isol, uv, pmr, cmr*)Makarieva, T.N. *et al.*, *J. Nat. Prod.*, 1995, **58**, 254 (*Varacin*)Greer, A. *et al.*, *J.A.C.S.*, 2001, **123**, 10379-10386 (*Varacin, struct*)**Lissoclinotoxin B****L-206**8,9,10,11-Tetrahydro-6-methoxy-1,2,3,4,5-pentathiepio[6,7-*f*]isoquinolin-7-ol, 9CI
[157536-34-0]C₁₀H₁₁NO₂S₅ 337.532Minor const. of the tunicate *Lissoclinum perforatum*. Exhibits antimicrobial activity. Antiplasmodial agent. Pale yellow powder. Fairly sol. MeOH; poorly sol. CHCl₃, CH₂Cl₂. Mp 310-313°. λ_{max} 215 (ε 20000); 246 (ε 6500) (MeOH) (Berdy).Litaudon, M. *et al.*, *Tetrahedron*, 1994, **50**, 5323 (*isol, uv, ir, pmr, cmr, struct*)**Lissoclinotoxin C****L-207**5-(2-Aminoethyl)-2-methoxy-3,4-bis(-methylthio)phenol, 9CI. 5-Hydroxy-4-methoxy-2,3-bis(methylthio)phenethylamine
[158761-15-0]C₁₁H₁₇NO₂S₂ 259.393Alkaloid from the tropical ascidian *Lissoclinum* sp.Searle, P.A. *et al.*, *J.O.C.*, 1994, **59**, 6600-6605 (*isol, pmr, struct*)**Lissoclinotoxin D****L-208**4,10-Bis(2-aminoethyl)-1,7-dimethoxydi-benzo[*c,g*][1,2,5,6]-tetrathiocin-2,8-diol, 9CI
[158761-16-1]C₁₈H₂₂N₂O₄S₄ 458.647Alkaloid from the tropical ascidian *Lissoclinum* sp. Exhibits antifungal activity. 3,3'-*Di-O-de-Me*, 4,4'-*di-Me ether*, N-

tetra-Me: Lissoclibadin 7


[930086-81-0]

C₂₂H₃₀N₂O₄S₄ 514.754Isol. from *Lissoclinum* cf. *badium*. λ_{max} 260 (log ε 4.03); 322 (log ε 3.88) (MeOH) (as bis(trifluoroacetate)).Searle, P.A. et al., *J.O.C.*, 1994, **59**, 6600-6605 (isol, pmr, struct)Nakazawa, T. et al., *J. Nat. Prod.*, 2007, **70**, 439-442 (*Lissoclibadin 7*)**Lissoclinotoxin E****L-209***Lissoclin*C₂₆H₃₈N₂O₄S₄ 570.861Struct. revised from, and isomeric with Lissoclibadin 3, L-198 in 2005. Isol. from *Lissoclinum* cf. *badium* and an unidentified Philippine ascidian. Cytotoxic. λ_{max} 210 (ε 15000); 232 (sh) (ε 9000); 254 (ε 7000); 272 (ε 12000) (MeOH).**5,10-Dioxide: Lissoclin disulfoxide**


[195303-92-5]

C₂₆H₃₈N₂O₆S₄ 602.86Isol. from a *Lissoclinum* sp. Inhibitor of interleukin-8 receptors. Yellow solid. λ_{max} 243; 266; 318 (MeOH).Patil, A.D. et al., *Nat. Prod. Lett.*, 1997, **10**, 225-229 (*Lissoclin disulfoxide*)Davies, R.A. et al., *Tetrahedron*, 2003, **59**, 2855-2859 (isol, pmr, cmr)Liu, H. et al., *Tetrahedron*, 2005, **61**, 8611-8615 (isol, struct)**Lissoclinotoxin F****L-210**C₂₆H₃₈N₂O₄S₅ 602.927Isol. from a Philippine didemnid ascidian. Cytotoxic. Light brown film (as bis-TFA salt). λ_{max} 210 (ε 13000); 238 (sh) (ε 11000); 268 (ε 11000); 320 (ε 3000) (MeOH).Davis, R.A. et al., *Tetrahedron*, 2003, **59**, 2855-2859 (isol, pmr, cmr)**Litebamine****L-211**

1,2,3,4-Tetrahydro-9,11-dimethoxy-2-methylnaphth[2,1-f]isoquinoline-8,12-diol, 9CI
[137031-56-2]

C₂₀H₂₁NO₄ 339.39Alkaloid from *Litsea cubeba* (mountain pepper) (Lauraceae). Cryst. (MeOH). Mp 218-220°.Wu, Y.-C. et al., *Tet. Lett.*, 1991, **32**, 4169 (isol, pmr, cmr, ir, uv, struct)Lee, S.-S. et al., *Tet. Lett.*, 1992, **33**, 6309 (synth)Hara, H. et al., *Tetrahedron*, 1995, **51**, 10189 (synth)**Lithosenine****L-212**

[159903-58-9]

C₂₀H₃₃NO₈ 415.483Alkaloid from aerial parts of *Lithospermum officinale* (Boraginaceae).**O^{3''}-Ac: Acetylithosenine**


[159690-29-6]

C₂₂H₃₅NO₉ 457.52Alkaloid from aerial parts of *Lithospermum officinale* (Boraginaceae).**3'-Deoxy, stereoisomer: Uluganine**

[55437-97-3]

C₂₀H₃₃NO₇ 399.483Alkaloid from *Ulugbekia tschinganica* (Boraginaceae) (preferred genus name *Arnebia*). Cryst. (Me₂CO). Mp 106-107°. [α]_D²⁵ -31.9 (c, 0.47 in Me₂CO). Stereochem. not detd.Khasanova, M.A. et al., *Khim. Prir. Soedin.*, 1974, **10**, 809; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 842 (*Uluganine*)Krenn, L. et al., *Phytochemistry*, 1994, **37**, 275 (*Lithosenine, Acetylithosenine*)**Lithospermoside****L-213**

[63492-69-3]

C₁₄H₁₉NO₈ 329.306Constit. of *Lithospermum officinale*, *Lithospermum caerulum* and *Thalictrum orientale*. Cryst. (MeOH aq.). Mp 278-279°. [α]_D²⁰ -156 (c, 0.99 in H₂O).**4-Me ether: Bauhinin**

[100757-58-2]

C₁₅H₂₁NO₈ 343.333Constit. of *Bauhinia championii*. Cryst. + 2H₂O (Me₂CO aq.). Mp 213-214°. [α]_D²⁵ -80 (c, 1.05 in MeOH). λ_{max} 258 (19700) (MeOH).**5-Deoxy: Menisdaurin**

[67765-58-6]

C₁₄H₁₉NO₇ 313.307Isol. from *Ilex warburgii*, *Menispermum dauricum* and *Semiaquilegia adoxoides*. Plates. Mp 175-176°. [α]_D¹⁵ -185.4 (c, 1.00 in MeOH).**E-Isomer, aglycone: [845536-74-5]**C₈H₉NO₃ 167.164Constit. of the roots of *Semiaquilegia adoxoides*. [α]_D²⁰ +54.3 (c, 1 in MeOH).**4-Epimer: Dasycarponin**

[72523-59-2]


Isol. from roots of *Thalictrum dasycarpum*. Rhombic cryst. (MeOH aq.). Mp 253-255°. [α]_D²⁷ -11 (c, 0.25 in H₂O).**4-Epimer, aglycone: Dasycarponilide†**Mp 164-165°. [α]_D²⁷ +370 (c, 0.01 in MeOH).**5-Epimer: [84799-31-5]**C₁₄H₁₉NO₈ 329.306Constit. of *Ilex warburgii* and *Semiaquilegia adoxoides*. Cryst. (MeOH aq.). Mp 221-223°. [α]_D²⁵ -247 (c, 0.611 in MeOH).**6-Epimer, 5-deoxy: Purshianin**

[162990-70-7]

C₁₄H₁₉NO₇ 313.307Isol. from *Purshia tridentata*, Plates (EtOH). Mp 167-169.5°. [α]_D²⁵ -90.2 (c, 0.45 in MeOH). λ_{max} 271 (log ε 3.99) (EtOH).**5,6-Diepimer, E-isomer, aglycone: Huazhongilexol**

[161407-80-3]

C₈H₉NO₃ 167.164Constit. of *Ilex centrocimensis*.**4,5,6-Triepimer: Griffonin**C₁₄H₁₉NO₈ 329.306Constit. of roots of *Griffonia simplicifolia*. Feathery cryst. (MeOH). Mp 263-265° dec. [α]_D²¹ +6 (c, 0.5 in H₂O). Griffonin is given the same reg. no. as Lithospermoside in CA but appears to be a distinct diastereoisomer. The opt. rotns. are v. different.Dwuma-Badu, D. et al., *J. Nat. Prod.*, 1976, **39**, 213 (*Griffonin*)Sosa, A. et al., *Phytochemistry*, 1977, **16**, 707 (isol, struct)Takahashi, K. et al., *Chem. Pharm. Bull.*, 1978, **26**, 1677 (*Menisdaurin*)Wu, J. et al., *J. Nat. Prod.*, 1979, **42**, 500 (*Dasycarponin*)Ueda, K. et al., *Chem. Lett.*, 1983, 149 (isol, struct)Chen, C.C. et al., *J. Nat. Prod.*, 1985, **48**, 933 (*Bauhinin*)Nakanishi, T. et al., *Chem. Pharm. Bull.*, 1994, **42**, 2251-2255 (*Menisdaurin, Purshianin*)Lin, L.D. et al., *Huaxue Xuebao*, 1995, **53**, 98; *CA*, **122**, 183194 (*Huazhongilexol*)Josien-Lefebvre, D. et al., *Helv. Chim. Acta*, 2001, **84**, 890-897; 2003, **86**, 661-672; 2007, **90**, 19-30 (synth, pmr, cmr)Erdemgil, F.Z. et al., *Z. Naturforsch., C*, 2003, **58**, 632-636 (isol, pmr, cmr)Zhang, H. et al., *Chin. J. Chem.*, 2004, **22**, 1200-1203 (*Semiaquilegia aglycone*)Seigler, D.S. et al., *Phytochemistry*, 2005, **66**, 1567-1580 (*Menisdaurin*)Niu, F. et al., *Chin. J. Chem.*, 2006, **24**, 1788-1791 (*5-epimer*)

Litsericine*Norhexahydromecambrine*C₁₇H₂₁NO₃ 287.358**(+)-form** [15650-04-1]

Alkaloid from *Neolitsea sericea*, *Neolitsea buisanensis* and *Neolitsea aurata* (Lauraceae). Mp 156°. [α]_D²⁵ +67 (c, 1.33 in EtOH).

O,N-Di-Ac: Mp 154°.

N-Me: N-Methylitsericine

[5890-36-8]

C₁₈H₂₃NO₃ 301.385

Alkaloid from *Neolitsea aurata* (Lauraceae). Mp 195-196° (185°). [α]_D²⁵ +32 (Me₂CO). [α]_D¹⁶ +53.2 (c, 1.09 in EtOH).

N,N-Di-Me: Mp 173-175° (as iodide).

10-Epimer: Lauformine

[91177-60-5]

C₁₇H₂₁NO₃ 287.358

Alkaloid from the bark of *Phoebe formosana* (Lauraceae). Needles (Me₂CO). Mp 235-237°. [α]_D²³ +62 (c, 1.0 in MeOH). λ_{max} 240 (sh) (log ε 3.32); 290 (log ε 3.36) (EtOH).

10-Epimer, N-Me: N-Methylauformine

[91199-02-9]

C₁₈H₂₃NO₃ 301.385

Alkaloid from the bark of *Phoebe formosana* (Lauraceae). Needles (Me₂CO). Mp 255-256°. [α]_D²³ +68 (c, 0.1 in MeOH). λ_{max} 240 (sh) (log ε 3.18); 292 (log ε 3.23) (EtOH).

(-)-form

N-Me: Hexahydromecambrine. *Hexahydrofugapavine*

C₁₈H₂₃NO₃ 301.385

Alkaloid from *Papaver lecoquii* and *Phoebe scortechinii*. Prisms (MeOH). Mp 266-267°. [α]_D²⁵ -44 (c, 0.09 in MeOH). λ_{max} 211 (log ε 4.5); 241 (sh) (log ε 3.61); 260 (log ε 3.13); 287 (log ε 3.63) (EtOH).

Nakasato, T. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 129; 134; 1205 (*isol, uv, pmr, struct, config*)


Slávik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 1558-1566; 1981, **46**, 2587-2593 (*Hexahydromecambrine, isol, uv, ord, struct, synth*)

Lu, S.T. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1975, **22**, 349; *CA*, **84**, 132659f (*N-Methylitsericine*)

Lu, S.T. *et al.*, *Heterocycles*, 1984, **22**, 1031-1033; 1323-1326 (*Litsericine, Lauformine, N-Methylauformine, isol, uv, ir, pmr, ms, struct*)

L-214**(+)-form****Littoraline**

[171828-68-5]


C₁₈H₂₁NO₅ 331.368

Alkaloid from bulbs of *Hymenocallis littoralis* and *Hymenocallis rotata* (Amaryllidaceae). Shows inhibitory activity of HIV reverse transcriptase. Powder. Sol. MeOH. Not obt. completely pure. Stereochem. at C-4a not absolutely proved, no further studies to 2001. λ_{max} 243 (ε 12590); 282 (ε 6456) (MeOH).

Lin, L.-Z. *et al.*, *Phytochemistry*, 1995, **40**, 1295-1298 (*isol, uv, pmr, cmr*)

Littorine

α-Hydroxybenzenepropanoic acid 8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester, 9CI. 3α-(2-Hydroxy-3-phenylpropionyl-oxy)tropane. Tropan-3α-yl phenyllactate

C₁₇H₂₃NO₃ 289.374**(R)-form** [21956-47-8]

Alkaloid from *Anthocercis littorea* and *Datura sanguinea* (Solanaceae). Mp 96-97°. [α]_D²⁶ -12.7 (c, 4.42 in EtOH).

Picrate:

Cryst. (EtOH). Mp 162-163°.

Methiodide: Mp 167-168°.**(±)-form**

Synthetic. Powerful mydriatic agent. Cryst. (also descr. as an oil). Mp 89-90°.

Hydrochloride: Mp 173-175°.**Picrate:**

Short yellow needles. Mp 159-160°.

[62083-53-8]


Jowett, H.A.D. *et al.*, *J.C.S.*, 1909, **95**, 1020 (*synth*)

Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 221 (*isol, struct, uv, ir*)

Takeuchi, Y. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2603 (*synth, ir, pmr*)

Lituarine C

[143621-77-6]

**L-215**C₃₈H₅₅NO₁₁ 701.853

Structures may need revision; spectral properties of synthetic compds. do not match those of natural Lituarines (2008). Isol. from the sea pen *Lituarina australasiae*. Shows antifungal, cytotoxic and antineoplastic activities. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 153-157°. λ_{max} 269 (ε 22419) (MeOH) (Berdy).

5-Ac: Lituarine B

[143621-76-5]

C₄₀H₅₇NO₁₂ 743.89

Isol. from *Lituarina australasiae*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 126-129°. λ_{max} 269 (ε 26604) (MeOH) (Berdy).

4,5-Dideoxy: Lituarine A

[143621-75-4]

C₃₈H₅₅NO₉ 669.854


Isol. from *Lituarina australasiae*. Cryst. Mp 83-85°. λ_{max} 270 (ε 15857) (MeOH) (Berdy).

Vidal, J.-P. *et al.*, *J.O.C.*, 1992, **57**, 5857-5860 (*isol, pmr, cmr*)

Smith, A.B. *et al.*, *J.A.C.S.*, 2008, **130**, 422-423 (*synth*)

Lobelanidine, 8CI**L-218**

1-Methyl-α,α'-diphenyl-2,6-piperidine-diethanol, 9CI. 2,6-Bis(2-hydroxy-2-phenylethyl)-1-methylpiperidine. 8,10-Diphenyllobelidol
[552-72-7]

**(2RS,6SR,8RS,10SR)-form**C₂₂H₂₉NO₂ 339.477

Two stereoisomers occur naturally. Strictly the name Lobelanidine refers to the *meso*-isomer. Other stereoisomers known synthetically.
▶ Toxic. TM7132400

(2RS,6SR,8RS,10SR)-form

Alkaloid from *Lobelia inflata*, *Lobelia hassleri*, *Lobelia stallfeldii*, other *Lobelia* spp. and *Isotoma longiflora* (preferred genus name *Laurentia*). Lobelanidine of undetd. stereochem. also isol. from *Sedum acre* (Campanulaceae, Crassulaceae). Prisms (EtOH). Mp 150°. Opt. inactive (*meso*-).

Di-Ac: Mp 214-215°.**O-β-D-Glucopyranoside:**C₂₈H₃₉NO₇ 501.619

Alkaloid from *Sedum acre* (Crassulaceae). Mp 233-236° (as penta-Ac). [α]_D²² -28 (MeOH) (penta-Ac).

O-β-D-Glucopyranoside, penta-Ac: Mp 233-236° dec. [α]_D²² -28 (MeOH).

Diketone: Lobelanine. 2,2'-(1-Methyl-2,6-piperidinediyl) bis[1-phenylethanone], 9CI. 1-Methyl-2,6-diphenacylpiperidine. 8,10-Diphenyllobelidione
[579-21-5]

C₂₂H₂₅NO₂ 335.445

Alkaloid from *Lobelia inflata* (Campanulaceae). Plates (petrol). Mp 99° (94-95°). Opt. inactive (*meso*-).

Diketone; hydrochloride:

Cryst. (EtOH). Mp 196° (188°).

Diketone, N-oxide: Mp 84-86°.

N-De-Me: Norlobelanidine. α, α' -Diphenyl-2,6-piperidinediethanol, 9CI. 8,10-Diphenylnorlobelidol [495-49-8]

C₂₁H₂₇NO₂ 325.45

Main alkaloid from *Lobelia polyphylla* (Campanulaceae). Needles (Et₂O). Mp 120°. Opt. inactive.

N-De-Me; hydrochloride:

Needles (EtOH aq.). Mp 254° (244°).

Diketone, N-de-Me: Norlobelanine. 2,6-Diphenacylpiperidine. 8,10-Diphenyl-norlobelidone. *Isolobelanine*

C₂₁H₂₃NO₂ 321.418

Alkaloid from *Lobelia* sp. (Campanulaceae). Prisms (Et₂O or EtOH aq.). Mp 120-121° (117-118°). Opt. inactive (*meso*-).

Diketone, N-de-Me; hydrochloride:

Cryst. (EtOH). Mp 201-202°.

Monoketone: see Lobeline, L-219

(2S,6R,8S,10S)-form

(-)-*cis-form*

Alkaloid from *Isotoma longiflora* (Campanulaceae). Prismatic cryst. (Et₂O). Mp 117°. [α]_D²⁰ -71.8 (c, 1.16 in CHCl₃).

Hydrate:

Needles (Me₂CO). Mp 216-218° dec.

Nitrate:

Needles (MeOH). Mp 216-217°.

Wieland, H. *et al.*, *Annalen*, 1925, **444**, 40; 1929, **473**, 83; 102; 118; 126 (*isol. struct. synth*)

Scheuing, G. *et al.*, *Annalen*, 1929, **473**, 126 (*synth*)

Ebnöther, A. *et al.*, *Helv. Chim. Acta*, 1958, **41**, 386

Parker, W. *et al.*, *J.C.S.*, 1959, 2433 (*synth. Lobelanine*)

Arthur, H.R. *et al.*, *J.C.S.*, 1963, 750 (*isol*)

Schöpf, C. *et al.*, *Annalen*, 1965, **682**, 206; **687**, 241 (*abs config*)

Weinges, K. *et al.*, *Annalen*, 1972, **756**, 177 (*Norlobelanidine*)

Krochmal, A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 303 (*isol*)

Shah, C.S. *et al.*, *Phytochemistry*, 1972, **11**, 2884 (*isol*)

O'Donovan, D.G. *et al.*, *J.C.S. Perkin 1*, 1975, 415 (*biosynth*)


Francis, L.P.S. *et al.*, *Planta Med.*, 1977, **32**, 268 (*isol*)

Moreira, E. *et al.*, *CA*, 1981, **95**, 183865q (*isol*)

Piccinni-Leopardi, C. *et al.*, *Bull. Soc. Chim. Belg.*, 1987, **96**, 97 (*Lobelanidine glycoside*)

Lobeline, BAN, INN L-219

2-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-1-phenylethanol, 9CI. 8,10-Diphenyllobelionol. α -Lobeline. *Inflatine. Lobelidine. Many other names*



(-)-form

C₂₂H₂₇NO₂ 337.461

Lobelidine was originally the racemate.

Antiasthmatic, antibronchitic, analeptic agent. Diuretic, respiratory stimulant. Possesses expectorant props. Used in antismoking preparations. Log P 3.74 (calc).

(-)-form [90-69-7]

Alkaloid from *Lobelia inflata*, *Lobelia nicotianaefolia*, *Lobelia hassleri*, *Lobelia stalfeldii* and several other *Lobelia* spp. Also isol. from seeds of *Campanula medium* (Campanulaceae). Used as a 0.1M aq. soln. for photometric detn. of Mo (λ_{\max} 465 nm, ϵ 13800). Needles (EtOH). Mp 130-131°. [α]_D¹⁵ -42.85 (EtOH). Log P 3.74 (calc). Oxidn. gives the symmetrical Lobelanine which is *meso*-.

▶ Adverse gastrointestinal and CNS effects by ingestion. LD₅₀ (mus, ipr) 107 mg/kg. OJ8480000

Hydrochloride: Zoolobelin. Lobran

[134-63-4]

Needles (EtOH). Mp 182°.

▶ LD₅₀ (mus, ipr) 40 mg/kg. OJ8490100

Sulfate (1:2): [134-64-5]

Cryst. (EtOH). [α]_D²⁵ -25 (c, 0.2 in CHCl₃).

▶ OJ8490170

Benzoyl: Mp 155-157°.

(±)-form [134-65-6]

Alkaloid from *Lobelia inflata* and *Lobelia salicifolia* (Campanulaceae). Prisms. Mp 110°.

Hydrochloride: Mp 170°.

[10122-32-4]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 47B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 868A (nmr)

Wieland, H. *et al.*, *Annalen*, 1929, **473**, 118 (*isol*)

Schöpf, C. *et al.*, *Annalen*, 1965, **682**, 206; **687**, 241 (*abs config*)

Cambar, P.J. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1969, **177**, 1 (*pharmacol*)

Korczyn, A.D. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1969, **182**, 370 (*pharmacol*)

Döpke, W. *et al.*, *Pharmazie*, 1970, **25**, 128 (*isol*)

Krochmal, A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 303 (*isol*)

Shah, C.S. *et al.*, *Phytochemistry*, 1972, **11**, 2884 (*isol, occur*)

Mansuri, S.M. *et al.*, *Arzneim.-Forsch.*, 1973, **23**, 1721 (*pharmacol*)

O'Donovan, D.G. *et al.*, *J.C.S. Perkin 1*, 1975, 415 (*biosynth*)

Alhadeff, M. *et al.*, *Drugs of Today (Barcelona)*, 1977, **13**, 236 (*rev*)

Nytko, K. *et al.*, *Microchem. J.*, 1980, **25**, 548 (*detn, Mo*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 6579

Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1990, **19**, 261 (*rev*)

Glaser, R. *et al.*, *J.C.S. Perkin 2*, 1992, 1071 (*cryst struct, abs config*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1227

Flammia, D. *et al.*, *J. Med. Chem.*, 1999, **42**, 3726-3731 (*sar, pharmacol*)

Felplin, F.-X. *et al.*, *J.O.C.*, 2002, **67**, 9192-9199 (*synth*)

Felplin, F.-X. *et al.*, *Tetrahedron*, 2004, **60**, 10127-10153 (*rev*)


Birman, V.B. *et al.*, *Org. Lett.*, 2007, **9**, 3237-3243 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LHY000; LHZ000

Lobinaline

L-220

Decahydro-1-methyl-5,7-diphenyl-6-(3,4,5,6-tetrahydro-2-pyridinyl)quinoline, 9CI [6887-36-1]



Relative Configuration

C₂₇H₃₄N₂ 386.579

Originally considered to be C₂₈H₃₈N₂O, then C₂₈H₃₆N₂. Abs stereochem. not reported to 1996 although it is reported in CAS. Major alkaloid from *Lobelia cardinalis*, also detected in *Lobelia elongata* (Campanulaceae). Cryst. (hexane or by subl.). Mp 108-110° (94-95°). [α]_D²⁰ +38 (CHCl₃).

Hydrochloride:

Cryst. + ½H₂O (CHCl₃/Me₂CO). Mp 220°.

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1938, **16**, 445 (*isol*)

Robison, M.M. *et al.*, *J.O.C.*, 1966, **31**, 3206-3213; 3220-3223 (*ms, pmr, struct*)


Clugston, D.M. *et al.*, *Can. J. Chem.*, 1967, **45**, 39-47 (*ms*)

Gupta, R.N. *et al.*, *Can. J. Chem.*, 1971, **49**, 384-397 (*biosynth*)

Lobinandinine

L-221

8-Ethyl-10-phenyldehydrolobelidol



Relative configuration

C₁₈H₂₇NO₂ 289.417

Posn. of the double bond not thoroughly established. Alkaloid from *Lobelia inflata* (Campanulaceae). Small plates (petrol). Mp 95°. [α]_D -120 (EtOH). Hydrogenation of the double bond gives a Lelobanidine, L-86 diastereoisomeric with the known alkaloids.

Hydrochloride: Mp 169°.

2'-Ketone: **Lobinine.** 8-Ethyl-10-phenyldehydrolobelionol

C₁₈H₂₅NO₂ 287.401

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Noncryst. Needles (EtOH/Et₂O)(as hydrochloride). Mp 144° (hydrochloride). [α]_D -106.1 (EtOH) (hydrochloride). Originally considered to be C₁₈H₂₇NO₂.

Diastereoisomer: Isolobinandinine

C₁₈H₂₇NO₂ 289.417

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Amorph. A 2,6-cis-isomer; hydrogenation gives (-)-Lelobanidine I, L-86.

Diastereoisomer, hydrochloride:

Cryst. + 2H₂O. Mp 111°. [α]_D -28.3 (H₂O).

Diastereoisomer, 2'-ketone: Isolobinine

C₁₈H₂₅NO₂ 287.401

Minor alkaloid from *Lobelia inflata* (Campanulaceae). Shows antiasthmatic, antitussive and hypertensive props. Cryst. (petrol). Mp 78°. A 2,6-*cis*-isomer giving (-)-Lelobanidine I, L-86 on hydrogenation.

Diastereoisomer, 2'-ketone, hydrochloride:

Cryst. + 1H₂O. Mp 154° (anhyd.). [α]_D -76 (H₂O).

Wieland, H. *et al.*, *Annalen*, 1931, **491**, 14; 1939, **540**, 103 (*isol, struct*)

Thomä, O. *et al.*, *Annalen*, 1939, **540**, 99 (*Isolobinine*)

Schöpf, C. *et al.*, *Annalen*, 1957, **608**, 88 (*struct, nomencl*)

Lobine**L-222**


C₂₃H₃₁N₃O₄ 413.516

Struct. unknown. Alkaloid from *Oxylobium parviflorum* (Fabaceae).

Mann, E.A. *et al.*, *Proc. R. Soc. London, B*, 1907, **79**, 485-491 (*isol*)

Lobscurinol**L-223**

[123118-12-7]



C₁₇H₂₅NO₂ 275.39

Minor alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Oil.

Ac: Acetyllobscurinol

C₁₉H₂₇NO₃ 317.427

Minor alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Opaque solid. [α]_D +110 (c, 0.16 in MeOH).

5-Epimer: Epilobscurinol

[123165-90-2]


C₁₇H₂₅NO₂ 275.39

Trace alkaloid from *Lycopodium obscurum* (Lycopodiaceae). No phys. props. reported.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lochneridine**L-224**

Methyl 2,16-didehydro-20-hydroxycuran-17-oate, 9CI



(-)-form

C₂₀H₂₄N₂O₃ 340.421

Originally considered to be C₁₉H₂₄N₂O₃. Publ. information on the nat. occurrence of Lochneridine stereoisomers is incomplete. Some isolates are of unreported opt. rotn.

(-)-form [5980-01-8]

Alkaloid from *Catharanthus roseus* (Apocynaceae). Prisms (MeOH). Mp 211-214°. [α]_D -608. p*K*_a 5.5. The original publication erroneously gave a (+)- optical rotn. λ_{\max} 206; 230; 298; 328 (EtOH).

(+)-form

Alkaloid from *Tabernaemontana pandacacui* and *Alstonia scholaris* (Apocynaceae). Mp 225°. [α]_D +650 (CHCl₃). Stereochem. not certain. It is likely that this is 20-*epi*(+)-Lochneridine, which is the probable alkaloid on biogenetic grounds.

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1961, **50**, 409-413 (*isol, ir, uv*)

Nakagawa, Y. *et al.*, *Chem. Ind. (London)*, 1962, 1986 (*struct*)

Djerassi, C. *et al.*, *Experientia*, 1963, **19**, 467-469 (*stereochem*)

Lathuilliere, P. *et al.*, *Ann. Pharm. Fr.*, 1966, **24**, 547-549 (*(+)-form, isol*)

Scott, A.I. *et al.*, *J.A.C.S.*, 1974, **96**, 2273-2274 (*biosynth*)

Mirand, C. *et al.*, *Tet. Lett.*, 1982, **23**, 1257-1258 (*20-Epilochneridine, synth*)

Banerji, J. *et al.*, *Indian J. Chem., Sect. B*, 1984, **23**, 455 (*isol*)

Bonjoch, J. *et al.*, *J.A.C.S.*, 1997, **119**, 7230-7240 (*20-Epilochneridine, synth*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1998, **63**, 9434-9439 (*synth, bibl*)

Lochnerivine**L-225**

[1359-07-5]

C₂₄H₂₈N₂O₅ 424.496

Struct. unknown. Alkaloid from the roots of *Catharanthus roseus*. Cryst. (MeOH). Mp 278-280°.

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (*isol, uv, ir*)

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1963, **52**, 407-408 (*isol*)

Lochrovicine**L-226**

[1359-08-6]

C₂₀H₂₂N₂O₂ 322.406

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Prisms (MeOH). Mp 234-238°. [α]_D²⁵ -345 (c, 1 in CHCl₃). p*K*_a 4.5 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1964, **27**, 203-213 (*isol, ir, uv*)

Lochroidine**L-227**

[1359-09-7]

C₂₂H₂₆N₂O₄ 382.458

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Prisms (EtOH). Mp 213-218°. [α]_D²⁵ -496 (c, 1 in CHCl₃). p*K*_a 5.6 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1964, **27**, 203-213 (*isol, ir, uv*)

Lochrovine**L-228**

[1359-10-0]

C₂₃H₃₀N₂O₃ 382.502

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Plates (MeOH). Mp 258-263°.

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1964, **27**, 203-213 (*isol, ir, uv*)

Locoine**L-229**


Struct. unknown. Presumed to be an alkaloid. Isol. from *Astragalus earlei* (Fabaceae).

Fraps, G.S. *et al.*, *CA*, 1936, **31**, 6807

Loesenerine**L-230**

9-Acetyl-2-(1-heptenyl)-1,5,9-triazacyclo-tridecan-4-one, 9CI

[110011-48-8]



C₁₉H₃₅N₃O₂ 337.504

The revised abs. config. of Dihydromyricoidine (1996) suggests that the other abs. configs. covered by this entry should also be reversed. Alkaloid from the leaves of *Maytenus loeseneri* (Celastraceae). Cryst. (EtOAc). Mp 117°. [α]_D²⁵ +45.4 (c, 0.52 in CHCl₃).

17,18Z-Didehydro: 17,18-Didehydroloesenerine

[114542-50-6]

C₁₉H₃₃N₃O₂ 335.489

Alkaloid from the leaves of *Maytenus loeseneri* (Celastraceae). Cryst. (Me₂CO). Mp 106°. [α]_D²² +51.3 (c, 0.52 in CHCl₃).

16,17E-Didehydro, 18ξ-hydroxy: 16,17-Didehydroloesenerin-18-ol

[114542-51-7]

C₁₉H₃₃N₃O₃ 351.488

Alkaloid from leaves of *Maytenus loeseneri* (Celastraceae). Cryst. (MeOH/Me₂CO). Mp 203-207°. [α]_D²² +176.8 (c, 0.39 in CHCl₃/EtOH, 19:1).

17,18Z-Didehydro, N-de-Ac: Myricoidine

[117611-57-1]

C₁₇H₃₁N₃O 293.451

Trace alkaloid from the whole plant of *Clerodendrum myricoides* (Verbenaceae). Oil. [α]_D²² +83 (c, 6 in MeOH).

N-De-Ac: Dihydromyricoidine

[117611-59-3]

C₁₇H₃₃N₃O 295.467

Trace alkaloid from the whole plant of *Clerodendrum myricoides* (Verbenaceae). Oil. [α]_D²² +77 (c, 5.3 in MeOH).


Dáz, M. *et al.*, *Phytochemistry*, 1987, **26**, 1847 (*isol, ir, pmr, cmr, ms, struct*)

Preiss, A. *et al.*, *Phytochemistry*, 1988, **27**, 589 (*isol, uv, ir, pmr, cmr, ms, ord, struct, derivs*)

- Bashwira, S. *et al.*, *Tetrahedron*, 1988, **44**, 4521 (isol, ir, pmr, cmr, ms, struct)
 Häusermann, U.A. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 1995 (synth, abs config, Dihydropyridine)
 Häusermann, U.A. *et al.*, *Tet. Lett.*, 1998, **39**, 257-260 (Myricoidine, synth)

Lolicine A

[201214-57-5]

C₃₈H₅₃NO₅ 603.84

Isol. from *Lolium perenne* infected with *Neotyphodium lolii* (formerly *Acremonium lolii*). Solid (as 11-propanoyl).

41-Oxo: Lolicine B

[201214-58-6]


C₃₈H₅₁NO₆ 617.824

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*. Solid (as 11-propanoyl).

Munday-Finch, S.C. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 590-598

Lolidine

[60092-32-2]


C₁₈H₂₇ClN₄O₄ 398.888

Alkaloid from *Lolium cuneatum* (Poaceae). Mp 225-226°. [α]_D²⁰ +146 (c, 2.0 in CHCl₃).

Batirov, E.Kh. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 63; *CA*, **85**, 74883s (isol, ir, ms, struct)

Lolilline

[185217-49-6]


C₃₇H₄₇NO₆ 601.781

Alkaloid from the seeds of *Lolium perenne* infected with *Acremonium lolii*. Mycotoxin.

Mundayfinch, S.C. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 199-204

Loline

Hexahydro-N-methyl-2,4-methano-4H-furo[3,2-b]pyrrol-3-amine, 9CI, 2,7-Epoxy-1-methylaminopyrrolizidine. *Festuca* [25161-91-5]

C₈H₁₄N₂O 154.211

Alkaloid from the grasses *Lolium cuneatum*, *Lolium temulentum* and *Festuca arundinacea* (Poaceae). Oil. d₄²⁰ 1.13. Bp₇₃₁ 229° Bp₅ 103°. n_D²² 1.5505. pK_a 2.5.

Hydrochloride (1:2): [25161-92-6] Needles (EtOH). Mp 256-257° dec. (237-242° dec.). [α]_D¹⁵ +6.2 (H₂O).

N-Formyl: N-Formyloline

[38964-33-9]

C₉H₁₄N₂O₂ 182.222

Alkaloid from *Lolium cuneatum* (Poaceae). Cryst. Mp 93-94°. [α]_D²⁰ +47.9 (c, 0.73 in CHCl₃).

N-Ac: Lolinine. N-Acetyloline

[4914-36-7]

C₁₀H₁₆N₂O₂ 196.249Mp 73°. [α]_D²⁰ +36.9.**N-Ac, N-oxide: Lolinine N-oxide. N-Acetyloline N-oxide**

[61391-09-1]

C₁₀H₁₆N₂O₃ 212.248

From *Lolium cuneatum* (Poaceae). Oil. [α]_D²⁰ +21.6 (c, 0.9 in MeOH).

N-Me: N-Methyloline

[22143-50-6]

C₉H₁₆N₂O 168.238

Alkaloid from *Lolium cuneatum* (Poaceae). Oil. d₄²⁰ 1.02. Bp₂ 90-91°. [α]_D³¹ +9.31 (c, 9.5 in MeOH). [α]_D²⁰ +13.7 (c, 2.44 in Me₂CO). n_D²² 1.4968.

N-Me; hydrochloride (1:2): Mp 246-247°.

N-De-Me: Norloline. N-Depropionyldecorticasine

[4839-19-4]

C₇H₁₂N₂O 140.185

Alkaloid from *Lolium cuneatum* and *Adenocarpus decorticans* (Poaceae, Fabaceae). Bp₅ 94-95°. [α]_D¹⁶ +15.1.

N-De-Me, N-formyl: N-FormylnorlolineC₈H₁₂N₂O₂ 168.195

From *Lolium cuneatum* (Poaceae). Liq. [α]_D²⁰ +31.3 (c, 0.96 in Me₂CO).

N-De-Me, N-Ac: N-Acetylnorloline

[38964-35-1]

C₉H₁₄N₂O₂ 182.222

From *Lolium cuneatum* (Poaceae). Oil. [α]_D²⁰ +49.8 (c, 2.23 in CHCl₃).

N-De-Me, N-Ac; hydrochloride (1:2): Mp 232-235° dec.

N-De-Me, N-propanoyl: Decorticasine

[1380-03-6]

C₁₀H₁₆N₂O₂ 196.249

Alkaloid from *Adenocarpus decorticans* (Fabaceae).

N-De-Me, N-butanoyl: N-Butyryl-N-depropionyldecorticasine**L-234**

Absolute Configuration

C₁₁H₁₈N₂O₂ 210.275

Alkaloid from *Adenocarpus decorticans* (Fabaceae). Mp 206° dec. (as picrate).

N-De-Me, N-(2-methylpropanoyl): N-Isobutyryl-N-depropionyldecorticasineC₁₁H₁₈N₂O₂ 210.275

Alkaloid from *Adenocarpus decorticans* (Fabaceae). Solid (EtOH). Mp 127°.

N-De-Me, N-(2-methylpropanoyl), picrate:

Cryst. (EtOH). Mp 241°.

N-De-Me, N-(3-methylbutanoyl): N-Isovaleryl-N-depropionyldecorticasineC₁₂H₂₀N₂O₂ 224.302

Alkaloid from *Adenocarpus decorticans* (Fabaceae). Cryst. (EtOH) (as picrate). Mp 241° dec. (picrate).

5,6-Didehydro, N-Ac: N-Acetyl-5,6-dehydrololine. 5,6-Dehydrololine

[194205-01-1]

C₁₀H₁₄N₂O₂ 194.233

Alkaloid from endophyte-infected *Festuca argentina*.

Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 298; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 252 (ms)

Aasen, A.J. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2021 (ir, pmr)

Soto, J.P. *et al.*, *CA*, 1970, **73**, 109943 (Decorticasine)

Robbins, J.D. *et al.*, *J. Agric. Food Chem.*, 1972, **20**, 1040 (isol, pmr, ms, derivs)

Bates, R.B. *et al.*, *Tet. Lett.*, 1972, 1629 (abs config, cryst struct, bibl)

Landa-Veloz, A. *et al.*, *An. Quim.*, 1974, **70**, 360 (isol, struct, derivs)

Batirov, E. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 60; 120 (isol, struct, derivs)

Wilson, S.R. *et al.*, *J.O.C.*, 1981, **46**, 3887 (synth)

Dannhardt, G. *et al.*, *Planta Med.*, 1985, 212 (isol)

Tufariello, J.J. *et al.*, *J.O.C.*, 1986, **51**, 3556 (synth)

Knoch, F. *et al.*, *Z. Kristallogr.*, 1993, **205**, 346 (cryst struct)

Casabuono, A.C. *et al.*, *J. Ethnopharmacol.*, 1997, **57**, 1-9 (Acetyldehydrololine)

Blakemore, P.R. *et al.*, *J.C.S. Perkin 1*, 2001, 1831-1845 (synth)


Lolididine**L-235**

Struct. unknown. Alkaloid from the seeds of *Lolium cuneatum*. Mp 215-216° (as hydrochloride). Bp₅ 85-93°.

Yunusov, S.Y. *et al.*, *Zh. Obshch. Khim.*, 1955, **25**, 1813-1820; *CA*, **50**, 7117d

Lolitrem B**L-236**

[81771-19-9]

C₄₂H₅₅NO₇ 685.899

Isol. from *Lolium perenne* infected with

Neotyphodium lolii (formerly *Acremonium lolii*). Potent neurotoxin affecting cattle. Mycotoxin. Tremorgenic agent. Mp 303-304°. Major component of Lolitrein complex. λ_{\max} 267 (ϵ 26800); 290 (ϵ 6700) (MeOH) (Berdy).

40,41-Dihydro: Lolitrein C

[109862-63-7]

$C_{42}H_{57}NO_7$ 687.915

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*. Tremorgenic neurotoxin. Tentative struct.

40,41-Epoxy: Lolitrein A

[162681-78-9, 162616-65-1]

$C_{42}H_{55}NO_8$ 701.898

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*. Amorph. solid.

Isol. as a mixt. of C-40 epimers. λ_{\max} 262 (ϵ 34000) (MeCN).

31-Epimer: 31-Epilolitrein B

$C_{42}H_{55}NO_7$ 685.899

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*.

35-Epimer: Lolitrein F

[179464-12-1]

$C_{42}H_{55}NO_7$ 685.899

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*. Mycotoxin. Tremorgenic agent. Neurotoxin. Amorph. solid. λ_{\max} 265 (MeOH) (Berdy).

31,35-Diepimer: 31-Epilolitrein F

$C_{42}H_{55}NO_7$ 685.899

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*.

Gallagher, R.T. et al., *Chem. Comm.*, 1984, 614-616 (isol. struct, pmr, cmr, props)

Gallagher, R.T. et al., *J. Chromatogr.*, 1985, **322**, 159-167 (hplc)

Weedon, C.M. et al., *Phytochemistry*, 1987, **26**, 969-971 (biosynth)

Miles, C.O. et al., *J. Agric. Food Chem.*, 1994, **42**, 1488-1492 (isol, Lolitrein B)


Munday-Finch, S.C. et al., *J. Agric. Food Chem.*, 1995, **43**, 1283-1288; 1996, **44**, 2782-2788 (Lolitrein A, Lolitrein F)

Munday-Finch, S.C. et al., *J. Agric. Food Chem.*, 1998, **46**, 590-598 (31-Epilolitrein B, 31-Epilolitrein F)

Lolitrein N

L-237

[201214-59-7]



$C_{37}H_{49}NO_7$ 619.797

Isol. from *Lolium perenne* infected with *Neotyphodium lolii* (formerly *Acremonium lolii*).

31-Epimer: 31-Epilolitrein N

[201214-60-0]

$C_{37}H_{49}NO_7$ 619.797

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*.

35-Epimer: Lolitriol. 35-Epilolitrein N

[138605-81-9]

$C_{37}H_{49}NO_7$ 619.797

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*.

35-Epimer, O²⁷-prenyl: Lolitrein E

[155964-95-7]

$C_{42}H_{57}NO_7$ 687.915

Isol. from ryegrass *Lolium perenne* infected with *Acremonium lolii*.

Amorph. solid.


Miles, C.O. et al., *J. Agric. Food Chem.*, 1994, **42**, 1488-1492 (Lolitrein E)

Munday-Finch, S.C. et al., *J. Agric. Food Chem.*, 1998, **46**, 590-598 (Lolitrein N, 31-Epilolitrein N, Lolitriol)

Lomaiviticin B

L-238

[349662-91-5]



$C_{54}H_{56}N_6O_{18}$ 1077.066


Prod. by the marine-derived *Micromonospora lomaivitiensis* (strain LL-371366). Potent antitumour agent and active against gram-positive bacteria. Amorph. red powder. $[\alpha]_D^{25}$ -71.4 (c, 0.07 in MeOH).

He, H. et al., *J.A.C.S.*, 2001, **123**, 5362-5363 (isol, pmr, cmr, ms)

Lombine

L-239

[935261-35-1]



$C_{21}H_{24}N_2O_3$ 352.432


Alkaloid from the bark of *Voacanga foetida*.

Hadi, S. et al., *Nat. Prod. Commun.*, 2006, **1**, 825-829 (isol)

Lombricine

L-240

Serine 2-[(aminoiminomethyl)amino]ethyl hydrogen phosphate (ester), 9CI. O³-(2-Guanidinoethoxyphosphinoserine). Lumbricine



(R)-form

$C_6H_{15}N_4O_6P$ 270.181

(R)-form

D-form

Constit. of *Ophelia bicornis* muscle. Mp 229-230°. $[\alpha]_D^{25}$ +14.4 (c, 0.917 in H₂O).

N^ω-Phosphonyl: **Phospholombricine**. N-Phosphoryllombricine. Lumbricine phosphate

[25540-15-2]

$C_6H_{16}N_4O_9P_2$ 350.161

Isol. from *Lumbricus terrestris* and other earthworms. High-energy phosphate regulator.

Characterised as a cryst. NH₄⁺ salt and as other salts.

(S)-form

L-form

[18416-85-8]

Constit. of the body wall muscle of the echiuroid worm *Thalassema neptuni*.

Cryst. (EtOH aq.). Mp 233°. $[\alpha]_D^{25}$ -11.3 (c, 0.380 in H₂O). pK_{a1} 2; pK_{a2} 8.9 (20°).

N², N²-Di-Me: **Thalassemine**

[40524-74-1]

$C_8H_{19}N_4O_6P$ 298.235

Constit. of viscera of *Thalassema neptuni*. Cryst. + 1H₂O (H₂O). Mp 184°. $[\alpha]_D^{26}$ -11.3 (c, 0.919 in H₂O).

N², N²-Di-Me, N^ω-phosphonyl: **Phosphothalassemine**

[39667-80-6]

$C_8H_{20}N_4O_9P_2$ 378.215

Constit. of viscera of *Thalassema neptuni*.

N²-L- α -Aspartyl: N²-Aspartyllombricine.

Bonellidine

[76848-19-6]

$C_{10}H_{20}N_5O_9P$ 385.27

Isol. from *Bonellia viridis*.

[18555-02-7]

Thoai, N.-V. et al., *Biochim. Biophys. Acta*, 1954, **14**, 76 (isol)

Pant, R. et al., *Biochem. J.*, 1959, **73**, 30 (isol)

Ennor, A.H. et al., *Biochem. J.*, 1962, **83**, 14

(Phospholombricine)

Dubey, S.S. et al., *Indian J. Chem.*, 1963, **1**, 453

(isol)

Beatty, I.M. et al., *J.C.S.*, 1965, 12 (synth,

Phospholombricine)

Thoai, N.-V. et al., *Biochemistry*, 1972, **11**,

3890-3895 (Thalassemine,

Phosphothalassemine)

Robin, Y. et al., *Oceanis*, 1980, **5**, 575-580; *CA*,

94, 118125p (Bonellidine)


Everby, M.R. et al., *J. Chromatogr.*, 1988, **445**,

433

Lomofungin, USAN

L-241

Methyl 6-formyl-4,7,9-trihydroxy-1-phenazinecarboxylate, 9CI. 5-Formyl-4,6,8-trihydroxy-1-methoxycarbonylphenazine. Lomodomycin. NSC 106995. U 24792. Antibiotic U 24792 [26786-84-5]



C₁₅H₁₀N₂O₆ 314.254


Isol. from *Streptomyces lomondensis*. Broad-spectrum antibiotic. Sol. bases, Me₂CO, DMF; poorly sol. H₂O, hexane, MeOH, EtOAc. Mp 320°. Log P 3.87 (calc). λ_{max} 219 (ε 25300); 258 (ε 26400); 271 (ε); 284 (ε 35200); 316 (ε 13200); 441 (ε 4550) (MeOH) (Derep). λ_{max} 221 (E1%/1cm 876); 270 (E1%/1cm 1605); 311 (E1%/1cm 434); 375 (E1%/1cm 300); 465 (E1%/1cm 428) (MeOH/HCl) (Berdy). λ_{max} 239 (E1%/1cm 622); 265 (E1%/1cm 1110); 299 (E1%/1cm 900); 342 (E1%/1cm 1113); 478 (E1%/1cm 151) (MeOH/NaOH) (Berdy).

▶ SG1576500

Tri-Me ether: Mp 215-217°.

Bergy, M.E. *et al.*, *J. Antibiot.*, 1969, **22**, 126 (isol, uv, ir, pmr, props)Tipton, C.D. *et al.*, *J.A.C.S.*, 1970, **92**, 1425 (struct)Haber, A. *et al.*, *CA*, 1978, **89**, 18515u (nmr)**Longamide A†****L-242**

6,7-Dibromo-3,4-dihydro-4-hydroxypyrrrolo[1,2-a]pyrazin-1(2H)-one, 9CI

C₇H₆Br₂N₂O₂ 309.945**(S)-form** [170891-48-2]

Isol. from the Caribbean sponge *Agelas longissima* and by *Homaxinella* sp. Shows moderate antibacterial activity. Sol. MeOH, butanol. [α]_D +86 (c, 0.001 in MeOH). λ_{max} 209 (ε 16200); 230 (ε 8100); 276 (ε 7700) (MeOH) (Berdy).

(±)-form [219783-00-3]Alkaloid from the sponge *Homaxinella* sp.**(ξ)-form**

6-Debromo: 6-Debromolongamide A.

Mukanadin C


[203579-34-4]

C₇H₇BrN₂O₂ 231.049

Isol. from the sponge *Axinella carteri* and *Agelas nakamurai*. Yellow powder. [α]_D -6 (c, 0.08 in MeOH). Stereochem. not determined for either isolate.

Cafieri, F. *et al.*, *Tet. Lett.*, 1995, **36**, 7893-7896 (isol, uv, ir, pmr, cmr, cd, struct)Li, C.-J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 387-389 (deriv, isol, pmr, cmr)Umeyama, A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1433-1434 (isol)Uemoto, H. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1581-1583 (*Mukanadin C*)Marchais, S. *et al.*, *Tet. Lett.*, 1999, **40**, 5519-5522 (synth)**Longamide B****L-243**

6,7-Dibromo-1,2,3,4-tetrahydro-1-oxopyrrolo[1,2-a]pyrazine-4-acetic acid, 9CI

C₉H₈Br₂N₂O₃ 351.982**(R)-form**


Me ester: [219782-99-7]

C₁₀H₁₀Br₂N₂O₃ 366.009Alkaloid from the sponge *Agelas ceylonica*. [α]_D +7 (c, 1 in MeOH).

7-Debromo, Me ester: [335022-79-2]

C₁₀H₁₁BrN₂O₃ 287.113Alkaloid from the sponge *Axinella tenuidigitata*. Solid. [α]_D²⁵ +8.2 (c, 0.5 in MeOH). λ_{max} 220 (ε 29600); 244 (ε 12100); 303 (ε 8130) (MeOH).**(±)-form** [200264-70-6]Isol. from the sponge *Agelas dispar*.Amorph. solid. λ_{max} 210 (ε 12100); 230 (ε 8500); 280 (ε 7700) (MeCN). λ_{max} 210 (ε 12100); 230 (ε 7700); 280 (ε 8500) (MeOH) (Berdy).Et ester: **Hanishin**

[196303-62-5]

C₁₁H₁₂Br₂N₂O₃ 380.035Alkaloid from the sponges *Acanthella carteri* and *Homaxinella* sp. Semisolid. λ_{max} 203 (ε 17800); 230 (ε 10000); 280 (ε 6500) (MeOH).Mancini, I. *et al.*, *Tet. Lett.*, 1997, **38**, 6271-6274 (*Hanishin*)Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 122-125 (*Hanishin*)Umeyama, A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1433-1434 (*Hanishin*)Reddy, N.S. *et al.*, *Biochem. Syst. Ecol.*, 2000, **28**, 1035-1037 (*Me ester, isol*)Reddy, N.S. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 971-972 (*7-debromo Me ester*)Patel, J. *et al.*, *J.O.C.*, 2005, **70**, 9081-9084 (synth, abs config)Trost, B.M. *et al.*, *Org. Lett.*, 2007, **9**, 2357-2359 (synth)Sun, X.-T. *et al.*, *Tet. Lett.*, 2007, **48**, 3459-3461 (synth)**Longanine****L-244**8,10-Epoxy-3,7-dimethoxy-17-methylhasubanan-4,6,8-triol, 9CI. *Longaninine* [78101-26-5]C₁₉H₂₅N₂O₆ 363.41Alkaloid from the roots and stems of *Stephania longa* (Menispermaceae). Mp 197-199°.6-O-E-Cinnamoyl: **Stephalonine D**

[863712-18-9]

C₂₈H₃₁NO₇ 493.555Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -17.9 (c, 0.28 in CHCl₃).O⁴-Me: **Stephasunoline**

[52309-77-0]

C₂₀H₂₇NO₆ 377.436Alkaloid from the stems and rhizomes of *Stephania japonica* (Menispermaceae). Cryst. (MeOH). Mp 233°. [α]_D²⁰ +121.4 (c, 1.44 in CHCl₃).O⁸-Me: **N-Methylstephuline**. O-Methylhermandine

[30660-43-6]

C₂₀H₂₇NO₆ 377.436Alkaloid from *Stephania hernandifolia* and *Stephania longa*. Mp 152-153°. [α]_D +125 (c, 0.4 in EtOH).O⁸-Me, 6-O-(2-methylbutanoyl): **Stephalonine A**

[863712-12-3]

C₂₅H₃₅NO₇ 461.554Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ +102.8 (c, 0.68 in CHCl₃). λ_{max} 225 (log ε 3.69); 285 (log ε 2.97) (MeOH).O⁸-Me, 6-benzoyl: **Stephalonine F**

[863712-22-5]

C₂₇H₃₁NO₇ 481.544Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -11.9 (c, 0.21 in CHCl₃). λ_{max} 282 (log ε 3.39) (MeOH).O⁸-Me, 6-O-(4-hydroxy-3-methoxybenzoyl): **Stephalonine H**

[863712-26-9]

C₂₈H₃₃NO₉ 527.57Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -100.7 (c, 0.15 in Py). λ_{max} 264 (log ε 3.9); 288 (log ε 3.73) (MeOH).O⁸-Me, 6-O-(3,4-dimethoxybenzoyl):**Stephalonine G**

[863712-24-7]

C₂₉H₃₅NO₉ 541.597Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -26.9 (c, 0.3 in CHCl₃). λ_{max} 262 (log ε 3.92); 287 (log ε 3.68) (MeOH).O⁸-Me, 6-O-E-cinnamoyl: **Stephalonine B**

[863712-14-5]

C₂₉H₃₃NO₇ 507.582Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ +7.7 (c, 0.77 in CHCl₃). λ_{max} 277 (log ε 4.25) (MeOH).O⁸-Me, N-de-Me: **Stephuline**

[30753-39-0]

C₁₉H₂₅NO₆ 363.41Alkaloid from *Stephania longa*. Mp 223-225°. [α]_D³² +93 (c, 0.55 in CHCl₃).O⁸-Me, N-de-Me, 6-O-(4-hydroxy-3-methoxybenzoyl): **Stephalonine I**

[863712-28-1]

C₂₇H₃₁NO₉ 513.543Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -18.3 (c, 0.11 in CHCl₃). λ_{max} 263 (log ε 3.91); 287 (log ε 3.74)

(MeOH).

*O*⁸-Me, *N*-de-Me, 6-*O*-E-cinnamoyl: **Stephalonine C**
[863712-16-7]
C₂₈H₃₁NO₇ 493.555

Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ -4.2 (c, 0.12 in MeOH). λ_{max} 277 (log ε 4.21) (MeOH).

*O*⁸-Me, *N*-de-Me, 6-*O*-(3-hydroxy-4-methoxycinnamoyl): **Stephisoferuline**.
Hernandifoline

[30511-63-8]
C₂₉H₃₃NO₉ 539.581

Alkaloid from *Stephania hernandifolia* (Menispermaceae). Cryst. (CHCl₃/Et₂O). Mp 133-135° (CHCl₃ solvate). [α]_D²⁵ +48 (c, 0.82 in MeOH). [α]_D²⁰ -25 (EtOH). Stephisoferuline and *Hernandifoline* appear to be the same but their opt. rotns. are v. different and comparison of lit. values for Mp's is confused by solvate formn.

*O*⁸-Me, *O*³-de-Me, *N*-de-Me, 6-*O*-(3-hydroxy-4-methoxycinnamoyl): ***O*³-Demethylhernandifoline**

[38940-54-4]
C₂₈H₃₁NO₉ 525.554

Alkaloid from *Stephania hernandifolia* (Menispermaceae). Mp 148-149°.

*O*⁴,*O*⁸-Di-Me, 6-*Ac*: **Dihydroepistephamiersine 6-acetate**

[57361-74-7]

C₂₃H₃₁NO₇ 433.5

Alkaloid from *Stephania abyssinica*

(Menispermaceae).

O-De-Me, Me ether (?): ***Hernandine***[†]

[32593-70-7]

C₁₉H₂₅NO₆ 363.41

Alkaloid from the aerial portion of *Stephania hernandifolia* (Menispermaceae). Mp 197-199°. [α]_D -33 (EtOH). May be identical with *Longanine* or with its *O*⁷-de-Me, *O*⁸-Me isomer. The C-7 config. was not definitely established. The reported Mp. corresponds to that of *Longanine*.

6-Ketone, *O*⁸-Me: ***Longanone***

[81525-51-1]

C₂₀H₂₅NO₆ 375.421

Alkaloid from the roots and stems of *Stephania longa* (Menispermaceae). Mp 161-163°. [α]_D +86.5 (CHCl₃).

16-*Oxo*, *O*⁴-Me: ***Oxostephasunoline***

[91897-38-0]

C₂₀H₂₅NO₇ 391.42

Alkaloid from *Stephania japonica* roots (Menispermaceae). Prisms (Me₂CO). Mp 217°. [α]_D¹⁴ +199.36 (c, 0.84 in MeOH).

16-*Oxo*, *O*⁸-Me, 6-*O*-E-cinnamoyl: **Stephalonine E**

[863712-20-3]

C₂₉H₃₁NO₈ 521.566

Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ +49.6 (c, 0.11 in CHCl₃). λ_{max} 279 (log ε 4.25) (MeOH).

1-Nitro, *O*⁸-Me, 6-*O*-(2*S*-methylbutanoyl): **Stephalonine J**

[908568-26-3]

C₂₅H₃₄N₂O₉ 506.552

Alkaloid from *Stephania longa*. Yellow powder. [α]_D²¹ +181.2 (c, 0.16 in CHCl₃). λ_{max} 285 (log ε 3.76) (MeOH).

1-Nitro, *O*⁸-Me, 6-*O*-E-cinnamoyl: **Stephalonine K**

[908568-28-5]

C₂₉H₃₂N₂O₉ 552.58

Alkaloid from *Stephania longa*. Yellow powder. [α]_D²¹ +354.6 (c, 0.13 in CHCl₃). λ_{max} 277 (log ε 4.22) (MeOH).

7-*Epimer*, 6-ketone, *O*⁸-Me: ***Isolonganone***

[863712-34-9]

C₂₀H₂₅NO₆ 375.421

Alkaloid from *Stephania longa*. Powder. [α]_D²⁰ +57.5 (c, 0.57 in CHCl₃). λ_{max} 284 (log ε 3.31) (MeOH).

Kupchan, S.M. *et al.*, *Tet. Lett.*, 1970, 4975 (*Stephuline*)

Fedenko, D.A. *et al.*, *Khim. Priro. Soedin.*, 1971, 7, 158; 180; 455; 1972, 8, 130; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, 7, 150; 171; 432; 1972, 8, 136 (*Hernandine*, *O*-Methylhernandine, *Hernandifoline*, *O*-Demethylhernandifoline)

Matsui, M. *et al.*, *Chem. Pharm. Bull.*, 1975, 23, 1323 (*Longanine*, *Stephasunoline*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Van Wyk, A.T. *et al.*, *J. S. Afr. Chem. Inst.*, 1975, 28, 284 (*Dihydroepistephamiersine 6-acetate*)

Lao, A. *et al.*, *Yaouxue Xuebao*, 1980, 15, 696; 1981, 16, 940; *C.A.*, 95, 49252j; 96, 196523x (*Longanone*)

Matsui, M. *et al.*, *J. Nat. Prod.*, 1984, 47, 465 (*Oxostephasunoline*)

Zhang, H. *et al.*, *J. Nat. Prod.*, 2005, 68, 1201-1207 (*cmr*, *Stephalonines*, *Stephuline*, *Stephisoferuline*, *Isolonganone*)


Zhang, H. *et al.*, *Chin. J. Chem.*, 2006, 24, 781-784 (*Stephalonines J-K*)

Longeracinylline A

L-245

Daphniyunnine C

[881388-89-2]



C₂₁H₂₇NO₂ 325.45

Alkaloid from the leaves of *Daphniphyllum longeracemosum* and from *Daphniphyllum yunnanense*. Prisms (MeOH) or amorph. powder. Mp 160-162°. [α]_D²⁵ -246.2 (c, 0.57 in CHCl₃). [α]_D²⁰ -147.1 (c, 0.14 in CHCl₃). λ_{max} 226; 242 (CHCl₃). λ_{max} 209 (log ε 4.24); 241 (log ε 4.02) (MeOH).

Zhang, H. *et al.*, *J. Nat. Prod.*, 2006, 69, 553-557 (*Daphniyunnine C*)


Di, Y.-T. *et al.*, *J. Nat. Prod.*, 2006, 69, 1074-1076 (*Logeracinylline A*, *cryst struct*)

Longeracinylline B

L-246

Daphniyunnine B

[881388-88-1]



C₂₁H₂₉NO₃ 343.465

Alkaloid from the leaves of *Daphniphyllum longeracemosum* and from *Daphniphyllum yunnanense*. Amorph. powder. [α]_D²⁵ +89.5 (c, 0.53 in CHCl₃). [α]_D²⁰ +67 (c, 0.1 in CHCl₃). λ_{max} 200; 247 (CHCl₃). λ_{max} 246 (log ε 3.75) (MeOH).


Zhang, H. *et al.*, *J. Nat. Prod.*, 2006, 69, 553-557 (*Daphniyunnine B*)

Di, Y.-T. *et al.*, *J. Nat. Prod.*, 2006, 69, 1074-1076 (*Longiracinylline B*)

Longetherine

L-247

[151271-86-2]




C₂₁H₂₉NO₄ 359.464

Alkaloid from *Stephania longa* (Menispermaceae). Needles (CHCl₃/petrol). Mp 184-185°. [α]_D²² -258 (c, 0.02 in MeOH).

Deng, J.-Z. *et al.*, *Phytochemistry*, 1993, 33, 941 (*isol*, *ir*, *pmr*, *cmr*, *cd*, *ms*, *struct*)

Longiberine

L-248



C₃₈H₄₀N₂O₇ 636.743

The first natural protoberberine-benzylisoquinoline dimer. Alkaloid from the roots of *Thalictrum longistylum*. Cryst. (MeOH/Et₂O). Mp 169-170°. [α]_D²⁵ +43.8 (c, 0.56 in MeOH). λ_{max} 221 (end) (log ε 4.77); 240 (sh) (log ε 4.42); 283 (log ε 4.14) (MeOH).

Me ether: ***O*-Methylongiberine**


C₃₉H₄₂N₂O₇ 650.77

Alkaloid from the roots of *Thalictrum longistylum*. Amorph. [α]_D²⁴ +31 (c, 0.21 in MeOH). λ_{max} 220 (end) (log ε 4.82); 242 (sh) (log ε 4.42); 282 (log ε 4.01) (MeOH).

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1999, 62, 1410-1414

Longicaudatine L-249

[85335-06-4]



Absolute Configuration

 $C_{38}H_{40}N_4O$ 568.76Alkaloid from *Strychnos longicaudata* and a number of other *Strychnos* spp. (Loganiaceae). Mp 350° dec. $[\alpha]_D +141$ (c, 0.5 in $CHCl_3$).**N^d-Oxide: Longicaudatine N-oxide**


[91377-16-1]

 $C_{38}H_{40}N_4O_2$ 584.76Alkaloid from stem bark of *Strychnos chrysophylla* and root bark of *Strychnos matopensis*.**19',20'β-Dihydro: Dihyrolongicaudatine**

[144525-18-8]

 $C_{38}H_{42}N_4O$ 570.776Alkaloid from root bark of *Strychnos potatorum* (Loganiaceae). $[\alpha]_D +83$ (c, 0.25 in MeOH).Massiot, G. *et al.*, *Tetrahedron*, 1983, **39**, 3645 (isol, uv, ir, pmr, cmr, ms, struct)Verpoorte, R. *et al.*, *J. Ethnopharmacol.*, 1984, **10**, 243 (oxide)Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304; 1992, **31**, 2873 (cmr, Dihyrolongicaudatine)Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (occur, *Strychnos*)**Longicaudatine Z** L-250

[119308-19-9]



Absolute Configuration

 $C_{38}H_{44}N_4O_2$ 588.791Alkaloid from the root bark of *Strychnos matopensis* (Loganiaceae). $[\alpha]_D -13$ (c, 0.16 in $CHCl_3$).**17'-Deoxy, 16',17'-didehydro: Longicaudatine Y**


[88733-18-0]

 $C_{38}H_{42}N_4O$ 570.776Alkaloid from the root bark and stem bark of *Strychnos longicaudata* and from *Strychnos matopensis* (Logania-ceae). Amorph. $[\alpha]_D +328$ (c, 0.4 in MeOH). Stereochem. not explicitly assigned but appears homologous with Longicaudatine Z.**17'-Deoxy, 16',17'-didehydro, 18'-hydroxy: Longicaudatine F**

[88733-17-9]

 $C_{38}H_{42}N_4O_2$ 586.775Isol. from the root bark and stem bark of *Strychnos longicaudata* and from *Strychnos matopensis* and *Strychnos minfiensis* (Loganiaceae). Amorph. $[\alpha]_D +268$ (c, 0.6 in MeOH). Stereochem. not explicitly fully assigned but appears homologous with Longicaudatine Z.**17'-Deoxy, 16',17'-didehydro, 19,20β-dihydro: Dihyrolongicaudatine Y**


[144548-38-9]

 $C_{38}H_{44}N_4O$ 572.792Alkaloid from root bark of *Strychnos potatorum* (Loganiaceae). $[\alpha]_D +262$ (c, 0.1 in MeOH).**17'-Deoxy, 3',4',5',6',16',17'-hexadecahydro: 3',4',5',6'-Tetradehydrolongicaudatine Y** $C_{38}H_{39}N_4O^{\oplus}$ 567.752Quaternary alkaloid from *Strychnos usambarensis*.Massiot, G. *et al.*, *Tetrahedron*, 1983, **39**, 3645-3656 (Longicaudatine Y)Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304; 1992, **31**, 2873-2876 (Longicaudatine Z, Dihyrolongicaudatine Y)Frederich, M. *et al.*, *Phytochemistry*, 1998, **48**, 1263-1266 (Tetradehydrolongicaudatine Y)**Longifolidine** L-251**1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-7,8-dimethoxy-1-methylisoquinoline**

(R)-form

 $C_{19}H_{23}NO_3$ 313.396**(R)-form** [77480-17-2]Alkaloid from *Cryptocarya longifolia* (Lauraceae). Pale yellow amorph. solid. $[\alpha]_D^{20} -72$ (c, 0.2 in EtOH).**N-Me: 8-O-Methylolongine**


[152230-58-5]

 $C_{20}H_{26}NO_3^{\oplus}$ 328.43Alkaloid from stems of *Litsea cubeba* (mountain pepper). Amorph. solid (as perchlorate). Mp 102° (perchlorate). $[\alpha]_D^{24} -12$ (c, 1.0 in MeOH).**(±)-form**Synthetic. Cryst. (Et₂O). Mp 166-168°.Grethe, G. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 874 (synth, uv, ir, pmr, ms)Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1981, **34**, 195 (isol, uv, ir, pmr, ms, struct)Lee, S.S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1971 (8-O-Methylolongine)**Longifolonine** L-252**3,4-Dihydro-7-hydroxy-1-(4-hydroxybenzoyl)-6-methoxyisoquinoline** [77410-37-8] $C_{17}H_{15}NO_4$ 297.31Alkaloid from *Cryptocarya longifolia*.Small prisms + 1CHCl₃ (MeOH/CHCl₃). Mp 143-145°.**4'-Me ether: Velucryptine**


[123825-69-4]

 $C_{18}H_{17}NO_4$ 311.337Alkaloid from leaves of *Cryptocarya velutinos*.**7-Me ether: Nelumstemine** $C_{18}H_{17}NO_4$ 311.337Alkaloid from the stems of *Nelumbo nucifera* (East Indian lotus). Orange cryst. (MeOH). Mp 272-273°.Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1981, **34**, 195-207 (Longifolonine, cryst struct)Leboeuf, M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 516-521 (Velucryptine)Duan, X.H. *et al.*, *Chin. Chem. Lett.*, 2008, **19**, 308-310 (Nelumstemine)**Longistrobine** L-253

[23544-51-6]

 $C_{17}H_{19}N_3O_3$ 313.355Alkaloid from the aerial parts of *Macrorungia longistrobus* (preferred genus name *Anisotes*) (Acanthaceae). Needles (Me₂CO/hexane). Mp 145-148°. λ_{max} 257 (ε 11300) (EtOH). λ_{max} 235 (ε 14900) (1% HCl/EtOH).Arndt, R.R. *et al.*, *Tetrahedron*, 1969, **25**, 2767-2799 (isol, ir, uv, pmr, ms)Wuonola, M.A. *et al.*, *Tetrahedron*, 1976, **32**, 1085-1095 (struct)**Longistylumphylline B** L-254

[857672-35-6]




Relative Configuration

 $C_{23}H_{33}NO_3$ 371.519

Alkaloid from *Daphniphyllum longistylum*. Powder. $[\alpha]_D^{20}$ -10 (c, 0.5 in MeOH/CHCl₃).

Chen, X. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 854-860 (*isol*, *pmr*, *cmr*)

Longistylumphylline C L-255
[857672-36-7]




Relative Configuration

C₂₃H₃₅NO₃ 373.534
Alkaloid from *Daphniphyllum longistylum*. Oil. $[\alpha]_D^{20}$ +19 (c, 1 in CHCl₃).

Chen, X. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 854-860 (*isol*, *pmr*, *cmr*, *ms*)

Longitubanine A L-256
N-(2,5-Dihydro-3-methyl-2-furanyl)-4-hydroxy-L-glutamine, 9CI
[155416-39-0]



C₁₀H₁₆N₂O₅ 244.247
Constit. of the fresh bulbs of *Hemerocallis longituba*. Needles. Mp 178-180°. $[\alpha]_D$ -8.7 (c, 2.5 in H₂O). Similar to Oxypinnatanine.

4-Deoxy: Longitubanine B
[155416-38-9]

C₁₀H₁₆N₂O₄ 228.247

Constit. of the fresh bulbs of *Hemerocallis longituba*. Needles. Mp 181-183°. $[\alpha]_D$ -10.5 (c, 1.6 in H₂O).

Yoshikawa, K. *et al.*, *Phytochemistry*, 1994, **35**, 1057-1058 (*isol*, *pmr*, *cmr*)

Lophanterine L-257

Struct. unknown. MW 352. Alkaloid from the leaves of *Lophanthera lactescens* (Malpighiaceae). Shows febrifugal props. Needles (EtOH aq.). Mp 175-177°. $[\alpha]_D^{24}$ -92.7 (EtOH). Extremely bitter taste.

Hydrochloride:

Glassy prisms (EtOH aq.). Mp 188-189°.

Sulfate:

Prisms. Mp 194-195°.

Ribeiro, O. *et al.*, *An. Assoc. Quim. Bras.*, 1946, **5**, 39-42; *CA*, **41**, 3109

Lophilacrine L-258

C₁₄H₂₅NO₂ 239.357

Struct. unknown. Approx. MF may be H₂₇. Alkaloid from *Lobelia syphilitica* (Campanulaceae). Bp_{0.001} 117-118°. $[\alpha]_D$ -105 (EtOH).

Hydrochloride:

Needles. Mp 210-211°.

Nitrate:

Needles. Mp 83-84°.

Steinogger, E. *et al.*, *Pharm. Acta Helv.*, 1952, **27**, 113-120; 207-211; 1955, **32**, 205-206; *CA*, **47**, 6954; 12753; **51**, 14202a (*isol*)

Tschesche, R. *et al.*, *Chem. Ber.*, 1961, **94**, 3227-3336 (*Lobelia syphilitica constits*)

Tschesche, R. *et al.*, *Tetrahedron*, 1964, **20**, 2885-2893 (*Lobelia syphilitica constits*)

Lophiline L-259

C₂₈H₃₆N₂O₂ 432.605


Struct. unknown. Approx. MF, may be H₃₈ and/or O₃. No further reports to 2007. Later more thorough investigations on this sp. are reported. Alkaloid from *Lobelia syphilitica* (Campanulaceae). Mp 210-211° (as dihydrochloride).

Steinogger, E. *et al.*, *Pharm. Acta Helv.*, 1952, **27**, 113-120; 207-211; 1955, **32**, 205-206 (*isol*)

Lophocerine L-260

1,2,3,4-Tetrahydro-6-methoxy-2-methyl-1-(2-methylpropyl)-7-isoquinolinol, 9CI

1,2,3,4-Tetrahydro-7-hydroxy-1-isobutyl-6-methoxy-2-methylisoquinoline
[19485-63-3]



C₁₅H₂₃NO₂ 249.352

(ξ)-form

Alkaloid from *Lophocereus schottii* (Cactaceae). Oil. Bp_{0.05} 150-225°.

Oxalate:

Prisms (EtOH). Mp 213-214° dec.

Picrate:

Yellow cryst. (EtOH). Mp 194-195°.

Me ether: Bp_{0.5} 137-139°.

Djerassi, C. *et al.*, *Tetrahedron*, 1958, **2**, 58 (*isol*)


Bobbitt, J.M. *et al.*, *J.O.C.*, 1959, **24**, 1106 (*synth*)

Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1963, **11**, 1477 (*synth*)

O'Donovan, D.G. *et al.*, *J.C.S. Perkin 1*, 1974, 2528 (*biosynth*)

Lophocine L-261

1,2,3,6a,7,8,9,12a-Octahydro-5,11-dimethoxy-1,7-dimethyl-6a,12a-bis(2-methylpropyl)-6,12-dioxo-1,7-diazadibenzo[def,mno]chryseno, 9CI
[74991-76-7]



Relative configuration


C₃₀H₄₀N₂O₄ 492.657

Alkaloid from *Lophocereus schottii* (Cactaceae). Monoclinic cryst. Mp 194-196°. Racemic. Prob. artifact.

Wani, M.C. *et al.*, *J. Chem. Res., Miniprint*, 1980, 0301 (*isol*, *uv*, *ir*, *pmr*, *cryst struct*)

Lorneamide A L-262

4-[2-(3-Hydroxy-1-hexenyl)-4-methylphenyl]-3-butenamide
[313951-44-9]



C₁₇H₂₃NO₂ 273.374

Alkaloid from a marine actinomycete (MST-MA190). Glass. $[\alpha]_D^{19}$ -7.2 (c, 0.2 in MeOH). λ_{max} 239 (log ε 4.09); 263 (log ε 3.96) (MeOH).

1',2'-Dihydro, 3'-ketone: 4-[4-Methyl-2-(3-oxohexyl)phenyl]-3-butenamide.

Lorneamide B

[313951-45-0]


C₁₇H₂₃NO₂ 273.374

Alkaloid from a marine actinomycete (MST-MA190). Glass. λ_{max} 209 (log ε 4); 253 (log ε 3.8) (MeOH).

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1682-1683 (*Lorneamides A,B*)

Loroquine L-263

2,3-Dihydro-7-(hydroxymethyl)-1H-pyrrolizin-1-one, 9CI
[27792-82-1]



C₈H₉NO₂ 151.165

Constit. of the roots of *Urechites karwinsky* (preferred genus name *Pentalinon*) (Apocynaceae). Cryst. (cyclohexane). Mp 77-78°.

O-β-D-Glucopyranoside: **Loroquine glucoside**

C₁₄H₁₉NO₇ 313.307

Alkaloid from the roots of *Ligularia cymbulifera*. Amorph. solid. $[\alpha]_D^{21}$ -20 (c, 1.3 in MeOH). λ_{max} 289 (log ε 1.3) (MeOH).

Deoxy: 2,3-Dihydro-7-methyl-1H-pyrrolizin-1-one. **Danaidone**
[6064-85-3]

C₈H₉NO 135.165

Pheromone from hair-pencil and wing sex gland secretions of butterflies in the genera *Lycorea* and *Danaus* where it arises by metab. of pyrroli-

zidine alkaloids from plants. Mp 74-75°.

Meinwald, J. *et al.*, *J.A.C.S.*, 1966, **88**, 1305-1310 (*Danaidone*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)

Borges del Castillo, J. *et al.*, *Tet. Lett.*, 1970, **11**, 1219-1220 (*Loroquine*, *isol*, *uv*, *ir*, *pmr*, *struct*)

Edgar, J.A. *et al.*, *Experientia*, 1971, **27**, 761 (*Danaidone*, *isol*)

Petty, R.L. *et al.*, *Experientia*, 1977, **33**, 1324 (*Danaidone*, *isol*)

Komai, H. *et al.*, *Agric. Biol. Chem.*, 1983, **47**, 157 (*Danaidone*)


Pereira, A.L. *et al.*, *Quim. Nova*, 1983, **6**, 74 (*Danaidone*, *synth*)

Röder, E. *et al.*, *Amalen*, 1993, 711-713 (*Loroquine*, *synth*, *ir*, *pmr*, *cmr*, *ms*)

Liu, C.-M. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 308-316 (*glucoside*)

Losbanine**L-264**

6,7-Seco-6-norangustilobine B
[126618-33-5]



$C_{19}H_{22}N_2O_3$ 326.394

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Prisms (MeOH). Mp 108-110°. $[\alpha]_D^{27} +191.9$ (c, 1.41 in MeOH).

N^b-Me: 6,7-Secoangustilobine B

[112464-23-0]

$C_{20}H_{24}N_2O_3$ 340.421

Alkaloid from *Alstonia angustiloba*, *Alstonia congensis* and *Alstonia scholaris* (Apocynaceae). Solid. $[\alpha]_D^{27} +232.9$ (c, 1.38 in MeOH).

19 α ,20 α -Epoxide: Angustilocine

[740811-98-7]

$C_{19}H_{22}N_2O_4$ 342.394

Alkaloid from the leaves of *Alstonia angustiloba*. Light yellow oil. $[\alpha]_D -546$ (c, 0.5 in $CHCl_3$). λ_{max} 216 (log ϵ 4.36); 273 (log ϵ 3.79); 282 (log ϵ 3.77); 291 (log ϵ 3.62) (EtOH).

19 α ,20 α -Epoxide, *N*^b-Me: 6,7-Seco-19,20-epoxyangustilobine

[112448-51-8]

$C_{20}H_{24}N_2O_4$ 356.421

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Amorph. solid. $[\alpha]_D^{29} +73.6$ (c, 1.25 in MeOH).

Zeches, M. *et al.*, *J. Nat. Prod.*, 1987, **50**, 714-720 (6,7-Secoangustilobine B)


Caron, C. *et al.*, *Phytochemistry*, 1989, **28**, 1241-1244 (6,7-Secoangustilobine B)

Yamauchi, T. *et al.*, *Phytochemistry*, 1990, **29**, 3221-3225; 3547-3352 (*Alstonia scholaris* *constits*)

Kam, T.-S. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 366-369 (*Angustilocine*)

Lotthanongine**L-265**

Afzelechin-(4 β →2'')-N-(*p*-coumaroyl)-6''-hydroxytryptamine
[444588-55-0]




$C_{34}H_{30}N_2O_8$ 594.62

Flavanoid-indole alkaloid. Alkaloid from the roots of *Trigonostemon reidioides*. Amorph. powder. $[\alpha]_D^{15} +106$ (c, 0.8 in MeOH).

Kanchanapoom, T. *et al.*, *Tet. Lett.*, 2002, **43**, 2941-2943 (*isol*, *pmr*, *cmr*)

Lotusanine B**L-266**

[164366-87-4]




$C_{37}H_{40}N_4O_5$ 620.747

Stereochem. uncertain. CAS assigns the abs. stereochem. shown, though there is no published evidence for this. Stated by the authors to be racemic, but with no evidence cited. Alkaloid from aerial parts of *Zizyphus lotus* (Rhamnaceae). Amorph. solid. λ_{max} 280 (log ϵ 3.12) (MeOH).

Abu-Zarga, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 504-511 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Lotusine A**L-267**

[149377-30-0]



$C_{30}H_{38}N_4O_4$ 518.655

Alkaloid from roots of *Paliurus ramosissimus* and root bark of *Zizyphus lotus* (Rhamnaceae). $[\alpha]_D -215$ (c, 1.0 in $CHCl_3$). λ_{max} 208 ; 250 (sh) (MeOH).

N-De-Me: Lotusine D

[149355-66-8]

$C_{29}H_{36}N_4O_4$ 504.628

Alkaloid from roots of *Paliurus ramosissimus* and root bark of *Zizyphus lotus* (Rhamnaceae). $[\alpha]_D -187$ (c, 0.5 in $CHCl_3$).

13'-Hydroxy: Ramosine C

[530092-41-2]

$C_{30}H_{38}N_4O_5$ 534.654


Constit. of the roots of *Paliurus ramosissimus*. $[\alpha]_D^{26} -39$ (c, 1 in MeOH). λ_{max} 218 (log ϵ 4.75); 285 (log ϵ 3.98) (MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1993, **32**, 1591-1594 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (*Ramosine C*)

Lotusine B**L-268**

[163136-12-7]




$C_{36}H_{49}N_5O_5$ 631.814

Alkaloid from root bark of *Zizyphus lotus* (Rhamnaceae). $[\alpha]_D -179$ (c, 0.32 in $CHCl_3$). λ_{max} 207 ; 245 (sh) (MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1995, **38**, 767-772 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Lotusine C**L-269**

[163136-14-9]



$C_{35}H_{47}N_5O_5$ 617.787

Alkaloid from root bark of *Zizyphus*


lotus (Rhamnaceae). $[\alpha]_D$ -168 (c, 0.5 in CHCl_3). λ_{max} 204 (MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1995, **38**, 767 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lotusine E

L-270

[163136-13-8]



$\text{C}_{36}\text{H}_{49}\text{N}_5\text{O}_6$ 647.813


Alkaloid from root bark of *Zizyphus lotus* (Rhamnaceae). $[\alpha]_D$ -106 (c, 1.0 in CHCl_3). λ_{max} 206 ; 268 ; 323 (MeOH).

Ghedira, K. *et al.*, *Phytochemistry*, 1995, **38**, 767-772 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lotusine G

L-271

[446035-33-2]



$\text{C}_{24}\text{H}_{34}\text{N}_4\text{O}_4$ 442.557


Alkaloid from the root bark of *Zizyphus lotus*. $[\alpha]_D$ -142.7 (c, 0.5 in CHCl_3). λ_{max} 229 (MeOH).

Le Crouéour, G. *et al.*, *Fitoterapia*, 2002, **73**, 63-68 (*isol, pmr, cmr, ms*)

Louisianin C

L-272

6,7-Dihydro-4-(2-propenyl)-5H-cyclopenta[c]pyridin-5-one, 9CI. 4-Allyl-6,7-dihydro-5H-2-pyridin-5-one [171784-05-7]



$\text{C}_{11}\text{H}_{11}\text{NO}$ 173.214

Prod. by *Streptomyces* sp. WK-4028. Antiandrogen. Angiogenesis inhibitor. Antihormone agent. Inhibits growth of SC 115 cells. Dark brown oil. Sol.

MeOH, EtOAc; poorly sol. H_2O . λ_{max} 205 (ϵ 9600); 231 (ϵ 3720); 296 (ϵ 2140) (MeOH) (Berdy).

***A'*-Isomer(E)-: Louisianin D**

[171784-06-8]

$\text{C}_{11}\text{H}_{11}\text{NO}$ 173.214

Prod. by *Streptomyces* sp. WK-4028. Antiandrogen. Angiogenesis inhibitor. Antihormone agent. Inhibits growth of SC 115 cells. Plates. Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 165-169°. λ_{max} 203 (ϵ 5500); 219 (ϵ 4400); 240 (ϵ 3100); 262 (ϵ 2880); 267 (ϵ 2680); 324 (ϵ 2050) (MeOH) (Berdy).

***1'*- ξ -Hydroxy, 2',3'-dihydro: 6,7-Dihydro-4-(1-hydroxypropyl)-5H-2-pyridin-5-one. Uluypyrinone**

$\text{C}_{11}\text{H}_{13}\text{NO}_2$ 191.229

Prod. by *Streptomyces spina*. Yellow oil. Photolabile. λ_{max} 204 (log ϵ 3.71); 230 (sh) ; 294 (log ϵ 3.12) (MeOH).

1-Hydroxy: Louisianin A

[171784-03-5]

$\text{C}_{11}\text{H}_{11}\text{NO}_2$ 189.213

Prod. by *Streptomyces* sp. WK-4028. Antiandrogen. Angiogenesis inhibitor. Antihormone agent. Inhibits growth of SC 115 cells. Needles. Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 189-191°. λ_{max} 222 (ϵ 17000); 346 (ϵ 5500) (MeOH) (Berdy).

1-Hydroxy, 5 ξ -alcohol: Louisianin B

[171784-04-6]

$\text{C}_{11}\text{H}_{13}\text{NO}_2$ 191.229

Prod. by *Streptomyces* sp. WK-4028. Antiandrogen. Angiogenesis inhibitor. Antihormone agent. Inhibits growth of SC 115 cells. Plates. Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 168-170°. λ_{max} 205 (ϵ 11450); 235 (ϵ 4300); 301 (ϵ 3470) (MeOH) (Berdy).

Komiyama, K. *et al.*, *J. Antibiot.*, 1995, **48**, 1086-1089; 1090-1094 (*Louisianins A-D, isol, pmr, cmr, props*)

Henne, P. *et al.*, *Liebigs Ann./Recl.*, 1997, 937-939 (*Uluypyrinone*)

Beierle, J.M. *et al.*, *J.O.C.*, 2003, **68**, 4970-4972 (*synth*)


Chang, C.-Y. *et al.*, *J.O.C.*, 2006, **71**, 6302-6304 (*Louisianin A, synth*)

Chen, H.-W. *et al.*, *Org. Lett.*, 2006, **8**, 3033-3035 (*synth*)

Catozzi, N. *et al.*, *Tet. Lett.*, 2008, **49**, 2865-2868 (*synth*)

Louludinium(1+)

L-273



$\text{C}_{19}\text{H}_{30}\text{N}^{\oplus}$ 272.453

Isol. from the blue-green alga *Lyngbya gracilis*. Cryst. (as chloride). $[\alpha]_D^{25}$ +97 (c, 0.6 in MeOH). Mp. >300° dec. (chloride). λ_{max} 204 (ϵ 1700); 218 (ϵ 1200); 278 (ϵ 800) (MeOH).

Yoshida, W.Y. *et al.*, *Heterocycles*, 1998, **47**, 1023-1027 (*isol, uv, ir, pmr, cmr*)

Loveraine

L-274

$\text{C}_{31}\text{H}_{47}\text{NO}_{11}$ 609.712


Steroidal ester alkaloid. Struct. unknown. Alkaloid from roots of *Veratrum lobelianum* (Liliaceae). Mp 181-184°. $[\alpha]_D^{20}$ -17.3 (c, 0.46 in CHCl_3).

Starostenko, A.G. *et al.*, *Biol. Nauki (Moscow)*, 1969, 89-91; *CA*, **71**, 109768y

Lucentamycin B

L-275

[949009-87-4]



Absolute Configuration


$\text{C}_{31}\text{H}_{45}\text{N}_7\text{O}_5$ 595.74

Prod. by *Nocardiopsis lucentensis* (strain CNR-712). Cytotoxic. Yellow oil. $[\alpha]_D^{25}$ -15 (c, 0.1 in MeOH). λ_{max} 206 (log ϵ 4.1); 209 (log ϵ 3.9); 282 (log ϵ 3) (MeOH).

Cho, J.Y. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1321-1328 (*isol, pmr, cmr*)

Lucidamine A

L-276



$\text{C}_{23}\text{H}_{23}\text{NO}_5$ 393.438


(S)-form [956012-27-4]

Alkaloid from the stem bark of *Garcinia lucida*. Yellowish powder (hexane/EtOAc). Mp 212°. $[\alpha]_D^{25}$ -133 (c, 0.2 in DMSO). Lucidamine B also obt. by semisynthesis from Lucidamine A. λ_{max} 286 (log ϵ 2.22); 326 (log ϵ 0.71) (DMSO).

Fotie, J. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1650-1653 (*isol, pmr, cmr, ms*)

Lucidine B

Serratanine A
[71384-23-1]



$C_{30}H_{49}N_3O$ 467.737

The previously isolated Serratanine was a mixt. of Lucidine B and Oxolucidine B. Alkaloid from *Lycopodium lucidulum* and *Lycopodium serratum* (Lycopodiaceae). Noncryst.

14 β -Hydroxy: Oxolucidine B. Serratanine B

[71384-25-3]
 $C_{30}H_{49}N_3O_2$ 483.736

Alkaloid from *Lycopodium lucidulum* and *Lycopodium serratum*. Also obt. by air oxidn. of Lucidine B (Lycopodiaceae). Cryst. (Me₂CO/EtOAc) (as perchlorate). Mp 259-262° dec. (perchlorate). $[\alpha]_D^{21}$ -30.7 (c, 0.55 in CHCl₃).

14,15,16,17-Tetrahydro: Dihydrolycolucine

[71384-29-7]
 $C_{30}H_{45}N_3O$ 463.705

Alkaloid from *Lycopodium lucidulum* (Lycopodiaceae).

10,11,14,15,16,17-Hexahydro: Lycolucine

[71384-28-6]
 $C_{30}H_{43}N_3O$ 461.689

Alkaloid from *Lycopodium lucidulum* (Lycopodiaceae). Mp 198-200°.

6',17-Diepimer: Lucidine A

[72254-08-1]
 $C_{30}H_{49}N_3O$ 467.737

Alkaloid from *Lycopodium lucidulum*. C-14 config. not determined.

6',17-Diepimer, 14 β -hydroxy: Oxolucidine A

$C_{30}H_{49}N_3O_2$ 483.736
Alkaloid from *Lycopodium lucidulum*.
 $[\alpha]_D^{21}$ -29.1 (c, 0.74 in CHCl₃).


Ayer, W.A. *et al.*, *Can. J. Chem.*, 1979, **57**, 1105; 1984, **62**, 298 (*isol, cryst struct, stereochem*)

Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1980, **100**, 672 (*isol, ir, pmr, ms, struct*)

Tori, M. *et al.*, *Phytochemistry*, 2000, **53**, 503-509 (*isol, pmr, cmr, cryst struct*)

Lucidinine

[95014-14-5]



L-277

$C_{19}H_{25}NO_3$ 315.411

Config. at C-6 revised in 1988. Trace alkaloid from the leaves of *Phelline lucida* (Phellinaceae). $[\alpha]_D^{+73}$ (c, 1.03 in CHCl₃).

Me ether: 6,7-Dihydrohomoerysotrine.

Comosidine. Alkaloid 2

[31689-98-2]
 $C_{20}H_{27}NO_3$ 329.438

Alkaloid from the leaves of *Phelline comosa* and *Phelline lucida* (Phellinaceae). Amorph. Mp 143-145° (as picrate). $[\alpha]_D^{+72}$ (c, 0.7 in CHCl₃).

Me ether, O¹⁶-de-Me: Isolucidinine

[128508-55-4]
 $C_{19}H_{25}NO_3$ 315.411

Trace alkaloid from leaves of *Phelline comosa* var. *robusta* (Phellinaceae). Tentative, struct., not obt. pure.

1,2-Dihydro, Me ether: 1,2-Dihydrocomosidine

[99664-95-6]
 $C_{20}H_{29}NO_3$ 331.454

Trace alkaloid from the leaves of *Phelline lucida* (Phellinaceae).

Langlois, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 3535 (*isol, uv, pmr, ir, ms, struct, deriv*)


Langlois, N. *et al.*, *Heterocycles*, 1984, **22**, 2453; 1990, **30**, 659 (*uv, cd, ir, pmr, ms, Isolucidinine*)

Razafimbelo, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1985, **301**, 519 (*1,2-Dihydrocomosidine*)

Langlois, N. *et al.*, *J. Nat. Prod.*, 1988, **51**, 499 (*config*)

Lucidioline

[22594-91-8]



Absolute Configuration

L-279

$C_{16}H_{25}NO_2$ 263.379

Alkaloid from *Lycopodium lucidulum*, *Lycopodium clavatum* var. *megastachyon*, *Lycopodium gnidioides* and *Lycopodium ophioglossoides* (Lycopodiaceae). Mp 230-233°.

Perchlorate:

Cryst. (MeOH). Mp 238-240° dec.

5-Ketone: Lycoposerramine K

[67723-12-5]
 $C_{16}H_{23}NO_2$ 261.363

Alkaloid from *Lycopodium serratum*. Amorph. powder.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1962, **40**, 2088-2100; 1969, **47**, 449-455; 499-502 (*isol, ir, pmr, ms, struct*)

Nyembo, L. *et al.*, *Bull. Soc. Chim. Belg.*, 1976, **85**, 595-604; *CA*, **86**, 90108u (*isol*)

Zhou, B.-N. *et al.*, *Phytochemistry*, 1993, **34**, 1425-1428 (*pmr, cmr*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1163-1169 (*Lycoposerramine K*)


Luciduline

Alkaloid L21

[21041-42-9]

[38142-93-7 ((\pm)-form), 240132-32-5]

L-280



(+)-form

$C_{13}H_{21}NO$ 207.315

Alkaloid from *Lycopodium lucidulum* and *Huperzia miyoshiana*. Liq. $[\alpha]_D^{+10}$ (c, 0.28 in EtOH).

2 α -Alcohol: Dihydrolyciduline

[21041-43-0]
 $C_{13}H_{23}NO$ 209.331

Alkaloid from *Lycopodium lucidulum* and *Huperzia miyoshiana*. Mp 65-70°.

11-Oxo: Lucidulinone

[21030-40-0]
 $C_{13}H_{19}NO_2$ 221.299

Alkaloid from *Lycopodium lucidulum*. $[\alpha]_D^{21} +185$ (c, 1.1 in CHCl₃).

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1968, **46**, 3631-3642 (*Luciduline, Dihydrolyciduline, isol, ir, ord, pmr, ms, cryst struct*)

Scott, W.L. *et al.*, *J.A.C.S.*, 1972, **94**, 4779 (*synth*)

Hall, D. *et al.*, *Cryst. Struct. Commun.*, 1973, **2**, 271-273 (*Dihydrolyciduline, cryst struct*)

Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 2755 (*synth, ir, pmr, ms*)

Szychowski, J. *et al.*, *Can. J. Chem.*, 1979, **57**, 1631-1637 (*synth*)

Schumann, D. *et al.*, *Annalen*, 1984, 1519-1528 (*synth*)

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241-298 (*rev*)

Comins, D.L. *et al.*, *Org. Lett.*, 1999, **1**, 229-231 (*synth*)

Tori, M. *et al.*, *Phytochemistry*, 2000, **53**, 503-509 (*Lucidulinone*)


Tong, X. *et al.*, *CA*, 2005, **142**, 435915 (*isol*)

Cypridina Luciferin

Cypridina luciferin

[7273-34-9]

L-281




$C_{22}H_{27}N_7O$ 405.502

Isol. from *Cypridina noctiluca* and *Vargula hilgendorffii*. Cryst. (MeOH) (as dihydrobromide). Mp 252-254° (dec.) (dihydrobromide). λ_{max} 218 (ϵ 27500); 270 (ϵ 17000); 310 (sh) (ϵ 11500); 435 (ϵ 9000) (MeOH aq.) (dihydrobromide). λ_{max} 221 (ϵ 29000); 243 (sh) (ϵ 12000); 275 (ϵ 14500); 307 (ϵ 20000); 320 (sh) (ϵ 6000) (MeOH/HBr aq.) (dihydrobromide).

- Shimomura, O. *et al.*, *Bull. Chem. Soc. Jpn.*, 1957, **30**, 929-933 (*isol, uv*)
 Kishi, Y. *et al.*, *Tet. Lett.*, 1966, 3427-3436; 3437-3444; 3445-3450 (*struct, synth*)
 Nakamura, H. *et al.*, *Tet. Lett.*, 2000, **41**, 2185-2188 (*synth*)
 Kato, S.-I. *et al.*, *Tetrahedron*, 2004, **60**, 11427-11434 (*biosynth, ms*)
 Wu, C. *et al.*, *Tet. Lett.*, 2006, **47**, 753-756 (*synth*)
 Teranishi, K. *et al.*, *Bioorg. Chem.*, 2007, **35**, 82-111 (*rev*)
 Kato, S. *et al.*, *Heterocycles*, 2007, **72**, 673-676 (*biosynth*)

Photinus Luciferin **L-282**
 4,5-Dihydro-2-(6-hydroxy-2-benzothiazolyl)-4-thiazolecarboxylic acid, 9CI. Firefly luciferin
 [20240-21-5]



$C_{11}H_8N_2O_3S_2$ 280.328

(S)-form

D-form

[2591-17-5]

Light-producing substance (luciferin) in the firefly *Photinus pyralis* and from *Luciola mingrelica*. Pale yellow cryst. (difficult to crystallize). Mp 204° (196°) dec. $[\alpha]_D^{24}$ -29 (DMF). Unstable to O_2 , light and acids.

[103404-75-7]

White, J.D. *et al.*, *J.A.C.S.*, 1963, **85**, 337 (*struct, synth, uv*)

Dennis, D. *et al.*, *Acta Cryst. B*, 1973, **29**, 1053 (*cryst struct*)


Rubin, B.A. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 293; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 303 (*synth, ms*)

Okada, K. *et al.*, *Chem. Comm.*, 1976, 32 (*biosynth*)

McCapra, F. *et al.*, *Chem. Comm.*, 1976, 153 (*biosynth*)

Orlova, G. *et al.*, *J.A.C.S.*, 2003, **125**, 6962-6971 (*bioluminescence*)

Lucilactaene **L-283**
 [386278-77-9]



$C_{22}H_{27}NO_6$ 401.458

Similar to Fusarin A, F-226. Prod. by *Fusarium* sp. RK97-94. Cell cycle inhibitor. Pale yellow solid. $[\alpha]_D$ +39.5 (c, 0.1 in MeOH). Readily undergoes racemisation. Isolable nat. prod. is racemic. λ_{max} 273 (ε 8800); 363 (ε 48200) (MeOH).

10'-Z-Isomer; 6 α -hydroxy: **Hydroxylucilactaene**


[1001568-32-6]

$C_{22}H_{27}NO_7$ 417.458

Prod. by *Fusarium acuminatum*. Called

- 13 α -Hydroxylucilactaene in the lit.
 Kakeya, H. *et al.*, *J. Antibiot.*, 2001, **54**, 850-853 (*isol, pmr, cmr*)
 Yamaguchi, J. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 3110-3115 (*synth, abs config*)
 Coleman, R.S. *et al.*, *J.A.C.S.*, 2005, **127**, 16038-16039 (*synth*)
 Bashyal, B.P. *et al.*, *Nat. Prod. Commun.*, 2007, **2**, 547-550 (*Hydroxylucilactaene*)

Lukianol A **L-284**
 3,7,8-Tris(4-hydroxyphenyl)-1H-pyrrolo[2,1-c][1,4]oxazin-1-one, 9CI
 [144398-24-3]



$C_{25}H_{17}NO_5$ 411.413

Isol. from an unidentified tunicate. Cytotoxic; MDR reversing agent. Amorph. powder (MeOH). Mp 264-266°. Mp refers to crystalline synthetic material. λ_{max} 206 (ε 33113); 282 (ε 35481); 344 (ε 14125) (MeOH) (Berdy).

3'-Iodo: Lukianol B

[144398-25-4]

$C_{25}H_{16}INO_5$ 537.31

Minor constit. from the unidentified tunicate. Cytotoxic. Off-white powder (MeOH).

Yoshida, W.Y. *et al.*, *Helv. Chim. Acta*, 1992, **75**, 1721 (*isol, uv, pmr, cmr, struct*)

Fürstner, A. *et al.*, *J.O.C.*, 1995, **60**, 6637 (*synth*)


Banwell, M.G. *et al.*, *Chem. Comm.*, 1997, 207 (*synth*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 54-62 (*synth*)

Gupton, J.T. *et al.*, *Tetrahedron*, 1999, **55**, 14515-14522 (*synth*)

Liu, J.-H. *et al.*, *J.O.C.*, 2000, **65**, 3587-3595 (*synth, pmr, cmr*)

Lumichrome **L-285**
 7,8-Dimethylbenzo[g]pteridine-2,4(1H,3H)-dione, 9CI. 7,8-Dimethylalloxazine
 [1086-80-2]
 [809279-55-8 (Isolumichrome)]



$C_{12}H_{10}N_4O_2$ 242.237

The 4-OH tautomer, called Isolumichrome in the lit., appears to be separately isolable. The reported Mp was the same. Irradiation prod. of Riboflavine. Prod. by *Micromonospora* sp. Tü 6368

and *Paecilomyces* sp. J300. Also isol. (as Isolumichrome) from the marine sponge *Cinachyrella australiensis*. Pale yellow cryst. ($CHCl_3$ or $AcOH$ aq.). Spar. sol. H_2O , $EtOH$, $CHCl_3$. Mp 300° dec. Blue fluor. in soln. λ_{max} 211 (log ε 4.53); 242 (log ε 4.54); 252 (sh) (log ε 4.47); 258 (log ε 4.66); 331 (log ε 3.93); 377 (log ε 3.87) (MeOH).

N^4 - α -Ribofuranosyl: N^4 - α -Ribofuranosyl-lumichrome

$C_{17}H_{18}N_4O_6$ 374.352

Prod. by *Micromonospora* sp. Tü 6368. Yellow solid. $[\alpha]_D^{20}$ +319 (c, 0.1 in MeOH). λ_{max} 214 (log ε 4.45); 254 (log ε 4.55); 340 (log ε 3.85) (MeOH). λ_{max} 219 (log ε 4.52); 253 (log ε 4.54); 331 (log ε 3.95); 383 (log ε 3.92) (MeOH/NaOH).

N^4 -Me: 1,7,8-Trimethylbenzo[g]pteridine-2,4(1H,3H)-dione. N^4 -Methyl-lumichrome

[18950-64-6]

$C_{13}H_{12}N_4O_2$ 256.263

Prod. by *Micromonospora* sp. Tü 6368. Yellow solid. λ_{max} 219 (log ε 4.43); 250 (log ε 4.44); 259 (sh) (log ε 3.78); 386 (log ε 3.76) (MeOH).

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 768B (*nmr*)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 894D (*ir*)

Karrer, P. *et al.*, *Helv. Chim. Acta*, 1934, **17**, 1010 (*isol*)

Seng, F. *et al.*, *Angew. Chem., Int. Ed.*, 1972, **11**, 1010 (*synth*)

Yoneda, F. *et al.*, *J.A.C.S.*, 1976, **98**, 830 (*synth*)

Villemin, D. *et al.*, *Synth. Commun.*, 1995, **25**, 2319 (*synth, ms*)

Sikorska, E. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 2003, **77**, 65-73 (*spectra*)


Kwon, H.C. *et al.*, *Arch. Pharmacol. Res.*, 2004, **27**, 604-609 (*isol, uv, pmr, cmr*)

Li, L. *et al.*, *CA*, 2004, **142**, 52432n (*isol, Isolumichrome*)

Antal, N. *et al.*, *J. Antibiot.*, 2005, **58**, 95-102; 103-110 (*isol, pmr, cmr, ribosyl, N-Me*)

Sigma-Aldrich Library of Stains, Dyes and Indicators, 430

α -Lumicolchicine **L-286**
 [7183-82-6]



$C_{44}H_{50}N_2O_{12}$ 798.885


Product of irradiation of Colchicine, C-569. Possible alkaloid artifact. Mp 165°. $[\alpha]_D$ +81.

Chapman, O.L. *et al.*, *J.A.C.S.*, 1961, **83**, 3916 (*ir, uv, pmr, struct*)

β-Lumicolchicine

L-287

Lumicolchicine I. Substance I
[6901-13-9]



C₂₂H₂₅NO₆ 399.443

Numbering systems vary. Alkaloid from many *Colchicum* spp. (Liliaceae) and other spp. as photoisom. prod. of Colchicine, C-569. May be a genuine alkaloid. Cryst. (EtOH/Et₂O). Mp 183° Mp 206° (dimorph.).

N-De-Ac, N-formyl: N-Deacetyl-N-formyl-β-lumicolchicine, 8CI
[18172-23-1]

C₂₁H₂₃NO₆ 385.416

Alkaloid from tubers of *Gloriosa superba* (Liliaceae). Mp 214-215°. [α]_D²⁰ +340 (c, 1 in CHCl₃).

N-De-Ac, N,N-di-Me: N-Deacetyl-N,N-dimethyl-β-lumicolchicine

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Colchicum speciosum*. Mp 158-160°. [α]_D +321.

N-De-Ac, N-Me, N-(2-hydroxybenzyl): β-Lumispeciosine

C₂₈H₃₁NO₆ 477.556

Alkaloid from *Colchicum speciosum*. Cryst. (Me₂CO). Mp 157-158°. [α]_D +132 (CHCl₃). λ_{max} 226 (sh); 265; 340

N-Me: N-Methyl-β-lumicolchicine

C₂₃H₂₇NO₆ 413.469

Alkaloid from *Colchicum speciosum* (Liliaceae). Mp 132-133°. [α]_D +410 (CHCl₃).

2-O-De-Me: 2-O-Demethyl-β-lumicolchicine

[18172-24-2]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Colchicum autumnale*, *Colchicum kesselringii*, *Gloriosa superba* and *Wurmbea* spp. Cryst. (EtOAc). Mp 198-200°. [α]_D²⁰ +337 (c, 0.45 in CHCl₃). λ_{max} 226; 264; 344 (MeOH).

3-O-De-Me: 3-O-Demethyl-β-lumicolchicine

[30632-52-1]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Colchicum* spp., *Gloriosa superba* and *Merendera raddeana*. Mp 235°. [α]_D²² +316 (CHCl₃).

2-O-De-Me, N-de-Ac, N-formyl: N-Deacetyl-N-formyl-2-O-demethyl-β-lumicolchicine

[18172-25-3]

C₂₀H₂₁NO₆ 371.389

Alkaloid from *Colchicum decaisnei* and *Gloriosa superba* (Liliaceae). Cryst. (MeOH/EtOAc). Mp 229-230°. [α]_D +240 (c, 0.12 in MeOH). λ_{max} 201; 221; 266 (MeOH). λ_{max} 201; 291 (MeOH/

NaOH).

8,12-Diepimer: γ-Lumicolchicine. Lumicolchicine II. Substance J

[6901-14-0]

C₂₂H₂₅NO₆ 399.443

Alkaloid from *Colchicum* spp., *Merendera* spp. and others, as artifact of illumination of Colchicine, C-569 and poss. as genuine alkaloid (Liliaceae). Cryst. (EtOAc/Et₂O, dioxan or anisole). Mp 268°. [α]_D¹⁸ -445 (c, 1 in CHCl₃). λ_{max} 265 (EtOH) (Berdy).

8,12-Diepimer, N-de-Ac, N-formyl: N-Deacetyl-N-formyl-γ-lumicolchicine
[18172-22-0]

C₂₁H₂₃NO₆ 385.416

Alkaloid from tubers of *Gloriosa superba* (Liliaceae). Cryst. (CHCl₃/MeOH). Mp 255-257°. [α]_D²⁰ -451 (c, 1 in CHCl₃).

8,12-Diepimer, 2-O-de-Me: 2-O-Demethyl-γ-lumicolchicine. Alkaloid K13
[26194-64-9]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Colchicum autumnale*, *Gloriosa superba* and *Merendera raddeana*. Mp 287-288°. [α]_D -420 (c, 0.40 in CHCl₃).

8,12-Diepimer, 3-O-de-Me: 3-O-Demethyl-γ-lumicolchicine. Alkaloid L6
[28101-27-1]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Colchicum luteum* (Liliaceae). Mp 291-293°. [α]_D²² -410 (CHCl₃).

8,12-Diepimer, 2-O-de-Me, N-Me, 2-O-β-D-glucopyranoside: Colchicum autumnale Alkaloid M
[29623-34-5]

C₂₈H₃₅NO₁₁ 561.585

Alkaloid from *Colchicum autumnale* (Liliaceae). Mp 310-314°. [α]_D²² -310 (c, 0.89 in MeOH aq.).

Grewe, R. et al., *Chem. Ber.*, 1951, **84**, 621-627 (synth, uv)

Gardner, P.D. et al., *J.A.C.S.*, 1957, **79**, 6334-6337 (struct)

Chapman, O.L. et al., *J.A.C.S.*, 1963, **85**, 803-806; 807-812 (struct, pmr)

Canonica, L. et al., *Chim. Ind. (Milan)*, 1967, **49**, 1304-1312; *CA*, **68**, 87424s (N-Deacetyl-N-formyl-lumicolchicine)

Severini Ricca, G. et al., *Gazz. Chim. Ital.*, 1969, **99**, 133-151 (pmr)

Canonica, L. et al., *Tet. Lett.*, 1969, 607-608 (struct)

Santavy, F. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 2857-2860 (Alkaloid M)

Chommadov, B.C. et al., *Khim. Prir. Soedin.*, 1970, **6**, 275; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 274-275 (3-O-Demethyl-γ-lumicolchicine, Deacetyldimethyl-lumicolchicine)

Turdikulov, K. et al., *Khim. Prir. Soedin.*, 1971, **7**, 541; 1972, **8**, 502-505; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 528; 1972, **8**, 494-496 (2-O-Demethyl-β-lumicolchicine, Alkaloid K13)

Hrbek, J. et al., *Coll. Czech. Chem. Comm.*, 1982, **47**, 2258-2279 (cd)

Meksuriyen, D. et al., *J. Nat. Prod.*, 1988, **51**, 88-93 (pmr, cmr, β-Lumicolchicine, γ-Lumicolchicine)

Chommadov, B. et al., *Khim. Prir. Soedin.*, 1990, **26**, 147-165; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 113-128 (N-Methyl-lumicolchicine)


Al-Tel, T.H. et al., *Phytochemistry*, 1991, **30**, 3081-3085 (Deacetylformyl-demethyl-β-lumicolchicine)

Shakirov, R. et al., *Khim. Prir. Soedin.*, 1996, **32**, 615-681; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 596-675 (β-Lumispeciosine)

Lunacridine

L-288

3-(2-Hydroxy-3-methylbutyl)-4,8-dimethoxy-1-methyl-2(1H)-quinolinone, 9CI



(R)-form

C₁₇H₂₃NO₄ 305.373

(R)-form [83-58-9]

Alkaloid from *Lunasia amara* bark (Rutaceae). Hypotensive agent. Mp 86-87°. [α]_D²⁴ +30. Dieterle and Beyl (1937) describe an alkaloid by this name; it was probably Lunamarine, L-290. Low toxicity (LD₅₀ =1100 mg/kg in mice).

Perchlorate: Mp 146-148° (solidifies and remelts at 193-195°).

3'-Hydroxy: 3-(2,3-Dihydroxy-3-methylbutyl)-4,8-dimethoxy-1-methyl-2(1H)-quinolinone. Balfourlone. Hydroxylunacridine
[478-68-2]

C₁₇H₂₃NO₅ 321.372

Artifact arising from ring opening of Balfouronin, B-14 during isol. Cryst. (CCl₄/hexane). Mp 99-100°. [α]_D -36. Abs. config. revised in 2002.

(S)-form

Synthetic. Viscous oil. [α]_D -28 (c, 0.84 in MeOH).

(±)-form

Mp 72-74°.

(ξ)-form

3',4'-Didehydro: Acutifoline

[145237-07-6]

C₁₇H₂₁NO₄ 303.357

Alkaloid from the leaves of *Zanthoxylum petiolare* (Rutaceae). Viscous oil. [α]_D +40 (c, 0.00025 in CHCl₃). Source originally misidentified.

3',4'-Didehydro, O-hexadecanoyl: Acutifoline palmitate

[145204-98-4]

C₃₃H₅₁NO₅ 541.77

Alkaloid from leaves of *Zanthoxylum petiolare* (Rutaceae). Viscous oil. [α]_D +15 (c, 0.00066 in CHCl₃).

3',4'-Didehydro, N-de-Me: Acutifolidine†
[145237-08-7]

C₁₆H₁₉NO₄ 289.33

Alkaloid from leaves of *Zanthoxylum petiolare* (Rutaceae). Amorph. solid.

Mp 121-123°. $[\alpha]_D +18.6$ (c, 0.00053 in CHCl_3).

3',4'-Didehydro, Me ether: O-Methylacutifoline

[145204-97-3]

$\text{C}_{18}\text{H}_{23}\text{NO}_4$ 317.384

Alkaloid from leaves of *Zanthoxylum petiolare* (Rutaceae). Viscous oil. $[\alpha]_D -6.4$ (c, 0.00156 in CHCl_3).

Goodwin, S. et al., *J.A.C.S.*, 1959, **81**, 1908 (uv, ir, struct)

Rapoport, H. et al., *J.A.C.S.*, 1959, **81**, 3738 (Balfourone)

Clarke, E.A. et al., *J.C.S.*, 1964, 438 (synth)

Bowman, R.M. et al., *J.C.S. Perkin 1*, 1973, 1051 (config)

Brown, N.M.D. et al., *Tetrahedron*, 1980, **36**, 3579 (cmr)

Arruda, M.S.P. et al., *Phytochemistry*, 1992, **31**, 3617; 1994, **36**, 1303 (Acutifoline, Acutifoline palmitate, Acutifolidine, O-Methylacutifoline)

Barr, S.A. et al., *Tetrahedron*, 1994, **50**, 11219 (synth)

Anand, R.C. et al., *J. Chem. Res., Synop.*, 1998, 6-7 (synth, abs config)

Boyd, D.R. et al., *Chem. Comm.*, 2002, 3070-3071 (Balfourone, config)

Lunamaridine

L-289

$\text{C}_{16}\text{H}_{15}\text{NO}_2$ 253.3


Struct. unknown. Alkaloid from the bark of *Lunasia amara*. Small needles (EtOH aq.). Mp 209-210°.

Steldt, F.A. et al., *J. Am. Pharm. Assoc.*, 1943, **32**, 107-111 (isol)

Lunamarine

L-290

7-Methoxy-1-methyl-2-(3,4-methylene-dioxyphenyl)-4(1H)-quinolinone



$\text{C}_{18}\text{H}_{15}\text{NO}_4$ 309.321

Alkaloid from *Lunasia amara* (Rutaceae). Hypotensive agent, weak smooth muscle stimulant. Mp 245-246°. Low toxicity in mice.

Steldt, F.A. et al., *J. Am. Pharm. Assoc.*, 1943, **32**, 107; *CA*, **37**, 3562 (isol)

Goodwin, S. et al., *J.A.C.S.*, 1959, **81**, 6209 (isol, ir, uv)

Lunariamine

L-291

$\text{C}_{24}\text{H}_{33}\text{N}_3\text{O}_4$ 427.542

Prob. spermidine-type alkaloid. Struct. unknown. Minor constit. of the seeds of *Lunaria biennis* and *Lunaria rediviva* (Brassicaceae). Yellow prisms ($\text{CHCl}_3/\text{EtOH}$). Mp 248° (tube) Mp 253-254° dec Mp 290° (block). $[\alpha]_D^{20} 0$ (c, 0.37 in AcOH).


Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1956, 1840-1842 (isol, uv, ir)

Huneck, S. et al., *Naturwissenschaften*, 1962, **49**, 233 (isol)

Lunaridine

[34340-56-2]

L-292



$\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_4$ 437.538

Isomeric with Lunarine, L-293. Alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Needles. Mp 265° (261°)(block). $[\alpha]_D +233$ (c, 0.25 in CHCl_3).

Hydroiodide:

Cryst. (H_2O). Mp 335° dec.

Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1956, 1840 (isol, uv, ir)

Potier, P. et al., *Bull. Soc. Chim. Fr.*, 1959, 456 (isol, uv)

Poupat, C. et al., *Tetrahedron*, 1972, **28**, 3103 (uv, ir, pmr, struct)


Nagao, Y. et al., *Chem. Comm.*, 1981, 286; *Heterocycles*, 1982, **17**, 537 (synth)

Fujita, E. et al., *Pure Appl. Chem.*, 1981, **53**, 1141 (synth)

Lunarine

[24185-51-1]

L-293



$\text{C}_{25}\text{H}_{31}\text{N}_3\text{O}_4$ 437.538

Major alkaloid from the seeds of *Lunaria biennis* and *Lunaria rediviva* (Brassicaceae). Needles (MeOH aq.). Mp 239-240° (block)(232-235°). $[\alpha]_D^{20} +291$ (c, 1.03 in CHCl_3).

Hydrochloride: Mp 340°.

3 α -Alcohol: Alkaloid LBY. Lunarinol II.

3-Deoxo-3-hydroxylunarine

[24185-52-2]

$\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_4$ 439.553

Minor alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Cryst. + $2\text{H}_2\text{O}$ (MeOH/Et₂O). Mp 268-273°. $[\alpha]_D^{20} +108$ (c, 0.43 in EtOH).

[79298-94-5 ((±)-form)]

Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1956, 1840 (isol, uv, ir)

Huneck, S. et al., *Naturwissenschaften*, 1962, **49**, 233 (isol)

Potier, P. et al., *Tet. Lett.*, 1963, 293 (uv, pmr, struct)

Jeffreys, J.A.D. et al., *J.C.S. (B)*, 1970, 826 (cryst struct)

Tamura, C. et al., *J.C.S. (B)*, 1970, 991 (cryst struct)

Poupat, C. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 433 (biosynth)

Doskotch, R.W. et al., *Experientia*, 1972, **28**, 382-383 (Alkaloid LBY)

Poupat, C. et al., *Tetrahedron*, 1972, **28**, 3087-3101 (Lunarine, Lunarinol II, pmr, struct)

Fujita, E. et al., *Pure Appl. Chem.*, 1981, **53**, 1141 (synth)

Nagao, Y. et al., *Heterocycles*, 1982, **17**, 537 (synth)

Sagner, S. et al., *Tet. Lett.*, 1997, **38**, 2443 (synth)


Sagner, S. et al., *Phytochemistry*, 1998, **47**, 375-387 (cd, pmr, cmr, biosynth)

Hamilton, C.J. et al., *J.C.S. Perkin 1*, 2002, 1115-1123 (synth)

Lundurine A

[162616-56-0]

L-294



$\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_4$ 366.416

Highly rearranged indole alkaloids prob. most closely related to the Aspidofractine group. Alkaloid from the leaves of *Kopsia tenuis*. Cytotoxic. Oil. $[\alpha]_D -90$ (c, 0.09 in CHCl_3), λ_{max} 208 (log ϵ 4.71); 250 (log ϵ 4.34); 298 (log ϵ 3.8) (EtOH).

3-Deoxo: Lundurine B

[162616-57-1]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from the leaves of *Kopsia tenuis*. Cytotoxic. Light yellow oil. $[\alpha]_D -34$ (c, 0.16 in CHCl_3), λ_{max} 211 (log ϵ 4.43); 252 (log ϵ 4.07); 299 (log ϵ 3.92) (EtOH).

3-Deoxo, 14,15-dihydro: Lundurine C

[162616-58-2]

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3$ 354.448

Alkaloid from the leaves of *Kopsia tenuis*. Light yellow oil. $[\alpha]_D -25$ (c, 0.067 in CHCl_3), λ_{max} 203 (log ϵ 4.71); 253 (log ϵ 4.42); 299 (log ϵ 3.92) (EtOH).

11-Methoxy, 3-deoxo: Lundurine D

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458


Alkaloid from the leaves of *Kopsia tenuis*. Light yellow oil. $[\alpha]_D -22$ (c, 0.01 in CHCl_3), λ_{max} 215 (log ϵ 4.38); 255 (log ϵ 3.94); 302 (log ϵ 3.71) (EtOH).

Kam, T.-S. et al., *Tetrahedron*, 2004, **60**, 10739-10745 (isol, uv, pmr, cmr, ms)

Lunidonine

[6899-80-5]

L-295



$\text{C}_{17}\text{H}_{19}\text{NO}_5$ 317.341

Alkaloid from the bark of *Lunasia amara* var. *repanda* (Rutaceae). Cryst. (MeOH). Mp 118-119°.

Semicarbazone:

Cryst. (EtOAc). Mp 209-211°.

N-De-Me: N-Demethylunidonine

[55396-51-5]

C₁₆H₁₇NO₅ 303.314
Alkaloid from *Orixa japonica* and *Ptelea trifoliata*. Powder. λ_{\max} 223 (ε 45800); 257 (ε 36600); 263 (ε 37200); 318 (ε 16200) (MeOH).

2'- ξ -Alcohol: **Lunidine**

[6899-79-2]
C₁₇H₂₁NO₅ 319.357
Alkaloid from the bark of *Lunasia amara* var. *repanda* (Rutaceae). Cryst. (diisopropyl ether). Mp 65-66.5°. $[\alpha]_{\text{D}}^{20}$ +28 (c, 0.4 in EtOH).

2'- ζ -Alcohol, Ac:

Cryst. (hexane/butanone). Mp 111-112°. $[\alpha]_{\text{D}}^{20}$ +85 (c, 0.4 in EtOH).

3'-Hydroxy: **Hydroxylunidonine**

[55812-53-8]
C₁₇H₁₉NO₆ 333.34
Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (hexane/EtOAc). Mp 202-205°. λ_{\max} 220; 265 (sh); 317; 330 (sh) (MeOH).

6-Methoxy: **6-Methoxylunidonine**

[55812-54-9]
C₁₈H₂₁NO₆ 347.367
Alkaloid from *Ptelea trifoliata* (Rutaceae). Needles (diisopropyl ether). Mp 123-125°.

Rüegger, A. et al., *Helv. Chim. Acta*, 1963, **46**, 2329 (isol, uv, ir, pmr, ms, struct, Lunidonine, Lunidine)


Mitscher, L.A. et al., *J. Nat. Prod.*, 1975, **38**, 117 (6-Methoxylunidonine)

Reisch, J. et al., *Phytochemistry*, 1975, **14**, 2722-2723 (Hydroxylunidonine)

Funayama, S. et al., *Phytochemistry*, 1994, **36**, 525-528; *CA*, **83**, 40169g (N-Demethylunidonine)

Lunine L-296

7,10-Dihydro-10-methyl-8-(1-methyl-ethyl)-1,3-dioxolo[4,5-h]furo[2,3-b]quinolin-6(8H)-one, 9CI, 3,9-Dihydro-2-isopropyl-7,8-methylenedioxy-9-methyl-furo[2,3-b]quinolin-4(2H)-one



C₁₆H₁₇NO₄ 287.315

(S)-form [518-60-5]

Alkaloid from leaves of *Lunasia amara* and other *Lunasia* spp. (Rutaceae). Prisms (EtOAc). Mp 228-229° (222-223°). $[\alpha]_{\text{D}}^{20}$ -39 (c, 1.5 in CHCl₃).
Hydrochloride: Mp 215-216° dec.

(ξ)-form

1',2'-Didehydro: **Ptelefolidone**

[58880-21-0]
C₁₆H₁₅NO₄ 285.299
Alkaloid from flowers of *Ptelea trifoliata* (Rutaceae). Cryst. (hexane/Me₂CO). Mp 152-154°.

Goodwin, S. et al., *J.A.C.S.*, 1959, **81**, 3065-3069; 6209-6213 (pmr, struct, ir, uv, isol)

McKay, M.F. et al., *Acta Cryst. B*, 1969, **25**, 1925-1932 (cryst struct)

Reisch, J. et al., *Phytochemistry*, 1975, **14**, 2722-2723 (Ptelefolidone)

Hammerum, S. et al., *Acta Chem. Scand., Ser. B*, 1977, **31**, 31-39 (ms)


Lunolone L-297

[1359-18-8]
Struct. unknown. Alkaloid from the leaves of *Lunasia amara*. Prisms (EtOAc/pentane). Mp 100-103°. $[\alpha]_{\text{D}}^{23}$ +20.6 (c, 1 in EtOH).

Goodwin, S. et al., *J.A.C.S.*, 1959, **81**, 6209-6213 (isol)

Luotonine A L-298

Quino[2',3':3,4]pyrrolo[2,1-b]quinazolin-11(13H)-one
[205989-12-4]



C₁₈H₁₁N₃O 285.304

Alkaloid from *Peganum nigellastrum*. Cytotoxic towards P388 cells. Active against human DNA topoisomerase I. Yellow needles (CHCl₃/Me₂CO). Mp 281-283° dec. λ_{\max} 212 (log ε 4.61); 248 (log ε 4.71); 298 (log ε 4); 326 (sh) (log ε 4.18); 341 (log ε 4.27); 358 (log ε 4.21) (MeOH).

7-Hydroxy: **Luotonine B**

[205989-13-5]
C₁₈H₁₁N₃O₂ 301.304
Alkaloid from *Peganum nigellastrum*. Cytotoxic towards P388 cells. Yellow cryst. Mp 237-240°. Racemic. λ_{\max} 212 (log ε 4.67); 249 (log ε 4.69); 296 (sh) (log ε 4.1); 322 (sh) (log ε 4.27); 337 (log ε 4.32); 352 (log ε 4.22) (MeOH).

7-Methoxy: **Luotonine E**

C₁₉H₁₃N₃O₂ 315.331
Alkaloid from *Peganum nigellastrum*. Pale yellow cryst. Mp 222-225°. Racemic. λ_{\max} 213 (log ε 4.67); 250 (log ε 4.66); 301 (log ε 4.13); 323 (sh) (log ε 4.21); 337 (log ε 4.28); 352 (log ε 4.18) (MeOH).

Ma, Z.-Z. et al., *Heterocycles*, 1997, **46**, 541-546; 1999, **51**, 1593-1596; 1883-1889 (Luotonines A,B,E, isol, uv, ir, pmr, cmr, ms, synth, activity)

Kelly, T.R. et al., *Tet. Lett.*, 1999, **40**, 2723-2724 (synth)

Toyota, M. et al., *Heterocycles*, 2002, **56**, 101-103 (synth)

Dallavalle, S. et al., *Tet. Lett.*, 2002, **43**, 1835-1837 (synth)

Yadav, J.S. et al., *Tet. Lett.*, 2002, **43**, 1905-1907 (synth)

Osborne, D. et al., *Tet. Lett.*, 2002, **43**, 5469-5470 (synth)

Cagir, A. et al., *J.A.C.S.*, 2003, **125**, 13628-13629 (activity)

Twin, H. et al., *Org. Lett.*, 2004, **6**, 4913-4916 (synth)

Chavan, S.P. et al., *Tetrahedron*, 2004, **60**, 9931-9935 (synth)

Harayama, T. et al., *Tetrahedron*, 2004, **60**, 10645-10649 (synth)

Mhaske, S.B. et al., *Tetrahedron*, 2006, **62**, 9787-9826 (rev, synth)


Zhou, H.-B. et al., *J.O.C.*, 2007, **72**, 6270-6272 (synth)

Bowman, W.R. et al., *Org. Biomol. Chem.*, 2007, **5**, 103-113 (synth)

Mason, J.J. et al., *Org. Biomol. Chem.*, 2007, **5**, 2486-2490 (synth)

Luotonine C L-299

9-Methoxy-5-methylcanthin-6-one
[261948-33-8]



R = CH₃

C₁₆H₁₂N₂O₂ 264.283

Alkaloid from *Peganum nigellastrum*. Pale yellow needles (Me₂CO). Mp 166-168°. λ_{\max} 209 (log ε 4.35); 265 (sh) (log ε 3.9); 273 (log ε 4.02); 308 (log ε 3.66); 346 (log ε 3.71); 358 (log ε 3.7); 376 (sh) (log ε 3.47) (MeOH).

Ma, Z.-Z. et al., *Phytochemistry*, 2000, **53**, 1075-1078 (isol, uv, pmr, cmr)

Luotonine D L-300

5-Ethyl-9-methoxycanthin-6-one

[261948-34-9]
As Luotonine C, L-299 with
R = CH₂CH₃


C₁₇H₁₄N₂O₂ 278.31

Alkaloid from *Peganum nigellastrum*. Pale yellow needles (Me₂CO). Mp 141-143°. λ_{\max} 210 (log ε 4.36); 266 (sh) (log ε 3.92); 273 (log ε 4.06); 310 (log ε 3.6); 347 (log ε 3.76); 358 (log ε 3.75); 376 (sh) (log ε 3.54) (MeOH).

Ma, Z.-Z. et al., *Phytochemistry*, 2000, **53**, 1075-1078 (isol, uv, pmr, cmr)

Luotonine F L-301

[244616-85-1]



C₁₈H₁₁N₃O₂ 301.304


Alkaloid from aerial parts of *Peganum nigellastrum*. Pale yellow needles. Mp 238-240°. λ_{\max} 212 (log ε 4.4); 255 (log ε 3.98); 303 (log ε 3.83); 316 (sh) (log ε 3.79); 328 (sh) (log ε 3.7) (MeOH).

Ma, Z.-Z. et al., *Heterocycles*, 1999, **51**, 1883-1889 (isol, synth, uv, ir, pmr, cmr, ms)

Mhaske, S.B. et al., *Synthesis*, 2002, 323-325 (synth, ir, pmr, cmr, ms)

Lupanine**L-302**

2-Oxo-11 α -sparteine. Dilupanine. Trilupanine. Hydrorhombinine. Tetrahydroanagryrine

**(+)-form**C₁₅H₂₄N₂O 248.367

Dilupanine was impure Lupanine. Trilupanine (originally assigned the struct. Lupanine di-N-oxide) was lupanine monohydrochloride dihydrate. Hydrorhombinine and Tetrahydroanagryrine were the (-)-form.

(+)-form [550-90-3]

Alkaloid from *Cytisus scoparius* and several other *Cytisus* spp., *Lupinus* spp.; also in *Leontice* spp., *Cadia purpurea*, *Ammopiptanthus mongolicus*, *Genista* spp., *Templetonia* spp., *Thermopsis chinensis* (Fabaceae, Leonticaceae). Toxic cause (together with other alkaloids including the (-)- and (\pm)-forms) of lupinosis in livestock fed on *Lupinus* spp. Shows antimicrobial activity. Mp 40°.

Bp_{0.08} 185-186°. [α]_D +61.4 (Me₂CO).

► LD₅₀ (rat, orl) 1440 mg/kg. OK5745000

Hydrochloride: Mp 269-271°.

Hydrochloride (1:2):

Cryst. + 1H₂O. Mp 162-163°.

Picrate: Mp 185°.

N¹⁶-Oxide: **Lupanine N-oxide**

[3019-47-4]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from the seeds of *Thermopsis lupinoides* (Fabaceae). Hygroscopic needles (C₆H₆). Mp 58°. [α]_D²⁴ +65.5 (EtOH).

(-)-form [486-88-4]

Alkaloid from *Lupinus pusillus*, *Lupinus macounii*, *Baptisia versicolor*, *Podalyria calyptata*, *Ammodendron* spp., *Leontice smirnovii* and *Leontice eversmannii* (Fabaceae, Leonticaceae). Liq. Bp₁ 186-188°. [α]_D -61 (Me₂CO).

Hydroiodide:

Cryst. + 2H₂O. Mp 190°. [α]_D -43.6 (H₂O).

(\pm)-form [4356-43-8]

Alkaloid from *Lupinus albus*, *Lupinus termis*, *Podalyria buxifolia*, and *Virgilia capensis* (Fabaceae). Mp 98°.

Hydrochloride:

Cryst. + 2H₂O. Mp 127-128° Mp 250-252° (anhyd.).

Picrate: Mp 183° dec.

N¹⁶-Oxide: [133163-04-9]

C₁₅H₂₄N₂O₂ 264.367

Constit. of *Lupinus termis*. Oil.

Clemo, G.R. et al., *J.C.S.*, 1928, 1811-1820; 1931, 429-437 (struct, resoln)

Couch, J.F. et al., *J.A.C.S.*, 1937, 59, 1469-1471 (isol)

Marion, L. et al., *J.O.C.*, 1948, 13, 780-781 (isol)

Marion, L. et al., *Can. J. Chem.*, 1952, 30, 386-387 (Dilupanine, Trilupanine)

Van Tamelen, E.E. et al., *J.A.C.S.*, 1956, 78, 2913-2914 (synth)

Clemo, G.R. et al., *J.C.S.*, 1956, 3390-3394 (synth, struct)

Bohlmann, F. et al., *Tet. Lett.*, 1965, 6, 2435-2440 (pmr)

Wiewirowski, M. et al., *Can. J. Chem.*, 1967, 45, 1447-1457 (ir, pmr, conformn)

Fales, H.M. et al., *J.A.C.S.*, 1970, 92, 1590-1597 (ms)

Barciszewski, J. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1971, 19, 545-548 (pmr)

Bohlmann, F. et al., *Chem. Ber.*, 1975, 108, 1043-1051 (cmr)

Skrzypczak-Jankun, E. et al., *Acta Cryst. B*, 1978, 34, 2651-2653 (cryst struct)

Golebiewski, W.M. et al., *Chem. Comm.*, 1983, 1509-1511 (biosynth)

Wink, M. et al., *Z. Naturforsch., C*, 1984, 39, 548-552 (activity)

Wippich, C. et al., *Experientia*, 1985, 41, 1477-1479 (activity)

Saito, K. et al., *Phytochemistry*, 1988, 27, 3715-3716 (isol, ir, pmr, cmr, ms, struct, oxide)

Saito, K. et al., *CA*, 1990, 114, 203603 (Lupanine N-oxide)

Mohamed, M.H. et al., *J. Nat. Prod.*, 1990, 53, 1578-1580 ((\pm)-N-oxide)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LIQ800

Lupilaxine**L-303**C₁₅H₂₄N₂O₂ 264.367

Quinolizidine alkaloid. Struct. unknown. Alkaloid from *Lupinus laxus* and *Lupinus sericeus* (Fabaceae). Prisms (Me₂CO). Mp 175°.

Couch, J.F. et al., *J.A.C.S.*, 1937, 59, 1469 (isol)

Marion, L. et al., *Can. J. Chem.*, 1953, 31, 181-185 (isol)


Lurenine**L-304**

Struct. unknown. Alkaloid from *Lobelia urens* (Campanulaceae). Mp 202-204°.

Steinegger, E. et al., *Pharm. Acta Helv.*, 1950, 25, 276-277; *CA*, 45, 3853

Lusitanine**L-305**

N-[(Hexahydro-2H-quinolizin-1(6H)-ylidene)methyl]acetamide, 9CI

**(R)-form**C₁₂H₂₀N₂O 208.303

Needles (Me₂CO).

(R)-form

Alkaloid from freshly-harvested young stems of *Maaackia amurensis* var. *buergeri* and *Maaackia tashiroi* (Fabaceae). Needles. Mp 185-186°. [α]_D²³ -4.6 (c, 0.60 in EtOH).

(S)-form [5121-36-8]

Alkaloid from *Genista lusitanica*, *Cadia ellisiana* bark and *Chamaecytisus austriacus* (Fabaceae). Mp 184-186° dec. (168-170°). [α]_D²⁷ +6.9 (c, 2.5 in EtOH).

Dihydro: Mp 96-100°.

epi-Dihydro: Mp 140-144°.

Steinegger, E. et al., *Pharm. Acta Helv.*, 1965, 40, 610 (isol, uv, ir, ms)

Wicky, K. et al., *Pharm. Acta Helv.*, 1965, 40, 658; 1982, 57, 215 (ir, pmr, uv, struct, synth, config)


Faugeras, G. et al., *Plant. Med. Phytother.*, 1975, 9, 37 (isol)

Daily, A. et al., *Planta Med.*, 1979, 36, 188 (isol)

Saito, K. et al., *Chem. Pharm. Bull.*, 1987, 35, 1308 (ir, pmr, cmr, ms, abs config)

Luteicine**L-306**

[1359-25-7]

C₂₀H₂₇NO₄ 345.438

Synthesised from Luteidine, L-307. Since the struct. of Luteidine, L-307 has been revised, Luteicine must be formulated as shown. Alkaloid from *Colchicum luteum* epigeal parts (Liliaceae). Free base rapidly oxid. in air.

Ac:

Cryst. (Me₂CO). Mp 210-211°.


Ketone: **Luteicinone**

Mp 265-266° (as hydrochloride).

Yusupov, M.K. et al., *Khim. Prir. Soedin.*, 1976, 359; *Chem. Nat. Compd. (Engl. Transl.)*, 313

Luteidine**L-307**

[55790-08-4]

C₂₀H₂₅NO₄ 343.422

Struct. revised in 1985. Isomeric 2,13-dimethoxy struct. originally proposed. Alkaloid from *Colchicum luteum* (Liliaceae). Mp 231-232°. [α]_D -96 (MeOH).

N-Oxide(α -): **Luteidine α -N-oxide**

C₂₀H₂₅NO₅ 359.421

Alkaloid from *Colchicum luteum*. Mp 179-180°.

N-Oxide(β -): **Luteidine β -N-oxide**

C₂₀H₂₅NO₅ 359.421

Alkaloid from *Colchicum luteum*.

Mukhamed'yarova, N.L. et al., *Khim. Prir. Soedin.*, 1976, 12, 354-359; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, 12, 308-312 (isol, uv, ir, pmr, cmr, ms)


Timbekov, E.K. et al., *Khim. Prir. Soedin.*, 1985, 21, 3-11; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, 21, 1-9 (ms, struct)

- Chommatov, B. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 810-815; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 770-773 (struct)
- Nazarov, G.B. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 348-352; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 321-325 (cryst struct)
- Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 761-863; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 737-858 (N-oxides)

Luteinine

L-308

[1359-27-9]

C₁₉H₂₅NO₃ 315.411

Alkaloid from *Colchicum luteum*. λ_{max} 220 (log ε 4.12); 246 (sh) (log ε 4.04); 278 (sh) (log ε 3.2) (no solvent reported).

Sadykov, A.S. *et al.*, *Zh. Prikl. Khim. (Leningrad)*, 1965, **38**, 222-225; *CA*, **62**, 12160d (isol)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 615-681; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 596-675 (rev)

Lutenurine

L-309

[12698-70-3]

Struct. unknown. Alkaloid complex from *Nuphar luteum*. Antiprotozoal and antitrichomonal agent. Main component appears to be Nuphleine, in T-380.


Rubinichik, M.A. *et al.*, *Antibiotiki (Moscow)*, 1969, **14**, 926-929

Vichkanova, S.A. *et al.*, *Farmatsiya (Moscow)*, 1969, **18**, 66-70

Lutessine

L-310

[102487-18-3]

C₁₉H₂₁NO₆ 359.378

Alkaloid from the bulbs of *Sternbergia lutea* (Amaryllidaceae). Cryst. (CHCl₃). Mp 114-118°. [α]_D²⁵ -13.4 (c, 1.24 in EtOH). The first lycorine-type alkaloid with a substituent on the D ring.

O-De-Ac: Deacetylutessine

[102487-20-7]

C₁₇H₁₉NO₅ 317.341


Alkaloid from the bulbs of *Sternbergia lutea* (Amaryllidaceae). Oil. [α]_D²⁵ -42.9 (c, 0.35 in CHCl₃).

Evidente, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 90 (isol, uv, ir, pmr, cmr, ms, struct, deriv)

Lutinine

L-311

4-(Phenylamino)-2-(phenylimino)-1(2H)-naphthalenone, 9CI [66855-48-9]

C₂₂H₁₆N₂O 324.381


Alkaloid from *Reseda luteola* (Reseda-ceae). Mp 177-179°.

Lutfullin, K.L. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 826; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 696 (ir, ms, isol)

Luzopeptin C

L-312

BBM 928C. Antibiotic BBM 928C [76110-01-5]

C₆₀H₇₄N₁₄O₂₂ 1343.325

Depsipeptide antibiotic. Dimer with twofold symmetry. Isol. from *Actinomadura luzonensis*. Active against gram-positive bacteria. Cytotoxic, inhibits HIV-1 reverse transcriptase. Sol. CHCl₃, CH₂Cl₂; fairly sol. C₆H₆, MeOH, butanol; poorly sol. H₂O, hexane. Mp 244-248°. [α]_D²⁵ -91 (c, 1 in CHCl₃). λ_{max} 230 (ε 80400); 256 (ε 109000); 330 (ε 16800); 383 (ε 16700) (EtOH/NaOH) (Derep). λ_{max} 235 (ε 83600); 264 (ε 59200); 345 (ε 23500) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) .81 mg/kg. CB9267000

Mono-O-Ac: Luzopeptin B. BBM 928B. Antibiotic BBM 928B [76149-24-1]

C₆₂H₇₆N₁₄O₂₃ 1385.362

From *Actinomadura luzonensis*. Active against gram-positive bacteria and tumours. Cytotoxic, inhibits HIV-1 reverse transcriptase. Sol. CHCl₃, CH₂Cl₂; fairly sol. C₆H₆, butanol, MeOH; poorly sol. H₂O, hexane. Mp 214-217°. [α]_D²⁵ -74 (c, 1 in CHCl₃). Acetylated at one of the alicyclic OH groups. λ_{max} 230 (ε 80400); 256 (ε 109000); 330 (ε 16800); 383 (ε 16700) (EtOH/NaOH) (Derep). λ_{max} 235 (ε 83600); 264 (ε 59200); 345 (ε 23500) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) .18 mg/kg. CB9266000

Di-O-Ac: Luzopeptin A. BBM 928A. Antibiotic BBM 928A [75580-37-9]

C₆₄H₇₈N₁₄O₂₄ 1427.4

From *Actinomadura luzonensis*. Active against gram-positive bacteria and tumours. Cytotoxic, inhibits HIV-1 reverse transcriptase. Mp 246-248°. [α]_D²⁵ -27 (c, 1 in CHCl₃). Acetylated at both alicyclic OH groups. λ_{max} 230 (ε 80400); 256 (ε 109000); 330 (ε 16800); 383 (ε 16700) (EtOH/NaOH) (Derep). λ_{max} 235 (ε 83600); 264 (ε 59200); 345 (ε 23500) (EtOH) (Derep).

▶ CB9391300

Dideoxy: Luzopeptin E₂. Dideoxyluzopeptin C [93079-89-1]

C₆₀H₇₄N₁₄O₂₀ 1311.326

Isol. from *Actinomadura luzonensis* ATCC31491. Active against gram-positive and acid-fast bacteria. Shows antitumour activity. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Lacks the two alicyclic OH groups. λ_{max} 230 (ε 80400); 256 (ε 109000); 330 (ε 16800); 383 (ε 16700) (EtOH/NaOH) (Derep). λ_{max} 235 (ε 83600); 264 (ε 59200); 345 (ε 23500) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) .034 mg/kg.

Ohkuma, H. *et al.*, *J. Antibiot.*, 1980, **33**, 1087

Tomita, K. *et al.*, *J. Antibiot.*, 1980, **33**, 1098

Konishi, M. *et al.*, *J.A.C.S.*, 1981, **103**, 1241

Rose, W.C. *et al.*, *Cancer Res.*, 1983, **43**, 1504

(props)

Netherlands Pat., 1984, 84 00 237; *CA*, **102**,

22795 (*Luzopeptin E₂*)

Ciufolini, M.A. *et al.*, *Tet. Lett.*, 1989, **30**, 3027

(synth)

Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 1098-

1099; 11375-11383; 2001, **123**, 561-568

(synth, activity)


Valognes, D. *et al.*, *Tet. Lett.*, 2001, **42**, 1907-

1909 (synth)

Lyadine

L-313

[52811-49-1]

C₂₁H₂₁N₃O₃ 363.415

Struct. in doubt (2004). Alkaloid from the roots of *Pauridiantha lyalli* (Rubiaceae). Cryst. (Et₂O). Mp 158-160°. [α]_D +335 (c, 1 in CHCl₃).


Levesque, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **278**, 959-961 (isol, uv, ir, pmr, ms)

Bennasar, M.-L. *et al.*, *J.O.C.*, 2004, **69**, 752-756 (synth, struct)

Lyalidine

L-314

[55856-50-3]



C₂₁H₁₉N₃O₃ 361.399
Minor alkaloid from the roots of *Pauridiantha lyalli* (Rubiaceae). Cryst. (Et₂O/MeOH). Mp 190-191°. [α]_D 0 (c, 1 in CHCl₃).

19,20-Dihydro, 19ξ-hydroxy: **Hydroxy-lyalidine**


[55856-51-4]
C₂₁H₂₁N₃O₄ 379.415

Minor alkaloid from roots of *Pauridiantha lyalli* (Rubiaceae). Noncryst.

Levesque, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **279**, 1053 (uv, ir, pmr, ms, struct)

Lyaline**L-315**

[52811-48-0]



C₂₁H₁₉N₃O₂ 345.4

Struct. in doubt (2004). Alkaloid from root bark of *Pauridiantha lyalli* (Rubiaceae). Cryst. (Et₂O). Mp 280-282°. [α]_D +650 (c, 1 in CHCl₃).


18,19-Dihydro: Mp 229-230°. [α]_D +600 (c, 1 in CHCl₃).

Levesque, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **278**, 959-961 (isol, uv, ir, ms, pmr)

Bennasar, M.-L. et al., *J.O.C.*, 2004, **69**, 752-756 (synth, struct)

Lyaloside**L-316**

[56021-85-3]



Absolute configuration

C₂₇H₃₀N₂O₉ 526.542

Alkaloid from *Palicourea adusta* and *Pauridiantha lyalli* (Rubiaceae). Cryst. (MeOH). Mp 168-169°. [α]_D²⁰ -202 (c, 1 in MeOH). λ_{max} 215 (ε 9425); 236 (ε 12300); 291 (ε 5660); 351 (ε 4510) (MeOH).

Tetra-O-Ac: Mp 156-157°. [α]_D²⁰ -78 (c, 1 in CHCl₃).

6'-(4-Hydroxy-3-methoxycinnamoyl) (E-): **6'-trans-Feruloyllyaloside**

[83404-71-1]

C₃₇H₃₈N₂O₁₂ 702.713

Alkaloid from the leaves of *Palicourea adusta* and *Pauridiantha lyalli*. λ_{max} 238 (ε 26990); 292 (ε 13260); 330 (ε 10360) (MeOH).

6'-(4-Hydroxy-3,5-dimethoxycinnamoyl) (E-): **6'-trans-Sinapoyllyaloside**

[83404-73-3]

C₃₈H₄₀N₂O₁₃ 732.74

Alkaloid from the leaves of *Palicourea adusta* and *Pauridiantha lyalli*. λ_{max}

206 (ε 57350); 238 (ε 76100); 292 (ε 28680); 335 (ε 27540) (MeOH).

N^b-Me: **3,4,5,6-Tetrahydrodolichantoside**

C₂₈H₃₃N₂O₉[⊕] 541.577

Quaternary alkaloid from *Strychnos mellodora*. Amorph. brown-yellow powder. Counterion not specified. λ_{max} 209 (log ε 3.96); 254 (log ε 3.69); 311 (log ε 3.52); 375 (log ε 2.93) (EtOH).

Parent acid: **Lyalosidic acid**

[104821-27-4]

C₂₆H₂₈N₂O₉ 512.515

Alkaloid from *Ophiorrhiza japonica* (Rubiaceae). Amorph. powder. [α]_D -151.7 (MeOH).

Parent acid, N^b-Me: **3,4,5,6-Tetrahydrodripalicoside**

C₂₇H₃₁N₂O₉[⊕] 527.55

Quaternary alkaloid from *Strychnos mellodora*. Amorph. brown-yellow powder. Counterion not specified. Presumably a zwitterion. λ_{max} 222 (log ε 3.97); 255 (log ε 3.06); 311 (log ε 2.85); 373 (log ε 2.31) (EtOH).

10-Hydroxy, parent acid: **10-Hydroxylyalosidic acid**

[104821-28-5]

C₂₆H₂₈N₂O₁₀ 528.515

Alkaloid from *Ophiorrhiza japonica* (Rubiaceae). Pale-yellow prisms. Mp 300°. [α]_D -127.7 (MeOH).

Levesque, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1975, **280**, 593-595 (uv, ir, ms, pmr, struct)

Levesque, J. et al., *Tetrahedron*, 1982, **38**, 1417-1424 (*Feruloyllyaloside*, *Sinapoyllyaloside*)


Aimi, N. et al., *Chem. Pharm. Bull.*, 1986, **34**, 3064-3066; 1992, **40**, 2588-2596 (*Lyalosidic acid*, *10-Hydroxylyalosidic acid*, synth, abs config)

Brandt, V. et al., *Phytochemistry*, 1999, **51**, 1171-1176 (*Tetrahydrodolichantoside*, *Tetrahydrodripalicoside*)

Valverde, J. et al., *Phytochemistry*, 1999, **52**, 1485-1489 (isol, uv, cd, pmr, cmr, ms)

Lycaconitine**L-317**

[25867-19-0]



C₃₆H₄₈N₂O₁₀ 668.783

Alkaloid from *Aconitum gigas*, *Aconitum lycoctonum*, *Aconitum umbrosum*, *Aconitum barbatum* and *Delphinium cashmirianum* (Ranunculaceae). Cryst. (MeOH). Mp 113°. [α]_D²⁷ +16 (c, 0.64 in EtOH).

N-De-Et, N-formyl, N-Deethyl-N-formyllycaconitine, N-Deethyllycaconitine-N-aldehyde

[460731-53-7]

C₃₅H₄₄N₂O₁₁ 668.739

Alkaloid from the roots of *Aconitum leave*. Amorph. [α]_D²⁵ +22.5 (c, 0.1 in CHCl₃). λ_{max} 230 (log ε 4.1); 280 (log ε 2.1) (MeOH).

O⁸-Me: **8-O-Methyllycaconitine**. *Oreacoinine*

[144049-69-4]

C₃₇H₅₀N₂O₁₀ 682.809

Alkaloid from the roots of *Aconitum leave*, *Aconitum septentrionale* and *Aconitum orientale*. Amorph. [α]_D +22.6 (c, 0.32 in CHCl₃).

O¹-De-Me: **Albiviolaconitine C**

[138822-62-5]

C₃₅H₄₆N₂O₁₀ 654.756

Alkaloid from roots of *Aconitum albiviolaaceum* (Ranunculaceae). [α]_D²⁸ +57.5 (c, 1.4 in CHCl₃).

O⁶-De-Me, 8-Ac: **Potanidine B**

[161068-59-3]

C₃₇H₄₈N₂O₁₁ 696.793

Alkaloid from the roots of *Delphinium potaninii*. Amorph. powder. [α]_D²⁴ +28.6 (c, 0.07 in CHCl₃).

O¹⁴-De-Me: **14-O-Demethyllycaconitine**

[460731-52-6]

C₃₅H₄₆N₂O₁₀ 654.756

Alkaloid from the roots of *Aconitum leave*. Amorph. [α]_D²⁵ +20 (c, 0.1 in CHCl₃). λ_{max} 225 (log ε 4.1); 278 (log ε 2.05) (MeOH).

O⁶, O¹⁴-Di-de-Me, O⁶-Ac: **Albiviolaconitine B**

[138822-61-4]

C₃₆H₄₆N₂O₁₁ 682.766

Alkaloid from roots of *Aconitum albiviolaaceum* (Ranunculaceae). [α]_D²⁸ +24.6 (c, 0.71 in CHCl₃).

Kuzovkov, A.D. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 2746-2749 (struct)

Pelletier, S.W. et al., *J. Nat. Prod.*, 1979, **42**, 615-623 (isol, ir, pmr)

Pelletier, S.W. et al., *J.A.C.S.*, 1981, **103**, 6536-6538 (config)

Edwards, O.E. et al., *Can. J. Chem.*, 1982, **60**, 2661-2667 (config)

Chen, D. et al., *Huaxue Xuebao*, 1992, **50**, 1211-1218; *CA*, **118**, 230146 (*Albiviolaconitines*)

Ross, S.A. et al., *Tetrahedron*, 1992, **48**, 1183-1192 (*8-O-Methyllycaconitine*)

Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1994, **30**, 829-831; *Chem. Nat. Compd. (Engl. Transl.)*, 1994, **30**, 776-777 (*Oreacoinine*)

Pu, H. et al., *Yaoxue Xuebao*, 1994, **29**, 689-692; *CA*, **122**, 122898 (*Potanidine B*)

Jacyno, J.M. et al., *J. Nat. Prod.*, 1996, **59**, 707-709 (synth, ir, pmr, cmr)

Ulubelen, A. et al., *Pharmazie*, 2002, **57**, 427-429 (*Aconitum leave alkaloids*)

Lyceamine**L-318**

[11055-27-9]

C₁₉H₄₂N₄O₇ 438.563

Struct. unknown. Alkaloid from *Lycium europaeum* (Solanaceae). Cryst. + 4H₂O (MeOH/Me₂CO). Mp 280-281°. [α]_D³¹ +1.5 (c, 0.79 in EtOH).


Hydrobromide: Mp 218-221°.

Picrate: Mp 180-181°.

Manzoor-i-Kuda, et al., *Pak. J. Sci. Ind. Res.*, 1968, **11**, 247-249; *CA*, **71**, 777z

Lyciumins

L-319



Lyciumin A [125708-06-7]

C₄₂H₅₁N₉O₁₂ 873.918
 Isol. from root bark of *Lycium chinense* (Chinese boxthorn) and from *Lycium barbarum*. Shows anti-ACE and antiretinin activity. Powder. [α]_D²⁵ +10.1. λ_{max} 273; 281; 291 (solvent not reported) (Derep).

Lyciumin B [125756-66-3]

C₄₄H₅₂N₁₀O₁₁ 896.955
 Isol. from *Lycium chinense* (Chinese boxthorn) and from *Lycium barbarum*. Shows anti-ACE and antiretinin activity. Powder. [α]_D²⁵ +3.5. λ_{max} 281; 291 (solvent not reported) (Derep).

Lyciumin C [150394-23-3]

C₄₉H₅₇N₉O₁₂ 964.042
 Isol. from *Lycium chinense* (Chinese boxthorn) and from *Lycium barbarum*. Powder. Sol. MeOH, CHCl₃; poorly sol. hexane, H₂O. [α]_D²⁴ -11.9 (c, 0.97 in DMSO). λ_{max} 230; 235; 292; 418; 438 (MeOH) (Berdy).

Lyciumin D [150415-40-0]

C₄₅H₅₇N₉O₁₁ 899.999
 Isol. from *Lycium chinense* (Chinese boxthorn) and from *Lycium barbarum*. Powder. [α]_D²⁵ -8.4 (c, 0.45 in DMSO).
 Yahara, S. *et al.*, *Tet. Lett.*, 1989, **30**, 6041 (*isol*)
 Schmidt, U. *et al.*, *Chem. Comm.*, 1992, 1353 (*synth*)
 Yahara, S. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 703 (*isol*, *pmr*, *cmr*, *struct*)
 Morita, H. *et al.*, *Tetrahedron*, 1996, **52**, 2795 (*pmr*, *conformn*, *config*)

Lycocaroline


L-320

C₃₂H₅₃N₃O 495.79
 Struct. unknown. Alkaloid from *Lycopodium carolinianum* (Lycopodiaceae).
Didehydro: Dehydrolycocaroline
 C₃₂H₅₁N₃O 493.774
 Alkaloid from *Lycopodium carolinianum* (Lycopodiaceae).
 MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Lycoceruinine

L-321

Alkaloid L33
 [6871-55-2]



C₁₆H₂₆N₂O₂ 278.394
 Rel. config. shown. CAS numbering. Alkaloid from *Lycopodium cernuum*, *Lycopodium carolinianum* and *Lycopodium inundatum* (Lycopodiaceae). Mp 230-231°. [α]_D²⁵ -24.5 (c, 0.1 in MeOH).
N¹³-Oxide: Lycoceruinine N-oxide
 [760947-80-6]
 C₁₆H₂₆N₂O₃ 294.393
 Alkaloid from *Lycopodium cernuum*. Amorph. solid. [α]_D²² -23 (c, 1 in MeOH).

Methodide: Mp 270-272°.

4,5-Didehydro: **Carolinianine**

[36101-39-0]
 C₁₆H₂₄N₂O₂ 276.378
 Alkaloid from *Lycopodium carolinianum* var. *affine* (Lycopodiaceae). Cryst. (Me₂CO). Mp 198.5-201.5°.

4,5-Didehydro, **Ac:** Mp 141-144°.

Deoxo: **Dihydrodesoxylycoceruinine**

[54307-43-6]
 C₁₆H₂₈N₂O 264.41
 Alkaloid from *Lycopodium carolinianum* var. *affine* (Lycopodiaceae).

Deoxy: **Ceruinine. Alkaloid L32**

[6880-84-8]
 C₁₆H₂₆N₂O 262.394
 Alkaloid from *Lycopodium cernuum* and *Lycopodium carolinianum* (Lycopodiaceae). Mp 103-104°. [α]_D²⁵ -20.5 (c, 0.1 in MeOH).

Deoxy, N¹³-oxide: **Ceruinine N-oxide**

[760947-79-3]
 C₁₆H₂₆N₂O₂ 278.394
 Alkaloid from *Lycopodium cernuum*. Amorph. solid. [α]_D²⁷ -18 (c, 1 in MeOH).

Deoxy, **methodide:** Mp 237-238°.

Deoxy, 3,3a-didehydro: **Anhydrolycoceruinine**

[34331-32-3]
 C₁₆H₂₄N₂O 260.378
 Alkaloid from *Lycopodium inundatum* and *Lycopodium carolinianum* var. *affine* (Lycopodiaceae). Mp 140-142°.

Deoxy, deoxo: **Dihydrodesoxylycoceruinine**


[15216-33-8]
 C₁₆H₂₈N₂ 248.411
 Alkaloid from *Lycopodium carolinianum* var. *affine* (Lycopodiaceae). Mp 64-65°.

Marion, L. *et al.*, *Can. J. Res., Sect. B*, 1948, **26**, 1 (*isol*)
 Ayer, W.A. *et al.*, *Can. J. Chem.*, 1967, **45**, 433; 445; 451 (*ir*, *pmr*, *ms*, *struct*)
 Miller, N. *et al.*, *Bull. Soc. Chim. Belg.*, 1971, **80**, 629; *CA*, **76**, 96981p (*Carolinianine*)
 Ho, Y.K. *et al.*, *Can. J. Chem.*, 1971, **49**, 3352 (*biosynth*)
 Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1971, **91**, 980; *CA*, **75**, 148492f (*Anhydrolycoceruinine*)
 Braekman, J.C. *et al.*, *Phytochemistry*, 1974, **13**, 2519-2528 (*Dihydrodesoxylycoceruinine*, *Anhydrolycoceruinine*, *Dihydrodesoxylycoceruinine*)
 Morita, H. *et al.*, *Tetrahedron*, 2004, **60**, 7015-7023 (*pmr*, *cmr*, *N-oxides*)
 Nishikawa, Y. *et al.*, *Org. Lett.*, 2008, **10**, 1987-1990 (*Ceruinine*, *synth*)

Lycoclavine

L-322

[6900-91-0]



Absolute Configuration

C₁₈H₂₉NO₃ 307.432
 Alkaloid from *Lycopodium clavatum* var. *megastachyon*, *Lycopodium alpinum*, *Lycopodium gnidioides* and *Lycopodium paniculatum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 212-213° (201-203°). [α]_D²⁵ -9 (95% EtOH).

Hydrochloride: Mp 281-285°.

Perchlorate: Mp 283-286°.

Ac: Acetyllycoclavine

[73907-66-1]
 C₂₀H₃₁NO₄ 349.469
 Alkaloid from *Lycopodium clavatum* var. *megastachyon* (Lycopodiaceae). Cryst. (petrol). Mp 144-145°.

O-De-Ac: **Deacetyllycoclavine. Alkaloid P4**

[23627-70-5]
 C₁₆H₂₇NO₂ 265.395
 Alkaloid from *Lycopodium paniculatum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 217°.

O-De-Ac, 5-O-[3-(4-hydroxy-3-methoxyphenyl)propanoyl], 6-Ac: **Lycoposerramine O**

[677723-23-8]
 C₂₈H₃₉NO₆ 485.619
 Alkaloid from *Lycopodium serratum*. Amorph. powder. [α]_D²³ -27.8 (c, 0.06 in CHCl₃).

O-De-Ac, O⁵-[3-(3,4-dimethoxyphenyl)propanoyl]: **Lycognidine**

[62023-84-1]
 C₂₇H₃₉NO₅ 457.609
 Alkaloid from *Lycopodium gnidioides* (Lycopodiaceae). Oil.

O-De-Ac, 5-O-(3,4-dimethoxycinnamoyl), 6-Ac: **Lyconesidine C**

[462065-75-4]
 C₂₉H₃₉NO₆ 497.63
 Alkaloid from *Lycopodium chinense*. Amorph. solid. [α]_D²⁵ -3 (c, 1 in MeOH). λ_{max} 204 (ε 16400); 217 (ε 14100); 236 (ε 12100); 300 (sh) (ε 13900); 327 (ε 18400) (MeOH).

5-Ketone, O-de-Ac: **6α-Hydroxylycoclavine**

dine. Alkaloid L20
 [21061-92-7]
 C₁₆H₂₅NO₂ 263.379
 Alkaloid from *Lycopodium lucidulum*, *Lycopodium selago*, *Lycopodium serratum* var. *longipetiolatum* (Lycopodiaceae). Cryst. (MeOH or Me₂CO). Mp 258-259° (250-254°).

5-Ketone, O-de-Ac, 6-Ac: Needles (petrol). Mp 143-144°. Formed at -10°.

8,15-Didehydro, O-de-Ac: **Lycoposerramine J. Miyoshianine B**

[623945-30-2]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Huperzia miyoshiana* and *Lycopodium serratum*. Prisms (Me₂CO/MeOH). Mp 264-266°. [α]_D²⁵ -117 (c, 0.25 in MeOH).

8,15-Didehydro, 5-ketone, O-de-Ac: **Lycoposerramine H**

[677723-08-9]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum*. Prisms (MeOH). Mp 227-228° subl.

6-Epimer, 5-ketone, O-de-Ac: **Lycoposerramine L**. 6β-Hydroxylycopodine

[677723-14-7]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium serratum*. Amorph. powder.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1962, **40**, 2088-2100; 1963, **41**, 649-657 (*Lycoclavine*, *Acetylylcochlorine*, *6-Hydroxylycopodine*)

Miller, N. *et al.*, *Phytochemistry*, 1971, **10**, 1931-1934 (*isol*)

Nyembo, L. *et al.*, *Bull. Soc. Chim. Belg.*, 1976, **85**, 595-604; *CA*, **86**, 90108u (*Lycognidine*)

Castillo, M. *et al.*, *Can. J. Chem.*, 1976, **54**, 2900-2908 (*Deacetylylcochlorine*)

Morales, G. *et al.*, *Phytochemistry*, 1979, **18**, 1719-1720 (*Deacetylylcochlorine*)

Muñoz, O.M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 200-203 (*pmr*, *Deacetylylcochlorine*)

Sun, C.M. *et al.*, *Planta Med.*, 1993, **59**, 467-471 (*6-Hydroxylycopodine*)

Hirasawa, Y. *et al.*, *Tetrahedron*, 2002, **58**, 5483-5488 (*Lycosidine C*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1163-1169 (*Lycoposerramines*)

Tong, X.-T. *et al.*, *Planta Med.*, 2003, **69**, 576-579 (*Miyoshianine B*)

Lycodiflexine

L-323

C₃₅H₅₀N₂O₄ 562.791


Struct. unknown. Alkaloid from *Lycopodium clavatum* var. *borbonicum* (Lycopodiaceae).

MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241

Lycodine

L-324

[20316-18-1]



Absolute Configuration

C₁₆H₂₂N₂ 242.363

Alkaloid from *Lycopodium annotinum*, *Lycopodium clavatum*, *Lycopodium fawcettii*, *Lycopodium obscurum*, *Lycopodium flabelliforme*, *Lycopodium lucidulum*, *Lycopodium serratum* var. *thunbergii* and *Lycopodium magellanicum* (Lycopodiaceae). Mp 118°. [α]_D -10 (c, 1 in EtOH).

Dipicrate: Mp 229-233° dec.

N-Ac, picrate: Mp 180-182.5°.

N-Me: **N-Methyllycodine**

[54301-02-9]

C₁₇H₂₄N₂ 256.39

Alkaloid from *Lycopodium complanatum*, *Lycopodium erythraeum* and *Lycopodium magellanicum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 91-92° (87-88°).

11R-Hydroxy: **11-Hydroxylycodine**C₁₆H₂₂N₂O 258.363

Alkaloid from *Lycopodium complanatum*. Cytotoxic. Amorph. solid. [α]_D -47 (c, 0.5 in MeOH). λ_{max} 269 (ε 3500) (MeOH).

Anet, F.A.L. *et al.*, *Can. J. Chem.*, 1958, **36**, 902 (*isol, ir, uv, pmr*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1960, **38**, 1823 (*N-Methyllycodine*)

Anet, F.A.L. *et al.*, *Tet. Lett.*, 1960, **No. 20**, 9 (*struct*)

Nakashima, T.T. *et al.*, *Can. J. Chem.*, 1975, **53**, 1936 (*cmr*)

Loyola, L.A. *et al.*, *Phytochemistry*, 1979, **18**, 1721 (*N-Methyllycodine*)

Kleinman, E. *et al.*, *Tet. Lett.*, 1979, 4125 (*synth*)

Heathcock, C.H. *et al.*, *J.A.C.S.*, 1982, **104**, 1054 (*synth*)


Kobayashi, J. *et al.*, *J.O.C.*, 2001, **64**, 5901-5904 (*11-Hydroxylycodine*)

Lycodoline

L-325

Alkaloid L8. Alkaloid L30

[6900-92-1]



Absolute Configuration

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium alopecuroides*, *Lycopodium annotinum*, *Lycopodium clavatum* various varieties, *Lycopodium carolinum*, *Lycopodium fawcettii*, *Lycopodium inundatum*, *Lycopodium lucidulum*, *Lycopodium phlegmaria*, *Lycopodium saururus*, *Lycopodium selago* and *Lycopodium serratum* various forms. Mp 180°. Alkaloid L35 was a mixt. of Lycodoline with Lycopodine, L-341 and Flabelliformine, F-73.

Perchlorate:

Cryst. (MeOH/EtOAc). Mp 318° dec.

N-Oxide: **Lycodoline N-oxide. Obscuramine A**

[849096-85-1]

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium obscurum*. Amorph. solid. [α]_D²⁵ -44 (c, 0.1 in MeOH).

6α-Hydroxy: **Serratezomine C**. 6-Hydroxylycodoline

[301677-01-0]

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium serratum* var. *serratum*. Amorph. solid. [α]_D -8 (c, 0.3 in MeOH).

6α-Acetoxy: **Lycoposerramine N**

[677723-20-5]

C₁₈H₂₇NO₄ 321.416

Alkaloid from *Lycopodium serratum*. Amorph. powder.

12-Epimer: **12-Epilycodoline**. *Isolycodoline*. *Pseudoselagine*. *12-Hydroxylycopodine*. Alkaloid L23. Alkaloid L25

[21061-90-5]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium lucidulum* and *Lycopodium selago* (Lycopodiaceae). Cryst. (Me₂CO). Mp 163-164°. [α]_D -43 (c, 0.13 in EtOH). Alkaloid L22 was a mixt. of Isolycodoline and Lycopodine, L-341.

12-Epimer, perchlorate:

Prisms (MeOH/EtOH). Mp 300°.

12-Epimer, N-oxide: **12-Epilycodoline N-oxide**

[791588-37-9]

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Huperzia serrata*. Prisms (Me₂CO). Mp 238-240°. [α]_D²⁵ -21.2 (c, 1.3 in CHCl₃).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 92-96; 1946, **24**, 57-62 (*isol, epimer*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1964, **42**, 2514-2522; 1969, **47**, 449-455; 1990, **68**, 1300-1304 (*isol, ir, ms, pmr, struct, 12-Epilycodoline*)

Nakashima, T.T. *et al.*, *Can. J. Chem.*, 1975, **53**, 1936-1942 (*cmr*)

Heathcock, C.H. *et al.*, *J.A.C.S.*, 1982, **104**, 1054-1068 (*synth*)

Morita, H. *et al.*, *J.O.C.*, 2000, **65**, 6241-6245 (*Serratezomine C*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1163-1169 (*Lycoposerramine N*)


Tan, C.-H. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1963-1967 (*12-Epilycodoline N-oxide*)

Morita, H. *et al.*, *Tetrahedron*, 2005, **61**, 1955-1960 (*Obscuramine A*)

Lycofawcine

L-326

[3175-90-4]

C₁₈H₂₉NO₄ 323.431

Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Oil.

Perchlorate: Mp 290-294° dec.

Methiodide:

Cryst. (Me₂CO). Mp 281-282°.

8-Ac: **Acetyllycofawcine**C₂₀H₃₁NO₅ 365.469

Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Mp 181-182°.


Burnell, R.H. *et al.*, *Can. J. Chem.*, 1960, **38**, 1927 (*Acetyllycofawcine*)

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1963, **41**, 3091 (*isol, ir*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1965, **43**, 328 (ms. struct)

Lycloflexine**L-327**

Lycobergine
[52485-00-4]



Absolute
Configuration

$C_{17}H_{25}NO_2$ 275.39

Alkaloid from *Lycopodium clavatum* var. *inflexum*, *Lycopodium clavatum* var. *borbonicum*, *Lycopodium carolinum*, *Lycopodium inundatum*, *Lycopodium phlegmaria* and *Lycopodium serratum* (Lycopodiaceae). Mp 130-131°.

Hydrochloride: Mp 225-226° dec.

Methiodide: Mp 243-244°.

2S-Hydroxy: Lycoposerramine D

[481048-29-7]

$C_{17}H_{25}NO_3$ 291.389

Alkaloid from *Lycopodium serratum*. Prisms (EtOAc/hexane). Mp 173-176°. λ_{max} 235 ; 310 (EtOH).

8R-Hydroxy: Lycoposerramine U

[481048-31-1]

$C_{17}H_{25}NO_3$ 291.389


Alkaloid from *Lycopodium serratum*. Amorph. powder. λ_{max} 237 (sh) ; 306 (sh) (EtOH).

Ayer, W.A. *et al.*, *Tet. Lett.*, 1973, 5045-5048 (isol, ir, pmr, ms, cryst struct)

Takayama, H. *et al.*, *Tet. Lett.*, 2002, **43**, 8307-8311 (cmr, Lycoposerramine D, Lycoposerramine U)

Lycofoline**L-328**

11,12-Didehydrolycopodane-5,8-diol, 9CI
[7163-08-8]



$C_{16}H_{25}NO_2$ 263.379

Alkaloid from *Lycopodium annotinum* and *Lycopodium obscurum* (Lycopodiaceae). Prisms (Et₂O). Mp 144-145°. $[\alpha]_D^{20}$ -75 (c, 2.0 in EtOH).

Hydrobromide:

Cryst. (MeOH/Et₂O). Mp 274-275°.

Methiodide:

Cryst. (MeOH). Mp 263-264°.

O⁵-Ac: Acetyllycofoline

$C_{18}H_{27}NO_3$ 305.416

Alkaloid from *Lycopodium fawcettii*. Cryst. (H₂O)(as perchlorate). Mp 280-

282° (perchlorate). $[\alpha]_D$ -61 (c, 1.10 in EtOH).

O⁸-Ac: By acetylation of Lycofoline. Mp 180-182° (159-160°). $[\alpha]_D$ -42.8 (c, 1.29 in EtOH).

Di-Ac: Diacetyllycofoline

$C_{20}H_{29}NO_4$ 347.453

Alkaloid from *Lycopodium fawcettii* (Lycopodiaceae). Cryst. (Et₂O). Mp 113-118°.

5-Ketone: Gnidiodine

[62023-83-0]

$C_{16}H_{23}NO_2$ 261.363

Alkaloid from *Lycopodium gnidioides* and *Lycopodium phlegmaria* (Lycopodiaceae). Oil which browns in air. $[\alpha]_D^{20}$ -20 (c, 0.43 in CHCl₃). Configs. of OH and CH₃ groups descr. as both equatorial, which appears to be as shown.

8-Ketone: Acrifoline†. Alkaloid L27. Alkaloid L29

[664-24-4]

[93412-90-9]

$C_{16}H_{23}NO_2$ 261.363

Alkaloid from *Lycopodium annotinum* and *Lycopodium selago* (Lycopodiaceae). Needles (Et₂O or hexane). Mp 103-104° (sinters at 95°). $[\alpha]_D^{20}$ -266.2 (c, 1 in CHCl₃). $[\alpha]_D^{20}$ -250 (c, 1 in EtOAc). $[\alpha]_D^{21}$ -264.8 (Me₂CO). Also isol. as 1:1 molecular complexes with other alkaloids; see Isolycopodine below and also Annotoxine in A-1062.

8-Ketone, hydroiodide: Mp 258-259° dec.**8-Ketone, perchlorate:**

Stout prisms (Me₂CO/EtOAc). Mp 266° (260°).

8-Ketone, 1:1 complex with Lycopodine,

L-341: Isolycopodine

$C_{32}H_{48}N_2O_3$ 508.743

Isol. from *Lycopodium annotinum* (Lycopodiaceae).

8-Ketone, 5-Ac: Acetylacrifoline. Alkaloid

L12

[6793-40-4]

$C_{18}H_{25}NO_3$ 303.4

Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Stout prisms (hexane). Mp 119°.

15-Epimer: Acrifolinol

$C_{16}H_{25}NO_2$ 263.379

Minor alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Cryst. (MeOH). Mp 191.1-192.6° (synthetic).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1943, **21**, 92 (*O*-Acetylacrifoline)

Manske, R.H.F. *et al.*, *J.A.C.S.*, 1947, **69**, 2126 (*Acrifoline*)

Bertho, A. *et al.*, *Chem. Ber.*, 1952, **85**, 663 (*Acrifoline*)

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1955, **29**, 509; 1956, **30**, 233 (*Acrifoline, isol*)

Perry, G.S. *et al.*, *Can. J. Chem.*, 1956, **34**, 1189 (*Acrifoline, Acrifolinol, synth*)

Anet, F.A.L. *et al.*, *Can. J. Chem.*, 1959, **37**, 1589; 1962, **40**, 236 (*Lycofoline*)

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1960, **38**, 1927 (*Acetyllycofoline, Diacetyllycofoline*)

French, W.N. *et al.*, *Can. J. Chem.*, 1961, **39**, 2100 (*Acrifoline, struct*)

Burnell, R.H. *et al.*, *Tetrahedron*, 1962, **18**, 1467 (*struct*)

Rodewald, W.J. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1967, **15**, 579; *CA*, **68**, 69166e (*Isolycopodine*)

Nyembo, L. *et al.*, *Bull. Soc. Chim. Belg.*, 1976, **85**, 595 (*Gnidiodine*)


Nyembo, L. *et al.*, *Can. J. Chem.*, 1978, **56**, 851 (*Gnidiodine*)

Wenkert, E. *et al.*, *Chem. Comm.*, 1984, 714 (*synth*)

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077; 1990, **68**, 1300 (*Acrifoline, Acrifolinol*)

Lycogalic acid**L-329**

3,4-Di-1H-indol-3-yl-1H-pyrrole-2,5-dicarboxylic acid, 9CI. *Chromopyrrolic acid* [150044-68-1]



$C_{22}H_{15}N_3O_4$ 385.378

Isol. from fruit bodies of the slime mould *Lycogala epidendrum*. Also prod. by *Chromobacterium violaceum* as a tryptophan metab. Cryst. Mp > 200°. Closely related to the Arcyriarubins and Arcyriaflavins (see for example Arcyriarubin A, A-1394).

Mono-Me ester:

$C_{23}H_{17}N_3O_4$ 399.405

Isol. from *Lycogala epidendrum*. Brown powder. λ_{max} 224 (ε 57000); 268 (ε 29000) (MeOH).

Di-Me ester: Lycogarubin C. Dimethyl lycogalate A

[150044-77-2]

$C_{24}H_{19}N_3O_4$ 413.432

Isol. from *Lycogala epidendrum*. Shows moderate anti-HSV-1 virus activity. Pale yellow needles (MeOH aq.). Sol. MeOH. Mp 126-127° dec. λ_{max} 226 (ε 58900); 270 (ε 33900) (EtOH) (Derep).

5'-Hydroxy, 1-Me ester:

$C_{23}H_{17}N_3O_5$ 415.404

Isol. from *Lycogala epidendrum*. Pale yellow powder. λ_{max} 225 (ε 38000); 268 (ε 22000) (MeOH).

5'-Hydroxy, di-Me ester: Lycogarubin B.

Dimethyl lycogalate B

[154071-67-7]

$C_{24}H_{19}N_3O_5$ 429.431

Isol. from *Lycogala epidendrum*. Sol. MeOH. λ_{max} 227 (ε 42700); 270 (ε 26900) (EtOH) (Derep).

5''-Hydroxy, 1-Me ester:


$C_{23}H_{17}N_3O_5$ 415.404

Isol. from *Lycogala epidendrum*. Pale yellow powder. λ_{max} 225 (ε 35000); 269 (ε 20000) (MeOH).

5',5''-Dihydroxy, di-Me ester: **Lycogarin A**. Dimethyl lycogalate C [154071-66-6]
 $C_{24}H_{19}N_3O_6$ 445.431
 Isol. from *Lycogala epidendrum*. λ_{max} 224 (ϵ 50100); 268 (ϵ 33900); 298 (sh) (ϵ 15800); 332 (sh) (ϵ 4270) (MeOH). λ_{max} 226 (ϵ 41700); 270 (ϵ 30200) (EtOH).

Hoshino, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 775-781 (Chromopyrrolic acid)
 Fröde, R. *et al.*, *Tet. Lett.*, 1994, **35**, 1689-1690 (Lycogalic acid)
 Hashimoto, T. *et al.*, *Tet. Lett.*, 1994, **35**, 2559-2560 (Lycogarubins)
 Fürtner, A. *et al.*, *Tetrahedron*, 2002, **58**, 6373-6380 (synth)
 Kamata, K. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 594-597 (mono-Me ester, 5'-hydroxy 1-Me ester, 5''-hydroxy 1-Me ester)
 Hinze, C. *et al.*, *Synthesis*, 2007, 608-612 (synth, di-Me ester)

Lyconadine A L-330
 [362512-63-8]




Relative Configuration

$C_{16}H_{20}N_2O$ 256.347
 Alkaloid from *Lycopodium complanatum*. Cytotoxic. Amorph. solid. $[\alpha]_D^{+14}$ (c, 0.35 in MeOH). λ_{max} 235 (ϵ 6300); 323 (ϵ 4800) (MeOH).

2,3-Dihydro: **Lyconadine B** [906477-40-5]
 $C_{16}H_{22}N_2O$ 258.363
 Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{+23}$ -66 (c, 0.5 in MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 2001, **64**, 5901-5904 (Lyconadine A)
 Ishiuchi, K. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 5995-6000 (Lyconadine B)
 Bisai, A. *et al.*, *J.A.C.S.*, 2008, **130**, 7222-7223 (synth)
 Beshore, D.C. *et al.*, *J.A.C.S.*, 2008, **130**, 13778-13789 (synth)

Lyconesidine A L-331
 [462065-73-2]




$C_{16}H_{25}NO_2$ 263.379
 Alkaloid from *Lycopodium chinense*. Solid. $[\alpha]_D^{-53}$ (c, 1 in MeOH).

16-Hydroxy: **Lyconesidine B** [462065-74-3]
 $C_{16}H_{25}NO_3$ 279.378
 Alkaloid from *Lycopodium chinense*. Needles (MeOH aq.). Mp 170-171°. $[\alpha]_D^{-71}$ (c, 1.8 in MeOH).

Hirasawa, Y. *et al.*, *Tetrahedron*, 2002, **58**, 5483-5488 (isol, pmr, cmr, cryst struct)


Lyconnotine L-332
 [6900-93-2]



$C_{17}H_{25}NO_3$ 291.389
 Alkaloid from *Lycopodium annotinum* (Lycopodiaceae). Mp 123°. $[\alpha]_D^{+20}$ +125.

Picrolonate: Mp 230-231°. Anet, F.A.L. *et al.*, *Tet. Lett.*, 1964, 751 (isol, uv, ir, ms, pmr, struct)


Lyconnotinol L-333
 [123145-46-0]



$C_{16}H_{25}NO_2$ 263.379
 Alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Off-white prisms (MeOH/Me₂CO). Mp 173-175°. $[\alpha]_D^{+103}$ (c, 0.43 in MeOH).

Di-Ac; methiodide: Cryst. (Me₂CO/Et₂O). Mp 176-177°. Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077 (isol, uv, ir, pmr, cmr, ms, struct)

Lycopaniculatine L-334
 Paniculatine† [58786-38-2]




$C_{17}H_{27}NO_2$ 277.406
 The name Lycopaniculatine is preferred since the name Paniculatine has also been used for an *Aconitum* alkaloid. See Paniculatine in H-181. Alkaloid from *Lycopodium paniculatum* (Lycopodiaceae). Mp 181°. $[\alpha]_D^{+66}$.

Hydrobromide: Mp 310°. Castillo, M. *et al.*, *Can. J. Chem.*, 1976, **54**, 2900 (isol, ir, pmr, ms, cryst struct)
 Sha, C.-K. *et al.*, *J.A.C.S.*, 1999, **121**, 9875-9876 (synth)

Kozaki, T. *et al.*, *J.O.C.*, 2007, **72**, 10147-10154 (synth)

Lycopecurine L-335
 [33293-82-2]



Absolute configuration

$C_{16}H_{25}NO$ 247.38
 Alkaloid from *Lycopodium alopecuroides* (Lycopodiaceae). Mp 239-241°.

Hydrobromide: Mp 305-307° dec. 5-Ketone: **Dehydrolycopecurine** [31147-30-5]
 $C_{16}H_{23}NO$ 245.364

Alkaloid from *Lycopodium inundatum* (Lycopodiaceae). Mp 57-59°.

2 α -Hydroxy: **Debenzoylalopecurine** $C_{16}H_{25}NO_2$ 263.379
 Alkaloid from *Lycopodium alopecuroides* (Lycopodiaceae). Cryst. (Me₂CO). Mp 230-232°.

2 α -Hydroxy; hydrobromide: [23247-02-1] Cryst. (Me₂CO/MeOH). Mp 249-251°.

2 α -Acetoxy: **Acetyldebenzoylalopecurine** [23247-04-3]
 $C_{18}H_{27}NO_3$ 305.416
 Alkaloid from *Lycopodium alopecuroides* (Lycopodiaceae). Cryst. (Me₂CO). Mp 238-240°.

2 α -Benzoyloxy: **Alopecurine** [23258-68-6]
 $C_{23}H_{29}NO_3$ 367.487
 Alkaloid from *Lycopodium alopecuroides* (Lycopodiaceae). Mp 244-245°. $[\alpha]_D^{-105}$ (c, 0.5 in MeOH).

2 α -Hydroxy, N-Me: **Lycopodatine A** [872610-90-7]
 $C_{17}H_{28}NO_2^{\oplus}$ 278.414
 Alkaloid from *Lycopodium inundatum*. Solid. $[\alpha]_D^{+33}$ -16 (c, 0.3 in MeOH). Counterion not specified.


2 β -Hydroxy, 5-ketone: **Inundatine** [34415-56-0]
 $C_{16}H_{23}NO_2$ 261.363
 Alkaloid from *Lycopodium inundatum* (Lycopodiaceae). Mp 174-177°.

2-Oxo: **Isoinundatine** [34324-37-3]
 $C_{16}H_{23}NO_2$ 261.363
 Alkaloid from *Lycopodium inundatum* (Lycopodiaceae). Mp 255°. 5-Config. not detd.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1968, **46**, 15; 1969, **47**, 2449; 1971, **49**, 524 (isol, ir, ms, cryst struct, Alopecurine, Lycopecurine)
 Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*, 1971, **80**, 83 (Dehydrolycopecurine, Inundatine, Isoinundatine)
 Morita, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1809-1812 (Lycopodatine A)


Lycoperine A

[873868-50-9]

Relative
ConfigurationC₃₁H₄₉N₃O₂ 495.747Alkaloid from *Lycopodium hamiltonii*. ACE inhibitor. Amorph. solid. $[\alpha]_D^{24}$ -238 (c, 0.1 in MeOH).Hirasawa, Y. *et al.*, *Org. Lett.*, 2006, **8**, 123-126 (*isol*, *pmr*, *cmr*)**Lycophlegmarine**


L-337

3,4,14,15-Tetradehydro-5-hydroxy-14-methoxy-17-methyl-4,17-secoserratinan-13-one, 9CI [80953-35-1]

Absolute
ConfigurationC₁₈H₂₇NO₃ 305.416Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Oil. $[\alpha]_D^{17}$ +258.5 (c, 0.94 in EtOH).4-Bromobenzoyl: Mp 140-142°. $[\alpha]_D^{17}$ +133.5 (c, 1.07 in EtOH).Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 3418 (*ir*, *pmr*, *cmr*, *cryst struct*)**Lycopladine A**


L-338

[886546-04-9]

Relative
ConfigurationC₁₆H₂₁NO₂ 259.347Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{23}$ +102 (c, 1 in MeOH). λ_{\max} 270 (ε 2800) (MeOH).Ishiuchi, K. *et al.*, *Tet. Lett.*, 2006, **47**, 3287-3289 (*isol*, *pmr*, *cmr*)**Lycopladine B**

L-339

[906477-37-0]


C₁₇H₂₃NO₃ 289.374Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{23}$ +202 (c, 0.2 in MeOH). λ_{\max} 315 (ε 1200) (MeOH).**Ac: Lycopladine C**

[906477-38-1]

C₁₉H₂₅NO₄ 331.411Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{23}$ +121 (c, 0.3 in MeOH). λ_{\max} 316 (ε 760) (MeOH).Ishiuchi, K. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 5995-6000 (*isol*, *pmr*, *cmr*)**Lycopladine D**


L-340

[906477-39-2]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Lycopodium complanatum*. Amorph. solid. $[\alpha]_D^{23}$ -17 (c, 1 in MeOH).Ishiuchi, K. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 5995-6000 (*isol*, *pmr*, *cmr*)**Lycopodine**

L-341

Alkaloid L24

C₁₆H₂₅NO 247.38**(-)-form** [466-61-5]

Alkaloid from a wide variety of *Lycopodium* spp. including *Lycopodium alopecuroides*, *Lycopodium alpinum*, *Lycopodium annotinum*, *Lycopodium annotinum* var. *acrifolium*, *Lycopodium carolinum*, *Lycopodium cernuum*, *Lycopodium clavatum*, *Lycopodium clavatum* var. *borbonicum*, *Lycopodium clavatum* var. *megastachyon*, *Lycopodium contiguum*, *Lycopodium densus*, *Lycopodium erythraeum*, *Lycopodium flabelliforme*, *Lycopodium inundatum*, *Lycopodium issleri*, *Lycopodium lucidulum*, *Lycopodium magellanicum*, *Lycopodium obscurum*, *Lycopodium obscurum* var. *dendroideum*, *Lycopodium paniculatum*, *Lycopodium phlegmaria*, *Lycopodium sabiniaefolium*, *Lycopodium saururus*, *Lycopodium selago*, *Lycopodium serratum*, *Lycopodium sitchense*, *Lycopodium thyoides*, *Lycopodium tristachyum* and *Lycopodium volubile* (Lycopodiaceae). Mp 116°. $[\alpha]_D^{26}$ -24.5 (c, 1.1 in EtOH). Alkaloid L28 was a mixt. with 5,15-Oxidolycopodane, O-163. Alkaloid L15 was a mixt. with Flabelliformine, F-73. Alkaloid L16 was a mixt. with Anhydrodihydrolycopodine (see below). ▶ LD₅₀ (mus, ivn) 27.58 mg/kg. OL2800600

Perchlorate: Mp 276°.

Oxime: Mp 262-264°.

5β-Alcohol: Dihydrolycopodine.

Complanatine†. Alkaloid L1. Alkaloid L26

[23768-74-3]

C₁₆H₂₇NO 249.395

Alkaloid from *Lycopodium carolinum*, *Lycopodium clavatum*, *Lycopodium clavatum* var. *megastachyon*, *Lycopodium clavatum* var. *borbonicum*, *Lycopodium clavatum* var. *inflexum*, *Lycopodium contiguum*, *Lycopodium flabelliforme*, *Lycopodium paniculatum*, *Lycopodium sabiniaefolium*, *Lycopodium saururus*, *Lycopodium thyoides* and *Lycopodium volubile* (Lycopodiaceae). Cryst. (Me₂CO/Et₂O). Mp 169°.

5β-Alcohol; perchlorate:Prisms (MeOH/Et₂O). Mp 223-224°. Forms monohydrate, Mp 194°.**5β-Alcohol, O-Ac: Acetyldihydrolycopodine.**

Alkaloid L2. Alkaloid L3

[52998-89-7]

C₁₈H₂₉NO₂ 291.433

Alkaloid from *Lycopodium clavatum*, *Lycopodium clavatum* var. *borbonicum*, *Lycopodium contiguum*, *Lycopodium flabelliforme*, *Lycopodium magellanicum*, *Lycopodium paniculatum* and *Lycopodium thyoides* (Lycopodiaceae). Mp 95-96°.

5β-Alcohol, O-Ac; perchlorate: Mp 246-247°.

8,15-Didehydro, 5β-alcohol: Selagoline

[790220-40-5]

C₁₆H₂₅NO 247.38

Alkaloid from the aerial parts of *Huperzia selago*. $[\alpha]_D^{25}$ -31 (c, 0.7 in CHCl₃).

Deoxo, 4,5-didehydro: Anhydrodihydrolycopodine.

Alkaloid L14

[54551-08-5]

C₁₆H₂₅N 231.38

Alkaloid from *Lycopodium tristachyum* (Lycopodiaceae).

Deoxo, 4,5-didehydro; perchlorate:

Stout prisms (MeOH/EtOAc). Mp 238°.

7-Hydroxy: 7-Hydroxylycopodine

[791588-38-0]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Huperzia serrata*. Amorph. powder. Mp 250-252°. $[\alpha]_D^{25}$ -29.2 (c, 0.37 in MeOH).

11α-Hydroxy: Lycoposerramine M.

11α-Hydroxylycopodine

[677723-17-0]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium serratum*. Amorph. solid.

11α-Hydroxy, 8,15-didehydro: Lycoposerramine I

[677723-10-3]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium serratum*. Amorph. powder.


(±)-form [18688-24-9]

Mp 130-131°.

Marion, L. *et al.*, *Can. J. Res., Sect. B*, 1944,


- 22, 1-4 (*Anhydrodihydrolycopodine*)
Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1946, **24**, 57-62 (*isol*)
Barclay, L.R.C. *et al.*, *Can. J. Chem.*, 1956, **34**, 1519-1527 (*isol*)
Harrison, W.A. *et al.*, *Can. J. Chem.*, 1961, **39**, 2086-2099 (*struct*)
Stork, G. *et al.*, *J.A.C.S.*, 1968, **90**, 1647-1648 (*synth*)
Ayer, W.A. *et al.*, *J.A.C.S.*, 1968, **90**, 1648-1650 (*synth*)
Marshall, W.D. *et al.*, *Can. J. Chem.*, 1975, **53**, 41-50 (*biosynth*)
Nakashima, T.T. *et al.*, *Can. J. Chem.*, 1975, **53**, 1936-1942 (*cmr*)
Ul-Haque, M. *et al.*, *J.C.S. Perkin 2*, 1975, 93-98 (*cryst struct, abs config*)
Kleinman, E. *et al.*, *Tet. Lett.*, 1979, 4125-4128 (*synth*)
Schumann, D. *et al.*, *Annalen*, 1982, 1700-1705 (*synth*)
Heathcock, C.H. *et al.*, *J.A.C.S.*, 1982, **104**, 1054-1068 (*synth*)
Wenkert, E. *et al.*, *Chem. Comm.*, 1984, 714-715 (*synth*)
MacLean, D.B. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 241-298 (*rev*)
Kraus, G.A. *et al.*, *J.A.C.S.*, 1985, **107**, 4341-4342 (*synth*)
Ayer, W.A. *et al.*, *Can. J. Chem.*, 1990, **68**, 1300-1304 (*Alkaloid L28, Alkaloid L15, Alkaloid L16*)
Muñoz, O.M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 200-203 (*pmr, cmr, Acetyldihydrolycopodine*)
Padwa, A. *et al.*, *J.O.C.*, 1997, **62**, 78-87 (*synth*)
Grieco, P.A. *et al.*, *J.A.C.S.*, 1998, **120**, 5128-5129 (*synth*)
Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1163-1169 (*Lycoposerramines*)
Tan, C.-H. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1963-1967 (*7-Hydroxylycopodine*)
Staerk, D. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 197-203 (*Selagoline*)
Yang, H. *et al.*, *J.A.C.S.*, 2008, **130**, 9238-9239 (*synth*)

Lycoposerramine A L-342
[392725-16-5]




- $C_{18}H_{29}N_3O_2$ 319.446
Alkaloid from *Lycopodium serratum*.
Needles (EtOAc). Mp 169-171°.
Takayama, H. *et al.*, *Org. Lett.*, 2001, **3**, 4165-4167 (*isol, ord, pmr, cmr*)

Lycoposerramine B L-343
[840489-33-0]




- $C_{17}H_{28}N_2O_2$ 292.42
Alkaloid from *Lycopodium serratum*.
Amorph. powder.
Katakawa, K. *et al.*, *J.O.C.*, 2005, **70**, 658-663 (*isol, cd, pmr, cmr*)

Lycoposerramine G L-344
4,12-Dihydroxylycopodine
[677723-05-6]




- $C_{16}H_{25}NO_3$ 279.378
Alkaloid from *Lycopodium serratum*.
Amorph. powder.
N-Oxide: **Lycoposerramine F. Miyoshianine A**
[623945-29-9]
 $C_{16}H_{25}NO_4$ 295.378
Alkaloid from *Huperzia miyoshiana* and *Lycopodium serratum*. Prisms (MeOH/EtOAc). Mp > 300° (Lycoposerramine G) Mp 216-218° (Miyoshianine A). $[\alpha]_D^{24}$ -15.2 (c, 0.06 in MeOH) (Lycoposerramine F). $[\alpha]_D^{25}$ -85 (c, 0.08 in MeOH) (Miyoshianine A).
Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1163-1169 (*isol, cd, pmr, cmr, cryst struct*)
Tong, X.-T. *et al.*, *Planta Med.*, 2003, **69**, 576-579 (*Miyoshianine A*)

Lycoposerramine S L-345
[480443-97-8]



- $C_{17}H_{30}N_2$ 262.437
Alkaloid from *Lycopodium serratum*.
Amorph. powder. $[\alpha]_D^{25}$ -37.8 (c, 0.3 in MeOH).
Takayama, H. *et al.*, *Tet. Lett.*, 2002, **43**, 8307-8311 (*isol, pmr, cmr*)


Lycoposerramine W L-346
[955937-63-0]



Absolute Configuration

- $C_{17}H_{26}N_2O$ 274.405
Related to Phlegmarine, P-361. Alkaloid from *Lycopodium serratum*. Amorph. powder. $[\alpha]_D^{25}$ +22.4 (c, 0.14 in $CHCl_3$).
Deoxy, N-de-Me: Lycoposerramine V
[955937-62-9]
 $C_{16}H_{24}N_2$ 244.379
Alkaloid from *Lycopodium serratum*. Amorph. powder. $[\alpha]_D^{23}$ +17.7 (c, 0.24 in $CHCl_3$).
Shigeyama, T. *et al.*, *Org. Lett.*, 2007, **9**, 4069-4072 (*isol, synth, pmr, cmr*)

Lycorenine L-347
[477-19-0]



- $C_{18}H_{23}NO_4$ 317.384
Alkaloid from *Lycoris radiata*, *Narcissus munozii-garmendiae* and many other spp. in the Amaryllidaceae. Mp 202°. $[\alpha]_D^{25}$ +125 (EtOH). $[\alpha]_D^{25}$ +180 ($CHCl_3$).
Hydrochloride: Mp 146-147° dec.
Me ether: O-Methyllycorenine
[87733-82-2]
 $C_{19}H_{25}NO_4$ 331.411
Isol. from the bulbs of *Lycoris radiata* and from whole plants of *Narcissus munozii-garmendiae* (Amaryllidaceae). Insect antifeedant. Mp 121-125°. $[\alpha]_D^{20}$ +164.1 (c, 0.53 in $CHCl_3$). $[\alpha]_D^{24}$ +198 (c, 1.04 in MeOH).

- Me ether, α-N-oxide: O-Methyllycorenine N-oxide**
[119308-27-9]
 $C_{19}H_{25}NO_5$ 347.41
Alkaloid from *Lapiedra martinezii* (Amaryllidaceae). Cryst. + H_2O . Mp 136°. $[\alpha]_D^{25}$ +86.8 (c, 0.1 in MeOH).

- Me ether, O⁹-de-Me: 9-O-Demethyl-7-O-methyllycorenine**
[955089-66-4]
 $C_{18}H_{23}NO_4$ 317.384
Alkaloid from *Hosta plantaginea*. Amorph. solid. $[\alpha]_D^{26}$ +122.6 (c, 0.65 in $CHCl_3$). λ_{max} 240 (log ε 3.69); 284 (log ε 3.47) ($CHCl_3$).

- Et ether, O⁹-de-Me: Eugenine. 9-O-De-methyl-7-O-ethyllycorenine**
[128562-43-6]
 $C_{19}H_{25}NO_4$ 331.411
Alkaloid from *Narcissus eugeniae* (Amaryllidaceae). Erroneous struct. diag. in paper.


- 7-Ketone (lactone): Homolycorenine. Narcipoetine**
[477-20-3]
 $C_{18}H_{21}NO_4$ 315.368
Alkaloid from *Narcissus poeticus* and from many other spp. in the Amaryllidaceae. Mp 175°. $[\alpha]_D$ +85 (95%)

- EtOH). λ_{\max} 226 (log ϵ 4.22); 230 (log ϵ 4.22); 270 (log ϵ 3.92); 306 (log ϵ 3.7) (EtOH).
- 7-Ketone (lactone), hydrochloride: Mp 285°.
- 7-Ketone (lactone), picrate: Mp 268°.
- 7-Ketone (lactone), α -N-oxide: **Homoly-corine N-oxide**
[119308-26-8]
C₁₈H₂₁NO₅ 331.368
Alkaloid from *Lapiedra martinezii*.
Cryst. + H₂O (MeOH aq.). Mp 134-136°. $[\alpha]_D^{25}$ +27.4 (c, 0.1 in MeOH).
- 7-Ketone (lactone), O⁹-de-Me: **9-O-Demethylhomolycorine**
[6879-81-8]
C₁₇H₁₉NO₄ 301.341
Alkaloid from *Lycoris radiata*, the bulbs of *Narcissus confusus* and *Crinum defixum* and the leaves of *Leucorum aestivum*. Also isol. from aerial parts of *Narcissus papyraceus* (Amaryllidaceae). Needles (EtOAc or H₂O). Mp 138-140° Mp 213-214° Mp 270-272°. $[\alpha]_D$ +96.4 (c, 0.28 in CHCl₃). Variation in Mp due to polymorphism. λ_{\max} 230 (log ϵ 4.25); 274 (log ϵ 3.84); 306 (log ϵ 3.65) (EtOH).
- 7-Ketone (lactone), O⁹-de-Me, α -N-oxide: **9-O-Demethylhomolycorine α -N-oxide**
[128517-02-2]
C₁₇H₁₉NO₅ 317.341
Alkaloid from aerial parts of *Narcissus papyraceus* (Amaryllidaceae). Cryst. (EtOH). Mp 153-154°. $[\alpha]_D^{23}$ +19 (c, 0.1 in MeOH). λ_{\max} 230 (log ϵ 4.43); 270 (log ϵ 4.28); 304 (log ϵ 3.9) (MeOH). λ_{\max} 208 (log ϵ 5.34); 248 (log ϵ 4.57); 274 (sh) (log ϵ 4.3); 340 (log ϵ 3.82) (MeOH/NaOH).
- 7-Ketone (lactone), O⁹-de-Me, 9-Ac: **9-O-Demethyl-9-O-acetylhomolycorine**
[139955-91-2]
C₁₉H₂₁NO₅ 343.379
Alkaloid from whole plants of *Narcissus vasconicus* (Amaryllidaceae). Cryst. Mp 186-188°. $[\alpha]_D^{20}$ +70.4 (c, 0.54 in EtOH). Erroneously descr. as 8-O-Acetylhomolycorine in the lit.
- 7-Deoxy: **Neruscine. Deoxylycorenine**
[13255-14-6]
C₁₈H₂₃NO₃ 301.385
Alkaloid from the bulbs of *Nerine corusca*, *Hippeastrum aulicum* var. *robustum* and from *Crinum powellii* (Amaryllidaceae). Mp 117-118°. $[\alpha]_D^{25}$ +92 (c, 0.5 in CHCl₃).
- 7-Deoxy, hydroiodide:
Prisms (EtOH aq.). Mp 247° dec. $[\alpha]_D^{25}$ +60 (c, 0.12 in EtOH).
- 7-Deoxy, perchlorate:
Cryst. (H₂O). Mp 212-213° dec.
- 17-Epimer, 7-ketone (lactone): **Epithomolycorine**
[66537-26-6]
C₁₈H₂₁NO₄ 315.368
Alkaloid from the bulbs of *Hippeastrum ananuca* (Amaryllidaceae).
- Stereoisomer, 17-ketone (lactone)(?):
Penarcine
C₁₈H₂₁NO₄ 315.368

- Isol. from *Narcissus cyclamineus* hybrid "Peeping Tom" (Amaryllidaceae). Mp 171-172°. $[\alpha]_D$ +110 (CHCl₃). Tentative struct., config. unknown.
- Boit, H.G. et al., *Chem. Ber.*, 1955, **88**, 133-136; 1957, **90**, 369-373 (*Homolycorine, Neruscine, isol*)
- Kitagawa, T. et al., *J.C.S.*, 1955, 1066-1068; 1959, 3741-3751 (*Lycorenine, Neruscine, Homolycorine, isol, uv, ir, struct, synth*)
- Boit, H.G. et al., *Naturwissenschaften*, 1958, **45**, 85; 262-263; 1960, **47**, 109; 159 (*Neruscine, Penarcine*)
- Uyeo, S. et al., *J.C.S.*, 1959, 172-177 (*9-Demethylhomolycorine*)
- Hawksworth, W.A. et al., *J.C.S.*, 1965, 1991-2001 (*pmr*)
- Ibuka, T. et al., *Tet. Lett.*, 1966, 4745-4748 (*ms*)
- Clardy, J. et al., *J.O.C.*, 1972, **37**, 49-51 (*cryst struct*)
- Fuganti, C. et al., *J.C.S. Perkin 1*, 1973, 954-956 (*biosynth*)
- Pacheco, P. et al., *Rev. Latinoam. Quim.*, 1978, **9**, 28-32; *CA*, **89**, 103746q (*Epithomolycorine*)
- Numata, A. et al., *Chem. Pharm. Bull.*, 1983, **31**, 2146-2149 (*Me ether, isol, ir, pmr*)
- Kobayashi, S. et al., *Chem. Pharm. Bull.*, 1985, **33**, 5258-5263 (*9-Demethylhomolycorine*)
- Jeffs, P.W. et al., *J.O.C.*, 1985, **50**, 1732-1737 (*9-Demethylhomolycorine*)
- Suau, R. et al., *Phytochemistry*, 1988, **27**, 3285-3287 (*Homolycorine, Homolycorine N-oxide, O-Methyllycorenine N-oxide*)
- Via, J. et al., *Acta Cryst. C*, 1989, **45**, 2020-2022 (*cryst struct, Eugenie*)
- Codina, C. et al., *Planta Med.*, 1989, **55**, 116 (*Eugenie*)
- Suau, R. et al., *Heterocycles*, 1990, **31**, 517-522 (*9-O-Demethylhomolycorine N-oxide*)
- Kihara, M. et al., *Chem. Pharm. Bull.*, 1991, **39**, 1849-1853 (*9-O-Demethylhomolycorine*)
- Bastida, J. et al., *J. Nat. Prod.*, 1992, **55**, 122-125 (*9-O-Demethyl-9-acetylhomolycorine*)
- Codina, C. et al., *Nat. Prod. Lett.*, 1992, **1**, 85-92 (*pmr, struct*)
- Codina, C. et al., *Phytochemistry*, 1993, **32**, 1354-1356 (*O-Methyllycorenine, Lycorenine*)
- Latvala, A. et al., *Tetrahedron: Asymmetry*, 1995, **6**, 361-364 (*9-O-Demethylhomolycorine*)
- Wagner, J. et al., *Tetrahedron*, 1996, **52**, 6591-6600 (*cd, Homolycorine, 9-O-Demethylhomolycorine*)
- Labrana, J. et al., *Phytochemistry*, 1999, **50**, 183-188 (*cryst struct, abs config*)
- Wang, Y.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1458-1461 (*9-O-Demethyl-7-O-methyllycorenine*)

Lycorine

Narcissine†. Galanthidine

L-348C₁₆H₁₇NO₄ 287.315**(+)-form**

Obt. by hydrol. of Poetaminine. Mp 262-264° dec.

1-Ac: **Poetaminine. (+)-1-O-Acetyllycorine**
[73543-66-5]**(-)-form**C₁₈H₁₉NO₅ 329.352
Alkaloid from the bulbs of *Narcissus poeticus* var. *ornatus* (Amaryllidaceae). Needles (MeOH). Mp 221-223°. $[\alpha]_D^{20}$ +100 (c, 0.2 in EtOH).

Di-Ac: Mp 215-217°.

(-)-form [476-28-8]Alkaloid of *Lycoris*, *Urginea*, *Crinum*, *Clivia*, *Narcissus*, *Hymenocallis*, *Hippeastrum*, *Pancreatum*, *Brunsvigia* spp. and of a very large number of spp. in the Amaryllidaceae. The most widespread of the Amaryllidaceae alkaloids. Causes scorbutic symptoms in exptl. animals. Respiratory stimulant. Phagocytosis modulator, antimalarial. Plant growth inhibitor by inhibition of protein synth. Shows mod. antitumour activity, prob. by clastogenicity. Antiviral agent, weak protozoicidal. Shows antifungal activity. Mp 280-281°. $[\alpha]_D^{26}$ -120 (EtOH). λ_{\max} 290 (EtOH) (Berdy).▶ Highly toxic, freq. cause of accidental human poisoning by *Narcissus* spp. LD₅₀ (dog, ivn) 41 mg/kg. OL2900000

Picrate: Mp 198-200°.

O¹-β-D-Glucopyranoside: Lycorine 1-O-β-D-glucoside

[93710-67-9]

C₂₂H₂₇NO₉ 449.457Alkaloid from the roots of *Pancreatum biflorum*, *Crinum asiaticum*, *Crinum augustum*, *Crinum latifolium* and *Crinum pratense* (Amaryllidaceae). Amorph. solid + 1H₂O. $[\alpha]_D^{20}$ -92.4 (c, 0.53 in MeOH).**O¹-(6-O-Hexadecanoyl-β-D-glucopyranoside): Lycoriside**

[99964-81-5]

C₃₈H₅₇NO₁₀ 687.869Alkaloid from the fruits of *Crinum asiaticum*. Also detected in the fleshy scale leaves and in the roots of this sp. (Amaryllidaceae). Amorph. powder. $[\alpha]_D^{25}$ -34.4 (c, 0.5 in MeOH).**O¹, O²-Di-O-β-D-glucopyranoside: Lycorine 1,2-di-O-β-D-glucoside**

[128701-12-2]

C₂₈H₃₇NO₁₄ 611.599Alkaloid from fruits of *Crinum asiaticum*. Straw-coloured solid + 1H₂O. $[\alpha]_D^{28}$ -27.5 (c, 0.58 in H₂O).**1-Ac: (-)-1-O-Acetyllycorine**

[7344-73-2]

C₁₈H₁₉NO₅ 329.352Alkaloid from the seeds of *Crinum moorei* and the bulbs of *Crinum latifolium* and *Nerine bowdenii* (Amaryllidaceae). Prisms (Me₂CO). Mp 220-221° (217-219°). $[\alpha]_D^{23}$ -96 (c, 1.11 in CHCl₃). $[\alpha]_D^{24}$ -66 (c, 1.18 in EtOH).**2-Ac: Aulamine. 2-Acetyllycorine**

[477-10-1]

C₁₈H₁₉NO₅ 329.352Alkaloid from the bulbs of *Hippeastrum aulicum* var. *robustum* (Amaryllidaceae). Shows antifungal activity. Mp 230-231°. $[\alpha]_D$ -38 (c, 0.2 in CHCl₃). $[\alpha]_D$ +20 (c, 0.2 in EtOH).

Di-Ac: 1,2-Di-O-acetyllycorine

[2492-05-9]

C₂₀H₂₁NO₆ 371.389

Alkaloid from the bulbs of *Crinum pratense* and *Crinum bulbispermum* (Amaryllidaceae). Shows antifungal activity. Platelets (Et₂O/MeOH). Mp 212-213° (208-210°). [α]_D²² -22 (c, 0.9 in CHCl₃).

O¹-Hexadecanoyl: Palmilycorine

[99965-60-3]

C₃₂H₄₇NO₅ 525.727

Alkaloid from the fruits of *Crinum asiaticum*. Also detected in the fleshy scale leaves and in the roots of this sp. (Amaryllidaceae). Amorph. powder. [α]_D²² -58.5 (c, 0.23 in MeOH).

O¹-Hexadecanoyl, O²-(1-O-hexadecanoyl-2-O-octadecanoyl)octadecanoyl-glycerophosphoryl: 1-O-Palmitoyl-2-O-(1-O-palmitoyl-2-O-stearoyl)oleoyl-glycerophosphoryllycorine

[128701-17-7, 128701-13-3]

Alkaloid from fruits of *Crinum asiaticum*. Straw-coloured amorph. powder. Mixt. contg. stearoyl and oleoyl alkyl residues.

O¹-(5,8,11,14-Eicosatetraenoyl) (all-Z): 1-O-Arachidonoyllycorine

[134253-41-1]

C₃₆H₄₇NO₅ 573.771

Alkaloid elaborated by the insect *Polytela gloriosa* which feeds on various Amaryllidaceae. Cream powder. [α]_D²⁸ -35.7 (c, 0.55 in CHCl₃).

2-O-(3-Acetoxybutanoyl): 2-O-(3-Acetoxybutanoyl)lycorine

[946570-72-5]

C₂₂H₂₅NO₇ 415.442

Alkaloid from *Galanthus nivalis*.

O²-Me: Hippamine

[6879-76-1]

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Hippeastrum* sp. Mp 162°. [α]_D²³ -72 (c, 1.5 in EtOH).

O²-(1-Hydroxyethyl), 1-Ac: Cliviasindhine

[92070-72-9]

C₂₀H₂₃NO₆ 373.405

Alkaloid from *Clivia miniata* (Amaryllidaceae). Long needles (Me₂CO). Mp 173-175°. [α]_D²⁴ -115 (c, 0.2 in CHCl₃).

O²-Glycerophosphoryl: 2-O-Glycerophosphoryllycorine

[109269-80-9]

C₁₉H₂₄NO₉P 441.374

Alkaloid from the flowers of *Zephyranthes flava* (Amaryllidaceae). Mp 300° dec.

O²-Phosphatidyl: Phosphatidyllycorine

[109269-84-3, 109269-81-0]

Alkaloid from flowers of *Zephyranthes flava* (Amaryllidaceae). Straw-coloured powder. Mp 280° dec. A mixt. of stearoyl and oleoyl homologues.

O²-Phosphatidyl, N-Me: Phosphatidylmethyllycorinium

[109269-87-6]

Isol. from flowers of *Zephyranthes flava* (Amaryllidaceae). Brown hygroscopic powder. Mixt. of α - and β -epimers at N.

2-Ketone, 1-Ac: JonquillineC₁₈H₁₇NO₅ 327.336

Alkaloid from *Narcissus jonquilla* Golden Sceptre hybrid. Needles (MeOH). Mp 188-190° dec. [α]_D²⁰ -325 (c, 0.5 in CHCl₃).

2-Epimer: 2-Epilycorine

[6835-91-2]

C₁₆H₁₇NO₄ 287.315

Alkaloid from the flower-stem fluid of *Clivia latifolium* (Amaryllidaceae). Mp 168-170°. [α]_D²⁸ -212.8 (c, 0.6 in MeOH).

2-Epimer, di-Ac:

Clusters of needles (petrol/MeOAc).

Mp 190-191°.

(±)-form [66816-51-1]

Mp 253-258° dec.

Takagi, S. *et al.*, *J.C.S.*, 1955, 4003 (*struct*)Lyle, R.E. *et al.*, *J.A.C.S.*, 1960, **82**, 2620

(Acetyllycorine, Diacetyllycorine)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960,**47**, 109 (Aulamine)Döpke, W. *et al.*, *Arch. Pharm. (Weinheim,**Ger.*), 1962, **295**, 920; *CA*, **58**, 11416a

(Hippamine)

Döpke, W. *et al.*, *Naturwissenschaften*, 1963,**50**, 354; 1965, **52**, 60; 61 (Poetamine,*Jonquilline)*Kotera, K. *et al.*, *Tet. Lett.*, 1966, 2009 (*pmr,**config*)Kinstle, T.H. *et al.*, *Tet. Lett.*, 1966, 4659 (*ms*)Bruce, I.T. *et al.*, *Chem. Comm.*, 1968, 207

(biosynth)

Sadikov, T. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**,134; *Chem. Nat. Compd. (Engl. Transl.)*,1972, **8**, 140 (*manuf*)Tsuda, Y. *et al.*, *Chem. Comm.*, 1975, 933

(synth)

Miyakado, M. *et al.*, *Phytochemistry*, 1975, **14**,2717 (*activity*)Gopalakrishna, E.M. *et al.*, *Cryst. Struct.**Commun.*, 1976, **5**, 795 (*cryst struct*)Moller, O. *et al.*, *Acta Chem. Scand., Ser. B*,1978, **32**, 98 (*synth*)Martin, S.F. *et al.*, *J.O.C.*, 1981, **46**, 3763;1982, **47**, 3634 (*synth*)Ghosal, S. *et al.*, *Phytochemistry*, 1981, **20**,2003; 1984, **23**, 1167; 1985, **24**, 2703; 1987,**26**, 823; 1989, **28**, 2535; 1990, **29**, 805

(Diacetyllycorine, Palmilycorine, Lycoriside,

*Phosphatidyllycorines,**glycerophosphoryllycorine, diglucoside,**glucoside, struct, epimer)*Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1984,**32**, 3015 (Acetyllycorine, Diacetyllycorine)Döpke, W. *et al.*, *Z. Chem.*, 1984, **24**, 209

(Cliviasindhine)

Suffness, M. *et al.*, *Alkaloids (Academic**Press)*, 1985, **25**, 200 (*occur, biol props*)Martin, S.F. *et al.*, *Alkaloids (Academic**Press)*, 1987, **30**, 251 (*rev*)Boeckman, R.K. *et al.*, *J.A.C.S.*, 1988, **110**,8250 (*synth*)Ghosal, S. *et al.*, *Indian J. Chem., Sect. B*,1991, **30**, 260 (1-Arachidonoyllycorine)Spohn, M. *et al.*, *Arch. Pharm. (Weinheim,**Ger.*), 1994, **327**, 123 (*cmr*)Schultz, A.G. *et al.*, *J.A.C.S.*, 1996, 571 (*synth*)Hoshino, O. *et al.*, *J.C.S. Perkin 1*, 1996, 571

(synth)

Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591

(cd)

Campbell, W.E. *et al.*, *Planta Med.*, 1998, **64**,91-93 (Diacetyllycorine, *pmr, cmr*)

Berkov, S. *et al.*, *Phytochemistry*, 2007, **68**,
1791-1798 (2-Acetoxybutanoyllycorine)

Lycoserramine**L-349**

[11052-59-8]

C₁₆H₂₅NO₂ 263.379

Struct. unknown. Alkaloid from *Lycopodium serratum* var. *serratum* f. *intermedium* (Lycopodiaceae). [α]_D -46.2 (c, 0.56 in MeOH).

Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1967, **87**,
1394-1404; *CA*, **68**, 47027h (*isol*)

Lycoserrine**L-350**

[11052-58-7]


C₁₆H₂₆N₂ 246.395

Struct. unknown. Alkaloid from *Lycopodium serratum* var. *serratum* f. *serratum* and *Lycopodium serratum* var. *serratum* f. *intermedium* (Lycopodiaceae). Mp 99-100°.

Inubushi, Y. *et al.*, *Yakugaku Zasshi*, 1967, **87**,
1394-1404; *CA*, **68**, 47027h (*isol*)

Lycosinine A**L-351**

[870273-47-5]

C₁₈H₂₁NO₃ 299.369

Alkaloid from the bulbs of *Lycoris aurea*. Powder. λ_{\max} 240 (log ϵ 3.57); 283 (log ϵ 3.28) (CHCl₃).

Aldehyde: Lycosinine B

[870273-48-6]


C₁₈H₁₉NO₃ 297.353

Alkaloid from the bulbs of *Lycoris aurea*. Powder. λ_{\max} 242 (log ϵ 4.39); 280 (log ϵ 4.13); 312 (log ϵ 3.97) (CHCl₃).

Yang, Y. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
2550-2553 (*isol, pmr, cmr*)

Lycovatine A**L-352**

[925689-16-3]

C₁₇H₂₈NO₂[⊕] 278.414


Alkaloid from *Lycopodium clavatum* var. *robustum*. Amorph. solid. [α]_D²³ -8 (c, 0.1

in MeOH). Counterion not specified.

Kubota, T. *et al.*, *Heterocycles*, 2006, **69**, 469-474 (*isol*, *pmr*, *cmr*)

Lydicamycin

[133352-27-9]



$C_{47}H_{74}N_4O_{10}$ 855.122

Prod. by *Streptomyces lydicus* SANK 60390. Active against gram-positive bacteria. Powder. Sol. MeOH, butanol; poorly sol. H_2O . Mp 161-166°. $[\alpha]_D^{18} +75.1$ (c, 1 in MeOH). λ_{max} 207 (€ 18900); 250 (sh) (€ 6600); 282 (€ 9900) (MeOH/HCl) (Derep). λ_{max} 208 (€ 19900); 245 (€ 9700); 281 (€ 9600) (MeOH/NaOH) (Derep). λ_{max} 207 (€ 19600); 245 (€ 9900); 282 (€ 10000) (MeOH) (Derep).

8-Deoxy-8-Deoxylydicamycin. Antibiotic TPU 0037D. TPU 0037D

$C_{47}H_{74}N_4O_9$ 839.123

Prod. by the marine-derived *Streptomyces platensis* TP-A0598. Active against gram-positive bacteria incl. MRSA. Powder. Mp 161-163°. $[\alpha]_D^{28} +50.3$ (c, 1 in MeOH). λ_{max} 203 (log € 4.31); 244 (log € 3.95); 282 (log € 3.96) (MeOH).

8-Deoxy, 14,15-didehydro-14,15-Didehydro-8-deoxylydicamycin. Antibiotic TPU 0037B. TPU 0037B

$C_{47}H_{72}N_4O_9$ 837.107

Prod. by the marine-derived *Streptomyces platensis* TP-A0598. Active against gram-positive bacteria incl. MRSA. Powder. Mp 162-164°. $[\alpha]_D^{28} +105.5$ (c, 1 in MeOH). λ_{max} 207 (log € 4.31); 244 (log € 3.95); 283 (log € 3.97) (MeOH).

30-Demethyl-30-Demethyllydicamycin.

Antibiotic TPU 0037A. TPU 0037A

$C_{46}H_{72}N_4O_{10}$ 841.096

Prod. by the marine-derived *Streptomyces platensis* TP-A0598. Active against gram-positive bacteria incl. MRSA. Powder. Mp 151-153°. $[\alpha]_D^{28} +57.5$ (c, 1 in MeOH). λ_{max} 204 (log € 4.31); 247 (log € 3.95); 282 (log € 3.95) (MeOH).

30-Demethyl, 8-deoxy-30-Demethyl-8-deoxylydicamycin. Antibiotic TPU 0037C. TPU 0037C

$C_{46}H_{72}N_4O_9$ 825.096

Prod. by the marine-derived *Streptomyces platensis* TP-A0598. Active against gram-positive bacteria incl. MRSA. Powder. Mp 161-163°. $[\alpha]_D^{28} +52$ (c, 0.76 in MeOH). λ_{max} 207 (log € 4.31); 244 (log € 3.96); 281 (log € 3.97) (MeOH).

Hayakawa, Y. *et al.*, *J. Antibiot.*, 1991, **44**, 282-287; 288-292 (*isol*, *props*)


Hayakawa, Y. *et al.*, *Tet. Lett.*, 1991, **32**, 213-216 (*pmr*, *cmr*, *ms*)

Furuma, T. *et al.*, *J. Antibiot.*, 2002, **55**, 873-880 (TPU 0037 antibiotics)

Lymphostin

Antibiotic LK6A. LK6A

[191474-39-2]



$C_{16}H_{14}N_4O_3$ 310.312

Prod. by *Streptomyces* sp. KY11783. Immunosuppressant and cytotoxic agent. Orange powder. Mp 275-277°. λ_{max} 218 (€ 15200); 256 (€ 19300); 284 (€ 16000); 344 (€ 11900); 412 (€ 6000); 467 (€ 8200) (MeOH).


Nagata, H. *et al.*, *J. Antibiot.*, 1997, **50**, 537-542; 543-545 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Nagata, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 2002, **66**, 501-507 (*activity*)

Tatsuta, K. *et al.*, *Tet. Lett.*, 2004, **45**, 2847-2850 (*synth*)

Lyngbyabellin A

[273202-85-0]



$C_{29}H_{40}Cl_2N_4O_7S_2$ 691.695

Depsipeptide antibiotic. Related to Dolabellin, D-901. Isol. from *Lyngbya majuscula*. Potent disruptor of the cellular microfilament network. Cytotoxic agent. Cryst. ($CH_2Cl_2/6$ -methylheptane). Mp 150-152°. $[\alpha]_D^{27} -74$ (c, 0.5 in $CHCl_3$). λ_{max} 202 (log € 4.48); 236 (log € 4.08) (MeOH).


Luesch, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 611-615 (*isol*)

Yokokawa, F. *et al.*, *Tetrahedron*, 2002, **58**, 9445-9458 (*synth*)

Lyngbyabellin B

Tortugamide

[300830-53-9]



$C_{28}H_{40}Cl_2N_4O_7S_2$ 679.684

Isol. from *Lyngbya majuscula*. Cytotoxic

L-354

and antifungal agent. Amorph. solid. $[\alpha]_D^{25} -152$ (c, 0.06 in $CHCl_3$). $[\alpha]_D +33$ (c, 0.2 in CH_2Cl_2). Isol. simultaneously by 2 groups. λ_{max} 200 (log € 4.49); 224 (log € 4.06); 236 (sh) (log € 3.93) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1437-1439


Milligan, K.E. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1440-1443

Yokokawa, F. *et al.*, *Tetrahedron*, 2002, **58**, 9445-9458 (*synth*)

Lyngbyabellin C

[479482-34-3]

L-357



$C_{24}H_{30}Cl_2N_2O_8S_2$ 609.547


Isol. from a *Lyngbya* sp. Cytotoxic. Amorph. solid. $[\alpha]_D^{25} -10$ (c, 0.1 in $CHCl_3$). λ_{max} 202 (log € 4.47); 235 (log € 4.04) (MeOH).

Luesch, H. *et al.*, *Tetrahedron*, 2002, **58**, 7959-7966 (*isol*, *pmr*, *cmr*)

Lyngbyabellin D

[532427-69-3]

L-358



$C_{38}H_{55}Cl_2N_3O_{13}S_2$ 896.902


Isol. from a *Lyngbya* sp. Cytotoxic. Powder. $[\alpha]_D^{25} +20$ (c, 0.4 in MeOH). λ_{max} 202 (log € 7.31); 223 (log € 6.81) (MeOH).

Williams, P.G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 595-598 (*isol*, *pmr*, *cmr*)

Lyngbyabellin E

[532427-69-3]

L-359




$C_{37}H_{51}Cl_2N_3O_{12}S_2$ 864.86

Isol. from *Lyngbya majuscula*. Amorph. solid. $[\alpha]_D^{26} -31$ (c, 0.7 in MeOH). λ_{max} 240 (log € 4.18) (MeOH).

21-Deoxy: Lyngbyabellin H

$C_{37}H_{51}Cl_2N_3O_{11}S_2$ 848.861
Isol. from *Lyngbya majuscula*.
Amorph. solid. $[\alpha]_D^{26}$ -53 (c, 0.08 in MeOH). λ_{max} 242 (log ϵ 4.15) (MeOH).

Han, B. *et al.*, *Tetrahedron*, 2005, **61**, 11723-11729 (isol, pmr, cmr)


Lyngbyabellin F**L-360**

$C_{38}H_{55}Cl_2N_3O_{13}S_2$ 896.902
Isol. from *Lyngbya majuscula*. Amorph. solid. $[\alpha]_D^{26}$ -6.5 (c, 0.2 in MeOH). λ_{max} 238 (log ϵ 4) (MeOH).

21-Deoxy: Lyngbyabellin I

$C_{38}H_{55}Cl_2N_3O_{12}S_2$ 880.903
Isol. from *Lyngbya majuscula*.
Amorph. solid. $[\alpha]_D^{26}$ -25 (c, 0.04 in MeOH). λ_{max} 238 (log ϵ 4.28) (MeOH).

Han, B. *et al.*, *Tetrahedron*, 2005, **61**, 11723-11729 (isol, pmr, cmr)

Lyngbyabellin G**L-361**


Absolute Configuration

$C_{23}H_{28}Cl_2N_2O_8S_2$ 595.52
Isol. from *Lyngbya majuscula*. Amorph. solid. $[\alpha]_D^{26}$ -26 (c, 0.2 in MeOH). λ_{max} 238 (log ϵ 3.75) (MeOH).

Han, B. *et al.*, *Tetrahedron*, 2005, **61**, 11723-11729 (isol, pmr, cmr)

Lyngbyapeptin A**L-362**

[221531-83-5]



Absolute Configuration

$C_{37}H_{55}N_5O_6S$ 697.937
Isol. from *Lyngbya bouillonii* and *Lyng-*


bya majuscula. Amorph. solid. $[\alpha]_D^{25}$ -235 (c, 0.58 in MeOH). λ_{max} 201 (log ϵ 4.55); 225 (log ϵ 4.3); 240 (sh) (log ϵ 4.09) (MeOH).

15-O-De-Me: 15-Norlyngbyapeptin A

[532427-68-2]
 $C_{36}H_{53}N_5O_6S$ 683.911
Isol. from a *Lyngbya* sp. Powder. $[\alpha]_D^{22}$ -31 (c, 0.4 in MeOH). λ_{max} 201 (log ϵ 4.2); 226 (log ϵ 4.02) (MeOH).
Klein, D. *et al.*, *Tet. Lett.*, 1999, **40**, 695-696 (isol, pmr, cmr)
Luesch, H. *et al.*, *J.O.C.*, 2000, **63**, 1437-1439 (abs config)
Williams, P.G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 595-598 (15-Norlyngbyapeptin A)

Lyngbyapeptin B**L-363**

[479482-35-4]



Absolute Configuration

$C_{38}H_{51}N_5O_7S$ 721.916
Isol. from a *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25}$ -50 (c, 0.23 in MeOH). λ_{max} 200 (log ϵ 4.53); 227 (log ϵ 4.24); 245 (sh) (log ϵ 4) (MeOH).

Luesch, H. *et al.*, *Tetrahedron*, 2002, **58**, 7959-7966 (isol, pmr, cmr)

Lyngbyapeptin C**L-364**

[479482-36-5]

As Lyngbyapeptin B, L-363 with


R = CH₂CH₃

$C_{39}H_{53}N_5O_7S$ 735.943
Isol. from a *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25}$ -52 (c, 0.09 in MeOH). λ_{max} 200 (log ϵ 4.53); 227 (log ϵ 4.26); 245 (sh) (log ϵ 4) (MeOH).

Luesch, H. *et al.*, *Tetrahedron*, 2002, **58**, 7959-7966 (isol, pmr, cmr)

Lyngbyastatin 4**L-365**

[928202-70-4]



Absolute Configuration

$C_{53}H_{68}N_8O_{18}S$ 1137.229
Depsipeptide antibiotic. Isol. from *Lyngbya confervoides*. Elastase and chymotrypsin inhibitor. Amorph. powder. $[\alpha]_D^{20}$ +8.4 (c, 0.25 in MeOH). λ_{max} 210 (log ϵ 4.57); 280 (sh) (log ϵ 3.6) (MeOH).

Desulfo: Lyngbyastatin 5
[957130-98-2]

$C_{53}H_{68}N_8O_{15}$ 1057.165
Isol. from *Lyngbya confervoides*. Elastase inhibitor. Amorph. solid. λ_{max} 210 (log ϵ 4.57); 280 (sh) (log ϵ 3.79) (MeOH).

11-Me ether: Lyngbyastatin 6

[957130-99-3]
 $C_{54}H_{70}N_8O_{18}S$ 1151.256
Isol. from a *Lyngbya* sp. Amorph. powder. λ_{max} 210 (log ϵ 4.48); 280 (sh) (log ϵ 3.65) (MeOH).

Matthew, S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 124-127 (Lyngbyastatin 4)

Taori, K. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1593-1600 (Lyngbyastatins 5,6)

Lyngbyastatin 7**L-366**

[957131-00-9]


As Somamide A, S-360 with
R¹ = -CH₂CH₂CONH₂, R² = -
(CH₂)₄CH₃

$C_{48}H_{66}N_8O_{12}$ 947.096
Isol. from a *Lyngbya* sp. Elastase inhibitor. Amorph. powder. $[\alpha]_D^{20}$ -7.4 (c, 0.27 in MeOH). λ_{max} 230 (log ϵ 3.8); 280 (sh) (log ϵ 3.12) (MeOH).

Taori, K. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1593-1600 (isol, pmr, cmr)

Lyngbyatoxin A**L-367**

Teleocidin A,
[70497-14-2]



$C_{27}H_{39}N_3O_2$ 437.624

Numbering as given by Aimi *et al.* Isol. from lipid extracts of a Hawaiian shallow-water variety of the marine blue-green alga *Lyngbya majuscula*. Also prod. by *Streptomyces medicidicus*. Nematocidal and acaricidal. Potent tumour promoter. Mp 61°. $[\alpha]_D$ -110 (MeOH). $[\alpha]_D$ -171 (c, 1.8 in CHCl₃). Closely related to Teleocidin B₁, T-63. λ_{max} 233 (ϵ 34700); 287 (ϵ 9770); 298 (sh) (ϵ 8130) (MeOH) (Derep).

► Highly inflammatory and vesicatory. Tumour promoter. Causes dermatitis. Very toxic. LD₅₀ (mus, orl) 2mg/kg. WY1981000

14-O-(2-Acetamido-2-deoxy-β-D-glucopyranoside): 14-O-(N-Acetylglucosaminyl)teleocidin A

$C_{35}H_{52}N_4O_7$ 640.818
Prod. by *Streptomyces* sp. MM216-87F4. Powder. $[\alpha]_D$ +130.2 (c, 0.15 in MeOH). Stereochem. not confirmed. λ_{max} 229 (ϵ 19000); 279 (ϵ 7416); 297 (sh) (ϵ 7200) (MeOH).

14-Ac: O-Acetyllyngbyatoxin A

[70497-15-3]
 C₂₉H₄₁N₃O₃ 479.661
 Isol. from the sea hare *Stylocheilus longicauda*. Oil. [α]_D²⁰ -4 (c, 0.4 in MeOH). λ_{max} 229 (ε 4260); 297 (ε 1270) (MeOH).

N-De-Me: N¹³-Demethylteleocidin A₁
 C₂₆H₃₇N₃O₂ 423.597
 From *Streptovorticillium blastmyceticum*. No phys. props. reported. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{max} 228 (ε 25900); 285 (ε 7600); 295 (ε 6800) (MeOH) (Derep).

► Tumour promoter.

Me ether: O-Methylteleocidin A₁
 [114774-86-6]
 C₂₈H₄₁N₃O₂ 451.651
 From *Streptovorticillium blastmyceticum*. Gummy solid. Sol. MeOH, Me₂CO, C₆H₆; poorly sol. H₂O. [α]_D¹⁸ -198 (c, 0.21 in CHCl₃). Numbering systems vary. λ_{max} 230 (ε 27000); 286 (ε 8400); 301 (ε 9200) (MeOH) (Berdy).

► Tumour promoter.

A²⁶-Isomer, 24ξ-hydroxy: Lyngbyatoxin B
 [133084-52-3]
 C₂₇H₃₉N₃O₃ 453.623
 Isol. from *Lyngbya majuscula*. TPA binding inhibitor. Amorph.

► Skin irritant.

A²⁴-Isomer, 26-hydroxy: Lyngbyatoxin C
 [133084-53-4]
 C₂₇H₃₉N₃O₃ 453.623
 Isol. from *Lyngbya majuscula*. TPA binding inhibitor. Amorph.

► Skin irritant.

19-Epimer: Teleocidin A₂
 [102209-77-8]
 C₂₇H₃₉N₃O₂ 437.624
 From *Streptovorticillium mediocidicus*. [α]_D¹⁸ -185.1 (c, 0.18 in MeOH). λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

► Potent tumour promoter.

19-Epimer, Ac: 14-O-Acetylteleocidin A₂
 C₂₉H₄₁N₃O₃ 479.661
 Prod. by *Actinomyces* sp. Stamm K 17/7. Yellow oil. [α]_D²⁰ -81 (c, 0.1 in MeOH). λ_{max} 229 (log ε 4.41); 288 (sh); 297 (log ε 3.93) (MeOH).

Takashima, M. *et al.*, *Agric. Biol. Chem.*, 1962, **26**, 660 (*isol, props*)

Cardellina, J.H. *et al.*, *Science (Washington, D.C.)*, 1979, **204**, 193 (*isol, pmr, cmr, cd, struct*)

Moore, R.E. *et al.*, *Pure Appl. Chem.*, 1982, **54**, 1919 (*rev*)

Sakai, S.-I. *et al.*, *Tet. Lett.*, 1986, **27**, 5219

Muratake, H. *et al.*, *Tet. Lett.*, 1987, **28**, 2265 (*synth*)

Hagiwara, N. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 641-648 (*O-Methylteleocidin A₁*)

Irie, K. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 3193 (*deriv*)

Aimi, N. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1593 (*Lyngbyatoxin B, Lyngbyatoxin C*)

Irie, K. *et al.*, *Tetrahedron*, 1990, **46**, 2773-2788 (*biosynth*)

Muratake, H. *et al.*, *Tetrahedron*, 1991, **47**, 8535; 8545; 8559 (*synth*)

Gallimore, W.A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1022-1026 (*O-Acetyllyngbyatoxin A*)

Ströck, K. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*14-O-Acetylteleocidin A₂*)

Nakae, K. *et al.*, *J. Antibiot.*, 2006, **59**, 11-17 (*14-Acetylglucosaminylteleocidin A*)

Lyophyllin L-368

Trimethyl diazenecarboxamide 2-oxide, 9CI

[88245-13-0]

Me₂NCON=N(O)Me

C₄H₉N₃O₂ 131.134


Isol. from the toadstool *Lyophyllum connatum*. Mp 27-28°.

► Prob. mutagenic, by analogy.

Fugmann, B. *et al.*, *Angew. Chem., Int. Ed.*, 1984, **23**, 72 (*isol, synth, pmr, cmr, ir, tox*)

Ye, Y. *et al.*, *Tet. Lett.*, 1997, **38**, 8013-8016 (*biosynth*)

Lysergene L-369



(*R*)-form

C₁₆H₁₆N₂ 236.316

(*R*)-form [478-91-1]

Alkaloid from ergot (*Claviceps purpurea*), and from *Argyria cuneata*, *Argyria nervosa* and *Rivea corymbosa* (Convolvulaceae). Mp 247-249°. [α]_D²⁰ +504 (c, 0.4 in PY).

(±)-form

Synthetic. Cryst. (Me₂CO). Mp 210-212° dec.

Abe, M. *et al.*, *Agric. Biol. Chem.*, 1961, **25**, 594 (*isol*)

Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 2473 (*cd, ord*)

Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435 (*occur*)

Kiguchi, T. *et al.*, *Heterocycles*, 1984, **22**, 43 (*synth*)

Wheeler, W.J. *et al.*, *Tet. Lett.*, 1986, **27**, 3469 (*synth*)


Ninomiya, I. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 23 (*synth*)

Harris, J.R. *et al.*, *Synth. Commun.*, 1992, **22**, 995 (*synth*)

Saá, C. *et al.*, *Synlett*, 1994, 487 (*synth*)

Lysergic acid L-370

9,10-Didehydro-6-methylergoline-8-carboxylic acid, 9CI



(+)-form

C₁₆H₁₆N₂O₂ 268.315

(+)-form [82-58-6]

Alkaloid from *Claviceps purpurea*; obt. by the hydrol. of the ergotaman group of alkaloids. 5-HT Receptor antagonist. Psychomimetic, mycotoxin, depressant. Sol. H₂O. Mp 238° dec. [α]_D²⁰ +40 (Py).

Hydrochloride: Mp 208-210° dec.

Me ester: Mp 168°.

Amide: Ergine. Lysergic acid amide.

Lysergamide

[478-94-4]

C₁₆H₁₇N₃O 267.33

Alkaloid from *Claviceps paspali* and several spp. in the genera *Argyria*, *Stictocardia*, *Ipomoea* and *Rivea* (Convolvulaceae). Hallucinogen.

Prisms (MeOH). Mp 242° dec. [α]_D²⁰ 0 (c, 0.5 in Py). [α]_D²⁰ +15 (c, 0.5 in Py).

► OL4025000

α-Hydroxyethylamide: Lysergic acid α-hydroxyethylamide. Lysergic acid methylcarbinolamide

C₁₈H₂₁N₃O₂ 311.383

Alkaloid from *Claviceps paspali* and several *Argyria*, *Rivea*, *Ipomoea* and *Stictocardia* spp. (Convolvulaceae). Mp 135° dec. [α]_D²⁰ +29 (c, 1 in DMF).

8-Hydroxy, amide: 8-Hydroxyergine. 8-Hydroxylysergic acid amide

C₁₆H₁₇N₃O₂ 283.329

Alkaloid from *Claviceps paspali* strain MG-6.

8-Epimer: Isoysergic acid

[478-95-5]

C₁₆H₁₆N₂O₂ 268.315

Isol. from *Claviceps purpurea* and *Claviceps paspali* (Convolvulaceae). Mp 218° dec. [α]_D²⁰ +281 (Py). pK_{a1} 3.33; pK_{a2} 8.46 (24°).

8-Epimer, amide: Isoergine. Erginine. Iso-lysergic acid amide. Isoysergamide

[2889-26-1]

C₁₆H₁₇N₃O 267.33

Alkaloid from *Claviceps paspali* and several *Argyria*, *Rivea* and *Ipomoea* spp. Readily formed by rearr. of Ergine, e.g. by recrystallisation of the latter from MeOH (Convolvulaceae). Prisms + 1MeOH (MeOH). Mp 132-134° dec. [α]_D²⁰ +480 (c, 0.5 in Py).

8-Epimer, α-hydroxyethylamide: Isoysergic acid α-hydroxyethylamide. Isoysergic acid methylcarbinolamide

C₁₈H₂₁N₃O₂ 311.383

Alkaloid from *Claviceps paspali* and several *Argyria*, *Rivea* and *Ipomoea* spp. (Convolvulaceae). Noncryst.

8-Epimer, 8-hydroxy, amide: 8-Hydroxyerginine

C₁₆H₁₇N₃O₂ 283.329

Alkaloid from *Claviceps paspali* strain MG-6.

(-)-form

Synthetic. Mp 235-240° dec. [α]_D²⁰ -40 (Py).

(±)-form [23953-76-6]


Mp 242-243° dec. pK_{a1} 3.32; pK_{a2} 7.82 (24°).

Stoll, A. *et al.*, *Helv. Chim. Acta*, 1949, **32**, 506; 1954, **37**, 2039 (*struct, uv, config, ir*)

- Kornfeld, E.C. *et al.*, *J.A.C.S.*, 1956, **78**, 3087 (synth, ir)
- Hofmann, A. *et al.*, *Experientia*, 1960, **16**, 414 (isol, uv)
- Arcamone, F. *et al.*, *Nature (London)*, 1960, **187**, 238 (amides, α -hydroxyethylamides)
- Stadler, P.A. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 2005 (abs config)
- Castagnoli, N. *et al.*, *Nature (London)*, 1966, **211**, 859 (occur)
- Inoue, T. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 409 (ms)
- Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435 (occur)
- Bach, N.J. *et al.*, *J.O.C.*, 1974, **39**, 1272 (cmr)
- Armstrong, V.W. *et al.*, *Tet. Lett.*, 1976, 4311 (synth)
- Bellatti, M. *et al.*, *Tetrahedron*, 1977, **33**, 1821 (biosynth)
- Kumbar, M. *et al.*, *NIDA Res. Monogr.*, 1978, **22**, 374 (conformn, props)
- Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 478 (synth)
- Ramage, R. *et al.*, *Tetrahedron*, 1981, **37**, 157 (synth)
- Rebek, J. *et al.*, *J.A.C.S.*, 1984, **106**, 1813 (synth)
- Ninomiya, I. *et al.*, *J.C.S. Perkin I*, 1985, 941 (synth)
- Kurihara, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 4793 (synth)
- Cacchi, S. *et al.*, *Tet. Lett.*, 1988, **29**, 3117 (synth)
- Flieger, M. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1003 (8-Hydroxyergine, 8-Hydroxyerginine)
- Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1395
- Moldvai, I. *et al.*, *J.O.C.*, 2004, **69**, 5993-6000 (synth)
- Hendrickson, J.B. *et al.*, *Org. Lett.*, 2004, **6**, 3-5 (synth)

Lysergol

[602-85-7]

L-371

Absolute Configuration

$C_{16}H_{18}N_2O$ 254.331
Alkaloid from ergot (*Claviceps purpurea*) and from several spp. in the genera *Argyriaea*, *Stictocardia*, *Rivea* and *Ipomoea* (Convolvulaceae). 5-HT receptor antagonist, mycotoxin. Cryst. (EtOH). Mp 253-255°. $[\alpha]_D^{20} +54$ (c, 0.3 in Py).

▶ KE6330000

Deoxy: Lysergine

[519-10-8]

 $C_{16}H_{18}N_2$ 238.332

Alkaloid from ergot (*Claviceps purpurea*). Cryst. (EtOH or Me_2CO). Mp 286-288°. $[\alpha]_D^{20} +65$ (c, 0.5 in Py).

8-Epimer: Isolysergol

[478-93-3]

 $C_{16}H_{18}N_2O$ 254.331

Alkaloid from ergot (*Claviceps purpurea*), *Argyriaea nervosa* and several

other *Argyriaea* spp. and *Ipomoea violacea* (Convolvulaceae).

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 168B (nmr)

Schreier, E. *et al.*, *Helv. Chim. Acta*, 1958, **41**, 1984-1997 (*Lysergine*, synth)

Abe, M. *et al.*, *Agric. Biol. Chem.*, 1961, **25**, 594-595 (*Lysergol*, *Lysergine*, synth)

Hofmann, A. *et al.*, *Planta Med.*, 1961, **9**, 354-367 (isol)


Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 2473-2476 (ord)

Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435-2440 (occur)

Ninomiya, I. *et al.*, *Heterocycles*, 1984, **22**, 1035-1038 (*Isolysergol*, synth)

Ninomiya, I. *et al.*, *J.C.S. Perkin I*, 1990, 707-713 (*Lysergol*, *Isolysergol*, synth)

Harris, J.R. *et al.*, *Synth. Commun.*, 1992, **22**, 995-999 (synth, pmr, cmr)

Lysergylvaline**L-372**

Absolute Configuration

 $C_{21}H_{25}N_3O_3$ 367.447**Me ester: Lysergylvaline methyl ester**

Alkaloid from ergot (*Claviceps purpurea*). Amorph. Mp 80-85°. $[\alpha]_D^{20} -103$ (c, 1.6 in $CHCl_3$). $[\alpha]_D^{20} -65$ (c, 1.4 in Py). Probably an artifact arising from fast methanolysis of Ergocristam, E-154 and related substances.

Me ester, hydrogen maleate:

Needles (Me_2CO). Mp 185° dec. $[\alpha]_D^{20} +38$ (c, 1.2 in EtOH/ H_2O 1:1).

Amide: Ergovalide. Lysergylvaline amide

[41645-63-0]

 $C_{21}H_{26}N_4O_2$ 366.462

Alkaloid from ergot. Cryst. (MeOH). Mp 152.5-154°. $[\alpha]_D^{20} -80.8$ (c, 0.47 in $CHCl_3$). Prob. an artifact arising from solvolytic cleavage of Lysergylvalylphenylalanylproline.

Amide; hydrochloride: Mp 273-275° dec.


Schlientz, W. *et al.*, *Experientia*, 1963, **19**, 397 (isol, uv, struct)

Stütz, P. *et al.*, *Experientia*, 1973, **29**, 936

Ban'kovskaya, A.N. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 134; *Chem. Nat. Compd. (Engl. Transl.)*, 139 (*Ergovalide*)

Lysianadiolic acid**L-373**

[3-(Aminoiminomethyl)propylidene]butanedioic acid. (3-Guanidinopropylidene)-succinic acid
[1010815-66-3]


 $C_8H_{13}N_3O_4$ 215.208

Alkaloid from *Lysiana subfalcata*. Carboxypeptidase B inhibitor. Amorph. solid. λ_{max} 198 (log ϵ 3.65) (MeOH).

Buchanan, M.S. *et al.*, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 1495-1497 (isol, pmr, cmr)

Lysolipin I, 9CI**L-374**

***A*₃-118D. Antibiotic *A*₃-118D**
[59113-57-4]



Absolute Configuration

 $C_{29}H_{24}ClNO_{11}$ 597.961

Isol. from *Streptomyces violaceoniger*. Active against fungi. Cytotoxic. Yellow needles. Mp 260-262°. $[\alpha]_D -50.2$ ($CHCl_3$). Related to Albofungin, A-244. λ_{max} 278 (ϵ 34100); 310 (ϵ 18600); 356 (ϵ 13000); 408 (sh) (ϵ 5500) ($CHCl_3$) (De-rep).

Drautz, H. *et al.*, *Arch. Microbiol.*, 1975, **106**, 175-190 (isol, ir, uv, ms, pmr, cmr)


Dobler, M. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 178-185 (cryst struct, pmr)

Yang, J. *et al.*, *CA*, 1992, **117**, 44149 (isol)

Bockholt, H. *et al.*, *J.O.C.*, 1994, **59**, 2064-2069 (biosynth, pmr, cmr)

Lysolipin X**L-375**

[59029-83-3]



Probable Absolute Configuration

 $C_{29}H_{26}ClNO_{12}$ 615.977

C-11 and C-12 configs. not certain. Isol. from *Streptomyces violaceoniger* and other *Streptomyces* spp. Active against gram-positive and -negative bacteria and fungi. Cytotoxic agent. Sol. MeOH, $CHCl_3$; poorly sol. hexane. λ_{max} 275 ; 312 ($CHCl_3$) (Berdy).


Drautz, H. *et al.*, *Arch. Microbiol.*, 1975, **106**, 175-190 (isol, uv, ir, pmr)

U.S. Pat., 1981, 4 277 478; *CA*, **95**, 130977 (isol, props)

Bockholt, H. *et al.*, *J.O.C.*, 1994, **59**, 2064-2069 (biosynth, pmr, cmr)

Lythramine**L-376**

[32420-56-7]

 $C_{29}H_{37}NO_5$ 479.615

Alkaloid from *Lythrum anceps* (Lythraceae). Also obt. on treatment of Lythranine in L-378 with HCHO. Cryst. + 0.5 Me₂CO (Me₂CO). Mp 150-152° (solvate). $[\alpha]_D^{20}$ -85 (c, 0.9 in dioxan).

Me ether:

Cryst. (Me₂CO/heptane). Mp 169-171°.

Fujita, E. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 2216 (*isol, uv, ir, pmr, struct, synth*)


Fujita, E. *et al.*, *J.C.S.(C)*, 1971, 1651 (*struct, abs config*)

Fuji, K. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 63-69 (*cryst struct*)

Lythrancine I

Lythrancine, 9CI
[32209-71-5]

L-377



C₂₇H₃₅NO₅ 453.577

Alkaloid from *Lythrum anceps* (Lythraceae). Powder. $[\alpha]_D^{20}$ +65 (c, 0.96 in CHCl₃).

O¹¹-Ac: Lythrancine II

[32209-72-6]

C₂₉H₃₇NO₆ 495.614

Alkaloid from *Lythrum anceps* (Lythraceae). Needles (MeOH/hexane). Mp 274-275°. $[\alpha]_D^{20}$ +125 (c, 0.73 in CHCl₃).

O³,O¹¹-Di-Ac: Lythrancine III

[32209-73-7]

C₃₁H₃₉NO₇ 537.652

Alkaloid from *Lythrum anceps* (Lythraceae). Needles (MeOH). Mp 134-136°. $[\alpha]_D^{20}$ +38 (c, 1.1 in CHCl₃).

Tri-Ac: Lythrancine IV

[32209-74-8]

C₃₃H₄₁NO₈ 579.689

Alkaloid from *Lythrum anceps* (Lythraceae). Needles (MeOH). Mp 237-238°. $[\alpha]_D^{20}$ +27 (c, 1.1 in CHCl₃).

4-Deoxy: Lythrancepine I. 9-Deoxylythrancine, 9CI

[32209-75-9]

C₂₇H₃₅NO₄ 437.578

Alkaloid from *Lythrum anceps* (Lythraceae). Cryst. (Et₂O). Mp 149-151°.

$[\alpha]_D^{20}$ +59 (c, 0.98 in CHCl₃).

4-Deoxy, O¹¹-Ac: Lythrancepine II

[32209-76-0]

C₂₉H₃₇NO₅ 479.615

Alkaloid from *Lythrum anceps* (Lythraceae). Cryst. (CHCl₃/Et₂O). Mp 187-189°. $[\alpha]_D^{20}$ +44 (c, 1.0 in CHCl₃).

4-Deoxy, di-Ac: Lythrancepine III

[32209-77-1]

C₃₁H₃₉NO₆ 521.652

Alkaloid from *Lythrum anceps* (Lythraceae). Cryst. (EtOH). Mp 175-177°. $[\alpha]_D^{20}$ +7 (c, 0.23 in CHCl₃).

3-Epimer, O³,O⁴-di-Ac: Lythrancine VI

[40179-99-5]

C₃₁H₃₉NO₇ 537.652

Minor alkaloid from *Lythrum anceps* (Lythraceae). Oil. $[\alpha]_D$ +25.5 (c, 1.8 in CHCl₃).

3-Epimer, O³, O¹¹-di-Ac: Lythrancine VII

[40180-00-5]

C₃₁H₃₉NO₇ 537.652

Minor alkaloid from *Lythrum anceps* (Lythraceae). Oil. $[\alpha]_D$ +101.5 (c, 0.37 in CHCl₃).

3-Epimer, tri-Ac: Lythrancine V

[40179-98-4]

C₃₃H₄₁NO₈ 579.689

Minor alkaloid from *Lythrum anceps* (Lythraceae). Cryst. (MeOH). Mp 133-134°. $[\alpha]_D$ +91 (c, 0.55 in CHCl₃).

Fujita, E. *et al.*, *J. Nat. Prod.*, 1971, **34**, 306 (*isol*)

Fujita, E. *et al.*, *J.C.S. Perkin 1*, 1972, 2141; 1973, 297; 301; 306 (*uv, ir, pmr, ms, abs config, synth*)

Barrow, M.J. *et al.*, *J.C.S. Perkin 2*, 1974, 1812 (*cryst struct, Lythrancepines*)

Hart, D.J. *et al.*, *J.O.C.*, 1985, **50**, 3670; 1987, **52**, 4665 (*synth, ir, pmr, ms, Lythrancepines II, III*)


Lythranidine

Deacetyllythramine

[17812-78-1]

[70832-04-1 ((±)-form)]

L-378



(-)-form

C₂₆H₃₅NO₄ 425.567

Alkaloid from *Lythrum anceps* (Lythraceae). Liq.; cryst. + 1/2 H₂O (as acetate salt). Mp 136-139° (acetate salt). $[\alpha]_D^{20}$ -71 (c, 1.7 in dioxan).

O¹¹-Ac: Lythranine

[32420-54-5]

C₂₈H₃₇NO₅ 467.604

Alkaloid from *Lythrum anceps* (Lythraceae). DMSO adduct, cryst. + 1H₂O (EtOAc or CCl₄). Mp 92-94°.

O¹¹-Ac; hydrochloride:

Cryst. + 1H₂O (MeOH). Mp 189-191° dec. $[\alpha]_D^{20}$ -40 (c, 1 in CHCl₃).

Tri-Ac: Mp 180-218° dec. (as hydrochloride). $[\alpha]_D^{25}$ -33 (c, 0.5 in CHCl₃).

Fujita, E. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 2216; 2393 (*isol, struct, ir, pmr, synth*)

Fujita, E. *et al.*, *J.C.S.(C)*, 1971, 1651 (*struct, abs config*)

McClure, R.J. *et al.*, *J.C.S. Perkin 2*, 1972, 2073 (*cryst struct, abs config, deriv*)

Fuji, K. *et al.*, *Tet. Lett.*, 1979, 361 (*synth, ir, pmr*)

Cox, P.J. *et al.*, *Acta Cryst. B*, 1982, **38**, 303 (*cryst struct, Lythranine*)


Fuji, K. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 63-69 (*cryst struct*)

Lythrumine

L-379

O⁶-Demethyl-9,12-dideoxy-12-oxolythrancine, 9CI

[51024-99-8]



C₂₆H₃₁NO₄ 421.535

Alkaloid from *Lythrum lanceolatum* (Lythraceae). Mp 214-216°. $[\alpha]_D^{26}$ -8 (MeOH).

O³-Ac: Monoacetyllythrumine

[50651-33-7]

C₂₈H₃₃NO₅ 463.572

Alkaloid from *Lythrum lanceolatum* (Lythraceae). Mp 184-185°. $[\alpha]_D^{26}$ -34 (MeOH).

Wright, H. *et al.*, *J.A.C.S.*, 1973, **95**, 6467 (*uv, ir, pmr, cryst struct, abs config*)

Maandrosine

M-1

[1359-36-0]

Struct. unknown. Alkaloid from the roots of *Catharanthus roseus* (Apocynaceae). Prisms (MeOH) (as sulfate salt). Mp 160-173° (sulfate).

Sulfate:


Prisms (MeOH). Mp 160-173°. pK_a 6.9 (33% DMF aq.).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (*isol, uv, ir*)

Macaridine

M-2

3-Benzyl-1,2-dihydro-1-hydroxy-4-pyridinecarboxaldehyde. 3-Benzyl-4-formyl-1,2-dihydro-1-hydroxypyridine
[405914-36-5]



C₁₃H₁₃NO₂ 215.251


Alkaloid from the tubers of *Lepidium meyenii* (maca). Amorph. solid. λ_{max} 208 (log ε 4.07); 255 (sh) (log ε 3.76); 294 (log ε 4.14) (MeOH).

Muhammad, I. *et al.*, *Phytochemistry*, 2002, **59**, 105-110 (*isol*)

Machilaminoside A

M-3

[922510-41-6]



C₃₇H₅₄N₂O₁₂ 718.84


Constit. of *Machilus yaoshansis*. Amorph. powder. [α]_D²⁰ -48.8 (c, 0.06 in MeOH).

Liu, M.-T. *et al.*, *Org. Lett.*, 2007, **9**, 129-132 (*isol, pmr, cmr, ms*)

Machilaminoside B

M-4

[922510-42-7]



C₄₆H₆₅N₅O₁₆ 944.043


Constit. of *Machilus yaoshansis*. Amorph. powder (MeOH). [α]_D²⁰ -90.3 (c, 0.06 in MeOH).

Liu, M.-T. *et al.*, *Org. Lett.*, 2007, **9**, 129-132 (*isol, pmr, cmr, ms*)

Mackinazolinone

M-5

6,7,8,9-Tetrahydropyrido[2,1-b]quinazolin-11-one
[2446-62-0]



C₁₂H₁₂N₂O 200.24

Alkaloid from the leaves of *Mackinlaya macrosciadia* and *Mackinlaya subulata* (Araliaceae). Cryst. (H₂O or hexane). Mp 100.5°.

Stephen, T. *et al.*, *J.C.S.*, 1956, 4694 (*synth*)
Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1966, **19**, 151 (*isol, pmr, ms*)

Kametani, T. *et al.*, *Heterocycles*, 1976, **4**, 1487 (*synth, ir, pmr*)


Liu, J.-F. *et al.*, *Org. Lett.*, 2005, **7**, 3363-3366 (*synth, pmr, cmr*)

Bowman, W.R. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 103-113 (*synth*)

Macleanine

M-6

[154963-00-5]



C₁₆H₂₆N₂ 246.395


Alkaloid from *Huperzia serrata* (*Lycopodium serratum*) (Lycopodiaceae). Mp 300° (dec.). [α]_D +12.5 (c, 0.24 in CHCl₃). Unique structural type among the Lycopodium alkaloids.

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1994, **72**, 128 (*isol, ir, pmr, cmr, ms, struct*)

Macralstonidine

M-7

[17801-01-3]



C₄₁H₄₈N₄O₃ 644.855

Absolute Configuration

Alkaloid from the bark of *Alstonia macrophylla*, *Alstonia somersetensis* and *Alstonia spectabilis* (Apocynaceae). Needles (EtOH). Mp 295-300° dec. [α]_D²² +166 (CHCl₃). [α]_D +174.5 (c, 1.02 in C₆H₆). λ_{max} 230 (log ε 4.8); 285 (log ε 4.26); 365 (sh) (log ε 3.75) (EtOH).

Dihydrochloride:

Needles (EtOH). Mp 326° dec. [α]_D +136.5 (c, 1.09 in H₂O).

Ac:

Cryst. (MeOH). Mp 205-210° dec.

Sharp, T.M. *et al.*, *J.C.S.*, 1934, 1227-1232 (*isol*)

Kishi, T. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 946-964 (*isol*)

Waldner, E.E. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 1926-1939 (*uv, ir, pmr, ms, struct*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2739-2741 (*isol*)


Das, B.C. *et al.*, *Tet. Lett.*, 1974, 4299-4302 (*cmr*)

Zhao, S. *et al.*, *J.O.C.*, 2003, **68**, 6279-6295 (*synth*)

Macralstonine

M-8

10-(20,21-Dihydro-21-hydroxy-21-methyl-18-noralstophyllan-19-yl)-11-methoxyalstophyllan-19-one, 9CI
[5594-42-3]



Absolute Configuration

C₄₃H₅₂N₄O₅ 704.908

Alkaloid from *Alstonia macrophylla*, *Alstonia muelleriana* and *Alstonia glaberriflora* (Apocynaceae). Cryst. (MeOH or CH₂Cl₂/EtOH). Mp 293° (276-278°, 279-280°) dec. [α]_D²⁰ +74.7 (c, 0.3 in CHCl₃).

Sulfate:

Cryst. (MeOH). Mp 263° (dec.). [α]_D -36.8 (c, 1.022 in H₂O).

Ac: O-Acetylmacralstonine

[6254-08-6]

C₄₅H₅₄N₄O₆ 746.945

Cytotoxic agent. Cryst. (MeOH/Et₂O). Mp 215-222°. [α]_D²⁰ +160 (c, 0.07 in CHCl₃).

Me ether: O-Methylmacralstonine

[5523-51-3]

C₄₄H₅₄N₄O₅ 718.934

Alkaloid from *Alstonia* spp. Cryst. (MeOH/Et₂O or Me₂CO/Et₂O). Mp 202-207°. [α]_D²⁰ +241.4 (c, 0.05 in CHCl₃). λ_{max} 229; 260; 293 (MeOH).

19-Deoxy, 19,20-didehydro, N^{1'}-de-Me: Des-N^{1'a}-methylanhydromacralstonine
[49776-73-0]

C₄₂H₄₈N₄O₄ 672.866

Minor alkaloid from the bark of *Alstonia muelleriana* (Apocynaceae). Needles (MeOH). Mp 240-248° dec. [α]_D +10.2 (c, 0.58 in CHCl₃).

Sharp, T.M. *et al.*, *J.C.S.*, 1934, 1227 (*isol*)

- Kishi, T. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 946 (isol, uv, ir, pmr, ms, struct)
- Cook, J.M. *et al.*, *Phytochemistry*, 1971, **10**, 437 (isol)
- Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2739 (isol)
- Burke, D.E. *et al.*, *Chem. Comm.*, 1972, 1346 (synth)
- Burke, D.E. *et al.*, *Phytochemistry*, 1973, **12**, 1467-1474 (Des-N^{1a}-methylanhidromacralstonine)
- Keawpradub, N. *et al.*, *Planta Med.*, 1997, **63**, 97-101; 1999, **65**, 311-315 (Methylmacralstonine, O-Acetylmacralstonine)
- Liao, X. *et al.*, *Org. Lett.*, 2005, **7**, 3501-3504 (synth)

Macranthine

M-9

[1359-38-2]

C₁₆H₁₉NO₅ 305.33

Struct. unknown. Amaryllidaceae alkaloid. Isol. from fresh leaf sheaths of *Crinum macrantherum* (Amaryllidaceae). Needles (EtOH). Mp 238-240° dec. [α]_D²⁵ -19 (c, 0.235 in CHCl₃). [α]_D²² +49 (c, 0.229 in EtOH).

O-Ac: O-AcetylmacranthineC₁₈H₂₁NO₆ 347.367

Isol. from *Crinum macrantherum* (Amaryllidaceae). Needles (Me₂CO). Mp 222-224°. [α]_D²² -26 (c, 0.213 in CHCl₃). [α]_D²² +18 (c, 0.213 in EtOH).

Di-O-Ac: O,O-DiacetylmacranthineC₂₀H₂₃NO₇ 389.404

Isol. from *Crinum macrantherum* (Amaryllidaceae). Fine needles (Me₂CO/Et₂O). Mp 219-221°. [α]_D²² +44 (c, 0.284 in CHCl₃). [α]_D²² +38 (c, 0.203 in EtOH).

Tri-Ac:


Prisms (C₆H₆/Et₂O). Mp 189-191°. [α]_D²² -2 (c, 0.22 in CHCl₃).

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 185-194 (isol, uv, ir, derivs)

Macrocaffrine

M-10

N^a,18-Didemethyl-19-hydroxy-N^b-methylsuaveoline. Alkaloid RCLM 12. 18-Nor-macrophylline [88660-05-3]

C₁₉H₁₉N₃O 305.379

Struct. originally assigned has been shown to be incorrect by synthesis. Alkaloid from leaves of *Rauwolfia caffra*. Amorph. powder. [α]_D +33.8 (c, 0.3 in MeOH). λ_{max} 225 (log ε 4.32); 273 (log ε 3.77); 283 (log ε 3.75); 291 (log ε 3.64) (MeOH).


Nasser, A.M.A.G. *et al.*, *Phytochemistry*, 1983, **22**, 2297-2300 (isol, pmr, ms)

Ohba, M. *et al.*, *Tetrahedron*, 2007, **63**, 12689-12694 (synth)

Macrocarpamine

M-11

Methyl 19,20-didehydro-2-(18,19-didehydroalstophyllan-18-yl)-2,7-dihydro-1,16-cyclocorynan-17-oate, 9CI [66408-46-6]



Absolute configuration

C₄₁H₄₆N₄O₃ 642.839

Alkaloid from the bark of *Alstonia macrophylla* (Apocynaceae). Cytotoxic. Amorph. [α]_D -16 (c, 3.3 in CHCl₃).

10-Methoxy: 10-Methoxymacrocarpamine

[120374-21-2]

C₄₂H₄₈N₄O₄ 672.866

Alkaloid from the leaves of *Alstonia angustifolia* (Apocynaceae).

10-Methoxy, 4'-N-oxide: 10-Methoxymacrocarpamine 4'-N-oxide

[120374-22-3]

C₄₂H₄₈N₄O₅ 688.865

Alkaloid from the leaves of *Alstonia angustifolia* (Apocynaceae). [α]_D -9 (c, 0.1 in MeOH).

Mayerl, F. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 337 (isol, uv, cd, ir, pmr, cmr, ms, struct)

Ghedira, K. *et al.*, *Phytochemistry*, 1988, **27**, 3955 (isol, uv, ir, pmr, cmr, ms, struct, derivs)


Gan, T. *et al.*, *J.O.C.*, 1998, **63**, 1478-1483 (synth)

Keawpradub, N. *et al.*, *Planta Med.*, 1999, **65**, 311-315 (isol, activity)

Macrocentrine

M-12

[113963-58-9]


C₂₂H₃₅NO₅ 393.522

Alkaloid from *Delphinium macrocentrum* (Ranunculaceae). Tablets (EtOH aq.). Mp 207-209°.

Benn, M.H. *et al.*, *Heterocycles*, 1987, **26**, 2331 (ir, pmr, cmr, ms, struct)

Macrocidin A

M-13



Relative Configuration

C₂₀H₂₃NO₅ 357.405

Tetramic acid deriv. Prod. by *Phoma macrostoma* on *Cirsium arvense*. Phytotoxin. Pale beige solid. [α]_D²⁵ +45 (c, 0.35 in MeOH). λ_{max} 202 ; 226 ; 245 (sh) ; 284 (MeCN).


13ξ-Hydroxy: Macrocidin BC₂₀H₂₃NO₆ 373.405

Prod. by *Phoma macrostoma* on *Cirsium arvense*. Phytotoxin. Pale beige solid.

Graupner, P.R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1558-1561 (isol, pmr, cmr, cryst struct)

Macrodasine A

M-14

C₂₆H₃₄N₂O₅ 454.565

Alkaloid from the bark of *Alstonia macrophylla*. Oil. [α]_D +36 (c, 0.36 in CHCl₃). λ_{max} 230 (log ε 3.88); 287 (log ε 3.17) (EtOH).

23-Ketone: Macrodasine BC₂₆H₃₂N₂O₅ 452.549


Alkaloid from the bark of *Alstonia macrophylla*. Oil. [α]_D +149 (c, 0.07 in CHCl₃). λ_{max} 230 (log ε 3.95); 287 (log ε 3.23) (EtOH).

Kam, T.-S. *et al.*, *Tetrahedron*, 2004, **60**, 3957-3966 (isol, pmr, cmr)

Macrogentine

M-15

[301521-94-8]



Absolute Configuration

C₂₁H₂₆N₂O₃ 354.448


A C16→C5 cyclomacroline alkaloid. Alkaloid from *Alstonia macrophylla*. [α]_D -21 (c, 0.15 in CHCl₃). λ_{max} 208 (log ε 4.44); 258 (log ε 3.74); 284 (log ε 3.15) (EtOH).

Kam, T.-S. *et al.*, *Tetrahedron*, 2000, **56**, 6143-6150


Macrolidine

M-16

[54371-00-5]

C₂₈H₃₂N₂O₁₀ 556.568

Alkaloid from the heartwood of *Adina rubescens* (Rubiaceae). Noncryst. [α]_D²⁵ +2

(CHCl₃) (as N⁴,O,O-tetra-Ac).Brown, R.T. *et al.*, *Chem. Comm.*, 1974, 553 (struct, ms, pmr)**Macroline** M-1720,21-Didehydro-17-hydroxy-1,4-dimethyl-4,21-secosarpagan-19(20H)-one, 9CI
[2269-93-4]


Absolute Configuration

C₂₁H₂₆N₂O₂ 338.449Important degradn. prod. of the dimeric *Alstonia* alkaloids, apparently not yet isol. as a native alkaloid. Mp 211-213° dec. [α]_D²² +19 (c, 0.406 in MeOH). λ_{max} 231 (log ε 4.56); 282 (log ε 3.67); 288 (log ε 3.66) (EtOH).

Ac:

Cryst. (MeOH/Et₂O). Mp 180°.Hesse, M. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 689-710; 1966, **49**, 1173-1182 (uv, ir, pmr, ms, struct)Esmond, R.W. *et al.*, *J.A.C.S.*, 1980, **102**, 7116-7117 (synth)Neukomm, G. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 90-96 (abs config)Zhang, L.H. *et al.*, *J.A.C.S.*, 1990, **112**, 4088-4090 (synth)Bi, Y. *et al.*, *J.A.C.S.*, 1994, **116**, 9027-9041 (synth)Tran, Y.S. *et al.*, *Org. Lett.*, 2005, **7**, 4289-4291 (synth)Liao, X. *et al.*, *J.O.C.*, 2006, **71**, 8884-8890 (synth)Lewis, S.E. *et al.*, *Tetrahedron*, 2006, **62**, 8655-8681 (rev, chem)**Macromerine** M-18

α-[(Dimethylamino)methyl]-3,4-dimethoxybenzenemethanol, 9CI. 1-(3,4-Dimethoxyphenyl)-2-dimethylaminoethanol



(R)-form

C₁₂H₁₉NO₃ 225.287

(R)-form [19751-75-8]

Alkaloid from *Coryphantha* (*Thelocactus*) *macromeris*, *Coryphantha cornifera*, *Coryphantha elephantidens* and *Coryphantha pectinata* (Cactaceae). Mp 66-67.5°. [α]_D²⁵ -42.6 (c, 2 in EtOH). [α]_D²⁵ -147 (c, 0.039 in CHCl₃).

N-De-Me: Normacromerine. N-Methyl-3,4-dimethoxy-β-hydroxyphenethylamine

[41787-64-8]

C₁₁H₁₇NO₃ 211.26Alkaloid from *Coryphantha macro-**meris*, *Coryphantha calipensis*, *Coryphantha greenwoodii*, *Desmodium tiliacifolium* and *Dolichothele longimamma* (Cactaceae). Cryst. or brown gum. Mp 101-103°. [α]_D²⁴ -38.4 (c, 0.41 in CHCl₃).N-De-Me; hydrochloride: Mp 131-132°. [α]_D²⁵ -51.3.

N-De-Me, N-formyl: N-[2-(3,4-Dimethoxyphenyl)-2-hydroxyethyl]-N-methylformamide, 9CI. N-Formylnormacromerine

[41688-37-3]

C₁₂H₁₇NO₄ 239.271Alkaloid from *Coryphantha macromeris* (Cactaceae).

Me ether, β,3,4-Trimethoxy-N,N-dimethylbenzeneethanamine, 9CI. O-Methylmacromerine

[52910-84-6]

C₁₃H₂₁NO₃ 239.314Alkaloid from *Coryphantha calipensis* (Cactaceae). Mp 178° (as hydrochloride). [α]_D²⁵ -94 (c, 0.01 in EtOH).

Me ether, N-de-Me: β,3,4-Trimethoxy-N-methylbenzeneethanamine, 9CI. Calipamine. O-Methylnormacromerine

[52910-83-5]

C₁₂H₁₉NO₃ 225.287Alkaloid from *Coryphantha greenwoodii* and *Coryphantha calipensis* (Cactaceae). Mp 213-214° (206-208°) (as hydrochloride). [α]_D²⁵ -92 (c, 0.01 in EtOH).

(S)-form [33066-27-2]

Synthetic. Mp 59-62°. [α]_D²⁶ +34 (c, 0.035 in EtOH).

N-Di-de-Me, N,O-di-E-cinnamoyl: Syncarpamide

C₂₈H₂₇NO₅ 457.525Alkaloid from the stems of *Zanthoxylum syncarpum*. Antiplasmodial agent. Light brown solid. [α]_D²⁵ +12.5 (c, 0.08 in CHCl₃). λ_{max} 242; 250; 278 (MeOH).

(±)-form [33122-27-9]

Mp 47.5-48.5°.

Picrate: Mp 157° (147-148°).

N-De-Me; hydrochloride: Mp 132-133°.

N-Di-de-Me: α-(Aminomethyl)-3,4-dimethoxybenzenemethanol. 2-Amino-1-(3,4-dimethoxyphenyl)ethanol

[4899-03-0]

C₁₀H₁₅NO₃ 197.233Needles (C₆H₆/petrol). Mp 89-91°.

N-Di-de-Me, N,O-di-E-cinnamoyl: (±)-Syncarpamide

C₂₈H₂₇NO₅ 457.525Alkaloid from the leaves of *Zanthoxylum syncarpum*. Pale yellow powder. λ_{max} 242 (log ε 4.83); 252 (log ε 4.69); 278 (log ε 4.94) (MeOH).

(ξ)-form

N-Di-de-Me, N-benzoyl: N-[2-(3,4-Dimethoxyphenyl)-2-hydroxyethyl]benzamide. Balsamide

[70216-31-8]

[128820-23-5]

C₁₇H₁₉NO₄ 301.341Isol. from *Amyris balsamifera* (Rutaceae). Prisms. Mp 135-136°.


Me ether, N-di-de-Me, N-benzoyl: N-[2-(3,4-Dimethoxyphenyl)-2-methoxyethyl]benzamide. O-Methylbalsamide

C₁₈H₂₁NO₄ 315.368Alkaloid from the stem bark of *Zanthoxylum hyemale*. Needles. Mp 110-112° (natural) Mp 124° (synthetic). [α]_D +8.8 (c, 0.01 in CHCl₃).

[33066-28-3, 56120-35-5, 33066-32-9]

Keller, W.J. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 147; 1973, **62**, 408 (derivs)Brown, S.D. *et al.*, *J.O.C.*, 1972, **37**, 773 (isol, ir, uv, pmr, ms, synth)Ghosal, S. *et al.*, *Phytochemistry*, 1973, **12**, 193 (isol, deriv)Butterick, J.R. *et al.*, *Can. J. Chem.*, 1974, **52**, 2873 (synth)Bruhn, J.G. *et al.*, *J. Pharm. Sci.*, 1974, **63**, 574 (isol, derivs)Ranieri, R.L. *et al.*, *J. Nat. Prod.*, 1976, **39**, 172 (deriv)Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (occur)Woodward, R.W. *et al.*, *Acta Chem. Scand., Ser. B*, 1978, **32**, 619 (cd, abs config, Calipamine)Burke, B. *et al.*, *Heterocycles*, 1979, **12**, 349-351 (Balsamide)Keller, W.J. *et al.*, *J. Pharm. Sci.*, 1979, **68**, 85 (biosynth)Quintard, J.P. *et al.*, *Synthesis*, 1984, 495 (synth)Estevez, J.C. *et al.*, *Synth. Commun.*, 1990, **20**, 503 (Balsamide, synth)Zhao, X. *et al.*, *Org. Prep. Proced. Int.*, 1995, **27**, 513-517 (di-N-de-Me)De Moura, N.F. *et al.*, *Planta Med.*, 2002, **68**, 534-538 (O-Methylbalsamide)Ross, S.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 88-90; 2005, **68**, 1297-1299 (Syncarpamide)**Macrophominol** M-19


2,3-Dihydro-3-(1-hydroxyethyl)pyrrolo[1,2-a]pyrazine-1,4-dione [172335-92-1]



Absolute Configuration

C₉H₁₀N₂O₃ 194.19Isol. from cultures of the phytopathogenic fungus *Macrophomina phaseolina*. Phytotoxic agent. Mp 116-118°. [α]_D²⁵ -25.24 (c, 0.103 in EtOH).Trigos, A. *et al.*, *Phytochemistry*, 1995, **40**, 1697-1698 (isol, pmr, cmr, ms, struct)**Macrophylline†** M-20

[76372-23-1]

C₂₀H₂₁N₃O 319.405Alkaloid from *Rauwolfia caffra* and

Rauwolfia macrophylla. Amorph. powder. $[\alpha]_D^{25} +33.8$ (c, 0.3 in MeOH). λ_{\max} 225 (log ϵ 4.32); 273 (log ϵ 3.77); 283 (log ϵ 3.75); 291 (log ϵ 3.64) (MeOH).

Amer, M.M.A. *et al.*, *Planta Med.*, Suppl., 1980, 8-12 (*isol*)

Nasser, A.M.A.G. *et al.*, *Phytochemistry*, 1983, 22, 2297-2300 (*isol, uv, ir*)

Macrophylline† M-21

C₄₅H₅₄N₄O₅ 730.945

Bisindole alkaloid. Struct. unknown. Alkaloid from *Alstonia macrophylla* (Apocynaceae). Mp 267-268° dec.

Picrate: Mp 199.5-200° dec.

Manas-Santos, F. *et al.*, *CA*, 1937, 31, 6243 (*isol*)

Macrophylline A† M-22

[1359-43-9]


C₂₀H₂₃N₂O₂[⊕] 323.414

Struct. unknown. Alkaloid from *Strychnos macrophylla* (Loganiaceae). Mp 300° (as picrate)(browns from 200°). Violet col. with Ce(SO₄)₂.

Iorio, M.A. *et al.*, *Gazz. Chim. Ital.*, 1956, 86, 923

Macropodumine A M-23

[908008-85-5]




C₂₁H₂₇NO₅ 373.448

Alkaloid from the stems of *Daphniphyllum macropodum*. Light yellow oil. $[\alpha]_D^{20} +30$ (c, 0.25 in MeOH). λ_{\max} 210 (ϵ 3260); 239 (ϵ 6245) (MeOH).

Zhang, W. *et al.*, *Chem. Eur. J.*, 2006, 12, 5122-5127 (*isol, pmr, cmr, ms*)

Macropodumine B M-24

[908008-86-6]




C₂₃H₂₇NO₅ 397.47

Zwitterion showing delocalisation of the negative charge. Alkaloid from the stems of *Daphniphyllum macropodum*. Light yellow cryst. (CHCl₃/MeOH). Mp 245-247°. $[\alpha]_D^{20} -235$ (c, 0.25 in MeOH). λ_{\max} 210 (ϵ 7940); 245 (ϵ 2980); 295 (ϵ 14890); 323 (ϵ 7150); 362 (ϵ 13300) (MeOH).

Zhang, W. *et al.*, *Chem. Eur. J.*, 2006, 12, 5122-5127 (*isol, pmr, cmr, cryst struct*)

Macropodumine C M-25

[908008-87-7]




C₂₃H₂₉NO₆ 415.485

Alkaloid from the stems of *Daphniphyllum macropodum*. Light yellow oil. $[\alpha]_D^{20} -156$ (c, 0.17 in MeOH). λ_{\max} 209 (ϵ 7260); 250 (ϵ 2700); 301 (ϵ 15150); 325 (ϵ 7055); 367 (ϵ 14110) (MeOH).

Zhang, W. *et al.*, *Chem. Eur. J.*, 2006, 12, 5122-5127 (*isol, pmr, cmr, ms*)

Macropodumine D M-26

[925462-82-4]



Relative Configuration


C₂₃H₃₁NO₅ 401.502

Alkaloid from *Daphniphyllum macropodum*. Oil. $[\alpha]_D^{20} -10.6$ (c, 0.24 in CHCl₃).

Li, Z.-Y. *et al.*, *Org. Lett.*, 2007, 9, 477-480 (*isol, pmr, cmr*)

Macropodumine E M-27

[925462-83-5]




C₂₇H₃₅NO₈ 501.575

Alkaloid from *Daphniphyllum macropodum*. Oil. $[\alpha]_D^{20} -51.1$ (c, 0.47 in CHCl₃).

Li, Z.-Y. *et al.*, *Org. Lett.*, 2007, 9, 477-480 (*isol, pmr, cmr*)

Macrorungine M-28

1,3-Dihydro-1-methyl-4-(2-quinoliny)-2H-imidazol-2-one, 9CI
[2552-98-9]



C₁₃H₁₁N₃O 225.249

Alkaloid from the aerial parts of *Macrorungia longistrobus* (preferred genus name

Anisotes) (Acanthaceae). Yellow prisms (MeOH). Mp 267-270° dec.

Hydrochloride:

Red needles (MeOH). Mp 290° dec.

Perchlorate:

Yellow plates + 0.5 H₂O (MeOH). Mp 290° dec.

Picrate:


Cryst. (MeOH). Mp 250° dec.

Arndt, R.R. *et al.*, *J.C.S.*, 1964, 5969 (*isol, uv, ir, struct*)

Joynt, V.P. *et al.*, *J.C.S. (B)*, 1966, 980 (*ms*)

Macrosalhin M-29

17,19-Epoxy-19,20-dihydro-19-hydroxy-1,4,21-trimethyl-18-norsarpaganium, 9CI
[16103-76-7]



Absolute Configuration

C₂₁H₂₇N₂O₂[⊕] 339.456

Minor quaternary alkaloid from the stem bark of *Alstonia macrophylla* (Apocynaceae).

Chloride: [16049-34-6]

Mp 284-286° dec. $[\alpha]_D^{24} +27$ (c, 0.467 in MeOH).

Thiocyanate: [16103-77-8]


Mp 300°. $[\alpha]_D^{25} +27.6$ (c, 0.284 in MeOH).

Khan, Z.M. *et al.*, *Helv. Chim. Acta*, 1967, 50, 1002-1010 (*isol, uv, ir, pmr, ms, struct*)

Wulf, H. *et al.*, *Helv. Chim. Acta*, 1967, 50, 1011-1012 (*cryst struct*)

Macrospetratine M-30

[113775-81-8 (chloride/monohydrochloride salt)]



Absolute Configuration

C₄₀H₄₅N₄O₃[⊕] 629.82

Quaternary alkaloid from the roots of *Rauwolfia verticillata* (Apocynaceae). Cryst.(as chloride hydrochloride). Mp 300° dec.(chloride hydrochloride). $[\alpha]_D^{17} +215$ (CHCl₃) (chloride hydrochloride).

Lin, X. *et al.*, *Jiegou Huaxue*, 1987, 6, 89-93; *CA*, 108, 167761j (*cryst struct*)

Lin, M. *et al.*, *Yaoxue Xuebao*, 1987, 22, 833-836; *CA*, 108, 164701d

C₂₀H₂₇NO₇ 393.436

Alkaloid from *Crotalaria agatiflora* and *Crotalaria rosenii* (Fabaceae). Prisms (EtOH). Mp 179-180° (158-160°). [α]_D²⁵ +68 (c, 0.60 in EtOH). [α]_D²⁰ +66 (c, 2 in CHCl₃).

(15*Z*)-Isomer, *O*⁷-Ac: **Acetyl-cis-madurensine**

C₂₀H₂₇NO₇ 393.436

Alkaloid from *Crotalaria agatiflora* (Fabaceae). Fine needles (petrol). Mp 156-157°. [α]_D²⁰ +102.3 (c, 0.39 in EtOH).

18-Hydroxy: **Crotaflorine**

[112156-54-4]

C₁₈H₂₅NO₇ 367.398

Alkaloid from *Crotalaria agatiflora* and *Crotalaria rosenii* (Fabaceae). Prisms (EtOH). Mp 179-180°. [α]_D²⁰ +50 (c, 0.51 in EtOH). [α]_D²⁵ +32 (c, 2 in CHCl₃). Ester of crotanecine with retronecic acid.

Atal, C.K. *et al.*, *Tet. Lett.*, 1966, 537 (*pmr*)
Culvenor, C.C.J. *et al.*, *Chem. Comm.*, 1970, 65 (*struct*)

Culvenor, C.C.J. *et al.*, *An. Quim.*, 1972, **68**, 883 (*isol, pmr, ms, derivs*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (*cd*)

Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417 (*cmr*)


Mackay, M.F. *et al.*, *Acta Cryst. C*, 1984, **40**, 1073 (*cryst struct*)

Abegaz, B. *et al.*, *Tetrahedron*, 1987, **43**, 3263 (*isol, ir, pmr, cmr, ms, derivs*)

Maedamine A

M-37

[310396-69-1]

C₂₃H₂₄Br₃N₃O₃ 630.173

Alkaloid from the sponge *Suberea* sp. Cytotoxic agent. Amorph. yellow solid. λ_{max} 285 (ε 4000); 350 (ε 1000) (MeOH).

N-De-Me: **Maedamine B**

[310396-70-4]

C₂₂H₂₂Br₃N₃O₃ 616.146


Alkaloid from *Suberea* sp. Cytotoxic agent. Amorph. yellow solid. λ_{max} 285 (ε 5000); 350 (ε 1200) (MeOH).

Hirano, K. *et al.*, *Tetrahedron*, 2000, **56**, 8107-8110 (*Maedamines A,B*)

Maeganedin A

M-38

[205247-54-7]

C₃₇H₅₂N₄O₂ 584.843


Related to Manzamine A, M-91. Alkaloid from the sponge *Amphimedon* sp. Amorph. solid. [α]_D²⁵ +47 (c, 0.4 in MeOH). λ_{max} 205 (ε 13000); 223 (ε 2500); 274 (ε 2500); 322 (ε 4000) (MeOH).

Tsuda, M. *et al.*, *Tet. Lett.*, 1998, **39**, 1207-1210 (*isol, uv, ir, pmr, cmr*)

Magallanesine

M-39

[75767-34-9]

C₂₁H₁₇NO₆ 379.368

Alkaloid from the stems and branches of *Berberis darwinii* (Berberidaceae). Yellow cryst. (MeOH). Mp 254-256°. The first isoindolobenzazocine alkaloid. Prob. an artifact.

Manikumar, G. *et al.*, *J.O.C.*, 1981, **46**, 386 (*synth*)

Valencia, E. *et al.*, *Tet. Lett.*, 1985, **26**, 993 (*uv, ir, pmr, cmr, ms, struct*)

Shamma, M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 398


Fang, F.G. *et al.*, *Tet. Lett.*, 1989, **30**, 2743 (*synth*)

Yoneda, R. *et al.*, *Tetrahedron*, 1996, **52**, 14563 (*synth*)

Magellanine†

M-40

[61273-75-4]



Absolute Configuration

C₁₇H₂₅NO₂ 275.39

Alkaloid from *Lycopodium magellanicum* (Lycopodiaceae). Mp 165-166°.

Methbromide: Mp 274°.

Ketone: **Magellaninone**

[73885-47-9]

C₁₇H₂₃NO₂ 273.374

Alkaloid from *Lycopodium magellanicum* (Lycopodiaceae). Oil; Cryst. (Me₂CO)(as perchlorate). Mp 244-245° (perchlorate).

Castillo, M. *et al.*, *Can. J. Chem.*, 1976, **54**, 2893 (*isol, ir, pmr, ms, cryst struct*)

Loyola, L.A. *et al.*, *Phytochemistry*, 1979, **18**, 1721 (*Magellaninone*)

Hirst, G.C. *et al.*, *J.A.C.S.*, 1993, **115**, 2992 (*synth*)

Paquette, L.A. *et al.*, *J.A.C.S.*, 1993, **115**, 4377 (*synth*)

Williams, J.P. *et al.*, *J.A.C.S.*, 1994, **116**, 4689 (*synth*)

Yen, C.-F. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 4090-4093 (*synth*)

Ishizaki, M. *et al.*, *Tetrahedron*, 2005, **61**, 4053-4065 (*synth*)

Kozaki, T. *et al.*, *J.O.C.*, 2007, **72**, 10147-10154 (*synth*)

Magnarcine

M-41

[1359-45-1]

C₁₇H₂₁NO₄ 303.357

Amaryllidaceae alkaloid. Struct. unknown. Isol. from several *Narcissus* hybrids and from *Galanthus nivalis* (Amaryllidaceae). Cryst. (MeOH or Me₂CO). Mp 221° Mp 253-254° dec.

Perchlorate:

Prisms (EtOH/Et₂O). Mp 285° dec.


Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 2197 (*isol*)

Boit, H.G. *et al.*, *Naturwissenschaften*, 1959, **46**, 228; 1960, **47**, 109 (*isol*)

Magnesidin

M-42

[52081-52-4]



n = 4,6

A mixt. of Mg salts of 2 tetramic acids. Obt. from *Pseudomonas magnesiorubra* ATCC 21856 and a marine *Vibrio gazogenes*. Active against gram-positive bacteria, particularly the spore bearers. Sol. MeOH, Et₂O; poorly sol. H₂O. Mol. formula C₂₈H₃₆MgN₂O₈/C₃₂H₄₄MgN₂O₈. λ_{max} 257 (ε 19800) (MeOH) (Derep). λ_{max} 261 (E1%/1cm 928) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 50 mg/kg. OM2457600

Gandhi, N.M. *et al.*, *J. Antibiot.*, 1973, **26**, 797 (*isol*)


U.K. Pat., 1977, 1 478 643; CA, **88**, 168480 (*synth, props*)

Imamura, N. *et al.*, *J. Antibiot.*, 1994, **47**, 257-261 (*isol, pmr, cmr, ms*)

Magniflorine

M-43

[139610-18-7]

C₂₀H₂₀N₂O₃ 336.39


Not to be confused with Magnoflorine in C-702. Alkaloid from the fruit of *Hamelia magniflora* (Rubiaceae). Prisms (EtOAc). Mp 168-170°. [α]_D²⁰ -271 (c, 1 in CHCl₃).

Rumbero, A. *et al.*, *Tet. Lett.*, 1991, **32**, 5153-5154 (*isol, uv, ir, pmr, cmr, ms, struct*)

Magnolamide

M-44

[212703-85-0]



C₂₀H₂₄N₂O₅ 372.42

Alkaloid from the leaves of *Magnolia coco*. Antioxidant. Pale yellow oil. λ_{\max} 202 (log ϵ 4.5); 236 (log ϵ 4.66); 294 (log ϵ 4.43) (MeOH).

Yu, H.-J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1017-1019 (*isol*, *pmr*, *cmr*)


Dong, Y. *et al.*, *Heterocycles*, 2002, **56**, 221-225 (*synth*, *pmr*, *cmr*)

Chiou, W.-F. *et al.*, *Heterocycles*, 2005, **65**, 1215-1220 (*synth*, *activity*)

Magnolamine

M-45

[573-73-9]



C₃₇H₄₂N₂O₇ 626.748

Alkaloid from the leaves of *Magnolia fuscata* and *Michelia fuscata* (Magnoliaceae). Needles (C₆H₆). Mp 117-119°. $[\alpha]_D$ +111.6 (EtOH).

Picrate: Mp 142-145° dec.

Picrolonate: Mp 163-164° dec.

Proskurnina, N. *et al.*, *Bull. Soc. Chim. Fr.*, Sér. 5, 1938, 1357 (*isol*)

Tomita, M. *et al.*, *Pharm. Bull.*, 1954, **2**, 115 (*struct*)


Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592 (*ms*)
Yakhontova, L.D. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 234; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 200 (*struct*, *ms*, *pmr*)

Tanaka, H. *et al.*, *Heterocycles*, 1981, **16**, 1275 (*occur*, *ir*, *pmr*, *synth*, *struct*, *bibl*)

Mahanimbicine

M-46

3,11-Dihydro-3,8-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole, 9CI. *Isomahanimbicine*



C₂₃H₂₅NO 331.457

(+)-*form* [28305-77-3]

Alkaloid from the leaves of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 142°. $[\alpha]_D^{30}$ +18.6 (c, 0.86 in CHCl₃). λ_{\max} 277 (log ϵ 4.71); 288 (log ϵ 5); 315 (log ϵ 3.84); 335 (log ϵ 3.99); 350 (log ϵ 3.97) (MeOH). λ_{\max} 239 (log ϵ 4.68); 291 (log ϵ 4.63); 339 (log ϵ 3.82); 353 (log ϵ 3.8) (EtOH).

7-Hydroxy: *Pyrafoline D*. *Isomahanine* [138876-26-3]

C₂₃H₂₅NO₂ 347.456

Alkaloid from the stem bark of *Murraya euchrestifolia* and seeds of *Murraya koenigii* (curryleaf tree) (Rutaceae). Shows cytotoxic activity. Pale brown powder, *cryst.* (CH₂Cl₂/petrol). Mp 184°. $[\alpha]_D$ 0 (MeOH). *Pyrafoline D* descr. as brown powder,

Isomahanine as *cryst.* with indicated Mp. Opt. rotn. refers to *Pyrafoline D*. The samples have not been compared. λ_{\max} 220 (log ϵ 4.51); 236 (log ϵ 4.48); 286 (sh) (log ϵ 4.29); 294 (log ϵ 4.52); 324 (log ϵ 3.68); 338 (log ϵ 3.74) (MeOH).

(±)-*form* [28305-78-4]

Synthetic. Mp 140°. Racemate.

8-Methoxy: *Murrayamine C*†

[134779-19-4]

C₂₄H₂₇NO₂ 361.483

Alkaloid from leaves of *Murraya euchrestifolia* (Rutaceae). Syrup. λ_{\max} 240; 274; 284; 340; 353 (MeOH).

[26871-46-5]

Kureel, S.P. *et al.*, *Chem. Ind. (London)*, 1970, 958 (*uv*, *ir*, *pmr*, *struct*, *synth*)

Joshi, B.S. *et al.*, *Tetrahedron*, 1970, **26**, 1475-1482 (*isol*, *uv*, *ir*, *pmr*, *struct*)

Narasimhan, N.S. *et al.*, *Indian J. Chem.*, 1975, **13**, 993-999 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Narasimhan, N.S. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 329-331 (*synth*)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1668-1671 (*Pyrafoline D*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1991, **30**, 1048-1051 (*Murrayamine C*)


Reisch, J. *et al.*, *Phytochemistry*, 1992, **31**, 2877-2879 (*Isomahanine*)

Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893-897 (*activity*)

Mahanine

M-47

3,11-Dihydro-3,5-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazol-9-ol, 9CI



(R)-*form*

C₂₃H₂₅NO₂ 347.456

Data on opt. rotn. and abs. configs. for these compds. is mostly unsatisfactory.

(R)-*form* [28360-49-8]

Alkaloid from the leaves of *Murraya koenigii* (curry leaf tree) (Rutaceae). Shows cytotoxic activity. Fluffy *cryst.* (petrol). Mp 98-100°. $[\alpha]_D$ -24.4. λ_{\max} 240 (ϵ 4.46); 263 (ϵ 3.89); 286 (ϵ 4.32); 295 (ϵ 4.57); 300 (ϵ 4); 340 (ϵ 3.73); 360 (ϵ 3.63) (MeOH).

(S)-*form* [134876-75-8]

From leaves of *Murraya euchrestifolia* (Rutaceae). Cytotoxic agent. Needles (Me₂CO). Mp 95-96°. $[\alpha]_D$ +34 (c, 2 in CHCl₃). λ_{\max} 221 (ϵ 74130); 240 (ϵ 81200); 285 (sh) (ϵ 47860); 294 (ϵ 64600); 325 (ϵ 4500); 342 (ϵ 4800); 357 (ϵ 3450) (MeOH).

(±)-*form*

Deoxy: (±)-*Mahanimbicine*

[24948-14-9]

C₂₃H₂₅NO 331.457

Obt. by heating (+)-mahanimbicine at 150°. Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). *Cryst.* (hexane). Mp 72-73° Mp 75-76° Mp 94-95°. λ_{\max} 225 (log

4.6); 238 (log ϵ 4.7); 287 (log ϵ 4.6); 326 (log ϵ 4) (EtOH).

Deoxy, 10-methoxy: 3,11-Dihydro-10-methoxy-3,5-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole.

Murrayamine B

[134779-18-3]

C₂₄H₂₇NO₂ 361.483

Alkaloid from leaves of *Murraya euchrestifolia* (Rutaceae). Syrup.

(±)-*form*

Deoxy: (+)-*Mahanimbicine*

[21104-28-9]

C₂₃H₂₅NO 331.457

Alkaloid from the fruits and leaves of *Murraya koenigii* (curryleaf tree) and the stem bark of *Murraya exotica* (Rutaceae). *Cryst.* (hexane). Mp 94-95°. $[\alpha]_D$ +40.6 (c, 2.1 in CHCl₃). $[\alpha]_D$ +52 (CHCl₃). Abs. config. not certain. λ_{\max} 223 (log ϵ 4.55); 239 (log ϵ 4.66); 288 (log ϵ 4.61); 330 (log ϵ 3.9); 343 (log ϵ 3.42) (EtOH).

Deoxy, N-Me:

Needles (EtOH). Mp 120°.

Deoxy, 1'-oxo: 3,11-Dihydro-3-methyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole-5-carboxaldehyde. *Murrayamine*

[53508-00-2]

C₂₃H₂₃NO₂ 345.44

Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 105°. Aldehyde corresponding to Mahanimbicine. λ_{\max} 234 (log ϵ 4.6); 280 (log ϵ 4.5); 301 (log ϵ 4.58); 312 (log ϵ 4.5) (EtOH).

Deoxy, 3',4'-dihydro, 4'-hydroxy: *Mahanimbicine*

[30048-24-9]

C₂₃H₂₇NO₂ 349.472

Alkaloid from leaves of *Murraya koenigii* (curryleaf tree) (Rutaceae). Mp 179°. λ_{\max} 238 (log ϵ 4.64); 288 (log ϵ 4.61); 329 (log ϵ 3.83); 344 (log ϵ 3.87); 359 (log ϵ 3.82) (EtOH).

Deoxy, Δ⁴-isomer, 3'-hydroxy: *Murrayamine C*†. *Mahanimboline*

[73385-58-7]

C₂₃H₂₅NO₂ 347.456

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Pale yellow oil. This *struct.* was formerly assigned erroneously to Mahanimbicine. The pmr data of the compd. *isol.* by Roy *et al.* (1979) are different from those reported for *Murrayamine C* (see Mahanimbicine, M-46), which has now been assigned this *struct.*

Kureel, S.P. *et al.*, *Experientia*, 1970, **26**, 1055 (*Mahanimbicine*)

Joshi, B.S. *et al.*, *Tetrahedron*, 1970, **26**, 1475-1482 (*Mahanimbicine*, *isol*, *uv*, *ir*, *pmr*, *struct*)

Chakraborty, D.P. *et al.*, *Chem. Ind. (London)*, 1974, 165-166 (*Murrayamine*, *isol*, *uv*, *ir*, *ms*, *struct*, *synth*)

Bandranayake, W.M. *et al.*, *J.C.S. Perkin 1*, 1974, 998-1007 (*Mahanimbicine*, *synth*)

Roy, S. *et al.*, *Phytochemistry*, 1974, **13**, 2893 (*Mahanimbicine*, *isol*, *uv*, *ir*, *pmr*)


Narasimhan, N.S. *et al.*, *Indian J. Chem.*, 1975, **13**, 993-999; *Indian J. Chem., Sect. B*, 1976,

14, 329-331; 430-433 (*Mahanine, Mahanimbine, isol, uv, ir, pms, ms, struct, synth*)

- Bhattacharyya, P. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 308-309 (*Mahanimbine*)
 Roy, S. *et al.*, *Chem. Ind. (London)*, 1979, 669-670 (*Mahanimboline*)
 Roy, S. *et al.*, *J. Indian Chem. Soc.*, 1980, **57**, 759 (*Murrayayinine, synth*)
 Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1988, **59**, 494-495 (*pmr, cmr*)
 Wu, T.-S. *et al.*, *Phytochemistry*, 1991, **30**, 1048-1051 (*Mahanine, Murrayamine B, isol, pmr*)
 Ito, C. *et al.*, *Phytochemistry*, 1992, **31**, 1083-1084 (*Murrayamine C*)
 Meragelman, K.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 427-428 (*Mahanimbine, isol, cmr*)
 Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893-897 (*activity*)

Maingayinine

[863890-49-7]




$C_{10}H_{15}N_3O_4$ 241.246
 Alkaloid from the twigs of *Mitrephora maingayi*. Amorph. powder. Mp 135-137°. $[\alpha]_D^{25}$ -12.8 (c, 0.2 in MeOH).

Yu, R. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 359-362 (*isol, pmr, cmr, ms*)

Maistemone

Protostemotinine

[137031-44-8]
 [169534-85-4 (Protostemotinine)]



Absolute Configuration

$C_{23}H_{29}NO_6$ 415.485
 Alkaloid from the roots of *Stemona mairei* and *Stemona sessilifolia*.

8 β -Hydroxy- **Oxymaistemone**

[137031-45-9]

 $C_{23}H_{29}NO_7$ 431.485Alkaloid from the roots of *Stemona mairei*.2'-Epimer: **Isomaistemone**

[263870-90-2]

 $C_{23}H_{29}NO_6$ 415.485Alkaloid from the roots of *Stemona japonica*.

Lin, W. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 369-370 (*Maistemone, Oxymaistemone*)

Cong, X. *et al.*, *Phytochemistry*, 1995, **40**, 615-617 (*Protostemotinine*)


Zou, C. *et al.*, *J. Chin. Pharm. Sci.*, 1999, **8**, 185-190 (*Isomaistemone*)

Greger, H. *et al.*, *Planta Med.*, 2006, **72**, 99-113 (rev)

Majdine

Herbavinine

[20497-42-1]



Absolute Configuration

$C_{23}H_{28}N_2O_6$ 428.484
 Alkaloid from *Vinca herbacea*, *Vinca elegantissima*, *Vinca erecta*, and *Vinca major* (Apocynaceae). Mp 190-192°. $[\alpha]_D^{24}$ -108 (Py). λ_{max} 225 (log ϵ 4.57); 248 (sh) (log ϵ 4.23); 285 (sh) (log ϵ 3.16) (EtOH).

O¹¹-De-Me: Vinerinine

[54347-88-5]

 $C_{22}H_{26}N_2O_6$ 414.457

Alkaloid from *Vinca erecta* (Apocynaceae). Amorph. (?). $[\alpha]_D^{20}$ -74. λ_{max} 224 (log ϵ 4.44) (EtOH).

3-Epimer: **Elegantissine. Herboxine. Herboksine**

[57495-67-7]

 $C_{23}H_{28}N_2O_6$ 428.484

Alkaloid from *Vinca elegantissima* and *Vinca herbacea* (Apocynaceae). Amorph. $[\alpha]_D$ +3.4 (CHCl₃). λ_{max} 224 (log ϵ 4.39); 250 (sh) (no solvent reported).

7-Epimer: **Isomajdine. Elegantine**

[20497-41-0]

 $C_{23}H_{28}N_2O_6$ 428.484

Alkaloid from the *Vinca herbacea* and *Vinca elegantissima* (Apocynaceae). Mp 208-210°. $[\alpha]_D^{24}$ -111 (Py). λ_{max} 225 (log ϵ 4.53); 248 (sh) (log ϵ 4.16); 285 (sh) (log ϵ 3.04) (EtOH).

3,7-Diepimer: **Isoelegantissine**

[57495-68-8]

 $C_{23}H_{28}N_2O_6$ 428.484

Alkaloid from *Vinca elegantissima* (Apocynaceae). Mp 190°. $[\alpha]_D$ +5.7 (CHCl₃).

Stereoisomer (1): **Herbavine**

[27274-48-2]

 $C_{23}H_{28}N_2O_6$ 428.484

Alkaloid from *Vinca herbacea* (Apocynaceae). Stereochem. unknown.

Stereoisomer (1), parent acid: **16-Carboxyherbavine**

[58031-34-8]

 $C_{22}H_{26}N_2O_6$ 414.457

Alkaloid from *Vinca herbacea* (Apocynaceae). Corresponds to Herbavine.

Stereoisomer (2): **Isoherbavine**

[56832-55-4]

 $C_{23}H_{28}N_2O_6$ 428.484

Alkaloid from *Vinca herbacea* (Apocynaceae). Stereochem. unknown.

Stereoisomer (2), parent acid: **16-Carboxyisoherbavine**

[74924-37-1]

 $C_{22}H_{26}N_2O_6$ 414.457

Alkaloid from *Vinca herbacea* (Apocynaceae). Corresponds to Isoherbavine.

Ognyanov, I. *et al.*, *Tetrahedron*, 1968, **24**, 4641-4648 (*Majdine, Isomajdine, uv, pmr, ms, struct, stereochem*)

M-50

Bhattacharyya, J. *et al.*, *Tet. Lett.*, 1972, 159-162 (*Elegantine*)

Khalmirzoev, M.M. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 411-412; 1977, **13**, 718-719; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 419; 1977, **13**, 605 (*Vinerinine*)

Ali, E. *et al.*, *Experientia*, 1975, **31**, 876-877 (*Elegantissine, Isoelegantissine, isol, cd, struct*)

Dzhakeli, E.Z. *et al.*, *CA*, 1976, **84**, 44499h; 1980, **93**, 146290m (*Herbavine, Isoherbavine*)

Chkhivadze, G.V. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 227-229; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 201-202

(*Herboxine, isol, uv, ir, pmr, ms, struct*)
 Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 217-224; *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 170-176 (*cmr*)

Majorine**M-51** $C_{24}H_{26}N_2O_3$ 390.481


Struct. unknown. Alkaloid from the aerial parts of *Vinca major* (Apocynaceae). Small needle-shaped cryst. (MeOH). Mp 247-249° dec. $[\alpha]_D^{20}$ -265 (c, 0.63 in CHCl₃).

Kaul, J.L. *et al.*, *J. Nat. Prod.*, 1966, **29**, 26-34 (*isol, uv, ir*)

Majovine**M-52**

Struct. unknown. Alkaloid from the aerial parts of *Vinca major* (Apocynaceae). Feathery cryst. (MeOH). Mp 227-229°. $[\alpha]_D^{20}$ +133 (c, 0.64 in CHCl₃).

Kaul, J.L. *et al.*, *J. Nat. Prod.*, 1966, **29**, 26-34 (*isol, uv, ir*)

Makalika ester**M-53** $C_{19}H_{30}ClNO_2$ 339.904

Isol. from the sea hare *Stylocheilus longicauda*. Oil. $[\alpha]_D$ -39 (c, 0.1 in MeOH). λ_{max} 214 (ϵ 45000); 220 (ϵ 45000); 223 (ϵ 11995); 229 (ϵ 45000); 245 (ϵ 12810) (MeOH).


5-Oxo: **Makalikone ester** $C_{19}H_{28}ClNO_3$ 353.888

Isol. from *Stylocheilus longicauda*. Oil. $[\alpha]_D$ -23 (c, 0.14 in MeOH). λ_{max} 203 (ϵ 15910); 207 (ϵ 13130); 234 (ϵ 7575); 357 (ϵ 4645) (MeOH).

Gallimore, W.A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1022-1026

Makaluvamine A**M-54**

[146555-78-4]

 $C_{11}H_{12}N_3O^{\oplus}$ 202.235Isol. from the sponges *Zyzzya fuliginosa*

and *Histodermella* sp. Also isol. from the myxomycete *Didymium bahiense*. Cytotoxic. Inhibitor of topoisomerase II. Green solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 220 (€ 17400); 330 (€ 16100) (MeOH/NaOH) (Derep). λ_{\max} 242 (€ 24000); 348 (€ 15500) (MeOH) (Derep).

N⁵-Me: Makaluvamine H. NSC 700647 [174232-34-9]
C₁₂H₁₄N₃O[⊕] 216.262
Metab. from the sponge *Zyzzya fuliginosa*. Red-brown solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 240 (€ 19400); 345 (€ 13500); 522 (€ 1100) (MeOH) (Berdy).

N⁹-(4-Hydroxyphenethyl): Makaluvamine K
[174232-36-1]
C₁₉H₂₀N₃O[⊕] 322.386
From *Zyzzya fuliginosa*. Shows cytotoxic activity. Red-brown solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 222 (€ 18000); 246 (€ 29800); 347 (€ 19200); 536 (€ 2600) (MeOH) (Berdy).

N⁹-(4-Hydroxyphenethyl), N³-Me: Makaluvamine P
C₂₀H₂₂N₃O[⊕] 336.413
Isol. from *Zyzzya fuliginosa*. Cytotoxic. Violet solid. Counterion not specified. λ_{\max} 222 (€ 7470); 248 (€ 7750); 361 (€ 5075) (MeOH).

N⁹-(4-Hydroxyphenylethyl)(E-): Makaluvamine E
[146555-82-0]
C₁₉H₁₈N₃O[⊕] 320.37
Isol. from *Zyzzya fuliginosa*. Cytotoxic. Inhibitor of topoisomerase II. Green solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 314 (€ 8600); 607 (€ 5000) (MeOH/NaOH) (Derep). λ_{\max} 226 (€ 6200); 278 (€ 8700); 333 (€ 6800); 448 (€ 4900); 626 (€ 5700) (MeOH) (Derep).

N⁹-(4-Hydroxyphenylethyl)(E-), N⁵-Me: Makaluvamine G
[152273-69-3]
C₂₀H₂₀N₃O[⊕] 334.397
Metab. from the Indonesian sponge *Histodermella* sp. and the micronesia sponge *Zyzzya fuliginosa*. Cytotoxic. Moderate inhibitor of topoisomerase I. Immunomodulator. Protein, DNA and RNA synthesis inhibitor. Green-black powder. Mp 250°. λ_{\max} 250 (€ 13850); 346 (€ 12590); 450 (€ 7940); 624 (€ 10000) (MeOH) (Berdy). λ_{\max} 286 (€ 12590); 612 (€ 12590) (MeOH/NaOH) (Berdy).

N¹-De-Me: Makaluvamine I
[138087-43-1]
C₁₀H₁₀N₃O[⊕] 188.208
Isol. from *Zyzzya fuliginosa*. Green solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 240 (€ 23900); 340 (€ 13500); 534 (€ 1000) (MeOH) (Berdy).

N¹-De-Me, N¹- β -D-ribofuranosyl: N¹- β -D-Ribofuranosylmakaluvamine I
C₁₃H₁₈N₃O[⊕] 320.324
Isol. from the sponge *Strongyloidesma alivaliensis*. Orange-brown solid. [α]_D²⁵ +5.3 (c, 0.19 in MeOH). λ_{\max} 242 (€

7815); 342 (€ 4110); 532 (€ 190) (MeOH).

N¹-De-Me, N⁵-Me: Makaluvamine C
[146555-80-8]
C₁₁H₁₂N₃O[⊕] 202.235
Isol. from *Zyzzya fuliginosa*. Cytotoxic. Inhibitor of topoisomerase II. Green solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 246 (€ 20400); 355 (€ 16900) (MeOH/NaOH) (Derep). λ_{\max} 241 (€ 25300); 358 (€ 19100) (MeOH) (Derep).

N¹-De-Me, N⁹-(4-hydroxyphenethyl): Makaluvamine D
[146555-81-9]
C₁₈H₁₈N₃O[⊕] 308.359
Isol. from *Zyzzya fuliginosa*. Cytotoxic. Inhibitor of topoisomerase II. Brown solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 243 (€ 16800); 339 (€ 12100) (MeOH/NaOH) (Derep). λ_{\max} 245 (€ 22000); 348 (€ 10600) (MeOH) (Derep).

N¹-De-Me, N⁹-(4-hydroxyphenethyl), N⁵-Me: Makaluvamine J
[174232-35-0]
C₁₉H₂₀N₃O[⊕] 322.386
Isol. from *Zyzzya fuliginosa*. Red-brown solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 220 (€ 11600); 241 (€ 18600); 354 (€ 13900); 534 (€ 1400) (MeOH) (Berdy).

N¹-De-Me, N⁹-(4-hydroxystyryl)(E-): Makaluvamine M
[174232-41-8]
C₁₈H₁₆N₃O[⊕] 306.343
Trace metab. from *Zyzzya fuliginosa*. Green solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 274 (€ 14900); 330 (€ 11000); 445 (€ 8400); 623 (€ 9200) (MeOH) (Berdy).

N¹-De-Me, N⁹-(4-hydroxystyryl)(E-), N⁵-Me: Makaluvamine L
[174232-37-2]
C₁₉H₁₈N₃O[⊕] 320.37
From *Zyzzya fuliginosa*. Green solid (as trifluoroacetate). Sol. MeOH. λ_{\max} 276 (€ 10500); 344 (€ 11700); 451 (€ 9200); 638 (€ 10400) (MeOH) (Berdy).

3,4-Didehydro: Makaluvamine B
[146555-79-5]
C₁₁H₁₀N₃O[⊕] 200.219
Isol. from *Zyzzya fuliginosa*. Cytotoxic. Inhibitor of topoisomerase II. Red solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 204 (€ 17200); 419 (€ 4800) (MeOH/NaOH) (Derep). λ_{\max} 228 (€ 9600); 442 (€ 4800) (MeOH) (Derep).


6-Bromo, N¹-de-Me: Makaluvamine N
[187964-02-9]
C₁₀H₉BrN₃O[⊕] 267.105
Isol. from *Zyzzya fuliginosa*. Cytotoxic against human colon tumour cells. Inhibitor of topoisomerase II. Reddish-brown solid. Counterion not specified. λ_{\max} 242 (€ 52966); 344 (€ 35116); 390 (€ 18806); 544 (€ 2274) (MeOH).

Radisky, D.C. *et al.*, *J.A.C.S.*, 1993, **115**, 1632 (isol, uv, ir, pmr, cmr, struct)
Carney, J.R. *et al.*, *Tetrahedron*, 1993, **49**, 8483 (Makaluvamine G)

White, J.D. *et al.*, *J.A.C.S.*, 1994, **116**, 1831 (synth, Makaluvamine D)
Izawa, T. *et al.*, *Tetrahedron*, 1994, **50**, 13593 (synth, Makaluvamines A-E)
Yamada, F. *et al.*, *Heterocycles*, 1995, **41**, 1905 (synth, Makaluvamine A)
Schmidt, E.W. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1861 (Makaluvamines H-M)
Sadanandan, E.V. *et al.*, *J.O.C.*, 1995, **60**, 1800 (synth, Makaluvamine D)
Fu, X. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1104-1106 (activity)
Peat, A.J. *et al.*, *J.A.C.S.*, 1996, **118**, 1028 (synth, Makaluvamine C)
Venables, D.A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 408 (Makaluvamine N)
Roberts, D. *et al.*, *J.O.C.*, 1997, **62**, 568 (synth, Makaluvamines A-D)
Kraus, G.A. *et al.*, *J.O.C.*, 1998, **63**, 9846-9849 (synth, Makaluvamine C)
Iwao, M. *et al.*, *Tetrahedron*, 1998, **54**, 8999-9010 (synth)
Ishibashi, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 108-110 (isol, pmr, cmr)
Casapullo, A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1354-1356 (Makaluvamine P)
Keyzers, R.A. *et al.*, *Tet. Lett.*, 2004, **45**, 9415-9418 (N¹-Ribofuranosylmakaluvamine I)
Dijoux, M.-G. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 6035-6044 (isol, activity)
Harayama, Y. *et al.*, *Curr. Org. Chem.*, 2005, **9**, 1567-1588 (rev, synth)

Makaluvamine F**M-55**

[146555-83-1]




C₁₈H₁₅BrN₃O₂S[⊕] 417.305
Metab. of the Fijian sponge *Zyzzya* cf. *marsailis*. Cytotoxic. Inhibitor of topoisomerase II. Orange solid (as trifluoroacetate). Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁵ -475.8 (c, 0.0248 in MeOH). λ_{\max} 324 (€ 17700) (MeOH/NaOH) (Derep). λ_{\max} 246 (€ 30200); 311 (€ 10800); 344 (€ 14100) (MeOH) (Derep).

Radisky, D.C. *et al.*, *J.A.C.S.*, 1993, **115**, 1632 (isol, uv, ir, pmr, cmr, struct)
Kita, Y. *et al.*, *Chem. Comm.*, 1999, 143-144 (synth)
Kita, Y. *et al.*, *Synthesis*, 1999, 885-897 (synth, ir, uv, pmr, cmr)

Makaluvic acid A**M-56**

[184023-18-5]



C₉H₁₀N₂O₃ 194.19
Alkaloid from the marine sponge *Zyzzya fuliginosa*. Cryst. (MeOH). Mp not reported. λ_{\max} 268 (€ 12684); 285 (€ 11086) (MeOH).

N⁵-De-Me: Makaluvic acid C

C₈H₈N₂O₃ 180.163

Alkaloid from the sponge *Strongylo-*
desma aliwaliensis. Brown solid. λ_{max}
200 (log ε 4.3); 260 (log ε 4.02); 289
(log ε 3.94) (MeOH).

N⁵-De-Me, N¹-β-D-ribofuranosyl: N¹-β-
D-Ribofuranosylmakaluvic acid C

C₁₃H₁₆N₂O₇ 312.279

Alkaloid from *Strongylo-*
desma ali-
waliensis. Brown solid. [α]_D²⁵ +41.2 (c,
0.0003 in MeOH). λ_{max} 210 (log ε
3.93); 260 (log ε 3.7); 286 (log ε 3.61)
(MeOH).

N¹-Me, N⁵-de-Me: **Makaluvic acid B**
[184023-37-8]

C₉H₁₀N₂O₃ 194.19

From *Zyzzya fuliginosa*. λ_{max} 268 (ε
12497); 288 (ε 11386) (MeOH).


Fu, X. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1104 (*isol*,
uv, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)

Keyzers, R.A. *et al.*, *J. Nat. Prod.*, 2005, **68**,
506-510 (*Makaluvic acid C*)

Malacitanine

M-57

5,8,13,13a-Tetrahydro-8-(hydroxy-
methyl)-2,10-dimethoxy-6H-diben-
zo[a,g]quinolizine-1,11-diol, 9CI
[132074-88-5]



Absolute
Configuration

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Ceratocarpus heterocarpa*
(Papaveraceae). Powder. Mp 116-118°. [α]_D
-87.3 (c, 0.05 in MeOH).


[131985-34-7, 131985-29-0]

Suau, R. *et al.*, *Tetrahedron*, 1990, **46**, 4421
(*isol*, *synth*, *struct*)

Malagashanine

M-58

[139682-33-0]



Absolute
Configuration

C₂₃H₃₀N₂O₄ 398.501

A 4,20-secocuran. Struct. revised in
1996. Alkaloid from *Strychnos mos-*
tueoides and *Strychnos myrto-*
ides (Loganiaceae). Shows antimalarial
potentiating activity.

12-Hydroxy: **12-Hydroxymalagashanine**
[184363-09-5]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from the stem bark of
Strychnos myrto-
ides.

19-Epimer, 12-hydroxy: **12-Hydroxy-19-**
epimalagashanine

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from the stem bark of
Strychnos myrto-
ides. Cryst. (hexane).

Mp 93-95°. [α]_D²⁰ -69.4 (c, 0.3 in
CHCl₃). λ_{max} 212 (log ε 4.39); 256 (log
ε 4); 291 (log ε 3.39) (MeOH).

3,20-Diepimer, 21-alcohol: **Malagashanol**
[243464-95-1]

C₂₂H₃₀N₂O₃ 370.491

Alkaloid from the stem bark of
Strychnos myrto-
ides. Cryst. (EtOAc/
hexane). [α]_D²⁰ +23.3 (c, 0.3 in CH₂Cl₂).
Dec. without melting. λ_{max} 251 (log ε
3.89); 278 (sh) (log ε 3.28); 285 (log ε
3.17) (MeOH).

Caira, M.R. *et al.*, *J. Chem. Crystallogr.*, 1995,
25, 725-729 (*cryst struct*)


Rasonaivo, P. *et al.*, *Gazz. Chim. Ital.*, 1996,
126, 517-519 (*struct*)

Martin, M.-T. *et al.*, *Phytochemistry*, 1999, **51**,
479-486 (*isol*, *pmr*, *cmr*)

Malarboreine

M-59

23-Norcona-4,6,18(22)-trienin-3-one.
18,20-Epiminopregna-4,6,18(11)-trien-3-
one
[15216-19-0]



C₂₁H₂₇NO 309.45

Minor alkaloid from *Malouetia arborea*
(Apocynaceae). Mp 173-176°. [α]_D +57
(EtOH). λ_{max} 283 (ε 24700) (no solvent
reported).

4,5α,6,7-Tetrahydro: **Malarborine**

[15216-18-9]

C₂₁H₃₁NO 313.482


Alkaloid from *Malouetia arborea*
(Apocynaceae). Mp 164-165.5°. [α]_D 0
(CHCl₃) (synthetic).

Sóti, F. *et al.*, *Tet. Lett.*, 1967, 1437 (*isol*,
struct, *synth*)

Malasseziacitrin

M-60

9,9a-Dihydro-3,9-di-1H-indol-3-ylfur-
o[3',2':3,4]cyclopent[1,2-b]indol-2(8H)-
one, 9CI
[863767-43-5]



Absolute
Configuration

C₂₉H₁₉N₃O₂ 441.488


Alkaloid from the yeast *Malassezia*
furfur. Mp > 200° dec. λ_{max} 222 (log ε
4.72); 269 (log ε 4.23); 285 (log ε 4.21);
357 (log ε 3.88) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
1472-1485 (*isol*, *pmr*, *cmr*, *ms*)

Malassezialactic acid

M-61

[863767-08-2]



Absolute
Configuration

C₂₀H₁₈N₂O₃ 334.374


Alkaloid from the yeast *Malassezia*
furfur. Powder. Mp 135-137°. [α]_D²⁵ -8 (c,
0.17 in MeOH). λ_{max} 209 (log ε 4.14); 225
(log ε 4.12); 276 (log ε 3.75) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
1472-1485 (*isol*, *pmr*, *cmr*, *ms*)

Malasseziazole A

M-62

5,11-Dihydro-α-oxoindolo[3,2-b]carba-
zole-6-acetic acid, 9CI
[454222-43-6]



R¹ = -COCOOH, R² = H

C₂₀H₁₂N₂O₃ 328.326

Alkaloid from the yeast *Malassezia*
furfur. Yellow solid. Mp > 250° dec. λ_{max}
211 (log ε 0.88); 225 (log ε 0.97); 255 (log
ε 1.01); 305 (log ε 0.31); 375 (log ε 0.38);
447 (log ε 0.19) (MeCN).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
1472-1485 (*isol*, *uv*, *pmr*, *cmr*, *ms*)

Malasseziazole B

M-63

[863767-09-3]

As Malasseziazole A, M-62 with
R¹ = -COCOOH, R² = -CHO

C₂₁H₁₂N₂O₄ 356.337

Alkaloid from the yeast *Malassezia*
furfur. Red solid. Mp > 300° dec. λ_{max}
209 (log ε 4.39); 230 (sh) (log ε 4.15); 242
(sh) (log ε 4.05); 275 (sh) (log ε 3.7); 306
(log ε 3.55); 409 (log ε 3.33); 482 (log ε
3.16) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
1472-1485 (*isol*, *uv*, *pmr*, *cmr*)

Malasseziazole C

M-64

[863767-15-1]

As Malasseziazole A, M-62 with
R¹ = -COOH, R² = -CHO


C₂₀H₁₂N₂O₃ 328.326

Alkaloid from the yeast *Malassezia*
furfur. Orange solid. Mp > 300° dec. λ_{max} 214
(log ε 3.76); 226 (sh) (log ε 3.71); 247 (log ε
3.66); 268 (log ε 3.51); 308 (log ε 3.23); 391
(log ε 3.19); 462 (log ε 2.95) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**,
1472-1485 (*isol*, *uv*, *pmr*, *cmr*, *ms*)

Malassezin M-65

2-[(1*H*-Indol-3-yl)methyl]-1*H*-indole-3-carboxaldehyde. 3-Formyl-2,3'-methylenebisindole. 2,3'-Methylenebisindole-3-carboxaldehyde
[352712-51-7]




C₁₈H₁₄N₂O 274.321
Alkaloid from the yeast *Malassezia furfur*. Agonist of the arylhydrocarbon receptor. Induces cytochrome P₄₅₀. Pinkish cryst. Mp 240°. λ_{max} 219 (log ε 4.73); 243 (log ε 4.23); 268 (log ε 4.2); 282 (log ε 4.15); 290 (log ε 4.17) (MeCN).

Wille, G. *et al.*, *Bioorg. Med. Chem.*, 2001, **9**, 955-960 (*isol, synth, uv, pmr, cmr, cryst struct*)

Malassezindole A M-66

1,2,3,4,5,6-Hexahydro-5-(1*H*-indol-3-yl)-4-oxoazepino[4,5-*b*]indole-2-carboxylic acid, 9CI
[863766-97-6]



C₂₁H₁₇N₃O₃ 359.384
Alkaloid from the yeast *Malassezia furfur*. Powder. Mp > 250°. [α]_D²⁵ -87 (c, 0.04 in MeOH). λ_{max} 222 (log ε 4.3); 280 (log ε 3.68) (MeOH).

5-Hydroxy: Malassezindole B

[863766-98-7]


C₂₁H₁₇N₃O₄ 375.383

Alkaloid from *Malassezia furfur*. Powder. Mp 213-215° dec. [α]_D²⁵ -15 (c, 0.12 in MeOH). λ_{max} 193 (log ε 4.23); 218 (log ε 4.1); 261 (log ε 3.56); 287 (log ε 3.48) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 1472-1485 (*isol, cd, pmr, cmr*)

Malasseziona M-67

1,3-Di-1*H*-indol-3-yl-2-propanone, 9CI. Malasseziaketone
[863767-46-8]




C₁₉H₁₆N₂O 288.348

Alkaloid from the yeast *Malassezia furfur*. Amorph. solid. Mp > 150° dec. λ_{max} 207 (log ε 3.04); 216 (sh) (log ε 3.01); 266 (log ε 2.57); 280 (log ε 2.56); 289 (sh) (log ε 2.54) (MeOH).

Irlinger, B. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 1472-1485 (*isol, pmr, cmr, ms*)

Malatyamine M-68

1,4-Dihydro-4-oxo-2-quinolinehexanoic acid, 9CI
[98752-01-3]




C₁₅H₁₇NO₃ 259.304

Alkaloid from *Haplophyllum cappadocicum* (whole plant) (Rutaceae). Isol. and elucidated as the Et ester.

Arar, G. *et al.*, *J. Nat. Prod.*, 1985, **48**, 642 (*isol, uv, ir, pmr, ms, struct*)

Malbrancheamide A M-69

[881216-55-3]



C₂₁H₂₃Cl₂N₃O 404.338

Prod. by *Malbranchea aurantiaca*. Calmodulin inhibitor. Phytotoxin. Cryst. (MeOH/CH₂Cl₂). Mp 321-324°. [α]_D +42 (c, 1 in MeOH). λ_{max} 233 (log ε 3.94); 293 (log ε 4.74) (MeOH).

12a-Epimer: Epimalbrancheamide
[934266-33-8]

C₂₁H₂₃Cl₂N₃O 404.338

Oil.


Valente, M.W.N. *et al.*, *Heterocycles*, 2006, **70**, 249-259 (*epimer, synth, pmr, cmr*)

Martinez-Luis, S. *et al.*, *Tetrahedron*, 2006, **62**, 1817-1822 (*isol, cd, pmr, cmr, ms, cryst struct*)

Miller, K.A. *et al.*, *J.O.C.*, 2008, **73**, 3116-3119 (*synth*)

Maleimycin M-70

Antibiotic IV-3. 5,6-Dihydro-4-hydroxycyclopenta[*c*]pyrrole-1,3(2*H*,4*H*)-dione, 9CI. 3-Aza-6-hydroxybicyclo[3.3.0]oct-1(5)-ene-2,4-dione



C₇H₇NO₃ 153.137

λ_{max} 225 (ε 14000); 230 (sh) (ε 13000) (H₂O) (Derep).

▶ LD₅₀ (mus, ipr) 6.25mg/kg.

(+)-*form* [50988-16-4]

Isol. from *Streptomyces showdoensis* and *Streptomyces filamentous*. Active against gram-positive and -negative organisms, acid-fast bacteria, leukaemia L-1210 cells and tumours. Sol. H₂O; fairly sol. EtOAc. Mp 116°. λ_{max} 225 (ε 14000) (H₂O) (Berdy).

(±)-*form* [62356-52-9]

Cryst. (C₆H₆/hexane). Mp 110-112°.

Elstner, E.F. *et al.*, *Biochemistry*, 1973, **12**,

4992 (*isol, uv, ir, ms, nmr, struct, biosynth*)

Singh, P. *et al.*, *Tetrahedron*, 1976, **32**, 2379

(*synth, ir, nmr*)

Ghiringhelli, D. *et al.*, *Biomed. Mass*

Spectrom., 1981, **8**, 155 (*ms*)

Zhu, B. *et al.*, *Kangshengsu*, 1986, **11**, 167; *CA*,


105, 94208k

Lee, C.J. *et al.*, *Synth. Commun.*, 1992, **22**, 803

(*synth*)

Malekulatine M-71

[87183-76-4]



C₃₉H₄₆N₂O₈ 670.801

Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). Shows selective antiparasitoid and anti-platelet aggregation activity. [α]_D²⁵ +156 (c, 0.14 in MeOH).

Bruneton, J. *et al.*, *J.O.C.*, 1983, **48**, 3957 (*isol, uv, cd, pmr, ms, struct*)

Chen, I.S. *et al.*, *Planta Med.*, 1995, **61**, 537-539 (*activity*)

Angerhoffer, C.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 59-66 (*activity*)

Malic acid, 8CI M-72

Hydroxybutanedioic acid, 9CI. Hydroxysuccinic acid. E296
[6915-15-7]



C₄H₆O₅ 134.088

Flavour enhancer and other uses in food. Acidulant, antioxidant. Flavouring agent, buffering agent and a chelating agent.

▶ Skin and severe eye irritant. ON7175000

(*S*)-*form*

L-*form*. FEMA 2655

[97-67-6]

Occurs naturally in apples and various other fruits. Cryst. (Me₂CO). V. sol. EtOH, spar. sol. Et₂O. Mp 104-105°

(100°). $[\alpha]_D^{21}$ -2.3 (c, 8.5 in H₂O). $[\alpha]_D^{20}$ -28.7 (c, 5.5 in Py) (>99.8% ee). pK_{a1} 3.46; pK_{a2} 5.1. Dec. at ca. 140°. Specific rotn. concn. dependent. See (±)-form for food use.

Dibenzylamide: N,N'-Dibenzyl-2-hydroxybutanediamide. N,N'-Dibenzylmalamide

C₁₈H₂₀N₂O₃ 312.368

Constit. of the stems of *Salvadora persica*. Mp 153-155°. $[\alpha]_D^{25}$ -24.2 (c, 0.05 in MeOH).


[3105-51-9, 68303-40-2, 149-61-1, 676-46-0, 39015-77-5, 4387-09-1, 22138-22-3]

Khalil, A.T. et al., *Arch. Pharmacol. Res.*, 2006, **29**, 952-956 (N,N'-Dibenzylmalamide, isol)

Malindine

M-73

[81161-28-6]



C₂₀H₂₂N₃[⊕] 304.414

Alkaloid from the stem bark of *Strychnos decussata* and the root bark of *Strychnos usambarensis* (Loganiaceae). Muscular relaxant. Mp 232-234° dec.

N-De-Me: Normalindine

[113122-64-8]

C₁₉H₁₉N₃ 289.379

Alkaloid from the root bark of *Strychnos johnsonii* (Loganiaceae) and leaves of *Ophiorrhiza filistipula* (Rubiaceae). Yellow plates (EtOAc/petrol). Mp 131-136°. $[\alpha]_D$ -210 (c, 0.1 in CHCl₃) (-103).

19-Epimer: Isomalindine

[91739-07-0]

C₂₀H₂₂N₃[⊕] 304.414

Alkaloid from root bark of *Strychnos usambarensis* (Loganiaceae).

19-Epimer, N-de-Me: Norepimalindine.

Norisomalindine

[113215-44-4]

C₁₉H₁₉N₃ 289.379

Alkaloid from the root of *Strychnos johnsonii* (Loganiaceae). $[\alpha]_D$ -276 (c, 0.1 in CHCl₃).

Olaniyi, A.A. et al., *Planta Med.*, 1981, **43**, 353-359 (isol, uv, ir, pmr, cmr, ms, cd, struct)
Caprasse, M. et al., *Planta Med.*, 1984, **50**, 27-30 (isol, uv, ir, pmr, ms, cd)

Massiot, G. et al., *Phytochemistry*, 1987, **26**, 2839-2846 (Normalindine, Norepimalindine)

Maiti, B.C. et al., *Heterocycles*, 1990, **31**, 847-850 (Normalindine, Norepimalindine, synth)

Rey, A.W. et al., *Heterocycles*, 1991, **32**, 1143-1151 (Normalindine, synth)

Arbain, D. et al., *Aust. J. Chem.*, 1993, **46**, 977-985 (isol, abs config, Normalindine)


Ohba, M. et al., *Tetrahedron*, 2000, **56**, 7751-7761 (Normalindine, synth)

Davis, F.A. et al., *J.O.C.*, 2006, **71**, 8761-8766 (Normalindine, synth)

Malonganenone A

M-74

[882403-69-2]



C₂₆H₃₈N₄O₂ 438.612

Isol. from *Leptogorgia gilchristi*. Cytotoxic. Glass. λ_{max} 221 (ε 30900); 252 (ε 28800) (no solvent reported).

A¹⁰E-Isomer: Malonganenone D

[945458-47-9]

C₂₆H₃₈N₄O₂ 438.612

Isol. from *Euplexaura nuttingi*. Oil.

11'E-Isomer: Malonganenone E

[945458-48-0]

C₂₆H₃₈N₄O₂ 438.612

Isol. from *Euplexaura nuttingi*. Oil.


Keyzers, R.A. et al., *Tetrahedron*, 2006, **62**, 2200-2206 (Malonganenone A)

Sorek, H. et al., *J. Nat. Prod.*, 2007, **70**, 1104-1109 (Malonganenones D,E)

Malonganenone B

M-75

[882403-70-5]



C₂₇H₄₂N₄O₃ 470.654

Isol. from *Leptogorgia gilchristi*. Cytotoxic. Glass. λ_{max} 238 (ε 14400) (MeOH).

A¹⁰E-Isomer: Malonganenone F

[945458-49-1]

C₂₇H₄₂N₄O₃ 470.654

Isol. from *Euplexaura nuttingi*. Oil.

11'E-Isomer: Malonganenone G

[945458-50-4]

C₂₇H₄₂N₄O₃ 470.654

Isol. from *Euplexaura nuttingi*. Oil.


Keyzers, R.A. et al., *Tetrahedron*, 2006, **62**, 2200-2206 (Malonganenone B)

Sorek, H. et al., *J. Nat. Prod.*, 2007, **70**, 1104-1109 (Malonganenones F,G)

Malonomicin

M-76

Antibiotic K 16A. K 16A
[38249-71-7]



C₁₃H₁₈N₄O₉ 374.307

Peptide antibiotic. Several tautomers possible. Shown in zwitterionic form in the lit. Produced by *Streptomyces rimo-*

sus. Active against gram-positive organisms. Microcryst. powder. Sol. acids, bases; fairly sol. H₂O, THF, dioxan, Py, DMSO, DMF; poorly sol. MeOH, hexane. $[\alpha]_D^{20}$ -64 (c, 1 in H₂O). Mp indefinite. λ_{max} 240 (ε 6600); 277 (ε 7600) (H₂O) (Berdy). λ_{max} 274 (ε 6700) (HCl) (Berdy).

▶ LD₅₀ (mus, ipr) 1000 mg/kg. TY1760000 [38249-70-6]

Batelaan, J.G. et al., *Tet. Lett.*, 1972, 3103; 3107 (struct)


van der Baan, J.L. et al., *Tetrahedron*, 1978, **34**, 223 (synth)

Schipper, D. et al., *J.C.S. Perkin 1*, 1979, 2017
Schipper, D. et al., *Tet. Lett.*, 1982, **23**, 1289; 1293 (biosynth, pmr)

Maltoxazine

M-77

1,2,3,3a,6,7-Hexahydrocyclopenta[d]pyrrolo[2,1-b][1,3]oxazin-8(5H)-one, 9CI.
Daechalkaloid A
[80933-73-9]
[111261-77-9]



C₁₀H₁₃NO₂ 179.218

Aroma substance isol. from malt. Alkaloid from fruit of *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Amorph. Mp 52°. $[\alpha]_D^{22}$ +0.3 (c, 0.82 in CHCl₃). Rapidly becomes resinous in soln. The small opt. rotn. and the Mp. were reported for Daechalkaloid A.


Tressl, R. et al., *Helv. Chim. Acta*, 1982, **65**, 483 (Maltoxazine)

Han, B.H. et al., *Tet. Lett.*, 1987, **28**, 3957 (Daechalkaloid A)

Malynamide W

M-78

[485388-83-8]



C₂₃H₃₉NO₅ 409.565


Isol. from *Lyngbya majuscula*. Pale yellow oil. $[\alpha]_D^{18}$ -15 (c, 0.06 in CHCl₃). λ_{max} 203 (log ε 4.08); 227 (log ε 3.88) (MeOH).

McPhail, K.L. et al., *J. Nat. Prod.*, 2003, **66**, 132-135 (isol, pmr, cmr)

Mamanine

M-79

6-[Octahydro-1-(hydroxymethyl)-2H-quinolizin-3-yl]-2-(1H)-pyridinone, 9CI
[60394-92-5]



C₁₅H₂₂N₂O₂ 262.351

Numbering schemes differ. Alkaloid from the bark of *Sophora chrysophylla* and from flowers of *Sophora flavescens* (Fabaceae). Mp 100° (remelts 171-172°). $[\alpha]_D^{25} +31.7$ (c, 2.32 in EtOH).

N-Oxide: Mamanine N-oxideC₁₅H₂₂N₂O₃ 278.35

Alkaloid from leaves of *Sophora chrysophylla* (Fabaceae). Amorph. $[\alpha]_D^{25} -9$ (c, 0.003 in EtOH).

8 β -Hydroxy: 13 β -HydroxymamanineC₁₅H₂₂N₂O₃ 278.35

Alkaloid from the stems of *Maackia amurensis* var. *buergeri* (Fabaceae). Amorph. solid. $[\alpha]_D^{25} +31.2$ (c, 0.107 in MeOH).

NH-form**O,N-Di-Ac:**

Yellow oil. Dec. on standing.

OH-form**O⁶-Me: Jussiaeiine B**C₁₆H₂₄N₂O₂ 276.378

Alkaloid from *Ulex jussiaei*. Oil. $[\alpha]_D^{25} +27.2$ (c, 0.27 in CHCl₃). λ_{\max} 217 (log ϵ 3.26); 268 (log ϵ 3.38); 273 (sh) (MeOH).

7 β -Hydroxy, O⁶-Me: Jussiaeiine DC₁₆H₂₄N₂O₃ 292.377

Alkaloid from *Ulex jussiaei*. Oil. $[\alpha]_D^{25} +14.5$ (c, 0.64 in CHCl₃). λ_{\max} 218 (log ϵ 3.2); 269 (log ϵ 3.29) (MeOH).

8 β -Hydroxy, O⁶-Me: Jussiaeiine CC₁₆H₂₄N₂O₃ 292.377

Alkaloid from *Ulex jussiaei*. Oil. $[\alpha]_D^{25} +32.2$ (c, 0.19 in CHCl₃). λ_{\max} 218 (log ϵ 3.29); 269 (log ϵ 3.43); 273 (sh) (MeOH).

Kadooka, M.M. *et al.*, *Tetrahedron*, 1976, **32**, 919 (*Mamanine*, *isol*, *ir*, *uv*, *pmr*, *cmr*, *cryst struct*)


Murakoshi, I. *et al.*, *Phytochemistry*, 1981, **20**, 1407; 1982, **21**, 2379; 1984, **23**, 887 (*Mamanine*, *Mamanine N-oxide*)

Saito, K. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 3982 (*13-Hydroxymamanine*)

Maximo, P. *et al.*, *J. Nat. Prod.*, 2000, **63**, 201-204 (*Jussiaeiines*)

Manadomanzamine A

M-80

C₃₉H₅₄N₄O₂ 610.881

Isol. from the Indonesian sponge *Acanthostrongylophora* sp. Active against HIV-1 and mycobacteria. Powder. $[\alpha]_D -19$ (c, 0.11 in MeOH). λ_{\max} 282 (ϵ 7700) (no solvent reported).

22-Epimer: Manadomanzamine BC₃₉H₅₄N₄O₂ 610.881

Isol. from *Acanthostrongylophora* sp.


Powder. $[\alpha]_D -18$ (c, 0.11 in MeOH).

λ_{\max} 282 (ϵ 7200) (no solvent reported).

Peng, J. *et al.*, *J.A.C.S.*, 2003, **125**, 13382-13386 (*isol*, *cd*, *pmr*, *cmr*)

Manibacanine

M-81

C₁₉H₂₁NO₃ 311.38**(R)-form [151757-07-2]**

Alkaloid from stem bark of *Aniba canelilla* (Lauraceae). $[\alpha]_D +176$ (c, 0.34 in MeOH).

(S)-form**O²-De-Me: Anibacanine**

[151757-06-1]

C₁₈H₁₉NO₃ 297.353


Alkaloid from stem bark of *Aniba canelilla* (Lauraceae). $[\alpha]_D -129$ (c, 0.3 in CHCl₃).

Oger, J.-M. *et al.*, *Can. J. Chem.*, 1993, **71**, 1128 (*isol*, *uv*, *pmr*, *cmr*, *ms*, *struct*)

Manicoline A

M-82

[79820-32-9]

C₁₅H₂₁NO₂ 247.336


Alkaloid from the root bark of *Dulacia guianensis* (Olacaceae). Pale-yellow cryst. (EtOAc). Mp 197-199°. $[\alpha]_D^{22} +69$ (c, 0.9 in CHCl₃ + 10% MeOH). λ_{\max} 252 (ϵ 32100); 266 (sh) (ϵ 21400); 277 (sh) (ϵ 10700); 340 (ϵ 13200); 420 (ϵ 10700) (EtOH) (Derep).

Polonsky, J. *et al.*, *Chem. Comm.*, 1981, 731 (*isol*, *uv*, *pmr*, *ms*, *cryst struct*)

Manicoline B

M-83

[79921-08-7]


C₁₅H₂₁N₃O₂ 275.35

Alkaloid from the root bark of *Dulacia guianensis* (Olacaceae). Cryst. (Me₂CO). Mp 199-200°. $[\alpha]_D^{22} +36$ (c, 0.23 in CHCl₃). Equimolar mixt. of two diastereoisomers, epimeric at C-4. λ_{\max} 228 (ϵ 2110); 272 (ϵ 1710) (EtOH) (Derep).

Polonsky, J. *et al.*, *Tet. Lett.*, 1984, **25**, 2359 (*uv*, *pmr*, *ms*, *cryst struct*)

Manilamine

M-84

C₁₈H₂₂N₂O 282.385


A secouline alkaloid (cf. Uleine, U-8). Alkaloid from the leaves of *Alstonia scholaris*. Beige solid. Mp 88-92°. $[\alpha]_D +15.5$ (c, 0.5 in MeOH). λ_{\max} 223 (log ϵ 2.69); 282 (log ϵ 2.38); 297 (log ϵ 2.49) (MeOH).

Macabeo, A.P.G. *et al.*, *Phytochemistry*, 2005, **66**, 1158-1162 (*isol*, *pmr*, *cmr*, *ms*)

Maniwamycin B

M-85

[3-(1-Hexenyl)-O,N,N-azoxy]-2-butanol, 9CI. Antibiotic KC 7367A. KC 7367A [122547-71-1]

C₁₀H₂₀N₂O₂ 200.28

Related to Antibiotic LL-BH872 α , A-1199. Prod. by *Streptomyces prasinopilosus*. Antifungal agent. Oil. $[\alpha]_D^{22} +108$ (c, 1 in CHCl₃). λ_{\max} 230 (ϵ 45000) (MeOH).

► LD₅₀ (mus, ivn) 100 mg/kg. EL4140000

2-Ketone: **Maniwamycin A**. Antibiotic KC 7367B. KC 7367B [122566-70-5]

C₁₀H₁₈N₂O₂ 198.264

From *Streptomyces prasinopilosus*. Antifungal agent. Oil. $[\alpha]_D^{22} -144$ (c, 1 in CHCl₃).

► LD₅₀ (mus, ivn) 10 mg/kg. EL8777000


Nakayama, M. *et al.*, *J. Antibiot.*, 1989, **42**, 1535-1540; 1541-1546 (*isol*, *struct*)

Takahashi, Y. *et al.*, *Tet. Lett.*, 1991, **32**, 1067-1068 (*synth*)

Nakata, M. *et al.*, *Tet. Lett.*, 1993, **34**, 6095-6098 (*synth*)

Mannopeptimycin β

M-86

C₄₂H₅₈N₁₂O₁₅ 970.992

Glycopeptide antibiotic. Prod. by *Streptomyces hygroscopicus* LL-AC98. Antibacterial agent. Amorph. powder. $[\alpha]_D -29.6$ (c, 0.6 in MeOH aq.).

4'-O-[α -L-Mannopyranosyl-(1 \rightarrow 4)- α -L-mannopyranoside]: **Mannopeptimycin α**
 $C_{54}H_{78}N_{12}O_{25}$ 1295.276
 Prod. by *Streptomyces hygroscopicus* LL-AC98. Amorph. powder. $[\alpha]_D$ -3.4 (c, 0.38 in MeOH aq.).

4'-O-[3-Methylbutanoyl-(\rightarrow 2)- α -L-mannopyranosyl-(1 \rightarrow 4)- α -L-mannopyranoside]: **Mannopeptimycin γ**
 $C_{59}H_{86}N_{12}O_{26}$ 1379.393
 Prod. by *Streptomyces hygroscopicus* LL-AC98. Amorph. powder. $[\alpha]_D$ -7.1 (c, 0.35 in MeOH aq.).


4'-O-[3-Methylbutanoyl-(\rightarrow 3)- α -L-mannopyranosyl-(1 \rightarrow 4)- α -L-mannopyranoside]: **Mannopeptimycin δ**
 $C_{59}H_{86}N_{12}O_{26}$ 1379.393
 Prod. by *Streptomyces hygroscopicus* LL-AC98. Amorph. powder. $[\alpha]_D$ -12.2 (c, 0.37 in MeOH aq.).

4'-O-[3-Methylbutanoyl-(\rightarrow 4)- α -L-mannopyranosyl-(1 \rightarrow 4)- α -L-mannopyranoside]: **Mannopeptimycin ϵ**
 $C_{59}H_{86}N_{12}O_{26}$ 1379.393
 Prod. by *Streptomyces hygroscopicus* LL-AC98. Amorph. powder. $[\alpha]_D$ -13.8 (c, 0.1 in MeOH aq.).

He, H. *et al.*, *J.A.C.S.*, 2002, **124**, 9729-9736 (isol, pmr, cmr)

Manumycin A M-87

Antibiotic TMC 1F. TMC 1F. UCF1C.
 Antibiotic UCF1C
 [52665-74-4]



$C_{31}H_{38}N_2O_7$ 550.65

Abs. config. revised in 1999. Isol. from *Streptomyces parvulus* Tü 64 and a marine *Streptomyces* sp. Active against gram-positive bacteria. Inhibits protein-farnesyl transferase. Shows antiprotozoal activity. Cryst. (MeOH aq.). Sol. MeOH, MeCN, C_6H_6 , bases; poorly sol. H_2O , hexane. Mp 139-141° dec. $[\alpha]_D^{20}$ -185 (c, 0.4 in $CHCl_3$). Similar to *Asukamycin*. λ_{max} 270 (ϵ 31700); 328 (ϵ 32300) (MeOH/HCl) (Derep). λ_{max} 261 (ϵ 40200) (MeOH/NaOH) (Derep). λ_{max} 278 (ϵ 36400); 314 (ϵ 34600) (MeOH) (Derep). λ_{max} 261 (ϵ 40200) (MeOH-NaOH) (Berdy).

Zeeck, A. *et al.*, *J. Antibiot.*, 1987, **40**, 1530; 1541; 1549 (isol, abs config, props, bibl)

Thiericke, R. *et al.*, *J.A.C.S.*, 1990, **112**, 3979 (biosynth)

Hara, M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 2281-2285 (UCF1C)

Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1212 (TMC 1F)

Alcaraz, L. *et al.*, *J.O.C.*, 1998, **63**, 3526-3527 (synth, config)

Sattler, I. *et al.*, *Nat. Prod. Rep.*, 1998, **15**, 221-240 (rev)

Alcaraz, L. *et al.*, *Tetrahedron*, 1999, **55**, 3707-3716 (synth, abs config)

Pat. Coop. Treaty (WIPO), 2001, 01 05 384; CA, **134**, 110443q (activity)

Li, F. *et al.*, *J. Nat. Prod.*, 2005, **68**, 349-353 (isol, cmr)


Manumycin C M-88

UCF1B. Antibiotic UCF1B. TMC 1G.

Antibiotic TMC 1G

[139023-59-9]

As Manumycin A, M-87 with



$C_{30}H_{36}N_2O_7$ 536.624

Prod. by *Streptomyces parvulus*. Mp 188-196° (dec.). $[\alpha]_D^{20}$ +32.4 (c, 0.4 in $CHCl_3$). λ_{max} 278 (ϵ 42700); 307 (ϵ 33080) (MeOH).

Sattler, I. *et al.*, *J.O.C.*, 1993, **58**, 6583 (isol, uv, ir, pmr, cmr, ms)

Hara, M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 2281-2285 (UCF1B)


Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1212-1220 (TMC 1G)

Sattler, I. *et al.*, *Nat. Prod. Rep.*, 1998, **15**, 221-240 (rev)

Manumycin D M-89

TMC 1E. Antibiotic TMC 1E

[149764-30-7]



$C_{31}H_{40}N_2O_7$ 552.666

Prod. by *Streptomyces parvulus*. Protein farnesylation inhibitor. Sol. MeOH, Me_2CO , MeCN, $CHCl_3$; poorly sol. H_2O , hexane. Mp 96°. $[\alpha]_D^{20}$ -25.3 (c, 0.3 in $CHCl_3/MeOH$). λ_{max} 205 (ϵ 52690); 245 (ϵ 38900); 309 (ϵ 43690) (MeOH). λ_{max} 205 (ϵ 52690); 245 (ϵ 38900); 309 (ϵ 43690) (MeOH) (Berdy). λ_{max} 207 (ϵ 32080); 325 (ϵ 24150) (MeOH-HCl) (Berdy). λ_{max} 211 (ϵ 24870); 261 (ϵ 25580); 302 (ϵ 24570) (MeOH-NaOH) (Berdy).


Sattler, I. *et al.*, *J.O.C.*, 1993, **58**, 6583 (isol, uv, ir, pmr, cmr, ms)

Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1212-1220 (TMC 1E)

Alcaraz, L. *et al.*, *Tetrahedron*, 1999, **55**, 3707-3716 (abs config)

Manzacidin B M-90

[134029-42-8]



$C_{12}H_{14}BrN_3O_5$ 360.164

Alkaloid from the marine sponge *Hymeniacidon* sp. Oil. $[\alpha]_D^{22}$ -71 (c, 0.43 in MeOH). λ_{max} 209 (ϵ 5100); 272 (ϵ 5800) (MeOH) (Derep).

5-Deoxy: **Manzacidin C**

[134107-38-3]

$C_{12}H_{14}BrN_3O_4$ 344.164

From the sponge *Hymeniacidon* sp.

Oil. $[\alpha]_D^{22}$ +37 (c, 0.23 in MeOH). λ_{max} 209 (ϵ 5100); 272 (ϵ 5800) (MeOH) (Derep).

5-Deoxy, N^1 -Me: **N^1 -Methylmanzacidin C**

[389120-67-6]

$C_{13}H_{16}BrN_3O_4$ 358.191

Isol. from the sponge *Axinella brevis-*

tyla. Antifungal and cytotoxic agent.

$[\alpha]_D^{23}$ +36.4 (c, 0.17 in MeOH). λ_{max} 202 (log ϵ 3.7); 220 (log ϵ 3.7); 273 (log ϵ 3.7) (MeOH).

6-Epimer, 5-deoxy: **Manzacidin A**

[134029-41-7]

$C_{12}H_{14}BrN_3O_4$ 344.164

From the sponge *Hymeniacidon* sp.

Oil. $[\alpha]_D^{27}$ -28 (c, 0.67 in MeOH). λ_{max} 209 (ϵ 5100); 272 (ϵ 5800) (MeOH) (Derep).

6-Epimer, 5-deoxy, debromo, N^1 -Me:

Manzacidin D

[192197-62-9]

$C_{13}H_{17}N_3O_4$ 279.295

Alkaloid from the sponge *Astrosclera willejana*. Viscous oil. $[\alpha]_D^{25}$ +14.5 (c, 0.3 in $CHCl_3$). λ_{max} 225 (ϵ 5980); 270 (ϵ 7980) (no solvent reported).

Kobayashi, J. *et al.*, *J.O.C.*, 1991, **56**, 4574-4576 (isol, uv, ir, pmr, cmr, ms)

Jahn, T. *et al.*, *Tet. Lett.*, 1997, **38**, 3883-3884 (Manzacidin D)

Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1576-1578 (N^1 -Methylmanzacidin C)

Wehn, P.M. *et al.*, *J.A.C.S.*, 2002, **124**, 12950-12951 (Manzacidin A, C, synth)

Drouin, C. *et al.*, *Tet. Lett.*, 2004, **45**, 7197-7199 (Manzacidin D, synth)

Lanter, J.C. *et al.*, *Org. Lett.*, 2005, **7**, 5905-5907 (Manzacidin C, synth)

Kano, T. *et al.*, *J.A.C.S.*, 2006, **128**, 2174-2175 (Manzacidin A, synth)

Shinada, T. *et al.*, *Org. Lett.*, 2007, **9**, 1765-1767 (abs config)


Hashimoto, T. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 829-835 (rev, synth)

Tran, K. *et al.*, *Org. Lett.*, 2008, **10**, 3165-3167 (Manzacidin C, synth)

Oe, K. *et al.*, *Tet. Lett.*, 2008, **49**, 7426-7429 (Manzacidin A, C, synth)

Manzamine A M-91

Keramamine A



C₃₆H₄₄N₄O 548.77

(+)-**form** [104196-68-1]

Alkaloid from the Japanese marine sponges *Haliclona* sp., *Xestospongia* sp. and *Pellina* sp. Isol. from *Acanthostromgylophora* aff. *ingens* as free base. Shows antitumour props. Exhibits weak cytotoxicity against KB cells, also insecticidal activity and activity against gram-positive and gram-negative bacteria. Exhibits strong antimalarial props. and activity against *Leishmania donovani*; also strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Cryst. (MeOH) (as hydrochloride). Mp 240° dec. (hydrochloride). [α]_D²⁰ +50 (c, 0.28 in CHCl₃) (hydrochloride). Usually isol. as hydrochloride salt. λ_{max} 238 (ε 15300); 278 (ε 9680); 290 (ε 8350); 346 (ε 4420); 358 (ε 4770) (hydrochloride). λ_{max} 219 (ε 22900); 236 (ε 18600); 280 (ε 10800); 290 (ε 11000); 346 (ε 5300); 357 (ε 5600) (MeOH).

2-N-Oxide: **Manzamine A N-oxide**

[184361-76-0]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from the Phillipine marine sponge *Xestospongia ashmorica*. Cytotoxic agent. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Yellow cryst. powder. [α]_D +18.6 (c, 0.35 in CHCl₃). λ_{max} 201 (ε 26000); 241 (ε 23000); 261 (ε 23000); 310 (ε 21000) (MeOH).

3,4-Dihydro: **3,4-Dihyromanzamine A**

[162465-80-7]

C₃₆H₄₆N₄O 550.786

Alkaloid from the Okinawan marine sponge *Amphimedon* sp. Cytotoxic against L-1210 and KB cells. Exhibits antibacterial activity. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Amorph. solid. Mp 237-241°. [α]_D²⁰ +86 (c, 0.25 in CHCl₃). λ_{max} 244 (ε 21000); 323 (ε 10000) (MeOH) (Berdy).

3,4-Dihydro, 2-N-oxide: **3,4-Dihyromanzamine A N-oxide**

[184361-75-9]

C₃₆H₄₆N₄O₂ 566.785

Alkaloid from *Xestospongia ashmorica*. Cytotoxic agent. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Yellow cryst. powder. [α]_D +34.1 (c, 0.59 in CHCl₃). λ_{max} 201 (ε 25000); 355 (ε 11000) (MeOH).

IR,2,3,4-Tetrahydro: **Manzamine D**

[116477-23-7]

C₃₆H₄₈N₄O 552.801

Alkaloid from the sponges *Haliclona*, *Ircinia* and *Amphimedon* spp. Cytotoxic agent. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁴ +44. λ_{max} 223 (ε 28800); 281 (ε 6700); 288 (ε 5400) (MeOH) (Berdy).

IS,2,3,4-Tetrahydro: **1-Epimanzamine D**

[302327-68-0]

C₃₆H₄₈N₄O 552.801

Alkaloid from a Palaun sponge. Cytotoxic agent. Shows strong activity against *Mycobacterium tuberculosis*

(H₃₇Rv). Amorph. powder. [α]_D²³ +77.3 (c, 0.16 in CHCl₃). λ_{max} 225 (log ε 4.23); 281 (log ε 3.79) (MeOH).

IS,2,3,4-Tetrahydro, N²-Me: **1-Epi-2-N-methylmanzamine D**

[302327-67-9]

C₃₇H₅₀N₄O 566.828

Alkaloid from a Palaun sponge. Cytotoxic agent. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Cryst. (MeOH). Mp 185-188°. [α]_D²³ +91.4 (c, 0.27 in CHCl₃). λ_{max} 223 (log ε 4.24); 281 (log ε 3.74) (MeOH).

31,34-Epoxy, 32,33-dihydro: **6-Deoxymanzamine X**

[184361-73-7]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from *Xestospongia ashmorica*. Cytotoxic agent. Active against *Leishmania donovani*. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Pale yellow amorph. powder. [α]_D +30.1 (c, 0.35 in CHCl₃). λ_{max} 210 (ε 26000); 260 (ε 11800); 312 (sh) (ε 10000); 378 (ε 3000) (MeOH).

31-Oxo, 32,33-dihydro: **Manzamine E**

[117631-50-2]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from a *Xestospongia* sponge. Shows activity against *Mycobacterium tuberculosis* and *Leishmania donovani*. Cryst. (MeCN). Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O, hexane. Mp 174-176°. [α]_D +63.7 (c, 2.51 in CHCl₃). λ_{max} 238 (ε 15300); 278 (ε 9680); 290 (ε 8350); 346 (ε 4420); 358 (ε 4770) (as hydrochloride) (Derep). λ_{max} 219 (ε 22900); 236 (ε 18600); 280 (ε 10800); 290 (ε 11000); 346 (ε 5300); 357 (ε 5600) (MeOH) (Derep). λ_{max} 220 (ε 35800); 237 (ε 28100); 279 (ε 18000); 346 (ε 7700); 359 (ε 9300) (MeOH) (Berdy).

6-Hydroxy: **6-Hydroxymanzamine A**

Manzamine Y

[162465-79-4]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from the Okinawan marine sponge *Amphimedon* sp. and *Haliclona* sp. Cytotoxic against L-1210 and KB cells. Exhibits antibacterial activity. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Yellowish amorph. solid. Mp 253°. [α]_D²⁰ +139 (c, 1.1 in MeOH). [α]_D¹⁹ +33.2 (c, 2.50 in CHCl₃).

6-Hydroxy, 3,4-dihydro: **3,4-Dihydro-6-hydroxymanzamine A**

[208392-25-0]

C₃₆H₄₆N₄O₂ 566.785

Alkaloid from *Amphimedon* sp. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Amorph. solid. [α]_D²⁵ +28 (c, 1.2 in MeOH). λ_{max} 207 (ε 9000); 225 (ε 6500); 250 (ε 3500); 337 (ε 2500) (MeOH).

6-Hydroxy, 31β,34β-epoxy, 32,33-dihydro: **Manzamine X**

[164301-23-9]

C₃₆H₄₄N₄O₃ 580.769

Alkaloid from sponge *Xestospongia* sp. Cytotoxic against L-1210 and KB cells.

Exhibits antibacterial activity. Yellow prisms (hexane/Me₂CO). Mp 250°. [α]_D¹⁹ +66.1 (c, 1.93 in CHCl₃). λ_{max} 215 (ε 29500); 300 (ε 17000); 378 (ε 4800) (MeOH) (Berdy).

6-Hydroxy, 31-oxo, 32,33-dihydro: **6-Hydroxymanzamine E**

C₃₆H₄₄N₄O₃ 580.769

Alkaloid from the Indonesian sponge *Acanthostromgylophora* sp. Yellow powder (CHCl₃). Fp ca. 198. [α]_D²⁵ +34.4 (c, 0.2 in CHCl₃). λ_{max} 218 (log ε 3.64); 239 (log ε 3.63); 280 (log ε 3.25); 288 (log ε 3.09); 346 (log ε 3.42) (MeOH).

6-Hydroxy, 35-oxo, 32,33-dihydro: **32,33-Dihydro-6-hydroxy-35-oxomanzamine A**

[581782-67-4]

C₃₆H₄₄N₄O₃ 580.769

Alkaloid from an Indonesian sponge. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Pale yellow powder. Mp > 200° dec. [α]_D²⁵ +10 (c, 1 in MeOH). λ_{max} 219; 248; 268; 356; 395 (MeOH).

8-Hydroxy: **Manzamine G**. 8-Hydroxymanzamine A

[154466-37-2]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from *Acanthostromgylophora* aff. *ingens*, *Amphimedon* sp. and *Pachypellina* sp. Exhibits antitumour, antimalarial and anti-HSV-II activities. Active against *Leishmania donovani*. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Pale yellow cryst. Mp 230° dec. [α]_D +118.5 (c, 1.94 in CHCl₃). Usually obt. as a salt. λ_{max} 222 (ε 32300); 245 (ε 30600); 268 (ε 14000); 360 (ε 8200) (EtOH) (Berdy).

8-Hydroxy, 1R,2,3,4-tetrahydro: **1,2,3,4-Tetrahydro-8-hydroxymanzamine A**. 8-Hydroxymanzamine D

[160070-84-8]

C₃₆H₄₈N₄O₂ 568.801

Alkaloid from the Papua New Guinea sponge *Petrosia contignata*. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv).

8-Hydroxy, 1R,2,3,4-tetrahydro, N²-Me: **1,2,3,4-Tetrahydro-2-N-methyl-8-hydroxymanzamine A**. 8-Hydroxy-2-N-methylmanzamine D

[160070-83-7]

C₃₇H₅₀N₄O₂ 582.828

Alkaloid from the Papua New Guinea sponges *Petrosia contignata* and *Cribrochalinia* sp. Cytotoxic to P388 leukemia cells. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Powder. [α]_D +46 (c, 0.006 in CH₂Cl₂). Props. refer to a salt (acid component not specified).

8-Hydroxy, 31-oxo, 32,33-dihydro: **Manzamine F**. **Keramamine B**

[107900-75-4]

C₃₆H₄₄N₄O₃ 580.769

Isol. from a *Xestospongia* sponge and from a *Petrosia* sponge. Cytotoxic. Shows antimicrobial activity and strong activity against *Mycobacterium*

tuberculosis (H₃₇Rv). Cryst. (MeCN). Mp 200° dec. $[\alpha]_D^{25} +59.9$ (c, 1.67 in CHCl₃). λ_{\max} 245 (ε 25600); 268 (ε 12100); 353 (ε 6800); 360 (ε 6820) (as AcOH salt) (Derep). λ_{\max} 220 (ε 36000); 244 (ε 31000); 265 (ε 13900); 355 (ε 8100) (MeOH) (Derep).

31β-Hydroxy, 32,33-dihydro: 32,33-Dihydro-31-hydroxymanzamine A

[581782-65-2]

C₃₆H₄₆N₄O₂ 566.785

Alkaloid from an Indonesian sponge. Cryst. (MeOH). $[\alpha]_D^{25} +34.4$ (c, 0.9 in CHCl₃). λ_{\max} 215; 248; 281; 291; 352; 359 (MeOH).

6,31β-Dihydroxy, 32,33-dihydro: 32,33-Dihydro-6,31-dihydroxymanzamine A

[581782-71-0]

C₃₆H₄₆N₄O₃ 582.784

Alkaloid from an Indonesian sponge. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Pale yellow powder. $[\alpha]_D^{25} +25.9$ (c, 0.5 in MeOH). λ_{\max} 218; 240; 280; 291; 356; 359 (MeOH).

12-Deoxy, 12,28α-epoxy: 12,28-Oxamanzamine A

[721429-59-0]

C₃₆H₄₂N₄O 546.754

Isol. from an *Acanthostrongylophora* sp. Amorph. solid. Mp 148° dec. $[\alpha]_D^{25} +38$ (c, 0.1 in CHCl₃). λ_{\max} 253 (log ε 3.81); 274 (log ε 3.66); 352 (log ε 3.41) (MeOH).

12-Deoxy, 12,34-epoxy: 12,34-Oxamanzamine A

C₃₆H₄₂N₄O 546.754

Alkaloid from an Indo-Pacific sponge. Shows activity against *Leishmania donovani*. Powder (MeOH). Mp 164° dec. $[\alpha]_D^{25} +40$ (c, 0.6 in CHCl₃). Not indexed in CA 138. λ_{\max} 252 (log ε 3.82); 271 (log ε 3.71); 358 (log ε 3.41) (MeOH).

12-Deoxy, 31-oxo, 12,28α-epoxy, 32,33-dihydro: 12,28-Oxamanzamine E

C₃₆H₄₂N₄O₂ 562.753

Alkaloid from an *Acanthostrongylophora* sp. Pale yellow solid. $[\alpha]_D^{25} +29.4$ (c, 0.2 in CHCl₃). λ_{\max} 250 (log ε 3.8); 281 (log ε 3.63); 352 (log ε 3.31); 361 (log ε 3.42) (MeOH).

12-Deoxy, 31-oxo, 12,34-epoxy, 32,33-dihydro: 12,34-Oxamanzamine E

C₃₆H₄₂N₄O₂ 562.753

Alkaloid from the Indonesian sponge *Acanthostrongylophora* sp. Pale yellow powder (MeOH). Mp 152° dec. $[\alpha]_D^{25} +44.3$ (c, 0.6 in CHCl₃). λ_{\max} 239 (log ε 3.38); 252 (log ε 3.82); 275 (log ε 3.65); 354 (log ε 3.41) (MeOH).

12-Deoxy, 6-hydroxy, 31-oxo, 12,34-epoxy, 32,33-dihydro: 6-Hydroxy-12,34-oxamanzamine E

C₃₆H₄₂N₄O₃ 578.753

Alkaloid from an *Acanthostrongylophora* sp. Yellow powder. Mp 163° dec. $[\alpha]_D^{25} +44.3$ (c, 0.4 in CHCl₃). λ_{\max} 252 (log ε 3.82); 275 (log ε 3.67); 356 (log ε 3.41) (MeOH).

12-Deoxy, 8-hydroxy, 12,28α-epoxy: 8-Hydroxy-12,28-oxamanzamine A

[721429-60-3]

C₃₆H₄₂N₄O₂ 562.753

Isol. from an *Acanthostrongylophora* sp. Amorph. powder (EtOH). Mp 160° dec. $[\alpha]_D^{25} +8$ (c, 0.1 in CHCl₃). λ_{\max} 251 (log ε 3.85); 273 (log ε 3.78); 356 (log ε 3.38) (MeOH).

(-)-form

8-Hydroxy: ent-Manzamine G

[403619-70-5]

C₃₆H₄₄N₄O₂ 564.769

Alkaloid from an undescribed sponge of the Petrosiidae. Exhibits activity against *Toxoplasma gondii* and strong antimalarial activity. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Yellowish powder (EtOH). Mp 196-198° dec. $[\alpha]_D^{25} -112$ (c, 0.12 in CHCl₃). λ_{\max} 266 (log ε 2.95); 282 (log ε 2.94); 390 (log ε 2.85) (MeOH).

8-Hydroxy, 31-oxo, 32,33-dihydro: ent-Manzamine F

[403619-71-6]

C₃₆H₄₄N₄O₃ 580.769

Alkaloid from an undescribed sponge of the Petrosiidae. Exhibits activity against *Toxoplasma gondii* and strong antimalarial activity. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Yellowish powder (EtOH). Mp 194° dec. $[\alpha]_D^{25} -44.6$ (c, 0.11 in CHCl₃). λ_{\max} 266 (log ε 3.04); 300 (log ε 3.02); 380 (log ε 2.92) (MeOH).

12-Deoxy, 31-oxo, 12,34-epoxy, 32,33-dihydro: ent-12,34-Oxamanzamine E

[479411-38-6]

C₃₆H₄₂N₄O₂ 562.753

Alkaloid from an Indo-Pacific sponge. Amorph. brown powder (CHCl₃). Mp 152° dec. $[\alpha]_D^{25} -54.6$ (c, 0.3 in CHCl₃). λ_{\max} 252 (log ε 3.82); 275 (log ε 3.65); 354 (log ε 3.41) (MeOH).

12-Deoxy, 8-hydroxy, 31-oxo, 12,34-epoxy, 32,33-dihydro: ent-12,34-Oxamanzamine F

[402935-15-3]

C₃₆H₄₂N₄O₃ 578.753

Alkaloid from an Indo-Pacific sponge. Shows strong activity against *Mycobacterium tuberculosis* (H₃₇Rv). Powder (MeOH). Mp 164° dec. $[\alpha]_D^{25} +40$ (c, 0.6 in CHCl₃). λ_{\max} 252 (log ε 3.82); 271 (log ε 3.71); 358 (log ε 3.41) (MeOH).

Sakai, R. *et al.*, *J.A.C.S.*, 1986, **108**, 6404-6405 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Nakamura, H. *et al.*, *Tet. Lett.*, 1987, **28**, 621-624 (isol, uv, pmr, cmr, cryst struct)

Eur. Pat., 1988, 272 056; *CA*, **109**, 129416p

(Manzamine D)

Ichiba, T. *et al.*, *Tet. Lett.*, 1988, **29**, 3083-3086 (Manzamines E, F)

Baldwin, J.E. *et al.*, *Tet. Lett.*, 1992, **33**, 2059-2062 (biosynth)

Ichiba, T. *et al.*, *J. Nat. Prod.*, 1994, **57**, 168-170 (8-Hydroxymanzamine A)

Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1737-1740 (3,4-Dihydromanzamine A, 6-Hydroxymanzamine A)

Crews, P. *et al.*, *Tetrahedron*, 1994, **50**, 13567-13574 (1,2,3,4-Tetrahydro-8-

hydroxymanzamine A, 1,2,3,4-Tetrahydro-2-N-methyl-8-hydroxymanzamine A)

Kobayashi, M. *et al.*, *Tetrahedron*, 1995, **51**, 3727-3736 (Manzamine X, Manzamine Y, isol, uv, ir, pmr, cmr, cryst struct)

Edrada, R.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1056-1060 (Manzamine A N-oxide, 3,4-Dihydromanzamine A N-oxide, 6-Deoxymanzamine X)

Tsuda, M. *et al.*, *Tetrahedron*, 1996, **52**, 2319-2324 (Manzamine D)

Tsuda, M. *et al.*, *Heterocycles*, 1997, **46**, 765-794 (rev)

Watanabe, D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 689-692 (Dihydrohydroxymanzamine A)

Winkler, J.D. *et al.*, *J.A.C.S.*, 1998, **120**, 6425-6426 (synth, Manzamine D)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (rev, synth)

Magnier, E. *et al.*, *Tetrahedron*, 1998, **54**, 6201-6258 (rev, synth)

Ang, K.K.H. *et al.*, *Antimicrob. Agents*

Chemother., 2000, **44**, 1645-1649 (activity)

Zhou, B.-N. *et al.*, *Tetrahedron*, 2000, **56**, 5781-5784 (1-Epimanzamine D, N-Me-1-Epimanzamine D)

El Sayed, K.A. *et al.*, *J.A.C.S.*, 2001, **123**,

1804-1808 (ent-Manzamine F, ent-Manzamine G, isol, activity)

Humphrey, J.M. *et al.*, *J.A.C.S.*, 2002, **124**, 8584-8592 (synth)

Yousaf, M. *et al.*, *Tetrahedron*, 2002, **58**, 7397-7402 (12,34-Oxamanzamine)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 2003, **66**, 823-828; 2004, **67**, 1314-1318 (32,33-Dihydrohydroxymanzamines, 6-Hydroxymanzamine E, 12,34-Oxamanzamine E)

Yousaf, M. *et al.*, *J. Med. Chem.*, 2004, **47**, 3512-3517 (12,28-Oxamanzamines)


Rao, K.V. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1034-1040 (12,28-Oxamanzamine E, 6-Hydroxy-12,34-oxamanzamine E)

Zhang, B. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 866-869 (isol, pmr, cmr, cryst struct)

Manzamine B

[112663-92-0]

M-92



C₃₆H₄₆N₄O 550.786

Alkaloid from the marine sponges *Haliclona*, *Amphimedon* and *Ircinia* spp. Shows activity against *Mycobacterium tuberculosis* (H₃₇Rv). Cryst. (EtOAc). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 198-203°. $[\alpha]_D^{20} +89$ (c, 1.8 in CHCl₃). λ_{\max} 212 (ε 18000); 235 (ε 22000); 240 (sh) (ε 20000); 250 (sh) (ε 15000); 282 (sh) (ε 6900); 288 (ε 11000); 338 (ε 3500); 351 (ε 3500) (MeOH) (Derep).

1β,2,3,4-Tetrahydro: 1,2,3,4-Tetrahydro-manzamine B

C₃₆H₅₀N₄O 554.817

Alkaloid from the sponge *Amphimedon* sp. Shows activity against *Mycobacterium tuberculosis* (H₃₇Rv). Amorph. solid. $[\alpha]_D^{25}$ -16 (c, 0.14 in MeOH). λ_{\max} 225 (ε 9700); 273 (ε 2500); 280 (ε 2600); 289 (ε 2100) (MeOH).

8-Hydroxy: 8-Hydroxymanzamine B

$C_{36}H_{46}N_4O_2$ 566.785

Alkaloid from an *Acanthostrongylophora* sp. Amorph. brown powder. $[\alpha]_D^{25}$ +39.6 (c, 0.2 in CHCl₃). λ_{\max} 241 (log ε 3.78); 252 (log ε 3.65); 278 (log ε 3.61); 291 (log ε 3.19); 334 (log ε 3.26); 352 (log ε 3.41) (MeOH).

Sakai, R. *et al.*, *Tet. Lett.*, 1987, **28**, 5493-5496 (*isol, uv, ir, pmr, cmr, cryst struct*)

Baldwin, J.E. *et al.*, *Tet. Lett.*, 1992, **33**, 2059-2062 (*biosynth*)

Tsuda, M. *et al.*, *Heterocycles*, 1997, **46**, 765-794 (*rev*)


Tsuda, M. *et al.*, *Heterocycles*, 1999, **50**, 485-488 (*Tetrahydromanzamine B*)

El Sayed, K.A. *et al.*, *J.A.C.S.*, 2001, **123**, 1804-1808 (*activity*)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1034-1040 (*8-Hydroxymanzamine B*)

Manzamine C

[112693-24-0]



$C_{22}H_{29}N_3$ 347.502

Alkaloid from the marine sponge *Haliclona* sp. and from *Amphimedon* sp. Shows activity against *Mycobacterium tuberculosis* (H₃₇Rv). Plates (CHCl₃/MeCN). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 77-82°. λ_{\max} 212 (ε 13500); 234 (ε 22000); 239 (sh) (ε 21000); 248 (sh) (ε 14000); 282 (sh) (ε 6100); 287 (ε 9500); 335 (ε 3000); 350 (ε 3000) (MeOH) (Derep).

1ξ,2,3,4-Tetrahydro: Keramamine C

[157207-88-0]

$C_{23}H_{33}N_3$ 351.534

Alkaloid from the Okinawan marine sponge *Amphimedon* sp. Plausible biogenetic precursor of Manzamine C, M-93. Oil. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{25}$ +20 (c, 0.92 in MeOH). λ_{\max} 225 (sh); 271 (ε 5600); 285 (sh); 290 (sh) (MeOH) (Derep).

Sakai, R. *et al.*, *Tet. Lett.*, 1987, **28**, 5493-5496 (*isol, uv, ir, pmr, cmr, cryst struct*)

Torisawa, Y. *et al.*, *Tetrahedron*, 1991, **47**, 8067-8078 (*synth*)

Baldwin, J.E. *et al.*, *Tet. Lett.*, 1992, **33**, 2059-2062 (*biosynth*)

Nowak, W. *et al.*, *Annalen*, 1993, 153-159 (*synth*)

Seki, H. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1173-1176 (*pmr, cmr*)

Tsuda, M. *et al.*, *Tet. Lett.*, 1994, **35**, 4387-4388 (*Keramamine C*)

Tsuda, M. *et al.*, *Heterocycles*, 1997, **46**, 765-794 (*rev*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (*rev, synth*)


Magnier, E. *et al.*, *Tetrahedron*, 1998, **54**, 6201-6258 (*rev, synth*)

MaGee, D.I. *et al.*, *Can. J. Chem.*, 2000, **78**, 1060-1066 (*synth*)

El Sayed, K.A. *et al.*, *J.A.C.S.*, 2001, **123**, 1804-1808 (*activity*)

Manzamine J

[139975-58-9]



$C_{36}H_{46}N_4O$ 550.786

Alkaloid from the Okinawan marine sponge *Ircinia* sp. and the Philippine sponge *Xestospongia ashmorica*. Exhibits cytotoxicity against L1210 murine leukaemia cells and KB human epidermoid carcinoma cells. Shows antimycobacterial activity. Mp 140°. $[\alpha]_D^{25}$ +47 (c, 2.0 in CHCl₃). λ_{\max} 238 (ε 15300); 278 (ε 9680); 290 (ε 8350); 346 (ε 4420); 358 (ε 4770) (as HCl salt) (Derep). λ_{\max} 219 (ε 22900); 236 (ε 18600); 280 (ε 10800); 290 (ε 11000); 346 (ε 5300); 357 (ε 5600) (MeOH) (Derep).

N²-Oxide: Manzamine J N-oxide

$C_{36}H_{46}N_4O_2$ 566.785

Alkaloid from *Xestospongia ashmorica*. Cytotoxic agent. Shows antimycobacterial activity. Insecticide. Yellow cryst. powder. $[\alpha]_D^{25}$ +15 (c, 0.40 in CHCl₃). λ_{\max} 261 (ε 25000); 325 (ε 19000) (MeOH).

3,4-Dihydro: 3,4-Dihydromanzamine J

[208392-24-9]

$C_{36}H_{48}N_4O$ 552.801

Alkaloid from *Amphimedon* sp. Shows antimycobacterial activity. Amorph. solid. $[\alpha]_D^{30}$ +50 (c, 0.1 in MeOH). λ_{\max} 209 (ε 13000); 242 (ε 9000); 322 (ε 4500) (MeOH).

1R,2,3,4-Tetrahydro: Manzamine H

[139975-57-8]

$C_{36}H_{50}N_4O$ 554.817

From *Ircinia* spp. and *Amphimedon* sp. Exhibits cytotoxicity against L1210 murine leukaemia cells and KB human epidermoid carcinoma cells. Shows antimycobacterial activity. Mp 145°. $[\alpha]_D^{25}$ +17 (c, 1.1 in CHCl₃). λ_{\max} 224 (ε 20000); 283 (ε 4600) (MeOH) (Derep). λ_{\max} 225 (ε 29000); 277 (ε 6600); 282 (ε 6800); 290 (ε 5500) (MeOH) (Derep).

1S,2,3,4-Tetrahydro: Manzamine L

M-94

[175274-50-7]

$C_{36}H_{50}N_4O$ 554.817

Alkaloid from the Okinawan marine sponge *Amphimedon* sp. Shows antimycobacterial activity. Amorph. solid. Mp 143°. $[\alpha]_D^{24}$ -15 (c, 0.42 in CHCl₃). λ_{\max} 223 (ε 34000); 283 (ε 6400) (MeOH) (Berdy).

8-Hydroxy: 8-Hydroxymanzamine J

$C_{36}H_{46}N_4O_2$ 566.785

Alkaloid from the Indonesian sponge *Acanthostrongylophora* sp. Pale yellow powder (CHCl₃). $[\alpha]_D^{25}$ +23.4 (c, 0.2 in CHCl₃). λ_{\max} 251 (log ε 3.62); 274 (log ε 3.68); 358 (log ε 3.39) (MeOH).

Kondo, K. *et al.*, *J.O.C.*, 1992, **57**, 2480 (*isol, uv, ir, pmr, cmr, ms, struct*)

Edrada, R.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1056 (*N-oxide*)

Tsuda, M. *et al.*, *Tetrahedron*, 1996, **52**, 2319 (*Manzamine L*)


Watanabe, D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 689-692 (*3,4-Dihydromanzamine J*)

El Sayed, K.A. *et al.*, *J.A.C.S.*, 2001, **123**, 1804-1808 (*activity*)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1314-1318 (*8-Hydroxymanzamine J*)

Manzamine M

M-95



$C_{36}H_{44}N_4O_2$ 564.769


Alkaloid from the sponge *Amphimedon* sp. Amorph. solid. $[\alpha]_D^{25}$ +16 (c, 0.48 in MeOH). λ_{\max} 205 (ε 14000); 233 (ε 3000); 276 (ε 1500); 359 (ε 1000) (MeOH).

Watanabe, D. *et al.*, *J. Nat. Prod.*, 1998, **61**, 689-692 (*isol, uv, ir, pmr, cmr*)

Mappicine

M-96

7-(1-Hydroxypropyl)-8-methylindolizino[1,2-b]quinolin-9(1H)-one



Probable
Absolute
Configuration

$C_{19}H_{18}N_2O_2$ 306.363

(S)-form [54318-59-1]

Minor alkaloid from *Mappia foetida* (Icacinaeae). Pale yellow cryst. + 2H₂O (MeOH). Mp 251-252°. $[\phi]_{500}$ -89° (c,

0.024 in dioxan). λ_{\max} 218 (ϵ 30200); 253 (ϵ 22387); 290 (ϵ 5495); 366 (ϵ 14790) (EtOH) (Berdy).

O- β -D-Glucopyranoside: *Mappicine glucoside*

[162109-68-4]
C₂₅H₂₈N₂O₇ 468.505

Alkaloid from trunk bark of *Nothapodytes foetida* (Icacinaeae). Cryst. (EtOH). Mp 184-185°. $[\alpha]_{\text{D}}^{25}$ -77.3 (c, 1.59 in H₂O).

O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: *Mappicine gentiobioside*

[162109-69-5]
C₃₁H₃₈N₂O₁₂ 630.647

From trunk bark of *Nothapodytes foetida* (Icacinaeae). Cryst. (EtOH). Mp 193-194°. $[\alpha]_{\text{D}}^{25}$ -110.7 (c, 0.66 in H₂O).

O-Ac:

Cryst. (MeOH). Mp 191-192°.

18-Ketone: *Mappicine ketone*. **Nothapodytine B**

[55854-89-2]
C₁₉H₁₆N₂O₂ 304.348

Alkaloid from stems of *Nothapodytes foetida*. Shows antiviral activity. Pale yellowish needles (CHCl₃). Mp 210-215°.

17-Hydroxy; O- β -D-glucopyranoside: **17-Hydroxymappicine glucoside**

[162109-70-8]
C₂₅H₂₈N₂O₈ 484.505

From trunk bark of *Nothapodytes foetida* (Icacinaeae). Cryst. (EtOH). Mp 164-165°. $[\alpha]_{\text{D}}^{25}$ -60.37 (c, 4.62 in H₂O).

19-Hydroxy: **19-Hydroxymappicine**

[123086-78-2]
C₁₉H₁₈N₂O₃ 322.363

Alkaloid from seeds of *Camptotheca acuminata*. Yellow cryst. (Me₂CO). Mp 245-248° dec.

9-Methoxy: **9-Methoxymappicine**

C₂₀H₂₀N₂O₃ 336.39

Alkaloid from *Nothapodytes foetida*. Yellow cryst. (MeOH). Mp 249-251°. $[\alpha]_{\text{D}}^{25}$ -9.6 (c, 0.62 in CHCl₃/MeOH). λ_{\max} 262 ; 319 ; 362 (MeOH).

9-Methoxy, O- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: **9-Methoxymappicine gentiobioside**

[162109-71-9]
C₃₂H₄₀N₂O₁₃ 660.674

From trunk bark of *Nothapodytes foetida* (Icacinaeae). Cryst. (EtOH). Mp 168-169°. $[\alpha]_{\text{D}}^{25}$ -27.7 (c, 1.3 in MeOH).

9-Methoxy, 18-ketone: **Nothapodytine A**

[142727-51-3]
C₂₀H₁₈N₂O₃ 334.374

Alkaloid from stems of *Nothapodytes foetida*. Pale yellowish needles (CHCl₃). Mp 235-238°.

(\pm)-**form** [54352-77-1]

Synthetic. Cryst. (MeOH). Mp 270-271° (264-266°).

Govindachari, T.R. *et al.*, *J.C.S. Perkin 1*, 1974, 1215 (*isol, uv, ir, pmr, ms, ord, struct*)
Adamovics, J.A. *et al.*, *Phytochemistry*, 1979, **18**, 1085 (*synth, pmr*)

Kingsbury, W.D. *et al.*, *Tet. Lett.*, 1988, **29**, 6847 (*synth, pmr*)

Lin, L.-Z. *et al.*, *Phytochemistry*, 1989, **28**, 1295 (*19-Hydroxymappicine*)

Pendrak, I. *et al.*, *J.O.C.*, 1994, **59**, 2623-2625; 1995, **60**, 3249-3251 (*activity*)

Pirillo, A. *et al.*, *J.C.S. Perkin 1*, 1995, 583 (*glycosides*)

Comins, D.L. *et al.*, *J.O.C.*, 1996, **61**, 9623 (*synth*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1996, **42**, 907 (*Nothapodytines*)

Josien, H. *et al.*, *Tetrahedron*, 1997, **53**, 8881 (*synth*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1998, **120**, 1218-1222 (*synth*)

Yadav, J.S. *et al.*, *Tetrahedron*, 1999, **55**, 5449-5456 (*Nothapodytine B, synth*)

Das, B. *et al.*, *Tetrahedron*, 1999, **55**, 7875-7880 (*synth*)

Mekouar, K. *et al.*, *J.O.C.*, 2000, **65**, 5212-5215 (*Nothapodytine B, synth*)

Toyota, M. *et al.*, *J.O.C.*, 2000, **65**, 7110-7113 (*synth*)

Das, B. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 135-140 (*9-Methoxymappicine*)

Das, B. *et al.*, *Synth. Commun.*, 2000, **30**, 3321-3325 (*Nothapodytines*)

Zhang, Q. *et al.*, *J.A.C.S.*, 2002, **124**, 5774-5781 (*synth*)

Carles, L. *et al.*, *J.O.C.*, 2002, **67**, 4304-4308 (*Nothapodytine B, synth*)

Raolji, G.B. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 5059-5061 (*Nothapodytine B, synth*)

Henegar, K.E. *et al.*, *J. Het. Chem.*, 2003, **40**, 601-605 (*synth, pmr, cmr, ms, Nothapodytine B*)


Kato, I. *et al.*, *J.O.C.*, 2003, **68**, 7983-7989 (*Nothapodytine B, synth*)

Chavan, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 3941-3943 (*synth*)

Marcanine A

M-97

4-Methylbenzo[g]quinoline-2,5,10(1H)-trione, **9CI**. 4-Methyl-1H-1-aza-2,9,10-anthracenetrione. *Griffiazanone B* [157463-84-8]



C₁₄H₉NO₃ 239.23

Alkaloid from *Annona glabra* (pond apple), *Goniothalamus griffithii* and *Goniothalamus marcanii*. Cytotoxic agent. Yellow needles. Mp 240-242° Mp 249-251° Mp > 300°.

3-Methoxy: 3-Methoxy-4-methylbenzo[g]quinoline-2,5,10(1H)-trione, **9CI**. **Dielsquinone**

[104696-15-3]

C₁₅H₁₁NO₄ 269.256

Alkaloid from *Guatteria dielsiana* and *Goniothalamus marcanii*. Cytotoxic agent. Yellow-green amorph. powder. Mp 250-252°.

3-Methoxy, N-Me: **Marcanine B**

[249285-58-3]

C₁₆H₁₃NO₄ 283.283

Alkaloid from the stem bark of *Goniothalamus marcanii*. Cytotoxic agent. Orange powder. λ_{\max} 272 (log ϵ 4.36);

299 (sh) (log ϵ 4.08); 320 (sh) (log ϵ 3.86); 419 (log ϵ 3.3) (MeOH).

3-Methoxy, 6-hydroxy: **Marcanine D**

[249285-60-7]

C₁₅H₁₁NO₅ 285.256

Alkaloid from the stem bark of *Goniothalamus marcanii*. Cytotoxic agent. Yellow powder. λ_{\max} 275 (log ϵ 4.2); 295 (log ϵ 4.05) (MeOH).

3-Methoxy, 9-hydroxy, N-Me: **Marcanine E**

[249285-61-8]

C₁₆H₁₃NO₅ 299.282

Alkaloid from the stem bark of *Goniothalamus marcanii*. Cytotoxic agent. Orange powder. λ_{\max} 274 (log ϵ 4.13); 300 (log ϵ 3.95) (MeOH).

3-Methoxy, 11-hydroxy, N-Me: **Marcanine C**

[249285-59-4]

C₁₆H₁₃NO₅ 299.282

Alkaloid from the stem bark of *Goniothalamus marcanii*. Cytotoxic agent. Orange powder. λ_{\max} 272 (log ϵ 4.23); 306 (sh) (log ϵ 3.97); 323 (sh) (log ϵ 3.82); 419 (log ϵ 3.19) (MeOH).

Goulart, M.O.F. *et al.*, *Phytochemistry*, 1986, **25**, 1691-1695 (*Dielsquinone*)

Marcos, A. *et al.*, *Tetrahedron*, 1994, **50**, 12941-12952 (*synth*)

Zhang, Y.J. *et al.*, *Chin. Chem. Lett.*, 1998, **9**, 1029-1032 (*Griffiazanone B*)

Soonthornchareonnon, N. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1390-1394 (*Goniothalamus marcanii* consists)


Chang, F.-R. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 913-920 (*isol, pmr, cmr*)

Brisach-Wittmeyer, A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 3609-3610 (*Dielsquinone, synth*)

Marcfortine B

M-98

[75789-29-6]



C₂₇H₃₃N₃O₄ 463.575

Alkaloid from *Penicillium roquefortii* B26. Mycotoxin. Antiparasitic and nematocidal insecticide. Anthelmintic agent. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 178-180°. $[\alpha]_{\text{D}}^{22}$ -67.7 (c, 1.77 in CHCl₃). λ_{\max} 229 (ϵ 2755) (MeOH).

N²-Me: **Marcfortine A**

[75731-43-0]

C₂₈H₃₅N₃O₄ 477.602

Alkaloid from *Penicillium roquefortii* B26 and *Penicillium* sp. UC7780. Mycotoxin. Antiparasitic and nematocidal insecticide. Anthelmintic agent. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 226 (ϵ 30080); 286 (ϵ 4625) (MeOH).

Polonsky, J. *et al.*, *Chem. Comm.*, 1980, 601-602 (*Marcfortine A, cryst struct*)

Prangé, T. *et al.*, *Tet. Lett.*, 1981, 1977-1980 (*uv, pmr, cmr, ms, struct*)

Lee, B.H. *et al.*, *Tet. Lett.*, 1994, **35**, 1135-1136 (*abs config*)

Kuo, M.S. *et al.*, *J. Antibiot.*, 1996, **49**, 1006-1013 (*biosynth*)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (rev, synth, biosynth)
Trost, B.M. *et al.*, *J.A.C.S.*, 2007, **129**, 3086-3087 (synth)

Marckidine**M-99**

[11050-74-1]

C₂₈H₃₅N₃O₃ 461.603


Struct. unknown. Alkaloid from the root bark of *Alangium lamarckii* (Alangiaceae). Mp 228°. [α]_D²⁵ -84 (Py).

Siddiqui, S. *et al.*, *Pak. J. Sci. Ind. Res.*, 1965, **8**, 161-165; *CA*, **68**, 899d

Raffauf, R.A. *et al.*, *Handb. Alkaloids Alkaloid-Containing Plants*, Wiley, 1970,

Maremycin A**M-100**

[165467-67-4]

C₁₇H₂₁N₃O₄S 363.437

Probable abs. config. depicted. Isol. from the marine *Streptomyces* sp. B9173. Sol. MeOH, DMSO; fairly sol. CHCl₃, EtOAc, Me₂CO, CH₂Cl₂; poorly sol. H₂O. Mp 229°. [α]_D²⁰ -120.95 (c, 0.21 in MeOH). λ_{\max} 209 (ε 33890); 257 (ε 3715) (MeOH). λ_{\max} 208 (ε 35450); 257 (ε 3715) (MeOH/HCl). λ_{\max} 212 (ε 27540); 248 (ε 3236) (MeOH/NaOH).

3'-Epimer: Maremycin B

[165877-95-2]

C₁₇H₂₁N₃O₄S 363.437

From *Streptomyces* sp. B9173. Shows sl. cytotoxicity. Sol. MeOH, DMSO; fairly sol. CHCl₃, CH₂Cl₂, EtOAc, Me₂CO; poorly sol. H₂O. Mp 216°. [α]_D²⁰ +2.94 (c, 0.21 in MeOH). λ_{\max} 209 (ε 29510); 258 (ε 4265) (MeOH). λ_{\max} 208 (ε 31620); 258 (ε 4265) (MeOH/HCl). λ_{\max} 212 (ε 25700); 252 (ε 7943) (MeOH/NaOH).

3'-Epimer, S-oxide(R-): Maremycin C₂

[333997-43-6]

C₁₇H₂₁N₃O₅S 379.436

Prod. by *Streptomyces* sp. strain GT 051237. Powder. [α]_D²⁰ +20 (c, 0.58 in DMSO). Data given is for mixt. with Maremycin C₁. λ_{\max} 215 (log ε 4.03); 258 (log ε 3.59); 293 (sh) (log ε 2.97) (EtOH).

3'-Epimer, S-oxide(S-): Maremycin C₁

[333997-39-0]


C₁₇H₂₁N₃O₅S 379.436

Prod. by *Streptomyces* sp. strain GT 051237. Obt. as a mixt. with Maremycin C₂.

Balk-Bindseil, W. *et al.*, *Annalen*, 1995, 1291-1294 (isol, uv, ir, pmr, cmr, ms, cd, struct)
Takayama, H. *et al.*, *Tetrahedron*, 1999, **55**, 6841-6846 (config)

Tang, Y.-Q. *et al.*, *Eur. J. Org. Chem.*, 2001, 261-267 (Maremycin B, Maremycin C, activity)

Ueda, T. *et al.*, *Org. Lett.*, 2008, **10**, 2043-2046 (synth)

Maremycin D₁**M-101**C₁₆H₁₇N₃O₄ 315.328

Prod. by *Streptomyces* sp. strain GT 051237. Powder. [α]_D²⁰ -30.2 (c, 0.9 in DMSO). Data shown is for mixt. with Maremycin D₂. λ_{\max} 218 (log ε 4.49); 253 (log ε 4.06); 293 (sh) (log ε 3.66) (EtOH).

3'-Epimer: Maremycin D₂C₁₆H₁₇N₃O₄ 315.328


Prod. by *Streptomyces* sp. strain GT 051237. Obt. as a mixt. with Maremycin D₁.

Tang, Y.-Q. *et al.*, *Eur. J. Org. Chem.*, 2001, 261-267 (isol, pmr, cmr)

Ueda, T. *et al.*, *Org. Lett.*, 2008, **10**, 2043-2046 (synth)

Margaritarine**M-102**

[138832-15-2]



Absolute Configuration


C₂₃H₂₇N₃O₂ 377.485

Alkaloid from the bark of *Margaritaria indica* (Euphorbiaceae). Amorph. solid. Mp 66-68° (softens). [α]_D²⁰ +106.9 (c, 0.06 in EtOH).

Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1991, 1863-1869 (isol, ir, pmr, cmr, ms, struct)

Margrapine A**M-103**

[185417-68-9]

C₂₁H₂₃NO₇ 401.415

Alkaloid from roots of Marsh grapefruit (*Citrus paradisi*). Yellow oil. [α]_D -16.7 (c, 0.036 in CHCl₃). λ_{\max} 225 (sh); 264; 295 (sh); 331; 381 (EtOH).

6-Me ether: Margrapine B

[185417-69-0]


C₂₂H₂₅NO₇ 415.442

From roots of Marsh grapefruit (*Citrus paradisi*). Yellow oil. [α]_D -37.5 (c, 0.008 in CHCl₃). λ_{\max} 226 (sh); 264; 331; 375 (EtOH).

Takemura, Y. *et al.*, *Heterocycles*, 1996, **43**, 2483 (isol, uv, ir, pmr, cmr, ms, struct)

Marinamide**M-104**

1,2-Dihydro-1-oxo-4-(1H-pyrrol-2-yl)-3-isoquinolinecarboxylic acid
[920020-06-0]

C₁₄H₁₀N₂O₃ 254.245

Prod. by mixed fermentation of two mangrove endophytic fungi (strain Nos. 1924 and 3893).


Me ester: [920020-07-1]C₁₅H₁₂N₂O₃ 268.271

Prod. by two mangrove endophytic fungi. Mp >° 300.

Zhu, F. *et al.*, *Chin. Sci. Bull.*, 2006, **51**, 1426-1430 (isol)


Marine Streptomyces**M-105****C₃₀H₃₈N₂O₅ lactam**

[184348-39-8]

C₃₀H₃₈N₂O₅ 506.641

Related to Aburatubolactam A, A-16. Prod. by *Streptomyces* sp. SCRC A20. Cytotoxic agent.

Japan. Pat., 1996, 96 245 625; *CA*, **126**, 17872d

Marinobactins**M-106**

Marinobactin A R = -CO(CH₂)₁₀CH₃
" B R = -CO(CH₂)₅CH=CH(CH₂)₅CH₃
" C R = -CO(CH₂)₁₂CH₃
" D₁ R = -CO(CH₂)₅CH=CH(CH₂)₅CH₃
" D₂ R = -CO(CH₂)₅CH=CH(CH₂)₇CH₃
" E R = -CO(CH₂)₁₄CH₃
" F R = octadecenoyl

Isol. from *Marinobacter* sp. strains DS40M6 and DS40M8. Siderophores.

Marinobactin A [265319-69-5]C₄₀H₆₉N₉O₁₆ 932.036**Marinobactin B** [265319-73-1]C₄₂H₇₁N₉O₁₆ 958.074**Marinobactin C** [265319-74-2]C₄₂H₇₃N₉O₁₆ 960.089**Marinobactin D₁** [265319-75-3]C₄₄H₇₅N₉O₁₆ 986.127

Marinobactin D₂ [265319-76-4]C₄₄H₇₅N₉O₁₆ 986.127**Marinobactin E** [265319-77-5]C₄₄H₇₇N₉O₁₆ 988.143**Marinobactin F** [960157-07-7]C₄₆H₇₉N₉O₁₆ 1014.181


Posn. of double bond in octadecenoyl residue not determined.

Martinez, J.S. *et al.*, *Science (Washington, D.C.)*, 2000, **287**, 1245-1247 (*isol, struct*)
 Martinez, J.S. *et al.*, *J. Inorg. Biochem.*, 2007, **101**, 1692-1698 (*isol, ms*)

Marinopyrrole A

M-107

[1010732-14-5]



Absolute Configuration

C₂₂H₁₂Cl₄N₂O₄ 510.159

Prod. by a marine-derived *Streptomyces* sp. (strain CNQ-418). Active against MRSA. [α]_D²⁰ -69 (c, 0.39 in MeOH). λ_{max} 264 (log ε 3.9); 315 (log ε 4) (MeOH).

4-Bromo: Marinopyrrole B

[1010732-15-6]


C₂₂H₁₁BrCl₄N₂O₄ 589.055

Prod. by *Streptomyces* sp. (strain CNQ-418). [α]_D²⁰ -72 (c, 0.2 in MeOH). λ_{max} 265 (log ε 4); 319 (log ε 3.9); 353 (log ε 4) (MeOH).

Hughes, C.C. *et al.*, *Org. Lett.*, 2008, **10**, 629-631 (*isol, cd, pmr, cmr, cryst struct*)

Maritidine

M-108



(+)-form

C₁₇H₂₁NO₃ 287.358**(+)-form** [22331-07-3]

Alkaloid from *Pancreatium maritimum*, *Hippeastrum ananuca*, *Narcissus tazetta* var. *chinensis*, *Narcissus papyraceus*, *Zephyranthes robusta*, *Zephyranthes sulphurea* and *Zephyranthes flava* (Amaryllidaceae). Prisms (EtOH/hexane). Mp 253-256° (263-265°). [α]_D²⁵ +31 (c, 0.09 in MeOH).

Picrate: Mp 192-194°.

Me ether: O-Methylmaritidine

[80550-22-7]

C₁₈H₂₃NO₃ 301.385

Minor alkaloid from the bulbs of *Narcissus tazetta* and from aerial parts of *Narcissus papyraceus* (Amaryllidaceae). Prisms (Et₂O). Mp 88-89°. [α]_D²³ +30.9 (c, 1 in CHCl₃).

Me ether, hydrochloride: Mp 243-245°.

Me ether, perchlorate: Mp 258-259°.

Me ether, dihydro:

Prisms (EtOAc). Mp 195-197°. [α]_D²³ +30 (c, 1.4 in CHCl₃).

O⁸-De-Me: Normaritimide. 8-O-De-methylmaritidine

[35182-60-6]

C₁₆H₁₉NO₃ 273.331

Alkaloid from the bulbs of *Hymenocallis rotata* (Amaryllidaceae). Prisms (Me₂CO). Mp 139-141°. [α]_D²³ +22.9 (c, 0.31 in MeOH).

O⁹-De-Me: 9-O-Demethylmaritidine

[120139-66-4]

C₁₆H₁₉NO₃ 273.331

Alkaloid from the whole plant of *Narcissus radiganorum* (Amaryllidaceae). Mp 238-239°.

3-Ketone: OxomaritimideC₁₇H₁₉NO₃ 285.342

Alkaloid from *Zephyranthes citrina*. Mp 140-142°. [α]_D²⁰ +55 (c, 0.29 in MeOH).

1,2-Dihydro, O⁸-de-Me: Maritinamine

[123931-00-0]

C₁₆H₂₁NO₃ 275.347

Alkaloid from *Sternbergia lutea* (Amaryllidaceae). Amorph. [α]_D²⁰ -20.3 (c, 1.8 in MeOH).

3-Epimer: 3-Epimaritimide

[28510-31-8]

C₁₇H₂₁NO₃ 287.358

Alkaloid from the fresh bulbs of *Zephyranthes rosea* (Amaryllidaceae). Microcryst. (CHCl₃ + trace MeOH). Mp 214-215°. [α]_D²² +83.2 (c, 0.47 in MeOH).

3-Epimer, O⁸-de-Me: Siculine

[123930-99-4]

C₁₆H₁₉NO₃ 273.331

Alkaloid from *Sternbergia sicula* (Amaryllidaceae). Amorph. [α]_D²⁰ -80 (c, 1.3 in CHCl₃).

3-Epimer, 1,2-dihydro, O⁸-de-Me: Epi-maritinamine

[123854-66-0]

C₁₆H₂₁NO₃ 275.347

Alkaloid from *Sternbergia lutea* (Amaryllidaceae). Amorph. [α]_D²⁰ -4 (c, 2.2 in MeOH).

3-Epimer, 1,2-dihydro, O⁹-de-Me, O³-Ac: Cantabricine

[171828-70-9]

C₁₈H₂₃NO₄ 317.384

Alkaloid from combined aerial parts and bulbs of *Narcissus cantabricus* (Amaryllidaceae). Mp 75-76°. [α]_D²⁰ -7.14 (c, 0.52 in MeOH).

(-)-form

O⁸-De-Me: (-)-Normaritimide. Macowine. 8-Hydroxy-9-methoxycrinine

C₁₆H₁₉NO₃ 273.331

Alkaloid from bulbs of *Crinum macowanii* and *Pancreatium maritimum*. Amorph. powder. Mp 115-117°. [α]_D²⁰ -34 (c, 0.23 in CHCl₃). The authors' semitrivial name based on crinine is confusing.

(±)-form [26722-76-9]

Synthetic. Mp 230-233° dec.

Sandberg, F. *et al.*, *J. Nat. Prod.*, 1963, **26**, 78

(isol, ir)

Wildman, W.C. *et al.*, *Alkaloids (Academic Press)*, 1968, 307 (*struct*)

DeAngelis, G.G. *et al.*, *Tet. Lett.*, 1969, 729 (*cd, ord*)

Schwartz, M.A. *et al.*, *J.A.C.S.*, 1970, **92**, 1090 (*synth, pmr*)

Kametani, T. *et al.*, *Tetrahedron*, 1971, **27**, 5441 (*synth*)

Kotani, E. *et al.*, *Chem. Comm.*, 1973, 550 (*synth*)

Yamada, S. *et al.*, *Tet. Lett.*, 1976, 57 (*synth, pmr*)

Tomioaka, K. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 2681 (*synth, ir, pmr*)

Zabel, V. *et al.*, *Cryst. Struct. Commun.*, 1979, **8**, 371 (*cryst struct*)

Rao, R.V.K. *et al.*, *Curr. Sci.*, 1979, **48**, 110 (*isol, ir*)

Tani, S. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 3381 (*O-Methylmaritidine*)

Ghosal, S. *et al.*, *Phytochemistry*, 1985, **24**, 635; 1986, **25**, 1975 (*isol*)

Kihara, M. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 1070 (*Normaritimide*)

Bastida, J. *et al.*, *Planta Med.*, 1988, **54**, 524 (*9-O-Demethylmaritidine*)

Pabuççoğlu, V. *et al.*, *J. Nat. Prod.*, 1989, **52**, 785 (*Siculine, Maritinamine, Epimaritinamine*)

Suau, R. *et al.*, *Heterocycles*, 1990, **31**, 517 (*O-Methylmaritidine*)

Sener, B. *et al.*, *J. Chem. Soc. Pak.*, 1994, **16**, 275-279 (*(-)-Normaritimide*)

Bastida, J. *et al.*, *Phytochemistry*, 1995, **40**, 1549 (*Cantabricine*)

Kita, Y. *et al.*, *J.O.C.*, 1996, **61**, 5857-5864 (*synth*)

Youssef, D.T.A. *et al.*, *Planta Med.*, 1998, **64**, 669-670 (*isol, pmr, cmr, ms*)

Ley, S.V. *et al.*, *J.C.S. Perkin 1*, 1999, 1251-1252 (*synth*)

Nair, J. *et al.*, *Phytochemistry*, 2000, **54**, 945-950 (*Macowine*)

Herrera, M.R. *et al.*, *Planta Med.*, 2001, **67**, 191-193 (*Oxomaritimide*)

Abou-Donia, A.H. *et al.*, *Planta Med.*, 2002, **68**, 379-381 (*Normaritimide, synth*)

Bru, C. *et al.*, *Org. Lett.*, 2003, **5**, 1845-1846 (*synth*)

Kodama, S. *et al.*, *Tetrahedron*, 2004, **60**, 4901-4907 (*Siculine, synth*)


Roe, C. *et al.*, *Org. Lett.*, 2008, **10**, 189-192 (*synth*)

Martefragine A

M-109

[200809-93-4]

[188546-50-1]



Absolute Configuration

C₂₀H₂₅N₃O₃ 355.436


Alkaloid from the marine alga *Martensia fragilis*. Inhibitor of lipid peroxidation in rat microsomes. Powder. Mp 147-148°. [α]_D²⁶ -20.3 (c, 0.76 in MeOH). λ_{max} 224 (ε 20200); 248 (sh) (ε 8600); 295 (sh) (ε 8800); 323 (ε 14200) (MeOH).

Takahashi, S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1527-1529 (*isol, uv, ir, pmr, cmr, cryst struct*)

Nishida, A. *et al.*, *Tet. Lett.*, 1998, **39**, 5983-5986 (*synth, cd, pmr, cmr, abs config*)

Martensine A M-110

2-(Hydroxy-1H-indol-3-ylmethyl)-1-methyl-5-(2-methylbutyl)-4-imidazolide, 9CI
[87168-35-2]



C₁₈H₂₅N₃O₂ 315.414

Alkaloid from the marine red alga *Martensia fragilis*. Shows antibiotic activity. [α]_D +42 (c, 0.34 in MeOH). λ_{max} 223 (ε 20900); 257 (sh) (ε 5200); 268 (sh) (ε 5500); 280 (ε 5800); 288 (ε 5200) (EtOH) (Derep). λ_{max} 260; 280; 288 (MeOH) (Berdy). λ_{max} 223 (ε 20900); 280 (ε 5800); 288 (ε 5200) (EtOH) (Berdy).

10-Epimer: 10-Epimartensine A

[87246-79-5]

C₁₈H₂₅N₃O₂ 315.414

Minor alkaloid from *Martensia fragilis*. λ_{max} 223 (ε 20900); 257 (sh) (ε 5200); 268 (sh) (ε 5500); 280 (ε 5800); 288 (ε 5200) (EtOH) (Derep).

10-Ketone: Martensine B

[87168-37-4]


C₁₈H₂₃N₃O₂ 313.399

Alkaloid from *Martensia fragilis*. Mp 184-186°. [α]_D -18 (c, 1.1 in Me₂CO). λ_{max} 267 (ε 10000); 335 (ε 10000) (EtOH/NaOH) (Derep). λ_{max} 215 (ε 10000); 244 (ε 11000); 261 (ε 8400); 303 (ε 10000) (EtOH) (Derep).

Kirkup, M.P. *et al.*, *Tet. Lett.*, 1983, **24**, 2087 (*isol, uv, ir, pmr, cmr, ms, struct*)

Martinelline M-111

[164178-52-3]



C₃₃H₅₂N₁₀O₂ 620.84

Ikeda *et al* suggest the difference in optical rotations between the synthetic and natural forms is due to the natural products being isolated in nearly racemic form. Alkaloid from roots of *Martinella iquitosensis* (Bignoniaceae). Bradykinin receptor antagonist. Pale yellow amorph. solid. [α]_D +9.4 (c, 0.02 in MeOH) (natural form). [α]_D²⁸ -108 (c, 0.09 in MeOH) (synthetic). λ_{max} 314 (ε 12500) (H₂O at pH 7.5) (Derep).

Parent acid: Martinellie acid

[164178-53-4]

C₂₇H₄₁N₇O₂ 495.667

Alkaloid from roots of *Martinella*

iquitosensis (Bignoniaceae). Bradykinin receptor antagonist. Pale yellow amorph. solid. [α]_D -8.5 (c, 0.01 in MeOH) (natural form). [α]_D²⁹ -164.3 (c, 0.14 in MeOH) (synthetic). λ_{max} 300 (ε 11600) (MeOH) (Derep).

Witherup, K.M. *et al.*, *J.A.C.S.*, 1995, **117**, 6682 (*isol, uv, ir, pmr, cmr, struct*)

Ma, D. *et al.*, *Org. Lett.*, 2001, **3**, 2189-2191 (*Martinellie acid, synth*)

Snider, B.B. *et al.*, *Org. Lett.*, 2001, **3**, 4217-4220 (*Martinellie acid, synth*)

Powell, D.A. *et al.*, *Org. Lett.*, 2002, **4**, 2913-2916 (*synth*)

Xia, C. *et al.*, *Tet. Lett.*, 2002, **43**, 9405-9409 (*synth*)

He, Y. *et al.*, *Tetrahedron*, 2006, **62**, 8755-8769 (*Martinellie acid, synth*)

Ikeda, S. *et al.*, *Chem. Comm.*, 2007, 504-506 (*synth, Martinellie acid*)


Badarinarayana, V. *et al.*, *Tet. Lett.*, 2007, **48**, 2607-2610 (*Martinellie acid, synth*)

Miyata, O. *et al.*, *Tetrahedron*, 2007, **63**, 10092-10117 (*synth*)

Shirai, A. *et al.*, *J.O.C.*, 2008, **73**, 4464-4475 (*Martinellie acid, synth*)

Massadine M-112

[662145-41-7]



C₂₂H₂₄Br₄N₁₀O₅ 828.112

Alkaloid from the sponge *Stylissa* aff. *massa*. Yellow powder. [α]_D¹⁷ -12 (c, 0.1 in MeOH). λ_{max} 278 (ε 20000) (MeOH).

14-Deoxy, 14α-chloro: Massadine chloride

[952523-62-5]


C₂₂H₂₃Br₄ClN₁₀O₄ 846.557

Isol. from *Stylissa* aff. *massa*. Powder. [α]_D²⁰ -14.9 (c, 0.5 in MeOH).

Nishimura, S. *et al.*, *Org. Lett.*, 2003, **5**, 2255-2257 (*isol, cd, pmr, cmr, ms*)

Grube, A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 6721-6724 (*Massadine chloride*)

Su, S. *et al.*, *J.A.C.S.*, 2008, **130**, 16490-16491 (*synth*)

Massadine aziridine M-113


C₂₂H₂₂Br₄N₁₀O₄ 810.096

There is indirect evidence for the occurrence of this alkaloid. Probably occurs in sponges. Precursor of pyrrole-imidazole dimers.

Grube, A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 6721-6724; 8107 (*occur*)

Matadine M-114

[138195-38-7]




C₁₉H₂₀N₂O 292.38

Abs. config. at C-15 tentatively assigned as *S*- (illus.) on biogenetic grounds. Alkaloid from the root bark of *Strychnos gossweileri* (Loganiaceae). Cytotoxic. λ_{max} 215 (ε 52600); 253 (ε 21950); 307 (ε 14200) (MeOH) (Berdy).

Quetin-Leclercq, J. *et al.*, *Phytochemistry*, 1991, **30**, 1697-1700 (*isol, struct*)

Matemone M-115

6-Bromo-1,2-dihydro-2-hydroxymethyl-2-methoxy-3H-indol-3-one
[288849-91-2]



C₁₀H₁₀BrNO₃ 272.098


Isol. from the sponge *Ietrochota purpurea*. Mild cytotoxic agent. Yellow-green film. [α]_D²⁵ +8.9 (c, 0.45 in MeOH). λ_{max} 228 (log ε 4.34); 245 (log ε 4.33); 273 (log ε 4.01); 399 (log ε 3.5) (MeOH).

Carletti, I. *et al.*, *J. Nat. Prod.*, 2000, **63**, 981-983 (*isol, pmr, cmr, uv*)

***Saccharomyces kluyveri* α-** M-116**Mating pheromone 1**

α^{sk1} Mating pheromone

[103488-73-9]




C₈₂H₁₀₆N₁₈O₂₀S 1695.914

Prod. by mating type α cells of *Saccharomyces kluyveri*. Mating pheromone.

Sakamoto, M. *et al.*, *Agric. Biol. Chem.*, 1986, **50**, 1279-1285

Matlystatin D

[140667-42-1]

R = (CH₂)₄CH₃

C₂₇H₄₄N₆O₆ 548.681
 Prod. by *Actinomadura atramentaria*
 SANK 61488. Collagenase inhibitor.
 Powder. Mp 122-126°. [α]_D²⁰ -20.3 (c, 1.14
 in MeOH). λ_{max} 229 (sh) (ε 7690)
 (MeOH) (Derep).

Ogita, T. et al., *J. Antibiot.*, 1992, **45**, 1723;
 1733; 1994, **47**, 1473 (*isol. struct.*
props)

Matlystatin E

[140638-26-2]

As Matlystatin D, M-117 with


R = (CH₂)₃CH₃

C₂₆H₄₂N₆O₆ 534.654
 Prod. by *Actinomadura atramentaria*
 SANK 61488. Collagenase inhibitor.
 Powder. Mp 89-94°. [α]_D²⁰ -21.95 (c, 0.41
 in MeOH). λ_{max} 229 (sh) (ε 7690)
 (MeOH) (Derep). λ_{max} 228 (ε 8770)
 (MeOH) (Berdy).

Ogita, T. et al., *J. Antibiot.*, 1992, **45**, 1723;
 1733 (*isol. struct. props*)

Matlystatin F

[140638-25-1]




C₂₇H₄₄N₆O₆ 548.681
 Prod. by *Actinomadura atramentaria*
 SANK 61488. Collagenase inhibitor.
 Powder. Mp 69-75°. [α]_D²⁰ +1.43 (c, 0.14 in
 MeOH). λ_{max} 229 (sh) (ε 7690) (MeOH)
 (Derep). λ_{max} 228 (ε 7030) (MeOH)
 (Berdy).

Ogita, T. et al., *J. Antibiot.*, 1992, **45**, 1723;
 1733; 1994, **47**, 1473 (*isol. struct.*
props)

M-117

Matopensine16,16'-Dihydro-17,17'-oxybisnordihydro-
toxiferine

[88761-31-3]



C₃₈H₄₂N₄O 570.776
 Alkaloid from *Strychnos matopensis*,
Strychnos kasengaensis, *Strychnos min-*
fiensis and *Strychnos panganensis*
 (Loganiaceae). Amorph. [α]_D +105 (c,
 0.6 in MeOH). λ_{max} 217 (log ε 4.46);
 292 (log ε 3.92); 315 (log ε 3.95)
 (EtOH).

Mono-N-oxide: Matopensine mono-N-
oxide. Matopensine N-oxide

[94898-71-2]

C₃₈H₄₂N₄O₂ 586.775

Alkaloid from *Strychnos kasengaensis*
 and *Strychnos matopensis* root bark
 (Loganiaceae). λ_{max} 217 ; 263 ; 313
 (EtOH).

18-Hydroxy: 18-Hydroxymatopensine

[119308-24-6]

C₃₈H₄₂N₄O₂ 586.775

Alkaloid from root bark of *Strychnos*
matopensis. [α]_D +68 (c, 0.3 in MeOH).
 λ_{max} 218 ; 263 ; 313 (MeOH).

18,18'-Dihydroxy: 18,18'-Dihydroxyma-
topensine

[119308-25-7]

C₃₈H₄₂N₄O₃ 602.775

Alkaloid from root bark of *Strychnos*
matopensis. λ_{max} 215 ; 263 ; 310
 (MeOH).

16'-Epimer, 16-methoxy: 16-Methoxyiso-
matopensine

[119308-22-4]

C₃₉H₄₄N₄O₂ 600.802

Alkaloid from root bark of *Strychnos*
matopensis. λ_{max} 217 ; 265 ; 315
 (MeOH).

16'-Epimer, 16-ethoxy: 16-Ethoxyiso-
matopensine

[119308-23-5]

C₄₀H₄₆N₄O₂ 614.829

Alkaloid from root bark of *Strychnos*
matopensis. [α]_D -13 (c, 0.3 in CHCl₃).
 λ_{max} 216 ; 257 ; 293 (MeOH).

Massiot, G. et al., *Heterocycles*, 1983, **20**,
 2339-2342 (*isol. struct. pmr, biosynth*)
 Thepenier, P. et al., *Phytochemistry*, 1984,
23, 2659-2663 (*isol. uv, ir, pmr, ms,*
oxide)


Massiot, G. et al., *Phytochemistry*, 1988, **27**,
 3293-3304 (*Strychnos matopensis*
constit)

Nuzillard, J.M. et al., *Phytochemistry*, 1996,
43, 897-902 (*isol*)

M-120

Matrine

Matridin-15-one, 9CI. Lupanidine. Dode-
 cahydro-1H,5H,10H-dipyrido[2,1-
 f:3',2',1'-ij][1,6]naphthyridin-10-one
 [519-02-8]

Absolute
ConfigurationC₁₅H₂₄N₂O 248.367

Stereoisomer of Sophoridine, S-380, Iso-
 matrine, I-244, Allomatrine, A-628 and
 Darvasamine, D-79. The Matrine skele-
 ton contains four chiral centres leading to
 a total of 8 possible diastereoisomers. Of
 these, 6 are currently known (4 or 5 of
 which are natural alkaloids). However,
cis-Matrine, having a *cis*-CD ring junction
 (diastereoisomeric at N) has recently
 been prepd. so that up to 32 diastereoi-
 somers may be possible. Alkaloid from
Sophora alopecuroides, *Sophora flaves-*
cens, *Sophora microphylla*, *Sophora pa-*
chycarpa, *Sophora tetraptera*, many other
Sophora spp., *Vexibia pachycarpa* (pre-
 ferred genus name *Sophora*) and *Eu-*
chresta horsfeldii (Fabaceae). Has
 antiulcer, antineoplastic and antibacterial
 props. Reverse transcriptase inhibitor.
 Sol. H₂O, C₆H₆, Et₂O, CHCl₃; fairly sol.
 hexane. Mp 77°. [α]_D¹⁵ +40.9 (H₂O). Log P
 0.58 (uncertain value) (calc).

▶ LD₅₀ (mus, ivn) 150 mg/kg; (rat, ipr) 125
 mg/kg.

Hydrobromide: Mp 272-275°.

Perchlorate: Mp 214.5-216°.

N¹-Oxide: Oxymatrine. Matrine N-oxide.
Ammothamnine. Pachycarpidine

[16837-52-8]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Sophora flavesces*, *Soph-*
ora macrocarpa, *Sophora alopecur-*
oides, *Ammothamnus lehmannii*
 (preferred genus name *Sophora*), *Eu-*
chresta horsfeldii and other plants
 (Fabaceae). Mp 208° (148-150°). [α]_D
 +47 (EtOH).

▶ OQ1770000

3α-Hydroxy: 3α-Hydroxymatrine

[81970-01-6]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from leaves of *Sophora*
macrocarpa (Fabaceae).

5-Hydroxy: see Sophoranol, S-377

9α-Hydroxy: 9α-Hydroxymatrine

[88509-92-6]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from the leaves of *Sophora*
macrocarpa (Fabaceae). Needles (pet-
 rol). Mp 158-159°. [α]_D²⁰ +25.4 (c, 1.0 in
 MeOH).

13α-Hydroxy: 13α-Hydroxymatrine

[148409-26-1]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Sophora viciifolia*.

Possesses antiulcer activity.

14 α -Hydroxy: 14 α -Hydroxymatrine

[225781-48-6]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Sophora tonkinensis*.

Mp 120°. [α]_D²⁵ +30.4.

14 α -Acetoxy: 14 α -Acetoxymatrine

[225781-46-4]

C₁₇H₂₆N₂O₃ 306.404

Alkaloid from *Sophora tonkinensis*.

Cryst. Mp 96°. [α]_D²⁵ +33 (EtOH).

14 β -Hydroxy: 14 β -Hydroxymatrine

[183074-18-2]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from roots of *Sophora tonkinensis*. Needles (CH₂Cl₂/hexane). Mp 69-70°. [α]_D²¹ -8.3 (c, 0.28 in EtOH).

14 β -Hydroxy, N¹-oxide: 14 β -Hydroxyoxymatrine. 14 β -Hydroxymatrine N¹-oxide

[914776-54-8]

C₁₅H₂₄N₂O₃ 280.366

Alkaloid from the roots and rhizomes of *Sophora tonkinensis*. Pale yellow oil. [α]_D²⁰ -5.3 (c, 0.3 in MeOH). λ _{max} 208 (log ϵ 3.66) (MeOH).

14 β -Acetoxy: 14 β -Acetoxymatrine

[225781-44-2]

C₁₇H₂₆N₂O₃ 306.404

Alkaloid from *Sophora tonkinensis*.

Cryst. Mp 113°. [α]_D²⁵ -26.7 (EtOH).

Stereoisomer: Albertidine

[56832-52-1]

C₁₅H₂₄N₂O 248.367

Alkaloid from epigeal parts of *Leontice albertii* (Leonticeaceae). Mp 70-71°. [α]_D¹⁸ +33.8 (c, 0.52 in EtOH). Config. unknown.

Stereoisomer, perchlorate: Mp 254°.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1958, **91**, 2176-2189; 2189-2193; 1975, **108**, 1043-1051 (*ir. struct, config, cmr*)

Schütte, H.R. *et al.*, *Annalen*, 1965, **685**, 194-199 (*biosynth*)

Mandell, L. *et al.*, *J.A.C.S.*, 1965, **87**, 5234-5236 (*synth*)

Aslanov, A. *et al.*, *CA*, 1967, **67**, 71108 (*oxide*)
Kamalitidinov, D.D. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 409-412; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 340-342 (*Albertidine*)

Kojima, R. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 2555-2563 (*pharmacol*)

Vul'ison, N.S. *et al.*, *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1974, **10**, 221-229 (*ms*)

Ueno, A. *et al.*, *Chem. Pharm. Bull.*, (footnote), 1975, **23**, 2560-2566 (*abs config*)

Morinaga, K. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 2483-2488 (*pmr*)

Leeper, F.J. *et al.*, *Can. J. Chem.*, 1981, **59**, 106-115 (*biosynth*)

Negrete, R.E. *et al.*, *Bol. Soc. Chil. Quim.*, 1982, **27**, 263 (*3 α -Hydroxymatrine*)

Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 71-75; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 66-69 (*rev. stereochem*)

Negrete, R. *et al.*, *Phytochemistry*, 1983, **22**, 2069-2072 (*9 α -Hydroxymatrine*)

Abdusalomov, B.A. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 3-12; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 1-9 (*biosynth*)

Sakamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 2018-2023 (*synth*)

Gonnella, N.C. *et al.*, *Magn. Reson. Chem.*, 1988, **26**, 185-190 (*pmr, cmr*)

Zhu, Z. *et al.*, *CA*, 1993, **119**, 20256s (*13 α -Hydroxymatrine*)

Xiao, P. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1951-1953; 1999, **47**, 448-450 (*14-Hydroxymatrine, 14-Acetoxymatrine*)


Zhang, Z.-T. *et al.*, *Huaxue Xuebao*, 2003, **61**, 1058-1064; *CA*, **140**, 5196n (*cryst struct*)

Ding, P.-L. *et al.*, *Planta Med.*, 2006, **72**, 854-856 (*14 β -Hydroxyoxymatrine, Oxymatrine*)

Mattogrossine

M-122

13-Acetyl-5,6,7a,8,8a,9,12,12a,12b,13-decahydro-7a-hydroxy-10-methyl-10H-furo[2,3-d]pyrrolo[3,4-a]carbazole-9-carboxaldehyde, 9CI
[131989-71-4]



Absolute Configuration

C₂₁H₂₅NO₅ 371.432


Apparently an unusual secocuran alkaloid lacking one N (cf. Malagashanine, M-58). Alkaloid from the roots and branches of *Strychnos mattogrossensis* (Loganiaceae). Powder.

Angenot, L. *et al.*, *Phytochemistry*, 1990, **29**, 2746 (*isol, uv, ir, cd, ms, pmr, cmr, struct*)

Mauritamide A

M-123

[155210-56-3]




C₁₅H₂₂Br₂N₆O₅S 558.25

Alkaloid from the Fijian sponge *Agelas mauritiana*. Protein-tyrosine kinase inhibitor. Amorph. solid. [α]_D²⁰ +1.3 (c, 0.003 in MeOH). λ _{max} 254 (ϵ 6200); 258 (ϵ 6110); 278 (ϵ 3910) (MeOH) (Derep).

Jiménez, C. *et al.*, *Tet. Lett.*, 1994, **35**, 1375 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mauritiamine

M-124



C₂₂H₂₀Br₄N₁₀O₃ 792.081

(\pm)-form [175669-27-9]

Alkaloid from the marine sponge *Agelas mauritiana*. Antifouling substance. Exhibits moderate antibacterial activity. Solid. λ _{max} 272; 278 (ϵ 18400) (MeOH) (Berdy).


Tsakamoto, S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 501-503 (*isol, uv, ir, pmr, cmr*)

Olofson, A. *et al.*, *J.O.C.*, 1997, **62**, 7918-7919 (*synth*)

Mauritine A

M-125

[38478-72-7]



Absolute Configuration

R = -CH(CH₃)₂

C₃₂H₄₁N₅O₅ 575.706

Alkaloid from the bark of *Zizyphus mauritiana*, *Zizyphus amphibia* and *Zizyphus spinachristi*, and the stem bark of *Zizyphus jujuba* (Chinese date) (Rhamnaceae). Rhombs (CH₂Cl₂/petrol). Mp 104°. [α]_D²⁰ -315 (c, 0.33 in MeOH).

N-De-Me: Mauritine F

[55609-23-9]

C₃₁H₃₉N₅O₅ 561.68

Alkaloid from the bark of *Zizyphus mauritiana* and the root bark of *Zizyphus nummularia* (Rhamnaceae). Needles (CHCl₃/petrol). Mp 222-225°. [α]_D²⁰ -285 (c, 0.15 in MeOH).

1''R-Hydroxy: Mauritine E

[55609-22-8]

C₃₂H₄₁N₅O₆ 591.706

Alkaloid from the bark of *Zizyphus mauritiana* (Rhamnaceae). Amorph. [α]_D²⁰ -243 (c, 0.11 in MeOH).

Tschesche, R. *et al.*, *Tet. Lett.*, 1972, 2609-2612 (*isol, ms, struct*)

Tschesche, R. *et al.*, *Annalen*, 1974, 1694-1701 (*Mauritines E-F*)

Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 1633; 1976, **15**, 541-542 (*isol*)

Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (*Mauritine F, isol*)


Kirfel, A. *et al.*, *Z. Naturforsch., B*, 1976, **31**, 279-280 (*cryst struct*)

Cristau, P. *et al.*, *Chem. Eur. J.*, 2005, **11**, 2668-2679 (*synth, pmr, cmr*)

Mauritine B

M-126

[38478-73-8]



Absolute Configuration


C₃₅H₄₇N₅O₅ 617.787
Alkaloid from the bark of *Zizyphus mauritiana* (Rhamnaceae). Amorph. [α]_D²⁰ -151 (c, 0.44 in MeOH).

Tschesche, R. *et al.*, *Tet. Lett.*, 1972, 2609-2612 (*isol, ms*)

Cristau, P. *et al.*, *Chem. Eur. J.*, 2005, **11**, 2668-2679 (*synth, pmr, cmr*)

Mauritine C
[54578-03-9]

M-127



Absolute Configuration

C₂₈H₃₄N₄O₄ 490.601
Alkaloid from *Zizyphus mauritiana*, *Zizyphus nummularia* and *Zizyphus spinachristi* (Rhamnaceae). [α]_D²⁰ -224 (c, 0.11 in MeOH).


Tschesche, R. *et al.*, *Annalen*, 1974, 1694-1701 (*isol, uv, ir, pmr, ms, struct*)

Tschesche, R. *et al.*, *Phytochemistry*, 1974, **13**, 1633 (*isol*)

Cristau, P. *et al.*, *Chem. Eur. J.*, 2005, **11**, 2668-2679 (*synth, pmr, cmr*)

Mauritine D
[55609-21-7]

M-128



Absolute Configuration

C₃₃H₅₁N₅O₅ 597.796
Alkaloid from the bark of *Zizyphus mauritiana*, *Zizyphus nummularia*, *Zizyphus oenoplia* and *Zizyphus xylopyra* (Rhamnaceae). Amorph. [α]_D²⁰ -259 (c, 0.16 in MeOH).

10,11-Dihydro:
Powder. [α]_D²⁰ -111 (c, 0.41 in MeOH).

Tschesche, R. *et al.*, *Annalen*, 1974, 1694-1701 (*isol, uv, ir, pmr, ms*)

Cristau, P. *et al.*, *Chem. Eur. J.*, 2005, **11**, 2668-2679 (*synth*)

Mauritine H
[64309-18-8]

M-129

As Mauritine A, M-125 with

R = -CH₂CH(CH₃)₂

C₃₃H₄₃N₅O₅ 589.733
Alkaloid from the bark of *Zizyphus mauritiana* (Rhamnaceae). Cryst. (MeOH/petrol). Mp 212-215°. [α]_D²⁰ -169 (c, 0.013 in MeOH).


10,11-Dihydro:
Cryst. (EtOH aq.). Mp 166-169°. [α]_D²⁰ -128 (c, 0.013 in MeOH).

Tschesche, R. *et al.*, *Phytochemistry*, 1977, **16**, 1025-1028 (*isol, uv, ir, pmr, ms, struct*)

C-Mavacurine

M-130

19,20-Didehydro-17-hydroxy-4-methyl-1,16-cyclocorynanium, 9CI. Mavacurine [6801-19-0]



Probable absolute configuration

C₂₀H₂₅N₂O⁺ 309.43
Alkaloid from *Strychnos nux-vomica*, *Strychnos melinoniana*, *Strychnos variabilis* root bark, *Strychnos macrophylla*, *Strychnos divaricans*, *Strychnos toxifera*, *Strychnos amazonica*, *Strychnos mitscherlichii* and other *Strychnos* spp.; present in Calabash curare (*Logania-ceae*).

Iodide:

C₂₀H₂₅IN₂O 436.335
Cryst. [α]_D +247 (c, 0.338 in 50% Me₂CO aq.). Carmine col. with Ce(IV).

Picrate:

Cryst. Mp 179-180°.

N-De-Me: Normavacurine

[38836-02-1]

C₁₉H₂₂N₂O 294.396

Alkaloid from *Strychnos minfiensis*, *Strychnos potatorum* and *Strychnos longicaudata* (Loganiaceae). Amorph. solid. [α]_D +127 (c, 0.67 in CHCl₃). Erroneously assigned the struct. 1,2-Dehydrodeacetylretuline when first isolated from *S. longicaudata*.

Wieland, T. *et al.*, *Chem. Ber.*, 1952, **85**, 731 (*isol*)

Asmis, H. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1968 (*isol*)

Hesse, M. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 878; 1965, **48**, 674 (*uv, ir, pmr, ms, struct*)

Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 1 (*isol*)

Angenot, L. *et al.*, *Planta Med.*, 1981, **42**, 364-370; 371-374 (*isol, ms, pmr*)

Calverley, M.J. *et al.*, *Tet. Lett.*, 1981, 1635 (*synth, bibl*)


Coune, C.A. *et al.*, *J. Pharm. Belg.*, 1982, **37**, 189 (*cmr*)

Massiot, G. *et al.*, *Heterocycles*, 1989, **29**, 1435 (*Normavacurine*)

Maxonine

M-131

5-Methylindolo[2',3':3,4]pyrido[1,2-b][2,7]naphthyridin-14(5H)one, 9CI [125797-63-9]



C₁₉H₁₃N₃O 299.331

Struct. revised in 1993. Alkaloid from the roots of *Simira maxonii* (Rubiaceae). Yellow oil. [α]_D²⁰ -85 (c, 0.2 in CHCl₃).


Hasbun, C.P. *et al.*, *Tet. Lett.*, 1989, **30**, 6199 (*isol, uv, ir, pmr, cmr, ms*)

Kelly, T.R. *et al.*, *Tet. Lett.*, 1993, **34**, 6173 (*synth, struct*)

Mayotamide A

M-132

[217449-23-5]



C₃₀H₄₃N₇O₄S₄ 693.978

Isol. from *Didemnum molle*. Cytotoxic agent. Amorph. powder. [α]_D +77 (c, 0.3 in MeOH).

Rudi, A. *et al.*, *Tetrahedron*, 1998, **54**, 13203-13210 (*isol, pmr, cmr, ms*)

Mayotamide B

M-133

[217449-24-6]

As Mayotamide A, M-132 with

R = CH₃

C₂₉H₄₁N₇O₄S₄ 679.951


Isol. from *Didemnum molle*. Cytotoxic agent. Amorph. powder. [α]_D +130 (c, 0.1 in MeOH).

Rudi, A. *et al.*, *Tetrahedron*, 1998, **54**, 13203-13210 (*isol, pmr, cmr, ms*)

Maysenine

M-134

[52978-30-0]



Absolute Configuration

C₂₇H₃₃ClN₂O₆ 517.02

Alkaloid from *Maytenus buchananii* (Celastraceae). Antitumour agent. Mp 184-185°. [α]_D³⁰ -57 (c, 0.056 in EtOH). pK_a 12.2.

4S,5S-Epoxide: Normaysine

[52978-29-7]

C₂₇H₃₃ClN₂O₇ 533.02

Alkaloid from *Maytenus buchananii* (Celastraceae). Cryst. + 1/2 H₂O (Me₂CO). Mp 187-188°. [α]_D -217 (c, 0.051 in EtOH). λ _{max} 226; 241; 252; 280; 288 (MeOH) (Berdy).

4S,5S-Epoxide, N²²-Me: Maysine†

[52978-28-6]


C₂₈H₃₅ClN₂O₇ 547.046

Alkaloid from *Maytenus buchananii*

(Celastraceae). Mp 137-141°. $[\alpha]_D^{30}$ -173 (c, 0.23 in EtOH). λ_{\max} 226; 241; 252; 280; 288 (MeOH) (Berdy).

- Kupchan, S.M. *et al.*, *J.O.C.*, 1977, **42**, 2349-2357 (*Maysenine, Normaysine, Maysine, isol, uv, ir, pmr, ms, struct*)
 Corey, E.J. *et al.*, *J.A.C.S.*, 1978, **100**, 2916 (*synth*)
 Meyers, A.I. *et al.*, *J.A.C.S.*, 1983, **105**, 5015 (*Maysine, synth*)
 Keeffe, J.R. *et al.*, *J.A.C.S.*, 1988, **110**, 1982 (*uv*)

Maytansine, 9CI, USAN **M-135**
Maïtansine, INN. MAYT. NSC 153858
 [35846-53-8]



$\text{C}_{34}\text{H}_{46}\text{ClN}_3\text{O}_{10}$ 692.204

Ansamycin-type antibiotic. Isol. from *Maytenus ovatus* and *Maytenus senata* (Celastraceae). Shows antileukaemic, cytotoxic, antitubulin and antimetabolic props. Most potent of the maytansinoids. Has been evaluated clinically against various human carcinomas but is not a practically effective antitumour agent. Shows antimetabolic, antigibberellin and auxin activity in plants. Sol. MeOH, Et₂O, CHCl₃; poorly sol. H₂O. Mp 171-172°. $[\alpha]_D^{26}$ -145 (c, 0.055 in CHCl₃). Log P 3.02 (calc). λ_{\max} 233 (€ 29800); 243 (sh) (€ 27100); 254 (€ 27200); 282 (€ 5690); 290 (€ 5520) (EtOH) (Derep).

- Gastrointestinal and CNS adverse effects when used therapeutically. LD₅₀ (rat, scu) 480 mg/kg. OQ2290000

N-De-Ac, N-propanoyl: Maytanprine
 [38997-09-0]

$\text{C}_{35}\text{H}_{48}\text{ClN}_3\text{O}_{10}$ 706.231

Isol. from *Maytenus buchananii* (Celastraceae). Antileukaemic antibiotic. Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 169-170°. $[\alpha]_D^{30}$ -125 (c, 0.0559 in CHCl₃). Log P 3.55 (calc). λ_{\max} 234 (€ 33100); 243 (sh) (€ 30400); 254 (€ 30500); 282 (€ 6430); 290 (€ 6380) (EtOH) (Derep). λ_{\max} 234 (€ 30700); 254 (€ 27800); 282 (€ 5870); 290 (€ 5800) (MeOH) (Berdy). λ_{\max} 233; 254; 282; 290 (EtOH) (Berdy).

N-De-Ac, N-(3-mercaptopropanoyl): N²-Deacetyl-N²-(3-mercaptopropanoyl)maytansine. DM1
 [139504-50-0]

$\text{C}_{35}\text{H}_{48}\text{ClN}_3\text{O}_{10}\text{S}$ 738.297

Used in the form of its conjugated humanised C242 monoclonal antibody, Cantuzumab mertansine, SB 408075, huC242-DM1 for the treatment of colorectal cancer, pancreatic cancer and other solid tumour types that express the C242 antigen.

N-De-Ac, N-mercaptopropanoyl; conjugated humanised C242 monoclonal antibody: Cantuzumab mertansine, INN, USAN. SB 408075. huC242-DM1
 Used for the treatment of colorectal cancer, pancreatic cancer and other solid tumour types that express the C242 antigen. Tumour-activated immunotoxin. Consists of a colon cancer-specific humanised antibody, C242 conjugated to the maytansine deriv. DM1.

N-De-Ac, N-(2-methylpropanoyl): Maytanbutine. NSC 165014
 [38997-10-3]

$\text{C}_{36}\text{H}_{50}\text{ClN}_3\text{O}_{10}$ 720.258

From *Maytenus buchananii* and *Colubrina texensis* (Celastraceae, Rhamnaceae). Antileukaemic antibiotic. Mp 170-171°. $[\alpha]_D^{30}$ -122 (c, 0.0492 in CHCl₃). Log P 3.85 (calc). λ_{\max} 234 (€ 33100); 243 (sh) (€ 30400); 254 (€ 30500); 282 (€ 6430); 290 (€ 6380) (EtOH) (Derep). λ_{\max} 234 (€ 33100); 254 (€ 30800); 282 (€ 6430); 290 (€ 6380) (MeOH) (Berdy). λ_{\max} 233; 254; 282; 290 (EtOH) (Berdy).

N-De-Ac, N²-(3-methylpentanoyl): Maytanvaline. NSC 219970
 [52978-27-5]

$\text{C}_{38}\text{H}_{54}\text{ClN}_3\text{O}_{10}$ 748.311

Alkaloid from *Maytenus buchananii* (Celastraceae). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 175-176.5°. $[\alpha]_D^{26}$ -135 (c, 0.95 in CHCl₃). λ_{\max} 234 (€ 33100); 243 (sh) (€ 30400); 254 (€ 30500); 282 (€ 6430); 290 (€ 6380) (EtOH) (Derep). λ_{\max} 233 (€ 29100); 254 (€ 26800); 280 (€ 5300); 288 (€ 5360) (MeOH) (Berdy).

N-De-Ac, N²-(3-methyl-2E-pentanoyl): Isomallotusin
 [159993-33-6]

$\text{C}_{38}\text{H}_{52}\text{ClN}_3\text{O}_{10}$ 746.296

Prod. by *Mallotus anomalus*. Cytotoxic agent.

N-De-Ac, N²-(3-methyl-2Z-pentanoyl): Mallotusin
 [159934-12-0]

$\text{C}_{38}\text{H}_{52}\text{ClN}_3\text{O}_{10}$ 746.296

Prod. by *Mallotus anomalus*. Cytotoxic agent.

N²-De-Ac, N²-de-Me, N²-(2-methylpropanoyl): N²-Demethylmaytanbutine
 $\text{C}_{35}\text{H}_{48}\text{ClN}_3\text{O}_{10}$ 706.231

Isol. from *Maytenus buchananii*. Amorph. solid. Mp 156-158°.

N²-De-Ac, N²-mercaptopropanoyl; conjugated humanised N901 monoclonal antibody: huN901-DM1

Compd. comprised of a CD56-targeted humanised N901 antibody conjugated to the cytotoxic agent DM1 for the potential treatment of cancers that

express CD56.

N²²-De-Me: Normaytansine
 [75983-74-3]

$\text{C}_{33}\text{H}_{44}\text{ClN}_3\text{O}_{10}$ 678.177

Minor alkaloid from *Maytenus buchananii* (Celastraceae). Active against P388 lymphocytic leukaemia and KB cell cultures; activity resembles that of Maytansine. Cryst. (CH₂Cl₂/hexane). Mp 166-167°. Unstable in soln. λ_{\max} 240 (€ 34100); 250 (€ 25800); 281 (€ 4100); 288 (€ 3900) (EtOH) (Derep).

N,N²-Di-de-Me, N²-de-Ac, N²-(2,3-dimethylcyclopropylcarbonyl): Normaytancyprine
 [84123-43-3]

$\text{C}_{36}\text{H}_{48}\text{ClN}_3\text{O}_{10}$ 718.242

Alkaloid from *Putterlickia verrucosa* and *Maytenus buchananii*. Cytotoxic agent. Pale tan cryst. (CHCl₃/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 143-145°. λ_{\max} 232 (€ 48700); 252 (€ 28250); 280 (€ 7450); 289 (€ 7420) (EtOH).

Kupchan, S.M. *et al.*, *J.A.C.S.*, 1972, **94**, 1354 (*uv, ir, ms, nmr, cryst struct*)

Bryan, R.F. *et al.*, *J.C.S. Perkin 2*, 1973, 897 (*cryst struct, abs config*)

Wolpert-Defilippes, M.K. *et al.*, *Biochem. Pharmacol.*, 1975, **24**, 751 (*pharmacol*)

Brinkmeyer, R.S. *et al.*, *Diss. Abstr. Int.*, **B**, 1975, **36**, 1704; *CA*, **84**, 74150 (*synth*)

Meyers, A.I. *et al.*, *Tet. Lett.*, 1975, 1745; 1749 (*synth, ir, uv, nmr*)

Götschi, E. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1416 (*synth*)

Kupchan, S.M. *et al.*, *J. Med. Chem.*, 1977, **21**, 31 (*derivis*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1977, **42**, 2349 (*isol, struct, uv, ir, ms, nmr, derivis*)

Jackson, D.V. *et al.*, *Appl. Methods Oncol.*, 1978, **1**, 277 (*rev, pharmacol*)

Snedden, A.T. *et al.*, *J. Nat. Prod.*, 1980, **43**, 637 (*Normaytansine*)

Corey, E.J. *et al.*, *J.A.C.S.*, 1980, **102**, 6613 (*synth*)

Snedden, A.T. *et al.*, *J. Nat. Prod.*, 1982, **45**, 624-628 (*Normaytancyprine*)

Wallace, W.A. *et al.*, *Org. Magn. Reson.*, 1982, **19**, 31 (*cmr*)

Reider, P.J. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 71 (*rev*)

Goodwin, T.E. *et al.*, *J. Chem. Educ.*, 1984, **61**, 511 (*synth*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 142 (*antitumour props*)

Chari, R.V.J. *et al.*, *Cancer Res.*, 1992, **52**, 127-131 (*DM1, immunconjugate, synth, pharmacol*)

Feng, S.C. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 743-746 (*Mallotusin, Isomallotusin*)

Liu, C. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1996, **93**, 8618-8623 (*huC242-DM1*)

Larson, G.M. *et al.*, *J. Nat. Prod.*, 1999, **62**, 361-363 (*N-Demethylmaytanbutine*)

Smith, S. *et al.*, *Curr. Opin. Mol. Ther.*, 2001, **3**, 198-203 (*cantuzumab mertansine*)

Tassone, P. *et al.*, *Cancer Res.*, 2004, **64**, 4629-4636 (*huN901-DM1, pharmacol*)

Cassady, J.M. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1-26 (*rev*)

Smith, S.V. *et al.*, *Curr. Opin. Mol. Ther.*, 2004, **6**, 666-674 (*cantuzumab mertansine, rev*)

Xie, H. *et al.*, *J. Pharmacol. Exp. Ther.*, 2004, **308**, 1073-1082 (*cantuzumab mertansine, metab, pharmacokinetic*)

Smith, S.V. *et al.*, *Curr. Opin. Mol. Ther.*, 2005, **7**, 394-401 (*huN901-DM1, rev*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MBU820

Maytansinol**M-136**

3-De[2-(acetylmethylamino)-1-oxopropoxy]-3-hydroxymaytansine, 9CI. Ansamitocin P₀. C 15003P0. Antibiotic C 15003P0

[57103-68-1]

As Maytansine, M-135 with

R = H

C₂₈H₃₇ClN₂O₈ 565.062

Constit. of *Putterlickia verrucosa* (Celastraceae). Shows pronounced antileukaemic props. No antiprotozoal activity. Mp 173-174.5°. [α]_D²³ -309 (c, 0.11 in CHCl₃).

3-Ac: **Maytanacine**. Ansamitocin P1. C 15003P1. Antibiotic C 15003P1. 3-O-Acetylmaytansinol

[57103-69-2]

C₃₀H₃₉ClN₂O₉ 607.099

Constit. of *Putterlickia verrucosa* (Celastraceae) and *Nocardia* spp. Active against fungi, tumours, dermatophytes and protozoa. Cryst. (EtOAc). Mp 234-237°. [α]_D -121 (c, 0.25 in CHCl₃). λ_{max} 233 (ε 30300); 242 (sh) (ε 28000); 252 (ε 27900); 281 (ε 5360); 289 (ε 5360) (EtOH) (Derep).

3-Propanoyl: **Ansamitocin P2**. 2'-De(acetylmethylamino)maytansine, 9CI. C 15003P2. Antibiotic C 15003P2

[57103-70-5]

C₃₁H₄₁ClN₂O₉ 621.126

From *Nocardia* sp. and *Actinosynnema pretiosum*. Active against fungi and tumours. Cryst. (EtOAc). Sol. MeOH, DMSO, EtOAc, THF, CHCl₃; fairly sol. C₆H₆, Et₂O; poorly sol. H₂O, hexane. Mp 188-190° dec. [α]_D -127 (c, 0.35 in CHCl₃). λ_{max} 233 (ε 30300); 242 (sh) (ε 28000); 252 (ε 27900); 281 (ε 5360); 289 (ε 5360) (EtOH) (Derep). λ_{max} 233 (ε 30240); 252 (ε 27650); 280 (ε 5740); 288 (ε 5710) (MeOH) (Berdy).

3-Butanoyl: **Ansamitocin P3'**. C 15003P3'. Antibiotic C 15003P3' [66547-09-9]

C₃₂H₄₃ClN₂O₉ 635.152

From *Nocardia* sp. and *Actinosynnema pretiosum*. Active against fungi and tumours. Potent antiprotozoal agent. Cryst. (EtOAc). Sol. MeOH, DMSO, EtOAc, CHCl₃, THF; fairly sol. C₆H₆, Et₂O; poorly sol. H₂O, hexane. Mp 182-185° dec. [α]_D -134 (c, 0.11 in CHCl₃). λ_{max} 233 (ε 30300); 240 (sh) (ε 28500); 252 (ε 27600); 280 (ε 5750); 288 (ε 5700) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 0.313 mg/kg. OQ2293000

3-O-(2-Methylpropanoyl): **Ansamitocin P3**. C 15003P3. Antibiotic C 15003P3 [66584-72-3]

C₃₂H₄₃ClN₂O₉ 635.152

From *Actinosynnema pretiosum* and *Nocardia* sp. Active against fungi and tumours. Potent antiprotozoal agent.

Antimitotic agent, tubulin polymerisation inhibitor. Cryst. (EtOAc). Sol. MeOH, EtOAc, THF, DMSO, CHCl₃; fairly sol. C₆H₆, Et₂O; poorly sol. H₂O, hexane. Mp 190-192° dec. [α]_D -136 (c, 0.375 in CHCl₃). λ_{max} 233 (ε 30300); 240 (sh) (ε 28500); 252 (ε 27600); 280 (ε 5750); 288 (ε 5700) (MeOH) (Derep).

▶ Potent mutagen. LD₅₀ (mus, ipr) 0.313 mg/kg. OQ2291000

3-O-(3-Methylbutanoyl): **Ansamitocin P4**. C 15003P4. Antibiotic C 15003P4 [66547-10-2]

C₃₃H₄₅ClN₂O₉ 649.179

From *Nocardia* sp. and *Actinosynnema pretiosum*. Active against fungi and tumours. Potent antiprotozoal agent. Cryst. (EtOAc). Sol. MeOH, DMSO, THF, EtOAc, CHCl₃; fairly sol. C₆H₆, Et₂O; poorly sol. H₂O, hexane. Mp 177-180° dec. [α]_D -142 (c, 0.52 in CHCl₃). λ_{max} 233 (ε 30300); 240 (sh) (ε 28500); 252 (ε 27600); 280 (ε 5750); 288 (ε 5700) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 0.313 mg/kg. OQ2295000

3-O-(3-Hydroxy-3-methylbutanoyl): 3-(3-Hydroxyisovaleroyl)maytansinol. C 15003 P 4 βHY. Antibiotic C 15003 P 4 βHY

[78619-41-7]

C₃₃H₄₅ClN₂O₁₀ 665.179

From *Nocardia* sp. and *Actinosynnema pretiosum*. Weakly active against *Tetrahymena pyriformis*. Needles (EtOAc). Mp 201-203° dec. λ_{max} 231 (ε 30100); 251 (ε 27500); 280 (ε 5650); 288 (ε 5630) (MeOH) (Berdy).

3-O-(3-Hydroxy-3-methylbutanoyl), N-de-Me: **N-Demethyl-3-(3-hydroxyisovaleroyl)maytansinol**. C 15003 PND 4 βHY. Antibiotic C 15003 PND 4 βHY

[78619-43-9]

C₃₂H₄₃ClN₂O₁₀ 651.152

From *Nocardia* sp. and *Actinosynnema pretiosum*. Weakly active against *Tetrahymena pyriformis*. Powder. λ_{max} 232 (ε 31500); 239 (ε 32000); 279 (ε 3800); 288 (ε 3700) (MeOH) (Berdy).

3-O-(4-Hydroxy-3-methylbutanoyl): 3-(4-Hydroxyisovaleroyl)maytansinol. C 15003 P 4 γHY. Antibiotic C 15003 P 4 γHY

[78709-93-0]

C₃₃H₄₅ClN₂O₁₀ 665.179

From *Nocardia* sp. and *Actinosynnema pretiosum*. Weakly active against *Tetrahymena pyriformis*. Mp 205-207° dec. λ_{max} 232 (ε 30000); 252 (ε 27300); 280 (ε 5630); 288 (ε 5610) (MeOH) (Berdy).

3-O-[2-(N-Methylacetamido)propanoyl]: see Maytansine, M-135

N-De-Me, 3-Ac: **N-Demethylansamitocin P1**. N-Demethylmaytanacine. Ansamitocin PND 1. C 15003 PND 1. Antibiotic C 15003 PND 1

[77353-67-4]

C₂₉H₃₇ClN₂O₉ 593.072

From *Nocardia* sp., *Actinosynnema pretiosum* and *Streptomyces* sp. Powder. Sol. MeOH, CHCl₃, THF,

DMSO, Py; poorly sol. H₂O, hexane. [α]_D -55.8 (EtOH). λ_{max} 232 (ε 31500); 239 (ε 32000); 279 (ε 3780); 288 (ε 3700) (MeOH).

▶ LD₅₀ (mus, ipr) 1-2 mg/kg.

N-De-Me, 3-propanoyl: **N-Demethylansamitocin P2**. Ansamitocin PND 2. C 15003 PND 2. Antibiotic C 15003 PND 2

[77353-68-5]

C₃₀H₃₉ClN₂O₉ 607.099

Prod. by *Nocardia* sp. and *Streptomyces* sp. Powder. [α]_D -56.3 (EtOH). λ_{max} 232 (ε 31000); 239 (ε 32000); 279 (ε 3800); 288 (ε 3760) (MeOH).

N-De-Me, 3-O-(2-methylpropanoyl): **N-Demethylansamitocin P3**. C 15003 PND 3. Antibiotic C 15003 PND 3. Ansamitocin PND 3

[77353-69-6]

C₃₁H₄₁ClN₂O₉ 621.126

From *Nocardia* sp., *Actinosynnema pretiosum* and *Streptomyces* sp. Powder. Sol. MeOH, CHCl₃, Py, THF, DMSO; poorly sol. H₂O, hexane. [α]_D -57.1 (EtOH). λ_{max} 239 (ε 33000); 279 (ε 3880); 288 (ε 3790) (MeOH).

▶ LD₅₀ (mus, ipr) 1-2 mg/kg.

N-De-Me, 3-O-(3-methylbutanoyl): **N-Demethylansamitocin P4**. C 15003 PND 4. Antibiotic C 15003 PND 4. Ansamitocin PND 4

[77353-70-9]

From *Nocardia* sp., *Actinosynnema pretiosum* and *Streptomyces* spp. Powder. Sol. MeOH, THF, DMSO, Py, CHCl₃; poorly sol. H₂O, hexane. [α]_D -56.6 (EtOH). λ_{max} 232 (ε 31500); 239 (ε 31100); 279 (ε 3690); 288 (ε 3760) (MeOH).

▶ LD₅₀ (mus, ipr) 1-2 mg/kg.

N²²-De-Me, 3-Ac, N²²-β-D-glucopyranosyl: **Ansamitocinoside P1**

[950187-24-3]

C₃₅H₄₇ClN₂O₁₄ 755.214

Prod. by *Actinosynnema pretiosum*. Pale yellow solid. [α]_D¹⁹ -28 (c, 6.5 in MeOH). Possible struct. revision required (cf. Ansamitocinoside P2 (2007). λ_{max} 203 (log ε 4.57); 230 (log ε 4.33); 253 (log ε 4.25); 282 (log ε 3.72) (MeOH).

N²²-De-Me, 3-O-propanoyl, N²²-(4-O-carbamoyl-β-D-glucopyranosyl): **Ansamitocinoside P2**

C₃₇H₅₀ClN₃O₁₅ 812.266

Prod. by *Actinosynnema pretiosum*. Indicated struct. shown to be the most probable in 2007.

4,5-Deepoxy, 4,5-didehydro, N-de-Me: **N-Demethyl-4,5-deepoxymaytansinol**. C 15003 QNDO. Antibiotic C 15003 QNDO

[78619-44-0]


C₂₇H₃₅ClN₂O₇ 535.035

From *Nocardia* sp. and *Actinosynnema pretiosum*. Weakly active against *Tetrahymena pyriformis*. Powder. λ_{max} 231 (ε 30300); 240 (ε 30000); 251 (ε 25800); 279 (ε 3600); 288 (ε 3560) (MeOH) (Berdy).

- 4,5-Deepoxy, 4E,5-didehydro, N-de-Me, 3-O-(2-methylpropanoyl): **Antibiotic AI-R2397**. AI-R2397 [115753-88-3] C₃₁H₄₁ClN₂O₈ 605.126 Prod. by a *Thermomonospora* sp. Sol. MeOH, DMSO; fairly sol. Me₂CO, CHCl₃, EtOAc; poorly sol. H₂O, hexane. λ_{max} 221 (ε 37730); 283 (ε 2400) (MeOH) (Berdy).
- 15R-Hydroxy: **Ansamitocin PHO 0**. 15-Hydroxyansamitocin P₀. C 15003 PHO 0. Antibiotic C 15003 PHO 0 [79082-29-4] [78709-92-9] C₂₈H₃₇ClN₂O₉ 581.061 Prod. by *Nocardia* sp., *Chainia* sp. and *Streptomyces sclerotialus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D -178 (EtOH). λ_{max} 233 (ε 29600); 252 (ε 25700); 289 (ε 4990) (MeOH).
- 15R-Hydroxy, 3-Ac: **Ansamitocin PHO 1**. 15-Hydroxyansamitocin P₁. C 15003 PHO 1. Antibiotic C 15003 PHO 1 [79056-16-9] C₃₀H₃₉ClN₂O₁₀ 623.098 Prod. by *Nocardia* sp., *Chainia* sp. and *Streptomyces sclerotialus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D -102 (EtOH). λ_{max} 233 (ε 27000); 252 (ε 23500); 281 (ε 4560); 289 (ε 4500) (MeOH).
- 15R-Hydroxy, 3-propanoyl: **Ansamitocin PHO 2**. 15-Hydroxyansamitocin P₂. C 15003 PHO 2. Antibiotic 15003 PHO 2 [79028-00-5] [62414-96-4] C₃₁H₄₁ClN₂O₁₀ 637.125 Prod. by *Nocardia* sp., *Chainia nigra*, *Streptosporangium roseum* and *Streptomyces sclerotialus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D -107 (EtOH). λ_{max} 233 (ε 26800); 252 (ε 23300); 281 (ε 4550); 289 (ε 4550) (MeOH).
- 15R-Hydroxy, 3-O-(2-methylpropanoyl): **Ansamitocin PHO 3**. 15-Hydroxyansamitocin P₃. C 15003 PHO 3. Antibiotic 15003 PHO 3 [79082-28-3] [80953-27-1] C₃₂H₄₃ClN₂O₁₀ 651.152 Prod. by *Nocardia* sp., *Actinosynnema pretiosum*, *Chainia* sp. and *Streptomyces sclerotialus*. Cryst. Sol. Py, DMSO, CHCl₃, Me₂CO, EtOAc, MeOH, THF, DMF; poorly sol. hexane, H₂O. [α]_D -96 (EtOH). λ_{max} 233 (ε 26600); 252 (ε 23100); 281 (ε 4520); 289 (ε 4520) (EtOH).
- 15R-Hydroxy, 3-O-(3-methylbutanoyl): **Ansamitocin PHO 4**. 15-Hydroxyansamitocin P₄. C 15003 PHO 4. Antibiotic C 15003 PHO 4 [79056-15-8] C₃₃H₄₅ClN₂O₁₀ 665.179 Prod. by *Nocardia* sp., *Streptosporangium roseum* and *Streptomyces sclerotialus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D -108 (EtOH). λ_{max} 233 (ε 26300); 252 (ε 22900); 281 (ε 4500); 289 (ε 4500) (MeOH).
- (MeOH).
- 15S-Hydroxy: **15-Epiansamitocin PHO 0** C₂₈H₃₇ClN₂O₉ 581.061 Prod. by *Streptomyces sclerotialus*.
- 15S-Hydroxy, 3-Ac: **15-Epiansamitocin PHO 1** [79027-99-9] C₃₀H₃₉ClN₂O₁₀ 623.098 Prod. by *Streptomyces sclerotialus*.
- 15S-Hydroxy, 3-O-(2-methylpropanoyl): **15-Epiansamitocin PHO 3** [79082-27-2] C₃₂H₄₃ClN₂O₁₀ 651.152 Prod. by *Nocardia* sp. and *Streptomyces sclerotialus*. Cryst. Sol. DMF, CHCl₃, MeOH, DMSO, Py; poorly sol. hexane, H₂O. λ_{max} 233 (ε 28100); 253 (ε 25400); 281 (ε 5220); 289 (ε 5220) (MeOH).
- 15S-Hydroxy, 3-O-(3-methylbutanoyl): **15-Epiansamitocin PHO 4** C₃₃H₄₅ClN₂O₁₀ 665.179 Prod. by *Streptomyces sclerotialus*.
- 30-Hydroxy, 3-Ac: **30-Hydroxyansamitocin P1**. C 15003 PHM1. Antibiotic C 15003 PHM1 [78619-40-6] C₃₀H₃₉ClN₂O₁₀ 623.098 Prod. by *Nocardia* sp., and *Actinosynnema pretiosum*. Active against *Tetrahymena pyriformis* and tumours. Cryst. Sol. Py, THF, DMF, MeOH; poorly sol. hexane, H₂O. λ_{max} 232 (ε 25800); 249 (ε 23900); 280 (ε 4210); 288 (ε 4250) (MeOH) (Berdy).
- 30-Hydroxy, 3-propanoyl: **30-Hydroxyansamitocin P2**. C 15003 PHM2. Antibiotic C 15003 PHM2 [78619-39-3] C₃₁H₄₁ClN₂O₁₀ 637.125 Prod. by *Nocardia* sp., *Streptosporangium roseum* and *Actinosynnema pretiosum*. Active against *Tetrahymena pyriformis* and tumours. Cryst. Sol. Py, THF, CHCl₃, DMF, DMSO; poorly sol. hexane, H₂O. λ_{max} 232 (ε 24500); 249 (ε 23600); 280 (ε 4200); 288 (ε 4240) (MeOH) (Berdy).
- 30-Hydroxy, 3-O-(2-methylpropanoyl): **30-Hydroxyansamitocin P3**. C 15003 PHM3. Antibiotic C 15003 PHM3 [78619-38-2] C₃₂H₄₃ClN₂O₁₀ 651.152 From *Nocardia* sp. Active against *Tetrahymena pyriformis* and tumours. Cryst. (EtOH). Sol. Py, MeOH, CHCl₃, THF, DMF, DMSO; poorly sol. hexane, H₂O. Mp 192-194° dec. [α]_D²⁴ -148 (c, 0.5 in EtOH). λ_{max} 232 (ε 25600); 249 (ε 23700); 280 (ε 4190); 288 (ε 4250) (MeOH) (Berdy).
- 30-Hydroxy, 3-O-(3-methylbutanoyl): **30-Hydroxyansamitocin P4**. C 15003 PHM4. Antibiotic C 15003 PHM4 [78630-36-1] C₃₃H₄₅ClN₂O₁₀ 665.179 Prod. by *Nocardia* sp., *Streptosporangium roseum* and *Actinosynnema pretiosum*. Active against *Tetrahymena pyriformis* and tumours. Cryst. Sol. Py, DMF, CHCl₃, MeOH, DMSO; poorly sol. hexane, H₂O. Mp 190-192°. [α]_D²⁴ -
- 149 (c, 0.23 in EtOH). λ_{max} 232 (ε 25400); 249 (ε 23600); 280 (ε 4160); 288 (ε 4220) (MeOH) (Berdy).
- 15ξ-Acetoxy, 3-O-(2-methylpropanoyl): **Maytanbutacine** [62414-95-3] C₃₄H₄₅ClN₂O₁₁ 693.189 Alkaloid from *Maytenus serrata* (Celastraceae). Cryst. (CH₂Cl₂/Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 253-255°. [α]_D³³ -90 (c, 0.055 in EtOH). Stereochem. of epoxide undetermined. λ_{max} 233 (ε 29800); 243 (sh) (ε 27100); 254 (ε 27200); 282 (ε 5690); 290 (ε 5520) (EtOH) (Derep). λ_{max} 233 (ε 27200); 253 (ε 24200); 282 (ε 5050); 290 (ε 5080) (EtOH) (Berdy).
- 15-Methoxy, 3-O-(2-methylpropanoyl): **15-Methoxyansamitocin P3** [117860-08-9] C₃₃H₄₅ClN₂O₁₀ 665.179 Isol. from the Japanese mosses *Isotheicum subdiversiforme* (Lembophyllaceae) and *Thamnobryum sandei* (Neckeraceae). Cytotoxic. Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O.
- Dechloro, 4,5-deepoxy, 4,5-didehydro: **Dechloro-N-deepoxymaytansinol**. C 15003 deClQO. Antibiotic C 15003 deClQO [78630-38-3] C₂₈H₃₈N₂O₇ 514.617 From *Nocardia* sp. Weakly active against *Tetrahymena pyriformis*. Powder.
- Dechloro, 4,5-deepoxy, 4,5-didehydro, N-de-Me: **Dechloro-4,5-deepoxy-N-de-methylmaytansinol**. C 15003 deClQNDQ. Antibiotic C 15003 deClQNDQ [78630-37-2] C₂₇H₃₆N₂O₇ 500.591 From *Nocardia* sp. and *Actinosynnema pretiosum*. Weakly active against *Tetrahymena pyriformis*. Powder. λ_{max} 218 (ε 39800); 243 (ε 35400); 280 (ε 2800); 288 (ε 2500) (MeOH) (Berdy).
- Kupchan, S.M. et al., *J.O.C.*, 1977, **42**, 2349-2357 (Maytansine, Maytanbutacine, isol. struct, uv, ir, ms, nmr, synth)
- Higashide, E. et al., *Nature (London)*, 1977, **270**, 721
- Asai, M. et al., *Tetrahedron*, 1979, **35**, 1079 (Ansamitocins)
- Tanida, S. et al., *J. Antibiot.*, 1980, **33**, 192; 1981, **34**, 489-495; 496-506 (derivs)
- Japan. Pat., 1981, 45 484; *CA*, **96**, 67249p (Ansamitocin PHO)
- Wallace, W.A. et al., *Org. Magn. Reson.*, 1982, **19**, 31 (cmr)
- Hatano, K. et al., *Agric. Biol. Chem.*, 1984, **48**, 1721 (Ansamitocins)
- Reider, P.J. et al., *Alkaloids (Academic Press)*, 1984, **23**, 71 (rev)
- Isobe, M. et al., *J.A.C.S.*, 1984, **106**, 3252 (synth)
- Suffness, M. et al., *Alkaloids (Academic Press)*, 1985, **25**, 145 (pharmacol)
- Sakai, K. et al., *J. Nat. Prod.*, 1988, **51**, 845 (15-Methoxyansamitocin P3)
- Rengaraju, S. et al., *CA*, 1989, **110**, 209019 (AI-R2397)
- Benechie, M. et al., *J.O.C.*, 1996, **61**, 7133 (synth)

- Cassady, J.M. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1-26 (rev)
 Lu, C. *et al.*, *J. Antibiot.*, 2004, **57**, 348-350
 (*Ansamitocinoside P2*)
 Widdison, W.C. *et al.*, *J. Med. Chem.*, 2006, **49**, 4392-4408 (*synth*)
 Ma, J. *et al.*, *Arch. Pharmacol. Res.*, 2007, **30**, 670-673 (*Ansamitocinosides P1,P2*)
 Snipes, C.E. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1578-1581 (*Ansamitocinoside P2*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, APE529

Mazethramycin A **M-137**
 [68373-96-6]



$C_{17}H_{19}N_3O_4$ 329.355

Anthramycin-type antibiotic. Isol. from *Streptomyces thioluteus*. Active against bacteria and tumours. Light-yellow amorph. powder (Me₂CO aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 181-193° dec. $[\alpha]_D^{21} +730$ (c, 0.062 in DMF). λ_{max} 235 (ε 18200); 333 (ε 31800) (MeCN) (Derep). λ_{max} 220 (ε 34000); 235 (sh) (ε 30000); 335 (ε 39400) (MeOH) (Derep). λ_{max} 235; 335 (ε 39400) (MeCN) (Berdy). ▶ LD₅₀ (mus, ivn) .8 mg/kg. UC6475850

Me ether: Mazethramycin B

[68373-95-5]
 $C_{18}H_{21}N_3O_4$ 343.382

From *Streptomyces thioluteus*. Active against gram-positive and -negative bacteria. Yellow needles (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 245-279° dec. $[\alpha]_D^{23} +900$ (c, 0.2 in DMF). λ_{max} 235 (ε 18200); 333 (ε 31800) (MeCN) (Derep). λ_{max} 220 (ε 34000); 235 (sh) (ε 30000); 335 (ε 39400) (MeOH) (Derep). λ_{max} 215 (ε 25600); 235 (ε 22200); 334 (ε 46100) (MeCN) (Berdy). λ_{max} 258 (ε 17200); 351 (ε 43400) (NaOH) (Berdy).

▶ LD₅₀ (mus, ivn) .8 mg/kg. UC6475920

Et ether: Mazethramycin C

[68373-94-4]
 $C_{19}H_{23}N_3O_4$ 357.408

From *Streptomyces thioluteus*. Active against gram-positive and -negative bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 216-223° dec. $[\alpha]_D^{21} +450$ (c, 0.067 in DMF). λ_{max} 235 (ε 18200); 333 (ε 31800) (MeCN) (Derep). λ_{max} 220 (ε 34000); 235 (sh) (ε 30000); 335 (ε 39400) (MeOH) (Derep). λ_{max} 217 (ε 25700); 235 (ε 19300); 333 (ε 43600) (MeCN) (Berdy).


11-Deoxy, 10,11-didehydro: Anhydromazethramycin

[68373-93-3]
 $C_{17}H_{17}N_3O_3$ 311.34

Semisynthetic. Active against gram-positive and -negative bacteria. Light-yellow cryst. powder. Mp 252-262° dec. $[\alpha]_D^{22} +1940$ (c, 0.05 in DMF). Log P -0.81 (uncertain value) (calc).

Kunimoto, S. *et al.*, *J. Antibiot.*, 1980, **33**, 665

Mearsamine **M-138**
 [960511-41-5]




$C_{22}H_{37}N_2O_3$ 377.546

Alkaloid from the leaves of *Peripentadenia mearsii*. Yellow gum (as trifluoroacetate). $[\alpha]_D^{23} -1.5$ (c, 0.07 in MeOH) (trifluoroacetate). λ_{max} 202 (log ε 3.14); 265 (log ε 3.38) (MeOH) (trifluoroacetate).

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1946-1950 (*isol, pmr, cmr, ms*)

Mearsine **M-139**
 3,8-Dimethyl-2-azabicyclo[2.2.2]oct-2-en-5-one, 9CI



(-)-form

$C_9H_{13}NO$ 151.208

Unique isoquinuclidine alkaloid.

(+)-form [127911-13-1]
 Synthetic. Mp 41.5-42°. $[\alpha]_D^{20} +32.1$ (c, 0.5 in CH₂Cl₂).

Picrate:

Yellow prisms (MeOH). Mp 209-210° dec.

(-)-form [92446-42-9]
 Minor alkaloid from *Peripentadenia mearsii* (Elaeocarpaceae). Mp 43-44°. $[\alpha]_D^{17} -34.5$ (c, 0.495 in CH₂Cl₂).


Picrate: Mp 212-213°.

Robertson, G.B. *et al.*, *Tet. Lett.*, 1984, **25**,

2695 (*isol, uv, ir, pmr, cryst struct*)

Crouse, J.R. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1227 (*synth, uv, ir, pmr, cmr, ms*)

Mecambridine **M-140**
Oreophiline



(S)-form

$C_{22}H_{25}NO_6$ 399.443

(S)-form [31098-60-9]
 Alkaloid from *Meconopsis cambrica* and some *Papaver* spp. (Papaveraceae). Mp 179-180°. $[\alpha]_D^{24} -243$ (c, 0.22 in CHCl₃).

Hydrochloride: Mp 224-225° dec.

Ac: Mp 119-120°.

N-Me: Mecambridine methosalt

$C_{23}H_{28}NO_6$ 414.477

Alkaloid from *Papaver pseudocane-cens* (Papaveraceae). Cryst. (MeOH/Et₂O) (as iodide). Mp 203-206° (iodide). $[\alpha]_D^{23} -169$ (c, 0.1 in MeOH).

O¹¹-De-Me: Aryapavine. 11-Demethyl-mecambrine

[57129-99-4]

$C_{21}H_{23}NO_6$ 385.416

Alkaloid from the dry latex of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (Me₂CO). Mp 156-157°. No opt. rotn. reported, abs. config. not confirmed.

(±)-form [27293-09-0]
 Synthetic. Mp 158-160°.

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 1720 (*isol, uv, ir*)

Pfeifer, S. *et al.*, *Tet. Lett.*, 1967, 83 (*uv, ir, ms, pmr, struct*)

Preininger, V. *et al.*, *Tet. Lett.*, 1969, 2109 (*struct*)

Novák, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 883 (*deriv*)


Kametani, T. *et al.*, *Heterocycles*, 1974, **2**, 55 (*Aryapavine, synth, pmr*)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1974, 1954 (*synth, pmr, ms*)

Shafiee, A. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 1570 (*Aryapavine, isol, struct*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr, cryst struct*)

Mecambrine **M-141**
Fugapavine



Absolute configuration

$C_{18}H_{17}NO_3$ 295.337

(S)-form [1093-07-8]

Alkaloid from *Meconopsis cambrica*, *Papaver fugax*, *Papaver dubium* and *Papaver caucasicum* (Papaveraceae). Increases gastrointestinal motility in exptl. animals. Also shows antihistaminic and hypertensive props.; respiratory stimulant. Mp 178.5-179.5°. $[\alpha]_D -116$ (-94) (CHCl₃).

▶ Toxic, LD₅₀ 4.1 mg/kg (mus). OQ9950000

Hydrochloride: Mp 269-270°.

(±)-form [41590-24-3]
 Synthetic. Mp 197-198°.

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1965, **30**, 914 (*isol, struct*)

Snatzke, G. *et al.*, *J.C.S.(C)*, 1966, 1681 (*config*)

Barton, D.H.R. *et al.*, *J.C.S.(C)*, 1967, 2134 (*biosynth*)


Kametani, T. *et al.*, *Tetrahedron*, 1971, **27**, 5375 (*synth, uv, ir, pmr, ms*)

Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 571 (*ms*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr, cryst struct*)

Sari, A. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 265-268 (*isol, cmr*)


Mechercharstatin A M-142
Mechercharmycin A. IB 01211
[822520-96-7]



$C_{35}H_{32}N_8O_7S$ 708.753
Prod. by the marine-derived *Thermoactinomyces* sp. YM3-251. Cytotoxic. Powder. $[\alpha]_D^{25} +110$ (c, 0.04 in DMSO). λ_{max} 223 (log ϵ 4.71); 260 (log ϵ 4.73); 300 (sh) (MeOH).

Kanoh, K. *et al.*, *J. Antibiot.*, 2005, **58**, 289-292 (*isol, pmr, cmr, cryst struct*)
Hernández, D. *et al.*, *Org. Lett.*, 2007, **9**, 809-811 (*synth*)


Mechercharstatin B M-143
Mechercharmycin B
[861231-28-9]



$C_{35}H_{36}N_8O_{10}$ 728.717
Prod. by the marine-derived *Thermoactinomyces* sp. YM3-251. Cytotoxic. Powder. $[\alpha]_D^{25} +56$ (c, 0.06 in DMSO). λ_{max} 220 (log ϵ 4.65); 261 (log ϵ 4.62); 310 (sh) (MeOH).

Kanoh, K. *et al.*, *J. Antibiot.*, 2005, **58**, 289-292 (*isol, pmr, cmr*)

Meconoquintupline M-144
Mecoquitupline
[623945-12-0]




Relative Configuration

$C_{19}H_{21}NO_4$ 327.379
Alkaloid from the whole plant of *Mecopopsis quintuplinervia*. Mp 192-194°. $[\alpha]_D^{22} -52.8$ (c, 0.11 in $CHCl_3$).

Shang, X.Y. *et al.*, *Chin. Chem. Lett.*, 2003, **14**, 597-598 (*isol, pmr, cmr*)


Medelline M-145
[102505-09-9]



$C_{37}H_{38}N_2O_6$ 606.717
Minor alkaloid from the bark of *Pseudoxandra* aff. *lucida* (Annonaceae). Amorph. $[\alpha]_D -38$ (c, 0.16 in MeOH). The first known example of a bisbenzylisoquinoline alkaloid with an intramolecular methyleneoxy bridge.

Ac: $[\alpha]_D -42$ (c, 0.60 in $CHCl_3$).
Cortes, D. *et al.*, *Heterocycles*, 1986, **24**, 607 (*uv, ir, pmr, cmr, ms, cd, struct*)

Megalanthone M-146
Platynecine 9-iridiflorate
[215930-93-1]




$C_{15}H_{27}NO_5$ 301.382
Alkaloid from *Heliotropium megalanthum*. Insecticide. Oil. $[\alpha]_D -5$ (c, 0.04 in EtOH).

5'-Hydroxy: **Leptanthine**[†]
 $C_{15}H_{27}NO_6$ 317.381
Alkaloid from *Onosma leptantha*. Light brown oil. $[\alpha]_D^{25} -2$ (c, 0.2 in MeOH).

5'-Hydroxy, N-oxide: **Leptanthine N-oxide**
 $C_{15}H_{27}NO_7$ 333.381
Alkaloid from *Onosma leptantha*. Amorph. powder. $[\alpha]_D^{25} +16.5$ (c, 0.2 in MeOH).
Reina, M. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1418-1420 (*Megalanthone*)
Kretsi, O. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 3136-3140 (*Leptanthine, Leptanthine N-oxide*)


Megastachine M-147
[72049-07-1]



Absolute configuration

$C_{20}H_{29}NO_3$ 331.454
Alkaloid from *Lycopodium megastachyum* (Lycopodiaceae). Mp 167-169°. Ac: Mp 140-142°.
Braekman, J.-C. *et al.*, *Can. J. Chem.*, 1979, **57**, 1691 (*isol, ir, ms, pmr, cryst struct*)

Megistophylline I M-148
[261915-32-6]



$C_{21}H_{23}NO_5$ 369.416
Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. solid. $[\alpha]_D^{25} +32.9$ (c, 0.26 in $CHCl_3$). λ_{max} 301 (log ϵ 3.95); 342 (log ϵ 3.88); 407 (sh) (MeOH).

Papageorgiou, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 385-386


Megistophylline II M-149

[261915-33-7]
As Megistophylline I, M-148 with
R = $-CH_2CH=C(CH_3)CH_2CH_2CH=C(CH_3)_2$

$C_{26}H_{31}NO_5$ 437.535
Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. solid. $[\alpha]_D^{25} +11.3$ (c, 0.1 in $CHCl_3$). λ_{max} 301 (log ϵ 4.36); 338 (log ϵ 3.88); 404 (sh) (MeOH).

Papageorgiou, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 385-386


Megistoquinone I M-150
[449732-09-6]



$C_{13}H_9NO_5$ 259.218
Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow solid. λ_{max} 258 (log ϵ 4.23); 291 (sh); 371 (log ϵ 3.37) (MeOH).

Fokialakis, N. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 413-414 (*isol, uv, pmr, cmr, ms*)

Megistoquinone II M-151
[449732-11-0]



$C_{14}H_{13}NO_6$ 291.26
Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. purple solid.


λ_{\max} 278 (log ϵ 3.54); 315 (sh); 515 (log ϵ 2.33) (MeOH).

Fokialakis, N. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 413-414 (*isol, uv, pmr, cmr, ms*)

Megistosarconine

M-152

[259822-41-8]



Relative Configuration

$C_{18}H_{19}NO_5$ 329.352

Alkaloid from *Sarcomelicope megistophylla*. Amorph. powder. $[\alpha]_D^{25} +162$ (c, 0.2 in $CHCl_3$). λ_{\max} 255 (log ϵ 4.26); 278 (sh) (MeOH).

8-Imide: Megistosarcimine

[259822-39-4]

$C_{18}H_{20}N_2O_4$ 328.367


Alkaloid from *Sarcomelicope megistophylla*. Amorph. powder. $[\alpha]_D^{25} +131$ (c, 0.2 in $CHCl_3$). V. unstable, readily converts to Megistosarconine. λ_{\max} 252 (log ϵ 4.25); 275 (sh) (MeOH).

Fokialakis, N. *et al.*, *Phytochemistry*, 1999, **52**, 1745-1748

Mehranine

M-153

[88840-03-3]

**(+)-form**

$C_{20}H_{26}N_2O$ 310.438

(+)-form

Alkaloid from the leaves of *Ervatamia coronaria* (Apocynaceae). Amorph. $[\alpha]_D^{25} +36$ ($CHCl_3$).

(-)-form [168396-00-7]

Alkaloid from leaves of *Tabernaemontana divaricata* (double flower variety) (Apocynaceae). Solid (Me_2CO). Mp 102-104°. $[\alpha]_D^{25} -49$ (c, 0.831 in $CHCl_3$). Tentative struct. assignment based on biogenetic grounds.

3-Oxo: 3-Oxomehranine

[219758-22-2]

$C_{20}H_{24}N_2O_2$ 324.422

Alkaloid from *Tabernaemontana bovina*. Oil. $[\alpha]_D^{25} -19.1$ (c, 0.4 in $CHCl_3$).

Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B.*, 1983, **38**, 1700-1701 (*isol, uv, ir, pmr, ms, struct*)

Kam, T.-S. *et al.*, *Phytochemistry*, 1995, **40**, 313-316 (*isol, uv, pmr, cmr, ms, struct*)


Lien, T.P. *et al.*, *Phytochemistry*, 1998, **49**, 1457-1461 (*3-Oxomehranine*)

Merzweiler, K. *et al.*, *J. Prakt. Chem.*, 1999, **341**, 69-71 (*cryst struct, abs config*)

Éles, J. *et al.*, *J.O.C.*, 2002, **67**, 7255-7260 (*synth*)

Melanocin A

M-154




$C_{18}H_{14}N_2O_5$ 338.319

Related to Xanthocillin X, X-7. Prod. by *Eupenicillium shearii* F80695. Inhibitor of mushroom tyrosinase and melanin biosynth. Antioxidant. Yellow powder. Mp 212°. λ_{\max} 220 (ϵ 32300); 336 (ϵ 13100) (MeOH).

Kim, J.-P. *et al.*, *J. Antibiot.*, 2003, **56**, 993-999; 1000-1003 (*isol, pmr, cmr, activity*)

Melanocin B

M-155




$C_{17}H_{15}NO_6$ 329.309

Prod. by *Eupenicillium shearii* F80695. Antioxidant. Dark brown powder. Mp 223°. λ_{\max} 253 (ϵ 31400); 341 (ϵ 8300) (MeOH).

Kim, J.-P. *et al.*, *J. Antibiot.*, 2003, **56**, 993-999; 1000-1003 (*isol, pmr, cmr, activity*)

Melanocin C

M-156



$C_{18}H_{14}N_2O_6$ 354.318


Prod. by *Eupenicillium shearii* F80695. Antioxidant. Dark brown powder. Mp 190-200° dec. λ_{\max} 241 (ϵ 42650); 341 (ϵ 5150) (MeOH).

Kim, J.-P. *et al.*, *J. Antibiot.*, 2003, **56**, 993-999; 1000-1003 (*isol, pmr, cmr, activity*)

Melanocrocine

M-157

[364356-73-0]



Absolute Configuration

$C_{29}H_{33}NO_4$ 459.584

Pigment from *Melanogaster broomeianus*. Amorph. orange solid. λ_{\max} 243 (log ϵ 3.88); 283 (log ϵ 3.89); 306 (log ϵ 3.91); 319 (log ϵ 3.9); 404 (log ϵ 4.31) (MeOH).


Aulinger, K. *et al.*, *Z. Naturforsch., C.*, 2001, **56**, 495-498

Melanostatin†

M-158

BYM 28566. Antibiotic BYM 28566

[134019-80-0]



$C_{19}H_{25}N_5O_5$ 403.437

Peptide antibiotic. Prod. by *Streptomyces clavifer*. Melanin synthesis inhibitor. Amorph. powder. Sol. H_2O , MeOH, EtOH; fairly sol. DMSO; poorly sol. Me_2CO , $CHCl_3$, EtOAc. $[\alpha]_D^{29} +50$ (c, 1 in H_2O). Dec. gradually above 155°. Similar to Feldamycin, F-33.

Oki, T. *et al.*, *J. Antibiot.*, 1991, **44**, 25; 76 (*isol, pmr, cmr, ir, props, synth, abs config*)


Melanoxidin

M-159

5-(4-Oxazolyl)-4-pentene-2,3-diol. 4-(3,4-Dihydroxy-1-pentenyl)oxazole. MR 93A. Antibiotic MR 93A. ATF 606.

Antibiotic ATF 606

[169397-87-9]



$C_8H_{11}NO_3$ 169.18

Most probable config. shown. Isol. from cultures of *Trichoderma harzianum* KCTC 0114BP and *Trichoderma* sp. ATF606. Melanin biosynthesis inhibitor. Needles (EtOAc/hexane). Mp 73-74°. $[\alpha]_D^{25} -6$ (c, 0.1 in MeOH). $[\alpha]_D^{24} -20$ (c, 0.1 in MeOH). λ_{\max} 227 (ϵ 19800) (MeOH) (Derep).

3'-Ketone: 4-Hydroxy-1-(4-oxazolyl)-1-penten-3-one. 4-(4-Hydroxy-3-oxo-1-pentenyl)oxazole

[197768-84-6]

$C_8H_9NO_3$ 167.164

Prod. by an unidentified fungus. Needles. Mp 52°. $[\alpha]_D^{25} -34$ (c, 1.2 in MeOH). λ_{\max} 279 (ϵ 14700) (MeOH).


Hashimoto, R. *et al.*, *J. Antibiot.*, 1995, **48**, 1052 (*Melanoxidin, isol, pmr, cmr*)

Lee, C. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1605 (*MR93A, isol, uv, cd, pmr, cmr, struct*)

Morimitsu, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 1428-1433 (*isol, uv, pmr, cmr, ms*)

Melanthioidine

M-160

**(R,R)-form**

C₃₈H₄₂N₂O₆ 622.76

Dimeric phenethylisoquinoline alkaloid.

(R,R)-form [4085-28-3]Alkaloid from *Androcymbium mel-anthioides* var. *stricta* (Liliaceae). Prisms (EtOAc or Me₂CO). Mp 142-144°. [α]_D²² -63 (c, 0.82 in CHCl₃).


Dibenzyl ether: Mp 154-155°.

(RS,RS)-form


Di-O-benzyl: Mp 192-194°.

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 255 (*isol, ir, uv*)Battersby, A.R. *et al.*, *J.C.S. (C)*, 1967, 1739 (*ms, pmr, ord, struct, synth*)Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1968, **16**, 663 (*synth, uv, ir, ms*)**Melaxillarinine****M-161**

[192331-99-0]

C₃₈H₄₈N₄O₂ 592.823Alkaloid from the roots of *Melodinus axillaris*. Pale red powder. [α]_D -36.2 (c, 0.15 in MeOH). λ_{\max} 210 (log ϵ 4.57); 230 (sh) (log ϵ 4.5); 290 (log ϵ 3.82) (no solvent reported).Yan, K.X. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 313-314**Meleagrinn****M-162**

[17151-77-4]



Absolute configuration

C₂₃H₂₃N₅O₄ 433.466Closely related to Neoxaline, N-149. Isol. from *Penicillium meleagrinnum* and a marine-derived *Penicillium chrysogenum*. Shows structural similarity to tremorgenic mycotoxins. Pale yellow leaves (CHCl₃ or CH₂Cl₂). Mp 250° dec. [α]_D -116 (c, 0.088 in CHCl₃). λ_{\max} 232 (ϵ 27500); 285 (sh) (ϵ 8320); 349 (ϵ 27500) (EtOH) (Derrep). λ_{\max} 230 (ϵ 24500); 345 (ϵ 26200) (MeOH) (Berdy).

O-Ac:

Pale yellow cryst. powder (C₆H₆/petrol).N¹⁴-Me:

Pale yellow needles (MeOH). Mp 212° dec.

Me ether: **Oxaline**


[55623-37-5]

C₂₄H₂₅N₅O₄ 447.493Isol. from cultures of the toxicogenic fungus *Penicillium oxalicum*, and from *Penicillium* sp. Fg-234. Also prod. by a marine-derived fungus isol. from *Gracilaria verrucosa*. Cryst. (Me₂CO or MeNO₂). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 220-221° Mp 230-232° dec. λ_{\max} 228 (ϵ 21300); 345 (ϵ 25200) (MeOH) (Derrep). λ_{\max} 228 (ϵ 20890); 347 (ϵ 24550) (MeOH) (Berdy).O,N¹⁴-Di-Me:


Needles (hexane). Mp 217-218° (214-216°).

Nagel, D.W. *et al.*, *Tetrahedron*, 1976, **32**, 2625 (*Oxaline, cryst struct*)Nozawa, K. *et al.*, *J. Nat. Prod.*, 1979, **42**, 374 (*isol*)Vlegaar, R. *et al.*, *Chem. Comm.*, 1980, 160; 1983, 560 (*cmr, biosynth*)Konda, Y. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2987 (*Oxaline*)Kawai, K. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 94 (*cryst struct, abs config*)Reshetilova, T.A. *et al.*, *Food Addit. Contam.*, 1995, **12**, 461-466 (*occur*)Li, Y. *et al.*, *Arch. Pharmacol. Res.*, 2003, **26**, 21-23 (*Oxaline, marine isol*)Bringmann, G. *et al.*, *Tetrahedron*, 2005, **61**, 7252-7265 (*marine isol*)Overy, D.P. *et al.*, *Biochem. Syst. Ecol.*, 2006, **34**, 345-348 (*isol, abs config*)**Melemeleone A****M-163**


[144587-55-3]

C₂₃H₃₃NO₅S 435.583Constit. of *Dysidea avara*. Red amorph. solid. Mp 110-115°. [α]_D²⁰ -20.1 (c, 0.006 in CH₂Cl₂). λ_{\max} 214 (ϵ); 290 (ϵ); 483 (ϵ) (MeOH) (Derrep).Alvi, K.A. *et al.*, *J.O.C.*, 1992, **57**, 6604 (*isol, pmr, cmr*)**Melemeleone B****M-164**



[144587-56-4]

C₂₃H₃₃NO₅S 435.583Constit. of *Dysidea avara*. Tyrosinekinase inhibitor. Red amorph. solid. Sol. MeOH. Mp 190-200°. [α]_D²⁰ -22 (c, 0.01 in CH₂Cl₂). λ_{\max} 214 (ϵ); 290 (ϵ); 483 (ϵ) (MeOH) (Derrep).Abvi, K.A. *et al.*, *J.O.C.*, 1992, **57**, 6604 (*isol, pmr, cmr*)**Melicarpine****M-165**

4,7,8,8-Tetramethoxy-5H-furo[2,3-b]quinolin-5(8H)-one [632335-15-0]

C₁₅H₁₅NO₆ 305.287Alkaloid from the leaves of *Melicope semecarpifolia*. Needles (MeOH). Mp 211-213°. λ_{\max} 207 (log ϵ 4.29); 242 (log ϵ 4.47); 287 (sh) (log ϵ 3.83) (EtOH).Chen, J.-J. *et al.*, *Planta Med.*, 2003, **69**, 542-546 (*isol, pmr, cmr, ms*)**Melicarpinone****M-166**


4-Methoxy-9-methyl-7H-furo[2,3-b]quinolin-7(9H)-one [479415-37-7]

C₁₃H₁₁NO₃ 229.235Alkaloid from the leaves of *Melicope semecarpifolia*. Pale yellow needles (CHCl₃/MeOH). Mp 145-147°. λ_{\max} 204 (log ϵ 4.32); 234 (log ϵ 4.42); 276 (log ϵ 4.48); 320 (log ϵ 3.86); 339 (log ϵ 3.61); 399 (log ϵ 4.11) (EtOH).Chen, J.-J. *et al.*, *Planta Med.*, 2002, **68**, 790-793 (*isol, pmr, cmr, ms*)**Melicobisquinolinone A****M-167**C₃₀H₃₀N₂O₄ 482.578Dimer of N-methylflindersine in F-90. Alkaloid from *Melicope ptelefolia*. Amorph. λ_{\max} 226 (log ϵ 4.92); 274 (log ϵ 4.14); 318 (log ϵ 4.13) (MeOH). λ_{\max} 226 (ϵ 83170); 274 (ϵ 15500); 328 (ϵ 13500) (MeOH) (Berdy).Kamperdick, C. *et al.*, *Phytochemistry*, 1999,

50, 177-181 (isol, uv, ir, pmr, cmr, ms)


Melicobisquinolinone B

M-168

C₂₇H₂₄N₂O₃ 424.498Alkaloid from *Melicope ptelefolia*.Amorph. Racemic. λ_{max} 229 (log ε 4.8); 322 (log ε 4.08) (MeOH). λ_{max} 229 (ε 63100); 322 (ε 12000) (MeOH) (Berdy).Kamperdick, C. et al., *Phytochemistry*, 1999, **50**, 177-181 (isol, uv, ir, pmr, cmr, ms)**Melicopine**

M-169

4,5-Dimethoxy-11-methyl-1,3-dioxolo[4,5-c]acridin-6(11H)-one, 9CI. 1,2-Dimethoxy-10-methyl-3,4-methylenedioxyacridone [568-01-4]

C₁₇H₁₅NO₅ 313.309Alkaloid from *Melicope fareana*, *Evodia alata*, *Acronychia baueri*, *Teclea natalensis* and *Bauerella simplicifolia* (preferred genus name *Acronychia*) (Rutaceae). Mp 181-182° (173-175°). λ_{max} 270 (ε 55000); 301 (ε 13500) (EtOH) (Derep).*O*¹-De-Me: 1-Hydroxy-2-methoxy-10-methyl-3,4-methylenedioxyacridone. **O-Normelicopine** [517-76-0]C₁₆H₁₃NO₅ 299.282Alkaloid from the bark of *Acronychia baueri* (Rutaceae). Orange needles. Mp 235°. Prob. an artifact.Price, J.R. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 249; *CA*, **46**, 4010d (isol)Crow, W.D. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 282; *CA*, **46**, 4014a (struct)Bowie, J. et al., *Aust. J. Chem.*, 1967, **20**, 1179 (ms)Prager, R.H. et al., *Aust. J. Chem.*, 1968, **21**, 229 (synth, pmr, Normelicopine)Prager, R.H. et al., *Aust. J. Chem.*, 1969, **22**, 2627 (biosynth)Pegel, K.H. et al., *J.C.S. (C)*, 1969, 2327 (isol, uv, ir, pmr)Mester, I. et al., *Z. Naturforsch., B*, 1979, **34**, 516 (cmr)Tillequin, F. et al., *J. Nat. Prod.*, 1980, **43**, 498 (isol, uv, ir, ms, pmr)**Melinonine I**

M-170

Struct. unknown. Alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Mp 160-170° dec. (as picrate). Mol. formula not recorded.Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167-1187 (isol, uv)**Melinonine H**

M-171

[1359-58-6]

C₂₀H₂₁N₂O⁺ 305.399Struct. unknown. Approx. MF given (may be H₂₃). Quaternary alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Mp 302-304° dec. (as perchlorate).Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167-1187 (isol, uv, ir)**Melinonine K**

M-172

Struct. unknown. Alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Mol. formula not recorded.*Chloride*: Mp 210-212° dec.*Picrate*: Mp 196-199° dec.Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167-1187 (isol, uv)**Melinonine L**

M-173


C₂₂H₂₆N₂O₄ 382.458Struct. unknown. Alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Mp 248-250°.*Methopicate*: Mp 156-160° Mp 241-243° (double Mp).*Ac*:Needles (CHCl₃/Et₂O). Mp 220-223°.Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167-1187 (isol, uv)**Melinonine M**

M-174

Struct. unknown. Alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Mp 242-245° dec. (as picrate). Mol. formula not recorded.Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167-1187 (isol, uv)**Melisemine**

M-175

1,2-Dihydro-4,6,7-trimethoxy-2-oxo-3-quinolinecarboxaldehyde. 3-Formyl-2-hydroxy-4,6,7-trimethoxyquinoline

C₁₃H₁₃NO₅ 263.249Alkaloid from the leaves of *Melicope semecarpifolia*. Pale yellow needles (CHCl₃/MeOH). Mp 189-191°. λ_{max} 219 (log ε 4.32); 235 (log ε 4.38); 260 (sh) (log ε 4.12); 316 (log ε 3.9); 387 (log ε 3.78) (EtOH).Chen, J.-J. et al., *Planta Med.*, 2002, **68**, 790-793 (isol, pmr, ms)**Melithiazole I**

M-176


[248938-52-5]

As Melithiazole B, M-177 with

R = -CH₂CH(CH₃)₂C₂₁H₂₈N₂O₄S₂ 436.595Prod. by *Myxococcus stipitatus*. Antifungal and cytotoxic. Oil. λ_{max} 223 ; 241 ; 312 (MeOH).Böhlendorf, B. et al., *Eur. J. Org. Chem.*, 1999, 2601-2608 (isol, struct)Sasse, F. et al., *J. Antibiot.*, 1999, **52**, 721-729 (isol, props)Takayama, H. et al., *Eur. J. Org. Chem.*, 2006, 644-649 (synth)**Melithiazole B**

M-177

[170894-31-2]



Absolute Configuration

C₂₀H₂₄N₂O₄S₂ 420.553Prod. by *Archangium gephyra* and *Melittangium lichenicola*. Antifungal and cytotoxic agent. Oil. λ_{max} 235 (log ε 4.38); 307 (log ε 3.83) (MeCN).*O*⁵-De-Me: **Cystothiazole D**

[214215-09-5]

C₁₉H₂₂N₂O₄S₂ 406.526Prod. by *Cystobacter fuscus*. Antifungal agent. Powder. [α]_D²⁵ +134 (c, 0.05 in CHCl₃). λ_{max} 235 (ε 37000); 306 (ε 11900) (MeOH).14,15-Epoxide: **Melithiazole N**

[248938-54-7]

C₂₀H₂₄N₂O₅S₂ 436.552Prod. by *Archangium gephyra*.11α,12-Dihydro: **Melithiazole A**

[170894-30-1]

C₂₀H₂₆N₂O₄S₂ 422.568Prod. by *Archangium gephyra* and *Melittangium lichenicola*. Antifungal and cytotoxic agent. Oil. [α]_D²² +0.3 (c, 1 in MeOH). λ_{max} 211 (sh) (log ε 4.33); 227 (log ε 4.47); 256 (sh) (log ε 4.28); 263 (sh) (log ε 4.11); 309 (sh) (log ε 3.27); 311 (sh) (log ε 3.27); 327 (sh) (log ε 3.07); 335 (sh) (log ε 2.83) (MeCN).11α,12-Dihydro, 14,15-epoxide: **Melithiazole K**

[248938-47-8]

C₂₀H₂₆N₂O₅S₂ 438.568Prod. by *Archangium gephyra*. Oil.λ_{max} 217 (log ε 4.07); 239 (log ε 4.12); 305 (sh) (log ε 3.16) (MeOH).14,15-Dihydro: **Cystothiazole A**. **Antibiotic YSI 40-2**. **Melithiazole E** [207399-36-8]C₂₀H₂₆N₂O₄S₂ 422.568Prod. by *Cystobacter fuscus* and *Myxococcus stipitatus*. Antifungal and cytotoxic agent. Needles. Mp 111-112°. [α]_D²⁵ +109 (c, 0.2 in CHCl₃). λ_{max} 223 (ε 38200); 242 (ε 34500); 310 (ε 12400) (MeOH).14,15-Dihydro, *O*⁵-de-Me: **Cystothiazole C**

[214215-07-3]

C₁₉H₂₄N₂O₄S₂ 408.542

Prod. by *Cystobacter fuscus*. Antifungal agent. Powder. $[\alpha]_D^{25} +145$ (c, 0.2 in CHCl_3). λ_{max} 222 (ϵ 33900); 242 (ϵ 30500); 312 (ϵ 10400) (MeOH).

14,15-Dihydro, 14-hydroxy: Cystothiazole B

[207399-38-0]
 $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_5\text{S}_2$ 438.568
 Prod. by *Cystobacter fuscus*. Powder. $[\alpha]_D +139$ (c, 0.09 in CHCl_3). λ_{max} 222 (ϵ 38400); 243 (ϵ 35700); 311 (ϵ 12000) (MeOH).

14,15-Dihydro, 15-hydroxy: Cystothiazole F

$\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_5\text{S}_2$ 438.568
 Prod. by *Cystobacter fuscus*. Antifungal agent. Powder. $[\alpha]_D^{25} +77$ (c, 0.07 in CHCl_3). λ_{max} 222 (ϵ 30400); 242 (ϵ 26800); 311 (ϵ 9990) (MeOH).

11 α ,12,14,15-Tetrahydro: Melithiazole D

[248938-46-7]
 $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_4\text{S}_2$ 424.584
 Prod. by *Archangium gephyra*. Oil. λ_{max} 215 ; 239 (MeOH).

11 α ,12,14,15-Tetrahydro, 15-methoxy: Melithiazole L

[248938-48-9]
 $\text{C}_{21}\text{H}_{30}\text{N}_2\text{O}_5\text{S}_2$ 454.61
 Prod. by *Archangium gephyra*. Oil. λ_{max} 217 (log ϵ 4.07); 240 (log ϵ 4.11); 280 (sh) (log ϵ 3.53) (MeOH).

Ojika, M. *et al.*, *J. Antibiot.*, 1998, **51**, 275-281 (Cystothiazoles)

Suzuki, Y. *et al.*, *Tetrahedron*, 1998, **54**, 11399-11404 (Cystothiazoles)

Boehlendorf, B. *et al.*, *Eur. J. Org. Chem.*, 1999, 2601-2608 (Melithiazoles)

Sasse, F. *et al.*, *J. Antibiot.*, 1999, **52**, 721-729 (Melithiazoles)

Söker, U. *et al.*, *Eur. J. Org. Chem.*, 2000, 1497-1502; 2021-2026 (synth)

Williams, D.R. *et al.*, *J.O.C.*, 2001, **66**, 8463-8469 (synth)

Suzuki, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 372-378 (biosynth)

DeRoy, P.L. *et al.*, *Org. Lett.*, 2003, **5**, 4163-4165 (synth)

Kato, K. *et al.*, *Tetrahedron*, 2003, **59**, 2679-2685 (synth)

Sasaki, T. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 770-771 (Cystothiazole B, synth, abs config)

Shao, J. *et al.*, *Org. Lett.*, 2004, **6**, 3083-3085 (synth)

Ojika, M. *et al.*, *Tetrahedron*, 2004, **60**, 187-194 (synth)

Akita, H. *et al.*, *Heterocycles*, 2005, **66**, 219-228 (Melithiazole B, synth)


Akita, H. *et al.*, *Tetrahedron*, 2006, **62**, 11592-11598 (synth)

Iwaki, Y. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 1610-1614 (synth)

Gebauer, J. *et al.*, *Eur. J. Org. Chem.*, 2008, 2701-2704 (Cystothiazole A, synth)

Melithiazole C **M-178**

[214420-47-0]



$\text{C}_{16}\text{H}_{21}\text{NO}_5\text{S}$ 339.412

Prod. by *Melittangium lichenicola*. Antifungal agent. Oil. Occurs as 6E/6Z-mixture which was not separated. λ_{max} 235 ; 329 (MeOH).

Boehlendorf, B. *et al.*, *Eur. J. Org. Chem.*, 1999, 2601-2608 (isol, pmr, cmr)

Sasse, F. *et al.*, *J. Antibiot.*, 1999, **52**, 721-729 (isol, activity)

Söker, U. *et al.*, *Eur. J. Org. Chem.*, 2000, 1497-1502; 2021-2026 (synth)

Takayama, H. *et al.*, *Heterocycles*, 2007, **71**, 75-85 (synth)

Gebauer, J. *et al.*, *Org. Lett.*, 2007, **9**, 3425-3427 (synth)

Melithiazole F **M-179**

[248938-49-0]

As Melithiazole B, M-177 with

R = $-\text{CH}_2\text{Ph}$

$\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_4\text{S}_2$ 470.612

Prod. by *Myxococcus stipitatus*. Antifungal and cytotoxic. Oil. λ_{max} 223 (log ϵ 4.43); 241 (log ϵ 4.36); 312 (log ϵ 3.9) (MeOH).

Böhlendorf, B. *et al.*, *Eur. J. Org. Chem.*, 1999, 2601-2608 (isol, struct)

Sasse, F. *et al.*, *J. Antibiot.*, 1999, **52**, 721-729 (isol, props)

Takayama, H. *et al.*, *Eur. J. Org. Chem.*, 2006, 644-649 (synth)

Melithiazole G **M-180**

[248938-50-3]

As Melithiazole B, M-177 with

R = $-\text{CH}_2(\text{CH}_3)\text{CH}_2\text{CH}_3$

$\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_4\text{S}_2$ 436.595

Prod. by *Myxococcus stipitatus*. λ_{max} 223 (log ϵ 4.28); 241 (log ϵ 4.24); 312 (log ϵ 3.8) (MeOH).

Böhlendorf, B. *et al.*, *Eur. J. Org. Chem.*, 1999, 2601-2608 (isol)

Sasse, F. *et al.*, *J. Antibiot.*, 1999, **52**, 721-729 (isol, props)

Melithiazole H **M-181**

Cystothiazole G

[248938-51-4]

As Melithiazole B, M-177 with

R = $-\text{C}^{14}\text{H}_2\text{CH}_3$

$\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_4\text{S}_2$ 408.542

Prod. by *Cystobacter fuscus* and *Myxococcus stipitatus*. Antifungal and cytotoxic. Solid. $[\alpha]_D^{25} +108$ (c, 0.04 in CHCl_3). λ_{max} 226 (ϵ 33400); 244 (ϵ 34000); 312 (ϵ 12500) (MeOH).

14-Oxo: Melithiazole M

[248938-53-6]

$\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_5\text{S}_2$ 422.525

Prod. by *Archangium gephyra*. Antifungal and cytotoxic. Oil. λ_{max} 217 (log ϵ 4.07); 239 (log ϵ 4.12); 305 (sh) (log ϵ 3.16) (MeOH).


Böhlendorf, B. *et al.*, *Eur. J. Org. Chem.*, 1999, 2601-2608 (Melithiazoles, isol)

Sasse, F. *et al.*, *J. Antibiot.*, 1999, **52**, 721-729 (Melithiazoles, isol, props)

Akita, H. *et al.*, *Tetrahedron*, 2004, **60**, 4735-4738 (Cystothiazole G)

Akita, H. *et al.*, *Heterocycles*, 2005, **66**, 219-228 (Melithiazole M, synth)

Melleumin A **M-182**



$\text{C}_{25}\text{H}_{29}\text{N}_3\text{O}_8$ 499.519

Prod. by the myxomycete *Physarum melleum*. Amorph. solid. $[\alpha]_D^{26} +27$ (c, 0.15 in MeOH). λ_{max} 225 (ϵ 13000); 254 (ϵ 15000) (MeOH).

Parent hydroxyacid, Me ester: Melleumin B


$\text{C}_{26}\text{H}_{33}\text{N}_3\text{O}_9$ 531.561

Prod. by the myxomycete *Physarum melleum*. Amorph. solid. λ_{max} 226 (ϵ 10000); 254 (ϵ 13000) (MeOH).

Nakatani, S. *et al.*, *Tet. Lett.*, 2005, **46**, 267-271 (isol, pmr, cmr)

Melochicorine **M-183**

2-Heptyl-1,2-dihydro-2-hydroxy-6-methoxy-4-methyl-3H-indol-3-one, 9CI. 2-Heptyl-2-hydroxy-6-methoxy-4-methylpseudooxindole
 [138615-21-1]




$\text{C}_{17}\text{H}_{25}\text{NO}_3$ 291.389

Alkaloid from the aerial parts of *Melochia corchorifolia* (Sterculiaceae). Viscous oil. $[\alpha]_D +56$ (CHCl_3).

Bhakuni, R.J. *et al.*, *Phytochemistry*, 1991, **30**, 3159 (isol, uv, ir, pmr, cmr, ms, struct)

Melochinine **M-184**

6-(11-Hydroxydodecyl)-3-methoxy-2-methyl-4-(1H)-pyridinone. 4-Hydroxy-6-(11-hydroxydodecyl)-3-methoxy-2-methylpyridine
 [70001-21-7]



$\text{C}_{19}\text{H}_{33}\text{NO}_3$ 323.475

Alkaloid from the leaves of *Melochia pyramidata*. Needles (MeOH/Et₂O). Mp 147°. $[\alpha]_D^{20} +11.2$ (c, 0.1 in EtOH).

O-β-D-Glucopyranoside: [77795-03-0]

$\text{C}_{25}\text{H}_{43}\text{NO}_8$ 485.617

Alkaloid from the leaves of *Melochia pyramidata* (Apocynaceae). Cryst. (MeOH/Et₂O). Mp 143-145°.

Ketone: Melochininone. Melochininone†

[70001-20-6]


$\text{C}_{19}\text{H}_{31}\text{NO}_3$ 321.459

Alkaloid from *Melochia pyramidata*.

Needles (Me₂CO). Mp 90°. Formerly named Melochinone, name changed to avoid confusion with Melochinone, M-185.

Medina, E. *et al.*, *Chem. Ber.*, 1979, **112**, 376; 1981, **114**, 814 (*isol, uv, ir, pmr, cmr, ms, struct*)
Medina, E. *et al.*, *Annalen*, 1981, 538 (*derivis*)
Spiteller, G. *et al.*, *Annalen*, 1981, 2096
Voss, G. *et al.*, *Annalen*, 1982, 1466 (*synth*)


Melochinone† **M-185**
[57609-68-4]



C₂₂H₂₁NO₂ 331.413
Alkaloid from *Melochia tomentosa* (Sterculiaceae). Mp 316-318° dec.

Kapadia, G.J. *et al.*, *J.A.C.S.*, 1975, **97**, 6814 (*isol, ir, uv, ms, pmr, cmr, cryst struct*)

Melofoline **M-186**
[107316-95-0]




C₂₆H₄₀N₄O₅ 488.626
Alkaloid from the aerial parts of *Melochia corchorifolia* (Apocynaceae). Cryst. (MeOH). Mp 305-307°. [α]_D²⁵ -252 (CHCl₃).

O-Ac: [107335-31-9]
Cryst. (MeOH). Mp 259-261°.

Bhakuni, R.S. *et al.*, *Phytochemistry*, 1987, **26**, 324 (*isol, ir, pmr, ms, struct*)


Melomorsine **M-187**
[156280-90-9]



C₄₁H₄₈N₄O₃ 644.855
Alkaloid from aerial parts of *Melodinus morsei* (Apocynaceae). Reddish cryst. Mp 70-71°. [α]_D²⁰ -197.64 (c, 0.635 in CHCl₃).

He, Y.-L. *et al.*, *J. Nat. Prod.*, 1994, **57**, 411 (*isol, uv, ir, pmr, cmr, ms, struct*)

Melonine **M-188**
2,11-Cyclo-11,12-secoaspido-spermidine,
9CI
[69771-89-7]



Absolute configuration

C₁₉H₂₆N₂ 282.428
Alkaloid from *Melodinus celastroides* (Apocynaceae). Mp 262° (as hydrochloride). [α]_D²² +82.


N^b-Oxide: N^b-Oxymelonine

[69771-90-0]
C₁₉H₂₆N₂O 298.427

Alkaloid from *Melodinus celastroides* (Apocynaceae). Mp 198°. [α]_D²² +110.

Baassou, S. *et al.*, *Tet. Lett.*, 1983, **24**, 761 (*uv, cmr, struct*)

Melonovine A **M-189**
[64408-09-9]



R = CH₂CH(CH₃)₂

C₂₇H₄₂N₄O₄ 486.653
CAS numbering shown. Alkaloid from the roots of *Melochia tomentosa* and from *Zizyphus jujuba* var. *inermis*. Cryst. (Me₂CO). Mp 295° dec. [α]_D -285 (CHCl₃).

Stereoisomer (1): **Pubescine A**

[75684-59-2]
C₂₇H₄₂N₄O₄ 486.653

Alkaloid from *Discaria pubescens* (Rhamnaceae). Needles (MeOH/petrol). Mp 247-250°. [α]_D -230 (c, 0.076 in MeOH). Conts. D-leucine (first report from a cyclopeptide alkaloid).

Stereoisomer (2): **Daechuine S5**

C₂₇H₄₂N₄O₄ 486.653

Alkaloid from stem bark of *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Mp 233-235°. [α]_D -421.3. No info. on stereochem. said in a review to be identical to Melonovine A.

Kapadia, G.J. *et al.*, *Phytochemistry*, 1977, **16**,

1431 (*isol, ir, pmr, ms, struct*)
Tschesche, R. *et al.*, *Phytochemistry*, 1980, **19**, 1000 (*Pubescine A*)
Han, B. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443 (*Daechuine S5*)
Tan, N.H. *et al.*, *Chem. Rev.*, 2006, **106**, 840-895 (*Daechuine S5, struct*)

Melonovine B **M-190**
[64408-10-2]

As Melonovine A, M-189 with
R = 4-Hydroxybenzyl

C₃₀H₄₀N₄O₅ 536.67
Alkaloid from the roots of *Melochia tomentosa* (Apocynaceae). Cryst. (Me₂CO/C₆H₆). Mp 200-206°.

Deoxy, 10',11'-dihydro, 11'-hydroxy:

Pandamine

[10233-40-6]

C₃₀H₄₂N₄O₅ 538.686


Alkaloid from *Panda oleosa* (Panda-ceae). Cryst. (EtOAc). Mp 272°. [α]_D -117 (CHCl₃).

Pais, M. *et al.*, *Ann. Chim. (Paris)*, 1966, **11**, 83; *CA*, **65**, 9018a (*Pandamine*)

Kapadia, G.J. *et al.*, *Phytochemistry*, 1977, **16**, 1431-1433 (*Melonovine B*)

Pais, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869-1872 (*Pandamine, cmr*)

Melophlins **M-191**



Melophlin A R¹ = -(CH₂)₁₄CH₃, R² = H
B R¹ = -CH₂CH₂CH(CH₃)(CH₂)₂CH₃, R² = CH₃
C R¹ = -(CH₂)₂CH(CH₃)(CH₂)₆CH₃, R² = CH₃
D R¹ = -(CH₂)₁₂CH₃, R² = H
E R¹ = -(CH₂)₁₂CH(CH₃)₂, R² = H
F R¹ = -(CH₂)₁₁CH(CH₃)CH₂CH₃, R² = H
G R¹ = -(CH₂)₁₂CH₃, R² = H
H R¹ = -(CH₂)₁₁CH(CH₃)₂, R² = H
I R¹ = -(CH₂)₁₀CH(CH₃)CH₂CH₃, R² = H
J R¹ = -(CH₂)₁₂CH₃, R² = CH₃
K R¹ = -(CH₂)₁₀CH(CH₃)₂, R² = H
L R¹ = -(CH₂)₁₁CH₃, R² = CH₃
M R¹ = -(CH₂)₁₀CH₃, R² = CH₃
N R¹ = -(CH₂)₈CH(CH₃)₂, R² = CH₃
O R¹ = -(CH₂)₈CH(CH₃)CH₂CH₃, R² = CH₃
P R¹ = -(CH₂)₁₄CH₃, R² = CH₃
Q R¹ = -(CH₂)₁₁CH(CH₃)₂, R² = CH₃
R R¹ = -(CH₂)₁₀CH(CH₃)CH₂CH₃, R² = CH₃
S R¹ = -(CH₂)₂CH(CH₃)(CH₂)₆CH₃, R² = CH₃

Tetramic acid derivs. Enolised β-triketones. Isol. from the marine sponge *Melophlus sarassinorum*.

Melophlin A
3-(1-Hydroxyhexadecylidene)-1-methyl-2,4-pyrrolidinedione, 9CI
[268745-62-6]

C₂₁H₃₇NO₃ 351.528
Cytotoxic agent. Oil. λ_{max} 244 (ε 9000); 284 (ε 15000) (MeOH).

Melophlin B
3-(1-Hydroxy-4-methyldodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[268745-63-7]

C₁₉H₃₃NO₃ 323.475
Cytotoxic. [α]_D -12.4 (c, 1.2 in MeOH). Has S-config. at C-5. λ_{max} 245 (ε 13800); 285 (ε 17000) (MeOH).

Melophlin C

3-(1-Hydroxy-5-methyldodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-46-8]

C₁₉H₃₃NO₃ 323.475
Antibacterial agent. Yellow oil. $[\alpha]_D^{20}$ -5.8 (c, 0.35 in MeOH). Isol. as a 1:2 mixt. of 5*R*- and 5*S*-enantiomers. λ_{\max} 248 (log ϵ 3.9); 286 (log ϵ 4.12) (MeOH).

Melophlin D

3-(1-Hydroxypentadecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-47-9]

C₂₀H₃₅NO₃ 337.501
Yellow oil. λ_{\max} 247 (log ϵ 3.74); 286 (log ϵ 3.93) (MeOH).

Melophlin E

3-(1-Hydroxy-14-methylpentadecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-48-0]

C₂₁H₃₇NO₃ 351.528
Yellow oil. λ_{\max} 246 (log ϵ 4.08); 286 (log ϵ 4.2) (MeOH).

Melophlin F

3-(1-Hydroxy-13-methylpentadecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-49-1]

C₂₁H₃₇NO₃ 351.528
Yellow oil. λ_{\max} 245 (log ϵ 3.06); 285 (log ϵ 3.15) (MeOH).

Melophlin G

3-(1-Hydroxytetradecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-50-4]

C₁₉H₃₃NO₃ 323.475
Yellow oil. λ_{\max} 247 (log ϵ 4); 285 (log ϵ 4.19) (MeOH).

Melophlin H

3-(1-Hydroxy-13-methyltetradecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-52-6]

C₂₀H₃₅NO₃ 337.501
Yellow oil. λ_{\max} 247 (log ϵ 4.01); 285 (log ϵ 4.25) (MeOH).

Melophlin I

3-(1-Hydroxy-12-methyltetradecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-55-9]

C₂₀H₃₅NO₃ 337.501
Yellow oil. $[\alpha]_D^{20}$ -2.6 (c, 0.34 in MeOH). λ_{\max} 245 (log ϵ 3.27); 286 (log ϵ 3.47) (MeOH).

Melophlin J

3-(1-Hydroxytetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-60-6]

C₂₀H₃₅NO₃ 337.501
Yellow oil. $[\alpha]_D^{20}$ -2.6 (c, 0.31 in MeOH). Isol. as a partial racemate, approx. 3:2 5*R*/5*S*-. λ_{\max} 246 (log ϵ 3.11); 285 (log ϵ 3.25) (MeOH).

Melophlin K

3-(1-Hydroxy-12-methyltridecylidene)-1-methyl-2,4-pyrrolidinedione
[499136-61-7]

C₁₉H₃₃NO₃ 323.475
Yellow oil. λ_{\max} 243 (log ϵ 3.04); 286 (log ϵ 3.09) (MeOH).

Melophlin L

3-(1-Hydroxytridecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-75-3]

C₁₉H₃₃NO₃ 323.475
Yellow oil. $[\alpha]_D^{20}$ -3.8 (c, 0.16 in MeOH). Isol. as a partial racemate, approx. 1:2 5*R*/5*S*-. λ_{\max} 245 (log ϵ 3); 286 (log ϵ 3.03) (MeOH).

Melophlin M

3-(1-Hydroxydodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-76-4]

C₁₈H₃₁NO₃ 309.448
Yellow oil. $[\alpha]_D^{20}$ -5 (c, 0.4 in MeOH). Isol. as a partial racemate, approx. 1:2 5*R*/5*S*-. λ_{\max} 248 (log ϵ 3.53); 286 (log ϵ 3.69) (MeOH).

Melophlin N

3-(1-Hydroxy-11-methyldodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-77-5]

C₁₉H₃₃NO₃ 323.475
Yellow oil. Isol. as a partial racemate, approx. 1:2 5*R*/5*S*-. λ_{\max} 248 (log ϵ 3.79); 286 (log ϵ 3.99) (MeOH).

Melophlin O

3-(1-Hydroxy-10-methyldodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione
[499136-81-1]

C₁₉H₃₃NO₃ 323.475
Yellow oil. Isol. a partial racemate, approx. 1:2 5*R*/5*S*-. λ_{\max} 248 (log ϵ 3.79); 286 (log ϵ 3.99) (MeOH).

Melophlin P

3-(1-Hydroxyhexadecylidene)-1,5-dimethyl-2,4-pyrrolidinedione

C₂₂H₃₉NO₃ 365.555
Yellowish oil. Racemate. λ_{\max} 245 (log ϵ 3.8); 282 (log ϵ 4.1) (MeOH).

Melophlin Q

3-(1-Hydroxy-13-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione

C₂₁H₃₇NO₃ 351.528
Yellowish oil. Racemate. λ_{\max} 246 (log ϵ 4.1); 283 (log ϵ 4.2) (MeOH).

Melophlin R

3-(1-Hydroxy-12-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione

C₂₁H₃₇NO₃ 351.528
Yellowish oil. Racemate. λ_{\max} 248 (log ϵ 4.2); 280 (log ϵ 4.2) (MeOH).

Melophlin S

3-(1-Hydroxy-5-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione


C₂₁H₃₇NO₃ 351.528
Yellowish oil. Racemate. λ_{\max} 248 (log ϵ 3.8); 283 (log ϵ 4.5) (MeOH).

Aoki, S. et al., *Tetrahedron*, 2000, **56**, 1833-1836 (*Melophlins A,B*)
Wang, C.-Y. et al., *J. Nat. Prod.*, 2003, **66**, 51-56 (*Melophlins C-O*)

Xu, J. et al., *Chem. Pharm. Bull.*, 2006, **54**, 852-854 (*Melophlins P-S*)

Meloscandonine

[28645-27-4]



Absolute Configuration

C₂₀H₂₀N₂O₂ 320.39

Alkaloid from *Melodinus scandens* (Apocynaceae). Mp 318-320° (after change of cryst. form) (>300°). $[\alpha]_D^{20}$ +72 (c, 1 in CHCl₃).

19-Epimer: 19-Epimeloscandonine

[151636-21-4]

C₂₀H₂₀N₂O₂ 320.39

Alkaloid from aerial parts of *Melodinus hemsleyanus* (Apocynaceae). Amorph. $[\alpha]_D^{23}$ +41.6 (c, 0.1 in CHCl₃).

Plat, M. et al., *Tet. Lett.*, 1970, 3395 (uv, ir, pmr, struct)

Mehri, H. et al., *Ann. Pharm. Fr.*, 1971, **29**, 291 (isol, uv, ir, ms)


Daudon, M. et al., *J.O.C.*, 1975, **40**, 2838 (cmr)

Daudon, M. et al., *Acta Cryst. B*, 1978, **34**, 232 (cryst struct)

Guo, L.-W. et al., *Phytochemistry*, 1993, **34**, 563 (*19-Epimeloscandonine*)

Meloscine

M-193



Absolute Configuration

C₁₉H₂₀N₂O 292.38

Numbering systems differ. Alkaloid from *Melodinus scandens* (Apocynaceae). Mp 178-180°.

Methiodide: Mp 240-245° dec.

17-Epimer: 3-Epimeloscine. 17-Epimeloscine

[24314-58-7]

C₁₉H₂₀N₂O 292.38

Alkaloid from *Melodinus scandens* (Apocynaceae). Noncryst. $[\alpha]_D^{25}$ +238 (c, 0.2 in EtOH).

17-Epimer, picrate: Mp 210-215°.

17-Epimer, N⁴-oxide: 3-Epimeloscine N-oxide

[34210-64-5]

C₁₉H₂₀N₂O₂ 308.379

Alkaloid from *Melodinus scandens* (Apocynaceae). Mp 203-207°. $[\alpha]_D^{22}$ +310.

17-Epimer, 9-hydroxy: 9-Hydroxyepimeloscine


[169700-70-3]

$C_{19}H_{20}N_2O_2$ 308.379
Alkaloid from leaves of *Melodinus scandens* (Apocynaceae). $[\alpha]_D^{20} +206$ (c, 1 in $CHCl_3$).

- Bernauer, K. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 1886 (*uv, ir, pmr, ms, ord*)
Oberhänsli, W.E. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 1905 (*cryst struct*)
Mehri, H. *et al.*, *Ann. Pharm. Fr.*, 1971, **29**, 291 (*oxide*)
Daudon, M. *et al.*, *J.O.C.*, 1975, **40**, 2838 (*cmr*)
Overman, L.E. *et al.*, *J.A.C.S.*, 1991, **113**, 2598 (*synth*)
Mehri, H. *et al.*, *Phytochemistry*, 1995, **40**, 1005 (*9-Hydroxyepimelosmine*)

Melosmine**M-194**

2,3-Dimethoxy-7,7-dimethyl-7H-dibenzo[de,g]quinoline-1,9-diol, 9CI
[81525-68-0]



$C_{20}H_{19}NO_4$ 337.374
Alkaloid from the stem bark of *Guatteria melosma* and *Guatteria ouregou* (Annonaceae). Light-yellow cuboidal cryst. (EtOH). Mp 104°.

O¹-Me: Melosmidine

[81525-72-6]
 $C_{21}H_{21}NO_4$ 351.401

Alkaloid from the stem bark of *Guatteria melosma* (Annonaceae). Yellow amorph. solid. Mp 170-171°.

4,5-Dihydro: Dihydromelosmine

[84391-88-8]
 $C_{20}H_{21}NO_4$ 339.39

Alkaloid from the stem bark of *Guatteria ouregou* (Annonaceae). Amorph.

Leboeuf, M. *et al.*, *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1982, **295**, 191 (*Dihydromelosmine*)

Zabel, V. *et al.*, *J. Nat. Prod.*, 1982, **45**, 94 (*Melosmine, Melosmidine, isol, uv, ir, pmr, ms, struct*)


Hocquemiller, R. *et al.*, *Tet. Lett.*, 1982, **23**, 4247 (*cmr*)

Leboeuf, M. *et al.*, *Tetrahedron*, 1982, **38**, 2889 (*pmr, cmr*)

Leboeuf, M. *et al.*, *Planta Med.*, 1983, **48**, 234 (*isol, uv, ir, pmr, struct, Dihydromelosmine*)

Melovinone**M-195**

[69905-17-5]



$C_{24}H_{29}NO_4$ 395.497

An open chain analogue of Melochinone, M-185. Alkaloid from the roots of


Melochia tomentosa (Sterculiaceae).

Cryst. (C_6H_6 /petrol). Mp 134-136°.

Kapadia, G.J. *et al.*, *Phytochemistry*, 1978, **17**, 1445 (*isol, ir, uv, ms, pmr, struct*)

Memnobotrin A**M-196**

[256235-08-2]



Absolute Configuration

$C_{25}H_{33}NO_5$ 427.539

Prod. by *Memnoniella echinata*. Cytotoxic agent. Pale yellow glass. $[\alpha]_D^{25} +9.6$ (c, 1.14 in MeOH). Dec. at 240-250°. λ_{max} 217 (ε 26200); 257 (ε 5600); 300 (ε 2300) (MeOH).

N-(2-Hydroxyethyl): Memnobotrin B

[256235-09-3]


$C_{27}H_{37}NO_6$ 471.592

Prod. by *Memnoniella echinata*. Cytotoxic agent. Cream powder. Mp 186-192°. $[\alpha]_D^{25} +11.7$ (c, 1.25 in MeOH). λ_{max} 218 (ε 37200); 259 (ε 10600); 302 (ε 3300) (MeOH).

Hinkley, S.F. *et al.*, *J. Antibiot.*, 1999, **52**, 988-997 (*isol, uv, ir, pmr, cmr*)

Menisarine**M-197**

[6415-97-0]



$C_{36}H_{34}N_2O_6$ 590.674

Alkaloid from *Cocculus laeabe* and *Cocculus sarmentosus* (Menispermaceae). Mp 203°. $[\alpha]_D +149$ ($CHCl_3$).

O-De-Me: Normenisarine. O-Demethylmenisarine

$C_{35}H_{32}N_2O_6$ 576.648

Alkaloid from *Cocculus trilobus* (Menispermaceae). Mp 223°. Location of OH not determined.

Kondo, H. *et al.*, *Yakugaku Zasshi*, 1930, **50**, 633-645; *Ann. Chim. Farm.*, 1935, **55**, 911; *CA*, **24**, 5302; **30**, 726 (*isol, Menisarine, Normenisarine*)

Sinha, A. *et al.*, *J. Proc. Inst. Chem. (India)*, 1960, **32**, 250-252; *CA*, **55**, 18886b (*occur*)
Tomita, M. *et al.*, *Tet. Lett.*, 1962, 635-636 (*struct*)

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 190; *CA*, **59**, 3971h (*occur*)

Menismine**M-198**

[1359-64-4]

$C_{21}H_{28}NO_4^{\oplus}$ 358.456

Struct. unknown. Quaternary alkaloid from the root bark of *Cissampelos pareira* (Menispermaceae).

Chloride:

Amorph.

Iodide:

Cryst. (MeOH/EtOAc). Mp 140-142°. $[\alpha]_D^{20} +24$ (c, 0.5 in MeOH).


Picrate:

Cryst. + $\frac{3}{4}H_2O$. Mp 116-117°.

Srivastava, R.M. *et al.*, *Chem. Ber.*, 1964, **97**, 2732-2741 (*isol, uv, ir*)

Menisporphine**M-199**

5,6,9-Trimethoxy-7H-dibenzo[de,h]quinolin-7-one, 9CI
[83287-02-9]



$C_{19}H_{15}NO_4$ 321.332

Alkaloid from *Menispermum dauricum* (Menispermaceae). Yellow needles (CH_2Cl_2/Me_2CO). Mp 199.5-200.5°. λ_{max} 254 (log ε 4.72); 288 (sh) (log ε 4.13); 319 (log ε 3.97); 368 (log ε 3.91); 420 (log ε 3.97) (EtOH).

▶HQ1787600**O⁹-De-Me: Daurioxoisoporphine D**

[356047-67-1]

$C_{18}H_{13}NO_4$ 307.305

Alkaloid from the rhizomes of *Menispermum dauricum*. Amorph. yellow powder. λ_{max} 210 (log ε 4.32); 250 (log ε 3.42); 310 (log ε 2.43); 360 (log ε 2.42); 406 (log ε 3.34); 420 (log ε 4.33); 458 (log ε 3.43) (MeOH).

2,3-Dihydro: 2,3-Dihydromenisporphine

[100009-83-4]

$C_{19}H_{17}NO_4$ 323.348

Alkaloid from the rhizomes of *Menispermum dauricum* (Menispermaceae). Yellow needles (Me_2CO). Mp 177-180°. λ_{max} 225 (log ε 4.37); 255 (sh) (log ε 4.31); 274 (log ε 4.42); 372 (log ε 3.76) (EtOH).

Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 2659-2660 (*uv, ir, pmr, ms, struct, synth*)

Kunitomo, J. *et al.*, *Tetrahedron*, 1983, **39**,

3261-3265 (*uv, ir, pmr, ms, struct, synth*)

Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1985,


33, 2778-2782 (*2,3-Dihydromenisporphine*)

Yu, B.-W. *et al.*, *J. Nat. Prod.*, 2001, **64**, 968-

970 (*Daurioxoisoporphine D*)

Mequinine**M-200**

[166990-19-8]




C₂₃H₂₆NO₇[⊖] 428.461
Alkaloid from *Meconopsis quintuplinervia* (Papaveraceae).

Wang, M. *et al.*, *Tianran Chanwu Yanjiu Yu Kaifa*, 1995, 7, 32; *CA*, **123**, 138808j

2-Mercapto-1H-indole-3-carboxaldehyde, 9CI M-201

3-Formyl-2-mercaptoindole
[183946-30-7]



C₉H₇NOS 177.226
Solid.

S-Me: 2-(Methylthio)-1H-indole-3-carboxaldehyde. 3-Formyl-2-(methylthio)indole. **Brassicinal A**
[113866-44-7]

C₁₀H₉NOS 191.253

Alkaloid from Chinese cabbage (*Brassica campestris* ssp. *pekinensis*) inoculated with *Pseudomonas cichorii*. Phytoalexin. Mp 210-213°.

6-Methoxy, *S*-Me: 6-Methoxy-2-(methylthio)-1H-indole-3-carboxaldehyde. **Capparin B**

C₁₁H₁₁NO₂S 221.279

Alkaloid from *Capparis himalayensis*. Amorph. powder (CHCl₃/MeOH). Mp 202-204° dec. λ_{max} 192 (log ε 3.29); 223 (log ε 3.17); 247 (log ε 3.17); 284 (log ε 2.97); 333 (log ε 2.83) (MeOH).


Monde, K. *et al.*, *Chem. Lett.*, 1990, 209 (*isol*, *uv*, *pmr*, *struct*)

Pedras, M.S.C. *et al.*, *Chem. Comm.*, 1998, 1565-1566 (*synth*)

Pedras, M.S.C. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 1196-1202 (*synth*, *pmr*, *cmr*, *ms*)

Li, Y.-Q. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 189-191 (*Capparin B*)

2-Mercapto-1H-indole-3-carboxylic acid, 9CI M-202



C₉H₇NO₂S 193.226

S-Me, *Me ester*: **Brassicinate A**
[330195-74-9]

C₁₁H₁₁NO₂S 221.279


Alkaloid from *Brassica napus* ssp. *rapifera* (rutabaga). Phytoalexin. Needles (CH₂Cl₂/hexane). Mp 109-110° (105-107°). λ_{max} 206 (log ε 4.4); 238 (log ε 4.4); 268 (log ε 3.9); 301 (log ε 4) (no solvent reported).

Somei, M. *et al.*, *Heterocycles*, 2001, **54**, 425-432 (*S*-Me *Me ester*, *synth*)

Pedras, M.S.C. *et al.*, *J.O.C.*, 2004, **69**, 4471-4476 (*S*-Me *Me ester*, *isol*)

5-Mercaptopentanoic acid, M-203 9CI

5-Mercaptovaleric acid, 8CI
[30247-98-4]



C₅H₁₀O₂S 134.199
Liq. or cryst. d₄²⁰ 1.12. Mp 23.5-25°. Bp_{0.3} 96°. n_D²⁵ 1.4873.

Et ester: [70124-55-9]

C₇H₁₄O₂S 162.252

Oil. d₄²⁰ 1.16. Bp₄ 82°. n_D²⁵ 1.5305. n_D²⁰ 1.4618.

S-Me: 5-(Methylthio)pentanoic acid, 9CI
[69082-95-7]

C₆H₁₂O₂S 148.226

Liq. Bp_{0.3} 107°. n_D²⁷ 1.4800.

S-Me, *S*-oxide: 5-(Methylsulfinyl)pentanoic acid, 9CI

[69082-96-8]

C₆H₁₂O₃S 164.225

Cryst. Mp 55-57°.

S-Me, *S,S*-dioxide: 5-(Methylsulfonyl)pentanoic acid, 9CI

[69082-97-9]

C₆H₁₂O₄S 180.224

Cryst. Mp 113-116°.

S-Me, *Me ester*: [69082-98-0]

C₇H₁₄O₂S 162.252

Liq. Bp₇ 92-95°.

S-Me, *Et ester*:

C₈H₁₆O₂S 176.279

Liq. Bp_{10.5} 65°. n_D²⁵ 1.4594.

S-Me, *Et ester*, *S,S*-dioxide:

C₈H₁₆O₄S 208.278

Liq. Bp_{0.4} 148°. n_D²⁵ 1.4608.

S-Me, nitrile: 5-(Methylthio)pentanenitrile. 1-Cyano-4-(methylthio)butane. 4-(Methylthio)butyl cyanide
[59121-25-4]

C₆H₁₁NS 129.226

Hydrolyt. product from *Brassica* spp.

S-Triphenylmethyl:

C₂₄H₂₄O₂S 376.518

Solid. Mp 124-125°.

Truce, W.E. *et al.*, *J.A.C.S.*, 1955, **77**, 5063-5067 (*S*-Me, *S*-Me *Et ester*)

Petrova, R.G. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1958, 290-295; *CA*, **52**, 12750d (*synth*, *Et ester*)

Buttery, R.G. *et al.*, *J. Agric. Food Chem.*, 1976, **24**, 829-832 (*S*-Me nitrile, *isol*)

Buckingham, D.A. *et al.*, *Chem. Comm.*, 1978, 705-707 (*S*-Me oxides)

Dawson, G.W. *et al.*, *J. Biol. Chem.*, 1993, **268**, 27154-27159 (*S*-Me nitrile, *synth*)

Kurahara, M. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1505-1510 (*synth*, *ir*, *pmr*, *Et ester*)

Chiang, W.C.K. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 1018 (*S*-Me nitrile, *isol*)

Majer, P. *et al.*, *J. Med. Chem.*, 2003, **46**, 1989-1996 (*S*-trityl, *synth*, *pmr*, *cmr*)

Jessing, M. *et al.*, *J.O.C.*, 2006, **71**, 6734-6741 (*synth*)

5-Mercapto-4-pentenoic acid M-204 HSCH=CHCH₂CH₂COOH

C₅H₈O₂S 132.183

(*E*)-form

Nitrile, *S*-[4-hydroxy-3-methoxy-*E*-cinnamoyl-(→6)-β-*D*-glucopyranosyl]:

[923593-04-8]

C₂₁H₂₅NO₈S 451.496

Constit. of the seeds of *Raphanus sativus* (radish). Yellow oil. λ_{max} 320 (log ε 3.8) (MeOH).

Nitrile, *S*-[4-hydroxy-3,5-dimethoxy-*E*-cinnamoyl-(→6)-β-*D*-glucopyranosyl]:

[923593-05-9]

C₂₂H₂₇NO₉S 481.523

Constit. of the seeds of *Raphanus sativus* (radish). Yellow oil. λ_{max} 279 (sh); 330 (log ε 3.81) (MeOH).

(*Z*)-form

Nitrile, *S*-[4-hydroxy-3,5-dimethoxy-*E*-cinnamoyl-(→6)-β-*D*-glucopyranosyl]:

[923593-06-0]

C₂₂H₂₇NO₉S 481.523


Constit. of the seeds of *Raphanus sativus* (radish). Yellow oil. λ_{max} 279 (sh); 330 (log ε 3.81) (MeOH).

Duan, L.-X. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 2953-2957 (*isol*, *pmr*, *cmr*)

Meridianin A

M-205

[213472-98-1]



C₁₂H₁₀N₄O 226.237

Alkaloid from the tunicate *Aplidium meridianum*. Protein kinase inhibitor. Yellow needles (MeOH aq.). Mp 164-168°. λ_{max} 248 (log ε 3.68); 356 (log ε 3.58) (CHCl₃). λ_{max} 248 (ε 4786); 356 (ε 3800) (MeOH) (Berdy).

6-Bromo: **Meridianin B**

[213472-99-2]

C₁₂H₉BrN₄O 305.133

Alkaloid from *Aplidium meridianum*.

Cytotoxic agent. Yellow powder

(EtOAc). Mp 190° dec. λ_{max} 246 (log ε 3.87); 354 (log ε 3.71) (CHCl₃). λ_{max} 246 (ε 7413); 354 (ε 5128) (MeOH) (Berdy).

7-Bromo: **Meridianin E**

[213473-03-1]

C₁₂H₉BrN₄O 305.133

Alkaloid from *Aplidium meridianum*.

Cytotoxic agent. Yellow cryst. (MeOH

aq.). Mp 172-175°. λ_{max} 224 (log ε 4.2); 358 (log ε 3.85) (MeOH).

Deoxy: **Meridianin G**

[289628-76-8]

C₁₂H₁₀N₄ 210.238

Alkaloid from *Aplidium meridianum*.

Yellow needles (MeOH aq.). Mp 215° dec. λ_{max} 220 (ε 8530); 331 (ε 3530) (MeOH).

Deoxy, 5-bromo: **Meridianin C**

[213473-00-8]

C₁₂H₉BrN₄ 289.134

Alkaloid from *Aplidium meridianum*.

Cytotoxic agent. Yellow powder (MeOH aq.). Mp 103-106°. λ_{\max} 244 (log ϵ 4.06); 324 (log ϵ 4.1) (CHCl₃). λ_{\max} 246 (ϵ 11480); 324 (ϵ 12590) (MeOH) (Berdy).

Deoxy, 6-bromo: Meridianin D

[213473-01-9]

C₁₂H₉BrN₄ 289.134Alkaloid from *Aplidium meridianum*.

Cytotoxic agent. Yellow powder (EtOAc/MeOH). Mp 218-221°. λ_{\max} 240 (log ϵ 4.17); 324 (log ϵ 4.14) (CHCl₃). λ_{\max} 240 (ϵ 14790); 324 (ϵ 13800) (MeOH) (Berdy).

Deoxy, 5,6-dibromo: Meridianin F

[690267-59-5]

C₁₂H₇Br₂N₄ 368.03Alkaloid from *Aplidium meridianum*.

Yellow needles (MeOH aq.). Mp 175°. λ_{\max} 201 (ϵ 30440); 215 (ϵ 28825); 274 (ϵ 7220); 292 (ϵ 6180); 329 (ϵ 8260) (MeOH).

Franco, L.H. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1130-1132 (*isol, uv, ir, pmr, cmr*)

Jiang, B. *et al.*, *Heterocycles*, 2000, **53**, 1489-1498 (*Meridianin D, synth*)

Fresneda, P.M. *et al.*, *Tetrahedron*, 2001, **57**, 2355-2363 (*synth*)

Gompel, M. *et al.*, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 1703-1707 (*activity*)


Karpov, A.S. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 6951-6956 (*synth*)

Simon, G. *et al.*, *J. Het. Chem.*, 2007, **44**, 793-801 (*Meridianin C, synth*)

Seldes, A.M. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 555-563 (*Meridianins F,G*)

Meridine*12(13H)-Meridinone*

[129722-90-3]

C₁₈H₉N₃O₂ 299.288

Tautomeric with the 12-oxo-13H-form, both tautomers isolable. Alkaloid from the ascidian *Amphicarpa meridiana* and the marine sponge *Corticium* sp. Cytotoxic. Exhibits antifungal activity. Yellow amorph. solid. Mp 250°.

Oxo-form*Isomeridine*

Separately isolable, tautom. to Meridine after 1-2 d. in CDCl₃ soln.

[129722-91-4]

Schmitz, F.J. *et al.*, *Pure Appl. Chem.*, 1990, **62**, 1393-1396 (*Isomeridine*)

Schmitz, F.J. *et al.*, *J.O.C.*, 1991, **56**, 804-808 (*isol, pmr, cmr, cryst struct*)

McCarthy, P.J. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1664-1668 (*isol, props*)


Kitahara, Y. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1363-1364 (*synth*)

Bontemps, N. *et al.*, *Tetrahedron*, 1997, **53**, 1743-1750 (*synth*)

Kitahara, Y. *et al.*, *Tetrahedron*, 1998, **54**, 8421-8432 (*synth, pmr, cmr*)

Merresectine C

M-207

R = -CH₂CH=C(CH₃)₂C₂₅H₃₅NO₃ 397.556

Alkaloid from the roots of *Merremia cissoides*, *Merremia dissecta* and *Merremia quinquefolia*. Oil.

O- β -D-Glucopyranoside: **Merresectine B** [219829-75-1]

C₃₁H₄₅NO₈ 559.698

Alkaloid from *Merremia cissoides*, *Merremia dissecta* and *Merremia quinquefolia*. Oil. [α]_D²⁰ -10.2 (c, 0.1 in MeOH).

3-Epimer: **3 β -Merresectine C**C₂₅H₃₅NO₃ 397.556Alkaloid from *Merremia* spp.

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (*isol, pmr, cmr, ms*)

Merresectine D

M-208

As Merresectine C, M-207 with

R = -OMe

C₂₁H₂₉NO₄ 359.464

Alkaloid from *Merremia guerichii*, *Merremia kentrocaulos* and *Merremia quinata*. Merresectine G and Merresectine H also isol. but structs. not fully elucidated.

O- β -D-Glucopyranoside: **Merresectine D glucoside**

C₂₇H₃₉NO₉ 521.606Alkaloid from *Merremia guerichii*. Oil.Demethoxy: **Merresectine E**C₂₀H₂₇NO₃ 329.438

Alkaloid from *Merremia dissecta* and *Merremia guerichii*.


Demethoxy, O- β -D-glucopyranoside:**Merresectine E glucoside**C₂₆H₃₇NO₈ 491.58Alkaloid from *Merremia* spp.3-Epimer: **3 β -Merresectine D**C₂₁H₂₉NO₄ 359.464

Alkaloid from *Merremia guerichii* and *Merremia kentrocaulos*.

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (*isol, pmr, cmr*)

Merresectine F


M-209

C₂₀H₂₅NO₃ 327.422Alkaloid from *Merremia* spp.3-Epimer: **3 β -Merresectine F**C₂₀H₂₅NO₃ 327.422Alkaloid from *Merremia* spp.

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (*isol, pmr, cmr*)

Mersicarpine

M-210

C₁₇H₂₀N₂O₂ 284.357

Poss. derived from Eburenine, E-3 by rearrangement and elimination of C-5 and C-6. Alkaloid from *Kopsia arborea* and *Kopsia fruticosa*. Oil. [α]_D -18 (c, 0.28 in CHCl₃). λ_{\max} 204; 239; 246; 266; 275; 329 (EtOH).


Kam, T.-S. *et al.*, *Tet. Lett.*, 2004, **45**, 5995-5998 (*isol, pmr, cmr*)

Magolan, J. *et al.*, *Org. Lett.*, 2008, **10**, 1437-1440 (*synth*)

Mersidasine C

M-211

[1011489-43-2]

C₂₅H₃₂N₂O₉ 504.536

Alkaloid from the leaves of *Kopsia singapurensis*. Pale yellow oil. [α]_D +23 (c, 0.22 in CHCl₃). λ_{\max} 209 (log ϵ 4.31); 234 (log ϵ 3.73); 284 (log ϵ 3.17) (EtOH).

16-Epimer: **Mersidasine D**

[1011489-44-3]

C₂₅H₃₂N₂O₉ 504.536


Alkaloid from the leaves of *Kopsia singapurensis*. Oil. [α]_D -205 (c, 0.12 in CHCl₃). λ_{\max} 218 (log ϵ 4.23); 244 (log ϵ 3.73); 284 (log ϵ 3.01) (EtOH).

Subramaniam, G. *et al.*, *Tetrahedron*, 2008, **64**, 1397-1408 (*isol, pmr, cmr, ms*)

Mersifoline A

M-212

[1011489-38-5]

C₂₃H₂₄N₂O₇ 440.452Alkaloid from the leaves of *Kopsia*

singaporensis. Oil. $[\alpha]_D$ -19 (c, 0.26 in CHCl_3). λ_{max} 206 (log ϵ 4.02); 230 (log ϵ 3.36); 271 (log ϵ 3) (EtOH).

11-Methoxy: Mersifoline B

[1011489-39-6]

 $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_8$ 470.478

Alkaloid from the leaves of *Kopsia singaporensis*. Oil. $[\alpha]_D$ -14 (c, 0.18 in CHCl_3). λ_{max} 211 (log ϵ 4.4); 229 (log ϵ 4.05); 270 (log ϵ 3.52) (EtOH).

20-Epimer: Mersifoline C

[1011489-40-9]

 $\text{C}_{23}\text{H}_{24}\text{N}_2\text{O}_7$ 440.452


Alkaloid from the leaves of *Kopsia singaporensis*. Oil. $[\alpha]_D$ +18 (c, 0.28 in CHCl_3). λ_{max} 211 (log ϵ 3.84); 230 (log ϵ 3.72); 273 (log ϵ 3.3) (EtOH).

Subramaniam, G. *et al.*, *Tetrahedron*, 2008, **64**, 1397-1408 (*isol, pmr, cmr, ms*)

Mersilongine

M-213

[705948-72-7]

 $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_8$ 444.44


Probable abs. config. based on speculative biosynthesis. Alkaloid from the leaves of a Malayan *Kopsia* sp. Oil. $[\alpha]_D$ +106 (c, 0.17 in CHCl_3). λ_{max} 220 ; 250 (sh) ; 291 (no solvent reported).

Kam, T.-S. *et al.*, *Tet. Lett.*, 2004, **45**, 3521-3524 (*isol, pmr, cmr*)

Mersilosine

M-214

[367943-78-0]



Absolute Configuration

 $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_{10}$ 502.477

Rearranged indole alkaloid. Alkaloid from the leaves of *Kopsia singaporensis*. Amorph. solid. Mp 185-188°. $[\alpha]_D$ -50 (c, 0.04 in CHCl_3). λ_{max} 215 (log ϵ 4.53); 242 (log ϵ 4.09); 291 (log ϵ 3.55) (EtOH).

15-Epimer: Mersilosine B $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_{10}$ 502.477

Alkaloid from the leaves of *Kopsia singaporensis*. Amorph. solid. Mp 202-204°. $[\alpha]_D$ +24 (c, 0.2 in CHCl_3). λ_{max} 212 (log ϵ 3.93); 244 (log ϵ 3.44); 287 (log ϵ 2.98) (EtOH).

16-Epimer: Mersilosine A $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_{10}$ 502.477

Alkaloid from the leaves of *Kopsia*


singaporensis. Oil. $[\alpha]_D$ -38 (c, 0.66 in CHCl_3). λ_{max} 213 (log ϵ 4.53); 235 (log ϵ 4.17); 289 (log ϵ 3.66) (EtOH).

Subramaniam, G. *et al.*, *Tetrahedron*, 2008, **64**, 1397-1408 (*isol, pmr, cmr, ms*)

Mersinaline

M-215

[953394-57-5]

 $\text{C}_{26}\text{H}_{28}\text{N}_2\text{O}_{11}$ 544.514


Alkaloid from *Kopsia singaporensis*. Pale yellow oil. $[\alpha]_D$ +98 (c, 0.15 in CHCl_3).

Subramaniam, G. *et al.*, *Tet. Lett.*, 2007, **48**, 6677-6680 (*isol, pmr, cmr*)

Mersingine A

M-216

[162616-54-8]



Absolute Configuration

 $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$ 363.415

Novel cage struct.; see also Kopsinitarine A, K-79. Alkaloid from leaves of *Kopsia teoi* (Apocynaceae). $[\alpha]_D$ +88 (c, 0.058 in CHCl_3). Artifact. λ_{max} 213 (log ϵ 4.54); 247 (log ϵ 3.85); 290 (log ϵ 3.28) (EtOH).

14,15-Dihydro, 15 α -hydroxy: Mersingine B [162665-99-8]

 $\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_4$ 381.43


From a Malaysian *Kopsia teoi* (Apocynaceae). $[\alpha]_D$ -82 (c, 0.022 in CHCl_3). Artifact. λ_{max} 213 (log ϵ 4.72); 246 (log ϵ 4.05); 290 (log ϵ 3.52) (EtOH).

Kam, T.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1109-1112 (*isol, uv, pmr, cmr, ms, struct*)

Mersinine A

M-217

[367943-76-8]



Absolute Configuration

 $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_9$ 500.504

Rearranged indole alkaloid. Alkaloid from the leaves of *Kopsia singaporensis*. Cryst. (petrol). Mp 204-205°. $[\alpha]_D$ -58 (c,

0.27 in CHCl_3). λ_{max} 213 (log ϵ 4.5); 239 (log ϵ 4.1); 289 (log ϵ 3.82) (EtOH).

14,15-Dihydro, 15 α -hydroxy: Mersidasine A

 $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_{10}$ 518.519

Alkaloid from the leaves of *Kopsia singaporensis*. Pale yellow oil. $[\alpha]_D$ -6 (c, 0.41 in CHCl_3). λ_{max} 211 (log ϵ 4.35); 236 (log ϵ 3.96); 286 (log ϵ 3.46) (EtOH).

16-Epimer: Mersinine B [367943-77-9]

 $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_9$ 500.504

Alkaloid from the leaves of *Kopsia singaporensis*. Oil. $[\alpha]_D$ -77 (c, 0.48 in CHCl_3). λ_{max} 214 (log ϵ 4.55); 241 (log ϵ 4.17); 287 (log ϵ 3.79) (EtOH).

16-Epimer, 14,15-dihydro, 15 α -hydroxy: Mersidasine B

 $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_{10}$ 518.519

Alkaloid from the leaves of *Kopsia singaporensis*. Amorph. solid. Mp 215-217°. $[\alpha]_D$ -7 (c, 0.47 in CHCl_3). λ_{max} 214 (log ϵ 4); 240 (log ϵ 3.6); 287 (log ϵ 3.04); 293 (log ϵ 3.07) (EtOH).

16-Epimer, 14,15-dihydro, 14 α ,15 β -dihydroxy: Mersidasine E

 $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_{11}$ 534.519

Alkaloid from the leaves of *Kopsia singaporensis*. Pale yellow oil. $[\alpha]_D$ -7 (c, 0.37 in CHCl_3). λ_{max} 213 (log ϵ 4.63); 239 (log ϵ 4.25); 287 (log ϵ 3.76) (EtOH).

20-Epimer, 14,15-dihydro, 15 α -hydroxy: Mersidasine F

 $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_{10}$ 518.519

Alkaloid from the leaves of *Kopsia singaporensis*. Oil. $[\alpha]_D$ +33 (c, 0.15 in CHCl_3). λ_{max} 213 (log ϵ 4.18); 237 (log ϵ 3.75); 290 (log ϵ 3.25) (EtOH).

16,20-Diepimer: Mersinine C

 $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_9$ 500.504

Alkaloid from the leaves of *Kopsia singaporensis*. Cryst. (petrol). Mp 108-110°. $[\alpha]_D$ +14 (c, 0.72 in CHCl_3). λ_{max} 217 (log ϵ 4.46); 244 (log ϵ 3.99); 288 (log ϵ 3.46) (EtOH).

16,20-Diepimer, 14,15-dihydro, 15 α -hydroxy: Mersidasine G

 $\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_{10}$ 518.519


Alkaloid from the leaves of *Kopsia singaporensis*. Oil. $[\alpha]_D$ -30 (c, 0.33 in CHCl_3). λ_{max} 217 (log ϵ 4.33); 240 (log ϵ 3.94); 288 (log ϵ 3.39) (EtOH).

Subramaniam, G. *et al.*, *Tetrahedron*, 2008, **64**, 1397-1408 (*isol, pmr, cmr, ms*)

Mersirachine

M-218

[953394-56-4]


 $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_9$ 500.504

Alkaloid from *Kopsia singapurensis*. Oil. $[\alpha]_D^{20}$ -50 (c, 0.05 in CHCl_3).

Subramaniam, G. *et al.*, *Tet. Lett.*, 2007, **48**, 6677-6680 (*isol*, *pmr*, *cmr*)

Mescaline M-219

3,4,5-Trimethoxybenzeneethanamine, 9CI. 3,4,5-Trimethoxyphenethylamine. *Mezcaline*. *Hallucinex*. *TMPEA* [54-04-6]



$\text{C}_{11}\text{H}_{17}\text{NO}_3$ 211.26

Alkaloid from mescal (*Lophophora williamsii*), *Trichocereus* spp., *Gymnocycium gibbosum*, *Opuntia cylindrica* and other spp. in the Cactaceae. Hallucinogenic; a psychotomimetic agent. Mp 35-36°. Bp₁₂ 180°. Log P 0.18 (calc).

- Adverse systemic effects by ingestion etc. esp. on CNS. LD₅₀ (mus, orl) 880 mg/kg, exp. reprod. and teratogenic effects. SI2625000

Hydrochloride: [832-92-8]

Mp 184°.

- LD₅₀ (mus, orl) 912 mg/kg. SI2800000

N-Formyl: **N-Formylmescaline**

$\text{C}_{12}\text{H}_{17}\text{NO}_4$ 239.271

Alkaloid from peyote cactus (*Lophophora williamsii*) (Cactaceae). Mp 68-69°.

N-Ac: **N-Acetylmescaline**. *N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]acetamide, 9CI

[4593-89-9]

$\text{C}_{13}\text{H}_{19}\text{NO}_4$ 253.297

Alkaloid from mescal (*Lophophora williamsii*). Metab. of Mescaline (Cactaceae). Mp 93-94°.

N-Benzoyl: **N-Benzoylmescaline**

$\text{C}_{18}\text{H}_{21}\text{NO}_4$ 315.368

Alkaloid from *Pothomorphe umbellata*. Antibacterial agent. Mp 123-125°. λ_{max} 213 (ε 23000); 225 (sh) (ε 20000) (MeOH).

N-(2,5-Dimethoxybenzoyl): *N*-(2,5-Dimethoxybenzoyl)mescaline. **Taiwanamide C**

[1000991-98-9]

$\text{C}_{20}\text{H}_{25}\text{NO}_6$ 375.421

Alkaloid from the stems of *Piper taiwanense*. Needles (hexane). Mp 95-97°. λ_{max} 239 (log ε 4.86); 314 (log ε 4.42) (MeOH).

N-Me: **N-Methylmescaline**

[4838-96-4]

$\text{C}_{12}\text{H}_{19}\text{NO}_3$ 225.287

Alkaloid from *Alhagi pseudalhagi* and *Pelecyphora aselliformis* (Fabaceae, Cactaceae). Cryst. (MeOH/Et₂O) (as hydrochloride). Mp 201-202° (hydrochloride).

N,N-Di-Me: **Trichocereine**

[529-91-9]

$\text{C}_{13}\text{H}_{21}\text{NO}_3$ 239.314

Alkaloid from *Trichocereus* (*Cereus terscheckii*) (Cactaceae). Shows plant

growth inhibiting activity. Tremorogenic and respiratory paralytic agent in rats. Weak hypotensive. No noticeable hallucinogenic props. in humans. Liq. Mp 172° (as picrate).

N-(Carboxymethyl): *N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]glycine. *N*-(Carboxymethyl)mescaline. **Mescaloxyllic acid**

[7738-40-1]

$\text{C}_{13}\text{H}_{19}\text{NO}_5$ 269.297

Trace constit. of peyote cactus *Lophophora williamsii* (*Anhalonium williamsii*) (Cactaceae). Cryst. (EtOH). Mp 187-189°.

N-(1-Carboxyethyl): *N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]alanine. *N*-(1-Carboxyethyl)mescaline. **Mescaloruvic acid**

[7738-43-4]

$\text{C}_{14}\text{H}_{21}\text{NO}_5$ 283.324

Trace constit. of peyote cactus *Lophophora williamsii* (*Anhalonium williamsii*) (Cactaceae). Cryst. (EtOH) (racemate). Mp 235-236.5° (racemate). Opt. rotn. not detd., identity with synthetic racemate detd. chromatographically and by ms.

Heffter, A. *et al.*, *Ber.*, 1896, **29**, 216 (*isol*)

Späth, E. *et al.*, *Ber.*, 1937, **70**, 2446; 1938, **71**, 1275 (*Acetylmescaline*, *Methylmescaline*)

Reti, L. *et al.*, *J.A.C.S.*, 1951, **73**, 1767

(*Trichocereine*)

Bahnholzer, K. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 1577 (*synth*)

Kapadia, G.J. *et al.*, *Chem. Comm.*, 1968, 1688 (*Formylmescaline*)

Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1699 (*rev*)

Lundström, J. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 3489 (*biosynth*)

Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 1172 (*Mescaloxyllic acid*, *Mescaloruvic acid*)

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*occur*, *rev*)

Doetsch, P.W. *et al.*, *J. Chromatogr.*, 1980, **189**, 79 (*occur*)

Van Peteghem, C. *et al.*, *Eur. J. Drug Metab. Pharmacokinet.*, 1982, **7**, 1 (*pharmacol*)

Bailey, K. *et al.*, *Org. Magn. Reson.*, 1983, **21**, 391 (*cmr*)


Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1388

Isobe, T. *et al.*, *Yakugaku Zasshi*, 2002, **122**, 291-294 (*N-Benzoylmescaline*)

Chen, I.-S. *et al.*, *Fitoterapia*, 2007, **78**, 414-419 (*Taiwanamide C*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MDI500; MDI750

Mescaline citrimide M-220




$\text{C}_{17}\text{H}_{21}\text{NO}_8$ 367.355

Alkaloid from mescal (*Lophophora williamsii*) (Cactaceae).

Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1699-1727 (*isol*, *rev*)

Mescaline isocitrimide lac-tone M-221




$\text{C}_{17}\text{H}_{19}\text{NO}_7$ 349.34

Alkaloid from mescal (*Lophophora williamsii*) (Cactaceae).

Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1699-1727

Mescaline succinimide M-222

1-[2-(3,4,5-Trimethoxyphenyl)ethyl]-2,5-pyrrolidinedione. *N*-(3,4,5-Trimethoxyphenethyl)succinimide



$\text{C}_{15}\text{H}_{19}\text{NO}_5$ 293.319

Alkaloids not named in the lit.; not registered by CA. Alkaloid from peyote cactus (*Lophophora williamsii*) (Cactaceae). Mp 125-126°.

3-Hydroxy: **Mescaline malimide**. *N*-(3,4,5-Trimethoxyphenethyl)malimide

$\text{C}_{15}\text{H}_{19}\text{NO}_6$ 309.318

Alkaloid from peyote cactus (*Lophophora williamsii*) (Cactaceae). Opt. rotn. not detd.

3,4-Didehydro: **Mescaline maleimide**. *N*-(3,4,5-Trimethoxyphenethyl)maleimide


$\text{C}_{15}\text{H}_{17}\text{NO}_5$ 291.303

From peyote cactus (*Lophophora williamsii*) (Cactaceae). Poss. artifact.

Kapadia, G.J. *et al.*, *Chem. Comm.*, 1968, 1688 (*occur*, *ms*, *synth*)

Mescengricin M-223

[191409-84-4]




$\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_8$ 428.398

First natural α-carboline. Prod. by *Streptomyces griseoflavus* 2853-SVS4. Neuronal cell protecting agent. L-Glutamate toxicity suppressor. Antioxidant. Sol. MeOH, Me₂CO, CHCl₃, EtOAc. Mp 247-249° dec. $[\alpha]_D^{20}$ -33 (c, 0.04 in MeOH). λ_{max} 210 (ε 10500); 256 (ε 5200); 284 (ε 6400); 400 (ε 4500) (MeOH). λ_{max} 238 (ε

- 10000); 410 (ε 4800) (MeOH/NaOH).
Kim, J.-S. *et al.*, *Tet. Lett.*, 1997, **38**, 3431-3434
(*isol, uv, pmr, cmr*)
Kazuo, S.-Y. *et al.*, *J. Asian Nat. Prod. Res.*,
2000, **2**, 121-132 (*isol, pmr, cmr*)
Martin, G.E. *et al.*, *J. Nat. Prod.*, 2000, **63**,
543-585 (*N-15 nmr*)

Mesembrenol M-224

3a-(3,4-Dimethoxyphenyl)-2,3,3a,6,7,7a-hexahydro-1-methyl-1H-indol-6-ol, 9CI



(+)-form

C₁₇H₂₃NO₃ 289.374

(+)-form [25516-15-8]

Alkaloid from *Sceletium strictum* (Aizoaceae). Prisms (EtOAc or Me₂CO). Mp 140°. [α]_D²⁵ +91 (c, 0.0176 in CHCl₃). λ_{max} 230 (ε 9210); 279 (ε 3515); 284 (ε 2990) (EtOH).

Ac: O-Acetylmesebrenol

[25516-16-9]
C₁₉H₂₅NO₄ 331.411

Alkaloid from *Sceletium strictum* (Aizoaceae). Oil.

N-De-Me: N-Demethylmesebrenol

[34084-31-6]
C₁₆H₂₁NO₃ 275.347

Alkaloid from *Sceletium strictum* (Aizoaceae). Mp 63-65°. [α]_D +86.

O^{4'}-De-Me: 4'-O-Demethylmesebrenol

[25516-14-7]
C₁₆H₂₁NO₃ 275.347

Alkaloid from *Sceletium strictum* (Aizoaceae). Prisms (MeOH). Mp 219-220°. [α]₃₀₀²⁵ +533 (c, 0.083 in EtOH). λ_{max} 225 (log ε 3.44); 280 (log ε 3.44) (EtOH).

6-Ketone: Synthetic. by oxidn. of (+)-Mesembrenol. Oil. CD [θ]₃₃₄ -4060°. V. readily racemised.

4,5-Dihydro, O^{4'}-de-Me: 4'-O-Demethylmesebrenol

[25516-13-6]
C₁₆H₂₃NO₃ 277.363

Alkaloid from *Sceletium strictum* (Aizoaceae). Prisms (MeOH). Mp 201°. [α]₃₀₀²⁵ -199 (c, 0.71 in EtOH). λ_{max} 229 (log ε 3.62); 280 (log ε 3.38) (EtOH).

4,5-Dihydro, 6-ketone: see Mesembrine, M-226

(±)-form

6-Ketone: Mesembrenone. Mesembrinine. Mesembrine

[80287-15-6]
[25516-12-5]

C₁₇H₂₁NO₃ 287.358

Alkaloid from *Sceletium namaquense* and *Sceletium strictum* (Aizoaceae). Mp 88°. Prob. arises by racemisation. The name Mesembrenone is preferred

as more descriptive of the struct.; it was originally called Mesembrenine (Mesembrinine).

6-Ketone, hydrochloride:

Plates (MeOH/Et₂O). Mp 192-193° (190-195°) dec.

6-Ketone, N-de-Me, N-formyl: N-De-methyl-N-formylmesebrenone

[61921-43-5]
C₁₇H₁₉NO₄ 301.341

Alkaloid from *Sceletium strictum* (Aizoaceae).

6-Ketone, O^{4'}-de-Me: 4'-O-Demethylmesebrenone

[51934-30-6]
C₁₆H₁₉NO₃ 273.331

Alkaloid present in *Sceletium strictum* and *Sceletium namaquense* (Aizoaceae). Oil. λ_{max} 220 (log ε 4.04); 280 (log ε 3.59) (95% EtOH).

Popelak, A. *et al.*, *Naturwissenschaften*, 1960, **47**, 231-232 (*Mesembrenone, isol*)

Jeffs, P.W. *et al.*, *J.O.C.*, 1970, **35**, 3512-3515; 1974, **39**, 2703-2710 (*Mesembrenol, Mesembrenone, O-Acetylmesebrenol, 4'-O-demethyl derivs, isol, uv, ir, pmr, ms, struct, synth, cd*)


Kruger, P.E.J. *et al.*, *J. S. Afr. Chem. Inst.*, 1971, **24**, 235-237 (*N-Demethylmesebrenol*)

Martin, N.H. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 1-19 (*ms*)

Karle, J.M. *et al.*, *Acta Cryst. B*, 1977, **33**, 185-188 (*N-Demethyl-N-formylmesebrenone*)

Δ⁷-Mesembrenone**M-225**

3a-(3,4-Dimethoxyphenyl)-1,2,3,3a,4,5-hexahydro-1-methyl-6H-indol-6-one, 9CI



C₁₇H₂₁NO₃ 287.358

(S)-form [35714-44-4]


Alkaloid from *Sceletium namaquense* (Aizoaceae). Oil.

Jeffs, P.W. *et al.*, *J.O.C.*, 1974, **39**, 2703 (*isol, pmr, ir, uv, ord, synth, struct*)

Martin, N.H. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 1 (*ms*)

Mesembrine**M-226**

3a-(3,4-Dimethoxyphenyl)octahydro-1-methyl-6H-indol-6-one, 9CI. *Mesembranone*



(-)-form

C₁₇H₂₃NO₃ 289.374
Sedative. Log P 1.25 (calc).

(-)-form [24880-43-1]

Alkaloid from *Sceletium tortuosum* and *Sceletium namaquense* (Aizoaceae). Liq. Bp_{0.3} 186-190°. [α]_D²⁰ -55.4 (MeOH).

Hydrochloride: Mp 204.5-206° dec. [α]_D -8.8 (c, 0.632 in MeOH).

Deoxo: Mesembrane

[510-80-5]

C₁₇H₂₅NO₂ 275.39

Minor alkaloid from *Sceletium namaquense* (Aizoaceae). Oil.

6α-Alcohol: Mesembranol. Mesembrinol

[23544-42-5]

C₁₇H₂₅NO₃ 291.389

Alkaloid from *Sceletium tortuosum* and *Sceletium strictum* (Aizoaceae), also obt. by redn. of (-)-Mesembrine. Mp 144-145°. [α]_D -32 (CHCl₃).

6α-Alcohol, N-de-Me: N-Demethylmesebrenol

C₁₆H₂₃NO₃ 277.363

Alkaloid from *Sceletium strictum* (Aizoaceae). Mp 175-185°. [α]_D -13.

(+)-form [468-53-1]

Synthetic. Pale yellow oil. [α]_D²⁰ +16.1 (c, 1.32 in MeOH).

Hydrochloride:

Needles (2-propanol). Mp 208-210.5°. [α]_D²⁰ +7.3 (c, 0.465 in MeOH).

(±)-form [6023-73-0]

Synthetic. Oil. Bp_{0.07} 178°.

Hydrochloride: Mp 179-181°.

Picrate:

Cryst. (EtOH/EtOAc). Mp 171.5-172.5°.

Popelak, A. *et al.*, *Naturwissenschaften*, 1960, **47**, 156; 231 (*struct*)

Smith, E. *et al.*, *Chem. Ind. (London)*, 1961, 402 (*Mesembranol*)

Stevens, R.V. *et al.*, *J.A.C.S.*, 1968, **90**, 5580 (*synth, ir, pmr*)

Keely, S.L. *et al.*, *J.A.C.S.*, 1968, **90**, 5584 (*synth*)

Jeffs, P.W. *et al.*, *J.A.C.S.*, 1969, **91**, 3831 (*abs config*)

Oh-ishi, T. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 299 (*synth, ir, pmr*)

Otani, G. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 2130 (*synth, cd, ord, abs config*)

Capps, T.M. *et al.*, *J.C.S. Perkin 2*, 1977, 1098 (*isol, uv, pmr, ms, cryst struct, Mesembrane*)

Jeffs, P.W. *et al.*, *Phytochemistry*, 1978, **17**, 719 (*biosynth*)

Martin, S.F. *et al.*, *J.O.C.*, 1979, **44**, 3391 (*synth*)

Strauss, H.F. *et al.*, *Tet. Lett.*, 1979, 4495 (*synth*)

Takano, S. *et al.*, *Tet. Lett.*, 1981, 4479 (*synth*)

Keck, G.E. *et al.*, *J.O.C.*, 1982, **47**, 1302 (*synth*)

Jeffs, P.W. *et al.*, *J.O.C.*, 1983, **48**, 3861 (*synth*)

Sánchez, I.H. *et al.*, *Tet. Lett.*, 1983, **24**, 551 (*synth*)

Howard, A.S. *et al.*, *Tet. Lett.*, 1983, **24**, 829 (*synth*)

Kochhar, K.S. *et al.*, *Tet. Lett.*, 1983, **24**, 4785 (*synth*)

Meyers, A.I. *et al.*, *J.A.C.S.*, 1985, **107**, 7776 (*synth*)

Gramain, J.-C. *et al.*, *Tet. Lett.*, 1985, **26**, 4083 (*synth*)

Hackett, S. *et al.*, *J.O.C.*, 1986, **51**, 1629 (*synth*)

- Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2734 (*synth*, *ir*, *pmr*)
 Winkler, J.D. *et al.*, *J.A.C.S.*, 1988, **110**, 4831 (*synth*)
 Shono, T. *et al.*, *Chem. Lett.*, 1989, 1963 (*synth*)
 Takano, S. *et al.*, *Chem. Lett.*, 1990, 1239 (*synth*)
 Bauermeister, S. *et al.*, *J.C.S. Perkin 1*, 1991, 561 (*synth*)
 Parkinson, C.J. *et al.*, *J.C.S. Perkin 1*, 1991, 1053 (*synth*)
 Ishibashi, H. *et al.*, *J.O.C.*, 1991, **56**, 95 (*synth*, *Mesembranol*)
 Gelas-Mialhe, Y. *et al.*, *Heterocycles*, 1992, **34**, 37 (*synth*)
 Michael, J.P. *et al.*, *Tet. Lett.*, 1992, **33**, 6023 (*synth*)
 Yokomatsu, T.Y. *et al.*, *Tet. Lett.*, 1992, **33**, 6999 (*synth*)
 Matsumura, Y. *et al.*, *Tetrahedron*, 1993, **49**, 8503 (*synth*)
 Kosugi, H. *et al.*, *Tetrahedron: Asymmetry*, 1993, **4**, 1409 (*synth*)
 Chida, N. *et al.*, *Chem. Comm.*, 1994, 901 (*synth*, *Mesembranol*)
 Nemoto, H. *et al.*, *J.O.C.*, 1995, **60**, 6785 (*synth*)
 Mori, M. *et al.*, *J.O.C.*, 1997, **62**, 3263 (*synth*)
 Kamikubo, T. *et al.*, *Chem. Comm.*, 1998, 783-784 (*synth*)
 Yamada, O. *et al.*, *Tet. Lett.*, 1998, **39**, 7747-7750 (*synth*)
 Dalko, P.I. *et al.*, *Tet. Lett.*, 1998, **39**, 8979-8982 (*synth*)
 Rigby, J.H. *et al.*, *Org. Lett.*, 2000, **2**, 1673-1675 (*synth*)
 Everts, J.B. *et al.*, *Tet. Lett.*, 2001, **42**, 3673-3675 (*Mesembranol*, *synth*)
 Hayashi, M. *et al.*, *Tet. Lett.*, 2002, **43**, 1461-1464 (*synth*)
 Kulkarni, M.G. *et al.*, *Tet. Lett.*, 2002, **43**, 2297-2298 (*synth*)
 Chavan, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 5263-5265 (*synth*)
 Taber, D.F. *et al.*, *J.O.C.*, 2005, **70**, 7711-7714 (*synth*)
 Paul, T. *et al.*, *Org. Lett.*, 2006, **8**, 4007-4010 (*synth*)
 Arns, S. *et al.*, *J.O.C.*, 2007, **72**, 9314-9322 (*synth*)
 Saito, M. *et al.*, *Tetrahedron*, 2007, **63**, 4865-4873 (*Mesembrane*, *synth*)
 Roe, C. *et al.*, *Tet. Lett.*, 2008, **49**, 650-653 (*synth*)


Mesembryanthemins M-227

Glycosides of Betanidin or Isobetanidin; see Betanidin, B-110. Structs. unknown. Pigments from *Mesembryanthemum conspicuum*, *Mesembryanthemum edule*, *Mesembryanthemum floribundum* and *Portulaca grandiflora* (Aizoaceae, Portulacaceae). Mesembryanthemins I-III recognised.

Piattelli, M. *et al.*, *Phytochemistry*, 1964, **3**, 547-557 (*occur*)

Metachromin G M-228

[143592-24-9]



R = -NHCH₂CH₂Ph

C₂₉H₃₇NO₃ 447.616
 Constit. of *Hippospongia metachroma*. Purple oil. [α]_D²⁰ -18 (c, 0.2 in C₆H₆). λ_{max} 211 (ε 29700); 324 (ε 14700); 508 (ε 800) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *J.O.C.*, 1992, **57**, 5773 (*isol*, *pmr*, *cmr*)

Metachromin H M-229

[143592-25-0]

As Metachromin G, M-228 with

R = -NHCH₂CH₂CH(CH₃)₂

C₂₆H₃₉NO₃ 413.599


Constit. of *Hippospongia metachroma*. Purple oil. [α]_D¹⁹ -9 (c, 0.2 in C₆H₆). λ_{max} 324 (ε 12800); 512 (ε 700) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *J.O.C.*, 1992, **57**, 5773 (*isol*, *pmr*, *cmr*)

Metacycloprodigiosin M-230

Streptorubin A. NSC 76779

[22869-99-4]



C₂₅H₃₃N₃O 391.555

Pigment from *Streptomyces longisporus ruber*. Orange-brown cryst. (petrol). Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O. Mp 208-209°. λ_{max} 273 (ε 3640); 297 (ε 6900); 362 (ε 5000); 500 (sh) (ε 31500); 530 (ε 75900) (MeOH/HCl) (Derep). λ_{max} 285 (ε 6050); 325 (ε 4850); 467 (ε 30600) (MeOH/KOH) (Derep). λ_{max} 273 (ε 3640); 297 (ε 6900); 362 (ε 5000); 500 (sh) (ε 31500); 530 (ε 75900) (hydrochloride) (Derep).

Hydrochloride: Mp 214-216°.

Wassermann, H.H. *et al.*, *J.A.C.S.*, 1969, **91**, 1263; 1264 (*struct*, *synth*)


Wassermann, H.H. *et al.*, *Tetrahedron*, 1976, **32**, 1855; 1867

Fürstner, A. *et al.*, *J.O.C.*, 1999, **64**, 8281-8286 (*synth*)

Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (*rev*)

Metacytofilin M-231

3-Hydroxy-6-(methylamino)-6-(2-methylpropyl)-3-(phenylmethyl)-2,5-morpholinedione, 9CI
 [145398-57-8]



Relative configuration

C₁₆H₂₂N₂O₄ 306.361


Prod. by *Metarhizium* sp. TA 2759. Immunomodulator. Prisms (MeOH aq.). Sol. MeOH, DMSO; poorly sol. CHCl₃,

H₂O. Mp 201-202°. [α]_D²⁷ +5.1 (c, 0.63 in MeOH). λ_{max} 252 (sh) (ε 182); 258 (sh) (ε 200); 264 (sh) (ε 151) (MeOH) (Derep).

Iijima, M. *et al.*, *J. Antibiot.*, 1992, **45**, 1553 (*isol*, *struct*, *props*)

Metanicotine M-232

N-Methyl-4-(3-pyridinyl)-3-buten-1-amine, 9CI. 3-[4-(Methylamino)-1-butenyl]pyridine, 8CI. **Rivanicline**, INN. RJR 2403. TC 2403
 [538-79-4]



C₁₀H₁₄N₂ 162.234

Alkaloid from leaves of *Duboisia hopwoodii* (Solanaceae). Also a pyrrol. prod. of nicotine alkaloids present in tobacco smoke. CNS selective nicotinic agonist. Potential agent for the treatment of Alzheimer's. Used in the treatment of ulcerative colitis. Bp_{3,7} 141°. (*E*)-form is pharmacol. active.

▶ Teratogen. UT5550000

Picrate: Mp 173-175°.

Galactarate: **Rivanicline galactarate**, **USAN**
 [675132-86-2]

[1129-68-6, 183288-99-5, 15585-43-0, 180915-56-4]

Swain, M.L. *et al.*, *J.A.C.S.*, 1949, **71**, 1341 (*synth*, *wv*)

Luanratana, O. *et al.*, *Phytochemistry*, 1982, **21**, 449 (*occur*, *ms*)

Bencherif, M. *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, **279**, 1413-1429; 1422-1429 (*pharmacol*)

Pat. Coop. Treaty (WIPO), 1996,

((*Reynolds*))96 20 599 (*synth*, *pharmacol*)


Bencherif, M. *et al.*, *CNS Drug Rev.*, 1997, **3**, 325-345 (*rev*, *pharmacol*)

Papke, R.L. *et al.*, *J. Pharmacol. Exp. Ther.*, 2002, **301**, 765-773 (*pharmacol*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MDM750

Metaphanine M-233

8,10-Epoxy-8-hydroxy-3,4-dimethoxy-17-methylhasubanan-7-one, 9CI



(-)-form

C₁₉H₂₃NO₅ 345.394

(-)-form [1805-86-3]

Alkaloid from the roots and stems of *Stephania japonica*. Also isol. from *Stephania abyssinica* and *Stephania japonica* var. *australis* (Menispermaceae). Cryst.

(Me₂CO/CHCl₃). Mp 233°. [α]_D²⁵ -21 (c, 1.0 in CHCl₃). pK_a 6.03.

Hydrochloride:

Cryst. + ½ H₂O (Me₂CO). Mp 226°.

Hydrotide:

Cryst. + ½ H₂O (THF). Mp 230° dec.

O⁴-De-Me: Stephabyssine. O⁴-Demethyl-metaphanine

[36871-84-8]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the roots and rhizomes of *Stephania abyssinica* and roots and stems of *Stephania longa*. Also isol. from *Stephania japonica* var. *australis* (Menispermaceae). Cryst. (EtOH aq.). Mp 178-180°. [α]_D -58.9 (c, 0.87 in CHCl₃).

O⁴-De-Me, hydrochloride:

Plates (CHCl₃/CCl₄). Mp 247-250° dec. [α]_D²⁵ -32.5 (c, 0.41 in 60% EtOH).

7 α -Alcohol, O⁴-de-Me: Isostephaboline

[863712-36-1]

C₁₈H₂₃NO₅ 333.383

Alkaloid from *Stephania longa*. Powder. [α]_D²⁵ +26.7 (c, 0.3 in MeOH). λ_{\max} 226 (log ϵ 3.63); 285 (log ϵ 3.23) (MeOH).

7 β -Alcohol, O⁴-de-Me: Stephaboline

[36871-86-0]

C₁₈H₂₃NO₅ 333.383

Alkaloid from the roots and rhizomes of *Stephania abyssinica* and the roots and stems of *Stephania longa* (Menispermaceae). Cryst. (MeOH aq.). Mp 186-188° dec. [α]_D²⁵ +34.7 (c, 0.47 in MeOH).

7 β -Alcohol, O⁴-de-Me, methiodide:

Cryst. (MeOH/EtOAc). Mp 230-232°.

[α]_D²⁵ +19.6 (c, 0.51 in MeOH).

Me enol ether: see Prometaphanine, P-639

(±)-form [38305-47-4]

Synthetic. Prisms (Et₂O/MeOH). Mp 205-206°.

Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 695 (*ir, pmr, ms, struct*)

De Waal, H.L. *et al.*, *Tet. Lett.*, 1966, 6169 (*uv, ir, pmr, struct*)

Kupchan, S.M. *et al.*, *J.O.C.*, 1973, **38**, 151

(*Stephaboline, Metaphanine, isol, synth*)

Ibuka, T. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 907 (*synth, ir, pmr*)


Matsui, M. *et al.*, *Phytochemistry*, 1979, **18**, 1087 (*Stephabyssine, isol*)

Zhang, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1201-1207 (*Stephaboline, Isostephaboline, Stephabyssine*)

Metenaticin C

M-234

[866954-60-1]



C₂₂H₂₃NO₁₀ 461.424

Prod. by *Streptomyces violaceoruber* Tü22.

15-O-(2,3,6-Trideoxy-β-L-threo-hexopyranoside): Metenaticin B

[866954-24-7]

C₂₈H₃₃NO₁₂ 575.568


Prod. by *Streptomyces violaceoruber* Tü22.

Pham, L.H. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 710-723 (*isol, uv, pmr, ms*)

Methanopterin

M-235

[79484-89-2]



C₃₀H₄₁N₆O₁₆P 772.658

Prod. by *Methanobacterium thermoautotrophicum*.

Van Beelen, P. *et al.*, *Eur. J. Biochem.*, 1984, **138**, 563 (*isol, struct*)

Keltjens, J.T. *et al.*, *Methods Enzymol.*, 1986, **122**, 412 (*rev*)

White, R.H. *et al.*, *Biochemistry*, 1990, **29**, 5397 (*biosynth*)

White, R.H. *et al.*, *Chirality*, 1996, **8**, 332 (*cd, abs config*)


Methoxatin

M-236

4,5-Dihydro-4,5-dioxo-1H-pyrrolo[2,3-f]quinoline-2,7,9-tricarboxylic acid.

Coenzyme PQQ. PQQ. Pyrroloquinoline-quinone

[72909-34-3]



C₁₄H₆N₂O₈ 330.21

Coenzyme of several bacterial alcohol dehydrogenases. Believed to be a mammalian redox-cofactor vitamin (B group). Oxidising agent. e.g. for amines and amino acids. Dark red solid. Sol. H₂O. λ_{\max} 250 ; 332 (H₂O) (Berdy).

Tri-Me ester: [74447-88-4]

Orange-red solid. Mp 260-263° (>250°).

[122628-50-6]

Anthony, C. *et al.*, *Biochem. J.*, 1967, **104**, 960 (*isol*)

Corey, E.J. *et al.*, *J.A.C.S.*, 1981, **103**, 5599 (*synth*)

Gainor, J.A. *et al.*, *J.O.C.*, 1982, **47**, 2833 (*synth*)

Buechi, G. *et al.*, *J.A.C.S.*, 1985, **107**, 5555 (*synth*)

Hendrickson, J.B. *et al.*, *J.O.C.*, 1985, **50**, 1688 (*synth*)

Itoh, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 1907 (*bibl*)

Duine, J.A. *et al.*, *FEMS Microbiol. Rev.*, 1986, **32**, 165 (*rev*)

MacKenzie, A.R. *et al.*, *Tetrahedron*, 1986, **42**, 3259 (*synth, bibl*)

Ameyama, M. *et al.*, *BioFactors*, 1988, **1**, 51 (*rev*)

Flückiger, R. *et al.*, *Biochem. Biophys. Res. Commun.*, 1988, **153**, 353-358 (*occur*)

Houen, G. *et al.*, *Tetrahedron*, 1989, **45**, 4235 (*use*)

Kumazawa, T. *et al.*, *Biochim. Biophys. Acta*, 1992, **1156**, 62-66 (*occur*)

Itoh, S. *et al.*, *J.C.S. Perkin 2*, 1992, 1245 (*uv, bibl*)

Martin, P. *et al.*, *Helv. Chim. Acta*, 1993, **76**, 988; 1996, **79**, 658 (*synth*)

Itoh, S. *et al.*, *Chem. Comm.*, 1995, 2077 (*ester, synth, pmr, cmr, uv*)

Itoh, S. *et al.*, *Nat. Prod. Rep.*, 1995, **12**, 45-53 (*rev*)


Kasahara, T. *et al.*, *Nature (London)*, 2003, **422**, 832 (*vitamin*)

Magnusson, O.T. *et al.*, *J.A.C.S.*, 2004, **126**, 5342-5343 (*biosynth*)

12-Methoxyaffinisine

M-237

[457613-93-3]



C₂₁H₂₆N₂O₂ 338.449

Alkaloid from the bark of *Rauwolfia bahiensis*. Amorph. yellow solid. [α]_D²⁰ +3 (c, 0.72 in CHCl₃). λ_{\max} 274 (log ϵ 3.25) (MeOH).

17-Aldehyde: 12-Methoxy-N⁴-methylvellosimine

[457613-91-1]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from the bark of *Rauwolfia bahiensis*. Amorph. yellow solid. [α]_D²⁰ +14 (c, 0.6 in CHCl₃). λ_{\max} 284 (log ϵ 3.1) (MeOH).

17-Aldehyde, N-de-Me: 12-Methoxyvellosimine

[457613-94-4]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from the bark of *Rauwolfia bahiensis*. Amorph. yellow solid. [α]_D²⁰ +5 (c, 0.6 in CHCl₃). λ_{\max} 230 (log ϵ 4.2); 254 (log ϵ 3.78) (MeOH).

17-Carboxylic acid, N⁴-Me, Et ester:

Fuchsiaeoline

[109269-75-2]

C₂₄H₃₁N₂O₅[±] 395.52


Quaternary alkaloid from the stem bark of *Peschiera fuchsiaeifolia*. Oil. [α]_D²⁵ -55.7 (c, 0.9 in MeOH). Counterion not specified. λ_{\max} 225 (log ϵ 4.66); 269 (log ϵ 3.9); 283 (log ϵ 3.8); 293 (log ϵ 3.73) (EtOH).

Braga, R.M. *et al.*, *Phytochemistry*, 1987, **26**, 833-836 (*Fuchsiaeoline*)

Kato, L. *et al.*, *Phytochemistry*, 2002, **60**, 315-320 (*isol, pmr, cmr, ms*)
 Zhou, H. *et al.*, *J.O.C.*, 2006, **71**, 251-259 (*synth*)

2-Methoxyatherosperminine M-238

2,3,4-Trimethoxy-N,N-dimethyl-1-phenanthreneethanamine, 9CI. 1-(2-Dimethylaminoethyl)-2,3,4-trimethoxyphenanthrene
 [5635-94-9]



C₂₁H₂₅NO₃ 339.433

Aporphine-type numbering shown. Alkaloid from the bark of *Atherosperma moschatum* and from the root and stem bark of *Meiocarpidium lepidotum* (Monimiaceae, Annonaceae). Oil. Rapidly darkens upon exp. to air.

Picrate:

Cryst. (Me₂CO/MeOH). Mp 161-162°. N-Me: Mp 243-245° dec. (as iodide).

N-Oxide: 2-Methoxyatherosperminine N-oxide

[67341-54-2]

C₂₁H₂₅NO₄ 355.433

Alkaloid from the root and stem bark of *Meiocarpidium lepidotum* (Annonaceae). Mp 189° (as picrate).

O¹,O³-Di-de-Me: *Stipitatine*

C₁₉H₂₁NO₃ 311.38

Alkaloid from *Unonopsis stipitata* (Annonaceae).

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1965, **18**, 1997 (*isol, uv, ir, pmr, struct, synth*)


Leboeuf, M. *et al.*, *Plant. Med. Phytother.*, 1977, **11**, 284 (*isol, uv, pmr, ms, struct, deriv*)

El-Tohami, M. *et al.*, *Chem. Biol. Isoquinoline Alkaloids, Int. Symp., Phytochem. Soc. Eur., Abstr. Pap.*, 1984, 17 (*Stipitatine*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1988, **51**, 389; 439 (*Stipitatine*)

10-Methoxybenzimidazo[6,7,1-def][1,6]naphthyridine, 9CI M-239

[597553-94-1]



C₁₃H₉N₃O 223.234


Alkaloid from a *Xestospongia* sp. Brown

gum. λ_{max} 209 (ε 5845); 237 (ε 5250); 258 (ε 2980); 269 (ε 2770); 355 (ε 2015) (EtOH).

Calcul, L. *et al.*, *Tetrahedron*, 2003, **59**, 6539-6544 (*isol, pmr, cmr*)

4-Methoxybenzoic acid, 9CI M-240

p-Anisic acid, 8CI. Umbellinic acid. Draconic acid. Badianic acid. FEMA 3945
 [100-09-4]
 [1335-08-6]



C₈H₈O₃ 152.149

Isol. from aniseed *Pimpinella anisum* (Apiaceae) and higher plants. Flavouring agent. Needles or prisms. Mp 184°. Bp 275-280°. pK_a 4.5 (25°).

▶ LD₅₀ (mus, scu) 400 mg/kg. BZ4395000

Amide: 4-Methoxybenzamide. Anisamide
 [3424-93-9]


C₈H₉NO₂ 151.165

Constit. of *Naravelia zeylanica*. Needles or tablets (H₂O). Mp 163°. Bp 295°.

Jaroszewski, J.W. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 291-294 (*amide, isol*)

8-Methoxy-9H-benzo[de][1,6]naphthyridin-9-one, 9CI M-241

Demethoxyaaptamine
 [88839-99-0]



C₁₂H₈N₂O₂ 212.207

Alkaloid from the Okinawan marine sponge *Aaptos aaptos* and a *Xestospongia* sp. Cytotoxic. Exhibits potent antimicrobial activity against gram-positive and -negative bacteria. Fine bright yellow rods (EtOAc). Mp 210-212° (198-200° dec.). λ_{max} 235 (ε 10600); 308 (ε 1740); 360 (sh) (ε 5540); 373 (ε 5040); 415 (sh) (ε 2310) (H₂O) (Derep).

Di-Me acetal: 8,9,9-Trimethoxy-9H-benzo[de][1,6]naphthyridine

C₁₄H₁₄N₂O₃ 258.276

Alkaloid from a *Xestospongia* sp. Brown gum. λ_{max} 205 (ε 7045); 229 (ε 11330); 350 (ε 4560) (EtOH).

Nakamura, H. *et al.*, *J.C.S. Perkin 1*, 1987, 173 (*isol, uv, ir, pmr, cmr, struct*)

Pelletier, J.C. *et al.*, *J.O.C.*, 1987, **52**, 616


(*synth, ir, pmr*)

Coutinho, A.F. *et al.*, *Heterocycles*, 2002, **57**, 1265-1272 (*isol, pmr, cmr*)

Calcul, L. *et al.*, *Tetrahedron*, 2003, **59**, 6539-6544 (*isol, pmr, cmr*)

1-(4-Methoxybenzoyl)-6,7-methylenedioxyisoquinoline Pulcheotine A M-242

[17656-61-0]



C₁₈H₁₃NO₄ 307.305

Alkaloid from the bark of *Ocotea pulchella* (Lauraceae). Cryst. (Me₂CO). Mp 152-153°.

α-Alcohol: 1-(*α*-Hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline

[151590-38-4]

C₁₈H₁₅NO₄ 309.321

Alkaloid from bark of *Ocotea pulchella* (Lauraceae). Cryst. (Me₂CO). Mp 138-139°. [α]_D²⁵ +125 (c, 1.2 in CHCl₃).

Unstable; converted spontaneously to the *α*-ketone in air.

1,2-Dihydro: 1,2-Dihydro-1-(4-methoxybenzoyl)-6,7-methylenedioxyisoquinoline

[151590-39-5]

C₁₈H₁₅NO₄ 309.321

Alkaloid from bark of *Ocotea pulchella* (Lauraceae). Unstable in air.

1,2-Dihydro, *α*-alcohol: 1,2-Dihydro-1-(*α*-hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline

[151590-40-8]

C₁₈H₁₇NO₄ 311.337

Alkaloid from bark of *Ocotea pulchella* (Lauraceae). Unstable in air.


Marsaioli, A.J. *et al.*, *Phytochemistry*, 1980, **19**, 995 (*cmr*)

Botega, C. *et al.*, *Phytochemistry*, 1993, **32**, 1331 (*isol, ir, pmr, cmr, ms, struct*)

Wakchaure, P.B. *et al.*, *Synthesis*, 2008, 2321-2322 (*synth, ir, pmr, cmr*)

1-Methoxyberberine M-243

[29133-52-6]



C₂₁H₂₀NO[⊕] 366.393


Alkaloid from *Dactylicapnos torulosa* (Papaveraceae). Yellow needles (EtOAc/MeOH) (as chloride). Mp 185-190° dec. (chloride). CAS no. refers to chloride.

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 1440 (*synth*)

Zhang, G.L. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 604; *CA*, **114**, 160667y (*isol*)

4-Methoxy-2,2'-bipyrrole-5-carboxaldehyde M-244

5-Formyl-4-methoxy-2,2'-bipyrrole.
Tambjamine aldehyde
[10476-41-2]



C₁₀H₁₀N₂O₂ 190.201


Isol. from the nembrothid nudibranchs *Roboastra tigris*, *Tambja eliora*, *Tambja abdere* and the bryozoan *Sessibugula translucens*. Cryst. (EtOH aq.). Mp 263-265° dec. Artifact. λ_{max} 251 (ε 14700); 361 (ε 40300) (MeOH).

Rapoport, H. *et al.*, *J.A.C.S.*, 1962, **84**, 635-642 (*synth*, *uv*)

Carté, B. *et al.*, *J.O.C.*, 1983, **48**, 2314-2318 (*isol*, *uv*, *ir*, *pmr*)

Methoxybrassinin A M-245

[142449-74-9]




C₁₃H₁₆N₂O₂S₂ 280.414

Stress metabolite isol. from cabbage (*Brassica oleracea* var. *capitata*) inoculated with *Pseudomonas cichorii*. Phytoalexin. Gum. λ_{max} 205 (ε 25200); 220 (ε 23800); 290 (ε 3800) (MeOH) (Berdy).

Monde, K. *et al.*, *Phytochemistry*, 1991, **30**, 3921-3922 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Methoxybrassinin B M-246

[142449-75-0]




C₁₃H₁₄N₂O₂S₂ 294.398

Stress metabolite isol. from cabbage (*Brassica oleracea* var. *capitata*) inoculated with *Pseudomonas cichorii*. Mp 73-74°.

Monde, K. *et al.*, *Phytochemistry*, 1991, **30**, 3921-3922 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

37-(Methoxycarbonyl)dictyodendrine E M-247

[149444-92-8]



C₄₃H₃₂N₂O₁₁S 784.799

Alkaloid from the marine sponge *Dic-*

tyodendrilla sp. Potent aldose reductase inhibitor. Antidiabetic agent. Purple solid. Sol. MeOH, EtOAc. Mp 300°. λ_{max} 231 (ε 41000); 281 (ε 21000); 482 (ε 23200) (EtOH) (Derep).

De(methoxycarbonyl): **Dictyodendrine E**
[510709-72-5 (Na salt)]
C₄₁H₃₀N₂O₉S 726.762
Alkaloid from *Dictyodendrilla verongiformis*. Amorph. red solid (as Na salt). λ_{max} 230 (ε 33900); 280 (ε 17700); 464 (ε 20600) (MeOH) (Na salt).


Sato, A. *et al.*, *J.O.C.*, 1993, **58**, 7632-7634 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Warabi, K. *et al.*, *J.O.C.*, 2003, **68**, 2765-2770 (*Dictyodendrin E*)

Fürstner, A. *et al.*, *J.A.C.S.*, 2006, **128**, 8087-8094 (*synth*)

5-Methoxy-2,6-dimethyl-2H-isoindole-4,7-dione, 9CI M-248

2,5-Dimethyl-6-methoxy-4,7-isoindolequinone
[79664-59-8]



C₁₁H₁₁NO₃ 205.213

Found in *Reniera* sp. Active against gram-positive bacteria and marine pseudomonads. Mp 153-154° Mp 167-168° (synthetic). First naturally-occurring isoindole. λ_{max} 224 (ε 13600); 233 (ε 12000); 275 (ε 11000); 362 (ε 3030) (MeOH) (Derep).


Frincke, J.M. *et al.*, *J.A.C.S.*, 1982, **104**, 265 (*isol*, *struct*)

Parker, K.A. *et al.*, *Tet. Lett.*, 1984, **25**, 4917 (*synth*)

Padwa, A. *et al.*, *J.O.C.*, 1985, **50**, 4006 (*synth*)
Schubert-Zsilavecz, M. *et al.*, *Annalen*, 1991, 973 (*synth*)

7-Methoxy-1,6-dimethyl-5,8-isoquinolinedione, 9CI M-249

[79664-58-7]



C₁₂H₁₁NO₃ 217.224

Isol. from an unidentified *Reniera* sponge and an undescribed *Xestospongia* sp. Shows insecticidal activity and weak activity against gram-positive bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 188-190° dec. λ_{max} 223; 276; 283; 292 (MeOH) (Derep).

Frincke, J.M. *et al.*, *J.A.C.S.*, 1982, **104**, 265 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)


Edrada, R.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 973 (*isol*, *props*)

Molina, P. *et al.*, *Synthesis*, 1997, 963-966 (*synth*)

Kuwabara, N. *et al.*, *Tetrahedron*, 2004, **60**, 2943-2952 (*synth*)

4-Methoxy-7,8-dimethyl-2H-pyranol[4,3-b]pyridin-2-one, 9CI M-250

Acuminatopyrone
[135038-52-7]



C₁₁H₁₁NO₃ 205.213


Struct. revised in 1994. Metab. of *Fusarium acuminatum*, *Fusarium chlamydosporum* and *Fusarium trincinum*. Prisms (C₆H₆ or EtOAc). Mp 203-204° dec.

Grove, J.F. *et al.*, *J.C.S. Perkin 1*, 1991, 997 (*isol*)

Visconti, A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 695 (*struct*)

5-Methoxy-2,2-dimethyl-2H-pyranol[2,3-b]quinoline, 9CI M-251

[88148-18-9]



C₁₅H₁₅NO₂ 241.289

Alkaloid from *Evodia pilulifera*. Needles (C₆H₆/petrol). Mp 69-70°.

3,4-Dihydro: 3,4-Dihydro-5-methoxy-2,2-dimethyl-2H-pyranol[2,3-b]quinoline, 9CI
[42522-05-4]

C₁₅H₁₇NO₂ 243.305

From *Evodia pilulifera*. Prisms (petrol). Mp 95-96°.

8,9-Dimethoxy: 5,8,9-Trimethoxy-2,2-dimethyl-2H-pyranol[2,3-b]quinoline, 9CI
[131524-75-9]

C₁₇H₁₉NO₄ 301.341

From *Evodia pilulifera*.

8,9-Dimethoxy, 3,4-dihydro: 3,4-Dihydro-5,8,9-trimethoxy-2,2-dimethyl-2H-pyranol[2,3-b]quinoline, 9CI
[131524-74-8]

C₁₇H₂₁NO₄ 303.357

From *Evodia pilulifera*.

3,4-Dihydro, 3-hydroxy: see Geibalansine, G-29


Bowman, R.M. *et al.*, *J.C.S. Perkin 1*, 1973, 1055 (*synth*)

Ramesh, M. *et al.*, *Heterocycles*, 1984, **22**, 125 (*synth*)

Goh, S.H. *et al.*, *Bull. Singapore Natl. Inst. Chem.*, 1989, **17**, 21; *C.A.*, **114**, 39186n (*isol*)

9-Methoxyellipticine M-252

9-Methoxy-5,11-dimethylpyrido[4,3-b]carbazole
[10371-86-5]



C₁₈H₁₆N₂O 276.337

Numbering systems vary. Alkaloid from *Ochrosia maculata*, *Ochrosia oppositifolia*, *Ochrosia elliptica*, *Strychnos dinklagei*, *Excavatia coccinea* (preferred genus name *Ochrosia*), *Bleekeria coccinea* (preferred genus name *Ochrosia*) and *Bleekeria vitiensis*. Powerful inhibitor of DNA synthesis (intercalating agent). Orally active antineoplastic agent. Antiparasitic. Mp 286-287° dec. Log P 4.32 (calc). λ_{max} 211 (ε 17000); 245 (ε 20400); 276 (ε 35500); 291 (ε 41700); 337 (ε 5130); 354 (ε 2820); 403 (ε 2750) (MeOH) (Derep). λ_{max} 211 ; 227 ; 237 ; 245 ; 275 ; 276 ; 286 ; 291 ; 295 ; 332 ; 337 ; 346 ; 354 ; 384 ; 403 (MeOH) (Berdy). λ_{max} 242 (ε 22400); 275 (ε 38600); 290 (ε 45700); 335 (ε 5130) (EtOH) (Berdy).

► LD₅₀ (mus, ipr) 150 mg/kg. UU8840000
Hydrochloride: Mp 288-290° (cryst. transition at 255-265°).

Picrate: Mp 273-275°.

O-De-Me: 9-Hydroxyellipticine

[51131-85-2]

C₁₇H₁₄N₂O 262.31

Alkaloid from the stem bark of *Strychnos dinklagei* and *Ochrosia* sp. Strongly active against gram-positive and gram-negative bacteria. Orange prisms (CHCl₃/MeOH). Mp > 300°. λ_{max} 225 (log ε 3.85); 245 (log ε 4.03); 263 (log ε 4.04); 278 (log ε 4.01); 317 (log ε 4.36); 356 (log ε 3.37); 366 (log ε 3.46) (EtOH/HCl). λ_{max} 260 (log ε 4.23); 272 (sh) (log ε 4.17); 298 (log ε 4.19); 311 (log ε 4.27) (EtOH/KOH). λ_{max} 245 (log ε 4.07); 277 (log ε 4.28); 294 (log ε 4.35); 310 (sh) (log ε 4.12); 338 (log ε 3.53); 352 (log ε 3.26); 400 (log ε 3.19) (EtOH).

► LD₅₀ (mus, ipr) 68 mg/kg. Exp. reprod. and teratogenic effects. UU8886500

1,2-Dihydro: 1,2-Dihydro-9-methoxyellipticine

[41628-51-7]

C₁₈H₁₈N₂O 278.353

Alkaloid from the leaves of *Ochrosia balansae* (Apocynaceae).

1,2,3,4-Tetrahydro: 1,2,3,4-Tetrahydro-9-methoxyellipticine

[41628-52-8]

C₁₈H₂₀N₂O 280.369

Alkaloid from the leaves of *Ochrosia balansae* (Apocynaceae).

Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 1903 (isol, uv)

Dalton, L.K. *et al.*, *Aust. J. Chem.*, 1967, **20**, 2715 (synth, pmr)

Bruneton, J. *et al.*, *Ann. Pharm. Fr.*, 1972, **30**, 629 (derivs)

Michel, M.M.G. *et al.*, *Rev. Inst. Pasteur Lyon*, 1975, **8**, 53-68 (9-Hydroxyellipticine, activity)

Rousselle, D. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1977, **284**, 377 (synth)

Ahond, A. *et al.*, *Tetrahedron*, 1978, **34**, 2385 (cmr)

Michel, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 489-494 (9-Hydroxyellipticine, isol, uv)


Miller, R.B. *et al.*, *J.O.C.*, 1983, **48**, 886 (synth, pmr)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 116 (rev, pharmacol)

Gribble, G.W. *et al.*, *J.O.C.*, 1992, **57**, 5891 (synth)


Plug, J.P.M. *et al.*, *Synthesis*, 1992, 1221 (9-Hydroxyellipticine, synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MEL775

1-Methoxy-8-hydroxy-methyl-3H-phenoxazin-3-one M-253
HalxazoneC₁₄H₁₁NO₄ 257.245

Prod. by *Streptomyces halstedii* 4029-SVS1. Neuronal cell protectant. Mp 201-202°. λ_{max} 204 (ε 18300); 254 (ε 13500); 365 (ε 10000); 456 (ε 4700) (MeOH).

Katoh, H. *et al.*, *J. Antibiot.*, 2002, **55**, 508-510 (isol, pmr, cmr)

5-Methoxy-1H-indole-4,7-dione, 9CI M-254
[17606-06-3]C₉H₇NO₃ 177.159


Alkaloid from the muricid gastropod *Drupella fragum*. Antimicrobial agent. Orange needles. Mp 198-200° (natural) Mp 205-208° (synthetic). λ_{max} 222 (ε 18200); 281 (ε 19400); 324 (ε 25400); 432 (ε 1120) (EtOH).

Fukuyama, Y. *et al.*, *Tetrahedron*, 1998, **54**, 10007-10016 (isol, synth, ir, uv, pmr, cmr)

6-Methoxy-1H-indole-4,7-dione, 9CI M-255
[137542-70-2]C₉H₇NO₃ 177.159


Alkaloid from the muricid gastropod *Drupella fragum*. Antimicrobial agent. Orange prisms. Mp 188-190° (natural) Mp 210-212° (synthetic). λ_{max} 222 (ε 17300); 282 (ε 16000); 326 (ε 3800); 424 (ε 880) (EtOH).

Fukuyama, Y. *et al.*, *Tetrahedron*, 1998, **54**, 10007-10016 (isol, synth, ir, uv, pmr, cmr)

5-Methoxy-5-(4-methoxy-phenyl)-3-methyl-2,4-imidazolidine-dione M-256C₁₂H₁₄N₂O₄ 250.254**(±)-form**

Metab. of Indian ocean ascidian *Polycarpa clavata*. Amorph. solid. Prob. artifact. CAS no. not found 13-14CI. λ_{max} 203 (log ε 4.01); 225 (log ε 3.79); 277 (log ε 3.28); 292 (log ε 3.19) (MeOH).

Kang, H. *et al.*, *Tet. Lett.*, 1996, **37**, 2369-2372 (isol, ir, uv, pmr, cmr, struct)

4-Methoxy-5-[(3-methoxy-5-pyrrol-2-yl-2H-pyrrol-2-ylidene)-methyl]-2,2'-bipyrrole, 8CI M-257
[19369-65-4]
[19369-64-3]C₁₉H₁₈N₄O₂ 334.377


Pyrrole antibiotic. Pigment from an Australian ascidian and from the bryozoan *Bugula dentata*, also prod. by a mutant strain of the bacterium *Serratia marcescens*, from *Streptomyces coelicolor* and from *Nembrotha kubaryana*. Shows mild activity against gram-positive bacteria. Blue cryst. (CH₂Cl₂/petrol)(as hydrochloride). Mp 300° dec. (hydrochloride). Red colour in basic soln. λ_{max} 555 (sh); 588 (HCl salt) (Derep). λ_{max} 555 (sh); 588 (HCl salt) (Derep). λ_{max} 326 (ε 26300); 556 (ε 2840); 599 (ε 103000) (CHCl₃) (Berdy).

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1968, 641 (isol, pmr, struct, synth)

Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1982, **35**, 215 (isol, uv, ir, pmr, cmr, ms)



Matsunaga, S. *et al.*, *Experientia*, 1986, **42**, 84 (isol, uv, pmr, cmr)

Karuso, P. *et al.*, *Molecules*, 2002, **7**, 1-6 (isol, Nembrotha)

10-Methoxy-11-[10-(11-methoxyvincamajinyl)]cathafoline M-258
[132242-27-4]C₄₅H₅₄N₄O₇ 762.944

Alkaloid from the stem bark of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae). Opt. inactive.

Morfaux, A.-M. *et al.*, *Phytochemistry*, 1990, **29**, 3345 (isol, uv, ir, pmr, cmr, ms, struct)

2-Methoxy-N-(3-methylbutyl)acetamide, 9CI M-259N-(Methoxyacetyl)-3-methylbutylamine
[81164-06-9]MeOCH₂CONHCH₂CH₂CH(CH₃)₂C₈H₁₇NO₂ 159.228Isol. from the rectal gland of male *Dacus* spp. Bp_{0.2} 78-80°.Baker, R. et al., *Experientia*, 1982, **38**, 232 (isol)Perkins, M.V. et al., *J.C.S. Perkin I*, 1990, 1111 (synth)**7-Methoxy-4-methyl-γ-carboline** M-2607-Methoxy-1-methyl-5H-pyrido[4,3-b]indole, 9CI. γ-Harmine
[128508-19-0]C₁₃H₁₂N₂O 212.251Alkaloid from the seeds of *Pegamum harmala* (Zygophyllaceae). Cryst. (EtOH). Mp 238-240°. First example of a naturally occurring γ-carboline alkaloid. λ_{max} 205 (log ε 4.27); 245 (log ε 4.45); 260 (sh) (log ε 4.16); 265 (sh) (log ε 3.99); 303 (log ε 3.98); 323 (log ε 4.1); 365 (log ε 3.63) (MeOH).Li, G. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 393-397; *CA*, **113**, 74723m (isol, pmr, ms)**2-Methoxy-4,5-methylene-dioxycinnamic acid** M-261C₁₁H₁₀O₅ 222.197**(E)-form**Pyrrolidide: 2-Methoxy-4,5-methylene-dioxycinnamoyl pyrrolidide
[82682-47-1]C₁₅H₁₇NO₄ 275.304Alkaloid from the roots of *Piper amalga* (Piperaceae). Yellowish cryst. Mp 178°.Piperidide: 2-Methoxy-4,5-methylene-dioxycinnamoyl piperidide
[67879-91-8]C₁₆H₁₉NO₄ 289.33Alkaloid from the roots of *Piper amalga* (Piperaceae). Yellowish plates (hexane/Me₂CO). Mp 120°.

2-Methylpropylamide: 2-Methoxy-4,5-methylenedioxcinnamoyl isobutylamide. 5'-Methoxyfagaramide

[259258-85-0]

C₁₅H₁₉NO₄ 277.319Alkaloid from the roots of *Piper amalga* var. *nigrinodum*. Cryst. Mp 150-152°.**(Z)-form**


Pyrrolidide: [105705-79-1]

Minor alkaloid from the leaves of *Piper peepuloides* and from *Piper hispidum* (Piperaceae). Needles (EtOAc/petrol). Mp 118°.

Piperidide: [73885-65-1]

Alkaloid from *Piper peepuloides*. Cryst. (EtOAc/petrol). Mp 99°. λ_{max} 240; 275; 333 (MeOH).Sehgal, C.K. et al., *Phytochemistry*, 1979, **18**, 1865-1867 (cis-Piperidide)Domínguez, X.A. et al., *Phytochemistry*, 1986, **25**, 239 (isol, uv, ir, pmr, ms, struct)Shah, S. et al., *Phytochemistry*, 1986, **25**, 1997-1998 (cis-pyrrolidide)Jacobs, H. et al., *J. Indian Chem. Soc.*, 1999, **76**, 713-717 (isobutylamide)**8-Methoxy-3,4-methylene-dioxy-10-nitro-1-phenanthrenecarboxylic acid** M-262

8-Methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. Aristolochic acid I. Aristolochic acid A. Aristolochic acid. Isoaristolochic acid. Aristolochin†. Aristinic acid. Aristolochia yellow. Descressept. Tardolyt. TR 1736 [313-67-7]

C₁₇H₁₁NO₇ 341.276Alkaloid from a wide range of *Aristolochia* spp. Also found in the Chinese drug Fang-chi, *Asarum canadense* var. *reflexum*, *Bragantia wallichii* (preferred genus name *Thottea*), and in the butterflies *Pachlioptera aristolochiae* and *Battus archidamas*. Phospholipase A₂ inhibitor. Antineoplastic, antiseptic, antiinflammatory and bactericidal agent. Platelet aggregation inhibitor. Cytotoxic. Cryst. (DMF/EtOH, EtOH or MeOH/Et₂O). Sol. MeOH, EtOH, bases; poorly sol. hexane, H₂O. Mp 281-286° dec. (271-273°, 275-278°). Log P 3.48 (calc). H₂SO₄ → green colour. λ_{max} 241; 305; 340; 450 (EtOH/NaOH) (Derep). λ_{max} 221 (ε 29500); 255 (ε 32400); 283; 314 (ε 11200); 389 (ε 6030) (EtOH) (Derep). λ_{max} 223 (ε 1174); 227 (ε 1148); 250 (ε 1445); 318 (ε 478); 390 (ε 251) (MeOH) (Berdy). λ_{max} 221 (ε 29500); 250 (ε 32200); 317 (ε 11200); 390 (ε 6000) (EtOH) (Berdy).▶ Nephrotoxic. Exp. carcinogen. LD₅₀ (rat, orl) 184 mg/kg. LD₅₀ (rat, ivn) 74 mg/kg. CF3325000

Me ester: Methyl aristolochate

[1169-60-4]

C₁₈H₁₃NO₇ 355.303Alkaloid from roots of *Asarum indica* and *Asarum argentina* (Aristolochiaceae). Cytotoxic. Yellow needles (CHCl₃/MeOH), cryst. (CHCl₃). Mp 285-286° (281-282°).

▶ SF8280000

O-De-Me: 8-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. Aristolochic acid Ia [38965-71-8]

C₁₆H₉NO₇ 327.25Alkaloid from the roots of *Aristolochia argentina*, and the leaves and tender stems of *Aristolochia chilensis*. Also found in the butterflies *Zerynthia polyxena* and *Battus archidamas* (Aristolochiaceae). Orange-coloured solid (MeOH). Mp 278° dec.

▶ SF8276000


O-De-Me, Me ester: Aristolochic acid Ia methyl ester

C₁₇H₁₁NO₇ 341.276Alkaloid from *Aristolochia kaempferi*. Yellow syrup. λ_{max} 224; 251; 270 (sh); 288 (sh); 321; 391 (MeOH).

[10190-99-5]

Krishnaswamy, P.R. et al., *J. Indian Chem. Soc.*, 1935, **12**, 476 (isol)Munjunath, B.L. et al., *J. Indian Chem. Soc.*, 1938, **15**, 646 (isol)Pailer, M. et al., *Monatsh. Chem.*, 1956, **87**, 249; 1957, **88**, 367 (isol, ir, uv, struct)Doskotch, R.W. et al., *J. Nat. Prod.*, 1967, **30**, 141 (isol)v. Eeuw, J. et al., *Isr. J. Chem.*, 1968, **6**, 659 (isol, ms)Kupchan, S.M. et al., *J.O.C.*, 1968, **33**, 3735 (isol, pmr)Comer, F. et al., *Can. J. Chem.*, 1969, **47**, 481 (isol, biosynth)Priestap, H.A. et al., *An. Asoc. Quim. Argent.*, 1971, **59**, 245; *CA*, **76**, 43943w (isol, uv, ir)Rothschild, M. et al., *Insect Biochem.*, 1972, **2**, 334 (Aristolochic acid Ia, isol, ms)Pakrashi, S.C. et al., *Phytochemistry*, 1977, **16**, 1103 (isol, ir, ms, struct, ester)Martindale, *The Extra Pharmacopoeia*, 28th edn., Pharmaceutical Press, 1982, 12391Nakanishi, T. et al., *Phytochemistry*, 1982, **21**, 1759 (pmr, ester)Priestap, H.A. et al., *Phytochemistry*, 1982, **21**, 2755 (Me ester, O-de-Me, isol, uv, ir, pmr, ms)Urzúa, A. et al., *Coll. Czech. Chem. Comm.*, 1983, **48**, 1513 (isol, uv, pmr, ms, O-de-Me)De Pascual Teresa, J. et al., *Phytochemistry*, 1983, **22**, 2745 (isol, ir, pmr, ms, synth)Rosenthal, M.D. et al., *Biochim. Biophys. Acta*, 1989, **1001**, 1 (pharmacol)Schmeiser, H.H. et al., *Cancer Res.*, 1990, **50**, 5464 (tox)Mengs, U. et al., *Arch. Toxicol.*, 1993, **67**, 307 (tox)Moreno, J.J. et al., *Immunopharmacology*, 1993, **26**, 1 (pharmacol)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1411Leu, Y.-L. et al., *Phytochemistry*, 1998, **48**, 734-745 (isol, uv, pmr, ms)Wu, T.S. et al., *Biol. Pharm. Bull.*, 2000, **23**, 1216-1219 (Aristolochic acid Ia methyl ester)Wu, T.S. et al., *Chem. Pharm. Bull.*, 2000, **48**, 1006-1009 (activity)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AQY250

8-Methoxy-3,4-methylene-dioxy-1-phenanthrenecarboxylic acid M-263
8-Methoxyphenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. *Aristolonic acid* [35142-05-3]



C₁₇H₁₂O₅ 296.279

Constit. of *Aristolochia indica*. Mp 292°.

► SF8278000

Me ester: [35142-06-4]

C₁₈H₁₄O₅ 310.306

Constit. of *Aristolochia indica*. Mp 172°.

► SF8279000

Amide: Aristolamide

[64543-56-2]

C₁₇H₁₃NO₄ 295.294

Constit. of *Aristolochia indica* and *Uvaria grandiflora*. Cryst. (Me₂CO). Mp 293-294°.

6-Hydroxy: 6-Hydroxy-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid. Aristofolin D
[64543-61-9]

C₁₇H₁₂O₆ 312.278

Constit. of *Aristolochia cucurbitifolia*. Yellow needles (MeOH). Mp >280°. λ_{max} 244 (sh); 259; 296 (sh); 331; 383 (MeOH).

6-Hydroxy, O-β-D-glucopyranoside: Aristofolin A

[214132-42-0]

C₂₃H₂₂O₁₁ 474.42

Constit. of *Aristolochia kaempferi* and *Aristolochia cucurbitifolia*. Pale yellow powder (CHCl₃/MeOH). Mp >280°. Isol. from *A. cucurbitifolia* as the Na salt. λ_{max} 224; 244 (sh); 259; 295; 316; 330; 360 (MeOH).

7-Hydroxy: 7-Hydroxy-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid. Aristofolin B

[214132-44-2]

C₁₇H₁₂O₆ 312.278

Constit. of *Aristolochia cucurbitifolia*. Yellow needles (MeOH). Mp >280°. λ_{max} 227; 275; 297; 323; 377 (MeOH).

6-Methoxy: 6,8-Dimethoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid. Aristolinic acid

[64543-59-5]

C₁₈H₁₄O₆ 326.305

Cryst. (CHCl₃/MeOH). Mp 284-285°.

6-Methoxy, Me ester: [64543-60-8]

C₁₉H₁₆O₆ 340.332

Constit. of *Aristolochia indica*. Cryst. (CHCl₃/MeOH). Mp 206-207°.

7-Methoxy, Me ester: Aristofolin C

[214132-45-3]

C₁₉H₁₆O₆ 340.332

Constit. of *Aristolochia cucurbitifolia*. Yellow needles (CHCl₃). Mp 165-168°. λ_{max} 230; 277; 330; 378 (MeOH).

Pakrashi, S.C. *et al.*, *Phytochemistry*, 1977, **16**, 1103-1104 (*isol, struct*)

Liao, Y. *et al.*, *Zhongcaoyao*, 1996, **27**, 524-525; *CA*, **125**, 323022c (*amide, isol*)


Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1301-1302 (*Aristolofolins*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1998, **49**, 2509-2510 (*Aristolofolin A*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AQX825; MGQ525

4-Methoxy-2-(3,4-methylenedioxyphenyl)quinoline M-264

2-(1,3-Benzodioxol-5-yl)-4-methoxyquinoline, 9CI. *Graveolinine* [4179-37-7]



C₁₇H₁₃NO₃ 279.295

Alkaloid of *Ruta graveolens* (rue) and *Lunasia amara* (Rutaceae). Spasmolytic agent. Prisms (pentane/EtOAc). Mp 116-117°. Log P 3.97 (calc).


Goodwin, S. *et al.*, *J.A.C.S.*, 1959, **81**, 6209 (*isol, ir, uv, struct*)

Chatterjee, A. *et al.*, *Chem. Ind. (London)*, 1962, 1982 (*isol, ir, uv, synth*)

Reisch, J. *et al.*, *Naturwissenschaften*, 1967, **54**, 200 (*synth*)

4-Methoxy-2-(3,4-methylenedioxystyryl)quinoline M-265

2-[2-(1,3-Benzodioxol-5-yl)ethenyl]-4-methoxyquinoline, 9CI



C₁₉H₁₅NO₃ 305.332

(E)-form [124902-96-1]

Alkaloid from the stem bark of *Galipea longiflora* (Rutaceae). Amorph. Mp 178-180° (synthetic).


Fournet, A. *et al.*, *Can. J. Chem.*, 1989, **67**, 2116 (*isol, uv, pmr, ms, struct*)

Caron, S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 138 (*synth, pmr, cmr, ms*)

3-(Methoxymethylene)-2-pyrrolidinethione M-266

Raphanusamide

[104730-65-6]




C₆H₉NOS 143.209

Originally assigned a 1,3-oxazepin-2-one struct. The 1997 *CA Index Guide* retains the erroneous nomencl. Isol. from Japanese radish (*Raphanus sativus* var. *hortensis* f. *gigantissimus*) (Brassicaceae). Growth inhibitor involved in phototropism of radish hypocotyls. Prisms (EtOAc/hexane). Mp 150-151°. λ_{max} 270 (ε 15800); 307 (ε 8200) (MeOH) (Derep).

Hasegawa, K. *et al.*, *Plant Physiol.*, 1986, **81**, 976 (*isol*)

Harada, N. *et al.*, *Tet. Lett.*, 1991, **32**, 6761 (*uv, ir, pmr, cmr, ms, cryst struct, bibl*)

3-Methoxy-2-methyl-5-(1-methylpropyl)pyrazine M-267



C₁₀H₁₆N₂O 180.249


(ξ)-form [870543-96-7]

Prod. by *Chondromyces crocatus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol, synth, pmr, cmr, ms*)

3-Methoxy-2-methyl-5-(2-methylpropyl)pyrazine M-268

5-Isobutyl-3-methoxy-2-methylpyrazine [36329-96-1]




C₁₀H₁₆N₂O 180.249

Prod. by *Chondromyces crocatus*.

Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (*isol, synth, pmr, cmr*)

3-Methoxy-4-methyl-5-(2-methyl-4-thiazolyl)-4-pentenoic acid M-269

[354817-91-7]



C₁₁H₁₅NO₃S 241.31


Prod. by *Sorangium cellulosum*. Precursor of Epithilonos. Amorph. solid. [α]_D²² +11 (c, 1.2 in MeOH). λ_{max} 210 (ε 15900); 249 (ε 11800) (MeOH).

Hardt, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 847-856

3-Methoxy-2-methyl-1-phenyl-9H-carbazole, 9CI M-270

Hyellazole

[74364-11-7]



C₂₀H₁₇NO 287.36

Alkaloid from the blue-green alga *Hyella caespitosa*. Mp 133-134°. λ_{max} 226 (sh) (ε 34000); 232 (ε 36500); 238 (sh) (ε 34500); 250 (sh) (ε 20000); 260 (sh) (ε 13500); 292 (sh) (ε 12500); 304 (ε 18500); 338 (ε 4500); 352 (ε 5000) (EtOH) (Derep).

6-Chloro-6-Chloro-3-methoxy-2-methyl-1-phenyl-9*H*-carbazole, 9*CI*. Chloro-hyellazole

[74364-10-6]

C₂₀H₁₆Cl₂NO 321.805

Alkaloid from *Hyella caespitosa*. Mp 163-164°. λ_{max} 220 (ε 32600); 235 (ε 32600); 242 (ε 32700); 255 (sh) (ε 20000); 272 (sh) (ε 14600); 300 (sh) (ε 13200); 310 (ε 18000); 348 (ε 4400); 360 (ε 4200) (EtOH) (Derep).

Cardellina, J.H. *et al.*, *Tet. Lett.*, 1979, 4915 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Kano, S. *et al.*, *Chem. Comm.*, 1980, 1241 (synth, cmr)

Takano, S. *et al.*, *Heterocycles*, 1981, **16**, 1479 (synth)

Kano, S. *et al.*, *J.O.C.*, 1981, **46**, 3856 (synth, pmr, ms)

Bergman, J. *et al.*, *Tetrahedron*, 1988, **44**, 5215 (synth)

Kawasaki, T. *et al.*, *Chem. Comm.*, 1989, 43 (synth)

Moody, C.J. *et al.*, *J.C.S. Perkin 1*, 1989, 376; 2463 (synth)

Kawasaki, T. *et al.*, *J.C.S. Perkin 1*, 1993, 1777 (synth)

Beccalli, E.M. *et al.*, *J.C.S. Perkin 1*, 1994, 579 (synth, Hyellazole, Chlorohyellazole)


Knölker, H.-J. *et al.*, *Tet. Lett.*, 1995, **36**, 5339 (synth)

Choshi, T. *et al.*, *J.O.C.*, 1997, **62**, 2535 (synth)

Knoelker, H.-J. *et al.*, *Tetrahedron*, 1999, **55**, 10391-10412 (synth)

3-Methoxy-2-(1-methylpropyl)-5-(2-methylpropyl)pyrazine M-271

[870543-98-9]

C₁₃H₂₂N₂O 222.33Prod. by *Chondromyces crocatus*.


Dickschat, J.S. *et al.*, *Eur. J. Org. Chem.*, 2005, 4141-4153 (isol, synth, pmr, cmr)

2-Methoxy-3-(1-methylpropyl)pyrazine M-272

9*CI*2-sec-Butyl-3-methoxypyrazine, 8*CI*. 2-Methoxy-3-sec-butylpyrazine. *FEMA*

3433

[24168-70-5]

C₉H₁₄N₂O 166.222

Volatile component of many vegetables, e.g. asparagus, carrot, celery, cucumber,

parsnip, bell peppers, *Pisum sativum* (pea). Found in galbanum and petitgrain oils and ginger. Prod. by *Chondromyces crocatus* and a marine *Streptomyces* sp. GWS-BW-H5. Fragrance and flavour ingredient.

(±)-form

Bp₁₀ 78-83°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 842C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 404C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1560D (ir)

Bramwell, A.F. *et al.*, *Tet. Lett.*, 1969, 3215

Murray, K.E. *et al.*, *Chem. Ind. (London)*, 1970, 897

Murray, K.E. *et al.*, *J. Sci. Food Agric.*, 1975, **26**, 973

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 459

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1698 (occur)

Schulz, S. *et al.*, *Tetrahedron*, 2004, **60**, 3863-3872 (isol)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, **2**, 837-865 (marine isol)

2-Methoxy-3-(2-methylpropyl)pyrazine M-273

9*CI*2-Isobutyl-3-methoxypyrazine, 8*CI*. Galbazine. *FEMA* 3132

[24683-00-9]

C₉H₁₄N₂O 166.222

Occurs in petitgrain and galbanum oils. Potent component of bell-pepper (*Capsicum annuum*). Found in human urine. Fragrance and flavouring ingredient with powerful earthy flavour. Odour threshold 2 parts per trillion in water.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 842D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 405A (nmr)

Buttery, R.G. *et al.*, *Chem. Ind. (London)*, 1969, 490-491 (isol, synth, ir, pmr, ms)

Murray, K.E. *et al.*, *Chem. Ind. (London)*, 1970, 897-898 (isol)

Seifert, R.M. *et al.*, *J. Agric. Food Chem.*, 1972, **20**, 135-137; *CA*, **76**, 44863g (props, struct)

Bramwell, A.F. *et al.*, *J.C.S. Perkin 1*, 1972, 2004-2007 (synth, pmr)


Calabretta, P.J. *et al.*, *Cosmet. Perfum.*, 1975, **90**, 74; 76; 79-80; *CA*, **83**, 183292c (synth, use)

Hawksworth, G.S. *et al.*, *Xenobiotica*, 1975, **5**, 389-399; *CA*, **83**, 188835s (metab)

3-Methoxy-1-methyl-2-propyl-4(1*H*)-quinolinone M-274

Leioquinine A

[132587-63-4]

C₁₄H₁₇NO₂ 231.294


Alkaloid from the leaves of *Esenbeckia leiocarpa* (Rutaceae). Shows antifeedant activity against *Pectinophora gossypiella*. Needles. Mp 89-91.2°.

Nakatsu, T. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1508 (isol, pmr, cmr, struct, synth)

4-Methoxy-1-methylquinolinium(1+) M-275

Echinorine

[18095-64-2]

C₁₁H₁₂NO⁺ 174.222

Obtd. from leaves of *Echinops commutatus*, fruit of *Echinops ritro* and in *Echinops sphaerocephalus* (Asteraceae).

Iodide:

C₁₁H₁₂INO 301.126

Mp 132-133°.

Perchlorate:

C₁₁H₁₂ClNO₅ 273.672

Mp 251-253°.

Schroeder, P. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1968, **301**, 39


Bereznegovskaya, L.N. *et al.*, *CA*, 1970, **73**, 95486g (biosynth)

Barlin, G.B. *et al.*, *J.C.S. Perkin 2*, 1975, 298 (uv, nmr)

5-Methoxy-2-(methylthio)-1,3-thiazino[6,5-*b*]indole M-276

Dehydro-4-methoxycyclobrassinin

[154933-74-1]

C₁₂H₁₀N₂OS₂ 262.356


Isol. from roots of the turnip *Brassica campestris* ssp. *rapa* (Brassicaceae) inoculated with *Pseudomonas cichorii*. Phytoalexin. Mp 142-143°.

Monde, K. *et al.*, *Heterocycles*, 1994, **38**, 263 (isol, uv, ir, pmr, cmr, ms, struct)

11-Methoxy-3*H*-naphthridino[6,5,4-*def*]quinoxalin-2-ol M-277

11-Methoxy-1*H*-naphthridino[6,5,4-*def*]-quinoxalin-2(3*H*)-one, 9*CI*

[597553-92-9]


C₁₄H₁₁N₃O₂ 253.26

CAS name refers to *NH*-tautomer. Alkaloid from a *Xestospongia* sp. Brown gum. λ_{\max} 210 (ϵ 5420); 222 (ϵ 4795); 238 (ϵ 4660); 260 (ϵ 2940); 271 (ϵ 2600); 357 (ϵ 1330); 371 (ϵ 1070) (EtOH).

Calcul, L. et al., *Tetrahedron*, 2003, **59**, 6539-6544 (*isol*, *pmr*, *cmr*)

11-Methoxyneooxydiabolone M-278

1-Acetyl-17,18:17,19-diepoxy-11-methoxycuran-20-ol, 9CI
[131984-92-4]



Absolute Configuration


$C_{22}H_{26}N_2O_5$ 398.458

Alkaloid from *Strychnos pungens* (Loganiaceae). $[\alpha]_D^{25} +149$ (c, 0.82 in $CHCl_3$).

Thépenier, P. et al., *Phytochemistry*, 1990, **29**, 2384-2386 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

13-Methoxyoxoberberine M-279

13-Methoxy-8-oxoberberine
[56470-41-8]



$C_{21}H_{19}NO_6$ 381.384


Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Cryst. ($CHCl_3$ /hexane). Mp 197-200° (194-195°).

Moniot, J.L. et al., *J.O.C.*, 1979, **44**, 4337 (*synth*, *ir*, *pmr*)

Valencia, E. et al., *Tetrahedron*, 1984, **40**, 3957 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

2-(4-Methoxyphenyl)ethylamine M-280

4-Methoxybenzeneethanamine, 9CI. 4-Methoxyphenethylamine, 8CI. **O-Methyltyramine**. Homoanisylamine
[55-81-2]



$C_9H_{13}NO$ 151.208

Alkaloid detected in *Coryphantha ottonis*, *Coryphantha poselgeniana* and *Coryphantha cornifera* (Cactaceae). Bp₂₀ 138-140° Bp₁₂ 127-130°.

►SH7875000

Hydrochloride: [645-58-9]
Mp 211°.

►DA0301000

Picrate: Mp 177-178°.

N-(3-Methyl-2-butenoyl): **O-Methyl-N-seneciolytyramine**

[924664-21-1]

$C_{14}H_{19}NO_2$ 233.31

Constit. of *Anabasis* sp.

N-Benzoyl: N-[2-(4-Methoxyphenyl)ethyl]benzamide, 9CI. **N-Benzoyltyramine methyl ether**. Dihydroalotamide
[3278-19-1]

$C_{16}H_{17}NO_2$ 255.316

Minor alkaloid from leaves of *Pleiospermium alatum* (Rutaceae) and from *Aniba riparia* and *Pleiospermium alatum*. Cryst. (Et_2O/C_6H_6). Mp 119-121°.

N-(2-Hydroxybenzoyl): **N-(2-Hydroxybenzoyl)-O-methyltyramine**

[112356-53-3]

$C_{16}H_{17}NO_3$ 271.315

Constit. of the fruit of *Aniba riparia*. Cryst. (C_6H_6). Mp 111-112°. λ_{\max} 225 (ϵ 5850); 240 (ϵ 3500); 305 (ϵ 1650) (MeOH).

N-(2,6-Dihydroxybenzoyl): N-(2,6-Dihydroxybenzoyl)-O-methyltyramine. **Riparin**

[112356-54-4]

$C_{16}H_{17}NO_4$ 287.315

Constit. of the fruit of *Aniba riparia* and from *Salvia albo-caerulea*. Antifungal and spasmolytic agent, smooth muscle relaxant. Cryst. (C_6H_6). Mp 131-133°. λ_{\max} 220 (ϵ 14350); 255 (ϵ 6300); 312 (ϵ 2150) (MeOH).

N-(2-Hydroxy-6-methoxybenzoyl): **N-(2-Hydroxy-6-methoxybenzoyl)-O-methyltyramine**

[112356-55-5]

$C_{17}H_{19}NO_4$ 301.341

Constit. of the bark of *Aniba riparia*. Cryst. (petrol). Mp 55-56°. λ_{\max} 220 (ϵ 18050); 255 (ϵ 7650); 312 (ϵ 2850) (MeOH).

N-(3,4-Dihydroxy-E-cinnamoyl): **N-trans-Caffeoyl-O-methyltyramine**

[189307-47-9]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Cuscuta reflexa*. α -Glucosidase inhibitor. Solid. λ_{\max} 225 (log ϵ 4.12); 290 (log ϵ 3.96); 320 (log ϵ 4.43) (MeOH).

N-Me: N-Methyl-4-methoxyphenethylamine. 4-Methoxy-N-methylbenzeneethanamine, 9CI

[4091-50-3]

$C_{10}H_{15}NO$ 165.235

Alkaloid in *Coryphantha macromeris*, *Coryphantha ramiflora* and other spp. in Cactaceae. Bp₁₉ 141-142°.

►SH8110000

N-Me; hydrochloride: [35803-88-4]
Mp 181-182°.

N-Me, N-(3,4-dimethoxy-E-cinnamoyl): **Beecheyamide**

$C_{21}H_{25}NO_4$ 355.433

Alkaloid from the stems of *Zanthoxylum beecheyanum*. Oil. λ_{\max} 245 (log ϵ 4.4); 284 (log ϵ 4.28); 295 (log ϵ 4.11); 324 (log ϵ 4.21) (MeOH).

N,N-Di-Me: N,N-Dimethyl-4-methoxyphenethylamine. 4-Methoxy-N,N-dimethylbenzeneethanamine, 9CI. **O-Methylhordenine**

[775-33-7]

$C_{11}H_{17}NO$ 179.261

Alkaloid from bark of *Teclea simplicifolia* (Rutaceae). Centrally acting anti-hypertensive agent. Bp₉ 117-120°.

N,N-Di-Me; hydrochloride: [50822-98-5]

Blades (EtOH). Mp 176.5° Mp 279-280° dec.

N,N,N-Tri-Me: **O-Methylcandicine**

[777-74-2]

$C_{12}H_{20}NO^{\oplus}$ 194.296

Alkaloid from *Coryphantha greenwoodii* (Cactaceae).

N,N,N-Tri-Me, chloride: [90706-47-1]

$C_{12}H_{20}ClNO$ 229.749

Mp 206-207°.

N,N,N-Tri-Me, iodide:

$C_{12}H_{20}INO$ 321.201

Prisms. Mp 214-215° (204-206°).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1289A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 610B (*nmr*)

Rosenmund, K.W. et al., *Ber.*, 1909, **42**, 4778-4783 (*synth*)

Barger, G. et al., *J.C.S.*, 1909, **95**, 1720-1724 (*synth*)

Badger, G.M. et al., *Aust. J. Chem.*, 1963, **16**,

734-735 (*N,N*-Dimethyl-4-methoxyphenethylamine)

Hornemann, K.M.K. et al., *J. Pharm. Sci.*,

1972, **61**, 41-45 (*occur*)

Smith, J.R.L. et al., *J.C.S. Perkin 1*, 1972, 228-

234 (*synth*, *nmr*)

McLauchlin, J.L. et al., *J. Pharm. Sci.*, 1973,

62, 408-411; 411-414 (*N-Me*, *isol*, *struct*)

Chatterjee, A. et al., *Aust. J. Chem.*, 1975, **28**,

457-460 (*N-Benzoyltyramine methyl ether*)

Bailey, K. et al., *Can. J. Pharm. Sci.*, 1975, **10**,

31-32 (*pmr*)

Meyer, B.N. et al., *J. Nat. Prod.*, 1983, **46**, 688-

693 (*O-Methylcandicine*)

Barbosa-Filho, J.M. et al., *Phytochemistry*,

1987, **26**, 2615-2617 (*N-2-hydroxybenzoyl*,

N-2,6-dihydroxybenzoyl, *N-2-hydroxy-6-methoxybenzoyl*)

Anis, E. et al., *Chem. Pharm. Bull.*, 2002, **50**,

112-114 (*N-Caffeoyl-O-methyltyramine*)

Cheng, M.-J. et al., *J. Chin. Chem. Soc.*

(*Taipei*), 2004, **51**, 1065-1072

(*Beecheyamide*)

Chen, H. et al., *Yaouxue Xuebao*, 2005, **40**, 248-

251 (*O-Methyl-N-seneciolytyramine*)

Lewis, R.J. et al., *Sax's Dangerous Properties*


of Industrial Materials, 8th edn., Van

Nostrand Reinhold, 1992, CNH375;

MFC500

6-(4-Methoxyphenyl)-N-(3-methylbutyl)-2-oxo-3,5-hexadienamide, 9CI

5-(4-Methoxyphenyl)-2,4-hexadienoic acid isopentylamide





$C_{18}H_{23}NO_3$ 301.385

(*E,E*)-form [75933-10-7]

Alkaloid from the aerial parts of *Ottonia ovata* (preferred genus name *Piper*) (Piperaceae).


Haensel, R. et al., *Planta Med.*, 1980, **40**, 161

(4-Methoxyphenyl)-N-methyl-2-oxothioacetamide M-282
4-Methoxy-N-methyl- α -oxobenzeneethanethioamideC₁₀H₁₁NO₂S 209.268Isol. from the ascidian *Polycarpa aurata*. Inhibitor of inosine monophosphate dehydrogenase. Yellow cryst. (CHCl₃/MeOH). Probable artifact. Not in CA.Abas, S.A. *et al.*, *J.O.C.*, 1996, **61**, 2709-2712 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)**6-Methoxy-3-(1-propenyl)-2-pyridinecarboxylic acid, 9CI** M-283
[103972-05-0]C₁₀H₁₁NO₃ 193.202Alkaloid from the aerial parts of *Melochia corchorifolia* (Apocynaceae). Mp 196°.*Me ester*: Mp 118°.Bhakuni, R.S. *et al.*, *Chem. Ind. (London)*, 1986, 464 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)**N-(2-Methoxy-2-propenyl)-trimethylammonium(1+)** M-284
(2-Methoxyallyl)trimethylammonium, 8CIH₂C=C(OMe)CH₂N⁺Me₃C₇H₁₆NO⁺ 130.21Isol. from the marine sponge *Raspailia* sp. Counterion of natural compd not specified.*Bromide*: **Meprochol**, **BAN**. *Esmodil* [590-31-8]C₇H₁₆BrNO 210.114

Muscarinic agonist. Cryst. powder.

Sol. H₂O, EtOH. Mp 169°.

▶ BR7167000

Ger. Pat., 1935, 614 462; *CA*, **29**, 5994 (*synth*)Asakawa, H. *et al.*, *Okayama Igakkai Zasshi*, 1939, **51**, 1020; *CA*, **37**, 682b (*pharmacol*)Capon, R.J. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 305-309 (*isol*, *synth*, *pmr*, *cmr*)**3-Methoxypyrazolo[4,3-e][1,2,4]triazine, 9CI** M-285
Fluviol A. Normethylpseudoiodinine. Norpseudoiodinin [37526-53-7]C₅H₅N₅O 151.127Prod. by *Pseudomonas fluorescens*. Yellow cryst. (MeOH). Mp 196-197°. Subl. at 150°. λ_{\max} 233 (log ϵ 4.16); 280 (log ϵ 3.56); 283 (log ϵ 3.56); 375 (log ϵ 3.16) (EtOH).**1H-form***N-Ac*: [37526-54-8]

Mp 147°.

N-Me: 3-Methoxy-1-methylpyrazolo[4,3-e][1,2,4]triazine




[37526-55-9]

C₆H₇N₅O 165.154

Mp 149°.

N-Hydroxymethyl: Fluviol B

[195154-66-6]

C₆H₇N₅O₂ 181.154Prod. by *Pseudomonas fluorescens*. Mp 141° dec. λ_{\max} 206 (log ϵ 3.66); 233 (log ϵ 4.22); 280 (log ϵ 3.6); 284 (log ϵ 3.6); 290 (sh) (log ϵ 3.5); 375 (log ϵ 3.2); 402 (sh) (log ϵ 3.1) (EtOH).**6H-form***N-Me*: 3-Methoxy-6-methyl-6H-pyrazolo[4,3-e][1,2,4]triazine, 9CI. **Fluviol E** [37526-57-1]C₆H₇N₅O 165.154Prod. by *Pseudomonas fluorescens*. Unstable violet pigment.**7H-form***N-Me*: 3-Methoxy-7-methyl-7H-pyrazolo[4,3-e][1,2,4]triazine. **Fluviol C**. **Pseudoiodinine** [37526-58-2]C₆H₇N₅O 165.154Prod. by *Pseudomonas fluorescens* and *Pseudomonas fluorescens* var. *pseudoiodinum*. Cryst. (Me₂CO/hexane). Mp 142-144° dec. Subl. at 120-130°. Struct. of Pseudoiodinine revised in 2006. λ_{\max} 236 (log ϵ 4.24); 291 (log ϵ 3.79); 298 (sh) (log ϵ 3.72); 401 (log ϵ 3.26); 430 (sh) (log ϵ 3.17) (EtOH).Korth, H. *et al.*, *Arch. Microbiol.*, 1971, **77**, 59Lindner, H.J. *et al.*, *Chem. Ber.*, 1972, **105**,1949-1955 (*synth*, *cryst struct*)Smirnov, V.V. *et al.*, *FEMS Microbiol. Lett.*, 1997, **153**, 357-361 (*isol*)Kelly, T.R. *et al.*, *J.A.C.S.*, 2006, **128**, 5646-5647 (*synth*)**2-[[3-Methoxy-4-(1H-pyrrol-1-yl)-2H-pyrrol-2-ylidene]methyl]-5-methyl-1-pentyl-1H-pyrrole, 9CI** M-286
[418767-73-4]C₂₀H₂₅N₃O 323.437Related to Prodigiosin, P-636. Red pigment prod. by *Micrococcus* sp. Oil. λ_{\max} 536 (MeOH).Variyar, P.S. *et al.*, *Indian J. Chem., Sect. B*, 2002, **41**, 232-233 (*isol*, *pmr*, *cmr*, *ms*)Subramanian, M. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 2480-2486 (*activity*)**N-4-Methoxystyrylcinnamide** M-287*N*-[2-(4-Methoxyphenyl)ethenyl]-3-phenyl-2-propenamide, 9CIC₁₈H₁₇NO₂ 279.338**(E,E)-form** [87596-53-0]Alkaloid from the leaves of *Aegle marmelos* (bael fruit) (Rutaceae). Yellow needles (MeOH). Mp 191°.Govindachari, T.R. *et al.*, *Phytochemistry*, 1983, **22**, 755 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)**5-Methoxytryptamine** M-288
5-Methoxy-1H-indole-3-ethanamine. 3-(2-Aminoethyl)-5-methoxyindole [608-07-1]C₁₁H₁₄N₂O 190.244

See also refs. under 5-Hydroxytryptamine, H-755. Mp 120-121°.

▶ LD₅₀ (mus, ipr) 176 mg/kg. NL4059000*Hydrochloride*: **Mexamine**

[66-83-1]

Cryst. (EtOH). Mp 246-248°.

N^b-Ac: *N*-[2-(5-Methoxy-1H-indol-3-yl)ethyl]acetamide, 9CI. *N*-Acetyl-5-methoxytryptamine. **Melatonin** [73-31-4]C₁₃H₁₆N₂O₂ 232.282Isol. from bovine pineal glands. Product of tryptophan metabolism by the pineal gland. Potent effects on the control of seasonal reproduction in mammals. Synthesis and secretion subject to photoperiodic control. May influence circadian rhythms in humans. Tumour-inhibiting props. and role in immune system under investigation. Pale yellow leaflets (C₆H₆). Mp 116-118°. Low toxicity, non-mutagenic.

▶ AC5955000

N^b-(4-Hydroxy-E-cinnamoyl): *N*-E-p-Coumaroyl-5-methoxytryptamine. **Centcyamine**

[366452-03-1]

C₂₀H₂₀N₂O₃ 336.39Alkaloid from the seeds of *Centaurea cyanus*.*N^b-(4-Hydroxy-Z-cinnamoyl)*: *N*-Z-p-Coumaroyl-5-methoxytryptamine. **cis-Centcyamine**

[365540-94-9]

C₂₀H₂₀N₂O₃ 336.39Alkaloid from the seeds of *Centaurea cyanus*.N^b,Me: 5-Methoxy-N-methyltryptamine

[2009-03-2]

C₁₂H₁₆N₂O 204.271Alkaloid from *Phalaris arundinacea* (Poaceae) and *Desmodium pulchellum* (Fabaceae). Prisms (MeOH/Et₂O) (as hydrochloride). Mp 166-167° (hydrochloride).N^b,N^b-Di-Me: 5-Methoxy-N,N-dimethyltryptamine. O-Methylbufotenine

[1019-45-0]

C₁₃H₁₈N₂O 218.298Alkaloid from *Phalaris tuberosa* (Poaceae), *Desmodium pulchellum* (Fabaceae) and many other spp. in many families. Toxic agent causing staggers-like poisoning of sheep fed on *Phalaris tuberosa* in Australia. Prisms (Et₂O/hexane). Mp 69°. Bp₄ 208-210°.▶ Toxic. LD₅₀ (mus, ipr) 115 mg/kg. NL7380000N^b,N^b-Di-Me, picrate: [4006-02-4]Orange-yellow cryst. (MeOH/Me₂CO). Mp 176-177°.N^b,N^b-Di-Me, N^b-oxide: O-Methylbufotenine N-oxideC₁₃H₁₈N₂O₂ 234.297Alkaloid from several *Desmodium* spp. (Fabaceae). Mp 158° (as picrate).N^b,N^b-Diisopropyl: N,N-Diisopropyl-5-methoxytryptamine. Foxy

[4021-34-5]

C₁₇H₂₆N₂O 274.405

Hallucinogen, mild euphoric, drug of abuse.

N^b,N^b-Diisopropyl, hydrochloride: [2426-63-3]Cryst. (EtOH/Et₂O). Mp 180-181°.

[66-83-1]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 663A


Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 677A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 134B; 156C (nmr)

Wilkinson, S. et al., *J.C.S.*, 1958, 2079-2081 (5-Methoxy-N-methyltryptamine, isol)Patchter, I.J. et al., *J.O.C.*, 1959, 24, 1285-1289 (5-Methoxy-N,N-dimethyltryptamine)Adlerova, I. et al., *Coll. Czech. Chem. Comm.*, 1960, 25, 784; *CA*, 54, 13095fSzmuzkovicz, J. et al., *J.O.C.*, 1960, 25, 857-859 (Melatonin, synth, uv)Julia, M. et al., *Bull. Soc. Chim. Fr.*, 1965, 1411-1417 (diisopropyl, synth)Ghosal, S. et al., *J.O.C.*, 1966, 31, 2284-2288 (5-Methoxy-N-methyltryptamine, 5-Methoxy-N,N-dimethyltryptamine, 5-Methoxy-N,N-dimethyltryptamine oxide)Quarles, W.G. et al., *Acta Cryst. B*, 1974, 30, 99-103 (Melatonin, cryst struct)Mattox, S.E. et al., *Biomed. Mass Spectrom.*, 1975, 2, 272 (synth, ms)Minneman, K.P. et al., *Life Sci.*, 1975, 17, 1189-1199 (rev, pharmacol)Glennon, R.A. et al., *Eur. J. Pharmacol.*, 1983, 86, 453-459 (diisopropyl, pharmacol)Skaltsounis, A.L. et al., *J. Nat. Prod.*, 1983, 46, 732-735 (N,N-di-Me, N,N-di-Me oxide)Arendt, J. et al., *Clin. Endocrinol. (Oxford)*, 1988, 29, 205 (Melatonin, rev)Neville, S. et al., *J. Pineal Res.*, 1989, 6, 73-76 (mutagenicity)Dubcovich, M.L. et al., *Adv. Pineal Res.*, 1991, 5, 343-349 (Melatonin, pharmacol)Reiter, R.J. et al., *Endocr. Rev.*, 1991, 12, 151-180 (Melatonin, rev)Guerrero, J.M. et al., *Endocr. Rev.*, 1992, 18, 91 (Melatonin)Vivien-Roels, B. et al., *Experientia*, 1993, 49, 642-647 (rev, Melatonin)Marais, W. et al., *Synth. Commun.*, 1998, 28, 3681-3691 (Melatonin)Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1599 (Melatonin)Hwang, K.-J. et al., *Synth. Commun.*, 1999, 29, 2099-2104 (Melatonin, synth, pmr)Somei, M. et al., *Chem. Pharm. Bull.*, 2001, 49, 87-89 (Melatonin, synth)Sarker, S.D. et al., *Phytochemistry*, 2001, 57, 1273-1276 (Centcyamine)Amat, M. et al., *Synthesis*, 2001, 267-275 (Melatonin)Hügel, H.M. et al., *Heterocycles*, 2003, 60, 2349-2354 (Melatonin, rev, synth)Thomson, D.W. et al., *Synth. Commun.*, 2003, 33, 3631-3641 (synth, Melatonin)Wilson, J.M. et al., *Forensic Sci. Int.*, 2005, 148, 31-36 (diisopropyl, pharmacol)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, MCB350; Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MFT000; MFS400

11-[10-(11-Methoxyvincamajinyl)]vincorine M-289

[142795-96-8]

C₄₅H₅₄N₄O₇ 762.944Alkaloid from the leaves of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae).

Ac: 11-[10-(11-Methoxyvincamedinyl)]vincorine

[142750-30-9]

C₄₇H₅₆N₄O₈ 804.981Alkaloid from leaves of *Tonduzia pittieri* (Apocynaceae). [α]_D -58 (c, 1 in CHCl₃).


17'-Epimer: 11-[10-(11-Methoxy-17-epi-vincamajinyl)]vincorine

[142750-29-6]

C₄₅H₅₄N₄O₇ 762.944Alkaloid from leaves of *Tonduzia pittieri* (Apocynaceae). [α]_D -37 (c, 1 in CHCl₃).Morfaux, A.-M. et al., *Phytochemistry*, 1992, 31, 1079 (isol, uv, ir, pmr, cmr, ms, struct)

4-Methoxy-1-vinyl-β-carboline M-290

1-Ethenyl-4-methoxy-9H-pyrido[3,4-b]indole, 9CI. Dehydrocrenatine [26585-13-7]

C₁₄H₁₂N₂O 224.262Alkaloid from *Picrasma javanica* and *Ailanthus malabarica* (Simaroubaceae). Antimalarial agent. Mp 146-147°.

Dihydro: 1-Ethyl-4-methoxy-β-carboline.

Crenatine†

[26585-14-8]

C₁₄H₁₄N₂O 226.277Alkaloid from *Aeschrina crenata* (preferred genus name *Picrasma*) and *Quassia undulata* (Simaroubaceae). Mp 181-183°. λ_{max} 243 (ε 28200); 265 (ε 58000); 274 (ε 6500); 284 (ε 11700); 331 (ε 5500); 348 (ε 6200) (MeOH) (Berdy).


6-Hydroxy: 6-Hydroxy-4-methoxy-1-vinyl-β-carboline. 6-Hydroxydehydrocrenatine

[115910-13-9]

C₁₄H₁₂N₂O₂ 240.261Alkaloid from *Picrasma javanica*. Antimalarial agent. Cryst. (EtOH).Johns, S.R. et al., *Aust. J. Chem.*, 1970, 23, 629 (isol, pmr, ms, struct)Sánchez, E. et al., *Phytochemistry*, 1971, 10, 2155 (deriv)Joshi, B.S. et al., *Heterocycles*, 1977, 7, 193 (isol, uv, pmr)Cain, M. et al., *J.O.C.*, 1982, 47, 4933 (synth)Koike, K. et al., *Org. Magn. Reson.*, 1984, 22, 471-473 (cmr)Pavanand, K. et al., *Phytother. Res.*, 1988, 2, 33-36 (6-Hydroxydehydrocrenatine)Hagen, T.J. et al., *J.O.C.*, 1989, 54, 2170 (synth, Crenatine)Holt, E.M. et al., *J. Crystallogr. Spectrosc. Res.*, 1990, 20, 261 (cryst struct, Crenatine)Murakami, Y. et al., *Chem. Pharm. Bull.*, 1991, 39, 2189 (synth, Crenatine)

10-Methoxyyohimbine M-291

Methyl 17-hydroxy-10-methoxyyohimbane-16-carboxylate. 16-Epiexcelsinine [15266-53-2]



Absolute Configuration

C₂₂H₂₈N₂O₄ 384.474Alkaloid from bark of *Aspidosperma*

pruinoseum (Apocynaceae). Amorph. solid. $[\alpha]_D^{25} +56.6$ (c, 1.2 in EtOH). λ_{\max} 225 (log ϵ 4.46); 279 (log ϵ 3.92); 294 (log ϵ 3.87) (EtOH).

Hydrochloride: [16218-82-9]

Mp 275° dec. $[\alpha]_D +77$ (EtOH).

16-Epimer: Excelsinine. 10-Methoxycorynanthine

[15218-17-4]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Aspidosperma excelsum* (Apocynaceae). Cryst. (MeOH). Mp 199-201° dec. $[\alpha]_D -63.4$ (Py).

16-Epimer: hydrochloride: [15218-18-5]

$[\alpha]_D -54.5$. No def. Mp.

16-Epimer, O-Ac: [16179-94-5]

Mp 225° dec. $[\alpha]_D -54.8$ (Py).

17-Epimer: 10-Methoxy- β -yohimbine

[88607-64-1]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. $[\alpha]_D +29$ (c, 0.31 in MeOH).

16,20-Diepimer: 10-Methoxy- α -yohimbine

[88668-49-9]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. $[\alpha]_D -13$ (c, 0.71 in MeOH).

17,20-Diepimer: 10-Methoxy-17-epialloyohimbine

[88644-43-3]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. $[\alpha]_D +4$ (c, 0.53 in MeOH). Rare stereochem. not yet encountered among the yohimbine stereoisomers.

3,16,20-Triepimer: 10-Methoxy-3-epi- α -yohimbine

[121154-13-0]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the stem bark of *Alstonia coriacea* (Apocynaceae). $[\alpha]_D -66$ (c, 1 in $CHCl_3$). λ_{\max} 227 ; 285 ; 310 ; 335 (MeOH).

16,17,20-Triepimer, O-10-de-Me: Powerine

[1360-57-2]

$C_{21}H_{26}N_2O_4$ 370.447

Alkaloid from leaves of *Ochrosia poweri* (Apocynaceae). Plates + 1MeOH (MeOH/Me₂CO). Mp 188-189° dec. $[\alpha]_D -216$ (c, 0.33 in Me₂CO).

Doy, F.A. et al., *Aust. J. Chem.*, 1962, **15**, 548-554 (*Powerine*)

Benoin, P.R. et al., *Can. J. Chem.*, 1967, **45**, 725-730 (*Excelsinine*)

Kowala, C. et al., *J. Cryst. Mol. Struct.*, 1977, **7**, 1 (*Powerine, cryst struct*)

Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 708-722 (*10-Methoxy- α -yohimbine, 10-Methoxy-17-epialloyohimbine, 10-Methoxy- β -yohimbine*)

Cherif, A. et al., *Phytochemistry*, 1989, **28**, 667-670 (*10-Methoxy-3-epi- α -yohimbine*)

Nunes, D.S. et al., *Phytochemistry*, 1992, **31**, 2507-2511 (*10-Methoxyyohimbine*)

11-Methoxyyohimbine

M-292

[65025-21-0]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from bark of *Rauwolfia capuroni* and from *Rauwolfia nitida* root bark (Apocynaceae). Mp 200-202° (block). $[\alpha]_D^{20} +90$ (c, 0.5 in Py).

O-Ac: Poweridine

[1360-56-1]

$C_{24}H_{30}N_2O_5$ 426.511

Alkaloid from leaves of *Ochrosia poweri* (Apocynaceae). Needles + 1MeOH (Me₂CO/MeOH). Mp 226° dec. $[\alpha]_D^{20} -4.9$ (c, 3.0 in Me₂CO).

3-Epimer: 11-Methoxypseudoyohimbine

[65025-20-9]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from bark of *Rauwolfia capuroni* (Apocynaceae). Mp 267-268° (block). $[\alpha]_D^{20} +30$ (c, 1 in Py).

16,20-Diepimer: 11-Methoxy- α -yohimbine

[84710-88-3]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the leaves of *Neisosperma glomerata* (Apocynaceae). Shows weak α_2 -adrenoceptor blocking activity. Cryst. (MeOH). Mp 218°. $[\alpha]_D^{20} -46$ (c, 0.5 in EtOH).

3,16,20-Triepimer: Quaternatine. 11-Methoxy-3-epi- α -yohimbine

[57499-04-4]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Alstonia quaternata* (Apocynaceae). Shows weak α_2 -adrenoceptor blocking activity. Cryst. (MeOH/Me₂CO). Mp 168°. $[\alpha]_D^{20} -105$ (c, 0.35 in Py). λ_{\max} 227 (log ϵ 4.39); 271 (log ϵ 3.7); 297 (log ϵ 3.79) (EtOH).

3,16,20-Triepimer, O-Ac:

Cryst. (Me₂CO). Mp 174°.

16,17,20-Triepimer: 17-Epi-11-methoxy- α -yohimbine. 11-Methoxy-17-epi- α -yohimbine

[84710-89-4]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the leaves of *Neisosperma glomerata* (Apocynaceae). Cryst. (MeOH). Mp 223°. $[\alpha]_D^{20} -68$ (c, 1 in EtOH).

Isomer: Alstonia quaternata Alkaloid 3

[57605-18-2]

$C_{22}H_{28}N_2O_4$ 384.474

Minor alkaloid from *Alstonia quaternata* (Apocynaceae). Shows weak α_2 -adrenoceptor blocking activity. Non-cryst. This is an *ar*-methoxyyohimbine of undetd. regio- and stereochemistry.

Doy, F.A. et al., *Aust. J. Chem.*, 1962, **15**, 548-554 (*Poweridine, isol, uv, ir, struct*)

Johns, S.R. et al., *Aust. J. Chem.*, 1975, **28**, 1627-1629 (*Poweridine, pmr, cmr, ms, stereochem*)

Mamatas-Kalamaras, S. et al., *Phytochemistry*, 1975, **14**, 1849-1854 (*Alstonia quaternata constits*)

Miet, C. et al., *Phytochemistry*, 1977, **16**, 803-805 (*11-Methoxyyohimbine, 11-Methoxypseudoyohimbine, isol, uv, ir, pmr, struct*)


Amer, M.A. et al., *Phytochemistry*, 1981, **20**, 2569-2573 (*isol*)

Seguin, E. et al., *J. Nat. Prod.*, 1982, **45**, 738-744 (*11-Methoxy- α -yohimbine, 11-Methoxy-17-epi- α -yohimbine*)

Methuenine

M-293

4-Ethylidene-1,3,4,4a,5,7,12,12a-octahydro-2-methylpyrido[3',4':4,5]cyclohept[1,2-b]indol-6(2H)-one, 9CI
[63425-00-3]



Absolute Configuration

$C_{19}H_{22}N_2O$ 294.396

Alkaloid from *Ervatamia officinalis*, *Hazunta* spp. and *Pandaca boiteaui*. Anticholinergic agent. Cryst. (MeOH). Mp 205°. $[\alpha]_D +21$ (c, 1 in $CHCl_3$). Log P 3.12 (calc). λ_{\max} 238 (ϵ 13800); 314 (ϵ 19700) (MeOH).

N¹⁸-Oxide: Methuenine N¹⁸-oxide

[96391-95-6]

$C_{19}H_{22}N_2O_2$ 310.395

Alkaloid from *Pterotaberna inconspicua*. Amorph. λ_{\max} 236 (log ϵ 4.14); 311 (log ϵ 4.27) (MeOH).

19,20 α -Dihydro: 20-Episilicine

$C_{19}H_{24}N_2O$ 296.411

Alkaloid from *Ervatamia officinalis* and *Hazunta modesta* var. *methuenii* subvar. *methuenii*. Cryst. (MeOH). Mp 190°. $[\alpha]_D^{20} -57.8$ (c, 0.12 in $CHCl_3$).

19,20 β -Dihydro: Silicine. 16-Demethoxycarbonyl-20-epiervatamine

[60426-50-8]

$C_{19}H_{24}N_2O$ 296.411

Alkaloid from *Rauwolfia discolor*, *Hazunta modesta* and *Hazunta siliciola*. Cryst. (MeOH). Mp 112°. $[\alpha]_D -18$ (c, 1 in $CHCl_3$). Revised struct. Originally descr. as a vobasine alkaloid. λ_{\max} 239 (ϵ 15200); 314 (ϵ 22300) (EtOH).

6,16-Didehydro, 19,20 α -dihydro: 6,16-Didehydro-20-episilicine

$C_{19}H_{22}N_2O$ 294.396

Alkaloid from *Ervatamia officinalis*. Yellow powder. $[\alpha]_D +76.7$ (c, 0.21 in $CHCl_3$). λ_{\max} 217 (log ϵ 4.4); 236 (log ϵ 4.4); 252 (log ϵ 4.45); 326 (log ϵ 4.06) (MeOH).

6-Oxo: 6-Oxomethuenine

[63425-01-4]

$C_{19}H_{20}N_2O_2$ 308.379

Alkaloid from *Hazunta* spp. Mp 260°. $[\alpha]_D -15$ (c, 1 in Py). λ_{\max} 227 (ϵ 20400); 260 (ϵ 19500); 336 (ϵ 14000) (EtOH).

6-Oxo, 19,20 β -dihydro: 6-Oxosilicine

[63492-71-7]

$C_{19}H_{22}N_2O_2$ 310.395

Alkaloid from *Hazunta modesta* and *Hazunta siliciola*. Mp 257-258°. $[\alpha]_D^{20} -21$ (c, 0.16 in Py). λ_{\max} 227 (ϵ 19200); 260 (ϵ 17700); 335 (ϵ 13200) (EtOH).

N¹-Methoxy: N¹-Methoxymethuenine

[130263-08-0]

$C_{20}H_{24}N_2O_2$ 324.422


Alkaloid from the leaves and stem bark of *Ervatamia malaccensis*.

16-Epimer: Isomethuenine. 16-Epimethuenine

- [72748-32-4]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Hazunta* spp., the root bark of *Pterotaberna inconspicua* and the leaves of *Tabernaemontana dichotoma*. Mp 251° (as hydrochloride). [α]_D²⁰ -178 (c, 0.1 in CHCl₃). λ_{max} 240 ; 315 (MeOH).
- 16-Epimer, N-oxide: 16-Epimethuenine N-oxide**
[84306-85-4]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from the root bark of *Pterotaberna inconspicua*. Amorph.
- 16-Epimer, 19,20α-dihydro: 16,20-Diepisilicine**
C₁₉H₂₄N₂O 296.411
Alkaloid from *Ervatamia officinalis*. Amorph. powder. [α]_D²⁰ -67.1 (c, 0.23 in CHCl₃). λ_{max} 236 (log ε 4.36); 312 (log ε 4.42) (MeOH).
- 16-Epimer, 19,20β-dihydro: 16-Episilicine**
Alkaloid J[†]
[169871-26-5]
C₁₉H₂₄N₂O 296.411
Alkaloid from stem bark, root bark and leaves of *Pandaca caducifolia*. Mp 153°. [α]_D²⁰ +20 (c, 1 in CHCl₃). λ_{max} 238 ; 314 (no solvent reported).
- 16-Epimer, 6-oxo, 19,20α-dihydro: 6-Oxo-16,20-diepisilicine**
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from *Ervatamia officinalis*. Yellow powder. [α]_D²⁰ -46.2 (c, 0.13 in CHCl₃). λ_{max} 220 (log ε 4.52); 257 (log ε 4.37); 330 (log ε 4.24) (MeOH).
- 16-Epimer, 6-oxo, 19,20β-dihydro: 16-Epi-6-oxosilicine. 6-Oxo-16-episilicine**
[76231-43-1]
[186246-58-2]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from root bark and stem bark of *Hazunta modesta* var. *modesta* subvar. *montana*.
- Combes, G. et al., *Phytochemistry*, 1968, **7**, 477-483 (*Silicine*, uv, ir, pmr, ms)
Zèches, M. et al., *Phytochemistry*, 1975, **14**, 1122-1124 (*Alkaloid J*)
Shafiee, A. et al., *Tet. Lett.*, 1976, 921-924 (*Silicine*, cryst struct)
Reis, F. et al., *Tet. Lett.*, 1976, 1085-1088 (6-Oxosilicine)
Riche, C. et al., *Acta Cryst. B*, 1977, **33**, 133-135 (*Silicine*, cryst struct)
Bui, A.-M. et al., *Phytochemistry*, 1977, **16**, 703; 1979, **18**, 1329-1331 (*Methuenine*, *Silicine*, isol, uv, ir, pmr, ms, struct, 6-oxo derivs)
Vecchietti, V. et al., *Phytochemistry*, 1978, **17**, 835-836 (*Silicine*, pmr, cmr)
Bui, A.M. et al., *Phytochemistry*, 1980, **19**, 1473-1475 (6-Oxo-16-episilicine)
Morfaux, A.M. et al., *Phytochemistry*, 1982, **21**, 1767-1769 (16-Epimethuenine, 16-Epimethuenine N-oxide, isol, uv, ir, pmr, ms, struct)
Perera, P. et al., *Planta Med.*, 1983, **47**, 148-150 (isol, uv, ir, pmr, ms, epimer)
Bakana, P. et al., *Planta Med.*, 1984, **50**, 331-334 (*Methuenine*, N-oxide)
Clivio, P. et al., *Phytochemistry*, 1990, **29**, 2693-2696; 1995, **40**, 987-990 (N¹-Methoxymethuenine, 16-Episilicine)
Bennasar, M.L. et al., *Eur. J. Org. Chem.*, 2000, 3919-3925 (6-Oxo-16-episilicine, synth)
Zhang, H. et al., *Helv. Chim. Acta*, 2005, **88**, 2537-2542 (*Ervatamia officinalis* alkaloids)

Methyl [(acetylamino)oxy]acetate, 9CI M-294
Methyl acethydroxamylacetate
[35404-96-7]
AcNHCH₂COOMe
C₅H₉NO₄ 147.13
Metab. of *Irpex pachyodon*. Cryst. Mp 89-91°.
Nano, G.M. et al., *Tet. Lett.*, 1972, 1195 (isol, synth, pmr, ir, ms)

Methyl 3-alkylpyrrole-2-carboxylates M-295



Isol. as a mixt. of n = 18 (29%), n = 19 (24%), n = 20 (40%) and n = 22 (7%).
Metab. from the marine sponge *Oscarella lobularis*.

Methyl 3-nonadecyl-1H-pyrrole-2-carboxylate [57992-59-3]
C₂₅H₄₅NO₂ 391.636
n = 18.

Methyl 3-icosyl-1H-pyrrole-2-carboxylate [57992-60-6]
C₂₆H₄₇NO₂ 405.663
n = 19.

Methyl 3-heneicosyl-1H-pyrrole-2-carboxylate [57992-61-7]
C₂₇H₄₉NO₂ 419.69
n = 20.

Didehydro: Methyl 3-heneicosenyl-1H-pyrrole-2-carboxylate
[58018-93-2]
C₂₇H₄₇NO₂ 417.674
Metab. from the marine sponge *Oscarella lobularis*. Posn. of double bond in side-chain not determined. Isol. in admixture with the tricosenyl homologue.


Methyl 3-tricosyl-1H-pyrrole-2-carboxylate [57992-62-8]
C₂₉H₅₃NO₂ 447.743
n = 22.

Didehydro: Methyl 3-tricosenyl-1H-pyrrole-2-carboxylate
[58018-92-1]
C₂₉H₅₁NO₂ 445.727
Metab. of *Oscarella lobularis*. Isol. in admixture with its lower homologue. Posn. of double bond in side-chain undetermined.

Tetradehydro: Methyl 3-(tricosadienyl)-1H-pyrrole-2-carboxylate
[58018-98-7]
C₂₉H₄₉NO₂ 443.712
Metab. from the marine sponge *Oscarella lobularis*. Posn. of 2 double bonds in side chain undetermined.

Cimino, G. et al., *Experientia*, 1975, **31**, 1387-1389 (isol, uv, ir, pmr, ms, struct)

Methyl chanofruticosinate M-296
Dimethyl 21-oxo-11,21-cycloaspido-permide-1,2-dicarboxylate, 9CI. Chanofruticosine methyl ester
[14050-92-1]



Absolute Configuration

C₂₃H₂₆N₂O₅ 410.469
Alkaloid from *Kopsia officinalis* (Apocynaceae). Mp 165-166°. [α]_D²⁰ +103.7 (CHCl₃). Forms MeOH solvate, softening at 140-143°.

N-De(methoxycarbonyl): **Methyl N¹-demethoxycarbonylchanofruticosinate**
[80151-89-9]

C₂₁H₂₄N₂O₃ 352.432
Obt. from *Kopsia officinalis* and leaves of *Kopsia arborea* (Apocynaceae). Mp 202-203°. [α]_D²⁰ +246.7 (CHCl₃).

3-Oxo, 14,15-didehydro, N-de(methoxycarbonyl): Methyl N¹-demethoxycarbonyl-3-oxo-Δ^{14,15}-chanofruticosinate
C₂₁H₂₀N₂O₄ 364.4
Alkaloid from the leaves of *Kopsia officinalis*. Powder. [α]_D²⁰ +167.5 (c, 0.24 in CHCl₃). λ_{max} 192 ; 241 ; 289 (CHCl₃).

15S-Hydroxy, N-de(methoxycarbonyl): Methyl 15-hydroxy-N¹-demethoxycarbonylchanofruticosinate. Flavisiamine A. Prunifoline D
[1007552-36-4]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from the leaves of *Kopsia arborea* and *Kopsia flavida*. Amorph. solid. [α]_D²⁰ +214 (c, 0.3 in CHCl₃). λ_{max} 205 (log ε 4.4); 236 (log ε 3.8); 293 (log ε 3.43) (EtOH).

12-Methoxy: Methyl 12-methoxychanofruticosinate
[352276-74-5]
C₂₄H₂₈N₂O₆ 440.495
Alkaloid from *Kopsia flavida*. λ_{max} 215 (log ε 4.24); 246 (log ε 3.65); 289 (log ε 2.9) (EtOH).

12-Methoxy, N-de(methoxycarbonyl): Methyl 12-methoxy-N¹-demethoxycarbonylchanofruticosinate
[352276-73-4]
C₂₂H₂₆N₂O₄ 382.458
Alkaloid from *Kopsia flavida*. λ_{max} 210 (log ε 4.53); 293 (log ε 3.52) (EtOH).

12-Methoxy, 14,15-didehydro, N-de(methoxycarbonyl): Methyl 12-methoxy-N¹-demethoxycarbonyl-Δ^{14,15}-chanofruticosinate. Prunifoline C
[605666-56-6]
C₂₂H₂₄N₂O₄ 380.443
Alkaloid from the leaves of *Kopsia arborea*. Oil. [α]_D²⁰ +226 (c, 0.51 in CHCl₃). λ_{max} 206 (log ε 4.48); 235 (log ε 3.88); 292 (log ε 3.39) (EtOH).

12-Methoxy, 3-oxo: Methyl 12-methoxy-3-oxochanofruticosinate. Prunifoline A
[1007552-34-2]

C₂₄H₂₆N₂O₇ 454.479Alkaloid from the leaves of *Kopsia arborea*. Oil. [α]_D +230 (c, 0.15 in CHCl₃); λ_{max} 214 (log ε 4.77); 250 (log ε 4.13); 290 (log ε 3.53) (EtOH).12-Methoxy, 3-oxo, 14,15-didehydro, N-de(methoxycarbonyl): **Methyl 12-methoxy-N¹-demethoxycarbonyl-3-oxo-Δ^{14,15}-chanofruticosinate**

[605666-54-4]

C₂₂H₂₂N₂O₅ 394.426Alkaloid from the leaves of *Kopsia flavida*. Amorph. yellow powder. λ_{max} 210 (log ε 4.33); 293 (log ε 3.3) (MeOH).12-Methoxy, 15S-hydroxy, N-de(methoxycarbonyl): **Methyl 15-hydroxy-12-methoxy-N¹-demethoxycarbonylchanofruticosinate. Flavisiamine C. Prunifoline F**

[1007552-38-6]

C₂₂H₂₆N₂O₅ 398.458Alkaloid from the leaves of *Kopsia arborea* and *Kopsia flavida*. Amorph. solid. [α]_D +261 (c, 0.12 in CHCl₃); λ_{max} 208 (log ε 4.38); 238 (log ε 3.68); 293 (log ε 3.19) (EtOH).11,12-Dimethoxy: **Methyl 11,12-dimethoxychanofruticosinate**

[352276-75-6]

C₂₅H₃₀N₂O₇ 470.521Alkaloid from *Kopsia flavida* (Apocynaceae). λ_{max} 221 (log ε 4.04); 279 (log ε 2.95) (EtOH).

11,12-Dimethoxy, 14,15-didehydro:

Methyl 11,12-dimethoxy-Δ^{14,15}-chanofruticosinate. Prunifoline B

[1007552-35-3]

C₂₅H₂₈N₂O₇ 468.505Alkaloid from the leaves of *Kopsia arborea*. Oil. [α]_D +51 (c, 0.07 in CHCl₃); λ_{max} 220 (log ε 4.45); 247 (log ε 3.93); 280 (log ε 3.22) (EtOH).11,12-Methylenedioxy: **Methyl 11,12-methylenedioxychanofruticosinate**

[80151-90-2]

C₂₄H₂₆N₂O₇ 454.479Alkaloid from *Kopsia officinalis* and leaves of *Kopsia arborea* (Apocynaceae). Mp 252-258° dec. [α]_D +145 (c, 0.27 in CHCl₃).11,12-Methylenedioxy, N-de(methoxycarbonyl): **Methyl 11,12-methylenedioxy-N¹-demethoxycarbonylchanofruticosinate**

[147089-77-8]

C₂₂H₂₄N₂O₅ 396.442Alkaloid from leaves of *Kopsia arborea* (Apocynaceae). [α]_D +193 (c, 0.14 in CHCl₃).11,12-Methylenedioxy, 14,15-didehydro, N-de(methoxycarbonyl): **Methyl 11,12-methylenedioxy-N¹-demethoxycarbonyl-Δ^{14,15}-chanofruticosinate**

[147089-78-9]

C₂₂H₂₂N₂O₅ 394.426Alkaloid from leaves of *Kopsia arborea* (Apocynaceae). [α]_D +198 (c, 0.32 in CHCl₃).11,12-Methylenedioxy, 3-oxo, 14,15-didehydro, N-de(methoxycarbonyl): **Methyl 11,12-methylenedioxy-N¹-de-****methoxycarbonyl-3-oxo-Δ^{14,15}-chanofruticosinate**

[605666-55-5]

C₂₂H₂₀N₂O₆ 408.41Alkaloid from the leaves of *Kopsia flavida*. Amorph. orange powder. λ_{max} 218 (log ε 4.37); 246 (log ε 3.65) (MeOH).11,12-Methylenedioxy, 15S-hydroxy, N-de(methoxycarbonyl): **Methyl 15-hydroxy-11,12-methylenedioxy-N¹-demethoxycarbonylchanofruticosinate. Flavisiamine D. Prunifoline E**

[1007552-37-5]

C₂₂H₂₄N₂O₆ 412.441Alkaloid from the leaves of *Kopsia arborea* and *Kopsia flavida*. Amorph. solid. [α]_D +166 (c, 0.33 in CHCl₃); λ_{max} 219 (log ε 4.16); 239 (log ε 3.93); 285 (log ε 3) (EtOH).Guggisberg, A. et al., *Helv. Chim. Acta*, 1966,


49, 2321-2337 (synth, uv, ir, ms)

Wei-Shin, C. et al., *Annalen*, 1981, 1886-1892(isol, 11,12-dimethoxy, N¹-

de(methoxycarbonyl))

Kam, T.-S. et al., *Phytochemistry*, 1993, 32,489-491 (*Kopsia arborea* constits)Husain, K. et al., *Phytochemistry*, 2001, 57,603-606 (*Kopsia flavida* constits)Husain, K. et al., *J. Asian Nat. Prod. Res.*,2003, 5, 63-67 (*Kopsia flavida* 3-oxo derivs)Zhou, H. et al., *Helv. Chim. Acta*, 2006, 89,515-519 (*Kopsia officinalis* 3-oxo deriv)Sekiguchi, M. et al., *Heterocycles*, 2008, 75,2283-2288 (*Flavisiamine A-D*)Lim, K.-H. et al., *Phytochemistry*, 2008, 69,558-561 (*Prunifolines A-F*)**Methyl 1,2-didehydro-6,7-epoxy-3-hydroxy-8-oxoaspidospermidine-3-carboxylate, 9CI**


16-Carbomethoxy-16-hydroxy-14,15-epoxy-3-oxo-1,2-didehydroaspidospermidine [67332-48-3]

C₂₁H₂₂N₂O₅ 382.415Alkaloid from *Amsonia elliptica* (Apocynaceae). Amorph. powder.Aimi, N. et al., *Chem. Pharm. Bull.*, 1978, 26,




1182 (isol, uv, ir, pmr, cd, ms, struct)

Methyl 16,17-didehydro-methoxy-19-methylxayohimban-16-carboxylate, 9CI

[49693-40-5]


C₂₂H₂₆N₂O₄ 382.458Alkaloid from *Rauwolfia suaveolens* (Apocynaceae). Full struct. not determined. Mol. formula incorrectly given as C₂₂H₂₂N₂O₄ in the paper. λ_{max} 214 ; 225 ; 284 (sh) (EtOH).Majumdar, S.P. et al., *Phytochemistry*, 1973, 12, 1167-1169 (isol, uv, ir, ms)**4-Methyl N-[(2,3-dihydro-2-oxo-1H-indol-3-yl)acetyl]aspartate** M-299

[115082-62-7]

C₁₅H₁₆N₂O₆ 320.301Isol. from pistils and stamens of *Hibiscus moscheutos*. Prisms. Mp 187-189°.Ohmoto, T. et al., *Chem. Pharm. Bull.*, 1988, 36, 578-581 (isol, uv, ir, pmr, cmr, ms, struct)**Methyl 4-[(1-hydroxymethyl-2-phenylethenyl)amino]-4-oxobutenoate** M-300C₁₄H₁₅NO₄ 261.277**(Z,Z)-form**Prod. by *Aspergillus niger* Ta1. Yellow powder. λ_{max} 205 ; 244 ; 291 (MeOH).Yuan, W. et al., *Nat. Prod. Res.*, 2006, 20, 573-577 (isol, pmr, cmr)**4-Methylnaconitane-1,2,3,6,8,13,14,15,16,18-decol** M-301C₁₉H₂₉NO₁₀ 431.439**(1α,2β,3α,5β,6α,14α,15α,16β)-form**O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Altaconitine. 2-Hydroxyxayohimban-16-carboxylate**

[171090-87-2]

C₃₄H₄₇NO₁₂ 661.745Alkaloid from aerial parts of *Aconitum altaicum*. Mp 235-237° dec.Butbayer, N. et al., *Khim. Prir. Soedin.*, 1993, 29, 47; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, 29, 38 (*Altaconitine*)

4-Methylnaconitane-
1,3,6,8,10,13,14,15,16,18-decolC₁₉H₂₉NO₁₀ 431.439(1 α ,3 α ,5 β ,6 α ,14 α ,15 α ,16 β)-formO¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 8-Ac: **Beiwutine**

[76918-93-9]

C₃₃H₄₅NO₁₂ 647.718Alkaloid from *Aconitum kusnezoffii* (Ranunculaceae). Mp 196-198°.

▶ CT3577000

O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 3,8-di-Ac: **3-O-Acetylbeiwutine**


[147677-09-6]

C₃₅H₄₇NO₁₃ 689.755Alkaloid from roots of *Aconitum liaotungense* (Ranunculaceae). Cryst. Mp 187°. [α]_D²⁵ +25.8 (c, 0.95 in EtOH).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **10-Hydroxyaconitine. Aconifine. Nagarin†**

[41849-35-8]

C₃₄H₄₇NO₁₂ 661.745Alkaloid from tubers of *Aconitum karakolicum* and roots of *Aconitum nagarum* var. *lasiandrum* (Ranunculaceae). Mp 198-200°. [α]_D¹⁹ +30.6 (c, 1.10 in CHCl₃).

▶ AR5550000

O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac, hydrobromide: Mp 209°.O⁸-Et, O¹,O⁶,O¹⁶,O¹⁸-tetra-Me, N-Me, 14-benzoyl: **Beiwucine**C₃₃H₄₇NO₁₁ 633.734Alkaloid from *Aconitum kusnezoffii*. Cryst. (Me₂CO). Mp 164-167°.Sultankhodzhaev, M.N. et al., *Khim. Prir. Soedin.*, 1973, **9**, 127; 1980, **16**, 665; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 129; 1980, **16**, 481 (Aconifine)Wang, Y.-G. et al., *Yaoxue Xuebao*, 1980, **15**, 526; *CA*, **94**, 117772k (Beiwutine)Wang, H. et al., *Huaxue Xuebao*, 1981, **39**, 869; *CA*, **97**, 107080f (Aconifine)Zhu, Y. et al., *Heterocycles*, 1982, **17**, 607 (struct, rev)Zhu, D.-Y. et al., *Phytochemistry*, 1993, **32**, 767 (3-O-Acetylbeiwutine)Yu, H. et al., *Yaoxue Xuebao*, 2000, **35**, 232-234 (Beiwucine)4-Methylnaconitane-
1,3,6,8,14,16,18-heptolC₁₉H₂₉NO₇ 383.441(1 α ,3 α ,6 α ,14 α ,16 β)-formO¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et: **Ezo-chasmanine**

[77757-07-4]

C₂₅H₄₁NO₇ 467.601Minor alkaloid from *Aconitum yesoense* (Ranunculaceae). Cryst. + 1/8H₂O (Et₂O). Mp 115-118°. [α]_D²¹ +40.3 (CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl: **Deacetylhemisleyanine. 13-Deoxy-ludaconitine**

[77757-11-0]

C₃₂H₄₅NO₈ 571.709Alkaloid from the roots of *Aconitum hemisleyanum* var. *pengzhouense*. Amorph. solid. [α]_D¹⁷ +22.2 (c, 0.5 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Hemisleyanine. 13,15-Dideoxyaconitine. 8-Acetyl-14-benzoylzechasmanine**

[77757-14-3]

C₃₄H₄₇NO₉ 613.747Alkaloid from roots of *Aconitum sungpanense* and from *Aconitum hemisleyanum* (Ranunculaceae). Cryst. Mp 167-169°. [α]_D²⁵ +16.4 (c, 0.07 in EtOH).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-(4-methoxybenzoyl): **Austroconitine B**

[128717-96-4]

C₃₃H₄₇NO₉ 601.736Alkaloid from the roots of *Aconitum austroyunnanense* (Ranunculaceae).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-(4-methoxybenzoyl), 8-Ac: **Vilmorrianine A. 3 α -Hydroxyforesaconitine**

[1361-96-2]

C₃₅H₄₉NO₁₀ 643.773Alkaloid from the roots of *Aconitum vilmorrianum* and *Aconitum kongboense* (Ranunculaceae). Prisms (EtOAc/hexane). Mp 169°. [α]_D²⁵ +24.1 (CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-(4-methoxybenzoyl), 3,8-di-Ac: **3-Acetyl-vilmorrianine A**

[80787-50-4]

C₃₇H₅₁NO₁₁ 685.81Alkaloid from *Aconitum* sp. (Ranunculaceae).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-(3,4-dimethoxybenzoyl): **Falconerine**

[102358-18-9]

C₃₄H₄₉NO₁₀ 631.762Alkaloid from the Indian crude drug Mitha telia (roots of *Aconitum falconeri*) (Ranunculaceae). Amorph. solid. [α]_D³⁰ +40.3 (c, 0.55 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-(3,4-dimethoxybenzoyl), 8-Ac: **Falconerine 8-acetate**

[102386-46-9]

C₃₆H₅₁NO₁₁ 673.799Alkaloid from *Aconitum falconeri* (Ranunculaceae). Cryst. (hexane). Mp 162-163°. [α]_D³⁰ +13.7 (c, 1.05 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 8-E-cinnamoyl, 14-Ac: **Leuantine A**

[499969-75-4]

C₃₆H₄₉NO₉ 639.784Alkaloid from the roots of *Aconitum hemisleyanum* var. *leuantinus*. Amorph. powder. [α]_D²⁰ +13.4 (c, 0.5 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-E-cinnamoyl: **Deacetylsungpaconitine**


[327028-19-3]

C₃₄H₄₇NO₈ 597.747Alkaloid from the roots of *Aconitum hemisleyanum* var. *pengzhouense*. Amorph. solid. [α]_D¹⁷ +38 (c, 0.45 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-E-cinnamoyl, 8-Ac: **Sungpaconitine**

[114358-55-3]

C₃₆H₄₉NO₉ 639.784Alkaloid from roots of *Aconitum sungpanense* (Ranunculaceae). Needles. Mp 127-129°. [α]_D²⁰ +17.4 (c, 0.87 in CHCl₃).18-Aldehyde, O⁶,O¹⁶-di-Me, N-Et, 8,14-di-Ac: **Staphisadrine**

[132236-91-0]

C₂₇H₃₉NO₉ 521.606Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Amorph. [α]_D²⁰ +6.3 (c, 0.175 in CHCl₃).Takayama, H. et al., *Heterocycles*, 1981, **15**, 403 (Ezo-chasmanine, synth, cd)Yang, C.-R. et al., *Huaxue Xuebao*, 1981, **39**, 147-152; 445; *CA*, **95**, 58094m; **96**, 82689a (Vilmorrianine A, 3-Acetylvilmorrianine A)Wang, F. et al., *Yaoxue Tongbao*, 1982, **17**, 395; *CA*, **97**, 195756j (Ezo-chasmanine)Wang, R. et al., *Planta Med.*, 1987, **53**, 544 (Sungpaconitine)Liu, L. et al., *Yaoxue Xuebao*, 1988, **23**, 460; *CA*, **109**, 208259t (13,15-Dideoxyaconitine)Jiang, Z. et al., *Yunnan Zhiwu Yanjiu*, 1989, **11**, 461; *CA*, **113**, 74751 (Austroconitine B)Liang, X. et al., *J. Nat. Prod.*, 1990, **53**, 1307-1311 (Staphisadrine)Ding, L. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1994, **36**, 901; *CA*, **122**, 235277x (Hemisleyanine)Peng, C.S. et al., *J. Asian Nat. Prod. Res.*, 2000, **2**, 245-249 (13-Deoxyhudaconitine, Deacetylsungpaconitine)Li, L.-Y. et al., *J. Nat. Prod.*, 2003, **66**, 269-271 (Leuantine A)4-Methylnaconitane-
1,3,8,13,14,16,18-heptolC₁₉H₂₉NO₇ 383.441(1 α ,3 α ,14 α ,16 β)-formO¹,O¹⁸-Di-Me, N-Et, 16-O-(4-methoxybenzoyl): **Circinasine C**

[939972-77-7]

C₃₁H₄₃NO₉ 573.682Alkaloid from the roots of *Aconitum*

hemsleyanum var. *circinatum*. Amorph. powder. $[\alpha]_D^{20} +43.8$ (c, 1 in CHCl_3).

O^1, O^{16}, O^{18} -Tri-Me, N-Et: **Circinadine B**
 $\text{C}_{24}\text{H}_{39}\text{NO}_7$ 453.575

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*.

Amorph. powder. Mp 92-93°. $[\alpha]_D^{20} -32.3$ (c, 1 in CHCl_3).

O^1, O^{16}, O^{18} -Tri-Me, N-Et, 14-O-(4-methoxybenzoyl): **Circinadine A**
 $\text{C}_{32}\text{H}_{45}\text{NO}_9$ 587.709


Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*.

Amorph. powder. Mp 102-103°. $[\alpha]_D^{20} -69.2$ (c, 1 in CHCl_3).

Gao, F. et al., *Chem. Pharm. Bull.*, 2006, **54**, 117-118 (*Circinadines A,B*)

Gao, F. et al., *J. Nat. Prod.*, 2007, **70**, 876-879 (*Circinasine C*)

4-Methylnaconitane-1,5,6,7,8,14,16-heptol M-305



$\text{C}_{19}\text{H}_{29}\text{NO}_7$ 383.441

(1 α ,5 β ,6 β ,14 α ,16 β)-form

7,8-Methylene, O^1, O^{14} -di-Me ether, N-Et, 6-Ac: **Nordhagenine C**

$\text{C}_{26}\text{H}_{39}\text{NO}_8$ 493.596

Alkaloid from *Delphinium nordhagenii*. Amorph. solid. $[\alpha]_D^{30} -4.3$ (c, 0.7 in CHCl_3).

7,8-Methylene, O^{14}, O^{16} -di-Me, N-Et:

Laxicymisine

[1008106-59-9]

$\text{C}_{24}\text{H}_{37}\text{NO}_7$ 451.559

Alkaloid from *Delphinium laxicymosum* var. *pilostachyum*. Cryst. Mp 161-163°. $[\alpha]_D^{30} -23.6$ (c, 0.5 in CHCl_3).

7,8-Methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Me: **Bonvalol**

[93961-23-0]

$\text{C}_{24}\text{H}_{37}\text{NO}_7$ 451.559

Alkaloid from the roots of *Delphinium bonvalotii* (Ranunculaceae). Mp 165-166°. $[\alpha]_D^{32} -26.3$ (c, 0.4 in CHCl_3).

7,8-Methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Me, 6-Ac: **Bonvalotine**

[93961-22-9]

$\text{C}_{26}\text{H}_{39}\text{NO}_8$ 493.596

Alkaloid from the roots of *Delphinium bonvalotii* (Ranunculaceae). Mp 218-220°. $[\alpha]_D^{32} -35.7$ (c, 0.6 in CHCl_3).

7,8-Methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Et: **Bonvalotidine B**

[929019-26-1]

$\text{C}_{25}\text{H}_{39}\text{NO}_7$ 465.586

Alkaloid from the roots of *Delphinium bonvalotii*.

7,8-Methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Et, 6-Ac: **Bonvalotidine A**

[929019-25-0]

$\text{C}_{27}\text{H}_{41}\text{NO}_8$ 507.623

Alkaloid from the roots of *Delphinium bonvalotii*.

6-Ketone, 7,8-methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Me: **Bonvalone**

[93961-24-1]

$\text{C}_{24}\text{H}_{35}\text{NO}_7$ 449.543

Alkaloid from the roots of *Delphinium bonvalotii* (Ranunculaceae). Mp 235-236°. $[\alpha]_D^{32} -89.3$ (c, 0.3 in CHCl_3).

6-Ketone, 7,8-methylene, O^1, O^{14}, O^{16} -tri-Me ether, N-Et: **Bonvalotidine C**

[929019-27-2]

$\text{C}_{25}\text{H}_{37}\text{NO}_7$ 463.57

Alkaloid from the roots of *Delphinium bonvalotii*.


Jiang, Q.P. et al., *Heterocycles*, 1984, **22**, 2429-2432 (*Bonvalol, Bonvalotine, Bonvalone*)

Shaheen, F. et al., *J. Nat. Prod.*, 2006, **69**, 823-825 (*Nordhagenine C*)

He, Y. et al., *Nat. Prod. Commun.*, 2006, **1**, 357-362 (*Bonvalotidines A-C*)

Tang, P. et al., *Chin. Chem. Lett.*, 2007, **18**, 700-703 (*Laxicymisine*)

4-Methylnaconitane-1,5,8,13,14,16,18-heptol M-306



$\text{C}_{19}\text{H}_{29}\text{NO}_7$ 383.441

(1 α ,14 α ,16 β)-form

O^1, O^{18} -Di-Me, N-Et: **Circinasine A**

[939972-75-5]

$\text{C}_{23}\text{H}_{37}\text{NO}_7$ 439.548

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. $[\alpha]_D^{20} -32$ (c, 1 in CHCl_3).

O^1, O^{18} -Di-Me, 14-O-(4-methoxybenzoyl), N-Et: **Isohemsleyanidine**. *Isohemsleyanisine* (*incorr.*)

[178960-60-6]

[939972-83-5]

$\text{C}_{31}\text{H}_{43}\text{NO}_9$ 573.682

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. $[\alpha]_D +14.6$ (c, 1 in CHCl_3). Struct. revised in 2007.

O^1, O^{18} -Di-Me, 16-O-(4-methoxybenzoyl), N-Et: **Hemsleyanidine**. *Hemsleyanisine* (*incorr.*)

[178960-59-3]

[939972-82-4]

$\text{C}_{31}\text{H}_{43}\text{NO}_9$ 573.682

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. $[\alpha]_D -8.9$ (c, 0.71 in CHCl_3). Struct. revised in 2007.

O^1, O^{16}, O^{18} -Tri-Me, N-Et: **Hemsleyanine B**

$\text{C}_{24}\text{H}_{39}\text{NO}_7$ 453.575

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph.

powder. Mp 92-93°. $[\alpha]_D^{20} -35.2$ (c, 1 in CHCl_3).

O^1, O^{16}, O^{18} -Tri-Me, 14-(4-hydroxybenzoyl), N-Et: **Hemsleyanine A**

$\text{C}_{31}\text{H}_{43}\text{NO}_9$ 573.682

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. Mp 126-127.5°. $[\alpha]_D^{20} +23.6$ (c, 1 in CHCl_3).

O^1, O^{16}, O^{18} -Tri-Me, 14-(4-methoxybenzoyl), N-Et: **Hemsleyadine**

[183245-57-0]

$\text{C}_{32}\text{H}_{45}\text{NO}_9$ 587.709


Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. Mp 106-107°. $[\alpha]_D^{20} +56.4$ (c, 1 in CHCl_3). Struct. revised in 2005.

Xu, Q.Y. et al., *Chin. Chem. Lett.*, 1996, **7**, 723-724; *CA*, **125**, 322994j (*Hemsleyadine*)

Gao, F. et al., *Heterocycles*, 2005, **65**, 365-370 (*Hemsleyanines A,B, Hemsleyadine, cryst struct*)

Gao, F. et al., *J. Nat. Prod.*, 2007, **70**, 876-879 (*Circinasine A, Hemsleyanidine, Isohemsleyanidine*)

4-Methylnaconitane-1,6,7,8,10,14,16-heptol M-307



$\text{C}_{19}\text{H}_{29}\text{NO}_7$ 383.441

(1 α ,6 β ,14 α ,16 α)-form

7,8-Methylene, O^1, O^{14} -di-Me, N-Et, 6-Ac: **Nordhagenine B**. 16-Epieliasine

$\text{C}_{26}\text{H}_{39}\text{NO}_8$ 493.596

Alkaloid from *Delphinium nordhagenii*. Cryst. (hexane/ Me_2CO). Mp 200-203°. $[\alpha]_D^{30} -3.7$ (c, 0.4 in CHCl_3).

(1 α ,6 β ,14 α ,16 β)-form

O^6, O^{16} -Di-Me, N-Et, 14-benzoyl: **Gadenine**

[89384-03-2]

$\text{C}_{30}\text{H}_{41}\text{NO}_8$ 543.656

Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 147-150°.

O^1, O^6, O^{16} -Tri-Me, N-Et: **10-Hydroxyundicaulidine**

[532993-24-1]

$\text{C}_{24}\text{H}_{39}\text{NO}_7$ 453.575

Alkaloid from *Delphinium excelsum*. $[\alpha]_D^{20} +26.3$ (CHCl_3).

O^1, O^{14}, O^{16} -Tri-Me, N-Et: **Demethylneddelidine**

[63596-60-1]

$\text{C}_{24}\text{H}_{39}\text{NO}_7$ 453.575

Alkaloid from roots of *Delphinium dictyocarpum* (Ranunculaceae). Mp 98-100°.

O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et: **Delterine**

[107814-38-0]

$\text{C}_{25}\text{H}_{41}\text{NO}_7$ 467.601


Alkaloid from the aerial parts of *Delphinium ternatum* (Ranunculaceae).

- Cryst. (Et₂O/hexane). Mp 73-75°.
- 7,8-Methylene, O¹, O¹⁴-di-Me, N-Et, 6-Ac: **Elasine**. 16-Demethyl-deltaline [123064-65-3]
C₂₆H₃₉NO₈ 493.596
Alkaloid from seeds of *Delphinium elatum* (Ranunculaceae). Cryst. (hexane/petrol). Mp 116-118°. [α]_D²⁵ -55.7 (c, 0.51 in CHCl₃).
- 7,8-Methylene, O¹, O¹⁴-di-Me, N-Et, 6,16-di-Ac: **Delretine**. Delpetine [123064-66-4]
C₂₈H₄₁NO₉ 535.633
Alkaloid from *Delphinium retropilosum*. Cryst. (hexane). Mp 218-219°. Synonym misspelt as Delpetine in CA abstract.
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et: **Dictyocarpinine** [50657-27-7]
C₂₄H₃₇NO₇ 451.559
Alkaloid from *Delphinium iliense* and the aerial parts of *Delphinium glaucescens*. Hydrol. prod. of Dictyocarpine (Ranunculaceae). Cryst. (hexane/Me₂CO). Mp 199.5-202°. [α]_D²¹ -4.6 (c, 0.8 in MeOH).
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et, 6-Ac: **Dictyocarpine**. 6-Acetyldictyocarpinine [59989-92-3]
C₂₆H₃₉NO₈ 493.596
Alkaloid from the aerial parts of *Delphinium dictyocarpum*, *Delphinium glaucescens* and *Aconitum columbianum* ssp. *columbianum* (Ranunculaceae). Cryst. (hexane/Me₂CO). Mp 214.5-216.5° (201-203°). [α]_D²⁴ -14.7 (c, 0.97 in CHCl₃). [α]_D²² -12.8 (c, 1.04 in MeOH).
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et, 6,14-di-Ac: **14-Acetyldictyocarpine** [75659-26-6]
C₂₈H₄₁NO₉ 535.633
Alkaloid from aerial parts of *Delphinium barbeyi* (Ranunculaceae). Amorph. solid. Mp 64-69.5°. [α]_D²⁸ -46.6 (c, 1.0 in CHCl₃). [α]_D¹⁹ -50.9 (c, 0.369 in CHCl₃) (natural).
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et, 14-O-(2-methylpropanoyl), 6-Ac: **Glauce-
cerine**. 14-Isobutyryldictyocarpine [78018-29-8]
C₃₀H₄₅NO₉ 563.687
Alkaloid from the aerial parts of *Delphinium glaucescens* (Ranunculaceae). Amorph. [α]_D²⁸ -48.5 (c, 1.5 in CHCl₃). [α]_D²⁷ -27.5 (c, 1.13 in EtOH).
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et, 14-O-(2R-methylbutanoyl), 6-Ac: **Glauce-
nine**. 14-(2-Methylbutyryl)dictyocarpine [78018-27-6]
C₃₁H₄₇NO₉ 577.714
Alkaloid from *Delphinium barbeyi*, *Delphinium glaucescens* and *Delphinium occidentale* (Ranunculaceae). Amorph.; cryst. (MeOH/hexane) (as perchlorate). Mp 227.5-232.5° (perchlorate). [α]_D²⁶ -45 (c, 0.58 in CHCl₃).
- 7,8-Methylene, O¹, O¹⁶-di-Me, N-Et, 14-benzoyl, 6-Ac: **Glaucephine**. 14-Ben-

- zoyldictyocarpine. 6-Acetyl-14-benzoyldictyocarpinine [78018-30-1]
C₃₃H₄₃NO₉ 597.704
Alkaloid from the aerial parts of *Delphinium glaucescens* and above-ground parts of *Delphinium dictyocarpum* (Ranunculaceae). Amorph. [α]_D²⁸ -33.6 (c, 0.76 in CHCl₃).
- 7,8-Methylene, O¹, O¹⁴, O¹⁶-tri-Me, N-Et: **Deltamine**. Eldelidine [6836-10-8]
C₂₅H₃₉NO₇ 465.586
Alkaloid from *Delphinium elatum*, *Delphinium bonvalotii* and *Delphinium dictyocarpum* (Ranunculaceae). Cryst. (EtOH). Mp 226-228°. [α]_D -17.08 (MeOH).
- 7,8-Methylene, O¹, O¹⁴, O¹⁶-tri-Me, N-Et, 6-Ac: **Deltaline**. Delpelatine. Eldeline [6836-11-9]
C₂₇H₄₁NO₈ 507.623
Alkaloid from *Delphinium occidentale*, *Delphinium elatum*, *Delphinium bonvalotii*, *Delphinium dictyocarpum*, *Delphinium glaucescens*, *Delphinium iliense* and *Aconitum columbianum* ssp. *columbianum* (Ranunculaceae). Cryst. (Et₂O). Mp 186.5-188° (180-181°). [α]_D²⁶ -30 (c, 1.1 in MeOH).
▶ AK5544000
- 7,8-Methylene, O¹, O⁶, O¹⁰, O¹⁴, O¹⁶-penta-Me, N-Et: **Terdeline** [119152-46-4]
C₂₇H₄₃NO₇ 493.639
Alkaloid from aerial parts of *Delphinium ternatum* (Ranunculaceae).
- 6-Ketone, 7,8-methylene, O¹, O¹⁴, O¹⁶-tri-Me, N-Et: **Dehydroeldelidine**. Dehydrodeltamine [51856-91-8]
C₂₅H₃₇NO₇ 463.57
Alkaloid from *Delphinium ternatum* (Ranunculaceae).
- Rabinovich, M.S. et al., *Zh. Obshch. Khim.*, 1952, **22**, 1702-1710; *CA*, **47**, 9336e (Deltamine)
- Kuzovkov, A.D. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 2746-2749 (Deltamine)
- Narzullaev, A.S. et al., *Khim. Prir. Soedin.*, 1972, **8**, 498-502; 1973, **9**, 443-444; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 491-493; 1973, **9**, 424-425 (isol, ir, pmr, struct, Dictyocarpine, Dictyocarpinine, Deltamine, Deltaline)
- Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1977, **13**, 128-129; 1978, **14**, 235-241; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 120-121; 1978, **14**, 194-198 (Demethyleneeldelidine)
- Pelletier, S.W. et al., *Can. J. Chem.*, 1980, **58**, 1875-1879 (cmr)
- Pelletier, S.W. et al., *J.A.C.S.*, 1981, **103**, 6536-6538 (config)
- Pelletier, S.W. et al., *J.O.C.*, 1981, **46**, 3284-3293 (*Delphinium glaucescens* constits)
- Finer-Moore, J. et al., *J.O.C.*, 1981, **46**, 3399-3406 (cmr, struct, Glaucephine, Glauce-
cerine, Dictyocarpine)
- Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1981, 530-531; *CA*, **96**, 100872m (isol, ir, pmr, ms, struct, Glaucephine)
- Edwards, O.E. et al., *Can. J. Chem.*, 1982, **60**, 2661-2667 (config)
- González, A.G. et al., *Heterocycles*, 1984, **22**, 17-20 (Gadenine)

- Pelletier, S.W. et al., *J. Nat. Prod.*, 1984, **47**, 643-647 (*Dictyocarpine, cryst struct*)
- Pelletier, S.W. et al., *Heterocycles*, 1985, **23**, 331-338 (*Deltaline, Dictyocarpine*)
- Matveev, V.M. et al., *Khim. Prir. Soedin.*, 1985, **21**, 131-132; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 133-134 (*Dehydroeldelidine*)
- Narzullaev, A.S. et al., *Khim. Prir. Soedin.*, 1986, **22**, 802-803; 1988, **32**, 739-742; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 745-746; 1988, **24**, 335-338 (*Deltaline, Terdeline*)
- Pelletier, S.W. et al., *Phytochemistry*, 1989, **28**, 1521-1525 (*14-Acetyldictyocarpine*)
- Pelletier, S.W. et al., *Tetrahedron*, 1989, **45**, 1887-1892 (*Elasine, Deltaline*)
- Saidkhodzhaeva, S.A. et al., *Khim. Prir. Soedin.*, 1996, **32**, 739-742; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 720-722 (*Delretine, Elasine*)
- Batbayar, N. et al., *Phytochemistry*, 2003, **62**, 543-550 (*10-Hydroxynudicaulidine, Deltaline, isol, cmr*)
- Shaheen, F. et al., *J. Nat. Prod.*, 2006, **69**, 823-825 (*Nordhagenine B*)

4-Methylnaconitane-1,6,7,8,12,14,16-heptol M-308




C₁₉H₂₉NO₇ 383.441

(1α,5β,6β,12β,14α,16β)-form

- O⁶, O¹⁶-Di-Me, N-Et, 14-benzoyl: **Gracine** [107040-83-5]
C₃₀H₄₁NO₈ 543.656
Minor alkaloid from *Delphinium gracile* (Ranunculaceae). Resin. [α]_D +37.6 (c, 0.15 in CHCl₃).
- González, A.G. et al., *Heterocycles*, 1986, **24**, 2753 (ir, pmr, cmr, ms, struct)

4-Methylnaconitane-1,6,7,8,14,16,18-heptol M-309



C₁₉H₂₉NO₇ 383.441

(1α,5β,6α,14α,16β)-form

- O⁸, O¹⁶-Di-Me, N-Et: **14-Deacetyl-18-demethylpubescence** [194149-13-8]
C₂₃H₃₇NO₇ 439.548
Alkaloid from aerial parts of *Aconitella hohenackeri* (*Consolida hohenackeri*). Cryst. (EtOAc/MeOH). Mp 139-140°. [α]_D -10.8 (c, 0.37 in MeOH).

- O^8, O^{16} -*Di-Me*, *N-Et*, *14-Ac*: **18-De-methylpubescenine**
[446859-91-2]
 $C_{25}H_{39}NO_8$ 481.585
Alkaloid from *Consolida orientalis*.
Amorph. solid. $[\alpha]_D^{23} +8$ (c, 0.1 in $CHCl_3$), λ_{max} 202 (log ϵ 3.72); 287 (log ϵ 2.93); 321 (sh) (log ϵ 2.75) (MeOH).
- O^8, O^{16} -*Di-Me*, *N-Et*, *18-benzoyl*: **14-Deacetyl-18-benzoyl-18-demethylpubescenine**
 $C_{30}H_{41}NO_8$ 543.656
Alkaloid from *Consolida orientalis*.
Amorph. solid. $[\alpha]_D^{25} +23.3$ (c, 0.15 in $CHCl_3$).
- O^8, O^{16}, O^{18} -*Tri-Me*, *N-Et*: **14-O-Deacetylpubescenine**
 $C_{24}H_{39}NO_7$ 453.575
Alkaloid from *Consolida orientalis*.
Light yellow oil. $[\alpha]_D^{25} +17.6$ (c, 1 in $CHCl_3$).
- O^8, O^{16}, O^{18} -*Tri-Me*, *N-Et*, *14-Ac*: **Pubescenine**[†]
[116339-93-6]
 $C_{26}H_{41}NO_8$ 495.612
Alkaloid from *Consolida pubescens* (Ranunculaceae) and *Consolida oliveriana*. Cryst. (EtOAc). Mp 227-229°. $[\alpha]_D -1.3$ (c, 0.15 in EtOH).
- $O^8, O^{14}, O^{16}, O^{18}$ -*Tetra-Me*, *N-Et*: **Consolidine**[†]. *14-Deacetylpubescenine 14-methyl ether*
[181798-46-9]
 $C_{25}H_{41}NO_7$ 467.601
Alkaloid from aerial parts of *Consolida oliveriana*. Amorph. $[\alpha]_D +11.8$ (c, 0.17 in $CHCl_3$).
- (1 α ,5 β ,6 β ,14 α ,16 β)-form**
- O^6, O^{16} -*Di-Me*, *N-Et*: **Takaosamine**. *18-O-Demethyldecosine*
[71239-56-0]
 $C_{23}H_{37}NO_7$ 439.548
Alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Mp 178-180°. $[\alpha]_D^{25} +61.3$ (c, 0.2 in $CHCl_3$).
- O^6, O^{16} -*Di-Me*, *N-Et*, *14-Ac*: **14-Acetyltakaosamine**
[725747-04-6]
 $C_{25}H_{39}NO_8$ 481.585
Alkaloid from the aerial parts of *Consolida orientalis*. Amorph. solid. $[\alpha]_D^{25} +25.3$ (c, 0.4 in $CHCl_3$).
- O^6, O^{16} -*Di-Me*, *N-Et*, *14-benzoyl*: **14-O-Benzoyltakaosamine**
 $C_{30}H_{41}NO_8$ 543.656
Alkaloid from *Consolida orientalis*.
Amorph. powder. $[\alpha]_D^{25} +34$ (c, 0.12 in $CHCl_3$).
- O^6, O^{18} -*Di-Me*, *N-Et*: **Macrocentridine**
[121747-79-3]
 $C_{23}H_{37}NO_7$ 439.548
Alkaloid from *Delphinium macrocentrum* (Ranunculaceae). Amorph. off-white solid. $[\alpha]_D^{23} +27$ (c, 1.0 in EtOH).
- O^{16}, O^{18} -*Di-Me*, *N-Et*: **Delphinifoline**
[78346-68-6]
 $C_{23}H_{37}NO_7$ 439.548
Minor alkaloid from *Aconitum delphinifolium* (Ranunculaceae). Mp 218-220°.
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*: **Delectine**
[58480-82-3]
 $C_{24}H_{39}NO_7$ 453.575
Alkaloid from *Delphinium dictyocarpum* (Ranunculaceae). Alkaline hydrolyt. product of Ajadine. Mp 167-169°. $[\alpha]_D +42$ (c, 0.67 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *14-Ac*: **14-O-Acetyldelectine**
[130756-24-0]
 $C_{26}H_{41}NO_8$ 495.612
Alkaloid from *Delphinium nuttallianum* (Ranunculaceae).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*: **Delectine**
[58485-71-5]
 $C_{31}H_{44}N_2O_8$ 572.697
Alkaloid from the roots and aerial parts of *Delphinium dictyocarpum* (Ranunculaceae). Mp 107-109°.
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*, *14-Ac*: **14-Acetyldelectine**. Andersonidine
[66408-08-0]
 $C_{33}H_{46}N_2O_9$ 614.734
Alkaloid from *Delphinium dictyocarpum* and *Delphinium andersonii*. Mp 127-130°. $[\alpha]_D^{27} +39.1$ (c, 0.565 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*, *14-O-(2-methylpropionyl)*: **Trifolioliasine A**
[740815-33-2]
 $C_{35}H_{50}N_2O_9$ 642.788
Alkaloid from *Delphinium trifoliolatum*. Cryst. Mp 125-127°. $[\alpha]_D^{20} +44.2$ (c, 0.53 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*, *14-O-(2R-methylbutanoyl)*: **Jiufengdine**
[486429-07-6]
 $C_{36}H_{52}N_2O_9$ 656.815
Alkaloid from the roots of *Delphinium potaninii* var. *jiufengshanense*. Amorph. powder. $[\alpha]_D +63.8$ (c, 0.5 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-acetamidobenzoyl)*: **N-Acetyldelectine**. *14-Deacetylajadine*
[63596-61-2]
 $C_{33}H_{46}N_2O_9$ 614.734
Alkaloid from the aerial parts of *Delphinium dictyocarpum* and *Delphinium formosum*. Mp 116-118°. $[\alpha]_D +8$ (c, 1.0 in $CHCl_3$). Opt. rotn. refers to the *Delphinium formosum* isolate which was erroneously claimed to be novel.
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-acetamidobenzoyl)*, *14-Ac*: **Ajadine**
[58480-81-2]
 $C_{35}H_{48}N_2O_{10}$ 656.772
Alkaloid from seeds of *Consolida ambigua* (formerly *Delphinium ajacis*). Cryst. (Me₂CO/hexane). Mp 134-136° dec. (sinters at 115°). $[\alpha]_D^{21} +43.9$ (c, 1.0 in EtOH).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-acetamidobenzoyl)*, *14-O-(2-methylpropionyl)*: **14-Deacetyl-14-isobutrylajadine**
[254876-72-7]
 $C_{37}H_{52}N_2O_{10}$ 684.825
Alkaloid from the roots of *Delphinium stapeliosum*. Amorph. $[\alpha]_D^{20} +22.9$ (c, 0.31 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-acetamidobenzoyl)*: **Ajanine**[†]
[152606-90-1]
 $C_{38}H_{54}N_2O_{11}$ 714.851
Alkaloid from leaves of *Delphinium ajacis* (Ranunculaceae). Amorph. $[\alpha]_D +47.37$ (c, 0.1 in $CHCl_3$).
- O^1, O^{14}, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*: **Acoseptrine**. *6-Demethylanthranoyllycoctonine*
[146001-07-2]
 $C_{31}H_{44}N_2O_8$ 572.697
Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Mp 220-222°. $[\alpha]_D -14.3$ (c, 0.063 in $CHCl_3$).
- O^1, O^6, O^{16} -*Tri-Me*, *N-Et*, *18-O-[2S-methylbutanoylamino]benzoyl]*, *14-Ac*: **Delajadine**
[152606-89-8]
 $C_{38}H_{54}N_2O_{10}$ 698.852
Alkaloid from leaves of *Delphinium ajacis* (Ranunculaceae). Amorph. $[\alpha]_D +54$ (c, 0.1 in $CHCl_3$).
- O^1, O^{16}, O^{18} -*Tri-Me*, *N-Et*: **Vaginatine**[†]
[106982-85-8]
 $C_{24}H_{39}NO_7$ 453.575
Alkaloid from the roots of *Aconitum scaposum* var. *vaginatum* (Ranunculaceae). Mp 86-88°. $[\alpha]_D^{28} +25.3$ (c, 0.1 in $CHCl_3$).
- O^1, O^{16}, O^{18} -*Tri-Me*, *N-Et*, *6-Ac*: **Alboviolaconitine A**
[138822-60-3]
 $C_{26}H_{41}NO_8$ 495.612
Alkaloid from roots of *Aconitum alboviolaceum* (Ranunculaceae). Mp 125-126°. $[\alpha]_D^{28} +16.2$ (c, 2.0 in $CHCl_3$).
- O^6, O^8, O^{16} -*Tri-Me*, *N-Et*: **Potanine**
[160568-11-6]
 $C_{24}H_{39}NO_7$ 453.575
Alkaloid from roots of *Delphinium potaninii*.
- O^6, O^{14}, O^{16} -*Tri-Me*, *N-Et*: **Gigactonine**. *Paniculine*[†]
[65967-20-6]
 $C_{24}H_{39}NO_7$ 453.575
Minor alkaloid from the roots of *Aconitum gigas* and from *Consolida regalis* ssp. *paniculata* (Ranunculaceae). Cryst. + 0.2 Me₂CO. Mp 168-169°. $[\alpha]_D^{22} +49$ (c, 1 in EtOH). Paniculine appears to be identical with Gigactonine (ref. inaccessible). The CA abstract omits the 14-OMe group.
- O^6, O^{14}, O^{16} -*Tri-Me*, *N-Et*, *18-Ac*: **1-O-Demethyltricornine**
 $C_{26}H_{41}NO_8$ 495.612
Alkaloid from *Consolida orientalis*. Amorph. solid. $[\alpha]_D^{25} +18.7$ (c, 0.16 in $CHCl_3$).
- O^6, O^{14}, O^{16} -*Tri-Me*, *N-Et*, *18-O-(2-aminobenzoyl)*: **Isodelectine**
[133034-09-0]
 $C_{31}H_{44}N_2O_8$ 572.697
Alkaloid from *Delphinium vestitum* (Ranunculaceae). Amorph. $[\alpha]_D +54$ (c, 0.345 in $CHCl_3$).
- O^6, O^{14}, O^{18} -*Tri-Me*, *N-Et*: **16-O-Demethyldecosine**

- [239479-43-7]
 $C_{24}H_{39}NO_7$ 453.575
 Alkaloid from *Delphinium fangshanense*. Amorph. powder. $[\alpha]_D^{25} +41.5$ (c, 0.4 in $CHCl_3$).
- O^6, O^{16}, O^{18} -Tri-Me, N-Et: **Delcosine**. Iliensine†. Lucaconine. Alkaloid C†. Takaobase I. Takawobase I. *Delphinamine* [545-56-2]
 $C_{24}H_{39}NO_7$ 453.575
 Alkaloid from *Delphinium consolida*, *Delphinium biternatum*, *Delphinium glaucescens*, *Delphinium tatsienense*, *Aconitum ibukiense*, *Aconitum japonicum*, *Consolida ambigua* (*Delphinium ajacis*), and others (Ranunculaceae). Fine cryst. (EtOH). Mp 204-205°. $[\alpha]_D^{25} +54.1$ (c, 0.8 in $CHCl_3$).
- ▶ AR5564000
 O^6, O^{16}, O^{18} -Tri-Me, N-Et, hydroiodide: Mp 196-197°.
- O^6, O^{16}, O^{18} -Tri-Me, N-Et, 14-Ac: **14-Acetyldelcosine**. Alkaloid B† [50676-21-6]
 $C_{26}H_{41}NO_8$ 495.612
 Alkaloid from the seeds of *Consolida ambigua* (formerly *Delphinium ajacis*) (Ranunculaceae). Cubes (Et₂O/petrol). Mp 193-194°. $[\alpha]_D +30.4$ (c, 1.0 in EtOH).
- O^6, O^{16}, O^{18} -Tri-Me, N-Et, 14-benzoyl: **14-Benzoyldelcosine** [66891-16-5]
 $C_{31}H_{43}NO_8$ 557.683
 Alkaloid from the aerial parts of *Delphinium biternatum* (Ranunculaceae). Mp 147-149°. $[\alpha]_D +50$.
- O^8, O^{16}, O^{18} -Tri-Me, N-Et: **Deacetyl-6-epipubescenine** [130774-03-7]
 $C_{24}H_{39}NO_7$ 453.575
 Minor alkaloid from *Delphinium nuttallianum*.
- O^8, O^{16}, O^{18} -Tri-Me, N-Et, 14-Ac: **6-Epipubescenine** [125263-90-3]
 $C_{26}H_{41}NO_8$ 495.612
 Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Amorph. solid. $[\alpha]_D^{25} +23$ ($CHCl_3$).
- O^{14}, O^{16}, O^{18} -Tri-Me, N-Et: **Exceconitine** [229159-16-4]
 $C_{24}H_{39}NO_7$ 453.575
 Alkaloid from *Aconitum excelsum*. Cryst. Mp 202-204°.
- O^{14}, O^{16}, O^{18} -Tri-Me, N-Et, 6-Ac: **6-O-Acetyl-14-O-methyldephiniifoline** [229159-13-1]
 $C_{26}H_{41}NO_8$ 495.612
 Alkaloid from the flowers of *Aconitum lycoctonum*. Oil.
- O^1, O^6, O^{14}, O^{16} -Tetra-Me: **Davidisine A**. N-Deethyldeisine. N-Deethyllycoctonine [26871-62-5]
 $C_{23}H_{37}NO_7$ 439.548
 Alkaloid from *Delphinium davidii*.
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-formyl: **Potanisine A**. **Davidisine B** [187653-15-2]
 [936116-23-3 (Davidisine B)]
- $C_{24}H_{37}NO_8$ 467.558
 Alkaloid from *Delphinium davidii* and *Delphinium potaninii*. Amorph. powder. $[\alpha]_D +53.9$ (c, 1.78 in $CHCl_3$).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et: **Lycocotinine**. **Deisine**. **Royline**. **Lycocotinine** [26000-17-9]
 $C_{25}H_{41}NO_7$ 467.601
 Alkaloid from *Delphinium barbeyi*, *Delphinium tricorne*, *Inula royleana*, *Aconitum lycoctonum* and many other *Delphinium* and *Aconitum* spp. (Ranunculaceae, Asteraceae). Mp 151-153°. $[\alpha]_D^{20} +50$ (+53) (EtOH).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, hydrochloride: Mp 165°.
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-Ac: **Tricornine** [26871-60-3]
 $C_{27}H_{43}NO_8$ 509.639
 Alkaloid from *Delphinium tricorne* (whole plant) (Ranunculaceae). Mp 187-189° dec. $[\alpha]_D^{22} +47.3$ (c, 1.3 in EtOH).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, 18-O-(2-aminobenzoyl): N-Deethylanthranolyllycoctonine. **Jiufengtine** [486429-11-2]
 $C_{30}H_{42}N_2O_8$ 558.67
 Alkaloid from the roots of *Delphinium potaninii* var. *jiufengshanense*. Amorph. powder. $[\alpha]_D +46$ (c, 0.5 in $CHCl_3$).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-(2-aminobenzoyl): **Anthranolyllycoctonine**. **Inuline†** [22413-78-1]
 $C_{32}H_{46}N_2O_8$ 586.724
 Alkaloid from *Inula royleana*, *Delphinium consolida*, *Delphinium barbeyi*, *Delphinium elisabethae*, *Delphinium grandiflorum*, *Delphinium tamarae*, *Delphinium biternatum*, *Delphinium cashmirianum*, *Delphinium dictyocarpum*, *Delphinium glaucescens*, *Delphinium confusum*, *Consolida ambigua* (*Delphinium ajacis*), *Aconitum umbrosum* and others (Asteraceae, Ranunculaceae). Amorph. solid. Mp 160-165° (132-135°, 145-154° dec.). $[\alpha]_D^{20.5} +50.6$ (c, 0.67 in $CHCl_3$). $[\alpha]_D^{27} +51.2$ (c, 0.519 in EtOH).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Ajacine** [509-17-1]
 $C_{34}H_{48}N_2O_9$ 628.761
 Alkaloid from *Delphinium orientale*, *Delphinium belladonna*, *Consolida ambigua* (*Delphinium ajacis*) and *Aconitum umbrosum* (Ranunculaceae). Cryst. (EtOH aq.). Mp 142-143° (137-139°). $[\alpha]_D +50$ (EtOH).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-[2-(ethoxycarbonylamino)benzoyl]: **Omeienine** [180338-87-8]
 $C_{35}H_{50}N_2O_{10}$ 658.787
 Alkaloid from roots of *Delphinium omeiense*. Amorph. powder. $[\alpha]_D +40.4$ (c, 0.5 in $CHCl_3$).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-(2-methylaminobenzoyl): **Septerine** [192992-36-2]
- $C_{33}H_{48}N_2O_8$ 600.751
 Alkaloid from *Aconitum septentrionale* (Ranunculaceae). Powder (as perchlorate).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-[2-(2-methylpropanoylamino)benzoyl]: **Delajacirine** [152606-88-7]
 $C_{36}H_{52}N_2O_9$ 656.815
 Alkaloid from leaves of *Delphinium ajacis* (Ranunculaceae). Amorph. $[\alpha]_D +39.62$ (c, 0.2 in $CHCl_3$).
- O^1, O^6, O^{14}, O^{16} -Tetra-Me, N-Et, 18-O-[2-(2S-methylbutanoylamino)benzoyl]: **Conambine**. **Delajacine** [152606-87-6]
 $C_{37}H_{54}N_2O_9$ 670.842
 Alkaloid from the seeds of *Consolida ambigua* and leaves of *Delphinium ajacis*. Cryst. Mp 127-129° Mp 73-76°. $[\alpha]_D +47.3$ (c, 0.3 in $CHCl_3$) (Conambine). $[\alpha]_D +58.9$ (c, 0.1 in $CHCl_3$) (Delajacine). Conambine and Delajacine not compared. Higher Mp refers to Conambine from *C. ambigua*.
- O^1, O^6, O^{14}, O^{18} -Tetra-Me, N-Et: **Delbiterine**. **16-Demethyldephatine** [66891-14-3]
 $C_{25}H_{41}NO_7$ 467.601
 Alkaloid from the roots of *Delphinium biternatum* (Ranunculaceae). Mp 137-138°.
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-formyl: **α -Oxobrowniine**. N-Deethyl-N-formylbrowniine [128341-39-9]
 $C_{24}H_{37}NO_8$ 467.558
 Alkaloid from the rhizomes of *Aconitum yesoense* var. *macrocysoense* (Ranunculaceae). Amorph. $[\alpha]_D +70.7$ (c, 0.15 in EtOH). The name α -Oxobrowniine is misleading.
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et: **Browniine** [5140-42-1]
 $C_{25}H_{41}NO_7$ 467.601
 Alkaloid from *Delphinium brownii*, *Delphinium biternatum*, *Delphinium carolinianum*, *Delphinium glaucescens*, *Delphinium iliense*, *Delphinium virescens*, *Delphinium tatsienense* and *Consolida ambigua* (formerly *Delphinium ajacis*) (Ranunculaceae). Amorph. solid; cryst. (as hydroiodide). Mp 193-196° hydroiodide. $[\alpha]_D^{25} +40.2$ (c, 2.5 in EtOH).
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-Ac: **14-Acetylbrowniine** [65601-04-9]
 $C_{27}H_{43}NO_8$ 509.639
 Alkaloid from the aerial portions of *Delphinium brownii* (both early shoots and mature flowering stages), the above-ground parts of *Delphinium oreophilum*, and from the seeds of *Consolida ambigua* (*Delphinium ajacis*) (Ranunculaceae). Cryst. (hexane). Mp 123-124° (115-116°). $[\alpha]_D^{27} +27.8$ (c, 1.0 in $CHCl_3$).
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, O^{14} -(2-methylpropanoyl): **Occidentalidine** [116199-47-4]


- C₂₉H₄₇NO₈ 537.692
Alkaloid from the aerial parts of *Delphinium occidentale* (Ranunculaceae). Mp 137-140°. [α]_D²² +31.3 (c, 0.32 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, O¹⁴-(2S-methylbutanoyl): **Glaucedine**. 14-(2-Methylbutyryl)browniine [78039-66-4]
C₃₀H₄₉NO₈ 551.719
Alkaloid from the aerial parts of *Delphinium glaucescens* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 117-120°. [α]_D²⁷ +36.4 (c, 0.81 in MeOH).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl: **14-Benzoylbrowniine** [66891-15-4]
C₃₂H₄₅NO₈ 571.709
Alkaloid from the aerial parts of *Delphinium biternatum* (Ranunculaceae).
- O¹, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et: **6-De-methyldephatine**. Demethylenedelcorine. Acosanine [52358-56-2]
C₂₅H₄₁NO₇ 467.601
Alkaloid from roots of *Aconitum septentrionale* and *Aconitum sajanense* (Ranunculaceae). Amorph. [α]_D +19.3 (c, 0.295 in CHCl₃).
- O¹, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et, 6-Ac: **Leucostine A**. 6-O-Acetyldemethylenedelcorine [173075-39-3]
C₂₇H₄₃NO₈ 509.639
Alkaloid from roots of *Aconitum leucostomum* and flowers of *Aconitum lycoctonum*. Amorph. powder or oil. Leucostine A and 6-Acetyldemethylenedelcorine not compared.
- O⁶, O⁸, O¹⁶, O¹⁸-Tetra-Me, N-Et: **Deltatine** [92631-66-8]
C₂₅H₄₁NO₇ 467.601
Alkaloid from *Consolida orientalis* and *Delphinium tatsienense*. Amorph. powder + 1H₂O. [α]_D²⁰ +28.6 (c, 2.4 in EtOH).
- O⁶, O⁸, O¹⁴, O¹⁶-Tetra-Me, N-Et, 18-O-(2-aminobenzoyl): **Delvestine** [99815-77-7]
C₃₂H₄₆N₂O₈ 586.724
Alkaloid from the aerial parts of *Delphinium vestitum* (Ranunculaceae). Mp 185-187°. [α]_D²⁷ +18.6 (c, 0.8 in CHCl₃).
- O⁶, O⁸, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-Ac: **Delbonine** [95066-33-4]
C₂₇H₄₃NO₈ 509.639
Alkaloid from *Consolida orientalis* and *Delphinium bonvalotii*. Amorph. solid. [α]_D¹⁶ +41.1 (c, 0.1 in CHCl₃).
- O⁶, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et: **Delso-line**. 14-O-Methyldecosine. 18-O-Methylgigactonine. Acomonine [509-18-2]
C₂₅H₄₁NO₇ 467.601
Alkaloid from *Delphinium consolida*, *Delphinium orientale*, *Delphinium biternatum*, *Delphinium yunnanense*, *Aconitum finetianum*, *Aconitum monticola*, *Aconitum karakolicum*, *Consolida ambigua* (*Delphinium ajacis*) and others (Ranunculaceae). Cryst. (MeOH). Mp 212-213°. [α]_D²⁴ +54.3 (c, 1.0 in CHCl₃).
- ▶ AR5569490
O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et: **Deacetylabiguine** [74690-96-3]
C₂₆H₄₃NO₇ 481.628
Alkaloid from roots of *Delphinium tatsienense*. Amorph. [α]_D²⁵ +36.6 (c, 0.38 in EtOH).
- O¹, O⁶, O⁸, O¹⁴, O¹⁶-Penta-Me, N-Et, 18-O-(2-aminobenzoyl): **Delvestidine** [99815-78-8]
C₃₃H₄₈N₂O₈ 600.751
Alkaloid from the aerial parts of *Delphinium vestitum* (Ranunculaceae). Amorph. [α]_D²⁵ +22.1 (c, 0.515 in CHCl₃).
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-Ac: **Ambiguine** [68388-52-3]
C₂₈H₄₅NO₈ 523.665
Minor alkaloid from seeds of *Consolida ambigua* (formerly *Delphinium ajacis*). Mp 106-108°. [α]_D²⁵ +38 (c, 1.0 in CHCl₃).
- O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-Penta-Me: N-**Deethyldephatine** [173559-78-9]
C₂₄H₃₉NO₇ 453.575
Alkaloid from aerial parts of *Aconitum orientale* (Ranunculaceae). [α]_D +52 (c, 1 in CHCl₃).
- O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-Penta-Me, N-Et: **Delphatine**. Delsonine. 18-O-Methyllycoctonine [25488-62-4]
C₂₆H₄₃NO₇ 481.628
Alkaloid from the roots, seeds and aerial parts of *Delphinium biternatum*, from *Delphinium consolida* and from the seeds of *Consolida ambigua*, (formerly *Delphinium ajacis*) (Ranunculaceae). Amorph. solid. [α]_D²⁹ +38.2 (c, 2.95 in CHCl₃). Identity of Delsonine (from *Delphinium consolida*) with Delphatine is not conclusively established.
- O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-Penta-Me, N-Et, perchlorate: Mp 212-214° (205-207°). [α]_D³² +23 (c, 0.412 in MeOH).
- O⁶, O⁷, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et: **Delosine** [68463-98-9]
C₂₆H₄₃NO₇ 481.628
Alkaloid from *Delphinium ajacis*.
- O⁶, O⁷, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-Ac: **14-Acetyldeposine** [67983-95-3]
C₂₈H₄₅NO₈ 523.665
Alkaloid from *Consolida ambigua*.
- O⁶, O⁸, O¹⁴, O¹⁶, O¹⁸-Penta-Me, N-Et: **Delbotine** [92631-67-9]
C₂₆H₄₃NO₇ 481.628
Alkaloid from the roots of *Delphinium bonvalotii*. Mp 155-157°. [α]_D¹⁸ +13.6 (c, 0.1 in CHCl₃).
- Hepta-Me ether*, N-Et: **Delcoline** [68463-99-0]
C₂₈H₄₇NO₇ 509.682
Alkaloid from *Delphinium confusum*.
- 7,8-Methylene, O⁶, O¹⁶-di-Me, N-Et: **Campyloine** [960201-80-3]
C₂₄H₃₇NO₇ 451.559
Alkaloid from *Delphinium campylocentrum*.
- 7,8-Methylene ether, O¹⁴, O¹⁸-di-Me, N-Et: **Delcorinine** [404945-91-1]
C₂₄H₃₇NO₇ 451.559
Alkaloid from *Delphinium corymbosum*. Cryst. (Me₂CO). Mp 226-228°.
- 7,8-Methylene ether, O¹⁶, O¹⁸-di-Me, N-Et: **Delbruninol** [244619-04-3]
C₂₄H₃₇NO₇ 451.559
Alkaloid from *Delphinium brunonianum*. [α]_D²⁵ -3.6 (c, 0.01 in CHCl₃).
- 7,8-Methylene ether, O¹, O¹⁶, O¹⁸-tri-Me, N-Et: **Delcoridine** [76971-29-4]
C₂₅H₃₉NO₇ 465.586
Alkaloid from the aerial parts of *Delphinium iliense* (Ranunculaceae). Mp 222-223° (as perchlorate).
- 7,8-Methylene ether, O⁶, O¹⁶, O¹⁸-tri-Me, N-Et: **Delbrunine** [106982-83-6]
C₂₅H₃₉NO₇ 465.586
Alkaloid from the whole plant of *Delphinium brunonianum* (Ranunculaceae). Mp 178°. [α]_D¹⁷ 0 (c, 0.085 in EtOH).
- 7,8-Methylene ether, O¹, O⁶, O¹⁶, O¹⁸-tetra-Me, N-Et: **Delbruline** [106982-82-5]
C₂₆H₄₁NO₇ 479.612
Alkaloid from *Delphinium brunonianum* (Ranunculaceae). Mp 129-131°. [α]_D¹⁷ 0 (c, 0.09 in CHCl₃).
- 7,8-Methylene ether, O¹, O¹⁴, O¹⁶, O¹⁸-tetra-Me, N-Me, 6-Ac: **Souline** [259653-97-9]
C₂₇H₄₁NO₈ 507.623
Alkaloid from *Delphinium souliei*. Cryst. (EtOH). Mp 90-91°. [α]_D²⁴ +35 (c, 0.2 in CHCl₃).
- 7,8-Methylene ether, O¹, O¹⁴, O¹⁶, O¹⁸-tetra-Me, N-Et: **Delcorine** [52358-55-1]
C₂₆H₄₁NO₇ 479.612
Alkaloid from the aerial parts of *Delphinium corumbosum* (*corymbosum*), *Delphinium iliense* and *Delphinium ternatum* (Ranunculaceae). Mp 200-202°. [α]_D -18 (CHCl₃).
- 7,8-Methylene ether, O¹, O¹⁴, O¹⁶, O¹⁸-tetra-Me, N-Et, 6-Ac: **Tortumine** [51856-89-4]
C₂₈H₄₃NO₈ 521.65
Alkaloid from *Consolida hohenackeri* (Ranunculaceae).
- 7,8-Methylene ether, O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-penta-Me, N-Et: **Delbrusine**. 6-O-Methyldecorine [76971-31-8]
C₂₇H₄₃NO₇ 493.639
Alkaloid from *Aconitum septentrionale* and *Delphinium brunonianum* (Ranun-

- culaceae). Mp 141°. $[\alpha]_D^{25} +16.8$ (c, 0.57 in CHCl_3).
- 6-Ketone, $\text{O}^1, \text{O}^{14}, \text{O}^{16}, \text{O}^{18}$ -tetra-Me, N-Et: **Dehydroacosanine** [192386-07-5] $\text{C}_{25}\text{H}_{39}\text{NO}_7$ 465.586
Alkaloid from *Aconitum sajanense* (Ranunculaceae) and *Aconitum leucostomum*. Cryst. (MeOH/hexane). Mp 140-141°.
- 6-Ketone, 7,8-methylene ether, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **Ilidine** [66921-48-0] $\text{C}_{25}\text{H}_{37}\text{NO}_7$ 463.57
Alkaloid from above ground parts of *Delphinium iliense* (Ranunculaceae). Mp 141-143°.
- 6-Ketone, 7,8-methylene ether, $\text{O}^{14}, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **Glandulosine**† [123715-13-9] $\text{C}_{25}\text{H}_{37}\text{NO}_7$ 463.57
Alkaloid from *Consolida glandulosum* (Ranunculaceae).
- 6-Ketone, 7,8-methylene ether, $\text{O}^1, \text{O}^{14}, \text{O}^{16}, \text{O}^{18}$ -tetra-Me, N-Et: **6-Dehydrodelcorine** [51856-90-7] $\text{C}_{26}\text{H}_{39}\text{NO}_7$ 477.597
Alkaloid from the aerial parts of *Delphinium iliense* (Ranunculaceae). Mp 133-134°.
- 14-Ketone, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **Vaginoline** [106982-86-9] $\text{C}_{24}\text{H}_{37}\text{NO}_7$ 451.559
Alkaloid from *Aconitum scaposum* var. *vaginatatum* (Ranunculaceae). Mp 209-213° dec. $[\alpha]_D^{25} +28.6$ (c, 0.1 in EtOH).
- 14-Ketone, $\text{O}^6, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **14-Dehydrodelcosine**. *Shimoburo* base II. 14-Dehydroilidensine [1361-18-8] $\text{C}_{24}\text{H}_{37}\text{NO}_7$ 451.559
Alkaloid from *Aconitum japonicum*, roots of *Aconitum ibukiense* and aerial parts of *Delphinium biternatum* (Ranunculaceae). Mp 208-213°. $[\alpha]_D +25.2$ (CHCl_3).
- 14-Ketone, $\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -tetra-Me, N-Et: **14-Dehydrobrowniine** [4829-56-5] $\text{C}_{25}\text{H}_{39}\text{NO}_7$ 465.586
Alkaloid from *Delphinium cardinale*, the aerial parts of *Delphinium biternatum* and *Delphinium glaucescens*, and roots of *Aconitum ibukiense* (Ranunculaceae). Small prisms (Et_2O). Mp 161-163°. $[\alpha]_D^{27} +19$ (EtOH).
- 6,14-Diketone, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **Vaginadine** [106982-87-0] $\text{C}_{24}\text{H}_{35}\text{NO}_7$ 449.543
Alkaloid from *Aconitum scaposum* var. *vaginatatum* (Ranunculaceae). Mp 147-149°. $[\alpha]_D^{18.5} -49.4$ (c, 0.1 in EtOH).
- (1β,5β,6β,14α,16β)-form**
7,8-Methylene ether, $\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -tetra-Me, N-Et: **Corepanine** [119459-59-5] $\text{C}_{26}\text{H}_{41}\text{NO}_7$ 479.612
Alkaloid from epigeal parts of *Consolida regalis* ssp. *paniculata* (Ranunculaceae). The CAS abstract diag. omits the 14-OH group.
- Goodson, J.A. et al., *J.C.S.*, 1945, 245-246 (*Delcosine, 14-Acetyldecosine, isol*)
Marion, L. et al., *J.A.C.S.*, 1947, **69**, 2010-2014 (*Delsonine*)
Cook, W.B. et al., *J.A.C.S.*, 1952, **74**, 1411-1415 (*Anthranoyllycoctonine*)
Taylor, W.I. et al., *Can. J. Chem.*, 1954, **32**, 780-784 (*14-Acetyldecosine, struct*)
Abubakirov, N.K. et al., *CA*, 1955, **49**, 5495 (*Delphatine*)
Przybylska, M. et al., *Can. J. Chem.*, 1956, **34**, 185-187 (*Lycocotinine*)
Edwards, O.E. et al., *Can. J. Chem.*, 1956, **34**, 1315-1328; 1959, **37**, 1187-1190; 1982, **60**, 2661-2667 (*Anthranoyllycoctonine, struct, config*)
Sparatore, F. et al., *Tetrahedron*, 1958, **4**, 157-168 (*Delsoline*)
Kuzovkov, A.D. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 2746-2749 (*Anthranoyllycoctonine, Ajacine*)
Skarić, V. et al., *Can. J. Chem.*, 1960, **38**, 2433-2440 (*Delcosine, Delsoline, struct*)
Benn, M.H. et al., *Can. J. Chem.*, 1963, **41**, 477-452; 1966, **44**, 1-8 (*Browniine, 14-Acetylbrowniine, Dehydrobrowniine*)
Yunusov, M.S. et al., *Khim. Prir. Soedin.*, 1970, **6**, 334-339; 1975, **11**, 770-775; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 9; 1975, **11**, 791-794 (*Delphatine, Delcosine, ir, pmr, ms, struct*)
Jones, A.J. et al., *Can. J. Chem.*, 1973, **51**, 486-499 (*Browniine, cmr*)
Narzullaev, A.S. et al., *Khim. Prir. Soedin.*, 1973, **9**, 497-501; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 468-471 (*Delcorine*)
Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1975, **11**, 665-666; 1977, **13**, 128-129; 1978, **14**, 235-241; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 704-705; 1977, **13**, 120-121; 1978, **14**, 194-198 (*Delectinine, Delectine, 14-Acetyldelectine*)
Pelletier, S.W. et al., *Heterocycles*, 1977, **7**, 327-329; 1983, **20**, 1347-1354 (*Delcosine, Delsoline, 14-Acetylbrowniine, Lycocotinine, cmr*)
Sakai, S. et al., *Heterocycles*, 1977, **8**, 207-210 (*Gigactonine*)
Zhamierashvili, M.G. et al., *Khim. Prir. Soedin.*, 1977, **13**, 836-838; 1980, **16**, 663-665; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 704-706; 1980, **16**, 479-480 (*Delcorine, Delcoridine, 6-Dehydrodelcorine, Ilidine*)
Kazlikhin, V.G. et al., *Khim. Prir. Soedin.*, 1977, **13**, 869; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 737-738 (*14-Acetylbrowniine*)
Pelletier, S.W. et al., *Phytochemistry*, 1977, **16**, 1464 (*Tricornine*)
Waller, G.R. et al., *CA*, 1978, **89**, 176315 (*Delosine*)
Pelletier, S.W. et al., *Heterocycles*, 1978, **9**, 463-468 (*Delectinine, cmr*)
Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1978, **14**, 106-111; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 84-88 (*14-Benzoylbrowniine, 14-Benzoyldecosine, 14-Dehydrodelcosine, Delbitrine*)
Aiyar, V.N. et al., *Phytochemistry*, 1978, **17**, 1453-1454 (*14-Acetylbrowniine*)
Pelletier, S.W. et al., *Can. J. Chem.*, 1979, **57**, 1652-1655 (*Browniine, cmr*)
Pelletier, S.W. et al., *J. Nat. Prod.*, 1979, **42**, 615-623; 1980, **43**, 395-406 (*14-Acetylbrowniine, 14-Acetyldecosine, Delphatine, Delsoline, Ambiguine, Ajacine, Anthranoyllycoctonine, Browniine*)
Sakai, S. et al., *Yakugaku Zasshi*, 1979, **99**, 647-656; *CA*, **91**, 105185w (*Takaosamine*)
Pelletier, S.W. et al., *J.A.C.S.*, 1981, **103**, 6536-6538 (*Delcorine, Delsoline, Deltatsine, cmr, config*)
Pelletier, S.W. et al., *J.O.C.*, 1981, **46**, 3284-3293 (*Glaucidine*)
Finer-Moore, J. et al., *J.O.C.*, 1981, **46**, 3399-3406 (*Delcosine, cmr, struct*)
Pelletier, S.W. et al., *Tet. Lett.*, 1981, **22**, 207-210 (*Delsoline, 14-Dehydrodelcosine*)
Aiyar, V.N. et al., *Tet. Lett.*, 1981, **22**, 483-484 (*Delphinifoline*)
Kerr, K.A. et al., *Acta Cryst. B*, 1982, **38**, 1237-1241 (*Delphinifoline, cryst struct*)
Pelletier, S.W. et al., *Heterocycles*, 1983, **20**, 1347-1354 (*Deacetylbrowniine*)
Matveev, V.M. et al., *Khim. Prir. Soedin.*, 1983, **19**, 657-658; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 627-628 (*Delcorine*)
Joshi, B.S. et al., *Heterocycles*, 1984, **22**, 2037-2042 (*Deltatsine*)
Pelletier, S.W. et al., *J. Nat. Prod.*, 1984, **47**, 643-647 (*Browniine, cryst struct*)
Sakai, S. et al., *Yakugaku Zasshi*, 1984, **104**, 222-237; *CA*, **101**, 69329k (*isol, Dehydrobrowniine*)
Jiang, Q.P. et al., *Heterocycles*, 1985, **23**, 11-15; 1986, **24**, 877-879 (*Delbonine, Delbotine, Vaginatine, Vaginaline, Vaginadine*)
Desai, H.K. et al., *Heterocycles*, 1985, **23**, 2483-2487 (*Delvestine, Delvestidine*)
Deng, W. et al., *Heterocycles*, 1986, **24**, 873-876 (*Delbruline, Delbrusine, Delbruline*)
Pelletier, S.W. et al., *Heterocycles*, 1987, **26**, 2835-2840 (*Delbruline, Delbruline, synth*)
Vaisov, Z.M. et al., *Khim. Prir. Soedin.*, 1987, **23**, 869-872; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 725-727 (*Delcoline*)
Sener, B. et al., *CA*, 1988, **110**, 111674g; **111**, 211906y; 1990, **112**, 175528 (*Paniculine, Glandulosine, Corepanine, Tortumine*)
Kulanthaivel, P. et al., *Heterocycles*, 1988, **27**, 339-342 (*Occidentalidine*)
De la Fuente, G. et al., *Tet. Lett.*, 1988, **29**, 2723-2736 (*Pubescenine, isol, ir, pmr, cmr, ms, cryst struct*)
Pelletier, S.W. et al., *Heterocycles*, 1989, **28**, 107-110 (*Andersonidine*)
Bai, Y. et al., *Heterocycles*, 1989, **29**, 1017-1021; 1990, **31**, 1233-1236 (*14-Acetyldelectinine, Pubescenine, 6-Epipubescenine, Deacetyl-6-epipubescenine*)
Wada, K. et al., *Heterocycles*, 1989, **29**, 2141-2148 (*α-Oxobrowniine*)
Benn, M.H. et al., *Phytochemistry*, 1989, **28**, 919-922 (*Macrocentridine*)
Bhandary, K.K. et al., *Acta Cryst. C*, 1990, **46**, 1704-1707 (*Delvestine, cryst struct*)
Desai, H.K. et al., *J. Nat. Prod.*, 1990, **53**, 1606-1608 (*Isodelectine*)
Chen, D. et al., *Huaxue Xuebao*, 1992, **50**, 1211-1218; *CA*, **118**, 230146m (*Albiviolalconitine A*)
Joshi, B.S. et al., *J. Crystallogr. Spectrosc. Res.*, 1992, **22**, 477-483 (*Delsoline, cryst struct*)
Sayed, H.M. et al., *J. Nat. Prod.*, 1992, **55**, 1595-1606 (*6-Deethyldephatine, Aceoseprinine*)
Vaisov, Z.M. et al., *Khim. Prir. Soedin.*, 1992, **28**, 531-534; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 463-466 (*6-Dehydroacosanine*)
Lu, J. et al., *J. Nat. Prod.*, 1993, **56**, 2098-2103 (*Ajanine, Delajacine, Delajadine, Delajacrine*)
Pu, H.Y. et al., *Chin. Chem. Lett.*, 1994, **5**, 939-940; *CA*, **122**, 106208b (*Potanine*)
Ulubelen, A. et al., *J. Nat. Prod.*, 1995, **58**, 1555-1561; 1996, **59**, 907-910 (*Lycocotnone, Consolidine, pmr, cmr*)
Xu, Q.Y. et al., *Chin. Chem. Lett.*, 1996, **7**, 555-556 (*Omeienine*)

- Yue, J. et al., *J. Nat. Prod.*, 1996, **59**, 277-279 (*Leucostines*)
- Usmanova, S.K. et al., *Khim. Prir. Soedin.*, 1996, **32**, 225-228; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 198-200 (*Septerine*)
- Ulubelen, A. et al., *Phytochemistry*, 1996, **41**, 957-961 (*N-Deethyldephatine*)
- Meriçli, F. et al., *Phytochemistry*, 1996, **42**, 1249-1251 (*N-Acetyldelectine*)
- Pu, H. et al., *Planta Med.*, 1996, **62**, 462-464 (*Potansine A*)
- Wei, X. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1996, **38**, 995-997 (*Dehydroacosanine*)
- Almanza, G. et al., *Phytochemistry*, 1997, **45**, 1079-1085 (*14-Deacetyl-18-demethylpubescence*)
- Desai, H.K. et al., *Heterocycles*, 1998, **48**, 1343-1346 (*Conambine*)
- Zhang, S.X. et al., *Chin. Chem. Lett.*, 1999, **10**, 133-134 (*Execonitine*)
- He, L. et al., *Chin. Chem. Lett.*, 1999, **10**, 1027-1028 (*Soulidine*)
- Ulubelen, A. et al., *Heterocycles*, 1999, **51**, 1897-1903 (*Delbruninol*)
- Chen, Y. et al., *J. Nat. Prod.*, 1999, **62**, 798-799 (*6-O-Acetyldemethylnedelcorine*, *6-O-Acetyl-14-O-methyldephinitofoline*)
- Zhang, S. et al., *Phytochemistry*, 1999, **51**, 333-336 (*16-Demethyldeolsone*)
- Wada, K. et al., *Chem. Pharm. Bull.*, 2000, **48**, 660-665 (*Delcosine*, *ms*)
- Shrestha, P.M. et al., *J. Nat. Prod.*, 2000, **63**, 2-5; 2004, **67**, 1574-1576 (*Ajadine*, *Deacetylisobutyryljadine*, *Delbonine*, *Deltatsine*, *cmr*)
- Salimov, B.T. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 272-273 (*Delcorinine*)
- Shen, X.-L. et al., *Chem. Pharm. Bull.*, 2002, **50**, 1265-1267 (*Glaucedine*, *Jiufengine*, *Jiufengdine*)
- Hohman, J. et al., *J. Nat. Prod.*, 2002, **65**, 1069-1072 (*Gigactonine*, *Takaosamine*, *18-Demethylpubescence*)
- Zhou, X.-L. et al., *Chem. Pharm. Bull.*, 2004, **52**, 381-383 (*Trifoliolasmine A*, *Glaucedine*, *Occidentalidine*, *cmr*)
- Alva, A. et al., *Chem. Pharm. Bull.*, 2004, **52**, 530-534 (*Delbonine*, *14-Acetyltakaosamine*)
- Alva, A. et al., *Helv. Chim. Acta*, 2004, **87**, 2110-2119 (*14-Deacetylpubescence*, *14-Deacetyl-18-benzoyl-18-demethylpubescence*, *1-Demethyltricornine*, *14-Benzoyltakaosamine*)
- Csupor, D. et al., *Helv. Chim. Acta*, 2004, **87**, 2125-2130 (*Lycocotinine*, *pmr*, *cmr*)
- Shrestha, P.M. et al., *J. Nat. Prod.*, 2004, **67**, 1574-1576 (*Delectinine*, *pmr*, *cmr*, *ms*)
- Shaheen, F. et al., *Phytochemistry*, 2005, **66**, 935-940 (*Lycocotinine*, *cmr*)
- Li, X. et al., *Chin. Chem. Lett.*, 2006, **17**, 1466-1468 (*Davidisines A, B*)
- Song, L. et al., *Chem. Pharm. Bull.*, 2007, **55**, 918-921 (*Delcorine*)
- Yan, L.-P. et al., *Youji Huaxue*, 2007, **27**, 976-980; *CA*, **148**, 74053r (*Campylotone*)

4-Methylnaconitane-1,6,8,9,14,16,18-heptol

M-310


C₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,6 α ,14 α ,16 β)-form

O¹⁶,O¹⁸-Di-Me, N-Et: **9-Hydroxysenbusine A**C₂₃H₃₇NO₇ 439.548Alkaloid from the roots of *Aconitum balfourii*. Amorph. solid. [α]_D +5.4 (CHCl₃).Khetwal, K.S. et al., *Nat. Prod. Res.*, 2004, **18**, 129-133 (*isol*)

4-Methylnaconitane-1,6,8,10,14,16,18-heptol

M-311



Absolute Configuration

C₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,6 α ,14 α ,16 β)-form

O⁶,O¹⁶,O¹⁸-Tri-Me, N-Et: **10-Hydroxyneoline**

[132362-42-6]


C₂₄H₃₉NO₇ 453.575Alkaloid from the roots of *Aconitum fukutomei* (Ranunculaceae). Cryst. (C₆H₆). Mp 75-79°. [α]_D²⁰ +35.8 (c, 0.41 in CHCl₃).O⁶,O¹⁶,O¹⁸-Tri-Me, N-Et, O¹⁴-Ac: **14-Acetyl-10-hydroxyneoline**

[132309-20-7]

C₂₆H₄₁NO₈ 495.612Alkaloid from the roots of *Aconitum fukutomei* (Ranunculaceae). Amorph. [α]_D²³ +37.1 (c, 0.52 in CHCl₃).Takayama, H. et al., *J. Nat. Prod.*, 1990, **53**, 936 (*isol*)

4-Methylnaconitane-1,6,8,13,14,16,18-heptol

M-312

C₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,6 α ,14 α ,16 β)-form

O¹,O¹⁶,O¹⁸-Tri-Me, N-Et: **Lasiansine**C₂₄H₃₉NO₇ 453.575Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrum*. Amorph. powder. Mp 94-96°. [α]_D²⁰ +5.5 (c, 0.5 in CHCl₃).O¹,O⁶,O¹⁶,N-Tetra-Me, 14-benzoyl, 8-Ac: **Delstaphigine**
[119347-25-0]C₃₂H₄₃NO₉ 585.693Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae).Amorph. [α]_D¹⁸ +4.2 (c, 0.4 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, 14-benzoyl, 8-Ac, N-formyl: **α -Oxodelphinine**

[466-25-1]

C₃₃H₄₃NO₁₀ 613.703Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae). Cryst. (CHCl₃/hexane). Mp 217.5-219.5°.[α]_D²⁷ -62.8 (c, 0.28 in EtOH).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, 14-O-(4-methoxybenzoyl), 8-Ac: **Habaenine B**

[947594-23-2]

C₃₃H₄₅NO₁₀ 615.719Alkaloid from *Aconitum habaense*.Amorph. solid. [α]_D¹⁵ -14.9 (c, 0.14 in CHCl₃). λ_{max} 259 (log ϵ 4.57); 336 (log ϵ 3.82); 387 (log ϵ 3.1) (CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et: **Bikhaconine**

[38146-48-4]

Semisynthetic, formed by alkaloid

hydrol. of Bikhaconitine. Gum. [α]_D²⁴

+33 (c, 0.60 in EtOH).

O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Chasmanitine**. N-Ethyl-N-desmethyldephinitine

[6846-46-4]

C₃₄H₄₇NO₉ 613.747Alkaloid from roots of *Aconitum chasmanthum*, *Aconitum ferox*, *Aconitum franchetii* and *Aconitum forrestii* (Ranunculaceae). Cryst. (Et₂O or hexane). Mp 181-182° dec. (167°).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl): **Forestine**

[91794-14-8]

C₃₃H₄₇NO₉ 601.736Alkaloid from the roots of *Aconitum forrestii* (Ranunculaceae). Amorph.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-O-acyl: **Lipofores-tine**. Lipo-14-O-anisoylbikhaconine

Alkaloid from processed tubers of

Aconitum carmichaeli. Oil. [α]_D²³ +18.7(c, 0.13 in CHCl₃). C-8 Acylating

group is a mixt. of C-16 and C-18

alkanoyl and alkenoyl groups.

O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **Crassicauline A**. **Crassicauline I**

[79592-91-9]

C₃₅H₄₉NO₁₀ 643.773Alkaloid from the roots of *Aconitum crassicaule*, *Aconitum forrestii*, *Aconitum pseudogeniculatum* and *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Needles (Me₂CO).Mp 162.5-164.5°. [α]_D²³ +31.5 (c, 0.57 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-hexadecanoyl: **Dolichotinine E**

[128341-12-8]

C₄₉H₇₇NO₁₀ 840.148Minor alkaloid from roots of *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Amorph.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 8,14-bis-O-(4-methoxybenzoyl): **Liwaconitine**


- [86408-15-3]
C₄₁H₅₃NO₁₁ 735.87
Alkaloid from the roots of *Aconitum forrestii* (Ranunculaceae). Cryst. (Et₂O). Mp 201-202.5°. [α]_D²² +133.3 (c, 0.63 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Veratroylbikhaconine**
[140188-63-2]
C₃₄H₄₉NO₁₀ 631.762
Alkaloid from *Aconitum balfourii* and *Aconitum ferox* (Ranunculaceae). Amorph. Mp 90-92°. [α]_D²⁵ +55.2 (c, 0.51 in CHCl₃) (+46.5).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl), 8-Ac: **Bikhaconitine**
[6078-26-8]
C₃₆H₅₁NO₁₁ 673.799
Alkaloid from the roots of *Aconitum spicatum*, *Aconitum ferox* and *Aconitum violaceum* (Ranunculaceae). Needles (Et₂O or EtOH aq.). Mp 105-110° (hydrate). [α]_D +16 (c, 1.6 in EtOH).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl), 8-Ac, hydrobromide:
Prisms (Et₂O). Mp 174-175°. [α]_D -8.5 (c, 1.2 in H₂O).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl), 8-Ac, perchlorate:
Needles (EtOH). Mp 240-242°. [α]_D -5.2 (c, 1.6 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl), 8-acyl: **Lipobikhaconitine**
Alkaloid from root tubers of *Aconitum ferox* (Ranunculaceae). Oil. [α]_D²⁵ +2 (c, 0.4 in CHCl₃). The esterifying group on C-8 is a mixt. of long-chain acyl (linoleoyl, palmitoyl and stearoyl).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-(E-cinnamoyl), 8-Ac: **Chasmanthinine**
[6846-47-5]
C₃₆H₄₉NO₉ 639.784
Alkaloid from the roots of *Aconitum chasmanthum* (Ranunculaceae). Needles (hexane or Et₂O). Mp 160-161° dec. [α]_D²⁵ +9.6 (c, 1.09 in EtOH).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, O⁸, N-di-Et, 14-benzoyl: **Kongboenine**. 14-O-Benzoyl-8-O-ethylbikhaconine
[129748-09-0]
C₃₄H₄₉NO₈ 599.763
Alkaloid from the *Aconitum chasmanthum* and *Aconitum kongboense* (Ranunculaceae). Prisms (MeOH). Mp 191-192°.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, O⁸, N-di-Et, 14-O-(4-methoxybenzoyl): **Acoforestine**
[110011-75-1]
C₃₅H₅₁NO₉ 629.789
Alkaloid from roots of *Aconitum forrestii* (Ranunculaceae). Mp 203-204°. [α]_D +23.4 (c, 0.56 in EtOH).
- O¹, O⁶, O¹⁶, O¹⁸, N-Penta-Me: **Delphinine**
[466-23-9]
Semisynthetic obt. by alkaline hydroly. of Delphinine or 14-O-Benzoyldelphinine. Mp 76-78°. [α]_D²⁴

- +37.5 (c, 3.2 in EtOH). [α]_D²⁷ +36 (c, 0.6 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸, N-Penta-Me, 14-benzoyl: **14-O-Benzoyldelphinine**. Deacetyldelphinine
[119347-23-8]
C₃₁H₄₃NO₈ 557.683
Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Amorph. [α]_D²⁴ +60.1 (c, 0.2 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸, N-Penta-Me, 14-benzoyl, 8-Ac: **Delphinine**
[561-07-9]
C₃₃H₄₅NO₉ 599.72
Alkaloid from the seeds of *Delphinium staphisagria* and the roots of *Atragene sibirica* (preferred genus name *Clematis*) (Ranunculaceae). Plates (EtOH). Mp 198-200°. [α]_D²⁵ +25 (EtOH).
- Poisonous.
- O¹, O⁶, O¹⁶, O¹⁸, N-Penta-Me, 14-benzoyl, 8-Ac, hydrochloride: Mp 208-210° dec.
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-benzoyl: **8-O-Deethyl-8-O-methylkongboenine**. 14-O-Benzoyl-8-O-methylbikhaconine
[212467-12-4]
C₃₃H₄₇NO₈ 585.736
Alkaloid from *Aconitum chasmanthum*.
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-O-(4-methoxybenzoyl): **Crassicausine**
[109305-85-3]
C₃₄H₄₉NO₉ 615.762
Alkaloid from roots of *Aconitum crassicaule* (Ranunculaceae). Amorph. [α]_D²³ +32.7 (c, 0.20 in CHCl₃).
- Dunstan, W.R. et al., *J.C.S.*, 1905, **87**, 1636 (isol, *Bikhaconitine*)
Jacobs, W.A. et al., *J. Biol. Chem.*, 1939, **127**, 361 (isol, *Delphinine*)
Craig, L.C. et al., *J. Biol. Chem.*, 1949, **154**, 293 (*Delphinine*)
Schneider, W. et al., *Chem. Ber.*, 1956, **89**, 768 (ir, *Delphinine*)
Jacobs, W.A. et al., *J.O.C.*, 1957, **22**, 1428 (*Delphinine*)
Wiesner, K. et al., *Tetrahedron*, 1960, **9**, 254 (struct, *Delphinine*)
Achmatowicz, O. et al., *Can. J. Chem.*, 1964, **42**, 154 (*Chasmanthinine*, *Chasmaconitine*)
Birnbaum, K.B. et al., *Acta Cryst. B*, 1972, **28**, 1551 (config)
Klásek, A. et al., *J. Nat. Prod.*, 1972, **35**, 55 (*Bikhaconitine*, *Chasmaconitine*)
Pelletier, S.W. et al., *J.A.C.S.*, 1976, **98**, 2626 (cmr)
Tiwari, K.P. et al., *J. Indian Chem. Soc.*, 1977, **54**, 924 (isol, *Bikhaconitine*)
Krasnov, E.A. et al., *Khim. Prir. Soedin.*, 1981, **17**, 806; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 598 (isol, *Delphinine*)
Wang, F. et al., *Planta Med.*, 1981, **42**, 375 (*Crassicauline A*)
Pelletier, S.W. et al., *J.O.C.*, 1982, **47**, 5290 (cryst struct, cmr, *Delphinine*)
Wang, C. et al., *Planta Med.*, 1983, **48**, 55 (*Livaconitine*)
Pelletier, S.W. et al., *J. Nat. Prod.*, 1984, **47**, 474 (*Forestine*)
Pelletier, S.W. et al., *Heterocycles*, 1987, **25**, 365 (*Acoforestine*)
Ross, S.A. et al., *Heterocycles*, 1987, **26**, 2895 (α -*Oxodelphinine*)
Wang, F.-P. et al., *J. Nat. Prod.*, 1987, **50**, 55 (*Crassicausine*)
Pelletier, S.W. et al., *Heterocycles*, 1988, **27**, 2467 (*14-O-Benzoyldelphinine*, *Delstaphigine*)

- Huiling, L. et al., *Heterocycles*, 1989, **29**, 2317 (*Crassicauline A*, *Dolichotine E*)
Yue, J.-M. et al., *Phytochemistry*, 1990, **29**, 2379-2380 (*Kongboenine*)
Khetwal, K.S. et al., *Heterocycles*, 1992, **34**, 441 (*Veratroylbikhaconine*)
Hanuman, J.B. et al., *J. Nat. Prod.*, 1993, **56**, 801; 1994, **57**, 105 (*Veratroylbikhaconine*, *Lipobikhaconitine*)
Nair, M.M. et al., *Spectrosc. Lett.*, 1997, **30**, 213-222 (pmr, cmr, *Delphinine*)
Parvez, M. et al., *Acta Cryst. C*, 1998, **54**, 125-126; 236-238; 790-792 (*Chasmaconitine*, *Chasmanthinine*, 8-Deethyl-8-methylkongboenine, *cryst struct*)
Shim, S.H. et al., *Chem. Pharm. Bull.*, 2003, **51**, 999-1002 (*Lipoforestine*)
Ji, H. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 619-624 (*Lasiansine*)
Yang, S. et al., *Helv. Chim. Acta*, 2007, **90**, 1160-1164 (*Habaenine B*)

4-Methylnaconitane-1,6,8,14,15,16,18-heptol

M-313

(1 α ,5 β ,6 α ,14 α ,15 α ,16 β)-formC₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,6 α ,14 α ,15 α ,16 β)-form

- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et: **Senbusine C**. 15-Hydroxyneoline. 15-Epinagarine. *Fuziline*
[80665-72-1]
C₂₄H₃₉NO₇ 453.575
Alkaloid from the roots of *Aconitum carmichaeli*, *Aconitum japonicum*, *Aconitum napellus* and *Aconitum ibukiense* (Ranunculaceae). Cryst. (MeOH, Me₂CO or EtOAc). Mp 214-216° (193-194°). [α]_D +6.7 (c, 0.21 in MeOH). [α]_D^{22.5} +19.3 (CHCl₃). [α]_D²⁶ +11.6 (c, 1.0 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 8-Ac: **Taurenine**. 8-Acetyl-15-hydroxyneoline
[172616-86-3]
C₂₆H₄₁NO₈ 495.612
Alkaloid from *Aconitum tauricum*. Also isol. from aphids feeding on *Aconitum napellus*. Cryst. (petrol/CHCl₃). Mp 100-102°. [α]_D²² +4.1 (c, 0.07 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 8-Ac: **Isodelphinine**
[1357-96-6]
C₃₃H₄₅NO₉ 599.72
Alkaloid from the roots of *Aconitum miyabei* and *Aconitum carmichaeli* (Ranunculaceae). Needles (Et₂O/petrol). Mp 167-168°. [α]_D²⁰ +20.1 (c, 2.05 in EtOH).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Penduline**†
[81362-34-7]
C₃₄H₄₇NO₉ 613.747
Alkaloid from roots of *Aconitum*

pendulum (Ranunculaceae). Mp 166-167°.

(1 α ,5 β ,6 α ,14 α ,15 β ,16 β)-form

O⁶,O¹⁶,O¹⁸-Tri-Me, N-Et: *Nagarine*†. 15-Episenbusine C. 15-Epifuziline. Crassicaulisine. 3-Deoxycrassicaulidine. *Bullatine F* [80665-73-2]

C₂₄H₃₉NO₇ 453.575

Alkaloid from roots of *Aconitum nazarum* var. *heterotrichum* and *Aconitum crassaule* (Ranunculaceae). Cryst. (Me₂CO). Mp 190-191°. [α]_D²¹ +55.4 (c, 1.0 in EtOH). [α]_D²¹ +20.4 (c, 0.88 in CHCl₃).

O⁶,O¹⁶,O¹⁸-Tri-Me, N-Et, 8,15-O-isopropylidene: *Bullatine E* [1354-88-7]

C₂₇H₄₃NO₇ 493.639

Alkaloid from *Aconitum bullatifolium* var. *homotrichum* (Ranunculaceae). Mp 182-183°. [α]_D²⁴ +79.6 (Me₂CO).

Suginome, H. et al., *Bull. Chem. Soc. Jpn.*, 1959, **32**, 604 (*Isodelphinine*, isol)

Katsui, N. et al., *Bull. Chem. Soc. Jpn.*, 1959, **32**, 774

Chu, J.-H. et al., *Huaxue Xuebao*, 1965, **31**, 222-228; *CA*, **63**, 16400a (*Bullatine E*)

Pelletier, S.W. et al., *Tet. Lett.*, 1977, 4027 (*Isodelphinine*, ir, pmr, cmr, struct)

Takayama, H. et al., *Chem. Pharm. Bull.*, 1981, **29**, 3078; 1982, **30**, 386 (pmr, cmr, ms, struct, synth)

Zhu, Y. et al., *Heterocycles*, 1981, **16**, 1723 (*Penduline*)

Mody, N.V. et al., *Heterocycles*, 1982, **17**, 91 (*Nagarine*)

Pelletier, S.W. et al., *Heterocycles*, 1982, **18**, 47 (ir, pmr, cmr, cryst struct)

Konno, C. et al., *J. Nat. Prod.*, 1982, **45**, 128-133 (*Senbusine C*)

Zhang, D. et al., *Zhongcaoyao*, 1982, **13**, 481; *CA*, **98**, 149496x (occur)

Hikino, H. et al., *J. Nat. Prod.*, 1983, **46**, 178; 1984, **47**, 190 (isol, ir, pmr)

Wang, F. et al., *Planta Med.*, 1983, **47**, 39 (*Nagarine*)

Liu, L. et al., *Yaoxue Xuebao*, 1983, **18**, 39; *CA*, **99**, 67495c (*Penduline*)

Sakai, S. et al., *Yakugaku Zasshi*, 1984, **104**, 222; *CA*, **101**, 69329k (isol)


Yang, P. et al., *Huaxue Xuebao*, 1988, **46**, 827; *CA*, **109**, 226747e (*Bullatine E*, struct)

Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1992, **28**, 108-112; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **28**, 91-94 (*Taurenine*)

Liu, H. et al., *J. Nat. Prod.*, 1996, **59**, 135-138 (*Taurenine*)

4-Methylnaconitane-1,6,8,14,16,18,19-heptol

M-314



C₁₉H₂₉NO₇ 383.441

(1 α ,6 α ,14 α ,16 β ,19S)-form

O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, N-Et: *Aconitilearine*


C₂₅H₄₁NO₇ 467.601

Alkaloid from the aerial parts of *Aconitum cochleare*. [α]_D +55 (c, 0.11 in CHCl₃).

Meriçli, A.H. et al., *Helv. Chim. Acta*, 2006, **89**, 210-217 (*Aconitilearine*)

4-Methylnaconitane-1,7,8,9,14,16,18-heptol

M-315



C₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,14 α ,16 β)-form

7,8-Methylene ether, O¹,O¹⁴,O¹⁸-tri-Me, N-Et: *Molline*

[163634-09-1]

C₂₅H₃₉NO₇ 465.586

Alkaloid from aerial parts of *Delphinium mollipilum* (Ranunculaceae).

7,8-Methylene ether, O¹,O¹⁴,O¹⁶O¹⁸-tetra-Me, N-Et: *Talistine B*

[110081-93-1]

C₂₆H₄₁NO₇ 479.612

Alkaloid from the roots of *Delphinium taliense* (Ranunculaceae). Also isol. from whole herb of *Delphinium caeruleum*. Cryst. (MeOH). Mp 98-100°.


Chen, S. et al., *Yunnan Zhiwu Yanjiu*, 1986, **8**, 81; *CA*, **107**, 112626t (*Talistine B*, struct)

Zhang, S.M. et al., *Chin. Chem. Lett.*, 1995, **6**, 101; *CA*, **123**, 5584m (*Molline*)

Wang, Y. et al., *Phytochemistry*, 1996, **42**, 569 (*Talistine B*)

4-Methylnaconitane-1,8,10,13,14,15,18-heptol

M-316



C₁₉H₂₉NO₇ 383.441

(1 α ,5 β ,14 α ,15 α)-form

O¹,O¹⁸-Di-Me, N-Et, 14-O-(4-methoxybenzoyl): *Guayewuanine A* [83145-44-2]

C₃₁H₄₃NO₉ 573.682


Alkaloid from the roots of *Aconitum hemisleyanum* (Ranunculaceae). Mp 120°.

Zhang, H. et al., *Zhiwu Xuebao (Acta Bot.*

Sin.), 1982, **24**, 259; *CA*, **97**, 159517y (isol, ir, pmr, ms, struct)

4-Methylnaconitane-1,3,8,14,16,18-hexol

M-317



C₁₉H₂₉NO₆ 367.441

(1 α ,3 α ,14 α ,16 β)-form

O¹,O¹⁶,O¹⁸-Tri-Me, N-Et: *3 α -Hydroxy-talatinzamine*


C₂₄H₃₉NO₆ 437.575

Alkaloid from above-ground parts of *Aconitum nasutum*. Amorph. [α]_D -14.3 (c, 0.3 in CHCl₃).

Meriçli, A.H. et al., *Phytochemistry*, 1996, **42**, 909-911 (isol, pmr, cmr)

4-Methylnaconitane-1,5,7,8,14,16-hexol

M-318



C₁₉H₂₉NO₆ 367.441

(1 α ,14 α ,16 α)-form

7,8-Methylene ether, O¹,O¹⁴,O¹⁶-tri-Me, N-Et: *Nordhagenine A*


C₂₅H₃₉NO₆ 449.586

Alkaloid from *Delphinium nordhagenii*. Cryst. (hexane/Me₂CO). Mp 178-180°. [α]_D³⁰ -3.2 (c, 0.4 in CHCl₃).

Shaheen, F. et al., *J. Nat. Prod.*, 2006, **69**, 823-825 (isol, pmr, cmr, cryst struct)

4-Methylnaconitane-1,5,8,14,16,18-hexol

M-319




C₁₉H₂₉NO₆ 367.441

(1 α ,5 β ,14 α ,16 β)-form

O¹,O¹⁸-Di-Me, N-Et: *Circinasine E* [939972-79-9]

C₂₃H₃₇NO₆ 423.548Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. [α]_D²⁰ +36.5 (c, 1 in CHCl₃).**O¹,O¹⁸-Di-Me, 14-O-(4-methoxybenzoyl), N-Et: Circinasine D**
[939972-78-8]C₃₁H₄₃NO₈ 557.683Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. [α]_D²⁰ +15.8 (c, 1 in CHCl₃).**O¹,O¹⁶,O¹⁸, N-Et: Hemsleyanine C**C₂₄H₃₉NO₆ 437.575Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. Mp 74-75°. [α]_D²⁰ -48.8 (c, 1 in CHCl₃).**O¹,O¹⁶,O¹⁸-Tri-Me, 14-O-(4-methoxybenzoyl), N-Et: Hemsleyanine D**C₃₂H₄₅NO₈ 571.709Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. Mp 89-90°. [α]_D²⁰ +12.8 (c, 1 in CHCl₃).Gao, F. et al., *Arch. Pharmacol. Res.*, 2007, **30**, 1497-1500 (*Hemsleyanines C,D*)Gao, F. et al., *J. Nat. Prod.*, 2007, **70**, 876-879 (*Circinasines D-E*)**4-Methylnaconitane-1,6,7,8,12,14-hexol**


M-320

C₁₉H₂₉NO₆ 367.441**(1α,6α,12β,14α)-form**N-Et, 14-benzoyl, 6,12-di-Ac: **Aconitinine**

[192200-74-1]

C₃₂H₄₁NO₉ 583.677Alkaloid from *Consolida aconiti*.Martin, G.D.L.P. et al., *Bol. Soc. Quim. Peru*, 1997, **63**, 44-49; *CA*, **127**, 106630m**4-Methylnaconitane-1,6,7,8,14,16-hexol**

M-321

**(1α,5β,6α,14α,16β)-form**C₁₉H₂₉NO₆ 367.441**(1α,5β,6α,14α,16β)-form****O¹⁶-Me, N-Et: Consolarine**

[205646-92-0]

C₂₂H₃₅NO₆ 409.522Alkaloid from *Consolida armeniaca* (Ranunculaceae). Amorph. [α]_D +0.6 (c, 0.8 in CHCl₃).**O⁸,O¹⁶-Di-Me, N-Et: 8-O-Methylconsolarine**C₂₃H₃₇NO₆ 423.548Alkaloid from *Consolida orientalis*. Light yellow oil. [α]_D²⁵ +15 (c, 0.3 in CHCl₃).**O⁸,O¹⁶-Di-Me, N-Et, 14-Ac: 18-De-methoxypubescenine**C₂₅H₃₉NO₇ 465.586Alkaloid from the aerial parts of *Consolida orientalis*. Amorph. solid. [α]_D²⁵ +1.1 (c, 0.4 in CHCl₃).**O¹,O⁷,O⁸,O¹⁶-Tetra-Me, N-Et, 14-benzoyl, 6-Ac: Delphiperegrine**

[142609-22-1]

C₃₄H₄₇NO₈ 597.747Alkaloid from aerial parts of *Delphinium peregrinum* (Ranunculaceae). [α]_D²² +5.6 (c, 0.2 in MeOH).**1-Ketone, O⁷,O⁸,O¹⁶-tri-Me, 6,14-di-Ac: Pergilone**

[142609-20-9]

C₂₆H₃₇NO₈ 491.58Alkaloid from aerial parts of *Delphinium peregrinum* (Ranunculaceae). [α]_D²⁵ +4.2 (c, 0.1 in MeOH).**(1α,5β,6β,14α,16β)-form****O⁶,O¹⁶-Di-Me, N-Et: Dihydrogadesine**

[70420-63-2]

C₂₃H₃₇NO₆ 423.548Alkaloid from *Delphinium pentagynum* and from the aerial parts of *Delphinium peregrinum* var. *elongatum* collected during the flowering period (Ranunculaceae). Cryst. (petrol/EtOAc). Mp 136-138°. [α]_D +54 (c, 0.1 in EtOH).**O⁶,O¹⁶-Di-Me, N-Et, 14-Ac: 14-Acetyldihydrogadesine**

[84306-87-6]

C₂₅H₃₉NO₇ 465.586Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Resin.**O⁶,O¹⁶-Di-Me, N-Et, 14-benzoyl: 14-Benzoyldihydrogadesine**

[103976-34-7]

C₃₀H₄₁NO₇ 527.656Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Cryst. (EtOAc). Mp 199-202°.**O¹,O⁶,O¹⁶-Tri-Me, N-Et: Nudicaulidine**

[99815-81-3]

C₂₄H₃₉NO₆ 437.575Alkaloid from *Delphinium nudicaule* (Ranunculaceae). Amorph. [α]_D +42 (c, 0.15 in CHCl₃).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-Ac: 14-Acetylnudicaulidine**

[99815-82-4]

C₂₆H₄₁NO₇ 479.612Alkaloid from *Delphinium andersonii* and *Delphinium confusum* (Ranunculaceae). Mp 206-208°. [α]_D²⁶ +18.9 (c, 0.54 in CHCl₃).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-O-(2-methylpropanoyl): 14-Isobutyrylnudi-****caulidine**

[195153-55-0]

C₂₈H₄₅NO₇ 507.666Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin. [α]_D +31 (c, 0.1 in EtOH).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-O-(2-methylbutanoyl): 14-(2-Methylbutanoyl)nudicaulidine**

[195246-59-4]

C₂₉H₄₇NO₇ 521.693Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin. [α]_D +44.5 (c, 0.05 in EtOH).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-benzoyl: 14-Benzoylnudicaulidine**

[195153-54-9]

C₃₁H₄₃NO₇ 541.683Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Cryst. (EtOAc/hexane). Mp 199-202°. [α]_D +48.6 (c, 0.22 in EtOH).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-E-cinnamoyl: 14-trans-Cinnamoylnudicaulidine**

[195153-57-2]

C₃₃H₄₅NO₇ 567.721Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin. [α]_D +72.1 (c, 0.26 in EtOH).**O¹,O⁶,O¹⁶-Tri-Me, N-Et, 14-Z-cinnamoyl: 14-cis-Cinnamoylnudicaulidine**

[195153-56-1]

C₃₃H₄₅NO₇ 567.721Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin. [α]_D +4 (c, 0.125 in EtOH).**O¹,O¹⁴,O¹⁶-Tri-Me, N-Et: Demethylene-delpheline**

[122279-79-2]

C₂₄H₃₉NO₆ 437.575Alkaloid from aerial parts of *Delphinium corumbosum* (Ranunculaceae). Solid (petrol). Mp 78-80°.**O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, N-Et: 18-Deoxylycoctonine**

[41547-37-9]

C₂₅H₄₁NO₆ 451.602Alkaloid from *Delphinium confusum* (Ranunculaceae). Mp 73-75°.**7,8-Methylene ether, O¹⁶-Me, N-Et: Delmenzine. Blacknine. 1,14-Didemethyldelpheline. O¹⁴-Demethyldelelatine. O¹⁴-Demethylpacidine**

[139768-08-4]

C₂₃H₃₅NO₆ 421.533Alkaloid from *Delphinium menziesii* and whole plants of *Delphinium elatum* var. *black night* (Ranunculaceae). Amorph. [α]_D -5.13 (c, 0.887 in CHCl₃).**7,8-Methylene ether, O¹,O¹⁴-di-Me, N-Et: Eladine**

[123064-69-7]

C₂₄H₃₇NO₆ 435.559Alkaloid from the seeds of *Delphinium elatum* (Ranunculaceae). Amorph. [α]_D²³ -57.5 (c, 0.89 in CHCl₃).**7,8-Methylene ether, O¹,O¹⁶-di-Me, N-Et: Delelatine**

[119371-58-3]

C₂₄H₃₇NO₆ 435.559

Alkaloid from seeds of *Delphinium elatum* and from *Delphinium tatsienense* (Ranunculaceae). Cryst. (Et₂O). Mp 84.5-86.5°.

7,8-Methylene ether, O¹⁴, O¹⁶-di-Me, N-Et: **Pacidine**

[142748-53-6]
C₂₄H₃₇NO₆ 435.559

Alkaloid from seeds of *Delphinium elatum* hybrid pacific giant (Ranunculaceae). Amorph. solid. [α]_D²⁰ -14.3 (c, 0.49 in CHCl₃).

7,8-Methylene ether, O¹⁴, O¹⁶-di-Me, N-Et, 6-Ac: **Tiantaishannine**

[959843-31-3]
C₂₆H₃₉NO₇ 477.597

Alkaloid from the roots of *Delphinium tiantaishanense*. Amorph. powder. Mp 206-208°. [α]_D²⁰ -23.9 (c, 0.73 in CHCl₃).

7,8-Methylene ether, O¹, O⁶, O¹⁶-tri-Me, N-Et: **Isodelpheline**. Tongoline

[123064-67-5]
C₂₅H₃₉NO₆ 449.586

Alkaloid from the seeds of *Delphinium elatum* and *Delphinium tongolense*. Cryst. Mp 217-218°. [α]_D²⁰ -1.2 (c, 1 in CHCl₃).

7,8-Methylene ether, O¹, O¹⁴, O¹⁶-tri-Me, N-Et: **Delpheline**

[509-28-4]
C₂₅H₃₉NO₆ 449.586

Alkaloid from the seeds of *Delphinium elatum* and from the roots and aerial parts of *Delphinium ternatum* (Ranunculaceae) also from *Delphinium occidentale*. Prisms (EtOH aq.). Mp 227° (210-212°, 216-219°). [α]_D¹⁵ -25.8 (c, 2 in CHCl₃).

7,8-Methylene ether, O¹, O¹⁴, O¹⁶-tri-Me, N-Et, hydrochloride:
Prisms + 1H₂O (Me₂CO aq.). Mp 219° (frothing). [α]_D²⁰ -42.8 (c, 2 in H₂O).

7,8-Methylene ether, O¹, O⁶, O¹⁴, O¹⁶-tetra-Me, N-Et: **Paciline**. 6-O-Methyldepheline

[122384-95-6]
C₂₆H₄₁NO₆ 463.613

Alkaloid from seeds of the *Delphinium* hybrid pacific giant (Ranunculaceae). Amorph. [α]_D -7.2 (c, 1.0 in CHCl₃).

6-Ketone, 7,8-methylene ether, O¹, O¹⁶-di-Me, N-Et: **Yunnadelphinine**

[90718-33-5]
C₂₄H₃₅NO₆ 433.544

Alkaloid from *Delphinium yunnanense* (Ranunculaceae).

6-Ketone, 7,8-methylene ether, O¹, O¹⁶-di-Me, N-Et, 14-Ac: **Barbinidine**

[123519-72-2]
C₂₆H₃₇NO₇ 475.581

Alkaloid from aerial parts of *Delphinium barbeyi* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 215-216°.

6-Ketone, 7,8-methylene ether, O¹, O¹⁴, O¹⁶-tri-Me, N-Et: **Pacinine**. 6-Dehydrodelpheline

[60223-97-4]
C₂₅H₃₇NO₆ 447.57

Alkaloid from seeds of the *Delphinium* hybrid pacific giant (Ranunculaceae).

Mp 133-135.5°. [α]_D -58 (c, 1.0 in MeOH).

(1β,5β,6β,14α,16β)-form

7,8-Methylene ether, O¹, O¹⁴, O¹⁶-tri-Me, N-Et, 6-Ac: **Souline** C. 6-Acetyldepheline (incorr.)

[42907-15-3]
C₂₇H₄₁NO₇ 491.623

Alkaloid from *Delphinium souliei*, *Delphinium barbeyi* and aerial parts of *Delphinium occidentale* (Ranunculaceae). Prisms (EtOH aq.). Mp 125° (120-122°) Mp 170-171°. [α]_D¹⁸ -44.5 (c, 0.67 in CHCl₃). [α]_D²⁰ -34.5 (c, 2 in EtOH).

Goodson, J.A. et al., *J.C.S.*, 1943, 139; 1944, 665 (*Delpheline, isol, struct*)

Cookson, R.C. et al., *J.C.S.*, 1956, 2689; 3121 (*Delpheline, isol, uv, struct*)

Edwards, O.E. et al., *Can. J. Chem.*, 1958, 36, 1097 (*Delpheline, struct*)

Carmack, M. et al., *J.A.C.S.*, 1958, 80, 497 (*struct*)

Pelletier, S.W. et al., *J.A.C.S.*, 1981, 103, 6536 (*Delpheline, config*)

González, A.G. et al., *Phytochemistry*, 1982, 21, 1781-1782 (*Dihydrogadesine, Acetyldihydrogadesine*)

Luo, S. et al., *CA*, 1984, 101, 35860m (*Yunnadelphinine*)

Jiang, Q.P. et al., *Heterocycles*, 1984, 22, 2429 (*Delpheline, cmr*)

Kulanthaivel, P. et al., *Heterocycles*, 1985, 23, 2515-2520 (*Nudicaulidine*)

González, A.G. et al., *Heterocycles*, 1986, 24, 1513-1516 (*Dihydrogadesine, Benzoyldihydrogadesine*)

Ross, S.A. et al., *Phytochemistry*, 1988, 27, 3719 (*Delpheline, isol, ir, pmr, cmr, ms, struct*)

Pelletier, S.W. et al., *Heterocycles*, 1989, 28, 107-110 (*14-Acetylnudicaulidine*)

Bando, H. et al., *Heterocycles*, 1989, 29, 1293 (*Paciline, Pacinine*)

Narzullaev, A.S. et al., *Khim. Prir. Soedin.*, 1989, 25, 48-50; 50-51; 1991, 27, 547-549; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, 25, 41-42; 43; 1991, 27, 478-479

(*Demethylnedelpheline, 14-Acetylnudicaulidine, 18-Deoxylycoctonine*)

Pelletier, S.W. et al., *Phytochemistry*, 1989, 28, 1521 (*Barbinidine*)

Pelletier, S.W. et al., *Tetrahedron*, 1989, 45, 1887 (*Isodelpheline, Eladine*)

Sun, F. et al., *Heterocycles*, 1991, 32, 1963 (*Delmenzine*)

Joshi, B.S. et al., *Tetrahedron*, 1991, 47, 4299 (*Delpheline, pmr, cmr*)

Ulubelen, A. et al., *Phytochemistry*, 1992, 31, 1019-1022 (*Delphiperegrine, Pergilone*)

Wada, K. et al., *Phytochemistry*, 1992, 31, 2135 (*Pacidine*)

Park, J.C. et al., *J. Nat. Prod.*, 1995, 58, 291 (*Blacknine*)

He, L. et al., *Chin. Chem. Lett.*, 1996, 7, 557-560 (*Tongoline*)

Reina, M. et al., *Phytochemistry*, 1997, 45, 1707-1711 (*14-acylnudicaulidines*)

Merclii, A.H. et al., *Heterocycles*, 1998, 47, 329-335 (*Consolarine*)

Zhang, K. et al., *Planta Med.*, 1998, 64, 580-581 (*Souline* C)


Alva, A. et al., *Chem. Pharm. Bull.*, 2004, 52, 530-534 (*18-Demethoxypubescenine*)

Alva, A. et al., *Helv. Chim. Acta*, 2004, 87, 2110-2119 (*8-O-Methylconsolarine*)

Li, J. et al., *Molecules*, 2007, 12, 353-360 (*Tiantaishannine*)

4-Methylnaconitane-1,6,8,10,14,16-hexol

M-322



C₁₉H₂₉NO₆ 367.441

(1α,5β,6α,14α,16β)-form

O⁶, O¹⁶-Di-Me, N-Et, 14-benzoyl: **Pentagyline**

[89384-05-4]
C₃₀H₄₁NO₇ 527.656

Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 199-200°.

(1α,5β,6β,14α,16β)-form

O¹, O⁸, O¹⁶-Tri-Me, N-Et: **6-Deacetyl-10-hydroxyperegrine**

[160884-37-7]
C₂₄H₃₉NO₆ 437.575

Alkaloid from *Delphinium peregrinum* (Ranunculaceae).

O¹, O⁸, O¹⁶-Tri-Me, N-Et, 6-Ac: **10-Hydroxyperegrine**

[167559-25-3]
C₂₆H₄₁NO₇ 479.612

Alkaloid from aerial parts of *Delphinium munzianum* (Ranunculaceae).

Amorph. [α]_D +15 (c, 0.18 in CHCl₃).


González, A.G. et al., *Heterocycles*, 1984, 22, 17 (*Pentagyline*)

Ulubelen, A. et al., *Nat. Prod. Lett.*, 1994, 5, 135 (*6-Deacetyl-10-hydroxyperegrine*)

de La Fuente, G. et al., *Phytochemistry*, 1995, 39, 1467 (*10-Hydroxyperegrine*)

4-Methylnaconitane-1,6,8,14,16,18-hexol

M-323



C₁₉H₂₉NO₆ 367.441

(1α,5β,6α,14α,16β)-form

O⁶, O¹⁶-Di-Me, N-Et: **Neolinine**

[112515-37-4]
C₂₃H₃₇NO₆ 423.548

Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Mp 226-228.5°. [α]_D²⁶ +29.3 (c, 0.19 in CHCl₃).

O⁶, O¹⁶-Di-Me, N-Et, 8,14-di-Ac: **Delstaphisagrine**

[102686-12-4]
C₂₇H₄₁NO₈ 507.623

Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae).

Amorph. [α]_D²³ +3.8 (c, 0.6 in EtOH).


O⁶, O¹⁸-Di-Me, N-Et, 8,14-di-Ac: **Delsta-**

- phisine**
[102686-13-5]
C₂₇H₄₁NO₈ 507.623
Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae). Plates (Me₂CO/hexane). Mp 182-184°. [α]_D²⁵ -11 (c, 1.35 in EtOH).
- O¹⁶, O¹⁸-Di-Me, N-Et: **Senbusine A. Bataconine**
[82202-95-7]
C₂₃H₃₇NO₆ 423.548
Alkaloid from the roots of *Aconitum carmichaeli*, *Aconitum napellus* and *Aconitum ibukiense* (Ranunculaceae). Amorph. powder or cryst. (Me₂CO). Mp 96-99°. Bataconine not proved identical with Senbusine A. Mp refers to Bataconine.
- O¹⁶, O¹⁸-Di-Me, N-Et, 14-Ac: **14-O-Acetylsenbusine A**
[158784-70-4]
C₂₅H₃₉NO₇ 465.586
Alkaloid from root tubers of *Aconitum ferox* (Ranunculaceae). Mp 88-90°. [α]_D²⁵ +6 (c, 0.5 in EtOH).
- O¹⁶, O¹⁸-Di-Me, N-Et, 8,14-di-Ac: **Delstaphisinine**
[116139-60-7]
C₂₇H₄₁NO₈ 507.623
Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae). Cryst. (Et₂O/hexane). Mp 158-160°. [α]_D²⁵ -14.1 (c, 0.15 in CHCl₃).
- O¹⁶, O¹⁸-Di-Me, N-Et, 1,6,14-tribenzoyl: **1,6,14-Tribenzoylsenbusine A**
C₄₄H₄₉NO₉ 735.872
Alkaloid from *Delphinium montanum*. CAS no. not found 8-14Cl.
- O¹, O⁶, O¹⁶-Tri-Me, N-Et: **Leucanthum-sine C**
[947591-61-9]
C₂₄H₃₉NO₆ 437.575
Alkaloid from the roots of *Aconitum sungpanense* var. *leucanthum*. Amorph. powder. Mp 152-154°. [α]_D²⁰ +53 (c, 1 in CHCl₃).
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et: **6-Epiforesticine**
[321338-28-7]
C₂₄H₃₉NO₆ 437.575
Alkaloid from *Aconitum hemsleyanum* var. *pengzhouense*. Needles (Et₂O/hexane). Mp 114-116°. [α]_D²⁴ +43 (c, 0.73 in CHCl₃).
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-benzoyl: **Kongboentine B**
C₃₁H₄₃NO₇ 541.683
Alkaloid from the roots of *Aconitum kongboense*. Amorph. powder. Mp 96-98°. CAS no. not found 8-14Cl.
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-benzoyl, 8-Ac: **Patentine**
[132185-72-9]
C₃₃H₄₅NO₈ 583.72
Alkaloid from the roots of *Aconitum vilnorrianum* var. *patentipilum* (Ranunculaceae). Amorph. solid.
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-O-(4-methoxybenzoyl): **Geniconitine**
[99713-74-3]
C₃₂H₄₅NO₈ 571.709
Alkaloid from roots of *Aconitum geniculatum* (Ranunculaceae).
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-E-cinnamoyl: **Leucantine D**
[499969-78-7]
C₃₃H₄₅NO₇ 567.721
Alkaloid from the roots of *Aconitum hemsleyanum* var. *leucanthus*. Amorph. powder. [α]_D²⁰ +34 (c, 0.5 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Me, 8,14-di-Ac: **Pictumine†**
[123213-56-9]
C₂₇H₄₁NO₈ 507.623
Alkaloid from the seeds of *Delphinium pictum* ssp. *pictum* (Ranunculaceae). Amorph.
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et: **Neoline. Bullatine B**
[466-26-2]
C₂₄H₃₉NO₆ 437.575
Alkaloid from *Aconitum bullatifolium* var. *homotrichum*, *Aconitum napellus*, *Aconitum nagarum*, *Aconitum sczukini*, *Aconitum soongoricum*, *Aconitum sachalinense* var. *compactum*, *Aconitum carmichaeli*, *Aconitum karakolicum*, *Aconitum teipeicum* and *Aconitum ibukiense* (Ranunculaceae). Mp 159-161°. [α]_D²⁶ +22 (c, 4.3 in EtOH).
- ▶ AR5569450
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 8-Ac: **Delphidine. 8-Acetylneoline**
[60158-58-9]
C₂₆H₄₁NO₇ 479.612
Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Mp 98-100° (slight softening at 90°). [α]_D²⁷ +16.6 (c, 1.3 in EtOH).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-Ac: **Bullatine C. 14-Acetylneoline. Delstaphisagnine**
[1354-86-5]
C₂₆H₄₁NO₇ 479.612
Alkaloid from *Aconitum bullatifolium* var. *homotrichum*, *Aconitum nagarum*, *Aconitum napellus*, *Aconitum jinyangense* and *Aconitum yesoense* (Ranunculaceae). Also descr. as Delstaphisagnine from *Delphinium staphisagria*. Needles (Me₂CO/Et₂O) or amorph. solid. Mp 200°. [α]_D²⁵ +20 (c, 0.85 in EtOH). Opt. rotn. refers to amorph. Delstaphisagnine.
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 1,14-di-Ac: **1,14-Diacetylneoline**
[60158-60-3]
C₂₈H₄₃NO₈ 521.65
Alkaloid from above-ground parts of *Aconitum napellus* ssp. *castellanum* (Ranunculaceae). Resin. [α]_D +15 (c, 0.68 in EtOH).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 8,14-di-Ac: **Delphisine. 8,14-Diacetylneoline**
[1399-07-1]
C₂₈H₄₃NO₈ 521.65
Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 122-123°. [α]_D²⁶ +7.1 (c, 4.0 in EtOH).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, tri-Ac: **1-Acetyldelphisine**
[55085-41-1]
C₃₀H₄₅NO₉ 563.687
- Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 151-153°. [α]_D²⁵ -8.8 (c, 0.63 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-benzoyl: **14-Benzoylneoline**
[99633-05-3]
C₃₁H₄₃NO₇ 541.683
Alkaloid from rhizomes of *Aconitum subneatum* (Ranunculaceae). Amorph. powder. [α]_D¹⁸ +9.1 (c, 0.11 in MeOH).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-(4-methoxybenzoyl): **14-Anisoylneoline. 14-(4-Methoxybenzoyl)neoline**
[640282-28-6]
C₃₂H₄₅NO₈ 571.709
Alkaloid from processed tubers of *Aconitum carmichaeli*. Amorph. powder (MeOH). [α]_D²² +22.3 (c, 0.7 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 8-O-(3,4-dimethoxybenzoyl): **Acotoxinine. 8-Veratroylneoline. 8-(3,4-Dimethoxybenzoyl)neoline**
C₃₃H₄₇NO₉ 601.736
Alkaloid from the roots of *Aconitum toxicum*. Amorph. solid. [α]_D²⁵ +32 (c, 0.05 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Geniculatine B. 14-Veratroylneoline. 14-(3,4-Dimethoxybenzoyl)neoline**
[359798-73-5]
C₃₃H₄₇NO₉ 601.736
Alkaloid from processed tubers of *Aconitum carmichaeli* and from *Aconitum geniculatum*. Amorph. powder (MeOH). [α]_D²³ +22.1 (c, 0.75 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **Geniculatine A**
[359798-72-4]
C₃₄H₄₇NO₉ 613.747
Alkaloid from the roots of *Aconitum geniculatum*. Amorph. solid. [α]_D +19.3 (c, 0.5 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 8-cinnamoyl: **8-O-trans-Cinnamoylneoline**
[674365-50-5]
C₃₃H₄₅NO₇ 567.721
Alkaloid from the flower buds of *Aconitum carmichaeli*. Amorph. powder. [α]_D²⁰ +20.4 (c, 1 in EtOH). λ_{max} 203 (log ε 4.48); 217 (log ε 4.46); 223 (log ε 4.44); 278 (log ε 4.6) (EtOH).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-E-cinnamoyl: **14-Cinnamoylneoline**
[640282-27-5]
C₃₃H₄₅NO₇ 567.721
Alkaloid from processed tubers of *Aconitum carmichaeli*. Amorph. powder (MeOH). [α]_D²³ +9.7 (c, 0.35 in CHCl₃).
- O¹, O⁶, O¹⁴, O¹⁶-Tetra-Me, N-Et: **Acoseptriginine**
[144049-70-7]
C₂₅H₄₁NO₆ 451.602
Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Amorph.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et: **Chasma-**

- nine. Tokoro-base II**
[5066-78-4]
C₂₅H₄₁NO₆ 451.602
Alkaloid from *Aconitum chasmanthum*, *Aconitum subcuneatum*, *Aconitum yesoense*, *Aconitum crassicaule*, *Aconitum franchetii*, *Aconitum teipecum*, *Aconitum sachalinense* var. *compactum*, *Aconitum forrestii*, *Aconitum pseudogeniculatum* and others (Ranunculaceae). Fine needles (hexane). Mp 90-91°. [α]_D²⁵ +23.6 (c, 2.5 in EtOH).
- ▶ AR5543000
O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-Ac: **14-Acetylchasanine**
[77757-06-3]
C₂₇H₄₃NO₇ 493.639
Alkaloid from *Delphinium uncinatum*.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 8,14-di-Ac: **1-O-Methyldeiphisine. 8,14-Diacetylchasanine**
[5066-80-8]
C₂₉H₄₅NO₈ 535.676
Alkaloid from aerial parts of *Consolida hellespontica* (Ranunculaceae). Mp 138-139°.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 8-benzoyl, 14-Ac: **Ezochasmanine. 14-Acetyl-8-benzoylchasanine**
[77757-04-1]
C₃₄H₄₇NO₈ 597.747
Minor alkaloid from *Aconitum yesoense* (Ranunculaceae). Prisms (Me₂CO). Mp 163-165°.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Crassicaudine. 8-Acetyl-14-benzoylchasanine**
[4296-54-2]
C₃₄H₄₇NO₈ 597.747
Alkaloid from underground parts of *Aconitum crassicaule*, *Aconitum sungpanense* and *Aconitum longtounense* (Ranunculaceae). Cryst. (Et₂O). Mp 148-150° (143-145°). [α]_D²⁷ +15 (c, 0.32 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 8-O-(4-methoxybenzoyl), 14-Ac: **Anisoezochasmanine. 14-Acetyl-8-anisoylchasanine**
[77757-05-2]
C₃₅H₄₉NO₉ 627.773
Minor alkaloid from *Aconitum yesoense* (Ranunculaceae). Cryst. (Et₂O). Mp 136-138.5°.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl): **Acoforesticine**
[110011-76-2]
C₃₃H₄₇NO₈ 585.736
Alkaloid from roots of *Aconitum forrestii* (Ranunculaceae). Amorph. [α]_D +40.5 (c, 0.92 in EtOH).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **Foresaconitine. Vilmorrianine C**
[73870-35-6]
C₃₅H₄₉NO₉ 627.773
Alkaloid from roots of *Aconitum forrestii*, *Aconitum vilmorrianum* and *Aconitum pseudogeniculatum* (Ranunculaceae). Cryst. (Me₂CO/MeOH). Mp 153-154°. [α]_D²⁰ +30.5 (CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-hexadecanoyl: **Dolichotine D. 8-Deacetyl-8-palmitoylvilmorrianine C**
[128364-24-9]
C₄₉H₇₇NO₉ 824.149
Minor alkaloid from roots of *Aconitum dolichorhynchum* var. *subfabratum* (Ranunculaceae). Amorph.
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Falconeridine. 14-Veratroylchasanine**
[121880-22-6]
C₃₄H₄₉NO₉ 615.762
Alkaloid from roots of *Aconitum falconeri* (Ranunculaceae). Amorph. [α]_D²³ +50.8 (c, 0.191 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl), 8-Ac: **Falconericine. 8-Acetyl-14-veratroylchasanine**
[121880-21-5]
C₃₆H₅₁NO₁₀ 657.8
Alkaloid from the roots of *Aconitum falconeri* (Ranunculaceae). Amorph. solid. [α]_D²⁴ +16.7 (c, 0.336 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 8-E-cinnamoyl, 14-Ac: **Leucantine B**
[499969-76-5]
C₃₆H₄₉NO₈ 623.785
Alkaloid from the roots of *Aconitum hemsleyanum* var. *leucanthus*. Amorph. powder. [α]_D²⁰ +19 (c, 0.5 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-E-cinnamoyl: **Leucanthumsine B**
[947591-60-8]
C₃₄H₄₇NO₇ 581.748
Alkaloid from the roots of *Aconitum sungpanense* var. *leucanthum*. Amorph. powder. Mp 97-98°. [α]_D²⁰ +41.7 (c, 1 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-E-cinnamoyl, 8-Ac: **Leucanthumsine A**
[947591-59-5]
C₃₆H₄₉NO₈ 623.785
Alkaloid from the roots of *Aconitum sungpanense* var. *leucanthum*. Amorph. powder. Mp 100-102°. [α]_D²⁰ +12.1 (c, 1 in CHCl₃).
- O¹, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et: **14-O-Methyl-6-epiforesticine**
[146028-68-4]
C₂₅H₄₁NO₆ 451.602
Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Cryst. (Me₂CO). Mp 127-129°. [α]_D +20.9 (c, 0.134 in CHCl₃).
- O¹, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et, 6-Ac: **Acoseptrigine. 6-Acetyl-14-O-methyl-foresticine**
[144074-86-2]
C₂₇H₄₃NO₇ 493.639
Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Amorph. [α]_D +20.4 (c, 0.35 in CHCl₃).
- O⁶, O⁸, O¹⁶, O¹⁸-Tetra-Me, N-Et: **Consolinine**
[226697-96-7]
C₂₅H₄₁NO₆ 451.602
Alkaloid from *Consolida hohneckeri*. [α]_D +184 (c, 2 in MeOH).
- O¹, O⁶, O⁸, O¹⁴, O¹⁶-Penta-Me, N-Et: **Ho-**
- heconsoline**
[226697-95-6]
C₂₆H₄₃NO₆ 465.629
Alkaloid from *Consolida hohneckeri*. [α]_D +27 (c, 1.2 in CHCl₃).
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et: **Homochasmanine**
[3512-83-2]
C₂₆H₄₃NO₆ 465.629
Alkaloid from the roots of *Aconitum chasmanthum* (Ranunculaceae). Prisms (hexane). Mp 105-107°. [α]_D²⁵ +19.2 (c, 2.4 in EtOH). Unstable and dec. gradually at r.t.
- O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-Penta-Me, N-Et: **6,14-Di-O-methyl-6-epiforesticine**
[91794-17-1]
C₂₆H₄₃NO₆ 465.629
Alkaloid from roots of *Aconitum leucostomum*. Amorph. solid.
- 1-Ketone, O⁶, O¹⁶, O¹⁸-tri-Me, N-Et, 6,14-di-Ac: **1-Dehydrodeiphisine. 1-Ketodeiphisine**
[55085-42-2]
C₂₈H₄₁NO₈ 519.634
Alkaloid from seeds of *Delphinium staphisagria* (Ranunculaceae), also obt. by oxidn. of Deiphisine. Prisms (Me₂CO/Et₂O). Mp 170-171° (168-170°).
- 16-Ketone, O⁶, O¹⁸-di-Me, N-Et: **Staphisadrinine**
[132210-59-4]
C₂₃H₃₅NO₆ 421.533
Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Amorph. [α]_D¹⁹ -19.1 (c, 0.15 in CHCl₃).
- (1α,5β,6β,14α,16β)-form**
O⁶, O¹⁶-Di-Me, N-Et: **6-Epineoline**
[125263-89-0]
C₂₃H₃₇NO₆ 423.548
Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Gum. [α]_D²³ +50 (CHCl₃). Plain positive dispersion curve to 365 nm.
- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et: **Foresticine. Longtouconitine B. 6β-Hydroxytalatzamine**
[91794-15-9]
C₂₄H₃₉NO₆ 437.575
Alkaloid from the roots of *Aconitum forrestii* and *Aconitum longtounense* (Ranunculaceae). Mp 79-80°. [α]_D²¹ -1.9 (c, 1 in CHCl₃). C-6 config. revised in 2000.
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et: **Subcusine. Deacetylsubcusine. 6-Epineoline**
[81037-25-4]
C₂₄H₃₉NO₆ 437.575
Alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 194-196°. [α]_D +33.3 (c, 0.3 in CHCl₃).
- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-Ac: **Subcusmine**
[119152-45-3]
C₂₆H₄₁NO₇ 479.612
Alkaloid from roots of *Aconitum japonicum* (Ranunculaceae). Cryst. (Me₂CO/hexane). Mp 200-202°. [α]_D +18.3 (c, 0.48 in CHCl₃).

- O^8, O^{16}, O^{18} -Tri-Me, N-Et, 14-Ac: **Nuttallianine**. 7-Deoxypubescenine [125263-87-8] $C_{26}H_{41}NO_7$ 479.612
Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Needles. Mp 84-86°. Plain -ve ORD dispersion curve. Config. of C-6 OH revised in 1994.
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et: **6-Epichasmanine** [178699-06-4] $C_{25}H_{41}NO_6$ 451.602
Alkaloid from roots of *Aconitum kusnezoffii*.
- (1 β ,5 β ,6 α ,14 α ,16 β)-form**
- O^6, O^{16}, O^{18} -Tri-Me, N-Et: **Delphirine**. 1-Epineoline [60208-28-8] $C_{24}H_{39}NO_6$ 437.575
Minor alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Cryst. (C_6H_6). Mp 95-100°. $[\alpha]_D^{25} +3.8$ (c, 1.0 in EtOH).
- O^6, O^{16}, O^{18} -Tri-Me, N-Et, 14-Ac: **14-Acetylpineoline** [123795-80-2] $C_{26}H_{41}NO_7$ 479.612
Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Amorph. $[\alpha]_D^{25} +4.8$ (c, 0.2 in EtOH aq.).
- O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **1-Epicrassicaudine** [163083-76-9] $C_{34}H_{47}NO_8$ 597.747
Alkaloid from roots of *Aconitum hemsleyanum* var. *pengshinense* (Ranunculaceae). CAS abstract diag. shows the 1,6-diepimeric struct.
- Achmatowicz, O. *et al.*, *Can. J. Chem.*, 1964, **42**, 154; 1965, **43**, 825-839; 1093-1095 (*Homochasmanine*, *Chasmanine*)
- Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1964, **30**, 139; *CA*, **61**, 8128e (*Neoline*, *Bullatine C*, *isol*)
- Pelletier, S.W. *et al.*, *J.A.C.S.*, 1974, **96**, 7817-7818; 1976, **98**, 2617-2625; 2626-2636 (*struct*, *abs config*, *stereochem*, *cmr*, *Neoline*, *Delphisine*, *Chasmanine*, *Homochasmanine*)
- Pelletier, S.W. *et al.*, *Chem. Comm.*, 1976, 253-254 (*Chasmanine*, *cryst struct*)
- Tsai, T.Y.R. *et al.*, *Heterocycles*, 1977, **7**, 217-226 (*Chasmanine*, *synth*)
- Pelletier, S.W. *et al.*, *Phytochemistry*, 1977, **16**, 404-405 (*Delphidine*)
- Wiesner, K. *et al.*, *Can. J. Chem.*, 1978, **56**, 1451-1454 (*Chasmanine*, *synth*)
- Chen, W. *et al.*, *Chem. Ber.*, 1981, **114**, 394-397 (*Foresaconitine*)
- Takayama, H. *et al.*, *Heterocycles*, 1981, **15**, 403-408 (*Ezoachasmanitine*, *Anisoezoachasmanitine*)
- Yang, C.R. *et al.*, *Huaxue Xuebao*, 1981, **39**, 147-152; *CA*, **95**, 58094m (*Foresaconitine*)
- Konno, C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 128-133 (*Senbusine A*, *isol*, *ir*, *pmr*, *cmr*, *struct*)
- Hikino, H. *et al.*, *J. Nat. Prod.*, 1983, **46**, 178-182; 1984, **47**, 190 (*Neoline*, *Senbusine A*, *14-Acetylpineoline*)
- Pelletier, S.W. *et al.*, *J. Nat. Prod.*, 1984, **47**, 474-477 (*Forensicine*)
- Sakai, S. *et al.*, *Yakugaku Zasshi*, 1984, **104**, 222-237; *CA*, **101**, 69329k (*Senbusine A*, *isol*)
- Wada, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3658-3661 (*14-Benzoylpineoline*)
- Pelletier, S.W. *et al.*, *Heterocycles*, 1985, **23**, 2873-2883 (*Delstaphisine*, *Delstaphisagnine*, *Delstaphisagrine*)
- Luo, S. *et al.*, *Huaxue Xuebao*, 1985, **43**, 577-580; *CA*, **103**, 157320x (*Longtouconitine B*)
- Hao, X. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1985, **27**, 504-509; *CA*, **104**, 31721f (*Geniconitine*)
- Luo, S. *et al.*, *Planta Med.*, 1986, 412-413 (*8-Acetyl-14-benzoylchasmamine*)
- Pelletier, S.W. *et al.*, *Heterocycles*, 1987, **25**, 365-376 (*Acoforesticine*)
- Ross, S.A. *et al.*, *Heterocycles*, 1987, **26**, 2895-2904 (*Neoline*)
- Wang, F.-D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 55-62 (*Crassicaudine*)
- Pelletier, S.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 381-385 (*1-Dehydrodelphisine*)
- De la Fuente, G. *et al.*, *Heterocycles*, 1988, **27**, 1109-1113 (*1,14-Diacetylpineoline*)
- Bando, H. *et al.*, *Heterocycles*, 1988, **27**, 2167-2174 (*Subcumine*, *Subcusine*)
- Ross, S.A. *et al.*, *J. Nat. Prod.*, 1988, **51**, 572-577 (*Delstaphisine*, *1-Acetyldephisine*)
- Batbayar, N. *et al.*, *Khim. Prir. Soedin.*, 1988, **24**, 237-239; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **24**, 200-202 (*Bataconine*)
- De la Fuente, G. *et al.*, *Heterocycles*, 1989, **29**, 205 (*Pictumine*)
- Desai, H.K. *et al.*, *Heterocycles*, 1989, **29**, 225-230 (*Falconeridine*, *Falconericine*)
- Bai, Y. *et al.*, *Heterocycles*, 1989, **29**, 1017-1021 (*Nuttallianine*, *6-Epineolinine*)
- Huiling, L. *et al.*, *Heterocycles*, 1989, **29**, 2317-2326 (*Dolichotine D*)
- Liang, X. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1307-1311 (*14-Acetylpineoline*, *Staphisadrinine*)
- Ding, L.-S. *et al.*, *Phytochemistry*, 1990, **29**, 3694-3696 (*Patentine*)
- Sayed, H.M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1595-1606 (*14-O-Methyl-6-epiforesticine*)
- Ross, S.A. *et al.*, *Tetrahedron*, 1992, **48**, 1183-1192 (*Acoseptrigine*, *Acoseptriginine*)
- Desai, H.K. *et al.*, *Heterocycles*, 1993, **36**, 1081-1089 (*1-O-Methyldephisine*)
- Yunosov, M.S. *et al.*, *Nat. Prod. Rep.*, 1993, **10**, 471-486 (*rev*, *1,6,14-Tribenzoylsenbusine A*)
- Hanuman, J.B. *et al.*, *Phytochemistry*, 1994, **36**, 1527-1535 (*14-O-Acetylsenbusine A*)
- Bai, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1717-1724 (*Senbusine A*, *Nuttallianine*, *config*)
- Wang, F.P. *et al.*, *Chin. Chem. Lett.*, 1995, **6**, 109; *CA*, **122**, 291232r (*1-Epicrassicaudine*)
- Wei, X. *et al.*, *Zhongcaoyao*, 1995, **26**, 344-346; *CA*, **124**, 170554e (*6,14-Di-O-methyl-6-epiforesticine*)
- Li, Z.B. *et al.*, *Chin. Chem. Lett.*, 1996, **7**, 443-444; *CA*, **125**, 86949a (*6-Epichasmanine*)
- Kolak, U.S. *et al.*, *Sci. Pharm.*, 1998, **66**, 381-385 (*14-Acetylpineoline*)
- Ulubelen, A. *et al.*, *Phytochemistry*, 1999, **50**, 909-912 (*Consolinine*, *Hoheconsoline*)
- Wada, K. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 660-668 (*ms*)
- Wang, F.P. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 1003-1004 (*Foresticine*, *6-Epiforesticine*)
- Li, Z.-B. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 131-137 (*Geniculatines A,B*)
- Shim, S.H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 999-1002 (*14-Anisoylpineoline*, *14-Cinnamoylpineoline*, *14-Veratroylpineoline*)
- Li, L.-Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 269-271 (*Leueantines B,D*)
- Taki, M. *et al.*, *Planta Med.*, 2003, **69**, 800-803 (*8-Cinnamoylpineoline*)
- Ping, A. *et al.*, *J. Asian Nat. Prod. Res.*, 2004, **6**, 151-154 (*Kongboentine B*)
- Csupor, D. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 2981-2986 (*Acotoxinine*)
- Yan, H. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 1133-1140 (*Leucanthumsines A-C*)

4-Methylnaconitane-1,7,8,9,14,16-hexol M-324

 $C_{19}H_{29}NO_6$ 367.441**(1 α ,5 β ,14 α ,16 β)-form** O^{16} -Me, N-Et: **Tatsinine**


[90038-21-4]

 $C_{22}H_{35}NO_6$ 409.522Alkaloid from the roots of *Delphinium tatsienense* (Ranunculaceae). Mp 163-164°. $[\alpha]_D +9$ (EtOH). O^{16} -Me, N-Et, perchlorate: Mp 132-133°.7,8-Methylene ether, O^{16} -Me, N-Et: **Tatsidine**

[126617-62-7]

 $C_{23}H_{35}NO_6$ 421.533Alkaloid from the roots of *Delphinium tatsienense* (Ranunculaceae). Cryst. (CH_2Cl_2 /hexane). Mp 191.5-193.5°.Glinski, J.A. *et al.*, *Tet. Lett.*, 1984, **25**, 1211(*Tatsinine*, *ir*, *pmr*, *cmr*, *struct*)Maddry, J.A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 674(*Tatsinine*, *cryst struct*)Snyder, J.K. *et al.*, *Magn. Reson. Chem.*, 1989, **27**, 1057 (*Tatsidine*, *pmr*, *cmr*)Joshi, B.S. *et al.*, *Phytochemistry*, 1990, **29**, 357 (*Tatsidine*, *isol*, *pmr*, *cmr*, *struct*)

4-Methylnaconitane-1,7,8,14,16,18-hexol M-325

 $C_{19}H_{29}NO_6$ 367.441**(1 α ,5 β ,14 α ,16 β)-form** O^{16} -Me, N-Et: **1-Demethylwinkleridine**.

18-Demethylvirescenine

[194149-12-7]

 $C_{22}H_{35}NO_6$ 409.522Alkaloid from aerial parts of *Aconitella hohenackeri* (*Consolida hohenackeri*). Cryst. (EtOAc/MeOH). Mp 204-205°. $[\alpha]_D +25.6$ (c, 0.43 in MeOH). O^1, O^{16} -Di-Me, N-Et: **Winkleridine**

[128700-88-9]

 $C_{23}H_{37}NO_6$ 423.548Alkaloid from *Delphinium winklerianum* (Ranunculaceae). Rhombic cryst. (MeOH). Mp 215-217°. O^8, O^{16} -Di-Me, N-Et, 14-Ac: **Ajadelpine**

[142808-41-1]

 $C_{25}H_{39}NO_7$ 465.586

Alkaloid from roots of *Delphinium ajacis* (Ranunculaceae). Amorph. $[\alpha]_D^{24} +2$ (c, 0.13 in CHCl_3).

O¹⁶, O¹⁸-Di-Me, N-Et: Virescine

[71609-78-4]
C₂₃H₃₇NO₆ 423.548

Alkaloid from *Delphinium virescens* (whole plant) (Ranunculaceae). Cryst. (Et₂O/hexane). Mp 68-70°. $[\alpha]_D^{24} +16.9$ (c, 1.0 in EtOH).

O¹⁶, O¹⁸-Di-Me, N-Et, 14-Ac: 14-Acetyl-virescine

[71609-79-5]
C₂₅H₃₉NO₇ 465.586

Alkaloid from *Delphinium virescens* (whole plant) (Ranunculaceae). Cryst. (Et₂O/hexane). Mp 157-159°. $[\alpha]_D^{24} +31.8$ (c, 0.5 in CHCl_3).

O¹, O¹⁶, O¹⁸-Tri-Me, N-Et: Racemuloline A

[463933-00-8]
C₂₄H₃₉NO₆ 437.575

Alkaloid from *Aconitum racemosum* var. *pengzhouense*. Amorph. powder. $[\alpha]_D^{24} +11.2$ (c, 0.5 in CHCl_3).

O¹⁴, O¹⁶, O¹⁸-Tri-Me, N-Et: Umbrosine.

Paniculatin†

[63201-50-3]
C₂₄H₃₉NO₆ 437.575

Alkaloid from the roots of *Aconitum umbrosum* and from *Consolida regalis* ssp. *paniculata* (Ranunculaceae). Mp 150-151°. Identity of *Paniculatin* with *Umbrosine* not apparently definitely establ. although CA gives them the same reg. no. Ref. for *Paniculatin* inaccessible: the CA abstract shows it with undefined C(1) config. and omits the C-14 OMe group.

7,8-Methylene, O¹⁶-Me, N-Et: Ajadelphinine

[142808-42-2]
C₂₃H₃₅NO₆ 421.533

Alkaloid from roots of *Delphinium ajacis* (Ranunculaceae). Amorph. $[\alpha]_D^{24} -22.6$ (c, 0.32 in CHCl_3).

7,8-Methylene, O¹⁶, O¹⁸-di-Me, N-Et:

Winkleriline

[128700-86-7]
C₂₄H₃₇NO₆ 435.559

Alkaloid from the whole plant of *Delphinium winklerianum* (Ranunculaceae). Rhombic cryst. (Me₂CO/petrol). Mp 131-134° (as di-Ac). $[\alpha]_D^{21} +89.7$ (c, 0.39 in CHCl_3).

7,8-Methylene, O¹, O¹⁴, O¹⁶-tri-Me, N-Et:

Hohenackerine

[126223-69-6]
C₂₅H₃₉NO₆ 449.586

Alkaloid from the aerial parts of *Consolida hohackeri* (Ranunculaceae).

7,8-Methylene, O¹, O¹⁴, O¹⁸-tri-Me, N-Et:

Corumdefine

[98752-03-5]
C₂₅H₃₉NO₆ 449.586

Alkaloid from *Delphinium corumbosum* (Ranunculaceae). Powder.

7,8-Methylene, O¹, O¹⁶, O¹⁸-tri-Me, N-Et:

Nudicaulamine

[99815-84-6]
C₂₅H₃₉NO₆ 449.586

Alkaloid from *Delphinium nudicaule*

(whole plant) (Ranunculaceae). Non-cryst.; cryst. (as perchlorate). Mp 212-214° (perchlorate). $[\alpha]_D^{24} -27$ (c, 0.36 in CHCl_3).

7,8-Methylene, O¹, O¹⁴, O¹⁶, O¹⁸-tetra-Me, N-Et: Deoxydelcorine

[54387-74-5]
C₂₆H₄₁NO₆ 463.613

Alkaloid from *Delphinium corumbosum* (Ranunculaceae). Mp 93-95°.

Narzullaev, A.S. et al., *Khim. Prir. Soedin.*, 1974, **10**, 412; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 420 (*Nudicaulamine, Deoxydelcorine*)

Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1976, **12**, 675; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 610 (*Umbrosine*)

Pelletier, S.W. et al., *Heterocycles*, 1979, **12**, 779-782 (*Virescine, 14-Acetylvirescine*)

Kulanthavil, P. et al., *Heterocycles*, 1985, **23**, 2515 (*Nudicaulamine, Deoxydelcorine*)

Salimov, B.T. et al., *Khim. Prir. Soedin.*, 1985, **21**, 95; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 91 (*Corumdefine*)

Sener, B. et al., *CA*, 1988, **110**, 111674g; 1990, **112**, 175528 (*Paniculatin, Hohenackerine*)

Chen, Y.Z. et al., *Phytochemistry*, 1990, **29**, 1016-1019 (*Winkleridine, Winkleriline*)


Pelletier, S.W. et al., *J. Nat. Prod.*, 1992, **55**, 736 (*Ajadelphine, Ajadelphinine*)

Almanza, G. et al., *Phytochemistry*, 1997, **45**, 1079-1085 (*1-Demethylwinkleridine*)

Peng, C.-S. et al., *Heterocycles*, 2002, **57**, 1117-1120 (*Racemuloline A*)

Khairitdinova, E.D. et al., *Russ. Chem. Bull. (Engl. Transl.)*, 2003, **52**, 2078-2080 (*Virescine, isol, cmr*)

4-Methylnaconitane-1,8,10,14,16,18-hexol M-326



C₁₉H₂₉NO₆ 367.441

(1 α , 14 α , 16 β)-form

O¹⁶, O¹⁸-Di-Me, N-Et: 10-Hydroxyisotalatizidine

[121687-87-4]
C₂₃H₃₇NO₆ 423.548

Alkaloid from aerial parts of *Aconitum sanyoense* var. *tonenze* (Ranunculaceae). Amorph. solid. $[\alpha]_D^{14} +7$ (c, 0.28 in CHCl_3).

O¹, O¹⁶, O¹⁸-Tri-Me, N-Et: 10-Hydroxy-talatizamine

[121687-88-5]
C₂₄H₃₉NO₆ 437.575

Alkaloid from aerial parts of *Aconitum sanyoense* var. *tonenze* (Ranunculaceae). Amorph. solid. $[\alpha]_D^{20} +6.4$ (c, 0.33 in CHCl_3).

14-Ketone, O¹⁶, O¹⁸-di-Me, N-Et: Genicine C

[287186-11-2]
C₂₃H₃₅NO₆ 421.533


Alkaloid from the roots of *Aconitum*

geniculatum. $[\alpha]_D^{24} -14.7$ (c, 1 in CHCl_3).

Takayama, H. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2927 (*10-Hydroxyisotalatizidine, 10-Hydroxytalatizamine*)

Wang, F.P. et al., *Huaxue Xuebao*, 2002, **58**, 576-579 (*Genicine C*)

4-Methylnaconitane-1,8,14,15,16,18-hexol M-327



C₁₉H₂₉NO₆ 367.441

(1 β , 5 β , 14 α , 15 α , 16 β)-form


O¹⁶, O¹⁸-Di-Me, N-Et: Senbusine B. 15-Hydroxyisotalatizidine

[82202-97-9]
C₂₃H₃₇NO₆ 423.548

Alkaloid from the roots of *Aconitum carmichaeli* (Ranunculaceae). Amorph. powder.

Konno, C. et al., *J. Nat. Prod.*, 1982, **45**, 128 (*isol, ir, pmr, cmr, struct*)

4-Methylnaconitane-6,7,8,14,16,18-hexol M-328



C₁₉H₂₉NO₆ 367.441

(5 β , 6 β , 14 α , 16 β)-form


O⁶, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et: Deoxydelsole


[58111-35-6]
C₂₅H₄₁NO₆ 451.602

Alkaloid from the aerial parts of *Aconitum monticola* (Ranunculaceae).


Ametova, E.F. et al., *Khim. Prir. Soedin.*, 1982, **18**, 504-507; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 472-474 (*Deoxydelsole*)

4-Methylnaconitane-1,2,3,6,8,13,14,16,18-nonol M-329



C₁₉H₂₉NO₉ 415.439**(1 α ,2 β ,3 α ,5 β ,6 α ,14 α ,16 β)-form**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Atropurpursine**
[1007889-96-4]C₃₄H₄₇NO₁₁ 645.745Alkaloid from the roots of *Aconitum hemsleyanum* var. *atropurpureum*. Amorph. powder. Mp 224-226°. [α]_D²⁰ +25.5 (c, 1 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **Transconitine B**
[175992-66-2]C₃₅H₄₉NO₁₂ 675.772Alkaloid from roots of *Aconitum transsectum*. Amorph. [α]_D²⁰ +12.3 (c, 0.02 in CHCl₃).Zheng, S. et al., *Phytochemistry*, 1997, **46**, 951-954 (*Transconitine B*)Tang, P. et al., *Chin. Chem. Lett.*, 2007, **18**, 704-707 (*Atropurpursine*)**4-Methylnaconitane-1,2,6,8,13,14,15,16,18-nonol M-330**C₁₉H₂₉NO₉ 415.439**(1 α ,2 α ,5 β ,6 α ,14 α ,15 α ,16 β)-form**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Karakonitine**. 2-Hydroxy-3-deoxyaconitine
[178960-46-8]C₃₄H₄₇NO₁₁ 645.745Alkaloid from roots of *Aconitum karakolium*. Mp 222-224°. [α]_D +7.96 (c, 0.47 in MeOH).Lao, A. et al., *Heterocycles*, 1996, **43**, 1267-1270 (*Karakonitine*)**4-Methylnaconitane-1,3,6,8,13,14,15,16,18-nonol M-331**

[107965-53-7]

C₁₉H₂₉NO₉ 415.439**(1 α ,3 α ,6 α ,14 α ,15 α ,16 β)-form**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, 14-benzoyl, 8-Ac: **N-Deethylaconitine**
[3327-35-3]C₃₂H₄₃NO₁₁ 617.692Alkaloid from the roots of *Aconitum napellus* ssp. *vulgare* (Ranunculaceae).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl: 14-Benzoylmesaconine. **Hokbusine A**
[86500-43-8]C₃₁H₄₃NO₁₀ 589.681Alkaloid from roots of *Aconitum napellus* and *Aconitum carmichaeli* (Ranunculaceae). Amorph. powder. [α]_D +11.4 (c, 0.23 in MeOH).**▶ AR5569200**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 8-Ac: **Mesaconitine**. N-Desethyl-N-methylnaconitine. *Japaconitine A*. *Japaconitine B*
[2752-64-9]C₃₃H₄₅NO₁₁ 631.719Alkaloid from *Aconitum manshuricum* and many other *Aconitum* spp. (Ranunculaceae). Mp 208-209° dec. [α]_D¹⁷ +25.4 (CHCl₃).**▶ AR5563000**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 8-acyl: **Lipomesaconitine**
[81941-16-4, 158584-86-2]From *Aconitum carmichaeli*(Ranunculaceae). Oil. [α]_D¹³ +13.8 (CHCl₃). Analogue of Mesaconitine with a mixt. of long-chain acyl groups replacing Ac.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 14-benzoyl, 3,8-di-Ac: **Tuberanine**
[121897-29-8]C₃₅H₄₇NO₁₂ 673.756Alkaloid from the roots of *Aconitum tuberosum*. Cryst. (Me₂CO). Mp 253-255°.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et: **Aconine**. *Jesaconine*
[509-20-6]C₂₅H₄₁NO₉ 499.6Alkaloid from the roots of *Aconitum soongoricum* (Ranunculaceae), also obt. by hydrol. of Aconitine, A-104. Possesses gastric anaesthetic props. Febrifuge. Amorph. Sol. H₂O, insol. Et₂O. Mp 132°. [α]_D +23 (H₂O). Log P -7.18 (uncertain value) (calc).**▶ Poisonous.**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, hydrochloride: Mp 175-176°. [α]_D -8 (H₂O).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 8-O-(9Z,12Z-octadecadienyl): **Aconine 8-linoleate**
[1005031-40-2]C₄₃H₇₁NO₁₀ 762.035Alkaloid from the roots of *Aconitum flavum*.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Me, 8,14-dibenzoyl: **Manshuritine**
C₃₈H₄₇NO₁₁ 693.789Alkaloid from the roots of *Aconitum manshuricum*. Amorph. powder. [α]_D +38 (c, 0.67 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl: 14-Benzoylconine. **Picraconitine**
[466-24-0]C₃₂H₄₅NO₁₀ 603.708Alkaloid from the roots of *Aconitum**carmichaeli* and *Aconitum**polyschistum* (Ranunculaceae). Also isol. from the processed aconite "Kako-bushi-matsu", produced by autoclaving the crude drug ("bushi") obt. from the roots of some *Aconitum* spp. Amorph. Mp 130°. [α]_D +5 (EtOH).**▶ TJ7849250**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 3-Ac: 3-Acetyl-14-benzoylconine. **Polyschistine D**
[119347-26-1]C₃₄H₄₇NO₁₁ 645.745Alkaloid from the roots of *Aconitum polyschistum* (Ranunculaceae). Needles (EtOAc). Mp 251-252°. [α]_D +11.4 (CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-Ac: see Aconitine, A-104O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-O-(3E-hexenoyl): **Brachyaconitine**
[173220-47-8]C₃₈H₅₃NO₁₁ 699.837Alkaloid from aphids *Brachycaudus aconiti* and *Brachycaudus napelli* feeding on *Aconitum napellus*. Oil. [α]_D²² 0 (c, 0.07 in CHCl₃).O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-(8-carboxyooctanoyl): **8-O-Azaloil-14-benzoylconine**
[869496-03-7]C₄₁H₅₉NO₁₃ 773.916Alkaloid from the roots of *Aconitum karacolicum*. Zwitterionic.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-benzoyl, 8-acyl: **Lipoaconitine**
[81941-14-2, 158584-72-6, 158584-73-7]

Alkaloid from Aconiti tuber

Aconitum carmichaeli (Ranunculaceae). Oil. [α]_D¹³ +6 (CHCl₃). Analogue of Aconitine, A-104 in which the esterifying group on C₈ is a mixt. of long-chain acyl (linoleoyl, palmitoyl, oleoyl, stearoyl, linolenoyl) instead of acetyl.O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl): **14-Anisoilaconine**
[121923-73-7]C₃₃H₄₇NO₁₁ 633.734

Alkaloid from the processed aconite "Kako-bushi-matsu". Amorph. Mp 183-184°.

▶ AR5562580O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **Jesaconitine**
[16298-90-1]C₃₅H₄₉NO₁₂ 675.772Alkaloid from *Aconitum fischeri*, *Aconitum subcuneatum*, *Aconitum sachalinense*, *Aconitum yesoense* and *Aconitum mitakense* (Ranunculaceae). Amorph. Mp 128-131°.**▶ AR5562540**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, 8,15-O-isopropylidene, N-Me, 14-benzoyl: **Tuberme-saconitine**
[405157-68-8]C₃₄H₄₇NO₁₀ 629.746Alkaloid from the roots of *Aconitum tuberosum*. Amorph. solid.

O^1, O^6, O^{16}, O^{18} -*Tetra-Me, 8,15-O-isopropylidene, N-Et, 14-benzoyl*: **Tuberacoinitine**

[405157-67-7]

$C_{35}H_{49}NO_{10}$ 643.773

Alkaloid from the roots of *Aconitum tuberosum*.

$O^1, O^6, O^8, O^{16}, O^{18}$ -*Penta-Me, N-Et, 14-benzoyl*: **14-Benzoyl-8-O-methylaconitine**

[93772-68-0]

$C_{33}H_{47}NO_{10}$ 617.735

Alkaloid from *Aconitum pseudostaphanium*. Component of Ni Yu Long Wu Tou. Amorph. $[\alpha]_D^{20}$ -2.4 (c, 0.72 in $CHCl_3$).

$O^1, O^6, O^8, O^{16}, O^{18}$ -*Penta-Me, N-Et, 14-O-(4-methoxybenzoyl)*: **Aljesaconitine A**

[101247-23-8]

$C_{34}H_{49}NO_{11}$ 647.761

Alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Also obtained from Jesaconitine + MeOH at r.t. Amorph. powder. $[\alpha]_D^{20}$ +7.5 (c, 0.93 in EtOH).

▶ AR5568500

O^8 -*Et, O^1, O^6, O^{16}, O^{18}*-*tetra-Me, N-Et, 14-benzoyl*: **Spicatinine A. 14-Benzoyl-8-O-ethylaconitine**

[124256-81-1]

$C_{34}H_{49}NO_{10}$ 631.762

Alkaloid from aerial parts of *Aconitum napellus* and the roots of *Aconitum spicatum* (Ranunculaceae). Amorph. powder. Mp 139-141°. $[\alpha]_D$ -3 (c, 0.10 in EtOH). $[\alpha]_D^{27}$ -1 (c, 1 in $CHCl_3$). Possibly an artifact.

O^8 -*Et, O^1, O^6, O^{16}, O^{18}*-*tetra-Me, N-Et, 14-benzoyl, 3-Ac*: **Polyschistine A. Szechenyine**

[96562-88-8]

$C_{36}H_{51}NO_{11}$ 673.799

Alkaloid from the roots of *Aconitum polyschistum* and *Aconitum szechenyianum* (Ranunculaceae). Mp 265-266°.

O^8 -*Et, O^1, O^6, O^{16}, O^{18}*-*tetra-Me, N-Et, 3,14,15-tribenzoyl*: **8-O-Ethyl-3,14,15-tribenzoyl-aconitine**

$C_{48}H_{57}NO_{12}$ 839.978

Alkaloid from *Delphinium montanum*.

O^8 -*Et, O^1, O^6, O^{16}, O^{18}*-*tetra-Me, N-Et, 14-O-(4-methoxybenzoyl)*: **Aljesaconitine B**

[101247-24-9]

$C_{35}H_{51}NO_{11}$ 661.788

Alkaloid from roots of *Aconitum japonicum* (Ranunculaceae). Also obtained from Jesaconitine + EtOH. Amorph. powder. $[\alpha]_D^{20}$ +5.8 (c, 1.18 in EtOH).

▶ AR5565200

(1β,3α,6α,14α,15α,16β)-form

O^1, O^6, O^{16}, O^{18} -*Tetra-Me, N-Et, 14-benzoyl*: **De-O-acetyl-1-epiaconitine**

[303154-28-1]

$C_{32}H_{45}NO_{10}$ 603.708

Alkaloid from the roots of *Aconitum nemorum*. Cryst. Mp 215-217°. $[\alpha]_D^{25}$ -21.2 (c, 0.25 in MeOH).

Makoshi, K. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1909, **247**, 243; *CA*, **3**, 2707; **5**, 674 (*Jesaconitine, isol*)

Morio, S. et al., *Annalen*, 1929, **476**, 181 (*Mesaconitine*)

Nath, B. et al., *J. Indian Chem. Soc.*, 1955, **32**, 75 (*Aconine, synth, ir*)

Schneider, W. et al., *Annalen*, 1959, **628**, 114 (*Aconine, struct*)

Keith, L.H. et al., *J.O.C.*, 1968, **33**, 2497 (*Jesaconitine, pmr, struct*)

Yunusov, M.S. et al., *Khim. Prir. Soedin.*, 1969, **5**, 515; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 434 (*Aconine, ms*)

Birnbaum, K.B. et al., *Tet. Lett.*, 1971, 867 (*config*)

Pelletier, S.W. et al., *J.A.C.S.*, 1976, **98**, 2626 (*Mesaconitine, cmr*)

Pelletier, S.W. et al., *Can. J. Chem.*, 1979, **57**, 1652 (*Aconine, cmr*)

Pelletier, S.W. et al., *Cryst. Struct. Commun.*, 1979, **8**, 299 (*Jesaconitine, crystal struct*)

Zhamierashvili, M.G. et al., *CA*, 1982, **97**, 107014n (*Aconine, occur*)

Kitagawa, I. et al., *Chem. Pharm. Bull.*, 1982, **30**, 758-761 (*Lipoaconitine, Lipomesaconitine*)

Hikino, H. et al., *J. Nat. Prod.*, 1983, **46**, 178; 1984, **47**, 190 (*Hokbusine A*)

Hikino, H. et al., *Shoyakugaku Zasshi*, 1983, **37**, 68; *CA*, **99**, 110568t (*Picraconitine*)

Bando, H. et al., *Chem. Pharm. Bull.*, 1985, **33**, 4717 (*Aljesaconitines*)

Wang, H. et al., *Heterocycles*, 1985, **23**, 803; 1988, **27**, 1615 (*Polyschistines*)

Arlandini, E. et al., *J. Nat. Prod.*, 1987, **50**, 937 (*N-Deethylaconitine*)

Kobayashi, K. et al., *Anal. Sci.*, 1989, **5**, 237; *CA*, **112**, 21173e (*Benzoyldeoxyaconine, crystal struct*)

Mori, T. et al., *Heterocycles*, 1989, **29**, 873 (*14-Anisoyl-aconine, Benzoyl-aconine*)

De la Fuente, G. et al., *Heterocycles*, 1989, **29**, 1577 (*8-O-Ethylbenzoyl-aconine*)

Desai, H.K. et al., *J. Nat. Prod.*, 1989, **52**, 720 (*Hokbusine A, synth*)

Sun, W.J. et al., *Yaoxue Xuebao*, 1989, **24**, 71; *CA*, **111**, 74778s (*Szechenyine*)

Yunusov, M.S. et al., *Nat. Prod. Rep.*, 1993, **10**, 471 (*8-O-Ethyl-3,14,15-tribenzoyl-aconine*)

Liu, H. et al., *J. Nat. Prod.*, 1996, **59**, 135 (*Brachyaconitine*)

Ying, C. et al., *Zhongcaoyao*, 1996, **27**, 5 (*14-Benzoyl-8-methyl-aconine*)

Sultankhodzhaev, M.N. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1999, **35**, 201-203 (*Tuberanine*)

Sun, W. et al., *Planta Med.*, 1999, **65**, 432-436 (*Lipoaconitine, Lipomesaconitine*)

Wei, X. et al., *Heterocycles*, 2000, **53**, 2027-2031 (*Deacetyl-1-epiaconitine*)

Boronova, Z.S. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 269-271 (*Tubermesaconitine, Tuberacoinitine*)

Chodoeva, A. et al., *Bioorg. Med. Chem.*, 2005, **13**, 6493-6501 (*8-Azeloxy-14-benzoyl-aconine*)

Gao, L.M. et al., *Chin. Chem. Lett.*, 2005, **16**, 475-478 (*Spicatinine A*)


Ishimi, K. et al., *J. Nat. Med. (Tokyo)*, 2006, **60**, 255-257 (*Manshuritine*)

Zhang, F. et al., *Zhongguo Yaoxue Zazhi*, 2006, **41**, 1851-1854; *CA*, **148**, 210217b (*Aconine 8-linoleate*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ADG500; PIC250

4-Methylnaconitane-1,6,8,10,13,14,15,16,18-nonol

M-332



$C_{19}H_{29}NO_9$ 415.439

(1α,6α,14α,15α,16β)-form

O^6, O^{16}, O^{18} -*Tri-Me, 14-benzoyl, 8-Ac*: **Flavaconitine**

[83145-43-1]

$C_{31}H_{41}NO_{11}$ 603.665

Alkaloid from the roots of *Aconitum flavum* (Ranunculaceae).

O^6, O^{16}, O^{18} -*Tri-Me, 14-benzoyl, 8-Ac, N-formyl*: **Flavaconidine. N-Formylflavaconitine**

[125286-17-1]

$C_{32}H_{41}NO_{12}$ 631.675

Alkaloid from *Aconitum flavum* (Ranunculaceae). Mp 186.5-189°. $[\alpha]_D$ -49 (c, 0.568 in DMF).

O^6, O^{16}, O^{18} -*Tri-Me, 14-benzoyl, 8,N-di-Ac*: **N-Acetylflavaconitine**

[125263-91-4]

$C_{33}H_{43}NO_{12}$ 645.702

Alkaloid from *Aconitum flavum* (Ranunculaceae). Mp 274.5-275.5°. $[\alpha]_D^{13}$ -45.6 (c, 1.54 in MeOH).

O^1, O^6, O^{16}, O^{18} -*Tetra-Me, N-Et, 14-benzoyl, 8-Ac*: **Polyschistine B. 10-Hydroxy-3-deoxyaconitine**

[96562-89-9]

$C_{34}H_{47}NO_{11}$ 645.745

Alkaloid from the roots of *Aconitum polyschistum* (Ranunculaceae). Mp 182-185°.


Liu, Y. et al., *Yaoxue Tongbao*, 1982, **17**, 243; *CA*, **97**, 141679t (*Flavaconitine*)

Wang, H. et al., *Heterocycles*, 1985, **23**, 803 (*Polyschistine B*)

Chen, Z. et al., *Heterocycles*, 1989, **29**, 997 (*Flavaconidine, N-Acetylflavaconitine*)

4-Methylnaconitane-1,2,6,8,13,14,16,18-octol

M-333



$C_{19}H_{29}NO_8$ 399.44

(1 α ,2 α ,5 β ,6 α ,14 α ,16 β)-form

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-methoxybenzoyl), 8-*Ac*: **Ouvrardiantine**
[951765-86-9]


C₃₅H₄₉NO₁₁ 659.772

Alkaloid from the roots of *Aconitum ouvardianum*. Amorph. powder. [α]_D²⁰ +25 (c, 0.8 in CHCl₃).

Hou, L.-H. et al., *Chem. Pharm. Bull.*, 2007, **55**, 1090-1092 (*Ouvrardiantine*)

4-Methylnaconitane-1,3,5,8,13,14,16,18-octol

M-334



C₁₉H₂₉NO₈ 399.44

(1 α ,3 α ,5 β ,14 α ,16 β)-form

O¹,O¹⁸-*Di-Me*, 14-O-(4-methoxybenzoyl), N-*Et*: **Circinasine B**
[939972-76-6]

C₃₁H₄₃NO₁₀ 589.681

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. [α]_D²⁰ +52.3 (c, 1 in CHCl₃).

O¹,O¹⁶,O¹⁸-*Tri-Me*, N-*Et*: **Circinasine F**
[939972-80-2]


C₂₄H₃₉NO₈ 469.574

Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Cubic cryst. (Me₂CO). Mp 239-240°. [α]_D²⁰ +24.8 (c, 1 in CHCl₃).

Gao, F. et al., *J. Nat. Prod.*, 2007, **70**, 876-879 (*isol. pmr, cmr, cryst struct*)

4-Methylnaconitane-1,3,6,8,13,14,15,16-octol

M-335



C₁₉H₂₉NO₈ 399.44

(1 α ,3 α ,6 α ,14 α ,15 α ,16 β)-form


O¹,O⁶,O⁸,O¹⁶-*Tetra-Me*, N-*Et*, 14-benzoyl: **Acofamine B**
C₃₂H₄₅NO₉ 587.709

Alkaloid from the aerial parts of *Aconitum karakolicum*. Amorph. [α]_D²⁰ +28 (c, 0.1 in CHCl₃). λ_{\max} 201 (log ϵ 4.18); 230 (log ϵ 4.24); 275 (log ϵ 2.86) (MeOH).

Atta-ur-Rahman, et al., *Nat. Prod. Res.*, 2005, **19**, 713-718 (*Acofamine B*)

4-Methylnaconitane-1,3,6,8,13,14,16,18-octol

M-336



C₁₉H₂₉NO₈ 399.44

► AR5566300

(1 α ,3 α ,6 α ,14 α ,16 α)-form

O⁶,O¹⁶,O¹⁸-*Tri-Me*, N-*Et*, 14-benzoyl, 8-*Ac*: **Leucanthumsine E**
[947412-91-1]

C₃₃H₄₅NO₁₀ 615.719

Alkaloid from the roots of *Aconitum sungpanense* var. *leucanthum*. Amorph. powder. Mp 95-96°. [α]_D²⁰ +13.2 (c, 1 in CHCl₃).

(1 α ,3 α ,6 α ,14 α ,16 β)-form

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, 14-O-(3,4-dimethoxybenzoyl), 8-*Ac*, N-formyl: **Faleoconitine**
[307495-69-8]

C₃₅H₄₇NO₁₃ 689.755

Alkaloid from *Aconitum falconeri*. Cryst. (MeOH). Mp 205°. [α]_D²⁵ -28 (c, 0.14 in MeOH). Exhibits atropisomerism. λ_{\max} 220 (log ϵ 3.29); 263 (log ϵ 3.01); 292 (log ϵ 2.75) (MeOH).

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*: **Pseudoconitine**
[38146-89-3]

C₂₅H₄₁NO₈ 483.601

Alkaloid from *Aconitum ferox*. Mp 90-92°.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-benzoyl: **Ludaconitine**
[82144-72-7]

C₃₂H₄₅NO₉ 587.709

Alkaloid from the roots of *Aconitum francheti* (Ranunculaceae).

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-benzoyl, 8-*Ac*: **Indaconitine**
[4491-19-4]

C₃₄H₄₇NO₁₀ 629.746

Alkaloid from roots of *Aconitum chasmanthum*, *Aconitum ferox*, *Aconitum falconeri*, *Aconitum franchetti* and *Aconitum violaceum* (Ranunculaceae). Cryst. (Et₂O). Mp 200-203° (190-191°). [α]_D²⁰ +18.3 (c, 2.0 in EtOH).

► AR5569430

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-benzoyl, 8-acyl: **Lipoindaconitine**

Alkaloid from root tubers of *Aconitum ferox* (Ranunculaceae). Oil. [α]_D²⁵ +1.1 (c, 0.5 in CHCl₃). The esterifying group on C-8 is a mixt. of long-chain acyl groups (linoleoyl, palmitoyl and stearoyl).

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-methoxybenzoyl): **8-Deacetylyunaconitine**
[93460-55-0]

C₃₃H₄₇NO₁₀ 617.735

Alkaloid from the roots of *Aconitum*

austroyunnanense, *Aconitum forrestii* and *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae).

Amorph. powder. Mp 101-105°.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-methoxybenzoyl), 8-*Ac*: **Yunaconitine**. *Guayewuanine B*. *Isoaconitine*. 8-Acetyl-14-anisoylpseudoaconitine. 3 α ,13-Dihydroxyforesaconitine. *Vilmorrianine B*
[70578-24-4]

C₃₅H₄₉NO₁₁ 659.772

Alkaloid from *Aconitum delavayi*, *Aconitum vilmorrianum*, *Aconitum crassicaule*, *Aconitum forrestii*, *Aconitum hemsleyanum*, *Aconitum teipecum*, *Aconitum pseudogeniculatum* and *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Mp 141-144° (98-103°). [α]_D²¹ +20 (c, 0.9 in CHCl₃). Identity with 3,13-Dihydroxyforesaconitine appears probable despite its lower Mp.

► AR5566500

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-methoxybenzoyl), 8-acyl: **Lipoyunaconitine**

Alkaloid from root tubers of *Aconitum ferox* (Ranunculaceae). Oil. [α]_D²⁵ +12 (c, 0.5 in CHCl₃). The esterifying group on C-8 is a mixt. of long-chain acyl groups (linoleoyl, palmitoyl and stearoyl).

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(4-methoxybenzoyl), 3,8-di-*Ac*: **3-Acetylyunaconitine**
[80787-51-5]

C₃₇H₅₁NO₁₂ 701.809

Alkaloid from *Aconitum* sp.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(3,4-dimethoxybenzoyl): **Veratrolylpseudoaconitine**
[53905-17-2]

C₃₄H₄₉NO₁₁ 647.761

Alkaloid from the roots of *Aconitum ferox* and *Aconitum falconeri* (Ranunculaceae). Prisms (MeOH/Et₂O). Mp 212-214°. [α]_D²⁰ +38.5 (c, 0.9 in EtOH). [α]_D²⁰ +54.1 (c, 1.5 in CHCl₃).

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(3,4-dimethoxybenzoyl), 8-*Ac*: **Pseudoaconitine**. *Pseudoaconitine*
[127-29-7]

C₃₆H₅₁NO₁₂ 689.798

Alkaloid from the roots of *Aconitum ferox*, *Aconitum falconeri*, *Aconitum balfourii*, *Aconitum deinorrhizum* and *Aconitum spicatum* (Ranunculaceae). Prisms (MeOH). Mp 211-213° (203-205°). [α]_D²⁰ +24 (c, 2.5 in CHCl₃).

► Highly toxic.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(3,4-dimethoxybenzoyl), 8-acyl: **Lipopsudaconitine**

Alkaloid from root tubers of *Aconitum ferox* (Ranunculaceae). Oil. [α]_D²⁵ +12.4 (c, 0.5 in CHCl₃). The esterifying group on C-8 is a mixt. of long-chain acyl groups (linoleoyl, palmitoyl and stearoyl).


O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-O-(3,4-dimethoxybenzoyl), 3,8,13-tri-*Ac*: **Diacetylpseudoaconitine**
[53905-18-3]

- C₄₀H₅₅NO₁₄ 773.873
Alkaloid from the roots of *Aconitum ferox* (Ranunculaceae). Needles (Me₂CO/hexane). Mp 230° (227-229°). [α]_D²⁵ +17 (c, 2.8 in CHCl₃).
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-O-(4-methoxybenzoyl): **Crassicautine** [109333-61-1]
C₃₄H₄₉NO₁₀ 631.762
Alkaloid from the roots of *Aconitum crassicaule* (Ranunculaceae). Amorph. [α]_D²⁴ +26.1 (c, 0.18 in CHCl₃).
- O¹, O⁶, O⁸, O¹⁶, O¹⁸-Penta-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **8-O-Methylveratroylpseudoaconine** [141695-34-3]
C₃₅H₅₁NO₁₁ 661.788
Alkaloid from *Aconitum balfourii* (Ranunculaceae). Amorph.
- O⁸-Et, O¹, O⁶, O¹⁶, O¹⁸-tetra-Me, N-Et, 14-O-(4-methoxybenzoyl): **Acoforestinine** [110011-77-3]
C₃₅H₅₁NO₁₀ 645.789
Alkaloid from the roots of *Aconitum forrestii* (Ranunculaceae). Amorph. [α]_D²⁵ +27.3 (c, 0.71 in EtOH).
- O⁸-Et, O¹, O⁶, O¹⁶, O¹⁸-tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **3'-Methoxyacoforestinine** [307321-72-8]
C₃₆H₅₃NO₁₁ 675.815
Alkaloid from *Aconitum falconeri*. Amorph. solid. [α]_D²⁵ +37 (c, 0.03 in MeOH). λ_{max} 207 (log ε 5.2); 218 (log ε 5.2); 260 (log ε 4.91) (MeOH).
- Dunstan, W.R. et al., *J.C.S.*, 1905, **87**, 1620-1636 (*Indaconitine, isol*)
- Henry, T.A. et al., *J.C.S.*, 1928, 1105-1121 (*Pseudoaconitine, isol*)
- Gilman, R.E. et al., *Tet. Lett.*, 1962, 923-925 (*Indaconitine, Pseudoaconitine, struct*)
- Tsuda, Y. et al., *Can. J. Chem.*, 1963, **41**, 1485-1489; 3055-3062 (*Pseudoaconitine, isol, struct, synth*)
- Miana, G.A. et al., *Phytochemistry*, 1971, **10**, 3320-3322 (*Indaconitine, isol, ir, pmr*)
- Birnbaum, K.B. et al., *Tet. Lett.*, 1971, 867-870 (*Ludaconitine*)
- Klásek, A. et al., *J. Nat. Prod.*, 1972, **35**, 55-66 (*Pseudoaconitine, isol, uv, ir*)
- Purusothaman, K.K. et al., *Phytochemistry*, 1974, **13**, 1975-1977 (*Veratroylpseudoaconine, Diacetylpseudoaconine*)
- Pelletier, S.W. et al., *Heterocycles*, 1977, **7**, 327-339 (*cmr*)
- Pelletier, S.W. et al., *Phytochemistry*, 1977, **16**, 623-624 (*Pseudoaconitine, Veratroylpseudoaconine*)
- Chen, S.-Y. et al., *Huaxue Xuebao*, 1979, **37**, 15-20; *CA*, **91**, 20833f (*Yunaconitine*)
- Yang, C. et al., *Huaxue Xuebao*, 1981, **39**, 445-452; *CA*, **96**, 82689a (*3-Acetylyunaconitine*)
- Wang, F. et al., *Planta Med.*, 1981, **42**, 375-379 (*Yunaconitine*)
- Zhang, H. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1982, **24**, 259-263; *CA*, **97**, 159517y (*Yunaconitine*)
- Chen, D. et al., *Zhongcaoyao*, 1982, **13**, 8-12; *CA*, **97**, 88655d
- Wang, C. et al., *Planta Med.*, 1983, **48**, 55 (*Yunaconitine, occur*)
- Pelletier, S.W. et al., *J. Nat. Prod.*, 1984, **47**, 474-477 (*Yunaconitine*)
- Chen, S. et al., *Yunnan Zhiwu Yanjiu*, 1984, **6**, 338-340; *CA*, **102**, 3240p (*8-Deacetylyunaconitine*)

- Chen, D. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1984, **26**, 82-84; *CA*, **101**, 51713w (*Yunaconitine*)
- Chen, W. et al., *Annalen*, 1985, 1297-1301 (*Dihydroxyforesaconitine*)
- Pelletier, S.W. et al., *Heterocycles*, 1987, **25**, 365-376 (*Acoforestinine*)
- Wang, F.-P. et al., *J. Nat. Prod.*, 1987, **50**, 55-62 (*Crassicautine*)
- Huiling, L. et al., *Heterocycles*, 1989, **29**, 2317-2326 (*Yunaconitine, 8-Deacetylyunaconitine*)
- Khetwal, K.S. et al., *Heterocycles*, 1992, **34**, 441-444 (*8-O-Methylveratroylpseudoaconine*)
- Hanuman, J.B. et al., *J. Nat. Prod.*, 1994, **57**, 105 (*Lipoindaconitine, Lipoyunaconitine, Lipopsuedaconitine*)
- Hanuman, J.B. et al., *Phytochemistry*, 1994, **36**, 1527-1535 (*Pseudoaconine*)
- Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000, **63**, 1393-1395 (*Faleoconitine, 3'-Methoxyacoforestinine*)
- Zhou, X.-L. et al., *Chem. Pharm. Bull.*, 2003, **51**, 592-594 (*Pseudoaconine, cmr*)
- Li, Z.Y. et al., *Helv. Chim. Acta*, 2004, **87**, 2085-2087 (*Yunaconitine, cmr*)
- Yan, H. et al., *Helv. Chim. Acta*, 2007, **90**, 1133-1140 (*Leucanthumsine E*)

4-Methylnaconitane-1,3,6,8,14,15,16,18-octol

M-337

C₁₉H₂₉NO₈ 399.44

(1α,3α,6α,14α,15α,16β)-form


- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl: **Acofamine A**
C₃₂H₄₅NO₉ 587.709
Alkaloid from the aerial parts of *Aconitum karakolicum*. Amorph. [α]_D²⁰ +21 (c, 0.06 in CHCl₃). λ_{max} 201 (log ε 4.13); 230 (log ε 4.18); 273 (log ε 2.94) (MeOH).

(1α,3α,6α,14α,15β,16β)-form

- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et: **Crassicaulidine**
[83704-07-8]
C₂₄H₃₉NO₈ 469.574
Alkaloid from the roots of *Aconitum crassicaule*. Mp 206-209°.
- Wang, F. et al., *Planta Med.*, 1983, **47**, 39-42 (*Crassicaulidine*)
- Atta-ur-Rahman, et al., *Nat. Prod. Res.*, 2005, **19**, 713-718 (*Acofamine A*)

4-Methylnaconitane-1,3,7,8,13,14,16,18-octol

M-338


C₁₉H₂₉NO₈ 399.44

(1α,3β,5β,14α,16β)-form

- O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-O-(4-methoxybenzoyl): **Hemsleyasine**
[189316-13-0]
C₃₂H₄₅NO₁₀ 603.708
Alkaloid from the roots of *Aconitum hemsleyanum* var. *circinatum*. Amorph. powder. [α]_D²⁵ +44.7 (c, 0.5 in CHCl₃).
- Xu, Q.Y. et al., *Chin. Chem. Lett.*, 1997, **8**, 137-140 (*isol, pmr, cmr*)

4-Methylnaconitane-1,6,7,8,10,14,16,18-octol

M-339


C₁₉H₂₉NO₈ 399.44

(1α,6β,14α,16β)-form

- O⁶, O¹⁶, O¹⁸-Tri-Me, N-Et: **Turcosine**
[168922-20-1]
C₂₄H₃₉NO₈ 469.574
Alkaloid from aerial parts of *Aconitum turczaninowi*. Cryst. (Me₂CO). Mp 206-208°.
- O¹, O⁶, O¹⁴, O¹⁶-Tetra-Me, N-Et: **Swatinine**
C₂₅H₄₁NO₈ 483.601
Alkaloid from the aerial parts of *Aconitum laeve*. Amorph. powder. [α]_D³⁰ +12.5 (c, 2 in CHCl₃).
- O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et: **Delcaroline**
[77996-00-0]
C₂₅H₄₁NO₈ 483.601
Major alkaloid from *Delphinium carolinianum* (Ranunculaceae). Mp 160-162° (as perchlorate). [α]_D^{23.5} +49.8 (c, 0.49 in MeOH).
- O⁶, O¹⁴, O¹⁶, O¹⁸-Tetra-Me, N-Et: **10-Hydroxydelsoiline. Turlosine**
[159394-69-1]
C₂₅H₄₁NO₈ 483.601
Alkaloid from above-ground parts of *Aconitum vulparia* ssp. *neapolitanum* (Ranunculaceae). Cryst. (hexane/EtOAc). Mp 240-243°. [α]_D²⁵ +51.4 (c, 0.14 in CHCl₃).
- O¹, O⁶, O¹⁴, O¹⁶, O¹⁸-Penta-Me, N-Et: **18-Methoxydelterine. 14-O-Methyl delcaroline**
C₂₆H₄₃NO₈ 497.628
Alkaloid from *Delphinium excelsum*. Amorph.
- Pelletier, S.W. et al., *Heterocycles*, 1981, **16**, 747 (*Delcaroline*)
- Pelletier, S.W. et al., *J.A.C.S.*, 1981, **103**, 6536-6538 (*config*)
- Batbayar, N. et al., *Khim. Prir. Soedin.*, 1992, **28**, 594; 1993, **29**, 60; *Chem. Nat. Compd.*

(Engl. Transl.), 1992, **28**, 521-522; 1993, **29**, 48 (*Turlosine, Turcosine*)
De la Fuente, G. et al., *Phytochemistry*, 1994, **37**, 271-274 (*10-Hydroxydelsoline*)
Batbayar, N. et al., *Phytochemistry*, 2003, **62**, 543-550 (*8-Methoxydelterine*)
Shaheen, F. et al., *Phytochemistry*, 2005, **66**, 935-940 (*Swatinine*)

4-Methylnaconitane-1,6,7,8,12,14,16,18-octol




$C_{19}H_{29}NO_8$ 399.44

(1 α ,5 β ,6 β ,12 α ,14 α ,16 β)-form

O^6, O^{16}, O^{18} -Tri-Me, N-Et: **Leucostine B**.
12-Hydroxydelcosine
[173075-40-6]
 $C_{24}H_{39}NO_8$ 469.574
Alkaloid from roots of *Aconitum leucostomum*. Needles (MeOH). Mp 263-264°.

Yue, J.M. et al., *J. Nat. Prod.*, 1996, **59**, 277-279

4-Methylnaconitane-1,6,7,8,14,15,16,18-octol



$C_{19}H_{29}NO_8$ 399.44


(1 α ,5 β ,6 β ,14 α ,15 β ,16 β)-form

$O^1, O^6, O^8, O^{14}, O^{16}$ -Penta-Me, N-formyl:
Potansine C
[182262-34-6]
 $C_{25}H_{39}NO_9$ 497.584
Alkaloid from roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D^{25} +45.5$ (c, 1.10 in $CHCl_3$).

$O^1, O^6, O^8, O^{14}, O^{16}$ -Penta-Me, N-formyl,
18-O-(2-aminobenzoyl): **Potansine D**
[182262-35-7]
 $C_{32}H_{44}N_2O_{10}$ 616.707
From roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D^{25} +32.1$ (c, 0.70 in $CHCl_3$).

Pu, H.-Y. et al., *Phytochemistry*, 1996, **43**, 287 (isol, pmr, cmr, struct)

4-Methylnaconitane-1,6,8,12,13,14,16,18-octol




$C_{19}H_{29}NO_8$ 399.44

(1 α ,5 β ,6 α ,12 β ,14 α ,16 β)-form

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Duclouxine**. *12-Hydroxy-chasmaconitine*
[96681-55-9]
 $C_{34}H_{47}NO_{10}$ 629.746
Alkaloid from roots of *Aconitum duclauxii* (Ranunculaceae). Mp 168-169°.

Wang, C. et al., *Yaoxue Xuebao*, 1984, **19**, 445; *CA*, **102**, 218381w

4-Methylnaconitane-1,6,8,13,14,15,16,18-octol



$C_{19}H_{29}NO_8$ 399.44

▶ AR5565000

(1 α ,6 α ,14 α ,15 α ,16 β)-form

O^6, O^{16}, O^{18} -Tri-Me, 14-benzoyl, 8-Ac:
Spicatine B. *10-Deoxyflavaconitine*. *10-Dehydroxyflavaconitine*
[863887-61-0]
 $C_{31}H_{41}NO_{10}$ 587.666
Alkaloid from the roots of *Aconitum spicatum* and *Aconitum nagarum*. Needles. Mp 130-132° (126-127°). $[\alpha]_D^{27} +39$ (c, 1 in $CHCl_3$). $[\alpha]_D^{25} +48$ (c, 0.1 in $CHCl_3$).

O^6, O^{16}, O^{18} -Tri-Me, 14-benzoyl, 8,N-di-Ac: **Flavaconijine**
[125263-92-5]
 $C_{33}H_{43}NO_{11}$ 629.703
Alkaloid from *Aconitum flavum* (Ranunculaceae). Mp 248.5-249.5°. $[\alpha]_D -55.5$ (c, 0.492 in DMF).

O^6, O^{16}, O^{18} -N-Tetra-Me, 14-benzoyl, 8-Ac: **1-Demethylhyaconitine**
[110081-95-3]
 $C_{32}H_{43}NO_{10}$ 601.692
Alkaloid from the roots of *Aconitum flavum* (Ranunculaceae). Mp 180-182°. $[\alpha]_D +19.9$ (c, 0.38 in $CHCl_3$).

O^1, O^6, O^{16}, O^{18} -Tetra-Me, 14-benzoyl, 8-Ac, N-formyl: **Aldohyaconitine**
[159602-66-1]

$C_{33}H_{43}NO_{11}$ 629.703

Alkaloid from O-Zhang-Ye fuzi (cultivated *Aconitum carmichaeli*) (Ranunculaceae). Cryst. Mp 262-264°. $[\alpha]_D^{20} -56.9$ (c, 0.3 in $CHCl_3$).

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 8-O-(9Z,12Z-octadecadienyl): **3-Deoxyaconitine 8-linoleate**
[1005031-38-8]

$C_{43}H_{71}NO_9$ 746.035
Alkaloid from the roots of *Aconitum flavum*.

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-benzoyl: **Benzoyldeoxyaconitine**
[78715-16-9]

$C_{32}H_{45}NO_9$ 587.709
Alkaloid from the roots of *Aconitum polyschistum* (Ranunculaceae). Mp 232-234°. $[\alpha]_D -15.4$ ($CHCl_3$).

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-benzoyl, 8-Ac: **Deoxyaconitine**
[3175-95-9]

$C_{34}H_{47}NO_{10}$ 629.746
Alkaloid from numerous *Aconitum* spp. (Ranunculaceae); contaminant in crude aconitine. Prisms (Et_2O). Mp 177-180°. $[\alpha]_D^{25} +15.5$ (c, 1.4 in EtOH).

▶ AR5569100

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-benzoyl, 8-acyl: **Lipodeoxyaconitine**
Lipoalkaloid from *Aconitum carmichaeli* (Ranunculaceae). Oil. $[\alpha]_D^{13} +12.4$ ($CHCl_3$). Analogue of Deoxyaconitine with a mixt. of long-chain acyl groups at C-8 in place of Ac.

O^1, O^6, O^{16}, O^{18} -Tetra-Me, N-Et, 14-(4-methoxybenzoyl), 8-Ac: **Deoxyjesaconitine**
[69787-27-5]

$C_{35}H_{49}NO_{11}$ 659.772
Alkaloid from the rhizomes of *Aconitum subcuneatum* (Ranunculaceae). Needles (Me_2CO). Mp 174-176°. $[\alpha]_D +52$ (c, 0.065 in MeOH).

▶ AR5569130

O^1, O^6, O^{16}, O^{18} -N-Penta-Me: **Hyaconine**
[63238-68-6]
 $C_{24}H_{39}NO_8$ 469.574
Alkaloid from *Aconitum* spp.

O^1, O^6, O^{16}, O^{18} -N-Penta-Me, 14-benzoyl: **Benzoylhyaconine**. *Deacetylhyaconitine*
[63238-66-4]
 $C_{31}H_{43}NO_9$ 573.682

Alkaloid from the roots of *Aconitum carmichaeli* (Ranunculaceae). Also isol. from the processed aconite "Kako-bushi-matsu", produced by autoclaving the crude drug ("bushi") obt. from the roots of some *Aconitum* spp.

▶ AR5569370

O^1, O^6, O^{16}, O^{18} -N-Penta-Me, 14-benzoyl, 8-Ac: **Hyaconitine**. *Deoxymesaconitine*. *Japaconitine B₁*. *Japaconitine C*. *Japaconitine C₁*
[6900-87-4]

$C_{33}H_{45}NO_{10}$ 615.719
Alkaloid from *Aconitum senanense*, *Aconitum carmichaeli*, *Aconitum koreanum*, *Aconitum bullatifolium* var. *homotrichum*, *Aconitum callianthum*,

Aconitum ibukiense, *Aconitum tortuosum*, *Aconitum hakusanense* and many other *Aconitum* spp. (Ranunculaceae). Prisms (MeOH). Mp 197-198° dec. $[\alpha]_D^{17} +22.4$ (CHCl₃).

O¹,O⁶,O¹⁶,O¹⁸,N-Penta-Me, 14-benzoyl, 8-Ac; hydrobromide: Cryst. + 2.5H₂O (EtOH/Et₂O). Mp 178-179° dec. $[\alpha]_D -19$ (H₂O).

O¹,O⁶,O¹⁶,O¹⁸,N-Penta-Me, 14-benzoyl, 8-acyl: **Lipohypaconitine** [81941-15-3]

From *Aconitum carmichaeli* (Ranunculaceae). Oil. $[\alpha]_D^{22} +13.5$ (CHCl₃). Analogue of Hypaconitine with a mixt. of long-chain acyl groups at C-8 in place of Ac.

O¹,O⁶,O⁸,O¹⁶,O¹⁸,N-Penta-Me, N-Et, 14-benzoyl: **Neojiangyouaconitine** [126266-35-1]

C₃₃H₄₇NO₉ 601.736
Alkaloid from *Aconitum carmichaeli*. Cryst. Mp 201-204°. $[\alpha]_D^{14} -9.5$ (c, 0.22 in MeOH).

O¹,O⁶,O⁸,O¹⁶,O¹⁸,N-Hexa-Me: **8-O-Methylhypaconine** [949118-97-2]

C₂₅H₄₁NO₈ 483.601
Alkaloid from an *Aconitum* sp. Needles (CHCl₃/MeOH). Mp 225-228°. $[\alpha]_D^{23} +4.4$ (c, 0.3 in MeOH).

(1β,6α,14α,15α,16β)-form

O¹,O⁶,O¹⁶,O¹⁸,N-Penta-Me, 14-benzoyl, 8-Ac: **Ipaconitine** [111612-41-0]

C₃₃H₄₅NO₁₀ 615.719
Alkaloid from *Aconitum napellus* (Ranunculaceae).

Majima, R. *et al.*, *Annalen*, 1929, **476**, 171-181 (*Hypaconitine*, *isol*)

Tsuda, Y. *et al.*, *Annalen*, 1964, **680**, 88-92 (*Hypaconitine*, *Deoxyhypaconitine*)

Birnbaum, K.B. *et al.*, *Tet. Lett.*, 1971, 867-870 (*config*)

Pelletier, S.W. *et al.*, *Can. J. Chem.*, 1979, **57**, 1652-1655 (*cmr*)

Kitagawa, I. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 758-761 (*Lipodeoxyhypaconitine*, *Lipohypaconitine*, *isol*)

Mori, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2884-2886 (*Deoxyhypaconitine*)

Hikino, H. *et al.*, *Shoyakugaku Zasshi*, 1983, **37**, 68-72; *CA*, **99**, 110568t (*Benzoylhypaconitine*)

Chen, Z., *et al.*, *Heterocycles*, 1987, **26**, 1455-1460; 1989, **29**, 997-1002 (*1-Demethylhypaconitine*, *Flavaconifine*)

Wang, H. *et al.*, *Heterocycles*, 1988, **27**, 1615-1621 (*Benzoyldeoxyhypaconitine*)

Mori, T. *et al.*, *Heterocycles*, 1989, **29**, 873-885 (*Benzoylhypaconitine*)

Zhang, W.D. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 670-673 (*Neojiangyouaconitine*)

Wang, X.K. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 671-672 (*Aldohypaconitine*)

Sun, W. *et al.*, *Planta Med.*, 1999, **65**, 432-436 (*Lipodeoxyhypaconitine*, *Lipohypaconitine*, *ms*)

Gao, L.M. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 475-478 (*Spicataine B*)


Zhang, F. *et al.*, *Chin. Chem. Lett.*, 2005, **16**, 1043-1046 (*10-Deoxyflavaconitine*)

Zhang, F. *et al.*, *Zhongguo Yaoxue Zazhi*, 2006, **41**, 1851-1854; *CA*, **148**, 210217b (*3-Deoxyhypaconitine 8-linoleate*)

Lee, S.Y. *et al.*, *Arch. Pharmacol. Res.*, 2007, **30**, 691-694 (*8-O-Methylhypaconitine*)

4-Methyloaconitane-1,7,8,9,10,14,16,18-octol

M-344

C₁₉H₂₉NO₅ 351.442C₁₉H₂₉NO₈ 399.44

(1α,5β,14α,16β)-form

7,8-Methylene ether, O¹⁴,O¹⁶,O¹⁸-tri-Me, N-Et, 1-Ac: **Talidine C** [110081-94-2]

C₂₇H₄₁NO₉ 523.622
Alkaloid from the roots of *Delphinium taliense* (Ranunculaceae).

7,8-Methylene ether, O¹,O¹⁴,O¹⁶,O¹⁸-tetra-Me, N-Et: **Talidine A** [110120-83-7]

C₂₆H₄₁NO₈ 495.612
Alkaloid from *Delphinium taliense* (Ranunculaceae).

Chen, S. *et al.*, *Yunnan Zhiwu Yanjiu*, 1986, **8**, 81; 1987, **9**, 365; *CA*, **107**, 112626t; **108**, 147146j (*isol*, *ir*, *pmr*, *struct*)


(1α,3α,5β,14α,16β)-form

O⁸,O¹⁶-Di-Me, N-Et: **Crispulidine** C₂₃H₃₇NO₅ 407.549
Alkaloid from *Delphinium crispulum*. Racemic.

Ulubelen, A. *et al.*, *Phytochemistry*, 1999, **50**, 513-516 (*isol*, *ir*, *pmr*, *cmr*, *ms*)

4-Methyloaconitane-1,6,8,14,16-pentol

M-347



(1α,5β,6α,14α,16β)-form

C₁₉H₂₉NO₅ 351.442

C-6 Configs. for alkaloids covered by this entry follows from the revised struct. of Peregrine based X-ray anal. of Peregrine alcohol (1994).

(1α,5β,6α,14α,16β)-form

O¹,O⁶,O¹⁶-Tri-Me, N-Et: **Royleinine** C₂₄H₃₉NO₅ 421.576
Alkaloid from *Delphinium roylei*. Amorph. powder.

(1α,5β,6β,14α,16β)-form

O¹⁶-Me, N-Et: **Bicolorine†** [41478-10-8]
C₂₂H₃₅NO₅ 393.522
Minor alkaloid from *Delphinium bicolor* (Ranunculaceae). Hydrol. prod. of Bicoloridine. Mp 190-191°. $[\alpha]_D^{24} +16$ (CHCl₃).

O¹⁶-Me, N-Et, 6-Ac: **Bicolorine 6-O-acetate**

C₂₄H₃₇NO₆ 435.559
Alkaloid from *Delphinium bicolor* (Ranunculaceae). Cryst. (CH₂Cl₂/hexane). Mp 165-167°. $[\alpha]_D +19$ (c, 0.2 in CHCl₃).

O¹⁶-Me, N-Et, 14-Ac: **Bicolorine 14-O-acetate**

[130774-04-8]
C₂₄H₃₇NO₆ 435.559
Alkaloid from *Delphinium nuttallianum* (Ranunculaceae).

O¹⁶-Me, N-Et, 1,8-di-Ac: **Souline A**


[244246-51-3]
C₂₆H₃₉NO₇ 477.597
Alkaloid from *Delphinium souliei*. Cryst.

O¹,O¹⁶-Di-Me, N-Et: **Regaline. Munzianine**

[119459-58-4]
C₂₃H₃₇NO₅ 407.549
Alkaloid from epigeal parts of *Consolida regalis* ssp. *paniculata* and *Delphinium munzianum* (Ranunculaceae). Cryst. (hexane/EtOAc). Mp 154-157°. $[\alpha]_D +3.8$ (c, 0.2 in CHCl₃). The CA abstract diag. omits the 14-OH group. Regaline and Munzianine not compared but same struct. assigned. Physical data refer to Munzianine.

4-Methyloaconitane-6,8,10,13,14,15,16,18-octol

M-345

C₁₉H₂₉NO₈ 399.44

(5β,6α,14α,15α,16β)-form


O⁶,O¹⁶,O¹⁸-Tri-Me, 14-benzoyl, 8-Ac: **Polyschistine C** [96562-90-2]

C₃₁H₄₁NO₁₀ 587.666
Alkaloid from the roots of *Aconitum polyschistum* (Ranunculaceae). Amorph. solid.

Wang, H. *et al.*, *Heterocycles*, 1985, **23**, 803 (*pmr*, *cmr*, *struct*)

4-Methyloaconitane-1,3,8,14,16-pentol

M-346




- O^6, O^{16} -Di-Me, N-Et: **Dihydropentagynine**
[84306-88-7]
 $C_{23}H_{37}NO_5$ 407.549
Alkaloid from *Delphinium pentagynum* (Ranunculaceae). Mp 150-154°. $[\alpha]_D^{25} +43$ (c, 0.12 in EtOH).
- O^8, O^{16} -Di-Me, N-Et: **Bicoloridine alcohol**. 6-Deacetylbicoloridine
 $C_{23}H_{37}NO_5$ 407.549
Alkaloid from epigeal parts of *Delphinium peregrinum* var. *elongatum* (Ranunculaceae). Resin. $[\alpha]_D -3.3$ (c, 0.24 in $CHCl_3$).
- O^8, O^{16} -Di-Me, N-Et, 6-Ac: **Bicoloridine**
[41710-20-7]
 $C_{25}H_{39}NO_6$ 449.586
Minor alkaloid from *Delphinium bicolor* (Ranunculaceae). Also isol. from the aerial parts of *Delphinium peregrinum* var. *elongatum* collected during flowering. Noncryst. Mp 240° (as hydroiodide). $[\alpha]_D^{25} +10$ ($CHCl_3$).
- O^1, O^8, O^{16} -Tri-Me, N-Et: **Peregrine alcohol**. 6-O-Deacetylperegrine
[113689-39-7]
 $C_{24}H_{39}NO_5$ 421.576
Alkaloid from aerial parts of *Delphinium peregrinum*, *Delphinium davidii* and *Delphinium gueneri* (Ranunculaceae). $[\alpha]_D^{25} +14$ (c, 0.12 in MeOH).
- O^1, O^8, O^{16} -Tri-Me, N-Et, 6-Ac: **Peregrine**
[113689-38-6]
 $C_{26}H_{41}NO_6$ 463.613
Alkaloid from aerial parts of *Delphinium peregrinum* var. *elongatum* (Ranunculaceae). Cryst. (EtOAc/hexane). Mp 124-125°. $[\alpha]_D +12$ (c, 0.34 in EtOH).
- O^1, O^8, O^{16} -Tri-Me, N-Et, 6,14-di-Ac: **14-O-Acetylperegrine**
 $C_{28}H_{43}NO_7$ 505.65
Alkaloid from aerial parts of *Delphinium munzianum* and from *Delphinium davisii* (Ranunculaceae). Resin. $[\alpha]_D -49.6$ (c, 0.24 in $CHCl_3$).
- O^1, O^8, O^{16} -Tri-Me, N-Et, 14-benzoyl, 6-Ac: **14-O-Benzoylperegrine**
 $C_{33}H_{45}NO_7$ 567.721
Alkaloid from aerial parts of *Delphinium munzianum* (Ranunculaceae). Cryst. (hexane/EtOAc). Mp 112-114°. $[\alpha]_D -21$ (c, 0.1 in $CHCl_3$).
- O^1, O^8, O^{14}, O^{16} -Tetra-Me, 6-Ac: N-**Deethyl-14-O-methylperegrine**
 $C_{25}H_{39}NO_6$ 449.586
Alkaloid from aerial parts of *Delphinium gueneri* (Ranunculaceae). $[\alpha]_D^{22} -38.4$ (c, 0.1 in MeOH).
- O^1, O^8, O^{14}, O^{16} -Tetra-Me, N-Et, 6-Ac: **14-O-Methylperegrine**
 $C_{27}H_{43}NO_6$ 477.64
Alkaloid from aerial parts of *Delphinium gueneri* (Ranunculaceae). $[\alpha]_D^{22} -28.9$ (c, 0.2 in MeOH).
- 6-Ketone, O^1, O^8, O^{16} -tri-Me, N-Et: **Munzianone**
[167782-16-3]
 $C_{24}H_{37}NO_5$ 419.56
Alkaloid from aerial parts of *Delphinium munzianum* (Ranunculaceae). Cryst. (hexane/EtOAc). Mp 117-118°.

$[\alpha]_D -25$ (c, 0.17 in $CHCl_3$).

- Jones, A.J. *et al.*, *Can. J. Chem.*, 1973, **51**, 486 (*Bicolorine, isol*)
Pelletier, S.W. *et al.*, *Tet. Lett.*, 1976, 3025 (*Bicolorine, cmr, struct*)
Coddling, P.W. *et al.*, *Acta Cryst. B*, 1981, **37**, 379 (*Bicoloridine, cryst struct*)
González, A.G. *et al.*, *Phytochemistry*, 1982, **21**, 1781 (*Dihydropentagynine*)
Kulanthaivel, P. *et al.*, *Phytochemistry*, 1986, **25**, 1511 (*Bicolorine 6-O-acetate*)
Sener, B. *et al.*, *CA*, 1988, **110**, 111674g (*Regaline*)
de la Fuente, G. *et al.*, *Heterocycles*, 1988, **27**, 1 (*Bicoloridine, Peregrine*)
Bai, Y. *et al.*, *Heterocycles*, 1990, **31**, 1233 (*Bicolorine 14-O-acetate*)
Ulubelen, A. *et al.*, *Phytochemistry*, 1992, **31**, 1019; 1993, **33**, 213 (*Peregrine alcohol, 14-O-Methylperegrine, N-Deethyl-14-O-methylperegrine*)
De la Fuente, G. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1768 (*Peregrine, struct*)
De la Fuente, G. *et al.*, *Phytochemistry*, 1995, **39**, 1459-1465; 1467-1473 (*Bicoloridine alcohol, Munzianone, Munzianone, 14-O-Acetylperegrine, 14-O-Benzoylperegrine*)
Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 300 (*14-O-Acetylperegrine, 6-O-Deacetylperegrine*)
Pan, X. *et al.*, *Chin. Chem. Lett.*, 1998, **9**, 57-59 (*Souline A*)
Ulubelen, A. *et al.*, *Heterocycles*, 2000, **53**, 2279-2282 (*Royleinine*)

4-Methylnaconitane-1,7,8,14,16-pentol M-348




$C_{19}H_{29}NO_5$ 351.442

(1 α ,5 β ,14 α ,16 β)-form

- O^{16} -Me, N-Et: **Leroyine**
[145249-70-3]
 $C_{22}H_{35}NO_5$ 393.522
Alkaloid from *Delphinium leroyi* (Ranunculaceae). Amorph. solid.
- O^{16} -Me, N-Et, 14-Ac: **Leroyine 14-acetate**
[145204-93-9]
 $C_{24}H_{37}NO_6$ 435.559
Alkaloid from whole plants of *Delphinium leroyi* (Ranunculaceae). Amorph. solid.
- O^8, O^{16} -Di-Me, N-Et: **Blacknidine**
[164176-13-0]
 $C_{23}H_{37}NO_5$ 407.549
Alkaloid from whole plants of *Delphinium elatum* var. "black night" (Ranunculaceae). Amorph. solid. $[\alpha]_D +8.05$ (c, 0.642 in $CHCl_3$).
- 7,8-Methylene ether, O^1, O^{14}, O^{16} -tri-Me, N-Et: **Occidentaline**. 6-Deoxydelpheline
[113707-10-1]
 $C_{25}H_{39}NO_5$ 433.587
Alkaloid from *Delphinium barbeyi* and *Delphinium occidentale* (Ranunculaceae).

- ceae). Mp 119.5-120.5° (112-115°).
Kulanthaivel, P. *et al.*, *Heterocycles*, 1988, **27**, 339-342 (*Occidentaline*)
Joshi, B.S. *et al.*, *Tet. Lett.*, 1988, **29**, 2397-2400 (*6-Deoxydelpheline*)
Bai, Y. *et al.*, *Phytochemistry*, 1992, **31**, 3243-3245 (*Leroyine, Leroyine 14-acetate*)
Park, J.C. *et al.*, *J. Nat. Prod.*, 1995, **58**, 291 (*Blacknidine*)

4-Methylnaconitane-1,8,10,14,16-pentol M-349




$C_{19}H_{29}NO_5$ 351.442

(1 α ,14 α ,16 β)-form

- O^{16} -Me, N-Et: **Karakolidine**. Karakolidine. 10 β -Hydroxykarakoline
[41655-13-4]
 $C_{22}H_{35}NO_5$ 393.522
Alkaloid from tubers of *Aconitum karakolicum* (Ranunculaceae). Cryst. (MeOH). Mp 222-224°.
- O^1, O^{16} -Di-Me, N-Et: **Genicinine B**
[287186-10-1]
 $C_{23}H_{37}NO_5$ 407.549
Alkaloid from *Aconitum geniculatum* and *Aconitum variegatum*. $[\alpha]_D^{24} +3.4$ (c, 0.6 in $CHCl_3$).
- O^1, O^{16} -Di-Me, N-Et, O^{14} -Ac: **14-O-Acetylgenicinine B**
[852381-85-2]
 $C_{25}H_{39}NO_6$ 449.586
Alkaloid from *Aconitum variegatum*. Amorph. solid. $[\alpha]_D^{20} +24.2$ (c, 0.55 in $CHCl_3$).
- Sultankhodzhaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 481; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 498 (*Karakolidine*)
Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 837-846 (*Genicinine B, 14-O-Acetylgenicinine B*)

4-Methylnaconitane-1,8,12,14,16-pentol M-350



(1 α ,12 β ,14 α ,16 β)-form

$C_{19}H_{29}NO_5$ 351.442

(1 α ,12 β ,14 α ,16 β)-form

- O^{14}, O^{16} -Di-Me, N-Et: **12 β -Hydroxykarasamine**
[161993-26-6]
 $C_{23}H_{37}NO_5$ 407.549
Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Amorph. solid.

O¹⁴, O¹⁶-Di-Me, N-Et, O⁸-Ac: **12β-Hydroxykarasamine 8-O-acetate**
[161993-23-3]
C₂₅H₃₉NO₆ 449.586
Minor alkaloid from *Delphinium nuttalianum*. Amorph. solid.


(1β,12β,14α,16β)-form

O¹⁴, O¹⁶-Di-Me, N-Et: **1-Epi-12β-hydroxykarasamine**
[161993-24-4]
C₂₃H₃₇NO₅ 407.549
Minor alkaloid from *Delphinium nuttalianum* (Ranunculaceae). Amorph. solid.

Bai, Y. et al., *Phytochemistry*, 1994, **37**, 1717-1724 (isol)

4-Methylnaconitane-1,8,13,14,16-pentol

M-351



C₁₉H₂₉NO₅ 351.442


(1α,5β,14α,16β)-form

O¹, O¹⁶-Di-Me, N-Et: **Conaconitine**
[128357-74-4]
C₂₃H₃₇NO₅ 407.549
Alkaloid from the roots of *Aconitum contortum* (Ranunculaceae).

Wang, S. et al., *Huaxue Xuebao*, 1989, **47**, 1101; *CA*, **113**, 55802

4-Methylnaconitane-1,8,14,16,18-pentol

M-352

**(1α,14α,16β)-form**

C₁₉H₂₉NO₅ 351.442

▶ AR5541000

(1α,14α,16β)-form

O¹⁶-Me, N-Et: **Columbianine**
[90579-23-0]
C₂₂H₃₅NO₅ 393.522
Minor alkaloid from the aerial parts of *Aconitum columbianum* (Ranunculaceae). Cryst. (MeOH). Mp 202-205°. [α]_D -6 (c, 0.5 in EtOH).

O¹⁶-Me, N-Et, 18-O-(2-aminobenzoyl): **Acoseptridine**
[146001-06-1]

C₂₉H₄₀N₂O₆ 512.645
Alkaloid from roots of *Aconitum septentrionale*. Amorph. solid. [α]_D +1.5 (c, 0.345 in CHCl₃).

O¹⁶-Me, N-Et, 8-O-(2-hydroxyphenyl), 18-O-[2-(methoxycarbonylamino)benzoyl]: **Linearilobine**
[914475-08-4]

C₃₇H₄₆N₂O₉ 662.778
Alkaloid from the roots of *Delphinium linearilobum*. Amorph. solid. λ_{max} 222 (log ε 4.5); 252 (log ε 3.6); 308 (log ε 2) (MeOH).

O¹, O¹⁶-Di-Me, N-Et: **Cammaconine**.

Aconoridine
[32152-70-8]
C₂₃H₃₇NO₅ 407.549

Alkaloid from the above-ground parts of *Aconitum variegatum*, roots of *Aconitum forrestii* and *Aconitum dolichorhynchum* var. *subglabratum* and the aerial parts of *Aconitum columbianum* (Ranunculaceae). Also isol from *Notopterygium incisum*. Rods (MeOH). Mp 136-138°. [α]_D 0 (c, 1.0 in CHCl₃). pK_a 9.55 (50% MeOH). Struct. revised in 1980.

O¹, O¹⁶-Di-Me, N-Et, hydroiodide: Mp 225-227°.

O¹, O¹⁶-Di-Me, N-Et, 18-Ac: **18-Acetylcammaconine**

C₂₅H₃₉NO₆ 449.586
Alkaloid from the roots of *Aconitum piepunense*. Amorph. powder. Mp 123-125°. [α]_D²⁰ -1.7 (c, 0.5 in CHCl₃).

O¹, O¹⁶-Di-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Geniculatine D**
[359798-76-8]

C₃₂H₄₅NO₈ 571.709
Alkaloid from *Aconitum geniculatum*. Amorph. solid. [α]_D +7.3 (c, 0.6 in CHCl₃).

O¹, O¹⁶-Di-Me, N-Et, 18-O-(2-aminobenzoyl): **Acobretine E**
[158734-29-3]

C₃₀H₄₂N₂O₆ 526.672
Alkaloid from roots of *Aconitum brevicelecaratum* (Ranunculaceae).

O¹, O¹⁶-Di-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Aconorine**
[58969-45-2]

C₃₂H₄₄N₂O₇ 568.709
Alkaloid from *Aconitum orientale* and *Aconitum columbianum* (Ranunculaceae). Amorph. Mp 237° (as perchlorate).

O⁸, O¹⁶-Di-Me, N-Et: **Raveyine**. 8-O-Methylcolumbianine

[199273-28-4]
C₂₃H₃₇NO₅ 407.549

Alkaloid from the aerial parts of *Consolida raveyi* and from *Consolida oliveriana* (Ranunculaceae). Amorph. solid. [α]_D -4.1 (c, 1 in CHCl₃). [α]_D²⁵ -2.9 (c, 0.1 in CHCl₃).

O¹⁶, O¹⁸-Di-Me, N-Et: **Isotalatizidine**

[7633-68-3]
C₂₃H₃₇NO₅ 407.549
Alkaloid from the roots of *Aconitum carmichaeli*, *Aconitum talassicum* and *Delphinium denudatum*, the aerial parts of *Aconitum columbianum*, and from

Aconitum japonicum (Ranunculaceae). Cryst. (Et₂O/hexane). Mp 116-117° Mp 140-142°.

O¹⁶, O¹⁸-Di-Me, N-Et, 14-Ac: **Condelpine**. 14-Acetylisotalatizidine

[7633-69-4]
C₂₅H₃₉NO₆ 449.586
Alkaloid from *Delphinium denudatum*, *Delphinium confusum*, *Delphinium oreophilum*, *Delphinium iliense*, *Delphinium poltoratzkii* and *Aconitum japonicum* (Ranunculaceae). Cryst. (Et₂O/MeOH). Mp 158-159°. [α]_D³⁰ +21.3 (CHCl₃).

O¹⁶, O¹⁸-Di-Me, N-Et, 8,14-di-Ac: **8-Acetylcondelpine**

[211358-75-7]
C₂₇H₄₁NO₇ 491.623
Alkaloid from *Delphinium pyramadale*. [α]_D -28.5 (c, 1.5 in CHCl₃).

O¹⁶, O¹⁸-Di-Me, N-Et, 8-O-(3,4-dimethoxybenzoyl), 14-Ac: **Geniculatine C**

[359798-74-6]
C₃₄H₄₇NO₉ 613.747
Alkaloid from *Aconitum geniculatum*. Amorph. solid. [α]_D +50 (c, 0.7 in CHCl₃).

O¹⁶, O¹⁸-Di-Me, N-Et, 14-O-(3,4-dimethoxycinnamoyl): **Gymnaconitine**
[103956-41-8]

C₃₄H₄₇NO₈ 597.747
Alkaloid from *Aconitum gymndrum* (Ranunculaceae). Cryst. Mp 110-111°. [α]_D¹⁷ +18.2.

O¹, O¹⁴, O¹⁶-Tri-Me, N-Et: **Scaconine**
[156310-13-3]

C₂₄H₃₉NO₅ 421.576
Alkaloid from *Aconitum scaposum* var. *vaginatatum* and *Aconitum brevicelecaratum*. Amorph.

O¹, O¹⁴, O¹⁶-Tri-Me, N-Et, 18-O-(2-aminobenzoyl): **N-Deacetylscaconitine**
[155653-05-7]

C₃₁H₄₄N₂O₆ 540.698
Alkaloid from *Aconitum scaposum* var. *vaginatatum* and *Aconitum brevicelecaratum*. Amorph.

O¹, O¹⁴, O¹⁶-Tri-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Scaconitine**
[156310-14-4]

C₃₃H₄₆N₂O₇ 582.736
Alkaloid from *Aconitum scaposum* var. *vaginatatum* and *Aconitum brevicelecaratum*. Amorph.

O¹, O¹⁶, O¹⁸-Tri-Me, N-Et: **Talatizamine**. *Talatisamine*

[20501-56-8]
C₂₄H₃₉NO₅ 421.576
Alkaloid from *Aconitum talassicum*, *Aconitum nemorum*, *Aconitum carmichaeli*, *Aconitum variegatum*, *Aconitum franchetii*, *Aconitum forrestii*, *Aconitum columbianum*, *Aconitum saposchnikovii*, *Aconitum pseudogeniculatum* and others. Mp 145-146° (142-143°). [α]_D 0 (c, 1.0 in EtOH).

O¹, O¹⁶, O¹⁸-Tri-Me, N-Et, 14-Ac: **14-Acetyltalatizamine**


[71239-55-9]
C₂₆H₄₁NO₆ 463.613
Alkaloid from *Aconitum japonicum*,

- the aerial parts of *Aconitum columbianum* and the roots of *Aconitum carmichaeli* (Ranunculaceae). Amorph. powder. $[\alpha]_D^{20} +19.7$ (c, 0.36 in CHCl_3).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 8-benzoyl, 14-Ac: **Transconitine A**. 14-Acetyl-8-benzoyltalatzimine
[175923-71-4]
 $\text{C}_{33}\text{H}_{45}\text{NO}_7$ 567.721
Alkaloid from roots of *Aconitum transectum* (Ranunculaceae). Amorph. $[\alpha]_D^{20} +16.9$ (c, 0.002 in CHCl_3).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 14-benzoyl: **14-Benzoyltalatzimine**
[11037-59-5]
 $\text{C}_{31}\text{H}_{43}\text{NO}_6$ 525.684
Alkaloid from roots of *Aconitum kongboense* (Ranunculaceae) and *Aconitum nemorum*. Cryst. (hexane/ Me_2CO). Mp 104-105°. Erroneously claimed to be novel by Sultankhodzhaev.
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 8-O-(4-methoxybenzoyl), 14-Ac: **Dolichotine A**. 14-Acetyl-8-anisoyltalatzimine
[128364-21-6]
 $\text{C}_{34}\text{H}_{47}\text{NO}_8$ 597.747
Minor alkaloid from roots of *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Amorph. $[\alpha]_D +15.2$ (CHCl_3).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 14-O-(4-methoxybenzoyl): **14-Anisoyltalatzimine**
[592528-66-0]
 $\text{C}_{32}\text{H}_{45}\text{NO}_7$ 555.71
Alkaloid from the roots of *Aconitum bulleyanum*. Amorph. $[\alpha]_D^{20} +12.5$ (c, 0.15 in CHCl_3).
- $\text{O}^1, \text{O}^{14}, \text{O}^{16}$ -Tri-Me, N-Et, 18-O-(2-aminobenzoyl), 8-Ac: **Acobretine A**
[156318-68-2]
[158734-30-6]
 $\text{C}_{33}\text{H}_{46}\text{N}_2\text{O}_7$ 582.736
Alkaloid from roots of *Aconitum brevicaratum* (Ranunculaceae).
- $\text{O}^1, \text{O}^{14}, \text{O}^{16}$ -Tri-Me, N-Et, 18-O-(2-acetamidobenzoyl), 8-Ac: **Acobretine D**
[155653-03-5]
 $\text{C}_{33}\text{H}_{48}\text{N}_2\text{O}_8$ 624.773
Alkaloid from roots of *Aconitum brevicaratum* (Ranunculaceae).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 14-O-(4-methoxybenzoyl), 8-Ac: **8-Acetyl-14-anisoyltalatzimine**
[592528-65-9]
 $\text{C}_{34}\text{H}_{47}\text{NO}_8$ 597.747
Alkaloid from the roots of *Aconitum bullayanum*. Amorph. $[\alpha]_D^{20} +14.3$ (c, 0.15 in CHCl_3).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 8-O-(3,4-dimethoxybenzoyl), 14-Ac: **Dolichotine B**. 14-Acetyl-8-veratroyltalatzimine. **Vilmorinine**
[128364-22-7]
 $\text{C}_{35}\text{H}_{49}\text{NO}_9$ 627.773
Minor alkaloid from roots of *Aconitum dolichorhynchum* var. *subglabratum* and of *Aconitum vilmorianum* (Ranunculaceae). Amorph. $[\alpha]_D$ 0 (CHCl_3). Dolichotine B and Vilmorinine appear identical. Props. refer to Dolichotine B.
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 14-O-E-cinnamoyl: 14-O-trans-Cinnamoyltalatzimine. **Leueantine C**
[499969-77-6]
 $\text{C}_{33}\text{H}_{45}\text{NO}_6$ 551.722
Alkaloid from the roots of *Aconitum hemisleyanum* var. *leueanthus*. Amorph. powder. $[\alpha]_D^{20} +34.6$ (c, 0.5 in CHCl_3).
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 8-O-E-cinnamoyl, 14-Ac: **Dolichotine C**. 14-Acetyl-8-trans-cinnamoyltalatzimine
[128364-23-8]
 $\text{C}_{35}\text{H}_{47}\text{NO}_7$ 593.759
Minor alkaloid from roots of *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Amorph.
- $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et, 14-O-(3,4-dimethoxycinnamoyl): **Methylgymnaconitine**
[103956-42-9]
 $\text{C}_{35}\text{H}_{49}\text{NO}_8$ 611.774
Alkaloid from *Aconitum gymndrum* (Ranunculaceae). Amorph. powder. $[\alpha]_D^{21} +33.2$.
- $\text{O}^8, \text{O}^{14}, \text{O}^{16}$ -Tri-Me, N-Et, 18-O-(2-methoxybenzoyl): **Cyphoplectine**
[244616-82-8]
 $\text{C}_{32}\text{H}_{45}\text{NO}_7$ 555.71
Alkaloid from *Delphinium cyphoplectrum*. Amorph. Racemic.
- $\text{O}^{14}, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et: **14-Methylsotalatzidine**
[114622-07-0]
 $\text{C}_{24}\text{H}_{39}\text{NO}_5$ 421.576
Alkaloid from aerial parts of *Delphinium confusum* (Ranunculaceae). Cryst. (Me_2CO). Mp 136-137°.
- $\text{O}^1, \text{O}^8, \text{O}^{16}, \text{O}^{18}$ -Tetra-Me, N-Et: **8-O-Methyltalatzimine**
[90579-28-5]
 $\text{C}_{25}\text{H}_{41}\text{NO}_5$ 435.603
Alkaloid from the aerial parts of *Aconitum columbianum* (Ranunculaceae). Amorph. $[\alpha]_D^{29.5} -4$ (c, 1.16 in CHCl_3).
- O^8 -Et, $\text{O}^1, \text{O}^{16}$ -di-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Talassicumine A**. 8-O-Ethylaconorine
[160824-50-0]
 $\text{C}_{34}\text{H}_{48}\text{N}_2\text{O}_7$ 596.762
Alkaloid from roots of *Aconitum talassicum* (Ranunculaceae). Amorph. powder.
- O^8 -Et, $\text{O}^1, \text{O}^{14}, \text{O}^{16}$ -tri-Me, N-Et, 18-O-(2-aminobenzoyl): **Acobretine B**
[156310-15-5]
 $\text{C}_{33}\text{H}_{48}\text{N}_2\text{O}_6$ 568.752
Alkaloid from roots of *Aconitum brevicaratum* (Ranunculaceae).
- O^8 -Et, $\text{O}^1, \text{O}^{14}, \text{O}^{16}$ -tri-Me, N-Et, 18-O-(2-acetamidobenzoyl): **Acobretine C**
[156310-16-6]
 $\text{C}_{35}\text{H}_{50}\text{N}_2\text{O}_7$ 610.789
Alkaloid from roots of *Aconitum brevicaratum* (Ranunculaceae).
- O^8 -Et, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **Columbidine**
[95690-58-7]
 $\text{C}_{26}\text{H}_{43}\text{NO}_5$ 449.629
Alkaloid from the aerial parts of *Aconitum columbianum* ssp. *columbianum* and roots of *Aconitum dolichorhynchum* var. *subglabratum* (Ranunculaceae). Amorph. $[\alpha]_D^{25} -6.4$ (c, 1.4 in CHCl_3).
- O^8 -Et, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et, 14-Ac: **Acoforine**. 14-Acetylcolumbidine
[95690-59-8]
 $\text{C}_{28}\text{H}_{45}\text{NO}_6$ 491.667
Alkaloid from the roots of *Aconitum forrestii* (Ranunculaceae). Amorph. $[\alpha]_D -35.3$ (c, 0.57 in EtOH).
- 14-Ketone, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me, N-Et: **14-Dehydrotalatzimine**
[36069-44-0]
 $\text{C}_{24}\text{H}_{37}\text{NO}_5$ 419.56
Alkaloid from *Aconitum saposhnikovii*. Mp 128-130° (124-127°).
- 19-Oxo, $\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -tri-Me: **Piepunensine A**
 $\text{C}_{22}\text{H}_{33}\text{NO}_6$ 407.506
Alkaloid from the roots of *Aconitum piepunense*. Amorph. powder. Mp 94-96°. $[\alpha]_D^{20} -16.9$ (c, 0.5 in CHCl_3).
- (1β,14α,16β)-form**
 $\text{O}^{16}, \text{O}^{18}$ -Di-Me, N-Et: **Talatzidine**. **Talatisidine**
[7633-67-2]
 $\text{C}_{23}\text{H}_{37}\text{NO}_5$ 407.549
Alkaloid from the roots of *Aconitum talassicum* and aerial parts of *Aconitum columbianum* (Ranunculaceae). Mp 218-219°. $[\alpha]_D -20$ (MeOH).
- $\text{O}^{16}, \text{O}^{18}$ -Di-Me, N-Et, hydrochloride: Mp 186.9°.
- Konowalowa, R.A. et al., *Bull. Soc. Chim. Fr.*, 1940, **7**, 95-105; *CA*, **34**, 5450 (Talatzidine, *isol*)
- Rabinovich, M.S. et al., *Zh. Obsheh. Khim.*, 1942, **12**, 329-335; *CA*, **37**, 3098 (Condelphine, *isol*)
- Khaimova, M.A. et al., *Dokl. Bulg. Akad. Nauk*, 1967, **20**, 193-196; *CA*, **67**, 54296c (Cammaconine, *isol*)
- Pelletier, S.W. et al., *J.A.C.S.*, 1967, **89**, 4146-4157 (Condelphine, Talatzidine, Isotalatzidine, occur, uv, ir, pmr, struct, synth)
- Brutko, L.I. et al., *CA*, 1968, **69**, 30133n (Condelphine)
- Yunusov, M.S. et al., *Khim. Prir. Soedin.*, 1969, **5**, 515-519; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 434-437 (Talatzimine, Talatzidine, ms)
- Khaimova, M.A. et al., *Tetrahedron*, 1971, **27**, 819-822 (Talatisamine)
- Wiesner, K. et al., *J.A.C.S.*, 1974, **96**, 4990-4992 (Talisamine, synth)
- Tel'nov, V.A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 814-815; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 830-831 (Aconorine)
- Wiesner, K. et al., *Pure Appl. Chem.*, 1975, **41**, 93-112 (Talisamine, synth, rev)
- Pelletier, S.W. et al., *J.A.C.S.*, 1976, **98**, 2626-2636 (cmr)
- Pelletier, S.W. et al., *Acta Cryst. B*, 1977, **33**, 716-722 (Condelphine, *cryst struct, abs config*)
- Sakai, S. et al., *Yakugaku Zasshi*, 1979, **99**, 647-656; *CA*, **91**, 105185w (14-Acetyltalatzimine)
- Mody, N.V. et al., *Heterocycles*, 1980, **14**, 1751-1752 (Cammaconine)

- Finer-Moore, J. *et al.*, *J.O.C.*, 1981, **46**, 3399-3406 (*cmr*)
- Konno, C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 128-133 (*14-Acetylatalatizamine*)
- Sultankhodzhaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 265-266; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 249-250 (*14-Dehydrotalatizamine*)
- Wang, C. *et al.*, *Planta Med.*, 1983, **48**, 55 (*Cammaconine, occur*)
- Boido, V. *et al.*, *Can. J. Chem.*, 1984, **62**, 778-784 (*Talatizidine, 8-O-Methyltalatizamine, 14-Acetylatalatizamine, 14-Dehydrotalatizamine, Columbianine*)
- Pelletier, S.W. *et al.*, *Heterocycles*, 1985, **23**, 331-338; 1987, **25**, 365-376 (*8-O-Methyltalatizamine, Columbidine, 14-Dehydrotalatizamine, Acoforine*)
- Hao, X.J. *et al.*, *Yunnan Zhiwu Yanjiu*, 1985, **7**, 217 (*Scaconine, Scaconitine, N-Deacetylscaconitine*)
- Liang, X.-T. *et al.*, *Pure Appl. Chem.*, 1986, **58**, 711-718 (*Gymnaconitine, Methylgymnaconitine*)
- Jiang, S. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 279-284; *CA*, **105**, 94502b (*Gymnaconitine, Methylgymnaconitine*)
- Vaisov, Z.M. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 869-872; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 725-727 (*14-Methylsotalatizidine*)
- Huiling, L. *et al.*, *Heterocycles*, 1989, **29**, 2317-2326 (*Dolichotines*)
- Yunusov, M.S. *et al.*, *Nat. Prod. Rep.*, 1991, **8**, 499-526 (*Scaconine, Scaconitine, N-Deacetylscaconitine*)
- Ding, L. *et al.*, *Huaxue Xuebao*, 1992, **50**, 405-408; *CA*, **117**, 108100f (*Vilmorinine*)
- Sayed, H.M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1595-1606 (*Acosepidinine*)
- Li, Y. *et al.*, *Huaxue Xuebao*, 1994, **52**, 204-208; *CA*, **121**, 78248y (*Scaconine, Scaconitine, N-Deacetylscaconitine, Acobretines A-C*)
- Yue, J.-M. *et al.*, *Phytochemistry*, 1994, **35**, 829-831; **37**, 1467-1470 (*14-Benzoyltalatizamine, Talassicumine A*)
- Li, Y. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1994, **36**, 148-152; *CA*, **121**, 251236t (*Acobretines D,E*)
- Sultankhodzhaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1995, 283-284; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, 233-234 (*14-Benzoyltalatizamine*)
- Gao, L.M. *et al.*, *Chin. Chem. Lett.*, 1996, **7**, 135-138 (*Transconitine A*)
- Merigli, A.H. *et al.*, *Heterocycles*, 1997, **45**, 1955-1965; 1999, **51**, 1843-1848 (*Raveyine, Cyphoplectine*)
- Zheng, S. *et al.*, *Phytochemistry*, 1997, **46**, 951-954 (*Transconitine A*)
- Ulubelen, A. *et al.*, *Phytochemistry*, 1998, **48**, 385-388 (*8-Acetylcondelphine, Isotalatizidine*)
- Merigli, A.H. *et al.*, *Heterocycles*, 1999, **51**, 1843-1848 (*Cyphoplectine*)
- Li, Z.-B. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 131-137 (*Geniculatines*)
- Grandez, M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 513-516 (*8-O-Methylcolumbianine*)
- Jiang, S.-H. *et al.*, *Planta Med.*, 2002, **68**, 1147-1149 (*14-Anisoyltalatizamine, 8-Acetyl-14-anisoyltalatizamine*)
- Li, L.-Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 269-271 (*Leueantine C*)
- Tashkhodzhaev, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 611-612 (*Talatizamine, cryst struct*)
- Cai, L. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 779-781 (*18-Acetylcammaconine, Cammaconine, Piepuensine A*)
- Kolak, U. *et al.*, *Phytochemistry*, 2006, **67**, 2170-2175 (*Linearilobine*)

4-Methylnaconitane-1,7,8,14-tetrol

M-353

C₁₉H₂₉NO₄ 335.442

(1 α ,5 β ,14 α)-form

N-Et: *Cardiopetalidine*

[75375-44-9]


C₂₁H₃₃NO₄ 363.496

Alkaloid from *Delphinium cardiopetalum* (*Delphinium verdunense*) (Ranunculaceae). Mp 223-227°. [α]_D +1.1 (c, 0.18 in EtOH).

González, A.G. *et al.*, *Tet. Lett.*, 1980, **21**, 1155-1158 (*ir, pmr, ms, cryst struct*)
 Khairitdinova, E.D. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2003, **52**, 2078-2080 (*isol, cmr*)

4-Methylnaconitane-1,8,10,14-tetrol

M-354

C₁₉H₂₉NO₄ 335.442

(1 α ,14 α)-form


O¹-Me, N-Et: *Racemuloline B*C₂₂H₃₅NO₄ 377.523

Alkaloid from *Aconitum racemosum* var. *pengzhouense*. Amorph. powder. [α]_D¹⁷ +20.5 (c, 0.55 in CHCl₃).

Peng, C.-S. *et al.*, *Heterocycles*, 2002, **57**, 1117-1120 (*isol, pmr, cmr*)

4-Methylnaconitane-1,8,14,16-tetrol

M-355

(1 α ,14 α ,16 α)-formC₁₉H₂₉NO₄ 335.442

▶ AR5569480

(1 α ,14 α ,16 α)-form

O¹-Me, N-Et: *Genicinine A*

[287186-09-8]

C₂₂H₃₅NO₄ 377.523

Alkaloid from the roots of *Aconitum geniculatum*. [α]_D²⁴ -6 (c, 0.6 in CHCl₃). There is confusion over the stereo-

chem. of the C-16 position.

(1 α ,14 α ,16 β)-form

N-Et: *16-Hydroxycardiopetaline*C₂₁H₃₃NO₄ 363.496

Alkaloid from *Aconitum variegatum*. Amorph. solid. [α]_D²⁰ -11.3 (c, 0.15 in CHCl₃).

N-Et, 16-Ac: *16-Acetoxyacardiopetaline*

[160884-40-2]

C₂₃H₃₅NO₅ 405.533

Alkaloid from seeds of *Aconitum napellus* ssp. *vulgare* (Ranunculaceae).

O¹-Me, N-Et: *Kongboensine. 16-De-**methylsachaconitine*

[163734-40-5]

C₂₂H₃₅NO₄ 377.523

Alkaloid from roots of *Aconitum kongboense* (Ranunculaceae).

O¹⁶-Me, 14-Ac: *Hokbusine B*

[86500-46-1]

C₂₂H₃₃NO₅ 391.506

Alkaloid from the roots of *Aconitum carmichaeli* (Ranunculaceae). Prisms (MeOH). Mp 183-185°.

O¹⁶-Me, N-Et: *Karakoline. Karacolone.**Carmichaeline*

[39089-30-0]

C₂₂H₃₅NO₄ 377.523

Alkaloid from the tubers of *Aconitum carmichaeli*, the tubers and aerial parts of *Aconitum karakolicum*, and from *Delphinium pentagynum* (Ranunculaceae). Cryst. (Me₂CO). Mp 183-184°. [α]_D¹⁷ -10 (c, 0.3 in MeOH).

O¹⁶-Me, N-Et, 14-Ac: *14-Acetylkarakoline*

[86538-27-4]

C₂₄H₃₇NO₅ 419.56

Alkaloid from the aerial parts of *Delphinium confusum* (Ranunculaceae). Cryst. (Me₂CO). Mp 99-100°.

O¹⁶-Me, N-Et, tri-Ac: Mp 165-169°.O¹, O¹⁶-Di-Me, N-Et: *Sachaconitine. Vil-**morrianine D*

[1361-02-0]

C₂₃H₃₇NO₄ 391.55

Alkaloid from roots of *Aconitum vilmorrianum* and *Aconitum miyabei* and aerial parts of *Aconitum columbianum* (Ranunculaceae). Plates (EtOH). Mp 129-130°. [α]_D²⁰ -13.08 (c, 2.35 in EtOH).

O¹, O¹⁶-Di-Me, N-Et; hydrochloride:

Needles (Me₂CO/EtOH). Mp 203-204°. [α]_D²⁰ -30.43 (c, 0.78 in EtOH).

O¹, O¹⁶-Di-Me, N-Et, 14-Ac: *14-Acetyl-**sachaconitine. Austroconitine A*

[102719-98-2]

C₂₅H₃₉NO₅ 433.587

Alkaloid from *Aconitum delphinifolium* (Ranunculaceae). Amorph. solid; cryst. (as perchlorate). Mp 182-184° (perchlorate). [α]_D²⁵ +24.8 (c, 0.6 in CHCl₃).

O¹, O¹⁶-Di-Me, N-Et, 14-benzoyl: *14-**Benzoylsachaconitine*C₃₀H₄₁NO₅ 495.658

Alkaloid from the roots of *Aconitum nagarum* var. *lasiantrum*. Amorph. powder. [α]_D²⁵ +33.3 (c, 0.005 in CHCl₃).

O¹,O¹⁶-Di-Me, N,O⁸-di-Et: 8-O-EthylsachaconitineC₂₅H₄₁NO₄ 419.603Alkaloid from *Aconitum variegatum*. Amorph. solid. [α]_D²⁵ -17.6 (c, 0.08 in CHCl₃).**O⁸,O¹⁶-Di-Me, N-Et: 8-O-Methylkarakoline. 8-Methoxykarakoline (incorr.)**

[1015699-82-7]

C₂₃H₃₇NO₄ 391.55Alkaloid from *Delphinium gracile*. Amorph. solid. [α]_D +12.5 (c, 0.032 in CHCl₃).**O¹⁴,O¹⁶-Di-Me, N-Et: Karasamine**

[84714-33-0]

C₂₃H₃₇NO₄ 391.55Alkaloid from the aerial parts of budding *Aconitum karakolicum* (Ranunculaceae).**O¹⁴,O¹⁶-Di-Me, N-Et, 8-Ac: 8-O-Acetylkarasamine**

[161993-25-5]

C₂₅H₃₉NO₅ 433.587Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Amorph. solid.**O¹⁴,O¹⁶-Di-Me, N-Et, 1-benzoyl: 1-Benzoylkarasamine**

[84732-43-4]

C₃₀H₄₁NO₅ 495.658Alkaloid from the aerial parts of budding *Aconitum karakolicum* (Ranunculaceae).**O¹,O⁸,O¹⁶-Tri-Me, N-Et: 8-O-Methylsachaconitine**


[195153-58-3]

C₂₄H₃₉NO₄ 405.576Alkaloid from *Delphinium cardiopetalum* (Ranunculaceae). Resin. [α]_D +7 (c, 0.06 in EtOH).**O⁸,O¹⁴,O¹⁶-Tri-Me, N-Et: 8-O-Methylkarasamine**

[125263-88-9]

C₂₄H₃₉NO₄ 405.576Minor alkaloid from *Delphinium nuttallianum* (Ranunculaceae). Amorph. solid. [α]_D²³ -5.**1-Deoxy, 1,2-didehydro, O¹⁶-Me, N-Et: Delpoline**C₂₂H₃₃NO₃ 359.508Alkaloid from the roots of *Delphinium poltoratskii*. Cryst. (Me₂CO). Mp 192-195°.Suginome, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1959, **32**, 604 (*Sachaconitine, isol, uv*)Iwasa, J. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 585; *CA*, **65**, 10629e (*Karakoline, isol*)Sultankhodzhaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 199; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 194 (*Karakoline, isol, ir, pmr, ms, struct*)Pelletier, S.W. *et al.*, *Tet. Lett.*, 1977, 4027 (*Sachaconitine, ir, pmr, cmr, struct*)Sultankhodzhaev, M.N. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 629-630 (*Karasamine, 1-Benzoylkarasamine*)Konno, C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 128 (*Karakoline, isol, ir, pmr, cmr*)Hikino, H. *et al.*, *J. Nat. Prod.*, 1983, **46**, 178 (*Hokbusine B*)González, A.G. *et al.*, *Tet. Lett.*, 1983, **24**, 959 (*occur*)Vaisov, Z.M. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 801; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 744 (*14-Acetylkarakoline*)Aiyar, V.N. *et al.*, *Phytochemistry*, 1986, **25**, 973 (*14-Acetylsachaconitine*)Jiang, Z. *et al.*, *Yunnan Zhiwu Yanjiu*, 1988, **10**, 317-323; *CA*, **110**, 151273g (*14-Acetylsachaconitine, Austroconitine A*)Bai, Y. *et al.*, *Heterocycles*, 1989, **29**, 1017 (*8-O-Methylkarasamine*)Liu, H. *et al.*, *Nat. Prod. Lett.*, 1994, **5**, 147 (*16-Acetoxycardiopetaline*)Bai, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1717 (*8-O-Acetylkarasamine*)Chen, Y. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1994, **36**, 970; *CA*, **123**, 5616y (*Kongboensine*)Yusupova, I.M. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 109-111; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 91-93 (*Karakoline, cryst struct*)Reina, M. *et al.*, *Phytochemistry*, 1997, **45**, 1707 (*8-O-Methylsachaconitine*)Boronova, Z.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2000, **36**, 390-392 (*Delpoline*)Dong, J.Y. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 1005-1006 (*14-Benzoylsachaconitine*)Wang, F.P. *et al.*, *Huaxue Xuebao*, 2000, **58**, 576-579; *CA*, **133**, 147501 (*Genicine A*)Zhou, X.-L. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 456-458 (*Genicine A, pmr, cmr*)Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 837-846 (*16-Hydroxycardiopetaline, 8-O-Ethylsachaconitine*)Reina, M. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1048-1055 (*8-O-Methylkarakoline*)**4-Methylnaconitane-1,8,14-triol**

M-356



(1α,5β,14α)-form

C₁₉H₂₉NO₃ 319.443**(1α,5β,14α)-form****N-Et: Cardiopetaline**


[75375-43-8]

C₂₁H₃₃NO₃ 347.497Minor alkaloid from *Delphinium cardiopetalum* (*Delphinium verdunense*) (Ranunculaceae). Mp 179-181°. [α]_D -16 (c, 0.22 in EtOH).**O¹-Me, N-Et: Giralidine I**C₂₂H₃₅NO₃ 361.523Alkaloid from the roots of *Delphinium giraldui*. Amorph. powder. Mp 78-80°. [α]_D²⁰ -15.6 (c, 1 in CHCl₃).**(1β,5β,14α)-form****O¹-Me, N-Et: Souline D**

[214191-29-4]

C₂₂H₃₅NO₃ 361.523Alkaloid from *Delphinium souliei*. Cryst. Mp 110-112°. [α]_D²⁴ -14.9 (c, 0.47 in CHCl₃). Incorrect name assigned in CA.González, A.G. *et al.*, *Tet. Lett.*, 1980, **21**, 1155 (*Cardiopetaline, ir, pmr, ms, cryst struct*)Zhang, K. *et al.*, *Planta Med.*, 1998, **64**, 580-581 (*Souline D*)Zhou, X.-L. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 456-458 (*Giralidine I*)**4-Methylnaconitane-1,14,16-triol**


M-357

C₁₉H₂₉NO₃ 319.443**(1α,14α,16β)-form****O¹⁶-Me, N-Et: Souline E**

[244616-33-9]

C₂₂H₃₅NO₃ 361.523Alkaloid from *Delphinium souliei*. Cryst. (EtOH). Mp 77-78°. [α]_D²⁴ -8 (c, 0.05 in CHCl₃).He, L. *et al.*, *Chin. Chem. Lett.*, 1999, **10**, 395-396 (*isol, pmr, cmr*)He, L. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 1285-1286 (*isol, pmr, cmr*)**4-Methylnaconit-19(N)-ene-1,6,7,8,10,14,16-heptol**


M-358

C₁₉H₂₇NO₇ 381.425**(1α,5β,6β,14α,16β)-form****7,8-Methylene ether, O¹,O¹⁴,O¹⁶-tri-Me, 6-Ac: Barbeline**





[116199-46-3]

C₂₅H₃₅NO₈ 477.553Alkaloid from *Delphinium barbeyi* (Ranunculaceae). Mp 265-267.5°. [α]_D²² +45.03 (c, 0.31 in CHCl₃).Joshi, B.S. *et al.*, *Tet. Lett.*, 1988, **29**, 2397 (*isol, ir, pmr, cmr, cryst struct*)**4-Methylnaconit-19(N)-ene-1,6,7,8,14,16,18-heptol**

M-359



C₁₉H₂₇NO₇ 381.425**(1α,5β,6β,14α,16β)-form****O¹,O⁶,O¹⁶-Tri-Me, 18-O-(2-acetamidobenzoyl), 14-Ac: Ajadine**

[157566-48-8]


C₃₃H₄₂N₂O₁₀ 626.702Alkaloid from seeds of *Delphinium ajacis* (Ranunculaceae). Amorph. [α]_D +83.7 (c, 0.166 in CHCl₃).O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me: **Acovulparine**
C₂₃H₃₅NO₇ 437.532Alkaloid from *Aconitum vulparia*. Amorph. solid. [α]_D²⁵ +25.3 (c, 2 in CHCl₃).O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, 18-O-(2-aminobenzoyl): **Pacifidine**
[142748-52-5]C₃₀H₄₀N₂O₈ 556.655Alkaloid from seeds of *Delphinium elatum* cv. Pacific giant (Ranunculaceae). Amorph. solid. [α]_D +40.7 (c, 2.47 in CHCl₃).O¹,O⁶,O¹⁴,O¹⁶-Tetra-Me, 18-O-[2-(4-amino-3-methyl-4-oxobutanoylamino)-benzoyl]: **Bulleyanine A**
[128717-97-5]C₃₅H₄₇N₃O₁₀ 669.77Alkaloid from the roots of *Delphinium bulleyanum* (Ranunculaceae). C-1 Config. revised in 1990.O¹,O⁶,O⁸,O¹⁴,O¹⁶-Penta-Me, 18-O-(2-aminobenzoyl): **Acosepridine**
[146028-67-3]C₃₁H₄₂N₂O₈ 570.681Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Amorph. [α]_D +65 (c, 0.1 in CHCl₃).O¹,O⁶,O⁸,O¹⁶,O¹⁸-Penta-Me, 14-Ac: **Olividine**C₂₆H₃₉NO₈ 493.596Alkaloid from *Consolida oliveriana*. Amorph. solid. [α]_D²⁵ +96 (c, 0.8 in CHCl₃).O¹,O⁶,O¹⁴,O¹⁶,O¹⁸-Penta-Me: **Olivimine**C₂₄H₃₇NO₇ 451.559Alkaloid from *Consolida oliveriana*. Amorph. solid. [α]_D²⁵ +77 (c, 1.1 in CHCl₃).Wei, X. et al., *Yunnan Zhiviu Yanjiu*, 1989, **11**, 453; *CA*, **113**, 74750 (*Bulleyanine A*)Joshi, B.S. et al., *J. Nat. Prod.*, 1990, **53**, 1028 (*Bulleyanine A*, config)Sayed, H.M. et al., *J. Nat. Prod.*, 1992, **55**, 1595 (*Acosepridine*)Wada, K. et al., *Phytochemistry*, 1992, **31**, 2135 (*Pacifidine*)Desai, H.K. et al., *J. Nat. Prod.*, 1994, **57**, 677 (*Ajadinine*)Grandez, M. et al., *J. Nat. Prod.*, 2002, **65**, 513-516 (*Olividine*, *Olivimine*)Csupor, D. et al., *Helv. Chim. Acta*, 2004, **87**, 2125-2130 (*Acovulparine*)**4-Methylnaconit-19(N)-ene-1,6,8,13,14,16,18-heptol** M-360C₁₉H₂₇NO₇ 381.425**(1 α ,5 β ,6 α ,14 α ,16 β)-form**O¹,O⁶,O¹⁶,O¹⁸-Tetra-Me, 14-O-(4-methoxybenzoyl), 8-Ac: **Macrorrhynine B**
[199339-82-7]C₃₃H₄₃NO₁₀ 613.703Alkaloid from the roots of *Aconitum macrorrhynchum*. Amorph. solid. [α]_D²⁵ -50.7 (c, 0.12 in CHCl₃). λ_{\max} 234 (log ϵ 4.4); 257 (log ϵ 4.5) (CHCl₃).Yang, X.-D. et al., *Helv. Chim. Acta*, 2008, **91**, 569-574 (isol, pmr, cmr, ms)**4-Methylnaconit-19(N)-ene-1,7,8,9,14,16,18-heptol** M-361C₁₉H₂₇NO₇ 381.425**(1 α ,5 β ,14 α ,16 β)-form**7,8-Methylene ether, O¹,O¹⁴,O¹⁶,O¹⁸-tetra-Me: **Caerunine**
[175923-73-6]C₂₄H₃₅NO₇ 449.543Alkaloid from *Delphinium caeruleum*. Amorph. powder. [α]_D²⁰ -4.2 (c, 0.25 in CHCl₃).Wang, Y. et al., *Phytochemistry*, 1996, **42**, 569 (isol, ir, pmr, cmr, struct)**4-Methylnaconit-2-ene-1,6,7,8,10,14,16-heptol** M-362C₁₉H₂₇NO₇ 381.425**(1 α ,5 β ,6 β ,14 α ,16 β)-form**7,8-Methylene, 1,14-di-Me ether, N-Me, 6-Ac: **Siwanine D**
[164230-59-5]C₂₅H₃₅NO₈ 477.553From *Delphinium siwanense* var. *leptogen* (Ranunculaceae). Cryst. (EtOH). Mp 226-228°. [α]_D +30.2 (c, 0.1 in CHCl₃).7,8-Methylene, 1,14-di-Me ether, N-Et, 6-Ac: **Siwanine B**
[159903-63-6]C₂₆H₃₇NO₈ 491.58From *Delphinium siwanense* var. *leptogen* (Ranunculaceae). Amorph. solid (EtOH). Mp 194-195°. [α]_D -35.4 (c, 5.9 in CHCl₃). Incorrect struct. in CA.7,8-Methylene, 1,14-di-Me ether, N-Et, 6,16-di-Ac: 16-Acetyl-2,3-dehydroelastine. **Siwanine E**
[195247-83-7]C₂₈H₃₉NO₉ 533.617Alkaloid from *Delphinium siwanense* (Ranunculaceae). Resin. [α]_D +20.8 (c, 0.7 in CHCl₃).7,8-Methylene, 1,16-di-Me ether, N-Et, 6-Ac: 2,3-Dehydrodictyocarpine. **Siwanine F**
[195631-62-0]C₂₆H₃₇NO₈ 491.58Alkaloid from *Delphinium siwanense* (Ranunculaceae). Powder. Mp 208-210°. [α]_D +34.1 (c, 1.5 in CHCl₃).7,8-Methylene, 1,14,16-tri-Me ether, N-Et: **Deacetylsiwanine A**
[297137-06-5]C₂₅H₃₇NO₇ 463.57Alkaloid from *Delphinium orthocentrum*. Needles. Mp 225-227°. [α]_D²⁰ +56 (c, 0.25 in CHCl₃).7,8-Methylene, 1,14,16-tri-Me ether, N-Et, 6-Ac: **Siwanine A**
[159903-64-7]C₂₇H₃₉NO₈ 505.607Alkaloid from *Delphinium siwanense* var. *leptogen* (Ranunculaceae). Amorph. powder (EtOH). Mp 172-174°. [α]_D +27.1 (c, 4.8 in CHCl₃).Zhang, S.M. et al., *Phytochemistry*, 1997, **45**, 1713-1716; 1998, **48**, 191-196 (*Siwanines*, pmr, cmr, ms, cryst struct)Ding, L.S. et al., *Zhiviu Xuebao (Acta Bot. Sin.)*, 2000, **42**, 523-525 (*Deacetylsiwanine A*)**4-Methylnaconit-2-ene-1,6,7,8,14,16,18-heptol** M-363C₁₉H₂₇NO₇ 381.425**(1 α ,5 β ,6 β ,14 α ,16 β)-form**O¹⁶,O¹⁸-Di-Me, N-Et: **Ibukinamine**. 2,3-Dehydro-6-O-demethyldelecosine
[88142-58-9]C₂₃H₃₅NO₇ 437.532Alkaloid from the roots of *Aconitum ibukiense* (Ranunculaceae). Mp 243-246°. [α]_D⁹ +71.7 (c, 0.12 in MeOH).O⁶,O¹⁶,O¹⁸-Tri-Me, N-Et: 2,3-Dehydrodelecosine
[119237-52-4]C₂₄H₃₇NO₇ 451.559Alkaloid from the roots of *Aconitum japonicum* var. *montanum* (Ranunculaceae). Amorph. solid. [α]_D¹² +86.2 (c, 0.09 in CHCl₃).O⁶,O¹⁴,O¹⁶,O¹⁸-Tetra-Me, N-Et: **Giraldine E**

C₂₅H₃₉NO₇ 465.586Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder. Mp 51-53°. [α]_D²⁰ +16 (c, 0.8 in CHCl₃).14-Ketone, O⁶, O¹⁶, O¹⁸-tri-Me, N-Et: **Takaonine**


[71264-39-6]

C₂₄H₃₅NO₇ 449.543Alkaloid from *Aconitum japonicum* and from the roots of *Aconitum ibukiense* (Ranunculaceae). Mp 186-187.5°. [α]_D¹⁹ +52 (c, 0.22 in CHCl₃).Sakai, S. et al., *Yakugaku Zasshi*, 1979, **99**, 647; 1984, **104**, 222 (*Takaonine*)Sakai, S. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3338 (*Ibukinamine*, ir, pmr, ms, cryst struct)Takayama, H. et al., *Chem. Pharm. Bull.*, 1988, **36**, 3210 (*2,3-Dehydrodelcosine*)Zhou, X.-L. et al., *Heterocycles*, 2004, **63**, 123-128 (*Giraldine E*)**4-Methylnaconit-2-ene-1,6,8,13,14,16,18-heptol** M-364C₁₉H₂₇NO₇ 381.425O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-(4-methoxybenzoyl), 8-Ac: **Liaconitine A**
C₃₅H₄₇NO₁₀ 641.757Alkaloid from *Aconitum episcopale*.Needles. Mp 146-147.3°. [α]_D¹¹ +53.8 (c, 0.26 in CHCl₃).O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 8,14-bis(4-methoxybenzoyl): **Liaconitine B**
[221347-29-1]C₄₁H₅₁NO₁₁ 733.854Alkaloid from *Aconitum episcopale*.Amorph. powder. [α]_D¹³ -2.3 (c, 0.77 in CHCl₃).O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N, O⁸-di-Et, 14-(4-methoxybenzoyl): **Liaconitine C**
C₃₅H₄₉NO₉ 627.773Alkaloid from *Aconitum episcopale*.Amorph. powder. [α]_D³ +46 (c, 0.36 in CHCl₃).Yang, J.H. et al., *Phytochemistry*, 1999, **50**, 345-348**4-Methylnaconit-8(15)-ene-1,3,6,13,14,16,18-heptol** M-365C₁₉H₂₇NO₇ 381.425(1 α , 3 α , 5 β , 6 α , 14 α , 16 β)-formO¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-benzoyl: **Mithaconitine**


[62926-58-3]

C₃₂H₄₃NO₈ 569.694Minor alkaloid from the roots of *Aconitum falconeri* (Ranunculaceae). Amorph. [α]_D²⁸ +94 (c, 1.0 in EtOH).O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Falacnitine**
[62926-57-2]C₃₄H₄₇NO₁₀ 629.746Alkaloid from the roots of *Aconitum falconeri* (Ranunculaceae). Amorph. [α]_D²⁸ +111.5 (c, 1.0 in EtOH).Pelletier, S.W. et al., *Chem. Comm.*, 1977, **12** (ir, pmr, cmr, struct)Pelletier, S.W. et al., *Heterocycles*, 1977, **7**, 327 (cmr)**4-Methylnaconit-15-ene-1,3,6,8,13,14,18-heptol** M-366C₁₉H₂₇NO₇ 381.425(1 α , 3 α , 5 β , 6 α , 14 α)-formO¹, O⁶, O¹⁸-Tri-Me, N-Et, 14-O-(3,4-dimethoxybenzoyl): **Balfourine**


[142808-40-0]

C₃₃H₄₅NO₁₀ 615.719Alkaloid from roots of *Aconitum balfourii* (Ranunculaceae). Amorph. First naturally occurring isopyrroditerepenoid alkaloid.Khwah, K.S. et al., *Heterocycles*, 1992, **34**, 441 (isol, pmr, cmr, struct)**4-Methylnaconit-19(N)-ene-1,3,6,8,14,16-hexol** M-367C₁₉H₂₇NO₆ 365.425(1 α , 3 α , 5 β , 6 α , 14 α , 16 β)-formO¹, O⁸, O¹⁴, O¹⁶-Tetra-Me, 6-Ac: **Guenerine**

[149998-33-4]

C₂₅H₃₇NO₇ 463.57Alkaloid from aerial parts of *Delphinium gueneri* (Ranunculaceae). [α]_D²² -23.3 (c, 0.1 in MeOH).Ulubelen, A. et al., *Phytochemistry*, 1993, **33**, 213 (isol, ir, pmr, cmr, ms, struct)**4-Methylnaconit-19(N)-ene-1,6,7,8,14,16-hexol** M-368C₁₉H₂₇NO₆ 365.425(1 α , 5 β , 6 β , 14 α , 16 β)-form7,8-Methylene ether, O¹, O¹⁴, O¹⁶-tri-Me, 6-Ac: **Tiantaishanmine**

[959843-32-4]


C₂₅H₃₅NO₇ 461.554Alkaloid from the roots of *Delphinium tiantaishanense*. Amorph. powder. Mp 251-253°. [α]_D²⁰ +24.2 (c, 0.26 in CHCl₃).Li, J. et al., *Molecules*, 2007, **12**, 353-360 (isol, pmr, cmr)**4-Methylnaconit-19(N)-ene-1,6,8,14,16,18-hexol** M-369C₁₉H₂₇NO₆ 365.425(1 α , 5 β , 6 α , 14 α , 16 β)-formO¹, O⁶, O¹⁶, O¹⁸-Tetra-Me: **Leucanthum-sine D**

[947591-62-0]

C₂₃H₃₅NO₆ 421.533Alkaloid from the roots of *Aconitum sungpanense* var. *leucanthum*. Amorph. powder. Mp 150-152°. [α]_D²⁰ +64.1 (c, 1 in CHCl₃).O¹, O⁶, O¹⁶, O¹⁸-Tetra-Me, 14-O-(4-methoxybenzoyl), 8-Ac: **Macrorhynine A**
[1019744-40-1]C₃₃H₄₃NO₉ 597.704Alkaloid from the roots of *Aconitum macrorhynchum*. Amorph. solid. [α]_D²⁵ -9.1 (c, 0.11 in CHCl₃). λ _{max} 232 (log ϵ 4.48); 259 (log ϵ 4.58) (CHCl₃).Yan, H. et al., *Helv. Chim. Acta*, 2007, **90**, 1133-1140 (*Leucanthumsine D*)

Yang, X.-D. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 569-574 (*Macrorrhynine A*)

4-Methylnaconit-19(N)-ene-1,7,8,14,16,18-hexol M-370



C₁₉H₂₇NO₆ 365.425


(1α,5β,14α,16β)-form

O¹,O¹⁴,O¹⁶,O¹⁸-*Tetra-Me*: **Tongolenine C** [198754-25-5]
C₂₃H₃₅NO₆ 421.533
Alkaloid from *Delphinium tongolense* (Ranunculaceae). Powder. [α]_D +44.4 (c, 0.5 in CHCl₃). Mp. >350°.

He, L. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 791-792 (*isol, pmr, cmr, struct*)

He, L. *et al.*, *Indian J. Chem., Sect. B*, 1997, **8**, 791-792

4-Methylnaconit-2-ene-1,6,7,8,14,16-hexol M-371



C₁₉H₂₇NO₆ 365.425

(1α,5β,6β,14α,16β)-form

O¹⁶-*Me*, N-*Et*: **Glabredelphinine** [132160-37-3]
C₂₂H₃₃NO₆ 407.506
Alkaloid from *Delphinium kamaonense* var. *glabrescens* (Ranunculaceae). Cryst. Mp 201-203°.

O⁶,O¹⁶-*Di-Me*, N-*Et*: **Giraldine A**
C₂₃H₃₅NO₆ 421.533
Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder (CHCl₃/MeOH). Mp 79-80°. [α]_D +75.3 (c, 0.55 in CHCl₃).

O⁶,O¹⁶-*Di-Me*, N-*Et*, 14-*Ac*: **Giraldine B**
C₂₅H₃₇NO₇ 463.57
Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder (CHCl₃/MeOH). Mp 76-78°. [α]_D +37.5 (c, 0.45 in CHCl₃).

O⁶,O¹⁶-*Di-Me*, N-*Et*, 14-*benzoyl*: **Giraldine C**
C₃₀H₃₉NO₇ 525.641
Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder (CHCl₃/

MeOH). Mp 58-59°. [α]_D +36.5 (c, 0.4 in CHCl₃).

O⁶,O⁸,O¹⁶-*Tri-Me*, N-*Et*: **Giraldine D**
C₂₄H₃₇NO₆ 435.559
Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder. Mp 159-160°. [α]_D +38.5 (c, 0.3 in CHCl₃).


7,8-*Methylene ether*, O¹,O¹⁴,O¹⁶-*tri-Me*, N-*Me*, 6-*Ac*: **Siwanine C**
[164176-43-6]
C₂₆H₃₇NO₇ 475.581
Alkaloid from *Delphinium siwanense* var. *leptogen*. Amorph. solid. [α]_D +9.6 (CHCl₃).

7,8-*Methylene ether*, O¹,O¹⁴,O¹⁶-*tri-Me*, N-*Et*: **Deacetyltatsiensine**
[86695-19-4]
C₂₇H₃₉NO₇ 489.608
Alkaloid from the roots of *Delphinium kamaonense*, *Delphinium siwanense* var. *leptogen* and *Delphinium tatsienense* (Ranunculaceae). Amorph. [α]_D +17.4 (c, 0.3 in EtOH).

14-*Ketone*, O⁶,O¹⁶-*di-Me*, N-*Et*: **Giraldine F**
C₂₃H₃₃NO₆ 419.517
Alkaloid from the roots of *Delphinium giraldii*. Amorph. powder. Mp 55-57°. [α]_D +12.3 (c, 0.7 in CHCl₃).

Pelletier, S.W. *et al.*, *Heterocycles*, 1983, **20**, 1347-1354 (*Tatsiensine*)
Ding, L.-S. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 438-440 (*Glabredelphinine*)
Zhang, S.M. *et al.*, *Phytochemistry*, 1998, **48**, 191-196 (*Siwanine C*, *Tatsiensine*)
Zhou, X.-L. *et al.*, *Chin. J. Chem.*, 2003, **21**, 871-874 (*Giraldines A-C*)
Zhou, X.-L. *et al.*, *Heterocycles*, 2004, **63**, 123-128 (*Giraldines D,F*)

4-Methylnaconit-8(15)-ene-1,6,13,14,16,18-hexol M-372



C₁₉H₂₇NO₆ 365.425

(1α,5β,6α,14α,16β)-form

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Me*, 14-*benzoyl*: **Pyrodelphinine**
[60050-12-6]
C₃₁H₄₁NO₇ 539.667

Alkaloid from the seeds of *Delphinium staphisagria* (Ranunculaceae). Also a pyrolysis prod. of Delphinine. Cryst. (MeOH). Mp 208-212°. Not an artifact.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-*benzoyl*: **Pyrochasmaconitine**
[156401-00-2]
C₃₂H₄₃NO₇ 553.694

Alkaloid from roots of *Aconitum kongboense* (Ranunculaceae). Cryst. (hexane/Me₂CO). Mp 159-160°.

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, N-*Et*, 14-*O*-(4-*methoxybenzoyl*): **Pyrocrassicauline A**
[156401-01-3]

C₃₃H₄₅NO₈ 583.72
Alkaloid from roots of *Aconitum kongboense* (Ranunculaceae). Cryst. (hexane/Me₂CO). Mp 132-134°.


Pelletier, S.W. *et al.*, *Heterocycles*, 1977, **7**, 327 (*cmr*)

Pelletier, S.W. *et al.*, *J.O.C.*, 1982, **47**, 5290 (*Pyrodelphinine, synth, cmr, cryst struct*)

Liang, X. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1307 (*Pyrodelphinine, isol*)

Yue, J.-M. *et al.*, *Phytochemistry*, 1994, **35**, 829 (*Pyrochasmaconitine, Pyrocrassicauline A*)

4-Methylnaconit-19(N)-ene-1,3,6,8,13,14,15,16,18-nonol M-373



C₁₉H₂₇NO₉ 413.424


(1α,3α,5β,6α,14α,15α,16β)-form

O¹,O⁶,O¹⁶,O¹⁸-*Tetra-Me*, 14-*benzoyl*, 8-*Ac*: **Merckonine**
C₃₂H₄₁NO₁₁ 615.676

Alkaloid from Aconitine Potent Merck (*Aconitum napellus*). Amorph. [α]_D -29.9 (c, 0.3 in CHCl₃).

Desai, H.K. *et al.*, *Heterocycles*, 1998, **48**, 1107-1110 (*isol, ir, pmr, cmr*)

4-Methylnaconit-19(N)-ene-1,6,8,14,16-pentol M-374



C₁₉H₂₇NO₅ 349.426

(1α,5β,6β,14α,16β)-form


O¹,O⁸,O¹⁶-*Tri-Me*, 6-*Ac*: **Peregrinine**[†]
[168010-53-5]

C₂₄H₃₅NO₆ 433.544
Alkaloid from epigeal parts of *Delphi-*

nium peregrinum var. *elongatum* (Ranunculaceae). Amorph. $[\alpha]_D^{25} +66.7$ (c, 0.1 in CHCl_3).

De la Fuente, G. *et al.*, *Phytochemistry*, 1995, **39**, 1459 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

4-Methylnaconit-19(N)-ene-1,8,14,16,18-pentol M-375



$\text{C}_{19}\text{H}_{27}\text{NO}_5$ 349.426

(1 α ,14 α ,16 β)-form

$\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me: *Nagadine*. *Pengshenine B*

$\text{C}_{22}\text{H}_{33}\text{NO}_5$ 391.506

Alkaloid from the roots of *Aconitum nagarum* var. *lasiandrum* and *Aconitum hemsleyanum* var. *penzhouense*. Amorph. powder. $[\alpha]_D^{25} +43$ (c, 0.003 in CHCl_3).

$\text{O}^1, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, 8-hexadecanoyl, 14-Ac: *Transconitine C* [175923-72-5]

$\text{C}_{40}\text{H}_{65}\text{NO}_7$ 671.956


Alkaloid from the roots of *Aconitum transsectum*. Amorph. $[\alpha]_D^{20} +40$ (c, 0.02 in CHCl_3).

Zheng, S. *et al.*, *Phytochemistry*, 1997, **46**, 951-954 (*Transconitine C*)

Dong, J.Y. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 1005-1006 (*Nagadine*)

Peng, C.S. *et al.*, *Chin. Chem. Lett.*, 2002, **13**, 233-236 (*Pengshenine B*)

4-Methylnaconit-19(N)-ene-7,8,10,14,16-pentol M-376



$\text{C}_{19}\text{H}_{27}\text{NO}_5$ 349.426


$\text{O}^8, \text{O}^{14}, \text{O}^{16}$ -Tri-Me: *Orthocentrine* [297137-05-4]

$\text{C}_{22}\text{H}_{33}\text{NO}_5$ 391.506

Alkaloid from *Delphinium orthocentrum*. Cryst. Mp 224-226°. $[\alpha]_D^{20} -94$ (c, 0.07 in CHCl_3).

Ding, L.-S. *et al.*, *Zhivuo Xuebao (Acta Bot. Sin.)*, 2000, **42**, 523-525 (*isol*, *pmr*, *cmr*, *ms*)

4-Methylnaconit-8(15)-ene-1,6,14,16,18-pentol M-377



$\text{C}_{19}\text{H}_{27}\text{NO}_5$ 349.426

(1 α ,5 β ,6 α ,14 α ,16 β)-form

$\text{O}^1, \text{O}^6, \text{O}^{16}, \text{O}^{18}$ -Tetra-Me, N-Et: *Pyrochasmamine*

[5066-77-3]


$\text{C}_{25}\text{H}_{39}\text{NO}_5$ 433.587

Alkaloid from *Aconitum yesoense* (Ranunculaceae). Prisms (hexane). Mp 126-129°.

Achmatowicz, O. *et al.*, *Can. J. Chem.*, 1965, **43**, 825 (*synth*, *ir*, *pmr*)

Takayama, H. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 245; *CA*, **97**, 36082v (*isol*)

4-Methylnaconit-19(N)-ene-1,8,14,16-tetrol M-378



$\text{C}_{19}\text{H}_{27}\text{NO}_4$ 333.427

(1 α ,14 α ,16 β)-form


$\text{O}^1, \text{O}^{16}$ -Di-Me: *N-Deethyl-N,19-didehydrosachaconitine*

$\text{C}_{21}\text{H}_{31}\text{NO}_4$ 361.48

Alkaloid from *Aconitum variegatum*. Amorph. solid. $[\alpha]_D^{25} +181.8$ (c, 0.11 in CHCl_3).

Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 837-846 (*isol*, *pmr*, *cmr*, *ms*)

4-Methylnaconit-2-ene-1,6,14,16-tetrol M-379



$\text{C}_{19}\text{H}_{27}\text{NO}_4$ 333.427

(1 α ,5 β ,6 β ,14 α ,16 β)-form


$\text{O}^{14}, \text{O}^{16}$ -Di-Me, N-Et, 6-Ac: *Caeruleine* [147468-47-1]

$\text{C}_{25}\text{H}_{37}\text{NO}_5$ 431.571

Minor alkaloid from *Delphinium caeruleum* (Ranunculaceae). Amorph. powder. $[\alpha]_D^{17} +13.5$ (c, 0.5 in MeOH).

Pan, Y.-J. *et al.*, *Planta Med.*, 1993, **59**, 83 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

4-Methylnaconit-8(15)-ene-1,14,16,18-tetrol M-380



$\text{C}_{19}\text{H}_{27}\text{NO}_4$ 333.427

(1 α ,5 β ,14 α ,16 β)-form


$\text{O}^{14}, \text{O}^{16}, \text{O}^{18}$ -Tri-Me, N-Et: *Aconasutine* [179598-81-3]

$\text{C}_{24}\text{H}_{37}\text{NO}_4$ 403.561

Alkaloid from aerial parts of *Aconitum nasutum*.

Merikli, A.H. *et al.*, *Turk. J. Chem.*, 1996, **20**, 164-167 (*isol*, *pmr*, *cmr*)

4-Methylnaconit-15-ene-1,14,18-triol M-381



$\text{C}_{19}\text{H}_{27}\text{NO}_3$ 317.427

(1 α ,5 β ,14 α)-form

O^1 -Me, N-Et, 18-O-(2-acetamidobenzoyl): *Talassicumine C* [160824-51-1]


$\text{C}_{31}\text{H}_{40}\text{N}_2\text{O}_5$ 520.667

Alkaloid from roots of *Aconitum talassicum* (Ranunculaceae). Amorph. powder.

Yue, J. *et al.*, *Phytochemistry*, 1994, **37**, 1467 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

10-Methylacridone M-382

10-Methyl-9(10H)-acridinone, 9CI. 10-Methyl-9-acridanone, 8CI. N-Methylacridone [719-54-0]



C₁₄H₁₁NO 209.247

Constit. of the roots of *Citrus deliciosa* (Italian tangerine) and *Thamnosma montana* (Rutaceae). Cryst. (MeOH). Mp 199-200°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 93A (ir)

Gilman, H. *et al.*, *J.O.C.*, 1952, 17, 860 (synth)
Schumann, D. *et al.*, *Chem. Ber.*, 1969, 102, 3192 (ms)

Chang, P.T.O. *et al.*, *J. Nat. Prod.*, 1976, 39, 134 (isol, uv, ir, pmr, ms)

Potescu, I.D. *et al.*, *J. Prakt. Chem.*, 1976, 318, 515 (synth)

Beak, P. *et al.*, *J.A.C.S.*, 1976, 98, 171 (uv)

Adams, J.H. *et al.*, *J.C.S. Perkin 1*, 1977, 2173 (synth)

Grignon-Dubois, M. *et al.*, *Synth. Commun.*, 1995, 25, 2999 (synth)

El-Shafae, A.M. *et al.*, *Pharmazie*, 1998, 53, 640-643 (cmr)

Methylamine, 8CI**M-383**

Methanamine, 9CI. Aminomethane. Carbinamine

[74-89-5]

MeNH₂CH₅N 31.057

Found in fish oils and plants, e.g.

Mercurialis annua, *Mercurialis perennis*.

Fairly widespread in fungi. Refrigerant, nucleophile in org. synth., solvent.

Flammable gas with strong ammoniacal odour. Freq. encountered as strong aq. soln. V. sol. H₂O. d₄¹⁰ 0.7. Mp -93.5°. Bp

-6.3° Bp₂₀₀ -32.4° Bp₁₀ -73.8°. Crit. pt. 156.9°/73.6 atm.

- ▶ Extremely flammable, fl. p. -18°, auto-ignition temp. 430°. Eye, skin and respiratory tract irritant (and aq. soln.). OES: long-term 10 ppm. PF6300000

Monohydrate:

Liq. d^{13.9} 0.9.

Trihydrate: Mp -35.8°.

Hydrochloride: [593-51-1] Used in tanning industry.

Plates (EtOH). Sol. EtOH; insol.

Me₂CO, Et₂O, CHCl₃. Mp 227-228°.

Bp₁₅ 225-230°.

- ▶ PA0603000

Di-Ac: Methyl diacetamide

[1113-68-4]

C₅H₉NO₂ 115.132

Misc. H₂O; insol. Et₂O. Bp 192°.

N-(2-Methylpropylidene): N-Isobutylidene-methylamine, 8CI

[6898-70-0]

C₅H₁₁N 85.149

Bp 69.5°.

Nitrate: Methylammonium nitrate

[22113-87-7]

CH₆N₂O₃ 94.07

Sensitiser for slurry and emulsion explosives containing ammonium nitrate. V. hygroscopic cryst. (EtOH). Mp 110.5-111.5°.

- ▶ Explosive.

[7436-22-8, 17000-00-9, 14779-52-3, 14779-55-6]

Aldrich Library of NMR Spectra, 2nd edn., 1983, 1, 237B (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 279A; 279B (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 363D (ir)

Org. Synth., Coll. Vol., 2, 1943, 374 (synth)

Tiollais, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1947, 708 (*Isobutylidene-methylamine*)

Cottrell, T.L. *et al.*, *J.C.S.*, 1951, 1798-1800 (*nitrate, synth*)

Hesse, G. *et al.*, *Annalen*, 1957, 607, 24 (*synth*)

von Kamienski, E.S. *et al.*, *CA*, 1958, 52, 5551,6502 (*occur*)

Suhr, H. *et al.*, *Chem. Ber.*, 1963, 96, 1720 (*pmr*)

Fr. Pat., 1968, 1 525 149; *CA*, 71, 80663 (*synth*)

Jayasooriya, U.A. *et al.*, *J. Chem. Soc., Faraday Trans.*, 1992, 1631-1636 (*nitrate, ir, Raman*)

Kurniadi, W. *et al.*, *J.O.C.*, 1994, 59, 5502-5505 (*nitrate, synth, props*)

Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn., (ed. Snyder, R.), Elsevier, Volume 2, 1990, 113

Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 0469


Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 825; 826

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MGC250

2-(Methylamino)benzoic acid, 9CI**M-384**

N-Methylantranilic acid, 8CI

[119-68-6]

C₈H₉NO₂ 151.165

Isol. from grapefruit peel oil. Plates (EtOH or petrol). Mp 178-179° (175-182°). Bp_{0.01} 80°. pK_{a1} 2; pK_{a2} 5.34 (22°, 0.1M KCl). Amphoteric. Sublimes.

- ▶ Eye irritant. CB3300000

Hydrochloride:

Needles. Mp 141°. No CAS no. found 8-14CI.

Me ester: Methyl N-methylantranilate.

FEMA 2718

[85-91-6]

C₉H₁₁NO₂ 165.191

Constit. of mandarin peel oil (*Citrus madurensis*), hyacinth oil, petitgrain oil and oil of *Kaempferia ethelae* bulbs.

Flavouring agent. Cryst. (petrol) with apple-like flavour at low conc. Mp 19°.

Bp 256° Bp₁₅ 130-131°.

- ▶ CB3500000

Me ester, hydrochloride:

Needles (EtOH). Mp 218°.

Et ester: [35472-56-1]

C₁₀H₁₃NO₂ 179.218

Mp 39°. Bp₄₅ 172-175°.

Ph ester: [31358-73-3]

C₁₄H₁₃NO₂ 227.262

Yellow needles (EtOH aq.). Mp 70-71°.

Amide: 2-(Methylamino)benzamide

[7505-81-9]

C₈H₁₀N₂O 150.18

Alkaloid from the fruit of *Evodia rutaecarpa* and the flowers of *Glycosmis pentaphylla*. Plates (EtOH). Mp 162-163°.

- ▶ CV5570450

N-Ac: [78944-67-9]

C₁₀H₁₁NO₃ 193.202

Needles. Mp 192-193°.

N-(3-Phenylpropanoyl): 2-[Methyl(3-phenylpropanoyl)amino]benzoic acid [207305-65-5]

C₁₇H₁₇NO₃ 283.326

Prod. by the marine *Streptomyces* sp.

B7747. Antimicrobial agent. Cryst.

(Et₂O). Mp 107°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 189B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 1066C (nmr)

Houben, J. *et al.*, *Ber.*, 1906, 39, 3233-3240;

1909, 42, 3188-3196 (*synth, Ph ester*)

Cosulich, D.B. *et al.*, *J.A.C.S.*, 1948, 70, 1922-

1926 (*synth*)

Tomita, K. *et al.*, *Yakugaku Zasshi*, 1951,

71, 1100-1112; *CA*, 46, 50456 (*synth, esters*)

Freifelder, M. *et al.*, *J. Phys. Chem.*, 1965, 69,

3645-3648 (*pmr*)

Pant, L.M. *et al.*, *Acta Cryst. B*, 1972, 28, 647-

649 (*cryst struct*)

Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, MGQ250

Sinhbabu, A. *et al.*, *Asian J. Chem.*, 1995, 7,

221-222 (*amide, isol*)

Biabani, M.A.F. *et al.*, *J. Antibiot.*, 1998, 51,

333-340 (*3-phenylpropanoyl*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley,

2000, MGQ000; MGQ250

[(Methylamino)carbonyl]carbamic acid**M-385**

ω-Methylallophanic acid

[114728-45-9]

MeNHCONHCOOH

C₃H₆N₂O₃ 118.092

Isol. from *Echinops echinatus*.

Me ester: Methyl [(methylamino)carbonyl]carbamate, 9CI. Monospermin

[83225-61-0]

C₄H₈N₂O₃ 132.119

Alkaloid from *Butea monosperma* seeds (Fabaceae). Cryst. (EtOH). Mp

161-163°. Formerly assigned an incorrect imidazolone struct.

N²-Hydroxy: Hydroxy[(methylamino)-carbonyl]carbamic acid, 9CI. 2-Hydroxy-ω-methylallophanic acid

[115909-20-1]

C₃H₆N₂O₄ 134.091

Isol. from *Butea monosperma*. Shining needles (MeOH) (as Me ester). Mp

171-172° (Me ester).

Mehta, B. *et al.*, *Chem. Ind. (London)*, 1981, 98 (*isol*)


Dhar, K.L. *et al.*, *Chem. Ind. (London)*, 1982, 862 (*struct*)

Mochalin, V.B. *et al.*, *Zh. Org. Khim.*, 1982, 18, 1202 (*synth*)

Porwal, M. *et al.*, *Indian J. Chem., Sect. B*, 1988, 27, 281 (*isol, deriv*)

2-(Methylamino)-1-phenyl-1-propanol M-386

α -[1-(Methylamino)ethyl]benzenemethanol, 9CI
[53214-57-6]



$C_{10}H_{15}NO$ 165.235

Resolving agent for aldehydes and ketones via chiral oxazolidine formn. Used as 0.2% soln. in $CHCl_3$ for photometric detn. of As. Orally active sympathomimetic agent. Weaker but longer-acting than adrenaline. Hypertensive, cardiac stimulant, bronchodilator, hyperglycaemic agent. Low toxicity. Has been used clinically against bronchial asthma, hay fever, whooping cough, myasthenia gravis, dysmenorrhea and heart block (Stokes-Adam syndrome). Log P 0.89 (calc).

► DO9500000

(1R,2R)-form

(-)- ψ -Ephedrine

[321-97-1]

Synthetic. Mp 118-118.5°. $[\alpha]_D^{20}$ -52 (EtOH).

► UL5750500

(1R,2S)-form

Ephedrine, BAN

[299-42-3]

Main alkaloid from the Chinese drug "Ma-Huang", and from many *Ephedra* spp., also in *Aconitum napellus*, *Catha edulis*, *Taxus baccata*, *Sida cordifolia*, *Roemeria refracta* and some other spp. (Ephedraceae, Ranunculaceae, Celastraceae, Taxaceae, Malvaceae, Papaveraceae). Sol. $CHCl_3$. Mp 40°. Bp 225°. $[\alpha]_D$ -6.3 (EtOH). pK_a 9.52 (25°). Log P 0.89 (calc). More pharmacol. active isomer.

► Adverse effects reported when used therapeutically. LD₅₀ (rat, orl) 600 mg/kg. Exp. teratogen. KB0700000

Hydrochloride: **Ephedrine hydrochloride**, JAN, USAN

[50-98-6]

Mp 218°. $[\alpha]_D^{20}$ -37 (H₂O). Component of Amesec, Bronkotabs, Primatene M, Primatene P, Quadrinal, Quibron Plus and Tedral.

► Adverse effects when used therapeutically. KB1750000

Sulfate: **Ephedrine sulfate**, USAN. Isofedrol

[134-72-5] Component of Wyanooids.

► Adverse effects when used therapeutically. LD₅₀ (rat, orl) 404 mg/kg. KB2625000

N-Me: see 2-Dimethylamino-1-phenyl-1-propanol, D-738

N-Ac: [52305-20-1]

$C_{12}H_{17}NO_2$ 207.272

Mp 85-86°. $[\alpha]_D^{20}$ +8 (EtOH). $[\alpha]_D^{20}$ -63 ($CHCl_3$).

Compd. with theophylline: see Theophylline, T-360

Nitrate salt: [81012-98-8]

Mp 126-128°. $[\alpha]_D^{21}$ -31 (c, 5 in H₂O).

(1S,2S)-form

ψ -Ephedrine. Pseudoephedrine, BAN,

INN. Isoephedrine

[90-82-4]

Alkaloid from *Ephedra vulgaris* var. *helvetica* and many other *Ephedra* spp., and from *Roemeria refracta* and *Sida cordifolia* (Ephedraceae, Papaveraceae, Malvaceae). Shows somewhat similar pharmacol. activity to Ephedrine but has less pressor activity and CNS effects. Used for illicit production of methamphetamine. Since 2005, banned for use in OTC formulations in USA. Mp 117-118°. $[\alpha]_D$ +52 (EtOH). pK_a 9.57 (30°). Log P 0.89 (calc). Also used as a copolymer with sulfonated styrene-divinylbenzene (Pseudoephedrine polistirex, USAN) as a nasal decongestant.

► Adverse effects when used therapeutically. LD₅₀ (rat, orl) 660 mg/kg. UL5800000

Hydrochloride: **Pseudoephedrine hydrochloride**, USAN

[345-78-8] Adrenergic (vasoconstrictor). Mp 182-182.5°. $[\alpha]_D^{20}$ -62 (H₂O).

Component of Actifed, Novafed, Sudafed, Sinutab, Demazin, Phenergan and numerous proprietary formulations. In some formulations, replaced by phenylephrine.

► Human systemic effects if ingested. LD₅₀ (mus, orl) 371 mg/kg. UL5950000

Sulfate: **Pseudoephedrine sulfate**, USAN.

Afrinol. D-Isoephedrine sulfate, JAN

[7460-12-0] Component of Disophrol and Drixoral.

N-De-Me: see 2-Amino-1-phenyl-1-propanol, A-869

N-Me: see 2-Dimethylamino-1-phenyl-1-propanol, D-738

(1S,2R)-form

(+)-Ephedrine

[321-98-2]

Synthetic. Resolving agent for aldehydes and ketones. Shows about 1/3 the pharmacol. activity of (-)-ephedrine. Mp 40-40.5°.

► LD₅₀ (mus, ipr) 255 mg/kg. KB0600000

Hydrochloride: [24221-86-1]

Mp 217-218°. $[\alpha]_D^{20}$ +34 (H₂O).

► KB1925000

(1RS,2RS)-form

(±)- ψ -Ephedrine. Racephedrine, BAN, INN

[4125-58-0]

Synthetic. Sympathomimetic agent.

Mp 118°.

(1RS,2SR)-form

(±)-Ephedrine

[90-81-3]

Mp 76°.

► KB0610000

Hydrochloride: [134-71-4]

Mp 188-189.5°.

► KB1575000

N-Ac:

$C_{12}H_{17}NO_2$ 207.272

Mp 77-78.5°.

[50906-05-3, 670-40-6, 56979-55-6]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 1277C; 1278A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 578C; 579A; 579B; 586A; 586B; 586C; 587A; 587B (nmr)

Emde, H. et al., *Helv. Chim. Acta*, 1929, 12, 365; 405 (struct, rel config)

Freudenberg, K. et al., *J.A.C.S.*, 1932, 54, 234

Freudenberg, K. et al., *Annalen*, 1934, 510, 223 (abs config)

Phillips, D.C. et al., *Acta Cryst.*, 1954, 7, 159 (rel config)

Kopp, J.F. et al., *Anal. Chem.*, 1973, 45, 1786 (use, Ephedrine)

Milne, J.W.A. et al., *Anal. Chem.*, 1973, 45, 1952 (ms)

Yamasaki, K. et al., *Phytochemistry*, 1973, 12, 2877 (biosynth)

Furlei, R.R. et al., *CA*, 1974, 80, 144 913z (ms)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1975, 5, 289; 1981, 9, 209;

1984, 11, 230; 1989, 14, 159; 1990, 15, 154 (use)

Ghosal, S. et al., *Phytochemistry*, 1975, 14, 830 (isol)

Lovgren, K. et al., *Acta Pharm. Suec.*, 1977, 14, 30 (pmr)

Smith, T.A. et al., *Phytochemistry*, 1977, 16, 9 (occur)

Benzezza, S.A. et al., *Anal. Profiles Drug Subst.*, 1979, 8, 489 (rev, uv, ir, pmr, ms, anal, Pseudoephedrine)

Baudet, M. et al., *Analyst (London)*, 1979, 12, 641 (cmr)

Ali, S.L. et al., *Anal. Profiles Drug Subst.*, 1986, 15, 255 (rev, uv, ir, ms, pmr, anal)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 4, 2321-2325 (use)

Reddy, G.V. et al., *Tet. Lett.*, 2000, 41, 953-954 (synth)

Kim, H.K. et al., *Chem. Pharm. Bull.*, 2003, 51, 1382-1385 (pmr, anal)


Eccles, R. et al., *Br. J. Clin. Pharmacol.*, 2007, 63, 10-14 (use)

Martindale, *The Extra Pharmacopoeia*, 35th edn., Pharmaceutical Press, 2007, 1417

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, EAW000; EAX000; EAY000; POH000; POH250;

EAW500; EAY500; POH500; EAX500

7-(Methylamino)-3H-pyrrolo[2,3-c]isoquinoline-6,9-dione M-387


$C_{12}H_9N_3O_2$ 227.222

Prod. by the marine-derived *Streptomyces* sp. B1848. Dark red solid. λ_{max} 233; 287; 295 (sh); 372; 422 (sh); 481 (sh) (MeOH).

Shaaban, M. et al., *Dissertation*, Univ. of Göttingen, 2004, (isol, pmr, cmr, ms)

8-Methylanibacanine M-388

[151757-09-4]



$C_{19}H_{21}NO_3$ 311.38


Alkaloid from the stem bark of *Aniba*

canelilla (Lauraceae). $[\alpha]_D$ -60 (c, 0.7 in MeOH). λ_{\max} 225 (log ϵ 3.98); 281 (log ϵ 3.61) (MeOH).

Oger, J.-M. *et al.*, *Can. J. Chem.*, 1993, **71**, 1128-1135 (*isol, uv, pmr, cmr, ms, struct*)

1-Methyl-9-azabicyclo[3.3.1]nonan-3-one, 9CI M-389

Euphococcinine
[45977-26-2]



$C_9H_{15}NO$ 153.224

(+)-form [15486-23-4]

Main alkaloid from *Euphorbia atoto* (Euphorbiaceae). Component of the defence secretion of the ladybird *Cryptolaemus montrouzieri* and the Mexican bean beetle *Epilachna varivestis*. Active against gram-negative bacteria. Mp 30° (28°). $[\alpha]_D$ +3 (c, 2.0 in $CHCl_3$). $[\alpha]_D$ +6 (c, 2.0 in MeOH).

Picrate: Mp 240° dec.

N-Me: 1,9-Dimethyl-9-azabicyclo[3.3.1]nonan-3-one, 9CI. **N-Methyleuphococcinine**. 1-Methylgranatan-3-one [219756-74-8]

$C_{10}H_{17}NO$ 167.25

Trace alkaloid from *Picea pungens*. $[\alpha]_D^{23}$ +22 (c, 0.2 in $CHCl_3$).

(-)-form [84026-78-8]

Synthetic. Powder (CH_2Cl_2). Mp 32°. $[\alpha]_D^{20}$ -6.5 (c, 1.80 in MeOH).

(±)-form

Oil. Mp 230° (as picrate). Bp_{0.001} 50°.

Alder, K. *et al.*, *Annalen*, 1959, **620**, 73-87 (*synth*)

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1967, **20**, 561-563 (*isol, ir, pmr, ms*)

Brown, W.V. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1255-1261 (*isol*)

Hill, R.K. *et al.*, *Tetrahedron*, 1982, **38**, 1958-1963 (*synth*)

Gnecco Medina, D.H. *et al.*, *Tet. Lett.*, 1983, **24**, 2099-2102 (*synth*)

Eisner, T. *et al.*, *Experientia*, 1986, **42**, 204-207 (*isol*)

Yue, C. *et al.*, *J.O.C.*, 1992, **57**, 4211-4214 (*synth, abs config*)

Tawara, J.N. *et al.*, *J.O.C.*, 1993, **58**, 4813-4818 (*activity*)


Davison, E.C. *et al.*, *Tet. Lett.*, 1995, **36**, 9047-9050 (*synth*)

Tawara, J.N. *et al.*, *J. Nat. Prod.*, 1999, **62**, 321-323 (*N-Methyleuphococcinine*)

Murahashi, S. *et al.*, *Heterocycles*, 2000, **52**, 557-561 (*synth*)

Mechelke, M.F. *et al.*, *Tet. Lett.*, 2000, **41**, 4339-4342 (*synth*)

8-Methyl-8-azabicyclo[3.2.1]octane-1,3-diol M-390



$C_8H_{15}NO_2$ 157.212

(1R,3R)-form

3-O-Benzoyl: **1-Hydroxytropacocaine**

[156497-23-3]

$C_{15}H_{19}NO_3$ 261.32

Alkaloid from the leaves of *Erythroxylum novogranatense* (2 varieties).

Moore, J.M. *et al.*, *Phytochemistry*, 1994, **36**, 357-360

8-Methyl-8-azabicyclo[3.2.1]octane-3,6-diol, 9CI M-391

3,6-Tropanediol. 3,6-Dihydroxytropane

[41164-06-1]



(3R,6R)-form

$C_8H_{15}NO_2$ 157.212

(1R,3R,6R)-form [7688-72-4]

Alkaloid from *Datura* spp. (Solanaceae). Mp 211°. $[\alpha]_D^{20}$ +24 (c, 2 in EtOH).

3-Ac: **3-Acetoxytropane-6-ol**

[34446-04-3]

$C_{10}H_{17}NO_3$ 199.249

Alkaloid from leaves of *Peripentadenia mearnsii* (Elaeocarpaceae). Needles (Me_2CO). Mp 105-106°. $[\alpha]_D$ +16 (c, 0.31 in $CHCl_3$).

3,6-Di-Ac:

Noncryst. Mp 192-193° (as picrate). $[\alpha]_D$ +17 (c, 2.4 in EtOH).

6-O-(3-Methylbutanoyl): **6-Isovaleroyloxytropane-3-ol**

[152128-83-1]

$C_{13}H_{23}NO_3$ 241.33

Alkaloid from the stem bark *Erythroxylum zambeziacum*.

6-Angeloyl: **6-Angeloyloxytropane-3-ol**

[77123-13-8]

$C_{13}H_{21}NO_3$ 239.314

Alkaloid from *Schizanthus hookeri* (Solanaceae). Oil. Mp 168-169° (as hydrobromide). $[\alpha]_D$ -18 (c, 0.2 in EtOH).

3-Tigloyl, 6-Ac: **6-Acetoxy-3-tigloyloxytropane**

[7688-71-3]

$C_{15}H_{25}NO_4$ 281.351

Alkaloid from leaves of *Datura sanguinea* (Solanaceae). Mp 184-184.5° (as picrate). $[\alpha]_D^{20}$ -11.5 (c, 12.4 in EtOH).

3-Tigloyl, 6-propanoyl: **6-Propanoyloxy-3-tigloyloxytropane**

[54354-59-5]

$C_{16}H_{25}NO_4$ 295.378

Alkaloid from *Datura innoxia* (Solanaceae). Gum. Mp 163° (as picrate). $[\alpha]_D^{23}$ 0.

6-Tigloyl: **6-Tigloyloxytropane-3-ol**

[20399-77-3]

$C_{13}H_{21}NO_3$ 239.314

Alkaloid from *Datura cornigera* leaves and from *Schizanthus hookeri* (Solanaceae). Gum; needles (as hydrobromide). Mp 185° (hydrobromide). $[\alpha]_D^{20}$ -28.1 (c, 1.64 in $CHCl_3$).

Ditigloyl: **3,6-Ditigloyloxytropane**. 3,6-Ditigloyltropanediol

[23517-33-1]

$C_{18}H_{27}NO_4$ 321.416

Alkaloid from *Datura innoxia*, *Datura stramonium*, *Mandragora autumnalis* and *Mandragora* sp. (Solanaceae).

Noncryst.; filamentous needles (as picrate). Mp 150-151° (picrate). $[\alpha]_D$ -21.5 (c, 3.1 in EtOH).

3-O-(3-Methyl-2-butenoyl): **3-Seneciolyloxytropane-6-ol**

[20399-78-4]

$C_{13}H_{21}NO_3$ 239.314

Alkaloid from *Schizanthus hookeri* (Solanaceae). Solid (EtOH aq.). Mp 79-80°. $[\alpha]_D$ -17.6 (c, 0.08 in EtOH).

6-O-(3-Methyl-2-butenoyl): **6-Seneciolyloxytropane-3-ol**

[20399-79-5]

$C_{13}H_{21}NO_3$ 239.314

Alkaloid from the leaves of *Schizanthus litoralis*. Oil. $[\alpha]_D$ -23 (c, 0.31 in EtOH).

3-O-Phenylacetyl: **Tropane-3,6-diol 3-phenylacetate**

[104086-64-8]

$C_{16}H_{21}NO_3$ 275.347

Alkaloid from root bark of *Erythroxylum hypericifolium* (Erythroxylaceae). Cryst. (EtOH). Mp 85°. $[\alpha]_D^{20}$ +4.3 (c, 1.5 in EtOH).

3-O-Phenylacetyl, 6-Ac: **Tropane 3,6-diol 3-phenylacetate 6-acetate**

[97345-82-9]

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from root bark of *Erythroxylum hypericifolium* (Erythroxylaceae). Mp 133° (as picrate).

3-O-(3-Methoxycarbonyl-2E-butenoyl), 6-O-(3-methyl-2-butenoyl): [182015-07-2]

$C_{19}H_{27}NO_6$ 365.425

Alkaloid from the leaves of *Schizanthus litoralis*. Brown oil. $[\alpha]_D^{25}$ -18.7 (c, 0.15 in $CHCl_3$). λ_{\max} 216 (log ϵ 4.09) (EtOH).

3-O-(3-Methoxycarbonyl-2E-butenoyl), 6-O-E-cinnamoyl: [182015-08-3]

$C_{23}H_{27}NO_6$ 413.469

Alkaloid from the leaves of *Schizanthus litoralis*.

3-Benzoyl: **3-Benzoyloxytropane-6-ol**

[99562-18-2]

$C_{15}H_{19}NO_3$ 261.32

Major alkaloid from the leaves of *Erythroxylum sideroxyloides* (Erythroxylaceae). Mp 110° (as picrate).

6-Benzoyl: **3-Hydroxy-6-benzoyloxytropane**

[77096-53-8]

$C_{15}H_{19}NO_3$ 261.32

Alkaloid from *Erythroxylum cuneatum*, *Erythroxylum zambeziacum* and *Knightia strobilina*. Amorph. $[\alpha]_D^{20}$ +8 ($CHCl_3$).

3-O-(3,4,5-Trimethoxybenzoyl): **3-(3,4,5-Trimethoxybenzoyloxy)tropane-6-ol**

[111509-20-7]

$C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Erythroxylum zambe-*

- siacum* (Erythroxylaceae). Rods (EtOH aq.) (as picrate). Mp 214° (picrate). $[\alpha]_D^{20} +3.5$ (c, 1.5 in EtOH).
- 3-O-(3,4,5-Trimethoxybenzoyl), 6-Ac: **Erythrozeylanine A**
C₂₀H₂₇NO₇ 393.436
Alkaloid from *Erythroxylum zeylanicum*. Semisolid. $[\alpha]_D^{25} -22.1$ (c, 0.3 in CHCl₃).
- 3-Cinnamoyl: 3-Cinnamoyloxytropan-6-ol [77062-74-9]
C₁₇H₂₁NO₃ 287.358
Alkaloid from the leaves of *Knightia strobilina* (Proteaceae). Viscous oil. Abs. config. not detd., may belong to the (3S,6S)-form.
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl): **Pervilleine G**
C₂₀H₂₇NO₆ 377.436
Alkaloid from the stem bark of *Erythroxylum pervillei*. Amorph. solid. $[\alpha]_D +24.5$ (c, 0.19 in CHCl₃). λ_{max} 311 (log ϵ 4.42) (CHCl₃).
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl), 6-benzoyl: **Tropane 3,6-diol 3-(3,4,5-trimethoxycinnamate) 6-benzoate**. Tropan-6-yl benzoate-3-yl-3,4,5-trimethoxycinnamate
[58721-83-8]
[60873-17-8, 60894-58-8]
C₂₇H₃₁NO₇ 481.544
Alkaloid from *Erythroxylum monogynum* and *Erythroxylum rotundifolium*. Amorph. solid. Mp 54-55°. $[\alpha]_D +39$ (c, 1 in CHCl₃). Abs. config. not determined. Synth. as racemate.
- 6-O-(3,4,5-Trimethoxy-E-cinnamoyl), 3-O-(3,4,5-trimethoxybenzoyl): **Pervilleine B**
C₃₀H₃₇NO₁₀ 571.623
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 40-42°. $[\alpha]_D -22.5$ (c, 0.25 in CHCl₃). Abs. config. not determined. λ_{max} 302 (log ϵ 4.31) (MeOH).
- 6-O-(3,4,5-Trimethoxy-Z-cinnamoyl), 3-O-(3,4,5-trimethoxybenzoyl): **cis-Pervilleine B**
C₃₀H₃₇NO₁₀ 571.623
Alkaloid from the stem bark of *Erythroxylum pervillei*. Amorph. solid. $[\alpha]_D -25.2$ (c, 0.28 in CHCl₃). λ_{max} 299 (log ϵ 4.16) (CHCl₃).
- 3-O-(4-Hydroxy-3-methyl-2E-butenoyl), 6-angeloyl: 6-Angeloyl-3-(trans-4-hydroxysenecieryl) tropane
C₁₈H₂₇NO₅ 337.415
Alkaloid from *Schizanthus tricolor*. ACE inhibitor.
- 3-O-(4-Hydroxy-3-methyl-2E-butenoyl), 6-O-(3-methyl-2-butenoyl): 3-(trans-4-Hydroxysenecieryl)-6-senecieryl tropane
C₁₈H₂₇NO₅ 337.415
Alkaloid from *Schizanthus tricolor*. ACE inhibitor.
- 3-Ketone: 6-Hydroxy-8-methyl-8-azabicyclo[3.2.1]octan-3-one. 6-Hydroxy-3-tropanone
[5932-53-6]
[110061-26-2]
C₈H₁₃NO₂ 155.196
- Alkaloid from the stem bark
Erythroxylum zambesiacum.
- 3-Ketone, 6-benzoyl: 6-Benzoyloxy-3-tropanone
[133613-86-2]
C₁₅H₁₇NO₃ 259.304
Alkaloid from the stem bark of *Erythroxylum zambesiacum*.
- (1R,3S,6S)-form** [4839-11-6]
Alkaloid in *Anthocercis* spp. (Solanaceae). Mp 212°. $[\alpha]_D^{27} -23.3$ (c, 2 in EtOH).
Di-Ac, hydrobromide: Mp 219-220°.
- 3-O-(2-Methylpropanoyl): 3-Isobutyryloxytropan-6-ol
[78886-99-4]
C₁₂H₂₁NO₃ 227.303
Alkaloid from aerial parts of *Anthocercis albicans* (Solanaceae).
- 3-O-(3-Methylbutanoyl): **Valeroidine**
[490-96-0]
C₁₃H₂₃NO₃ 241.33
Alkaloid from *Datura sanguinea* and *Duboisia myoporoides* (Solanaceae). Mp 85°. $[\alpha]_D^{20} -9$ (c, 5 in EtOH).
- 3-O-(3-Methylbutanoyl); hydrobromide: Mp 170-172°. $[\alpha]_D^{20} +5$ (c, 20 in H₂O).
- 6-O-Benzoyl: Alkaloid from the stems of *Erythroxylum rotundifolium*. Amorph. solid. Mp 155-157°. $[\alpha]_D -11$ (c, 0.7 in CHCl₃). λ_{max} 217 (log ϵ 3.89); 272 (log ϵ 3.41); 294 (log ϵ 3.35) (MeOH).
- 3-O-E-Cinnamoyl, 6-Ac: C₁₉H₂₃NO₄ 329.395
Alkaloid from *Erythroxylum zeylanicum*.
- 3-O-Z-Cinnamoyl, 6-Ac: **Erythrozeylanine C**
C₁₉H₂₃NO₄ 329.395
Alkaloid from *Erythroxylum zeylanicum*. Isol. as a mixt. with the (E)-isomer. Poss. artifact.
- Bis-O-(3,4,5-trimethoxy-E-cinnamoyl): **Pervilleine C**
[392232-10-9]
C₃₂H₃₉NO₁₀ 597.661
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 55-57°. $[\alpha]_D +29$ (c, 0.1 in CHCl₂).
- (1RS,3RS,6RS)-form** [65941-67-5]
Synthetic. Mp 179-180°.
Hydrobromide: Mp 267° dec.
- 3-Ac: [7688-76-8]
C₁₀H₁₇NO₃ 199.249
Alkaloid from *Bellendena montana* (Proteaceae). Noncryst. Mp 182-184° (as picrate). Opt. inactive.
- 3-O-(2-Methylpropanoyl), 6-Ac: **6-Acetoxy-3-isobutyryloxytropane**
[72362-51-7]
[72362-53-9]
C₁₄H₂₃NO₄ 269.34
Alkaloid from *Bellendena montana* (Proteaceae). Mp 145-147° (as picrate). Opt. inactive.
- 6-O-(2-Methylpropanoyl), 3-Ac: **3-Acetoxy-6-isobutyryloxytropane**
[72362-49-3]
[72362-50-6]
C₁₄H₂₃NO₄ 269.34
Alkaloid from *Bellendena montana* (Proteaceae). Noncryst. Mp 113-115° (as picrate). Opt. inactive.
- 3-O-(3-Hydroxy-2-phenylpropanoyl): see 6-Hydroxyhyoscyamine, H-521
- 3-Tigloyl: (\pm)-3-O-Tigloyloxytropan-6 β -ol
[7688-74-6]
C₁₃H₂₁NO₃ 239.314
Alkaloid from *Duboisia suaveolens* and from *Schizanthus hookeri* (stereochem. undetd.) (Solanaceae). Mp 167-171° (as picrate).
- 3-Tigloyl, 6-O-(2-methylbutanoyl): **6-(α -Methylbutyryloxy)-3-tigloyloxytropane**
C₁₈H₂₉NO₄ 323.431
Alkaloid from *Duboisia suaveolens* (Solanaceae). Noncryst. Mp 168-170° (as picrate). No opt. rotn. mentioned, may be opt. active.
- Dibenzoyl: **3,6-Dibenzoyloxytropane**
[117005-28-4]
C₂₂H₂₃NO₄ 365.428
Principal alkaloid from leaves of *Erythroxylum cuneatum* (Erythroxylaceae). Plates (EtOH aq.) (as picrate). Mp 240° (picrate).
- 3-O-(3,4-Dimethoxybenzoyl), 6-Ac: **Convalline**
[169626-32-8]
C₁₉H₂₅NO₆ 363.41
Alkaloid from *Convolvulus subhirsutus*.
- 3-O-(4-Methoxyphenylacetyl): **Physo-chlaine**
[54357-41-4]
C₁₇H₂₃NO₄ 305.373
Alkaloid from *Physochlaine alaica* (Solanaceae). Mp 75-76°. Opt. inactive.
- (1R,3R,6R)-form**
- 6-Tigloyl, 3-formyl: 3-Formyloxy-6-tigloyloxytropane
[855784-81-5]
C₁₄H₂₁NO₄ 267.324
Alkaloid from *Datura stramonium*.
- 3-O-(Phenylacetyl), 6-O-(3,4,5-trimethoxy-E-cinnamoyl): **Pervilleine F**
C₂₈H₃₃NO₇ 495.571
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 95-97°. $[\alpha]_D +12$ (c, 0.12 in CHCl₃). λ_{max} 304 (log ϵ 4.24) (MeOH).
- 3-O-(Phenylacetyl), 6-O-(3,4,5-trimethoxy-Z-cinnamoyl): **cis-Pervilleine F**
C₂₈H₃₃NO₇ 495.571
Alkaloid from the stem bark of *Erythroxylum pervillei*. Amorph. solid. $[\alpha]_D +4$ (c, 0.23 in CHCl₃). λ_{max} 308 (log ϵ 4.18) (CHCl₃).
- 3-O-(3-Hydroxyphenylacetyl), 6-O-(3,4,5-trimethoxy-E-cinnamoyl): **Pervilleine E**
C₂₈H₃₃NO₈ 511.571
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 82-84°. $[\alpha]_D +28$ (c, 0.12 in CHCl₃). λ_{max} 302 (log ϵ 4.25) (MeOH).
- Bis-O-(4-methoxybenzoyl): **Merredissine**
C₂₄H₂₇NO₆ 425.48
Alkaloid from *Merremia dissecta*. Oil. $[\alpha]_D^{20} -8.6$ (c, 0.2 in MeOH).

- 3-O-(3,4,5-Trimethoxybenzoyl), 6-O-benzoyl: **Catuabine B**
[65862-79-5]
C₂₅H₂₉NO₇ 455.507
Alkaloid from *Erythroxylum vacciniifolium*.
- 3-O-(3,4,5-Trimethoxy-Z-cinnamoyl), 6-O-benzoyl:
C₂₇H₃₁NO₇ 481.544
Alkaloid from the stems of *Erythroxylum rotundifolium*. Amorph. solid.
Mp 33-35°. [α]_D +4 (c, 0.17 in CHCl₃). λ_{max} 207 (log ε 4.49); 228 (log ε 4.54); 304 (log ε 3.85) (MeOH).
- 3-Phenylacetyl, 6-propanoyl: **Tropane-3,6-diol 3-phenylacetate 6-propanoate**
[855784-80-4]
C₁₉H₂₅NO₄ 331.411
Alkaloid from *Datura stramonium*.
- 3-O-(1-Methyl-2-pyrrolocarbonyl): **Catuabine I**
C₁₄H₂₀N₂O₃ 264.324
Alkaloid from the bark of *Erythroxylum vacciniifolium*.
- 3-O-(1-Methyl-2-pyrrolocarbonyl), 6-O-(2-pyrrolocarbonyl): **Catuabine D**
C₁₉H₂₃N₃O₄ 357.408
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.
[α]_D -32 (c, 0.58 in EtOH). λ_{max} 267 (log ε 4.38) (EtOH).
- 6-O-(1-Methyl-2-pyrrolocarbonyl): **Catuabine H**
C₁₄H₂₀N₂O₃ 264.324
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.
[α]_D²⁰ -9.6 (c, 0.46 in EtOH). λ_{max} 267 (log ε 4.1) (EtOH).
- 6-O-(1-Methyl-2-pyrrolocarbonyl), 3-O-(3,4,5-trimethoxybenzoyl): **Catuabine A**
[65862-78-4]
C₂₄H₃₀N₂O₇ 458.51
Alkaloid from *Erythroxylum vacciniifolium*.
- 6-O-(1-Methyl-2-pyrrolocarbonyl), 3-O-(4-hydroxy-3,5-dimethoxybenzoyl): **Catuabine F**
C₂₃H₂₈N₂O₇ 444.483
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.
[α]_D -32.8 (c, 0.46 in EtOH). λ_{max} 221 (log ε 4.27); 270 (log ε 4.3); 293 (sh) (log ε 3.85) (EtOH).
- 6-O-(1-Methyl-2-pyrrolocarbonyl), 3-O-(2-pyrrolocarbonyl): **Catuabine C**
[65862-80-8]
C₁₉H₂₃N₃O₄ 357.408
Alkaloid from *Erythroxylum vacciniifolium*.
- Bis-O-(1-methyl-2-pyrrolocarbonyl): **Catuabine E**
C₂₀H₂₅N₃O₄ 371.435
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.
[α]_D -35.4 (c, 0.57 in EtOH). λ_{max} 268 (log ε 4.38) (EtOH).
- Bis-O-(1-methyl-2-pyrrolocarbonyl), N-oxide: **Catuabine E N-oxide**
C₂₀H₂₅N₃O₅ 387.435
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.

[α]_D²⁰ -7.3 (c, 0.33 in EtOH). λ_{max} 268 (log ε 4.43) (EtOH).

(1R,3R,6S)-form

6-O-(1-Methyl-2-pyrrolocarbonyl): **3-Epicatuabine H**. 3αH-Catuabine H
C₁₄H₂₀N₂O₃ 264.324

Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid.
[α]_D²⁰ -8 (c, 0.41 in EtOH). λ_{max} 267 (log ε 3.96) (EtOH).

Fodor, G. *et al.*, *J.C.S.*, 1953, 2341; 1961, 3219; 1965, 6830 (synth, abs config, Valeroidine, Ditioglyoxytropone, tigloyloxy)

Evans, W.C. *et al.*, *J.C.S.*, 1958, 1991; 1963, 4348 (tigloyl esters)

Evans, W.C. *et al.*, *J.C.S. (C)*, 1966, 1621 (Tigloyloxyacetoxytropone)

Mandava, N. *et al.*, *Can. J. Chem.*, 1968, **46**, 2761 (pmr)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1971, **24**, 2399 (acetate)

Evans, W.C. *et al.*, *Phytochemistry*, 1972, **11**, 3293; 1981, **20**, 497 (isol, pmr, ms, esters)

Mirzamatov, R.T. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 415; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 426 (Physochlaine)

Beresford, P.J. *et al.*, *Phytochemistry*, 1974, **13**, 1249; 1975, **14**, 2205

(Propanoyloxytigloyloxytropone, biosynth)
Agar, J.T.H. *et al.*, *J. Pharm. Pharmacol.*, 1975, **27**, 85P (benzoate-trimethoxycinnamate)

Agar, J.T.H. *et al.*, *J.C.S. Perkin 1*, 1976, 1550 (trimethoxycinnamate-benzoate)

Graf, E. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 139-152 (Catuabines A-C)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1979, **32**, 1827 (acetoxisobutyryloxytropans)

Lounasmaa, M. *et al.*, *Phytochemistry*, 1980, **19**, 949 (3-cinnamate, 6-benzoate)

Gambaro, V. *et al.*, *Phytochemistry*, 1980, **19**, 2007; 1983, **22**, 1838 (tiglate, angelate, seneciolate)

Al-Said, M.S. *et al.*, *J.C.S. Perkin 1*, 1986, 957 (phenylacetates)

Al-Said, M.S. *et al.*, *Phytochemistry*, 1986, **25**, 851 (3α-Benzoyloxytropanol)

El-Iman, Y.M.A. *et al.*, *Phytochemistry*, 1987, **26**, 2385-2389; 1988, **27**, 2181-2184 (trimethoxybenzoate, 6-benzoate)

Aripova, S.F. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 88-90; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 74-75 (Convolacine)

Christen, P. *et al.*, *Phytochemistry*, 1993, **34**, 1147-1151 (*Erythroxylum zambesiacum* tropans)

Muñoz, O. *et al.*, *Phytochemistry*, 1996, **43**, 709-713 (*Schizanthus litoralis* tropans)

Bringmann, G. *et al.*, *Phytochemistry*, 2000, **53**, 409-416 (*Erythrozylanines*)

Silva, G.L. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1514-1520 (Pervilleines)

Chavez, D. *et al.*, *J. Nat. Prod.*, 2002, **65**, 606-610 (*Erythroxylum rotundifolium* esters)

Zanolari, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 497-502; 2005, **68**, 1153-1158 (Catuabines)

Kletter, C. *et al.*, *Planta Med.*, 2004, **70**, 993-1000 (Catuabine D)

Doncheva, T. *et al.*, *CA*, 2005, **143**, 74802f (*Datura stramonium* constits)

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (Merredissine)

Chin, Y.-W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 414-417 (Pervilleine G, cis-Pervilleines B,F)


Humam, M. *et al.*, *Planta Med.*, 2006, **72**, 969 (*Schizanthus tricolor* constits)

Humam, M. *et al.*, *Chirality*, 2008, **20**, 20-25 (3-4-Hydroxysenecioid-6-senecioidtropone, cd, abs config)

Kulkarni, K. *et al.*, *Synlett*, 2008, 2209-2212 (Pervilleine C, synth, abs config)

8-Methyl-8-azabicyclo[3.2.1]octane-3,6,7-triol, 9CI

3,6,7-Tropanetriol. 3,6,7-Trihydroxytropone



(1R*,3S*,6R*,7R*)-form

C₈H₁₅NO₃ 173.211

The alkaloids are named here as given in the lit., but the numbering of positions 6 and 7, which is irregular in the lit., has been corrected. Mp 169-170° (hydrate).

(1R,3S,6R,7R)-form

(3α,6α,7β)-form

Chiral stereoisomer.

3-O-(3,4,5-Trimethoxy-E-cinnamoyl), 6-O-(3,4,5-trimethoxybenzoyl): **Erythrorotundine**

[307334-87-8]

C₃₀H₃₇NO₁₁ 587.622

Alkaloid from *Erythroxylum rotundifolium*. Cryst. (EtOH). Mp 176°. [α]_D²⁵ -20.8 (c, 0.1 in EtOH).

(1R,3S,6S,7R)-form

(3α,6β,7β)-form. Teloidine

[575-62-2]

Mp 169-172° (hydrate). A meso-stereoisomer but 6-mono- and 3,6-disubst. derivs. are chiral. There is currently no data on abs. configs.

Hydrochloride:

Prisms (EtOH). Mp 300°.

Hydrobromide: Mp 295°.

3-O-(2-Methylbutanoyl): **3-(2-Methylbutanoyloxy)tropane-6,7-diol**

[152187-24-1]

C₁₃H₂₃NO₄ 257.329

Alkaloid from *Erythroxylum monogynum* and *Erythroxylum zambesiacum*.

3-O-Tigloyl: **Meteloidine**

[526-13-6]

C₁₃H₂₁NO₄ 255.313

Alkaloid from *Datura meteloides*, *Datura suaveolens*, *Datura ferox* and many other *Datura* spp. and *Erythroxylum australe* (Solanaceae, Erythroxyllaceae). Mp 141-142°. Meso-. Shows no marked physiological activity.

3-Tigloyl, 6-propanoyl: **7-Propanoyloxy-3-tigloyloxytropone-6-ol**

C₁₆H₂₅NO₅ 311.377

Alkaloid from hairy root cultures of *Datura stramonium*.

3-O-Tigloyl, 6-O-(3-methylbutanoyl): **7-Isovaleryloxy-3-tigloyloxytropone-6-ol**

[21030-84-2]

[21030-91-1]

C₁₈H₂₉NO₅ 339.431

Alkaloid from *Datura sanguinea* (Solanaceae). Prisms (Et₂O/EtOH) (as hydrobromide). Mp 182-883° (hydrobromide).

6-O-Tigloyl: **Isometeloidine**. 7-Tigloyloxy-

tropane-3,6-diol

[39972-69-5]

C₁₃H₂₁NO₄ 255.313

- Alkaloid from *Datura candida* and *Datura suaveolens* (Solanaceae). Mp 247° (as hydrobromide).
- 3,6-Di-O-tigloyl: 3,6-Ditigloyloxytropan-7-ol**
[7159-86-6]
C₁₈H₂₇NO₅ 337.415
Alkaloid present in *Datura suaveolens*, *Datura cornigera*, *Datura meteloides* and *Cyphomandra betacea* (tree tomato) (Solanaceae). Mp 215° (as hydrobromide).
- 3-O-Benzoyl: 6,7-Dihydroxytropan-3-yl benzoate. 3-O-Benzoylteloidine**
[66759-80-6]
C₁₅H₁₉NO₄ 277.319
Alkaloid from the roots of *Erythroxylum australe* (Erythroxylaceae). Mp 290° dec. (as hydrobromide). *Meso*-.
- 6-O-Benzoyl: 7-Benzoyloxytropane-3,6-diol**
[111509-23-0]
C₁₅H₁₉NO₄ 277.319
Alkaloid from the root bark of *Erythroxylum zambeziacum* (Erythroxylaceae). Plates (EtOH/Et₂O) (as picrate). Mp 213° (picrate).
- 3,7-Dibenzoyl: Alaternifoline. 3,7-Dibenzoyloxytropan-6-ol**
[307334-86-7]
C₂₂H₂₃NO₅ 381.427
Alkaloid from *Erythroxylon alaternifolium*. Cryst. (EtOH). Mp 220°. [α]_D²⁵ +10 (c, 0.1 in EtOH).
- 3-O-(3,4,5-Trimethoxybenzoyl): Tropane-3,6,7-triol 3-(3,4,5-trimethoxybenzoate)**
[60820-76-0]
C₁₈H₂₅NO₇ 367.398
Alkaloid from the root bark of *Erythroxylum monogynum* and *Erythroxylum zambeziacum* (Erythroxylaceae). Cryst. solid. Mp 112°. *Meso*-.
- 3-O-Phenylacetyl: Tropane-3,6,7-triol 3-phenylacetate**
[104086-63-7]
C₁₆H₂₁NO₄ 291.346
Minor alkaloid from the root bark of *Erythroxylum hypericifolium* (Erythroxylaceae). Cryst. *Meso*-.
- 3-O-(3-Hydroxyphenylacetyl): Catuabine G**
C₁₆H₂₁NO₅ 307.346
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. *Meso*-. λ_{max} 218 (log ε 4.35); 276 (log ε 3.85) (EtOH).
- 3-O-(2-Hydroxy-3-phenylpropanoyl) (R-): 6,7-Dihydroxylittorine**
[62083-54-9]
C₁₇H₂₃NO₅ 321.372
Minor alkaloid from leaves of *Datura candida* (Solanaceae). Feathery needles (Et₂O/EtOH) (as hydrobromide). Mp 184° (hydrobromide). [α]_D²⁵ -6 (c, 5.0 in H₂O).
- 6-O-(3ξ-Hydroxy-2ξ-methyl-3-phenylpropanoyl): 6-(3-Hydroxy-2-methyl-3-phenylpropanoyloxy)-3,7-tropanediol**
C₁₈H₂₅NO₅ 335.399
Alkaloid from the leaves of *Erythroxylum moonii*. Light yellow powder.
- [α]_D²⁵ -27.4 (c, 0.17 in CHCl₃). λ_{max} 216 (log ε 4.1); 273 (log ε 2.38) (MeOH).
- 3-O-Cinnamoyl: 3-Cinnamoyloxy-6,7-tropanediol**
[117005-30-8]
C₁₇H₂₁NO₄ 303.357
Alkaloid from root bark of *Erythroxylum australe* (Erythroxylaceae). Stereochem. of cinnamoyl residue not determined. Detected by ms.
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl): Pervilleine H**
C₂₀H₂₇NO₇ 393.436
Alkaloid from the stem bark of *Erythroxylum pervillei*. Amorph. solid. λ_{max} 312 (log ε 4.15) (CHCl₃).
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl), 6-O-benzoyl: 7-Benzoyloxy-3-(3,4,5-trimethoxycinnamoyloxy)tropan-6-ol**
[111509-24-1]
C₂₇H₃₁NO₈ 497.544
Alkaloid from *Erythroxylum rotundifolium* and *Erythroxylum zambeziacum* (Erythroxylaceae). Needles. Mp 184-187°. [α]_D +16 (c, in CHCl₃).
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl), 6-O-benzoyl, 7-Ac: 6-Acetoxy-7-benzoyloxy-3-(3,4,5-trimethoxycinnamoyloxy)tropane**
[111534-76-0]
C₂₉H₃₃NO₉ 539.581
Alkaloid from *Erythroxylum rotundifolium* and *Erythroxylum zambeziacum*. Cryst. Mp 56-58°. [α]_D +2.1 (c, 0.9 in CHCl₃).
- 3-O-(3,4,5-Trimethoxy-E-cinnamoyl), 6-O-(3,4,5-trimethoxybenzoyl): 7-(3,4,5-Trimethoxycinnamoyloxy)tropan-6-ol**
C₃₀H₃₇NO₁₁ 587.622
Alkaloid from the stems of *Erythroxylum rotundifolium*. Amorph. solid. Mp 179-180°. [α]_D +9 (c, 0.1 in CHCl₃). λ_{max} 214 (log ε 4.65); 233 (sh) (log ε 4.3); 272 (log ε 4.18); 306 (log ε 4.3) (MeOH).
- 6-O-(3,4,5-Trimethoxy-E-cinnamoyl), 3-O-(3,4,5-trimethoxybenzoyl): Pervilleine A**
C₃₀H₃₇NO₁₁ 587.622
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. [α]_D -0.6 (c, 0.18 in CHCl₃). Softens at 46-48°. λ_{max} 302 (log ε 4.39) (MeOH).
- 6-O-(3,4,5-Trimethoxy-E-cinnamoyl), 3-O-(3,4,5-trimethoxybenzoyl), N-oxide: Pervilleine A N-oxide**
C₃₀H₃₇NO₁₂ 603.622
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 118-121°. [α]_D +1.5 (c, 0.28 in CHCl₃). Softens at 114°. λ_{max} 303 (log ε 4.24) (MeOH).
- 3,6-Bis-O-(3,4,5-trimethoxy-E-cinnamoyl): Pervilleine D**
C₃₂H₃₉NO₁₁ 613.66
Alkaloid from the roots of *Erythroxylum pervillei*. Amorph. solid. Mp 59-61°. [α]_D +9 (c, 0.1 in CHCl₃). λ_{max} 305 (log ε 4.5) (MeOH).
- 3-O-(1-Methyl-2-pyrrolicarboxyl): 7β-Hydroxycatuabine I**
C₁₄H₂₀N₂O₄ 280.323
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. λ_{max} 267 (log ε 4.13) (EtOH).
- 3-O-(1-Methyl-2-pyrrolicarboxyl), 6-O-(2-pyrrolicarboxyl): 7β-Hydroxycatuabine D**
C₁₉H₂₃N₃O₅ 373.408
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D -2.8 (c, 0.4 in EtOH). λ_{max} 267 (log ε 3.92) (EtOH).
- 6-O-(1-Methyl-2-pyrrolicarboxyl): 7β-Hydroxycatuabine H**
C₁₄H₂₀N₂O₄ 280.323
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D²⁰ -1.5 (c, 0.53 in EtOH). λ_{max} 267 (log ε 4.04) (EtOH).
- 6-O-(1-Methyl-2-pyrrolicarboxyl), 3-O-(4-hydroxy-3,5-dimethoxybenzoyl): 7β-Hydroxycatuabine F**
C₂₃H₂₈N₂O₈ 460.483
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D -2.6 (c, 0.44 in EtOH). λ_{max} 221 (log ε 4.16); 269 (log ε 4.13); 293 (sh) (log ε 3.71) (EtOH).
- 3,6-Bis-O-(1-methyl-2-pyrrolicarboxyl): 7β-Hydroxycatuabine E**
C₂₀H₂₅N₃O₅ 387.435
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D +0.8 (c, 0.57 in EtOH). λ_{max} 267 (log ε 4.41) (EtOH).
- 3,6-Bis-O-(1-methyl-2-pyrrolicarboxyl), 7-Ac: 7β-Acetoxycatuabine E**
C₂₂H₂₇N₃O₆ 429.472
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D -36.8 (c, 0.56 in EtOH). λ_{max} 268 (log ε 4.45) (EtOH).
- (1R*,3S*,6S*,7S*)-form**
- 3-O-Tigloyl, 6-O-propanoyl:** [821787-78-4]
C₁₆H₂₅NO₅ 311.377
Alkaloid from *Datura innoxia* and *Datura stramonium*.
- 6-O-(1-Methyl-2-pyrrolicarboxyl): 7α-Hydroxycatuabine H**
C₁₄H₂₀N₂O₄ 280.323
Alkaloid from the bark of *Erythroxylum vacciniifolium*. Amorph. solid. [α]_D²⁰ -40.7 (c, 0.57 in EtOH). λ_{max} 268 (log ε 4.11) (EtOH).
- (1ξ,3ξ,6ξ,7ξ)-form**
- 3-O-(2-Hydroxy-3-phenylpropanoyl), 6-tigloyl, N-de-Me: 6-Hydroxy-7-tigloyloxyntropan-3-yl 2-hydroxy-3-phenylpropionate**
[66759-84-0]
C₂₁H₂₇NO₆ 389.447
Alkaloid from aerial parts of *Erythroxylum australe*. Mp 225° (as hydrobromide). [α]_D²⁵ +5 (c, 0.03 in H₂O). Tentative struct.
- [21030-92-2, 62058-54-2]
Schöpf, C. et al., *Annalen*, 1947, **558**, 109 (synth)
Sheehan, J.C. et al., *J.A.C.S.*, 1952, **74**, 3825 (synth)

- Zeile, K. et al., *Z. Naturforsch., B*, 1957, **12**, 661 (synth, Meteloidine)
- Evans, W.C. et al., *J.C.S. (C)*, 1968, 2775-2778 (Isovaleryloxytigloyloxytropone)
- Evans, W.C. et al., *Phytochemistry*, 1972, **11**, 3293 (derivs)
- Beresford, P.J. et al., *Phytochemistry*, 1975, **14**, 2205 (biosynth)
- Griffin, W.J. et al., *Aust. J. Chem.*, 1976, **29**, 2329-2332; 1978, **31**, 1161-1165 (Teloidine, Isometeloidine, Benzoyteloidine, Dihydroxytelitorine, Hydroxytigloyloxynortropanyl 2-hydroxy-3-phenylpropionate)
- Agar, J.T.H. et al., *J.C.S. Perkin 1*, 1976, 1550 (isol, pmr, ms, trimethoxybenzoate)
- Al-Said, M.S. et al., *J.C.S. Perkin 1*, 1986, 957-959 (phenylacetate)
- El-Imam, Y.M.A. et al., *Phytochemistry*, 1987, **26**, 2385-2389; 1988, **27**, 2181-2184 (3-cinnamoyl, 3-benzoyl, 6-benzoyl, 3-tigloyl)
- Christen, P. et al., *Phytochemistry*, 1993, **34**, 1147-1151 (2-methylbutanoyl)
- Payo-Hill, A.L. et al., *Phytochemistry*, 2000, **54**, 927-932 (Alaternifoline, Erythrorotundine)
- Silva, G.L. et al., *J. Nat. Prod.*, 2001, **64**, 1515-1520 (Pervilleines, 3-phenylacetate)
- Chavez, D. et al., *J. Nat. Prod.*, 2002, **65**, 606-610 (Erythroxyllum rotundifolium ester)
- Khattak, K.F. et al., *J. Nat. Prod.*, 2002, **65**, 929-931 (3-hydroxy-2-methyl-3-phenylpropanoate)
- Zanolari, B. et al., *J. Nat. Prod.*, 2003, **66**, 497-502 (Catuabines)
- Berkov, S. et al., *Z. Naturforsch., C*, 2003, **58**, 42-45 (7-Propanoyl-3-tigloyltropan-6-ol)
- Kletter, C. et al., *Planta Med.*, 2004, **70**, 993-1000 (7-Hydroxycataubine E)
- Berkov, S. et al., *Z. Naturforsch., C*, 2004, **59**, 184-186 (3-tigloyl 6-propanoyl)
- Zanolari, B. et al., *J. Nat. Prod.*, 2005, **68**, 1153-1158 (7-Hydroxycataubines H,I)
- Chin, Y.-W. et al., *J. Nat. Prod.*, 2006, **69**, 414-417 (Pervilleine H)

8-Methyl-8-azabicyclo[3.2.1]octan-3-ol, 9CI M-393
3-Tropanol. 3-Hydroxytropone



$C_8H_{15}NO$ 141.213

(1R,3R)-form

endo-form. **Tropine**. *Tropeine*
[120-29-6]

Alkaloid from *Atropa belladonna*, *Hyoscyamus niger*, *Datura* spp. and several other genera, nearly in all the Solanaceae; from hydrolysis of ester alkaloids such as Tropane tropate, T-632. Mp 63°. Bp 229°. pK_a 10.33 (25°).

▶ Highly toxic. YM3875000

Picrate: Mp 290-295° (275°).

Ac: **Tropine acetate**

[3423-27-6]

$C_{10}H_{17}NO_2$ 183.25

Alkaloid from *Duboisia myoporoides*, *Datura sanguinea* and *Bruguiera* spp. (Solanaceae, Rhizophoraceae). Bp 235-237°.

O-Propanoyl: **Tropine propionate**

$C_{11}H_{19}NO_2$ 197.277

Alkaloid from *Bruguiera sexangula* and *Bruguiera exaristata* (Rhizophor-

aceae). Isol. by glc and identified by glc and ms comparison with an authentic sample.

O-(2-Methylpropanoyl): **Butropine**. *Tropine isobutyrate*

[495-80-7]

$C_{12}H_{21}NO_2$ 211.303

Alkaloid from *Duboisia leichhardtii* (Solanaceae). Oil; prisms (as hydrobromide). Mp 242° (hydrobromide). n_D^{15} 1.4726. Confused in one review with Poroidine in A-1597.

O-(2-Phenylpropanoyl): see Apotropane, A-1359

O-Butanoyl: **3-Butyryloxytropone**. *Tropine butyrate*

[19038-34-7]

$C_{12}H_{21}NO_2$ 211.303

Alkaloid from a ssp. of *Anthocercis albicans*, also detected in *Bruguiera sexangula* and *Bruguiera exaristata* (Solanaceae, Rhizophoraceae). Mp 194° (as picrate).

O-Nonanoyl: **Tropine nonanoate**

[146018-90-8]

$C_{17}H_{31}NO_2$ 281.437

Alkaloid from leaves of *Duboisia myoporoides* (Solanaceae).

O-(S-2-Methylbutanoyl): **Valtropine**

[495-82-9]

$C_{13}H_{23}NO_2$ 225.33

Alkaloid from *Duboisia leichhardtii* and other *Duboisia* spp. (Solanaceae). Oil. Mp 212° (as hydrobromide). $[\alpha]_D^{20}$ +8.3 (neat). n_D^{15} 1.4725. Confused in one review with Isoporoidine in A-1597.

O-(3-Methylbutanoyl): **Tropan-3-yl isovalerate**. *Tropine isovalerate*

[23768-69-6]

$C_{13}H_{23}NO_2$ 225.33

Alkaloid from the root-bark of *Erythroxyllum deckindtii*, also detected (probably) in *Bruguiera sexangula* and *Bruguiera exaristata* (Erythroxyllaceae, Rhizophoraceae). Plates (EtOH aq.) (as picrate). Mp 230° (picrate).

O-(3-Methyl-2-butenoyl): **Tropanyl seneciolate**. *3-Seneciolyloxytropone*

[81943-54-6]

$C_{13}H_{21}NO_2$ 223.314

Alkaloid from aerial parts of *Schizanthus hookeri* and *Schizanthus grahamii* (Solanaceae). Antihypothermic agent of potential utility against parkinsonism. Oil.

O-Tigloyl: see 3-Tigloyloxytropone, T-404

O-(2-Hydroxy-3-phenylpropanoyl): see Littorine, L-216

O-(3-Hydroxy-2-phenylpropanoyl): see Tropane tropate, T-632

O-(Phenylacetyl): **Tropanyl phenylacetate**

[1690-22-8]

$C_{16}H_{21}NO_2$ 259.347

Alkaloid from the root-bark of *Erythroxyllum deckindtii* (Erythroxyllaceae). Not obt. pure; picrate obt. as 1:1 mixt. with the picrate of the isovalerate.

O-(3-Hydroxyphenylacetyl): **Tropan-3-yl**

3-hydroxyphenylacetate

[104086-65-9]

$C_{16}H_{21}NO_3$ 275.347

Alkaloid from the root bark of *Erythroxyllum hypericifolium* (Erythroxyllaceae). Mp 157° (as picrate).

O-(4-Hydroxyphenylacetyl): **3-(4-Hydroxyphenylacetoxyl)tropane**

$C_{16}H_{21}NO_3$ 275.347

Alkaloid from the root bark of *Erythroxyllum australe* (Erythroxyllaceae).

O-Benzoyl: **Benzoyltropein**. *Tropine benzoate*

[19145-60-9]

$C_{15}H_{19}NO_2$ 245.321

Alkaloid from *Erythroxyllum coca*, *Crossostylis sebertii*, *Bruguiera sexangula* and *Bruguiera exaristata* (Erythroxyllaceae, Rhizophoraceae). Mp 41-42° (anhyd.) Mp 56° (dihydrate).

O-(3-Hydroxybenzoyl): m-Hydroxybenzoyltropine. 3-(m-Hydroxybenzoyloxy)tropane. **Cochlearine**

[52418-07-2]

$C_{15}H_{19}NO_3$ 261.32

Alkaloid from *Cochlearia arctica* (Brassicaceae). Mp 235-236°.

O-(3-Hydroxybenzoyl), hydrochloride: Cryst. (EtOH aq.). Mp 307° dec.

O-(4-Hydroxybenzoyl): **3 α -(4-Hydroxybenzoyloxy)tropane**

$C_{15}H_{19}NO_3$ 261.32

Alkaloid from *Merremia* spp.

O-(4-Methoxybenzoyl): **Datumetine**

[67078-20-0]

$C_{16}H_{21}NO_3$ 275.347

Alkaloid from *Datura metel* and *Merremia dissecta*. Prismatic plates (CHCl₃/Et₂O). Mp 84°.

O-(3,4-Dimethoxybenzoyl): see Convolvine, C-633

O-E-Cinnamoyl: **Tropanyl trans-cinnamate**

[35721-92-7]

$C_{17}H_{21}NO_2$ 271.358

Alkaloid from *Crossostylis sebertii* and *Erythroxyllum hypericifolium*.

O-Z-Cinnamoyl: **Tropanyl cis-cinnamate**

$C_{17}H_{21}NO_2$ 271.358

Alkaloid from the leaves of *Erythroxyllum moonii*. Amorph. powder. λ_{max} 275 (log ϵ 4.36) (MeOH).

O-(4-Hydroxy-3-methoxycinnamoyl):

Tropanyl 3-hydroxy-4-methoxycinnamate. *Tropan-3-yl ferulate*

[86702-58-1]

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from *Crossostylis multiflora* (Rhizophoraceae).

O-(3,4,5-Trimethoxycinnamoyl): **Tropine 3,4,5-trimethoxycinnamate**

[27303-60-2]

$C_{20}H_{27}NO_5$ 361.437

Alkaloid from *Erythroxyllum ellipticum* leaves and *Erythroxyllum monogynum* root bark (Erythroxyllaceae). Prisms (Me₂CO). Mp 165-166°.

O-(2-Furoyl): **Tropanyl furoate**

[72594-46-8]

[72594-47-9]

- C₁₃H₁₇NO₃ 235.282
Alkaloid from the root bark of *Erythroxylum deckindtii* (Erythroxylaceae). Mp 66°.
- O-(3-Pyridinecarbonyl): **3 α -Nicotinoyloxypipropene**
C₁₄H₁₈N₂O₂ 246.308
Alkaloid from *Cochlearia* spp. and *Merremia vitifolia*.
- O-(1,2-Thiolane-3-carboxylate): see Brugine, B-369
- (1R,3SR)-form**
exo-form. Pseudotropine. ψ -3-Tropanol. **ψ -Tropine** [135-97-7]
Alkaloid from *Datura innoxia*, several other *Datura* spp. and from several other genera, nearly all in the Solanaceae. Mp 108-109°. Bp 240-241°. pK_a 9.86 (25°).
▶YM3875010
Hydrochloride: Mp 280-282° dec. (sinters at 250°).
- Ac: **3 α -Acetoxypipropene** [3423-26-5]
C₁₀H₁₇NO₂ 183.25
Alkaloid from *Datura sanguinea* leaves and stems (Solanaceae). Mp 217° (as picrate).
- O-Tigloyl: see 3-Tigloyloxypipropene, T-404
- O-Benzoyl: **Tropacocaine** [537-26-8]
C₁₅H₁₉NO₂ 245.321
Alkaloid from *Erythroxylum coca*, other *Erythroxylum* spp., *Babylonia japonica* and *Peripentadenia mearsii* (Erythroxylaceae, Elaeocarpaceae). Antinicotinic and mydriatic agent, neurotoxic poison, sodium ion influx blocker. Mp 49°. pK_a 9.88 (15°).
▶Neurotoxin. LD₅₀ (mus, ivn) 69 mg/kg. YM4200000
O-Benzoyl; hydrochloride: [637-23-0]
Mp 283° dec. λ_{\max} 220 ; 282 (MeOH) (Berdy).
▶YM4201000
O-Benzoyl, picrate: Mp 243-245°.
- O-(4-Hydroxybenzoyl): **3 β -(4-Hydroxybenzoyloxypipropene**
C₁₅H₁₉NO₃ 261.32
Alkaloid from *Merremia* spp.
- O-(4-Hydroxy-3-methoxybenzoyl): **Concneorine**. 3 β -Vanillyloxypipropene
C₁₆H₂₁NO₄ 291.346
Alkaloid from *Convolvulus concneorum* and *Merremia* spp.
- O-E-Cinnamoyl:
C₁₇H₂₁NO₂ 271.358
Alkaloid from *Erythroxylum hypericifolium* and *Erythroxylum zeylanicum*. Cryst. (as picrate). Mp 215° (picrate).
- O-Z-Cinnamoyl: **Erythrozeylanine B**
C₁₇H₂₁NO₂ 271.358
Alkaloid from *Erythroxylum zeylanicum*. Isol. as a mixt. with (E)-isomer. Possible artifact.
- O-(2S,3-Dihydroxy-2-phenylpropanoyl): **3-(2-Hydroxytropoyloxypipropene**
C₁₇H₂₃NO₄ 305.373
Alkaloid from the seeds of *Datura stramonium*. Config. not confirmed.

- O-(3-Hydroxy-2-methylbutanoyl): **Astrimalvine B** [953384-48-0]
C₁₃H₂₃NO₃ 241.33
Alkaloid from the root bark of *Astrimopoea malvacea*. Yellow oil.
- O-(2-Methyl-3-tigloyloxypipropene), N-oxide: **Astrimalvine A N-oxide** [953384-47-9]
C₁₈H₂₉NO₅ 339.431
Alkaloid from the root bark of *Astrimopoea malvacea*. Yellow oil.
- Willstätter, R. et al., *Ber.*, 1901, **34**, 3165; 1902, **35**, 1870 (synth)
Hardegger, E. et al., *Helv. Chim. Acta*, 1953, **36**, 1186 (config)
Leete, E. et al., *Can. J. Chem.*, 1954, **32**, 1116 (biosynth)
Rosenblum, E.I. et al., *J. Pharm. Pharmacol.*, 1954, **6**, 410 (*Butropine*, *Valtropine*)
Evans, W.C. et al., *J.C.S.*, 1959, 1406 (*isol*)
Platonova, T.F. et al., *CA*, 1964, **60**, 8350 (*Cochlearine*)
Evans, W.C. et al., *J.C.S. (C)*, 1966, 1621 (*3 α -Acetoxypipropene*)
Schenk, H. et al., *Acta Cryst.*, 1967, **23**, 423 (*cryst struct*)
Loder, J.W. et al., *Aust. J. Chem.*, 1969, **22**, 1271 (*isol*, *esters*)
Johns, S.R. et al., *Aust. J. Chem.*, 1970, **23**, 421 (*trimethoxycinnamate*)
Chappell, G.S. et al., *J. Pharm. Sci.*, 1973, **62**, 414 (*pmr*)
Libisch, H.W. et al., *Z. Chem.*, 1973, **13**, 372 (*Cochlearine*, *biosynth*)
Agar, J.T.H. et al., *J.C.S. Perkin 1*, 1976, 1551 (*trimethoxycinnamate*)
Hamor, T.A. et al., *J.C.S. Perkin 2*, 1976, 1359-1363 (*O-benzoyl*, *cryst struct*)
Hanish, P. et al., *J.C.S. Perkin 2*, 1977, 1202 (*cmr*, *conformn*)
Grützmacher, H.-F. et al., *Chem. Ber.*, 1978, **111**, 1962 (*ms*)
Hayakawa, Y. et al., *J.A.C.S.*, 1978, **100**, 1786 (*synth*)
Tufariello, J.J. et al., *J.A.C.S.*, 1979, **101**, 2435 (*synth*)
Sharova, S.F. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 487 (*Phyllalbine*)
Evans, W.C. et al., *Phytochemistry*, 1981, **20**, 497 (*butyrate*)
Gambaro, V. et al., *Bol. Soc. Chil. Quim.*, 1982, **27**, 296; *CA*, **97**, 3737u (*seneciolate*)
Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 422; 12406
Gnecco Medina, D.H. et al., *J. Nat. Prod.*, 1983, **46**, 398 (*cinnamate*, *3-hydroxy-4-methoxycinnamate*)
Iida, H. et al., *J.O.C.*, 1985, **50**, 1818 (*synth*)
Siddiqui, S. et al., *J. Nat. Prod.*, 1986, **49**, 511 (*Datumetine*)
Al-Said, M.S. et al., *J.C.S. Perkin 1*, 1986, 957 (*3-hydroxyphenylacetate*)
Lounasmaa, M. et al., *Alkaloids (Academic Press)*, 1988, **33**, 1 (*synth*, *rev*)
El-Imam, Y.M.A. et al., *Phytochemistry*, 1988, **27**, 2181 (*4-hydroxyphenylacetate*)
Al-Said, M.S. et al., *Phytochemistry*, 1989, **28**, 3211 (*E-cinnamoyl*)
Shukla, Y.N. et al., *Phytochemistry*, 1992, **31**, 4389 (*Tropine nonanoate*)
Bringmann, G. et al., *Phytochemistry*, 2000, **53**, 409-416 (*Erythrozeylanine B*)
Khattak, K.F. et al., *J. Nat. Prod.*, 2002, **65**, 929-931 (*Tropanyl cis-cinnamate*)
Phillipov, S. et al., *Z. Naturforsch., C*, 2002, **57**, 559-561 (*3-(2-Hydroxytropoyloxypipropene*)
Jenett-Siems, K. et al., *Phytochemistry*, 2005, **66**, 1448-1464 (*Datumetine*, *Phyllalbine*, *Concneorine*, *Merremia constits*)


- Ott, S.C. et al., *Z. Naturforsch., B*, 2007, **62**, 285-288 (*Astrimalvines*)
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TNS200

8-Methyl-8-azabicyclo[3.2.1]octan-3-one, 9CI M-394
Tropinone. 3-Tropanone
[532-24-1]



- C₈H₁₃NO 139.197
Alkaloid from *Nicandra physaloides* (Solanaceae). Needles (petrol). Mp 42°. Bp₁₃ 103-104°. Co-occurs with Hygrine, H-759.
Hydrochloride: [4827-85-4]
Mp 188-0° 188-189° dec.
▶LD₅₀ (mus, ipr) 430 mg/kg. YM6235000
Picrate: Mp 217-218°.
- N-Oxide:
C₈H₁₃NO₂ 155.196
Mp 217-218°.
- Oxime:
C₈H₁₄N₂O 154.211
Mp 114°. Chiral, has been resolved.
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 725A (*nmr*)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 462D (*ir*)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 547D (*ir*)
Robinson, R. et al., *J.C.S.*, 1917, **111**, 706 (*synth*)
Romeike, A. et al., *Naturwissenschaften*, 1966, **53**, 82 (*isol*)
Singh, H. et al., *Tet. Lett.*, 1966, 3243; 6502 (*resoln*, *oxime*)
Ohashi, M. et al., *Chem. Comm.*, 1971, 34 (*pmr*)
Halpern, A.M. et al., *J.A.C.S.*, 1976, **98**, 3242 (*uv*)
Katritzky, A.R. et al., *Org. Magn. Reson.*, 1979, **12**, 357 (*cmr*, *conformn*)
Lounasmaa, M. et al., *Alkaloids (Academic Press)*, 1988, **33**, 1 (*rev*)

8-Methyl-8-azabicyclo[3.2.1]oct-2-ene-2-carboxylic acid M-395
Anhydroecgonine. *Ecgonidine*



- C₉H₁₃NO₂ 167.207
Topical anaesthetic. pK_{a1} 3.8; pK_{a2} 9.8.
- (1R)-form**
(-)-form
[484-93-5]
Obt. semisynthetically from Cocaine, C-537. Cryst. (MeOH/EtOH). Mp 235° dec. $[\alpha]_D^{25}$ -84.6.
Hydrochloride: [74242-55-0]
Needles (EtOH) or tablets (H₂O). Mp 240-241°. $[\alpha]_D^{25}$ -62.7 (H₂O).
Me ester: **Anhydroecgonine methyl ester** [43021-26-7]
C₁₀H₁₅NO₂ 181.234

Alkaloid from *Erythroxylum coca* and *Erythroxylum emarginatum*. Amorph. powder. Bp₇ 107°. [α]_D²⁵ -23 (c, 1.1 in CHCl₃).

Me ester, N-oxide: **Anhydroecgonine methyl ester N-oxide**

C₁₀H₁₅NO₃ 197.233

Alkaloid from the bark of *Erythroxylum emarginatum*. Amorph. powder. [α]_D²⁵ +36 (c, 0.1 in CHCl₃).

Et ester: [73045-45-1]

C₁₁H₁₇NO₂ 195.261

Sol. cold H₂O, spar. sol. hot. Bp₁₆ 136-139°.

(1S)-form

(+)-form

Cryst. + 1H₂O (MeOH or MeOH/Et₂O). Sol. H₂O; spar. sol. EtOH. Mp 226-230°.

Me ester: [50373-10-9]

Bp_{1,9} 87°. n_D^{23,2} 1.5026.

Matchett, J.R. *et al.*, *J.A.C.S.*, 1941, **63**, 2444-2446 (*Me ester, isol*)


Grundman, C. *et al.*, *Annalen*, 1957, **605**, 24 (*synth, resoln, bibl*)

Holmquist, C.R. *et al.*, *Org. Prep. Proced. Int.*, 1997, **29**, 308 (*Me ester, synth*)

Nishiyama, Y. *et al.*, *J. Nat. Med. (Tokyo)*, 2007, **61**, 56-58 (*isol, pmr, cmr*)

4-Methyl-15-azasterol M-396

4-Methyl-15-aza-D-homoergosta-8,14,24(28)-trien-3-ol, 9CI. *Azx. Antibiotic Azx* [128500-61-8]




C₂₉H₄₇NO 425.696

The CA name is defective. Isol. from *Calcarisporium thermophilum* ATCC11485. Antifungal agent.

Chrip, P. *et al.*, *Z. Naturforsch., C*, 1990, **45**, 179 (*isol*)

Methylazoxymethanol M-397

(Methyl-ONN-azoxy)methanol, 9CI. *MAM*



C₂H₆N₂O₂ 90.082

▶ PC2625000

(Z)-form [590-96-5]

Toxic constit. of the nuts of *Cycas circinalis*. Bp_{0,6} 51°.

▶ Exp. carcinogen and teratogen. PC2625000

O-β-D-Glucopyranoside: **Cycasin**. β-D-Glucosyloxymethane

[14901-08-7]

C₈H₁₆N₂O₇ 252.224

Alkaloid from seeds of the false sago *Cycas circinalis* and sago cycas *Cycas revoluta* (Cycadaceae). Carcinogen of significance in human nutrition. Mp 154° dec. (144-145° dec.). [α]_D¹⁸ -44 (c, 0.62 in H₂O). λ_{max} 218 (ε 7240); 275 (sh) (ε 50) (H₂O) (Derep).

▶ Exp. teratogen. Possible human carcinogen (IARC 2B). LZ5950000

O-β-D-Glucopyranoside, tetra-O-Ac: Mp 137°. [α]_D¹⁸ -27 (c, 1 in CHCl₃).

O-[[β-D-Xylopyranosyl-(1→6)]-D-glucopyranoside]: **Macrozamin** [6327-93-1]

C₁₃H₂₄N₂O₁₁ 384.339

Constit. of *Macrozamia riedlei* and *Macrozamia spiralis*. Prisms (EtOH aq.). Mp 202-203° dec. [α]_D²³ -71.3 (c, 0.4 in H₂O). λ_{max} 217 (ε 7940); 275 (sh) (ε 50) (H₂O) (Derep).

O-[[β-D-Glucopyranosyl-(1→3)]-β-D-glucopyranoside]: **Neocycasin A** [2288-32-6]

C₁₄H₂₆N₂O₁₂ 414.366

Isol. from *Cycas revoluta*. Cryst. + 1H₂O. Mp 162-163° dec. [α]_D²⁹ -35.1. λ_{max} 217 (ε 7940); 275 (sh) (ε 50) (H₂O) (Derep).

▶ QO7500000

O-[[β-D-Glucopyranosyl-(1→4)]-β-D-glucopyranoside]: **Neocycasin E** [2288-32-6]

C₁₄H₂₆N₂O₁₂ 414.366

Alkaloid from *Cycas revoluta* (Cycadaceae). Mp 156-158° dec. [α]_D⁵ -29.2 (c, 0.9 in H₂O).

O-[[α-D-Glucopyranosyl-(1→6)]-β-D-glucopyranoside]: **Neocycasin B₂** [97673-88-6]

C₁₄H₂₆N₂O₁₂ 414.366

Isol. from seeds of *Cycas revoluta*.

O-[[β-D-Glucopyranosyl-(1→6)]-β-D-glucopyranoside]: **Neocycasin B** [2697-20-3]

C₁₄H₂₆N₂O₁₂ 414.366

Isol. from *Cycas revoluta*. λ_{max} 217 (ε 7940); 275 (sh) (ε 50) (H₂O) (Derep).

O-[[β-D-Glucopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→3)]-β-D-glucopyranoside]: **Neocycasin C** [2288-31-5]

C₂₆H₄₆N₂O₂₂ 738.65

Isol. from *Cycas revoluta*.

O-[[β-D-Glucopyranosyl-(1→4)]-β-D-glucopyranosyl-(1→3)]-β-D-glucopyranoside]: **Neocycasin H** [102770-23-0]

C₂₀H₃₆N₂O₁₇ 576.508

Isol. from *Cycas revoluta*. Artifact formed from Neocycasin A by enzymic transglycosylation.

O-[[β-D-Glucopyranosyl-(1→6)]-β-D-glucopyranosyl-(1→3)]-β-D-glucopyranoside]: **Neocycasin G** [2288-28-0]

C₂₀H₃₆N₂O₁₇ 576.508

Isol. from seeds of *Cycas circinalis*.

O-[[β-D-Glucopyranosyl-(1→6)]-[[β-D-glucopyranosyl-(1→3)]-β-D-glucopyranoside]: **Neocycasin I**

[102770-22-9]

C₂₀H₃₆N₂O₁₇ 576.508

Artifact of transglycosylation by incubation of *Cycas* sp. with β-glucosidase.

O-[[β-D-Glucopyranosyl-(1→3)]-[[β-D-xylopyranosyl-(1→6)]-β-D-glucopyranoside]: **Neocycasin J** [102770-21-8]

C₁₉H₃₄N₂O₁₆ 546.481

Isol. from leaves of *Cycas revoluta*.

Ac: [592-62-1]

C₄H₈N₂O₃ 132.119

Bp 191° Bp_{0,45} 49°.

▶ LD₅₀ (rat, orl) 270 mg/kg. Possible human carcinogen. Exp. carcinogen and teratogen. PC2800000

[106251-55-2]

Riggs, N.V. *et al.*, *Chem. Ind. (London)*, 1956, 926 (*isol, uv, struct*)

Nagahama, T. *et al.*, *Agric. Biol. Chem.*, 1961, **25**, 937 (*Neocycasin E*)

Korsch, B.H. *et al.*, *Tet. Lett.*, 1964, 523 (*pmr, struct*)

Kobayashi, A. *et al.*, *Arch. Biochem. Biophys.*, 1965, **110**, 373 (*isol*)

Palekar, R.S. *et al.*, *Nature (London)*, 1965, **206**, 1363

IARC Monog., 1976, **10**, 131; *Suppl.*, 7, 66 (*tox, rev, Ac*)

Baldwin, J.E. *et al.*, *J.O.C.*, 1978, **43**, 2427 (*synth*)

Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1980, **33**, 2229 (*Macrozamin*)

Moretti, A. *et al.*, *Phytochemistry*, 1981, **20**, 1415 (*occur*)

Yagi, F. *et al.*, *Agric. Biol. Chem.*, 1983, **47**, 137; 1985, **49**, 1531; 2985 (*Neocycasins*)

Matsumoto, H. *et al.*, *CRC Handb. Nat. Occurring Food Toxicants*, 1983, 39 (*rev*)

Zedeck, M.S. *et al.*, *ACS Monogr.*, No. 182, 1984, (*rev*)

Hoffmann, G.R. *et al.*, *Environ. Mutagen.*, 1984, **6**, 103 (*rev*)

Tate, M.E. *et al.*, *Aust. J. Chem.*, 1995, **48**, 1059 (*cryst struct*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, COU000; HMG000; MGS750

3-Methylbenzoic acid M-398

m-Toluic acid

[99-04-7]

[25567-10-6]



C₈H₈O₂ 136.15

Prisms (MeOH or H₂O). Mod. sol. hot H₂O. Mp 111°. Bp 263°. pK_a 4.25 (25°). Sublimes, steam-volatile.

▶ LD₅₀ (mus, orl) 1630 mg/kg. XU1200000

Diethylamide: Diethyltoluamide, BAN, BSI, ISO, USAN. Metadelphene.

Detamide†. Deet, ESA. Off

[134-62-3]

[26545-51-7]

C₁₂H₁₇NO 191.272

Isol. from the female pink bollworm moth *Pectinophora gossypiella*. Insect repellent esp. effective against

Culicidae. Colourless to amber liq.

Insol. H₂O. d₄²⁰ 1. Bp₁₉ 160° Bp₁ 111°.

n_D^{20} 1.5206.

► Irritant. Can affect CNS by skin absorption. LD₅₀ (rat, orl) 2000 mg/kg. XS3675000

McCabe, E.T. *et al.*, *J.O.C.*, 1954, **19**, 493 (synth, diethylamide)

Jones, W.A. *et al.*, *Science (Washington, D.C.)*, 1968, **159**, 99-100 (diethylamide, isol)


Shioiri, T. *et al.*, *Tetrahedron*, 1976, **32**, 2211 (synth, ir, pmr, diethylamide)

McIver, S.B. *et al.*, *J. Med. Entomol.*, 1981, **18**, 357 (rev, diethylamide)

Robbins, P.J. *et al.*, *J. Toxicol. Environ. Health*, 1986, **18**, 503 (diethylamide)

LeFevre, J.W. *et al.*, *J. Chem. Educ.*, 1990, **67**, A278 (diethylamide)

4-Methylbenzo[g]quinoline-5,10-dione, 9CI M-399
Cleistopholine
[96889-94-0]

C₁₄H₉NO₂ 223.231

Systematic numbering shown. Has also been numbered as an anthraquinone. Alkaloid from *Cananga odorata*, *Meiogyne virgata*, *Annona hayesii*, root bark of *Cleistopholis patens*, seeds of *Annona cherimolia* and stem bark of *Oncodostigma monosperma* (Annonaceae). Pale yellow amorph. solid. Mp 185-190°.

7-Methoxy: 7-Methoxy-4-methylbenzo[g]quinoline-5,10-dione. 7-Methoxycleistopholine. 6-Methoxycleistopholine [190127-27-6]

C₁₅H₁₁NO₃ 253.257

Alkaloid from *Porcelia macrocarpa*. Amorph. powder. λ_{max} 234 (log ε 3.41); 262 (log ε 3.44); 302 (log ε 3.46); 366 (log ε 3.22) (MeOH).

7-Methoxy, 6-hydroxy: 6-Hydroxy-7-methoxy-4-methylbenzo[g]quinoline-5,10-dione. 6-Hydroxy-7-methoxycleistopholine. 5-Hydroxy-6-methoxycleistopholine [330579-53-8]

C₁₅H₁₁NO₄ 269.256

Alkaloid from *Porcelia macrocarpa*. Amorph. powder. λ_{max} 214 (log ε 2.43); 246 (log ε 3.65); 260 (log ε 3.49); 430 (log ε 2.75) (MeOH).

6,7-Dimethoxy: 6,7-Dimethoxy-4-methylbenzo[g]quinoline-5,10-dione. 6,7-Dimethoxycleistopholine. 5,6-Dimethoxycleistopholine [330579-50-5]

C₁₆H₁₃NO₄ 283.283

Alkaloid from *Porcelia macrocarpa*. Amorph. powder. λ_{max} 208 (log ε 3.76); 234 (log ε 3.79); 282 (log ε 3.91); 374 (log ε 2.95) (MeOH).

7,8-Dimethoxy, 6-hydroxy: 6-Hydroxy-7,8-dimethoxy-4-methylbenzo[g]quinoline-5,10-dione. 6-Hydroxy-7,8-dimethoxycleistopholine. 5-Hydroxy-6,7-

dimethoxycleistopholine

[330579-56-1]

C₁₆H₁₃NO₅ 299.282

Alkaloid from *Porcelia macrocarpa*. Amorph. solid. λ_{max} 212 (log ε 4.3); 242 (log ε 4.27); 282 (log ε 4.35); 400 (log ε 3.8) (MeOH).

Waterman, P.G. *et al.*, *Phytochemistry*, 1985, **24**, 523-527 (isol, uv, ir, pmr, cmr, ms, struct)

Bracher, F. *et al.*, *Annalen*, 1989, 87-88 (synth, uv, ir, pmr, cmr, ms)


Koyama, J. *et al.*, *Heterocycles*, 1989, **29**, 1649-1654 (synth)

Bou-Abdallah, E. *et al.*, *J. Nat. Prod.*, 1989, **52**, 273-278 (isol)

Rios, J.L. *et al.*, *Planta Med.*, 1989, **55**, 321-323 (isol)

Chaves, M.H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 240-242 (*Porcelia* derivs)

2-Methylbenzothiazole, 9CI M-400
[120-75-2]

C₈H₇NS 149.216

Prod. by a *Micrococcus* sp. found in *Tedania ignis*. Also known as an aroma constit. of tea leaves. Mp 14°. Bp 238° Bp₁₅ 150-151°. pK_a 2.06 (H₂O). pK_a 8.63 (MeCN).

► LD₅₀ (mus, ipr) 300 mg/kg. DL5600000
Picrate: Mp 153.5°.

N-Oxide: [23808-61-9]

C₈H₇NOS 165.215

Light yellow cryst. + 2H₂O (MeCN aq.). Mp 46-48°.

N-Et: 1-Ethyl-2-methylthiazolium(1+)

[3119-93-5 (iodide)]

C₁₀H₁₂NS⁺ 178.278

Needles (EtOH) (as iodide). Mp 196° (iodide).

N-Propyl: 2-Methyl-1-propylthiazolium(1+)

[60126-29-6 (iodide)]

C₁₁H₁₄NS⁺ 192.304

Precursor in synth. of carbocyanine dyes. Mp 173-176° (as iodide).

[14933-76-7]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 699D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 200B; 201B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1511B (ir)

Hofmann, A.W. *et al.*, *Ber.*, 1880, **13**, 8-22; 1223-1238 (synth)

Mills, W.H. *et al.*, *J.C.S.*, 1922, **121**, 455-466 (N-Et)

Clark, L.M. *et al.*, *J.C.S.*, 1928, 2313-2320 (synth)

Brooker, L.G.S. *et al.*, *J.A.C.S.*, 1935, **57**, 2480 (N-propyl, synth, use)

Millard, B.J. *et al.*, *Org. Mass Spectrom.*, 1968, **1**, 285 (ms)

Schweizer, E.E. *et al.*, *J.O.C.*, 1978, **43**, 2972 (synth)

Sawhney, S.N. *et al.*, *J.O.C.*, 1979, **44**, 1136 (cmr)

Peyronel, G. *et al.*, *Spectrochim. Acta A*, 1981, **37**, 71 (ir)

Bowman, W.R. *et al.*, *Tet. Lett.*, 1982, **23**, 5093 (synth)

Wanner, K.T. *et al.*, *Annalen*, 1984, 1100 (synth)

Stefaniak, L. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 215 (N-15 nmr)

Kolb, B. *et al.*, *J. Agric. Food Chem.*, 1985, **33**, 999 (isol)

Alyea, E.C. *et al.*, *J. Het. Chem.*, 1985, **22**, 1325 (synth)


Stierle, A.A. *et al.*, *Tet. Lett.*, 1991, **32**, 4847 (isol)

Kamila, S. *et al.*, *Heterocycles*, 2005, **65**, 2119-2126 (synth, pmr, cmr)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MHI750

N-Methylbicycloatalaphyl- line M-401

3,4,6,7,8,9-Hexahydro-10-hydroxy-2,2,6,6,9-pentamethyl-2H,14H-dipyrido[2,3-a:2',3'-c]acridin-14-one, 9CI [35096-35-6]


C₂₄H₂₇NO₄ 393.482

Alkaloid from the root bark of *Atalantia monophylla* (Rutaceae). Yellow needles (C₆H₆). Mp 185°.

Basu, D. *et al.*, *J.O.C.*, 1972, **37**, 3035 (isol, uv, ir, pmr, ms, struct)

α-Methylbiotin, 8CI M-402

Hexahydro-2-oxo-α-methyl-1H-thieno[3,4-d]imidazole-4-pentanoic acid, 9CI [30868-27-0]

C₁₁H₁₈N₂O₃S 258.341


Prod. by *Streptomyces lydicus*. Antimetabolite antibiotic. Cryst. Sol. H₂O; fairly sol. Me₂CO. Mp 251.5-252°.

Martin, D.G. *et al.*, *Tet. Lett.*, 1971, 3791 (isol, struct, ms, pmr)

Hanka, L.J. *et al.*, *Antimicrob. Agents Chemother.*, 1972, **1**, 135 (isol, biosynth)

4-Methyl-2,3'-bipyridine, 9CI M-403

[38840-05-0]

C₁₁H₁₀N₂ 170.213

Present in burley tobacco (*Nicotiana tabacum*). Liq. Bp₁ 166°.

Picrate:


Cryst. (EtOH). Mp 202-203°.

Frank, R.L. *et al.*, *Bull. Soc. Chim. Fr.*, 1958, 419 (*synth*)Warfield, A.H. *et al.*, *Phytochemistry*, 1972, **11**, 3371 (*isol, synth, ir, ms*)Ishikura, M. *et al.*, *Synthesis*, 1984, 936 (*synth*)**4-Methyl-3,3'-bipyridine, 9CI** M-404


[38840-06-1]

C₁₁H₁₀N₂ 170.213Isol. from burley tobacco (*Nicotiana tabacum*).Warfield, A.H. *et al.*, *Phytochemistry*, 1972, **11**, 3371 (*isol, synth, ir, pmr*)**5-Methyl-2,3'-bipyridine** M-405


[26844-80-4]

C₁₁H₁₀N₂ 170.213Alkaloid from burley tobacco *Nicotiana tabacum* (Solanaceae).Warfield, A.H. *et al.*, *Phytochemistry*, 1972, **11**, 3371 (*isol, uv, ir, pmr, ms, struct, synth*)**1-Methyl-3,3-bis(3-methyl-2-butenyl)-2,4(1H,3H)-quinolinedione, 9CI** M-406*1-Methyl-3,3-diprenyl-2,4-quinolinedione* [57931-85-8]C₂₀H₂₅NO₂ 311.423Alkaloid from the wood of *Esenbeckia flava*. Oil. λ_{\max} 235 (ϵ 3400); 258 (ϵ 450); 342 (ϵ 250) (EtOH).Dreyer, D.L. *et al.*, *Phytochemistry*, 1980, **19**, 941-944 (*isol, uv, ir, pmr, ms*)Nunes, F.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 180-183 (*pmr, cmr*)**6-(3-Methyl-1,3-butadienyl)-1H-indole, 9CI** M-407

[36948-68-2]

C₁₃H₁₃N 183.252**(E)-form** [40863-49-8]Alkaloid from seeds of *Monodora tenuifolia* (Annonaceae). Leaflets (hexane). Mp 124-127°.**(Z)-form** [85864-10-4]

Synthetic. Unstable oil.

Nwaji, M.N. *et al.*, *Chem. Comm.*, 1972, 327*(isol, pmr, uv, struct)*Ishii, H. *et al.*, *Tet. Lett.*, 1973, 355 (*synth, config*)Onyiruka, S.O. *et al.*, *J. Chem. Res., Synop.*, 1983, 21 (*synth*)Somei, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 4109 (*synth, pmr*)**6-(3-Methyl-1,3-butadienyl)-1H-indole-3-carboxylic acid** M-408C₁₄H₁₃NO₂ 227.262**(E)-form***Antibiotic TMC 205. TMC 205*

[403646-00-4]

Prod. by the unidentified fungal strain TC 1630. Transcriptional up-regulator of SV40 promoter. Cytotoxic. Pale yellow powder. λ_{\max} 205 (log ϵ 4.03); 210 (sh) (log ϵ 4.02); 271 (log ϵ 4.27); 305 (log ϵ 4.15) (MeOH).Sakurai, M. *et al.*, *J. Antibiot.*, 2001, **54**, 628-634**3-Methylbutanoic acid** M-409*Isovaleric acid. Viburnum acid. Baldrianic acid. Phoenicic acid. Isopropylacetic acid. Isopentanoic acid. FEMA 3102*

[503-74-2]

(H₃C)₂CHCH₂COOHC₅H₁₀O₂ 102.133Constit. of tobacco, hops, cheese etc. Widespread in plants, in free state and esp. as esters. Prod. by *Xanthomonas campestris* (Oryzae). Flavouring agent. Simple esters are used in flavourings and perfumes. Histone deacylase inhibitor. Phytotoxic. Oil with disagreeable, rancid-cheese aroma, acid taste. Mod. sol. H₂O. d_4^{18} 0.93. Mp -37.6°. Bp 176.7° (183°). n_D^{15} 1.4064. pK_a 4.8.► Skin and eye irritant. LD₅₀ (rat, orl) 2000 mg/kg. Fl. p. 79°, autoignition temp. 416/418°. NY1400000*Amide: 3-Methylbutanamide. Isovaleramide. USAN. NPS 1776*

[541-46-8]

C₅H₁₁NO 101.148Constit. of *Rhizoclonium hieroglyphicum*. Anticonvulsant agent used to treat epilepsy and other neurological and psychiatric disorders. Mp 131°.

► EJ3650000

Dembitsky, V.M. *et al.*, *Phytochemistry*, 2000, **54**, 965-967 (*isol, amide*)**N-(3-Methyl-2-butenyl)guanidine** M-410*N-Prenylguanidine. Galegine*


[543-83-9]

HN=C(NH₂)NHCH₂CH=C(CH₃)₂C₆H₁₃N₃ 127.189Toxic principle from *Verbesina encelioides*, *Galega officinalis* (goat's rue) etc. (Asteraceae, Fabaceae). Antihyperglycaemic agent. Shows antidiabetic props. V. hygroscopic cryst. Mp 60-65°. Log P 0 (calc).

► Pulmonary toxicant. MF3687000

*Sulfate:*Needles (H₂O). Mp 223-225°.Tanret, G. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1914, **158**, 1182Schreiber, K. *et al.*, *Annalen*, 1964, **671**, 147; 154 (*isol, synth*)Reuter, G. *et al.*, *Planta Med.*, 1965, **13**, 494 (*biosynth*)Eichholzer, J.V. *et al.*, *Phytochemistry*, 1982, **21**, 97-99 (*isol, pmr, ms*)Huxtable, C.R. *et al.*, *Aust. Vet. J.*, 1993, **70**, 169 (*isol, tox*)Monache, G.D. *et al.*, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 3249-3254 (*isol, pmr, cmr*)**3-(3-Methyl-2-butenyl)-1H-indole, 9CI** M-411*3-(γ,γ -Dimethylallyl)indole. 3-Prenylindole*

[17771-42-5]

C₁₃H₁₅N 185.268Alkaloid from the stem bark of *Monodora tenuifolia* (Annonaceae). Antibacterial and antifungal agent. Cryst. (hexane). Sol. MeOH, CHCl₃; fairly sol. hexane; poorly sol. H₂O. Mp 45-47° (43-45°). λ_{\max} 225 ; 282 ; 292 (MeOH) (Berdy).Bocchi, V. *et al.*, *Tetrahedron*, 1978, **34**, 929 (*synth*)Adeoye, A.O. *et al.*, *J. Nat. Prod.*, 1986, **49**, 534 (*isol, uv, ir, pmr, cmr, ms, synth*)**5-(3-Methyl-2-butenyl)-1H-indole, 9CI** M-412*5-Prenylindole*

[32962-26-8]

C₁₃H₁₅N 185.268Alkaloid from roots of *Esenbeckia leio-carpa*. Oil. *Δ^3 -Isomer, 1',2'-didehydro: 5-(3-Methyl-1,3-butadienyl)-1H-indole*C₁₃H₁₃N 183.252Alkaloid from the stem bark of *Isolona cauliflora*.Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (*isol, uv, pmr, cmr, ms*)Nkunya, M.H.H. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 253-258 (*Δ^3 -isomer didehydro*)**6-(3-Methyl-2-butenyl)-1H-indole** M-413*6-Prenylindole. Antibiotic 434B*

[23158-16-9]

C₁₃H₁₅N 185.268

Alkaloid from the liverworts *Riccardia sinuata* and *Riccardia incurvata* (Hepaticae) and from roots of *Esenbeckia leiocarpa*. Prod. by *Streptomyces hygrosopicus* and *Streptomyces* sp. TP-A595. Antitumour agent. Mp 33-34° (30.5-31°). Bp₂ 120-130°. Log P 4.13 (calc). Also descr. as pale brown oil. λ_{\max} 269; 279; 293 (MeOH) (Berdy). λ_{\max} 269; 280; 291 (MeOH/HCl) (Berdy). λ_{\max} 270; 280; 292 (MeOH/NaOH) (Berdy).

1,3,5-Trinitrobenzene complex: Mp 106°.

Benešová, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1807 (*isol, uv, ir, pmr, struct*)

Pleiningner, H. *et al.*, *Chem. Ber.*, 1971, **104**, 2027 (*synth*)

Huneck, S. *et al.*, *CA*, 1972, **79**, 29523 (*isol*)

Ishii, H. *et al.*, *Tetrahedron*, 1975, **31**, 933 (*synth*)

Japan. Pat., 1979, 79 135 300; *CA*, **92**, 126908 (*isol*)

Achenbach, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1985, **318**, 1147 (*synth*)

Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435 (*isol*)

Sasaki, T. *et al.*, *J. Antibiot.*, 2002, **55**, 1009-1012 (*isol, synth*)

7-(3-Methyl-2-butenyl)-1H-indole M-414

7-Prenylindole
[23158-14-7]

C₁₃H₁₅N 185.268

Alkaloid from the liverworts *Riccardia sinuata* and *Riccardia incurvata*, and from the stem bark of *Anonidium mannii* (Annonaceae) and roots of *Esenbeckia leiocarpa* (Rutaceae). Mp 43-44° (ca. 20°).

1,3,5-Trinitrobenzene complex: Mp 110.5-111°.

1'-Oxo: 1-(1H-Indol-7-yl)-3-methyl-2-buten-1-one, 9CI. 7-(3-Methyl-2-butenyl)-1H-indole. **Leiocarpone**
[105205-52-5]

C₁₃H₁₃NO 199.252

Isol. from the roots of *Esenbeckia leiocarpa*. Air- and light-sensitive cryst.

1'-Hydroxy, 3',4'-epoxide: α -(3,3-Dimethyloxiran-1-yl)-1H-indole-7-methanol, 9CI. **Leiocarpol**
[126365-18-2]

C₁₃H₁₅NO₂ 217.267

Alkaloid from roots of *Esenbeckia leiocarpa*. Amorph. solid. $[\alpha]_{\text{D}}^{20}$ -1.5 (c, 1.1 in CHCl₃).

Benešová, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1807 (*isol, uv, ir, pmr*)

Achenbach, H. *et al.*, *Heterocycles*, 1985, **23**, 2075 (*isol, uv, ir, pmr, cmr, ms*)

Muratake, H. *et al.*, *Heterocycles*, 1986, **24**, 261 (*synth*)

Moyer, M.P. *et al.*, *J.O.C.*, 1986, **51**, 5106 (*synth, pmr, Leiocarpone*)


Somei, M. *et al.*, *Heterocycles*, 1987, **26**, 1783 (*synth*)

Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 435; 1990, **120**, 387 (*Leiocarpol, Leiocarpone*)

Xiong, X. *et al.*, *J.O.C.*, 2007, **72**, 5832-5834 (*synth*)

5-(3-Methyl-2-butenyl)-1H-indole-2,3-dione, 9CI M-415

5-Prenylisatin. 5-Isopentenylisatin. **Prenisatin**. 2-Hydroxy-5-(3-methyl-2-butenyl)-3H-indol-3-one
[177957-18-5]




C₁₃H₁₃NO₂ 215.251

Likely to be tautomeric with the Δ^1 -enone form. Prod. by *Chaetomium globosum*. Orange solid. λ_{\max} 247 (log ϵ 4.28); 299 (log ϵ 3.59); 422 (log ϵ 2.82) (MeOH).

Breinholt, J. *et al.*, *Acta Chem. Scand.*, 1996, **50**, 443-445

6-(3-Methyl-2-butenyl)-1H-indole-2,3-dione, 9CI M-416

6-Prenylisatin. 6-Isopentenylisatin. 2-Hydroxy-6-(3-methyl-2-butenyl)-3H-indol-3-one
[101023-73-8]




C₁₃H₁₃NO₂ 215.251

Isol. from *Streptomyces albus*. Shows weak activity against gram-positive bacteria. Yellow cryst. (MeOH). Sol. EtOAc, MeOH, CHCl₃; poorly sol. H₂O. Mp 109-110°.

Graefe, U. *et al.*, *J. Antibiot.*, 1986, **39**, 162 (*isol*)

6-(3-Methyl-2-butenyl)-1-phenazinecarboxylic acid M-417

6-Prenyl-1-phenazinecarboxylic acid
[85223-60-5]



C₁₈H₁₆N₂O₂ 292.337

Isol. from *Streptomyces cinnamomensis*. Weakly active against gram-positive bacteria. Cryst. (CHCl₃/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 168-170°. λ_{\max} 255 (ϵ 65000); 367 (ϵ 14000) (MeOH) (Berdy).

6-Deoxy- α -L-talopyranosyl ester: **Phenazoviridin**
[155233-15-1]

C₂₄H₂₆N₂O₆ 438.479

Prod. by *Streptomyces* sp. HR04. Free radical scavenger, lipid peroxidation

inhibitor, shows antiedementia props. Pale greenish powder. Sol. MeOH, Me₂CO, CHCl₃, EtOAc, EtOH; poorly sol. H₂O, hexane. Mp 71-73° dec. λ_{\max} 253 (ϵ 71700); 348 (ϵ 11600); 366 (ϵ 16900) (MeOH).

$\Delta^{1'}$ -Isomer(E-), 3''-hydroxy: 6-(3-Hydroxy-3-methyl-1-butenyl)-1-phenazinecarboxylic acid
[129083-43-8]

C₁₈H₁₆N₂O₃ 308.336

Prod. by *Streptomyces cinnamomensis*.

Tax, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1983, **48**, 527 (*isol, struct, spectra*)

Beran, M. *et al.*, *Folia Microbiol. (Prague)*, 1990, **35**, 172-175 (*Streptomyces cinnamomensis constits*)

Kato, S. *et al.*, *J. Antibiot.*, 1993, **46**, 1485-1493 (*Phenazoviridin*)

9-(3-Methyl-2-butenyl)-1-phenazinecarboxylic acid M-418

9-Prenyl-1-phenazinecarboxylic acid. **Endophenazine A**

C₁₈H₁₆N₂O₂ 292.337

Prod. by various strains of the endosymbiotic *Streptomyces anulatus*. Active against gram-positive bacteria and some fungi. Yellow solid. λ_{\max} 255 (log ϵ 4.81); 365 (log ϵ 4.06) (MeOH).

5,10-Dihydro, N³-Me: 5,10-Dihydro-5-methyl-9-(3-methyl-2-butenyl)-1-phenazinecarboxylic acid. **Endophenazine C**

C₁₉H₂₀N₂O₂ 308.379


Prod. by *Streptomyces anulatus*. Active against gram-positive bacteria and some fungi. Green solid (probably as a complex with a paramagnetic metal). Orange solid (as Me ester). λ_{\max} 254 (log ϵ 4.47); 410 (log ϵ 3.77) (MeOH) (complex). λ_{\max} 248 (log ϵ 4.09); 323 (log ϵ 2.6); 432 (log ϵ 3.28) (MeOH) (Me ester).

Gebhardt, K. *et al.*, *J. Antibiot.*, 2002, **55**, 794-800; 801-806 (*isol, pmr, cmr, activity*)

Bringmann, G. *et al.*, *J.O.C.*, 2007, **72**, 4198-4202 (*biosynth*)

4-(3-Methyl-2-butenyl)tryptophan M-419

4-(3,3-Dimethylallyl)tryptophan. 4-Prenyltryptophan
[5017-33-4]



C₁₆H₂₀N₂O₂ 272.346

(S)-form [29702-35-0]

Occurs in a *Pennisetum* type of ergot strain producing mainly Elymoclavine. Important intermed. in biosynth. of ergot alkaloids. Cryst. (DMF aq.). Mp 210°.

N^z-Me: N-Methyl-4-dimethylallyltryptophan

[75917-16-7]

C₁₇H₂₂N₂O₂ 286.373Isol. from *Claviceps fusiformis* and *Lotus subglauca*. Needles (MeOH). Mp 232°. λ_{max} 274 ; 280 ; 295 (MeOH).**4'-Hydroxy: 4-(4-Hydroxy-3-methyl-2-butenyl)tryptophan**

[53774-59-7]

C₁₆H₂₀N₂O₃ 288.346Alkaloid from cultures of *Claviceps purpurea*. Config. of double bond not established. Identified by ms.Plieninger, H. *et al.*, *Chem. Ber.*, 1963, **96**, 1618-1629 (synth)Aguire, S. *et al.*, *Tet. Lett.*, 1968, 5127-5128 (occur, ms)Anderson, J.A. *et al.*, *Tet. Lett.*, 1974, 2107-2108 (N^z-Me, isol)Barrow, K.D. *et al.*, *Tet. Lett.*, 1975, 4269-4270 (N^z-Me, isol)Petroski, R.J. *et al.*, *FEBS Lett.*, 1977, **82**, 55-57 (biosynth)**3-Methyl-1-butylamine M-420****3-Methyl-1-butanamine, 9CI. 1-Amino-3-methylbutane. Isoamylamine. Isopentylamine. FEMA 3219**

[107-85-7]

(H₃C)₂CHCH₂CH₂NH₂C₅H₁₃N 87.164Present in roasted cocoa beans. Widespread in fungi and higher plants e.g. apple, banana, wine grape, swede, rhubarb, porcini (*Boletus edulis*). Protein degradn. product. Liq. Misc. H₂O. Bp 95° Bp 232°.

▶ Highly flammable, fl. p. <21°, irritant.

N-Ac: N-(3-Methylbutyl)acetamide. N-Isopentylacetamide. N-Isoamylacetamide

[13434-12-3]

C₇H₁₅NO 129.202Constit. of the pheromones of *Dacus* and *Vespa* spp. (wasp). Also found in wines and tobacco. Bp 238°.**N-Me: [4104-44-3]**C₆H₁₅N 101.191Found in tobacco. Bp₇₅₉ 108-110°.**N-Me; hydrochloride: [2419-59-2]**Plates (Me₂CO). V. sol. H₂O, insol.Et₂O. Mp 181° (178°).**N-Et: [21035-52-9]**C₇H₁₇N 115.218

Bp 127°.

N-Et; hydrochloride: Mp 205-206°.**N-Propyl: 3-Methyl-N-propylbutylamine, 9CI. Isopentylpropylamine**


[78579-58-5]

C₈H₁₉N 129.245Bp 148-149° Bp₂₀₀ 106-107°.*Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, **1**, 286D (ir)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 458C (nmr)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 370C (ir)Erickson, J.L.E. *et al.*, *Ber.*, 1926, **59**, 2665 (synth)Nicholas, H.O. *et al.*, *J.A.C.S.*, 1926, **48**, 2174 (synth)Campbell, K.N. *et al.*, *J.A.C.S.*, 1944, **66**, 82 (N-propyl)Dornow, A. *et al.*, *Annalen*, 1954, **588**, 62-71 (N-Et)Cope, A.C. *et al.*, *J.A.C.S.*, 1957, **79**, 4750 (N-propyl)von Kamienski, E.S. *et al.*, *CA*, 1958, **52**, 5551; 6502 (occur)Richardson, M. *et al.*, *Phytochemistry*, 1966, **5**, 23 (biosynth)Demole, E. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 1866-1882 (N-Ac, isol)Eggert, H. *et al.*, *J.A.C.S.*, 1973, **95**, 3710 (cmr)*Ger. Pat.*, 1976, 2 456 006; *CA*, **85**, 62631v


(synth)

Heath, R.R. *et al.*, *Experientia*, 1988, **44**, 82 (isol, synth)Perkins, M.V. *et al.*, *J.C.S. Perkin 1*, 1990, 1111-1117 (N-Ac, isol, bibl)Granvogel, M. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 1730-1739 (detn, cocoa)Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials, 5th edn.*, Van Nostrand Reinhold, 1979, 748**1-(3-Methylbutyl)-3-pyrroli-dinemethanol, 9CI M-421****3-Hydroxymethyl-1-isopentylpyrrolidine**

[163315-02-4]

C₁₀H₂₁NO 171.282Trace alkaloid from poison glands of the ants *Harpagoxenus sublaevis*, *Leptothorax acervorum* and *Leptothorax muscorum*. Bp₁ 52°.Reder, E. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 73 (isol, pmr, cmr, ms, synth, struct)**2-(3-Methylbutyl)-1*H*-pyrrolo[2,3-*b*]pyridine M-422****2-Isopentyl-7-azaindole**


[946-18-9]

C₁₂H₁₆N₂ 188.272

Constit. of chicken eggs. Antifungal agent. Cryst. Mp 150-151°.

Tsuji, K. *et al.*, *Yakugaku Zasshi*, 1976, **96**, 475-478 (isol)Landor, S.R. *et al.*, *J.C.S. Perkin 1*, 1979, 2289-2293 (synth)**1-Methyl-9*H*-carbazole, 9CI M-423**

[6510-65-2]

C₁₃H₁₁N 181.237Isol. from the sponge *Tedania ignis*. also from *Ailanthus melabrica*. Toxic to brine shrimp, shows insecticidal props. Plates(petrol). Mp 120.5°. λ_{max} 242 (ε 10715); 250 (ε 11700); 262 (ε 12300); 283 (ε 8317); 305 (ε 8317); 375 (ε 7050) (MeOH) (Berdy).Campaigne, E. *et al.*, *J.O.C.*, 1959, **24**, 478 (synth)Kuroki, M. *et al.*, *J. Het. Chem.*, 1981, **18**, 709 (synth, bibl)Clancy, M.G. *et al.*, *J.C.S. Perkin 1*, 1984, 429 (synth)Dillman, R.L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1056 (isol, pmr, cmr)**3-Methyl-9*H*-carbazole, 9CI M-424**


[4630-20-0]

C₁₃H₁₁N 181.237Alkaloid from the roots of *Clausena heptaphylla* and *Clausena indica* and from the rootbark of *Glycosmia pentaphylla* (Rutaceae). Plates (CH₂Cl₂/hexane). Mp 206-208°. Pale green soln. in H₂SO₄, which becomes deep green on addn. of HNO₃.

Picrate: Mp 176°.

N-Formyl: 3-Methyl-9*H*-carbazole-9-carboxaldehyde. 9-Formyl-3-methyl-9*H*-carbazoleC₁₄H₁₁NO 209.247Alkaloid from the roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Off-white solid. Mp 58-60°. λ_{max} 222 ; 267 ; 291 (sh) ; 315 (EtOH).**N-(Ethoxycarbonyl): Ethyl 3-methyl-9*H*-carbazole-9-carboxylate. 9-Carboethoxy-3-methyl-9*H*-carbazole**C₁₆H₁₅NO₂ 253.3Alkaloid from the roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Needles (MeOH). Mp 122-123°. λ_{max} 229 ; 251 ; 262 ; 278 (sh) ; 287 ; 304 ; 315 (EtOH).Campaigne, E. *et al.*, *J.O.C.*, 1959, **24**, 478Islam, A. *et al.*, *Chem. Comm.*, 1972, 537 (synth)Joshi, B.S. *et al.*, *Indian J. Chem.*, 1974, **12**, 437 (isol, uv)Roy, S. *et al.*, *Phytochemistry*, 1974, **13**, 1017 (isol, uv, ir)Åkermark, B. *et al.*, *J.O.C.*, 1975, **40**, 1365 (synth)Clancy, M.G. *et al.*, *Chem. Comm.*, 1980, 1112 (synth)Bhattacharyya, P. *et al.*, *Chem. Comm.*, 1984, 1668 (synth)Chowdhury, B.K. *et al.*, *Phytochemistry*, 1987, **26**, 2138 (isol)Luo, J.K. *et al.*, *J. Het. Chem.*, 1989, **26**, 1213 (synth, pmr)Chakrabarty, M. *et al.*, *Phytochemistry*, 1997, **46**, 751-755 (isol, derivs, pmr, cmr)Liu, Z. *et al.*, *Tetrahedron*, 2007, **63**, 347-355 (synth)**3-Methyl-1*H*-carbazole-1,4(9*H*)-dione, 9CI M-425****Murrayaquinone A**

[100108-66-5]




C₁₃H₉NO₂ 211.22

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Also obt. by photooxidn. of Murrayafoline A or oxidn. of 1-Hydroxy-3-methylcarbazole with Fremy's salt. Cytotoxic agent. Exhibits cardiotoxic activity. Brown prisms (Me₂CO). Mp 246–247° (240–241°).

- Wu, T.-S. *et al.*, *Heterocycles*, 1983, **20**, 1267 (uv, ir, pmr, cmr, struct)
 Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4132 (isol, uv, ir, pmr, cmr, synth, struct)
 Ramesh, K. *et al.*, *Chem. Ind. (London)*, 1986, 614 (synth)
 Martin, T. *et al.*, *J.C.S. Perkin 1*, 1988, 235 (synth, uv, ir, pmr, ms)
 Takeya, K. *et al.*, *Eur. J. Pharmacol.*, 1989, **169**, 137 (pharmacol)
 Yogo, M. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 328 (synth)
 Miki, Y. *et al.*, *Synlett*, 1993, 333 (synth)
 Wada, A. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 416 (synth)
 Matsuo, K. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1325 (synth)
 Hagiwara, H. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1948–1949 (synth)
 Murakami, Y. *et al.*, *Heterocycles*, 1998, **49**, 127–132 (synth)
 Murphy, W.S. *et al.*, *J.C.S. Perkin 1*, 1998, 4115–4119 (synth)
 Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 893–897 (activity)
 Knölker, H.-J. *et al.*, *Heterocycles*, 2003, **60**, 1049–1052 (synth)
 Scott, T.L. *et al.*, *Tetrahedron*, 2003, **59**, 6232–6232 (synth)
 Mal, D. *et al.*, *Tetrahedron*, 2007, **63**, 3768–3781 (synth)

1-Methyl- β -carboline M-426

1-Methyl-9H-pyrido[3,4-b]indole, 9CI. **Harman**. Loturine. Passiflorine†. Zygofabagine. Aribine [486-84-0]

C₁₂H₁₀N₂ 182.224

Present in tobacco smoke and espresso coffee. Alkaloid from *Arariba rubra* (preferred genus name *Simira*) (Rubiaceae), *Passiflora incarnata* (Passifloraceae) and many other *Passiflora* spp., and a wide range of spp. in several families. Prod. by *Nocardia* sp. and *Streptomyces* sp. Also a trace constit. of the fluorescent dinoflagellate *Noctiluca miliaris*. Isol. from *Costaticella hastata*. Cytotoxic intercalating agent. Plant growth and enzyme inhibitor. Sedative. Hallucinogen. Mp 237–238° (228°). Log P 3.06 (calc).

- ▶ A comutagen with aromatic amines active towards bacteria. LD₅₀ (mus, ipr) 50 mg/kg. UV0280000

Hydrochloride: Mp 292–295°.

Picrate: Mp 250–255°.

N-Oxide: Harmanine

[2506-09-4]

C₁₂H₁₀N₂O 198.224

Alkaloid from *Calligonum minimum* and *Ophiorrhiza rosacea*. Needles

(EtOAc/MeOH); cryst. (H₂O) (as picrate). Mp 182–184° subl Mp 239–248° (picrate).

N^b. Me: Melinonine F

[6801-22-5]

C₁₃H₁₃N₂⁺ 197.259

Quaternary alkaloid from *Strychnos melinoniana* and *Strychnos usambarensis* (Loganiaceae). Antimitotic agent. Poorly sol. hexane. Mp 288° dec. (as chloride).

3,4-Dihydro: 4,9-Dihydro-1-methyl-3H-pyrido[3,4-b]indole. 3,4-Dihydro-1-methyl- β -carboline. Harmalan. Dihydroharman

[525-41-7]

C₁₂H₁₂N₂ 184.24

Alkaloid from *Flindersia laevis*, *Elaeagnus angustifolia* (Russian olive), *Strychnos dale* and *Burkea africana* (Flindersiaceae, Elaeagnaceae, Loganiaceae, Fabaceae). Cream needles (Me₂CO). Mp 183–185° dec. λ_{\max} 237 (log ϵ 3.89); 241 (sh); 319 (log ϵ 3.86); 346 (sh) (log ϵ 3.62) (MeOH).

3,4-Dihydro, picrate:

Yellow needles (CHCl₃/MeOH). Mp 245° dec. (234–235° dec.).

3,4-Dihydro, N^b-oxide: Harmalan N^b-oxide

C₁₂H₁₂N₂O 200.24

Alkaloid from stem bark of *Burkea africana*. Probable struct.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 682A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 165B (nmr)

Manske, R.H.F. *et al.*, *J.C.S.*, 1927, 1–14 (*Harmalan, synth*)

Bächli, E. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 1167–1187 (*Melinonine F, isol, uv, ir*)

Kanaoka, Y. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 101–107 (*Harmalan, synth, uv, ir*)

Kametani, T. *et al.*, *J.C.S. (C)*, 1968, 1006–1007 (*synth, pmr*)

Ferreira, M.A. *et al.*, *Garcia de Orta, Ser. Farmacogn.*, 1973, **2**, 23–32; *CA*, **81**, 166304z (*Harmalan oxide*)

Phillipson, J.D. *et al.*, *J. Chromatogr.*, 1975, **105**, 163–178 (*tlc, glc, uv, ms*)

Picker, K. *et al.*, *Aust. J. Chem.*, 1976, **29**, 2023–2036 (*Harmalan, isol, uv, ir, pmr, ms*)

Inoue, S. *et al.*, *Chem. Lett.*, 1980, 297–298 (*isol, dinoflagellate*)

Han, B.H. *et al.*, *Arch. Pharmacol. Res.*, 1986, **9**, 21; *CA*, **105**, 57927y (*N^b-Formylharman, isol*)

Yomosa, K. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 921–922 (*isol, bibl*)

Hardiman, J. *et al.*, *Bioorg. Chem.*, 1987, **15**, 213–223 (*cryst struct*)

Blackman, A.J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 494–496 (*isol, Costaticella*)

Bracher, F. *et al.*, *Annalen*, 1993, 1335–1337 (*synth, pmr, cmr*)

Seki, H. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1169–1172 (*pmr, cmr*)

Rocca, P. *et al.*, *Tetrahedron*, 1993, **49**, 3325–3342 (*synth*)

Dachriyanus, *et al.*, *Aust. J. Chem.*, 2000, **53**, 221–224 (*oxide*)

Kitajima, M. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1410–1412 (*Harmalan, pmr, cmr*)

Nakano, K. *et al.*, *Mutat. Res.*, 2000, **470**, 141–146 (*comutagenicity, pmr*)

Ivanov, I. *et al.*, *Heterocycles*, 2005, **65**, 2483–2492 (*synth, Harmalan, pmr*)


Alves, R.C. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 1832–1838 (*isol, coffee*)

Huang, W. *et al.*, *Synth. Commun.*, 2007, **37**, 2137–2143 (*synth, ir, pmr*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MPA050

1-Methyl- β -carboline-3-carboxylic acid M-427

1-Methyl-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI. **Harman-3-carboxylic acid** [22329-38-0]

C₁₃H₁₀N₂O₂ 226.234

Alkaloid from the bark of *Aspidosperma polyneuron* and *Aspidosperma exalatum* (Apocynaceae). Mp 305–306°.

Me ester: Methyl harman-3-carboxylate. 3-Carbomethoxyharman [16641-82-0]

C₁₄H₁₂N₂O₂ 240.261

Alkaloid from bark of *Nauclea diderichii* (Rubiaceae). Cryst. (MeOH). Mp 252–253° (245°, 249–253° dec.).

▶ UU9780400

Et ester: Ethyl harman-3-carboxylate. 3-Carbethoxyharman [33821-71-5]

C₁₅H₁₄N₂O₂ 254.288

Alkaloid from bark of *Aspidosperma exalatum* (Apocynaceae).

Snyder, H.R. *et al.*, *J.A.C.S.*, 1948, **70**, 219 (*synth*)

Antonaccio, L.D. *et al.*, *Monatsh. Chem.*, 1962, **93**, 962 (*isol*)

Sanchez, W.E.L. *et al.*, *Ann. Acad. Bras. Cinc.*, 1971, **43**, 603–605; *CA*, **78**, 55312a (*isol, deriv*)


McLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**, 1478 (*isol, ir, pmr, ms*)

Cardoso, C.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1882–1885 (*isol, cmr*)

Zhao, M. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 6996–7010 (*synth, ir, pmr*)

4-Methylcinoline, 9CI M-428

[14722-38-4]

C₉H₈N₂ 144.176

Identified in cigarette smoke condensate and yeast extract. Pale yellow needles (petrol), orange needles (hexane). Mp 76–77° (73–74°). Deep red colour in acid soln.

▶ GE3630000

Picrate:

Aggregates of green needles (EtOH). Mp 179–180° (176–177°).

1-Oxide: [5580-86-9]

C₉H₈N₂O 160.175

Mp 95-96°.

2-Oxide: [5580-85-8]
 $C_9H_8N_2O$ 160.175
 Mp 150-151° (147-148°).


1,2-Dioxide:
 $C_9H_8N_2O_2$ 176.174
 Mp 171-172°.

1,2-Dihydro:
 $C_9H_{10}N_2$ 146.191
 Needles (petrol). Mp 64-65°. Bp_{0.2} 94-99°. Also descr. as a liq.

[82000-09-7, 66169-64-0]

Atkinson, C.M. *et al.*, *J.C.S.*, 1947, 808 (*synth. derivs*)Castle, R.N. *et al.*, *J.O.C.*, 1953, **18**, 1706 (*synth*)Castle, R.N. *et al.*, *J. Am. Pharm. Assoc.*, 1959, **48**, 135 (*ir*)Klemm, L.H. *et al.*, *J. Chromatogr.*, 1966, **23**, 428 (*tlc*)Palmer, M.H. *et al.*, *J.C.S.(C)*, 1968, **21**, 2621 (*synth, pmr, derivs*)Elkins, J.R. *et al.*, *J. Het. Chem.*, 1968, **5**, 639 (*ms, pmr*)Palmer, M.H. *et al.*, *Org. Mass Spectrom.*, 1969, **2**, 1265 (*ms*)Witanowski, M. *et al.*, *Tetrahedron*, 1971, **27**, 3129 (*N-14 nmr*)Van Hummel, G.J. *et al.*, *Acta Cryst. B*, 1979, **35**, 516 (*cryst struct*)Castellano, A. *et al.*, *J. Chem. Res., Synop.*, 1979, **2**, 70 (*epi*)


Methylcyclodecylprodigiosin M-429
Cyclomethyldecylprodiginine
 [34852-33-0]

 $C_{25}H_{33}N_3O$ 391.555

Pyrrole antibiotic. Isol. from *Actinomadura pelletieri* (*Nocardia pelletieri*). Active against *Staphylococcus aureus*, fungi and protozoa. λ_{max} 539 (EtOH/HCl) (Berdy). λ_{max} 549 (hexane) (Berdy).

Gerber, N.N. *et al.*, *Appl. Microbiol.*, 1969, **18**, 1; *J. Antibiot.*, 1971, **24**, 636; *J. Het. Chem.*, 1973, **10**, 925 (*isol, uv, ir, pmr, ms*)

Methylcyclooctylprodigiosin M-430
R 39FF. Antibiotic R 39FF
 [52589-16-9]

 $C_{23}H_{29}N_3O$ 363.502Pyrrole antibiotic. Prod. by *Actinoma-*

dura madurae. Active against gram-positive bacteria. Yellow solid. λ_{max} 551 (CHCl₃) (Berdy). λ_{max} 543 (EtOH/HCl) (Berdy).

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1973, **10**, 925-929 (*isol*)

N-(9-Methyldecyl)sulfamic acid M-431
acid


[1025324-20-2]
 $(H_3C)_2CH(CH_2)_7CH_2NHSO_3H$

 $C_{11}H_{25}NO_3S$ 251.389Isol. from *Daphnia pulex*. Kairomone.

Yasumoto, K. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 133-136 (*isol, pmr*)

α -Methyldethiobiotin M-432

α ,5-Dimethyl-2-oxo-4-imidazolidinehexanoic acid, 9CI, 8CI. Libramycin A
 [31602-99-0]

 $C_{11}H_{20}N_2O_3$ 228.291

From *Streptomyces lydicus*, *Streptomyces lavendulae* and *Streptomyces* sp. 72192.

Antimetabolite antibiotic. Cryst. (Me₂CO). Mp 161.5-162.5°.

[51746-00-0]


Martin, D.G. *et al.*, *Tet. Lett.*, 1971, 3791 (*isol, struct, ms, pmr*)

Hanka, L.J. *et al.*, *Antimicrob. Agents Chemother.*, 1972, **1**, 135 (*isol, biosynth*)

Okami, Y. *et al.*, *J. Antibiot.*, 1974, **27**, 143; 656

7-Methyl-1,5-diazacyclotetradecane M-433

Haliclovensin
 [202747-72-6]

*(R)*-form $C_{13}H_{28}N_2$ 212.378

Struct. revised in 2001.

(R)-form
 Oil. $[\alpha]_D^{20}$ +20 (c, 2 in MeOH).

(S)-form
 Alkaloid from the marine sponge *Haliclona tulearensis*. Oil. $[\alpha]_D^{20}$ -2.2 (c, 1.3 in MeOH) (natural). $[\alpha]_D^{20}$ -18.5 (c, 0.6 in MeOH) (synthetic). Nat. prod. is a partial racemate of 75% o.p.


Koren-Goldshlager, G. *et al.*, *J. Nat. Prod.*, 1998, **61**, 282 (*isol, pmr, cmr, ms*)

Heinrich, M.R. *et al.*, *Tetrahedron*, 2001, **57**, 9973-9978 (*synth, cd, pmr, cmr, struct*)

Zheng, J.-F. *et al.*, *Org. Lett.*, 2004, **6**, 1139-1142 (*synth*)

2-(Methyldithio)-1H-indole-3-carboxaldehyde M-434

3-Formyl-2-(methyldithio)indole. 3-Formylindolyl methyl disulfide. *Caulilexin A*

 $C_{10}H_9NOS_2$ 223.319

Alkaloid from *Brassica oleracea* var. *botrytis* (cauliflower). Phytoalexin. Pale yellow solid. Mp 114-116°. λ_{max} 213 (log ϵ 4.4); 252 (log ϵ 4.2); 307 (log ϵ 4) (MeOH).

Pedras, M.S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1503-1509 (*isol, synth, pmr, cmr, ms*)

2-Methyl-2,6-eicosadienoic acid M-435

$H_3C(CH_2)_{12}CH=CHCH_2CH_2CH=C(CH_3)COOH$

 $C_{21}H_{38}O_2$ 322.53*(2E,6E)*-form

(2-Hydroxyethyl)amide: Semiplenamide A

[630100-41-3]

 $C_{23}H_{43}NO_2$ 365.598Alkaloid from *Lyngbya semiplena*.

Amorph. solid. λ_{max} 206 (ϵ 5100) (MeOH).

(2-Acetoxyethyl)amide: Semiplenamide B


[630100-42-4]

 $C_{25}H_{45}NO_3$ 407.635Alkaloid from *Lyngbya semiplena*. Oil. λ_{max} 203 (ϵ 5900) (MeOH).

Han, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1364-1368 (*isol, pmr, cmr*)

3,3'-Methylenebisindole, 9CI M-436

3,3'-Methylenediindole. Di-3-indolyl-methane. *Arundine*
 [1968-05-4]

 $C_{17}H_{14}N_2$ 246.311

Alkaloid from the roots of *Arundo donax*. Prod. by the marine bacterium *Vibrio parahaemolyticus* Bio249. Major digestive prod. of 1H-Indole-3-methanol, I-98. Exhibits various potent anticancer props. Needles (EtOH aq.). Mp 165-166°.

N-Me: *Aridine*

[174545-80-3]

 $C_{18}H_{16}N_2$ 260.338Alkaloid from *Arundo donax*.

Amorph.

N,N'-Di-Me:

 $C_{19}H_{18}N_2$ 274.365


Plates (EtOH). Mp 110-112°.

Thesing, J. *et al.*, *Chem. Ber.*, 1954, **87**, 692-699 (*synth*)

Jackson, A.H. *et al.*, *J.C.S. Perkin 1*, 1987, 2543-2551 (*synth, pmr*)


Khuzaev, V.U. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1994, **30**, 635; 1995, **31**, 277; 277; 1996, **32**, 190 (*isol. Arundine, Ardine*)
 Veluri, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1520-1523 (*isol*)
 Anderton, M.J. *et al.*, *Clin. Cancer Res.*, 2004, **10**, 5233-5241 (*pharmacokinetic*)
 Pradhan, P.K. *et al.*, *Synthesis*, 2005, 1779-1782 (*synth, ir, pmr, cmr*)
 Wang, Z. *et al.*, *Mol. Cancer Ther.*, 2007, **6**, 2757-2765; 2008, **7**, 341-349 (*pharmacol*)
 Minich, D.M. *et al.*, *Nutr. Rev.*, 2007, **65**, 259-267 (*rev*)

Methylenebismehranine **M-437**
 [220327-18-4]



$C_{41}H_{52}N_4O_2$ 632.887
 Alkaloid from *Tabernaemontana bovina*.
 Oil. $[\alpha]_D^{28}$ -5.9 (c, 0.5 in MeOH).
 Lien, T.P. *et al.*, *Phytochemistry*, 1998, **49**, 1797-1799 (*isol, pmr, cmr*)


**N^b,N^{b'}-Methylenebismeloni-
 nium(2+)** **M-438**
 [135626-51-6]



$C_{39}H_{54}N_4^{2+}$ 578.882
 Alkaloid from *Melodinus cestroides* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ +71 (c, 1 in $CHCl_3$). Poss. artifact. CAS no. refers to dichloride.


Mehri, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 372-379 (*isol, uv, ir, pmr, ms, struct*)

**10,10'-Methylenebis[norval-
 lesamidine]** **M-439**
 [135626-50-5]




$C_{39}H_{52}N_4$ 576.867
 Alkaloid from *Melodinus cestroides* (Apocynaceae). $[\alpha]_D^{20}$ +140 (c, 1 in $CHCl_3$).
 Mehri, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 372 (*isol, uv, ir, pmr, cmr, struct*)

**1,1'-Methylenebis[2-phenazi-
 nol], 9CI** **M-440**
Di(2-hydroxy-1-phenazinylo)methane
 [73749-96-9]



$C_{25}H_{16}N_4O_2$ 404.427
 Isol. from *Pseudomonas aureofaciens*. Mp 300°.
 Neuenhaus, W. *et al.*, *Z. Naturforsch., B*, 1980, **35**, 385 (*synth*)


12,12'-Methylenedicytisine **M-441**
 [121290-68-4]



$C_{23}H_{28}N_4O_2$ 392.5
 Alkaloid from *Maackia amurensis*.

Kubo, H. *et al.*, *Thai J. Pharm. Sci.*, 1993, **17**, 171-173 (*isol*)
 Wang, Y.-H. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 641-645 (*isol*)

3,4-Methylenedioxybenzoic **M-442**
acid
1,3-Benzodioxole-5-carboxylic acid, 9CI.
Piperonylic acid. Heliotropic acid
 [94-53-1]



$C_8H_6O_4$ 166.133
 Isol. from Paracoto bark, from *Cinnamomum kanahirai*, *Piper longum* (long pepper) and *Melicope simplex*.
 Needles (EtOH), *cryst.* (H_2O). Mp 229°.
 pK_a 4.5 (25°, 5% EtOH). Sublimes in prisms.

►DF4912765
Me ester: [326-56-7]
 $C_9H_8O_4$ 180.16
 Needles or leaflets (petrol). Mp 273-274° part. dec. Steam-volatile.
Et ester: [6951-08-2]
 $C_{10}H_{10}O_4$ 194.187
 Prisms. Mp 18.5°. Bp 285°.

Amide: *3,4-Methylenedioxybenzamide*
 [4847-94-3]
 $C_8H_7NO_3$ 165.148
 Constit. of *Naravelia zeylanica*. Prisms or needles (H_2O). Mp 169°.


2-Phenylethylamide: *3,4-Methylenedioxy-N-phenethylbenzamide. Nantoamide*
 $C_{16}H_{15}NO_3$ 269.299
 Alkaloid from the rhizomes of *Begonia nantoensis*. Syrup. λ_{max} 293 (log ϵ 2.88) (MeOH).

Chloride: [25054-53-9]
 $C_8H_5ClO_3$ 184.578
 Cryst. Mp 80°. Bp₂₅ 155°.

Nitrile: *4-Cyano-1,2-methylenedioxybenzene*
 [4421-09-4]
 $C_8H_5NO_2$ 147.133
 Needles (H_2O). Mp 95°.

►TO2645000
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 212C; 457B (*ir*)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1111A; 1349B; 1533C (*nmr*)
Org. Synth., Coll. Vol., **2**, 1943, 538 (*synth*)
 Briggs, L.H. *et al.*, *J.C.S.*, 1950, 2376 (*isol*)
 Dobrowsky, A. *et al.*, *Monatsh. Chem.*, 1955, **86**, 325 (*synth*)
 Byrne, M.M. *et al.*, *J.C.S. (B)*, 1968, 809
 Šantavý, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 2418 (*uv*)
 Kobayashi, M. *et al.*, *CA*, 1990, **112**, 171755k (*Asarumin D*)
 Wu, P.-L. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 345-349 (*Nantoamide*)
 Jiang, X. *et al.*, *J.O.C.*, 2004, **69**, 2327-2331 (*amide, synth, pmr, cmr*)
 Jaroszewski, J.W. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 291-294 (*amide, isol, synth*)
 Li, J.-T. *et al.*, *Acta Cryst. E*, 2006, **62**, 1893-1894 (*cryst struct*)

3,4-Methylenedioxcinnamic **M-443**
acid 2-phenylethylamide




$C_{18}H_{17}NO_3$ 295.337

(E)-form [95925-01-2]
 Constit. of *Critoniella acuminata*. Cryst. (Et_2O /petrol). Mp 131°.

(Z)-form [95925-02-3]
 Constit. of *Critoniella acuminata*. Oil.
 Bohlmann, F. *et al.*, *Planta Med.*, 1984, **50**, 187-188 (*isol, uv, ir, pmr, ms*)

**3,4-Methylenedioxcinna-
 moyl piperidide** **M-444**
1-[3-(1,3-Benzodioxol-5-yl)-1-oxo-2-propenyl]piperidine, 9CI. 1-[3,4-(Methylenedioxy)cinnamoyl]piperidine, 8CI
 [23434-86-8]



C₁₅H₁₇NO₃ 259.304
Anticonvulsant. Shows serotonergic activity. Log P 2.1 (calc).


(E)-form

Ilepcimide, USAN. Antiepilepsirine
[82857-82-7]
Alkaloid from the wood of *Piper novae-hollandiae* (Piperaceae). Mp 89° (80-82°).
▶ LD₅₀ (mus, orl) >1000 mg/kg.
TM4568500

(Z)-form [111479-04-0]


Constit. of the roots of *Piper nigrum* (pepper).
Scholtz, M. *et al.*, *Ber.*, 1895, **28**, 1187 (synth)
Loder, J.W. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1531 (isol, uv)
Lin, G. *et al.*, *CA*, 1982, **97**, 102148z (cryst struct)
Lin, G.Q. *et al.*, *Biochem. Pharmacol.*, 1984, **33**, 3883 (pharmacol)
Yan, Q.-S. *et al.*, *J. Pharmacol. Exp. Ther.*, 1992, **261**, 652 (pharmacol)
Shen, Y. *et al.*, *Synth. Commun.*, 1992, **22**, 657 (synth, pmr, ir)
Wang, L. *et al.*, *Epilepsy Res.*, 1993, **15**, 1 (pharmacol)
Pei, Y.Q. *et al.*, *Res. Commun. Pharm. Toxicol.*, 1999, **4(II)**, 13-23 (rev, pharmacol)
Wei, K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1005-1009 (isol)

1,2-Methylenedioxy-6a,7-dehydroaporphine-10,11-quinone M-445
6,7-Dihydro-7-methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline-11,12-dione, 9CI
[38973-13-6]



C₁₈H₁₃NO₄ 307.305
Constit. of *Corydalis cava* (Papaveraceae). Blue-green cryst. Mp 218-220°.
Cava, M.P. *et al.*, *Tetrahedron*, 1972, **28**, 4299 (synth)
Preininger, V. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 294 (isol)

6,7-Methylenedioxyisoquinoline M-446
line
1,3-Dioxolo[4,5-g]isoquinoline, 9CI, 8CI.
Papraline
[269-44-3]




C₁₀H₇NO₂ 173.171
Alkaloid from aerial parts of *Fumaria indica* (Papaveraceae). Mp 127-128°.
Picrate: Mp 240-241°.
N-Oxide: [62761-43-7]
C₁₀H₇NO₃ 189.17

Beige flakes + 0.5 H₂O. Mp 221-224° (hemihydrate).

N-Me: 6,7-Methylenedioxy-N-methylisoquinolinium(1+)
C₁₁H₁₀NO₂[⊕] 188.206
Alkaloid from the stem bark of *Cryptocarya chinensis*. Yellow syrup. Counterion not specified. λ_{max} 253 ; 287 ; 296 ; 338 (MeOH).
1,2,3,4-Tetrahydro-N-Me: see Hydrohydrastinine, H-398
Späth, E. *et al.*, *Monatsh. Chem.*, 1929, **51**, 190
Boger, D.L. *et al.*, *Tetrahedron*, 1981, **37**, 3977 (synth)
Peet, N.P. *et al.*, *J. Het. Chem.*, 1987, **24**, 715 (ms)
Vollhardt, P.C. *et al.*, *Synthesis*, 1993, 579
Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1995, **40**, 593 (isol)
Lin, F.-W. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 157-159 (N-Me)

3,4-Methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid M-447
6-Nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. **Aristolochic acid II**. *Aristolochic acid B*. *Noraristolochic acid* [475-80-9]



C₁₆H₉NO₆ 311.25
Alkaloid from *Aristolochia argentina*, *Aristolochia clematitis*, *Aristolochia esperanzae*, *Aristolochia longa*, *Aristolochia manshuriensis*, *Aristolochia rotunda*, *Aristolochia pallida*, *Aristolochia siphon* (*Aristolochia durior*) and from the Chinese drugs Ching-Mu-Hsiang (roots of *Aristolochia debilis*) and Fang-chi (*Aristolochiaceae*). Yellow plates (MeOH/EtOAc), orange-yellow cryst. (Me₂CO). Mp 270-272° (264-268° dec.). λ_{max} 241; 305; 340; 450 (EtOH/NaOH) (Derep). λ_{max} 221 (ε 29500); 255 (ε 32400); 283; 314 (ε 11200); 389 (ε 6030) (EtOH) (Derep).

▶ SF8281600


Me ester: Aristolochic acid II methyl ester
[3580-54-9]
C₁₇H₁₁NO₆ 325.277
Alkaloid from the roots of *Aristolochia argentina* (*Aristolochiaceae*). Yellow needles (MeOH), cryst. (CHCl₃). Mp 277° (274-275°).

Alanine amide: 2-(Phenanthro[3,4-d]-1,3-dioxole-6-nitro-5-carboxamido)propanoic acid, 9CI
C₁₉H₁₄N₂O₇ 382.329
Alkaloid from the roots of *Aristolochia longa* (*Aristolochiaceae*). Pale yellow cryst. (MeOH)(as Me ester). Mp 252-253° dec. (Me ester). [α]_D -133 (c, 0.57 in CHCl₃) (Me ester).

Tseng, K.-F. *et al.*, *Huaxue Xuebao*, 1957, **23**, 156; *CA*, **52**, 13188g (isol)
Pailer, M. *et al.*, *Monatsh. Chem.*, 1957, **88**, 367; 1965, **96**, 863 (uv, ms, struct)

Kupchan, S.M. *et al.*, *J.O.C.*, 1965, **30**, 3792 (synth, uv, ir, ester)
Schunack, W. *et al.*, *Pharmazie*, 1965, **20**, 685 (isol)
Carboni, S. *et al.*, *Gazz. Chim. Ital.*, 1966, **96**, 641 (isol, ir)
Priestap, H.A. *et al.*, *An. Asoc. Quim. Argent.*, 1971, **59**, 245; *CA*, **76**, 43943w (isol)
Ruecker, G. *et al.*, *Planta Med.*, 1975, **27**, 68 (isol)
Cisowski, W. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1977, **51**, 2125 (isol, uv, ir)
Halm, I. *et al.*, *Herba Hung.*, 1980, **19**, 179; *CA*, **93**, 245336n (isol)
Podolov, B. *et al.*, *Acta Pharm. Jugosl.*, 1981, **31**, 249; *CA*, **98**, 82732j (isol, uv)
Priestap, H.A. *et al.*, *Phytochemistry*, 1982, **21**, 2755 (isol, ester)
De Pascual Teresa, J. *et al.*, *Phytochemistry*, 1983, **22**, 2745 (isol, ir, pmr, ms, synth, amide)
Nascimento, I.R. *et al.*, *Phytochemistry*, 2003, **63**, 953-957 (isol, pmr, cmr)

8,9-Methylenedioxyphenanthridine M-448
[1,3]Dioxolo[4,5-j]phenanthridine, 9CI.
Trisphaeridine. *Trisphaeridine*
[224-11-3]



C₁₄H₉NO₂ 223.231
Alkaloid from the bulbs and leaves of *Ungernia trisphaera*, aerial parts of *Pancreatum maritimum* and from *Pancreatum foetidum* (*Amaryllidaceae*). Mp 140-141° (138°). λ_{max} 210 ; 270 ; 337 ; 375 (MeOH) (Berdy).

N-Me: 5-Methyl[1,3]dioxolo[4,5-j]phenanthridinium(1+), 9CI. *5-Methyl-8,9-methylenedioxyphenanthridinium(1+)*. **Bicolorine**[†]
[129225-25-8]
C₁₅H₁₂NO₂[⊕] 238.265
Alkaloid from bulbs of *Narcissus bicolor* and bulbs and leaves of *Lapiedra martinii* (*Amaryllidaceae*). Yellow cryst. + 2H₂O (MeOH/Me₂CO) (as chloride). Mp 250-252° (chloride). Counterion not specified. The prev. descr. alkaloid 5,6-Dihydrobicolorine has been shown identical with Ismine, I-186.

3-Hydroxy: 3-Hydroxy-8,9-methylenedioxyphenanthridine. 3-Hydroxytrisphaeridine

C₁₄H₉NO₃ 239.23
Alkaloid from whole plants of *Crinum firmifolium* var. *hygrophilum* (*Amaryllidaceae*). Needles (MeOH). Mp 215-218° dec.


3-Methoxy: 3,4-dihydro-3-methoxy-8,9-methylenedioxyphenanthridine. 3,4-Dihydro-3-methoxytrisphaeridine

C₁₅H₁₃NO₃ 255.273
Alkaloid from the bulbs of *Hymenocallis x festalis*. Amorph. solid. [α]_D²⁵ -13 (c, 0.01 in CHCl₃). λ_{max} 252 (log ε 4.29); 300 (log ε 3.63); 312 (log ε 3.64); 329 (sh) (log ε 3.48); 346 (sh) (log ε

3.35) (MeOH).

- Allayarov, K. *et al.*, *Uzb. Khim. Zh.*, 1964, **8**, 46; *CA*, **61**, 4407a (isol)
- Kessar, S.V. *et al.*, *Tetrahedron*, 1973, **29**, 177 (synth)
- Krohn, K. *et al.*, *Annalen*, 1978, 608 (synth, uv, pmr)
- Prabhakar, S. *et al.*, *Chem. Comm.*, 1978, 884 (synth)
- Vdovin, A.D. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 356; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 279 (isol, uv, ir, pmr, ms)
- Viladomat, F. *et al.*, *Phytochemistry*, 1990, **29**, 1307 (*Bicolorine*)
- Suau, R. *et al.*, *Phytochemistry*, 1990, **29**, 1710 (*Bicolorine*)
- Banwell, M.G. *et al.*, *Aust. J. Chem.*, 1994, **47**, 2235 (synth)
- Cowden, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1746
- Razafimbelo, J. *et al.*, *Phytochemistry*, 1996, **41**, 323 (*3-Hydroxytrisphaeridine*)
- Shao, H.W. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 493-496 (synth)
- Harayama, T. *et al.*, *J.C.S. Perkin 1*, 2001, 523-528 (synth)
- Hohmann, J. *et al.*, *Fitoterapia*, 2002, **73**, 749-751 (*3,4-Dihydro-3-methoxytrisphaeridine*)


2-[4-(3,4-Methylenedioxyphenyl)butyl]-4(1H)-quinolinone M-449
[17889-77-9]

C₂₀H₁₉NO₃ 321.375Alkaloid from *Ruta graveolens* (rue) (Rutaceae). Cryst. (MeOH). Mp 224°.

- Novák, I. *et al.*, *Pharmazie*, 1965, **20**, 654 (isol, ir, pmr, ms, struct)
- Reisch, J. *et al.*, *Naturwissenschaften*, 1967, **54**, 517 (synth)

1-[12-(3,4-Methylenedioxyphenyl)-2,4,9,11-dodecatetraenoyl]-pyrrolidine M-450

12-(3,4-Methylenedioxyphenyl)-2,4,9,11-dodecatetraenoic acid pyrrolidide

C₂₃H₂₇NO₃ 365.471


(2E,4E,9E,11Z)-form [144205-85-6]

Alkaloid from *Piper guineense*.

- Gbewonyo, W.S.K. *et al.*, *J. Chromatogr.*, 1992, **607**, 105-111 (isol)

1-[17-(3,4-Methylenedioxyphenyl)-16-heptadecenoyl]pyrrolidine M-451

17-(3,4-Methylenedioxyphenyl)-16-heptadecenoic acid pyrrolidide

C₂₈H₄₃NO₃ 441.653


(E)-form [102965-08-2]

Alkaloid from *Piper amalago*.

- Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (isol)

1-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienoyl]pyrrolidine M-452

1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,6-heptadienyl]pyrrolidine, 9CI. 7-(3,4-Methylenedioxyphenyl)-2,6-heptadienoic acid pyrrolidide



(2E,6E)-form

C₁₈H₂₁NO₃ 299.369

(2E,6E)-form

Piperamide-C7:2 (2E,6E).*Sarmentosine*†

[112448-68-7]

Constit. of the fruits of pepper (*Piper nigrum*) and cha-plu (*Piper sarmentosum*) (Piperaceae). Exhibits larvicidal activity. Antihepatotoxic agent. Mp 77.5-79.5°.

2,3-Dihydro: 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-6-heptenyl]pyrrolidine, 9CI. 1-[7-(3,4-Methylenedioxyphenyl)-6-heptenoyl]pyrrolidine. 7-(3,4-Methylenedioxyphenyl)-6-heptenoic acid pyrrolidide. Piperamide-C7:1 (6E)

[117137-66-3]

C₁₈H₂₃NO₃ 301.385

Constit. of pepper fruits (*Piper nigrum*) (Piperaceae). Exhibits larvicidal activity. Oil.


- Likhitwitayawuid, K. *et al.*, *Tetrahedron*, 1987, **43**, 3689 (*Sarmentosine*)

Kiuchi, F. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2452 (isol, uv, ir, pmr, ms, struct)

Strunz, G.M. *et al.*, *Phytochemistry*, 1995, **39**, 731 (synth, *Sarmentosine*)

Bari, S.S. *et al.*, *J. Indian Chem. Soc.*, 1996, **73**, 520 (synth, *Sarmentosine*)

2-[6-(3,4-Methylenedioxyphenyl)hexyl]-4(1H)-quinolinone M-453

C₂₂H₂₃NO₃ 349.429

Alkaloid from the roots of *Ruta chalepensis*. Needles (CHCl₃). Mp 163°. λ_{max} 235 (log ε 2.61); 290 (log ε 1.97); 315 (log ε 2.14); 328 (log ε 2.12) (MeOH).

N-Me: 1-Methyl-2-[6-(3,4-methylenedioxyphenyl)hexyl]-4(1H)-quinolinone

C₂₃H₂₅NO₃ 363.455

Alkaloid from the leaves of *Ruta graveolens* (rue). Antifungal agent. Cryst. Mp 167-168°.

OH-form

Me ether: 4-Methoxy-2-[6-(3,4-methyle-

*nedioxyphenyl)hexyl]quinoline*C₂₃H₂₅NO₃ 363.455

Alkaloid from the roots of *Ruta chalepensis*. Oil. λ_{max} 227 (log ε 3.49); 285 (log ε 3.43); 311 (log ε 2.64) (MeOH).


El Sayed, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 995-997 (isol, pmr, cmr)

Oliiva, A. *et al.*, *J. Agric. Food Chem.*, 2003, **51**, 890-896 (*N-Me*)

Back, T.G. *et al.*, *J.O.C.*, 2003, **68**, 2223-2233 (synth)

9-(3,4-Methylenedioxyphenyl)-2,4-nonadienoic acid M-454

9-Piperonyl-2,4-nonadienoic acid

C₁₆H₁₈O₄ 274.316

(E,E)-form

2-Methylpropylamide:C₂₀H₂₇NO₃ 329.438

Isol. from root and stems of *Ottonia anisum* (preferred genus name *Piper*).

2,3,4,5-Tetrahydro: 1,3-Benzodioxole-5-nonanoic acid. 9-(3,4-Methylenedioxyphenyl)nonanoic acid. 9-Piperonylnonanoic acid

C₁₆H₂₂O₄ 278.347

Constit. of *Viola surinamensis*. Viscous oil.


Giesbrecht, A.M. *et al.*, *Planta Med.*, 1981, **43**, 375; *CA*, **96**, 119009

Lopes, N.P. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1255-1257

(*Methylenedioxyphenylnonanoic acid*)

1-[9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienoyl]pyrrolidine M-455

1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,8-nonatrienyl]pyrrolidine, 9CI. 9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienoic acid pyrrolidide

C₂₀H₂₃NO₃ 325.407

(2E,4E,8E)-form

Piperamide-C9:3 (2E,4E,8E)

[117137-68-5]

Constit. of pepper fruits (*Piper nigrum*, Piperaceae). Yellow needles. Mp 105-106°.

4,5-Dihydro: 1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,8-nonadienyl]pyrrolidine, 9CI. 1-[9-(3,4-Methylenedioxyphenyl)-2,8-nonadienoyl]pyrrolidine. 9-(3,4-Methylenedioxyphenyl)-2,8-nonadienoic acid pyrrolidide. Piperamide-C9:2 (2E,8E).

Brachyamide B

[117137-67-4]

C₂₀H₂₅NO₃ 327.422

Constit. of pepper fruits (*Piper nigrum*) and *Piper brachystachyum* (Piperaceae). Exhibits larvicidal activity. Oil.

2,3,4,5-Tetrahydro: 1-[9-(1,3-Benzodiox-

*ol-5-yl)-1-oxo-8-nonenyl]pyrrolidine, 9CI. 1-[9-(3,4-Methylenedioxyphenyl)-8-nonenyl]pyrrolidine. 9-(3,4-Methylenedioxyphenyl)-8-nonenol acid pyrrolidide. **Piperamide-C9:1 (8E)**. Tricholein [62510-52-5] C₂₀H₂₇NO₃ 329.438*

Constit. of pepper (*Piper nigrum*) and *Piper trichostachyon* (Piperaceae). Exhibits larvicidal activity. Oil.

Singh, J. *et al.*, *Phytochemistry*, 1976, **15**, 2018 (*Piperamide-C9:1*)


Vig, O.P. *et al.*, *Indian J. Chem., Sect. B*, 1979, **17**, 560 (*Piperamide-C9:1*)

Kiuchi, F. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2452 (*isol, uv, ir, pmr, cmr, ms, struct*)

Koul, S.K. *et al.*, *Phytochemistry*, 1988, **27**, 3523 (*Piperamide-C9:2*)

Strunz, G.M. *et al.*, *Can. J. Chem.*, 1996, **74**, 419 (*synth*)

1-[5-(3,4-Methylenedioxyphenyl)-2-pentenol]pyrrolidine M-456
5-(3,4-Methylenedioxyphenyl)-2-pentenoic acid pyrrolidide



C₁₆H₁₉NO₃ 273.331

(E)-form Piperamide-C5:1 (2E)

[117137-65-2]


Constit. of pepper fruits (*Piper nigrum*, Piperaceae). Oil.

Kiuchi, F. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2452; 2647 (*isol, uv, ir, pmr, ms, struct*)

Alecio, A.C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 637-639 (*isol, pmr, cmr*)

3-(3,4-Methylenedioxyphenyl)-2-propenoic acid M-457

3-(1,3-Benzodioxol-5-yl)-2-propenoic acid, 9CI. 3,4-Methylenedioxyacrylic acid, 8CI. 3-(3,4-Methylenedioxyphenyl)acrylic acid. Piperonylideneacetic acid [2373-80-0]



(E)-form

C₁₀H₈O₄ 192.171

(E)-form [38489-76-8]

Isol. from leaves of *Piper tuberculatum*. Also isol. from *Brombya platynema*. Needles (EtOH). Mp 247° (242°). pK_a 4.6 (25°).

Me ester: [40918-96-5] C₁₁H₁₀O₄ 206.198

Plates (MeOH). Mp 133-134°.

Amide:

C₁₀H₉NO₃ 191.186

Needles (EtOH). Mp 184-185°.

2-Methylpropylamide: 3-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2-propenamide. N-Isobutyl-3,4-methylenedioxyacrylamide. trans-Fagaramide [495-86-3] C₁₄H₁₇NO₃ 247.293

Alkaloid from the bark of *Zanthoxylum macrophylla* and *Anthocleista vogelii*, the root of *Fagara tessmannii*, the root bark of *Fagara zanthoxyloides* and the wood of *Piper novae-hollandiae* (Rutaceae, Piperaceae). Plates (EtOAc). Mp 119.5° (softens at 105°).

Penylamide: N-(3,4-Methylenedioxyphenyl)pentylamine. 3,4-Methylenedioxy-N-pentylcinnamide [73073-94-6] C₁₅H₁₉NO₃ 261.32

Alkaloid from *Piper amalago*. Config. not determined but most probably *E*-.

3-Methylbutylamide: N-(3,4-Methylenedioxyphenyl)isopentylamine. N-Isopentyl-3,4-methylenedioxyacrylamide [23795-28-0] C₁₅H₁₉NO₃ 261.32

Alkaloid from *Piper amalago*. Config. not determined but most probably *E*-.

Piperidide: see 3,4-Methylenedioxyacrylamoyl piperidide, M-444

(Z)-form

Cryst. (CS₂). Mp 99-100°. pK_a 4.94 (25°).

Amide:

Plates (Et₂O). Mp 131°.

2-Methylpropylamide: cis-Fagaramide

[92449-51-9] C₁₄H₁₇NO₃ 247.293

Alkaloid from the leaves of *Zanthoxylum schinifolium*, *Fagara macrophylla* and *Fagara zanthoxyloides*. Light yellow needles (Et₂O). Mp 113-115°. λ_{max} 217 (log ε 3.72); 274 (log ε 3.56); 309 (log ε 3.58) (MeOH).

[16386-34-8]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 183D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1057A (*nmr*)

Thoms, H. *et al.*, *Ber.*, 1911, **44**, 3717 (*isol, struct, synth, derivs*)

Roth, W.A. *et al.*, *Ber.*, 1913, **46**, 260

Goodson, J.A. *et al.*, *Biochem. J.*, 1921, **15**, 123 (*Fagaramide*)

Pearl, I.A. *et al.*, *J.O.C.*, 1951, **16**, 216 (*synth, uv*)

Simmonds, N.W. *et al.*, *Nature (London)*, 1956, **178**, 752 (*isol*)

Loder, J.W. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1531 (*Fagaramide*)

Santavy, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 1825 (*uv*)

Takemoto, T. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 1161 (*synth, ir, pmr*)

Okorie, D.A. *et al.*, *Phytochemistry*, 1976, **15**, 1799 (*Fagaramide, isol, ir, pmr, ms*)

Coutrot, P. *et al.*, *Synthesis*, 1978, 133 (*synth*)

Nagao, Y. *et al.*, *Tet. Lett.*, 1980, **21**, 841 (*Fagaramide, synth*)

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (*isol, amides*)

Terada, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2437 (*synth*)

Patra, A. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 734 (*cmr*)

Parsons, I.C. *et al.*, *Phytochemistry*, 1993, **33**, 479 (*isol, pmr*)

Cheng, M.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2002, **49**, 125-128 (*cis-Fagaramide*)


Chaaib, F. *et al.*, *Planta Med.*, 2003, **69**, 316-320 (*cis-Fagaramide*)

2-(3,4-Methylenedioxyphenyl)quinoline M-458

2-(1,3-Benzodioxol-5-yl)quinoline, 9CI.

Dubamine

[6808-65-7]



C₁₆H₁₁NO₂ 249.268

Alkaloid from *Haplophyllum dubium* and *Haplophyllum latifolium* (Rutaceae).

Cryst. (MeOH). Mp 96-97°.

Sidyakin, G.P. *et al.*, *Uzb. Khim. Zh.*, 1962, **6**, 56-59; *CA*, **58**, 4608 (*isol, synth*)

Koyama, J. *et al.*, *Chem. Express*, 1991, **6**, 197-200; 1992, **7**, 321-324 (*synth*)

Ali, N.M. *et al.*, *Tetrahedron*, 1992, **48**, 8117-8126 (*synth*)

Mahanty, J.S. *et al.*, *Tetrahedron*, 1997, **53**, 13397-13418 (*synth, ir, uv, pmr*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1038-1039 (*synth, ir, pmr*)

2-Methylene-4-oxopentane-dioic acid M-459

α-Keto-γ-methyleneglutamic acid

[55601-65-5]

H₂C=C(COOH)CH₂COCOOH

C₆H₆O₅ 158.11

Found in tulip *Tulipa gesneriana*, *Lilium* spp. and peanut (*Arachis hypogaea*).

2,4-Dinitrophenylhydrazone: Mp 228° dec.

1-Amide:

C₆H₇NO₄ 157.126


Found in *Tulipa gesneriana*. Called the 5-amide in the original lit.

Towers, G.H.N. *et al.*, *J.A.C.S.*, 1954, **76**, 1959

Fowden, L. *et al.*, *Biochem. J.*, 1955, **59**, 228

11-Methylenepregnane-3,20-diamine M-460


3,20-Diamino-11-methylenepregnane



C₂₂H₃₈N₂ 330.556

(3α,5α,20S)-form


N³,N²⁰-Tri-Me, N³-benzoyl: **11-Methylenepachysamine H** [282726-60-7]

C₃₂H₄₈N₂O 476.744Alkaloid from *Pachysandra procumbens*. Needles (MeOH). Mp 185°. [α]_D²⁰ +60.2 (c, 0.04 in CHCl₃). λ_{\max} 241 (log ϵ 3.86) (MeOH).Chang, L.C. *et al.*, *Tetrahedron*, 2000, **56**, 3133-3138 (*11-Methylenepachysamine H*)**1-Methylenepyrrolizidine M-461***Hexahydro-1-methylene-1H-pyrrolizine, 8CI*
[6029-70-5]

(S)-form

C₈H₁₃N 123.197**(S)-(?) -form [34539-78-1]**Alkaloid from *Crotalaria anagyroides*, *Crotalaria damarensis*, *Crotalaria rhodesiae*, *Crotalaria stolzii*, *Crotalaria verrucosa*, *Crotalaria lachnophora* and *Crotalaria natalitia* (Fabaceae). Hygroscopic liq. Bp₁₇₀ 120°. [α]_D²⁰ -100 (c, 0.68 in EtOH). Alkaloid from *C. damarensis* was opt. pure, but samples from *C. anagyroides* were partly racemic (Fabaceae).*Picrate*: Mp 217-218°. [α]_D -14.9 (c, 0.94 in Me₂CO).**N-Oxide: 1-Methylenepyrrolizidine N-oxide**

[81649-43-6]

C₈H₁₃NO 139.197Alkaloid from seeds of *Crotalaria anagyroides* (Fabaceae).**(±)-form**Bp₁₆₈ 114-116°. n_D^{20} 1.4880.Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1959, **12**, 255; 1962, **15**, 328 (*isol, ir, uv, resoln*)Kochetkov, N.K. *et al.*, *Tet. Lett.*, 1961, 92 (*synth, resoln*)Sethi, M.L. *et al.*, *Planta Med.*, 1964, **12**, 173 (*isol*)Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3652 (*cd*)Rastogi, R. *et al.*, *Pharmazie*, 1982, **37**, 75 (*isol, oxide*)Keusenkothen, P.F. *et al.*, *J.C.S. Perkin 1*, 1994, 2485 (*synth*)**6-(1-Methylethyl)-3-(2-****methylpropyl)-2(1H)-pyrazinone M-462***3-Isobutyl-6-isopropyl-2(1H)-pyrazinone. Deoxydehydromutaaspergillilic acid*C₁₁H₁₈N₂O 194.276Metab. of *Aspergillus flavus*. Mp 111°.*1-Hydroxy: 1-Hydroxy-6-(1-methylethyl)-3-(2-methylpropyl)-2(1H)-pyrazinone. 1-Hydroxy-3-isobutyl-6-isopropyl-2(1H)-pyrazinone*C₁₁H₁₈N₂O₂ 210.275Metab. from *Aspergillus sojae*. Mp 94-95°. λ_{\max} 232; 326 (MeOH) (Berdy).► LD₅₀ (mus, ipr) 62.5 mg/kg.**1'-Hydroxy: Deoxymutaaspergillilic acid**C₁₁H₁₈N₂O₂ 210.275Metab. of *Aspergillus sojae*. Mp 133-134.5°.**1,1'-Dihydroxy: 1-Hydroxy-6-(1-hydroxy-1-methylethyl)-3-(2-methylpropyl)-2(1H)-pyrazinone. Mutaaspergillilic acid**

[15272-17-0]

C₁₁H₁₈N₂O₃ 226.275Metab. of *Aspergillus oryzae*. Antibiotic. Pale yellow needles. Sol. MeOH, C₆H₆; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 173-174° (167-168°) dec. pK_a 4.8. λ_{\max} 232 (ϵ 12500); 332 (ϵ 7300) (MeOH) (Berdy). λ_{\max} 233 (ϵ 6100); 334 (ϵ 7200) (EtOH) (Berdy). λ_{\max} 242 (ϵ 5100); 335 (ϵ 8600) (H₂O) (Berdy).► LD₅₀ (mus, ipr) 100 mg/kg. UQ4405500Sugiyama, M. *et al.*, *Tet. Lett.*, 1967, 845(Mutaaspergillilic acid, *struct, synth*)Ohta, A. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 20 (*synth*)Yamamoto, Y. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 185 (*rev, bibl, derivs*)Okada, Y. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2259-2262 (*synth*)**N-Methylformamide, 9CI, M-463***Formylmethylamine. NSC 3051*

[123-39-7]


HCONHMe

C₂H₅NO 59.068Isol. from the red alga *Erythrophyllum delesserioides*. Antineoplastic agent. Sol. H₂O, EtOH; insol. Et₂O. d_4^{19} 1.01. Mp -3.8°. Bp 180-185°. n_D^{20} 1.4319.► Highly flammable, fl. p. <22°. Eye irritant. LD₅₀ (rat, orl) 4000 mg/kg. Exp. reprod. and teratogenic effects. Hepatotoxic. LQ3000000*N-Bromo: N-Bromo-N-methylformamide*C₂H₄BrNO 137.964


Mp 81-83°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 753D (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 1227C (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 778D (*ir*)Gautier, A. *et al.*, *Annalen*, 1869, **151**, 241 (*synth*)U.K. Pat., 1926, 252 460; CA, **21**, 2273 (*synth*)Russell, R.A. *et al.*, *Spectrochim. Acta*, 1956, **8**, 138 (*ir*)Suzuki, I. *et al.*, *Bull. Chem. Soc. Jpn.*, 1962, **35**, 540 (*ir*)LaPlanche, L.A. *et al.*, *J.A.C.S.*, 1964, **86**, 337 (*pmr*)Nielsen, E.B. *et al.*, *J. Phys. Chem.*, 1967, **71**, 2297 (*uv*)Dorman, D.E. *et al.*, *J.O.C.*, 1973, **38**, 1719 (*cmr*)Warren, J.P. *et al.*, *J. Phys. Chem.*, 1974, **78**, 2507 (*N-15 nmr*)Izac, R.R. *et al.*, *Phytochemistry*, 1982, **21**, 229 (*isol*)Ohtaki, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1986, **59**, 271 (*struct, props, bibl*)Ettinger, D.S. *et al.*, *Invest. New Drugs*, 1990, **8**, 183-185 (*phase II study, pharmacol*)McGuire, W.P. *et al.*, *Invest. New Drugs*, 1990, **8**, 191-194; 195-197 (*phase II study, pharmacol*)Fujisaki, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1993, **66**, 2426 (*N-bromo*)*Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn.*, (ed. Snyder, R.), Elsevier, Volume 2, 1990, 169Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 0813Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MKG500**4-Methyl-10H-furo[3,2-****a]carbazole, 9CI M-464***Fuostifoline*

[133740-44-0]

C₁₅H₁₁NO 221.258Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Cryst. (cyclohexane). Mp 174-175° (*synthetic*).Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1548 (*isol, uv, ir, ms, pmr, struct*)Hagiwara, H. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1948-1949 (*synth, pmr*)Beccalli, E.M. *et al.*, *Tetrahedron*, 1998, **54**, 11675-11682 (*synth*)Soos, T. *et al.*, *Tet. Lett.*, 1999, **40**, 8607-8609 (*synth*)Knölker, H.-J. *et al.*, *Synthesis*, 2000, 2131-2136 (*synth, ir, uv, pmr, cmr, ms*)Yasuhara, A. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 143-145 (*synth*)**9-Methyl-6H-furo[2,3-c]carbazole, 9CI M-465***Eustifoline D*

[133740-43-9]

C₁₅H₁₁NO 221.258Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.Ho, C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1548 (*isol, pmr, cmr, struct*)Lebold, T.P. *et al.*, *Org. Lett.*, 2007, **9**, 1883-1886 (*synth*)**N-Methylguanidine M-466**

[471-29-4]

HN=C(NH₂)NHMeC₂H₇N₃ 73.097

Prod. of putrefaction. Solid. Easily dec. Strong base.

► MF3683000

Hydrochloride: [22661-87-6]

V. hygroscopic cryst.

Nitrate: [546-82-7]

Cryst. (EtOH). Mp 151°.

Sulfate salt (2:1): [598-12-9]
Cryst. (H₂O). Mp 239-240°.

Picrate: [1609-02-5]
Yellow plates or needles (H₂O). Mp 200°.

Ac:
C₄H₉N₃O 115.135
Mp 172° (as hydrochloride).

N-Nitroso: N-Methyl-N-nitrosoguanidine
[4262-56-0]

C₂H₆N₄O 102.096
In former use as synth. precursor of
Diazomethane.

► Exp. carcinogen.

N-(4-Methoxybenzoyl): N-(4-Methoxybenzoyl)-N'-methylguanidine
[387867-99-4]

C₁₀H₁₃N₃O₂ 207.232
Isol. from the ascidian *Polycarpa aurata*. Oil. λ_{max} 276 (ε 4754) (MeOH).

[21770-81-0]

Aldrich Library of FT-IR Spectra, 1st edn.,
1985, 1, 818D (ir)

Aldrich Library of 13C and 1H FT NMR
Spectra, 1992, 1, 1328B (nmr)

Werner, E.A. et al., J.C.S., 1922, 121, 1790-
1794 (synth)

Phillipi, E. et al., Ber., 1927, 60, 2120-2122
(synth)

Davis, T.L. et al., J.A.C.S., 1937, 59, 2112-
2115 (synth)

Spasskaya, R.I. et al., CA, 1968, 68, 39059
(synth)


Beynon, J.H. et al., Org. Mass Spectrom.,
1968, 1, 169-187 (ms)

Wessels, M. et al., J. Nat. Prod., 2001, 64,
1556-1558 (4-methoxybenzoyl, isol)

Lewis, R.J. et al., Sax's Dangerous Properties
of Industrial Materials, 10th edn., J. Wiley,
2000, MK1750

1-Methylguanaine, 8CI M-467

2-Amino-1,7-dihydro-1-methyl-6H-purin-
6-one, 9CI
[938-85-2]



C₆H₇N₅O 165.154
Isol. from RNA hydrolysates, e.g from
yeast, *Saccharomyces cerevisiae*. Present
in Chinese cabbage, *Brassica chinensis*,
and other plants. Plates or needles
(AcOH aq.). Mp 350°.

7H-form

7-Me: 1,7-Dimethylguanaine
[26758-00-9]
C₇H₉N₅O 179.181
Cryst. (EtOH aq.). Mp 338-340° (330-
331°).

9H-form

9-Me: 1,9-Dimethylguanaine
[42484-34-4]
C₇H₉N₅O 179.181
Needles (EtOH). Mp 287°.
Traube, W. et al., Ber., 1913, 46, 3845
(synth)

Dunn, D.B. et al., Biochem. J., 1959, 72, 294;
1963, 88, 34P (isol)

Pfleiderer, W. et al., Annalen, 1961, 647, 167
(synth)

Elion, G.B. et al., J.O.C., 1962, 27, 2478
(synth)


Rice, J.M. et al., J.A.C.S., 1967, 89, 2719 (ms)

Le Breton, P.R. et al., J.A.C.S., 1990, 112,
2138 (synth, pmr, uv, cmr)

Kasende, O.E. et al., Spectrosc. Lett., 1997, 30,
415-431 (Dimethylguanines, ir, Raman)

3-Methylguanaine, 8CI M-468

2-Amino-3,7-dihydro-3-methyl-6H-purin-
6-one, 9CI
[2958-98-7]



C₆H₇N₅O 165.154
Cryst. (H₂O). Mp 375-377°.

7H-form

7-Me: 3,7-Dimethylguanaine
[19143-67-0]
C₇H₉N₅O 179.181
Isol. from the sponge *Zyzzya fuliginosa*.
Cryst. (EtOH aq.). Mp 327-333°
dec.

9H-form

9-Me: 3,9-Dimethylguanaine
[67513-75-1]
C₇H₉N₅O 179.181
Cryst. (H₂O). Mp 326°.

Townsend, L.B. et al., J.A.C.S., 1962, 84, 3008
(synth, uv)

Elion, G.B. et al., J.O.C., 1962, 27, 2478
(synth)

Rice, J.M. et al., J.A.C.S., 1967, 89, 2719 (ms)

Wiley, D.W. et al., J.O.C., 1976, 41, 1889
(synth, uv, nmr)


Abola, J.E. et al., Tet. Lett., 1976, 3483 (cryst
struct)

Itaya, T. et al., Chem. Pharm. Bull., 1982, 30,
3392 (synth, uv, nmr)

Tasdemir, D. et al., Chem. Pharm. Bull., 2001,
49, 1628-1630 (3,7-Dimethylguanaine, isol,
pmr, cmr, N-15 nmr)

7-Methylguanaine, 9CI M-469

2-Amino-1,7-dihydro-7-methyl-6H-purin-
6-one, 9CI. Epiguanine
[578-76-7]



C₆H₇N₅O 165.154
Present in biol. systems, prob. as a dec.
prod. of 7-Methylguanosine. Urinary
marker of exposure to methylating
agents. Needles (H₂O). Mp 370°.

► MF8464000

N²,N²-Di-Me: N²,N²,7-Trimethylguanaine
[92333-92-1]

C₈H₁₁N₅O 193.208
Isol. from the ascidian *Lissoclinum
notti*. Cryst. (H₂O). Mp 320° dec.
λ_{max} 252 (log ε 3.8); 294 (log ε 3.6)
(MeOH).

Fischer, E. et al., Ber., 1898, 31, 544 (synth)

Gulland, J.M. et al., J.C.S., 1938, 692
(synth)


Bredereck, H. et al., Chem. Biol., 1962, 95,
1812-1819 (N²,N²,7-Trimethylguanaine,
synth)

Rice, J.M. et al., J.A.C.S., 1967, 89, 2719 (ms)

Shuker, D.E.G. et al., Arch. Toxicol., Suppl.,
1989, 13, 55 (use)

Pearce, A.N. et al., Nat. Prod. Lett., 2001, 15,
237-241 (N²,N²,7-Trimethylguanaine, isol)

2-Methyl-6-(8-heptadecenyl)-piperidine M-470




C₂₃H₄₅N 335.615
Minor component of the venoms of
Solenopsis spp. ants. See also refs. under
2-Methyl-6-tridecylpiperidine, M-580
and 2-Methyl-6-undecylpiperidine, M-
584.

Jones, T.H. et al., Tetrahedron, 1982, 38, 1949
(isol, struct, bibl)

8-(5-Methylhexyl)-1H,3H-[1,3,4]oxadiazolo[3,4-a][1,2,4]triazin-1-one, 9CI M-471

[897363-61-0]




C₁₂H₁₉N₃O₂ 237.301
Prod. by a myxobacterium sp. HK1.
Cytotoxic.

Lee, H.-K. et al., J. Microbiol. Biotechnol.,
2005, 15, 734-739 (isol, struct)

N-Methylhydrastine M-472

3-[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]-6,7-dimethoxy-
1(3H)-isobenzofuranone, 9CI



C₂₂H₂₃NO₆ 397.427

(E)-form [55922-35-5]

Alkaloid from *Fumaria parviflora*, *Fumaria vaillantii*, *Fumaria schleicheri* and *Corydalis lutea* (Papaveraceae). Mp 156°.

May be an artifact.

(Z)-form [76202-53-4]


Cryst. (CHCl₃). Mp 155-156°.

Forgacs, P. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1973, **276**, 105; 1974, **279**, 855 (synth, occur)

Blaskó, G. et al., *J.O.C.*, 1982, **47**, 880 (synth, uv, ir, pmr, ms, struct)

4(5)-Methylimidazole, 9CI M-473

[822-36-6]



C₄H₆N₂ 82.105

Isol. from flowers of *Sophora flavescens*. Cryst. Sol. H₂O, EtOH. Mp 56°. Bp 263° Bp_{0.2} 120-125°. pK_a 7.52 (25°). Prob. an artifact.

▶ LD₅₀ (mus, orl) 370 mg/kg. NI7350000

Picrate:

Yellow cryst. (H₂O). Mp 159-160°.

1H-form

N-Benzoyl:

C₁₁H₁₀N₂O 186.213

Needles. V. sol. EtOH, Me₂CO, C₆H₆, sol. petrol. Mp 54-55°.

N-Me: 1,4-Dimethyl-1H-imidazole

[6338-45-0]

C₅H₈N₂ 96.132

Bp 198-200°.

l-Me; hydrochloride: Mp 168-169°.

N-Hydroxy: 1-Hydroxy-4-methyl-1H-imidazole

C₄H₆N₂O 98.104

Semi cryst.

3H-form

N-Me: 1,5-Dimethyl-1H-imidazole

[10447-93-5]

C₅H₈N₂ 96.132

Bp 220-222°.

N-Me; hydrochloride: Mp 194-195°.

N-Hydroxy: 1-Hydroxy-5-methyl-1H-imidazole

C₄H₆N₂O 98.104

Cryst. Mp 159-160°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 613A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 76C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1484C (ir)

Gabriel, S. et al., *Ber.*, 1893, **26**, 2205

Pyman, F.L. et al., *J.C.S.*, 1922, **121**, 2621 (synth, deriv)

Turner, R.A. et al., *J.A.C.S.*, 1949, **71**, 2801

Bowie, J.H. et al., *Aust. J. Chem.*, 1967, **20**, 1613 (ms)

Martin, P.K. et al., *J.O.C.*, 1968, **33**, 3758 (synth, pmr, ms, derivs)

Matthews, H.R. et al., *J.A.C.S.*, 1973, **95**, 2297 (nmr, derivs)

Sattler, H.J. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1975, **308**, 795 (nmr, tautom)

Murakoshi, I. et al., *Phytochemistry*, 1982, **21**, 2379 (isol)


Eriksen, B.L. et al., *J.O.C.*, 2001, **66**, 8344-8348 (*N*-hydroxy derivs, synth, pmr, cmr)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MKU000

3-Methyl-1H-indole, 9CI M-474

β-Methylindole. Scatole. Skatole. FEMA 3019

[83-34-1]



C₉H₉N 131.177

Occurs in coal tar, tobacco smoke, various plants and faeces. Ubiquitous as a bacterial dec. prod. of proteins; may also occur in plants as a genuine alkaloid. Odour enhancer in perfumery. A major contributor to boar taint in meat products from uncastrated male pigs. Gut flora tryptophan metabolite. Causes acute bovine pulmonary oedema and emphysema. Leaflets (petrol) with faecal odour at high levels becoming warm and fruity at low concns. Mp 95°. Bp₇₅₅ 265-266°. pK_{a1} -4.55; pK_{a2} 16.6 (25°, acid, KOH aq.).

▶ LD₅₀ (rat, orl) 3450 mg/kg. Exp. pneumotoxic and splenotoxic. Ruminants are particularly sensitive. NM0350000

Picrate: Mp 174°.

N-Ac: [23543-66-0]

C₁₁H₁₁NO 173.214

Mp 68°.

▶ NL3677500

N-Me: 1,3-Dimethyl-1H-indole, 9CI

[875-30-9]

C₁₀H₁₁N 145.204

Liq. Bp 257-260° Bp₇ 119-120°. pK_a -3.37 (25°, HClO₄ aq.).

N-Me, picrate:

Cryst. Mp 145°.

N-Ph: 3-Methyl-1-phenyl-1H-indole

[112817-88-6]

C₁₅H₁₃N 207.274

Bp₂ 150°.

N-Carboxy: 3-Methyl-1H-indole-1-carboxylic acid, 9CI

[65610-64-2]

C₁₀H₉NO₂ 175.187

Cryst. Mp 129° dec. Decarboxylates at Mp.

N-Methoxycarbonyl:

C₁₁H₁₁NO₂ 189.213

Bp 289° Bp_{0.5} 101-102°.

N-Ethoxycarbonyl:

C₁₂H₁₃NO₂ 203.24

Bp_{0.5} 105-110°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 653D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 122A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1497D (ir)

Fischer, E. et al., *Annalen*, 1886, **236**, 126 (synth)

Snyder, H.R. et al., *J.A.C.S.*, 1948, **70**, 1703 (*N-Me*)

Andrisano, R. et al., *Gazz. Chim. Ital.*, 1957, **87**, 949 (uv)

Stowe, B.B. et al., *Prog. Chem. Org. Nat. Prod.*, 1959, **17**, 254 (occur)

Rees, C.W. et al., *J.C.S.*, 1964, 938 (*N*-carboxy)

Safe, S. et al., *Org. Mass Spectrom.*, 1972, **6**, 33 (ms)

Opdyke, D.L.J. et al., *Food Cosmet. Toxicol.*, 1976, **14**, 863 (rev, tox)

Bergman, J. et al., *J. Het. Chem.*, 1977, **14**, 1123 (*N*-carboxy)

Janda, M. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 3278 (*N-Me*)

Ito, Y. et al., *Bull. Chem. Soc. Jpn.*, 1984, **57**, 73 (synth)

Carlson, J.R. et al., *J. Toxicol., Toxin Rev.*, 1986, **5**, 217 (rev)

Nishio, T. et al., *J.O.C.*, 1988, **53**, 1323 (deriv, synth, ir, pmr, cmr)

Yost, G.S. et al., *Chem. Res. Toxicol.*, 1989, **2**, 273 (tox, rev)

Tsuji, Y. et al., *J.O.C.*, 1990, **55**, 580 (synth, pmr, cmr)

Updyke, L.W. et al., *Toxicol. Appl. Pharmacol.*, 1991, **109**, 375 (tox)

Black, D.St.C. et al., *Aust. J. Chem.*, 1992, **45**, 1879 (*N-Me*, pmr, ms)

Bray, T.M. et al., *Annu. Rev. Pharmacol. Toxicol.*, 1994, **34**, 91 (rev)

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 726

Tokmakov, G.P. et al., *Tetrahedron*, 1995, **51**, 2091 (*N-Ph*)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2513-2514

Baek, C. et al., *J. Agric. Food Chem.*, 1997, **45**, 2332-2340 (occur)

Larock, R.C. et al., *J.O.C.*, 1998, **63**, 7652-7662 (*N-Ac*, synth, cmr)


Barluenga, J. et al., *Chem. Eur. J.*, 2002, **8**, 2034-2046 (*N-Me*, synth, pmr, cmr)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MKV750

7-Methyl-1H-indole-3-carboxaldehyde M-475

3-Formyl-7-methylindole

[4771-50-0]



C₁₀H₉NO 159.187

Cryst. (CH₂Cl₂/EtOAc or EtOH). Mp 212-214° (206-208°).

N-Me: 1,7-Dimethyl-1H-indole-3-carboxaldehyde. 3-Formyl-1,7-dimethylindole [164353-61-1]


C₁₁H₁₁NO 173.214

Metab. of the marine cyanobacterium *Lyngbya majuscula*.

Boyd, W.J. et al., *Biochem. J.*, 1935, **29**, 555-561 (synth)


Belg. Pat., 1964, 628 441; *CA*, **60**, 13226 (synth)
 Heath-Brown, B. *et al.*, *J.C.S.*, 1965, 7165-7178 (synth)
 Todd, J.S. *et al.*, *J. Nat. Prod.*, 1995, **58**, 586 (isol, uv, ir, pmr, cmr, struct, deriv)

3-Methyl-1H-indole-2-carboxylic acid, 9CI M-476
Skatole-2-carboxylic acid
 [10590-73-5]



$C_{10}H_9NO_2$ 175.187
 Prod. by *Micromonospora* sp. P1068. Needles (AcOH). Sol. EtOH, Et₂O, C₆H₆, spar. sol. H₂O, petrol. Mp 164-165°.
Me ester: [104711-29-7]
 $C_{11}H_{11}NO_2$ 189.213
 Cryst. (MeOH). Mp 148°.
Et ester: [26304-51-8]
 $C_{12}H_{13}NO_2$ 203.24
 Needles (EtOH). Sol. Et₂O, C₆H₆. Mp 133-134°.
N-Me, Me ester: [69913-90-2]
 $C_{12}H_{13}NO_2$ 203.24
 Cryst. solid (petrol). Mp 76-78°.
N-Benzyl, Me ester:
 $C_{18}H_{17}NO_2$ 279.338
 Cryst. solid (petrol). Mp 43-45°.
 Perkin, W.H. *et al.*, *J.C.S.*, 1921, **119**, 1602-1642 (synth)
 Whalley, W.B. *et al.*, *J.C.S.*, 1954, 1651-1653 (synth)
 Andrisano, R. *et al.*, *Gazz. Chim. Ital.*, 1957, **87**, 949-970 (uv)
 Jones, C.D. *et al.*, *J.O.C.*, 1972, **37**, 3624-3625 (synth)
 Wadia, M.S. *et al.*, *Synthesis*, 1987, 401-404 (*Et ester, synth, ir, pmr*)
 Katritzky, A.R. *et al.*, *Synth. Commun.*, 1988, **18**, 1151-1158 (*Me ester, synth, ir, pmr*)
 Murakami, Y. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1739-1744 (*Et ester, synth, ir, pmr*)
 Bashford, K.E. *et al.*, *J.C.S. Perkin 1*, 2002, 1672-1687 (*N-Me Me ester, N-benzyl Me ester, synth, ir, pmr*)
 Gutierrez-Lugo, M.-T. *et al.*, *Nat. Prod. Res.*, 2005, **19**, 645-652 (isol, pmr, cmr)

5-Methyl-1H-indole-4,7-dione, 9CI M-477
 [213040-66-5]




$C_9H_7NO_2$ 161.16
 Alkaloid from the muricid gastropod

Drupella fragum. Antimicrobial agent. Orange prisms. Mp 202-204°. λ_{max} 228 (ε 8060); 250 (ε 2010); 316 (ε 4180); 470 (ε 790) (EtOH).


Fukuyama, Y. *et al.*, *Tetrahedron*, 1998, **54**, 10007-10016 (isol, uv, ir, pmr, cmr)

2-Methyl-1H-indole-4-methanol, 9CI M-478
4-(Hydroxymethyl)-2-methyl-1H-indole
 [154523-21-4]



$C_{10}H_{11}NO$ 161.203
 Alkaloid from fruit bodies of the mushrooms *Tricholoma sciodes* and *Tricholoma virgatum*. Yellow oil (natural). Mp 80-81° (synthetic). λ_{max} 222 (ε 25400); 279 (ε 6100) (EtOH).
Me ether: 4-(Methoxymethyl)-2-methylindole
 [154523-22-5]
 $C_{11}H_{13}NO$ 175.23
 From *Tricholoma sciodes* and *Tricholoma virgatum*. Yellow oil.
 Garlaschelli, L. *et al.*, *Tetrahedron*, 1994, **50**, 3571-3574 (isol, uv, ir, pmr, cmr, ms)
 Söderberg, B.C. *et al.*, *J.O.C.*, 1999, **64**, 9731-9734 (synth)

5-Methyl-3-isoxazolecarboxylic acid, 9CI M-479
 [3405-77-4]



$C_5H_5NO_3$ 127.099
 Prisms or plates (H₂O). Mp 176°.

Me ester: Premnazole
 [19788-35-3]
 $C_6H_7NO_3$ 141.126
 Constit. of the leaves of *Gmelina arborea* and *Premna integrifolia*. Anti-inflammatory. Cryst. (C₆H₆). Mp 98-99°.


Et ester: [3209-72-1]
 $C_7H_9NO_3$ 155.153
 Bp₁₅ 140°.

Amide: [3445-52-1]
 $C_5H_6N_2O_2$ 126.115
 Mp 166°.

Nitrile: 3-Cyano-5-methylisoxazole
 [57351-99-2]
 $C_5H_4N_2O$ 108.099
 Bp₂₅ 87-88°.

Good, R.H. *et al.*, *J.C.S. Perkin 1*, 1972, 2441
 Barik, B.R. *et al.*, *Fitoterapia*, 1992, **63**, 295 (*Premnazole*)
 Khanna, I.K. *et al.*, *J. Med. Chem.*, 2000, **43**, 3168-3185 (*nitrile, synth, pmr*)

3-Methyl-2-[6-(3-methyl-2-butenyl)-1H-indol-3-yl]-1,3-butane-diol, 9CI M-480
3-(2-Hydroxy-1-hydroxymethyl-2-methylpropyl)-6-prenylindole
 [93822-67-4]



$C_{18}H_{25}NO_2$ 287.401
 λ_{max} 227 (ε 50120); 283 (ε 7940); 293 (ε 7940) (MeOH) (Berdy).

(R)-form [165604-40-0]
 Alkaloid from the stem bark of *Hexalobus crispiflorus* (Annonaceae). Oil. $[\alpha]_D^{21}$ -16 (c, 0.8 in CHCl₃). Poss. artifact.

O^{2'}-Hexadecanoyl: 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1H-indol-3-yl]butyl hexanoate, 9CI
 [165604-36-4]
 [93822-69-6]
 $C_{34}H_{55}NO_3$ 525.813

Alkaloid from the stem bark of *Hexalobus crispiflorus* and *Isolona maitlandii* (Annonaceae). Oil. $[\alpha]_D^{21}$ -19 (c, 1.3 in CHCl₃). Poss. artifact. λ_{max} 227 (ε 50118); 283 (ε 7940); 293 (ε 7940) (MeOH) (Berdy).

O^{2'}-(9Z-Octadecenoyl): 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1H-indol-3-yl]butyl 9Z-octadecenoate, 9CI
 [93822-70-9]


$C_{36}H_{57}NO_3$ 551.851
 Alkaloid from the stem bark of *Hexalobus crispiflorus* and *Isolona maitlandii* (Annonaceae). Oil. $[\alpha]_D^{21}$ -20 (c, 1.9 in CHCl₃). Poss. artifact.

O^{2'}-(9Z,12Z-Octadecadienyl): 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1H-indol-3-yl]butyl 9Z,12Z-octadecadienoate, 9CI
 [93822-71-0]
 $C_{36}H_{55}NO_3$ 549.835

Alkaloid from the stem bark of *Hexalobus crispiflorus* and *Isolona maitlandii* (Annonaceae). Oil. $[\alpha]_D^{21}$ -20 (c, 1.7 in CHCl₃). Poss. artifact.

Achenbach, H. *et al.*, *Annalen*, 1995, 1327-1337 (isol, uv, ir, pmr, cmr, ms, struct)
 Achenbach, H. *et al.*, *Phytochemistry*, 1995, **40**, 967-973 (isol, ms)

2-Methyl-N-(2-methylbutyl)butanamide, 9CI M-481
 [158964-24-0]



$C_{10}H_{21}NO$ 171.282
 All 4 stereoisomers have been synthesised.

(S,S)-form [179826-31-4]


Sex pheromone of the longhorn beetle *Migdolus fryanus*. Oil. $[\alpha]_D^{20} +27.8$ (c, 3.2 in Et₂O).

Leal, W.S. *et al.*, *Experientia*, 1994, **50**, 853-856 (isol, ms)

Santangelo, E.M. *et al.*, *Synth. Commun.*, 2001, **31**, 3685-3698 (stereoisomers, synth)

2-Methyl-4-(2-methylpropyl)-2,4-azetidinedicarboxylic acid, 9CI M-482

2-Isobutyl-4-methyl-2,4-azetidinedicarboxylic acid. *Monascus acid*



C₁₀H₁₇NO₄ 215.249

(2R,4S)-form

Isol. from *Monascus pilosus*-fermented rice (red-mould rice). Needles. Mp 122-123°. $[\alpha]_D^{25} +3.7$ (c, 0.39 in Me₂CO).


(2S,4R)-form

Isol. from *Monascus pilosus*-fermented rice (red-mould rice). Needles. Mp 122-123°. $[\alpha]_D^{25} -4.4$ (c, 0.36 in Me₂CO).

Akihisa, T. *et al.*, *J. Nat. Prod.*, 2004, **67**, 479-480; 2005, **68**, 1444 (isol, pmr, cmr, ms)

2-Methyl-3-(2-methylpropyl)pyrazine, 9CI M-483

2-Isobutyl-3-methylpyrazine, 8CI. FEMA 3133 [13925-06-9]



C₉H₁₄N₂ 150.223

Present in seed oils of red pepper and pumpkin. Bp₁₀ 74°.

Flament, I. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 694-705 (isol, synth, ms)

Takken, H.J. *et al.*, *J. Agric. Food Chem.*, 1975, **23**, 638-642 (props, occur)

U.S. Pat., 1976, 3 989 713; CA, **86**, 43556d (synth, use)


Stofberg, J. *et al.*, *Perfum. Flavor.*, 1984, **9**, 53-56; 58-59; 62; 65-72; 78-83 (use)

Buchbauer, G. *et al.*, *Ernaehrung (Vienna)*, 1998, **22**, 246-249 (detn, pumpkin oil)

Jung, M.Y. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 1700-1704 (detn, pepper oil)

2-Methyl-4-(3-methyl-1-pyrrolidinyl)-1-butanol M-484

β,3-Dimethyl-1-pyrrolidinebutanol, 9CI [163315-01-3]




C₁₀H₂₁NO 171.282

Trace alkaloid from poison glands of the ants *Harpagoxenus sublaevis*, *Leptothorax acervorum*, *Leptothorax muscorum* and *Doronomyrmex goesswaldi*. Oil.

Reder, E. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 73 (isol, pmr, cmr, ms, synth, struct)

Koob, R. *et al.*, *Helv. Chim. Acta*, 1997, **80**, 267-272 (config)

N-[15-Methyl-3-(13-methyl-4-tetradecenoyloxy)hexadecanoyl]-glycine M-485



C₃₄H₆₃NO₅ 565.875

(R)-(Z)-form [193825-66-0]

Prod. by a marine *Cytophaga* sp. N-type calcium channel blocker. Leaflets (hexane/CH₂Cl₂). Mp 71-72°. $[\alpha]_D^{25} +0.45$ (c, 8 in CHCl₃).

4',5'-Dihydro: N-[15-Methyl-3-(13-methyltetradecanoyloxy)hexadecanoyl]glycine. **Topostin B 567**. *Cytolipin* [103196-73-2]

C₃₄H₆₅NO₅ 567.891

Prod. by the marine *Cytophaga johnsonae* and *Flexibacter topostinus*. N-type calcium channel blocker. Topoisomerase inhibitor. Leaflets (hexane/CH₂Cl₂). $[\alpha]_D^{22} +1.9$ (c, 0.2 in CHCl₃).

(±)-(Z)-form [183167-45-5]

Cryst. (hexane/CH₂Cl₂). Mp 84-85°.

Suzuki, K. *et al.*, *J. Antibiot.*, 1990, **43**, 154-157; 158-162 (*Topostin*)


Morishita, T. *et al.*, *J. Antibiot.*, 1997, **50**, 457-468 (isol, synth, ir, pmr, cmr)

Nemoto, T. *et al.*, *Tetrahedron*, 1998, **54**, 2683-2690 (isol, *Topostin*)

Shioiri, T. *et al.*, *Tetrahedron*, 1998, **54**, 15701-15710 (synth)

3-Methyl-1-[3-(methylthio)propyl]pyrrolidine, 9CI M-486

[163315-03-5]



(R)-form

C₉H₁₉NS 173.322

(R)-form


Trace alkaloid from the poison glands of the ants *Harpagoxenus sublaevis*, *Leptothorax acervorum* and *Leptothorax muscorum*. Oil.

Reder, E. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 73 (isol, pmr, cmr, ms, synth, struct)

Koob, R. *et al.*, *Helv. Chim. Acta*, 1997, **80**, 267-272 (abs config)

4-Methyl-2,6-naphthyridine, 9CI M-487

[31121-65-0]




C₉H₈N₂ 144.176

Constit. of *Antirrhinum majus* and several other *Antirrhinum* spp. Needles (EtOH). Mp 78°.

Harkins, K.J. *et al.*, *Planta Med.*, 1971, **20**, 108 (isol, spectra)

Taurins, A. *et al.*, *Can. J. Chem.*, 1974, **52**, 843 (synth)

1-Methyl-2,7-naphthyridine-4-carboxylic acid M-488



C₁₀H₈N₂O₂ 188.185

Me ester: 4-Methoxycarbonyl-1-methyl-2,7-naphthyridine. **Neozeylanicine** [112561-62-3]

C₁₁H₁₀N₂O₂ 202.212

Alkaloid from the wood of *Neonauclea zeylanica* (Rubiaceae) and from *Strychnos cocculoides* and *Strychnos spinosa*. Light yellow amorph. solid. Mp 110°.

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1988, **54**, 461-462 (*Me ester*, isol, uv, ir, pmr, ms, struct)

Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1992, **61**, 429-440 (*Me ester*, isol)

Bracher, F. *et al.*, *Annalen*, 1995, 645-647 (synth, ir, pmr, cmr, ms)

5-Methyl-2,7-naphthyridine-4-carboxylic acid M-489

C₁₀H₈N₂O₂ 188.185

Me ester: 5-Methoxycarbonyl-4-methyl-2,7-naphthyridine

[149155-02-2]


C₁₁H₂₀N₂O₂ 212.291

Alkaloid from *Strychnos cocculoides* and *Strychnos spinosa*. No phys. props. reported.

Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1992, **61**, 429-440 (*Me ester*, isol, pmr, ms)

3-Methyl-2-[(2-nonene-4,6,8-triynoyl)amino]butanoic acid M-490

N-(2-Nonene-4,6,8-triynoyl)valine



C₁₄H₁₃NO₃ 243.262

(S,E)-form

L-trans-form


Metab. of *Poria sinuosa*.

Me ester:

Unstable oil.

Cambie, R.C. et al., *J.C.S.*, 1963, 2056 (*isol, uv, ir, struct*)**3-Methyl-2-(2-nonenyl)-4(1H)-quinolinone** M-491

4-Hydroxy-3-methyl-2-(2-nonenyl)quinolinone. 3-Methyl-2-(2-nonenyl)-4-quinolinol. *Antibiotic SF 2420B. SF 2420B. PSC-D. NMQ. PC 3* [147348-69-4]

C₁₉H₂₅NO 283.413

Prod. by an unknown strain SF 2420B and by *Pseudomonas cepacia* PC II. Lipoygenase inhibitor. Cryst. (MeOH aq.). Mp 197-198°. λ_{max} 214 ; 243 ; 321 ; 334 (MeOH) (Berdy).

2',3'-Dihydro-3-methyl-2-nonyl-4(1H)-quinolinone. PSC-EC₁₉H₂₇NO 285.428

From *Pseudomonas cepacia* PC II. Cryst. (MeOH aq.). Mp 215-216°. λ_{max} 214 ; 239 ; 322 ; 335 (MeOH) (Berdy).


NH-form [130772-03-1]

N-Hydroxy: 1-Hydroxy-3-methyl-2-(2-nonenyl)-4(1H)-quinolinone. *Antibiotic YM 30059. YM 30059. YL 02729S.*

[162382-62-9]

C₁₉H₂₅NO₂ 299.412

Prod. by *Arthrobacter* sp. Antibacterial agent. Cryst. (formerly descr. as yellow syrup). Mp 101-103°. λ_{max} 250 (ε 18000); 337 (ε 6300); 348 (ε 6200) (MeOH).

Homma, Y. et al., *Soil Biol. Biochem.*, 1989, **21**, 723-728 (*isol*)Minowa, N. et al., *CA*, 1993, **118**, 212735(*synth*)Kamigiri, K. et al., *J. Antibiot.*, 1996, **49**, 823-825 (*YM 30059*)Moon, S.-S. et al., *Phytochemistry*, 1996, **42**, 365-368 (*isol, uv, ir, pmr, cmr, ms, struct*)Tatsuta, K. et al., *J. Antibiot.*, 2000, **53**, 418-421 (*synth, pmr, cmr*)**2-Methyl-6-nonylpiperidine** M-492C₁₅H₃₁N 225.417

Minor component of venoms of *Solenopsis* spp. ants. See also references under 2-Methyl-6-tridecylpiperidine, M-580 and 2-Methyl-6-undecylpiperidine, M-584.

N-Me: N,2-Dimethyl-6-nonylpiperidine

[33444-22-3]

C₁₆H₃₃N 239.443Isol. from the venom of *Solenopsis**steinheili* and *Solenopsis* sp. PR.

Stereochem. not detd.

[52084-40-9 ; 52084-39-6]


Jones, T.H. et al., *Tetrahedron*, 1982, **38**, 1949 (*isol, struct, bibl*)**N-(8-Methylnonyl)sulfamic acid** M-493

[1040631-00-2]

(H₃C)₂CH(CH₂)₆CH₂NHSO₃HC₁₀H₂₃NO₃S 237.363Isol. from *Daphniapulex*. Kairomone.Yasumoto, K. et al., *Chem. Pharm. Bull.*, 2008, **56**, 133-136 (*isol, pmr*)**8-Methylnorchelerythrine** M-494

1,2-Dimethoxy-13-methyl[1,3]benzodioxolo[5,6-c]phenanthridine, 9CI

[154490-59-2]

C₂₁H₁₇NO₄ 347.37

Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Prisms (CHCl₃/MeOH). Mp 198-200°.


Chen, I.-S. et al., *J. Nat. Prod.*, 1994, **57**, 1206 (*isol, uv, ir, pmr, ms, struct*)**7-Methyl-2,4-octadienoic acid** M-495(H₃C)₂CHCH₂CH=CHCH=CHCOOHC₉H₁₄O₂ 154.208**(2E,4E)-form**

Amide: 7-Methyl-2,4-octadienamide

[918637-54-4]

C₉H₁₅NO 153.224


Prod. by the marine-derived *Streptomyces* strain CNQ-085. Solid. λ_{max} 262 (log ε 4.12) (MeOH).

Asolkar, R.N. et al., *J. Nat. Prod.*, 2006, **69**, 1756-1759 (*isol, pmr, cmr*)**2-(6-Methyl-2,4-octadienyl)-1H-pyrrole** M-496*Axinellamine A*⁺C₁₃H₁₉N 189.3**(2'E,4'E,6R)-form** [213269-06-8]

Alkaloid from the marine sponge *Axinella* sp. Pale yellow gum. [α]_D²⁵ -45.4 (c, 0.2 in CHCl₃). λ_{max} 227 (ε 8000) (MeOH).

Bascombe, K.C. et al., *Heterocycles*, 1998, **48**, 1461-1464 (*isol, uv, ir, pmr, cmr, ms*)Seki, M. et al., *Eur. J. Org. Chem.*, 2001, 503-506 (*synth, abs config*)**N-(7-Methyloctyl)sulfamic acid** M-497


[1040631-01-3]

(H₃C)₂CH(CH₂)₅CH₂NHSO₃HC₉H₂₁NO₃S 223.336Isol. from *Daphnia pulex*. Kairomone.Yasumoto, K. et al., *Chem. Pharm. Bull.*, 2008, **56**, 133-136 (*isol, pmr, cmr*)**1-Methyl-2-(9-oxodecyl)-4(1H)-quinolinone, 9CI** M-498C₂₀H₂₇NO₂ 313.439

Alkaloid from *Ruta montana*. λ_{max} 215 (log ε 4.1); 236 (log ε 4.5); 320 (log ε 3.8); 330 (log ε 3.7) (MeOH).

Touati, D. et al., *Phytochemistry*, 2000, **53**, 277-279**5-Methyl-2-oxo-4-imidazolidine-dihexanoic acid, 8CI** M-499

7,8-Ureylenenonanoic acid. *Desthiobiotin. Desthiobiotin* [15720-25-9]

HOOC(CH₂)₅C₁₀H₁₈N₂O₃ 214.264**(4R,5S)-form**

(+)-cis-form

[533-48-2]

Prod. by various microorganisms, e.g. *Penicillium chrysogenum*, *Corynebacterium xerosis* and *Phycomyces blakesleeanus*. Intermed. in biotin biosynth. Cryst. Mp 160-162°. [α]_D²⁵ +10.4 (c, 0.57 in EtOH).

Me ester: [6020-51-5]Cryst. Mp 69-70°. [α]_D²³ +2.6 (c, 2 in CHCl₃).**(4RS,5SR)-form**

(±)-cis-form

[636-20-4]

Cryst. Mp 162-163°.

Me ester:Cryst. Mp 72°. Bp_{0.5} 170°.**(4RS,5RS)-form**

(±)-trans-form

[34879-36-2]


Cryst. Mp 156°.

du Vigneaud, V. et al., *J. Biol. Chem.*, 1942, **146**, (*synth*)Duschinsky, R. et al., *J.A.C.S.*, 1945, **67**, 2079 (*synth*)Kuzuhara, H. et al., *Tet. Lett.*, 1970, 1185 (*synth*)Parry, R.J. et al., *Chem. Comm.*, 1975, 321 (*synth*)Guillerm, G. et al., *J.O.C.*, 1977, **42**, 3776 (*synth, pmr, ms*)

Okabe, N. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 2143 (*cryst struct*)

Lucet, D. *et al.*, *Eur. J. Org. Chem.*, 2000, 3575-3579 (*synth*)

8-(4-Methyl-2-oxopentyl)dihydrochelerythrine M-500
[112899-81-7]




$C_{27}H_{29}NO_5$ 447.53

Alkaloid from the stem bark of a *Zanthoxylum* sp. collected in Grand Cayman (*Zanthoxylum spinosum* or *Zanthoxylum coriaceum*) (Rutaceae).

Ng, K.M. *et al.*, *Phytochemistry*, 1987, **26**, 3251-3254 (*isol, uv, ir, pmr, cmr, ms, struct*)

18-O-[2-(2-Methyl-4-oxo-4H-quinazolin-3-yl)benzoyl]lycoctonine M-501



$C_{41}H_{51}N_3O_9$ 729.869

Alkaloid from the roots of *Aconitum pseudo-laeve* var. *erectum*. Amorph. powder (MeOH). $[\alpha]_D^{21} +65$ (c, 3.5 in MeOH).


6-Demethoxy, 7-deoxy, O^{14} -de-Me, O^8 -Me, 14-Ac: **14-O-Acetyl-8-O-methyl-18-O-[2-(2-methyl-4-oxo-4H-quinazolin-3-yl)benzoyl]cammaconine**

$C_{42}H_{51}N_3O_8$ 725.88

Alkaloid from the roots of *Aconitum pseudo-laeve* var. *erectum*. Amorph. powder (MeOH). $[\alpha]_D^{21} -10.4$ (c, 1.3 in MeOH).

Shim, S.H. *et al.*, *J. Nat. Prod.*, 2006, **69**, 400-402 (*isol, pmr, cmr, ms*)

17-Methylparsonsianidine M-502
[135637-68-2]



$C_{23}H_{35}NO_9$ 469.531

Alkaloid from leaves of *Parsonsia laevigata* (Apocynaceae). Cryst. Mp 195-205° dec. $[\alpha]_D^{20} +31.3$ (c, 0.3 in MeOH).

Stereoisomer (?): **Spiranine**

[77156-23-1]

$C_{23}H_{35}NO_9$ 469.531

Alkaloid from *Parsonsia spiralis* (Apocynaceae). Identified by nmr and ms. Same gross struct. as 17-methylparsonsianidine but with less stereochem. defined. May be identical with it (no reference to comparison of spectra).

24- ξ -Hydroxy: **Spiracine**

[77156-24-2]

$C_{23}H_{35}NO_{10}$ 485.53

Alkaloid from *Parsonsia spiralis* (Apocynaceae). Identified by nmr and ms. No stereochem. attributed to necic acid residue.

16-Deoxy, *stereoisomer*: **Heterophylline**†

[77156-22-0]

$C_{23}H_{35}NO_8$ 453.531


Alkaloid from *Parsonsia heterophylla* and *Parsonsia spiralis* (Apocynaceae). Noncryst. $[\alpha]_D^{20} +57.6$ (c, 0.38 in MeOH). No stereochem. attributed to necic acid residue.

Edgar, J.A. *et al.*, *Tet. Lett.*, 1980, 2657

(*Spiranine, Spiracine, Heterophylline*)

Abe, F. *et al.*, *Phytochemistry*, 1991, **30**, 1737 (*17-Methylparsonsianidine*)

16-Methylpendolmycin M-503
[138590-60-0]




$C_{23}H_{33}N_3O_2$ 383.533

Isol. from a *Nocardioopsis* sp. Protein kinase C inhibitor. Mp 174-178°. $[\alpha]_D -76$ (c, 0.57 in MeOH). λ_{max} 230 (ϵ 18000); 289 (ϵ 6200); 299 (ϵ 6500) (MeOH) (Berdy).

Sun, H.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1440 (*isol, uv, ir, pmr, cmr, cd, struct*)

2-Methyl-6-pentadecylpiperidine, 9CI M-504



$C_{21}H_{43}N$ 309.577

See also refs. under 2-Methyl-6-tridecylpiperidine, M-580 and 2-Methyl-6-undecylpiperidine, M-584.

(**2R*,6R***)-form
trans-form. *Solenopsine C*

[32778-79-3]

Major component of venoms of both red and black forms of the fire ant (*Solenopsis saevissima*). Also present in venom of *Solenopsis invicta*. Oil.

6',7'-Didehydro(Z-): 2-Methyl-6-(6-pentadecenyl)piperidine, 9CI. **Dehydroso-lenopsin C**

[32778-78-2]

Major component of venom of red form of *Solenopsis saevissima*, only present in minor amounts in black form. Also from *Solenopsis invicta* venom. Bp_{0.01} 80°.

(**2R*,6S***)-form

cis-form

[35285-28-0]

Minor component of venoms of red and black forms of *Solenopsis saevissima*. Also present in venom of *Solenopsis invicta*. Oil.

6',7'-Didehydro(Z-): Minor component of venom of red form of *Solenopsis saevissima*.

[35285-32-6]

MacConnell, J.G. *et al.*, *Tetrahedron*, 1971, **27**, 1129 (*isol, synth, ir, ms, pmr*)

Jouvenaz, D.P. *et al.*, *Antimicrob. Agents Chemother.*, 1972, **2**, 291

Brand, J.M. *et al.*, *Toxicol.*, 1972, **10**, 259 (*ms*)

Jones, T.H. *et al.*, *Tetrahedron*, 1982, **38**, 1949 (*isol, struct, bibl*)

4-Methylpentanoic acid, 9CI M-505

4-Methylvaleric acid. Isocaproic acid.

Isohexanoic acid. Isobutylacetic acid.

FEMA 3463

[646-07-1]

$(H_3C)_2CHCH_2CH_2COOH$

$C_6H_{12}O_2$ 116.16

Found in bananas and lime oil. Oil, with sour penetrating odour. d_4^{20} 0.92. Mp -33°. Bp 199.1° Bp_{2.5} 69°. n_D^{20} 1.4144.

▶ Fl. p. 89°. Eye and skin irritant.

NR2975000

Amide: 4-Methylpentanamide

[1119-29-5]

$C_6H_{13}NO$ 115.175


Isol. from *Rhizoctonium hieroglyphicum*. Mp 120-121°.

Dembitsky, V.M. *et al.*, *Phytochemistry*, 2000, **54**, 965-967 (*isol, amide*)

3-Methyl-3-pentenoic acid, 9CI M-506

3-Ethylidenebutyric acid. 2-Methyl-2-butene-1-carboxylic acid

[16313-37-4]



$C_6H_{10}O_2$ 114.144

(**E**)-form

trans-form

[41653-93-4]

Mp 35°.

Me ester: [41654-12-0]

$C_7H_{12}O_2$ 128.171

Bp₅₀ 74°.

Amide:

C₆H₁₁NO 113.159
Isol. from *Senecio rivularis*. Cryst. (EtOAc). Mp 130-132°.

(Z)-form

cis-form
[41653-94-5]
Oil.

Et ester:

C₈H₁₄O₂ 142.197
Bp₁₃ 61°.

Amide:

Plates (C₆H₆/pentane). Mp 126-127°.

Kon, G.A.R. *et al.*, *J.C.S.*, 1925, **127**, 623

(*synth*)

Wagner, R.B. *et al.*, *J.A.C.S.*, 1949, **71**, 3214


(*synth*)

Klásek, A. *et al.*, *Chem. Ind. (London)*, 1969, 987

1-Methyl-4-(2-penten-4-ynyl)quinolizidine

M-507

Octahydro-1-methyl-4-(2-penten-4-ynyl)-2H-quinolizidine, 9CI. Dendrobates *Alkaloid 217A*. *Quinolizidine 217A* [151805-13-9]



C₁₅H₂₃N 217.353

Alkaloid from the frogs *Epipedobates tricolor*, *Mantella madagascariensis* and *Mantella baroni*. [α]_D +75 (c, 0.15 in MeOH). λ_{max} 287 (ε 16900) (MeOH).

1-Epimer: [219796-70-0]

C₁₅H₂₃N 217.353

Occurs in a Madagascan frog. Pale yellow oil. Unpublished work.

Jain, P. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1174-1178 (*isol*)


Pearson, W.H. *et al.*, *J.O.C.*, 1998, **63**, 9910-9918 (*synth*)

Maloney, K.M. *et al.*, *Org. Lett.*, 2005, **7**, 3115-3118 (*synth*)

3-Methyl-2-pentyl-4(1H)-quinolinone

M-508

4-Hydroxy-3-methyl-2-pentylquinoline. 3-Methyl-2-pentyl-4-quinolinol. PSC-A [178955-99-2]



C₁₅H₁₉NO 229.321


Alkaloid from *Pseudomonas cepacia* PC-II. Shows plant growth promoting and fungicidal activity. Cryst. (MeOH aq.). Mp 222-223°. λ_{max} 213 ; 239 ; 321 ; 334 (MeOH) (Berdy).

Moon, S.-S. *et al.*, *Phytochemistry*, 1996, **42**, 365 (*isol, uv, ir, pmr, cmr, ms, struct*)

N-Methyl-N-(2-phenyleth-nyl)-3-phenyl-2-oxiranecarboxamide

M-509

SB 204900
[173220-67-2]



Relative configuration

C₁₈H₁₇NO₂ 279.338

Isol. from leaves of *Clausena lansium* (wampee). Mp 59-60°. [α]_D²⁵ +14 (c, 1.0 in CHCl₃).

Milner, P.H. *et al.*, *J. Nat. Prod.*, 1996, **59**, 400


(*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Stefanutti, I. *et al.*, *Tet. Lett.*, 2000, **41**, 3735-3738 (*synth*)

1-Methyl-3-(2-phenylethyl)-2,4(1H,3H)-quinazolidinone, 9CI

M-510

[75652-62-9]



C₁₇H₁₆N₂O₂ 280.326

Alkaloid from the leaves and seed husks of *Zanthoxylum arborescens* (Rutaceae). Cryst. (MeOH or EtOAc/petrol). Mp 100-102°.

4'-Methoxy: 3-[2-(4-Methoxyphenyl)ethyl]-1-methyl-2,4(1H,3H)-quinazolidinone, 9CI

[75652-63-0]

C₁₈H₁₈N₂O₃ 310.352

Alkaloid from leaves, bark and seed husks of *Zanthoxylum arborescens* (Rutaceae). Cryst. (MeOH aq. or EtOAc/petrol). Mp 135-136°.

Dreyer, D.L. *et al.*, *Phytochemistry*, 1980, **19**, 935-939 (*isol, synth, uv, ir, pmr, cmr, ms*)

Grina, J.A. *et al.*, *J.O.C.*, 1982, **47**, 2648 (*isol*)


Rivero, I.A. *et al.*, *Molecules*, 2004, **9**, 609-616

(*synth, pmr, cmr*)

4-Methyl-5-phenyl-2-oxazolidinone

M-511

[54418-69-8]



(4*R*,5*R*)-form

C₁₀H₁₁NO₂ 177.202

(4*R*,5*R*)-form

(-)-*trans-form*

[125133-96-2]

Cryst. (EtOAc/hexane). Mp 117-119°.

[α]_D²¹ -29.5 (c, 2.3 in CHCl₃).

(4*R*,5*S*)-form

(+)-*cis-form*

[77943-39-6] Chiral auxiliary.

Cryst. (toluene or petrol/CH₂Cl₂). Mp

121-122°. [α]_D +177.2 (c, 2.21 in CHCl₃).

(4*S*,5*R*)-form

(-)-*cis-form*

[16251-45-9]

Needles (hexane). Mp 116-117°. [α]_D²⁰ -170 (c, 1.2 in CHCl₃). [α]_D²⁵ -168 (c, 2 in CHCl₃) (98% ee).

N-Me: 3,4-Dimethyl-5-phenyl-2-oxazolidinone. **Ephedroxane**

[16251-46-0]

Alkaloid from the aerial parts of

Ephedra intermedia (Ephedraceae).

Needles (EtOH). Mp 79-81°.

(4*S*,5*S*)-form

(+)-*trans-form*

[17097-67-5]

Mp 120-121°. [α]_D²⁶ +25.9 (c, 0.34 in CHCl₃).

N-Me: [16251-47-1]

Prisms (EtOH). Mp 50-51°. [α]_D +36.7

(c, 0.98 in CHCl₃).

(4*R*,5*SR*)-form

(±)-*cis-form*

[39663-75-7]

Cryst. (C₆H₆). Mp 148-149°.

(4*R*,5*RS*)-form

(±)-*trans-form*

[149055-02-7]

[28044-23-7]

Cryst. (EtOH). Mp 96-96.5° (96°).

[28044-22-6]

Stratton, J.M. *et al.*, *J.C.S.*, 1932, 1133 (*synth*)

Close, W.J. *et al.*, *J.O.C.*, 1950, **15**, 1131 (*synth*)

Fodor, G. *et al.*, *Monatsh. Chem.*, 1967, **98**,

1027 (*synth, N-Me*)

Auerbach, R.A. *et al.*, *Org. Mass Spectrom.*,

1970, **4**, 41 (*ms*)

Spassov, S.L. *et al.*, *Chem. Ber.*, 1972, **105**,

2462 (*pmr*)

Caccia, G. *et al.*, *J.O.C.*, 1973, **38**, 2264 (*synth*)

Konno, C. *et al.*, *Phytochemistry*, 1979, **18**,

697-698 (*N-Me, isol, ir, pmr, synth*)

Klyne, W. *et al.*, *Tetrahedron*, 1979, **35**, 2009-

2012 (*cd, abs config*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1980, **45**, 1066

(*synth, pmr*)

Wainer, I.W. *et al.*, *J. Chromatogr.*, 1983, **268**,

107-111 (*chromatog, resoln*)

Evans, D.A. *et al.*, *J.O.C.*, 1985, **50**, 1830

(4*R*,5*S*-form, *synth, ir, pmr, cmr*)

Kano, S. *et al.*, *Tet. Lett.*, 1987, **28**, 6331

(*synth*)

Jephcote, V.J. *et al.*, *J.C.S. Perkin 1*, 1989, 1529

(4*R*,5*R*-form, *synth*)

Ku, T.W. *et al.*, *J.O.C.*, 1989, **54**, 3487 (*synth,*

pmr)

Brimble, M.A. *et al.*, *Aust. J. Chem.*, 1990, **43**,

1035 (4*R*,5*S*-form, *synth*)

Robinson, P.D. *et al.*, *Acta Cryst. C*, 1993, **49**,

1238 (4*S*,5*S*-form, *cryst struct*)

Moreno-Manas, M. *et al.*, *J. Het. Chem.*, 1993,

30, 1235-1239 (*N-Me*)

Cainelli, G. *et al.*, *Tetrahedron*, 1993, **49**, 3809-

3826 (*synth, ms, ir, pmr, cmr*)

Davies, S.G. *et al.*, *Tetrahedron: Asymmetry*,

1993, **4**, 2513 (4*R*,5*S*-form, *ir, pmr, ms*)

Pettit, G.R. *et al.*, *Synthesis*, 1996, 719 (4*R*,5*S*-

form, synth, pmr, ir)

Cutugno, S. *et al.*, *Eur. J. Org. Chem.*, 2001,

517-522 (4*S*,5*R*-form, *synth*)


Alouane, N. *et al.*, *Synthesis*, 2006, 885-889

(4*S*,5*R*-form, 4*S*,5*S*-form, *N-Me, synth, ir,*

pmr, cmr)

3-(2-Methylphenyl)-3-oxopropanamide M-512

3-Hydroxy-3-(2-methylphenyl)-2-propenamide, 9CI. 2-Methyl-β-oxobenzenepropanamide. USF 142A. Antibiotic USF 142A [156281-39-9]




C₁₀H₁₁NO₂ 177.202

Partially enolised. Prod. by *Streptomyces* sp. USF-142. Antioxidant. Needles (EtOAc/hexane). Mp 83-84°. λ_{max} 245 (ε 3980); 271 (ε 8690) (MeOH).

Hirota, A. et al., *Biosci., Biotechnol., Biochem.*, 1994, **58**, 792 (isol, uv, ir, pmr, cmr)

3-(2-Methylphenyl)-2-propenoic acid, 9CI M-513

o-Methylcinnamic acid, 8CI. 3-o-Tolylacrylic acid [2373-76-4]



(E)-form

C₁₀H₁₀O₂ 162.188
pK_a 4.5.

(E)-form [939-57-1]

Cryst. (EtOH or C₆H₆). Mp 175-176°.

Et ester: [24393-48-4]

C₁₂H₁₄O₂ 190.241

Oil. Bp_{2.2} 114-117° Bp_{1.2} 148°.

Amide: 3-(2-Methylphenyl)-2-propenamide. U 77863. Antibiotic U 77863

C₁₀H₁₁NO 161.203

Prod. by *Streptomyces griseoluteus*.

Exhibits antitumour props. Cryst. (MeOH). Mp 151-153°. λ_{max} 208 (ε 17300); 220 (ε 14900); 225 (sh) (ε 12000); 276 (ε 16500) (MeOH) (Derrep). λ_{max} 208 (ε 17320); 220 (ε 14900); 276 (ε 16540) (MeOH) (Berdy).

(Z)-form [41397-71-1]

Mp 91-92° (89-90°).

[83716-65-8, 16366-23-7]

Cope, A.C. et al., *J.A.C.S.*, 1957, **79**, 240 (synth)

Craig, J. et al., *Aust. J. Chem.*, 1959, **12**, 447 (synth)

Norcross, B.E. et al., *J.O.C.*, 1977, **42**, 369 (synth)

Schaldach, B. et al., *Org. Mass Spectrom.*, 1980, **15**, 182 (ms)

Galamb, V. et al., *Tet. Lett.*, 1983, **24**, 2965 (synth)

Suzuki, H. et al., *Chem. Lett.*, 1986, 403 (ester)

Liu, J.-M. et al., *J.O.C.*, 1986, **51**, 1120 (synth, pmr)


Robinson, C.N. et al., *J.O.C.*, 1986, **51**, 3535 (cmr, ester)

Harper, D.E. et al., *J. Antibiot.*, 1992, **45**, 1827 (U 77863)

Berry, J.M. et al., *J.C.S. Perkin 1*, 1997, 1147-1156 (synth, pmr)

3-Methyl-4-phenyl-1H-pyrrole M-514

[116267-86-8]



C₁₁H₁₁N 157.215

Isol. from the ant *Anochetus kemphi*. Oil.


Van Leusen, D. et al., *J.O.C.*, 1992, **57**, 2245 (synth, ir, pmr)

Jones, T.H. et al., *J. Nat. Prod.*, 1999, **62**, 1343-1345 (isol, synth)

1-Methylpiperidine, 9CI M-515

Methylpiperidylamine

[626-67-5]



C₆H₁₃N 99.175

Isol. from *Girgensohnia diptera* and *Girgensohnia oppositiflora* (Chenopodiaceae). d₄²⁰ 0.82. Bp 107°. n_D²² 1.4378. pK_a 10.38 (25°, 0.2M NaCl).

▶ Highly flammable, fl. p. 3°. LD₅₀ (mus, ipr) 400 mg/kg. TN1225000

Hydrochloride: [17874-59-8]

V. hygroscopic needles. Mp 185°.

▶ TN1275600

Methiodide:

Prisms (EtOH). Dec. at 334°.

Picrate: [18953-36-1]

Mp 223-224° (148°).

N-Oxide: N-Methylpiperidine oxide

[17206-00-7]

C₆H₁₃NO 115.175

Alkaloid from *Vandopsis longicaulis* (Orchidaceae). Hygroscopic solid by subl. pK_a 5.26.

N-Oxide; hydrochloride:

Cryst. (CHCl₃). Mp 180-200° dec.

N-Oxide, picrate:

Yellow plates (Me₂CO). Mp 194-208° dec.

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **1**, 564A (nmr)

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 360B (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase,

1989, **3**, 450C (ir)

Yurashevskii, N.K. et al., *CA*, 1946, **40**, 6754 (isol)

Lukeš, R. et al., *CA*, 1947, **41**, 4150a (synth)

Weitkamp, H. et al., *Chem. Ber.*, 1962, **95**, 2896 (pmr)

Bild, N. et al., *Helv. Chim. Acta*, 1967, **50**, 1885 (ms, oxide)

Ferris, J.P. et al., *J.O.C.*, 1968, **33**, 3493

Allan, G.G. et al., *Tappi*, 1968, **51**, 224 (synth)

Lanum, W.J. et al., *J. Chem. Eng. Data*, 1969, **14**, 93 (props)

Brandänge, S. et al., *Acta Chem. Scand.*, 1970, **24**, 353 (isol, synth, oxide)

Fr. Pat., 1971, 2 017 634; *CA*, **74**, 124791f

Katritzky, A.R. et al., *J.C.S. (B)*, 1971, 1330 (synth, conformn, oxide)

Vernot, A. et al., *Bull. Soc. Chim. Fr.*, 1972, 4736 (synth)

Eliel, E.L. et al., *J.A.C.S.*, 1975, **97**, 2424 (nmr)


Lambert, J.B. et al., *J.A.C.S.*, 1976, **98**, 3778 (cmr, oxide)

Chastanet, J. et al., *J.O.C.*, 1985, **50**, 2910 (oxide, synth, ms, pmr, use)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MOG500

2-Methyl-3,4-piperidinediol, 9CI M-516

3,4-Dihydroxy-2-methylpiperidine



C₆H₁₃NO₂ 131.174

(2R,3R,4R)-form 6-Deoxyfagomine


[197449-09-5]

Alkaloid from the roots of *Lycium chinense* (Chinese boxthorn). [α]_D -11.1 (c, 0.1 in H₂O).

Asano, N. et al., *Eur. J. Biochem.*, 1997, **248**, 296-303 (6-deoxyfagomine, isol, pmr, cmr, ms)

2-Methyl-3,4,5-piperidine-triol, 9CI M-517

3,4,5-Trihydroxy-2-methylpiperidine. 1,5-Imino-1,5,6-trideoxyhexitol



C₆H₁₃NO₃ 147.174

(2R,3R,4R,5S)-form

1,5-Imino-1,5,6-trideoxy-D-glucitol. 1,6-Dideoxyojirimycin

Alkaloid from the pods of *Angylocalyx pynaertii*. [α]_D +37.8 (c, 0.26 in H₂O).


Yasuda, K. et al., *J. Nat. Prod.*, 2002, **65**, 198-202 (isol, pmr, cmr)

Rengasamy, R. et al., *J.O.C.*, 2008, **73**, 2898-2901 (synth)

4-(1-Methyl-2-piperidinyl)-β-carboline M-518

4-(1-Methyl-2-piperidinyl)-9H-pyrido[3,4-b]indole, 9CI

[71005-06-6]




C₁₇H₁₉N₃ 265.357

In the review by Allen *et al* this compd. is shown. with a C-1 Me group and named Homobrevicolline. Alkaloid from *Carex brevicollis* (Cyperaceae).

Sharipov, I.N. *et al.*, *CA*, 1979, **91**, 87295a
Allen, J.R.F. *et al.*, *Phytochemistry*, 1980, **19**, 1573

1-(6-Methyl-2-piperidinyl)-2-propanone, 9CI M-519

C₉H₁₇NO 155.239


(R*,R*)-form

cis-form
[83285-66-9]

Alkaloid from the Australian mealybug ladybird *Cryptolaemus montrouzieri*.

Brown, W.V. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1255

O-Methylprechilenine M-520

C₂₁H₁₉NO₇ 397.384

Me ether of the hitherto hypothetical alkaloid prechilenine.


(±)-form [95585-80-1]

Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Amorph. Unstable.

Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957
(*isol, uv, ir, pmr, ms, struct*)

1-(2-Methyl-1-propenyl)-β-carboline M-521

1-(2-Methyl-1-propenyl)-9H-pyrido[3,4-b]indole. **Vulcanine**. 1-Isobutenyl-β-carboline
[185140-85-6]

C₁₅H₁₄N₂ 222.289

Alkaloid from *Haplophyllum vulcanium*. Mp 103° (as hydrochloride). λ_{max} 212 (log ε 4.58); 239 (log ε 4.7); 260 (sh) (log ε 4.43); 292 (log ε 4.36); 345 (sh) (log ε 3.98); 355 (log ε 4.01) (MeOH).

1,2,3,4-Tetrahydro, N^b-Me: 2,3,4,9-Tetrahydro-2-methyl-1-(2-methyl-1-propenyl)-

nyl)-1H-pyrido[3,4-b]indole. 1,2,3,4-Tetrahydro-1-isobutenyl-2-methyl-β-carboline. **Borrerine**

[51076-19-8]

C₁₆H₂₀N₂ 240.347

Alkaloid from *Borreria verticillata* (Rubiaceae). Mp 107-108° (102-103°) (synthetic) Mp 73° (natural). Racemic. λ_{max} 223 (log ε 4.69); 283 (log ε 4.01) (EtOH).

1,1',2,2',3,4-Hexahydro(±): 1,2,3,4-Tetrahydro-1-(2-methylpropyl)-9H-pyrido[3,4-b]indole. 1,2,3,4-Tetrahydro-1-isobutyl-β-carboline
[6649-77-0]

[26962-07-2]

C₁₅H₂₀N₂ 228.336

Alkaloid from the root bark of *Elaeagnus commutata* (Elaeagnaceae). Cryst. (MeOH)(as hydrochloride). Mp 257-259° (hydrochloride). Racemic. λ_{max} 223; 273; 279; 289 (EtOH) (as hydrochloride).

1,1',2,2',3,4-Hexahydro, N^b-Me: Mp 92°.

Δ²-Isomer, 1,2,3,4-tetrahydro, N^b-Me:

Isoborrerine

[76177-24-7]

C₁₆H₂₀N₂ 240.347

Alkaloid from *Flindersia fourmieri* (Flindersiaceae). Amorph. [α]_D²⁰ 0 (CHCl₃). Racemic. λ_{max} 225 (log ε 4.06); 282 (log ε 3.51); 290 (log ε 3.48) (EtOH).

Slywka, G.W.A. *et al.*, *Tet. Lett.*, 1969, 4635-4638 (*Tetrahydroisobutylcarboline*)

Pousset, J.L. *et al.*, *Phytochemistry*, 1973, **12**, 2308-2310 (*Borrerine, isol, uv, ms, pmr, struct*)

Tillequin, F. *et al.*, *Phytochemistry*, 1980, **19**, 1282-1283 (*Isoborrerine, isol, uv, ir, pmr, ms, struct, synth*)


Yamanaka, E. *et al.*, *Heterocycles*, 1984, **22**, 371-374 (*Borrerine, synth, pmr, cmr, ms*)

Somei, M. *et al.*, *Heterocycles*, 1988, **27**, 1585-1587 (*Borrerine, synth*)

Goetzler, T. *et al.*, *Phytochemistry*, 1996, **43**, 1425-1426 (*Vulcanine, isol, uv, ir, pmr, cmr, ms, cryst struct*)

Somei, M. *et al.*, *Heterocycles*, 2007, **73**, 537-554 (*synth*)

2-Methyl-6-(1-propenyl)piperidine, 9CI M-522



(2R,6R)-form

C₉H₁₇N 139.24

(2R,6R)-form

Pinidine

[501-02-0]

Alkaloid from *Pinus sabiniana*, *Pinus jeffreyi* and *Pinus torreyana* (Pinaceae). Bp₇₅₁ 176-177°. [α]_D²⁵ -10.5.

Hydrochloride: [55399-22-9]

Mp 244-246°. [α]_D²⁴ -9.6 (c, 0.25 in EtOH).

Methiodide: Mp 214-218°.

(2S,6S)-form

ent-3-(+)-Pinidine

[55448-42-5]

Synthetic. Oil. [α]_D²⁴ +10.2 (c, 0.32 in EtOH).

Hydrochloride: [55399-24-1]

Needles (EtOH/Et₂O). Mp 246-248°.

(2RS,6RS)-form [55399-23-0]

Bp₇₆₂ 175-177°.

Hydrochloride: Mp 192-193°.

[96894-80-3]

Tallent, W.H. *et al.*, *J.A.C.S.*, 1955, **77**, 6361; 1956, **78**, 4467 (*isol, ir, struct*)

Hill, R.K. *et al.*, *Tetrahedron*, 1965, **21**, 147; 1977, **33**, 1569 (*struct, abs config*)

Leete, E. *et al.*, *J.O.C.*, 1975, **40**, 2151 (*synth*)

Leete, E. *et al.*, *Tet. Lett.*, 1975, 3779

(*biosynth*)

Arseniyadis, S. *et al.*, *Tet. Lett.*, 1985, **26**, 729 (*synth*)

Yamazaki, N. *et al.*, *J.A.C.S.*, 1989, **111**, 1396 (*synth, pmr, cmr, ms*)

Dolle, R.E. *et al.*, *Tet. Lett.*, 1991, **32**, 5029 (*synth*)

Takahata, H. *et al.*, *Tetrahedron: Asymmetry*, 1992, **3**, 607 (*synth*)

Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1993, **76**, 957 (*synth*)


Kirihara, M. *et al.*, *Tetrahedron*, 1999, **55**, 2911-2926 (*synth*)

Davison, E.C. *et al.*, *J.C.S. Perkin 1*, 2002, 1494-1514 (*synth*)

2-Methyl-5-(1-propenyl)pyrazine, 9CI M-523

[108653-51-6]

[55138-66-4, 18217-82-8]

C₈H₁₀N₂ 134.18

Constit. of roasted green tea, roast peanut, coffee and cooked beef. Present in okra pods. Characterised spectroscopically.

Bondarovich, H.A. *et al.*, *J. Agric. Food Chem.*, 1967, **15**, 1093-1099 (*occur, coffee, ms, ir, uv*)

Tas, A.C. *et al.*, *Riechst., Aromen, Koerperpflegem.*, 1974, **24**, 326; 328; 331-332; *CA*, **82**, 57645m (*synth, ms, pmr, ir*)

Ames, J.M. *et al.*, *Phytotherapy*, 1990, **29**, 1201-1207 (*occur, okra*)

2-Methyl-6-(1-propenyl)pyrazine, 9CI M-524

[104638-11-1]

[18217-81-7, 55138-67-5]

C₈H₁₀N₂ 134.18

Constit. of coffee, roasted sesame, roasted green tea and cooked beef. Present in okra pods. Characterised spectroscopically.

Bondarovich, H.A. *et al.*, *J. Agric. Food Chem.*, 1967, **15**, 1093-1099 (*occur, coffee, ir, ms, uv*)

Tas, A.C. *et al.*, *Riechst., Aromen, Koerperpflegem.*, 1974, **24**, 326; 328; 331-332; *CA*, **82**, 57645m (*synth, ms, pmr, ir*)

Shimoda, M. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 3193-3196 (*occur, sesame*)

2-Methyl-1-propylamine M-525

2-Methyl-1-propanamine, 9CI. Isobutylamine, 8CI. Valamine†
[78-81-9]
(H₃C)₂CHCH₂NH₂

C₄H₁₁N 73.138

Found in foodstuffs and tobacco smoke. Present in fungi. Liq. Misc. H₂O, EtOH, Et₂O. d₄²⁰ 0.74. Fp -84.6. Bp 68-69°. n_D²⁰ 1.3970. pK_a 10.48 (25°).

- Highly flammable, fl. p. -9°, autoignition temp. 378°. Very irritating to eyes, skin and mucous membranes. A vesicant. Inhalation can cause headache and dryness of nose and throat. LD₅₀ (rat, orl) 228 mg/kg. NP9900000

Hydrochloride: [5041-09-8]

Cryst. (EtOH/Et₂O). Mp 177-178° (168-171°).

N-Propyl: 2-Methyl-N-propyl-1-propylamine, 9CI. N-Propylisobutylamine. Isobutylpropylamine
[39190-66-4]

C₇H₁₇N 115.218

Found in foodstuffs and is a metabolite of *Streptococcus lactis*. Liq. with odour resembling fusel oil. Spar. sol. H₂O. Bp 123-125° (120°).

N-Propyl, hydrochloride:

Plates (EtOH/Et₂O). Mp 135°.

Dornow, A. et al., *Annalen*, 1954, **588**, 62 (derivs)

Cope, A.C. et al., *J.A.C.S.*, 1957, **79**, 4720 (derivs)

Feeny, J. et al., *J.C.S.*, 1961, 1123 (pmr)


Faure, R. et al., *Org. Magn. Reson.*, 1980, **14**, 20 (cmr)

Harrison, A.G. et al., *Can. J. Chem.*, 1986, **64**, 1652 (ms)

Smirnov, Y.D. et al., *Zh. Org. Khim.*, 1992, **28**, 461; *J. Org. Chem. USSR (Engl. Transl.)*, 1992, **28**, 374 (N-alkyl derivs)

N-(2-Methylpropyl)-3-(3-oxo-1-cyclopenten-1-yl)propanamide M-526

[878633-76-2]




C₁₂H₁₉NO₂ 209.288

Prod. by *Streptomyces* sp. GT-20026114 isol. from the mangrove plant *Aegiceras comiculatum*. Amorph. powder. λ_{max} 242 (MeOH).

Lin, W. et al., *J. Antibiot.*, 2005, **58**, 594-598 (isol, pmr, cmr)

2-(1-Methylpropyl)-5-phenyl-1H-imidazole M-527

Catharsitoxin E



C₁₃H₁₆N₂ 200.283

Named in CAS as the 3H- (4-phenyl) isomer. In the paper it is shown as the 5-Ph isomer illus.


(ξ)-form [376587-14-3]

Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.

Suenaga, K. et al., *Tet. Lett.*, 2001, **42**, 7079-7081 (isol, pmr)

2-(2-Methylpropyl)-5-phenyl-1H-imidazole M-528

2-Isobutyl-5-phenyl-1H-imidazole. Catharsitoxin D
[376587-13-2]




C₁₃H₁₆N₂ 200.283

Named in CAS as the 3H (4-phenyl) isomer. In the paper it is shown as the 5-Ph isomer illus. Isol. from the Chinese remedy Qiung laug prepd. from the beetle *Catharsius molossus*. Oil.

Suenaga, K. et al., *Tet. Lett.*, 2001, **42**, 7079-7081 (isol, synth, pmr, cmr)

2-Methyl-6-propylpiperidine, 9CI M-529

[68170-79-6]



(2R,6R)-form

C₉H₁₉N 141.256

(2R,6R)-form

Epidihydropinidine

[135910-40-6]

Alkaloid from *Picea* spp. and *Pinus engelmannii*. Free base is unstable.

Hydrochloride: [135968-13-7]

Cryst. (2-propanol/EtOAc). Mp 171-173° (162-163°). [α]_D²⁰ +5.1 (c, 0.4 in EtOH).

(2R,6S)-form [65266-41-3]

Oil. Bp 175-180°.

Hydrochloride: [115014-58-9]

Mp 245-247°. [α]_D²⁰ +14.2 (c, 1.05 in EtOH).

(2S,6R)-form

Dihydropinidine

[65337-41-9]

Alkaloid from the beetle *Epilachna varivestis*. Oil. Bp 175-180°.

Hydrochloride: [65375-67-9]

Needles (Et₂O). Mp 239-240°. [α]_D -13 (c, 0.07 in EtOH).

(2RS,6RS)-form

(±)-trans-form

[84367-22-6]

[151851-20-6]

Hydrochloride: [175478-17-8]

Cryst. (Et₂O/MeOH). Mp 134-135°.

(2RS,6SR)-form

(±)-cis-form

[65337-42-0]

[109583-97-3]

Oil. Bp 176-177°.

Hydrochloride: [121963-72-2]

[109583-96-2]

Needles (EtOAc/EtOH). Mp 215-217° (212-213°).

Waelchli, P.C. et al., *Helv. Chim. Acta*, 1978, **61**, 921-928 (cd)

Bonin, M. et al., *Tet. Lett.*, 1982, 3369-3372 (synth)

Husson, H.P. et al., *J. Nat. Prod.*, 1985, **48**, 894-906 (rev. synth)

Ryckman, D.M. et al., *J.O.C.*, 1987, **52**, 4274-4279 (2RS,6SR-form, synth, ir, pmr, cmr, ms)

Watanabe, Y. et al., *J.O.C.*, 1989, **54**, 4088-4097 (synth)

Momose, T. et al., *Chem. Lett.*, 1990, 1319-1322 (2R,6S-form, synth)

Schneider, M.J. et al., *J. Nat. Prod.*, 1991, **54**, 905-909 (2R,6R-form, isol, pmr, cmr)

Comins, D.L. et al., *J.O.C.*, 1991, **56**, 2506-2512 (synth, bibl)

Tawara, J.N. et al., *J.O.C.*, 1993, **58**, 4813-4818 (2R,6R-form, isol, cryst struct, abs config)

Chênevert, R. et al., *J.O.C.*, 1996, **61**, 3332 (synth)

Amat, M. et al., *Tetrahedron: Asymmetry*, 1996, **7**, 977 (synth, bibl)

Takahara, H. et al., *Tetrahedron: Asymmetry*, 1996, **7**, 3047-3054 (2R,6R-form, synth, ir, pmr, cmr)

Muraoka, O. et al., *J.C.S. Perkin 1*, 1997, 113-119 (2R,6S-form, synth)

Weymann, M. et al., *Synthesis*, 1997, 1151-1160 (synth, pmr, cmr)

Bubnov, Y.N. et al., *Tet. Lett.*, 1997, **38**, 4631 (synth)

Yamauchi, T. et al., *Chem. Pharm. Bull.*, 1998, **46**, 384-389 (2S,6R-form, synth, pmr)

Davis, F.A. et al., *Tet. Lett.*, 1998, **39**, 5951-5954 (synth)

Adamo, M.F.A. et al., *Synth. Commun.*, 1999, **29**, 1747-1756 (2R,6R-form, synth)

Wilkinson, T.J. et al., *Org. Lett.*, 2000, **2**, 155-158 (2R,6R-form, synth)

Loh, T.-P. et al., *Tet. Lett.*, 2000, **41**, 7779-7783 (synth)

Dondas, H.A. et al., *Tetrahedron*, 2002, **58**, 161-173 (synth)


Shu, C. et al., *J.A.C.S.*, 2003, **125**, 2878-2879 (2S,6R-form, synth)

Yamauchi, S. et al., *Biosci., Biotechnol., Biochem.*, 2004, **68**, 676-684 (synth)

Wang, X. et al., *J.O.C.*, 2005, **70**, 1897-1900 (2S,6R-form, synth)

3-(2-Methylpropyl)piperidine M-530

3-Isobutylpiperidine



C₉H₁₉N 141.256

(ξ)-form

N-Et: 1-Ethyl-3-(2-methylpropyl)piperi-

dine. 1-Ethyl-3-isobutylpiperidine. *Nor-stenusine*
[934416-51-0]
C₁₁H₂₃N 169.309
Found in beetles *Stenus* spp.


(±)-form

N-Et: Bp₁₅ 69°.

Gedig, T. et al., *Tetrahedron*, 2007, **63**, 2670-2674 (synth, pmr, cmr, ms)

2-(1-Methylpropyl)-4(1H)-quinolinone M-531

2-(1-Methylpropyl)-4-quinolinol. 4-Hydroxy-2-(1-methylpropyl)quinoline



C₁₃H₁₅NO 201.268

NH-form

N-Me: 1-Methyl-2-(1-methylpropyl)-4(1H)-quinolinone

[1008099-97-5]


C₁₄H₁₇NO 215.294

Alkaloid from *Boronia ternata* var. *elongata*. Brownish oil. [α]_D +2 (c, 1 in CHCl₃). λ_{max} 214 (log ε 4.62); 238 (log ε 4.67); 322 (log ε 4.14); 330 (log ε 3.97) (MeOH).

Agier, C. et al., *Nat. Prod. Res.*, 2007, **21**, 698-703 (isol, pmr, cmr, ms)

4-Methyl-6-propylquinolizidine M-532

Octahydro-4-methyl-6-propyl-2H-quinolizine, 9CI



Relative Configuration

C₁₃H₂₅N 195.347

(4S,6R,9aR)-form

Dendrobates Alkaloid 195C. Quinolizidine 195C

[120037-60-7]

[232602-07-2, 232602-08-3, 232602-06-1]

Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae) and from the Madagascar frog *Mantella madagascariensis* (Ranidae, subfamily Mantelinae).

Daly, J.W. et al., *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (ms, rev)

Jones, T.H. et al., *J. Chem. Ecol.*, 1999, **25**, 1179-1193 (synth, struct)

Garraffo, H.M. et al., *Rapid Commun. Mass Spectrom.*, 1999, **13**, 1553-1563 (ms, struct)

2-(2-Methylpropyl)thiazole, 9CI M-533

2-Isobutylthiazole, 8CI
[18640-74-9]



C₇H₁₁NS 141.237

Preorbital gland secretion of grey duiker *Sylvicapra grimmia*. Occurs in tomatoes. Liq. with tomato leaf odour. d₄²⁵ 1. Bp 180° (172°) Bp₁₀ 66°. n_D²⁵ 1.4939.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 109B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1492C (ir)

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 1232D (ir)

Cottet, R. et al., *Bull. Soc. Chim. Fr.*, 1967, 4499-4502 (synth, ir, pmr, uv)

Viani, R. et al., *Helv. Chim. Acta*, 1969, **52**, 887-891 (isol)

Buttery, R.G. et al., *J. Agric. Food Chem.*, 1971, **19**, 524-529; *CA*, **75**, 18920n (isol)


Tabacchi, R. et al., *Helv. Chim. Acta*, 1974, **57**, 324-336 (ms)

Pittet, A.O. et al., *J. Agric. Food Chem.*, 1974, **22**, 264-269; *CA*, **80**, 120827y (synth, ms, props)

Dubs, P. et al., *Synthesis*, 1974, 294-295 (synth, ir, pmr, ms)

Dirinck, P. et al., *J. Sci. Food Agric.*, 1976, **27**, 499-508 (isol)

Burger, B.V. et al., *Z. Naturforsch., C*, 1988, **43**, 731-736 (isol)

8-Methylpseudomanibacnine M-534

(8R,13aS)-form

C₁₉H₂₁NO₃ 311.38


(8R,13aS)-form [151757-08-3]

Alkaloid from the stem bark of *Aniba canelilla* (Lauraceae). [α]_D -61 (c, 1.1 in MeOH).

(8S,13aS)-form [151851-40-0]

Alkaloid from stem bark of *Aniba canelilla* (Lauraceae). [α]_D -62 (c, 0.3 in MeOH).

Oger, J.-M. et al., *Can. J. Chem.*, 1993, **71**, 1128 (isol, uv, pmr, cmr, ms, struct)

6-Methylpurine M-535

7H-form

C₆H₆N₄ 134.14

7H-Form predominates. Cryst. Sol. H₂O, butanol. Mp 236-237°.

▶ LD₅₀ (mam, skn) 200 mg/kg. UO8700000

1H-form [2004-03-7]

1-Me: 1,6-Dimethyl-1H-purine, 9CI

[80888-49-9]

C₇H₈N₄ 148.167

Solid (as hydroiodide). Mp 239-241° (hydroiodide).

7H-form

Isol. from *Collybia maculata* and *Collybia dryophila*. Melanin biosynthesis inhibitor. Mp 235°. Isol. paper draws prod. as 9H-form.

7-Me: 6,7-Dimethyl-7H-purine, 9CI

[66122-65-4]

C₇H₈N₄ 148.167

Cryst. Mp 228-230°.

9H-form

9-Me: 6,9-Dimethyl-9H-purine, 9CI

[14675-46-8]

C₇H₈N₄ 148.167

Cryst. Mp 108-108.5°. pK_a 3.16 (20°).

9-Ph: 6-Methyl-9-phenyl-9H-purine, 9CI

[140421-33-6]

C₁₂H₁₀N₄ 210.238

Needles. Mp 155-156°.

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 582B (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 707D (ir)

Gabriel, S. et al., *Ber.*, 1901, **34**, 1234-1257 (synth)

Heiss, J. et al., *Org. Mass Spectrom.*, 1970, **3**, 181-190 (ms)

Thorpe, M.C. et al., *J. Magn. Reson.*, 1974, **15**, 98-112 (cmr)

Taylor, E.C. et al., *J.A.C.S.*, 1974, **96**, 8095-8102 (synth)

Leonhardt, K. et al., *Z. Naturforsch., C*, 1987, **42**, 420-424 (isol)


Tanji, K. et al., *Chem. Pharm. Bull.*, 1992, **40**, 227-229 (9-Ph)

Thines, E. et al., *Z. Naturforsch., C*, 1995, **50**, 813-819 (isol, activity)

Methylpyrazine, 9CI, 8CI M-536

FEMA 3309

[109-08-0]



C₅H₆N₂ 94.116

Isol. from an arctic marine bacterium. Flavouring agent. Liq. with disagreeable picoline-like odour. Misc. H₂O, EtOH, Et₂O. d₄²⁰ 1.03. Bp 136-137°. n_D²⁰ 1.4953. pK_{a1} 1.41; pK_{a2} -4.89 (25°, H₂O).

▶ Emits highly toxic fumes when heated to dec. Flammable, fl. p. 50° (oc). UQ3675000

Picrate:

Yellow prisms (EtOH). Mp 133°.

Methodide: [34260-05-4]

Leaflets and needles (EtOH). V. sol.

H₂O. Mp 129-130°.

1-Oxide: [31396-35-7]

C₅H₆N₂O 110.115

Mp 93-95°.

4-Oxide: [25594-37-0]

C₅H₆N₂O 110.115

Mp 69-71°.

l,4-Dioxide: [32046-26-7]C₅H₆N₂O₂ 126.115

Cryst. (MeOH). Mp 230-231°.


Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 840A (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, 3, 397C (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, 3, 1555D (*ir*)Jones, R.G. *et al.*, *J.A.C.S.*, 1950, 72, 3539 (*synth*)Kitchen, L.J. *et al.*, *J.A.C.S.*, 1951, 73, 1838 (*synth*)Cox, R.H. *et al.*, *J. Phys. Chem.*, 1968, 72, 1646 (*pmr*)Uchimaru, F. *et al.*, *J. Het. Chem.*, 1971, 8, 99 (*ms*)Sanyal, N.K. *et al.*, *J. Mol. Spectrosc.*, 1979, 78, 335 (*uv*)Matsuo, M. *et al.*, *Org. Magn. Reson.*, 1980, 13, 172 (*cmr*)Gumbley, S.J. *et al.*, *J. Het. Chem.*, 1985, 22, 1143 (*props*)Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, 2, 318-353 (*marine isol*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MOW750**2-Methylpyridine, 9CI**

M-537

α-Picoline

[109-06-8]

[1333-41-1]

C₆H₇N 93.128Isol. from leaves of *Rumex obtusifolius* (Polygonaceae). Found in coal tar oil.Manuf. by gas-phase catalytic reaction of acetaldehyde and ammonia. Used in manuf. of 2-Vinylpyridine and pharmaceuticals. Oil with strong unpleasant odour. d_4^{15} 0.95. Mp -70°. Bp 129°. n_D^{17} 1.5029. pK_a 7.98 (25°). pK_a 5.96 (25°).

- ▶ Flammable, fl. p. 28°, autoignition temp. 535/538°. Toxic, irritant. Skin and severe eye irritant. Adverse CNS effects by inhalation. LD₅₀ (rat, orl) 790 mg/kg. LD₅₀ (rbt, skn) 410 mg/kg. TJ4900000

Hydrochloride: [14401-91-3]Hygroscopic cryst. + ½H₂O. Mp 200° (anhyd.).*Hydrobromide, Br₂ addn. compd.*:

[135862-25-8]

Red cryst. Mp 76°.

Methodide: 1,2-Dimethylpyridinium iodide, 9CI. 2-Picoline methodide

[872-73-1]

C₇H₁₀IN 235.067

Reagent for the detn. of aromatic aldehydes (forms 2-stilbazole methodide). Needles (EtOH). Mp 229-231°.

- ▶ TJ7700000

Ethobromide: 1-Ethyl-2-methylpyridinium bromide, 9CI

[32353-50-7]

Red solid. Mp 65° Mp 97°.

Ethiodide: 1-Ethyl-2-methylpyridinium iodide, 9CI

[19760-15-7]

Mp 123°.

- ▶ TJ7696000

Picrate: [15938-03-1]

Needles. Mp 169-171°.

N-Oxide: [931-19-1]C₆H₇NO 109.127

Hygroscopic oil or solid. Mp 48-50°.

Bp₁₅ 123-124°. Anhydrous samples crystallise to a low-melting solid, which rapidly liquefies on exp. to air.


[18241-33-3]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 241C (*nmr*)*Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, 2, 912D (*ir*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, 3, 1514D; 1575C (*ir, N-oxide*)Heap, T. *et al.*, *J.A.C.S.*, 1921, 43, 1936 (*isol*)Wilkie, K. *et al.*, *J. Soc. Chem. Ind., London*, 1927, 46, 469 (*purifn*)Cartwright, C.H. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1935, 200, 914 (*ir*)Cullinane, N.M. *et al.*, *J. Soc. Chem. Ind., London*, 1948, 67, 142 (*synth*)Boekelheide, V. *et al.*, *J.A.C.S.*, 1954, 76, 1286 (*N-oxide, synth*)Coulson, E.A. *et al.*, *J.C.S.*, 1954, 1957 (*purifn*)Kosower, E.M. *et al.*, *J.A.C.S.*, 1956, 78, 3493 (*synth, methodide*)Wilkinson, S. *et al.*, *Nature (London)*, 1958, 181, 636 (*isol*)Weissberger, A. *et al.*, *Chem. Heterocycl. Compd.*, 1960, 14, 99Salsmans, R. *et al.*, *Org. Mass Spectrom.*, 1974, 8, 357 (*ms*)Bist, H.D. *et al.*, *Indian J. Pure Appl. Phys.*, 1983, 21, 236 (*ir, Raman*)Denisov, A.Y. *et al.*, *Khim. Geterotsikl. Soedin.*, 1984, 1223; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1984, 1000 (*pmr, cmr*)Kosmidis, A. *et al.*, *J. Mol. Spectrosc.*, 1986, 120, 11 (*uv*)Tarek, M. *et al.*, *Anal. Lett.*, 1987, 20, 2049; 1989, 22, 3091 (*use, methodide*)Cook, I.B. *et al.*, *Aust. J. Chem.*, 1989, 42, 1493 (*cmr*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MOY000*Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn.*, (ed. Snyder, R.), Elsevier, Volume 2, 1990, 218Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 2211Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 868**1-(5-Methyl-3-pyridinyl)-1-**

M-538


decanone*3-Decanoyl-5-methylpyridine*

[194345-50-1]

C₁₆H₂₅NO 247.38Constit. of *Piper retrofractum* (Javanese long pepper). Cryst. (CHCl₃). Mp 208°.Pande, A. *et al.*, *Indian J. Chem., Sect. B*, 1997, 36, 377-379 (*isol, ir, pmr, ms*)**2-Methyl-5-pyrimidinecarboxylic acid, 9CI**

M-539

[5194-32-1]

C₆H₆N₂O₂ 138.126Light tan solid or solid (H₂O). Mp 197-198°.*Me ester*: [5571-03-9]C₇H₈N₂O₂ 152.152

Cryst. Mp 78°.

Et ester: [2134-38-5]C₈H₁₀N₂O₂ 166.179Bp_{0.45} 70-71°.*Amide*: [5194-33-2]C₆H₇N₃O 137.141

Solid (butanol). Mp 218-219°.

*Methylamide: N,2-Dimethyl-5-pyrimidinecarboxamide*C₇H₉N₃O 151.168

Prod. by an unidentified marine bacterium.

Nitrile: 5-Cyano-2-methylpyrimidine

[5506-97-8]

C₆H₅N₃ 119.126


Solid by subl. Mp 72°.

May, C.D. *et al.*, *J.C.S. (C)*, 1966, 649-653(*synth, Me ester, Et ester, amide, nitrile*)Kress, T.J. *et al.*, *Heterocycles*, 1994, 38, 1375-1382 (*synth, uv, pmr*)Laatsch, H. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*methylamide*)**5-Methyl-2,4,6(1H,3H,5H)-pyrimidinetrione, 9CI**

M-540


5-Methylbarbituric acid

[2417-22-3]

C₅H₆N₂O₃ 142.114Metab. of Thymine catabolism in bacteria. Cryst. (H₂O). Mp 203°.Doumas, B. *et al.*, *J. Biol. Chem.*, 1962, 237, 2306 (*synth*)Arct, B. *et al.*, *CA*, 1967, 67, 54098 (*synth*)**4-Methyl-1H-pyrrole-2-carboxylic acid, 9CI**

M-541

[18711-59-6]

C₆H₇NO₂ 125.127Mp 203-204°. pK_{a1} 4.6.*Me ester: Methyl 4-methylpyrrole-2-carboxylate*

[34402-78-3]

C₇H₉NO₂ 139.154Trail pheromone of *Atta texana*. Cryst.

(hexane). Mp 73-74°.

Et ester: [40611-85-6]

C₈H₁₁NO₂ 153.18

Cryst. (EtOH aq.). Mp 37-38°.

Rapoport, J. et al., *J.O.C.*, 1964, **29**, 2727

Tumlinson, J.H. et al., *Nature (London)*, 1971, **234**, 348

Sonnet, P.E. et al., *J. Med. Chem.*, 1972, **15**, 97

Alonso Garrido, D.O. et al., *J.O.C.*, 1988, **53**, 403 (ester, synth, cmr)

Cornforth, J. et al., *J.C.S. Perkin 1*, 1990, 1459

(synth, Me ester, pmr)

Lash, T.D. et al., *J. Het. Chem.*, 1991, **28**, 1671

(synth)


Zimmer, R. et al., *Annalen*, 1992, 709 (synth, Me ester)

Tehrani, K.A. et al., *Tetrahedron*, 1999, **55**, 4133-4152 (synth)

3-Methyl-1H-pyrrole-2,5-dione, 9CI

M-542

2-Methylmaleimide. Citraconimide, 8CI [1072-87-3]



C₅H₅NO₂ 111.1

Prod. of chromic acid oxidation of tetrapyrroles. Cryst. (C₆H₆ or Et₂O/petrol). Mp 103-104° Mp 108° (107°). N-substituted derivs. undergo radical homo- and copolym.

5-Oxime: 3-Methylmaleimide 5-oxime

C₅H₆N₂O₂ 126.115

Isol. from the sponges *Cliona patera* and *Pseudoceratina purpurea*. Cryst. (MeOH). Mp 170-172°.

N-Me: 1,3-Dimethyl-1H-pyrrole-2,5-dione, 9CI

[4050-34-4]

C₆H₇NO₂ 125.127

Oil. Bp₁₀ 86°. n_D²⁰ 1.4970. N-Alkylcitraconimides polym. by a radical mechanism to give linear systems with the imide rings intact. Q/e values for radical polym., Q 0.87, e +1.59.

N-Ph: 3-Methyl-1-phenyl-1H-pyrrole-2,5-dione, 9CI

[3120-04-5]

C₁₁H₉NO₂ 187.198

Mp 99°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1454B (nmr)

Protopopescu, L. et al., *CA*, 1944, **38**, 3958h (synth)

Brown, P.M. et al., *J.C.S.*, 1957, 2882 (synth)

Sheremeteva, T.V. et al., *CA*, 1959, **53**, 4127h (N-Me)

Kojima, K. et al., *J. Polym. Sci., Part A*, 1966, **4**, 1121 (synth)

Sheremeteva, T.V. et al., *J. Polym. Sci., Part C*, 1967, **16**, 1631 (polym, derivs)

Ellsworth, R.K. et al., *Arch. Biochem.*

Biophys., 1968, **124**, 358

Chenon, M.T. et al., *J. Het. Chem.*, 1973, **10**, 427 (cmr)

Gossauer, A. et al., *Annalen*, 1974, 1496 (synth)

Hebenbrock, K.-F. et al., *Annalen*, 1978, 320 (derivs)

Earl, R.A. et al., *J. Het. Chem.*, 1978, **15**, 1479 (synth, pmr, imide)

Greenley, R.Z. et al., *J. Macromol. Sci., Part A: Chem.*, 1980, **14**, 427 (Q/e values, deriv)

Pyriadi, T.M. et al., *J. Polym. Sci., Polym.*

Chem. Ed., 1980, **18**, 2535 (derivs)

Martin, J. et al., *Tetrahedron*, 1980, **36**, 3261 (synth, ms)

Oishi, T. et al., *Polym. J. (Tokyo)*, 1981, **13**, 65 (polym, reactivity, derivs)

Yogo, M. et al., *J.C.S. Perkin 1*, 1984, 2097 (synth, ir, uv, pmr, imides)

Bestmann, H.J. et al., *Chem. Ber.*, 1985, **118**, 2640 (deriv)

Otsu, T. et al., *Polym. Bull. (Berlin)*, 1990, **24**, 475 (polym, derivs)


Gill, G.B. et al., *J.C.S. Perkin 1*, 1993, 2567 (synth, ir, pmr, cmr)

Kijjoo, A. et al., *Z. Naturforsch., B*, 2005, **60**, 904-908 (5-oxime, isol, pmr, cmr)

1-Methylpyrrolidine, 9CI

M-543

[120-94-5]



C₅H₁₁N 85.149

Minor alkaloid from tobacco (*Nicotiana tabacum*), also found in *Atropa belladonna* (Solanaceae). Liq. Misc. H₂O. Bp₇₅₀ 80-81°. n_D²⁰ 1.4311. pK_a 10.32 (25°). Volatile in Et₂O vapour.

► Highly flammable, fl. p. -14°. Corrosive and irritating to skin, eyes and mucous membranes. UY1420500

Methobromide: [23827-15-8]

Cryst. (EtOAc/EtOH). Mp 225°.

Picrate:

Yellow plates (EtOH). Mp 224°.

N-Oxide: [7529-17-1]

C₅H₁₁NO 101.148

Cryst. Mp 60°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 558A (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 354B (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 443A (ir)

Späth, E. et al., *Ber.*, 1939, **72**, 1809 (isol)

Org. Synth., Coll. Vol., **3**, 1955, 159 (synth)

Yasumatsu, N. et al., *CA*, 1965, **63**, 18668a (isol)

U.S. Pat., 1966, 3 239 535; *CA*, **64**, 17815f (oxide)

Ferles, M. et al., *Coll. Czech. Chem. Comm.*, 1968, **33**, 2121 (synth)

Osman, S. et al., *Org. Mass Spectrom.*, 1969, **2**, 977 (ms)

Hawthorne, D.G. et al., *Aust. J. Chem.*, 1976, **29**, 515 (cmr)

Leung, J. et al., *Org. Magn. Reson.*, 1977, **9**, 333 (cmr)


Beecroft, R.A. et al., *J.C.S. Perkin 2*, 1985, 1063 (uv)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MPB250

3-Methylpyrrolidine, 9CI

M-544

[34375-89-8]



(R)-form

C₅H₁₁N 85.149

(R)-form

Bp 103°. [α]_D²² +17.2 (c, 0.9 in EtOH). n_D²² 1.4420. Abs. config. assignment contradictory before 1978.

Hydrochloride: [α]_D²⁰ +6.9 (c, 1.67 in MeOH) (>98% ee).

N-Benzoyl:

C₁₂H₁₅NO 189.257

Solid (hexane). Mp 64-65°. [α]_D²⁰ +70.2 (c, 1.03 in CH₂Cl₂).

N-(2-Phenylethyl): 3-Methyl-1-(2-phenylethyl)pyrrolidine

[163315-04-6]

C₁₃H₁₉N 189.3

Trace alkaloid from poison glands of *Harpagoxenus sublaevis*, *Leptothorax acervorum*, *Leptothorax muscorum* and *Doronomyrmex goesswaldi*. Oil.

N-(2-Methylbutyl): 3-Methyl-1-(2-methylbutyl)pyrrolidine

[163315-00-2]

C₁₀H₂₁N 155.283

Trace alkaloid from poison glands of ants *Harpagoxenus sublaevis*, *Leptothorax acervorum* and *Leptothorax muscorum*. Abs. config. of the side-chain not known (1997).

N-(3-Methylbutyl): 3-Methyl-1-(3-methylbutyl)pyrrolidine. **Leptothoracine** [163314-99-6]

C₁₀H₂₁N 155.283

Trace alkaloid from poison glands of the ants *Harpagoxenus sublaevis*, *Leptothorax acervorum*, *Leptothorax muscorum* and *Doronomyrmex goesswaldi*. Bp₃₀ 88°.

(S)-form [69498-25-5]

Bp 103°. [α]_D²² -18.7 (c, 1.4 in EtOH).

N-Nitroso: [α]_D²² -42 (c, 1 in EtOH). n_D²² 1.4813.

► Exp. carcinogen.

(±)-form

Liq. with odour resembling piperidine. Bp 103-105°. V. air-sensitive.

Picrate:

Cryst. (EtOH). Mp 106°.

N-Me: 1,3-Dimethylpyrrolidine

[45470-22-2]

C₆H₁₃N 99.175

Liq. d₄⁵ 0.79. Mp 181-182° (picrate)

Mp 110-115° (picrate). Bp 96-97°.

Picrate is dimorphous.

N-Isopropyl: 1-Isopropyl-3-methylpyrrolidine. 3-Methyl-1-(1-methylethyl)pyrrolidine, 9CI

C₈H₁₇N 127.229

Liq. Sol. Et₂O. Bp 150-165°.

N-Isopropyl, picrate:

Yellow needles (EtOH). Mp 140-141°.

Löffler, K. et al., *Ber.*, 1910, **43**, 2044 (N-Me)

Elderfield, R.C. et al., *J.A.C.S.*, 1950, **72**, 1334 (N-isopropyl)

Ringdahl, B. et al., *Acta Pharm. Suec.*, 1978, **15**, 255 (abs config)

Lin, T. et al., *J. Het. Chem.*, 1984, **21**, 1871 (synth, ir, pmr, derivs)

Budrham, R.S. et al., *Org. Prep. Proced. Int.*, 1986, **18**, 295 (synth, pmr)

Reder, E. et al., *Helv. Chim. Acta*, 1995, **78**, 73 (alkaloids)

Koob, R. et al., *Helv. Chim. Acta*, 1997, **80**, 267-272 (abs config, alkaloids)

- Veith, H.J. *et al.*, *Liebigs Ann./Recl.*, 1997, 391 (synth, pmr)
 Andrés, C. *et al.*, *J.O.C.*, 1999, **64**, 4273-4281 (R-form, S-form, synth, pmr, cmr)
 Zhang, X. *et al.*, *Nat. Prod. Lett.*, 2002, **16**, 53-56 (Leptothoracine, synth)

5-Methyl-2-pyrrolidinecarboxylic acid **M-545**

5-Methylproline, 9CI
 [16639-15-9]



C₆H₁₁NO₂ 129.158

(2S,5R)-form

Et ester:

C₈H₁₅NO₂ 157.212
 Oil. [α]_D²⁵ -38.9 (c, 0.95 in CHCl₃).

(2S,5S)-form

tert-Butyl ester: [599191-13-6]

C₁₀H₁₉NO₂ 185.266
 Oil. [α]_D²³ -11.8 (c, 0.74 in CHCl₃).

(2R*,5R*)-form

cis-form

Isol. from hydrolysates of Actinomycin Z.

(2RS,5RS)-form

(±)-cis-form

Et ester: Bp₃ 59°.

(2RS,5SR)-form

(±)-trans-form

Mp 207° (197-199°).

Sakurai, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1969, **42**, 3524 (synth)

Katz, E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1973, **52**, 819 (isol)

Collado, I. *et al.*, *J.O.C.*, 1995, **60**, 5011 (Et ester, synth, ir, pmr, cmr)


Mota, A.J. *et al.*, *Eur. J. Org. Chem.*, 2003, 4187-4198 (2S,5S-form, tert-butyl ester)

1-Methyl-2-pyrrolidinone, **M-546**
9CI

N-Methylpyrrolidone

[872-50-4]

[51013-18-4]



C₅H₉NO 99.132

Alkaloid from *Melicope hayesii*. Dipolar aprotic solvent, with low oral acute tox. Weakly basic, good proton acceptor from strong acids. Sol. H₂O, most org. solvs. d₄²⁰ 1.03. Mp -17° (-24°). Bp 202° Bp₁₀ 85°. n_D²⁵ 1.4690. pK_a -0.92 (H₂SO₄ aq.). Steam-volatile. Forms hydrates.

- Fl. p. 96° (oc), autoignition temp. 346°. Severe eye irritant. Exp. reprod. and teratogenic effects. OES: long-term 100 ppm. UY5790000

Hydrochloride: [16889-93-3]

Mp 79-81°.

Oxime: [35197-40-1]

C₅H₁₀N₂O 114.147

Mp 98-99°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 789C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1286A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 788D (ir)

Fery, L.P.A. *et al.*, *Bull. Soc. Chim. Belg.*, 1960, **69**, 63 (synth)

Blicke, F.F. *et al.*, *J.O.C.*, 1961, **26**, 1826 (synth)

Metzger, H. *et al.*, *Angew. Chem., Int. Ed.*, 1963, **2**, 624 (synth)

Falbe, J. *et al.*, *Chem. Ber.*, 1965, **98**, 1928 (synth)

Basch, H. *et al.*, *J. Chem. Phys.*, 1968, **49**, 5007 (uv)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1969, **2**, 281 (use)

Fronza, G. *et al.*, *J.C.S. Perkin 2*, 1977, 1746 (N-15 nmr, cmr)

Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, **19**, 514

Cerioni, G. *et al.*, *Spectrochim. Acta A*, 1983, **39**, 929 (pmr, cmr)

Ruostesuo, P. *et al.*, *Spectrochim. Acta A*, 1985, **41**, 739 (cmr, N-15 nmr, O-17 nmr)

Ermakov, A.I. *et al.*, *Zh. Org. Khim.*, 1985, **21**, 1986 (ms)

Tyman, J. *et al.*, *Synth. Commun.*, 1989, **19**, 179 (use)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **5**, 3580-3581 (use)

Latip, J. *et al.*, *Phytochemistry*, 1999, **51**, 107-110 (isol, pmr, cmr)

Herler, S. *et al.*, *Acta Cryst. E*, 2007, **63**, o3991 (cryst struct)


Chemical Hazards of the Workplace, 2nd edn., (eds. Proctor, N.H. *et al.*), J.B. Lippincott, 1988, 344

Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 869

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, MPF200

3-(1-Methyl-2-pyrrolidinyl)-1,2-dithiolan-4-ol, **M-547**
9CI

1-Methyl-2-(4-hydroxy-1,2-dithiolan-3-yl)pyrrolidine



C₈H₁₅NOS₂ 205.345

CA numbering shown.

(2'R*,3R*,4R*)-form

cis,erythro-form. *Guinesine A*

[121702-91-8]

Alkaloid from the bark of *Cassipourea guianensis* (Rhizophoraceae). Exhibits insecticidal activity. Yellowish oil. [α]_D²⁴ +80.5 (c, 0.71 in CHCl₃). λ_{max} 242; 330 (solvent not reported) (Derep).

(2R*,3S*,4S*)-form

cis,threo-form. *Guinesine B*

[121702-92-9]

Alkaloid from the bark of *Cassipourea guianensis* (Rhizophoraceae). Exhibits insecticidal activity. Needles. Mp 61-62°. [α]_D²⁴ -36.5 (c, 0.34 in CHCl₃). λ_{max} 242; 330 (solvent not reported) (Derep).

(2'R*,3R*,4S*)-form

trans,erythro-form. *Guinesine C*

[121702-93-0]


Alkaloid from the bark of *Cassipourea guianensis* (Rhizophoraceae). Needles. Mp 76-77°. [α]_D²⁴ -4.8 (c, 0.48 in CHCl₃). λ_{max} 242; 330 (solvent not reported) (Derep).

Kato, A. *et al.*, *Tet. Lett.*, 1989, **30**, 3671 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Mitsudera, H. *et al.*, *J. Het. Chem.*, 1990, **27**, 1361 (synth)

2',3-(N-Methylpyrrolidinyl)-hygrine **M-548**

1-(1,1'-Dimethyl[2,3'-bipyrrolidin]-2'-yl)-2-propanone, 9CI



C₁₃H₂₄N₂O 224.345

Alkaloid from *Merremia* spp. Oil.


Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (isol, synth, pmr, cmr, ms)

2',4-(N-Methylpyrrolidinyl)-hygrine **M-549**

1-(1,1'-Dimethyl[2,3'-bipyrrolidin]-5'-yl)-2-propanone, 9CI

[110024-53-8]

[130404-24-9, 137328-59-7]



C₁₃H₂₄N₂O 224.345

Alkaloid from *Datura innoxia*, *Datura candida*, *Erythroxylum lucidum*, *Hyoscyamus albus* and *Merremia* spp. Oil. Isol. as a mixture of stereoisomers.


Witte, L. *et al.*, *Planta Med.*, 1987, **53**, 192-197 (isol, ms)

Brachet, A. *et al.*, *Phytochemistry*, 1997, **46**, 1439-1442 (isol)

Jenett-Siems, K. *et al.*, *Phytochemistry*, 2005, **66**, 1448-1464 (isol, synth, pmr, cmr, ms)

20-(2-Methyl-1-pyrrolin-5-yl)pregn-4-en-3-one **M-550**

20-(3,4-Dihydro-5-methyl-2H-pyrrol-2-yl)pregn-4-en-3-one, 9CI. 22,25-Epimino-27-norcholesta-4,24-dien-3-one [55486-07-2]




C₂₆H₃₉NO 381.6

Alkaloid from aerial parts of *Veratrum album* ssp. *lobelianum* (Liliaceae). Shows *in vitro* cytotoxicity against leukaemia P388 cells. Cryst. (Et₂O). Mp 153-157°. [α]_D²⁴ +107 (c, 0.64 in CHCl₃). Stereochem. assigned from subsequently determined x-ray struct. of Veracintine, V-57.

Vassová, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1975, **40**, 695-698 (*isol*, *pmr*, *uv*, *struct*)
Tomko, J. *et al.*, *CA*, 1981, **94**, 205391r (*isol*)
Fuska, J. *et al.*, *CA*, 1982, **96**, 173914m (*pharmacol*)

3-Methyl-1H-pyrrolo[3,2- M-551**b]pyridine, 9CI****4-Azaskatole**

[25796-94-5]

C₈H₈N₂ 132.165

Isol. from fish meal tar. Antifungal agent. Cryst. (C₆H₆). Mp 228-230°. λ_{\max} 225 (ε 21950); 292 (ε 6950) (MeOH) (Berdy).


Kelly, A.H. *et al.*, *J.C.S. (C)*, 1970, 303-307 (*synth*)

Wakabayashi, K. *et al.*, *Yakugaku Zasshi*, 1978, **98**, 898-903 (*isol*)

Fontan, R. *et al.*, *Heterocycles*, 1981, **16**, 1473-1474 (*synth*)

4-Methyl-3H-pyrrolo[2,3- M-552**c]quinoline****Marinoquinoline A**

[920317-36-8]

C₁₂H₁₀N₂ 182.224


Alkaloid from the marine-derived gliding bacterium *Rapidithrix thailandica* GB009. Acetylcholinesterase inhibitor. Needles (Me₂CO/CHCl₃/hexane).

Kanjana-opas, A. *et al.*, *Acta Cryst. E*, 2006, **62**, o2728-o2730 (*cryst struct*)

Sangnoi, Y. *et al.*, *Mar. Drugs*, 2008, **6**, 578-586 (*activity*)

2-Methyl-1-(1H-pyrrol-1-yl)- M-553**8-decene-1,3-dione****1-(2-Methyl-3-oxo-8-decenoyl)-1H-pyrrole, 9CI**

[217461-35-3]

C₁₅H₂₁NO₂ 247.336

Alkaloid from *Penicillium brevicompactum*.

tum. Oil. [α]_D²⁰ +23 (c, 0.2 in CHCl₃).

2,3-Dihydro: 1-(2,3-Dihydro-1H-pyrrol-1-yl)-2-methyl-8-decene-1,3-dione. 2,3-Dihydro-1-(2-methyl-3-oxo-8-decenoyl)-1H-pyrrole, 9CI

[220568-62-7]

C₁₅H₂₃NO₂ 249.352

Prod. by *Penicillium brevicompactum*.

8',9'-Dihydro: 2-Methyl-1-(1H-pyrrol-1-yl)-1,3-decanedione, 9CI. N-(2-Methyl-3-oxodecanoyl)-1H-pyrrole

[217461-49-9]

C₁₅H₂₃NO₂ 249.352

Prod. by *Penicillium brevicompactum*.

Oil. [α]_D¹⁹ +14 (c, 0.14 in CHCl₃).

Cantin, A. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 4748-4753 (*isol*, *8',9'-dihydro*)


Cantin, A. *et al.*, *Eur. J. Org. Chem.*, 1999, 221-226 (*2,3-dihydro, isol, ir, pmr, cmr, ms*)

Castillo, M.-A. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 2120-2124 (*2,3-dihydro*)

Clark, D. *et al.*, *Tetrahedron*, 2000, **56**, 6181-6184 (*synth*)

4-Methylquinazoline, 9CI M-554

[700-46-9]

C₉H₈N₂ 144.176

Alkaloid prod. by the bacteria *Pseudomonas aeruginosa*, *Sarcina lutea* and (in trace amounts) *Pseudomonas fluorescens* and *Pseudomonas putida*. Pale-yellow prisms. Mp 36-37°. Bp 260° Bp_{0.15} 78°. pK_{a1} 2.35; pK_{a2} -4.4 (20°, H₂SO₄ aq.).

Picrate:

Yellow-green flakes (EtOH). Mp 182-183.5°.

Bogert, M.T. *et al.*, *J.A.C.S.*, 1924, **46**, 1702; 1932 (*synth*)

Elderfield, R.C. *et al.*, *J.O.C.*, 1951, **16**, 1669 (*synth*)


Schofield, K. *et al.*, *J.C.S.*, 1952, 1924 (*synth*)

Mann, S. *et al.*, *Arch. Mikrobiol.*, 1967, **56**, 324

Batterham, T.J. *et al.*, *J.C.S. (B)*, 1967, 892 (*ms*)

Kant, J. *et al.*, *J. Het. Chem.*, 1985, **22**, 1313 (*synth*)

Uff, B.C. *et al.*, *J.C.S. Perkin 1*, 1986, 2295 (*synth, pmr*)

4-Methyl-2-quinazolinecar- M-555**boxylic acid**C₁₀H₈N₂O₂ 188.185

Trace metab. prod. by the bacterium *Pseudomonas aeruginosa*.

Amide: 4-Methyl-2-quinazolinecarboxamide. 2-Carbamoyl-4-methylquinazoline

[13535-92-7]

C₁₀H₉N₃O 187.201

Metab. of *Pseudomonas aeruginosa*.

Cryst. (EtOH). Mp 235.5°.


Bogert, M.T. *et al.*, *J.A.C.S.*, 1924, **46**, 1702

(synth)

Mann, S. *et al.*, *Arch. Mikrobiol.*, 1967, **56**, 324 (*occur*)

1-Methyl-4(1H)-quinazoli- M-556**none, 9CI****Glycorine**

[3476-68-4]

C₉H₈N₂O 160.175

Minor alkaloid from the leaves of *Glycosmis arborea* (Rutaceae). Mp 145-147°.

▶ VA3676800

Hydrochloride:

Needles + ²/₃ H₂O (EtOH/Me₂CO). Mp 242° dec.

Picrate: Mp 249° dec.

Pakrashi, S.C. *et al.*, *Tetrahedron*, 1963, **19**, 1011-1026 (*uv, ir, pmr, ms, struct, synth*)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 2347-2349 (*synth*)

Bhattacharyya, P. *et al.*, *Chem. Ind. (London)*, 1978, 532 (*synth*)


Bhattacharyya, J. *et al.*, *Heterocycles*, 1980, **14**, 1469-1473 (*cmr*)

Ozaki, K. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2234-2243 (*synth*)

2-Methylquinoline, 9CI M-557**Quinaldine, 8CI**

[91-63-4]

[27601-00-9]

C₁₀H₉N[⊕] 143.188

Present in coal tar and shale oil. Chemical scent constituent of the urine of the red fox *Vulpes vulpes*. Used as soln. in dil. H₂SO₄ for gravimetric detn. of Bi. Oil; turns reddish brown in air. Insol. H₂O; sol. Et₂O, CHCl₃. d₄²⁰ 1.06. Mp -2°. Bp 248° Bp₁₀₀ 171.5°. n_D²⁰ 1.6126. pK_a 5.71 (22°). pK_a 5.65 (25°).

▶ Fl. p. 110°. Skin and severe eye irritant. LD₅₀ (rat, orl) 1230 mg/kg. UZ9625000

Hydrochloride: [62763-89-7]

Needles (EtOH/Et₂O). Mp 224°.

Hydrobromide:

Cryst. (EtOH). Mp 227°.

Hydroiodide: [43101-13-9]

Needles (EtOH). Mp 186°.

Perchlorate: [41439-52-5]

Mp 131°.

N-Et: [606-55-3]

C₁₂H₁₄N 172.249

Mp 244° dec. (iodide). CAS no. refers to iodide.

▶ VA0740000

Picrate: [1773-07-5]

Yellow needles. Spar. sol. H₂O, EtOH. Mp 193-194°.

N-Oxide: [1076-28-4]

C₁₀H₉NO 159.187Needles (H₂O) (hydrated). Sol. dil. acids. Mp 77-78°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 853C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 421B; 421C (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1565B (ir)

Doebner, O. et al., Ber., 1883, 16, 2464 (synth)

Mills, W.H. et al., J.C.S., 1921, 1294 (synth)

Welcher, F.J. et al., Organic Analytical Reagents, Van Nostrand, New York, 1947, 3, 62 (detn, Bi)

Cervinka, O. et al., Coll. Czech. Chem. Comm., 1963, 28, 535 (synth)

Draper, P.M. et al., Can. J. Chem., 1968, 46, 1487 (ms)

Ghersesti, S. et al., Spectrosc. Lett., 1973, 6, 167 (ir)

Johns, S.R. et al., Aust. J. Chem., 1976, 29, 1617 (cmr)

Jorgenson, J.W. et al., Science (Washington, D.C.), 1978, 199, 796

Whitten, W.K. et al., J. Chem. Ecol., 1980, 6, 49

Song, Z. et al., J. Het. Chem., 1993, 30, 17 (synth)


Horaguchi, T. et al., J. Het. Chem., 2002, 39, 61-67 (synth, pmr, cmr)

Kitamura, M. et al., Synthesis, 2003, 2415-2426 (synth, ir, pmr, cmr)

Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold, 1992, QEJ000

1-Methyl-4(1H)-quinolinone, 9CI M-558**Echinopsine**

[83-54-5]

C₁₀H₉NO 159.187Alkaloid from the seeds of *Echinops ritro* and several other *Echinops* spp. (Asteraceae). Needles (C₆H₆). Mp 152°. pK_a 2.36 (30°).

▶ VC8618000

Hydrochloride:Cryst. (MeOH/Me₂CO). Mp 185-186°.**Picrate:**

Cryst. (EtOH). Mp 223-224°.

Imine: EchinopsidineC₁₀H₁₀N₂ 158.202Alkaloid from the aerial parts of *Echinops echinatus* (Asteraceae). Amorph.; cryst. (MeOH) (as hydrochloride). Mp 214-215° dec. (hydrochloride).

Greshoff, M. et al., Rec. Trav. Chim. (J. R. Neth. Chem. Soc.), 1900, 19, 360 (isol)

Späth, E. et al., Monatsh. Chem., 1922, 43, 469 (struct, synth)


Avramova, B. et al., Khim. Prir. Soedin., 1970, 6, 98; Chem. Nat. Compd. (Engl. Transl.), 1970, 6, 92 (synth, Echinopsidine)

Merchant, J.R. et al., Chem. Ind. (Dekker), 1979, 320 (synth)

Chaudhuri, P.K. et al., Phytochemistry, 1987, 26, 587 (Echinopsidine)

N-Methylsecoglaucine M-559

3,4,6,7-Tetramethoxy-N,N-dimethyl-1-phenanthreneethanamine, 9CI. 1-(2-Dimethylaminoethyl)-3,4,6,7-tetramethoxyphenanthrene. Glaucine methyl methine [66396-10-9]

C₂₂H₂₇NO₄ 369.46Aporphine-type numbering shown. Alkaloid from *Platycapnos spicata* and *Sarcocapnos enneaphylla* (Papaveraceae). Amorph. solid; cryst. (MeOH/Et₂O) (as hydrochloride). Mp 249-251° (hydrochloride).N-Me: [10313-69-6]
Mp 278° (273-274°).N-De-Me: 1-(2-Methylaminoethyl)-3,4,6,7-tetramethoxyphenanthrene. **Coryphenanthrine. Secoglaucine** [66190-61-2]C₂₁H₂₅NO₄ 355.433Alkaloid from the aerial parts of *Corydalis yanhusuo* (Papaveraceae).O¹-De-Me: **Thaliporphinemethine**

[100009-99-2]

C₂₁H₂₅NO₄ 355.433Alkaloid from *Illigera pentaphylla* (Hernandiaceae). Amorph.O⁹-De-Me: **Secoxanthoplanine**

[104385-49-1]

C₂₁H₂₅NO₄ 355.433Alkaloid from leaves of *Dehaasia triandra*. Amorph. solid.N,O⁹-Di-de-Me, N-Ac: 1-(N-Acetyl-N-methylamino)ethyl-7-hydroxy-3,4,6-trimethoxyphenanthrene. **N-Acetylseco-N-methylaurotetanine**

[143084-45-1]

C₂₂H₂₅NO₅ 383.443Alkaloid from heartwood of *Aromadendron elegans* (preferred genus name *Magnolia*) (Magnoliaceae). Needles (EtOAc/hexane). Mp 210-211°.O¹,O²-Di-de-Me, 1,2-methylene ether:**Dicentrine methine**C₂₁H₂₃NO₄ 353.417Alkaloid from *Atherosperma* sp. Cytotoxic agent. Shows antiplatelet and vasorelaxing activity.N,N,O¹-Tri-de-Me: 1-(2-Aminoethyl)-3,6,7-trimethoxy-4-phenanthrenol.**Hemiargine C**

[710948-56-4]

C₁₉H₂₁NO₄ 327.379Constit. of the leaves of *Croton hemiargyreus* var. *gymnodiscus*.

Bremner, J.B. et al., Aust. J. Chem., 1978, 31, 313 (synth)

Ross, S.A. et al., J. Nat. Prod., 1985, 48, 835 (Thaliporphinemethine)

Hu, T. et al., Nanjing Yaoxueyuan Xuebao, 1985, 16, 7; CA, 103, 175443u

Wu, Y.C. et al., Planta Med., 1989, 55, 163-165 (Dicentrine methine)

Blanco, O. et al., Heterocycles, 1990, 31, 1077 (isol, struct, synth)

Tojo, E. et al., Phytochemistry, 1991, 30, 1005 (isol)

Goh, S.H. et al., Phytochemistry, 1992, 31, 2495 (deriv)

Blanco, O.M. et al., Phytochemistry, 1993, 32, 1055 (isol, pmr, cmr, ms)


Chen, K.-S. et al., J. Nat. Prod., 1996, 59, 531-534 (Dicentrine methine, activity)

Lee, S.-S. et al., Tetrahedron, 1996, 52, 6561 (Secoxanthoplanine)


Lin, W.-H. et al., J. Chin. Pharm. Sci., 2003, 12, 117-122 (Hemiargine C)

Methylstemofoline

M-560

C₁₉H₂₃NO₅ 345.394Lower homologue of Stemofoline, S-515. Alkaloid from an unidentified *Stemona* sp. Yellow gum. [α]_D²³ +125 (c, 0.24 in CHCl₃).

Sastrarajji, T. et al., J. Nat. Prod., 2005, 68, 1763-1767 (isol, pmr, cmr)

9-Methylstreptimidone M-5614-(2-Hydroxy-5,7-dimethyl-4-oxo-6,8-decadienyl)-2,6-piperidinedione, 9CI. S 632A₂. TS 885. Antibiotic S 632A₂. Antibiotic TS 885 [51867-94-8]C₁₇H₂₅NO₄ 307.389Glutarimide-type antibiotic. Prod. by *Streptomyces hygroscopicus* S632 and *Streptomyces* sp. S-885. Antiviral and antifungal agent. NF-κB inhibitor. Pale-yellow oil. Sol. MeOH, EtOAc, CH₂Cl₂, CHCl₃, bases, C₆H₆; fairly sol. Et₂O; poorly sol. H₂O, acids, hexane, bases. Bp₁ 135-140°. [α]_D²⁰ +109.6 (c, 15 in CHCl₃). λ_{max} 232 (ε 23100); 291 (ε 790) (EtOH) (Derep). λ_{max} 231 (ε 15350); 283 (ε 1260); 291 (ε 1230) (EtOH) (Berdy).▶ LD₅₀ (mus, ivn) 210 mg/kg, LD₅₀ (mus, ipr) 280 mg/kg. TM75100008,9-Epoxyde: 8,9-Epoxy-9-methylstreptimidone. **Antibiotic S 632B₁**. S 632B₁ [121995-32-2, 122620-15-9]C₁₇H₂₅NO₅ 323.388Prod. by *Streptomyces hygroscopicus* S632. Active against *Saccharomyces* spp. Cytotoxic. Pale yellow oil. Sol. MeOH, CHCl₃, EtOAc; poorly sol.

H₂O. $[\alpha]_D^{25} +68$ (c, 1 in CHCl₃). λ_{\max} 206 (ε 14500); 282 (ε 450) (EtOH) (Derep).

5-Hydroxy: 5-Hydroxy-9-methylstreptimidone

[162901-85-1]

C₁₇H₂₅NO₅ 323.388

Prod. by a *Streptomyces* sp. HIL Y-9065404. Shows herbicidal props. Sol. CHCl₃, MeOH, EtOAc. $[\alpha]_D -52.3$ (c, 0.4 in CHCl₃). λ_{\max} 238 (ε 23000); 290 (ε 790) (MeOH) (Derep).

8E-Isomer: Antibiotic S 632A₃. S 632A₃

C₁₇H₂₅NO₄ 307.389

Prod. by *Streptomyces hygroscopicus* S632. Antitumour agent. Pale yellow oil. $[\alpha]_D^{28} +58.7$ (c, 0.15 in CHCl₃). λ_{\max} 233 ; 289 (MeOH).

Stereoisomer, 8,9-epoxide: Antibiotic S 632B₂. S 632B₂

C₁₇H₂₅NO₅ 323.388

From *Streptomyces hygroscopicus* S632. Similar activity as S 632B₁. Pale yellow oil. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{25} +73$ (c, 1 in CHCl₃). λ_{\max} 206 (ε 14500); 282 (ε 450) (EtOH) (Derep).

Saito, N. *et al.*, *J. Antibiot.*, 1974, **27**, 206-214 (isol, struct)

Allen, M.S. *et al.*, *Aust. J. Chem.*, 1976, **29**, 673 (isol, uv, ir, pmr, ms, biosynth)

Becker, A.M. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2393 (ir, uv, ms, pmr, abs config)

Otani, T. *et al.*, *J. Antibiot.*, 1989, **42**, 647-653; 654-661 (S 632B)

Chatterjee, S. *et al.*, *J. Antibiot.*, 1995, **48**, 271-273 (5-Hydroxy-9-methylstreptimidone)

Wang, Z.C. *et al.*, *Heterocycles*, 2006, **69**, 377-383 (isol, pmr, cmr, activity)

Cheng, C.-L. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 55-60 (S 632A₃)

Methylsuccirubine M-562

[1361-39-3]

C₂₀H₂₆N₂O₃ 342.437

Struct. unknown. Alkaloid from the bark of *Cinchona succirubra* (Rubiaceae). Mp 171-172°. $[\alpha]_D^{30} -87$ (EtOH).

Qudrat-i-Khuda, M. *et al.*, *Sci. Res. (Dacca)*, 1965, **2**, 1-7; *CA*, **63**, 12004g

10-(Methylsulfonyl)decanoic acid, 9CI M-563

MeSO₂(CH₂)₉COOH

C₁₁H₂₂O₄S 250.358

▶ HD6420000

Amide: 10-(Methylsulfonyl)decanamide.

Rorifamide

[53078-91-4]

C₁₁H₂₃NO₅S 249.374

Alkaloid from *Rorippa montana*.

Nitrile: 10-(Methylsulfonyl)decanenitrile.

9-Cyanononyl methyl sulfone. Rorifone

Isol. from *Rorippa montana*. Antitussive, expectorant. Chinese folk remedy. Prisms (EtOH). Mp 45-46°. Bp₁ 188-192°. Log P 1.2 (calc).

▶ LD₅₀ (mus, orl) 355 mg/kg.

Shanghai Inst. of Materia Medica, *et al.*, *Sci. Sin. (Engl. Edn.)*, 1973, **16**, 506; *CA*, **80**, 112554s (amide, nitrile)

Liang, Y. *et al.*, *Zhongguo Yaoli Xuebao (Acta Pharmacol. Sin.)*, 1987, **8**, 147; *CA*, **106**, 207133y (nitrile, occur)

Liang, Y. *et al.*, *Drugs of the Future*, 1988, **13**, 128 (nitrile, rev)

9-(Methylsulfonyl)nonylamine M-564

9-(Methylsulfonyl)nonanamine, 9CI. 9-Aminononyl methyl sulfone. Roripamine

[166546-98-1]

MeSO₂(CH₂)₈CH₂NH₂

C₁₀H₂₃NO₂S 221.363

Isol. from the whole herb of *Rorippa indica* (Brassicaceae). Pale yellow needles. Mp 153-155°.

N-Ac: [166406-57-1]


C₁₂H₂₅NO₃S 263.4

Mp 130-132°.

Lin, Y.-L. *et al.*, *Phytochemistry*, 1995, **39**, 919 (isol, ir, pmr, ms, synth, struct)

4-Methyl-1,2,6,8-tetraazacycloundeca-4,9-diene-3,7,11-trione, 9CI M-565

[246167-60-2]




C₈H₁₀N₄O₃ 210.192

Isol. from the marine brown alga *Sargassum vachellianum*. Cryst. (MeOH). Mp 228-230°.

Xu, S.H. *et al.*, *Chin. Chem. Lett.*, 1999, **10**, 401-402

12-Methyl-1-tetradecylamine M-566

12-Methyl-1-tetradecanamine, 9CI. 1-Amino-12-methyltetradecane. Medelamine B



C₁₅H₃₃N 227.432

(S)-form [168706-10-3]


Prod. by *Streptomyces* sp. NK14819.

Cytotoxic agent. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O.

Morino, T. *et al.*, *J. Antibiot.*, 1995, **48**, 904 (isol, props)

(2-Methyltetrahydrofuran)so-lanidine M-567

[868273-72-7]




Absolute Configuration

C₃₀H₄₇NO₂ 453.707

Related to Solanidine, S-345. Alkaloid from the aerial parts of *Solanum cornifolium*.

Luis, E.C.S. *et al.*, *Rev. Colomb. Quim.*, 2004, **33**, 7-12; *CA*, **143**, 418926m (isol)

α-Methyl-1-(2,3,4,9-tetrahydro-2-methyl-1H-carbazol-1-yl)-1H-indole-3-butanoic acid, 9CI M-568



C₂₆H₂₈N₂O₂ 400.519

Me ester: [131669-95-9]

C₂₇H₃₀N₂O₂ 414.546


Alkaloid from the roots of *Murraya glaberrima* (Rutaceae). Gum. $[\alpha]_D -7.5$.

Kumar, V. *et al.*, *Tet. Lett.*, 1990, **31**, 5217 (isol, struct)

4-Methylthiazole, 9CI M-569

FEMA 3716

[693-95-8]



C₄H₅NS 99.156

Present in asparagus and coffee. Flavouring ingredient. Liq. with nutty, green odour. Misc. H₂O. d_4^{25} 1.11. Bp 130-133° Bp₅₉ 70-71°. n_D^{25} 1.5234. pK_a 3.15 (25°).

▶ XJ5096000

Picrate: Mp 181°.

N-Me:

C₅H₈NS⁺ 114.191

Cryst. (EtOH/Et₂O) (as iodide). Mp 117-118.5° (iodide).

N-Et:

C₆H₁₀NS⁺ 128.218

Cryst. (EtOH/Et₂O) (as iodide). Mp 169-171° (iodide).

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 510D (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 643A (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1491D (ir)

Bergen, G. *et al.*, *Acta Chem. Scand.*, 1966, **20**, 2593 (pmr)

Vincent, E.J. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 3524 (nmr)

Colebourne, N. *et al.*, *J.C.S.(C)*, 1967, 685 (synth)

Gisele, D. *et al.*, *Spectrochim. Acta A*, 1967, **23**, 2669 (ir, Raman)


Haag, A. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 511 (ms)

Galy, J.P. *et al.*, *Can. J. Chem.*, 1978, **56**, 46 (cmr)

Amato, J.S. *et al.*, *Heterocycles*, 1984, **22**, 1947 (synth)
 Bordwell, F.G. *et al.*, *J.A.C.S.*, 1991, **113**, 985 (deriv. synth)
 Pavlik, J.W. *et al.*, *J.O.C.*, 1993, **58**, 3407 (synth)
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1861-1862 (occur, props)

2-Methylthiazolidine, 9CI M-570

Tetrahydro-2-methylthiazole. 2-Methyltetrahydrothiazole
 [24050-16-6]



C₄H₉NS 103.188

► LD₅₀ (mus, ipr) 250 mg/kg. XJ6061000

(±)-form

Used as aroma additive for food and tobacco.

Liq. Bp₃₆ 76-78° Bp₅ 37°.

(ξ)-form

Sex pheromone of cockroach *Nauphoeta cinerea*.

Surzur, J.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1969, **268**, 2109 (synth)

Hillers, S. *et al.*, *CA*, 1969, **71**, 70554y (synth)

Asinger, F. *et al.*, *Monatsh. Chem.*, 1970, **101**, 1298 (synth)

Yasuhara, A. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 2273 (synth, ms, pmr)


Barbry, D. *et al.*, *J.C.S. Perkin 2*, 1990, 133 (pmr, cmr, N-15 nmr)

Sirugue, D. *et al.*, *J. Chem. Ecol.*, 1992, **18**, 2261 (occur)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MPR000

4-(Methylthio)canthin-6-one M-571

4-Methylthio-6H-indolo[3.2.1-de][1,5]-naphthyridin-6-one, 9CI. Alkaloid C† [157770-30-4]



C₁₅H₁₀N₂OS 266.323

Alkaloid from the bark of *Pentaceras australis* and root bark of *Quassia africana*. Isol. from the mushroom *Boletus curtisii*. Antineoplastic agent. Needles (CHCl₃/EtOH). Mp 257-258°. Log P 3.07 (uncertain value) (calc). Incorr. indexed by CAS as 5-methylthio (1993).

Ayafor, J.F. *et al.*, *Bull. Chem. Soc. Ethiop.*, 1993, **7**, 121; *CA*, **121**, 175195m (isol, pmr, cmr, ms)

Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, ir, pmr, cmr, ms)

5-(Methylthio)canthin-6-one M-572

5-Methylthio-6H-indolo[3.2.1-de][1,5]-naphthyridin-6-one, 9CI

C₁₅H₁₀N₂OS 266.323

Alkaloid from the mushroom *Boletus curtisii*. Yellow solid. Mp 198-200°. λ_{max} 252 (log ε 4.27); 258 (log ε 4.28); 306 (log ε 3.9); 344 (sh) (log ε 3.77); 360 (log ε 4.07); 378 (log ε 4.23); 396 (log ε 4.05) (CHCl₃).

Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, ir, pmr, cmr, ms)

9-(Methylthio)canthin-6-one M-573

9-Methylthio-6H-indolo[3.2.1-de][1,5]-naphthyridin-6-one, 9CI

C₁₅H₁₀N₂OS 266.323

Alkaloid from the mushroom *Boletus curtisii*. Yellow solid. λ_{max} 208 (log ε 1); 284 (log ε 0.47); 318 (log ε 0.33); 362 (log ε 0.4) (CHCl₃).

Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, pmr, cmr, ms)

11-(Methylthio)canthin-6-one M-574

11-Methylthio-6H-indolo[3.2.1-de][1,5]-naphthyridin-6-one, 9CI


C₁₅H₁₀N₂OS 266.323

Alkaloid from the mushroom *Boletus curtisii*. Lemon-yellow solid. Mp 211-214°. λ_{max} 244 (log ε 4.04); 268 (log ε 4.05); 328 (sh) (log ε 3.72); 340 (log ε 3.75); 368 (log ε 3.75); 386 (log ε 3.87); 400 (sh) (log ε 3.71) (CHCl₃).

Bröckelmann, M.G. *et al.*, *Eur. J. Org. Chem.*, 2004, 4856-4863 (isol, uv, ir, pmr, cmr, ms)

2-(Methylthio)-3H-indol-3-one M-575

[35524-66-4]



C₉H₇NOS 177.226


Constit. of egg masses of muricid molluscs, *Ceratosoma erinaceum* and *Trunculariopsis trunculus*. Red needles (petrol). Mp 133-134°. λ_{max} 249 (ε 30300); 284 (ε 6500); 328 (ε 1440); 344 (ε 500); 428 (ε 1970); 450 (ε 2030) (no solvent reported).

Baker, J.T. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2467-2475 (synth, uv, ir)

Benkendorff, K. *et al.*, *Molecules*, 2001, **6**, 70-78 (isol)

2-(Methylthio)-3H-phenoazin-3-one, 9CI M-576

[115695-20-0]



C₁₃H₉NO₂S 243.286

Isol. from cultures of the mushroom *Calocybe gambosa* (St George's mushroom).

Kuchen, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 363-365

3-(Methylthio)-2-propenoic acid, 9CI M-577

3-(Methylthio)acrylic acid, 8CI

[26995-94-8]

MeSCH=CHCOOH

C₄H₆O₂S 118.156

Possesses antimicrobial props.

(E)-form [26398-93-6]

Metab. of *Streptomyces lincolnensis*. Cryst. (CCl₄). Mp 138-141° Mp 137°.

S-Oxide, (±)-form: 3-(Methylsulfinyl)-2-propenoic acid [121949-97-1]

C₄H₆O₃S 134.156

S,S-Dioxide: 3-(Methylsulfonyl)-2-propenoic acid

[63068-43-9]

C₄H₆O₄S 150.155

Cryst. (EtOAc). Mp 111-113°.

Benzyl ester: *Isotachin A*

[98259-97-3]

C₁₁H₁₂O₂S 208.281

Constit. of *Isotachis japonica* and *Balantiopsis rosea*.

2-Methoxybenzyl ester: *Isotachin C*

[107584-57-6]

C₁₂H₁₄O₃S 238.307

From *Balantiopsis rosea*. Oil.

2-Phenylethyl ester: *Isotachin B*

[98259-98-4]

C₁₂H₁₄O₂S 222.307

From *Balantiopsis rosea*.

4-Bromophenacyl ester: Mp 110-111°.

Amide: 3-(Methylthio)-2-propenamamide. 3-(Methylthio)acrylamide

[36846-63-6]

C₄H₇NOS 117.171

Alkaloid from *Clinacanthus siamensis* and metab. of *Streptomyces siyoaensis*. Needles (CHCl₃). Mp 117-118°.

Methylamide: N-Methyl-3-(methylthio)-2-propenamamide. *Penangine*

[150071-78-6]

C₅H₉NOS 131.198

Alkaloid from the leaves of *Glycosmis chlorosperma* and *Glycosmis citrifolia*. Cryst. (Et₂O). Mp 70-71°. λ_{max} 230 (sh); 261 (Et₂O).

Methylamide, S-oxide: *Glycothiomine A*

[167504-55-4]

C₅H₉NO₂S 147.198

Isol. from leaves of *Glycosmis citrifolia* (Rutaceae). Needles (CHCl₃). Mp 160-161°. λ_{max} 205; 260 (MeOH).

(Z)-form [70424-09-8]

Mp 99-100° Mp 123-123.9°.

Methylamide: *Isopenangine*

[150036-28-5]

Alkaloid from the leaves of *Glycosmis chlorosperma*. Cryst. (Et₂O). Mp 120-121°.

Methylamide, S-oxide: Glycothiomin B


[167504-56-5]

From leaves of *Glycosmis citrifolia* (Rutaceae). Yellow oil. λ_{\max} 211 ; 251 (sh) ; 293 (sh) (MeOH).

[63068-05-3]

Stoll, A. *et al.*, *Experientia*, 1956, **12**, 361 (synth)Novotny, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 2182 (synth)Hart, N.K. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1259Yagi, S. *et al.*, *Agric. Biol. Chem.*, 1972, **36**, 336 (isol)Surette, R. *et al.*, *J. Antibiot.*, 1976, **29**, 646 (biosynth)Topek, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 2395 (synth)Asakawa, Y. *et al.*, *Phytochemistry*, 1985, **24**, 1505; 1986, **25**, 2543 (Isotachins)Ikegami, F. *et al.*, *Phytochemistry*, 1989, **28**, 881 (oxide)Deshmukh, A.R.A.S. *et al.*, *Synth. Commun.*, 1990, **20**, 2259 (synth, Isotachin C)De Medeiros, E.F. *et al.*, *J.C.S. Perkin 1*, 1991, 2725 (synth)Greger, H. *et al.*, *Phytochemistry*, 1993, **32**, 933-936 (Penangin, Isopenangin)Hinterberger, S. *et al.*, *Tetrahedron*, 1994, **50**, 6279-6286 (Penangin, Isopenangin, synth)Wu, T.S. *et al.*, *Phytochemistry*, 1995, **39**, 1453-1457 (Glycothiomin)Hinterberger, S. *et al.*, *Tetrahedron*, 1998, **54**, 487-496 (dioxide, synth, pmr, cmr, ms)Tuntiwachwuttikul, P. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 1423-1425 (amide)Karrer, P. *et al.*, *Helv. Chim. Acta*, 1950, **33**, 1237-1245; 1951, **34**, 1392-1399 (synth, R-S-oxide)Dose, K. *et al.*, *Chem. Ber.*, 1957, **90**, 1251 (synth)Suyama, T. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1012; *CA*, **62**, 6552 (synth)Cheung, K.K. *et al.*, *Chem. Comm.*, 1965, 100-102 (R-S-oxide, abs config)Hartmann, T. *et al.*, *Naturwissenschaften*, 1968, **55**, 391 (isol)Pesch, R. *et al.*, *Org. Mass Spectrom.*, 1974, **9**, 861 (ms)Dalgaard, L. *et al.*, *Phytochemistry*, 1977, **16**, 931-932 (R-S-oxide, isol)Granvogel, M. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 1730-1739 (detn, cocoa)**12-Methyl-1-tridecylamine M-579***12-Methyl-1-tridecanamine, 9CI. 1-Amino-12-methyltridecane. Medelamine A*

[168706-09-0]

(H₃C)₂CH(CH₂)₁₀CH₂NH₂C₁₄H₃₁N 213.406Prod. by *Streptomyces* sp. NK14819.Cytotoxic agent. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O.▶ LD₅₀ (mus, ipr) 25-50 mg/kg.Morino, T. *et al.*, *J. Antibiot.*, 1995, **48**, 904 (isol, props)**2-Methyl-6-tridecylpiperidine, 9CI M-580**(2*R**,6*R**)-formC₁₉H₃₉N 281.524**(2*R**,6*R**)-form**trans-form. *Solenopsine B*

[32778-77-1]


Minor component of the venom of the red form of the fire ant (*Solenopsis saevissima*). Also present in the venom of *Solenopsis invicta*. Oil.*4',5'-Didehydro (Z)-: 2-Methyl-6-(4-tridecenyloxy)piperidine. Dehydrosolenopsin B*

[32778-76-0]

C₁₉H₃₇N 279.508Major component of venom of *Solenopsis saevissima*, also found in *Solenopsis xyloni* and *Solenopsis geminata*. Dehydrosolenopsin B is the 2,6-trans-form but the cis-form also occurs.**(2*R**,6*S**)-form**

cis-form


[35285-26-8]

Minor component of the venoms of both red and black forms of *Solenopsis saevissima*, *Solenopsis xyloni* and *Solenopsis geminata*. Trace component of the venom of *Solenopsis aurea*. Oil.MacConnell, J.G. *et al.*, *Tetrahedron*, 1971, **27**, 1129 (isol, synth, ms)Maruoka, K. *et al.*, *J.A.C.S.*, 1983, **105**, 2831 (synth)Taber, D.F. *et al.*, *J.O.C.*, 1988, **53**, 2968 (synth, ir, pmr, cmr, ms, *Solenopsin B*)Kotsuki, H. *et al.*, *Tet. Lett.*, 1991, **32**, 4159 (synth)**2-Methyl-4-(3,6,7-trimethoxy-2-naphthalenyl)-2-azabicyclo[2.2.1]heptan-3-one M-581**

Relative Configuration

C₂₀H₂₃NO₄ 341.406**(+)-form [288847-45-0]**Alkaloid from *Delphinium caeruleum*.Pan, Y.-J. *et al.*, *CA*, 2000, **133**, 190514f (isol)**N-Methyltryptamine M-582***N-Methyl-1H-indole-3-ethanamine, 9CI.**3-(2-Methylaminoethyl)indole. Dipterine*

[61-49-4]

C₁₁H₁₄N₂ 174.245Alkaloid from *Virola theiodora*, *Girgensohnia diptera*, *Acacia maidenii*, *Arthrophyllum leptocladum*, *Zanthoxylum arborescens* and others (Myristicaceae, Chenopodiaceae, Fabaceae, Rutaceae). Prisms (petrol). Mp 90° (87-88°). λ_{\max} 229 ; 275 ; 284 ; 292 (EtOH).**▶ NMO450000***Hydrochloride*: Mp 180°.*Picrate*:

Red needles. Mp 193-195° (190°).

N^b-Formyl: *N^b-Formyl-N^b-methyltryptamine*

[54268-27-8]

C₁₂H₁₄N₂O 202.255Isol. from *Acacia simplicifolia*, *Testulea gabonensis* and *Virola sebifera*. Oil; orange needles (as picrate). Mp 107.5-109.5° (picrate).*N^b-Ac*: *N^b-Acetyl-N^b-methyltryptamine*

[91821-04-4]

C₁₃H₁₆N₂O 216.282Alkaloid from stems of *Limonia acidissima* (wood apple) (Rutaceae). Amorph.*N^{tb}-(3-Methylbutanoyl)*: *Methylmadugin*

[190850-86-3]

C₁₆H₂₂N₂O 258.363Alkaloid from leaves of *Clausena indica*. Oil. λ_{\max} 224 ; 274 (sh) ; 283 ; 292 (Et₂O).*N^b-(2-Methylaminobenzoyl)*: *Evodia-mide*. *N-Methyl-N-(2-methylaminobenzoyl)tryptamine*

[116965-70-9]

C₁₉H₂₁N₃O 307.394Alkaloid from the fruits of *Evodia rutaecarpa*. Prisms (C₆H₆). Mp 208-209°. λ_{\max} 221 (log ϵ 4.65); 250 (log ϵ 4); 284 (log ϵ 3.84); 291 (log ϵ 3.82); 311 (log ϵ 3.44) (EtOH).**3-(Methylthio)propylamine, M-578 8CI***3-(Methylthio)propanamine, 9CI. 3-Methylmercaptopropylamine. 3-Amino-propyl methyl sulfide*

[4104-45-4]

H₂NCH₂CH₂CH₂SMeC₄H₁₁NS 105.204Constit. of roasted cocoa beans. Formed by *Streptomyces* from Methionine. Constit. of *Iberis amara* and from the brown alga *Desmarestia aculeata*. Steam-volatile liq. Misc. H₂O, EtOH, Et₂O. Bp 170° Bp₁₄ 66-67°.*Hydrochloride*: [2419-62-7]Hygroscopic needles (Me₂CO). Mp 144°.**▶ UI4100000***Picrate*:

Cryst. Mp 126-127°.

S-Oxide (R-): 3-(Methylsulfinyl)propylamine

[15813-61-3]

[3470-43-7]

C₄H₁₁NOS 121.203Found in *Iberis amara*. Cryst. Mp 27°. Bp_{0.04} 90-91°. $[\alpha]_D^{19}$ +117.8 (c, 1.04 in EtOH).*S,S-Dioxide: 3-(Methylsulfonyl)propylamine*

[26209-83-6]

C₄H₁₁NO₂S 137.202Hygroscopic cryst. Mp 44°. Bp₆ 165-168°.Schneider, W. *et al.*, *Annalen*, 1910, **375**, 207-254 (*S,S-dioxide, synth*)


N^b-Hydroxy: N^b-Hydroxy-N^b-methyl-tryptamine

[57383-99-0]

C₁₁H₁₄N₂O 190.244Alkaloid from the roots of *Desmanthus illinoensis* (Fabaceae). Prisms (C₆H₆). Mp 100-101°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 661A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 132A (nmr)

Yurashevskii, N.K. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1940, 10, 1781-1783; *CA*, 35, 4016 (isol, struct)Cohen, L.A. et al., *J.A.C.S.*, 1960, 82, 2184-2187 (synth, pmr)Fitzgerald, J.S. et al., *Aust. J. Chem.*, 1965, 18, 433-434 (isol, struct)Nakagawa, M. et al., *Tetrahedron*, 1974, 30, 2591-2600 (N-formyl)Nakagawa, M. et al., *J.A.C.S.*, 1975, 97, 6496-6501 (N-Hydroxy-N-methyltryptamine)Poupat, C. et al., *Phytochemistry*, 1976, 15, 2019-2020 (N-formyl)Grina, J.A. et al., *J.O.C.*, 1982, 47, 2648-2651 (isol, uv, pmr, ms)Zarga, M.H.A. et al., *J. Nat. Prod.*, 1986, 49, 901-904 (N^b-Acetyl-N^b-methyltryptamine)Thompson, A.C. et al., *J. Agric. Food Chem.*, 1987, 35, 361-365 (N-Hydroxy-N-methyltryptamine)Shoji, N. et al., *J. Nat. Prod.*, 1988, 51, 791-792 (Evodiamide)Riemer, B. et al., *Phytochemistry*, 1997, 45, 337-341 (Methylmadugin)**13a-Methyltylohirsutine****M-583***(S)*-formC₂₅H₂₉NO₄ 407.508**(S)-form****O⁴-De-Me: 14-Desoxy-13a-methyltylo-hirsutinidine**C₂₄H₂₇NO₄ 393.482Minor alkaloid from aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Light-yellow flakes (Me₂CO). Mp 180-182°. [α]_D¹⁸ +14.66 (c, 0.2 in MeOH).O⁴-De-Me, Ac: Mp 187-188°.**O⁴,O⁶-Di-de-Me: Tyloindicine C**

[126624-15-5]

C₂₃H₂₅NO₄ 379.455Alkaloid from the aerial parts of *Tylophora indica* (Asclepiadaceae). Brown beads (Me₂CO). Mp 223-225°. [α]_D²⁵ +16 (c, 0.25 in MeOH).O⁴,O⁶-Di-de-Me, di-Ac: Mp 176-177°.

14ξ-Hydroxy: Mp 200-201°.


14ξ-Hydroxy, O⁴-de-Me: 13a-Methyltylo-hirsutinidine

[95066-43-6]

C₂₄H₂₇NO₅ 409.481Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Flakes (Me₂CO). Mp 213-214°. [α]_D²⁵ +120 (c, 0.6 in CHCl₃).**(ξ)-form**Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Light-yellow flakes (Me₂CO). Mp 196-198°. Prob. racemic. λ_{max} 254 (log ε 4.6); 262 (log ε 4.4); 278 (log ε 3.4); 284 (log ε 3.1); 304 (log ε 2.6); 315 (log ε 2.5); 343 (log ε 1.9); 360 (log ε 1.8) (MeOH).Bhutani, K.K. et al., *Phytochemistry*, 1984, 23, 1765-1769 (13a-Methyltylohirsutine, 13a-Methyltylohirsutinidine)Ali, M. et al., *Phytochemistry*, 1987, 26, 2089-2092; 1989, 28, 3513-3517 (14-Desoxy-13a-methyltylohirsutinidine, Tyloindicine C)**2-Methyl-6-undecylpiperidine, 9CI****M-584**

[83709-88-0]

[35285-24-6, 76094-26-3, 92619-72-2]

C₁₇H₃₅N 253.47**(2R,6R)-form****trans-form. Solenopsine A**

[137038-57-4]

One of the major components of the venoms of fire ants spp. *Solenopsis xyloni*, *Solenopsis geminata*, *Solenopsis aurea*. Minor component of venoms of both red and black forms of *Solenopsis saevissima*. Also present in *Solenopsis invicta*. Oil. [α]_D²⁰ -1.3 (c, 1.3 in MeOH). Hydrochloride: Mp 147-150° (141-142°). [α]_D²⁰ -7.7 (c, 0.51 in CHCl₃).**1,2-Didehydro: 2,3,4,5-Tetrahydro-6-methyl-2-undecylpyridine. 1-Methyl-6-undecyl-Δ²-piperidine**C₁₇H₃₃N 251.454From *Solenopsis* spp.**(2R,6S)-form****cis-form. Isosolenopsin A**

[137038-58-5]

One of the major components of the venoms of *Solenopsis xyloni*, *Solenopsis geminata* and *Solenopsis aurea*. Minor component in venoms of both red and black forms of *Solenopsis saevissima*. Oil. [α]_D²⁵ -10.6 (c, 0.33 in CHCl₃) (as hydrochloride).**(2RS,6RS)-form***(±)*-trans-form

[28720-60-7]

[63950-17-4]

Oil. Mp 114° (as hydrochloride).

(2RS,6SR)-form*(±)*-cis-form

[63950-16-3]


Oil. Bp_{0.02} 91-96°.

Hydrochloride: [128561-86-4]

Cryst. (hexane/EtOH). Mp 155°.

MacConnell, J.G. et al., *Tetrahedron*, 1971, 27, 1129-1139 (isol, synth, ms, ir, pmr)Moriyama, Y. et al., *Tet. Lett.*, 1977, 825-828 (synth, ms, cmr)Maruoka, K. et al., *J.A.C.S.*, 1983, 105, 2831 (synth)Wasserman, H.H. et al., *Tet. Lett.*, 1988, 29, 4977 (synth)Nagasaka, T. et al., *Heterocycles*, 1989, 29, 2157-2166 (synth)Adams, D.R. et al., *J.C.S. Perkin 1*, 1989, 1507-1513 (synth)Comins, D.L. et al., *J.O.C.*, 1991, 56, 2506-2512 (synth)Jefford, C.W. et al., *Tet. Lett.*, 1993, 34, 2911-2914 (synth, nmr)Leclerq, S. et al., *Tetrahedron*, 1994, 50, 8465-8478 (synth, cmr, abs config)Chackalamannil, S. et al., *Tetrahedron*, 1997, 53, 11203-11210 (synth)Reding, M.T. et al., *J.O.C.*, 1998, 63, 6344-6347 (synth)Poerwono, H. et al., *Tetrahedron*, 1998, 54, 13955-13970 (synth)Amat, M. et al., *Tetrahedron: Asymmetry*, 1998, 9, 2419-2422 (synth)Wilkinson, T.J. et al., *Org. Lett.*, 2000, 2, 155-158 (synth)Wang, X. et al., *J.O.C.*, 2005, 70, 1897-1900 (synth)**4-Methyl-5-vinylthiazole, 8CI****M-585**

5-Ethenyl-4-methylthiazole. FEMA 3313 [1759-28-0]

C₆H₇NS 125.194Found in passion fruit and garlic. Meat flavouring ingredient. d₂₅²⁵ 1.09. Bp₂₅ 78-80° Bp₁₉ 78-82° (lit. gives a pressure range).▶ LD₅₀ (mus, ori) 400 mg/kg. XJ5104000**Picrate:**

Needles (AcOH). Mp 162-163°.


Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 643B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 111A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1493C (ir)

Buchman, E.R. et al., *J.A.C.S.*, 1945, 67, 395-399 (synth)Bachmann, G.B. et al., *J.A.C.S.*, 1949, 71, 1985-1988 (synth)Murav'eva, K.M. et al., *Zh. Obshch. Khim.*, 1963, 33, 3723-3725; *J. Gen. Chem. USSR (Engl. Transl.)*, 1963, 33, 3655-3657 (synth)Schilling, C.L. et al., *Macromolecules*, 1968, 1, 445-451 (pmr)*Fr. Pat.*, 1974, 2 329 309; *CA*, 82, 170885c (synth)Strauss, C.R. et al., *Chem. Ind. (London)*, 1978, 232-233 (occur)*Fenaroli's Handbook of Flavor Ingredients*, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 2, 583 (rev)*Encyclopedia of Food and Color Additives*, (ed. Burdock, G.A.), CRC Press, 1997, 1884 (rev)**1'-Methylzeatin****M-586**

2-Methyl-4-(1H-purin-6-ylamino)-2-penten-1-ol, 9CI. 6-(4-Hydroxy-1,3-dimethyl-2-butenylamino)purine



C₁₁H₁₅N₅O 233.272

(R)-form [101512-26-9]

Isol. from the culture filtrate of *Pseudomonas syringae* p.v. *savastanoi*. Sol. H₂O. Mp 201-202°. [α]_D²⁶ -109 (c, 0.153 in EtOH) (-52.6). λ_{\max} 270 (€ 11359) (EtOH) (Berdy). λ_{\max} 269 (€ 11842) (H₂O) (Berdy). λ_{\max} 272 (€ 17900); 275 (€ 11912) (pH 10 buffer) (Berdy). λ_{\max} 275 (€ 17400) (pH 1 buffer) (Berdy).

9- β -D-Ribofuranosyl: [98211-30-4]

C₁₆H₂₃N₅O₅ 365.388

Isol. from *Pseudomonas syringae* p.v. *savastanoi*. Cytokinin. Cryst. + 1/2 H₂O. Sol. MeOH. Mp 130-132°. [α]_D⁴ -117 (c, 0.102). λ_{\max} 265 (€ 19200) (pH 1 H₂O) (Derep). λ_{\max} 269 (€ 19400) (pH 13 H₂O) (Derep). λ_{\max} 270 (€ 18400) (95% EtOH) (Derep).

2-Hydroxy-2-Hydroxy-1'-methylzeatin

[137592-13-3]

C₁₁H₁₅N₅O₂ 249.272

Prod. by *Alternaria brassicae* and isol. from marine green algae (NIO-143). Shows cytokinin activity. Plant growth stimulator. Sol. MeOH. Mp 300°. [α]_D¹⁹ +41.6 (c, 0.288 in MeOH). Darkens at 275°. Probable abs. config. λ_{\max} 232 ; 242 ; 262 ; 282 (MeOH) (Berdy). λ_{\max} 256 ; 288 (MeOH/HCl) (Berdy). λ_{\max} 257 ; 288 (MeOH/NaOH) (Berdy).

(±)-form [127516-55-6]

Prisms (H₂O). Mp 175.5-176.5°.

Surico, G. *et al.*, *Phytochemistry*, 1985, **24**, 1499-1502 (isol, pmr, cmr, ms, uv)

Evidente, A. *et al.*, *Phytochemistry*, 1986, **25**, 525-526 (isol, pmr, cmr, uv, ms)

Itaya, T. *et al.*, *Tet. Lett.*, 1986, **27**, 6349 (synth, cd, abs config)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 3119-3121; 1990, **38**, 2702-2706 (synth, uv, pmr, cmr)

Ohba, M. *et al.*, *Heterocycles*, 1990, **31**, 599-602 (cmr)

Farooqi, A.H.A. *et al.*, *Phytochemistry*, 1990, **29**, 2061-2063 (isol, 2-hydroxy)


Dahiya, J.S. *et al.*, *Phytochemistry*, 1991, **30**, 2825-2828 (isol, 2-hydroxy)

Fujii, T. *et al.*, *Heterocycles*, 1992, **34**, 21-24 (2-hydroxy, synth, abs config)

Fujii, T. *et al.*, *Heterocycles*, 1997, **46**, 659-671 (rev)

Mevashuntin**M-587**

[848940-01-2]




C₂₂H₂₃NO₇S 445.492

Prod. by *Streptomyces prunicolor* 1884-SVT2 treated with Pravastatin. Orange powder. Mp 166-167°. [α]_D²¹ -415.5 (c, 0.03 in CHCl₃). λ_{\max} 245 (€ 22200); 264 (€

31100); 326 (€ 9300); 434 (€ 6500) (MeOH). λ_{\max} 242 (€ 33200); 290 (sh) (€ 9300); 326 (€ 9300); 420 (sh) (€ 2900); 565 (€ 6700) (MeOH/NaOH).

Shin-ya, K. *et al.*, *Tet. Lett.*, 2005, **46**, 1273-1276 (isol, uv, pmr, cmr)

Micacocidins**M-588**

C₂₇H₃₉N₃O₄S₃ 565.821

The Micacocidins are octahedral η^6 metallocomplexes of the organic ligand shown (in an anionic form). Antifungal agent.

Zn complex: **Micacocidin A. Antibiotic B057-250A. B057-250A**

[176950-36-0]

C₂₇H₃₇N₃O₄S₃Zn 629.195

Prod. by a *Pseudomonas* sp. No. 57-250. Active against *Mycoplasma* spp. Orange powder or rhombic cryst. (MeOH/EtOAc). Sol. MeOH, CH₂Cl₂, CHCl₃, EtOH; fairly sol. EtOAc; poorly sol. H₂O, butanol. Mp 226-228°. [α]_D²⁴ +53.2 (c, 1 in MeOH). λ_{\max} 238 (€ 19900); 264 (sh) (€ 12900); 336 (€ 5530) (MeOH). λ_{\max} 221 ; 265 (MeOH/HCl) (Berdy). λ_{\max} 275 ; 375 (MeOH/NaOH) (Berdy).

Cu complex: **Micacocidin B. Antibiotic B057-250B. B057-250B**

[176950-35-9]

C₂₇H₃₇CuN₃O₄S₃ 627.351

Prod. by *Pseudomonas* sp. No. 57-250. Active against *Mycoplasma* spp. Green cryst. or orange powder. Sol. MeOH, CHCl₃, CH₂Cl₂, EtOH, DMSO; fairly sol. EtOAc; poorly sol. H₂O, hexane, butanol. Mp 251-253°. [α]_D²⁴ -1369.2 (c, 0.1 in MeOH). λ_{\max} 223 (€ 22000); 275 (€ 14000); 375 (€ 2400) (MeOH). λ_{\max} 221 ; 265 (MeOH/HCl) (Berdy). λ_{\max} 275 ; 375 (MeOH/NaOH) (Berdy).

Fe complex: **Micacocidin C. Antibiotic B057-250C. B057-250C**

[176950-37-1]

C₂₇H₃₆FeN₃O₄S₃ 618.644

Prod. by *Pseudomonas* sp. No. 57-250. Active against *Mycoplasma* spp. Red-brown powder. Sol. MeOH, CHCl₃, EtOH, CH₂Cl₂, DMSO; fairly sol. EtOAc; poorly sol. H₂O, hexane, butanol. λ_{\max} 218 ; 239 (MeOH). λ_{\max} 221 ; 242 (MeOH/HCl) (Berdy). λ_{\max} 219 ; 239 (MeOH/NaOH) (Berdy).

Co complex: **Antibiotic B057-250D. B057-250D**

[176950-38-2]

C₂₇H₃₇CoN₃O₄S₃ 622.738

Prod. by a *Pseudomonas* sp. No. 57-250. Yellow powder. Sol. MeOH,

CHCl₃, EtOH, CH₂Cl₂; poorly sol. H₂O, hexane, butanol. Mp 251-253°. [α]_D²³ +28.4 (c, 0.95 in MeOH). λ_{\max} 202 (€ 22800); 237 (€ 25500); 265 (€ 18300); 337 (€ 5600) (MeOH). λ_{\max} 221 ; 265 (MeOH/HCl) (Berdy). λ_{\max} 275 ; 375 (MeOH/NaOH) (Berdy).

Ni complex: **Antibiotic B057-250E. B057-250E**

[176950-39-3]

C₂₇H₃₇N₃NiO₄S₃ 622.495

Prod. by a *Pseudomonas* sp. No. 57-250. Yellow powder. Sol. MeOH, EtOH, CHCl₃, CH₂Cl₂; poorly sol. H₂O, butanol, hexane. Mp 287-288°. [α]_D³ +197.6 (c, 0.49 in MeOH). λ_{\max} 202 (€ 21300); 235 (€ 24800); 265 (€ 15600); 370 (€ 4700) (MeOH). λ_{\max} 221 ; 265 (MeOH/HCl) (Berdy). λ_{\max} 275 ; 375 (MeOH/NaOH) (Berdy).


Pat. Coop. Treaty (WIPO), 1996, 96 4 262;

CA, **125**, 8683v

Kobayashi, S. *et al.*, *J. Antibiot.*, 1998, **51**, 323-327; 328-332 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Ino, A. *et al.*, *Tetrahedron*, 1999, **55**, 10271-10282; 10283-10294 (synth)

Kobayashi, S. *et al.*, *J. Antibiot.*, 2000, **53**, 532-539 (synth, activity)

Michellamine**M-589**

C₄₆H₄₈N₂O₈ 756.894

(1R,1''R,3R,3''R,5R,5''R)-form Michellamine C

[143168-23-4]

Alkaloid from *Ancistrocladus korupensis* (Ancistrocladaceae). Shows anti-HIV-cytopathic activity. [α]_D -16.8 (c, 0.14 in MeOH).

(1R,1''R,3R,3''R,5R,5''S)-form Michellamine B

[137893-48-2]

Alkaloid from *Ancistrocladus abbreviatus* and *Ancistrocladus korupensis* (Ancistrocladaceae). Shows anti-HIV-cytopathic activity. [α]_D -14.8 (c, 0.74 in MeOH). Stereoisomeric (atropisomeric) with Michellamine A at one of the biaryl functions. λ_{\max} 230 (€ 25118); 262 (€ 12590); 287 (€ 6310); 312 (€ 6310); 333 (€ 6310); 344 (€ 6310) (MeOH) (Berdy).

Diacetate salt: NSC 661755

[185310-27-4, 185310-28-5] Anti-HIV agent.

8'''-Me ether: Michellamine D

[191539-52-3]

C₄₇H₅₀N₂O₈ 770.921

Alkaloid from *Ancistrocladus korupensis* (Ancistrocladaceae). Amorph. solid. [α]_D -13.7 (c, 0.04 in MeOH) (as acetate salt). λ_{max} 207 (log ε 4.86); 230 (log ε 4.82); 264 (log ε 4.55); 331 (log ε 4.24); 344 (log ε 4.25) (MeOH).

1''',2'''-Didehydro, 8'''-Me ether: **Michellamine F**

[191539-54-5]

C₄₇H₄₈N₂O₈ 768.905

Alkaloid from *Ancistrocladus korupensis*. [α]_D +55 (c, 0.06 in MeOH). λ_{max} 205 (log ε 4.89); 233 (log ε 4.9); 238 (log ε 4.9); 261 (log ε 4.68); 331 (log ε 4.45); 347 (log ε 4.51) (MeOH).

(1R,1'''R,3R,3'''R,5S,5'''S)-form

Michellamine A

[137793-81-8]

Alkaloid from *Ancistrocladus abbreviatus* and *Ancistrocladus korupensis* (Ancistrocladaceae). Exhibits anti-HIV-cytopathic activity. [α]_D -10.5 (c, 0.38 in MeOH). λ_{max} 230 (ε 25000); 262 (ε 12500); 287 (ε 6300); 312 (ε 6300); 331 (ε 6300); 344 (ε 6300) (MeOH) (Berdy).

(1R,1'''R,3S,3'''R,5R,5'''S)-form

N²-Me: **Michellamine E**

[191539-53-4]

C₄₇H₅₀N₂O₈ 770.921

Alkaloid from *Ancistrocladus korupensis*. Light brown solid. [α]_D +33.6 (c, 0.07 in MeOH). λ_{max} 207 (log ε 4.84); 230 (log ε 4.8); 263 (log ε 4.54); 331 (log ε 4.22); 345 (log ε 4.24) (MeOH).

Manfredi, K.P. *et al.*, *J. Med. Chem.*, 1991, **34**, 3402 (*isol, uv, ir, pmr, cmr, struct*)

Boyd, M.R. *et al.*, *J. Med. Chem.*, 1994, **37**, 1740 (*Michellamine C, abs config*)

Kelly, T.R. *et al.*, *Tet. Lett.*, 1994, **35**, 7621 (*synth*)

Hoye, T.R. *et al.*, *Tet. Lett.*, 1994, **35**, 8747 (*synth*)

Bringmann, G. *et al.*, *Tetrahedron*, 1994, **50**, 7807 (*cd*)

Supko, J.G. *et al.*, *Antimicrob. Agents Chemother.*, 1995, **39**, 9-14 (*Michellamine B, pharmacol*)

Bringmann, G. *et al.*, *Annalen*, 1996, 2045-2058 (*synth*)

Hobbs, P.D. *et al.*, *Chem. Comm.*, 1996, 923 (*synth*)

Hallock, Y.F. *et al.*, *J. Nat. Prod.*, 1997, **60**, 677 (*isol, uv, ir, cd, pmr, cmr*)

Bringmann, G. *et al.*, *J.O.C.*, 1998, **63**, 1090-1097 (*synth*)

Hoye, T.R. *et al.*, *J.O.C.*, 1999, **64**, 7184-7201 (*synth*)


Bringmann, G. *et al.*, *Synthesis*, 2000, 1843-1847 (*Michellamine A, Michellamine C*)

Bringmann, G. *et al.*, *Tetrahedron*, 2003, **59**, 1245-1253 (*cd, conformn*)

Michigazone

M-590

8-(Hydroxymethyl)-2,4-dimethoxy-3H-phenoxazin-3-one, 9CI
[62267-63-4]



C₁₅H₁₃NO₅ 287.271

Isol. from *Streptomyces michiganensis*. Purple-red needles (MeOH). λ_{max} 226 (ε 28800); 261 (ε 13800); 384 (ε 20000); 460 (ε 6460) (MeOH) (Derep).

4-Demethoxy-8-(Hydroxymethyl)-2-methoxy-3H-phenoxazin-3-one, 9CI. **4-Demethoxymichigazone**

C₁₄H₁₁NO₄ 257.245

Prod. by *Streptomyces halstedii*. Neuronal cell protecting agent. Orange powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 193-194°. λ_{max} 222 (ε 17600); 251 (ε 10600); 390 (ε 11400); 444 (ε 5900) (MeOH) (Berdy).


Achenbach, H. *et al.*, *Chem. Ber.*, 1977, **110**, 12 (*pmr, synth*)

Kunigami, T. *et al.*, *J. Antibiot.*, 1996, **49**, 312 (*Demethoxymichigazone*)

Micranthine

M-591

[36104-64-0]



Absolute Configuration

C₃₄H₃₂N₂O₅ 548.637

Alkaloids covered by this entry belong to the enantiomeric series with respect to those in Isotrilobine, I-339 and are diastereomeric with those in Apateline, A-1331. Alkaloid from the bark of *Daphnandra micrantha* and from the leaves and terminal twigs of an unnamed *Daphnandra* sp. (Monimiaceae). Oedematous agent on injection, vasodilator, respiratory paralytic. Cryst. (MeOH). Mp 190-194° dec. [α]_D¹⁶ -221 (c, 1.3 in CHCl₃).

Me ether: **O-Methylmicranthine**

[40225-93-2]

C₃₅H₃₄N₂O₅ 562.664

Alkaloid from the bark of *Daphnandra micrantha* and *Daphnandra* sp. Dt-7, and from the leaves and terminal twigs of an unnamed *Daphnandra* sp. (Monimiaceae). Cryst. + ½ H₂O (C₆H₆). Mp 163-165° dec. [α]_D²⁰ -208 (CHCl₃).

Me ether, N-Ac:

Cryst. + 0.67 CCl₄ (CHCl₃/CCl₄). Mp 174-179° dec. [α]_D²² -203 (CHCl₃).

Me ether, N-Me: N,O-Dimethylmicranthine

[36296-04-5]

C₃₆H₃₆N₂O₅ 576.691

Alkaloid from the bark of *Daphnandra micrantha* and *Daphnandra* sp. Dt-7, and from the leaves and terminal twigs of an unnamed *Daphnandra* sp. (Monimiaceae). Cryst. + 0.25CHCl₃. Mp 210-214° dec. [α]_D¹⁹ -230 (CHCl₃).

Bick, I.R.C. *et al.*, *J.C.S. Perkin I*, 1972, 2884-2889 (*Micranthine, O-Methylmicranthine*)

Bick, I.R.C. *et al.*, *Tet. Lett.*, 1975, 2219-2220 (*N,O-Dimethylmicranthine*)


Bhakuni, D.S. *et al.*, *J.C.S. Perkin I*, 1978, 121-125 (*biosynth*)

Bick, I.R.C. *et al.*, *Heterocycles*, 1981, **16**, 2105-2108 (*Micranthine*)

Microcarpine[†]

M-592

[93552-71-7]



C₂₂H₂₃NO₇ 413.426

Alkaloid from *Fumaria microcarpa* (Papaveraceae).

Sener, B. *et al.*, *Int. J. Crude Drug Res.*, 1984, **22**, 45; *CA*, **102**, 3212f

Microcin B17

M-593

McB17

[84286-90-8]

Polypeptide antibiotic. Large depsipeptide containing 20 aminoacid residues interspersed with 8 heterocyclic residues (4 oxazole, 4 thiazole), including oxazole-thiazole bonds. Prod. by *Escherichia coli* LP 17. DNA gyrase inhibitor. Active against gram-negative bacteria.


Bayer, A. *et al.*, *Angew. Chem., Int. Ed.*, 1993, **32**, 1336-1339 (*struct*)

Videnov, G. *et al.*, *Angew. Chem., Int. Ed.*, 1996, **35**, 1506-1508 (*synth*)

Microcin SF608

M-594

[248582-51-6]



C₃₂H₄₄N₆O₆ 608.736

Isol. from *Microcystis* sp. Trypsin inhibitor. [α]_D²⁵ -19.1 (c, 1 in MeOH). λ_{max} 229 (ε 4360); 278 (ε 1040) (MeOH).

Banker, R. *et al.*, *Tetrahedron*, 1999, **55**, 10835-10844

Valls, N. *et al.*, *J.O.C.*, 2002, **67**, 4945-4950 (*synth*)

Micrococcin M

M-595

[57036-15-4]

[57036-14-3, 57036-16-5]


Thiazole-peptide antibiotic complex. Struct. unknown. Prod. by *Staphylococcus aureus*. Mp 221-224°. [α]_D +66.6 (EtOH). λ_{max} 216; 345 (EtOH).

Breiter, J. *et al.*, *Arzneim.-Forsch.*, 1975, **25**, 1244-1248

Hameister, W. *et al.*, *Arzneim.-Forsch.*, 1975, **25**, 1365-1369

Micrococcin P₁

[67401-56-3]



Absolute Configuration

C₄₈H₄₉N₁₃O₉S₆ 1144.393

Struct. revised in 1999. Isol. from *Bacillus pumilus*. Peptide antibiotic. Shows broad spectrum *in vitro* activity. Cryst. Sol. MeOH, Py, CHCl₃, EtOH, Me₂CO, EtOH; poorly sol. H₂O, C₆H₆, Et₂O, EtOAc, hexane. Mp 252° dec. [α]_D²¹ +116 (c, 0.10 in 90% EtOH aq.). Major component of Micrococcin P complex. Similar to Thiocillin I, T-383. λ_{max} 344 (MeOH) (Berdy). λ_{max} 346 (E1%/1cm 210) (EtOH) (Berdy). λ_{max} 320 (ε 57500) (HCl) (Berdy).

▶ LD₅₀ (mus, scu) 1500 mg/kg. PY4591000**17'-Ketone: Micrococcin P₂**

[67401-55-2]


C₄₈H₄₇N₁₃O₉S₆ 1142.377

From *Bacillus pumilus*. λ_{max} 345 (EtOH) (Berdy).

[1392-46-7]

Fuller, A.T. et al., *Nature (London)*, 1955, **175**, 722 (isol)Pestka, S. et al., *Antibiotics (N.Y.)*, 1975, **3**, 480-486 (rev)Walker, J. et al., *Chem. Comm.*, 1977, 706-708 (cmr)Bycroft, B.W. et al., *Chem. Comm.*, 1978, 256-258 (struct, pmr, cmr, biosynth)Kelly, T.R. et al., *Tet. Lett.*, 1991, **32**, 4263-4266 (synth)Shin, C.-G. et al., *Chem. Lett.*, 1998, 139-140 (synth)Okumura, K. et al., *Bull. Chem. Soc. Jpn.*, 1999, **72**, 1561-1569; 2483-2490 (synth, pmr, cmr)Ciufolini, M. et al., *Org. Lett.*, 1999, **1**, 1843-1846 (struct)Fenet, B. et al., *Tet. Lett.*, 2002, **43**, 2367-2370 (pmr, cmr, struct)Bagley, M.C. et al., *J. Antibiot.*, 2004, **57**, 829-831 (stereochem)Bagley, M.C. et al., *Chem. Rev.*, 2005, **105**, 685-714 (rev)**Microcyclamide**


[297730-40-6]




Absolute Configuration

C₂₆H₃₀N₈O₄S₂ 582.706

Abs. config. revised in 2002. Isol. from *Microcystis aeruginosa*. Moderate cytotoxic agent. Amorph. solid. [α]_D²³ -46.3 (c,

M-5960.1 in MeOH). λ_{max} 231 (ε 26500) (MeOH).Ishida, K. et al., *J. Nat. Prod.*, 2000, **63**, 1315-1317 (isol, pmr, cmr)Fujii, K. et al., *Tetrahedron*, 2002, **58**, 6873-6879 (isol, pmr, abs config)Ziemert, N. et al., *Appl. Environ. Microbiol.*, 2008, **74**, 1791-1797 (biosynth)**Micromide****M-598**C₄₉H₇₃N₇O₇S 904.224


Isol. from a *Symploca* sp. Cytotoxic. Amorph. powder. [α]_D²¹ -28 (c, 0.5 in MeOH). λ_{max} 201 (log ε 4.4); 225 (log ε 3.8); 274 (log ε 2.4) (MeOH).

Williams, P.G. et al., *J. Nat. Prod.*, 2004, **67**, 49-53 (isol, pmr, cmr)**Micromonosporin A****M-599**C₂₆H₃₇NO₄ 427.583

Prod. by *Micromonospora* sp. (strain TT1-11). Off-white powder. Mp 146° dec. [α]_D²⁵ -248.4 (c, 0.1 in MeOH). λ_{max} 281 (log ε 4.7); 291 (log ε 4.77); 306 (log ε 4.51); 322 (log ε 4.33) (MeOH).

Thawai, C. et al., *Chem. Biodiversity*, 2004, **1**, 640-645 (isol, biosynth, pmr, cmr)**Micropine****M-600**


6-(1,3,5-Decatrienyl)-3-hydroxy-1-methyl-2-piperidinemethanol, 9CI. 6-(1,3,5-Decatrienyl)-3-hydroxy-2-hydroxy-methyl-1-methylpiperidine [129011-71-8]



Absolute Configuration

C₁₇H₂₉NO₂ 279.422

Alkaloid from the leaves of *Microcos philippinensis*. Off-white plates (MeOH). Mp 146-148°. [α]_D²⁰ -63 (c, 0.15 in EtOH). λ_{max} 250 (ε 24000); 258 (ε 39000); 268 (ε 47000); 280 (ε 414000); 299 (ε 4200); 315 (ε 3300) (EtOH).

Aguinaldo, A.M. et al., *Phytochemistry*, 1990, **29**, 2309-2313 (isol, uv, ir, pmr, cmr, ms)Bayquen, A.V. et al., *Tetrahedron*, 1996, **52**, 13467-13482 (synth, abs config)**Microsphaerone B****M-601**C₁₇H₂₁NO₅ 319.357


Prod. by the fungus *Microsphaeropsis* sp. isol. from the sponge *Aplysina aerophoba*. Yellow powder (MeOH). λ_{max} 211 (log ε 3.65); 231 (log ε 3.63); 313 (log ε 3.71) (MeOH).

N-(3S-Carboxybutanoyl): Microsphaerone AC₂₂H₂₇NO₈ 433.457

Prod. by a *Microsphaeropsis* sp. from *Aplysina aerophoba*. Yellow powder (MeOH). [α]_D²⁰ -8.9 (c, 0.56 in MeOH). λ_{max} 229 (log ε 3.77); 312 (log ε 3.63) (MeOH).

Wang, C.-Y. et al., *J. Nat. Prod.*, 2002, **65**, 772-775 (isol, pmr, cmr, ms)**Microsporin B****M-602**

[945491-39-4]



Absolute Configuration

C₂₈H₄₂N₄O₅ 514.664


Prod. by the marine-derived fungus *Microsporium* cf. *gypseum*. Histone deacetylase inhibitor. Oil. [α]_D -39.8 (c, 0.44 in CH₂Cl₂). λ_{max} 242 (ε 13260) (CHCl₃).

6'-Ketone: Microsporin A

[945491-38-3]

C₂₈H₄₀N₄O₅ 512.648

Prod. by *Microsporium* cf. *gypseum*. Oil. [α]_D +11.6 (c, 0.17 in CH₂Cl₂). λ_{max} 242 (ε 1670); 270 (ε 3510) (CHCl₃).


Gu, W. et al., *Tetrahedron*, 2007, **63**, 6535-6541 (isol, synth, pmr, cmr)**Microxine****M-603**C₈H₁₁N₅O₄S 273.272

Isol. from the sponge *Microxina* sp. Weak cdc2 kinase inhibitor. Amorph. solid. λ_{max} 210 (ε 12890); 274 (ε 8410) (H₂O).

Killday, K.B. et al., *J. Nat. Prod.*, 2001, **64**, 525-526 (isol)

Midpacamide **M-604**

4,5-Dibromo-1-methyl-N-[3-(1-methyl-2,5-dioxo-4-imidazolidinyl)propyl]-1H-pyrrole-2-carboxamide, 9CI
[66067-05-8]



$C_{13}H_{16}Br_2N_4O_3$ 436.102

Isol. from an unidentified marine sponge and from the sponge *Agelas mauritiana*. Cryst. (MeCN). Mp 93-95°. λ_{max} 206 (€ 5010); 239 (€ 7940); 276 (€ 11200) (MeOH) (Derep).

5-Debromo- 5-Debromomidpacamide

[114216-89-6]

$C_{13}H_{17}BrN_4O_3$ 357.206


Isol. from the marine sponge *Agelas mauritiana*. Mp 116-119°.

Chevolot, L. et al., *Heterocycles*, 1977, 7, 891 (isol, pmr, cmr, struct)

Fathi-Afshar, R. et al., *Can. J. Chem.*, 1988, 66, 45 (isol, ir, pmr, cmr, ms, deriv)

Lindel, T. et al., *Liebigs Ann./Recl.*, 1997, 1525-1528 (synth, pmr, cmr, ms)


Fresneda, P.M. et al., *Tet. Lett.*, 2001, 42, 851-854 (synth)

Miersine**M-605**

$C_{20}H_{27}NO_6$ 377.436

Minor alkaloid from *Stephania japonica* (Menispermaceae). Mp 222°.

Thorner, C.W. et al., *Phytochemistry*, 1970, 9, 157 (ir, uv, pmr, struct)

Migrastatin**M-606**

Absolute Configuration

$C_{27}H_{39}NO_7$ 489.608


Prod. by *Streptomyces* sp. MK929-43F1. Strictly a shunt metab. of Isomigrastatin, I-245. Inhibitor of tumour cell migration. Powder. Mp 54-55°. $[\alpha]_D^{27} +17.9$ (c, 3.2 in MeOH). Related to Antibiotic BR 040, A-1139.

Nakae, K. et al., *J. Antibiot.*, 2000, 53, 1130-

1136; 1228-1230
Takemoto, Y. et al., *J. Antibiot.*, 2001, 54, 1104-1107 (activity)
Nakamura, H. et al., *J. Antibiot.*, 2002, 55, 442-444 (abs config)
Gaul, C. et al., *J.A.C.S.*, 2003, 125, 6042-6043 (synth)
Ju, J. et al., *J.A.C.S.*, 2005, 127, 1622-1623 (biosynth)
Ju, J. et al., *Org. Lett.*, 2006, 8, 5865-5868 (synth)
Reymond, S. et al., *Tetrahedron*, 2007, 63, 5918-5929 (synth)

Milbemycin α 1**M-607**

B 41A₃. Antibiotic B 41A₃. Milbemycin A₃. Milbectin
[51596-10-2]



$R^1 = R^2 = CH_3$

$C_{31}H_{44}O_7$ 528.684

Macrolide antibiotic. Isol. from *Streptomyces hygroscopicus*. Anthelmintic agent. Cryst. (hexane). Mp 193-195°. $[\alpha]_D^{20} +106$ (c, 0.25 in Me₂CO). λ_{max} 238 (sh) (€ 27800); 244 (€ 30500); 253 (sh) (€ 22000) (EtOH) (Derep).

5-Me ether: Milbemycin α 2. B 41B₂.

Antibiotic B 41B₂. Milbemycin B₂

[51596-12-4]

$C_{32}H_{46}O_7$ 542.711

Prod. by *Streptomyces hygroscopicus*. Anthelmintic agent. Mp 139-142°. $[\alpha]_D^{20} +131$ (c, 0.25 in Me₂CO). λ_{max} 238 (sh) (€ 27800); 244 (€ 30500); 253 (sh) (€ 22000) (EtOH).

5-Ketone: Milbemycin J

[86691-98-7]

$C_{31}H_{42}O_7$ 526.669

Prod. by a *Streptomyces hygroscopicus* ssp. *aureolacrimosus* mutant. Shows insecticidal and acaricidal props. Mp 213-215°. $[\alpha]_D^{27} +40$ (c, 0.25 in Me₂CO). λ_{max} 240 (€ 28000) (EtOH).

23S-Hydroxy, 5-Me ether: 23-Hydroxy-milbemycin α 2

[106313-99-9]

$C_{32}H_{46}O_8$ 558.711

Prod. by *Streptomyces* sp. NCIB 11867. Anthelmintic, pesticide. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O. λ_{max} 239; 244 (MeOH) (Berdy).

26-Hydroxy: Milbemycin α 26. 26-Hydroxymilbemycin α 1

[86629-93-8]

$C_{31}H_{44}O_8$ 544.684

Prod. by *Streptomyces* sp. KSB 1939 and *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder. Sol. MeOH, Me₂CO; poorly sol. H₂O. λ_{max} 244 (MeOH).

26-Hydroxy, 5-Me ether: Milbemycin α 24. 26-Hydroxymilbemycin α 2

[133656-16-3]

$C_{32}H_{46}O_8$ 558.711

Prod. by *Streptomyces* sp. KSB 1939 and *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Insecticide, acaricide. Amorph. powder. Sol. MeOH, Me₂CO; poorly sol. H₂O. λ_{max} 244 (MeOH).

26-(Propanoyloxy): Milbemycin α 22

$C_{34}H_{48}O_9$ 600.748

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder.

26-(3-Methylbutanoyloxy): Milbemycin α 12

[123071-17-0]

$C_{36}H_{52}O_9$ 628.801

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. $[\alpha]_D^{25} +118.3$ (c, 1 in CHCl₃). Stereochem. not explicitly given but by implication is the same as for other milbemycins. λ_{max} 238 (€ 750); 244 (€ 810); 253 (sh) (EtOH).

26-(Tigloyloxy): Milbemycin α 20

$C_{36}H_{50}O_9$ 626.786

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder.

26-(3-Methyl-2-butenoyloxy): Milbemycin α 11. MI 198Z1. KSB 1939H3.

Antibiotic KSB 1939H3. Antibiotic MI 198Z1

[119053-92-8]

$C_{36}H_{50}O_9$ 626.786

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. $[\alpha]_D^{25} +104.3$ (c, 1.05 in CHCl₃). Stereochem. not explicitly given but by implication is the same as for other milbemycins. λ_{max} 230 (sh); 238 (€ 990); 244 (€ 990); 252 (sh) (EtOH).

26-(3-Methyl-2-pentenoyloxy): Milbemycin α 13. MI 19827. KSB 193944.

Antibiotic KSB 193944. Antibiotic MI 19827

[123123-62-6]

$C_{37}H_{52}O_9$ 640.812

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. $[\alpha]_D^{25} +91.6$ (c, 0.89 in CHCl₃). Stereochem. not explicitly given but by implication is the same as for other milbemycins.

26-(2-Pyrrolylcarbonyloxy): Milbemycin α 9. B 41C₁. Antibiotic B 41C₁. Milbemycin C₁

[51596-14-6]

$C_{36}H_{47}NO_9$ 637.769

Prod. by *Streptomyces hygroscopicus*. Anthelmintic. $[\alpha]_D^{20} +57$ (c, 0.25 in Me₂CO). λ_{max} 238 (€ 31300); 245 (€ 35000); 253 (€ 31000); 266 (sh) (€ 15000) (EtOH).

Ger. Pat., 1973, 2 329 486; CA, 80, 131682

(synth)

Japan. Pat., 1975, 75 29 742; CA, 83, 127530

(synth)

U.S. Pat., 1976, 3 992 551; CA, 86, 70107

(synth)

U.S. Pat., 1977, 4 144 352; CA, 90, 210144

(props)

Takiguchi, Y. et al., *J. Antibiot.*, 1980, 33,

1120-1127 (Milbemycins α 2,B2, isol, uv, ir, ms)

- Ono, M. *et al.*, *J. Antibiot.*, 1983, **36**, 438-441; 509-515; 980-990 (*Milbemycins* α 2, J)
U.K. Pat., 1986, 2 170 499; *CA*, **106**, 133823z
 (23-Hydroxymilbemycin α 2)
Japan. Pat., 1991, 91 17 082; *CA*, **114**, 245968x
 (26-Hydroxymilbemycins)
 Takahashi, S. *et al.*, *J. Antibiot.*, 1993, **46**,
 1364-1371 (*Milbemycins* α 11, α 12, α 13)
 Ley, S.V. *et al.*, *Tet. Lett.*, 1993, **34**, 7479-7482
 (*synth*)
 Nonaka, K. *et al.*, *J. Antibiot.*, 2000, **53**, 694-
 704 (*Milbemycins* α 20, α 22, α 24, α 26)
 Tsukiyama, T. *et al.*, *J. Antibiot.*, 2002, **55**, 993-
 1003 (*synth*)

Milbemycin α 3 **M-608**

B 41A₄. Antibiotic B 41A₄. Milbemycin A₄
 [51596-11-3]
 As Milbemycin α 1, M-607 with
 $R^1 = \text{CH}_2\text{CH}_3$, $R^2 = \text{CH}_3$
 $\text{C}_{32}\text{H}_{46}\text{O}_7$ 542.711
 Macrolide antibiotic. Isol. from *Streptomyces hygroscopicus*. Anthelmintic agent. Cryst. Sol. MeOH, hexane; poorly sol. H_2O . Mp 212-215°. $[\alpha]_{\text{D}}^{20} +106$ (c, 0.25 in Me_2CO). λ_{max} 238 (sh) (ϵ 27800); 244 (ϵ 30500); 253 (sh) (ϵ 22000) (EtOH) (Derep).

5-Me ether: Milbemycin α 4. B 41B₃. Antibiotic B 41B₃. Milbemycin B₃
 [51596-13-5]
 $\text{C}_{33}\text{H}_{48}\text{O}_7$ 556.738
 Prod. by *Streptomyces hygroscopicus*. Anthelmintic agent. Amorph. powder. $[\alpha]_{\text{D}}^{20} +126$ (c, 0.25 in Me_2CO). λ_{max} 238 (sh) (ϵ 27800); 244 (ϵ 30500); 253 (sh) (ϵ 22000) (EtOH).

5-Ketone: Milbemycin K
 [86691-97-6]
 $\text{C}_{32}\text{H}_{44}\text{O}_7$ 540.695
 Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus* mutant. Shows insecticidal and acaricidal props. Cryst. (hexane) or amorph. Mp 215-218°. $[\alpha]_{\text{D}}^{25} +42$ (c, 0.25 in Me_2CO). λ_{max} 240 (ϵ 28000) (EtOH).

23S-Hydroxy: Milbemycin α 19. 23-Hydroxymilbemycin α 3
 [122006-28-4]
 $\text{C}_{32}\text{H}_{46}\text{O}_8$ 558.711
 Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Antiparasitic agent, miticide. Sol. MeOH, Me_2CO , MeCN; fairly sol. hexane; poorly sol. H_2O . λ_{max} 244 (ϵ 27500) (EtOH) (Berdy).

23S-Hydroxy, 5-ketone: Milbemycin α 18
 [122006-27-3]
 $\text{C}_{32}\text{H}_{44}\text{O}_8$ 556.695
 Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Antiparasitic agent, miticide. Powder. Sol. MeOH, Me_2CO , MeCN; fairly sol. hexane; poorly sol. H_2O . λ_{max} 240 (ϵ 29500) (EtOH). λ_{max} 240 (ϵ 29500) (EtOH) (Berdy).

26-Hydroxy: Milbemycin α 27. 26-Hydroxymilbemycin α 3
 [86629-94-9]
 $\text{C}_{32}\text{H}_{46}\text{O}_8$ 558.711
 Prod. by *Streptomyces* sp. KSB 1939 and *Streptomyces hygroscopicus* ssp.

aureolacrimosus. Insecticide, acaricide. Amorph. powder. Sol. MeOH, Me_2CO ; poorly sol. H_2O . λ_{max} 244 (MeOH) (Berdy).

26-Hydroxy, 5-Me ether: Milbemycin α 25
 [205525-19-5]
 $\text{C}_{33}\text{H}_{48}\text{O}_8$ 572.737

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder.

26-(Propanoyloxy): Milbemycin α 23
 [205525-18-4]
 $\text{C}_{35}\text{H}_{50}\text{O}_9$ 614.775

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder.

26-(3-Methylbutanoyloxy): Milbemycin α 15
 [125602-42-8]
 $\text{C}_{37}\text{H}_{54}\text{O}_9$ 642.828

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. λ_{max} 230 (sh); 238 (sh); 244 (990); 252 (sh) (EtOH).

26-(Tigloyloxy): Milbemycin α 21
 [119053-97-3]
 $\text{C}_{37}\text{H}_{52}\text{O}_9$ 640.812

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. Amorph. powder.

26-(3-Methyl-2-butenoyloxy): Milbemycin α 14
 [119053-94-0]
 $\text{C}_{37}\text{H}_{52}\text{O}_9$ 640.812

Prod. by *Streptomyces hygroscopicus* ssp. *aureolacrimosus*. λ_{max} 230 (sh); 238 (ϵ 990); 244 (ϵ 990); 252 (sh) (EtOH).

26-(2-Pyrrolylcarbonyloxy): Milbemycin α 10. BC 41C₂. Antibiotic BC 41C₂.

Milbemycin C₂
 [51596-15-7]
 $\text{C}_{37}\text{H}_{49}\text{NO}_9$ 651.795

Prod. by *Streptomyces hygroscopicus*. Anthelmintic. Amorph. powder. $[\alpha]_{\text{D}}^{20} +54$ (c, 0.25 in Me_2CO). λ_{max} 238 (ϵ 31300); 245 (ϵ 35000); 266 (sh) (ϵ 15000) (EtOH).

Ger. Pat., 1973, 2 329 486; *CA*, **80**, 131682 (*synth*)

Japan. Pat., 1975, 75 29 742; *CA*, **83**, 127530 (*synth*)

U.S. Pat., 1976, 3 992 551; *CA*, **86**, 70107 (*synth*)

U.S. Pat., 1977, 4 144 352; *CA*, **90**, 210144 (*props*)

Takiguchi, Y. *et al.*, *J. Antibiot.*, 1980, **33**, 1120-1127 (*Milbemycin α 4, isol, uv, ir, ms*)

Ono, M. *et al.*, *J. Antibiot.*, 1983, **36**, 438-441; 509-515 (*Milbemycins* α 4, K , *isol*)

Japan. Pat., 1988, 88 227 590; *CA*, **111**, 76530d (*Milbemycins* α 18, α 19)

Japan. Pat., 1991, 91 17 082; *CA*, **114**, 245968x (*26-Hydroxymilbemycin α 3*)

Takahashi, S. *et al.*, *J. Antibiot.*, 1993, **46**, 1364-1371 (*Milbemycins* α 14, α 15)

Nonaka, K. *et al.*, *J. Antibiot.*, 1999, **52**, 109-116 (*Milbemycin α 14, biosynth*)

Nonaka, K. *et al.*, *J. Antibiot.*, 2000, **53**, 694-704 (*Milbemycin α 21, α 23, α 25, α 27*)

Tsukiyama, T. *et al.*, *J. Antibiot.*, 2002, **55**, 993-1003 (*synth*)

Milbemycin D **M-609**

B 41D. Antibiotic B 41D
 [77855-81-3]

As Milbemycin α 1, M-607 with
 $R^1 = -\text{CH}(\text{CH}_3)_2$, $R^2 = \text{CH}_3$

$\text{C}_{33}\text{H}_{48}\text{O}_7$ 556.738

Macrolide antibiotic. Isol. from *Streptomyces hygroscopicus*. Antiparasitic and pesticidal agent. Needles (hexane/EtOAc). Sol. MeOH, EtOAc; fairly sol. hexane; poorly sol. H_2O . Mp 186-188°. $[\alpha]_{\text{D}}^{27} +107$ (c, 0.25 in Me_2CO). λ_{max} 238 (sh) (ϵ 27800); 244 (ϵ 30500); 253 (sh) (ϵ 22000) (EtOH) (Derep). λ_{max} 243 (ϵ 30500); 244 (ϵ 31000) (EtOH) (Berdy).

▶PY5788000

O⁵-Me: Milbemycin G. B 41G. Antibiotic B 41G

[83471-31-2]

$\text{C}_{34}\text{H}_{50}\text{O}_7$ 570.765

Prod. by *Streptomyces hygroscopicus*. Antiparasitic and pesticidal agent.

Amorph. $[\alpha]_{\text{D}}^{27} +108$ (c, 0.25 in Me_2CO). λ_{max} 238 (sh) (ϵ 27800); 244 (ϵ 30500); 253 (sh) (ϵ 22000) (EtOH).

26-O-(2-Pyrrolylcarbonyloxy): Milbemycin F. B 41F. Antibiotic B 41F

[83177-49-5]

$\text{C}_{38}\text{H}_{51}\text{NO}_9$ 665.822

Prod. by *Streptomyces hygroscopicus*. Antiparasitic and pesticidal agent.

Amorph. λ_{max} 238 (ϵ 31300); 253 (ϵ 31000); 254 (ϵ 35000); 266 (sh) (ϵ 15000) (EtOH).

Takeguchi, Y. *et al.*, *J. Antibiot.*, 1983, **36**, 438; 502; 980; 991 (*isol, struct, uv, ir, cmr*)


Ono, M. *et al.*, *J. Antibiot.*, 1983, **36**, 438; 502; 980 (*Milbemycin F, Milbemycin G*)

Mrozik, H. *et al.*, *Tet. Lett.*, 1983, 5333 (*synth, uv, pmr*)

Crimmins, M.T. *et al.*, *J.A.C.S.*, 1996, **118**, 7513 (*synth*)

Bailey, S. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 3654-3677 (*Milbemycin G, synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MQT600

Militarinone A**M-610**

$\text{C}_{26}\text{H}_{37}\text{NO}_6$ 459.581

Isol. from *Paecilomyces militaris* RCEF 0095. Neurotrophic agent. Yellow solid. $[\alpha]_{\text{D}}^{25} -14.6$ (c, 1.3 in MeOH). λ_{max} 261 (log ϵ 3.98); 388 (log ϵ 4.65) (MeOH).

N-Deoxy: N-Deoxymilitarinone A

$\text{C}_{26}\text{H}_{37}\text{NO}_5$ 443.582


Isol. from *Paecilomyces farinosus* RCEF 0097. Amorph. yellow solid.

$[\alpha]_{\text{D}}^{24} +33.3$ (c, 1.8 in MeOH). λ_{max} 253 (log ϵ 4.02); 391 (log ϵ 4.46) (MeOH).

Schmidt, K. *et al.*, *Org. Lett.*, 2002, **4**, 197-199 (*isol, pmr, cmr*)

Schmidt, K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 378-383 (*biosynth*)

Cheng, Y. *et al.*, *J. Nat. Prod.*, 2006, **69**, 436-438 (*N-Deoxymilitarinone A*)


Militarinone B
M-611

 $C_{26}H_{33}NO_5$ 439.55

Tetramic acid deriv. Isol. from *Paecilomyces militaris* RCEF 0095. Amorph. yellow powder. Mp 135-139°. $[\alpha]_D^{25}$ -553.2 (c, 0.19 in MeOH). Occurs in soln. as equilibrium mixture with (7Z)-isomer. λ_{max} 259 (log ϵ 4.14); 358 (log ϵ 4.53) (MeOH).

6-Deoxy- Milliamine C
 $C_{26}H_{33}NO_4$ 423.551

Isol. from *Paecilomyces militaris* RCEF 0095. Amorph. yellow powder. Mp 86-92°. $[\alpha]_D^{25}$ -430.2 (c, 0.17 in MeOH). Occurs in soln. as equilibrium mixture with (7Z)-isomer. λ_{max} 258 (log ϵ 4.14); 359 (log ϵ 4.57) (MeOH).

Schmidt, K. et al., *J. Nat. Prod.*, 2003, **66**, 378-383 (isol, uv, pmr, cmr, ms)


Militarinone D
M-612

 $C_{26}H_{31}NO_4$ 421.535

Isol. from *Paecilomyces militaris* RCEF 0095. Cytotoxic. Amorph. yellow powder. Mp 132-138°. $[\alpha]_D^{25}$ -74.4 (c, 0.24 in MeOH). λ_{max} 253 (log ϵ 4.43); 384 (log ϵ 4.55) (MeOH).

Schmidt, K. et al., *J. Nat. Prod.*, 2003, **66**, 378-383 (isol, uv, pmr, cmr, ms)

Millaurine†
M-613

[152606-57-0]


 $R = CH_3, R' = OH$
 $C_{14}H_{17}N_3O_2$ 259.307

Rel. configs. only detd. Alkaloid from the seeds of *Milletia laurentii*. Yellow needles (MeOH). Mp 248-249°. $[\alpha]_D^{21}$ +20.6 (c, 0.18 in MeOH).

8-Ac- O-Acetylmillaurine

[152307-09-0]

 $C_{16}H_{19}N_3O_3$ 301.344

Alkaloid from seeds of *Milletia laurentii*. Yellow needles (EtOH). Mp 218-219°. $[\alpha]_D^{21}$ +51.3 (c, 0.34 in MeOH).

9a β -Hydroxy, 5a β ,9a-dihydro: **5a,9a-Di-**

hydro-5a-hydroxymillaurine

[158150-01-7]

 $C_{14}H_{19}N_3O_3$ 277.322

Alkaloid from seeds of *Milletia laurentii*. Needles (EtOH). Mp 259-260°. $[\alpha]_D^{21}$ +6.8 (c, 0.2 in MeOH).

Ngamga, D. et al., *J. Nat. Prod.*, 1993, **56**, 2126-2132; 1994, **57**, 1022-1024 (isol, ir, pmr, cmr, ms, cryst struct)

Millaurine A†
M-614

[946051-22-5]

As Millaurine, M-613 with


 $R = H, R' = OMe$
 $C_{14}H_{17}N_3O_2$ 259.307

Alkaloid from the seeds of *Milletia laurentii*. Needles. Mp 136-137°. $[\alpha]_D$ +46.7 (c, 0.2 in MeOH).

Ngamga, D. et al., *Fitoterapia*, 2007, **78**, 276-277 (isol, pmr, cmr, ms)

Millettonine
M-615

[159509-32-7]



 $C_{14}H_{17}N_3O_3$ 275.307

Alkaloid from the stem bark of *Milletia laurentii* (Fabaceae). Needles (hexane). Mp 170°. $[\alpha]_D^{21}$ +46.2 (c, 0.71 in MeOH).

Kamnaing, P. et al., *Phytochemistry*, 1994, **36**, 1561 (isol, ir, pmr, cmr, ms, struct)

Milliamine C
M-616

[49620-09-9]


 $C_{43}H_{47}N_3O_9$ 749.859

Alkaloid from *Euphorbia milii* (Euphorbiaceae). Amorph. solid. $[\alpha]_D^{22}$ +11 (c, 1 in $CHCl_3$). λ_{max} 220 (c 48170); 260 (c 19130); 312 (c 14060) (MeOH).

5-Ac: Milliamine E

[88524-60-1]

 $C_{45}H_{49}N_3O_{10}$ 791.896

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 135-138°. $[\alpha]_D$ +34.6 (c, 0.24 in $CHCl_3$). λ_{max} 226 (c 45200); 258 (c 19200); 309 (c 15820) (MeOH).

20-Ac: Milliamine A

[34391-10-1]

 $C_{45}H_{49}N_3O_{10}$ 791.896

Main alkaloid from *Euphorbia milii* (Euphorbiaceae). Mp 134-136°. $[\alpha]_D$ +11.6 (c, 1 in $CHCl_3$). CAS no. refers to the hydrochloride. λ_{max} 232 (c 37300); 260 (c 21900); 315 (c 14100); 342 (c 4800) (MeOH) (Berdy). λ_{max} 219 (c 44930); 260 (c 17640); 314 (c 14110) (EtOH) (Berdy). λ_{max} 225 (c 35590); 260 (c 15820); 308 (c 13450) (MeOH).

► DG8736000
20-Ac, N-de-Me: Milliamine J

[152135-64-3]

 $C_{44}H_{47}N_3O_{10}$ 777.869

Alkaloid from *Euphorbia milii*.

Amorph. yellow powder. Mp 112-116°. $[\alpha]_D$ +9.1 (c, 0.3 in $CHCl_3$). λ_{max} 223 (c 44130); 256 (c 19900); 310 (c 18800); 365 (c 8080) (MeOH).

Nⁱ-Deacyl, 20-Ac: Milliamine L

[152135-61-0]

 $C_{36}H_{40}N_2O_9$ 644.72

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 108-111°. $[\alpha]_D$ +3 (c, 1.4 in $CHCl_3$). λ_{max} 224 (c 50500); 264 (c 17870); 347 (c 14700) (MeOH).

Nⁱ-Deacyl, Nⁱ-tigloyl, 20-Ac: Milliamine K

[153133-63-2]

 $C_{41}H_{46}N_2O_{10}$ 726.822

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 120-123°. λ_{max} 221 (c 54900); 266 (c 23800); 313 (c 21000) (MeOH).

De(dimethylamino): Milliamine G

[88524-63-4]

 $C_{41}H_{42}N_2O_9$ 706.791

Alkaloid from *Euphorbia milii*.

Amorph.

3-O-Deacyl-5-O-acyl isomer, 20-Ac:
Milliamine D

[88524-59-8]

 $C_{45}H_{49}N_3O_{10}$ 791.896

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 136-139°. $[\alpha]_D$ -15 (c, 0.24 in $CHCl_3$). λ_{max} 228 (c 46800); 260 (c 23730); 315 (c 18270) (MeOH).

3-O-Deacyl-5-O-acyl isomer, Nⁱ-deacyl,
20-Ac: Milliamine M

[152135-62-1]

 $C_{36}H_{40}N_2O_9$ 644.72

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 112-115°. $[\alpha]_D$ +12.6 (c, 1 in $CHCl_3$). λ_{max} 224 (c 50500); 264 (c 19400); 348 (c 19420) (MeOH).

3-O-Deacyl-5-O-acyl isomer, Nⁱ-deacyl,
3,20-di-Ac: Milliamine N

[152135-63-2]

 $C_{38}H_{42}N_2O_{10}$ 686.757

Alkaloid from *Euphorbia milii*.

Amorph. powder. Mp 116-120°. $[\alpha]_D^{25}$ -25.2 (c, 1 in CHCl_3). λ_{max} 225 (ε 42880); 265 (ε 15350); 349 (ε 13050) (MeOH).

3-O-Deacyl-5-O-acyl isomer, de(dimethylamino), 20-Ac: **Milliamine F** [88524-61-2]

$\text{C}_{43}\text{H}_{44}\text{N}_2\text{O}_{10}$ 748.828

Alkaloid from *Euphorbia milii*. Amorph.

3-O-Deacyl-20-O-acyl isomer: **Milliamine B**

[34420-84-3]

$\text{C}_{43}\text{H}_{47}\text{N}_3\text{O}_9$ 749.859

Alkaloid from *Euphorbia milii* (Euphorbiaceae). λ_{max} 227 (ε 43800); 260 (ε 18800); 315 (ε 14600); 342 (ε 5000) (MeOH) (Berdy). λ_{max} 217 (ε 34480); 260 (ε 12950); 312 (ε 10940) (EtOH) (Berdy).


Uemura, D. et al., *Bull. Chem. Soc. Jpn.*, 1977, **50**, 2005-2009 (isol, uv, ir, pmr, struct)

Marston, A. et al., *Planta Med.*, 1983, **47**, 141-147; 1984, **48**, 319-322 (isol, uv, ir, pmr, ms)

Zani, C.L. et al., *Phytochemistry*, 1993, **34**, 89-95 (isol, uv, ir, pmr, cmr, ms)

Milliamine H M-617

[91197-53-4]



$\text{C}_{37}\text{H}_{42}\text{N}_2\text{O}_9$ 658.747

Alkaloid from *Euphorbia milii*. Pale yellow glass. λ_{max} 212 (ε 47300); 228 (ε 53460); 257 (ε 19530); 325 (ε 6890); 368 (ε 12130) (MeOH).

2'-De(methylamino), 2'-hydroxy: **Milliamine I**

[91197-54-5]


$\text{C}_{36}\text{H}_{39}\text{NO}_{10}$ 645.705

Alkaloid from *Euphorbia milii*. Glass. λ_{max} 212 (ε 42090); 237 (ε 23080); 260 (ε 12220); 322 (ε 12060) (MeOH).

Marston, A. et al., *Planta Med.*, 1984, **50**, 319-322

Millingtonine M-618

[172924-27-5]




$\text{C}_{32}\text{H}_{45}\text{NO}_{14}$ 667.706

Alkaloid from flower buds of *Millingtonia hortensis* (Bignoniaceae). Powder + $1\text{H}_2\text{O}$. Sol. MeOH, CHCl_3 ; poorly sol. cyclohexane. Racemic. λ_{max} 209 (E1%/1cm 362); 226 (E1%/1cm 371); 257 (E1%/1cm 329); 497 (E1%/1cm 169) (MeOH) (Berdy).

Hase, T. et al., *Phytochemistry*, 1996, **41**, 317-321 (isol, pmr, cmr, struct)

Milnamide A M-619

[156430-97-6]



$\text{C}_{31}\text{H}_{46}\text{N}_4\text{O}_4$ 538.729

Peptide antibiotic. Alkaloid from Papua New Guinean collections of the marine sponge *Auletta cf. constricta*. Cytotoxic agent, microtubule formation inhibitor. Amorph. solid. Sol. MeOH, CHCl_3 , $[\alpha]_D^{27}$ +28.8 (c, 0.5 in CH_2Cl_2). λ_{max} 223 (ε 14700); 285 (ε 2810); 294 (ε 3070); 304 (ε 3030); 380 (ε 1260) (MeOH).

1-(2N)-Dehydro: **Milnamide D**

$\text{C}_{31}\text{H}_{45}\text{N}_4\text{O}_4^{\oplus}$ 537.721

Quaternary alkaloid from a *Cymbastela* sp. Cytotoxic agent. CAS No. not found 8-14Cl.

1-Oxo: **Milnamide C**

[675875-88-4]

$\text{C}_{31}\text{H}_{44}\text{N}_4\text{O}_5$ 552.712

Alkaloid from an *Auletta* sp. Cytotoxic. Yellow solid.

Crews, P. et al., *J.O.C.*, 1994, **59**, 2932-2934 (isol, uv, ir, pmr, cmr)


Chevallier, C. et al., *Org. Lett.*, 2003, **5**, 3737-3739 (Milnamide D)

Liu, C. et al., *Angew. Chem., Int. Ed.*, 2004, **43**, 5951-5954 (synth)

Sonnenschein, R.N. et al., *Org. Lett.*, 2004, **6**, 779-782 (Milnamide C)

Miltanthine M-620

[201136-33-6]




$\text{C}_{21}\text{H}_{28}\text{NO}_4^{\oplus}$ 358.457

Quaternary alkaloid from *Papaver triniifolium*. Amorph. solid (as iodide). $[\alpha]_D^{+97}$ (c, 0.4 in CHCl_3). λ_{max} 218 (log ε 4.13); 283 (log ε 3.34) (MeOH). λ_{max} 283 (log ε 3.34) (iodide).

Wang, B.-G. et al., *Planta Med.*, 1997, **63**, 577-578 (isol, uv, ir, pmr)

Miltanthoridine M-621



$\text{C}_{20}\text{H}_{23}\text{NO}_4$ 341.406

Alkaloid from *Papaver triniifolium*. Amorph. yellow solid. λ_{max} 280; 332; 420 (MeOH).

α -Oxo: **Miltanthoridinone**


$\text{C}_{20}\text{H}_{21}\text{NO}_5$ 355.39

Alkaloid from *Papaver triniifolium*. Orange solid. λ_{max} 286; 342; 432 (MeOH).

Sari, A. et al., *Nat. Prod. Res.*, 2006, **20**, 493-496 (isol, pmr, cmr, ms)

Mimocin M-622

[76177-28-1]



$\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_5$ 302.286

Benzoquinone antibiotic. Isol. from *Streptomyces lavendulae*. Active against *Bacillus subtilis*. Yellow prisms (Et_2O). Mp 189-191° dec. λ_{max} 243 (ε); 322 (ε) (MeOH) (Derep). λ_{max} 243; 322 (MeOH) (Berdy).

Kubo, A. et al., *Tet. Lett.*, 1980, **21**, 3207-3208 (uv, ir, pmr, ms, struct, synth)

Matsuo, K. et al., *Chem. Pharm. Bull.*, 1982, **30**, 4170-4174 (synth)


Kubo, A. et al., *Chem. Pharm. Bull.*, 1983, **31**, 341-343; 1985, **33**, 823-830; 1988, **36**, 4355-4363 (synth, cmr)

Saito, N. et al., *Chem. Pharm. Bull.*, 1989, **37**, 1493-1499 (synth)

Kuwabara, N. et al., *Tetrahedron*, 2004, **60**, 2943-2952 (synth)

Mimopudine M-623

1,1'-[(Hydroxyimino) bis(1-amino-2-hydroxy-2,1-ethanediy)] bis[4(1H)-pyridinone], 9CI [221056-52-6]



$\text{C}_{14}\text{H}_{19}\text{N}_5\text{O}_5$ 337.335


Alkaloid from *Mimosa pudica*. Leaf-opening substance. Yellow powder.

Ueda, M. et al., *Tetrahedron*, 1999, **55**, 10937-10948 (isol, pmr, cmr, ms)

- Boit, H.-G. *et al.*, *Naturwissenschaften*, 1961, **48**, 603 (*Clivimine, isol, ir*)
 Döpke, W. *et al.*, *Tet. Lett.*, 1967, 451; 1970, 745 (*Clivimine, Miniatine, pmr, ms, struct, stereochem, bibl*)
 Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **38**, 1827 (*Clivimine, isol, ir, pmr*)
 Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591 (*Clivimine, cd*)

Minovincine M-630

Methyl 2,3-didehydro-20-oxoaspidospermidine-3-carboxylate, 9CI. Minoricine



$C_{21}H_{24}N_2O_3$ 352.432

(+)-form [15622-69-2]

Alkaloid from *Tabernaemontana riedelii* (Apocynaceae). Noncryst. $[\alpha]_D^{26}$ +340 (c, 1 in EtOH).

(-)-form [6792-12-7]

Alkaloid from *Vinca minor* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -504 (c, 0.5 in EtOH). $[\alpha]_D$ -534 (MeOH).

11-Methoxy: Minoricine. 11-Methoxyminovincine [22341-30-6]

Alkaloid from *Vinca minor* (Apocynaceae). Amorph. Mp 141°. $[\alpha]_D^{20}$ -414 (c, 0.03 in EtOH).

Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 2237 (*isol, uv, ir, pmr, ms, struct, deriv*)

Zachystalová, D. *et al.*, *Chem. Ind. (London)*, 1963, 610 (*Minoricine*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 1055 (*isol, uv, ir, pmr, ms, deriv*)


Döpke, W. *et al.*, *Tet. Lett.*, 1968, 605; 1971, 1287 (*Minoricine*)

Kuehne, M. *et al.*, *Tetrahedron*, 1983, **39**, 3707; 3715 (*synth*)

Kalaus, G. *et al.*, *J.O.C.*, 1997, **62**, 9188-9191 (*synth*)

Minovincinine M-631

Methyl 2,3-didehydro-20-hydroxyaspidospermidine-3-carboxylate, 9CI. Alkaloid V3



$C_{21}H_{26}N_2O_3$ 354.448

(+)-form [6801-25-8]

Alkaloid from *Alstonia venenata* (Apocynaceae). Noncryst. $[\alpha]_D$ +338 (CHCl₃). $[\alpha]_D$ +441 (EtOH).

O-Ac: Echitovenine [4697-85-2]

$C_{23}H_{28}N_2O_4$ 396.485

Alkaloid from *Alstonia venenata* (Apocynaceae). Mp 170-172°. $[\alpha]_D$ +690 (CHCl₃).

19-Epimer, O-(3,4,5-trimethoxybenzoyl):

19-Epi-(+)-echitoveniline

$C_{31}H_{36}N_2O_7$ 548.635

Alkaloid from the leaves of *Alstonia venenata* (Apocynaceae). Prisms (EtOAc/petrol). Mp 168-169°. $[\alpha]_D$ +462 (EtOH).

(-)-form

Alkaloid from *Vinca minor*, *Catharanthus longifolius* and *Catharanthus trichophyllus* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -418 (c, 1 in EtOH). $[\alpha]_D^{22}$ -473 (c, 0.85 in EtOH) (synthetic). λ_{max} 225 (log ϵ 4.02); 297 (log ϵ 4); 328 (log ϵ 4.09) (no solvent reported).

O-(3-Methyl-2-butenoyl): Echitovenidine [7222-35-7]

$C_{26}H_{32}N_2O_4$ 436.55

Alkaloid from *Alstonia venenata* (Apocynaceae). Mp 162-163°. $[\alpha]_D$ -580 (CHCl₃). Cooccurrence with Echitovenine is remarkable.

► C0145000

O-(3,4,5-Trimethoxybenzoyl): Echitoveniline

[72855-79-9]

$C_{31}H_{36}N_2O_7$ 548.635

Alkaloid from *Alstonia venenata* (Apocynaceae). Amorph. $[\alpha]_D$ -263 (CHCl₃).

O-(3-Methoxy-4,5-methylenedioxybenzoyl): Echitoserpidine

[52718-36-2]

$C_{30}H_{32}N_2O_7$ 532.592

Alkaloid from the fruits of *Alstonia venenata* (Apocynaceae). Cryst. (MeOH). Mp 110°. $[\alpha]_D$ -427 (CHCl₃).

11-Methoxy: 11-Methoxyminovincine

[22341-28-2]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Vinca minor* (Apocynaceae). Amorph. $[\alpha]_D^{24}$ -395 (c, 0.2 in CHCl₃).

11-Methoxy, O-Ac: Echitovenaldine

[52358-57-3]

$C_{24}H_{30}N_2O_5$ 426.511

Minor alkaloid from the leaves of *Alstonia venenata* (Apocynaceae). Mp 148°. $[\alpha]_D$ -485 (CHCl₃).

11-Methoxy, O-(3,4,5-trimethoxybenzoyl): 11-Methoxyechitoveniline. 16-Methoxyechitoveniline

[72855-80-2]

Alkaloid from *Alstonia venenata* (Apocynaceae). Amorph. $[\alpha]_D$ -388 (EtOH).

11-Methoxy, O-(3-Methyl-2-butenoyl): 11-Methoxyechitovenidine

[72855-81-3]

$C_{27}H_{34}N_2O_5$ 466.576

Alkaloid from *Alstonia venenata* (Apocynaceae). Plates (MeOH). Mp 140°. $[\alpha]_D$ -325 (CHCl₃).

11-Methoxy, O-(3-methoxy-4,5-methylenedioxybenzoyl): Echitoserpine

[55870-82-1]

$C_{31}H_{34}N_2O_8$ 562.618

Alkaloid from the fruits of *Alstonia venenata* (Apocynaceae). Cryst. (MeOH). Mp 154°. $[\alpha]_D$ -444.5 (CHCl₃).

19-Epimer. 19-Epiminovincinine

[26568-42-3]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from *Vinca minor* (Apocynaceae). Mp 135°. $[\alpha]_D$ -580 (c, 0.2 in EtOH).

19-Epimer, 14-oxo: Baloxine. 20S-Hydroxy-7-oxovincadifformine

[38225-09-1]

$C_{21}H_{24}N_2O_4$ 368.432

Minor alkaloid from *Melodinus balsae* (Apocynaceae). Noncryst. $[\alpha]_D$ -120 (c, 0.1 in CHCl₃).

[34020-05-8]

Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 2237-2241 (*isol, uv, ir, pmr, ms, struct*)

Das, B. *et al.*, *Tet. Lett.*, 1966, 2483 (*uv, ir, pmr, ms, struct*)

Döpke, W. *et al.*, *Tet. Lett.*, 1968, 6065; 1970, 749; 1971, 1287 (*16-Methoxyminovincinine, 16-Epiminovincinine*)

Mehri, M.H. *et al.*, *Bull. Soc. Chim. Fr.*, 1972, 3291-3292 (*Baloxine*)

Majumder, P.L. *et al.*, *Chem. Ind. (London)*, 1973, 1032 (*Echitovenaldine*)

Majumder, P.L. *et al.*, *Phytochemistry*, 1974, **13**, 645 (*Echitoserpidine*)

Majumder, P.L. *et al.*, *Tetrahedron*, 1974, **30**, 2761; *tet.*, 1979, **35**, 1151; 1981, **37**, 1243 (*16-Methoxyminovincinine, Echitoserpine, 11-Methoxyechitovenidine, Echitoveniline, 19-Epi-(+)-echitoveniline*)

Langlois, N. *et al.*, *J.O.C.*, 1979, **44**, 2468 (*synth*)

Caron, C. *et al.*, *Heterocycles*, 1981, **16**, 645-646 (*Baloxine, synth, abs config*)

Minpeimine**M-632**

$C_{27}H_{43}NO_2$ 413.642


Struct. unknown. Alkaloid from the Chinese drug Pei-mu (*Fritillaria* sp.) (Liliaceae). Mp 210-212°. Contains a carbonyl group.

Chu, T.-T. *et al.*, *CA*, 1959, **53**, 647

Minpeiminine**M-633**

Struct. unknown. Alkaloid from the Chinese drug Pei-mu (*Fritillaria* sp.) (Liliaceae), occurring with Minpeimine, M-632. Mp 215-216°.

Chu, T.T. *et al.*, *CA*, 1959, **53**, 647

Mirabazole A**M-634**

$R^1 = H, R^2 = CH_3$

$C_{18}H_{24}N_4S_4$ 424.678

4',5'-Didehydro: Didehydromirabazole A
[135432-51-8]
C₁₈H₂₂N₄S₄ 422.663
Isol. from the terrestrial blue-green alga *Scytonema mirabile*. Cytotoxic agent. [α]_D -26 (c, 0.44 in CHCl₃). May be an artifact formed from Mirabazole A. Config. of ring-A stereocentre revised in 1994.
Carmeli, S. *et al.*, *Tet. Lett.*, 1991, **32**, 2593-2596 (*isol, pmr, struct*)
Pattenden, G. *et al.*, *J.C.S. Perkin 1*, 1993, 1629-1636 (*synth, Didehydromirabazole A*)
Boyce, R.J. *et al.*, *Synlett*, 1994, 587-588 (*synth, config, Didehydromirabazole A*)

Mirabazole B M-635


[135459-34-6]
As Mirabazole A, M-634 with
R¹ = R² = CH₃
C₁₉H₂₆N₄S₄ 438.705
Isol. from the blue-green alga *Scytonema mirabile*. Cytotoxic agent. [α]_D -166 (c, 0.09 in CHCl₃). 12CI name is incorrect.
Carmeli, S. *et al.*, *Tet. Lett.*, 1991, **32**, 2593 (*isol, pmr, struct*)
Walker, M.A. *et al.*, *J.O.C.*, 1992, **57**, 5566 (*synth*)
Parsons, R.L. *et al.*, *Tet. Lett.*, 1994, **35**, 1383 (*synth*)
Kuriyama, N. *et al.*, *Tetrahedron*, 1997, **53**, 8323 (*synth, pmr*)

Mirabazole C M-636

[135432-52-9]
As Mirabazole A, M-634 with
R¹ = CH₃, R² = H
C₁₈H₂₄N₄S₄ 424.678
Config. of ring-A stereocentre revised in 1994. Isol. from the terrestrial blue-green alga *Scytonema mirabile*. Cytotoxic. [α]_D -113 (c, 0.024 in CHCl₃). 12CI name is incorrect.
Carmeli, S. *et al.*, *Tet. Lett.*, 1991, **32**, 2593 (*isol, pmr, struct*)
Walker, M.A. *et al.*, *J.O.C.*, 1992, **57**, 5566 (*synth*)
Parsons, R.L. *et al.*, *Tet. Lett.*, 1994, **35**, 1379 (*config, synth*)
Akaji, K. *et al.*, *J.O.C.*, 1996, **61**, 3350 (*synth*)

Mirabilin† M-637


[1009368-84-6]



C₇₆H₁₂₃N₃O₂₂ 1430.815
Alkaloid from *Siliquariaspongia mirabilis*. Antitumour agent. Pale yellow powder. [α]_D²⁵ -14 (c, 0.05 in MeOH). λ_{max} 350 (log ε 3.96) (MeOH).
Plaza, A. *et al.*, *J. Nat. Prod.*, 2008, **71**, 473-477 (*isol, pmr, cmr*)

Mirabilin B M-638

[182266-75-7]

R = -CH₂CH₂CH₂CH₃

C₁₅H₂₃N₃ 245.367
Alkaloid from the sponges *Arenochalina mirabilis*, *Batzella* sp. and *Monanchora unguifera*. Antifungal agent. Cryst. [α]_D +41.6 (c, 0.48 in MeOH). λ_{max} 236; 302; 375 (MeOH).

N-Ac:

Pale yellow oil. [α]_D +89 (c, 0.40 in CHCl₃). λ_{max} 244 (ε 11800); 275 (ε 400) (MeOH).
Barrow, R.A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 767-773 (*isol, pmr, cmr, ms*)
Patil, A.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 704-707 (*isol, pmr, cmr*)
Hua, H.-M. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 6461-6464 (*isol*)
Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*synth, pmr, cmr*)

Mirabilin C M-639

[182266-76-8]

As Mirabilin B, M-638 with
R = -CH₂CH=CHCH₂CH₂CH₃ (Z)

C₁₇H₂₅N₃ 271.405
Alkaloid from the Australian marine sponge *Arenochalina mirabilis*. Pale yellow oil (as N-Ac). [α]_D +153 (c, 0.57 in CHCl₃) (N-Ac). Isol. and characterised as the monoacetate. λ_{max} 244 (ε 12400); 274 (ε 4500) (MeOH) (N-Ac) (Berdy).

10-Epimer: **Mirabilin A**


[182145-37-5]

C₁₇H₂₅N₃ 271.405
From *Arenochalina mirabilis*. Pale yellow oil (as N-Ac). [α]_D +12.8 (c, 0.74 in CHCl₃) (N-Ac). λ_{max} 206 (ε 5500); 244 (ε 1300); 275 (ε 4400) (MeOH) (N-Ac) (Berdy).

Barrow, R.A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 767 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mirabilin D M-640

[182145-57-9]

R = -CH₂CH=CHCH₂CH₂CH₃ (Z)

C₁₇H₂₇N₃O 289.42
Alkaloid from the Australian marine sponge *Arenochalina mirabilis*. Pale yellow oil (as di-Ac). [α]_D +285 (c, 0.3 in CHCl₃) (di-Ac). λ_{max} 204 (ε 4600); 246 (ε 3600); 287 (ε 3750) (MeOH) (di-Ac) (Berdy). λ_{max} 246 (ε 3600); 287 (ε 3750) (EtOH) (di-Ac).

Relative Configuration

Relative Configuration

Relative Configuration

Barrow, R.A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 767 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mirabilin E M-641

[182145-71-7]


As Mirabilin D, M-640 with
R = -CH₂CH₂CH₂CH₃

C₁₅H₂₅N₃O 263.382
Alkaloid from the Australian marine sponge *Arenochalina mirabilis*. Pale yellow oil (as di-Ac). [α]_D +332 (c, 0.05 in CHCl₃) (di-Ac). Relative stereochem. about C9 not detd. λ_{max} 202 (ε 2100); 241 (ε 1650); 285 (ε 1250) (EtOH) (di-Ac).

Barrow, R.A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 767 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mirabilin F M-642

[182145-74-0]

R = -CH=CHCH₂CH₃ (E)C₁₅H₂₃N₃ 245.367

Appears to be a didehydro deriv. of Ptilocaulin, P-773; shown with different tautomerism of the cyclic guanidine system. Alkaloid from the Australian marine sponge *Arenochalina mirabilis*. Pale yellow oil (as N-Ac). [α]_D +384 (c, 0.2 in CHCl₃) (N-Ac). λ_{max} 241 (ε 3278); 295 (ε 3150) (no solvent reported) (Berdy). λ_{max} 241 (ε 5278); 295 (ε 3150) (EtOH) (N-Ac).

Barrow, R.A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 767-773 (*isol, uv, ir, pmr, cmr, ms, struct*)


Mirabilin G M-643As Mirabilin F, M-642 with
R = -CH=CH(CH₂)₃CH₃ (E)C₁₇H₂₇N₃ 273.42

Alkaloid from the Australian marine sponge *Clathria* sp. Pale yellow oil. [α]_D +49 (c, 0.1 in CHCl₃). λ_{max} 241 (log ε 3.7); 295 (log ε 3.4) (EtOH).

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 643-644 (*isol, uv, pmr, cmr, ms*)

Mirabimide E M-644

[159903-50-1]


C₁₈H₂₆Cl₄N₂O₅ 492.225

Isol. from the terrestrial blue-green alga *Scytonema mirabile*. Cytotoxic. [α]_D +6.5 (c, 0.9 in CHCl₃).

Paik, S. *et al.*, *J.A.C.S.*, 1994, **116**, 8116 (*isol, ir, pmr, cmr, synth, struct*)

Miraxanthin I M-645

4-[[[1-Carboxy-3-(methylsulfinyl)propyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI. 4-[[N-[1-Carboxy-3-(methylsulfinyl)propyl]formimido]methylene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 8CI [5296-79-7]




$C_{14}H_{18}N_2O_7S$ 358.371

Yellow pigment of flowers of *Mirabilis jalapa* (Nyctaginaceae). Yellow solid.

Piattelli, M. *et al.*, *Phytochemistry*, 1965, **4**, 817 (isol, struct, uv, synth)

Miraxanthin II M-646

4-[[[(1,2-Dicarboxyethyl)imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI. Aspartic acid betaxanthin [5375-63-3]




$C_{13}H_{14}N_2O_8$ 326.262

Yellow pigment of flowers of *Mirabilis jalapa* (Nyctaginaceae). Yellow solid.

Piattelli, M. *et al.*, *Phytochemistry*, 1965, **4**, 817 (isol, struct, uv, synth)

Miraxanthin III M-647

1,2,3,4-Tetrahydro-4-[[[2-(4-hydroxyphenyl)ethyl]imino]ethylidene]-2,6-pyridinedicarboxylic acid, 9CI. Tyraminebetaxanthin [5589-85-5]



$C_{17}H_{18}N_2O_5$ 330.34

Yellow pigment of flowers of *Mirabilis jalapa* (Nyctaginaceae). Yellow solid. L-Config. probable on biosynth. grounds.

3'-Hydroxy: Dopaminebetaxanthin. **Mir-**

axanthin V

[5375-64-4]

$C_{17}H_{18}N_2O_6$ 346.339

Yellow pigment from *Mirabilis jalapa* (Nyctaginaceae). Yellow solid.

3'-Methoxy: 3-Methoxytyraminebetaxanthin

$C_{18}H_{20}N_2O_6$ 360.366

Yellow pigment from *Celosia argentea*.

Piattelli, M. *et al.*, *Phytochemistry*, 1965, **4**, 817 (isol, struct, uv, synth)

Kobayashi, N. *et al.*, *Phytochemistry*, 2001, **56**, 429-436 (biosynth)


Schliemann, W. *et al.*, *Phytochemistry*, 2001, **58**, 159-165 (3-Methoxytyraminebetaxanthin)

Stintzing, F.C. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 1008-1016 (Miraxanthin V, pmr, cmr)

Miraziridine A

M-648

[312322-94-4]



Absolute Configuration

$C_{30}H_{52}N_8O_9$ 668.789

Isol. from the marine sponge *Theonella* aff. *mirabilis*. Cysteine protease inhibitor. Also amino-proteinase and serine-proteinase inhibitor. $[\alpha]_D^{20}$ -74 (c, 0.085 in MeOH). λ_{max} 205 (ϵ 14000) (MeOH).

Nakao, Y. *et al.*, *J.A.C.S.*, 2000, **122**, 10462-10463 (isol, cmr, pmr)


Schaschke, N. *et al.*, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 855-857 (pharmacol)

Konno, H. *et al.*, *Tetrahedron*, 2007, **63**, 9502-9513 (synth)

Misenine

M-649

[202075-41-0]



$m + n = 6$

Relative Configuration

$C_{33}H_{54}N_2O$ 494.802


Alkaloid from the sponge *Reniera* sp. Oil. $[\alpha]_D$ +6.4 (c, 2.2 in $CHCl_3$).

Guo, Y. *et al.*, *Tetrahedron*, 1998, **54**, 541-550 (isol, ir, pmr, cmr)

Misramine

M-650

[94801-27-1]



$C_{19}H_{25}NO_4$ 331.411

Alkaloid from the above-ground parts of *Roemeria hybrida* and *Roemeria dodecandra* (Papaveraceae). Cryst. (MeOH). Mp 103-105°. $[\alpha]_D^{25}$ -23 (c, 0.1 in MeOH). $[\alpha]_D^{25}$ -26 (c, 0.1 in C_6H_6). $[\alpha]_D^{25}$ -22 (c, 0.1 in Me_2CO). The first pentacyclic proaporphine alkaloid. Probably derived biogenetically from a diastereomer of Orientalinone, O-112.

O²-De-Me: Labrandine. Misrametine

[128581-42-0]

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. $[\alpha]_D$ -120 (c, 0.22 in MeOH). $[\alpha]_D$ -145 (c, 0.12 in $CHCl_3$).

O²-De-Me, di-Ac:

Amorph. $[\alpha]_D$ -120 (c, 0.12 in MeOH).

El-Masry, S. *et al.*, *J.O.C.*, 1985, **50**, 729 (uv, ir, pmr, ms, cd, struct)


El-Masry, S. *et al.*, *Alexandria J. Pharm. Sci.*, 1990, **4**, 90; *CA*, **114**, 160639 (Misrametine)

Gözler, B. *et al.*, *Heterocycles*, 1990, **31**, 149 (Labrandine)

Mitiphylline

M-651

[22214-87-5]



Absolute Configuration

$C_{31}H_{49}NO_6$ 531.731

Alkaloid from leaves of *Holarrhena mitis* (Apocynaceae). Noncryst. $[\alpha]_D$ +20 (c, 1 in $CHCl_3$).

N-Ac: N-Acetylmitiphylline

$C_{33}H_{51}NO_7$ 573.768

Alkaloid from leaves of *Holarrhena mitis* (Apocynaceae). Cryst. (Me_2CO). Mp 126-128°. $[\alpha]_D$ +38 (c, 2 in $CHCl_3$).

N-De-Me: N-Demethylmitiphylline

$C_{30}H_{47}NO_6$ 517.704

Alkaloid from leaves of *Holarrhena mitis* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ +9 ($CHCl_3$).

N-De-Me, N-Ac:

Cryst. (MeOH). Mp 253-256°. $[\alpha]_D^{20}$ +16 ($CHCl_3$). Cryst. with difficulty.


Janot, M.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1968, **267**, 1050 (isol, uv, ir, pmr, ms, struct)

Leboeuf, M. *et al.*, *Phytochemistry*, 1972, **11**, 843 (isol, struct, derivs)

Mitromycin A

M-652

[27164-43-8]



Absolute configuration

C₁₆H₁₇N₃O₅ 331.327

Isol. from *Streptomyces verticillatus*. Antibiotic active against gram-positive and -negative bacteria and certain tumours. Red-purple needles. Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 124-126°. Related to the Mitomycins. λ_{max} 227 (ε 8920); 322 (ε 7580); 530 (ε 1030) (MeOH) (Berdy).

► CN1065000

Lefemine, D.V. et al., *J.A.C.S.*, 1962, **84**, 3184 (isol, uv)


Morton, G.O. et al., *J.A.C.S.*, 1970, **92**, 2588 (struct, ir, pmr, ms)

Kono, M. et al., *J. Antibiot.*, 1991, **44**, 309 (abs config)

Mitomycin A

Mitiomycin B

[4055-39-4]



Absolute Configuration

C₁₆H₁₉N₃O₆ 349.343

Isol. from *Streptomyces caespitosus*. Antitumour agent, selectively inhibits DNA synthesis. Antibiotic. Purple needles. Mp 159-161° dec. [α]_D²⁵ -143 (c, 0.1 in MeOH). λ_{max} 235; 285; 335; 435 (MeOH/HCl) (Derep). λ_{max} 360; 555 (MeOH/NaOH) (Derep). λ_{max} 215 (ε 15000); 317 (ε 9300); 528 (ε 1300) (MeOH) (Derep).

► LD₅₀ (mus, ivn) 1 mg/kg. CN1050000

1-N-Me: **Mitomycin F**. 1-Methylmitomycin A

[18209-14-8]

C₁₇H₂₁N₃O₆ 363.369

From *Streptomyces caespitosus*. Antitumour agent. Reddish-purple cryst. Mp 145°. λ_{max} 220 (ε 12050); 324 (ε 7580); 520 (ε 850) (MeOH) (Berdy).

► LD₅₀ (mus, ivn) 4.3 mg/kg.

De(carbamoyloxy), 8,15-didehydro, 1-N-Me: **Mitomycin K**

[74148-45-1]

C₁₆H₁₈N₂O₄ 302.329

Prod. by *Streptomyces caespitosus*. Antitumour agent. Purple cryst.

8-Epimer, 1-N-Me: **Mitomycin J**

[74985-82-3]

C₁₇H₂₁N₃O₆ 363.369

From *Streptomyces caespitosus*. Antitumour agent. Sol. MeOH, CHCl₃. λ_{max} 219 (ε 12550); 324 (ε 10200); 539 (ε 1230) (MeOH) (Berdy).

► CN0736000

8-Epimer, O⁷-de-Me, 1-N-Me: **Mitomycin B**. Mitiomycin C

[4055-40-7]

C₁₆H₁₉N₃O₆ 349.343

Isol. from *Streptomyces verticillatus* and *Streptomyces caespitosus*. Antitumour agent. Purple-blue needles. [α]_D -835 (c, 0.012 in MeOH). Dec. without melting. λ_{max} 235; 285; 335; 435 (MeOH/HCl) (Derep). λ_{max} 360; 555 (MeOH/NaOH) (Derep). λ_{max} 215 (ε 15000); 317 (ε 9300); 528 (ε 1300) (MeOH) (Derep).

► LD₅₀ (mus, ivn) 3 mg/kg. CN0875000

Stereoisomer: **Antibiotic A**

[87139-13-7]

C₁₆H₁₉N₃O₆ 349.343

From *Streptomyces caespitosus*. Active against gram-positive and -negative bacteria. Cryst. Sol. MeOH, Me₂CO; poorly sol. H₂O, hexane. Mp 130° grad. dec. [α]_D²⁶ -27 (c, 0.5 in CHCl₃). Stereochem. undefined. λ_{max} 215 (ε 10000); 288 (ε 6000) (MeOH) (Berdy).

Demethoxy, 7,8-didehydro, 1-N-Me: 7-Methoxy-1,2-(N-methylaziridino)mitosene

[37394-29-9]

C₁₆H₁₇N₃O₅ 331.327

Biologically active redn. prod. from Mitomycin B. Antitumour antibiotic. Orange plates (DMF or Py). [α]_D²⁵ +36.4 (c, 0.99 in DMSO). Does not melt but loses birefringence at 230°.

Webb, J.S. et al., *J.A.C.S.*, 1962, **84**, 3185-3187; 3187-3188 (struct, ir, uv, ms, nmr)

Tulinsky, A. et al., *J.A.C.S.*, 1962, **84**, 3188-3190; 1967, **89**, 2905-2911 (cryst struct)

Patrick, J.B. et al., *J.A.C.S.*, 1964, **86**, 1889-1890 (7-Methoxy-1,2-(N-methylaziridino)mitosene)

v. Lear, G.E. et al., *Tetrahedron*, 1970, **26**, 2587-2597 (struct, ms)

Kinoshita, S. et al., *J. Med. Chem.*, 1971, **14**, 103-107 (synth, activity)

Hornemann, U. et al., *J. Antibiot.*, 1975, **28**, 841-843 (biosynth)

Yahashi, R. et al., *J. Antibiot.*, 1976, **29**, 104-106 (cryst struct, Mitomycin B)

Fukuyama, T. et al., *Tet. Lett.*, 1977, 4295-4298 (synth)

Urakawa, C. et al., *J. Antibiot.*, 1981, **34**, 243-244; 1152-1156 (1-N-Me, derivs)

Shirahata, K. et al., *J.A.C.S.*, 1983, **105**, 7199-7200 (abs config)


Japan. Pat., 1983, 83 89 189; CA, **99**, 138160 (Antibiotic A)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, MQX500; MQX750

Mitomycin C, JAN

M-654

Mitomycin, BAN, INN, USAN. Mitiomycin S. Mitiomycin E. Ametycine. Mitocin C. Mutamycin. NSC 26980 [50-07-7]



Absolute Configuration

C₁₅H₁₈N₄O₅ 334.331

Benzoquinone antibiotic. Struct. revised in 1983. Isol. from *Streptomyces caespitosus*. Antineoplastic antibiotic. Deep blue-violet cryst. Sol. MeOH, CHCl₃, Me₂CO, acids; fairly sol. H₂O, C₆H₆, Et₂O; poorly sol. hexane. Mp 360°. Log P -3.04 (uncertain value) (calc). Mitomycin S was shown to be a mixt. of Mitomycin C with NaCl. λ_{max} 217 (ε 24600); 245 (sh) (ε 11000); 360 (ε 23000); 558 (ε 209) (MeOH) (Derep). λ_{max} 216 (ε 21880); 254; 360 (ε 21400); 420; 555 (ε 235) (MeOH) (Berdy).

► Adverse systemic effects reported when

used therapeutically incl. bone-marrow depression, gastrointestinal, renal, pulmonary and cardiac toxicity. A possible human carcinogen. LD₅₀ (rat, orl) 30 mg/kg. LD₅₀ (mus, ivn) 5 mg/kg. LD₅₀ (mus, ipr) 9 mg/kg. Exp. carcinogen. Exp. reprod. and teratogenic effects. CN0700000

N-Me: **Porfiromycin**, BAN, INN, USAN. Mitiomycin D. NSC 56410. Methylmitomycin C

[801-52-5]

C₁₆H₂₀N₄O₅ 348.358

Isol. from the fermentation broth of *Streptomyces ardens*. Antibacterial and antineoplastic agent. Purple triclinic cryst. Fairly sol. H₂O, C₆H₆, CHCl₃, Me₂CO; poorly sol. Et₂O, hexane. Mp 201-201.5° dec. [α]_D²⁵ +275 (c, 0.1 in MeOH). Log P -2.18 (uncertain value) (calc). λ_{max} 217 (ε 24600); 245 (sh) (ε 11000); 360 (ε 23000); 558 (ε 209) (MeOH) (Derep).

► LD₅₀ (rat, orl) 68 mg/kg; LD₅₀ (mus, ipr) 20 mg/kg. CN0525000

N⁴-(3-Oxobutyl): **Antibiotic MMC 4**.

MMC 4. N-(3-Oxobutyl)mitomycin C [183997-07-1]

C₁₉H₂₄N₄O₆ 404.422

Prod. by *Streptomyces ardens* sp. A1143. Antitumour agent. Red-violet cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D +270 (MeOH). λ_{max} 216 (E1%/1cm 547); 360 (E1%/1cm 505) (MeOH) (Berdy).

O-De-Me, N-Me: **Mitomycin D**. 6-Amino-6-demethoxymitomycin B

[10169-34-3]

C₁₅H₁₈N₄O₅ 334.331

Also semisynthetic. From *Streptomyces caespitosus*. Shows antileukaemic props. Dark bluish-purple prisms (C₆H₆/MeOH). Mp 234-236°. λ_{max} 217; 361; 558 (MeOH) (Berdy).

► CN0726000

8-Epimer, N-Me: **Mitomycin E**

[74707-94-1]

C₁₆H₂₀N₄O₅ 348.358

Semisynthetic. Shows antileukaemic props. λ_{max} 217 (ε 17800); 361 (ε 19100); 558 (ε 215) (MeOH) (Berdy).

Stereoisomer: [87139-12-6]

C₁₅H₁₈N₄O₅ 334.331

Prod. by *Streptomyces caespitosus* and *Micromonospora* sp. Active against gram-positive and -negative bacteria. Cryst. Sol. MeOH, Me₂CO; poorly sol. H₂O, hexane. Mp 120° (grad. dec.). [α]_D²⁵ +34.7 (c, 0.3 in CHCl₃). λ_{max} 243 (ε 5000); 344 (ε 6300) (MeOH) (Berdy).

[52081-33-1]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 1068A (ir)

Webb, J.S. et al., *J.A.C.S.*, 1962, **84**, 3185-3187; 3187-3188 (struct, ir, uv, ms, pmr)

Uzu, K. et al., *Agric. Biol. Chem.*, 1964, **28**, 394-402 (isol, struct)

Carter, S.K. et al., *Cancer Chemother. Rep., Suppl.*, 1968, **1**, 81 (Porfiromycin, rev, pharmacol)


v. Lear, G.E. et al., *Tetrahedron*, 1970, **26**, 2587-2597 (struct, ms)

Lown, J.W. et al., *Can. J. Chem.*, 1974, **52**, 2331-2336 (ir, uv, ms, pmr, cmr, struct)

- Hornemann, U. *et al.*, *J. Antibiot.*, 1975, **28**, 841-843 (*biosynth*)
IARC Monog., 1976, **10**, 171; *Suppl.* 7, 67 (*rev. tox*)
 Nakatsubo, F. *et al.*, *J.A.C.S.*, 1977, **99**, 8115-8116 (*Porfiromycin, synth, ir, pmr*)
 Fukuyama, T. *et al.*, *Tet. Lett.*, 1977, 4295-4298 (*synth*)
 Crooke, S.T. *et al.*, *Cancer Chemother.*, 1981, **3**, 49 (*rev. pharmacol*)
 Iyengar, B.S. *et al.*, *J. Med. Chem.*, 1981, **24**, 975-981 (*analogs, bibl*)
 Andrews, P.A. *et al.*, *J. Chromatogr.*, 1983, **262**, 231 (*Porfiromycin, hplc, ms*)
 Shirahata, K. *et al.*, *J.A.C.S.*, 1983, **105**, 7199-7200 (*abs config*)
Japan. Pat., 1985, 85 89 189; *CA*, **99**, 138160 (*stereoisomer*)
 Danishefsky, S.J. *et al.*, *J.A.C.S.*, 1986, **108**, 4648-4650 (*props*)
 Beijnen, J.H. *et al.*, *Anal. Profiles Drug Subst.*, 1987, **16**, 361-401 (*rev. pharmacol, synth, anal*)
 Fukuyama, T. *et al.*, *J.A.C.S.*, 1989, **111**, 8303-8304 (*synth*)
Japan. Pat., 1996, 96 245 626; *CA*, **126**, 6558e (*MMC 4*)
 Martindale, *The Extra Pharmacopoeia*, 32nd edn., *Pharmaceutical Press*, 1999, 552
 Abraham, L.M. *et al.*, *Drugs*, 2006, **66**, 321-340 (*rev*)
 Gruschow, S. *et al.*, *J.A.C.S.*, 2007, **129**, 6470-6476 (*biosynth*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., *J. Wiley*, 2000, AHK500; MLY000

Mitomycin G **M-655**

10-Decarbamoyloxy-9-dehydroporfiromycin
 [74148-46-2]




Absolute Configuration

- $C_{15}H_{17}N_3O_3$ 287.318
 Isol. from cultures of *Streptomyces caespitosus*. Green needles. Mp 270°. λ_{max} 222 (ϵ 10500); 289 (ϵ 10700); 373 (ϵ 17800); 602 (ϵ 234) (MeOH) (Berdy).
 ▶ CN0731000
Japan. Pat., 1980, 80 118 396; 80 120 587; *CA*, **94**, 82164; 103338 (*isol, synth*)

Mitomycin H **M-656**


10-Decarbamoyloxy-9-dehydromitomycin B
 [74148-44-0]



Absolute Configuration

- $C_{15}H_{16}N_2O_4$ 288.302
 Quinone antibiotic. Isol. from *Streptomyces caespitosus*. Active against gram-positive bacteria and tumours. Bluish-purple needles. Mp 125°. λ_{max} 226 (ϵ 12000); 291 (ϵ 11700); 578 (ϵ 1100) (MeOH) (Berdy).
 ▶ LD₅₀ (mus, ipr) 210 mg/kg. CN0740650

- Japan. Pat.*, 1980, 80 118 396; *CA*, **94**, 82164 (*isol*)
 Urakawa, C. *et al.*, *J. Antibiot.*, 1981, **34**, 243; 1152-1156 (*isol*)

Mitragynine **M-657**
 [4098-40-2]

Absolute configuration

- $C_{23}H_{30}N_2O_4$ 398.501
 Alkaloid from *Mitragyna speciosa* and from the genus *Uncaria* (Rubiaceae). Analgesic and antitussive agent. General depressant, anaesthetic. Noncryst. $[\alpha]_D^{25}$ -127 (CHCl₃). Log P 3.93 (calc).

Hydrochloride: Mp 243°.

Picrate: Mp 223-224°.

3,N-Didehydro-3-Dehydromitragynine

- [107286-93-1]
 $C_{23}H_{26}N_2O_4^{\oplus}$ 397.493
 Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae).

3,5,6-N-Tetrahydro-3,4,5,6-Tetrahydromitragynine

- $C_{23}H_{27}N_2O_4^{\oplus}$ 395.477
 Quaternary alkaloid from the leaves of *Mitragyna speciosa*. Dark orange powder. $[\alpha]_D^{18}$ -410 (c, 0.4 in CHCl₃). λ_{max} 220 ; 258 ; 314 ; 366 (MeOH).

A¹-Isomer, 7S-hydroxy: 7-Hydroxymitragynine

- $C_{23}H_{30}N_2O_5$ 414.5
 Alkaloid from leaves of *Mitragyna speciosa* (Rubiaceae). Amorph. powder. $[\alpha]_D^{25}$ +47.9 (c, 0.55 in CHCl₃).

3-Epimer, A¹-isomer, 7β-hydroxy: 7-Hydroxyspeciociliatine

- Alkaloid from the fruit of *Mitragyna speciosa*. Amorph. yellow solid. λ_{max} 221 (log ϵ 4.49); 240 (sh) (log ϵ 4.25); 294 (log ϵ 3.67) (MeOH).

3,20-Diepimer: Mitraciliatine


- [14509-92-3]
 $C_{23}H_{30}N_2O_4$ 398.501
 Alkaloid from *Mitragyna ciliata*, *Mitragyna tubulosa*, *Mitragyna speciosa* and *Uncaria* spp. (Rubiaceae). Mp 140-141°.

3,20-Diepimer, perchlorate:

Yellow prisms (Et₂O/EtOH). Mp 229-230°.

- Beckett, A.H. *et al.*, *J. Pharm. Pharmacol., Suppl.*, 1963, **15**, 166T; 267T (*Mitraciliatine*)
 Zacharias, D.E. *et al.*, *Acta Cryst.*, 1965, **18**, 1039 (*cryst struct*)
 Lee, C.M. *et al.*, *Tetrahedron*, 1967, **23**, 375 (*config, uv, ir, pmr, cd, ord*)
 Beckett, A.H. *et al.*, *Tetrahedron*, 1969, **25**, 5961 (*ms*)
 Phillipson, J.D. *et al.*, *J. Chromatogr.*, 1975, **105**, 163 (*occur*)
 Nakagura, N. *et al.*, *J.C.S. Perkin 1*, 1979, 2308 (*biosynth*)

- Houghton, P.J. *et al.*, *Phytochemistry*, 1986, **25**, 2910 (*3-Dehydromitragynine*)
 Ponglux, D. *et al.*, *Planta Med.*, 1994, **60**, 580 (*7-Hydroxy-7H-mitragynine*)
 Takayama, H. *et al.*, *Tet. Lett.*, 1995, **36**, 9337 (*synth*)
 Takayama, H. *et al.*, *Tetrahedron*, 1998, **54**, 8433-8440 (*Dehydromitragynine*)
 Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 916-928 (*rev*)
 Kitajima, M. *et al.*, *J. Nat. Med. (Tokyo)*, 2006, **60**, 28-35 (*7-Hydroxyspeciociliatine*)
 Ma, J. *et al.*, *Org. Lett.*, 2007, **9**, 3491-3494 (*synth*)

Mitrajavine **M-658**
 [13062-04-9]

Absolute Configuration

- $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from *Mitragyna javanica*, *Mitragyna hirsuta* and *Uncaria* spp. (Rubiaceae). Mp 117°. $[\alpha]_D^{23}$ -37.6 (c, 0.26 in CHCl₃). λ_{max} 227 (log ϵ 4.47); 265 (log ϵ 3.89); 292 (log ϵ 3.73) (EtOH).


Hydrobromide:

Cryst. + 1H₂O. Mp 230°.

3,20-Diepimer, O-de-Me, O-β-D-glucopyranoside: 9-β-D-Glucopyranosyloxy-tetrahydroalstonine

- [191481-02-4]
 $C_{27}H_{34}N_2O_9$ 530.574
 Alkaloid from seeds of *Tabernaemontana cymosa*. Amorph. $[\alpha]_D^{25}$ -16 (c, 0.2 in MeOH). λ_{max} 226 (log ϵ 4.42); 266 (sh) (log ϵ 3.8); 291 (sh) (log ϵ 3.66) (MeOH).


- Shellard, E.J. *et al.*, *J. Pharm. Pharmacol.*, 1966, **18**, 553-555 (*uv, ir, pmr, struct*)
 Phillipson, J.D. *et al.*, *Phytochemistry*, 1973, **12**, 1507 (*isol, ms*)
 Phillipson, J.D. *et al.*, *J. Chromatogr.*, 1975, **105**, 163-178 (*occur*)
 Achenbach, H. *et al.*, *Phytochemistry*, 1997, **45**, 325-335 (*9-Glucosyloxytetrahydroalstonine*)

Mitralactonal **M-659**
 [212264-09-0]

- $C_{21}H_{20}N_2O_4$ 364.4
 Abs. config. not definitely determined, but assigned as prob. S- (illus.) on biogenetic grounds. Alkaloid from the

leaves of *Mitragyna speciosa*. Dark orange powder. $[\alpha]_D^{25}$ -50 (c, 0.06 in CHCl_3). λ_{max} 258; 265; 363; 496 (MeOH).

Takayama, H. *et al.*, *Tetrahedron*, 1998, **54**, 8433-8440 (isol, uv, cd, pmr, cmr, ms)

Mitrilactonine**M-660**

$\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_4$ 364.4

(R)-form [223264-23-1]

Alkaloid from *Mitragyna speciosa*. Amorph. orange solid. $[\alpha]_D^{26}$ -16.5 (c, 0.17 in CHCl_3). λ_{max} 223 (log ϵ 4.24); 262 (sh) (log ϵ 3.79); 344 (log ϵ 3.78); 434 (sh) (log ϵ 4.37); 460 (log ϵ 4.49) (MeOH).

9-Methoxy-9-Methoxymitrilactonine

$\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_5$ 394.426


Alkaloid from *Mitragyna speciosa*. Dark orange powder. $[\alpha]_D^{25}$ -123 (c, 0.19 in CHCl_3) (natural). $[\alpha]_D$ -838 (c, 0.1 in CHCl_3) (synthetic). Partial racemate. λ_{max} 220 (log ϵ 4.21); 267 (sh); 340 (log ϵ 3.72); 435 (sh); 462 (log ϵ 4.51) (MeOH).

Takayama, H. *et al.*, *J.O.C.*, 1999, **64**, 1772-1773 (isol, synth, uv, pmr, cmr, ms)

Takayama, H. *et al.*, *Tetrahedron*, 2000, **56**, 3145-3151 (9-Methoxymitrilactonine)

Mitraphylline**M-661**

Ajmalicine oxindole B. Rubradimine [509-80-8]



Absolute Configuration

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Mitragyna hirsuta*, some other *Mitragyna* spp. and some *Uncaria* spp. (Rubiaceae). Hypotensive agent, general depressant. Mp 275-276° (265-266°). $[\alpha]_D$ -7 (CHCl_3) (-3). $[\alpha]_D$ -10 (EtOH). $[\alpha]_D$ +11 (Py). $[\alpha]_D^{24}$ -3 (c, 1.3 in CHCl_3). λ_{max} 243 (log ϵ 4.22); 280 (sh) (log ϵ 3.18) (EtOH).

Perchlorate: Mp 240°.

Picrate: Mp 207-209°.

N-Oxide: Mitraphylline N-oxide

[51920-99-1]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Alkaloid from *Uncaria orientalis* and *Uncaria longiflora* (Rubiaceae).

16 ξ ,17-Dihydro, 17 β -hydroxy: 16,17-Dihydro-17-hydroxymitraphylline

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_5$ 386.447

Alkaloid from the leaves of *Mitragyna parvifolia*. Viscous solid. λ_{max} 209; 256

; 286; 321 (no solvent reported).

Parent acid: Mitraphyllic acid

[10126-00-8]

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4$ 354.405

Alkaloid from stems of *Uncaria sinensis* (Rubiaceae). Mp 234-236° dec. $[\alpha]_D$ -32.8 (c, 0.07 in MeOH). λ_{max} 242 (log ϵ 4.27); 280 (log ϵ 3.5) (EtOH).

Parent acid, β -D-glucopyranosyl ester:**Mitraphyllic acid glucopyranosyl ester**

[152696-77-0]

$\text{C}_{26}\text{H}_{32}\text{N}_2\text{O}_9$ 516.547

Alkaloid from leaves of *Uncaria sinensis* (Nauclaceae).

7-Epimer: Isomitraphylline. Ajmalicine oxindole A

[4963-01-3]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Mitragyna* and *Uncaria* spp. (Rubiaceae). Amorph. $[\alpha]_D$ +18 (CHCl_3). pK_a 4.6.

7-Epimer, picrate:

Yellow prisms (MeOH). Mp 223° dec.

7-Epimer, N-oxide: Isomitraphylline N-oxide

[51920-98-0]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Alkaloid from *Mitragyna hirsuta*, *Uncaria longiflora* and *Uncaria orientalis*.

7-Epimer, 16 ξ ,17-dihydro, 17 β -hydroxy: 16,17-Dihydro-17-hydroxyisomitraphylline

[51920-98-0]

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_5$ 386.447

Alkaloid from the leaves of *Mitragyna parvifolia*. Viscous solid. λ_{max} 211; 256; 286; 322 (no solvent reported).

7-Epimer, parent acid: Isomitraphyllic acid

[152887-08-6]

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4$ 354.405

From leaves of *Uncaria sinensis* (Nauclaceae).

7-Epimer, parent acid, β -D-glucopyranosyl ester: Isomitraphyllic acid β -D-glucopyranosyl ester

[152595-58-9]

$\text{C}_{26}\text{H}_{32}\text{N}_2\text{O}_9$ 516.547

From leaves of *Uncaria sinensis* (Nauclaceae). Mp 206-209°.

19-Epimer: Uncarine B. Formosanine

[6883-35-8]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Uncaria* spp. and *Ouroparia formosana* (preferred genus name *Uncaria*) (Rubiaceae). Plates or needles. Mp 215-216°. $[\alpha]_D$ +91 (CHCl_3).

19-Epimer, hydrochloride: Mp 227-228° dec.**20-Epimer: Uncarine C. Isospeciophylline. Pteropodine**

[5629-60-7]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Mitragyna* and *Uncaria* spp. (Rubiaceae). Fine needles (C_6H_6). Mp 212-213°. $[\alpha]_D$ -109 (CHCl_3).

20-Epimer, picrate: Mp 143-144°.**20-Epimer, N-oxide: Pteropodine N-oxide**

[40242-13-5]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Isol. from *Uncaria bernaysii*, *Uncaria orientalis* and *Uncaria longiflora* (Rubiaceae). Identified by tlc only.

20-Epimer, parent acid: Maruquine. Pteropodic acid

[13897-75-1]

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4$ 354.405

Alkaloid from the leaves of *Hamelia patens* and stems of *Uncaria sinensis*. Cryst. (EtOH). Mp 227-229° dec. (Pteropodic acid) Mp 245-247° (Maruquine). $[\alpha]_D$ -126 (c, 0.1 in MeOH) (Pteropodic acid). $[\alpha]_D^{20}$ +247 (c, 0.16 in EtOH) (Maruquine). The radically different opt rotns. reported indicate that the two isolates are different. λ_{max} 244 (log ϵ 4.24); 280 (log ϵ 3.24) (EtOH) (Pteropodic acid).

3,20-Diepimer: Uncarine F

[14019-66-0]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Mitragyna* and *Uncaria* spp. (Rubiaceae). Amorph. $[\alpha]_D$ +85 (CHCl_3).

3,20-Diepimer, N-oxide: Uncarine F N-oxide

[40242-12-4]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Isol. from *Mitragyna parvifolia*, *Uncaria bernaysii* and *Uncaria orientalis* (Rubiaceae). Identified by tlc only.

7,19-Diepimer: Uncarine A. Isoformosanine

[6899-73-6]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Uncaria* spp. and *Ouroparia formosana* (Rubiaceae). Amorph. Mp 120-130°. $[\alpha]_D$ +106.5.

7,19-Diepimer, hydrochloride:

Needles + $\frac{1}{2}$ H_2O . Mp 231° dec.

7,20-Diepimer: Uncarine E. Isopteropodine

[5171-37-9]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Mitragyna parvifolia* and *Uncaria* spp. (Rubiaceae). Also isol. from the marine mollusc *Nerita albicilla*. Needles (C_6H_6 or MeOH). Mp 209-211°. $[\alpha]_D$ -111 (CHCl_3).

7,20-Diepimer, hydrochloride: Mp 181-183°. $[\alpha]_D$ -124.**7,20-Diepimer, N-oxide: Isopteropodine N-oxide**

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Isol. from *Uncaria bernaysii*, *Uncaria orientalis* and *Uncaria longiflora* (Rubiaceae). Identified by tlc only.

7,20-Diepimer, parent acid: Isomaruquine. Isopteropodic acid

[13897-74-0]

$\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4$ 354.405

Alkaloid from the leaves of *Hamelia patens* and stems of *Uncaria sinensis*. Cryst. (hexanol). Mp 223-224° dec. (Isopteropodic acid) Mp 188-189° (Isomaruquine). $[\alpha]_D$ -130.2 (c, 0.04 in MeOH) (Isopteropodic acid). $[\alpha]_D^{20}$ -78.9 (c, 0.136 in CHCl_3) (Isomaruquine). Props. refer to Isopteropodic acid. The two isolates not compared. λ_{max} 242 (log ϵ 4.03); 280 (log ϵ 3.11) (MeOH) (Isopteropodic acid).

- 3,7,20-Triepimer: **Uncarine D. Speciohylline**
[4697-68-1]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from *Mitragyna* and *Uncaria* spp. Needles (EtOAc or Et₂O). Mp 183-184°. [α]_D²¹ +91 (c, 0.21 in CHCl₃).
- 3,7,20-Triepimer, picrate:
Cryst. (EtOH). Mp 215°.
- 3,7,20-Triepimer, N-oxide: **Speciophylline N-oxide**
[40242-11-3]
C₂₁H₂₄N₂O₅ 384.431
Alkaloid from *Uncaria bernaysii*, *Uncaria orientalis*, *Uncaria longiflora* and *Mitragyna parvifolia* (Rubiaceae). Identified by tlc only.
- 7,19,20-Triepimer: **Rauniticine oxindole A**
[17507-91-4]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from the leaves of *Uncaria elliptica* (Rubiaceae). λ_{max} 212 ; 242 ; 290 (MeOH).
- Stereoisomer (1): Gambirdine**
[19775-85-0]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from *Uncaria gambir* (Rubiaceae). Mp 199-201°. [α]_D²¹ +84.8 (c, 0.02 in CHCl₃). Stereochem. unknown. λ_{max} 244 (log ε 4.19); 280 (sh) (log ε 3.13) (EtOH).
- Stereoisomer (2): Isogambirdine**
[19775-86-1]
C₂₁H₂₄N₂O₄ 368.432
Alkaloid from *Uncaria gambir* (Rubiaceae). Oil. Mp 179-181° dec. (as hydrochloride). [α]_D²¹ +115.5 (c, 0.02 in CHCl₃). Stereochem. unknown. λ_{max} 244 (log ε 4.24); 280 (sh) (log ε 3.18) (EtOH).
- Stereoisomer, 21-oxo: Hamelia patens Alkaloid A**
[82564-37-2]
C₂₁H₂₂N₂O₅ 382.415
Alkaloid from the aerial parts of *Hamelia patens* (Rubiaceae). Mp 210-212°. λ_{max} 209 ; 242 (sh) ; 280 (EtOH).
- Seaton, J.C. et al., *Can. J. Chem.*, 1958, **36**, 1031-1038 (*Mitraphylline*)
- Finch, N. et al., *J.A.C.S.*, 1962, **84**, 3871-3877 (*Isomitraphylline*)
- Gilbert, B. et al., *J.A.C.S.*, 1963, **85**, 1523-1528 (ms)
- Chan, K.C. et al., *J.C.S. (C)*, 1966, 2245-2249 (*Pteropodine*, *Isopteropodine*, *isol*, *pmr*, *ir*)
- Johns, S.R. et al., *Tet. Lett.*, 1966, 4883-4888 (*pmr*, *config*)
- Shamma, M. et al., *J.A.C.S.*, 1967, **89**, 1739-1740 (*stereochem*, *pmr*)
- Beecham, A.F. et al., *Aust. J. Chem.*, 1968, **21**, 491-504 (*Uncarines A-F*, *stereochem*)
- Chan, K.C. et al., *Tet. Lett.*, 1968, 3403-3406 (*Gambirdine*, *Isogambirdine*)
- Winterfeldt, E. et al., *Chem. Ber.*, 1969, **102**, 3558-3572 (*Formosanine*, *synth*)
- Phillipson, J.D. et al., *Phytochemistry*, 1973, **12**, 1481; 2791-2794; 1975, **14**, 1855-1863 (*oxides*)
- Shellard, E.J. et al., *Planta Med.*, 1974, **25**, 172-174 (*Speciophylline N-oxide*, *Uncarine F N-oxide*)
- Phillipson, J.D. et al., *J. Chromatogr.*, 1975, **105**, 163-178 (*tlc*, *glc*, *ms*)
- Ban, Y. et al., *Chem. Pharm. Bull.*, 1976, **24**, 736-751 (*synth*)

- Borges del Castillo, J. et al., *An. Quim., Ser. C*, 1980, **76**, 294-295; 1982, **78**, 126-128; 180-183; *CA*, **94**, 171040h; **97**, 39200t; 141710w (*Maruquine*, *Isomaruquine*, *Hamelia patens Alkaloid A*)
- Phillipson, J.D. et al., *Phytochemistry*, 1983, **22**, 1809-1813 (*Rauniticine oxindole A*)
- Martin, G.E. et al., *J. Nat. Prod.*, 1986, **49**, 406-411 (*Isopteropodine*, *isol*, *pmr*, *cmr*)
- Seki, H. et al., *Chem. Pharm. Bull.*, 1993, **41**, 2077-2086 (*stereoisomers*, *pmr*, *cmr*, *conformn*)
- Liu, H.-M. et al., *Phytochemistry*, 1993, **33**, 707-710 (*Mitraphyllin acid*, *Pteropodic acid*, *Isopteropodic acid*)
- Liu, H.-M. et al., *Yaoxue Xuebao*, 1993, **28**, 849; *CA*, **120**, 102001q (*Mitraphyllin acid*, *Isomitraphyllin acid*, *glucosides*)
- Laus, G. et al., *Helv. Chim. Acta*, 2003, **86**, 181-187 (*Mitraphylline*, *Speciophylline*, *cryst struct*)
- García, R. et al., *Z. Naturforsch., C*, 2005, **60**, 385-388 (*Uncarine E*, *cmr*, *ms*)
- Pandey, R. et al., *Phytochemistry*, 2006, **67**, 2164-2169 (*Dihydrohydroxymitraphylline*, *Dihydrohydroxymitraphylline*)
- Kitajima, M. et al., *J. Nat. Med. (Tokyo)*, 2007, **61**, 192-195 (*Isomitraphylline N-oxide*)

Mitraspicine


M-662

- C₂₈H₃₆N₂O₅ 480.603
Struct. unknown. Alkaloid from *Mitragyna speciosa* bark (Rubiaceae). Platelets (MeOH). Mp 244-245°. [α]_D²³ -59.15.
Conts. 3 OMe groups.
Picrate: Mp 136°.
Denis, P. et al., *Bull. Cl. Sci., Acad. R. Belg.*, 1938, **24**, 653-658; *CA*, **33**, 1741 (*isol*)
Raymond-Hamet, M. et al., *Ann. Pharm. Fr.*, 1950, **8**, 482-490

Mitrasulgynine

M-663

[212264-12-5]



Absolute Configuration

- C₂₃H₂₈N₂O₇S 476.549
Alkaloid from the leaves of *Mitragyna speciosa*. Yellow powder. [α]_D¹⁸ +413 (c, 0.09 in CHCl₃). Abs. config. proposed on biogenetic grounds. λ_{max} 241 ; 359 ; 438 (MeOH).
Takayama, H. et al., *Tetrahedron*, 1998, **54**, 8433-8440 (*isol*, *uv*, *cd*, *pmr*, *cmr*, *ms*)

Mitraversine


M-664

[1400-34-6]

- C₂₂H₂₆N₂O₄ 382.458
Struct. unknown. Alkaloid from *Mitragyna diversifolia* (Rubiaceae). Mp 237°
Mp 263.5-264.5°. Conts. 2 OMe groups.
Hydrochloride:
Leaflets. Mp 208-210°.
Field, E. et al., *J.C.S.*, 1921, **119**, 887-891
Raymond-Hamet, et al., *J. Pharm. Chim.*, 1937, **25**, 391-398; *Analyst (London)*, 1937, **62**, 472

Miyaconitine

M-665

2-(Acetyloxy)-7,9-dihydroxy-21-methyl-6,21-secohetisan-6,13-dione, 9CI
[28330-59-8]

- C₂₃H₂₉NO₆ 415.485
Alkaloid from *Aconitum miyabei* (Ranunculaceae). Mp 218° dec. [α]_D -87.8 (CHCl₃).

7-Ketone: Miyaconitnone

[28604-00-4]

C₂₃H₂₇NO₆ 413.469Alkaloid from *Aconitum miyabei* (Ranunculaceae). Mp 285° dec. [α]_D -27.6 (AcOH).**9-Deoxy, 7-ketone: Vilmorrianone. Delphatamine**

[135626-83-4]


C₂₃H₂₇NO₅ 397.47Alkaloid from the roots of *Aconitum vilmorrianum* and *Delphinium denudatum*. Cryst. (MeOH). Mp 253-255°. [α]_D -22 (c, 0.2 in MeOH).

[36059-80-0]

- Kakimoto, S. et al., *Bull. Chem. Soc. Jpn.*, 1959, **32**, 1153 (*uv*, *ir*, *struct*)
Ichinohe, Y. et al., *Tet. Lett.*, 1970, 2323 (*uv*, *ir*, *pmr*, *struct*)
Shimanouchi, H. et al., *Tet. Lett.*, 1970, 2327 (*cryst struct*)
Ding, L. et al., *Planta Med.*, 1991, **57**, 275-277 (*Vilmorrianone*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)
Atta-ur-Rahman, et al., *Turk. J. Chem.*, 1997, **21**, 13-20 (*Delphatamine*)

Modestanine

M-666

Methyl 2,3-didehydro-6,21-epoxyaspido-permidine-3-carboxylate. Deoxoapodine. Demethoxyvandrikine
[57932-88-4]

Absolute Configuration

- C₂₁H₂₄N₂O₃ 352.432
Alkaloid from *Hazunta modesta* and *Tabernaemontana armeniaca* (Apocynaceae). Amorph.

18R-Hydroxy: Hedrantherine

[38739-18-3]

C₂₁H₂₄N₂O₄ 368.432Alkaloid from the leaves of *Hedranthera barteri* (Apocynaceae). Noncryst. [α]_D²⁴ -459 (c, 1 in DMF). [α]_D²⁴ -484 (c, 0.81 in DMSO).**11,18R-Dihydroxy: 11-Hydroxyhe-**

dranthine. 17-Hydroxyhedranthine
[38741-26-3]
C₂₁H₂₄N₂O₅ 384.431
Alkaloid from the leaves of *Hedranthera barteri* (Apocynaceae).
Cryst. (CHCl₃/Me₂CO). Mp 245° dec.

18-Oxo (lactone): Apodine

[57932-86-2]
C₂₁H₂₂N₂O₄ 366.416
Alkaloid from *Tabernaemontana armeniaca* (Apocynaceae). Noncryst.
18-Oxo, hydrochloride: Mp 176° dec. [α]_D²² -520 (c, 2 in MeOH).

18-Oxo (lactone), 14ξ-hydroxy: Apodinine

[77017-02-8]
C₂₁H₂₂N₂O₅ 382.415
Alkaloid from *Tabernaemontana apoda*.

11-Methoxy: Vandrikine

[50656-91-2]
C₂₂H₂₆N₂O₄ 382.458
Minor alkaloid from a number of *Aspidosperma* spp. (Apocynaceae).

11-Methoxy, 18R-hydroxy: [38741-28-5]

Noncryst. [α]_D²² -502 (c, 1.26 in DMSO).
11-Methoxy, 19ξ-hydroxy: 19-Hydroxyvandrikine. 20-Hydroxyvandrikine
[76989-82-7]
C₂₂H₂₆N₂O₅ 398.458
Alkaloid from the root bark of *Craspidospermum verticillatum* (Apocynaceae). Mp 214°. [α]_D²⁰ -388 (c, 1 in CHCl₃).

Naranjo, J. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 1849 (*Hedrantherine*, 11-Hydroxyhedrantherine)

Wenkert, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4990 (*Vandrikine*)

Iglesias, R. *et al.*, *Rev. CENIC, Cienc. Fis.*, 1975, **6**, 135; 141; *CA*, **84**, 44502d; 44503e (*Apodine*)

Iglesias, R. *et al.*, *Rev. CENIC, Cienc. Fis.*, 1979, **10**, 357-362; *CA*, **94**, 136123 (*Apodinine*)

Lounasmaa, M. *et al.*, *Acta Chem. Scand., Ser. B*, 1980, **34**, 379 (*19-Hydroxyvandrikine*)


Bui, A.M. *et al.*, *Phytochemistry*, 1980, **19**, 1473 (*Modestanine*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1991, **113**, 2598 (*Apodine*, *Deoxoapodine*, *synth*)

Moenjodaramine

M-667

[36357-45-6]



C₂₈H₄₆N₂O 426.684

It appears that these alkaloids all have the 4S-config. shown, although the structs. of Harappamine, N-Formylharappamine and Papillamidine were publ.

as 4R- and have not since been amended. The C-28 oxygenation pattern (4S-) is usual for steroidal alkaloids (Sangare *et al.*). Alkaloid from *Buxus papillosa* and *Buxus hyrcana*. Acetylcholinesterase inhibitor. Mp 177°. [α]_D +33.3 (CHCl₃). [α]_D +61 (synthetic).

N²⁰-De-Me: Harappamine

[84679-85-6]
C₂₇H₄₄N₂O 412.657
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Gum.

N²⁰-De-Me, N²⁰-formyl: N-Formylharappamine

[115713-24-1]
C₂₈H₄₄N₂O₂ 440.668
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. [α]_D +40 (c, 0.50 in CHCl₃). λ_{max} 225 (log ε 3.96); 238 (log ε 4.04); 245 (log ε 4.03); 254 (log ε 3.9) (MeOH).

N³,N²⁰-Di-de-Me: Papilinine

[96910-88-2]
C₂₆H₄₂N₂O 398.631
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Gum. [α]_D +29.4 (CHCl₃). C-20-Config. not determined but almost certainly as given.

N²⁰,N²⁰-Di-de-Me: N^b-Demethylharappamine

[145385-76-8]
C₂₆H₄₂N₂O 398.631
Alkaloid from roots of *Buxus papillosa* (Buxaceae). Amorph. solid. [α]_D²⁰ +12.4 (CHCl₃). λ_{max} 238 ; 246 (MeOH).

N²⁰,N²⁰-Di-de-Me, N²⁰-formyl: Papillamidine. N-Formyl-N-demethylharappamine

[113689-37-5]
C₂₇H₄₂N₂O₂ 426.641
Alkaloid from the leaves of *Buxus papillosa* (Buxaceae). Amorph. [α]_D 0 (CHCl₃). λ_{max} 228 ; 238 ; 246 ; 254 (MeOH).

Khuong-Huu, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 558-560 (*Moenjodaramine*, *synth*, *pmr*)

Sangare, M. *et al.*, *Tet. Lett.*, 1975, **16**, 1791-1794 (*stereochem*)

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1983, **49**, 126 (*Moenjodaramine*)

Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 1984, **39**, 524-527; 1985, **40**, 565-566 (*Harappamine*, *Papilinine*)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1988, **27**, 89-92 (*Papillamidine*)


Choudhary, M.I. *et al.*, *Phytochemistry*, 1988, **27**, 1561-1562 (*N-Formylharappamine*)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1992, **55**, 1063-1066 (*N^b-Demethylharappamine*)

Moiramide A

M-668

[124731-96-0]



C₁₇H₁₉NO₃ 285.342

Metab. of a marine *Pseudomonas fluorescens*. Amorph. solid. Sol. MeOH, EtOAc. λ_{max} 294 (ε 27421) (MeOH). λ_{max}


293 (ε 27421) (MeOH) (Berdy).

Needham, J. *et al.*, *J.O.C.*, 1994, **59**, 2059-2063 (*Moiramide A*)

Moiramide B

M-669

[155233-31-1]



C₂₅H₃₁N₃O₅ 453.537

Metab. of a marine *Pseudomonas fluorescens*. Amorph. solid. Similar to Andrimide, A-1004.


Needham, J. *et al.*, *J.O.C.*, 1994, **59**, 2059-2063 (*Moiramide B*)

Davies, S.G. *et al.*, *J.C.S. Perkin 1*, 1998, 2635-2643 (*synth*)

Mollenine A

M-670

[205693-72-7]



C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Eupenicillium molle* NRRL13062. Exhibits moderate antibacterial and cytotoxic properties. Oil. [α]_D -410 (c, 1 in CH₂Cl₂). λ_{max} 240 (ε 4900); 295 (ε 1700) (CH₂Cl₂).

N⁷-Formyl: Mollenine B


[205692-57-5]
C₂₃H₂₈N₂O₄ 396.485
Alkaloid from *Eupenicillium molle* NRRL13062. Oil. [α]_D -180 (c, 1 in CH₂Cl₂). λ_{max} 251 (ε 9000); 282 (ε 2000) (CH₂Cl₂).

Wang, H.-J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 804-807 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Molliclavine

M-671

9-Hydroxyelymoclavine
[6858-92-0]



C₁₆H₁₈N₂O₂ 270.33

Tentative struct.; seems improbable (an enol). Alkaloid from ergot (*Claviceps purpurea*), *Argyria nervosa*, *Rivea cor-*

ymbosa and *Ipomoea muelleri* (Convolvulaceae). Mp 253° dec. $[\alpha]_D^{15} +30$ (c, 0.2 in Py).


Abe, M. *et al.*, *Nippon Nogeï Kagaku Kaishi*, 1959, **33**, 249-252; *CA*, **59**, 2879d (*isol, struct*)

Chao, J.-M. *et al.*, *J. Polym. Sci.*, 1973, **62**, 588-591 (*occur*)

Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435-2440 (*occur*)

Mollinedine M-672

5-(1,3-Benzodioxol-5-ylmethyl)-1,3-dioxolo[4,5-g]isoquinoline, 9CI. 6,7-Methylenedioxy-1-(3,4-methylenedioxybenzyl)-isoquinoline
[16658-48-3]



$C_{18}H_{13}NO_4$ 307.305

Alkaloid from the wood of *Mollinedia costaricensis* (Monimiaceae). Needles (C_6H_6 or MeOH). Mp 170-172° (158-160°).

N-Me: Escholamine

[30331-75-0]
 $C_{19}H_{16}NO_4^{\oplus}$ 322.34

Quaternary alkaloid from the above-ground parts of *Eschscholtzia* sp. (?*Eschscholtzia oregana*) (Papaveraceae). Pale yellow needles (H_2O) (as iodide). Mp 265-266° (iodide).

Buck, J.S. *et al.*, *J.C.S.*, 1925, 1462 (*synth*)
Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1949, **23**, 1880 (*synth*)

Slavková, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1966, **31**, 3362 (*Escholamine, isol, uv, ir, struct*)


Birch, A.J. *et al.*, *J.C.S. Perkin I*, 1974, 2190 (*Escholamine, synth, pmr*)

López, J.A. *et al.*, *J. Nat. Prod.*, 1988, **51**, 754 (*isol, uv, ir, pmr, cmr, ms, synth*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*Escholamine, N-15 nmr*)

Molliorin A M-673

[63693-18-5]




$C_{32}H_{49}NO_2$ 479.745

Constit. of the marine sponge *Cacospongia mollior*. Mp 102-105°. $[\alpha]_D +51.7$ (c, 1 in $CHCl_3$).

Cafieri, F. *et al.*, *Tet. Lett.*, 1977, 477 (*isol, ir, pmr, uv, ms, struct*)

Molliorin B M-674

[64421-20-1]



$C_{58}H_{84}N_2O_4$ 873.313

Constit. of the marine sponge *Cacospongia mollior*. Cryst. (petrol). Mp 173-174°. $[\alpha]_D +14.6$ (c, 1.2 in $CHCl_3$).

Cafieri, F. *et al.*, *Experientia*, 1977, **33**, 994 (*isol, uv, synth, struct*)

Molliorin C M-675

[66648-53-1]

As Molliorin A, M-673 with

R = $-CH_2CH(CH_3)_2$

$C_{31}H_{47}NO_2$ 465.718

Minor alkaloid from the marine sponge *Cacospongia mollior*. Amorph. solid. $[\alpha]_D -46.9$ (c, 0.9 in $CHCl_3$). λ_{max} 257 (ε 11900) (EtOH).

Cafieri, F. *et al.*, *Experientia*, 1978, **34**, 300 (*isol, ms, pmr, synth*)

Molliorin D M-676

[70022-71-8]

As Molliorin A, M-673 with

R = $-CH_2CH_2Ph$

$C_{35}H_{47}NO_2$ 513.762

Minor alkaloid from the marine sponge *Cacospongia mollior*. $[\alpha]_D +11.6$. λ_{max} 262 (ε 11760) (EtOH).

Cafieri, F. *et al.*, *Experientia*, 1979, **35**, 157 (*isol, struct, ir, pmr, ms, synth*)

Molliorin E M-677

[69819-82-5]

As Molliorin A, M-673 with

R = Me


$C_{28}H_{41}NO_2$ 423.637

Minor alkaloid from the marine sponge *Cacospongia mollior*. $[\alpha]_D +6.5$. λ_{max} 263 (ε 11900) (EtOH).

Cafieri, F. *et al.*, *Experientia*, 1979, **35**, 157 (*isol, struct, ir, pmr, ms, synth*)

Mollisine† M-678

[160568-08-1]



$C_{19}H_{13}NO_5$ 335.315

Alkaloid from bark of *Alphonsea mollis* (Annonaceae).

8-O-De-Me: 8-O-Demethylmollisine

$C_{18}H_{11}NO_5$ 321.289


Alkaloid from *Alphonsea mollis*. Orange needles. Mp 264-267° (dec.).

Xie, N. *et al.*, *Zhongguo Yaoke Daxue Xuebao*, 1994, **25**, 205; *CA*, **122**, 101554b

Xie, N. *et al.*, *Chin. Chem. Lett.*, 1999, **10**, 675-676 (8-O-Demethylmollisine)

Molokaiaimine M-679

4-(3-Aminopropoxy)-3,5-dibromobenzeneethanamine, 9CI
[151345-09-4]



$C_{11}H_{16}Br_2N_2O$ 352.068

Metab. from the marine sponge *Pseudoceratina purpurea*, also from *Psammaphysilla* sp. and *Aplysinella* sp. Cytotoxic. Off-white powder (MeOH) (as dihydrochloride). λ_{max} 206 (ε 28000); 284 (ε 600) (MeOH).

N^{3''}- (Cyanofornyl): Ceratinamine. Pseudoceramine

[175993-07-4]

$C_{13}H_{15}Br_2N_3O_2$ 405.088

Metab. from *Pseudoceratina purpurea* and *Aplysinella* sp. Cytotoxic against P388 murine leukaemia cells. Anti-fouling substance. Solid (presumably as hydrochloride). Sol. MeOH. First report of a cyanofornamide nat. prod. λ_{max} 207 (ε 37800); 276 (ε 850) (MeOH) (Berdy).

N^{2'}-Me: N'-Methylmolokaiaimine

$C_{12}H_{18}Br_2N_2O$ 366.095

Isol. from a *Hexadella* sp. Powder. λ_{max} 277 (ε 340); 284 (ε 330) (MeOH).

N^{2'}-Me, N^{3''}- (cyanofornyl): N-Methylceratinamine

[256448-59-6]

$C_{14}H_{17}Br_2N_3O_2$ 419.115

Metab. of an undescribed Verongid sponge. λ_{max} 278 (log ε 3.76); 285 (log ε 3.77) (MeOH).

N^{3''}-Me, N^{2'}-methoxycarbonyl:

$C_{14}H_{20}Br_2N_2O_3$ 424.132

Alkaloid from *Psammaphysilla purpurea*. Brown semisolid. λ_{max} 250 (ε 2210) (MeOH).

N^{3''}, N^{3''}-Di-Me: Purplealidin E

[260437-59-0]

$C_{13}H_{20}Br_2N_2O$ 380.122

Metab. of the sponge *Psammaphysilla purpurea*. Oil. Mp 148° (as 2'-Ac, hydrochloride). Not indexed in *CA* **141**. λ_{max} 223 (ε 5770); 275 (ε 950); 280 (ε 944) (MeOH) (2'-Ac, hydrochloride).

N^{3''}, N^{3''}-Di-Me, N^{2'}-Ac: [215942-07-7]

$C_{15}H_{22}Br_2N_2O_2$ 422.159

Isol. from *Psammaphysilla purpurea*.

$N^{3''}, N^{3''}$ -Di-Me, $N^{2''}$ -methoxycarbonyl: *Methyl 3,5-dibromo-4-[[3-(dimethylamino)propoxy]phenylethyl]carbamate* $C_{15}H_{22}Br_2N_2O_3$ 438.158
Alkaloid from *Psammaphysilla purpurea*. Brown solid. Mp 160-162°. Also isol. as the monohydrochloride, Mp 170-173°. Not indexed in CA 141.

$N^{2''}, N^{2''}, N^{3''}$ -Tri-Me: **Purealidin F**
[145205-29-4]
 $C_{14}H_{22}Br_2N_2O$ 394.149
Alkaloid from the Okinawan sponge *Psammaphysilla purpurea*. Inhibitor of Na/K-ATPase. Oil (as dihydrochloride). Sol. MeOH. CAS no. refers to hydrochloride. λ_{max} 218 (€ 12000); 274 (€ 800); 285 (€ 1000) (MeOH) (hydrochloride). λ_{max} 218 (€ 12000); 274 (€ 800); 285 (€ 1000) (MeOH) (Berdy).

$N^{2''}, N^{3''}, N^{3''}$ -Tri-Me: **Purealidin G**
[145196-30-1]
 $C_{14}H_{22}Br_2N_2O$ 394.149
Alkaloid from *Psammaphysilla purpurea*. Na/K-ATPase inhibitor. Sol. MeOH. CAS no. refers to dihydrochloride. λ_{max} 214 (€ 14000); 276 (€ 800); 288 (€ 1100) (MeOH) (hydrochloride).

$N^{2''}, N^{2''}, N^{3''}, N^{3''}$ -Tetra-Me: **Aplysamine I**
[159026-30-9]
[125547-38-8]
 $C_{15}H_{24}Br_2N_2O$ 408.175
Metab. of an Australian marine sponge *Aplysina* sp. Pale yellow semicryst. solid (as hydrochloride). Mp 122-123° (hydrochloride). Biol. inactive against several test bacteria and fungi.

$N^{2''}$ -Cyano, $N^{2''}$ -Me: **N'-Cyano-N'-methylmoloakaamine**
 $C_{13}H_{17}Br_2N_3O$ 391.105
Isol. from a *Hexadella* sp. Amorph. solid. λ_{max} 277 (€ 440); 284 (€ 440) (MeOH).

$1''$ -Hydroxy, $N^{3''}$ -(cyanocarbonyl): **7-Hydroxyceratinamine**
[230295-94-0]
 $C_{13}H_{15}Br_2N_3O_3$ 421.088
Isol. from *Aplysina* sp. Amorph. solid. $[\alpha]_D^{25} +3$ (c, 0.8 in MeOH).

$1''\xi$ -Hydroxy, $N^{3''}, N^{3''}$ -di-Me: **Purpurealidin F**
[799246-90-5]
 $C_{13}H_{20}Br_2N_2O_2$ 396.121
Isol. from *Psammaphysilla purpurea*. Oil. λ_{max} 277 (€ 925); 282 (€ 950) (MeOH).

$1''\xi$ -Hydroxy, $N^{3''}, N^{3''}$ -di-Me, $N^{2''}$ -propionyl: **Purpurealidin G**
[799246-91-6]
 $C_{16}H_{24}Br_2N_2O_3$ 452.185
Isol. from *Psammaphysilla purpurea*. Oil. λ_{max} 277 (€ 925); 282 (€ 950) (MeOH).

Kynas, R. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1427-1433 (*Aplysamine I*)
Tsuda, M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1325-1327 (*Purealidins*)
Hamann, M.T. *et al.*, *J.O.C.*, 1993, **58**, 6565-6569 (*isol, uv, pmr, cmr, ms, N-Methylceratinamine*)
Tsukamoto, S. *et al.*, *J.O.C.*, 1996, **61**, 2936-2937 (*Ceratinamine*)

Tsukamoto, S. *et al.*, *Tetrahedron*, 1996, **52**, 8181-8186 (*isol, props*)
Venkateswarlu, Y. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1388-1389; 1999, **62**, 893-894 (*Psammaphysilla purpurea constits*)
Schoenfeld, R.C. *et al.*, *Tet. Lett.*, 1998, **39**, 4147-4150 (*synth*)
Rao, M.R. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 1301-1303 (*N-di-Me-N^{2''}-Ac*)
Fu, X. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1072-1073 (*7-Hydroxyceratinamine*)
Lacy, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 119-121 (*N-Methylceratinamine*)
Schoenfeld, R.C. *et al.*, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 823-825 (*synth*)
Tilvi, S. *et al.*, *Tetrahedron*, 2004, **60**, 10207-10215 (*Purpurealidin A-H*)
Ravinder, K. *et al.*, *ARKIVOC*, 2005, **iii**, 51-55 (*N-Me N'-methoxycarbonyl*)
Matsunaga, S. *et al.*, *J.O.C.*, 2005, **70**, 1893-1896 (*N'-Methylmoloakaamine, N'-Cyano-N'-methylmoloakaamine*)
Yoshida, M. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 1362-1363 (*Aplysamine I, Purpurealidin E, synth*)


Mololipids

A series of fatty acid lipids with a core containing Moloakaamine, M-679. Isol. from a Hawaiian sponge of the order Verongidae. Active against HIV-1. Waxy solid. λ_{max} 240 (€ 1900); 277 (€ 650); 284 (€ 620) (MeOH).

Ross, S.A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 501-503 (*isol*)

Molybdenum cofactor

Molybdoenzyme molybdenum cofactor, 9CI. Molybdopterin. MOCO



Molybdenum cofactor can be released from molybdoenzymes by treatment with denaturing reagents. Catalyses oxidative hydroxylation or reductive dehydroxylation at C, N, or S atoms. During these redox reactions Mo undergoes changes between oxidn. states of Mo^{VI} and Mo^{IV} , possibly linked to changes in the pyrazine ring oxidn. level. Molybdenum cofactor functions as a prosthetic metal in the 12 or so Mo-containing enzymes discovered, except nitrogenase which has its own FebMo cofactor. Air-sensitive ($t_{1/2}$ 15 min, 3.5°). Oxidative degradation involves labilisation of the metal and oxidn. of the reduced pterin. Stable under anaerobic conditions in presence of stabilising agents, e.g. sodium dithionite, or in DMSO (activity loss 3%/24 h at 20-25°).

Dioxomolybdate-form [104676-61-1]

[73508-07-3]

X = O. Isol. from purified milk xanthine oxidase. Molybdenum cofactor functions as a prosthetic metal in the 12 or so Mo-containing enzymes discovered, except

nitrogenase which has its own FebMo cofactor.

Stable at pH 5.9 under anaerobic conditions.


Oxothioxomolybdate-form [104676-62-2]

X = S.

Johnson, J.L. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1982, **79**, 6856-6860 (*isol, struct*)
Hawkes, T.R. *et al.*, *Biochem. J.*, 1984, **219**, 481-493 (*anal*)
Johnson, J.L. *et al.*, *J. Biol. Chem.*, 1984, **259**, 5414-5422 (*isol, props*)
Burgmayer, S.J.N. *et al.*, *J. Chem. Educ.*, 1985, **62**, 943 (*Dioxomolybdate, rev*)
Kramer, S.P. *et al.*, *J. Biol. Chem.*, 1987, **262**, 16357-16363 (*struct, props*)
Taylor, E.C. *et al.*, *J.O.C.*, 1988, **53**, 5839-5847 (*Dioxomolybdate, synth, bibl*)
Deistung, J. *et al.*, *Biochem. J.*, 1989, **263**, 477-483 (*isol, props, bibl*)
Taylor, E.C. *et al.*, *J.A.C.S.*, 1989, **111**, 285-291 (*Dioxomolybdate, synth, bibl*)
Larsen, L. *et al.*, *J.C.S. Perkin I*, 1989, 2311-2316; 2317-2327 (*Dioxomolybdate, synth, bibl*)
Rajagopalan, K.V. *et al.*, *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1991, **64**, 215-289 (*rev*)
Stiefel, E.I. *et al.*, *ACS Symp. Ser.*, 1993, **535**, (rev)
Taylor, E.C. *et al.*, *Adv. Exp. Med. Biol.*, 1993, **338**, 363-367 (*Dioxomolybdate, rev, synth*)
Goswami, S. *et al.*, *Heterocycles*, 1993, **35**, 1551-1570 (*Dioxomolybdate, rev*)
Bradshaw, B. *et al.*, *J.C.S. Perkin I*, 2001, 3232-3238; 3239-3244 (*Dioxomolybdate, synth*)
Metal Ions in Biology, Vol. 7: Molybdenum Enzymes, (ed. Spiro, T.G.), John Wiley, N.Y., 1985, (*Dioxomolybdate, book*)
Coughlan, M.P. *et al.*, *Molybdenum Molybdenum-Containing Enzymes*. 1980, Pergamon, N.Y., 1980, (*Dioxomolybdate, book*)

Monamidocin

M-682
N-(5-Guanidino-2-hydroxypentanoyl)phenylalanine
[171864-90-7]




$C_{15}H_{22}N_4O_4$ 322.363

Dipeptide antibiotic. Prod. by *Streptomyces* sp. NR 0637. Fibrinogen receptor antagonist, platelet aggregation inhibitor. Powder (as monohydrochloride). Sol. MeOH, H_2O , DMSO; poorly sol. hexane. $[\alpha]_D^{24} +21.5$ (c, 0.5 in MeOH) (HCl salt). Kamiyama, T. *et al.*, *J. Antibiot.*, 1995, **48**, 1221; 1226 (*isol, ir, pmr, cmr, synth*)

Monanchorin

[723335-95-3]




$C_{11}H_{21}N_3O$ 211.306

Isol. from the sponge *Monanchora unguiculata*. Light yellow oil. $[\alpha]_D^{25} +39$ (c, 3.9 in MeOH).

Meragelman, K.M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1165-1167 (*isol*, *pmr*, *cmr*)

Monascopyridine A **M-684**
[604786-54-1]



Absolute Configuration

R = $-\text{CO}(\text{CH}_2)_4\text{CH}_3$

$\text{C}_{21}\text{H}_{25}\text{NO}_4$ 355.433

Similar to Rubropunctamine, R-152. Prod. by *Monascus purpureus* DSM1379 and DSM1603 on rice. Needles (EtOH). Mp 161°.

Wild, D. *et al.*, *J. Agric. Food Chem.*, 2003, **51**, 5493-5496 (*isol*, *cd*, *pmr*, *cmr*, *ms*)

Monascopyridine B **M-685**

[604786-55-2]

As Monascopyridine A, M-684 with

R = $-\text{CO}(\text{CH}_2)_6\text{CH}_3$


$\text{C}_{23}\text{H}_{29}\text{NO}_4$ 383.486

Related to Monascorubramine, M-686. Prod. by *Monascus purpureus* DSM1379 and DSM1603 on rice. Needles (EtOH). Mp 124°.

Wild, D. *et al.*, *J. Agric. Food Chem.*, 2003, **51**, 5493-5496 (*isol*, *cd*, *pmr*, *cmr*, *ms*)

Monascorubramine **M-686**

9a-Methyl-3-(1-oxooctyl)-6-(1-propenyl)furo[3,2-g]isoquinoline-2,9(7H,9aH)-dione, 9CI. Monascamine [3627-51-8]



$\text{C}_{23}\text{H}_{27}\text{NO}_4$ 381.471

Fungal pigment of *Monascus* spp. Purple needles (C_6H_6). Mp 198° dec.

N-(1R-Carboxyethyl): [186649-31-0]

$\text{C}_{26}\text{H}_{31}\text{NO}_6$ 453.534

Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20} -3100$ (c, 0.003 in MeOH). λ_{max} 202 ; 248 ; 295 ; 420 ; 510 (MeOH).

N-(1S-Carboxyethyl): [186649-30-9]

$\text{C}_{26}\text{H}_{31}\text{NO}_6$ 453.534

Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20} -2900$ (c, 0.003 in MeOH). λ_{max} 202 ; 248 ; 292 ; 420 ; 498 (MeOH).

N-(1R,2-Dicarboxyethyl): [186649-35-4]

$\text{C}_{27}\text{H}_{31}\text{NO}_8$ 497.544

Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20} -2600$ (c, 0.003 in MeOH). λ_{max} 204 ; 250 ; 295 ; 418 ; 505 (MeOH).

N-(1S,2-Dicarboxyethyl): [186649-34-3]

$\text{C}_{27}\text{H}_{31}\text{NO}_8$ 497.544

Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20} -2000$ (c, 0.003 in MeOH). λ_{max} 204 ; 248 ; 278 ; 418 ; 490 (MeOH).

(Z)-Isomer, N-(2-hydroxyethyl): N-(2-Hydroxyethyl)monascorubramine. PP-R

[340014-04-2]

$\text{C}_{25}\text{H}_{31}\text{NO}_5$ 425.524

Prod. by *Penicillium* sp. AZ. Amorph. red solid.

Fielding, B.C. *et al.*, *J.C.S.*, 1961, 4579 (*isol*)

Nakanishi, K. *et al.*, *Tetrahedron*, 1962, **18**,

1171; 1185 (*struct*)


Sato, K. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**,

227-229 (*N-carboxyethyl derivs*)

Ogihara, J. *et al.*, *J. Biosci. Bioeng.*, 2001, **91**,

44-47 (*PP-R*)

Monocillinol A **M-687**



Relative Configuration

$\text{C}_{11}\text{H}_{13}\text{NO}_5$ 239.227

Prod. by a *Monocillium* sp. Amorph. solid. $[\alpha]_D -33.5$ (c, 0.02 in MeOH). λ_{max} 205 (log ϵ 4.08); 218 (log ϵ 4.14) (EtOH).

A⁷-Isomer: **Monocillinol B**


$\text{C}_{11}\text{H}_{13}\text{NO}_5$ 239.227

Prod. by a *Monocillium* sp. Amorph. solid. $[\alpha]_D +53$ (c, 0.02 in MeOH). λ_{max} 207 (log ϵ 4.06); 217 (log ϵ 4.01) (EtOH).

Biswas, M.H.U. *et al.*, *Tet. Lett.*, 2000, **41**, 7177-7180

Monocrotaline **M-688**

14,19-Dihydro-12,13-dihydroxy-20-nor-crotalanan-11,15-dione, 9CI [315-22-0]



Absolute configuration

$\text{C}_{16}\text{H}_{23}\text{NO}_6$ 325.361

Cyclic ester of retronecine in T-188 with Monocrotalic acid. Alkaloid from *Crotalaria retusa*, *Crotalaria spectabilis*, *Crotalaria aegyptiaca*, *Crotalaria burhia* and *Lindelofia spectabilis* (Fabaceae, Boraginaceae). Hepatotoxic. Causative agent of much seneciosis, e.g. accidental poisoning by *S.* by weed residues in bread, and characterised by venooculo-sive disease. Shows antineoplastic activity esp. vs. adenocarcinoma. Male insect sterilant. Used in China as an antineoplastic agent. Prisms (EtOH). Mp 202-

203° (197-198° dec.). $[\alpha]_D^{26} -54.7$ (CHCl_3). Log P -1.44 (uncertain value) (calc). λ_{max} 217 (ϵ 20500) (MeOH) (Berdy).

► Exp. carcinogen. Possible human carcinogen. LD₅₀ (rat, orl) 66 mg/kg. Hepatotoxic. QB3140000

Hydrochloride:

Prisms (MeOH/ CHCl_3). Mp 184° dec. $[\alpha]_D^{28} -38.4$ (H_2O).

Methiodide:

Prisms (MeOH/ CHCl_3). Mp 205° dec. $[\alpha]_D^{28} +23.4$ (c, 3.1 in MeOH).

N-Oxide: **Monocrotaline N-oxide**

[35337-98-5]

$\text{C}_{16}\text{H}_{23}\text{NO}_7$ 341.36

Isol. from *Crotalaria sagittalis* (Fabaceae).

► RC1295000

O¹³-Ac: **Spectabiline**†. *Acetylmonocrotaline*

[520-55-8]

$\text{C}_{18}\text{H}_{25}\text{NO}_7$ 367.398

Prisms (EtOH). Mp 185.5-186°. $[\alpha]_D^{20} +121$ (c, 1.7 in CHCl_3). $[\alpha]_D^{20} +143$ (c, 1.38 in EtOH). Spectabiline is also a synonym for Lemobiline, L-88.

► RC1260000

O¹³-(2-Methylbutanoyl): **Grahamine**†

[24583-56-0]

$\text{C}_{21}\text{H}_{31}\text{NO}_7$ 409.478

Alkaloid from *Crotalaria grahamiana* seeds (Fabaceae). Mp 163°. $[\alpha]_D^{20} +100$ (c, 1.05 in EtOH).

12,13-Cyclic acetaldehyde acetal: **Monocrotalinine**

[64595-67-1]

$\text{C}_{18}\text{H}_{25}\text{NO}_6$ 351.399

Alkaloid from *Crotalaria grahamiana* (Fabaceae). Cryst. (Et_2O). Mp 160-161° dec. $[\alpha]_D +125.2$ (EtOH). Prob. artifact.

3,7a-Didehydro: *Dehydromonocrotaline*.

Monocrotaline pyrrole

[23291-96-5]

$\text{C}_{16}\text{H}_{21}\text{NO}_6$ 323.345

Metab. of monocrotaline. Pneumotoxic agent. Needles (petrol). Mp 85-90° dec. $[\alpha]_D^{15} +109.4$ (c, 1.1 in EtOH).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 681B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1080C (*nmr*)

Neal, W.M. *et al.*, *J.A.C.S.*, 1935, **57**, 2560 (*isol*)

Adams, R. *et al.*, *J.A.C.S.*, 1939, **61**, 2815; 2819; 1950, **72**, 158 (*struct*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1957, **10**, 474; 1958, **11**, 97 (*Spectabiline*)

Neuner-Jehle, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 321 (*ms*)

Atal, C.K. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1773 (*Grahamine*)

Simanek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1970, **23**, 1853-1857 (*Dehydromonocrotaline*)

Robins, D.J. *et al.*, *J.C.S. (C)*, 1970, 1334; 1336 (*config*)


Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653 (*cd*)

Willette, R.E. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 122 (*Monocrotaline N-oxide*)

Robins, D.J. *et al.*, *J.C.S. Perkin 1*, 1974, 2082 (*biosynth*)

- IARC Monog.*, 1976, **10**, 291; *Suppl.* 7, 67 (rev. *tox*)
- Rajagopalan, T.R. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 455 (*Monocrotalinine*)
- Stoekli-Evans, H. *et al.*, *Acta Cryst. B*, 1979, **35**, 231 (*cryst struct*)
- Matsumoto, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 3329 (*synth, abs config*)
- Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417 (*cmr*)
- Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)
- Wang, S. *et al.*, *CA*, 1982, **97**, 19843o (*cryst struct, Spectabiline*)
- Molyneux, R.J. *et al.*, *Phytochemistry*, 1982, **21**, 439 (*cmr*)
- Mackay, M.F. *et al.*, *Acta Cryst. C*, 1984, **40**, 473-476 (*cryst struct, Dehydromonocrotaline*)
- Mattocks, A.R. *et al.*, *Chemistry and Toxicology of Pyrrolizidine Alkaloids*, 1986, (*tox, rev*)
- Vedejs, E. *et al.*, *J.O.C.*, 1987, **52**, 3937 (*synth*)
- Niwa, H. *et al.*, *Tet. Lett.*, 1988, **29**, 5139 (*synth*)
- Jackson, G.E. *et al.*, *Spectrosc. Lett.*, 1990, **23**, 971 (*pmr, cmr*)
- Niwa, H. *et al.*, *Tetrahedron*, 1992, **48**, 10531 (*synth*)
- Yan, C.C. *et al.*, *Toxicol. Appl. Pharmacol.*, 1994, **127**, 58-63 (*rev, tox*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MRH000

Monodontamide B **M-689**
[152273-62-6]



$C_{26}H_{32}N_4O_7$ 512.561
Alkaloid from the marine gastropod mollusc *Monodonta labio*. Serine protease inhibitor. Amorph. solid. λ_{max} 231 (ϵ 26600); 259 (ϵ 9830); 318 (ϵ 3930) (MeOH) (Berdy).

Deformyl: Monodontamide C


[152273-63-7]
 $C_{25}H_{32}N_4O_6$ 484.551
From *Monodonta labio*. Serine protease inhibitor. Amorph. solid. λ_{max} 226 (ϵ 32300); 256 (ϵ 6990); 281 (ϵ 3720); 365 (ϵ 6000) (MeOH) (Berdy).

21-Demethoxy, 22-deoxy: Monodontamide A

[152273-61-5]
 $C_{25}H_{30}N_4O_5$ 466.536
Alkaloid from *Monodonta labio*. Serine protease inhibitor. Amorph. solid. λ_{max} 229 (ϵ 15300); 258 (ϵ 6020); 318 (ϵ 2900) (MeOH) (Berdy).

Niwa, H. *et al.*, *Tetrahedron*, 1994, **50**, 6805-6818 (*isol, uv, ir, pmr, cmr, struct, synth*)

Monodontamide D **M-690**
[156401-06-8]




$C_{27}H_{31}N_5O_5$ 505.572
Alkaloid from the marine gastropod

mollusc *Monodonta labio*. Amorph. solid.

Deformyl: Monodontamide E

[156401-05-7]
 $C_{26}H_{31}N_5O_4$ 477.562
From *Monodonta labio*. Amorph. solid.
Niwa, H. *et al.*, *Tetrahedron*, 1994, **50**, 6805-6818 (*isol, uv, ir, pmr, cmr, struct, synth*)

Monodontamide F **M-691**
[156401-04-6]




$C_{25}H_{27}N_5O_5$ 477.519
Alkaloid from the marine gastropod mollusc *Monodonta labio*. Plates (MeOH). Mp 129-132°.

Niwa, H. *et al.*, *Tetrahedron*, 1994, **50**, 6805-6818 (*isol, uv, ir, pmr, cmr, struct, synth*)

Monodoroindeole **M-692**

6,6'-[3-Methyl-5-(2-methyl-1-propenyl)-2-pentene-1,5-diyl]bis-1H-indole, 9CI. 1,5-Di-6-indolyl-3,7-dimethyl-2,6-octadiene
[110892-13-2]




$C_{26}H_{28}N_2$ 368.521
Alkaloid from the seeds of *Monodora myristica* (Annonaceae).

Muhammad, I. *et al.*, *J. Bangladesh Acad. Sci.*, 1987, **11**, 1-7; *CA*, **107**, 172452t

Monogagaine, 9CI **M-693**

Alkaloid 6F. Alkaloid 20A
[98360-63-5]




$C_{39}H_{44}N_4O_2$ 600.802
Minor alkaloid from the root bark of *Tabernaemontana chippii* and from the stem bark of *Tabernaemontana dichotoma*. Also isol. from a callus culture of *Tabernaemontana elegans* (Apocynaceae).

Sol. MeOH; poorly sol. H_2O . λ_{max} 222 ; 288 (MeOH) (Berdy).

van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400 (*isol, uv, pmr, ms*)
Perera, P. *et al.*, *Phytochemistry*, 1985, **24**, 2097 (*isol, uv, ir, pmr, ms*)
van Beek, T.A. *et al.*, *Z. Naturforsch., B*, 1985, **40**, 693 (*uv, pmr, cmr, ms, cd, struct*)
van der Heijden, R. *et al.*, *Phytochemistry*, 1986, **25**, 843 (*isol*)

Monomargine **M-694**

6H-Azepino[3',2':4,5]cyclopent[1,2-g]indole-7,11-dione, 9CI
[148717-63-9]




$C_{15}H_8N_2O_2$ 248.24
Pigment from the stem bark of *Monocarpia marginalis*. Cytotoxic. Orange amorph. solid. λ_{max} 229 ; 445 (MeOH) (Berdy).

Mahmood, K. *et al.*, *Tet. Lett.*, 1993, **34**, 1795-1796 (*isol, uv, ir, pmr, cmr*)

Monspessulanine **M-695**

[3921-73-1]



$C_{15}H_{22}N_2O$ 246.352
Stereoisomer of Aphyllidine, A-1336. Alkaloid from *Cytisus monspessulanus* (Fabaceae). Cryst. (Me_2CO aq.). Mp 102°. $[\alpha]_D^{20} +168$ (c, 1 in EtOH). $[\alpha]_D +176$ (EtOH).

Perchlorate:

Cryst. (H_2O). Mp 213°.

Picrate:

Cryst. (EtOH). Mp 242° dec.


White, E.P. *et al.*, *N.Z. J. Sci. Technol., Sect. B*, 1946, **27**, 339; 1951, **33**, 54 (*isol*)

White, E.P. *et al.*, *J.C.S.*, 1964, 4613 (*ir, uv, pmr, struct*)

Bohlmann, F. *et al.*, *Tet. Lett.*, 1965, 2433 (*synth*)

Montabuphine **M-696**

[168611-77-6]




$C_{17}H_{19}NO_4$ 301.341
Structure has been shown to require revision. Alkaloid from bulbs of *Boo-*

phone flava. Mp 162-164°. $[\alpha]_D^{22} +157$ (c, 0.106 in EtOH).

Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 307-311 (*isol, ir, pmr, cmr, ms, cd, struct*)
Matveenko, M. *et al.*, *Org. Lett.*, 2008, **10**, 4693-4696 (*struct*)

Montamine **M-697**

[917098-78-3]




$C_{40}H_{38}N_4O_8$ 702.762

Dimer of Moschamine in H-755. Alkaloid from the seeds of *Centaurea montana*. Cytotoxic. Gum. λ_{max} 213 ; 271 ; 274 (MeOH).

Shoeb, M. *et al.*, *Tetrahedron*, 2006, **62**, 11172-11177 (*isol, pmr, cmr*)

Montiporyne E **M-698**

1,5,6,7-Tetrahydro-4-(1,3-undecadiynyl)-2H-azepin-2-one



$C_{17}H_{23}NO$ 257.375

Isol. from the stony coral *Montipora* sp. Cytotoxic agent. Pale yellow gum.


Bae, B.H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1511-1514

Collison, C.G. *et al.*, *Synthesis*, 2006, 2319-2322 (*synth, ir, pmr, cmr*)

Montipyridine **M-699**

1-Carboxymethyl-4-methoxy-2-(8-nonenyl)pyridinium (1+)

[352230-21-8]



$C_{17}H_{25}NO_3$ 291.389


Alkaloid from the stony coral *Montipora* sp. Light brown solid.

Alam, N. *et al.*, *J. Nat. Prod.*, 2001, **64**, 956-957 (*isol, pmr, cmr*)

Fürstner, A. *et al.*, *J.A.C.S.*, 2002, **124**, 13856-13863 (*synth*)

Mooniine A

M-700



$C_{36}H_{44}N_2O_6$ 600.753

Alkaloid from the leaves of *Erythroxylum moonii*. Solid. λ_{max} 201 ; 227 ; 272 (CHCl₃).

N⁸-Me, O-depropanoyl, Ac: **Mooniine B**

$C_{36}H_{44}N_2O_6$ 600.753


Alkaloid from the leaves of *Erythroxylum moonii*. Solid. λ_{max} 200 ; 221 ; 272 (CHCl₃).

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1998, **48**, 377-383 (*isol, uv*)

Mooreine

M-701

[342008-06-4]



$C_{17}H_{18}NO_4^{\oplus}$ 300.334


Tentative struct. Quaternary alkaloid from *Crinum moorei*. Amorph. Counterion not specified.

Elgorashi, E.E. *et al.*, *Phytochemistry*, 2001, **56**, 637-640

Moroidin

M-702

[104041-75-0]



Absolute Configuration

$C_{47}H_{66}N_{14}O_{10}$ 987.126

Constit. of *Celosia argentea* and *Laportea moroides*. Tubulin polymn. inhibitor. Antimitotic agent. Needles (MeOH aq.). $[\alpha]_D^{24} -55$ (c, 0.3 in MeOH aq.). Dec. at 280°. λ_{max} 215; 275 (solvent not reported) (Derep).

1-Epimer, L-aspartic acid amide: **Celogentin E**

[602308-61-2]

$C_{51}H_{71}N_{15}O_{13}$ 1102.215

Constit. of the seeds of *Celosia argentea*. Tubulin polymn. inhibitor. Amorph. solid. $[\alpha]_D^{22} -39$ (c, 0.5 in MeOH aq.). λ_{max} 225 (log ϵ 4.2); 282 (log ϵ 3.6) (MeOH).

1-Epimer, L-arginine amide: **Celogentin F**

[602308-62-3]

$C_{53}H_{78}N_{18}O_{11}$ 1143.313

Constit. of the seeds of *Celosia argentea*. Tubulin polymn. inhibitor. Amorph. solid. $[\alpha]_D^{22} -31$ (c, 0.5 in MeOH aq.). λ_{max} 225 (log ϵ 4.2); 282 (log ϵ 3.6) (MeOH).

[119528-95-9, 119528-96-0, 119616-50-1, 119616-51-2]

Leung, T.-W.C. *et al.*, *Tetrahedron*, 1986, **42**, 3333-3348 (*isol, pmr, cmr*)

Kahn, S.D. *et al.*, *J.O.C.*, 1989, **54**, 1901-1904; 2000, **65**, 8406 (*pmr, struct*)

Morita, H. *et al.*, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 469-471 (*activity, isol, pmr, cmr, abs config*)


Suzuki, H. *et al.*, *Tetrahedron*, 2003, 5307-5315 (*Celogentins*)

Suzuki, H. *et al.*, *Tetrahedron*, 2004, **60**, 2489-2495 (*cryst struct*)

ψ -Morphine

M-703

7,7',8,8'-Tetrahydro-4,5:4',5'-diepoxy-17,17'-dimethyl[2,2'-bimorphinan], 9CI. Pseudomorphine. 2,2'-Bimorphine. Dehydromorphine. Oxydimorphine. Oxymorphine [125-24-6]



Absolute configuration

$C_{34}H_{36}N_2O_6$ 568.668

Alkaloid from opium poppies and opium, also obt. by mild oxidn. of Morphine, M-704. Rods + 1H₂O. Insol. MeOH. Mp 330° dec.

Tetra-Ac:

Prisms (MeOH). Mp 294-296°.

3,3'-Di-Me ether Mp 155-156°.


Bentley, K.W. *et al.*, *J.C.S.*, 1959, 2574 (*synth, struct, uv*)

Mushinskaya, S.Kh. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 855; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 845 (*isol*)

Morphine, BAN

M-704

7,8-Didehydro-4,5-epoxy-17-methylmorphinan-3,6-diol, 9CI. Meconium. Morfine



(-)-form

$C_{17}H_{19}NO_3$ 285.342


Log P 0.24 (uncertain value) (calc).

(-)-form [57-27-2]

- Principal alkaloid of opium (*Papaver somniferum*) (Papaveraceae). Also found in traces in other plants (e.g. lettuce) and mammalian tissues, apparently as a result of genuine biosynthesis. Powerful opioid analgesic. Psychotomimetic agent, anti-tussive, antiemetic. A drug of abuse. Acts as a stimulant to horses, has been used in horse doping. Prisms + 1H₂O (EtOH aq.), anhyd. prisms (anisole). V. sol. EtOAc, Me₂CO; sol. EtOH; spar. sol. H₂O, Et₂O, CHCl₃. Mp 254-256.4° dec. (anhyd.). $[\alpha]_D^{23}$ -130.9 (MeOH). pK_{a1} 8.2; pK_{a2} 9.9 (25°).
- Gastrointestinal effects when used therapeutically. Human and exp. reprod. effects. Abuse leads to habituation or addiction. Larger doses produce respiratory depression leading to coma. Can cause death. LD₅₀ (rat, orl) 335 mg/kg. QC7875000
- Hydrochloride*: [52-26-6] [6055-06-7]
Silky needles + 3H₂O (dil. HCl). Sol. H₂O; v. spar. sol. EtOH. Mp 200°. $[\alpha]_D^{18}$ -97.9 (H₂O).
- Abuse leads to habituation or addiction. Exp. reprod. and teratogenic effects. LD₅₀ (rat, orl) 335 mg/kg. QC8575000
- Sulfate*: **Morphine sulfate, JAN, USAN**. *Astramorph P/F. Duramorph. Infumorph. MS Contin. Oramorph SR* [6211-15-0]
Solid + 5H₂O.
- N-Oxide**: **Morphine N-oxide. Genomorphine** [639-46-3]
C₁₇H₁₉NO₄ 301.341
Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Mp 272-273°. Metab. of Morphine, M-704.
- 3-O-β-D-Glucuronopyranoside**: [20290-09-9]
C₂₃H₂₇NO₉ 461.468
Needles + 2.5 H₂O (H₂O). Mp 243-246° dec. (hydrate). $[\alpha]_D^{28}$ -132 (c, 0.5 in H₂O). Human metabolite of morphine.
- 6-O-α-D-Glucuronopyranoside**: [150949-28-3]
C₂₃H₂₇NO₉ 461.468
Mp 310° dec. $[\alpha]_D^{20}$ -86.9 (c, 1 in DMSO).
- 6-O-β-D-Glucuronopyranoside: Morphine glucuronide, INN. M6G** [20290-10-2]
C₂₃H₂₇NO₉ 461.468
Active metab. of morphine associated with less respiratory depression than morphine. Analgesic. Cryst. + 2H₂O (EtOH aq.). Mp 254-256° dec. (dihydrate). $[\alpha]_D^{28}$ -172 (c, 0.5 in H₂O). Human metabolite of morphine.
- 3,6-Di-O-β-D-glucuronopyranoside**: [64947-41-7]
C₂₉H₃₅NO₁₅ 637.593
Cryst. + 3H₂O (MeOH aq.). Mp 243-244° dec. (trihydrate). $[\alpha]_D$ -188 (c, 10 in H₂O).
- 6-Ac**: [2784-73-8]
Mp 187° (as hydrochloride). $[\alpha]_D^{23}$ -163 (H₂O).
- 3-Benzoyl**: [105092-16-8]
C₂₄H₂₃NO₄ 389.45
Cryst. (EtOH). Mp 180-182°.
- 6-Benzoyl**: Mp 269-270°.
- 3-Me ether**: see Codeine, C-555
- 6-Me ether: Heterocodeine** [639-47-4]
C₁₈H₂₁NO₃ 299.369
Narcotic analgesic, preanaesthetic, antitussive, antiperistaltic agent. Mp 242-243°.
- 6-Me ether, 3-Ac**:
Cryst. (EtOH). Mp 133.5-134°.
- Di-Me ether: 6-O-Methylcodeine** [2859-16-7]
C₁₉H₂₃NO₃ 313.396
Impurity in Codeine, C-555 prod. by methylation of morphine. Minor alkaloid of *Papaver somniferum* (opium poppy). Prisms or plates. Mp 140-141°. $[\alpha]_D$ -188.5 (c, 1.09 in 95% EtOH).
- 3-Et ether: Ethylmorphine, BAN** [76-58-4]
C₁₉H₂₃NO₃ 313.396
Analgesic, antitussive, mydriatic. Mp 199-201°. Log P 1.35 (uncertain value) (calc).
- Addictive. LD₅₀ (rat, orl) 810 mg/kg. QD0850000
- 3-Et ether; hydrochloride: Codethyline. Dionina** [125-30-4]
Bitter-tasting powder. Mp 123° dec.
- LD₅₀ (rat, orl) 950 mg/kg. QD0874000
- (+)-form** [65165-99-3]
Synthetic. $[\alpha]_D^{25}$ +118 (c, 0.25 in CHCl₃). [16206-77-2, 64-31-3]
- Bentley, K.W. et al., *Chemistry of the Morphine Alkaloids*, Oxford Univ. Press, 1954, (isol, bibl, uv)
- Gates, M. et al., *J.A.C.S.*, 1956, **78**, 1380-1393 (synth)
- Kelentei, B. et al., *Arzneim.-Forsch.*, 1957, **7**, 594-597 (oxide, synth, pharmacol)
- Brochmann-Hanssen, E. et al., *J. Pharm. Sci.*, 1965, **54**, 1393 (di-Me ether, isol, ir, pmr)
- Wheeler, D.M.S. et al., *J.A.C.S.*, 1967, **89**, 4494-4501 (ms)
- Yoshimura, H. et al., *Chem. Pharm. Bull.*, 1968, **16**, 2114-2119 (glucuronosiduronates)
- Fridrichsons, J. et al., *Tet. Lett.*, 1968, 2887-2890 (cryst struct, abs config)
- Fennessy, M.R. et al., *Eur. J. Pharmacol.*, 1969, **8**, 261 (pharmacol)
- Kametani, T. et al., *J.C.S.(C)*, 1969, 2030-2033 (synth)
- de Angelis, G.G. et al., *Tetrahedron*, 1969, **25**, 5099-5112 (ord)
- Bentley, K.W. et al., *Alkaloids (Academic Press)*, 1971, **13**, 1-163 (rev, bibl)
- Heimans, R.L.H. et al., *J. Pharm. Pharmacol.*, 1971, **23**, 831-836 (metab)
- Barber, R.B. et al., *J. Med. Chem.*, 1975, **18**, 1074-1077 (di-Me ether, Heterocodeine, synth)
- Battersby, A.R. et al., *Tet. Lett.*, 1975, 1873-1876 (biosynth)
- Bye, E. et al., *Acta Chem. Scand., Ser. B*, 1976, **30**, 549-554 (cryst struct)
- Carroll, F.I. et al., *J.O.C.*, 1976, **41**, 996-1001 (cmr)
- Phillipson, J.D. et al., *Phytochemistry*, 1976, **15**, 1297-1301 (oxide)
- Thenot, J.P.G. et al., *Methods Biochem. Anal.*, 1977, **24**, 1-38 (rev, glc)
- Factors Affecting Action Narc.* [Proc. Meet.], Raven Press, N.Y., 1978, 103; 233; 565; 595 (rev, pharmacol)
- Brochmann-Hanssen, E. et al., *J. Nat. Prod.*, 1980, **43**, 731-735 (Heterocodeine)
- Hazum, E. et al., *Science (Washington, D.C.)*, 1981, **213**, 1010-1012 (pharmacol)
- Allen, A.C. et al., *J.O.C.*, 1983, **48**, 3951-3954 (oxide, synth)
- Rice, K.C. et al., *Chem. Biol. Isoquinoline Alkaloids, Int. Symp., Phytochem. Soc. Eur., Abstr. Pap.*, 1985, 191 (rev, synth)
- Goldstein, A. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1985, **82**, 5203-5207 (isol)
- Toth, J.E. et al., *J.O.C.*, 1987, **52**, 473-475 (synth)
- Muhtadi, F.J. et al., *Anal. Profiles Drug Subst.*, 1988, **17**, 259 (rev)
- Toth, J.E. et al., *J.O.C.*, 1988, **53**, 4694-4708 (synth)
- Parker, K.A. et al., *J.A.C.S.*, 1992, **114**, 9688-9689 (synth)
- Hong, C.Y. et al., *J.A.C.S.*, 1993, **115**, 11028-11029 (synth)
- Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms, 7th edn., Akademie-Verlag*, 1994, 5800 (synonyms)
- Hong, C.Y. et al., *Tet. Lett.*, 1994, **35**, 3453-3456 (synth)
- Opium Poppy: Botany, Chemistry and Pharmacology*, Pharmaceutical Products Press, 1995,
- Mulzer, J. et al., *Angew. Chem., Int. Ed.*, 1996, **35**, 2830-2832 (synth)
- White, J.D. et al., *J.O.C.*, 1997, **62**, 5250-5253 (synth)
- Mulzer, J. et al., *Chirality*, 1999, **11**, 475-482 (synth)
- Meuzelaar, G.J. et al., *Eur. J. Org. Chem.*, 1999, 2315-2321 (synth)
- Martindale, *The Extra Pharmacopoeia, 32nd edn.*, Pharmaceutical Press, 1999, 26-27; 56-60
- Rukhman, I. et al., *Synthesis*, 2000, 1241-1246 (3-benzoyl)
- Brown, R.T. et al., *Tetrahedron*, 2000, **56**, 7591-7594 (3,6-diglucuronoside)
- Rukhman, I. et al., *Tetrahedron*, 2001, **57**, 1083-1092 (glucuronosides)
- Blakemore, P.R. et al., *Chem. Comm.*, 2002, 1159-1168 (rev)
- Taber, D.F. et al., *J.A.C.S.*, 2002, **124**, 12416-12417; 15399 (synth)
- Romberg, R. et al., *Anesthesiology*, 2003, **99**, 788-798 (6-glucuronopyranoside, pharmacol)
- Okura, T. et al., *Br. J. Pharmacol.*, 2003, **140**, 211-217 (6-glucuronopyranoside, pharmacol)
- Trost, B.M. et al., *J.A.C.S.*, 2005, **127**, 14785-14803 (synth)
- Parker, K.A. et al., *J.O.C.*, 2006, **71**, 449-455 (synth)
- Uchida, K. et al., *Org. Lett.*, 2006, **8**, 5311-5313 (synth)
- Tanimoto, H. et al., *Tet. Lett.*, 2008, **49**, 358-362 (synth)
- Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, ENK000; MRO500; MRO0750

Morpholine, 9CI**M-705**

1-Oxa-4-azacyclohexane. Tetrahydro-1,4-oxazine. Drevamine [110-91-8]




C₄H₉NO 87.121

- Solvent for resins, waxes and dyes. Permitted (FDA) in edible coatings for fruit and vegetables. Synthetic reagent. Anal. reagent for α,β -unsatd. compds. and for anhydrides. Food contaminant arising from its use as a boiler water additive. Hygroscopic oil with ammoniacal odour and caustic props. Misc. H_2O . d_4^{20} 1. Mp -4.9°. Bp 128.9° Bp₆ 20°. n_D^{20} 1.4545. pK_a 8.49 (25°). Steam-volatile.
- Flammable, fl. p. 37°(oc)/38°(oc), auto-ignition temp. 310°/290°. Corrosive and irritating to skin, eyes and mucous membranes. LD₅₀ (rbt, orl) 1050 mg/kg. LD₅₀ (rbt, skn) 500 mg/kg. Exp. nephrotoxic. OES: long-term 20 ppm; short-term 30 ppm (Sk). QD6475000
- Hydrochloride*: [10024-89-2]
Cryst. (HCl aq.). Mp 175-176°.
- QE5075000
Picrate: Mp 225°.
- N-Formyl: 4-Morpholinecarboxaldehyde* [4394-85-8]
 $C_5H_9NO_2$ 115.132
Formylating reagent. Liq. Mp 17.5°. Bp 234°.
- QD9694000
N-Ac: [1696-20-4]
 $C_6H_{11}NO_2$ 129.158
Sol. H_2O . d_4^{20} 1.12. Mp 14.5°. Bp₅₀ 152° Bp₁₂ 118°. n_D^{20} 1.4827.
- Eye irritant. QD7000000
- N-Propanoyl: 1-Morpholino-1-propanone* [30668-14-5]
 $C_7H_{13}NO_2$ 143.185
Liq. Bp_{1.1} 83-86°.
- N-Benzoyl*: [1468-28-6]
 $C_{11}H_{13}NO_2$ 191.229
Cryst. (Et₂O). Mp 74-75°.
- QD8050000
N-(4-Methylbenzenesulfonyl):
 $C_{11}H_{15}NO_3S$ 241.31
Mp 147°.
- N-Me: N-Methylmorpholine* [109-02-4]
 $C_5H_{11}NO$ 101.148
Isol. from seeds of *Cassia occidentalis*. Base used in mixed anhydride peptide synth. which minimises racemisation. Bp 116-117°.
- Corrosive and irritating to skin, eyes and mucous membranes. LD₅₀ (rat, orl) 1970 mg/kg. QE5775000
- N-Me; hydrochloride*: [3651-67-0]
Mp 207-211°.
- N-Me, N-oxide*: [7529-22-8]
 $C_5H_{11}NO_2$ 117.147
Catalyst for oxidn. of alcohols. Dipolar aprotic solv.; in conjunction with DMSO completely dissolves plant wall polysaccharides. Used in the manuf. of Tencel from cellulose (Courtaulds). Oxidises halides to alcohols/ketones. Monohydrate. Mp 73-76°.
- N-Et*: [100-74-3]
 $C_6H_{13}NO$ 115.175
Solvent. Dissolves LiAlH₄. Bp 138-139°.
- Skin and severe eye irritant. LD₅₀ (rat, orl) 1780 mg/kg. QE4025000
- N-Et, borane complex: (4-Ethylmorpholino-N⁴)trihydroboron* [88996-22-9]
 $C_6H_{16}BNO$ 129.01
d 0.94. n_D^{20} 1.4680. Mp -10 to -8°.
- N-(2-Propenyl)*: [696-57-1]
 $C_7H_{13}NO$ 127.186
 d_4^{25} 0.93. Mp 118° (as picrate). Bp 156-158°.
- N-Butyl*: [1005-67-0]
 $C_8H_{17}NO$ 143.228
Bp₉₈ 110-115°.
- N-(2-Methylpropyl): 4-(2-Methylpropyl)morpholine, 9CI* [10315-98-7]
 $C_8H_{17}NO$ 143.228
Bp₁₉ 66°. n_D^{20} 1.4391.
- N-(2-Methylpropyl), picrate*: [10315-99-8]
Needles (EtOH). Mp 139°.
- N-Ph*: [92-53-5]
 $C_{10}H_{13}NO$ 163.219
Dehydrohalogenating reagent. Mp 57°. Bp 270°.
- Toxic by skin absorption. Eye irritant. LD₅₀ (rat, orl) 930 mg/kg. LD₅₀ (rbt, skn) 360 mg/kg. QE8575000
- N-(2-Hydroxyethyl): 4-Morpholineethanol, 9CI, 8CI* [622-40-2]
 $C_6H_{13}NO_2$ 131.174
Liq. d_4^{20} 1.08. Bp₇₅₇ 227°.
- QE3510000
N-(2-Hydroxypropyl): α -Methyl-4-morpholineethanol, 9CI [2109-66-2]
 $C_7H_{15}NO_2$ 145.201
Bp₂₀ 98-102°.
- N-(2-Chloroethyl)*: [3647-69-6]
 $C_6H_{12}ClNO$ 149.62
Mp 184-186° (as hydrochloride). CAS no. refers to hydrochloride.
- QE0260000
N-Nitroso: [59-89-2]
 $C_4H_8N_2O_2$ 116.119
- Possible human carcinogen (IARC 2B). LD₅₀ (rat, orl) 282 mg/kg. Exp. carcinogen. QE7525000
- N-Nitro: 4-Nitromorpholine, 9CI* [4164-32-3]
 $C_4H_8N_2O_3$ 132.119
Cryst. Mp 52-54°.
- QE7392000
N-Amino: 4-Morpholinamine, 9CI. 4-Aminomorpholine, 8CI [4319-49-7]
[56677-61-3]
 $C_4H_{10}N_2O$ 102.136
Liq.; cryst. (as hydrochloride). Mp 167° (hydrochloride). Bp 163-165°.
- Aldrich Library of NMR Spectra, 2nd edn.*, 1983, 2, 1009C
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 382A; 382D; 383C; 383D; 383C; 764D; 765A; 1311A; 2, 907C; 1129B (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 597A; 597C; 598A; 599C; 1250C; 1251A; 2, 653C (nmr)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 472A; 472C; 472D; 473D; 785D; 1178B (ir)
- Jones, L.W. et al., *J.A.C.S.*, 1925, 47, 2966 (synth)
Medard, M.L. et al., *Bull. Soc. Chim. Fr.*, 1936, 1343 (deriv. synth)
Hampton, B.L. et al., *J.A.C.S.*, 1936, 58, 2338 (synth)
Butler, G.B. et al., *J.A.C.S.*, 1949, 71, 3120 (N-2-propenyl)
Reynolds, D.D. et al., *J.A.C.S.*, 1950, 72, 1597 (N-butyl)
Adv. Chem. Ser., 1955, 15, 515 (props)
Johnson, J.B. et al., *Anal. Chem.*, 1955, 27, 1464 (use)
Diamond, L.H. et al., *J.A.C.S.*, 1955, 77, 3131 (N-amino)
Critchfield, F.E. et al., *Anal. Chem.*, 1956, 28, 76 (use)
Opitz, G. et al., *Annalen*, 1963, 665, 101 (N-2-methylpropyl)
Chapman, N.B. et al., *J.C.S.*, 1963, 1385 (N-2-hydroxypropyl)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1967, 1, 383; 705; 1969, 2, 278; 1979, 7, 244 (N-Me, N-Et, use)
Potnis, S.P. et al., *Chem. Process. Eng. (Bombay)*, 1969, 3, 27 (rev)
Kim, H.L. et al., *J. Agric. Food Chem.*, 1971, 19, 198 (isol)
Bark, L.S. et al., *Analyst (London)*, 1972, 97, 783 (use)
Org. Synth., Coll. Vol., 5, 1973, 839 (nitro)
Takeuchi, Y. et al., *J.C.S. Perkin 2*, 1974, 1927 (cmr)
Van Rheenen, V. et al., *Tet. Lett.*, 1976, 1973 (N-Me N-oxide)
Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, 2, 295 (rev)
Org. Synth., 1978, 58, 43 (deriv)
Calderbank, K.E. et al., *J.C.S. Perkin 2*, 1979, 869 (nitro)
Otargaliev, T. et al., *Uzb. Khim. Zh.*, 1981, 81; CA, 95, 24951y (N-2-hydroxyethyl)
Brown, H.C. et al., *Inorg. Chem.*, 1984, 23, 2746 (N-Et borane complex, synth)
Lunn, G. et al., *J.O.C.*, 1984, 49, 3470-3473 (N-amino, synth, ir, pmr, cmr, ms)
Olah, G.A. et al., *J.O.C.*, 1984, 49, 3856 (deriv, use)
IARC Monog., 1989, 47, 199 (rev, tox)
Griffith, W.P. et al., *Synth. Commun.*, 1992, 22, 1967 (use, deriv)
Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 5, 3545-3547 (N-Me N-oxide, use)
Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 5, 3671-3673 (use)
Turnbull, M.M. et al., *Acta Cryst. C*, 1997, 53, 818-820 (cryst struct, N-Me hydrochloride)
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1912
Org. Synth., 2007, 84, 22-31 (N-propanoyl)
IARC Monog. (Web), (N-nitroso)
Patty's Ind. Hyg. Toxicol. (3rd Rev. edn.), Vol. 2, Wiley, 1980, 2693
Ethel Browning's Toxicity and Metabolism of Industrial Solvents, 2nd edn., (ed. Snyder, R.), Elsevier, Volume 2, 1990, 244
Bretherick, L. et al., *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 1555
Luxon, S.G. et al., *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 879
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, ACR750; NKZ000; ENL000; CHE000; MMA250; MRQ500; MRQ600; PFS750; MRP750

Mortivinacin B

M-706

C₁₃H₁₇NO₅ 267.281

Prod. by *Mortierella vinacea*. Pale yellow glass. λ_{\max} 212 (ϵ 5200); 258 (ϵ 1200); 304 (ϵ 400) (MeOH). λ_{\max} 212 (ϵ 5200); 258 (ϵ 1200); 304 (ϵ 400) (MeOH) (Berdy).

Soman, A.G. *et al.*, *J. Nat. Prod.*, 1999, **62**, 386-388 (*isol, uv, ir, pmr, cmr, ms*)

Mortivinacin C

M-707

As Mortivinacin B, M-706 with R = CH₃

C₁₄H₁₉NO₅ 281.308


Prod. by *Mortierella vinacea*. Pale yellow glass. $[\alpha]_D$ -16 (c, 1.9 in CHCl₃). λ_{\max} 210 (ϵ 3300); 258 (ϵ 1300); 304 (ϵ 400) (MeOH). λ_{\max} 210 (ϵ 3300); 258 (ϵ 1300); 304 (ϵ 400) (MeOH) (Berdy).

Soman, A.G. *et al.*, *J. Nat. Prod.*, 1999, **62**, 386-388 (*isol, uv, ir, pmr, cmr, ms*)

Morusimic acid B

M-708

[455949-43-6]

C₁₈H₃₅NO₄ 329.479

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ +8.8 (c, 0.42 in MeOH).

3-O- β -D-Glucopyranoside: **Morusimic acid A**

[455949-42-5]

C₂₄H₄₅NO₉ 491.621

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ +15.3 (c, 0.18 in MeOH).


Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192 (*isol, pmr, cmr*)

Bouillon, M.E. *et al.*, *Tetrahedron*, 2007, **63**, 2712-2723 (*synth*)

Morusimic acid D

M-709

[455949-45-8]

C₁₈H₃₅NO₄ 329.479

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ -14.6 (c, 0.25 in MeOH).

3-O- β -D-Glucopyranoside: **Morusimic acid C**

[455949-44-7]

C₂₄H₄₅NO₉ 491.621

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ -20.3 (c, 0.24 in MeOH).

6'-Hydroxy, 3-O- β -D-glucopyranoside:**Morusimic acid E**

[455949-46-9]

C₂₄H₄₅NO₁₀ 507.62

Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ -17.2 (c, 0.61 in MeOH).

4'-Epimer: **Morusimic acid F**

[455949-47-0]

C₁₈H₃₅NO₄ 329.479


Alkaloid from the fruit of *Morus alba*. Powder. $[\alpha]_D$ +6.4 (c, 0.28 in MeOH).

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 185-192 (*isol, pmr, cmr*)

Moschamide

M-710

[205234-95-3]

C₂₀H₁₈N₂O₆ 382.372


Alkaloid from the seeds of *Centaurea moschata*. Amorph. yellow solid. λ_{\max} 236 (log ϵ 3.93); 271 (log ϵ 3.53); 320 (log ϵ 2.98); 354 (log ϵ 3.01) (MeOH).

Sarker, S.D. *et al.*, *Tet. Lett.*, 1998, **39**, 1421-1424 (*isol, uv, ir, pmr, cmr, ms*)

Moschamindolol

M-711

[193225-06-8]

C₂₀H₂₀N₂O₅ 368.388


Alkaloid from the seeds of *Centaurea moschata* (Asteraceae). Amorph. yellow solid. Related to Serotobenine, S-252. λ_{\max} 228 (log ϵ 4.11); 286 (log ϵ 3.85); 311 (log ϵ 3.6); 322 (log ϵ 3.54) (MeOH).

Sarker, S.D. *et al.*, *Nat. Prod. Lett.*, 1997, **9**, 189-199 (*isol, uv, ir, pmr, cmr, ms*)

Moschatoline

M-712

2-Hydroxy-1,3-dimethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI
[5574-23-2]

C₁₈H₁₃NO₄ 307.305

Alkaloid from *Atherosperma moschatum* (Monimiaceae). Noncryst.

Ac:

Yellow cryst. Mp 190-200°.

Me ether: **O-Methylmoschatoline**. *Homo-moschatoline*. *Liridine*

[5140-38-5]

C₁₉H₁₅NO₄ 321.332

Alkaloid from *Guatteria subsessilis* and several other genera in the Annonaceae (*Annona*, *Duguetia*, *Cleistopholis*, *Enantia*, *Polyalthia*), also from the Magnoliaceae (*Liriodendron*) and Menispermaceae (*Abuta*, *Triclisia*). Mp 188° dec.

Me ether, O³-De-Me: **Isomoschatoline**

[83375-15-9]

C₁₈H₁₃NO₄ 307.305

Alkaloid from the stem bark of *Guatteria melosma* and from *Cleistopholis patens* (Annonaceae). Red cuboidal cryst. (MeOH). Mp 245°.

Me ether, O³-De-Me, Ac:

Fine yellow needles. Mp 193-194° dec.

O³-De-Me, O²-Me, N-Me: **1-Hydroxy-2,3-dimethoxy-6-methyl-7-oxo-7H-dibenzo[de,g]quinolinium inner salt**. **Teliglazine**

[199333-21-6]

C₁₉H₁₅NO₄ 321.332

Alkaloid from *Telitoxicum glaziovii* (Menispermaceae). Dark green needles (CHCl₃). Mp 229-232°. Inner salt with negative charge on O³. λ_{\max} 246 (sh) (log ϵ 3.24); 318 (log ϵ 3.43); 430 (log ϵ 2.53); 648 (log ϵ 2.55) (EtOH).

11-Hydroxy, O³-de-Me, O²-Me, N-Me:

SciaferineC₁₉H₁₅NO₅ 337.331

Alkaloid from *Sciadotenia toxifera*. Dark green needles (CH₂Cl₂/MeOH). Mp >293°. λ_{\max} 240 (sh) (log ϵ 2.96); 318 (log ϵ 2.84); 422 (log ϵ 2.09); 632 (log ϵ 1.94) (EtOH).

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1403 (*Moschatoline*)

Cava, M.P. *et al.*, *Tetrahedron*, 1975, **31**, 1667 (*synth*)

Marsaioli, A.J. *et al.*, *Phytochemistry*, 1980, **19**, 995 (*O-Methylmoschatoline*)

Abd-El Atti, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 476 (*Isomoschatoline*)

Tavanaiepour, I. *et al.*, *Acta Cryst. C*, 1987, **43**, 2230 (*O-Acetylisosmoschatoline, cryst struct*)


Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1328-1330 (*Teliglazine*)

Freyer, A.J. *et al.*, *Heterocycles*, 1999, **51**, 2221-2226 (*Sciaferine*)

Mossambine

M-713


Methyl 2,16,19,20-tetrahydro-14-hydroxycuran-17-oate, 9CI. **14-Hydroxykuanammicine**. **Diplorrhincine**
[30735-22-9]



Absolute Configuration

C₂₀H₂₂N₂O₃ 338.405Alkaloid from *Diplorhynchus condylocarpon* ssp. *mossambicensis* (Apocynaceae). Cryst. (MeOH). Mp 240°. [α]_D²⁰ -498 (CHCl₃). [α]_D²⁰ -470 (c, 0.40 in CHCl₃).

Hydrochloride: Mp 225-228° dec.

Stauffer, D. et al., *Helv. Chim. Acta*, 1961, **44**, 2006 (*isol, ir, uv*)Monseur, X. et al., *Bull. Soc. Chim. Fr.*, 1962, 1088 (*uv, ir, ms, pmr, struct*)Kuehne, M.E. et al., *J.O.C.*, 1996, **61**, 7873 (*synth, resoln*)**Motuporamine A** M-714N-(3-Azacyclotridec-1-ylpropyl)-1,3-propanediamine, 9CI
[211566-77-7]C₁₈H₃₉N₃ 297.526Alkaloid from the sponge *Xestospongia exigua*. Cytotoxic agent. Oil (as di-Ac).Goldring, W.P.D. et al., *Org. Lett.*, 1999, **1**, 1471-1473 (*synth, struct*)Baldwin, J.E. et al., *Tet. Lett.*, 1999, **40**, 5401-5404 (*synth*)Williams, D.E. et al., *J.O.C.*, 2002, **67**, 245-258 (*isol, pmr, cmr, activity*)**Motuporamine B** M-715N-(3-Azacyclotetradec-1-ylpropyl)-1,3-propanediamine
[211566-78-8]


As Motuporamine A, M-714 with n = 13

C₁₉H₄₁N₃ 311.553Alkaloid from the sponge *Xestospongia exigua*. Cytotoxic agent. Oil (as di-Ac).5Z,6-Didehydro: **Motuporamine D**



[398144-67-7]

C₁₉H₃₉N₃ 309.537Alkaloid from *Xestospongia exigua*. Oil.Williams, D.E. et al., *J.O.C.*, 1998, **63**, 4838-4841; 2002, **67**, 245-258 (*Motuporamines B, D, isol, pmr, cmr, activity*)Goldring, W.P.D. et al., *Org. Lett.*, 1999, **1**, 1471-1473 (*synth, struct*)Baldwin, J.E. et al., *Tet. Lett.*, 1999, **40**, 5401-5404 (*synth*)**Motuporamine C** M-716

[211569-34-5]

C₂₀H₄₁N₃ 323.564Alkaloid from the sponge *Xestospongia exigua*. Cytotoxic agent. Oil (as di-Ac).N⁸-Formyl: **Motuporamine F**
[398144-69-9]C₂₁H₄₁N₃O 351.574Alkaloid from *Xestospongia exigua*. Oil (as di-Ac).9Z,10-Didehydro: **Motuporamine E**

[398144-68-8]

C₂₀H₃₉N₃ 321.548Alkaloid from *Xestospongia exigua*. Oil.Williams, D.E. et al., *J.O.C.*, 1998, **63**, 4838-4841; 2002, **67**, 245-258 (*Motuporamines C, E, F, isol, pmr, cmr, activity*)Goldring, W.P.D. et al., *Org. Lett.*, 1999, **1**, 1471-1473 (*synth, struct*)Fuerstner, A. et al., *J.O.C.*, 2000, **65**, 2608-2611 (*synth*)**Mubironine C** M-717C₁₇H₂₉NO₃ 295.421Alkaloid from *Dendrobium* Snowflake "Red Star". Solid. [α]_D²³ -5 (c, 0.6 in MeOH).Morita, H. et al., *Tetrahedron*, 2000, **56**, 5801-5805**Mucronine A** M-71812-(Dimethylamino)-15-methoxy-9-(1-methylpropyl)-6-(phenylmethyl)-4,7,10-triazabicyclo[12.3.1]octadeca-1(18),2,14,16-tetraene-5,8,11-trione, 9CI
[38840-25-4]C₂₉H₃₈N₄O₄ 506.644Alkaloid from the bark, leaves and branches of *Zizyphus mucronata* and the bark of *Zizyphus abyssinica* (Rhamnaceae). Needles (CHCl₃/Et₂O or CH₂Cl₂/petrol). Mp 235°. [α]_D²⁰ -28.3 (c, 0.06 in CHCl₃).

Dihydro:

Needles (Me₂CO). Mp 237-239°. [α]_D²⁵ +57 (c, 0.06 in CHCl₃).N-De-Me: **Mucronine B**

[38840-26-5]

C₂₈H₃₆N₄O₄ 492.617Alkaloid from the bark of *Zizyphus mucronata* and *Zizyphus abyssinica* (Rhamnaceae). Fine needles (Me₂CO/petrol). Mp 222-224°. [α]_D²⁵ +175 (c, 0.2 in CHCl₃).

N-De-Me, dihydro:

Needles (Me₂CO). Mp 245-247°. [α]_D²⁵ +32.5 (c, 0.2 in CHCl₃).Bis-N-de-Me: **Mucronine H**

[55856-96-7]

C₂₇H₃₄N₄O₄ 478.59Alkaloid from bark and leaves of *Zizyphus mucronata* (Rhamnaceae). Shows antibacterial and antifungal props. Amorph. [α]_D²⁰ +5 (c, 0.1 in MeOH).

Bis-N-de-Me, N-Ac:

Needles (Me₂CO). Mp 315-317°. [α]_D²⁰ -24.3 (c, 0.074 in MeOH/CHCl₃ 1:1).Fehlhaber, H.-W. et al., *Annalen*, 1972, **759**, 195 (*isol, ir, uv, cd, ms, pmr, struct*)Tschesche, R. et al., *Annalen*, 1974, 1915 (*isol, uv, ir, pmr, ms, struct, Mucronine H*)Tschesche, R. et al., *Phytochemistry*, 1974, **13**, 2328 (*occur, Mucronine H*)Schmidt, U. et al., *Angew. Chem., Int. Ed.*, 1983, **22**, 152 (*synth, Mucronine B*)**Mucronine C** M-71912-(Dimethylamino)-15-methoxy-9-(1-methylpropyl)-6-(2-methylpropyl)-4,7,10-triazabicyclo[12.3.1]octadeca-1(18),2,14,16-tetraene-5,8,11-trione, 9CI
[38840-27-6]As Mucronine A, M-718 with R = CH₂CH(CH₃)₂C₂₆H₄₀N₄O₄ 472.626Alkaloid from the leaves and branches of *Zizyphus mucronata* and the bark of *Zizyphus abyssinica* (Rhamnaceae). Needles (CH₂Cl₂/petrol). Mp 257°. [α]_D²⁰ -39.4 (c, 0.09 in CHCl₃).N-De-Me: **Abyssenine A**. **Abyssinine A**
[54519-14-1]C₂₅H₃₈N₄O₄ 458.6Alkaloid from the bark of *Zizyphus mucronata* and *Zizyphus abyssinica* and from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Needles (Me₂CO/petrol). Mp 237-239°. [α]_D²⁰ +160 (c, 0.22 in CHCl₃). [α]_D²⁰ -58 (c, 0.1 in MeOH).17-Methoxy, N-de-Me: **Mucronine E**

[55856-87-6]

C₂₆H₄₀N₄O₅ 488.626Alkaloid from the bark of *Zizyphus mucronata*. Needles (Me₂CO/petrol). Mp 232-234°. [α]_D²⁰ -89 (c, 0.084 in MeOH). Stereochem. determined in 2007.17-Methoxy, N-di-de-Me: **Mucronine F**


[55856-90-1]

C₂₅H₃₈N₄O₅ 474.599Alkaloid from the bark of *Zizyphus mucronata* (Rhamnaceae). Cryst. (CHCl₃/MeOH). Mp 208-214°. [α]_D²⁰ +17.4 (c, 0.092 in MeOH).Fehlhaber, H.-W. et al., *Annalen*, 1972, **759**, 195-207 (*Mucronine C*)Tschesche, R. et al., *Annalen*, 1974, 1915-1928 (*Mucronines E, F, Abyssinine A*)Tschesche, R. et al., *Phytochemistry*, 1974, **13**, 2328 (*Abyssinine A, occur*)Cassels, B.K. et al., *Tetrahedron*, 1974, **30**, 2461-2466 (*Zizyphus oenoplia consti*)Toumi, M. et al., *J.O.C.*, 2007, **72**, 9003-9009 (*Abyssenine A, synth*)Wang, J. et al., *Tet. Lett.*, 2007, **48**, 6717-6721 (*Mucronine E, synth*)

Mucronine D

Daechuine S9
[38496-00-3]

M-720



$C_{37}H_{51}N_5O_6$ 661.84

Alkaloid from the stem or root bark of *Zizyphus mucronata*, *Zizyphus mucronata*, *Zizyphus sativa*, *Zizyphus nummularia*, *Zizyphus jujuba* and *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Amorph. $[\alpha]_D^{20}$ -487 (c, 0.12 in $CHCl_3$).

N-De-Me: Nummularine A. N-Demethylnummularine D
[53947-95-8]

$C_{36}H_{49}N_5O_6$ 647.813

Alkaloid from the root bark of *Zizyphus nummularia* and the stem bark of *Zizyphus jujuba* (Rhamnaceae). Needles (MeOH). Mp 235-240° dec. $[\alpha]_D^{20}$ -397 (c, 0.2 in $CHCl_3$).

O-De-Me: O-Demethylmucronine D
[155656-07-8]

$C_{36}H_{49}N_5O_6$ 647.813

Alkaloid from roots of *Zizyphus mucronata* (Rhamnaceae). $[\alpha]_D^{20}$ -191 (c, 0.3 in $CHCl_3$).

N²-Deacyl, N², N²-di-Me: [155656-06-7]
 $C_{28}H_{42}N_4O_5$ 514.664

Alkaloid from roots of *Zizyphus mucronata* (Rhamnaceae). Needles (MeOAc/hexane). Mp 153-154°. $[\alpha]_D^{20}$ -418 (c, 1.1 in $CHCl_3$).

10,11-Dihydro:

Needles (CH_2Cl_2/Et_2O). Mp 250-252° (247°). $[\alpha]_D^{20}$ -225 (c, 0.17 in $CHCl_3$).

Tschesche, R. *et al.*, *Chem. Ber.*, 1972, **105**, 3106-3114; 1974, **107**, 3180-3185 (*Mucronine D, Nummularine A, isol, uv, cd, ir, pmr, ms, struct*)

Tschesche, R. *et al.*, *Phytochemistry*, 1976, **15**, 541-542; 1979, **18**, 702-704 (*isol*)


Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443-448 (*Nummularine A*)

Barboni, L. *et al.*, *Phytochemistry*, 1994, **35**, 1579-1582 (*O-Demethylmucronine D, N-deacyl-N-di-Me*)

Mucronine J

[177714-94-2]

M-721



Absolute
Configuration

$C_{27}H_{40}N_4O_4$ 484.637

Alkaloid from root bark of *Zizyphus mucronata* and roots of *Paliurus ramosissimus*. Amorph. powder. $[\alpha]_D^{21}$ -236 (c, 1 in $CHCl_3$). λ_{max} 219 (log ϵ 4.5) (MeOH).

Auvin, C. *et al.*, *J. Nat. Prod.*, 1996, **59**, 676-678 (*isol, ir, pmr, cmr, ms, struct*)

Lim, H.Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (*isol, config*)

Mucudaine

M-722

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Lemon-yellow pungent liq. which gradually turns brown. Bp 239-240°. $[\alpha]_D^{30}$ +2.5 (EtOH). n_D^{30} 1.446.

Picrate:

Cryst. (EtOH). Mp 110-111°.

Majumdar, D.N. *et al.*, *Indian Pharm.*, 1954, **10**, 79-84; *CA*, **49**, 9881a (*isol*)

Mucudanine

M-723

Struct. unknown. MW 215.6. Alkaloid from *Mucuna pruriens* (Fabaceae). Light-brown pungent liq. Mp 5-7°. Bp 300°. $[\alpha]_D^{30}$ +3.1 (EtOH). n_D^{30} 1.459.

Picrate: Mp 114-115°.

Majumdar, D.N. *et al.*, *Indian Pharm.*, 1954, **10**, 79-84; *CA*, **49**, 9881a (*isol*)

Mucudanine

M-724

Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Flakes. Mp 300°.

Picrate: Mp 106-107°.

Majumdar, D.N. *et al.*, *Indian Pharm.*, 1954, **10**, 79-84; *CA*, **49**, 9881a (*isol*)

Mucunadine

M-725

Struct. unknown. Alkaloid from the seeds of *Mucuna pruriens* (Fabaceae). Brown powder ($CHCl_3$). Mp 201-202°.

Hydrochloride:

Microcryst. powder. Dec. >300°.

Santra, D.K. *et al.*, *Indian J. Pharm.*, 1953, **15**, 60-61; *CA*, **48**, 8793h (*isol*)

Mucunine

M-726

Struct. unknown. Alkaloid from the seeds of *Mucuna pruriens* (Fabaceae). Brown amorph. solid (Et_2O). Mp 220-221°.

Hydrochloride:

Needles. Mp 300°.

Picrate:


Yellowish powder. Mp 245-247°.

Santra, D.K. *et al.*, *Indian J. Pharm.*, 1953, **15**, 60-61; *CA*, **48**, 8793h (*isol*)

Mukanadin B

M-727

[250782-32-2]



$C_{11}H_{11}BrN_4O_3$ 327.137

Similar to Midpacamide, M-604. Isol. from the sponges *Agelas nakamura* and *Axinella verrucosa*. Amorph. solid. λ_{max} 274 (ϵ 6700) (MeOH).

9-Hydroxy: 9-Hydroxymukanadin B

$C_{11}H_{11}BrN_4O_4$ 343.136

Isol. from *Axinella verrucosa*.

2-Bromo: Mukanadin D

[193619-42-0]

$C_{11}H_{10}Br_2N_4O_3$ 406.033

Isol. from the sponge *Didiscus oxeata*.

Lindel, T. *et al.*, *Liebigs Ann./Recl.*, 1997, 1525-1528 (*synth*)

Uemoto, H. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1581-1583 (*isol, pmr, cmr*)


Hu, J.-F. *et al.*, *J. Chem. Res., Synop.*, 2005, 427-428 (*Mukanadin D*)

Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 17-24 (*9-Hydroxymukanadin B*)

Mukoenine A

M-728

2-Hydroxy-3-methyl-1-prenylcarbazole. 3-Methyl-1-(3-methyl-2-butenyl)-9H-carbazol-2-ol, 9CI. Girinimbilol
[155519-81-6]



$C_{18}H_{19}NO$ 265.354

Alkaloid from stem bark and roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Needles (hexane/ CH_2Cl_2) or prisms. Mp 106° (104-107°). Mukoenine A and Girinimbilol have not been directly compared but they appear to be identical.

O-Ac:

Needles (hexane/ CH_2Cl_2). Mp 172-174°.


Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 2096 (*isol, uv, ir, pmr, ms, struct*)

Reisch, J. *et al.*, *Phytochemistry*, 1994, **36**, 1073 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mukoenine B

M-729

1-(3,7-Dimethyl-2,6-octadienyl)-2-hydroxy-9H-carbazole-3-carboxaldehyde, 9CI. 3-Formyl-1-geranyl-2-hydroxycarbazole. Clausenatine A
[155519-82-7]



$C_{23}H_{25}NO_2$ 347.456

Alkaloid from *Clausena excavata* and *Murraya koenigii* (curryleaf tree) (Rutaceae). Yellowish powder (Me₂CO). Mp >280°.

8-Hydroxy: 3-Formyl-1-geranyl-2,8-dihydroxycarbazole. Clausoline F
[185508-06-9]


C₂₃H₂₅NO₃ 363.455

Alkaloid from the stem bark of *Clau-sena excavata*. Yellow powder. λ_{\max} 204 ; 216 ; 242 ; 276 ; 291 ; 300 (sh) ; 354 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 2096; 1996, **44**, 2231-2235 (*isol, uv, ir, pmr, ms, struct*)

Wu, T.S. *et al.*, *Phytochemistry*, 1999, **52**, 523-527 (*Clausenatine A*)

7,9(11)-Multifloradiene-3,29-diol M-730

C₃₀H₄₈O₂ 440.708

3 α -form

Karounidiol

[118117-31-0]

Constit. of *Trichosanthes kirilowii*. Cryst. Mp 201-203°.

3-O-Benzoyl: [118117-32-1]

C₃₇H₅₂O₃ 544.816

Constit. of *Trichosanthes kirilowii*. Cryst. Mp 119-122°.

Dibenzoyl: [389122-01-4]

C₄₄H₅₆O₄ 648.924

Constit. of *Gymnopetalum integrifolium* fruits. Cryst. (hexane). Mp 149-152°. λ_{\max} 228 (log ϵ 4.48) (hexane).

3-O-(4-Aminobenzoyl): **Debenzoylzucchini factor B**

[308811-93-0]

C₃₇H₅₃NO₃ 559.831

Constit. of *Cucurbita pepo* seeds. Gum. $[\alpha]_D^{25}$ -111 (c, 0.9 in MeOH). λ_{\max} 290 (MeOH).

3-O-(4-Aminobenzoyl), 29-O-benzoyl:

Zucchini factor B

[246248-10-2]

C₄₄H₅₇NO₄ 663.939

Constit. of seeds of pumpkin (*Cucurbita maxima*) and squash (*Cucurbita pepo*). Powder (Et₂O). Mp 204-206°. $[\alpha]_D^{25}$ -130 (c, 0.9 in CHCl₃). λ_{\max} 240 (sh) ; 290 (log ϵ 4.06) (EtOH).

3 β -form

3-Epikarounidiol

Constit. of *Trichosanthes kirilowii*. Cryst. Mp 225-226°. λ_{\max} 232 (ϵ 11200); 236 (ϵ 12600); 247 (ϵ 3180) (EtOH) (Derep).

Akihisa, T. *et al.*, *J.C.S. Perkin 1*, 1988, 439 (*isol, pmr, cmr, cryst struct*)

Akihisa, T. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1101 (*isol, pmr, cmr*)

Appendino, G. *et al.*, *Phytochemistry*, 1999, **51**, 1021-1026 (*Cucurbita ester*)


Appendino, G. *et al.*, *Fitoterapia*, 2000, **71**, 258-263 (*Debenzoylzucchini factor B*)

Sekine, T. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 645-648 (*dibenzoyl*)

Multiflorine

M-731

4-Oxo-2,3-didehydrosparteine. Base LV1. 1,7,7a,8,9,10,11,13,14,14a-Decahydro-7,14-methano-2H,6H-dipyrido[1,2-a:1',2'-e][1,5]diazocin-2-one, 9CI [529-80-6]



Absolute configuration

C₁₅H₂₂N₂O 246.352

Alkaloid from *Lupinus multiflorus*, *Lupinus varius*, *Lupinus albus*, *Lupinus mutabilis* and *Cadia ellisiana* (Fabaceae). CNS depressant. Mp 108-109°. $[\alpha]_D^{22}$ -317 (c, 0.4 in MeOH).

Perchlorate: Mp 160-162° dec.

Picrolonate: Mp 204°.

16-N-Oxide: **Multiflorine N-oxide**

[137453-21-5]

C₁₅H₂₂N₂O₂ 262.351

Constit. of *Lupinus hirsutus*. Oil. $[\alpha]_D^{23}$ -145.8 (c, 0.07 in EtOH).

5,6-Didehydro: **5-Dehydromultiflorine**

[66216-62-4]

C₁₅H₂₀N₂O 244.336

Constit. of *Lupinus hirsutus* and *Lupinus termis*. Oil. $[\alpha]_D^{25}$ -94.4 (c, 0.015 in CH₂Cl₂).

Didehydro: **Dehydromultiflorine**

C₁₅H₂₀N₂O 244.336

Alkaloid present in seeds and leaves of *Lupinus albuscens* (Fabaceae). Provisional identification. Posn. of double bond unknown (ms spectrum differs significantly from that of 5-dehydromultiflorine above). CAS No. not found to 2008.

13 α -Hydroxy: **13 α -Hydroxymultiflorine**.

Alkaloid X₁

[71657-64-2]

C₁₅H₂₂N₂O₂ 262.351

Minor alkaloid from *Lupinus albus* and *Lupinus cosentinii* seeds (as esters) (Fabaceae). Highly hygroscopic gum. $[\alpha]_D^{25}$ -335 (c, 1 in MeOH). Obt. from *L. cosentinii* as an ester mixt. (Alkaloid LC3) of which the benzoate was the only positively identified component.

13 α -Tigloyloxy: **13 α -Tigloyloxymultiflorine**

[136396-56-0]

C₂₀H₂₈N₂O₃ 344.453

Alkaloid from *Lupinus hirsutus* (Fabaceae). Oil. $[\alpha]_D^{23}$ -292 (c, 0.1 in EtOH).

13 α -Hydroxy, 5,6-didehydro: **A⁵-Dehydro-13-hydroxymultiflorine**. Alkaloid X₂

[220672-15-1]

C₁₅H₂₀N₂O₂ 260.335

Alkaloid from *Lupinus albus* (Fabaceae). Highly hygroscopic, obt. only in small amt.

13 α -Hydroxy, 5,6-didehydro, N¹⁶-oxide: **A⁵-Dehydro-13-hydroxymultiflorine N-oxide**

[220672-13-9]

C₁₅H₂₀N₂O₃ 276.335Alkaloid from *Lupinus albus*.

13 β -Hydroxy: **13 β -Hydroxymultiflorine** [197297-28-2]

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Lupinus varius* (Fabaceae). Yellow oil. $[\alpha]_D^{25}$ -304 (c, 0.1 in CHCl₃). λ_{\max} 327 (log ϵ 4.14) (MeOH).

13-Methoxy: **13-Methoxymultiflorine**

[149817-78-7]

C₁₆H₂₄N₂O₂ 276.378

Trace alkaloid in seeds and leaves of *Lupinus albuscens* (Fabaceae). Provisional identification.

[74867-85-9, 74902-86-6]

Crow, W.D. *et al.*, *Aust. J. Chem.*, 1955, **8**, 136-139; 1959, **12**, 474-482 (*isol, uv, ir, struct, pmr*)

Comin, J. *et al.*, *Aust. J. Chem.*, 1959, **12**, 468-473 (*isol, uv, ir, struct*)

Wiewiorowski, M. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1961, **9**, 715-719 (*isol, ir, struct, 13 α -hydroxy, 13 α -hydroxy 5,6-didehydro*)

Goldberg, S.I. *et al.*, *J.O.C.*, 1976, **32**, 1832-1837 (*cd, abs config*)

Beck, A.B. *et al.*, *J. Nat. Prod.*, 1979, **42**, 385-398 (*13-Hydroxymultiflorine*)

Pyzalska, D. *et al.*, *Acta Cryst. B*, 1980, **36**, 1602-1606; 1685-1687 (*cryst struct, 5-Dehydromultiflorine*)

Wink, M. *et al.*, *Planta Med.*, 1981, **43**, 342-352 (*isol, ms*)

Hatzold, T. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 934-938 (*Lupinus mutabilis constiit*)

Wysocka, W. *et al.*, *Croat. Chem. Acta*, 1989, **62**, 109 (*cd*)

Wyrzykiewicz, E. *et al.*, *Org. Mass Spectrom.*, 1990, **25**, 453-456 (*ms*)

Takamatsu, S. *et al.*, *Heterocycles*, 1991, **32**, 1167-1171 (*N-oxide*)

Takamatsu, S. *et al.*, *J. Nat. Prod.*, 1991, **54**, 477-482 (*13-Tigloyloxymultiflorine*)

Planchuelo-Ravelo, A.M. *et al.*, *Z. Naturforsch., C*, 1993, **48**, 414-416 (*13-Methoxymultiflorine, Dehydromultiflorine*)


Mohamed, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 365-369 (*13-Hydroxymultiflorines*)

Mohamed, M.H. *et al.*, *CA*, 1999, **130**, 179920c (*Dehydrohydroxymultiflorine N-oxide*)

Munroniamide

M-732

[632340-19-3]


C₃₁H₃₉N₃O₉ 597.664

Alkaloid from *Munronia henryi*. Moderate insect antifeedant. Powder. Mp 338-339°. $[\alpha]_D^{25}$ -3.7 (c, 0.27 in Py). λ_{\max} 209 (log ϵ 4.39) (MeOH).

Qi, S.-H. *et al.*, *J. Agric. Food Chem.*, 2003, **51**, 6949-6952 (*isol, pmr, cmr*)

Munronin D

[590400-77-4]


C₂₈H₃₃NO₈ 511.571

Constit. of *Munronia henryi*. Powder.
[α]_D²⁶ +5.9 (c. 0.34 in Py). Mp >350°. λ_{max}
208 (log ε 4) (MeOH).

Qi, S.-H. *et al.*, *Tetrahedron*, 2003, **59**, 4193-
4199 (*isol*, *pmr*, *cmr*)

Muramine

5,7,8,14-Tetrahydro-3,4,10,11-tetra-
methoxy-6-methylidibenz[c,g]azecin-
13(6H)-one, 9Cl. Cryptopalmatine
[2292-20-8]

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Papaver nudicaule* var.
amurense, several other *Papaver* spp.,
Glaucium vitellinum, *Argemone munita*
and *Argemone squarrosa* (Papaveraceae).
Cryst. (Me₂CO or MeOH). Mp 176-177°.

Hydroiodide: Mp 176-177° dec.

O⁹-De-Me: Protothalipine

[60332-15-2]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Thalictrum rugosum*
(Ranunculaceae). Needles (MeOH).
Mp 195-196° dec.

O²,O³-Di-de-Me: Vaillantine

[53964-96-8]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Fumaria vaillantii*
(Papaveraceae). Cryst. (Me₂CO/
MeOH). Mp 165-167°.

13-Oxo: 13-Oxomuramine. Alpinone†C₂₂H₂₅NO₆ 399.443

Trace alkaloid from *Papaver nudicaule*
var. *croceum* (Papaveraceae). Yellow
prisms (Me₂CO). Mp 173-180° Mp
204-205°.

Haworth, R.D. *et al.*, *J.C.S.*, 1927, 2261 (*synth*)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960,
47, 180 (*isol*, *ir*)

Stermitz, F.R. *et al.*, *J.O.C.*, 1966, **31**, 2925 (*isol*)

Flentje, H. *et al.*, *Pharmazie*, 1966, **21**, 321 (*uv*,
ir, 13-Oxomuramine)

Giacoppello, D. *et al.*, *Tet. Lett.*, 1966, 2859
(*struct*, *synth*)

Hanuš, V. *et al.*, *Coll. Czech. Chem. Comm.*,
1967, **32**, 1759 (*synth*, *ms*, 13-Oxomuramine)

Hruban, L. *et al.*, *Coll. Czech. Chem. Comm.*,
1967, **32**, 3414 (*uv*)

M-733

Stermitz, F.R. *et al.*, *J. Pharm. Sci.*, 1967, **56**,
760 (*isol*)

Nakashima, T.T. *et al.*, *Org. Magn. Reson.*,
1973, **5**, 9 (*cmr*)

Ibragimova, M.U. *et al.*, *Khim. Prir. Soedin.*,
1974, **10**, 476; *Chem. Nat. Compd. (Engl.*
Transl.), 1974, **10**, 481 (*isol*, *uv*, *ir*, *pmr*, *ms*,
struct, *Vaillantine*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 65
(*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *Protothalipine*)


Shafiee, A. *et al.*, *J. Nat. Prod.*, 1977, **40**, 352
(*isol*)

Moniot, J.L. *et al.*, *Heterocycles*, 1978, **9**, 145
(*pmr*, *Muramine*, *Protothalipine*)

Murayaanthraquinone

[162382-63-0]

M-735


C₂₆H₁₆N₂O₇ 468.422

Metab. from *Streptomyces murayamaen-*
sis. Orange-yellow cryst.

Hassan, A.M. *et al.*, *Bioorg. Med. Chem. Lett.*,
1995, **5**, 191 (*isol*, *ir*, *pmr*, *cmr*, *cryst struct*)

Muraymycin A

M-736

**Muraymycin A₁**C₅₂H₉₀N₁₄O₁₉ 1215.365

Prod. by a *Streptomyces* sp. Peptidogly-
can biosynth. inhibitor.

13'''-N-Deoxy: Muraymycin A₃C₅₂H₉₀N₁₄O₁₈ 1199.366

Prod. by a *Streptomyces* sp.

2^A-O-De-Me: Muraymycin A₄C₅₁H₈₈N₁₄O₁₉ 1201.339

Prod. by a *Streptomyces* sp.

3-O-Deglycosyl: Muraymycin A₅C₄₆H₇₉N₁₃O₁₆ 1070.208

Prod. by a *Streptomyces* sp.

Muraymycin A₂C₅₀H₈₆N₁₄O₁₉ 1187.312

Prod. by a *Streptomyces* sp. Peptidogly-
can biosynth. inhibitor.

McDonald, L.A. *et al.*, *J.A.C.S.*, 2002, **124**,
10260-10261 (*struct*)

Yamashita, A. *et al.*, *Bioorg. Med. Chem.*
Letts., 2003, **13**, 3345-3350 (*synth*, *sar*)

Muraymycin C₁

M-737

As Muraymycin A, M-736 with
R = OH

C₃₈H₆₃N₁₁O₁₇ 945.979

Prod. by a *Streptomyces* sp. Peptidogly-
can biosynth. inhibitor.

3''-O-(6-Methylheptanoyl): Muraymycin B₆C₄₆H₇₇N₁₁O₁₈ 1072.177

Prod. by a *Streptomyces* sp.

3''-O-(6-Methyloctanoyl): Muraymycin B₃C₄₇H₇₉N₁₁O₁₈ 1086.204

Prod. by a *Streptomyces* sp.

3''-O-(7-Methyloctanoyl): Muraymycin B₄C₄₇H₇₉N₁₁O₁₈ 1086.204

Prod. by a *Streptomyces* sp.

3''-O-(8-Methylnonanoyl): Muraymycin B₂C₄₈H₈₁N₁₁O₁₈ 1100.231

Prod. by a *Streptomyces* sp.

3''-O-(8-Methyldecanoyl): Muraymycin B₁C₄₉H₈₃N₁₁O₁₈ 1114.258

Peptidoglycan biosynth. inhibitor.

2^A-O-De-Me: Muraymycin C₂C₃₇H₆₁N₁₁O₁₇ 931.952

Prod. by a *Streptomyces* sp.

2^A-O-De-Me, 3''-O-(6-methylheptanoyl): Muraymycin B₇C₄₅H₇₅N₁₁O₁₈ 1058.15

Prod. by a *Streptomyces* sp.

2^A-O-De-Me, 3''-O-(7-methyloctanoyl): Muraymycin B₅C₄₆H₇₇N₁₁O₁₈ 1072.177

Prod. by a *Streptomyces* sp.

3''-Deoxy: Muraymycin D₁C₃₈H₆₃N₁₁O₁₆ 929.98

Prod. by a *Streptomyces* sp. Peptido-
glycan biosynth. inhibitor.

3''-Deoxy, 2^A-O-de-Me: Muraymycin D₂C₃₇H₆₁N₁₁O₁₆ 915.953

Prod. by a *Streptomyces* sp.

2^A-Demethoxy: Muraymycin C₃C₃₇H₆₁N₁₁O₁₆ 915.953

Prod. by a *Streptomyces* sp.

2^A-Demethoxy, 3''-deoxy: Muraymycin D₃C₃₇H₆₁N₁₁O₁₅ 899.954

Prod. by a *Streptomyces* sp.

3-O-Deglycosyl: Muraymycin C₄C₃₂H₅₂N₁₀O₁₄ 800.821

Prod. by a *Streptomyces* sp.


McDonald, L.A. *et al.*, *J.A.C.S.*, 2002, **124**,
10260-10261 (*struct*)

Yamashita, A. *et al.*, *Bioorg. Med. Chem.*
Letts., 2003, **13**, 3345-3350 (*synth*, *sar*)

Mureidomycin E

M-738

[126049-02-3]



$C_{39}H_{48}N_8O_{12}S$ 852.921

Nucleoside peptide antibiotic. Prod. by *Streptomyces flavidovirens*. Active against *Pseudomonas aeruginosa*. Powder. Sol. H_2O , MeOH; fairly sol. Me_2CO ; poorly sol. EtOAc, hexane. $[\alpha]_D -34.2$ (c, 1.17 in MeOH aq.). Amphoteric. Similar to Mureidomycin A. λ_{max} 258 (€ 21000) (pH 2) (Derep). λ_{max} 240 (€ 36800); 265 (sh) (€ 20000); 295 (sh) (€ 6820) (pH 9) (Derep). λ_{max} 258 (€ 21500) (pH 7) (Derep). λ_{max} 258 (E1%/1cm 252) (H_2O) (Berdy). λ_{max} 258 (E1%/1cm 247) (HCl) (Berdy). λ_{max} 240 (E1%/1cm 432) (NaOH) (Berdy).

6'-Hydroxy, 8'-deoxy: Mureidomycin F.

Napsamycin A

[126049-03-4]

$C_{39}H_{48}N_8O_{12}S$ 852.921

Prod. by *Streptomyces flavidovirens*, *Streptomyces candidus* and *Streptomyces* spp. HIL Y82, 11372. Active against *P. aeruginosa*. Powder. Mp 190° dec. $[\alpha]_D -40.3$ (c, 1.05 in MeOH aq.). Amphoteric. λ_{max} 258 (€ 21000) (pH 2) (Derep). λ_{max} 240 (€ 36800); 265 (sh) (€ 20000); 295 (sh) (€ 6820) (pH 9) (Derep). λ_{max} 258 (€ 21500) (pH 7) (Derep).

6'-Hydroxy, 8'-deoxy, 5'',6''-dihydro:

Napsamycin C

[144379-26-0]

$C_{39}H_{50}N_8O_{12}S$ 854.936

Prod. by *Streptomyces candidus* and *Streptomyces* spp. HIL Y82, 11372. Active against *P. sp.* Sol. H_2O , DMSO, MeOH- H_2O ; poorly sol. EtOAc, hexane. λ_{max} 256 (H_2O) (Berdy). λ_{max} 254 (HCl) (Berdy). λ_{max} 242 ; 294 (NaOH) (Berdy).

[144408-85-5]


Eur. Pat., 1992, 487 756; *CA*, **117**, 232207 (*Napsamycins*)

Isono, F. *et al.*, *J. Antibiot.*, 1993, **46**, 1203 (*Mureidomycins*)

Chattarjee, S. *et al.*, *J. Antibiot.*, 1994, **47**, 595 (*isol*, *pmr*, *ms*)

Murexine M-739

2-[[3-(1*H*-Imidazol-4-yl)-1-oxo-2-propenyl]oxy]-*N,N,N*-trimethylethanaminium(1+), 9*Cl*. Choline imidazole-4-acrylate. Choline urocanate [20284-40-6]



$C_{11}H_{18}N_3O_2^{\oplus}$ 224.282

Found in large amts. in the hypobranchial glands of molluscs, esp. *Murex* spp., incl. *Murex trunculus*, *Murex fulvescens* and *Murex brandaris*, and also *Concholepas concholepas* and *Tri-*

tonalia erinacea. Muscle relaxant with strong curariform action. Cholinesterase inhibitor. Sol. H_2O . pK_a 4.6. λ_{max} 285 (€ 16700) (pH 4.5 buffer) (Berdy).

Chloride: [6209-43-4]

$C_{11}H_{18}ClN_3O_2$ 259.735

Hygroscopic cryst. + $1H_2O$. Unstable, instantly hydrol. by H_2O .

Chloride; hydrochloride:

Hygroscopic microcryst. powder + $1/2 H_2O$. Mp 219-221° dec.

Dipicrate:

Yellow needles (H_2O). Mp 221-222° dec.

***N*-Me: N-Methylmurexine**

$C_{12}H_{20}N_3O_2^{\oplus}$ 238.309

Reported to occur in the marine gastropod mollusc *Nucella emarginata*. Struct. assignment incorr. A recent unambiguous synth. of the two possible *N*-Me derivs. of Murexine indicates that the nat. compd. is neither of these (Duke *et al* 1981).

1-Me, iodide: [70346-56-4]

Synthetic. Cryst. (MeOH). Mp 224-225°.

3-Me, iodide: [70346-59-7]

Synthetic. Mp 224°.

Erspamer, V. *et al.*, *Arch. Int. Pharmacodyn. Ther.*, 1947, **74**, 263-285; *CA*, **43**, 1491b (*isol*)

Pasini, C. *et al.*, *Annalen*, 1952, **578**, 6-10 (*synth*, *uv*)

Roseghini, M. *et al.*, *Eur. J. Biochem.*, 1970, **12**, 468 (*isol*)

Bender, J.A. *et al.*, *Comp. Gen. Pharmacol.*, 1974, **5**, 191 (*N*-Me)

Duke, C.C. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1739-1744 (*N*-Me, *synth*)

Muricine M-740


$C_{19}H_{21}NO_4$ 327.379

Struct. unknown. Alkaloid from *Annona muricata* (soursop) (Annonaceae). Conts. 3OMe groups and no NMe group. Suggested to be identical with De-*N*-methylcorydine (see Norcorydine in C-682).

Hydrobromide: Mp 242-243°.

Manske, R.H.F. *et al.*, *Alkaloids (Academic Press)*, 1954, **4**, 142

Murrafoline A M-741




$C_{41}H_{42}N_2O_2$ 594.795

(±)-**form** [88939-34-8]

Alkaloid from *Murraya euchrestifolia* (Rutaceae). Mp 260-262° (sinters at 170°).

McPhail, A.T. *et al.*, *Tet. Lett.*, 1983, **24**, 5377 (*uv*, *ir*, *pmr*, *ms*, *cryst struct*)

Murrafoline B M-742




$C_{32}H_{30}N_2O_2$ 474.601

(±)-**form** [98260-38-9]

Alkaloid from *Murraya euchrestifolia* (Rutaceae). Needles (MeOH). Mp 234-237°.

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 2611 (*uv*, *pmr*, *ms*, *struct*)

Murrafoline C M-743




$C_{36}H_{34}N_2O_2$ 526.677

(±)-**form** [98260-37-8]

Alkaloid from *Murraya euchrestifolia* (Rutaceae). Oil.

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 2611 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Murrafoline D M-744



$C_{32}H_{30}N_2O_2$ 474.601


(±)-**form** [98260-39-0]

Alkaloid from *Murraya euchrestifolia* (Rutaceae).

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 2611 (*uv*, *pmr*, *synth*)

Murrafoline E M-745

3,11-Dihydro-5-[(1-methoxy-3-methyl-9H-carbazol-9-yl)methyl]-3,3-dimethyl-pyrano[3,2-a]carbazole, 9CI
[117591-98-7]




$C_{32}H_{28}N_2O_2$ 472.585

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1988, **36**, 2377 (isol, uv, ir, pmr, ms, struct)

Murrafoline F M-746


1-Methoxy-2-[(9-methoxy-9H-carbazol-3-yl)methyl]-3-methyl-9H-carbazole, 9CI
[117592-00-4]



$C_{28}H_{24}N_2O_2$ 420.51

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1988, **36**, 2377 (isol, uv, ir, pmr, cmr, ms, struct)

Murrafoline G M-747

$C_{32}H_{30}N_2O_2$ 474.601


(±)-form [98260-40-3]

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Furukawa, H. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1249 (isol, uv, pmr, cmr, struct)

Murrafoline H M-748

1,2,3,3',11,11'-Hexahydro-3,3,3',5,5,5'-hexamethyl-1,8'-bipyrano[3,2-a]carbazole



$C_{36}H_{34}N_2O_2$ 526.677


(±)-form [98260-36-7]

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Furukawa, H. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1249 (isol, uv, pmr, ms, struct)

Murranimbine M-749

[161068-68-4]




$C_{36}H_{34}N_2O_2$ 526.677

Unsymmetrical dimer of Girinimbine, G-81. Alkaloid from *Murraya euchrestifolia* (Rutaceae). Pale yellow oil. Racemic.

Ito, C. et al., *Chem. Pharm. Bull.*, 1991, **39**, 1355 (isol, pmr, cmr, struct)

Murrapanine M-750

8-(1H-Indol-3-yl)-2-methoxy-6-methyl-1,4-naphthalenedione, 9CI
[128508-22-5]



$C_{20}H_{15}NO_3$ 317.343

First example of a naturally occurring indole-naphthoquinone alkaloid. Alkaloid from the root bark of *Murraya paniculata* var. *omphalocarpa* (Rutaceae). Exhibits significant cytotoxicity. Deep purplish prisms (Et₂O). Mp 278-280°. λ_{max} 221 (ε 36300); 246 (ε 17800); 281 (ε 20900); 323 (ε 5370) (MeOH) (Derep).


Wu, T.-S. et al., *Tet. Lett.*, 1989, **30**, 6649 (isol, uv, ir, pmr, cmr, ms, synth, cryst struct)
Sheu, J.-H. et al., *J.O.C.*, 1993, **58**, 5784 (synth)

Sheu, J.H. et al., *Heterocycles*, 1996, **43**, 1751 (synth)

Sheu, J.H. et al., *J.C.S. Perkin 1*, 1998, 1959-1965 (synth)

Murrastifoline A M-751

1',8-Dimethoxy-3',6-dimethyl-3,9'-bi-9H-carbazole, 9CI
[129748-48-7]



$C_{28}H_{24}N_2O_2$ 420.51


The N,6'-dimer of 1-Hydroxy-3-methyl-9H-carbazole, H-583. Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)

Kitawaki, T. et al., *Heterocycles*, 2005, **65**, 1561-1567 (synth)

Murrastifoline B M-752

1,1'-Dimethoxy-3'-methyl-3,9'-bi-9H-carbazole, 9CI
[129748-49-8]




$C_{27}H_{22}N_2O_2$ 406.483

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil.

Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)


Murrastifoline C M-753

3,11-Dihydro-5-[(1-methoxy-3-methyl-9H-carbazol-9-yl)methyl]-3-methyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole, 9CI
[129777-58-8]



C₃₇H₃₆N₂O₂ 540.704Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil. Racemic.Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143 (isol, uv, ir, pmr, ms, struct)**Murrastifoline D** M-754


1,2,3,11-Tetrahydro-1-(1-methoxy-3-methyl-9H-carbazol-9-yl)-3,3,5-trimethylpyrano[3,2-a]carbazol-2-ol, 9CI [129777-59-9]

C₃₂H₃₀N₂O₃ 490.601Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil. Racemic.3'-Deoxy: **Murrastifoline E**


[133740-45-1]

C₃₂H₃₀N₂O₂ 474.601Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Oil. [α]_D -5.7 (c, 0.035 in CHCl₃).Ito, C. et al., *Chem. Pharm. Bull.*, 1990, **38**, 1143; 1548 (isol, uv, ir, pmr, cmr, ms, struct)**Murrastifoline F** M-755


1,1'-Dimethoxy-3,3'-dimethyl-4,9'-bi-9H-carbazole, 9CI [155519-85-0]

C₂₈H₂₄N₂O₂ 420.51Alkaloid from roots of *Murraya koenigii* (curryleaf tree) (Rutaceae). Oil. Isol. as a mixture of atropisomers.Ito, C. et al., *Chem. Pharm. Bull.*, 1993, **41**, 2096 (isol, uv, ir, pmr, ms, struct)Bringmann, G. et al., *J.A.C.S.*, 2001, **123**, 2703-2711 (synth, ed)**Murrayacarine** M-756

[125288-14-4]

C₁₄H₁₃NO₃ 243.262Alkaloid from root bark of *Murraya**paniculata* var. *omphalocarpa*. Pale yellow needles (CHCl₃). Mp 146-148°. λ_{max} 208 (log ε 4.17); 247 (log ε 3.73); 268 (log ε 3.68); 275 (sh) (log ε 3.65) (MeOH).Wu, T.-S. et al., *Phytochemistry*, 1989, **28**, 2873-2874 (isol, uv, pmr, ms)Boumendjel, A. et al., *Bull. Soc. Chim. Fr.*, 1990, 645 (synth, cmr, config)Johnson, A.-L. et al., *Heterocycles*, 2006, **68**, 2165-2170 (synth)**Murrayacine** M-757

3,11-Dihydro-3,3-dimethylpyrano[3,2-a]carbazole-5-carboxaldehyde, 9CI [27300-29-4]

C₁₈H₁₅NO₂ 277.322CAS numbering shown. Alternative numbering freq. used in which C-9 becomes C-7. Alkaloid from the stem bark of *Murraya koenigii* (curryleaf tree) and the roots of *Clausena heptaphylla* (Rutaceae). Cryst. (EtOAc). Mp 244-245°. Bp_{0.05} 180° subl. λ_{max} 226 (log ε 4.6); 282 (log ε 4.57); 301 (log ε 4.58) (EtOH).

2,4-Dinitrophenylhydrazone: Mp 280°.

1,2-Dihydro: Cycloheptaphylline. Dihydromurrayacine [17750-37-7]

Cryst. (MeOH). Mp 250° (176°).

1,2-Dihydro, 2ξ,10-dihydroxy: **Clauszoline E** [185508-05-8]C₁₈H₁₇NO₄ 311.337Alkaloid from the stem bark of *Clausena excavata*. Yellow powder. [α]_D²⁵ 0 (c, 0.1 in CHCl₃). λ_{max} 204; 213; 242; 273; 292; 358 (MeOH).10-Hydroxy: **Clauszoline G**

[185508-07-0]

C₁₈H₁₅NO₃ 293.321Alkaloid from the stem bark of *Clausena excavata*. Yellow powder. λ_{max} 204; 234; 245 (sh); 278; 300; 361 (MeOH).10-Hydroxy, 1,2-dihydro: **Heptazolicine**.

Cycloheptazoline

[32042-36-7]

C₁₈H₁₇NO₃ 295.337Alkaloid from the roots of *Clausena heptaphylla* (Rutaceae). Cryst.(CHCl₃). Mp 285° dec. λ_{max} 242 (log ε 4.65); 275 (log ε 4.62); 300 (log ε 4.4) (EtOH).


10-Hydroxy, 1,2-dihydro, O-Ac:

Cryst. (C₆H₆). Mp 230°.9-Methoxy: **7-Methoxymurrayacine**


[132160-51-1]

C₁₉H₁₇NO₃ 307.348Alkaloid from *Murraya siamensis* (Rutaceae). Mp 211-213°. Numbering systems vary. λ_{max} 306 (log ε 4.58); 354 (log ε 4.09) (EtOH).Chakraborty, D.P. et al., *J.O.C.*, 1971, **36**, 725-727; 1973, **38**, 2728-2729 (isol, uv, ir, pmr, ms, struct, synth)Anwer, F. et al., *Indian J. Chem.*, 1973, **11**, 1314-1315 (synth)Oikawa, Y. et al., *Heterocycles*, 1976, **5**, 233-238 (synth, ms)Roy, S. et al., *Phytochemistry*, 1976, **15**, 356 (isol, uv, ir)Bhattacharyya, P. et al., *Phytochemistry*, 1984, **23**, 2409-2410 (Heptazolicine, isol, uv, ir, pmr, ms, struct)Chakraborty, D.P. et al., *J. Indian Chem. Soc.*, 1985, **62**, 602-603 (Heptazolicine, synth)Ruangrunsi, N. et al., *J. Nat. Prod.*, 1990, **53**, 946-952 (7-Methoxymurrayacine)Ito, C. et al., *Chem. Pharm. Bull.*, 1996, **44**, 2231-2235 (Clauszoline)Knölker, H.J. et al., *Tet. Lett.*, 1996, **37**, 7947-7950 (synth)**Murrayacatine** M-758


2,3-Epoxy-2,3-dihydro-2-hydroxy-1H-indole-3-acetic acid. 1a,2-Dihydro-1a-hydroxy-6bH-oxireno[b]indole-6b-acetic acid, 9CI [159934-09-5]

C₁₀H₉NO₄ 207.185Alkaloid from fresh flowers of *Murraya paniculata* (Rutaceae). Needles (Me₂CO). Mp 168-170°. [α]_D +17.2 (c, 0.027 in CHCl₃).Wu, T.-S. et al., *Phytochemistry*, 1994, **37**, 287 (isol, uv, ir, pmr, cmr, ms, struct)**Murrayafoline B** M-759

7-Methoxy-3-methyl-8-(3-methyl-2-but-1-enyl)-9H-carbazol-1-ol, 9CI. 1-Hydroxy-7-methoxy-3-methyl-8-prenylcarbazole [100108-68-7]

C₁₉H₂₁NO₂ 295.38Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Syrup.Furukawa, H. et al., *Chem. Pharm. Bull.*, 1985, **33**, 4132 (isol, uv, ir, pmr, ms, struct)**Murrayaline C** M-760


7-Hydroxy-2-methoxy-9H-carbazole-1,6-dicarboxaldehyde [139726-43-5]

C₁₅H₁₁NO₄ 269.256

Alkaloid from the stem bark of *Murraya euchrestifolia*. Pale yellow powder. λ_{\max} 239 (sh) ; 251 (sh) ; 267 ; 307 (sh) ; 329 (MeOH).

Ito, C. et al., *Chem. Pharm. Bull.*, 1991, **39**, 2525-2528

Murrayamine F **M-761**
[183130-86-1]



$C_{24}H_{27}NO_2$ 361.483
Alkaloid from leaves of *Murraya euchrestifolia*. Oil. λ_{\max} 221 ; 244 ; 256 (sh) ; 303 ; 316 (sh) ; 328 (sh) (MeOH).

Demethoxy: Murrayamine G


[183130-89-4]

$C_{23}H_{25}NO$ 331.457

From leaves of *Murraya euchrestifolia*. Needles (MeOH). Mp 173-176°. λ_{\max} 221 ; 241 ; 262 ; 303 ; 307 (MeOH).

Wu, T.-S. et al., *Phytochemistry*, 1996, **43**, 785 (isol, uv, ir, pmr, ms, struct)

Murrayamine J **M-762**
[175669-20-2]



$C_{23}H_{23}NO_2$ 345.44
Alkaloid from leaves of *Murraya euchrestifolia*. Oil.


1-Methoxy: Murrayamine N

$C_{24}H_{25}NO_3$ 375.466

From leaves of *Murraya euchrestifolia*. Oil. Erroneously descr. as C_{23} in paper.

Wu, T.-S. et al., *Phytochemistry*, 1996, **41**, 1433-1435 (isol, uv, ir, pmr, ms, struct)

Murrayamine M **M-763**
[175669-22-4]




$C_{23}H_{23}NO_2$ 345.44

Closely related to Bicyclomahanimbine, B-122. Alkaloid from leaves of *Murraya euchrestifolia*. Oil.

Wu, T.-S. et al., *Phytochemistry*, 1996, **41**, 1433 (isol, uv, ir, pmr, ms, struct)

Murrayamine O **M-764**
[166990-10-9]



$C_{23}H_{27}NO_2$ 349.472
Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Plates (MeOH). Mp 130-132°. $[\alpha]_D$ -137.6 (c, 0.07 in $CHCl_3$).

3'-Epimer: Murrayamine P


[167077-71-6]

$C_{23}H_{27}NO_2$ 349.472

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Plates (MeOH). Mp 116-118°. $[\alpha]_D$ +92 (c, 0.015 in $CHCl_3$).

Wu, T.-S. et al., *Tet. Lett.*, 1995, **36**, 5385 (isol, uv, ir, pmr, cmr, ms, struct)

Murrayanine† **M-765**
[500725-15-5]



Relative Configuration


$C_{30}H_{31}NO_7$ 517.577

Not to be confused with Murrayanine in H-447. Alkaloid from *Murraya koenigii*. Brown powder. $[\alpha]_D^{25}$ +8 (c, 0.74 in MeOH). λ_{\max} 206 ; 225 ; 238 ; 300 ; 342 (MeOH).

Wang, Y.-S. et al., *J. Nat. Prod.*, 2003, **66**, 416-418 (isol, pmr, cmr, ms)

Murrayaquinone B **M-766**

7-Methoxy-3-methyl-8-(3-methyl-2-butenyl)-1H-carbazole-1,4(9H)-dione, 9CI. 7-Methoxy-3-methyl-8-prenyl-1,4-carbazolequinone [86695-16-1]



$C_{19}H_{19}NO_3$ 309.364
Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Also obt. by photooxidn. of Murrayafoline B. Deep

purple needles (Me_2CO). Mp 221-223°.

O-De-Me: Murrayaquinone E

[139726-47-9]

$C_{18}H_{17}NO_3$ 295.337

Alkaloid from the stem bark of *Murraya euchrestifolia*. Brown oil. λ_{\max} 232 ; 262 ; 406 (MeOH).

Wu, T.-S. et al., *Heterocycles*, 1983, **20**, 1267 (uv, ir, pmr, cmr, struct)


Furukawa, H. et al., *Chem. Pharm. Bull.*, 1985, **33**, 4132 (isol, uv, ir, pmr, cmr, ms, synth, struct)

Martin, T. et al., *J.C.S. Perkin I*, 1988, 241 (synth, uv, ir, pmr, ms)

Ito, C. et al., *Chem. Pharm. Bull.*, 1991, **39**, 2525-2528 (Murrayaquinone E)

Murrayaquinone D **M-767**

8-(3,7-Dimethyl-2,6-octadienyl)-7-hydroxy-3-methyl-1H-carbazole-1,4(9H)-dione, 9CI. 8-Geranyl-7-hydroxy-3-methylcarbazole-1,4-quinone [100108-67-6]



$C_{23}H_{25}NO_3$ 363.455

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Violet needles. Mp 164-168°.

Me ether: Murrayaquinone C. 8-(3,7-Dimethyl-2,6-octadienyl)-7-methoxy-3-methyl-1H-carbazole-1,4(9H)-dione, 9CI

[86695-17-2]

$C_{24}H_{27}NO_3$ 377.482

Alkaloid from root bark of *Murraya euchrestifolia* (Rutaceae). Violet needles (Et_2O). Mp 158-159°.


Wu, T.S. et al., *Heterocycles*, 1983, **20**, 1267 (occur, pmr, struct)

Furukawa, H. et al., *Chem. Pharm. Bull.*, 1985, **33**, 4132 (isol, uv, ir, pmr, ms, struct, synth)

Muscaaurin I **M-768**

4-[[[Carboxy(2,3-dihydro-3-oxo-5-isoxazolyl)methyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI

[52012-51-8]



C₁₄H₁₃N₃O₈ 351.272


Pigment from *Amanita muscaria*. Hydrol. gives Betalamic acid, B-109 and Ibotenic acid, I-8. λ_{\max} 245 (sh); 295 (sh); 465; 475 (no solvent reported).

Doeppe, H. *et al.*, *Chem. Ber.*, 1973, **106**, 3473 (isol, uv, cd)

Doeppe, H. *et al.*, *Annalen*, 1982, 254 (struct)

Muscaaurin II M-769

4-[[[1-Carboxy-2-(6-carboxy-2-oxo-2H-pyran-4-yl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI
[12624-17-8]

C₁₈H₁₆N₂O₁₀ 420.332


Pigment from *Amanita muscaria*. λ_{\max} 478, 462 sh, 301, 261, λ_{\min} 360, 278, 242 nm. Hydrol. gives Betalamic acid and Stizolobic acid. Further Muscaaurins having different residues are descr. but owing to misprints it is not clear what their structs. are and they are not indexed by CA.

Doeppe, H. *et al.*, *Chem. Ber.*, 1973, **106**, 3473 (isol, uv, cd)

Doeppe, H. *et al.*, *Annalen*, 1982, 254 (struct)

Muscaflavin M-770

4-Formyl-2,3-dihydro-1H-azepine-2,7-dicarboxylic acid, 9CI
[12624-18-9]

C₉H₉NO₅ 211.174

Unstable pigment from the skin of *Amanita muscaria*. Yellow cryst. Sol. H₂O. An incorrect struct. was previously assigned. λ_{\max} 238; 420 (H₂O) (Berdy).

Di-Me ester: [70991-04-7]


Cryst. (MeOH/Et₂O). Mp 98-98.5°.

Barth, H. *et al.*, *Annalen*, 1981, 2164 (struct, synth)

Muscapurpurin

[12624-19-0]

M-771

C₁₈H₁₄N₂O₁₀ 418.316


The struct. shown is given in CA but there are no refs. on the CA database to support it. Violet pigment from *Amanita muscaria*. λ_{\max} 540, 303, λ_{\min} 380, 262 nm.

Döpp, H. *et al.*, *Chem. Ber.*, 1973, **106**, 3473 (isol, uv, ir, cd)

Muscaridine

M-772

4,5-Dihydroxy-N,N,N-trimethyl-1-hexanaminium(1+), 9CI

C₉H₂₂NO₂[⊕] 176.278

Present in *Amanita muscaria*. Mp 129-131° dec. (as tetrachloroaurate). $[\alpha]_D^{20}$ +20.5 (tetrachloroaurate). Stereochem. of the nat. prod. could not be confirmed. It was claimed to have the erythro-config. but props. do not coincide with those of the well-characterised synthetic stereoisomers later obt.

(2R,3R)-form

D-threo-form

Iodide: Mp 98-100°. $[\alpha]_D$ +14.2 (c, 2.8 in H₂O).

Tetrachloroaurate: Mp 98-101°. $[\alpha]_D^{20}$ +6.2 (c, 0.5 in H₂O).

(2R,3S)-form

D-erythro-form

Iodide:

C₉H₂₂INO₂ 303.183

Mp 101-103°. $[\alpha]_D$ +14.7 (c, 2.3 in H₂O).

Tetrachloroaurate: Mp 128-130°. $[\alpha]_D$ -9.

(2S,3R)-form

L-erythro-form

Iodide: Mp 101-103°. $[\alpha]_D^{20}$ +14.7 (c, 1 in H₂O).

(2S,3S)-form

L-threo-form

Iodide: Mp 97-98°. $[\alpha]_D$ -14.9 (c, 1.2 in H₂O).

(2RS,3RS)-form

(±)-threo-form

Tetrachloroaurate: Mp 97-102°.

(2RS,3SR)-form

(±)-erythro-form

[6801-43-0]

Tetrachloroaurate: Mp 127-131° dec.


Salemink, C.A. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1963, **82**, 21 (isol, bibl, ir, synth, resoln)

Pedersen, C. *et al.*, *Acta Chem. Scand.*, 1993, **47**, 885 (synth, cmr, bibl)

Muscarine†

M-773

Tetrahydro-4-hydroxy-N,N,N,5-tetramethyl-2-furanmethanaminium(1+), 9CI
[7619-12-7]

C₉H₂₀NO₂[⊕] 174.262

All 8 stereoisomers have been synthesised.

(2S,4R,5S)-form [300-54-9]

Toxic constit. of the fly fungus *Amanita muscaria* and various *Inocybe*, *Boletus* and *Clitocybe* spp. Cholinergic agent. V. powerful parasymphathomimetic agent causing hypotension and bronchoconstriction. Sol. EtOH; fairly sol. CHCl₃, Et₂O. V. hygroscopic. Stereoisomers show only a fraction of the biol. activity.

▶ V. toxic if swallowed. Can cause convulsions, coma, death. LD₅₀ (mus, ivn) 0.23 mg/kg. Antidote atropine sulfate.

QG3325000

Chloride: [2303-35-7]

C₉H₂₀ClNO₂ 209.715

Needles. Mp 181.5-182°, (179-180°).

$[\alpha]_D$ +7.4 (c, 3.1 in H₂O).

▶ QG3500000

Iodide: [24570-49-8]

C₉H₂₀INO₂ 301.167

Needles (MeCN/Me₂CO). Mp 149-

149.5°. $[\alpha]_D$ +6.3 (c, 1.1 in H₂O). Turns yellow within a few hours.

Tetraphenylborate: [104487-60-7]

C₃₃H₄₀BNO₂ 493.495

Mp 193°. $[\alpha]_D$ +9.7 (c, 1.34 in Me₂CO).

Ketone: Tetrahydro-N,N,N,5-tetramethyl-4-oxo-2-furanmethanaminium(1+).

Muscarone

[4780-69-2]

C₉H₁₈NO₂[⊕] 172.247

Acetylcholine mimic.

(2R,4R,5S)-form

5-Allomuscarine

[6252-44-4]

[35119-38-1]

Constit. of numerous Agaricales incl. *Amanita* spp. and *Clitocybe* spp. Shows similar biol. activity to Muscarine. Mp 130.8° (as iodide). $[\alpha]_D$ -37.7 (c, 0.8 in H₂O).

(2R,4S,5R)-form [92981-62-9]

[35119-35-8]

Mp 179–180° (as chloride). $[\alpha]_D$ -8.4 (EtOH).**(2R,4S,5S)-form**

3-Epi-5-allomuscarine

[6836-06-2]

[5487-32-1]

Mp 199.2° (as iodide). $[\alpha]_D$ 0 (c, 1 in H₂O).**(2S,4S,5S)-form**

3-Epimuscarine

[6836-08-4]

[93226-49-4]

Constit. of various Agaricales incl. *Amanita* spp. and *Clitocybe* spp. Cryst. (2-propanol) (as iodide). Mp 170.4° (iodide). $[\alpha]_D$ +43.2 (c, 0.6 in EtOH).**(2RS,4SR,5RS)-form** [71-06-7]

[6032-87-7]


Mp 119–120° (as tetrachloroaurate).

[2936-25-6]

Hardegger, E. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 2383–2389 (2S,4R,5S-form, synth, abs config)Wilkinson, S. *et al.*, *Q. Rev., Chem. Soc.*, 1961, **15**, 153–171 (rev)Eugster, C.H. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 708–715 (occur, glc)Matsumoto, T. *et al.*, *Tetrahedron*, 1969, **25**, 5889–5892 ((±)-forms, synth)Bollinger, H. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 2704–2730 (synth, stereoisomers, abs config, cd)Whiting, J. *et al.*, *Can. J. Chem.*, 1972, **50**, 3322–3325 (synth, pmr)Stadelmann, R.J. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2432–2436 (occur, stereoisomers)Nitta, K. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1747–1753 (biosynth)Wang, P.-C. *et al.*, *Alkaloids (Academic Press)*, 1984, **23**, 327–380 (rev)Mulzer, J. *et al.*, *Annalen*, 1987, 7–14Pirrung, M.C. *et al.*, *Tet. Lett.*, 1988, **29**, 159–162 (synth, bibl)Takano, S. *et al.*, *Chem. Comm.*, 1989, 1371–1372 (2S,4R,5S-form, synth)Adams, J. *et al.*, *Tet. Lett.*, 1989, **30**, 1753 (synth, pmr, cmr)De Amici, M. *et al.*, *J.O.C.*, 1991, **56**, 67–72 (synth, stereoisomers, bibl)Frydenvang, K. *et al.*, *Acta Cryst. C*, 1992, **48**, 469; 1993, **49**, 985; 1834 (cryst struct, Muscarone)Chan, T.H. *et al.*, *Can. J. Chem.*, 1992, **70**, 2726–2729 (2S,4R,5S-form, synth)Mantell, S.J. *et al.*, *J.C.S. Perkin 1*, 1992, 3023–3027 (2S,4R,5S-form synth, bibl)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1391Knight, D.W. *et al.*, *Synlett*, 1994, 295–296 ((-)-Muscarine, synth)Norild, J.C. *et al.*, *Synthesis*, 1997, 1128–1130 (Allomuscarine)Popsavin, V. *et al.*, *Carbohydr. Lett.*, 1998, **3**, 1–8 (synth, Epimuscarine)Hartung, J. *et al.*, *Eur. J. Org. Chem.*, 2000, 1677–1683 (synth)Kang, K.H. *et al.*, *Tet. Lett.*, 2000, **41**, 8137–8140 (synth)Popsavin, V. *et al.*, *Tetrahedron*, 2000, **56**, 5929–5940 (synth)Knight, D.W. *et al.*, *Tet. Lett.*, 2002, **43**, 6771–6773 (synth)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, MRW250**Muscopyridine**

M-774

3-Methyl-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene, 9CI

C₁₆H₂₅N 231.38

CA numbering shown.

(R)-form [501-08-6]Alkaloid from the scent gland of the musk deer (*Moschus moschiferus*). Liq. Bp₁₂ 155–160°. $[\alpha]_D^{23}$ +17.4 (c, 1.92 in CHCl₃).

Picrolonate: Mp 163–166° dec.

11-Hydroxy: Hydroxymuscopyridine A

[89368-39-8]

C₁₆H₂₅NO 247.38Alkaloid from *Moschus moschiferus*.**2-Hydroxy: Hydroxymuscopyridine B**

[89368-40-1]


C₁₆H₂₅NO 247.38Alkaloid from *Moschus moschiferus*.**(±)-form** [56912-83-5]

Synthetic. Mp 163–166° dec. (as picrolonate).

Schinz, H. *et al.*, *Helv. Chim. Acta*, 1946, **29**, 1524 (isol, uv)Biemann, K. *et al.*, *J.A.C.S.*, 1957, **79**, 5558 (synth, struct)Tamao, K. *et al.*, *J.A.C.S.*, 1975, **97**, 4405 (synth)Hiyama, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 2747 (synth)Yu, D. *et al.*, *Planta Med.*, 1983, **49**, 183 (Hydroxymuscopyridines)Hadj-Abo, F. *et al.*, *Helv. Chim. Acta*, 1992, **75**, 1834 (synth)Olofsson, K. *et al.*, *J.O.C.*, 2000, **65**, 7235–7239 (synth, pmr, cmr, abs config)Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 308–311 (synth)Suwa, K. *et al.*, *Tet. Lett.*, 2008, **49**, 1510–1513 (synth)**Muscorigide A**

M-775


[164672-63-3]

C₂₈H₄₀N₄O₅ 512.648Alkaloid from the terrestrial freshwater cyanobacterium *Nostoc muscorum*.Shows weak antibacterial activity. Amorph. solid. $[\alpha]_D^{23}$ -89.4 (c, 0.70 in MeOH). λ_{max} 252 (ε 14600) (MeOH) (Derep). λ_{max} (MeOH) (Berdy).Nagatsu, A. *et al.*, *Tet. Lett.*, 1995, **36**, 4097 (isol, ir, pmr, cmr, struct)Muir, J.C. *et al.*, *Synthesis*, 1998, 613–618

(synth)

Coqueron, P.-Y. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 1411–1414 (synth)**Mycalazoles**

M-776



Mycalazole

R

1	(CH ₂) ₅ (CH=CHCH ₂) ₆ CH ₃
2	(CH ₂) ₁₀ (CH=CHCH ₂) ₃ CH ₃
3	(CH ₂) ₉ (CH=CHCH ₂) ₄ CH ₃
4	(CH ₂) ₁₂ (CH=CH(CH ₂) ₃)CH ₃
7	(CH ₂) ₁₄ CH(CH ₃) ₂
10	(CH ₂) ₁₂ CH=CH(CH ₂) ₅ CH ₃

All double bonds have Z-config. Metabs. from the sponge *Mycale micracanthoxea*. Cytotoxic agents.**Mycalazole 1**

1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-7,10,13,16,19,22-pentacosahexaen-1-one, 9CI

[185387-05-7]

C₃₀H₄₃NO₂ 449.675Oil. λ_{max} 201 (ε 4170); 247 (ε 1581); 294 (ε 7656) (MeOH).7',8'-Dihydro: 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-10,13,16,19,22-pentacosapentaen-1-one, 9CI. **Mycalazole 5** [185389-23-5]C₃₀H₄₅NO₂ 451.691Oil. λ_{max} 204 (ε 11230); 250 (ε 2122); 296 (ε 10003) (MeOH).7',8',10',11'-Tetrahydro: 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-13,16,19,22-pentacosatetraen-1-one, 9CI. **Mycalazole 9**

[185389-50-8]

C₃₀H₄₇NO₂ 453.707Oil. λ_{max} 203 (ε 10145); 251 (ε 1580); 293 (ε 13110) (MeOH).**Mycalazole 2**

1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-12,15,18-heneicosatrien-1-one, 9CI [185387-07-9]

C₂₆H₄₁NO₂ 399.615Oil. λ_{max} 202 (ε 8490); 238 (ε 6912); 296 (ε 16584) (MeOH).18',19'-Dihydro: 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-12,15-heneicosadien-1-one, 9CI. **Mycalazole 6**

[185389-27-9]

C₂₆H₄₃NO₂ 401.631Oil. λ_{max} 204 (ε 7452); 294 (ε 9470) (MeOH).**Mycalazole 3**

1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-11,14,17,20-tricosatetraen-1-one, 9CI [185388-42-5]

C₂₈H₄₃NO₂ 425.653Oil. λ_{max} 206 (ε 11045); 246 (ε 2332); 295 (ε 10106) (MeOH).

11',12'-Dihydro- 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-14,17,20-tricosatrien-1-one, 9CI. Mycalazole 8
[185389-47-3]
C₂₈H₄₅NO₂ 427.669
Oil. λ_{max} 201 (ε 9128); 240 (ε 4312); 295 (ε 16348) (MeOH).

11',12',20',21'-Tetrahydro- 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-14,17-tricosadien-1-one, 9CI. Mycalazole 12
[185389-63-3]
C₂₈H₄₇NO₂ 429.685
Oil. λ_{max} 203 (ε 7852); 295 (ε 8864) (MeOH).

Mycalazole 4

1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-14-nonadecen-1-one, 9CI
[185389-09-7]

C₂₄H₄₁NO₂ 375.593
Cryst. (MeOH). Mp 68-70°. λ_{max} 206 (ε 10492); 252 (ε 1106); 292 (ε 13105) (MeOH).

14',15'-Dihydro- 1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-1-nonadecanone, 9CI. Mycalazole 11
[185389-61-1]
C₂₄H₄₃NO₂ 377.609
Amorph. powder. λ_{max} 204 (ε 7452); 294 (ε 9470) (MeOH).

Mycalazole 7

1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-16-methyl-1-heptadecanone, 9CI
[185389-45-1]

C₂₃H₄₁NO₂ 363.582
Oil. λ_{max} 204 (ε 7452); 294 (ε 9470) (MeOH).

Mycalazole 10


1-[5-(Hydroxymethyl)-1H-pyrrol-2-yl]-14-heneicosen-1-one, 9CI
[185389-60-0]

C₂₆H₄₅NO₂ 403.647
Cryst. (MeOH). Mp 74-76°. λ_{max} 204 (ε 10978); 252 (ε 925); 296 (ε 12940) (MeOH).

Ortega, M.J. *et al.*, *Tetrahedron*, 1997, **53**, 331-340 (*isol, uv, ir, pmr, cmr, ms, struct*)
Nabbs, B.K. *et al.*, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 505-508 (*Mycalazole 11, synth*)
Hansen, T.V. *et al.*, *Tet. Lett.*, 2004, **45**, 2809-2811 (*Mycalazole 5, synth*)

Mycalexime

[290825-58-0]




C₂₂H₃₄N₂O₃ 374.522
Isol. from the sponge *Mycale tenuispiculata*. Viscous liq. λ_{max} 232 (ε 8228); 304 (ε 19640) (MeCN).

Venketesham, U. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1318-1320 (*isol, ir, uv, pmr, cmr, ms*)

Mycalisine B

5-Cyano-7-(3-O-methyl-5-deoxy-β-D-erythro-pent-4-enofuranosyl)pyrrolo[2,3-d]pyrimidin-4-one
[98890-72-3]




C₁₃H₁₂N₄O₄ 288.262
Isol. from *Mycale* sp., Japanese marine sponge. Oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²¹ -75.9 (c, 0.26 in EtOH). Unstable at r.t. λ_{max} 214 (ε 3000); 265 (ε 3000) (pH 2 EtOH) (Derep). λ_{max} 265 (sh) (ε) (pH 12 EtOH) (Derep). λ_{max} 210 (ε 6000); 265 (ε 3000) (pH 7 EtOH) (Derep).

Kato, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 3483 (*isol, ir, pmr, cmr, uv*)

Mycalolide A

[121038-36-6]



C₄₇H₆₄N₄O₁₄ 909.041
Isol. from the sponge *Mycale* sp. Cytotoxic and antifungal agent. Yellowish gum. Sol. MeOH, Et₂O; poorly sol. H₂O. [α]_D -60.3 (c, 0.5 in CHCl₃). λ_{max} 230 (ε 30000) (MeOH) (Derep).

O-De-Ac: 32-Hydroxymycalolide A

[216770-61-5]
C₄₅H₆₂N₄O₁₃ 867.004
Isol. from *Mycale magellanica*. Cytotoxic agent. [α]_D²⁷ -90 (c, 0.1 in MeOH). Misleading synonym. λ_{max} 230 (ε 26000) (MeOH).

30R-Alcohol: 30-Hydroxymycalolide A

[216770-57-9]
C₄₇H₆₆N₄O₁₄ 911.057
Isol. from *Mycale magellanica*. Cytotoxic agent. [α]_D²⁷ -86.9 (c, 0.1 in MeOH). Misleading synonym. λ_{max} 230 (ε 27000) (MeOH).

30R-Alcohol, 30-O-(2R,3-dimethoxyprop-2-yl): Mycalolide B

[122752-21-0]
C₅₂H₇₄N₄O₁₇ 1027.173
Isol. from the sponge *Mycale* sp. Cytotoxic and antifungal agent. Yellowish gum. [α]_D -53.4 (c, 1.3 in CHCl₃). λ_{max} 230 (ε 31000) (MeOH).

30R-Alcohol, 30-O-(3R-hydroxy-2-methoxyprop-2-yl): 38-Hydroxymycalolide B

M-778

[216770-65-9]
C₅₁H₇₂N₄O₁₇ 1013.146
Isol. from the sponge *Mycale magellanica*. Cytotoxic agent. [α]_D²⁷ -80.9 (c, 0.1 in MeOH). Misleading synonym. λ_{max} 230 (ε 25000) (MeOH).

30R-Alcohol, 30-O-(2R-methoxyprop-2-yl): Mycalolide C

[122752-20-9]
C₅₁H₇₂N₄O₁₆ 997.147
Isol. from the sponge *Mycale* sp. and from the stony coral *Tubastrea faulkneri*. Cytotoxic and antifungal agent. Yellowish gum. [α]_D -62.1 (c, 3.7 in CHCl₃). λ_{max} 230 (ε 31000) (MeOH).

30R-Alcohol, O-de-Ac: 30,32-Dihydroxymycalolide A

[442690-43-9]
C₄₅H₆₄N₄O₁₃ 869.02
Isol. from *Mycale izuensis*. [α]_D²² -68.7 (c, 0.1 in MeOH). λ_{max} 231 (ε 33055) (MeOH).

Fusetani, N. *et al.*, *Tet. Lett.*, 1989, **30**, 2809-2812 (*isol, uv, ir, pmr, cmr, struct*)


Rashid, M.A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1120-1125 (*Mycalolides C-E, isol*)
Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1164-1167 (*Mycale magellanica constits*)
Matsunaga, S. *et al.*, *J.A.C.S.*, 1999, **121**, 5605-5606; 8969 (*abs config*)

Liu, P. *et al.*, *J.A.C.S.*, 2000, **122**, 1235-1236 (*synth*)

Phuwapraisirisan, P. *et al.*, *J. Nat. Prod.*, 2002, **65**, 942-943 (*30,32-Dihydroxymycalolide A*)

Mycalolide D

[168301-18-6]




C₅₀H₇₂N₄O₁₇ 1001.135
Isol. from the stony coral *Tubastrea faulkneri*. Cytotoxic. Gum. [α]_D -19.5 (c, 0.5 in CHCl₃). λ_{max} 228 (ε 47700); 260 (MeOH) (Berdy).

Rashid, M.A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1120 (*isol, uv, ir, pmr, cmr, struct*)

Mycalolide E

[168301-19-7]




C₄₆H₆₂N₄O₁₃ 879.015
Minor const. of the stony coral *Tubastrea faulkneri*. Cytotoxic. Gum. [α]_D -39 (c, 0.1 in CHCl₃). λ_{max} 231 (ε 33200); 254 (ε 30800) (MeOH) (Berdy).

Rashid, M.A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1120 (*isol, uv, ir, pmr, struct*)

Mycapolyols

M-782



Isol. from the marine sponge *Mycalca izuensis*. Cytotoxic.

Mycapolyol A

$C_{55}H_{94}N_2O_{18}$ 1071.35
Powder. $[\alpha]_D^{21} +101.1$ (c, 0.34 in MeOH aq.). λ_{max} 259 (log ϵ 4.15) (MeOH aq.).

Mycapolyol B

$C_{53}H_{90}N_2O_{17}$ 1027.297
Powder. $[\alpha]_D^{27} +104.7$ (c, 0.22 in MeOH aq.). λ_{max} 259 (log ϵ 4.6) (MeOH aq.).

Mycapolyol C

$C_{51}H_{86}N_2O_{16}$ 983.244
Powder. $[\alpha]_D^{27} +105.7$ (c, 0.32 in MeOH aq.). λ_{max} 259 (log ϵ 4.09) (MeOH aq.).

Mycapolyol D

$C_{49}H_{82}N_2O_{15}$ 939.191
Powder. $[\alpha]_D^{27} +100.6$ (c, 0.25 in MeOH aq.). λ_{max} 259 (log ϵ 4.18) (MeOH aq.).

Mycapolyol E

$C_{47}H_{78}N_2O_{14}$ 895.138
Powder. $[\alpha]_D^{21} +117.7$ (c, 0.27 in MeOH aq.). λ_{max} 258 (log ϵ 4.15) (MeOH aq.).

Mycapolyol F


$C_{45}H_{74}N_2O_{13}$ 851.085
Powder. $[\alpha]_D^{21} +120.1$ (c, 0.2 in MeOH aq.). λ_{max} 257 (log ϵ 4) (MeOH aq.).

Phuwapraisirisan, P. *et al.*, *Org. Lett.*, 2005, 7, 2233-2236 (isol, pmr, cmr, ms)

Myceliothermophin A

M-783

[955083-86-0]



$C_{26}H_{39}NO_3$ 413.599
Prod. by *Myceliophthora thermophila*. Cytotoxic. Powder. $[\alpha]_D^{20} -135$ (c, 0.2 in $CHCl_3$).

5'-Me ether: Myceliothermophin C

[955083-88-2]
 $C_{27}H_{41}NO_3$ 427.626

Prod. by *Myceliophthora thermophila*. Cytotoxic. Powder. $[\alpha]_D^{20} -120$ (c, 0.1 in $CHCl_3$).

5'-Deoxy, 5',6'-didehydro: Myceliothermophin E

[955083-90-6]
 $C_{26}H_{37}NO_2$ 395.584

Prod. by *Myceliophthora thermophila*. Powder. $[\alpha]_D^{20} -40$ (c, 0.2 in $CHCl_3$).

5'-Epimer: Myceliothermophin B

[955083-87-1]
 $C_{26}H_{39}NO_3$ 413.599

Prod. by *Myceliophthora thermophila*. Powder. $[\alpha]_D^{20} -100$ (c, 0.4 in $CHCl_3$).

5'-Epimer, 5'-Me ether: Myceliothermophin D

[955083-89-3]
 $C_{27}H_{41}NO_3$ 427.626


Prod. by *Myceliophthora thermophila*. Powder. $[\alpha]_D^{20} -55$ (c, 0.2 in $CHCl_3$).

Yang, Y.-L. *et al.*, *Chem. Eur. J.*, 2007, 13, 6985-6991 (isol, pmr, cmr, ms)

Mycenarubin A

M-784

[939403-53-9]



$C_{14}H_{15}N_3O_4$ 289.29

Alkaloid from *Mycena haematopus* (blood mycena) and *Mycena rosea*. Red solid. Mp 126° dec. $[\alpha]_D^{25} +1190$ (c, 0.008 in H_2O). λ_{max} 244 (log ϵ 4.12); 358 (log ϵ 3.94); 539 (log ϵ 2.82) (H_2O).

3'-Deamino, 3'-hydroxy: Mycenarubin F


$C_{14}H_{14}N_2O_5$ 290.275
Alkaloid from *Mycena haematopus* (blood mycena). Red solid.

Peters, S. *et al.*, *Eur. J. Org. Chem.*, 2007, 1571-1576; 2008, 319-323 (isol, cd, pmr, cmr, ms)

Mycenarubin B

M-785

[939403-54-0]



$C_{28}H_{28}N_6O_7$ 560.565


Isol. from *Mycena rosea*. Red solid. $[\alpha]_D^{25} +170$ (c, 0.006 in H_2O). λ_{max} 245 (log ϵ 4.19); 354 (log ϵ 3.89); 534 (log ϵ 3.01) (H_2O).

Peters, S. *et al.*, *Eur. J. Org. Chem.*, 2007, 1571-1576 (isol, cd, pmr, cmr, ms)

Mycenarubin D

M-786

[1032121-81-5]



$C_{14}H_{16}N_4O_3$ 288.305

Alkaloid from *Mycena haematopus* (blood mycena). Red solid. $[\alpha]_D^{25} +630$ (c, 0.007 in H_2O). λ_{max} 242 (log ϵ 4.06); 368 (log ϵ 3.96); 544 (log ϵ 2.85) (H_2O).

3'-Deamino, 3'-hydroxy: Mycenarubin E

[1032121-82-6]

$C_{14}H_{15}N_3O_4$ 289.29


Alkaloid from *Mycena haematopus* (blood mycena). Red solid. $[\alpha]_D^{25} +1010$ (c, 0.005 in H_2O). λ_{max} 243 (log ϵ 4.04); 368 (log ϵ 3.94); 548 (log ϵ 2.86) (H_2O).

Peters, S. *et al.*, *Eur. J. Org. Chem.*, 2008, 319-323 (isol, cd, pmr, cmr, ms)

Mycobactin

M-787

[1400-46-0]



A $R^1 = R^2 = CH_3$, $R^3 = H$,
 $R^4 = CH=CH(CH_2)_{10}CH_3$,
J $R^1 = H$, $R^2 = CH(CH_3)_2$, $R^3 = CH_3$,
 $R^4 = CH=CH(CH_2)_{12}CH_3$,
P $R^1 = R^3 = CH_3$, $R^2 = CH_2CH_3$,
 $R^4 = CH=CH(CH_2)_{14}CH_3$,
R $R^1 = H$, $R^2 = CH_2CH_3$, $R^3 = CH_3$,
 $R^4 = CH=CH(CH_2)_{16}CH_3$,
S $R^1 = R^3 = H$, $R^2 = CH_3$,
 $R^4 = CH=CH(CH_2)_{18}CH_3$

A family of modified lipopeptide side-ropores with varying stereochem. Prod. by *Mycobacterium johnei*, *Mycobacterium paratuberculosis*, *Mycobacterium phlei* and other *Mycobacterium* spp. Growth factor complex able to chelate metals (shows strong selectivity for Fe).

Mycobactin A [28338-75-2]

$C_{41}H_{63}N_5O_{10}$ 785.976

Mycobactin J

Mycobactin J1
[80941-50-0]

$C_{45}H_{71}N_5O_{10}$ 842.083

Mycobactin P [1264-66-0]

$C_{47}H_{75}N_5O_{10}$ 870.137

Major component of complex. Powder. Mp 165-165.5°. $[\alpha]_D^{25} -19$ (c, 4.9 in $CHCl_3$). λ_{max} 250; 311 (MeOH) (Berdy).

Picrate:

Noncryst. Mp 150-152°.

Di-Ac:

Noncryst. Mp 134-135.5°. $[\alpha]_D^{15} -6.7$ (c, 4 in $CHCl_3$).

Mycobactin R [28338-74-1]

$C_{48}H_{77}N_5O_{10}$ 884.164

Mycobactin S [26769-11-9]

$C_{44}H_{69}N_5O_{10}$ 828.057

[78245-01-9, 28338-74-1, 78245-02-0, 28338-75-2, 84028-50-2]

Snow, G.A. *et al.*, *J.C.S.*, 1954, 2588; 4080

White, A.J. *et al.*, *Biochem. J.*, 1968, 108, 593 (isol, ms, struct)

Snow, G.A. *et al.*, *Biochem. J.*, 1969, 115, 1031 (isol, props)


- Greatbanks, D. *et al.*, *Biochem. J.*, 1969, **115**, 1047 (*nmr*)
 Snow, G.A. *et al.*, *Bacteriol. Rev.*, 1970, **34**, 99 (*rev*)
 Hough, E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1974, **57**, 73 (*cryst struct*)
 McCullough, W.G. *et al.*, *Curr. Microbiol.*, 1982, **7**, 337 (*Mycobactin J*)
 Barclay, R. *et al.*, *J. Bacteriol.*, 1985, **164**, 896 (*Mycobactin S*)
 MacCordick, H.J. *et al.*, *Nouv. J. Chim.*, 1985, **9**, 535 (*struct*)
 Hu, J. *et al.*, *J.A.C.S.*, 1997, **119**, 3462-3468 (*Mycobactin S, synth*)
 Schwartz, B.D. *et al.*, *Tet. Lett.*, 2001, **42**, 3653-3655 (*Mycobactin J, abs config*)

Mycobactin T M-788

As Mycobactin, M-787 with
 $R^1 = R^3 = H$, $R^2 = CH_3$, $R^4 = C_{17}-C_{20}$
 Prod. by *Mycobacterium tuberculosis*.
 Siderophore and growth factor complex.
 White, A.J. *et al.*, *Biochem. J.*, 1968, **108**, 593-597
 Hu, J. *et al.*, *J.A.C.S.*, 1997, **119**, 3462-3468

Mycosporin glutamicol M-789

5-Hydroxy-4-[[5-hydroxy-5-(hydroxymethyl)-2-methoxy-3-oxo-1-cyclohexen-1-yl]amino]pentanoic acid, 9CI
 [77101-66-7]



- $C_{13}H_{21}NO_7$ 303.311
 Metab. of the fungi *Ascochyta* sp.,
Gnomonia sp., *Morchella* sp. and *Pyronema* sp.
 5'-O-β-D-Glucopyranoside: [81543-07-9]
 $C_{19}H_{31}NO_{12}$ 465.453
 Metab. of *Ascochyta* spp.
Cladosporium sp. and *Septoria* sp.

Amide: Mycosporin glutaminol

[85769-64-8]
 $C_{13}H_{22}N_2O_6$ 302.327
 Widespread mycosporine reduced glutamine. Unstable.

1'-Amide, 5'-O-β-D-glucopyranoside:

[85769-65-9]
 $C_{19}H_{32}N_2O_{11}$ 464.469
 Widespread mycosporine reduced glutamine.

5'-Carboxylic acid: Mycosporin-Glu

[83031-20-3]
 $C_{13}H_{19}NO_8$ 317.295
 Metab. of *Glomerella* sp. and *Helvella* sp.

5'-Carboxylic acid, 1'-amide: Mycosporin-Gln

[92632-66-1]
 $C_{13}H_{20}N_2O_7$ 316.31
 Metab. of *Glomerella* sp. and *Pyronema* sp.

5'-Carboxylic acid, 1'-amide, O-de-Me: Normycosporin-Gln


[77101-67-8]
 $C_{12}H_{18}N_2O_7$ 302.283

Metab. of Pyronema omphalodes.

- [126643-34-3]
 Lunel, M.-C. *et al.*, *Tet. Lett.*, 1980, **21**, 4715 (*Normycosporin-Gln*)
 Bouillant, M.L. *et al.*, *Phytochemistry*, 1981, **20**, 2705-2707; 2709-2710 (*isol, pmr*)
 Young, H. *et al.*, *Phytochemistry*, 1982, **21**, 1075 (*Mycosporine-Glu*)
 Pittet, J.-L. *et al.*, *Tet. Lett.*, 1983, **24**, 65 (*amide*)
 Bernillon, J. *et al.*, *Phytochemistry*, 1984, **23**, 1083 (*isol, glc, ms*)
 Favre-Bonvin, J. *et al.*, *Phytochemistry*, 1987, **26**, 2509 (*isol, amide*)
 White, J.D. *et al.*, *J.O.C.*, 1995, **60**, 3601
 Bandaranayake, W.M. *et al.*, *Nat. Prod. Rep.*, 1998, **15**, 159-172 (*rev*)

Mycothiazole M-790

[114582-75-1]



$C_{22}H_{32}N_2O_3S$ 404.572


Isol. from the marine sponge *Cacospongia mycofijiensis* and the sea slug *Chromodoris lochi*. Exhibits anthelmintic activity (*in vitro*). Highly toxic to mice. Viscous oil. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{20}$ -3.8 (c, 2.9 in $CHCl_3$). $[\alpha]_D^{25}$ -13.7 (c, 0.6 in MeOH). C-14 config. revised in 2006. λ_{max} 235 (ε 5270); 290 (ε 1780) (MeOH) (Berdy).

4,19-Dihydro, 4ξ,19-dihydroxy: Mycothiazole-4,19-diol

- $C_{22}H_{34}N_2O_5S$ 438.587
 Isol. from *Cacospongia mycofijiensis*.
 Powder. $[\alpha]_D^{27}$ -2.5 (c, 0.04 in MeOH).
 Crews, P. *et al.*, *J.A.C.S.*, 1988, **110**, 4365-4368 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Sugiyama, H. *et al.*, *Tetrahedron*, 2003, **59**, 6579-6593 (*synth, abs config*)
 Le Flohic, A. *et al.*, *Org. Lett.*, 2005, **7**, 339-342 (*synth*)
 Sonnenschein, R.N. *et al.*, *J. Nat. Prod.*, 2006, **69**, 145-147 (*Mycothiazole-4,19-diol*)

Mycothioli bimane M-791

[158761-05-8]




- $C_{27}H_{40}N_4O_{14}S$ 676.697
 Prod. by *Streptomyces clavuligerus* and other *Streptomyces* spp., *Micrococcus* sp., *Nocardia* sp., *Rhodococcus* sp., *Actinomadura* sp. and *Nocardiopsis* sp. Antioxi-

idant. Formaldehyde dehydrogenase cofactor.

- Spies, H.S.C. *et al.*, *Eur. J. Biochem.*, 1994, **224**, 203-213 (*isol*)
 Nicholas, G.M. *et al.*, *J.A.C.S.*, 2002, **124**, 3492-3493 (*synth, cd*)
 Mahadevan, J. *et al.*, *J.O.C.*, 2003, **68**, 3380-3386 (*pmr, conformm*)

Myosmine M-792

3-(3,4-Dihydro-2H-pyrrol-5-yl)pyridine, 9CI. 3-(1-Pyrrolin-2-yl)pyridine, 8CI. 2-(3-Pyridyl)-1-pyrroline [532-12-7]



$C_9H_{10}N_2$ 146.191

Present in tobacco smoke. Alkaloid from *Nicotiana* spp. (Solanaceae). Present in hazelnuts, peanuts, potato, tomato, carrot, apple, pineapple, kiwi fruit, cow's milk and cereals. Mp 40.5-42°. Bp_{0.1} 55-60°. pK_a 5.26 (24°).

▶ LD₅₀ (rat, orl) 1875 mg/kg. UT7660000


Dipicrate: Mp 184-185°.

Dipicronate: Mp 213°.

- Swain, M.L. *et al.*, *J.A.C.S.*, 1949, **71**, 1341 (*uv*)
 Eddy, C.R. *et al.*, *Anal. Chem.*, 1954, **26**, 1428 (*ir*)
 Witkop, B. *et al.*, *J.A.C.S.*, 1954, **76**, 5597 (*ir, struct*)
 Pople, J.A. *et al.*, *High Resolution Nuclear Magnetic Resonance*, McGraw-Hill, N.Y., 1959, 281 (*pmr*)
 Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 2926 (*ms*)
 Leete, E. *et al.*, *J.O.C.*, 1972, **37**, 4465 (*synth*)
 Fejer-Kossey, O. *et al.*, *Phytochemistry*, 1972, **11**, 415 (*isol*)
 Brandänge, S. *et al.*, *Acta Chem. Scand., Ser. B*, 1976, **30**, 93 (*synth*)
 Spitzner, D. *et al.*, *Synthesis*, 1977, 242 (*synth*)
 Glenn, D.F. *et al.*, *J.O.C.*, 1978, **43**, 2860 (*ms*)
 Mahboobi, S. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 175 (*synth, pmr*)
 Zwickenflug, W. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 2703-2706; 2002, **50**, 4909-4915 (*glc, anal, occur*)


Myrianthine A M-793

2-(Dimethylamino)-4-methyl-N-[7-(1-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]pentanamide, 9CI
 [24532-76-1]



C₃₁H₄₂N₄O₄ 534.697Alkaloid from the leaves of *Myrianthus arboreus* and from *Discaria americana*. Mp 286°. [α]_D -263 (c, 1 in CHCl₃).Marchand, J. et al., *Ann. Pharm. Fr.*, 1968, **26**, 771-778; *CA*, **71**, 42203q (isol, uv, ir, pmr, struct)**Myrianthine C****M-794**


N-[3,7-Bis(1-methylethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-2-(dimethylamino)-4-methylpentanamide, 9CI [24532-78-3]

C₂₇H₄₂N₄O₄ 486.653Alkaloid from the leaves of *Myrianthus arboreus* (Urticaceae) and aerial parts of *Plectronia odorata* (preferred genus name *Olinia*) (Oliniaceae). Mp 294° dec. [α]_D -228 (CHCl₃).

N-De-Me: N-Demethylmyrianthine C [123115-06-0]

C₂₆H₄₀N₄O₄ 472.626Alkaloid from aerial parts of *Plectronia odorata* (Oliniaceae). Noncryst. [α]_D -103 (c, 1 in CHCl₃).Marchand, J. et al., *Ann. Pharm. Fr.*, 1968, **26**, 771-778; *CA*, **71**, 42203q (isol, uv, ir, pmr, struct)Gournelis, D. et al., *J. Nat. Prod.*, 1989, **52**, 306-316 (N-Demethylmyrianthine C)**Myrianthosine****M-795**


[934766-78-6]



Absolute Configuration

C₂₇H₃₆N₂O₉ 532.589Alkaloid from the aerial parts of *Psychotria myriantha*. Amorph. powder. [α]_D²⁵ +80.8 (c, 0.1 in MeOH).Simões-Pires, C.A. et al., *Nat. Prod. Commun.*, 2006, **1**, 1101-1106 (isol, pmr, cmr)**Myrioneurinol**


[957199-81-4]



Absolute Configuration

C₁₆H₂₇NO₂ 265.395Alkaloid from the leaves of *Myrioneuron nutans*. Cryst. (Et₂O/EtOH). Mp 145-146°. [α]_D²⁰ +63 (c, 1 in MeOH).Pham, V.C. et al., *Tetrahedron*, 2007, **63**, 11244-11249 (isol, pmr, cmr, ms)**Myrionine**


[951681-61-1]



Absolute Configuration

C₁₅H₂₆N₂O 250.383Alkaloid from the leaves of *Myrioneuron nutans*. Oil. [α]_D²⁰ -16.1 (c, 1 in MeOH).Pham, V.C. et al., *Org. Lett.*, 2007, **9**, 3531-3534 (isol, synth, pmr, cmr, cryst struct)**Myrmicarins 663**

[183127-15-3]




Relative Configuration

C₄₅H₆₅N₃O 664.028Alkaloid from the poison gland secretion of the African ant *Myrmecaria opaciventris*.Schröder, F. et al., *Tetrahedron*, 1996, **52**, 13539 (isol, ms)Schröder, F. et al., *Angew. Chem., Int. Ed.*, 1997, **36**, 77 (pmr, cmr)**Myrmicarins 233A**

[183185-10-6]

C₁₅H₂₃NO 233.353Struct. unknown. Prob. an indolizidine alkaloid and a struct. has been tentatively proposed. Trace alkaloid from the poison gland secretion of the African ant *Myrmecaria opaciventris*.**M-799****M-796**Schröder, F. et al., *Tetrahedron*, 1996, **52**, 13539 (isol, ms)Michael, J.P. et al., *Nat. Prod. Rep.*, 1998, **15**, 571-594 (struct)**Myrmicarins 430A**


[183122-49-8]

M-800C₃₀H₄₂N₂ 430.675Constit. of poison gland secretion of the African ant *Myrmecaria opaciventris*.

Dec. rapidly in air.

Schröder, F. et al., *Chem. Comm.*, 1996, 2139 (isol, pmr, cmr)Schröder, F. et al., *Tetrahedron*, 1996, **52**, 13539 (isol, ms)**Myrmicarins 213B**

[183254-01-5]

M-801C₁₅H₁₉N 213.322Alkaloid from the poison gland secretion of the African ant *Myrmecaria opaciventris*.*Z*-Isomer: **Myrmicarins 213A**

[183126-24-1]

C₁₅H₁₉N 213.322From *Myrmecaria opaciventris*.6,7-Dihydro: **Myrmicarins 215B**

[183253-97-6]

C₁₅H₂₁N 215.338From *Myrmecaria opaciventris*. Isol. as a 1:2 inseparable mixt. with Myrmicarins 215A.6,7-Dihydro, *Z*-isomer: **Myrmicarins 215A**C₁₅H₂₁N 215.338From *Myrmecaria opaciventris*. Isol. as an inseparable mixt. with Myrmicarins 215B.1',2'-Dihydro: **Myrmicarins 215C**

[183124-40-5]

C₁₅H₂₁N 215.338From *Myrmecaria opaciventris*.1',2',6,7-Tetrahydro: **Myrmicarins 217**


[183122-85-2]

C₁₅H₂₃N 217.353From *Myrmecaria opaciventris*.Schröder, F. et al., *Tetrahedron*, 1996, **52**,


13539 (*isol, pmr, cmr, ms, struct*)
Schroeder, F. *et al.*, *Tetrahedron*, 1998, **54**,
5259-5264 (*synth, Myrmicaridin 217*)
Sayah, B. *et al.*, *J.O.C.*, 2000, **65**, 2824-2826
(*synth, Myrmicaridin 217*)
Movassaghi, M. *et al.*, *Org. Lett.*, 2005, **7**,
4423-4426 (*synth*)

Myrobotinol**M-802**

[960510-92-3]

Absolute
ConfigurationC₂₀H₃₂N₂O₂ 332.485Alkaloid from the leaves of *Myrioneuron nutans*. Cryst. (hexane/Et₂O). Mp 225-226°. [α]_D²⁰ +36.4 (c, 0.4 in MeOH).Pham, V.C. *et al.*, *J.O.C.*, 2007, **72**, 9826-9829
(*isol, biosynth, pmr, cmr, ms*)**Myrtoidine****M-803**

[243464-91-7]

Absolute
ConfigurationC₂₃H₂₆N₂O₅ 410.469Alkaloid from the stem bark of *Strychnos myrtoides*. Cryst. (EtOAc/cyclohexane). [α]_D²⁰ -46.6 (c, 0.5 in CH₂Cl₂). Dec. without melting. λ_{max} 220 (log ε 4.1); 242 (sh) (log ε 3.72); 289 (log ε 3.18) (MeOH).**Demethoxy: 11-Demethoxymyrtoidine**

[243464-93-9]

C₂₂H₂₄N₂O₄ 380.443Alkaloid from the stem bark of *Strychnos myrtoides*. Cryst. (EtOAc/hexane). Mp 237-240°. [α]_D²⁰ -28.9 (c, 0.4 in CH₂Cl₂). λ_{max} 207 (log ε 4); 241 (log ε 3.74); 277 (sh) (log ε 2.96) (MeOH).**3-Epimer: 3-Epimyrtoidine**

[349493-74-9]

C₂₃H₂₆N₂O₅ 410.469Alkaloid from *Strychnos diplotricha* and *Strychnos myrtoides*. Cryst. (EtOAc/cyclohexane). Mp 105-107°. [α]_D²⁰ +24.7 (c, 0.2 in CHCl₃). λ_{max} 220 (log ε 4.48); 241 (log ε 4.1); 294 (log ε 3.67) (MeOH).**3-Epimer, demethoxy: 11-Demethoxy-3-epimyrtoidine**


[349493-75-0]

C₂₂H₂₄N₂O₄ 380.443Alkaloid from *Strychnos diplotricha*and *Strychnos myrtoides*. Cryst. (EtOAc/cyclohexane). Mp 103-105°. [α]_D²⁰ +37.6 (c, 0.2 in CHCl₃). λ_{max} 210 (log ε 4.45); 244 (log ε 4.18); 276 (log ε 3.5) (MeOH).**3-Epimer, demethoxy, 12-hydroxy: 12-Hydroxy-11-demethoxy-3-epimyrtoidine**

[349493-76-1]

C₂₃H₂₄N₂O₅ 396.442Alkaloid from *Strychnos diplotricha* and *Strychnos myrtoides*. Cryst. (EtOAc/cyclohexane). Mp 230-232°. [α]_D²⁰ -184.5 (c, 0.2 in CHCl₃). λ_{max} 217 (log ε 4.43); 247 (log ε 3.96); 288 (log ε 3.44) (MeOH).Martin, M.-T. *et al.*, *Phytochemistry*, 1999, **51**, 479-486 (*isol, uv, cd, pmr, cmr, ms*)Rasoanaivo, P. *et al.*, *Phytochemistry*, 2001, **56**, 863-867 (*isol, 3-epimers*)**Myrtopsine****M-804**


[60623-06-5]

Relative
configurationC₁₅H₁₇NO₄ 275.304Alkaloid from *Myrtopsis selligii*, *Haplophyllum foliosum* and *Zanthoxylum integrifolium* (Rutaceae). Beige cryst. (CHCl₃). Mp 208°. [α]_D -5 (c, 1 in MeOH).**O³-Ac:**Cryst. (Et₂O). Mp 178°.**Di-Ac:**

Noncryst.

7,8-Dimethoxy: 7,8-Dimethoxymyrtopsine

[142741-28-4]

C₁₇H₂₁NO₆ 335.356Alkaloid from stem bark of *Dutailliea baudouinii* (Rutaceae).**3-Deoxy:** see Platydesmine, P-515Hifnawy, M.S. *et al.*, *Phytochemistry*, 1977, **16**, 1035-1039 (*isol*)Akhmedzhanova, V.I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 447-449 (*isol, struct*)Ishii, H. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 182; *C.A.*, **97**, 69240y (*isol*)Muyard, F. *et al.*, *Phytochemistry*, 1992, **31**, 1087-1089 (*7,8-Dimethoxymyrtopsine*)Chen, I.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1143-1147 (*7,8-Dimethoxymyrtopsine, cmr*)
Snider, B.B. *et al.*, *Heterocycles*, 2006, **70**, 279-294 (*synth*)**Myxalamides****M-805**Myxalamide A: R = CH(CH₃)CH₂CH₃
B: R = CH(CH₃)₂
C: R = CH₂CH₃
D: R = CH₃**Myxalamide A** [86934-09-0]C₂₆H₄₁NO₃ 415.615Prod. by *Myxococcus xanthus* Mx x12 (GBF) and *Stigmatella aurantiaca*. Sol. MeOH, CH₂Cl₂, CHCl₃; fairly sol. CCl₄, Et₂O, C₆H₆; poorly sol. hexane, H₂O. [α]_D²⁰ -71.2 (c, 0.2 in MeOH). λ_{max} 207 (ε 24000); 263 (ε 11000); 340 (sh); 357 (ε 46800); 370 (sh) (EtOH).

▶RG0181000

Myxalamide B [86934-10-3]C₂₅H₃₉NO₃ 401.588Prod. by *Myxococcus xanthus* Mx x12 (GBF) and *Stigmatella aurantiaca*. Electron transport inhibitor. Sol. MeOH, CH₂Cl₂; fairly sol. CCl₄, C₆H₆, Et₂O; poorly sol. hexane, H₂O. [α]_D²⁰ -61.6 (c, 0.25 in MeOH). λ_{max} 207 (ε 24000); 263 (ε 11000); 340 (sh); 357 (ε 46800); 370 (sh) (EtOH).▶LD₅₀ (mus, scu) 211 mg/kg, LD₅₀ (mus, orl) 613 mg/kg. MI3000000**Myxalamide C** [86934-11-4]C₂₄H₃₇NO₃ 387.561Prod. by *Myxococcus xanthus* Mx x12 (GBF) and *Stigmatella aurantiaca*. Electron transport inhibitor. Sol. MeOH, CHCl₃, CH₂Cl₂; fairly sol. CCl₄, C₆H₆; poorly sol. hexane, H₂O. [α]_D²⁰ -35.7 (c, 0.4 in CHCl₃). λ_{max} 207 (ε 24000); 263 (ε 11000); 340 (sh); 357 (ε 46800); 370 (sh) (EtOH).▶LD₅₀ (mus, scu) 211 mg/kg, LD₅₀ (mus, orl) 613 mg/kg. MI3010000**Myxalamide D** [86934-12-5]C₂₃H₃₅NO₃ 373.534Prod. by *Myxococcus xanthus* Mx x12 (GBF) and *Stigmatella aurantiaca*. Electron transport inhibitor. Sol. MeOH, CHCl₃, CH₂Cl₂; fairly sol. CCl₄, C₆H₆; poorly sol. H₂O, hexane. [α]_D²⁰ -14.6 (c, 0.25 in CHCl₃). λ_{max} 207 (ε 24000); 263 (ε 11000); 340 (sh); 357 (ε 46800); 370 (sh) (EtOH).


▶ML8000000

2'-Me ether: 2'-O-Methylmyxalamide DC₂₄H₃₇NO₃ 387.561Isol. from *Cystobacter fuscus*. Pale yellow oil. [α]_D²¹ -27 (c, 0.04 in MeOH). λ_{max} 260 (ε 6000); 340 (sh); 355 (ε 30000) (MeOH).**6E-Isomer, 2'-Me ether: (6E)-2'-O-Methylmyxalamide D**C₂₄H₃₇NO₃ 387.561Isol. from *Cystobacter fuscus*. Pale yellow oil. [α]_D²¹ -31 (c, 0.02 in MeOH). λ_{max} 257 (ε 3000); 340 (sh); 356 (ε 30000) (MeOH).**(6E,10Z)-Isomer, 2'-Me ether: (6E,10Z)-2'-O-Methylmyxalamide D**C₂₄H₃₇NO₃ 387.561Isol. from *Cystobacter fuscus*. Pale yellow oil. [α]_D²¹ -12 (c, 0.03 in MeOH). λ_{max} 260 (ε 6000); 340 (sh); 355 (ε 30000) (MeOH).Jansen, R. *et al.*, *Annalen*, 1983, 1081-1095; 1984, 78-84 (*isol, pmr, cmr, abs config*)

- Mapp, A.K. *et al.*, *J.O.C.*, 1999, **64**, 23-27 (synth)
 Kundim, B.A. *et al.*, *Tetrahedron*, 2004, **60**, 10217-10221 (2'-O-Methylmyxalamide D isomers)
 Feng, Z. *et al.*, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 699-705 (2'-O-Methylmyxalamide D, biosynth)
 Coleman, R.S. *et al.*, *J.A.C.S.*, 2007, **129**, 3826-3827 (2'-O-Methylmyxalamide D, synth)

Myxochelin A M-806

N,N' -[1-(Hydroxymethyl)-1,5-pentanediy]bis[2,3-dihydroxybenzamide], 9CI.
 N^2,N^6 -Bis(2,3-dihydroxybenzoyl)lysineol [120243-02-9]



$C_{20}H_{24}N_2O_7$ 404.419

Prod. by *Angiococcus disciformis* strain An d30 and *Nonomuraea pusilla* TP-A0861. Siderophore. Amorph. solid. Sol. MeOH, $CHCl_3$, EtOAc; fairly sol. Et_2O ; poorly sol. hexane. $[\alpha]_D^{25}$ -16.5 (c, 0.85 in MeOH). Related to N^2,N^6 -Bis(2,3-dihydroxybenzoyl)lysine. λ_{max} 210 (€ 57500); 245 (sh) (€ 20000); 310 (€ 7940) (MeOH) (Derep).

10-Deoxy, 10-amino: Myxochelin B

[201535-34-4]

[201612-28-4]

$C_{20}H_{25}N_3O_6$ 403.434

Prod. by *Myxococcus xanthus*. Siderophore. $[\alpha]_D^{20}$ -10 (c, 1 in 6M HCl).

10-Deoxy, 10-(2,3-dihydroxybenzoylamino): Myxochelin C

[179751-76-9]

[201535-64-0]

$C_{27}H_{29}N_3O_9$ 539.541

Prod. by *Myxococcus xanthus*. Siderophore. Plates. Mp 115°. $[\alpha]_D^{20}$ +9.1 (c, 0.35 in MeOH).

Kunze, B. *et al.*, *J. Antibiot.*, 1989, **42**, 14-17 (Myxochelin A, isol, activity)


Ambrosi, H.-D. *et al.*, *Eur. J. Org. Chem.*, 1998, 541-551 (Myxochelins B, C)

Miyayama, S. *et al.*, *J. Antibiot.*, 2006, **59**, 698-703 (Myxochelin A, isol, synth, pmr, cmr, abs config)

Li, Y. *et al.*, *J.A.C.S.*, 2008, **130**, 7554-7555 (biosynth)

Myxothiazole A M-807

7-[2'-(1,6-Dimethyl-2,4-heptadienyl)[2,4'-bithiazol]-4-yl]-3,5-dimethoxy-4-methyl-2,6-heptadienamamide, 9CI [76706-55-3]



$C_{25}H_{33}N_3O_3S_2$ 487.686

Isol. from *Myxococcus fulvus*. Active against filamentous fungi. Needles (Et_2O /heptane). Mp 79°. $[\alpha]_D^{25}$ +43.4 (c, 6.0 in MeOH). λ_{max} 234 (€ 47000); 313 (€ 10500) (MeOH) (Derep).

►QH7580000**17,18-Epoxyde: 17,18-Epoxyomyxothiazole**

$C_{25}H_{33}N_3O_4S_2$ 503.685

Prod. by *Myxococcus fulvus*. Amorph. solid. $[\alpha]_D^{25}$ +28 (c, 0.1 in MeOH). Possesses trans-epoxide config. λ_{max} 208 (€ 9700); 224 (€ 8910); 310 (€ 1600) (MeOH).

 O^3 -De-Me, 2,3-dihydro, 3-ketone: Desmethylmyxothiazole A. Desmethylmyxothiazole A

$C_{24}H_{31}N_3O_3S_2$ 473.659

Prod. by *Myxococcus fulvus*. Amorph. solid. λ_{max} 245 (€ 31600); 312 (€ 10450) (MeOH).

Parent acid, Me ester: Myxothiazole Z.

Antibiotic KR 025. KR 025

[222850-18-2]

$C_{26}H_{34}N_2O_4S_2$ 502.698

Isol. from *Myxococcus fulvus*. Cytotoxic. Oil. $[\alpha]_D^{25}$ +152 (c, 0.67 in MeOH). λ_{max} 232 (log € 4.63); 312 (log € 3.99) (MeOH).

17Z-Isomer: 17Z-Myxothiazole A

$C_{25}H_{33}N_3O_3S_2$ 487.686

Prod. by *Myxococcus fulvus*. Amorph. solid. $[\alpha]_D^{25}$ +77.4 (c, 0.2 in MeOH). Struct. incorrectly drawn in ref. λ_{max} 245 (€ 33880); 312 (€ 11480) (MeOH).

Gerth, K. *et al.*, *J. Antibiot.*, 1980, **33**, 1474-1479; 1480-1490 (isol, struct)

Thierbach, G. *et al.*, *Antimicrob. Agents Chemother.*, 1981, **19**, 504-507 (props)

Trowitzsch, W. *et al.*, *Tet. Lett.*, 1981, **22**, 3829-3832 (abs config)

Trowitzsch-Kienast, W. *et al.*, *Annalen*, 1986, 93-98 (biosynth)

Conradt, P. *et al.*, *J. Antibiot.*, 1989, **42**, 1158-1162 (props)

Martin, B.J. *et al.*, *Tet. Lett.*, 1993, **34**, 5151-5154 (synth)

Ahn, J.-W. *et al.*, *J. Nat. Prod.*, 1999, **62**, 495-496 (KR 025)

Backhaus, D. *et al.*, *Tet. Lett.*, 2000, **41**, 2087-2090 (synth)

Steinmetz, H. *et al.*, *Tetrahedron*, 2000, **56**, 1681-1684 (Myxothiazole Z)


Clough, J.M. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 2906-2911 (synth)

Ahn, J.-W. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 477-479 (17Z-Myxothiazole, 17,18-Epoxyomyxothiazole)

Iwaki, Y. *et al.*, *Tet. Lett.*, 2008, **49**, 7024-7026 (synth)

Myxovalargin

M-808



Polypeptide antibiotic complex. Struct. of Myxovalargin A shown. Other components are lower homologues of A but their structs. are not fully known. Prod. by *Myxococcus fulvus*. Protein biosynthesis inhibitor. Antibacterial agent.

Myxovalargin A [82658-75-1]

$C_{81}H_{137}N_{21}O_{17}$ 1677.104

Powder. Sol. MeOH, butanol, DMSO; fairly sol. H_2O . λ_{max} 226 (MeOH). λ_{max} 226 (E1%/1cm 1.47) (MeOH) (Berdy).

►LD₅₀ (mus, scu) 10 mg/kg.**Myxovalargin B** [85568-63-4]

Powder. Sol. MeOH, DMSO, butanol; fairly sol. H_2O . λ_{max} 226 (MeOH). λ_{max} 222 ; 275 (MeOH) (Berdy).

Myxovalargin C [85568-64-5]

Sol. MeOH, butanol. λ_{max} 226 (MeOH). λ_{max} 222 ; 275 (MeOH) (Berdy).

Myxovalargin D [85568-65-6]


Sol. MeOH, butanol. λ_{max} 226 (MeOH). λ_{max} 222 ; 275 (MeOH) (Berdy).

Irschik, H. *et al.*, *J. Antibiot.*, 1983, **36**, 6-12; 1985, **38**, 1237-1245 (isol, activity)

Steinmetz, H. *et al.*, *Chem. Pept. Proteins*, 1989, **4**, 13-18 (pmr, cmr, ms, struct)

Naamidine A

[110189-06-5]

C₂₃H₂₃N₅O₄ 433.466

Extensive tautomerism possible in all Naamidines. Alkaloid from the Red Sea sponge *Leucetta chagosensis* and from *Notodoris citrina*. Antagonist of the epidermal growth factor receptor. Yellow foaming oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 240 (ε 1160); 281 (ε 4330); 366 (ε 10500); 392 (ε 13500); 416 (ε 10700) (CHCl₃) (Derep). λ_{max} 225 (ε 21400); 276 (ε 4800); 384 (ε 10060) (MeOH) (Berdy).

Zn complex: **Bis(naamidinato A)zinc(II)**

[171114-02-6]

C₄₆H₄₄N₁₀O₈Zn 930.306

From *Leucetta* sp. Yellow amorph. solid. MF erroneously given as

C₄₆H₄₄N₁₀O₁₀Zn in CAS.N³-Me: **Naamidine C**

[121819-69-0]

C₂₄H₂₅N₅O₄ 447.493

Alkaloid from *Leucetta chagosensis*. Exists as Δ^{4(N)}-isomer.

O¹¹-Me: **Naamidine G**

[171784-01-3]

C₂₄H₂₅N₅O₄ 447.493

From *Leucetta* sp. Yellow solid (EtOAc). Mp 94°.

O¹¹-Me, Zn complex: **Bis(naamidinato G)zinc(II)**

[171114-03-7]

C₄₈H₄₈N₁₀O₈Zn 958.359

From *Leucetta* sp. Yellow amorph. solid. MF erroneously given as

C₄₈H₄₈N₁₀O₁₀Zn in CAS.O¹¹-Me, Zn complex with naamidine A: **(Naamidinato A)(naamidinato G)zinc(II)**

[171114-04-8]

C₄₇H₄₆N₁₀O₈Zn 944.333

From *Leucetta* sp. Yellow amorph. solid. First naturally occurring mixed-ligand metal complex. MF erroneously given as C₄₇H₄₆N₁₀O₁₀Zn in CAS.

O¹¹-Me, N³-De-Me: **Naamidine D**

[121819-71-4]

C₂₃H₂₃N₅O₄ 433.466

Trace alkaloid from *Leucetta chagosensis*. λ_{max} 240 (ε 8800); 268 (ε 305); 386 (ε 6500) (CHCl₃/MeOH) (Derep). λ_{max} 240 (ε 8800); 268 (ε 3050); 386 (ε 6500) (MeOH-CHCl₃) (Berdy).

10-Hydroxy, O¹¹-Me: **Naamidine B**

[121819-68-9]

C₂₄H₂₅N₅O₅ 463.492

Alkaloid from *Leucetta chagosensis*.

14-Hydroxy, O¹¹-Me: **14-Hydroxy-naamidine G**

[171114-06-0]

N-1

C₂₄H₂₅N₅O₅ 463.492

From *Leucetta* sp. [α]_D²⁵ -8 (c, 0.2 in MeOH).

14-Hydroxy: **14-Hydroxy-naamidine A**

[171114-05-9]

C₂₃H₂₃N₅O₅ 449.465

Alkaloid from a calcareous sponge, *Leucetta* sp., of the Coral Sea. [α]_D²⁵ +6 (c, 0.3 in MeOH).

14-Methoxy: **14-Methoxy-naamidine A**

[171114-07-1]

C₂₄H₂₅N₅O₅ 463.492

From *Leucetta* sp. Isol. as a 12:88 mixt. with Naamidine A.

14-Methoxy, O¹¹-Me: **14-Methoxy-naamidine G**

[171114-08-2]

C₂₅H₂₇N₅O₅ 477.519

From *Leucetta* sp. [α]_D²⁵ +4 (c, 0.3 in MeOH).

10,12-Dimethoxy: **Naamidine H**

[952512-59-3]

C₂₅H₂₇N₅O₆ 493.518

Isol. from *Leucetta chagosensis*.

Amorph. yellow solid. λ_{max} 225 (sh) (log ε 4.27); 278 (log ε 3.77); 382 (log ε 3.97) (MeOH).

10,12-Dimethoxy, 2'-(N-methylimide): **Naamidine I**

[952512-60-6]

C₂₆H₃₀N₆O₅ 506.56

Isol. from *Leucetta chagosensis*.

Amorph. yellow solid. λ_{max} 225 (sh) (log ε 4.29); 275 (log ε 3.85); 384 (log ε 3.91) (MeOH).

14-Oxo, O¹¹-Me: **14-Oxonaamidine G**

[171114-09-3]

C₂₄H₂₃N₅O₅ 461.476

From *Leucetta* sp.

Carmely, S. *et al.*, *Tetrahedron*, 1989, **45**, 2193-2200 (*isol, uv, ir, pmr, cmr, ms, struct*)

Mancini, I. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 1178 (*derivs*)

Copp, B.R. *et al.*, *J. Med. Chem.*, 1998, **41**, 3909-3911 (*isol, use*)


Ohta, S. *et al.*, *Heterocycles*, 2000, **53**, 1939-1955 (*synth*)

Aberle, N.S. *et al.*, *Org. Lett.*, 2006, **8**, 419-421 (*synth*)

Tsukamoto, S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1658-1660 (*Naamidines H, I*)

Naamidine F

[152273-85-3]

C₂₅H₂₅N₅O₇ 507.502

Alkaloid from the sponge *Leucetta* sp. Weakly cytotoxic. Yellow needles. Mp 172-174°.


Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1993, **46**, 1229 (*isol, uv, ir, pmr, cmr, ms, struct*)

Naamine A

N-3

4-[2-Amino-4-[(4-methoxyphenyl)-methyl]-1-methyl-1H-imidazol-5-yl]-methyl]phenol, 9CI. 2-Amino-5-(p-hydroxybenzyl)-4-(p-methoxybenzyl)-1-methylimidazole

[110189-03-2]

C₁₉H₂₁N₃O₂ 323.394

Alkaloid from the Red Sea sponge *Leucetta chagosensis* and from *Notodoris citrina*. Also obt. by acid treatment of Naamidine A, N-1. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 185-187.5° (as picrate). λ_{max} 226 (ε 17000); 260 (ε 11600); 282 (ε 6000); 290 (ε 5500); 302 (ε 4700) (dioxan) (Derep).


Carmely, S. *et al.*, *Tetrahedron*, 1989, **45**, 2193 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ohta, S. *et al.*, *Heterocycles*, 2000, **53**, 1939-1955 (*synth*)

Naamine B

N-4

[121849-79-4]

C₂₁H₂₅N₃O₃ 367.447

Alkaloid from the Red Sea sponge *Leucetta chagosensis* with *Notodoris citrina*.


Carmely, S. *et al.*, *Tetrahedron*, 1989, **45**, 2193 (*isol, ir, pmr, cmr, ms, struct*)

Kawasaki, I. *et al.*, *J.C.S. Perkin I*, 2001, 3095-3099 (*synth*)

Naamine C

N-5

[189825-81-8]



$C_{21}H_{25}N_3O_4$ 383.446
Alkaloid from the Micronesian sponge *Leucetta chagosensis*. Yellow powder. λ_{max} 228 ; 276 ; 368 (MeOH).

2'-Deoxy, 5'-hydroxy, 3'-O-de-Me:
Naamine E

[454472-13-0]
 $C_{20}H_{23}N_3O_4$ 369.419

Alkaloid from *Leucetta cf. chagosensis*. Amorph. yellow solid. λ_{max} 233 (€ 21600); 277 (€ 6090) (MeOH).

5'-Methoxy, 2'-deoxy, 4'-O-de-Me:

Naamine G
[700813-14-5]

$C_{21}H_{25}N_3O_4$ 383.446
Alkaloid from *Leucetta chagosensis*. Antifungal agent. Yellow-brown oil. λ_{max} 230 ; 276 (MeOH).

2'-Deoxy, 4'-O-de-Me: **Naamine F**

$C_{20}H_{23}N_3O_3$ 353.42
Alkaloid from *Leucetta chagosensis*. Dark brown solid. CAS No. not found 8-14Cl. λ_{max} 226 ; 279 (MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1997, **60**, 497-498 (*Naamine C*)


Gross, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1190-1193 (*Naamine E*)

Nakamura, S. *et al.*, *J.C.S. Perkin 1*, 2002, 1061-1066 (*synth, pmr*)

Hassan, W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 817-822 (*Naamines F,G*)

Naamine D N-6

4,5-Bis[(4-methoxyphenyl)methyl]-1H-imidazol-2-amine, 9CI
[316814-71-8]



$C_{19}H_{21}N_3O_2$ 323.394
Alkaloid from the sponge *Leucetta cf. chagosensis*. Antifungal and nitric oxide synthase inhibitory agent. Amorph. powder. λ_{max} 230 (€ 15400); 262 (€ 1290); 299 (€ 3300) (CHCl₃).

N¹,N³-Di-Me: N¹,N³-Dimethylnaamine D

[496954-15-5]
 $C_{21}H_{25}N_3O_2$ 351.447


Alkaloid from a *Leucetta* sp. Amorph. amber solid. λ_{max} 205 (log € 4.25); 281 (log € 3.75); 312 (log € 3.58) (MeOH).

Dunbar, D.C. *et al.*, *Tetrahedron*, 2000, **56**, 8795-8798 (*isol, pmr, cmr, ir, uv*)

Crews, P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 177-182 (*N,N-Dimethylnaamine D*)

Nagelamide B N-7

[690627-55-5]



$C_{22}H_{22}Br_4N_{10}O_3$ 794.097

Alkaloid from the sponge *Agelas* sp. Antibacterial agent. Amorph. solid. λ_{max} 202 (€ 19200); 282 (€ 16000) (MeOH).

9-Deoxy: **Nagelamide A**

[690627-54-4]
 $C_{22}H_{22}Br_4N_{10}O_2$ 778.098

Alkaloid from an *Agelas* sp. Antibacterial agent. Amorph. solid. λ_{max} 202 (€ 29800); 279 (€ 27800) (MeOH).

9-Deoxy, 9,10-didehydro(E-): **Nagelamide C**

[690627-56-6]
 $C_{22}H_{20}Br_4N_{10}O_2$ 776.082

Alkaloid from an *Agelas* sp. Antibacterial agent. Amorph. solid. λ_{max} 202 (€ 21200); 280 (€ 19700) (MeOH).

9-Deoxy, 9',10'-dihydro: **Nagelamide D**


[690627-57-7]
 $C_{22}H_{24}Br_4N_{10}O_2$ 780.113

Alkaloid from an *Agelas* sp. Antibacterial agent. Amorph. solid. λ_{max} 202 (€ 16500); 280 (€ 14900) (MeOH).

Endo, T. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1262-1267 (*isol, pmr, cmr*)

Nagelamide H N-8

[690627-61-3]




$C_{24}H_{25}Br_4N_{11}O_5S$ 899.214

Related to Mauritamine, M-124. Alkaloid from the sponge *Agelas* sp. Amorph. solid. λ_{max} 221 (€ 34900); 279 (€ 28300) (MeOH).

Endo, T. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1262-1267 (*isol, pmr, cmr*)

Nagstatin N-9

8-(Acetylamino)-5,6,7,8-tetrahydro-6,7-dihydroxy-5-(hydroxymethyl)imidazo[1,2-a]pyridine-2-acetic acid
[126844-81-3]



$C_{12}H_{17}N_3O_6$ 299.283

Prod. by *Streptomyces amakusaensis*. Inhibitor of *N*-acetyl-β-D-glucosaminidase. Immunopotentiator. Powder. Sol. H₂O, DMSO, MeOH; fairly sol. butanol; poorly sol. CHCl₃, hexane. Mp 190-195°. $[\alpha]_D^{25} +46.2$ (c, 0.5 in H₂O). λ_{max} 225 (€ 3310) (MeOH) (Derep). λ_{max} 225 (€ 3311) (H₂O) (Berdy).

▶ NJ5198400

Aoyagi, T. *et al.*, *J. Antibiot.*, 1992, **45**, 1404; 1557 (*isol, pmr, cmr, struct, props*)

Tatsuta, K. *et al.*, *Tet. Lett.*, 1995, **36**, 6721 (*synth*)

Nairomycin N-10

Niromycin

Glutarimide-type antibiotic complex. Struct. unknown. Prod. by *Streptomyces albus*. Antiviral agent.

Nairomycin A

Cryst. Sol. MeOH, CHCl₃, H₂O; fairly sol. Et₂O, C₆H₆; poorly sol. hexane. Mp 98-105°.

▶ LD₅₀ (mus, ivn) 40 - 60 mg/kg.

Nairomycin B


Cryst. Sol. MeOH, H₂O, CHCl₃; fairly sol. Et₂O, C₆H₆; poorly sol. hexane. Mp 47-67°.

▶ LD₅₀ (mus, ivn) 48 mg/kg.

Japan. Pat., 1960, 60 16997; *CA*, **55**, 20325i (*isol*)

Nakodomarin A N-11

[199483-01-7]



Relative Configuration

$C_{26}H_{36}N_2O$ 392.583

Alkaloid from the sponge *Amphimedon* sp. Cyclin-dependent kinase inhibitor. Amorph. solid. $[\alpha]_D^{25} -16$ (c, 0.1 in MeOH). λ_{max} 206 (€ 11000); 228 (€ 10000) (MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 1997, **62**, 9236-9239 (*isol, uv, ir, pmr, cmr*)


Nagata, T. *et al.*, *J.A.C.S.*, 2003, **125**, 7484-7485; 13618 (*synth*)

Ono, K. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 2020-2023 (*synth*)

Young, I.S. *et al.*, *J.A.C.S.*, 2007, **129**, 1465-1469 (*synth*)

Nakamuric acid N-12

[247145-39-7]



Relative configuration

C₂₀H₂₁Br₂N₇O₄ 583.238

Isol. from the sponge *Agelas nakamurai*. Antibacterial agent. Amorph. brown solid. $[\alpha]_D^{20}$ -9.9 (c, 0.26 in MeOH). Incorr. MF in ref. λ_{\max} 213 (log ϵ 4.02); 269 (log ϵ 4.08) (MeOH).

Me ester: [247113-88-8]


C₂₁H₂₃Br₂N₇O₄ 597.265

Isol. from *Agelas nakamurai*. Amorph. brown solid. $[\alpha]_D^{20}$ -4.1 (c, 0.3 in MeOH). Possible artifact. λ_{\max} 213 (log ϵ 4.04); 269 (log ϵ 4.09) (MeOH).

Eder, C. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1295-1297 (*isol, pmr, cmr, uv*)O'Malley, D.P. *et al.*, *J.A.C.S.*, 2007, **129**, 4762-4775; 7702 (*synth*)**Nakijinol**

[166990-20-1]

N-13


C₂₂H₂₉NO₃ 355.476

Constit. of an Okinawan sponge of the family Spongiidae. Amorph. powder. $[\alpha]_D^{20}$ -172 (c, 0.03 in MeOH). λ_{\max} 300 (ε 3600); 332 (ε 2000) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *Tet. Lett.*, 1995, **36**, 5589 (*isol, pmr, cmr, uv, ir*)**Nakijiquinone A**

[157207-60-8]

N-14

R = CH₂COOHC₂₃H₃₁NO₅ 401.502

Constit. of an Okinawan marine sponge. Protein-tyrosine kinase inhibitor. Red solid. Mp 156-158°. $[\alpha]_D^{20}$ -71.7 (c, 1 in MeOH). Related to Avarol, A-1560. λ_{\max} 317 (ε 11800); 488 (ε 860) (MeOH) (Berdy).

Shigemori, H. *et al.*, *Tetrahedron*, 1994, **50**, 8347 (*isol, pmr, cmr*)Stahl, P. *et al.*, *J.A.C.S.*, 2001, **123**, 11586-11593 (*synth*)**Nakijiquinone B**

[157232-59-2]

N-15

As Nakijiquinone A, N-14 with


R = CH(COOH)CH(CH₃)₂(S-)C₂₆H₃₇NO₅ 443.582

Constit. of an Okinawan marine sponge. Protein-tyrosine kinase inhibitor. Red solid. $[\alpha]_D^{20}$ -282.3 (c, 0.13 in CHCl₃). λ_{\max} 320 (ε 12000); 492 (ε 910) (MeOH) (Berdy).

Shigemori, H. *et al.*, *Tetrahedron*, 1994, **50**, 8347 (*isol, pmr, cmr*)Stahl, P. *et al.*, *J.A.C.S.*, 2001, **123**, 11586-11593 (*synth*)**Nakijiquinone D**

[169438-44-2]

N-16

C₂₅H₃₅NO₆ 445.555

Constit. of a marine sponge. Protein-tyrosine kinase inhibitor. Red amorph. solid. $[\alpha]_D^{20}$ -172 (c, 0.2 in EtOH). λ_{\max} 317 (ε 12600); 490 (ε 1000) (MeOH) (Berdy).

25-Demethyl: *Nakijiquinone C*

[169438-43-1]


C₂₄H₃₃NO₆ 431.528

Constit. of a marine sponge. Protein-tyrosine kinase inhibitor. Red amorph. solid. $[\alpha]_D^{20}$ -73 (c, 0.03 in EtOH). λ_{\max} 321 (ε 12100); 498 (ε 920) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**, 10867 (*isol, pmr, cmr*)Stahl, P. *et al.*, *J.A.C.S.*, 2001, **123**, 11586-11593 (*synth*)**Nakinadine A**

[945979-01-1]

N-17




Relative Configuration

C₄₅H₆₇N₃O₂ 682.043

Alkaloid from an *Amphimedon* sp. (SS-1059). Oil. $[\alpha]_D^{25}$ -3 (c, 1 in CHCl₃). λ_{\max} 257 (ε 3700); 263 (ε 4100); 269 (ε 3600) (MeOH).

Kubota, T. *et al.*, *Tet. Lett.*, 2007, **48**, 4983-4985 (*isol, pmr, cmr*)**Nakinadine B**


N-18

C₂₇H₄₀N₂O₂ 424.625**(S)-form** [1021487-41-1]

Alkaloid from an *Amphimedon* sp. (SS-1059). Cytotoxic. Oil. λ_{\max} 257 (ε 3100); 263 (ε 3300); 270 (ε 2900) (MeOH).

Nishi, T. *et al.*, *Tetrahedron*, 2008, **64**, 3127-3132 (*isol, pmr, cmr*)**Nakinadine C**

N-19


C₂₈H₄₀N₂O₂ 436.636**(S)-form** [1021486-53-2]

Alkaloid from an *Amphimedon* sp. (SS-1059). Cytotoxic. Oil. λ_{\max} 258 (ε 3200); 263 (ε 3300); 269 (ε 3000) (MeOH).

Nishi, T. *et al.*, *Tetrahedron*, 2008, **64**, 3127-3132 (*isol, pmr, cmr, ms*)**Nakinadine D**

[1021486-55-4]

N-20



Relative Configuration


C₄₃H₆₅N₃O₂ 656.005

Alkaloid from an *Amphimedon* sp. (SS-1059). Oil. λ_{\max} 258 (ε 4700); 263 (ε 5000); 270 (ε 4400) (MeOH).

Nishi, T. *et al.*, *Tetrahedron*, 2008, **64**, 3127-3132 (*isol, pmr, cmr, ms*)**Nakinadine E**

[1021486-57-6]

N-21

C₄₄H₆₇N₃O₂ 670.032


Alkaloid from an *Amphimedon* sp. (SS-1059). Oil. λ_{\max} 259 (ϵ 4900); 263 (ϵ 5100); 269 (ϵ 4500) (MeOH).

Nishi, T. *et al.*, *Tetrahedron*, 2008, **64**, 3127-3132 (*isol, pmr, cmr, ms*)

Nakinadine F

N-22

[1021486-59-8]



$C_{46}H_{67}N_3O_2$ 694.054


Alkaloid from an *Amphimedon* sp. (SS-1059). Oil. λ_{\max} 258 (ϵ 4900); 264 (ϵ 5200); 270 (ϵ 4600) (MeOH).

Nishi, T. *et al.*, *Tetrahedron*, 2008, **64**, 3127-3132 (*isol, pmr, cmr, ms*)

Nakirodine A

N-23

[473304-17-5]



$C_{17}H_{24}Br_2N_2O_4$ 480.196


Isol. from a Verongid sponge. Oil. $[\alpha]_D^{25}$ +35 (c, 0.1 in MeOH). λ_{\max} 283 (ϵ 880) (MeOH).

Tsuda, M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1670-1671 (*isol, pmr, cmr, ms*)

Nandazurine

N-24

[49679-20-1]



$C_{19}H_{13}NO_5$ 335.315

Alkaloid from the bark of *Nandina domestica* and from *Corydalis bulbosa* (Nandiniaceae, Papaveraceae). Dark-green amorph. powder. Mp 350°.

Kunitomo, J. *et al.*, *Experientia*, 1973, **29**, 518 (*uv, ir, pmr, struct*)


Kupchan, S.M. *et al.*, *Chem. Comm.*, 1973, 915 (*synth*)

Kiryakov, H.G. *et al.*, *Planta Med.*, 1981, **43**, 51 (*isol, ir, pmr, synth*)

Nandigerine

N-25

6,7,7a,8-Tetrahydro-12-methoxy-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-11-ol, 9CI. 10-Hydroxy-11-methoxy-1,2-methylenedioxy-noraporphine. *Hernandigerine*



$C_{18}H_{17}NO_4$ 311.337

(S)-form [31520-97-5]

Alkaloid from *Hernandia ovigera*, *Hernandia cordigera*, *Hernandia jamaicensis*, *Hernandia catalpifolia*, *Hernandia papuana* and *Laurus nobilis* (Hernandiaceae). Shows anti-platelet aggregation activity. Needles (MeOH). Mp 176-177°. $[\alpha]_D$ +248 (EtOH). Forms MeOH solvate, Mp 99-100°.

Hydrochloride: Mp 245-247° dec.

Hydrobromide:

Cryst. (MeOH/Me₂CO). Mp 227-230° dec.

N-Formyl: N-Formylnandigerine.

N-Formylhernandigerine

[176679-16-6]

$C_{19}H_{17}NO_5$ 339.347

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Prisms + $\frac{1}{2}H_2O$ (MeOH). Mp 233-235°. $[\alpha]_D^{24}$ +461 (c, 0.10 in CHCl₃). Exists in soln. as a mixt. of amide bond rotamers.

N-Me: N-Methylnandigerine. 10-Hydroxy-11-methoxy-1,2-methylenedioxy-noraporphine. N-Methylhernandigerine.

Isobulbocapnine

[5544-68-3]

$C_{19}H_{19}NO_4$ 325.363

Alkaloid from *Lindera oldhamii*, *Hernandia ovigera*, *Hernandia jamaicensis*, *Hernandia catalpifolia* and *Hernandia peltata* (Lauraceae, Hernandiaceae). Shows antiplatelet aggregation activity. Mp 169-170°. $[\alpha]_D^{32}$ +300 (c, 1 in CHCl₃).

N-Me, hydrobromide: Mp 243-245° dec. $[\alpha]_D$ +170 (H₂O).

N-Me, N-oxide (β-): N-Methylnandigerine β-N-oxide

[122331-85-5]

$C_{19}H_{19}NO_5$ 341.363

Alkaloid from the leaves of *Polyalthia longifolia* (Annonaceae). Light greyish needles (MeOH). Mp 218-220°. $[\alpha]_D$ +252 (c, 0.1 in MeOH).

N-Hydroxy: N-Hydroxynandigerine. N-Hydroxyhernandigerine

$C_{18}H_{17}NO_5$ 327.336

Alkaloid from trunk bark of *Hernandia nymphaeifolia* (Hernandiaceae). Prisms (CHCl₃/MeOH). Mp 99-101°.

$[\alpha]_D^{24}$ +182 (c, 0.08 in CHCl₃). λ_{\max} 223 (log ϵ 4.39); 269 (log ϵ 4.11); 309 (log ϵ 3.77) (EtOH).

(±)-form

N-Me: Synthetic. Tablets (EtOH/Et₂O). Mp 165-166°.

Hey, D.H. *et al.*, *J.C.S.*, 1954, 2246 (*N-Me, synth*)

Cava, M.P. *et al.*, *Tet. Lett.*, 1966, 1577; 4279 (*Nandigerine, N-Methylnandigerine, uv, pmr, struct*)

Cava, M.P. *et al.*, *Tetrahedron*, 1971, **27**, 2639 (*isol*)

Lu, S.-T. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 910; *CA*, **77**, 101949m (*N-Methylnandigerine*)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (*cd*)

Wu, Y.-C. *et al.*, *Heterocycles*, 1989, **29**, 463 (*N-Methylnandigerine β-N-oxide*)

Chen, I.-S. *et al.*, *Planta Med.*, 1995, **61**, 537-539 (*Nandigerine, N-Methylnandigerine, activity*)


Chen, I.-S. *et al.*, *Heterocycles*, 1996, **43**, 799 (*N-formyl*)

Chen, I.-S. *et al.*, *Planta Med.*, 1997, **63**, 154-157 (*N-hydroxy*)

Nangustine

N-26

[474794-31-5]



Absolute Configuration

$C_{16}H_{17}NO_4$ 287.315


Alkaloid from fresh bulbs of *Narcissus angustifolius* ssp. *transcarpathicus*. Mp 261°. $[\alpha]_D^{20}$ -69.6 (c, 0.35 in MeOH).

Labraña, J. *et al.*, *Phytochemistry*, 2002, **60**, 847-852 (*isol, ir, cd, pmr, cmr, ms, abs config*)

Kokas, O.J. *et al.*, *Tetrahedron*, 2008, **64**, 6444-6451 (*synth*)

Nankakurine A

N-27



$C_{17}H_{30}N_2$ 262.437

Stereochem. revised in 2006. Alkaloid from *Lycopodium hamiltonii*. Amorph. solid. $[\alpha]_D^{21}$ +16 (c, 0.4 in MeOH).

N²-Me: Nankakurine B

$C_{18}H_{32}N_2$ 276.464

Alkaloid from *Lycopodium hamiltonii*. Amorph. solid. $[\alpha]_D^{19}$ +12 (c, 1 in MeOH).

Hirasawa, Y. *et al.*, *Org. Lett.*, 2004, **6**, 3389-3391 (*isol, pmr, cmr*)


Hirasawa, Y. *et al.*, *Heterocycles*, 2006, **68**, 2357-2362 (*isol, pmr, cmr, abs config*)

Nilsson, B.L. *et al.*, *J.A.C.S.*, 2008, **130**, 11297-11299 (*synth, abs config*)

Nannochelin C

N-28

[133705-26-7]

C₃₆H₄₄N₄O₁₃ 740.763

Isol. from the myxobacterium *Nannocystis exedens*. Siderophore. Sol. H₂O, MeOH, EtOAc, Me₂CO; fairly sol. CHCl₃, Et₂O; poorly sol. H₂O. [α]_D -0.5 (c, 0.5 in MeOH aq.). λ_{max} 280 (ε 25100) (MeOH) (Derep). λ_{max} 280 (ε 25000) (MeOH) (Berdy).

1-Me ester: Nannochelin B

[133705-27-8]

C₃₇H₄₆N₄O₁₃ 754.789

Isol. from *Nannocystis exedens*. Siderophore. [α]_D -5.3 (c, 1 in MeOH aq.). λ_{max} 280 (ε 25100) (MeOH) (Derep).

1,1'-Di-Me ester: Nannochelin A

[133705-25-6]

C₃₈H₄₈N₄O₁₃ 768.816

Isol. from *Nannocystis exedens*. Siderophore. [α]_D -13 (c, 0.9 in MeOH). Possibly an artifact. λ_{max} 280 (ε 25100) (MeOH) (Derep).

Kunze, B. *et al.*, *J. Antibiot.*, 1992, **45**, 147 (isol, uv, ir, props)


Mulqueen, G.C. *et al.*, *Tetrahedron*, 1993, **49**, 9137 (synth)

Sakamoto, T. *et al.*, *J.O.C.*, 1996, **61**, 8496 (synth)

Nantentine

N-29

5,6,6a,7-Tetrahydro-1,2-dimethoxy-6-methyl-4H-benzo[de][1,3]benzodioxolo[5,6-g]quinoline, 9Cl. 1,2-Dimethoxy-9,10-methylenedioxyaporphine. Domesticine. Epidicentrine. O-Methyldomesticine [42971-15-3]



(S)-form

C₂₀H₂₁NO₄ 339.39

► CE0450000

(R)-formMp 138-139°. [α]_D¹⁸ -101 (CHCl₃).**(S)-form**

Alkaloid from *Nandina domestica* and *Cassytha pubescens*, also from *Ocotea*, *Corydalis* and *Papaver* spp. (Nandinaaceae, Lauraceae, Papaveraceae). Mp 139-141°. [α]_D +93 (c, 0.17 in CHCl₃). λ_{max} 217 (log ε 4.52); 234 (sh) (log ε 4.3);

273 (sh) (log ε 3.92); 282 (log ε 3.96); 309 (log ε 4.05); 318 (sh) (log ε 4.01) (95% EtOH).

N-Me: N-Methylnantentine

[56799-50-9]

C₂₁H₂₄NO₄[⊕] 354.425

Quaternary alkaloid from *Thalictrum polygamum* and *Thalictrum sachalinense* (Ranunculaceae). Mp 213-214° (as chloride). [α]_D +39 (EtOH) (chloride). λ_{max} 225 (log ε 4.24); 278 (sh) (log ε 3.65); 285 (log ε 3.78); 310 (log ε 3.94); 320 (sh) (log ε 3.84) (EtOH).

N-De-Me: 1,2-Dimethoxy-9,10-methylenedioxyaporphine. Nornantentine

[15401-66-8]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Nandina domestica*, *Cassytha racemosa*, *Laurelia sempervirens*, *Laurelia philippiana*, *Xylopia danguyella*, *Hernandia nymphaeifolia*, *Hernandia cordigera* and *Hernandia peltata* (Nandinaaceae, Lauraceae, Annonaceae, Hernandiaceae). Shows antimicrobial activity. Prisms (Me₂CO/petrol). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 163-164°. [α]_D +85 (c, 0.75 in CHCl₃). λ_{max} 218 (log ε 4.61); 282 (log ε 4.06); 308 (log ε 4.15) (no solvent reported).

N-De-Me, N-formyl: N-Formylnornantentine

[123715-09-3]

C₂₀H₁₉NO₅ 353.374

Alkaloid from *Cyclea atjehensis* (Menispermaceae). Cryst. (MeOH). Mp 232°. [α]_D +292 (c, 0.13 in MeOH). Exists in CDCl₃ soln. as a mixt. of rotamers about the amide bond.

N-De-Me, N-Ac: N-Acetylnornantentine

[15401-67-9]

C₂₁H₂₁NO₅ 367.401

Alkaloid from the heartwood of *Liriodendron tulipifera* (Magnoliaceae). Needles (EtOH). Mp 294° (283-284°). [α]_D +349 (c, 0.44 in CHCl₃). λ_{max} 215 (log ε 4.61); 281 (log ε 4.04); 307 (log ε 4.12) (MeOH).

O¹-De-Me: 1-Hydroxy-2-methoxy-9,10-methylenedioxyaporphine. Domesticine

[476-71-1]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Nandina domestica* and *Cassytha pubescens*, also present in *Corydalis* and *Glaucaum* spp. (Nandinaaceae, Lauraceae, Papaveraceae). Mp 115-117°. [α]_D +44.4 (c, 0.6 in CHCl₃). (R)-enantiomer also known (synthetic). λ_{max} 221 (log ε 4.5); 283 (log ε 4.01); 310 (log ε 4.17) (EtOH).

O¹-De-Me, N-Me: N-Methyl domesticine

[96400-62-3]

C₂₀H₂₂NO₄[⊕] 340.398

Quaternary alkaloid from the aerial parts of *Glaucaum oxylobum* (Papaveraceae). Thin prisms (MeOH) (as iodide). Mp 255-256° (iodide). CAS no. refers to iodide. λ_{max} 224 (log ε 4.64); 284 (log ε 3.95); 311 (log ε 4.06) (no solvent reported) (as iodide).

O¹-De-Me, N-de-Me: 1-Hydroxy-2-methoxy-9,10-methylenedioxyaporphine**phine. Nordomesticine**

[14787-35-0]

C₁₈H₁₇NO₄ 311.337

Alkaloid from *Corydalis pubescens* (Papaveraceae). Amorph. [α]_D +31.4 (c, 0.83 in CHCl₃).

O¹-De-Me, N-de-Me, N,O-di-Ac:

Cryst. (EtOH). Mp 252-254° (235-236°). [α]_D +326.4 (c, 0.4 in CHCl₃).

O²-De-Me: 2-Hydroxy-1-methoxy-9,10-methylenedioxyaporphine. Isodomesticine

[70560-83-7]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Nandina domestica* and from the leaves of *Laurus nobilis* (bay laurel) (Nandinaaceae, Lauraceae). Amorph. Mp 85°. λ_{max} 216 (log ε 4.55); 283 (log ε 3.99); 310 (log ε 4.07) (EtOH).

O²-De-Me, N-de-Me: 2-Hydroxy-1-methoxy-9,10-methylenedioxyaporphine. Norisodomesticine

[80151-84-4]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the leaves of *Xylopia danguyella* and *Laurus nobilis* (bay laurel) (Annonaceae). Amorph.

6a,7-Didehydro: Dehydronantentine. Alkaloid CC 1

[55898-15-2]

C₂₀H₁₉NO₄ 337.374

Alkaloid from *Ocotea macrophylla*, *Nandina domestica*, *Corydalis cava*, *Corydalis marshalliana*, *Corydalis bulbosa* and *Corydalis slivenensis* (Lauraceae, Nandinaaceae, Papaveraceae). Yellow needles (C₆H₆/cyclohexane). Mp 197°. λ_{max} 258 (log ε 4.51); 295 (log ε 3.95); 333 (log ε 3.82) (EtOH).

6a,7-Didehydro, N-de-Me, N-formyl:**6a,7-Dehydro-N-formylnornantentine**

[163404-70-4]

C₂₀H₁₇NO₅ 351.358

Alkaloid from rhizomes of *Aristolochia brevipes* (Aristolochiaceae). Cryst. Mp 223-227°. Exhibits E/Z-stereoisomerism of the N-formyl group.

3-Chloro, N-de-Me, N-formyl: 3-Chloro-N-formylnornantentine

[276689-73-7]

C₂₀H₁₈ClNO₅ 387.819

Alkaloid from *Lindera glauca*. Needles (CHCl₃). [α]_D²⁴ +165 (c, 0.1 in MeOH). Mp >300°. λ_{max} 272 (sh); 281; 302; 312 (sh) (no solvent reported).

4R-Hydroxy, N-de-Me: 4-Hydroxynornantentine

[69128-21-8]

C₁₉H₁₉NO₅ 341.363

Alkaloid from the bark of *Laurelia philippiana*. Also present in the wood of *Laurelia sempervirens* (Peruvian nutmeg) (Monimiaceae). Oil which darkens rapidly on contact with air; cryst. (EtOH) (di-Ac). Mp 179-181° (as di-Ac).

7-Hydroxy: Hydroxynantentine

[53694-46-5]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Nandina domestica* (Nandinaaceae).

(±)-form

Mp 142°.

Hydrochloride: Mp 265-270°.N-*De-Me*: [65732-35-6]Synthetic. Cryst. (Me₂CO). Mp 160°.N-*De-Me*, N-*Ac*: [40042-35-1]


Synthetic. Needles (EtOH). Mp 255-256° (244-245°).

O¹-*De-Me*: [24316-05-0]

Synthetic. Needles (MeOH). Mp 185-186° (181-183°).

O¹-*De-Me*, N-*de-Me*: Synthetic. Cryst. (CH₂Cl₂/hexane). Mp 205-207°.O²-*De-Me*: Synthetic. Cream-coloured cryst. (Et₂O/hexane). Mp 180-183° dec.O²-*De-Me*, *hydrochloride*:Tan-coloured cryst. + ½ MeOH (MeOH/Et₂O). Mp 245-250° dec.Tomita, M. *et al.*, *Yakugaku Zasshi*, 1961, **81**,1090-1093 (*Domesticine*, *Nantentine*, *isol*)Craig, J.C. *et al.*, *Tetrahedron*, 1965, **21**, 395-399 (*uv*, *abs config*)Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, **19**,2331-2338; 1967, **20**, 1457-1462(*Nordomesticine*, *Nantentine*, *Nornantentine*, *isol*, *pmr*, *uv*, *struct*)Govindachari, T.R. *et al.*, *Indian J. Chem.*,*Sect. B*, 1969, **7**, 841-843; 1970, **8**, 16-18(*Domesticine*, *Isodomesticine*, *synth*)Kametani, T. *et al.*, *J.C.S. (C)*, 1971, 2446-2448; 2712-2714 (*Domesticine*, *synth*)Ghosal, C.R. *et al.*, *Chem. Ind. (London)*,1972, 889-890 (*N-Acetylnornantentine*,*synth*)Hufford, C.D. *et al.*, *J. Pharm. Sci.*, 1974, **63**,1338-1339; 1975, **64**, 789-798 (*N-**Acetylnornantentine*)Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1974, **94**,1149-1153 (*Hydroxynantentine*)Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1975,**23**, 2048-2053; 1978, **26**, 3920 (*Domesticine*,*Isodomesticine*, *synth*)Shamma, M. *et al.*, *Heterocycles*, 1975, **3**, 297-300 (*N-Methylnantentine*)Franca, N.C. *et al.*, *Phytochemistry*, 1975, **14**,1671-1672 (*Dehydronantentine*, *isol*, *struct*)Manikumar, G. *et al.*, *Indian J. Chem., Sect. B*,1977, **15**, 740-741 (*Nornantentine*, *synth*)Umarova, D. *et al.*, *Khim. Prir. Soedin.*, 1978,**14**, 594-597; *Chem. Nat. Compd. (Engl.**Transl.)*, 1978, **14**, 511-513 (*N-**Methylnantentine*)Urzúa, A. *et al.*, *Tet. Lett.*, 1978, 2649-2652 (*4-**Hydroxynornantentine*)Zabel, V. *et al.*, *Acta Cryst. B*, 1979, **35**, 3126-3129 (*4-Hydroxynornantentine*, *cryst struct*)Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1979, **42**,325; 329 (*Domesticine*, *rev*, *cmr*)Hoquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**,551-556 (*Norisodomesticine*)Pech, B. *et al.*, *J. Nat. Prod.*, 1982, **45**, 560-563(*Isodomesticine*)Villar, A. *et al.*, *Farm. Tijdschr. Belg.*, 1984, **61**,300 (*activity*)Slavíková, L. *et al.*, *Coll. Czech. Chem. Comm.*,1985, **50**, 854-860 (*N-Methyl-domesticine*)Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1989, **52**,652-654 (*N-Formylnornantentine*)Ribár, B. *et al.*, *Acta Cryst. C*, 1991, **47**, 2500-2501 (*cryst struct*)Ozaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**,481-483 (*Nantentine*, *synth*)Achenbach, H. *et al.*, *Planta Med.*, 1995, **61**,189-190 (*6a,7-Dehydro-N-**formylnornantentine*)Hara, H. *et al.*, *Tetrahedron: Asymmetry*, 1995,**6**, 1683-1692 (*Nordomesticine*, *synth*)Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**,165-171 (*N-Methylnantentine*, *cmr*)Chang, Y.-C. *et al.*, *J. Chin. Chem. Soc.*(Taipei), 2000, **47**, 373-380 (*3-Chloro-N-**formylnornantentine*)**Napelline****N-30****21-Ethyl-4-methyl-16-methylene-7,20-cyclovecatchane-1,12,15-triol, 9CI. Luciculine**

[5008-52-6]



Absolute Configuration

C₂₂H₃₃NO₃ 359.508Alkaloid from *Aconitum napellus*, *Aconitum karakolicum* and *Aconitum yesoense* (Ranunculaceae). Cryst. + 1H₂O. Mp 117-118.5°. [α]_D²¹ -13 (MeOH).*Hydrochloride*: Mp 220-222° dec. [α]_D²² -93.9 (c, 5 in H₂O). Probably contains solvent of crystallisation.*Hydrobromide*:Cryst. (MeOH/Et₂O). Mp 229° dec. (darkens at 200°). [α]_D²³ -42.7 (c, 6 in H₂O).N-*Oxide*: **Napelline N-oxide. Flavamine. Luciculine N-oxide**

[74047-88-4]

C₂₂H₃₃NO₄ 375.507Alkaloid from the above-ground parts of *Aconitum karakolicum* (Ranunculaceae) and *Aconitum flavum*. Mp 224-226°. [α]_D²² -30.5 (c, 0.61 in MeOH).**1-Ac: 1-Acetylnapelline. 1-Acetyl-****luciculine**

[68676-52-8]

C₂₄H₃₅NO₄ 401.545Alkaloid from *Aconitum yesoense*. Amorph. [α]_D²¹ +3.6 (c, 0.28 in MeOH).**12-Ac: 12-Acetylnapelline. 12-Acetyl-****luciculine**

[62511-84-6]

C₂₄H₃₅NO₄ 401.545Alkaloid from *Aconitum karakolicum*

(Ranunculaceae). Mp 205-206°.

12-Ac, N-oxide: 12-Acetylnapelline N-**oxide**

[83019-08-3]

C₂₄H₃₅NO₅ 417.544Alkaloid from the above-ground parts of *Aconitum karakolicum* (Ranunculaceae). Mp 235°.**15-Ac: Lucidusculine**

[5008-49-1]

C₂₄H₃₅NO₄ 401.545Alkaloid from the roots of *Aconitum lucidusculum* (Ranunculaceae). Plates (MeOH). Mp 170-171°. [α]_D -95.5 (CHCl₃).**15-Ac, hydrochloride**: Mp 245-265° dec.**15-Ac, N-oxide: Flavadin**

[119188-45-3]

C₂₄H₃₅NO₅ 417.544Alkaloid from the roots of *Aconitum flavum* (Ranunculaceae). Mp 198-200°.**12,15-Di-Ac: 12-Acetylucidusculine**C₂₆H₃₇NO₅ 443.582Alkaloid from roots of *Aconitum flavum*, and rhizomes of *Aconitum yesoense* var. *macroyesoense* (Ranunculaceae). Mp 132-134° Mp 144-147°. [α]_D -19.2 (c, 0.5 in EtOH).N-*De-Et*, N-*Me*: **Finetianine**

[112543-24-5]

C₂₁H₃₁NO₃ 345.481Alkaloid from roots of *Aconitum finetianum* (Ranunculaceae).**12-Ketone: Songorine. Napellonine. Bul-****latine G. Shimoburo base I. Zongorine**

[509-24-0]

C₂₂H₃₁NO₃ 357.492Alkaloid from the roots of *Aconitum soongoricum* and *Aconitum monticola*, the above-ground parts of *Aconitum karakolicum*, and from the Chinese drug "Fuzi" (*Aconitum carmichaeli*) (Ranunculaceae). Cryst. (MeOH). Mp 212° (201-203°). [α]_D -140.▶ LD₅₀ (rat, ipr) 408 mg/kg. WG0780000**12-Ketone, hydrochloride:**Cryst. (MeOH). Mp 257-258°. [α]_D -114 (H₂O).**12-Ketone, N-oxide: Songorine N-oxide**

[66921-56-0]

C₂₂H₃₁NO₄ 373.491Alkaloid from above-ground parts of *Aconitum monticola* (Ranunculaceae). Mp 253-255°.**12-Ketone, N-de-Et: Norsongorine. N-Deethylsongorine**

[29722-71-2]

C₂₀H₂₇NO₃ 329.438Alkaloid from the roots of *Aconitum monticola* (Ranunculaceae). Mp 284-286°. [α]_D -86 (MeOH).**12-Ketone, 1 or 15-Ac: Acetylsongorine**

[82621-08-7]

C₂₄H₃₃NO₄ 399.529Alkaloid from the roots of *Aconitum soongoricum*.**12-Ketone, O¹-Me: Liangshanone**

[137031-52-8]

C₂₃H₃₃NO₃ 371.519Alkaloid from *Aconitum liangshanum* (Ranunculaceae). Amorph. [α]_D^{22.5} -101 (c, 1.1 in CHCl₃).**1,12-Diketone: 1-Dehydrosongorine. 1-****Deoxy-1-oxosongorine**

[112515-31-8]

C₂₂H₂₉NO₃ 355.476Alkaloid from roots of *Aconitum finetianum* (Ranunculaceae).**16β,17-Dihydro, 12-ketone: Dihydro-****songorine**

[29603-32-5]

C₂₂H₃₃NO₃ 359.508Alkaloid from aerial parts of *Aconitum karakolicum* (Ranunculaceae). Cryst. (EtOH). Mp 202-204°.**11β-Hydroxy: Turpelline**

[169626-33-9]

C₂₂H₃₃NO₄ 375.507

Alkaloid from above-ground parts of *Aconitum turczaninowii*. Mp 268-271°.

1-Epimer: 1-Epinapelline

[110064-70-5]

C₂₂H₃₃NO₃ 359.508

Alkaloid from roots of *Aconitum flavum* (Ranunculaceae). Mp 87-89°. [α]_D²⁰ -11.7 (c, 1.19 in MeOH).

12-Epimer: 12-Epinapelline

[110064-71-6]

C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Aconitum flavum* and aerial parts of *Aconitum karakolicum* (Ranunculaceae). Also present in aerial parts of *Aconitum baicalense*. Cryst. (Me₂CO). Mp 72-73.5°. [α]_D²⁵ -40.2 (c, 1.03 in CHCl₃).

12-Epimer, N-oxide: 12-Epinapelline N-oxide

[169336-56-5]

[169336-57-6]

C₂₂H₃₃NO₄ 375.507

Alkaloid from aerial parts of *Aconitum baicalense*. Cryst. (H₂O) (as perchlorate). Mp 224-225° (perchlorate).

12-Epimer, 12-Ac: 12-Acetyl-12-epinapelline

C₂₄H₃₅NO₄ 401.545

Alkaloid from *Aconitum soongoricum*. Cryst. (Me₂CO). Mp 198.5-200.5°.

12-Epimer, 15-Ac: 12-Epilucidusculine

[137120-52-6]

C₂₄H₃₅NO₄ 401.545

Alkaloid from *Aconitum liangshanium* (Ranunculaceae). Prisms. Mp 160-164°. [α]_D^{21.5} -100 (c, 0.91 in CHCl₃).

12-Epimer, N-de-Et, N-Me: 12-Epifinetianine, N-Deethyl-N-methyl-12-epinapelline

C₂₁H₃₁NO₃ 345.481

Alkaloid from the roots of *Aconitum nagorum* var. *lasiandrum*. Powder. Mp 187-188°. [α]_D²⁰ -40 (c, 0.1 in CHCl₃).

12-Epimer, O'-Me: Liangshanine†

[137031-43-7]

C₂₃H₃₅NO₃ 373.534

Alkaloid from *Aconitum liangshanium* (Ranunculaceae). Amorph. [α]_D^{10.5} -16.4 (c, 1.5 in CHCl₃).

12-Epimer, 16β,17-dihydro, 15-ketone:

Karakomine

[142768-81-8]

C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Aconitum karakolicum* (Ranunculaceae). Yellow powder. [α]_D²⁰ -18.7 (c, 0.27 in EtOH).

12-Epimer, 3α-hydroxy: 3α-Hydroxy-12-epinapelline

C₂₂H₃₃NO₄ 375.507

Alkaloid from the roots of *Aconitum brunneum*. Cryst. Mp 201-202°. [α]_D²⁷ -11.6 (c, 0.5 in CHCl₃).

Majima, R. *et al.*, *Ber.*, 1932, **65**, 599

(*Lucidusculine, isol*)

Freudenberg, W. *et al.*, *J.A.C.S.*, 1937, **59**, 2572 (*isol*)

Sugasawa, T. *et al.*, *Chem. Pharm. Bull.*, 1956, **4**, 6; 1961, **9**, 889; 897 (*Songorine, struct*)

Wiesner, K. *et al.*, *Experientia*, 1958, **14**, 167 (*Songorine, Napelline, struct*)

Kuzovkov, A.D. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1958, **28**, 2320; 1959, **29**, 1706 (*Songorine, struct*)

Suginome, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1959, **32**, 352; 824 (*Lucidusculine, struct*)

Amiya, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1959, **32**, 1133; 1960, **33**, 1175; 1961, **34**, 898 (*Lucidusculine, struct, uv, ir*)

Suzuki, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1961, **34**, 455 (*Lucidusculine, pmr*)

Okamoto, T. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 1270 (*cryst struct, abs config*)

Yoshino, A. *et al.*, *Acta Cryst.*, 1966, **21**, 57 (*Lucidusculine, cryst struct, abs config*)

Yunusov, M.S. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 101; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 95 (*Songorine, isol, ms, synth*)

Wiesner, K. *et al.*, *Can. J. Chem.*, 1973, **51**, 3978; 1974, **52**, 2355 (*synth*)

Nezhevenko, V.E. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 409; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 416 (*Songorine, Norsongorine*)

Sultankhodzaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 681; 1978, **14**, 479; 1982, **18**, 265; 660; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 617; 1978, **14**, 407; 1982, **18**, 249; 629 (*isol, struct, derivs*)

Ametova, E.F. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 867; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 735 (*Songorine N-oxide*)

Zhamerashvili, M.E. *et al.*, *CA*, 1982, **97**, 107014n (*Acetylsongorine*)

Pelletier, S.W. *et al.*, *Heterocycles*, 1982, **18**, 47 (*occur*)

Sultankhodzaev, M.N. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 265; 660; 1987, **23**, 917; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 249; 629; 1987, **23**, 772 (*Songorine, Dihydrosongorine*)

Takayama, H. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 245-247 (*1-Acetyl napelline*)

Bando, H. *et al.*, *Heterocycles*, 1987, **20**, 2623 (*Lucidusculine, 12-Acetyl lucidusculine, isol, ir, ms, pmr, struct*)

Chen, Z. *et al.*, *Heterocycles*, 1987, **26**, 1455 (*1-Epinapelline, 12-Epinapelline*)

Tian, R. *et al.*, *Huaxue Xuebao*, 1987, **45**, 776; *CA*, **108**, 72040j (*Finetianine, 1-Dehydrosongorine*)

Chen, Z.-G. *et al.*, *Planta Med.*, 1988, **54**, 318 (*Flavadine*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1644 (*Liangshanine, Liangshanone, 12-Epilucidusculine*)

Huang, B. *et al.*, *Heterocycles*, 1991, **32**, 2429 (*Karakomine*)

Batbayar, N. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 740; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 658 (*Turpelline*)

Zhapova, Ts. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 888; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 791 (*12-Epinapelline N-oxide, isol, pmr, cmr, ir*)

Salimov, B.T. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2004, **40**, 151-155 (*12-Acetyl-12-epinapelline*)

Gao, L. *et al.*, *Heterocycles*, 2004, **63**, 1181-1184 (*3α-Hydroxy-12-epinapelline*)

Atta-ur-Rahman, *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 609-610 (*12-Epinapelline, cryst struct*)


Zhang, F. *et al.*, *Planta Med.*, 2005, **71**, 1073-1076 (*12-Epifinetianine*)

Csupor, D. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 2981-2986 (*Songorine, synth*)

Naphthisoazole A

N-31

9,9a-Dihydronaphtho[2,3-c]isoxazol-3(1H)-one



C₁₁H₉NO₂ 187.198

(±)-form


Alkaloid from the roots of *Glehnia littoralis*. Yellowish oil. [α]_D²⁵ -62.6 (c, 0.57 in MeOH). λ_{max} 213; 278 (MeOH).

Li, G.Q. *et al.*, *Fitoterapia*, 2008, **79**, 238-239 (*isol, pmr, cmr*)

Naphthyridinomycin A

N-32

[54913-26-7]



Absolute Configuration

C₂₁H₂₇N₃O₆ 417.461

Benzoquinone-type antibiotic. Poss. artifact. Further attempts to reisolate this compd. gave Bioxalomyacin β2 in B-140. U. Isol. from *Streptomyces luctitanus* NRRL 8034. Antitumour agent. Ruby-red orthorhombic cryst. (Et₂O). Sol. MeOH, Et₂O, H₂O; poorly sol. hexane. Mp 108-110° dec. [α]_D²⁵ +69.4 (c, 1 in CHCl₃). λ_{max} 270 (ε 10400) (MeOH) (Derep).

O-De-Me: Antibiotic SF 1739HP. SF 1739HP

[72146-18-0]

C₂₀H₂₅N₃O₆ 403.434

Prod. semisynthetically from Antibiotic SF 1739, A-1254 on treatment with mineral acid. Shows marked activity against gram-negative bacteria and tumours with low toxicity. Stable form of SF 1739. Purple powder. [α]_D²⁵ +64 (c, 0.5 in MeOH). λ_{max} 275 (E1%/1cm 165); 330 (sh) (MeOH).

Cluepfel, D. *et al.*, *J. Antibiot.*, 1975, **25**, 497-502 (*isol, uv, ir, pmr, struct*)

Syngusch, J. *et al.*, *Acta Cryst. B*, 1976, **32**, 1139-1142 (*cryst struct*)

Zmijewski, M.J. *et al.*, *Antimicrob. Agents Chemother.*, 1982, **21**, 787-793 (*props*)

Itoh, I. *et al.*, *J. Antibiot.*, 1982, **35**, 642-644 (*SF 1739HP*)

Zmijewski, M.J. *et al.*, *J.A.C.S.*, 1982, **104**, 4969-4971 (*biosynth*)

Fukuyama, T. *et al.*, *Adv. Heterocycl. Nat. Prod. Synth.*, 1992, **2**, 189-249 (*rev, synth, abs config*)

Zaccardi, J. *et al.*, *J.O.C.*, 1994, **59**, 4045-4047 (*abs config, bibl*)

Scott, J.D. *et al.*, *Chem. Rev.*, 2002, **102**, 1669-1730 (*rev*)

Naphthridinomycin B N-33


[62046-88-2]

Struct. unknown. Prod. by *Streptomyces lusitanus*. Antibacterial agent. Yellow powder. λ_{\max} 282 (MeOH).

U.S. Pat., 1977, 4 003 902; *CA*, **86**, 119256d (isol)

Napsamycin B N-34

[144408-86-6]



$C_{40}H_{50}N_8O_{12}S$ 866.947

Prod. by *Streptomyces candidus* and *Streptomyces* sp. HIL Y82, 11372. Active against *Pseudomonas* sp. Sol. H_2O , DMSO, MeOH- H_2O ; poorly sol. EtOAc, hexane. Mp 190° (dec.). λ_{\max} 240 (ϵ 36000); 292 (ϵ 6800) (dil. NaOH) (Derep). λ_{\max} 256 (ϵ 21000) (H_2O) (Derep). λ_{\max} 256 (HCl) (Berdy).

5'',6''-Dihydro: Napsamycin D

[144379-27-1]


$C_{40}H_{52}N_8O_{12}S$ 868.963

Prod. by *Streptomyces candidus* and *Streptomyces* sp. HIL Y82, 11372. Active against *P.* sp. Sol. H_2O , DMSO, MeOH- H_2O ; poorly sol. EtOAc, hexane. Mp 190° (dec.). λ_{\max} 240 (ϵ 36000); 292 (ϵ 6800) (dil. NaOH) (Derep). λ_{\max} 256 (ϵ 21000) (H_2O) (Derep). λ_{\max} 256 (HCl) (Berdy).

Eur. Pat., 1992, 487 756; *CA*, **117**, 232207 (isol) Chattarjee, S. et al., *J. Antibiot.*, 1994, **47**, 595 (isol, pmr, ms)

Narceine N-35

6-[[6-[2-(Dimethylamino)ethyl]-4-methoxy-1,3-benzodioxol-5-yl]acetyl]-2,3-dimethoxybenzoic acid, 9CI [131-28-2]



$C_{23}H_{27}NO_8$ 445.468

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Respiratory stimulant, antitussive agent. Shows

no analgesic activity. Antihypertensive agent. Intestinal smooth muscle. Cryst. (MeOH/Et₂O or H_2O). Mp 145° (anhyd.) Mp 176-178° (trihydrate). pK_{a1} 3.5; pK_{a2} 9.3 (15°). Log P -1.79 (uncertain value) (calc).

▶ YX5210000

Hydrochloride: Mp 192-193°.

Picrate: Mp 195°.

N-Me: Mp 207° (as iodide).

N-De-Me: Nornarceine. Oxynarcotine [483-89-6]

$C_{22}H_{25}NO_8$ 431.441

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Cryst. (EtOH). Mp 229° (223-225°). Prob. artifact of demethylation of Narcotine. λ_{\max} 209 (ϵ 55700); 230 (sh) (ϵ 22000); 273 (ϵ 14500) (EtOH).

1-Oxo: Narceinone

[125187-48-6]

$C_{23}H_{25}NO_9$ 459.452

Alkaloid from unlanded dried capsules of *Papaver somniferum* (opium poppy) (Papaveraceae). Cryst. (MeOH). Mp 162°.

8-Demethoxy: N-Methylhydrasteine

[31971-15-0]

$C_{22}H_{25}NO_7$ 415.442

Alkaloid from *Fumaria parviflora*, *Fumaria vaillantii*, *Fumaria schleicheri*, *Corydalis lutea*, *Dactylicapnos torulosa* and *Corydalis solida* (Papaveraceae). Cryst. (EtOH, H_2O or $CHCl_3$ /MeOH). Mp 146-147° Mp 150° (hydrate) Mp 223°. Poss. artifact. A uv spectrum has been reported but contains a typographical error. See also Coryrutine, C-698.

8-Demethoxy, Me ester: N-Methylhydrasteine methyl ester

[80243-94-3]

$C_{23}H_{27}NO_7$ 429.469

Isol. from *Dactylicapnos torulosa*. Powder. Mp 93-94°. May be an artifact. λ_{\max} 208 (log ϵ 4.45); 226 (log ϵ 4.33); 273 (log ϵ 4.11); 293 (log ϵ 4.13); 377 (log ϵ 3.78) (MeOH).

8-Demethoxy, 1-oxo: N-Methyloxohydrasteine

[41515-85-9]

$C_{22}H_{23}NO_8$ 429.426

Alkaloid from *Fumaria microcarpa* and *Fumaria officinalis* (Papaveraceae). Cryst. ($CHCl_3$ /MeOH, H_2O or MeOH). Mp 234-235° Mp 203° (hydrate) Mp 168° (MeOH solvate). See also Oxocoryrutine in C-698.

8-Demethoxy, 1-oxo, Et ester: Papracine

[144525-63-3]

$C_{24}H_{27}NO_8$ 457.479

Alkaloid from whole plants of *Fumaria indica* (Papaveraceae).

Rabe, P. et al., *Annalen*, 1910, **377**, 223-258 (struct, bibl)

Addinall, C.R. et al., *J.A.C.S.*, 1933, **55**, 1202-1209; 2153-2163 (isol, synth)

Briggs, L.H. et al., *Anal. Chem.*, 1957, **29**, 904-911 (ir)

Klötzer, W. et al., *Monatsh. Chem.*, 1972, **103**, 1210-1212 (*Nornarceine, synth, uv, ir, pmr*)

Forgacs, P. et al., *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1973, **276**, 105-108; 1974, **279**, 855-857 (*N-Methylhydrasteine, N-Methyloxohydrasteine*)

Blaskó, G. et al., *J.O.C.*, 1982, **47**, 880-885 (synth, uv, pmr)

Chaudhuri, P.K. et al., *Phytochemistry*, 1989, **28**, 2002-2003 (*Narceinone*)


Atta-ur-Rahman, et al., *Fitoterapia*, 1992, **63**, 129-135 (*Papracine*)

Rucker, G. et al., *Phytochemistry*, 1994, **36**, 519-523 (*N-Methylhydrasteine methyl ester*)

Baggio, R. et al., *Acta Cryst. C*, 1996, **52**, 703-705 (cryst struct)

Narceine imide N-36

3-[[6-[2-(Dimethylamino)ethyl]-4-methoxy-1,3-benzodioxol-5-yl]methylene]-2,3-dihydro-6,7-dimethoxy-1H-isoindol-1-one, 9CI [38952-67-9]



(Z)-form

$C_{23}H_{26}N_2O_6$ 426.468

(E)-form [75794-87-5]

Cryst. (Me₂CO/cyclohexane). Mp 142-144°.

(Z)-form [55968-77-9]

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Cryst. (Me₂CO). Mp 151-152°. May be an artifact of work-up.

Hydrochloride:

Cryst. (H_2O). Mp 239-240° Mp 249-251°.

Picrate:

Cryst. (EtOH). Mp 249-251°.

N-Oxide: [75794-90-0]

Cryst. (Me₂CO aq.). Mp 158-159°.

(E)-form**8-Demethoxy: Fumaridine. Hydrastine imide**

[30341-98-1]

$C_{22}H_{24}N_2O_5$ 396.442

Alkaloid from *Fumaria parviflora*, *Fumaria schleicheri* and *Fumaria vaillantii* (Papaveraceae). Mp 191°. May be an artifact of work-up.

8-Demethoxy, picrate: Mp 204-205°.

Hodková, J. et al., *J. Nat. Prod.*, 1972, **35**, 61 (isol, ir, uv, pmr, synth)

Shamma, M. et al., *Chem. Comm.*, 1975, 89 (*Fumaridine, uv, ir, pmr, struct*)


Trojánek, J. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 681 (synth, uv, ir)

Proksa, B. et al., *Coll. Czech. Chem. Comm.*, 1980, **45**, 2125 (uv, ir, ms, struct)

Hussain, S.F. et al., *J. Nat. Prod.*, 1981, **44**, 169 (*Fumaridine, isol, pmr, ms*)

Rys, V. et al., *Tetrahedron*, 2003, **59**, 6615-6619 (*Fumaridine, synth*)

Lamblin, M. et al., *Org. Biomol. Chem.*, 2007, **5**, 1466-1471 (synth)

NarciclasineLycoricidinol
[29477-83-6]Absolute
configurationC₁₄H₁₃NO₇ 307.259

Alkaloid from *Narcissus incomparabilis*, all *Narcissus* spp. examined, and present in many other Amaryllidaceae spp. Also found in the Texas grasshopper *Bra-chystola magna*. Protein synthesis inhibitor. Antimitotic agent. Possesses antiviral props. Most active antineoplastic agent of the Amaryllidaceae alkaloids. Light yellowish needles with yellow-green fluor. Mp 232-234° dec. $[\alpha]_D^{25} +145$ (c, 1.5 in EtOH). Log P -2.75 (uncertain value) (calc). Easily dehydrates in acid. λ_{max} 252 (ε 25600); 303 (ε 6400); 330 (sh) (EtOH) (Derep).

▶ LD₅₀ (mus, scu) 5 mg/kg. JI50000002-O-β-D-Glucopyranoside: **Kalbreclasine**
[98900-01-7]C₂₀H₂₃NO₁₂ 469.401

Alkaloid from roots of *Haemanthus kalbreyeri* (Amaryllidaceae). Hygroscopic solvated straw-coloured solid; microcryst. (Me₂CO)(as hexa-O-Ac). Mp 198-201° (hexa-O-Ac). $[\alpha]_D^{22} +78.14$ (c, 0.7 in MeOH).

4-O-β-D-Glucopyranoside: **Narciclasine 4-O-glucoside**

[141544-37-8]

C₂₀H₂₃NO₁₂ 469.401

Alkaloid from the bulbs of *Pancreatium maritimum* (Amaryllidaceae). Needles (MeOH). Mp 210-211°. λ_{max} 252; 285 (MeOH) (Berdy).

1,10bβ-Dihydro: **trans-Dihydronarciclasine**

[40042-05-5]

C₁₄H₁₃NO₇ 309.275

Alkaloid from *Zephyranthes candida* (Amaryllidaceae). Cytotoxic agent. Cryst. (MeOH). Mp 290-291°. $[\alpha]_D^{20} +4.7$ (c, 0.27 in THF). Data given is for synthetic compd. λ_{max} 231 (ε 10960); 239 (ε 10200); 280 (ε 5620); 310 (ε 2140) (MeOH) (Derep).

1,10bβ-Dihydro: **cis-Dihydronarciclasine**

[40042-04-4]

Synthetic. Cryst. (MeOH). Mp 293-294°. $[\alpha]_D^{20} +56$ (c, 0.18 in THF).7-Deoxy: **Lycoricidine**. 7-Deoxynarciclasine. **Margetine**

[19622-83-4]

[58207-40-2]

C₁₄H₁₃NO₆ 291.26

Alkaloid from *Hymenocallis littoralis* and *Lycoris radiata*. Cytotoxic; inhibits division of Ehrlich ascites cells. Plant growth regulator. Mp 214.5-215.5° dec. $[\alpha]_D^{20} +180$ (c, 0.45 in Py) (synthetic). λ_{max} 233 (ε 13800); 248 (ε 14100); 302 (ε 5620) (MeOH).

N-37

7-Deoxy, O,O,O-tri-Ac: Mp 200-201°.

7-Deoxy, O,O,O,N-tetra-Ac: Mp 229-231°.


7-Deoxy, 1,10bα-dihydro: **7-Deoxy-trans-dihydronarciclasine**C₁₄H₁₅NO₆ 293.276

Alkaloid from bulbs of *Hymenocallis littoralis*, *Hymenocallis caribaea*, *Hymenocallis latifolia* and an unidentified *Hymenocallis* sp. Cytotoxic. Also shows activity against a series of RNA viruses. Needles. Mp 303-304° (320-322°, synthetic). $[\alpha]_D^{25} +138$ (c, 0.96 in DMSO). λ_{max} 217 (ε 5000); 260 (ε 2800); 304 (ε 3460) (EtOH).

Okamoto, T. et al., *Chem. Pharm. Bull.*, 1968, **16**, 1860; 1976, **24**, 2977 (*Lycoricidine, isol, uv, ir, pmr, struct, synth*)Piozzi, F. et al., *Tetrahedron*, 1968, **24**, 1119 (*uv, ir, pmr, ms, struct*)Piozzi, F. et al., *Phytochemistry*, 1969, **8**, 1745 (*isol*)Fuganti, C. et al., *Chem. Comm.*, 1972, 239 (*biosynth*)Immirzi, A. et al., *Chem. Comm.*, 1972, 240 (*cryst struct*)Mondon, A. et al., *Chem. Ber.*, 1975, **108**, 445 (*pmr, config*)Paulsen, H. et al., *Annalen*, 1983, 535(*Lycoricidine, synth*)Suffness, M. et al., *Alkaloids (Academic Press)*, 1985, **25**, 205 (*antitumour props*)Ghosal, S. et al., *Phytochemistry*, 1985, **24**, 1825 (*Kalbreclasine*)Pettit, G.R. et al., *J. Nat. Prod.*, 1990, **53**, 176; 1993, **56**, 1682 (*Dihydronarciclasine, 7-Deoxydihydronarciclasine*)Abou-Donia, A.H. et al., *Phytochemistry*, 1991, **30**, 3445 (*4-glucoside*)Evidente, A. et al., *Planta Med.*, 1991, **57**, 293 (*pmr, cmr*)Gabrielson, B. et al., *J. Nat. Prod.*, 1992, **55**, 1569 (*7-Deoxydihydronarciclasine, props*)Hudlicky, T. et al., *J.A.C.S.*, 1992, **114**, 9694 (*Lycoricidine, synth*)Martin, S.F. et al., *Heterocycles*, 1993, **35**, 85 (*Lycoricidine, synth*)Chida, N. et al., *J.O.C.*, 1993, **58**, 4441(*Lycoricidine, synth*)Chida, N. et al., *Heterocycles*, 1996, **43**, 1385 (*7-Deoxydihydronarciclasine, synth*)Rigby, J.H. et al., *J.A.C.S.*, 1997, **119**, 12655-12656; 2000, **122**, 6624-6628 (*synth*)Keck, G.E. et al., *J.A.C.S.*, 1999, **121**, 5176-5190 (*synth*)Gonzalez, D. et al., *Tet. Lett.*, 1999, **40**, 3077-3080 (*synth*)Elango, S. et al., *J.O.C.*, 2002, **67**, 6954-6959 (*synth*)Hudlicky, T. et al., *J.O.C.*, 2002, **67**, 8726-8743 (*synth*)Elango, S. et al., *Tetrahedron*, 2002, **58**, 7335-7338 (*Lycoricidine, synth*)Pettit, G.R. et al., *J. Nat. Prod.*, 2005, **68**, 207-211; 1256-1258; 2006, **69**, 7-13 (*isol, synth, Lycoricidine, cryst struct*)Zhang, H. et al., *Org. Lett.*, 2006, **8**, 247-250 (*Lycoricidine, synth*)Shin, I.-J. et al., *Angew. Chem., Int. Ed.*, 2007, **46**, 2303-2305 (*Dihydronarciclasine, synth*)Wang, Y.-H. et al., *J. Nat. Prod.*, 2007, **70**, 1458-1461 (*7-Deoxydihydronarciclasine*)Matveenko, M. et al., *Org. Lett.*, 2007, **9**, 3683-3685 (*Lycoricidine, synth*)Kornienko, A. et al., *Chem. Rev.*, 2008, **108**, 1982-2014 (*rev, chem, bio*)Cho, Y.-S. et al., *Tetrahedron*, 2008, **64**, 2172-2177 (*Dihydronarciclasine, synth*)Matveenko, M. et al., *Tetrahedron*, 2008, **64**, 4817-4826 (*synth*)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LJC000**Narcicrinine**

[92138-21-1]

N-38


C₁₄H₁₇NO₅ 279.292Alkaloid from the leaves of *Crinum oliganthum* (Amaryllidaceae).Döpke, W. et al., *Z. Chem.*, 1984, **24**, 184 (*uv, ir, ms, struct*)**Narcicriptine**

N-39

Amaryllidaceae alkaloid. Struct. unknown. Alkaloid from *Narcissus* sp. (Amaryllidaceae). No other information recorded.Fuganti, C. et al., *Chim. Ind. (Milan)*, 1967, **49**, 1196-1197**Narcidine**


N-40

[139083-10-6]

C₁₇H₂₁NO₄ 303.35711-Config. is R- (difficult to show unambiguously). Alkaloid from the bulbs of *Narcissus pseudonarcissus* (Amaryllidaceae). $[\alpha]_D +16$ (c, 0.11 in MeOH).Tojo, E. et al., *J. Nat. Prod.*, 1991, **54**, 1387 (*isol, uv, ir, pmr, ms, struct*)**Narcisone**

N-41

[107894-72-4]

C₁₈H₂₁NO₄ 315.368Alkaloid from bulbs of *Narcissus tazetta* (Amaryllidaceae). Needles (MeOH). Mp 158-160°. $[\alpha]_D^{20} -18$ (c, 0.5 in CHCl₃).Abdallah, O.M. et al., *Phytochemistry*, 1993, **34**, 1447-1448 (*isol, uv, ir, pmr, cmr, struct*)**Narcisline**


N-42

Amaryllidaceae alkaloid. Struct. unknown. Alkaloid from *Narcissus* sp. (Amaryllidaceae). No other information recorded.

Fuganti, C. *et al.*, *Chim. Ind. (Milan)*, 1967, **49**, 1196-1197

Narcissidine

[27857-07-4]



Absolute configuration

C₁₈H₂₃NO₅ 333.383

Alkaloid from *Narcissus poeticus* and many other spp. in the Amaryllidaceae. Mp 218-219° dec. [α]_D -32 (CHCl₃).

Picrate: Mp 192° dec.

O³-Ac: 3-O-Acetylnarcissidine

[137760-70-4]

C₂₀H₂₅NO₆ 375.421

Alkaloid from *Hippeastrum puniceum* and bulbs of *Leucojum autumnale* (Amaryllidaceae). Cubes (C₆H₆/CHCl₃) or amorph. solid. Mp 191-192.5° (cubes). [α]_D²⁵ -154.2 (c, 0.25 in CHCl₃). A lower opt. rotn. of -46 was recorded for the amorphous isolate.

O³-Ac, N-oxide: 3-O-Acetylnarcissidine N-oxide

[162521-23-5]

C₂₀H₂₅NO₇ 391.42

Alkaloid from bulbs of *Leucojum autumnale* (Amaryllidaceae). Pillars (Me₂CO/MeOH). Mp 150-152° dec. [α]_D²² -102.3 (c, 0.11 in EtOH).

Di-Ac: Mp 170-172° dec.

N-(Chloromethyl): N-Chloromethylnarcissidinium

[132406-98-5]

C₁₉H₂₅ClNO₅⁺ 382.863

Alkaloid from *Lapiedra martinezii* (Amaryllidaceae). Prob. an artifact of the isol. process. CAS No. refers to the chloride.

O²-De-Me: Parkacine

[27857-08-5]

C₁₇H₂₁NO₅ 319.357

Alkaloid from *Amaryllis parkeri*. Fine needles (Me₂CO). Mp 222-223° dec. [α]_D²⁴ +58 (c, 0.2 in CHCl₃).

O²-De-Me, perchlorate:Prisms (H₂O). Mp 240° dec.**O²-De-Me, Ac: Mp 198° dec.**

Fales, H.M. *et al.*, *J.A.C.S.*, 1958, **80**, 4395 (*uv struct*)

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 645 (*Parkacine*)

Kinstle, T.H. *et al.*, *Tet. Lett.*, 1966, 4659 (*ms struct*)

Clardy, J.C. *et al.*, *J.A.C.S.*, 1970, **92**, 1781 (*cryst struct, ir*)

Fuganti, C. *et al.*, *Chem. Comm.*, 1974, 350 (*biosynth*)

Suau, R. *et al.*, *An. Quim.*, 1990, **86**, 672; *C.A.*, **114**, 98281n (*N-Chloromethylnarcissidinium*)


Quirion, J.C. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1112 (*3-O-Acetylnarcissidine*)

Kihara, M. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 318 (*3-O-Acetylnarcissidine, 3-O-Acetylnarcissidine N-oxide*)

Narcotine

N-44

6,7-Dimethoxy-3-(5,6,7,8-tetrahydro-4-methoxy-6-methyl-1,3-dioxolo[4,5-g]isoquinolin-5-yl)-1(3H)-isobenzofuranone, 9CI. Noscapine, BAN, INN. Anarkotin. Gnoscopine. Opiatine. Tusscapine. NSC 5366. Many other names



(1R,9R)-form

C₂₂H₂₃NO₇ 413.426p*K*_a 5.86 (24°).**(1R,9R)-form**

(-)-β-Narcotine

[3860-46-6]

Synthetic. Mp 181-182° (176°). [α]_D²² -100 (c, 1 in dioxan). [α]_D²⁰ -87.5 (c, 1 in CHCl₃).

▶ LD₅₀ (mus, orl) 840 mg/kg. NP7224500**(1R,9S)-form**

(-)-α-Narcotine. Longatin

[128-62-1]

Alkaloid from *Corydalis cava* (*Corydalis tuberosa*), *Papaver fugax*, *Papaver paeoniflorum*, *Papaver persicum* and *Papaver somniferum* (opium poppy). Byprod. in extraction of morphine from opium. One of the first alkaloids to be isol. (Papaveraceae).

Reference material used in elemental microanalysis. Antitussive agent, of similar potency to and poss. with some advantages over codeine. Mod. effective smooth muscle relaxant. Stout needles (EtOH). Mp 176°. [α]_D -200 (CHCl₃). Log P 2.74 (uncertain value) (calc). Polymorphic. This is prob. the only nat. occurring stereoisomer but it readily epimerises to (-) β-Narcotine which may be isol. as an artifact. Opt. rotn. highly solv. and concn. dependent. λ_{max} 209 (ε 72500); 291 (ε 4000); 309 (ε 4900) (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 83 mg/kg. LD₅₀ (rat, orl) 1520 mg/kg. Mutagenic props. RD2625000

Hydrochloride: [912-60-7]

▶ LD₅₀ (mus, orl) 853 mg/kg. Mutagenic props. NP7225000

Picrate: Mp 174°.

N-Oxide:

Hygroscopic solid. V. sol. H₂O. [α]_D +135 (CHCl₃).

O⁸-De-Me: Narcotoline

[521-40-4]

C₂₁H₂₁NO₇ 399.399

Alkaloid from straw of *Papaver somniferum* (opium poppy) (Papaveraceae). Rods (MeOH aq.). Mp 202° (189°). [α]_D -189 (CHCl₃).

O⁸-De-Me, Ac: Mp 208-209°.**(1S,9R)-form**

(+) -α-Narcotine

[35933-64-3]

Obt. by resoln. of synthetic (±)-narcotine. Cryst. (EtOH). [α]_D +199.9.

N-Me: (+)-N-Methyl-α-narcotineC₂₃H₂₆NO₇⁺ 428.461

Alkaloid from *Papaver cylindricum* (Papaveraceae).

N-Me, iodide: [51606-51-0]

C₂₃H₂₆INO₇ 555.365Amorph. [α]_D +42 (c, 0.16 in MeOH).**(1R,9RS)-form**

(±) -β-Narcotine. β-Gnoscopine

[35933-63-2]

Synthetic. Cryst. (EtOH). Mp 184-186° (180°).

Hydrochloride: Mp 224-226°. Epimerises readily.

(1R,9SR)-form

(±) -α-Narcotine. α-Gnoscopine

[6035-40-1]

Formed in opium, prob. as an artifact. Cryst. (EtOH). V. spar. sol. EtOH. Mp 230-233° (227-230°).

▶ LD₅₀ (rat, ipr) 750 mg/kg. QN5600000

Hydrochloride:

Cryst. + 3H₂O. Mp 220° Mp 238°.

Picrate: Mp 188° (185°).

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1304B (*nmr*)

Robiquet, M. *et al.*, *Ann. Chim. (Paris)*, 1817, **5**, 275 (*isol*)

Perkin, W.H. *et al.*, *J.C.S.*, 1911, **99**, 775 (*synth, resoln*)

Ohta, M. *et al.*, *Tet. Lett.*, 1963, 859 (*abs config*)

Ohta, M. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1080 (*isol*)

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 1087 (*abs config*)

Snatzke, G. *et al.*, *Tetrahedron*, 1969, **25**, 5059 (*ord*)

Analyst (London), 1972, **97**, 740 (*microanal*)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1977, 374 (*synth*)

Al-Yahya, M.A. *et al.*, *Anal. Profiles Drug Subst.*, 1982, **11**, 407 (*rev. synth, props, anal*)

Blaskó, G. *et al.*, *J. Nat. Prod.*, 1982, **45**, 105 (*occur*)

Shono, T. *et al.*, *J.O.C.*, 1983, **48**, 1621 (*synth*)

Moss, D.S. *et al.*, *Acta Cryst. C*, 1984, **40**, 1960 (*cryst struct*)

Johansson, M. *et al.*, *J. Chromatogr.*, 1988, **459**, 301 (*hplc*)

Sariyar, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1302 (*N-Methyl-α-narcotine*)

Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1990, **29**, 3331 (*pmr, cmr, conformn*)

Porter, R. *et al.*, *Mutagenesis*, 1992, **7**, 205 *Martindale, The Extra Pharmacopoeia, 30th edn.*, Pharmaceutical Press, 1993, 750

Tiveron, C. *et al.*, *Mutagenesis*, 1993, **8**, 311


Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms, 7th edn.*, Akademie-Verlag, 1994, 8884 (*synonyms*)

Baradarani, M.M. *et al.*, *Tet. Lett.*, 1999, **40**, 7403-7406 (*epimerisation*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, NBP275; NOA000; NOA500

Narcotinediol

[2725-94-2]

C₂₂H₂₇NO₇ 417.458

Protoberberine-type numbering shown. Minor alkaloid from the capsules of *Papaver pseudo-orientale* (Papaveraceae). Amorph. $[\alpha]_D^{25}$ -14.1 (c, 0.08 in MeOH). $[\alpha]_D$ +53.1 (c, 0.16 in CHCl₃). λ_{\max} 236 (log ϵ 4.02); 281 (log ϵ 3.52) (MeOH).

13-Ac: Papaveroxinoline

[106982-93-8]

C₂₄H₂₉NO₈ 459.495

Minor alkaloid from capsules of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (MeOH). Mp 177-180° Mp 181-184°. $[\alpha]_D$ -7.3 (c, 1.04 in MeOH) (natural). $[\alpha]_D^{18}$ +15.8 (c, 1 in MeOH) (synthetic). Synthetic material is dextrorotatory in contrast to the nat. prod. for which a negative specific rotn. has been recorded. Yet the identity of these 2 compds. seems to be beyond any doubt. λ_{\max} 234 (log ϵ 4.18); 281 (log ϵ 3.56) (MeOH).

8-Aldehyde, 13-Ac: Papaveroxine

[106982-92-7]

C₂₄H₂₇NO₈ 457.479

Main alkaloid from aerial parts of *Papaver fugax* (Papaveraceae). Cryst. (Et₂O). Mp 130-131°. $[\alpha]_D^{25}$ -35 (c, 0.13 in MeOH). λ_{\max} 238 (log ϵ 3.95); 282 (log ϵ 3.54) (MeOH).

8-Carboxylic acid, 13-Ac: Papaveroxidine

[116988-08-0]

C₂₄H₂₇NO₉ 473.479

Alkaloid from the stems, leaves and capsules of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (Et₂O). Mp 164°. $[\alpha]_D$ -139.5 (c, 0.13 in MeOH). λ_{\max} 237 (sh) (log ϵ 4.09); 283 (log ϵ 4.09) (MeOH).

13-Deoxy: Macrantaline

[62818-76-2]

C₂₂H₂₇NO₆ 401.458

A major alkaloid from the aerial parts of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (MeOH). Mp 140-141° Mp 144.5-145.5°. $[\alpha]_D$ +13.3 (c, 0.16 in MeOH). $[\alpha]_D^{18}$ +18.4 (c, 1 in MeOH). λ_{\max} 238 (sh) (log ϵ 4.04); 385 (log ϵ 3.64) (MeOH).

13-Deoxy, 8-Ac:Cryst. (Et₂O). Mp 105°.**13-Deoxy, O¹-de-Me: Narcotolinol**

[106982-94-9]

C₂₁H₂₅NO₆ 387.432

Minor alkaloid from capsules of *Papaver pseudo-orientale* (Papavera-

N-45

ceae). Cryst. (Et₂O). Mp 190-191°. $[\alpha]_D^{25}$ -13 (c, 0.1 in MeOH). λ_{\max} 232 (log ϵ 4.15); 283 (log ϵ 3.56) (MeOH).

13-Deoxy, 8-aldehyde: Macrantaldehyde

[106982-91-6]

C₂₂H₂₅NO₆ 399.443

Alkaloid from capsules of *Papaver pseudo-orientale* (Papaveraceae). Amorph. $[\alpha]_D^{25}$ +4.8 (c, 0.4 in MeOH). λ_{\max} 241 (log ϵ 3.98); 283 (log ϵ 3.59) (MeOH).

13-Deoxy, 8-carboxylic acid: Macrantoridine

[65560-20-5]

C₂₂H₂₅NO₇ 415.442

Minor alkaloid from the aerial parts of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (CHCl₃/heptane). Mp 114°.

Sariyari, G. et al., *Phytochemistry*, 1977, **16**,2009-2013; 1986, **25**, 2403-2406


(*Macrantaline, Narcotinediol, Papaveroxinoline, Papaveroxine, Narcotolinol*)

Sariyari, G. et al., *J. Nat. Prod.*, 1988, **51**, 802-803 (*Papaveroxidine*)

Rozwadowska, M.D. et al., *Annalen*, 1991, 1357-1359 (*Macrantaline, Papaveroxinoline, synth, cmr*)

Narcotinehemiacetal

N-46

C₂₂H₂₅NO₇ 415.442


Alkaloid from the aerial parts of *Papaver fugax* and the capsules of *Papaver pseudo-orientale* (Papaveraceae). Cryst. (MeOH). Mp 155-158°. $[\alpha]_D^{25}$ -220 (c, 0.092 in MeOH). $[\alpha]_D^{25}$ -286 (c, 0.2 in CHCl₃).

Sariyari, G. et al., *Phytochemistry*, 1986, **25**, 2403 (*isol, pmr, ms, struct*)

Nareline

N-47

Methyl 4,5-epoxy-5-hydroxy-6,21-cyclo-4,5-secoakuammilan-17-oate, 9CI
[63950-46-9]



Absolute Configuration

C₂₀H₂₀N₂O₄ 352.389

Alkaloid from the leaves of *Alstonia scholaris*. Prisms (CHCl₃/MeOH). Mp 275° dec. $[\alpha]_D$ -71 (c, 0.12 in CHCl₃/MeOH 1:1).

O-Ac:

Cryst. (MeOH). Mp 186°.

Me ether: O-Methylnareline

[63964-36-3]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from leaves of *Alstonia scholaris*. Cryst. (MeOH). Mp 223°.

Et ether: O-Ethylnareline

[194091-57-1]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from leaves of *Alstonia scholaris*. Needles (CHCl₃/EtOAc). Mp 225-227°. $[\alpha]_D$ -72 (c, 0.066 in CHCl₃). λ_{\max} 216 (log ϵ 3.93); 256 (log ϵ 3.45) (EtOH).

10,11-Dimethoxy: 10,11-Dimethoxynareline

[683239-97-6]

C₂₂H₂₄N₂O₆ 412.441

Alkaloid from the leaves of *Alstonia macrophylla*. Light yellowish oil. $[\alpha]_D$ -56 (c, 1.6 in CHCl₃). λ_{\max} 232 (log ϵ 4.38); 294 (log ϵ 3.81) (EtOH).

10,12-Dimethoxy: 10,12-Dimethoxynareline

[162289-01-2]

C₂₂H₂₄N₂O₆ 412.441

Alkaloid from twigs and leaves of *Tabernaemontana glandulosa*. Cryst. (CHCl₃/MeOH, 1:1). Mp 224-228°. $[\alpha]_D$ -89 (c, 0.9 in CHCl₃).

5-Epimer, Et ether: 5-Epinareline ethyl ether

[194091-58-2]

C₂₂H₂₄N₂O₄ 380.443

From leaves of *Alstonia scholaris*. Light yellow oil. $[\alpha]_D$ -57 (c, 0.037 in CHCl₃). λ_{\max} 216 (log ϵ 4.07); 256 (log ϵ 3.58) (EtOH).

Morita, Y. et al., *Helv. Chim. Acta*, 1977, **60**, 1419-1432 (*isol, ord, uv, ir, pmr, cmr, ms, cryst struct*)

Achenbach, H. et al., *Phytochemistry*, 1994, **37**, 1737-1743 (*10,12-Dimethoxynareline*)

Kam, T.-S. et al., *Phytochemistry*, 1997, **45**, 1303-1305 (*Et ethers*)


Kam, T.-S. et al., *J. Nat. Prod.*, 2004, **67**, 547-552 (*10,11-Dimethoxynareline*)

Nargenicin B₁

N-48

CP 51467. Antibiotic CP 51467

[75923-01-2]

R¹ = R² = OMe, R³ = OHC₂₉H₃₉NO₁₀ 561.628

Cyclic polyketide related to Nargenicin A₁ in N-276. Prod. by *Nocardia argentinensis* ATCC31306. Active against *Staphylococcus*. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{25}$ +80.9 (c, 1 in MeOH). λ_{\max} 264 (MeOH).

Tone, J. et al., *Intersci. Conf. Antimicrob. Agents Chemoth.*, 20th, 1980, Abstr. 62
U.S. Pat., 1980, 4 224 314; *CA*, **94**, 28881

Nargenicin B₂ N-49

CP 52726. Antibiotic CP 52726
[75923-02-3]

As Nargenicin B₁, N-48 with
R¹ = OMe, R² = OH, R³ = H

C₂₈H₃₇NO₉ 531.602

Cyclic polyketide. Prod. by *Nocardia argentinensis* ATCC31306. Active against *Staphylococcus*. Off-white amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 116-124°. λ_{max} 265 (MeOH) (Berdy).

Tone, J. et al., *Intersci. Conf. Antimicrob. Agents Chemoth.*, 20th, 1980, Abstr. 62
U.S. Pat., 1980, 4 224 314; *CA*, **94**, 28881

Nargenicin B₃ N-50

CP 52748. Antibiotic CP 52748
[75923-03-4]

As Nargenicin B₁, N-48 with
R¹ = H, R² = OMe, R³ = OH


C₂₈H₃₇NO₉ 531.602

Prod. by *Nocardia argentinensis* ATCC31306. Active against *Staphylococcus*. Off-white amorph. solid. λ_{max} 235 (MeOH) (Berdy).

Tone, J. et al., *Intersci. Conf. Antimicrob. Agents Chemoth.*, 20th, 980, Abstr. 62
U.S. Pat., 1980, 4 224 314; *CA*, **94**, 28881

Narlumidine N-51

6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]carbonyl]furo[3,4-e]-1,3-benzodioxol-8(6H)-one, 9CI. Paparine
[73710-86-8]



C₂₁H₁₉NO₇ 397.384

Struct. of Paparine given as the enol form of Narlumidine - claimed as a new alkaloid. Isol. from same plant. Alkaloid from *Fumaria indica* (Papaveraceae). Mp 248-250°.

8-Alcohol: **Narlumicine**

[73710-85-7]

C₂₁H₂₁NO₇ 399.399

Alkaloid from stems of *Fumaria indica* (Papaveraceae). Also obt. by NaBH₄ redn. of Narlumidine. Granules (MeOH). Mp 163-165°.

Seth, K.K. et al., *Chem. Ind. (London)*, 1979, 744 (uv, ir, ms, pmr, struct)

Tripathi, V.K. et al., *Phytochemistry*, 1992, **31**, 2188 (Narlumicine)

Atta-ur-Rahman, et al., *Phytochemistry*, 1995, **40**, 593 (Paparine)

Chrzanoska, M. et al., *Heterocycles*, 1998, **47**, 501-504 (Narlumicine, synth)

Tykariska, E. et al., *Pol. J. Chem. (Rocz. Chem.)*, 1998, **72**, 433-438 (cryst struct, Narlumicine)

Narzettine N-52

C₂₀H₂₃NO₆ 373.405

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Narcissus* sp. (Amaryllidaceae). Cryst. (MeOH or EtOH). Mp 224-226°. [α]_D²⁵ -90 (c, 0.2 in CHCl₃).


Methiodide:

Needles (H₂O). Mp 256-258°.

Boit, H.-G. et al., *Chem. Ber.*, 1956, **89**, 2462-2465 (isol)

Natalinine N-53

[101219-60-7]



C₂₅H₂₃NO₅ 417.46

Alkaloid from *Berberis empetrifolia* (Berberidaceae).

1''-Alcohol: **Natalamine**

[123715-10-6]

C₂₅H₂₅NO₅ 419.476

Alkaloid from above-ground parts of *Berberis empetrifolia* (Berberidaceae). [α]_D -5 (c, 0.15 in MeOH).

Fajardo, V. et al., *Rev. Latinoam. Quim.*, 1985, **16**, 59; *CA*, **104**, 145471e (Natalinine)

Hussain, S.F. et al., *J. Nat. Prod.*, 1989, **52**, 644 (Natalamine)

Natrine N-54

C₂₃H₃₈NO 344.559


Struct. unknown. Alkaloid from the branches of *Solanum crispum* (Solanaceae). Cryst. (Me₂CO). Mp 172-173°. Opt. inactive.

Hydrochloride: Mp 288-289°.

Methiodide:

Platelets (MeOH). Mp 272-274°.


Bianchi, E. et al., *Ann. Chim. (Rome)*, 1953, **43**, 208-212; *CA*, **48**, 12142f (isol, uv)

Naucleactonine A N-55

C₁₉H₁₄N₂O₃ 318.331

Alkaloid from *Nauclea officinalis*. Yellowish powder. λ_{max} 255 (log ε 4.35); 313 (log ε 4.34); 348 (log ε 4.26); 416 (log ε 4.26) (MeOH).

Xuan, W.-D. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 719-722 (isol, pmr, cmr, ms)

Naucleactonine B N-56


C₂₀H₁₈N₂O₄ 350.373

Alkaloid from *Nauclea officinalis*. Yellowish powder. [α]_D²⁰ -1.9 (c, 0.52 in MeOH). λ_{max} 217 (log ε 4.03); 309 (log ε 4.1); 423 (log ε 4) (MeOH).

Xuan, W.-D. et al., *J. Asian Nat. Prod. Res.*, 2006, **8**, 719-722 (isol, pmr, cmr, ms)

Nauclealine A N-57

[359795-07-6]



C₂₀H₁₄N₂O₃ 330.342

Alkaloid from the bark of *Nauclea orientalis*. Amorph. yellow solid (MeOH). Mp 267-268°. [α]_D²² -11.4. λ_{max} 206 (log ε 3.28); 288 (log ε 2.67); 424 (log ε 3.01) (MeOH).

12-β-D-Glucopyranosyloxy, 18,19,20,21-tetrahydro, 19,20-didehydro (E-): **Naucleofficine B**

[1023699-68-4]

C₂₆H₂₆N₂O₉ 510.499


Alkaloid from the stems of *Nauclea officinalis*. Orange-yellow powder. Mp 178-179°. [α]_D²⁰ -11.4 (c, 0.21 in MeOH). λ_{max} 225 (log ε 4.53); 296 (log ε 3.57) (MeOH).

Zhang, Z. et al., *J. Nat. Prod.*, 2001, **64**, 1001-1005 (isol, pmr, cmr)

Sun, J. et al., *Phytochemistry*, 2008, **69**, 1405-1410 (Naucleofficine B)

Nauclealine B N-58

[359795-08-7]



C₁₇H₁₆N₂O₃ 296.325


Truncated indole alkaloid, original skeleton uncertain. Alkaloid from the bark of *Nauclea orientalis*. Amorph. yellow solid (MeOH). Mp 222-223°. $[\alpha]_D^{22}$ -11.4 (c, 0.07 in MeOH). λ_{\max} 214 (log ϵ 4.98); 290 (log ϵ 3.19); 374 (log ϵ 3.62); 394 (log ϵ 3.61) (MeOH).

Zhang, Z. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1001-1005

Naucleamide A

N-59

[623531-11-3]



$C_{20}H_{24}N_2O_3$ 340.421

Alkaloid from the bark and wood of *Nauclea latifolia*. Amorph. solid. $[\alpha]_D^{23}$ -113 (c, 0.33 in MeOH). λ_{\max} 225 (log ϵ 4.2); 282 (log ϵ 3.6) (MeOH).

3,14,15,16-Tetrahydro: Naucleamide D
[623531-14-6]

$C_{20}H_{20}N_2O_3$ 336.39

Alkaloid from the bark and wood of *Nauclea latifolia*. Amorph. solid. λ_{\max} 215 (log ϵ 4.18); 372 (log ϵ 4.08); 389 (log ϵ 4) (MeOH).

3,14,15,16-Tetrahydro, 21-O- β -D-glucopyranoside: Naucleofficine A

$C_{26}H_{30}N_2O_8$ 498.532

Alkaloid from the stems of *Nauclea officinalis*. Orange-yellow powder. Mp 185-187°. $[\alpha]_D^{25}$ -12.3 (c, 0.24 in MeOH). λ_{\max} 220 (log ϵ 4.45); 282 (log ϵ 3.51) (MeOH).

16-Epimer: Naucleamide B

[623531-12-4]

$C_{20}H_{24}N_2O_3$ 340.421

Alkaloid from the bark and wood of *Nauclea latifolia*. Amorph. solid. $[\alpha]_D^{23}$ -42 (c, 0.27 in MeOH). λ_{\max} 226 (log ϵ 4.26); 282 (log ϵ 3.6) (MeOH).


Shigemori, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 58-61 (*Naucleamides*)

Sun, J. *et al.*, *Phytochemistry*, 2008, **69**, 58-61 (*Naucleofficine A*)

Naucleamide C

N-60

[623531-13-5]



Relative Configuration

$C_{20}H_{20}N_2O_3$ 336.39

Enolised β -diketone. A heteroyohimbinoïd. Alkaloid from the bark and

wood of *Nauclea latifolia*. Amorph. solid. $[\alpha]_D^{23}$ -47 (c, 0.2 in MeOH). λ_{\max} 224 (log ϵ 4.2); 281 (log ϵ 3.53) (MeOH).

16 β ,17 β -Dihydro: Naucleofficine D

$C_{20}H_{22}N_2O_3$ 338.405

Alkaloid from the stems of *Nauclea officinalis*. Yellowish cryst. Mp 253-255°. $[\alpha]_D^{25}$ -22.4 (c, 0.13 in MeOH). λ_{\max} 221 (log ϵ 4.42); 277 (log ϵ 3.48) (MeOH).


Shigemori, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 58-61 (*Naucleamide C*)

Sun, J. *et al.*, *Phytochemistry*, 2008, **69**, 1405-1410 (*Naucleofficine D*)

Naucleamide E

N-61

[623531-15-7]




$C_{20}H_{22}N_2O_3$ 338.405

A secoyohimbinoïd. Alkaloid from the bark and wood of *Nauclea latifolia*. Amorph. solid. $[\alpha]_D^{24}$ -40 (c, 0.13 in MeOH). λ_{\max} 225 (log ϵ 4.26); 275 (log ϵ 3.6) (MeOH).

Shigemori, H. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 58-61 (*isol, pmr, cmr*)

Naucleorine

N-62



$C_{27}H_{30}N_2O_9$ 526.542

Alkaloid from the stems of *Nauclea orientalis*. Amorph. yellow powder. $[\alpha]_D^{20}$ -54.7 (c, 0.15 in MeOH). λ_{\max} 208 (log ϵ 4.15); 280 (log ϵ 2.17); 369 (log ϵ 3.65); 387 (log ϵ 3.47) (MeOH).

19-Epimer: 19-Epinaucleorine. Epi-methoxynaucleorine

$C_{27}H_{30}N_2O_9$ 526.542


Alkaloid from the stems of *Nauclea orientalis*. Amorph. yellow powder. $[\alpha]_D^{20}$ +52.5 (c, 0.18 in MeOH). λ_{\max} 215 (log ϵ 4.03); 285 (log ϵ 2.01); 367 (log ϵ 3.53); 386 (log ϵ 3.43) (MeOH).

He, Z.-D. *et al.*, *Chem. Biodiversity*, 2005, **2**, 1378-1386 (*isol, pmr, cmr, ms*)

Nauclechine

N-63

[38940-73-7]



Probable Structure Relative Configuration

$C_{21}H_{21}N_3O_3$ 363.415

Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Cryst. (EtOH). Mp 108-114°. λ_{\max} 224 (ϵ 40700); 272 (ϵ 10050); 288 (sh) (MeOH).

Deoxy: 16-Carbomethoxy-18,19-dihydro-naufoline

[149180-46-1]

$C_{21}H_{21}N_3O_2$ 347.416

Alkaloid from leaves of *Nauclea diderrichii* (Rubiaceae).

19-Deoxy, 18,19-didehydro: 16-Carbomethoxynaufoline

[149155-05-5]

$C_{21}H_{19}N_3O_2$ 345.4

Alkaloid from leaves of *Nauclea diderrichii* (Rubiaceae).

De(methoxycarbonyl): Decarbomethoxy-nauclechine

[59785-76-1]

$C_{19}H_{19}N_3O$ 305.379

Alkaloid from the root bark of *Nauclea latifolia* (Rubiaceae). Cryst. (EtOAc). Mp 222°.

De(methoxycarbonyl), 19-deoxy, 18,19-didehydro: Naufoline

[59785-75-0]

$C_{19}H_{17}N_3$ 287.363

Alkaloid from the root bark of *Nauclea latifolia* (Rubiaceae). Cryst. (MeOH). Mp 252°. λ_{\max} 225 (log ϵ 4.48); 285 (log ϵ 4); 291 (log ϵ 4.03) (no solvent reported).

Murray, D.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 1486-1495 (*Nauclechine*)

Hotellier, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1976, **282**, 595-597 (*Naufoline, Decarbomethoxynauclechine*)


Shariff, A. *et al.*, *Tet. Lett.*, 1982, **23**, 4895 (*Decarbomethoxynauclechine, synth*)

Richard, B. *et al.*, *Bull. Soc. R. Sci. Liege*, 1992, **61**, 423-428; *CA*, **119**, 91170v (*16-Carbomethoxynaufoline, 16-Carbomethoxy-18,19-dihydro-naufoline*)

Naucleosidine

N-64

[121880-13-5]




$C_{25}H_{26}N_2O_9$ 498.488

Alkaloid from the stems of *Nauclea officinalis* (Rubiaceae). Cryst. Mp 200-202°.

[121880-12-4]

Lin, M. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 32-36;
CA, **111**, 74776q (*isol, struct*)

Nauclederine N-65
[38940-72-6]



$C_{19}H_{19}N_3O_2$ 321.378
Indole alkaloid of uncertain biogenetic origin.


(ξ)-form

Alkaloid from *Nauclea diderrichii* (Rubiaceae). Mp 102-124° Mp 172-183° (double Mp, partial remelting). $[\alpha]_D^{26}$ -13 (c, 0.8 in $CHCl_3$) (0.0).

Murray, D.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 1486 (*uv, ir, ms, pmr, struct*)

MacLean, S. *et al.*, *Can. J. Chem.*, 1976, **54**, 1262 (*synth*)

Naucleidine N-66
[26238-84-6]




$C_{18}H_{15}N_3O_2$ 305.335
Indole alkaloid of uncertain biogenetic origin. Alkaloid from *Nauclea diderrichii* (Rubiaceae). Pale-yellow needles (MeOH). Mp 84-90° dec.

McLean, S. *et al.*, *Can. J. Chem.*, 1970, **48**, 867 (*ir, ms, uv, pmr*)

Murray, D.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 1486 (*ms, pmr, synth*)

Naucleficine N-67
5,7,8,13-Tetrahydro-5-oxobenz[*g*]indolo[2,3-*a*]quinolizine-1-carboxaldehyde, 9CI
[96400-54-3]



$C_{20}H_{14}N_2O_2$ 314.343
Alkaloid from the stems of *Nauclea officinalis* (Rubiaceae). Orange-red cryst. (CH_2Cl_2 /MeOH). Mp 290-291°.

19-Hydroxy: *Nauclequinine*


[192192-96-4]
 $C_{20}H_{14}N_2O_3$ 330.342
Alkaloid from the roots of *Nauclea pobequinii* (Rubiaceae). Orange-red cryst. (CH_2Cl_2 /MeOH). Mp 291-292°. λ_{max} 260 (log ϵ 4.25); 290 (log ϵ 3.92); 366 (log ϵ 4.39); 375 (log ϵ 4.39); 408 (log ϵ 4.18) (EtOH).

Lin, M. *et al.*, *Planta Med.*, 1984, **50**, 459-461 (*isol, uv, ir, pmr, ms, struct*)

Naito, T. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 2216-2218 (*synth*)

Anam, E.M. *et al.*, *Indian J. Chem., Sect. B*, 1997, **36**, 54-56 (*Nauclequinine*)

Nauclefidine N-68
[96400-52-1]




$C_{16}H_{12}N_2O_2$ 264.283
Struct. revised in 1994. Alkaloid from the stems of *Nauclea officinalis* (Rubiaceae). Prob. derived biogenetically from Vincosamide, V-126. Orange-yellow cryst. Mp 307-309°. λ_{max} 439 (ϵ 4.44) (no solvent reported).

Lin, M. *et al.*, *Planta Med.*, 1984, **50**, 459-461 (*isol, uv, ir, pmr, ms*)

Takayama, H. *et al.*, *Tet. Lett.*, 1994, **35**, 8813-8816 (*synth, cmr, struct*)

Manna, R.K. *et al.*, *Synth. Commun.*, 1998, **28**, 9-16 (*synth, ir, pmr, ms*)


Nauclefiline N-69
[102358-19-0]



$C_{20}H_{20}N_2O_2$ 320.39
Alkaloid from *Nauclea officinalis* (Rubiaceae). Cryst. Mp 315-317°.

Lin, M. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 902-905 (*isol, struct*)

Nauclefine N-70
8,13-Dihydroindolo[2',3':3,4]pyrido[1,2-b][2,7]naphthyridin-5(7H)-one, 9CI. Parvine
[57103-51-2]



$C_{18}H_{13}N_3O$ 287.32
Alkaloid from the root bark of *Nauclea latifolia*, the bark of *Nauclea parva* and from *Nauclea orientalis* (Rubiaceae). Shows antileukaemic activity. Orange-coloured solid (MeOH aq.). Mp 292-294°. λ_{max} 203 (ϵ 31200); 250 (ϵ 10000); 288 (ϵ 78000); 364 (ϵ 78000) (EtOH) (Berdy).

Hotellier, F. *et al.*, *Phytochemistry*, 1975, **14**, 1407 (*isol, uv, ir, pmr, ms, struct*)

Sainsbury, M. *et al.*, *Phytochemistry*, 1975, **14**, 2691 (*isol, uv, ir, pmr, ms, struct, synth*)


Repke, D.B. *et al.*, *Tetrahedron*, 1989, **45**, 2541 (*synth*)

Laronze, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1992, **129**, 303 (*synth*)

Vohra, R. *et al.*, *Heterocycles*, 1994, **39**, 445 (*synth*)

Lavilla, R. *et al.*, *Eur. J. Org. Chem.*, 1999, 373-378 (*synth*)


Nauclefoline† N-71
[81149-29-3]



$C_{21}H_{21}N_3O_3$ 363.415
Alkaloid from the leaves of *Nauclea latifolia* (Rubiaceae). Mp 164-167°. $[\alpha]_D^{20}$ +468 (c, 0.08 in EtOH).

Hotellier, F. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1981, **293**, 577 (*isol*)

Nauclefoline† N-72
[96400-51-0]




$C_{19}H_{16}N_2O_3$ 320.347
Struct. poss. erroneous. Resembles heteroyohimbinoïd alkaloids such as Cadamine, C-6 but the E-ring skeleton is biogenetically unexpected and not derivable from Secologanin. Alkaloid from the stems of *Nauclea officinalis* (Rubiaceae). Pale-yellow cryst. (Me₂CO). Mp 270-272°. λ_{max} 244 (log ϵ 4.26); 295 (log ϵ 4.08); 306 (log ϵ 4.14); 348 (log ϵ 4.33); 366 (log ϵ 4.26) (EtOH).


Lin, M. *et al.*, *Planta Med.*, 1984, **50**, 459-461 (*isol, uv, ir, pmr, ms, struct*)

Nauclefolinine

[774603-00-8]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from the roots of *Nauclea latifolia*. Mp 115-116°. [α]_D²⁵ +3 (c, 0.6 in CH₂Cl₂).Ngnokam, D. *et al.*, *Bull. Chem. Soc. Ethiop.*, 2003, **17**, 173-176; *CA*, **141**, 346483**Naucleidinal**


[77513-45-2]

C₂₀H₂₀N₂O₃ 336.39Alkaloid from the root bark of *Nauclea latifolia* and the stems of *Nauclea officinalis* (Rubiaceae). Needles (CHCl₃). Mp 203° (192-194°). [α]_D²⁰ +124 (c, 1.06 in EtOH). λ_{\max} 226 (log ϵ 4.47); 282 (sh) (log ϵ 3.85); 290 (log ϵ 3.8) (EtOH).**17-Epimer: 17-Epinaucleidinal**

[77513-46-3]

C₂₀H₂₀N₂O₃ 336.39Alkaloid from *Nauclea latifolia* (Rubiaceae). Noncryst. Named 19-Epinaucleidinal in the paper.Hotellier, F. *et al.*, *Phytochemistry*, 1980, **19**, 1884-1885 (*isol, uv, ir, pmr, ms, struct, epimer*)Lin, M. *et al.*, *Planta Med.*, 1984, **50**, 459-461 (*isol, uv, ir, pmr, ms*)Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 2192-2194 (*synth, cmr, cd*)**Naucleofficine C**


[1023699-70-8]

C₂₅H₂₈N₂O₈ 484.505Alkaloid from the stems of *Nauclea officinalis*. Orange-yellow powder. Mp 175-177°. [α]_D²⁰ -13.1 (c, 0.19 in MeOH). λ_{\max} 222 (log ϵ 4.49); 285 (log ϵ 3.5) (MeOH).Sun, J. *et al.*, *Phytochemistry*, 2008, **69**, 1405-1410 (*isol, pmr, cmr*)

N-73

Naucleonidine

Alkaloid ND 363C

C₂₁H₂₁N₃O₃ 363.415Indole alkaloid of unknown biogenesis. prob. secologanin-derived but details not known. β -Carboline numbering shown. Both natural and synthetic alkaloids were a mixt. of the two epimers (α - and β -Naucleonidines).**(1R*,2'R*)-form** β -Naucleonidine

[55221-44-8]

Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Noncryst. Not separately characterised, obt. only as a mixt. with α -Naucleonidine.**De(methoxycarbonyl): β -Naucleonine**

[55252-84-1]

C₁₉H₁₉N₃O 305.379Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Obt. only as a mixt., both naturally and synthetically with α -Naucleonine above.**(1R,2'S)-form** α -Naucleonidine

[37304-94-2]

Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Not separately characterised.**De(methoxycarbonyl): α -Naucleonine.**

Alkaloid ND 305B


[37303-54-1]

C₁₉H₁₉N₃O 305.379Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Syrup. Both natural and synthetic alkaloids were a mixt. of α - and β -Naucleonines.Murray, D.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 1486 (*uv, ir, pmr, ms*)MacLean, S. *et al.*, *Can. J. Chem.*, 1976, **54**, 1262 (*struct, synth*)

N-75

Nauclexine


[38940-74-8]

C₁₈H₁₇N₃O₂ 307.351Alkaloid from the bark of *Nauclea diderrichii* (Rubiaceae). Needles (CH₂Cl₂/MeOH). Mp 229-232°.Murray, D.G. *et al.*, *Can. J. Chem.*, 1972, **50**, 1486-1495 (*uv, ir, pmr, ms, struct*)


N-76

Naulafine


[70503-66-1]

C₂₀H₁₃N₃O 311.342Indole alkaloid, prob. strictosidine-derived. Alkaloid from leaves of *Nauclea latifolia* (Rubiaceae). Red-orange cryst. (CH₂Cl₂). Mp 300°.Hotellier, F. *et al.*, *Planta Med.*, 1979, **35**, 242 (*isol, uv, ir, pmr, ms, struct*)Repke, D.B. *et al.*, *Tetrahedron*, 1989, **45**, 2541 (*synth*)Lavilla, R. *et al.*, *Eur. J. Org. Chem.*, 1999, 373-378 (*synth*)**Navirine**

[773850-94-5]

C₃₀H₄₀N₂O₂ 460.658Isol. from *Aconitum naviculare*. Needles. Mp 175-176°. [α]_D¹⁹ +22 (c, 0.6 in CHCl₃).Gao, L. *et al.*, *J. Chem. Res., Synop.*, 2004, 307-308 (*isol, pmr, cmr, ms*)

N-79

Nazlinine2,3,4,9-Tetrahydro-1H-pyridof[3,4-b]indole-1-butanamide, 9Cl. 1-(4-Aminobutyl)tetrahydro- β -carboline [136945-81-8]C₁₅H₂₁N₃ 243.351

Struct. revised in 1993.


(\pm)-formAlkaloid from *Nitraria schoberi* (Zygophyllaceae). Possesses serotonergic activity. Amorph. powder. Mp 75-78°. λ_{\max} 271 ; 280 ; 288 (MeOH) (Derep).Corthout, J. *et al.*, *Planta Med.*, 1989, **55**, 605 (*isol, ir, pmr*)Ustunes, L. *et al.*, *J. Nat. Prod.*, 1991, **54**, 959-966 (*isol, uv, ir, pmr, cmr, ms*)Wanner, M.J. *et al.*, *Chem. Comm.*, 1993, 174-175 (*synth, struct*)Mahboobi, S. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1994, **327**, 463-465 (*struct*)

Diker, K. *et al.*, *J. Nat. Prod.*, 1997, **60**, 790-793 (*synth, ir, pmr, cmr*)

Neamphine

N-81

5-Methylimidazo[4,5-e]-1,2-thiazin-4(5H)-one, 9CI
[135384-35-9]



C₆H₅N₃OS 167.191


Metab. of the marine sponge *Neamphius huxleyi*. Cytotoxic. Needles (hexane/CHCl₃).

de Silva, E.D. *et al.*, *Tet. Lett.*, 1991, **32**, 2707 (*isol, cryst struct*)

Nebularine

N-82

9-β-D-Ribofuranosyl-9H-purine, 9CI, 8CI. Purinosine
[550-33-4]



C₁₀H₁₂N₄O₄ 252.229

Nucleoside antibiotic. Isol. from the mushroom *Clitocybe nebularis* (clouded agaric), from *Streptomyces yokosukaensis* and *Microbispora* sp. nov. Shows tuberculostatic and antimetabolic activity. Adenosine deaminase inhibitor. Investigated as an antileukaemic agent in combination therapies. Needles (MeOH). Mp 182-183°. [α]_D²⁵ -46.8 (c, 2.0 in H₂O). λ_{max} 263 (ε 7020) (H₂). λ_{max} 263 (ε 5900) (1M HCl). λ_{max} 263 (ε 7110) (pH 9.2).

► LD₅₀ (rat, scu) 220 mg/kg. UO9100000

2',3',5'-Tri-Ac: [15981-63-2]

[α]_D²⁶ -10.8 (c, 1.5 in MeOH).

Isono, K. *et al.*, *J. Antibiot., Ser. A*, 1960, **13**, 270-272 (*isol*)

Cory, J.G. *et al.*, *Biochemistry*, 1965, **4**, 1729-1732 (*pharmacol*)

Iwamura, H. *et al.*, *J.O.C.*, 1968, **33**, 1796-1799 (*synth, tri-Ac*)

Takeda, T. *et al.*, *Acta Cryst. B*, 1974, **30**, 825-827 (*cryst struct*)

Ohno, M. *et al.*, *J.A.C.S.*, 1974, **96**, 4326-4327 (*tri-Ac*)

Chenon, M.T. *et al.*, *J.A.C.S.*, 1975, **97**, 4627-4636 (*cmr*)

Westhof, E. *et al.*, *Z. Naturforsch., C*, 1975, **30**, 131-140 (*pmr*)

Gupta, P.K. *et al.*, *Indian J. Chem., Sect. B*, 1981, **20**, 534-537 (*synth*)

Zavgorodnii, S.G. *et al.*, *Bioorg. Khim.*, 1984, **10**, 1496-1507 (*conformn*)

Nair, V. *et al.*, *Synthesis*, 1984, 401-402 (*synth, uv, pmr, cmr, ms*)

Cooper, R. *et al.*, *J. Ind. Microbiol.*, 1986, **1**, 275-276 (*isol*)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711-1739 (*rev*)

Worner, K. *et al.*, *Helv. Chim. Acta*, 1999, **82**, 2094-2104 (*synth, pmr*)

Francom, P. *et al.*, *J.O.C.*, 2002, **67**, 6788-6796 (*tri-Ac, synth, pmr, ms*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, RJF000

Neburnamine

N-83

[1359-86-0]


Prob. a dimeric indole alkaloid. Struct. unknown. Alkaloid from *Hunteria eburnea* (Apocynaceae). Cryst. (CH₂Cl₂). Mp 290-292°. [α]_D -199 (MeOH).

Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 2197-2199

Necatorine

N-84

11-Hydroxy-2H-benzo[*c*]pyrano[2,3-*h*]cinnolin-2-one, 9CI
[89915-35-5]



C₁₅H₈N₂O₃ 264.24

Alkaloid from the mushroom *Lactarius necator*. Cryst. Mp 220-225°. *L. necator* is considered edible in some countries e.g. Finland, however necatorine is only partially deactivated under cooking conditions.

► Highly mutagenic. DJ2622000

Suortti, T. *et al.*, *J. Chromatogr.*, 1983, **255**, 529; 1984, **301**, 303-307 (*isol*)


Suortti, T. *et al.*, *Phytochemistry*, 1983, **22**, 2873 (*pmr, ms, struct*)

Suortti, T. *et al.*, *Food Chem. Toxicol.*, 1984, **22**, 579-581 (*stability*)

Necatorone

N-85

5,10-Dihydroxy-6H-pyrido[4,3,2-*kl*]acridin-6-one, 9CI
[92631-70-4]



C₁₅H₈N₂O₃ 264.24

Alkaloidal pigment from the fruit-bodies of the gilled toadstool *Lactarius necator* and *Lactarius atroviridis*. Red needles (MeOH). Sol. DMSO, bases; fairly sol. MeOH, Me₂CO; poorly sol. C₆H₆, hexane. Mp 360°. λ_{max} 248; 270 (sh); 327; 400; 520; 610 (sh) (1M NH₃) (Derep). λ_{max} 212 (sh) (ε 24000); 233 (ε 39810); 265 (sh) (ε 13500); 293 (ε 7586); 310 (sh) (ε 7080); 431 (ε 13490) (MeOH) (Derep). λ_{max} 223 (ε 39810); 293 (ε 7595); 431 (ε 13490) (MeOH) (Berdy).

► Mutagenic agent.

4,4'-Dimer: 4,4'-Binectorone
[126647-31-2]

C₃₀H₁₄N₄O₆ 526.464

Alkaloidal pigment from fruiting bodies of the toadstools *Lactarius necator* and *Lactarius atroviridis*. Dark brown or red cryst. (MeOH). Sol. MeOH, DMSO. Mp 360°. λ_{max} 236; 438 (MeOH) (Berdy).

► Mutagenic agent.

4,4'-Dimer, 10-deoxy: 10-Deoxy-4,4'-binectorone

[126624-08-6]

C₃₀H₁₄N₄O₅ 510.464

Alkaloidal pigment from *Lactarius necator* and *Lactarius atroviridis*. Dark brown cryst. (MeOH). Sol. MeOH, DMSO. Mp 360°. λ_{max} 226; 264; 420 (MeOH) (Berdy).

► Mutagenic agent.

4,4'-Dimer, 10,10'-dideoxy: 10,10'-Dideoxy-4,4'-binectorone

[126647-32-3]

C₃₀H₁₄N₄O₄ 494.465

Major alkaloid from *Lactarius atroviridis*. Blackish-green or orange cryst. (MeOH). Sol. MeOH, DMSO. Mp 360°. λ_{max} 223; 264; 358; 428 (MeOH) (Berdy).

► Mutagenic agent.

Fugmann, B. *et al.*, *Tet. Lett.*, 1984, **25**, 3575 (*isol, uv, ir, pmr, cmr, ms, struct*)

Hilger, C.S. *et al.*, *Tet. Lett.*, 1985, **26**, 5975 (*synth*)

Klamann, J.-D. *et al.*, *Phytochemistry*, 1989, **28**, 3519 (*isol, uv, ir, pmr, cmr, ms, dimers*)

Senecio doria Necine base

N-86

C₇H₁₃NO₂ 143.185


Pyrrrolizidine alkaloid. Struct. unknown. Alkaloid from *Senecio doria* (aerial parts in bloom) (Asteraceae). Sol. H₂O. Mp 179-180°.

Constantinescu, E. *et al.*, *Farmacia (Bucharest)*, 1961, **9**, 131; *CA*, **56**, 14396

Neferine

N-87

12'-O-Methylinsinine
[2292-16-2]



C₃₈H₄₄N₂O₆ 624.775

Alkaloid from the seed embryo of *Nelumbo nucifera* (East Indian lotus) (Nelumbonaceae). Tumour promotor. Oil. [α]_D²⁵ -37.8 (c, 1.43 in CHCl₃). [α]_D²⁷ -100 (c, 0.24 in MeOH).

► Tumour promotor. SM9495000

Me ether, styphnate: Mp 139-140° (133-135°). [α]_D²⁵ -93.3 (c, 0.6 in Me₂CO) (-81.5).

Methodide: Carcinogen promotor. Mp 179-181°.

O⁷-De-Me: *Isolesinine*
[6817-41-0]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from the roots of *Cissampelos*

mucronata an the seed embryo of *Nelumbo nucifera* (East Indian lotus) (Nelumbonaceae). Tumour promotor. Oil. $[\alpha]_D^{22} +49.3$ (Me₂CO). $[\alpha]_D^{25} -43.3$ (CHCl₃).

► Tumour promotor.

O⁷-De-Me; hydrochloride (1:2):

Cryst. + 4H₂O. Mp 185-186°.

O⁷-De-Me; perchlorate (1:2):

Cryst. + 1H₂O. Mp 200-203°. $[\alpha]_D^{22} -70$ (Me₂CO).

O¹²-De-Me; Liensinine

[2586-96-1]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from seed embryo of *Nelumbo nucifera* (East Indian lotus) (Nelumbonaceae). Mp 95-99°. $[\alpha]_D^{31} +15.85$ (c, 0.883 in Me₂CO).

O¹²-De-Me; perchlorate (1:2): Mp 212-214°.

O¹²-De-Me, di-Ac: Mp 124°.

Chao, Y.C. et al., *Sci. Sin. (Engl. Edn.)*, 1962, **11**, 215-219; *CA*, **57**, 7383i (Liensinine)

Hsieh, Y.Y. et al., *Sci. Sin. (Engl. Edn.)*, 1964, **12**, 2018-2019; *CA*, **62**, 9183h (Liensinine, abs config)

Tomita, M. et al., *Tet. Lett.*, 1964, **5**, 2637-2642 (Isoliensinine)

Furukawa, H. et al., *Yakugaku Zasshi*, 1965, **85**, 335-338; 353-354; *CA*, **63**, 4351 (isol, ir, pmr, struct)

Yang, T.-H. et al., *J. Chin. Chem. Soc. (Taipei)*, 1970, **17**, 235-242; *CA*, **74**, 100254g (isol, uv, ir, pmr)

Kametani, T. et al., *J. Het. Chem.*, 1970, **7**, 181-186 (isol, pmr)

Tshibangu, J.N. et al., *Phytochem. Anal.*, 2003, **14**, 13-22 (Isoliensinine, isol, hplc)

Yang, J. et al., *Magn. Reson. Chem.*, 2004, **42**, 994-997 (Neferine, Isoliensinine, pmr, cmr)

Neflexine

N-88

[1359-87-1]

C₁₈H₂₁NO₃ 299.369

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine flexuosa* (Amaryllidaceae). Mp 249-250° dec. $[\alpha]_D +65$ (c, 0.25 in CHCl₃).


Picrate: Mp 195-196° dec.

Boit, H.-G. et al., *Naturwissenschaften*, 1960, **47**, 109 (isol)

Nemorine

N-89

[50906-96-2]



Absolute configuration

C₁₈H₂₇NO₅ 337.415

Alkaloid from *Senecio nemorensis* (several varieties) (Asteraceae). Cryst. (cyclohexane). $[\alpha]_D^{24} -58$ (c, 2.35 in CHCl₃).

N-Oxide: *Oxynemorine* [74209-54-4]

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio nemorensis* var. *subdecurrens* (Asteraceae). Cryst. (EtOAc). Mp 160-163°. $[\alpha]_D^{20} -35$ (c, 1.0 in CHCl₃).

11,14-Diepimer: *Mulgediifoline*

[172018-19-8]

C₁₈H₂₇NO₅ 337.415

Alkaloid from leaves of *Senecio mulgediifolius* (Asteraceae). Cryst. (hexane). Mp 102-104°. $[\alpha]_D -32.5$ (c, 0.28 in CHCl₃).

11,14-Diepimer, 1,2-didehydro: *Retrososenine*. Alkaloid SN-A

[62018-78-4]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio nemorensis* and *Senecio mulgediifolius* (Asteraceae). Mp 127°. $[\alpha]_D^{25} +118$ (c, 0.69 in CHCl₃).

11,14-Diepimer, 1,2-didehydro, N-oxide: *Oxyretrososenine*. Retrososenine N-oxide

[171927-39-2]

C₁₈H₂₅NO₆ 351.399

Alkaloid from leaves of *Senecio mulgediifolius* (Asteraceae). Light brown cryst. (EtOAc/hexane). Mp 128-131°. $[\alpha]_D +31.4$ (c, 0.4 in CHCl₃).

11,14-Diepimer, 1,2-didehydro, 12S-hydroxy: *12S-Hydroxyretrososenine*

C₁₈H₂₅NO₆ 351.399

Alkaloid from *Senecio helodes* and *Senecio roseus* (Asteraceae). Cryst. (hexane/EtOAc). Mp 167-170°. $[\alpha]_D^{25} +88$ (c, 2 in CHCl₃).

11,14-Diepimer, 1,2-didehydro, 13R-hydroxy: *13R-Hydroxyretrososenine*

C₁₈H₂₅NO₆ 351.399

Alkaloid from *Senecio roseus* (Asteraceae). Cryst. (hexane/Me₂CO). Mp 187-190°. $[\alpha]_D^{25} +116.4$ (c, 2 in CHCl₃).

Klásek, A. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 2504; 1980, **45**, 548 (isol, ir, pmr, cmr, ms, struct, oxide)

Nghia, N.T. et al., *Coll. Czech. Chem. Comm.*, 1975, **40**, 2952; 1976, **41**, 2952 (Retrososenine)

Gencheva, E. et al., *CA*, 1979, **90**, 127593j (chromatog)

Klein, L.L. et al., *J.A.C.S.*, 1985, **107**, 2573 (synth)

Dillon, M.P. et al., *Chem. Comm.*, 1995, 1645 (abs config)

De Vivar, A.R. et al., *Tetrahedron*, 1995, **51**, 12521 (*Oxyretrososenine*, *Mulgediifoline*)

Perez-Castorena, A.-L. et al., *J. Nat. Prod.*, 1997, **60**, 1322-1325 (*Hydroxyretrososenines*, *Retrososenine*, abs config)

Nemorine

N-90

[1359-88-2]

C₂₄H₃₉NO₄ 405.576

Struct. unknown. Alkaloid from the tubers of *Aconitum nemorum* (Ranunculaceae). Mp 126-128°. $[\alpha]_D^{17} -4.5$.

Oxalate: Mp 175-176°.

Picrate: Mp 105-106°.


O-Ac: Mp 110-113°.

Abubakirov, N.K. et al., *Zh. Obshch. Khim.*, 1959, **29**, 2454-2456; *J. Gen. Chem. USSR (Engl. Transl.)*, 1959, **29**, 2418

Nemuarine, 9CI

N-91

[38739-62-7]



Absolute configuration

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from leaves and twigs of *Nemuaron vieillardii* (Monimiaceae). Prisms + 1H₂O (MeOH). Mp 222-223° dec. $[\alpha]_D^{20} -42.7$ (CHCl₃).


Me ether: Mp 154-156°. $[\alpha]_D^{20} -43.8$ (CHCl₃).

Bick, I.R.C. et al., *Aust. J. Chem.*, 1973, **26**, 455 (isol, uv, ms, pmr, struct)

Neocrimarine A

N-92

[149301-45-1]



C₄₀H₄₃NO₉ 681.781

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) × Dancy tangerine (*Citrus tangerina*)] (Rutaceae). Yellow cubes (Me₂CO). Mp 225-230°. $[\alpha]_D 0$ (CHCl₃).

O⁶-De-Me: *Neocrimarine B*

[149301-47-3]

C₃₉H₄₁NO₉ 667.754


Alkaloid from Yalaha (Rutaceae). Yellow cubes (Me₂CO). Mp 240-243°. $[\alpha]_D 0$ (Py).

Takemura, Y. et al., *Heterocycles*, 1993, **36**, 675 (isol, uv, ir, pmr, cmr, struct)

Neocrimarine C

N-93

[155519-79-2]



Relative configuration

C₃₃H₂₉NO₉ 583.593

Alkaloid from roots of *Citrus hassaku* (Rutaceae). Yellow cubes. Mp 111-113°. $[\alpha]_D +42.3$ (c, 0.026 in CHCl_3).

10-Deoxy: Neocrimarine H


[217199-06-9]
 $\text{C}_{33}\text{H}_{29}\text{NO}_8$ 567.594

Alkaloid from the roots of *Citrus paradisi* (grapefruit). Yellow oil. $[\alpha]_D +80.6$ (c, 0.07 in CHCl_3). λ_{max} 204; 265; 286; 294 (sh); 318; 343 (sh); 410 (MeOH).

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1757-1759; 1998, **46**, 1518-1521 (*isol, uv, ir, pmr, cmr, ms, struct*)

Neocrimarine D N-94

[155519-80-5]




$\text{C}_{38}\text{H}_{37}\text{NO}_8$ 635.712

Alkaloid from roots of *Citrus hassaku* (Rutaceae). Yellow oil. Opt. inactive.

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1757 (*isol, uv, ir, pmr, cmr, ms, struct*)

Neocrimarine E N-95

[158182-13-9]




$\text{C}_{35}\text{H}_{35}\text{NO}_9$ 613.663

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) and Dancy tangerine (*Citrus tangerina*)] (Rutaceae). Yellow cubes (Me_2CO). Mp 212-215°. $[\alpha]_D -21.6$ (c, 0.3 in CHCl_3).

Takemura, Y. *et al.*, *Heterocycles*, 1994, **38**, 1937 (*isol, uv, ir, pmr, cmr, ms, struct*)

Neocrimarine F N-96

[195057-36-4]



$\text{C}_{29}\text{H}_{25}\text{NO}_9$ 531.518

Alkaloid from the roots of *Citrus paradisi* (grapefruit) x *Citrus tangerina* (tangerine) (Rutaceae). Yellow cubes. Mp 218-220°. $[\alpha]_D +130.9$ (c, 0.2 in MeOH). λ_{max} 220 (sh); 264; 295 (sh); 330; 369 (EtOH).

N-De-Me: Neocrimarine J

[217199-08-1]
 $\text{C}_{28}\text{H}_{23}\text{NO}_9$ 517.491

Alkaloid from the roots of *Citrus paradisi* (grapefruit). Yellow oil. $[\alpha]_D -131.9$ (c, 0.05 in CHCl_3). λ_{max} 203; 218 (sh); 263; 292 (sh); 327; 373 (sh) (MeOH).

6-Me ether, N-de-Me: Neocrimarine I

[217199-07-0]
 $\text{C}_{29}\text{H}_{25}\text{NO}_9$ 531.518

Alkaloid from the roots of *Citrus paradisi* (grapefruit). Yellow oil. $[\alpha]_D -157.3$ (c, 0.27 in CHCl_3). λ_{max} 205; 255 (sh); 266; 292 (sh); 326; 381 (MeOH).

6-Deoxy: Neocrimarine G


[195057-37-5]
 $\text{C}_{29}\text{H}_{25}\text{NO}_8$ 515.518

Alkaloid from the roots of *Citrus paradisi* (grapefruit) (Rutaceae). Yellow oil. $[\alpha]_D +130.2$ (c, 0.03 in CHCl_3). λ_{max} 220 (sh); 262; 280 (sh); 315; 324 (MeOH).

Takemura, Y. *et al.*, *Heterocycles*, 1997, **45**, 1411-1414 (*isol, uv, ir, pmr, cmr, ms*)
 Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1518-1521 (*isol, uv, ir, pmr, cmr, ms*)

Neocrimarine K N-97

[217199-09-2]




$\text{C}_{31}\text{H}_{29}\text{NO}_9$ 559.571

Alkaloid from the roots of *Citrus paradisi* (grapefruit). Yellow oil. Racemic. λ_{max} 204; 222 (sh); 261 (sh); 269; 334; 393 (MeOH).

Takemura, Y. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1518-1521 (*isol, uv, ir, pmr, cmr, ms*)

Neodifoline N-98

[250721-70-1]




$\text{C}_{22}\text{H}_{20}\text{N}_2\text{O}_7$ 424.409

Alkaloid from the heartwood of *Adina cordifolia*. Pale yellow cryst. (CHCl_3). Mp. >350°.

Balazs, B. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 751-753

Neoamphimedine N-99

10-Methyl-8H-benzo[b]pyrido[4,3,2-de][1,8]phenanthroline-8,9(10H)-dione, 9CI
 [221456-55-9]



$\text{C}_{19}\text{H}_{11}\text{N}_3\text{O}_2$ 313.315

Alkaloid from *Xestospongia* cf. *carbonaria* and *Xestospongia* cf. *exigua*. Topoisomerase II inhibitor which catenates DNA. Yellow solid. Mp >300°. λ_{max} 205 (ϵ 26370); 226 (ϵ 23740); 278 (ϵ 16870); 371 (ϵ 10190) (EtOH).

5-Methoxy: 5-Methoxyneoamphimedine


[485817-61-6]
 $\text{C}_{20}\text{H}_{13}\text{N}_3\text{O}_3$ 343.341

Alkaloid from *Xestospongia* cf. *carbonaria* and *Xestospongia* cf. *exigua*. Yellow solid.

De Guzman, F.S. *et al.*, *J.O.C.*, 1999, **64**, 1400-1402 (*Neoamphimedine*)
 Thale, Z. *et al.*, *J.O.C.*, 2002, **67**, 9384-9391 (*5-Methoxyneoamphimedine*)
 LaBarbera, D.V. *et al.*, *J.O.C.*, 2007, **72**, 8501-8505 (*synth*)

Neoamphimedine Y N-100

[486992-27-2]



$\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_4$ 349.345

Alkaloid from *Xestospongia* cf. *carbonaria*. Purple solid. Turns brown on standing.

11,12-Di-Me ether: Neoamphimedine Z
 [486992-28-3]

$\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_4$ 377.399

Alkaloid from *Xestospongia* cf. *carbonaria*. Purple solid. Turns brown on standing.

Thale, Z. *et al.*, *J.O.C.*, 2002, **67**, 9384-9391 (*isol, pmr, cmr*)

Neoberninamycin

N-101

Antibiotic EM 5596. EM 5596

[115537-10-5]


Peptide antibiotic. Struct. not determined. Prod. by *Micrococcus luteus*. Active against gram-positive bacteria. Amorph. solid. Sol. MeOH, butanol; poorly sol. H₂O. λ_{\max} 240 (MeCN). λ_{\max} 230 ; 250 (MeCN) (Berdy).

Biskupiak, J.E. et al., *J. Antibiot.*, 1988, **41**, 684-688 (isol, uv, ir, pmr, props)

Neobudofficide B

N-102

[219998-44-4]

C₃₄H₄₀NO₁₅[⊖] 702.688


Alkaloid from the flowers of *Buddleia officinalis*. Isol. as the chloride, to which registry number refers.

Li, J. et al., *J. Chin. Pharm. Sci.*, 1997, **6**, 178-181

Neocarazostatin A

N-103

1-(1,2-Dihydroxypropyl)-4-hydroxy-3-methoxy-2-methyl-6-prenyl-9H-carbazole [137767-81-8]

C₂₂H₂₇NO₄ 369.46

Prod. by *Streptomyces* sp. strain GP38. Free radical scavenger. Mp 141-143° dec. $[\alpha]_{\text{D}}^{25}$ -36 (c, 0.1 in MeOH). λ_{\max} 230 (ε 43700); 250 (ε 65900); 271 (ε 23800); 292 (ε 26800); 331 (ε 6900); 345 (ε 8400) (MeOH) (Derep).

l'-Deoxy: **Neocarazostatin B**

[137767-82-9]

C₂₂H₂₇NO₃ 353.46

From a *Streptomyces* sp. strain GP38. Free radical scavenger. Mp 55-57° dec. $[\alpha]_{\text{D}}^{25}$ -24 (c, 0.1 in MeOH). Possesses (R)- abs. config. λ_{\max} 230 (ε 43700); 250 (ε 65900); 271 (ε 23800); 292 (ε 26800); 331 (ε 6900); 345 (ε 8400) (MeOH) (Derep).

l'-Me ether: **Neocarazostatin C**

[137714-93-3]

C₂₃H₂₉NO₄ 383.486

From a *Streptomyces* sp. strain GP38. Free radical scavenger. Mp 80-82° dec. $[\alpha]_{\text{D}}^{25}$ -92 (c, 0.1 in MeOH). λ_{\max} 230 (ε 43700); 250 (ε 65900); 271 (ε 23800); 292 (ε 26800); 331 (ε 6900); 345 (ε 8400) (MeOH) (Derep).

Kato, S. et al., *J. Antibiot.*, 1991, **44**, 903-907 (isol, uv, pmr, cmr)


Czerwonka, R. et al., *Chem. Comm.*, 2006, 711-713 (*Neocarazostatin B*, synth, abs config)

Neocaryachine

N-104

5,6,11,12-Tetrahydro-9-methoxy-14-methylbenzo[5,6]cycloocta[1,2-f]-1,3-benzodioxol-5,11-imin-10-ol, 9CI. 1-Hydroxy-2-methoxy-8,9-methylenedioxy-pavine

[131985-46-1]

C₁₉H₁₉NO₄ 325.363

Has been numbered the other way round the symmetrical ring-system, as 7-hydroxy-8-methoxy-2,3-methylenedioxy. Alkaloid from the bark of *Cryptocarya chinensis* (Lauraceae). Amorph. Mp 95-97°. $[\alpha]_{\text{D}}^{25}$ -197 (c, 0.36 in MeOH).

N-Me: **Neocaryachine N-methosalt**. *N*-MethylneocaryachiniumC₂₀H₂₂NO₄[⊕] 340.398

Quaternary alkaloid from callus cultures of *Cryptocarya chinensis*. $[\alpha]_{\text{D}}^{25}$ -40.1 (c, 0.01 in MeOH) (as perchlorate). λ_{\max} 206 (log ε 4.53); 288 (log ε 3.64) (MeOH) (as perchlorate).

Me ether, *N*-Me: **7-O-Methylneocaryachine N-methosalt**C₂₁H₂₄NO₄[⊕] 354.425

Quaternary alkaloid from *Eschscholtzia californica*. Amorph. solid. $[\alpha]_{\text{D}}^{25}$ -170 (c, 0.14 in MeOH).

Lee, S.-S. et al., *J. Nat. Prod.*, 1990, **53**, 1267-1271 (isol, uv, pmr, cmr, ms)

Chang, W.-T. et al., *Phytochemistry*, 1998, **48**, 119-124 (*N*-metho salt)

Gafner, S. et al., *J. Nat. Prod.*, 2006, **69**, 432-435 (*Me* ether *N*-Me)

Neochondocurarine

N-105

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from Curare.

Nitrate: $[\alpha]_{\text{D}}^{20}$ +185 (c, 2 in H₂O).


Chloride: Mp 268° dec. $[\alpha]_{\text{D}}^{20}$ +179 (c, 2 in H₂O).

Bodendorf, K. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1954, **287**, 555-561; *C.A.*, **52**, 10119h (isol)

Neocochliodinol, 9CI

N-106

[79921-14-5]

C₃₂H₃₀N₂O₄ 506.6

Isomeric with Cochliodinol, C-547. Pigment isol. from *Chaetomium amygdalisporum*. Mycotoxin and cytotoxic. Violet cryst. powder (C₆H₆/CH₂Cl₂). Poorly sol. hexane. Mp 285° dec. Related to Asterriquinones. λ_{\max} 290 (ε 30200); 470 (ε 4270) (EtOH) (Derep).

Di-O-Ac:Violet needles (C₆H₆). Mp 214°.*Di*-Me ether:

Violet cryst. powder. Mp 237°.


Sekita, S. et al., *Can. J. Microbiol.*, 1981, **27**, 766-772 (isol, props)

Sekita, S. et al., *Chem. Pharm. Bull.*, 1983, **31**, 2998-3001 (uv, ir, pmr, struct)

Neocoprogen I

N-107

[94141-24-9]

R = —CH=C(CH₃)CH₂CH₂OHC₃₁H₄₇FeN₆O₁₂ 751.592

Isol. from *Curvularia lunata*. Siderophore. Cryst. λ_{\max} 428 (MeOH) (Berdy). λ_{\max} 428 (ε 2730) (H₂O) (Berdy).

Hossain, M.B. et al., *J.A.C.S.*, 1987, **109**, 4948 (isol, struct)

Neocoprogen II N-108


[105250-32-6]
As Neocoprogen I, N-107 with
R = CH₃
C₂₇H₄₁FeN₆O₁₁ 681.501
Isol. from *Curvularia lunata*. Sidero-
phore. Cryst. λ_{max} 422 (MeOH) (Berdy).
λ_{max} 422 (ε 2612) (H₂O) (Berdy).
Hossain, M.B. *et al.*, *J.A.C.S.*, 1987, **109**, 4948
(*isol, struct*)

Neodaphniphylline N-109

Struct. unknown. Alkaloid from the
leaves and bark of *Daphniphyllum*
macropodium (Daphniphyllaceae). Mp
242-244° (as hydrochloride).
Sakabe, N. *et al.*, *Tet. Lett.*, 1966, 963 (*isol*)

Neodichroine N-110

3-(Octahydro-9-hydroxy-2-oxo-2H-quinoliz-
izin-3-yl)-4-(3H)-quinazolinone, 9CI.
Hydrachine A



(+)-form

C₁₇H₁₉N₃O₃ 313.355
Struct. of Hydrachine A revised in 2003.

(+)-form [317832-22-7]

Alkaloid from the leaves of *Dichroa*
febrifuga. [α]_D +198.8 (c. 0.17 in MeOH).
Stereochem. possibly incorrect.


(-)-form [354819-61-7]

Alkaloid from the roots of
Hydrangea chinensis. Semisolid. [α]_D -
25.3 (c. 0.2 in MeOH). λ_{max} 224 (log ε
1.47); 230 (log ε 1.35); 233 (log ε 1.3);
265 (log ε 0.59); 302 (log ε 0.28)
(MeOH).

Deng, Y. *et al.*, *J. Chin. Pharm. Sci.*, 2000, **9**,
116-118 (*isol*)
Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2003, **66**,
1245-1248 (*struct*)

Neodihydrothebaine N-111

[15228-73-6]




C₁₉H₂₃NO₃ 313.396
Alkaloid from the capsules of
Papaver bracteatum (Papaveraceae). Mp
107°.

Theuns, H.G. *et al.*, *Phytochemistry*, 1984, **23**,
1157 (*isol, pmr, ms, struct, synth*)

Neodysiherbaine A N-112

[342410-85-9]




Absolute Configuration

C₁₁H₁₇NO₈ 291.257
Isol. from *Dysidea herbacea*. Neurotoxin.
Pale yellow solid. [α]_D²³ -6.5 (c. 0.75 in
H₂O).

Sakai, R. *et al.*, *Org. Lett.*, 2001, **3**, 1479-1482
(*isol, synth, cd, pmr, cmr*)
Lygo, B. *et al.*, *Tet. Lett.*, 2005, **46**, 6629-6632
(*synth*)
Takahashi, K. *et al.*, *J.O.C.*, 2006, **71**, 4227-
4231 (*synth*)
Shoji, M. *et al.*, *J.O.C.*, 2006, **71**, 5208-5220
(*synth*)

Neoenactin† N-113

2-Amino-N-(3,10-dioxohexadecyl)-N,3-
dihydroxypropanamide, 9CI



Neoenactin A R = CH₂CH₂CH₃
B₁ R = CH(CH₃)CH₂CH₃
B₂ R = CH₂CH(CH₃)₂
NL₁ R = CH₂CH₃
NL₂ R = CH₂CH₂CH₂CH₃

Antibiotic complex. Isol. from *Strepto-*
verticillium olivoreticuli neoenacticus.
Antifungal agent. Sol. MeOH, Et₂O;
poorly sol. H₂O, hexane. Not related to
the nucleoside Neoenactins. Neoenactin
M₁ is identical with Lipoxamycin, L-189.
λ_{max} 208 (E1%/1cm 166) (MeOH) (Ber-
dy). λ_{max} 208 (E1%/1cm 161) (MeOH/
HCl) (Berdy).

Neoenactin A*Enactin IVb*

[88231-85-0]

C₁₉H₃₆N₂O₅ 372.504
From *Streptomyces roseoviridis*. Fairly
sol. MeOH, EtOH, H₂O; poorly sol.
Me₂CO, Et₂O. λ_{max} 211 (ε 5900) (MeOH)
(Berdy). λ_{max} 211 (ε 5900) (MeOH/HCl)
(Berdy). λ_{max} 238 (ε 5900) (MeOH/
NaOH) (Berdy).

Sulfate (2:1): Mp 144-145° dec.*10'-Alcohol*: Neoenactin M₂, *Enactin Vb*
[108334-69-6]

C₁₉H₃₈N₂O₅ 374.52
From *Streptomyces olivoreticuli*
ssp. *neoenactus* and *Streptomyces ro-*
seoviridis.

3'-Alcohol; sulfate (2:1): Mp 166°.**Neoenactin B₁** [88231-86-1]C₂₀H₃₈N₂O₅ 386.531

From *Streptomyces roseoviridis*. Sol.
MeOH, acids; poorly sol. hexane,

H₂O. λ_{max} 211 (ε 5000) (MeOH)
(Berdy). λ_{max} 211 (ε 5000) (MeOH/HCl)
(Berdy). λ_{max} 238 (ε 5800) (MeOH/
NaOH) (Berdy).

Sulfate (2:1): Mp 139.4-141° dec.*15'-Hydroxy: Neactin Ib*

[137252-26-7]

C₂₀H₃₈N₂O₆ 402.53
Isol. from *Streptomyces roseoviridis*.
Sol. MeOH, H₂O. Mp 128-129° dec.
(as sulfate salt). Enactin Ib₁ and
Enactin Ib₂ are diastereoisomers at C-
15. λ_{max} 238 (ε 5550) (MeOH/NaOH)
(Derep). λ_{max} 211 (ε 5500) (MeOH)
(Derep). λ_{max} 213 (ε 4400) (MeOH)
(Berdy).

Neoenactin B₂*Enactin VIb*

[88231-87-2]

C₂₀H₃₈N₂O₅ 386.531
From *Streptomyces roseoviridis*. Sol.
MeOH, acids; poorly sol. hexane, H₂O.
λ_{max} 211 (ε 5400) (MeOH) (Berdy).
λ_{max} 211 (ε 5500) (MeOH/HCl) (Berdy).
λ_{max} 238 (ε 6000) (MeOH/NaOH)
(Berdy).

Sulfate (2:1):

C₂₀H₃₈N₂O₅ 386.531
Mp 141-143° dec.

15'-Hydroxy: Neactin Ia

[130640-30-1]

C₂₀H₃₈N₂O₆ 402.53
Isol. from *Streptomyces roseoviridis*.
Sol. MeOH, H₂O. Mp 134.5° dec. (as
sulfate salt). λ_{max} 238 (ε 5550) (MeOH/
NaOH) (Derep). λ_{max} 211 (ε 5500)
(MeOH) (Derep). λ_{max} 212 (ε 4100)
(MeOH) (Berdy).

Neoenactin NL₁ [117193-00-7]C₁₈H₃₄N₂O₅ 358.477

Biosynthetic prod. Sol. MeOH, acids;
poorly sol. H₂O, hexane. λ_{max} 211 (ε
5200) (MeOH) (Berdy).

Sulfate (2:1):

Needles (MeOH). Mp 145°.

Neoenactin NL₂ [117193-01-8]C₂₀H₃₈N₂O₅ 386.531

Biosynthetic prod. Sol. MeOH, acids;
poorly sol. H₂O, hexane. λ_{max} 238 (ε
5550) (MeOH/NaOH) (Derep). λ_{max} 211
(ε 5500) (MeOH) (Derep).

Sulfate (2:1):

Needles (MeOH). Mp 145°.

[130668-40-5]

Kondo, H. *et al.*, *J. Antibiot.*, 1979, **32**, 13
(*isol*)

Nishio, M. *et al.*, *J. Antibiot.*, 1983, **36**, 1399
(*isol, uv, ir*)

Roy, S.K. *et al.*, *J. Antibiot.*, 1986, **39**, 717;
1987, **40**, 266 (*isol, struct*)

Okada, H. *et al.*, *J. Antibiot.*, 1989, **42**, 276
(NL₁, NL₂)


Yamamoto, K. *et al.*, *J. Antibiot.*, 1990, **43**,
1012 (*isol*)

Yamamoto, K. *et al.*, *Chem. Pharm. Bull.*,
1991, **39**, 1436 (*isol, pmr, cmr, Enactin I*)

Darwish, I.S. *et al.*, *J.O.C.*, 1994, **59**, 451
(*synth*)

Neoengleromycin


N-114

C₄₅H₈₄N₂O₇ 765.168Prod. by fruiting bodies of *Engleromyces goetzii*. Oily solid. [α]_D²⁰ +3.3 (c, 0.3 in CHCl₃).N-De-Et: *Vibratilicin*C₄₃H₈₀N₂O₇ 737.114Prod. by *Cortinarius vibratilis*. Yellow oily solid. [α]_D¹⁹ +8.6 (c, 1.3 in CHCl₃).Jikai, L. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1439-1442 (*Neoengleromycin*)Wang, F. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 1912-1915 (*Vibratilicin*)

Neofolitispate 1

N-115

[229160-50-3]

C₄₀H₆₇N₃O₆ 685.986Related to Ptilomyalin A, P-774. Alkaloid from the sponge *Neofolitispa dianchora*. Antiviral agent.Lower homologue (n = 14): *Neofolitispate 3*

[229160-52-5]

C₃₈H₆₃N₃O₆ 657.932Alkaloid from *Neofolitispa dianchora*. Antiviral agent.Lower homologue (n = 15): *Neofolitispate 2*. *Crambescidin 657 methyl ester*

[229160-51-4]

C₃₉H₆₅N₃O₆ 671.959Alkaloid from *Neofolitispa dianchora*. Antiviral agent. Oil. [α]_D -18 (c, 1 in CHCl₃).

Lower homologue (n = 15), parent acid:

Crambescidin 657

[214215-58-4]


C₃₈H₆₃N₃O₆ 657.932Alkaloid from *Crambe crambe*. Cytotoxic agent. Zwitterionic.Lower homologue (n = 15), 13,14,15-triepipimer, parent acid: *Isocrambescidin 657*

[214215-60-8]

C₃₈H₆₃N₃O₆ 657.932Alkaloid from *Crambe crambe*. Zwitterionic.Venkateswarlu, Y. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 254-256 (*isol*)Coffey, D.S. *et al.*, *J.A.C.S.*, 2000, **122**, 4893-4903; 4904-4914 (*synth*)

Neogermin


N-116

4,7-Epoxy-3,14,16,20-cevanetrol
[99211-68-4]C₂₇H₄₃NO₅ 461.64Alkaloid from the roots and rhizomes of *Veratrum japonicum*. Amorph. solid. Mp 188-190°. [α]_D²⁰ -13.8 (c, 0.003 in CHCl₃).Cong, Y. *et al.*, *Planta Med.*, 2007, **73**, 1588-1591 (*isol*, *pmr*, *cmr*)

Neoharringtonine

N-117

[142748-51-4]

C₃₀H₃₃NO₈ 535.593Alkaloid from *Cephalotaxus fortunei* and from *Cephalotaxus harringtonia* var. *drupacea*. Shows significant antileukaemic activity. Oil. [α]_D -148 (c, 0.48 in MeOH). λ_{max} 291 (log ε 3.61) (MeOH). λ_{max} 291 (ε 4073) (MeOH) (Berdy).


1''S-Hydroxy: 1''-Hydroxyneoharringtonine. 3'-Hydroxyneoharringtonine

C₃₀H₃₃NO₉ 551.592Isol. from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Shows antileukaemic activity. Oil. [α]_D -126 (c, 0.20 in MeOH). λ_{max} 291 (log ε 3.66) (MeOH). λ_{max} 291 (ε 4743) (MeOH) (Berdy).Homologue (n=2): *Homoneoharringtonine*C₃₁H₃₅NO₈ 549.619Isol. from leaves and stems of *Cephalotaxus harringtonia* var. *drupacea*. Shows antileukaemic activity. Oil. [α]_D -114 (c, 0.07 in MeOH). λ_{max} 291 (log ε 3.39) (MeOH). λ_{max} 291 (ε 3143) (MeOH) (Berdy).Wang, D.Z. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 173; *CA*, **117**, 86698cTakano, I. *et al.*, *Phytochemistry*, 1997, **44**, 735 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *derivs*)

Neohuperzidine

N-118


[677335-27-2]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*. Cryst. (CHCl₃/Me₂CO). Mp 203-205°. [α]_D²² -48.2 (c, 0.1 in EtOH). λ_{max} 202 (log ε 3.7) (no solvent reported).Yuan, S.-Q. *et al.*, *Yaoxue Xuebao*, 2002, **37**, 946-949 (*isol*, *pmr*, *cmr*, *ms*)

Neohydroxylunine

N-119


[55812-52-7]

C₁₆H₁₇NO₅ 303.314Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (MeOH/Et₂O). Mp 228-231°.Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1975, **38**, 117 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)Neville, C.F. *et al.*, *J.C.S. Perkin 1*, 1991, 259 (*synth*)

Neokauluamine

N-120


[334491-54-2]

C₇₂H₈₈N₈O₆ 1161.537Alkaloid from an undescribed sponge of the Petrosiidae. Cytotoxic. Needles (EtOH). Mp 184°. [α]_D²⁵ +94.6 (c, 0.1 in CHCl₃). λ_{max} 252 (log ε 4.2); 357 (log ε 3.85) (MeOH).El Sayed, K.A. *et al.*, *J.A.C.S.*, 2001, **123**, 1804-1808 (*isol*, *pmr*, *cmr*)

Neolamellarin A

N-121

[959860-11-8]

C₂₄H₁₉NO₄ 385.418Alkaloid from *Dendrilla nigra*. Needles. Mp 144-146°. λ_{max} 210 (log ε 4.53); 250 (log ε 4.41); 285 (log ε 4.27) (MeOH).**7-Hydroxy: 7-Hydroxyneolamellarin A**
[959662-26-1]C₂₄H₁₉NO₅ 401.418Alkaloid from *Dendrilla nigra*. Yellow oil. Racemic. λ_{max} 285 (log ε 4.34) (MeOH).**2,5-Dihydro, 2-oxo: Neolamellarin B**

[959860-12-9]


C₂₄H₁₉NO₅ 401.418Alkaloid from *Dendrilla nigra*. Yellow oil. λ_{max} 224 (log ε 4.34); 330 (log ε 4.06) (MeOH).**2,5-Dihydro, 2-oxo, 5-ξ-hydroxy: 5-Hydroxyneolamellarin B**

[959662-29-4]

C₂₄H₁₉NO₆ 417.417Alkaloid from *Dendrilla nigra*. Yellow powder. [α]_D²⁵ +9.2 (c, 0.06 in Me₂CO). λ_{max} 224 (log ε 4.47); 285 (log ε 4.14); 330 (log ε 4.18) (MeOH).Liu, R. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1741-1745 (*isol*, *pmr*, *cmr*)**Neolaugerine**

N-122

[149405-98-1]

C₁₉H₂₄N₂O₂ 312.411Indole alkaloid plausibly derived from and Akuammicine type by bond cleavages and rearrangement. Alkaloid from root bark of *Neolaugeria resinosa* (Rubiaceae). Amorph. orange-red powder. [α]_D²⁵ +89 (c, 0.60 in MeOH).**9E-Isomer: Isonelaugerine**

[149355-67-9]

C₁₉H₂₄N₂O₂ 312.411Alkaloid from root bark of *Neolaugeria resinosa* (Rubiaceae). Amorph.orange-red powder. [α]_D²⁵ +34 (c, 0.80 in MeOH).**9E-Isomer, 15-hydroxy: 15-Hydroxyisonelaugerine**

[149355-68-0]

C₁₉H₂₄N₂O₃ 328.41Alkaloid from root bark of *Neolaugeria resinosa* (Rubiaceae). Amorph. orange-red powder. [α]_D²⁵ +3 (c, 0.20 in MeOH).Weniger, B. *et al.*, *Phytochemistry*, 1993, **32**, 1587 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)**Neoleurocristine**

N-123

[1359-90-6]

C₄₆H₅₆N₄O₁₂ 856.968

Bisindole alkaloid. Struct. unknown.

Alkaloid from *Catharanthus roseus* (Apocynaceae). Plates (MeOH). Mp 188-196° dec. [α]_D²⁶ -57.87 (c, 1 in CHCl₃).Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*isol*, *uv*, *ir*)**Neoleurosidine**

N-124


[1359-91-7]

C₄₈H₆₂N₄O₁₁ 871.038

Bisindole alkaloid. Struct. unknown.

Alkaloid from *Catharanthus roseus* (Apocynaceae). Rods (MeOH). Mp 219-225° dec. [α]_D²⁶ +41.6 (c, 1 in CHCl₃).Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*isol*, *uv*, *ir*)**Neolitsine**

N-125

1,2:9,10-Bis(methylenedioxy)aporphineC₁₉H₁₇NO₄ 323.348**(S)-form** [2466-42-4]Alkaloid from the leaves of *Neolitsea pulchella* and *Laurus nobilis* (bay laurel), also from *Cassipoupa americana* (*Cassipoupa filiformis*) (Lauraceae). Needles (Me₂CO). Mp 149-150°. [α]_D +56.5 (c, 1.57 in CHCl₃).**Hydrochloride:**

Cryst. (EtOH). Mp 219° (chars).

Picrate:

Cryst. (EtOH). Mp 186° dec.

N-De-Me: Cryptodorine. 1,2:9,10-Bis(methylenedioxy)noraporphine. Norneolitsine

[41787-55-7]

C₁₈H₁₅NO₄ 309.321Alkaloid from the trunk bark of *Cryptocarya odorata* and leaves of *Laurus nobilis* (bay laurel) (Lauraceae). Unstable cryst.; cryst. + 2H₂O (assulfate). Mp 219-221° (sulfate). [α]_D¹⁹ +61 (CHCl₃).**6a,7-Didehydro: Dehydroneolitsine**

[40158-86-9]

C₁₉H₁₅NO₄ 321.332Alkaloid from *Gutteria goudotiana* (Annonaceae). Yellow cryst. (EtOH). Mp 201-203° (196-198°).**3-Methoxy, N-de-Me: 3-Methoxy-1,2:9,10-bis(methylenedioxy)noraporphine. Cassythidine**

[6081-07-8]

C₁₉H₁₇NO₅ 339.347Alkaloid from *Cassipoupa filiformis* (*Cassipoupa americana*) (Lauraceae). Prisms (CHCl₃/MeOH). Mp 206-207°. [α]_D +15 (c, 0.64 in CHCl₃).**(±)-form**Synthetic. Needles (Me₂CO). Mp 148-150°.N-De-Me: Synthetic. Cryst. (Me₂CO/Et₂O). Mp 169° dec.Hui, W.H. *et al.*, *J.C.S.*, 1965, 2285 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, **19**, 297 (*Cassipoupa*)Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2443 (*isol*, *uv*)Bick, I.R.C. *et al.*, *Bull. Soc. Chim. Fr.*, 1972, 4596 (*isol*, *uv*, *pmr*, *ms*, *struct*, *Cryptodorine*)Moltrasio, G.Y. *et al.*, *J.C.S. Perkin 1*, 1973, 349 (*synth*, *uv*)Pai, B.R. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 1042 (*synth*, *uv*, *pmr*, *ms*, *Cryptodorine*)Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (*cd*)Pech, B. *et al.*, *J. Nat. Prod.*, 1982, **45**, 560 (*isol*)Castedo, L. *et al.*, *Phytochemistry*, 1991, **30**, 2781 (*Dehydroneolitsine*)**Neolitsinine**


N-126

[1359-92-8]

C₁₉H₂₃NO₃ 313.396Struct. unknown. Alkaloid from the leaves of *Neolitsea pulchella* (Lauraceae). Cryst. (Me₂CO). Mp 214-215°. [α]_D +94.7 (c, 2.31 in CHCl₃).Hui, W.H. *et al.*, *J.C.S.*, 1965, 2285-2286**Neonaucleoside A**

N-127

[539825-80-4]




Absolute Configuration

C₄₄H₅₈N₂O₁₈ 902.945Alkaloid from the dried roots of *Neonauclea sessilifolia*. Amorph. powder. [α]_D²⁸ -133 (c, 1 in MeOH). λ_{max} 227 (log ε 4.54); 273 (log ε 3.73); 283 (sh) (log ε 3.72); 290 (sh) (log ε 3.64) (MeOH).Itoh, A. *et al.*, *Phytochemistry*, 2003, **62**, 359-369 (*isol*, *pmr*, *cmr*)

Neonaucleoside B

[539825-81-5]


Absolute
ConfigurationC₄₅H₅₄N₂O₂₀ 942.923

Alkaloid from the dried roots of *Neonauclea sessilifolia*. Amorph. powder. $[\alpha]_D^{28}$ -212 (c, 1 in MeOH). λ_{\max} 238 (log ϵ 4.58); 263 (log ϵ 4.43); 290 (sh) (log ϵ 3.89); 304 (log ϵ 3.73); 335 (sh) (log ϵ 3.56); 349 (log ϵ 3.65); 387 (sh) (log ϵ 3.21) (MeOH).

Itoh, A. et al., *Phytochemistry*, 2003, **62**, 359-369 (isol, pmr, cmr)

Neonaucleoside C

[539825-82-6]

Absolute
ConfigurationC₄₄H₅₄N₂O₂₀ 930.912


Alkaloid from the dried roots of *Neonauclea sessilifolia*. Cryst. (H₂O). $[\alpha]_D^{28}$ -326 (c, 0.2 in Py). Mp >300°. λ_{\max} 220 (log ϵ 4.61); 243 (log ϵ 4.34); 282 (sh) (log ϵ 3.84); 288 (log ϵ 3.74) (MeOH).

Itoh, A. et al., *Phytochemistry*, 2003, **62**, 359-369 (isol, pmr, cmr)

Neonocintine

N-130

10-(3-Aminopropyl)octadecahydro-2H-pyrido[1,2-f][1,6]diazacyclooctadecin-11(6H)-one, 9CI

C₂₃H₄₅N₃O 379.628**(ξ)-form** [53602-28-1]

Alkaloid from the stem bark of *Oncinotis nitida* (Apocynaceae). Obt. only in admixture with Oncinotine, O-91.

[62107-29-3 ((±)-form)]

Guggisberg, A. et al., *Helv. Chim. Acta*, 1974, **57**, 414; 1976, **59**, 3013 (struct, synth, ir, ms)


N-128

Hou, D.-R. et al., *J.O.C.*, 2004, **69**, 6094-6099 (synth)

Neooxazolomycin

N-131

Antibiotic AIL-B
[96851-27-3]

C₃₄H₄₇N₃O₉ 641.76

Polyene-type antibiotic. Prod. by *Streptomyces* sp. Y32026 and a *Penicillium* sp. Active against Ehrlich ascites tumour *in vivo*. Sol. CHCl₃, EtOAc, MeOH. λ_{\max} 230 (ϵ 32000); 265 (ϵ 28000); 275 (ϵ 34000); 285 (ϵ 27000) (MeOH) (Derep). [99745-19-4]

Japan. Pat., 1985, 85 81 184; CA, **104**, 49830

(isol, struct, props)

Takahashi, K. et al., *Tet. Lett.*, 1985, **26**, 1077 (struct)

Kawai, S. et al., *Agric. Biol. Chem.*, 1989, **53**, 1127 (isol)


Kende, A.S. et al., *J.A.C.S.*, 1990, **112**, 4070 (synth)

Onyango, E.O. et al., *Angew. Chem., Int. Ed.*, 2007, **46**, 6703-6705 (synth)

Neooxygambirtannine

N-132

[23594-97-0]

C₂₁H₁₆N₂O₃ 344.369

Alkaloid from *Uncaria gambier* (*Ourouparia gambier*) (Rubiaceae). Red. Mp 220°. Artifact.

9-Methoxy, 5,6-didehydro: **Anhydroalstonatine**

[63524-06-1]

C₂₂H₁₆N₂O₄ 372.379

Alkaloid from the bark of *Alstonia venenata* (Apocynaceae). Red. Cryst. Mp 270-272°. λ_{\max} 248 (log ϵ 4.5); 262 (log ϵ 4.49); 312 (log ϵ 4.52) (EtOH).


Merlini, L. et al., *Tetrahedron*, 1967, **23**, 3129-3145 (*Neooxygambirtannine*)

Chatterjee, A. et al., *Indian J. Chem., Sect. B*, 1977, **15**, 183-184 (*Anhydroalstonatine*)

Neopeltolide

N-133

[934220-90-3]



Absolute Configuration

C₃₁H₄₆N₂O₉ 590.712

Stereochem. revised in 2007. Isol. from a sponge of the Neopeltidae. Cytotoxic and antifungal agent. Oil. $[\alpha]_D^{24}$ +24 (c, 0.24 in MeOH). λ_{\max} 218 (log ϵ 4.54); 262 (log ϵ 4.29) (EtOH).

Youngsaye, W. et al., *Angew. Chem., Int. Ed.*, 2007, **46**, 9211-9214 (synth, abs config)

Wright, A.E. et al., *J. Nat. Prod.*, 2007, **70**, 412-416 (isol, pmr, cmr)


Fuwa, H. et al., *Angew. Chem., Int. Ed.*, 2008, **47**, 4737-4739 (synth)

Custar, D.W. et al., *J.A.C.S.*, 2008, **130**, 804-805 (synth, abs config)

Neopine

N-134

8,14-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-ol, 9CI. β -Codeine [467-14-1]

Absolute
configurationC₁₈H₂₁NO₃ 299.369

Minor alkaloid of opium (*Papaver somniferum*), also obt. from *Papaver bracteatum* (Papaveraceae). Has analgesic activity, less than that of Morphine, M-704. Long needles (cyclohexane). Mp 127.5-127.8°. $[\alpha]_D^{26}$ -28.1 (c, 7.7 in CHCl₃).

Hydrobromide:

Cryst. (H₂O). Mod. sol. H₂O. Mp 280-285° dec. (darkens at 240°). $[\alpha]_D^{20}$ +17.1 (c, 5.2 in H₂O).

Sulfate:

Cryst. (Me₂CO). Sol. EtOH; v. sol. H₂O. Mp 166-167° dec. $[\alpha]_D^{28}$ +16.4 (c, 3 in H₂O). Slowly dec. in air.

Homeyer, A.H. et al., *J.O.C.*, 1947, **12**, 356 (isol)

Small, L. et al., *J.O.C.*, 1947, **12**, 359 (struct)

Conroy, H. et al., *J.A.C.S.*, 1955, **77**, 5960 (synth)

Brochmann-Hanssen, E. et al., *J. Pharm. Sci.*, 1964, **53**, 1549 (glc, ilc)

Batterham, T.J. et al., *Aust. J. Chem.*, 1965, **18**, 1799 (pmr)

Weiss, U. et al., *Bull. Soc. Chim. Fr.*, 1965, 3707 (ord)


Wheeler, D.M.S. et al., *J.A.C.S.*, 1967, **89**, 4494 (ms)

Kuppers, F.J.E.M. et al., *Phytochemistry*, 1976, **15**, 443 (isol)

Neopolyalthenol

N-135

[99615-97-1]

C₂₃H₃₁NO 337.504


Alkaloid from *Polyalthia suaveolens* (Annonaceae). Cryst. (petrol). Mp 89°.

$[\alpha]_D^{25}$ -19 (c, 0.9 in EtOH).

Kunesch, N. *et al.*, *Tet. Lett.*, 1985, **26**, 4937
(*uv, ir, pmr, cmr, ms, struct*)

Neopyrrolomycin N-136

3,4,5-Trichloro-2-(2,3-dichloro-1H-pyrrol-1-yl)phenol, 9Cl. 2,3-Dichloro-1-(2,3,4-trichloro-6-hydroxyphenyl)pyrrole [131956-34-8]



$C_{10}H_4Cl_5NO$ 331.411

Prod. by *Streptomyces* sp. MI424-38F1. Active against gram-positive and -negative bacteria and some fungi. Needles (CHCl₃/heptane) (as K salt). Sol. MeOH, Me₂CO, EtOAc, CHCl₃; poorly sol. H₂O. Mp 70-72° (K salt). $[\alpha]_D^{25}$ -4.8 (c, 0.5 in CHCl₃). λ_{max} 319 (ε 6030) (MeOH/NaOH) (Derep). λ_{max} 299 (ε 3715); 319 (ε 6025) (MeOH) (Berdy).


► LD₅₀ (mus, ipr) 75 mg/kg. SL0533500 [132226-97-2]

Nogami, T. *et al.*, *J. Antibiot.*, 1990, **43**, 1193
(*isol, struct*)

Tatsuta, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 1994, **67**, 1449 (*synth*)

Neosalvianene N-137

[790673-00-6]




$C_{21}H_{21}NO_2$ 319.402

Alkaloid from the roots of *Salvia miltiorrhiza*. Needles (EtOAc/hexane). Mp 230-231° (213-215°). λ_{max} 201 (log ε 4.9); 255 (log ε 5.34); 263 (log ε 5.43) (MeOH).

Don, M.-J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1066-1070 (*isol, synth, pmr, cmr, ms*)

Neosidomycin N-138

[72033-44-4]



$C_{17}H_{20}N_2O_6$ 348.355

Nucleoside-type antibiotic. Metab. from *Streptomyces hygroscopicus*. Weakly active against gram-negative bacteria. Amorph. powder + ½H₂O. Sol. MeOH, butanol. Mp 93-103°. $[\alpha]_D^{26}$ +51 (c, 0.48 in MeOH). Similar to Antibiotic SF 2140, A-1255. λ_{max} 270 (ε 8130); 279 (ε 7760); 283 (sh) (ε 7410); 298 (ε 5750) (MeOH) (Derep).

Di-O-Ac: Mp 78-80°.

4-Methoxy-4-Methoxyneosidomycin. Kahakamide A

[362051-29-4]

$C_{18}H_{22}N_2O_7$ 378.381

Prod. by the marine *Nocardiopsis dassonvillei*. Stereochem. not confirmed. λ_{max} 222 ; 266 ; 286 ; 295 (no solvent reported).

4-Methoxy, 6'-parent acid, 6'-amide: Kahakamide B

[362051-30-7]

$C_{17}H_{21}N_3O_6$ 363.369

Prod. by the marine *Nocardiopsis dassonvillei*. Stereochem. not confirmed.

Furuta, R. *et al.*, *Tet. Lett.*, 1979, 1701-1704
(*isol, uv, ir, pmr, ms*)


Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711-1739 (*rev*)

Buchanan, J.G. *et al.*, *J.C.S. Perkin I*, 1994, 1417-1426 (*synth*)

Schumacher, R.W. *et al.*, *Tet. Lett.*, 2001, **42**, 5133-5135 (*Kahakamides*)

Neostemofoline N-139

[1002109-44-5]




$C_{22}H_{29}NO_5$ 387.475

Alkaloid from *Stemona japonica*. Oil. $[\alpha]_D^{23}$ +230 (c, 0.11 in MeOH). λ_{max} 305 (log ε 4.32) (MeOH).

Tang, C.-P. *et al.*, *J. Nat. Prod.*, 2008, **71**, 112-116 (*isol, pmr, cmr*)

Neostemonine N-140

[164124-85-0]



$C_{18}H_{25}NO_4$ 319.4

Alkaloid from the roots of *Stemona japonica*. Cryst. (EtOH) (as hydrochloride). Mp 198-200° (hydrochloride). $[\alpha]_D^{25}$ +245 (c, 1.3 in EtOH) (hydrochloride).

1,2,3,9a-Tetrahydro: *Bisdehydroneoste-*

monine

[145701-16-2]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid from the roots of *Stemona japonica*. Cryst. (Et₂O). Mp 218-221°. $[\alpha]_D^{25}$ +187 (c, 0.33 in EtOH).

Demethoxy, 11S,12R-dihydro: Isonestemocochinine

[1003567-59-6]

$C_{17}H_{25}NO_3$ 291.389

Alkaloid from the roots of *Stemona cochinchinensis*. Isol. as a 1:1 mixt. with Neostemocochinine.

Demethoxy, 11S,12S-dihydro: Neostemocochinine

[1003567-58-5]

$C_{17}H_{25}NO_3$ 291.389

Alkaloid from the roots of *Stemona cochinchinensis*. Isol. as a 1:1 mixt. with Isonestemocochinine.


Ye, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1205-1208 (*isol, ir, pmr, cmr, ms*)

Lin, L.-G. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2167-2175 (*Neostemocochinine, Isonestemocochinine*)

Neosurugatoxin

N-141

[80680-43-9]



$C_{30}H_{34}BrN_5O_{15}$ 784.527

Toxin from the mid-gut gland of *Babylonia japonica* (Japanese ivory shell) of dietary origin and from *Corynebacterium* sp. Shows powerful anti-mydriatic activity. Insecticide, nematocide. Prisms + 1H₂O (H₂O). Sol. H₂O-MeOH. Mp 331-335° dec. Extremely unstable in alkaline medium and fairly heat-labile. λ_{max} 220 (ε 44700); 282 (ε 15100); 310 (sh) (ε 9120); 325 (sh) (ε 6310) (H₂O) (Derep).

5'-O-Dexylosyl: Prosurugatoxin

[99102-40-6]

$C_{25}H_{26}BrN_5O_{11}$ 652.411


Isol. from *Babylonia japonica*.

Ganglion blocking agent, phycotoxin, mydriatic agent. λ_{max} 220 (ε 44700); 282 (ε 15100); 310 (sh) (ε 9120); 325 (sh) (ε 6310) (H₂O) (Derep).

Kosuge, T. *et al.*, *Tet. Lett.*, 1981, **22**, 3417-3420 (*cryst struct*)

- Kosuge, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 3255-3259; 1985, **33**, 2890-2895; 3059-3061 (*isol*, *Prosurगतoxin*)
 Inoue, S. *et al.*, *CA*, 1983, **100**, 191639 (*synth*)
 Hayashi, E. *et al.*, *J. Neurochem.*, 1984, **42**, 1491 (*props*)
 Inoue, S. *et al.*, *Tet. Lett.*, 1986, **27**, 5225 (*synth*)
 Inoue, S. *et al.*, *Yakugaku Zasshi*, 1987, **107**, 645 (*rev, synth*)
 Wada, A. *et al.*, *Methods Neurosci.*, 1992, **8**, 311 (*rev, props*)
 Inoue, S. *et al.*, *Tetrahedron*, 1994, **50**, 2753 (*synth*)


Neosutchueneine **N-142**
 [151271-91-9]



$C_{36}H_{40}N_2O_6$ 596.722
 Alkaloid from roots of *Cyclea sutchuensis* (Menispermaceae). Powder. $[\alpha]_D^{26} +7.8$ (c, 0.158 in EtOH).

Wang, X.-K. *et al.*, *Phytochemistry*, 1993, **33**, 1253 (*isol, uv, ir, pmr, cmr, ms, struct*)


Neosymbioimine **N-143**
 [865093-33-0]



$C_{21}H_{27}NO_5S$ 405.514
 Alkaloid from the marine dinoflagellate *Symbiodinium* sp. Oil. $[\alpha]_D^{23} +149$ (c, 0.1 in MeOH).

Kita, M. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 5253-5258 (*isol, pmr, cmr*)
 Varseev, G.N. *et al.*, *Org. Lett.*, 2007, **9**, 1461-1464 (*synth*)

Neothalfine **N-144**




$C_{38}H_{36}N_2O_8$ 648.711
 Alkaloid from the roots of *Thalictrum atriplex*. Amorph. powder. Mp 132-

135°. $[\alpha]_D +36.4$ (c, 0.5 in $CHCl_3$).
 λ_{max} 210 ; 240 (sh) ; 256 ; 340 (MeOH).

Gao, G.-Y. *et al.*, *J. Asian Nat. Prod. Res.*, 2005, **7**, 805-809 (*isol, cd, pmr, cmr, ms*)

Neothramycin A, 9CI **N-145**
 1,2,3,11a-Tetrahydro-3,8-dihydroxy-7-methoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI
 [59593-16-7]



Absolute Configuration

$C_{13}H_{14}N_2O_4$ 262.265
 Isol. from *Streptomyces* spp.
 Antineoplastic agent showing less toxicity than other anthramycins. Amorph. powder. Sol. MeOH, dioxan, DMF, DMSO, $CHCl_3$; fairly sol. H_2O ; poorly sol. C_6H_6 , hexane. Mp 132-147° dec. $[\alpha]_D^{26} +272$ (c, 0.52 in dioxan). Log P -0.64 (uncertain value) (calc). Coml. prod. Neothramycin is mixt. of epimers. λ_{max} 223 (ε 23200); 240 (sh) (ε); 265 (ε 7600); 320 (ε 3640) (90% MeOH/HCl) (Derep). λ_{max} 228 (ε 16700); 254 (ε 14800); 291 (ε 11100); 324 (ε 10800) (MeOH aq./NaOH) (Derep). λ_{max} 223 (ε 22400); 240 (sh) (ε); 265 (ε 7600); 318 (ε 4100) (90% MeOH) (Derep). λ_{max} 223 (E1%/1cm 936); 310 (E1%/1cm 178) (MeOH) (Berdy).

► LD₅₀ (mus, ivn) 20 - 30 mg/kg, LD₅₀ (mus, ipr) 20 - 50 mg/kg. UY8546000

3-Epimer: Neothramycin B
 [59593-15-6]


From *Streptomyces* spp. Antineoplastic agent. Amorph. powder. Sol. MeOH, DMF, DMSO, dioxan, $CHCl_3$; fairly sol. H_2O ; poorly sol. C_6H_6 , hexane. Mp 144-151° dec. $[\alpha]_D^{26} +314$ (c, 0.48 in dioxan). Log P -0.64 (uncertain value) (calc). λ_{max} 223 (ε 23200); 240 (sh); 265 (ε 7600); 320 (ε 3640) (90% MeOH/HCl) (Derep). λ_{max} 228 (ε 16700); 254 (ε 14800); 291 (ε 11100); 324 (ε 10800) (MeOH aq./NaOH) (Derep). λ_{max} 223 (ε 22400); 240 (sh); 265 (ε 7600); 318 (ε 4100) (90% MeOH) (Derep). λ_{max} 223 (E1%/1cm 980); 310 (E1%/1cm 178) (MeOH) (Berdy). λ_{max} 224 (ε 26200); 320 (ε 4080) (MeOH-HCl) (Berdy). λ_{max} 228 (ε 20900); 254 (ε 19000); 291 (ε 11900); 324 (ε 12200) (MeOH-NAOH) (Berdy).

► LD₅₀ (mus, ivn) 20 - 30 mg/kg, LD₅₀ (mus, ipr) 28 - 40 mg/kg. UY8545700 [67298-49-1]

Takeuchi, T. *et al.*, *J. Antibiot.*, 1976, **29**, 93 (*isol, nmr, ir, uv, ms*)
 Ger. Pat., 1977, ((*Microbiochemical Research Foundation*))2 645 528; *CA*, **87**, 51624k (*manuf*)

- Miyamoto, M. *et al.*, *J. Antibiot.*, 1977, **30**, 340 (*synth, struct, uv, pmr, cmr*)
 Kusama, T. *et al.*, *J. Antibiot.*, 1981, **34**, 845 (*hplc*)
 Fujita, H. *et al.*, *J. Antibiot.*, 1982, **35**, 1093 (*pharmacokinetic, exp, human*)
 Tanaka, N. *et al.*, *Antibiotics (N.Y.)*, 1983, **6**, 136 (*rev*)
 Mori, M. *et al.*, *Chem. Comm.*, 1986, 841 (*synth*)
 Andriamialisoa, R.Z. *et al.*, *Tet. Lett.*, 1986, 841 (*synth*)
 Fukuyama, T. *et al.*, *J.A.C.S.*, 1990, **112**, 7050 (*synth*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NCM275


Neotrilobine **N-146**
 [158371-93-8]



$C_{27}H_{26}N_2O_5$ 458.513
 Alkaloid from roots of *Cocculus trilobus* (Menispermaceae).

Chen, H.S. *et al.*, *Chin. Chem. Lett.*, 1994, **5**, 385; *CA*, **121**, 226368e (*isol, struct*)

Neovasipyrindone **N-147**



$C_{16}H_{25}NO_3$ 279.378

N-Et: Neovasipyrindone E
 [184777-35-3]

$C_{18}H_{29}NO_3$ 307.432
 Isol. from *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20} +178$ (c, 0.5 in EtOH). λ_{max} 201 (log ε 3.82); 262 (log ε 3.72); 322 (log ε 3.7) (EtOH).

N-(2-Methylpropyl): Neovasipyrindone C
 [172430-71-6]

$C_{20}H_{33}NO_3$ 335.486
 From *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20} +383$ (c, 1.0 in EtOH).

N-(2-Methylbutyl): Neovasipyrindone A
 [172430-69-2]

$C_{21}H_{35}NO_3$ 349.512
 Metab. from the fungus *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20} +366$ (c, 1.0 in EtOH).

N-(3-Methylbutyl): Neovasipyrindone B
 [172430-70-5]

$C_{21}H_{35}NO_3$ 349.512
 From *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20} +392$ (c, 1.0 in EtOH).

N-(2-Phenylethyl): Neovasipyrindone D
 [184777-33-1]

$C_{24}H_{33}NO_3$ 383.53

From *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20}$ +188 (c, 1.0 in EtOH). λ_{max} 202 (log ϵ 3.91); 262 (log ϵ 3.88); 323 (log ϵ 3.93) (EtOH).

N-(3-Carbamoylpropyl): **Neovasipyridone F**

[184777-36-4]

$C_{20}H_{32}N_2O_4$ 364.484


From *Neocosmospora vasinfecta*. Oil. $[\alpha]_D^{20}$ +260 (c, 0.2 in EtOH). λ_{max} 202 (log ϵ 3.61); 262 (log ϵ 3.91); 322 (log ϵ 3.9) (EtOH).

Nakajima, H. *et al.*, *Phytochemistry*, 1995, **40**, 1643; 1996, **43**, 1015

Neoverataline A

N-148

[603137-25-3]



$C_{27}H_{41}NO_8$ 507.623

Alkaloid from the roots and rhizomes of *Veratrum taliense*. Antifungal agent. Cryst. (MeOH/EtOAc). Mp 220-222°. $[\alpha]_D^{25}$ +91.7 (c, 0.36 in MeOH).

7 α -Hydroxy: **Neoverataline B**

[603137-26-4]

$C_{27}H_{41}NO_9$ 523.622


Alkaloid from the roots and rhizomes of *Veratrum taliense*. Antifungal agent. Needles. Mp 265-267°. $[\alpha]_D^{25}$ -46.8 (c, 0.32 in MeOH).

Zhou, C.-X. *et al.*, *Tetrahedron*, 2003, **59**, 5743-5747 (*isol*, *pmr*, *cmr*)

Neoxaline

N-149

[71812-10-7]



Absolute Configuration

$C_{23}H_{25}N_5O_4$ 435.482

Closely related to Meleagrins, M-162. Prod. by *Aspergillus japonicus* Fg-551. Weak CNS stimulant, mycotoxin. Needles (C_6H_6). Mp 202° dec. $[\alpha]_D^{24}$ -16.3 (c, 1 in $CHCl_3$). λ_{max} 237 (ϵ 16000); 300 (sh); 310 (ϵ 28000); 320 (sh) (MeOH/HCl) (Derep). λ_{max} 245 (ϵ 17000); 310 (sh); 320 (ϵ 35000); 330 (sh) (MeOH/KOH) (Derep). λ_{max} 237 (ϵ 17600); 330 (ϵ 29600) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 200 - 400 mg/kg. NJ5835000

9-Epimer: 9-Epineoxaline

$C_{23}H_{25}N_5O_4$ 435.482

Prod. by *Penicillium tulipae* IBT 3458. $[\alpha]_D$ +26.5 (c, 0.5 in MeOH). $[\alpha]_D$ +48 (c, 0.5 in $CHCl_3$).

Hirano, A. *et al.*, *J. Antibiot.*, 1979, **32**, 781-785 (*isol*, *uv*, *ir*, *pmr*)

Konda, Y. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 2987-2993 (*cd*, *pmr*, *cmr*)

Overy, D.P. *et al.*, *Biochem. Syst. Ecol.*, 2006, **34**, 345-348 (9-Epineoxaline)

Neoyuzurimine

N-150


Struct. unknown. Alkaloid from the leaves and bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Mp 195-198° (as picrate).

Irikawa, H. *et al.*, *Tet. Lett.*, 1966, 5363 (*isol*)

Nepetalactam

N-151

[115421-73-3]



$C_{10}H_{15}NO$ 165.235

Isol. from a commercial sample of catnip oil (*Nepeta cataria*) (Lamiaceae). Excellent crystalline deriv. of nepetalactone. Cryst. Mp 95-96°. $[\alpha]_D^{25}$ -46.5 (c, 5 in $CHCl_3$). Artifact.


Eisenbraun, E.J. *et al.*, *J.O.C.*, 1988, **53**, 3968-3972 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Nephilatoxin 1

N-152

NPTX I

[119688-54-9]



$C_{29}H_{48}N_8O_5$ 588.749

Isol. from the venom of the Joro spider (*Nephila clavata*).

Deoxy: **Nephilatoxin 8**

[114355-41-8]

$C_{29}H_{48}N_8O_4$ 572.749

Isol. from *Nephila clavata* venom.

Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)

Asani, T. *et al.*, *Pept. Chem.*, 1989, 833 (*synth*)


Miyashita, M. *et al.*, *Tet. Lett.*, 1997, **38**, 8297-8298 (*synth*)

Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)

Nephilatoxin 2

N-153

[119613-48-8]



$C_{37}H_{63}N_{13}O_7$ 801.988


Isol. from the venom of the Joro spider (*Nephila clavata*).

Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)

Nephilatoxin 3

N-154

[119613-49-9]



$C_{38}H_{65}N_{11}O_7$ 788.001


Isol. from the venom of the Joro spider (*Nephila clavata*).

Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)

Nephilatoxin 4

N-155

[119613-50-2]



$C_{45}H_{79}N_{13}O_8$ 930.201


Isol. from the venom of the Joro spider (*Nephila clavata*).










Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)

Nephilatoxin 5



N-156

[119613-51-3]



C₄₀H₆₉N₁₁O₇ 816.055Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)
Saito, H. *et al.*, *Tet. Lett.*, 1998, **39**, 6479-6482 (*synth*)**Nephilatoxin 6** N-157
[119613-52-4]C₄₇H₈₃N₁₃O₈ 958.255Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)
Saito, H. *et al.*, *Tet. Lett.*, 1998, **39**, 6479-6482 (*synth*)**Nephilatoxin 7** N-158
[114355-40-7]C₃₈H₆₄N₁₀O₇ 772.986Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 75; 421 (*isol*)**Nephilatoxin 9** N-159
[114355-42-9]C₃₀H₄₉N₁₁O₅ 643.788Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 75; 421 (*isol*)Miyashita, M. *et al.*, *Tet. Lett.*, 1992, **33**, 2833 (*synth*)**Nephilatoxin 10** N-160
NPTX 10. Neurotoxin NPTX 10
[119613-53-5]C₃₂H₅₃N₁₁O₅ 671.841Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)
Miyashita, M. *et al.*, *Tet. Lett.*, 1992, **33**, 2833 (*synth*)**Nephilatoxin 11** N-161
[119613-54-6]C₂₄H₃₇N₇O₄ 487.601Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)
Miyashita, M. *et al.*, *Tet. Lett.*, 1992, **33**, 2837 (*synth*)**Nephilatoxin 12** N-162
[119613-55-7]C₂₆H₄₁N₇O₄ 515.654Isol. from the venom of the Joro spider (*Nephila clavata*).Toki, T. *et al.*, *Biomed. Res.*, 1988, **9**, 421 (*isol*)
Miyashita, M. *et al.*, *Tet. Lett.*, 1992, **33**, 2837 (*synth*)**Nephilatoxin 473** N-163
NPTX 473C₂₄H₄₁N₇O₂ 459.634Isol. from the venom glands of the spider *Nephila* sp.Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)**Nephilatoxin 501** N-164
NPTX 501C₂₅H₃₉N₇O₄ 501.628Isol. from the venom glands of the spider *Nephila* sp.Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)**Nephilatoxin 565** N-165
Neurotoxin JSTX-3. JSTX-3. NPTX 565
[112163-33-4]C₂₇H₄₇N₇O₆ 565.712Isol. from the venom of the Joro spider *Nephila clavata*. Neurotoxin.

[113954-18-0]

Aramaki, Y. *et al.*, *Biomed. Res.*, 1987, **8**, 241 (*isol*)Hashimoto, Y. *et al.*, *Tet. Lett.*, 1987, **28**, 3511 (*synth*, *pmr*)Nason, D.M. *et al.*, *Tet. Lett.*, 1989, **30**, 2337 (*synth*)Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)**Nephilatoxin 594** N-166
NPTX 594C₂₈H₅₀N₈O₆ 594.753Isol. from the Joro spiders *Nephila clavipes* and *Nephila madagascariensis*. [α]_D¹⁹ -7.9 (c, 1 in H₂O) (as hydrochloride).Wakamiya, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 2001, **74**, 1743-1749 (*synth*, *pmr*, *ms*)**Nephilatoxin 643** N-167
NPTX 643
[205673-37-6]C₃₂H₅₃N₉O₅ 643.828

Isol. from the venom of the Madagascar


spider *Nephilengys borbonica*. Neurotoxin. $[\alpha]_D^{19}$ -1.9 (c, 0.1 in H₂O).

Itagaki, Y. *et al.*, *Nat. Toxins*, 1997, **5**, 1-13 (isol)

Miyashita, M. *et al.*, *Heterocycles*, 1998, **47**, 171-175 (synth)

Neplanocin B N-168

3-(6-Amino-9H-purin-9-yl)-1-(hydroxymethyl)-6-oxabicyclo[3.1.0]hexane-2,4-diol, 9CI
[72877-49-7]



C₁₁H₁₃N₅O₄ 279.255

Nucleoside antibiotic. Isol. from *Ampullariella regularis*. Weakly active against tumours. Platelets. Sol. H₂O, DMSO; poorly sol. EtOAc, hexane. Mp 269-272° dec. $[\alpha]_D^{24}$ -3.5 (c, 1 in DMSO). λ_{max} 260 (ε 15100) (H₂O at pH 2) (Derep). λ_{max} 262 (ε 15800) (H₂O) (Derep). λ_{max} 263 (E1%/1cm 522.7) (NaOH aq.) (Berdy).

▶ RN8990000

Ger. Pat., 1979, 2 917 000; CA, **92**, 109108 (isol, props)


Hayashi, M. *et al.*, *Nucleic Acids Symp. Ser.*, 1980, **8**, 565 (struct, props)

Japan. Pat., 1981, 81 51 414; CA, **95**, 175791 (isol, props)

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (rev)

Nerbowdine N-169

Haemanthine. Hemanthine. Buphanitine
[4673-18-1]



C₁₇H₂₁NO₅ 319.357

Alkaloid from the bulbs of *Boophone disticha*, *Nerine bowdenii*, *Crinum erubescens* and from unidentified *Brunsvigia* sp. (Amaryllidaceae). Cryst. (CHCl₃/Me₂CO or EtOH). Mp 230-232° Mp 244-245° dec. (dimorph.).

Hydrochloride:

Fine needles (EtOH aq.). Mp 254-265° dec. $[\alpha]_D^{23}$ -86.1 (c, 0.87 in H₂O).

Nitrate: Mp 222-224°.

3-Ac: 3-O-Acetylnerbowdine

[100196-22-3]

C₁₉H₂₃NO₆ 361.394

Alkaloid from the bulbs of *Brunsvigia disticha* and *Nerine crispa* (Amaryllidaceae). Cryst. (Et₂O). Mp 207-209°. $[\alpha]_D^{21}$ -116 (c, 0.40 in CHCl₃). $[\alpha]_D^{21}$ -45 (c, 0.395 in EtOH).

Di-Ac: Mp 151-152°. $[\alpha]_D^{25}$ -30.5 (c, 1.05 in CHCl₃).

Demethoxy: **Cribetamine**

[97673-68-2]

C₁₆H₁₉NO₄ 289.33

Alkaloid isol. from *Crinum oliganthum*.

Stereochem. at C-1 and C-3 not certain.

Lyle, R.E. *et al.*, *J.A.C.S.*, 1960, **82**, 2620-2625 (isol, uv)

Goosen, A. *et al.*, *J.C.S.*, 1960, 1094 (isol)

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 491-502 (isol, deriv)

Lloyd, H.A. *et al.*, *J.O.C.*, 1962, **27**, 373-377 (struct)

Döpke, W. *et al.*, *Naturwissenschaften*, 1962, **49**, 469 (3-Acetylnerbowdine)

Hauth, H. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 491-502 (3-Acetylnerbowdine)

Wildman, W.C. *et al.*, *J.O.C.*, 1968, **33**, 3749-3753 (isol)

Trimino, Z. *et al.*, *Rev. Cubana Quim.*, 1997, **9**, 123-127 (Cribetamine)

Nerifline

N-170

[1359-99-5]

C₁₇H₂₁NO₅ 319.357

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine flexuosa* (Amaryllidaceae). Mp 152°.


Hydroiodide: Mp 162-163°.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109 (isol)

Nerinine

N-171

[481-44-7]



C₁₉H₂₅NO₅ 347.41

Alkaloid from *Nerine sarniensis* and *Zephyranthes candida* (Amaryllidaceae). Cryst. (Me₂CO). Mp 209-210°. $[\alpha]_D$ +155 (CHCl₃). Oxid. to Albomaculine.

7-Ketone (lactone): **Albomaculine**

[668-63-3]

C₁₉H₂₃NO₅ 345.394

Alkaloid from the bulbs of *Haemanthus albomaculatus* (Amaryllidaceae). Cryst. (EtOAc). Mp 178-179°. $[\alpha]_D^{16}$ +78 (c, 1.0 in CHCl₃).

7-Ketone (lactone), perchlorate:

Prisms (MeOH). Mp 285-289° dec.

Stereoisomer (?): **Urceoline**

C₁₉H₂₅NO₅ 347.41

Alkaloid from bulbs of *Urceolina miniata* and *Hippeastrum brachyanthrum* (Amaryllidaceae). Prisms (Me₂CO). Mp 189-190°. $[\alpha]_D^{25}$ +180 (c, 0.2 in CHCl₃). Tentative struct.

Stereoisomer, 7-ketone (lactone), (?):

Urmidine

C₁₉H₂₃NO₅ 345.394

Alkaloid from bulbs of *Urceolina miniata* and *Amaryllis parkeri* (Amaryllidaceae). Prisms (Me₂CO). Mp 177-179°. $[\alpha]_D^{25}$ -40 (c, 0.2 in CHCl₃).

Tentative struct.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1954, **87**, 1704; 1955, **88**, 1590; 1957, **90**, 57 (isol)

Briggs, C.K. *et al.*, *J.A.C.S.*, 1956, **78**, 2899 (Albomaculine)

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 1827; 1959, **92**, 2578; 2582 (Urceoline, Urmidine)

Hawksworth, W.A. *et al.*, *J.C.S.*, 1965, 1991 (struct, stereochem)

Jeffs, P.W. *et al.*, *J.O.C.*, 1966, **31**, 189 (struct)

Clardy, J. *et al.*, *J.O.C.*, 1972, **37**, 49 (config)

Nerispine

N-172

[1360-00-5]

C₁₇H₁₉NO₄ 301.341

Struct. unknown. Possibly the 8-methoxy isomer of Falcatine, F-7. Alkaloid from the bulbs of *Nerine undulata* (Amaryllidaceae). Cryst. (Me₂CO). Mp 194-195°. $[\alpha]_D^{22}$ -210 (c, 0.2 in CHCl₃).

Perchlorate:

Prisms (H₂O). Mp 230-235° dec.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129-1134 (isol)

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109

Nerundine

N-173

[1360-01-6]

C₁₈H₂₁NO₅ 331.368

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Nerine undulata* (Amaryllidaceae). Cryst. + 1H₂O. Mp 256° dec. $[\alpha]_D$ -95 (c, 0.2 in CHCl₃).

Perchlorate: Mp 221° dec.

Picrate: Mp 168° dec.


Methiodide: Mp 286° dec.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1959, **46**, 228 (isol)

Netamine A

N-174

[910454-37-4]



Absolute Configuration

R¹ = -(CH₂)₅CH₃, R² = -CH₂CH₂CH₃

C₁₉H₃₅N₃ 305.506

Stereochem. revised in 2008. Related to Ptilocaulin, P-773. Isol. from the sponge *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{20}$ +2.2 (c, 0.65 in CH₂Cl₂).

2',3'-Didehydro(-Z-): **Netamine D**

[910454-40-9]

C₁₉H₃₃N₃ 303.49

Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{25}$ -5.8 (c, 0.03 in MeOH).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (isol, pmr, cmr)

Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (synth, abs config)

Netamine B N-175

[910454-38-5]
As Netamine A, N-174 with
 $R^1 = -CH(CH_3)(CH_2)_4CH_3$, $R^2 = -CH_2CH_3$

$C_{19}H_{35}N_3$ 305.506

Stereochem. revised in 2008. Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{20} +5.3$ (c, 0.05 in CH_2Cl_2).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (*isol, pmr, cmr*)

Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*config*)

Netamine C N-176

[910454-39-6]
As Netamine A, N-174 with
 $R^1 = -(CH_2)_5CH_3$, $R^2 = CH_3$

$C_{17}H_{31}N_3$ 277.452


Stereochem. revised in 2008. Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{25} +3.7$ (c, 0.09 in MeOH).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (*isol, pmr, cmr*)

Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*synth, config*)

Netamine E N-177

[910454-41-0]



Absolute Configuration

$C_{17}H_{29}N_3$ 275.436


Stereochem. revised in 2008. Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{21} +35$ (c, 0.8 in CH_2Cl_2).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (*isol, pmr, cmr*)

Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*synth, config*)

Netamine F N-178

[910454-42-1]



Absolute Configuration

$R^1 = -CH_2CH_3$, $R^2 = CH_3$

$C_{13}H_{19}N_3$ 217.313

Stereochem. revised in 2008. Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{20} +108$ (c, 0.05 in CH_2Cl_2).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (*isol, pmr, cmr*)

Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*config*)

Netamine G N-179

[910454-43-2]

As Netamine F, N-178 with

$R^1 = -(CH_2)_3CH_3$, $R^2 = -CH_2CH_2CH_3$
 $C_{17}H_{27}N_3$ 273.42

Stereochem. revised in 2008. Isol. from *Biemna laboutei*. Pale yellow oil. $[\alpha]_D^{21} +27$ (c, 0.2 in CH_2Cl_2).

Sorek, H. *et al.*, *Tetrahedron*, 2006, **62**, 8838-8843 (*isol, pmr, cmr*)


Yu, M. *et al.*, *J.O.C.*, 2008, **73**, 9065-9074 (*synth, config*)

Netropsin N-180

N'-(2-Carbamoyl-ethyl)-4-[(2-guanidinoacetimidoyl)amino]-1,1'-dimethyl-N,4'-bi[pyrrole-2-carboxamide], 8Cl.

Congocidine. Sinanomycin. Antibiotic K 117. Antibiotic IA 887. Antibiotic T 1384. Antibiotic 2814A. Antibiotic CH 777A. K 117. IA 887. T 1384. CH 777A

[1438-30-8]



$C_{18}H_{26}N_{10}O_3$ 430.469

Oligopeptide antibiotic. Lexitropsin (information-reading agent). Closely related to Distamycin A, D-866. Prod. by *Streptomyces netropsis*, *Streptomyces chromogenes* and *Streptomyces ambofaciens*. Shows antiviral and antimicrobial props. DNA minor groove-binding ligand with specificity for adenine and thymine base pairs. λ_{max} 235 (ϵ 25000); 297 (ϵ 27500) (0.1M HCl) (Derep). λ_{max} 301 (0.1M NaOH) (Derep).

►DW2973000

Hydrochloride:

Hygroscopic prisms. Mp 168-172° dec.

Sulfate:

Needles. Mp 224-225°.

Finlay, A.C. *et al.*, *J.A.C.S.*, 1951, **73**, 341 (*isol*)

Julia, M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1963, **257**, 1115 (*struct, synth, nmr*)

Patel, D.J. *et al.*, *Eur. J. Biochem.*, 1979, **99**, 369-378 (*bibl, pharmacol*)

Lown, J.W. *et al.*, *J.O.C.*, 1985, **50**, 3774 (*synth*)

Debart, F. *et al.*, *J. Med. Chem.*, 1989, **32**, 1074-1083 (*bibl, sar*)

Lown, J.W. *et al.*, *Antiviral Res.*, 1992, **17**, 179-196 (*rev*)

Chen, A.Y. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 8131-8135 (*pharmacol*)

Lown, J.W. *et al.*, *J. Mol. Recognit.*, 1994, **7**, 79-88 (*lexitropsins, rev*)


Numm, C.M. *et al.*, *Biochemistry*, 1997, **36**, 4792-4799 (*DNA complex, cryst struct*)

Singh, S.B. *et al.*, *J.A.C.S.*, 1999, **121**, 3267-3271 (*DNA binding*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NCP875

***Nephila maculata* Neurotoxin-in 3** N-181

Neurotoxin NSTX 3. NSTX 3 [107288-22-2]



$C_{30}H_{52}N_{10}O_7$ 664.804

Toxic principle in the venom of *Nephila maculata*. $[\alpha]_D^{15} +4.3$ (c, 0.51 in H_2O).

[125739-24-4]

Aromaki, Y. *et al.*, *Pept. Chem.*, 1987, 163 (*isol*)


Nason, D.M. *et al.*, *Tet. Lett.*, 1989, **30**, 2337 (*synth*)

Teshima, T. *et al.*, *Tetrahedron*, 1991, **47**, 3305 (*synth*)

Nihei, K. *et al.*, *Tetrahedron*, 2006, **62**, 8335-8350 (*synth*)

***Nephila clavata* Neurotoxin 1 JSTX-1** N-182

JSTX-1 [133698-34-7] [112111-39-4]



$C_{22}H_{38}N_6O_5$ 466.579


Isol. from the venom of the spider *Nephila clavata*.

Hashimoto, Y. *et al.*, *Tet. Lett.*, 1987, **28**, 3511-3514 (*synth, pmr*)

Toki, T. *et al.*, *CA*, 1991, **114**, 201423m (*isol, struct*)

***Nephila clavata* Neurotoxin 2 JSTX-2** N-183

JSTX-2 [133658-52-3]




$C_{36}H_{64}N_{10}O_8$ 764.964

Isol. from the venom of the spider *Nephila clavata*.

Toki, T. *et al.*, *CA*, 1991, **114**, 201423m (*isol*)

***Nephila clavata* Neurotoxin 4 JSTX-4** N-184

JSTX-4 [133689-61-9]




$C_{28}H_{48}N_{10}O_7$ 636.75

Isol. from the venom of the spider *Nephila clavata*.

Toki, T. *et al.*, *CA*, 1991, **114**, 201423m (*isol. struct*)

Newbouldine† N-185

3a,4,5,6-Tetrahydro-3-phenyl-3H-pyrrolo[1,2-b]pyrazole
[155416-35-6]



C₁₂H₁₄N₂ 186.256

Alkaloid from the root bark of *Newbouldia laevis* (Bignoniaceae). Oil. Racemic.

4'-Hydroxy: 4'-Hydroxynewbouldine

[155416-36-7]

C₁₂H₁₄N₂O 202.255

Alkaloid from root bark of *Newbouldia laevis* (Bignoniaceae). Cryst. (Me₂CO). Mp 155-156°.

4'-Methoxy: 4'-Methoxynewbouldine

C₁₃H₁₆N₂O 216.282


Alkaloid from the root bark of *Newbouldia laevis*. λ_{max} 227 (log ε 4.02); 278 (sh) (MeOH).

Adesanya, S.A. *et al.*, *Phytochemistry*, 1994, **35**, 1053 (*isol. uv, ir, pmr, cmr, ms, struct*)

Aladesanmi, A.J. *et al.*, *Planta Med.*, 1998, **64**, 90-91 (*4'-Methoxynewbouldine*)

Ngouniensiine N-186

[83348-21-4]



Absolute Configuration

C₁₉H₂₄N₂ 280.412

Related to the Uleine-dasycarpidan group VX 5880 (3,16-cyclo). Main alkaloid from the stem bark and root bark of *Strychnos ngouniensi* (Loganiaceae). Amorph. [α]_D -44 (c, 1 in CHCl₃).

10- or 11-Glucosyloxy: Glucosylngouniensiine

[88852-55-5]

C₂₅H₃₄N₂O₆ 458.553

Alkaloid from root bark of *Strychnos ngouniensi* (Loganiaceae). [α]_D -107 (c, 0.26 in MeOH). Stereochem. not certain.

20-Epimer: Epingouniensiine

[88721-00-0]

C₁₉H₂₄N₂ 280.412

Alkaloid from the stem bark and root bark of *Strychnos ngouniensi* (Loganiaceae). [α]_D -32 (c, 0.5 in MeOH).

Stereoisomer, 10- or 11-glucosyloxy: Epiglucosylngouniensiine

[88775-09-1]

C₂₅H₃₄N₂O₆ 458.553

Alkaloid from the root bark of *Strychnos ngouniensi* (Loganiaceae). [α]_D -80 (c, 0.29 in MeOH).

Massiot, G. *et al.*, *Tetrahedron*, 1983, **39**, 3645-3656 (*isol. uv, ir, cmr, ms, derivs, biosynth*)

Bosch, J. *et al.*, *Tet. Lett.*, 1987, **28**, 231 (*synth*)

Bennasar, M.L. *et al.*, *Heterocycles*, 1989, **29**, 381 (*synth*)

Hugel, G. *et al.*, *J.O.C.*, 1997, **62**, 578 (*abs config*)

Nicaeensin N-187

N-[4-[[[(Aminoiminomethyl)amino]carbonyl]amino]butyl]-N-methylacetamide, 9CI. 1-(3-Amidinoureido)-4-(N-methylacetamido)butane

[131669-98-2]

[131669-99-3 (*picrate*)]

HN=C(NH₂)NHCONH(CH₂)₄NMeAc


C₉H₁₉N₅O₂ 229.281

Isol. from the red alga *Schottera nicaeensis*. Powder; yellow cryst. (EtOH aq.) (as picrate). Mp 171-173° (*picrate*).

Chillemi, R. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1220 (*isol. pmr, cmr, ms, struct*)

Nicotaxine N-188

[126585-94-2]



C₃₀H₃₇NO₉ 555.624

Alkaloid from the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +112 (c, 0.21 in CHCl₃).

O⁵-(3-Amino-3-phenylpropanoyl): 5-(3-Amino-3-phenylpropanoyl)nicotaxine

[126617-17-2]

C₃₉H₄₆N₂O₁₀ 702.8

From the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +93 (c, 0.41 in CHCl₃).

O⁵-(3-Methylamino-3-phenylpropanoyl): 5-(3-Methylamino-3-phenylpropanoyl)nicotaxine

[126585-95-3]

C₄₀H₄₈N₂O₁₀ 716.827


From the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +103 (c, 0.12 in CHCl₃).

Ettouati, L. *et al.*, *Bull. Soc. Chim. Fr.*, 1989, 687-694 (*isol. pmr, cmr, struct*)

Nicotinamide ribonucleoside N-189

3-(Aminocarbonyl)-1-β-D-ribofuranosylpyridinium(1+), 9CI. Nicotinamide riboside

[1341-23-7]



C₁₁H₁₅N₂O₅[⊕] 255.25

Bromide: [78687-39-5]

C₁₁H₁₅BrN₂O₅ 335.154

Amorph. hygroscopic powder. [α]_D²⁰ -33 (c, 1.42 in H₂O).

5-O-Phosphate inner salt: Nicotinamide ribonucleotide. Nicotinamide ribotide. Nicotinamide mononucleotide

[1094-61-7]

C₁₁H₁₅N₂O₈P 334.222

Intermed. in synth. of nicotinamide adenine nucleotide analogues. Subcomponent of coenzymes I and II. Amorph. powder. [α]_D²⁰ -31.4 (c, 1.94 in H₂O).

5'-O-(2ξ-Acetoxy-3ξ-methylpentanoyl):

[291525-83-2]

C₁₉H₂₇N₂O₈[⊕] 411.431

Isol. from the chrysolimid beetle *Platyphora opima*. [α]_D²⁵ -1.6 (c, 0.19 in MeOH). Counterion not specified. λ_{max} 210 (log ε 3.8); 260 (log ε 3.44) (MeOH).

[75414-16-3]

Haynes, L.J. *et al.*, *J.C.S.*, 1957, 3727-3732 (*synth. phosphate*)

Mikhailopulo, I.A. *et al.*, *Synthesis*, 1981, 388-389 (*synth. pmr, bibl. phosphate*)

Walt, D.R. *et al.*, *J.A.C.S.*, 1984, **106**, 234-239 (*synth. use, phosphate, bibl*)


Micheli, V. *et al.*, *Arch. Biochem. Biophys.*, 1990, **283**, 40-45 (*metab. phosphate*)

Liu, R. *et al.*, *Nucleosides Nucleotides*, 1994, **13**, 1215-1216 (*synth. phosphate*)

Plasman, V. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1261-1264 (*isol*)

Nicotine, BSI, ISO N-190

3-(1-Methyl-2-pyrrolidinyl)pyridine, 9CI. 1-Methyl-2-(3-pyridyl)pyrrolidine
[75202-10-7]



(S)-form

C₁₀H₁₄N₂ 162.234

▶ Fl. p. >104°, autoignition temp. 240/244°.

(R)-form [25162-00-9]

Synthetic. Liq. Bp₇₂₉ 245.5-246.5°.

▶ QS5260000

Dipicrate: Mp 224-225°.

N-De-Me: see 3-(2-Pyrrolidinyl)pyridine, P-957

(S)-form [54-11-5]

Alkaloid from *Nicotiana tabacum* and other *Nicotiana* spp., *Asclepias syriaca*, *Lycopodium* spp., and other spp. (Solanales, Asclepiadaceae, Crassulaceae). Rare spread of occurrence between angiosperms and cryptogametes. Ganglion blocking agent. Horticultural insecticide. Used in chewing gum and transdermally in cigarette smoking cessation programmes. Bp_{730.5} 246.1°. [α]_D -166. Log P 1.32 (calc).

▶ OES: long-term 1 mg m⁻³; short-term 3 mg m⁻³ (Sk). V. toxic in contact with skin. Causes respiratory paralysis. Adverse effects at lower doses esp. gastrointestinal and CNS. Human and exp. teratogen.

Exp. reprod. effects. LD₅₀ (rat, orl) 50 mg/kg. LD₅₀ (rbt, skn) 50 mg/kg.

QS5250000

Hydroiodide: [6012-23-3]

Mp 195°.

Tartrate (1:2): **Nicotine bitartrate**, USAN

[65-31-6]

Used in the treatment of smoking withdrawal syndrome. Cryst. + 2H₂O. Mp 90° (dihydrate). [α]_D²⁰ +26 (dihydrate).

Dipicrate: Mp 224° (218°).

2-Hydroxybenzoate salt: **Eudermol**, **Nicotinyl salicylate**

[29790-52-1]

Parasiticide for scabies. Hexagonal plates. Mp 118°. [α]_D²⁰ +13.

► QS9600000

Compd. with methacrylic acid/divinyl benzene polymer: **Nicotine polacrilex**, USAN. **Nicorette**

[96055-45-7]

Used as a smoking deterrent.

1'-N-Oxide: **Nicotine 1'-N-oxide**. **Oxynicotine**

[491-26-9]

C₁₀H₁₄N₂O 178.233

Alkaloid identified in the leaves, stems and roots of *Nicotiana tabacum*, *Nicotiana affinis* and *Nicotiana sylvestris* (Solanaceae). Red cryst. or oil. Mp 5-10°. Both diastereoisomers identified.

► QS9604100

N-De-Me: see 3-(2-Pyrrolidinyl)pyridine, P-957

(±)-**form** [22083-74-5]

Liq. pK_{a1} 8.02; pK_{a2} 3.12 (25°, as dihydrochloride).

► QS5257500

Dipicrate: Mp 218°.

Methiodide: Mp 219°.

N-De-Me: see 3-(2-Pyrrolidinyl)pyridine, P-957

[6505-86-8, 2624-48-8, 65-31-6, 65-30-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 768B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 294B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1537B (ir)

Karrer, P. et al., *Helv. Chim. Acta*, 1925, 8, 364 (abs config)

Swan, M.L. et al., *J.A.C.S.*, 1949, 71, 1341 (uv)

Witkop, B. et al., *J.A.C.S.*, 1954, 76, 5597 (ir)

Hellmann, H. et al., *Annalen*, 1964, 672, 97 (synth)

Duffield, A.M. et al., *J.A.C.S.*, 1965, 87, 2926 (ms)

Breuer, E. et al., *Tet. Lett.*, 1969, 3595 (synth)

Kim, H.S. et al., *Acta Cryst. B*, 1971, 27, 1123 (cryst struct, salicylate)

Crain, W.O. et al., *J.A.C.S.*, 1971, 93, 990 (cmr)

Phillipson, J.D. et al., *Phytochemistry*, 1975, 14, 2683 (oxide)

Hutchinson, C.R. et al., *J.A.C.S.*, 1976, 98, 6006 (biosynth)

Whidby, J.F. et al., *J.O.C.*, 1976, 41, 1585 (pmr, config)

Enzell, C.R. et al., *Prog. Chem. Org. Nat. Prod.*, 1977, 34, 44 (rev)

Wigle, I.D. et al., *Chem. Comm.*, 1982, 662 (biosynth)

Banerjee, A. et al., *Life Sci. Adv.*, 1982, 1, 327 (rev, pharmacol)

Schievelbein, H. et al., *Pharmacol. Ther.*, 1982, 18, 233 (rev, metab)

Aceto, M.D. et al., *CRC Handb.*

Stereoisomers: Drugs Psychopharmacol., 1984, 67 (rev)

Seeman, J.I. et al., *Heterocycles*, 1984, 22, 165 (rev)

Slaven, R.W. et al., *J. Het. Chem.*, 1984, 21, 1329 (cmr, pmr)

Domino, E.F. et al., *Psychopharmacol. Bull.*, 1986, 22, 870 (rev, pharmacol)

Brunner, H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, 321, 73 (synth, ir, pmr)

Mahboobi, S. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, 321, 175 (synth, ir, pmr)

Kyerematen, G.A. et al., *Drug Metab. Dispos.*, 1988, 16, 125 (pharmacol)

Pesticide Manual, 9th edn., 1991, 8900

Agrochemicals Handbook, 3rd edn., Royal Society of Chemistry, 1992, A298

Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1393

Levin, E.D. et al., *Med. Chem. Res.*, 1993, 2, 509; 522 (revs, pharmacol)

Nicotine and Related Alkaloids: Absorption, Distribution, Metabolism and Excretion, (Gorrod, J.W., ed.), Chapman and Hall, 1993, (book)

Schmidt, B. et al., *Synthesis*, 1998, 42-44 (N-oxide)

Djuric, V.J. et al., *Physiol. Behav.*, 1999, 67, 533-537 (pharmacol)

Girard, S. et al., *Tet. Lett.*, 2000, 41, 9245-9249 (R-isomer, synth)

Fel'pin, F.-X. et al., *J.O.C.*, 2001, 66, 6305-6312 (synth)

Welter, C. et al., *Org. Biomol. Chem.*, 2005, 3, 3266-3268 (synth)


Wagner, F.F. et al., *Tetrahedron*, 2007, 63, 8065-8082 (rev, synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NDN000; NDR000; NDR500; NDS500

Nigeglanine

N-191

6,7,8,9-Tetrahydro-1-hydroxy-3-methylpyridazino[1,2-a]indazol-5-ium inner salt, 9CI



C₁₂H₁₄N₂O 202.255

Alkaloid from the seeds of *Nigella glandulifera*. Solid. Mp 289-290°. λ_{max} 224 ; 270 ; 320 (MeOH).


Liu, Y.-M. et al., *Chem. Pharm. Bull.*, 2004, 52, 454-455 (isol, pmr, cmr)

Elliot, E.L. et al., *Org. Lett.*, 2005, 7, 2449-2451 (synth)

Nigellicine

N-192

[98063-20-8]



C₁₃H₁₄N₂O₃ 246.265

Alkaloid from the seeds of *Nigella sativa* (black cumin) (Ranunculaceae). Yellow cryst. (EtOH). Dec. over a wide temp. range. λ_{max} 240 (ε 20000); 288 (ε 3310); 296 (ε 3240); 353 (ε 3390) (EtOH) (Derep).


Atta-ur-Rahman, et al., *Tet. Lett.*, 1985, 26, 2759-2762 (isol, pmr, cmr, ms, cryst struct)

Elliot, E.L. et al., *Org. Lett.*, 2005, 7, 2449-2451 (synth)

Nigellidine

N-193

6,7,8,9-Tetrahydro-1-hydroxy-11-(4-hydroxyphenyl)-3-methylpyridazino[1,2-a]indazol-5-ium inner salt, 9CI [120993-86-4]



C₁₈H₁₈N₂O₂ 294.352


Alkaloid from seeds of *Nigella sativa* (black cumin) (Ranunculaceae). Related to Nigellicine, N-192. λ_{max} 230 ; 280 ; 328 (MeOH) (Derep).

Atta-ur-Rahman, et al., *Tet. Lett.*, 1995, 36, 1993 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Nigerloxin

N-194

2-(Aminocarbonyl)-3-hydroxy-6-methoxy-5-methyl-4-(1-propenyl)benzoic acid



C₁₃H₁₅NO₅ 265.265


Prod. by *Aspergillus niger* CFR-W-105. Aldose reductase and lipoxygenase inhibitor. Antioxidant. Amorph. yellow powder. Dec. at 253°. λ_{max} 235 (ε 20700); 292 (ε 11600); 358 (ε 4400) (MeOH).

Rao, K.C.S. et al., *J. Antibiot.*, 2002, 55, 789-793 (isol, pmr, cmr, ms, activity)

Nigramide I

N-195

[847552-90-3]




Relative Configuration

$C_{32}H_{44}N_2O_4$ 520.711
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 208 (log ϵ 4.43); 270 (log ϵ 3.89) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide A
[847552-82-3]

N-196




$C_{32}H_{36}N_2O_6$ 544.646
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 207 (log ϵ 4.5); 287 (log ϵ 3.83) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide B
[847552-83-4]

N-197



$C_{34}H_{38}N_2O_6$ 570.684
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 211 (log ϵ 4.54); 271 (log ϵ 4.14) (MeOH).


1',2'-Dihydro: Nigramide F
[847552-87-8]

$C_{34}H_{40}N_2O_6$ 572.7
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. λ_{max} 206 (log ϵ 4.61); 287 (log ϵ 3.96) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide C
[847552-84-5]

N-198




$C_{32}H_{36}N_2O_6$ 544.646
Alkaloid from the roots of *Piper nigrum* (pepper). Prisms (MeOH). Mp 182-183°. Racemic. λ_{max} 210 (log ϵ 4.44); 287 (log ϵ 3.85) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*, *cryst struct*)

Nigramide D
[847552-85-6]

N-199




$C_{36}H_{40}N_2O_6$ 596.722
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 208 (log ϵ 4.51); 285 (log ϵ 4.29) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide E
[847552-86-7]

N-200




$C_{36}H_{40}N_2O_6$ 596.722
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 211 (log ϵ 4.63); 264 (log ϵ 4.31) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide G
[847552-88-9]

N-201




$C_{33}H_{38}N_2O_6$ 558.673
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 206 (log ϵ 4.6); 287 (log ϵ 3.96) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide H
[847552-89-0]

N-202




Relative
Configuration

$C_{30}H_{42}N_2O_4$ 494.673
Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{max} 206 (log ϵ 4.32); 286 (log ϵ 3.58) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide J
[847552-91-4]

N-203



$C_{30}H_{42}N_2O_4$ 494.673
Alkaloid from the roots of *Piper nigrum*


(pepper). Oil. λ_{\max} 205 (log ϵ 4.39); 287 (log ϵ 3.49) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide K

N-204

[847552-92-5]



$C_{29}H_{42}N_2O_4$ 482.662


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 205 (log ϵ 4.26); 286 (log ϵ 3.58) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide L

N-205

[847552-93-6]



$C_{32}H_{44}N_2O_4$ 520.711


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 209 (log ϵ 4.45) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide M

N-206

[847552-94-7]



$C_{31}H_{42}N_2O_4$ 506.684

Alkaloid from the roots of *Piper nigrum*


(pepper). Oil. Racemic. λ_{\max} 208 (log ϵ 4.41) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide N

N-207

[847552-95-8]



$C_{31}H_{44}N_2O_4$ 508.7


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 206 (log ϵ 4.52) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide O

N-208

[847552-96-9]



$C_{28}H_{50}N_2O_2$ 446.715


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 216 (log ϵ 4.12) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide P

N-209

[847552-97-0]



$C_{32}H_{36}N_2O_6$ 544.646


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 207 (log ϵ 4.72); 287 (log ϵ 4.08) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide Q

N-210

[847552-98-1]



$C_{32}H_{36}N_2O_6$ 544.646


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 209 (log ϵ 4.61); 273 (log ϵ 4.16) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide R

N-211

[847552-99-2]



$C_{34}H_{38}N_2O_6$ 570.684


Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 213 (log ϵ 4.56); 267 (log ϵ 4.3) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigramide S

N-212

[847553-00-8]




$C_{36}H_{40}N_2O_6$ 596.722

Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic. λ_{\max} 211 (log ϵ 4.44); 274 (log ϵ 4.1) (MeOH).

Wei, K. *et al.*, *J.O.C.*, 2005, **70**, 1164-1176 (*isol*, *pmr*, *cmr*)

Nigrifactin N-213
2-(1,3,5-Heptatrienyl)-3,4,5,6-tetrahydropyridine, 9CI
[23943-03-5]



$C_{12}H_{17}N$ 175.273
Metab. from several *Streptomyces* strains. Shows antihistaminic props. and affects blood pressure. Extremely unstable. λ_{max} 354 (MeOH/HCl) (Derep). λ_{max} 300 (MeOH at pH 9.2) (Derep). λ_{max} 354 (ϵ 36700) (MeOH) (Derep). λ_{max} 354 (ϵ 36700) (MeOH) (Berdy).

Picrate: Mp 175.5-176° dec. $[\alpha]_D$ 0 (c, 0.5 in MeOH).

Kaneko, Y. *et al.*, *Agric. Biol. Chem.*, 1968, **32**, 783 (*isol*)


Terashima, T. *et al.*, *Tet. Lett.*, 1969, 2535 (*uv*, *ir*, *pmr*, *ms*, *struct*)

Gshwend, H.W. *et al.*, *Tet. Lett.*, 1970, 2711 (*synth*, *config*)

Pailer, M. *et al.*, *Monatsh. Chem.*, 1970, **101**, 508 (*synth*)

Terashima, T. *et al.*, *Chem. Comm.*, 1973, 75 (*biosynth*)

Nigrinodine N-214
1-[7-(3,4-Methylenedioxyphenyl)-2,4-heptadienoyl]pyrrolidine. 7-(3,4-Methylenedioxyphenyl)-2,4-heptadienoic acid pyrrolidide



$C_{18}H_{21}NO_3$ 299.369

(2E,4E)-form [259654-51-8]
Alkaloid from the roots of *Piper amalago* var. *nigrinodum*. Yellow oil.

(2Z,4Z)-form [205507-49-9]
Alkaloid from the leaves of *Piper hispidum*. Antifungal agent. Amorph. solid. λ_{max} 262 (log ϵ 1.9) (MeOH). λ_{max} 262 (MeOH) (Berdy).

Alecio, A.C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 637-639 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Jacobs, H. *et al.*, *J. Indian Chem. Soc.*, 1999, **76**, 713-717 (*isol*, *pmr*, *cmr*)

Nikanine N-215
 $C_{18}H_{27}NO_5$ 337.415
Struct. unknown. Alkaloid from *Trichodesma incanum* (Boraginaceae).
Hydrochloride: Mp 199°.


Nitrate: Mp 182°.

Chloroplatinate: Mp 154°.

N-Oxide: Nikanine N-oxide
 $C_{18}H_{27}NO_6$ 353.414
Alkaloid from *Trichodesma incanum* (Boraginaceae).

Yunusov, S.Yu. *et al.*, *Dokl. Akad. Nauk UzSSR*, 1957, No. 4, 31-34; *CA*, **52**, 13017d

Nikkomycin X N-216
Neopolyoxin A
[72864-26-7]



$C_{20}H_{25}N_5O_{10}$ 495.445
Nucleoside antibiotic. *Isol.* from *Streptomyces tendae* and *Streptomyces cacaoi*. Chitin synthase inhibitor. Active against insects and fungi. Powder. Mp 204-207°. $[\alpha]_D^{21}$ +24.3 (c, 0.325 in H_2O). pK_{a1} 2.9; pK_{a2} 4.3; pK_{a3} 7.3; pK_{a4} 8.7; pK_{a5} 10.3. Amphoteric. λ_{max} 287 (MeOH) (Derep).

▶ BA2928600
4-Carboxylic acid: Neopolyoxin B
[75005-71-9]

$C_{20}H_{25}N_5O_{11}$ 511.444
Nucleoside antibiotic. Prod. by *Streptomyces cacaoi* ssp. *asoensis*. Active against fungi and insects. Mp 192-196°. $[\alpha]_D^{21}$ +19 (c, 0.722 in H_2O). pK_{a1} 2.4; pK_{a2} 3; pK_{a3} 4.4; pK_{a4} 7.7; pK_{a5} 9.1; pK_{a6} 11.1. Has -COOH replacing -CHO.

▶ BA2927800
5'''-Deoxy: Nikkomycin L_X
 $C_{20}H_{25}N_5O_9$ 479.446
Prod. by a genetically engineered *Streptomyces tendae*. Antifungal agent. Lacks the OH group on the pyridine ring.

[75044-70-1]

Dahn, U. *et al.*, *Arch. Microbiol.*, 1976, **107**, 143 (*isol*)

Hagenmaier, H. *et al.*, *Annalen*, 1979, 1494 (*isol*, *struct*)

Brillinger, G.V. *et al.*, *Arch. Microbiol.*, 1979, **121**, 71 (*props*)

Agric. Biol. Chem., 1980, 1709 (*Neopolyoxins*)
Ger. Pat., 1980, 2 900 591; *CA*, **95**, 19726 (*isol*, *use*)

Japan. Pat., 1980, 80 92 395; 1981, 80 26 900; *CA*, **93**, 202702; **95**, 22946

(*Neopolyoxins*)
Fielder, H. *et al.*, *Biotechnol. Lett.*, 1981, **3**, 303 (*hplc*)


Uramoto, M. *et al.*, *Tetrahedron*, 1982, **38**, 1599 (*Neopolyoxins*, *biosynth*, *ir*, *pmr*, *cmr*, *ms*)

Kraimer, E. *et al.*, *Biopolymers*, 1990, **29**, 1297 (*conformm*)

Baumberg, S. *et al.*, *Tet. Lett.*, 1995, **36**, 2351 (*biosynth*)


Bormann, C. *et al.*, *J. Antibiot.*, 1999, **52**, 582-585 (*Nikkomycin L_X*)

Nikkomycin Q_Z N-217
[95259-50-0]



$C_{24}H_{32}N_6O_{12}$ 596.55
Nucleoside antibiotic. From *Streptomyces tendae*. Exhibits lower activity than other Nikkomycins. λ_{max} 264; 287 (sh) (MeOH) (Derep).
Bormann, C. *et al.*, *J. Antibiot.*, 1985, **38**, 9 (*isol*, *props*)
Isono, K. *et al.*, *J. Antibiot.*, 1991, **41**, 1711 (*rev*)

Nikkomycin Z N-218
Nikkomycin. Neopolyoxin C
[59456-70-1]



$C_{20}H_{25}N_5O_{10}$ 495.445
Nucleoside-type antibiotic. Metab. of *Streptomyces tendae* and from *Streptomyces cacaoi*. Antifungal agent which also inhibits chitin biosynth. Mp 194-197°. $[\alpha]_D^{21}$ +49.5 (c, 1 in H_2O). pK_{a1} 2.7; pK_{a2} 4.1; pK_{a3} 7.1; pK_{a4} 8.5; pK_{a5} 9.7. λ_{max} 264; 287 (sh) (MeOH) (Derep).

▶ BA2928200
5'''-Deoxy: Nikkomycin L_Z
 $C_{20}H_{25}N_5O_9$ 479.446
Prod. by a genetically engineered *Streptomyces tendae*. Antifungal agent.

[75044-69-8]

Daehn, U. *et al.*, *Arch. Microbiol.*, 1976, **107**, 143 (*isol*, *ir*, *uv*, *ms*, *nmr*, *struct*)

Koenig, W.A. *et al.*, *Adv. Mass Spectrom.*, 1978, **7B**, 1530 (*ms*)

Kobinata, K. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 1709 (*Neopolyoxin C*)

Koenig, W.A. *et al.*, *Annalen*, 1980, 1728; 1984, 1216 (*config*, *struct*)

Hagenmaier, H. *et al.*, *Annalen*, 1980, 1728 (*config*)

Fiedler, H. *et al.*, *Biotechnol. Lett.*, 1981, **3**, 303 (*hplc*)

Hass, W. *et al.*, *Annalen*, 1982, 1615 (*synth*)

Uramoto, M. *et al.*, *Tetrahedron*, 1982, **38**, 1599 (*Neopolyoxin C*)


Fiedler, H.P. *et al.*, *Clin. Dermatol.*, 1993, **7**, 325 (*rev*)

Bormann, C. *et al.*, *J. Antibiot.*, 1999, **52**, 582-585 (*Nikkomycin L_Z*)

Nilgirine

N-219

12-Hydroxy-18-norsenecionan-11,16-dione, 9CI
[121009-05-2]



C₁₇H₂₃NO₅ 321.372
Alkaloid from *Crotalaria mucronata*,
Crotalaria naragutensis and *Crotalaria striata* (Fabaceae). Mp 127-128°. [α]_D¹⁸
+31.5 (EtOH).

Perchlorate: Mp 261-263° dec.

Ac: Crotastriatine

[11051-94-8]

C₁₉H₂₅NO₆ 363.41

From *Crotalaria striata* (Fabaceae).
Mp 133°.

Ac; perchlorate: Mp 245° dec.

Atal, C.K. *et al.*, *Tet. Lett.*, 1968, 5605-5608
(*isol*, *pmr*, *ms*, *struct*)

Sawhney, R.S. *et al.*, *Planta Med.*, 1972, **21**,
435-437 (*stereochem*)


Batra, V. *et al.*, *Indian J. Chem.*, 1975, **13**, 989-
990 (*Crotastriatine*)

Mattocks, A.R. *et al.*, *Phytochemistry*, 1988,
27, 3289-3291 (*isol*, *pmr*, *ms*)

Ningalin A

N-220

[188111-67-3]

C₁₈H₉NO₈ 367.271

Alkaloid from the ascidian *Didemnum* sp.
Amorph. yellow solid. λ_{max} 262 (ε 8650);
303 (ε 8450); 325 (sh); 370 (ε 2160)
(DMSO/MeOH).


Kang, H. *et al.*, *J.O.C.*, 1997, **62**, 3254 (*isol*, *uv*,
ir, *pmr*, *cmr*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 54-62
(*synth*)

Ningalin B

N-221

[188111-68-4]

C₂₅H₁₉NO₈ 461.427

Alkaloid from the ascidian *Didemnum* sp.
Multidrug-resistant reversal agent. Dark
yellow solid. Mp 303° dec. λ_{max} 204 (ε
46000); 236; 289 (ε 10000); 332 (ε 8700)
(MeOH).

Kang, H. *et al.*, *J.O.C.*, 1997, **62**, 3254-3262
(*isol*, *uv*, *ir*, *pmr*, *cmr*)

Boger, D.L. *et al.*, *J.O.C.*, 2000, **65**, 2479-2483
(*synth*)

Bullington, J.L. *et al.*, *J.O.C.*, 2002, **67**, 9439-
9442 (*synth*)

Iwao, M. *et al.*, *Tet. Lett.*, 2003, **44**, 4443-4446
(*synth*)


Gupton, J.T. *et al.*, *Tetrahedron*, 2003, **59**, 207-
215 (*synth*)

Peschko, C. *et al.*, *Synthesis*, 2006, 3048-3057

Ningalin C

N-222

[188111-69-5]

C₃₂H₂₃NO₁₀ 581.534

Alkaloid from the ascidian *Didemnum* sp.
Amorph. red solid. λ_{max} 204 (ε 34000);
289 (ε 9400); 301 (sh); 355 (ε 5800); 450
(sh) (MeOH).

Kang, H. *et al.*, *J.O.C.*, 1997, **62**, 3254 (*isol*, *uv*,
ir, *pmr*, *cmr*)


Peschko, C. *et al.*, *Tet. Lett.*, 2000, **41**, 9477-
9481 (*synth*)

Namsa-aid, A. *et al.*, *Org. Lett.*, 2002, **4**, 2633-
2635 (*synth*)

Ningpeisine

N-223

[117695-02-0]

C₂₈H₄₇NO₂ 429.685

Alkaloid from the bulbs of *Fritillaria
ningguoensis* (Liliaceae). Cryst. Mp 228-
230°. [α]_D²⁰ +20 (c, 0.5 in EtOH).

O-β-D-Glucopyranoside: **Ningpeisinoid**
[139742-29-3]

C₃₄H₅₇NO₇ 591.827


From bulbs of *Fritillaria ningguoensis*
(Liliaceae). Mp 284-286°. [α]_D²⁰ +24 (c,
0.24 in CHCl₃/MeOH).

Li, Q.H. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 415-
421; 1991, **26**, 794-797

Niphatesine B

N-224

16-(3-Pyridinyl)-7-hexadecyn-1-amine,
9CI. 3-(16-Amino-9-hexadecynyl)pyri-
dine
[132923-11-6]

C₂₁H₃₄N₂ 314.513

Alkaloid from the marine sponge *Ni-
phates* sp. Antineoplastic. Oil. λ_{max} 206 (ε
7800); 228 (ε 3300); 257 (ε 3100); 264 (ε
3400); 269 (ε 2500); 313 (ε 300) (MeOH)
(Berdy).


Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1990, 3301
(*isol*, *uv*, *ir*, *ms*, *pmr*, *cmr*, *struct*)

Rao, A.V.R. *et al.*, *Tet. Lett.*, 1993, **34**, 8329
(*synth*)

Niphatesine C

N-225

β-Methyl-3-pyridinedodecanamine, 9CI.
2-Methyl-12-(3-pyridinyl)dodecylamine.
3-(12-Amino-11-methyldodecyl)pyridine

C₁₈H₃₂N₂ 276.464**(R)-form** [170716-95-7]Synthetic. [α]_D²⁰ -3.2 (c, 3.2 in MeOH).**(S)-form** [132923-12-7]

Alkaloid from the marine sponge *Ni-
phates* sp. Antineoplastic. Oil. [α]_D²⁵ +9.4
(c, 0.053 in MeOH). [α]_D²⁰ +2.4 (c, 0.5 in
MeOH) (synthetic). λ_{max} 206 (ε 2600);
258 (ε 2400); 264 (ε 2700); 269 (ε 2200)
(MeOH) (Berdy).

Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1990, 3301
(*isol*, *uv*, *ir*, *ms*, *pmr*, *cmr*, *struct*)

Rao, A.V.R. *et al.*, *Tet. Lett.*, 1993, **34**, 8329
(*synth*, *abs config*)

Bracher, F. *et al.*, *J.C.S. Perkin 1*, 1995, 2323
(*synth*, *abs config*)


Wang, Y. *et al.*, *J.O.C.*, 2003, **68**, 3090-3098
(*synth*)

Krauss, J. *et al.*, *Arch. Pharm. (Weinheim,
Ger.)*, 2004, **337**, 371-375 (*synth*)

Niphatesine D

N-226

γ-Methyl-3-pyridinedodecanamine, 9CI. 3-
Methyl-12-(3-pyridinyl)dodecylamine. 3-
(12-Amino-10-methyldodecyl)pyridine
[132923-13-8]

C₁₈H₃₂N₂ 276.464

The assignment of the configuration has
been shown to be unreliable
(2004). Alkaloid from the marine sponge
Niphates sp. Antineoplastic. Oil. [α]_D²⁵
+4.4 (c, 0.045 in MeOH).
λ_{max} 207 (ε 4300); 259 (ε 2800); 264
(ε 3200); 270 (ε 2400) (MeOH)
(Berdy).

Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1990, 3301
(*isol*, *uv*, *ir*, *ms*, *pmr*, *cmr*, *struct*)


Rao, A.V.R. *et al.*, *Tet. Lett.*, 1993, **34**, 8329

(*synth, abs config*)

Romeril, S.P. *et al.*, *Tet. Lett.*, 2004, **45**, 3273-3277 (*config*)

Niphatesine E N-227

14-(3-Pyridinyl)-11-tetradecynal O-methyloxime
[143052-05-5]



$C_{20}H_{30}N_2O$ 314.47

Isol. as a *ca.* 1.4:1 mixt. of *E* and *Z* isomers. Alkaloid from the Okinawan marine sponge *Niphates* sp. Exhibits cytotoxicity against murine leukaemia L1210 cells and human epidermoid carcinoma KB cells *in vitro*. Also shows antimicrobial activity against some fungi and gram-positive bacteria.

Oil. λ_{max} 262 (ϵ 19000) (MeOH) (Berdy).

20,N-Dihydro: **Niphatesine H**. N-Methoxy-14-(3-pyridinyl)-11-tetradecyn-1-amine, 9CI

[143052-08-8]

$C_{20}H_{32}N_2O$ 316.486


Alkaloid from *Niphates* sp. Exhibits cytotoxic and antimicrobial activities. Oil. λ_{max} 262 (ϵ 2500) (MeOH) (Berdy).

[143052-09-9]

Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1992, 1291-1294 (*Niphatesines E,H, struct*)

Niphatesine F N-228

16-(3-Pyridinyl)-7-hexadecynal O-methyloxime, 9CI
[143052-06-6]



$C_{22}H_{34}N_2O$ 342.523

Isol. as a 1.7:1 mixt. of *E* and *Z* isomers. Alkaloid from the Okinawan marine sponge *Niphates* sp. Exhibits cytotoxicity against murine leukaemia L1210 cells and human epidermoid KB cells *in vitro*. Also shows antimicrobial activity against some fungi and gram-positive bacteria. Oil. λ_{max} 263 (ϵ 3100) (MeOH) (Berdy).


22,N-Dihydro: see Niphatyne A, N-233

[143052-10-2]

Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1992, 1291 (*isol, uv, ir, pmr, cmr, ms, struct*)

Niphatesine G N-229

α -Methyl-3-pyridinetridecanal O-methyloxime, 9CI
[143052-07-7]



$C_{20}H_{34}N_2O$ 318.501

Isol. as a 3.2:1 mixt. of *E* and *Z* isomers. Alkaloid from the Okinawan marine sponge *Niphates* sp. Exhibits cytotoxicity against murine leukaemia L1210 cells and human epidermoid carcinoma KB cells *in vitro*. Also shows antimicrobial activity against some fungi and gram-positive bacteria. Oil. λ_{max} 263 (ϵ 1500) (MeOH) (Berdy).


[143120-50-7]

Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1992, 1291 (*isol, uv, ir, pmr, ms, struct*)

Niphatoxin A N-230

3-[10-(3-Pyridinyl)-7-deceny]-1-[11-(3-pyridinyl)-8-deceny]pyridinium(1+), 9CI

[142808-48-8]



$C_{35}H_{48}N_3^+$ 510.784

Incorrect struct. given in ref. Alkaloid from the Red Sea sponge *Niphates* sp. Ichthyotoxic and cytotoxic. Counterion not specified.

Talpir, R. *et al.*, *Tet. Lett.*, 1992, **33**, 3033-3034 (*isol, pmr, cmr*)

Niphatoxin B N-231

3-[10-(3-Pyridinyl)-7-deceny]-1-[11-(3-pyridinyl)-9-undeceny]pyridinium(1+), 9CI

[142808-49-9]

As Niphatoxin A, N-230 with $n = 8$

$C_{36}H_{50}N_3^+$ 524.811


Incorrect struct. given in 1992 ref. Alkaloid from the Red Sea sponge *Niphates* sp. Ichthyotoxic and cytotoxic. Counterion not specified.

Talpir, R. *et al.*, *Tet. Lett.*, 1992, **33**, 3033-3034 (*isol, pmr, cmr*)

Kaiser, A. *et al.*, *J.O.C.*, 1999, **64**, 3778-3782 (*synth, ir, pmr, cmr*)

Niphatoxin C N-232

[1001439-38-8]



$C_{36}H_{50}N_3^+$ 524.811

Alkaloid from a *Callyspongia* sp. Cytotoxic. Amorph. solid. Counterion not

specified. λ_{max} 203 ($\log \epsilon$ 4.14); 263 ($\log \epsilon$ 3.84) (MeOH).

Buchanan, M.S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2040-2041 (*isol, pmr, cmr*)

Niphatyne A N-233

N-Methoxy-16-(3-pyridinyl)-7-hexadecyn-1-amine, 9CI
[109741-36-8]



$x = 7, y = 6$

$C_{22}H_{36}N_2O$ 344.539

Isol. from the sponge *Niphates* sp. Cytotoxic agent. Sol. MeOH, $CHCl_3$. λ_{max} 260 (ϵ 2500); 265 (ϵ 3100); 270 (ϵ 2000) (MeOH) (Derep).

Quiñoà, E. *et al.*, *Tet. Lett.*, 1987, **28**, 2467 (*isol, uv, ir, pmr, cmr, ms, struct*)

Niphatyne B N-234

N-Methoxy-16-(3-pyridinyl)-5-hexadecyn-1-amine, 9CI
[109741-37-9]

As Niphatyne A, N-233 with $x = 9, y = 4$

$C_{22}H_{36}N_2O$ 344.539

Isol. from the marine sponge, *Niphates* sp. Cytotoxic agent. Sol. MeOH, $CHCl_3$. Obt. contaminated with Niphatyne A, N-233. λ_{max} 260 (ϵ 2500); 265 (ϵ 3100); 270 (ϵ 2000) (MeOH) (Derep).

Quiñoà, E. *et al.*, *Tet. Lett.*, 1987, **28**, 2468 (*isol, uv, ir, pmr, cmr, ms, struct*)

Niranin N-235

S-Methyl methyl(2-phenylethyl)carbamothioate, 9CI
[92886-90-3]
 $Ph^3CH_2^+CH_2NMeCOSMe$

$C_{11}H_{15}NOS$ 209.312

Isol. from leaves of *Glycosmis* cf. *mauritianana*.

2',3'-Didehydro (*E*-): **Dehydroniranin A**
[184101-44-8]

$C_{11}H_{13}NOS$ 207.296

From leaves of *Glycosmis* cf. *cyano-carpa*. Exhibits pronounced insecticidal activity. Author's numbering. λ_{max} 216 ($\log \epsilon$ 4.11); 279 (sh) ($\log \epsilon$ 4.22); 286 ($\log \epsilon$ 4.23); 302 (sh) ($\log \epsilon$ 4.07) (Et_2O).

2',3'-Didehydro (*Z*-): **Dehydroniranin B**

[184101-51-7]

$C_{11}H_{13}NOS$ 207.296

From leaves of *Glycosmis* cf. *cyano-carpa*. Exhibits pronounced antifungal and insecticidal activity. λ_{max} 216 ($\log \epsilon$ 4.05); 271 ($\log \epsilon$ 4.1) (Et_2O).

2'-Oxo: S-Methyl methyl(phenylacetyl)-carbamothioate, 9CI. **Ritigalin**
[163046-80-8]


$C_{11}H_{13}NO_2S$ 223.295

Isol. from the leaves of *Glycosmis mauritianana* and *Glycosmis parviflora*. Oil. λ_{max} 231 (MeOH). λ_{max} 210; 226 (Et_2O).


Hofer, O. *et al.*, *Monatsh. Chem.*, 1995, **126**,

365-368 (*Ritigalin*)Greger, H. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1163-1168 (*isol, uv, ir, pmr, cmr, ms, struct*)Hinterberger, S. *et al.*, *Tetrahedron*, 1998, **54**, 487-496 (*synth*)**Nirurine†****N-236**


[105801-14-7]

Relative
ConfigurationC₁₂H₁₃NO₃ 219.24Related to the Securinega alkaloid skeleton. Alkaloid from the aerial parts of *Phyllanthus niruri* (Euphorbiaceae).Cryst. (CHCl₃/2-propanol). Mp 205-209°. [α]_D +196.Petchnaree, P. *et al.*, *J.C.S. Perkin 1*, 1986, 1551 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)Magnus, P. *et al.*, *J.A.C.S.*, 1992, **114**, 382 (*synth*)Magnus, P. *et al.*, *Tetrahedron*, 1993, **49**, 8059 (*synth*)**Nishindine****N-237**C₁₅H₂₁NO 231.337Struct. unknown. Alkaloid from *Vitex negundo* (Verbenaceae). Mp 266°.Basu, N.K. *et al.*, *Q. J. Pharm. Pharmacol.*, 1947, **20**, 136-137**Nitaphylline****N-238**

[190962-64-2]

C₄₈H₅₄N₄O₁₄ 910.973Alkaloid from the leaves of *Kopsia teoi* (Apocynaceae). Light yellow oil. [α]_D +136 (c, 1 in CHCl₃). λ_{\max} 217 (log ϵ 4.55); 254 (log ϵ 4.05); 282 (log ϵ 3.51); 289 (log ϵ 3.49) (EtOH).Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **50**, 171-175 (*isol, uv, pmr, cmr, ms*)**Nitidine****N-239**

2,3-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-c]phenanthridinium (1+), 9CI. Angolimine [6872-57-7]

C₂₁H₁₈NO₄[⊕] 348.377Benzo[*c*]phenanthridine alkaloid numbering shown. Alkaloid from a wide variety of *Zanthoxylum* and *Fagara* spp. (Rutaceae). Antitumour agent showing unacceptable toxicity profile for clinical use. Inhibitor of reverse transcriptase activity in RNA oncogenic viruses; also inhibits other enzymes. Shows antifungal activity. λ_{\max} 232 (ϵ 25700); 234 (ϵ 24550); 270 (ϵ 46770); 272 (ϵ 30200); 290 (ϵ 41680); 299 (ϵ 40740); 327 (ϵ 39800); 329 (ϵ 24000); 380 (ϵ 11750) (MeOH) (Berdy). λ_{\max} 228 (ϵ 35300); 278 (ϵ 40700); 311 (ϵ 19600) (EtOH) (Berdy).

Chloride:

C₂₁H₁₈ClNO₄ 383.83Yellow needles (MeOH or EtOH). Mp 220° Mp 238-240° Mp 277-278° (monohydrate) Mp 284-286° dec. (di-hydrate). λ_{\max} 232 (ϵ 21380); 236 (ϵ 22910); 272 (ϵ 40740); 292 (ϵ 37150); 301 (ϵ 36310); 330 (ϵ 36310); 386 (ϵ 1000) (MeOH) (Derep).

Nitrate:

C₂₁H₁₈N₂O₇ 410.382Yellowish-green needles (6M HNO₃/EtOH). Mp 276-278° (239°, 269-270°).

Iodide:

C₂₁H₁₈INO₄ 475.282

Pale yellow needles (EtOH). Mp 284-285°.

N-De-Me: 2,3-Dimethoxy[1,3]benzodioxolo[5,6-c]phenanthridine, 9CI. 8,9-Dimethoxy-2,3-methylenedioxybenzo[*c*]phenanthridine. **Nornitidine**. N-Desmethylnitidine. N-Demethylnitidine [18034-03-2]C₂₀H₁₅NO₄ 333.343Alkaloid from the bark of *Zanthoxylum microcarpum* and *Zanthoxylum myriacanthum*. Needles (Py or Pyl/EtOH). Mp 283-285°. λ_{\max} 229 (log ϵ 4.36); 274 (log ϵ 4.73); 278 (sh) (log ϵ 4.71); 311 (log ϵ 4.15); 330 (sh) (log ϵ 3.89); 348 (log ϵ 3.6); 367 (log ϵ 3.46) (EtOH).O¹⁰-De-Me: **Terihanine**

[112239-71-1]

[111181-33-0]

C₂₀H₁₆NO₄[⊕] 334.351Alkaloid from *Zanthoxylum nitidum*, *Zanthoxylum ovalifolium* and *Fagara nitida*. Yellow prisms (as chloride). Mp 280° (chloride). Melts and resolidifies at 240-245°.O¹¹-De-Me: **Isoterihanine**

[112239-70-0]


[111154-43-9]

C₂₀H₁₆NO₄[⊕] 334.351Alkaloid from *Zanthoxylum ovalifolium*. Yellow prisms (as chloride). Mp 243-247° (chloride).N,O¹⁰-Di-de-Me: **8-O-Demethylnornitidine**

[868247-40-9]

C₁₉H₁₃NO₄ 319.316Alkaloid from *Zanthoxylum caudatum* and *Zanthoxylum quinduensis*. Pale yellow semi-solid. Mp 235-237°. λ_{\max} 280 (log ϵ 6.29) (EtOH).Arthur, H.R. *et al.*, *J.C.S.*, 1959, 1840-1845;4010-4012 (*isol, struct, synth*)Gopinath, K.W. *et al.*, *J.C.S.*, 1959, 4012-4014 (*Nornitidine, synth, uv*)Kuck, A.M. *et al.*, *Phytochemistry*, 1967, **6**, 1541-1550 (*isol*)Calderwood, J.M. *et al.*, *Phytochemistry*, 1970, **9**, 675 (*occur, uv*)Fish, F. *et al.*, *Phytochemistry*, 1971, **10**, 3322-3324; 3325-3327 (*isol*)Vallejos, R.H. *et al.*, *Biochem. Pharmacol.*, 1972, **21**, 1160-1167 (*activity*)Kametani, T. *et al.*, *J. Het. Chem.*, 1973, **10**, 31-33 (*synth*)Zee-Cheng, K.-Y. *et al.*, *J. Het. Chem.*, 1973, **10**, 85-88 (*synth, uv, ir, Nornitidine*)Begley, W.J. *et al.*, *J.C.S. Perkin 1*, 1977, 2324-2328 (*synth*)Cushman, M. *et al.*, *J.O.C.*, 1978, **43**, 286-288 (*synth, pmr, ms*)Phillips, S.D. *et al.*, *J. Het. Chem.*, 1981, **18**, 223-232 (*rev*)Boulware, R.T. *et al.*, *J. Nat. Prod.*, 1981, **44**, 200-205 (*Nornitidine, isol, uv, pmr, ms*)Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4139-4151 (*synth, bibl*)Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2348-2354 (*synth, uv, pmr, ms*)Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2717-2725 (*Terihanine, Isoterihanine, synth*)Martin, G. *et al.*, *J.O.C.*, 1992, **57**, 5907-5911 (*synth*)Pérez, D. *et al.*, *J.O.C.*, 1992, **57**, 5911-5917 (*synth*)Minami, T. *et al.*, *Tet. Lett.*, 1995, **36**, 9505-9508 (*synth*)Green, G.R. *et al.*, *J.C.S. Perkin 1*, 1996, 1647-1648 (*Nornitidine, synth*)Harayama, T. *et al.*, *Heterocycles*, 1998, **48**, 1989-1992 (*synth*)Nissanka, A.P.K. *et al.*, *Phytochemistry*, 2001, **56**, 857-861 (*8-O-Demethylnornitidine*)Seckárová, P. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 147-152 (*pmr, cmr*)Halstead, C.W. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 940-945 (*Terihanine, Isoterihanine*)Luo, Y. *et al.*, *Tetrahedron*, 2006, **62**, 9131-9134 (*Nornitidine, synth*)**Nitrabirine****N-240**

6',7'-Dihydrospiro[cyclohexane-1,8'-(5'H)-imidazo[1,2-a]pyridin]-2-ol, 9CI [86629-99-4]

C₁₂H₁₈N₂O 206.287Alkaloid from the aerial parts of *Nitraria sibirica* (Zygophyllaceae). Cryst. (MeOH). Mp 184-185°.

Hydrochloride: [86630-00-4]

Cryst. (EtOH). Mp 224-225°.

4'-N-Oxide: **Nitrabirine N-oxide**

[450336-23-9]

C₁₂H₁₈N₂O₂ 222.286Alkaloid from the aerial parts of *Nitraria sibirica*. Cryst. (EtOH). Mp 235-236°. λ_{\max} 213 (log ϵ 3.81) (EtOH). [86630-01-5]

[86630-01-5]

Ibragimov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 213-216; 1988, **24**, 82-91; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 202-


205; 1988, **24**, 71-78 (*isol. struct. pmr, cmr, abs config*)

Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2001, **37**, 556-558 (*N-oxide*)

Nitramine

2-Azaspiro[5.5]undecan-7-ol

N-241



C₁₀H₁₉NO 169.266

Rel. configs. only appear to be known. The enantiomer shown is assigned by CAS.

(+)-form [49620-06-6]

Alkaloid from *Nitraria schoberi* (Zygophyllaceae). Liq. [α]_D²⁰ +16.5 (c, 4.85 in CHCl₃).

Nitrate: [81149-19-1]

Mp 204-206°.

7-Epimer: Isonitramine

[65620-67-9]

C₁₀H₁₉NO 169.266

Alkaloid from *Nitraria sibirica* (Zygophyllaceae). Needles. Mp 101-103° subl. [α]_D²⁵ -30 (c, 1.36 in CHCl₃).

(-)-form

N-Me: Sibirine

[83023-77-2]

C₁₁H₂₁NO 183.293

Alkaloid from the epigeal parts of *Nitraria sibirica* (Zygophyllaceae). [α]_D²⁵ -22.5 (c, 0.81 in CHCl₃).

N-Me; hydrochloride: [83057-62-9]

Cryst. (Me₂CO/EtOH). Mp 191-192°.

(±)-form [82227-98-3]

Alkaloid from *Nitraria sibirica* leaves and stems (Zygophyllaceae).

Hydrochloride: [89398-21-0]

Mp 225-228° (synthetic).

[81149-20-4, 89460-82-2, 98302-88-6, 81149-21-5, 81202-54-2]

Novgorodova, N.Y. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 196; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 191 (*isol*)

Osmanov, Z. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 720; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 607 (*isol. Isonitramine*)

Ibragimov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 623; 1982, **18**, 126; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 458; 1982, **18**, 121 (*cmr, cryst struct*)

Osmanov, Z. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 225; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 206 (*Sibirine*)

Snider, B.B. *et al.*, *J.O.C.*, 1984, **49**, 1688 (*synth*)

Kozikowski, A.P. *et al.*, *Chem. Comm.*, 1985, 847 (*synth*)

Tanner, D. *et al.*, *Tetrahedron*, 1989, **45**, 4309 (*synth*)

Imanishi, T. *et al.*, *Chem. Comm.*, 1991, 1409 (*synth. Sibirine*)

Fujii, T. *et al.*, *Chem. Lett.*, 1992, 1493 (*synth. Isonitramine, Sibirine*)


Keppens, M. *et al.*, *Synlett*, 1994, 285 (*synth. Sibirine*)

Senboku, H. *et al.*, *Heterocycles*, 1997, **46**, 413-420 (*synth*)

Nitraraidine

[619282-56-3]

N-242



C₂₀H₂₅N₂[⊕] 293.431


Quaternary alkaloid from *Nitraria sibirica*. Cryst. (EtOH). Mp 308-310°. Opt. inactive. Counterion not specified. λ_{max} 224 (log ε 4.05); 275 (log ε 3.59); 288 (sh) (log ε 3.47) (EtOH).

Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2002, **38**, 602-604 (*isol, pmr, cmr*)

Nitraraine

N-243

16,17-Didehydroyohimban-17-methanol, 9CI. 16-Hydroxymethyl-yohimb-16-ene [101222-61-1]



C₂₀H₂₄N₂O 308.422

Alkaloid from *Nitraria schoberi*. Mp 280-281°. Racemic. Has been synthesised chirally.

O-Ac: O-Acetylnitraraine

C₂₂H₂₆N₂O₂ 350.46

Alkaloid from *Nitraria schoberi*. Cryst. (MeOH). Mp 93-95°. Racemic. λ_{max} 227 (log ε 4.55); 281 (log ε 4.01); 292 (sh) (log ε 3.91) (EtOH).

16α,17-Dihydro: Dihydranitraraine

[101144-41-6]

C₂₀H₂₆N₂O 310.438

Alkaloid from *Nitraria schoberi*. Cryst. by subl. Mp 286-287°. Racemic. Has been synthesised chirally.

Ibragimov, A.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 502-509; 510-512 (*isol, ir, ms, struct, Dihydranitraraine*)

Takano, S. *et al.*, *Chem. Lett.*, 1989, 1777-1780 (*Nitraraine, Dihydranitraraine, synth*)

Yamaguchi, R. *et al.*, *Synlett*, 1991, 719-720 (*synth*)

Paivio, E. *et al.*, *Heterocycles*, 2000, **53**, 2241-2246 (*Dihydranitraraine, synth*)


Sakagami, H. *et al.*, *Heterocycles*, 2001, **54**, 43-47 (*synth*)

Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2005, **41**, 578-579 (*O-Acetylnitraraine*)

Nitramine

[57912-25-1]

N-244



1380

C₁₅H₂₄N₂O 248.367

Struct. revised in 1985. Alkaloid from epigeal parts of *Nitraria schoberi*. Mp 85-86°. Opt. inactive. Reported *isol.* of 1-Epinitramine has been shown to be incorrect (2005).

N⁵-Oxide: Nitramine N-oxide. Nitroxine

[57912-26-2]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Nitraria schoberi* and *Nitraria komarovii*. Cryst. (EtOH). Mp 251-252° (220-221°). [α]_D 0.

N⁴-(Phenylacetyl), N⁵-oxide: Nitramidine

[619282-52-9]

C₂₃H₃₀N₂O₃ 382.502

Alkaloid from *Nitraria sibirica*. Cryst. (CHCl₃/EtOH). Mp 225-226°. λ_{max} 209 (log ε 4.02); 249 (log ε 2.05) (EtOH).

Novgorodova, N.Yu. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 455 (*isol, pmr, ms*)

Tashkhodzhaev, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 649-655 (*cryst struct*)

Ibragimov, A.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 620 (*Nitroxine*)

Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 31 (*Nitramine N-oxide*)

Wanner, W.J. *et al.*, *J.O.C.*, 1995, **60**, 5634 (*synth, pmr, cmr, ms*)


Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2002, **38**, 602-604 (*Nitramidine*)

Gravel, E. *et al.*, *Org. Lett.*, 2005, **7**, 2497-2499 (*synth*)

Nitraricine

[176181-92-3]

N-245



C₂₀H₂₁N₃O 319.405


Alkaloid from aerial parts of *Nitraria komarovii*. Cryst. (EtOH/Me₂CO). Mp 214-215°. [α]_D 0. λ_{max} 208 (log ε 4.16); 253 (log ε 4.13); 310 (log ε 4.02); 365 (log ε 3.41) (EtOH).

Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1995, **95**; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 76

Nitrarine

[20069-03-8]

N-246



Relative Configuration

C₂₀H₂₅N₃ 307.438

Antispasmodic, antihypertensive agent.
Log P 2.18 (calc).

(±)-form

Alkaloid from *Nitraria schoberi* (Zygophyllaceae). Mp 256-257°. λ_{\max} 226 (log ϵ 4.52); 286 (log ϵ 4.01) (no solvent reported).

Hydrochloride (1:2): Mp 267°.

N¹⁶-Me: N-Methylnitrarine

C₂₁H₂₇N₃ 321.464

Alkaloid from the aerial parts of *Nitraria schoberi*. Cryst. (EtOH). Mp 263-264°. λ_{\max} 220 (log ϵ 4.68); 269 (log ϵ 3.88); 280 (sh) (log ϵ 3.85); 288 (log ϵ 3.52) (EtOH).

N¹⁶-(2-Propenyl): N-Allylnitrarine

C₂₃H₂₉N₃ 347.502

Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (EtOH). Mp 269-271°. λ_{\max} 222 (log ϵ 4.72); 268 (log ϵ 3.85); 280 (log ϵ 3.84); 388 (log ϵ 3.52) (EtOH).

1,3-Didehydro: Nitramidine. 1,3-Didehydronitrarine, 9CI

[56775-82-7]

C₂₀H₂₃N₃ 305.422

Alkaloid from *Nitraria schoberi* (Zygophyllaceae). Noncryst.; yellow cryst. (as hydrochloride). Mp 251-253° (hydrochloride). λ_{\max} 250 (log ϵ 3.95); 263 (log ϵ 4.27) (EtOH).

5,6,15,16-Tetradehydro, N⁴-oxide: Nitrarisine. Nitrazine

[176181-93-4]

C₂₀H₂₁N₃O 319.405

Alkaloid from aerial parts of *Nitraria komarovii*. Cryst. (EtOH/Me₂CO). Mp 265-267°. $[\alpha]_D^{20}$.

3-Epimer: Isonitrarine. 3-Epinitrarine

[57969-03-6]

C₂₀H₂₅N₃ 307.438

Alkaloid from *Nitraria schoberi* (Zygophyllaceae). Cryst. Mp 209°. λ_{\max} 226 (log ϵ 4.46); 286 (log ϵ 3.84) (EtOH).


3-Epimer; hydrochloride: Mp 239°.

Ibragimov, A.A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 273-274; 275; 276; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 293-294; 295-296; 298 (*Nitrarine, Nitramidine, Isonitrarine*)

Nasirov, S.-M. et al., *Khim. Prir. Soedin.*, 1976, **12**, 334-345; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 294-301 (*cryst struct*)

Vakhabov, A. et al., *CA*, 1980, **92**, 209116 (*rev. pharmacol*)

Tulyaganov, T.S. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 95-395; 2000, **36**, 393-395; 396-398 (*Nitrarisine, N-Allylnitrarine, N-Methylnitrarine*)

16,23-Nitroloicycloarta-7,16(N),17(20),22-tetraene-3,11,24,25-tetrol

C₃₀H₄₃NO₄ 481.674

(3 β ,11 β ,24S)-form

3-O- β -D-Xylopyranoside: **Cimicifine A.**

Cimicifugadine

[936924-56-0]

C₃₅H₅₁NO₈ 613.79

Alkaloid from the rhizomes of *Cimicifuga foetida*. Amorph. solid. Mp 246-248° (*Cimicifugadine*) Mp 275-277° (*Cimicifine A*). $[\alpha]_D^{24}$ -12.7 (c, 0.2 in MeOH) (*Cimicifine A*). $[\alpha]_D^{25}$ +13 (c, 0.095 in MeOH) (*Cimicifugadine*). *Cimicifine A* and *Cimicifugadine* assigned same struct., although props. not identical. λ_{\max} 271 (MeOH).

Sun, L.-R. et al., *Helv. Chim. Acta*, 2007, **90**, 1313-1318 (*Cimicifine A*)


Dan, C. et al., *Org. Lett.*, 2007, **9**, 1813-1816 (*Cimicifugadine*)

3,3',3''-[Nitrotris(methylene)] N-248

ne]tris[dihydro-2(3H)furanone], 9CI

Tri(α -methylene- γ -butyrolactonyl)amine

[102779-02-2]



C₁₅H₂₁NO₆ 311.334

Minor base from *Bellendena montana* (Proteaceae). Mp 186-187°. $[\alpha]_D^{19}$ +80 (CHCl₃).

Picrate: Mp 209-211°.

Diastereoisomer (?):

C₁₅H₂₁NO₆ 311.334

Artifact isol. from *Garnieria spathulaefolia* (Proteaceae). Cryst. (MeOH). Mp 184-186° dec.

Bick, I.R.C. et al., *Aust. J. Chem.*, 1979, **32**, 1827 (*isol*)


Lounasmaa, M. et al., *Heterocycles*, 1985, **23**, 939 (*isol, pmr, ir, cmr, ms, struct*)

Bick, I.R.C. et al., *Phytochemistry*, 1986, **25**, 972 (*uv, ir, pmr, cmr, struct*)

3-(12-Nitrododecyl)pyridine N-249

1-Nitro-12-(3-pyridinyl)dodecane. Unteine B

[174624-22-7]



C₁₇H₂₈N₂O₂ 292.42


Alkaloid from the Okinawan marine sponge *Callyspongia* sp. Exhibits potent anti-microfouling activity.

Wang, G.-Y.-S. et al., *Tet. Lett.*, 1996, **37**, 1813 (*isol, pmr, cmr, struct*)

2-Nitroimidazole, 9CI

Azomycin. Amicin

[527-73-1]



C₃H₃N₃O₂ 113.076

Produced by *Nocardia* spp., *Streptomyces eurocidicus* and *Pseudomonas fluorescens*. Antibiotic active against gram-positive organisms. Pale yellow cryst. (EtOH). Sol. MeOH, EtOAc, bases; fairly sol. CHCl₃, Et₂O; poorly sol. H₂O, CHCl₃, Me₂CO, hexane, Et₂O. Mp 282° dec Mp 287-288°. pK_{a1} -0.81; pK_{a2} 7.15 (50% MeOH). Log P -0.09 (calc). λ_{\max} 223 (ϵ 3500); 365 (ϵ 10700) (EtOH/NaOH) (*Derep*). λ_{\max} 219 (ϵ 3900); 315 (ϵ 8600) (95% EtOH) (*Derep*).

▶ LD₅₀ (mus, orl) 316 mg/kg. LD₅₀ (mus, ivn) 80 mg/kg. Mutagenic props. NI7875000

[36877-68-6]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 618B (*ir*)

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 489D (*nmr*)

Nakamura, S. et al., *J. Antibiot., Ser. A*, 1955, **8**, 66 (*isol*)

Beaman, A.G. et al., *J.A.C.S.*, 1965, **87**, 389 (*synth*)

Nakane, A. et al., *J. Biol. Chem.*, 1977, **252**, 5267 (*biosynth*)

Agrawal, K.C. et al., *J. Med. Chem.*, 1979, **22**, 583 (*synth*)

Luijten, W.C. et al., *Org. Mass Spectrom.*, 1982, **17**, 299 (*ms*)

Nair, M.D. et al., *Prog. Drug Res.*, 1983, **27**, 163 (*rev. pharmacol*)

Larsen, I.K. et al., *Acta Cryst. C*, 1984, **40**, 285 (*cryst struct*)


Shoji, T. et al., *J. Antibiot.*, 1989, **42**, 1513 (*isol*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, NHG000

3-Nitro-1H-indole

N-251

[4770-03-0]



C₈H₆N₂O₂ 162.148

Yellow needles (C₆H₆). Mp 213-214°. pK_a 10.12 (25°). Prod. by the marine-derived *Salegentibacter* sp. T436.

N-Ac:

C₁₀H₈N₂O₃ 204.185

Tan cryst. (hexane/CH₂Cl₂). Mp 167-168°.

N-Me: [36728-89-9]

C₉H₈N₂O₂ 176.174

Cryst. (C₆H₆/hexane). Mp 156-157° (154°). Subl. 2 140.

N-Et: [55543-24-3]

C₁₀H₁₀N₂O₂ 190.201

Yellow needles (petrol). Mp 102°.

N-Benzyl:

C₁₅H₁₂N₂O₂ 252.272

Tan cryst. (hexane/CH₂Cl₂). Mp 118-119°.

Noland, W.E. *et al.*, *J.O.C.*, 1966, **31**, 70-77 (synth, uv)

Berti, G. *et al.*, *J.C.S.(C)*, 1968, 2145-2151 (synth)

Pelkey, E.T. *et al.*, *Synthesis*, 1999, 1117-1122 (synth, derivs, ir, pmr, cmr, uv, ms)


Gribble, G.W. *et al.*, *Tetrahedron*, 2000, **56**, 10133-10140 (synth, pmr, cmr, ms)

Al-Zereini, W. *et al.*, *J. Antibiot.*, 2007, **60**, 301-308 (isol)

Nitropolyzonamine

N-252

Hexahydro-2,2-dimethyl-7'-nitrospiro[cyclopentane-1,1'-[1H]pyrrolizine], 9CI. Spiropyrrrolizidine 238 [57308-84-6]



Absolute configuration

C₁₃H₂₂N₂O₂ 238.329

Component of the defence secretion of the millipede *Polyzonium rosalbum*. Mp 65.5-66.5°. [α]_D²⁰ +12 (CHCl₃). Prob. formed from Loline, L-234 *in vivo*.

Meinwald, J. *et al.*, *Tet. Lett.*, 1975, 2367 (isol, ir, ms, cryst struct)

Miller, R.W. *et al.*, *J. Chem. Res., Synop.*, 1978, 76 (cryst struct, abs config)

Mori, K. *et al.*, *Tet. Lett.*, 2000, **41**, 6623-6625 (synth)

Nitrosoxacin A

N-253

N-Hydroxy-14-methyl-*N*-nitrosopentadecanamine, 9CI. *N*-(14-Methylpentadecyl)-*N*-nitrosohydroxylamine [147317-96-2] (H₃C)₂CH(CH₂)₁₃N(OH)NO

C₁₆H₃₄N₂O₂ 286.457

Prod. by *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. DMSO, CHCl₃; fairly sol. MeOH, hexane; poorly sol. H₂O. Mp 44-44.5°. λ_{max} 230 (ε 6300) (MeOH/HCl) (Derep). λ_{max} 249 (ε 9900) (MeOH/NaOH) (Derep).

Nishio, M. *et al.*, *J. Antibiot.*, 1993, **46**, 193 (isol, struct)

Nitrosoxacin B

N-254

N-Hydroxy-*N*-nitrosohexadecanamine, 9CI. *N*-Hexadecyl-*N*-nitrosohydroxylamine [147317-97-3] H₃C(CH₂)₁₅N(OH)NO

C₁₆H₃₄N₂O₂ 286.457

Prod. by *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. DMSO, CHCl₃; fairly sol. MeOH, hexane; poorly sol. H₂O. Mp 38.5-39.5°. λ_{max} 230 (ε 6300) (MeOH/HCl) (Derep). λ_{max} 249 (ε 9900) (MeOH/NaOH) (Derep).

Nishio, M. *et al.*, *J. Antibiot.*, 1993, **46**, 193 (isol, struct)

Nitrosoxacin C

N-255

N-Hydroxy-12-methyl-*N*-nitrosotridecanamine, 9CI. *N*-(12-Methyltridecyl)-*N*-nitrosohydroxylamine [147317-95-1]

(H₃C)₂CH(CH₂)₁₁N(OH)NO

C₁₄H₃₀N₂O₂ 258.403


Prod. by *Streptomyces* sp. 5-Lipoxygenase inhibitor. Powder. Sol. DMSO, CHCl₃; fairly sol. MeOH, hexane; poorly sol. H₂O. Mp 35.5-36°. λ_{max} 230 (ε 6300) (MeOH/HCl) (Derep). λ_{max} 249 (ε 9900) (MeOH/NaOH) (Derep).

Nishio, M. *et al.*, *J. Antibiot.*, 1993, **46**, 193 (isol, struct)

3-(14-Nitro-3-tetradecenyl)-pyridine

N-256

1-Nitro-14-(3-pyridinyl)-11-tetradecene



C₁₉H₃₀N₂O₂ 318.458

(Z)-form

Utenine A

[174756-37-7]

Alkaloid from the Okinawan marine sponge *Callyspongia* sp. Exhibits potent anti-microfouling activity.

3',4'-Didehydro: 3-(14-Nitro-3-tetradecenyl)pyridine. 1-Nitro-14-(3-pyridinyl)-11-tetradecyne. Utenine C [174756-38-8]

C₁₉H₂₈N₂O₂ 316.442


From *Callyspongia* sp. Exhibits potent anti-microfouling activity.

Wang, G.-Y.-S. *et al.*, *Tet. Lett.*, 1996, **37**, 1813 (isol, pmr, cmr, synth, struct)

4-Nitrotryptophan, 9CI

N-257

2-Amino-3-(4-nitro-1H-indol-3-yl)propanoic acid



C₁₁H₁₁N₃O₄ 249.226

(S)-form

L-form

Prod. by *Streptomyces scabies*. Yellowish solid.

N-Ac: *N*-Acetyl-4-nitrotryptophan

[168111-98-6]

C₁₃H₁₃N₃O₅ 291.263

Prod. by *Streptomyces scabies*. Yellow solid.

N-Ac, *Me ester*: [168029-89-8]

Yellow prisms. Mp 205-206°. λ_{max} 344 (ε 3480); 385 (ε 4070) (EtOH).

N-Me: *N*-Methyl-4-nitrotryptophan

[168029-91-2]

C₁₂H₁₃N₃O₄ 263.252

Prod. by *Streptomyces scabies*. Yellow solid.

N-Me, *Me ester*: [168029-90-1]

Yellow solid. λ_{max} 346 (ε 3360); 384 (ε 3980) (EtOH).

(±)-form

Cryst. Mp 200° dec.

Hydrochloride:

Cryst. (EtOH). Mp 233-235°.

Et ester: [84590-30-7]

C₁₃H₁₅N₃O₄ 277.279

Cryst. Mp 96-98°.

N-Ac:

C₁₃H₁₃N₃O₅ 291.263

Cryst. (EtOH). Mp 248-250°.

N-Ac, *Et ester*: [84590-31-8]

C₁₅H₁₇N₃O₅ 319.316

Yellow needles (EtOAc). Mp 162-164°.

Da Settimo, A. *et al.*, *Ann. Chim. (Rome)*, 1962, **52**, 17 (synth)

Endo, Y. *et al.*, *Tetrahedron*, 1986, **42**, 5905 (*Et ester*, synth, ir, pmr)


King, R.R. *et al.*, *Phytochemistry*, 1995, **40**, 41 (isol, *N*-Ac, *N*-Me)

King, R.R. *et al.*, *Phytochemistry*, 2003, **64**, 1091-1096 (isol, pmr)

Nitrotyrasanguinarine

N-258

[123787-75-7]



C₂₈H₂₂N₂O₇ 498.491


Alkaloid from *Hypocoum imberbe*, *Hypocoum pendulum* and *Hypocoum procumbens*. Amorph. Racemic. λ_{max} 232 (log ε 4.34); 281 (log ε 4.31); 324 (log ε 3.91) (MeOH).

Pabuççuoğlu, V. *et al.*, *J. Nat. Prod.*, 1989, **52**, 716-719 (isol, uv, ir, pmr, cmr, ms, struct)

Njaoamine A

N-259

[934220-47-0]



C₄₀H₅₆N₄O₂ 624.908

Alkaloid from a *Reniera* sp. Cytotoxic. Yellowish gum. [α]_D²⁵ +61.2 (c, 0.11 in MeOH). λ_{max} 204; 252; 314 (MeOH).

40-Deoxy: Njaoamine D

[934220-51-6]

C₄₀H₅₆N₄O 608.909Alkaloid from a *Reniera* sp. Yellowish gum. $[\alpha]_D^{25} +51.6$ (c, 0.7 in MeOH). λ_{\max} 204 ; 251 ; 312 (MeOH).**15-Methyl: Njaoamine B**

[934220-49-2]

C₄₁H₅₈N₄O₂ 638.935Alkaloid from a *Reniera* sp. Yellowish gum. $[\alpha]_D^{25} +51.1$ (c, 0.11 in MeOH). λ_{\max} 204 ; 252 ; 312 (MeOH).**41-Demethyl, 40-deoxy: Njaoamine C**

[934220-50-5]

C₃₉H₅₄N₄O 594.882Alkaloid from a *Reniera* sp. Yellowish gum. $[\alpha]_D^{25} +65.7$ (c, 0.6 in MeOH). λ_{\max} 204 ; 252 ; 313 (MeOH).**41-Demethyl, 40-deoxy, 35,36-dihydro, 36,36,37,37-tetradehydro: Njaoamine H**


[954418-81-6]

C₃₉H₅₂N₄O 592.866Alkaloid from a *Neopetrosia* sp. Pale yellow oil. $[\alpha]_D^{21} +20$ (c, 0.8 in MeOH).**41-Demethyl, 9,40-dideoxy, 35,36-dihydro, 36,36,37,37-tetradehydro: Njaoamine G**


[954418-80-5]

C₃₉H₅₂N₄ 576.867Alkaloid from a *Neopetrosia* sp. Cytotoxic. Pale yellow oil. $[\alpha]_D^{21} +22$ (c, 0.8 in MeOH).Sorek, H. *et al.*, *Tet. Lett.*, 2007, **48**, 7691-7694 (*Njaoamines G,H*)Reyes, F. *et al.*, *Tetrahedron*, 2007, **63**, 2432-2438 (*Njaoamines A-D*)**Njaoamine E****N-260**


[934220-52-7]

C₄₁H₅₄N₄O 618.904Alkaloid from a *Reniera* sp. Cytotoxic. Yellowish gum. $[\alpha]_D^{25} +24.7$ (c, 0.1 in MeOH). λ_{\max} 204 ; 252 (MeOH).Reyes, F. *et al.*, *Tetrahedron*, 2007, **63**, 2432-2438 (*isol, pmr, cmr*)**Njaoamine F****N-261**

[934220-53-8]

C₄₃H₅₆N₄O 644.942Alkaloid from a *Reniera* sp. Yellowish gum. $[\alpha]_D^{25} +20.1$ (c, 0.1 in MeOH). λ_{\max} 204 ; 252 (MeOH).Reyes, F. *et al.*, *Tetrahedron*, 2007, **63**, 2432-2438 (*isol, pmr, cmr*)**Nobiline†****N-262***Nobilonine*

[4684-24-6]




Absolute Configuration

C₁₇H₂₇NO₃ 293.405Alkaloid from *Dendrobium nobile*, *Dendrobium hildebrandii* and *Dendrobium findlayanum* (Orchidaceae). Cryst. (hexane). Mp 86.5-88°. $[\alpha]_D^{23} +21$ (c, 0.48 in CHCl₃). pK_a 8.5 (H₂O).*Hydrochloride*: Mp 206-208°.*Picrate*: Mp 172-175°.*Methiodide*: Mp 275-276°.**6-Hydroxy: 6-Hydroxynobiline**


[32562-94-0]

C₁₇H₂₇NO₄ 309.405Alkaloid from *Dendrobium hildebrandii* (Orchidaceae). Needles (C₆H₆). Mp 158-159.5°. $[\alpha]_D^{23} +62$ (c, 0.58 in CHCl₃).Yamamura, S. *et al.*, *Tet. Lett.*, 1964, 79 (*ir, pmr, ms, ord, struct*)Onaka, T. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 745 (*ir, pmr, ord, struct, abs config, synth*)Graneli, I. *et al.*, *Acta Chem. Scand.*, 1970, **24**, 1209 (*isol*)Elander, M. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 717 (*isol, pmr, ir, ms, struct, abs config, deriv*)Suzuki, M. *et al.*, *Chem. Lett.*, 1975, 611 (*synth*)**δ-Nobilonine****N-263**

[40170-44-3]

C₁₇H₂₇NO₃ 293.405Constit. of *Dendrobium nobile*. Cryst. Mp 210-225°. $[\alpha]_D^{23} -76$ (c, 0.45 in MeOH).Behr, D. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 3196-3202; *Acta Chem. Scand., Ser. B*, 1976, **30**, 279-280 (*δ-Nobilonine*)**Nocardimicin I****N-264***BU 2313B. Antibiotic BU 2313B*

[78339-49-8]

C₂₆H₃₃NO₉ 503.548Isol. from *Microtetraspora caesia* and *Nocardia* sp. Active against gram-positive and -negative bacteria, tumours and trichomonads. Mp 160-162°. $[\alpha]_D^{26} -69.9$ (c, 0.3 in MeOH). λ_{\max} 238 (ε 8550); 355 (ε 33600); 359 (sh); 370 (sh) (ε 29200) (EtOH/HCl) (Derep). λ_{\max} 255 (ε 17400); 286 (ε 20700); 295 (sh); 332 (ε 23500) (EtOH/NaOH) (Derep). λ_{\max} 239 (ε 8550); 295 (sh) (ε 11300); 352 (ε 26400); 370 (sh) (ε 21100) (EtOH) (Derep).**8-Alcohol: Nocardimicin II**


[78112-69-3]

C₂₆H₃₅NO₉ 505.564From *Microtetraspora caesia* and *Nocardia* sp. Active against gram-positive and -negative bacteria, tumours and trichomonads. Pale-yellow cryst. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 238 (ε 8550); 355 (ε 33600); 359 (sh) (ε 29200) (EtOH/HCl) (Derep). λ_{\max} 255 (ε 17400); 286 (ε 20700); 295 (sh); 332 (ε 23500) (EtOH/NaOH) (Derep). λ_{\max} 239 (ε 8550); 295 (sh) (ε 11300); 352 (ε 26400); 370 (sh) (ε 21100) (EtOH) (Derep). λ_{\max} 230 (ε 9600); 352 (ε 36300) (EtOH) (Berdy).**N-Me: Antibiotic BU 2313A. BU 2313A**

[69774-86-3]

C₂₇H₃₅NO₉ 517.575From *Microtetraspora caesia*. Active against gram-positive and -negative bacteria and trichomonads. Pale-yellow cryst. Mp 116-118°. $[\alpha]_D^{26} -58$ (c, 0.5 in MeOH). λ_{\max} 238 (ε 8550); 355 (ε 33600); 359 (sh); 370 (sh) (ε 29200) (EtOH/HCl) (Derep). λ_{\max} 255 (ε 17400); 286 (ε 20700); 295 (sh); 332 (ε 23500) (EtOH/NaOH) (Derep). λ_{\max} 239 (ε 8550); 295 (sh) (ε 11300); 352 (ε 26400); 370 (sh) (ε 21100) (EtOH) (Derep).**▶CB9394500**

[63748-09-4, 69774-87-4]

Horvath, G. *et al.*, *J. Antibiot.*, 1979, **32**, 555
Tsukiura, H. *et al.*, *J. Antibiot.*, 1980, **33**, 157; 166; 173; 1491 (*isol, struct, synth, props*)
Brazhnikova, M.G. *et al.*, *Bioorg. Khim.*, 1981, **7**, 298 (*struct*)**Nocardimicin****N-265**

Nocardimicin	A	R = -(CH ₂) ₈ CH ₃
	B	R = -(CH ₂) ₁₀ CH ₃
	D	R = -(CH ₂) ₁₂ CH ₃
	F	R = -(CH ₂) ₁₄ CH ₃

Peptide antibiotic complex. Related to Amamistatin A, A-672 and Mycobactin, M-787. Prod. by *Nocardia* sp. TP-A0674. Siderophores. Muscarinic M3 receptor inhibitors.**Nocardimicin A [863507-42-0]**C₃₇H₅₅N₅O₁₀ 729.869

43700); 272 (€ 44000); 322 (€ 6900); 335 (€ 6700) (MeOH) (Na salt).

Deoxo, 3',4'-dihydro, 3'-hydroxy (3'R,4'S-): Nodulisporic acid B₂

[478796-09-7]

C₄₃H₅₇NO₇ 699.926

Prod. by *Nodulisporium* sp. [α]_D²³ -30 (c, 0.4 in MeOH) (as Na salt). λ_{\max} 264 (€ 47000); 272 (€ 49300); 322 (€ 7200); 335 (€ 7000) (MeOH) (Na salt).

Hensens, O.D. *et al.*, *Tet. Lett.*, 1999, **40**, 5455-5458 (*isol, uv, ir, pmr, cmr*)


Ondeyka, J.G. *et al.*, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 2941-2944 (*deoxo derivs*)

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*biosynth, config*)

Nodulisporic acid A N-269

Antibiotic L 954967. L 954967

[163120-03-4]



C₄₃H₅₃NO₆ 679.895

Prod. by *Nodulisporium* sp. MF5954. Insecticidal agent. Mp 250-255°. [α]_D²³ +13 (c, 0.4 in CHCl₃). λ_{\max} 241 (€ 43900); 265 (€ 49800); 385 (€ 7360) (MeOH).

1'',2''-Dihydro: Nodulisporic acid A₄

[183161-27-5]

C₄₃H₅₅NO₆ 681.911

Prod. by *Nodulisporium* sp. MF6265. [α]_D²³ +5.9 (c, 0.5 in MeOH). λ_{\max} 236; 274; 396 (MeCN aq. / TFA).

Deoxo: Nodulisporic acid B

[478796-07-5]

C₄₃H₅₃NO₅ 665.911

Prod. by a *Nodulisporium* sp. [α]_D²³ -40 (c, 0.4 in MeOH) (as Na salt). λ_{\max} 266 (€ 64700); 273 (€ 69000); 322 (€ 8600); 335 (€ 8000) (MeOH) (Na salt).

24-Deoxy, 23,24-didehydro, 1'',2''-dihydro: A²³-Nodulisporic acid A₄

[766545-96-4]

C₄₃H₅₃NO₅ 663.895

Prod. by *Nodulisporium* sp. mutants. Yellow powder. λ_{\max} 212 (€ 27250); 267 (€ 25415); 316 (sh); 429 (sh) (MeOH).

[315715-92-5 (Nodulisporic acid A Me ester), 478705-56-5 (Nodulisporic acid B Me ester)]

Ondeyka, J.G. *et al.*, *J.A.C.S.*, 1997, **119**, 8809-8816 (*Nodulisporic acid A*)


Ondeyka, J.G. *et al.*, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 2941-2944 (*Nodulisporic acid B*)

Byrne, K.M. *et al.*, *J.A.C.S.*, 2002, **124**, 7055-7060 (*biosynth*)

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*Nodulisporic acids A₄, A²³-A₄*)

Nodulisporic acid C₁ N-270

[204205-72-1]



C₄₃H₅₇NO₆ 683.926

Prod. by *Nodulisporium* sp. MF5954 mutant. Insecticide. Off-white powder (as Na salt). [α]_D²³ -30 (c, 0.4 in MeOH) (Na salt). λ_{\max} 260 (€ 51980); 266 (€ 57300); 316 (€ 6830); 336 (€ 6400) (MeOH) (Na salt).

3',4'-Dihydro, 3'-hydroxy (3'R,4'S-): Nodulisporic acid C₂

[204205-71-0]


C₄₃H₅₉NO₇ 701.942

Prod. by *Nodulisporium* sp. MF5954 mutant. Off-white powder (as Na salt). [α]_D²³ -42.5 (c, 0.4 in MeOH). λ_{\max} 262 (€ 56560); 266 (€ 61100); 316 (€ 7240); 336 (€ 6340) (MeOH) (Na salt).

Ondeyka, J.G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 121-124; 2004, **67**, 1496-1506 (*isol, pmr, cmr, biosynth*)

Nodulisporic acid C N-271

[204205-66-3]



C₄₃H₅₇NO₅ 667.927

Prod. by *Nodulisporium* sp. MF5954 mutant. Insecticide. Off-white powder (as Na salt). [α]_D²³ -42.5 (c, 0.4 in MeOH) (Na salt). λ_{\max} 262 (€ 72400); 267 (€ 78800); 316 (€ 7240); 336 (€ 6460) (MeOH) (Na salt).

24-Deoxy, 23,24-didehydro, 1'',2''-dihydro: A²³-Nodulisporic acid C₄

[766545-97-5]

C₄₃H₅₇NO₄ 651.928


Prod. by a *Nodulisporium* sp. mutant. Yellow powder. [α]_D²³ +26.2 (c, 2.3 in MeOH). λ_{\max} 212; 284; 306 (sh); 420 (sh) (MeCN aq./TFA).

Ondeyka, J.G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 121-124 (*Nodulisporic acid C*)

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*A²³-Nodulisporic acid C₄*)

Nodulisporic acid D₁ N-272

[204205-64-1]



C₃₈H₄₉NO₅ 599.809

Prod. by *Nodulisporium* sp. MF6227. Amorph. solid. [α]_D²³ -3 (c, 2 in MeOH). λ_{\max} 214 (€ 27280); 260 (€ 46150); 312 (€ 6920); 324 (€ 6000) (MeOH).

3',4'-Dihydro, 3'-hydroxy (3'R,4'S-):

Nodulisporic acid D₂

[204205-65-2]

C₃₈H₅₁NO₆ 617.824

Prod. by *Nodulisporium* sp. MF6227. Amorph. solid. [α]_D²³ -15.7 (c, 2.9 in


MeOH). λ_{\max} 210 (€ 18050); 260 (€ 45250); 310 (€ 7400); 336 (sh) (MeOH).

[766545-99-7 (Nodulisporic acid D₁ Me ester), 766546-00-3 (Nodulisporic acid D₂ Me ester)]

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*isol, pmr, cmr*)

Nodulisporic acid D N-273

[204205-60-7]



C₃₈H₄₉NO₄ 583.809

Prod. by *Nodulisporium* sp. MF6227.

Amorph. solid. [α]_D²³ -5.2 (c, 1.5 in MeOH). λ_{\max} 206 (€ 18900); 262 (€ 58380); 316 (sh); 336 (sh) (MeOH).

7-Ketone: Nodulisporic acid D₃

[766545-94-2]

C₃₈H₄₇NO₄ 581.794


Prod. by *Nodulisporium* sp. MF6227.

Amorph. solid. [α]_D²³ -53.8 (c, 0.13 in MeOH). λ_{\max} 208 (€ 14550); 262 (€ 51060); 315 (sh); 330 (sh) (MeOH).

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*isol, pmr, cmr*)

Nodulisporic acid E N-274

[766545-95-3]



C₃₈H₅₃NO₃ 571.842

Prod. by *Nodulisporium* sp. MF6227.


Amorph. solid. [α]_D²³ +11.2 (c, 1.87 in MeOH). λ_{\max} 234 (€ 41540); 280 (€ 9300) (MeOH).

[766546-01-4 (Nodulisporic acid E Me ester)]

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*isol, pmr, cmr*)

Nodulisporic acid F N-275

[204205-61-8]



C₂₈H₃₇NO₃ 435.605

Prod. by *Nodulisporium* sp. MF5954.


Powder. [α]_D²³ +7.2 (c, 1.7 in MeOH). λ_{\max} 230; 282 (MeCN aq./TFA).

Singh, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1496-1506 (*isol, pmr, cmr*)

Smith, A.B. *et al.*, *J.O.C.*, 2007, **72**, 4596-4610 (*synth*)

Nodusmicin, 9CI N-276

U 59761. Antibiotic U 59761
[76265-48-0]
[78207-19-9]



C₂₃H₃₄O₇ 422.517

Cyclic polyketide. Isol. from *Saccharopolyspora hirsuta* and *Nocardia argentinensis*. Active against gram-positive and anaerobic bacteria. Cryst. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O, hexane. Mp 201-206°. [α]_D +121 (c, 0.76 in MeOH).

18-Ac: 18-O-Acetylnodusmicin

[88343-90-2]
C₂₅H₃₆O₈ 464.555

Prod. by *Nocardia argentinensis*. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O.

9-(1H-Pyrrole-2-carboxylate): Nargenicin A₁

CP 47444. Antibiotic CP 47444. U 59760. Antibiotic U 59760
[70695-02-2]
[80210-04-4]

C₂₈H₃₇NO₈ 515.602

Macrolide antibiotic. Prod. by *Nocardia argentinensis* and *Saccharopolyspora hirsuta*. Active against gram-positive bacteria. Amorph. solid. Mp 128-132°. [α]_D +49 (c, 1 in MeOH). λ_{max} 265 (ε 16500) (MeOH) (Derep).

9-(1H-Pyrrole-2-carboxylate), 18-Ac: 18-O-Acetylnargenicin A₁

[74686-27-4]
C₃₀H₃₉NO₉ 557.639

Prod. by *Nocardia argentinensis*. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O.

18-Deoxy, 9-(1H-pyrrole-2-carboxylate): 18-Deoxynargenicin A₁

Antibiotic U 61732
C₂₈H₃₇NO₇ 499.603

Prod. by *Saccharopolyspora hirsuta*. Active against gram-positive bacteria. Solid. [α]_D²⁵ +71 (c, 0.975 in MeOH).

19,20-Dimethoxy, 9-(1H-pyrrole-2-carboxylate): Nargenicin C₁

Antibiotic CP 57820
[90928-02-2]

C₃₀H₄₁NO₁₀ 575.655

From *Nocardia* sp. Active against gram-positive bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{max} 265 (E1%/1cm 291) (MeOH) (Berdy). λ_{max} 265 (E1%/1cm 290) (MeOH/HCl) (Berdy). λ_{max} 266 (E1%/1cm 291) (MeOH/NaOH) (Berdy).

U.S. Pat., 1979, 4 148 883; CA, 91, 54595 (isol)

Tone, J. et al., Intersci. Conf. Antimicrob. Agents Chemoth., 20th, 1980, Abstr. 62

Celmer, W.D. et al., J.A.C.S., 1980, 102, 4203 (Nargenicin, struct, pmr)

Whaley, H.A. et al., Tet. Lett., 1980, 21, 3659 (struct)

Snyder, W.C. et al., J.A.C.S., 1984, 106, 784-787; 787-789 (biosynth, cmr)

U.S. Pat., 1984, 4 436 747; CA, 101, 53332 (isol)

Cane, D.E. et al., J. Antibiot., 1985, 38, 423 (biosynth, abs config)


Ikeda, Y. et al., J. Antibiot., 1985, 38, 436 (isol)

Plata, D.J. et al., J.A.C.S., 1988, 110, 4041-4042 (synth)

Kallmerten, J. et al., Stud. Nat. Prod. Chem., 1995, 17, 283 (rev)

Noelaquinone

[220503-29-7]



C₂₁H₁₅N₃O₅ 389.367

Alkaloid from a sponge *Xestospongia* sp. Yellow solid. [α]_D²³ +53.6 (c, 0.25 in MeOH). Slow dec. >300°. λ_{max} 206 (ε 45000); 215 (ε 16660); 254 (ε 10620) (MeOH).

Zhu, Y. et al., Heterocycles, 1998, 49, 355-360

Noformicin

N-278

5-Amino-N-(3-amino-3-iminopropyl)-3,4-dihydro-2H-pyrrole-2-carboxamide, 9CI. N-(2-Amidinoethyl)-5-imino-2-pyrrolidonecarboxamide, 8CI. MK 61. Antibiotic MK 61

[155-38-4]



C₈H₁₅N₅O 197.239

Pyrrole antibiotic.

(+)-form

Produced by *Nocardia formica*. Shows antiviral and antibacterial props. Sol. MeOH, H₂O; poorly sol. Me₂CO, hexane. pK_a 9.4 (25°, H₂O).

► Toxic, LD₅₀ (mus, ivn) 5 mg/kg; LD₅₀ (mus, scu) 10 mg/kg.

Hydrochloride (1:2): [42373-21-7]

Cryst. (MeOH). Sol. H₂O. Mp 265° dec. [α]_D²⁵ +7 (c, 1 in H₂O). [α]_D²⁵ +8.8 (c, 1 in MeOH).

(±)-form

Hydrochloride (1:2): [42373-18-2]

Mp 253-254°.

Harris, D.A. et al., Antibiot. Annu., 1953, 609 (isol)


Gray, R.A. et al., Phytochemistry, 1955, 45, 281 (isol)

Diana, G.D. et al., J. Med. Chem., 1973, 16, 857 (synth)

Nominine†

N-279

[120991-21-1]



Relative configuration

C₂₈H₃₉NO 405.622

Isol. from the sclerotia of *Aspergillus nomius*, *Aspergillus flavus* NRRL 13462 and *Petromyces alliaceus*. Has insecticidal props. Off-white powder. Mp 54-55°. [α]_D +23.6 (c, 0.85 in MeOH). Biogenetically related to the Aflavinines. λ_{max} 224 (ε 20000); 251 (ε 3600); 285 (ε 3380) (MeOH) (Derep).

Gloer, J.B. et al., J.O.C., 1989, 54, 2530 (isol, pmr, cmr, ms, struct)

2,4-Nonadien-8-ynoic acid

N-280

HC≡CCH₂CH₂CH=CHCH=CHCOOH

C₉H₁₀O₂ 150.177

(2E,4E)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2,4-nonadien-8-ynamide. 2,4-Nonadien-8-ynoic acid isobutylamide

[18679-26-0]


C₁₃H₁₉NO 205.299

Constit. of the dried leaves of *Acmella caulirhiza*.

Crouch, N.R. et al., S. Afr. J. Bot., 2005, 71, 228-230 (isol)

1,2,3,4,6,8,9,12,14-Nona-hydroxydihydro-β-agarofuran

N-281



C₁₅H₂₆O₁₀ 366.364

(1α,2α,3β,4β,6β,8α,9α,11α)-form

Euonyminol

[33376-82-8]

Formed by hydrol. of alkaloids from *Tripterygium wilfordii*. Cryst. (MeOH/Me₂CO). Mp 250°.

8-Ketone, 12-(3-pyridinecarbonyl), 3-(2-methylbutanoyl), 1,2,6,9,14-penta-Ac: [339176-20-4]

C₃₆H₄₅NO₁₇ 763.748

Constit. of *Tripterygium wilfordii*.

Amorph. powder. [α]_D²⁵ +14.1 (c, 0.3 in

(MeOH). λ_{\max} 214 (log ϵ 4.31); 263 (log ϵ 3.71) (MeOH).

Duan, H.-Q. *et al.*, *Phytochemistry*, 2001, **56**, 341-346 (*Tripterygium wilfordii constii*)

2-Nonene-6,8-diynoic acid, 9CI N-282

$\text{HC}\equiv\text{CC}\equiv\text{CCH}_2\text{CH}_2\text{CH}=\text{CHCOOH}$
 $\text{C}_9\text{H}_8\text{O}_2$ 148.161

(E)-form

2-Phenylethylamide: N-(2-Phenylethyl)-2-nonene-6,8-diynamide
 [99615-84-6]
 $\text{C}_{17}\text{H}_{17}\text{NO}$ 251.327
 Isol. from *Acmella ciliata*.

(Z)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2-nonene-6,8-diynamide. 2-Nonene-6,8-diynoic acid isobutylamide
 [96602-65-2]
 $\text{C}_{13}\text{H}_{17}\text{NO}$ 203.283
 Isol. from roots of *Spilanthes oleracea*.
 Oil.

2-Phenylethylamide: [75872-74-1]
 $\text{C}_{17}\text{H}_{17}\text{NO}$ 251.327
 Isol. from *Spilanthes alba* and *Salmea scandens*. Oil. (E-) and (Z)-forms characterised from *S. alba*.

[75872-75-2, 75872-77-4]

Bohlmann, F. *et al.*, *Phytochemistry*, 1980, **19**, 1535; 1985, **24**, 595 (*isol, struct*)

Greger, H. *et al.*, *Monatsh. Chem.*, 1985, **116**, 273 (*isol, struct, uv, ir, cmr, pmr*)

Martin, R. *et al.*, *Phytochemistry*, 1985, **24**, 2295 (*isol, pmr*)

2-Nonene-4,6,8-triynoic acid, 9CI N-283

$\text{HC}\equiv\text{CC}\equiv\text{CC}\equiv\text{CCH}=\text{CHCOOH}$
 $\text{C}_9\text{H}_4\text{O}_2$ 144.129

(E)-form [38152-85-1]

Metab. of *Poria sinuosa*, *Psilocybe sarcocephala* and *Lepista diemii*. Pale yellow cryst. ($\text{CH}_2\text{Cl}_2/\text{CCl}_4$). Dec. at 70° with ignition.

Me ester: [54981-65-6]

$\text{C}_{10}\text{H}_6\text{O}_2$ 158.156

Isol. from *Lepista glaucocana*. Pale yellow needles (hexane) or cryst. (petrol). Sol. C_6H_6 , Et_2O . Mp 20° (dec. at 80° with ignition).

4-Carboxyanilide: 4-[(2-Nonene-4,6,8-triynoyl)amino]benzoic acid

$\text{C}_{16}\text{H}_9\text{NO}_3$ 263.252

Prod. by *Baeospora myosura*. Antibacterial agent. λ_{\max} 248; 261; 325; 347 (MeOH).

Jones, E.R.H. *et al.*, *Proc. Chem. Soc., London*, 1960, 180

Bohlmann, F. *et al.*, *Chem. Ber.*, 1963, **96**, 2586 (*synth, ester*)

Cambie, R.C. *et al.*, *J.C.S.*, 1963, 2056 (*isol, ester, ir, uv*)


Thaller, V. *et al.*, *J.C.S. Perkin 1*, 1972, 2032 (*isol*)

Hearn, M.T. *et al.*, *J.C.S. Perkin 1*, 1974, 2335 (*isol*)

Parish, C.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1900-1902 (*carboxyanilide*)

3-(8-Nonenyl)-5-(1-propenyl)pyrrolizidine N-284

[102492-05-7]




$\text{C}_{19}\text{H}_{33}\text{N}$ 275.476

Venom from the ant *Chelaner antarcticus*.

Jones, T.H. *et al.*, *J.O.C.*, 1986, **51**, 2712 (*isol, pmr, ms, struct, synth*)

Nonylprodigiosin N-285

4-Methoxy-3,24,25-triazatetracyclo[18.2.1.1^{2,5}.1^{7,10}]pentacos-2(25),3,5,7,9,20,22-heptaene, 9CI. Cyclononylprodigiosin. Cyclononylprodigine
 [34852-35-2]



$\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}$ 363.502

Pyrrole antibiotic. Isol. from *Actinomadura madurae* (*Nocardia madurae*). Active against *Staphylococcus aureus*, fungi and protozoa. Deep red solid (as hydrochloride). λ_{\max} 551 (CHCl_3) (Berdy). λ_{\max} 539 (EtOH/HCl) (Berdy). λ_{\max} 549 (hexane) (Berdy).

[52288-99-0, 249575-95-9]

Gerber, N.N. *et al.*, *Tet. Lett.*, 1970, 809 (*isol, pmr, ms*)

Gerber, N.N. *et al.*, *J. Antibiot.*, 1971, **24**, 636 (*isol, ir*)


Gerber, N.N. *et al.*, *J. Het. Chem.*, 1973, **10**, 925 (*isol, uv, ms*)

Fürstner, A. *et al.*, *ChemBioChem*, 2001, **1**, 60-68 (*synth, activity*)

Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (*rev*)

3-Nonyl-1H-pyrazole, 9CI N-286

[72738-01-3]



$\text{C}_{12}\text{H}_{22}\text{N}_2$ 194.319

Alkaloid from *Houttuynia cordata* (Yu Xing Cao). Antifungal agent. Cryst. Sol. Et_2O . Mp 47.5-49.5°. Bp₂₂ 195° Bp₉ 181°.


Kosuge, T. *et al.*, *Yakugaku Zasshi*, 1952, **72**, 1227-1231; 1954, **74**, 819-820 (*synth*)

Kosuge, T. *et al.*, *J. Biochem. (Tokyo)*, 1954, **41**, 183-186 (*isol*)

U.S. Pat., 1959, 2 883 392; CA, **53**, 16155e (*synth*)

Noractinidine N-287

6,7-Dihydro-7-methyl-5H-2-pyridine, 9CI



$\text{C}_9\text{H}_{11}\text{N}$ 133.193

(S)-form [22324-97-6]

Alkaloid from *Pedicularis macrochila* (Scrophulariaceae) and *Tecoma stans* (Bignoniaceae). Oil. $[\alpha]_D^{24}$ +3 (c, 2.34 in CHCl_3). $[\alpha]_D$ +22 (EtOH).


Picrate:

Cryst. (MeOH). Mp 116-117° Mp 135-137° (dimorph).

Dickinson, E.M. *et al.*, *Tetrahedron*, 1969, **25**, 1523

Norbelladine N-288

4-[[[2-(4-Hydroxyphenyl)ethyl]amino]methyl]-1,2-benzenediol, 9CI. 4-[[[p-Hydroxyphenethyl]amino]methyl]pyrocatechol, 8CI. N-(3,4-Dihydroxybenzyl)-tyramine
 [6053-00-5]



$\text{C}_{15}\text{H}_{17}\text{NO}_3$ 259.304

Biosynth. precursor of Amaryllidaceae alkaloids. Has not been reported as an isol. alkaloid.

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 175-176°.

Picrate:

Cryst. (H₂O). Mp 102-117°.

4'-Me ether: **O-Methylnorbelladine**

[4579-60-6]

$\text{C}_{16}\text{H}_{19}\text{NO}_3$ 273.331

Alkaloid from *Crinum asiaticum* and *Crinum augustum*, also obt. by reduction of Craugsodine, C-730. Biosynth. intermed. for Amaryllidaceae alkaloids. Mp 161-164°.

4'-Me ether, N-Me: **O,N-Dimethylnorbelladine**

$\text{C}_{17}\text{H}_{21}\text{NO}_3$ 287.358

Alkaloid from bulbs of *Pancreatium maritimum* (Amaryllidaceae). Mp 130-132°. Incorrectly indexed in CA.

3',4'-Di-Me ether: **Latisodine**. 4-[[[3,4-Dimethoxyphenyl]methyl]amino]ethylphenol, 9CI
 [88205-07-6]

$\text{C}_{17}\text{H}_{21}\text{NO}_3$ 287.358

Alkaloid from the flower-stem exudate

of *Crinum latifolium*. Mp 205-207°. λ_{\max} 220 (log ϵ 4.02); 277 (log ϵ 3.71); 288 (sh) (log ϵ 3.65) (MeOH).

3',4'-Di-Me ether, O''- β -D-glucopyranoside: **Latisoline**
[88205-08-7]
C₂₃H₃₁NO₈ 449.5
Alkaloid from the flower-stem exudate of *Crinum latifolium*. Amorph. solid. $[\alpha]_D^{22}$ -48.5 (c, 0.52 in MeOH). λ_{\max} 275 (log ϵ 3.69); 280 (log ϵ 3.61) (MeOH).

Tri-Me ether: **N-Demethylbelladine**
C₁₈H₂₃NO₃ 301.385
Alkaloid from the bulbs of *Nerine filifolia*. Light brown oil. λ_{\max} 220 (log ϵ 14.63); 240 (log ϵ 5.95); 260 (log ϵ 3.82); 280 (log ϵ 5.11) (MeOH).

O,O,O,N-Tetra-Me: N-[(3,4-Dimethoxyphenyl)methyl]-4-methoxy-N-methylbenzeneethanamine, 9CI. (3,4-Dimethoxybenzyl) (4-methoxyphenethyl)methylamine. **Belladine**
[501-06-4]
C₁₉H₂₅NO₃ 315.411
Alkaloid widespread in the family Amaryllidaceae. Liq. Mp 128-129.5° (as perchlorate).

3''-Methoxy, tri-Me ether: **Ryllistine**
[24997-85-1]
C₁₉H₂₅NO₄ 331.411
Alkaloid from the flower-stem fluid of *Amaryllis vittata*. Mp 86-88° (79-80°). λ_{\max} 234 (log ϵ 4.3); 280 (log ϵ 3.77); 285 (sh) (log ϵ 3.68) (MeOH).

Surrey, A.R. *et al.*, *J.A.C.S.*, 1949, **71**, 2421-2422 (synth)

Warnhoff, E.W. *et al.*, *Chem. Ind. (London)*, 1957, 1385-1386 (*Belladine*, *uv*, *ir*, *struct*)

Barton, D.H.R. *et al.*, *J.C.S.*, 1963, 4545-4558 (*Norbelladine*, *synth*)

Battersby, A.R. *et al.*, *J.C.S.*, 1964, 1595-1609 (*Norbelladine*, *synth*)

Zetta, L. *et al.*, *J.C.S. Perkin 2*, 1973, 1180-1184 (*Belladine*, *cmr*)

Fuganti, C. *et al.*, *Alkaloids (Academic Press)*, 1975, **15**, 146-147 (*biochem*, *rev*)

Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1983, 238-239; 1984, 412-413 (*Latisoline*, *Ryllistine*)


Ghosal, S. *et al.*, *Phytochemistry*, 1985, **24**, 2141-2156 (*O-Methylnorbelladine*)

Vázquez Tato, M.P. *et al.*, *Heterocycles*, 1988, **27**, 2833-2838 (*O,N-Dimethylnorbelladine*)

Eichhorn, J. *et al.*, *Phytochemistry*, 1998, **49**, 1037-1047 (*O-Methylnorbelladine*, *synth*, *biochem*, *bibl*)

Nair, J.J. *et al.*, *Phytochemistry*, 2005, **66**, 373-382 (*Belladine*, *N-Demethylbelladine*)

Norcassaidide **N-289**
[52579-69-8]



C₂₃H₃₉NO₄ 393.565
Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Needles (MeOH/Et₂O). Mp 244°. $[\alpha]_D^{22}$ -61 (c, 0.9 in EtOH).


3-Ac: **Norcassaidide 3-acetate**
C₂₅H₄₁NO₅ 435.603
Alkaloid from *Erythrophleum guineense* (Fabaceae).

7-Ketone: **Norcassaidide**. *Cassaide*
[35314-33-1]
C₂₃H₃₇NO₄ 391.55
Alkaloid from the bark of *Erythrophleum ivorense* (Fabaceae). Cryst. (CHCl₃/hexane). Mp 205-206°. The name Norcassaidide is preferred since the alkaloid contains 1C less than the isomeric Cassaine.

7-Ketone, 3-O-(3-hydroxy-3-methylbutanoyl): **Norcoumingide**
[59035-77-7]
C₂₈H₄₅NO₆ 491.667
Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Cryst. (Et₂O/petrol) (as acetate). Mp 114-115° (acetate).


6-Oxo: **Norerythrosuaveolide**
C₂₃H₃₇NO₅ 407.549
Alkaloid from the stem bark of *Erythrophleum suaveolens*.
Cronlund, A. *et al.*, *Acta Pharm. Suec.*, 1971, **8**, 351 (*Norcassaidide*)
Cronlund, A. *et al.*, *Planta Med.*, 1973, **24**, 371 (*Norcassaidide 3-acetate*)
Loder, J.W. *et al.*, *Aust. J. Chem.*, 1974, **27**, 179 (*Norcassaidide*)
Cronlund, A. *et al.*, *Acta Pharm. Suec.*, 1975, **12**, 467 (*Norcoumingide*)
Ngounou, F.N. *et al.*, *Bull. Chem. Soc. Ethiop.*, 2005, **19**, 221-226 (*Norerythrosuaveolide*)

Norchalciporol **N-290**




C₁₃H₁₇NO 203.283
O-Propanoyl: **Norchalciporol propionate**
[112448-75-6]
C₁₆H₂₁NO₂ 259.347
Pungent principle from fruit-bodies of the edible pfefferröhrling mushroom (*Chalciporus piperatus*). Sl. yellow oil. Sol. MeOH, Et₂O; poorly sol. H₂O. $[\alpha]_D^{22}$ -290 (c, 0.2 in Et₂O). λ_{\max} 246 (ϵ 15900) (EtOH) (Berdy).
Sternier, O. *et al.*, *Tetrahedron*, 1987, **43**, 1075 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Norcoronatine **N-291**
[98043-42-6]



C₁₇H₂₃NO₄ 305.373
Prod. by *Pseudomonas syringae* pv. *glycinea*. Phytotoxin. Chlorosis inducer.
Mitchell, R.E. *et al.*, *Phytochemistry*, 1985, **24**, 1485-1487 (*isol*, *pmr*, *ms*)


Norcotarnine **N-292**
7,8-Dihydro-4-methoxy-1,3-dioxolo[4,5-g]isoquinoline, 9CI. 3,4-Dihydro-8-methoxy-6,7-methylenedioxyisoquinoline
[484-30-0]



C₁₁H₁₁NO₃ 205.213
Yellow flocks (petrol). Mp 91-92°. *Picrate*:
Yellow needles (EtOH). Mp 182-184°. *N-Me*: **Cotarnine**†
[20276-45-3]
C₁₂H₁₄NO₃ 220.247
Alkaloid from *Papaver pseudo-orientale* (Papaveraceae). Mp 184-186° dec. (as iodide).


[30936-27-7]
Decker, H. *et al.*, *Annalen*, 1913, **395**, 328
Beke, D. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1958, **16**, 439; *CA*, **53**, 5267b
Schneider, W. *et al.*, *Annalen*, 1958, **615**, 34
Korbonits, D. *et al.*, *Chem. Ber.*, 1966, **99**, 273; 1970, **103**, 3605
Fleischhacker, W. *et al.*, *Monatsh. Chem.*, 1989, **120**, 765 (*synth*, *Cotarnine*)
Sariyar, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1302 (*isol*, *pmr*, *struct*)
Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 519

19-Nor-4-dehydrocassaidine **N-293**
[61017-49-0]



C₂₃H₃₇NO₄ 391.55
Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Needles (petrol). Mp 116°.
Oguakwa, J.U. *et al.*, *J. Nat. Prod.*, 1976, **39**, 248 (*isol*, *ir*, *uv*, *pmr*, *ms*, *struct*)


4-Nor-7,8-dehydro-10-hydroxyskytanthine **N-294**




Relative Configuration

C₁₀H₁₇NO 167.25Alkaloid from the roots of *Argyria radiata*. Amorph. solid.Bianco, A. *et al.*, *Nat. Prod. Lett.*, 2002, **16**, 77-80 (*isol, pmr, ms*)**13-Nor-7(11)-dehydropilocarpine** N-295

3-Methyl-4-[(1-methyl-1H-imidazol-5-yl)methyl]-2(5H)-furanone, 9CI. 7(11)-Dehydro-13-norpilocarpine [177857-31-7]


C₁₀H₁₂N₂O₂ 192.217Alkaloid from *Pilocarpus microphyllus* and *Pilocarpus trachylophus*. Mp 156.1-156.3° (as nitrate).Andrade-Neto, M. *et al.*, *Phytochemistry*, 1996, **42**, 885-887 (*isol, pmr, cmr, ms, struct*)Abreu, I.N. *et al.*, *Rapid Commun. Mass Spectrom.*, 2007, **21**, 1205-1213 (*isol, ms*)**Nordine**

N-296


C₄₆H₄₆N₄O₄ 718.894Red pigment from the bark of the Mexican tree Sangre de Drago (prob. *Croton lechleri*). Red solid. Mp 292°. Struct. not well founded.Pallares, E.S. *et al.*, *Arch. Biochem.*, 1946, **10**, 235 (*isol, struct*)**Nordomoic acid**

N-297

2-Carboxy-4-(5-carboxy-1-methyl-1,3-pentadienyl)pyrrolidineacetic acid, 9CI [101899-43-8]

C₁₄H₁₉NO₆ 297.307Constit. of the red alga *Chondria armata*. Insecticidal agent.Maeda, M. *et al.*, *CA*, 1986, **104**, 183260t (*isol*)**Norepinephrine, INN**

N-298

4-(2-Amino-1-hydroxyethyl)-1,2-benzenediol, 9CI. α -(Aminomethyl)-3,4-dihydroxybenzyl alcohol, 8CI. 2-Amino-1-(3,4-dihydroxyphenyl)ethanol. 4-(β -Amino- α -hydroxyethyl)catechol. **Noradrenaline**, **BAN**. Arterenol

(R)-form

C₈H₁₁NO₃ 169.18

Log P -0.99 (calc).

(R)-form

Levarterenol. Adrenor. Levophed. Norartrinal. Nor-Epirenan

[51-41-2] α , β -1-Adrenoceptor agonist, bronchodilator. Sympathomimetic. Vasoconstrictor. Sol. H₂O. Mp 216.5-218°. [α]_D²⁵ -37.3. pK_{a1} 8.6; pK_{a2} 9.8; pK_{a3} 12 (25°). Pharmacol. active isomer.▶ LD₅₀ (mus, orl) 20 mg/kg; LD₅₀ (mus, ivn) 0.55 mg/kg. DN5950000

Hydrochloride: [329-56-6]

Cryst. (2-propanol). Mp 145-146°. [α]_D²⁵ -40.Tartrate (1:1): **Norepinephrine bitartrate**,**USAN**

[51-40-1]

▶ Causes tissue necrosis if administered scu or ims. LD₅₀ (rat, ivn) 0.21 mg/kg. DN6750000N-Ac: **N-Acetylnoradrenaline**

[30959-88-7]

[67083-59-4, 34649-26-8]

C₁₀H₁₃NO₄ 211.217

Found in insect cuticle. Biol. precursor of sclerotised cuticle. Mp 77-79°.

N-(3-Aminopropanoyl): **N- β -Alanyl noradrenaline**

[68299-73-0]

C₁₁H₁₆N₂O₄ 240.258

Degradn. prod. of Papiliochrome II. Found in insect cuticle. Biol. precursor of sclerotised cuticle. Mp 134-137°.

N-Me: see Adrenaline, A-152

3'-Me ether: see Normetanephrine, N-303

(S)-form [149-95-1]Mp 215-217° dec. [α]_D²⁵ +37.4.

▶ DN6125000

N-(4-Methoxy-E-cinnamoyl):

C₁₈H₁₉NO₅ 329.352Alkaloid from *Salsola foetida*. Tyrosinase inhibitor and antioxidant. Amorph. yellow powder. Mp 218-220°. [α]_D²⁵ -16 (c, 0.1 in MeOH). λ _{max} 217 (log ϵ 4.85); 324 (log ϵ 4.51); 327 (log ϵ 4.75) (MeOH).

N-(3,4-Dimethoxy-E-cinnamoyl):

C₁₉H₂₁NO₆ 359.378Constit. of *Salsola foetida*. Tyrosinase inhibitor and antioxidant. Amorph. yellow powder. Mp 225-227°. [α]_D²⁵ -24 (c, 0.06 in MeOH). λ _{max} 219 (log ϵ 4.6); 324 (log ϵ 4.29); 327 (log ϵ 4.35) (MeOH).

4'-Me ether, N-(4-methoxy-E-cinnamoyl):

C₁₉H₂₁NO₅ 343.379Constit. of *Salsola foetida*. Tyrosinase inhibitor and antioxidant. Amorph. yellow powder. Mp 255-257°. [α]_D²⁵ -20 (c, 0.04 in MeOH). λ _{max} 204 (log ϵ 4.56); 319 (log ϵ 4.48); 327 (log ϵ 4.53) (MeOH).**(±)-form** [138-65-8]Sl. sol. H₂O. Mp 191° dec.

▶ DN6300000

Hydrochloride: [55-27-6]


Mp 191°.

▶ LD₅₀ (mus, ipr) 15.6 mg/kg. Exp. reprod. effects. DN6650000**(ξ)-form**Alkaloid in *Albizia julibrissin*, *Mimosa pudica*, *Phaseolus multiflorus*, *Samanea saman* (preferred genus name *Albizia*), *Musa paradisiaca*, *Passiflora quadrangularis*, *Portulaca oleracea*, *Prunus domestica*, *Citrus sinensis*, *Aconitum napellus*, *Aconitum paniculatum* and *Solanum tuberosum* (Fabaceae, Musaceae, Passifloraceae, Portulacaceae, Rosaceae, Rutaceae, Ranunculaceae, Solanaceae). [69815-49-2, 5794-08-1]Biel, J.H. *et al.*, *J.A.C.S.*, 1954, **76**, 3149-3153 (*synth, pharmacol*)Waalke, T.P. *et al.*, *Science (Washington, D.C.)*, 1958, **127**, 648-650 (*isol*)Udenfriend, S. *et al.*, *Arch. Biochem. Biophys.*, 1959, **85**, 487-490 (*detn, occur*)Pratesi, P. *et al.*, *J.C.S.*, 1959, 4062-4065 (*abs config*)Levy, B. *et al.*, *Drill's Pharmacol. Med.*, 4th edn., McGraw-Hill, New York, 1971, 627 (*rev, pharmacol*)Karlson, P. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1972, **327**, 86-94 (*isol, synth, N-Ac*)Applewhite, P.B. *et al.*, *Phytochemistry*, 1973, **12**, 191-192 (*occur*)Donike, M. *et al.*, *Chromatographia*, 1974, **7**, 651-654 (*ms*)Andersen, A.M. *et al.*, *Acta Chem. Scand., Ser. B*, 1975, **29**, 871-876 (*cryst struct*)Rajan, K.S. *et al.*, *Bioinorg. Chem.*, 1976, **6**, 93-117 (*uv*)Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9-18 (*occur, bibl*)Rembold, H. *et al.*, *Z. Naturforsch., C*, 1978, **33**, 498-503 (*N-3-aminopropanoyl*)Wilson, T.D. *et al.*, *Anal. Profiles Drug Subst.*, 1982, **11**, 555 (*rev*)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1249Khan, K.M. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 457-464 (*Salsola foetida cinnamates*)Nagy, P.I. *et al.*, *J.A.C.S.*, 2003, **125**, 2770-2785 (*conformn*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ARL500; NNP050; NNO500; NNO699; ARL750

Norerythrostachamide

N-299

[52579-70-1]

C₂₄H₃₉NO₆ 437.575

Alkaloid from the bark of *Erythrophleum chlorostachys* and *Erythrophleum africanum* (Fabaceae). Rearrangement product of Norerythrostachamine, N-300. Cryst. (MeOH/Et₂O). Mp 207°. [α]_D²⁴ -31 (c, 0.6 in EtOH).

7-Ketone: Norerythrophlamide. *Erythrophlamide* (obsol.)

[35314-34-2]

C₂₄H₃₇NO₆ 435.559

Alkaloid from the bark of *Erythrophleum africanum*, *Erythrophleum chlorostachys*, *Erythrophleum ivorense* and *Erythrophleum suaveolens* (Fabaceae). Rearrangement product of Norerythrophlamine in N-300. Cryst. (CHCl₃/hexane). Mp 163-164°. The name Norerythrophlamide is preferred since it contains 1C less than the isomer Erythrophlamine.

3-Deoxy: Norcassamidide. *Norcassamidide* (obsol.)†

[40445-00-9]

C₂₄H₃₉NO₅ 421.576

Alkaloid from the bark of *Erythrophleum guineense* and *Erythrophleum chlorostachys* (Fabaceae). Rearrangement product of Norcassamidine in N-300. Cryst. (EtOH/Et₂O). Mp 115°. [α]_D²³ -18.5 (c, 0.3 in EtOH). Originally named Norcassamidine based on an incorrect isomeric (ester) struct. assignment. The genuine Norcassamidine was later isol.

3-Deoxy, 7-ketone: Norcassamide. *Norcassamide* (obsol.). *Cassamide* (obsol.)

[35314-35-3]

C₂₄H₃₇NO₅ 419.56

Alkaloid from the bark of *Erythrophleum ivorense* and *Erythrophleum guineense* (Fabaceae). Cryst. (CHCl₃/hexane). Mp 134°. Originally erroneously thought to be an ester and named Norcassamine. Name then changed to Cassamide but Norcassamide is now preferred since it contains 1C less than Cassamine.

6-Oxo: 3β-Hydroxynorerythrosuamide

[58189-27-8]

C₂₄H₃₇NO₇ 451.559

Alkaloid from the bark of *Erythrophleum couminga* (Fabaceae). Cryst. (Et₂O/petrol) (as tri-Ac). Mp 171-172° (tri-Ac).

6-Oxo, 3-deoxy: Norerythrosuamide.

Norerythrosuamide (obsol.)

[36150-74-0]

C₂₄H₃₇NO₆ 435.559

Minor alkaloid from the bark of *Erythrophleum guineense* (Fabaceae). Amorph. [α]_D²⁰ -71.9 (c, 0.44 in EtOH). Originally, Norerythrosuamide and its dehydro deriv. were erroneously assigned ester structs. and named Norerythrosuamide and Dehydronorerythrosuamide.

6-Oxo, 3-deoxy, 7-ketone: Dehydronorerythrosuamide. *Dehydronorerythrosuamide*

[41758-73-0]

C₂₄H₃₅NO₆ 433.544

Minor alkaloid from the bark of *Erythrophleum guineense* (Fabaceae). Amorph. [α]_D²⁰ -41 (c, 0.3 in EtOH).

Cronlund, A. et al., *Acta Pharm. Suec.*, 1971, **8**, 351-360; 1975, **12**, 467-478; *CA*, **76**, 11995y; **84**, 135892x (*Norcassamide*,

Norerythrophlamide, 3-
Hydroxynorerythrosuamide)

Friedrich-Fiechtl, J. et al., *Chem. Ber.*, 1971,

104, 3535-3548 (*Norcassamidide*,
Norerythrosuamide,
Dehydronorerythrosuamide, *Norcassamide*)

Loder, J.W. et al., *Tet. Lett.*, 1972, 3645-3646;

5069-5072 (struct, nomencl)

Loder, J.W. et al., *Aust. J. Chem.*, 1974, **27**,

179-185 (*Norcassamidide*,
Norerythrophlamide, *Norerythrostachamide*)

Falkiner, M.J. et al., *Aust. J. Chem.*, 1975, **28**,

645-650 (*Norcassamidide*, isol, pmr)

Jansson, S. et al., *Acta Pharm. Suec.*, 1976, **13**,

51-54; *CA*, **85**, 13671z (*Norerythrophlamide*,
Norerythrostachamide, isol, ir)


Cronlund, A. et al., *Planta Med.*, 1976, **29**,

123-128 (*Norerythrophlamide*, occur)

Norerythrostachamine

N-300

[52579-68-7]

C₂₄H₃₉NO₆ 437.575

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Antineoplastic agent. Glass. Log P 2.4 (calc). Rearranges to the isomeric Norerythrostachamide, N-299 on standing or after chromatography on alumina.

3-Ac: Norerythrostachamine 3-acetate

[55394-76-8]

C₂₆H₄₁NO₇ 479.612

Tentatively identified as a nat. product from *Erythrophleum chlorostachys* (Fabaceae).

19-Aldehyde: Norerythrostachaldine. 19-

Oxonorcassaldine

[55729-25-4]

[55394-74-6 (hydrochloride)]

C₂₃H₃₇NO₅ 407.549

Alkaloid from the bark of *Erythrophleum chlorostachys*

(Fabaceae). Strongly cytotoxic to KB cells. Gum; prisms (2-propanol/Me₂CO)(as hydrochloride). Mp 156° (as hydrochloride). [α]_D¹⁶ -93 (c, 1 in H₂O) (hydrochloride). Has -CHO in place of -COOMe. λ_{max} 225 (ε 16200) (H₂O) (Berdy).

19-Aldehyde, 3-Ac: Norerythrostachaldine 3-acetate

[765842-22-6]

[55729-26-5 (hydrochloride)]

C₂₅H₃₉NO₆ 449.586

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Strongly cytotoxic to KB cells. Gum; cryst. (EtOH/Me₂CO)(as hydrochloride). Mp 231-232° (hydrochloride). [α]_D¹⁶ -75 (c, 0.4 in H₂O) (hydrochloride). λ_{max} 225 (log ε 4.18) (EtOH) (hydrochloride).

7-Ketone: Norerythrophlamine

[55394-77-9]

C₂₄H₃₇NO₆ 435.559

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Gum; cryst. (Me₂CO) as hydrochloride. Mp 212° (hydrochloride).

7-Ketone, 3-Ac: Norerythrophlamine 3-acetate

[55394-79-1]

C₂₆H₃₉NO₇ 477.597

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae).

7-Ketone, N-Me: Erythrophlamine

[511-00-2]

C₂₅H₃₉NO₆ 449.586

Alkaloid from the bark of *Erythrophleum guineense*, *Erythrophleum ivorense*, *Erythrophleum suaveolens*, *Erythrophleum africanum* and *Erythrophleum couminga* (Fabaceae). Antiseptic, cardiotoxic agent. Needles (Et₂O/petrol). Mp 149-151°. [α]_D²⁰ -62.5 (c, 0.911 in EtOH). Log P 2.41 (calc).

3-Deoxy: Norcassamidine†. *Erythrophleine*

[36150-73-9]

C₂₄H₃₉NO₅ 421.576

Alkaloid from the bark of *Erythrophleum chlorostachys* (Fabaceae). Antiseptic, cardiotoxic agent. Vasodilator, thermolytic agent. Glass. Log P 3.69 (calc).

▶ SF7325000

3-Deoxy, N-Me: Cassamidine

[22260-35-1]

C₂₅H₄₁NO₅ 435.603

Alkaloid from the bark of *Erythrophleum guineense*, *Erythrophleum chlorostachys*, *Erythrophleum couminga*, *Erythrophleum ivorense* and *Erythrophleum africanum* (Fabaceae). Glass or oil; cryst. as hydrochloride. Mp 235-236° (as hydrochloride). [α]_D -47 (c, 1.0 in EtOH) (hydrochloride).

3-Deoxy, 7-ketone, N-Me: Cassamine

[471-71-6]

C₂₅H₃₉NO₅ 433.587

Alkaloid from the bark of *Erythrophleum guineense*, *Erythrophleum couminga* and *Erythrophleum ivorense* (Fabaceae). Antiseptic, cardiac stimu-

lant. Cryst. (pentane). Mp 86-87°. $[\alpha]_D^{19}$ -56 (c, 0.84 in EtOH). Log P 3.7 (calc).

3-Deoxy, 7-ketone, N-Me, hydrochloride: Mp 214-217° dec. $[\alpha]_D^{17}$ -48 (c, 0.65 in H₂O).

3-Deoxy, 6 α -hydroxy, 7-ketone, N-Me: Erythrophleguine. 6-Hydroxycassamine [4829-28-1]

C₂₅H₃₉N₂O₆ 449.586

Alkaloid from the bark of *Erythrophleum guineense*, *Erythrophleum coumanga*, *Erythrophleum ivorense* and *Erythrophleum suaveolens* (Fabaceae). Mp 77-78°. $[\alpha]_D$ -38 (c, 1.6 in EtOH).

Engel, B.G. et al., *Helv. Chim. Acta*, 1949, **32**, 2364-2381 (*Cassamine*, *Erythrophlamine*, *isol*, *uv*)

Mathieson, D.W. et al., *Experientia*, 1960, **16**, 404-406 (*Cassamine*, *Erythrophlamine*, *struct*)

Lindwall, O. et al., *Tet. Lett.*, 1965, 4203-4208 (*Erythrophleguine*)

Thorell, A. et al., *Acta Chem. Scand.*, 1968, **22**, 2835-2844 (*Cassamidine*)

Cronlund, A. et al., *Acta Pharm. Suec.*, 1971, **8**, 351-360; 1975, **12**, 467-478; *CA*, **76**, 11995y; **84**, 135892x (*Cassamine*, *Cassamidine*)

Clarke, R.L. et al., *Phytochemistry*, 1971, **10**, 851-856 (*isol*, *uv*)

Loder, J.W. et al., *Tet. Lett.*, 1972, 5069-5072; 1975, 2497-2498 (*struct*, *nomencl*, *cmr*)


Loder, J.W. et al., *Aust. J. Chem.*, 1974, **27**, 179-185; 1975, **28**, 651-656 (*Cassamidine*, *Norerythrostachaldine 3-acetate*, *Norcassamidine*, *Norerythrostachamine*)

Falkiner, M.J. et al., *Aust. J. Chem.*, 1975, **28**, 645-650 (*Norerythrostachamine 3-acetate*, *Norerythrostachaldine*, *Norerythrophlamine*, *Norerythrophlamine 3-acetate*)

Cronlund, A. et al., *Planta Med.*, 1976, **29**, 123-128 (*Erythrophleguine*, *occur*)

Nor-C-fluorocurarine N-301

2,16,19,20-Tetrahydrocuran-17-al, 9CI. Vincanine. Vinervidine. Metvin



(-)-form

C₁₉H₂₀N₂O 292.38

The synonym Vinervidine was applied to the (±)-form. Metvin (Russian) refers to the hydrochloride.

(-)-form [6880-54-2]

Alkaloid from *Diplorhynchus condylocarpon* ssp. *mossambicensis*, *Vinca erecta*, *Amsonia tabernaemontana*, *Rhazya stricta*, *Leuconotis griffithii*, *Ochrosia elliptica*, *Strychnos dolichothyrsa*, *Tabernaemontana eglandulosa*, *Tabernaemontana ventricosa* and *Alstonia sphaerocapitata* (Apocynaceae, Loganiaceae). Cryst. (Me₂CO). Mp 184-186° dec. $[\alpha]_D^{20}$ -1248 (c, 0.027 in CHCl₃).

N^b-Oxide: Vincanine N-oxide. Norfluorocurarine N-oxide [52845-16-6]

C₁₉H₂₀N₂O₂ 308.379

Alkaloid from *Vinca erecta* and *Leuconotis griffithii*. Also isol. from *Ervatamia hirta* (Apocynaceae).

Amorph. $[\alpha]_D$ -690 (c, 1 in MeOH). Identity of all samples not certain. Opt. rotn. refers to a sample from *E. hirta*.

N^b-Oxide; hydrobromide:

Cryst. (MeOH). Mp 203-205°.

N^b-Me: C-Fluorocurarine. C-Curarine III [6866-39-3]

C₂₀H₂₃N₂O⁺ 307.414

Calabash curare alkaloid; also present in *Strychnos mitscherlichii* and many other *Strychnos* spp. (Loganiaceae). Weak curarising agent. Mp 270-274° dec. (as chloride). $[\alpha]_D^{20}$ -937 (H₂O).

N^b-Me, picrate: Mp 189°.

17-Alcohol, N¹-hydroxyacetyl: 2,16-Didehydro-23-hydroxyretuline. 23-Hydroxy-2,16-dehydroretuline

[88721-03-3]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from stem bark of *Strychnos longicaudata* (Loganiaceae). λ_{max} 213; 255; 283; 293 (EtOH).

10-Methoxy: 10-Methoxynor-C-fluorocurarine

[84575-23-5]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Alstonia lanceolifera* stem bark (Apocynaceae). $[\alpha]_D$ -405 (c, 0.25 in CHCl₃). Yellow colour with Ce(IV).

11-Methoxy, 12-hydroxy: 12-Hydroxy-11-methoxynor-C-fluorocurarine

[182745-06-8]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from root bark of *Strychnos panganensis*. $[\alpha]_D$ -688 (c, 0.47 in CHCl₃).

11-Methoxy, 12-hydroxy, N-Ac, imine:

12-Hydroxy-11-methoxy-N-acetyl nor-C-fluorocurarine

[182745-05-7]

C₂₂H₂₅N₃O₃ 379.458

Alkaloid from root bark of *Strychnos panganensis*. $[\alpha]_D$ -478 (c, 0.65 in CHCl₃).

12-Hydroxy: Vincanidine. 12-Hydroxy-vincanine

[1630-41-7]

C₁₉H₂₀N₂O₂ 308.379

Alkaloid from roots of *Vinca erecta*. Major alkaloid of leaves of *Hazunta membranacea* (Apocynaceae). Cryst. (MeOH or EtOH). $[\alpha]_D^{30}$ -848.6 (c, 0.853 in MeOH). Chars without melting at 250-280°. Originally considered to be 11-hydroxyvincanine. λ_{max} 242 (log ϵ 3.95); 291 (log ϵ 3.26); 375 (log ϵ 4.13) (EtOH).

12-Hydroxy, methiodide:

Cryst. (EtOH). Mp 310-313°.

12-Methoxy: Vincanicine

[32258-16-5]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Vinca erecta* (Apocynaceae). Amorph. $[\alpha]_D^{24}$ -438 (c, 0.6 in CHCl₃). Struct. revised in 1974. λ_{max} 248 (log ϵ 3.9); 293 (log ϵ 3.29); 376 (log ϵ 4.04) (EtOH).

12-Methoxy; methiodide:

Cryst. (MeOH). Mp 280-282°. $[\alpha]_D^{27}$ -615 (c, 0.715 in H₂O).

18-Hydroxy: 18-Hydroxynorfluorocurarine. Strychnofluorine

[88721-06-6]

C₁₉H₂₀N₂O₂ 308.379

Alkaloid from stem bark of *Strychnos goudouensis* and root bark of *Strychnos gossweileri*, also from *Strychnos angolensis* (Loganiaceae). Exhibits low cytotoxic activity vs. B16 melanoma cells. $[\alpha]_D$ -280 (c, 0.3 in MeOH). λ_{max} 243 (ϵ 5750); 291 (ϵ 2090); 299 (ϵ 2400); 365 (ϵ 10200) (MeOH) (Berdy).

18-Acetoxy: 18-Acetoxy norfluorocurarine [88721-05-5]

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from stem bark of *Strychnos goudouensis* (Loganiaceae).

(±)-form [27565-46-4]

Alkaloid from *Vinca erecta*. Mp 183-185°.

[1669-04-1]

Wieland, H. et al., *Annalen*, 1941, **547**, 140-155 (*C-Curarine III*, *isol*)

Schmid, H. et al., *Helv. Chim. Acta*, 1952, **35**, 1864-1879; 1953, **36**, 102-121 (*C-Fluorocurarine*, *isol*, *uv*)

Stauffer, D. et al., *Helv. Chim. Acta*, 1961, **44**, 2006-2015 (*Nor-C-fluorocurarine*, *isol*, *ir*, *uv*)

Fritz, H. et al., *Annalen*, 1963, **663**, 150-156 (*Nor-C-fluorocurarine*, *synth*)

Yuldashev, P.Kh. et al., *Khim. Prir. Soedin.*, 1965, **1**, 34-42; 1974, **10**, 260-261; 1983, **19**, 210-212; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 25-42; 1974, **10**, 276-277; 1983, **19**, 199-201 (*Vincanidine*, *Vincanicine*, *cmr*, *struct*, *bibl*)

Pyuskyulev, B. et al., *Tet. Lett.*, 1967, 4559-4562 (*pmr*, *ms*)

Rakhimov, D.A. et al., *Khim. Prir. Soedin.*, 1969, **5**, 461-462; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **4**, 386-387 (*Vincanicine*)

Crawley, G.C. et al., *Chem. Comm.*, 1971, 685-686 (*C-Fluorocurarine*, *synth*)

Sharipov, M. et al., *Khim. Prir. Soedin.*, 1974, **10**, 263-264; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 281-283 (*Vincanine N-oxide*)

Ahmad, Y. et al., *J.A.C.S.*, 1977, **99**, 1943-1946 (*Nor-C-fluorocurarine*, *isol*, *synth*)

Bui, A.M. et al., *Phytochemistry*, 1977, **16**, 703-706 (*Vincanidine*, *isol*)

Coune, C.A. et al., *Herba Hung.*, 1980, **19**, 189-193; *CA*, **93**, 217906e (*18-Hydroxynorfluorocurarine*)

Ravao, T. et al., *Phytochemistry*, 1982, **21**, 2160-2161 (*10-Methoxynorfluorocurarine*)

Massiot, G. et al., *Tetrahedron*, 1983, **39**, 3645-3656 (*18-Hydroxynorfluorocurarine*, *18-Acetoxy norfluorocurarine*, *23-Hydroxy-2,16-dehydroretuline*)

Goh, S.H. et al., *Tet. Lett.*, 1984, **25**, 3483-3484 (*Nor-C-fluorocurarine N-oxide*)

Clivio, P. et al., *Phytochemistry*, 1991, **30**, 3785-3792 (*Vincanine N-oxide*)

Quetin-Leclercq, J. et al., *Phytochemistry*, 1992, **31**, 4347-4349 (*18-Hydroxynorfluorocurarine*)

Sole, D. et al., *J.O.C.*, 1996, **61**, 4194-4195 (*Nor-C-fluorocurarine*, *synth*)

Nuzillard, J.-M. et al., *Phytochemistry*, 1996, **43**, 897-902 (*12-Hydroxy-11-methoxynor-C-*


fluorocararine, 12-Hydroxy-11-methoxy-N-acetyl-nor-C-fluorocararine)

Delaude, C. et al., *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-186 (23-Hydroxy-2,16-dehydroretuline)

Bonjoch, J. et al., *J.A.C.S.*, 1997, **119**, 7230-7240 (*Nor-C-fluorocararine, synth*)

Norisotuboflavine**N-302**

6-Methyl-4H-indolo[3,2,1-de]-1,5-naphthyridin-4-one, 9CI. 6-Methyl-canthin-4-one
[3464-65-1]



C₁₅H₁₀N₂O 234.257

Alkaloid from *Pleiocarpa mutica* (Apocynaceae). Mp 298-300° (282-284°).

6-Ethyl analogue: 6-Ethylcanthin-4-one.

Isotuboflavine

[3464-64-0]

C₁₆H₁₂N₂O 248.284

Alkaloid from the stem bark of *Pleiocarpa mutica* (Apocynaceae). Yellow needles (MeOH). Mp 263-265°.


Achenbach, H. et al., *J.A.C.S.*, 1965, **87**, 4177 (*isol, uv, ir, ms, struct*)

Rosenkranz, H.J. et al., *Helv. Chim. Acta*, 1968, **51**, 565 (*synth*)

McEvoy, F.J. et al., *J.O.C.*, 1969, **34**, 4199 (*synth, uv, ms*)

Normetanephrine**N-303**

α -(Aminomethyl)-4-hydroxy-3-methoxybenzenemethanol, 9CI. α -(Aminomethyl)vanillyl alcohol, 8CI. 2-Amino-1-(4-hydroxy-3-methoxyphenyl)ethanol. Normetadrenaline

**(R)-form**

C₉H₁₃NO₃ 183.207

Metab. of Norepinephrine, N-298. Used as a marker of autonomic nervous system dysfunction, and of phaeochromocytoma.

(R)-form

[13015-71-9 (hydrochloride)]

Cryst. (MeOH/Et₂O). Mp 192-194° dec. (as hydrochloride).

N-Me: **Metanephrine**. Metadrenaline. O³-Methyladrenaline

[2282-54-4, 105870-52-8]

C₁₀H₁₅NO₃ 197.233

Alkaloid from *Coryphantha macromeris* var. *runyonii* (Cactaceae). Also obt. from biol. sources e.g. urine. Sympathomimetic agent. Mp 158-159°. Opt. rotn. of nat. alkaloid not detd., but abs. config. v. prob. (R-).

(±)-form

Mp 192-195° (hydrochloride).

N-(4-Hydroxy-3-methoxy-E-cinnamoyl):

N-trans-FeruloylnormetanephrineC₁₉H₂₁NO₆ 359.378

Alkaloid from the leaves of *Aptenia cordifolia*. Oil. λ_{\max} 226 (log ϵ 2.9); 298 (log ϵ 1.2); 320 (log ϵ 1.1) (MeOH).

N-Me:

[881-95-8 (hydrochloride)]

Prisms (EtOH/Et₂O) (as hydrochloride).

Mp 175° (hydrochloride).

N-Isopropyl: **Metiprenaline**. 3-Methoxyisoprenaline. Egypt 402. Moxa

[1212-03-9]

[1420-27-5]

C₁₂H₁₉NO₃ 225.287

Metab. of Isoprenaline. Antiallergic drug. Sympathomimetic agent. Mp 152° (as hydrochloride). Log P 0.6 (calc).

► LD₅₀ (mus, ivn) 305 mg/kg. YW6120000

(ξ)-form [97-31-4]

Alkaloid from the leaves of potato *Solanum tuberosum*.

[1011-74-1, 709-52-4, 2282-53-3, 35778-41-7, 5001-33-2]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 625A (*nmr*)

Langecker, H. et al., *Naunyn-Schmiedeberg Arch. Exp. Pathol. Pharmacol.*, 1955, **226**, 493-504 (*pharmacol, tox*)

Axelrod, J. et al., *J. Biol. Chem.*, 1958, **233**, 697-701 (*isol, synth*)

Heacock, R.A. et al., *Chem. Ind. (London)*, 1961, 595 (*synth*)

Cession-Fossion, A. et al., *Arch. Int. Pharmacodyn. Ther.*, 1964, **150**, 160 (*pharmacol*)

Breese, G.R. et al., *J. Pharmacol. Exp. Ther.*, 1968, **165**, 9-13 (*metab*)

Pyrin, L. et al., *Pathol. Biol.*, 1968, **16**, 447-455 (*occur*)

Keller, W.J. et al., *J. Pharm. Sci.*, 1973, **62**, 408-411 (*N-Me, isol*)

Anhoury, M.-L. et al., *J.C.S. Perkin 1*, 1974, 1015-1017 (*synth, pmr*)

Manusadzhyan, V.G. et al., *Zh. Org. Khim.*, 1974, **10**, 145-149; *J. Org. Chem. USSR (Engl. Transl.)*, 1974, **10**, 147-150 (*ms*)

Pattanayek, R.R. et al., *Acta Cryst. C*, 1984, **40**, 294-297 (*cryst struct*)

Albert, L.S.R. et al., *J. Chromatogr.*, 1984, **312**, 357-385 (*isol, tlc*)

Hassel, P.G. et al., *CA*, 1987, **106**, 207727 (*hplc*)

Bryan, L.J. et al., *J. Chromatogr.*, 1989, **487**, 29-39 (*hplc*)

Gamache, P.H. et al., *Clin. Chem. (Winston-Salem, N.C.)*, 1993, **39**, 1825-1830 (*hplc*)

Peaston, R.T. et al., *J. Clin. Pathol.*, 1993, **46**, 734 (*use*)

Eisenhofer, G. et al., *Clin. Sci.*, 1995, **88**, 533-542 (*pharmacol*)

Szopa, J. et al., *Phytochemistry*, 2001, **58**, 315-320 (*Normetanephrine, ξ-form, isol, ms*)

Lenders, J.W.M. et al., *Ann. N.Y. Acad. Sci.*, 2002, **970**, 29-40 (*pharmacol, rev*)

Swiedrych, A. et al., *Plant Physiol. Biochem. (Paris)*, 2004, **42**, 593-600 (*ξ-form, isol*)


Oelmann, T. et al., *CA*, 2005, **142**, 152734 (*rev, pharmacol*)

Mariola, J. et al., *J. Hypertens.*, 2006, **24**, 2331-2339 (*pharmacol, rev*)

DellaGreca, M. et al., *Tetrahedron*, 2006, **62**, 2877-2882 (*N-Feruloylnormetanephrine*)

Norrufescine**N-304**

4,5,6-Trimethoxyindeno[1,2,3-ij]isoquinolin-9-ol, 9CI. 9-Hydroxy-4,5,6-trimethoxyindeno[1,2,3-ij]isoquinoline [58189-34-7]



C₁₈H₁₅NO₄ 309.321

Alkaloid from the stems of *Abuta imene*, *Abuta rufescens* and *Telitoxicum peruvianum*. Also isol. from the roots and wood of *Cissampelos pareira* (Menispermaceae). Exhibits cytotoxicity against P388 cells. Orange-yellow cryst. (MeOH or MeOH/CHCl₃). Mp 235-238° dec. (232-234°). λ_{\max} 208 (ϵ 25800); 252 (ϵ 32000); 300 (ϵ 21300); 308 (ϵ 19400); 340 (ϵ 2400) (MeOH) (Berdy).

Me ether: Rufescine

[38366-04-0]

C₁₉H₁₇NO₄ 323.348

Alkaloid from *Abuta imene* and *Abuta rufescens* (Menispermaceae). Bright-yellow needle-shaped prisms (hexane/Et₂O). Mp 88-90°.

Cava, M.P. et al., *Tetrahedron*, 1975, **31**, 1667 (*isol, uv, ir, pmr, struct*)

Menachery, M.D. et al., *J. Nat. Prod.*, 1981, **44**, 320 (*isol*)


Menachery, M.D. et al., *Heterocycles*, 1982, **19**, 2255 (*synth*)

Boyer, D.L. et al., *J.O.C.*, 1984, **49**, 4050 (*synth, ir, pmr, ms, Rufescine*)

Morita, H. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1307 (*isol, uv, ir, pmr, cmr*)

Norruspoline**N-305**

2-Methoxy-4-[2-(2-pyrrolidinyl)ethenyl]phenol, 9CI. 2-(4-Hydroxy-3-methoxytyryl)pyrrolidine

**(R)-form**

C₁₃H₁₇NO₂ 219.283

(R)-form [132340-25-1]

Synthetic. Pale orange cryst. Mp 142°. [α]_D +132.9 (c, 0.8 in EtOH).

(±)-form [67270-82-0]

Alkaloid from seeds of *Ruspolia hypercrateriformis* (Acanthaceae). Red-brown gum.

N,O-Di-Ac:

Oil.

Me ether:


Oil.

Roessler, F. et al., *Helv. Chim. Acta*, 1978, **61**, 1200 (*isol, uv, ms, pmr, struct*)

Wanner, K.T. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1990, **323**, 977 (*synth*)

Brown, D.S. *et al.*, *Tetrahedron*, 1991, **47**, 1311 (synth)

Norsecurinic acid N-306
[940274-57-7]




Absolute Configuration

$C_{12}H_{15}NO_3$ 221.255
Alkaloid from the roots of *Flueggea virosa*.

Gan, L.-S. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 819-823 (isol, pmr, cmr)

Norsecurinine N-307
A-Norsecurinan-11-one, 9CI



(+)-form

$C_{12}H_{13}NO_2$ 203.24

(+)-form [25472-13-3]

Alkaloid from *Phyllanthus niruri* (Euphorbiaceae). Pale yellow oil. $[\alpha]_D^{25} +213$ (CHCl₃). $[\alpha]_D^{25} +255.8$ (c, 4.2 in MeOH).

Hydrochloride:

Plates (MeOH/Et₂O). Mp 275-276°. $[\alpha]_D^{20} +188.6$ (c, 0.93 in H₂O).

4ξ-Methoxy: 4-Methoxynorsecurinine

$C_{13}H_{15}NO_3$ 233.266

Alkaloid from *Phyllanthus niruri* (Euphorbiaceae). $[\alpha]_D$ -47 (c, 0.3 in MeOH).

4ξ-Methoxy, 14,15-dihydro: 4-Methoxy-dihydronorsecurinine

[130968-99-9]

$C_{13}H_{17}NO_3$ 235.282

Alkaloid from *Phyllanthus niruri* (Euphorbiaceae). Yellow granules (EtOH). Mp 110-111°. Abs. config. not detd., assigned here on biogenetic probability.

(-)-form [2650-35-3]

Alkaloid from the root bark of *Securinega virosa* (Euphorbiaceae). Needles (petrol) (also descr. as water-sol. viscous liq.). Mp 81-82°. $[\alpha]_D^{20} -19.5$ (c, 0.2 in EtOH). $[\alpha]_D -272$ (c, 6.9 in EtOH). The low sp. rotn. quoted by Iketubosin *et al* suggests that their specimen was largely racemic.

Hydrochloride: Mp 223-225° dec. $[\alpha]_D^{20} -309$ (c, 1.24 in EtOH).

Picrate:

Yellow needles (DMF). Mp 258-260° dec.

14S,15R-Epoxyde: 14,15-Epoxy-norsecurinine

$C_{12}H_{13}NO_3$ 219.24

Alkaloid from *Flueggea virosa*.

14,15-Dihydro: Dihydronorsecurinine.

14,15-Dihydro-A-norsecurinan-11-one,

9CI. Virosine†

[2650-34-2]

$C_{12}H_{15}NO_2$ 205.256

Minor alkaloid from the root bark of *Securinega virosa* (Euphorbiaceae). Rhombs (EtOAc). Mp 135-136°. $[\alpha]_D -13$ (c, 1.0 in dioxan).

14,15-Dihydro; hydrochloride:

Needles (EtOH/Et₂O). Mp 228-230° dec.

14,15-Dihydro, 15ξ-hydroxy: Fluggeainol

[101899-47-2]

$C_{12}H_{15}NO_3$ 221.255

Alkaloid from *Flueggea virosa* (Euphorbiaceae).

14,15-Dihydro, 15ξ-hydroxy, dimeric ether: Fluggeaine ether

[101899-48-3]

$C_{24}H_{28}N_2O_5$ 424.496

Alkaloid from *Flueggea virosa* (Euphorbiaceae).

Iketubosin, G.O. *et al.*, *J. Pharm. Pharmacol.*, 1963, **15**, 810 (isol, uv, ir, pmr, ms, struct)

Saito, S. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1126; *CA*, **62**, 5498d (isol,

Dihydronorsecurinine)

Saito, S. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 614; 786 (*Norsecurinine,*

Dihydronorsecurinine)

Rouffiac, R. *et al.*, *Plant. Med. Phytother.*, 1969, **3**, 220 (isol, uv, struct)

Mulchandani, N.B. *et al.*, *Planta Med.*, 1984, **50**, 104 (*4-Methoxynorsecurinine*)

Chen, M. *et al.*, *Zhǐwǔ Xuebao (Acta Bot. Sin.)*, 1985, **27**, 625; *CA*, **104**, 183282b

(*Fluggeainol, Fluggeaine ether*)

Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 614

(isol, uv, ir, pmr, cmr, ms, cd, ord, cryst struct)

Heathcock, C.H. *et al.*, *Heterocycles*, 1987, **25**, 75 (synth)

Hassarajani, S.A. *et al.*, *Indian J. Chem., Sect. B*, 1990, **29**, 801 (*4-Methoxydihydronorsecurinine*)

Jacobi, P.A. *et al.*, *J.A.C.S.*, 1991, **113**, 5384

(synth)

Magnus, P. *et al.*, *J.A.C.S.*, 1992, **114**, 382

(synth)

Magnus, P. *et al.*, *Tetrahedron*, 1993, **49**, 8059

(synth)

Dehmlow, E.V. *et al.*, *Phytochemistry*, 1999, **52**, 1715-1716 (*14,15-Epoxy-norsecurinine*)

Han, G. *et al.*, *J.O.C.*, 2000, **65**, 6293-6306


(synth)

Alibés, R. *et al.*, *Org. Lett.*, 2005, **7**, 5107-5109

(synth)

Norsegoline N-308

[117694-98-1]



$C_{18}H_{14}N_2O_3$ 306.32

Alkaloid from the Red Sea tunicate *Eudistoma* sp. λ_{max} 225 (ε 18900); 264 (ε 12000); 295 (ε 10900); 366 (ε 1500); 385 (ε 2300); 402 (ε 2300); 515 (ε 850) (MeOH/HCl) (Derep). λ_{max} 225 (ε

17300); 264 (ε 12700); 340 (ε 3000); 367 (ε 1500); 400 (ε 2800); 440 (ε 1200) (MeOH) (Derep).


Rudi, A. *et al.*, *J.O.C.*, 1989, **54**, 5331 (isol, uv, ir, pmr, cmr, struct)

Dunn, S.H. *et al.*, *J.C.S. Perkin 1*, 1993, 879 (synth)

Kitahara, Y. *et al.*, *Nat. Prod. Lett.*, 1993, **2**, 159 (synth)

Nakahara, S. *et al.*, *Heterocycles*, 2005, **65**, 1925-1929 (synth)

Norsenecaudatin 9-O-senecioate N-309




$C_{15}H_{21}NO_5$ 295.335

Alkaloid from *Senecio caudatus* (Asteraceae). The trivial name is somewhat misleading; compared with Senecicaudatin it lacks a -C(CH₃)₂OH group.

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151 (isol, ir, ms, struct)

Norstaminolactone A N-310

[460741-44-0]



$C_{38}H_{45}NO_{12}$ 707.773

Constit. of *Orthosiphon stamineus*. Amorph. solid. $[\alpha]_D^{25} -18.3$ (c, 0.25 in CHCl₃).


Awale, S. *et al.*, *Tetrahedron*, 2002, **58**, 5503-5512 (isol, pmr, cmr)

Nortenuazonic acid N-311

Valyltenuazonic acid

[16820-44-3]

[707-84-6, 2113-91-9]




$C_9H_{13}NO_3$ 183.207

Tetramic acid deriv. Prod. by *Pyricularia oryzae*. Phytotoxin. $[\alpha]_D -160$ (c, 0.2 in H₂O) (as Na salt). λ_{max} 240 (log ε 4.04); 280 (log ε 4.14) (H₂O) (Na salt). λ_{max} 240 (ε 11000); 280 (ε 13800) (H₂O) (Berdy).

Lebrun, M.H. *et al.*, *Phytochemistry*, 1988, **27**, 77-84; 1990, **29**, 3777-3783 (isol, synth)

Nortopsentin A N-312

3,3'-(1*H*-Imidazole-2,4-diyl)bis[6-bromo-1*H*-indole], 9*CI*. 2,4-Bis(6-bromo-3-indolyl)imidazole
[134029-43-9]



C₁₉H₁₂Br₂N₄ 456.139

Alkaloid from the marine sponges *Spongosorites ruetzleri* and *Halichondria* sp. Exhibits cytotoxic and antifungal activities. Oil. λ_{max} 207 (ε 50300); 236 (ε 42300); 277 (ε 26400); 310 (sh) (MeOH) (Derep).

6-Debromo: Nortopsentin B

[134029-44-0]

C₁₉H₁₃BrN₄ 377.242

Alkaloid from *Spongosorites ruetzleri* and *Halichondria* sp. Exhibits cytotoxic and antifungal activities. Plates (EtOAc/CHCl₃). Dec. at 250-270°. λ_{max} 206 (ε 50700); 232 (ε 45200); 278 (ε 25600); 310 (sh) (MeOH) (Derep).

6-Debromo: Nortopsentin C

[134029-45-1]

C₁₉H₁₃BrN₄ 377.242

Alkaloid from *Spongosorites ruetzleri* and *Halichondria* sp. Exhibits cytotoxic and antifungal activities. Oil. λ_{max} 206 (ε 50700); 232 (ε 45200); 278 (ε 25600); 310 (sh) (MeOH) (Derep). λ_{max} 207 (ε 50300); 280 (MeOH) (Berdy).

6,6'-Bisdebromo: 3,3'-(1H-Imidazole-2,4-diyl)bis-1H-indole, 9CI. Nortopsentin D†

[134779-34-3]

C₁₉H₁₄N₄ 298.346

Prod. by *Halichondria* sp. λ_{max} 207 (ε 50300); 280 (MeOH) (Berdy).

Sun, H.H. *et al.*, *CA*, 1991, **115**, 35701z


Sakemi, S. *et al.*, *J.O.C.*, 1991, **56**, 4304 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kawasaki, I. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1831 (*synth*)

Miyake, F.Y. *et al.*, *Org. Lett.*, 2000, **2**, 2121-2123 (*synth*)

Nortopsentin D†

N-313



C₂₃H₁₇Br₂N₇O 567.242


Alkaloid from the sponge *Dracmacidon* sp. Amorph. solid. Not to be confused with Nortopsentin D in N-312. λ_{max} 276 (ε 28000); 294 (ε 22000); 360 (ε 15400) (MeOH). λ_{max} 276 (ε 2800); 334 (ε 210000) (MeOH/HCl) (Berdy). λ_{max} 283 (ε 28000); 352 (ε 6000) (MeOH/

NaOH) (Berdy).

Mancini, I. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 2075-2082

Nortrachelogenin amide

N-314



C₂₁H₂₇NO₇ 405.447


Alkaloid from *Trachelospermum jasminoides*. Amorph. solid. [α]_D²⁵ -9 (c, 0.55 in MeOH).

Tan, X.-Q. *et al.*, *Planta Med.*, 2005, **71**, 93-95 (*isol, pmr, cmr, ms*)

Nortryptoquivaline

N-315

Tryptoquivaline D. Fumitremorgin D
[60676-56-4]



Absolute Configuration

C₂₈H₂₈N₄O₇ 532.552

The stereochemical information on some of the derivs. is not certain. Toxic metab. from *Aspergillus clavatus* and *Aspergillus fumigatus*. Tremorgenic toxin. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 256-258°. [α]_D²⁵ +170 (c, 0.64 in CHCl₃). λ_{max} 228 (ε 3600); 279 (ε 10200); 306 (ε 4500); 319 (ε 3500) (MeOH) (Berdy).

Ac: [60676-59-7]

Prisms (Et₂O). Mp 155-157°. [α]_D²⁵ +159 (c, 0.16 in CHCl₃).

27-Ketone, O-de-Ac: Nortryptoquivalone.

Tryptoquivaline B. Tryptoquivalone.

FTB

[55387-46-7]

C₂₆H₂₄N₄O₆ 488.499

Metab. of *Aspergillus clavatus*. Tremorgenic toxin. Prisms (CH₂Cl₂/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 208-209°. [α]_D²⁵ +255 (c, 0.3 in CHCl₃). λ_{max} 234 (ε 35000); 292 (ε 9550); 320 (ε 6300) (EtOH) (Derep).

▶ Toxic.

N-Deoxy: N-Deoxynortryptoquivaline

[60676-58-6]

C₂₈H₂₈N₄O₆ 516.552

Metab. of *Aspergillus clavatus*. Prisms (Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 158-160°. [α]_D²⁵ +69.5 (c, 0.82 in CHCl₃). λ_{max} 228 (ε 43900); 268 (ε 11700); 305 (ε 4100); 317 (ε 3300) (MeOH) (Berdy).

N-Deoxy, 27-ketone, O-de-Ac: Tryptoquivaline N. Fumitremorgin N. Deoxynortryptoquivalone

[60676-61-1]

C₂₆H₂₄N₄O₅ 472.499

Isol. from *Aspergillus fumigatus* and *Aspergillus clavatus*. Tremorgenic toxin. Needles (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 193-197°. [α]_D²³ +127 (c, 0.06 in DMSO). [α]_D²⁵ +171 (c, 0.79 in CHCl₃). λ_{max} 232 (ε 32400); 288 (ε 9250) (MeOH) (Berdy).

▶ Toxic.

N-Acetoxy, N-deoxy, O-de-Ac: Norisotryptoquivaline

[61897-84-5]

C₂₈H₂₈N₄O₇ 532.552

Metab. of *Aspergillus fumigatus*. Mp 224-225° dec. [α]_D²⁸ +115 (c, 0.23 in CHCl₃).

12-Epimer: Tryptoquivaline M. Fumitremorgin M

[69575-59-3]

C₂₈H₂₈N₄O₇ 532.552

Isol. from *Aspergillus fumigatus*. Tremorgenic toxin. Plates (MeOH aq.). Mp 157-164°. [α]_D²⁴ -154 (c, 0.5 in CHCl₃).

▶ Toxic.

12-Epimer, O-Ac: [71658-07-6]

Amorph. solid. [α]_D²¹ -148 (c, 0.28 in CHCl₃).

27-Epimer: 27-Epinortryptoquivaline

[182967-48-2]

C₂₈H₂₈N₄O₇ 532.552

Metab. from *Corynascus setosus*. Tremorgenic agent. Needles (MeOH). Mp 236-238°. [α]_D¹⁸ +196 (c, 0.292 in CHCl₃). λ_{max} 209 (log ε 4.64); 290 (sh) (log ε 3.8); 307 (sh) (log ε 3.53); 318 (sh) (log ε 3.43) (MeOH).

Yamazaki, M. *et al.*, *Tet. Lett.*, 1976, 2861-2864 (*Norisotryptoquivaline*)

Büchi, G. *et al.*, *J.O.C.*, 1977, **42**, 244-246

(*Nortryptoquivaline*,

Deoxynortryptoquivaline,

Nortryptoquivalone, isol, struct, uv, ir, pmr, ms, cd)

Yamazaki, M. *et al.*, *Chem. Pharm. Bull.*,

1979, **27**, 1611-1617 (*Tryptoquivalines M,N*)

Springer, J.P. *et al.*, *Tet. Lett.*, 1979, **20**, 339-342

(*Nortryptoquivaline, cryst struct, abs config*)

Fujimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1996,

44, 1843-1848 (*27-Epinortryptoquivaline*)

Handbook of Secondary Fungal Metabolites,


(ed. Cole, R.J. *et al.*), Academic Press, 2003,

1, 383; 408

Norushinsunine

N-316

Michelalbine. Normicheline A



(-)-form

C₁₇H₁₅NO₃ 281.31

(+)-form

N-Me, N-oxide(β-): (+)-Norushinsunine

N-oxideC₁₈H₁₇NO₄ 311.337

Alkaloid from *Cananga odorata* (ylang ylang). Cryst. Mp 177-179°. [α]_D²⁴ +50.7 (c, 0.05 in CHCl₃). λ_{\max} 271 (log ϵ 4.03); 283 (log ϵ 3.98); 320 (log ϵ 3.61) (EtOH).

(-)-form [3175-84-6]

Alkaloid from a wide variety of genera in the Annonaceae (*Annona*, *Asimina*, *Desmos*, *Melodorum*, *Polyalthia*), Magnoliaceae (*Liriodendron*, *Magnolia*, *Michelia*, *Elmerrillia*), Menispermaceae (*Sinomenium*), Eupomatiaceae (*Eupomatia*) and Monimiaceae (*Laurelia*). Shows antimicrobial activity. Mp 206-207°. [α]_D²⁵ -103 (c, 0.60 in CHCl₃).

N,O-Di-Ac: [3327-33-1]

Mp 243-245°.

N-Me: Ushinsunine. Micheline A

[3175-89-1]

C₁₈H₁₇NO₃ 295.337

Alkaloid from a variety of genera in the Annonaceae (*Asimina*, *Cananga*, *Polyalthia*), Lauraceae (*Litsea*), Magnoliaceae (*Michelia*) and Menispermaceae (*Stephania*). Shows antiplasmodial, antiamoebic and cytotoxic activities. Mp 122-123°. [α]_D -122 (c, 0.37 in MeOH).

▶ CE1055600

N-Me, N-oxide(β -): Ushinsunine β -N-oxideC₁₈H₁₇NO₄ 311.337

Alkaloid from the rhizomes of *Stephania venosa* (Menispermaceae). Amorph. [α]_D -52 (c, 0.06 in MeOH).

N,N-Di-Me: N-Methylushinsunine

[61619-44-1]

C₁₉H₂₀NO₃⁺ 310.372


Quaternary alkaloid from the bark of *Elmerrillia papuana*. Cryst. (Me₂CO aq.) (as iodide). Mp 235-237° (iodide). [α]_D -118 (c, 1 in CHCl₃).

(\pm)-form

Synthetic. Mp 210-212° (202-203°).

N-Me:Synthetic. Mp 144-145°.Yang, S.-S. *et al.*, *Hua Hsueh*, 1961, **144**; *CA*, **56**, 1489c (*N-Methylushinsunine*)Yang, T.-H. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 811; 1963, **83**, 216 (*isol*)Tomita, M. *et al.*, *Yakugaku Zasshi*, 1965, **85**, 77 (*isol*, *uv*, *ir*, *ms*, *pmr*)Lu, S.T. *et al.*, *Yakugaku Zasshi*, 1969, **89**, 1313 (*uv*, *ir*, *pmr*, *deriv*)Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 1502 (*synth*, *abs config*)Cleaver, L. *et al.*, *Aust. J. Chem.*, 1976, **29**, 2003 (*N-Methylushinsunine*)Kessar, S.V. *et al.*, *Tet. Lett.*, 1980, **21**, 3307 (*synth*, *Norushinsunine*, *Ushinsunine*)Charles, B. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1113 (*Ushinsunine N-oxide*)Yang, T.H. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1988, **35**, 305; *CA*, **110**, 111669j (*Ushinsunine N-oxide*)Simeon, S. *et al.*, *Pharmazie*, 1989, **44**, 593-597 (*activity*)Hsieh, T.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1999, **46**, 607-612 ((+)-*Ushinsunine N-oxide*)Wright, C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1638-1640 (*activity*)**Norzoanthamine**

[164991-65-5]



Absolute Configuration

C₂₉H₃₉NO₅ 481.631

Alkaloid from the colonial zoanthid *Zoanthus* sp. Cytotoxic agent. Active against osteoporosis. Cryst. Mp 282-285°. [α]_D +1.6 (c, 1 in CHCl₃).

3 β -Hydroxy: 3-Hydroxynorzoanthamine

[231945-74-7]

C₂₉H₃₉NO₆ 497.63

Alkaloid from a zoanthid. Amorph.

solid. [α]_D²⁵ +12 (c, 0.075 in Et₂O).**11 ξ -Hydroxy: 11-Hydroxynorzoanthamine**

[231945-76-9]

C₂₉H₃₉NO₆ 497.63

Alkaloid from a zoanthid. Amorph.

solid. [α]_D²⁵ -11.5 (c, 0.13 in Et₂O).**11-Oxo: Norzoanthaminone**

[164991-66-6]

C₂₉H₃₇NO₆ 495.614Alkaloid from the zoanthid *Zoanthus* sp. Interleukin-6 inhibitor, osteoporosis inhibitor.**30-Hydroxy: 30-Hydroxynorzoanthamine**

[231945-75-8]

C₂₉H₃₉NO₆ 497.63


Alkaloid from a zoanthid. Amorph.

solid. [α]_D²⁵ -1.1 (c, 0.27 in CHCl₃).**A^{15,26}-Isomer, 17S-alcohol: Epinorzoanthamine**

[159509-37-2]

C₂₉H₄₁NO₅ 483.647Alkaloid from *Zoanthus* sp. Misleading name.Fukuzawa, S. *et al.*, *Heterocycl. Commun.*, 1995, **1**, 207-214 (*isol*)Kuramoto, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1998, **71**, 771-779 (*isol*, *pmr*, *abs config*, *Norzoanthaminone*, *Epinorzoanthamine*)Daranas, A.H. *et al.*, *Tetrahedron*, 1999, **55**, 5539-5546 (*Hydroxyzoanthamines*)Miyashita, M. *et al.*, *Science (Washington, D.C.)*, 2004, **305**, 495-499 (*rev*, *synth*)Juhl, M. *et al.*, *J.O.C.*, 2007, **72**, 4644-4654 (*synth*)**Norzoanemonin****N-318**

4-Carboxy-1,3-dimethyl-1H-imidazolium hydroxide inner salt, 9CI, 1,3-Dimethyl-4-imidazolecarboxylic acid betaine
[51800-34-1]

C₆H₈N₂O₂ 140.141

Isol. from the sponges *Axinyssa* sp. and *Niphates* sp.; also from the Caribbean gorgonian *Pseudopterogorgia americana* and the hydroid *Tubularia larynx*. Also from *Cacospongia scalaris* and *Astrosciera willelyana*. Cryst. (MeOH/Me₂CO). Mp 260-263° (250-251°). λ_{\max} 212 (ϵ 2005) (MeOH).

Hydrochloride:

Cryst. (MeOH). Mp 216-219° (213-217°).


Me ester:C₇H₁₁N₂O₂⁺ 155.176

Isol. from various sponges incl. *Axinyssa* sp., *Axinella* sp., *Echinodictyum* sp. and *Niphates* sp. Counterion not specified. λ_{\max} 264 (MeOH).

Weinheimer, A.J. *et al.*, *Tetrahedron*, 1973, **29**, 3135-3136 (*isol*, *synth*, *ir*, *pmr*, *ms*)Gupta, K.C. *et al.*, *Experientia*, 1977, **33**, 1556 (*isol*, *cryst struct*)Jahn, T. *et al.*, *Tet. Lett.*, 1997, **38**, 3883-3884 (*isol*, *pmr*, *cmr*)Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 118-122 (*Me ester*)**Nosiheptide, 9CI, BAN,****N-319****INN, USAN**

Nosiheptide. Multithiomycin. Primofax. RP 9671

[56377-79-8]

C₅₁H₄₃N₁₃O₁₂S₆ 1222.377


Prod. by *Streptomyces actuosus*, *Streptomyces antibioticus*, *Streptomyces glaucogriseus* and *Actinoplanes* sp. Shows broad-spectrum antibacterial activity *in vitro*. Veterinary growth stimulant. Yellow powder or prisms. Mp 310-320°. λ_{\max} 292 (ϵ 31100); 406 (ϵ 16100) (MeOH/NaOH) (Derep). λ_{\max} 328 (ϵ 26900); 420 (sh) (ϵ 2440) (MeOH) (Derep).

▶ RD2685000

Pascard, C. *et al.*, *J.A.C.S.*, 1977, **99**, 6418-6423 (*cryst struct*)Depaire, H. *et al.*, *Tet. Lett.*, 1977, **18**, 1395-1396; 1397-1402; 1401-1406; 1403 (*struct*, *nmr*, *ir*, *uv*)Endo, T. *et al.*, *J. Antibiot.*, 1978, **31**, 623-625 (*struct*, *ir*, *cmr*)Benazet, F. *et al.*, *Experientia*, 1980, **36**, 414-416 (*isol*, *props*, *activity*)Japan. Pat., 1981, 81 127 092; *CA*, **96**, 33357 (*isol*)Iwakawa, M. *et al.*, *Chem. Lett.*, 1982, 1975-1978 (*synth*)Eur. Pat., 1983, 73 329; *CA*, **99**, 37103 (*isol*)Houck, D.R. *et al.*, *J.A.C.S.*, 1987, **109**, 1250-1252; 1988, **110**, 5800-5806 (*biosynth*)

Mocek, U. *et al.*, *J. Antibiot.*, 1989, **42**, 1643-1648 (*pmr, cmr, struct*)
Kirk-Othmer Encycl. Chem. Technol., 4th edn., Wiley, 1991, **3**, 281-282 (*rev*)
Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 188


Nostocine A **N-320**
 2,7-Dihydro-7-methyl-3H-pyrazolo[4,3-*e*][1,2,4]triazin-3-one, 9CI
 [180128-24-9]



$C_5H_5N_5O$ 151.127
 Metab. of the freshwater cyanobacterium *Nostoc spongiaeforme*. Exhibits a broad spectrum of growth inhibitory activity. Algicide. Phytotoxin. Allelopathic agent. Black needles (MeOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 171-172°. λ_{max} 239 (ϵ 6400); 277 (sh) (ϵ 3000); 555 (ϵ 900 br. (540-570 nm)) (MeOH).

Hirata, K. *et al.*, *Heterocycles*, 1996, **43**, 1513 (*isol, uv, ir, pmr, cryst struct*)
 Kelly, T.R. *et al.*, *J.A.C.S.*, 2006, **128**, 5646-5647 (*synth*)

Nostocyclamide **N-321**
 [170129-43-8]



R = -CH(CH₃)₂

$C_{20}H_{22}N_6O_4S_2$ 474.564
 Isol. from the freshwater cyanobacterium *Nostoc* sp. 31. Anticyanobacterial and antialgal agent. Toxic against the freshwater rotifer *Brachionus calyciflorus*. Cryst. (EtOAc/petrol). Sol. CHCl₃, CH₂Cl₂; fairly sol. MeOH; poorly sol. H₂O. Mp 255.8-256.9° dec Mp 259-260° (synthetic). $[\alpha]_D^{25} +25$ (CHCl₃) (natural). $[\alpha]_D^{19} +51.3$ (c, 0.84 in CHCl₃) (synthetic). λ_{max} 202 (ϵ 30900); 223 (ϵ 25120) (MeOH) (Berdy).


Todorova, A.K. *et al.*, *J.O.C.*, 1995, **60**, 7891-7895 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)
 Moody, C.J. *et al.*, *J.C.S. Perkin 1*, 1998, 601-607 (*synth, pmr, cmr*)
 Bertram, A. *et al.*, *Heterocycles*, 2002, **58**, 521-561 (*synth*)

Nostocyclamide M **N-322**
 As Nostocyclamide, N-321 with R = CH₂CH₂SMe

$C_{20}H_{22}N_6O_4S_3$ 506.63
 Isol. from the freshwater cyanobacterium *Nostoc* sp. 31. Allelopathic agent. Needles. λ_{max} 220 (ϵ 41000) (MeOH).

Juettner, F. *et al.*, *Phytochemistry*, 2001, **57**, 613-619 (*isol, pmr, cmr*)


Nostodione A **N-323**
 3,4-Dihydro-3-[(4-hydroxyphenyl)methylene]cyclopent[b]indole-1,2-dione, 9CI
 [158182-28-6]



$C_{18}H_{11}NO_3$ 289.29
 Isol. from the terrestrial blue-green alga *Nostoc commune*. Antimitotic agent. Orange powder. Mp 320° (dec.). λ_{max} 278 (ϵ 7500); 300 (ϵ 9000); 382 (ϵ 25000) (MeOH).

Kobayashi, A. *et al.*, *Z. Naturforsch., C*, 1994, **49**, 464-470 (*isol, uv, ir, pmr, cmr*)

Notoamide C **N-324**
 [937251-97-3]



Absolute Configuration


$C_{26}H_{31}N_3O_4$ 449.549
 Prod. by a marine-derived *Aspergillus* sp. $[\alpha]_D^{25} +23$ (c, 0.25 in MeOH). λ_{max} 247 (log ϵ 4.3); 283 (sh) (log ϵ 3.9); 294 (sh) (log ϵ 3.7); 319 (sh) (log ϵ 3.4) (MeOH).

17-Hydroxy: Norgeamide B
 [873934-44-2]
 $C_{26}H_{31}N_3O_5$ 465.548
 Prod. by a marine-derived *Aspergillus* sp.

17-Methoxy: Norgeamide A
 [873934-43-1]
 $C_{27}H_{33}N_3O_5$ 479.575
 Prod. by a marine-derived *Aspergillus* sp.

Kato, H. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2254-2256 (*isol, struct*)
 Grubbs, A.W. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2257-2261; 2262-2265 (*synth*)

Notoamide D **N-325**
 [937251-98-4]




$C_{26}H_{31}N_3O_4$ 449.549
 Prod. by a marine-derived *Aspergillus* sp. $[\alpha]_D^{27} -163$ (c, 0.32 in MeOH). λ_{max} 237 (log ϵ 4.5); 286 (log ϵ 4); 331 (log ϵ 3.8) (MeOH).

17-Epimer: Norgeamide C
 [873934-45-3]
 $C_{26}H_{31}N_3O_4$ 449.549
 Prod. by a marine-derived *Aspergillus* sp.

17-Epimer, 17-hydroxy: Norgeamide D
 [873934-46-4]
 $C_{26}H_{31}N_3O_5$ 465.548
 Prod. by a marine-derived *Aspergillus* sp.

Kato, H. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2254-2256 (*isol, pmr, cmr*)
 Grubbs, A.W. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2257-2261; 2262-2265 (*synth*)


Noyaine **N-326**
 [110815-78-6]



$C_{21}H_{23}NO_7$ 401.415
 Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. The first C-ring secocularine alkaloid.

Boente, J.M. *et al.*, *Tet. Lett.*, 1986, **27**, 5535 (*uv, ir, pmr, cmr, ms, synth, struct*)

Nudicaulin aglycone **N-327**
 19-(4-Hydroxyphenyl)-10H-1,10-ethenochromeno[2,3-*b*]indole-6,8,18-triol



$C_{23}H_{15}NO_5$ 385.375

6-O-β-D-Glucopyranoside, 18-O-[β-D-glucopyranosyl-(1→2)-β-D-glucopyranoside]: Nudicaulin I
 [875766-61-3]

$C_{41}H_{45}NO_{20}$ 871.801
 Constit. of the flowers of *Papaver nudicaule*.

6-O-β-D-Glucopyranoside, 18-O-[β-D-glucopyranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]: Nudicaulin IV
 [875766-64-6]
 Constit. of the flowers of *Papaver nudicaule*.

6-O-(6-O-Malonyl-β-D-glucopyranoside), 18-O-[β-D-glucopyranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]: **Nudicaulin VII**
[875766-65-7]
C₄₇H₄₉NO₂₆ 1043.895
Constit. of the flowers of *Papaver nudicaule*. [α]_D²⁶ -354 (c, 0.002 in MeOH aq.). λ_{max} 254 (log ε 4.17); 384 (log ε 3.96); 470 (log ε 3.68); 509 (log ε 3.52) (MeOH aq.). λ_{max} 224 (log ε 4.4); 256 (log ε 4.32); 338 (log ε 3.8); 461 (log ε 4.45) (AcOH/MeOH aq.).

10-Epimer, 6-O-β-D-glucopyranoside, 18-O-[β-D-glucopyranosyl-(1→2)-β-D-glucopyranoside]: **Nudicaulin II**
[875766-62-4]
C₄₁H₄₅NO₂₀ 871.801
Constit. of the flowers of *Papaver nudicaule*.


10-Epimer, 6-O-β-D-glucopyranoside, 18-O-[β-D-glucopyranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]: **Nudicaulin III**
[875766-63-5]
C₄₄H₄₇NO₂₃ 957.848
Constit. of the flowers of *Papaver nudicaule*.

10-Epimer, 6-O-(6-O-malonyl-β-D-glucopyranoside), 18-O-[β-D-glucopyranosyl-(1→2)-6-O-malonyl-β-D-glucopyranoside]: **Nudicaulin VIII**
[875766-66-8]
C₄₇H₄₉NO₂₆ 1043.895
Constit. of the flowers of *Papaver nudicaule*. [α]_D²⁶ +370 (c, 0.002 in MeOH aq.). λ_{max} 250 (log ε 4.06); 386 (log ε 3.9); 470 (log ε 3.64); 508 (log ε 3.45) (MeOH aq.). λ_{max} 228 (log ε 4.28); 256 (log ε 4.32); 337 (log ε 3.81); 464 (log ε 4.47) (AcOH/MeOH aq.).

Schliemann, W. *et al.*, *Phytochemistry*, 2006, **67**, 191-201 (isol, cd, uv, pmr, cmr, ms)

Nuevamine

N-328

C₁₉H₁₇NO₅ 339.347

(±)-form [96443-59-3]

Minor alkaloid from *Berberis darwinii* (Berberidaceae). Cryst. (MeOH). Mp 212°. Struct. revised in 1985.

Valencia, E. *et al.*, *Tet. Lett.*, 1984, **25**, 599 (uv, ir, pmr, ms)


Alonso, R. *et al.*, *Tet. Lett.*, 1985, **26**, 2925 (struct, synth)

Moreau, A. *et al.*, *Tetrahedron*, 2004, **60**, 6169-6176 (synth)

Numismine

[38143-08-7]

N-329



C₂₅H₃₃N₃O₄ 439.553
Alkaloid from the seeds of *Lunaria biennis* (Brassicaceae). Cryst. (Me₂CO). Mp 278-280°.


Potier, P. *et al.*, *Bull. Soc. Chim. Fr.*, 1959, 201 (uv, ir)

Poupat, C. *et al.*, *Tetrahedron*, 1972, **28**, 3087 (uv, ir, pmr, struct)

Nummularine C

N-330

1-[2-(Dimethylamino)-1-oxo-3-phenylpropyl]-1,2,3,3a,12,14,15,16a-octahydro-8-methoxy-14-(2-methylpropyl)-5,9-metheno-9H-pyrrolo[3,2-b][1,5,8]oxadiazacyclopentadecine-13,16-dione, 9CI [53947-97-0]




C₃₁H₄₀N₄O₅ 548.681
Alkaloid from the root bark of *Zizyphus nummularia* (Rhamnaceae). Needles (MeOH). Mp 278-280°. [α]_D²⁰ -371 (c, 0.2 in CHCl₃).

Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 3180 (isol, uv, ir, pmr, ms, struct)

Nummularine E

N-331

2-(Dimethylamino)-3-hydroxy-N-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]butanamide, 9CI [58775-95-4]

R¹ = -CH(OH)CH₃R² = -CH₂CH(CH₃)₂

C₂₉H₃₈N₄O₅ 522.643
Alkaloid from the root bark of *Zizyphus nummularia* and the bark of *Zizyphus hysodrica* (Rhamnaceae). Cryst.

(MeOH). Mp 278-279°. [α]_D²⁰ +12 (c, 0.02 in MeOH).

O-Ac:

Cryst. (CHCl₃). Mp 230-232°.

Dihydro:

Cryst. (MeOH). Mp 282°. [α]_D²⁰ +14 (c, 0.02 in MeOH).


Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (isol, uv, ir, pmr, ms, struct)

Tschesche, R. *et al.*, *Pharmazie*, 1981, **36**, 511 (isol, uv, ir, pmr, ms)

Nummularine F

N-332

1-[(Dimethylamino)acetyl]-2,3,3a,13,14,15a-hexahydro-13-(1-methylpropyl)-5,8-ethenopyrrolo[3,2-b][1,5,8]oxadiazacyclopentadecine-12,15(1H,11H)-dione, 9CI [58775-98-7]



Absolute Configuration

C₂₃H₃₂N₄O₄ 428.53
Alkaloid from the root bark of *Zizyphus nummularia* (Rhamnaceae). Cryst. (MeOH). Mp 120°. [α]_D²⁰ -204 (c, 0.2 in MeOH).

Dihydro:

Cryst. (MeOH). Mp 175°. [α]_D²⁰ -159 (c, 0.23 in MeOH).


Tschesche, R. *et al.*, *Tetrahedron*, 1975, **31**, 2944-2947 (isol, uv, ir, pmr, ms, struct)

Heffner, R.J. *et al.*, *J.A.C.S.*, 1992, **114**, 10181-10189 (synth, cmr)

Nummularine G

N-333

4-[3-Methyl-4-(2-methylpropyl)-5-oxo-1-imidazolidinyl]-7-(2-methylpropyl)-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraene-5,8-dione, 9CI [64309-21-3]

C₃₁H₄₀N₄O₄ 532.681

Alkaloid from the bark of *Zizyphus nummularia* (Rhamnaceae). Light-yellow needles (CHCl₃/petrol). Mp 174-175°. [α]_D²⁰ -133 (c, 0.085 in MeOH). Related to Sativanine B, S-93.


Dihydro:

Powder. Mp 206-209°. [α]_D²⁰ -98 (c, 0.2 in MeOH).

Tschesche, R. *et al.*, *Chem. Ber.*, 1977, **110**, 2649 (isol, uv, ir, pmr, ms, struct)

Nummularine K N-334

α -(Dimethylamino)-N-[3-(1-methyl-ethyl)-7-(2-methylpropyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-1H-indole-3-propanamide, 9CI
[64350-62-5]



$C_{33}H_{43}N_5O_4$ 573.734
Alkaloid from the bark of *Zizyphus nummularia* and from *Zizyphus xylopyra* and *Discaria longispina*. Powder (MeOH). Mp 237-239°. $[\alpha]_D^{20}$ -45 (c, 0.04 in MeOH).

Dihydro:

Powder. Mp 283-285°. $[\alpha]_D^{20}$ +12 (c, 0.08 in MeOH).

Stereoisomer(?): Discarine X

[164577-53-1]

$C_{33}H_{43}N_5O_4$ 573.734

Alkaloid from root bark of *Discaria longispina* (Rhamnaceae). Mp 295-298°. $[\alpha]_D^{25}$ -184 (c, 0.5 in MeOH). No stereochem. determined. May be identical with Nummularine K.

Tschesche, R. et al., *Chem. Ber.*, 1977, **110**, 2649 (Nummularine K)

Machado, E.C. et al., *J. Nat. Prod.*, 1995, **58**, 548 (Discarine X)

Nummularine M N-335

[93754-86-0]

As Nummularine E, N-331 with

$R^1 = R^2 = -CH(CH_3)CH_2CH_3$


$C_{31}H_{42}N_4O_4$ 534.697

Alkaloid from the stem bark of *Zizyphus nummularia* (Rhamnaceae). Amorph. powder. Mp 263-265°. $[\alpha]_D$ -46.66 (c, 0.1 in $CHCl_3$).

Pandey, V.B. et al., *Phytochemistry*, 1984, **23**, 2118 (isol, uv, ir, ms, struct)

Nummularine N N-336

[93754-85-9]




$C_{32}H_{41}N_5O_6$ 591.706

Alkaloid from the stem bark of *Zizyphus nummularia* (Rhamnaceae). Cryst. Mp 243-245°. $[\alpha]_D$ not reported. Mol. formula given as $C_{31}H_{41}N_5O_6$, apparently incorrectly, in the paper.

Pandey, V.B. et al., *Phytochemistry*, 1984, **23**, 2118 (isol, uv, ir, pmr, ms, struct)

Nummularine P N-337

[110187-25-2]



$C_{29}H_{43}N_5O_6$ 557.689

Alkaloid from the stem bark of *Zizyphus nummularia* and *Zizyphus rugosa* (Rhamnaceae). Mp 143-144°.

N^o-Formyl: Rugosanine A

[110200-25-4]

$C_{30}H_{43}N_5O_7$ 585.699


Alkaloid from stem bark of *Zizyphus nummularia* and *Zizyphus rugosa* (Rhamnaceae). Granules (MeOH). Mp 237-240° dec. Not correlated with Nummularine P, could differ stereochemically.

Dwivedi, S.P.D. et al., *J. Nat. Prod.*, 1987, **50**, 235 (isol, uv, ir, pmr, struct)

Pandey, V.B. et al., *Phytochemistry*, 1988, **27**, 1915 (Rugosanine A)

Nummularine R N-338

[113836-69-4]



$C_{33}H_{41}N_5O_5$ 587.717

Alkaloid from *Zizyphus nummularia* and *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Mp 134-135°.

Stereoisomer (?): Daechuine S10

[123089-24-7]

$C_{33}H_{41}N_5O_5$ 587.717

Alkaloid from stem bark of *Zizyphus*


jujuba var. *inermis* (Rutaceae). Mp 126-128°. $[\alpha]_D$ -381.5. Could be identical with Nummularine R (similar Mp). Incorrectly registered by CAS with Tyr sidechain in place of Try.

Devi, S. et al., *Phytochemistry*, 1987, **26**, 3374-3375 (Nummularine R)

Han, B.H. et al., *Pure Appl. Chem.*, 1989, **61**, 443-448 (Daechuine S10)

Nuphacristine N-339

6-(3-Furanyl)-1,6,7,8,9,9a-hexahydro-9-(hydroxymethyl)-2H-quinolizine-3-carboxaldehyde, 9CI
[119459-68-6]




$C_{15}H_{19}NO_3$ 261.32

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). $[\alpha]_D^{20}$ +65.9 (EtOH).

Cybulski, J. et al., *Phytochemistry*, 1988, **27**, 3339 (isol, pmr, cmr, ms, struct)

Nupharamine N-340

6-(3-Furanyl)- $\alpha,\alpha,3$ -trimethyl-3-piperidinepropanol, 9CI
[17812-38-3]
[53956-48-2 (\pm)-form]



Absolute Configuration

$C_{15}H_{25}NO_2$ 251.368

Alkaloid from *Nuphar japonicum* (Nymphaeaceae). Bp₁ 130-134°. $[\alpha]_D$ -35.4.

Picolonate: Mp 168°.

Me ether: O-Methylnupharamine

$C_{16}H_{27}NO_2$ 265.395

Minor alkaloid from *Nuphar japonicum* (Nymphaeaceae). Mp 48-50°. $[\alpha]_D^{25}$ -52.56 (EtOH).

Et ether: O-Ethylnupharamine

$C_{17}H_{29}NO_2$ 279.422

Minor alkaloid from *Nuphar japonicum* (Nymphaeaceae). Bp_{0.0001} 95-100°. $[\alpha]_D^{25}$ -48.25 (EtOH).

3''-Deoxy, 2'',3''-didehydro: Nuphenine

[4850-01-5]

$C_{15}H_{23}NO$ 233.353

Major alkaloid from the rhizomes of the North American yellow water lily *Nuphar variegatum* (Nymphaeaceae). $[\alpha]_{H_2O}^{25}$ -23 (MeOH).

3''-Deoxy, 4''-hydroxy (3''-R): 6-(3-Furanyl)- β ,3-dimethyl-2-piperidinebutanol.

Secodihydrocastoramine

[56283-43-3]

C₁₅H₂₅NO₂ 251.368Alkaloid from the roots of *Nuphar japonicum*.3''-Deoxy, 2'',3''-didehydro, 4''-hydroxy (2''E-): 4-[6-(3-Furanyl)-3-methyl-2-piperidinyl]-2-methyl-2-buten-1-ol. **Nupharamine**

[1630-49-5]

C₁₅H₂₃NO₂ 249.352Alkaloid from *Nuphar japonicum* and *Nuphar luteum* ssp. *variegatum* (Nymphaeaceae). Viscous liq. Bp₃ 150-153°. [α]_D¹⁹ -60.5 (CHCl₃).3-Epimer: **3-Epinupharamine**

[31146-50-6]

[53956-49-3 (\pm)-form]C₁₅H₂₅NO₂ 251.368Alkaloid from the rhizomes of *Nuphar variegatum* (Nymphaeaceae). Pale yellow oil. [α]_D²³ -24.8 (c, 1.45 in MeOH).

3-Epimer, N-Ac:

Cryst. (Me₂CO). Mp 107-109°.

3-Epimer, 3''-deoxy, 2'',3''-didehydro:

Anhydronupharamine

[4849-88-1]

C₁₅H₂₃NO 233.353Alkaloid from the rhizomes of *Nuphar japonicum*. Dehydration product of (-)-Nupharamine (Nymphaeaceae). Bp₄ 108-109°. [α]_D²⁵ -62.5 (c, 1.0 in CHCl₃). [α]_D -59 (solvent not specified).3-Epimer, 3''-deoxy, 2'',3''-didehydro, 4''-hydroxy: **3-Epinupharamine**

[29073-35-6]

C₁₅H₂₃NO₂ 249.352Minor alkaloid from the rhizomes of *Nuphar luteum* ssp. *variegatum* (Nymphaeaceae). [α]_D²⁵ -47.5 (c, 0.52 in 95% EtOH).Arata, Y. et al., *Yakugaku Zasshi*, 1959, **79**, 127-128; 729-734; 734-737; 1967, **87**, 1094-1095 (*Nupharamine*, *Anhydronupharamine*, *isol*, *ir*, *struct*)Kawasaki, I. et al., *Bull. Chem. Soc. Jpn.*, 1963, **36**, 1474 (*config*)Kotake, M. et al., *Nippon Kagaku Zasshi*, 1963, **84**, 160-162; *CA*, **60**, 6891a (*derivis*)Arata, Y. et al., *Chem. Pharm. Bull.*, 1965, **13**, 1247; 1365 (*Nupharamine*)Barchet, R. et al., *Tet. Lett.*, 1965, 4229-4232 (*Nupharine*)Wong, C.F. et al., *Phytochemistry*, 1970, **9**, 1851 (*3-Epinupharamine*)Forrest, T.P. et al., *Can. J. Chem.*, 1971, **49**, 1774-1775 (*3-Epinupharamine*)Khaleque, A. et al., *CA*, 1975, **82**, 28643d; 1979, **90**, 204316a (*Secodihydrocastoramine*)Itatani, Y. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2521-2524 (*cmr*)Szychowski, J. et al., *Can. J. Chem.*, 1977, **55**, 3105-3110 (*synth*)Sabat, M. et al., *Can. J. Chem.*, 1977, **55**, 3111-3117 (*3-Epinupharamine*, *cryst struct*)Shimizu, I. et al., *Chem. Lett.*, 1990, 777-778 (*synth*, *cmr*)Leniewski, A. et al., *Coll. Czech. Chem. Comm.*, 1991, **56**, 1309-1316 (*synth*)Aoyagi, S. et al., *Tet. Lett.*, 1991, **32**, 4325-4328 (*synth*, *abs config*)Honda, T. et al., *Chem. Comm.*, 1994, 499-500 (*synth*)Miyazawa, M. et al., *J. Agric. Food Chem.*, 1998, **46**, 1059-1063 (*Nupharine*, *isol*, *pmr*, *cmr*, *ms*)Barluenga, J. et al., *J.O.C.*, 1999, **64**, 3736-3740 (*synth*)Davies, F.A. et al., *J.O.C.*, 2006, **71**, 4222-4226 (*synth*) **α -Nupharidine**

N-341

C₁₅H₂₃NO 233.353*Nuphar* alkaloid; probably identical with Deoxynupharidine, D-227. Props. of characterisation derivs. agree, but status of 'dihydro deriv.' is not clear. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Viscous liq. Bp₂ 121-121.5°. [α]_D¹⁸ -112.1.*Hydrochloride*: Mp 258-259°.*Picrate*: Mp 165-167°.*Dihydro*: Mp 240-242° (as hydrochloride).Achmatowicz, O. et al., *Pol. J. Chem. (Rocz. Chem.)*, 1939, **19**, 493-506 (*isol*) **β -Nupharidine**

N-342


C₁₅H₂₃NO 233.353*Nuphar* alkaloid. Struct. unknown. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Viscous liq. Bp₂ 140°.*Hydrochloride*: Mp 269-270°.*Picrate*: Mp 152-153°.Achmatowicz, O. et al., *Pol. J. Chem. (Rocz. Chem.)*, 1939, **19**, 493-506 (*isol*)**Nupharine[†]**

N-343

[1400-60-8]


C₁₈H₂₄N₂O₂ 300.4*Nuphar* alkaloid. Struct. unknown. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Amorph. Mp 65°.Grüning, W. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1882, **20**, 589 (*isol*)Novikova, S.I. et al., *CA*, 1964, **60**, 2041g (*isol*)**Nuttingine A**

N-344

N⁷-(3,7,11,15-Tetramethyl-13-oxo-2,6,10-hexadecatrienyl)theophylline [945458-41-3]C₂₇H₄₀N₄O₃ 468.638Alkaloid from *Euplexaura nuttingi*. Oil.*A*^{11'}-Isomer (Z-): **Nuttingine B** [945458-42-4]C₂₇H₄₀N₄O₃ 468.638Alkaloid from *Euplexaura nuttingi*. Oil.Sorek, H. et al., *J. Nat. Prod.*, 2007, **70**, 1104-1109 (*Nuttingins A,B*)**Nuttingine D**

N-345

[945458-44-6]

C₂₇H₄₂N₄O₂ 454.654Alkaloid from *Euplexaura nuttingi*. Oil.N,2-Didehydro: **Nuttingine F**

[945458-46-8]

C₂₇H₄₁N₄O₂[⊕] 453.647Alkaloid from *Euplexaura nuttingi*. Oil. Positive charge delocalised over N-1 and N-3.*A*^{10'}-Isomer (E-): **Nuttingine C**

[945458-43-5]


C₂₇H₄₂N₄O₂ 454.654Alkaloid from *Euplexaura nuttingi*. Oil.*11'*Z-Isomer: **Nuttingine E**

[945458-45-7]

C₂₇H₄₂N₄O₂ 454.654Alkaloid from *Euplexaura nuttingi*. Oil.Sorek, H. et al., *J. Nat. Prod.*, 2007, **70**, 1104-1109 (*isol*, *pmr*, *cmr*)**Nybomycin**

N-346

8-(Hydroxymethyl)-6,11-dimethyl-2H,4H-oxazolo[5,4,3-ij]pyrido[3,2-g]quinoline-4,10(11H)-dione, 9CI, 8CI [30408-30-1]

C₁₆H₁₄N₂O₄ 298.298Prod. by Streptomycete A 717 *isol*. from Missouri soil. Antibiotic showing anti-phage and antibacterial props. Needles (AcOH). Sol. conc. acids; fairly sol. DMF; poorly sol. H₂O, MeOH, hexane. Mp 325-330°. Subl. 15 250. λ_{\max} 222 (ϵ 18400); 236 (sh) (ϵ 15000); 264 (ϵ 11150); 286 (ϵ 40500); 296 (sh) (ϵ 32100); 353 (ϵ 11900); 367 (ϵ 12200) (MeOH) (Derep). ▶LD₅₀ (mus, ipr) 650 mg/kg.*Ac*:Cryst. (CHCl₃/EtOH). Mp 236-237°.*Deoxy*: **Deoxynybomycin**. 6,8,11-Trimethyl-2H,4H-oxazolo[5,4,3-ij]pyrido[3,2-g]quinoline-4,10(11H)-dione, 9CI [27259-98-9]C₁₆H₁₄N₂O₃ 282.298From *Streptomyces hyalimum*. Antibiotic active against gram-positive organisms. Inhibits topoisomerase I. Apoptosis inducer. Selective anti-tumor agent. Needles. Fairly sol. AcOH; poorly sol. H₂O, MeOH, DMF, DMSO, hexane. Mp 335° dec.

λ_{\max} 220 (ϵ 27550); 264 (ϵ 36300); 285 (ϵ 46800); 294 (sh) (ϵ 38900); 353 (ϵ 1200); 368 (ϵ 1290) (MeOH) (Derep).
 ► LD₅₀ (mus, ipr) 650 mg/kg. RQ5836000

***l'*-Hydroxy: Hydroxybomycin**

[63582-81-0]

C₁₆H₁₄N₂O₅ 314.297

From *Streptomyces* spp. Antibiotic active against gram-positive organisms. Pale yellow cryst. (DMF). Mp 360°.

Strelitz, F. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1955, **41**, 620-624 (*isol*)

Rinehart, K.L. *et al.*, *J.A.C.S.*, 1961, **83**, 3729-3731; 1970, **92**, 6994-6995 (*struct, ir, pmr*)

Naganawa, H. *et al.*, *J. Antibiot.*, 1970, **23**, 365-368 (*Deoxybomycin, isol*)

Forbis, R.M. *et al.*, *J.A.C.S.*, 1973, **95**, 5003-5013 (*synth, uv, ir, ms, nmr, Deoxybomycin*)

Nazdan, A.M. *et al.*, *J. Antibiot.*, 1977, **30**, 523-524 (*Hydroxybomycin*)


Nazdan, A.M. *et al.*, *J.A.C.S.*, 1977, **99**, 4647-4654 (*biosynth, cmr*)

Egawa, K. *et al.*, *Biol. Pharm. Bull.*, 2000, **23**, 1036-1040 (*activity*)

Nymphaedaline

[385366-59-6]

N-347

C₂₇H₂₉NO₇ 479.529

Alkaloid from the bark of *Hernandia nymphaeifolia*. Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +34 (c, 0.11 in CHCl₃). λ_{\max} 204 (log ϵ 4.77); 230 (log ϵ 4.43); 279 (log ϵ 4.07); 309 (sh) (log ϵ 3.74) (EtOH).

Chen, J.-J. *et al.*, *Planta Med.*, 2001, **67**, 593-598

Nympheine

N-348

*Nymphaeine*C₁₄H₂₃NO₂ 237.341


Struct. unknown. Alkaloid from the roots of *Nymphaea alba* (Nymphaeaceae). Amorph. Mp 76-77°.

Hydrochloride: Dec. at 230° without melting.

Bureš, E. *et al.*, *Cas. Cesk. Lek.*, 1934, **14**, 129-135; *CA*, **28**, 5460 (*isol*)

Obaberine

6,6',7,12'-Tetramethoxy-2,2'-dimethoxy-yacanthan, 9Cl. O-Methoxyacanthine [1263-80-5]



$C_{38}H_{42}N_2O_6$ 622.76

Alkaloids in this entry have the 1R,1'S-configuration (Shamma-Moniot numbering convention; older CAS numbering put the primes on the opposite rings). Alkaloids in diastereomeric series are given in other entries; for enantiomeric alkaloids see Gyrolidine, G-235. Alkaloid from *Thalictrum lucidum*, *Thalictrum minus* Race B, *Thalictrum minus* var. *microphyllum*, *Thalictrum rugosum*, *Pycnarrhena longifolia*, *Berberis tschonoskyana*, *Berberis laurina*, *Dehaasia triandra*, *Mahonia repens* and *Albertisia papuana* (Ranunculaceae). Mp 139-140° Mp 199-200°. $[\alpha]_D^{26} +169$ (c, 0.147 in MeOH). λ_{max} 280 (MeOH).

Hydrochloride (1:2):

Needles + 3H₂O. Mp 260-261°.

Dipicrate:

Cryst. + 1H₂O. Mp 178-180° dec.

N²,N^{2'}-Di-Me:

Needles + 2H₂O. Mp 256-258° dec.

N²-De-Me, N²-β-oxide: 2-Norobaberine 2'-β-N-oxide

[149155-48-6]

$C_{37}H_{40}N_2O_7$ 624.732

Alkaloid from seeds of *Anisocycla cymosa* (Menispermaceae). Amorph. powder. $[\alpha]_D^{20} +158$ (c, 0.31 in CHCl₃). λ_{max} 212 ; 284 (MeOH).

N²-De-Me: 2'-Norobaberine

[6016-52-0]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the tubers of *Stephania pierrii* (Menispermaceae).

O⁶-De-Me: Stephibaberine

[123930-97-2]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the tubers of *Stephania pierrii* (Menispermaceae). $[\alpha]_D +207$ (c, 0.17 in CHCl₃). λ_{max} 241 (sh) (log ε 4.36); 283 (log ε 4) (MeOH).

O⁷-De-Me: Homoaromoline. Thlrugosamine. Homothalicrine. N-Methyl-daphnandrine

[17132-74-0]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the stems of *Albertisia papuana* and *Abuta splendida*, the bark of *Doryphora aromatica*, the stems and roots of *Pycnarrhena longifolia* and *Arcangelisia flava*, the roots of *Thalictrum thunbergii*, *Thalictrum rugosum* and *Limaciopsis loangensis* and the tubers of *Stephania erecta* (Menispermaceae, Monimiaceae, Ranunculaceae). Cryst. (MeOH). Mp 235-237°. $[\alpha]_D^{19} +409$ (CHCl₃). Epimer of Limacusine, L-175. Identity with

Thlrugosamine established in 1984.

λ_{max} 282 (log ε 3.91) (MeOH).

►MS9272000

O⁷-De-Me, N²-de-Me: Daphnandrine.

12-O-Methyl-daphnoline

[1183-76-2]

$C_{36}H_{38}N_2O_6$ 594.706

Alkaloid from the bark of *Daphnandra micrantha* and *Doryphora aromatica* (Monimiaceae). Cryst. (CHCl₃, MeOH or MeOH/Et₂O). Mp 280° (270° dec.). $[\alpha]_D^{25} +480$ (c, 1.2 in CHCl₃). λ_{max} 284 (ε 8800) (MeOH) (Berdy).

O¹²-De-Me: Oxyacanthine

[548-40-3]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from a wide variety of *Berberis* and *Mahonia* spp., including *Berberis aquifolium*, *Berberis floribunda*, *Berberis integerrima*, *Berberis julianae*, *Berberis lambertii*, *Berberis oblonga*, *Berberis orthobotrys*, *Berberis thunbergii*, *Berberis tschonoskyana*, *Berberis vulgaris* (barberry), *Mahonia acanthifolia*, *Mahonia borealis*, *Mahonia fortunei*, *Mahonia griffithii*, *Mahonia leschenaultii*, *Mahonia manipurensis*, *Mahonia repens*, *Mahonia sikkimensis* and *Mahonia simonsii*; also occurs in *Albertisia papuana*, *Cocculus laeabe*, *Magnolia compressa*, *Thalictrum lucidum* and *Xanthorhiza simplicissima* (Berberidaceae, Menispermaceae, Magnoliaceae, Ranunculaceae). Shows antibacterial and antitubercular activity, and *in vitro* antineoplastic activity against HeLa-S₃ cells. Symphatholytic agent, adrenaline antagonist, vasodilator. Cryst. (petrol). Mp 212-214°. $[\alpha]_D^{20} +285.6$ (c, 0.5 in CHCl₃). Log P 7.72 (uncertain value) (calc). Epimer of Repandine, R-46. λ_{max} 223 ; 283 (log ε 3.95) (MeOH) (Berdy).

►LD₅₀ (mus, ipr) 50 mg/kg. RR6475000

O¹²-De-Me, N²-Me: Oblongamine

[63511-70-6]

[63504-42-7]

$C_{38}H_{43}N_2O_6^{\oplus}$ 623.767

Alkaloid from *Berberis oblonga* (Berberidaceae). Mp 198-200° (as iodide).

O¹²-De-Me, N²-de-Me: Sepeerine. Oco-

teamine. 7'-O-Methyl-daphnoline

[6787-93-5]

$C_{36}H_{38}N_2O_6$ 594.706

Alkaloid from the bark of *Ocotea rodiaei* (Lauraceae). Exhibits antiparasitic and cytotoxic activities. Platelets (MeOH). Mp 221.5-222.5°. $[\alpha]_D^{25} +392$ (c, 0.29 in CHCl₃).

O¹²-De-Me, N²-de-Me: 2'-Noroxycanthine

$C_{36}H_{38}N_2O_6$ 594.706

Alkaloid from *Thalictrum cultratum*. Amorph. $[\alpha]_D +125$ (c, 0.1 in MeOH).

O⁶, O¹²-Di-de-Me: Baluchistine

[72154-62-2]

$C_{36}H_{38}N_2O_6$ 594.706

Alkaloid from *Berberis baluchistanica* (Berberidaceae). Cryst. (MeOH). Mp 222-224°. $[\alpha]_D^{25} +333$ (c, 0.075 in

MeOH). λ_{max} 283 (log ε 3.67) (EtOH).

O⁷, O¹²-Di-de-Me: Aromoline. 2-N-

Methyl-daphnoline. Thalicrine

[519-53-9]

Alkaloid from *Daphnandra aromatica*, *Daphnandra tenuipes*, *Doryphora aromatica*, *Triclisia patens*, *Thalictrum lucidum*, *Thalictrum thunbergii*, *Thalictrum rugosum*, *Berberis orthobotrys*, *Abuta splendida*, *Albertisia papuana* and *Pycnarrhena longifolia* (Monimiaceae, Menispermaceae, Ranunculaceae, Berberidaceae). Plates + 1CHCl₃ or needles (CHCl₃). Mp 174-175° (solvate) Mp 198-202° (anhyd.). $[\alpha]_D^{22} +249.5$ (c, 1.0 in Py). λ_{max} 208 (log ε 4.94); 228 (sh) (log ε 4.69); 285 (log ε 3.95) (MeOH).

►RR6425000

O⁷, O¹²-Di-de-Me, N²-de-Me:

Daphnoline†. Trilobamine

[479-36-7]

$C_{35}H_{36}N_2O_6$ 580.679

Alkaloid from the bark of *Daphnandra micrantha*, *Daphnandra aromatica* and *Doryphora aromatica*, the roots and stems of *Pycnarrhena longifolia*, the stems of *Albertisia papuana*, and from *Cocculus trilobus* (Monimiaceae). Vasodilator, oedematous agent, CNS depressant and respiratory paralytic. Cryst. (CHCl₃). Mp 196-197°. $[\alpha]_D^{15} +356.6$ (AcOH). λ_{max} 285 (log ε 3.95) (MeOH).

O⁷, O¹²-Di-de-Me, di-N-de-Me: N,N'-Bisnoraromoline

[38962-93-5]

$C_{34}H_{34}N_2O_6$ 566.652

Alkaloid from leaves and stems of *Pycnarrhena ozantha* (Menispermaceae). Needles (Me₂CO). Sol. CHCl₃-MeOH; fairly sol. Me₂CO. Mp 206° (*in vacuo*). $[\alpha]_D^{25} +177$ (c, 0.5 in 1M HCl). λ_{max} 285 (ε 7520) (EtOH).

Bick, I.R.C. et al., *J.C.S.*, 1949, 2767-2774; 1960, 4928-4931; 1961, 1896-1903

(*Daphnandrine, Daphnoline, Aromoline, isol, uv, pmr, struct*)

Fujita, E. et al., *Yakugaku Zasshi*, 1952, **72**, 213-216; *CA*, **47**, 6429a (*Obaberine, Oxyacanthine, struct, synth*)

Inubushi, Y. et al., *Pharm. Bull.*, 1955, **3**, 384-387 (*Trilobamine*)

Fujita, E. et al., *Bull. Inst. Chem. Res., Kyoto Univ.*, 1964, **42**, 235; *CA*, **62**, 5310 (*Aromoline, Homoaromoline*)

Hearst, P.J. et al., *J.O.C.*, 1964, **29**, 466-470 (*Sepeerine*)

Battersby, A.R. et al., *J.C.S.*, 1965, 2239-2247 (*Obaberine, Oxyacanthine, ord*)

Knapp, J.E. et al., *J. Pharm. Sci.*, 1967, **56**, 139-141 (*Oxyacanthine, uv, ir, pmr*)

Falco, M.R. et al., *Tet. Lett.*, 1968, 1953-1959 (*pmr*)

Loder, J.W. et al., *Aust. J. Chem.*, 1972, **25**, 2193-2197 (*Bisnoraromoline*)

Mitscher, L.A. et al., *Experientia*, 1972, **28**, 500 (*Thlrugosamine, uv, cd, pmr, ms*)

Baldas, J. et al., *J.C.S. Perkin I*, 1972, 592-596 (*Obaberine, Oxyacanthine, ms*)

Kuroda, H. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2413-2420 (*Oxyacanthine, pharmacol*)

Wu, W.-N. et al., *J. Nat. Prod.*, 1976, **39**, 378-379; 1978, **41**, 257-270; 1980, **43**, 143-150


(*Aromoline, Thlrugosamine, isol, synth*)

Saá, J.M. et al., *J.O.C.*, 1976, **41**, 317

(*Aromoline, Homoaromoline, ir, pmr, ms*)


Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 1636-1644 (*synth, pmr, ms*)
 Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 80-83; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 68-70 (*Oblongamine*)
 Bhakuni, D.S. *et al.*, *J.C.S. Perkin 1*, 1978, 1318-1321 (*biosynth*)
 Miana, G.A. *et al.*, *Experientia*, 1979, **35**, 1137-1138 (*Baluchistine*)
 Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1980, **33**, 225-228 (*Aromoline, Daphnoline, Daphnandrine, Homoaromoline*)
 Siwon, J. *et al.*, *Phytochemistry*, 1981, **20**, 323 (*Aromoline, Homoaromoline, Daphnoline*)
 Koike, L. *et al.*, *J.O.C.*, 1982, **47**, 4351-4353 (*cmr*)
 Nakova, E. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 86-91; 91-95; *Chem. Nat. Compd. (Engl. Transl.)*, 83-87; 88-91 (*synth*)
 Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 721-725 (*2'-Noroxyacanthine*)
 Tantisewie, B. *et al.*, *J. Nat. Prod.*, 1989, **52**, 846-851 (*2'-Norobaberine, Stephibaberine*)
 Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1993, **56**, 618-620 (*2-Norobaberine-2'-β-N-oxide*)
 Sivý, J. *et al.*, *Acta Cryst. C*, 1996, **52**, 1479-1482 (*Oxyacanthine, cryst struct*)
 Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*Thalrugosamine, pmr, cmr*)
 Mahiou, V. *et al.*, *Phytochemistry*, 2000, **54**, 709-716 (*Sepeerine, activity*)
 Chen, C.-K. *et al.*, *Chin. Pharm. J. (Taipei)*, 2003, **55**, 35-47 (*Aromoline, Daphnandrine, Daphnoline, Homoaromoline*)

Obesine O-2
 [143114-83-4]



$C_{16}H_{17}NO_4$ 287.315
 Alkaloid from aerial parts and bulbs of *Narcissus ovesus* (Amaryllidaceae). $[\alpha]_D^{20}$ -5.6 (c, 0.69 in MeOH).
 Viladomat, F. *et al.*, *J. Nat. Prod.*, 1992, **55**, 804 (*isol, pmr, cmr, ms, struct*)

Obovamine O-3
 [172670-51-8]




Absolute Configuration

$C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from stem bark of *Stemmadenia obovata*. Amorph. $[\alpha]_D^{25}$ +134 (c, 0.10 in $CHCl_3$).
 Madinaveitia, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 185-189 (*isol, uv, ir, pmr, cmr, ms, struct*)

Obscuridine O-4
 Struct. unknown. Alkaloid from the root bark of *Rauwolfia obscura* (Apocynaceae). Cryst. (EtOH). Mp 228° dec.

Roland, M. *et al.*, *J. Pharm. Belg.*, 1959, **14**, 347-375; *CA*, **54**, 15834h

β-Obscurine O-5
 [467-79-8]



Absolute configuration

$C_{17}H_{24}N_2O$ 272.389
 Alkaloid from *Lycopodium selago*, *Lycopodium annotinum*, *Lycopodium flabelliforme* and *Lycopodium obscurum* var. *dendroideum* (Lycopodiaceae). Mp 322-323°.

Perchlorate: Mp 322° dec.

Picrate: Mp 254° dec.

N-De-Me: De-N-methyl-β-obscurine [107965-41-3]

$C_{16}H_{22}N_2O$ 258.363
 Alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Mp 290-293°. $[\alpha]_D$ -34 (c, 0.21 in MeOH).

2,3-Dihydro: α-Obscurine

[596-55-4]
 $C_{17}H_{26}N_2O$ 274.405
 From *Lycopodium annotinum*, *Lycopodium clavatum*, *Lycopodium flabelliforme*, *Lycopodium obscurum* var. *dendroideum* and *Lycopodium selago* (Lycopodiaceae). Mp 283-284°.

2,3-Dihydro, picrate: Mp 135°.

2,3-Dihydro, N-de-Me: De-N-methyl-α-obscurine [34399-44-5]

$C_{16}H_{24}N_2O$ 260.378
 Alkaloid from *Lycopodium clavatum*, *Lycopodium fawcettii*, *Lycopodium flabelliforme* and *Lycopodium alpinum* (Lycopodiaceae). Needles (Me₂CO). Mp 266-268°.

12-Hydroxy, 2,3-dihydro, N-de-Me: Hydroxy-de-N-methyl-α-obscurine

$C_{16}H_{24}N_2O_2$ 276.378
 Alkaloid from *Lycopodium flabelliforme*. Cryst. (MeOH/Me₂CO). Mp 300-305° dec.

12-Epimer, 2,3-dihydro: Sauroxine

[3279-74-1]
 $C_{17}H_{26}N_2O$ 274.405
 From *Lycopodium saururus*. Mp 200-201°. $[\alpha]_D^{20}$ -72 (c, 0.79 in EtOH). p*K*_a 8.1 (50% MeOH).

Moore, B.P. *et al.*, *Can. J. Chem.*, 1953, **31**, 952 (*isol, uv, ir*)

Ayer, W.A. *et al.*, *Tetrahedron*, 1962, **18**, 567; 1965, **21**, 2169 (*isol, ir, pmr, struct, Sauroxine*)

Alam, S.N. *et al.*, *Can. J. Chem.*, 1964, **42**, 2456 (*Hydroxy-de-N-methyl-α-obscurine*)
 Schumann, D. *et al.*, *Annalen*, 1983, 220 (*synth, α-Obscurine, De-N-methyl-α-obscurine*)


Ayer, W.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 1077 (*isol, pmr, ms, Des-N-methyl-β-obscurine*)

Obscurine† O-6

Struct. unknown. Alkaloid from the root bark of *Rauwolfia obscura* (Apocynaceae). Fine needles (EtOH/NH₃). Mp 255° dec. $[\alpha]_D^{21}$ +250.4 (c, 0.5 in $CHCl_3$).

Roland, M. *et al.*, *J. Pharm. Belg.*, 1959, **14**, 347-375; *CA*, **54**, 15834h

Obscurinervidine O-7
 [7168-67-4]



R = CH₃

$C_{24}H_{28}N_2O_5$ 424.496
 Alkaloid from *Aspidosperma obscurinervium* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 206-207°. $[\alpha]_D^{25}$ -39 (c, 0.62 in $CHCl_3$).

14,15-Dihydro: Dihydroobscurinervidine

$C_{24}H_{30}N_2O_5$ 426.511
 From *Aspidosperma obscurinervium* (Apocynaceae). Mp 189-190° dec. $[\alpha]_D^{27}$ -44 (c, 0.85 in $CHCl_3$).

10-Demethoxy: Neblinine, 8CI

[6872-64-6]
 $C_{23}H_{26}N_2O_4$ 394.469
 Alkaloid from *Aspidosperma neblinae* (Apocynaceae). Cryst. (hexane). Mp 256-258° dec. $[\alpha]_D^{27}$ -14 (c, 1.19 in $CHCl_3$).

Brown, K.S. *et al.*, *J.A.C.S.*, 1964, **86**, 2451

(*Neblinine, isol, uv, ir, pmr, ms, struct*)

Kahrl, J. *et al.*, *Tet. Lett.*, 1971, 2527

(*Obscurinervidine, cryst struct*)

Obscurinervine O-8
 [7097-00-9]

As Obscurinervidine, O-7 with

R = CH₂CH₃

$C_{25}H_{30}N_2O_5$ 438.522
 Alkaloid from *Aspidosperma obscurinervium* (Apocynaceae). Mp 203-205° dec. $[\alpha]_D^{27}$ -54 (c, 0.95 in $CHCl_3$).

14,15-Dihydro: Dihydroobscurinervine

$C_{25}H_{32}N_2O_5$ 440.538
 Alkaloid from *Aspidosperma obscurinervium* (Apocynaceae). Cryst. (Me₂CO/hexane). Mp 184-185° dec. $[\alpha]_D^{25}$ -61 (c, 0.92 in $CHCl_3$).


Brown, K.S. *et al.*, *J.A.C.S.*, 1964, **86**, 2451

(*isol, uv, ir, pmr, ms, struct*)

Harper, J.K. *et al.*, *J. Crystallogr. Spectrosc. Res.*, 1993, **23**, 1005 (*cryst struct, abs config, cmr*)

Harper, J.K. *et al.*, *J.C.S. Perkin 2*, 1996, 91 (*pmr, cmr*)

Obscurinine O-9
 [113883-30-0]



C₁₇H₂₄N₂O 272.389


Alkaloid from *Lycopodium obscurum* (Lycopodiaceae). Off-white plates (Me₂CO/hexane). Mp 128–130° (124–125.2°). [α]_D +242 (c, 0.5 in MeOH). Possibly an artifact.

Hu, T. et al., *Tet. Lett.*, 1987, **28**, 5993 (*isol, ir, pmr, cmr, ms, cryst struct*)

Ayer, W.A. et al., *Can. J. Chem.*, 1989, **67**, 1077 (*isol, uv, ir, pmr, cmr, ms*)

Obscurolide C_{2z}

[154272-84-1]

C₁₅H₁₉NO₅ 293.319

Prod. by *Streptomyces viridochromogenes*. Phosphodiesterase inhibitor. Yellow solid. Sol. MeOH, EtOH.

7-Epimer: **Obscurolide C_{2β}**

[154334-89-1]


C₁₅H₁₉NO₅ 293.319

Prod. by *Streptomyces viridochromogenes*. Phosphodiesterase inhibitor. Yellow solid. Sol. MeOH, EtOH.

Ritzau, M. et al., *J. Antibiot.*, 1993, **46**, 1625–1628 (*isol, struct*)

Obtusine

O-11

C₂₄H₃₉NO₅ 421.576

3-O-[6-Deoxy-3-O-methyl-β-D-glucopyranosyl-(1→4)-2,6-dideoxy-3-O-methyl-β-D-ribo-hexopyranoside]: [263839-28-7]


C₃₈H₆₃NO₁₂ 725.915

Alkaloid from the roots of *Cryptolepis obtusa*. Powder.

Paulo, A. et al., *Phytochemistry*, 2000, **53**, 417–422

Obtusolactam

O-12

C₂₄H₃₇NO₄ 403.561

Parent not isol.

3-O-[6-Deoxy-3-O-methyl-β-D-glucopyr-

anosyl-(1→4)-2,6-dideoxy-3-O-methyl-β-D-ribo-hexopyranoside]: [263839-29-8]


C₃₈H₆₁NO₁₁ 707.9

Alkaloid from the roots of *Cryptolepis obtusa*. Powder.

Paulo, A. et al., *Phytochemistry*, 2000, **53**, 417–422

Obyanamide

O-13



Absolute Configuration

C₃₀H₄₁N₅O₆S 599.75


Depsipeptide antibiotic. Stereochem. revised in 2006. Isol. from *Lynghya confervoides*. Cytotoxic. Powder. [α]_D²⁷ +20 (c, 0.04 in MeOH). λ_{max} 210 (log ε 5.2); 222 (log ε 4.32) (MeOH).

Williams, P.G. et al., *J. Nat. Prod.*, 2002, **65**, 29–31 (*isol, pmr, cmr*)

Zhang, W. et al., *Tetrahedron*, 2006, **62**, 9966–9972 (*synth, struct*)

Oceanalin A

O-14

C₄₁H₇₂N₂O₉ 737.028


Isol. from the sponge *Oceanapia* sp. Antifungal agent. Amorph. solid. [α]_D -5.7 (c, 0.14 in EtOH). λ_{max} 238 (ε 7600); 288 (ε 7850) (MeOH).

Makariev, T.N. et al., *Org. Lett.*, 2005, **7**, 2897–2900 (*isol, pmr, cmr*)

Oceanapamine

O-15

[162857-98-9]

C₂₀H₃₃N₃ 315.501


Alkaloid from the Philippine sponge *Oceanapia* sp. Shows antimicrobial activity. Oil (as trifluoroacetate). [α]_D²² -6.4 (c, 3.1 in MeOH) (as trifluoroacetate). λ_{max} 202 (ε 14290); 220 (ε 6690) (MeOH) (Berdy).

Boyd, K.G. et al., *J. Nat. Prod.*, 1995, **58**, 302 (*isol, uv, ir, pmr, cmr, struct*)

Ochotensine

O-16

3',4',6,8-Tetrahydro-7'-methoxy-2'-methyl-6-methylenespiro[7H-indeno[4,5-d]-1,3-dioxole-7,1'(2'H)-isoquinolin]-6'-ol, 9CI. Alkaloid F17



(S)-form

C₂₁H₂₁NO₄ 351.401

(S)-form [4959-88-0]

Alkaloid from *Corydalis ochotensis*, *Corydalis ochotensis* var. *raddeana*, *Corydalis sibirica*, *Corydalis solida*, *Corydalis vaginans* and *Dicentra cucullaria* (Papaveraceae). Prisms (CHCl₃). Mp 252°. [α]_D²⁴ +51.7 (c, 0.2 in CHCl₃). [α]_D²³ +63.9 (c, 2 in 0.1M HCl).

LD₅₀ (mus, ivn) 10600 μg/kg. WH1312400

Methiodide:

Cryst. (MeOH). Mp 215°. Mp not well defined and depends on the rate of heating.

Me ether: Ochotensimine. Alkaloid F48

[4829-36-1]

C₂₂H₂₃NO₄ 365.428

Alkaloid from *Corydalis ochotensis*, *Corydalis ochotensis* var. *raddeana* and *Corydalis solida* (Papaveraceae). Syrup. [α]_D²⁰ +46.3 (c, 0.54 in MeOH).

Me ether; hydroiodide: Mp 190°.**Me ether; perchlorate:** Mp 172°.**Me ether; methiodide:**

Cryst. (Et₂O/MeOH). Mp 225° dec. [α]_D²⁵ +49.2 (c, 0.2 in MeOH).

O-De-Me, O³-Me: Isoochotensine

[112642-44-1]

C₂₁H₂₁NO₄ 351.401

Alkaloid from *Corydalis ochotensis* (Papaveraceae).

(±)-form [21008-78-6]

Synthetic. Prisms (Py/EtOH). Mp 241–243.5°.

Hydrochloride:

Cryst. (EtOH). Mp 205–212°.

Me ether: [19893-94-8]

Synthetic. Oil; cryst. (as perchlorate). Mp 235–238° (perchlorate).

Manske, R.H.F. et al., *Can. J. Res., Sect. B*, 1936, **14**, 354; 1938, **16**, 81; 1940, **18**, 75 (*isol*)

McLean, S. et al., *Can. J. Chem.*, 1966, **44**, 2449 (*ir, ms, struct*)

MacDonald, A.C. et al., *J.C.S.(B)*, 1966, 929 (*cryst struct*)

Irie, H. et al., *J.C.S.(C)*, 1968, 3051 (*synth, pmr*)

Kelly, R.B. et al., *Can. J. Chem.*, 1969, **47**, 2501 (*synth, uv, ir*)

Shamma, M. et al., *Chem. Comm.*, 1972, 310 (*abs config*)

Holland, H.L. et al., *Can. J. Chem.*, 1974, **52**, 2818; 1979, **57**, 1588–1597 (*biosynth*)


Lu, S.T. et al., *J.C.S. Perkin 1*, 1976, 63 (*isol*)

- Hughes, D.W. *et al.*, *Can. J. Chem.*, 1977, **55**, 3304 (*cmr*)
 Kametani, T. *et al.*, *J.C.S. Perkin I*, 1977, 390 (*isol*)
 Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1978, **56**, 383 (*isol*)
 Margvelashvili, N.N. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 592; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 509 (*occur*)
 Wu, T.S. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1987, **34**, 157; *CA*, **108**, 72118r (*Isoochotensine*)
 Meyers, A.I. *et al.*, *Tet. Lett.*, 1997, **38**, 4195 (*synth, Isoochotensine*)

Ochrindole A

O-17

[157414-83-0]

 $C_{29}H_{28}N_2O_3$ 452.552

Isol. from the sclerotia of *Aspergillus ochraceus*. Shows moderate anti-insectan activity. $[\alpha]_D^{25} +9.2$ (c, 0.13 in MeOH). λ_{max} 231 (ϵ 14000) (MeOH) (Berdy).

O¹-De-Me: Ochrindole C

[157414-85-2]

 $C_{28}H_{26}N_2O_3$ 438.525From *Aspergillus ochraceus*.**10-Hydroxy: Ochrindole B**

[157414-84-1]

 $C_{29}H_{28}N_2O_4$ 468.551


From *Aspergillus ochraceus*. Shows moderate anti-insectan activity. $[\alpha]_D^{25} +10$ (c, 0.1 in MeOH). λ_{max} 225 (ϵ 25000); 284 (ϵ 9400) (MeOH) (Berdy).

De Guzman, F.S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 634 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ochrindole D

O-18

[157414-86-3]

 $C_{27}H_{22}N_2O_3$ 422.482


Alkaloid from the sclerotia of *Aspergillus ochraceus*. Shows moderate anti-insectan activity. λ_{max} 226 (ϵ 27000); 272 (ϵ 2000) (MeOH) (Berdy).

De Guzman, F.S. *et al.*, *J. Nat. Prod.*, 1994, **57**, 634 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ochrobactin C

O-19

[906064-18-4]

 $C_{38}H_{64}N_4O_{13}$ 784.943

Isol. from *Ochrobactrum* sp. SP18 and *Vibrio* sp. DS40M5.

N-Deacyl, N-octanoyl: Ochrobactin B

[906064-17-3]

 $C_{36}H_{62}N_4O_{13}$ 758.905

Isol. from *Ochrobactrum* sp. SP18 and *Vibrio* sp. DS40M5.

N-Deacyl, N-(2E-octenoyl): Ochrobactin A

[906064-16-2]


 $C_{36}H_{60}N_4O_{13}$ 756.889

Isol. from *Ochrobactrum* sp. SP18. Siderophore.

Martin, J.D. *et al.*, *J. Biol. Inorg. Chem.*, 2006, **11**, 633-641 (*isol, uv, pmr, cmr, ms*)

Ochrobirine, 9CI

O-20



Absolute Configuration

 $C_{20}H_{19}NO_6$ 369.373

Diastereoisomer of Severzinine, S-279.

(+)-form [24181-64-4]

Alkaloid from *Corydalis lutea*, *Corydalis ochroleuca*, *Corydalis sibirica* and *Corydalis vaginans* (Papaveraceae). Cryst. (MeOH or EtOH). Mp 204° (198°). $[\alpha]_D^{21} +35.9$ (c, 0.4 in $CHCl_3$). Forms a methanolate, Mp 138-139°. λ_{max} 205 (log ϵ 4.8); 240 (log ϵ 3.94); 291 (log ϵ 3.91) (MeOH).

Di-Ac:

Cryst. (MeOH). Mp 179°.

N-De-Me, O⁸-Ac: Lederine

[76193-61-8]

 $C_{21}H_{19}NO_7$ 397.384

Alkaloid from *Corydalis ledebouriana* and *Dicentra peregrina* (Papaveraceae). Mp 208-209°. $[\alpha]_D^{25} +13$ (c, 0.84 in $CHCl_3$).

(±)-form [33805-02-6]

Synthetic. Cryst. (C_6H_6 /hexane or MeOH). Mp 185-187° Mp 210° Mp 235° (prob. polymorph.).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1939, **17**, 89-94; 95-98 (*isol*)


Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1969, **47**, 3589-3592 (*uv, ir, pmr, struct*)

Kametani, T. *et al.*, *Chem. Comm.*, 1971, 925-926 (*synth*)

- Nalliah, B.C. *et al.*, *Can. J. Chem.*, 1972, **50**, 1819-1824; 1979, **57**, 1545-1549 (*synth*)
 Shamma, M. *et al.*, *Chem. Comm.*, 1972, 310-311 (*uv, abs config*)
 Margvelashvili, N.N. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 848 (*isol*)
 Mathew, M. *et al.*, *Acta Cryst. B*, 1975, **31**, 2899-2901 (*cryst struct*)
 Hughes, D.W. *et al.*, *Can. J. Chem.*, 1977, **55**, 3304-3311 (*cmr*)
 Israilov, I.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **16**, 392-393 (*Lederine*)
 Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 3446-3449 (*synth, ir, pmr, ms*)

Ochrolifuanines

O-21



(3S,17R,20R)-form

 $C_{29}H_{34}N_4$ 438.614**(3S,17R,20R)-form****Ochrolifuanine A**

[35527-46-9]

Alkaloid from *Ochrosia lifuana*, *Ochrosia miana*, *Ochrosia confusa*, *Aspidosperma excelsum* and *Dyera costulata* (Apocynaceae). Also from *Strychnos potatorum*. Amorph. Sol. MeOH, $CHCl_3$, C_6H_6 ; poorly sol. H_2O , hexane. $[\alpha]_D^{20} +15$ (c, 1 in EtOH). λ_{max} 228 (log ϵ 4.73); 284 (log ϵ 4.2); 291 (log ϵ 4.13) (MeOH).

►RC1224000**18,19-Didehydro: 18-Dehydroochrolifuanine A** $C_{29}H_{32}N_4$ 436.599

Isol. from the leaves of *Dyera costulata* (Apocynaceae). $[\alpha]_D^{25} +4$ (c, 0.7 in EtOH). λ_{max} 226; 283; 290 (EtOH).

(3S,17S,20R)-form**Ochrolifuanine B**

[35471-11-5]

Alkaloid from the leaves of *Ochrosia lifuana* and from *Ochrosia miana* (Apocynaceae). Amorph. $[\alpha]_D^{20} +0.1$ (c, 1 in EtOH).

►RC1225000**N²-Me: Nigritanine. 18,19-Dihydrousambarine**

[67657-99-2]

 $C_{30}H_{36}N_4$ 452.641

Alkaloid from *Strychnos usambarensis*, *Strychnos nigritana*, *Strychnos potatorum* and *Strychnos barteri* (Loganiaceae). Shows cytotoxic and antimitotic activity. Cryst. (EtOAc/hexane). Mp 202-204°. Not correlated with Usambarine. λ_{max} 226 (log ϵ 4.76); 282 (log ϵ 4.16); 290 (log ϵ 4.09) (MeOH).

18,19-Didehydro, N²-Me: Usambarine.**18,19-Dehydronigritanine**

[35226-29-0]

 $C_{30}H_{34}N_4$ 450.625

Alkaloid from leaves of *Strychnos usambarensis*, also from *Strychnos nigritana*, *Strychnos barteri* and *Rauwolfia obscura* (Loganiaceae).

Apocynaceae). Microprisms (Et₂O) or cryst. (EtOAc/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 215-218° (block) Mp 226-228°. λ_{max} 226 (log ε 4.77); 282 (log ε 4.16); 290 (log ε 4.09) (MeOH).

10-Hydroxy, N²-Me: 18,19-Dihydro-usambaridine Vi. 10-Hydroxy-18,19-dihydrousambarine. 10-Hydroxynigritanine [67658-02-0]
C₃₀H₃₆N₄O 468.641
Alkaloid from leaves of *Strychnos barteri* and *Strychnos nigrifolia* (Loganiaceae). Cryst. (EtOAc). Mp 181-183°. OH group originally placed at the 12-posn. The Vi suffix refers to a violet colour reaction of the alkaloid. λ_{max} 226 (log ε 4.75); 282 (log ε 4.17); 290 (log ε 4.1); 310 (log ε 3.76) (MeOH).

10-Hydroxy, 18,19-didehydro, N²-Me: 10-Hydroxyusambarine. Usambaridine Vc. Usambaridine Vi. 10-Hydroxy-18,19-dehydronigritanine. 18-Dehydro-10-hydroxynigritanine [42814-43-7]
C₃₀H₃₄N₄O 466.625
Alkaloid from leaves of *Strychnos usambarensis*, also from *Strychnos barteri*, *Strychnos nigrifolia* (Loganiaceae). Antimitotic agent. Cryst. (EtOAc/hexane). Mp 174-176°. OH group originally placed at the 12-posn. The Vc and Vi suffixes refer to colour reactions. λ_{max} 226 (log ε 4.74); 281 (log ε 4.2); 290 (log ε 4.12); 310 (log ε 3.76) (MeOH).

10-Hydroxy, 18,19-didehydro, N²,N⁴-di-Me: N⁴-Methyl-10-hydroxyusambarine [88607-20-9]
C₃₁H₃₇N₄O⁺ 481.659
Alkaloid from leaves and stem bark of *Strychnos usambarensis*.

10-Methoxy, 18,19-didehydro: Obscurifoline. Alkaloid B1 [55538-31-3]
C₃₀H₃₄N₄O 466.625
Alkaloid from leaves of *Rauwolfia obscura* (Apocynaceae). Small cryst. Mp 160-162°. λ_{max} 226 (log ε 4.72); 280 (log ε 4.23); 289 (sh) (log ε 4.15); 307 (sh) (MeOH).

11-Hydroxy, N²-Me: 18,19-Dihydro-usambaridine Br. 11-Hydroxy-18,19-dihydrousambarine [67658-01-9]
C₃₀H₃₆N₄O 468.641
Alkaloid from leaves of *Strychnos usambarensis* (Loganiaceae). The Br subscript refers to a colour reaction of the alkaloid.

11-Hydroxy, 18,19-didehydro, N²-Me: Usambaridine Br. 11-Hydroxyusambarine [67658-00-8]
C₃₀H₃₄N₄O 466.625
Alkaloid from leaves of *Strychnos usambarensis* (Loganiaceae). The Br label refers to a colour reaction of the alkaloid.

11-Hydroxy, 18,19-didehydro, N²,N⁴-di-

Me: N⁴-Methyl-11-hydroxyusambarine [88607-19-6]
C₃₁H₃₇N₄O⁺ 481.659
Alkaloid from leaves of *Strychnos usambarensis*.

(3S,17S,20S)-form
Ochrolifuanine C
51820-25-8]
Synthetic. [α]_D -62 (c, 1 in EtOH). λ_{max} 227 ; 283 ; 290 (EtOH).

(3S,17R,20S)-form
Ochrolifuanine D
51820-26-9]
Synthetic. [α]_D -23 (c, 1 in EtOH). λ_{max} 226 ; 282 ; 289 (EtOH).

(3R,17ξ,20ξ)(1)-form
Ochrolifuanine E
[87479-88-7]
Alkaloid from the leaves of *Dyera costulata* (Apocynaceae). Also from *Strychnos potatorum*. Active against gram-positive bacteria. [α]_D +50 (c, 0.86 in EtOH). λ_{max} 227 ; 283 ; 290 (EtOH) (Berdy).

18,19-Didehydro: 18-Dehydroochrolifuanine E
[87419-58-7]
C₂₉H₃₂N₄ 436.599
Isol. from the leaves of *Dyera costulata* (Apocynaceae). [α]_D +80 (c, 0.64 in EtOH).

(3R,17ξ,20ξ)(2)-form
Ochrolifuanine F
[87479-89-8]
Alkaloid from the leaves of *Dyera costulata* (Apocynaceae). Active against gram-positive and -negative bacteria. [α]_D -37 (c, 0.92 in EtOH). λ_{max} 227 ; 283 ; 290 (EtOH) (Berdy).

18,19-Didehydro: 18-Dehydroochrolifuanine F
[87419-57-6]
C₂₉H₃₂N₄ 436.599
Isol. from the leaves of *Dyera costulata* (Apocynaceae). Active against gram-positive and gram-negative bacteria. [α]_D -57 (c, 1.06 in EtOH). λ_{max} 227 ; 283 ; 290 (EtOH) (Berdy).

Koch, M. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 753-754 (*Usambarine, isol, uv, ir, pmr, struct*)

Peube-Locou, N. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 905-906 (*Ochrolifuanines A,B, uv, ir, pmr, ms*)

Bruneton, J. et al., *Phytochemistry*, 1972, **11**, 2618 (*isol*)

Koch, M. et al., *Ann. Pharm. Fr.*, 1973, **31**, 45-48 (*Usambarine, isol, uv, ir, pmr, struct*)

Koch, M. et al., *Bull. Soc. Chim. Fr.*, 1973, 2868-2869 (*cd, struct*)

Préaux, N. et al., *Plant. Med. Phytother.*, 1974, **8**, 250-254; *CA*, **83**, 75339q (*isol*)

Koch, M. et al., *J.O.C.*, 1975, **40**, 2836-2838 (*cmr*)

Timmins, P. et al., *Planta Med.*, 1975, **27**, 105-111 (*Obscurifoline*)

Oguakwa, J.U. et al., *Gazz. Chim. Ital.*, 1978, **108**, 615-618 (*Strychnos nigrifolia constits*)

Coune, C. et al., *J. Pharm. Belg.*, 1978, **33**, 11-23; 284-286; 1982, **37**, 189-194 (*isol, cmr, struct, Strychnos usambarensis constits*)

Richard, C. et al., *Phytochemistry*, 1978, **17**, 539-541 (*Ochrolifuanines C,D, synth, uv, pmr, ms*)

Angenot, L. et al., *Phytochemistry*, 1978, **17**, 1687-1689 (*Usambarine, isol, uv, ir, pmr, ms, struct*)

Seguin, E. et al., *Planta Med.*, 1979, **37**, 175-177 (*Usambarine, synth, uv, ir, pmr, ms, struct*)

Nicoletti, M. et al., *Fitoterapia*, 1980, **51**, 131-134 (*Usambarine, Nigritanine, cmr*)

Caprasse, M. et al., *J. Pharm. Belg.*, 1983, **38**, 211-218; *CA*, **100**, 64988u (*N⁴-Methyl-10-hydroxyusambarine, N⁴-Methyl-11-hydroxyusambarine*)

Mirand, C. et al., *Phytochemistry*, 1983, **22**, 577-579 (*Dyera costulata constits*)


Caron, C. et al., *Planta Med.*, 1988, **54**, 409-412 (*activity*)

Quetin-Leclercq, J. et al., *Planta Med.*, 1991, **57**, 501 (*N⁴-Methyl-10-hydroxyusambarine*)

Massiot, G. et al., *Phytochemistry*, 1992, **31**, 2873-2876 (*isol, Strychnos constits*)

Ochromianoxine O-22

[55322-95-7]




C₂₀H₂₈N₂O₃ 344.453
Alkaloid from the bark of *Ochrosia miana* (Apocynaceae). [α]_D²⁰ +29 (c, 1 in EtOH). CAS assigns the 20-config. as 20*R*-, although it does not seem to have been determined. λ_{max} 226 (log ε 4.22); 258 (log ε 3.65); 285 (log ε 3.58); 294 (log ε 3.47) (EtOH).

Preaux, N. et al., *Phytochemistry*, 1974, **13**, 2607-2609 (*uv, ir, ms, cd, struct*)

Ochroprosinine oxindole O-23

[77794-97-9]



C₂₁H₃₀N₂O₄ 374.479
Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{max} 217 ; 274 ; 304 (sh) (EtOH).

Ahond, A. et al., *J. Nat. Prod.*, 1981, **44**, 193-199 (*isol, uv, ir, pmr, ms, cd, struct*)

Ocodemerine O-24


C₃₇H₄₀N₂O₆ 608.733

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the bark of *Nectandra rodiaei* (*Ocotea rodiaei*) (Lauraceae). Cryst. + 1.5H₂O (as dihydrochloride). Mp 275° (dihydrochloride). [α]_D -170 (c, 1.0 in H₂O) (dihydrochloride).

Hearst, P.J. et al., *J.O.C.*, 1964, **29**, 466 (*isol*)

Ookryptine**O-25**

11-Hydroxy-1,10-dimethoxy-2,3-methylenedioxyaporphine
[19893-96-0]



$C_{20}H_{21}NO_5$ 355.39


Alkaloid from an unidentified *Ocotea* sp. (Lauraceae). Cryst. (Et₂O). Mp 160-161°. $[\alpha]_D^{27} +164$ (c, 0.4 in CHCl₃).

Me ether: 1,10,11-Trimethoxy-2,3-methylenedioxyaporphine
Mp 170-171°. $[\alpha]_D^{27} +156$ (c, 1.1 in CHCl₃).

Cava, M.P. et al., *Tet. Lett.*, 1968, 2437 (uv, pmr, struct)

Ocominarone**O-26**

4,9,10,11-Tetramethoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI
[72170-13-9]




$C_{21}H_{17}NO_7$ 395.368

Alkaloid from the leaves of *Ocotea minarum* (Lauraceae). Mp 268-270°.

Vecchietti, V. et al., *Farmaco, Ed. Sci.*, 1979, 34, 829 (isol, uv, pmr, ms, struct)

Ocoteine**O-27**

3,9,10-Trimethoxy-1,2-methylenedioxyaporphine. *Thalicmine*

*(S)*-form

$C_{21}H_{23}NO_5$ 369.416

(S)-form [3246-21-7]

Alkaloid from *Ocotea puberula* and *Thalictrum minus* (Lauraceae, Ranunculaceae). Antitussive, hypotensive, spasmolytic and adrenergic agent in exptl. animals. Also shows antimicrobial and antiinflammatory activity. Mp 140-142°.

$[\alpha]_D +33.3$ (c, 1.02 in CHCl₃).

Hydroiodide: Mp 198-200°. $[\alpha]_D^{28} +22$ (c, 0.215 in 50% EtOH).

N-De-Me: 3,9,10-Trimethoxy-1,2-methylenedioxyaporphine. **O-Methylcassifiline**. *Northalicmine*
[3984-10-9]

$C_{20}H_{21}NO_5$ 355.39

Alkaloid from *Cassytha filiformis* and *Thalictrum strictum* (Lauraceae, Ranunculaceae). Cryst. (Et₂O). Mp 150-152°. $[\alpha]_D^{15} +16.4$ (c, 1 in CHCl₃).

O⁹-De-Me: 9-Hydroxy-3,10-dimethoxy-1,2-methylenedioxyaporphine
Mp 210-211°. $[\alpha]_D^{21} +24.6$ (c, 0.63 in CHCl₃).

O⁹-De-Me, N-de-Me: 9-Hydroxy-3,10-dimethoxy-1,2-methylenedioxyaporphine. **Cassythine**
[4030-51-7]

$C_{19}H_{19}NO_5$ 341.363

Alkaloid from *Cassytha filiformis* (*Cassytha americana*) (Lauraceae). Cryst. (CHCl₃/EtOH). Mp 217-219° dec. $[\alpha]_D +24$ (c, 0.56 in CHCl₃).

O⁹-De-Me, N-de-Me, N-(methoxycarbonyl): **Cathaformine**. *N-Methoxycarbonylcassifiline*

$C_{21}H_{21}NO_7$ 399.399

Alkaloid from *Cassytha filiformis* (Lauraceae). Amorph. brown powder. Mp 122-124°. $[\alpha]_D +116.4$ (c, 0.06 in CHCl₃). λ_{max} 237 ; 286 ; 307 (EtOH).

6a,7-Didehydro: **Dehydroocoteine**. *Dehydrothalicmine*. *Didehydroocoteine*[†]
[38366-01-7]

$C_{21}H_{21}NO_5$ 367.401

Alkaloid from the bark of *Ocotea puberula* and the roots of *Thalictrum isopyroides*. Also present in the bark of *Nectandra saligna* (Lauraceae, Ranunculaceae). Shows antiplasmodial and amoebicidal activities. Mp 203-204° (190-191°).

4,5,6a,7-Tetrahydro: 4,10,11-Trimethoxy-7-methyl-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline.

Didehydroocoteine[†]. *Tetrahydroocoteine*
[38366-02-8]

$C_{21}H_{19}NO_5$ 365.385

Alkaloid from *Ocotea puberula* (Lauraceae).

(±)-form

Hydroiodide: Mp 225°.

Iacobucci, G.A. et al., *Cienc. Invest.*, 1951, 7, 48 (*Ocoteine, isol*)

Vernengo, M.J. et al., *Experientia*, 1963, 19, 294 (*Ocoteine, struct*)

Govindachari, T.R. et al., *Tetrahedron*, 1964, 20, 2895 (*Ocoteine, synth, uv, ir, pmr*)

Tomita, M. et al., *Yakugaku Zasshi*, 1965, 85, 827 (*Cassythine*)

Johns, S.R. et al., *Aust. J. Chem.*, 1966, 19, 297 (*Cassythine, isol, uv, pmr, struct*)

Cava, M.P. et al., *J.O.C.*, 1968, 33, 2443 (*O-Methylcassifiline*)

Ismailov, Z.F. et al., *Khim. Prir. Soedin.*, 1968, 4, 196; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, 4, 169 (*Ocoteine, ms*)

Maekh, S.Kh. et al., *Khim. Prir. Soedin.*, 1971, 7, 381; 1976, 12, 560; *Chem. Nat.*

Compd. (Engl. Transl.), 1971, 7, 363; 1976, 12, 507 (*Dehydroocoteine, O-Methylcassifiline*)

Baralle, F. et al., *Experientia*, 1972, 28, 875 (*Dehydroocoteine, Didehydroocoteine*)

Gorovoi, P.G. et al., *Khim. Prir. Soedin.*, 1975, 11, 533; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, 11, 568 (*O-Methylcassifiline*)

Ringdahl, B. et al., *J. Nat. Prod.*, 1981, 44, 80 (*cd*)

Dickman, D.A. et al., *Tet. Lett.*, 1986, 27, 1465 (*synth*)

Wu, Y.C. et al., *Phytochemistry*, 1997, 46, 181 (*Cathaformine*)


Nimgirawath, S. et al., *Aust. J. Chem.*, 2000, 53, 527-529 (*Cathaformine, synth*)

Wright, C.W. et al., *J. Nat. Prod.*, 2000, 63, 1038-1040 (*Dehydroocoteine, activity*)

Stévigny, C. et al., *Planta Med.*, 2002, 68, 1042-1044 (*isol, pmr, cmr*)

Ocotosine**O-28**

[18529-51-6]



$C_{37}H_{38}N_2O_6$ 606.717

(S)-form

Alkaloid from the seeds of *Ocotea rodiaei* (*Nectandra rodiaei*) (Lauraceae). Plates (Me₂CO). $[\alpha]_D^{24} +291$ (c, 1.0 in CHCl₃). The probable (S)-config. is assigned here on the basis of its (+)ve opt. rotn. (see Saez et al).

O¹²-De-Me, O^{12'}-Me: **Guattamine**

[116064-69-8]

$C_{37}H_{38}N_2O_6$ 606.717

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). $[\alpha]_D +142$ (c, 0.67 in CHCl₃). Exists in CDCl₃ soln. as a 70:30 mixt. of two conformers.

α' -Ketone, O¹²-de-Me, O^{12'}-Me: **Guattaminone**

[116084-91-4]

$C_{37}H_{36}N_2O_7$ 620.701

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). $[\alpha]_D +78$ (c, 0.26 in CHCl₃).

(R)-form

O⁶-De-Me: **Cordobimine**

[121892-86-2]

$C_{36}H_{36}N_2O_6$ 592.69

Alkaloid from the stem bark of an unidentified *Crematosperma* sp. (Annonaceae). Amorph. $[\alpha]_D^{20} -185$ (c, 0.5 in CHCl₃).

Chan, K.C. et al., *J.C.S.(C)*, 1967, 2479 (*Ocotosine*)

Berthou, S. et al., *Tetrahedron*, 1988, 44, 2193 (*Guattamine, Guattaminone*)

Saez, J. et al., *Can. J. Chem.*, 1989, 67, 275 (*Cordobimine*)

2,4-Octadecadienoic acid**O-29**

[76282-17-2]

$H_3C(CH_2)_{12}CH=CHCH=CHCOOH$

$C_{18}H_{32}O_2$ 280.45

(2E,4E)-form [59404-49-8]*Me ester*: [86120-41-4]C₁₉H₃₄O₂ 294.476

Oil.

2-Methylpropylamide: N-Isobutyl-2,4-octadecadienamide


[54794-70-6]

C₂₂H₄₁NO 335.572Constit. of *Piper guineense* fruits and whole plants of *Piper argyrophyllum*. Mp 78–80° Mp 92–94°.**(2E,4Z)-form****2-Methylpropylamide: Pipericine**C₂₂H₄₁NO 335.572Constit. of the fruit of *Piper nigrum* (pepper). Oil. λ_{max} 257 (EtOH).**(2Ě,4Ě)-form****Pyrrolidide: 2,4-Octadecadienoic acid****pyrrolidide**


[102934-33-8]

C₂₂H₃₉NO 333.556Constit. of leaves and fruit of *Piper guineense*.**Piperidide: 2,4-Octadecadienoic acid piperidide**

[151391-73-0]

C₂₃H₄₁NO 347.583Isol. from the leaves and fruit of *Piper guineense*.Okogun, J.I. *et al.*, *J.C.S. Perkin 1*, 1974, 2195 (*isol, ms*)Vig, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 1358 (*synth, deriv*)Akimoto, A. *et al.*, *Angew. Chem., Int. Ed.*, 1981, **20**, 90 (*synth*)Cardillo, G. *et al.*, *Tetrahedron*, 1986, **42**, 917 (*deriv, synth, ir, pmr, cmr, ms*)Banerji, A. *et al.*, *Indian J. Chem., Sect. B*, 1988, **27**, 163 (*isol, uv, ir, pmr*)Strunz, G.M. *et al.*, *Can. J. Chem.*, 1996, **74**, 419 (*synth, isobutylamide*)Siddiqui, B.S. *et al.*, *Phytochemistry*, 1997, **45**, 1617 (*Pipericine*)Adesina, S.K. *et al.*, *Pharmazie*, 2003, **58**, 423–425 (*pyrrolidide, piperidide*)**2,12-Octadecadienoic acid O-30**C₁₈H₃₂O₂ 280.45**(2E,12Z)-form****4-Methylpentylamide: N-(4-Methylpentyl)-2,12-octadecadienamide. Pipzorine**

[622405-49-6]

C₂₄H₄₅NO 363.626Alkaloid from the dried seeds of *Piper nigrum*. Amorph. powder. λ_{max} 209 (ε 6160) (MeOH).Siddiqui, B.S. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2760–2767 (*Pipzorine*)**2,4,8,10,12,16-Octadecahexaenoic acid O-31**C₁₈H₂₄O₂ 272.386**2-Methylpropylamide: 2,4,8,10,12-Octadecapentaenoic isobutylamide. Heliopsin**C₂₂H₃₃NO 327.509Isol. from seeds of *Heliopsis helianthoides* var. *scabra*. Light-yellow viscous oil. Bp_{0.08} 198–200° part dec. Struct. not certain.Jacobson, M. *et al.*, *J.A.C.S.*, 1957, **79**, 356**Octadecanoic acid, 9CI O-32***Stearic acid. Bassinic acid. Lactic acid. Stearophanic acid. Talcic acid. FEMA 3035*

[57-11-4]


H₃C(CH₂)₁₆COOHC₁₈H₃₆O₂ 284.481Manuf. by hydrolysis of fats and oils. Constit. of most vegetable and animal fats, including the lipids of *Physalia physalis* (Portuguese-man-of-war). Acid and derivatives widely used as additives to industrial preparations. Used as EtOH soln. for nephelometric detn. of Ca. Used (with heptadecanoic acid) for amino acid sequencing in peptides. Used for surfactants, soaps and plasticisers. Pharmaceutical excipient (tablet lubricant; Ca and Mg salts are used similarly). Leaflets. Sol. EtOH, C₆H₆, CHCl₃; sl. sol. H₂O. Mp 69.7°. Bp 386° Bp₅ 213°. Polymorphic, at least 4 cryst. forms are known.

► Skin irritant. Fl. p. 196°, autoignition temp. 395°. WI2800000

*Amide: Octadecanamide. Stearamide.**Amide C-18. Hidorin M7. Kemamide S* [124-26-5]C₁₈H₃₇NO 283.496Alkaloid from *Zostera marina* (Zosteraceae) and *Rhizoclonium hieroglyphicum*. Cryst. (EtOH). Mp 109°. Bp₁₂ 250–251°.

► RG0182000

[4568-28-9, 593-29-3]

Kawasaki, W. *et al.*, *Phytochemistry*, 1998, **47**, 27–29 (*amide*)**2,4,12-Octadecatrienoic acid O-33**C₁₈H₃₀O₂ 278.434**(2E,4E,12Z)-form****2-Methylpropylamide: 2,4,12-Octadecatrienoic acid isobutylamide. N-Isobutyl-2,4,12-octadecatrienamide**


[151391-69-4]

C₂₂H₃₉NO 333.556Alkaloid from *Piper retrofractum* (Javanese long pepper). Oil. λ_{max} 263 (log ε 4.04) (MeOH).**4-Methylpentylamide: N-(4-Methylpentyl)-2,4,12-octadecatrienamide. Pipnoohine**C₂₄H₄₃NO 361.61Alkaloid from the fruit of *Piper nigrum* (pepper). Amorph. powder. λ_{max} 261 (ε 24500) (MeOH).**Piperidide: 2,4,12-Octadecatrienoic acid piperidide. 1-(2,4,12-Octadecatrienoyl)piperidine**


[151391-71-8]

C₂₃H₃₉NO 345.567Alkaloid from *Piper retrofractum* (Javanese long pepper).Kikuzaki, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 1329–1333 (*isol, amides*)
Siddiqui, B.S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1349–1352 (*Pipnoohine*)**2,4,14-Octadecatrienoic acid O-34**

[25448-03-7]

C₁₈H₃₀O₂ 278.434**(2E,4E,14Z)-form****Piperidide: Piperoctadecalidine**

[144525-15-5]

C₂₃H₃₉NO 345.567Alkaloid from fruits of *Piper retrofractum* (Piperaceae). Oil.Ahn, J.W. *et al.*, *Phytochemistry*, 1992, **31**, 3609**9-Octadecene-1,17-diamine, 9CI O-35****1,17-Diamino-9-octadecene**

Absolute Configuration

C₁₈H₃₈N₂ 282.512**(9Z,17R)-form****Harmonine**Defence alkaloid from *Harmonia lewis conformis* and other ladybug spp. Yellow oil. [α]_D -3.9 (c, 1 in C₆H₆) (>97%ee).Braconnier, M.F. *et al.*, *Experientia*, 1985, **41**, 519–521 (*isol*)Enders, D. *et al.*, *Annalen*, 1991, 569–574 (*synth*)**2-Octadecenoic acid O-36**

[5340-63-6]

C₁₈H₃₄O₂ 282.465**(E)-form** [2825-79-8]

Cryst. (EtOH or petrol). Mp 58.5°.

Me ester: [14663-11-7]C₁₉H₃₆O₂ 296.492

Mp 36°.

Et ester: [97975-07-0]C₂₀H₃₈O₂ 310.519

Cryst. (MeOH). Mp 25–26°. Bp 360°.

Amide:C₁₈H₃₅NO 281.481

Cryst. (EtOH). Mp 107–108°.

Piperidide: 1-(1-Oxo-2-octadecenyl)piperidine, 9CI. N-(2-Octadecenoyl)piperidine. Piperitine

[74267-81-5]

C₂₃H₄₃NO 349.599Alkaloid from the dried seeds of black pepper (*Piper nigrum*). Amorph. powder. λ_{max} 210 (ε 6450) (MeOH).**(Z)-form** [2825-66-3]


Cryst. (petrol). Mp 50.5°.

Barve, J.A. *et al.*, *Chem. Phys. Lipids*, 1971, **7**, 311–323 (*E-form, Z-form, synth*)

- Davies, J.E.D. *et al.*, *J.C.S. Perkin 2*, 1972, 1557-1561 (*ir*, *Raman*)
 Howton, D.R. *et al.*, *Org. Prep. Proced. Int.*, 1974, **6**, 175-178 (*E-form*, *synth*)
 Gunstone, F.D. *et al.*, *Chem. Phys. Lipids*, 1977, **18**, 115-129 (*cmr*)
 Plate, M. *et al.*, *Synthesis*, 1998, 1255-1258 (*Et ester*)
 Siddiqui, B.S. *et al.*, *Heterocycles*, 2002, **57**, 1653-1658 (*Piperitine*)

9-Octadecenoic acid, 9CI O-37

[2027-47-6]
 [26764-26-1]



$\text{C}_{18}\text{H}_{34}\text{O}_2$ 282.465

Present in lipids of *Physalia physalis* (Portuguese-man-of-war).

(E)-form

Elaidic acid

[112-79-8]

[27251-59-8]

Minor constit. of plant oils. Cryst. (EtOH). Mp 45-45.5°. Bp₁₅ 234° Bp_{0.8} 180-185°. Dimorphic.

► LD₅₀ (mus, ivn) 100 mg/kg. JX6125000

Amide: Elaidamide

[4303-70-2]

$\text{C}_{18}\text{H}_{35}\text{NO}$ 281.481

Cryst. (C_6H_6). Mp 93-94°.

(Z)-form

Oleic acid. Rapinic acid. Pamolyn 100. MYX 6121. FEMA 2815

[112-80-1]

[27104-13-8]

Major constit. of plant oils e.g. olive oil (ca. 80%), almond oil (ca. 80%) and many others, mainly as glyceride.

Constit. of tall oil and present in fruits.

Food additive. Used in manuf. of surfactants, soaps, plasticisers.

Emulsifying agent in foods and pharmaceuticals. Skin penetrant.

Herbicide, insecticide, fungicide. Oil.

Insol. H_2O ; sol. EtOH, Et₂O. d_4^{20} 0.9.

Mp 12° (labile form) Mp 16° (stable form)

Bp₅ 203-205°. n_D^{20} 1.4595. Log P 7.75 (uncertain value) (calc).

¹²⁵I and ¹³¹I labelled cpds. are used as radioactive agents (Oleotope). Undergoes BF₃-catalysed dimerisation.

► Skin irritant. LD₅₀ (rat, orl) 74000 mg/kg. RG2275000

Amide: Oleamide. Kemamide O

[301-02-0]

$\text{C}_{18}\text{H}_{35}\text{NO}$ 281.481

Isol. from the cerebrospinal fluid of humans, cats and rats. Isol. from lichen *Stereocaulon alpinum* and from *Desmos cochinchinensis*. Sleep inducer.

Shows activity against cyclooxygenase.

Cannabinoid CB₁ receptor agonist.

TRPV1 vanilloid receptor activator.

Vasodilator. Scales (EtOH). Sol.

MeOH. Mp 75-76°.

[2717-15-9, 1555-53-9, 557-07-3, 142-17-6, 2462-84-2, 10402-16-1, 544-60-5, 688-37-9]

Lerner, R. *et al.*, *Science (Washington, D.C.)*, 1995, **268**, 1506 (*isol*, *amide*)

Ingolfssdottir, K. *et al.*, *Phytomedicine*, 1997, **4**, 331-334; *CA*, **128**, 158800g (*Oleamide*, *activity*)

2,4-Octadien-1-amine O-38

1-Amino-2,4-octadiene



$\text{C}_8\text{H}_{15}\text{N}$ 125.213

(E,E)-form

N-(2*R*-Hydroxyheptyl): *Pseudodistamine* [356522-64-0]

$\text{C}_{15}\text{H}_{29}\text{NO}$ 239.4

Isol. from a *Pseudodistoma* sp. Gum.

[α]_D²⁰ -3 (c, 0.1 in MeOH). λ_{max} 225 (log ϵ 3.72) (MeOH).

Rashid, M.A. *et al.*, *Tetrahedron*, 2001, **57**, 5751-5755 (*Pseudodistamine*, *isol*, *pmr*, *cmr*, *uv*)

2,4-Octadienoic acid, 9CI O-39

[90435-12-4]



$\text{C}_8\text{H}_{12}\text{O}_2$ 140.182

(2E,4E)-form [22329-75-5]

Leaflets (petrol). Mp 76°.

Me ester: [74418-28-3]

$\text{C}_9\text{H}_{14}\text{O}_2$ 154.208

Bp₁₅ 101°.

Amide: 2,4-Octadienamido

[205311-23-5]

$\text{C}_8\text{H}_{13}\text{NO}$ 139.197

Alkaloid from *Phyllanthus fraternus*.

Cryst. Mp 136-137°. λ_{max} 276 (log ϵ 4.44) (MeOH). λ_{max} 276 (ϵ 27540)

(MeOH) (Berdy).

2-Methylpropylamide: N-Isobutyl-2,4-octadienamido

N-(2-Methylpropyl)-2,4-octadienamido. 2,4-Octadienoic acid isobutylamide

[23512-47-2]

$\text{C}_{12}\text{H}_{21}\text{NO}$ 195.304

Alkaloid from the wood of *Piper novae-hollandiae* and from the leaves and stems of *Piper banksii* (Piperaceae). Cryst. (petrol). Mp 94° (88-90°).

2-Methylpropylamide: N-Isobutyl-2,4-octadienamido

N-(2-Methylpropyl)-2,4-octadienamido. 2,4-Octadienoic acid isobutylamide

[23512-47-2]

$\text{C}_{12}\text{H}_{21}\text{NO}$ 195.304

Alkaloid from the wood of *Piper novae-hollandiae* and from the leaves and stems of *Piper banksii* (Piperaceae). Cryst. (petrol). Mp 94° (88-90°).

(2E,4Z)-form [68676-76-6] Has been synthesised.

Et ester: [39924-38-4]

$\text{C}_{10}\text{H}_{16}\text{O}_2$ 168.235

Liq. Bp_{0.05} 53°.

2-Methylpropylamide: [99615-85-7]

[100452-77-5]

Constit. of *Acmella ciliata*.

(2Z,4E)-form

2-Phenylethylamide: N-(2-Phenethyl)-2,4-octadienamido. 2,4-Octadienoic acid 2-phenylethylamide

[99615-86-8]

$\text{C}_{16}\text{H}_{21}\text{NO}$ 243.348

Constit. of *Acmella ciliata*.

[108965-85-1, 39924-38-4, 60388-61-6, 64713-73-1, 88738-80-1, 108965-83-9, 39924-43-1, 81798-18-7]

Jacobson, M. *et al.*, *J.A.C.S.*, 1956, **78**, 5084 (*synth*, *uv*)

Loder, J.W. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1531 (*isobutylamide*)

Loder, J.W. *et al.*, *Phytochemistry*, 1972, **11**, 2645 (*isobutylamide*)

Baumann, M. *et al.*, *Synthesis*, 1977, 681 (*synth*)

Sakai, T. *et al.*, *J.O.C.*, 1982, **47**, 1101 (*synth*, *ir*, *pmr*, *cmr*)

Tsuboi, S. *et al.*, *J.O.C.*, 1982, **47**, 4478 (*synth*, *ir*, *pmr*, *cmr*)

Martin, R. *et al.*, *Phytochemistry*, 1985, **24**, 2295-2300 (*phenylethylamides*)

Concepcion, A.B. *et al.*, *Tetrahedron*, 1995, **51**, 4011 (*synth*, *pmr*)

Bellina, F. *et al.*, *Synth. Commun.*, 1996, **26**, 3297 (*Me ester*)


Sittie, A.A. *et al.*, *Planta Med.*, 1998, **64**, 192-193 (*isol*, *amide*)

1',3,4,4',5,5',6,6'-Octahydro-2,3'-bipyridine, 9CI, 8CI O-40

Hystrine

[18017-50-0]

[18017-52-2]



$\text{C}_{10}\text{H}_{16}\text{N}_2$ 164.25

Alkaloid from *Lupinus formosus*, *Genista hystrix* and *Genista hystrix* spp. *legionensis* (Fabaceae). Prisms (MeOH/Me₂CO) (as hydrochloride). Mp 209° (hydrochloride).

N-Ac: N-Acetylhystrine

$\text{C}_{12}\text{H}_{18}\text{N}_2\text{O}$ 206.287

Alkaloid in *Lupinus formosus* (Fabaceae). Oil, rapidly yellowing in air.

Steinegger, E. *et al.*, *Pharm. Acta Helv.*, 1967, **42**, 177 (*isol*)

Steinegger, E. *et al.*, *Helv. Chim. Acta*, 1968, **51**, 206 (*synth*, *ir*, *uv*)

Steinegger, E. *et al.*, *Pharm. Acta Helv.*, 1970, **45**, 157; *CA*, **72**, 97301w (*biosynth*)

Fitch, W.L. *et al.*, *J.A.C.S.*, 1974, **96**, 4917 (*ms*)

Fitch, W.L. *et al.*, *J.O.C.*, 1974, **39**, 2974 (*N-Acetylhystrine*)

Octahydro-3-butyl-5-propyl-1H-indolizine O-41

1H-indolizine

3-Butyl-5-propylindolizidine. Indolizidine 223AB. Dendrobates Alkaloid 223AB. Gephyrotoxin 223AB

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]


[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

[67217-85-0]

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401



$\text{C}_{15}\text{H}_{29}\text{N}$ 223.401

(-)-form [67217-85-0]

Minor alkaloid from skin extracts of the neotropical poison-frog *Dendrobates histrionicus* (various populations) (Dendrobates)

batidae). [α]_D²⁰ -35 (c, 0.49 in MeOH).

[α]_D²⁰ -101 (c, 2.3 in hexane).

4'-Hydroxy: Indolizidine 239CD. Dendrobates Alkaloid 239CD. Gephyrotoxin 239CD

[67217-87-2]

$\text{C}_{15}\text{H}_{29}\text{NO}$ 239.4

Major alkaloid in *Dendrobates histrionicus* and *Dendrobates occulator*. $[\alpha]_D^{25}$ -52 (c, 0.19 in MeOH).

3'-Hydroxy: **Indolizidine 239AB**. *Dendrobates Alkaloid 239AB*. *Gephyrotoxin 239AB*

[67217-86-1]
[142925-69-7]

C₁₅H₂₉NO 239.4

Major to minor constit. of various populations of *Dendrobium histrionicus*; major constit. of *Dendrobium occulator*. $[\alpha]_D^{16}$ -38 (c, 1.0 in MeOH).

1'-Oxo: see 3-Butyloctahydro-5-propaenylindolizine, B-436

(±)-**form** [81076-50-8]

Synthetic. Noncryst.

Hydrochloride:

Flat needles (EtOAc/Et₂O). Mp 131°.

Hydrobromide:

Flat needles (Me₂CO). Mp 147-150°.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)

Spande, T.F. *et al.*, *Experientia*, 1981, **37**, 1242 (*ms, cmr, struct*)

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)

Iida, H. *et al.*, *J.A.C.S.*, 1985, **107**, 5534 (*synth, cmr, ms*)

Royer, J. *et al.*, *Tet. Lett.*, 1985, **26**, 1515 (*synth, cmr, ms, bibl*)

Pryzbylska, M. *et al.*, *Acta Cryst. C*, 1986, **42**, 832 (*cryst struct*)

Daly, J.W. *et al.*, *J. Nat. Prod.*, 1986, **49**, 265 (*isol, ms*)

Broka, C.A. *et al.*, *J.O.C.*, 1986, **51**, 5043 (*synth, cmr, ms*)

Tokuyama, T. *et al.*, *Tetrahedron*, 1986, **42**, 3453 (*isol, cmr*)

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1988, **66**, 1163 (*synth, pmr, cmr, ms*)

Brandi, A. *et al.*, *J.O.C.*, 1989, **54**, 1748 (*synth, ir, pmr, cmr, ms*)

Watanabe, Y. *et al.*, *J.O.C.*, 1989, **54**, 4088 (*synth, ir, pmr, cmr, ms*)

Machinaga, N. *et al.*, *Chem. Comm.*, 1991, 405 (*synth, abs config, 239CD*)

Yang, T.-K. *et al.*, *Heterocycles*, 1994, **38**, 1711 (*synth*)

Pilli, R.A. *et al.*, *J.O.C.*, 1995, **60**, 717-722 (*synth, bibl*)

Momose, T. *et al.*, *J.C.S. Perkin 1*, 1997, 1315 (*synth*)

Celimene, C. *et al.*, *Tetrahedron*, 1998, **54**, 10457-10468 (*synth*)

Clark, R.B. *et al.*, *Org. Lett.*, 1999, **1**, 349-351 (*synth, Indolizidine 239CD*)

Lee, E. *et al.*, *Org. Lett.*, 2000, **2**, 2169-2171 (*synth*)


Kiewel, K. *et al.*, *Tet. Lett.*, 2001, **42**, 6621-6623 (*synth*)

Toyooka, N. *et al.*, *Heterocycles*, 2005, **66**, 549-555 (*synth*)

Smith, A.B. *et al.*, *J.O.C.*, 2006, **71**, 2547-2557 (*synth*)

Octahydro-5,7-dimethylindolizine, 9CI O-42

5,7-Dimethylindolizidine



(5R,7S,8aR)-form

C₁₀H₁₉N 153.267

(5R,7S,8aR)-**form**

Dendroprimine

[40169-68-4]

Alkaloid from *Dendrobium primulinum* (Orchidaceae). Oil. $[\alpha]_D$ -38 (c, 1.0 in CHCl₃). n_D^{25} 1.4732.

Hydrochloride: Mp 194-195°.

Hydrobromide: Mp 206-207°.

Methiodide: Mp 263-264°.

(5R,7S,8aS)-**form**

8a-Epidendroprimine

Pale yellow oil (as hydrochloride). $[\alpha]_D^{20}$ -45.8 (c, 0.72 in CHCl₃). Could not be crystallised.

Lüning, B. *et al.*, *Acta Chem. Scand.*, 1965, **19**, 1607-1611 (*isol, ir, pmr, ms, struct*)

Blomqvist, L. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 3203-3206 (*abs config*)


Diederich, M. *et al.*, *Synthesis*, 1999, 286-289 (8a-Epidendroprimine, *synth*)

De Saboulin Bollena, A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1029-1031 (*synth, pmr, cmr*)

Kobayashi, T. *et al.*, *Org. Lett.*, 2006, **8**, 3813-3816 (*synth*)

2-[Octahydro-4,7-dimethyl-1-oxocyclopent[a]c[pyran-3-yl]nepetalactam O-43

[115421-74-4]




C₂₀H₂₉NO₃ 331.454

Mixed dimer of Nepetalactam, N-151 and Nepetalic acid. Isol. from a commercial sample of catnip oil (*Nepeta cataria*) (Lamiaceae). Cryst. (2-methylpentane). Mp 116-117°. $[\alpha]_D^{25}$ +3.8 (c, 5 in CHCl₃). Artifact.

Eisenbraun, E.J. *et al.*, *J.O.C.*, 1988, **53**, 3968-3972 (*isol, uv, ir, pmr, cmr, ms, struct*)

Octahydro-5,8-dipropyl-1H-indolizine, 9CI O-44

5,8-Dipropylindolizidine. **Indolizidine 209I**. *Dendrobates Alkaloid 209I*



(5R,8R,8aS)-form

C₁₄H₂₇N 209.374

(5R,8R,8aS)-**form**

Oil. $[\alpha]_D^{24}$ -126.5 (c, 0.2 in Me₂CO). $[\alpha]_D^{22}$ -92.1 (c, 1 in CHCl₃).

(5ξ,8ξ,8αξ)-**form**

Alkaloid from a member of the Dendrobatidae. Isolation details inaccessible,


stereochem. of nat. isolate not clear or may be a mixture.

Michel, P. *et al.*, *J.O.C.*, 2000, **65**, 8908-8918 (*synth, pmr, cmr*)

Hu, S. *et al.*, *J.O.C.*, 2005, **70**, 7364-7370 (*synth, pmr, cmr*)

1,2,3,4,6,7,8,9-Octahydrodi-pyrido[1,2-a:1',2'-c]imidazol-10-ium(1+), 8CI O-45

Anosmine



C₁₁H₁₇N₂[⊕] 177.269

Quaternary alkaloid from *Dendrobium anosmum* and *Dendrobium parishii* (Orchidaceae).

Chloride: [18813-73-5]

Mp 86-88.5° Mp 153-156° (double Mp). Hygroscopic.

Bromide: [18813-72-4]

Needles (Me₂CO). Mp 164-165°.


Leander, K. *et al.*, *Tet. Lett.*, 1968, 905 (*isol, ir, uv, pmr, synth, struct*)

Söderberg, E. *et al.*, *Acta Chem. Scand.*, 1970, **24**, 397 (*cryst struct*)

Hemscheidt, T. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1281 (*biosynth*)

Octahydro-1,2-indolizinediol, 9CI O-46

1,2-Dihydroxyindolizidine. **1-Azabicyclo[4.3.0]nonane-7,8-diol**



(1S,2S,8aS)-form

C₈H₁₅NO₂ 157.212

Considerable confusion regarding the abs. config. of natural lentiginosine. The compd. isol. by Pastuszak *et al* ($[\alpha]_D$ -3.3) was attributed the (1S,2S,8aS) config. based on biogenetic reasoning. The (1S,2S,8aS) enantiomer synthesised independently by Yoda *et al* and Cordero *et al* had opt. rotns. of $[\alpha]_D$ = +0.19 and +3.2 respectively. The most recent research (Brandi *et al*, 1995) based on synth. of both enantiomers and their inhibition of amyloglucosidases, demonstrates that natural lentiginosine has the (1S,2S,8aS) config. and that it is dextrorotatory.

(1R,2R,8aR)-**form**

Synthetic. Mp 106-107°. $[\alpha]_D$ -2.6 (c, 1.0 in MeOH). $[\alpha]_D^{23}$ -1.6 (c, 0.24 in MeOH).

(1S,2R,8aR)-**form**

2,8-Diepilentiginosine [108866-44-0]

Synthetic. Cryst. (hexane). Mp 109-111°. $[\alpha]_D^{24}$ -37.1 (c, 0.55 in MeOH).

1,2-O-Isopropylidene: *Octahydro-2,2-di-*

methyl-1,3-dioxolo[4,5-a]indolizine,
9CI
C₁₁H₁₉NO₂ 197.277
Solid. Mp 89-91°. [α]_D²³ -38 (c, 0.95 in MeOH).

(1S,2R,8aS)-form
2-Epilentiginosine

Alkaloid from leaves of *Astragalus lentiginosus* and shoots of *Astragalus oxyphyphus* (Fabaceae). Also isol. from the fungus *Rhizoctonia leguminicola*. Oil. [α]_D²⁴ -32.5 (c, 0.13 in CHCl₃).

(1S,2S,8aR)-form

Solid (C₆H₆). Mp 137-138°. [α]_D²⁷ -5.8 (c, 0.88 in MeOH).

(1S,2S,8aS)-form

Lentiginosine

[125279-72-3]

Alkaloid from leaves of *Astragalus lentiginosus* (Fabaceae). Potent inhibitor of the fungal α -glucosidase, amyloglucosidase (twice as potent as Castanospermine, C-188), but does not inhibit other α -glucosidases (e.g. sucrase, maltase, yeast α -glucosidase) nor any other glycosidases. Mp 108-109°. [α]_D²⁵ +3.2 (c, 0.6 in MeOH).

Harris, T.M. *et al.*, *J.O.C.*, 1987, **52**, 3094-3098

(*isol, cmr, 2,8-Diepilentiginosine, synth*)

Harris, C.M. *et al.*, *Tet. Lett.*, 1988, **29**, 4815-4818 (*isol*)

Pastuszak, I. *et al.*, *Biochemistry*, 1990, **29**, 1886 (*isol, pmr, cmr, ms*)

Yoda, H. *et al.*, *Tetrahedron: Asymmetry*, 1993, **4**, 1455 (*synth*)

Cordero, F.M. *et al.*, *Tet. Lett.*, 1994, **35**, 949 (*synth*)

Gurjar, M.K. *et al.*, *Tet. Lett.*, 1994, **35**, 8871 (*synth, abs config*)

Brandi, A. *et al.*, *J.O.C.*, 1995, **60**, 6806 (*synth, abs config*)

Paolucci, C. *et al.*, *Synthesis*, 1997, 1415-1419 (*1S,2S,8aR-form*)

Ha, D.-C. *et al.*, *J.O.C.*, 2000, **65**, 621-623 (*synth*)

Paolucci, C. *et al.*, *J.O.C.*, 2001, **66**, 4787-4794 (*2,8-Diepilentiginosine, synth, pmr, cmr*)

Rasmussen, M.O. *et al.*, *J.O.C.*, 2001, **66**, 5438-5443 (*synth, pmr, cmr, bibl*)

Yoda, H. *et al.*, *Tet. Lett.*, 2001, **42**, 2509-2512 (*synth*)

Chandra, K.L. *et al.*, *J.O.C.*, 2002, **67**, 4630-4633 (*Lentiginosine, synth*)

Feng, Z.-X. *et al.*, *Tet. Lett.*, 2003, **44**, 497-498 (*Lentiginosine, synth*)

Sha, C.-K. *et al.*, *Tet. Lett.*, 2003, **44**, 499-501 (*synth*)

Ayad, T. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 2626-2631 (*Lentiginosine, synth*)

Kim, I.S. *et al.*, *Org. Lett.*, 2006, **8**, 4101-4104 (*synth*)

Chaudhari, V.D. *et al.*, *Tetrahedron*, 2006, **62**, 4349-4354 (*synth*)

Cardona, F. *et al.*, *Eur. J. Org. Chem.*, 2007, 1551-1565 (*Lentiginosine, rev*)

Angle, S.R. *et al.*, *J.O.C.*, 2007, **72**, 5592-5597 (*Lentiginosine, synth*)

Chen, M.-J. *et al.*, *Tet. Lett.*, 2007, **48**, 6271-6274 (*synth*)

Alam, M.A. *et al.*, *Tet. Lett.*, 2008, **49**, 5534-5536 (*Lentiginosine, synth*)


Shi, G.-F. *et al.*, *Tetrahedron*, 2008, **64**, 5005-5012 (*synth*)

Chandrasekhar, S. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 746-750 (*Lentiginosine, synth*)

Octahydro-8-indolizine-methanol, 9CI

O-47

8-Hydroxymethylindolizidine
[126371-98-0, 154905-41-6]



(8R,8aS)-form

C₉H₁₇NO 155.239

(8R,8aS)-form [111975-29-2]

Pale yellow oil. [α]_D²² -25.9 (c, 1.16 in EtOH).

(8S,8aR)-form

Amorph. solid. Mp 35°. [α]_D²⁰ +44.7 (c, 1.1 in EtOH).

(8R*,8aS*)-form

Tashiromine

[128573-77-3]

Alkaloid from the stems of *Maackia tashiroi* (Fabaceae). Oil. No opt. rotn. was reported for the nat. prod., therefore the abs. config. cannot be assigned.

Ohmiya, S. *et al.*, *Heterocycles*, 1990, **30**, 537 (*isol, pmr, cmr, ms, struct*)

Nagao, Y. *et al.*, *J.O.C.*, 1990, **55**, 1148-1156 (*synth, ir, pmr, cmr*)

Paulvannan, K. *et al.*, *J.O.C.*, 1994, **59**, 1613 (*synth*)

Gage, J.L. *et al.*, *Tet. Lett.*, 1997, **38**, 7007-7010 (*synth*)

David, O. *et al.*, *J.O.C.*, 1999, **64**, 3122-3131 (*synth, pmr, cmr*)

David, O. *et al.*, *Heterocycles*, 2001, **55**, 1689-1701 (*Tashiromine, synth*)

Bates, R.W. *et al.*, *J.C.S. Perkin 1*, 2001, 654-656 (*Tashiromine, synth*)

Dieter, R.K. *et al.*, *Tet. Lett.*, 2002, **43**, 7725-7728 (*synth*)


Bélanger, G. *et al.*, *J.O.C.*, 2006, **71**, 704-712 (*synth*)

Pohmakotr, M. *et al.*, *Tetrahedron*, 2008, **64**, 2339-2347 (*synth*)

1,2,3,4,6,7,12,12b-Octahydroindolo[2,3-a]quinolizine, 9CI

O-48

Indoloquinolizidine. Tetramethylenetetrahydro- β -carboline
[4802-79-3]



(S)-form

C₁₅H₁₈N₂ 226.321

Indole alkaloid, not secologanin-derived.

(R)-form [15051-70-4]

Mp 149-151°. [α]_D +84.8 (MeOH).

(S)-form [10252-12-7]

Main alkaloid from leaves of *Dracontomelon mangiferum* (Anacardiaceae). Also from *Nitraria schoberi*. Mp 149-151°. [α]_D -86.5 (MeOH). The alkaloid is partly racemic. λ_{\max} 228 (log ϵ 4.43); 284 (log ϵ 3.5); 292 (log ϵ 3.43) (EtOH).

10-Bromo: Arborescidine A

[147395-92-4]

C₁₅H₁₇BrN₂ 305.217

Alkaloid from the marine tunicate *Pseudodistoma arborescens*. Cryst. (MeOH). Mp 202°. [α]_D -85 (c, 1 in CHCl₃).

(±)-form [46798-86-1]

Alkaloid from *Dracontomelon mangiferum* (Anacardiaceae). Inhibitor of spontaneous motor activity, antiphlogistic agent. Mp 153-155° (148-151.5°). Many syntheses reported.

Hydrochloride: Mp 311-312°.

N³-Oxide (cis-): [82484-93-3]

[82484-94-4]

C₁₅H₁₈N₂O 242.32

Alkaloid from *Nitraria komarovii*. Cryst. (Me₂CO). Mp 213-214°.

Groves, J.H. *et al.*, *J.C.S.*, 1952, 650-661 (*synth*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1951-1954 (*S-form, isol, uv, ir, pmr, ms, struct*)

Pospišek, J. *et al.*, *Chem. Ind. (London)*, 1969, 25-26 (*resoln, ord, config*)

Pakhritdinov, B.M. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 641-642; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 663-664 (*S-form, isol, uv*)

Akimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 2614-2623 (*synth*)

Gribble, G.W. *et al.*, *J.O.C.*, 1974, **39**, 1845-1850; 1975, **40**, 3720-3725 (*ms, cmr*)

Nakagawa, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 304-312 (*synth, bibl*)

Rosenmund, P. *et al.*, *Annalen*, 1979, 1643-1656 (*synth, uv*)

Meyers, A.I. *et al.*, *J.O.C.*, 1986, **51**, 3108-3112 (*synth*)

Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1990, **26**, 560-561; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 478-479 (*N-oxide*)

Hua, D.H. *et al.*, *J.O.C.*, 1991, **56**, 6998-7007 (*synth*)

Chbani, M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 99-104 (*Arborescidine A*)

Cheng, E. *et al.*, *Tetrahedron*, 1996, **52**, 6725-6732 (*synth*)

Diker, K. *et al.*, *J. Nat. Prod.*, 1997, **60**, 790-793 (*synth, ir, uv, pmr, cmr*)

Burm, B.E.A. *et al.*, *Tetrahedron*, 1998, **54**, 6135-6146 (*Arborescidine A, synth*)

Santos, L.S. *et al.*, *J.O.C.*, 2004, **69**, 1283-1289 (*Arborescidine A, synth*)

Allin, S.M. *et al.*, *Eur. J. Org. Chem.*, 2005, 4179-4186 (*S-form, synth, pmr, cmr*)


King, F.D. *et al.*, *J. Het. Chem.*, 2007, **44**, 1459-1463 (*synth, pmr, cmr*)

Szawalko, J. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 406-413 (*R-form, synth*)

1,2,3,4,6,7,12,12b-Octahydroindolo[2,3-a]quinolizine-4-carboxylic acid

O-49

Indoloquinolizidine-4-carboxylic acid



C₁₆H₁₈N₂O₂ 270.33

(4R,12bS)-form


Nitrile: 1,2,3,4,6,7,12,12b-Octahydroindolo[2,3-a]quinolizine-4-carbonitrile.

4-Cyanoindoloquinolizidine. Phoebe-grandine E


[909118-77-0]

C₁₆H₁₇N₃ 251.33Alkaloid from the leaves of *Phoebe grandis*. Amorph. solid. [α]_D²³ -38.4 (c, 0.17 in CHCl₃). λ_{\max} 233 (log ϵ 3.95); 300 (log ϵ 3.6); 339 (log ϵ 3.87) (MeOH).Awang, K. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 567-572 (*isol, pmr, cmr, ms*)**1,1',4,4',5,5',6,6'-Octahydro-5-methoxy-2,3'-bipyridine, 9CI**


[139339-33-6]

C₁₁H₁₈N₂O 194.276**(ξ)-form**Alkaloid from the exudate of *Jatropha gossypifolia* (Euphorbiaceae). Mp 201-203°.Ahmad, M.U. *et al.*, *Indian J. Chem., Sect. B*, 1992, **31**, 67 (*isol, ir, pmr, cmr, ms, struct*)**Octahydro-8-methyl-5-(6-nonen-8-ynyl)-1H-indolizine, 9CI****8-Methyl-5-(6-nonen-8-ynyl)indolizidine. Indolizidine 259B.** *Dendrobates Alkaloid 259B*

[150044-82-9]

C₁₈H₂₉N 259.434Alkaloid from the poison frog *Dendrobates pumilio*.Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 357-373 (*isol, ms*)**Octahydro-8-methyl-5-(4-pentenyl)-1H-indolizine****8-Methyl-5-(4-pentenyl)indolizidine.***Dendrobates Alkaloid 207A. Indolizidine 207A*

[141724-48-3]

C₁₄H₂₅N 207.358Alkaloid from skin extracts of *Dendrobates histrionicus* and *Dendrobates speciosus* (Dendrobatidae) and from the Madagascar frog *Mantella madagascariensis* (Ranidae, subfamily Mantellinae). Oil. [α]_D -85.25 (c, 0.174 in EtOH). Opt. rotn. refers to synthetic material.**Dihydro: Octahydro-8-methyl-5-pentylindolizine. 8-Methyl-5-pentylindolizidine. Dendrobates Alkaloid 209B. Indolizidine 209B**

[117959-79-2]

[131319-70-5]


C₁₄H₂₇N 209.374Trace alkaloid from skin extracts of an undescribed *Dendrobates* sp. from Panama (Dendrobatidae). Mp 184-185° (as hydrochloride). [α]_D²² -94.3 (c, 1.85 in MeOH). Enantiomeric excess judged to be >95%. Opt. rotn. of natural 209B not reported.**4',5'-Didehydro: Octahydro-8-methyl-5-(4-pentynyl)-1H-indolizine. 8-Methyl-5-(4-pentynyl)indolizidine. Dendrobates Alkaloid 205A. Indolizidine 205A**

[109175-46-4]

[130979-70-3, 118015-96-6]

C₁₄H₂₃N 205.342Alkaloid from skin extracts of the Panamanian poison frog *Dendrobates pumilio*. [α]_D -35 (c, 0.24 in MeOH).**2',3',Z,4',5'-Tetrahydro: Octahydro-8-methyl-5-(2-penten-4-ynyl)-1H-indolizine. 8-Methyl-5-(2-penten-4-ynyl)indolizidine. Dendrobates Alkaloid 203A. Indolizidine 203A**

[120328-23-6]

C₁₄H₂₁N 203.327Alkaloid from skin extracts of *Dendrobates auratus*. [α]_D -23.3 (c, 0.3 in CHCl₃).Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163 (*isol, ms, rev*)Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205 (*ms, rev*)Daly, J.W. *et al.*, *Toxicol.*, 1984, **22**, 905 (*isol, ms, nomencl*)Tokuyama, T. *et al.*, *Tetrahedron*, 1987, **43**, 643; 1991, **47**, 5401 (*isol, pmr, cmr, ms, struct, derivs*)Edwards, M.W. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1188 (*isol, pmr, ms*)Smith, A.L. *et al.*, *J.A.C.S.*, 1988, **110**, 8696 (*synth*)Gneccho, D. *et al.*, *Chem. Comm.*, 1991, 625-626 (*synth*)Holmes, A.B. *et al.*, *J.O.C.*, 1991, **56**, 1393 (*synth*)Taber, D.F. *et al.*, *J.O.C.*, 1995, **60**, 529 (*synth*)Comins, D.L. *et al.*, *J.O.C.*, 1997, **62**, 8182-8187 (*Indolizidine 205A, synth*)Toyooka, N. *et al.*, *Tetrahedron*, 1997, **53**, 9553 (*synth*)Michael, J.P. *et al.*, *J.C.S. Perkin 1*, 2000, 1919-1928 (*Indolizidine 209B, synth*)Back, T.G. *et al.*, *J.O.C.*, 2000, **65**, 4543-4552 (*synth*)Michel, P. *et al.*, *J.O.C.*, 2000, **65**, 8908-8918 (*synth*)Shu, C. *et al.*, *J.A.C.S.*, 2001, **123**, 12477-12487 (*Indolizidine 209B, synth*)Song, Y. *et al.*, *Tet. Lett.*, 2002, **43**, 8635-8637 (*Indolizidine 209B, synth*)Davis, F.A. *et al.*, *Org. Lett.*, 2003, **5**, 5011-5014 (*Indolizidine 209B, synth*)Toyooka, N. *et al.*, *Chem. Pharm. Bull.*, 2005, **53**, 555-560 (*synth*)**Octahydro-4-methyl-2H-quinolizin-2-one, 9CI****(4R,9aR)-form**C₁₀H₁₇NO 167.25**(4R,9aR)-form****Myrtiline**

[66835-10-7]

Alkaloid from *Vaccinium myrtillus* (bilberry) aerial parts (Ericaceae). Oil. [α]_D²⁰ +3.1 (c, 2.1 in CHCl₃) (natural). [α]_D²³ +11.3 (c, 2.7 in CHCl₃) (synthetic). Isol. sample has only 25% opt. purity. This is not due to isom. during isol.**(4R,9aS)-form****Epimyrtine**

[82111-02-2]

Alkaloid from *Vaccinium myrtillus* (bilberry) (Ericaceae). Oil. [α]_D²⁰ -2.5 (c, 1.2 in CHCl₃) (natural). [α]_D²³ -18 (c, 5.2 in CHCl₃) (synthetic). Low opt. purity.**Hydrochloride:** Mp 211-213° dec. [α]_D²³ -31.7.**(4RS,9aRS)-form** [66900-13-8]

[82083-37-2]

Mp 196-198° dec. (as hydrochloride).


(4RS,9aSR)-form [66900-14-9]

[82083-36-1]

Mp 200-201° dec. (as hydrochloride).

Slosse, P. *et al.*, *Tetrahedron*, 1981, **37**, 4287 (*isol, ir, ms, pmr, cmr, synth, resoln*)Comins, D.L. *et al.*, *J.O.C.*, 1992, **57**, 5807 (*synth*)Gelas-Miakhe, Y. *et al.*, *Tet. Lett.*, 1992, **33**, 73 (*synth*)Pilli, R.A. *et al.*, *J.O.C.*, 1995, **60**, 717 (*synth*)Davis, F.A. *et al.*, *J.O.C.*, 2003, **68**, 8061-8064 (*Epimyrtine, synth*)Back, T.G. *et al.*, *J.O.C.*, 2005, **70**, 967-972 (*synth*)Amorde, S.M. *et al.*, *Org. Lett.*, 2005, **7**, 2031-2033 (*synth*)**1,2,3,4,6,7,8,9-Octahydrophenazine****Polycartine B**

[4006-50-2]

C₁₂H₁₆N₂ 188.272Minor constit. of scent gland of the Canadian beaver (*Castor fiber*). Also isol. from the fruits of *Idesia polycarpa* (Flacourtiaceae). Flavour enhancer. Mp 112-113° (106-108°). Bp₁₁ 135-170°. λ_{\max} 260; 323 (MeOH) (Derep).

Hydrochloride: [42383-40-4]
Mp 160-162°.

N,N'-Dioxide: [24716-05-0]
C₁₂H₁₆N₂O₂ 220.271
Cryst. Mp 230°.

Schäffler, A. *et al.*, *Chem. Ber.*, 1955, **88**, 767
(*synth, uv*)

Fellman, J.H. *et al.*, *J.O.C.*, 1956, **21**, 713
(*synth*)

Wilens, S.H. *et al.*, *Chem. Ind. (London)*, 1969,
237 (*synth*)

Smith, H.E. *et al.*, *Chem. Comm.*, 1970, 1112
(*uv*)


Scheinbaum, M.L. *et al.*, *J.O.C.*, 1970, **35**,
2790 (*oxide*)

Maurer, B. *et al.*, *Helv. Chim. Acta*, 1976, **59**,
1169 (*isol*)

Moritake, M. *et al.*, *Tet. Lett.*, 1987, **28**, 1425
(*isol*)

Octahydro-3-propylindolizine O-55

3-Propylindolizidine



C₁₁H₂₁N 167.294


(3R*,8aS*)-form [929631-64-1]

Isol. from the venom of *Myrmecaria melanogaster*.

Jones, T.H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 160-
168 (*isol, synth, ir, ms*)

Octahydro-5-propylindolizine, 9CI O-56

5-Propylindolizidine. Dendrobates *Alkaloid 167B*. Gephyrotoxin *167B*. *Indolizidine 167B*



(5R,8aR)-form

C₁₁H₂₁N 167.294

(5R,8aR)-form

(-)-trans-form
[120057-35-4]

Trace alkaloid from skin extracts of an undescribed *Dendrobates sp.* from Panama (Dendrobatidae). Oil. [α]_D²⁰ -106.3 (c, 0.80 in hexane).

(5S,8aR)-form

(-)-cis-form

[129893-55-6]

Oil. [α]_D -1.7 (c, 1.1 in CH₂Cl₂).

(5S,8aS)-form

(+)-trans-form

[168610-27-3]

Oil. [α]_D²⁶ +28.1 (c, 1.125 in CH₂Cl₂).

(5RS,8aRS)-form

(±)-trans-form

[130884-95-6 (hydrochloride)]
Mp 167-169° (as hydrochloride).

[134357-40-7]

Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*,
1982, **41**, 205 (*ms, rev*)

Smith, A.L. *et al.*, *J.A.C.S.*, 1988, **110**, 8696
(*synth, struct*)

Polniaszek, R.P. *et al.*, *J.O.C.*, 1990, **55**, 4688-
4693 (*ir, pmr, cmr, synth*)

Jefford, C.W. *et al.*, *J.A.C.S.*, 1991, **113**, 3513
(*synth, ms, pmr, cmr*)

Holmes, A.B. *et al.*, *J.O.C.*, 1991, **56**, 1393
(*synth, ir, pmr, cmr, ms*)

Zeller, E. *et al.*, *Synlett*, 1991, 44 (*synth*)

Takahata, H. *et al.*, *Heterocycles*, 1995, **41**,
1797 (*synth, ir, pmr*)

Angle, S.R. *et al.*, *J.O.C.*, 1997, **62**, 8549-8552
(*synth*)

Weymann, M. *et al.*, *Synthesis*, 1997, 1151-
1160 (*synth, pmr, cmr*)

Michael, J.P. *et al.*, *Eur. J. Org. Chem.*, 1998,
865-870 (*synth*)

Chenevert, R. *et al.*, *Heterocycles*, 1999, **51**,
593-598 (*synth*)

Chalard, P. *et al.*, *Tet. Lett.*, 1999, **40**, 1661-
1664 (*synth*)

Back, T. *et al.*, *J.O.C.*, 2000, **65**, 4543-4552
(*synth*)

Yamazaki, N. *et al.*, *Org. Lett.*, 2000, **2**, 465-
467 (*synth*)

Yoda, H. *et al.*, *Tet. Lett.*, 2001, **42**, 2509
(*synth*)

Peroche, S. *et al.*, *Tet. Lett.*, 2001, **42**, 4617-
4619 (*synth*)

Kiewel, K. *et al.*, *Tet. Lett.*, 2001, **42**, 6621-
6626 (*synth*)

Kim, G. *et al.*, *Tetrahedron: Asymmetry*, 2001,
12, 2073-2076 (*synth*)

Carbonnel, S. *et al.*, *Heterocycles*, 2002, **57**,
1807-1830 (*synth*)

Corvo, M.C. *et al.*, *Tet. Lett.*, 2002, **43**, 455-
458 (*synth*)

Zaminer, J. *et al.*, *Tet. Lett.*, 2002, **43**, 6739-
6741 (*synth*)

Reddy, P.G. *et al.*, *J.O.C.*, 2004, **69**, 3093-3101
(*synth*)


Guazzelli, G. *et al.*, *Synthesis*, 2005, 3119-3123
(*synth*)

Chang, M.-Y. *et al.*, *Heterocycles*, 2007, **71**,
933-940 (*synth*)

Kuhakarn, C. *et al.*, *Tetrahedron*, 2008, **64**,
1663-1670 (*synth*)

Pohmakotr, M. *et al.*, *Tetrahedron*, 2008, **64**,
2339-2347 (*synth*)

Octahydro-1H,3H-pyrido[3,2,1-ij]benzoxazine O-57



(3aR,6aS,10aR)-form

C₁₁H₁₉NO 181.277

(3aR,6aS,10aR)-form

Myrioxazine A

[481012-19-5]

Alkaloid from the leaves of *Myrioneuron nutans*. Oil. [α]_D²⁰ +21 (c, 1 in MeOH).

(3aS,6aS,10aR)-form

Myrioxazine B

[481012-21-9]


Alkaloid from the leaves of *Myrioneuron nutans*. Oil. [α]_D²⁰ +9 (c, 1

in MeOH).

Pham, V.C. *et al.*, *Tet. Lett.*, 2002, **43**, 7565-
7568 (*isol, synth, pmr, cmr, abs config*)

Octahydro-2H-quinolizine-1-methanol, 9CI O-58

1-Hydroxymethylquinolizidine. *Lupinine*



(-)-form

C₁₀H₁₉NO 169.266

Insect antifeedant. Antiinflammatory, antiarrhythmic agent. Oxytocic agent. Possesses cathartic, diuretic props. Log P 0.85 (calc).

(+)-form

Synthetic. Mp 68°. [α]_D +19.9.

(-)-Tartrate salt: Mp 167-168°. [α]_D -
15.8 (H₂O).

(-)-form [486-70-4]

Alkaloid from *Anabasis aphylla*, *Lupinus luteus*, *Lupinus palmeri* and a few other *Lupinus* spp. (Chenopodiaceae, Fabaceae). Mp 70-71°. Bp 255-257°. [α]_D¹⁷ -20.4 (EtOH). Pharmacol. active isomer.

▶LD₅₀ (mus, ivn) 15 mg/kg. OK5802000

Hydrochloride: [6113-09-3]

Mp 212-213° (207-209°). [α]_D -14
(H₂O).

Picrate: [6113-13-9]

Mp 136-137° Mp 196-197° (dimorph.).

Butanoyl: *Butanoyllupinine*

[38976-65-7]

C₁₄H₂₅NO₂ 239.357

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

2-Methylbutanoyl: [135531-66-7]

C₁₅H₂₇NO₂ 253.384

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

Benzoyl: [40179-92-8]

Mp 49-50°.

O-4-Hydroxy-E-cinnamoyl: (*4-Hydroxycinnamoyl*)lupinine

C₁₉H₂₅NO₃ 315.411

Alkaloid from fresh seedlings of *Lupinus luteus* (Fabaceae).

Highly viscous oil. Mp 130-131° (as 4-nitrobenzoyl). The *Z*-isomer of this alkaloid (and its relatives) are present as artifacts in samples of the alkaloids.

O-[β-D-Glucopyranosyl-(1→4)-4-hydroxy-E-cinnamoyl]: (*4'-β-D-Glucopyranosyloxycinnamoyl*)lupinine

[70155-14-5]

C₂₅H₃₅NO₈ 477.553

Alkaloid from *Lupinus luteus* seedlings (Fabaceae). Amorph. solid. [α]_D²² -58.9 (c, 0.3 in EtOH).

O-[α-L-Rhamnopyranosyl-(1→4)-4-hydroxy-E-cinnamoyl]: (*4'-α-L-Rhamnopyranosyloxycinnamoyl*)lupinine

[65526-78-5]

C₂₅H₃₅NO₇ 461.554

Alkaloid from *Lupinus luteus* seedlings (Fabaceae). $[\alpha]_D^{22}$ -105 (c, 1.16 in MeOH).

O-(4-Hydroxy-3-methoxy-E-cinnamoyl): ***o*-Feruloyloxylupinane**
[18161-87-0]
C₂₀H₂₇NO₄ 345.438
Alkaloid from green parts of *Lupinus luteus* (Fabaceae). Cryst. (C₆H₆). Mp 118° Mp 130° (dimorph.).

O-[β-D-Glucopyranosyl-(1→4)-4-hydroxy-3-methoxy-E-cinnamoyl]: (**4'-O-β-D-Glucopyranosyloxy-3'-methoxycinnamoyl**)lupinine
[71926-09-5]
C₂₆H₃₇NO₉ 507.58
Alkaloid from fresh seedlings of *Lupinus luteus* (Fabaceae). Amorph. solid. $[\alpha]_D^{22}$ -48.9 (c, 0.06 in EtOH).

O-[α-L-Rhamnopyranosyl-(1→4)-4-hydroxy-3-methoxy-E-cinnamoyl]: (**4'-α-L-Rhamnosyloxy-3'-methoxycinnamoyl**)lupinine
[70155-16-7]
C₂₆H₃₇NO₈ 491.58
Alkaloid from seedlings of *Lupinus luteus* (Fabaceae). $[\alpha]_D$ -78.3.

5-Epimer: see Epilupinine, E-95

(±)-form [10248-30-3]

Mp 63-64° (59°).

Picrate: Mp 127°.

Cassola, M. et al., *Annalen*, 1835, **13**, 308 (isol)

Karrer, P. et al., *Helv. Chim. Acta*, 1928, **11**, 1062 (isol, struct)

Couch, J.F. et al., *J.A.C.S.*, 1934, **56**, 2434 (isol)

Clemo, G.R. et al., *J.C.S.*, 1937, 965; 1938, 1574 (synth, resoln)

Cookson, R.C. et al., *Chem. Ind. (London)*, 1953, 337 (abs config)

Thomas, A.F. et al., *Can. J. Chem.*, 1955, **33**, 1290 (ir)

Podkowinska, H. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Biol.*, 1965, **13**, 623; 1967, **15**, 467; *CA*, **64**, 18025h (Feruloyloxylupinane)

Goldberg, S.I. et al., *J.O.C.*, 1967, **32**, 1046 (synth)

Kappe, Th. et al., *Monatsh. Chem.*, 1967, **98**, 1852 (synth)

Schöpf, C. et al., *Annalen*, 1968, **712**, 168 (synth)

Ishbaev, A.I. et al., *Khim. Prir. Soedin.*, 1972, **8**, 328; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 322 (ord)

Bohlmann, F. et al., *Chem. Ber.*, 1975, **108**, 1043 (cmr)

Bohlmann, F. et al., *Phytochemistry*, 1975, **14**, 155 (biosynth)

Murakoshi, I. et al., *Phytochemistry*, 1975, **14**, 2714; 1977, **16**, 2046; 1978, **17**, 1817; 1979, **18**, 699 (cinnamoyl derivs)

Gerrans, G.C. et al., *Tet. Lett.*, 1975, 4147 (synth)

Tufariello, J.J. et al., *Tet. Lett.*, 1976, 4037 (synth)

Murakoshi, I. et al., *Chem. Pharm. Bull.*, 1977, **25**, 527 (cinnamoyl derivs, biosynth)

Murakoshi, I. et al., *Chem. Pharm. Bull.*, 1979, **27**, 144 (glucosylcinnamoyl derivs)

Iwashita, T. et al., *J.O.C.*, 1982, **47**, 230 (synth, ir, pmr)

Kozioł, A.E. et al., *Acta Cryst. C*, 1983, **39**, 1375 (cryst struct, abs config)

Podkowinska, H. et al., *Org. Magn. Reson.*, 1984, **22**, 379 (cmr)

Golebiewski, W.M. et al., *Can. J. Chem.*, 1985, **63**, 2707 (biosynth)

Takahata, H. et al., *Chem. Pharm. Bull.*, 1986, **34**, 4523 (synth)

Aslanov, Kh.A. et al., *Alkaloids (Academic Press)*, 1987, **31**, 117 (rev)

Matsubara, Y. et al., *Chem. Pharm. Bull.*, 1988, **36**, 1597 (synth)

Grieco, P.A. et al., *J.O.C.*, 1988, **53**, 3325 (synth, ir, pmr)

Célérier, J.P. et al., *Tetrahedron*, 1989, **45**, 6161 (synth)

Morley, C. et al., *Tetrahedron: Asymmetry*, 1990, **1**, 147 (synth)

Veen, G. et al., *Phytochemistry*, 1991, **30**, 1891 (esters)

Hua, D.H. et al., *Synthesis*, 1991, 970 (synth)

Rycroft, D.S. et al., *Magn. Reson. Chem.*, (Spec. Issue), 1992, **30**, S15 (pmr)


Gesson, J.P. et al., *Tet. Lett.*, 1992, **33**, 3633 (synth)

Paulvannan, K. et al., *Tet. Lett.*, 1993, **34**, 215 (synth)

Morley, C. et al., *J.C.S. Perkin 1*, 1994, 2903 (synth)

Mangeny, P. et al., *Tetrahedron*, 1998, **54**, 10349-10362 (synth)

5-[[[(Octahydro-2H-quinolizin-1-yl)methyl]amino]-5-oxopentanoic acid, **9CI** **O-59**



C₁₅H₂₆N₂O₃ 282.382

(1R,9aR)-form

Me ester: 11-(4-Carboxybutyramido)-11-deoxy-1-epilupinine methyl ester. 5-[(3-Methoxycarbonylbutyryl)aminomethyl]quinolizidine
[18688-43-2]
C₁₆H₂₈N₂O₃ 296.409

Alkaloid from *Lamprolobium fruticosum* (Fabaceae). Cryst. Mp 77-78°. Considered to be an artifact derived from Lamprolobine, L-26 on MeOH treatment.

(1R,9aS)-form

Me ester, N-oxide: 5-(3-Methoxycarbonylbutyryl)aminomethyl-trans-quinolizidine N-oxide
C₁₆H₂₈N₂O₄ 312.408

Trace alkaloid from *Sophora tomentosa* (Fabaceae). Amorph. solid. Prob. artifact derived from Epilamprolobine N-Oxide in L-26.

(1ξ,9aξ)-form

Butyl ester: *Sophorine*†

[81037-26-5]

C₁₉H₃₄N₂O₃ 338.489


Alkaloid from *Sophora alopecuroides* (Fabaceae). Prob. biosynth. precursor of matrine and sparteine groups. Mp 59-60°. $[\alpha]_D^{23}$ -18.9 (c, 0.98 in EtOH). Probable struct., based on biogenetic considerations.

Hart, N.K. et al., *Aust. J. Chem.*, 1968, **21**, 1619 (Me ester)

Kuchkarov, S. et al., *Khim. Prir. Soedin.*, 1979, **15**, 413; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 364 (*Sophorine, isol*)

Kamaev, F.G. et al., *Khim. Prir. Soedin.*, 1981, **17**, 439; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 439 (*Sophorine, struct, pmr, cmr*)
Murakoshi, I. et al., *Phytochemistry*, 1981, **20**, 1725 (Me ester N-oxide)


7-[(1,2,3,4,4a,7,8,8a-Octahydro-1,2,4a,5-tetramethyl-1-naphthalenyl)methyl]-6-benzoxazolol, **9CI** **O-60**
[193157-85-6]



C₂₂H₂₉NO₂ 339.477

Not named in paper. Related to Avarol, A-1560. Constit. of a *Dysidea* sp. Oil.
Stewart, M. et al., *Aust. J. Chem.*, 1997, **50**, 341-347 (isol, pmr, cmr)

2,6,7,10,11,14,15,27-Octahydroxydotriacontanoic acid **O-61**



C₃₂H₆₄O₁₀ 608.852

Amide, 15-O-sulfate: *Turbinamide*


[366788-20-7]

C₃₂H₆₅NO₁₂S 687.931

Isol. from the tunicate *Sidnyum turbinatum*. Selective cytotoxic agent. Solid (as Na salt). $[\alpha]_D^{25}$ +7.1 (c, 0.003 in MeOH) (Na salt).

Aiello, A. et al., *Org. Lett.*, 2001, **3**, 2941-2944 (isol, pmr, cmr)

5-Octanoylamino-2,3,6(1H)-pyridinetrione **O-62**
N-(1,2,5,6-Tetrahydro-2,5,6-trioxo-3-pyridyl)octanamide




C₁₃H₁₈N₂O₄ 266.296

Oxidn. prod. of pigment from *Pseudomonas lemmonieri*. Mp 162°. λ_{max} 208 (log ε 4.02), 261(4.14) and 348 nm (3.8) (EtOH).

Ferguson, G. et al., *Chem. Comm.*, 1965, 640 (struct, ms, ir, pmr)

Octopamine, INN O-63

α -(Aminomethyl)-4-hydroxybenzenemethanol, 9CI. α -(Aminomethyl)-p-hydroxybenzyl alcohol, 8CI. 2-Amino-1-(4-hydroxyphenyl)ethanol. p-(β -Amino- α -hydroxyethyl)phenol. Norden. Norphen. Noroxedrine. Norsympathol. Norsynephrine. WIN 5512. Many other names [104-14-3]



C₈H₁₁NO₂ 153.18

The racemate and its enantiomers melt with dec. around 160° to form 2,5-bis(4-hydroxyphenyl)piperazine, which remelts around 250° or above.

▶ LD₅₀ (rat, orl) 1240 mg/kg. LD₅₀ (mus, ipr) 600 mg/kg. DN7350000

(R)-form [876-04-0]

Alkaloid from leaves of *Capsicum frutescens*, *Citrus* spp., *Cyperus rotundus*, *Cyperus papyrus*. Occurs in many animal tissues; found in high concs. in the posterior salivary gland of *Octopus vulgaris* and in *Octopus apollyon*, *Octopus bimaculatus*, *Rhizoglyphus echinopus*, *Fascaplysinopsis reticulata*. Invertebrate neurotransmitter. In vertebrates, prob. a cotransmitter with noradrenaline. Adrenergic vasodilator. Cryst. (H₂O). Mp 160° approx. [α]_D²⁵ -37.4 (c, 1 in H₂O). [α]_D²⁵ -56 (c, 1 in 1M HCl).

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyl-R-octopamine**
C₁₈H₁₉NO₅ 329.352
Constit. of the root bark of *Lycium chinense* (Chinese boxthorn). Antifungal agent.

N-Me: see Synephrine, S-659

O⁴-Me: **4-Methoxy- β -hydroxyphenethylamine**. α -(Aminomethyl)-4-methoxybenzenemethanol, 9CI. Alkaloid NP [46084-23-5]

[55275-61-1 ((±)-form)]

C₉H₁₃NO₂ 167.207

Minor alkaloid from *Coryphantha cornifera*, also detected in *Pereskia grandiflora* and *Pereskia chapistle* (Cactaceae). Shows weak vasoconstrictor and cardiac depressant props. Tentative identification, not obt. cryst. Opt. rotn. and abs. config. not detd.

(S)-form [826-01-7]

Synthetic. Less potent than (R)-form. Mp 160°. [α]_D²⁵ +37.2 (c, 1 in H₂O). [α]_D²⁵ +56.1 (c, 1 in 0.1N HCl).

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyl-S-octopamine**

C₁₈H₁₉NO₅ 329.352

Alkaloid from the leaves of *Aptenia cordifolia*. Oil. [α]_D²⁵ -3 (c, 0.12 in MeOH). λ_{\max} 222 (log ϵ 3.9); 294 (log ϵ 2.5); 316 (log ϵ 2.4) (MeOH).

β -Me ether, N-(4-hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyl- β -O-**

methyloctopamine

C₁₅H₂₁NO₅ 343.379

Alkaloid from *Isodon excisus*. Amorph. yellow powder. [α]_D²⁵ -2 (c, 1 in MeOH). λ_{\max} 221 (log ϵ 4.44); 294 (log ϵ 4.36); 318 (log ϵ 4.43) (MeOH).

β -Propyl ether, N-(4-hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyl- β -O-propyloctopamine**

C₂₁H₂₅NO₅ 371.432

Alkaloid from the leaves of *Aptenia cordifolia*. Oil. [α]_D²⁵ -2 (c, 0.08 in MeOH). λ_{\max} 280 (log ϵ 3.1) (MeOH).

(±)-form [1915-83-9]

Mp 156-158°.

Hydrochloride: [770-05-8]

Cryst. Mp 177-179° dec.

N-Me: see Synephrine, S-659

4-Me ether: [55275-61-1]

Cryst. (C₆H₆/petrol). Mp 70° (sealed tube).

O⁴,N-Di-Me: [125722-39-6]

[58777-87-0 (non-stereospecific), 57286-93-8 (R-form)]

C₁₀H₁₅NO₂ 181.234

Mp 106.5-107.5°.

Di-Me ether: 2-Methoxy-2-(4-methoxyphenyl)ethylamine

[31367-42-7]

C₁₀H₁₅NO₂ 181.234

Bp₂₀ 158-160°.

Di-Me ether, N-Me:

C₁₁H₁₇NO₂ 195.261

Bp₅ 129-130°.

(ξ)-form

N-(E-Cinnamoyl): N-[2-Hydroxy-2-(4-hydroxyphenyl)ethyl]-3-phenyl-2-propenamide. N-[2-Hydroxy-2-(4-hydroxyphenyl)ethyl]cinnamide. N-Cinnamoyl-2,4'-dihydroxyphenethylamine. N-Cinnamoyloctopamine

[87596-52-9]

C₁₇H₁₇NO₃ 283.326

Alkaloid from the leaves of *Aegle marmelos* (bael) (Rutaceae). Needles (MeOH). Mp 158° (solidifies and remelts at 228-231°).

N-(4-Hydroxy-E-cinnamoyl): **N-trans-p-Coumaroyloctopamine**

[66648-45-1]

[152433-78-8, 88700-34-9]

C₁₇H₁₇NO₄ 299.326

Alkaloid from roots of bell pepper (*Capsicum annuum* var. *grossum*), *Ophiopogon japonicus* and *Solanum khasianum*. Cryst. Mp 214-215°.

N-(4-Hydroxy-Z-cinnamoyl): **N-cis-p-Coumaroyloctopamine**

[180050-82-2]

C₁₇H₁₇NO₄ 299.326

Alkaloid from *Solanum khasianum*.

N-(4-Hydroxy-3-methoxy-E-cinnamoyl): **N-trans-Feruloyloctopamine**

[66648-44-0]

C₁₈H₁₉NO₅ 329.352

Constit. of *Antidesma membranaceum*, roots of bell pepper (*Capsicum annuum* var. *grossum*) and *Solanum khasianum*. Powder. Mp 164-165° (as tri-Ac).

N-(4-Hydroxy-3-methoxy-Z-cinnamoyl):

N-cis-Feruloyloctopamine

[180050-83-3]

[152433-77-7]

C₁₈H₁₉NO₅ 329.352

Constit. of *Antidesma membranaceum* and *Solanum khasianum*.

4-O-(3-Methyl-2-butenyl), N-cinnamoyl: N-[2-Hydroxy-2-[4-(3-methyl-2-butenyl)oxy]phenyl]ethyl]-3-phenyl-2-propenamide. N-2-Hydroxy-2-[4-(3,3-dimethylallyloxy)phenyl]ethylcinnamide. **Marmeline**. N-Cinnamoyl-O-propyloctopamine

[81532-00-5]

C₂₂H₂₅NO₃ 351.444

Alkaloid from the leaves and unripe fruits of *Aegle marmelos* (bael) (Rutaceae). Needles (MeOH or CHCl₃/hexane). Mp 159-161° (156-157°).

4-O-(3-Methyl-2-butenyl), β -Me ether, N-(E-cinnamoyl): *Aegle marmelos*

Alkaloid C

[70546-92-8]

C₂₃H₂₇NO₃ 365.471

Alkaloid from leaves of *Aegle marmelos* (bael) (Rutaceae). Cryst. (C₆H₆/hexane). Mp 110°.

Aldrich Library of FT-IR Spectra, 1st edn.,

1985, **1**, 1290D (ir)

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, **2**, 613B (nmr)

Quelet, R. et al., *C. R. Hebd. Seances Acad. Sci.*, 1948, **226**, 1020-1022 (di-Me ether, di-Me ether N-Me, synth)

Bretschneider, H. et al., *Monatsh. Chem.*,

1949, **80**, 530 (resoln)

Bergmann, E.D. et al., *J.O.C.*, 1951, **16**, 84-89 (O⁴,N-di-Me, synth)

Corrigan, J.R. et al., *J.A.C.S.*, 1953, **75**, 6258 (synth)

Ascher, M. et al., *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1953, **84**, 1021 (synth)

Adityachaudhury, N. et al., *J. Indian Chem. Soc.*, 1959, **36**, 585-589 (4-Me ether, synth)

Kappe, T. et al., *J. Med. Chem.*, 1964, **7**, 569-571 (resoln, mp)

Hardwick, B.C. et al., *Plant Physiol.*, 1969, **44**, 1745 (ir, uv)

Wheaton, T.A. et al., *Lloydia*, 1970, **33**, 244-254 (occur)

Kelley-Hornemann, K.M. et al., *J. Pharm. Sci.*, 1972, **61**, 41-45 (O⁴-Me, occur)

Brandau, K. et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1972, **273**, 123-133 (biosynth)

Hengstmann, J.H. et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1974, **283**, 93 (metab)

Paxton, K. et al., *Acta Cryst. B*, 1977, **33**, 2143 (cryst struct)

Axelrod, J. et al., *Nature (London)*, 1977, **265**, 501 (rev)

Smith, T.A. et al., *Phytochemistry*, 1977, **16**, 9 (occur)

Manandhar, M.D. et al., *Phytochemistry*,

1978, **17**, 1814 (*Aegle marmelos* Alkaloid C)

Doetsch, P.W. et al., *J. Chromatogr.*, 1980, **189**, 79-85 (Me ether, occur)

Yoshihara, T. et al., *Agric. Biol. Chem.*, 1981, **45**, 2593-2598 (cinnamoyl derivs)

Sharma, B.R. et al., *Phytochemistry*, 1981, **20**, 2606 (*Marmeline*)

Govindachari, T.R. et al., *Phytochemistry*,


1983, **22**, 755 (N-cinnamoyl)

Redouane, K. et al., *Agressologie*, 1984, **25**, 3 (rev)

- Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 945
- Midgley, J.M. et al., *J.C.S. Perkin 2*, 1989, 963-964 (*cryst struct, abs config, bibl*)
- Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1250
- Matsuda, F. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 625-627 (*N-Coumaroyloctopamine*)
- Lee, C. et al., *J. Nat. Prod.*, 2001, **64**, 659-660 (*N-Feruloyl-O-methyloctopamine*)
- Lee, D.G. et al., *Biotechnol. Lett.*, 2004, **26**, 1125-1130 (*N-Feruloyl-R-octopamine*)
- Lundell, K. et al., *Tetrahedron: Asymmetry*, 2004, **15**, 3723-3729 (*R-form, S-form*)
- Sadyandy, R. et al., *ARKIVOC*, 2005, **iii**, 36-43 (*R-form, synth*)
- DellaGreca, M. et al., *Tetrahedron*, 2006, **62**, 2877-2882 (*N-Feruloyl-S-octopamine*)
- Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AKT250

2-Octylpyrrole, 9CI

[158697-97-3]

C₁₂H₂₁N 179.305


No phys. props. reported.

N-Sulfo: 2-Octylpyrrole sulfamate
[587875-51-2]C₁₂H₂₁NO₃S 259.369Alkaloid from the marine annelid *Cirriformia tentaculata*. Glass (as Na salt). CAS no. refers to Na salt.Garrido, D.O.A. et al., *J.O.C.*, 1984, **49**, 2619-2622 (*synth*)Barsby, T. et al., *J. Nat. Prod.*, 2003, **66**, 1110-1112 (*sulfamate*)

Ocular age pigment A2-E

A2-E


[173449-96-2]

C₄₂H₅₈NO⁺ 592.926

Orange fluorescent pigment isol. from human eyes (>40 years).

Sakai, N. et al., *J.A.C.S.*, 1996, **118**, 1559Ren, R.X.-F. et al., *J.A.C.S.*, 1997, **119**, 3619 (*synth*)Tanaka, K. et al., *Org. Lett.*, 2000, **2**, 373-375 (*synth*)Sicre, C. et al., *Org. Lett.*, 2005, **7**, 5737-5739 (*synth*)

Odoline

Stevensine
[99102-22-4]C₁₁H₉Br₂N₅O 387.033Metab. from the New Caledonian sponge *Pseudaxinysa cantharella* and from *Ptilocaulis walpersi* and *Teichaxinella morchella*. Also isol. (as Stevensine) from an unidentified Micronesian sponge. Amorph. [α]_D +28 (c, 0.84 in MeOH).4ξ,5-Dihydro: *Hymenine*[†]

[105748-62-7]

C₁₁H₁₁Br₂N₅O 389.049Isol. from the Okinawan marine sponge *Hymeniacidon* sp. Potent α-adrenoceptor blocking agent. Amorph. solid. [α]_D²⁵ -15 (c, 0.5 in MeOH). λ_{max} 274 (ε 9200) (MeOH) (Derep).

2-Debromo: 2-Debromodoline. 2-Debromostevensine


C₁₁H₁₀BrN₅O 308.137Isol. from the sponge *Stylissa carteri*. Amorph. orange solid. λ_{max} 232 (log ε 3.84); 263 (log ε 3.62); 288 (log ε 3.37) (MeOH).

2-Debromo, 4ξ,5-dihydro: 2-Debromohymenine

C₁₁H₁₂BrN₅O 310.153Isol. from the sponge *Stylissa carteri*. Amorph. brown solid. λ_{max} 220 (log ε 3.86); 269 (log ε 3.64) (MeOH).Albizati, K.F. et al., *J.O.C.*, 1985, **50**, 4163-4164 (*isol, uv, ir, pmr, cmr*)de Nanteuil, G. et al., *Tetrahedron*, 1985, **41**, 6019-6033 (*isol, cd, uv, ir, pmr, cmr, ms*)Kobayashi, J. et al., *Experientia*, 1986, **42**, 1064-1065 (*Hymenine*)Xu, Y.-Z. et al., *J.O.C.*, 1997, **62**, 456-464 (*Hymenine, Stevensine, synth*)Eder, C. et al., *J. Nat. Prod.*, 1999, **62**, 184-187 (*2-Debromodoline, 2-Debromohymenine*)Andrade, P. et al., *Tet. Lett.*, 1999, **40**, 4775-4778 (*biosynth*)

Odorine

O-67

2-Methyl-N-[1-(1-oxo-3-phenyl-2-propenyl)-2-pyrrolidiny]butanamide, 9CI. N-Cinnamoyl-2-(2-methylbutanoylamino)-pyrrolidine. Roxburghiline
[72755-20-5]C₁₈H₂₄N₂O₂ 300.4

(+)-form

O-66

λ_{max} 283 (ε 15800) (EtOH) (Derep). λ_{max} 283 (ε 15850) (MeOH) (Berdy).

(+)-form

Alkaloid from the leaves of *Aglaia odorata*. Also isol. from *Aglaia roxburghiana* (Meliaceae). Needles (C₆H₆). Mp 218-219°. [α]_D²⁵ +72.4 (c, 0.03 in CHCl₃).2,3-Didehydro: *Dehydroodorine*C₁₈H₂₂N₂O₂ 298.384Alkaloid from leaves of *Aglaia formosana* (Meliaceae). Exhibits cytotoxicity against P-388 lymphocytic leukaemia system in cell culture. Needles (MeOH). Mp 167-168°. [α]_D¹⁸ +42.5 (c, 0.01 in CHCl₃). λ_{max} 282 (ε 66000) (MeOH) (Berdy).2-Hydroxy: (-)-*Odorinol*

[85761-62-2]

C₁₈H₂₄N₂O₃ 316.399Alkaloid from *Aglaia odorata* (Meliaceae). Mp 209-211°. [α]_D -20 (EtOH). [α]_D -35 (CHCl₃). λ_{max} 283 (ε 15800) (EtOH) (Derep).2'-Epimer: *Epidorine*

[73069-04-2]

C₁₈H₂₄N₂O₂ 300.4Alkaloid from *Aglaia odorata*, *Aglaia pyramidata* and *Aglaia roxburghiana*. Cytotoxic agent. Cryst. [α]_D +34. Called 5'-epi in the lit.

(-)-form

2-Hydroxy: (+)-*Odorinol*. *Hydroxyroxburghiline*

[72755-22-7]

C₁₈H₂₄N₂O₃ 316.399Alkaloid from *Aglaia odorata* and *Aglaia roxburghiana* (Meliaceae). Inhibitor of P-388 lymphocytic leukaemia in mice. Needles (petrol). Sol. EtOH, CHCl₃; poorly sol. H₂O. Mp 166-168°. [α]_D²⁵ +40.5 (c, 0.01 in CHCl₃). λ_{max} 283 (ε 15800) (EtOH) (Derep). λ_{max} 283 (ε 17780) (EtOH) (Berdy).

(±)-form

Synthetic. Needles (C₆H₆). Mp 187-190°.N²-Deacyl, N²-(3-methylbutanoyl):*Aglamide C*

[922149-09-5]

C₁₈H₂₄N₂O₂ 300.4Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. λ_{max} 205 (log ε 4.16); 219 (log ε 4.21); 283 (log ε 4.33) (MeOH).


2-Epimer:

Synthetic. Needles (C₆H₆). Mp 174-177°.Purushothaman, K.K. et al., *J.C.S. Perkin 1*, 1979, 3171-3174Shienhthong, D. et al., *Tet. Lett.*, 1979, 2247 (*uv, ir, pmr, ms, struct*)Babidge, P.J. et al., *Aust. J. Chem.*, 1980, **33**, 1841 (*isol, synth*)Hayashi, N. et al., *Phytochemistry*, 1982, **21**, 2371-2373 (*uv, ir, pmr, cmr, ms, cryst struct, synth*)Nagasaka, T. et al., *Heterocycles*, 1988, **27**, 2219 (*synth, ir, pmr*)Saifah, E. et al., *J. Nat. Prod.*, 1993, **56**, 473-477 (*5'-Epidorine*)

Duh, C.-Y. *et al.*, *Phytochemistry*, 1993, **34**, 857 (Dehydroodorine)
 Kim, S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1769-1775 (Aglamide C)

Oduline

[477-18-9]



$C_{17}H_{19}NO_4$ 301.341
 Alkaloid from the bulbs of *Narcissus jonquilla* hybrids “Golden sceptre” and *Narcissus odorus* var. *rugulosus* (*Narcissus jonquilla* x *Narcissus poeticus*) (Amaryllidaceae). Prisms (Me₂CO). Mp 168°. $[\alpha]_D^{25} +239$ (c, 0.35 in CHCl₃).

Picrate:

Cryst. + ½ H₂O (Me₂CO aq.). Mp 221° dec.

Me ether: O-Methyloduline

[162411-70-3]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid from bulbs of *Narcissus pseudonarcissus* ssp. *pseudonarcissus* cv. Carlton (Amaryllidaceae). Mp 164° (as picrate). $[\alpha]_D^{25} +148$ (c, 0.13 in MeOH).

7-Ketone (lactone): Masonine

[568-40-1]

$C_{17}H_{17}NO_4$ 299.326

Alkaloid from the bulbs of *Nerine masonorum*, from *Narcissus jonquilla* hybrid Golden Sceptre, and from *Galanthus nivalis* (Amaryllidaceae). Cryst. (EtOAc/Et₂O). Mp 180°. $[\alpha]_D^{25} +140$ (c, 0.2 in CHCl₃).

7-Ketone, N-de-Me: N-Demethylmasonine

[98764-82-0]

$C_{16}H_{15}NO_4$ 285.299

Alkaloid from the bulbs of *Narcissus pseudonarcissus* cv. Carlton. Mp 229° (as picrate). $[\alpha]_D^{25} +9.1$ (c, 0.4 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 725 (isol, ir)

Boit, H.G. *et al.*, *Naturwissenschaften*, 1958, **45**, 85 (Masonine, isol, ir, struct, occur)

Döpke, W. *et al.*, *Naturwissenschaften*, 1965, **52**, 60 (Masonine, occur)

Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1966, **299**, 994 (Masonine, struct, stereochem)

Döpke, W. *et al.*, *Pharmazie*, 1966, **21**, 323; *CA*, **65**, 10632d (struct)

Schnoes, H.K. *et al.*, *Tetrahedron*, 1968, **24**, 2825 (ms)

Clardy, J. *et al.*, *J.O.C.*, 1972, **37**, 49 (config)


Battersby, A. *et al.*, *J.C.S. Perkin I*, 1973, 1609 (biosynth, ir, pmr)

Kreh, M. *et al.*, *Phytochemistry*, 1995, **38**, 1533 (O-Methyloduline, N-Demethylmasonine)

Okaramine A

AK-40-I

[115444-43-4]



Absolute Configuration

$C_{32}H_{32}N_4O_3$ 520.63

Isol. from *Penicillium simplicissimum* (AK-40; ATCC90288). Mycotoxin. Insecticidal agent. Pale green needles (toluene). Sol. MeOH, EtOAc; fairly sol. toluene; poorly sol. H₂O. Mp 210-212° dec. $[\alpha]_D^{30} +101$ (c, 0.09 in MeOH). λ_{max} 229 (€ 29900); 255 (€ 21100); 284 (€ 17400); 374(€ 19400) (MeOH) (Berdy).

Ac:

$C_{34}H_{34}N_4O_4$ 562.667

Pale yellow needles (MeOH). Mp 190-193° dec. $[\alpha]_D^{22} +482$ (c, 0.075 in MeOH).

N-De(1,1-dimethylpropenyl): Okaramine I

[120717-00-2]

$C_{27}H_{24}N_4O_3$ 452.512

Prod. by *Aspergillus aculeatus* and *Penicillium simplicissimum*. Pale yellow powder. Mp 266-269°. $[\alpha]_D^{20} +645$ (c, 0.09 in MeOH). λ_{max} 233 (€ 31000); 285 (€ 16600); 371 (€ 16600) (MeOH).

Hayashi, H. *et al.*, *ABC*, 1989, **53**, 461-469; (isol, pmr, cmr, cryst struct)


Hayashi, H. *et al.*, *JNP*, 1999, **62**, 315-317 (cmr, Okaramine I)

Hewitt, P.R. *et al.*, *OBCH*, 2004, **2**, 2415-2417 (synth)

Okaramine B

AK-40-II

[117332-63-5]



Absolute Configuration

$C_{33}H_{34}N_4O_5$ 566.655

Isol. from *Penicillium simplicissimum* AK-40. Shows insecticidal activity. Toxic against *Artemia salina* (brine shrimp). Pale yellow needles (MeOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 295-298° dec. $[\alpha]_D^{20} +570$ (c, 0.24 in Py).

O-68a

λ_{max} 233 (€ 27000); 288 (€ 12900); 375 (€ 17900) (MeOH) (Berdy).

12-Hydroxy: Okaramine D

[162413-54-9]

$C_{33}H_{34}N_4O_6$ 582.655

Prod. by *Penicillium simplicissimum* AK-40; ATCC90288. Shows insecticidal props. Pale yellow needles (CHCl₃). Mp 265° dec. $[\alpha]_D^{20} +480$ (c, 0.16 in MeOH). λ_{max} 234 (€ 27500); 288 (€ 14500); 381 (€ 17200) (MeOH).

Demethoxy: Okaramine Q

[263905-01-7]

$C_{32}H_{32}N_4O_4$ 536.629

Prod. by *Penicillium simplicissimum* ATCC90288. Yellow powder. Mp 243-246°. $[\alpha]_D^{20} +516$ (c, 0.31 in MeOH). λ_{max} 234 (€ 26600); 288 (€ 13900); 376 (€ 15500) (MeOH).

Hayashi, H. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 2131-2133; 1989, **53**, 461-469 (isol, pmr, cmr, uv, ir, ms, cd, cryst struct)

Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 246-250 (Okaramine D)


Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1997, **61**, 914-916 (Okaramine B, activity)

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 103-110 (Okaramine Q)

Okaramine C

O-70

[142677-16-5]



Absolute Configuration

$C_{32}H_{36}N_4O_3$ 524.661

Prod. by *Penicillium simplicissimum* AHU8402. Insecticide. Powder. $[\alpha]_D^{25} +19$ (c, 0.2 in MeOH). λ_{max} 223 (€ 41400); 260 (€ 11100); 284 (€ 9210); 291 (€ 8690) (MeOH) (Derep).

N-Dealkyl, N-(3-methyl-2-butenyl):

Okaramine L

[244071-73-6]

$C_{32}H_{36}N_4O_3$ 524.661

Prod. by *Penicillium simplicissimum*. Powder. Mp 130-133°. $[\alpha]_D^{20} +69$ (c, 0.03 in MeOH). λ_{max} 223 (€ 48300); 284 (€ 11700); 291 (€ 11500) (MeOH).

1',2'-Didehydro: Okaramine G

[208719-70-4]

$C_{32}H_{34}N_4O_3$ 522.646

Prod. by *Penicillium simplicissimum* ATCC90288. Insecticide. Cryst. (CCl₄). Mp 145-147°. $[\alpha]_D^{20} +137$ (c, 0.1 in MeOH). λ_{max} 223 (€ 31600); 246 (€ 19500); 286 (€ 9200); 290 (€ 9300); 340 (€ 10600) (MeOH).


Hayashi, H. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 3143-3145 (isol, pmr, cmr, struct)

Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 804-806 (Okaramine G)

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1910-1920 (Okaramine L)

Okaramine E

[162413-55-0]

Absolute
ConfigurationC₃₂H₃₂N₄O₄ 536.629

Prod. by *Penicillium simplicissimum*. Pale yellow powder. $[\alpha]_D^{18} +500$ (c, 0.01 in MeOH). λ_{\max} 234 (ε 24200); 284 (ε 12600); 373 (ε 13300) (MeOH).

2,3-Didehydro- Okaramine F

[162413-56-1]


C₃₂H₃₀N₄O₄ 534.613

Prod. by *Penicillium simplicissimum*. Yellow powder. $[\alpha]_D^{18} -490$ (c, 0.01 in MeOH). λ_{\max} 230 (ε 26400); 289 (ε 14100); 402 (ε 15900) (MeOH).

Hayashi, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 246-250

Okaramine H

[220525-10-0]


Absolute
ConfigurationC₃₂H₃₂N₄O₃ 520.63

Prod. by *Aspergillus aculeatus* KF428 and *Penicillium simplicissimum*. Yellow needles (toluene). Mp 204-206°. $[\alpha]_D^{20} +559$ (c, 0.22 in MeOH). λ_{\max} 233 (ε 30300); 286 (ε 18600); 374 (ε 19400) (MeOH).

Hayashi, H. *et al.*, *J. Nat. Prod.*, 1999, **62**, 315-317 (isol, uv, ir, pmr, cmr, ms)

Okaramine J

[244071-71-4]

Absolute
Configuration

O-71

C₃₂H₃₆N₄O₃ 524.661

Prod. by *Penicillium simplicissimum*. Powder. Mp 151-153°. $[\alpha]_D^{20} +31$ (c, 0.33 in MeOH). λ_{\max} 220 (sh); 284 (ε 11800); 291 (ε 11300) (MeOH).

1',2'-Didehydro- Okaramine K

[244071-72-5]

C₃₂H₃₄N₄O₃ 522.646


Prod. by *Penicillium simplicissimum*. Powder. Mp 148-151°. $[\alpha]_D^{20} +100$ (c, 0.06 in MeOH). λ_{\max} 225 (sh); 290 (ε 9400); 342 (ε 11600) (MeOH).

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1910-1920

Roe, J.M. *et al.*, *Org. Lett.*, 2003, **5**, 2825-2827 (synth)

Okaramine M

[244071-74-7]


Absolute
ConfigurationC₂₉H₃₀N₄O₃ 482.581

Prod. by *Penicillium simplicissimum*. Powder. Mp 157-160°. $[\alpha]_D^{20} -39$ (c, 0.33 in MeOH). λ_{\max} 247 (ε 12500); 279 (ε 6900); 290 (ε 5200) (MeOH).

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1910-1920

Okaramine O

[244071-76-9]

R¹ = -C(CH₃)₂CH=CH₂, R² = HC₃₂H₃₄N₄O₄ 538.645

The abs. config. was assigned as shown on biogenetic grounds only, but is the opposite to those assigned to other closely related okaramines. Prod. by *Penicillium simplicissimum* ATCC 90288. Powder. Mp 190-193°. $[\alpha]_D^{20} +275$ (c, 0.04 in MeOH). λ_{\max} 254 (ε

10800); 284 (ε 6800); 291 (ε 6600) (MeOH).

1'-Deoxy- Okaramine N

[244071-75-8]

C₃₂H₃₄N₄O₃ 522.646

Prod. by *Penicillium simplicissimum* ATCC 90288. Powder. Mp 192-195°. $[\alpha]_D^{20} +284$ (c, 0.15 in MeOH). λ_{\max} 254 (ε 12000); 284 (ε 8200); 292 (ε 7600) (MeOH).

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 103-110 (isol)

Baran, P.S. *et al.*, *J.A.C.S.*, 2003, **125**, 5628-5629 (Okaramine N, synth)

Okaramine P

[244071-77-0]

As Okaramine O, O-75 with


R¹ = H, R² = -CH₂-CH=C(CH₃)₂C₃₂H₃₄N₄O₄ 538.645

Prod. by *Penicillium simplicissimum* ATCC 90288. Powder. Mp 188-190°. $[\alpha]_D^{20} +254$ (c, 0.22 in MeOH). λ_{\max} 284 (ε 9300); 290 (ε 8600) (MeOH).

Shiono, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 2000, **64**, 103-110

Okaramine R

[263905-02-8]


C₃₂H₃₂N₄O₄ 536.629

Prod. by *Penicillium simplicissimum* ATCC 90288. Yellow powder. Mp 185-187°. $[\alpha]_D^{20} m109$ (c, 0.02 in MeOH). λ_{\max} 232 (ε 29400); 286 (ε 16200) (MeOH).

Shiono, Y. *et al.*, *BBB.*, 2000, **64**, 103-110 (isol, ms, uv, pmr, cmr)

Oldhamine A

[1037404-79-7]

C₂₁H₂₃NO₄ 353.417

Alkaloid from the twigs of *Daphniphyllum oldhami*. Light yellow cryst. $[\alpha]_D^{20} +24.4$ (c, 0.43 in MeOH). λ_{\max} 229 (ε 1500); 295 (ε 5220); 360 (ε 2680) (MeOH).

Tan, C. *et al.*, *Tet. Lett.*, 2008, **49**, 3376-3379 (isol, pmr, cmr, cryst struct)

O-76

O-74


O-75

O-76a


O-77

Oldhamiphylline A

[848490-57-3]

Relative
ConfigurationC₂₂H₃₁NO₃ 357.492Alkaloid from *Daphniphyllum oldami*.
Amorph. solid. $[\alpha]_D^{20}$ -41 (c, 1.5 in MeOH).Chen, X. *et al.*, *Chem. Biodiversity*, 2004, **1**,
1513-1518 (*isol*, *pmr*, *cmr*)**Olefinin, 8CI**




[12764-54-4]

C₃₄H₄₇NO₉ 613.747Polyene antibiotic. Prod. by *Streptomyces parvulus*. Active against gram-positive organisms and Yoshida sarcoma. Dark red-black solid. Sol. MeOH, Et₂O; fairly sol. CCl₄; poorly sol. hexane, H₂O. Mp 96-98°. $[\alpha]_D^{30}$ -182 (c, 0.2 in EtOH). λ_{max} 262; 445; 475; 485 (MeOH/HCl) (Derep). λ_{max} 262; 415 (0.1N NaOH) (Derep). λ_{max} 262; 420 (€ 58200) (MeOH) (Derep).▶ LD₅₀ (mus, ivn) 40 mg/kg. UX9641800Gyimesi, J. *et al.*, *J. Antibiot.*, 1971, **24**,
277; 1978, **31**, 626 (*isol*, *ir*, *uv*, *pmr*, *ms*, *struct*)
Horváth, G. *et al.*, *Tet. Lett.*, 1973, 3643 (*ir*, *uv*,
ms, *nmr*, *struct*)
Kovac, L. *et al.*, *Biochim. Biophys. Acta*, 1982,
721, 349 (*props*)**Oleracins**

[57903-85-2, 57903-84-1]

Structs. unknown. Betacyanin pigments from *Portulaca oleracea* (purslane) (Portulacaceae) stems. Oleracins I and II recognised.Piatelli, M. *et al.*, *Phytochemistry*, 1964, **3**,
547-557 (*occur*)Imperato, F. *et al.*, *Phytochemistry*, 1975, **14**,
2091-2092 (*Oleracins I,II*)**Oliganone**

[85967-03-9]

**O-81**C₂₀H₂₄N₂O₃ 340.421Alkaloid from *Crinum oliganthum*
(Amaryllidaceae).Döpke, W. *et al.*, *Z. Chem.*, 1983, **23**, 101 (*isol*,
ir, *pmr*, *ms*, *struct*)**Oligophylline**3,5-Dihydro-2,3,3,5-tetramethylfuro[3,2-c]quinolin-4(2H)-one, 9CI
[86702-40-1]**O-82**C₁₅H₁₇NO₂ 243.305Alkaloid from the roots of *Acronychia oligophylebia* and the heartwood of *Euxylophora paraënsis* (Rutaceae). Needles. Mp 90°. Opt. inactive. λ_{max} 218 (€ 46700); 231 (€ 48800); 282 (€ 7850); 292 (€ 9200); 317 (€ 8150); 330 (€ 6600) (MeOH) (Berdy).N-De-Me: 2,3,3-Trimethyl-2,3,4,5-tetrahydrofuran[3,2-c]quinolin-4-one. N-Demethyloligophylline
[86702-39-8]C₁₄H₁₅NO₂ 229.278Trace alkaloid from the heartwood of *Euxylophora paraënsis* (Rutaceae). Glistening needles (MeOH). Mp 200-201°.6-Methoxy, N-de-Me.: 3,5-Dihydro-6-methoxy-2,3,3-trimethylfuro[3,2-c]quinolin-4(2H)-one, 9CI. Oligophylline
[94482-45-8]C₁₅H₁₇NO₃ 259.304Alkaloid from the roots of *Acronychia oligophylebia* (Rutaceae). λ_{max} 213 (€ 36200); 236 (€ 36200); 282 (€ 10200); 292 (€ 11000); 320 (€ 4400); 333 (€ 3000) (MeOH) (Berdy).6,7-Dimethoxy, N-de-Me.: 3,5-Dihydro-6,7-dimethoxy-2,3,3-trimethylfuro[3,2-c]quinolin-4(2H)-one. Oligophyllidine
[94443-42-2]C₁₆H₁₉NO₄ 289.33Alkaloid from the roots of *Acronychia oligophylebia* (Rutaceae).Jurd, L. *et al.*, *Aust. J. Chem.*, 1983, **36**, 759
(*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*, *deriv*)Xu, W. *et al.*, *Huaxue Xuebao*, 1984, **42**, 899;
CA, **102**, 59257s**Olivacine**1,5-Dimethyl-6H-pyrido[4,3-b]carbazole, 9CI. Guatambuine
[484-49-1]**O-83**C₁₇H₁₄N₂ 246.311Alkaloid from several *Aspidosperma* spp., from *Tabernaemontana psychotrifolia*, *Ochrosiaspp.* and some other spp. in Apocynaceae. Antineoplastic agent. Shows v. strong cytotoxic activity vs. human carcinoma KB cells and L1210 leukaemia. Also shows antiprotozoal activity. Mp 318-326° dec. (315-320°). Log P 4.37 (calc). λ_{max} 224 (€ 30200); 237 (sh) (€ 26900); 245 (sh) (€ 21900); 275 (€ 57500); 285 (€ 75900); 293 (€ 70800); 331 (€ 7080); 346 (€ 5010); 380 (€ 3800); 400 (€ 3800) (EtOH) (Derep). λ_{max} 224; 238; 267; 276; 286; 292; 314; 318; 343 (EtOH) (Berdy). λ_{max} 242; 300; 348 (EtOH-HCl) (Berdy).

▶ UU8824900

N-Oxide: Olivacine N-oxide

[2122-22-7]

C₁₇H₁₄N₂O 262.31Alkaloid from bark of *Aspidosperma nigricans* (Apocynaceae). Shows antineoplastic activity vs. walker 256 carcinosarcoma cells. Mp 304-305° dec. Log P 2.28 (calc).

9-Methoxy: 9-Methoxyolivacine

[16101-08-9]

C₁₈H₁₆N₂O 276.337Alkaloid from the bark of *Aspidosperma vargasii* (Apocynaceae). Pale yellow cryst. (MeOH). Mp 291-293° dec. Probable struct.



9-Methoxy; methiodide: Mp 340° dec.

1,2,3,4-Tetrahydro: see Guatambuine, G-211

Marini-Bettolo, G.B. *et al.*, *Helv. Chim. Acta*, 1959, **42**, 2146 (*uv*, *ir*, *struct*)Schmutz, J. *et al.*, *Helv. Chim. Acta*, 1960, **43**, 793 (*synth*, *ir*)Wenkert, E. *et al.*, *J.A.C.S.*, 1962, **84**, 94
(*synth*)Gilbert, B. *et al.*, *Tetrahedron*, 1965, **21**, 1141
(*oxide*)Burnell, R.H. *et al.*, *Can. J. Chem.*, 1967, **45**, 89 (9-Methoxyolivacine)Oikawa, Y. *et al.*, *J.C.S. Perkin I*, 1976, 1479
(*synth*)Ahond, A. *et al.*, *Tetrahedron*, 1978, **34**, 2385
(*cmr*)Kano, S. *et al.*, *Heterocycles*, 1981, **19**, 1673
(*synth*)Besselièvre, R. *et al.*, *Tetrahedron, Suppl.*, No. 1, 1981, 241 (*synth*)Kutney, J.P. *et al.*, *Can. J. Chem.*, 1982, **60**, 2426 (*synth*)Murakami, Y. *et al.*, *Tet. Lett.*, 1983, **24**, 2189
(*synth*)Wanner, M.J. *et al.*, *Tetrahedron*, 1983, **39**, 3673 (*synth*)Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 89 (*rev*, *antitumour props*)
Narasimhan, N.S. *et al.*, *Chem. Comm.*, 1985, 86 (*synth*)Danieli, B. *et al.*, *Alkaloids (Academic Press)*, 1986, **27**, 120 (*pharmacol*)Naito, T. *et al.*, *J.C.S. Perkin I*, 1986, 99
(*synth*, *ir*, *pmr*)Hibino, S. *et al.*, *J. Het. Chem.*, 1990, **27**, 1751
(*synth*)Yokoyama, Y. *et al.*, *J.C.S. Perkin I*, 1990, 1319 (*synth*)Bäckvall, J.-E. *et al.*, *J.O.C.*, 1990, **55**, 4528
(*synth*)Gribble, G.W. *et al.*, *J.O.C.*, 1992, **57**, 5891
(*synth*)Tilve, S.G. *et al.*, *Synth. Commun.*, 1996, **26**, 1921 (*synth*)

Oliveramine

[14645-64-1]

C₂₀H₂₀N₂O₄ 352.389Alkaloid from *Gentiana olivieri* (Gentianaceae). Mp 144-145°. λ_{max} 273 (log ε 3.48) (EtOH).Rakhmatullaev, T.U. et al., *Khim. Prir. Soedin.*, 1973, 9, 64-67; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, 9, 56-58 (ir, uv, pmr, ms, isol, struct)**Oliveridine†**7-Hydroxy-1,2-methylenedioxy-9-methoxyaporphine
[53602-02-1]Absolute
Configuration

R = OMe

C₁₉H₁₉NO₄ 325.363Alkaloid from the trunk bark and leaves of *Polyalthia oliveri* and the trunk bark of *Polyalthia suaveolens*; also isol. from *Isolona campanulata* (Annonaceae). Causes transient hypotension followed by hypertension in rats. Needles (MeOH). Mp 95-100°. [α]_D -28 (c, 0.9 in CHCl₃).*Hydrochloride*:Cryst. (MeOH/Et₂O). Mp 251-253° dec.*N-Oxide: Oliveridine N-oxide*

[62173-00-6]

C₁₉H₁₉NO₅ 341.363Alkaloid from the stem bark of *Enantia pilosa* and *Greenwayodendron suaveolens* (*Polyalthia suaveolens*) (Annonaceae). Mp 208-209°. [α]_D -51 (c, 0.6 in EtOH).*Ac*: Mp 121-122°. [α]_D +26 (c, 0.5 in CHCl₃).*N-De-Me: Noroliveridine*

[64234-39-5]

C₁₈H₁₇NO₄ 311.337Alkaloid from the leaves of *Polyalthia oliveri* (Annonaceae). Air- and light-sensitive amorph. powder.*Me ether: Oliverine†*

[53602-03-2]

C₂₀H₂₁NO₄ 339.39Alkaloid from trunk and stem barks of *Polyalthia suaveolens* and trunk bark and leaves of *Polyalthia oliveri*; also from *Isolona campanulata* (Annonaceae) and *Enantia pilosa*. Effective antiinflammatory agent. Noncryst.; cryst.(MeOH/Et₂O) (as

O-84

hydrochloride). Mp 250° dec. (hydrochloride). [α]_D +70 (c, 0.7 in EtOH). λ_{max} 222 (log ε 4.22); 284 (log ε 4.2) (EtOH).*Me ether, N-oxide: Oliverine N-oxide*

[62172-99-0]

C₂₀H₂₁NO₅ 355.39Alkaloid from the stem and root barks of *Enantia pilosa*, the stem bark of *Polyalthia suaveolens* and the bark of *Isolona campanulata* (Annonaceae). Fine beige needles (CH₂Cl₂). Mp 134° (110-114°). [α]_D +111 (c, 0.9 in EtOH).*Me ether, N-de-Me: Noroliverine*

[67627-74-1]

C₁₉H₁₉NO₄ 325.363Alkaloid from the trunk bark of *Polyalthia suaveolens* (Annonaceae). Noncryst.; cryst. (MeOH/Et₂O) (as hydrochloride). Mp 280-282° dec. (hydrochloride). [α]_D +65 (c, 0.5 in EtOH).*O-De-Me: Roemerolidine*

[112494-61-8]

C₁₈H₁₇NO₄ 311.337Alkaloid from *Duguetia spixiana* (Annonaceae). [α]_D -23 (c, 0.24 in EtOH).*O-De-Me, O⁷-Me: Polysuavine*

[67627-72-9]

C₁₉H₁₉NO₄ 325.363Alkaloid from the trunk bark and stem bark of *Polyalthia suaveolens* (Annonaceae). Noncryst.*7-Epimer, N-de-Me: Michelanugine*


[51059-63-3]

C₁₈H₁₇NO₄ 311.337Alkaloid from the trunk bark of *Michelia lanuginosa*. Amorph.; cryst. (as hydrochloride). Mp 275° dec. (hydrochloride). [α]_D -105 (c, 0.62 in EtOH).Hamonnière, M. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, 278, 921-924 (uv, pmr, ms, abs config)Talapatra, S.K. et al., *Tetrahedron*, 1975, 31, 1105-1107 (*Michelanugine*)Nieto, M. et al., *J. Nat. Prod.*, 1976, 39, 350-356 (*Oliverine N-oxide, Oliveridine N-oxide*)Hamonnière, M. et al., *Phytochemistry*, 1977, 16, 1029-1034 (isol, pmr, *Oliverine, Noroliveridine*)Hocquemiller, R. et al., *Plant. Med. Phytother.*, 1978, 12, 230-234 (isol, *Oliveridine, Oliverine, Oliverine oxide*)Cavé, A. et al., *Planta Med.*, 1978, 33, 243-250 (*Noroliverine, Polysuavine*)Jackman, L.M. et al., *J. Nat. Prod.*, 1979, 42, 437-449 (cmr)Kessar, S.V. et al., *Indian J. Chem., Sect. B*, 1981, 20, 984; 1983, 22, 321-324 (synth, uv, pmr, ms)Hasan, C.M. et al., *J.C.S. Perkin I*, 1982, 2807-2812 (oxides)Rasamizafy, S. et al., *J. Nat. Prod.*, 1987, 50, 674-679 (*Roemerolidine*)**Oliverine†**

O-86

C₂₀H₁₆N₂O₄ 348.357Monoterpene alkaloid. Struct. unknown. Alkaloid from the epigeous part of *Gentiana olivieri* (Gentianaceae). Cryst. (MeOH). Mp 206-207°.Rakhmatullaev, T.U. et al., *Khim. Prir.**Soedin.*, 1969, 5, 608; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, 5, 531-532 (isol)Rakhmatullaev, T.U. et al., *Khim. Prir. Soedin.*, 1972, 350-353; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, 8, 344-345 (isol)**Oliveroline**

O-87

7-Hydroxy-1,2-methylenedioxyaporphine
[62560-99-0]Absolute
ConfigurationC₁₈H₁₇NO₃ 295.337Alkaloid from the trunk bark and leaves of *Polyalthia oliveri*, the trunk bark of *Polyalthia suaveolens*, and the stem and root barks of *Pachypodanthium confine* (Annonaceae). Shows antiparkinsonian activity. Causes transient hypotension followed by hypertension in rats. Cryst. (MeOH). Mp 152°. [α]_D -64 (c, 0.5 in CHCl₃).*β-N-Oxide: Oliveroline β-N-oxide*

[64190-95-0]

C₁₈H₁₇NO₄ 311.337Alkaloid from the leaves of *Polyalthia oliveri* and from *Polyalthia longifolia* (Annonaceae). Also isol. from *Guatteria sagotiana* (Annonaceae). Microcryst. powder (MeOH). Mp 138°. [α]_D -154 (c, 0.4 in EtOH). Stereochem. at *N*-oxide centre unequivocally detd. in 1989.*N-De-Me: 7-Hydroxy-1,2-methylenedioxyaporphine. Noroliveroline*

[83730-14-7]

C₁₇H₁₅NO₃ 281.31Alkaloid from the bark and leaves of *Polyalthia acuminata* (Annonaceae). Amorph.*N-De-Me, N,O-di-Ac: N,O-Diacetylnoroliveroline*

[83730-15-8]

C₂₁H₁₉NO₅ 365.385Alkaloid from the rhizomes of *Lysichiton camtschaticense*. Light brown powder. [α]_D²⁵ -221.3 (c, 0.18 in CHCl₃). λ_{max} 211 (log ε 4.36); 276 (log ε 3.98); 319 (log ε 3.44) (EtOH).*Me ether: 7-Methoxy-1,2-methylenedioxyaporphine. N-Methylpachypodanthine*

[60033-07-0]

C₁₉H₁₉NO₃ 309.364Alkaloid from the bark of *Polyalthia staudtii* (Annonaceae). Mp 250° dec. (as hydrochloride). [α]_D +24 (c, 0.8 in EtOH).*Me ether, N-oxide: N-Methylpachypodanthine N-oxide*

[64190-96-1]

C₁₉H₁₉NO₄ 325.363Alkaloid from the leaves and stem bark of *Polyalthia oliveri* (Annonaceae). Noncryst.*Me ether, N-de-Me: 7-Methoxy-1,2-methylenedioxyaporphine. Pachypodanthine*

[60033-05-8]

C₁₈H₁₇NO₃ 295.337

Alkaloid from the trunk bark and root bark of *Pachypodanthium staudtii*, the leaves of *Polyalthia oliveri* and the trunk bark of *Polyalthia suaveolens* (Annonaceae). Needles (Et₂O). Mp 127°. [α]_D +38 (c, 0.8 in EtOH).

Me ether, N-de-Me, hydrochloride: Mp 255° dec.

Me ether, N-de-Me, N-Ac: N-Acetyl-pachypodanthine

[60033-06-9]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the bark of *Pachypodanthium confine*. Needles (MeOH) (synthetic); yellowish powder (natural). Mp 212-213° (synthetic). [α]_D -204 (c, 0.8 in EtOH) (synthetic). [α]_D²⁰ -64.3 (c, 0.28 in EtOH) (natural). λ_{max} 277 (log ε 3.92) (MeOH) (natural).

Bévalot, F. et al., C. R. Hebd. Seances Acad. Sci. Ser. C, 1976, **282**, 865 (*Pachypodanthine*, uv, pmr, ms, struct)

Bévalot, F. et al., Ann. Pharm. Fr., 1977, **35**, 65; CA, **86**, 185953e (isol)

Hamonnière, M. et al., Phytochemistry, 1977, **16**, 1029 (*Oliveroline β-N-oxide*, N-Methylpachypodanthine oxide)

Bévalot, F. et al., Plant. Med. Phytother., 1977, **11**, 315 (*N-Methylpachypodanthine*)

Cavé, A. et al., Planta Med., 1978, **33**, 243 (isol)

Kessar, S.V. et al., Tet. Lett., 1980, **21**, 3307 (*Oliveroline*, *Noroliveroline*, synth)

Zarga, M.H.A. et al., J. Nat. Prod., 1982, **45**, 471 (*Noroliveroline*)

Rasamizafy, S. et al., J. Nat. Prod., 1986, **49**, 1078 (*oxide*)

Wu, Y.-C. et al., Heterocycles, 1989, **29**, 463 (*config*, *oxide*)

Takatsu, H. et al., J. Nat. Prod., 2005, **68**, 430-431 (*N,O-Diacetylnoroliveroline*)


Brastianos, H.C. et al., J. Nat. Prod., 2007, **70**, 287-288 (*activity*)

Mathouet, H. et al., Phytochemistry, 2007, **68**, 1813-1818 (*N-Acetyl-pachypodanthine*)

Olivoretin C

O-88

[90599-28-3]

C₂₉H₄₃N₅O₂ 465.678

Isol. from *Streptovorticillium olivoreticuli* and *Streptovorticillium blastmyceticus*. Possesses strong vesicatory activity and possible tumour-promoting props. Prisms (EtOH). Sol. MeOH, CHCl₃. Mp 305-307°. [α]_D³² -256.6 (c, 0.07 in CHCl₃). λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep). λ_{max} 233 (ε 36300); 289 (ε 9550) (MeOH) (Berdy).

O-De-Me: Des-O-methylolivoretin C

[95013-42-6]

C₂₈H₄₁N₅O₂ 451.651

Minor metab. of *Streptovorticillium olivoreticuli*. Plates (C₆H₆). Mp 268-270°. λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).


Aimi, N. et al., Chem. Pharm. Bull., 1984, **32**, 354; 358; 3774; 4233 (*isol*, pmr, cmr, struct)

Irie, K. et al., Agric. Biol. Chem., 1985, **49**, 221 (*isol*)

Olivoretin E

O-89

[110187-21-8]

C₂₉H₄₃N₅O₂ 465.678

Prod. by *Streptovorticillium olivoreticuli*. Tumour promotor. Prisms. Mp 266-269° dec. Similar to Teleocidin B₁, T-63. λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).


Sakai, S. et al., Chem. Pharm. Bull., 1986, **34**, 4883 (*isol*, struct)

Irie, K. et al., Tetrahedron, 1990, **46**, 2773 (*biosynth*)

Olomoucine

O-90

2-[[9-Methyl-6-[(phenylmethyl)amino]-9H-purin-2-yl]amino]ethanol, 9CI. 6-(Benzylamino)-2-(2-hydroxyethylamino)-9-methylpurine [101622-51-9]

C₁₅H₁₈N₆O 298.347

Constit. of the cotyledons of radish *Raphanus sativus*. Cytokinin. Cyclin-dependent kinase inhibitor. Cryst. (MeOH/EtOAc/petrol). Mp 126°. λ_{max} 232 (ε 23260); 290 (ε 12880) (EtOH aq.).

Parker, C.W. et al., Phytochemistry, 1986, **25**, 303-310 (*synth*)

Tao, G.Q. et al., J. Plant Growth Regul., 1991, **10**, 179-185 (*isol*)

Schulze-Gahmen, U. et al., Proteins, 1995, **22**, 378-391 (*cryst struct with ATP*)

Havlicek, L. et al., J. Med. Chem., 1997, **40**, 408-412 (*synth*, *activity*)

Nugiel, D.A. et al., J.O.C., 1997, **62**, 201-203 (*synth*)

Krystof, V. et al., Cell Mol. Life Sci., 2005, **62**, 1763-1771 (*pharmacol*)


Wan, Z.-K. et al., Org. Lett., 2006, **8**, 2425-2428 (*synth*)

Tian, D.S. et al., Brain Res., 2007, **1135**, 177-185; **1154**, 206-214 (*pharmacol*)

Oncinotine

O-91

4-(4-Aminobutyl)octadecahydropyridido[1,2-e][1,5]diazacycloheptadecine-5(6H)-one, 9CI

C₂₃H₄₅N₃O 379.628**(R)-form** [21008-79-7]

Alkaloid from the stem bark of *Oncinotis nitida* (Apocynaceae). Oil. [α]_D -33 (c, 0.106 in MeOH). [α]_D -29 (CHCl₃).

11-Oxo: Oncinotin-11-one

[181486-83-9]

C₂₃H₄₃N₃O₂ 393.612

Minor alkaloid from leaves of *Oncinotis tenuiloba*. Has (S)-config.

12-Oxo: Oncinotin-12-one

[181486-84-0]

C₂₃H₄₃N₃O₂ 393.612

From leaves of *Oncinotis tenuiloba*. Has (S)-config.

(±)-form [53648-98-9]

Synthetic. Noncryst.

Badawi, M.M. et al., Helv. Chim. Acta, 1968, **51**, 1813 (*struct*)

Guggisberg, A. et al., Helv. Chim. Acta, 1974, **57**, 414; 1976, **59**, 3013 (*isol*, uv, ord, ir, ms, struct, synth, abs config)

Schneider, F. et al., Helv. Chim. Acta, 1974, **57**, 434 (*synth*, ir, ms)

Bienz, S. et al., Helv. Chim. Acta, 1988, **71**, 1708 (*synth*)

Doll, M.K.-H. et al., Helv. Chim. Acta, 1996, **79**, 973; 1379 (*Oncinotin-11-one*, *Oncinotin-12-one*)

Ina, H. et al., J.O.C., 1996, **61**, 1023 (*synth*, pmr, cmr)


Hou, D.-R. et al., J.O.C., 2004, **69**, 6094-6099 (*synth*)

Onetine

O-92

15,20-Dihydro-12,15,20-trihydroxy-4-methyl-4,8-secosenecionan-8,11,16-trione, 9CI

[41451-67-6]



Absolute configuration

C₁₉H₂₉NO₈ 399.44

Cyclic otonecine diester. Closely related to the Jacoline group. Alkaloid from *Senecio othonnae* (Asteraceae). Cryst. (EtOH). Mp 192-193°. [α]_D²⁰ +73 (c, 2.5 in CHCl₃).

Flavianate:

Cryst. (EtOH). Mp 238-240°.

O¹²-Ac: Floridanine

[16958-31-9]

C₂₁H₃₁NO₉ 441.477

Alkaloid from *Senecio erraticus*, *Senecio orthomnae*, *Senecio aureus* and *Cacalia floridana* (Asteraceae). Prisms + CHCl₃, Mp 195-196°. [α]_D +66.5 (c, 0.8 in CHCl₃).

▶ VS3595000

O¹², O²⁰, Di-Ac: Floraline

[16958-32-0]

C₂₃H₃₃NO₁₀ 483.514

Alkaloid from *Cacalia floridana* (Asteraceae). Solved prisms (C₆H₆). Mp 120-122° Mp 177-178° (dried).

20-Deoxy, 20-chloro: Desacetyldoronine

[120481-77-8]

C₁₉H₂₈ClNO₇ 417.885

Alkaloid from *Senecio inaequidens* (Asteraceae).

20-Deoxy, 20-chloro, O¹²-Ac: Doronine**6'-Chlorodeoxyfloridanine**

[60367-00-2]

C₂₁H₃₀ClNO₈ 459.923

Alkaloid from *Doronicum macrophyllum* roots and from *Senecio abrotanifolius* and *Senecio clevelandii* (Asteraceae). Cryst. (C₆H₆/cyclohexane). Mp 113-114° dec. [α]_D +45.4 (c, 1.1 in CHCl₃).

▶ VS3585000

20-Deoxy, 20-chloro, O¹²-Ac, picrate: Mp 235° dec.

[30512-07-3, 7079-67-6, 30484-73-2, 7097-02-1]

Daniilova, A.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1962, **32**, 638 (isol)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 3570 (*Floridanine, Floraline*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (*cd, stereochem*)

Alieva, Sh.A. *et al.*, *Khim. Prir. Soedin.*, 1976, **194**; *Chem. Nat. Compd. (Engl. Transl.)*, 173 (*Doronine*)


Roeder, E. *et al.*, *Planta Med.*, 1983, **49**, 57 (*isol, ms, pmr, cmr, Floridanine*)

Wong, R.Y. *et al.*, *Acta Cryst. C*, 1984, **40**, 163 (*cryst struct, Doronine*)

Bicchi, C. *et al.*, *J. Nat. Prod.*, 1989, **52**, 32 (*occur, ms, Desacetyldoronine*)

Onychine†**O-93**

4-Methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 1-Methyl-4-azafluorenone
[58787-04-5]

C₁₃H₉NO 195.22

Alkaloid from the trunkwood of *Onychopetalum amazonicum* and *Guatteria dielsiana*, and from the root bark of *Cleistopholis patens* (Annonaceae). Shows antiyeast activity. Pale yellow needles (EtOAc/petrol). Mp 136-138° (133-135°, 125-129°). Struct. firmly established by 1979. The erroneous for-

mula, 4-methyl-1-azafluorenone, was retained several years later when Onychine was reisolated and the structs. of several other more oxygenated azafluorenones were based on it. See also Dielsinol, D-385 and 8,9-Dihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, D-626. λ_{max} 253 (log ε 4.62); 279 (log ε 3.85); 289 (log ε 3.88); 308 (log ε 3.3) (EtOH).

3-Hydroxy: 3-Hydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 2-Hydroxy-1-methyl-4-azafluorenone. 2-Hydroxyonychine

[942472-64-2]

C₁₃H₉NO₂ 211.22Alkaloid from *Polyalthia nemoralis*.

Amorph. yellow powder. λ_{max} 214 (log ε 3.99); 249 (log ε 4.1); 269 (log ε 4.12) (MeOH).

8-Hydroxy: 8-Hydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 6-Hydroxy-1-methyl-4-azafluorenone. 6-Hydroxyonychine. Oxylopinine

C₁₃H₉NO₂ 211.22

Alkaloid from *Oxandra xylopioides* (Annonaceae) and *Unonopsis spectabilis* (Annonaceae). Yellow needles. Mp 243-245°.

8-Methoxy: 8-Methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 6-Methoxy-1-methyl-4-azafluorenone. 6-Methoxyonychine

[105418-67-5]

C₁₄H₁₁NO₂ 225.246

Alkaloid from the trunkwood of *Guatteria dielsiana*. Cryst. (EtOH) or gum. Mp 95-98° Mp 130-132°. Struct. revised in 1987; erroneously described as 6-methoxy-4-methyl-1-azafluorenone in the original lit. λ_{max} 252 (log ε 4.22); 280 (log ε 3.79); 292 (log ε 3.75); 312 (sh) (log ε 3.43) (EtOH).

Bowden, B.F. *et al.*, *Aust. J. Chem.*, 1975, **28**, 2681-2701 (*synth, uv, ir, pmr, ms*)

De Almeida, M.E.L. *et al.*, *Phytochemistry*, 1976, **15**, 1186-1187 (*isol, uv, ir, pmr, ms*)

Koyama, J. *et al.*, *Heterocycles*, 1979, **12**, 1017-1019 (*synth, pmr*)

Waterman, P.G. *et al.*, *Phytochemistry*, 1985, **24**, 523-527 (*isol, pmr, cmr*)

Goulart, M.O.F. *et al.*, *Phytochemistry*, 1986, **25**, 1691-1695 (*6-Methoxyonychine, isol, uv, ir, pmr, ms*)

Zhang, J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 800-806 (*6-Hydroxyonychine, isol, synth*)

Hufford, C.D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 961-964 (*activity*)

Tadić, D. *et al.*, *Phytochemistry*, 1987, **26**, 1551-1552 (*6-Methoxyonychine, struct, synth, uv, ir, pmr, ms*)

Irie, H. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 3134-3137 (*synth, 6-Methoxyonychine*)

Alves, T. *et al.*, *Tet. Lett.*, 1988, **29**, 2135-2136 (*Onychine, 6-Methoxyonychine, synth*)

Cassels, B.K. *et al.*, *J. Nat. Prod.*, 1989, **52**, 420-422 (*cmr*)

Nitta, M. *et al.*, *J.C.S. Perkin 1*, 1991, 1115-1118 (*synth*)

Bracher, F. *et al.*, *Synlett*, 1991, 95-96 (*synth, 6-Methoxyonychine*)

Tong, T.H. *et al.*, *Synth. Commun.*, 1992, **22**, 1773-1782 (*synth*)


Bracher, F. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1994, **327**, 371-375 (*synth, 6-Methoxyonychine*)

Padwa, A. *et al.*, *J.O.C.*, 2000, **65**, 2368-2378 (*synth, ir, pmr*)

He, X.-F. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 783-791 (*2-Hydroxyonychine*)

Ophiocarpine**O-94**

5,8,13,13a-Tetrahydro-9,10-dimethoxy-6H-benzo[*g*]-1,3-benzodioxolo[5,6-*a*]quinolizin-13-ol, 9CI. 9,10-Dimethoxy-2,3-(methylenedioxy)berbin-13-ol, 8CI. 13-Hydroxycanadine



(-)-form

C₂₀H₂₁NO₅ 355.39**(-)-form [478-13-7]**

Alkaloid from *Corydalis cheilanthifolia*, *Corydalis ophiocarpa*, *Corydalis govani* and *Corydalis campulicarpa* (Papaveraceae). Shows cytotoxic and antimicrobial props. Prisms (MeOH). Mp 188°. [α]_D²⁴ -283 (c, 1 in CHCl₃).

▶ DR9900000

β-N-Oxide: Carpoxydine. Ophiocarpine N-oxide

[66408-19-3]

C₂₀H₂₁NO₆ 371.389

Alkaloid from *Corydalis ophiocarpa* (Papaveraceae). Cryst. (MeOH). Mp 213-215° (207-208°). [α]_D -110 (c, 0.5 in CHCl₃). [α]_D²⁵ -185 (c, 0.77 in C₆H₆/MeOH 2:1).

N-Me: Mp 272-273° dec. (as iodide). [α]_D -166 (c, 0.50 in EtOH) (as iodide).

Ac:

Cryst. (CH₂Cl₂/petrol). Mp 141-143° Mp 165-167°.

13-Ketone: Ophiocarpinone

[83532-42-7]

C₂₀H₁₉NO₅ 353.374

Minor alkaloid from *Cocculus pendulus* (Menispermaceae). Cryst. (MeOH). Mp 217-218°. [α]_D +265 (c, 1.0 in CHCl₃).

13-Epimer: Epiophiocarpine

[5308-61-2]

Synthetic. Mp 161-162°. [α]_D -282 (c, 1.0 in CHCl₃).

(±)-form [18090-55-6]

Mp 248-252°.

Ac: Mp 176-177°.**13-Ketone: [83572-34-3]**

Mp 232-234°.

13-Epimer:

Cryst. (CHCl₃/MeOH). Mp 184-185° (178-179°).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1939, **17**, 51 (*isol*)

Govindachari, T.R. *et al.*, *J.C.S.*, 1957, 557 (*synth*)


Ohta, M. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1072 (*ir, pmr, abs config, synth*)

- Elliott, I.W. *et al.*, *J. Het. Chem.*, 1967, **4**, 639 (synth)
 Jeffs, P.W. *et al.*, *J.O.C.*, 1975, **40**, 644 (ord)
 Taira, Z. *et al.*, *Cryst. Struct. Commun.*, 1978, **7**, 71 (cryst struct, oxide)
 Tani, C. *et al.*, *Yakugaku Zasshi*, 1978, **98**, 1243 (isol, oxide)
 Preininger, V. *et al.*, *Planta Med.*, 1979, **36**, 213 (isol, oxide)
 Bhakuni, D.S. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 389 (Ophiocarpinone, isol, ir, pmr, cd, abs config, synth)
 Iwasa, K. *et al.*, *J.O.C.*, 1982, **47**, 4275 (cmr, pmr, conformn)

Ophiorine A

O-95

[99615-91-5]



$C_{26}H_{28}N_2O_9$ 512.515
 Alkaloid from the leaves of *Ophiorrhiza japonica* and *Ophiorrhiza kuroi* and from the bark of *Sickingia tinctoria* and *Sickingia williamsii* (preferred genus name *Simira*) (Rubiaceae). Pale-yellow needles. Mp 217-219°. $[\alpha]_D^{25} +51$. Unique among the naturally occurring glucoindole alkaloids in possessing a $C_{17}-N_b$ linkage.

16-Epimer: Ophiorine B

[99631-26-2]

 $C_{26}H_{28}N_2O_9$ 512.515

Alkaloid from the leaves of *Ophiorrhiza japonica* and *Ophiorrhiza kuroi* (Rubiaceae). Pale-yellow needles. Mp 188-191°. $[\alpha]_D^{25} +18.2$.


Aimi, N. *et al.*, *Tet. Lett.*, 1985, **26**, 5299 (isol, uv, pmr, struct)

Aquino, R. *et al.*, *Phytochemistry*, 1994, **37**, 1471 (isol, cmr)

Ophiorrhizine

O-96

11-Hydroxy-4,17-cyclocorynanium(1+)
 [142750-49-0]

 $C_{19}H_{25}N_2O^+$ 297.419

Erroneously named 10-Hydroxy-4,17-cyclocorynanium in CAS. Alkaloid from aerial parts of *Ophiorrhiza major* (Rubiaceae). Plates + 1H₂O (MeOH/EtOAc) (as chloride). Mp 162-164° dec. (chloride). $[\alpha]_D^{20} -60$ (c, 0.001 in MeOH). CAS no. refers to chloride.


Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1992, 663-664 (isol, uv, pmr, cmr, cryst struct)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 49-52 (synth, cd, abs config)

Ophiorrhizine-12-carboxylate

O-97

[214403-96-0]

 $C_{20}H_{24}N_2O_3$ 340.421


Alkaloid from *Ophiorrhiza blumeana*. Prisms + 2H₂O (MeOH/EtOAc). Mp 278-281° dec. $[\alpha]_D^{27} -52$ (c, 0.001 in MeOH). λ_{max} 229 (ε 10800); 291 (ε 2900); 320 (ε 2900) (MeOH).

Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1998, 2537-2540 (isol, uv, ir, cd, pmr, cmr, ms, cryst struct)

Ophiuroidine

O-98

4,8,9-Trihydroxyindolo[2,1-b]quinazoline-6,12-dione
 [944822-48-4]

 $C_{15}H_8N_2O_5$ 296.239


Isol. from *Ophiocoma riisei*. Orange powder. Mp > 300° dec. λ_{max} 224 (log ε 4.4); 254 (log ε 4.2); 284 (log ε 4.2); 312 (log ε 4.05); 320 (log ε 3.77); 399 (log ε 3.96) (EtOH). λ_{max} 240 (log ε 4.48); 320 (log ε 4.2); 500 (log ε 4.18) (EtOH/KOH).

Utkina, N.K. *et al.*, *Tet. Lett.*, 2007, **48**, 4445-4447 (isol, uv, pmr, cmr, ms)

OPHR 17

O-99

[486394-19-8]



Absolute Configuration

 $C_{27}H_{28}N_2O_9$ 524.526

Alkaloid from hairy root cultures of *Ophiorrhiza pumila*. Amorph. λ_{max} 219; 254; 280 (sh); 363 (MeOH).

17-Epimer: OPHR 23

[486394-18-7]

 $C_{27}H_{28}N_2O_9$ 524.526

Alkaloid from hairy root cultures of *Ophiorrhiza pumila*. Amorph. λ_{max} 219; 254; 290; 362


(MeOH).

Kitajima, M. *et al.*, *Tetrahedron*, 2002, **58**, 9169-9178 (isol, uv, cd, pmr, cmr)

Opuntine B

O-100

3-[(4-Hydroxyphenyl)methyl]-1-phenyl-1H-pyrrole-2,5-dione. 2-(4-Hydroxybenzyl)-N-phenylmaleimide
 [850008-92-3]

 $C_{17}H_{13}NO_3$ 279.295


Alkaloid from the stems of *Opuntia vulgaris*. Cryst. (Me₂CO). Mp 213°. λ_{max} 229 (log ε 0.34); 322 (log ε 0.67) (MeOH).

Jiang, J.-Q. *et al.*, *Yaoxue Xuebao*, 2003, **38**, 677-679 (isol, pmr, cmr)

Orbiclamide A

O-101

[137041-28-2]

 $C_{46}H_{62}BrN_9O_{10}$ 980.954

Cyclic peptide antibiotic. Constit. of the marine sponge *Theonella* sp. Cytotoxic agent. Powder. $[\alpha]_D^{23} -60$ (c, 0.005 in MeOH). λ_{max} 203 (ε 14100); 220 (ε 10200); 269 (ε 6900); 303 (ε 1500) (MeOH) (Derep).

Fusetani, N. *et al.*, *J.A.C.S.*, 1991, **113**, 7811 (isol, uv, pmr, cmr)

Orcein

O-102

Archil. Orchil. C.I. Natural Red 28
 [1400-62-0]

Reddish-brown dye; a mixt. of 14 compds. with a phenoxazone struct. Prepd. from the lichen *Rocella tinctoria*; obtained from 5-Methyl-1,3-benzenediol by the action of aq. NH₃ and air. Biological stain. Dark brown cryst.


Musso, H. *et al.*, *Chem. Ber.*, 1956, **89**, 1659-1673; 1957, **90**, 1808-1814; 1968, **101**, 1510-1518 (isol, struct)

Musso, H. *et al.*, *Planta Med.*, 1960, **8**, 431-446 (struct)

Beecken, H. *et al.*, *Angew. Chem.*, 1961, **73**, 665-688 (*rev. struct*)

Oreobeiline

[97400-76-5]



$C_{19}H_{27}NO_4$ 333.427

Alkaloid from the stem bark and wood of *Beilschmiedia oreophila* (Lauraceae). Prisms ($CHCl_3$). Mp 116-118°. $[\alpha]_D^{20} +39$ (c, 1 in $CHCl_3$).

6-Epimer: 6-Epioreobeiline

[97400-78-7]

$C_{19}H_{27}NO_4$ 333.427

Alkaloid from the stem bark and wood of *Beilschmiedia oreophila* (Lauraceae). Needles (MeOH). Mp 207-209°. $[\alpha]_D^{20} +59$ (c, 1 in $CHCl_3$).

Tillequin, F. *et al.*, *Heterocycles*, 1985, **23**, 1357 (*uv, pmr, ms, abs config, epimer*)

Oreoline†

[11004-92-5]

$C_{26}H_{43}NO_7$ 481.628

Diterpene alkaloid, struct. unknown, similar to Lycotoline in M-309. Conts. NMe, $-OCH_2O-$, OH and 4 OMe groups. Alkaloid from underground parts of the Central Asiatic larkspur, *Delphinium oreophilum* (Ranunculaceae). Cryst. (Me_2CO or EtOH). Mp 211-213°. $[\alpha]_D -28$ (c, 1.0 in $CHCl_3$). $[\alpha]_D +55.1$ (c, 1.0 in EtOH).

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 186-188°.


Perchlorate:

Cryst. (EtOH/Et₂O). Mp 139-142°.

Bocharnikova, A.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1958, **28**, 2918 (*isol, ir*)

Oriamide

[200182-07-6]




$C_{44}H_{55}N_9O_{15}S_2$ 1014.102

Cyclic peptide antibiotic. Isol. from the sponge *Theonella* sp. Cytotoxic agent. Amorph. powder (as Na salt).

Chill, L. *et al.*, *Tetrahedron*, 1997, **53**, 16147-16152 (*isol, pmr, cmr, ms*)

Oriciacridone A

O-106



$C_{36}H_{32}N_2O_9$ 636.657

Alkaloid from the stem bark of *Oriopsis glaberrima*. Yellow cryst. Mp 249°. $[\alpha]_D^{25} -45.7$ (c, 0.07 in DMSO). λ_{max} 269 (log ϵ 4.84); 272 (log ϵ 4.85); 285 (log ϵ 4.62); 323 (log ϵ 4.28); 352 (log ϵ 4.15); 401 (log ϵ 4.14) (DMSO).

11-Hydroxy: Oriciacridone B


$C_{36}H_{32}N_2O_{10}$ 652.656

Alkaloid from the stem bark of *Oriopsis glaberrima*. Yellow cryst. Mp 309°. $[\alpha]_D^{25} -85.3$ (c, 0.07 in DMSO). λ_{max} 239 (log ϵ 4.48); 257 (log ϵ 4.6); 275 (log ϵ 4.4); 278 (log ϵ 4.4); 295 (log ϵ 4.12); 312 (log ϵ 4.13); 372 (log ϵ 3.8); 404 (log ϵ 3.88) (DMSO).

Wansi, J.D. *et al.*, *Phytochemistry*, 2006, **67**, 475-480 (*isol, pmr, cmr, ms*)

Oriciacridone D

[886227-04-9]



$C_{18}H_{15}NO_4$ 309.321


(R)-form

Alkaloid from the stem bark of *Oriopsis glaberrima*. Yellow cryst. (MeOH). Mp 265-266°. $[\alpha]_D^{25} +87.4$ (c, 0.5 in MeOH). λ_{max} 255 (log ϵ 6.05); 279 (log ϵ 5.62); 291 (log ϵ 5.6); 295 (log ϵ 5.67); 300 (log ϵ 5.58); 346 (log ϵ 4.58) (MeOH).

Wansi, J.D. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 292-296 (*isol, uv, cd, pmr, cmr*)

Oriciacridone F

[886227-06-1]



O-108

$C_{36}H_{32}N_2O_8$ 620.657


Alkaloid from the stem bark of *Oriopsis glaberrima*. α -Glucosidase inhibitor. Yellow powder (MeOH). Mp 187-189°. $[\alpha]_D^{25} +35.6$ (c, 0.62 in MeOH). λ_{max} 233 (log ϵ 1.69); 254 (log ϵ 2.06); 277 (log ϵ 1.84); 281 (log ϵ 1.85); 295 (log ϵ 1.62); 303 (log ϵ 1.64); 348 (log ϵ 0.93); 389 (log ϵ 1.18) (MeOH).

Wansi, J.D. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 292-296 (*isol, uv, pmr, cmr, ms*)

Oridine

Oreoline†

O-109



(-)-form

$C_{17}H_{23}NO_3$ 289.374

(-)-form [17366-36-8]

Alkaloid from *Papaver oreophilum* (Papaveraceae). Mp 237-240°. $[\alpha]_D -88$ (c, 0.88 in MeOH).

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 214-220°. $[\alpha]_D^{21} -71.6$ (c, 1.06 in H₂O).

N-Me: N-Methyloreoline. N-Methyloridine

$C_{18}H_{25}NO_3$ 303.4

Alkaloid from *Papaver oreophilum* (Papaveraceae). Mp 192-193° (186-188°). $[\alpha]_D -59.7$ (c, 1 in $CHCl_3$) (synthetic).

Me ether, N-Me: Mp 265-270° subl. $[\alpha]_D -39.6$ (c, 0.91 in H₂O).

O-De-Me, O¹-Me: Isooridine. Oridine 2

[78416-82-7]

$C_{17}H_{23}NO_3$ 289.374

Alkaloid from *Papaver oreophilum* (Papaveraceae). Mp 166-168°. $[\alpha]_D^{20} -57.3$ (c, 0.55 in MeOH). No stereochem. indicated but gives the same Me ether as Oridine.

(±)-form

N-Me: [58166-04-4]

Synthetic. Cryst. (EtOAc). Mp 189-192°.

N-Me, di-Ac:

Cryst. (Et₂O). Mp 170-171°.

Bernauer, K. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 2122 (*synth*)

Pfeifer, S. *et al.*, *Pharmazie*, 1966, **21**, 251; 1967, **22**, 124 (*isol*)

Mann, I. *et al.*, *Pharmazie*, 1967, **22**, 124 (*isol, uv, struct, Oreoline, N-Methyloreoline*)

Santavý, F. *et al.*, *Planta Med.*, 1967, **15**, 311 (*isol, ms, uv, pmr, struct*)

Casagrande, C. *et al.*, *J.C.S. Perkin 1*, 1975, 1652 (*synth, uv, ir, pmr, N-Methyloreoline*)


Bindra, J.S. *et al.*, *J.O.C.*, 1977, **42**, 910 (*synth, pmr, uv, ms, N-Methyloreoline*)

Věžník, F. *et al.*, *Phytochemistry*, 1981, **20**, 347 (*Isooridine*)

Orientalidine

O-110

9,10,15b,16-Tetrahydro-5,15-dimethoxy-1H,7H-[1,3]-benzodioxino[5,6-g][1,3]benzodioxolo[5,6-a]quinolizine, 9CI. Bractavine



(S)-form

C₂₂H₂₃NO₆ 397.427

(S)-form [23943-90-0]

Alkaloid from *Papaver orientale*, *Papaver pseudo-orientale*, *Papaver bracteatum* and other *Papaver* spp. (Papaveraceae). Needles (EtOAc). Mp 196-198°. [α]_D²² -253 (c, 0.444 in CHCl₃). [α]_D²¹ -254 (c, 0.310 in MeOH).

(±)-form [22325-09-3]

Needles (EtOAc). Mp 190-192°.

Preininger, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 875 (*isol, ir, pmr, uv, ms, struct*)

Šimánek, V. *et al.*, *Tet. Lett.*, 1969, 2109 (*struct*)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1975, 1822 (*synth, pmr*)

Denisenko, O.N. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 547; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 456 (*isol*)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 474; 1984, **20**, 81; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 402; 1984, **20**, 76 (*isol*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1980, **45**, 914 (*isol*)


Sariyar, G. *et al.*, *Planta Med.*, 1980, **38**, 378 (*isol*)

Phillipson, J.D. *et al.*, *Planta Med.*, 1981, **43**, 261 (*isol*)

Kerekes, P. *et al.*, *CA*, 1982, **97**, 127857t (*synth*)

Orientaline†

O-111



(R)-form

C₁₉H₂₃NO₄ 329.395

(R)-form

Synthetic.

Hydrochloride:

Noncryst. [α]_D²¹ -53.4 (c, 1.26 in H₂O).

Perchlorate: Mp 124-125°.

(S)-form [27003-74-3]

Alkaloid from seeds and pod-walls of *Erythrina arborescens* and leaves of *Erythrina abyssinica* (Fabaceae). Biosynthetic precursor of Isothebaine.

Hydrochloride:

Noncryst. [α]_D²¹ +53.5 (c, 1.28 in H₂O).

Perchlorate: Mp 128-130°.

Di-Me ether: see Laudanosine, L-69

N-De-Me: **N-Nororientaline**

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Erythrina poeppigiana*, *Erythrina herbacea* and *Erythrina x. bidwillii* (Fabaceae). Cryst. (EtOAc). Mp 145-147°. [α]_D +42 (c, 0.18 in CHCl₃).

N-De-Me; hydrochloride: Mp 247-249°.

(±)-form [20938-53-8]

Synthetic.

Perchlorate: Mp 127°.

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1961, **81**, 1253 (*synth*)

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 4550

(*resoln*)


Ghosal, S. *et al.*, *Phytochemistry*, 1972, **11**, 2101 (*isol*)

Barton, D.H.R. *et al.*, *J.C.S. Perkin I*, 1973, 874 (*isol, deriv*)

Tiwari, K.P. *et al.*, *Phytochemistry*, 1979, **18**, 704 (*isol*)

Orientalinone

O-112



(-)-form

C₁₉H₂₁NO₄ 327.379

(+)-form

Synthetic. Mp 184-186° dec. [α]_D²¹ +62.3 (c, 0.74 in CHCl₃).

(-)-form [2689-17-0]

Alkaloid from *Papaver orientale* (Papaveraceae). Intermediate in the biosynth. of aporphine alkaloids. Mp 183-184° dec. [α]_D²¹ -62.6 (c, 0.84 in CHCl₃).

Me ether: **Roemerinalinone**

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph.

11,12-Dihydro: 11,12-Dihydroorientalinone

[112652-49-0]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. [α]_D -81 (c, 0.08 in MeOH).

13-Epimer: Isoorientalinone

[30389-03-8]

C₁₉H₂₁NO₄ 327.379

Trace alkaloid from *Roemeria hybrida*, in admixture with orientalinone.

13-Epimer, Me ether: Isoroemerinalinone

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Roemeria hybrida*. Amorph. [α]_D -47 (c, 0.08 in CHCl₃).

13-Epimer, 8,9-dihydro: 8,9-Dihydroisoorientalinone

[6392-31-0]

C₁₉H₂₃NO₄ 329.395

Minor alkaloid from *Papaver orientale* (Papaveraceae). Amorph. [α]_D +37 (c, 0.15 in CHCl₃) (+50). Originally de-

scribed by Battersby *et al* as (+)-Dihydroorientalinone.

13-Epimer, 8,9-dihydro, Me ether: 8,9-Dihydroisoroemerinalinone

C₂₀H₂₅NO₄ 343.422

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. [α]_D +23 (c, 0.07 in MeOH).

(±)-form [33458-68-3]

Synthetic. Mp 227-229° dec.

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 4550-4556

(*synth, ir, pmr*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1966, 170-171 (*isol*)


Kametani, T. *et al.*, *J.C.S. Perkin I*, 1972, 1435-1441 (*synth*)

Kametani, T. *et al.*, *J.A.C.S.*, 1977, **99**, 3805-3808 (*synth*)

Gözler, B. *et al.*, *Tetrahedron*, 1987, **43**, 1765-1770 (*11,12-Dihydroorientalinone, Isoorientalinone, Roemerinalinone, Isoroemerinalinone, Dihydroisoroemerinalinone, pmr, cd, abs config*)

Orixine

O-113



(S)-form

C₁₇H₂₁NO₆ 335.356

(S)-form [17232-53-0]

Abs. config. revised in 2002. Alkaloid from the root bark of *Orixia japonica*. Mp 152.5°. [α]_D¹⁷ +83.3 (CHCl₃).

3'-Me ether: 3'-O-Methylorixine

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Orixia japonica*. Oil.

[α]_D²³ +10.5 (c, 0.1 in CHCl₃). λ_{max} 254 (ε 43600); 297 (ε 3400); 316 (ε 3400) (MeOH).

O²-De-Me: Nororixine

C₁₆H₁₉NO₆ 321.329

Alkaloid from the root bark of *Orixia japonica* (Rutaceae). Needles or prisms (EtOH). Mp 199-200°. Exists as 2(1H)-quinolinone tautomer.

3'-Deoxy, 2'-ketone: Orixinone

[39027-00-4]

C₁₇H₁₉NO₅ 317.341

Alkaloid from the stems of *Orixia japonica*. Pale-yellow needles (Et₂O/petrol). Mp 102-103°.

Terasaka, M. *et al.*, *Chem. Pharm. Bull.*, 1960, **8**, 523; 1142 (*isol, uv, ir, struct, Orixine, Nororixine*)

Bowman, R.M. *et al.*, *J.C.S.(C)*, 1967, 2368

(*pmr, synth*)

Donnelly, W.J. *et al.*, *J.C.S. Perkin I*, 1972, 2116 (*isol, ir, pmr, ms, struct, synth, Orixinone*)


Funayama, S. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1885-1889 (*abs config*)

Noshita, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2001, **65**, 710-713 (*3'-O-Methylorixine*)

Boyd, D.R. *et al.*, *Chem. Comm.*, 2002, 3070-3071 (*abs config*)

Ormojanine

16,17-Didehydropanamine, 9CI
[4697-87-4]



Relative
configuration

C₂₀H₃₁N₃ 313.485

Probable struct. CAS gives an absolute config., but there does not seem to be evidence for this. Alkaloid from *Ormosia jamaicensis* (Fabaceae). Mp 126°. [α]_D -143 (EtOH).

Perchlorate (1:2): Mp 256-257°.


Tripicrate: Mp 169-170°.

Hassall, C.H. *et al.*, *J.C.S.*, 1964, 2657-2663 (*isol, uv*)

Valenta, Z. *et al.*, *Can. J. Chem.*, 1966, **44**, 2525-2538; 2539-2551 (*struct, pmr, stereochem*)

Ormosanine

Piptamine



(+)-form

C₂₀H₃₅N₃ 317.517

Stereoisomeric with Dasycarpine, D-85, Templetine, T-68 and Piptanthine, P-476.

(+)-form

Alkaloid from *Ormosia panamensis* (Fabaceae). Prisms. Mp 167-168°. [α]_D²⁵ +3.3 (c, 1.04 in CHCl₃). (+)-Ormosanine has only been isol. once and in view of the weak opt. activity is doubtful. CAS No. not found to 2008.

Hydroiodide (1:2): Mp 249° dec.

16-Epimer: **16-Epiormosanine**

[58560-21-7]

C₂₀H₃₅N₃ 317.517

Minor alkaloid from *Hovea linearis* (Fabaceae), also by hydrogenation of (-) Podopetaline. Prisms (Me₂CO). Mp 113-114°. [α]_D 0 (c, 0.30 in CHCl₃). [α]_D -2.3 (c, 0.30 in EtOH).

6,16-Diepimer: **6-Epi-16-epiormosanine**

C₂₀H₃₅N₃ 317.517

Minor alkaloid from *Hovea linearis*. Mp 154-155°. [α]_D -28 (MeOH). Tentative struct. Rel. config. only. CAS No. not found to 2008.

(-)-form [5001-21-8]

Alkaloid from *Podopetalum ormondii*, (preferred genus name *Ormosia*), *Ormosia semicastrata* and *Ormosia jamaicensis* (Fabaceae). Mp 178-179°. [α]_D -9. Occurs as partial racemates and forms a quasir-

acemate with (-)-Podopetaline, in which form it was isol. from *P. ormondii*.

18-Epimer: **18-Epiormosanine**

[33792-80-2]

Alkaloid from *Ormosia semicastrata* (Fabaceae). Mp 267° dec. [α]_D -10. Rel. config. only known.

(±)-form [34429-31-7]

Alkaloid from *Ormosia* spp. and from *Piptanthus nanus* (Fabaceae). Mp 183-184°.

Hydroiodide (1:2): Mp 249° dec.

[54292-99-8, 58917-72-9]

Lloyd, H.A. *et al.*, *J.A.C.S.*, 1958, **80**, 1506-1510 (*isol*)

Naegeli, P. *et al.*, *Tet. Lett.*, 1963, 2069-2073 (*struct*)

Hassall, C.H. *et al.*, *J.C.S.*, 1964, 2657-2663 (*isol*)

McLean, S. *et al.*, *Can. J. Chem.*, 1971, **49**, 1976-1978; 1972, **50**, 1639-1641 (*isol, struct, epimer*)

Cannon, J.R. *et al.*, *Tet. Lett.*, 1974, **15**, 1683-1686 (*abs config*)

Lamberton, J.A. *et al.*, *Tet. Lett.*, 1975, **16**, 3875-3876 (*16-Epiormosanine*)

Liu, H.-J. *et al.*, *Can. J. Chem.*, 1976, **54**, 97-109 (*synth*)

Mackay, M.F. *et al.*, *J. Cryst. Mol. Struct.*, 1976, **6**, 125-137 (*16-Epiormosanine, cryst struct*)

McLean, S. *et al.*, *Can. J. Chem.*, 1981, **59**, 34-37 (*isol, abs config, biosynth*)

Lamberton, J.A. *et al.*, *Aust. J. Chem.*, 1982, **35**, 2577-2582 (*16-Epiormosanine, 6-Epi-16-epiormosanine*)

Bhacca, N.S. *et al.*, *J.A.C.S.*, 1983, **105**, 2538-2544 (*cmr, pmr*)

Le, P.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 283-293 (*pmr, cmr*)

Ormosine

[1400-66-4]

C₂₀H₃₃N₃ 315.501

Poss. identical with Panamine, P-53. Struct. unknown. Alkaloid from seeds of *Ormosia dasycarpa* (Fabaceae). Shows analgesic props. Cryst. + 3-4H₂O. Mp 85-87°.

Dipicrate: Mp 178°.


Methiodide: Mp 245-250°.

Hess, K. *et al.*, *Ber.*, 1919, **52**, 1976-1983 (*isol*)
Lloyd, H.A. *et al.*, *J.A.C.S.*, 1958, **80**, 1506-1510

Ormosinine

21-Ormosinin-20-ylpanamine, 9CI

[14350-67-5]



Absolute
Configuration

C₄₀H₆₆N₆ 631.002

Dimer containing Panamine, P-53 and Ormosanine, O-115 units. Alkaloid from the seeds of *Ormosia dasycarpa*, *Ormosia panamensis* and *Ormosia jamaicensis*, detected in other *Ormosia* spp. (Fabaceae). Needles (EtOAc). Mp 219-

220° (203-205°). [α]_D²⁵ +8.9 (CHCl₃).

Dipicrate: Mp 146-148°. Could not be recryst.

Stereoisomer: **Ormojine**

[11004-95-8]

C₄₀H₆₆N₆ 631.002

Alkaloid from *Ormosia jamaicensis* (Fabaceae). Mp 154-155°. [α]_D²⁰ +24 (c, 1.46 in EtOH). Of unknown stereochem.; dimer of Ormosanine in P-53.

Hess, K. *et al.*, *Ber.*, 1919, **52**, 1976-1983 (*isol*)
Lloyd, H.A. *et al.*, *J.A.C.S.*, 1958, **80**, 1506-1510 (*isol*)

Hassall, C.H. *et al.*, *J.C.S.*, 1964, 2657-2663 (*isol, Ormojine*)


Deslongchamps, P. *et al.*, *Can. J. Chem.*, 1966, **44**, 2539-2551 (*struct*)

Davies, A.P. *et al.*, *Tet. Lett.*, 1966, 6291-6294 (*struct, Ormojine*)

Bhacca, N.S. *et al.*, *J.A.C.S.*, 1983, **105**, 2538-2544 (*cmr, pmr, struct*)

Ornazidine

[69086-73-3]



Tentative structure

C₁₆H₁₉NO₃ 273.331

Alkaloid from the bulbs of *Crinum ornatum* (Amaryllidaceae). Mp 217°.

O¹¹-Ac: **Ornazamine**

[69086-74-4]

C₁₈H₂₁NO₄ 315.368

Isol. from *Crinum ornatum* (Amaryllidaceae). Mp 218-220° dec.

3,4-Didehydro, di-Me ether: **Ornamine**

[69086-75-5]


C₁₈H₂₁NO₃ 299.369

Isol. from *Crinum ornatum* (Amaryllidaceae). Mp 212-213°.

Onyiriuka, O.S. *et al.*, *Isr. J. Chem.*, 1978, **17**, 185 (*isol, ms, struct, derivs*)

Ornibactin C4

[154071-68-8]



R = CH₃

C₂₆H₄₈N₈O₁₃ 680.711

Prod. by *Pseudomonas cepacia*. Siderophore.

Stephan, H. *et al.*, *Biol. Met.*, 1993, **6**, 93

Ornibactin C6

[154071-69-9]

As Ornibactin C4, C-119 with

R = -CH₂CH₂CH₃

C₂₈H₅₂N₈O₁₃ 708.765

Prod. by *Pseudomonas cepacia*. Siderophore.

Stephan, H. *et al.*, *Biol. Met.*, 1993, **6**, 93 (*isol*)

Ornibactin F O-121

Ornibactin C8

[147363-90-4]

As Ornibactin C4, O-119 with

R = $-(\text{CH}_2)_4\text{CH}_3$

$\text{C}_{30}\text{H}_{56}\text{N}_8\text{O}_{13}$ 736.818

Gross structs. of Ornibactin C8 and Ornibactin F are the same but the stereochem. at C-3' may vary. Prod. by *Pseudomonas* sp. TVV 69. Siderophore.

[154170-72-6]

Stephan, H. *et al.*, *Annalen*, 1993, 43 (*isol, struct*)

Stephan, H. *et al.*, *Biol. Met.*, 1993, **6**, 93 (*isol, struct*)

Orobanchamine O-122


$\text{C}_{20}\text{H}_{31}\text{NO}_{14}$ 509.63

Struct. unknown. Alkaloid from *Orobanche lutea* (Orobanchaceae). Cryst. (CHCl_3 /petrol). Mp 193-194°. $[\alpha]_{\text{D}}^{25} +33.92$ (EtOH). No active hydrogens.

Rubinshtein, M.M. *et al.*, *Zh. Obshch. Khim.*, 1953, **23**, 156; *CA*, **48**, 696

Oroidin O-123

N-[3-(2-Amino-1H-imidazol-4-yl)-2-propenyl]-4,5-dibromo-1H-pyrrole-2-carboxamide, 9CI
[34649-22-4]



$\text{C}_{11}\text{H}_{11}\text{Br}_2\text{N}_5\text{O}$ 389.049

Alkaloid from the sponges *Agelas oroides*, *Agelas confifera*, *Agelas longissima*, *Agelas mauritiana*, *Agelas clathrodes*, *Agelas wiedenmayeri*, *Agelas sceptrum*, *Axinella verrucosa*, *Axinella damicornis*, *Hymeniacidon* sp., *Pseudaxinysa cantharella*, *Acanthella carteri* and *Acanthella aurantiaca*. Antiserotonergic, adrenergic antagonist, serotonin antagonist, anti-muscarinic, antifouling, anti-histamine. Key intermed. in the biosynth. of the marine pyrrole-imidazole alkaloids. Noncryst. Poorly sol. hexane. λ_{max} 296 (ϵ 24500) (MeOH/KOH) (Derep). λ_{max} 278 (ϵ 21000) (MeOH) (Derep).

N-Ac: Mp 256-258°.

N¹-Me: **Sventrine**

[392315-92-3]

$\text{C}_{12}\text{H}_{13}\text{Br}_2\text{N}_5\text{O}$ 403.076

Alkaloid from the sponge *Agelas sventres*. Light yellow powder. λ_{max} 274 (log ϵ 4.27) (MeOH).

2-Debromo: **Hymenidine**

[107019-95-4]

$\text{C}_{11}\text{H}_{12}\text{BrN}_5\text{O}$ 310.153

Isol. from the sponge *Hymeniacidon* sp. and *Agelas clathrodes*. Serotonin receptor antagonist. Antihypertensive, antimuscarinic, smooth muscle contractant. Amorph. solid. Sol. MeOH,

butanol. λ_{max} 269 (ϵ 21400) (MeOH) (Derep). λ_{max} 270 (ϵ 23000) (MeOH) (Berdy).

2,3-Didebromo: **Clathrodine**

[135383-64-1]

$\text{C}_{11}\text{H}_{13}\text{N}_5\text{O}$ 231.257

Isol. from the sponge *Agelas clathrodes*. Cholinergic receptor blocker, antiadrenergic, antimuscarinic neurotoxin. Amorph. solid. λ_{max} 272 (ϵ 24500) (MeOH) (Derep).

Forenza, S. *et al.*, *Chem. Comm.*, 1971, 1129-1130 (*isol, uv, ir, pmr*)

Garcia, E.E. *et al.*, *Chem. Comm.*, 1973, 78-79 (*struct*)

Walker, R.P. *et al.*, *J.A.C.S.*, 1981, **103**, 6772-6773 (*isol*)

Kobayashi, J. *et al.*, *Experientia*, 1986, **42**, 1176-1177 (*Hymenidine*)

Morales, J.J. *et al.*, *J. Nat. Prod.*, 1991, **54**, 629-631 (*Clathrodine*)

Daninos-Zeghal, S. *et al.*, *Tetrahedron*, 1997,

53, 7605-7614 (*Hymenidine, synth*)

Koenig, G.M. *et al.*, *Planta Med.*, 1998, **64**,

443-447 (*isol, pmr, cmr, ms*)

Lindel, T. *et al.*, *J.O.C.*, 2000, **65**, 2806-2809 (*synth*)

Assmann, M. *et al.*, *J. Nat. Prod.*, 2001, **64**,

1593-1595 (*Sventrine*)

Berré, F. *et al.*, *Tet. Lett.*, 2002, **43**, 4935-4938 (*synth*)

Hoffmann, H. *et al.*, *Synthesis*, 2003, 1753-1783 (*rev*)

Breckle, G. *et al.*, *Z. Naturforsch., B*, 2003, **58**, 451-456 (*Sventrine, synth*)


Orosomycin O-124

4,5-Dihydro-6-(N-methylformamido)-5-oxo-1,2-dithiolo[4,3-b]pyrrole. Thioaurin. HA 9. vD 844. Antibiotic HA 9. Antibiotic vD 844. Heyden Antibiotic 9. AK-Ps.

Antibiotic AK-Ps

[21787-65-5]

[67527-37-1]



$\text{C}_7\text{H}_6\text{N}_2\text{O}_2\text{S}_2$ 214.269

Identity with Thioaurin and with Ak-Ps is not certain. Isol. from an unidentified *Streptomyces* sp. and from *Streptomyces flavochromogenes iwayaensis* (in the case of AK-Ps). Active against gram-positive and -negative bacteria and fungi. Yellow needles (EtOAc). Mp 181-182°. λ_{max} 300 (ϵ 16500) (MeOH/NaOH) (Derep). λ_{max} 231 (ϵ 6600); 300 (sh) (ϵ 2500); 368 (ϵ 15800) (MeOH) (Derep).

► LQ2000600

N-De-Me: **Antibiotic vD 846**. vD 846

[21787-66-6]

$\text{C}_6\text{H}_4\text{N}_2\text{O}_2\text{S}_2$ 200.242

From a *Streptomyces* sp. Active against gram-positive and -negative bacteria. Orange prisms (Me_2CO). Sol. EtOAc, MeOH, CHCl_3 ; fairly sol. H_2O ; poorly sol. hexane. Mp 270-280° dec. λ_{max} 247 (ϵ 5800); 301 (ϵ 3350); 387 (ϵ 11000) (EtOH) (Derep).

► LD₅₀ (mus, ivn) 5 - 10 mg/kg. LQ3164500

Jensen, B. *et al.*, *J. Antibiot.*, 1969, **22**, 231 (*cryst struct*)

v. Daehne, W. *et al.*, *J. Antibiot.*, 1969, **22**, 233 (*isol, uv, struct*)


Jensen, B. *et al.*, *Acta Cryst. B*, 1971, **27**, 392

Fabian, J. *et al.*, *Z. Chem.*, 1972, **12**, 348 (*uv*)

Japan. Pat., 1978, 78 47 589; *CA*, **89**, 105898q (*AK-Ps*)

Orthidine A O-125

[1042101-21-2]



$\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4$ 384.394

Dimer of Tubastrine, T-656. Alkaloid from *Aplidium orthium*. Pale yellow gum. Racemic. λ_{max} 207 (log ϵ 4.56); 226 (log ϵ 4.17); 284 (log ϵ 4.02); 311 (sh) (log ϵ 3.73) (MeOH).

2-Epimer: **Orthidine B**

[1042101-23-4]


$\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4$ 384.394

Alkaloid from *Aplidium orthium*. Pale yellow gum. Racemic. λ_{max} 205 (log ϵ 4.55); 226 (log ϵ 4.16); 284 (log ϵ 4.08); 308 (sh) (log ϵ 3.83) (MeOH).

Pearce, A.N. *et al.*, *Tetrahedron*, 2008, **64**, 5748-5755 (*isol, pmr, cmr*)

Orthidine C O-126

[1042101-25-6]



Relative Configuration

$\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4$ 384.394

Dimer of Tubastrine, T-656. Alkaloid from *Aplidium orthium*. Pale yellow gum. Racemic. λ_{max} 205 (log ϵ 4.59); 224 (log ϵ 4.32); 284 (log ϵ 4.23); 311 (sh) (log ϵ 3.97) (MeOH).

2-Epimer: **Orthidine D**

[1042101-28-9]


$\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4$ 384.394

Alkaloid from *Aplidium orthium*. Pale yellow gum. λ_{max} 206 (log ϵ 4.59); 227 (log ϵ 4.23); 287 (log ϵ 4.25); 310 (sh) (log ϵ 4.01) (MeOH).

Pearce, A.N. *et al.*, *Tetrahedron*, 2008, **64**, 5748-5755 (*isol, pmr, cmr*)

Orthidine E O-127

[1042101-31-4]



Relative Configuration


$\text{C}_{18}\text{H}_{22}\text{N}_6\text{O}_4$ 386.41

Dimer of Tubastrine, T-656. Alkaloid from *Aplidium orthium*. Pale yellow gum. λ_{\max} 206 (log ϵ 4.45); 230 (log ϵ 3.83); 285 (log ϵ 3.45) (MeOH).

Pearce, A.N. *et al.*, *Tetrahedron*, 2008, **64**, 5748-5755 (*isol*, *pmr*, *cmr*)

Oscillatorin**O-128**


[182261-00-3]



$C_{60}H_{85}N_{15}O_{14}$ 1240.424

Cyclic peptide antibiotic. *Isol.* from the cyanobacterium *Oscillatoria agardhii*. Chymotrypsin inhibitor. Amorph. solid. λ_{\max} 283 (log ϵ 3.4) (H_2O).

Sano, T. *et al.*, *Tet. Lett.*, 1996, **37**, 6873-6876 (*isol*, *uv*, *pmr*, *cmr*)

Osoic acid**O-129**

$C_{15}H_{12}O_8$ 320.255

2',3-Di-Me ether, 1'-amide, 1-Me ester: *Neoplaether*


$C_{18}H_{19}NO_7$ 361.351

Prod. by *Neoplaconema napellum* IFB-E016. Cytotoxic. Cryst. (DMSO). Mp 298-299°.

Feng, W. *et al.*, *FEMS Microbiol. Lett.*, 2006, **261**, 218-223 (*Neoplaether*)

Osornine**O-130**

[89412-88-4]



$C_{38}H_{42}N_2O_7$ 638.759


Represents a member of the oxyacanthan series diastereomeric with Thalrugosaminine, T-346 and derivs. Alkaloid from *Berberis buxifolia* (Berberidaceae). Mp 244-245°. $[\alpha]_D^{25}$ -151 (c, 0.36 in $CHCl_3$).

[89412-89-5]

Leet, J.E. *et al.*, *J. Nat. Prod.*, 1983, **46**, 908-912 (*isol*, *uv*, *cd*, *pmr*, *ms*, *struct*)

Ostreogrycin A**O-131**

Mikamycin A. *Pristinamycin II_A*. *Staphylomycin M₁*. *Streptogramin A*. *Syncothrecin A*. *Synergistin A₁*. *Virginiamycin M₁*. *Vernamycin A*. *E129A*. *PA 114A*. *1745Z_{3A}*. *547C*. *Antibiotic 14752-2*. *Antibiotic E129A*. *Antibiotic PA 114A*. *Antibiotic 1745Z_{3A}*. *Antibiotic 547C*. *Factor M* [21411-53-0]



$C_{28}H_{35}N_3O_7$ 525.6

CAS numbering shown. *Isol.* from *Streptomyces pristinaespiralis*, *Streptomyces virginiae* and other *Streptomyces* spp. Active against gram-positive organisms. Cryst. (EtOAc). Mp 200° dec. $[\alpha]_D^{23}$ -209 (c, 1 in EtOH). Log P 0.39 (uncertain value) (calc). λ_{\max} 303 (ϵ 15800) (EtOH/HCl) (Derep). λ_{\max} 293 (ϵ 21900) (EtOH/NaOH) (Derep). λ_{\max} 228 (ϵ 32400); 272 (sh) (ϵ 10000) (95% EtOH) (Derep).

Mixture with ca. 25% other *Virginiamycins*: *Virginiamycin*, *BAN*, *INN*, *USAN*. *Cebin V*. *Eskalin V*. *Eskamicin*. *Stafac*. *Staphylomycin*. *Mikamycin*. *Ostreogrycin*. *Patricin*. *Pristinamycin*. *Streptogramin*. *Vernamycin*

[11006-76-1] Antibacterial food additive. Powder. $[\alpha]_D$ -124 (c, 0.05 in EtOH). Dec. at 138-140°.

► LD₅₀ (mus, orl) 2100 mg/kg. ZA4850000

26,27S-Dihydro: *Ostreogrycin G*. *Pristinamycin II_B*. *Virginiamycin M₂* [21102-49-8]

$C_{28}H_{37}N_3O_7$ 527.616

Prod. by the soil organism *Streptomyces ostreogriseus*. Active against gram-positive bacteria. Pale-yellow solid. Sol. MeOH, Et₂O; fairly sol. C₆H₆; poorly sol. H₂O, hexane. Mp 122-127° dec. (after sintering at 90°). $[\alpha]_D^{25}$ +78 (c, 1.36 in EtOH). λ_{\max} 215 (E1%/1cm 650) (MeOH) (Berdy). λ_{\max} 235 (ϵ 15135); 295 (ϵ 11480) (EtOH/NaOH) (Berdy).

Deoxy: *Antibiotic L 156588*. *L 156588*

[118693-65-5]

$C_{28}H_{35}N_3O_6$ 509.601

From *Streptomyces olivaceus*. Gastrin and brain cholecystokinin antagonist. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 212 (ϵ 28400); 275 (ϵ 38900) (MeOH) (Derep).

Deoxy, 14,15-epoxy: *Antibiotic L 156587*. *L 156587*

[119465-45-1]

$C_{28}H_{33}N_3O_7$ 523.585

From *Streptomyces olivaceus*. Gastrin and brain cholecystokinin antagonist. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. λ_{\max} 212 (ϵ 27700); 325 (ϵ 21100) (MeOH) (Derep).

Deoxy, 14,15-epoxy, 16-alcohol: *Antibiotic L 156586*. *L 156586*

$C_{28}H_{35}N_3O_7$ 525.6

From *Streptomyces olivaceus*. Gastrin and brain cholecystokinin antagonist. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. The C-15 epimer Antibiotic L 156906 is semisynthetic. λ_{\max} 212 (ϵ 28400); 275 (ϵ 38900) (MeOH) (Derep).

[118693-64-4, 118759-89-0]

Ball, S. *et al.*, *Biochem. J.*, 1958, **68**, 24P (*isol*)
Kingston, D.G.I. *et al.*, *J.C.S.(C)*, 1966, 1669; 1856 (*nmr*, *ir*, *ms*, *struct*)

Delpierre, G.R. *et al.*, *Tet. Lett.*, 1966, 369 (*struct*)

Preud'Homme, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 585 (*isol*, *ms*, *struct*)

Crooy, P. *et al.*, *J. Antibiot.*, 1972, **25**, 371

Bycroft, B.W. *et al.*, *J.C.S. Perkin 1*, 1977, 2464 (*cmr*, *pmr*, *conformn*)

Ogata, K. *et al.*, *J. Antibiot.*, 1978, **31**, 1313 (*isol*, *ir*, *nmr*)

Kingston, D.G.I. *et al.*, *J.A.C.S.*, 1983, **105**, 5106 (*biosynth*)

Biot, A.M. *et al.*, *Drugs Pharm. Sci.*, 1984, **22**, 695 (*rev*)

Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, VRF000

Paris, J.M. *et al.*, *Recent Prog. Chem. Synth. Antibiot.*, 1990, 183 (*rev*)

Shepherd, M.J. *et al.*, *Food Contaminants: Sources and Surveillance*, Royal Society of Chemistry, 1991, 175 (*anal*)

Lam, Y.K.T. *et al.*, *J. Antibiot.*, 1991, **44**, 613 (*derivs*)

Kingston, D.G.I. *et al.*, *Environ. Sci. Res.*, 1992, **44**, 105 (*rev*, *biosynth*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 224

Pechere, J.-C. *et al.*, *Drugs*, (Suppl. 1), 1996, **51**, 13 (*rev*)


Schlessinger, R.H. *et al.*, *J.A.C.S.*, 1996, **118**, 3301 (*synth*)

Breuilles, P. *et al.*, *Tet. Lett.*, 1998, **39**, 3145-3148; 3149-3152 (*synth*)

Dang, J. *et al.*, *Org. Biomol. Chem.*, 2004, 2919-2924 (*pmr*, *cmr*, *conformn*)

Mortensen, M.S. *et al.*, *Org. Lett.*, 2007, **9**, 3105-3108 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VRF000

Ostrerine A**O-132**


$C_{22}H_{30}N_2O_6$ 558.556

Isol. from the marine mollusc *Ostrea rivularis*. Amorph. powder. $[\alpha]_D^{21}$ +21 (c, 0.7 in MeOH).

Ouyang, M.-A. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 79-83 (*isol*, *pmr*, *cmr*, *ms*)

Otanthusic acid O-133

7-(2-Thienyl)-2,4,6-heptatrienoic acid.
Otanthus acid



C₁₁H₁₀O₂S 206.265

(E,E,E)-form

2-Methylpropylamide: **N-Isobutylothanthusic acid amide**. N-(2-Methylpropenyl)-7-(2-thienyl)-2,4,6-heptatrienamamide, 9CI

[52657-11-1]

C₁₅H₁₉NO₂S 277.387

Alkaloid from the roots of *Otanthus maritimus* and from *Anacyclus tomentosus* (Asteraceae). Yellow cryst. (Et₂O). Mp 183°.

Piperidide: **Otanthus acid piperidide**

[52657-10-0]

C₁₆H₁₉NOS 273.398

Alkaloid from the roots of *Otanthus maritimus* (Asteraceae). Yellow cryst. (Et₂O). Mp 96°.

2,3-Didehydropiperidide: **Otanthusic acid 2,3-didehydropiperidide**

[52657-09-7]

C₁₆H₁₇NOS 271.382


Alkaloid from the roots of *Otanthus maritimus* (Asteraceae). Yellow cryst. (Et₂O/petrol). Mp 116°.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1974, **107**, 1038 (*isol, uv, ir, ms, pmr, struct*)

Bellassoued, M. *et al.*, *Synth. Commun.*, 1997, **27**, 3103-3117 (*piperidide, synth*)

Oteromycin O-134

Antibiotic MF 5810. MF 5810
[170591-45-4]



C₃₂H₄₁NO₃ 487.681

Tetramic acid deriv. Prod. by unidentified fungal strains (ATCC 74201 and ATCC 74202). Endothelin receptor antagonist. Powder. [α]_D²⁵ -60 (c, 0.5 in MeOH). λ_{max} 209 (ε 16250); 235 (sh) (MeOH).

5'-Deoxy, 4'-hydroxy, 5',7''-didehydro(Z), 16,17-dihydro: **Talaroconvolutin A**

C₃₂H₄₁NO₃ 487.681

Prod. by *Talaromyces convolutus*. Antifungal agent. Yellow prisms (cyclohexane). Mp 142-144°. [α]_D²⁰ -111 (c, 0.45 in CHCl₃). λ_{max} 255 (log ε 4.01); 414 (log ε 4.44) (EtOH).

Stereoisomer, 4'-hydroxy, 16,17-dihydro:

Talaroconvolutin B

C₃₂H₄₃NO₄ 505.696

Prod. by *Talaromyces convolutus*. Antifungal agent. Microcryst. (cyclohexane). Mp 112-114°. [α]_D²⁰ -57 (c, 0.11 in EtOH). Stereoisomeric with ZG 1494α at C-5' or C-16 or both. λ_{max} 226 (log ε 3.84); 273 (log ε 3.09) (EtOH).

4''-Hydroxy, 16,17-dihydro: **Antibiotic ZG 1494α**. ZG 1494α. Antibiotic 1494A. 1494A

[182320-33-8]

C₃₂H₄₃NO₄ 505.696

Prod. by *Penicillium rubrum* and *Talaromyces convolutus*. Platelet-activating factor and acetyltransferase inhibitor. Antifungal agent. Powder. [α]_D²⁵ -133 (c, 1 in EtOH). λ_{max} 206; 221; 266 (EtOH).

Singh, S.B. *et al.*, *J.O.C.*, 1995, **60**, 7040-7042 (*isol, uv, ir, pmr, cmr*)

West, R.R. *et al.*, *J. Antibiot.*, 1996, **49**, 967-973 (ZG 1494z)

U.S. Pat., 1996, 5 550 148; *CA*, **125**, 257170w (ZG 1494z)

Suzuki, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 768-772 (*Talaroconvolutins, activity*)

Otocamine O-135

[11004-99-2]

C₃₇H₄₀N₂O₆ 608.733

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the bark of *Nectandra (Ocotea) rodiaei* (Lauraceae). Cryst. + 1H₂O (as dihydrochloride). Mp 281° (dihydrochloride). [α]_D +268 (c, 1.0 in H₂O) (dihydrochloride).


Hearst, P.J. *et al.*, *J.O.C.*, 1964, **29**, 466-470 (*isol*)

Otonecine O-136

2,3,5,7a-Tetrahydro-1,7a-dihydroxy-7-(hydroxymethyl)-4-methyl-1H-pyrrolizinium, 9CI

[6887-34-9]

[7201-79-8]



C₉H₁₅NO₃ 185.222

Necine base from several pyrrolizidine alkaloids. Hygroscopic cryst. (as hydrochloride). Mp 145-147° (137-138°) (hydrochloride). [α]_D²³ -13 (c, 0.96 in EtOH). Hydrochloride difficult to crystallise.

► Highly toxic.

Zhdanovich, E.S. *et al.*, *Zh. Obshch. Khim.*, 1941, **11**, 835

Briggs, L.H. *et al.*, *J.C.S.*, 1965, 2492 (*ir*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1967, **20**, 801 (*struct, pmr*)

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)


Niwa, H. *et al.*, *Tet. Lett.*, 1983, **24**, 5731

(*synth*)

Vedejs, E. *et al.*, *J.A.C.S.*, 1998, **120**, 3613-3622 (*synth*)

Otosenine O-137

Tomentosine†. Othosenine
[16958-29-5]



Absolute configuration

C₁₉H₂₇NO₇ 381.425

Cyclic ester of Otonecine, O-136 and Jaconecic acid. Stereoisomeric with Petasitenine, P-277. Alkaloid from *Senecio cineraria*, *Senecio floridanus (Brachyglottis floridiana)*, *Senecio fluviatilis*, *Senecio othonnae*, *Senecio renardi* and *Cacalia floridana* (Asteraceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 219° (232°). [α]_D +20.8 (+14) (CHCl₃). The (-)-enantiomer was reported from *S. jacobaea* but without proper identification.

► WH1306400

Ac: **Floroseneine**

[16958-30-8]

C₂₁H₂₉NO₈ 423.462

Alkaloid from *Senecio fluviatilis* and *Cacalia floridana* (Asteraceae). Flakes (cyclohexane/C₆H₆). Mp 101-103°. [α]_D +31.9 (c, 1.38 in CHCl₃).

► VS3595300

[981-42-0, 30484-74-3]

Zhdanovich, E.S. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1941, **11**, 835; *CA*, **36**, 4123 (*isol*)

Wunderlich, J.A. *et al.*, *Chem. Ind. (London)*, 1962, 2089 (*struct*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 3570

(*Floroseneine*)

Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 258; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 221 (*-)-form*)

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832 (*uv*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (*cd*)


Perez-Salazar, A. *et al.*, *Acta Cryst. B*, 1977, **33**, 3525 (*cryst struct*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173

(*cmr*)

Wiedenfeld, H. *et al.*, *Acta Cryst. C*, 1990, **46**, 1345 (*cryst struct*)

Loizzo, M.R. *et al.*, *Arch. Pharmacol. Res.*, 2007, **30**, 701-707 (*isol, pmr, cmr*)

Oubatchensine O-138


C₁₉H₁₇NO₆ 355.346

Secodibenzopyrrocoline. Alkaloid from the bark of *Cryptocarya oubatchensis*. Yellow-brown amorph. powder.

Toribio, A. *et al.*, *Org. Lett.*, 2006, **8**, 3825-3828 (*isol, pmr, cmr, ms*)

Ouregidione

[107633-70-5]

O-139

C₁₉H₁₅NO₅ 337.331

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae) and stem bark of *Artabotrys zeylanicus* (Annonaceae). Mp 280° (262-264°).

N-Me: **3-Methoxycepharadione B**. 1,2,3-Trimethoxy-4,5-dioxo-6a,7-dehydroa-porphine

[104380-50-9]

C₂₀H₁₇NO₅ 351.358

Alkaloid from the bark of *Pseuduvaria macrophylla* and *Mitrephora cf. maingayi* (Annonaceae). Yellow needles (MeOH). Mp 198-201°.

8-Methoxy: **8-Methoxyouregidione**

[180128-37-4]

C₂₀H₁₇NO₆ 367.357

Alkaloid from stem bark of *Artabotrys zeylanicus*. Orange powder (MeOH). Mp 276-277°.

Cortes, D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 878 (*isol, uv, ir, pmr, ms, struct*)

Mahmood, K. *et al.*, *Phytochemistry*, 1986, **25**, 1509-1510 (*N-Me, isol, uv, ir, pmr, ms, struct*)


Wijeratne, E.M.K. *et al.*, *Phytochemistry*, 1996, **42**, 1703 (*8-Methoxyouregidione*)

Lee, N.H.S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1158-1159 (*N-Me, isol*)

Hoarau, C. *et al.*, *Eur. J. Org. Chem.*, 2001, 2559-2567 (*N-Me, synth*)

Oourouparine**O-140**

*8,13-Dihydro-1-(methoxycarbonyl)-7H-benz[*g*]indolo[2,3-*a*]quinolizin-6-ium(1+)*, 9CI. 16-Carboxy-3,4,14,15,16,17,18,19,20,21-decadehydrohimbanium methyl ester, 9CI. De-methoxylstoniline [23594-96-9]



C₂₁H₁₇N₂O₂⁺ 329.377

Alkaloid from *Uncaria gambier*

(*Oourouparia gambier*) and *Gelsemium sempervirens*. Yellowish powder. Mp > 280°. λ_{max} 242 ; 256 ; 284 ; 369 (MeOH).

Iodide: [18111-07-4]

C₂₁H₁₇IN₂O₂ 456.282

Orange needles (EtOH). Mp > 300° dec.

Perchlorate: [81362-36-9]

C₂₁H₁₇ClN₂O₆ 428.828

Mp 285-290° dec. (synthetic).

Picrate: [27655-44-3]

Mp 267°.

11-Methoxy: **Alstoniline**

[6714-03-0]

C₂₂H₁₈N₂O₃ 358.396

Alkaloid from the bark of *Alstonia constricta*. Fine yellow-brown needles or fine yellow-brown needles + 1H₂O (MeOH aq.). Mp 372° (356°) dec. Autoxidises readily to the *N*-oxide, Mp 212-213°.

11-Methoxy, hydrochloride:

Fine red needles + 1H₂O (MeOH). No def. Mp.

11-Methoxy, picrate:

Red plates. Mp 294° dec.

Elderfield, R.C. *et al.*, *J.O.C.*, 1942, **7**, 573-580; 1954, **19**, 683-692; 693-697 (*Alstoniline, isol, uv, struct*)

Ban, Y. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1296-1301 (*Alstoniline, uv, ir, synth*)

Taylor, W.I. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. D.* 1966, **262**, 1141-1143 (*uv, struct*)

Merlini, L. *et al.*, *Gazz. Chim. Ital.*, 1967, **97**, 1915-1920 (*synth*)


Beisler, J.A. *et al.*, *Tetrahedron*, 1970, **26**, 1961-1965 (*synth*)

Frostell, E. *et al.*, *Acta Chem. Scand., Ser. B*, 1981, **35**, 671-672 (*synth*)

Kogure, N. *et al.*, *J. Nat. Med. (Tokyo)*, 2007, **61**, 208-212 (*isol, synth, pmr, cmr*)

Ouvrardiane A**O-141**

[951765-87-0]



C₂₈H₃₇NO₇ 499.603

Alkaloid from the roots of *Aconitum ouvrardianum*. Amorph. powder. [α]_D²⁰ -62 (c, 0.3 in CHCl₃).

15-O-Deacyl, 15-benzoyl: **Ouvrardiane B**

[951765-88-1]


C₃₀H₃₃NO₇ 519.593

Alkaloid from the roots of *Aconitum ouvrardianum*. Amorph. powder. [α]_D²⁰ -17.6 (c, 0.2 in CHCl₃).

Hou, L.-H. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 1090-1092 (*isol, pmr, cmr*)

Ovigeridimerine**O-142**

[178765-67-8]




C₃₇H₂₈N₂O₉ 644.636

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Light brownish prisms (MeOH). Mp 207-210°. [α]_D²² +197 (c, 0.12 in CHCl₃).

Chen, J.-J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (*isol, uv, ir, pmr, ms, struct*)

Ovigerine**O-143**

6,7,7a,8-Tetrahydro-5H-bis[1,3]benzodioxolo[6,5,4-de:4',5'-g]quinoline, 9CI. 1,2:10,11-Bismethylenedioxy-noraporphine. Hernovine†

*(S)-form*

C₁₈H₁₅NO₄ 309.321

(S)-form [6410-87-3]

Alkaloid from *Hernandia ovigera*, *Hernandia jamaicensis*, *Hernandia catalpifolia*, *Hernandia cordigera* and *Hernandia peltata* (Hernandiaceae). Shows anti-platelet aggregation activity. Amorph. Mp 300° dec. (as hydrochloride). [α]_D +177 (H₂O).

N-Formyl: **N-Formylovigerine**

[176679-15-5]

C₁₉H₁₅NO₅ 337.331

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Prisms + 1/3H₂O (MeOH). Mp 105-107°. [α]_D²⁴ +321 (c, 0.11 in CHCl₃). Exists in soln. as a mixt. of amide bond rotamers.

N-Me: **1,2:10,11-Bis(methylenedioxy)aporphine. N-Methylovigerine**

[13087-94-0]

C₁₉H₁₇NO₄ 323.348

Alkaloid from *Lindera oldhamii*, *Hernandia ovigera*, *Hernandia jamaicensis* and *Hernandia catalpifolia* (Hernandiaceae). Amorph., cryst. (as hydrobromide). Mp 243-245° dec. (hydrobromide). [α]_D²⁸ +161 (c, 0.4 in MeOH).

N,N-Di-Me:

C₂₀H₂₁NO₄⁺ 339.39

Mp 252-253° dec. (as iodide).

N-Hydroxy: **N-Hydroxyovigerine**

[176679-17-7]

C₁₈H₁₅NO₅ 325.32

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Greyish prisms + 3/4H₂O (CHCl₃/MeOH). Mp 102-104°.

$[\alpha]_D^{24} +184$ (c, 0.10 in CHCl_3).

6a,7-Didehydro, N-formyl: N-Formyldehydroovigerine

$\text{C}_{19}\text{H}_{13}\text{NO}_5$ 335.315

Alkaloid from the trunk bark of *Hernandia nymphaeifolia* (Hernandaceae). Prisms (MeOH). Mp 176-178°. λ_{max} 204 (log ϵ 4.52); 228 (log ϵ 4.28); 255 (sh) (log ϵ 4.26); 263 (log ϵ 4.34); 314 (log ϵ 3.73); 343 (log ϵ 3.66); 377 (log ϵ 3.27); 398 (log ϵ 3.28) (EtOH).

3-Methoxy: Oduocine

[155944-21-1]

$\text{C}_{19}\text{H}_{17}\text{NO}_5$ 339.347

Alkaloid from roots of *Lindera myrrha* (Lauraceae). Needles (H_2O) (as hydrochloride). Mp 220° (hydrochloride). $[\alpha]_D^{20} +140$ (c, 0.30 in MeOH).

(±)-form

N-Me: [30048-26-1]

Synthetic. Amorph.

N,N-Di-Me: [30048-27-2]

$\text{C}_{20}\text{H}_{20}\text{NO}_5^{\oplus}$ 338.382

Cryst. (MeOH) (as iodide). Mp 241-243° dec. (iodide). CAS no. refers to iodide.

Cava, M.P. *et al.*, *Tet. Lett.*, 1966, 1577; 4279 (uv, pmr, struct, deriv)

Cava, M.P. *et al.*, *Tetrahedron*, 1970, **26**, 4649; 1971, **27**, 2639 (deriv, synth)

Lu, S.-T. *et al.*, *Yakugaku Zasshi*, 1972, **92**, 910; *CA*, **77**, 101949m (deriv)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 80 (cd)

Lavault, M. *et al.*, *Planta Med.*, 1981, **42**, 50; 1982, **46**, 119 (isol)

Ba Hung Phan, *et al.*, *Phytochemistry*, 1994, **35**, 1363 (Oduocine)

Chen, I.-S. *et al.*, *Planta Med.*, 1995, **61**, 537-539 (Ovigerine, activity)


Chen, I.-S. *et al.*, *Heterocycles*, 1996, **43**, 799 (N-Formylovigerine, N-Hydroxyovigerine)

Chen, I.-S. *et al.*, *Planta Med.*, 1997, **63**, 154-157 (N-Formyldehydroovigerine)

Ovihernangerine

[187530-46-7]

O-144



$\text{R}^1, \text{R}^2 = \text{---CH}_2\text{---}$

$\text{R}^3 = \text{H}$

$\text{R}^4 = \text{Me}$

$\text{C}_{37}\text{H}_{30}\text{N}_2\text{O}_9$ 646.652

Minor alkaloid from trunk bark of *Hernandia nymphaeifolia*. Prisms (MeOH). Mp 194-196°. $[\alpha]_D^{24} +310$ (c, 0.08 in CHCl_3); λ_{max} 228 (log ϵ 4.84); 272 (log ϵ 4.49); 311 (log ϵ 4.2) (EtOH). λ_{max} 228 (ϵ 69180); 272 (ϵ 31000); 311 (ϵ 15720) (MeOH) (Berdy).

Chen, J.-J. *et al.*, *Planta Med.*, 1996, **62**, 528 (isol, uv, ir, pmr, ms, struct)

Oviisocorydine

[187669-80-3]

As Ovihernangerine, O-144 with $\text{R}^1 = \text{R}^2 = \text{R}^3 = \text{Me}$, $\text{R}^4 = \text{H}$

$\text{C}_{38}\text{H}_{34}\text{N}_2\text{O}_9$ 662.695

Minor alkaloid from trunk bark of *Hernandia nymphaeifolia*. Prisms (MeOH). Mp 168-170°. $[\alpha]_D^{24} +254$ (c, 0.08 in CHCl_3); λ_{max} 225 (log ϵ 4.9); 271 (log ϵ 4.51); 309 (log ϵ 4.21) (EtOH). λ_{max} 225 (ϵ 79430); 271 (ϵ 32360); 309 (ϵ 16220) (MeOH) (Berdy).


Chen, J.-J. *et al.*, *Planta Med.*, 1996, **62**, 528 (isol, uv, ir, pmr, ms, struct)

7-Oxa-3-azabicyclo[4.1.0]-heptan-2-one, 9CI

3,4-Epoxy-2-piperidone. 3,4-Epoxyvalerolactam. Tedanalactam

[956710-49-9]

[159934-17-5 (3R,4R)-form]



$\text{C}_5\text{H}_7\text{NO}_2$ 113.116

Isol. from the sponge *Tedania ignis* and the leaves of *Piper crassinervum*. Oil. $[\alpha]_D -8.9$ (c, 0.3 in MeOH).

(-)-form

Isol. from the sponge *Tedania ignis*. Oil. $[\alpha]_D -8.9$ (c, 0.3 in MeOH).


Cronan, J.M. *et al.*, *Nat. Prod. Lett.*, 1994, **5**, 85-88; 141-146 (isol, ir, pmr, cmr)

Lago, J.H.G. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 910-914 (isol, pmr, cmr)

Oxachelin

[853013-90-8]

O-147



$\text{C}_{27}\text{H}_{37}\text{N}_7\text{O}_{11}$ 635.63


Prod. by *Streptomyces* sp. GW9/1258. Siderophore. Powder. λ_{max} 203; 245; 249; 255 (sh); 304 (MeCN aq.).

Sontag, B. *et al.*, *J. Antibiot.*, 2006, **59**, 659-663 (isol, pmr, cmr)

Oxalicine A

[125309-96-8]

O-148



$\text{C}_{30}\text{H}_{33}\text{NO}_6$ 503.594

Prod. by *Penicillium oxalicum* and *Penicillium thiersii*. Mp 193-198° (189-194°). $[\alpha]_D +142$ (c, 0.3 in CH_2Cl_2); λ_{max} 232 (ϵ 17800); 329 (ϵ 6920) (MeOH) (Derep).

23-Hydroxy: Oxalicine B

$\text{C}_{30}\text{H}_{33}\text{NO}_7$ 519.593

Prod. by *Penicillium thiersii*. Antiinsectan. Cryst. Mp 143-146°. $[\alpha]_D +92$ (c, 0.1 in CH_2Cl_2); λ_{max} 232 (ϵ 17400); 271 (ϵ 6400); 317 (ϵ 4000) (MeOH).

15-Deoxy: 15-Deoxyoxalicine A

$\text{C}_{30}\text{H}_{33}\text{NO}_5$ 487.594

Prod. by *Penicillium thiersii*. Oil. $[\alpha]_D +54$ (c, 0.1 in CH_2Cl_2); λ_{max} 229 (ϵ 10200); 271 (ϵ 3900); 316 (ϵ 2200) (MeOH).

15-Deoxy, 23-hydroxy: 15-Deoxyoxalicine B

$\text{C}_{30}\text{H}_{33}\text{NO}_6$ 503.594

Prod. by *Penicillium decaturense*. Cryst. (MeCN aq.). Mp 165-167°. $[\alpha]_D +34$ (c, 1 in CH_2Cl_2); λ_{max} 205 (ϵ 16000); 235 (ϵ 14000); 335 (ϵ 5600) (MeOH).

Ubillas, R. *et al.*, *Chem. Comm.*, 1989, 1618-1619 (isol, uv, ir, pmr, cmr, cryst struct)


Zhang, Y. *et al.*, *Org. Lett.*, 2003, **5**, 773-776 (15-Deoxyoxalicine B)

Li, C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 319-322 (isol, pmr, cmr)

Oxandrine

[104778-24-7]

O-149



$\text{C}_{37}\text{H}_{38}\text{N}_2\text{O}_7$ 622.716

Alkaloid from the bark of *Pseudoxandra* aff. *lucida* (Annonaceae). Amorph. $[\alpha]_D -11$ (c, 0.9 in CHCl_3). $[\alpha]_D +9$ (c, 0.96 in MeOH).

12'-Me ether: Oxandrinine

[104777-91-5]


$\text{C}_{38}\text{H}_{40}\text{N}_2\text{O}_7$ 636.743

Alkaloid from *Pseudoxandra* aff. *lucida* (Annonaceae). Amorph. $[\alpha]_D +60$ (c, 0.5 in CHCl_3).

Cortes, D. *et al.*, *Can. J. Chem.*, 1986, **64**, 1390 (isol, uv, ir, pmr, cmr, ms, cd, struct, deriv)

Oxasetin

O-150



$\text{C}_{21}\text{H}_{29}\text{NO}_4$ 359.464


Tetramic acid deriv. Related to Equisetin,

E-142. Prod. by *Vaginatispora aquatica* HK1821. Antibacterial agent. Oil. $[\alpha]_D^{25}$ -147.5 (c, 0.37 in MeOH). λ_{\max} 225 ; 261 ; 345 (MeOH).

He, H. *et al.*, *J. Antibiot.*, 2002, **55**, 821-825 (isol, ir, pmr, cmr)

Oxathinecarboxylic acid O-151

6-Hydroxy-2-methyl-5-(1-oxopropyl)-2H-1,4-oxazine-3-carboxylic acid, 9CI. *P* 725C. Antibiotic *P* 725C [87979-96-2]



C₉H₁₁NO₅ 213.19

Isol. from *Penicillium thomii* IFO7984. Inhibits prostaglandin synthetase. Anti-inflammatory and analgesic. Cryst. Sol. EtOAc, MeOH; fairly sol. H₂O. Mp 115-120° dec. λ_{\max} 289 (E1%/1cm 452); 352 (E1%/1cm 382) (MeOH) (Berdy). λ_{\max} 283 (E1%/1cm 475) (MeOH/HCl) (Berdy). λ_{\max} 289 (E1%/1cm 463); 352 (E1%/1cm 390) (MeOH/NaOH) (Berdy).

▶ RP6198010


Ca salt:

Cryst. + 4H₂O. $[\alpha]_D^{20}$ +550 (c, 0.1 in MeOH).

Japan. Pat., 1983, 83 134 085; *CA*, **100**, 4785 (isol)

2H-1,4-Oxazine-3-carboxylic acid O-152

Codipiloic acid [136945-75-0]




C₅H₅NO₃ 127.099

Constit. of *Codonopsis pilosula*.

Wang, H. *et al.*, *Zhongcaoyao*, 1991, **22**, 195; *CA*, **115**, 214619

Oxazinine 1 O-153



Absolute Configuration

C₂₂H₂₁N₃O₄ 391.426

Stereochem. revised in 2006. Isol. from the digestive glands of *Mytilus galloprovincialis*. Cytotoxic. $[\alpha]_D^{25}$ +9 (MeOH). λ_{\max} 216 (€ 46000); 270 (€ 5700); 277 (€ 5500); 286 (€ 3500) (MeOH).

O-De(cyanoethyl): **Oxazinine 2**

C₁₉H₁₈N₂O₄ 338.362

Isol. from *Mytilus galloprovincialis*. $[\alpha]_D^{25}$ +8.7 (MeOH). λ_{\max} 216 (€ 20400); 275 (€ 3200); 377 (€ 700) (MeOH).

6-Epimer: **Oxazinine 4**

C₂₂H₂₁N₃O₄ 391.426

Isol. from *Mytilus galloprovincialis*.


$[\alpha]_D^{25}$ +60 (c, 0.2 in MeOH).

Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2001, 49-53 (isol, pmr, cmr, ms)

Ciminiello, P. *et al.*, *Tetrahedron*, 2006, **62**, 7738-7743 (Oxazinine 4, abs config)

Oxazinine 3 O-154

5-[(4-Hydroxyphenyl)methyl]-2-(1H-indol-3-yl)-3-morpholinone, 9CI [331836-05-6]



Absolute Configuration

C₁₉H₁₈N₂O₃ 322.363

Isol. from the digestive glands of *Mytilus galloprovincialis*. $[\alpha]_D^{25}$ +12 (MeOH). λ_{\max} 220 (€ 20600); 270 (€ 3100); 375 (€ 750) (MeOH).

7R-Hydroxy, 4'-O-(2-cyanoethyl): **Oxazinine 6**

[1001439-32-2]

C₂₂H₂₁N₃O₄ 391.426

Isol. from *Mytilus galloprovincialis*.

2-Epimer, 7R-hydroxy, 4'-O-(2-cyanoethyl): **Oxazinine 5**

[1001439-31-1]

C₂₂H₂₁N₃O₄ 391.426

Isol. from *Mytilus galloprovincialis*.


Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2001, 49-53 (Oxazinin 3, isol, pmr, cmr)

Couladouros, E.A. *et al.*, *Tet. Lett.*, 2004, **45**, 7779-7781 (Oxazinin 3, synth, abs config)

Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2007, 5434-5439 (Oxazinins 5,6)

Oxazolomycin O-155

Resistaphyllin. Diffusomycin [89808-64-0]



C₃₅H₄₉N₃O₉ 655.787

Polyene-type antibiotic. Numbering systems vary. Prod. by *Streptomyces* sp. Active against gram-positive bacteria and P388 leukaemia. Oil. $[\alpha]_D^{18}$ +56.1 (c, 1 in MeOH). Identity of Oxazolomycin with Resistaphyllin not conclusively established. λ_{\max} 230 (€ 33000); 267 (sh) (€ 28000); 275 (€ 34000); 285 (sh) (€ 27000) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 10.6 mg/kg.

(6'E)-Isomer: **Oxazolomycin C**

[208342-70-5]

C₃₅H₄₉N₃O₉ 655.787

Prod. by *Streptomyces albus* JA3453.

$[\alpha]_D^{22}$ +12.7 (c, 0.4 in MeOH). λ_{\max} 230 (€ 18200); 265 (€ 10600); 275 (€ 13400); 285 (€ 11000) (MeOH).

(4'E,6'E)-Isomer: **Oxazolomycin B**

[208342-69-2]

C₃₅H₄₉N₃O₉ 655.787

Prod. by *Streptomyces albus* JA3453.

$[\alpha]_D^{22}$ -3.7 (c, 0.3 in MeOH). λ_{\max} 230 (€ 18100); 265 (€ 12400); 275 (€ 15700); 285 (€ 12800) (MeOH).

13'-Methyl, 16-(methoxymethyl): **Triedimycin A**

[135094-12-1]

C₃₈H₅₅N₃O₁₀ 713.867

Prod. by a *Streptomyces* sp. Antitumour agent. Powder. Sol. MeOH, Me₂CO, EtOAc, CHCl₃; poorly sol. C₆H₆, H₂O. Mp 70-80° (grad. dec.). $[\alpha]_D^{21}$ +71 (c, 1 in MeOH). λ_{\max} 230 (E1%/1cm 670); 276 (E1%/1cm 668) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 6.25-12.5 mg/kg. HE1780000

16R-Methyl: **16R-Methyloxazolomycin**.

Antibiotic KSM 2690B. KSM 2690B

C₃₆H₅₁N₃O₉ 669.814

Prod. by *Streptomyces* sp. KSM-2690. Pale yellow powder. Four other KSM 2690 components were isol. but not characterised. λ_{\max} 230 ; 267 (sh) ; 278 ; 288 (sh) (no solvent reported).

16R-Methyl, 6'E-isomer: **16R-Methyloxazolomycin C**. Antibiotic KSM 2690C.

KSM 2690C

C₃₆H₅₁N₃O₉ 669.814

Prod. by *Streptomyces* sp. KSM-2690. Pale yellow powder.

16S-Methyl: **16S-Methyloxazolomycin**

[202863-86-3]

C₃₆H₅₁N₃O₉ 669.814

Prod. by a *Streptomyces* sp. Cytotoxic agent. Pale yellow powder. $[\alpha]_D^{23}$ +3.6 (c, 0.5 in MeOH). λ_{\max} 228 (€ 23000); 266 (€ 27200); 275 (€ 32000); 285 (€ 27000) (MeOH).

13',16-Dimethyl: **Triedimycin B**

[135094-13-2]

C₃₇H₅₃N₃O₉ 683.84

Prod. by a *Streptomyces* sp. Antitumour agent. Sol. MeOH, CHCl₃, Me₂CO, EtOAc; poorly sol. C₆H₆, H₂O. Mp 85-95° (grad. dec.). $[\alpha]_D^{23}$ +75 (c, 0.8 in MeOH). λ_{\max} 226 (€ 39300); 266 (sh) (€ 32900); 275 (€ 39300); 284 (sh) (€ 30300) (MeOH) (Derep).

▶ HE1785000

[89808-66-2]

Takahashi, K. *et al.*, *CA*, 1984, **100**, 167719 (props, cryst struct)

Mori, T. *et al.*, *Tet. Lett.*, 1985, **26**, 1073-1076 (struct, nmr)

Kawai, S. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 1127 (isol, props)

Ikeda, Y. *et al.*, *J. Antibiot.*, 1991, **44**, 453 (Triedimycin)

Graefe, U. *et al.*, *Annalen*, 1992, 429-432 (biosynth)


Ryu, G. *et al.*, *J. Antibiot.*, 1997, **50**, 1064-1066; 1999, **52**, 193-197 (16-Methyloxazolomycin)

Kanzaki, H. *et al.*, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 438-442 (Oxazolomycin B, Oxazolomycin C)

Otani, T. *et al.*, *J. Antibiot.*, 2000, **53**, 1397-1400 (KSM 2690 derivs)

4-(4-Oxazolyl)-3-(hydroxymethyl)-3-buten-2-ol O-156

2-[(4-Oxazolyl)methylene]-1,3-butane-diol. **Antibiotic MR 93B**. MR 93B [192212-11-6]



C₈H₁₁NO₃ 169.18

Metab. of *Trichoderma harzianum*. Tyrosinase inhibitor, melanin biosynth. inhibitor. Oil. [α]_D²⁵ -4 (c, 0.1 in MeOH). Related to Melanoxidin, M-159. λ_{max} 230 (ε 15140) (MeOH).

l'-Aldehyde: 3-Hydroxy-2-(4-oxazolylmethylene)butanal, 9CI. 2-(1-Hydroxyethyl)-3-(4-oxazolyl)-2-butenal.

Melanoxazol

[178275-98-4]

C₈H₉NO₃ 167.164


Prod. by *Trichoderma* sp. ATF-451. Melanin biosynth. inhibitor. Active against mushroom tyrosinase. Powder. Sol. MeOH, CHCl₃, EtOH, Me₂CO, EtOAc; poorly sol. H₂O, hexane. Mp 70-73°. [α]_D²⁴ +38.9 (c, 0.1 in MeOH). λ_{max} 274 (ε 7100) (MeOH).

Takahashi, S. *et al.*, *J. Antibiot.*, 1996, **49**, 513-518 (*Melanoxazol*)

Lee, C.H. *et al.*, *J. Antibiot.*, 1997, **50**, 469-473; 474-478 (MR 93B, *isol.*, *uv.*, *ir.*, *pmr.*, *cmr*)

3-(5-Oxazolyl)-1H-indole, 9CI O-157

5-(3-Indolyl)oxazole. **Antibiotic SF 2583B**. SF 2583B [120191-50-6]



C₁₁H₈N₂O 184.197

Prod. by *Streptomyces* sp. SF 2583. Antiparasitic and nematocidal agent. Powder. Sol. MeOH, EtOAc. Mp 170-172°. λ_{max} 222; 265; 280 (MeOH).

N-Me: [848488-06-2]

C₁₂H₁₀N₂O 198.224

Waxy solid.

N-Benzyl: [848488-08-4]

C₁₈H₁₄N₂O 274.321

Orange cryst. (petrol/CH₂Cl₂). Mp 106-108°.

N-(4-Methylbenzenesulfonyl): [848488-10-8]

C₁₈H₁₄N₂O₃S 338.386

Pale yellow cryst. (petrol/CH₂Cl₂). Mp 144°.

4-Chloro: 3-(4-Chloro-5-oxazolyl)-1H-indole, 9CI. 4-Chloro-5-(3-indolyl)ox-

azole. **Antibiotic SF 2583A**. SF 2583A [120191-51-7]

C₁₁H₇ClN₂O 218.642

Prod. by *Streptomyces* sp. SF 2583.


Active against *Caenorhabditis elegans*.

Watabe, H. *et al.*, *Meiji Seika Kenkyu Nenpo*, 1988, 55 (*isol.*, *4'-chloro*)

Chakrabarty, M. *et al.*, *Tetrahedron*, 2005, **61**, 1793-1801 (*synth.*, *1-Me*, *1-benzyl*, *1-tosyl*)

Oxepinamide A

O-158



Relative Configuration

R = CH(CH₃)CH₂CH₃,
R' = OH

C₁₇H₂₁N₃O₅ 347.37

Isol. from an *Acremonium* sp. obt. from the tunicate *Ecteinascidia turbinata*. Antiinflammatory agent. Yellow oil. [α]_D +43 (c, 0.001 in CHCl₃). λ_{max} 250 (ε 6100); 345 (ε 4900) (MeOH).

11-Epimer: **Oxepinamide B**

C₁₇H₂₁N₃O₅ 347.37

Isol. from an *Acremonium* sp. obt. from *Ecteinascidia turbinata*. Yellow oil. [α]_D +52 (c, 0.001 in CHCl₃). λ_{max} 252 (ε 4900); 347 (ε 2600) (MeOH).

Belofsky, G.N. *et al.*, *Chem. Eur. J.*, 2000, **6**, 1355-1360

Oxepinamide C

O-159

As Oxepinamide A, O-158 with R = CH₂CH(CH₃)₂, R' = OMe

C₁₈H₂₃N₃O₅ 361.397


Isol. from an *Acremonium* sp. obt. from *Ecteinascidia turbinata*. Yellow oil. [α]_D -35 (c, 0.001 in CHCl₃). λ_{max} 250 (ε 3000); 345 (ε 2100) (MeOH).

Belofsky, G.N. *et al.*, *Eur. J. Org. Chem.*, 2000, **6**, 1355-1360

Oxerine

O-160

6,7-Dihydro-7-methyl-5H-2-pyridine-5,7-diol, 9CI [137181-67-0]



C₉H₁₁NO₂ 165.191

Absolute configuration is not certain. Alkaloid from the aerial parts of *Oxera morieri* (Verbenaceae). Amorph. solid. [α]_D²⁰ -11 (c, 0.2 in MeOH).

Benkrief, R. *et al.*, *Planta Med.*, 1991, **57**, 79-80 (*isol.*, *uv.*, *ir.*, *pmr.*, *ms.*, *struct*)

Aoyagi, Y. *et al.*, *Tetrahedron*, 1994, **50**, 13575-13582 (*synth*)


Jones, K. *et al.*, *Tetrahedron*, 2000, **56**, 397-406 (*synth*)

Ohba, M. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 63-67 (*synth.*, *bibl*)

6,7-Oxidodeoxynupharidine dimer

O-161

[41758-68-3]



C₃₀H₄₂N₂O₄ 494.673


Alkaloid from the rhizomes of *Nuphar luteum* subsp. *macrophyllum* (Nymphaeaceae). Mp 165-172°. [α]_D²⁵ -93 (c, 0.26 in CH₂Cl₂).

LaLonde, R.T. *et al.*, *J.A.C.S.*, 1972, **94**, 8522 (*isol.*, *ir.*, *ord.*, *pmr.*, *ms.*, *struct*)

3,6-Oxidoiboxygaine

O-162

[77808-84-5]



Absolute Configuration

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the trunk bark of *Tabernanthe pubescens* (Apocynaceae). Cryst. (MeOH). Mp 226°. [α]_D -5.6 (c, 0.53 in CHCl₃).

Deoxy: **3,6-Oxidoibogaine**

[77795-11-0]


C₂₀H₂₄N₂O₂ 324.422

Alkaloid from root bark of *Tabernanthe pubescens* (Apocynaceae). Amorph. [α]_D -18.7 (c, 0.309 in CHCl₃).

Mulamba, T. *et al.*, *J. Nat. Prod.*, 1981, **44**, 184 (*isol.*, *uv.*, *ir.*, *pmr.*, *ms.*, *struct*)

5,15-Oxidolycopodane

O-163




C₁₆H₂₅NO 247.38

A component of Alkaloids L28 and L31. Alkaloid from *Lycopodium annotinum* var. *acrifolium* (Lycopodiaceae). [α]_D +51 (c, 0.019 in MeOH).

Ayer, W.A. *et al.*, *Can. J. Chem.*, 1990, **68**, 1300 (*pmr.*, *cmr.*, *ms.*, *struct*)

Oximidine II

[220578-36-9]

Absolute
ConfigurationC₂₃H₂₄N₂O₆ 424.452

Prod. by *Pseudomonas* sp. Q52002. Antitumour agent. Powder. Mp 113-115°. [α]_D²⁵ -141 (c, 0.75 in MeOH). λ_{\max} 282 (ε 30300) (MeOH). λ_{\max} 203 (ε 138000); 279 (ε 28200) (0.01 M NaOH/MeOH).

12 α ,13 α -Epoxide: Oximidine I

[220625-58-1]

C₂₃H₂₄N₂O₇ 440.452

Prod. by *Pseudomonas* sp. Q52002. Antitumour agent. Powder. Mp 104-106°. [α]_D²² -286 (c, 1 in MeOH). λ_{\max} 272 (ε 30500) (MeOH). λ_{\max} 203 (ε 112000); 270 (ε 30500) (0.1 M NaOH/MeOH).

17E-Isomer, 14-deoxy, 12 α ,13 α -epoxide:**Oximidine III**C₂₃H₂₄N₂O₆ 424.452

Prod. by *Pseudomonas* sp. QN05727. Antitumour agent. Powder. Mp 87-89°. [α]_D²² -80 (c, 0.03 in MeOH). λ_{\max} 210 (ε 23400); 272 (ε 28000) (0.1 M NaOH/MeOH). λ_{\max} 276 (ε 26600) (MeOH).


Kim, J.W. *et al.*, *J.O.C.*, 1999, **64**, 153-155

(Oximidines I-II)

Hayakawa, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 889-904; 905-908 (Oximidine III)**3-Oxiranyl-1H-indole, 9CI**

3-Indolylethylene oxide

O-165


C₁₀H₉NO 159.187**(-)-form [129225-30-5]**

Alkaloid from leaves of *Moricandia arvensis* (Brassicaceae). [α]_D²² -40 (c, 1 in H₂O).

Belkhir, A. *et al.*, *Phytochemistry*, 1990, **29**, 1315 (isol, uv, pmr, ms, struct)**Oxobaicaline**

[105418-66-4]

O-166


C₂₀H₁₅NO₆ 365.342

Alkaloid from the stems of *Thalictrum baicalense* (Ranunculaceae). Large needles (MeOH). Mp 240° dec.

Maekh, S.Kh. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 251; *Chem. Nat. Compd. (Engl. Transl.)*, 238 (isol, uv, pmr, struct)**8-(2-Oxobutyl)dihydrochelerythrine**

O-167

8-(Propionylmethyl)dihydrochelerythrine [71627-29-7]


C₂₅H₂₅NO₅ 419.476

Probably an artifact. Alkaloid from the bark of *Fagara mayu*. Also obt. by heating Chelerythrine chloride in 2-butanone containing Na₂CO₃ (Rutaceae). Cryst. (EtOAc). Mp 206-208.5°.

Assem, E.M. *et al.*, *Phytochemistry*, 1979, **18**, 511-512 (isol, ir, pmr, ms, struct, synth)**Oxobuxifoline**

O-168

4,10-Dimethoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI [89368-27-4]


C₁₉H₁₃NO₅ 335.315

Minor alkaloid from the stem bark of *Duguetia obovata* (Annonaceae). Also isol. from stem bark of *Artabotrys zeylanicus* (Annonaceae) and from leaves and stems of *Cissampelos glaberrima*. Cryst. (MeOH). Mp 268°.

Roblot, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 862 (isol, uv, ir, pmr, ms, struct)Wijeratne, E.M.K. *et al.*, *Phytochemistry*, 1996, **42**, 1703 (isol)Barbosa-Filho, J.M. *et al.*, *Phytochemistry*, 1997, **44**, 959 (isol, cmr)**Oxocompostelline**

O-169

6-Methoxy-13H-[1,3]dioxolo[7,8][1]-benzoxepino[2,3,4-ij]isoquinolin-13-one, 9CI [86817-61-0]

C₁₈H₁₁NO₅ 321.289

Compostelline is currently unknown. Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Yellow cryst. (EtOH). Mp 259°.

O-De-Me: Oxocularicine

[133084-02-3]


C₁₇H₉NO₅ 307.262

Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. The trivial name is somewhat misleading (Cularicine, C-804 is 1,2,3,4-tetrahydro).

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2295 (uv, ir, pmr, ms, struct, synth)Allais, D.P. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1280 (Oxocularicine)**Oxocrebaine**

O-170

9,10-Dimethoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. 1,2-Methylenedioxy-8,9-dimethoxyoxoaporphine. 8,9-Dimethoxylirodentine [38826-42-5]

C₁₉H₁₃NO₅ 335.315

Obt. by air oxidn. of Dehydrocrebaine in C-733 in the presence of an alkali catalyst, and from uthongine by chromatography on silica gel. Alkaloid from *Stephania sasakii* and from the stem-xylem of *Hernandia ovigera*. Tentatively identified in bark of *Uvariopsis guineensis*. Orange prisms (EtOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 265-269° (260-265°). λ_{\max} 214 ; 238 ; 245 ; 375 (MeOH/NaOH) (Berdy).

N-Me: Uthongine

[81255-04-1]

C₂₀H₁₆NO₅⁺ 350.35

Quaternary alkaloid from the tuberous roots of *Stephania venosa* (*Stephania rotunda*). Partially decomposed by chromatography on silica gel to Oxocrebaine.

9-O-De-Me: Fissiceine

[330832-88-7]

C₁₈H₁₁NO₅ 321.289


Alkaloid from *Fissistigma glaucescens*. Red needles (CHCl₃). Mp 238-240°. λ_{\max} 261 (log ε 4.5); 292 (log ε 4.13); 361 (log ε 3.99) (EtOH).

Leboeuf, M. *et al.*, *Phytochemistry*, 1972, **11**, 2833 (isol)Yang, T.H. *et al.*, *T'ai-wan Yao Hsueh Tsa Chih*, 1973, **25**, 8; *CA*, **86**, 127150e (isol, struct)Kunitomo, J. *et al.*, *Phytochemistry*, 1980, **19**, 2735 (synth, uv, ir, pmr, ms)Guinaudeau, H. *et al.*, *Chem. Comm.*, 1981, 1118 (Uthongine, synth)Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1981, **101**, 431 (isol, uv, ir, pmr, ms)Lo, W.-L. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 1251-1256 (Fissiceine)

Oxocularine

O-171

6,9,10-Trimethoxy-12H-[1]benzoxepino[2,3,4-ij]isoquinolin-12-one, 9CI [3395-23-1]



$C_{19}H_{15}NO_5$ 337.331

The trivial name is somewhat misleading (Cularine, C-805 is 1,2,3,4-tetrahydro). Alkaloid from *Corydalis claviculata* (Papaveraceae). Yellow cryst. (EtOH). Mp 198-199°.

O⁷-De-Me: Oxocularidine

[120481-42-7]

$C_{18}H_{13}NO_5$ 323.304

Alkaloid from *Sarcocapnos crassifolia* (Papaveraceae). Amorph. yellow powder. λ_{\max} 254 (log ϵ 3.86); 266 (log ϵ 3.67); 304 (sh) (log ϵ 3.28); 350 (log ϵ 3.18); 410 (log ϵ 3.2) (EtOH).

Kametani, T. *et al.*, *Yakugaku Zasshi*, 1965, **85**, 667; *CA*, **63**, 13229c (*synth*, *uv*, *ir*)

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1983, **46**, 881 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2295 (*pmr*, *ms*)


Castedo, L. *et al.*, *Phytochemistry*, 1989, **28**, 251 (*Oxocularidine*)

Garcia, A. *et al.*, *Tetrahedron*, 1995, **51**, 8585 (*synth*)

17-Oxocycloprotobuxine

O-172

[231628-57-2]



$C_{23}H_{37}NO$ 343.551


Misleading name assigned. Alkaloid from the roots of *Buxus sempervirens*. Gum. $[\alpha]_D^{20}$ +48 (c, 0.73 in $CHCl_3$). λ_{\max} 203 (log ϵ 1.85) (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1999, **62**, 665-669 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Oxocyclostylidol

O-173

[910450-17-8]



$C_{11}H_{10}BrN_5O_3$ 340.136

Related to Cyclooroidin, C-887. Isol. from *Stylissa caribica*. Yellow powder. $[\alpha]_D^{25}$ -12 (c, 1 in MeOH). λ_{\max} 224; 263; 320 (MeOH).

Grube, A. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1212-1214 (*isol*, *cd*, *pmr*, *cmr*)

6-Oxo-2,4-decadienoic acid

O-174

$H_3C(CH_2)_3COCH=CHCH=CHCOOH$

$C_{10}H_{14}O_3$ 182.219

(2E,4E)-form

Piperidine: 1-(1,6-Dioxo-2,4-decadienyl)-piperidine. 6-Oxo-2,4-decadienoic acid piperidine. N-(6-Oxo-2,4-decadienyl)-piperidine

$C_{15}H_{23}NO_2$ 249.352


Alkaloid from the roots of *Piper nigrum*. Oil. λ_{\max} 275 (log ϵ 4.32) (MeOH).

Wei, K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1005-1009 (*isol*, *pmr*, *cmr*)

Oxodesmosine

O-175

[147044-49-3]



$C_{23}H_{37}N_5O_7$ 495.575

Cross-linking amino acid. Isol. from bovine aorta elastin.

Suyama, K. *et al.*, *Bioorg. Med. Chem. Lett.*, 1992, **2**, 1767-1770 (*isol*, *pmr*)

**12-Oxo-2,4,8,10-dodecate-
traenoic acid**

O-176

11-Formyl-2,4,8,10-undecatetraenoic acid

$OHCCH=CHCH=CHCH_2CH_2CH=CHCH=CHCOOH$

$C_{12}H_{14}O_3$ 206.241

(all-E)-form

2-Methylpropylamide: N-Isobutyl-12-oxo-2,4,8,10-dodecatetraenamide. **Lanyuamide V** [350811-07-3]

$C_{16}H_{23}NO_2$ 261.363


Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. λ_{\max} 259 (sh) (log ϵ 4.23); 275 (log ϵ 4.3) (EtOH).

Tsai, I.-L. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 830-833

8-Oxoepiberberine

O-177

8-Oxodihydroepiberberine [19716-60-0]



$C_{20}H_{17}NO_5$ 351.358


Alkaloid from the rhizomes of *Coptis japonica*. Powder. λ_{\max} 226; 256; 331; 345; 377 (MeOH).

Min, Y.D. *et al.*, *Arch. Pharmacol. Res.*, 2006, **29**, 757-761 (*isol*, *pmr*, *cmr*)

Oxoglaucine

O-178

1,2,9,10-Tetramethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI. Liriodendron base. O-Methylatheroline [5574-24-3]



$C_{20}H_{17}NO_5$ 351.358

Alkaloid from *Liriodendron tulipifera* (Magnoliaceae), *Annona purpurea* (sontocoya) (Annonaceae) and trunk wood of *Phoebe cinnamomifolia* (Lauraceae); also from the Papaveraceae and Menispermaceae. Shows antimicrobial activity. Mp 227-229° (195-197°).

O¹-De-Me, N¹-Me, inner salt: **Corunnine. Glauvine** [34421-18-6]

[34421-18-6]

$C_{20}H_{17}NO_5$ 351.358

Alkaloid from *Glauicum flavum* var. *vestitum*, the discoloured sapwood of *Liriodendron tulipifera* and the roots of *Thalictrum foetidum* (Papaveraceae, Magnoliaceae, Ranunculaceae). Violet needles (EtOH) (also descr. as green). Mp 255-257°.

O¹-De-Me, N¹-Me, perchlorate:

Purple needles. Mp 293-295°.

O⁹-De-Me: **Atheroline**

[5140-35-2]

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from *Atherosperma moschatum*, *Dryadodaphne novoguineensis* and *Guatteria* spp. (Monimiaceae, Annonaceae). Mp 250-260° dec. λ_{\max} 252 (ϵ 50100); 299 (ϵ 39800); 323 (ϵ 39800); 400 (ϵ 15800) (EtOH/KOH) (Derep). λ_{\max} 242 (ϵ 39800); 272 (ϵ 39800); 292 (ϵ 25100); 352 (ϵ 12600); 380 (ϵ 10000); 433 (ϵ 5010) (EtOH) (Derep).

▶RB6035000

*O*⁹-De-Me, Ac:


Yellow needles (Py). Mp 190-195°.

- Buchanan, M.A. *et al.*, *J.O.C.*, 1960, **25**, 1389 (isol, uv)
 Cohen, J. *et al.*, *J.O.C.*, 1961, **26**, 4143 (isol, uv, synth)
 Bick, I.R.C. *et al.*, *Tet. Lett.*, 1965, 2399 (Atheroline, isol, ir, pmr, uv)
 Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1403 (ms)
 Tomita, M. *et al.*, *Yakugaku Zasshi*, 1968, **88**, 1143 (isol, pmr)
 Ribas, I. *et al.*, *Tet. Lett.*, 1971, 3093; 1973, 3617 (Corunnine, uv, ir, pmr, struct, synth)
 Cava, M.P. *et al.*, *J.O.C.*, 1972, **37**, 2936 (Atheroline, synth, ir, pmr, uv)
 Yakhontova, L.D. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 214; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 212 (Corunnine, uv, pmr)
 Kupchan, S.M. *et al.*, *Chem. Comm.*, 1973, 915 (Corunnine, synth)
 Castedo, L. *et al.*, *Heterocycles*, 1975, **3**, 449; 1980, **14**, 1131 (Corunnine, synth, struct)
 Chen, C.-L. *et al.*, *Phytochemistry*, 1976, **15**, 1161 (Corunnine, isol, uv, pmr)
 Kametani, T. *et al.*, *Tetrahedron*, 1977, **33**, 1069 (Atheroline, synth)
 Hufford, C.D. *et al.*, *J. Pharm. Sci.*, 1980, **69**, 1180-1182 (activity)
 Marsaioli, A.J. *et al.*, *Phytochemistry*, 1980, **19**, 995 (cmr)
 Saá, C. *et al.*, *Tet. Lett.*, 1985, **26**, 4559 (synth)
 Martinez, E. *et al.*, *Planta Med.*, 1988, **54**, 361 (isol, uv, ir, pmr, ms)
 Atanes, N. *et al.*, *J.O.C.*, 1991, **56**, 2984 (synth)

7-Oxohernandaline

O-179

[187530-48-9]

C₂₈H₂₃NO₈ 501.492

Minor alkaloid from trunk bark of *Hernandia nymphaeifolia* and from *Hernandia ovigera*. Yellowish prisms (EtOH). Mp 197-199°. λ_{max} 210 (log ε 4.65); 239 (log ε 4.57); 273 (log ε 4.51); 317 (log ε 4.14); 375 (sh) (log ε 3.86); 415 (sh) (log ε 3.68) (EtOH). λ_{max} 210 (ε 44668); 239 (ε 37150); 273 (ε 32360); 317 (ε 13800); 375; 415 (MeOH) (Berdy).

3-Methoxy: 9-(2-Formyl-4,5-dimethoxyphenoxy)-1,2,3,10-tetramethoxyoxoaporphine

C₂₉H₂₅NO₉ 531.518

Alkaloid from *Thalictrum elegans*. Orange-yellow powder (MeOH). Mp 136-138°.

4-Methoxy: 4-Methoxyoxohernandaline [178765-69-0]

C₂₉H₂₅NO₉ 531.518

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Orange-reddish prisms (CHCl₃/MeOH). Mp 255-259°.

Chen, J.J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (4-Methoxyoxohernandaline)


Chen, J.-J. *et al.*, *Planta Med.*, 1996, **62**, 528 (Oxohernandaline)

Liang, Z.Y. *et al.*, *Chin. J. Chem.*, 2005, **23**, 895-897 (3-methoxy)

7-Oxohernangerine

O-180

[172274-07-6]

C₁₈H₁₁NO₅ 321.289

Alkaloid from stem bark of *Hernandia nymphaeifolia*. Orange prisms (MeOH). Mp 256-258°.

Ac: [172274-09-8]

Yellowish needles (CHCl₃/MeOH). Mp 213-215°.

Me ether: O-Methyl-7-oxohernangerine

C₁₉H₁₃NO₅ 335.315

Alkaloid from the trunk bark of *Hernandia nymphaeifolia*. Orange-yellow prisms (CH₂Cl₂/Me₂CO). Mp 236-238°. λ_{max} 222 (log ε 4.72); 255 (log ε 4.56); 268 (sh) (log ε 4.5); 310 (sh) (log ε 3.99); 358 (log ε 4.13); 409 (log ε 4.12) (EtOH).

3-Methoxy: 3-Methoxy-7-oxohernangerine. Lindechunine A

C₁₉H₁₃NO₆ 351.315

Alkaloid from the roots of *Lindera chinii*. Yellow powder. λ_{max} 220; 265; 365 (MeOH).

Chen, J.-J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 156-158 (isol, uv, ir, pmr, ms, struct)

Chen, J.-J. *et al.*, *Planta Med.*, 2001, **67**, 593-598 (O-Methyl-7-oxohernangerine)

Zhang, C.-F. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1195-1200 (Lindechunine A)

6-Oxo-2,4-hexadienoic acid

O-181

OHCC=CH=CHCOOH

C₆H₆O₃ 126.112

(2E,4E)-form

2-Phenylethylamide: [96917-24-7]

C₁₄H₁₅NO₂ 229.278

Constit. of *Salmea scandens*. Oil.

[88973-46-0]


Bohlmann, F. *et al.*, *Phytochemistry*, 1985, **24**, 595-596 (isol, amide)

α-Oxo-1H-indole-3-acetic acid, 9CI

O-182

2-(3-Indolyl)glyoxylic acid

[1477-49-2]

C₁₀H₇NO₃ 189.17

Yellow cryst. Mp 215° dec.

Me ester: [18372-22-0]

C₁₁H₉NO₃ 203.197

Alkaloid from a *Spongosorites* sp. Cryst. (MeOH). Mp 230°.

Me ester, N-Ac: [66946-85-8]

C₁₃H₁₁NO₄ 245.234

Mp 130-132°.

Et ester: [51079-10-8]

C₁₂H₁₁NO₃ 217.224

Cryst. Mp 186°.

Chloride: [22980-09-2]

C₁₀H₆ClNO₂ 207.616

Mp 138-139°.

Amide: α-Oxo-1H-indole-3-acetamide

[5548-10-7]

C₁₀H₈N₂O₂ 188.185

Alkaloid from a *Spongosorites* sp. Mp 248° Mp 252° dec.

Nitrile: α-Oxo-1H-indole-3-acetonitrile.

3-(Cyanocarbonyl)indole

[19194-62-8]

C₁₀H₆N₂O 170.17

Mp 224-226°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 668D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 140A (nmr)

Baker, J.W. *et al.*, *J.C.S.*, 1940, 458 (synth)

Behr, D. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 2411-2414 (Et ester)


Janosik, T. *et al.*, *Tetrahedron*, 2002, **58**, 2813-2819 (nitrile)

Bao, B. *et al.*, *Mar. Drugs*, 2007, **5**, 31-39 (Me ester, amide, isol, pmr, cmr, ms)

Oxoisocalycine

O-183

10-Hydroxy-12-methoxy-8H-benzo[g]-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. 9-Hydroxy-11-methoxy-1,2-methylenedioxyoxoaporphine [91174-08-2]

C₁₈H₁₁NO₅ 321.289

Alkaloid from the stem bark of *Guatteria discolor*. Cryst. (MeOH). Mp 280° dec.

9-Me ether: Oxodiscoguttine

[921936-13-2]

C₁₉H₁₃NO₅ 335.315

Alkaloid from the stems of *Fissistigma oldhamii*. Amorph. red powder. λ_{max} 230 (log ε 4.62); 265 (log ε 4.45); 330 (log ε 4.06); 356 (log ε 4.03) (MeOH).

O¹¹-De-Me. 9-Me ether: Oxocalycine

[921936-14-3]

C₁₈H₁₁NO₅ 321.289


Alkaloid from the stems of *Fissistigma oldhamii*. Amorph. red powder. λ_{max} 225 (log ε 4.5); 267 (log ε 4.28); 356

(log ϵ 4.01) (MeOH).

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 353-362 (*Oxoisocalycinine*)
 Zhang, Y.-N. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 988-996 (*Oxocalycinine*, *Oxodiscoguatine*)

5-Oxo-2(5H)-isoxazoleacetic acid, 9CI O-184

2-Carboxymethyl-3-isoxazolin-5-one
 [51581-01-2]

C₅H₅NO₄ 143.099Constit. of *Lathyrus odoratus*.

v. Rompuy, L. *et al.*, *Biochem. Biophys. Res. Commun.*, 1974, **56**, 199

5-Oxo-2(5H)-isoxazolepropanoic acid, 9CI O-185

2-(2-Carboxyethyl)-3-isoxazolin-5-one
 [92279-63-5]

C₆H₇NO₄ 157.126

Alkaloid from the immature seeds and pods of *Lathyrus odoratus* (Fabaceae). Needles (Me₂CO aq.). Mp not recorded.

Nitrile: 5-Oxo-2(5H)-isoxazolepropanoic nitrile, 9CI. 2-(2-Cyanoethyl)-3-isoxazolin-5-one

[51580-99-5]

C₆H₆N₂O₂ 138.126


Constit. of *Lathyrus odoratus* and *Pisum sativum* (peas).

v. Rompuy, L. *et al.*, *J.C.S. Perkin 1*, 1973, 2503 (*synth*)

Ikegami, F. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 2450 (*isol, uv, ir, pmr, ms, struct*)

Oxojanussine

O-186


C₃₀H₃₂N₄O₃ 496.608

Isol. from *Strychnos johnsonii*. May be an oxidn. prod. of the Janussines (see Janussine A, J-22).

Massiot, G. *et al.*, *Tet. Lett.*, 1985, **26**, 2441-2444 (*pmr, struct*)

Oxolaureline

10-Methoxyliriodenine. *Lauterine*
 [28200-65-9]

C₁₈H₁₁NO₄ 305.289

Alkaloid from the stem bark of *Guatteria elata* and *Laurelia novae-zelandiae* and from *Magnolia soulangeana* (Annonaceae, Monimiaceae, Magnoliaceae). Also isol. from roots of *Miliusa cf. banacea* (Annonaceae). Yellow needles (CHCl₃). Mp 301-303° dec. λ_{\max} 247; 309; 343; 405 (MeOH) (Berdy).

O-De-Me: 10-Hydroxyliriodenine. Demethylloxolaureline. Demethyllauterine
 [154659-89-9]

C₁₇H₉NO₄ 291.262

Alkaloid from roots of *Miliusa cf. banacea* and from *Polyalthia* sp. (Annonaceae). Topoisomerase inhibitor. Orange amorph. solid. λ_{\max} 246; 269; 280; 354; 420 (MeOH).

Govindachari, T.R. *et al.*, *Indian J. Chem.*, 1970, **8**, 475 (*synth, uv*)

Ziyayev, R. *et al.*, *Khim. Priir. Soedin.*, 1975, **11**, 528; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 560 (*isol*)

Urzúa, A. *et al.*, *Heterocycles*, 1976, **4**, 1881 (*isol, ir, pmr*)

Hsu, C.C. *et al.*, *J. Nat. Prod.*, 1977, **40**, 152 (*isol, uv, ir, pmr, ms*)


Harrigan, G.G. *et al.*, *J. Nat. Prod.*, 1994, **57**, 68 (*10-Hydroxyliriodenine*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1033 (*10-Hydroxyliriodenine*)

17-Oxolupanine

O-188

2,17-Dioxosparteine
 [4697-83-0]

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Chamaecytisus* spp., *Lupinus polyphyllus*, *Lupinus angustifolius* and other *Lupinus* spp. (Fabaceae). Cryst. (petrol). Mp 155.5°. [α]_D²³ +139.

13- ξ -Hydroxy: 13-Hydroxy-17-oxolupanine

[152406-63-8]

C₁₅H₂₂N₂O₃ 278.35

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

13-Angeloyloxy: 13-Angeloyloxy-17-oxolupanine

[152406-64-9]

C₂₀H₂₈N₂O₄ 360.452**O-187**

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

13-Tigloyloxy: 13-Tigloyloxy-17-oxolupanine

[152406-65-0]

C₂₀H₂₈N₂O₄ 360.452

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

13-Benzoyloxy: 13-Benzoyloxy-17-oxolupanine

[152406-66-1]

C₂₂H₂₆N₂O₄ 382.458

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

13-(E)-Cinnamoyloxy: trans-13-Cinnamoyloxy-17-oxolupanine

[152406-67-2]

C₂₄H₂₈N₂O₄ 408.496

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

13-(Z)-Cinnamoyloxy: cis-13-Cinnamoyloxy-17-oxolupanine

[152434-59-8]

C₂₄H₂₈N₂O₄ 408.496

Minor alkaloid in leaves of *Lupinus linearis* and *Lupinus gibertianus* (Fabaceae). Tentative identification by glc/ms.

15 β -Hydroxy: 15-Hydroxy-17-oxolupanine

[162229-97-2]

C₁₅H₂₂N₂O₃ 278.35

Alkaloid from seeds of *Lupinus albus* (Fabaceae). Fine needles. Mp 85-87°. [α]_D²⁵ +4.5 (c, 0.15 in CH₂Cl₂).

Bratele, M.D. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1959, **33**, 1187 (*isol*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1960, **93**, 1956 (*isol*)

Schumann, D. *et al.*, *Monatsh. Chem.*, 1968, **99**, 390 (*ms*)

Cho, Y.D. *et al.*, *Anal. Biochem.*, 1971, **44**, 49 (*glc, ms*)

Daily, A. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 889 (*occur*)

Kinghorn, A.D. *et al.*, *Phytochemistry*, 1980, **19**, 1705 (*occur*)

Hatzold, T. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 934 (*isol*)

Planchuelo-Ravelo, A. *et al.*, *Z. Naturforsch., C*, 1993, **48**, 702 (*13-Hydroxy-17-oxolupanine, esters*)

Mohamed, M.H. *et al.*, *Phytochemistry*, 1994, **37**, 1751 (*15 β -Hydroxy-17-oxolupanine*)

5-Oxo-6,8-octadecadienoic acid O-189

H₃C(CH₂)₈CH=CHCH=CHCO(CH₂)₃COOH

C₁₈H₃₀O₃ 294.433**(E,E)-form**

Constit. of the tubers of *Lepidium meyenii*. Gum. λ_{\max} 224 (log ϵ 3.7); 274 (log ϵ 3.49) (MeOH).

Benzylamide: N-Benzyl-5-oxo-6,8-octa-

decadienamide. **Macamide A**

C₂₅H₃₇NO₂ 383.573

Alkaloid from the tubers of *Lepidium meyenii* (maca). Gum. λ_{max} 210 (log ε 4.08); 276 (log ε 3.99) (MeOH).

Muhammad, I. et al., *Phytochemistry*, 2002, **59**, 105-110 (*isol*, *pmr*, *cmr*)

9-Oxo-12,15-octadecadienoic acid O-190

H₃CCH₂CH=CHCH₂CH=CHCH₂CH₂CO(CH₂)₇COOH

C₁₈H₃₀O₃ 294.433

(12Z,15Z)-form

Benzylamide: N-Benzyl-9-oxo-12,15-octadecadienamide

C₂₅H₃₇NO₂ 383.573

Alkaloid from the tubers of *Lepidium meyenii*. Powder. λ_{max} 210 (log ε 3.96); 274 (log ε 3.18) (MeOH).

Zhao, J.-P. et al., *J. Agric. Food Chem.*, 2005, **53**, 690-693 (*isol*, *pmr*, *cmr*)

13-Oxo-9,11-octadecadienoic acid O-191

[31385-09-8]

[26474-39-5, 79790-32-2, 119051-89-7]

H₃C(CH₂)₄COCH=CHCH=CH(CH₂)₇COOH

C₁₈H₃₀O₃ 294.433

(9E,11E)-form [29623-29-8]

Constit. of the processed leaves of *Artemisia argyi*. Also from *Monnina emarginata*. Powder. λ_{max} 276 (log ε 4.21) (MeOH).

2,3-Dihydroxypropyl ester:

C₂₁H₃₆O₅ 368.512

Isol. from the mushroom *Clitocybe clavipes*. Oil.

Benzylamide: N-Benzyl-13-oxo-9,11-octadecadienamide

C₂₅H₃₇NO₂ 383.573

Alkaloid from the tubers of *Lepidium meyenii* (maca). Gum. λ_{max} 208 (log ε 4.02); 276 (log ε 4.04) (MeOH).

(9Z,11E)-form [54739-30-9]

Constit. of the processed leaves of *Artemisia argyi*. Also from the alga *Gracilariaopsis lemaneiformis*. Powder. λ_{max} 278 (log ε 4.15) (MeOH).

Phillips, B.E. et al., *Biochim. Biophys. Acta*, 1970, **210**, 353 (*isol*, *uv*, *ir*, *pmr*)

Jiang, Z.D. et al., *Phytochemistry*, 1991, **30**, 1187 (*isol*)

Yoshikawa, M. et al., *Chem. Pharm. Bull.*, 1996, **44**, 1656 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Kawagishi, H. et al., *J. Nat. Prod.*, 2002, **65**, 1712-1714 (*dihydroxypropyl ester*)

Zhao, J.-P. et al., *J. Agric. Food Chem.*, 2005, **53**, 690-693 (*benzylamide*)

12-Oxooctadecanoic acid O-192

12-Oxostearic acid. **Lycaonic acid** [925-44-0]

H₃C(CH₂)₅CO(CH₂)₁₀COOH

C₁₈H₃₄O₃ 298.465

Constit. of *Gracilaria verrucosa*. Mp 81.5°.

Semicarbazone: Mp 122°.

Me ester: [2380-27-0]

C₁₉H₃₆O₃ 312.492

Mp 46-48°. Bp_{0.8} 178-180°.

Pyrrolidide: **Lycaonic acid pyrrolidide**. 1-(1,12-Dioxooctadecyl)pyrrolidine

[83182-47-2]

C₂₂H₄₁NO₂ 351.571

Alkaloid from the roots of *Achillea* spp. (Asteraceae). Mp 44°.

Piperidide: **Lycaonic acid piperidide**. 1-(1,12-Dioxooctadecyl)piperidine

[83182-45-0]

C₂₃H₄₃NO₂ 365.598

Alkaloid from the roots of *Achillea lycaonica* (Asteraceae). Mp 41°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 1105C (*nmr*)

Belopol'skii, M.P. et al., *CA*, 1937, **31**, 7397 (*synth*)

Tulloch, A.P. et al., *Can. J. Chem.*, 1977, **55**, 1135 (*cmr*)

Tulloch, A.P. et al., *Lipids*, 1977, **12**, 92 (*synth*)

Greger, H. et al., *Phytochemistry*, 1982, **21**, 1071 (*isol*, *ir*, *pmr*, *struct*, *pyrrolidide*, *piperidide*)

Bohlmann, F. et al., *Tetrahedron*, 1983, **39**, 123 (*synth*)

Dasaradlin, L. et al., *J.C.S. Perkin 2*, 1995, 221 (*Me ester*, *synth*)

Dang, H.T. et al., *J. Nat. Prod.*, 2008, **71**, 232-240 (*isol*, *ms*)

9-Oxo-12-octadecenoic acid O-193

H₃C(CH₂)₄CH=CHCH₂CH₂CO(CH₂)₇COOH

C₁₈H₃₂O₃ 296.449

(Z)-form

Benzylamide: N-Benzyl-9-oxo-12-octadecenoamide

C₂₅H₃₉NO₂ 385.589

Alkaloid from the tubers of *Lepidium meyenii*. Powder. λ_{max} 212 (log ε 3.88); 274 (log ε 2.98) (MeOH).

Zhao, J.-P. et al., *J. Agric. Food Chem.*, 2005, **53**, 690-693 (*isol*, *pmr*, *cmr*)

12-Oxo-2-octadecenoic acid O-194

2,3-Dehydrolycaonic acid

H₃C(CH₂)₅CO(CH₂)₈CH=CHCOOH

C₁₈H₃₂O₃ 296.449

(E)-form

Pyrrolidide: **2,3-Dehydrolycaonic acid pyrrolidide**. 1-(1,12-Dioxo-2-octadecenyl)pyrrolidine

[83182-46-1]

C₂₂H₃₉NO₂ 349.556

Alkaloid from the roots of *Achillea lycaonica* (Asteraceae). Mp 64°.

Piperidide: **2,3-Dehydrolycaonic acid piperidide**. 1-(1,12-Dioxo-2-octadecenyl)piperidine

[83182-44-9]

C₂₃H₄₁NO₂ 363.582

Alkaloid from the roots of *Achillea lycaonica* (Asteraceae). Mp 43°.


Greger, H. et al., *Phytochemistry*, 1982, **21**, 1071 (*isol*, *ir*, *pmr*, *ms*, *struct*)

Bohlmann, F. et al., *Tetrahedron*, 1983, **39**, 123 (*synth*)

Oxophoebine

O-195

1,2,3-Trimethoxy-7H-benzo[de][1,3]-benzodioxolo[5,6-g]quinolin-7-one, 9CI. 1,2,3-Trimethoxy-9,10-methylenedioxyoxoaporphine [109175-37-3]



C₂₀H₁₅NO₆ 365.342

Alkaloid from the leaves of *Phoebe valeriana* and from *Xylopi aethiopica* (Lauraceae). Amorph. red solid. λ_{max} 211; 240; 274; 379; 477 (MeOH) (Berdy).

3-Demethoxy: 1,2-Dimethoxy-9,10-methylenedioxyoxoaporphine. **Oxonantenine**

[15358-38-0]

C₁₉H₁₃NO₅ 335.315

Alkaloid from *Cassitha racemosa*, *Corydalis bulbosa*, *Nandina domestica*, *Laurelia sempervirens* (Peruvian nutmeg), *Siparuna gilgiana* and *Hernandia nymphaefolia* (Lauraceae, Papaveraceae, Nandinaceae, Monimiaceae, Hernandiaceae). Shows antimicrobial activity. Glistening yellow needles (EtOH). Mp 215-218° dec Mp 225-227° dec Mp 297-299°. λ_{max} 228; 242; 270; 285; 318; 360 (MeOH) (Berdy).

1,3-Bis(demethoxy): **Hernanymphine**

[172924-21-9]

C₁₈H₁₁NO₄ 305.289

Alkaloid from stem bark of *Hernandia nymphaefolia*. Yellowish prisms (CHCl₃/MeOH). Mp 263-265°. Rare substitution pattern (no oxygenation at C-1).

Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1457 (*Oxonantenine*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Hufford, C.D. et al., *J. Pharm. Sci.*, 1980, **69**, 1180-1182 (*Oxonantenine*, *activity*)

Chiu, S.Y.C. et al., *J. Nat. Prod.*, 1982, **45**, 229 (*Oxonantenine*, *isol*, *uv*, *ir*, *pmr*, *ms*)


Castro, O. et al., *J. Nat. Prod.*, 1986, **49**, 1036 (*Oxophoebine*, *isol*, *uv*, *pmr*, *ms*, *struct*)

Harrigan, G.G. et al., *J. Nat. Prod.*, 1994, **57**, 68-73 (*Oxophoebine*, *isol*)

Chen, J.J. et al., *J. Nat. Prod.*, 1996, **59**, 156 (*Hernanymphine*)

4-Oxo-2-piperidinecarboxylic acid, 9CI O-196

4-Piperidone-2-carboxylic acid. 4-Oxopipelicolic acid. 4-Ketopipelicolic acid [190005-81-3]



(S)-form

C₆H₉NO₃ 143.142

Exists predominantly as the covalent hydrate in aq. soln.

(S)-form*L*-form

[65060-18-6]

Amino acid present in hydrolysates of the *Ostreogrycin* A, O-131 family of cyclo-depsipeptide antibiotics. Prisms + 1H₂O (2-propanol). Mp 240° dec. [α]_D²³ -19.2 (c, 0.5 in H₂O).*Hydrochloride*: [116049-23-1]

[166043-02-3]

Plates (2-propanol). Mp 204° (191°). [α]_D²² +4.1 (c, 0.55 in H₂O).*N*-tert-*Butyloxycarbonyl*: [198646-60-5]C₁₁H₁₇NO₅ 243.259Mp 127° dec. [α]_D²⁵ -19 (c, 1 in CHCl₃).*N*-*Benzyloxycarbonyl*: [3956-66-9]C₁₄H₁₅NO₅ 277.276Bright yellow oil. [α]_D²⁵ -19.6 (c, 1 in CHCl₃).**(±)-form***Hydrochloride*: [119945-10-7]

[99979-55-2]

Cryst. (Et₂O). Mp 139-142° dec.*N*-*Me*, *Me ester*: [86251-51-6]C₈H₁₃NO₃ 171.196


Oil.

[180675-07-4, 189516-49-2]

Eastwood, F.W. *et al.*, *J.C.S.*, 1960, 2286-2292 (*isol*)Clark-Lewis, J.W. *et al.*, *J.C.S.*, 1961, 189-201 (*synth*)Essawy, M.Y. *et al.*, *J. Het. Chem.*, 1983, **20**, 477-478 (*N*-*Me Me ester*, *synth*, *pmr*)Hartmann, P. *et al.*, *Synth. Commun.*, 1988, **18**, 553-557 (*synth*, *pmr*)Machetti, F. *et al.*, *Tet. Lett.*, 1996, **37**, 4205-4208 (*synth*)Golubev, A.S. *et al.*, *Tetrahedron*, 1996, **52**, 14757-14776 (*synth*, *pmr*)Badorrey, R. *et al.*, *Tet. Lett.*, 1997, **38**, 2547-2550 (*synth*)Bonsquet, Y. *et al.*, *Tetrahedron*, 1997, **53**, 15671-15680 (*N*-*tert*-*butyloxycarbonyl*)*Org. Synth.*, *Coll. Vol.*, 9, 1998, 526-530 (*synth*, *ir*, *pmr*)Machetti, F. *et al.*, *Tetrahedron*, 2001, **57**, 4995-4998 (*synth*)Jung, J.-C. *et al.*, *Tetrahedron: Asymmetry*, 2006, **17**, 2479-2486 (*S*-*form*, *benzyloxycarbonyl*)**8-[(2-Oxo-3-piperidinyl)amino]-no]-8-oxooctyl 5,9-anhydro-2,3,8-trideoxy-8-(5-hydroxy-4-methyl-2-hexenyl)-3-methyl-DL-glycero-LD-allo-non-2-enoate, 9CI**

O-197

[147024-92-8]


C₃₀H₅₀N₂O₉ 582.733Constit. of the marine bacterium *Alteromonas rava*. Antimicrobial agent. Oil. [α]_D -1.8 (c, 0.003 in MeOH). Related to Pseudomonic acid A. λ_{max} 220 (ε 14000) (MeOH) (Berdy).*4*-*Deoxy*: BP 2. *NSC* 646282

[147024-96-2]

C₃₀H₅₀N₂O₈ 566.734Constit. of *Alteromonas* sp. Antimicrobial agent. Oil. [α]_D -3.1 (c, 0.005 in MeOH). λ_{max} 220 (ε 14000) (MeOH) (Berdy).Stierle, D.B. *et al.*, *Experientia*, 1992, **48**, 1165 (*isol*, *pmr*, *cmr*)Shiozawa, H. *et al.*, *Annu. Rep. Sankyo Res. Lab.*, 1999, **51**, 45-72 (*rev*)**Oxopropaline D**

[152752-59-5]

O-198

C₁₅H₁₄N₂O₃ 270.287

Abs. config. revised in 2003.

(R)-formProd. by *Streptomyces* sp. G324. Cytotoxic agent. Pale yellow powder. Sol. MeOH, MeCN; poorly sol. hexane, H₂O. Mp 182-183° (synthetic). [α]_D²⁰ +30 (c, 0.1 in MeOH). λ_{max} 219 (ε 31000); 264 (ε 10600); 287 (ε 16300); 381 (ε 6200) (MeOH).*2*'-*O*-*α*-*L*-*Rhamnopyranoside*: **Oxopropaline A**

[152752-57-3]

C₂₁H₂₄N₂O₇ 416.43Prod. by *Streptomyces* sp. G324. Pale yellow powder. [α]_D²⁰ +56 (c, 0.05 in MeOH). λ_{max} 220 (ε 43800); 265 (ε 15500); 288 (ε 22800); 381 (ε 9600) (MeOH).*3*'-*O*-*α*-*L*-*Rhamnopyranoside*: **Oxopropaline B**

[152752-58-4]

C₂₁H₂₄N₂O₇ 416.43Prod. by *Streptomyces* sp. G324. Pale yellow powder. Sol. MeOH, MeCN; poorly sol. hexane, H₂O. [α]_D²⁰ -39 (c, 0.02 in MeOH). λ_{max} 218 (ε 44900); 265 (ε 15100); 287 (ε 22300); 378 (ε 8000) (MeOH).*2*'-*Deoxy*: **Oxopropaline G**

[152752-61-9]


C₁₅H₁₄N₂O₂ 254.288Prod. by *Streptomyces* sp. G324. Shows cytotoxic activity. Pale yellow powder. Sol. MeOH, MeCN; poorly sol. hexane, H₂O. Mp 155-157° (synthetic). λ_{max} 218 (ε 33700); 263 (ε 12500); 284 (ε 18900); 375 (ε 7200) (MeOH).*2*'-*Deoxy*, *3*'-*O*-*α*-*L*-*rhamnopyranoside*:**Oxopropaline E**

[152752-60-8]

C₂₁H₂₄N₂O₆ 400.43Prod. by *Streptomyces* sp. G324. Pale yellow powder. [α]_D²⁰ -39 (c, 0.02 in MeOH). λ_{max} 218 (ε 32800); 262 (ε 11000); 284 (ε 16800); 377 (ε 6300) (MeOH).Abe, N. *et al.*, *J. Antibiot.*, 1993, **46**, 1672-1677; 1678-1686Choshi, T. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 108-113; 2003, **51**, 20-23 (*synth*, *abs config*, *activity*)**3-(2-Oxopropyl)coronaridine**

O-199

[39967-45-8]



Absolute Configuration

C₂₄H₃₀N₂O₃ 394.513Alkaloid from *Ervatamia coronaria* and *Tabernaemontana dichotoma* (Apocynaceae).*10*-*Methoxy*: **3-(2-Oxopropyl)voacangine**

[145130-77-4]

C₂₅H₃₂N₂O₄ 424.539Minor alkaloid from roots of *Tabernaemontana pauciflora*. Powder. Mp 68-70°. [α]_D²⁰ -20 (c, 0.16 in EtOH). Poss. artifact.*19R*-*Hydroxy*: **3-Ketopropyl-19R-heyneanine**


[99326-72-4]

C₂₄H₃₀N₂O₄ 410.512Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Probably an artifact. λ_{max} 226; 286; 293 (EtOH) (Berdy).*Δ*¹-*Isomer*, *7α*-*hydroxy*: **7-Hydroxy-Δ**¹-**3-(2-oxopropyl)coronaridine**

[909120-05-4]

C₂₄H₃₀N₂O₄ 410.512Alkaloid from the roots of *Ervatamia hainanensis*. Called 9*S*-hydroxy-Δ¹⁶ in the lit.Delle Monache, G. *et al.*, *Atti Accad. Naz. Lincei. Cl. Sci. Fis., Mat. Nat., Rend.*, 1972, **52**, 375; *CA*, **78**, 40468a (3-(2-Oxopropyl)coronaridine)Perera, P. *et al.*, *Phytochemistry*, 1985, **24**, 2097-2104 (3-Ketopropyl-19R-heyneanine)Okuyama, E. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 2075-2079 (2-Oxopropylvoacangine)Huang, J.P. *et al.*, *Chin. Chem. Lett.*, 2006, **17**, 779-782 (7-Hydroxy-Δ¹-oxopropylcoronaridine)**5-(2-Oxopropyl)coronaridine**

O-200

*(5R)*-*form*


C₂₄H₃₀N₂O₃ 394.513**(5R)-form** [145106-97-4]

Minor alkaloid from roots of *Tabernaemontana pauciflora*. Powder. Mp 52-57°. [α]_D²⁰ -60 (c, 0.12 in EtOH). Poss. artifact. λ_{\max} 225 (log ϵ 4.5); 286 (log ϵ 3.9); 293 (log ϵ 3.87) (EtOH).

(5S)-form [145106-98-5]

Minor alkaloid from roots of *Tabernaemontana pauciflora*. Powder. Mp 169-171°. [α]_D²⁰ -72 (c, 0.051 in EtOH). Poss. artifact. λ_{\max} 225 (log ϵ 4.53); 283 (log ϵ 3.9); 291 (log ϵ 3.85) (EtOH).

Delle Monache, G. et al., *Atti Accad. Naz. Lincei, Cl. Sci. Fis., Mat. Nat., Rend.*, 1972, **52**, 375-380; *CA*, **78**, 40468a (isol)
Okuyama, E. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2075-2079 (isol, uv, ir, pmr, cmr)

3,3'-(Oxopropyl)diconaridine O-201C₄₅H₅₄N₄O₅ 730.945


Minor alkaloid from roots of *Tabernaemontana pauciflora*. Powder. Mp 136-138°. [α]₄₀₀²⁶ +23 (c, 0.13 in EtOH). Poss. artifact. CAS name and MF are defective.

[145106-99-6]

Okuyama, E. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2075-2079

8-Oxopseudopalmatine O-202

5,6-Dihydro-2,3,10,11-tetramethoxy-8H-dibenzo[a,g]quinolizin-8-one, 9CI. 8-Oxypseudopalmatine [10211-78-6]

C₂₁H₂₁NO₅ 367.401

Alkaloid from the roots of *Stephania suberosa* (Menispermaceae). Orange cryst. (MeOH). Mp 198-199° (187-188°, 192-194°).

13,13'-Dimer: Illicifoline

[177481-46-8]

C₄₂H₄₀N₂O₁₀ 732.785

Alkaloid from roots of *Berberis ilicifolia*. Yellow cryst. Mp 260°.

O²-De-Me: Cerasonine

[186424-51-1]

C₂₀H₁₉NO₅ 353.374

Alkaloid from stem bark of *Polyalthia cerasoides*. Amorph. solid. Mp > 200°. λ_{\max} 229 (log ϵ 3.5); 260 (log ϵ 3.42); 333 (log ϵ 3.35) (EtOH).

O²,O¹⁰-Di-de-Me: Cerasodine

[186424-27-1]

C₁₉H₁₇NO₅ 339.347

Alkaloid from stem bark of *Polyalthia cerasoides*. Oil. λ_{\max} 238 (log ϵ 3.61); 262 (log ϵ 3.62); 336 (log ϵ 3.6) (EtOH).

Lenz, G.R. et al., *J.O.C.*, 1974, **39**, 2839 (synth, uv, ir, pmr)

Ninomiyama, I. et al., *J.C.S. Perkin 1*, 1975, 1720 (synth, ir, pmr)

Kametani, T. et al., *Heterocycles*, 1976, **4**, 927 (synth, uv, ir, pmr)

Trifonov, L.S. et al., *Tet. Lett.*, 1985, 3159 (synth)

Patra, A. et al., *Phytochemistry*, 1987, **26**, 547 (isol, pmr)

Cobas, A. et al., *J.O.C.*, 1992, **57**, 6765 (synth)


Fajardo, V. et al., *Heterocycles*, 1996, **43**, 949 (Illicifoline)

González, M.C. et al., *J. Nat. Prod.*, 1997, **60**, 108 (Cerasodine, Cerasonine)

Le, T.N. et al., *Chem. Pharm. Bull.*, 2008, **56**, 1026-1029 (Cerasonine, synth)

Oxopurpureine O-203

1,2,3,9,10-Pentamethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI. 1,2,3,9,10-Pentamethoxyoxoaporphine [32845-27-5]

C₂₁H₁₉NO₆ 381.384

Alkaloid from *Annona purpurea*, *Liriodendron tulipifera* and *Phoebe cinnamomifolia*. Shows borderline antineoplastic activity. Cryst. + 1/2 toluene; yellow needles (EtOH). Mp 198-202° dec. (192-194°). Log P 2.34 (calc). λ_{\max} 251 ; 282 ; 354 ; 394 ; 456 (MeOH) (Berdy). λ_{\max} 215 (ϵ 3200); 248 (ϵ 2180); 278 (ϵ 3320); 389 (ϵ 640) (EtOH) (Berdy).

9-O-De-Me, 9-O-(2-hydroxy-4,5-dimethoxybenzyl): 9-(2-Hydroxy-4,5-dimethoxybenzyl)-1,2,3,10-tetramethoxyoxoaporphine

C₂₉H₂₇NO₉ 533.534

Alkaloid from *Thalictrum elegans*. Cryst. (MeOH). Mp 140-142°.


Sonnet, P.E. et al., *J. Pharm. Sci.*, 1971, **60**, 1254 (isol, uv, ir, pmr, struct)

Martinez, E. et al., *Planta Med.*, 1988, **54**, 361 (isol, uv, ir, pmr, ms)

Liang, Z.-Y. et al., *Chin. J. Chem.*, 2005, **23**, 895-897 (*Thalictrum elegans* alkaloid)

Oxoputerine

Oxo-O-methylpukateine [64040-72-8]

C₁₈H₁₁NO₄ 305.289

Alkaloid from the trunk wood of *Duguetia eximia*, the stem bark of *Duguetia calycina* and *Guatteria elata* and from the bark of *Laurelia novae-zelandiae* (Annonaceae, Monimiaceae). Shows antimicrobial activity. Orange-yellow needles (CHCl₃). Mp 241-243° dec.

O-De-Me: Oxopukateine. 11-Hydroxyliriodenine

[67951-19-3]

C₁₇H₉NO₄ 291.262

Alkaloid from trunk wood of *Duguetia stelichantha* (Annonaceae). Yellow cryst. (C₆H₆). Mp 255-257°.

3-Methoxy: 3-Methoxyoxoputerine

[158018-14-5]

C₁₉H₁₃NO₅ 335.315

Alkaloid from stem bark of *Guatteria foliosa* (Annonaceae). Amorph.

Hsu, C.C. et al., *J. Nat. Prod.*, 1977, **40**, 152 (isol, uv, ir, pmr, ms, struct)

Gottlieb, O.R. et al., *Phytochemistry*, 1978, **17**, 837 (isol, uv, ir, pmr, ms, struct)

Roblot, F. et al., *Plant. Med. Phytother.*, 1978, **12**, 259; *CA*, **91**, 2517b (isol)

Marsaioli, A.J. et al., *Phytochemistry*, 1980, **19**, 995 (cmr)


Urzúa, A. et al., *Phytochemistry*, 1982, **21**, 773 (isol, ir)

Villar, A. et al., *Planta Med.*, 1986, **52**, 556-557 (Oxoputerine, activity)

Mahiou, V. et al., *J. Nat. Prod.*, 1994, **57**, 890 (3-Methoxyoxoputerine)

2-Oxo-3-(4-pyridinyl)propanoic acid O-205

2-Hydroxy-3-(4-pyridinyl)-2-propenoic acid. (4-Pyridinyl)pyruvic acid. *Ascosonchine*

C₈H₇NO₃ 165.148


Exists mainly as Z-enol form. Prod. by *Ascochyta sonchi*. Selective phytotoxin. Amorph. solid. λ_{\max} 230 (log ϵ 3.66); 305 (log ϵ 3.61) (MeOH).

Evidente, A. et al., *Phytochemistry*, 2004, **65**,

475-480 (isol, pmr, cmr, ms)

5-Oxo-2-pyrrolidinecarboxylic acid **O-206**

5-Oxoproline, 9CI. **Pidolic acid, BAN, INN.** α -Aminoglutaric acid lactam. Glutimic acid lactam. Glutimic acid. Glutiminic acid. Pyroglutamic acid. 2-Pyrrolidone-5-carboxylic acid



(R)-form

C₅H₇NO₃ 129.115

Antihyperlipidaemic agent. Log P -1.27 (calc).

(S)-form

L-form

[98-79-3]

Found in substantial amounts in brain tissue and other tissue in bound form, esp. skin. Also present in plant tissues. Isol. from the tunicate *Botryllus schlosseri*. Used in resolin. of amines. Cryst. (H₂O). Mp 156-157°. [α]_D²⁰ -11.45 (c, 4.44 in H₂O).

Butyl ester: [4931-68-4]

C₉H₁₅NO₃ 185.222Constit. of the bulbs of *Ornithogalum caudatum*.

2-Methylpropyl ester: Isobutyl pidolate [849443-95-4]

C₉H₁₅NO₃ 185.222Constit. of the roots of *Panax ginseng*.**(ξ)-form**

N-(2,3-Dibromo-4,5-dihydroxybenzyl):

C₁₂H₁₁Br₂NO₅ 409.031

Isol. from the red alga *Rhodomela confervoides*. Cryst. (MeOH aq.). Mp 172-174°. [α]_D²⁰ +24 (c, 5.7 in Me₂CO).

N-(2,3-Dibromo-4,5-dihydroxybenzyl), Me ester:

C₁₃H₁₃Br₂NO₅ 423.057

Isol. from *Rhodomela confervoides*. Cryst. (MeOH aq.). Mp 183-185°. [α]_D²⁰ +7 (c, 5 in Me₂CO).

N-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl], Me ester:


C₂₀H₁₈Br₃NO₇ 624.077

Isol. from *Rhodomela confervoides*. Cryst. (MeOH aq.). Mp 196-198°. [α]_D²⁰ -30.9 (c, 9.7 in Me₂CO).

[28874-51-3]

Hardegger, E. et al., *Helv. Chim. Acta*, 1955, **38**, 312-320 (S-form, synth, abs config)Gibian, H. et al., *Annalen*, 1961, **640**, 145-156 (synth, bibl)Orlowski, M. et al., *The Enzymes*, 3rd Ed., Academic Press, N.Y., 1971, **4**, 123-151 (rev)Cervinka, O. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 897-901 (cd)Pattabhi, V. et al., *J.C.S. Perkin 2*, 1974, 1085-1089 (cryst struct)Voelter, W. et al., *Monatsh. Chem.*, 1974, **105**, 1110-1135 (cmr)Taira, Z. et al., *Acta Cryst. B*, 1977, **33**, 3823-3827 (cryst struct)Rigo, B. et al., *J. Het. Chem.*, 1995, **32**, 1489-1492 (occur)Tang, Y. et al., *J. Chin. Pharm. Sci.*, 2001, **10**, 169-171 (butyl ester)Tan, N. et al., *Yunnan Zhiwu Yanjiu*, 2003, **25**, 366-368; *CA*, **142**, 370710m (isobutyl ester, isol)Zhao, J. et al., *J. Nat. Prod.*, 2005, **68**, 691-694 (*Rhodomela dibromobenzyl constits*)**Oxosarcophylline**

[95377-97-2]


C₁₈H₁₃NO₅ 323.304

The trivial name is somewhat misleading (Sarcophylline, S-80 is 1,2,3,4-tetrahydro). Alkaloid from *Sarcocapnos enneaphylla* and *Sarcocapnos crassifolia* (Papaveraceae). Yellow needles (EtOH). Mp 170-171°.

Campello, M.J. et al., *Tet. Lett.*, 1984, **25**, 5933 (uv, ir, pmr, ms, struct)**17-Oxosparteine****O-208**

Oxopachycarpine, 8CI. Oxosparteine.

Oxysparteine



(-)-form

C₁₅H₂₄N₂O 248.367

Log P 1.38 (calc).

▶WG6390000

(+)-form [29882-72-2]

Alkaloid from aerial parts of *Genista involucrata* and *Genista albida*. Mp 82-84°. [α]_D +20 (EtOH). This appears to be the commonest form, but opt. rotns. of most samples are low or unrecorded.


(-)-form [489-72-5]

Alkaloid from leaves and fruits of *Chamaecytisus austriacus* and other *Chamaecytisus* spp. Also from *Thermopsis montana*, *Genista monosperma* and *Sophora pachycarpa* (Fabaceae). Hypertensive agent. Antiarrhythmic, oxytocic agent. Shows cathartic, diuretic activity. Mp 84-86°. [α]_D²⁵ -5.6 (c, 0.47 in EtOH). Pharmacol. active isomer.

Sadykov, A.S. et al., *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1957, **15**; *CA*, **53**, 9261White, E.P. et al., *N.Z. J. Sci. Technol., Sect. B*, 1957, **38**, 712 (isol, struct)Wiewiorowski, M. et al., *Can. J. Chem.*, 1967, **45**, 1447 (ir, pmr, conformn)Keller, W.J. et al., *J. Nat. Prod.*, 1969, **32**, 498 (isol)Iskandarov, S. et al., *Khim. Prir. Soedin.*, 1970, **6**, 494; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 519 (ord)Cho, Y.D. et al., *Arch. Mass Spectral Data*, 1971, **2**, 736 (ms)Bohlmann, F. et al., *Chem. Ber.*, 1975, **108**, 1043 (cmr)Daily, A. et al., *Planta Med.*, 1977, **32**, 380; 1979, **35**, 188 (isol)Daily, A. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 889Katrusiak, A. et al., *Acta Cryst. B*, 1980, **36**, 2442 (cryst struct)Tosun, F. et al., *Planta Med.*, 1987, **53**, 499 (isol)Katrusiak, A. et al., *Pol. J. Chem. (Roc. Chem.)*, 1994, **68**, 65 (cryst struct)**Oxostephanine****O-209**

9-Methoxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one, 9CI. NSC 383228

[58262-58-1]

C₁₈H₁₁NO₄ 305.289

Alkaloid from *Stephania japonica*, *Stephania venosa* and *Greenwayodendron (Polyalthia) suaveolens* (Menispermaceae, Annonaceae). Shows antibacterial activity. Yellow pillars (CHCl₃). Mp 270-272°.

N-Me: 9-Methoxy-7-methyl-8-oxo-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolinium(1+). **Thailandine**

[81255-03-0]

C₁₉H₁₄NO₄⁺ 320.324


Quaternary alkaloid from the tuberous roots of *Stephania venosa (Stephania rotunda)* (Menispermaceae). Partially dec. by chromatog. on silica gel to Oxostephanine.

O-De-Me: Oxostephanosine

[98672-45-8]

C₁₇H₉NO₄ 291.262Alkaloid from leaves of *Stephania venosa* (Menispermaceae). Yellow.Watanabe, Y. et al., *Phytochemistry*, 1975, **14**, 2522 (uv, ir, pmr, ms, struct)Cavé, A. et al., *Planta Med.*, 1978, **33**, 243 (isol)Guinaudeau, H. et al., *Chem. Comm.*, 1981, 1118 (*Oxostephanine, Thailandine, isol*)Hasan, C.M. et al., *J.C.S. Perkin 1*, 1982, 2807 (isol)Pharadai, K. et al., *J. Nat. Prod.*, 1985, **48**, 658 (*Oxostephanosine*)Ferdous, A.J. et al., *Fitoterapia*, 1992, **63**, 549-550 (*Oxostephanine, activity*)**3-Oxotabersonine****O-210**

[67249-35-8]



Absolute Configuration

C₂₁H₂₂N₂O₃ 350.416

Constit. of the seeds of *Amsonia elliptica* (Apocynaceae). Prisms (EtOAc). Mp 151.5–153°. [α]_D²⁵ -77.4 (CHCl₃). λ_{max} 297 (log ε 4.12); 331 (log ε 4.23) (MeOH).

14β,15β-Epoxy: 3-Oxopachysiphine

[116498-60-3]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from seeds of *Stemmadenia grandiflora*. Oil. [α]_D²¹ -205 (c, 0.14 in CHCl₃). May be identical with the subsequently descr. 3-Oxopachysiphine below. λ_{max} 296 (log ε 4.02); 328 (log ε 4.13) (MeOH).

14ξ,15ξ-Epoxy: 14,15-Epoxy-3-oxovincadifformine

[67249-36-9]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from seeds of *Amsonia elliptica* (Apocynaceae). Amorph. powder. Racemises on heating in soln. via a retro Diels-Alder cycloaddn. λ_{max} 222 (log ε 4.18); 297 (log ε 3.99); 328 (log ε 4.11) (MeOH).

14,15-Dihydro: 3-Oxovincadifformine

[55528-27-3]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from seeds of *Stemmadenia grandiflora* (Apocynaceae). Oil. [α]_D²¹ -315 (c, 0.12 in CHCl₃).

19-Oxo, 14,15-dihydro: 3-OxominovincineC₂₁H₂₂N₂O₄ 366.416

Alkaloid from *Tabernaemontana riedelii* (Apocynaceae). Needles (MeOH). Mp 257.5–259.5°. [α]_D²⁵ +268 (c, 0.1 in EtOH).

11-Methoxy: 11-Methoxy-3-oxotaberstonine

[102719-84-6]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from stems and leaves of *Alstonia yunnanensis*. Light yellow cryst. [α]_D³⁰ -67.4 (CHCl₃).

19S-Acetoxy: 19-Acetoxy-3-oxotaberstonine

[108906-71-4]

C₂₃H₂₄N₂O₅ 408.453

Alkaloid from the endocarp and seeds of *Melodinus scandens* (Apocynaceae).

Aimi, N. et al., *Chem. Pharm. Bull.*, 1978, **26**, 1182–1187 (8-Oxotaberstonine, 14,15-Epoxy-3-oxovincadifformine)

Levy, J. et al., *Tet. Lett.*, 1978, 1579–1580 (8-Oxotaberstonine)

Chen, W. et al., *Yaoxue Xuebao*, 1985, **20**, 906–912; 1986, **21**, 187–190; *CA*, **105**, 3520s; 3558k (19-Hydroxy-11-methoxytaberstonine)

Chazelet, I. et al., *Ann. Pharm. Fr.*, 1987, **44**, 355–362 (19-Acetoxy-3-oxotaberstonine)

Torrenegra, R. et al., *Phytochemistry*, 1988, **27**, 1843 (3-Oxopachysiphine)

Kalaus, G. et al., *J.O.C.*, 1997, **62**, 9188–9191 (3-Oxominovincine)

8-Oxo-2,4-tetradecadienoic acid O-211 acid

H₃C(CH₂)₅COCH₂CH₂CH=CHCH=CHCOOH

C₁₄H₂₂O₃ 238.326**(E,E)-form**

2-Methylpropylamide: N-Isobutyl-8-oxo-2,4-tetradecadienamamide. **Lanyuamide I**

C₁₈H₃₁NO₂ 293.448

Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. λ_{max} 259 (log ε 4) (EtOH).

2-Hydroxy-2-methylpropylamide: N-(2-Hydroxy-2-methylpropyl)-8-oxo-2,4-tetradecadienamamide. **Hydroxylanyuamide I**

C₁₈H₃₁NO₃ 309.448

Constit. of the root bark of *Zanthoxylum ailanthoides*. Oil. λ_{max} 259 (log ε 3.57) (MeOH).

Chen, I.-S. et al., *Phytochemistry*, 1999, **52**, 357–360 (**Lanyuamide I**)

Cheng, M.-J. et al., *J. Chin. Chem. Soc. (Taipei)*, 2003, **50**, 1241–1246 (**Hydroxylanyuamide I**)

12-Oxo-2,4,8-tetradecatrienoic acid O-212

H₃CCH₂COCH₂CH₂CH=CHCH₂CH₂CH=CHCH=CHCOOH

C₁₄H₂₀O₃ 236.31**(2E,4E,8Z)-form**

2-Methylpropylamide: N-Isobutyl-12-oxo-2,4,8-tetradecatrienamamide. **Lanyuamide II**

[249898-39-3]

C₁₈H₂₉NO₂ 291.433

Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. λ_{max} 259 (log ε 3.58) (EtOH).


2-Hydroxy-2-methylpropylamide: N-(2-Hydroxy-2-methylpropyl)-12-oxo-2,4,8-tetradecatrienamamide. **Hydroxylanyuamide II**

C₁₈H₂₉NO₃ 307.432

Constit. of the root bark of *Zanthoxylum ailanthoides*. Oil. λ_{max} 260 (log ε 4.01) (MeOH).

Chen, I.-S. et al., *Phytochemistry*, 1999, **52**, 357–360 (**Lanyuamide II**)

Cheng, M.-J. et al., *J. Chin. Chem. Soc. (Taipei)*, 2003, **50**, 1241–1246 (**Hydroxylanyuamide II**)

8-Oxotetrahydropalmatine O-213

(S)-form

C₂₁H₂₃NO₅ 369.416**(S)-form**

Alkaloid from the stem and roots of *Anamirta cocculus* (Menispermaceae).

13,13a-Didehydro: **Oxypalmatine**

[19716-59-7]

C₂₁H₂₁NO₅ 367.401

Alkaloid from stems of *Coscinium fenestratum* (Menispermaceae). Non-cryst.

13,13a-Didehydro, 8ξ-alcohol: **8-Hydroxypalmatine**. 7,8-Dihydro-8-hydroxypalmatine

C₂₁H₂₃NO₅ 369.416

Alkaloid from *Berberis heteropoda* (Berberidaceae) and *Enantia chlorantha*. Yellow cryst. (MeOH). Mp 129–130° Mp 188–190°. [α]_D -35 (c, 0.08 in CHCl₃). Higher Mp. and opt rotn. value refer to 7,8-dihydro-8-hydroxypalmatine from *E. chlorantha*. λ_{max} 358 (log ε 3.97); 434 (log ε 4.17) (no solvent reported).

(±)-form

Synthetic. Pale yellow cryst. Mp 167–168°.

Verpoorte, R. et al., *J. Nat. Prod.*, 1981, **44**, 221 (8-Oxotetrahydropalmatine)

Pinho, P.M.M. et al., *Phytochemistry*, 1992, **31**, 1403 (**Oxypalmatine, isol, ir, pmr, cmr, ms, struct**)

Nimgirawath, S. et al., *Aust. J. Chem.*, 1994, **47**, 951 (8-Oxotetrahydropalmatine, synth)

Shakirov, R. et al., *Khim. Prir. Soedin.*, 1996, **32**, 244–242; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 320 (8-Hydroxypalmatine)


Wafo, P. et al., *Phytochemistry*, 1999, **50**, 279–281 (8-Hydroxypalmatine)

Marek, R. et al., *J. Nat. Prod.*, 2003, **66**, 481–486 (8-Hydroxypalmatine, pmr, cmr, N-15 nmr)

Wang, X.-L. et al., *Acta Cryst. E*, 2006, **62**, 498–499 (**cryst struct**)

Oxothalicarpine O-214

[64234-41-9]

C₄₀H₄₀N₂O₉ 692.764

Alkaloid from *Hernandia ovigera* (Hernandiaceae). Mp 219–220° dec. [α]_D²⁵ +115 (c, 0.1 in CHCl₃).

O¹-De-Me, N⁶-Me, inner salt: **Vilaportine** [102487-23-0]

C₄₀H₄₀N₂O₉ 692.764


Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae).

Yang, T.-H. et al., *J. Chin. Chem. Soc. (Taipei)*, 1977, **24**, 91; *CA*, **87**, 130466p (**Oxothalicarpine**)

Chalandre, M.C. et al., *Can. J. Chem.*, 1986, **64**, 123 (**Vilaportine**)

Oxotuberostemonine O-215

[20675-62-1]

C₂₂H₃₁NO₅ 389.491

Alkaloid from the roots of *Stemona tuberosa* and *Stemona sessilifolia* (Stemonaceae). Also prod. by autoxidn. of

Tuberostemonine, T-664. Needles (MeOH). Mp 222° (217°).

Perchlorate: Mp 200°.


Ac: Mp 233-234°.

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1962, **40**, 455 (*isol, uv, ir*)

Huber, C.P. *et al.*, *Tet. Lett.*, 1968, 4081; *Acta Cryst. B*, 1972, **28**, 2015 (*cryst struct*)

Oxyberberine

Berberine
[549-21-3]



$C_{20}H_{17}NO_5$ 351.358

Alkaloid from *Arcangelisia gusanlung*, *Berberis lambertii*, other *Berberis* spp. and *Mahonia japonica*. Yellow needles (MeOH). Mp 198-200°. Genus name erroneously given as *Acanthopanax*.

▶ DS0180000

O⁹-De-Me: Oxyberberubine

[29580-82-3]

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from stems of *Arcangelisia gusanlung* (Menispermaceae). Reddish-brown needles. Mp 240-241°.

O¹⁰-De-Me: Oxythalifendine

[164920-59-6]

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from stems of *Arcangelisia gusanlung* (Menispermaceae). Light brown needles. Mp 286-287°.

8ξ-Alcohol: 7,8-Dihydro-8-hydroxyberberine. Berberinol

[10134-52-8]

[84094-52-0]

$C_{20}H_{19}NO_5$ 353.374

Alkaloid from *Arcangelisia flava* and *Mahonia japonica*. Yellow needles (MeOH). Mp 188-190° (145°). Possibly an artifact. Props refer to synthetic racemate. Not clear if nat. alkaloid is a racemate.

▶ DR9899000

8ξ-Alcohol, 8-Me ether: 7,8-Dihydro-8-methoxyberberine

$C_{21}H_{21}NO_5$ 367.401

Alkaloid from the stems of *Mahonia japonica*. Yellow needles (MeOH). Mp 140-142°. λ_{max} 255 ; 365 (EtOH).

Haworth, R.D. *et al.*, *J.C.S.*, 1927, 548 (*synth*)

Pavelka, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3654 (*uv, ir*)

Cushman, M. *et al.*, *J.O.C.*, 1979, **44**, 407 (*synth, ir, pmr, ms*)

Möhrle, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1982, **315**, 919 (*Berberinol, synth*)

Verpoorte, R. *et al.*, *J. Nat. Prod.*, 1982, **45**, 582 (*Berberinol, isol, uv, pmr, ms*)


Zhang, J.-S. *et al.*, *Phytochemistry*, 1995, **39**, 439-442 (*pmr, cmr, Oxyberberubine, Oxythalifendine*)

Hsieh, T.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2004, **51**, 443-446 (*Mahonia japonica berberines*)

Le, T.N. *et al.*, *Bull. Korean Chem. Soc.*, 2006, **27**, 2093-2096 (*synth*)

5,5'-Oxybis[1,5-dihydro-3-methyl-2H-pyrrol-2-one], 9CI

O-217
5,5'-Oxydi[3-methyl-3-pyrrolin-2-one]
[134296-37-0]



$C_{10}H_{12}N_2O_3$ 208.216

(5*SR*,5'*SR*)-form


meso-form

Alkaloid from the bulbs of *Lilium candidum* and *Lilium hansonii* (Liliaceae). Prisms (MeOH). Mp 213-215°.

Haladova, M. *et al.*, *Coll. Czech. Chem.*

Comm., 1991, **56**, 436 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ori, K. *et al.*, *Phytochemistry*, 1992, **31**, 2767 (*isol, uv, ms*)

7,7'-Oxybis[6-methyl-5,8-dioxo-1-isoquinolinemethanol]

$C_{22}H_{16}N_2O_7$ 420.378

Antibacterial agent. Red cryst. λ_{max} 231 (ε 31850); 256 (ε 29970); 304 (ε 6250); 324 (ε 8000); 379 (ε 5140); 485 (ε 2170) (MeOH).

Diangeloyl: [153585-64-9]


$C_{32}H_{28}N_2O_9$ 584.581

Alkaloid from the sponge *Haliclona* sp.

Parameswaran, P.S. *et al.*, *Oceanogr. Indian Ocean*, 1992, 417; *C.A.*, **120**, 187482v

3,3'-[Oxybis(methylene)]-bis[9-methoxy-9H-carbazole], 9CI

O-219
Bis(9-methoxy-9H-carbazol-3-ylmethyl) ether




$C_{28}H_{24}N_2O_3$ 436.509

Alkaloid from the stem bark of *Murraya koenigii*. Brown gum. λ_{max} 237 (log ε 4.87); 264 (log ε 4.55); 293 (log ε 4.42); 329 (sh) (log ε 3.79) (MeOH).

Rahman, M.M. *et al.*, *Phytochemistry*, 2005, **66**, 1601-1606 (*isol, pmr, cmr*)

10,11-Oxy-10,12a-cyclo-10,11-secocolchicine

O-220
10,11-Epoxycolchicine (obsol.)
[61014-70-8]



$C_{22}H_{25}NO_7$ 415.442

The synonym 10,11-Epoxycolchicine was based on an incorrect struct. assignment. Alkaloid from the seeds of *Colchicum latifolium* (Liliaceae). Cryst. (EtOAc or 2-propanol). Mp 251-253° (dec.). $[\alpha]_D^{22}$ -237 (c, 1.68 in $CHCl_3$). $[\alpha]_D^{25}$ -211 (c, 1 in $CHCl_3$).

Potěšilová, H. *et al.*, *Coll. Czech. Chem.*

Comm., 1976, **41**, 3146; 1977, **42**, 1571 (*isol, uv, ir, pmr, ms*)


Brossi, A. *et al.*, *Helv. Chim. Acta*, 1980,

63, 406 (*synth, uv, ir, pmr, cmr, cryst struct*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1982, **47**, 2258 (cd)

Oxydimurrayafoline

O-221
3,3'-[Oxybis(methylene)]bis[1-methoxy-9H-carbazole], 9CI
[107903-17-3]




$C_{28}H_{24}N_2O_3$ 436.509

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Oil. First example of a binary carbazole alkaloid having an ether linkage.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 450 (*uv, pmr, ms, struct*)

Cypridina Oxyluciferin


O-222

Cypridina oxyluciferin
[17297-78-8]C₂₁H₂₇N₇O 393.491Isol. from *Cypridina luciferin*. Chemiluminescent agent.


[29066-77-1, 29066-76-0]

Goto, T. *et al.*, *Tet. Lett.*, 1968, 4035-4038
(*struct.*, *uv.*, *fluorescence*)Sugiura, S. *et al.*, *Yakugaku Zasshi*, 1970, **90**,
711-715 (*synth*)Suzuki, N. *et al.*, *Heterocycles*, 1983, **20**, 1027-
1030 (*fluorescence*)**Watasenia Oxyluciferin**

O-223


Watasenia oxyluciferin
[54028-46-5]C₂₅H₂₁N₃O₉S₂ 571.588Isol. from the light organs of the squid *Watasenia scintillans*. Possibly involved in bioluminescence.Goto, T. *et al.*, *Tet. Lett.*, 1974, 2321 (*isol.*, *uv.*,
pmr., *struct.*, *synth*)**Oxypterine**

O-224

4-Chloro-1-(chloromethyl)octahydro-5-methyleneazocine, 9Cl
[143114-89-0]C₉H₁₅Cl₂N 208.13**(+)-form**Alkaloid from the leaves and twigs of *Lotononis oxyptera*. Off-white cryst. solid. Mp 127-129°. [α]_D²² +19.6 (c, 2.6 in CHCl₃).Verdoorn, G.H. *et al.*, *Phytochemistry*, 1992, **31**, 1029-1032 (*isol.*, *ir.*, *pmr.*, *cmr.*, *ms*)**Oxysceptrine**

O-225

[117417-62-6]

C₂₂H₂₄Br₂N₁₀O₃ 636.305Alkaloid from the marine sponge *Agelas cf. nemochinata*, *Agelas clathrodes*, *Agelas mauritiana* and *Agelas confiera*. Actomyosin ATPase activator. Possesses antiviral and antibacterial props. Antifouling agent. Amorph. solid. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D -19.7 (c, 1.02 in MeOH). λ_{\max} 203 (€ 35100); 221 (€ 30700); 267 (€ 28100) (MeOH) (Derep). λ_{\max} 214 (€ 21000); 268 (€ 21000) (MeOH) (Derep). λ_{\max} 206 (€ 35100); 221 (€ 30100); 267 (€ 23100) (MeOH).*Di-Ac*: [117417-63-7][α]_D²⁵ +4.5 (c, 1.37 in MeOH).**Monobromo: Debromooxysceptrine**

[132911-41-2]

[117472-97-6]


C₂₂H₂₅BrN₁₀O₃ 557.409Alkaloid from *Agelas confiera*, *Agelas clathrodes* and *Agelas mauritiana*.Possesses antiviral and antibacterial props. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. [α]_D²⁵ -25 (c, 0.108 in MeOH) (as di-Ac). λ_{\max} 203 (€ 32400); 220 (€ 24600); 265 (€ 26600) (MeOH) (Derep).

[117468-53-8]

Kobayashi, J. *et al.*, *Experientia*, 1991, **47**, 301-
304 (*isol.*, *uv.*, *ir.*, *pmr.*, *cmr.*, *ms*)Keifer, P.A. *et al.*, *J.O.C.*, 1991, **56**, 2965-2975
(*isol.*, *pmr.*, *cmr*)O'Malley, D.P. *et al.*, *J.A.C.S.*, 2007, **129**,
4762-4775; 7702 (*synth*)**Oxysporidinone**

O-226

[184871-55-4]




Relative Configuration

C₂₈H₄₃NO₆ 489.651Prod. by *Fusarium oxysporum* strains CBS 330.95 and EPH2R_{AA}. Antifungal agent. Glass. [α]_D +97 (c, 0.2 in EtOH). Related to Funiculosin, F-197 and Sambutoxin, S-48. λ_{\max} 291 (log € 3.68) (MeOH).**4'-Di-Me acetal:**C₃₀H₄₉NO₇ 535.72Prod. by *Fusarium oxysporum* strain N17B. Amorph. solid. [α]_D²⁶ -30.6 (c, 0.1 in CHCl₃). λ_{\max} 216 (log € 3.62); 290 (log € 3.45) (CHCl₃).**6'-Deoxy: 6'-Deoxyoxysporidinone**C₂₈H₄₃NO₅ 473.651Prod. by *Fusarium oxysporum* strain EPH2R_{AA}. Cryst. Mp 130-131°. [α]_D²⁵ -72 (c, 0.15 in MeOH). λ_{\max} 220 (log € 4.46); 296 (log € 3.66) (MeOH).**2'-Epimer: 2'-Epioxysporidinone**C₂₈H₄₃NO₆ 489.651Prod. by *Fusarium oxysporum* strain N17B. Antifungal agent. Needles (MeOH). Mp 176-178°. [α]_D²⁶ -86.9 (c, 0.1 in CHCl₃). λ_{\max} 220 (log € 3.64); 296 (log € 3.45) (CHCl₃).Breinholt, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 33-
35 (*isol.*, *uv.*, *ir.*, *pmr.*, *cmr.*, *ms*)Jayasinghe, L. *et al.*, *J. Nat. Prod.*, 2006, **69**,
439-442 (*2'-epimer.*, *di-Me acetal*)Zhan, J. *et al.*, *J. Nat. Prod.*, 2007, **70**, 227-232
(*isol.*, *pmr.*, *cmr.*, *ms*)**Oxystemokerrilactone**


O-227

[949092-27-7]

Relative
ConfigurationC₁₆H₂₅NO₄ 295.378Alkaloid from the roots of *Stemona saxorum*. Light yellow powder. [α]_D²⁰ -34 (c, 0.14 in MeOH).Wang, Y.-Z. *et al.*, *J. Nat. Prod.*, 2007, **70**,
1356-1359 (*isol.*, *pmr.*, *cmr*)


Pachychaline A

P-1

C₅₄H₁₀₀N₄²⁺ 805.411Isol. from a *Pachychalina* sp. Oil. Counterion not specified. λ_{max} 267 (ε 6600) (MeOH).Laville, R. *et al.*, *Eur. J. Org. Chem.*, 2008, 121-125 (*isol, synth, pmr, cmr, ms*)


Pachychaline B

P-2

C₅₈H₁₁₀N₆²⁺ 891.547Isol. from a *Pachychalina* sp. Oil. Counterion not specified. λ_{max} 267 (ε 6800) (MeOH).Laville, R. *et al.*, *Eur. J. Org. Chem.*, 2008, 121-125 (*isol, pmr, cmr, ms*)

Pachychaline C


P-3

C₇₁H₁₂₈N₅³⁺ 1051.826Isol. from a *Pachychalina* sp. Oil. Counterion not specified. λ_{max} 267 (ε 9700) (MeOH).Laville, R. *et al.*, *Eur. J. Org. Chem.*, 2008, 121-125 (*isol, pmr, cmr, ms*)

Pachyconfine

P-4

5,6,6a,7-Tetrahydro-1-methoxy-6-methyl-4H-dibenzo[de,g]quinoline-2,7-diol, 9Cl. 2,7-Dihydroxy-1-methoxyyaporphine. 6-Hydroxyasimilobine [62951-75-1]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Pachypodanthium confine* (Annonaceae). Mp 198° (as hydrochloride). [α]_D -150 (c, 0.7 in CHCl₃).

N-Oxide: **Pachyconfine N-oxide**

[112494-70-9]

C₁₈H₁₉NO₄ 313.352

Alkaloid from bark of Colombian

Duguetia spixiana (Annonaceae).Cryst. (MeOH). Mp 176-180°. [α]_D -164 (c, 0.41 in CHCl₃).

N-De-Me: 2,7-Dihydroxy-1-methoxy-norpachyconfine. **Norpachyconfine**

[112571-90-1]

C₁₇H₁₇NO₃ 283.326Alkaloid from *Duguetia spixiana* (Annonaceae). Amorph. [α]_D -281 (c, 0.07 in CHCl₃).

O²-Me: 7-Hydroxy-1,2-dimethoxyyaporphine. **Nuciferidine**

[107882-18-8]

C₁₉H₂₁NO₃ 311.38

Alkaloid from the trunk bark of

Gutteria sagotiana (Annonaceae).

Obt. impure. Negative opt. rotn.

O²-Me, N-de-Me: **Nornuciferidine**

[112494-69-6]

C₁₈H₁₉NO₃ 297.353Alkaloid from bark of *Duguetia spixiana* (Annonaceae). [α]_D -80 (c, 0.16 in EtOH).

O²-Me, N-de-Me, O⁷,N-di-Ac: **N,O-Diacetylnornuciferidine**

C₂₂H₂₃NO₅ 381.427Alkaloid from the rhizomes of *Lysichiton camtschatcense*. Yellow powder.[α]_D²⁵ -227.7 (c, 0.24 in CHCl₃). λ_{max} 211 (log ε 4.39); 273 (log ε 4); 313 (log ε 3.4) (EtOH).

6-Epimer, N-de-Me: **Anaxagoreine**

[77410-38-9]

C₁₇H₁₇NO₃ 283.326Alkaloid from *Anaxagorea dolichocarpa*, *Anaxagorea primoides* and *Cananga odorata* (ylang ylang). Amorph. Mp 198-200°. [α]_D -174 (c, 0.5 in EtOH).

Bévalot, F. *et al.*, *Ann. Pharm. Fr.*, 1977, **35**, 65; *CA*, **86**, 185953e (*isol, uv, pmr, ms, struct*)
Hocquemiller, R. *et al.*, *Planta Med.*, 1981, **41**, 48 (*Anaxagoreine*)

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1078; 1987, **50**, 674 (*Nuciferidine*, *Nornuciferidine*)

Debouges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664 (*Norpachyconfine*, *Pachyconfine N-oxide*)


Kessar, S.V. *et al.*, *Indian J. Chem., Sect. B*, 1994, **33**, 818 (*synth, Nuciferidine*)

Hsieh, T.J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1999, **46**, 607-612 (*Anaxagoreine*)

Takatsu, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 430-431 (*N,O-Diacetylnornuciferidine*)

Pachydermin

P-5

C₁₃H₈ClNO₆ 309.662


Tetramic acid deriv. Prod. by *Chamonixia pachydermis*. Amorph. yellow solid. λ_{max} 203 ; 254 ; 357 (TFA/MeCN aq.).

Lang, G. *et al.*, *J. Nat. Prod.*, 2006, **69**, 151-153 (*isol, pmr, cmr*)

Pachysantermine A

P-6

[15027-63-1]

C₂₉H₄₈N₂O₂ 456.71

Minor alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 260-263°. [α]_D +43 (CHCl₃).

N³-Me: Mp 258-261°.

Kikuchi, T. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 571 (*ir, uv, pmr, struct*)

Pachysterine

P-7

[1360-19-6]

Steroidal alkaloid. MF C₂₈₋₂₉H₄₈₋₅₀N₂O₂. Struct. unknown. Alkaloid from *Pachysandra terminalis*. Mp 220-224°.


[α]_D +24 (CHCl₃).

Kikuchi, T. *et al.*, *Tet. Lett.*, 1964, 1817-1823

Pachystermine A

P-8

[6156-99-6]



Absolute Configuration

C₂₉H₄₈N₂O₂ 456.71

Major alkaloid of *Pachysandra terminalis* (Buxaceae). Cytotoxic for P388 cells. Mp 220-224°. [α]_D +24 (CHCl₃).

▶CM4418000

4β-Alcohol: **Pachystermine B**

[6157-00-2]

C₂₉H₅₀N₂O₂ 458.726

Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 256-259°. [α]_D²⁰ -18 (CHCl₃).

2,3-Didehydro: 20-(Dimethylamino)-3-(3-isopropyl-2-oxo-1-azetidiny)pregn-2-en-4-one

[214402-19-4]

Alkaloid from *Pachysandra procumbens*. Needles. Mp 208°. [α]_D²⁰ +41.7 (c, 0.08 in CHCl₃). λ_{max} 221 (log ε 3.96); 270 (log ε 3.75) (MeOH).

16α-Hydroxy: 20-(Dimethylamino)-16-

hydroxy-3-(3-isopropyl-2-oxo-1-azetidiny)pregnan-4-one. 20-(Dimethylamino)-16-hydroxy-3-(3-isopropylactam)-pregnan-4-one

$C_{29}H_{48}N_2O_3$ 472.71

Alkaloid from *Pachysandra procumbens*. Needles. Mp 205°. $[\alpha]_D^{20} +67.4$ (c, 0.05 in $CHCl_3$). λ_{max} 242 (log ϵ 3.39) (MeOH).

16 α -Hydroxy, 2,3-didehydro: 20-(Dimethylamino)-16-hydroxy-3-(3-isopropyl-2-oxo-1-azetidiny)pregn-2-en-4-one

[214402-20-7]

$C_{29}H_{46}N_2O_3$ 470.694

Alkaloid from *Pachysandra procumbens*. Needles. Mp 205°. $[\alpha]_D^{20} +26.1$ (c, 0.07 in $CHCl_3$). λ_{max} 220 (log ϵ 3.83); 275 (log ϵ 3.5) (MeOH).

Kikuchi, T. et al., *Chem. Pharm. Bull.*, 1967, **15**, 549 (ir, ms, pmr, struct)

Kikuchi, T. et al., *Tet. Lett.*, 1968, 909 (synth)

Chang, L.C. et al., *J. Nat. Prod.*, 1998, **61**,

1257 (2,3-didehydro derivs)

Funayama, S. et al., *Biol. Pharm. Bull.*, 2000,

23, 262 (isol, activity)


Chang, L.C. et al., *Tetrahedron*, 2000, **56**,

3133-3138 (16-hydroxy)

Paeciloxazine

P-9

[677333-77-6]



$C_{29}H_{38}N_2O_7$ 526.628


Related to Antibiotic CJ 12662, A-1141. Prod. by *Paecilomyces* sp. strain BAUA3058. Moderate nematocidal agent. Powder. Mp 158°. λ_{max} 248 (ϵ 7400); 285 (ϵ 1400) (MeOH).

Kanai, Y. et al., *J. Antibiot.*, 2004, **57**, 24-28 (isol, pmr, cmr, activity)

Pagicerine

P-10

[99831-97-7]



Absolute Configuration

$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the stem bark of *Pagiantha cerifera* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. solid. $[\alpha]_D^{20} -130$ (c, 0.7 in $CHCl_3$).

De(methoxycarbonyl): **Amerovolficine.**

16-Demethoxycarbonylpagicerine

[123828-66-0]

$C_{20}H_{22}N_2O_2$ 322.406

Alkaloid from the stem bark of *Rauwolfia cubana* (Apocynaceae). Cryst. (MeOH). Mp. not recorded.

De(methoxycarbonyl), 10-methoxy: 10-Methoxy-16-demethoxycarbonylpagicerine. **10-Methoxyamerovolficine**

$C_{21}H_{24}N_2O_3$ 352.432

Alkaloid from the roots of *Rauwolfia yunnanensis*. Amorph. yellow solid.

$[\alpha]_D^{24} -97$ (c, 0.6 in Py). λ_{max} 209 (log ϵ 4.29); 322 (log ϵ 3.98) (MeOH).

Bert, M. et al., *Heterocycles*, 1985, **23**, 2505 (Pagicerine, struct)


Martinez, J.A. et al., *Planta Med.*, 1989, **55**, 283 (Amerovolficine)

Hu, X.-J. et al., *Helv. Chim. Acta*, 2006, **89**, 1344-1350 (10-Methoxyamerovolficine)

Pagisulfine

P-11

[103956-35-0]



$C_{23}H_{31}N_3O_2S$ 413.583

Alkaloid from the stem bark of *Pagiantha cerifera* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. solid. $[\alpha]_D^{20} +42$ (c, 1 in EtOH).


Bert, M. et al., *Heterocycles*, 1986, **24**, 1567 (uv, ir, pmr, ms, struct, synth)

Pahutoxin

P-12

2-[[[3-(Acetyloxy)-1-oxohexadecyl]oxy]-N,N,N-trimethylethanaminium(1+)], 9CI. 3-Acetoxyhexadecanoic acid choline ester

[122767-82-2]



$C_{23}H_{46}NO_4^{\oplus}$ 400.621

(S)-form [27742-14-9]

Poison secreted by the blue dogfish (*Ostracion lentiginosus*). Needles (Me₂CO) (as chloride). Mp 74-75° (chloride). $[\alpha]_D^{22} +3.05$ (c, 2.3 in MeOH). CAS no. refers to chloride.

► Toxic.

O-De-Ac, O-propanoyl: **Homopahutoxin**

[109777-68-6]

$C_{24}H_{48}NO_4^{\oplus}$ 414.648

Isol. from the boxfish *Ostracion immaculatus*. Haemolytic agent.

(ξ)-form [116613-97-9]

Constit. of toxic skin secretions of the trunkfish, *Anoplocapros lenticularis*, *Araucana ornata*, *Lactoria fornasini*, *Ostracion cubicus*, *Rhinesomas reipublicae* and *Strophurichthys robustus*. CAS no. refers

to chloride. Genus name incorr. given as *Anoplocapros*.

O-De-Ac: [116613-93-5]

$C_{21}H_{44}NO_3$ 358.584

Constit. of toxic skin secretions of the trunkfish *Anoplocapros lenticularis*, *Araucana ornata*, *Lactoria fornasini* and *Ostracion cubicus*. CAS no. refers to chloride.

O-De-Ac, O-propanoyl: [116613-96-8]

$C_{24}H_{48}NO_4$ 414.648

Trace constit. of toxic skin secretions of the trunkfish *Lactoria fornasini* and *Rhinesomas reipublicae*. CAS no. refers to chloride.

O-De-Ac, O-butanoyl: [116613-95-7]

$C_{25}H_{50}NO_4$ 428.674

Constit. of toxic skin secretions of the trunkfish *Anoplocapros lenticularis* and *Strophurichthys robustus*. CAS no. refers to chloride.

O-De-Ac, O-pentanoyl: [116613-94-6]

$C_{26}H_{52}NO_4$ 442.701

Constit. of toxic skin secretions of the trunkfish *Anoplocapros lenticularis*, *Rhinesomas reipublicae* and *Strophurichthys robustus*. CAS no. refers to chloride.

Boylan, D.B. et al., *Science (Washington, D.C.)*, 1967, **155**, 52-56

Fusetani, N. et al., *Toxicol.*, 1987, **25**, 459-461 (Homopahutoxin, isol)

Goldberg, A.S. et al., *Toxicol.*, 1988, **26**, 651-663 (trunkfish constit)

Yoshikawa, M. et al., *Agric. Biol. Chem.*, 1989, **53**, 37-40 (synth, ir, pmr, abs config)

Pahybrine

P-13

[93585-77-4]

$C_{22}H_{30}N_2O_4$ 386.49

Struct. unknown. Alkaloid from *Papaver hybridum*. Amorph. yellow solid. Mp 204° dec. (as dihydrobromide). $[\alpha]_D^{25} +45$ (H₂O) (dihydrobromide).

Platonova, T.F. et al., *Zh. Obshch. Khim.*, 1956, **26**, 173-180; *CA*, **50**, 13960e

Paipunine

P-14

$C_{24}H_{37}NO_4$ 403.561

Struct. unknown. Alkaloid from the Chinese drug "Pai pu", stated to be derived from an unidentified *Stemona* sp. (Stemonaceae). Prisms (2-propanol). Mp 105.5-106.5°. $[\alpha]_D^{25} -53.7$ (Me₂CO).

Hydrochloride:


► LD₅₀ (mus, ivn) 39 mg/kg. RT1310000

Lee, H.M. et al., *J. Am. Pharm. Assoc.*, 1940, **29**, 391-394 (isol, pharmacol)

Pakistanamine

P-15

[36506-66-8]



Absolute configuration

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from *Berberis valdiviana*, the roots of *Berberis baluchistanica*, *Berberis calliobotrys* and *Berberis orthobotrys*, and from the seeds of *Berberis julianae* (Berberidaceae). Mp 93-94°. [α]_D²⁵ +135 (c, 0.5 in MeOH). Dec. on heating >40° in soln. Rearr. in 3M HCl at 70° into 1-*O*-Methylpakistanine, 1-*O*-Methylkalashine and the monomeric (+)-Armepevine. In methanolic HCl, also forms 1,10-Di-*O*-methylpakistanine by methanolysis.

Hydrochloride:

Cryst. (MeOH). Mp 215°. [α]_D²⁵ +20 (c, 0.34 in MeOH).

Picrate:

Yellow cryst. (EtOH). Mp 158-162° dec.

O¹-De-Me: Valdivianine

[84423-01-8]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from *Berberis valdiviana* and *Berberis empetrifolia* (Berberidaceae). Amorph. [α]_D²⁵ +120 (c, 0.2 in MeOH).

O⁷-De-Me: Patagonine

[84423-06-3]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from *Berberis valdiviana* and *Berberis empetrifolia* (Berberidaceae). Amorph. [α]_D²⁵ +192 (c, 0.2 in MeOH).

O¹,O⁶-Di-de-Me: Berbivaldine

[84423-04-1]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis valdiviana* (Berberidaceae). Amorph. [α]_D +140 (c, 0.4 in MeOH). Undergoes acid-cat. rearr. to Porveniramine.

O¹,O⁷-Di-de-Me: Valdiberine

[84472-23-1]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis valdiviana* (Berberidaceae). Amorph. [α]_D²⁵ +91 (c, 0.4 in MeOH).

13-Epimer, O¹-de-Me: Epivaldivianine

[98604-30-9]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stems of *Berberis valdiviana* (Berberidaceae). [α]_D²⁵ +69.4 (c, 0.1 in MeOH).

13-Epimer, O¹,O⁶-di-de-Me: Epiberbivaldine

[96245-07-7]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis actinacantha* (Berberidaceae). Amorph. [α]_D²⁵ +45.7 (c, 0.12 in CHCl₃).

13-Epimer, O¹,O⁷-di-de-Me: Epivaldiberine

[84472-22-0]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis valdiviana* (Berberidaceae). Amorph. [α]_D²⁵ +31 (c, 0.1 in MeOH). Undergoes acid-cat. rearr. to Khyberine.

Shamma, M. *et al.*, *J.A.C.S.*, 1972, **94**, 1381; 1973, **95**, 5742 (*isol, uv, ir, pmr, ms, ord, struct*)

Kametani, T. *et al.*, *Heterocycles*, 1974, **2**, 159 (*synth*)

Hussain, S.F. *et al.*, *Tet. Lett.*, 1980, **21**, 723; 3315; 4573

Hussain, S.F. *et al.*, *Heterocycles*, 1981, **15**, 191 (*occur*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 274 (*isol*)


Guinaudeau, H. *et al.*, *Chem. Comm.*, 1982, 1122 (*occur, stereochem, cd, pmr, derivs*)

Weiss, I. *et al.*, *Heterocycles*, 1984, **22**, 2231 (*Epiberbivaldine*)

Firdous, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 664 (*Epivaldivianine*)

Pakistanine

[36506-69-1]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the roots of *Berberis baluchistanica*, *Berberis calliobotrys* and *Berberis orthobotrys*, and from the stems and above-ground wood of *Berberis empetrifolia*, and from *Berberis valdiviana* (Berberidaceae). Needles (EtOH). Mp 154-156°. [α]_D²⁵ +106 (c, 0.57 in MeOH).

N²-De-Me: 2'-Norpakistanine

[98618-06-5]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the stems of *Berberis valdiviana* (Berberidaceae). Cryst. (MeOH). Mp 148°.

O¹-Me: 1-O-Methylpakistanine

[36418-13-0]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from the roots of *Berberis calliobotrys* and *Berberis orthobotrys* (Berberidaceae), also obt. by rearr. of Pakistanine, P-15. Cryst. (Et₂O). Mp 117°. [α]_D²⁵ +85 (c, 0.40 in MeOH).

Di-Me ether: Formed by heating Pakistanine, P-15 in MeOH/HCl. Fine needles (Et₂O). Mp 139-141°. [α]_D²⁵ +66 (c, 0.40 in MeOH).

O⁶-De-Me: Porveniramine

[84423-08-5]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis empetrifolia* (Berberidaceae), also obt. by acid-cat. rearr. of Berbivaldine (see Pakistanine, P-15). [α]_D²⁵ +40 (c, 0.1 in MeOH).

O⁶-De-Me, O¹-Me: Waziristanine. 1-O-Methylporveniramine

[96497-83-5]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from root bark of *Berberis waziristanica* (Berberidaceae). [α]_D²⁵ +38 (c, 0.06 in MeOH).

O⁷-De-Me: Chitraline

[77754-91-7]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Berberis empetrifolia* and roots of *Berberis calliobotrys* and *Berberis orthobotrys* (Berberidaceae). Also obt. by dienone-phenol rearr. of

Valdiberine (see Pakistanine, P-15). Amorph. [α]_D²⁵ +136 (c, 0.172 in MeOH).

O⁷-De-Me, O¹-Me: 1-O-Methylchitraline

[84423-09-6]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Also obt. by acid rearr. of Patagonine (see under Pakistanine, P-15). Amorph. [α]_D²⁵ +29 (c, 0.4 in MeOH).

Shamma, M. *et al.*, *J.A.C.S.*, 1972, **94**, 1381; 1973, **95**, 5742 (*isol, uv, ord, pmr, ms, struct, abs config*)

Hussain, S.F. *et al.*, *Tet. Lett.*, 1980, **21**, 4573 (*isol, uv, pmr*)

Hussain, S.F. *et al.*, *Heterocycles*, 1981, **15**, 191 (*isol, uv, cd, pmr, ms, struct, Chitraline*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 274 (*isol, uv, pmr*)

Guinaudeau, H. *et al.*, *Chem. Comm.*, 1982, 1122 (*occur, cd, pmr, synth*)

Fajardo, V. *et al.*, *Tet. Lett.*, 1982, **23**, 39 (*occur*)

Firdous, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 664 (*O-Methylchitraline, 2'-Norpakistanine*)


Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1992, **31**, 1835 (*Waziristanine*)

Palasimide

P-17

4,7-Epoxyhexahydro-3a-methyl-2-phenyl-1H-isoindole-1,3(2H)-dione

[54382-61-5]

C₁₅H₁₅NO₃ 257.288

Alkaloid from the pods of *Butea monosperma*. Off-white fibrous needles (C₆H₆/petrol). Mp 204-205°. [α]_D -58.4 (CHCl₃).

Bochis, R.J. *et al.*, *Tet. Lett.*, 1968, 1971 (*synth*)


Peter, M.G. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 32 (*synth*)

Guha, P.K. *et al.*, *Phytochemistry*, 1990, **29**, 2017 (*isol, struct*)

Palauamine

P-18

[148717-58-2]



Absolute Configuration

C₁₇H₂₂ClN₉O₂ 419.873

Stereochem. revised in 2007. Alkaloid from the sponge *Stylotella aurantium* and *Stylotella agminata*. Cytotoxic. Possesses immunosuppressive, antibiotic and

antifungal activities. Off-white amorph. powder. $[\alpha]_D^{24}$ -45.2 (c, 3.0 in MeOH). λ_{\max} 273 (ε 4700) (MeOH) (Derep). λ_{\max} 224 (ε 7800); 272 (ε 7900) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 13 mg/kg.

4-Bromo-4-Bromopalauamine

[206069-58-1]

C₁₇H₂₁Br₂ClN₉O₂ 498.769

Alkaloid from *Stylorella aurantium*.

Amorph. tan solid. $[\alpha]_D$ -64.4 (c, 2.6 in MeOH).

4-Bromo, N¹⁹-(4,5-dibromo-2-pyrrolicarbonyl): Konbuacidin B

[934338-16-6]

[934338-17-7 (TFA salt)]

C₂₂H₂₂Br₃ClN₁₀O₃ 749.646

Alkaloid from *Stylyssa flabellata*.

Amorph. solid. $[\alpha]_D^{24}$ -24 (c, 0.04 in MeOH). λ_{\max} 281 (log ε 3.91) (MeOH).

4,5-Dibromo: 4,5-Dibromopalauamine

[206069-59-2]

C₁₇H₂₀Br₂ClN₉O₂ 577.665

Alkaloid from *Stylorella aurantium*

and *Stylyssa flabellata*. Solid. $[\alpha]_D$ -

115.3 (c, 2.7 in MeOH). Dec. on

heating.

12,17-Diepimer, 4-bromo, N¹⁹-(4,5-dibromo-2-pyrrolicarbonyl): Konbuacidin A

[200207-22-3]

C₂₂H₂₂Br₃ClN₁₀O₃ 749.646

The imidazole ring systems are delocalised.

Alkaloid from the sponge

Hymeniacidon sp. Cyclin dependent

kinase 4 inhibitor. Amorph. solid. $[\alpha]_D^{24}$

-45 (c, 0.5 in MeOH). λ_{\max} 277 (ε

13000) (MeOH).

Kobayashi, J. *et al.*, *Tetrahedron*, 1997, **53**,

15681-15684 (*Konbuacidin A*)

Kinnel, R.B. *et al.*, *J.O.C.*, 1998, **63**, 3281-3286

(*isol, uv, ir, pmr, cmr, ms, cd, struct, 4-bromo,*

dibromo)

Jacquot, D.E.N. *et al.*, *Curr. Org. Chem.*, 2005,

9, 1551-1566 (*rev, synth*)

Wang, S. *et al.*, *Tetrahedron*, 2006, **62**, 7155-

7161 (*synth*)

Grube, A. *et al.*, *Angew. Chem., Int. Ed.*, 2007,

46, 2320-2324 (*config*)

Köck, M. *et al.*, *Angew. Chem., Int. Ed.*, 2007,

46, 6586-6594 (*rev*)

Buchanan, M.S. *et al.*, *J.O.C.*, 2007, **72**,

2309-2317 (*Konbuacidin B, isol, config,*


dibromo)

Buchanan, M.S. *et al.*, *Tet. Lett.*, 2007, **48**,

4573-4574 (*abs config*)

Palauimide

P-19



C₂₅H₃₆N₂O₄ 428.57

Isol. from a *Lyngbya* sp. Cytotoxic.

Amorph. solid. $[\alpha]_D^{25}$ +50 (c, 0.33 in

MeOH). CAS no. not found to CA 138.

λ_{\max} 201 (log ε 3.78); 252 (log ε 3.44)


(MeOH).

Luesch, H. *et al.*, *Tetrahedron*, 2002, **58**, 7959-

7966 (*isol, pmr, cmr*)

Pali clavine

[52052-66-1]



C₁₆H₂₀N₂O 256.347

(+)-form

Alkaloid from *Claviceps paspali*. Cryst. (Me₂CO). Mp 177-179° dec. $[\alpha]_D^{20}$ +3 (c, 0.2 in Py). Yields Paspaclavine, P-106 on treatment with acetaldehyde.

O-Ac: Mp 149-150°. $[\alpha]_D^{20}$ -43 (c, 0.3 in Py).

(±)-form

Synthetic. Cryst. (Me₂CO). Mp 175-177°.

Tscherter, H. *et al.*, *Helv. Chim. Acta*, 1974, **57**,

113 (*isol, uv, ir, pmr, struct*)

Fehr, T. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 2484

(*synth*)

Oppolzer, W. *et al.*, *Tetrahedron*, 1983, **39**,

3695 (*synth, ir, pmr, ms*)

Kozikowski, A.P. *et al.*, *Tetrahedron*, 1984, **40**,

2345 (*synth*)

Palidimine

P-21

Dimeric alkaloid, prob. containing


Palimine in P-135 as one of its units.

Stuart, K.L. *et al.*, *Tet. Lett.*, 1974, 3853-3856

Palinurine A†

P-22

[254901-26-3]



C₂₇H₃₉NO₅ 457.609

Microbial metab. of Palinurin prod. by

Cunninghamella sp. NRRL 5695. Yel-

lowish oil. $[\alpha]_D^{25}$ +69.9 (c, 0.25 in CHCl₃).

λ_{\max} 246 (log ε 2.87); 268 (log ε 2.67)

(MeOH).

2-Deoxy, 5-oxo: Palinurine B

[254901-27-4]

C₂₇H₃₉NO₅ 457.609

Prod. by *Cunninghamella* sp. NRRL

5695. Yellowish oil. $[\alpha]_D^{25}$ +58.5 (c, 0.38

in CHCl₃). λ_{\max} 246 (log ε 2.87); 268


(log ε 2.67) (MeOH).

ElSayed, K.A. *et al.*, *J.O.C.*, 1999, **64**, 9258-9260

Paliurine B

P-23

[149183-88-0]



Absolute Configuration

C₃₆H₄₉N₅O₆ 647.813

Constit. of *Paliurus ramossissimus*. Ionophore. Amorph. solid. Mp 111-112°. $[\alpha]_D^{26}$ -391.3 (c, 0.76 in MeOH). λ_{\max} 216 (log ε 4.81); 268 (log ε 4.71); 319 (log ε 4.56) (MeOH).

2''-N-Me: Paliurine A

[298183-94-5]

C₃₇H₅₁N₅O₆ 661.84

Constit. of *Paliurus ramossissimus*.

Amorph. powder. $[\alpha]_D^{26}$ -345 (c, 1 in

MeOH). λ_{\max} 216 (log ε 4.84); 269 (log

ε 4.75); 318 (log ε 4.56) (MeOH).


Lin, H.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1338-

1343 (*isol, uv, pmr, cmr*)

Paliurine C

P-24

[298183-95-6]



Absolute Configuration

C₃₇H₅₁N₅O₆ 661.84

Alkaloid from the roots of *Paliurus*

ramossissimus. Amorph. solid. $[\alpha]_D^{26}$ -311

(c, 1 in MeCN). λ_{\max} 214 (log ε 4.8);

268 (log ε 4.62); 318 (log ε 4.48)

(MeOH).

2''-N-De-Me: Paliurine D

[298183-96-7]

C₃₆H₄₉N₅O₆ 647.813

Alkaloid from the roots of *Paliurus*

ramossissimus. $[\alpha]_D^{26}$ -164 (c, 1 in

MeCN). λ_{\max} 216 (log ε 4.56); 234 (log

ε 4.31); 269 (log ε 4.34); 318 (log ε 4.13)

(MeOH).

Lin, H.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1338-

1343 (*isol, pmr, cmr, struct*)


Paliurine E

P-25

Daechuine S6

[298184-60-8]

[123089-21-4]



C₃₁H₄₀N₄O₅ 548.681

Stereochemical identity of Paliurine E

and *Daechuine S6* is not certain.

Alkaloid from the stem bark of the

Daechu tree (*Zizyphus jujuba* var.

inermis) and from *Paliurus ramossissimus*

(Rhamnaceae). Mp 192°. $[\alpha]_D^{25}$ -393.5.

O-De-Me: Daechuine S26. Daechucyclopeptide I

[115610-63-4]

$C_{30}H_{38}N_4O_5$ 534.654

Alkaloid from the stem bark of *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Mp 114°. Daechuine S26 and Daechucyclopeptide I are given the same (planar) struct. Mp refers to Daechucyclopeptide I.

N,O-Di-de-Me: Lotusine F

[163136-15-0]

$C_{29}H_{36}N_4O_5$ 520.627

Alkaloid from root bark of *Zizyphus lotus* (Rhamnaceae). $[\alpha]_D^{25}$ -244 (c, 0.5 in $CHCl_3$).

Han, B.H. *et al.*, *Arch. Pharmacol Res.*, 1987, **10**, 208-211 (*Daechuine S26*)

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**, 443-448 (*Daechuine S26*)

Ghedira, K. *et al.*, *Phytochemistry*, 1995, **38**, 767-772 (*Lotusine F*)


Lin, H.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1338-1343 (*Paliurine E*)

Toumi, M. *et al.*, *J.O.C.*, 2008, **73**, 1270-1281 (*synth*)

Paliurine F

[298183-97-8]

P-26



$C_{34}H_{53}N_5O_6$ 627.823

Alkaloid from the roots of *Paliurus ramosissimus*. $[\alpha]_D^{26}$ -323 (c, 1 in MeCN). λ_{max} 217 (log ϵ 4.62); 269 (log ϵ 4.56); 318 (log ϵ 4.38) (MeOH).


Lin, H.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1338-1343 (*isol, ir, cd, pmr, cmr*)

Toumi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 572-575 (*synth*)

Paliurine G

[403842-59-1]

P-27



$C_{36}H_{49}N_5O_6$ 647.813

Alkaloid from the stems of *Paliurus ramosissimus*. Amorph. powder. $[\alpha]_D^{30}$ -335 (c, 0.33 in MeOH). λ_{max} 262 (log ϵ 4.79); 316 (log ϵ 4.18) (MeOH).

Stereoisomer(?) : Jubanine D

[164230-54-0]

$C_{36}H_{49}N_5O_6$ 647.813

Alkaloid from the root bark of *Zizyphus jujuba*. Same gross struct. as Paliurine G assigned, no stereochem. determined. Incorrect struct. shown in CAS.


Khokhar, I. *et al.*, *CA*, 1995, **123**, 29540v (*Jubanine D*)

Lee, S.-S. *et al.*, *Phytochemistry*, 2001, **58**, 1271-1276 (*Paliurine G*)

Paliurine H

[403842-57-9]

P-28



$C_{33}H_{51}N_5O_6$ 613.796


Alkaloid from the stems of *Paliurus ramosissimus*. Amorph. powder. $[\alpha]_D^{30}$ -412 (c, 0.24 in MeOH). λ_{max} 264 (log ϵ 4.32); 317 (log ϵ 4.05) (MeOH).

Lee, S.-S. *et al.*, *Phytochemistry*, 2001, **58**, 1271-1276 (*isol, pmr, cmr*)

Paliurine I

[403842-58-0]

P-29



$C_{36}H_{49}N_5O_6$ 647.813

Alkaloid from the stems of *Paliurus ramosissimus*. Amorph. powder. $[\alpha]_D^{30}$ -374.3 (c, 1.1 in MeOH). λ_{max} 266 (log ϵ 4.09); 317 (log ϵ 3.93) (MeOH).

N²-Deacyl: Nummularine S

[120396-83-0]

$C_{29}H_{36}N_4O_5$ 520.627

Alkaloid from the stem bark of *Zizyphus nummularia* (Rhamnaceae). Shows antibacterial and antifungal props. Mp 210-211°. Stereochem. not determined. May have different stereochem. to Paliurine I.

N²-Deacyl, N²-di-Me: Xylopyrine A

[1010795-95-5]

$C_{31}H_{40}N_4O_5$ 548.681

Alkaloid from the root bark of *Zizyphus xylopyra*. Cryst. (MeOH). $[\alpha]_D^{25}$ -270 (c, 0.21 in $CHCl_3$). λ_{max} 270 (log ϵ 2.8); 320 (log ϵ 2.26) (MeOH).

Shah, A.H. *et al.*, *Phytochemistry*, 1989, **28**, 305 (*Nummularine S*)


Lee, S.-S. *et al.*, *Phytochemistry*, 2001, **58**, 1271-1276 (*Paliurine I*)

Singh, A.K. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 1114-1120 (*Xylopyrine A*)

Pallidiflorine†

[133740-60-0]

P-30



$C_{34}H_{40}N_2O_7$ 588.699


Alkaloid from the whole plant of *Narcissus pallidiflorus* (Amaryllidaceae). Yellow cryst.

Codina, C. *et al.*, *Phytochemistry*, 1990, **29**, 2685 (*isol, ir, pmr, cmr, ms, struct*)

Pallidinine

5,6-Didehydro-2-hydroxy-3,6-dimethoxy-17-methylmorphinan-7-one, 9CI

P-31



(+)-form

$C_{19}H_{23}NO_4$ 329.395

(+)-form

Alkaloid from the leaves of *Dehaasia longipedicellata*. $[\alpha]_D^{25}$ +45 (c, 1 in $CHCl_3$).

λ_{\max} 205 (log ϵ 4.45); 228 (sh) ; 260 (log ϵ 3.99) (MeOH).

(-)-form [77485-11-1]

Alkaloid from *Ocotea acutangula* (Lauraceae). Mp 234-236°. $[\alpha]_{\text{D}}^{20}$ -80 (c, 0.5 in CHCl_3).

Me ether: O-Methylpallidine

[77485-12-2]

$\text{C}_{20}\text{H}_{25}\text{NO}_4$ 343.422

Alkaloid from *Ocotea acutangula* (Lauraceae). Noncryst. Mp 195-200° (as hydrochloride). $[\alpha]_{\text{D}}^{20}$ -50 (MeOH) (hydrochloride).

(±)-form

Me ether: [88199-99-9]

Synthetic. Mp 198-202° (as hydrochloride).

Vecchiotti, V. et al., *J.C.S. Perkin I*, 1981, 578 (isol, uv, ir, pmr, ms, struct)

McMurry, J.E. et al., *J.O.C.*, 1984, **49**, 3803 (deriv, synth, ir, pmr, cmr)


Kano, S. et al., *J.A.C.S.*, 1986, **108**, 6746 (deriv, synth)

Hanada, K. et al., *Org. Lett.*, 2002, **4**, 4515-4517 (synth)

Mukhtar, M.R. et al., *Fitoterapia*, 2004, **75**, 792-794 ((+)-form, pmr, cmr)

Pallimamine**Taurine†**

[123064-70-0]



Relative Configuration

$\text{C}_{23}\text{H}_{27}\text{NO}_5$ 397.47

(-)-form


Alkaloid from *Corydalis solidia* ssp. *tauricola*. Yellow cryst. Mp 179-182°. $[\alpha]_{\text{D}}^{22}$ -21.4 (c, 0.02 in CHCl_3).

(±)-form

Alkaloid from the whole herb of *Corydalis pallida* var. *sparsimamma*. Pale yellow prisms. Mp 203-207°.

Lu, S.-T. et al., *Phytochemistry*, 1989, **28**, 1245 (isol, uv, ir, pmr, ms, cryst struct)

Sener, B. et al., *Planta Med.*, 1990, **56**, 510 (isol)

Palmanine

$\text{C}_{21}\text{H}_{21}\text{NO}_7$ 399.399

(±)-form [95456-42-1]

Alkaloid from the stems and twigs of *Berberis actinacantha* (Berberidaceae). Amorph.

Valencia, E. et al., *Tetrahedron*, 1984, **40**, 3957 (isol, uv, ir, pmr, ms, struct)

Palmatine


5,6-Dihydro-2,3,9,10-tetramethoxydibenzo[*a,g*]quinolinizinium (1+), 9CI.

7,8,13,13a-Tetrahydro-2,3,9,10-tetrahydroxyberbinium (1+), 8CI. Berbericinine.

Hindarinine. Gindarinine

[3486-67-7]

[10605-02-4]



$\text{C}_{21}\text{H}_{22}\text{NO}_4^{\oplus}$ 352.409

Hindarinine is a variant of Gindarinine, via a Russian paper. Quaternary alkaloid from *Jateorhiza palmata* (Menispermaceae) and a large number of other spp., mostly in *Berberis* and *Mahonia* (Berberidaceae), but spread into many families. Uterine contractant. Shows bactericidal activity. Produces antiarrhythmic, inotropic, adrenocorticotrophic, anticholinesterase and analgesic effects in exp. animals. Yellow needles (as chloride). Mp 205° dec. (chloride). λ_{\max} 228 (ϵ 24400); 240 (sh) (ϵ 21000); 268 (ϵ 24100); 276 (sh) (ϵ 24000); 343 (sh) (ϵ 24400); 350 (ϵ 25000); 433 (ϵ 5100) (EtOH) (chloride).

▶ DR9890000

O²-De-Me: Dehydroisocorypalmine. Columbamine

[3621-36-1]

$\text{C}_{20}\text{H}_{20}\text{NO}_4^{\oplus}$ 338.382

Quaternary alkaloid from *Jateorhiza palmata*, several *Berberis* spp. (Menispermaceae, Berberidaceae) and some other spp. in several different families. Butyrylcholinesterase inhibitor; shows anti-HIV activity. Antihypertensive agent.

Possesses bactericidal, CNS depressant props. Shows antiplasmodial, antiameobic and cytotoxic activities.

Mp 239-240° (as chloride). Log P -0.57 (uncertain value) (calc). λ_{\max} 225 (ϵ 6025); 263 (ϵ 5750); 275 (ϵ 5000); 346 (ϵ 4500); 425 (ϵ 1250) (MeOH) (Berdy).

O³-De-Me: Dehydrocorypalmine. Jatrorrhizine. Jateorhizine. Neprotine. Yatrotrazine

[3621-38-3]

$\text{C}_{20}\text{H}_{20}\text{NO}_4^{\oplus}$ 338.382

Quaternary alkaloid from the root of *Jateorhiza palmata*, many *Berberis* and *Mahonia* spp. and many other spp. in several families (Menispermaceae, Berberidaceae). Butyrylcholinesterase inhibitor. Shows antibacterial activity vs. *Mycobacterium smegmatis*. Antiinflammatory, antihypertensive agent. Shows CNS stimulant props. Copper-coloured needles (as chloride). Mp 206° (chloride). Log P -0.57 (uncertain value) (calc). λ_{\max} 228 (ϵ 23400); 241 (sh) (ϵ 20800); 267

(ϵ 22600); 275 (sh) (ϵ 21000); 352 (ϵ 25200); 440 (ϵ 5200) (EtOH) (chloride).

O⁹-De-Me: Palmatrubine

[16176-68-4]

[16705-04-7]

$\text{C}_{20}\text{H}_{20}\text{NO}_4^{\oplus}$ 338.382

Alkaloid from *Fibraurea chloroleuca* and *Stephania glabra* tubers

(Menispermaceae). First known as a demethylation prod. of Palmatine, P-34. Shows anticancer activity.

Derivs. are specific xanthine oxidase inhibitors. Mp 290-295° (as chloride).

O¹⁰-De-Me: Dehydrocorydalmine

[6877-27-6]

$\text{C}_{20}\text{H}_{20}\text{NO}_4^{\oplus}$ 338.382

Alkaloid from a *Corydalis* spp.

(Papaveraceae) and from tubers of *Stephania glabra* (Menispermaceae).

Cryst. + 1H₂O (as chloride). Mp 220-221° dec. (chloride).

O²,O³-Di-de-Me: Demethyleneberberine

[25459-91-0]

[16705-03-6]

$\text{C}_{19}\text{H}_{18}\text{NO}_4^{\oplus}$ 324.355

Quaternary alkaloid from the stem, roots and tubers of *Stephania miyiensis* (Menispermaceae) and *Thalictrum javanicum*

(Ranunculaceae). Mp 225° (as

chloride). λ_{\max} 235 (log ϵ 4.63); 270 (log ϵ 4.52); 325 (log ϵ 4.49)

(MeOH) (chloride). λ_{\max} 225 (log ϵ 4.68); 263 (log ϵ 4.56); 340 (log ϵ 4.51) (MeOH/HCl) (chloride). λ_{\max}

238 (log ϵ 4.61); 274 (log ϵ 4.48); 330 (log ϵ 4.54) (MeOH/NaOH)

(chloride).

O²,O⁹-Di-de-Me: Dehydroscoulerine

[102694-43-9]

$\text{C}_{19}\text{H}_{18}\text{NO}_4^{\oplus}$ 324.355

Alkaloid from stem bark of *Pseuduvaria indochinensis* (Annonaceae).

Orange-red needles (MeOH) (counterion not specified). Mp 275-276° dec.

(counterion not specified). λ_{\max} 229 (log ϵ 4.22); 279 (log ϵ 4.27); 350 (log ϵ 4.21) (MeOH).

O²,O¹⁰-Di-de-Me: Stephanarine. Dehydrostepholidine

[17369-30-1]

[13509-87-0]

$\text{C}_{19}\text{H}_{18}\text{NO}_4^{\oplus}$ 324.355

Alkaloid from tubers of *Stephania glabra* and from *Stephania intermedia*

and *Tinospora capillipes*

(Menispermaceae). Cryst. + 1H₂O (as chloride). Mp 274-275° (chloride).

O³,O⁹-Di-de-Me: Jatrorrhizubine. Jatrorrubine

[38695-43-1]

[78673-37-7]

Synthetic.

O³,O¹⁰-Di-de-Me: Dehydrodiscretamine

[78134-82-4]

$\text{C}_{19}\text{H}_{18}\text{NO}_4^{\oplus}$ 324.355

Alkaloid from *Corydalis tashiroi*

whole plants and twigs of *Fissistigma balansae* (Annonaceae). Pale orange

needles (as chloride). Mp 217-220° dec. (chloride). λ_{\max} 206 ; 230 ; 286 ;


356 (MeOH).

- Feist, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1925, **263**, 294-305 (*Palmatrubine*)
 Späth, E. *et al.*, *Ber.*, 1925, **58**, 1939-1946 (*Columbamine, Jatrorrhizine, struct*)
 Haworth, R.D. *et al.*, *J.C.S.*, 1927, 548-554 (*Palmatine, synth*)
 Cava, M.P. *et al.*, *J. Nat. Prod.*, 1965, **28**, 73-83 (*isol, Columbamine, Jatrorrhizine*)
 Doskotch, R.W. *et al.*, *J.O.C.*, 1967, **32**, 3253-3254 (*Dehydrocorydalmine, Stepharanine*)
 Skerl, A.R. *et al.*, *Phytochemistry*, 1971, **10**, 2719-2721 (*biosynth*)
 Jewers, K. *et al.*, *J.C.S. Perkin 2*, 1972, 1393-1396 (*pmr*)
 Pavelka, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3157-3169 (*uv*)
 Patra, A. *et al.*, *Planta Med.*, 1980, **40**, 333-336 (*Palmatrubine, Stepharanine*)
 Tani, C. *et al.*, *Planta Med.*, 1981, **41**, 403-405 (*Dehydrodiscretamine*)
 Bahadur, S. *et al.*, *J. Nat. Prod.*, 1983, **46**, 454-457 (*Demethyleneberberine*)
 Bhakuni, D.S. *et al.*, *Tetrahedron*, 1983, **39**, 4003 (*Stepharanine, biosynth*)
 Zhong, S.-M. *et al.*, *Phytochemistry*, 1988, **27**, 4004-4005 (*Dehydroscoulerine*)
 Hussain, R.A. *et al.*, *Heterocycles*, 1989, **29**, 2257-2260 (*pmr, cmr*)
 Tan, G.T. *et al.*, *J. Nat. Prod.*, 1991, **54**, 143-154 (*Columbamine, anti-HIV activity*)
 Ghosh, R. *et al.*, *Acta Cryst. C*, 1993, **49**, 1665-1667 (*Jatrorrhizine, cryst struct*)
 Chia, Y.-C. *et al.*, *Phytochemistry*, 1998, **48**, 367-369 (*Dehydrodiscretamine*)
 Wright, C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1638-1640 (*Columbamine, activity*)
 Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr*)
 Grycová, L. *et al.*, *Phytochemistry*, 2007, **68**, 150-175 (*rev*)

Palmatine chloroform

P-35

[50932-23-5]



$C_{22}H_{22}Cl_3NO_4$ 470.778
 Alkaloid artifact from the roots of *Berberis lycium* (Berberidaceae). Cryst. (C_6H_6). Mp 184°.

2,3-Di-O-de-Me, 2,3-methylene ether: *Berberine chloroform*. 8-Trichloromethylidihydroberberine [50932-22-4]

$C_{21}H_{18}Cl_3NO_4$ 454.736
 Alkaloid artifact from the roots of *Berberis lycium*, *Thalictrum longistylum*, *Thalictrum lucidum* and *Thalictrum podocarpum* and from stem bark of *Phellodendron wilsonii* (Berberidaceae, Ranunculaceae, Rutaceae). Yellow cryst. ($CHCl_3$). Mp 182-184° dec. (178°).

- Miana, G.A. *et al.*, *Phytochemistry*, 1973, **12**, 1822 (*isol, uv, pmr, struct*)
 Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 204; 249; 1977, **40**, 281; 384 (*isol, uv, ir, pmr, deriv*)

- Khamidov, I.I. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 889-893; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 876-879 (*Berberine chloroform, cryst struct*)
 Marek, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 481-486 (*pmr, cmr, N-15 nmr*)

Palmatisine

P-36


[1360-20-9]

$C_{22}H_{25}NO_6$ 399.443
 Struct. unknown. Diterpenoid alkaloid. MF may be $C_{24}H_{27}NO_7$. Alkaloid from the roots of *Aconitum palmatum* (Ranunculaceae). Mp 285-286°. *Picrate*: Mp 164°. *Reineckate*: Mp 202° dec. *AuCl₃ complex*: Mp 220-225°. Singh, N. *et al.*, *J. Indian Chem. Soc.*, 1965, **42**, 49-50 (*isol, uv, ir*)

Palmerolide A

P-37

[863116-48-7]



$C_{33}H_{48}N_2O_7$ 584.751
 Macrolide antibiotic. Structure revised in 2007 by 3 different groups. Isol. from *Synocicum adareanum*. Cytotoxic. Amorph. solid. $[\alpha]_D^{24}$ -1.6 (c, 0.5 in MeOH). λ_{max} 224 (ε 2670); 242 (ε 2800); 296 (ε 1775) (MeOH).

- Diyabalanage, T. *et al.*, *J.A.C.S.*, 2006, **128**, 5630-5631 (*isol, pmr, cmr*)
 Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 5896-5899 (*synth, struct*)
 Jiang, X. *et al.*, *J.A.C.S.*, 2007, **129**, 6386-6387 (*synth, struct*)
 Lebar, M.D. *et al.*, *Tet. Lett.*, 2007, **48**, 8009-8010 (*struct*)

Palmidrol, INN

P-38

N-(2-Hydroxyethyl)hexadecanamide, 9CI. N-(2-Hydroxyethyl)palmitamide. *Impulsin*. *Palmitan*. *PEA* [544-31-0]

$H_3C(CH_2)_{14}CONHCH_2CH_2OH$
 $C_{18}H_{37}NO_2$ 299.496
 Isol. from soybean lecithin, egg yolk and peanut meal. Antiinflammatory agent, immunostimulant. Cryst. (EtOH aq.). Mp 98-99°. Log P 6 (calc).


- Roe, E.T. *et al.*, *J.A.C.S.*, 1952, **74**, 3442 (*synth*)
 Kuehl, F.A. *et al.*, *J.A.C.S.*, 1957, **79**, 5577 (*isol*)
 Benvenuti, F. *et al.*, *Boll. Soc. Ital. Biol. Sper.*, 1968, **44**, 809; *CA*, **69**, 85237e (*pharmacol*)
 Perlik, F. *et al.*, *Acta Phytopathol. Acad. Sci. Hung.*, 1971, **39**, 395 (*pharmacol*)
 Miyasaka, T. *et al.*, *Chem. Lett.*, 1985, **6**, 701 (*synth*)

Martindale, The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1397

Paludosine

P-39

[34137-24-1]



Absolute configuration

$C_{31}H_{45}NO_8$ 559.698
 Alkaloid from *Hammarbya paludosa* (Orchidaceae). Amorph. $[\alpha]_D^{23}$ +9 (c, 0.58 in EtOH). Hydrol. → Lindelofine in T-443.

N-Me: *Kumokirine*

[21284-20-8]
 $C_{32}H_{48}NO_8^{\oplus}$ 574.733
 Quaternary alkaloid from *Liparis kumokiri* and *Liparis kumokiri* (Orchidaceae). $[\alpha]_D^{20}$ -23.4 (as chloride).

2''-O-L-Arabinofuranosyl: *Nervosine*†

[23179-26-2]
 $C_{36}H_{53}NO_{12}$ 691.814
 Alkaloid from *Liparis nervosa*. Cryst. + 1H₂O (as picrate). Mp 130-131° (picrate). Conts. an arabinofuranosyl(1→2)glucosyl residue of undetd. anomeric config.

1-Epimer: *Auriculine*†


[22595-00-2]
 $C_{31}H_{45}NO_8$ 559.698
 Alkaloid from *Liparis auriculata* and *Liparis loeselii* (Orchidaceae). Amorph. $[\alpha]_D^{20}$ -19.9 (MeOH). $[\alpha]_D^{20}$ -14 (c, 0.56 in EtOH). Hydrol. → Laburnine in H-629.

- Leander, K. *et al.*, *Tet. Lett.*, 1967, 3477 (*isol*)
 Nishikawa, K. *et al.*, *Tetrahedron*, 1969, **25**, 2723 (*isol, struct*)
 Lindström, B. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 895 (*isol, struct*)

Palustrine†

P-40

1,4,5,6,7,8,9,10,11,13,14,16a-Dodecahydro-13-(1-hydroxypropyl)pyrido[2,1-d][1,5,9]triazacyclotridecin-2(3H)-one, 9CI [22324-44-3]



$C_{17}H_{31}N_3O_2$ 309.451
 Struct. revised in 1984. Alkaloid from *Equisetum palustre*, *Equisetum arvense*, *Equisetum limosum*, *Equisetum silvaticum*

and *Equisetum ramossissimum* (Equisetaceae). Prisms (Et₂O). Mp 120-122°. [α]_D¹⁸ +15.8 (c, 1.2 in H₂O). [α]_D²² +19.4 (c, 1.6 in EtOH).

Monohydrochloride:

Cryst. + 1H₂O. Mp 150-152°.

Dihydrochloride: Mp 188-190° dec. [α]_D¹⁸ +8.3 (c, 1.48 in H₂O).

Dipicrate:

Yellow prisms (Me₂CO aq.). Mp 150-150.5°.

N⁸-Formyl: Palustridine

[22324-43-2]

C₁₈H₃₁N₃O₃ 337.461

Alkaloid from *Equisetum palustre* (Equisetaceae). Cryst. (EtOH) (as hydrochloride). Mp 204° dec. (hydrochloride). [α]_D¹⁹ +50.2 (c, 1.235 in H₂O) (hydrochloride).

Deoxy: Deoxypalustrine. Alkaloid P3†

C₁₇H₃₁N₃O 293.451

Minor alkaloid from *Equisetum palustre* (Equisetaceae). Struct. prob. follows from the revised struct. of Palustrine.

15,16-Dihydro, 13,14-didehydro, 1'-ketone: Albizzine A

[163634-03-5]

C₁₇H₂₉N₃O₂ 307.435

Alkaloid from bark of *Albizia myriophylla* (Fabaceae). Oil. λ_{\max} 237 (no solvent reported).

Eugster, C.H. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 1387-1400 (*isol, ir*)

Green, C.L. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 673-678 (*Palustridine, ms, ir, pmr*)

Eugster, C.H. *et al.*, *Heterocycles*, 1976, **4**, 51-105 (*Deoxypalustrine*)

Rüedi, P. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 899-904 (*cmr, struct*)

Mayer, C. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 905-921 (*struct*)

Wälchli, P.C. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 921-928 (*abs config*)

Natsume, M. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3789-3791 (*synth, struct*)


Ito, A. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1966-1967 (*Albizzine A*)

Hirai, Y. *et al.*, *J.O.C.*, 1997, **62**, 776-777 (*synth*)

Palythazine

P-41

1,3,4,6,8,9-Hexahydrodipyrano[3,4-b:3',4'-c]pyrazine-3,8-dimethanol, 9CI [72681-96-0]



C₁₂H₁₆N₂O₄ 252.269

Isol. from the zoanthid *Palythoa tuberculosa*. Fp 169-170 (natural) Mp 223-225° (synthetic). [α]_D²⁵ -199 (MeOH) (synthetic). The abs. config. of (-)-Palythazine is as shown but it is not known whether this is the natural enantiomer.


Uemura, D. *et al.*, *Chem. Lett.*, 1979, 1481-1482 (*isol, uv, cmr, struct, synth*)

Jarglis, P. *et al.*, *Angew. Chem., Int. Ed.*, 1982, **21**, 141-142 (*struct, synth*)

Palytoxin

NSC 714361

[77734-91-9]



C₁₂₉H₂₂₃N₃O₅₄ 2680.168

Prod. by *Palythoa tuberculosa*, *Palythoa toxica* and *Radianthus macrodactylus*; also found in a variety of marine organisms that graze on coral reefs. Toxic. Na/K-ATPase inhibitor, sperm motility inhibitor, cardiotoxic and haemolytic agent, coronary vasoconstrictor. Sol. Py, DMSO, H₂O; fairly sol. MeOH, EtOH; poorly sol. CHCl₃, Me₂CO. 2 forms of Palytoxin are known, the one shown and an acetal form, C₁₂₉H₂₂₁N₃O₅₃. Struct. shown is for Palytoxin from a Tahitian *P. sp.* Palytoxins from *P. toxica* have slightly differing structs. λ_{\max} 233 ; 263 (€ 23600) (H₂O) (Berdy).

► Highly toxic, LD₅₀ 62.5 ng/kg (crab); LD₅₀ (mus, ivn) 0.45 mg/kg; LD₅₀ (mus, ipr) 0.05 mg/kg.

73-Deoxy: Deoxypalytoxin

[96391-17-2]

C₁₂₉H₂₂₃N₃O₅₃ 2664.169

Prod. by *Palythoa tuberculosa*. λ_{\max} 233 ; 263 (H₂O).

Homologue (n = 3): Homopalytoxin

[96391-18-3]

C₁₃₀H₂₂₅N₃O₅₄ 2694.195

Prod. by *Palythoa tuberculosa*. λ_{\max} 233 ; 263 (H₂O).

Homologue (n = 4): Bishomopalytoxin

[96411-43-7]

C₁₃₁H₂₂₇N₃O₅₄ 2708.222

Prod. by *Palythoa tuberculosa*. λ_{\max} 233 ; 263 (H₂O).

[11077-03-5]

Moore, R.E. *et al.*, *J.A.C.S.*, 1981, **103**, 2491-2494; 1982, **104**, 3776-3779 (*struct, abs config*)

Beress, L. *et al.*, *Toxicol.*, 1983, **21**, 285 (*isol*)
Uemura, D. *et al.*, *Tetrahedron*, 1985, **41**, 1007-1017 (*isol, struct, rev*)

Armstrong, R.W. *et al.*, *J.A.C.S.*, 1989, **111**, 7525; 7530 (*synth*)

Kishi, Y. *et al.*, *Pure Appl. Chem.*, 1993, **65**, 771-778 (*rev, conformn*)

Gleibs, S. *et al.*, *Toxicol.*, 1995, **33**, 1531-1537; 1999, **37**, 1521-1527 (*occur, distribution*)

Nicolaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods, VCH*, 1996, 711 (*bibl, synth*)

Food Sci. Technol., Seafood and Freshwater Toxins, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (revs)

Kan, Y. *et al.*, *Tet. Lett.*, 2001, **42**, 3197-3202 (*pmr, cmr*)

Kishi, Y. *et al.*, *Tetrahedron*, 2002, **58**, 6239-6258 (*rev*)

P-42

Inuzuka, T. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 897-899 (*struct*)


Inuzuka, T. *et al.*, *Tetrahedron*, 2008, **64**, 7718-7723 (*pmr, conformn*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PAF000

Pamamycin

P-43

[71892-94-9]



Pamamycin 593 R¹ = R³ = R⁴ = CH₃, R² = R⁵ = H

593B R¹ = CH₂CH₃, R² = R⁵ = H, R³ = R⁴ = CH₃, N-de-Me

607 R¹ = R² = R³ = R⁴ = CH₃, R⁵ = H

621A R¹ = R² = R³ = R⁴ = R⁵ = CH₃

621B R¹ = CH₂CH₃, R² = R³ = R⁴ = R⁵ = CH₃

621C R¹ = R² = R⁴ = CH₃, R³ = CH₂CH₃, R⁵ = H

621D R¹ = CH₂CH₃, R² = R³ = R⁴ = CH₃, R⁵ = H

635A R¹ = R² = R⁴ = R⁵ = CH₃, R³ = CH₂CH₃

635B R¹ = R² = R³ = R⁵ = CH₃, R⁴ = CH₂CH₃

635C R¹ = CH₂CH₃, R² = R³ = R⁴ = R⁵ = CH₃

635D R¹ = R⁴ = CH₂CH₃, R² = R³ = CH₃, R⁵ = H

635E R¹ = R² = CH₃, R³ = R⁴ = CH₂CH₃, R⁵ = H

635F R¹ = R² = CH₂CH₃, R³ = R⁴ = CH₃, R⁵ = H

649A R¹ = R³ = R⁴ = CH₂CH₃, R² = CH₃, R⁵ = H

649B R¹ = R³ = CH₂CH₃, R² = R⁴ = R⁵ = CH₃

Prod. by *Streptomyces* spp. Active against gram-positive and mycobacteria and fungi. Inhibits the membrane transportation of nucleosides. Ionophore. Sol. MeOH, hexane; poorly sol. H₂O.

Pamamycin 593 [171370-57-3]

C₃₄H₅₉NO₇ 593.843

Prod. by *Streptomyces alboniger*.

Pamamycin 593B

N-Demethylpamamycin 593B

[226722-70-9]

C₃₄H₅₉NO₇ 593.843

Prod. by *Streptomyces alboniger*.

Misleading synonym.

Pamamycin 607 [100905-89-3]

C₃₅H₆₁NO₇ 607.869

From *Streptomyces alboniger*. Oil.

Sol. MeOH, C₆H₆, MeCN, Py;

poorly sol. H₂O. [α]_D³³ +22.8 (c, 0.26

in MeOH). λ_{\max} 215 (€ 470)

(MeOH).

N-De-Me: N-Demethylpamamycin 593A.

Pamamycin 593A

[226722-69-6]

C₃₄H₅₉NO₇ 593.843

Prod. by *Streptomyces alboniger*. Misleading synonym.

Pamamycin 621A [149146-57-6]

C₃₆H₆₃NO₇ 621.896

Prod. by *Streptomyces aurantiacus* and *Streptomyces alboniger*.

Pamamycin 621B [171370-58-4]

C₃₆H₆₃NO₇ 621.896

Prod. by *Streptomyces alboniger*.

Pamamycin 621C [171370-59-5]

C₃₆H₆₃NO₇ 621.896

Prod. by *Streptomyces alboniger*.

Pamamycin 621D [171370-60-8]

C₃₆H₆₃NO₇ 621.896
Prod. by *Streptomyces alboniger*.

Pamamycin 635A [135702-84-0]

C₃₇H₆₅NO₇ 635.923
From *Streptomyces alboniger*. Sol.
EtOAc; poorly sol. H₂O.

Pamamycin 635B [135702-85-1]

C₃₇H₆₅NO₇ 635.923
From *Streptomyces alboniger*. Sol.
EtOAc; poorly sol. H₂O.

Pamamycin 635C [171370-61-9]

C₃₇H₆₅NO₇ 635.923
Prod. by *Streptomyces alboniger*.

Pamamycin 635D [171370-62-0]

C₃₇H₆₅NO₇ 635.923
Prod. by *Streptomyces alboniger*.

Pamamycin 635E [171370-63-1]

C₃₇H₆₅NO₇ 635.923
Prod. by *Streptomyces alboniger*.

Pamamycin 635F [171370-64-2]

C₃₇H₆₅NO₇ 635.923
Prod. by *Streptomyces alboniger*.

Pamamycin 649A [135702-86-2]

C₃₈H₆₇NO₇ 649.95
From *Streptomyces alboniger*. Sol.
EtOAc; poorly sol. H₂O.

Pamamycin 649B [135702-87-3]

C₃₈H₆₇NO₇ 649.95
From *Streptomyces alboniger*. Sol.
EtOAc; poorly sol. H₂O.

Pamamycin 663

C₃₉H₆₉NO₇ 663.977
Prod. by *Streptomyces* sp. HKI-0118.
Struct. not fully known. A higher
homologue with variation in the N-
containing side-chain.

Pamamycin 677

C₄₀H₇₁NO₇ 678.003
Prod. by *Streptomyces* sp. HKI-0118.
Struct. not fully known. A higher
homologue with variation in the N-
containing side-chain.

Pamamycin 691

C₄₁H₇₃NO₇ 692.03
Prod. by *Streptomyces* sp. HKI-0118.
Struct. not fully known. A higher
homologue with variation in the N-
containing side-chain.

Pamamycin 705

C₄₂H₇₅NO₇ 706.057
Prod. by *Streptomyces* sp. HKI-0118.
Struct. not fully known. A higher
homologue with variation in the N-
containing side-chain.

McCann, P.A. *et al.*, *J. Antibiot.*, 1979, **32**, 673
(*Pamamycin*)
Chou, N.G. *et al.*, *Antimicrob. Agents
Chemother.*, 1981, **20**, 443 (*Pamamycin*)
U.S. Pat., 1981, 4 283 391; *CA*, **95**, 148706
(*Pamamycin*)

Kondo, S. *et al.*, *Tet. Lett.*, 1987, **28**, 5861

(*Pamamycin 607*)
Kondo, S. *et al.*, *J. Antibiot.*, 1988, **41**, 1196

(*Pamamycin 607*)
Natsume, M. *et al.*, *Chem. Comm.*, 1989, 1911

(*Pamamycin 607*)
Natsume, M. *et al.*, *Tet. Lett.*, 1991, **32**, 3087

(*isol, struct*)
Graefe, U. *et al.*, *Nat. Prod. Lett.*, 1993, **3**, 265

(*Pamamycin 621*)
Natsume, M. *et al.*, *J. Antibiot.*, 1995, **48**,

1159-1164 (*isol*)
Haertl, A. *et al.*, *J. Antibiot.*, 1998, **51**, 1040-

1046 (*homologues*)
Kozone, I. *et al.*, *J. Antibiot.*, 1999, **52**, 329-331

(*N-Demethylpamamycin 593*)
Lee, E. *et al.*, *J.A.C.S.*, 2001, **123**, 10131-10132

(*Pamamycin 607, synth*)
Germay, O. *et al.*, *Tet. Lett.*, 2001, **42**, 4969-

4974 (*Pamamycin 607, synth*)
Wang, Y. *et al.*, *Tet. Lett.*, 2001, **42**, 7801-7804

(*Pamamycin 607, synth*)
Kang, S.H. *et al.*, *Angew. Chem., Int. Ed.*,

2002, **41**, 1392-1395 (*Pamamycin 607, synth*)
Jeong, E.J. *et al.*, *J.A.C.S.*, 2002, **124**, 14655-

14662 (*Pamamycin 607, synth*)
Fischer, P. *et al.*, *Angew. Chem., Int. Ed.*, 2005,

44, 6231-6234 (*Pamamycin 621A, 635B,
synth*)

Hashimoto, M. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 2005, **69**, 315-320 (*biosynth*)


Hashimoto, M. *et al.*, *J. Antibiot.*, 2005, **58**,

722-730 (*biosynth*)

Panacosmine

P-44

[250343-49-8]


C₂₁H₃₁N₃O 341.495

Alkaloid from the seeds of *Acosmium
panamense*. Amorph. yellow solid. [α]_D²⁵
+45 (c, 1 in CHCl₃). λ_{max} 245 ; 318
(MeOH).

Nuzillard, J.-M. *et al.*, *Tetrahedron*, 1999, **55**,
11511-11518

Panaefluoroline A

P-45

R = -CH(COO[⊖])CH(OH)CH₃C₁₉H₂₃NO₅ 345.394

Isol. from the cultured mycobiont of the
lichen *Amygdalaria panaeola*. Yellowish-
green solid. [α]_D²² +265.6 (c, 0.4 in
MeOH). λ_{max} 222 (log ε 4.08); 255 (log ε
4.14); 304 (log ε 3.46); 315 (log ε 3.38);
412 (log ε 3.71) (MeOH).

Kinoshita, K. *et al.*, *Tet. Lett.*, 2003, **44**, 8009-
8011 (*isol, pmr, cmr*)

Panaefluoroline B

P-46

As Panaefluoroline A, P-45 with

R = -CH₂COO[⊖]C₁₇H₁₉NO₄ 301.341

Isol. from the cultured mycobiont of the
lichen *Amygdalaria panaeola*. Yellowish-
green solid. [α]_D²² +52.3 (c, 0.57 in
MeOH). λ_{max} 218 (log ε 4.06); 256 (log ε
4.15); 305 (log ε 3.29); 317 (log ε 3.33);
412 (log ε 3.65) (MeOH).

Kinoshita, K. *et al.*, *Tet. Lett.*, 2003, **44**, 8009-
8011 (*isol, pmr, cmr*)

Panaefluoroline C

P-47

As Panaefluoroline A, P-45 with

R = -CH(CH₃)COO[⊖]C₁₈H₂₁NO₄ 315.368

Isol. from the cultured mycobiont of the
lichen *Amygdalaria panaeola*. Yellowish-
green solid. [α]_D²² +264.7 (c, 1 in MeOH).

Kinoshita, K. *et al.*, *Tet. Lett.*, 2003, **44**, 8009-
8011 (*isol, pmr, cmr*)

Panaefluoroline D

P-48

As Panaefluoroline A, P-45 with

R = -CH(COO[⊖])CH₂OHC₁₈H₂₁NO₅ 331.368

Isol. from the cultured mycobiont of
the lichen, *Amygdalaria panaeola*. Yellow
cryst. (CHCl₃/MeOH/Me₂CO). Mp
152° dec. [α]_D²⁵ +286.6 (c, 0.61 in
MeOH). Relative config. has been
determined. Fluorescent. λ_{max} 218 (log
ε 4.49); 258 (log ε 4.55); 304 (log ε 3.7);
315 (log ε 3.67); 412 (log ε 4.01)
(MeOH).

Kinoshita, K. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1723-1727 (*isol, pmr, cmr, cryst struct*)

Panaefluoroline E

P-49

As Panaefluoroline A, P-45 with

R = -CH(COO[⊖])CH₂CH(CH₃)₂C₂₁H₂₇NO₄ 357.449

Isol. from the cultured mycobiont of the
lichen *Amygdalaria panaeola*. Amorph.
yellow-green solid. [α]_D²⁵ +197.4 (c, 0.27 in
MeOH). Fluorescent. λ_{max} 211 (log ε
4.2); 260 (log ε 4.03); 304 (log ε 3.26); 317
(log ε 3.16); 413 (log ε 3.42) (MeOH).

Kinoshita, K. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1723-1727 (*isol, pmr, cmr*)

Panaefluoroline F

P-50

As Panaefluoroline A, P-45 with

R = -CH(COO[⊖])CH(CH₃)₂C₂₀H₂₅NO₄ 343.422

Isol. from the cultured mycobiont of the
lichen *Amygdalaria panaeola*. Amorph.
yellow-green solid. [α]_D²⁵ +228.9 (c, 0.28 in
MeOH). Fluorescent. λ_{max} 212 (log ε
4.21); 248 (sh) (log ε 3.97); 263 (log ε
4.03); 305 (sh) (log ε 3.35); 317 (sh) (log ε
3.22); 415 (log ε 3.45) (MeOH).

Kinoshita, K. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1723-1727 (*isol, pmr, cmr*)

Panaefluoroline G

P-51


As Panaefluoroline A, P-45 with
R = -CH(COO[⊖])CH₂C₆H₄OH(*p*-)C₂₄H₂₅NO₅ 407.465Isol. from the cultured mycobiont of
the lichen *Amygdalaria panaeola*.
Amorph. yellow-green solid. [α]_D²⁴ -110.3
(c, 0.31 in MeOH). Fluorescent. λ_{max}
214 (log ε 4.16); 258 (log ε 4.1); 313
(log ε 3.33); 414 (log ε 3.47)
(MeOH).Kinoshita, K. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1723-1727 (*isol, pmr, cmr*)**Panaefluoroline H**

P-52

As Panaefluoroline A, P-45 with
R = -CH(COO[⊖])CH₂CONH₂C₁₉H₂₂N₂O₅ 358.393Isol. from the cultured mycobiont of
the lichen *Amygdalaria panaeola*.
Amorph. yellow-green solid. [α]_D²⁵
+132.7 (c, 0.22 in MeOH).Fluorescent. λ_{max} 215 (log ε 4.27); 258
(log ε 4.26); 304 (sh) (log ε 3.35); 315
(sh) (log ε 3.3); 415 (log ε 3.62)
(MeOH).Kinoshita, K. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1723-1727 (*isol, pmr, cmr*)**Panamine, 9CI**

P-53

[2448-27-3]



(-)-form


C₂₀H₃₃N₃ 315.501Alkaloid from *Ormosia* spp. (Fabaceae).
Oil. Mp 285-287° (as diperchlorate). [α]_D
-11. Autoxidises in air.N-Me: Mp 103°. [α]_D +5.**Stereoisomer: Ormosajine**

[11004-96-9]


C₂₀H₃₃N₃ 315.501Alkaloid from *Ormosia jamaicensis*
(Fabaceae). Cryst. + 4H₂O. Mp 38-41°.
[α]_D +23. Config. unknown.**Stereoisomer, dipicrate:**Cryst. + H₂O. Mp 178°.Lloyd, H.A. *et al.*, *J.A.C.S.*, 1958, **80**, 1506-
1510 (*isol*)Hassall, C.H. *et al.*, *J.C.S.*, 1964, 2657-2663
(*isol, Ormosajine*)Wilson, E.M. *et al.*, *Tetrahedron*, 1965, **21**,
2561-2571 (*struct*)Karle, I.L. *et al.*, *Tet. Lett.*, 1966, 1659-1661
(*cryst struct*)Davies, A.P. *et al.*, *Tet. Lett.*, 1966, 6291-6294
(*struct, Ormosajine*)Cannon, J.R. *et al.*, *Tet. Lett.*, 1974, 1683-1686
(*abs config*)Liu, H.-J. *et al.*, *Can. J. Chem.*, 1976, **54**, 97-
109 (*synth*)Marcei, T.H. *et al.*, *J. Magn. Reson.*, 1982, **48**,
158-163 (*cmr*)Bhacca, N.S. *et al.*, *J.A.C.S.*, 1983, **105**, 2538-
2544 (*cmr, pmr*)**Pancoridine**

P-54


[39945-37-4]

C₁₉H₁₅NO₄ 321.332Alkaloid from the roots of *Corydalis*
paniculigera and *Corydalis stricta*, and
from the bark of *Popowia pisocarpa*
(Papaveraceae, Annonaceae). Orange
needles (CHCl₃/Et₂O). Mp 214° dec Mp
236-238°.Kupchan, S.M. *et al.*, *J.A.C.S.*, 1973, **95**, 4062
(*synth, uv, ir, pmr, ms*)Cava, M.P. *et al.*, *J.O.C.*, 1973, **38**, 2394 (*synth*,
uv, ir, pmr, ms)Alimova, M. *et al.*, *Khim. Prir. Soedin.*, 1982,
18, 727; *Chem. Nat. Compd. (Engl. Transl.)*,
1982, **18**, 689 (*isol*)Irgashev, T. *et al.*, *Khim. Prir. Soedin.*, 1983,
19, 490; *Chem. Nat. Compd. (Engl. Transl.)*,
1983, **19**, 461 (*isol*)Jossang, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1028
(*isol, pmr*)**Pancorine**

P-55


14-Methoxy[1,3]benzodioxolo[5,6-c]-
1,3-dioxolo[4,5-i]phenanthridine, 9CI. 6-
Methoxy-2,3;7,8-bis(methylenedioxy)-
benzo[c]phenanthridine. 8-Methoxynor-
sanguinarine
[80559-46-2]C₂₀H₁₃NO₅ 347.326Doubtful structural assignment; spectral
and physical props. of authentic synthetic
8-Methoxynorsanguinarine differ from
those reported for natural Pancorine.
Alkaloid from *Argemone mexicana* and
Corydalis paniculigera. Cryst. (CHCl₃/
EtOH). Mp 254-256°.N-Me: **8-Methoxysanguinarine**. N-**Methylpancorine**C₂₁H₁₆NO₅[⊕] 362.361Quaternary alkaloid from the rhizomes
of *Zanthoxylum nitidum*. Yellow pow-
der (as chloride). Mp 172-183° (chlor-
ide). λ_{max} 235 (log ε 4.35); 285 (log ε
4.58); 324 (sh) (log ε 4.17); 343 (log ε
4.2); 355 (log ε 4.35); 419 (log ε 2.99);
508 (log ε 3.23) (EtOH).Alimova, M. *et al.*, *Khim. Prir. Soedin.*, 1981,
17, 671; *CA*, **96**, 48970aSuau, R. *et al.*, *An. Quim., Ser. C*, 1988, **84**,
294Chang, Y.-C. *et al.*, *Z. Naturforsch., C*, 2003,
58, 521-526 (*isol, pmr, ms*)Cui, X.-G. *et al.*, *Helv. Chim. Acta*, 2008, **91**,
155-158 (8-Methoxysanguinarine)**Pancorinine**

P-56

7-Amino-2,9,10-trimethoxy-1H-dibenzo[*f*-
de,g]quinolin-1-one, 9CI
[85011-55-8]C₁₉H₁₆N₂O₄ 336.346Unusual 7-aminoaporphine alkaloid.
However studies by Kilmer *et al.* (2003)
suggest a struct. revision is possible
similar to that of Telazoline, T-62.Alkaloid from the roots of *Corydalis*
paniculigera and *Corydalis stricta*. Cryst.
(CHCl₃/MeOH). Mp 289-290°.Alimova, M. *et al.*, *Chem. Nat. Compd. (Engl.*
Transl.), 1982, **18**, 689 (*isol*)Kilmer, L. *et al.*, *J. Nat. Prod.*, 2003, **66**, 115-
118 (*struct*)**Pancracine**

P-57

[21416-14-8]



(-)-form

C₁₆H₁₇NO₄ 287.315Alkaloid from *Pancretium maritimum*,
Narcissus poeticus, *Hippeastrum bifida*
and *Hippeastrum vittatum* (Amaryllida-
ceae). Elongated prisms (MeOH). Mp
272-273°. [α]_D²⁵ -74 (c, 0.02 in MeOH).
λ_{max} 241 (log ε 3.64); 292 (log ε 3.72)
(EtOH).**Perchlorate:** Mp 163-166°.**Picrate:** Mp 249-252°.**Di-Ac:**Cryst. (C₆H₆/petrol). Mp 163-165°.**O²-Me: Montanine**†. **O²-Methylpancra-****cine**

[642-52-4]

C₁₇H₁₉NO₄ 301.341Alkaloid from the bulbs of *Haem-*
anthus montanus, *Haemanthus amar-*
ylloides, *Haemanthus coccineus* and
two unidentified *Haemanthus* spp.,
Haemanthus tigrinus and *Hippeastrum*
bifida (Amaryllidaceae). Weak hypo-
tensive and convulsive agent (dog). Oil;
microscopic needles + 1H₂O (H₂O).
Mp 88-89° (hydrate). [α]_D²⁶ -87.6 (c, 0.57
in CHCl₃). [α]_D²⁵ -70.2 (c, 1.03 in
CHCl₃). Also forms Me₂CO and
CHCl₃ solvates, Mp 57-60° and 59-65°
resp. λ_{max} 244 (log ε 3.65); 297 (log ε
3.71) (no solvent reported).▶ LD₅₀ (dog, ivn) 42 mg/kg.

O²-Me, perchlorate:
Prisms. Mp 249-250° dec. $[\alpha]_D^{26}$ -18.17
(c, 1.23 in MeOH).

O²-Me, picrate:
Yellow needles (H₂O). Mp 225-226°
dec.

O²-Me, Ac: **O-Acetylmontanine**
C₁₉H₂₁NO₅ 343.379
Trace alkaloid from the bulbs of
Hippeastrum bifida and *Rhodophiala*
bifida (Amaryllidaceae). Mp 210-213°
(as hydrochloride). CAS No not assigned to
natural isomer to 14CI but refers to racemic
material.

Di-Me ether: **Manthine**
[606-51-9]
C₁₈H₂₁NO₄ 315.368
Alkaloid from *Haemanthus amarylloides*
and *Haemanthus tigrinus* (Amaryllidaceae).
Cryst. (Et₂O). Mp 114-116°. $[\alpha]_D^{26}$ -71.3 (c, 0.47 in
CHCl₃). λ_{\max} 243 (log ϵ 3.64); 294 (log ϵ 3.71)
(no solvent reported).

2-Epimer, O²-Me: **Coccinine**
[485-57-4]
C₁₇H₁₉NO₄ 301.341
Alkaloid from *Haemanthus coccineus* and a
few other *Haemanthus* spp. (Amaryllidaceae).
Shows convulsive action in high dose.
Mp 162-163°. $[\alpha]_D^{27}$ -189 (c, 1.9 in EtOH).
 λ_{\max} 213 (log ϵ 4.25); 244 (log ϵ 3.58);
296 (log ϵ 3.64) (no solvent reported).

► Toxic, LD₅₀ (dog, ivn) 17.5 mg/kg.

2-Epimer, O²-Me, perchlorate: Mp 254-
255° dec.

2-Epimer, O²-Me, N-Me:
C₁₈H₂₂NO₄⁺ 316.376
Mp 223-224° (as iodide). $[\alpha]_D^{25}$ -60.5 (c,
1.4 in H₂O) (iodide).

3-Epimer: **Brunsvigine**
[1354-81-0]
C₁₆H₁₇NO₄ 287.315
Alkaloid from *Brunsvigia cooperi* and
Brunsvigia radulosa (Amaryllidaceae).
Needles by subl. Mp 243°. $[\alpha]_D^{20}$ -76.6
(c, 1 in EtOH).

3-Epimer, picrate: Mp 219° (190°, 195-
196° dec.).

3-Epimer, di-Ac:
Needles (Et₂O/petrol). Mp 184°.

3-Epimer, O²-Me: **Manthidine**
[43208-82-8]
C₁₇H₁₉NO₄ 301.341
Isol. from *Hippeastrum coccineus* and
from two unidentified *Hippeastrum*
spp. Cryst. by subl. Mp 269-270°. $[\alpha]_D^{21.5}$
-26.6 (c, 0.6 in CHCl₃). λ_{\max} 240
(log ϵ 3.63); 294 (log ϵ 3.72) (no solvent
reported).

Wildman, W.C. et al., *J.A.C.S.*, 1955, **77**, 1248-
1252 (*Manthine*, *Coccinine*, *Manthidine*,
Montanine, *isol*)

Briggs, L.H. et al., *Anal. Chem.*, 1957, **29**, 904-
911 (*Montanine*, *Manthidine*, *ir*)

Dry, L.J. et al., *J.C.S.*, 1958, 4701-4704
(*Brunsvigine*, *isol*)

Inubushi, Y. et al., *J.O.C.*, 1960, **25**, 2153-2164
(*Manthine*, *Montanine*, *Brunsvigine*, *struct*)

Duffield, A.M. et al., *J.A.C.S.*, 1965, **87**, 4902-
4912 (*Coccinine*, *Montanine*, *ms*)

Wildman, W.C. et al., *Pharmazie*, 1967, **22**, 725
(*O-Acetylmontanine*)

Wildman, W.C. et al., *J.A.C.S.*, 1968, **90**, 6439-
6446 (*Pancracine*, *Montanine*, *isol*, *uv*, *ms*,
struct, *bibl*)

Fuganti, C. et al., *Chem. Comm.*, 1973, 430-
431 (*Manthidine*, *biosynth*)

Clark, R.C. et al., *Tetrahedron*, 1975, **31**, 1855-
1859 (*Brunsvigine*, *struct*)

Ali, A.A. et al., *Planta Med.*, 1984, **50**, 188-189
(*uv*, *pmr*, *cmr*, *cd*, *stereochem*)

Overman, L.E. et al., *J.O.C.*, 1991, **56**, 5005-
5007 (*synth*)

Ishizaki, M. et al., *J.O.C.*, 1992, **57**, 7285-7295
(*Pancracine*, *Brunsvigine*, *Manthine*,
Coccinine, *Montanine*, *O-Acetylmontanine*,
synth)

Ishizaki, M. et al., *J.C.S. Perkin 1*, 1993, 101-
110 (*Pancracine*, *Coccinine*, *Montanine*,
synth)

Overman, L.E. et al., *J.O.C.*, 1993, **58**, 4662-
4672 (*synth*)

Jin, J. et al., *J.A.C.S.*, 1997, **119**, 5773-5784
(*Montanine*, *Coccinine*, *Brunsvigine*, *synth*)

Labrana, J. et al., *Phytochemistry*, 2002, **60**,
847-852 (*isol*, *cmr*)

Pandey, G. et al., *Org. Lett.*, 2005, **7**, 3713-
3716 (*synth*)


Banwell, M.G. et al., *Org. Lett.*, 2007, **9**, 3503-
3506 (*synth*)

Hong, A.-W. et al., *J.O.C.*, 2008, **73**, 7580-7585
(*Brunsvigine*, *Manthine*, *synth*)

Pancratistatin

NSC 349156
[96203-70-2]
[120962-18-7 ((±)-form)]

P-58



C₁₄H₁₅NO₈ 325.274
Alkaloid from the roots and bulbs of
Hymenocallis littoralis (*Pancratium lit-*
torale) and bulbs of *Haemanthus kal-*
breyeri. Also isol. from the Texas
grasshopper *Brachystola magna*. Pos-
sesses antineoplastic activity. Shows
good activity against P-388 leukaemia
cells and M5076 sarcoma. Cryst.
(DMF/MeOH/Et₂O). Sol. DMSO; fairly
sol. MeOH; poorly sol. Me₂CO, hexane.
Mp 322-324°. $[\alpha]_D^{34}$ +48 (c, 1.0 in
DMSO). λ_{\max} 209 (sh) ; 219 (sh) ; 233
(ϵ 20900); 278 (ϵ 8130); 308 (sh)
(MeOH) (Derep).

2-O- β -D-Glucopyranoside: **Pancratistide**
[121849-70-5]

C₂₀H₂₅NO₁₃ 487.416
Alkaloid from the bulbs of *Ha-*
emanthus kalbreyeri (Amaryllidaceae).
Light brown hygroscopic solid. Mp
203-218° (foaming, dec.). $[\alpha]_D^{28}$ +22.5
(c, 1.2 in H₂O).

Penta-Ac: Mp 162-166°.

1-O-(3-Hydroxybutanoyl): **1-(3-Hydro-**
xybutanoyl)pancratistatine
[212953-15-6]

C₁₈H₂₁NO₁₀ 411.365
Alkaloid from the bulbs of *Zephyr-*

anthes carinata. Cytotoxic agent. $[\alpha]_D^{25}$
+19.2 (c, 0.2 in MeOH). λ_{\max} 231
(log ϵ 4.05); 279 (log ϵ 3.7)
(MeOH).

1-O-(3- β -D-Glucopyranosyloxybutanoyl):
1-(3-Glucopyranosyloxybutanoyl)pan-
cratistatine
[212953-16-7]

C₂₄H₃₁NO₁₅ 573.507
Alkaloid from the bulbs of *Zephyr-*
anthes carinata. Cytotoxic agent. $[\alpha]_D^{25}$
+20.7 (c, 0.5 in MeOH). λ_{\max} 235 (log ϵ
3.84); 272 (log ϵ 3.57) (MeOH).

O⁷-Me: Mp 294-298° dec. $[\alpha]_D^{25}$ +289.9 (c,
0.69 in DMSO).

7-Deoxy: **7-Deoxypancratistatine**
[83603-30-9]

C₁₄H₁₅NO₇ 309.275
Alkaloid from *Haemanthus kalbreyeri*
(Amaryllidaceae). Phytotoxin. Straw-
coloured solid. Mp 310-314° dec. $[\alpha]_D^{28}$
+53.7 (c, 0.42 in DMSO). λ_{\max} 222 (ϵ
19950); 262 (sh) (ϵ 4700); 270 (ϵ
7580); 305 (ϵ 3235) (MeOH)
(Berdy).

7-Deoxy, 1-O-(2-acetamido-2-deoxy- β -D-
glucopyranoside): **Telastaside**
[133056-61-8]

C₂₂H₂₈N₂O₁₂ 512.469
Alkaloid from the insect *Polytela*
gloriosa which feeds on various Amaryl-
lidaceae. Mp 214-218° (dec.) (as per-
Ac). $[\alpha]_D^{28}$ +78.5 (c, 0.8 in H₂O). λ_{\max}
222 (sh) ; 233 (log ϵ 4.02); 260 (sh) ; 272
(log ϵ 3.58); 305 (log ϵ 3.14) (MeOH).

Pettit, G.R. et al., *J. Nat. Prod.*, 1986, **49**, 995-
1002; 1995, **58**, 37-43 (*Pancrastatin*, *isol*, *uv*,
ir, *pmr*, *ms*, *synth*, *cryst struct*)

Danishefsky, S. et al., *J.A.C.S.*, 1989, **111**,
4829-4837 (*synth*, *ir*, *pmr*)

Ghosal, S. et al., *Phytochemistry*, 1989, **28**,
611-613 (*Pancrastide*, 7-
Deoxypancratistatine)

Ghosal, S. et al., *J. Chem. Res., Synop.*, 1990,
334-335 (*Telastaside*)

Tian, X. et al., *J.A.C.S.*, 1995, **117**, 3643-3644
(*synth*)

Trost, B.M. et al., *J.A.C.S.*, 1995, **117**, 10143-
10144 (*synth*)

Tian, X. et al., *Synlett*, 1995, 1125-1126 (*synth*,
7-*Deoxypancratistatine*)

Chida, N. et al., *Heterocycles*, 1996, **43**, 1385-
1390 (7-*Deoxypancratistatine*, *synth*)

Doyle, T.J. et al., *Tetrahedron*, 1997, **53**, 11153-
11170 (*synth*)

Keck, G.E. et al., *J.O.C.*, 1998, **63**, 9164-9165
(7-*Deoxypancratistatine*, *synth*)

Kojima, K. et al., *Phytochemistry*, 1998, **48**,
1199-1202 (*butanoyl derivs*)

Magnus, P. et al., *Tetrahedron*, 1998, **54**, 15509-
15524 (*synth*)

Grubb, L.M. et al., *Tet. Lett.*, 1999, **40**, 2691-
2694 (*synth*, *bibl*)

Rigby, J.H. et al., *J.A.C.S.*, 2000, **122**, 6624-
6628 (*synth*)

Pettit, G.R. et al., *J.O.C.*, 2001, **66**, 2583-2587
(*synth*)

Hudlicky, T. et al., *J.O.C.*, 2002, **67**, 8726-8743
(*synth*)

Ko, H. et al., *J.O.C.*, 2004, **69**, 112-121 (*synth*)

Pettit, G.R. et al., *J. Nat. Prod.*, 2005, **68**,
1256-1258 (*isol*, *cryst struct*)


Li, M. et al., *Tet. Lett.*, 2006, **47**, 3707-3710
(*synth*)

Pettit, G.R. et al., *J. Nat. Prod.*, 2007, **70**, 417-
422 (*synth*)

Manpadi, M. et al., *Org. Prep. Proced. Int.*,
2008, **40**, 109-161 (rev)

Pandamarilactam 3x P-59

1-[4-(4-Methyl-5-oxo-2(5H)-furanlydene)butyl]-2-pyrrolidinone, 9CI
[189179-33-7]



C₁₃H₁₇NO₃ 235.282
Alkaloid from *Pandanus amaryllifolius*.
Food flavouring.

(Z)-Isomer: Pandamarilactam 3y

[189179-35-9]

C₁₃H₁₇NO₃ 235.282


Alkaloid from *Pandanus amaryllifolius*.
Food flavouring.

Sjaifullah, A. et al., *CA*, 1997, **126**, 297535w
(isol)

Takayama, H. et al., *Heterocycles*, 1999, **50**,
75-78 (synth)

Pandamarilactone 1 P-60

[152606-63-8]




C₁₈H₂₃NO₄ 317.384
Alkaloid from leaves of *Pandanus amaryllifolius* (Pandanaceae). Food flavouring. Yellow amorph. solid. [α]_D²³ -33 (c, 0.1 in MeOH).

Nonato, M.G. et al., *Phytochemistry*, 1993, **34**,
1159 (isol, uv, ir, pmr, cmr, struct)

Pandamarilactone 31 P-61

[152606-64-9]

R¹ = OMe, R² = CH₃C₁₉H₂₅NO₄ 331.411

Alkaloid from leaves of *Pandanus amaryllifolius* (Pandanaceae). Food flavouring. Amorph. solid. [α]_D²³ -2 (c, 0.1 in MeOH).

Nonato, M.G. et al., *Phytochemistry*, 1993, **34**,
1159 (isol, uv, ir, pmr, cmr, struct)

Pandamarilactone 32 P-62

[152606-65-0]

As Pandamarilactone 31, P-61 with
R¹R² = CH₂


C₁₈H₂₁NO₃ 299.369

Major alkaloid from leaves of *Pandanus amaryllifolius* (Pandanaceae). Food flavouring. Amorph. solid.

Nonato, M.G. et al., *Phytochemistry*, 1993, **34**,
1159 (isol, uv, ir, pmr, cmr, struct)

Pandamarilactonine A P-63

[303008-80-2]



Absolute
Configuration

C₁₈H₂₃NO₄ 317.384

Relative config. revised in 2002. Alkaloid from *Pandanus amaryllifolius*. Food flavouring. Amorph. powder. [α]_D²³ +35 (c, 4.4 in CHCl₃). Partial racemate. λ_{max} 207 (log ε 1.82); 232 (sh); 275 (log ε 2.43) (MeOH).

N-Dealkyl: 3-Methyl-5-(2-pyrrolidinyl)-2(5H)-furanone. **Norpandamarilactonine A**

[367941-40-0]

C₉H₁₃NO₂ 167.207

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder.
Racemic. λ_{max} 207 (log ε 2.29); 252
(log ε 0.35); 274 (log ε 0.44) (MeOH).

5E-Isomer: **Pandamarilactonine C**

[477332-25-5]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder.
[α]_D²³ +26.2 (c, 1 in CHCl₃). λ_{max} 201
(log ε 3.67); 220 (sh); 275 (log ε 4.23)
(MeOH).

15-Epimer: **Pandamarilactonine B**

[468097-76-9]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder.
Racemic. λ_{max} 207 (log ε 1.72); 232 (sh)
; 275 (log ε 2.49) (MeOH).

15-Epimer, N-oxide: **Pandamarilactonine B N-oxide**C₁₈H₂₃NO₅ 333.383

Alkaloid from *Pandanus amaryllifolius*.
Amorph. solid. Dec. readily. Probable

artifact.

15-Epimer, N-dealkyl: **Norpandamarilactonine B**

[367941-41-1]

C₉H₁₃NO₂ 167.207

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder.
Racemic. λ_{max} 207 (log ε 2.58); 253
(log ε 0.29); 274 (log ε 0.36) (MeOH).

15-Epimer, 5E-isomer: **Pandamarilactonine D**

[477332-26-6]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder.
Racemic. λ_{max} 203 (log ε 4.3); 234 (sh)
; 276 (log ε 4.38) (MeOH).

Takayama, H. et al., *J.A.C.S.*, 2000, **122**, 8635-
8639 (isol)

Takayama, H. et al., *J. Nat. Prod.*, 2001, **64**,
1224-1225 (Norpandamarilactonines)


Takayama, H. et al., *Chem. Pharm. Bull.*,
2002, **50**, 1303-1304 (isol, synth, pmr, cmr)

Blanco, P. et al., *Eur. J. Org. Chem.*, 2004, 48-
53 (synth)

Salin, A.A. et al., *J. Nat. Prod.*, 2004, **67**, 54-57
(Pandamarilactonine B N-oxide)

Takayama, H. et al., *Tet. Lett.*, 2005, **46**, 5795-
5797 (synth, abs config)

Honda, T. et al., *Tet. Lett.*, 2006, **47**, 6251-
6254 (synth)

Pandamarine P-64C₁₈H₂₅N₃O₂ 315.414**(±)-form** [145940-69-8]


Major alkaloid from leaves of
Pandanus amaryllifolius (Pandanaceae).
Plates (MeOH/EtOAc/Et₂O). Mp 210-
211°.

Byrne, L.T. et al., *Aust. J. Chem.*, 1992, **45**,
1903 (isol, uv, ir, pmr, cmr, ms, cryst
struct)

Pandanamine P-65

5,5'-(Iminodi-4-butanyl-1-ylidene)bis[3-
methyl-2(5H)-furanone], 9CI

[302914-53-0]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Pandanus amaryllifolius*.
Food flavouring. Amorph. powder. λ_{max}

203 (log ϵ 1.86); 242 (sh); 273 (log ϵ 1.15) (MeOH).

(E,E)-Isomer: **(E,E)-Pandamine**

[657389-58-7]
C₁₈H₂₃NO₄ 317.384

Minor alkaloid from *Pandanus amar-
yllifolius*.

Takayama, H. *et al.*, *J.A.C.S.*, 2000, **122**, 8635-8639 (*synth*)


Takayama, H. *et al.*, *Tet. Lett.*, 2001, **42**, 2995-2996 (*Z,Z-isomer*)

Salim, A.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 54-57 (*isol*)

Pandicine

P-66

[76282-39-8]



Probable absolute configuration

C₄₄H₅₀N₄O₇ 746.902


Alkaloid from the leaves of *Pandacastrum saccharatum* (preferred genus name *Tabernaemontana*) (Apocynaceae). Brown amorph. solid.

Kan-Fan, C. *et al.*, *J.O.C.*, 1981, **46**, 1481 (*uv, ir, pmr, cmr, ms, struct*)

Pandine

P-67

[55855-14-6]



Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Pandaca caducifolia* and other *Pandaca* spp.; also from *Ervatamia lifuana* and *Ervatamia orientalis* (Apocynaceae). Mp 108-113°. [α]_D +273 (MeOH).

Hoizey, M.J. *et al.*, *Phytochemistry*, 1974, **13**, 1994 (*isol, ir, uv, pmr, ms*)

LeMen, J. *et al.*, *Tet. Lett.*, 1974, 3119 (*ir, uv, pmr, cmr, struct*)


Ducruix, A. *et al.*, *Acta Cryst. B*, 1977, **33**, 1990 (*cryst struct*)

Bruneton, J. *et al.*, *Planta Med.*, 1980, **39**, 180 (*isol*)

Panganensine X

P-68

[182745-08-0]



Absolute Configuration

C₃₈H₄₆N₄O₂ 590.807

Alkaloid from root bark of *Strychnos panganensis*. [α]_D +14 (c, 0.18 in CHCl₃). The 16-configs. are opposite in the two halves of the molecule. λ _{max} 220 ; 247 ; 300 (MeOH).

19-Epimer: **Panganensine Y**

[182966-44-5]
C₃₈H₄₆N₄O₂ 590.807


From root bark of *Strychnos panganensis*. [α]_D +34 (c, 0.48 in CHCl₃). λ _{max} 208 ; 247 ; 301 (MeOH).

Nuzillard, J.-M. *et al.*, *Phytochemistry*, 1996, **43**, 897-902 (*isol, uv, ir, pmr, cmr, ms, struct*)

Panganensine R

R-Panganensine

[182817-43-2]



Absolute Configuration

C₃₈H₄₄N₄O₂ 588.791

Curan dimer. Alkaloid from root bark of *Strychnos panganensis*. [α]_D +66 (c, 0.28 in CHCl₃). λ _{max} 207 ; 247 ; 292 (MeOH).

19'-Epimer: **Panganensine S**. S-Panganensine

[182969-74-0]
C₃₈H₄₄N₄O₂ 588.791


From root bark of *Strychnos panganensis*. [α]_D +71 (c, 1.58 in CHCl₃). λ _{max} 206 ; 248 ; 295 (MeOH).

Nuzillard, J.-M. *et al.*, *Phytochemistry*, 1996, **43**, 897-902 (*isol, uv, ir, pmr, cmr, ms, struct*)

Paniculamine

P-70

[174545-79-0]



C₂₂H₃₅NO₅ 393.522

Alkaloid from roots of *Aconitum paniculatum*. Cryst. (MeOH or Py). Mp 222-224°.

17-O-(3,4-Dimethoxybenzoyl), N-deoxy:

C₃₁H₄₃NO₇ 541.683

Alkaloid from *Aconitum variegatum*. Amorph. solid. [α]_D²⁵ +30 (c, 0.11 in CHCl₃).

Yusupova, I.M. *et al.*, *Khim. Prir. Soedin.*,


1995, **31**, 277; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 228

Diaz, J.G. *et al.*, *Phytochemistry*, 2005, **66**, 837-846 (*Aconitum variegatum alkaloid*)

Paniculine†

P-71

10-Hydroxyacetyldihydrolycopodine. Alkaloid P2 [61328-14-1]



C₁₈H₂₉NO₃ 307.432

Alkaloid from *Lycopodium paniculatum* and *Lycopodium confertum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 65-66°.

O-De-Ac: **Deacetylpaniculine**

[73907-65-0]
C₁₆H₂₇NO₂ 265.395

Alkaloid from *Lycopodium paniculatum* and *Lycopodium confertum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 172-174°.

Deacetoxy, 4,5-didehydro: **Anhydrodeacetylpaniculine**. Alkaloid P5

[61328-12-9]
C₁₆H₂₅NO 247.38

Alkaloid from *Lycopodium paniculatum* (Lycopodiaceae). Cryst. (Me₂CO). Mp 175-185° dec.

Castillo, M. *et al.*, *Can. J. Chem.*, 1976, **54**, 2900 (*isol, ir, pmr, ms, Anhydrodeacetylpaniculine*)

Morales, G. *et al.*, *Phytochemistry*, 1979, **18**, 1719 (*Deacetylpaniculine*)

Muñoz, O. *et al.*, *Heterocycles*, 1982, **19**, 2287 (*pmr, cmr, struct*)

Garland, M.T. *et al.*, *J. Appl. Crystallogr.*, 1982, **15**, 112 (*Deacetylpaniculine, cryst struct*)


Manriquez, V. *et al.*, *Acta Cryst. C*, 1988, **44**, 165 (*cryst struct*)

Muñoz, O.M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 200 (*pmr, cmr, Paniculine, Deacetylpaniculine*)

Pannellin

P-72

[217481-12-4]



C₂₈H₂₆O₉ 506.508

Flavagline compd. Constit. of *Aglaia elaeagnoides* and *Aglaia edulis*. Shows insecticidal and antifungal activities. Mp 102-104°. [α]_D²⁰ -50 (c, 0.3 in

CHCl₃). λ_{max} 228 (sh) (log ε 4.11); 285 (sh) (log ε 3.64); 298 (log ε 3.73) (MeOH).

1-Ac: 1-O-Acetylpannellin

[217466-04-1]

C₃₀H₂₈O₁₀ 548.545

Constit. of *Aglaia elaeagnoidea*. Shows insecticidal activity. [α]_D²⁰ -65 (c, 0.1 in CHCl₃). λ_{max} 228 (sh) (log ε 4.08); 285 (sh) (log ε 3.63); 298 (log ε 3.71) (MeOH).

Parent acid, dimethylamide: Aglaroxin A

[176785-75-4]

C₂₉H₂₉NO₈ 519.55

Constit. of *Aglaia edulis* and *Aglaia roxburghiana*. Shows insecticidal and antifungal activities. [α]_D²⁰ -81 (c, 0.4 in CHCl₃).

Parent acid, dimethylamide, 1-Ac: 1-O-Acetylglaroxin A

C₃₁H₃₁NO₉ 561.587

Constit. of the bark of *Aglaia edulis*. Cytotoxic. Amorph. powder. [α]_D²² -69 (c, 0.1 in EtOH). λ_{max} 205 (log ε 4.84); 297 (log ε 3.73) (EtOH).

3'-Methoxy: 3'-Methoxypannellin

[217466-05-2]

C₂₉H₂₈O₁₀ 536.534

Constit. of *Aglaia elaeagnoidea*. [α]_D²⁰ -45 (c, 0.1 in CHCl₃). λ_{max} 228 (sh) (log ε 3.74); 285 (log ε 3.75); 298 (sh) (log ε 3.67) (MeOH).

3'-Methoxy, parent acid, dimethylamide: Aglaroxin B

[176785-76-5]

C₃₀H₃₁NO₉ 549.576

Constit. of *Aglaia roxburghiana*.

3'-Methoxy, parent acid, dimethylamide, 1-Ac: 1-O-Acetyl-3'-methoxyglaroxin A

C₃₂H₃₃NO₁₀ 591.613

Constit. of the bark of *Aglaia edulis*. Cytotoxic. Amorph. powder. [α]_D²² -132 (c, 0.2 in MeOH). λ_{max} 208 (log ε 4.52); 284 (log ε 3.56) (MeOH).

3'-Methoxy, 2'-hydroxy, parent acid, dimethylamide: 2'-Hydroxyglaroxin B. Aglaroxin F

[251982-68-0]

C₃₀H₃₁NO₁₀ 565.576

Constit. of the stems of *Aglaia oligophylla* and *Aglaia roxburghiana*. Amorph. solid.

Pat. Coop. Treaty (WIPO), 1996, 96 04 284; *CA*, **124**, 335673h (*Aglaroxin B*)

Brader, G. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1482-1490 (*isol, uv, ir, pmr, cmr, ms, 1-Ac*)

Molloyres, L.-P. *et al.*, *Pestic. Sci.*, 1999, **55**, 486-503 (*Aglaroxins*)

Bacher, M. *et al.*, *Phytochemistry*, 1999, **52**, 253-263 (*Aglaroxin A*)

Engelmeier, D. *et al.*, *J. Agric. Food Chem.*, 2000, **48**, 1400-1404 (*isol, activity*)

Dreyer, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 415-420 (*Aglaroxin A, abs config*)


Bringmann, G. *et al.*, *J. Nat. Prod.*, 2003, **66**, 80-85 (*2'-Hydroxyglaroxin B*)

Koul, O. *et al.*, *Entomol. Exp. Appl.*, 2005, **114**, 197-204 (*Aglaroxin A, activity*)

Kim, S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1769-1775 (*1-Acetylglaroxin A, 1-Acetyl-3'-methoxyglaroxin A*)

Pantherinine

[152606-66-1]



C₁₅H₈BrN₃O 326.152

Alkaloid from the ascidian *Aplidium pantherinum*. Exhibits mild cytotoxicity against P388 murine leukaemia cells. Purple powder. λ_{max} 254 (ε 16110); 316 (ε 3950) (MeOH) (Berdy).

Ac:

Red glass.


Kim, J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1813-

1816 (*isol, uv, ir, pmr, cmr, ms, struct*)

Nakahara, S. *et al.*, *Tet. Lett.*, 1998, **39**, 5531-5522 (*synth*)

Pantocin A₂

[152606-66-1]




C₁₆H₂₁N₅O₆ 379.372

Minor metab. of *Pantoea agglomerans*. Antibiotic. [α]_D²⁵ -202 (c, 0.5 in H₂O). λ_{max} 201 (ε 9550); 273 (ε 4040) (no solvent reported).

Jin, M. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2898-2901; 2902-2905 (*isol, pmr, cmr, ms*)

Pantocin A†

[152606-66-1]



Absolute Configuration


C₁₃H₁₆N₄O₅ 308.293

Prod. by *Pantoea agglomerans* (formerly *Erwinia herbicola*). Antibiotic. [α]_D²⁵ -291.1 (c, 1.8 in H₂O). λ_{max} 200 (ε 12090); 272 (ε 5450) (H₂O).

Jin, M. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2898-2901; 2902-2905 (*isol, pmr, cmr, N-15 nmr*)

Panurensine

[55701-99-0]



P-73

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stems of *Abuta panurensis* (Menispermaceae). Cryst. (CH₂Cl₂/hexane). Mp 156-158°. [α]_D -245.6 (c, 0.5 in CHCl₃). Rapidly turns yellow on exp. to light and air. λ_{max} 225 (ε 25500); 240 (ε 39000); 288 (ε 20700) (EtOH/NaOH) (Derep). λ_{max} 224 (ε 24000); 239 (ε 35400); 286 (ε 16800) (EtOH) (Derep).

N²-De-Me: Norpanurensine. 2'-De-methylpanurensine

[55702-00-6]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Abuta panurensis* (Menispermaceae). Needles (MeOH). Mp 175°. [α]_D -250 (c, 0.1 in CHCl₃). λ_{max} 225 (ε 25500); 240 (ε 39000); 288 (ε 20700) (EtOH/NaOH) (Derep). λ_{max} 224 (ε 24000); 239 (ε 35400); 286 (ε 16800) (EtOH) (Derep).

Me ether:


Prisms (EtOAc/hexane). Mp 124-125°. [α]_D -210 (c, 0.05 in CHCl₃).

Cava, M.P. *et al.*, *J.O.C.*, 1975, **40**, 2647 (*isol, uv, ir, pmr, ms, struct*)

Papaveraldine

P-77

6,7-Dimethoxy-1-(3,4-dimethoxybenzoyl)isoquinoline. Xanthaline†
[522-57-6]



C₂₀H₁₉NO₅ 353.374

Isol. from *Papaver somniferum* preparations (opium) (Papaveraceae). Mp 210°. Probably an artifact.

Hydrochloride: Mp 198-199°.

Picrate: Mp 208-209°.

N-Me: N-Methylpapaveraldinium. N-Methylxanthalinium

[7668-80-6]

C₂₁H₂₂NO₅[⊕] 368.408

Quaternary alkaloid from *Stephania sasakii* (Menispermaceae). Mp 159.5-161° (as chloride). CAS no. refers to chloride.

3'-O-De-Me, N-Me: Thalprzewalskiinone
[219915-63-6]

C₂₀H₂₀NO₅[⊕] 354.382

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow cryst. (as iodide). Mp 196-197° (iodide). Revised struct. (2001). λ_{max} 256 (log ε 4.38); 302 (log ε 3.89); 326 (log ε 3.94) (MeOH).

Noller, C.R. *et al.*, *J.A.C.S.*, 1950, **72**, 17 (*uv*)


Guthrie, D.A. *et al.*, *Can. J. Chem.*, 1955, **33**, 729 (*synth*)

Kunitomo, J. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 456; *CA*, **65**, 10633g (*N-Methylpapaveraldinium*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1980, **43**, 143 (*ir, pmr, ms*)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 869 (*synth*)

Al-Howiriny, T.A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 823-826 (*Thalprzewalskiinone*)

Papaverberbine**P-78**

$C_{21}H_{23}NO_6$ 385.416

N-Me (α): N-Methylpapaverberbine

[131723-86-9]

$C_{22}H_{26}NO_6^{\oplus}$ 400.45


Alkaloid from *Papaver pseudo-orientale* (Papaveraceae). Amorph. (as iodide). $[\alpha]_D^{25}$ -133 (c, 0.09 in MeOH). CA no. refers to iodide.

[107118-04-7]

Sariyar, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1302 (*isol, uv, pmr, ms, struct*)

Papaverine, BAN**P-79**

6,7-Dimethoxy-1-(3,4-dimethoxybenzyl)-isoquinoline. Papaveroline tetramethyl ether. NSC 35443
[58-74-2]



$C_{20}H_{21}NO_4$ 339.39

Alkaloid from *Papaver somniferum* (opium poppy) and *Rauwolfia serpentina* (Papaveraceae, Apocynaceae). Used as 0.05M soln. in $CHCl_3$ for extraction-photometric detn. of Zr (λ_{max} 580 nm, ϵ 51000); flotation-photometric detn. of Te (with bromide). Antianginal, anticholinergic agent. Smooth muscle relaxant, vasodilator, antiasthmatic agent. Oral antispasmodic agent which has been used for treating gastrointestinal spasm. Shows anti-HIV activity. Prisms (EtOH/Et₂O). Mp 147-148°. pK_a 6.4 (25°). Log P 3.37 (calc). λ_{max} 239 (ϵ 67500); 279 (ϵ 7380); 314 (ϵ 4000); 327 (ϵ 4660) (EtOH) (Berdy).

► Gastrointestinal and other adverse effects reported when used therapeutically, can cause drowsiness and dizziness by ingestion. LD₅₀ (rat, orl) 325mg/kg. NW8450000

Hydrochloride: Papaverine hydrochloride, USAN. Cerespan. Pavabid. Brovon
[61-25-6]

Rods (H₂O). Mp 221-222° dec. Component of Copavin.

► Adverse therapeutic effects similar to

papaverine. LD₅₀ (rat, orl) 360 mg/kg. Exp. teratogen. NW8575000

N-Me: Densiberine. N-Methylpapaverine

[19716-71-3]

$C_{21}H_{24}NO_4^{\oplus}$ 354.425

Quaternary alkaloid from *Berberis densiflora*. Cryst. (EtOH) (as chloride). Mp 156-157° (chloride). CAS no. refers to chloride. λ_{max} 222 ; 256 ; 317 (EtOH).

α -Oxo: see Papaveraldine, P-77

α -Methoxy: **Setigerine. α -Methoxypapaverine**

[180386-76-9]

$C_{21}H_{23}NO_5$ 369.416

Minor alkaloid from *Papaver setigerum*. Racemic.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 878B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 457C (*nmr*)

Hofmann, A. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 849-865 (*uv*)

Guthrie, D.A. *et al.*, *Can. J. Chem.*, 1955, **33**, 729-742 (*synth*)

Popp, F.D. *et al.*, *J.A.C.S.*, 1957, **79**, 3773-3777 (*synth*)

Saferstein, R. *et al.*, *CA*, 1974, **80**, 44458m (*ms*)

Preininger, V. *et al.*, *Alkaloids (Academic Press)*, 1975, **15**, 209-261 (*pharmacol, tox*)

Brochmann-Hanssen, E. *et al.*, *J.C.S. Perkin 1*, 1975, 1531-1537 (*biosynth*)

Neuman, M. *et al.*, *Drugs of Today*

(*Barcelona*), 1976, **12**, 278 (*rev*)

Shtokalo, M.I. *et al.*, *Zh. Anal. Khim.*, 1976, **31**, 2366-2370; *J. Anal. Chem. USSR (Engl. Transl.)*, 1976, **31**, 1739-1742 (*detn, Zr*)

Whipple, G.H. *et al.*, *Angiology*, 1977, **28**, 737-749 (*rev*)

Tomimatsu, T. *et al.*, *CA*, 1977, **87**, 23573z (*pmr*)

Marsaioli, A.J. *et al.*, *Phytochemistry*, 1978, **17**, 1655-1658 (*cmr*)

Wilen, G. *et al.*, *J. Pharm. Pharmacol.*, 1982, **34**, 264-266 (*metab*)

Skripechuk, V.G. *et al.*, *Zh. Anal. Khim.*, 1983, **38**, 2198-2201; *J. Anal. Chem. USSR (Engl. Transl.)*, 1983, **38**, 1696-1699 (*detn, Te*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms, 6th edn.*, Akademie-Verlag, 1987, 5642 (*synonyms*)

Hifnawy, M.S. *et al.*, *Anal. Profiles Drug Subst.*, 1988, **17**, 367-447 (*rev*)

Turano, A. *et al.*, *AIDS Res. Hum. Retroviruses*, 1989, **5**, 183-192 (*anti-HIV activity*)

Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1989, **28**, 2833-2839 (*pmr, cmr*)

Hirsenkorn, R. *et al.*, *Tet. Lett.*, 1991, **32**, 1775-1778 (*synth*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1996, **61**, 1047-1052 (*Setigerine*)

Marek, J. *et al.*, *Z. Kristallogr., Kristallgeom., Kristallphys., Kristalchem.*, 1996, **211**, 649-650; 651-652 (*cryst struct, Papaverine, Setigerine*)

Khamidov, I.I. *et al.*, *Khim. Prir. Soedin.*, 1997, **33**, 420-423; *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 323-325 (*Densiberine*)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Martindale, *The Extra Pharmacopoeia, 32nd edn.*, Pharmaceutical Press, 1999, 1614-1615

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, PAH000; PAH250

Papaveroline, BAN, INN**P-80**

1-[(3,4-Dihydroxyphenyl)methyl]-6,7-isoquinolinediol, 9CI. 1-(3,4-Dihydroxybenzyl)-6,7-dihydroxyisoquinoline. Modus†. Udiaci active substance
[574-77-6]



$C_{16}H_{13}NO_4$ 283.283

Vasodilator; also t-RNA transferase inhibitor (potential antineoplastic agent). Cryst. powder + 2H₂O, becoming anhyd. at ca. 100°. Mp 150° dec. Log P 2.79 (calc).

Hydrochloride:

Needles + 1H₂O. Sol. hot H₂O. Mp 250° (dec.).

Hydroiodide:

Needles + 2H₂O (H₂O). Mp 230°.

6-Me ether: 1-(3,4-Dihydroxybenzyl)-7-hydroxy-6-methoxyisoquinoline. Turcomanine

[60372-11-4]

$C_{17}H_{15}NO_4$ 297.31

Alkaloid from *Berberis turcomanica* (Berberidaceae). Mp 247-248° (as hydrobromide). λ_{max} 236 ; 282 ; 312 (EtOH) (hydrobromide).

3',6-Di-Me ether: 7-Hydroxy-1-(4-hydroxy-3-methoxybenzyl)-6-methoxyisoquinoline. Cristadine

[82463-46-5]

$C_{18}H_{17}NO_4$ 311.337

Alkaloid from the leaves of *Erythrina crista-galli* (Leguminosae). Pale yellow. Mp 210-211°. Mp. initially reported as 128-129° which is incorrect.

4',6-Di-Me ether, N-Me: Tetradehydroreticuline

[267006-18-8]

$C_{19}H_{20}NO_4^{\oplus}$ 326.371

Quaternary alkaloid from *Stephania cepharantha*. Cryst. (MeOH) (as perchlorate). Mp 231-233°. λ_{max} 232 (log ϵ 4.39); 257 (log ϵ 4.59); 280 (log ϵ 4.36); 318 (log ϵ 3.86); 367 (3.49) (MeOH).

6,7-Di-Me ether: 1-(3,4-Dihydroxybenzyl)-6,7-dimethoxyisoquinoline. Turcomanidine

[16637-56-2]

$C_{18}H_{17}NO_4$ 311.337

Alkaloid from *Berberis turcomanica*. Cryst. (MeOH). Mp 199-200°.

6,7-Di-Me ether, hydrochloride: [16637-68-6]

Cryst. (EtOH). Mp 232-233°. λ_{\max} 227 (sh) (ϵ 25400); 253 (ϵ 55200); 288 (ϵ 6800); 311 (ϵ 8300); 327 (sh) (ϵ 6600); 345 (sh) (ϵ 5000) (EtOH).

6,7-Di-Me ether, di-Ac: Mp 224-226°.

4',6,7-Tri-Me ether: 1-(3-Hydroxy-4-methoxybenzyl)-6,7-dimethoxyisoquinoline. **Palaudine** [18694-10-5]

$C_{19}H_{19}NO_4$ 325.363

Minor alkaloid from opium (*Papaver somniferum*) (Papaveraceae). Prisms (EtOH). Mp 175-176°.

4',6,7-Tri-Me ether, N-Me: N-Methylpalaudinium

$C_{20}H_{22}NO_4^+$ 340.398

Quaternary alkaloid from *Thalictrum polygamum*. Needles (MeOH/Me₂CO) (as chloride). Mp 232-233° (chloride).

Tetra-Me ether: see Papaverine, P-79

Bis(methylene) ether: see Mollinedine, M-672

[23740-72-9]

Kitasato, Z. *et al.*, *J.C.S.*, 1932, 785 (*synth*)

Vitali, T. *et al.*, *Chimica*, 1952, 7, 409 (*synth*)

Brochmann-Hanssen, E. *et al.*, *J. Pharm. Sci.*, 1968, 57, 940 (*Palaudine*)

Brossi, A. *et al.*, *J.O.C.*, 1970, 35, 1684-1687 (*Turcomanidine*, *synth*, *uv*, *ir*, *pmr*, *derivs*)

Shamma, M. *et al.*, *J. Pharm. Sci.*, 1972, 61, 295-297 (*N-Methylpalaudinium*)

Cheng, C.C. *et al.*, *J. Pharm. Sci.*, 1972, 61, 645 (*use*)

Data, P.G. *et al.*, *Farmaco, Ed. Sci.*, 1981, 36, 302 (*pharmacol*)

Ju-ichi, M. *et al.*, *Heterocycles*, 1982, 19, 849 (*Cristadine*)

Kovács, L. *et al.*, *Acta Chim. Hung.*, 1985, 120, 103 (*synth*, *Cristadine*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., *Pharmaceutical Press*, 1993, 1312


Khamidov, I. *et al.*, *Khim. Prir. Soedin.*, 1996, 32, 74-76; 885-888; *Chem. Nat. Compd.*

(*Engl. Transl.*), 1996, 32, 59-61; 873-875 (*Turcomanine*, *Turcomanidine*)

Tanahashi, T. *et al.*, *Chem. Pharm. Bull.*, 2000, 48, 370-373 (*Tetrahydroreticuline*)

Papavoriendine P-81

7-Methoxy[1,3]benzodioxolo[4,5-c]-1,3-dioxolo[4,5-c]phenanthridine, 9CI [309275-04-5]




$C_{20}H_{13}NO_5$ 347.326

Alkaloid from the roots of *Papaver orientale*. Yellow-brown amorph. powder. Mp 158-160°. λ_{\max} 215 ; 220 ; 240 ; 368 (EtOH).

Ma, W.G. *et al.*, *Fitoterapia*, 2000, 71, 527-534

Papavorientine P-82

5,8,13,13a-Tetrahydro-2,11-dimethoxy-6H-dibenzo[a,g]quinolizin-9-ol



$C_{19}H_{21}NO_3$ 311.38


(S)-form

Alkaloid from the roots of *Papaver orientale*. Amorph. brown powder. Mp 169-171°. $[\alpha]_D^{20}$ -189 (c, 0.03 in EtOH). λ_{\max} 206 ; 220 ; 229 ; 258 (EtOH).

Ma, W.G. *et al.*, *Fitoterapia*, 2000, 71, 527-534

Papillozine C P-83

[207793-40-6]




$C_{27}H_{44}N_2O$ 412.657

Alkaloid from the leaves of *Buxus papillosa*. Amorph. solid. $[\alpha]_D^{20}$ +23 (c, 0.5 in CHCl₃). λ_{\max} 219 ; 237 ; 244 (MeOH).

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1998, 48, 519-528 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Papuamine P-84

[112455-84-2]



Absolute Configuration

$C_{25}H_{40}N_2$ 368.604

Alkaloid from a marine sponge *Haliclona* sp. Antifungal. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 167.5-169°. $[\alpha]_D$ -150 (c, 1.5 in MeOH). C₂-symmetric struct. λ_{\max} 241 (ϵ 3000) (MeOH) (Derep). λ_{\max} 236 ; 240 (ϵ 3000) (MeOH) (Berdy). λ_{\max} 240 (MeOH/HCl) (Berdy).

Hydrochloride (1:2): Mp 230° dec. $[\alpha]_D$ -140 (c, 1.3 in MeOH).

5a-Epimer: Haliclondiamine

[117065-24-4]

$C_{25}H_{40}N_2$ 368.604

Alkaloid from a marine sponge *Haliclona* sp. Shows antimicrobial activity. Oil; cryst. as di-Ac. Sol. MeOH. Mp 96-98° (di-Ac). $[\alpha]_D$ -18.2. λ_{\max} 241 (ϵ 3000) (MeOH) (Derep). λ_{\max} 241 (ϵ 10700) (MeOH) (Berdy).

Baker, B.J. *et al.*, *J.A.C.S.*, 1988, 110, 965 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Fahy, E. *et al.*, *Tet. Lett.*, 1988, 29, 3427 (*Haliclondiamine*)

Borzilleri, R.M. *et al.*, *J.A.C.S.*, 1995, 117, 10905 (*synth*, *abs config*)

Barrett, A.G.M. *et al.*, *J.O.C.*, 1996, 61, 685 (*synth*, *enantiomer*)

McDermott, T.S. *et al.*, *J.O.C.*, 1996, 61, 700 (*synth*)

Taber, D.F. *et al.*, *J.A.C.S.*, 1997, 119, 22 (*synth*, *Haliclondiamine*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, 30, 3-51 (*rev. synth*)

Adlington, R.M. *et al.*, *Tetrahedron*, 2000, 56, 623-628 (*synth*)

Papyraceabromine A P-85

[110313-17-2]

$C_{22}H_{17}Br_4N_3O_5$ 723.009


Struct. unknown. Metab. from the marine bryozoan *Fustra papyracea* (*Chartella papyracea*). Wide-spectrum *in vitro* antibacterial alkaloid. Unstable. Minor alkaloids of structurally different type, Papyraceabromine B (with 5 bromine atoms) and Papyraceabromine C, were also isol.

[110313-18-3 (Papyraceabromine B), 110313-19-4 (Papyraceabromine C)]

Pietra, F. *et al.*, *Gazz. Chim. Ital.*, 1985, 115, 443-485 (*rev*)

Papyramine P-86

[81149-33-9]



$C_{18}H_{23}NO_4$ 317.384

Mixt. of C-6 epimers at r.t. Alkaloid from the bulbs and aerial parts of *Narcissus papyraceus* (Amaryllidaceae).

Me ether: O-Methylpapyramine

[128517-01-1]

$C_{19}H_{25}NO_4$ 331.411

Alkaloid from aerial parts of *Narcissus papyraceus* (Amaryllidaceae). Amorph. $[\alpha]_D^{24}$ -31 (c, 0.1 in MeOH). Config. at C-6 is β (trans. to the ethane bridge).

3-Epimer: 3-Epipapyramine. 6-Hydroxy-3-O-methylpimaritidine

$C_{18}H_{23}NO_4$ 317.384

Alkaloid from the bulbs of *Narcissus tazetta* var. *chinensis* (Amaryllidaceae). Pale-yellow cryst. $[\alpha]_D^{20}$ -10.2 (c, 0.45 in CHCl₃). Isol. as an inseparable mixt. of C-6 epimers.

Song, G. *et al.*, *Fenxi Huaxue*, 1981, 9, 520; *CA*, 96, 162994m (*pmr*, *config*)

Hung, S. *et al.*, *Huaxue Xuebao*, 1981, 39, 529; *CA*, 96, 139653r (*isol*, *struct*)


Ma, G.-E. *et al.*, *Heterocycles*, 1986, 24, 2089 (*epimer*, *uv*, *ir*, *pmr*, *ms*)

Suau, R. *et al.*, *Heterocycles*, 1990, 31, 517 (*O-Methylpapyramine*)

Paraensidimerine A

P-87

[80450-21-1]

C₃₀H₃₀N₂O₄ 482.578

Alkaloid from the heartwood of *Euxylophora paraënsis* (Rutaceae). Needles (Me₂CO or CHCl₃/MeOH). Mp 311-312°. Racemic.

16,16a-Didehydro: Paraensidimerine G

[86168-91-4]

C₃₀H₂₈N₂O₄ 480.562

Isol. from *Euxylophora paraënsis* (Rutaceae). Needles (CHCl₃/MeOH). Mp 280-281°.

1,2,10,11-Tetramethoxy: Vepridimerine A

[82841-71-2]

C₃₄H₃₈N₂O₈ 602.683

Alkaloid from the bark of *Vepris louisii* (Rutaceae). Mp 343-345°. Racemic.

1,2,10,11-Tetramethoxy, 16,16a-didehydro: Araliopdimerine C

[119089-42-8]

C₃₄H₃₆N₂O₈ 600.667

Alkaloid from the bark of *Araliopsis tabouensis* (Rutaceae). Needles (hexane/CHCl₃). Mp 254-255°. Racemic.

6a-Epimer: Paraensidimerine F

[86196-11-4]

C₃₀H₃₀N₂O₄ 482.578

Isol. from *Euxylophora paraënsis* (Rutaceae). Small rhombic cryst. (CHCl₃/MeOH). Mp 310°. Racemic.

16a-Epimer: Paraensidimerine C

[80450-23-3]

C₃₀H₃₀N₂O₄ 482.578

Isol. from *Euxylophora paraënsis* (Rutaceae). Cryst. (C₆H₆); cryst. + 1H₂O (CHCl₃/MeOH). Mp 210° (rapid heat) (hydrate). Racemic.

16a-Epimer, N¹⁸-de-Me: N-Demethylparaensidimerine C

[953393-39-0]

C₂₉H₂₈N₂O₄ 468.551

Alkaloid from the stem wood of *Zanthoxylum integrifolium*. Needles (CHCl₃/MeOH). Mp 197-199°. Racemic. λ_{max} 228 (log ε 4.65); 276 (log ε 3.85); 285 (log ε 3.84); 318 (log ε 3.86) (MeOH).

16a-Epimer, 1,2,10,11-tetramethoxy: Vepridimerine B

[82864-60-6]

C₃₄H₃₈N₂O₈ 602.683

Alkaloid from the bark of *Vepris louisii* and *Oricia renieri* (Rutaceae). Mp 278-279°. Racemic.

6a,16a-Diepimer: Paraensidimerine E

[80450-24-4]

C₃₀H₃₀N₂O₄ 482.578

Isol. from *Euxylophora paraënsis*

(Rutaceae). Needles + 1H₂O

(Py/MeOH). Mp 289-290°. Racemic.

Jurd, L. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1625; 1982, **35**, 2505; 1983, **36**, 759 (*isol, cryst struct, ir, pmr, cmr, ms*)

Ngadjui, T.B. *et al.*, *Tet. Lett.*, 1982, **23**, 2041 (*Vepridimerines*)

Ayafor, J.F. *et al.*, *Tet. Lett.*, 1985, **26**, 4529 (*Vepridimerines, synth*)

Ngadjui, B. *et al.*, *Phytochemistry*, 1988, **27**, 2979 (*Araliopdimerine C*)


Ngadjui, B.T. *et al.*, *J. Nat. Prod.*, 1989, **52**, 300 (*synth, Paraensidimerines A, C, F*)

Chen, J.-J. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1444-1448 (*N-Demethylparaensidimerine C*)

Paraensidimerine B

P-88

[80450-22-2]

C₃₀H₃₂N₂O₅ 500.593

Alkaloid from the heartwood of *Euxylophora paraënsis* (Rutaceae). Needles (CHCl₃/MeOH). Mp 286-287°. Possibly formed by the addition of water to the exocyclic double bond of Paraensidimerine D, below.

1'-Dehydro, 2'-Deoxy: Paraensidimerine D

[80357-91-1]

C₃₀H₃₀N₂O₄ 482.578

Alkaloid from *Euxylophora paraënsis* (Rutaceae). Needles (CHCl₃/MeOH). Mp 259°. [α]_D²⁰ -0.38 (c, 1.3 in CHCl₃).


Jurd, L. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1625; 1982, **35**, 2505; 1983, **36**, 759 (*isol, ir, pmr, cmr, ms, struct*)

Ngadjui, B.T. *et al.*, *J. Nat. Prod.*, 1989, **52**, 300 (*synth*)

Paraensine†

P-89

3,9,10,15-Tetrahydro-5-methoxy-3,3-dimethyl-7H-indolo[2',3':3,4]pyrido[2,1-b]pyrano[2,3-h]quinazolin-7-one, 9CI [38750-10-6]

C₂₄H₂₁N₃O₃ 399.448

Alkaloid from the bark of *Euxylophora paraënsis* (Rutaceae). Cryst. (C₆H₆). Mp 281-282°.

Dihydro: [38750-12-8]


Cryst. (diisopropyl ether). Mp 305°.

Danieli, B. *et al.*, *Experientia*, 1972, **28**, 249 (*uv, ir, pmr, ms, struct, synth*)

Paraherquamide A

P-90

VM 29919. Antibiotic VM 29919 [77392-58-6]



Absolute Configuration

R¹ = OH, R² = ³⁰CH₃C₂₈H₃₅N₃O₅ 493.602

Alkaloid from *Penicillium paraherquei*, *Penicillium charlesii* and *Penicillium* sp. IMI332995. Nematocide, anthelmintic and insecticidal mycotoxin. Sol. MeOH, DMSO, EtOAc. Mp 244-247° dec. [α]_D²² -28 (c, 0.43 in MeOH). λ_{max} 226 (ε 32400); 260 (ε 6100); 290 (ε 1600) (EtOH) (Derep). λ_{max} 225 (log ε 4.5) (MeOH).

▶ Toxic to mice. RV0545000

N¹⁷-Oxide: Paraherquamide A N¹⁷-oxide.

VM 55596. Antibiotic VM 55596

[141137-59-9]

C₂₈H₃₅N₃O₆ 509.601

Prod. by a *Penicillium* sp. λ_{max} 226 (ε 32400); 260 (ε 6100); 290 (ε 1600) (EtOH) (Derep).

14-Deoxy: Paraherquamide E. VM 54159.

Antibiotic VM 54159

[125600-53-5]

C₂₈H₃₅N₃O₄ 477.602

Prod. by a *Penicillium* sp. Anthelmintic; possesses nematocidal props. λ_{max} 226 (ε 32400); 260 (ε 6100); 290 (ε 1600) (MeOH) (Derep). λ_{max} 225 (log ε 4.4) (MeOH).

14-Deoxy, N-de-Me: Antibiotic SB

200437. SB 200437

[142449-76-1]

C₂₇H₃₃N₃O₄ 463.575

Prod. by *Aspergillus* sp. IMI 337664. Anthelmintic agent.

14-Deoxy, 4-hydroxy: Antibiotic SB

203105. SB 203105

[199723-64-3]

C₂₈H₃₅N₃O₅ 493.602

Prod. by *Aspergillus* sp. IMI 337664. Anthelmintic agent.

14-Deoxy, 14,30-didehydro: Paraherquamide C

[125439-04-5]

C₂₈H₃₃N₃O₄ 475.586

Alkaloid from *Penicillium charlesii*. Nematocide, insecticide, mycotoxin. Sol. MeOH, EtOAc, DMSO. λ_{max} 226 (ε 32400); 260 (ε 6100); 290 (ε 1600) (MeOH) (Derep). λ_{max} 225 (MeOH).

14-Deoxy, 14,30-epoxy: Paraherquamide D

[125600-52-4]

C₂₈H₃₃N₃O₅ 491.586

Alkaloid from *Penicillium charlesii*. Acaricide, insecticide, nematocide, mycotoxin. Sol. MeOH, EtOAc, DMSO. λ_{max} 226 (ε 32400); 260

(ϵ 6100); 290 (ϵ 1600) (EtOH) (Derep).
 λ_{\max} 225 (MeOH).

16-Oxo: 16-Oxoparaherquamide A. VM 55597. Antibiotic VM 55597

[120312-86-9]

$C_{28}H_{33}N_3O_6$ 507.585

Prod. by *Penicillium* sp. IMI332995.

Antiparasitic agent, mycotoxin. λ_{\max} 225 (MeOH).

16-Oxo, 14-deoxy: Paraherquamide H

[887478-34-4]

$C_{28}H_{33}N_3O_5$ 491.586

Prod. by *Penicillium cluniae* (CECT 2888). Amorph. solid. $[\alpha]_D^{20}$ -30 (c, 0.03 in MeOH). λ_{\max} 222 (log ϵ 1) (MeOH).

16-Oxo, 14-deoxy, 14,15-didehydro:

Paraherquamide I

[887478-35-5]

$C_{28}H_{31}N_3O_5$ 489.57

Prod. by *Penicillium cluniae* (CECT 2888). Amorph. solid. $[\alpha]_D^{20}$ -25.1 (c, 0.04 in MeOH). λ_{\max} 225 (log ϵ 1.01) (MeOH).

14-Demethyl, 14-deoxy: Paraherquamide B. UK 111866. Antibiotic UK 111866

[125600-51-3]

$C_{27}H_{33}N_3O_4$ 463.575

Alkaloid from *Penicillium charlesii* and *Penicillium roqueforti*. Nematocide.

λ_{\max} 226 (ϵ 32400); 260 (ϵ 6100); 290 (ϵ 1600) (MeOH) (Derep). λ_{\max} 225 (MeOH).

2-Deoxo: Derquantel, USAN. 2-Deoxoparaherquamide A. PF 00520904. PNU 141962

[187865-22-1]

$C_{28}H_{37}N_3O_4$ 479.618

Anthelmintic. Characterised by pmr.

Yamazaki, M. *et al.*, *CA*, 1980, **95**, 19321 (propos)

Yamazaki, M. *et al.*, *Tet. Lett.*, 1981, 135-136 (uv, ir, pmr, cryst struct)

Eur. Pat., 1989, 322 937; *CA*, **112**, 117340 (isol)

Blizzard, T.A. *et al.*, *J.O.C.*, 1989, **54**, 2657-2663 (abs config)

Ondeyka, J.G. *et al.*, *J. Antibiot.*, 1990, **43**, 1375-1379 (Paraherquamides B-E, isol)

Liesch, J.M. *et al.*, *J. Antibiot.*, 1990, **43**, 1380-1386 (Paraherquamides B-E, struct, pmr, cmr)

Blanchflower, S.E. *et al.*, *J. Antibiot.*, 1991, **44**, 492; 1993, **46**, 1355-1363 (Paraherquamide E, VM55596, VM55597)

Banks, R.M. *et al.*, *J. Antibiot.*, 1997, **50**, 840-846 (Paraherquamide C, SB 203105, SB 200437)

Lee, B.H. *et al.*, *J.O.C.*, 1997, **62**, 1795-1798 (synth)

Pat. Coop. Treaty (WIPO), 1997, ((Upjohn)97 039 88; *CA*, **126**, 211972 (2-deoxo, synth, pharmacol)

Stocking, E.M. *et al.*, *J.A.C.S.*, 2000, **122**, 9089-9098 (biosynth)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (rev, synth, biosynth)

Lee, B.H. *et al.*, *Curr. Top. Med. Chem.*, 2002, **2**, 779-793 (2-deoxo, rev)

Mauragis, M.A. *et al.*, *Org. Process Res. Dev.*, 2002, **6**, 192-196 (2-deoxo, synth)

Williams, R.M. *et al.*, *J.A.C.S.*, 2003, **125**, 12172-12178 (synth)

Domingo, L.R. *et al.*, *J.O.C.*, 2003, **68**, 2895-2902 (biosynth)


Johnson, S.S. *et al.*, *J. Vet. Pharmacol. Ther.*, 2004, **27**, 169-181 (2-deoxo, pharmacol)

López-Gresa, M.P. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 2921-2925 (Paraherquamides H,I)

Paraherquamide G

VM 54158. Antibiotic VM 54158

[125600-55-7]



$C_{28}H_{35}N_3O_4$ 477.602

Alkaloid from *Penicillium charlesii*. Possesses nematocidal props. λ_{\max} 245 (ϵ 28800) (MeOH) (Derep).

14-Deoxy: Paraherquamide F. VM 55594.

Antibiotic VM 55594

[125600-54-6]

$C_{28}H_{35}N_3O_3$ 461.603

From *Penicillium charlesii*. Possesses nematocidal props. λ_{\max} 245 (ϵ 28800) (MeOH) (Derep).


Ondeyka, J.G. *et al.*, *J. Antibiot.*, 1990, **43**, 1375; 1380 (isol, pmr, cmr)

Blanchflower, S.E. *et al.*, *J. Antibiot.*, 1991, **44**, 492 (isol, props)

Williams, R.M. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 711-740 (rev, synth, biosynth)

Paratunamide D

[914778-26-0]



$C_{26}H_{32}N_2O_{11}$ 548.546

Alkaloid from the bark of *Cinnamodendron axillare*. Amorph. solid. $[\alpha]_D^{24}$ -24.9 (c, 0.43 in MeOH). λ_{\max} 205 (ϵ 30700); 232 (ϵ 13600) (MeOH).

3-Me ether: Paratunamide A

[914778-23-7]

$C_{27}H_{34}N_2O_{11}$ 562.572

Alkaloid from the bark of *Cinnamodendron axillare*. Amorph. solid. $[\alpha]_D^{24}$ -98.5 (c, 0.15 in MeOH). λ_{\max} 209 (ϵ 23600); 239 (ϵ 14700) (MeOH).

3-Ketone: Paratunamide B

[914778-24-8]

$C_{26}H_{30}N_2O_{11}$ 546.53

Alkaloid from the bark of *Cinnamodendron axillare*. Amorph. solid. $[\alpha]_D^{24}$ -71.4 (c, 0.73 in MeOH). λ_{\max} 212 (ϵ 22800); 249 (ϵ 12200) (MeOH).

7-Epimer, 3-ketone: Paratunamide C

[914778-25-9]

$C_{26}H_{30}N_2O_{11}$ 546.53

Alkaloid from the bark of *Cinnamodendron axillare*. Amorph. solid. $[\alpha]_D^{24}$ -63.8 (c, 0.77 in MeOH). λ_{\max} 208 (ϵ 37600); 246 (ϵ 14700) (MeOH).

Kagata, T. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1517-1521 (isol, cd, pmr, cmr)


P-91

Paravallarine

20-Hydroxy-3-(methylamino)pregn-5-en-

18,20-olide, 9CI

[510-31-6]



Absolute Configuration

$C_{22}H_{33}NO_2$ 343.508

Alkaloid from *Paravallis microphylla* (Apocynaceae). Mp 181°. $[\alpha]_D^{20}$ -52 (c, 0.32 in $CHCl_3$).

Hydrochloride: Mp 295-298° dec. $[\alpha]_D^{20}$ -25.2 (c, 0.77 in MeOH).

N-Ac: Mp 258-260°.

N-Me: N-Methylparavallarine

[3837-09-0]

$C_{23}H_{35}NO_2$ 357.535

Alkaloid from *Paravallis microphylla* (Apocynaceae). Cryst. (EtOH). Mp 140°. $[\alpha]_D^{20}$ -37.7 (c, 0.398 in $CHCl_3$).

3-Epimer, N-de-Me: 3-Aminopregn-5-en-

18,20-olide. Gitingensine

[13084-70-3]

$C_{21}H_{31}NO_2$ 329.481

Alkaloid from the leaves of *Kibatalia gitingensis* (Apocynaceae). Shows antispasmodic activity. Prisms (Et₂O). Mp 161°. $[\alpha]_D^{31}$ -65 ($CHCl_3$).

7 α -Hydroxy: 7 α -Hydroxyparavallarine

[16137-30-7]

$C_{22}H_{33}NO_3$ 359.508

Alkaloid from *Paravallis microphylla* (Apocynaceae). Mp 213°. $[\alpha]_D^{20}$ -113 ($CHCl_3$).

7 β -Hydroxy: 7 β -Hydroxyparavallarine

[24411-84-5]

$C_{22}H_{33}NO_3$ 359.508

Alkaloid from *Paravallis microphylla* (Apocynaceae). Mp 210°. $[\alpha]_D^{20}$ -13.5 ($CHCl_3$).

11 α -Hydroxy: 11 α -Hydroxyparavallarine

[16137-32-9]

$C_{22}H_{33}NO_3$ 359.508

Alkaloid from *Paravallis microphylla* (Apocynaceae). Mp 235°. $[\alpha]_D^{20}$ -55 ($CHCl_3$).

16 α -Hydroxy: Paravallarinine

[5716-99-4]

$C_{22}H_{33}NO_3$ 359.508

Alkaloid from *Paravallis microphylla* (Apocynaceae). Mp 231°. $[\alpha]_D^{20}$ -48 ($CHCl_3$).

16 α -Hydroxy, N-Me: N-Methylparavallarinine

[25099-43-8]

$C_{23}H_{35}NO_3$ 373.534

Alkaloid from *Paravallis microphylla* (Apocynaceae). Cryst. (EtOH). Mp 200°. $[\alpha]_D^{20}$ -49 (c, 0.29 in $CHCl_3$).

3-Epimer, N-Me: Kibataline

[1244-02-6]

$C_{23}H_{35}NO_2$ 357.535

Alkaloid from *Kibatalia gitingensis* (Apocynaceae). Cryst. (hexane). Mp 171°. $[\alpha]_D^{20}$ -42 (c, 1.4 in $CHCl_3$).

20-Epimer, N-Me: 20-Epi-N-methylpara-

vallarine. 20-*epi*-Kibataline

[96443-60-6]

C₂₃H₃₅NO₂ 357.535

Alkaloid from *Kibatalia gitingensis*. Major alkaloid from leaves of *Paravallaris macrophylla* (Apocynaceae). Mp 191° Mp 220-224°. [α]_D -30 (c, 0.5 in CHCl₃).

Le Men, J. et al., *Bull. Soc. Chim. Fr.*, 1960, 860-864 (*isol, ir, struct*)

Le Men, J. et al., *Bull. Soc. Chim. Fr.*, 1963, 597-603; 1306-1310 (*Paravallaridine, N-Methylparavallaridine*)

Cavé, A. et al., *Bull. Soc. Chim. Fr.*, 1964, 2415-2418 (*Kibataline*)

Cavé, A. et al., *Bull. Soc. Chim. Fr.*, 1965, 2502-2505 (*20-Epi-N-methylparavallarine*)

Aguilar-Santos, G. et al., *Philipp. J. Sci.*, 1965, **94**, 217 (*Gitingensine*)

Aguilar-Santos, G. et al., *J.O.C.*, 1967, **32**, 2642-2644 (*Gitingensine, struct*)

Husson, H.P. et al., *Bull. Soc. Chim. Fr.*, 1969, 3162-3166 (*7-hydroxy, 11-hydroxy*)


Ngoc, P.H. et al., *Planta Med.*, 1984, **50**, 269-270 (*cryst struct, 20-epi-Kibataline*)

Kutschabsky, L. et al., *Cryst. Res. Technol.*, 1985, **20**, 365-369; *CA*, **103**, 71585a (*cryst struct, 20-epi-Kibataline*)

Parazoanthoxanthin A

P-94

4-Methyl-1*H*-cyclohepta[1,2-d:4,5-d']diimidazole-2,6-diamine, 9*CI*. 2,6-Diamino-4-methyl-1*H*-cyclohepta[1,2-d:4,5-d']diimidazole
[53823-11-3]

C₁₀H₁₀N₆ 214.229

Complex tautomerism possible with 1*H* and 3*H* tautomers. The parent compd. is symmetrical by tautomerism between the two imidazole rings but substitution can destroy this symmetry. Fluorescent pigment from *Parazoanthus axinellae adriaticus*, *Zoanthus sociatus* and *Zoanthus* cf. *pacificus*. Yellow cryst. (EtOH). Mp 310°.

3-Me: Parazoanthoxanthin B

[53571-91-8]

C₁₁H₁₂N₆ 228.256

Isol. from *Parazoanthus axinellae adriaticus* and *Zoanthus* cf. *pacificus*.

3,N⁶-Di-Me: Parazoanthoxanthin G

[71827-19-5]

C₁₂H₁₄N₆ 242.283

Isol. from *Parazoanthus axinellae adriaticus* and *Zoanthus sociatus*.

N⁶,N⁶-Di-Me: Parazoanthoxanthin D

[53941-25-6]

C₁₂H₁₄N₆ 242.283

Constit. of *Parazoanthus axinellae adriaticus*, *Zoanthus sociatus*, *Zoanthus* cf. *pacificus*, *Palythoa mammilosa* and *Palythoa tuberculosa*. Yellow needles (H₂O). Mp 303-304° dec. λ_{\max} 255 (€ 13200); 300 (€ 39800); 394 (€ 15100) (MeOH/HCl) (Derep). λ_{\max} 296 (sh) (€ 28800); 306 (€ 36300); 415 (€ 17800) (MeOH) (Derep).

N,N'-Di-Me: Parazoanthoxanthin CC₁₂H₁₄N₆ 242.283

Isol. from *Parazoanthus axinellae*. Exact struct. not determined. λ_{\max} 302; 412 (MeOH). λ_{\max} 297; 392 (MeOH/acid).

1,N⁶,N⁶-Tri-Me: Palyzoanthoxanthin A

[55084-57-6]

C₁₃H₁₆N₆ 256.31

Pigment from *Palythoa mammilosa*, *Palythoa tuberculosa* and *Zoanthus pacificus*. Cryst. (MeOH). Mp 310°.

3,N⁶,N⁶-Tri-Me: Zoanthoxanthin

[40451-47-6]

C₁₃H₁₆N₆ 256.31

Isol. from *Parazoanthus axinellae*. Yellow needles (MeOH). Mp 275-276°. Highly fluorescent. λ_{\max} 259 (€ 7410); 293 (€ 17800); 392 (€ 8710) (1*M* HCl) (Derep). λ_{\max} 293 (€ 33100); 427 (€ 22400) (MeOH) (Derep).

► GU3100500

N²,N²,N⁶-Tri-Me: Epizoanthoxanthin A

[55084-60-1]

C₁₃H₁₆N₆ 256.31

Pigment from *Parazoanthus axinellae* and *Zoanthus pacificus*. Orange-yellow needles (H₂O). Mp 191-192°.

1,3,N⁶,N⁶-Tetra-Me: Parazoanthoxanthin F

[55084-58-7]

C₁₄H₁₈N₆ 270.336

Isol. from *Parazoanthus axinellae adriaticus* and *Palythoa mammilosa*. Cryst. (EtOH). Mp 310° (darkens above 220°). Conts. =NH at C(2).

1,N²,N²,N⁶-Tetra-Me: Epizoanthoxanthin B

[55827-10-6]

C₁₄H₁₈N₆ 270.336

Pigment from *Parazoanthus axinellae* and *Palythoa mammilosa*. Amorph. yellow powder.

3,N²,N⁶,N⁶-Tetra-Me: Parazoanthoxanthin E

[55084-59-8]

C₁₄H₁₈N₆ 270.336

Isol. from *Parazoanthus axinellae*. Yellow cryst. (EtOH). Mp > 310° (darkens >220°). CAS gives incorrect struct.

N²,N²,N⁶,N⁶-Tetra-Me: Palyzoanthoxanthin B

[55084-61-2]

C₁₄H₁₈N₆ 270.336

Pigment from *Palythoa tuberculosa*. Cryst. (EtOH). Mp 310° (darkens > 220°).

3,N²,N²,N⁶,N⁶-Penta-Me: Palyzoanthoxanthin C

[71827-20-8]

C₁₅H₂₀N₆ 284.363

Pigment from *Palythoa mammilosa*. Cryst. (EtOH). Mp 180-181°.

Cariello, L. et al., *Experientia*, 1974, **30**, 849 (*Zoanthoxanthin, cryst struct, Parazoanthoxanthins A-D*)

Cariello, L. et al., *Tetrahedron*, 1974, **30**, 3281-3287; 3611-3614; 4191-4196 (*Parazoanthoxanthins E-F, Palyzoanthoxanthins, Epizoanthoxanthins*)

Braun, M. et al., *J.A.C.S.*, 1976, **98**, 3049 (*Parazoanthoxanthin A*)

Cariello, L. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1979, **63**, 77 (*occur*)

Jiménez, C. et al., *J. Nat. Prod.*, 1993, **56**, 9-14 (*Zoanthoxanthin, cmr*)

Paricine

P-95

[1400-76-6]

C₁₆H₁₈N₂O 254.331


Struct. unknown. Alkaloid from *Buena hexandra* (preferred genus name *Gonzalagunia*), *Cinchona lutea* and *Cinchona succirubra* (Rubiaceae). Amorph. yellow powder. Sol. H₂O. Mp 116°.

Hesse, O. et al., *Annalen*, 1873, **166**, 217 (*bibl*)

Parsonsianidine

P-96

[135601-81-9]

C₂₂H₃₃NO₉ 455.504

Alkaloid from the leaves of *Parsonsia laevigata* (Apocynaceae). Cryst. Mp 230-245° dec. [α]_D²⁵ +17.3 (c, 0.4 in MeOH).

14-Deoxy: 14-Deoxyparsonsianidine

[136997-62-1]

C₂₂H₃₃NO₈ 439.505

Alkaloid from leaves of *Parsonsia laevigata* (Apocynaceae). Fine needles (MeOH). Mp 185-192° dec. [α]_D²⁹ +25.3 (c, 0.50 in MeOH).

Abe, F. et al., *Chem. Pharm. Bull.*, 1991, **39**, 1576 (*14-Deoxyparsonsianidine*)


Abe, F. et al., *Phytochemistry*, 1991, **30**, 1737 (*isol, pmr, cmr, struct*)

Parsonsianine

P-97

16-Hydroxy-22-norparsonsine, 9CI

[131683-36-8]

C₂₁H₃₁NO₉ 441.477

CAS numbering shown. Alkaloid from the leaves of *Parsonsia laevigata* (Apocynaceae). Prisms (MeOH). Mp 235-245° dec. [α]_D²⁴ +37.9 (c, 0.7 in MeOH).

16-Deoxy: 22-Norparsonsine, 9CI. 16-Deoxyparsonsianine. 14-Deoxyparsonsianine. Ideamine B

[136997-61-0]

C₂₁H₃₁NO₈ 425.478


Alkaloid from leaves of *Parsonsia laevigata* (Apocynaceae). Fine prisms (MeOH). Mp 159-164° dec. [α]_D³⁰ +33.5 (c, 0.80 in MeOH). [α]_D³¹ +21 (c, 1 in EtOH).

16-Deoxy, N-oxide: **Ideamine B N-oxide**
[137760-63-5]
C₂₁H₃₁NO₉ 441.477
Isol. from adult bodies of the Apocynaceae-feeding danaine butterfly *Idea leuconoe*. [α]_D²⁰ +24 (c, 3.7 in MeOH).

Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 2127-2129 (*isol, pmr, cmr, ms, cryst struct*)
Nishida, R. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 1787-1792 (*Ideamine B*)
Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1576-1577 (*14-Deoxyparsonianine*)

Parsonsine, 9CI**P-98**

[72213-98-0]



Absolute configuration

C₂₂H₃₃NO₈ 439.505


Retronecine cyclic diester of unusual type. Alkaloid from *Parsonsia heterophylla* and *Parsonsia spiralis* (Apocynaceae). Plates (C₆H₆). Mp 158° Mp 196° (dimorph.). See also 17-Methylparsonianidine, M-502.

N-Oxide: **Parsonsine N-oxide**[137782-07-1]
C₂₂H₃₃NO₉ 455.504

Isol. from adult bodies of the Apocynaceae-feeding danaine butterfly *Idea leuconoe*. [α]_D²⁰ -21 (c, 0.75 in MeOH).

16ξ-Hydroxy: **Spiraline**[77156-25-3]
C₂₂H₃₃NO₉ 455.504

Alkaloid from *Parsonsia spiralis* (Apocynaceae). Characterised by ms and nmr, stereochem. not proven.

Gainsford, G.J. *et al.*, *Cryst. Struct. Commun.*, 1979, **8**, 597; 1980, **9**, 173 (*cryst struct*)Edgar, J.A. *et al.*, *Tet. Lett.*, 1980, 2657 (*isol, struct, pmr*)Nishida, R. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 1787 (*oxide*)**Parthenolidine****P-99**C₁₅H₂₃NO₃ 265.352N-Ac: **N-Acetylparthenolidine**[119766-96-0]
C₁₇H₂₅NO₄ 307.389

Constit. of *Michelia rajaniana* (Magnoliaceae). Oil. [α]_D²⁴ +23 (c, 0.67 in CHCl₃).

8α-Hydroxy, N-Ac: **N-Acetyl-8-hydroxy-parthenolidine**[119766-97-1]
C₁₇H₂₅NO₅ 323.388


Isol. from bark of *Michelia rajaniana* (Magnoliaceae). Mp 188-190°. [α]_D²⁰ +51 (c, 0.75 in CHCl₃).

Ruangrungsri, N. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1220 (*isol, pmr, cmr*)**Parvifagarine****P-100**C₂₃H₂₁NO₄ 375.423


Struct. unknown. Alkaloid from the root bark of *Fagara parvifolia*. Needles. Mp 165-166° Mp 185-186° (as hydrochloride).

Paris, R. *et al.*, *Ann. Pharm. Fr.*, 1948, **6**, 409-421; *CA*, **43**, 5546e (*isol*)**Parvineostemonine****P-101**

[539883-61-9]

C₁₇H₂₅NO₂ 275.39


Alkaloid from the stems and leaves of *Stemona parviflora*. Amorph. pale yellow solid.

Ke, C.Q. *et al.*, *Chin. Chem. Lett.*, 2003, **14**, 173-175 (*isol, pmr, cmr*)**Parvinine****P-102**C₁₉H₂₁NO₄ 327.379**(R)-form** [920757-19-3]

Alkaloid from *Xylopi parviflora*. Amorph. powder. [α]_D²² +20 (c, 0.28 in CHCl₃). λ_{max} 278 (log ε 3.36); 284 (sh) (CHCl₃).

Nishiyama, Y. *et al.*, *Phytochemistry*, 2006, **67**, 2671-2675 (*isol, pmr, cmr, ms*)**Parvistemoamide****P-103**

[139742-26-0]




Absolute Configuration

C₁₂H₁₉NO₄ 241.286

Alkaloid from roots of *Stemona parviflora*.

Lin, W. *et al.*, *Huaxue Xuebao*, 1991, **49**, 927; *CA*, **116**, 148198e**Parvistemoline****P-104**


[139766-99-7]

C₁₇H₂₅NO₄ 307.389

Alkaloid from roots of *Stemona parviflora*.

Lin, W. *et al.*, *Huaxue Xuebao*, 1991, **49**, 927; *CA*, **116**, 148198e**Parvistemonine****P-105**


Tetrahydro-3,5-dimethyl-6-[octahydro-3-(tetrahydro-4-methyl-5-oxo-2-furanyl)-1H-pyrrolo[1,2-a]azepin-9-yl]furo[3,2-b]furan-2(3H)-one, 9CI
[131653-95-7]

C₂₂H₃₃NO₅ 391.506

Alkaloid from the roots of *Stemona parviflora* (Stemonaceae).

Lin, W. *et al.*, *Huaxue Xuebao*, 1990, **48**, 811-814; *CA*, **114**, 78639g (*isol, pmr, struct*)Kaltenegger, E. *et al.*, *Phytochemistry*, 2003, **63**, 803-816 (*isol, pmr, cmr*)**Paspaclavine****P-106**

[52052-65-0]




Absolute Configuration


C₁₈H₂₂N₂O 282.385

Metab. of *Claviceps paspali*. Cryst. (Me₂CO). Mp 204-206° dec. [α]_D²⁰ +88 (c, 0.2 in Py). Cleaved in acidic medium to Paliclavine, P-20 and acetaldehyde.

Tscherter, H. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 113 (*isol, uv, ir, pmr, ms, struct*)Kozikowski, A.P. *et al.*, *Tetrahedron*, 1984, **40**, 2345 (*synth*)

Paspalic acid**P-107**6-Methyl-8-ergolene-8-carboxylic acid
[5516-88-1]Absolute
ConfigurationC₁₆H₁₆N₂O₂ 268.315Metab. from a strain of *Claviceps paspali* and *Claviceps purpurea*. Mp 245-247° dec. [α]_D -208 (c, 0.4 in 0.1M NaOH).*Hydrochloride*: Mp 257-258° dec. [α]_D -176 (c, 0.4 in 0.1M HCl).**10-Hydroxy, amide: 10-Hydroxy-trans-paspalic acid amide**C₁₆H₁₇N₃O₂ 283.329Alkaloid from *Claviceps paspali*.**10-Hydroxy, 10-epimer, amide: 10-Hydroxy-cis-paspalic acid amide**C₁₆H₁₇N₃O₂ 283.329Alkaloid from *Claviceps paspali*.Kobel, H. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1052 (*isol, uv, ir, pmr, struct*)Castagnoli, N. *et al.*, *Nature (London)*, 1966, **211**, 859 (*occur*)Flieger, M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 810 (*10-Hydroxypaspalic acid amide*)**Paspaline****P-108**

[11024-56-9]

C₂₈H₃₉NO₂ 421.622Isol. from *Claviceps paspali*, *Albophoma yamaushiensis* Fo2546-Bp4406, *Penicillium paxilli* and the mangrove-derived *Penicillium* sp. HKI0459. Tremorogenic mycotoxin. Inhibitor of acyl-CoA:cholesterol acyltransferase (ACAT). Cryst. (MeOH or by subl.). Mp 264° (254°). [α]_D -23 (c, 0.36 in CHCl₃). [α]_D -38.5 (c, 0.47 in C₆H₆). λ_{\max} 229 (ε 25120); 282 (ε 7940); 291 (ε 6300) (EtOH) (Berdy).**1'α-Hydroxy: Terpendole E**

[167427-23-8]

C₂₈H₃₉NO₃ 437.621Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Cryst. Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 174-176°. [α]_D²⁵ -36.4 (c, 1 in MeOH). λ_{\max} 228 (ε 38600); 280 (ε 8700) (MeOH) (Berdy).**1'α,19-Dihydroxy: Terpendole F**

[167427-24-9]

C₂₈H₃₉NO₄ 453.62Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆; poorly sol. H₂O. [α]_D²⁸ -35.8 (c, 1 in MeOH). λ_{\max} 228 (ε 48000); 280 (ε 10900) (MeOH) (Berdy).**19-Oxo: Paspaline B**


[173268-84-3]

C₂₈H₃₇NO₃ 435.605Isol. from *Penicillium paxilli*. Solid.**19-Oxo, 1'α-hydroxy: Terpendole G**

[167427-25-0]

C₂₈H₃₇NO₄ 451.605Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 248-250° dec. [α]_D²⁸ -28 (c, 1 in MeOH). λ_{\max} 228 (ε 49400); 280 (ε 11800) (MeOH) (Berdy).Fehr, T. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 1907 (*isol, uv, ir, pmr*)Springer, J.P. *et al.*, *Tet. Lett.*, 1980, 231 (*cryst struct*)Smith, A.B. *et al.*, *J.A.C.S.*, 1989, **111**, 5761 (*synth*)Mewshaw, R.E. *et al.*, *J.O.C.*, 1989, **54**, 3449 (*synth, ir, pmr*)Tomoda, H. *et al.*, *J. Antibiot.*, 1995, **48**, 793 (*Terpendoles*)Munday-Finch, S.C. *et al.*, *Phytochemistry*, 1996, **41**, 327-332 (*Paspaline B*)Xu, M. *et al.*, *Tetrahedron*, 2007, **63**, 435-444 (*isol, biosynth*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 390**Paspalinine****P-109**

[63722-91-8]



R = R' = H

C₂₇H₃₁NO₄ 433.546Isol. from *Claviceps paspali*, *Penicillium digitatum*, *Eupenicillium shearii* and *Aspergillus flavus*. Tremorogenic mycotoxin, shows insecticidal props. Sol. MeOH, CHCl₃. λ_{\max} 232 (ε 25000); 250 (ε 14800); 274 (ε 8000) (MeOH) (Berdy).**Deoxy: Paspalicine**

[11024-55-8]

C₂₇H₃₁NO₃ 417.547Isol. from *Claviceps paspali*. Tremorogenic mycotoxin. Large needles (Me₂CO). Mp ca. 240° (dec.). [α]_D +173 (c, 0.5 in CHCl₃). λ_{\max} 231 (ε 47800); 275 (ε 12600) (EtOH) (Berdy).

▶ Toxic.

14α-Hydroxy: 14-Hydroxypaspalinine

[151341-77-4]

C₂₇H₃₁NO₅ 449.546Isol. from the sclerotia of *Aspergillus nomius*. Tremorogenic mycotoxin. Exhibits insecticidal activity. [α]_D²⁹ +127 (c, 0.11 in MeOH).**14α-Hydroxy, 14-O-(2S-dimethylamino-3-methylbutanoyl): 14-(N,N-Dimethyl-L-valyloxy)paspalinine**

[151341-78-5]

C₃₄H₄₄N₂O₆ 576.731Isol. from the sclerotia of *Aspergillus nomius*. Tremorogenic mycotoxin, shows insecticidal props. [α]_D²⁶ +102 (c, 0.44 in MeOH). λ_{\max} 228 (ε 15620); 279 (ε 3110) (MeOH) (Berdy).Fehr, T. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 1907 (*isol, uv, ir, pmr*)Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 826; 1197; 1981, **29**, 293 (*isol, deriv*)Springer, J.P. *et al.*, *Tet. Lett.*, 1980, 231 (*cryst struct*)Gallagher, R.T. *et al.*, *Tet. Lett.*, 1980, 235 (*cryst struct, uv, ir, pmr, cmr*)Nozawa, K. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 626 (*isol*)Smith, A.B. *et al.*, *J.A.C.S.*, 1990, **112**, 8197 (*synth, Paspalinine, Paspalinine*)Staub, G.M. *et al.*, *Tet. Lett.*, 1993, **34**, 2569 (*14-Hydroxypaspalinine, 14-(N,N-Dimethyl-L-valyloxy)paspalinine*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 394; 398**Paspalitrem A****P-110****(3-Methyl-2-butenyl)paspalinine**

[63722-90-7]

As Paspalinine, P-109 with R = (H₃C)₂CH=CH'CH₂-, R' = HC₃₂H₃₉NO₄ 501.664Prod. by *Claviceps paspali* and the mangrove-derived *Penicillium* sp.HKI0459. Tremorogenic mycotoxin. λ_{\max} 227; 248 (MeOH) (Berdy). λ_{\max} 305; 335 (EtOH) (Berdy).▶ LD₅₀ (mus, ipr) < 14 mg/kg.**Δ^{1'}-Isomer, 3'-hydroxy: Paspalitrem B.****(3-Methyl-3-hydroxy-1-butenyl)paspalinine**

[63764-58-9]

C₃₂H₃₉NO₅ 517.664Isol. from *Claviceps paspali*. Tremorogenic mycotoxin.▶ LD₅₀ (mus, ipr) < 14 mg/kg.Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 1197-1201 (*Paspalitrem A-B*)Betina, V. *et al.*, *Mycotoxins: Chemical, Biological and Environmental Aspects*, Elsevier, 1989, 383 (*tox*)Xu, M. *et al.*, *Tetrahedron*, 2007, **63**, 435-444 (*Paspalitrem A, isol, biosynth*)Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 402**Paspalitrem C****P-111****4-(3-Methyl-2-butenyl)paspalinine**


[90866-61-8]

As Paspalinine, P-109 with

R = H, R' = (H₃C)₂CH=CHCH₂-C₃₂H₃₉NO₄ 501.664From *Claviceps paspali*. Tremorogenic mycotoxin. λ_{\max} 234 (ε 35500); 278 (ε 10500) (MeOH) (Berdy).Dorner, J.W. *et al.*, *J. Agric. Food Chem.*, 1984, **32**, 1069-1071 (*isol, pmr, cmr*)

Patchouliopyridine**P-112**

6,7,8,9-Tetrahydro-2,5,10,10-tetramethyl-5,8-methano-5H-cyclohepta[b]pyridine, 9CI
[6517-97-1]



$C_{15}H_{21}N$ 215.338

Alkaloid from the essential oil of *Pogostemon patchouly* (Lamiaceae). Waxy needles. Mp 24–26.5°. $[\alpha]_D$ -31.3 (c, 5.06 in EtOH).


Büchi, G. *et al.*, *J.A.C.S.*, 1966, **88**, 3109 (*isol, uv, ir, pmr*)

Cren, M.-C. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 3020 (*synth, ms*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 481 (*synth, pmr, cmr*)

Pateamine A**P-113**

[139220-18-1]



$C_{31}H_{45}N_3O_4S$ 555.78

Rare dilactone functionality. *Isol.* from the New Zealand marine sponge *Mycale* sp. Potent cytotoxin and immunosuppressant. $[\alpha]_D$ -253 (MeOH). λ_{max} 274 (ϵ 47000); 285 (ϵ 42000) (MeOH) (Berdy).

Northcote, P.T. *et al.*, *Tet. Lett.*, 1991, **32**, 6411-6414 (*isol, uv, cd, ir, pmr, cmr, ms, struct*)

Stirling, D.J. *et al.*, *Ph.D. Thesis*, 1996, (*activity*)


Remuñán, M.J. *et al.*, *Tet. Lett.*, 2000, **41**, 7367-7371 (*synth*)

Pattenden, G. *et al.*, *Can. J. Chem.*, 2004, **82**, 353-365 (*synth*)

Romo, D. *et al.*, *J.A.C.S.*, 2004, **126**, 10582-10588 (*synth*)

Patellamide A**P-114**

[81120-73-2]



$C_{35}H_{50}N_8O_6S_2$ 742.962

Struct. revised in 1985. Cyclopeptide from the marine tunicate *Lissoclinum patella*. Shows selective metal binding props. Cytotoxic, shows antineoplastic props. MDR inhibitor. Cryst. (C_6H_6). Sol. MeOH, EtOAc; poorly sol. hexane, H_2O . Mp 228–229°. $[\alpha]_D^{24}$ +140.7 (c, 0.27 in $CHCl_3$) (+113.9). λ_{max} 232 (ϵ 21000) (MeOH) (Derep).

Ireland, C.M. *et al.*, *J.O.C.*, 1982, **47**, 1807-1811 (*isol, ir, pmr, cmr*)

Biskupiak, J.E. *et al.*, *J.O.C.*, 1983, **48**, 2302-2304 (*abs config*)

Hamada, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 6501-6504 (*synth*)

In, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1686-1690 (*conformn, cryst struct*)

In, Y. *et al.*, *Acta Cryst. C*, 1994, **50**, 432-434 (*cryst struct*)

Morris, L.A. *et al.*, *Tetrahedron*, 2001, **57**, 3185-3197 (*activity*)


Salomon, C.E. *et al.*, *J. Nat. Prod.*, 2002, **65**, 689-692 (*isol*)

Schmidt, E.W. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2005, **102**, 7315-7320 (*biosynth*)

Garcia-Reynaga, P. *et al.*, *Org. Lett.*, 2008, **10**, 4621-4623 (*synth*)

Patellamide B**P-115**

[81098-23-9]



R = $CH_2CH(CH_3)_2$

$C_{38}H_{48}N_8O_6S_2$ 776.979

Cyclopeptide from the marine tunicate *Lissoclinum patella*. Cytotoxic, shows antineoplastic props. MDR inhibitor. Sol. MeOH, $CHCl_3$; poorly sol. hexane, H_2O . $[\alpha]_D$ +29.4 (c, 0.34 in CH_2Cl_2). λ_{max} 248 (ϵ 8000) (MeOH) (Derep).

Ireland, C.M. *et al.*, *J.O.C.*, 1982, **47**, 1807-1811 (*isol, ir, pmr, cmr*)

Biskupiak, J.E. *et al.*, *J.O.C.*, 1983, **48**, 2302-2304 (*abs config*)

Hamada, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 5155-5158; 5159-5162 (*synth, struct*)

Schmidt, U. *et al.*, *Tet. Lett.*, 1986, **27**, 163-166; 179 (*synth, struct*)

Salomon, C.E. *et al.*, *J. Nat. Prod.*, 2002, **65**, 689-692 (*isol*)

Patellamide C**P-116**

[81120-74-3]

As Patellamide B, P-115 with

R = $-CH(CH_3)_2$

$C_{37}H_{46}N_8O_6S_2$ 762.952

Cyclopeptide from the marine tunicate *Lissoclinum patella*. Shows selective metal binding props. Antineoplastic agent. MDR inhibitor. Sol. MeOH, $CHCl_3$;

poorly sol. H_2O , hexane. $[\alpha]_D$ +19 (c, 0.21 in CH_2Cl_2). λ_{max} 248 (ϵ 8000) (MeOH) (Derep).

Ireland, C.M. *et al.*, *J.O.C.*, 1982, **47**, 1807-1811; 1983, **48**, 2302-2304 (*isol, ir, pmr, cmr, abs config*)

Hamada, Y. *et al.*, *Tet. Lett.*, 1985, **26**, 5155-5158; 5159-5162 (*synth, struct*)

Morris, L.A. *et al.*, *Tetrahedron*, 2001, **57**, 3185-3197 (*activity*)

Salomon, C.E. *et al.*, *J. Nat. Prod.*, 2002, **65**, 689-692 (*isol*)

Schmidt, E.W. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2005, **102**, 7315-7320 (*biosynth*)

Patellamide D**P-117**

[120853-15-8]

As Patellamide B, P-115 with

R = $-CH(CH_3)CH_2CH_3$

$C_{38}H_{48}N_8O_6S_2$ 776.979


Cyclopeptide from the ascidian *Lissoclinum patella*. Cytotoxic. Mp 144–145°. $[\alpha]_D$ +32 (c, 0.37 in $CHCl_3$).

Degnan, B.M. *et al.*, *J. Med. Chem.*, 1989, **32**, 1349-1354 (*isol, pmr, cmr, struct*)

Schmitz, F.J. *et al.*, *J.O.C.*, 1989, **54**, 3463-3472 (*isol, ir, pmr, cmr, cryst struct*)

Patellamide E**P-118**

[140430-46-2]




$C_{39}H_{50}N_8O_6S_2$ 791.006

Cyclic peptide antibiotic. *Isol.* from the ascidian *Lissoclinum patella*. Weakly cytotoxic. Amorph. $[\alpha]_D^{25}$ +48.6 (c, 0.58 in $CHCl_3$). λ_{max} 235 (ϵ 12300) (MeOH) (Berdy).

McDonald, L.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 376-379 (*isol, cmr, struct*)

Patellamide F**P-119**


[164803-18-3]



C₃₇H₄₆N₈O₆S₂ 762.952
Cyclopeptide from the marine tunicate *Lissoclinum patella*. Cytotoxic. Amorph. solid. Sol. MeOH. [α]_D +40 (c, 0.1 in MeOH). λ_{max} 235 (ε 13700) (MeOH) (Berdy).

Rashid, M.A. *et al.*, *J. Nat. Prod.*, 1995, **58**, 594-597 (*isol, ir, pmr, cmr, struct*)


Patellamide G P-120
[218916-91-7]



C₃₈H₅₀N₈O₇S₂ 794.994
Cyclic peptide. *Isol.* from the ascidian *Lissoclinum patella*. Amorph. solid. [α]_D +40.6 (c, 0.35 in MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1547-1551 (*isol, ir, pmr, cmr, ms*)


Patellin 1 P-121
[181758-78-1]



C₃₇H₆₀N₆O₇S 732.983
Cyclic hexapeptide. *Isol.* from the ascidian *Lissoclinum patella*. Glass. λ_{max} 220 (ε 3800) (MeOH).

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1996, **49**, 659-667 (*isol, uv, ir, pmr, cmr, ms*)

Patellin 3 P-122
[181758-79-2]



C₄₈H₇₈N₈O₉S 943.258

R¹ = CH₃, R² = R³ = -CH₂CH(CH₃)₂

Cyclic octapeptide. *Isol.* from the ascidian *Lissoclinum patella*. Oil. [α]_D -63 (c, 0.09 in CHCl₃). λ_{max} 242 (ε 10100) (EtOH).

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1996, **49**, 659-667 (*isol, uv, ir, ms, struct*)

Patellin 4 P-123

[181758-80-5]
As Patellin 3, P-122 with
R¹ = CH₃, R² = -CH(CH₃)₂, R³ = -CH₂CH(CH₃)₂

C₄₇H₇₆N₈O₉S 929.232

Cyclic octapeptide, stereochem. not assigned. *Isol.* from the ascidian *Lissoclinum patella*.

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1996, **49**, 659-667 (*isol*)

Patellin 5 P-124

[181758-81-6]
As Patellin 3, P-122 with
R¹ = H, R² = -CH₂Ph, R³ = -CH(CH₃)₂

C₄₉H₇₂N₈O₉S 949.222

Cyclic octapeptide. *Isol.* from the ascidian *Lissoclinum patella*. Oil. [α]_D -36 (c, 0.05 in CHCl₃). λ_{max} 245 (ε 12500) (EtOH).

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1996, **49**, 659-667 (*isol, ur, ir, pmr, cmr, ms*)

Patellin 6 P-125


[181758-82-7]
As Patellin 3, P-122 with
R¹ = CH₃, R² = -CH(CH₃)₂, R³ = -CH₂Ph
C₅₀H₇₄N₈O₉S 963.249

Cyclic octapeptide. *Isol.* from the ascidian *Lissoclinum patella*. Oil. [α]_D -50 (c, 0.04 in CHCl₃). λ_{max} 248 (ε 12400) (EtOH).

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1996, **49**, 659-667 (*isol, uv, ir, pmr, cmr, ms*)

Patungensine P-126

[951212-48-9]



Absolute Configuration

C₂₂H₃₃NO₅ 391.506

Alkaloid from *Lysimachia patungensis*. Needles (MeOH). Mp 111.5-112.5°. [α]_D +12.8 (c, 0.2 in MeOH). λ_{max} 221 (log ε 4.4); 257 (log ε 4.27); 290 (log ε 3.85) (EtOH).

Huang, X. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 170-172 (*isol, pmr, cmr, ms*)


Paucaline P-127

C₁₈H₂₇NO₈ 385.413

Pyrrrolizidine alkaloid. Struct. unknown. Alkaloid from *Senecio pauciligulatus* (Asteraceae). Fine needles (EtOH). V. sol. H₂O. Mp 184° dec.

Pretorius, T.P. *et al.*, *CA*, 1950, **44**, 3217f

Paucidactine A P-128
[178253-98-0]



Absolute Configuration

C₂₃H₂₂N₂O₉ 470.435

Alkaloid from *Kopsia pauciflora*. Light yellow cryst. Mp 279-280°. [α]_D +53.8 (c, 0.156 in CHCl₃).

21-Deoxy: Paucidactine B


[178253-99-1]

C₂₃H₂₂N₂O₈ 454.435

Minor alkaloid from *Kopsia pauciflora*. [α]_D +1.2 (c, 0.017 in CHCl₃).

Kam, T.-S. *et al.*, *Tet. Lett.*, 1996, **37**, 3603-3606 (*isol, uv, pmr, cmr, ms, cryst struct*)

Paucidactine C P-129
[949932-23-4]




Absolute Configuration

C₂₄H₂₆N₂O₈ 470.478

Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D +250 (c, 0.01 in CHCl₃). λ_{max} 220 (log ε 4.64); 247 (log ε 3.93); 280 (log ε 3.36) (EtOH).

Lim, K.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*isol, pmr, cmr*)

Pauciflorine A P-130
[181486-81-7]



R¹ R² = -CH₂-

C₂₄H₂₆N₂O₈ 470.478

A 20,21-secoaspidofractine alkaloid. Alkaloid from leaves of *Kopsia pauciflora*. Melanin biosynthesis inhibitor. Amorph. [α]_D -50.7 (c, 2.0 in CHCl₃). λ_{max} 227 (log ε 4.62); 248 (log ε 4.17); 283 (log ε 3.14); 293 (log ε 3.02) (EtOH).

Kam, T.-S. *et al.*, *Tet. Lett.*, 1996, **37**, 5765 (*isol, uv, pmr, cmr, ms, struct*)

Kuehne, M.E. *et al.*, *Org. Lett.*, 1999, **1**, 1751-1753 (*synth*)

Pauciflorine B

P-131

[181486-82-8]
As Pauciflorine A, P-130 with
 $R^1 = R^2 = \text{Me}$

$\text{C}_{25}\text{H}_{30}\text{N}_2\text{O}_8$ 486.521

A secoaspidofractine alkaloid. Alkaloid from leaves of *Kopsia pauciflora*. Melanin biosynthesis inhibitor. $[\alpha]_{\text{D}}^{25}$ -25 (c, 0.4 in CHCl_3). λ_{max} 224 (log ϵ 4.49); 253 (log ϵ 3.83); 282 (log ϵ 3.04); 290 (log ϵ 3.01) (EtOH).

Kam, T.-S. et al., *Tet. Lett.*, 1996, **37**, 5765 (isol, uv, pmr, cmr, ms, struct)


Kuehne, M.E. et al., *Org. Lett.*, 1999, **1**, 1751-1753 (synth)

Magnus, P. et al., *Tetrahedron*, 2002, **58**, 3423-3443 (synth)

Paucivenine

P-132

[69734-97-0]



$\text{C}_{40}\text{H}_{48}\text{N}_4\text{O}_2$ 616.845


Alkaloid from the leaves and stems of *Melodinus balansae* var. *paucivenosus* (Apocynaceae). Amorph. λ_{max} 233 (log ϵ 3.52); 295 (log ϵ 2.98) (no solvent reported).

Mehri, M.H. et al., *Phytochemistry*, 1978, **17**, 1451-1452 (uv, ir, ms, struct)

Pauridianthine

P-133

1-[4-(9H-Pyrido[3,4-b]indol-1-ylcarbo-nyl)-3-pyridinyl]ethanone, 9CI [31482-18-5]



$\text{C}_{19}\text{H}_{13}\text{N}_3\text{O}_2$ 315.331

Alkaloid from the roots of *Pauridiantha callicarpoides* (Rubiaceae). Yellow prisms (MeOH). Mp 271°.

1''- ξ -Alcohol: **Pauridiantholol** [52908-86-8]

$\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_2$ 317.346


Alkaloid from the root and stem bark of *Pauridiantha lyalli* (Rubiaceae).

Pousset, J.-L. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **272**, 665 (uv, ir, ms, pmr, struct)

Pousset, J.-L. et al., *Plant. Med. Phytother.*, 1974, **7**, 51 (*Pauridiantholol*)

Pauridianthine

P-134



$\text{C}_{19}\text{H}_{13}\text{N}_3\text{O}_2$ 315.331

(ξ)-form [31460-98-7]


Alkaloid from the root bark of *Pauridiantha callicarpoides* (Rubiaceae).

Pousset, J.-L. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **272**, 665 (uv, ir, pmr, ms, struct)

Pauridianthoside

P-135

14-Oxolyaloside [63238-69-7]



$\text{C}_{27}\text{H}_{28}\text{N}_2\text{O}_{10}$ 540.526

The name 14-Oxolyaloside is apparently incorrect because the stereochemistries are different, but this has not been confirmed. Alkaloid from the leaves of *Pauridiantha lyalli* (Rubiaceae). Cryst. (Me₂CO). Mp 208-210°. $[\alpha]_{\text{D}}^{20}$ -264 (c, 0.5 in MeOH).

Tetra-O-Ac: Mp 143-145°. $[\alpha]_{\text{D}}$ -175 (CHCl_3).

20-Epimer: **Isopauridianthoside**

[88644-38-6]

$\text{C}_{27}\text{H}_{28}\text{N}_2\text{O}_{10}$ 540.526

Alkaloid from the leaves of *Pauridiantha lyalli* (Rubiaceae). Fine needles (Et₂O/MeOH) (as tetra-O-Ac). Mp 167-169° (as tetra-O-Ac).

Stereoisomer, 17-hydroxy, 16,17-dihydro:

Palinine

[55856-84-3]

$\text{C}_{27}\text{H}_{32}\text{N}_2\text{O}_{10}$ 544.557

Alkaloid from *Palicourea alpina* (Rubiaceae). Mp 166.5-168°. $[\alpha]_{\text{D}}^{28}$ -252.3 (MeOH). No stereochem. determined.

Stuart, K.L. et al., *Tet. Lett.*, 1974, 3853-3856 (*Palinine*)

Lévesque, J. et al., *Fitoterapia*, 1977, **48**, 5-7 (*Pauridianthoside*)

Lévesque, J. et al., *J. Nat. Prod.*, 1983, **46**, 619-625 (*Isopauridianthoside*)

Pavanoline

P-136

[12706-41-1]

$\text{C}_{21}\text{H}_{25}\text{NO}_6$ 387.432

Struct. unknown. Alkaloid from *Papaver*


anomalum (Papaveraceae). Mp 242-244°. $[\alpha]_{\text{D}}^{20}$ -226 (c, 0.16 in CHCl_3).

Pfeifer, S. et al., *Pharmazie*, 1967, **22**, 454; *CA*, **68**, 6130t

Paxdaphnidine A

P-137

[676365-61-0]



Relative Configuration

$\text{C}_{23}\text{H}_{35}\text{NO}_2$ 357.535

Alkaloid from *Daphniphyllum paxianum*. Oil. $[\alpha]_{\text{D}}^{20}$ -17 (c, 0.27 in MeOH).

14,15-Diepimer: **Caldaphnidine B**

[871261-41-5]

$\text{C}_{23}\text{H}_{35}\text{NO}_2$ 357.535

Alkaloid from *Daphniphyllum calycinum*. Oil. $[\alpha]_{\text{D}}^{20}$ -39 (c, 0.6 in MeOH).


Zhan, Z.-J. et al., *J.O.C.*, 2004, **69**, 1726-1729 (*Paxdaphnidine A*)

Zhan, Z.-J. et al., *Tetrahedron*, 2005, **61**, 11038-11045 (*Caldaphnidine B*)

Paxdaphnine A

P-138

[919111-61-8]



Absolute Configuration

$\text{C}_{22}\text{H}_{31}\text{NO}_3$ 357.492

Alkaloid from *Daphniphyllum longercemosum* and *Daphniphyllum paxianum*. Gum. $[\alpha]_{\text{D}}^{20}$ -71 (c, 0.18 in MeOH).


Li, C. et al., *Tet. Lett.*, 2007, **48**, 2737-2740 (isol, pmr, cmr)

Fan, C.Q. et al., *Tetrahedron*, 2007, **63**, 115-119 (isol, pmr, cmr, ms, cryst struct)

Paxdaphnine B

P-139

[919283-74-2]



Absolute Configuration

$\text{C}_{21}\text{H}_{31}\text{NO}_3$ 345.481

Alkaloid from the seeds of *Daphniphyllum paxianum*. Gum. $[\alpha]_{\text{D}}^{20}$ +23 (c, 0.12 in MeOH).

O-Ac: **O-Acetylpaxdaphnine B**

[960604-06-2]

$\text{C}_{23}\text{H}_{33}\text{NO}_4$ 387.518

Alkaloid from the fruit of *Daphniphyllum macropodum*. Oil. $[\alpha]_{\text{D}}^{20}$ +49.7

(c, 0.67 in MeOH).

N-Hydroxy: N-Hydroxypaxidaphnine B
[960604-04]

$C_{21}H_{31}NO_4$ 361.48

Alkaloid from the fruit of *Daphni-
phyllum macropodium*. Amorph. solid.
[α]_D²⁰ +38.6 (c, 0.72 in MeOH).

Lu, Y.B. *et al.*, *Acta Cryst. E*, 2007, **63**, o589-
o591 (*Paxdaphnine B*, *cryst struct*)


Wang, X.-N. *et al.*, *Helv. Chim. Acta*, 2007, **90**,
2156-2162 (*O-Ac, N-hydroxy*)

Fan, C.-Q. *et al.*, *Tetrahedron*, 2007, **63**, 115-
119 (*Paxdaphnine B*)

Paxilline

P-140

[57186-25-1]



$C_{27}H_{33}NO_4$ 435.562

CAS numbering shown. Several
schemes have been used. Metab. from
Penicillium paxilli, *Eupenicillium shearii*,
Acremonium loliae, *Emericella foveolata*,
Emericella desertorum, *Emericella striata*
and a marine-derived *Scopulariopsis*
brevicaulis. Tremorgenic mycotoxin,
muscle contractant. Plates (Me₂CO).
Mp 252°. λ_{max} 231 (ε 41700); 280 (ε
9330); 290 (sh) (ε 8320) (EtOH) (De-
rep).

▶ LD₅₀ (mus, orl) >227 mg/kg; produces
tremors. DJ2830000

l'-Ac: l'-O-Acetylpaxilline

[121998-08-1]

$C_{29}H_{35}NO_5$ 477.599

Isol. from the mycelium of *Emericella*
striata. Mycotoxin. Tremorgenic agent.
Needles (EtOAc). Subl. 262-265. λ_{max}
231 (ε 41700); 280 (ε 9330); 290 (sh) (ε
8320) (EtOH) (Derep).

3β-Alcohol, 3-Ac: PC-M5'

[133613-75-9]

$C_{29}H_{37}NO_5$ 479.615

Metab. of *Penicillium crustosum*. Cryst.
powder. Mp 162-164°.

4b-Deoxy: Dehydroxypaxilline

[112900-05-7]

$C_{27}H_{33}NO_3$ 419.563

Metab. of *Emericella striata*. Leaflets
(MeOH). Mp 232-234°.

**4b-Deoxy, 3β-alcohol: PC-M6. 10-Hy-
droxy-13-deoxypaxilline. 3-Hydroxy-4-
deoxypaxilline**

[133613-76-0]

$C_{27}H_{35}NO_3$ 421.578

Metab. of *Penicillium crustosum*, *Peni-
cillium paxilli* and *Acremonium lolii*.
Tremorgenic mycotoxin. Cryst. pow-
der. Mp 260-263° dec. Numbering
systems vary.

14a-Hydroxy: 14a-Hydroxypaxilline

[157530-30-8]

$C_{27}H_{33}NO_5$ 451.561

Metab. from *Penicillium paxilli*. Also

identified chromatographically in
Acremonium lolii. Apparently not tre-
morgenic.

**14a-Hydroxy, 4b-deoxy: 14a-Hydroxy-4b-
deoxypaxilline**

[157530-29-5]

$C_{27}H_{33}NO_4$ 435.562

Metab. of *Penicillium paxilli* and
Acremonium lolii.

Cole, R.J. *et al.*, *Can. J. Microbiol.*, 1974, **20**,
1159 (*isol, uv, ir, pmr, ms*)

Springer, J.P. *et al.*, *Tet. Lett.*, 1975, 2531;
1980, 231 (*pmr, cd, cryst struct, abs*
config)

Weedon, C.M. *et al.*, *Phytochemistry*, 1987, **26**,
969

Nozawa, K. *et al.*, *J.C.S. Perkin 1*, 1988, 2607
(*Dehydroxypaxilline*)

Nozawa, K. *et al.*, *Chem. Pharm. Bull.*, 1989,
37, 1387 (*l'-O-Acetylpaxilline*)

Betina, V. *et al.*, *Mycotoxins, Chemical,
Biological and Environmental Aspects*,
Elsevier, 1989, 365 (*rev*)

Hosoe, T. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**,
3473 (*PC-M5'*, *PC-M6*)

Antkowiak, R. *et al.*, *Alkaloids (Academic
Press)*, 1991, **40**, 249 (*isol, derivs*)

Mantle, P.G. *et al.*, *Phytochemistry*, 1994, **36**,
1209 (*7-Hydroxypaxilline, 7-Hydroxy-13-
deoxypaxilline, 10-Hydroxy-13-
deoxypaxilline*)


Fueki, S. *et al.*, *Org. Lett.*, 2004, **6**, 2697-2700
(*biosynth*)

Cole, R.J. *et al.*, *Handbook of Toxic
Fungal Metabolites*, Academic Press, 1981,
386

Paxinorol

P-141

[168482-45-9]



$C_{24}H_{27}NO_3$ 377.482


Formally derived from Paxilline,
P-140 by loss of acetone. Prob. an
artifact prod. during prolonged
storage prior to isol. Trace contaminant
of a paxilline-containing extract from
the fungus *Penicillium paxilli*. Solid.
Mp not detd. due to paucity of
material.

Miles, C.O. *et al.*, *J.O.C.*, 1995, **60**, 6067 (*isol,
pmr, cmr, ms, struct*)

Paxiphylline A

P-142

[1000410-97-8]



$C_{24}H_{27}NO_4$ 393.482

Relative
Configuration


Alkaloid from the leaves of *Daphni-
phyllum paxianum*. Pale yellow solid. [α]_D²³ -
221.7 (c, 0.4 in MeOH). λ_{max} 275 (ε
9710); 354 (ε 10960) (MeOH).

Zhang, Y. *et al.*, *Tet. Lett.*, 2007, **48**, 9104-9107
(*isol, pmr, cmr*)

Paxiphylline B

P-143

[1000410-99-0]



$C_{23}H_{25}NO_5$ 395.454


Alkaloid from the leaves of *Daphni-
phyllum paxianum*. Pale yellow solid.
[α]_D²² -326.7 (c, 0.17 in MeOH). λ_{max}
275 (ε 26350); 331 (ε 4480)
(MeOH).

Zhang, Y. *et al.*, *Tet. Lett.*, 2007, **48**, 9104-9107
(*isol, pmr, cmr*)

PC-M4

P-144

[148717-76-4]



$C_{37}H_{49}NO_4$ 571.798


Metab. from *Penicillium crustosum*.
Crystalline powder. Mp 269-272° dec.

Yamaguchi, T. *et al.*, *Phytochemistry*, 1993, **32**,
1177 (*isol, uv, ir, pmr, cmr, cd, struct*)

PC-M5

P-145

[148717-75-3]



$C_{32}H_{41}NO_4$ 503.68


Metab. from *Penicillium crustosum*. Pale
yellow needles (EtOAc). Mp 218-220°
dec.

Yamaguchi, T. *et al.*, *Phytochemistry*, 1993, **32**,
1177 (*isol, uv, ir, pmr, cmr, cd, struct*)

PD 125375

P-146

1,2,10,10a-Tetrahydro-2-ethylidene-10-hydroxydipyrrolo[1,2-c:2',1'-f]pyrimidin-5-one
[106039-74-1]



$C_{12}H_{14}N_2O_2$ 218.255


Structurally related to Tomaymycin, T-424. Isol. from a *Streptomyces* sp. Shows no antimicrobial or antitumour props. Needles (EtOAc). Sol. MeOH; poorly sol. H_2O . Mp 181-183°. $[\alpha]_D^{20} +89.8$ (c, 0.52 in MeOH). λ_{max} 231 (ϵ 9505); 273 (ϵ 12100) (MeOH) (Berdy).

Rithner, C.D. *et al.*, *J.O.C.*, 1987, **52**, 298 (isol, *cryst struct*)

Peceylanine

P-147

[76202-54-5]



$C_{44}H_{54}N_4O_7$ 750.933

The epoxide and olefinic groups may be interchanged. Alkaloid from *Petchia ceylanica* (Apocynaceae). Cryst. (MeOH). Mp 157-158°. $[\alpha]_D^{20} -237$ (c, 1 in $CHCl_3$).


Kunesch, N. *et al.*, *Tet. Lett.*, 1980, **21**, 1727 (*cmr, struct*)

Cavé, A. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1178 (isol, *uv, ir, pmr, struct*)

Peceyline

P-148

[76202-55-6]



$C_{42}H_{48}N_4O_6$ 704.864

The epoxide and olefinic groups may be interchanged. Alkaloid from *Petchia ceylanica* (Apocynaceae). Cryst. (MeOH/ Et_2O). Mp 310° dec. $[\alpha]_D^{20} -355$ (c, 1 in

$CHCl_3$). λ_{max} 218 ($\log \epsilon$ 4.42); 245 ($\log \epsilon$ 4.38); 294 ($\log \epsilon$ 4.2); 340 (sh) ($\log \epsilon$ 4.15); 366 ($\log \epsilon$ 4.41) ($EtOH$).

N^1 -De-Me: **Demethylpeceyline**

[115028-51-8]

$C_{41}H_{46}N_4O_6$ 690.838

Alkaloid from leaves of *Petchia ceylanica* (Apocynaceae). Amorph. $[\alpha]_D^{20} -120$ (MeOH).


Kunesch, N. *et al.*, *Tet. Lett.*, 1980, **21**, 1727-1730 (*cmr, struct*)

Cavé, A. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1178-1182 (isol, *uv, ir, pmr, struct*)

Atta-ur-Rahman, *et al.*, *Planta Med.*, 1988, **54**, 37-40 (*Demethylpeceyline*)

Pecrassipine A

P-149



$C_{26}H_{27}NO_5$ 433.503

(S)-form

Alkaloid from the bark of *Phaeanthus crassipetalus*. Exhibits a vasorelaxant activity on isol. rat aorta ring. Brownish solid. $[\alpha]_D^{26} +27$ (c, 0.4 in MeOH). λ_{max} 298 (ϵ 8200) (MeOH).

(ξ)-form

Me ether: **Vietnamine**

[139232-16-9]

$C_{27}H_{29}NO_5$ 447.53

Alkaloid from the leaves of *Phaeanthus vietnamensis* (Annonaceae).


Nguyen Thi Nghia, *et al.*, *Fitoterapia*, 1991,

62, 315-318 (*Vietnamine*)

Awang, K. *et al.*, *Heterocycles*, 2007, **71**, 2055-2061 (*Pecrassipine A*)

Pecrassipine B

P-150



$C_{26}H_{27}NO_5$ 433.503

(R)-form [953395-30-7]


Alkaloid from the bark of *Phaeanthus crassipetalus*. Brownish solid. $[\alpha]_D^{26} -21$ (c, 0.8 in MeOH). λ_{max} 296 (ϵ 8100) (MeOH).

Awang, K. *et al.*, *Heterocycles*, 2007, **71**, 2055-2061 (isol, *cd, pmr, cmr*)

Pedatisectine A

P-151

6-[(6-Amino-9H-purin-9-yl)methyl]-3-pyridinol, 9CI
[103823-31-0]



$C_{11}H_{10}N_6O$ 242.24


Alkaloid from *Pinellia pedatisecta* (Araceae). Mp 282-284°.

Qin, W. *et al.*, *Zhongcaoyao*, 1986, **17**, 197; *CA*, **105**, 94501a

Pedatisectine C

P-152

[103805-66-9]



$C_{17}H_{14}N_2O_3$ 294.309


Alkaloid from *Pinellia pedatisecta* (Araceae). Mp 162-164°.

Qin, W. *et al.*, *Zhongcaoyao*, 1986, **17**, 197; *CA*, **105**, 94501a

Pedatisectine D

P-153

4-(3-Methylpyrazinyl)-1,2,3-butanetriol, 9CI. 2-Methyl-3-(2,3,4-trihydroxybutyl)pyrazine
[163046-71-7]



$R^1 = R^3 = H$, $R^2 = CH_3$

$C_9H_{14}N_2O_3$ 198.221

Alkaloid from the rhizomes of *Pinellia pedatisecta* (Araceae). Cryst. Mp 110-112°.

Qin, W. *et al.*, *Zhongcaoyao*, 1995, **26**, 3; *CA*, **122**, 298789m

Pedatisectine E

P-154

1-(5-Methylpyrazinyl)-1,2,3,4-butanetetr-ol, 9CI. 2-Methyl-5-(1,2,3,4-tetrahydroxybutyl)pyrazine
[163181-57-5]

As Pedatisectine D, P-153 with $R^1 = OH$, $R^2 = H$, $R^3 = CH_3$


$C_9H_{14}N_2O_4$ 214.221

Alkaloid from the rhizomes of *Pinellia pedatisecta* (Araceae). Cryst. Mp 202-203°. $[\alpha]_D^{27} -87.8$ (c, 0.16 in DMSO).

Qin, W. *et al.*, *Zhongcaoyao*, 1995, **26**, 3; *CA*, **122**, 298789m

Pedatisectine F P-155

1-(6-Methylpyrazinyl)-1,2,3,4-butanetetrol, 9CI. 2-Methyl-6-(1,2,3,4-tetrahydroxybutyl)pyridine [206757-32-6]




C₉H₁₄N₂O₄ 214.221
Alkaloid from the rhizomes of *Pinellia pedatisecta*.

Wang, R. *et al.*, *CA*, 1998, **128**, 306181y

Pedicularidine P-156

[41645-65-2]



C₁₀H₁₁NO 161.203
Alkaloid from *Pedicularis olgae* (Scrophulariaceae). Mp 211-212°. [α]_D²⁰ +67.7.

Carboxylic acid: **Pedicularine**

[28330-57-6]

C₁₀H₁₁NO₂ 177.202


Alkaloid from *Pedicularis olgae* (Scrophulariaceae). Mp 208-209° dec. [α]_D²⁰ -15.3 (c, 0.78 in MeOH).

Khakimdzhanov, S. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 457; 1970, **6**, 142; 1973, **9**, 132; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 383; 1970, **6**, 142; 1973, **9**, 137 (isol, uv, ir, ms, pmr, struct)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431; 468 (rev)

Pediculidine P-157

5,6-Dihydro-7H-cyclohepta[c]pyridin-7-one, 9CI [33579-98-5]



C₁₀H₉NO 159.187
Alkaloid from *Pedicularis olgae* (Scrophulariaceae). Mp 74-75°. Struct. not firmly established.

Picrate: Mp 211-212°.

8,9-Dihydro, 7-alcohol: **Pedculimine**

[33605-28-6]

C₁₀H₁₃NO 163.219

Alkaloid from *Pedicularis olgae* and

Verbascum nobile (Scrophulariaceae). Mp 133-135°.


Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 304; 306; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 292; 294 (Pediculidine, Pediculimine, uv, ir, pmr, ms, isol, struct)

Ninova, P. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 540; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 526 (Pediculimine, isol)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (rev)

Pedculine

[19772-84-0]



C₁₀H₁₁NO 161.203


An impossible struct. has been assigned, and that shown is considered possible. Alkaloid from *Pedicularis olgae* (Scrophulariaceae). Mp 188-189°. [α]_D²⁵ +61.5 (c, 0.95 in EtOH).

Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 136-137; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 117 (isol, uv, ir, ms)

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1977, **16**, 431 (rev)

Pedoxin

3,6-Dihydro-10-hydroxy-9-methyl-4H-pyrindol[4,3-f][1,4]oxazocin-4-one, 9CI [158365-17-4]



C₁₀H₁₀N₂O₃ 206.201


Isol. from the sea urchin *Toxopneustes pileolus*. Prosthetic group of the protein toxin, Peditoxin. Sedative, anaesthetic. Yellow cryst. λ_{max} 256 ; 330 (pH 2 buffer) (Berdy). λ_{max} 334 (pH 7 buffer) (Berdy).

► LD₅₀ (mus, scu) 250 mg/kg.

Kuwabara, S. *et al.*, *J. Biol. Chem.*, 1994, **269**, 26734-26738

Peduncin

P-160



C₂₉H₃₇NO₄ 463.616


Alkaloid from *Pueraria peduncularis*. Platelets. Mp > 300°.

Li, N. *et al.*, *Chin. Chem. Lett.*, 2001, **12**, 899-902

Peduncularidine

P-161

[168922-22-3]



C₄₃H₅₄N₄O₈ 754.922

Alkaloid from leaves of *Ervatamia peduncularis* (Apocynaceae). Purple amorph. solid. λ_{max} 204 (sh) ; 208 ; 263 ; 307 ; 337 (MeOH).

Zèches-Hanrot, M. *et al.*, *Phytochemistry*, 1995, **40**, 587-591 (isol, uv, ir, pmr, cmr, ms, struct)


Peduncularine

P-162

Isopeduncularine

[34964-75-5]

[99396-15-3]



Absolute configuration

C₂₀H₂₄N₂ 292.423

Before 1992 Peduncularine was confused with the hydrochloride. Iso-peduncularine was eventually found to be identical with Peduncularine. Alkaloid from the stem and root of *Aris-totelia peduncularis* (Elaeocarpaceae). Shows weak antitumour activity. Small needles (CHCl₃). Mp 113-114°. [α]_D¹⁹ -53 (c, 1.2 in MeOH). [α]_D¹⁹ -40 (c, 4.1 in CHCl₃).

Hydrochloride:

Small needles (CHCl₃). Mp 155-157°.

7-Epimer: [119241-74-6]

C₂₀H₂₄N₂ 292.423

Mp 118-125°. [α]_D²⁰ +4.1 (c, 0.435 in CHCl₃). [α]_D²⁰ -13 (c, 0.425 in MeOH).

This struct. was erroneously assigned to Isopeduncularine in 1985.

Bick, I.R.C. *et al.*, *Chem. Comm.*, 1971, 1155-1156 (pmr, ms)

Ros, H.-P. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 481-487 (isol, uv, ir, pmr, cmr, ms, struct)

Klaver, W.J. *et al.*, *J.A.C.S.*, 1989, **111**, 2588-2595 (synth, ir, pmr, cmr, abs conf, bibl)

Dragar, C. *et al.*, *Phytochemistry*, 1992, **31**, 3601-3603 (pmr, cmr)


Roberson, C.W. *et al.*, *J.A.C.S.*, 2002, **124**, 11342-11348 (synth)

Washburn, D.G. *et al.*, *Org. Lett.*, 2003, **5**, 3523-3525 (synth)

Kitamura, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 2006, **79**, 1552-1560 (synth)

Peepuloidine**P-163**

1-[3-(6,7-Dimethoxy-1,3-benzodioxol-5-yl)-1-oxo-2-propenyl]pyrrolidine, 9CI. 1-[2,3-Dimethoxy-4,5-(methylenedioxy)-cinnamoyl]pyrrolidine, 8CI. 2,3-Dimethoxy-4,5-methylenedioxy-cinnamic acid pyrrolidide

C₁₆H₁₉NO₅ 305.33**(E)-form** [25990-52-7]


Alkaloid from the leaves of *Piper peepuloides* (Piperaceae). Cryst. (C₆H₆/petrol or EtOH), Mp 149-150°.

Atal, C.K. *et al.*, *Tet. Lett.*, 1968, 1397 (*uv, ir, pmr, ms, struct*)

Dallacker, F. *et al.*, *Chem. Ber.*, 1971, **104**, 1706 (*synth, uv, ir*)

Peganidine**P-164**

1-(1,2,3,9-Tetrahydro-3-hydroxypyrrolo[2,1-b]quinazolin-9-yl)-2-propanone, 9CI [28463-17-4]

C₁₄H₁₆N₂O₂ 244.293

Alkaloid from *Peganum harmala* (Zygophyllaceae). Mp 189-190°.

Oxime: Mp 85-87°.

Semicarbazone: Mp 204-206°.

Deoxy: 1-(1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazolin-9-yl)-2-propanone, 9CI.

Deoxypeganidine

[42405-57-2]

C₁₄H₁₆N₂O 228.293

Alkaloid from *Peganum harmala* (Zygophyllaceae). Noncryst. Mp 76-79°.

Deoxy, picrate: Mp 176°.

Deoxy, oxime: Mp 177-179°.

Stereoisomer: Isopeganidine

[53448-58-1]

C₁₄H₁₆N₂O₂ 244.293

Alkaloid from *Peganum harmala* (Zygophyllaceae). Mp 169-170°. Said to be a racemic diastereoisomer of Peganidine.


Khashimov, K.N. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 599; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 518 (*isol, uv, ir, pmr, ms, struct*)

Zharekeev, B.K. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 279; 1974, **10**, 264-265; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 272; 1974, **10**, 282 (*Deoxypeganidine, Isopeganidine*)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 249-256 (*pmr*)

Peganine**P-165**

1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazolin-3-ol, 9CI. 1,2,3,9-Tetrahydro-3-hydroxypyrrolo[2,1-b]quinoline. *Vasicine*. *Linarine*† [50591-64-5]

**(R)-form**C₁₁H₁₂N₂O 188.229

Abs. config. reversed in 1996. Bronchodilator and respiratory stimulant. Shows antihypertensive props. Uterine stimulant and abortifacient. Expectorant.

(R)-form

Alkaloid from *Galega officinalis* (Fabaceae). Mp 203-205° dec. [α]_D²⁴ +203 (+188, +163).

(S)-form [6159-55-3]

Alkaloid from the leaves of *Adhatoda vasica*, the roots of *Sida cordifolia*, and from several other *Sida* spp. and *Lunaria* spp. Also present in the seeds of *Peganum harmala* (Acanthaceae, Malvaceae, Brassicaceae, Zygophyllaceae). Mp 211-212°. [α]_D¹⁴ -254 (c, 2.44 in CHCl₃). Pharmacol. active isomer. Racemises slowly in acid or upon repeated subl. ▶LD₅₀ (rat, orl) 640 mg/kg. Exp. reprod. effects. UY9454000

Hydrochloride:Cryst. + 2H₂O. Mp 208° (anhyd.).*Picrate*: Mp 199° dec.

7-Hydroxy: see Vasicinol, V-37

(±)-form [6159-56-4]

Major alkaloid of *Anisotes sessiliflorus*. Also present in the above-ground parts of *Peganum harmala* and the foliage of *Peganum nigellastrum* (Acanthaceae, Zygophyllaceae). Cryst. (Et₂O or MeOH/C₆H₆). Mp 211-212° (204-206°).

N⁴-Oxide: Peganine N-oxide

[170712-17-1]

C₁₁H₁₂N₂O₂ 204.228Alkaloid from *Nitraria komarovii*.Cryst. (EtOH/Me₂CO). Mp 207-208°.**5-Methoxy: 1,2,3,9-Tetrahydro-5-methoxypyrrolo[2,1-b]quinazolin-3-ol, 9CI. 5-Methoxypeganine**

[100477-91-6]

C₁₂H₁₄N₂O₂ 218.255

Alkaloid from the leaves of *Adhatoda vasica* (Acanthaceae). Cryst. (EtOH). Mp 224-225°.

[7174-27-8, 79378-22-6]

Späth, E. *et al.*, *Ber.*, 1935, **68**, 1384 (*resoln*)
Leonard, J.N. *et al.*, *Tet. Lett.*, 1960, **No. 25**, 44 (*synth*)

Schreiber, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 271; *CA*, **57**, 12632e (*isol*)

Mehta, D.R. *et al.*, *J.O.C.*, 1963, **28**, 445 (*isol*)
Bhatnagar, A.K. *et al.*, *Indian J. Chem.*, 1966, **4**, 291 (*ms*)

Arndt, R.R. *et al.*, *Tetrahedron*, 1967, **23**, 3521 (*isol, uv, pmr*)

Liljegren, D.R. *et al.*, *Phytochemistry*, 1971, **10**, 2661 (*biosynth*)

Johne, S. *et al.*, *Pharmazie*, 1973, **28**, 403 (*biosynth*)

Ghosal, S. *et al.*, *Phytochemistry*, 1975, **14**, 830 (*isol*)

Szulzewsky, K. *et al.*, *J. Prakt. Chem.*, 1976, **318**, 463 (*cryst struct, abs config*)

Johne, S. *et al.*, *J. Prakt. Chem.*, 1977, **319**, 919 (*cmr*)

Sharma, R.L. *et al.*, *Indian J. Chem., Sect. B*, 1979, **18**, 449 (*synth, pmr*)

Al-Shamma, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 745 (*isol*)

Chowdhury, B.K. *et al.*, *Phytochemistry*, 1985, **24**, 3080 (*5-methoxy*)

Johne, S. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 129 (*rev, pharmacol*)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1991, **32**, 7131 (*synth*)

Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1994, **30**, 780-783; *Chem. Nat. Compd. (Engl. Transl.)*, 1994, **30**, 727-729 (*N-oxide*)

Turgunov, K.K. *et al.*, *Khim. Prir. Soedin.*, 1995, **31**, 426-430; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 353-356 (*cryst struct*)

Joshi, B.S. *et al.*, *Tetrahedron: Asymmetry*, 1996, **7**, 25 (*cryst struct, abs config*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VGA000

Peimidine**P-166**C₂₇H₄₅NO₂ 415.658

Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria roylei* (Liliaceae), occurring with Peimisine, P-168. Mp 222°. [α]_D²⁶ -74 (EtOH).

Chou, T.-Q. *et al.*, *J. Am. Pharm. Assoc., Sci. Ed.*, 1947, **36**, 215-217

Peimiphine**P-167**C₂₇H₄₆NO₃ 432.665


Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria roylei* (Liliaceae), *isol.* with Peimisine, P-168. Mp 127°. [α]_D²¹ -69 (EtOH).

Hydrochloride: Mp 318°.

Chou, T.Q. *et al.*, *J. Am. Pharm. Assoc., Sci. Ed.*, 1947, **36**, 215-217

Peimisine**P-168***Alkaloid FR 5, Ebeiensine*

[19773-24-1]

C₂₇H₄₁NO₃ 427.626

Alkaloid from *Fritillaria thunbergii*, *Fritillaria roylei*, *Fritillaria siechuanica* and *Petilium eduardii* (Liliaceae), and (as Ebeiensine) from *Fritillaria ebeiensis*, *Fritillaria pugiensis* and *Fritillaria ebeiensis* var. *purpurea*. Needles (MeOH). Mp 270° (255°). [α]_D -33.2 (c, 1 in CHCl₃). [α]_D²⁵ -44.6 (c, 0.58 in EtOH).

Hydrochloride: Mp 257° (250-252°).

Di-Ac:

Fine needles (EtOAc/hexane). Mp 231-232°. $[\alpha]_D^{25} +7.2$ (c, 1 in CHCl_3).

Cho, T.Q. *et al.*, *J. Am. Pharm. Assoc., Sci. Ed.*, 1947, **36**, 215-217 (*isol*)

Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 412-413; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 345-346 (*pmr, struct*)

Katajima, J. *et al.*, *Heterocycles*, 1981, **15**, 791-796 (*isol, struct, ms*)

Wu, J. *et al.*, *CA*, 1992, **116**, 170082j; 1993, **118**, 19193s (*Ebeinsine*)

Wang, I.P. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 273-278 (*struct*)

Peimitidine**P-169**

$\text{C}_{27}\text{H}_{44}\text{NO}_3$ 430.649

Steroidal alkaloid. Struct. unknown.

Alkaloid from *Fritillaria roylei* (Liliaceae), occurring with Peimisine, P-168. Mp 188°. $[\alpha]_D^{20} -68$ (EtOH).

Chou, T.Q. *et al.*, *J. Am. Pharm. Assoc., Sci. Ed.*, 1947, **36**, 215-217

Peimunine**P-170**

$\text{C}_{19}\text{H}_{30}\text{NO}_2$ 304.451


Tentative mol. formula, may be $\text{C}_{19}\text{H}_{32}\text{NO}_2$. Prob. a steroidal alkaloid. Struct. unknown. Alkaloid from the Chinese drug Pei-mu. Smooth muscle relaxant. Mp 223-224°. $[\alpha]_D^{26} -27.6$.

Liu, S.K. *et al.*, *CA*, 1937, **31**, 754

Li, S.Y. *et al.*, *CA*, 1942, **36**, 1940

Peinamine**P-171**

[64625-88-3]



Absolute Configuration

$\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706

Alkaloids covered by this entry (1*S*,1'*R*-series) are enantiomeric with those in Isotetrandrine, I-328 and diastereomeric with those in Tetrandrine, T-286 and Phaeanthine, P-301. Alkaloid from *Abuta grisebachii* (Menispermaceae) and a curare of the upper Orinoco. Mp 170-171°. $[\alpha]_D^{20} -109$ (CHCl_3).

*O*⁷-*De-Me*: **7-O-Demethylpeinamine**

[66254-50-0]

$\text{C}_{33}\text{H}_{36}\text{N}_2\text{O}_6$ 580.679

Alkaloid from *Abuta grisebachii* (Menispermaceae). Mp 205-206°. $[\alpha]_D^{20} -86$ (MeOH).

*O*⁷-*De-Me*, *N*²-*Me*: **N-Methyl-7-O-demethylpeinamine**

[66254-51-1]


$\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706

Alkaloid from *Abuta grisebachii* (Menispermaceae). Mp 187-190°. $[\alpha]_D^{20} -259$ (CHCl_3).

Galeffi, C. *et al.*, *Farmacol. Ed. Sci.*, 1977, **32**, 665; 853; *CA*, 1980683c; **89**, 6465q (*isol, uv, ord, ms, struct*)

Pelankine**P-172**

[76202-56-7]



Absolute Configuration

$\text{C}_{42}\text{H}_{48}\text{N}_4\text{O}_6$ 704.864


The epoxide and olefinic groups may be interchanged. Alkaloid from *Petchia ceylanica* (Apocynaceae). Amorph. $[\alpha]_D -214$ (c, 0.44 in CHCl_3). λ_{max} 218 (log ϵ 4.34); 238 (log ϵ 4.29); 285 (log ϵ 4.08); 330 (log ϵ 4.05); 358 (log ϵ 4.19) (EtOH).

Kunesch, N. *et al.*, *Tet. Lett.*, 1980, **21**, 1727-1730 (*cmr, struct*)

Cavé, A. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1178-1182 (*isol, uv, ir, pmr, struct*)

Pelletierine**P-173**

1-(2-Piperidinyl)-2-propanone, 9CI. 2-Acetyl-piperidine. Isopelletierine. 8-Methylnorlobelone [4396-01-4]



(*R*)-form

$\text{C}_8\text{H}_{15}\text{NO}$ 141.213

Pelletierine was originally given the erroneous struct. of 3-(2-piperidinyl)propanal. The early use of the names Pelletierine and Isopelletierine is confusing. Anthelmintic agent. Log P 0.56 (calc).

(*R*)-form [2858-66-4]

Alkaloid from *Punica granatum* (pomegranate) (Punicaceae). Liq. $[\alpha]_D^{23} -18.1$ (c, 8.18 in EtOH). The isoln. of the (*R*)-form is doubtful in view of its ready racemisation and of the fact that the opt. rotn. recorded for the isolate exceeded that of the resolved base.

►RX9625000

Picrate: Mp 131-132°.

N-Me: $[\alpha]_D^{14} -51.5$ (c, 4.8 in EtOH).

(*S*)-form [2858-67-5]

Synthetic. Liq. $[\alpha]_D^{23} +17.9$ (c, 9.41 in EtOH aq.).

(±)-form [539-00-4]

Alkaloid from pomegranate (*Punica granatum*), *Duboisia myoporoides* and *Sedum acre* (Punicaceae, Solanaceae, Crassulaceae). Bp₁₄ 91-92°. p*K*_a 9.45 (15°). Readily oxid. in air.

►RX9635000

Hydrochloride: [5984-61-2]

Needles. Mp 143°.

Picrate: Mp 150-151° (147-149°).

N-Benzoyl:

$\text{C}_{15}\text{H}_{19}\text{NO}_2$ 245.321

Prisms or plates. Mp 75°.

N-Me: Methyloisopelletierine. α-N-Methylpelletierin-β-one. Methylpelletierine

$\text{C}_9\text{H}_{17}\text{NO}$ 155.239

Alkaloid from pomegranate (*Punica granatum*), *Lupinus formosus*, *Sedum purpureum*, *Sedum hybridum* and *Sedum aizoon* (Fabaceae, Crassulaceae). Misc. H_2O . Bp₂₆ 114-117°. Strong base. The isoln. of this alkaloid in opt. active form was reported by Tanret (1880), but this has not been repeated.

N-Me; hydrochloride: Mp 158°.

N-Me; hydrobromide: Mp 151-153°.

N-Me, picrate: Mp 158°.

Hess, K. *et al.*, *Ber.*, 1917, **50**, 1192 (*isol*)

Galinowsky, F. *et al.*, *Monatsh. Chem.*, 1951, **82**, 551; 1953, **84**, 1221; 1954, **85**, 1012 (*struct*)

Mortimer, P.I. *et al.*, *J.C.S.*, 1957, 3967 (*isol*)

Franck, B. *et al.*, *Chem. Ber.*, 1958, **91**, 2803 (*isol*)

Kuwata, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1960, **33**, 1668; 1672 (*isol, struct, bibl*)

Gilman, R.E. *et al.*, *Bull. Soc. Chim. Fr.*, 1961, 1993 (*struct*)

Khanna, K.L. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 1194 (*isol*)

Drillien, G. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2393 (*rev*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1967, **86**, 80 (*abs config*)

Fitch, W.L. *et al.*, *J.O.C.*, 1974, **39**, 2974 (*isol, deriv*)

Hanaoka, M. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 531; *CA*, **81**, 13670r (*synth*)



Krasnov, E.A. *et al.*, *Khim. Prir. Soedin.*, 1977, 585; *Chem. Nat. Compd. (Engl. Transl.)*, 492 (*isol, deriv*)

Cymerman Craig, J. *et al.*, *J.O.C.*, 1978, **43**, 347 (*cd*)

Quick, J. *et al.*, *J.O.C.*, 1979, **44**, 573 (*synth*)

Nagasaka, T. *et al.*, *Heterocycles*, 1989, **29**, 155 (*synth, ir, pmr*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PAO500

Penaramides**P-174****Penaresamides**


Inseparable mixt. The simplest component Penaramide A has been synthesised. Isol. from the marine sponge *Penares* aff. *incrustans*. Inhibits binding of ω-Conotoxin GVIA to N-type Ca channels. Yellowish oil. $[\alpha]_D^{20} -11$ (c, 0.39 in $\text{CHCl}_3/\text{MeOH}$ 1:1).

Ushio-Sata, N. *et al.*, *Tet. Lett.*, 1996, **37**, 225 (*isol, ir, pmr, cmr, synth, struct*)

Penaresidin B

P-175

3-Hydroxy-4-(hydroxymethyl)- α -(2-methylpropyl)-2-azetidinedecanol, 9CI
[135574-63-9]




$C_{19}H_{39}NO_3$ 329.522

Alkaloid from the marine sponge *Penares* sp. Potent ATPase activator. Isol. as an inseparable mixt. with Penaresidin A.

- Kobayashi, J. *et al.*, *J.C.S. Perkin 1*, 1991, 1135-1137 (*isol, struct*)
 Kobayashi, J. *et al.*, *Tet. Lett.*, 1996, **37**, 6775-6776 (*abs config*)
 Takikawa, H. *et al.*, *J.C.S. Perkin 1*, 1997, 97-111 (*synth*)
 Yoda, H. *et al.*, *Tet. Lett.*, 1997, **38**, 3283-3284 (*synth*)
 Liu, D.-G. *et al.*, *Tet. Lett.*, 1999, **40**, 337-340 (*synth*)
 Yoda, H. *et al.*, *Tet. Lett.*, 2003, **44**, 977-979 (*synth*)

Penarolide A₁

P-176



$C_{35}H_{65}NO_6$ 595.902

Tri-O-sulfate: **Penarolide sulfate A₁**
[329019-71-8]


$C_{35}H_{65}NO_{15}S_3$ 836.094

Isol. from the marine sponge *Penares* sp. α -Glucosidase inhibitor. Amorph. solid (as tri-Na salt). $[\alpha]_D^{29}$ -25.7 (c, 0.5 in MeOH) (tri-Na salt). λ_{max} 204 (ϵ 9100) (MeOH) (tri-Na salt).

Nakao, Y. *et al.*, *Tetrahedron*, 2000, **56**, 8977-8987

Penarolide A₂

P-177



$C_{35}H_{65}NO_6$ 595.902

Tri-O-sulfate: **Penarolide sulfate A₂**
[329019-70-7]


$C_{35}H_{65}NO_{15}S_3$ 836.094

Isol. from the marine sponge *Penares* sp. α -Glucosidase inhibitor. Amorph. solid (as tri-Na salt). $[\alpha]_D^{29}$ -25.2 (c, 0.3 in MeOH) (tri-Na salt). λ_{max} 205 (ϵ 9000) (MeOH) (tri-Na salt).

Nakao, Y. *et al.*, *Tetrahedron*, 2000, **56**, 8977-8987

Penasulfate A

P-178



$C_{36}H_{69}NO_{11}S_2$ 756.073


Isol. as a 4:1 mixt. of 2*R*- and 2*S*-stereoisomers. Alkaloid from the sponge *Penares* sp. α -Glucosidase inhibitor. Amorph. solid (as di-Na salt). $[\alpha]_D^{29}$ +10 (c, 0.03 in MeOH) (di-Na salt). λ_{max} 203 (ϵ 9700) (MeOH) (di-Na salt).

Nakao, Y. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1346-1350

Pencolide

P-179

α -Ethylidene-2,5-dihydro-3-methyl-2,5-dioxo-1*H*-pyrrole-1-acetic acid, 9CI



$C_9H_9NO_4$ 195.174

(*Z*)-form [61464-52-6]


Fungal metab. of *Penicillium multicolor* and *Trichoderma album*. Rosettes (Et₂O/petrol). Mp 135.5° (113-118°). Formerly assigned the (*E*)-config.

- Birkinshaw, J.H. *et al.*, *Biochem. J.*, 1963, **86**, 237 (*isol, uv, ir, pmr*)
 Strunz, G.M. *et al.*, *Can. J. Chem.*, 1976, **54**, 2862 (*synth, config, bibl*)
 Ren, W.Y. *et al.*, *Diss. Abstr. Int., B*, 1978, **39**, 1293 (*isol, synth*)

Pendolmycin

P-180

[119375-01-8]



$C_{22}H_{31}N_3O_2$ 369.506

Isol. from *Nocardopsis* sp. SA 1715. Induces erythema in human skin. Exhibits weak antibacterial activity. Tumour promoter. Protein kinase activator. Cryst. Sol. MeOH, CHCl₃, Me₂CO; poorly sol. H₂O, hexane. Mp 124-126°. $[\alpha]_D$ -154 (c, 0.1 in MeOH). Related to Lyngbyatoxin A, L-367 and Teleocidin B₁, T-63. λ_{max} 214 (ϵ 11800); 230 (ϵ 19400); 287 (ϵ 7900); 298 (ϵ 7970) (MeOH) (Derep).


Yamashita, T. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1184 (*isol, pmr, cmr, struct*)

Okabe, K. *et al.*, *Tetrahedron*, 1990, **46**, 5113; 1991, **47**, 7615 (*synth*)

Muratake, H. *et al.*, *Tetrahedron*, 1991, **47**, 8535; 8545; 8559 (*synth*)

Pendulamine A

P-181



$C_{19}H_{19}NO_5$ 341.363

(*S*)-form

Alkaloid from the roots of *Polyalthia longifolia* var. *pendula*. Brown solid. $[\alpha]_D^{26}$ -25 (c, 0.4 in CHCl₃). λ_{max} 204; 226; 282; 322; 366 (MeOH).

13,14-Didehydro: **Pendulamine B**

$C_{19}H_{17}NO_5$ 339.347


Alkaloid from the roots of *Polyalthia longifolia* var. *pendula*. Reddish-brown gum. λ_{max} 207; 224; 262; 335; 366; 388 (MeOH).

Faizi, S. *et al.*, *Planta Med.*, 2003, **69**, 350-355 (*isol, pmr, cmr, ms*)

Pengbeimine A

P-182

[909114-88-1]



$C_{27}H_{43}NO_3$ 429.642

Alkaloid from the bulbs of *Fritillaria monatha*. Needles (Me₂CO).

N-Me: **Pengbeimine B**

[1016260-11-9]

$C_{28}H_{45}NO_3$ 443.668

Alkaloid from the bulbs of *Fritillaria monatha*. Cryst. (MeOH). Mp 275-276°.

20-Epimer, N-Me: **Pengbeimine D**

[1016260-12-0]

$C_{28}H_{45}NO_3$ 443.668

Alkaloid from the bulbs of *Fritillaria monatha*. Cryst. Mp 259-261°.


Liu, H.-N. *et al.*, *Chin. Chem. Lett.*, 2006, **17**, 631-634 (*Pengbeimine A, cryst struct*)

Liu, H.-N. *et al.*, *J. Asian Nat. Prod. Res.*, 2007, **9**, 563-567 (*Pengbeimines B,D*)

Pengbeimine C

P-183


[909114-89-2]



C₂₈H₄₅NO₃ 443.668
Alkaloid from the bulbs of *Fritillaria monatha*. Needles (Me₂CO).


Liu, H.-N. *et al.*, *Chin. Chem. Lett.*, 2006, 17, 631-634 (isol, pmr, cmr, cryst struct)

Pengshenine A P-184



C₂₄H₃₇NO₆ 435.559
Alkaloid from the roots of *Aconitum hemslayanum* var. *penzhouense*. Needles. Mp 189-191°. [α]_D²⁰ +26 (c, 0.5 in CHCl₃). Peng, C.S. *et al.*, *Chin. Chem. Lett.*, 2002, 13, 233-236

Peniamidienone P-185




C₁₉H₂₃NO₃ 313.396
Prod. by *Penicillium* sp. No. 13. Yellow oil. [α]_D²⁵ -141 (c, 0.41 in MeOH). λ_{max} 204 (log ε 3.66); 227 (log ε 3.8); 260 (log ε 3.16); 370 (log ε 2.72) (EtOH).

Kimura, Y. *et al.*, *Phytochemistry*, 2000, 53, 829-831

Penicidone A P-186

[1005327-87-6]



Absolute Configuration

C₁₈H₁₇NO₅ 327.336
Prod. by *Penicillium* sp. IFB-E022. Cytotoxic. Amorph. powder. [α]_D²⁰ -98.3 (c, 0.11 in CHCl₃). λ_{max} 206 (log ε 4.6); 237 (log ε 3.1) (MeOH).


O⁶-De-Me: Penicidone B

[1005327-88-7]
C₁₇H₁₅NO₅ 313.309
Prod. by *Penicillium* sp. IFB-E022. Cytotoxic agent. Amorph. powder. [α]_D²⁰ -87.4 (c, 0.2 in CHCl₃). λ_{max} 209 (log ε 4.4); 239 (log ε 4) (MeOH).

Ge, H.M. *et al.*, *Phytochemistry*, 2008, 69, 571-576 (isol, pmr, cmr)

Penicidone C P-187

[1005327-89-8]




C₁₉H₁₉NO₆ 357.362
Prod. by *Penicillium* sp. IFB-E022. Cytotoxic. Amorph. powder. λ_{max} 215 (log ε 4.1); 253 (log ε 4.3) (MeOH).

Ge, H.M. *et al.*, *Phytochemistry*, 2008, 69, 571-576 (isol, pmr, cmr)

Penicillenol A₁ P-188

[1039041-76-3]



C₁₆H₂₇NO₄ 297.394
Tetramic acid deriv. Prod. by *Penicillium* sp. GQ-7. Cytotoxic. Yellow oil. [α]_D²⁵ -864.5 (c, 0.155 in MeOH). λ_{max} 223; 279 (MeOH).

6',7'-Didehydro(E-): Penicillenol C₁

[1039041-80-9]
C₁₆H₂₅NO₄ 295.378
Prod. by *Penicillium* sp. GQ-7. Yellow oil. [α]_D²⁵ -47.2 (c, 0.12 in MeOH). λ_{max} 224; 280 (MeOH).

6-Deoxy, 5,6-didehydro(E-): Penicillenol B₂

[1039041-79-6]
C₁₆H₂₅NO₃ 279.378
Prod. by *Penicillium* sp. GQ-7. Yellow oil. [α]_D²⁵ -15.9 (c, 0.13 in MeOH). λ_{max} 256 (MeOH).

6-Deoxy, 5,6-didehydro(Z-): Penicillenol B₁

[1039041-78-5]
C₁₆H₂₅NO₃ 279.378
Prod. by *Penicillium* sp. GQ-7. Cytotoxic. Yellow oil. [α]_D²⁵ -7.8 (c, 0.23 in MeOH). λ_{max} 256 (MeOH).

5-Epimer: Penicillenol A₂

[1039041-77-4]
C₁₆H₂₇NO₄ 297.394
Prod. by *Penicillium* sp. GQ-7. Yellow oil. [α]_D²⁵ +386.7 (c, 0.13 in MeOH). λ_{max} 223; 279 (MeOH).

5-Epimer, 6',7'-didehydro(Z-): Penicillenol C₂


[1039041-81-0]
C₁₆H₂₅NO₄ 295.378
Prod. by *Penicillium* sp. GQ-7. Yellow oil. [α]_D²⁵ +32 (c, 0.2 in MeOH). λ_{max}

224; 280 (MeOH).

Lin, Z.-J. *et al.*, *Chem. Pharm. Bull.*, 2008, 56, 217-221 (isol, cd, pmr, cmr)

Penidiamide P-189

2-Amino-N-[2-[[2-(1H-indol-3-yl)ethenyl]amino]-2-oxoethyl]benzamide, 9CI
[206049-48-1]




C₁₉H₁₈N₄O₂ 334.377
Prod. by *Penicillium* sp. 62-92. Inhibitor of synovial phospholipase A₂. Oil. λ_{max} 279 (ε 13900); 303 (ε 16300) (MeOH).

Witter, L. *et al.*, *Z. Naturforsch., C*, 1998, 53, 60-64 (isol, pmr, cmr)

Penidilamine P-190

3,3'-Iminobis[5-methyl-2(5H)-furanone], 9CI
[279679-13-9]




C₁₀H₁₁NO₄ 209.201
Prod. by *Penicillium* sp. No. 13. Cryst. (hexane/Me₂CO). Mp 187-189°. [α]_D²⁵ +33 (c, 0.31 in MeOH). λ_{max} 207 (log ε 3.83); 260 (log ε 3.5); 278 (log ε 3.64) (EtOH).

Kimura, Y. *et al.*, *Phytochemistry*, 2000, 53, 829-831 (isol)

Penigequinolone A P-191

[180045-91-4]

[184046-64-8]



Relative Configuration

C₂₇H₃₃NO₆ 467.561
Isol. as an inseparable mixt. with Penigequinolone B (to which data refers). Prod. by *Penicillium* sp. No. 410 and *Penicillium* cf. *simplicissimum*. Pollen growth inhibitor. Powder. Mp 111-113°. [α]_D²⁰ +97.8 (c, 0.4 in EtOH).

23-Hydroxy: Yaecinolone D

C₂₇H₃₃NO₇ 483.56
Prod. by *Penicillium* sp. FKI-2140. Insecticidal agent. Pale yellow powder. [α]_D²⁵ +56 (c, 0.1 in EtOH). Stereochem. of side chain not determined. λ_{max} 221 (ε 20700); 279 (ε 12600); 288 (ε 9500); 324 (ε 10700) (EtOH).

19-Epimer: Penigequinolone B

[180185-69-7]

[184046-66-0]

C₂₇H₃₃NO₆ 467.561Prod. by *Penicillium* sp. No. 410 and *Penicillium* cf. *simplicissimum*. Pollen growth inhibitor.Kimura, Y. et al., *Tet. Lett.*, 1996, **37**, 4961-4964 (*isol, ir, pmr, cmr, struct*)Kusano, M. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2559-2568 (*isol, pmr, cmr*)Uchida, R. et al., *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*Yaequinolone D*)**Penimide****P-192****N-Methyl-N-[3-(methylthio)-1-oxo-2-propenyl]benzeneacetamide, 9CI**
MeSCH=CHCONMeCOCH₂PhC₁₃H₁₅NO₂S 249.333


Originally reported as the isomeric compds. with the phenyl and methylthio groups exchanged.

(E)-form**Penimide A**

[150036-29-6]

Isol. from leaves of *Glycosmis chlorosperma* (Rutaceae). Oil.**(Z)-form****Penimide B**


[150036-30-9]

From leaves of *Glycosmis chlorosperma* (Rutaceae). Mp 85-87°.Greger, H. et al., *Phytochemistry*, 1993, **32**, 933-936 (*isol, uv, ir, pmr, cmr, ms*)Hinterberger, S. et al., *Tetrahedron*, 1994, **50**, 6279-6286 (*synth, struct*)**Peniprequinolone****P-193**

Relative Configuration

C₂₂H₂₅NO₅ 383.443Prod. by *Penicillium* cf. *simplicissimum* and a marine-derived *Penicillium janczewski*. Cytotoxic. Pale yellowish solid. Mp 191-194°. [α]_D²⁰ -0.3 (c, 0.6 in CHCl₃). λ_{max} 206 (ε 74000); 222 (ε 65000); 302 (ε 19000) (EtOH).**A^{3'}-Isomer, 1',2'-didehydro(E)-: Yaequinolone E**C₂₂H₂₃NO₅ 381.427Prod. by *Penicillium* sp. FKI-2140. Insecticidal agent. Pale yellow powder. [α]_D²³ +51.2 (c, 0.1 in EtOH). λ_{max} 223 (ε 18800); 280 (ε 8400); 291 (ε 8300); 324 (ε 6800) (EtOH).Kusano, M. et al., *Biosci., Biotechnol., Biochem.*, 2000, **64**, 2559-2568 (*isol, cd, pmr, cmr, ms*)Hu, J. et al., *J. Nat. Prod.*, 2005, **68**, 1397-1399 (*isol*)Uchida, R. et al., *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*Yaequinolone E*)**Penitrem C**

[37318-84-6]



Absolute Configuration

C₃₇H₄₄ClNO₄ 602.212Metab. from cultures of *Penicillium crustosum*, *Penicillium granulatum*, *Penicillium puberulum*, *Penicillium palitans* and *Penicillium cycloptium* NRRL3476. Neurotoxic, tremorgenic substance. Insecticide. Amorph. solid. λ_{max} 232 (ε 38200); 292 (ε 12500) (MeOH) (Derep).**23α,24α-Epoxyde: Penitrem F**

[78213-65-7]

C₃₇H₄₄ClNO₅ 618.211Isol. from *Penicillium crustosum*. Tremorgenic agent. Amorph. powder. λ_{max} 233 (ε 37000); 295 (ε 11600) (MeOH) (Derep). λ_{max} 232; 292 (MeOH) (Berdy).**15-Hydroxy, 23α,24α-epoxyde: Penitrem A. Tremortin A**

[12627-35-9]

[147382-80-7]

C₃₇H₄₄ClNO₆ 634.21Metab. from cultures of *Penicillium crustosum*. Potassium BK_{Ca} channel blocker. Tremorgenic metab. Amorph. solid. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O, hexane. Mp 237-239°. λ_{max} 233 (ε 37000); 295 (ε 11600) (MeOH) (Derep). λ_{max} 233 (ε 31500); 295 (ε 10200) (MeOH) (Berdy).▶ Toxic. LD₅₀ (mus, ipr) 1.1 mg/kg. RY7535000**Bromo analogue, 15-hydroxy, 23α,24α-epoxyde: 6-Bromopenitrem E. 6-Bromo-6-dechloropenitrem A**

[86271-57-0]

C₃₇H₄₄BrNO₆ 678.662Isol. from *Penicillium simplicissimum* AK-40. λ_{max} 233; 295 (MeOH) (Berdy).**Dechloro: Penitrem D**

[78213-64-6]

C₃₇H₄₅NO₄ 567.767Metab. from cultures of *Penicillium crustosum*, *Penicillium palitans* and *Penicillium cycloptium* NRRL3476. Tremorgenic mycotoxin. Needles (C₆H₆). Mp 300° dec. λ_{max} 220 (ε 31900); 286 (ε 11900) (MeOH) (Derep).**Dechloro, 23α,24α-epoxyde: Penitrem B**

[11076-67-8]

C₃₇H₄₅NO₅ 583.766Isol. from *Penicillium crustosum* and *Penicillium palitans*. Tremorgenic agent. Needles (EtOH). Mp 185-195° dec. λ_{max} 220 (ε 31900); 286 (ε 11900) (MeOH) (Derep). λ_{max} 227; 286 (MeOH) (Berdy). λ_{max} 222; 286; 297 (EtOH) (Berdy).▶ LD₅₀ (mus, ipr) 5.8 mg/kg. RY7537000**P-194****Dechloro, 15-hydroxy, 23α,24α-epoxyde:****Penitrem E. 6-Dechloropenitrem A**

[78213-66-8]

C₃₇H₄₅NO₆ 599.766Isol. from *Penicillium crustosum*. Tremorgenic. Amorph. powder. λ_{max} 220 (ε 31900); 286 (ε 11900) (MeOH) (Derep).**Dechloro, 19-hydroxy: Penitrem G**C₃₇H₄₅NO₅ 583.766Prod. by *Penicillium crustosum*.Amorph. solid. [α]_D²⁵ -67.5 (c, 1.1 in CHCl₃). λ_{max} 231 (log ε 4.06); 285 (log ε 3.82) (MeOH).**Dechloro, 10-oxo, 11,33-dihydro: Penitremone C**

[164124-59-8]

C₃₇H₄₅NO₅ 583.766Metab. from *Penicillium* sp. (IBT 13163). Tremorgenic mycotoxin.The paper erroneously omits a double bond in the penitremones and shows them as 2,3-dihydroindoles. This causes registration and MF errors in CAS. λ_{max} 270 (MeOH) (Berdy).**Dechloro, 10-oxo, 11α,33-dihydro,****23α,24α-epoxyde: 11,33-Dihydro-10-oxo-penitrem B. Penitremone A**

[151434-31-0]

[164124-57-6 (Penitremone A)]


C₃₇H₄₅NO₆ 599.766Prod. by *Aspergillus sulphureus* and *Penicillium* sp. IB 13163.Tremorgenic agent. Light yellow solid. [α]_D -78.6. Identity of the two isolates based on uv and chromatographic props.**Dechloro, 10-oxo, 15-hydroxy, 11,33-dihydro,****23α,24α-epoxyde: Penitremone B**

[164124-58-7]

C₃₇H₄₅NO₇ 615.765From *Penicillium* sp. (IBT 13163).Tremorgenic mycotoxin. See note under Penitremone C above. λ_{max} 260; 288 (MeOH) (Berdy).Wilson, B.J. et al., *Nature (London)*, 1968, **220**, 77-78 (*Penitrem A, isol*)Norms, P.J. et al., *J. Neurochem.*, 1980, **34**, 33 (*Penitrem A, activity*)Fellows, P.N. et al., *Org. Mass Spectrom.*, 1981, **16**, 403-404 (*Penitrem A, Penitrem C, ms*)Mantle, P.G. et al., *Appl. Environ. Microbiol.*, 1983, **45**, 1486-1490 (*biosynth*)De Jesus, A.E. et al., *J.C.S. Perkin 1*, 1983, 1847-1856; 1857-1861; 1863 (*Penitrem, isol, biosynth, uv, ir, cd, pmr, cmr, abs config*)Hayashi, H. et al., *Chem. Express*, 1993, **8**, 233-236; *CA*, **118**, 251158g (6-*Bromopenitrem E*)Laakso, J.A. et al., *J. Agric. Food Chem.*, 1993, **41**, 973-975 (*Dihydrooxopenitrem B*)Naik, J.T. et al., *J.C.S. Perkin 1*, 1995, 1121-1125 (*Penitremones*)Gonzalez, M.C. et al., *J. Agric. Food Chem.*, 2003, **51**, 2156-2160 (*Penitrem G*)Smith, A.B. et al., *J.A.C.S.*, 2003, **125**, 8228-8237 (*Penitrem D, synth*)Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 382; 385Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, PAR250

Pennigritrem

[139682-30-7]


C₃₇H₄₄ClNO₆ 634.21

Prod. by *Penicillium nigricans*. Tremorgenic toxin; less active than Penitrem A in P-194. Amorph. solid. λ_{\max} 235 (€ 35700); 295 (€ 10300) (MeOH) (Berdy).

Penn, J. *et al.*, *J.C.S. Perkin 1*, 1992, 23 (*isol, pmr, cmr*)

Pensylpavoline

[53416-83-4]

C₃₉H₄₂N₂O₈ 666.769

Alkaloid from *Thalictrum polygamum* (Ranunculaceae). Cryst. (Et₂O). Mp 145-146°. $[\alpha]_D^{25}$ -245 (c, 0.66 in MeOH).

*Di-Ac:*Cryst. (Et₂O). Mp 188-189°.*O¹-Me: Pensylpavine*

[53416-82-3]

C₄₀H₄₄N₂O₈ 680.796


Alkaloid from *Thalictrum polygamum* (Ranunculaceae). Cryst. (Et₂O). Mp 122-123°. $[\alpha]_D^{25}$ -174 (c, 0.6 in MeOH).

*O¹-Me, Ac:*Cryst. (Et₂O). Mp 203-204°.

Shamma, M. *et al.*, *J.A.C.S.*, 1974, **96**, 3338 (*uv, pmr, ms, struct*)

Penochalasin C

[173792-72-8]



P-195

C₃₂H₃₅N₃O₃ 509.647

Prod. by a marine *Penicillium* sp. Mycotoxin. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 173-178°. $[\alpha]_D$ -6.2 (c, 0.1 in CHCl₃). λ_{\max} 208 (log € 4.5); 222 (log € 4.59); 284 (log € 3.92); 293 (log € 3.96); 315 (log € 3.87) (EtOH).

Δ^{5,6}-Isomer: Penochalasin B

[173792-71-7]

C₃₂H₃₅N₃O₃ 509.647

Prod. by a marine *Penicillium* sp. Mycotoxin. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 177-179°. $[\alpha]_D$ -6.2 (c, 0.2 in CHCl₃). λ_{\max} 206 (log € 4.1); 222 (log € 4.1); 284 (log € 4.08); 293 (log € 4.5); 312 (log € 4.45) (EtOH).

*7-Deoxy, 6,12-dihydro, 6β,7β-epoxide:***Penochalasin A**

[173792-70-6]

C₃₂H₃₅N₃O₃ 509.647

Prod. by a marine *Penicillium* sp. Mycotoxin. Needles (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 222-224°. $[\alpha]_D$ -10 (c, 0.2 in CHCl₃). λ_{\max} 206 (log € 4.58); 222 (log € 4.66); 285 (log € 4.4); 292 (log € 4.09); 313 (log € 4.13) (EtOH).

*7-Deoxy, 6,12,20β,21-tetrahydro, 6β,7β-epoxide: Penochalasin D*C₃₂H₃₇N₃O₃ 511.663

Prod. by a marine *Penicillium* sp. Cytotoxic. Oil. $[\alpha]_D$ +10.8 (c, 0.2 in CHCl₃). λ_{\max} 221 (log € 4.33); 242 (sh) (log € 3.72); 280 (log € 3.58); 292 (log € 3.48) (EtOH).

Numata, A. *et al.*, *J.C.S. Perkin 1*, 1996, 239-245 (*Penochalasin A-C*)


Iwamoto, C. *et al.*, *Tetrahedron*, 2001, **57**, 2997-3004 (*Penochalasin D*)

1,11,19,27,35-Pentaamino-3-tetracontanol, 9CI

P-198

Echin I

[187844-18-4]

C₄₀H₈₇N₅O 654.16

Prod. by *Erwinia chrysanthemi*. Antibiotic.

Liu, J. *et al.*, *CA*, 1997, **126**, 197207x (*isol, uv, ir, pmr, cmr, ms*)

4,8,12,16,20-Pentaazatricosane-1,23-diamine, 9CI

P-199

Caldoseptamine

[100009-88-9]

H₂N(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH₂

C₁₈H₄₅N₇ 359.6


Prod. by *Thermotoga maritima*.

[106395-90-8]

Zellner, G. *et al.*, *Arch. Microbiol.*, 1993, **159**, 473-476 (*isol*)

2,4,4',6,6'-Pentabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole

P-200

C₁₈H₁₁Br₅N₂S₂ 718.95

Alkaloid from *Laurencia brongniartii*. Plates (hexane/EtOAc). Mp 243-245°. λ_{\max} 231 (€ 83300); 302 (€ 28300) (MeOH).

*5-Bromo-2,4,4',5,6,6'-Hexabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole*C₁₈H₁₀Br₆N₂S₂ 797.846

Alkaloid from *Laurencia brongniartii*. λ_{\max} 233 (€ 51800); 303 (€ 16200) (MeOH).

*5'-Bromo-2,4,4',5',6,6'-Hexabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole*C₁₈H₁₀Br₆N₂S₂ 797.846

Alkaloid from *Laurencia brongniartii*. λ_{\max} 233 (€ 89400); 300 (€ 26700) (MeOH).


*5,5'-Dibromo-2,4,4',5,5',6,6'-Heptabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole*C₁₈H₉Br₇N₂S₂ 876.743

Alkaloid from *Laurencia brongniartii*. λ_{\max} 236 (€ 43700); 298 (€ 10200) (MeOH).

Kubota, N.K. *et al.*, *Heterocycles*, 2005, **65**, 2675-2682 (*isol, pmr, cmr, ms, cryst struct*)

3,3',4,4',5-Pentabromo-5'-chloro-2,2'-bi-1H-pyrrole

P-201

C₈H₂Br₅ClN₂ 561.09

N,N'-Di-Me: 3,3',4,4',5-Pentabromo-5'-chloro-1,1'-dimethyl-2,2'-bi-1H-pyrrole. **DBP-Br₅Cl**

[400767-00-2]

C₁₀H₆Br₅ClN₂ 589.144

Marine natural product of unknown biogenic origin. Present in marine or freshwater fish, shrimps, seabirds and seal blubber.

Tittlemier, S.A. *et al.*, *Environ. Pollut.*, 2002, **116**, 85-93 (*N,N'*-di-Me, occur, detn)

Tittlemier, S.A. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 2010-2015 (*N,N'*-di-Me, occur)

2,9-Pentadecadiene-12,14-diyonic acid

P-202

HC≡CC≡CCH₂CH=CH(CH₂)₅CH=CHCOOH

C₁₅H₁₈O₂ 230.306

(2E,9Z)-form

2-Methylpropylamide: **2,9-Pentadecadiene-12,14-diyonic acid isobutylamide**.

N-(2-Methylpropyl)-2,9-pentadecadiene-12,14-diyonamide, 9CI
[87797-75-9]

C₁₉H₂₇NO 285.428

Alkaloid from *Echinacea angustifolia* and *Echinacea purpurea*. Gum.

2-Hydroxy-2-methylpropylamide: **2,9-Pentadecadiene-12,14-diyonic acid (2-hydroxyisobutyl)amide**. N-(2-Hydroxy-2-methylpropyl)-2,9-pentadecadiene-12,14-diyonamide, 9CI

[87797-77-1]

C₁₉H₂₇NO₂ 301.428

Alkaloid from *Echinacea purpurea* (Asteraceae). Gum.

Bohlmann, F. et al., *Phytochemistry*, 1983, **22**, 1173-1175 (isol, ir, pmr, ms)

Bauer, R. et al., *Phytochemistry*, 1989, **28**, 505-508 (isol, pmr)

1-Pentadecylamine P-203

1-Pentadecylamine, 9CI. 1-Aminopentadecane

[2570-26-5]

H₃C(CH₂)₁₃CH₂NH₂

C₁₅H₃₃N 227.432

Cryst. Mp 36.5° (33.5°). Bp 298-301° Bp₁ 120.5°. pK_a 10.61 (25°).

▶ LD₅₀ (mus, orl) 520 mg/kg. RZ2120000

Hydrochloride: [1838-05-7]

Cryst. (EtOH). Mp 199°.

▶ RZ2134000

N-Ac:

C₁₇H₃₅NO 269.47

Cryst. (petrol). Mp 72°.

N-Benzyl, N-Me: N-Benzyl-N-methyl-1-pentadecylamine. **Piptamine**

C₂₃H₄₁N 331.584

Isol. from the mushroom *Piptoporus betulinus*. Active against various bacteria, yeasts and fungi. Waxy solid.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 454B (nmr)

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 167A (ir)

Naegeli, C. et al., *Helv. Chim. Acta*, 1929, **12**, 227 (synth)

Ralston, A.W. et al., *Ind. Eng. Chem.*, 1940, **32**, 1093

Murr, B.L. et al., *J.A.C.S.*, 1955, **77**, 1684 (synth)


Krivschich, S.Y. et al., *Toksikol. Nov. Prom. Khim. Veshchestv*, 1968, **10**, 124; *CA*, **71**, 47807 (tox)

Schlegel, B. et al., *J. Antibiot.*, 2000, **53**, 973-974 (Piptamine)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PBA000

4-Pentadecylpyridine P-204

[98771-51-8]



C₂₀H₃₅N 289.503


Isol. from plums infected with *Taphrina pruni*. Competitive polyphenoloxidase inhibitor. Bp_{0.01} 150°.

Hardy, D.R. et al., *Rapid Commun. Mass Spectrom.*, 1988, **2**, 16-18 (ms)

Fuchs, C.T. et al., *Z. Naturforsch. C*, 1995, **50**, 766-768; 1997, **52**, 761-765 (isol, synth, pmr)

1,2,3,4,5-Pentahydroxyacridone P-205

1,2,3,4,5-Pentahydroxy-9H-acridin-9-one



C₁₃H₉NO₆ 275.217

2,3,4-Tri-Me ether, N-Me: 1,5-Dihydroxy-2,3,4-trimethoxy-10-methylacridone. **Citbrasine**

[86680-34-4]

C₁₇H₁₇NO₆ 331.324

Alkaloid from the root bark of *Citrus sinensis* var. *brasiliensis* (navel orange) (Rutaceae). Red plates (Et₂O). Mp 154-156°.

Wu, T.-S. et al., *Chem. Pharm. Bull.*, 1983, **31**, 901 (isol, uv, ir, pmr, ms, struct)

1,2,3,5,6-Pentahydroxyacridone P-206

1,2,3,5,6-Pentahydroxy-9H-acridin-9-one

C₁₃H₉NO₆ 275.217

2,3-Di-Me ether, N-Me: **2,3-Dimethoxy-1,5,6-trihydroxy-10-methylacridone**

[126622-17-1]

C₁₆H₁₅NO₆ 317.298

Alkaloid from the root bark of *Pleiospermium alatum* (Rutaceae). Yellow cryst. Mp 118-119°.

2,5-Di-Me ether, N-Me: 1,3,6-Trihydroxy-2,5-dimethoxy-10-methylacridone. **Citramine**

[119459-67-5]

C₁₆H₁₅NO₆ 317.298

Alkaloid from roots of *Citrus natsuda-daidai* and roots of Ogonkan (a *Citrus* hybrid) (Rutaceae). Light-yellow prisms. Mp 277-279°.

2,3,5-Tri-Me ether, N-Me: **1,6-Dihydroxy-2,3,5-trimethoxy-10-methylacridone**

[101330-64-7]

C₁₇H₁₇NO₆ 331.324

Alkaloid from the stems of *Pleiospermium alatum* (Rutaceae). Yellow needles (CHCl₃/petrol). Mp 181-182°.

2,5,6-Tri-Me ether, N-Me: 1,3-Dihydroxy-2,5,6-trimethoxy-10-methylacridone. **Atalafoline**

[107259-49-4]

C₁₇H₁₇NO₆ 331.324

Alkaloid from the chinese drug Tung-Feng-Jie (roots of *Atalantia buxifolia* (Rutaceae)).

2,3,5,6-Tetra-Me ether: 1-Hydroxy-2,3,5,6-tetramethoxyacridone. **Cuspa-**

nine

[143438-94-2]

C₁₇H₁₇NO₆ 331.324

Alkaloid from leaves of *Angostura paniculata* (Rutaceae). Exhibits moderate molluscicidal activity and moderate cytotoxicity against several carcinoma cell lines. Yellow cryst. (MeOH). Mp 164-166°. λ_{max} 212 (ε 6950); 256 (ε 70000); 324 (ε 4000); 384 (ε 6000) (MeOH) (Berdy).

2,3,5,6-Tetra-Me ether, N-Me: **1-Hydroxy-2,3,5,6-tetramethoxy-10-methylacridone**

[101330-63-6]

C₁₈H₁₉NO₆ 345.351

Alkaloid from the root bark of *Pleiospermium alatum* (Rutaceae). Yellow needles (CHCl₃/petrol); cryst. (hexane/EtOAc). Mp 198-200° (170°).

1,2,3,5,6-Penta-Me ether: 1,2,3,5,6-Pentamethoxyacridone. **Cusculine**

[143438-95-3]

C₁₈H₁₉NO₆ 345.351

Alkaloid from leaves of *Atalantia paniculata* (Rutaceae). Exhibits moderate molluscicidal activity and moderate cytotoxicity against several carcinoma cell lines. Yellow cryst. (MeOH). Mp 223-224°. λ_{max} 220 (ε 7400); 264 (ε 74000); 312 (ε 3800); 372 (ε 6350) (MeOH) (Berdy).

Bowen, I.H. et al., *Phytochemistry*, 1986, **25**, 429 (isol, uv, ir, pmr, ms, struct)

Qin, D. et al., *Yaoxue Xuebao*, 1986, **21**, 683; *CA*, **106**, 135228h (Atalafoline)

Ju-Ichi, M. et al., *Heterocycles*, 1988, **27**, 2197 (Citramine)

Bandara, B.M.R. et al., *Phytochemistry*, 1990, **29**, 297 (isol, uv, ir, pmr, ms, struct)

Vieira, P.C. et al., *J. Nat. Prod.*, 1992, **55**, 1112 (Cuspanine, Cusculine)

1,3,4,5,6-Pentahydroxyacridone P-207

C₁₃H₉NO₆ 275.217

3,6-Di-Me ether, N-Me: 1,4,5-Trihydroxy-3,6-dimethoxy-10-methylacridone. **Atalafoline B**

[114216-93-2]

C₁₆H₁₅NO₆ 317.298

Alkaloid from the roots of *Atalantia buxifolia* (Rutaceae).

4,5-Di-Me ether, N-Me: 1,3,6-Trihydroxy-4,5-dimethoxy-10-methylacridone. **Buxifolidine H**

[263007-72-3]

C₁₆H₁₅NO₆ 317.298

Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 215-217°. λ_{max} 221 (log ε 2.94); 234 (log ε 2.6); 260 (log ε 3.45); 269 (sh) (log ε 3.41); 313 (log ε 2.67); 332 (log ε 2.93); 390 (log ε 2.53) (MeOH).

3,4,5-Tri-Me ether, N-Me: 1,6-Dihydroxy-3,4,5-trimethoxy-10-methylacridone. **Glycofolinine**

[172335-06-7]


C₁₇H₁₇NO₆ 331.324

Alkaloid from root bark and stem bark of *Glycosmis citrifolia* (Rutaceae). Yellow prisms. Mp 161-163°.

- Gu, G. *et al.*, *Yaouxue Xuebao*, 1987, **22**, 886-888; *CA*, **108**, 183612s
 Ono, T. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1629-1631 (*Glycofolinine*)
 Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*Buxifoliadine H*)

1,2,3,9,10-Pentahydroxyaporphine P-208

5,6,6a,7-Tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-1,2,3,9,10-pentol, 9CI



$C_{17}H_{17}NO_5$ 315.325

(S)-form

- 1,2,10-Tri-Me ether: 3,9-Dihydroxy-1,2,10-trimethoxyaporphine. Delporphine**
 [67253-86-5]
 $C_{20}H_{23}NO_5$ 357.405
 Alkaloid from *Delphinium dictyocarpum*. Mp 116-117°. $[\alpha]_D^{25} +68$ (c, 0.25 in MeOH).
- 2,3,10-Tri-Me ether, N-de-Me: 1,9-Dihydroxy-2,3,10-trimethoxynoraporphine. Nordelporphine**
 [109175-40-8]
 $C_{19}H_{21}NO_5$ 343.379
 Alkaloid from the leaves of *Phoebe valeriana* (Lauraceae). Amorph. solid. Opt. rotn. not recorded.
- 1,2,3,10-Tetra-Me ether: 9-Hydroxy-1,2,3,10-tetramethoxyaporphine. Thaliosopynine. Thaliosopinine**
 [24314-86-1]
 $C_{21}H_{25}NO_5$ 371.432
 Alkaloid from the roots *Thalictrum isopyroides* (Ranunculaceae). $[\alpha]_D^{25} +45$ (c, 0.13 in MeOH).
- 1,2,9,10-Tetra-Me ether: 3-Hydroxy-1,2,9,10-tetramethoxyaporphine. Thalbaicalidine. O-Demethylpurpureine. 3-Hydroxyglauoine**
 [66186-95-6]
 $C_{21}H_{25}NO_5$ 371.432
 Alkaloid from the stems and leaves of *Annona purpurea* (?), from *Ocotea bucherii* leaves and twigs and from *Thalictrum baicalense* (Annonaceae, Lauraceae, Ranunculaceae). Cryst. (Et₂O or MeOH aq.). Mp 193-195°. $[\alpha]_D^{25} +74.1$ (c, 0.6 in MeOH). Identity of 3-Hydroxyglauoine from *O. bucherii* and Thalbaicalidine with O-Demethylpurpureine from *A. purpurea*, which was not fully characterised, is not certain.
- 1,2,9,10-Tetra-Me ether, N-de-Me: 3-Hydroxy-1,2,9,10-tetramethoxynoraporphine. Thalbaicaline**
 [88187-19-3]

- $C_{20}H_{23}NO_5$ 357.405
 Alkaloid from *Thalictrum baicalense* (Ranunculaceae). Cryst. (EtOAc). $[\alpha]_D^{25} +61$ (MeOH).
- 2,3,9,10-Tetra-Me ether: 1-Hydroxy-2,3,9,10-tetramethoxyaporphine. Preocoteine**
 [18161-86-9]
 $C_{21}H_{25}NO_5$ 371.432
 Alkaloid from *Thalictrum fendleri* (whole plant) and the roots of *Thalictrum strictum*, also present in the bark of *Phoebe mollicella* (Ranunculaceae, Lauraceae). Oil. $[\alpha]_D^{25} +26$ (EtOH).
- 2,3,9,10-Tetra-Me ether, N-oxide: Preocoteine N-oxide**
 [41607-14-1]
 $C_{21}H_{25}NO_6$ 387.432
 Alkaloid from the roots of *Thalictrum minus* (Ranunculaceae). Mp 199-200° dec.
- 2,3,9,10-Tetra-Me ether, N-de-Me: 1-Hydroxy-2,3,9,10-tetramethoxynoraporphine. Norpreocoteine**
 [89412-82-8]
 $C_{20}H_{23}NO_5$ 357.405
 Alkaloid from the bark of *Phoebe mollicella* (Lauraceae). Brownish oil.
- Penta-Me ether: 1,2,3,9,10-Pentamethoxyaporphine. Thalicsimidine. Purpureine†. O-Methylpreocoteine**
 [19775-47-4]
 $C_{22}H_{27}NO_5$ 385.459
 Alkaloid from the roots of *Thalictrum simplex* and *Thalictrum strictum*. Also present in *Thalictrum filamentosum* and in the stems and leaves of *Annona purpurea* and the bark of *Phoebe mollicella* (Ranunculaceae, Annonaceae, Lauraceae). Cryst. (Me₂CO). Mp 131-132°. $[\alpha]_D^{25} +66.9$ (c, 1.42 in EtOH).
- Penta-Me ether, picrate**: Mp 141° dec. (131-132°).
- Penta-Me ether, N-oxide: Thalicsimidine N-oxide**
 [177413-36-4]
 $C_{22}H_{27}NO_6$ 401.458
 Alkaloid from aerial parts of *Thalictrum simplex*. $[\alpha]_D^{25} +46$ (c, 0.02 in MeOH).
- Penta-Me ether, N-Me: N-Methylthalicsimidine. N-Methylpurpureine**
 Quaternary alkaloid from the bark and roots of *Xylopija parviflora*. Mp 227-229° (as iodide) Mp 187.5-190° (as perchlorate). $[\alpha]_D^{25} +25.1$ (0.5 in MeOH) (perchlorate). λ_{max} 281 (log ϵ 4.01); 303 (log ϵ 4) (MeOH).
- Penta-Me ether, N-de-Me: 1,2,3,9,10-Pentamethoxynoraporphine. Norpurpureine**
 [34272-09-8]
 $C_{21}H_{25}NO_5$ 371.432
 Alkaloid from the stems and leaves of *Annona purpurea* and the bark of *Phoebe mollicella* (Annonaceae, Lauraceae). Mp 115-117°. $[\alpha]_D^{25} -70.4$ (c, 0.2 in CHCl₃).
- Penta-Me ether, N-de-Me, N-formyl: N-Formylnorpurpureine. N-Formylpurpureine (incorr.)**
 $C_{22}H_{25}NO_6$ 399.443

Alkaloid from the leaves of *Annona purpurea*. Brown amorph. powder. Mp 212-213°. $[\alpha]_D^{25} -135$ (c, 0.2 in CHCl₃). Exists as a mixt. of E- and Z-hydroximine zwitterions. λ_{max} 225; 268; 290; 316 (EtOH).

- Penta-Me ether, N-de-Me, N-methoxycarbonyl: Romucosine G**
 $C_{23}H_{27}NO_7$ 429.469
 Alkaloid from *Annona purpurea*. Needles. Mp 135-136°. $[\alpha]_D^{25} +44$ (c, 0.08 in CHCl₃). λ_{max} 233 (log ϵ 4.33); 282 (log ϵ 4.04); 303 (log ϵ 3.54) (EtOH).
- 6a,7-Didehydro, penta-Me ether: Dehydrothalicsimidine**
 [125236-63-7]
 $C_{22}H_{25}NO_5$ 383.443
 Alkaloid from *Thalictrum inchengense* (Ranunculaceae).
- 7-Oxo, penta-Me ether: 7-Hydroxydehydrothalicsimidine**
 $C_{22}H_{25}NO_6$ 399.443
 Alkaloid from *Annona purpurea*. Platelet aggregation inhibitor. Amorph. green powder. Mp 223-224°. Exists in the enol (7-hydroxy-6a,7-didehydro) form. λ_{max} 215; 256; 270 (sh); 340 (EtOH).
- 1,2-Methylene, 3,9,10-tri-Me ether**: see Ocoteine, O-27


(±)-form

- Penta-Me ether**:
 Synthetic. Cryst. + ¼ H₂O or yellowish-brown viscous syrup. Mp 121-122°.
- Penta-Me ether, perchlorate**:
 Prisms (MeOH/Et₂O). Mp 220-225°.
- Shamma, M. *et al.*, *Tetrahedron*, 1967, **23**, 2887 (*Preocoteine*)
 Shamma, M. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 262 (*Preocoteine*)
 Ismailov, Z.F. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 136; 196; 202; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 117; 169; 175 (*struct. ms. pnr*)
 Umarov, K.S. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 224; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 219 (*isol*)
 Sonnet, P.E. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1254 (*isol, uv, pnr, derivs*)
 Kametani, T. *et al.*, *J.C.S. (C)*, 1971, 1032; 3617 (*synth, uv, pnr*)
 Doskotch, R.W. *et al.*, *J.O.C.*, 1971, **36**, 2409 (*synth, uv, pnr*)
 Khozdaev, V.G. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 631; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 599 (*Preocoteine N-oxide*)
 Maekh, S.K. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 560; 1983, **19**, 537; 1986, **22**, 251; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 238; 1983, **19**, 507; 1986, **22**, 511 (*Thalbaicaline, Thalbaicalidine*)
 Umarova, D. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 788; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 706 (*isol*)
 Mollov, N. *et al.*, *Chem. Ber.*, 1978, **111**, 554 (*synth, uv, pnr*)
 Salimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 235; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 194 (*Delporphine*)
 Abdizhabbarova, S. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 472; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 400 (*Thaliosopynine*)
 Rönsch, H. *et al.*, *Annalen*, 1983, 744 (3-Hydroxyglauoine)
 Stermitz, F.R. *et al.*, *J. Nat. Prod.*, 1983, **46**, 913 (*Norpurpureine, Norpreocoteine, cmr*)

- Hara, H. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1946 (*synth*)
- Castro, O. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1036 (*Nordelporphine*)
- Wu, Z. *et al.*, *Shoyakugaku Zasshi*, 1989, **43**, 195; *CA*, **112**, 73781w (*Dehydrothalicisimidine*)
- Velcheva, M.P. *et al.*, *Phytochemistry*, 1996, **42**, 535 (*oxide*)
- Chang, F.R. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1457-1461 (*N-Formylnorpurpureine*)
- Chang, F.-R. *et al.*, *Phytochemistry*, 1998, **49**, 2015-2018 (*7-Hydroxydehydrothalicisimidine*)
- Chang, F.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 746-748 (*Romucosine G*)
- Nishiyama, Y. *et al.*, *Phytochemistry*, 2004, **65**, 939-944 (*N-Methylthalicisimidine*)

1,2,3,10,11-Pentahydroxyaporphine P-209

5,6,6a,7-Tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-1,2,3,10,11-pentol, 9CI



(S)-form

C₁₇H₁₇NO₅ 315.325

(S)-form

- 1,2,10-Tri-Me ether: 3,11-Dihydroxy-1,2,10-trimethoxyaporphine. **N-Methyl-danguyelline** [122297-36-3] C₂₀H₂₃NO₅ 357.405 Alkaloid from the whole plant of *Thalictrum pedunculatum* (Ranunculaceae). [α]_D²⁰ +96 (c, 0.25 in CHCl₃).
- 1,2,10-Tri-Me ether, N-de-Me: 3,11-Dihydroxy-1,2,10-trimethoxynoraporphine. **Danguyelline** [80151-80-0] C₁₉H₂₁NO₅ 343.379 Alkaloid from the trunk bark of *Xylopia danguyella* (Annonaceae). Cryst. (MeOH). Mp 190°.
- 1,2,3,10-Tetra-Me ether: 11-Hydroxy-1,2,3,10-tetramethoxyaporphine. **Oconovine** [19893-97-1] C₂₁H₂₅NO₅ 371.432 Alkaloid from an unidentified *Ocotea* sp. and from *Thalictrum urbaini* (Lauraceae, Ranunculaceae). Amorph. [α]_D²⁷ +156 (c, 0.2 in CHCl₃).
- 1,2,3,10-Tetra-Me ether, N-Me: Mp 204-205° (as iodide). [α]_D²⁷ +70 (c, 0.2 in CHCl₃) (iodide).
- 1,2,3,10-Tetra-Me ether, N-de-Me: 11-Hydroxy-1,2,3,10-tetramethoxynoraporphine. **Noroconovine** [69477-65-2] C₂₀H₂₃NO₅ 357.405 Alkaloid from the trunk bark of *Polyalthia oligosperma* (Annonaceae). λ_{max} 221 ; 274 ; 307 (sh) (EtOH).
- 2,3,10,11-Tetra-Me ether: 1-Hydroxy-

2,3,10,11-tetramethoxyaporphine. **Isoconovine**

[72170-11-7]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the leaves of *Ocotea minarum* (Lauraceae). Mp 98-99°. [α]_D²⁰ +155 (c, 1.0 in MeOH).

1,2-Methylene, 3,10-di-Me ether, N-de-Me: 11-Hydroxy-3,10-dimethoxy-1,2-methylenedioxyaporphine. **Cassyformine**

[208252-24-8]

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Cassytha filiformis*. Amorph. yellow powder (CHCl₃). Mp 135-137°. [α]_D²⁵ +30 (c, 0.2 in CHCl₃). λ_{max} 215 (log ε 4.03); 284 (log ε 4.1); 302 (log ε 3.77) (MeOH).

1,2-Methylene, 3,11-di-Me ether, N-de-Me: 10-Hydroxy-3,11-dimethoxy-1,2-methylenedioxyaporphine.

Hernandine[†]

[15583-41-2]

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Hernandia bivalvis*. Mp 240-241°. [α]_D²⁷ +347.

1,2-Methylene, 3,10,11-tri-Me ether, N-de-Me: 3,10,11-Trimethoxy-1,2-methylenedioxyaporphine. **O-Methylcassyformine**

[14050-90-9]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Cassytha filiformis*. λ_{max} 226 ; 277 (MeOH).

(±)-form

1,2,3,10-Tetra-Me ether: [23740-41-2]

Synthetic.

1,2,3,10-Tetra-Me ether, N-Me: Mp 228° (as iodide).

1,2-Methylene, 3,11-di-Me ether, N-de-Me: Synthetic. Mp 227-228° dec.

Soh, K.S. *et al.*, *Tet. Lett.*, 1966, **7**, 5279-5283; 1968, **9**, 19-22 (*Hernandine*, *pmr*, *struct*, *synth*)

Cava, M.P. *et al.*, *J.O.C.*, 1970, **35**, 1867-1869 (*Oconovine*)

Guinaudeau, H. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 166-172 (*Noroconovine*)

Vecchietti, V. *et al.*, *Farmaco, Ed. Sci.*, 1979, **34**, 829-840 (*Isoconovine*)


Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551-556 (*Danguyelline*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1989, **52**, 428-429 (*Danguyelline*, *N-Methyl-danguyelline*)

Chang, F.-R. *et al.*, *J. Nat. Prod.*, 1998, **61**, 863-866 (*Cassyformine*)

Tsai, T.-H. *et al.*, *J. Nat. Prod.*, 2008, **71**, 289-291 (*O-Methylcassyformine*)

1,2,8,9,10-Pentahydroxyaporphine P-210



1480

C₁₇H₁₇NO₅ 315.325

(S)-form

1,8,9,10-Tetra-Me ether: 2-Hydroxy-1,8,9,10-tetramethoxyaporphine.

Acutifolidine[†]

[126595-93-5]

C₂₁H₂₅NO₅ 371.432

Alkaloid from roots of *Thalictrum acutifolium*. Mp 183-184°. [α]_D²⁰ +77 (c, 1 in MeOH).

1,2:8,9-Bis(methylene), 10-Me ether: 10-Methoxy-1,2:8,9-bis(methylenedioxy)aporphine. **Ocominarine**

[72170-09-3]

C₂₀H₁₉NO₅ 353.374


Alkaloid from the leaves of *Ocotea minarum* (Lauraceae). Mp 133-135°. [α]_D²⁰ +65 (c, 0.5 in CHCl₃).

Vecchietti, V. *et al.*, *Farmaco, Ed. Sci.*, 1979, **34**, 829 (*Ocominarine*, *isol*, *uv*, *pmr*, *ms*, *struct*)

Lin, C. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 449; *CA*, **112**, 175576a (*Acutifolidine*, *isol*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1033 (*Acutifolidine*, *rev*, *uv*, *ir*, *pmr*, *ms*)

1,2,9,10,11-Pentahydroxyaporphine P-211



C₁₇H₁₇NO₅ 315.325

(S)-form


1,2:9,10-Bis(methylene) ether, N-de-Me: 11-Hydroxy-1,2:9,10-bis(methylenedioxy)aporphine. **Lindechunine B**


C₁₈H₁₅NO₅ 325.32

Alkaloid from the roots of *Lindera chinii*. Greyish powder. [α]_D²⁶ +43 (c, 0.01 in MeOH). λ_{max} 222 ; 280 (MeOH).


Zhang, C.-F. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1195-1200 (*Lindechunine B*)

3,7,14,16,20-Pentahydroxycevan-6-one, 9CI P-212





C₂₇H₄₃NO₆ 477.64**(3β,5α,7α,16β)-form***Pingpeimine C. Pengbeimine C*
[128585-96-6]Alkaloid from the bulbs of *Fritillaria ussuriensis* (Liliaceae). Mp 171.5-173°. [α]_D²⁵ -24.6 (c, 1.1 in MeOH).Xu, D.M. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 127-130; *CA*, **113**, 74794 (*isol, struct*)**3,3',4,4',5-Pentahydroxy-2,7'-cyclo lign-7-ene-9,9'-dioic acid** P-213C₁₈H₁₄O₉ 374.303**(7'R,8'S)-form**

3,5-Di-Me ether, bis[2-(4-hydroxyphenyl)ethylamide]: [200135-06-4]

C₃₆H₃₆N₂O₉ 640.688Alkaloid from *Porcelia macrocarpa*. Amorph. powder. [α]_D²⁵ -12 (c, 0.08 in MeOH). λ_{max} 203 (log ε 4.85); 249 (log ε 4.49); 283 (log ε 4.22); 332 (log ε 4.24) (EtOH).Chaves, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 879-881 (*isol, uv, pmr, cmr*)**1,2,6,7,9-Pentahydroxy-dibenz[cd,f]indol-4(5H)-one, 9CI** P-214C₁₅H₉NO₆ 299.239


CAS numbering shown. Alternative (phenanthrene) numbering freq. encountered in which 1,2,6,7,9- are 4,3,9,8,6 respectively.

1,2-Methylene, 6,7,9-tri-Me ether: **9-Methoxyaristolactam IV**
[133462-31-4]C₁₉H₁₅NO₆ 353.331Alkaloid from *Asarum petilotii* and *Aristolochia auricularia*. Pale yellow solid.Houghton, P.J. *et al.*, *Phytochemistry*, 1991, **30**, 253Ding, Z. *et al.*, *Yunnan Zhiwu Yanjiu*, 1994, **16**, 305; *CA*, **122**, 101604t**1,2,4,6,9-Pentahydroxy-dihydro-β-agarofuran** P-215
2,4-DihydroxycyclobicolC₁₅H₂₆O₆ 302.367**(1α,2β,4β,6β,9β)-form**6-(3-Pyridinecarbonyl), 1-cinnamoyl, 9-Ac: **Rzedowskin D**
[117677-26-6]C₃₂H₃₇NO₉ 579.646Alkaloid from the aerial parts of *Orthosphenia mexicana* and *Rzedowskia tolantonguensis* (Celastraceae). Amorph. solid. Mp 143-145°.Jiménez, M. *et al.*, *Phytochemistry*, 1988, **27**, 2213 (*Rzedowskins*)**1,2,6,8,9-Pentahydroxy-dihydro-β-agarofuran** P-216

(1α,2α,6β,8α,9α)-form

C₁₅H₂₆O₆ 302.367**(1α,2α,6β,8α,9α)-form**1-(3-Pyridinecarbonyl), 2,9-dibenzoyl, 8-(2-methylbutanoyl), 6-Ac: **Triptogelin A5**
[135118-41-1]C₄₂H₄₇NO₁₁ 741.833Constit. of *Tripterygium wilfordii*. Amorph. powder. [α]_D²⁵ +200.1 (c, 0.17 in CHCl₃).1-(3-Pyridinecarbonyl), 2,8,9-tribenzoyl, 6-Ac: **Triptogelin A10**
[135118-33-1]C₄₄H₄₃NO₁₁ 761.824Constit. of *Tripterygium wilfordii* var. *regelii*. Amorph. powder. [α]_D²⁵ +77.6 (c, 0.52 in MeOH).8-(3-Pyridinecarbonyl), 9-benzoyl, 6-Ac: **Triptogelin A7**
[135118-43-3]C₃₀H₃₅NO₉ 553.608Constit. of *Tripterygium wilfordii*. Amorph. powder. [α]_D²³ +63.1 (c, 0.5 in MeOH).8-(3-Pyridinecarbonyl), 1,9-dibenzoyl, 2-hexanoyl, 6-Ac: **Triptogelin A9**
[135118-45-5]C₄₃H₄₉NO₁₁ 755.86Constit. of *Tripterygium wilfordii*. Amorph. powder. [α]_D²³ +12.1 (c, 1 in MeOH).8-(3-Pyridinecarbonyl), 1,2,9-tribenzoyl, 6-Ac: **Triptogelin A6**
[135118-42-2]C₄₄H₄₃NO₁₁ 761.824Constit. of *Tripterygium wilfordii*.Amorph. powder. [α]_D²⁵ +31.5 (c, 0.21 in CHCl₃).**(1α,2α,6β,8β,9β)-form**

2-(3-Pyridinecarbonyl), 1,9-dibenzoyl, 6,8-di-Ac: [389865-20-7]

C₃₉H₄₁NO₁₁ 699.753Constit. of *Maytenus magellanica*.Amorph. solid. [α]_D²⁵ +31 (c, 0.19 in CHCl₃). λ_{max} 229; 264 (EtOH).Takaishi, Y. *et al.*, *Phytochemistry*, 1990, **29**, 3869-3873; 1991, **30**, 1561-1566; 1567-1572 (*Triptogelins*)Kennedy, M.L. *et al.*, *J. Med. Chem.*, 2001, **44**, 4668-4676 (*Maytenus magellanica constit*)**1,2,6,9,14-Pentahydroxy-dihydro-β-agarofuran** P-217

(1α,2α,6β,9α)-form

C₁₅H₂₆O₆ 302.367**(1α,2α,6β,9β)-form**

3,4-Dihydroxymaytol

Amorph. solid. [α]_D²³ -2.5 (c, 0.12 in MeOH).

2-(3-Pyridinecarbonyl), 6,9-dibenzoyl, 1,14-di-Ac: [389865-23-0]


C₃₉H₄₁NO₁₁ 699.753Constit. of *Maytenus chubutensis*.Amorph. solid. [α]_D²⁵ +70 (c, 0.39 in CHCl₃). λ_{max} 229; 264 (EtOH).

9-(3-Pyridinecarbonyl), 1,2,6,14-tetra-Ac:

C₂₉H₃₇NO₁₁ 575.611Constit. of *Euonymus nanus* and *Euonymus sachalinensis*. Amorph. solid.9-(3-Pyridinecarbonyl), 1-O-(2S-methylbutanoyl), 2,6,14-tri-Ac: **Ever I**C₃₂H₄₃NO₁₁ 617.692Alkaloid from *Euonymus verrucosus* (Celastraceae). Cryst. (MeOH aq.). Mp 189-192°.9-(3-Pyridinecarbonyl), 2-benzoyl, 1,6-di-Ac: **Maymyrsine**
[93772-06-6]C₃₂H₃₇NO₁₀ 595.645Alkaloid from the fruits of *Maytenus myrsinoides* (Celastraceae). Prisms (EtOH). Mp 185-187°. [α]_D +93 (c, 1 in CHCl₃).9-(3-Pyridinecarbonyl), 2-benzoyl, 1,6,14-tri-Ac: **Acetylmaymyrsine**
[93772-07-7]C₃₄H₃₉NO₁₁ 637.682Alkaloid from the fruits of *Maytenus myrsinoides* (Celastraceae). Amorph. solid. [α]_D²⁰ +92 (c, 1 in CHCl₃).

- Baudouin, G. et al., *Heterocycles*, 1984, **22**, 2221-2226 (*Maymyrsine, Acetylmyrsine, cryst struct*)
 Begley, M.J. et al., *J.C.S. Perkin 1*, 1986, 535-539 (*Ever 1, cryst struct*)
 González, A.G. et al., *J. Nat. Prod.*, 1990, **53**, 474-478 (*Maytenus chubutensis constits*)
 Hohmann, J. et al., *J. Nat. Prod.*, 1995, **58**, 1192-1199 (*Euonymus sachalinensis constits*)
 Kennedy, M.L. et al., *J. Med. Chem.*, 2001, **44**, 4668-4676 (*Maytenus chubutensis constits*)

1,2,8,9,14-Pentahydroxydihydro-β-agarofuran P-218



(1α,2α,8α,9β)-form

C₁₅H₂₆O₆ 302.367

(1α,2α,8α,9β)-form

14-(3-Pyridinecarbonyl), 9-benzoyl, 1,2-di-Ac: **Cathedulin E8**. *Cathedulin Y8* [61166-34-5]
 C₃₂H₃₇NO₁₀ 595.645
 Alkaloid from *Catha edulis* (Celastraceae). Gum. May be an artifact of isol. since the 4-nicotinoyl group of *Cathedulin E2* is labile.

8,14-Bis(3-pyridinecarbonyl), 9-benzoyl, 1,2-di-Ac: **Cathedulin E2**. O⁴-Nicotinoylcathedulin E8. *Cathedulin 2* [61231-06-9]
 C₃₈H₄₀N₂O₁₁ 700.741
 Alkaloid from *Catha edulis* (Celastraceae). Cryst. (Et₂O/petrol). Mp 149-151° (transition at 137-140°). [α]_D²⁵ -74 (c, 0.22 in CHCl₃).

(1α,2α,8β,9α)-form


1-Benzoyl, 2,8,9,14-tetra-Ac: **Kupitengester 3** [149252-08-4]
 C₃₀H₃₈O₁₁ 574.624
 Constit. of *Celastrus angulatus*.
 9-Benzoyl, 1,2,8,14-tetra-Ac: **Angulatueoid B** [145042-01-9]
 C₃₀H₃₈O₁₁ 574.624
 Constit. of *Celastrus angulatus*. Cryst. (MeOH). Mp 219-219.5° (214 - 215°). [α]_D²³ +59 (c, 0.505 in CHCl₃).

14-(2-Pyridinecarbonyl), 9-benzoyl, 1,2,8-tri-Ac: **Angulatueoid A**. *Angulatusine A* [144425-15-0]
 C₃₄H₃₉NO₁₁ 637.682
 Constit. of *Celastrus angulatus* (Celastraceae). Mp 227-230°. Unusual 2-pyridinecarbonyl residue, struct. proved by x-ray crystallography.

14-(3-Pyridinecarbonyl), 9-benzoyl, 1,2,8-tri-Ac: [154741-50-1]
 C₃₄H₃₉NO₁₁ 637.682
 Constit. of *Celastrus angulatus*. Cryst. (Me₂CO/petrol). Mp 213-214°. λ_{max} 218 (log ε 5.04); 229 (log ε 5.04); 264 (log ε 3.68) (MeOH).

- 9,14-Bis(3-pyridinecarbonyl), 1,2,8-tri-Ac: [147167-66-6]
 C₃₃H₃₈N₂O₁₁ 638.67
 Constit. of *Celastrus angulatus*. Cryst. (MeOH). Mp 183-184°. [α]_D²³ +64.8 (c, 0.515 in CHCl₃).
 Baxter, R.L. et al., *J.C.S. Perkin 1*, 1979, 2965-2970; 2972-2975 (*Cathedulins*)
 Wu, D.G. et al., *J. Nat. Prod.*, 1992, **55**, 982-985 (*Angulatusine A, cryst struct*)
 Chunquan, C. et al., *Phytochemistry*, 1992, **31**, 2777-2780 (*Angulatueoids, cryst struct*)
 Wang, G. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1992, **34**, 777-780 (*Kupitengester 3*)
 Tu, Y. et al., *Chin. Chem. Lett.*, 1993, **4**, 219 (*deriv, cryst struct*)
 Yongqing, T. et al., *Phytochemistry*, 1993, **32**, 458-459 (*Celastrus angulatus constits*)

1,4,6,8,9-Pentahydroxydihydro-β-agarofuran P-219



(1α,6β,8α,9α)-form

C₁₅H₂₆O₆ 302.367

(1α,6β,8β,9α)-form

6-O-(3-Pyridinecarbonyl), 1-benzoyl, 9-Ac: **9-Acetoxy-1-benzoyloxy-4,8-dihydroxy-6-nicotinoyloxydihydro-β-agarofuran** [110559-19-8]
 C₃₀H₃₅NO₉ 553.608
 Alkaloid from the aerial parts of *Orthosphenia mexicana* (Celastraceae). Mp 184-186°.

6-O-(3-Pyridinecarbonyl), 1-benzoyl, 8,9-di-Ac: **8,9-Diacetoxy-1-benzoyloxy-4-hydroxy-6-nicotinoyloxydihydro-β-agarofuran** [110559-18-7]
 C₃₂H₃₇NO₁₀ 595.645
 Alkaloid from the root bark of *Rzedowskia tolantonguensis* (Celastraceae). Mp 114-116°.

6-O-(3-Pyridinecarbonyl), 1,9-dibenzoyl, 4,8-di-Ac: [110559-18-7]
 C₃₉H₄₁NO₁₁ 699.753
 Constit. of *Rzedowskia tolantonguensis*.

(1α,6β,8β,9β)-form

6-O-(3-Pyridinecarbonyl), 1-benzoyl, 9-Ac: [110559-18-7]
 C₃₀H₃₅NO₉ 553.608
 Alkaloid from the aerial parts of *Orthosphenia mexicana* and the root bark of *Rzedowskia tolantonguensis* (Celastraceae). Mp 139-148°.

6-O-(1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 9-benzoyl, 8-Ac: **Reissantin G** [910651-21-7]
 C₃₁H₃₇NO₁₀ 583.634
 Constit. of *Reissantia buchananii*. Powder. Mp 168-169°. [α]_D²⁴ +33.6 (c, 0.55 in CHCl₃). λ_{max} 277 (log ε 3.66) (MeOH).

6-O-(1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 9-benzoyl, 1,8-di-Ac: **Reissantin A** [628290-35-7]
 C₃₃H₃₉NO₁₁ 625.671
 Constit. of *Reissantia buchananii*. Amorph. powder. Mp 218-220°. [α]_D²⁵ +42.44 (c, 0.69 in CHCl₃). λ_{max} 211 (log ε 4.62); 234 (log ε 4.2); 263 (log ε 4.37); 300 (log ε 3.72) (MeOH).


8-O-(1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 1,9-bis(3-furancarboxonyl), 6-Ac: **Reissantin C** [628290-37-9]
 C₃₄H₃₇NO₁₃ 667.665
 Constit. of *Reissantia buchananii*. Wax. [α]_D²⁵ +36.4 (c, 0.37 in CHCl₃). λ_{max} 200 (log ε 3.94); 221 (log ε 3.91); 267 (log ε 4.13) (MeOH).

1-Ketone: 4,6,8,9-Tetrahydroxydihydro-β-agarofuran-1-one
 C₁₅H₂₄O₆ 300.351

1-Ketone, 8-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 9-O-(2-hydroxybenzoyl), 6-benzoyl: **Reissantin H** [910651-22-8]
 C₃₆H₃₇NO₁₁ 659.688
 Constit. of *Reissantia buchananii*. Oil. [α]_D²¹ -50.1 (c, 0.36 in MeOH). λ_{max} 268 (log ε 3.52) (MeOH).

González, A.G. et al., *Phytochemistry*, 1997, **46**, 309-311 (*Rzedowskia tolantonguensis ester*)
 Chang, F.-R. et al., *J. Nat. Prod.*, 2003, **66**, 1416-1420 (*Reissantins*)
 Chang, F.-R. et al., *Planta Med.*, 2006, **72**, 92-96 (*Reissantins G,H*)

1,6,8,9,14-Pentahydroxydihydro-β-agarofuran P-220



(1α,6β,8α,9α)-form

C₁₅H₂₆O₆ 302.367

(1α,6β,8α,9α)-form

3-Pyridinecarbonyl, benzoyl, tri-Ac: **Ejap 13**
 C₃₄H₃₉NO₁₁ 637.682
 Isol. from *Euonymus japonicus*. Amorph. Posn. of ester groups not detd.

(1α,6β,8α,9β)-form

9-(3-Pyridinecarbonyl), 1,6,8,14-tetra-Ac: **Ejap 14** [122475-51-8]
 C₂₉H₃₇NO₁₁ 575.611
 Isol. from *Euonymus japonicus*. Amorph.

(1α,6β,8β,9α)-form

14-(3-Pyridinecarbonyl), 6-benzoyl, tri-Ac: **Kupitengester 1**


[135063-89-7]
C₃₄H₃₉NO₁₁ 637.682

Constit. of *Euonymus kuptiensis*.

Rózsa, Z. et al., *J.C.S. Perkin 1*, 1989, 1079-1087 (*Ejaps*)

Wang, G. et al., *Chin. Sci. Bull.*, 1991, **36**, 206; *CA*, **115**, 71924h (*Kuptiengester 1*)

1,2,10,11,12-Pentahydroxyhomoaporphine P-221



(*R*)-form

C₁₈H₁₉NO₅ 329.352

(*R*)-form

2,10,11-Tri-Me ether: Alkaloid CC 24

[35320-73-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the corms of *Colchicum cornigerum* (Liliaceae). Mp 245-249°. λ_{max} 221 (log ε 4.45); 281 (log ε 3.6) (EtOH).

2,10,12-Tri-Me ether: Multifloramine

[22324-15-8]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Kreysigia multiflora* and *Colchicum szovitsii* (Liliaceae). Mp 215-216°. [α]_D²⁵ -111.5 (MeOH). λ_{max} 221 (log ε 4.45); 259 (log ε 3.95); 289 (log ε 3.62); 298 (log ε 3.55) (EtOH).

2,11,12-Tri-Me ether: Floramultine

[21305-36-2]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Kreysigia multiflora* (Liliaceae). Mp 230-231° dec. [α]_D¹⁸ -77 (c, 1.19 in CHCl₃). λ_{max} 220 (ε 45100); 259 (ε 13400); 287 (ε 6500) (no solvent reported).

2,10,11,12-Tetra-Me ether: (-)-Kreysigine

[23117-57-9]

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Colchicum cornigerum* and *Bulbocodium vernum* (Liliaceae). Mp 112-113° (122-124°). [α]_D¹⁵ -65 (c, 0.40 in CHCl₃). λ_{max} 218 (log ε 4.62); 257 (log ε 4.1); 293 (log ε 3.67) (EtOH).

(*S*)-form

1,2,11-Tri-Me ether: Baytopine

[120947-99-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the leaves and flowers of *Merendera kurdica* (Liliaceae). Amorph. solid. [α]_D²⁰ +74 (c, 0.28 in CHCl₃). λ_{max} 215 (log ε 4.55); 257 (log ε 4.02); 287 (sh) (log ε 3.67); 296 (sh) (log ε 3.62) (EtOH).

1,10,12-Tri-Me ether: Merobustinine

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Merendera robusta*. Cryst. (Me₂CO). Mp 216-218°. [α]_D +42 (c, 0.50 in CHCl₃). CAS no. not found 8-14Cl. λ_{max} 258 ; 292

(MeOH).

1,11,12-Tri-Me ether: Androbine

[125326-08-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the bulbs of *Androcymbium palaestinum* (Liliaceae). [α]_D +39 (c, 0.12 in MeOH). λ_{max} 215 (log ε 4.44); 262 (log ε 3.98); 289 (sh) (log ε 3.65) (MeOH).

1,11,12-Tri-Me ether, N-de-Me: Norandrobine

[125326-10-5]

C₂₀H₂₃NO₅ 357.405

Minor alkaloid from the bulbs of *Androcymbium palaestinum* (Liliaceae). [α]_D +20 (c, 0.10 in MeOH). λ_{max} 218 (log ε 3.89); 258 (log ε 3.42); 291 (sh) (log ε 3.1) (MeOH).

2,10,11-Tri-Me ether: Merobustine. Decaisnine

[170554-73-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Merendera robusta* and tubers of *Colchicum decaisnei*. Cryst. (Me₂CO). Mp 241-242°. [α]_D +76 (c, 0.6 in CHCl₃). Enantiomer of CC 24. λ_{max} 219 ; 288 (MeOH).

2,10,12-Tri-Me ether: (+)-Multifloramine

[22324-16-9]

Alkaloid from the tubers of *Colchicum decaisnei*. Mp 214-215° (synthetic). [α]_D²⁵ +112 (synthetic). [α]_D +112 (c, 0.2 in CHCl₃) (natural).

2,11,12-Tri-Me ether: Merenderine. Bechuanine

[1354-66-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Iphigenia bechuanica*, *Iphigenia indica*, *Iphigenia pallida*, *Merendera raddeana* and *Merendera trigina* (Liliaceae). Mp 232-235° dec. [α]_D²² +76 (c, 0.79 in CHCl₃). Enantiomer of Floramultine. λ_{max} 218 (log ε 4.6); 260 (log ε 4.11); 291 (log ε 3.71) (95% EtOH).

2,11,12-Tri-Me ether, N-oxide: Merenderine N-oxide

[142542-98-1]

C₂₁H₂₅NO₆ 387.432

Alkaloid from leaves and stems of *Merendera raddeana* (Liliaceae). Cryst. (Me₂CO). Mp 251-252°. [α]_D +125 (c, 0.94 in MeOH). λ_{max} 260 (log ε 4.16); 290 (log ε 3.77) (MeOH).

1,2,10,11-Tetra-Me ether: Szovitsamine

[56596-02-2]

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Colchicum decaisnei*, *Colchicum szovitsii* and *Androcymbium palaestinum* (Liliaceae). Cryst. (Et₂O). Mp 188-190°. [α]_D +55 (c, 0.10 in MeOH). [α]_D²⁰ +86 (c, 1.4 in CHCl₃). λ_{max} 218 (log ε 4.29); 260 (log ε 3.78); 286 (sh) (log ε 3.5) (MeOH).

1,2,10,11-Tetra-Me ether, N-oxide(β-): Szovitsamine N-oxide

[172685-49-3]

C₂₂H₂₇NO₆ 401.458

Alkaloid from *Colchicum ritchii*. Amorph. [α]_D²⁵ +54 (c, 0.35 in MeOH). λ_{max} 217 (log ε 4.87); 222 (log ε 4.83);

292 (log ε 4.1) (MeOH).

1,2,11,12-Tetra-Me ether: Androcimine

[125326-07-0]

C₂₂H₂₇NO₅ 385.459

Alkaloid from the bulbs of *Androcymbium palaestinum* (Liliaceae). [α]_D +51 (c, 0.11 in MeOH). λ_{max} 218 (log ε 4.47); 260 (log ε 3.94); 291 (sh) (log ε 3.6) (MeOH).

1,10,11,12-Tetra-Me ether: Androcine

[125326-06-9]

C₂₂H₂₇NO₅ 385.459

Alkaloid from the bulbs of *Androcymbium palaestinum* (Liliaceae). [α]_D +39 (c, 0.10 in MeOH). λ_{max} 217 (log ε 4.47); 260 (log ε 3.99); 292 (sh) (log ε 3.93) (MeOH).

2,10,11,12-Tetra-Me ether: (+)-Kreysigine

[26287-94-5]

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Androcymbium palaestinum* and *Colchicum decaisnei* (Liliaceae). Mp 112-113°. [α]_D¹⁵ +60 (c, 0.40 in CHCl₃). [α]_D +64 (c, 0.11 in MeOH). λ_{max} 216 (log ε 3.94); 257 (log ε 3.49); 287 (sh) (log ε 3.24) (MeOH).

2,10,11,12-Tetra-Me ether, N-oxide(β-): Kreysigine N-oxide

[142542-96-9]

C₂₂H₂₇NO₆ 401.458

Alkaloid from *Merendera raddeana* (Liliaceae) and *Colchicum ritchii*. Mp 143-145°. [α]_D²⁵ +32 (c, 0.42 in MeOH). Oxide stereochem. is probable. λ_{max} 219 (log ε 4.9); 258 (log ε 4.34); 294 (log ε 4.03) (MeOH).

Penta-Me ether: O-Methylkreysigine

[58116-30-6]

C₂₃H₂₉NO₅ 399.486

Alkaloid from *Androcymbium palaestinum* and *Merendera raddeana* (Liliaceae). Oil. [α]_D +68 (c, 0.10 in MeOH). [α]_D +81 (CHCl₃). Szovitsine was an inseparable mixt. of homoaporphine isomers methylated to this penta-Me ether. λ_{max} 220 (log ε 4.44); 260 (log ε 3.95); 289 (sh) (log ε 3.46) (MeOH).

Penta-Me ether, N-oxide: O-Methylkreysigine N-oxide

[142542-97-0]

C₂₃H₂₉NO₆ 415.485

Alkaloid from *Merendera raddeana* (Liliaceae). Mp 161-163°.

Penta-Me ether, N-de-Me: Nor-O-methylkreysigine

[125326-09-2]

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Androcymbium palaestinum* (Liliaceae). [α]_D +27 (c, 0.07 in MeOH). λ_{max} 218 (log ε 4.37); 259 (log ε 3.88); 292 (sh) (log ε 3.38) (MeOH).

(±)-form

2,10,12-Tri-Me ether: [16845-28-6]

Mp 190-192° dec.

2,10,11,12-Tetra-Me ether: (±)-Kreysigine

[31735-22-5]

C₂₂H₂₇NO₅ 385.459
Alkaloid from *Kreysigia multiflora* (Liliaceae). Cryst. (EtOH). Mp 187-188°. λ_{max} 221 (ε 53890); 260 (ε 15600); 293 (ε 6600) (no solvent reported).

Penta-Me ether: [32689-00-2]
Mp 152.3° (as methiodide).

(ξ)-form

1,10,11-Tri-Me ether: *Szovitsinine*

[65022-65-3]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Colchicum szovitsii* (Liliaceae). No phys. props. reported.

2,11,12-Tri-Me ether, N-Me: *N-Methylmerenderine*

[664994-43-8]

C₂₂H₂₈NO₅⁺ 386.467

Alkaloid from *Merendera robusta*. Amorph. (as hydroxide).

Badger, G.M. et al., *J.C.S.*, 1960, 445-447 (*Floramultine, isol, uv, ir*)

Potesilová, H. et al., *Coll. Czech. Chem. Comm.*, 1969, **34**, 3540-3552 (*Alkaloid CC 24*)

Brossi, A. et al., *Helv. Chim. Acta*, 1969, **52**, 678-689 (*Multifloramine, synth, ir, uv, abs config*)

Šantavý, F. et al., *Helv. Chim. Acta*, 1971, **54**, 1084-1095 (*Kreysigine, isol, ir, uv, struct*)

Battersby, A.R. et al., *J.C.S. (C)*, 1971, 3514-3518 (*Alkaloid CC 24, uv, ir, pmr, struct*)

Kametani, T. et al., *J.C.S. Perkin 1*, 1972, 2160-2162 (*Kreysigine, synth*)

Santavý, F. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 1712-1713 (*Floramultine, Bechuanine, isol, uv*)

Hoshino, O. et al., *Chem. Pharm. Bull.*, 1974, **22**, 1307-1312 (*Kreysigine, synth, ir, pmr, ms*)

Battersby, A.R. et al., *J.C.S. Perkin 1*, 1974, 1394-1399; 1399-1402 (*Kreysigine, Multifloramine, Floramultine, isol, uv, ir, pmr, ms, biosynth, synth, struct*)

Yusupov, M.K. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 127-128 (*Szovitzamine*)

Trozyan, A.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 557 (*Merenderine*)

Kasimov, A.K. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 197-200 (*Szovitsinine, Szovitsamine, ms, pmr, struct*)

Kupchan, S.M. et al., *J.O.C.*, 1978, **43**, 4076-4081 (*Multifloramine, synth*)

Tojo, E. et al., *J. Nat. Prod.*, 1989, **52**, 1055-1059 (*O-Methylkreysigine, Nor-O-methylkreysigine, Androbine, Androcimine, Androcine, Norandrobine, Szovitsamine, isol, uv, ir, pmr, ms, cd, struct*)

Husek, A. et al., *Phytochemistry*, 1989, **28**, 3217-3219 (*Baytopine, isol, uv, ir, pmr, cmr, ms, cd, struct*)

Yusupov, M.K. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1991, **27**, 75-79 (*Kreysigine N-oxide, O-Methylkreysigine N-oxide, Merenderine N-oxide, isol*)

Alikulov, R.V. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 767-770 (*Merobustine, Merobustinine*)

Abu Zarga, M.H. et al., *J. Prakt. Chem.*, 1995, **337**, 675-679 (*Colchicum decaisnei constits*)


Herbert, R.B. et al., *Tet. Lett.*, 1995, **36**, 5649-5650 (*Kreysigine, synth*)

Abu Zarga, M.H. et al., *Z. Naturforsch., B*, 1995, **50**, 1424-1427 (*Kreysigine N-oxide, Methylkreysigine N-oxide, Szovitsamine N-oxide*)

Alikulov, R.V. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2003, **39**, 212-214 (*N-Methylmerenderine*)

Itoh, T. et al., *Chem. Pharm. Bull.*, 2003, **51**, 951-955 (*O-Methylkreysigine, synth*)

3,5,8,9,15-Pentahydroxy-6(17),11-jatrophadien-14-one P-222



C₂₀H₃₂O₆ 368.469

(2β,3β,5α,8α,9α,11E,13β,15β)-form

9-(3-Pyridinecarbonyl), 3-benzoyl, 5,8-di-Ac: **Euphocharacin E**

[777896-19-2]

C₃₇H₄₃NO₁₀ 661.747

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -16.7 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3-benzoyl, 5,8,15-tri-Ac: **Euphocharacin F**

[777896-21-6]

C₃₉H₄₅NO₁₁ 703.785

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -2.1 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3-(2-methylpropanoyl), 5,8-di-Ac: **Euphocharacin G**

[777896-23-8]

C₃₄H₄₅NO₁₀ 627.73

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -25 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3-(2-methylpropanoyl), 5,8,15-tri-Ac: **Euphocharacin H**

[777896-25-0]

C₃₆H₄₇NO₁₁ 669.767

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -17.3 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3-propanoyl, 5,8,15-tri-Ac: **Euphocharacin I**

[777896-27-2]

C₃₅H₄₅NO₁₁ 655.741

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -22 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3,5,8,15-tetra-Ac: **Euphocharacin J**

[777896-29-4]

C₃₄H₄₃NO₁₁ 641.714

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -46 (c, 0.1 in CHCl₃).

9-(3-Pyridinecarbonyl), 3-(2-methylbutanoyl), 8,15-di-Ac: **Euphocharacin K**

[777896-31-8]


C₃₅H₄₇NO₁₀ 641.757

Constit. of *Euphorbia characias*.

Amorph. solid. [α]_D²⁵ -19.33 (c, 0.1 in CHCl₃).

Corea, G. et al., *Planta Med.*, 2004, **70**, 657-665 (*isol, pmr, cmr*)

3,8,12,14,17-Pentahydroxypregn-5-en-20-one P-223



(3β,12β,14β,17βOH)-form

C₂₁H₃₂O₆ 380.48

(3β,12β,14β,17βOH)-form

Deacetylmataplexigenin

[3513-04-0]

Constit. of *Cynanchum wallichii* and *Marsdenia globifera*. Cryst. (Me₂CO). Mp 220-224°.

12-(3-Pyridinecarbonyl): **Rostratamine**.

12-O-Nicotinoyldeacetylmataplexigenin [50299-46-2]

C₂₇H₃₅NO₇ 485.576

Alkaloid from *Marsdenia rostrata* and *Leptadenia hastata*. Needles (MeOH/Et₂O). Mp 277-279° (198-203°). [α]_D²² -32 (c, 1.0 in MeOH). [α]_D²⁵ +178 (c, 1.0 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: [174097-87-1]

C₄₈H₇₁NO₁₆ 918.086

Constit. of *Cynanchum caudatum*.

Amorph. powder. [α]_D²⁰ -3.8 (c, 1.2 in MeOH).

12-O-(3-Pyridinecarbonyl), 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: [869768-73-0]

C₄₈H₇₁NO₁₇ 934.085

Constit. of *Stephanotis mucronata*.

Amorph. powder.

12-(3-Pyridinecarbonyl), 3-O-[α-L-cymaropyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: [174097-98-4]

C₅₅H₈₃NO₁₉ 1062.256

Constit. of *Cynanchum caudatum*.

Amorph. powder. [α]_D²⁰ -34.2 (c, 0.59 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: [175667-85-3]

C₅₅H₈₃NO₂₀ 1078.255

Constit. of *Leptadenia hastata*.

Amorph. powder. [α]_D²⁵ +157 (c, 1 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-digitoxopyranoside]: [267422-68-4]

C₅₂H₇₇NO₁₉ 1020.176

Constit. of *Asclepias incarnata*.

Amorph. powder. [α]_D²⁴ -4.6 (c, 0.61 in MeOH).

MeOH). λ_{\max} 218 (log ϵ 3.97); 258 (log ϵ 3.39); 263 (log ϵ 3.42); 270 (sh) (MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β -D-oleandropyranosyl-(1 \rightarrow 4)- β -D-oleandropyranosyl-(1 \rightarrow 4)- β -D-cymaropyranosyl-(1 \rightarrow 4)- β -D-cymaropyranoside]: [174097-95-1]
C₅₅H₈₃NO₁₉ 1062.256
Constit. of *Cynanchum caudatum*. Amorph. powder. $[\alpha]_D^{20}$ -6.1 (c, 0.4 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β -D-thevetopyranosyl-(1 \rightarrow 4)- β -D-cymaropyranosyl-(1 \rightarrow 4)- β -D-cymaropyranosyl-(1 \rightarrow 4)- β -D-cymaropyranoside]: [646063-90-3]
C₅₅H₈₃NO₂₀ 1078.255
Constit. of *Araujia sericifera*. Amorph. powder. $[\alpha]_D^{23}$ +10 (c, 0.36 in MeOH). λ_{\max} 202 (log ϵ 4.25); 217 (log ϵ 4.18); 250 (log ϵ 3.71); 255 (log ϵ 3.74); 260 (log ϵ 3.7) (MeOH).

Gellert, E. et al., *Aust. J. Chem.*, 1973, **26**, 1835 (*Rostratine*)

Warashina, T. et al., *Chem. Pharm. Bull.*, 1995, **43**, 977-982; 1734-1737; 1996, **44**, 358-363 (*Cynanchum caudatum saponins*)

Aquino, R. et al., *J. Nat. Prod.*, 1995, **58**, 672; 1996, **59**, 555 (*Rostratine glycosides*)


Aquino, R. et al., *J. Nat. Prod.*, 1996, **59**, 555-564 (*Leptadenia hastata saponins*)

Warashina, T. et al., *Chem. Pharm. Bull.*, 2000, **48**, 99-107 (*Asclepias incarnata saponins*)

Warashina, T. et al., *Phytochemistry*, 2000, **53**, 485-488 (*Asclepias incarnata saponins*)

Warashina, T. et al., *Chem. Pharm. Bull.*, 2003, **51**, 1036-1045 (*Araujia sericifera saponins*)

5,7,9,13,14-Pentahydroxy-4(20),11-taxadien-10-one P-224



C₂₀H₃₀O₆ 366.453

(5 α ,7 β ,9 α ,13 α ,14 β)-form

7,9-Di-Ac: [126585-89-5]

C₂₄H₃₄O₈ 450.528

From the bark of *Austrotaxus spicata*.

Amorph. $[\alpha]_D$ +11 (c, 0.59 in CHCl₃).

5-O-(3 ξ -Dimethylamino-3-phenylpropanoyl), 7,9,13,14-tetra-Ac: **2'-Deacetox-yaustrotaxine**
[119777-74-1]
C₃₉H₅₁NO₁₁ 709.832

From the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. $[\alpha]_D$ -37 (c, 0.5 in CHCl₃).

5-O-(3 ξ -Dimethylamino-2 ξ -hydroxy-3-phenylpropanoyl), 7,9-di-Ac: **2',13,14-Trideacetylaustrotaxine**
[126585-88-4]
C₃₅H₄₇NO₁₀ 641.757

From the bark of *Austrotaxus spicata*

(Taxaceae). Amorph. $[\alpha]_D$ -21 (c, 0.86 in CHCl₃).

5-O-(3 ξ -Dimethylamino-2 ξ -hydroxy-3-phenylpropanoyl), 7,9,13,14-tetra-Ac: **2'-Deacetylaustrotaxine**
[119777-73-0]
C₃₉H₅₁NO₁₂ 725.831


From the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. $[\alpha]_D$ -41 (c, 0.3 in CHCl₃).

5-O-(2 ξ -Acetoxy-3 ξ -dimethylamino-3-phenylpropanoyl), 7,9,13,14-tetra-Ac: **Austrotaxine**
[119789-82-1]
C₄₁H₅₃NO₁₃ 767.869

Alkaloid from the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. $[\alpha]_D$ -49 (c, 0.4 in CHCl₃).

Etouati, L. et al., *Bull. Soc. Chim. Fr.*, 1988, 749-755; 1989, 687-694 (*isol, pmr, cmr, struct*)

4,9,12,13,20-Pentahydroxy-1,6-tigliadien-3-one P-225



(4 α ,9 α ,12 β ,13 α)-form

C₂₀H₂₈O₆ 364.438

(4 α ,9 α ,12 β ,13 α)-form

Isophorbol. 4 α -Phorbol

[26241-63-4]

Constit. of croton oil. Hygroscopic cryst. (EtOAc). Mp 135-150°.

12-O-(2-Methylaminobenzoyl), 13-Ac:

[82042-11-3]

C₃₀H₃₇NO₈ 539.624

Constit. of *Sapium indicum*.

[71295-86-8, 16675-05-1, 25405-85-0, 57716-89-9, 24928-17-4, 37558-16-0, 24928-15-2, 61557-88-8]


Taylor, S.E. et al., *J. Nat. Prod.*, 1981, **44**, 729-731 (*Sapintoxin D*)

Taylor, S.E. et al., *J. Pharm. Pharmacol., Suppl.*, 1981, **33**, 54P (*Sapintoxin D, activity*)

Evans, F.J. et al., *Prog. Chem. Org. Nat. Prod.*, 1983, **44**, 1-99 (rev)

Naturally Occurring Phorbol Esters, (ed., Evans, F.J.), CRC Press, 1986, (book)

5,9,12,13,20-Pentahydroxy-1,6-tigliadien-3-one P-226



C₂₀H₂₈O₆ 364.438

Esters show varying degrees of antiplatelet activity.

(4 β ,5 β ,9 α ,12 β ,13 α)-form

12-(2,4,6-Decatrienoyl) (2Z,4E,6 ξ -), 13-Ac: **Sapatoxin B**

[82345-49-1]

C₃₂H₄₂O₈ 554.679

Constit. of *Sapium indicum*.

12-O-(2-Methylaminobenzoyl), 13-Ac:

Sapintoxin B

[81345-29-1]

C₃₀H₃₇NO₈ 539.624

Constit. of *Sapium indicum*.

Taylor, S.E. et al., *J. Pharm. Pharmacol.*, 1981, **33** (Suppl.), 54P (*Sapintoxin B, isol*)

Taylor, S.E. et al., *Phytochemistry*, 1981, **20**, 2749-2751; 1982, **21**, 405-407 (*Sapatoxin B, Sapintoxin B, isol, uv, ir, pmr*)

Edwards, M.C. et al., *Acta Pharmacol. Toxicol.*, 1983, **53**, 177-187 (*isol, pps*)

Pentalupine P-227


C₁₆H₃₀N₂O 266.426

Struct. unknown. Alkaloid from *Lupinus palmeri* (Fabaceae). Yellow oil with strong greenish fluor. Bp₂ 175-182°. $[\alpha]_D$ -3.2 (EtOH) (impure). n_D^{20} 1.5155. No cryst. derivs. could be obt.

Couch, J.F. et al., *J.A.C.S.*, 1934, **56**, 2434-2436

2,3,4,6,7-Pentamethoxyphenanthroindolizidine P-228

9,11,12,13,13a,14-Hexahydro-2,3,4,6,7-pentamethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinoline, 8CI. Alkaloid E†
[30062-26-1]



(S)-form

C₂₅H₂₉NO₅ 423.508

(S)-form

14S-Hydroxy, β -N-oxide: **Ficuseptine A**

C₂₅H₂₉NO₇ 455.507

Alkaloid from the leaves of *Ficus septica*.

Cytotoxic. Pale yellow powder.

Mp 210° dec. $[\alpha]_D$ +30.3 (c, 0.03 in MeOH).

λ_{\max} 214 (log ϵ 4.35); 263 (log ϵ 4.19); 282 (log ϵ 4); 359 (log ϵ 3.11); 405 (log ϵ 2.79) (MeOH).

(ξ)-form

Alkaloid from *Tylophora crebriflora* (Asclepiadaceae).

14 ξ -Hydroxy: **14-Hydroxy-2,3,4,6,7-pentamethoxyphenanthroindolizidine**. Alkaloid D†
[30062-25-0]

C₂₅H₂₉NO₆ 439.507

Alkaloid from *Tylophora crebriflora*

(Asclepiadaceae). Cryst. (Et₂O/diisopropyl ether) (as Ac). Mp 188-190° (Ac).

Rao, K.V. et al., *J. Pharm. Sci.*, 1970, **59**, 1608-1611 (*Alkaloid D, Alkaloid E*)

Wu, P.-L. *et al.*, *Heterocycles*, 2002, **57**, 2401-2408 (*Ficuseptine A*)

1,5-Pentanediamine, 9CI P-229

Pentamethylenediamine. 1,5-Diaminopentane. **Cadaverine**

[462-94-2]
H₂NCH₂(CH₂)₃CH₂NH₂

C₅H₁₄N₂ 102.179

Found in nature as bacterial decarboxylation prod. of Lysine, e.g. in putrefaction. Anal sac secretion of the red fox *Vulpes vulpes*. Occurs in *Anthracidaris crassispina* and *Halocynthia roretzi*.

Monomer for polyamides. Syrupy fuming liq. d₂₀²⁰ 0.84. Bp 178-180°. n_D²⁰ 1.4582. pK_{a1} 10.92; pK_{a2} 10.05 (25°, 0.5M KNO₃).

► Skin irritant. SA0200000

Hydrochloride (1:2): [1476-39-7]
Needles (H₂O). Mp 226° Mp 260-262°.

N-Benzoyl: [29833-52-1]

C₁₂H₁₈N₂O 206.287

Oil. Bp_{0.5} 202°.

N,N'-Dibenzoyl: [31991-79-4]

C₁₉H₂₂N₂O₂ 310.395

Cryst. (C₆H₆). Mp 135°.

N,N'-Bis(4-nitrobenzoyl): [81852-10-0]

Cryst. (Me₂CO aq.). Mp 185-185.5°.

N-tert-Butyloxycarbonyl: [51644-96-3]

C₁₀H₂₂N₂O₂ 202.296

Oil.

N-Me: [32752-52-6]

C₆H₁₆N₂ 116.206

Liq. Bp 177-178° Bp₁₂ 55°.

N,N'-Di-Me: [56992-95-1]

C₇H₁₈N₂ 130.233

Liq. Bp 190° Bp₁₀ 74°.

N,N,N-Tri-Me: 5-Amino-N,N,N-trimethylpentanaminium(1+), 9CI. **Asco-phylline**

[83524-44-1]

[4175-13-7, 30835-02-0]

C₈H₂₁N₂[⊕] 145.267

Constit. of the alga *Ascophyllum nodosum* and isol. from the beetle *Platyphora opima*.

N,N,N',N'-Tetra-Me: [44994-28-7]

C₉H₂₂N₂ 158.286

Liq. Bp 192-194° Bp₁₅ 85-95°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 292B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 466A (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 376B (*ir*)

Putochin, N. *et al.*, *Ber.*, 1926, **59**, 625-630 (*synth*)

Korshak, V.V. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1955, 756; *Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.)*, 1955, 673 (*polyamides*)

Ger. Pat., 1971, 2 043 141; *CA*, **75**, 6884 (*N-Me*)

Sarneski, J.E. *et al.*, *Anal. Chem.*, 1975, **47**, 2116-2124 (*cmr*)

Mayerl, F. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 127-133 (*ms*)

Albone, E.S. *et al.*, *J. Chem. Ecol.*, 1976, **2**, 167-175 (*occur*)

Richter, R. *et al.*, *J.O.C.*, 1978, **43**, 4150-4154 (*synth*)

Battersby, A.R. *et al.*, *J.C.S. Perkin I*, 1982, 449-453 (*biosynth*)

Blunden, G. *et al.*, *Magn. Reson. Chem.*, 1986, **24**, 965-971 (*Ascophylline*)

Hamana, K. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-62 (*occur*)

Blunden, G. *et al.*, *Biochem. Syst. Ecol.*, 1992, **20**, 373-388 (*Ascophylline, occur*)

Thalladi, V.R. *et al.*, *Angew. Chem., Int. Ed.*, 2000, **39**, 918-922 (*cryst struct*)


Gardner, R. *et al.*, *J. Med. Chem.*, 2004, **47**, 4933-4940 (*N-Boc*)

Pospieszna-Markiewicz, I. *et al.*, *Acta Cryst. C*, 2006, **62**, o399-o401 (*dihydrochloride, cryst struct*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, PBK500

Racophorus Pentapteridinyl skin pigment P-230

[116360-71-5]




C₃₅H₁₅N₂₅O₁₀ 945.665

Proposed struct. shown. Isol. from the skin of the frog of *Racophorus* sp. Black-violet pigment. λ_{max} 480 ; 500 (HCOOH).

Suga, T. *et al.*, *J. Nat. Prod.*, 1988, **51**, 713 (*isol, uv, ir, pmr*)

Pentasin P-231

[108133-63-7]



C₃₀H₄₉N₆O₁₀[⊕] 653.751

Cross-linking amino acid isol. from bovine ligament elastin.

Starcher, B. *et al.*, *Connect. Tissue Res.*, 1987, **16**, 15-25; *CA*, **106**, 191359p

1-Pentene-1,5-diamine P-232

1,5-Diamino-1-pentene

H₂NCH₂CH₂CH₂CH=CHNH₂

C₅H₁₂N₂ 100.163

(E)-form

N,N,N-Tri-Me: 5-Amino-N,N,N-trimethyl-1-pentenaminium(1+). *N,N,N-Trimethyl-1,2-dehydrocadaverine*

C₈H₁₉N₂[⊕] 143.252


Isol. from the beetle *Platyphora opima*. Solid. Counterion not specified. λ_{max} 204 (log ε 3.52) (MeOH).

Plasman, V. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1261-1264

Pentosidine P-233

α-Amino-2-[(4-amino-4-carboxybutyl)amino]-4H-imidazo[4,5-b]pyridine-4-hexanoic acid, 9CI

[124505-87-9]



C₁₇H₂₇N₆O₄[⊕] 379.438


Isol. from the human extracellular matrix of skin and tendon. Produces pentose-mediated protein crosslinking. [α]_D²⁴ +16.5 (c, 0.26 in MeOH) (as trifluoroacetate). Fluorescent.

Sell, D.R. *et al.*, *J. Biol. Chem.*, 1989, **264**, 21597-21602 (*isol, activity*)

Yokokawa, F. *et al.*, *Tetrahedron*, 2001, **57**, 4759-4766 (*synth, pmr, cmr*)

Pentouregine P-234

[99964-90-6]




C₁₈H₁₇NO₃ 295.337

Alkaloid from the leaves of *Gutteria ouregou* (Annonaceae). Amorph. powder. [α]_D -61 (c, 0.14 in EtOH). λ_{max} 224 (log ε 4.31); 284 (log ε 4.14); 303 (sh) (log ε 3.98) (EtOH).

Cortès, D. *et al.*, *Phytochemistry*, 1985, **24**, 2776-2777 (*isol, pmr, ms, struct*)

1-Pentyl-9-azabicyclo[3.3.1]nonan-3-one P-235

Adaline
[41267-60-1]



Absolute Configuration

C₁₃H₂₃NO 209.331

Alkaloid from *Adalia bipunctata* (European ladybug). Mp 204-205° (as hydrochloride). [α]_D -13 (CHCl₃).

Tursch, B. *et al.*, *Bull. Soc. Chim. Belg.*, 1973, **82**, 699 (*abs config, synth*)


Tursch, B. *et al.*, *Tet. Lett.*, 1973, 201 (*ir, pmr, isol, cryst struct*)

Hill, R.K. *et al.*, *Tetrahedron*, 1982, **38**, 1959-1963 (*synth*)

Gnecco Medina, D.H. *et al.*, *Tet. Lett.*, 1983, **24**, 2099-2102 (*synth*)


- Yue, C. *et al.*, *J.O.C.*, 1992, **57**, 4211-4214
(*synth, abs config*)
Davison, E.C. *et al.*, *Tet. Lett.*, 1995, **36**, 9047-9050 (*synth*)
Laurent, P. *et al.*, *Tetrahedron*, 2001, **57**, 3403-3412 (*biosynth*)
Itoh, T. *et al.*, *Org. Lett.*, 2002, **4**, 2469-2472 (*synth*)

6-Pentyl-3-piperidinol, 9CI **P-236**
5-Hydroxy-2-pentylpiperidine
[220088-35-7]



$C_{10}H_{21}NO$ 171.282
Alkaloid from *Conium maculatum*.
Lang, D.G. *et al.*, *CA*, 1999, **130**, 136554u

2-Pentylpyridine, 9CI **P-237**
FEMA 3383
[2294-76-0]




$C_{10}H_{15}N$ 149.235
Present in cooked foods. Present in bell peppers and coriander seed oil. Oil with fatty tallow-like odour and throat-catching greasy taste. d_4^{20} 0.88. Bp_{12} 86-88° $Bp_{8.5}$ 82.5-84.5°. n_D^{16} 1.4890. n_D^{20} 1.4834. Flavour threshold 1.2×10^{-5} ppm in H_2O .

Picrate: [23692-26-4]
Mp 73°.

[65307-85-9]

- Osuch, C. *et al.*, *J.A.C.S.*, 1956, **78**, 1723-1725 (*synth*)
Cuvigny, T. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 2000-2009; 1965, 1881-1888 (*synth*)
Miyadera, T. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 1344-1351 (*synth*)
Waldradt, J.P. *et al.*, *J. Agric. Food Chem.*, 1971, **19**, 972-979 (*occur, peanuts*)
Buttery, R.G. *et al.*, *J. Agric. Food Chem.*, 1977, **25**, 1227-1229 (*occur, glc, ms, lamb fat*)
Tang, J. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 1287-1292 (*occur*)
Mottram, D.S. *et al.*, *J. Sci. Food Agric.*, 1985, **36**, 377-382 (*detn, pork*)
Lamparsky, D. *et al.*, *Perfum. Flavor.*, 1988, **13**, 17-18 (*occur, coriander oil*)
Sutherland, M.M. *et al.*, *J. Sci. Food Agric.*, 1995, **69**, 403-413 (*occur*)
Takeoka, G. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 654-660 (*occur*)
Kim, Y.-S. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3906-3908 (*formn*)
Lambrechts, S. *et al.*, *Flavour Fragrance J.*, 1997, **12**, 439-442 (*synth, props*)
Boatright, W.L. *et al.*, *J. Am. Oil Chem. Soc.*, 1997, **74**, 1575-1581; 1998, **75**, 1379-1383 (*occur, glc, soybean*)
Kim, Y.-S. *et al.*, *J. Food Lipids*, 1997, **4**, 239-244; 1998, **5**, 173-182 (*occur*)

2-Pentylquinoline, 9CI **P-238**
[93005-16-4]



$C_{14}H_{17}N$ 199.295
Constit. of alkaloids of *Cusparia trifoliata* (preferred genus name *Angostura*) and from stems of *Galipea bracteata* (Rutaceae). Molluscicide, plant growth inhibitor. Oil. Bp_8 150-154°.

Picrate:

Cryst. (MeOH). Mp 108.5°.

1',2'-Didehydro: 2-(1-Pentenyl)quinoline
[128396-36-1]

$C_{14}H_{15}N$ 197.279

Alkaloid from *Galipea bracteata* stems. Molluscicide, plant growth inhibitor. Oil.

1,2,3,4-Tetrahydro, N-Me: 1,2,3,4-Tetrahydro-1-methyl-2-pentylquinoline. **Angustureine**
[246511-34-2 ((-)-form)]
 $C_{15}H_{23}N$ 217.353
Alkaloid from the bark of *Galipea officinalis*. Oil. $[\alpha]_D$ -7.2 (CHCl₃). λ_{max} 259 (log ϵ 3.7); 311 (log ϵ 3.15) (CHCl₃).

Späth, E. *et al.*, *Monatsh. Chem.*, 1930, **55**, 352-357 (*isol*)

Gautier, J.A. *et al.*, *Bull. Soc. Chim. Fr.*, 1961, 2092-2098 (*synth*)

Tertov, B.A. *et al.*, *CA*, 1963, **59**, 9978 (*synth*)
Vieira, P.C. *et al.*, *Phytochemistry*, 1990, **29**, 813-815 (*isol, pmr, cmr, ir, ms*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 332-334 (*synth*)

Jacquemond-Collet, I. *et al.*, *Phytochemistry*, 1999, **51**, 1167-1169 (*Angustureine*)

Cho, C.S. *et al.*, *Tetrahedron*, 2003, **59**, 7997-8002 (*synth, pmr, cmr*)

O'Byrne, A. *et al.*, *Tetrahedron*, 2008, **64**, 8067-8072 (*Angustureine, synth*)


Pepuline **P-239**
Struct. unknown. Alkaloid from the bark of *Zanthoxylum rhetsa* (Rutaceae). Spasmodic and cardiotonic agent. Yellow cryst. Mp 179-180°.

Hydrochloride:

Cryst. (EtOH). Mp 222°.

Mehta, C.R. *et al.*, *Curr. Sci.*, 1960, **29**, 95 (*isol*)

Perakine **P-240**
17-(Acetyloxy)-1,2-didehydro-1-dimethyl-21-methyl-18-norajmalan-19-al, 9CI. **Raucaffrine**
[4382-56-3]



Absolute Configuration

$C_{21}H_{22}N_2O_2$ 350.416

Alkaloid from *Rauwolfia perakensis*, *Rauwolfia caffra*, *Rauwolfia vomitoria*, *Rauwolfia volkensii* and *Voacanga africana*

(Apocynaceae). Cryst. (Me₂CO or EtOAc/petrol). Mp 186-189°. $[\alpha]_D^{24}$ +132. $[\alpha]_D^{24}$ +112 (c, 0.85 in CHCl₃). Possibly an artifact.

Di-Me acetal: Perakine dimethyl acetal $C_{23}H_{28}N_2O_4$ 396.485
From leaves of *Rauwolfia sellowii* (Apocynaceae). No phys. data reported. Prob. an artifact of the isol. procedure.

19-Alcohol: **Raucaffrinoline**

[36285-11-7]

$C_{21}H_{24}N_2O_3$ 352.432

Alkaloid from root bark of *Rauwolfia caffra* and *Rauwolfia nitida* and leaves of *Rauwolfia vomitoria* (Apocynaceae). Rods (CH₂Cl₂). Mp 236° dec Mp 249-251°.

10-Methoxy: **10-Methoxyperakine**

[163461-45-8]

$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from aerial parts of *Vinca major* (Apocynaceae). $[\alpha]_D$ -43 (c, 0.003 in CHCl₃).

Kiang, A.K. *et al.*, *J.C.S.*, 1960, 1394 (*isol, uv, ir, pmr, struct*)

Ulshafer, P.R. *et al.*, *Tet. Lett.*, 1961, 363 (*isol, uv, ir, pmr, struct*)

Taylor, W.I. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 611 (*synth*)

Thomas, D.W. *et al.*, *J. Nat. Prod.*, 1968, **31**, 1 (*isol, ms*)

Khan, M.A. *et al.*, *Experientia*, 1972, **28**, 127 (*Perakine, Raucaffrinoline, isol, uv, ir, pmr, ms, struct*)

Habib, M.S. *et al.*, *Phytochemistry*, 1974, **13**, 661 (*isol, uv, ir, ms*)

Libot, F. *et al.*, *Phytochemistry*, 1980, **19**, 989 (*Raucaffrinoline, uv, cmr, struct*)

Amer, M.M. *et al.*, *Phytochemistry*, 1980, **19**, 1833; 1981, **20**, 2569 (*Perakine, Raucaffrinoline, isol*)

Khan, M.A. *et al.*, *Z. Naturforsch., B*, 1982, **37**, 494 (*isol, cmr*)


Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1995, **38**, 1057 (*10-Methoxyperakine*)

Batista, C.V.F. *et al.*, *Phytochemistry*, 1996, **41**, 969 (*Perakine dimethyl acetal*)

Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1456-1460 (*rev*)

Sabino, J.R. *et al.*, *Acta Cryst. E*, 2006, **62**, 3181-3183 (*Raucaffrinoline, cryst struct*)

Peraksine **P-241**
17,19-Epoxy-19,20-dihydro-21-methyl-18-norsarpagan-17-ol, 9CI. **Vomifoline**. **Alkaloid RP5**
[15527-80-7]



Absolute Configuration

$C_{19}H_{22}N_2O_2$ 310.395

Alkaloid from *Rauwolfia perakensis* and several other *Rauwolfia* spp. (Apocynaceae). Cryst. (MeOH). Mp 196-198° (186°). $[\alpha]_D^{25}$ +44 (c, 0.86 in EtOH). Probable artifact.

O-Ac: Mp 142°.

17-Epimer: **Alkaloid RB20**
[20911-77-7]

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from *Rauwolfia verticillata* (Apocynaceae). Prisms (CHCl₃/MeOH trace). Mp 196-197°. [α]_D +19 (c, 0.48 in MeOH).

10-Hydroxy, N4-Me: Verticillatine†

[98243-58-4]

C₂₀H₂₅N₂O₃[⊕] 341.429


Quaternary alkaloid from the roots of *Rauwolfia verticillata* f. *rubrocarpa* (Apocynaceae). Ganglionic blocking agent. Cryst. (as chloride). Mp 324-326° dec. (chloride).

Kiang, A.K. *et al.*, *Tetrahedron*, 1966, **22**,3293-3300 (*uv, ms, pmr, cryst struct*)Arthur, H.R. *et al.*, *Aust. J. Chem.*, 1968, **21**,1399-1401 (*ms, pmr, struct, Alkaloid RB20*)Habib, M.S. *et al.*, *Phytochemistry*, 1974, **13**,661-662 (*isol, uv, ir, ms*)Pousset, J.-L. *et al.*, *Phytochemistry*, 1977, **16**,153-154 (*uv, ir, pmr, ms, struct*)Lin, M. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 198-202; *CA*, **103**, 138514x (*Verticillatine, cryst struct*)Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 2000, **63**,1456-1460 (*rev*)**Peramine****P-242**


3-(1,2-Dihydro-2-methyl-1-oxopyrrolo[1,2-a]pyrazin-3-yl)propylguanidine, 9CI

[102482-94-0]

[105749-70-0 (di-Ac)]

C₁₂H₁₇N₅O 247.299

Isol. from *Lolium perenne* infected *Acremonium loliae*. Insect deterrent and antifeedant. Cryst. (as di-Ac). Sol. MeOH; fairly sol. Me₂CO. Mp 141-142° (di-Ac). λ_{max} 231 (ε 33100); 237; 285 (MeOH) (Berdy).

Rowan, D.D. *et al.*, *Chem. Comm.*, 1986, 935(*uv, ir, pmr, cmr, ms, struct*)Rowan, D.D. *et al.*, *J. Nat. Prod.*, 1989, **52**, 193(*isol*)Brimble, M.A. *et al.*, *J.C.S. Perkin 1*, 1990, 311(*synth*)**Per(deacyl)alatusamine****P-243**C₂₆H₃₅NO₁₁ 537.563

1,2,6,8,9,14-Hexa-Ac: **Alatusamine** [89945-80-2]

C₃₈H₄₇NO₁₇ 789.786

Alkaloid from the fruits of *Euonymus alatus* forma *striatus* (Celastraceae). Amorph. solid. [α]_D²⁵ -11 (c, 3.0 in CHCl₃).

1,2,6,8,9,14-Hexa-Ac, picrate: Mp 138-144°.

1-Benzoyl, 6,8,9,14-tetra-Ac: **Euojaponin J**

[128397-46-6]

C₄₁H₄₇NO₁₆ 809.819

Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Needles (EtOH). Mp 243°. [α]_D²⁵ +26.2 (c, 0.042 in CHCl₃).

2-Benzoyl, 1,6,8,9,14-penta-Ac: **Neowilforine**. **Chuchuhuanine WI**

[121880-18-0]

C₄₃H₄₉NO₁₇ 851.857


Alkaloid from *Tripterygium wilfordii* and root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 149-154°. [α]_D +14.8 (c, 0.68 in CHCl₃).

1-(3-Pyridinecarbonyl), 2-benzoyl, 6,8,9,14-tetra-Ac: **Euojaponin N**

[199442-94-9]

C₄₇H₅₀N₂O₁₇ 914.915

Alkaloid from *Euonymus japonica*. Mp 161-163°. [α]_D²⁸ +31.6 (c, 0.1 in EtOH).

Ishiwata, H. *et al.*, *Phytochemistry*, 1983, **22**,2839 (*Alatusamine*)He, Z. *et al.*, *Huaxue Xuebao*, 1989, **47**, 178;*CA*, **111**, 74780m (*Neowilforine*)Han, B.H. *et al.*, *J. Nat. Prod.*, 1990, **53**, 909(*Euojaponin J*)Shirota, O. *et al.*, *Heterocycles*, 1994, **38**, 2219(*Norwilforine*)Ryu, J.-H. *et al.*, *CA*, 1998, **128**, 20569g(*Euojaponin N*)**Per(deacyl)chuchuhuanine****P-244**C₂₆H₃₅NO₁₁ 537.563

24 and 25 configs. not defined for the Chuchuhuanines. Given here as (24S,25S) in uniformity with related alkaloids.

1,2,8,9,14-Penta-Ac: **Chuchuhuanine EIV**

C₃₆H₄₅NO₁₆ 747.749

From root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 162-166°. [α]_D -9.3 (c, 0.51 in CHCl₃).

1,6,8,9,14-Penta-Ac: **Chuchuhuanine EV**

C₃₆H₄₅NO₁₆ 747.749

From root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 154-160°. [α]_D -18.3 (c, 0.45 in CHCl₃).

1,2,6,8,9,14-Hexa-Ac: **Chuchuhuanine EI**

C₃₈H₄₇NO₁₇ 789.786

Alkaloid from root bark of *Maytenus chuchuhuasca* (Celastraceae). Plates. Mp 247-250°. [α]_D -23.3 (c, 0.77 in CHCl₃).

1-Benzoyl, 2,6,8,9,14-penta-Ac: **Chuchuhuanine EII**

C₄₃H₄₉NO₁₇ 851.857

From root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 158-160°. [α]_D -8.5 (c, 0.46 in CHCl₃).

2-Benzoyl, 1,6,8,9,14-penta-Ac: **Chuchuhuanine EIII**

C₄₃H₄₉NO₁₇ 851.857

From root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 156-160°. [α]_D -412.5 (c, 0.04 in CHCl₃).

1,2-Dibenzoyl, 6,8,9,14-tetra-Ac: **Chuchuhuanine EVI**

C₄₈H₅₁NO₁₇ 913.927

Alkaloid from the bark of *Maytenus chuchuhuasca*. Amorph. solid. λ_{max} 229 (log ε 4.48); 265 (log ε 3.69) (MeOH).

1,8-Dibenzoyl, 2,6,9,14-tetra-Ac: **Chuchuhuanine EVII**

C₄₈H₅₁NO₁₇ 913.927

Alkaloid from the bark of *Maytenus chuchuhuasca*. Amorph. solid. λ_{max} 230 (log ε 4.49); 265 (log ε 3.67) (MeOH).

8-Epimer, 1,2,6,8,9,14-hexa-Ac: **4-Deoxyeuonymine**


C₃₈H₄₇NO₁₇ 789.786

From root bark of *Maytenus chuchuhuasca* (Celastraceae). Amorph. solid. Mp 138-142°. [α]_D -13.6 (c, 1.41 in CHCl₃).

8-Epimer, 1-benzoyl, 2,6,8,9,14-penta-Ac: **Ebenifoline EIV**. 4-Deoxymayteine

C₄₃H₄₉NO₁₇ 851.857

Alkaloid from stem bark of *Maytenus ebenifolia* (Celastraceae). Amorph. solid. Mp 143-146°. [α]_D -7.8 (c, 0.25 in CHCl₃).

Itokawa, H. *et al.*, *J.C.S. Perkin 1*, 1993, 1247(*Ebenifoline EIV*)Shirota, O. *et al.*, *Heterocycles*, 1994, **38**, 2219(*isol, uv, ir, pmr, ms, cd, struct*)Shirota, O. *et al.*, *Heterocycles*, 2004, **63**, 1891-1896 (*Chuchuhuanines EVI, EVII*)**Per(deacyl)-24-demethylemarginatine****P-245**

C₃₂H₃₈N₂O₁₄ 674.657

Parent struct. related to Per(deacyl)emarginatine, P-246 lacking the C-24 methyl group.

1,6,8,9,14-Penta-Ac:

C₄₂H₄₈N₂O₁₉ 884.843

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 177-184°. [α]_D²⁵ +1.8 (c, 0.66 in CHCl₃). λ_{max} 269 (log ε 4.43) (no solvent reported).

14-(2-Methylpropanoyl), 1,6,8,9-tetra-Ac:

C₄₄H₅₂N₂O₁₉ 912.897

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 178-192°. [α]_D²⁵ +4.2 (c, 0.7 in CHCl₃). λ_{max} 269 (log ε 4.39) (no solvent reported).

8,14-Bis(2-methylpropanoyl), 1,6,9-tri-Ac:

C₄₆H₅₆N₂O₁₉ 940.95


Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 161-164°. [α]_D²⁵ +2.6 (c, 0.4 in CHCl₃). λ_{max} 268 (log ε 4.33) (no solvent reported).

Furukawa, M. et al., *Phytochemistry*, 2002, **59**, 767-777

Per(deacyl)emarginatine

P-246

Perdeacylhippocrateine



C₃₃H₄₀N₂O₁₄ 688.684

1,6,8,9-Tetra-Ac: **Emarginatine H**

[168434-00-2]

C₄₁H₄₈N₂O₁₈ 856.833

Alkaloid from leaves of *Maytenus diversifolia* (Celastraceae). Amorph. powder. Mp 313-316°. [α]_D²⁵ +35 (c, 0.05 in CHCl₃).

1,6,8,14-Tetra-Ac: **Emarginatine C**

[156280-96-5]

C₄₁H₄₈N₂O₁₈ 856.833

Alkaloid from stems of *Maytenus emarginata* (Celastraceae). Needles. Mp 312-315°. [α]_D²⁵ +16 (c, 0.05 in CHCl₃).

6,8,9,14-Tetra-Ac: **Emarginatine D**

[156280-97-6]

C₄₁H₄₈N₂O₁₈ 856.833

Alkaloid from stems of *Maytenus emarginata* (Celastraceae). Needles. Mp 192-202°. [α]_D²⁵ +51 (c, 0.033 in CHCl₃).

1,6,8,9,14-Penta-Ac: **Emarginatine A**

[125227-50-1]

C₄₃H₅₀N₂O₁₉ 898.87

Alkaloid from *Hippocratea excelsa* and

Maytenus emarginata (Hippocrataceae). Cytotoxic agent. Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 312-313°. [α]_D²⁵ +70 (c, 0.32 in CHCl₃). λ_{max} 266 (MeOH) (Berdy).

8-(2-Methylpropanoyl), 1,6,9,14-tetra-Ac:

C₄₅H₅₄N₂O₁₉ 926.924

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 300-307°. [α]_D²⁵ +15.3 (c, 1 in CHCl₃). λ_{max} 268 (log ε 4.3) (no solvent reported).

9-(2-Methylpropanoyl), 6,8,14-tri-Ac:

C₄₃H₅₂N₂O₁₈ 884.886

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 194-198°. [α]_D²⁵ -4.3 (c, 1 in CHCl₃). λ_{max} 269 (log ε 4.31) (no solvent reported).

14-(2-Methylpropanoyl), 1,6,8,9-tetra-Ac:

C₄₅H₅₄N₂O₁₉ 926.924

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 246-249°. [α]_D²⁵ +15.5 (c, 3 in CHCl₃). λ_{max} 268 (log ε 4.34) (no solvent reported).

1,9-Bis(2-methylpropanoyl), 6,8,14-tri-Ac:

C₄₇H₅₈N₂O₁₉ 954.977

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 177-183°. [α]_D²⁵ -2.3 (c, 0.9 in CHCl₃). λ_{max} 268 (log ε 4.37) (no solvent reported).

1-(2-Methyl-2-butenoyl), 6,8,9,14-tetra-Ac: **Emarginatine G**

[155944-13-1]

C₄₆H₅₄N₂O₁₉ 938.935

Alkaloid from stems of *Maytenus emarginata* (Celastraceae). Needles. Mp 315-318°. [α]_D²⁰ +200 (c, 0.02 in CHCl₃). Incorrect struct. given in CA.

1-O-Benzoyl, 6,8,9,14-tetra-Ac: **Hippocrateine I**

[132185-31-0]

C₄₈H₅₂N₂O₁₉ 960.941

Alkaloid from roots and stem bark of *Hippocratea excelsa* (Hippocrataceae). Cryst. Mp 300°. [α]_D²⁵ +50 (c, 0.33 in CHCl₃).

1-O-Benzoyl, 14-O-(2-methylbutanoyl), 6,8,9-tri-Ac: **Hippocrateine II**

[132209-62-2]

C₅₁H₅₈N₂O₁₉ 1003.021

Alkaloid from the roots and stem bark of *Hippocratea excelsa* (Hippocrataceae). Mp 308°. λ_{max} 265 (CHCl₃) (Berdy).

8-Epimer, 6,8,14-tri-Ac: **Emarginatine E**

[156280-98-7]

C₃₉H₄₆N₂O₁₇ 814.796

Alkaloid from stems of *Maytenus emarginata* (Celastraceae). Exhibits cytotoxicity against human KB cells. Needles. Mp 240-242°. [α]_D²⁵ -36 (c, 0.07 in CHCl₃). λ_{max} 266 (ε 13400) (MeOH) (Berdy).

8-Epimer, 1-benzoyl, 6,9,14-tri-Ac:

Emarginatine F

[155944-12-0]

C₄₆H₅₀N₂O₁₈ 918.904

Alkaloid from stems of *Maytenus emarginata* (Celastraceae). Cytotoxic. Needles. Mp 221-223°. [α]_D²⁰ +600 (c,

0.01 in CHCl₃). λ_{max} 266 (ε 12200) (MeOH) (Berdy).

8-Epimer, 9-benzoyl, 1,6,8,14-tetra-Ac:

Emarginatine B

[128718-53-6]

C₄₈H₅₂N₂O₁₉ 960.941

Alkaloid from *Maytenus emarginata*. Cytotoxic agent. Needles. Mp 191-194°. [α]_D²⁵ +26 (c, 0.03 in CHCl₃). λ_{max} 266 (ε 12200) (no solvent reported).

[132209-61-1, 125227-49-8]

Kuo, Y.H. et al., *Heterocycles*, 1989, **29**, 1465 (*Emarginatine A*)

Kuo, Y.-H. et al., *J. Nat. Prod.*, 1990, **53**, 422; 1994, **57**, 263; 1995, **58**, 1103 (*Emarginatines B, F, G, H*)

Mata, R. et al., *J. Nat. Prod.*, 1990, **53**, 1212 (*Hippocrateines*)


Kuo, Y.H. et al., *Phytochemistry*, 1994, **35**, 803 (*Emarginatines C, D, E*)

Furukawa, M. et al., *Phytochemistry*, 2002, **59**, 767-777 (*Hippocratea excelsa* alkaloids)

Li, Y. et al., *Heterocycles*, 2006, **68**, 1241-1247 (*Emarginatine A, cmr*)

Per(deacyl)euojaponin

P-247



C₂₆H₃₅NO₁₂ 553.562

1,2,8,9,14-Penta-Ac: **Neoeuonymine**

[33510-25-7]

C₃₆H₄₅NO₁₇ 763.748

Alkaloid from *Euonymus sieboldianus* (Celastraceae). Cryst. (MeOH). Mp 259-262°. [α]_D²⁰ -11 (c, 0.49 in CHCl₃).

1,6,8,9,14-Penta-Ac: **4-Hydroxy-7-epichuhuanine EV**

C₃₆H₄₅NO₁₇ 763.748

Alkaloid from the roots of *Peritassa campestris*. Amorph. solid. [α]_D²⁰ -37.4 (c, 0.002 in CHCl₃). λ_{max} 201 (log ε 3.92); 223 (log ε 3.61); 265 (log ε 3.24) (MeOH).

1,2,6,8,9,14-Hexa-Ac: **Euonymine†**

[33458-82-1]

C₃₈H₄₇NO₁₈ 805.785

Alkaloid from *Euonymus alatus* and fruits of *Euonymus sieboldianus* (Celastraceae). Shows antitumour promoting activity. Amorph.

8-O-(2-Methylpropanoyl), 1,6,9,14-tetra-Ac: **Paniculatine A**

[906067-33-2]

C₃₈H₄₉NO₁₇ 791.802

Alkaloid from the stems of *Celastrus paniculatus*. Amorph. powder (CHCl₃). Mp 199-201°. [α]_D²⁵ -9.5 (c, 0.6 in


- CHCl₃). λ_{\max} 218 (log ϵ 5.12); 264 (log ϵ 2.81) (MeOH).
- 9-O-(2-Hydroxy-2-methylpropanoyl), 1,6,8-tri-Ac: **Cathedulin K15** [68419-30-7] C₃₆H₄₇NO₁₇ 765.764 Alkaloid from *Catha edulis* (Celastraceae). Cryst. (Et₂O/hexane). Mp 191-194°.
- 9-O-(2-Acetoxy-2-methylpropanoyl), 1,6,8-tri-Ac: **Cathedulin K6** [68419-28-3] C₃₈H₄₉NO₁₈ 807.801 Alkaloid from *Catha edulis* (Celastraceae). Mp 176-180°.
- 9-O-(2-Acetoxy-2-methylpropanoyl), 1,6,8,14-tetra-Ac: **Cathedulin K2** [68419-27-2] C₄₀H₅₁NO₁₉ 849.838 Major alkaloid from *Catha edulis* (Celastraceae). Needles (Et₂O/hexane). Mp 181-184°. $[\alpha]_{\text{D}}^{20}$ -17.8 (c, 1.9 in CHCl₃).
- 9-O-(2-Acetoxy-2-methylpropanoyl), 1,2,6,8,14-penta-Ac: **Cathedulin K1**. **Cathedulin Y1** [68423-03-0] C₄₂H₅₃NO₂₀ 891.875 Alkaloid from *Catha edulis* (Celastraceae). Mp 165-168°.
- 1-O-(2-Methyl-2-butenoyl) (ζ -), 2,6,8,9,14-penta-Ac: **Laevisine A** C₄₁H₅₁NO₁₈ 845.85 Alkaloid from the bark of *Maytenus laevis*. Amorph. solid. Mp 171-173°. $[\alpha]_{\text{D}}^{25}$ +97.3 (c, 0.1 in CHCl₃). λ_{\max} 223 (log ϵ 3.61); 265 (log ϵ 3.24) (EtOH).
- 1-Benzoyl, 2,8,9,14-tetra-Ac: **Euojaponin A** [133740-14-4] C₄₁H₄₇NO₁₇ 825.819 Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Mp 273-275°. $[\alpha]_{\text{D}}^{25}$ +24.67 (c, 0.3 in EtOH).
- 1-Benzoyl, 6,8,9,14-tetra-Ac: **Ebenifoline EI** [143439-10-5] C₄₁H₄₇NO₁₇ 825.819 Alkaloid from stem bark of *Maytenus ebenifolia* (Celastraceae). Amorph. solid (EtOH). Mp 183-185°.
- 1-Benzoyl, 2,6,8,9,14-penta-Ac: **Mayteine** [104736-05-2] C₄₃H₄₉NO₁₈ 867.856 Alkaloid from the roots of *Maytenus guaianensis* and stem bark of *Maytenus ebenifolia* (Celastraceae). Amorph. solid. Mp 172-175°. $[\alpha]_{\text{D}}^{25}$ -9.36 (c, 0.47 in CHCl₃).
- 2-Benzoyl, 1,6,8,9,14-penta-Ac: **Forrestine** [146439-78-3] C₄₃H₄₉NO₁₈ 867.856 Isol. from *Tripterygium forrestii*. Exhibits anti-HIV activity. Cryst. (MeOH). Mp 113-115°.
- 6-Benzoyl, 1,8,9,14-tetra-Ac: **Wilforinine F** C₄₁H₄₇NO₁₇ 825.819 Alkaloid from *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_{\text{D}}$ +18.2 (c, 0.4 in MeOH). λ_{\max} 229 (log ϵ 4.16); 264 (log ϵ 3.54) (MeOH).
- 6-Benzoyl, 1,2,8,9,14-penta-Ac: **Cangorinine EI** [155944-24-4] C₄₃H₄₉NO₁₈ 867.856 Alkaloid from root bark of *Maytenus ilicifolia* (Celastraceae). Amorph. solid. Mp 98-101°. $[\alpha]_{\text{D}}$ -25.4 (c, 0.36 in CHCl₃). C-24 and C-25 config. not defined.
- 9-Benzoyl, 1,2,6,8,14-penta-Ac: **Horridine** [102865-30-5] C₄₃H₄₉NO₁₈ 867.856 Alkaloid from the root bark of *Maytenus horrida* (Celastraceae). Amorph. powder.
- 14-Benzoyl, 1,6,8,9-tetra-Ac: **2-De-O-acetylhyponine C** C₄₁H₄₇NO₁₇ 825.819 Alkaloid from the bark of *Maytenus laevis*. Amorph. powder. $[\alpha]_{\text{D}}$ +4.3 (c, 0.2 in CHCl₃/MeOH).
- 14-Benzoyl, 1,2,6,8,9-penta-Ac: **Hyponine C** [192998-84-8] C₄₃H₄₉NO₁₈ 867.856 Alkaloid from root bark of *Tripterygium hypoglaucum*. Amorph. powder. $[\alpha]_{\text{D}}^{25}$ -15.5 (c, 0.8 in CHCl₃). λ_{\max} 228 (ϵ 15000); 265 (ϵ 3600) (MeOH).
- 1,6-Dibenzoyl, 8,9,14-tri-Ac: **Euojaponin C** [133740-15-5] C₄₆H₄₉NO₁₇ 887.89 Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Mp 172-175°. $[\alpha]_{\text{D}}^{25}$ +11.4 (c, 0.5 in MeOH).
- 1,6-Dibenzoyl, 2,8,9,14-tetra-Ac: **Ebenifoline EII**. **Euoverrine A** [133740-16-6] C₄₈H₅₁NO₁₈ 929.927 Alkaloid from stem bark of *Maytenus ebenifolia* (Celastraceae) and from *Euonymus verrucosides*. Amorph. solid (EtOH). Mp 174-177°. $[\alpha]_{\text{D}}$ -13.2 (c, 2.777 in CHCl₃). The C-24 config. in these two isolates is not clear.
- 1,6-Dibenzoyl, 8-propanoyl, 9,14-di-Ac: **Euophelline** [496865-96-4] C₄₇H₅₁NO₁₇ 901.916 Alkaloid from the root bark of *Euonymus phellomana*. Amorph. powder. Mp 168-169°. $[\alpha]_{\text{D}}^{24}$ +2.2 (c, 0.45 in CHCl₃). λ_{\max} 231 ; 265 (MeOH).
- 1,9-Dibenzoyl, 2,8,14-tri-Ac: **Ebenifoline EV** [150375-28-3] C₄₆H₄₉NO₁₇ 887.89 Alkaloid from stem bark of *Maytenus ebenifolia* (Celastraceae). Amorph. solid. Mp 165-169°. $[\alpha]_{\text{D}}$ -27.5 (c, 0.27 in CHCl₃).
- 1,9-Dibenzoyl, 2,6,8,14-tetra-Ac: **Ebenifoline EIII** [150375-26-1] C₄₈H₅₁NO₁₈ 929.927 Alkaloid from *Euonymus ebenifolia* (Celastraceae). Amorph. solid. Mp 174-177°. $[\alpha]_{\text{D}}$ -25.8 (c, 0.67 in CHCl₃).
- 1,6,9-Tribenzoyl, 2,8,14-tri-Ac: **Celahnine A** [172889-43-9] C₅₃H₅₃NO₁₈ 991.998 Alkaloid from stems of *Celastrus hindsii*. Amorph. Mp > 300°.
- 2-(3-Furancarboxyl), 1,6,8,9,14-penta-Ac: **Hyponine F** C₄₁H₄₇NO₁₉ 857.818 Alkaloid from the root bark of *Tripterygium hypoglaucum*. Amorph. powder. $[\alpha]_{\text{D}}^{25}$ +7 (c, 1 in MeOH). λ_{\max} 227 (log ϵ 4); 260 (log ϵ 3.58) (MeOH).
- 6-(3-Furancarboxyl), 1,2,8,9,14-penta-Ac: **Hyponine A** [192998-39-3] C₄₁H₄₇NO₁₉ 857.818 Alkaloid from root bark of *Tripterygium hypoglaucum*. Exhibits anti-HIV activity. Amorph. powder. $[\alpha]_{\text{D}}^{25}$ -27.5 (c, 0.7 in CHCl₃). λ_{\max} 222 (ϵ 10000); 253 (ϵ 5000) (MeOH).
- 14-(3-Furancarboxyl), 1,2,6,8,9-penta-Ac: **Hyponine B** [192998-73-5] C₄₁H₄₇NO₁₉ 857.818 Alkaloid from root bark of *Tripterygium hypoglaucum*. Exhibits anti-HIV activity. Amorph. powder. $[\alpha]_{\text{D}}^{25}$ -19.5 (c, 1.0 in CHCl₃). λ_{\max} 222 (ϵ 8800); 249 (ϵ 4000) (MeOH).
- 1-(3-Pyridinecarbonyl), 6,8,9,14-tetra-Ac: **Euojaponin M** [133740-19-9] C₄₀H₄₆N₂O₁₇ 826.807 Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Mp 188-190°. $[\alpha]_{\text{D}}^{25}$ +12.57 (c, 0.46 in EtOH).
- 1-(3-Pyridinecarbonyl), 2,6,8,9,14-penta-Ac: **Euojaponin I** [133740-17-7] C₄₂H₄₈N₂O₁₈ 868.844 Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Mp 168-171°. $[\alpha]_{\text{D}}^{25}$ +4.35 (c, 0.6 in EtOH).
- 1-(3-Pyridinecarbonyl), 6-benzoyl, 8,9,14-tri-Ac: **Euojaponin L** [133740-18-8] C₄₅H₄₈N₂O₁₇ 888.877 Alkaloid from the root bark of *Euonymus japonica* (Celastraceae). Mp 195-197°. $[\alpha]_{\text{D}}^{25}$ +11.24 (c, 0.5 in EtOH).
- 1-(3-Pyridinecarbonyl), 6-benzoyl, 2,8,9,14-tetra-Ac: **Paniculatine B** [906067-34-3] C₄₇H₅₀N₂O₁₈ 930.915 Alkaloid from the stems of *Celastrus paniculatus*. Amorph. powder (CHCl₃). Mp 178-181°. $[\alpha]_{\text{D}}^{25}$ -11 (c, 0.8 in CHCl₃). λ_{\max} 221 (log ϵ 4.39); 266 (log ϵ 3.83) (MeOH).
- 2-(3-Pyridinecarbonyl), 8-(2-methylpropanoyl), 1,6,9,14-tetra-Ac: **Angulatamine** [148245-79-8] C₄₄H₅₂N₂O₁₈ 896.897 Constit. of *Catha angulatus* (Celastraceae). Amorph. solid. $[\alpha]_{\text{D}}^{25}$ +13.7 (c, 0.5 in MeOH).
- 2-(3-Pyridinecarbonyl), 6-benzoyl, 1,8,9,14-tetra-Ac: **Hyponine D** C₄₇H₅₀N₂O₁₈ 930.915 Alkaloid from the root bark of *Tripterygium hypoglaucum*. Amorph.

- powder. $[\alpha]_D^{25} +5.2$ (c, 1.1 in MeOH). λ_{\max} 226 (log ϵ 4.39); 264 (log ϵ 3.83) (MeOH).
- 8-(3-Pyridinecarbonyl), 6-(3-furancarbo-
nyl), 1,2,9,14-tetra-Ac: **Hyponine E**
C₄₅H₄₈N₂O₁₉ 920.876
Alkaloid from the root bark of *Tri-
pterygium hypoglaucum*. Amorph. pow-
der. $[\alpha]_D^{25} -4.2$ (c, 1 in MeOH). λ_{\max} 222
(log ϵ 4.25); 263 (log ϵ 3.79) (MeOH).
- 8-(3-Pyridinecarbonyl), 1,2,6,9,14-penta-
Ac: **Wilformine G**. Putterine A
C₄₂H₄₈N₂O₁₈ 868.844
Alkaloid from *Tripterygium wilfordii*
and the roots of *Maytenus putterlick-
oides*. Amorph. powder. $[\alpha]_D +11$ (c,
0.7 in MeOH). λ_{\max} 222 (log ϵ 4.26);
260 (log ϵ 3.8) (MeOH).
- 8-(3-Pyridinecarbonyl), 2-(2-methylpro-
panoyl), 1,6,9,14-tetra-Ac: **Putterine B**
C₄₄H₅₂N₂O₁₈ 896.897
Alkaloid from the roots of *Maytenus
putterlickoides*. Amorph. solid.
- 14-(3-Pyridinecarbonyl), 9-(2-acetoxy-2-
methylpropanoyl), 8-(3,4,5-trimethox-
ybenzoyl), 1,2,6-tri-Ac: **Cathedulin K12**
[68419-29-4]
C₅₄H₆₂N₂O₂₃ 1107.083
Alkaloid from *Catha edulis* (Celastra-
ceae). Cryst. (Et₂O). Mp 268-272°.
- 14-(3-Pyridinecarbonyl), 9-(2-acetoxy-2-
methylpropanoyl), 8-(3,4,5-trimethox-
ybenzoyl), 1-benzoyl, 2-Ac: **Cathedulin
E6**
[61230-23-7]
C₅₇H₆₂N₂O₂₂ 1127.117
Alkaloid from *Catha edulis* (Celastra-
ceae). Amorph. solid.
- 14-(3-Pyridinecarbonyl), 9-(2-acetoxy-2-
methylpropanoyl), 8-(3,4,5-trimethox-
ybenzoyl), 1-benzoyl, 2,6-di-Ac: **Ca-
thedulin E5**
[61230-24-8]
C₅₉H₆₄N₂O₂₃ 1169.154
Alkaloid from *Catha edulis* (Celastra-
ceae). Amorph. solid.
- 1-(1,6-Dihydro-1-methyl-6-oxo-3-pyridi-
necarbonyl), 6,8,9,14-tetra-Ac:
C₄₁H₄₈N₂O₁₈ 856.833
Alkaloid from the stem bark of
Hippocratea excelsa. Powder. Mp 192-
199°. $[\alpha]_D -22.8$ (c, 0.3 in CHCl₃).
 λ_{\max} 268 (log ϵ 4.34) (no solvent
reported).
- 1-(1,6-Dihydro-1-methyl-6-oxo-3-pyridi-
necarbonyl), 2,6,8,9,14-penta-Ac:
C₄₃H₅₀N₂O₁₉ 898.87
Alkaloid from the stem bark of
Hippocratea excelsa. Powder. Mp 154-
178°. $[\alpha]_D -11.6$ (c, 0.7 in CHCl₃).
 λ_{\max} 268 (log ϵ 4.1) (no solvent
reported).
- 8-Ketone: see Per(deacyl)evonine, P-248
- 8-Epimer, 1,2,9,14-tetra-Ac: **Acanthotha-
mine**
[107651-98-9]
C₃₄H₄₃NO₁₆ 721.711
Alkaloid from the stems of
Acanthothamnus aphyllus (Celastra-
ceae). Cryst. (CH₂Cl₂). Mp 287-290°.
- 8-Epimer, 1,6,9,14-tetra-Ac: **Aquifoliunine
EII**

- C₃₄H₄₃NO₁₆ 721.711
Alkaloid from the root bark of *May-
tenus aquifolium* (tea). Amorph. $[\alpha]_D -$
17.2 (c, 3 in CHCl₃).
- 8-Epimer, 1,2,6,9,14-penta-Ac: **Aquifoliu-
nine EIII**
[220751-20-2]
C₃₆H₄₅NO₁₇ 763.748
Alkaloid from *Maytenus aquifolium*
(tea). Amorph. solid. $[\alpha]_D -20.8$ (c, 3.3
in CHCl₃).
- 8-Epimer, 8-benzoyl, 1,2,6,9,14-penta-Ac:
Aquifoliunine EI
C₄₃H₄₉NO₁₈ 867.856
Alkaloid from the root bark of *May-
tenus aquifolium* (tea). Amorph. solid.
 $[\alpha]_D -3.1$ (c, 3.2 in CHCl₃).
- 8-Epimer, 8-(3-pyridinecarbonyl),
1,2,6,9,14-penta-Ac: **Aquifoliunine EIV**
[220751-22-4]
C₄₂H₄₈N₂O₁₈ 868.844
Alkaloid from *Maytenus aquifolium*
(tea). $[\alpha]_D -15.2$ (c, 2.9 in CHCl₃).
- Sugiura, K. *et al.*, *Tet. Lett.*, 1971, 2733
(Euonymine)
- Yamada, K. *et al.*, *Tetrahedron*, 1977, **33**,
1725; 1978, **34**, 1915 (Euonymine,
Neoeuonymine)
- Baxter, R.L. *et al.*, *J.C.S. Perkin 1*, 1979, 2965;
2976; 2982; 2986 (Cathedulins)
- González, A.G. *et al.*, *Heterocycles*, 1986, **24**,
1295 (Horridine)
- Soriano-García, M. *et al.*, *J. Crystallogr.
Spectrosc. Res.*, 1986, **16**, 507; *CA*, **106**,
156743m (cryst struct, Acancothamine)
- De Sousa, J.R. *et al.*, *Phytochemistry*, 1986, **25**,
1776 (Mayteine)
- Sánchez, A.A. *et al.*, *Phytochemistry*, 1986, **25**,
2647 (Acancothamine)
- Han, B.H. *et al.*, *Phytochemistry*, 1990, **29**,
2303 (Euojaponins)
- Itokawa, H. *et al.*, *Heterocycles*, 1992, **34**, 885
(Ebenifolines)
- Chunquan, C. *et al.*, *Phytochemistry*, 1992, **31**,
4391 (Forrestine)
- Itokawa, H. *et al.*, *J.C.S. Perkin 1*, 1993, 1247
(Ebenifolines)
- Jikai, L. *et al.*, *Phytochemistry*, 1993, **32**, 487
(Angulatamine)
- Shirota, O. *et al.*, *Heterocycles*, 1994, **38**, 2219
(Cangorinine)
- Kuo, Y.H. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1735-
1738 (Celahinine A)
- Duan, H. *et al.*, *Phytochemistry*, 1997, **45**, 617-
621; 1999, **52**, 1735-1738 (Hyponines)
- Corsino, J. *et al.*, *Phytochemistry*, 1998, **48**,
137-140; **49**, 2181-2183 (Aquifoliunines)
- Piacente, S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 161-
163 (Laevisine A)
- Gonzalez, A.G. *et al.*, *Bioorg. Med. Chem.*,
2000, **8**, 1773-1778 (activity, Euonymine)
- Duan, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 357
(isol, activity)
- Duan, H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 582-
587 (Wilformine F,G)
- Schaneberg, B.T. *et al.*, *J. Nat. Prod.*, 2001, **64**,
624-626 (Putterines)
- Liao, L.M. *et al.*, *Phytochemistry*, 2001, **58**,
1205-1207 (4-Hydroxy-7-epichuchuluanine
EV)
- Furukawa, M. *et al.*, *Phytochemistry*, 2002, **59**,
767-777 (Hippocratea excelsa alkaloids)
- Jinbo, Z. *et al.*, *Phytochemistry*, 2002, **61**, 699-
704 (Euoverrine A, Euophelline)
- Nakagawa, H. *et al.*, *J. Nat. Prod.*, 2004, **67**,
1919-1924 (2-Deacetylhyponine C)
- Lu, Y. *et al.*, *Heterocycles*, 2006, **68**, 1241-1247
(Euonymine, Wilformine F, Paniculatin
A,B)

Per(deacyl)evonine

P-248


O¹,O²,O⁶,O⁹,O⁵-Pentadeacylevonine, 9CI
[33376-79-3]

- C₂₆H₃₃NO₁₂ 551.546
1,2,14-Tri-Ac: **Evozine**
[35721-62-1]
C₃₂H₃₉NO₁₅ 677.658
Alkaloid from the seeds of *Euonymus
europaeus* (Celastraceae). Cryst.
(MeOH/Et₂O). Mp 288-290°. $[\alpha]_D^{22} +13$
(c, 0.92 in CHCl₃).
- 1,9,14-Tri-Ac: **O²,O⁶-Dideacetyllevonine**
[36072-68-1]
C₃₂H₃₉NO₁₅ 677.658
Minor alkaloid from the seeds of
Euonymus europaeus (Celastraceae).
Cryst. (EtOH). Mp 141°.
- 2,9,14-Tri-Ac: **1,6-Dideacetyllevonine**
C₃₂H₃₉NO₁₅ 677.658
Alkaloid from the seeds of *Euonymus
europaeus*. Oil. $[\alpha]_D^{25} +20$ (c, 0.13 in
CHCl₃).
- 1,2,9,14-Tetra-Ac: **Evorine**. Neoeuonine
[33540-08-8]
C₃₄H₄₁NO₁₆ 719.695
Alkaloid from *Euonymus alatus*, the
fruits of *Euonymus sieboldianus* and
the seeds of *Euonymus europaeus* (Cela-
straceae). Cryst. (MeOH, Et₂O/petrol-
or petrol/Me₂CO). Mp 266-268°
(257-259°). $[\alpha]_D^{22} +20$ (c, 1.11 in
CHCl₃). $[\alpha]_D^{25} +41$ (c, 1.13 in MeOH).
- 1,6,9,14-Tetra-Ac: **O²-Deacetyllevonine**
[36017-57-9]
C₃₄H₄₁NO₁₆ 719.695
Minor alkaloid from the seeds of
Euonymus europaeus (Celastraceae).
Cryst. (EtOH). Mp 135°.
- 1,2,4,9,14-Penta-Ac: **Evonine, 9CI**
[33458-64-9]
C₃₆H₄₃NO₁₇ 761.732
Alkaloid from *Euonymus alatus*, *Eu-
onymus europaeus* and *Euonymus sie-
boldianus* (Celastraceae). Cryst. (EtOH
or petrol/Me₂CO). Mp 184-190°. $[\alpha]_D -$
8.4 (c, 1.5 in CHCl₃). $[\alpha]_D +21.1$ (c,
1.46 in EtOH).
- 4-Deoxy, 1,6,9,14-tetra-Ac: **Evonoline. 4-
Desoxyevonine**
[36017-50-2]
C₃₆H₄₃NO₁₆ 745.733
Alkaloid from the seeds of *Euonymus
europaeus* (Celastraceae). Cryst. (petrol/
Me₂CO). Mp 150-158°. $[\alpha]_D^{20} +6$ (c,
3.2 in CHCl₃).
- Klásek, A. *et al.*, *Helv. Chim. Acta*, 1971, **54**,
2144 (isol, uv, ir, ms)

- Pailer, M. et al., *Monatsh. Chem.*, 1971, **102**, 1873 (isol, uv, ir, pmr, ms, struct, synth)
- Wada, H. et al., *Tet. Lett.*, 1971, 3131 (stereochem)
- Sasaki, K. et al., *J.C.S. Perkin 2*, 1972, 1268 (Evonine, Evorine, *cryst struct, abs config*)
- Klásek, A. et al., *Tet. Lett.*, 1972, 941 (Evorine, Dideacetylevonine, pmr, struct)
- Budzikiewicz, H. et al., *Z. Naturforsch., B*, 1972, **27**, 800 (Evonine, Evonoline, Evorine, isol, uv, ir, pmr, ms, struct)
- Dubravkova, L. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 2132 (Evonine, Evonoline, isol, uv, ord, ir, pmr, ms)
- Crombie, L. et al., *Phytochemistry*, 1973, **12**, 703 (Evonine, Dideacetylevonine, O²-Deacetylevonine)
- Shizuri, Y. et al., *Tetrahedron*, 1973, **29**, 1773 (Evonine, Evorine, isol, uv, ir, pmr, struct)
- Sugiura, K. et al., *Chem. Lett.*, 1975, 579 (synth)
- Popik, P. et al., *Pharmacol. Rev.*, 1995, **47**, 235 (rev, pharmacol, history)
- Descoins, C. et al., *Chem. Pharm. Bull.*, 2002, **50**, 199-202 (1,6-Dideacetylevonine)

Per(deacyl)hypoglaunine

P-249

C₂₆H₃₅NO₁₃ 569.561

1,6,8,9,14-Penta-Ac: Wilfordlongine

C₃₆H₄₅NO₁₈ 779.747Alkaloid from *Tripterygium wilfordii*. Mp 179-181°. λ_{max} 233 (log ε 3.92); 278 (log ε 3.59) (EtOH).

1,2,6,8,9,14-Hexa-Ac: Wilfordinine B

C₃₈H₄₇NO₁₉ 821.785Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D²⁵ +43.5 (c, 1.1 in MeOH). λ_{max} 262 (log ε 3.39) (MeOH).

1-Benzoyl, 2,6,8,9,14-penta-Ac: Wilfordinine C

C₄₃H₄₉NO₁₉ 883.855Alkaloid from *Tripterygium wilfordii*. Amorph. solid. [α]_D²⁵ +39.1 (c, 1 in MeOH). λ_{max} 228 (log ε 4.11); 263 (log ε 3.5) (MeOH).

2-Benzoyl, 1,6,8,9,14-penta-Ac: Hypoglaunine C. Wilfordsine

[220750-99-2]

[171090-83-8]

C₄₃H₄₉NO₁₉ 883.855Alkaloid from *Tripterygium hypoglaucum* and *Tripterygium wilfordii*. Amorph. powder. [α]_D²⁵ +56.7 (c, 1 in MeOH). λ_{max} 230 (log ε 4.26); 265 (log ε 3.65) (MeOH).

2,6-Dibenzoyl, 1,8,9,14-tetra-Ac: Wilfor-

dinine I

C₄₈H₅₁NO₁₉ 945.926Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D +63 (c, 0.2 in MeOH). λ_{max} 230 (log ε 4.53); 262 (log ε 3.77) (MeOH).

2-O-(2-Furancarboxyl), 1,6,8,9,14-penta-Ac: Hypoglaunine B

[220750-98-1]

C₄₁H₄₇NO₂₀ 873.817Alkaloid from *Tripterygium hypoglaucum*. Amorph. powder. [α]_D²⁵ +46.2 (c, 0.8 in MeOH). λ_{max} 262 (log ε 3.82) (MeOH).

2-O-(3-Furancarboxyl), 1,6,8,9,14-penta-Ac: Isowilfortrine

[159028-40-7]

C₄₁H₄₇NO₂₀ 873.817Alkaloid from *Tripterygium wilfordii*. Cryst. Mp 329-331°.

8-O-(3-Furancarboxyl), 1,6,9,14-tetra-Ac: Wilfordconine

C₃₉H₄₅NO₁₉ 831.78Alkaloid from *Tripterygium wilfordii*. Cryst. (CH₂Cl₂/MeOH). Mp 192-193°. MF and struct. do not match in ref. λ_{max} 228 (ε 8760); 258 (ε 48010) (EtOH).

14-O-(3-Furancarboxyl), 1,2,6,8,9-penta-Ac: Hypoglaunine A. Hypoglaunine

[220750-97-0]

C₄₁H₄₇NO₂₀ 873.817Alkaloid from *Tripterygium hypoglaucum*. Amorph. powder. [α]_D²⁵ +42.8 (c, 1 in MeOH). λ_{max} 256 (log ε 3.86) (MeOH).

23-Deoxy, 1,2,8,9,14-penta-Ac: Tripfordine C

C₃₆H₄₅NO₁₇ 763.748Alkaloid from the roots of *Tripterygium wilfordii*. Amorph. powder. Mp 137-141°. [α]_D²⁵ -5.8 (c, 0.02 in MeOH). λ_{max} 237 (log ε 2.93) (MeOH).

23-Deoxy, 1,6,8,9,14-penta-Ac: Wilfordinine A

C₃₆H₄₅NO₁₇ 763.748Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D²⁵ -11 (c, 0.9 in MeOH). λ_{max} 220 (log ε 3.89); 263 (log ε 3.5) (MeOH).

23-Deoxy, 2,6,8,9,14-penta-Ac: Wilfordinine J

C₃₆H₄₅NO₁₇ 763.748Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D -8.1 (c, 0.7 in MeOH). λ_{max} 222 (log ε 3.85); 263 (log ε 3.46) (MeOH).

23-Deoxy, 1,2,6,8,9,14-hexa-Ac: Peritassine A

[150881-01-9]

C₃₈H₄₇NO₁₈ 805.785Alkaloid from stems and bark of *Peritassa compta* (Celastraceae). Cryst. Mp 116-117°. [α]_D +24.6 (c, 0.7 in CHCl₃). The abs. config. at C-22/C-23 in the peritassines need not be the same as in the hypoglaunines.

23-Deoxy, 8-O-(2-methylpropanoyl), 1,2,6,9,14-penta-Ac:

C₄₀H₅₁NO₁₈ 833.839Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 162-164°. [α]_D -36.8 (c, 0.5 in CHCl₃). λ_{max} 266 (log ε 3.46) (no solvent reported).

23-Deoxy, 8-benzoyl, 1,2,6,9,14-penta-Ac: Peritassine B

[150881-29-1]

C₄₃H₄₉NO₁₈ 867.856Alkaloid from stems and bark of *Peritassa compta* (Celastraceae). Amorph. powder. Mp 148-150°. [α]_D -39.2 (c, 0.13 in CHCl₃).

23-Deoxy, 14-O-(2-furancarboxyl), 1,2,6,8,9-penta-Ac: Hypoglaunine D

[220751-00-8]

C₄₁H₄₇NO₁₉ 857.818Alkaloid from *Tripterygium hypoglaucum*. Amorph. powder. [α]_D²⁵ -15.3 (c, 0.9 in MeOH). λ_{max} 226 (log ε 3.82); 264 (log ε 3.42) (MeOH).

23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 8-O-(2-methylpropanoyl), 1,6,9,14-tetra-Ac:

C₄₅H₅₄N₂O₁₉ 926.924Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 189-196°. [α]_D +2.3 (c, 1.1 in CHCl₃). λ_{max} 269 (log ε 3.97) (no solvent reported).

23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 8,14-bis(2-methylpropanoyl), 1,6-di-Ac:

C₄₅H₅₆N₂O₁₈ 912.94Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 171-176°. [α]_D +12.5 (c, 0.73 in CHCl₃). λ_{max} 270 (log ε 4.44) (no solvent reported).

23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 8,14-bis(2-methylpropanoyl), 1,6,9-tri-Ac:

C₄₇H₅₈N₂O₁₉ 954.977Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 179-187°. [α]_D +6.6 (c, 1.1 in CHCl₃). λ_{max} 269 (log ε 4.39) (no solvent reported).

23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 1,9,14-tris(2-methylpropanoyl), 6,8-di-Ac:

C₄₉H₆₂N₂O₁₉ 983.031Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 206-210°. [α]_D -16 (c, 1.1 in CHCl₃). λ_{max} 268 (log ε 4.35) (no solvent reported).

23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 8,9,14-tris(2-methylpropanoyl), 6-Ac:

C₄₇H₆₀N₂O₁₈ 940.994Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 189-194°. [α]_D -21.7 (c, 0.9 in CHCl₃). λ_{max} 269 (log ε 4.38) (no solvent reported).


23-Deoxy, 2-O-(1,6-dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 8,9,14-tris(2-methylpropanoyl), 1,6-di-Ac:

C₄₉H₆₂N₂O₁₉ 983.031Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 186-190°. [α]_D -18.6 (c, 0.86 in CHCl₃). λ_{max} 268 (log ε 3.83) (no solvent reported).Klass, J. et al., *J. Nat. Prod.*, 1993, **56**, 946-948 (Peritassines)

- Lin, S. et al., *Yaoxue Xuebao*, 1994, **29**, 599-602; 1995, **30**, 513-516 (*Isowilfordine*, *Wilfordine*)
 Duan, H. et al., *Phytochemistry*, 1998, **49**, 2185-2189 (*Hypoglaunines*)
 Li, W. et al., *Phytochemistry*, 1999, **50**, 1091-1093 (*isol*, *pmr*, *cmr*)
 Duan, H. et al., *J. Nat. Prod.*, 2000, **63**, 357-361; 2001, **64**, 582-587 (*Wilfordines*)
 Lin, S. et al., *Yaoxue Xuebao*, 2001, **36**, 116-119; 2002, **37**, 128-130 (*Wilfordconine*, *Wilfordlongine*)
 Furukawa, M. et al., *Phytochemistry*, 2002, **59**, 767-777 (*Hippocratea excelsa* alkaloids)
 Horiuchi, M. et al., *J. Nat. Prod.*, 2006, **69**, 1271-1274 (*Tripfordine C*)

Per(deacyl)isowilfordine

P-250

C₂₆H₃₅NO₁₃ 569.5611,6,8,9,14-Penta-Ac: *Wilfordsuine*C₃₆H₄₅NO₁₈ 779.747

Alkaloid from *Tripterygium wilfordii*.
 Mp 173-175°. [α]_D²⁵ -3.8 (c, 1.1 in Me₂CO). λ_{max} 234 (log ε 3.94); 278 (log ε 3.6) (EtOH).

2-Benzoyl, 1,6,8,9,14-penta-Ac: *Isowilfordine*

[134306-16-4]

C₄₃H₄₉NO₁₉ 883.855

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Cryst. (MeOH). Mp 165-167°.

2-(3-Furancarboxonyl), 1,6,8,9,14-penta-Ac: *Wilfordinine H*C₄₁H₄₇NO₂₀ 873.817

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Amorph. powder. [α]_D²⁵ +1.4 (c, 0.7 in MeOH). λ_{max} 227 (log ε 4.1); 262 (log ε 3.57) (MeOH).

8-Ketone, 1,2,6,9,14-penta-Ac: *Wilfordinine G*C₃₆H₄₃NO₁₈ 777.732

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Amorph. powder. [α]_D²⁵ +4.3 (c, 1.1 in MeOH). λ_{max} 221 (log ε 3.89); 263 (log ε 3.47) (MeOH).

26-Deoxy, 1,2,6,8,9,14-penta-Ac: *Wilfordinine E*C₃₈H₄₇NO₁₈ 805.785

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Amorph. powder. [α]_D²⁵ -0.6 (c, 0.7 in MeOH). λ_{max} 225 (log ε 3.98); 264 (log ε 3.5) (MeOH).

26-Deoxy, 2-benzoyl, 1,6,8,9,14-penta-Ac: *Wilfordinine F*C₄₃H₄₉NO₁₈ 867.856

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Amorph. powder. [α]_D²⁵

+16.2 (c, 0.9 in MeOH). λ_{max} 230 (log ε 4.32); 265 (log ε 3.55) (MeOH).

26-Deoxy, 2-(3-furancarboxonyl),


1,6,8,9,14-penta-Ac: *Wilfordinine D*C₄₁H₄₇NO₁₉ 857.818

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Amorph. powder. [α]_D²⁵ +14.6 (c, 0.4 in MeOH). λ_{max} 227 (log ε 4.06); 261 (log ε 3.57) (MeOH).

Ya, L. et al., *Phytochemistry*, 1991, **30**, 719 (*Isowilfordine*)Duan, H. et al., *Chem. Pharm. Bull.*, 1999, **47**, 1664-1667 (*Wilfordines*)Lin, S. et al., *Zhixu Xuebao (Acta Bot. Sin.)*, 2001, **43**, 647-649 (*Wilfordsuine*)

Per(deacyl)oppositine A

P-251

C₂₇H₃₇NO₁₂ 567.589

The ethyl group config. need not necessarily be the same in all of these alkaloids.

1-Benzoyl, 2,6,8,9,14-penta-Ac: *Oppositine A*

[917952-11-5]

C₄₄H₅₁NO₁₈ 881.883

Alkaloid from *Pleurostylyla opposita*.
 Pale yellow glass. [α]_D²⁶ +20.3 (c, 0.13 in MeOH). λ_{max} 206 (log ε 4.08); 230 (log ε 4.14); 270 (log ε 3.47) (MeOH).

8-Ketone, 1,2,6,9,14-penta-Ac: *Orthosphenine*

[126622-36-4]

C₃₇H₄₅NO₁₇ 775.759

Minor alkaloid from the roots of *Orthosphenia mexicana* (Celastraceae).
 Pale-yellow amorph. solid.

4-Deoxy, 1-benzoyl, 2,6,8,9,14-penta-Ac: *Oppositine B*

[917952-12-6]

C₄₄H₅₁NO₁₇ 865.883

Alkaloid from *Pleurostylyla opposita*.
 Pale yellow glass. [α]_D²⁶ +19.5 (c, 0.07 in MeOH). λ_{max} 204 (log ε 4.27); 228 (log ε 4.35); 268 (log ε 3.68) (MeOH).

4-Deoxy, 6-benzoyl, 1,2,8,9,14-penta-Ac: *Cassinine*

[62948-58-7]


C₄₄H₅₁NO₁₇ 865.883

Alkaloid from the root bark of *Cassine metabelica*. Mp 295-298°.

Wagner, H. et al., *Tet. Lett.*, 1977, **18**, 125-128 (*Cassinine*)González, A.G. et al., *J. Nat. Prod.*, 1989, **52**, 1338 (*Orthosphenine*)Whitson, E.L. et al., *J. Nat. Prod.*, 2006, **69**, 1833-1835 (*Oppositines*)

Per(deacyl)wilforine

P-252

C₂₆H₃₅NO₁₂ 553.562

▶ LF0195000

1,2,6,9,14-Penta-Ac: 8-O-Deacetylleuonine

C₃₆H₄₅NO₁₇ 763.748

Constit. of the roots of *Austroplenckia populnea* var. *ovata*. Amorph. solid (CHCl₃/MeOH). Mp 153-155°.

1,2,8,9,14-Penta-Ac: 6-O-Deacetylleuonine. *Tripfordine B*C₃₆H₄₅NO₁₇ 763.748

Alkaloid from the roots of *Tripterygium wilfordii*. Amorph. solid. Mp 132-133°. [α]_D²⁵ -15.4 (c, 0.13 in MeOH). λ_{max} 267 (log ε 3.55) (MeOH).

1,6,8,9,14-Penta-Ac: *Wilforjine*. 2-O-Deacetylleuonine

[147362-40-1]

C₃₆H₄₅NO₁₇ 763.748

Alkaloid from the roots of *Austroplenckia populnea* var. *ovata* and *Tripterygium wilfordii*. Mp 156-158° (147-149°).

1,2,6,8,9,14-Hexa-Ac: *Euonine*. *Wilforimine*

[41758-69-4]

C₃₈H₄₇NO₁₈ 805.785

Minor alkaloid from the seeds of *Euonymus sieboldianus* and *Tripterygium wilfordii* (Celastraceae). Mp 149-153°. [α]_D -2.5 (c, 6.4 in CHCl₃).

1-Benzoyl, 2,8,9,14-tetra-Ac: *Euojaponin D*

[128397-42-2]

C₄₁H₄₇NO₁₇ 825.819

Alkaloid from *Euonymus japonica* (Celastraceae). Mp 253°. [α]_D²⁵ +28.6 (c, 0.287 in CHCl₃).

1-Benzoyl, 6,8,9,14-tetra-Ac: *Euojaponin K*

[128397-44-4]

C₄₁H₄₇NO₁₇ 825.819

Alkaloid from root bark of *Euonymus japonica* (Celastraceae). Powder. Mp 188°. [α]_D²⁵ +26.2 (c, 0.37 in CHCl₃).

1-Benzoyl, 2,6,8,9,14-penta-Ac: *Euojaponin F*

[128397-43-3]

C₄₃H₄₉NO₁₈ 867.856

Alkaloid from root bark of *Euonymus japonica* (Celastraceae). Powder. Mp 142°. [α]_D²⁵ +9 (c, 0.3 in CHCl₃).

2-Benzoyl, 1,8,9,14-tetra-Ac: *Wilforzine*

[37239-46-6]

C₄₁H₄₇NO₁₇ 825.819

Alkaloid from the roots of *Tripterygium wilfordii* (Celastraceae). Needles (Me₂CO/MeOH). Mp 177-178°. [α]_D²⁵ +6 (Me₂CO).

2-Benzoyl, 1,6,8,9,14-penta-Ac:

Wilfortrine[†]

[11088-09-8]

C₄₃H₄₉NO₁₈ 867.856

Alkaloid from *Tripterygium wilfordii* and *Maytenus senegalensis* (Celastraceae). Shows insecticidal props. Plates (Me₂CO/MeOH). Mp 169-170°. [α]_D²⁵ +30 (Me₂CO).

1,2-Dibenzoyl, 8,9,14-tri-Ac: **Ebenifoline WII**

[150384-28-4]

C₄₆H₄₉NO₁₇ 887.89

Alkaloid from stem bark of *Maytenus ebenifolia* (Celastraceae). Amorph. solid. Mp 118-122°. [α]_D +61.1 (c, 0.18 in CHCl₃).

1,2-Dibenzoyl, 6,8,9,14-tetra-Ac: **Ebenifoline WI**

[143439-09-2]

C₄₈H₅₁NO₁₈ 929.927

Alkaloid from stem bark of *Maytenus ebenifolia* and *Peritassa compta* (Celastraceae). Amorph. solid (EtOH). Mp 222-224°. [α]_D +47.1 (c, 0.29 in CHCl₃).

2,6-Dibenzoyl, 1,8,9,14-tetra-Ac: **Cangorinine WI**

[155964-94-6]

C₄₈H₅₁NO₁₈ 929.927

Alkaloid from root bark of *Maytenus ilicifolia* (Celastraceae). Amorph. solid. Mp 149-151°. [α]_D +14.4 (c, 0.57 in CHCl₃).

2-O-(3-Furancarboxyl), 1,6,8,9,14-penta-Ac: **Wilforgine**

[37239-47-7]

C₄₁H₄₇NO₁₉ 857.818

Alkaloid from the roots of *Tripterygium hypoglaucum* (Celastraceae). Shows insecticidal props. Plates (Me₂CO/MeOH). Mp 211°. [α]_D²⁵ +25 (Me₂CO).

▶ LF0195500

1-O-(3-Pyridinecarbonyl), 2,6,8,9,14-penta-Ac: **Laevisine B**C₄₂H₄₈N₂O₁₈ 868.844

Alkaloid from the bark of *Maytenus laevis*. Amorph. solid. Mp 148-150°. [α]_D²⁵ +31.2 (c, 0.1 in CHCl₃). λ_{max} 220 (log ε 4.2); 262 (log ε 3.74) (EtOH).

2-O-(3-Pyridinecarbonyl), 1,6,8,9,14-penta-Ac: **Wilfortrine**. 2-Debenzoyl-2-nicotinoylwilfortrine

[112899-84-0]

C₄₂H₄₈N₂O₁₈ 868.844

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Immunosuppressant. Mp 177-178°.

6-O-(3-Pyridinecarbonyl), 1-benzoyl, 2,8,9,14-tetra-Ac: **Euojaponin G**

[128397-45-5]

C₄₇H₅₀N₂O₁₈ 930.915

Constit. of *Euonymus japonica*.

6-O-(3-Pyridinecarbonyl), 2-benzoyl, 1,8,9,14-tetra-Ac: **Cangorinine WII**

[155944-25-5]

C₄₇H₅₀N₂O₁₈ 930.915

Alkaloid from bark of *Maytenus ilicifolia* (Celastraceae). Amorph. solid. Mp 153-158°. [α]_D +7.9 (c, 0.24 in CHCl₃).

2-O-(1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarbonyl), 1,6,8,9,14-penta-Ac: C₄₃H₅₀N₂O₁₉ 898.87

Alkaloid from the stem bark of *Hippocratea excelsa*. Powder. Mp 178-182°. [α]_D +23.6 (c, 1 in CHCl₃). λ_{max} 269 (log ε 4.3) (no solvent reported).

8-Ketone, 1,2,6,9,14-penta-Ac: **Evonimine**.**Isoevonine**

[41758-54-7]

C₃₆H₄₃NO₁₇ 761.732

Alkaloid from the seeds of *Euonymus europaeus* and *Euonymus sieboldianus* (Celastraceae). Amorph. [α]_D +21 (c, 1.5 in CHCl₃). [α]_D +30.5 (c, 0.9 in EtOH).

9-Deoxy, 8-ketone, 6,14-di-Ac: **Austronine**C₃₀H₃₇NO₁₃ 619.621

Alkaloid from the roots of *Austroplenckia populnea* var. *ovata*. Amorph. solid (CHCl₃/MeOH). Mp 169-172°.


8-Epimer, 1-benzoyl, 2,6,8,9,14-penta-Ac: **Euoverrine B**C₄₃H₄₉NO₁₈ 867.856

Alkaloid from the root barks of *Euonymus fortunei* and *Euonymus verucosides*. Amorph. powder. Mp 148-149°. [α]_D²⁴ +10.9 (c, 0.55 in CHCl₃). λ_{max} 232 ; 266 (MeOH).

Beroza, M. et al., *J.A.C.S.*, 1951, **73**, 3656;1952, **74**, 1585; 1953, **75**, 2136 (*isol. uv.**Wilfortrine*)Tin-Wa, M. et al., *J. Nat. Prod.*, 1971, **34**, 79 (*isol*)Dubrakova, L. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 2132 (*Evonimine*)Sugiura, K. et al., *Tet. Lett.*, 1973, 113 (*Evonimine, Euonine*)Zhang, Z. et al., *Jieqou Huaxue*, 1986, **5**, 83;*CA*, **107**, 237078x (*Wilforgine*)He, Z. et al., *Huaxue Xuebao*, 1987, **45**, 510;*CA*, **107**, 130906pDeng, F. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1987, **29**, 523; *CA*, **108**, 87738s (*Wilfortrine*)Han, B.H. et al., *Arch. Pharmacol. Res.*, 1989, **12**, 306-309; *CA*, **113**, 55832e (*Euojaponins*)Ya, L. et al., *Can. J. Chem.*, 1990, **68**, 371 (*Wilfortrine*)Han, B.H. et al., *J. Nat. Prod.*, 1990, **53**, 909 (*Euojaponins*)Shi, J. et al., *Chin. Sci. Bull.*, 1991, **36**, 1266-1270; *CA*, **116**, 255847n (*Euonine, crystal*)Itokawa, H. et al., *Heterocycles*, 1992, **34**, 885 (*Ebenifoline WI*)Deng, F. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1992, **34**, 618-621 (*Wilfortrine*)Klass, J. et al., *J. Nat. Prod.*, 1993, **56**, 946 (*Ebenifoline WI*)Itokawa, H. et al., *J.C.S. Perkin 1*, 1993, 1247 (*Ebenifoline WII*)Shirota, O. et al., *Heterocycles*, 1994, **38**, 383 (*Cangorinines*)Piacente, S. et al., *J. Nat. Prod.*, 1999, **62**, 161-163 (*Laevisine B*)Furukawa, M. et al., *Phytochemistry*, 2002, **59**, 767-777 (*Hippocratea excelsa alkaloid*)Jinbo, Z. et al., *Phytochemistry*, 2002, **61**, 699-704 (*Euoverrine B*)De Sousa, J.R. et al., *J. Nat. Prod.*, 2006, **69**, 1225-1227 (*Austronine, 2- and 8-Deacetylauonine*)Horiuchi, M. et al., *J. Nat. Prod.*, 2006, **69**, 1271-1274 (*Tripfordine B*)

Per(deacyl)wilfortrine

P-253

C₂₆H₃₅NO₁₃ 569.561

CAS numbering shown; some authors' schemes differ.

▶ LF0195550

1,6,8,9,14-Penta-Ac: **Wilfordine**. *Tripfordine A*

[98618-76-9]

C₃₆H₄₅NO₁₈ 779.747

Alkaloid from *Tripterygium wilfordii*. Amorph. solid. Mp 172-174°. [α]_D²⁵ +4.5 (c, 0.03 in MeOH). λ_{max} 260 (log ε 2.91) (MeOH).

1,2,6,8,9,14-Hexa-Ac: **Alatusinine**

[89945-79-9]

C₃₈H₄₇NO₁₉ 821.785

Alkaloid from the fruits of *Euonymus alatus* (Celastraceae). Amorph. solid. [α]_D²⁶ -16 (c, 0.6 in CHCl₃).

2-Benzoyl, 6,8,9,14-tetra-Ac: **1-Deacyl-wilfordine**

[128666-73-9]

C₄₁H₄₇NO₁₈ 841.818

Alkaloid from *Tripterygium wilfordii* (Celastraceae). Mp 182-184°.

2-Benzoyl, 1,6,8,9,14-penta-Ac: **Wilfordine**

[37239-51-3]

C₄₃H₄₉NO₁₉ 883.855

Alkaloid from *Tripterygium wilfordii* and *Euonymus alatus* (Celastraceae). Shows insecticidal props. Cryst. (Me₂CO/MeOH). Mp 170-176°. [α]_D²² +5 (c, 5.0 in CHCl₃). λ_{max} 232 (ε 22600); 271 (ε 4400) (MeOH) (Berdy).

6-Benzoyl, 1,8,9,14-tetra-Ac: **5-Benzoyl-5-deacetylwilfordine**

[183008-61-9]

C₄₁H₄₇NO₁₈ 841.818

Alkaloid from stemwood of *Maytenus buchananii*. Cryst. (EtOAc). Mp 187-189°.

26-Benzoyl, 1,2,8,9,14-penta-Ac: **Wilfortrine B**C₄₃H₄₉NO₁₉ 883.855


Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D -17.8 (c, 0.8 in

- MeOH). λ_{\max} 228 (log ϵ 4.17); 265 (log ϵ 3.6) (MeOH).
- 26-Benzoyl, 1,2,6,8,9,14-hexa-Ac: Wilfor-nine A**
 $C_{45}H_{51}NO_{20}$ 925.893
 Alkaloid from *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{20}$ -41.9 (c, 1 in MeOH). λ_{\max} 228 (log ϵ 4.27); 267 (log ϵ 3.68) (MeOH).
- 1,2-Dibenzoyl, 6,8,9,14-tetra-Ac: Chiapenine ES I**
 $C_{48}H_{51}NO_{19}$ 945.926
 Alkaloid from the leaves of *Maytenus chiapensis*. Lacquer. $[\alpha]_D^{20}$ +15 (c, 1.3 in $CHCl_3$). λ_{\max} 201 (log ϵ 4.57); 229 (log ϵ 4.5); 268 (log ϵ 3.82) (EtOH).
- 6,26-Dibenzoyl, 1,2,8,9,14-penta-Ac: Wilfor-nine C**
 $C_{50}H_{53}NO_{20}$ 987.963
 Alkaloid from *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{20}$ -50 (c, 1.1 in MeOH). λ_{\max} 231 (log ϵ 4.51); 267 (log ϵ 3.74) (MeOH).
- 2-O-(3-Furancarboxyl), 6,8,9,14-tetra-Ac: 1-Desacetylwilfortrine**
 [128638-35-7]
 $C_{39}H_{45}NO_{19}$ 831.78
 Alkaloid from *Tripterygium wilfordii* (Celastraceae). Mp 179-182°.
- 2-O-(3-Furancarboxyl), 1,6,8,9,14-penta-Ac: Wilfortrine**
 [37239-48-8]
 $C_{41}H_{47}NO_{20}$ 873.817
 Alkaloid from the roots of *Tripterygium wilfordii* (Celastraceae). Shows insecticidal props. Plates ($Me_2CO/MeOH$). Mp 237.5-238°. $[\alpha]_D^{25}$ +10 (Me_2CO).
- 2-O-(3-Furancarboxyl), 26-O-benzoyl, 1,6,8,9,14-penta-Ac: Triptonine A†. 18-O-Benzylwilfortrine**
 [168009-84-5]
 $C_{48}H_{51}NO_{21}$ 977.925
 Alkaloid from root bark of *Tripterygium wilfordii* (Celastraceae). Needles (EtOH). Mp 163-164°. $[\alpha]_D^{27}$ -36 (c, 0.21 in Me_2CO). λ_{\max} 224; 264 (MeOH) (Berdy).
- 26-(3-Furancarboxyl), 1,2,6,8,9,14-hexa-Ac: Wilfor-nine D**
 $C_{43}H_{49}NO_{21}$ 915.854
 Alkaloid from *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{20}$ -21.1 (c, 0.8 in MeOH).
- 2,26-Bis-O-(3-furancarboxyl), 1,6,8,9,14-penta-Ac: Triptonine B†. 18-O-(3-Fur-oyl)wilfortrine**
 [168009-85-6]
 $C_{46}H_{49}NO_{22}$ 967.887
 Alkaloid from root bark of *Tripterygium wilfordii* (Celastraceae). Needles (EtOH). Mp 160-161°. $[\alpha]_D^{30}$ -13 (c, 0.20 in Me_2CO).
- 2-(1,6-Dihydro-1-methyl-6-oxo-3-pyridi-necarbonyl), 1,6,8,9,14-penta-Ac: Emarginatinine**
 [156280-99-8]
 $C_{43}H_{50}N_2O_{20}$ 914.869
 Alkaloid from *Maytenus emarginata*. Cytotoxic agent. Needles. Mp 238-241°. $[\alpha]_D^{20}$ -130 (c, 0.3 in $CHCl_3$). λ_{\max}

- 266 (ϵ 12600) (MeOH).
- 8-Ketone, 1,6,9,14-tetra-Ac: Chiapenine ES IV**
 $C_{34}H_{41}NO_{17}$ 735.694
 Alkaloid from the leaves of *Maytenus chiapensis*. Lacquer. $[\alpha]_D^{20}$ +7.9 (c, 0.43 in $CHCl_3$). λ_{\max} 201 (log ϵ 4.11); 220 (log ϵ 3.89); 264 (log ϵ 3.52) (EtOH).
- 8-Ketone, 1,2,6,9,14-penta-Ac: Wilfor-nine E**
 $C_{36}H_{43}NO_{18}$ 777.732
 Alkaloid from *Tripterygium wilfordii*. Amorph. powder. $[\alpha]_D^{20}$ +1.1 (c, 0.5 in MeOH). λ_{\max} 222 (log ϵ 3.82); 267 (log ϵ 3.45) (MeOH).
- 8-Ketone, 1-benzoyl, 6,9,14-tri-Ac: Chiapenine ES III**
 $C_{39}H_{43}NO_{17}$ 797.765
 Alkaloid from the leaves of *Maytenus chiapensis*. Lacquer. $[\alpha]_D^{20}$ +29.3 (c, 0.41 in $CHCl_3$). λ_{\max} 201 (log ϵ 4.11); 230 (log ϵ 3.98); 269 (log ϵ 3.37) (EtOH).
- 8-Ketone, 2-benzoyl, 1,9,14-tri-Ac: Neoal-atamine**
 [89945-78-8]
 $C_{39}H_{43}NO_{17}$ 797.765
 Alkaloid from the fruits of *Euonymus alata* forma *striatus* (Celastraceae). Amorph. solid. $[\alpha]_D^{26}$ +42 (c, 1.1 in $CHCl_3$).
- 8-Ketone, 2-benzoyl, 1,6,9,14-tetra-Ac: Alatamine**
 [41855-33-8]
 $C_{41}H_{45}NO_{18}$ 839.802
 Alkaloid from *Euonymus alatus* (Celastraceae). Cryst. (MeOH). Mp 185-193°. $[\alpha]_D^{25}$ +44 (c, 1.08 in $CHCl_3$).
- 8-Ketone, 1,2-dibenzoyl, 6,9,14-tri-Ac: Chiapenine ES II**
 $C_{46}H_{47}NO_{18}$ 901.873
 Alkaloid from the leaves of *Maytenus chiapensis*. Lacquer. $[\alpha]_D^{20}$ +60.9 (c, 4.71 in $CHCl_3$). λ_{\max} 201 (log ϵ 4.5); 230 (log ϵ 4.5); 269 (log ϵ 3.8) (EtOH).
- 26-Deoxy: see Per(deacyl)wilforine, P-252**
- Beroza, M. *et al.*, *J.A.C.S.*, 1951, **73**, 3656; 1952, **74**, 1585 (*Wilfordine, Wilfortrine, isol, ir, uv*)
- Shizuri, Y. *et al.*, *Tet. Lett.*, 1973, 741 (*Wilfordine*)
- Yamada, K. *et al.*, *Tetrahedron*, 1978, **34**, 1915 (*Alatamine*)
- Ishiwata, H. *et al.*, *Phytochemistry*, 1983, **22**, 2839 (*Alatusinine, Neoalatamine*)
- He, Z. *et al.*, *Huaxue Xuebao*, 1985, **43**, 593; *CA*, **103**, 157321y (*Wilfordidine*)
- He, Z. *et al.*, *Huaxue Xuebao*, 1989, **47**, 178; *CA*, **111**, 74780m
- Ya, L. *et al.*, *Can. J. Chem.*, 1990, **68**, 371 (*1-Deacetylwilfortrine, 1-Deacetylwilfordine*)
- Kuo, Y.H. *et al.*, *Phytochemistry*, 1994, **35**, 803-807 (*Emarginatinine*)
- Morota, T. *et al.*, *Phytochemistry*, 1995, **39**, 1219 (*Triptonines*)
- Sekar, K.V.S. *et al.*, *Planta Med.*, 1996, **62**, 368-370 (*5-Benzoyl-5-deacetylwilfordine*)
- Duan, H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 582-587 (*Wilforines*)
- Nunez, M.J. *et al.*, *J. Nat. Prod.*, 2004, **67**, 14-18 (*Chiapenines*)
- Horiuch, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1271-1274 (*Wilfordine*)

Perfamine

P-254
 4,8-Dimethoxy-8-(3-methyl-2-butenyl)-furo[2,3-b]quinolin-7(8H)-one, 9CI



$C_{18}H_{19}NO_4$ 313.352

(R)-form [59557-95-8]

Alkaloid from *Haplophyllum perforatum* and *Haplophyllum glabrinum* (Rutaceae). Prisms (hexane/ Me_2CO). Mp 175-178° (164-165°). $[\alpha]_D^{26.5}$ -20 (c, 1.00 in $CHCl_3$). $[\alpha]_D^{25}$ +53.4.

5,6-Dihydro: Dihydroperfamine

[105705-87-1]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid from the roots of *Haplophyllum glabrinum* (Rutaceae). Prisms (hexane/ Me_2CO). Mp 179-183°. $[\alpha]_{H_2O}^{26.5}$ -640 (c, 1.00 in $CHCl_3$).

(S)-form**5,6-Dihydro: (+)-Dihydroperfamine**

[141902-17-2]

$C_{18}H_{21}NO_4$ 315.368

Alkaloid from *Haplophyllum tuberculatum* (Rutaceae).

(E)-form**5-Hydroxy: Sarcomegistine**

[161068-67-3]

$C_{18}H_{19}NO_5$ 329.352

Alkaloid from aerial parts of *Sarcomelicope megistophylla*.

Razakova, D.M. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 812; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 828 (*isol*)

Grina, J.A. *et al.*, *J.O.C.*, 1982, **47**, 2648 (*struct*)


Rózsa, Zs. *et al.*, *Phytochemistry*, 1986, **25**, 2005 (*isol, uv, pmr, cmr, ms, deriv*)

Al-yahah, M.A. *et al.*, *Int. J. Pharm.*, 1991, **29**, 268; *CA*, **117**, 23293w (*(+)-Dihydroperfamine*)

Skaltsounis, A.L. *et al.*, *Nat. Prod. Lett.*, 1995, **5**, 281 (*Sarcomegistine*)

Perfragilin A

P-255
 7-Amino-2-methyl-6-(methylthio)-3,5,8(2H)-isoquinolinetrione, 9CI
 [129722-94-7]



$C_{11}H_{10}N_2O_3S$ 250.278

Alkaloid from the bryozoan *Membranipora perfragilis*. Cytotoxic agent. Red needles. Mp 219-220°. λ_{\max} 212 (ϵ 12616); 248 (ϵ 7000); 330 (ϵ 15140); 362

(ϵ 8785); 440 (ϵ 2224) (EtOH) (Berdy).

Deamino: 2-Methyl-6-(methylthio)-3,5,8(2H)-isoquinolinetrione

$C_{11}H_9NO_3S$ 235.263

Isol. from bryozoan *Biflustra perfragilis*. Yellow powder. λ_{max} 367 (ϵ 10594) (no solvent reported).

7-Deamino, 7-(methylthio): 2-Methyl-6,7-bis(methylthio)-3,5,8(2H)-isoquinolinetrione, 9CI. Perfragilin B

[146764-79-6]

$C_{12}H_{11}NO_3S_2$ 281.356

Alkaloid from *Membranipora perfragilis* and *Biflustra perfragilis*. Cytotoxic (considerably more potent than Perfragilin A). Red needles or orange plates. Mp 163° Mp 170-172°. Also obt. as a metastable green powder. λ_{max} 214 (ϵ 14941); 235 (ϵ 10504); 332 (ϵ 8227); 382 (ϵ 5946); 465 (ϵ 1848) (EtOH) (Berdy).

Rizvi, S.K. *et al.*, *Acta Cryst. C*, 1993, **49**, 151 (*cryst struct*)


Blackman, A.J. *et al.*, *Aust. J. Chem.*, 1993, **46**, 213-220 (7-deamino, 7-deamino-7-methylthio, *isol, cryst struct*)

Choi, Y.-H. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1431-1433 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ferreira, V.F. *et al.*, *Tetrahedron*, 2003, **59**, 1349-1357 (*synth*)

Perhentinine

P-256



$C_{43}H_{52}N_4O_5$ 704.908


Alkaloid from the bark of *Alstonia macrophylla*. Cytotoxic. Light yellow oil. $[\alpha]_D$ -61 (c, 1.2 in $CHCl_3$). λ_{max} 231 (log ϵ 4.25); 298 (log ϵ 3.45) (EtOH).

Kam, T.-S. *et al.*, *Tetrahedron*, 2004, **60**, 3957-3966 (*isol, pmr, cmr, ms*)

Pericidine

P-257

[926927-39-1]



Absolute Configuration

$C_{19}H_{20}N_2O_2$ 308.379


Indole alkaloid prob. most closely related to the Apparicine group. Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D$ +85 (c, 0.12 in $CHCl_3$). λ_{max} 223 (log ϵ 4.4); 283 (log ϵ 3.76) (EtOH).

Lim, K.-H. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 31-35 (*isol, pmr, cmr, ms*)

Periconin B

P-258

[145569-96-6]



$C_{20}H_{29}Cl_3N_4O_8$ 559.829


Prod. by the fungal pathogen *Periconia circinata*. λ_{max} 215 (ϵ 1300) (H_2O).

Macko, V. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 9574-9578 (*isol, uv, cd, ir, pmr, cmr*)

Perimargine

P-259

[64667-16-9]



R undetermined. MW of R group 149, total MW 423. Alkaloid from the leaves of *Peripterygia marginata* (Celastraceae). Amorph. Obt. only in admixture with Dihydroperimargine.

Dihydro: Dihydroperimargine

Alkaloid from the leaves of *Peripterygia marginata* (Celastraceae). Amorph. MW 425.

Hocquemiller, R. *et al.*, *Tetrahedron*, 1977, **33**, 645-651 (*isol, uv, ir, pmr, ms*)

Perimivine

P-260

[1360-37-8]

$C_{21}H_{22}N_2O_4$ 366.416

Vobasine alkaloid. Struct. unknown. Alkaloid from the roots of *Catharanthus lanceus* and *Catharanthus roseus* (Apocynaceae). Needles (MeOH), blades ($CHCl_3$). Mp 292-293° dec. $[\alpha]_D^{25}$ -98.71 (c, 1 in $CHCl_3$). Isomer of Perivine, P-268.


Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1964, **27**, 203-213 (*isol, ir*)

Blomster, R.N. *et al.*, *J. Nat. Prod.*, 1964, **27**, 480-485 (*isol*)

Perinadine A

P-261

[866924-03-0]



Absolute Configuration

$C_{28}H_{37}NO_7$ 499.603

Prod. by a marine-derived *Penicillium citrinum* strain N055. Amorph. solid.

$[\alpha]_D^{22}$ -33 (c, 1 in $CHCl_3$). λ_{max} 215 (ϵ 16000); 252 (ϵ 3600); 315 (ϵ 1700) (MeOH).


Sasaki, M. *et al.*, *Org. Lett.*, 2005, **7**, 4261-4264 (*isol, pmr, cmr, ms*)

Peripentadine

P-262

N-[3-[2-[2-(2-Hydroxy-6-methylphenyl)-2-oxoethyl]-1-pyrrolidinyl]propyl]-hexanamide, 9CI

[87201-37-4]



$C_{22}H_{34}N_2O_3$ 374.522

Major alkaloid from the bark and leaves of *Peripentadenia mearsii* (Elaeocarpaceae).

N-Dehexanoyl, N-butanoyl: Dinorperipentadine

[100595-84-4]

$C_{20}H_{30}N_2O_3$ 346.469

Minor alkaloid from the bark of *Peripentadenia mearsii* (Elaeocarpaceae). Yellow gum.

Lamberton, J.A. *et al.*, *J. Nat. Prod.*, 1983, **46**, 235-247 (*isol, uv, ir, pmr, ms, struct*)

Bick, I.R.C. *et al.*, *Tetrahedron*, 1985, **41**, 5627-5631 (*Dinorperipentadine, cmr*)


Michael, J.P. *et al.*, *Tetrahedron*, 1996, **52**, 2199-2216 (*Peripentadine, Dinorperipentadine, synth*)

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1946-1950 (*isol, pmr, cmr*)

Peripentonine A

P-263

[960511-38-0]



$C_{22}H_{36}N_2O_3$ 376.538

Tautomeric. Alkaloid from the leaves of *Peripentadenia mearsii*. Yellow gum (as trifluoroacetate). $[\alpha]_D^{23}$ +19 (c, 0.13 in MeOH) (trifluoroacetate). Isol. as a mixt. with Peripentonine B to which data refers. λ_{max} 226 (log ϵ 3.45); 339 (log ϵ 2.81) (MeOH) (trifluoroacetate).

2-Epimer: Peripentonine B

[960511-39-1]


$C_{22}H_{36}N_2O_3$ 376.538

Alkaloid from the leaves of *Peripentadenia mearsii*.

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1946-1950 (*isol, pmr, cmr, ms*)

Peripentonine C**P-264**

[960511-40-4]


C₁₅H₂₈N₂O₃ 284.398

Alkaloid from the leaves of *Peripentadenia mearnsii*. Yellow gum (as trifluoroacetate). $[\alpha]_D^{23}$ -5.8 (c, 0.07 in MeOH) (trifluoroacetate). λ_{\max} 204 (log ϵ 2.62); 269 (log ϵ 2.05); 339 (log ϵ 1.73) (MeOH) (trifluoroacetate).

Katavic, P.L. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1946-1950 (*isol*, *pmr*, *cmr*)

Periphylline**P-265**

9-(1-Oxo-3-phenyl-2-propenyl)-4-phenyl-1,5,9-triazacyclotridec-12-en-2-one, 9CI. 9-Cinnamoyl-4-phenyl-1,5,9-triazacyclotridec-12-en-2-one [11082-66-9]

C₂₅H₂₉N₃O₂ 403.523

Alkaloid from leaves of *Peripterygia marginata* (Celastraceae). Cryst. (MeOH or Me₂CO). Mp 165°. $[\alpha]_D^{20}$ -291 (c, 1 in CHCl₃).

2,3-Dihydro: Dihydroperiphylline

[65094-32-8]

C₂₅H₃₁N₃O₂ 405.539

Alkaloid from the leaves of *Peripterygia marginata* (Celastraceae). $[\alpha]_D^{23}$ +3.5 (c, 0.4 in CHCl₃) (synthetic). The originally reported opt. rotn. value (-21°) for natural material is erroneous.

(Z)-Cinnamoyl isomer: Isoperiphylline

[11082-67-0]

C₂₅H₂₉N₃O₂ 403.523

Alkaloid from the leaves of *Peripterygia marginata* (Celastraceae). Cryst. (Me₂CO). Mp 197°. $[\alpha]_D^{20}$ -120 (c, 1 in CHCl₃).

2Z-Isomer: Neoperiphylline

[65094-31-7]

C₂₅H₂₉N₃O₂ 403.523

Alkaloid from the leaves of *Peripterygia marginata* (Celastraceae). Amorph. $[\alpha]_D^{20}$ -34 (c, 0.5 in CHCl₃). Struct. revised in 2003.

Hocquemiller, R. *et al.*, *Tetrahedron*, 1977, **33**, 645-651; 653-656 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Wasserman, H.H. *et al.*, *J.A.C.S.*, 1981, **103**, 461-462 (*Dihydroperiphylline*, *synth*, *ir*, *pmr*)

Kaseda, T. *et al.*, *Tet. Lett.*, 1989, **30**, 4539-4542 (*Dihydroperiphylline*, *synth*)

Begley, M.J. *et al.*, *J.C.S. Perkin 1*, 1993, 2027-2046 (*Dihydroperiphylline*, *synth*)


Ishihara, K. *et al.*, *Synlett*, 1995, 41-42 (*Dihydroperiphylline*, *synth*)

Matsuyama, H. *et al.*, *Chem. Lett.*, 2000, 1104-1105 (*Dihydroperiphylline*, *synth*)

Sergeyev, S.A. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 161-167; 2003, **86**, 465-473 (*Dihydroperiphylline*, *Neoperiphylline*, *struct*, *synth*)

Peritoxin A**P-266**

[145585-98-4]

C₂₀H₂₉Cl₃N₄O₉ 575.829

Prod. by the fungal pathogen *Periconia circinata*. Mycotoxin. λ_{\max} 215 (ε 4400) (H₂O).

4-Deoxy: Periconin A

[145586-00-1]

C₂₀H₂₉Cl₃N₄O₈ 559.829

Prod. by *Periconia circinata*. λ_{\max} 215 (ε 1500) (H₂O). λ_{\max} 215 (ε 11100) (H₂O) (Berdy).

7'-Deoxy: Peritoxin B

[145585-99-5]

C₂₀H₂₉Cl₃N₄O₈ 559.829

Prod. by *Periconia circinata*. Mycotoxin. λ_{\max} 215 (ε 2100) (H₂O).

Macko, V. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, **89**, 9574-9578 (*isol*, *uv*, *ir*, *cd*, *pmr*, *cmr*)
Dunkle, L.D. *et al.*, *Can. J. Bot.*, Suppl. 1, 1995, **73**, 444-452 (*rev*)

Perividine**P-267**

[1360-38-9]

C₂₀H₂₂N₂O₄ 354.405


Poss. a 2-acylindole alkaloid. Struct. unknown. Minor alkaloid from *Catharanthus roseus* (Apocynaceae). Blades (MeOH). Mp 271-279° dec. λ_{\max} 240 (ε 17800); 286 (ε 29600); 315 (ε 15600) (EtOH) (Berdy).

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 243-246 (*isol*, *ir*, *uv*)

Weisbach, J.A. *et al.*, *J. Nat. Prod.*, 1964, **27**, 374-388

Perivine**P-268**

Methyl 4-demethyl-3-oxovobasan-17-oate, 9CI [2673-40-7]



Absolute Configuration

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Catharanthus roseus*, *Catharanthus longifolius*, *Gabunia eglan-dulosa* and several *Tabernaemontana* spp. Antineoplastic agent. Mod. cytotoxic agent. Local anaesthetic potentiator. Active against gram-positive bacteria. Mp 180-181°. $[\alpha]_D$ -121 (c, 1 in CHCl₃).

Log P 1.97 (calc). λ_{\max} 314 (E1%/1cm 2.67) (MeOH) (Berdy).

▶ ZB1740000

N⁴-Formyl: **Periformyline**. Methyl 3,22-dioxovobasan-17-oate, 9CI

[2779-18-2]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from *Catharanthus lanceus*, *Catharanthus roseus* and *Catharanthus trichophyllus*. Cryst. (MeOH). Mp 206-209° dec. λ_{\max} 240 (ε 16600); 315 (ε 20800) (EtOH) (Berdy).

N⁴-Me: Vobasine

[2134-83-0]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Voaacanga africana* and a wide range of spp. in the Apocynaceae. Weak CNS depressant, also showing analgesic and antipyretic action. Mp 111-113°. $[\alpha]_D^{23}$ -158 (c, 1 in CHCl₃).

N⁴-Me; hydrochloride: Mp 245-248° dec. $[\alpha]_D^{22}$ -120 (c, 1 in MeOH).

N⁴-Me, N⁴-oxide: Vobasine N⁴-oxide

[51666-18-3]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the leaves and stem bark of *Ervatamia polyneura* (Apocynaceae). $[\alpha]_D$ -130 (c, 1 in MeOH).

N¹, N⁴-Di-Me: Ochropamine

[2134-97-6]

C₂₂H₂₆N₂O₃ 366.459

Alkaloid from *Cabucala erythrocarpa*, *Striolaria* sp. and the stem bark of *Ochrosia poweri*. Rhombs (MeOH aq.). Mp 134°. $[\alpha]_D^{22}$ -158 (c, 1 in Me₂CO). λ_{\max} 243 ; 315 (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 39 mg/kg.**N¹, N⁴-Di-Me, picrate:**

Bright yellow plates (Me₂CO/EtOH). Mp 225° dec.

11-Methoxy, N¹, N⁴-di-Me: Ochropine

[2299-25-4]

C₂₃H₂₈N₂O₄ 396.485

Alkaloid from the stem bark of *Ochrosia poweri* (Apocynaceae). Rhombs (MeOH aq.). Mp 146°. $[\alpha]_D^{22}$ -229 (c, 1 in Me₂CO).

11-Methoxy, N¹, N⁴-di-Me, picrate:

Orange-yellow rhombs (Me₂CO/EtOH). Mp 235° dec.

16-Epimer, N⁴-Me: 16-Epivobasine

[6836-00-6]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from leaves and stems of *Ervatamia polyneura* (Apocynaceae). Mp 175-178°. $[\alpha]_D$ -50 (c, 0.66 in CHCl₃).

16-Epimer, parent acid, N⁴-Me: 16-Epivobasinic acid

[13281-24-8]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Tabernaemontana psychotrifolia* (Apocynaceae). Needles (MeOH). Mp 195° dec.

19,20-Dihydro, N⁴-Me: see Dregamine, D-935


Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1958, **47**, 834 (*isol*, *uv*, *ir*)

Budzikiewicz, H. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 1899 (ms)

- Renner, U. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 2186 (*Vobasine, isol, uv, ir, ms, struct*)
 Cava, M.P. *et al.*, *Tet. Lett.*, 1963, 53 (*pmr*)
 Douglas, B. *et al.*, *Aust. J. Chem.*, 1964, **17**, 246 (*Ochropamine, Ochropine*)
 Jaggi, H. *et al.*, *Chimia*, 1964, **18**, 173 (*Vobasine, cryst struct*)
 Gorman, M. *et al.*, *Tet. Lett.*, 1964, 3105 (*uv, pmr, struct*)
 Maloney, E.M. *et al.*, *J. Pharm. Sci.*, 1965, **54**, 1166 (*Periformylone, isol, uv, ir, ms, struct, synth*)
 Abraham, D.J. *et al.*, *Tet. Lett.*, 1965, 317 (*Periformylone, struct*)
 Battersby, A.R. *et al.*, *Chem. Comm.*, 1966, 810; 812; 888; 890 (*biosynth*)
 Burnell, R.H. *et al.*, *Can. J. Chem.*, 1971, **49**, 307 (*16-Epivobasinic acid*)
 Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 929 (*Periformylone, Vobasine, Ochropamine, cd, uv, abs config*)
 Taylor, W.I. *et al.*, *Catharanthus Alkaloids*, Marcel Dekker, 1975, (*rev*)
 Ahond, A. *et al.*, *J.O.C.*, 1976, **41**, 1878 (*Vobasine, Ochropamine, cmr*)
 Kingston, D.G.I. *et al.*, *J. Pharm. Sci.*, 1978, **67**, 272 (*cytotoxicity*)
 Hernandez, N.M.R. *et al.*, *Rev. Cubana Med. Trop.*, 1979, **31**, 199-204 (*activity*)
 Sakai, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 3454 (*Ochropine, synth, abs config*)
 Mukhopadhyay, S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 335 (*Periformylone, isol, uv, ir, pmr, ms*)
 Clivio, P. *et al.*, *Phytochemistry*, 1990, **29**, 3007 (*Vobasine N⁴-oxide, 16-Epivobasine*)
 Pereira, P.S. *et al.*, *Quim. Nova*, 2008, **31**, 20-24 (*Vobasine, pmr, cmr*)

Perloline P-269


6-(3,4-Dimethoxyphenyl)-3,4-dihydro-4-oxobenzo[c][2,7]naphthyridinium (1+), 9CI [7344-94-7]



- $C_{20}H_{17}N_2O_3^{\oplus}$ 333.366
 Alkaloid from *Lolium perenne* (perennial rye grass) (Poaceae). Mp 270° dec. (slow heat)(as perchlorate) Mp 280-290° dec. (rapid heat)(perchlorate). pK_a 8.54 (18°). Forms a covalent hydroxide.
 Jeffreys, J.A.D. *et al.*, *J.C.S.*, 1964, 4504 (*isol, uv, pmr, ir, struct*)
 Ferguson, G. *et al.*, *J.C.S.(B)*, 1966, 454 (*cryst struct*)
 Ridley, A.B. *et al.*, *Aust. J. Chem.*, 1987, **40**, 631 (*synth*)

Perlolirine P-270

5-(9H-Pyrido[3,4-b]indol-1-yl)-2-furanmethanol, 9CI. 2-(β -Carbolin-1-yl)-5-hydroxymethylfuran. *Tribulusterine*. Alkaloid YS [29700-20-7] [234750-74-4]




- $C_{16}H_{12}N_2O_2$ 264.283
 Struct. of *Tribulusterine* shown to be same as *Perlolirine* in 2004. Alkaloid from *Lolium perenne*, *Lolium chuanxiang* and *Tribulus terrestris*. Found in Korean ginseng and Japanese soy sauce. Yellow cryst. Mp 183° (179-181°). λ_{max} 233 (sh); 252 (sh); 288; 366; 375 (MeOH).

Hydrochloride:

- Yellow needles. Mp 204-233° dec.
 Jeffreys, J.A.D. *et al.*, *J.C.S.(C)*, 1970, 1091 (*isol, uv, ir, pmr, ms, cryst struct, synth*)
 Nakatsuka, S. *et al.*, *Tet. Lett.*, 1986, **27**, 3399 (*isol, pmr*)
 Gessner, W.P. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 95 (*synth, uv*)
 Bracher, F. *et al.*, *Annalen*, 1992, 1315 (*synth, ir, pmr, cmr, ms*)
 Bremner, J.B. *et al.*, *Aust. J. Chem.*, 2004, **57**, 273-276 (*synth, pmr, cmr, struct*)

Peronatin B P-271

1,1',2,2'-Tetrahydro-2,2',4,4'-tetramethyl[2,2'-bi-3H-indole]-3,3'-dione, 9CI [157536-39-5]




- $C_{20}H_{20}N_2O_2$ 320.39
 Isol. from injured fruit bodies of *Collybia peronata* (sometimes also present in small amounts in intact specimens). Yellow cryst. Mp 213° dec.
 7-Hydroxy, 7'-methoxy: **7-Hydroxy-7'-methoxyperonatin B** [157201-78-0]
 $C_{21}H_{22}N_2O_4$ 366.416
 Isol. from injured fruit bodies of *Tricholoma scalpturatum* (not present in intact specimens). Yellow solid. Mp 203-204°.
 7,7'-Dimethoxy: **7,7'-Dimethoxyperonatin B** [157201-79-1]
 $C_{22}H_{24}N_2O_4$ 380.443
 From injured fruit bodies of *Tricholoma scalpturatum* (not present in intact specimens). Yellow solid. Mp 224-225°.

2'-Epimer: Peronatin A

- [157536-38-4]
 $C_{20}H_{20}N_2O_2$ 320.39
 From injured fruit bodies of *Collybia peronata* (sometimes also present in small amounts in intact specimens). Yellow oil.
 Pang, Z. *et al.*, *J. Nat. Prod.*, 1994, **57**, 852-857 (*isol, uv, ir, pmr, cmr, struct*)
 Stachel, S.J. *et al.*, *J.O.C.*, 1997, **62**, 4756-4762 (*synth*)

Perphoramidine P-272

[474779-75-4]




- $C_{21}H_{17}BrCl_2N_4$ 476.202
 Alkaloid from the Philippine ascidian *Perophora namei*. Cytotoxic. Amorph. off-white solid. λ_{max} 220 (ϵ 13300); 240 (sh) (ϵ 10300); 298 (sh) (ϵ 5800); 308 (ϵ 6800); 338 (ϵ 4600) (MeOH).

- Verbitski, S.M. *et al.*, *J.O.C.*, 2002, **67**, 7124-7126 (*isol, uv, cd, pmr, cmr, ms*)
 Fuchs, J.R. *et al.*, *J.A.C.S.*, 2004, **126**, 5068-5069 (*synth*)

Persine P-273

Struct. unknown. Alkaloid from the roots of *Catharanthus roseus* (Apocynaceae). Blades (MeOH)(as sulfate salt). Mp 219-225° (sulfate).

- Svoboda, G. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141 (*isol, uv, ir*)

Perspicamide A P-274

- $C_{18}H_{14}N_2O_4$ 322.32
 Isol. from *Botrylloides perspicum*. Pale yellow solid. λ_{max} 230 (ϵ 15830); 246 (ϵ 16000); 314 (ϵ 6620); 349 (ϵ 7460) (MeOH).


Z-Isomer: Perspicamide B

- $C_{18}H_{14}N_2O_4$ 322.32
 Isol. from *Botrylloides perspicum*. Pale yellow solid. λ_{max} 221 (ϵ 17725); 246 (ϵ 16645); 315 (ϵ 6800); 347 (ϵ 7680) (MeOH).

- McKay, M.J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1776-1778 (*isol, pmr, cmr, ms*)

Peruvianine P-275

9-Hydroxy-1,2-dimethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI [78416-87-2]



- $C_{18}H_{13}NO_4$ 307.305


Alkaloid from the woody stems of *Telitoxicum peruvianum* (Menispermaceae). Red-orange prisms (MeOH). Mp 252-255°.

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 320 (*isol, uv, pmr, struct*)

Buck, K.T. *et al.*, *Heterocycles*, 1993, **36**, 2489 (*synth*)

Peshawarine P-276

7-[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]-6,7-dihydro-9H-1,3-dioxolo[4,5-h][2]benzopyran-9-one, 9CI



(S)-form

C₂₁H₂₁NO₆ 383.4

(S)-form [61224-21-3]

Alkaloid from *Hypecoum parviflorum* (Papaveraceae). Cryst. (MeOH). Mp 190-191°. [α]_D²⁵ -109 (c, 0.2 in MeOH).

(±)-form [63950-18-5]

Synthetic. Cryst. (MeOH). Mp 196-198° (182-183°).

N-Me:

Cryst. (MeOH)(as iodide). Mp 210° (iodide).

Šimánek, V. *et al.*, *Heterocycles*, 1977, **6**, 711 (*synth, ms*)


Shamma, M. *et al.*, *Tetrahedron*, 1978, **34**, 635 (*isol, uv, cd, ir, pmr, ms, struct, synth*)

Chrzanowska, M. *et al.*, *Tetrahedron*, 1986, **42**, 6021 (*synth, ir, pmr, ms*)

Fitzgerald, J.J. *et al.*, *J.O.C.*, 1994, **59**, 4117 (*synth, pmr, cmr*)

Petasinenine P-277

Fukinotoxin
[60102-37-6]



C₁₉H₂₇NO₇ 381.425

Cyclic ester of Otonecine, O-136 and Petasineic acid. Stereoisomer of Otosenine, O-137. Alkaloid from *Petasites japonicus* (sweet coltsfoot) (Asteraceae). Cryst. (C₆H₆/hexane or Me₂CO). Mp 129-130°. [α]_D +63.8 (CHCl₃).

►Carcinogen, hepatotoxin. SE5085000

Ac: Neopetasinenine

[60409-51-0]

[61303-14-8]

C₂₁H₂₉NO₈ 423.462

Alkaloid from *Petasites japonicus* (sweet coltsfoot) (Asteraceae).

Amorph. solid; cryst. (as hydrochloride). Mp 212° (as hydrochloride). [α]_D¹⁹ +49 (c, 1.19 in EtOH).

[60132-19-6 , 60409-53-2]

Yamada, K. *et al.*, *Chem. Lett.*, 1976, 461 (*isol, struct*)


Furuya, T. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 1120 (*isol, ir, pmr, cd, cryst struct*)

Iwao, H. *et al.*, *J. Natl. Cancer Inst.*, 1977, **58**, 1155; *CA*, **87**, 976u (*tox*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PCQ750

Petchicine P-278

[125547-41-3]



Absolute Configuration


C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the stem bark of *Petchia ceylanica* (Apocynaceae). Amorph. solid. [α]_D -380 (c, 0.1M MeOH).

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1989, **28**, 3221 (*isol, uv, ir, pmr, cmr, ms, struct*)

Petididine P-279

21-(3,4-Dihydro-3-methyl-2H-pyrrol-5-yl)-3-hydroxy-20-methylpregnane-6,21-dione, 9CI. 3-Hydroxy-23,26-epiminocholest-23(N)-ene-6,22-dione [79805-74-6]



C₂₇H₄₁NO₃ 427.626

Alkaloid from *Petilium raddeana* and bulbs of *Fritillaria persica* (Liliaceae). Cryst. (Me₂CO). Mp 150-152°. [α]_D²⁹ +18.4 (c, 0.50 in CHCl₃).

O-β-D-Glucopyranoside: [145867-02-3]

C₃₃H₅₁NO₈ 589.768
From bulbs of *Fritillaria persica* (Liliaceae). Amorph. powder + 1H₂O. [α]_D²⁹ -19.6 (c, 0.50 in MeOH).

3-Ketone: Petididinone

[107316-97-2]

C₂₇H₃₉NO₃ 425.61

Alkaloid from aerial parts of *Petilium raddeana* (Liliaceae). Cryst. (hexane/Me₂CO). Mp 217-219°. [α]_D 0 (c, 0.169 in CHCl₃).

20-Epimer: [145920-88-3]

C₂₇H₄₁NO₃ 427.626

Isol. from the bulbs of *Fritillaria persica* (Liliaceae). Amorph. powder. [α]_D²⁹ +1.6 (c, 0.50 in CHCl₃).

20-Epimer, O-β-D-glucopyranoside:

[145867-03-4]

C₃₃H₅₁NO₈ 589.768

From bulbs of *Fritillaria persica* (Liliaceae). Needles + 2H₂O (MeOH). Mp 214-215°. [α]_D²⁹ -13.2 (c, 0.50 in MeOH).

Nabiev, A. *et al.*, *Khim. Prir. Soedin.*, 1982, 528-529; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **18**, 502-503 (*isol, uv, ms, pmr, struct*)


Nabiev, A. *et al.*, *Khim. Prir. Soedin.*, 1986, 620-622; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 583-584 (*Petididinone*)

Ori, K. *et al.*, *Phytochemistry*, 1992, **31**, 4337-4341 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ori, K. *et al.*, *Phytochemistry*, 1992, **31**, 4337-4341 (*isol, uv, ir, pmr, cmr, ms, struct*)

Petidininone P-280

3,23-Dihydroxy-16,28-secosolanida-22,24,26(28)-trien-6-one, 9CI. 3-Hydroxy-20-(5-hydroxy-3-methyl-6-pyridinyl)-pregnan-6-one [89783-63-1]



C₂₇H₃₉NO₃ 425.61

Alkaloid from aerial parts and bulbs of *Petasites raddeana* (Asteraceae). Cryst. (MeOH). Mp 290-292°.

3-Ac: Verdinine

C₂₉H₄₁NO₄ 467.647

Alkaloid from *Veratrum lobelianum*. Cryst. (Me₂CO). Mp 265-267°. Has not been interconverted with Petidininone.

3-Epimer: Fetsininone

[571167-92-5]

C₂₇H₃₉NO₃ 425.61

Alkaloid from the bulbs of *Fritillaria imperialis*. Amorph. powder. [α]_D²⁵ -118 (c, 0.1 in MeOH). λ_{max} 202 (log ε 3.85); 220 (log ε 3.38); 287 (log ε 3.48) (MeOH).

Nakhatov, I. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 710-712 (*Petidininone*)


Shakirov, R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 79-82 (*Verdinine*)

Akhtar, M.N. *et al.*, *Phytochemistry*, 2003, **63**, 115-122 (*Fetsininone*)

Petisine

P-281

3-Hydroxy-16,28-secosolanid-22(28)-ene-6,23-dione, 9Cl. *Petizine* [79805-75-7]



C₂₇H₄₁NO₃ 427.626

Alkaloid from *Petilium raddeana* (Liliaceae). Cryst. (Me₂CO). Mp 221-222°.

3-O-β-D-Glucopyranosyl: *Petisinine*. *Petizinine* [80981-47-1]

C₃₃H₅₁NO₈ 589.768

Alkaloid from *Petilium raddeana* (Liliaceae). Cryst. (MeOH). Mp 232-234°. [α]_D²⁵ -35 (c, 0.48 in CHCl₃).

Nakhatov, I. et al., *Khim. Prir. Soedin.*, 1981, **17**, 616; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 450

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1986, **22**, 345-347; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 319-321 (*config*)

Petomine

P-282

[1360-40-3]

C₁₇H₂₁NO₆ 335.356


Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Narcissus* sp. and *Amaryllis parkeri* (Amaryllidaceae). Cryst. (Me₂CO or MeOH/Et₂O). Mp 253-254° dec. [α]_D²⁵ 0 (c, 0.1 in CHCl₃).

Boit, H.-G. et al., *Chem. Ber.*, 1957, **90**, 2197-2202; 1959, **92**, 2578-2582 (*isol*)

Petrobactin

P-283

[545434-89-7]



C₃₄H₅₀N₆O₁₁ 718.803

Struct. revised in 2003; formerly assigned as bis(2,3-dihydroxybenzoyl). Prod. by *Marinobacter hydrocarbonoclasticus*. Photoreactive siderophore.

2'-Sulfo: *Petrobactin sulfonate*

C₃₄H₅₀N₆O₁₄S 798.867

Prod. by *Marinobacter hydrocarbonoclasticus*. Amorph. solid. [α]_D²⁰ -2.5 (c, 0.013 in DMSO). λ_{max} 229 (log ε 3.95); 253 (log ε 3.83); 291 (log ε 3.74) (DMSO).

Barbeau, K. et al., *J.A.C.S.*, 2002, **124**, 378-379 (*isol, pmr, cmr, ms*)

Bergeron, R.J. et al., *Tetrahedron*, 2003, **59**, 2007-2014 (*synth, struct*)

Hickford, S.I.H. et al., *J. Nat. Prod.*, 2004, **67**, 1897-1899 (*Petrobactin sulfonate*)

Gardner, R.A. et al., *J.O.C.*, 2004, **69**, 3530-3537; 2007, **72**, 3158 (*synth, pmr, cmr*)


Abergel, R.J. et al., *J.A.C.S.*, 2008, **130**, 2124-2125 (*coord chem*)

Koppisch, A.T. et al., *J.O.C.*, 2008, **73**, 5759-5765 (*biosynth*)

Petromindole

P-284

[198992-51-7]



C₂₈H₃₉NO 405.622


Alkaloid from the ascostromata of *Petromyces muricatus*. Plates (Me₂CO). Mp >300°. λ_{max} 226 (log ε 4.24); 282 (log ε 3.65); 293 (log ε 3.56) (dioxan).

Ooike, M. et al., *Chem. Pharm. Bull.*, 1997, **45**, 1694-1696 (*isol, uv, ir, pmr, cmr, cryst struct*)

Petromurin A

P-285

[194608-26-9]



C₃₁H₃₂N₂O₅ 512.604

Isol. from the ascostromata of *Petromyces muricatus*. Microcryst. (CDCl₃). Mp 100° dec. λ_{max} 225 (log ε 4.66); 278 (log ε 4.32) (MeOH).

5'-Deoxy: *Petromurin B*

[194608-28-1]

C₃₁H₃₂N₂O₄ 496.605

Isol. from the ascostromata of *Petromyces muricatus*. Microcryst. (MeOH). Mp 200°. λ_{max} 224 (log ε 4.76); 282 (log ε 4.39) (MeOH).

Ooike, M. et al., *Can. J. Chem.*, 1997, **75**, 625-628 (*isol, uv, ir, pmr, cmr*)

Petromurin C

P-286

[194608-29-2]

As Petromurin A, P-285 with

R = H

C₂₆H₂₄N₂O₅ 444.486

Isol. from the ascostromata of *Petromyces muricatus*. Microcryst. (CDCl₃). Mp 240° dec. λ_{max} 223 (log ε 4.76); 281 (log ε 4.33) (MeOH).

5''-Methoxy: *Petromurin D*

[194608-30-5]

C₂₇H₂₆N₂O₆ 474.512


Isol. from the ascostromata of *Petromyces muricatus*. Microcryst. (CDCl₃). Mp 127-128°. λ_{max} 223 (log ε 4.7); 278 (log ε 4.36) (MeOH).

Ooike, M. et al., *Can. J. Chem.*, 1997, **75**, 625-628 (*isol, uv, ir, pmr, cmr*)

Petromyzonamine

P-287

[871332-07-9]



C₃₄H₆₀N₂O₃ 544.86

7,24-Disulfate: *Petromyzonamine disulfate*

[871543-67-8]

C₃₄H₆₀N₂O₉S₂ 704.988

Pheromone of the sea lamprey (*Petromyzon marinus*).


Sorensen, P.W. et al., *Nat. Chem. Biol.*, 2005, **1**, 324-328 (*isol*)

Hoye, T.R. et al., *J.O.C.*, 2007, **72**, 7544-7550 (*pmr, cmr, struct*)

Petrosamine

P-288

[113403-19-3]



C₂₁H₁₈BrN₃O₂[⊕] 424.296

Minor struct. revision in 2005. Originally descr. as the 5-oxo monocation tautomer. Props. given refer to the chloride, which should presumably be reassigned as dichloride (or chloride-hydrochloride). Alkaloidal pigment from the marine sponge *Petrosia* sp. Dark green cryst. (CH₂Cl₂/MeOH) (as dichloride). Mp 330° (chloride). λ_{max} 284 (ε 32000); 345 (ε 10900); 574 (ε 4700) (H₂O). λ_{max} 289 (ε 42600); 346 (ε 12400); 414 (sh) (ε 6900); 595 (ε 5300) (MeOH).

Debromo, 3-bromo: *Petrosamine B*

C₂₁H₁₈BrN₃O₂[⊕] 424.296


Alkaloid from the Australian sponge *Oceanapia* sp. Weak inhibitor of aspartyl semialdehyde dehydrogenase. Blue solid (as bis(trifluoroacetate)). λ_{max} 205 (ε 34610); 288 (ε 30460); 378 (ε 15845); 609 (ε 4385) (MeOH) (bis(trifluoroacetate)).

Molinski, T.F. et al., *J.O.C.*, 1988, **53**, 1340-1341 (*isol, uv, ir, pmr, cmr, cryst struct*)

Carroll, A.R. et al., *J. Nat. Prod.*, 2005, **68**, 804-806 (*tautom, Petrosamine B*)

Petrosaspongiolide L P-289

[194148-99-7]




$C_{24}H_{35}NO_2$ 369.546
Alkaloid from the sponge *Petrosaspongia nigra*. Cytotoxic agent. $[\alpha]_D^{25}$ -33.3 (c, 0.001 in MeOH). λ_{max} 210 (ε 5545); 266 (ε 2375) (no solvent reported).

N-(Carboxymethyl): *Spongidine C*
 $C_{26}H_{38}NO_4^{\oplus}$ 428.591
Alkaloid from a *Spongia* sp. Inhibits human phospholipase A₂. $[\alpha]_D^{25}$ -10 (c, 0.01 in MeOH). Counterion not specified. CAS no. not found 14CI.

Paloma, L.G. *et al.*, *Tetrahedron*, 1997, **53**, 10451-10458 (isol, uv, pmr, cmr)
De Marino, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 323-326 (*Spongidine C*)

Petrosine P-290

[84679-41-4]



$C_{30}H_{50}N_2O_2$ 470.737
Alkaloid from the Papua-New Guinean marine sponge *Petrosia seriata* and from an Okinawan sponge *Xestospongia* sp. Ichthyotoxin. Mp 215-216°. Petrosine C and Petrosine D are synthetic diastereoisomers of Petrosin A and B.

1,9-Diepimer: *Petrosine B*
[95189-04-1]
 $C_{30}H_{50}N_2O_2$ 470.737
From *Petrosia seriata*. Ichthyotoxin. Sol. MeOH, CHCl₃; poorly sol. H₂O. $[\alpha]_D^{25}$ -12 (c, 0.79 in CH₂Cl₂).


1,3,9,10-Tetraepimer: *Petrosine A*
[95189-03-0]
 $C_{30}H_{50}N_2O_2$ 470.737
From *Petrosia seriata* and the *Xestospongia* sp. Ichthyotoxin. Sol. MeOH, CHCl₃; poorly sol. H₂O. Struct. revised in 1988. Originally described as the 1,1',9,9'-tetraepimer. Opt. inactive (*meso* compd.). The low opt. rotn. originally reported refers to a sample contaminated with traces of Petrosine B.

3,3',9,10-Tetraepimer: *Xestosine A*
[290820-78-9]
 $C_{30}H_{50}N_2O_2$ 470.737
Alkaloid from *Xestospongia exigua*. Prisms (CH₂Cl₂/hexane). Mp 175-177°.

Braekman, J.C. *et al.*, *Tet. Lett.*, 1982, **23**, 4277-4280 (isol)
Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*, 1984, **93**, 941-944; 1988, **97**, 519-524 (isol, pmr, cmr, cryst struct)
Kobayashi, M. *et al.*, *Tet. Lett.*, 1989, **30**, 4149-4152 (isol, pmr, cmr, struct, Petrosine, Petrosine A)
Heathcock, C.H. *et al.*, *J.O.C.*, 1998, **63**, 5001-5012; 5013-5030 (synth)
Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (rev, synth)
Iwagawa, T. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1310-1311 (*Xestosine A*)

Petunianine A P-291

[140447-10-5]




$C_{34}H_{45}NO_6$ 563.733
Alkaloid from the leaves of *Petunia inflata* (Solanaceae). Cryst. (heptane/EtOAc). Mp 198-200°. $[\alpha]_D^{25}$ +76 (c, 0.25 in CHCl₃).

Elliger, C.A. *et al.*, *J.C.S. Perkin 1*, 1992, **5** (isol, uv, ir, pmr, cmr, cryst struct)
Elliger, C.A. *et al.*, *Phytochemistry*, 1993, **33**, 471 (isol, pmr, cmr)

Petunianine B P-292

[140475-14-5]




$C_{33}H_{43}NO_6$ 549.706
Alkaloid from the leaves of *Petunia inflata* (Solanaceae). Possesses insecticidal activity. Cryst. (EtOAc/2,3,3-trimethylpentane). Mp 260-262°. $[\alpha]_D^{25}$ +30 (c, 1 in CHCl₃).

Elliger, C.A. *et al.*, *J.C.S. Perkin 1*, 1992, **5** (isol, uv, ir, pmr, cmr, struct)
Elliger, C.A. *et al.*, *Phytochemistry*, 1993, **33**, 471 (isol, pmr, cmr)

Petunianine C P-293

[149725-28-0]




$C_{36}H_{49}NO_7$ 607.786
Constit. of *Petunia inflata*. Cryst. (EtOAc/heptane). Mp 182-184°. $[\alpha]_D^{25}$ +80 (c, 1 in CHCl₃).

Elliger, C.A. *et al.*, *Phytochemistry*, 1993, **33**, 471 (isol, pmr, cmr)

Petuniasterone B P-294

1-Acetoxy-24,25-epoxy-7,22-dihydroxytergost-4-en-3-one
[114176-00-0]



$C_{30}H_{46}O_6$ 502.69
Constit. of *Petunia hybrida*. Cryst. (EtOAc/heptane). Mp 191-192°. $[\alpha]_D^{25}$ +88.8 (CHCl₃).

22-Ac: *Petuniasterone B* 22-O-acetate
 $C_{32}H_{48}O_7$ 544.727
Constit. of *Petunia hybrida*. Cryst. (EtOAc/heptane). Mp 195-196°. $[\alpha]_D^{25}$ +39.5 (CHCl₃).

22-O-[(Methylthio)carbonyl]acetyl: *Petuniasterone B* 22-O-[(Methylthio)carbonyl]acetate
 $C_{34}H_{50}O_8S$ 618.83
Constit. of *Petunia hybrida*. Cryst. (MeOH). Mp 182-183°. $[\alpha]_D^{25}$ +65.7 (CHCl₃).


22-(3-Pyridinecarbonyl): *Petuniasterone B* 22-nicotinate
 $C_{36}H_{49}NO_7$ 607.786
Constit. of *Petunia inflata*. $[\alpha]_D^{25}$ +68 (c, 1 in CHCl₃).

7,22-Bis-(3-pyridinecarbonyl): *Petuniasterone B* 7,22-dinicotinate
 $C_{42}H_{52}N_2O_8$ 712.881
Constit. of *Petunia inflata*. $[\alpha]_D^{25}$ +21 (c, 0.45 in CHCl₃).

Elliger, C.A. *et al.*, *J.C.S. Perkin 1*, 1988, 711
Elliger, C.A. *et al.*, *Phytochemistry*, 1993, **33**, 471 (derivs)

Petuniasterone C **P-295**

24,25-Epoxy-7,22-dihydroxyergosta-1,4-dien-3-one, 9CI
[114176-01-1]



C₂₈H₄₂O₄ 442.637

Constit. of *Petunia hybrida*. Cryst. (EtOAc/heptane). Mp 183-185°. [α]_D²¹ +31.7 (CHCl₃).

22-Ac: [114176-06-6]

C₃₀H₄₄O₅ 484.675

Constit. of *Petunia hybrida*. Oil. Sol. MeOH. [α]_D²¹ +40.3 (CHCl₃). λ_{max} 246 (ε 21900) (MeOH) (Berdy).

7,22-Di-Ac:

C₃₂H₄₆O₆ 526.712

Constit. of *Petunia inflata*. [α]_D -27 (c, 0.13 in CHCl₃).

22-(Methylmalonyl): [201229-21-2]

C₃₂H₄₆O₇ 542.711

Constit. of *Petunia hybrida*.

22-O-[(Methylthio)carbonyl]acetyl: *Petuniasterone C* 22-O-[(methylthio)carbonyl]acetate [114176-05-5]

C₃₂H₄₆O₆S 558.778

Constit. of *Petunia hybrida*. Cryst. (EtOAc/heptane). Mp 141-142°. [α]_D²¹ +17.4 (CHCl₃).

22-(3-Pyridinecarbonyl): *Petuniasterone C* 22-nicotinate

C₃₄H₄₅NO₅ 547.733

Constit. of *Petunia inflata*. [α]_D²⁰ +25 (c, 0.2 in CHCl₃).

22-(3-Pyridinecarbonyl), 7-Ac: *Petuniasterone C* 22-nicotinate-7-acetate

C₃₆H₄₇NO₆ 589.77

Constit. of *Petunia inflata*. [α]_D -15 (c, 0.5 in CHCl₃).

7,22-Bis-(3-pyridinecarbonyl): *Petuniasterone C* 7,22-dinicotinate

C₄₀H₄₈N₂O₆ 652.829


Constit. of *Petunia inflata*. [α]_D +24 (c, 0.25 in CHCl₃).

Elliger, C.A. et al., *J.C.S. Perkin I*, 1988, 711
Elliger, C.A. et al., *Phytochemistry*, 1993, 33, 471 (derivs)

Moser, D. et al., *Pharm. Pharmacol. Lett.*, 1997, 7, 148-151; *CA*, 128, 99856e (methylmalonyl)

Peyoglunal **P-296**

5-Hydroxymethyl-1-[2-(3,4,5-trimethoxyphenyl)ethyl]-2-pyrrolecarboxaldehyde




C₁₇H₂₁NO₅ 319.357

Alkaloid from mescal (*Lophophora williamsii*) (Cactaceae).

Kapadia, G.J. et al., *J. Pharm. Sci.*, 1970, 59, 1699-1727 (isol, rev)

Peyoglutam

P-297



C₁₄H₁₇NO₄ 263.293

Alkaloid from peyote (*Lophophora williamsii*) (Cactaceae). Mp 217-219°.

Me ether: Mescalotam

C₁₅H₁₉NO₄ 277.319


Alkaloid from *Lophophora williamsii* (Cactaceae).

Kapadia, G.J. et al., *Chem. Comm.*, 1968, 1688

Peyonine

P-298

1-[2-(3,4,5-Trimethoxyphenyl)ethyl]-1H-pyrrole-2-carboxylic acid, 9CI [19717-25-0]



C₁₆H₁₉NO₅ 305.33


Alkaloid from peyote (*Lophophora williamsii* and other *Lophophora* spp.) (Cactaceae). Mp 131-133.5°.

Kapadia, G.J. et al., *J. Pharm. Sci.*, 1968, 57, 191 (isol, uv, ir, ms, pmr, synth)

Peyorvic acid

P-299

1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-methyl-1-isoquinolinecarboxylic acid, 9CI [29194-00-1]



C₁₃H₁₇NO₅ 267.281

(±)-form

Synthetic. Needles (MeOH). Mp 233-234° dec.

Me ether: [41303-72-4]

Synthetic. Mp 245-246° dec.

(ξ)-form

Trace component of *Lophophora williamsii* (peyote cactus) (Cactaceae). The nat. prod. was identified by glc-mass spectrometry in comparison with synthetic racemic matl.

Me ether: 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-methyl-1-isoquinolinecarboxylic acid, 9CI. **O-Methylpeyorvic acid**

C₁₄H₁₉NO₅ 281.308

Trace component of *Lophophora williamsii* (Cactaceae).

Kapadia, G.J. et al., *J.A.C.S.*, 1970, 92, 6943


(isol, ir, pmr, ms, synth, struct)

Kapadia, G.J. et al., *J. Het. Chem.*, 1973, 10, 135 (O-Methylpeyorvic acid)

Phaeantharine

P-300

Phaeantharine
[27670-80-0]



C₃₉H₄₀N₂O₆²⁺ 632.755

Struct. revised in 1983. Alkaloid from the bark of *Phaeanthus ebracteolatus* (Annonaceae). λ_{max} 228 ; 256 ; 282 ; 316 (MeOH) (Berdy). λ_{max} 228 ; 256 ; 282 ; 316 (MeOH-NaOH) (Berdy).

Dipicrate:

Cryst. (EtOH). Mp 180-184°.

Dipicrate:

Yellow amorph. solid. Mp 124-130°.

v. Bruchhausen, F. et al., *Arch. Pharm.*

(Weinheim, Ger.), 1957, 290, 232 (isol)

Van Beek, T.A. et al., *J. Nat. Prod.*, 1983, 46,

226 (struct, uv, pmr, cmr, ms)


Knabe, J. et al., *Arch. Pharm.* (Weinheim,

Ger.), 1984, 317, 92; 1986, 319, 950 (synth)

Phaeanthine

P-301

Phaeanthine
[1263-79-2]



Absolute Configuration

C₃₈H₄₂N₂O₆ 622.76

Alkaloids covered by this entry (abs. config. *R,R*-) are enantiomeric with those included in Tetrandrine, T-286 and diastereomeric with those included under Isotetrandrine, I-328 (1*R*,1'*S*-) and Peinamine, P-171 (1*S*,1'*R*-). Alkaloid from *Gyrocarpus americanus* (*Gyrocarpus jacquini*), *Phaeanthus ebracteolatus*, an unidentified *Phaeanthus* sp. (possibly *Phaeanthus macropodus*), *Triclisia patens*, *Pycnarrhena manillensis* and *Pycnarrhena novoguineensis* (Gyrocarpaceae, Annonaceae, Menispermaceae). Needles (Me₂CO or MeOH). Mp 222-224° (210°, 213-216°). [α]_D¹⁷ -284 (c, 4.015 in CHCl₃). Log P 8.3 (uncertain value) (calc). λ_{max} 218 (ε 31600); 240 (sh) (ε 15800); 282 (ε 12600) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 60 mg/kg. SF6850000

Dipicrate: Mp 263° dec. (*in vacuo*).

N²-α-Oxide: **Phaeanthine 2'α-N-oxide**

C₃₈H₄₂N₂O₇ 638.759


Alkaloid from roots and stems of *Pycnarrhena manillensis* (Menispermaceae). Needles (Me₂CO). Mp 193-195°.

- $[\alpha]_D^{27}$ -253 (c, 0.17 in CHCl_3). λ_{max} 282 (log ϵ 3.95) (MeOH).
- N^2, N^2 -Di-de-Me: 2,2'-Bisnorphaeanthine**
[110360-02-6]
 $\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706
Alkaloid from *Albertisia* cf. *Albertisia papuana* (Menispermaceae). $[\alpha]_D$ -272 (c, 0.1 in CHCl_3).
- N^2, N^2 -Di-Me:** Mp 267-268° dec. (as diiodide).
- O^6 -De-Me: Gyroamericine**
[102487-17-2]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733
Alkaloid from the stem bark of *Gyrocarpus americanus* (Hernandiaceae). Cryst. (MeOH). Mp 210°. $[\alpha]_D$ -238 (c, 1 in CHCl_3).
- O^7 -De-Me: Limacine**
[10172-02-8]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733
Alkaloid from *Limacia cuspidata*, *Limacia oblonga*, *Cyclea barbata*, *Pycnarrhena longifolia*, *Pycnarrhena novoguineensis*, *Colubrina faralaostra*, *Arcangelisia flava*, *Curarea candicans* and from a *Phaeanthus* sp. (possibly *Phaeanthus macropodus*) (Menispermaceae, Annonaceae). Cryst. ($\text{C}_6\text{H}_6/\text{Me}_2\text{CO}$). Mp 154-156°. $[\alpha]_D$ -212 (CHCl_3). λ_{max} 284 (MeOH).
- O^7 -De-Me, N^2 - β -oxide: Limacine 2 β -N-oxide**
[99883-66-6]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_7$ 624.732
Alkaloid from the roots of *Curarea candicans* (Menispermaceae). $[\alpha]_D$ -191 (c, 0.09 in CHCl_3).
- O^7 -De-Me, N^2 - α -oxide: Limacine 2' α -N-oxide**
[99877-67-5]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_7$ 624.732
Alkaloid from the roots of *Curarea candicans* (Menispermaceae). $[\alpha]_D$ -170 (c, 0.16 in CHCl_3).
- O^7 -De-Me, N^2 - β -oxide: Limacine 2' β -N-oxide**
[99877-66-4]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_7$ 624.732
Alkaloid from the roots of *Curarea candicans* (Menispermaceae). $[\alpha]_D$ -154 (c, 0.1 in CHCl_3).
- O^7 -De-Me, N^2 -de-Me: 2-Norlimacine**
 $\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706
Alkaloid from *Caryomene olivascens* (Menispermaceae). $[\alpha]_D$ -193 (c, 0.13 in CHCl_3).
- O^7 -De-Me, N^2 -de-Me: 2'-Norlimacine**
[152697-37-5]
 $\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706
Alkaloid from roots of *Cyclea barbata* (Menispermaceae). $[\alpha]_D$ -125 (c, 0.13 in CHCl_3).
- O^{12} -De-Me: Pycnamine**
[569-16-4]
 $\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733
Alkaloid from the bark of *Gyrocarpus americanus* (*Gyrocarpus jacquini*), the roots of *Pycnarrhena manillensis*, the stems of *Pycnarrhena novoguineensis*, and the stems and roots of *Triclisia patens* (Hernandiaceae, Menispermaceae). Cryst. (EtOH). Mp 186-187°.

- $[\alpha]_D^{25}$ -283 (CHCl_3).
- O^{12} -De-Me, dipicrate:**
Amorph. solid + $2\text{H}_2\text{O}$. Mp 218-220°.
- O^{12} -De-Me, N^2, N^2 -di-Me:**
Prisms + $6\text{H}_2\text{O}$ (H_2O). Mp 248-250°.
 $[\alpha]_D^{30}$ -164.4 (c, 0.365 in MeOH).
- $\text{O}^7, \text{O}^{12}$ -Di-de-Me: Krukovine**
[57377-42-1]
 $\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_6$ 594.706
Alkaloid from the stems of *Abuta splendida* and roots of *Pycnarrhena longifolia* (Menispermaceae). Prisms (CHCl_3). Mp 182-183° (173-178° dec.). $[\alpha]_D$ -180 (c, 0.06 in CHCl_3).
- McKenzie, A.W. et al., *Aust. J. Chem.*, 1953, **6**, 180-185 (Phaeanthine)
- Kidd, D.A.A. et al., *J.C.S.*, 1954, 669-677 (struct)
- Row, L.R. et al., *J. Sci. Ind. Res., Sect. B*, 1962, **21**, 581-582; *CA*, **59**, 690h (Phaeanthine)
- Battersby, A.R. et al., *J.C.S.*, 1965, 2239-2247 (ord, cd)
- Johns, S.R. et al., *Aust. J. Chem.*, 1968, **21**, 1387-1388 (Phaeanthine)
- Baldas, J. et al., *J.C.S. Perkin I*, 1972, 592-596 (ms)
- Saá, J.M. et al., *J.O.C.*, 1976, **41**, 317-319 (Krukovine)
- Koike, L. et al., *Tet. Lett.*, 1979, 3765-3768 (cmr)
- Bhakuni, D.S. et al., *Phytochemistry*, 1980, **19**, 2347-2350 (biosynth)
- Moulis, C. et al., *J. Nat. Prod.*, 1981, **44**, 101-103 (isol, uv, ir, pmr, ms)
- Siwon, J. et al., *Phytochemistry*, 1981, **20**, 323-325 (Limacine)
- Lavault, M. et al., *J. Chem. Res., Synop.*, 1985, 248-249; *J. Chem. Res., Miniprint*, 1985, 2786-2793 (Limacine oxides)
- Lavault, M. et al., *Chem. Pharm. Bull.*, 1986, **34**, 1148-1152 (2-Norlimacine)
- Chalandre, M.C. et al., *J. Nat. Prod.*, 1986, **49**, 101-105 (Gyroamericine)
- Lavault, M. et al., *Can. J. Chem.*, 1987, **65**, 343-347 (Bisnorphaeanthine)
- Regalado, J.C. et al., *Heterocycles*, 1987, **26**, 2573-2578 (Phaeanthine 2'-oxide)
- Guinaudeau, H. et al., *J. Nat. Prod.*, 1993, **56**, 1989-1992 (2'-Norlimacine)
- Wang, G. et al., *Life Sci.*, 1994, **56**, 295 (pharmacol, rev)
- Liu, Q.Y. et al., *J. Pharmacol. Exp. Ther.*, 1995, **273**, 32 (pharmacol)
- Schiff, P.L. et al., *J. Nat. Prod.*, 1997, **60**, 934-953 (Limacine, pmr, cmr)
- Mousa, S.A. et al., *Cardiovasc. Drug Rev.*, 1998, **16**, 48-61 (rev, pharmacol)


Phaeosphaeride A P-302

3,4,6,7-Tetrahydro-3,4-dihydroxy-6-methoxy-3-methyl-7-methylene-2-pentylpyrrol-2,3-c]pyrrol-5(2H)-one, 9CI
[910050-51-0]



- $\text{C}_{15}\text{H}_{23}\text{NO}_5$ 297.35
Similar to Curvupallide A, C-816. Prod. by an endophytic fungus closely related to *Phaeosphaeria avenaria*. Inhibitor of STAT3-dependent signalling. Yellow glass. $[\alpha]_D^{25}$ -93.6 (c, 2 in CH_2Cl_2). λ_{max} 262 (MeCN).
- 2-Epimer: **Phaeosphaeride B**

- [910050-52-1]
 $\text{C}_{15}\text{H}_{23}\text{NO}_5$ 297.35
Prod. by an endophytic fungus closely related to *Phaeosphaeria avenaria*. Orange glass. $[\alpha]_D^{25}$ -157.2 (c, 2 in CH_2Cl_2). λ_{max} 262 (MeCN).
- Maloney, K.N. et al., *Org. Lett.*, 2006, **8**, 4067-4070 (isol, pmr, cmr)


Phakellistatin 3, 9CI P-303
[153603-87-3]

- $\text{C}_{42}\text{H}_{54}\text{N}_8\text{O}_9$ 814.937
Isol. from the Western Indian ocean sponge *Phakellia carteri*. Shows significant cytotoxicity against murine P388 lymphocytic leukaemia. Amorph. powder. Sol. MeOH, CHCl_3 . Mp 178-180°. $[\alpha]_D^{24}$ -147 (c, 0.22 in MeOH). New type of cycloheptapeptide containing an amino acid unit apparently derived from a photooxidn. prod. of Tryptophan, T-640. λ_{max} 211 (ϵ 25700); 238 (ϵ 8130); 295 (ϵ 2340) (no solvent reported) (Derep). λ_{max} 214 (ϵ 14800); 234 (ϵ 11000); 291 (ϵ 3310) (MeOH) (Derep).

cis-Isomer: Isophakellistatin 3

- [153667-40-4]
 $\text{C}_{42}\text{H}_{54}\text{N}_8\text{O}_9$ 814.937
From *Phakellia carteri*. Cryst. (Me_2CO). Sol. MeOH, CHCl_3 . Mp 218-220°. $[\alpha]_D^{23}$ -138 (c, 0.21 in MeOH). Isomeric at the photo-Trp indole ring juncture. Does not show cytotoxicity against P388 cell line. λ_{max} 211 (ϵ 25700); 238 (ϵ 8130); 295 (ϵ 2340) (no solvent reported) (Derep). λ_{max} 214 (ϵ 14800); 234 (ϵ 11000); 291 (ϵ 3310) (MeOH) (Derep). λ_{max} 211 (ϵ 25700); 238 (ϵ 8130); 295 (ϵ 2344) (MeOH) (Berdy).

- Pettit, G.R. et al., *J.O.C.*, 1994, **59**, 1593 (isol, uv, ir, cryst struct)
- Greeman, K.L. et al., *Org. Lett.*, 2004, **6**, 1713-1716 (synth)

Phalaenopsines P-304

- $\text{C}_{20}\text{H}_{27}\text{NO}_5$ 361.437

Diastereoisomers formed by esterification of isomeric hydroxymethylpyrrolizidines with (*R*)-methyl 2-benzylmalate.

(1*R*,8*S*)-form**Phalaenopsine T**

[23412-97-7]

Alkaloid from *Phalaenopsis amabilis*, *Phalaenopsis aphrodite*, *Phalaenopsis fimbriata*, *Phalaenopsis sanderiana*, *Phalaenopsis stuartiana* and other *Phalaenopsis* spp. (often as mixt. with *Phalaenopsis* La) (Orchidaceae). Plant growth inhibitor. Cryst. (EtOH). Mp 104-105°. $[\alpha]_D^{20}$ -15 (c, 1.6 in EtOH). Derived from Trachelanthamidine.

Hydrochloride: [23412-98-8]

Mp 155-156°.

(1*S*,8*R*)-form**Phalaenopsine La**

[23412-99-9]

Alkaloid from *Phalaenopsis manni*, *Phalaenopsis amboinensis*, *Phalaenopsis sanderiana*, *Phalaenopsis schilleriana*, other *Phalaenopsis* spp. and from *Kingiella taenialis* (preferred genus name *Kingidium*) (Orchidaceae). Cryst. (EtOH). Mp 125-135°. $[\alpha]_D^{20}$ +10 (c, 1.1 in EtOH). Derived from Laburnine. Readily transesterifies during isol.

Hydrochloride: [23413-00-5]

Mp 131-132°.

(1*S*,8*S*)-form**Phalaenopsine Is**

[41451-64-3]

Alkaloid from *Phalaenopsis equestris* (Orchidaceae). $[\alpha]_D^{22}$ -42 (c, 1.5 in EtOH). Derived from Isoretroecanol.

Brandänge, S. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 1151-1154; 1970, **24**, 354; 1972, **26**, 2558-2560 (*Phalaenopsines La and Is, isol, struct, uv, ms, pmr*)


Brandänge, S. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 3668-3672 (*abs config*)

Fujieda, K. *et al.*, *Phytochemistry*, 1988, **27**, 1564-1566 (*Phalaenopsine T, isol, pmr, cmr, props*)

Phalarine**P-305**

[237078-85-2]

[934839-53-9 ((±)-form)]



Relative Configuration

C₂₄H₂₈N₄O₂ 404.511


Alkaloid from *Phalaris coerulescens*. Pale yellow oil. $[\alpha]_D^{20}$ -92 (c, 0.007 in MeOH).

Anderton, N. *et al.*, *Phytochemistry*, 1999, **51**, 153-157 (*isol, pmr, cmr, ms*)

Li, C. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 1448-1450 (*synth*)

Phallacidin, 9CI

[26645-35-2]



C₃₇H₅₀N₈O₁₃S 846.914

Cyclic peptide. A phallotoxin. Toxin from *Amanita phalloides* and *Amanita virosa*.

▶ Hepatotoxin, LD₅₀ (mus, ipr) 2 mg/kg. Human toxic effects occur within 2-5 hr of administration (*cf.* amatoxins).

GT8943000

Wieland, T. *et al.*, *Annalen*, 1962, **657**, 218 (*isol, uv, struct*)

Wieland, Th. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1967, **25**, 214 (*rev*)

Faulstich, H. *et al.*, *Eur. J. Biochem.*, 1971, **22**, 79 (*ord*)

Faulstich, H. *et al.*, *Z. Naturforsch., C*, 1974, **29**, 86 (*anal*)

Faulstich, H. *et al.*, *Annalen*, 1975, 2324 (*isol*)
Malak, S.H.A. *et al.*, *Planta Med.*, 1976, **29**, 80 (*occur*)

Wieland, T. *et al.*, *Crit. Rev. Biochem.*, 1978, **5**, 185 (*rev*)

Beutler, J.A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 422 (*anal*)

Wieland, T. *et al.*, *Peptides of Poisonous Amanita Mushrooms*, Springer-Verlag, 1986, (*book*)

Bönzli, P. *et al.*, *J.A.C.S.*, 1990, **112**, 3719 (*pmr, conformn*)

Antkowiak, R. *et al.*, *Alkaloids (Academic Press)*, 1991, **40**, 189 (*rev*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 12350 (*mushroom toxins*)

Phallacin**P-307**

3-(4-Hydroxy-D-leucine)phallacidin, 9CI [53568-40-4]

As Phallacidin, P-306 with
R¹ = COOH, R² = CH(CH₃)₂, R³ = CH(CH₃)CH₂OH, R⁴ = OH

C₃₇H₅₀N₈O₁₂S 830.914

A phallotoxin. Toxin from *Amanita phalloides*.

▶ Human toxic effects occur within 2-5 hr of administration (*cf.* amatoxins). Hepatotoxic, gastrointestinal effects by ingestion. LD₅₀ (mus, ipr) 2 mg/kg. SE9744000

Faulstich, H. *et al.*, *Z. Naturforsch., C*, 1974, **29**, 86

Faulstich, H. *et al.*, *Annalen*, 1975, 2324 (*isol*)
Wieland, T. *et al.*, *Crit. Rev. Biochem.*, 1978, **5**, 185 (*rev*)

Wieland, T. *et al.*, *Peptides of Poisonous Amanita Mushrooms*, Springer-Verlag, 1986, (*book*)

Antkowiak, R. *et al.*, *Alkaloids (Academic Press)*, 1991, **40**, 189 (*rev*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1390 (*mushroom toxins*)

Phallin B**P-308**

[57107-81-0]

As Phalloidin, P-311 with

R¹ = CH₃, R² = -CH₂Ph, R³ = -C(OH)(CH₃)₂, R⁴ = H

P-306C₄₁H₅₂N₈O₉S 832.976

Struct. found in CA Index Guide (1994). Exptl. evidence for struct. could not be found in lit. Isol. from *Amanita phalloides*.

Commercially available and/or commonly cited compound for which no useful literature data are available, 1998,

Phallisacin**P-309**

3-[4,5-Dihydroxy-4-(hydroxymethyl)-L-norvaline]phallacidin, 9CI. Phallisacin [58286-46-7]

As Phallacidin, P-306 with
R¹ = COOH, R² = CH(CH₃)₂, R³ = C(OH)(CH₂OH)₂, R⁴ = OH

C₃₇H₅₀N₈O₁₄S 862.913

Phallisacin appears to be a CA misprint. Toxin from *Amanita phalloides*.

▶ SE9740000

Faulstich, H. *et al.*, *Annalen*, 1975, 2324 (*isol, struct*)

Wieland, T. *et al.*, *Crit. Rev. Biochem.*, 1978, **5**, 185 (*rev*)

Phallisn**P-310**

7-[4,5-Dihydroxy-4-(hydroxymethyl)-L-norvaline]phalloidin, 9CI

[19774-69-7]

As Phallacidin, P-306 with
R¹ = R² = CH₃, R³ = -C(OH)(CH₂OH)₂, R⁴ = OH

C₃₅H₄₈N₈O₁₂S 804.877

A phallotoxin. Toxin from *Amanita phalloides*, also found in *Amanita virosa*.

▶ LD₅₀ (mus, ipr) 3 mg/kg. Gastrointestinal effects within 2-5h of ingestion (*cf.* amatoxins). Hepatotoxic. NJ8331000

Wieland, T. *et al.*, *Crit. Rev. Biochem.*, 1978, **5**, 185 (*rev*)

Lampe, K.F. *et al.*, *Annu. Rev. Pharmacol. Toxicol.*, 1979, **19**, 85 (*tox, rev*)


Wieland, T. *et al.*, *Peptides of Poisonous Amanita Mushrooms*, Springer-Verlag, 1986, (*book*)

Antkowiak, R. *et al.*, *Alkaloids (Academic Press)*, 1991, **40**, 189 (*rev*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1390

Phalloidin**P-311**

[17466-45-4]



Absolute Configuration

C₃₅H₄₈N₈O₁₁S 788.877

Phallotoxin. Isol. from *Amanita phalloides*, *Amanita verna* and *Amanita virosa*. Potassium (K_v) channel blocker. Pharmacol. tool. Cryst. +6 H₂O (H₂O). Sol. MeOH, H₂O, Py, EtOH. Mp 280-282°. Log P -1.64 (uncertain value) (calc). λ_{max} 290 (MeOH) (Berdy). λ_{max} 295 (H₂O) (Berdy).

► Adverse human effects by ingestion occur within 2-5 hr of administration (*cf.* amatoxins). Hepatotoxic, gastrointestinal effects by ingestion. LD₅₀ (mus, ipr) 2 mg/kg. LD₅₀ (mus, ims) 0.003 mg/kg. SE9800000

- Lynen, F. *et al.*, *Annalen*, 1938, **533**, 93 (*isol*)
 Wieland, T. *et al.*, *Annalen*, 1955, **593**, 157; 1962, **657**, 225 (*isol, uv, struct*)
 Faulstich, H. *et al.*, *Eur. J. Biochem.*, 1971, **22**, 79 (*ord*)
 Patel, D.J. *et al.*, *J. Mol. Biol.*, 1973, **79**, 185 (*pmr, conformn*)
 Faulstich, H. *et al.*, *Z. Naturforsch., C*, 1974, **29**, 86 (*anal*)
 Malak, S.H.A. *et al.*, *Planta Med.*, 1976, **29**, 80 (*occur*)
 Munekata, E. *et al.*, *J.A.C.S.*, 1977, **99**, 6151 (*cd, synth, analogues*)
 Wieland, T. *et al.*, *Crit. Rev. Biochem.*, 1978, **5**, 185 (*rev*)
 Beutler, J.A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 422 (*anal*)
 Fuerlinger, E. *et al.*, *Biochim. Biophys. Acta*, 1983, **760**, 411 (*uv, fluorescence*)
 Wieland, T. *et al.*, *Int. J. Pept. Protein Res.*, 1983, **21**, 3 (*analogues, synth*)
 Faulstich, H. *et al.*, *Klin. Wochenschr., Supp. 7*, 1986, **64**, 66 (*tox*)
 Wieland, T. *et al.*, *Peptides of Poisonous Amanita Mushrooms*, Springer-Verlag, 1986, (*book*)
 Wieland, T. *et al.*, *Naturwissenschaften*, 1987, **74**, 367 (*rev*)
 Enjalbert, F. *et al.*, *J. Chromatogr.*, 1989, **462**, 442 (*anal*)
 Antkowiak, R. *et al.*, *Alkaloids (Academic Press)*, 1991, **40**, 189 (*rev*)
 Kessler, H. *et al.*, *Annalen*, 1991, 179 (*struct, bibl, nmr, conformn*)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1390
 Anderson, M.O. *et al.*, *J.O.C.*, 2005, **70**, 4578-4584 (*synth*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PCU350

Phalloin P-312

7-(4-Hydroxy-L-leucine)phalloidin, 9CI [28227-92-1]
 As Phallacidin, P-306 with
 $R^1 = R^2 = \text{CH}_3$, $R^3 = -\text{C}(\text{CH}_3)_2\text{OH}$, $R^4 = \text{OH}$

$\text{C}_{35}\text{H}_{48}\text{N}_8\text{O}_{10}$ 772.878
 Toxin from *Amanita phalloides*. Cryst. (H₂O). Mp 250-280° dec.

► SF0350000

- Wieland, T. *et al.*, *Angew. Chem.*, 1957, **69**, 389 (*isol, uv*)
 Wieland, T. *et al.*, *Annalen*, 1958, **617**, 152; 1962, **657**, 225 (*isol, struct*)
 Faulstich, H. *et al.*, *Eur. J. Biochem.*, 1971, **22**, 79 (*isol*)
 Faulstich, H. *et al.*, *Z. Naturforsch., C*, 1974, **29**, 86 (*anal*)
 Beutler, J.A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 422 (*anal*)
 Wieland, T. *et al.*, *Annalen*, 1981, 2318 (*cd*)

Phascoline P-313

3-[(Aminoiminomethyl)amino]-N-(2-hydroxyheptyl)propanamide, 9CI. N-(3-Guanidinopropionyl)-2-hydroxyheptylamine
 $\text{H}_3\text{C}(\text{CH}_2)_4\text{CH}(\text{OH})\text{CH}_2\text{NHCOCH}_2\text{C}-\text{H}_2\text{NHC}(\text{NH}_2)=\text{NH}$
 $\text{C}_{11}\text{H}_{24}\text{N}_4\text{O}_2$ 244.336

(-)-form [50767-84-5]

Constit. of the sipunculid worm *Phascoscion strombi*. Oil; yellow needles (as picrate). Mp 73° (picrate). $[\alpha]_{\text{D}}^{25} -8.7$ (H₂O).
 Guillou, Y. *et al.*, *J. Biol. Chem.*, 1973, **248**, 5668 (*isol*)

Phascosomine P-314


3-[(Aminoiminomethyl)amino]-N-(2-methoxyheptyl)-2-methylpropanamide, 9CI. N-(3-Guanidinoisobutyryl)-2-methoxyheptylamine
 $\text{H}_3\text{C}(\text{C}_2)_4\text{CH}(\text{OMe})\text{CH}_2\text{NHCOCH}(\text{C}-\text{H}_3)\text{CH}_2\text{NHC}(\text{NH}_2)=\text{NH}$
 $\text{C}_{13}\text{H}_{28}\text{N}_4\text{O}_2$ 272.39

(-)-form

[50767-87-8 (picrate)]
 Constit. of the marine sipunculan worm *Phascoscion strombi*. Oil; yellow needles (as picrate). Mp 152° (picrate). $[\alpha]_{\text{D}}^{25} -37$ (H₂O).
 Guillou, Y. *et al.*, *J. Biol. Chem.*, 1973, **248**, 5668 (*isol*)

Phellibiline P-315

1,6-Didehydro-12,17-dihydro-3-hydroxy-C-homo-16(15H)-oxaerythrinan-15-one, 11CI. O-Demethyl-2,7-dihydro-C-homo- α -erythroidine, 10CI, 9CI
 [28305-80-8]



Absolute Configuration

$\text{C}_{16}\text{H}_{21}\text{NO}_3$ 275.347
 Alkaloid from *Phelline billiardieri* (Phellinaceae). Mp 160-161°. $[\alpha]_{\text{D}} +260$ (CHCl₃). Possibly an artifact.

A¹²-Isomer: Iso-phellibiline

[30519-45-0]
 $\text{C}_{16}\text{H}_{21}\text{NO}_3$ 275.347
 Alkaloid from *Phelline billiardieri* (Phellinaceae). Mp 186-188°. $[\alpha]_{\text{D}} +143$ (CHCl₃). Isom. to Phellibiline on chromatog. over alumina.

A¹²-Isomer, Me ether: O-Methylisophellibiline

2,7-Dihydrohomo- β -erythroidine
 [30519-68-7]
 $\text{C}_{17}\text{H}_{23}\text{NO}_3$ 289.374
 Alkaloid from *Phelline billiardieri* and from leaves of *Phelline* sp. aff. *Phelline lucida* (Phellinaceae). Mp 133°. $[\alpha]_{\text{D}} +140$ (CHCl₃).

12-Hydroxy, O³-Me: Phellibilidine


[76177-26-9]
 $\text{C}_{17}\text{H}_{23}\text{NO}_4$ 305.373
 Alkaloid from *Phelline billiardieri* (Phellinaceae). Cryst. (Me₂CO). Mp 114°. $[\alpha]_{\text{D}} -11$ (c, 1.1 in CHCl₃).

Hoang Nhu Mai, *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1970, **270**, 2154-2156 (*Phellibiline, Iso-phellibiline, O-Methylisophellibiline, ir, ms, struct*)
 Riche, C. *et al.*, *Acta Cryst. B*, 1974, **30**, 1386-1390 (*cryst struct, stereochem*)
 Séguineau, M.-F. *et al.*, *Phytochemistry*, 1980, **19**, 1279-1281 (*Phellibilidine, isol, uv, ir, cmr, pmr, ms, struct*)

Langlois, N. *et al.*, *Heterocycles*, 1984, **22**, 2453-2457 (*O-Methylisophellibiline*)

Phenalamide A1 P-316

Stipiamide
 [135383-02-7]



$\text{C}_{32}\text{H}_{45}\text{NO}_3$ 491.712
 Polyene-type antibiotic. Prod. by *Myxococcus stipitatus*. Inhibits the growth of KB cells. Exhibits anti-HIV and anti-fungal activities. Yellowish oil. $[\alpha]_{\text{D}}^{23} -115.1$ (c, 1.45 in MeOH). $[\alpha]_{\text{D}}^{25} -189$ (c, 1 in MeOH). λ_{max} 359 (ε 58900) (MeOH) (Derep). λ_{max} 209 (ε 32200); 264 (ε 6780); 359 (ε 31300) (MeOH/NaOH) (Derep). λ_{max} 207 (ε 9820); 264 (ε 6730); 359 (ε 31500) (MeOH) (Derep).

(6E)-Isomer: Phenalamide A2

[140695-41-6]
 $\text{C}_{32}\text{H}_{45}\text{NO}_3$ 491.712
 Prod. by *Myxococcus stipitatus*. Oil. Sol. MeOH, EtOAc. $[\alpha]_{\text{D}}^{25} +3$ (c, 1 in MeOH). λ_{max} 355 (log ε 4.78) (MeOH). λ_{max} 355 (ε 60256) (MeOH) (Berdy).

(4Z,6E)-Isomer: Phenalamide A3

[140695-42-7]
 $\text{C}_{32}\text{H}_{45}\text{NO}_3$ 491.712
 Prod. by *Myxococcus stipitatus*. Oil. Sol. MeOH, EtOAc. $[\alpha]_{\text{D}} -40.3$ (c, 0.3 in MeOH). λ_{max} 353 (log ε 4.76) (MeOH). λ_{max} 353 (ε 57540) (MeOH) (Berdy).


10ξ,11ξ-Epoxye (6E-): Phenalamide C

[140633-83-6]
 $\text{C}_{32}\text{H}_{45}\text{NO}_4$ 507.712
 Prod. by *Myxococcus stipitatus*. Oil. Sol. MeOH, EtOAc. $[\alpha]_{\text{D}}^{25} -56$ (c, 0.5 in MeOH). λ_{max} 328 (log ε 4.63) (MeOH). λ_{max} 328 (ε 42657) (MeOH) (Berdy).

Kim, Y.J. *et al.*, *J. Antibiot.*, 1991, **44**, 553 (*isol, pmr, cmr, struct*)
 Trowitzsch-Kienast, W. *et al.*, *Annalen*, 1992, 659 (*isol, uv, ir, pmr, cmr, ms*)
 Andrus, M.B. *et al.*, *J.A.C.S.*, 1997, **119**, 2327; 1999, 12159-12169 (*synth*)
 Hoffmann, R.W. *et al.*, *Org. Lett.*, 1999, **1**, 1713-1715 (*synth, Phenalamide A2*)

Phenalamide B P-317


[140633-82-5]



$\text{C}_{33}\text{H}_{47}\text{NO}_3$ 505.739
 Prod. by *Myxococcus stipitatus*. Oil. Sol. MeOH, EtOAc. $[\alpha]_{\text{D}}^{25} +60.9$ (c, 0.7 in MeOH). λ_{max} 350 (log ε 4.41) (MeOH). λ_{max} 349 (ε 25700) (MeOH) (Berdy).
 Trowitzsch-Kienast, W. *et al.*, *Annalen*, 1992, 659 (*isol, uv, ir, pmr, cmr, ms*)

Phenatic acid A

P-318

C₁₅H₁₉NO₅ 293.319


Prod. by *Streptomyces* sp. K03-0132. Potentiator of miconazole antifungal activity. Needles. Mp 147-149°. [α]_D²⁶ +2.5 (c, 0.1 in MeOH). λ_{\max} 217 (ε 12500); 262 (ε 6700); 345 (ε 2300) (MeOH).

Fukuda, T. et al., *J. Antibiot.*, 2005, **58**, 252-259 (isol, pmr, cmr)

1-Phenazinecarboxylic acid, 9CI

P-319

Tubermycin B. X-Pigment
[2538-68-3]

C₁₃H₈N₂O₂ 224.218

Prod. by *Pseudomonas* spp., *Streptomyces cinnamomensis*, *Streptomyces misakiensis*, *Bacillus* sp. B-6 and *Actinomadura dassonvillei*. Weakly active against gram-positive bacteria and *Candida albicans*. Inhibits acyl-CoA synthetase. Greenish-yellow needles (EtOH). Mp 241-242° (237-239°). λ_{\max} 251 (ε 97000); 364 (sh) (ε 10000); 370 (ε 19000) (CH₂Cl₂/pH 2) (Derep). λ_{\max} 250 (ε 84900); 364 (ε 13400) (MeOH/pH 11) (Derep). λ_{\max} 248 (ε 65000); 354 (sh) ; 364 (ε 13400) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 400 mg/kg. SG1576000

5-Oxide: [27210-90-8]

C₁₃H₈N₂O₃ 240.218

Photographic bleaching agent. Mp 223°.

10-Oxide:

C₁₃H₈N₂O₃ 240.218

Yellow needles (AcOH). Mp 233-234°.

Me ester: Methyl 1-phenazinecarboxylate
[3225-19-2]

C₁₄H₁₀N₂O₂ 238.245

Isol. from *Streptomyces luteoreticuli*. Active against mycobacteria. Mp 123-124°.

Amide: 1-Phenazinecarboxamide. **Oxy-chlororaphine**. Xanthoraphine
[550-89-0]

C₁₃H₉N₃O 223.234

Prod. by *Pseudomonas chlororaphis* and *Pseudomonas aeruginosa*. Active against phytopathogenic fungi and *Candida albicans*. Yellow needles. Mp 245-246° (237°).

► SG1575530

Nitrile: 1-Cyanophenazine

C₁₃H₇N₃ 205.218

Needles. Mp 171-172°.

5,10-Dihydro, carboxymethyl ester:
Carboxymethyl 5,10-dihydro-1-

phenazinecarboxylate. **Endophenazine D**

C₁₅H₁₂N₂O₄ 284.271

Prod. by various strains of the endosymbiotic *Streptomyces anulatus*. Active against gram-positive bacteria. Amorph. solid. λ_{\max} 293 (log ε 3.82); 351 (log ε 3.63) (MeOH).

5,10-Dihydro, amide; amide (1:3 complex): **Chlororaphin**

[13397-28-9]

C₅₂H₃₈N₁₂O₄ 894.95

Isol. from *Pseudomonas chloroaphis* and *Pseudomonas aeruginosa*. Active against gram-positive organisms. Green cryst. Mp 227°. Quinhydrone-type complex.

Kögl, F. et al., *Annalen*, 1930, **480**, 280 (deriv)

Birkofer, L. et al., *Chem. Ber.*, 1947, **80**, 212 (Chlororaphin, isol, synth)

Vivian, D.L. et al., *J.O.C.*, 1955, **20**, 797 (synth)

Isono, K. et al., *J. Antibiot., Ser. A*, 1958, **11**, 264 (struct)

Carter, R.E. et al., *J.A.C.S.*, 1961, **83**, 495 (Chlororaphin, biosynth)

Morgan, L.R. et al., *J.O.C.*, 1962, **27**, 4092 (Chlororaphin, biosynth)

Chaim, S. et al., *Spectrochim. Acta*, 1963, **19**, 1625 (ir)

Levitch, M.E. et al., *Arch. Biochem. Biophys.*, 1964, **106**, 194 (biosynth)

Mokrushin, V.S. et al., *CA*, 1970, **72**, 31750 (synth)

Hollstein, U. et al., *J.O.C.*, 1972, **37**, 3510 (synth)

Breitmaier, E. et al., *J.O.C.*, 1976, **41**, 2104 (cmr)

Kipriano, E.A. et al., *CA*, 1978, **88**, 11570 (use)

Kamer, D. et al., *J. Bacteriol.*, 1978, **134**, 690 (Chlororaphin, isol)

Tax, J. et al., *Coll. Czech. Chem. Comm.*, 1983, **48**, 527 (isol)

Brisbane, P.G. et al., *Antimicrob. Agents Chemother.*, 1987, **31**, 1967 (isol, struct)

Rewcastle, G.W. et al., *Synth. Commun.*, 1987, **17**, 1171 (synth)

Jones, G.P. et al., *Acta Cryst. C*, 1988, **44**, 2220 (cryst struct)

Ji, X. et al., *Acta Cryst. C*, 1992, **48**, 109 (cryst struct)

Jayatilake, G.S. et al., *J. Nat. Prod.*, 1996, **59**, 293-296 (isol, pmr, cmr)

Kim, K.-J. et al., *Yakhak Hoeji*, 1998, **42**, 552-557; *CA*, **130**, 14939z (isol, activity)

Kim, K.-J. et al., *J. Biochem. Mol. Biol.*, 2000, **33**, 332-336 (occur, activity)

Gebhardt, K. et al., *J. Antibiot.*, 2002, **55**, 794-800; 801-806 (Endophenazine D)


Sax, N.I. et al., *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 898

Hazards in the Chemical Laboratory, 3rd edn., (ed. Bretherick, L.), Royal Society of Chemistry, 1981, 437

1,6-Phenazinedicarboxylic acid

P-320

[23462-25-1]

C₁₄H₈N₂O₄ 268.228

Phenazine antibiotic. Fermentation prod. of *Pseudomonas* and *Streptomyces* spp. Inhibits xanthine oxidase. Shows

antibiotic props. by interacting with RNA. Intermed. in the biosynth. of Lomofungin, L-241 and other phenazines. Green cryst. Sol. EtOH, CHCl₃. Mp 325° dec. λ_{\max} 250 ; 370 (MeOH) (Berdy).

Mono-Me ester: **Phencomycin**

C₁₅H₁₀N₂O₄ 282.255

Prod. by a *Streptomyces* sp. Shows antibacterial and antitumor props. Yellow cryst. or yellow-green needles (CHCl₃/MeOH). Sol. MeCN, CHCl₃, MeOH, CH₂Cl₂, EtOAc, DMSO; fairly sol. MeOH; poorly sol. acids, hexane. Mp 263°. λ_{\max} 256 (log ε 4.87); 368 (log ε 4.2) (MeOH). λ_{\max} 250 ; 371 (MeOH/HCl) (Berdy). λ_{\max} 258 ; 366 (MeOH/NaOH) (Berdy).

Di-Me ester: [23531-24-0]

C₁₆H₁₂N₂O₄ 296.282

Mp 228-230°.

5,10-Dihydro, di-Me ester: **5,10-Dihydrophencomycin methyl ester**

C₁₆H₁₄N₂O₄ 298.298

Prod. by a marine *Streptomyces* sp. Orange needles (CHCl₃/MeOH). Mp 231°. λ_{\max} 216 (log ε 4.51); 245 (log ε 4.48); 450 (log ε 4.04) (MeOH).

Gerber, N.N. et al., *J. Het. Chem.*, 1969, **6**, 297 (isol, synth, uv)

Breitmaier, E. et al., *J.O.C.*, 1976, **41**, 2104 (synth, uv, cmr)

Kimura, Y. et al., *Tet. Lett.*, 1977, 4515 (biosynth)

Messenger, A.J. et al., *Biochem. Soc. Trans.*, 1978, **6**, 1326 (biosynth)

Buckland, P.R. et al., *J. Chem. Res., Synop.*, 1981, 362 (biosynth)

Roemer, A. et al., *Org. Magn. Reson.*, 1982, **19**, 118 (nmr)

Chatterjee, S. et al., *J. Antibiot.*, 1995, **48**, 1353 (Phencomycin)

Pusecker, K. et al., *J. Antibiot.*, 1997, **50**, 479 (Phencomycin, Dihydrophencomycin)


Maul, C. et al., *J. Antibiot.*, 1999, **52**, 1124-1134 (activity)

1,6-Phenazinediol, 9CI

P-321

1,6-Dihydroxyphenazine. **Crystalloiodinine B**

[69-48-7]

C₁₂H₈N₂O₂ 212.207

Isol. from *Streptomyces thioluteus*, *Brevibacterium crystalliodinum*, *Pseudomonas iodina* and various other bacteria. Active against gram-positive bacteria and fungi. Golden-yellow prisms (dioxan or EtOAc). Mp 280-281° (274°). λ_{\max} 262 (ε 13000); 291 (ε 30000) (HCl) (Derep). λ_{\max} 291 (ε 91160); 525 (ε 3820) (0.1N NaOH) (Derep). λ_{\max} 272 (ε 135000); 337 (ε 3130); 371 (ε 5200); 442 (ε 3500) (MeOH) (Derep).

► SG1579650

5-Oxide: [69-86-3]

C₁₂H₈N₂O₃ 228.207

Isol. from cultures of *Brevibacterium iodinum*, *Streptomyces thioluteus*,

Microbispora aerata, *Nocardioopsis dassonvillei* and a *Nocardia* strain. Red triangular platelets or cryst. (dioxan aq.). Sol. MeOH, Me₂CO, C₆H₆; poorly sol. H₂O. Mp 248-250° (218°) dec. λ_{max} 299; 575 (EtOH/NaOH) (Derep). λ_{max} 282 (ε 112000); 395 (ε 1710); 485 (ε 4260) (EtOH) (Derep). λ_{max} 283 (E1%/1cm 4900); 393 (E1%/1cm 75); 487 (E1%/1cm 107) (DMSO) (Berdy).

5,10-Dioxide: 1,6-Phenazinediol 5,10-dioxide. Iodinine. Crystalloiodinine A [68-81-5]

C₁₂H₈N₂O₄ 244.206

Pigment in *Chromobacterium iodinium*, *Streptomyces thioluteus*, *Nocardioopsis dassonvillei*, *Arthrobacter paraffinens*, *Streptosporangium album*, *Streptosporangium amethystogenes*, *Microbispora amethystogenes*, *Microbispora parva*, *Microbispora aerata*, *Brevibacterium stationis*, *Brevibacterium crystalloiodinum*, *Actinomadura dassonvillei*, *Waksmania aerata*, *Pseudomonas iodina*, *Sorangium* sp. and other microorganisms. Active against gram-positive bacteria, fungi and yeasts. Purple cryst. (CHCl₃). Sol. C₆H₆, toluene, bases, EtOAc, CHCl₃; poorly sol. MeOH, EtOH, butanol, AcOH, H₂O, hexane, acids. Mp 236° dec. Stable to acid, dec. by bases. λ_{max} 303; 368; 415; 615 (EtOH/NaOH) (Derep). λ_{max} 290 (ε 91000); 351 (ε 6600); 534 (ε 6100) (MeOH) (Berdy). λ_{max} 291 (E1%/1cm 6100); 352 (E1%/1cm 250); 535 (E1%/1cm 245) (CHCl₃) (Berdy).

► May explode when dry.

Di-Ac: [14031-12-0]

C₁₆H₁₂N₂O₄ 296.282

Yellow needles (EtOH). Mp 233-235° Mp 240-241°.

Mono-Me ether: 1-Hydroxy-6-methoxyphenazine. 6-Methoxy-1-phenazinol [13129-58-3]

C₁₃H₁₀N₂O₂ 226.234

From *Streptomyces thioluteus*. Active against mycobacteria. Yellow cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 190-193°. λ_{max} 271 (ε 67300); 371 (ε 2780); 438 (ε 1880) (EtOH) (Berdy).

Mono-Me ether, 5,10-dioxide: 6-Methoxy-1-phenazinol 5,10-dioxide. Myxin. 3C-Antibiotic [13925-12-7]

C₁₃H₁₀N₂O₄ 258.233

Isol. from *Sorangium* spp., *Lysobacter antibioticus* and *Pseudomonas* sp. Broad spectrum antibacterial and antifungal agent. Red needles (Me₂CO). Sol. MeOH, Et₂O; fairly sol. H₂O, hexane. Mp 130-135° dec. λ_{max} 283 (ε 97000); 340 (ε 5400); 505 (ε 6500) (0.1M HCl) (Derep). λ_{max} 278 (ε 25200); 304 (ε 26200); 410 (ε 7440); 605 (ε 3360) (EtOH/NaOH) (Derep). λ_{max} 285 (ε 70400); 350 (ε 6500); 390 (sh) (ε 5000); 407 (ε 2900); 502 (ε 6500) (EtOH) (Derep).

► May explode on drying with heat. LD₅₀ (mus, ipr) 133 mg/kg; LD₅₀ (mus, ipr) 40 mg/kg. SG1660000

Di-Me ether: 1,6-Dimethoxyphenazine

[13398-79-3]

C₁₄H₁₂N₂O₂ 240.261

From *Streptomyces thioluteus* and *Streptomyces luteoreticuli*. Active against *Sarcina lutea* and mycobacteria. Yellow needles. Mp 249-250°.

Di-Me ether, 5,10-dioxide:

C₁₄H₁₂N₂O₄ 272.26

Orange-red cryst. (C₆H₆/pentane). Mp 180-182°.

Clemo, G.R. *et al.*, *J.C.S.*, 1950, 1481-1485

(*synth, di-Me ether*)

Serebryani, S.B. *et al.*, *J. Gen. Chem. USSR*

(*Engl. Transl.*), 1952, **22**, 765 (*synth, di-Ac*)

Yoshioka, I. *et al.*, *Yakugaku Zasshi*, 1952, **72**,

847-848 (*synth, di-Ac, di-Me ether*)

Akabori, H. *et al.*, *J. Antibiot., Ser. A*, 1959,

12, 17 (*isol*)

Irie, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1960, **33**,

1057-1059 (*isol, 5,10-dioxide isol, di-Ac,*

synth)

Gerber, N.N. *et al.*, *Biochemistry*, 1964, **3**, 598;

1965, **4**, 176; 1966, **5**, 3824; 1970, **9**, 4611

(*isol*)

Peterson, E.A. *et al.*, *Can. J. Microbiol.*, 1966,

12, 221 (*isol, Myxin*)

Edwards, O.E. *et al.*, *Tet. Lett.*, 1966, 4867;

6502 (*struct, uv, ir, pmr, synth, Myxin*)

Grunberg, E. *et al.*, *Chemotherapy*, 1967, **12**,

272 (*activity, Myxin*)

Sigg, H.P. *et al.*, *Helv. Chim. Acta*, 1967, **50**,

716-718 (*mono-Me ether 5,10-dioxide, di-Me*

ether 5,10-dioxide, synth)

Gerber, N.N. *et al.*, *J.O.C.*, 1967, **32**, 4055 (*isol,*

mono-Me ether, uv)

Leimgruber, W. *et al.*, *Proc. Int. Symp. Drug*

Res., 1967, 1967, 240; 242 (*rev, Myxin*)

Weigele, M. *et al.*, *Tet. Lett.*, 1967, 715 (*synth,*

uv, pmr, ms, Myxin)

Hanson, A.W. *et al.*, *Acta Cryst. B*, 1968,

24, 1084; 1969, **25**, 1766 (*cryst struct, oxides*)

Japan. Pat., 1969, 79 86 694; *CA*, **91**, 156043

(*isol, Myxin*)

Breitmaier, E. *et al.*, *J.O.C.*, 1976, **41**, 2104-

2108 (*cmr*)

Albini, A. *et al.*, *J.C.S. Perkin 1*, 1978, 299-303

(*di-Ac, synth*)

Herbert, R.B. *et al.*, *J.C.S. Perkin 1*, 1979,

2411 (*biosynth*)

Tsujibo, H. *et al.*, *Agric. Biol. Chem.*, 1988, **52**,

301 (*isol*)

Alonso, A.M. *et al.*, *Org. Biomol. Chem.*, 2005,

3, 2832-2841 (*synth, pmr*)

1,8-Phenazinediol, 9CI

P-322

1,8-Dihydroxyphenazine

[18258-40-7]

C₁₂H₈N₂O₂ 212.207

Obt. from bacterial fermentation (*Pseudomonas phenazinium*). Yellow-brown solid. Mp 236° (230°).

10-Oxide: [23448-76-2]

C₁₂H₈N₂O₃ 228.207

Prod. by *Pseudomonas phenazinium*. Dark red solid. Sol. MeOH. Mp 235-240° dec. λ_{max} 285; 405; 540 (EtOH). λ_{max} 283; 403; 470 (EtOH/HCl) (Berdy). λ_{max} 295; 440; 550 (EtOH/NaOH) (Berdy).

Di-Ac: [18336-66-8]

C₁₆H₁₂N₂O₄ 296.282

Mp 178-181°.

Di-Me ether: 1,8-Dimethoxyphenazine

[14265-35-1]

C₁₄H₁₂N₂O₂ 240.261

Yellow cryst. Mp 154-155° (151-152°).

Yoshioka, I. *et al.*, *Pharm. Bull.*, 1953, **1**, 66;

CA, **49**, 12492

Gerber, N.N. *et al.*, *J. Het. Chem.*, 1969, **6**, 297

(*isol*)

Byng, G.S. *et al.*, *Biochem. Soc. Trans.*, 1975,

3, 742-744 (*oxide*)

1,9-Phenazinediol, 9CI

P-323

1,9-Dihydroxyphenazine

[23663-24-3]

C₁₂H₈N₂O₂ 212.207

Produced by bacteria. Orange needles or orange-red plates. Mp 296-297° dec.

Di-Ac: [23663-39-0]

C₁₆H₁₂N₂O₄ 296.282

Yellow cryst. (C₆H₆). Mp 258-259°.

Di-Me ether: 1,9-Dimethoxyphenazine

C₁₄H₁₂N₂O₂ 240.261

Yellow needles. Mp 254° Mp 259-260°.

Clemo, G.R. *et al.*, *J.C.S.*, 1950, 1481

Serebryanyi, S.B. *et al.*, *Zh. Obshch. Khim.*,

1952, **22**, 702; *CA*, **47**, 5408

Otomasu, H. *et al.*, *Chem. Pharm. Bull.*, 1958,

6, 77

2,3-Phenazinediol, 9CI

P-324

2,3-Dihydroxyphenazine

[19220-18-9]

C₁₂H₈N₂O₂ 212.207

Isol. from *Pseudomonas aureofaciens*.

Orange needles (EtOH). Mp 249-251°

Mp 340°.

5,10-Dioxide: [24890-65-1]

C₁₂H₈N₂O₄ 244.206

Mp 250°.

Di-Ac: [29453-85-8]

C₁₆H₁₂N₂O₄ 296.282

Yellow plates (C₆H₆ or xylene). Mp

230° (239-240°).

Di-Me ether: 2,3-Dimethoxyphenazine

C₁₄H₁₂N₂O₂ 240.261

Cryst. (petrol). Mp 226°.

Fischer, O. *et al.*, *Ber.*, 1890, **23**, 841 (*synth*)

Slack, P. *et al.*, *Nature (London)*, 1947, **160**,

437 (*synth*)

Yoshioka, I. *et al.*, *Pharm. Bull.*, 1954, **2**, 53

(*synth*)

Chernetskii, V.P. *et al.*, *Zh. Obshch. Khim.*,

1957, **27**, 2888; *CA*, **52**, 8147a (*synth*)

Otomasu, H. *et al.*, *Chem. Pharm. Bull.*, 1958,

6, 77 (*synth*)


Römer, A. *et al.*, *Tet. Lett.*, 1979, 509 (*isol, uv,*

pmr)

1,2,6-Phenazinetriol

P-325

1,2,6-Trihydroxyphenazine



C₁₂H₈N₂O₃ 228.207

Tri-Me ether: 1,2,6-Trimethoxyphenazine,

9CI. Mycomethoxin A

[39039-75-3]

C₁₅H₁₄N₂O₃ 270.287


Isol. from *Streptomyces luteoreticuli*.

Active against mycobacteria. Orange

needles. Mp 187-188°.

Yamanaka, S. *et al.*, *CA*, 1972, **77**, 162986

(Mycomethoxin A)

2,3,7-Phenazinetriol, 9CI P-3262,3,7-Trihydroxyphenazine
[71670-85-4]C₁₂H₈N₂O₃ 228.207Isol. from *Pseudomonas aureofaciens*.Römer, A. et al., *Tet. Lett.*, 1979, 509-512 (*pmr, struct*)**1-Phenazinol, 9CI** P-3271-Hydroxyphenazine. Hemipyocyanine
[528-71-2]C₁₂H₈N₂O 196.208

Prod. by *Pseudomonas aeruginosa*, *Pseudomonas pyocyanea*, *Pseudomonas aurantiaca* and *Streptomyces thioluteus*. Active against gram-positive bacteria and fungi. Shows some antiviral activity. Algicide. Fluffy yellow needles (EtOH aq.). Sol. MeOH, bases, Et₂O, Py, phenol; fairly sol. EtOH, CHCl₃; poorly sol. H₂O, hexane. Mp 157-159°. p*K*_{a1} 1.61; p*K*_{a2} 8.33 (15°). λ_{max} 264; 368 (MeOH) (Berdy). λ_{max} 264; 352; 360; 384; 425 (hexane) (Berdy). λ_{max} 263; 358 (EtOH) (Berdy). λ_{max} 298 (MeOH/NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 500 mg/kg. SG1647000

5-Oxide: [6479-82-9]

C₁₂H₈N₂O₂ 212.207

Yellow. Mp 190-191°.

10-Oxide: 1-Hydroxyphenazine 10-oxide
[14994-67-3]C₁₂H₈N₂O₂ 212.207

Prod. by microorganisms, e.g. *Brevibacterium iodinum*, *Microbispora aerata*, *Actinomadura dassonvillei*. Active against gram-positive bacteria. Orange cryst. Sol. CHCl₃; poorly sol. hexane. Mp 165-167°. λ_{max} 279; 326; 334; 368; 380; 387; 468 (ε 2120) (EtOH) (Berdy).

5,10-Dioxide: [18274-55-0]

C₁₂H₈N₂O₃ 228.207

Red. Mp 185-186°.

Ac: [6033-10-9]

C₁₄H₁₀N₂O₂ 238.245

Mp 120°.

Benzoyl: [6055-50-1]

C₁₉H₁₂N₂O₂ 300.316

Mp 173°.

5-Me: see Pyocyanine, P-833

Me ether: 1-Methoxyphenazine

[2876-17-7]

C₁₃H₁₀N₂O 210.235

From *Streptomyces luteoreticuli*. Active against mycobacteria. Cryst. (Py). Mp 167-169°. λ_{max} 261 (ε 60250); 395 (ε 8310); 407 (ε 3020) (MeOH) (Berdy).

Me ether, 5-Me: 1-Methoxy-5-methylphenazinium (1+)

[65162-13-2]

C₁₄H₁₃N₂O⁺ 225.27

Synthetic electron carrier for redox

mediated enzymatic reactions. Red cryst. (EtOAc/EtOH) (methanesulfonate). Mp 171-172° (as methanesulfonate salt). CAS no. refers to the methanesulfonate salt.

Et ether: 1-Ethoxyphenazine

[23169-51-9]

C₁₄H₁₂N₂O 224.262

Mp 126-127°.

(Methoxycarbonylmethyl) ether: (Methoxycarbonylmethoxy)phenazine. Antibiotic K₃-Ye. K₃-Ye

[159068-58-3]

C₁₅H₁₂N₂O₃ 268.271

Isol. from the culture of a genetically engineered *Streptomyces* sp. Inhibitor of nucleoside transport.

5-Me: see Pyocyanine, P-833

Hays, E.E. et al., *J. Biol. Chem.*, 1945, **159**, 725 (*isol*)Badger, G.M. et al., *J.C.S.*, 1951, 3204

Org. Synth., Coll. Vol., 3, 1955, 753-756 (1-methoxy-5-Me)

Gerber, N.N. et al., *Biochemistry*, 1966, **5**, 3824-3829 (10-oxide)Gerber, N.N. et al., *J.O.C.*, 1967, **32**, 4055 (*isol, props*)Wake, S. et al., *Tet. Lett.*, 1970, 2415 (*synth*)Kiprianova, E.A. et al., *Mikrobiol. Zh. (Kiev)*, 1971, **33**, 12; *CA*, **84**, 130344 (*isol, props*)Breitmaier, E. et al., *J.O.C.*, 1976, **41**, 2104 (*cmr*)Hisada, R. et al., *J. Biochem. (Tokyo)*, 1977, **82**, 1469-1473 (1-methoxy-5-Me, *synth, pmr, props*)Issidorides, C.H. et al., *Tetrahedron*, 1979, **34**, 217 (*synth*)Chen, W.P. et al., *Yaoxue Xuebao*, 1994, **29**, 585 (K₃-Ye)**2-Phenazinol, 9CI** P-328

2-Hydroxyphenazine. 2(10H)-Phenazone

[4190-95-8]

C₁₂H₈N₂O 196.208Formed by *Pseudomonas aureofaciens*.

Dark red cryst. + 1H₂O (EtOH), anhyd. at 100°. Mp 253-254° dec. (>200° dec.). p*K*_{a1} 2.47; p*K*_{a2} 7.24 (15°).

10-Oxide: [18274-41-4]

C₁₂H₈N₂O₂ 212.207

Mp 258° dec.

OH-form

Ac: [18258-47-4]

C₁₄H₁₀N₂O₂ 238.245Cryst. (EtOH or C₆H₆). Mp 151-152°.

Benzoyl: [35605-70-0]

C₁₉H₁₂N₂O₂ 300.316

Mp 144-145°.

Me ether: 2-Methoxyphenazine

[2876-18-8]

C₁₃H₁₀N₂O 210.235Cryst. (H₂O). Mp 126°.

O-(3ξ,7,11,15,19-Pentamethyl-6E,10E,14E,18-eicosatetraenyl):

Methanophenazine

[207606-29-9]

C₃₇H₅₀N₂O 538.815Isol. from *Methanosarcina mazei* Go1.Cofactor. [α]_D -3.7 (c, 1 in C₆H₆)

(synthetic S-enantiomer).

NH-form

N-Me: 10-Methyl-2(10H)-phenazinone, 9CI. Methylaposafranone

[63508-56-5]


C₁₃H₁₀N₂O 210.235

Redox indicator. Dark red needles (H₂O). Mp 202-204°. E° + 0.197 V (30°).

Priesler, P.W. et al., *J.A.C.S.*, 1937, **59**, 141 (*N-Me, synth, ind*)Levitch, M.E. et al., *Biochemistry*, 1966, **5**, 689 (*isol, uv, ir, synth*)Pietra, S. et al., *Ann. Chim. (Rome)*, 1971, **61**, 290 (*synth*)Inoue, H. et al., *Bull. Chem. Soc. Jpn.*, 1971, **44**, 3101 (*synth*)Flood, M.E. et al., *J.C.S. Perkin 1*, 1972, 622 (*isol*)Neuenhaus, W. et al., *Z. Naturforsch., B*, 1980, **35**, 385 (*synth*)Sugimoto, A. et al., *J. Het. Chem.*, 1999, **36**, 1057-1064 (*N-Me*)Beifuss, U. et al., *Tet. Lett.*, 2000, **41**, 9759-9763 (*Methanophenazine*)**Phenazinomycin**

P-329

[122898-63-9]


C₂₇H₃₂N₂O 400.563

Antibiotic from the mycelial extracts of *Streptomyces* sp. WK-2057. Exhibits antibacterial activity against gram-positive bacteria *in vitro*, direct cytotoxic activity against HeLaS₃, P388 and P388 doxorubicin-resistant cells *in vitro* and antitumour activity against experimental murine tumours *in vivo*. Dark-blue needles. Sol. MeOH, EtOAc, Me₂CO, CHCl₃; poorly sol. H₂O, hexane. Mp 113-118°. [α]_D²⁵ -49 (c, 0.45 in MeOH). Related to Lavanducyanin, L-76. λ_{max} 240 (ε 24500); 321 (ε 27200); 745 (ε 6100) (MeOH) (Derep). λ_{max} 238 (E1%/1cm 2500); 321 (E1%/1cm 2800); 730 (E1%/1cm 800) (MeOH) (Berdy).

Omura, S. et al., *J. Antibiot.*, 1989, **42**, 1037 (*isol, uv, ir, cmr, props*)Funayama, S. et al., *Tet. Lett.*, 1989, **30**, 3151 (*uv, ir, pmr, cmr, ms, struct*)Kinoshita, Y. et al., *Tet. Lett.*, 1997, **38**, 4993 (*synth*)**Phenazostatin A**

P-330

[185614-22-6]




C₂₈H₂₀N₄O₃ 460.491

Prod. by *Streptomyces* sp. 833. Neuronal cell protecting agent and free radical scavenger. Yellow cryst. Racemic. λ_{\max} 252 (ε 86000); 365 (ε 18000) (MeOH).

Kim, W.-G. *et al.*, *J. Antibiot.*, 1997, **50**, 715-721 (*isol*, *uv*, *ir*, *pmr*)

Phenazostatin B**P-331**

Dimethyl 6,6'-(1,2-dimethyl-1,2-ethanediyil)bis[1-phenazinecarboxylate], 9CI [73649-05-5]



Relative Configuration

C₃₂H₂₆N₄O₄ 530.582

Prod. by *Streptomyces* sp. 833 and *Streptomyces* sp. ME679m4. Neuronal cell protecting agent and free radical scavenger. Phosphodiesterase inhibitor. Yellow cryst. Racemic. λ_{\max} 253 (ε 89000); 366 (ε 19000) (MeOH).

Epimer: Phenazostatin D

[639512-19-9]


C₃₂H₂₆N₄O₄ 530.582

Prod. by the marine *Pseudonocardia* sp. B6273. Yellow solid. *meso*-form. λ_{\max} 256 (log ε 3.95); 350 (sh) (log ε 3.16); 366 (log ε 3.34); 385 (sh) (log ε 2.92) (CHCl₃).

Japan. Pat., 1980, 80 03733; *CA*, **92**, 213541p (*isol*)

Kim, W.-G. *et al.*, *J. Antibiot.*, 1997, **50**, 715-721 (*isol*, *uv*, *ir*, *pmr*)

Maskey, R.P. *et al.*, *Z. Naturforsch., B*, 2003, **58**, 692-694 (*Phenazostatin D*)


Phenazostatin C**P-332**C₃₀H₂₂N₄O₄ 502.528

Prod. by *Streptomyces* sp. 833. Neuronal cell protecting agent. Antioxidant. Yellow cryst. (MeOH). Racemic. λ_{\max} 249 (ε 109000); 365 (ε 26000) (MeOH).

Kim, W.-G. *et al.*, *J. Antibiot.*, 1999, **52**, 758-761

Phenopyrrozin**P-333**

5,6,7,7a-Tetrahydro-2-hydroxy-1-phenyl-3H-pyrrolizin-3-one. Antibiotic FO 2047. FO 2047



(R)-form

C₁₃H₁₃N₂O 215.251**(R)-form** [172923-76-1]

Prod. by *Penicillium* sp. FO-2047 and the marine-derived *Chromocleista* sp. strain R721. Radical scavenger. Antioxidant. Powder. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. Mp 147-152°. $[\alpha]_D^{25}$ -10.2 (c, 0.6 in MeOH). λ_{\max} 206 (ε 7960); 215 (ε 7850); 225 (sh) (ε 5810); 240 (ε 3870); 295 (ε 3810); 340 (sh) (ε 650) (EtOH).

(S)-form

4'-Hydroxy-5,6,7,7a-Tetrahydro-2-hydroxy-1-(4-hydroxyphenyl)-3H-pyrrolizin-3-one. **p-Hydroxyphenopyrrozin** [887602-12-2]

C₁₃H₁₃N₂O₃ 231.251

Prod. by the marine-derived *Chromocleista* sp. strain R721. Amorph. solid. $[\alpha]_D^{25}$ +34 (c, 0.34 in MeOH). λ_{\max} 198 (log ε 4.5); 222 (log ε 1.3); 307 (log ε 1.8) (MeOH).


Shiomi, K. *et al.*, *J. Antibiot.*, 1995, **48**, 1413-1418 (*isol*, *uv*, *pmr*, *cmr*)

Japan. Pat., 1996, 96 127 583; *CA*, **125**, 112919n

Park, Y.C. *et al.*, *J. Nat. Prod.*, 2006, **69**, 580-584 (*isol*, *Chromocleista*)

Phenoxan**P-334**

3-Ethyl-6-methoxy-5-methyl-2-[2-(3-methyl-4-phenyl-3-butenyl)-4-oxazolyl]-4H-pyran-4-one, 9CI [134332-63-1]

C₂₃H₂₅NO₄ 379.455

Isol. from *Polyangium* sp. Inhibitor of HIV-1 infection. NADH: ubiquinone oxidoreductase (Class II) inhibitor. Cryst. Mp 92-93°. Log P 4.19 (calc). λ_{\max} 204 (ε 75300); 229 (ε 53800); 244 (ε 42700); 252 (ε 42700); 262 (sh); 274 (sh) (MeOH) (Derp). λ_{\max} 204; 229 (ε 53900); 244 (ε 42600); 252 (ε 42600) (MeOH) (Berdy).

Jansen, R. *et al.*, *Annalen*, 1991, 707 (*isol*, *pmr*, *struct*)

Kunze, B. *et al.*, *J. Antibiot.*, 1992, **45**, 1549 (*props*)


Friedrich, L.T. *et al.*, *Eur. J. Biochem.*, 1994, **219**, 691 (*activity*)

Garey, D. *et al.*, *J.O.C.*, 1996, **61**, 4853 (*synth*, *pmr*, *cmr*)

Zhang, X. *et al.*, *J. Het. Chem.*, 1997, **34**, 1061-1064 (*synth*)

3H-Phenoxazin-3-one, 9CI**P-335**

3-Phenoxazone [1916-63-8]

C₁₂H₇NO₂ 197.193

Isol. from *Pycnoporus sanguineus*. Orange cryst. Mp 217-219° dec.

Kehrmann, F. *et al.*, *Ber.*, 1902, **35**, 341 (*synth*)

Musso, H. *et al.*, *Chem. Ber.*, 1957, **90**, 1814 (*uv*)

Corbett, J.F. *et al.*, *Spectrochim. Acta*, 1965, **21**, 1411 (*synth*, *ir*)

Musso, H. *et al.*, *Chem. Ber.*, 1978, **111**, 3012 (*synth*)

Bolognese, A. *et al.*, *J. Het. Chem.*, 1986, **23**, 1003 (*synth*, *uv*, *pmr*)

Barret, R. *et al.*, *Synth. Commun.*, 1990, **20**, 1543 (*synth*)

Achenbach, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 3 (*isol*)

Bolognese, A. *et al.*, *J. Organomet. Chem.*, 2002, **45**, 5205-5216 (*synth*, *uv*, *pmr*)

Phenylacetic acid**P-336**

Benzeneacetic acid, 9CI. *Phenylethanoic acid*. *α-Toluic acid*. FEMA 2878 [103-82-2]

PhCH₂COOHC₈H₈O₂ 136.15

Found in essential oils, e.g. neroli, rose oil, free and as esters, and in many fruits. Also prod. by microorganisms, e.g. by *Fusarium oxysporum*, *Glomerella cingulata* and a marine derived *Streptomyces* sp. Volatile component of tail gland secretion from red deer *Cervus elaphus*. Used as 1M soln. in CHCl₃ for selective extraction separation of Cu and U (CHCl₃). Important industrial intermediate. Perfumery and flavouring ingredient. Incorporation in fermentation of penicillin results in production of Benzylpenicillin. Phytotoxin, germination inhibitor. Plates (petrol) with sweet taste at low conc. and rose-like odour. Mp 77-78.5°. Bp 265.5° Bp_{0.01} 65-70°. pK_a 4.31 (H₂O).

► Fl. p. 102°. LD₅₀ (rat, orl) 2250 mg/kg. Exp. teratogenic effects. AJ2430000

Amide: Benzeneacetamide, 9CI. *Phenacetamide* [103-81-1]

C₈H₉NO 135.165

Produced by *Actinomyces* spp., sponge *Halichondria* sp., *Streptomyces albogiger*, *Streptoverticillium olivoreticuli*, *Tedania anhelans*. Constit. of the leaves of *Allophylus cobbe*. Plant growth regulator. Plates or leaflets (C₆H₆). Mp 157-158°.

► AC7705000

Diethylamide: N,N-Diethylbenzeneacetamide, 9CI [2431-96-1]

C₁₂H₁₇NO 191.272

Found in various plant oils, e.g. garden cress (*Lepidium sativum*). Mp 86°. Bp 297°.

► AB7410000

Benzylamide: N-Benzylbenzeneacetamide. *N-Benzyl-2-phenylacetamide* [7500-45-0]

C₁₅H₁₅NO 225.29

Constit. of the stems of *Salvadora persica*. Cryst. (EtOH). Mp 121.5°.

Nitrile: Benzeneacetonitrile, 9CI. *Benzyl cyanide*. *Benzyl nitrile*. (*Cyano-methyl*)/benzene. *Cyano-phenylmethane*.

Phenylacetoneitrile

[140-29-4]

C₈H₇N 117.15

Isol. from thick-tailed galago *Galago crassicaudatus*, oil of *Lepidium sativum* and other plant oils. Used in the detection of nitro compds. Liq. d_{15}^{25} 1.02. Bp 234° Bp₁₂ 107°. n_D^{25} 1.5211.

- Explosive reaction with NaOCl + H⁺. Skin irritant. LD₅₀ (rat, orl) 270 mg/kg. LD₅₀ (rbt, skn) 270 mg/kg. AM1400000 [114-70-5]

Kotera, A. *et al.*, *J.A.C.S.*, 1955, **77**, 6183-6186 (benzylamide)

Org. Synth., *Coll. Vol.*, **4**, 1963, 760 (amide)

Crewe, R.M. *et al.*, *J. Chem. Ecol.*, 1979, **5**, 861-868 (nitrile, isol)

Strauss, C.R. *et al.*, *Org. Prep. Proced. Int.*, 1995, **27**, 552 (amide)

Khalil, A.T. *et al.*, *Arch. Pharmacol. Res.*, 2006, **29**, 952-956 (benzylamide, isol)

N-(Phenylacetyl)-2-butenediamide, 9CI P-337

PhCH₂CONHCOCH=CHCONH₂C₁₂H₁₂N₂O₃ 232.238

(E)-form

N-Phenylacetylfulfaramide. **Fumaramid-mycin**. C 9154. Ro 09-0049. Antibiotic C 9154. Antibiotic Ro 09-0049 [57687-92-0]

Isol. from *Streptomyces kurssanovii* (grown on agar plates, not obt. from submerged cultures). Active against gram-positive and -negative bacteria. Mp 202-203° dec. λ_{\max} 212 (sh) (ϵ 20000); 225 (ϵ 23400) (EtOH) (Derep).


- LD₅₀ (mus, orl) 1250 mg/kg. EM4300000 Maruyama, H.B. *et al.*, *J. Antibiot.*, 1975, **28**, 636; 648 (isol, struct) Suhara, Y. *et al.*, *J. Antibiot.*, 1975, **28**, 648 Hasegawa, T. *et al.*, *J. Antibiot.*, 1975, **28**, 713 Japan. Pat., 1976, 76 32 789; CA, **85**, 61412n (synth)

Phenylahistin P-338

NSCL-96F037

[200815-37-8]

[200925-27-5, 200815-38-9]



(S)-form

C₂₀H₂₂N₄O₂ 350.419

Prod. by *Aspergillus ustus* NSC-F038. Cytotoxic agent (S-isomer only). Mammalian cell cycle inhibitory agent. Inhibits tubulin polym. Amorph. powder. Mp 233-236°. $[\alpha]_D^{25}$ +123 (c, 0.1 in MeOH). Obt. as a partial racemate, R:S 3:1. λ_{\max} 202 (log ϵ 4.38); 233 (sh) (log ϵ 4.05); 320 (log ϵ 4.43) (MeOH).

6,7-Didehydro: Dehydrophenylahistin

C₂₀H₂₀N₄O₂ 348.404

Synth. by enzymic dehydrogenation. Potent cytotoxic agent. Mixture of E/Z-isomers.

Kanoh, K. *et al.*, *Bioorg. Med. Chem. Lett.*,


1997, **7**, 2847-2852 (isol, uv, ir, pmr, cmr) Kanoh, K. *et al.*, *J. Antibiot.*, 1999, **52**, 134-141 (activity)

Hayashi, Y. *et al.*, *J.O.C.*, 2000, **65**, 8402-8405 (synth)

Kanzaki, H. *et al.*, *J. Antibiot.*, 2002, **55**, 1042-1047 (Dehydrophenylahistin)

Phenylalanyl-N-[1-(1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methyl-1-isoquinolinyl)ethyl]glycinamide, 9CI P-339

[147471-63-4]

C₂₄H₃₂N₄O₄ 440.541

Tentative struct. Similar to Amphibine I, A-934. Alkaloid from flowers of *Hibiscus rosa* (Malvaceae).

O⁷-De-Me, O⁶-Me: α -Amino-N-[2-oxo-2-[[1-(1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-2-methyl-1-isoquinolinyl)ethyl]amino]ethyl]benzenepropanamide, 9CI

[147455-43-4]

C₂₄H₃₂N₄O₄ 440.541


From flowers of *Hibiscus rosa* (Malvaceae). Tentative struct.

Khokhar, I. *et al.*, *Sci. Int. (Lahore)*, 1992, **4**, 147; CA, **118**, 230140e (isol, struct)

2-Phenyl-1-benzazocin-8-ol, 9CI P-340

8-Hydroxy-2-phenyl-1-benzazocine. **Hy-podematine**

[134746-11-5]

C₁₇H₁₃NO 247.296

Alkaloid from fern *Hypodematum sinense*. Light yellow cryst. Mp 156-158°.


Chen, Y. *et al.*, *Chin. Chem. Lett.*, 1990, **1**, 221; CA, **115**, 131977q (isol, pmr, cmr, struct)

Chen, Y. *et al.*, *J. Chin. Pharm. Sci.*, 1992, **1**, 1; CA, **118**, 209429w

2-(2-Phenylethenyl)indole, 9CI P-341

2-Styrylindole

[60986-51-8]



(E)-form

C₁₆H₁₃N 219.285

(E)-form [29475-88-5]

Cryst. (EtOAc/petrol). Mp 211-212° (207°).

(Z)-form

Ipomine A

[128263-26-3]

Alkaloid from the hairy roots of *Ipomoea batatas*.

Alper, H. *et al.*, *Chem. Comm.*, 1976, 483 (synth)

Frannicolo, F. *et al.*, *Tet. Lett.*, 1984, **25**, 3101 Eitel, M. *et al.*, *Synthesis*, 1989, 364-367 (synth, pmr)

Nagarathnam, D. *et al.*, *Synthesis*, 1992, 743-745 (synth)

Hudkins, R.L. *et al.*, *J.O.C.*, 1995, **60**, 6218 (synth, pmr)

Yuan, S. *et al.*, *Yaoxue Xuebao*, 2004, **39**, 818-830 (isol)

2-Phenylethylamine P-342

Benzeneethanamine, 9CI. Phenethylamine, 8CI. β -Aminoethylbenzene. FEMA3220. β -Phenethylamine

[64-04-0]

PhCH₂CH₂NH₂C₈H₁₁N 121.182

Very widely distributed alkaloid, esp. in *Acacia* spp. and *Crataegus* spp. Also present in animal tissues, some algae, fungi and cacti (Fabaceae, Rosaceae, Cactaceae). Metab. of *Prosopis alba*. Prod. by *Streptomyces* 699-A3. Shows DNA binding activity. Monoamine oxidase inhibitor. Liq. with a fish odour. d_4^{25} 0.96. Bp 197-198° Bp₇ 70-71°. n_D^{25} 1.5290. Absorbs CO₂ from air.

- Fl. p. 84°. Skin irritant and possible sensitiser. SG8750000

Hydrochloride: [156-28-5]

Leaflets (EtOH). Mp 223-224° (217°).

- SH7175000

N-Formyl: (2-Phenylethyl)formamide,

9CI. 2-Phenethylformamide

[23069-99-0]

C₉H₁₁NO 149.192

Oil.

N-Ac: N-(2-Phenylethyl)acetamide. N-Phenethylacetamide

[877-95-2]

C₁₀H₁₃NO 163.219

Prod. by *Streptomyces* sp. Lz531 and a limnic bacterial strain. Needles. Mp 51-52°. Bp 305-306° Bp₂ 154°.

- AC7740000

N-Propanoyl: N-(2-Phenylethyl)propanamide. N-Phenethylpropanamide

[6283-04-1]

C₁₁H₁₅NO 177.246

Prod. by the limnic bacterial strain GW90a. Amorph. Mp 52°.

N-(2-Methylpropanoyl): 2-Methyl-N-(2-phenylethyl)propanamide. N-Phenethylisobutyramide

[71022-62-3]

C₁₂H₁₇NO 191.272

Prod. by the limnic bacterial strain GW90a. Amorph. Mp 80-81°.

N-(2-Methylbutanoyl): 2-Methyl-N-(2-phenylethyl)butanamide. 2-Methyl-N-

- phenethylbutyramide*
[53181-99-0]
C₁₃H₁₉NO 205.299
Prod. by a limnic bacterial strain
GW90a. Amorph. Mp 57-58°.
- N-(3-Methylbutanoyl): 3-Methyl-N-(2-phenylethyl)butanamide. 3-Methyl-N-phenethylbutyramide. N-Phenethylisovaleramide
C₁₃H₁₉NO 205.299
Prod. by a North sea bacterium
Cytophaga marinoflava sp. AM13.1.
Also prod. by a limnic bacterial strain
GW90a. Amorph. Mp 64°.
- N-Hexanoyl: N-(2-Phenylethyl)hexanamide. N-Phenethylhexanamide
[10264-26-3]
C₁₄H₂₁NO 219.326
Prod. by the limnic bacterial strain
GW90a. Amorph. Mp 57-58°.
- N-(13-Methyltetradecanoyl): 13-Methyl-N-(2-phenylethyl)tetradecanamide
[921607-14-9]
C₂₃H₃₉NO 345.567
Prod. by *Streptomyces* sp. Lz531.
Powder. [α]_D²³ +7.1 (c, 0.6 in CHCl₃).
Unexplained opt. rotation.
- N-(7-Methoxy-4-tetradecenoyl) (4E,7S-): **Hermitamide A**
[286012-62-2]
C₂₃H₃₇NO₂ 359.551
Alkaloid from *Lyngbya majuscula*.
Cytotoxic agent. Pale yellow oil. [α]_D²⁶ -9.3 (c, 0.45 in CHCl₃). λ_{max} 216 (ε 5700) (EtOH).
- N-Hexadecanoyl: N-(2-Phenylethyl)hexadecanamide. Hexadecanoic acid 2-phenylethylamide
[10015-69-7]
C₂₄H₄₁NO 359.594
Alkaloid from the edible mushroom
Laetiporus sulphureus. Amorph. powder. Mp 180-181°. λ_{max} 259 (MeOH).
- N-(9-Hydroxyhexadecanoyl): 9-Hydroxy-N-(2-phenylethyl)hexadecanamide
[172303-93-4]
C₂₄H₄₁NO₂ 375.593
Isol. from *Teleso riisei*. Mildly cytotoxic to murine leukaemia cells (P-388) in culture. Cryst. Mp 78°. [α]_D²⁵ -2.68 (c, 0.41 in CHCl₃). Erroneous struct. diag. in paper.
- N-(9-Oxohexadecanoyl): 9-Oxo-N-(2-phenylethyl)hexadecanamide
[172303-92-3]
C₂₄H₃₉NO₂ 373.578
Isol. from the coelenterate *Teleso riisei*. Mildly cytotoxic to murine leukaemia cells (P-388) in culture. Cryst. Mp 85°. Erroneous struct. diag. in paper.
- N-Benzoyl: N-(2-Phenylethyl)benzamide. N-Phenethylbenzamide
[3278-14-6]
C₁₅H₁₅NO 225.29
Alkaloid from *Haplophyllum tuberculatum*, *Oxytropis muriceta* and *Oxytropis pseudoglandulosa*. Leaflets (EtOH). Mp 117-118°.
- N-(4-Hydroxy-E-cinnamoyl): N-trans-p-Coumaroyl-2-phenylethylamine
[103197-54-2]
C₁₇H₁₇NO₂ 267.327
Alkaloid from *Anomianthus dulcis*.
Cryst. Mp 146-148°. λ_{max} 217 (log ε 4.49); 223 (log ε 4.47); 277 (log ε 4.51); 330 (sh) (log ε 4.12) (MeOH).
- N-Me: N-Methylphenethylamine
[589-08-2]
C₉H₁₃N 135.208
Main alkaloid from *Acacia adunca*, also present in other *Acacia* spp., *Eria jarensis* and in *Arthropytum leptocladum* (Fabaceae, Chenopodiaceae). Bp 205° Bp₃₈ 112.5-115° (lit. gives a pressure range).
▶ SH9625000
N-Me, hydrochloride: [4104-43-2]
Plates (cyclohexane). Mp 162-164°.
- N,N-Di-Me: N,N-Dimethylphenethylamine
[1126-71-2]
C₁₀H₁₅N 149.235
Isol. from defensive secretions of Daddy long legs *Sclerobunus robustus*. Alkaloid from *Eria jarensis* (Orchidaceae). Bp 205° Bp₆ 66-68°.
- N,N-Di-Me, hydrochloride: [10275-21-5]
Flakes (EtOH/EtOAc/hexane). Mp 164-165°.
- N,N,N-Tri-Me: N,N,N-Trimethylbenzeneethanaminium(1+), 9CI. Trimethylphenethylammonium(1+), 8CI
[772-63-4]
C₁₁H₁₈N[⊕] 164.27
Quaternary alkaloid from *Eria jarensis* (Orchidaceae).
- N-Et: [22002-68-2]
C₁₀H₁₅N 149.235
Bp₁₄ 99°.
- N,N-Di-Et: [5300-21-0]
C₁₂H₁₉N 177.289
Bp 226° Bp₂ 88-90°.
- N-Benzyl: Benzyl-2-phenylethylamine. Benethanamine
[3647-71-0]
C₁₅H₁₇N 211.306
Bp₁₆ 162-165°.
- [62510-53-6, 1077-11-8, 6068-85-5, 75337-06-3]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 1269D; 1270A; 1275D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 2, 570C; 571A; 583C; 584A (nmr)
Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1164C; 1166D (ir)
Sadtler Standard C-13 NMR Spectra, 68 (cmr)
Org. Synth., Coll. Vol., 3, 1955, 720-722 (synth)
Fitzgerald, J.S. et al., *Aust. J. Chem.*, 1964, 17, 160-162 (isol, pmr; N-Methylphenethylamine)
Hedman, K. et al., *Acta Chem. Scand.*, 1969, 23, 3261 (*Eria jarensis* constitits)
Mosnaim, A. et al., *Clin. Chim. Acta*, 1973, 46, 407-413 (ir, ms)
Smith, T.A. et al., *Phytochemistry*, 1977, 16, 9-18 (rev, occur)
Iwase, H. et al., *Chem. Pharm. Bull.*, 1979, 27, 1009-1014 (glc, ms)
Doetsch, P.W. et al., *J. Chromatogr.*, 1980, 189, 79-85 (occur)
Keller, W.J. et al., *Phytochemistry*, 1982, 21, 2851-2852 (N-Methylphenethylamine, biosynth)
Ekpa, O. et al., *Tet. Lett.*, 1984, 25, 1315-1318 (*Dimethylphenethylamine, isol*)
- Purchase, C.F. et al., *J.O.C.*, 1991, 56, 457-459 (synth, ir, pmr)
Ding, C.Z. et al., *J. Med. Chem.*, 1993, 36, 1711-1715 (*Dimethylphenethylamine, synth, pmr, cmr*)
Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 2, 648
Liyanaage, G.K. et al., *J. Nat. Prod.*, 1996, 59, 148-151 (9-oxohexadecanoyl, 9-hydroxyhexadecanoyl)
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2155-2156
Sinz, A. et al., *Phytochemistry*, 1999, 50, 1069-1072 (*Coumaroylphenylethylamine*)
Luis-Astudillo, S. et al., *Planta Med.*, 1999, 65, 161-162 (activity)
Tan, L.T. et al., *J. Nat. Prod.*, 2000, 63, 952-955 (*Hermitamide A*)
Al-Rehaily, A.J. et al., *Phytochemistry*, 2001, 57, 597-602 (*N-Phenethylbenzamide, isol*)
Maskey, R.P. et al., *J. Antibiot.*, 2002, 55, 643-649 (*phenethylamides, isol*)
Elliot, M.C. et al., *Org. Biomol. Chem.*, 2003, 1, 3038-3047 (*N-formyl, synth, pmr, cmr*)
Shabaan, M. et al., *Dissertation*, Univ. of Göttingen, 2004, (*Cytophaga marinoflava amide*)
Shiono, Y. et al., *Nat. Prod. Res.*, 2005, 19, 363-366 (*N-hexadecanoyl*)
Virolleaud, M.-A. et al., *Tet. Lett.*, 2006, 47, 5127-5130 (*Hermitamide A, synth*)
Zhao, P.-J. et al., *Chem. Biodiversity*, 2007, 4, 899-904 (*13-methyltetradecanoyl, Ac*)
Frost, C.G. et al., *Org. Biomol. Chem.*, 2008, 6, 4340-4347 (*Hermitamide A, synth*)
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, PDE250; PDE500

N-(2-Phenylethyl)carbamic acid P-343

[152435-36-4]

PhCH₂CH₂NHCOOHC₉H₁₁NO₂ 165.191*Me ester: Methyl N-(2-phenylethyl)carbamate*

[26011-68-7]

C₁₀H₁₃NO₂ 179.218Solid (petrol) or oil. Mp 30°. Bp_{0.8} 120-123°.*Et ester: Ethyl N-(2-phenylethyl)carbamate*

[6970-83-8]

C₁₁H₁₅NO₂ 193.245Isol. from a culture medium of the marine bacterium *Cytophaga* sp. Inhibitor of biofilm formation and marine fouling. Isol. and props. not fully desc.*Butyl ester: Butyl N-(2-phenylethyl)carbamate. TMC 49A. Antibiotic TMC 49A*

[205594-32-7]

C₁₃H₁₉NO₂ 221.299Prod. by *Streptomyces* sp. AS1345. Transcriptional up-regulator of low density lipoprotein receptor. Oil. Sol. MeOH, EtOAc, DMSO, CHCl₃; poorly sol. H₂O, Me₂CO. Mp 6-7°. Bp 120-122°. λ_{max} 260 (ε 270) (MeOH).Rinkes, I.J. et al., *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1927, 40, 268-277 (*Me ester*)
Shriner, R.L. et al., *J.A.C.S.*, 1952, 74, 549-550 (*butyl ester, synth*)Yamada, A. et al., *Bull. Chem. Soc. Jpn.*, 1997, 70, 3061-3069 (*Et ester, isol*)

Koguchi, Y. *et al.*, *J. Antibiot.*, 1998, **51**, 107-111 (*butyl ester, isol, pmr, cmr, ms*)
 Salvatore, R.N. *et al.*, *Tetrahedron*, 2002, **58**, 3329-3347 (*Me ester*)

N-2-Phenylethylcinnamamide P-344

3-Phenyl-N-(2-phenylethyl)-2-propenamide, 9CI. *N-Cinnamoylphenethylamine* [55030-23-4]
 $\text{PhCH}=\text{CHCONHCH}_2\text{CH}_2\text{Ph}$
 $\text{C}_{17}\text{H}_{17}\text{NO}$ 251.327

(E)-form [103188-43-8]

Alkaloid from the leaves of *Spilanthes ocyimifolia* and aerial parts of *Simsia cronquistii* (Asteraceae), from leaves of *Clausena indica* and from *Oxytropis pseudoglandulosa* (Fabaceae). Needles (cyclohexane). Mp 125-126°.

N-Me: Lansiumamide C. *N-Methyl-N-phenylethylcinnamamide* [121817-38-7]
 $\text{C}_{18}\text{H}_{19}\text{NO}$ 265.354

Alkaloid from seeds of *Clausena lansium* (wampee) (Rutaceae). Yellowish plates (Et₂O). Mp 58-59°.

2,3-Dihydro: 3-Phenyl-N-(2-phenylethyl)propanamide [10264-31-0]
 $\text{C}_{17}\text{H}_{19}\text{NO}$ 253.343
 Isol. from *Simsia cronquistii*. Cryst. Mp 94-96°.

Borges-del-Castillo, J. *et al.*, *Phytochemistry*, 1984, **23**, 2671 (*isol, ir, pmr, cmr, ms, struct*)
 Huneck, S. *et al.*, *Fitoterapia*, 1986, **57**, 523 (*isol*)

Lin, J.H. *et al.*, *Phytochemistry*, 1989, **28**, 621 (*Lansiumamide C*)

Maldonado, E. *et al.*, *Phytochemistry*, 1992, **31**, 1413 (*3-Phenyl-N-(2-phenylethyl)propanamide*)

Riemer, B. *et al.*, *Phytochemistry*, 1997, **45**, 337 (*isol, uv, ir, pmr*)

Stefanuti, I. *et al.*, *Tet. Lett.*, 2000, **41**, 3735-3738 (*synth, Lansiumamide C*)

16-Phenyl-2,4-hexadecadienoic acid P-345

$\text{Ph}(\text{CH}_2)_{11}\text{CH}=\text{CHCH}=\text{CHCOOH}$
 $\text{C}_{22}\text{H}_{32}\text{O}_2$ 328.494


(2E,4E)-form

2-Methylpropylamide: N-Isobutyl-16-phenyl-2,4-hexadecadienamide [108331-97-1]
 $\text{C}_{26}\text{H}_{41}\text{NO}$ 383.616
 Constit. of *Piper brachystachyum* (*Piper mullesua*). Viscous solid. λ_{max} 218 ; 260 (MeOH).

Srivastava, S. *et al.*, *J. Indian Chem. Soc.*, 2000, **77**, 305-306 (*isol, pmr, cmr*)

5-Phenyl-2-oxazolidinone P-346

Barbarin. Resedinine



$\text{C}_9\text{H}_9\text{NOS}$ 179.242

(R)-form [3433-15-6]

Isol. from *Barbarea* spp. Hydrol. prod.

from Glucobarbarin. Tyrosinase inhibitor. Prisms (EtOH aq.). Mp 128-129°.
 $[\alpha]_{\text{D}}^{23}$ -69.3 (c, 2 in MeOH). λ_{max} 243 (log ϵ 4.26) (MeOH).

(S)-form [31456-67-4]


Hydrol. prod. isol. from genera *Reseda* and *Sibara*. Mp 126°. $[\alpha]_{\text{D}}^{23}$ +71 (c, 2 in MeOH).

(±)-form [85549-01-5]

Mp 139°.
 Kjaer, A. *et al.*, *Acta Chem. Scand.*, 1957, **11**, 906; 1958, **12**, 1693; 1970, **24**, 3031
 Tazhibaev, M.M. *et al.*, *CA*, 1976, **85**, 108843 (*isol*)
 Hirao, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1982, **55**, 1163 (*synth, ir, pmr*)
 Mieczkowski, J.B. *et al.*, *Bull. Pol. Acad. Sci., Chem.*, 1986, **34**, 109; *CA*, **107**, 78122p (*synth*)
 Moreno-Mañas, M. *et al.*, *J. Het. Chem.*, 1993, **30**, 1235 (*synth, pmr*)
 Seo, B. *et al.*, *Planta Med.*, 1999, **65**, 683-686 (*isol*)

5-Phenyl-2-oxazolidinone, P-347

9CI
Resedine
 [7693-77-8]



$\text{C}_9\text{H}_9\text{NO}_2$ 163.176
 ▶ LD₅₀ (mus, ivn) 316 mg/kg. RQ3710000

(±)-form [60426-44-0]

Constit. of *Reseda luteola*. Cryst. (C₆H₆), needles (EtOAc/hexane). Mp 88-89° Mp 120-121°.

3-Ph: 3,5-Diphenyl-2-oxazolidinone

[7426-72-4]
 $\text{C}_{15}\text{H}_{13}\text{NO}_2$ 239.273
 Needles (CHCl₃/petrol). Mp 78-79°.

Poos, G. *et al.*, *J. Med. Chem.*, 1963, **6**, 266 (*synth*)
 Foglia, T.A. *et al.*, *J.O.C.*, 1967, **32**, 75 (*synth*)
 Irwin, W.J. *et al.*, *J.C.S. (C)*, 1971, 3166 (*deriv, synth, ir, ms*)

Tazhibaev, M.M. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 270; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 245 (*isol*)

Fileenko, N.I. *et al.*, *CA*, 1979, **91**, 39376 (*synth*)
 Pirkle, W.H. *et al.*, *J.O.C.*, 1983, **48**, 2520 (*synth, pmr*)

Shibata, I. *et al.*, *J.O.C.*, 1986, **51**, 2177 (*deriv, synth*)
 Shen, Y.-M. *et al.*, *Eur. J. Org. Chem.*, 2004, 3080-3089 (*synth, pmr, cmr*)

3-Phenylloxiranecarboxylic acid, 9CI P-348

2,3-Epoxy-3-phenylpropanoic acid. β-Phenylglycidic acid [5694-02-0]
 [121-39-1]



$\text{C}_9\text{H}_8\text{O}_3$ 164.16

(2ξ,3ξ)-form

Methylamide: N-Methyl-3-phenylloxiranecarboxamide, 9CI. **Clausamide II** [189315-54-6]
 $\text{C}_{10}\text{H}_{11}\text{NO}_2$ 177.202
 Alkaloid from the leaves of *Clausena lansium*.

N-Methyl-N-(2-phenylethyl)amide:


Clausamide I [189315-53-5]
 [183812-58-0, 183812-57-9]
 $\text{C}_{18}\text{H}_{17}\text{NO}_2$ 279.338
 Alkaloid from the leaves of *Clausena lansium*.

[1566-68-3, 73610-79-4]

Li, S.H. *et al.*, *Chin. Pharm. J. (Taipei)*, 1996, **48**, 367-373 (*Clausamides*)

4-(5-Phenylpentyl)-1H-indole-2,3-dione P-349

4-(5-Phenylpentyl)isatin. Melosatin B [64838-03-5]



$\text{C}_{19}\text{H}_{19}\text{NO}_2$ 293.365
 Alkaloid from *Melochia tomentosa* (Sterculiaceae). Noncryst.

7-Methoxy: Melosatin C

[77410-47-0]
 $\text{C}_{20}\text{H}_{21}\text{NO}_3$ 323.391
 From *Melochia tomentosa* (Sterculiaceae). Red needles (C₆H₆/petrol). Mp 124-125°.

6,7-Dimethoxy: Melosatin A

[64838-04-6]
 $\text{C}_{21}\text{H}_{23}\text{NO}_4$ 353.417
 From *Melochia tomentosa* (Sterculiaceae). Mp 119-121°.

6,7-Dimethoxy, 5-hydroxy: Melosatin D

[155210-49-4]
 $\text{C}_{21}\text{H}_{23}\text{NO}_5$ 369.416
 From *Melochia tomentosa* (Sterculiaceae). Red needles. Mp 128-130°.

Kapadia, G.J. *et al.*, *Chem. Comm.*, 1977, 535 (*uv, ir, pmr, ms, struct*)

Kapadia, G.J. *et al.*, *Tetrahedron*, 1980, **36**, 2441 (*Melosatin C*)

Kapadia, G.J. *et al.*, *Planta Med.*, 1993, **59**, 568 (*Melosatin D*)

3-Phenylpropanoic acid P-350

Benzenepropanoic acid, 9CI. **Benzylacetic acid. Hydrocinnamic acid. Homotoluic acid. Dihydrocinnamic acid** [501-52-0]

$\text{PhCH}_2\text{CH}_2\text{COOH}$
 $\text{C}_9\text{H}_{10}\text{O}_2$ 150.177
 Prod. by *Clostridium butyricum*, also isol. from cultures of *Nicotiana tabacum*. Volatile component of tail gland secretion of red deer *Cervus elaphus*. Prisms (petrol). V. spar. sol. H₂O. Mp 48.5°. Bp 280° Bp₆ 125-129°. pK_a 4.66 (H₂O). Steam-volatile.

▶ LD₅₀ (mus, ivn) 3000-5000 mg/kg. DA8600000

Nitrile: Benzenepropanenitrile. Hydrocinnamonitrile. 1-Cyano-2-phenylethane

[645-59-0]

C₉H₉N 131.177

Constit. of *Brassica napus* and *Nasturtium officinale* (preferred genus name *Rorippa*). Bp 261° Bp₁₈ 114-118°.
 ▶ LD₅₀ (mus, oral) 116 mg/kg. MW5604750
 Gould, S.J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1239-1245 (nitrite, isol)

N-(3-Phenylpropanoyl)pyrrole P-351

1-(1-Oxo-3-phenylpropyl)-1H-pyrrole, 9CI. 3-Phenylpropanoic acid pyrrolidide [112448-69-8]

C₁₃H₁₃NO 199.252

Alkaloid from the fruit of *Piper sarmenosum*. Mp 48.5-50°.

4''-Hydroxy: 1-[3-(4-Hydroxyphenyl)-1-oxopropyl]-1H-pyrrole. N-[3-(4-Hydroxyphenyl)propanoyl]pyrrole. 3-(4-Hydroxyphenyl)propanoic acid pyrrolidide. **Piperlotine E**

[958296-14-5]

C₁₃H₁₃NO₂ 215.251

Alkaloid from the leaves of *Piper lolot*. Syrup. λ_{max} 229 ; 235 (sh) (MeOH).

3'',4''-Dimethoxy: 1-[3-(3,4-Dimethoxyphenyl)-1-oxopropyl]-1H-pyrrole. N-[3-(3,4-Dimethoxyphenyl)propanoyl]pyrrole. 3-(3,4-Dimethoxyphenyl)propanoic acid pyrrolidide [204587-76-8]

C₁₅H₁₇NO₃ 259.304

Alkaloid from *Piper brachystachyum*. Needles (petrol/CHCl₃). Mp 108°. λ_{max} 285 (MeOH).

Likhitwitayawuid, K. *et al.*, *Tetrahedron*, 1987, **43**, 3689-3694 (isol, synth, uv, ir, pmr, cmr, ms)


Kumar, R. *et al.*, *Acta Cryst. C*, 1998, **54**, 363-365 (dimethoxy deriv)

Parmar, V.S. *et al.*, *Phytochemistry*, 1998, **49**, 1069-1078 (dimethoxy deriv)

Li, C.-Y. *et al.*, *J. Agric. Food Chem.*, 2007, **55**, 9436-9442 (*Piperlotine E*)

4-Phenylpyridine, 9CI P-352

[939-23-1]

C₁₁H₉N 155.199

Occurs in tea and peppermint. Plates (H₂O). Sol. hot H₂O. Mp 77-78°. Bp 274-275°. pK_a 5.35 (25°).

▶ LD₅₀ (mus, ivn) 89 mg/kg. UT7141000

Picrate:

Orange needles. Mp 195-196°.

l-Oxide: [1131-61-9]C₁₁H₉NO 171.198Prisms (C₆H₆). Mp 152°.

l-Me: see 1-Methyl-4-phenylpyridinium(1+), #GVC30-V

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 739A; 917B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 249B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1519B (ir)

Elks, J. *et al.*, *J.C.S.*, 1943, 441 (synth)


Saeki, S. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 1780 (synth)

Godin, C.S. *et al.*, *Drug Metab. Dispos.*, 1989, **17**, 180 (metab)

Hassan, M.N. *et al.*, *Adv. Neurol.*, 1990, **53**, 219-223 (pharmacol)

Gromov, S.P. *et al.*, *Eur. J. Org. Chem.*, 2002, 4123-4126 (synth, pmr)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, PGE750


4-Phenyl-3-(2-pyridinyl)-2-buten-1-ol P-353C₁₅H₁₅NO 225.29**(E)-form** [915138-94-2]

Prod. by *Streptomyces* sp. KACC91010. Antiproliferative agent.

Shin, C. *et al.*, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 5643-5645 (isol, pmr, cmr)

2-Phenylquinoline, 9CI P-354

[612-96-4]

C₁₅H₁₁N 205.259

Alkaloid from the stem bark of *Galipea longiflora* (Rutaceae). Needles (EtOH aq.), cryst. (hexane). Sol. Et₂O, C₆H₆; spar. sol. H₂O; v. spar. sol. petrol. Mp 84°. Bp₆ 194-198°. Triboluminescent.

Methiodide: [14886-84-1]

Orange-red cryst. (EtOH/Et₂O). Mp 200°.

Picrate: [5278-66-0]

Yellow leaflets (EtOH). Mp 191-192°.

N-Oxide: [5659-33-6]C₁₅H₁₁NO 221.258

Cryst. (EtOAc). Mp 144°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 425A (nmr)

Filippi, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 268-275 (synth, uv, ir)

Koch, C.W. *et al.*, *J. Het. Chem.*, 1974, **11**, 475-479 (ms)

Crisp, G.T. *et al.*, *Aust. J. Chem.*, 1989, **42**, 279-285 (synth, cmr)

Fournet, A. *et al.*, *Can. J. Chem.*, 1989, **67**, 2116-2118 (isol, uv, pmr, ms)

Koyama, J. *et al.*, *Chem. Express*, 1991, **6**, 197-200; 1992, **7**, 321-324 (synth)

Ali, N.M. *et al.*, *Tetrahedron*, 1992, **37**, 8117-8126 (synth)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1038-1039 (synth, ir, pmr)

Horaguchi, T. *et al.*, *J. Het. Chem.*, 2002, **39**, 61-67 (synth, pmr, cmr)

Shi, D. *et al.*, *Synthesis*, 2005, 717-724 (synth, ir, pmr, cmr)

Han, R. *et al.*, *Heterocycles*, 2006, **68**, 1675-1684 (synth, ir, pmr, cmr)

Korn, T.J. *et al.*, *Synthesis*, 2006, 3547-3574 (synth, ir, pmr, cmr, ms)

Okuma, K. *et al.*, *Bull. Chem. Soc. Jpn.*, 2007, **80**, 1824-1827 (synth)

Hill, M.D. *et al.*, *Synthesis*, 2007, 1115-1119 (synth, ir, pmr)

Cho, S.-D. *et al.*, *Tetrahedron*, 2007, **63**, 1345-1352 (synth)

3-Phenylquinoline, 9CI P-355

[1666-96-2]

C₁₅H₁₁N 205.259

Alkaloid from *Peganum nigellastrum*. Plates (Et₂O). Sol. EtOH, Et₂O, CHCl₃, C₆H₆. Mp 49-52°. Bp 205-207° Bp_{0.1} 134°. Spar. steam-volatile.

Picrate:

Yellow cryst. (EtOH). Mp 205°.

Methiodide:

Yellow needles (MeOH). Mp 224°.

Cadogan, J.I.G. *et al.*, *J.C.S.*, 1962, 4257-4258 (synth)

Sobczyk, L. *et al.*, *CA*, 1963, **59**, 7345a (uv)


Sakamoto, T. *et al.*, *Tetrahedron*, 1993, **49**, 9713-9720 (synth, pmr)

Cacchi, S. *et al.*, *Tetrahedron*, 1996, **52**, 10225-10240 (synth, pmr, ir)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 332-334 (synth, ir, pmr)

Ma, Z.-Z. *et al.*, *Phytochemistry*, 2000, **53**, 1075-1078 (isol)

Fahkfeh, M.A. *et al.*, *Synth. Commun.*, 2002, **32**, 2863-2875 (synth, ir, pmr, cmr)

2-Phenyl-1,5,9-triazacyclotridecan-4-one, 9CI P-356C₁₆H₂₅N₃O 275.393**(S)-form***N*⁹-Ac: **N-Acetyl-N-deoxymayfoline**C₁₈H₂₇N₃O₂ 317.43

Alkaloid from leaves of *Maytenus buxifolia* (Celastraceae). Cryst. (EtOAc). Mp 177-178°. [α]_D²² -17.8 (c, 1.01 in CHCl₃). Called *N*¹-acetyl-*N*¹-deoxy in the lit.

*N*⁹-Benzoyl: **Celabenzine**

[53938-08-2]

C₂₃H₂₉N₃O₂ 379.501

Alkaloid from *Tripterygium wilfordii* and *Maytenus mossambicensis* var. *mossambicensis* (Celastraceae). Cryst. (EtOAc). Mp 163-167° (156-158°). No detectable opt. activity. λ_{max} 258 (sh) (ε 1390); 264 (sh) (ε 935); 270 (sh) (ε 685) (95% EtOH) (Derep).

*N*⁹-E-Cinnamoyl: **Celacinnine**

[53938-05-9]

C₂₅H₃₁N₃O₂ 405.539

Alkaloid from *Maytenus arbutifolia*,

Maytenus serrata, *Maytenus heterophylla*, *Tripterygium wilfordii* and *Pleurostyliya africana* (Celastraceae). Needles (hexane/CHCl₃). Mp 203-204°. [α]_D²⁵ +19 (c, 0.16 in CHCl₃). λ_{\max} 223 (sh) (ϵ 16000); 277 (ϵ 23000) (MeOH) (Derep).

N⁹-E-Cinnamoyl, N¹-Ac: Viburnine

[171236-03-6]
C₂₇H₃₃N₃O₃ 447.576
Alkaloid from *Viburnum rhytidophyllum*. Mp 115-118°. [α]_D²⁴ +67.7 (c, 0.939 in CHCl₃) (synthetic). [α]_D -2.6 (MeOH) (natural).

N⁹-Z-Cinnamoyl: Celalocinnine

[53990-48-0]
C₂₅H₃₁N₃O₂ 405.539
Alkaloid from *Maytenus arbutifolia*, *Maytenus serrata* and *Pleurostyliya africana* (Celastraceae). Needles (EtOAc/Et₂O/hexane). Mp 172-173°. [α]_D²⁵ -24 (c, 0.23 in CHCl₃). λ_{\max} 255 (ϵ 11800); 264 (sh) (MeOH) (Derep).

N⁹-(3-Furancarboxyl): Celafurine

[53938-09-3]
[79435-41-9]
C₂₁H₂₇N₃O₃ 369.463
Alkaloid from the roots of *Tripterygium wilfordii* (Celastraceae). Cryst. (EtOAc). Mp 154-155°. [α]_D²⁵ -11 (c, 0.11 in CHCl₃). λ_{\max} 222 (sh) (ϵ 232 (sh); 285 (ϵ 1230) (95% EtOH) (Derep).

N⁹-Hydroxy: Mayfoline

[74133-16-7]
C₁₆H₂₅N₃O₂ 291.392
Alkaloid from the aerial parts of *Maytenus buxifolia* (Celastraceae). Cryst. (EtOAc). Mp 200-204°. [α]_D²² -52.3 (c, 0.636 in CHCl₃) (the original reported value of +10.6 is considered erroneous).

3S-Hydroxy, N⁹-E-cinnamoyl: 3-Hydroxycelacinnine

[145199-54-8]
C₂₅H₃₁N₃O₃ 421.538
Minor alkaloid from leaves of *Pleurostyliya opposita* (Celastraceae). Amorph. [α]_D 0 (c, 1 in CHCl₃). Abs. config. is (2R,3S) owing to change in R,S-priorities.

(±)-form

N⁹-Benzoyl: [73465-25-5]

Synthetic. Cryst. (EtOAc). Mp 190-194° (softens then solidifies at 147-148°) (170-173°).

N⁹-E-Cinnamoyl:

Synthetic. Cryst. (EtOAc/hexane or C₆H₆). Mp 166-169° Mp 178-181°.

N⁹-Z-Cinnamoyl: [79435-40-8]

Synthetic. No Mp recorded.

Kupchan, S.M. *et al.*, *J.O.C.*, 1977, **42**, 3660 (Celabenzine, Celafurine, Celacinnine, *isol*, *uv*, *cd*, *ir*, *pmr*, *ms*, *struct*)

McManis, J.S. *et al.*, *J.O.C.*, 1980, **45**, 2041 (Celabenzine, Celacinnine, *synth*)

Ripperger, H. *et al.*, *Phytochemistry*, 1980, **19**, 162 (Mayfoline)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1980, **21**, 3493 (Celacinnine, *synth*, *ir*, *pmr*)

Wagner, H. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 283; 1982, **65**, 739 (Celabenzine, Celacinnine, *uv*, *ir*, *pmr*, *cmr*, *ms*, *synth*)

Yamamoto, H. *et al.*, *J.A.C.S.*, 1981, **103**, 6133 (*synth*, *ir*, *pmr*, *ms*)

Diaz, M. *et al.*, *Phytochemistry*, 1982, **21**, 255 (Mayfoline, N-Acetyl-N-deoxymayfoline)

Iida, H. *et al.*, *Tet. Lett.*, 1986, **27**, 207

(Celabenzine, *synth*, *ir*, *pmr*, *cd*)

Séguineau, C. *et al.*, *Helv. Chim. Acta*, 1992, **75**, 2283 (3-Hydroxycelacinnine)

Tawil, B.F. *et al.*, *Tetrahedron*, 1992, **48**, 3775

(N-Acetyl-N-deoxymayfoline, *synth*)

Begley, M.J. *et al.*, *J.C.S. Perkin 1*, 1993, 2027

(Celabenzine, *synth*)

Itoh, N. *et al.*, *Heterocycles*, 1995, **41**, 415

(Celacinnine, *synth*)

Abdallah, O.M. *et al.*, *CA*, 1996, **124**, 4885u

(Viburnine)

Kuehne, P. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 1085; 1997, **80**, 1802-1808 (Celabenzine,

Mayfoline, N-Acetyl-N-deoxymayfoline,

Celafurine, Celacinnine, *synth*)

Matsuyama, H. *et al.*, *J.C.S. Perkin 1*, 2001,

2924-2930 (Celacinnine, *synth*)

Khanjin, N.A. *et al.*, *Helv. Chim. Acta*, 2003,


86, 2028-2057 (3-Hydroxycelacinnine, *synth*,

abs config)

Phidolopin

P-357

3,7-Dihydro-7-[(4-hydroxy-3-nitrophenyl)methyl]-1,3-dimethyl-1H-purine-2,6-dione, 9CI. N⁷-(4-Hydroxy-3-nitrobenzyl)theophylline [92014-27-2]



C₁₄H₁₃N₅O₅ 331.287

Found in the marine bryozoan

Phidolopora pacifica. Shows

antifungal and anti-algal props. Cryst.

(MeOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 226-227°. λ_{\max} 275 (ϵ 16800); 353 (ϵ 3300) (MeCN) (Derep).

N¹,N³-Di-de-Me: Desmethylphidolopin

[103545-34-2]

C₁₂H₉N₅O₅ 303.234

From the marine bryozoans *Phidolopora pacifica* and *Diaperoecia californica*. Antifungal agent. Yellow amorph. powder.

Ayer, S.W. *et al.*, *J.O.C.*, 1984, **49**, 3869-3870 (*isol*, *uv*, *ir*, *pmr*, *ms*, *cryst struct*)

Hirota, K. *et al.*, *Tet. Lett.*, 1985, **26**, 2355

(*synth*)

Tischler, M. *et al.*, *Comp. Biochem. Physiol. B: Comp. Biochem.*, 1986, **84**, 43 (*deriv*)


Avasthi, K. *et al.*, *Indian J. Chem., Sect. B*, 1996, **35**, 437 (*synth*, *pmr*, *cmr*, *ms*)

Philanthotoxin 433

P-358

δ Philanthotoxin. PhTX-433

[115976-91-5]



C₂₃H₄₁N₅O₃ 435.609

Isol. from the venom of the solitary digger wasp *Philanthus triangulum*. Glutamate receptor inhibitor. Oil or amorph. powder.

Eldefrawi, A.T. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1988, **85**, 4910 (*isol*, *pharmacol*)

Rozental, R. *et al.*, *J. Pharmacol. Exp. Ther.*,

1989, **249**, 123-130 (*pharmacol*)

Nakanishi, K. *et al.*, *Pure Appl. Chem.*, 1990,

62, 1223 (*isol*, *synth*)

Goodnow, R. *et al.*, *Tetrahedron*, 1990, **46**,

3267 (*synth*)

Wang, F. *et al.*, *Org. Lett.*, 2000, **2**, 1581-1583

(*synth*)

Chhabra, S.R. *et al.*, *Tet. Lett.*, 2000, **41**, 1095-

1098; 1099-1102 (*synth*)


Stromgard, K. *et al.*, *Magn. Reson. Chem.*,

2006, **44**, 1013-1022 (*cmr*)

Philippinamide

P-359

11-(4-Methoxyphenyl)-N-(2-methylpropyl)-2,4,10-undecatrienamide. 11-(4-Methoxyphenyl)-2,4,10-undecatrienoic acid isobutylamide



C₂₂H₃₁NO₂ 341.492


(E,E,E)-form [949102-31-2]

Alkaloid from the stems of *Piper philippinum*. Needles (MeOH). Mp 135.5-136.5°. λ_{\max} 261 (log ϵ 3.62) (MeOH).

Chen, Y.-C. *et al.*, *Phytochemistry*, 2007, **68**, 2101-2111 (*isol*, *pmr*, *cmr*, *ms*)

Philogaline

P-360



C₃₅H₃₂N₂O₆ 576.648


Alkaloid from *Guatteria boliviana*. Amorph. λ_{\max} 205 (log ϵ 4.38); 231 (sh) (log ϵ 4.26); 284 (log ϵ 3.89) (EtOH).

Mahiou, V. *et al.*, *Phytochemistry*, 2000, **54**, 709-716

Phlegmarine

P-361

[66834-87-5]



Probable relative configuration

C₁₆H₃₀N₂ 250.426Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Noncryst.N^a-Me: N^a-Methylphlegmarine

[66834-88-6]

C₁₇H₃₂N₂ 264.453Alkaloid from *Lycopodium cernuum* (Lycopodiaceae). Oil.N^b-Me: N^b-Methylphlegmarine

[66834-89-7]

C₁₇H₃₂N₂ 264.453Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Oil.N^b-Me, N^a-Ac: N^a-Acetyl-N^b-methylphlegmarineC₁₉H₃₄N₂O 306.49Alkaloid from *Lycopodium clavatum* var. *borbonicum* (Lycopodiaceae). Oil. [α]_D²⁵ -11 (c, 0.7 in CHCl₃).

N,N'-Di-Me: N,N'-Dimethylphlegmarine


[66834-90-0]

C₁₈H₃₄N₂ 278.48Alkaloid from *Lycopodium phlegmaria*, also present in *Lycopodium clavatum*. Oil. [α]_D²⁵ -21 (c, 1.13 in CHCl₃) (synthetic). Alkaloid L18 from *L. clavatum* was a mixt. of N,N'-Dimethylphlegmarine and an isomeric compd.Nyembo, L. et al., *Can. J. Chem.*, 1978, **56**, 851 (isol, ir, pmr, ms, struct)Leniewski, A. et al., *Can. J. Chem.*, 1981, **59**, 2479; 2695 (N^a-Methylphlegmarine, synth, config)Inubushi, Y. et al., *Yakugaku Zasshi*, 1982, **102**, 434; *CA*, **97**, 107030q (N,N'-Dimethylphlegmarine)Ayer, W.A. et al., *Can. J. Chem.*, 1990, **68**, 1300 (Alkaloid L18)Comins, D.L. et al., *J.O.C.*, 1999, **64**, 2184-2185 (synth)

Phlegmariurine A

P-362

[93754-82-6]



Absolute Configuration

C₁₆H₂₃NO₂ 261.363Alkaloid from *Phlegmariurus fordii* whole plant (preferred genus name *Huperzia*) (Lycopodiaceae).

4-Hydroxy: Lycoposerramine E


[481048-27-5]

C₁₆H₂₃NO₃ 277.363Alkaloid from *Lycopodium serratum*. Needles (EtOH). Mp 259-262° subl.Tong, S. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1984, **26**, 411-415; *CA*, **102**, 3253v (isol, struct)Wan, Z. et al., *Kexue Tongbao (Foreign Lang. edn.)*, (Foreign Lang. Ed.), 1986, **31**, 489-492; *CA*, **106**, 5290m (cryst struct, abs config)Takayama, H. et al., *Tet. Lett.*, 2002, **43**, 8307-8311 (cmr, Lycoposerramine E)

Phlegmariurine B

P-363

[93754-83-7]

C₁₆H₂₃NO₂ 261.363Alkaloid from *Phlegmariurus fordii* (preferred genus name *Huperzia*) and *Huperzia serrata*. Prisms (Me₂CO). Mp 110-111°.

2α-Hydroxy: 2α-Hydroxyphlegmariurine B

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*. Prisms (Me₂CO). Mp 240-241°. [α]_D²⁵ -0.2 (c, 0.17 in CHCl₃).

7-Hydroxy: 7-Hydroxyphlegmariurine B

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*. Prisms (Me₂CO). Mp 214-216°. [α]_D²⁵ -2.2 (c, 0.17 in MeOH).

7-Hydroperoxy: 7-Hydroperoxyphlegmariurine B

C₁₆H₂₃NO₄ 293.362Alkaloid from *Huperzia serrata*. Prisms (Me₂CO/MeOH). Mp 188-190°. [α]_D²⁵ -1.8 (c, 0.67 in MeOH).

8β-Hydroxy: 8β-Hydroxyphlegmariurine B

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*. [α]_D²⁵ -205 (c, 0.2 in EtOH).

11α-Hydroxy: 11α-Hydroxyphlegmariurine B

C₁₆H₂₃NO₃ 277.363Alkaloid from *Huperzia serrata*. Prisms (Me₂CO/MeOH). Mp 246-248°. [α]_D²⁵ -1.2 (c, 0.67 in MeOH).

11α-Hydroperoxy: 11α-Hydroperoxyphlegmariurine B

C₁₆H₂₃NO₄ 293.362Alkaloid from *Huperzia serrata*. Prisms (Me₂CO/MeOH). Mp 164-166°. [α]_D²⁵ -2.9 (c, 0.58 in MeOH).

7,11α-Dihydroxy: 7,11α-Dihydroxyphlegmariurine B

C₁₆H₂₃NO₄ 293.362Alkaloid from *Huperzia serrata*. Prisms (Me₂CO/MeOH). Mp 265-267°. [α]_D²⁵ -1.7 (c, 0.42 in MeOH).

2-Oxo: 2-Oxophlegmariurine B

C₁₆H₂₁NO₃ 275.347Alkaloid from *Huperzia serrata*. Amorph. powder. [α]_D²⁵ -2.2 (c, 0.03 in CHCl₃).


11-Oxo: 11-Oxophlegmariurine B

C₁₆H₂₁NO₃ 275.347Alkaloid from *Huperzia serrata*. Prisms (Me₂CO/petrol). Mp 233-235°. [α]_D²⁵ -5.1 (c, 0.35 in CHCl₃).Tan, C.-H. et al., *J. Asian Nat. Prod. Res.*, 2002, **4**, 227-231 (isol, cd, pmr, cmr)Tan, C.-H. et al., *Planta Med.*, 2002, **68**, 188-190 (isol, cd, pmr, cmr)Tan, C.-H. et al., *Acta Bot. Sin.*, 2003, **45**, 118-121 (7-Hydroperoxyphlegmariurine B, 11-Hydroperoxyphlegmariurine B)Yuan, S.-Q. et al., *Yaoxue Xuebao*, 2003, **38**, 596-598 (8-Hydroxyphlegmariurine B)

Phlegmariurine C

P-364

[115491-58-2]




Relative Configuration

C₃₂H₄₄N₂O₄ 520.711The full struct. shown does not appear to have been publ., but is given on a Scripps internal website. Alkaloid from whole plants of *Phlegmariurus fordii* (preferred genus name *Huperzia*). Mp 151-152°.Chu, B.M. et al., *Yaoxue Xuebao*, 1988, **23**, 115-121; *CA*, **109**, 70347m (isol)

Phlegmariurine M

P-365


[125287-10-7]

C₁₇H₂₀N₂O₃ 300.357Alkaloid from *Phlegmariurus fordii* (preferred genus name *Huperzia*) (Lycopodiaceae).Miao, Z. et al., *Huaxue Xuebao*, 1989, **47**, 702; *CA*, **112**, 115700b

Phloeodictyne A

P-366

Phloeodictine A



Naming system and numbering revised in 2004. Under the revised numbering scheme, the number before the comma indicates *m*, the number after the comma indicates *n*, and the suffix *a* or *i* indicates the end-group *R* (*a* = allyl, -CH₂CH=CH₂, *i* = isopropyl, -CH(CH₃)₂). The +ve charge is delocalised over the two ring nitrogens. Isol. from the New Caledonian deep water sponge *Phloeodictyon* sp. and shallow-water sponge *Oceanapia fistulosa* (*Phloeodictyon fistulosa*). Shows *in vitro* antibacterial activity and mod. cytotoxicity against KB cells. Shows antiplasmodial props.

Phloeodictyne 4,5a

Phloeoictyne A4
[155070-26-1]

$C_{22}H_{41}N_5O$ 391.599
 λ_{max} 224 (ϵ 6700); 274 (ϵ 2200) (MeOH)
(Berdy).

Phloeoictyne 4,5i

$C_{22}H_{43}N_5O$ 393.615

Phloeoictyne 4,6a

$C_{23}H_{43}N_5O$ 405.626

Phloeoictyne 4,6i

$C_{23}H_{45}N_5O$ 407.641

Phloeoictyne 4,7a

Phloeoictyne A2
[155070-28-3]

$C_{24}H_{45}N_5O$ 419.652
Amorph. solid (as dichloride). Mixt. with
Phloeoictyne A1. λ_{max} 224 (ϵ 6700); 274
(ϵ 2200) (MeOH) (Berdy).

Phloeoictyne 4,7i

$C_{24}H_{47}N_5O$ 421.668

Phloeoictyne 4,8a

$C_{25}H_{47}N_5O$ 433.679

Phloeoictyne 4,8i

Phloeoictyne A7
[155070-24-9]

$C_{25}H_{49}N_5O$ 435.695
 λ_{max} 224 (ϵ 6700); 274 (ϵ 2200) (MeOH)
(Berdy).

Phloeoictyne 4,9a

Phloeoictyne A (obsol.)†
[142260-80-8]

$C_{26}H_{49}N_5O$ 447.706
Amorph. solid (as dichloride). λ_{max} 224
(ϵ 6700); 274 (ϵ 2200) (MeOH) (Derep).

Phloeoictyne 4,10a

$C_{27}H_{51}N_5O$ 461.733

Phloeoictyne 4,10i

$C_{27}H_{53}N_5O$ 463.749

Phloeoictyne 4,11a

$C_{28}H_{53}N_5O$ 475.76

Phloeoictyne 5,4a

Phloeoictyne A5
[155070-27-2]

$C_{22}H_{41}N_5O$ 391.599
 λ_{max} 224 (ϵ 6700); 274 (ϵ 2200) (MeOH)
(Berdy).

Phloeoictyne 5,4i

$C_{22}H_{43}N_5O$ 393.615

Phloeoictyne 5,5a

Phloeoictyne A3
[155112-61-1]

$C_{23}H_{43}N_5O$ 405.626
Amorph. solid (as dichloride). Initially
isol. as a *ca.* 2.6:0.7:0.3 insep. mixt.
with Phloeoictyenes A4 and A5. λ_{max}
224 (ϵ 6700); 274 (ϵ 2200) (MeOH)
(Berdy).

Phloeoictyne 5,5i

$C_{23}H_{45}N_5O$ 407.641

Phloeoictyne 5,6i

$C_{24}H_{47}N_5O$ 421.668

Phloeoictyne 5,7a

Phloeoictyne A1
[155070-29-4]

$C_{25}H_{47}N_5O$ 433.679
Amorph. solid (as dichloride). Isol. as a
ca. 2.6:1 insep. mixt. with Phloeoictyne
A2. λ_{max} 224 (ϵ 6700); 274 (ϵ 2200)
(MeOH) (Berdy).

Phloeoictyne 5,7i

$C_{25}H_{49}N_5O$ 435.695

Phloeoictyne 5,8a [690654-91-2]

$C_{26}H_{49}N_5O$ 447.706

Phloeoictyne 5,8i

Phloeoictyne A6
[155070-25-0]

$C_{26}H_{51}N_5O$ 449.722
Amorph. solid (as dichloride).
Initially isol. as a *ca.* 1:1.4 insep.
mixt. with Phloeoictyne A7. λ_{max} 224
(ϵ 6700); 274 (ϵ 2200) (MeOH)
(Berdy).

Phloeoictyne 5,9a

$C_{27}H_{51}N_5O$ 461.733

Phloeoictyne 5,9i

$C_{27}H_{53}N_5O$ 463.749

Phloeoictyne 5,10a

$C_{28}H_{53}N_5O$ 475.76


Phloeoictyne 5,10i

$C_{28}H_{55}N_5O$ 477.775

Kourany-Lefoll, E. *et al.*, *Tetrahedron*, 1994,
50, 3415-3426 (*isol*)
Neubert, B.J. *et al.*, *Org. Lett.*, 2003, **5**, 765-
768 (*synth*)
Mancini, I. *et al.*, *Org. Biomol. Chem.*, 2004, **2**,
783-787 (*isol, ms*)

Phloeoictyne B

[142260-81-9]




$C_{27}H_{53}N_8OS^{3+}$ 537.835

Isol. from a New Caledonian sponge
Phloeoictyon sp. Amorph. solid (as
trichloride). CAS no. refers to trichloride.
 λ_{max} 202 (ϵ 5500); 279 (ϵ 7100) (MeOH)
(Derep).

Kourany-Lefoll, E. *et al.*, *J.O.C.*, 1992, **57**,
3832 (*isol, uv, pmr, cmr, cd, struct*)

Phloeoictyne C

P-368



Phloeoictyne C1, $n = 5$
C2, $n = 4$

Exhibits *in vitro* antibacterial activity and
is moderately cytotoxic against KB cells.

Phloeoictyne C1 [155070-23-8]

$C_{28}H_{57}N_8OS$ 553.877
Alkaloid from the New Caledonian
sponge *Phloeoictyon* sp. Amorph. solid
(as trichloride) (mixt. with C2). Isol. as a
ca. 1:1 insep. mixt. of homologues. λ_{max}
219 (ϵ 9100) (MeOH) (Berdy).

Phloeoictyne C2 [155070-22-7]


$C_{27}H_{55}N_8OS$ 539.85
From *Phloeoictyon* sp. Amorph. solid
(as trichloride) (mixt. with C1). λ_{max} 219
(ϵ 9100) (MeOH) (Berdy).

Kourany-Lefoll, E. *et al.*, *Tetrahedron*, 1994,
50, 3415 (*isol, uv, ir, pmr, cmr, struct*)

Phoebine

P-369

*1,2,3-Trimethoxy-9,10-methylenedioxy-
porphine*



$C_{21}H_{23}NO_5$ 369.416

(S)-form

Major alkaloid from the leaves of *Phoebe
valeriana* (Lauraceae). Cryst. (Me_2CO).
Mp 114°. $[\alpha]_D^{24} +49.6$ (c, 6.2 in $CHCl_3$).

N-Me: N-Methylphoebine

$C_{22}H_{26}NO_5^{\oplus}$ 384.451
Quaternary alkaloid from the bark and
roots of *Xylopia parviflora*. Amorph.
powder (as perchlorate). $[\alpha]_D^{22} +37.1$ (c,
0.7 in MeOH). λ_{max} 283 (log ϵ 4.03);
309 (log ϵ 4.02) (MeOH).

**N-De-Me, N-Ac: 6-Acetyl-1,2,3-tri-
methoxy-9,10-methylenedioxy-norapor-
phine. 3-Methoxy-N-acetyl-norantennine**
[53452-28-1]

$C_{22}H_{23}NO_6$ 397.427
Alkaloid from the heartwood of *Lir-
iodendron tulipifera* (Magnoliaceae).
Cryst. (EtOH). Mp 216-217°. $[\alpha]_D^{26}$
+271 (c, 0.37 in $CHCl_3$).

**O³,N-Di-de-Me: 3-Hydroxy-1,2-
dimethoxy-9,10-methylenedioxy-nora-
porphine**

C₁₉H₁₉NO₅ 341.363

Alkaloid from the leaves of *Phoebe valeriana* (Lauraceae). Amorph. solid. Opt. rotn. not reported. It is likely that the *N*-Me deriv. of this alkaloid is also present in the plant.

6a,7-Didehydro: 1,2,3-Trimethoxy-9,10-methylenedioxydehydroaporphine

C₂₁H₂₁NO₅ 367.401

Alkaloid from the leaves of *Phoebe valeriana* (Lauraceae). Amorph. solid.

(±)-form

N-De-Me, N-Ac: [57236-56-3]

Synthetic. Cryst. (EtOH). Mp 174-175°.

(ξ)-form

N-De-Me: 1,2,3-Trimethoxy-9,10-methylenedioxyaporphine

[95523-04-9]

C₂₀H₂₁NO₅ 355.39

Alkaloid from the wood of *Phoebe pittieri* (Lauraceae). Oil. Not opt. rotn. recorded.

O¹,N-Di-de-Me: 1-Hydroxy-2,3-dimethoxy-9,10-methylenedioxyaporphine. 3-MethoxynordomesticineC₁₉H₁₉NO₅ 341.363

Alkaloid from bark of *Nectandra sinuata* (Lauraceae). Yellow oil. No info. on opt. rotn. or abs. config.

O²,N-Di-de-Me: 2-Hydroxy-1,3-dimethoxy-9,10-methylenedioxyaporphine. Xyloguyelline

[80151-83-3]


C₁₉H₁₉NO₅ 341.363

Alkaloid from the trunk bark of *Xylopia danguyella* (Annonaceae). Tentative struct; the 2-OMe, 3-OH struct. (see above) cannot be excluded.

Hufford, C.D. *et al.*, *J. Pharm. Sci.*, 1974, **63**, 1338; 1975, **64**, 789 (*isol, uv, ir, pmr, cd, struct, 3-Methoxy-N-acetylnornantene*)Hufford, C.D. *et al.*, *J.O.C.*, 1976, **41**, 375 (*synth, 3-Methoxy-N-acetylnornantene*)Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551 (*Xyloguyelline*)Castro, C.O. *et al.*, *Phytochemistry*, 1985, **24**, 203 (*1,2,3-Trimethoxy-9,10-methylenedioxyaporphine*)Castro, O. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1036 (*isol, uv, pmr, cmr, ms, struct, derivs*)Castro, O. *et al.*, *Fitoterapia*, 1991, **62**, 72 (*3-Methoxynordomesticine*)Nishiyama, Y. *et al.*, *Phytochemistry*, 2004, **65**, 939-944 (*N-Methylphoebine*)

Phoenicanthusine

P-370



Relative Configuration


C₃₈H₃₂N₂O₆ 612.681

Dimeric aporphine. Alkaloid from the stem bark of *Phoenicanthus obliqua*. Pale yellow cryst. Mp 261-263°. λ_{max} 214 (log ε 4.24); 264 (log ε 4.6); 328 (log ε 3.28) (EtOH).

Kithsiri Wijeratne, E.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1465-1467

Phomacin A

[187847-10-5]

C₂₅H₃₇NO₅ 431.571

Isol. from the fungus *Phoma* sp. Powder. Sol. MeOH, EtOAc. Mp 130-133°. [α]_D -91 (c, 1 in CHCl₃).

17-Deoxy, 25-hydroxy: Phomacin B

[187847-11-6]


C₂₅H₃₇NO₅ 431.571

Isol. from a *Phoma* sp. Powder. Mp 98-100°. [α]_D -51.2 (c, 1 in CHCl₃).

Alvi, K.A. *et al.*, *J.O.C.*, 1997, **62**, 2148-2151 (*isol, ir, pmr, cmr*)

Phomacin C

[187847-12-7]


C₂₅H₃₇NO₄ 415.572

Isol. from the fungus *Phoma* sp. Powder. Mp 112-114°. [α]_D -74.6 (c, 1 in CHCl₃).

Alvi, K.A. *et al.*, *J.O.C.*, 1997, **62**, 2148-2151 (*isol, ir, pmr, cmr*)

Phomasetin

P-373

C₂₅H₃₅NO₄ 413.556

Tetramic acid antibiotic. Related to Equisetin, E-142, but possessing opposite stereochemistry. Enolised triketone. Prod. by a *Phoma* sp. Inhibitor of HIV-1 integrase. Cryst. (C₆H₆). Sol. MeOH,

P-371

CHCl₃; poorly sol. H₂O. Mp 140-143°.


[α]_D²² +93.9 (c, 1 in CHCl₃). λ_{max} 235 (ε 7540); 290 (ε 5420) (MeOH). λ_{max} 235 (ε 7541); 290 (ε 5422) (MeOH) (Berdy).

Singh, S.B. *et al.*, *Tet. Lett.*, 1998, **39**, 2243-2246 (*isol, uv, ir, cd, pmr, cmr, ms*)Hazuda, D. *et al.*, *Antiviral Chem. Chemother.*, 1999, **10**, 63-70 (*isol, activity*)

Phomazarine

P-374

[10088-99-0]

C₁₉H₁₇NO₈ 387.345

Struct. has been revised twice. Metab. of fungus *Phoma terrestris*. Orange needles (Py). Mp 196° dec. λ_{max} 231 (ε 34673); 277 (ε 58884); 430 (ε 8317) (MeOH) (Berdy).

Me ester:

Orange-bronze needles. Mp 213°.

Tri-Ac:

Cryst. (petrol). Mp 160-170° dec.


Kogl, F. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1940, **59**, 1180; 1944, **63**, 251; 1945, **64**, 33Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 2471-2477 (*synth, struct*)

Phomoenamide

P-375

N,N'-(1,3-Butadiene-1,4-diyl)bis[2-hydroxy-3-methylbutanamide]

[1015078-56-4]

C₁₄H₂₄N₂O₄ 284.355


Prod. by *Phomopsis* sp. PSU-D15. Gum. [α]_D²⁶ +32.6 (c, 0.13 in CHCl₃). λ_{max} 260 (log ε 3.97) (MeOH).

Rukachaisirikul, V. *et al.*, *Phytochemistry*, 2008, **69**, 783-787 (*isol, pmr, cmr, ms*)

Phomopsichalasin

P-376

[163597-03-3]



Relative Configuration

C₃₂H₄₁NO₄ 503.68

Unique cytochalasin-type struct. Isol. from an endophytic *Phomopsis* sp. Antimicrobial agent, mycotoxin. Oil. $[\alpha]_D^{25}$ -7.16. λ_{\max} 225 (ϵ 1152); 245 (ϵ 2782) (MeOH) (Berdy).

8,13,16,21,22-Pentaepimer: **Diaporthichalasin**


C₃₂H₄₁NO₄ 503.68

Isol. from *Diaporthe* sp. Bkk3. Potent CYP3A4 inhibitor. Mp 169-170°. $[\alpha]_D^{20}$ -135 (c, 0.14 in MeOH). λ_{\max} 233 (log ϵ 3.51); 277 (log ϵ 3.04) (MeOH).

Horn, W.S. *et al.*, *Tetrahedron*, 1995, **51**, 3969-3978 (*Phomopsichalasin*)

Pornpakakul, S. *et al.*, *Tet. Lett.*, 2007, **48**, 651-655 (*Diaporthichalasin*)


Phomopsin A† P-377
4-Hydroxy-1-(3-hydroxyphenyl)-3(2H)-isoquinolinone

C₁₅H₁₁NO₃ 253.257

Prod. by the mangrove fungus *Phomopsis* sp. ZZ08. Amorph. solid. Mp 250-252°. λ_{\max} 222 (log ϵ 3.62); 286 (log ϵ 2.91); 308 (log ϵ 2.91); 318 (2.99); 330 (log ϵ 2.86) (MeOH).

Tao, Y. *et al.*, *Magn. Reson. Chem.*, 2008, **46**, 501-505 (*isol, pmr, cmr*)

Phorbasin E P-378
[1023993-34-1]

C₄₂H₆₃NO₈S 742.028

Alkaloid from a *Phorbas* sp. Amorph. solid. $[\alpha]_D^{22}$ -27.9 (c, 0.16 in MeOH). λ_{\max} 242 (log ϵ 4.52) (MeOH).

1-Ac: **Phorbasin F**


[1023993-35-2]

C₄₄H₆₅NO₉S 784.065

Alkaloid from a *Phorbas* sp. Amorph. solid. $[\alpha]_D^{22}$ -10.1 (c, 0.1 in MeOH). λ_{\max} 242 (log ϵ 4.39) (MeOH).

Zhang, H. *et al.*, *Org. Lett.*, 2008, **10**, 1959-1962 (*isol, pmr, cmr*)

Phorbazole D P-379
[156280-94-3]

C₁₃H₉ClN₂O₂ 260.679

Isol. from the marine sponge *Phorbas* aff. *clathrata*. Immunomodulator. Amorph. powder. Mp 210°.

3-Chloro: **Phorbazole C**

[156280-93-2]

C₁₃H₈Cl₂N₂O₂ 295.124

From *Phorbas* aff. *clathrata*. Immunomodulator. Amorph. powder. Mp 240°.

2,3-Dichloro: **Phorbazole B**

[156280-92-1]

C₁₃H₇Cl₃N₂O₂ 329.569

From *Phorbas* aff. *clathrata*. Immunomodulator. Amorph. off-white powder. Mp 270°.

3,9-Dichloro: **Phorbazole A**


[156280-91-0]

C₁₃H₇Cl₃N₂O₂ 329.569

From *Phorbas* aff. *clathrata*. Immunomodulator. Powder. Mp 240°.

Rudi, A. *et al.*, *Tet. Lett.*, 1994, **35**, 2589 (*isol, ir, pmr, cmr, struct*)

Phorbaxazole A P-380
[165883-76-1]

C₅₃H₇₁BrN₂O₁₃ 1024.054

Isol. from the sponge *Phorbas* sp. Antifungal and cytostatic agent. Pale yellow solid. $[\alpha]_D$ +44.8 (c, 1 in MeOH). λ_{\max} 235 (log ϵ 4.9) (MeOH).

13-Epimer: **Phorbaxazole B**

[165689-31-6]

C₅₃H₇₁BrN₂O₁₃ 1024.054

Isol. from *Phorbas* sp. Antifungal and cytostatic agent. Pale yellow solid. $[\alpha]_D$ +44.4 (c, 1 in MeOH). λ_{\max} 235 (log ϵ 4.8) (MeOH).

Searle, P.A. *et al.*, *J.A.C.S.*, 1995, **117**, 8126 (*isol, uv, ir, pmr, cmr, ms*)

Searle, P.A. *et al.*, *J.A.C.S.*, 1996, **118**, 9422 (*abs config*)

Molinski, T.F. *et al.*, *Tet. Lett.*, 1996, **37**, 7879 (*abs config*)

Evans, D.A. *et al.*, *J.A.C.S.*, 2000, **122**, 10033-10046 (*synth*)

Smith, A.B. *et al.*, *J.A.C.S.*, 2001, **123**, 10942-10953 (*synth, bibl*)

Williams, D.R. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 1258-1262 (*synth*)

Haustedt, L.O. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 2711-2716 (*rev, synth*)

Pattenden, G. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 4173-4208 (*synth*)


Li, D.-R. *et al.*, *Chem. Eur. J.*, 2006, **12**, 1185-1204 (*synth*)

White, J.D. *et al.*, *Org. Lett.*, 2006, **8**, 6043-6046 (*synth*)

Lucas, B.S. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 769-772 (*Phorbaxazole B, synth*)

Smith, A.B. *et al.*, *J.O.C.*, 2008, **73**, 1192-1200; 1201-1208 (*synth*)

Photobactin P-381


C₂₂H₂₅N₃O₇ 443.455

Prod. by the bacterium *Photobacterium luminescens*; associated with the nematode *Heterorhabditis bacteriophora*. Catechol siderophore.

Ciche, T.A. *et al.*, *Appl. Environ. Microbiol.*, 2003, **69**, 4706-4713 (*isol, pmr, cmr*)

Photolumazine B P-382

1'-Deoxy-1-[1,3,4,7-tetrahydro-6-(hydroxymethyl)-2,4,7-trioxo-8(2H)-pteridinyl]-D-ribose, 9CI
[50868-45-6]

C₁₂H₁₆N₄O₈ 344.28

Prod. by *Photobacterium phosphoreum*. Inhibitor of riboflavine synthetase.

1'-Deoxy: **Photolumazine C** 7-Oxolumazine

[17879-89-9]

C₁₂H₁₆N₄O₇ 328.281

Prod. by *Photobacterium phosphoreum*, *Pseudomonas ovalis* and *Russula* sp. Inhibitor of riboflavine synthetase. Needles. Mp 260° (dec.).

Rowan, T. *et al.*, *J.C.S. (C)*, 1968, 452-458 (*synth, uv*)


Suzuki, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1971, **44**, 1869-1872 (*isol*)

Suzuki, A. *et al.*, *Biochim. Biophys. Acta*, 1973, **313**, 229-234 (*isol*)

Iten, P.X. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 550-569 (*isol, uv, cmr*)

Phthalimide, 8CI

1*H*-Isoindole-1,3(2*H*)-dione, 9CI
[85-41-6]



C₈H₅NO₂ 147.133

Constit. of egg masses of muricid mollusc *Ceratostoma erinaceum*. Source of “protected” ammonia in primary amine synth. (Gabriel). Needles (H₂O); prisms (AcOH); leaflets by subl. Insol. C₆H₆, petrol. Mp 238°. Forms salts with bases.

▶ LD₅₀ (mus, orl) 5000 mg/kg. Exp. reprod. and teratogenic effects. TI3920000

N-Benzyl: N-Benzylphthalimide

[2142-01-0]


C₁₅H₁₁NO₂ 237.257

Isol. from curare *Chondrodendron tomentosum*. Cryst. (EtOH). Mp 115–116°. Considered not to be an artifact.

Bick, I.R.C. *et al.*, *J.C.S.*, 1960, 2402-2407 (*N*-benzyl, isol)

Phyrrine

[148139-97-3]



C₁₆H₂₈N₂O₂ 280.409

Alkaloid from roots and aerial parts of *Physalis alkekengi* (winter cherry), the famine food *Physalis angulata* (cutleaf ground cherry), *Physalis philadelphica* (tomatillo), *Physalis peruviana* (Cape gooseberry), *Physalis minima*, *Physalis pubescens* (ground cherry), *Physalis viscosa* and *Physalis pruinosa* (strawberry tomato) (Solanaceae). Gum.

Dipicrate: [148218-52-4]


Mp 189° dec.

Basey, K. *et al.*, *Phytochemistry*, 1992, **31**, 4173 (isol, ir, pmr, cmr, ms, struct)

Phyllanthimide

P-385

3-(*Dimethylamino*)-1-(2-phenylethyl)-2,6-piperidinedione, 9CI




C₁₅H₂₀N₂O₂ 260.335

(±)-*form* [116174-65-3]

Alkaloid from the leaves and stems of *Phyllanthus sellowianus* (Euphorbiaceae). Tempesta, M.S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 617 (isol, uv, ir, pmr, cmr, ms, struct)

Phyllanthine†

4-Methoxysecurinan-11-one, 9CI. 4-Methoxysecurinine
[20072-02-0]



Absolute Configuration

C₁₄H₁₇NO₃ 247.293

Alkaloid from the roots of *Phyllanthus discoides* (Euphorbiaceae). Yellow cryst. (Me₂CO/diisopropyl ether). Mp 96-98°. [α]_D -898 (c, 0.98 in CHCl₃). pK_a 5.28.

O-De-Me: 4-Hydroxysecurinine

[130969-00-5]

C₁₃H₁₅NO₃ 233.266

Alkaloid from *Phyllanthus niruri* (Euphorbiaceae). Gum. C4-config. not determined.

12,13β,14,15-Tetrahydro: 4-Methoxytetrahydrosecurinine

[131063-47-3]

C₁₄H₂₁NO₃ 251.325

Alkaloid from *Phyllanthus niruri* (Euphorbiaceae). Orange-yellow powder (EtOH). Mp 59-60°. C4-config. not determined.

12,13β,14,15-Tetrahydro: Mp 66-68°.

[α]_D -8.5 (c, 0.94 in CHCl₃).

4-Epimer: 4-Epiphyllanthine

[138876-93-4]

C₁₄H₁₇NO₃ 247.293

Alkaloid from bark of *Margaritaria indica* (Euphorbiaceae). Yellow needles (EtOAc/hexane). Mp 82-84°. [α]_D²⁰ -753 (c, 0.06 in EtOH).

2,4-Diepimer: Securitinine

[13861-71-7]

C₁₄H₁₇NO₃ 247.293

Alkaloid from the root bark of *Securinea suffruticosa* (Euphorbiaceae). Yellow plates (EtOAc). Mp 129-130°. [α]_D -952.3 (c, 1 in EtOH).

2,4-Diepimer, 14,15-dihydro:

Needles (diisopropyl ether). Mp 118-119°.

2,4-Diepimer, 14,15-dihydro, 15α-methoxy: Secuamamine B

[1004529-76-3]

C₁₅H₂₁NO₄ 279.335

Alkaloid from the wood of *Securinea suffruticosa* var. *amamiensis*. Amorph. solid. [α]_D²⁵ +42.6 (c, 0.54 in CHCl₃). λ_{max} 230 (log ε 3.27); 262 (sh) (log ε 2.92) (MeOH).

Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1966, **14**, 917-918; 1967, **15**, 1633-1640 (*Phyllanthine*, *Securitinine*, isol, uv, ord, ir, pmr, ms, struct)

Parello, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 1117-1129 (isol, uv, ir, pmr, ms, struct)

Hassarajani, S.A. *et al.*, *Indian J. Chem., Sect. B*, 1990, **29**, 801 (*4-Hydroxysecurinine*, *4-Methoxytetrahydrosecurinine*)

Arbain, D. *et al.*, *J.C.S. Perkin I*, 1991, 1863-1869 (*4-Epiphyllanthine*)


Ohsaki, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2003-2005 (*Secuamamine B*)

P-386

Phyllantidine

P-387

10,11,12,12a-Tetrahydro-9*H*-6,12*b*-methanofuro[2,3-*d*]pyrido[1,2-*b*][1,2]oxazocin-2(6*H*)-one, 9CI



(+)-*form*

C₁₃H₁₅NO₃ 233.266

(+)-*form* [146074-64-8]

Alkaloid from *Breynia coronata* and *Securinea suffruticosa*. Pale yellow needles (Et₂O). Mp 166-168°. [α]_D +333 (CHCl₃). This enantiomer has been referred to as *ent*-Phyllantidine.

(-)-*form* [38836-07-6]

Alkaloid from the roots of *Phyllanthus discoides*. Mp 169-170°. [α]_D -450 (c, 0.33 in CHCl₃). λ_{max} 258 (log ε 4.2) (EtOH).

4α-Methoxy: Secuamamine D

[1004529-78-5]

C₁₄H₁₇NO₄ 263.293

Alkaloid from *Securinea suffruticosa* var. *amamiensis*. Amorph. solid. [α]_D²⁵ -303.9 (c, 0.26 in CHCl₃). λ_{max} 212 (sh) (log ε 3.33); 261 (log ε 3.62) (MeOH).

Parello, J. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1965, **260**, 337-340 (isol)

Horii, Z. *et al.*, *Tet. Lett.*, 1972, **13**, 1877-1880 (*synth*, uv, ir, pmr, ms)

Lajis, N.H. *et al.*, *Aust. J. Chem.*, 1992, **45**, 1893-1897 ((+)-*form*)


Carson, C.A. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 6560-6563 (*synth*)

Ohsaki, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2003-2005 (*Secuamamine D*)

Yuan, W. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 234-242 ((+)-*form*, isol, pmr, cmr)

Phyllocryptonine

P-388



(+)-*form*

C₂₀H₂₄NO₅[⊕] 358.413

(+)-*form* [126262-26-8]

Alkaloid from the leaves, twigs, roots, fruit and bark of *Cryptocarya phyllostemon* (Lauraceae). Fine cryst. (Me₂CO) (as chloride). Mp 230-231° (chloride). [α]_D +36 (c, 1.0 in EtOH).


(±)-*form*

Synthetic. Fine cryst. (Et₂O)(as chloride). Mp 189° (chloride).

Cavè, A. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2243 (isol, uv, pmr, cmr, ms, cd, synth, struct)

Phyllospadine P-389

4',5,7-Trihydroxy-6-methoxy-8-(1-methyl-2-pyrrolidyl)flavone
[76540-48-2]




C₂₁H₂₁NO₆ 383.4

Alkaloid from the sea-grass *Phyllospadix ivatensis*. Needles (as tri-Ac). Mp 199° (tri-Ac).

Takagi, M. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 3019 (*isol, uv, pmr, struct*)

Phyllosteminine P-390

[126262-23-5]




C₂₁H₂₁NO₄ 351.401

Alkaloid from the leaves, twigs, roots and fruit of *Cryptocarya phyllostemon* (Lauraceae). Cryst. (CHCl₃). Mp 207-209° dec. [α]_D²⁵ -49 (c, 0.4 in EtOH).

Cavé, A. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2243-2263 (*isol, uv, ir, pmr, ms, struct*)

Phyllostictine A P-391

[1007346-21-5]




C₁₇H₂₇NO₅ 325.404

Prod. by *Phyllosticta cirsii*. Phytotoxin. Oil. [α]_D²⁵ -87.5 (c, 0.2 in CHCl₃). λ_{max} 263 (log ε 4.07) (MeCN).

Evidente, A. *et al.*, *Tetrahedron*, 2008, **64**, 1612-1619 (*isol, pmr, cmr, ms*)

Phyllostictine B P-392

[1007346-22-6]




C₁₅H₂₃NO₅ 297.35

Prod. by *Phyllosticta cirsii*. Oil. [α]_D²⁵ -99.8 (c, 0.07 in CHCl₃). λ_{max} 262 (log ε 4.13) (MeCN).

Evidente, A. *et al.*, *Tetrahedron*, 2008, **64**, 1612-1619 (*isol, pmr, cmr, ms*)

Phyllostictine C P-393

[1007346-23-7]




C₁₇H₂₇NO₆ 341.403

Prod. by *Phyllosticta cirsii*. Oil. [α]_D²⁵ -45.5 (c, 0.1 in CHCl₃). λ_{max} 262 (log ε 3.6) (MeCN).

Evidente, A. *et al.*, *Tetrahedron*, 2008, **64**, 1612-1619 (*isol, pmr, cmr, ms*)

Phyllostictine D P-394


[1007346-24-8]



C₁₇H₂₅NO₆ 339.388

Prod. by *Phyllosticta cirsii*. Oil. [α]_D²⁵ -70.2 (c, 0.2 in CHCl₃). λ_{max} 262 (log ε 3.25) (MeCN).

Evidente, A. *et al.*, *Tetrahedron*, 2008, **64**, 1612-1619 (*isol, pmr, cmr, ms*)

Physarigin A P-395

C₂₅H₂₈N₂O₅ 436.507

Isol. from the myxomycete *Physarum rigidum*. Amorph. yellow solid. λ_{max} 403 (ε 26000); 426 (ε 22000) (MeOH).

2,3-Dihydro, 3-hydroxy: **Physarigin C**

C₂₅H₃₀N₂O₆ 454.522

Isol. from *Physarum rigidum*. Amorph. yellow solid. [α]_D²² +15 (c, 1.2 in MeOH). λ_{max} 351 (ε 9200); 368 (ε 12000); 390 (ε 10000) (MeOH).

Misono, Y. *et al.*, *Tet. Lett.*, 2003, **44**, 4479-4481 (*isol, pmr, cmr*)

Physarigin B P-396

As Physarigin A, P-395 with R = CH₃

C₂₆H₃₀N₂O₅ 450.533


Isol. from the myxomycete *Physarum rigidum*. Amorph. yellow solid. [α]_D²² -6.8

(c, 0.5 in MeOH). λ_{max} 403 (ε 23000); 426 (ε 20000) (MeOH).

Misono, Y. *et al.*, *Tet. Lett.*, 2003, **44**, 4479-4481 (*isol, pmr, cmr*)

Physarorubicin acid A P-397

[196621-49-5]



C₁₈H₁₉NO₆ 345.351


Tetramic acid deriv. Pigment from the slime mold *Physarum polycephalum*. Amorph. orange-red powder. Similar to Fuligorubin A, F-165. λ_{max} 252 (log ε 3.46); 404 (log ε 3.96) (MeOH).

Nowak, A. *et al.*, *Liebigs Ann./Recl.*, 1997, 1817-1821 (*isol, uv, ir, cd, pmr, cmr, ms*)

Dixon, D.J. *et al.*, *J.C.S. Perkin 1*, 1999, 2231-2232 (*synth*)

Physarorubicin acid B P-398

[220422-56-0]




C₁₆H₁₇NO₆ 319.313

Enolised β-tricarboxyl compd. Isol. from *Physarum polycephalum*. [α]_D²⁵ -146.8 (c, 0.02 in MeOH). λ_{max} 247 (log ε 3.67); 286 (log ε 3.67); 393 (log ε 4.37); 409 (sh) (MeOH).

Nowak, A. *et al.*, *Angew. Chem., Int. Ed.*, 1998, **37**, 3139-3141 (*isol, uv, ir, pmr, cmr*)

Physoperuvine P-399

8-Methyl-8-azabicyclo[3.2.1]octan-1-ol, 9Cl. 4-(Methylamino)cycloheptanone, 9Cl. 1-Hydroxytropane. 1-Tropanol
[60723-27-5]



C₈H₁₅NO 141.213

Free base tautomeric. Alkaloid from *Physalis peruviana* (Cape gooseberry) (Solanaceae). Mp 47-48° Mp 68-70°. May be a partial racemate.

► Mod. toxic.

Hydrochloride: Mp 153°. [α]_D²⁵ -0.8 (c, 1 in MeOH). Tropane bicyclic struct.

N-Benzoyl:

C₁₅H₁₉NO₂ 245.321

Mp 136-137°. Monocyclic struct.

[82748-86-5]


Ray, A.B. *et al.*, *Heterocycles*, 1982, **19**, 1233 (*cryst struct, cmr, pmr, cd, abs config*)

- McPhail, A.T. *et al.*, *Tetrahedron*, 1984, **40**, 1661 (*cryst struct, synth, resoln*)
 Boyer, F.-D. *et al.*, *Synlett*, 1992, 357 (*synth*)
 Justice, D.E. *et al.*, *J.C.S. Perkin 1*, 1994, 2559 (*synth*)
 Hiroya, K. *et al.*, *Chem. Comm.*, 1995, 2205 (*synth*)
 Majewski, M. *et al.*, *Synlett*, 1996, 785 (*synth*)
 Hodgson, D.M. *et al.*, *Tet. Lett.*, 1999, **40**, 8637-8640 (*synth*)
 Hall, A. *et al.*, *J.C.S. Perkin 1*, 2000, 329-343 (*synth*)

Physostigmine, BAN, USAN P-400

1,2,3,3a,8,8a-Hexahydro-1,3a,8-trimethylpyrrolo[2,3-b]indol-5-ol methylcarbamate (ester), 9CI. *Eserine*.

Fysostigmine. Physostol. Cogmine. Synapton
 [57-47-6]



$C_{15}H_{21}N_3O_2$ 275.35
 Main alkaloid from the Calabar bean (*Physostigma venenosum*) (Fabaceae). Also prod. by *Streptomyces* sp. AH-4 and *Streptomyces pseudogriseolus*. Anticholinesterase used as miotic and to decrease intraocular pressure in glaucoma. Cognition activator. Mp 86-87° Mp 105-106° (dimorph.). $[\alpha]_D$ -82 (CHCl₃). Log P 2.21 (calc). The claimed alkaloid Calabarine was prob. a mixt. of decomposition prods. of Physostigmine. λ_{max} 253 (ε 12100); 311 (ε 2800) (EtOH).

- ▶ Adverse effects reported when used therapeutically. Can cross blood-brain barrier. Exp. reprod. effects. LD₅₀ (mus, orl) 3 mg/kg.

Hydrobromide (1:2): Mp 224-226°.

Sulfate (2:1): **Physostigmine sulfate, USAN**
 [64-47-1]
 Scales. Mp 140°.

- ▶ UY8585000

Salicylate (1:1): **Physostigmine salicylate, USAN. Antilirium. Eserocil. Esromistin. Fisostin. Zem-Eserine. Many other names**
 [57-64-7]
 Mp 184-186°.

- ▶ Adverse systemic effects when used therapeutically. LD₅₀ (mus, orl) 2.5 mg/kg. Exp. reprod. effects. TJ2450000

Methiodide: Mp 188°.

N¹-De-Me, N¹-Ac: Antibiotic TAN 1169A. TAN 1169A
 [132663-94-6]

$C_{16}H_{21}N_3O_3$ 303.36
 Prod. by *Streptomyces* sp. (AI2052; IFO14833; BP2328). Acetylcholinesterase inhibitor. Shows antiamnesic prods. Cryst. $[\alpha]_D$ -376 (MeOH). Abs. config. not certain. λ_{max} 254 (E1%/1cm 473); 310 (E1%/1cm 105) (MeOH) (Berdy).

N¹-De-Me, N¹-(methylaminocarbonyl):

Eseramine

[6091-57-2]

$C_{16}H_{22}N_4O_3$ 318.375

Alkaloid from *Physostigma venenosum* (Fabaceae). Acetylcholinesterase inhibitor. Needles (EtOH). Mp 240-242° dec. (216-218°). $[\alpha]_D^{23}$ -289 (EtOH). λ_{max} 256 (ε 13050); 311 (ε 2680) (EtOH).

N⁸-De-Me: N⁸-Norphysostigmine

[19573-10-5]

$C_{14}H_{19}N_3O_2$ 261.323

Isol. from *Streptomyces* sp. AH-4 and *Physostigma venenosum*. Shows insecticidal prods. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 151.5°. $[\alpha]_D^{30}$ -64.4 (c, 0.45 in MeOH). Log P 1.35 (calc). λ_{max} 245 (ε 4900); 303 (ε 1300) (MeOH) (Berdy).

N¹,N⁸-Di-de-Me, N¹-Ac: Antibiotic TAN 1169B. TAN 1169B

[132663-95-7]

$C_{15}H_{19}N_3O_3$ 289.333

Prod. by *Streptomyces* sp. (AI2052; IFO14833; BP2328). Acetylcholinesterase inhibitor. Shows antiamnesic prods. Cryst. $[\alpha]_D$ -382 (MeOH). Abs. config. not certain. λ_{max} 243 (E1%/1cm 345); 300 (E1%/1cm 98) (MeOH) (Berdy).

[8006-41-5 (Calabarine)]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 1065C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 612B (*nmr*)

Harley-Mason, J. *et al.*, *J.C.S.*, 1954, 3651-3654 (*synth*)

Robinson, B. *et al.*, *Chem. Ind. (London)*, 1964, 459-460 (*Eseramine, uv, ir, pmr, ms struct*)

Robinson, B. *et al.*, *J.C.S.*, 1965, 3336-3339 (*Eseramine, synth*)

Bild, N. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 1885-1892 (*ms*)

Robinson, R. *et al.*, *Alkaloids (Academic Press)*, 1968, **10**, 383-400 (*rev*)

Newkome, G.R. *et al.*, *Chem. Comm.*, 1969, 385 (*pmr, config*)

Longmore, R.B. *et al.*, *Chem. Ind. (London)*, 1969, 622-623 (*Eseramine, abs config*)

Hill, R.K. *et al.*, *Tetrahedron*, 1969, **25**, 1249-1260 (*config*)

Pauling, P. *et al.*, *J.C.S. Perkin 2*, 1973, 1342-1345 (*cryst struct*)

U.S. Pat., 1973, 3 734 832; CA, **79**, 30496q (*manuf*)

Crooks, P.A. *et al.*, *Phytochemistry*, 1974, **15**, 1092-1093 (*Eseramine, cmr*)

Granacher, R.P. *et al.*, *Clin. Neuropharmacol.*, 1976, **1**, 63-79 (*rev, pharmacol*)

Slenberg, V.I. *et al.*, *J. Het. Chem.*, 1977, **14**, 407-410 (*cmr, N-15 nmr*)

Bose, A.K. *et al.*, *Anal. Biochem.*, 1978, **89**, 284-291 (*ms*)

Wijnberg, J.B.P.A. *et al.*, *Tetrahedron*, 1978, **34**, 2399-2404 (*synth*)

Iwasa, T. *et al.*, *CA*, 1981, **95**, 148651 (*isol*)

Stoeckel, H. *et al.*, *CA*, 1982, **97**, 138642 (*book*)

Nieminen, A.O.K. *et al.*, *J. Het. Chem.*, 1983, **20**, 515-517 (*cmr*)

Murao, S. *et al.*, *Agric. Biol. Chem.*, 1986, **50**, 523-524 (*isol, N⁸-Norphysostigmine, activity*)

Yu, Q.-S. *et al.*, *Heterocycles*, 1988, **27**, 745-750; 1709-1712 (*synth*)

Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1989, **18**, 289-350 (*rev*)

Japan. Pat., 1990, 90 270 875; CA, **114**, 120278r (*TAN 1169A, TAN 1169B*)

Horne, S. *et al.*, *J.C.S. Perkin 1*, 1991, 3047-3051 (*synth*)

Lee, T.B.K. *et al.*, *J.O.C.*, 1991, **56**, 872-875 (*synth*)

Takano, S. *et al.*, *J.O.C.*, 1991, **56**, 5982-5984 (*synth*)

Marino, J.P. *et al.*, *J.A.C.S.*, 1992, **114**, 5566-5572 (*synth*)

Grieco, P.A. *et al.*, *Tet. Lett.*, 1992, **33**, 4401-4404 (*synth*)

Ashimori, A. *et al.*, *J.O.C.*, 1993, **58**, 6949-6951 (*synth*)

Yu, Q.-S. *et al.*, *Heterocycles*, 1994, **39**, 519-525 (*synth*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms, 7th edn.*, Akademie-Verlag, 1994, 4767 (*synonyms*)

Brossi, A. *et al.*, *Aust. J. Chem.*, 1996, **49**, 171-181 (*synth, rev*)

Node, M. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 715-719 (*synth*)

Morales-Rós, M.S. *et al.*, *Tetrahedron*, 1996, **52**, 5339-5348 (*synth*)

Triggler, D.J. *et al.*, *CNS Drug Rev.*, 1998, **4**, 87-136

Witkop, B. *et al.*, *Heterocycles*, 1998, **49**, 9-27 (*rev*)

Matsuura, T. *et al.*, *J.A.C.S.*, 1998, **120**, 6500-6503 (*synth*)

Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 1395; 1396

Kawahara, M. *et al.*, *Org. Lett.*, 2000, **2**, 675-678; 953-955 (*synth*)

ElAzab, A.S. *et al.*, *Org. Lett.*, 2000, **2**, 2757-2759 (*synth*)

Ishibashi, H. *et al.*, *Tetrahedron*, 2000, **56**, 1469-1473 (*synth*)

Santos-Sanchez, N.F. *et al.*, *Magn. Reson. Chem.*, 2001, **39**, 696-700 (*pmr, conformn*)

Tanaka, K. *et al.*, *Tet. Lett.*, 2001, **42**, 1049-1052 (*synth*)

Robinson, B. *et al.*, *Heterocycles*, 2002, **57**, 1327-1352 (*rev, synth*)

Morales-Rós, M.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 136-141 (*synth*)

Rege, P.D. *et al.*, *J.O.C.*, 2003, **68**, 6133-6139 (*synth*)

Santos, P.F. *et al.*, *Tetrahedron*, 2005, **61**, 9147-9156 (*synth*)

Trost, B.M. *et al.*, *J.A.C.S.*, 2006, **128**, 4590-4591 (*synth*)


Mukai, C. *et al.*, *Org. Lett.*, 2006, **8**, 83-86 (*synth*)

Asakawa, K. *et al.*, *Heterocycles*, 2008, **76**, 183-190 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, PIA500; PIB000; PIA750

Physovenine**P-401**

3,3a,8,8a-Tetrahydro-3a,8-dimethyl-2H-furo[2,3-b]indol-5-ol methyl carbamate (ester), 9CI



$C_{14}H_{18}N_2O_3$ 262.308

Log P 1.71 (calc).

(-)-form [6091-05-0]

Alkaloid from *Physostigma venenosum* (Fabaceae). Powerful miotic agent, ACH potentiator. Mp 123°. $[\alpha]_D^{22}$ -92 (EtOH). Pharmacol. active isomer. λ_{\max} 252 (ϵ 13200); 310 (ϵ 3300) (EtOH).

(+)-form [29347-11-3]

Synthetic. Also obt. by resoln. of the racemate. Yellow cryst. (EtOAc). Mp 127-128° (122-124°). $[\alpha]_D$ +93.5 (c, 0.5 in EtOH).

(±)-form [2520-34-5]


Synthetic. Prisms (Et₂O/petrol). Mp 142-143° (with subl. from 139°).

- Robinson, B. *et al.*, *J.C.S.*, 1964, 1503-1506 (*isol, uv, ir, pmr, struct*)
 Longmore, R.B. *et al.*, *Chem. Ind. (London)*, 1966, 1638-1639; 1969, 622-623 (*synth, pmr, abs config*)
 Crooks, P.A. *et al.*, *Phytochemistry*, 1976, **15**, 1092-1093 (*cmr*)
 Luo, Y. *et al.*, *Heterocycles*, 1990, **31**, 283-287 (*synth*)
 Shishido, K. *et al.*, *Tet. Lett.*, 1990, **31**, 219-220 (*synth*)
 Yu, Q.-S. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 761-766 (*synth, resoln*)
 Horne, S. *et al.*, *J.C.S. Perkin 1*, 1991, 3047-3051 (*synth*)
 Takano, S. *et al.*, *J.O.C.*, 1991, **56**, 5982-5984 (*synth*)
 Clark, A.J. *et al.*, *Tetrahedron*, 1992, **48**, 6875-6882 (*synth*)
 Yu, Q.-S. *et al.*, *Heterocycles*, 1994, **39**, 519-525 (*synth*)
 Node, M. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 715-719 (*synth*)
 Matsuura, T. *et al.*, *J.A.C.S.*, 1998, **120**, 6500-6503 (*synth*)
 Ishibashi, H. *et al.*, *Tetrahedron*, 2000, **56**, 1469-1473 (*synth*)
 Tanaka, K. *et al.*, *Tet. Lett.*, 2001, **42**, 1049-1052 (*synth*)
 Morales-Rios, M.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 136-141 (*synth*)
 Zhang, T.Y. *et al.*, *Tet. Lett.*, 2002, **43**, 1363-1365 (*synth*)
 Sunazuka, T. *et al.*, *Tet. Lett.*, 2005, **46**, 1459-1461 (*synth*)

Piccolamine

P-402

[139742-31-7]

C₁₈H₂₇NO 273.417

Isol. from the Senegalese gorgonian *Leptogorgia piccola*. $[\alpha]_D$ -22 (c, 0.022 in CHCl₃). λ_{\max} 204 (ϵ 10500); 208 (ϵ 8600); 228 (ϵ 3250) (MeOH).

N-Oxide: Piccolamine N-oxide

[139742-32-8]

C₁₈H₂₇NO₂ 289.417


Isol. from the gorgonian *Leptogorgia piccola*. $[\alpha]_D$ -47 (c, 0.012 in CHCl₃). λ_{\max} 204 (ϵ 11200); 207 (ϵ 3100) (MeOH).

Roussis, V. *et al.*, *New J. Chem.*, 1991, **15**, 959-961 (*isol, pmr, cmr, struct*)

Picranitine

P-403

[534592-56-8]



Absolute Configuration


C₂₁H₂₄N₂O₅ 384.431Constit. of the seeds of *Picralima nitida*.

Tane, P. *et al.*, *Bull. Chem. Soc. Ethiop.*, 2002, **16**, 165-168 (*isol*)

Picraphylline

P-404

[2447-72-5]



Absolute Configuration

C₂₂H₂₆N₂O₄ 382.458

Minor alkaloid from leaves of *Picralima nitida* (Apocynaceae). Mp 255°. $[\alpha]_D^{20}$ -37 (CHCl₃). λ_{\max} 238 (log ϵ 4.14); 313 (log ϵ 3.98) (no solvent reported).

11-Methoxy: 11-Methoxypicraphylline

[77809-92-8]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{\max} 223; 267 (sh); 281; 291; 336 (EtOH). λ_{\max} 225; 270 (sh); 282; 289; 299 (sh) (EtOH/HCl)

10,11-Dimethoxy: 10,11-Dimethoxypicraphylline

[36151-14-1]

C₂₄H₃₀N₂O₆ 442.511

Alkaloid from *Aspidosperma marcgravianum* and *Ochrosia balansae* (Apocynaceae). Cryst. (Et₂O). Mp 183-185°. $[\alpha]_D$ -50 (c, 1 in CHCl₃). λ_{\max} 214 (log ϵ 4.46); 342 (log ϵ 4.2) (EtOH).

Lévy, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1964, 1917-1924 (*uv, ir, ms, pmr, struct*)

Le Men, J. *et al.*, *J. Nat. Prod.*, 1964, **27**, 456-469 (*isol*)

Bruneton, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, **273**, 442-444 (*Dimethoxypicraphylline*)


Ahond, A. *et al.*, *J. Nat. Prod.*, 1981, **44**, 193-199 (*11-Methoxypicraphylline*)

Verpoorte, R. *et al.*, *Planta Med.*, 1982, **46**, 149-152 (*Dimethoxypicraphylline*)

Picrasidine X

P-405

[155416-26-5]

C₁₄H₁₂N₂O₃ 256.26

Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Pale yellow solid (MeOH). Mp 180-181°. $[\alpha]_D^{22}$ -37 (c, 0.4 in Py). Config. at 2' posn. not determined. λ_{\max} 250 (log ϵ 4.58); 290 (log ϵ 4.23); 303 (sh) (log ϵ 4.27); 352 (log ϵ 3.79); 370 (sh) (log ϵ 3.71) (MeOH).

1'-Hydroxy: Picrasidine Y

[155416-27-6]

C₁₄H₁₂N₂O₄ 272.26


From wood of *Picrasma quassioides* (Simaroubaceae). Pale yellow solid (MeOH). Mp 184-185°. $[\alpha]_D^{22}$ -33.2 (c, 0.5 in Py). Stereochem. at 1' and 2' posn. not determined. λ_{\max} 250 (log ϵ 4.51); 290 (log ϵ 4.21); 302 (log ϵ 4.27); 352 (log ϵ 3.6); 370 (sh) (log ϵ 3.51) (MeOH).

Li, H.-Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 1807-1811 (*isol, uv, ir, pmr, cmr, ms, struct*)

Picrasidine A

P-406

3-(4,8-Dimethoxy-9H-pyrido[3,4-b]indol-1-yl)-1-(9H-pyrido[3,4-b]indol-1-yl)-1-propanone, 9CI. β -Carbolin-1-yl 4,8-dimethoxy- β -carbolin-1-ylethyl ketone [82652-20-8]

C₂₇H₂₂N₄O₃ 450.496


Alkaloid from the roots of *Picrasma quassioides* (Simaroubaceae). Cryst. (CHCl₃/MeOH). Mp 263-264° dec.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1204; 1984, **32**, 3579 (*pmr, cmr, ms, struct, nomencl*)

Picrasidine B

P-407

4,8-Dimethoxy-1-(2-methoxyethyl)-9H-pyrido[3,4-b]indole, 9CI. 4,8-Dimethoxy-1-(2-methoxyethyl)- β -carboline [82652-19-5]


C₁₆H₁₈N₂O₃ 286.33

Alkaloid from the roots of *Picrasma quassioides* (Simaroubaceae). Prisms (Me₂CO). Mp 201-202°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1204 (*isol, uv, ir, pmr, cmr, ms, struct*)

Picrasidine C P-408

β -Carboline-1-yl 3-(4,8-dimethoxy- β -carboline-1-yl)-1-methoxypropyl ketone [88142-61-4]




$C_{29}H_{26}N_4O_4$ 494.549
Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Cryst. (MeOH). Mp 121-122°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 3198; 1984, **32**, 3579 (*isol, uv, ir, pmr, cmr, ms, struct, nomencl*)


Picrasidine E P-409

[94530-77-5]



$C_{18}H_{16}N_2O_5$ 340.335
Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Yellow needles (Me₂CO). Mp 220°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 3579 (*isol, uv, ir, pmr, cmr, ms, struct*)

Picrasidine F P-410

$C_{29}H_{27}N_4O_3^{\oplus}$ 479.557
Plates + $\frac{1}{2}$ MeOH (MeOH) (as chloride). Mp 265-267° dec. (chloride).


(\pm)-form [109225-38-9]

Quaternary alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae).

Koike, K. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 3228 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Picrasidine H

[105608-30-8]



$C_{28}H_{24}N_4O_4$ 480.522

Revised struct; probable struct (1997). Alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae). Needles (MeOH). Mp 218-220° dec. $[\alpha]_D^{25}$ -63.1 (c, 1.3 in DMSO).

O-Ac:

Needles (MeOH). Mp 158-159°.


Koike, K. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 2090 (*isol, uv, ir, pmr, cmr, ms*)

Suzuki, H. *et al.*, *Heterocycles*, 1997, **46**, 57-60 (*struct*)

Picrasidine K

P-412

1-[2-(Dimethylamino)ethyl]-4-methoxy-9H-pyrido[3,4-b]indol-8-ol, 9Cl. 1-(2-Diethylaminoethyl)-8-hydroxy-4-methoxy- β -carboline [100234-63-7]



$C_{18}H_{23}N_3O_2$ 313.399

Alkaloid from the bark of *Picrasma quassioides* (Simaroubaceae). Plates (MeOH). Mp 169-171°.

O-Ac; hydrochloride:

Pale-yellow needles. Mp 189-190°.

Me ether; hydrochloride:


Pale-yellow needles. Mp 223-225°.

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3356 (*isol, uv, ir, pmr, cmr, ms, struct*)

Picrasidine L

P-413

3-Methyl-3H-indolo[3,2,1-de][1,5]-naphthyridine-5,6-dione, 9Cl. 3-Methyl-canthin-5,6-dione [96405-70-8]



$C_{15}H_{10}N_2O_2$ 250.256

Struct. revised in 1985. Alkaloid from the wood and root bark of *Picrasma quassioides* (Simaroubaceae). Orange-red needles. Mp 330°. λ_{max} 242 (log ϵ 4.33); 250 (log ϵ 4.3); 287 (log ϵ 3.87); 345 (log ϵ 3.55); 444 (log ϵ 4.1); 462 (log ϵ 4.1) (EtOH).

4-Methoxy: 4-Methoxy-3-methyl-3H-indolo[3,2,1-de][1,5]naphthyridine-5,6-dione. 4-Methoxy-3-methyl-5,6-canthin-5,6-dione. **Picrasidine O** [101219-63-0]

$C_{16}H_{12}N_2O_3$ 280.282

Alkaloid from the root wood of *Picrasma quassioides* (Simaroubaceae). Red needles. Mp 274° dec. λ_{max} 242 (log ϵ 4.28); 250 (sh) (log ϵ 4.27); 290 (log ϵ 4.2); 360 (log ϵ 3.54); 480 (log ϵ 4.04); 500 (log ϵ 4) (EtOH).

9-Methoxy: 9-Methoxy-3-methyl-5,6-canthin-5,6-dione

[119935-09-0]

$C_{16}H_{12}N_2O_3$ 280.282

Alkaloid from the wood of *Eurycoma longifolia* (Simaroubaceae). Yellow needles. Mp 300°.


Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1204-1209; 1985, **33**, 3847-3851; 4901 (*Picrasidines, isol, uv, ir, pmr, cmr, ms, struct, synth*)

Mitsunaga, K. *et al.*, *Phytochemistry*, 1994, **35**, 799 (9-methoxy)

Picrasidine M

P-414

[99964-79-1]



$C_{29}H_{22}N_4O_4$ 490.517

Alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae). Orange needles (DMSO). Mp 294-295° dec. λ_{max} 240 (log ϵ 4.3); 284 (log ϵ 3.65); 338 (log ϵ 3.42); 444 (log ϵ 3.52); 464 (log ϵ 3.52) (EtOH).

4-Methoxy: **Picrasidine U**

[118636-90-1]

$C_{30}H_{24}N_4O_5$ 520.543

Alkaloid from root wood of *Picrasma quassioides*. Red needles (MeOH). Mp 199-200° dec. λ_{max} 221 (log ϵ 4.73); 286 (log ϵ 4.19); 344 (log ϵ 3.93); 472 (log ϵ 3.87); 500 (log ϵ 3.83) (EtOH).

8'-Demethoxy, N^{9'}-methoxy: **Picrasidine N**

[101219-62-9]

$C_{29}H_{22}N_4O_4$ 490.517

Alkaloid from the root wood of *Picrasma quassioides* (Simaroubaceae). Orange needles (CHCl₃/MeOH). Mp 171-172° dec. λ_{max} 232 (log ϵ 4.25); 241 (log ϵ 4.25); 284 (log ϵ 3.87); 342 (log ϵ 3.87)


3.52); 442 (log ϵ 3.59); 466 (log ϵ 3.59) (EtOH).

Ohmoto, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3847-3851; 4901-4905 (*Picrasidines M,N*)
Koike, K. *et al.*, *Phytochemistry*, 1988, **27**, 3029-3030 (*Picrasidine U*)

Picrasidine R

P-415

[106154-68-1]



$C_{30}H_{26}N_4O_6$ 538.559


Alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae). Needles (MeOH). Mp 297-300° dec.

Koike, K. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 2090 (*isol, uv, ir, pmr, cmr, ms, struct*)

Picrasidine T

P-416

[113808-03-0]



$C_{28}H_{25}N_4O_4^{\oplus}$ 481.53

(±)-form

Alkaloid from the bark of *Picrasma quassioides* (Simaroubaceae). Pale yellow needles (MeOH) (as chloride). M.p. >300° (chloride). CAS no. refers to chloride.

Di-Me ether: Picrasidine S

[112503-87-4]

$C_{30}H_{29}N_4O_4^{\oplus}$ 509.583

Alkaloid from the root bark of *Picrasma quassioides* (Simaroubaceae). Mp 215-217° dec. (chloride). Pale yellow needles (MeOH) (as chloride) (as chloride).

Dideoxy: Picrasidine G

[112503-90-9]

$C_{28}H_{25}N_4O_2^{\oplus}$ 449.531

Alkaloid from the root bark of *Picrasma quassioides* and from *Picrasma javanica* (Simaroubaceae). Shows anti-ulcer props. Pale yellow needles (MeOH) (as chloride). Mp 272-273° dec. (chloride). λ_{max} 248 (ϵ 22387); 316 (ϵ 5754); 342 (ϵ 3715); 372 (ϵ 1996) (EtOH) (Berdy). λ_{max} 240 (ϵ 19054); 279 (ϵ 28153); 328 (ϵ 4786); 400 (ϵ 1737) (EtOH-NaOH) (Berdy).

Koike, K. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 3305 (*isol, uv, ir, pmr, cmr, ms, struct*)


Koike, K. *et al.*, *Phytochemistry*, 1987, **26**, 3375 (*isol, uv, ir, pmr, ms, struct*)

Liu, J. *et al.*, *Magn. Reson. Chem.*, 1993, **31**, 1091 (*pmr, cmr, Picrasidine G*)

Picrinine

P-417

Methyl 2,5-epoxy-1,2-dihydroakuammilan-17-oate, 9CI. Deacetyldeformylpicrasiline. Vincaridine
[4684-32-6]



$C_{20}H_{22}N_2O_3$ 338.405

Alkaloid from *Amsonia brevifolia*, *Alstonia scholaris*, *Alstonia macrophylla*, *Vinca minor*, *Vinca erecta*, *Vinca libanotica*, *Rauwolfia vomitoria*, *Rauwolfia cumminsi* and *Rauwolfia oreogiton*. Needles (Me₂CO). Mp 223-225° dec. $[\alpha]_D^{25}$ -44 (CHCl₃). pK_a 5.72 (EtOH aq.). λ_{max} 237 (ϵ 7950); 287 (ϵ 3230) (EtOH).

Picrate: Mp 172-174° dec.

Methiodide: Mp 235-237° dec.

N¹-Me: Ervincine

[18223-72-8]

$C_{21}H_{24}N_2O_3$ 352.432

Alkaloid from *Vinca erecta* (Apocynaceae). Mp 156-157°. $[\alpha]_D^{25}$ +93 (c, 0.2 in CHCl₃). λ_{max} 246 (log ϵ 3.39); 295 (log ϵ 3.5) (EtOH).

10-Methoxy: Vincaricine. 12-Demethoxytabernulosine

[22223-13-8]

[82260-04-6]

$C_{21}H_{24}N_2O_4$ 368.432

Alkaloid from *Vinca erecta* and *Tabernaemontana glandulosa*. Cryst. (C₆H₆) (Vincaricine) or oil (12-Demethoxytabernulosine). Mp 187-189° (Vincaricine). $[\alpha]_D^{22}$ 0 (c, 0.6 in CHCl₃) (Vincaricine). $[\alpha]_D^{20}$ -21 (c, 0.2 in CHCl₃) (12-Demethoxytabernulosine). Struct. revised in 1985. Same struct. assigned to Vincaricine and 12-Demethoxytabernulosine although the reported props. are different. λ_{max} 236 (log ϵ 3.85); 308 (log ϵ 3.48) (EtOH). λ_{max} 237 (log ϵ 3.65); 306 (log ϵ 3.3) (MeOH).

ar-Methoxy, N¹-Me: Quaternidine

[57515-71-6]

$C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from *Alstonia quaternata* (Apocynaceae). Oil. $[\alpha]_D^{20}$ -85 (c, 0.4 in CHCl₃). Exact posn. of OMe subn. not detd. λ_{max} 241 (log ϵ 4.2); 308 (log ϵ 3.79) (EtOH).

10,11-Dimethoxy: Volkensine†. N^a-Demethylquaternine. Norquaternine

[74991-69-8]

$C_{22}H_{26}N_2O_5$ 398.458

Alkaloid from the leaves of *Rauwolfia oreogiton* and *Rauwolfia volkensii* and the trunk bark of *Alstonia legouixiae* (Apocynaceae). Needles. Mp 280°. $[\alpha]_D^{20}$ -63.4 (c, 0.1 in CHCl₃). λ_{max} 213 (log ϵ 4.3); 244 (log ϵ 4); 303 (log ϵ 3.9) (MeOH).

10,11-Dimethoxy, N¹-Me: Quaternine.*Alstipicalramine*

[57499-02-2]

$C_{23}H_{28}N_2O_5$ 412.485

Alkaloid from *Alstonia quaternata*, the leaves of *Rauwolfia oreogiton*, *Rauwolfia volkensii*, the trunk bark of *Alstonia legouixiae* and from leaves of *Alstonia macrophylla* (Apocynaceae). Cryst. (MeOH). Mp 153°. $[\alpha]_D^{20}$ -27 (c, 0.4 in CHCl₃). $[\alpha]_D^{20}$ +3.33. Samples not compared. The +ve opt. rotn. refers to Alstipicalramine. λ_{max} 249 (log ϵ 3.92); 304 (log ϵ 3.78) (EtOH). λ_{max} 232 (log ϵ 3.9); 272 (log ϵ 3.3) (MeOH).

10,12-Dimethoxy: Tabernulosine

[74627-72-8]

$C_{22}H_{26}N_2O_5$ 398.458

Alkaloid from the stems and leaves of *Tabernaemontana glandulosa* (Apocynaceae). Platelets (MeOH). Mp 190-191° dec. $[\alpha]_D^{20}$ -27 (c, 1.0 in CHCl₃). λ_{max} 240 (log ϵ 3.9); 301 (log ϵ 3.55) (MeOH).

16-Epimer: Picralstonine

[38801-86-4]

$C_{20}H_{22}N_2O_3$ 338.405

Alkaloid from the leaves of *Alstonia macrophylla* (Apocynaceae). Granules (Me₂CO). Mp 200°. $[\alpha]_D^{25}$ -90 (CHCl₃). λ_{max} 234 (log ϵ 3.81); 286 (log ϵ 3.4) (EtOH).

[27501-23-1]

Britten, A.Z. *et al.*, *J.C.S.*, 1963, 3850-3854 (*Picrinine, synth, uv, ir*)

Chatterjee, A. *et al.*, *Tet. Lett.*, 1965, 3633-3637 (*Picrinine, uv, pmr, ms*)

Rakhimov, D.A. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 354-355; 1969, **5**, 521-524; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 300-301; 1969, **5**, 440-443 (*Ervincine, Vincaridine*)

Il'yasova, Kh.T. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 327-328; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 278 (*Vincaricine*)

Rakhimov, D.A. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 521-524; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 440-443 (*Vincaricine*)

Banerji, A. *et al.*, *Phytochemistry*, 1972, **11**, 2605-2607 (*Picralstonine*)

Grossmann, E. *et al.*, *Phytochemistry*, 1973, **12**, 2058 (*uv, ms, pmr*)

Mamatas-Kalamaras, S. *et al.*, *Phytochemistry*, 1975, **14**, 1849-1854 (*Quaternidine, Quaternine*)

Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 307-311 (*Volkensine, Quaternine*)

Lewin, G. *et al.*, *Ann. Pharm. Fr.*, 1981, **39**, 273-275 (*Volkensine*)

Achenbach, H. *et al.*, *Annalen*, 1982, 830-844 (*Tabernulosine, 12-Demethoxytabernulosine*)

Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 129-130; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 131-132 (*Vincaricine*)


Cherif, A. *et al.*, *Heterocycles*, 1987, **26**, 3055-3058 (*Norquaternine, cmr*)

Ghosh, R. *et al.*, *Acta Cryst. C*, 1988, **44**, 2151-2154 (*cryst struct*)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1988, **27**, 961-965 (*Alstipicalramine*)

Pictamine**P-418**

Octahydro-4-methyl-6-(1,3-octadienyl)-2H-quinolizin-3-ol acetate, 9CI
[136945-62-5]



Absolute Configuration


C₂₀H₃₃NO₂ 319.486

Alkaloid from the tunicate *Clavelina picta*. Oil. $[\alpha]_D^{20}$ -87 (c, 0.1 in EtOAc). λ_{\max} 230 (ε 19000) (EtOH) (Derep). λ_{\max} 231 (MeOH) (Berdy).

Kong, F. *et al.*, *Tet. Lett.*, 1991, **32**, 3667-3668 (isol, uv, ir, pmr, cmr, ms, struct)
Toyooka, N. *et al.*, *Tetrahedron*, 1999, **55**, 15209-15224 (synth, abs config)
Yu, S. *et al.*, *Org. Lett.*, 2006, **8**, 3179-3182 (synth)

Pictonamine**P-419**

[95456-40-9]

C₂₁H₁₉NO₆ 381.384

Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Yellow cryst. (MeOH). Mp 231-232° (228-230°). Methyl enol ether of 13-Deoxychilenine in C-375.

Moniot, J.L. *et al.*, *J.O.C.*, 1979, **44**, 4347 (synth, uv, pmr)
Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957 (isol, uv, ir, pmr, ms)


Pictumine†**P-420**

[39292-54-1]

C₁₄H₂₃NO₄ 269.34

Pyrrolizidine alkaloid. Struct. unknown. Alkaloid from *Cynoglossum pictum* (Boraginaceae). Cryst. (Me₂CO). Mp 158-159°. Contains α,β-unsatd. ester and tertiary OH groups.

Man'ko, I.V. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 655-656; *Chem. Nat. Compd. (Engl. Transl.)*, 630

Piericidin**P-421**

Relative Configuration

Piericidin A₁, R¹ = H, R² = CH₃
A₂, R¹ = R² = CH₃
A₃, R¹ = H, R² = CH(CH₃)₂
A₄, R¹ = CH₃, R² = CH(CH₃)₂
A₅, R¹ = H, R² = CH₂CH₃
A₆, R¹ = CH₃, R² = CH₂CH₃
Antibiotic IT 143A, R¹ = CH₃, R² = -C(CH₃)=CHCH₂(E)-
IT 143B, R¹ = H, R² = -C(CH₃)=CHCH₂(E)-

Pyridine antibiotic family. Sol. MeOH,

hexane; poorly sol. H₂O. λ_{\max} 205 (ε 40800); 237 (ε 41800); 267 (ε 6600) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 0.87 mg/kg.

Piericidin A₁

Shaoguanmycin B. MT 1882-I. Antibiotic MT 1882-I. SN 198E. Antibiotic SN 198E. IT 143D. Antibiotic IT 143D [2738-64-9]

C₂₅H₃₇NO₄ 415.572

Isol. from *Streptomyces pactum* and other spp. Inhibits NADH oxidase of ox-heart mitochondria in the absence of bovine serum albumen. Also shows insecticidal activity. Viscous oil. Sol. MeOH, hexane; poorly sol. H₂O. $[\alpha]_D^{20}$ -0.9. λ_{\max} 232 (ε 39500); 239 (ε 40500); 268 (ε 5300) (MeOH) (Derep).

▶ YD4588000

4-O-D-Glucopyranoside: Glucopiericidin B [108073-61-6]

C₃₁H₄₇NO₉ 577.714

From *Streptomyces pactum*. More biol. active than Piericidin A₁. Amorph. Sol. MeOH, Et₂O, CHCl₃, Py, DMSO; poorly sol. H₂O. $[\alpha]_D^{20}$ -10 (c, 0.9 in MeOH). λ_{\max} 203 (ε 39700); 232 (ε 40050); 237 (ε 40200); 275 (ε 8300) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 10-30 mg/kg. LZ5902300

10'-O-β-D-Glucopyranoside: Glucopiericidin A. MT 1882-II. Antibiotic MT 1882-II [108073-65-0]

C₃₁H₄₇NO₉ 577.714

From *Streptomyces pactum* and *Streptomyces* sp. OM-5689. Hypotensive agent, antibody formation Inhibitor, bleb formation Inhibitor. More biol. active than Piericidin A₁. Amorph. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{20}$ -26.4 (c, 1 in CHCl₃). λ_{\max} 203 (ε 45400); 233 (ε 40000); 237 (ε 40800); 267 (ε 6100) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 10-30 mg/kg. LZ5901300

4-O-L-Rhamnopyranoside: 4-Rhamnopericidin A₁. SN 198C. Antibiotic SN 198C [131622-62-3]

C₃₁H₄₇NO₈ 561.714

From *Streptomyces* sp. SN-198. Active against gram-negative bacteria and fungi. Shows antitumour props. Amorph. Sol. MeOH, DMSO, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{25}$ -44 (c, 0.1 in MeOH). Grad. changes to oil > 82°. Named as 3'-rhamnosyl deriv. in paper. λ_{\max} 231 (ε 39300); 237 (ε 37700); 276 (ε 7200) (MeOH) (Berdy). λ_{\max} 232 (ε 37000); 238 (ε 37800); 277 (ε 6700) (EtOH) (Berdy).

4-O-(6-Deoxy-β-D-talopyranoside): 3'-Deoxytalopericidin A₁. DTPA [134876-72-5]

C₃₁H₄₇NO₈ 561.714

Isol. from an unconfirmed *Streptomyces* sp. Antitumour agent. Pale yellow oil. Poorly sol. H₂O. Lit. uses different numbering system. λ_{\max} 235; 275 (MeOH) (Berdy).

11',12'-Epoxide: Piericidin C₁ [58947-79-8]

C₂₅H₃₇NO₅ 431.571

Isol. from *Streptomyces pactum*. NADH oxidase inhibitor. Sol. MeOH, Et₂O, hexane; poorly sol. H₂O. λ_{\max} 239 (ε 40000); 269 (ε 5000) (MeOH) (Berdy). λ_{\max} 235 (ε 40000); 273 (ε 12000) (MeOH/HCl) (Berdy).

▶ LD₅₀ (mus, ipr) 3.24 mg/kg.

10'-Me ether: Piericidin B₁ [16891-54-6]

C₂₆H₃₉NO₄ 429.598

No phys. props. reported. λ_{\max} 232 (ε 31300); 237 (ε 31900); 267 (ε 5101) (MeOH).

10'-Me ether, 11',12'-epoxide: Piericidin D₁ [58947-82-3]

C₂₆H₃₉NO₅ 445.598

From *Streptomyces pactum*. NADH oxidase inhibitor. Sol. MeOH, hexane; poorly sol. H₂O. λ_{\max} 232 (ε 38000); 239 (ε 40000); 270 (ε 9000) (MeOH) (Berdy). λ_{\max} 235 (ε 35000); 275 (ε 12000) (MeOH/HCl) (Berdy).

▶ LD₅₀ (mus, ipr) 3.62 mg/kg.

O⁵-De-Me: Antibiotic NK 170204A. NK 170204A [161161-45-1]

C₂₄H₃₅NO₄ 401.545

Prod. by *Streptomyces* sp. NK 170204. Relationship with Piericidin A is tentative; stereochem. not fully descr. and their interconversion not reported.

13'-Hydroxy, 10'-O-β-D-glucopyranoside: 13-Hydroxyglucopiericidin A [132150-13-1]

C₃₁H₄₇NO₁₀ 593.713

From *Streptomyces* sp. OM-5689. Shows antitumour props. Yellow oil. Sol. EtOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{20}$ +20 (c, 0.2 in MeOH). λ_{\max} 239 (ε 40000); 268 (ε 5300) (MeOH) (Derep).

5-Demethoxy: Antibiotic Mer-A 2026B. Mer-A 2026B [144357-07-3]

C₂₄H₃₅NO₃ 385.545

From *Streptomyces karnatakensis* and *Streptomyces pactum*. Vasodilator, vasodepressor. Yellow oil. Sol. MeOH, butanol; poorly sol. H₂O. $[\alpha]_D^{24}$ -1.1 (c, 0.4 in MeOH). Abs. config. not determined. λ_{\max} 204 (ε 32700); 235 (ε 35000); 274 (ε 12000) (MeOH/HCl) (Derep). λ_{\max} 235 (sh) (ε 40500); 240 (ε 42000); 272 (ε 5800) (MeOH/NaOH) (Derep). λ_{\max} 223 (ε 33400); 235 (sh) (ε 35000); 238 (ε 37600); 264 (sh) (ε 11000) (MeOH) (Derep).

Piericidin A₂

IT 143C. Antibiotic IT 143C [58947-74-3]


C₂₆H₃₉NO₄ 429.598

From *Streptomyces pactum*. Inhibits NADH oxidase. Sol. MeOH, hexane; poorly sol. H₂O. λ_{\max} 235 (ε 20500); 272 (ε 10000) (MeOH/HCl) (Derep). λ_{\max} 238 (sh) (ε 24000) (MeOH/NaOH) (Derep). λ_{\max} 225 (ε 20500); 265 (sh) (ε 6000) (MeOH) (Derep). λ_{\max} 204 (ε 42400); 229 (ε 23200) (MeOH) (Berdy).

10'-Me ether: Piericidin B₂ [58947-76-5]

- $C_{27}H_{41}NO_4$ 443.625
No phys. props. reported. λ_{max} 225 (€ 20500); 265 (sh) (€ 6000) (MeOH).
- 11',12'-Epoxide: Piericidin C₂**
[58947-81-2]
 $C_{26}H_{39}NO_5$ 445.598
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. Sol. MeOH, Et₂O; poorly sol. H₂O. λ_{max} 235 (€ 20500); 272 (€ 10000) (MeOH/HCl) (Derep). λ_{max} 238 (sh) (€ 24000) (MeOH/NaOH) (Derep). λ_{max} 225 (€ 20500); 265 (sh) (€ 6000) (MeOH) (Derep). λ_{max} 235 (€ 21000); 275 (€ 11000) (MeOH/HCl) (Berdy).
- LD₅₀ (mus, ipr) 2.26 mg/kg.
10'-Me ether, 11',12'-epoxide: Piericidin D₂
[58947-72-1]
 $C_{27}H_{41}NO_5$ 459.625
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. Sol. MeOH, hexane; poorly sol. H₂O. λ_{max} 235 (€ 20500); 272 (€ 10000) (MeOH/HCl) (Derep). λ_{max} 238 (sh) (€ 24000) (MeOH/NaOH) (Derep). λ_{max} 225 (€ 20500); 265 (sh) (€ 6000) (MeOH) (Derep). λ_{max} 225 (€ 20000) (MeOH) (Berdy).
- Piericidin A₃**
Shaoguanmycin A
[58947-78-7]
 $C_{27}H_{41}NO_4$ 443.625
From *Streptomyces pactum* and *Streptomyces flaveolus shaoguanensis*. Inhibits NADH oxidase. λ_{max} 232 (€ 39500); 239 (€ 40500); 268 (€ 5300) (MeOH) (Derep).
- 10'-Me ether: Piericidin B₃**
[58947-71-0]
 $C_{28}H_{43}NO_4$ 457.652
No phys. props. reported.
- 11',12'-Epoxide: Piericidin C₃**
[58947-84-5]
 $C_{27}H_{41}NO_5$ 459.625
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. Sol. MeOH, Et₂O; poorly sol. H₂O. λ_{max} 239 (€ 40000); 269 (€ 5000) (MeOH) (Berdy). λ_{max} 235; 272 (EtOH) (Berdy).
- LD₅₀ (mus, ipr) 2.26 mg/kg.
10'-Me ether, 11',12'-epoxide: Piericidin D₃
[58947-75-4]
 $C_{28}H_{43}NO_5$ 473.651
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. Sol. MeOH, hexane; poorly sol. H₂O. λ_{max} 232; 239 (€ 40000); 270 (MeOH) (Berdy).
- 5-Demethoxy: Antibiotic Mer-A 2026A**
Mer-A 2026A
[144357-08-4]
 $C_{26}H_{39}NO_3$ 413.599
From *Streptomyces karnatakensis* and *Streptomyces pactum*. Vasodilator. Yellow oil. Sol. MeOH, butanol; poorly sol. H₂O. $[\alpha]_D^{22}$ -12.3 (c, 1.8 in MeOH). Abs. config. not determined. λ_{max} 204 (€ 32700); 235 (€ 35000); 274 (€ 12000) (MeOH/HCl) (Derep).
- λ_{max} 235 (sh) (€ 40500); 240 (€ 42000); 272 (€ 5800) (MeOH/NaOH) (Derep). λ_{max} 223 (€ 33700); 235 (sh) (€ 35000); 239 (€ 37400) (MeOH) (Derep).
- LD₅₀ (mus, ivn) 0.5 mg/kg, LD₅₀ (mus, ivn) 250-500 mg/kg.
- Piericidin A₄** [58947-77-6]
 $C_{28}H_{43}NO_4$ 457.652
From *Streptomyces pactum* and *Streptomyces piericidicus*. Inhibits NADH oxidase. Sol. MeOH, hexane; poorly sol. H₂O. λ_{max} 225 (€ 20000); 269 (MeOH) (Berdy).
- 11',12'-Epoxide: Piericidin C₄**
[58947-80-1]
 $C_{28}H_{43}NO_5$ 473.651
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. λ_{max} 225 (€ 20000) (MeOH) (Berdy). λ_{max} 235 (€ 21000); 275 (€ 11000) (MeOH/HCl) (Berdy).
- LD₅₀ (mus, ipr) 3.62 mg/kg.
10'-Me ether: Piericidin B₄
[58966-77-1]
 $C_{29}H_{45}NO_4$ 471.679
No phys. props. reported.
- 10'-Me ether, 11',12'-epoxide: Piericidin D₄**
[58947-73-2]
 $C_{29}H_{45}NO_5$ 487.678
From *Streptomyces pactum* and *Streptomyces piericidicus*. NADH oxidase inhibitor. Sol. MeOH, hexane; poorly sol. H₂O. λ_{max} 225 (€ 20000) (MeOH) (Berdy).
- Piericidin A₅**
 $C_{26}H_{39}NO_4$ 429.598
This component not yet characterised.
- 10'-Me ether: Piericidin B₅**
[149598-65-2]
 $C_{27}H_{41}NO_4$ 443.625
Pale yellow oil. $[\alpha]_D^{24}$ -12.5 (c, 0.2 in MeOH). λ_{max} 232 (€ 31300); 237 (€ 31900); 267 (€ 5010) (MeOH).
- 10'-Me ether, N-oxide: Piericidin B₅N-oxide**
[149665-01-0]
From a *Streptomyces* sp. Pale yellow oil. $[\alpha]_D^{24}$ -8 (c, 0.2 in MeOH). λ_{max} 226 (€ 34100); 238 (€ 29700); 246 (sh) (€ 20800); 267 (€ 7570) (MeOH).
- 11'R,12'R-Epoxide: Piericidin C₅**
 $C_{26}H_{39}NO_5$ 445.598
Prod. by a marine-derived *Streptomyces* sp. Cell division inhibitor. Pale yellow oil. $[\alpha]_D^{25}$ +7 (c, 0.12 in MeOH). λ_{max} 231 (€ 32000); 236 (€ 32300); 267 (€ 5100) (MeOH).
- Piericidin A₆**
 $C_{27}H_{41}NO_4$ 443.625
Component not yet characterised.
- 11'R,12'R-Epoxide: Piericidin C₆**
 $C_{27}H_{41}NO_5$ 459.625
Prod. by a marine-derived *Nocardioidea* sp. Cell division inhibitor. Pale yellow oil. $[\alpha]_D^{25}$ +34 (c, 0.03 in MeOH). λ_{max} 225 (€ 20600); 267 (€ 6000) (MeOH).
- Antibiotic IT143A**
IT 143A
[183485-32-7]
 $C_{29}H_{43}NO_4$ 469.663
Prod. by *Streptomyces* sp. IT 143. Pale yellow oil. $[\alpha]_D$ -13.9 (c, 0.2 in MeOH). Stereochem. not confirmed. λ_{max} 204 (€ 45600); 235 (€ 34300) (MeOH).
- 11S*,12S*-Epoxide: Piericidin C₈**
[939994-97-5]
 $C_{29}H_{43}NO_5$ 485.662
Prod. by the marine-derived *Streptomyces* sp. YM14-060. Cytotoxic. Oil. $[\alpha]_D^{23}$ +13 (c, 0.37 in MeOH). λ_{max} 225 (€ 21600); 267 (€ 6000) (MeOH).
- Antibiotic IT 143B**
IT 143B
[183485-34-9]
 $C_{28}H_{41}NO_4$ 455.636
Prod. by *Streptomyces* sp. IT 143. Pale yellow oil. $[\alpha]_D$ -26.3 (c, 0.1 in MeOH). Stereochem. not confirmed. λ_{max} 204 (€ 47600); 239 (€ 48100) (MeOH).
- 11S*,12S*-Epoxide: Piericidin C₇**
[939994-96-4]
 $C_{28}H_{41}NO_5$ 471.636
Prod. by a marine-derived *Streptomyces* sp. YM14-060. Cytotoxic. Oil. $[\alpha]_D^{23}$ +7 (c, 0.21 in MeOH). λ_{max} 231 (€ 30000); 237 (€ 30400); 269 (€ 3800) (MeOH).
- 5-Demethoxy: 5-Demethoxy IT 143B**
Antibiotic JBIR-02 *JBIR-02*
 $C_{27}H_{39}NO_3$ 425.61
Prod. by *Actinomyces* sp. Stamm K 17/9 and *Streptomyces* sp. ML55. Nuclear export inhibitor. Brown oil or solid. Mp 41-50°. $[\alpha]_D^{20}$ -17 (c, 0.1 in MeOH). λ_{max} 223 (log € 4.4); 240 (log € 4.45); 265 (sh); 294 (sh) (MeOH).
- Takahashi, N. *et al.*, *Agric. Biol. Chem.*, 1963, **27**, 583; 798 (*isol*)
Gutman, M. *et al.*, *Biochem. Biophys. Res. Commun.*, 1969, **37**, 615
Tanabe, M. *et al.*, *J.O.C.*, 1970, **35**, 2087 (*biosynth*)
Yoshida, S. *et al.*, *Agric. Biol. Chem.*, 1977, **41**, 587; 849; 855 (*isol*, *pmr*, *cmr*, *ms*)
Jansen, R. *et al.*, *Tet. Lett.*, 1983, **24**, 5485 (*stereochem*)
Liu, J. *et al.*, *CA*, 1984, **100**, 171237 (*isol*, *struct*)
Matsumoto, M. *et al.*, *J. Antibiot.*, 1987, **40**, 149-156 (*Glucopiericidins A-B*)
Mori, H. *et al.*, *J. Antibiot.*, 1990, **43**, 1329-1331 (*13-Hydroxyglucopiericidin A*)
Kimura, K. *et al.*, *J. Antibiot.*, 1990, **43**, 1341-1343 (*4'-Rhampopiericidin A₁*)
Iwasaki, H. *et al.*, *J. Antibiot.*, 1991, **44**, 451-452 (*3'-Deoxytalo piericidin A₁*)
Nishioka, H. *et al.*, *J. Antibiot.*, 1991, **44**, 1283-1285; 1993, **46**, 564-568 (*Piericidin B₁ N-oxide, Piericidin B₅, Piericidin B₅ N-oxide*)
Japan. Pat., 1994, 94 293 736; *CA*, **122**, 158767c (*NK 170204A*)
Kominato, K. *et al.*, *J. Antibiot.*, 1995, **48**, 99; 103 (*Mer-A 2026*)
Kimura, K. *et al.*, *J. Antibiot.*, 1996, **49**, 697 (*SN 198*)
Urakawa, A. *et al.*, *J. Antibiot.*, 1996, **49**, 1052-1055 (*IT 143*)
Kubota, N.K. *et al.*, *Bioorg. Med. Chem.*, 2003, **11**, 4569-4575 (*Piericidins C₅, C₆*)
Strösch, K. *et al.*, *Dissertation*, Univ. of Göttingen, 2003, (*5-Demethoxy IT 143B*)

- Schnermann, M.J. *et al.*, *J.A.C.S.*, 2006, **128**, 11799-11807 (*Piericidins A₁, B₁*, *synth*)
 Hayakawa, Y. *et al.*, *J. Antibiot.*, 2007, **60**, 196-200; 201-203 (*Piericidins C₇, C₈*)
 Ueda, J. *et al.*, *J. Antibiot.*, 2007, **60**, 459-462 (*JBIR-02*)

Piericidinol A₁**P-422**C₂₅H₃₇NO₅ 431.571**10-O-β-D-Glucopyranoside: Glucopiericidinol A₁**

[125535-05-9]

C₃₁H₄₇NO₁₀ 593.713

From *Streptomyces* sp. OM-5689. Active against *Piricularia oryzae*. Cytotoxic. Yellow oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D¹⁸ +38 (c, 0.08 in CHCl₃). λ_{max} 226; 290 (MeOH/HCl) (Derep). λ_{max} 212; 234 (sh); 290 (MeOH/NaOH) (Derep). λ_{max} 220; 240 (sh); 293 (MeOH) (Derep). λ_{max} 222; 296 (MeOH) (Berdy). λ_{max} 230; 290 (MeOH-HCl) (Berdy). λ_{max} 218; 280 (MeOH-NAOH) (Berdy).

3-Epimer, 10-O-β-D-glucopyranoside:**Glucopiericidinol A₂**

[125591-38-0]

C₃₁H₄₇NO₁₀ 593.713

Prod. by *Streptomyces* sp. OM-5689. Active against *P. oryzae*. Cytotoxic. Yellow oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D¹⁸ -20 (c, 0.06 in CHCl₃). λ_{max} 226; 290 (MeOH/HCl) (Derep). λ_{max} 212; 234 (sh); 290 (MeOH/NaOH) (Derep). λ_{max} 220; 240 (sh); 293 (MeOH) (Derep). λ_{max} 216; 290 (MeOH) (Berdy). λ_{max} 230; 290 (MeOH-HCl) (Berdy). λ_{max} 218; 292 (MeOH-NaOH) (Berdy).

Funayama, S. *et al.*, *J. Antibiot.*, 1989, **42**, 1734-1740 (*isol, struct*)


Pillijanine**P-423**C₁₅H₂₀N₂O 244.336

Struct. unknown. Alkaloid from *Lycopodium saururus* (Lycopodiaceae). Mp 64-65°.

Arata, P.N. *et al.*, *Gazz. Chim. Ital.*, 1892, **22**, 146-157 (*isol*)

Pilocarpine, BAN, USAN**P-424**

3-Ethylidihydro-4-(1-methyl-1H-imidazol-5-yl)methyl-2(3H)-furanone, 9CI. Ocularpiline. Ocusert P 20. Ocusert Pilo 40. Pilocarpol. Syncarpine
 [92-13-7]

C₁₁H₁₆N₂O₂ 208.26

Alkaloid from *Pilocarpus microphyllus* and several other *Pilocarpus* spp. (Ruta-

ceae). Cholinergic muscarinic agonist used as a miotic, especially for glaucoma, and in the treatment of nephritis. Exp. convulsant. Used in animal models of epilepsy. Mp 34°. Bp₅ 260°. [α]_D²⁰ +100.5 (H₂O). pK_{a1} 7.15; pK_{a2} 12.57 (20°). Log P -0.2 (calc).

► Adverse ocular effects reported when used therapeutically. LD₅₀ (rat, orl) 402 mg/kg.

Hydrochloride: Pilocarpine hydrochloride. USAN. Akarpine. Almocarpin. Pilo-miotin. Pilopine V.S. Pilovisc
 [54-71-7]

Cryst. (EtOH). Mp 204-205°. [α]_D¹⁸ +91. Component of Pilocar.

► Human systemic effect from ocular administration. Adverse ocular effects reported when used therapeutically. LD₅₀ (mus, orl) 200 mg/kg, exp. reprod. and teratogenic effects. TK1450000

Nitrate: Licarpin. Pilofrin. Pilagan

[148-72-1]

Mp 173-174° (dec.). [α]_D +77.

► TK1455000

Picrate: Mp 159-160°.

N-De-Me: Pilocarpidine

[127-67-3]

C₁₀H₁₄N₂O₂ 194.233

Alkaloid from the leaves of *Pilocarpus jaborandi* (Rutaceae). Pharmacol. resembles that of Pilocarpine, but weaker. Syrup. Sol. H₂O. [α]_D +81.3 (H₂O). Dec. on dist.

N-De-Me; nitrate: Mp 137°. [α]_D +73.2 (H₂O).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 624C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 86C (*nmr*)

Jowett, H.A.D. *et al.*, *J.C.S.*, 1900, **77**, 473; 851 (*struct*)

Späth, E. *et al.*, *Ber.*, 1925, **58**, 513 (*struct, Pilocarpidine*)

Burtles, R. *et al.*, *J.C.S.*, 1925, **127**, 581 (*struct, Pilocarpidine*)

Preobrashenski, N.A. *et al.*, *Ber.*, 1933, **66**, 1536; 1936, **69**, 1835 (*synth, Pilocarpidine*)

Hill, R.K. *et al.*, *Tetrahedron*, 1966, **22**, 2889 (*abs config*)

Ben-Bassat, A. *et al.*, *Isr. J. Chem.*, 1972, **10**, 385 (*uv, ir, pmr, ms*)

de Graw, J.I. *et al.*, *Tetrahedron*, 1972, **28**, 967 (*synth*)

Simeral, L. *et al.*, *Org. Magn. Reson.*, 1974, **6**, 26 (*cmr*)

Brochmann-Hanssen, E. *et al.*, *Planta Med.*, 1975, **28**, 1-5 (*biosynth*)

Noordam, A. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1981, **100**, 441 (*synth*)

Martindale, *The Extra Pharmacopoeia, 28th edn.*, Pharmaceutical Press, 1982, 4526; 4525

Al-Badr, A.A. *et al.*, *Anal. Profiles Drug Subst.*, 1983, **12**, 385 (*rev, uv, ir, pmr, cmr, ms, anal*)

Turski, L. *et al.*, *Synapse (N.Y.)*, 1989, **3**, 154 (*pharmacol, rev*)

Belletiere, J.L. *et al.*, *J. Nat. Prod.*, 1992, **55**, 194 (*synth*)

Shapiro, G. *et al.*, *Tet. Lett.*, 1992, **33**, 2447 (*synth*)

Gaggelli, E. *et al.*, *Can. J. Chem.*, 1993, **71**, 738 (*pmr, cmr*)

Horne, D.A. *et al.*, *J.O.C.*, 1993, **58**, 62 (*synth*)

Dener, J.M. *et al.*, *J.O.C.*, 1993, **58**, 1159 (*synth*)

Wiseman, L.R. *et al.*, *Drugs*, 1995, **49**, 143 (*rev*)

Lei, A. *et al.*, *J.A.C.S.*, 2002, **124**, 8198-8199 (*synth*)

Abreu, I.N. *et al.*, *Rapid Commun. Mass Spectrom.*, 2007, **21**, 1205-1213 (*biosynth, ms*)


Sawaya, A.C.H.F. *et al.*, *Molecules*, 2008, **13**, 1518-1529 (*ms, chromatog*)

Csuk, R. *et al.*, *Tetrahedron*, 2008, **64**, 9384-9387 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, PIF000; PIF250; PIF500

Pilocereine**P-425**

[2552-47-8]

C₄₅H₆₅N₃O₆ 744.025

Alkaloid from *Lophocereus australis*, *Lophocereus schottii* (various forms), *Lophocereus gatesii*, *Pachycereus marginatus* and *Pilocereus sargentianus* (said to be synonymous with *Cereus sargentianus* but taxonomically difficult to reconcile) (Cactaceae). Antineoplastic agent. Cryst. (EtOAc). Mp 176.5-177°. Log P 10.25 (uncertain value) (calc). Opt. inactive. λ_{max} 284 (ε 7000) (MeOH) (Berdy).

Hydrochloride (1:3):

Cryst. + 3H₂O. Mp 228-232° dec.

O-Ac:

Cryst. (Et₂O/Me₂CO). Mp 186-186.5°.

Me ether:

Cryst. (hexane). Mp 103-105° Mp 153.5-154.5° (double Mp).

Diastereoisomer: Piloceredine

C₄₅H₆₅N₃O₆ 744.025

Alkaloid from *Lophocereus schottii* (Cactaceae). Cryst. (Me₂CO). Mp 165-166°. Opt. inactive.

Diastereoisomer, Ac:

Cryst. (petrol). Mp 133-134°.

Diastereoisomer, Me ether:

Cryst. (hexane). Mp 150-152°.

Heyl, G. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1901, **239**, 451 (*isol*)

Djerassi, C. *et al.*, *J.A.C.S.*, 1953, **75**, 3632; 1954, **76**, 3215; 1956, **79**, 2203; 1962, **84**, 3210 (*isol, uv, ir, struct*)

Djerassi, C. *et al.*, *Tetrahedron*, 1958, **2**, 58 (*Piloceredine*)


Tomita, M. *et al.*, *Chem. Pharm. Bull.*, 1963, **11**, 1477; 1484 (*uv, ir, struct*)

Bessho, K. *et al.*, *Chem. Pharm. Bull.*, 1963, **11**, 1491; 1500; 1504; 15 (*struct*)

- Franck, B. *et al.*, *Annalen*, 1965, **685**, 207
(*synth, uv, ir, pmr, ms*)
O'Donovan, D.G. *et al.*, *J.C.S. (C)*, 1971, 2398
(*biosynth*)
West, L.G. *et al.*, *Phytochemistry*, 1975, **14**, 291
(*isol*)

Piloceanine P-426

3-(2-Hydroxy-3-methylbutyl)-8-methoxy-1-methyl-4(1H)-quinolinone



$C_{16}H_{21}NO_3$ 275.347
Alkaloid from the leaves of *Platydesma campanulata* (Rutaceae). Noncryst.; cryst. (EtOH)(as picrate). Mp 216° dec.(picrate).

Werny, F. *et al.*, *Tetrahedron*, 1963, **19**, 1293
(*isol, uv, ir, struct*)

Pilosamycin P-427


Closely related to Ferrimycin A₁, F-37. Prod. by *Streptomyces galilaeus* and *Streptomyces griseoflavus*. Siderophore. Antibacterial agent. Orange powder. Sol. MeOH, DMF, H₂O; fairly sol. EtOAc; poorly sol. butanol, hexane. λ_{max} 229 (E1%/1cm 317); 319 (E1%/1cm 28.3); 430 (E1%/1cm 26.4) (H₂O) (Berdy).

► LD₅₀ (mus, scu) 1200 mg/kg.

Ger. Pat., 1960, 1 089 122; *CA*, **55**, 27768f

Pilosine P-428

Carpidine
[13640-28-3]



$C_{16}H_{18}N_2O_3$ 286.33
Alkaloid from *Pilocarpus microphyllus* (Rutaceae). Narcotic to mammals. Other pharmacol. resembles that of Pilocarpine, P-424 but much weaker. Mp 101-104° (dihydrate) Mp 171-172° (anhyd.) (187°). This alkaloid was difficult to purify and pharmacol. activity was prob. investigated on impure samples.

Sulfate (2:1): Mp 194-195°. $[\alpha]_D +21$ (c, 4.5 in H₂O).

6-Ketone: Pilosinone

[942060-99-3]

$C_{16}H_{16}N_2O_3$ 284.314

Alkaloid from *Pilocarpus microphyllus*. Stereochem. not determined.

6-Deoxy, 5,6-didehydro(Z-): **Anhydriposine**

[53336-50-8]

$C_{16}H_{16}N_2O_2$ 268.315

Constit. of *Pilocarpus microphyllus*.

5-Epimer: Isopilosine. Carpiline

[491-88-3]

$C_{16}H_{18}N_2O_3$ 286.33

Alkaloid from *Pilocarpus microphyllus* (Rutaceae). Cryst. (EtOH). Mp 182-183°. $[\alpha]_D^{20} +37.6$ (c, 1 in EtOH).

5,6-Diepimer: Episopilosine

[38993-92-9]

$C_{16}H_{18}N_2O_3$ 286.33

Alkaloid from *Pilocarpus microphyllus* (Rutaceae). Cryst. (MeOH). Mp 179-180°. $[\alpha]_D^{20} -44$ (c, 1 in EtOH).

Link, H. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 1053; 1974, **57**, 2199 (*synth, ir, pmr, ms, abs config, bibl*)

Löwe, W. *et al.*, *Annalen*, 1973, 1476 (*struct, pmr*)

Tedeschi, E. *et al.*, *J.O.C.*, 1974, **39**, 1864 (*uv, pmr, ms, struct*)


Sarel, S. *et al.*, *Tet. Lett.*, 1975, 97 (*ir, struct, Isopilosine*)

Abreu, I.N. *et al.*, *Rapid Commun. Mass Spectrom.*, 2007, **21**, 1205-1213 (*Anhydriposine, Pilosinone*)

Sawaya, A.C.H.F. *et al.*, *Molecules*, 2008, **13**, 1518-1529 (*ms, chromatog*)

Pilosinone P-429

[38993-86-1]



Absolute Configuration

$C_9H_{12}N_2O_2$ 180.206


Alkaloid from *Pilocarpus microphyllus*.

Shapiro, G. *et al.*, *Tet. Lett.*, 1992, **33**, 2447-2450 (*synth*)

Abreu, I.N. *et al.*, *Rapid Commun. Mass Spectrom.*, 2007, **21**, 1205-1213 (*isol*)

Piloturine P-430

[69433-49-4]



$C_{16}H_{18}N_2O_3$ 286.33

Related to Pilosine, P-428. Alkaloid from *Pilocarpus microphyllus*. Isoln. not certain.

5,6-Diepimer: Episopiloturine

[69460-80-6]

$C_{16}H_{18}N_2O_3$ 286.33

Alkaloid from *Pilocarpus microphyllus*. Cryst. (EtOH). Mp 218-219°. $[\alpha]_D^{20} -11$ (c, 0.5 in EtOH).

Voigtlaender, H.W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 927-935

Pimpinellidine P-431

[42613-57-0]


$C_{27}H_{45}NO_3$ 431.657

Steroidal alkaloid, prob. of Solanum type. Struct. unknown but may be 23-Hydroxytomatidine. Aglycone from hydrol. of leaves and stems of *Lycopersicon pimpinellifolium* (currant tomato) and from *Solanum eryanthum* (Solanaceae). Needles (MeOH aq.). Mp 172-173°. $[\alpha]_D^{20} +14.3$ (c, 0.7 in MeOH).

Schreiber, K. *et al.*, *Phytochemistry*, 1966, **5**, 707 (*isol, ir*)

Pingbeidinose P-432

[125309-97-9]



$C_{34}H_{57}NO_9$ 623.826


Alkaloid from the stem and leaf of *Fritillaria ussuriensis*. Cryst. Mp 242-243°. $[\alpha]_D^{25} +6.9$ (c, 0.14 in MeOH).

Xu, D.M. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 668-672; *CA*, **112**, 95507p (*isol, struct*)

Pingbeinine P-433

28-Methyl-16,28-secosolanid-5-ene-3,16,25-triol, 9CI

[131984-89-9]



$C_{28}H_{47}NO_3$ 445.684

Alkaloid from the leaves of *Fritillaria ussuriensis* (Liliaceae). Needles (MeOH). Mp 223-235°. $[\alpha]_D -32.8$ (c, 0.09 in MeOH).

3-O-β-D-Glucopyranoside: Pingbeininoside

[131984-90-2]


$C_{34}H_{57}NO_8$ 607.826

Alkaloid from the leaves of *Fritillaria ussuriensis* (Liliaceae). Needles (MeOH). Mp 244-246°. $[\alpha]_D -4.57$ (c, 0.164 in MeOH).

Xu, D.-M. *et al.*, *J. Nat. Prod.*, 1990, **53**, 549 (*isol, ir, pmr, cmr, ms, struct*)

Pingbeinone

[125292-91-3]

C₂₆H₄₁NO₃ 415.615


Alkaloid from *Fritillaria ussuriensis* (Liliaceae). Needles (MeOH). Mp 200-202°. [α]_D -22 (c, 0.8 in CHCl₃). New subgroup of steroidal alkaloids.

[125356-74-3]

Kitamura, Y. *et al.*, *Tetrahedron*, 1989, **45**, 7281 (*isol, pmr, cmr, struct*)

Pinnaic acid

[178115-91-8]

C₂₃H₃₆ClNO₄ 425.994

Alkaloid from the Okinawan bivalve *Pinna muricata*. Phospholipase A2 inhibitor. Antiinflammatory agent.

2-Sulfoethylamide: Tauopinnaic acid

[178115-92-9]

C₂₅H₄₁ClN₂O₆S 533.127

Alkaloid from *Pinna muricata*. Phospholipase A2 inhibitor. Antiinflammatory agent.

Chou, T. *et al.*, *Tet. Lett.*, 1996, **37**, 3871-3874 (*isol, pmr, cmr, struct*)

Carson, M.W. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 4450-4452; 4453-4456 (*synth, abs config*)

Hayakawa, I. *et al.*, *Heterocycles*, 2003, **59**, 441-444 (*synth*)

Kuramoto, M. *et al.*, *Mar. Drugs*, 2004, **2**, 39-54 (*rev, biosynth*)

Matsumura, Y. *et al.*, *Org. Lett.*, 2004, **6**, 965-968 (*synth*)


Andrade, R.B. *et al.*, *Org. Lett.*, 2005, **7**, 5733-5735 (*synth*)

Xu, S. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 5746-5749 (*synth*)

Wu, H. *et al.*, *Tetrahedron*, 2007, **63**, 6454-6461 (*synth*)

Pinnamine

[298197-30-5]

C₁₃H₁₉NO₂ 221.299

Absolute Configuration

P-434

Alkaloid from the Okinawan bivalve *Pinna muricata*. Toxin. Oil. [α]_D²⁷ +71.2 (c, 0.04 in MeOH) (synthetic).

► Highly toxic, LD₉₉ 0.5mg/kg (mus).


Takada, N. *et al.*, *Tet. Lett.*, 2000, **41**, 6425-6428 (*isol, abs config*)

Kigoshi, H. *et al.*, *Tet. Lett.*, 2001, **42**, 7469-7471 (*synth*)

Kuramoto, M. *et al.*, *Mar. Drugs*, 2004, **2**, 39-54 (*tox*)

Pinnatoxin A

[160759-36-4]



R = COOH

Absolute Configuration

C₄₁H₆₁NO₉ 711.934

Exists as zwitterion. Neutral form shown. Principal toxin from the bivalves *Pinna muricata*, *Pinna attenuata* and *Pinna atropurpurea*. Calcium channel activating agent, smooth muscle contractant showing tetrodotoxin-like effect. λ_{\max} 216 (EtOH) (Berdy).

► Highly toxic, LD₅₀ 2.7 μ g/MU (mus, ip).

Uemura, D. *et al.*, *J.A.C.S.*, 1995, **117**, 1155-1158 (*isol, pmr, cmr*)

Chou, T. *et al.*, *Tet. Lett.*, 1996, **37**, 4023-4026 (*stereochem*)

McCauley, J.A. *et al.*, *J.A.C.S.*, 1998, **120**, 7647-7648 (*synth*)

Falk, M. *et al.*, *Tetrahedron*, 2001, **57**, 8659-8665 (*abs config*)

Sakamoto, S. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 6505-6510 (*synth*)

Kuramoto, M. *et al.*, *Mar. Drugs*, 2004, **2**, 39-54 (*rev, tox*)

Stivala, C.E. *et al.*, *J.A.C.S.*, 2008, **130**, 3774-3776 (*synth*)

Pinnatoxin B

P-438

[160903-44-6]

As Pinnatoxin A, P-437 with

R = -³⁴CH(NH₂)COOH(S-)C₄₂H₆₄N₂O₉ 740.976

Exists as a zwitterion. Minor toxin from *Pinna muricata*.

► Highly toxic, LD₅₀ (mus, ip) 0.99 μ g/MU as mixture with Pinnatoxin C.

34-Epimer: Pinnatoxin C

[167228-66-2]

C₄₂H₆₄N₂O₉ 740.976Minor toxin from *Pinna muricata*.

► Highly toxic.

Takada, N. *et al.*, *Tet. Lett.*, 2001, **42**, 3491-3494 (*isol*)


Kuramoto, M. *et al.*, *Mar. Drugs*, 2004, **2**, 39-54 (*rev, tox*)

Matsuura, F. *et al.*, *Org. Lett.*, 2006, **8**, 3327-3330 (*synth, abs config*)

Pinnatoxin D

P-439

[167228-67-3]



Absolute Configuration

C₄₅H₆₇NO₁₀ 782.025

Exists as zwitterion. Neutral form shown. Toxin from the bivalves *Pinna muricata*, *Pinna attenuata* and *Pinna atropurpurea*. Calcium channel activating agent. Less toxic to mammals than the other pinnatoxins but the most potent against murine leukaemia. Solid. [α]_D +42.5 (c, 0.5 in MeOH). λ_{\max} 226 (MeOH).

► Highly toxic, LD₅₀ (mus, ip) >10 μ g/MU.

Chou, T. *et al.*, *Tet. Lett.*, 1996, **37**, 4023-4026; 4027-4030 (*isol, uv, pmr, cmr, struct*)


Falk, M. *et al.*, *Tetrahedron*, 2001, **57**, 8659-8665 (*abs config*)

Kuramoto, M. *et al.*, *Mar. Drugs*, 2004, **2**, 39-54 (*rev, tox*)

Pinolinone

P-440

3,4-Dihydro-3,4-dihydroxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone



Relative Configuration


C₁₅H₁₉NO₃ 261.32

Alkaloid from the roots of *Boronia pinnata*. Oil. [α]_D -9.4 (c, 0.13 in CHCl₃). λ_{\max} 211 ; 258 (MeOH).

Ito, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1344-1348

Pipbinine

P-441


C₃₄H₃₈N₂O₆ 570.684

Alkaloid from the fruit of *Piper nigrum* (pepper). Amorph. powder. λ_{\max} 250 (ϵ 11600); 266 (ϵ 12800); 298 (ϵ 25200); 316 (ϵ 26900); 323 (ϵ 33400); 355 (ϵ 40500); 372 (ϵ 45000) (MeOH).

Siddiqui, B.S. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 660-666 (*isol, pmr, cmr, ms*)

Piperarborenine C

P-442

C₃₃H₃₄N₂O₁₀ 618.639

Alkaloid from the stems of *Piper arborescens*. Brownish oil. Racemic. λ_{\max} 219 (log ϵ 4.58); 260 (sh) (log ϵ 3.94); 280 (sh) (log ϵ 3.56) (MeOH).

3'-Demethoxy: Piperarborenine E


C₃₂H₃₂N₂O₉ 588.613

Alkaloid from the stems of *Piper arborescens*. Brownish oil. Racemic. λ_{\max} 210 (log ϵ 4.5); 280 (sh) (log ϵ 3.56) (MeOH).

Tsai, I.-L. *et al.*, *Planta Med.*, 2005, **71**, 535-542 (*isol, pmr, cmr, ms*)

Piperarboresine

P-443

C₃₃H₃₄N₂O₁₀ 618.639


Alkaloid from the stems and leaves of *Piper arborescens*. Needles (MeOH). Mp 182-184°. λ_{\max} 213 (log ϵ 4.55); 240 (sh) (log ϵ 4.11) (MeOH).

Lee, F.-P. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 463-468 (*isol, pmr, cmr*)

Piperazinomycin

P-444

13-Oxa-4,20-diazatetracyclo[12.2.2.2^{3,6}.1^{8,12}]heneicosane-8,10,12(19),14,16,17-hexaen-11-ol, 9CI. Neoenactin C [83858-82-6]

C₁₈H₂₀N₂O₂ 296.368

Isol. from *Streptoverticillium olivoreticuli* subsp. *neoenacticus*. Active against fungi

and yeasts esp. *Trichophyton*. Amorph. powder. Sol. MeOH, Py, 4-methyl-2-pentanone; fairly sol. Me₂CO, CHCl₃; poorly sol. H₂O, hexane. Mp 102-104°. $[\alpha]_D^{23}$ +31.1 (c, 0.74 in MeOH). λ_{\max} 214 (ϵ 11200); 280 (ϵ 2040); 290 (sh) (ϵ 1690) (MeOH) (Derrep).

Tamai, S. *et al.*, *J. Antibiot.*, 1982, **35**, 1130 (*isol*)
Kameda, M. *et al.*, *J. Antibiot.*, 1982, **35**, 1137 (*cryst struct*)


Jung, M.E. *et al.*, *J.O.C.*, 1985, **50**, 4909 (*synth*)
Nishiyama, S. *et al.*, *Tet. Lett.*, 1986, **27**, 4481 (*synth*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1993, **115**, 11426; 1994, **116**, 1601 (*synth*)

Piperazirum

P-445

6-(1-Methylethyl)-3,5-bis(2-methylpropyl)-2-piperazinone, 3,5-Diisobutyl-6-isopropyl-2-piperazinone [943985-04-4]



Relative Configuration

C₁₅H₃₀N₂O 254.415


Alkaloid from the leaves of *Arum palaestinum*. Cytotoxic. Amorph. powder. $[\alpha]_D$ +50 (c, 0.02 in MeOH).

El-Desouky, S.K. *et al.*, *Tet. Lett.*, 2007, **48**, 4015-4017 (*isol, pmr, cmr*)

Pipercollosidine

P-446

[83029-38-3]



n = 1

C₁₈H₂₅NO₃ 303.4

Alkaloid from roots of *Piper collosum* (Piperaceae). Cryst. (cyclohexane). Mp 80-82°.

Pring, B.G. *et al.*, *J.C.S. Perkin I*, 1982, 1493 (*isol, synth, uv, ir, pmr, ms, struct*)

Pipercollosine

P-447

[83029-39-4]

As Pipercollosidine, P-446 with n = 2

C₂₀H₂₇NO₃ 329.438


Alkaloid from roots of *Piper collosum* (Piperaceae). Cryst. (cyclohexane or EtOH aq.). Mp 114-115°.

Pring, B.G. *et al.*, *J.C.S. Perkin I*, 1982, 1493 (*isol, synth, uv, ir, ms, pmr, struct*)

Piperchabamide C

P-448

1-[13-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,12-tridecatrienyl]piperidine, 9CI. 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienic acid piperidide

C₂₅H₃₃NO₃ 395.541


Alkaloid from the fruit of *Piper chiba* (Javanese long pepper). Oil. λ_{\max} 265 (log ϵ 4.52) (EtOH).

Morikawa, T. *et al.*, *Planta Med.*, 2004, **70**, 152-159 (*isol, pmr, cmr, ms*)

Pipericyclamide

P-449

5-Hydroxy-1-(2-methylpropyl)-5-(1-oxohexyl)-2-pyrrolidinone, 5-Hexanoyl-5-hydroxy-N-isobutyl-2-pyrrolidinone


C₁₄H₂₅NO₃ 255.356

Alkaloid from the roots of *Piper nigrum* (pepper). Oil. Racemic.

Wei, K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1005-1009 (*isol, pmr, cmr*)

Pipericyclobutanamide A

P-450

Relative Configuration
n = 1C₃₄H₃₈N₂O₆ 570.684

Alkaloid from the fruit of *Piper nigrum* (pepper). Amorph. powder. λ_{\max} 265; 271; 289 (hexane).

Fujiwara, Y. *et al.*, *Tet. Lett.*, 2001, **42**, 2497-2499

Pipericyclobutanamide B

P-451

As Pipericyclobutanamide A, P-450 with n = 2

C₃₆H₄₀N₂O₆ 596.722


Alkaloid from the fruit of *Piper nigrum* (pepper). Amorph. powder. λ_{\max} 285; 296; 309 (hexane).

Fujiwara, Y. *et al.*, *Tet. Lett.*, 2001, **42**, 2497-2499

Piperettine



P-452

1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,6-heptatrienyl]piperidine, 9CI. 1-[7-(3,4-Methylenedioxyphenyl)heptatrienyl]piperidine [583-34-6]

C₁₉H₂₁NO₃ 311.38

Alkaloid from the dried fruits of *Piper aurantiacum* and *Piper nigrum* (pepper) (Piperaceae). Yellow needles (C₆H₆/hexane). Mp 148°.


4,5-Dihydro- 1-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]piperidine.

Pipersintenamamide. 4,5-DihydropiperettineC₁₉H₂₃NO₃ 313.396Alkaloid from the whole plant of *Piper sintonense*. Cytotoxic. Needles (MeOH). Mp 116-118°. λ_{max} 215 (log ε 4.27); 261 (log ε 4.06) (EtOH).6,7-Dihydro: 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4-heptadienyl]piperidine, 9CI. 1-[7-(3,4-Methylenedioxyphenyl)-2,4-heptadienyl]piperidine. **Piperdardine**. *Piperarboricoline*. 6,7-Dihydropiperettine [188426-70-2]C₁₉H₂₃NO₃ 313.396Constit. of *Piper sintonense* and *Piper tuberculatum* var. *tuberculatum*. Pale yellow oil. λ_{max} 205 (log ε 3.9); 266 (log ε 4.04) (EtOH).Spring, F.S. *et al.*, *J.C.S.*, 1950, 1177 (*isol. struct., synth*)Rao, J.M. *et al.*, *Curr. Sci.*, 1974, **43**, 76 (*isol*)
Dehmlow, E.V. *et al.*, *J. Chem. Res., Synop.*, 1981, 106 (*synth*)De Araujo-Junior, J.X. *et al.*, *Phytochemistry*, 1997, **44**, 559 (*Piperdardine*)De Araujo-Junior, J.X. *et al.*, *Synth. Commun.*, 2001, **31**, 117-123 (*Piperdardine*)Chen, J.-J. *et al.*, *Planta Med.*, 2002, **68**, 980-985 (*Pipersintenamamide, Piperarboricoline*)**1-Piperettylpyrrolidine** P-4531-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,6-heptatrienyl]pyrrolidine, 9CI. 7-(3,4-Methylenedioxyphenyl)-2,4,6-heptatrienoic acid pyrrolidide. **Piperettyline**C₁₈H₁₉NO₃ 297.353**(E,E,E)-form** [52812-40-5]Alkaloid from the stems of *Piper tri-chostachyon* (Piperaceae). Yellow needles (EtOAc/petrol). Mp 147-149°.Singh, J. *et al.*, *Phytochemistry*, 1974, **13**, 677 (*isol. uv. ir. pmr. ms. struct*)Vig, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 1015 (*synth*)Likhitwitayawuid, K. *et al.*, *Tetrahedron*, 1987, **43**, 3689 (*isol. uv. ir. pmr. cmr. ms. struct*)Kaga, H. *et al.*, *Synlett*, 1994, 607 (*synth*)**Piperidine, 9CI** P-454*Hexahydropyridine*. *Azacyclohexane*. *FEMA 2908* [110-89-4]C₅H₁₁N 85.149

Present in free state in plants, e.g.

Psilocaulon absimile, *Petrosimonia monandra*, *Glycine max.* Residue present in piperidine alkaloids. Present in black pepper (*Piper nigrum*). Useful base in org. synth. Used for detn. of Co, Ni, Pt, Pd, Rh, Ir. Liq. with characteristic odour and a burning, peppery taste. Misc. H₂O; sol. org. solvs. d₄²⁰ 0.86. Mp -9°. Bp 106° Bp₂₀ 17.7°. n_D²⁰ 1.4534. pK_a 11.12 (25°). Forms a hydrate, Mp -14°.► Skin, eye and respiratory tract irritant. LD₅₀ (rat, orl) 400 mg/kg. LD₅₀ (rbt, skn) 320 mg/kg. Exp. reprod. and teratogenic effects (by inhalation). Highly flammable, fl. p. 3/16°. OES: long-term 1 ppm (Sk). TM3500000*Hydrochloride*: [6091-44-7]


Prisms (EtOH). Mp 244-245°.

► LD₅₀ (rat, orl) 133 mg/kg. TN0400000
Späth, E. *et al.*, *Ber.*, 1935, **68**, 2218 (*occur*)
Lambert, J.B. *et al.*, *J.O.C.*, 1983, **48**, 3982 (*cmr*)Rubiralta, M. *et al.*, *Piperidine: Structure, Preparation, Reactivity and Synthetic Applications of Piperidine*, (Std. Org. Chem. Vol. 43), Elsevier, Amsterdam, 1991, (*book*)**1-Piperidinetri-carboxaldehyde, 9CI, 8CI** P-455*N-Formylpiperidine* [2591-86-8]C₆H₁₁NO 113.159Alkaloid from *Piper nigrum*. Liq. Bp 222° Bp₁₀ 97-100°.

► TN0380000

Hydrobromide: Mp 103.5°.*Oxime*: [18600-89-0]C₆H₁₂N₂O 128.174

Mp 49-50°.

Di-Me acetal: 1-(*Dimethoxymethyl*)piperidine [5211-86-9]C₈H₁₇NO₂ 159.228Liq. Bp₁₅ 83°.*Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, **1**, 763D (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **1**, 1248A (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 785A (*ir*)de Benneville, P.L. *et al.*, *J.O.C.*, 1956, **21**, 772 (*synth*)Nefedov, B.L. *et al.*, *Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.)*, 1973, 2652 (*synth, bibl*)De Brauwere, J. *et al.*, *Bull. Soc. Chim. Belg.*, 1975, **84**, 167-177 (*isol*)Nair, R.S. *et al.*, *Fundam. Appl. Toxicol.*, 1992, **18**, 96 (*tox*)*Encyclopaedia of Reagents for Organic Synthesis*, (ed. Paquette, L.A.), Wiley, 1995, **4**, 2599-2600 (*use*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, FOH000**2-Piperidinetri-carboxylic acid, 9CI** P-456*Pipecolic acid*. *Hexahydropicolinic acid*. *Pipecolinic acid*. *Homoprolin*e† [535-75-1]*(S)-form*C₆H₁₁NO₂ 129.158


► TK6021000

(S)-form*L-form*

[3105-95-1]

Present in beans and other legumes, and in lesser quantities in other plants including barley, hops, teak and mushrooms. Also from the red alga *Amphiroa beauvoisii*. Needles (MeOH/Et₂O). Sol. H₂O. Mp 259-260°. [α]_D²⁵ -30.6 (c, 1 in H₂O).*Hydrochloride*: [2133-33-7]Mp 261-264°. [α]_D²⁵ -10.3 (c, 0.2 in H₂O) (91% ee).*N-γ-Glutamyl*: *γ-L-Glutamyl-L-pipecolic acid*C₁₁H₁₈N₂O₅ 258.274Isol. from the seeds of *Gleditsia caspica* (Caspian locust). Cryst. (H₂O). [α]_D²⁰ -66.8 (c, 0.5 in H₂O).*N-(4-Hydroxycinnamoyl)* (E)-: *Cirsiumamide*. *N-p-Coumaroylpipecolic acid* [158238-37-0]C₁₅H₁₇NO₄ 275.304Alkaloid from *Cirsium brevicaulis-irmutiense*. Shows antioxidant props.

[41994-45-0]

Dardenne, G. *et al.*, *Phytochemistry*, 1974, **13**, 1515 (*Glutamylpipecolic acid*)Sawaguchi, K. *et al.*, *CA*, 1994, **121**, 251185a (*Cirsiumamide*)Agami, C. *et al.*, *Synth. Commun.*, 2000, **30**, 2565-2572 (*S-form, synth, pmr*)Watanabe, L.A. *et al.*, *Tetrahedron: Asymmetry*, 2005, **16**, 903-908 (*R-form, S-form, synth*)Hou, D.-R. *et al.*, *Tetrahedron: Asymmetry*, 2005, **16**, 3858-3864 (*R-form, S-form, synth, pmr, cmr*)Chang, M.-Y. *et al.*, *Heterocycles*, 2006, **68**, 2365-2373 (*S-form, synth*)**2,4,5-Piperidinetri-carboxylic acid, 9CI** P-457*Hexahydroberberonic acid*

Relative Configuration

C₈H₁₁NO₆ 217.178


Various stereoisomers have been synthesised.

(2R*,4S*,5R*)-form [160699-23-0]Isol. from fruit bodies of the poisonous mushroom *Clitocybe acromelaga*.

Yamano, K. *et al.*, *Z. Naturforsch., C*, 1994, **49**, 707-711 (*isol*, *pmr*, *cmr*, *struct*)
 Hashimoto, K. *et al.*, *Heterocycles*, 1997, **46**, 581-588 (*synth*, *pmr*)

3,4,5-Piperidinetriol, 9CI P-458

3,4,5-Trihydroxypiperidine. 1-Deoxy-5-nornojirimycin. 1,5-Dideoxy-1,5-iminopentitol
 [152689-94-6]

(3 α ,4 α ,5 α)-formC₅H₁₁NO₃ 133.147**(3 α ,4 α ,5 α)-form**

cis,cis-form. 1,5-Dideoxy-1,5-iminoribitol
 [172588-13-5]
 [176597-75-4] β -Glucosidase inhibitor.
 Cryst. (MeOH aq.) (as hydrochloride).
 Mp 161-163° (dec.) Mp 185-187°
 (hydrochloride). *meso*-.

N-Benzyl: [177767-33-8]

C₁₂H₁₇NO₃ 223.271

Cryst. (EtOAc). Mp 133.8-134.8°.

[3R-(3 α ,4 α ,5 β)]-form

(-)-*cis,trans*-form. 1,5-Dideoxy-1,5-imino-D-arabinitol. 1,5-Dideoxy-1,5-imino-D-lyxitol

[130549-94-9] α -L-Fucosidase inhibitor.
 Cryst. (MeOH/Et₂O). Mp 182-185° (dec.). $[\alpha]_D^{25}$ -71 (c, 0.6 in MeOH).

Hydrochloride: [187144-38-3]

Cryst. (MeOH). Mp 194-195°. $[\alpha]_D^{20}$ -22
 (c, 0.8 in MeOH).

[3S-(3 α ,4 α ,5 β)]-form

(+)-*cis,trans*-form. 1,5-Dideoxy-1,5-imino-L-arabinitol. 1,5-Dideoxy-1,5-imino-L-lyxitol

[130693-66-2]

[187144-40-7]

Cryst. (MeOH aq.) (as hydrochloride). Mp
 195.5-196°. $[\alpha]_D^{20}$ +22.7 (c, 0.8 in MeOH).

N-Me: 1-Methyl-3,4,5-piperidinetriol,
 9CI. 3,4,5-Trihydroxy-N-methylpiperidine

[260354-31-2]

C₆H₁₃NO₃ 147.174

Constit. of the leaves of *Eugenia uniflora* (nangapiry). Viscous oil. $[\alpha]_D$
 +19.1 (c, 1.6 in H₂O).

[\pm -(3 α ,4 α ,5 β)]-form(±)-*cis,trans*-form

[130114-77-1]

Oil.

Tetra-Ac: [130271-87-3]

C₁₃H₁₉NO₇ 301.296

Cryst. (EtOAc/diisopropyl ether). Mp
 102-103°.

N-Me: [155549-47-6]

C₆H₁₃NO₃ 147.174

Oil.

(3 α ,4 β ,5 α)-form

trans,trans-form. 1,5-Dideoxy-1,5-iminoxytilol

[13042-55-2]

[4451-19-8] β -Glucosidase inhibitor.

Cryst. (MeOH/EtOAc/H₂O). Mp 126.5-127.5°. *meso*-.

Paulsen, H. *et al.*, *Chem. Ber.*, 1967, **100**, 2467-2473 (3 α ,4 β ,5 α -form)

Bernotas, R.C. *et al.*, *Tet. Lett.*, 1990, **31**, 3393-3396 (*synth*, *pmr*, *cmr*)

Tschamber, T. *et al.*, *Tetrahedron*, 1994, **50**, 1135-1152 (3 α ,4 α ,5 β -form, *synth*, *ir*, *pmr*, *tetra-Ac*, *N-Me*)

Legler, G. *et al.*, *Carbohydr. Res.*, 1995, **272**, 17-30 (3R-(3 α ,4 α ,5 β)-form, *synth*, *pmr*, *cmr*)

Godskesen, M. *et al.*, *Bioorg. Med. Chem.*, 1996, **4**, 1857-1865 (*synth*, *pmr*, *cmr*, *biochem*)

Igarashi, Y. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 553-558 (3 α ,4 α ,5 α -form, *synth*, *pmr*, *cmr*, *biochem*)


Kim, D.K. *et al.*, *J.C.S. Perkin 1*, 1996, 803 (3 α ,4 α ,5 α -form, *synth*, *pmr*, *cmr*)

Matsumura, T. *et al.*, *Pharm. Biol.*, 2000, **38**, 302-307 (N-Me)

Reddy, M.S. *et al.*, *Tetrahedron*, 2007, **63**, 291-298 (3R-(3 α ,4 α ,5 β)-form, *synth*, *pmr*, *cmr*)

4-Piperidinone, 9CI γ -Piperidone

[41661-47-6]

C₅H₉NO 99.132

Alkaloid from leaves and twigs of *Dichilus strictus*, *Dichilus reflexus*, *Dichilus lebeckioides*, *Dichilus pilosus* and *Dichilus gracilis* (Fabaceae). Yellow oil. pK_a 8.6 (25°, H₂O). Cannot be distilled.

Covalent hydrate: 4,4-Piperidinediol

[40064-34-4]

C₅H₁₁NO₂ 117.147

Mp 94-96° (as hydrochloride).

Hydrochloride: [41979-39-9]

Cryst. + 1/2 EtOH (EtOH/Et₂O) or
 cryst. + H₂O. Mp 147-149° (anhyd.).

[73390-11-1]


McElvain, S.M. *et al.*, *J.A.C.S.*, 1949, **71**, 901 (*props*)

Dickerman, S.C. *et al.*, *J.O.C.*, 1949, **14**, 530 (*synth*)

Van Wyk, B.-E. *et al.*, *Biochem. Syst. Ecol.*, 1988, **16**, 471 (*occur*)

3-(2-Piperidinyl)pyridine, P-460**9CI**2-(3-Pyridinyl)piperidine. *Anabasine*,*JMAF*. *Nicotimine*. *Neonicotine*

[40774-73-0]



(S)-form

C₁₀H₁₄N₂ 162.234

The name *neonicotine* refers primarily to the racemate.

(R)-form [34366-21-7]Isol. from *Malacocarpus crithmifolius*.

Oil. Bp₁₅ 145-150°. $[\alpha]_D^{20}$ +10.1 (neat). n_D²⁰
 1.5412. Clearly of low opt. purity.

(S)-form [494-52-0]

Alkaloid from *Anabasis aphylla*, *Nicotiana tabacum*, many other *Nicotiana* spp. and many other spp. in several families. Also in marine ribbon worms and a component of the venom of ants *Aphaenogaster* spp. and *Messor* spp. Insecticide. Now superseded. Sol. H₂O. Mp 25-30°. Bp 276° Bp₂ 104-105°. $[\alpha]_D^{24}$ -79.2 (c, 0.5 in MeOH) (synthetic). $[\alpha]_D^{25}$ -41.5 (c, 1.0 in C₆H₆). pK_{a1} 11; pK_{a2} 3.21 (30°).

► Exp. reprod. effects and teratogen. LD₅₀ (gpg, scu) 22 mg/kg. BV4375000

Dipicrate: Mp 205°.

N-Ac: [3350-86-5]

C₁₂H₁₆N₂O 204.271

Used as 0.5M aq. soln. for extraction-photometric detn. of Ti. Cryst. Sol. EtOH; sl. sol. H₂O.

N-Me: N-Methylanabasine

[24380-92-5]

C₁₁H₁₆N₂ 176.261

Occurs in traces in crude nicotine and in *Anabasis aphylla*. Oil. Bp₁₂ 127-128°. $[\alpha]_D^{15}$ -85.1.

N-Me, picrate:

Cryst. (EtOH). Mp 287-288° dec.

1,2,3,4-Tetrahydro: Tetrahydroanabasine

[64191-30-6]

C₁₀H₁₈N₂ 166.266

Alkaloid from *Adenocarpus complicatus* ssp. *aureus* (Fabaceae). Important biosynth. intermed. for Nitraria and lupine alkaloids.

(±)-form [13078-04-1]

Alkaloid from *Duboisia myoporoides*, *Nicotiana glauca* and other spp. (Solana-ceae). Bp₇₇₅ 280-282° Bp 106-108°.

Dipicrate: Mp 213-214°.

N-Me: [2055-12-1]

Bp_{0.4} 62°.

N-Me, dipicrate:

Yellow needles (EtOH). Mp 237-239°.

[15251-47-5]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 295A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1537D (ir)

Orechhoff, A. *et al.*, *Ber.*, 1931, **64**, 266-274; 1932, **65**, 232-234 (*isol*)

Späth, E. *et al.*, *Ber.*, 1937, **70**, 70-72 (*resoln*)

Otroshchenko, O.S. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1954, **24**, 1847-1849; *Zh. Obshch. Khim.*, 1954, **24**, 1884-1886 (N-Ac, *synth*)

Linnell, R.H. *et al.*, *J.A.C.S.*, 1954, 1391-1393 (*uv*)

Lukeš, R. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 751-756 (*config*)

Talipov, S.T. *et al.*, *Zh. Anal. Khim.*, 1963, **18**, 178-181; *J. Anal. Chem. USSR (Engl. Transl.)*, 1963, **18**, 161-164 (N-Ac, *use*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 2926-2932 (*ms*)

Zharekeev, V.Kh. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 538-539; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 524 (R-form, *isol*)

Aslanov, Kh.A. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 324-328; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 319-321 (*isol*, *manuf*)

- Testa, B. *et al.*, *Mol. Pharmacol.*, 1973, **9**, 10-16 (*cd*)
- Fitch, W.L. *et al.*, *J.O.C.*, 1974, **39**, 2974-2979 (*S*-form, *isol*)
- Leete, E. *et al.*, *Chem. Comm.*, 1975, 9-10 (*biosynth*)
- Nehme, M. *et al.*, *An. Quim.*, 1977, **73**, 307-308; *CA*, **87**, 114689h (*Tetrahydroanabasine*)
- Alberici, G.F. *et al.*, *Tet. Lett.*, 1983, **24**, 1937-1940 (*synth*)
- Strunz, G.M. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 89-183 (*rev*)
- Pfrengele, W. *et al.*, *J.O.C.*, 1989, **54**, 4261-4263 (*S*-form, *synth*)
- Pesticide Manual*, 9th edn., 1991, No. 380 (*Anabasine*)
- Hattori, K. *et al.*, *Tetrahedron*, 1993, **49**, 1749-1760 (*S*-form, *synth*)
- Wanner, M.J. *et al.*, *J.O.C.*, 1996, **61**, 5581-5586 (*Tetrahydroanabasine*, *biosynth*, *bibl*)
- Wanner, M.J. *et al.*, *Pure Appl. Chem.*, 1996, **68**, 2051-2056 (*Tetrahydroanabasine*)
- Deo, N.M. *et al.*, *Tet. Lett.*, 1996, **37**, 1137-1140 (*synth*)
- Yang, C.-M. *et al.*, *Can. J. Chem.*, 1997, **75**, 616-620 (*synth*, *ir*, *pmr*, *cmr*, *ms*)
- Leclercq, S. *et al.*, *J. Chem. Ecol.*, 2001, **27**, 945-952 (*isol*, *abs config*)
- Felpin, F.-X. *et al.*, *J.O.C.*, 2001, **66**, 6305-6312 (*synth*)
- Castro, A. *et al.*, *Heterocycles*, 2007, **71**, 2699-2708 (*synth*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, AON875


Piperlongine P-461

Struct. unknown. Constit. of *Piper longum* roots (long pepper) (Piperaceae). Oil. Bp₃ 110-115° (lit. gives a pressure range). Not characterised. May not be an alkaloid.

Chatterjee, A. *et al.*, *Tetrahedron*, 1967, **23**, 1769-1781 (*isol*)

Piperlonguminine P-462

5-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,4-pentadienamide, 9CI. 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid isobutylamide



(2E,4E)-form

C₁₆H₁₉NO₃ 273.331

(2E,4E)-form [5950-12-9]

Alkaloid from *Ottonia corcovadensis* roots (preferred genus name *Piper*), *Piper longum*, *Piper nepalense*, *Piper sylvaticum* and other *Piper* spp. (Piperaceae). Cryst. (C₆H₆/petrol). Mp 167-169°.

2,3-Dihydro: 5-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-4-pentenamide, 9CI. N-Isobutyl-5-(3,4-methylenedioxyphenyl)-4-pentenamide. **Isodihydropiperlonguminine**

C₁₆H₂₁NO₃ 275.347

Alkaloid from the fruits of *Piper longum* (long pepper). Yellow solid. Mp 78°.

4,5-Dihydro: 5-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2-pentenamide, 9CI. N-Isobutyl-5-(3,4-methylenedioxyphenyl)-2-pentenamide, 8CI. **4,5-Dihydropiperlonguminine**. A^{α,β}.

Dihydropiperlonguminine

[23512-53-0]

C₁₆H₂₁NO₃ 275.347

Alkaloid from fruits of *Piper guineense* and *Piper longum* (long pepper). Also *isol.* from *Fagara macrophylla*. Insect growth inhibitor. Cryst. (C₆H₆ or C₆H₆/petrol). Mp 90-94° (natural) Mp 110° (synthetic).

2''-Acetoxy: Corcovadine

[79097-20-4]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the roots of *Ottonia corcovadensis* (preferred genus name *Piper*). Mp 141-144°. λ_{max} 244 (log ε 4.19); 252 (log ε 4.19); 307 (log ε 4.4); 342 (log ε 4.65) (MeOH).

6'-Methoxy: 5-(6-Methoxy-3,4-methylenedioxyphenyl)-2,4-pentadienoic acid isobutylamide. **6-Methoxypiperoyliso-butylamine**

[102934-25-8]

C₁₇H₂₁NO₄ 303.357

Alkaloid from *Piper amalago*.

(2E,4Z)-form

Scutifoliamide A

[1000390-09-9]

Alkaloid from the leaves of *Piper scutifolium*. Amorph. solid. λ_{max} 241 (log ε 3.2); 293 (log ε 3); 308 (log ε 3); 337 (log ε 3) (CHCl₃).

2''-Acetoxy: Scutifoliamide B

[1000390-10-2]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the leaves of *Piper scutifolium*. Amorph. solid. λ_{max} 241 (log ε 3.7); 296 (log ε 3.5); 310 (log ε 3.5); 342 (log ε 3.6) (CHCl₃).

(2Z,4E)-form

Hoffmannseggiamide A

[1000390-08-8]

Alkaloid from the leaves of *Piper hoffmannseggianum*. Amorph. solid.

2''-Acetoxy: Hoffmannseggiamide B

[1000390-11-3]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the leaves of *Piper hoffmannseggianum*. Amorph. solid.

(2Z,4Z)-form

Isopiperlonguminine

[79097-19-1]

Alkaloid from the roots of *Ottonia corcovadensis* (Piperaceae). Mp 140-143°.

2''-Acetoxy: Isocorcovadine

[79097-21-5]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the roots of *Ottonia corcovadensis*. Mp 80-85°. λ_{max} 245 (log ε 4.39); 254 (log ε 4.41); 310 (log ε 4.38); 327 (log ε 4.48) (MeOH).

Chatterjee, A. *et al.*, *Tetrahedron*, 1967, **23**, 1769 (*isol*, *uv*, *pmr*, *ms*, *synth*)

Loder, J.W. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1531 (*synth*, 4,5-Dihydropiperlonguminine)

Gupta, O.P. *et al.*, *Phytochemistry*, 1972, **11**, 2646 (*isol*)

Mahanta, P.K. *et al.*, *J. Pharm. Sci.*, 1974, **63**, 1160 (*uv*, *ir*, *pmr*)

Okogun, J.I. *et al.*, *J.C.S. Perkin 1*, 1974, 2195 (*isol*, *uv*, *ir*, *pmr*, *ms*, *synth*)

Patra, A. *et al.*, *Phytochemistry*, 1974, **13**, 2889 (*isol*)

Dwuma-Badu, D. *et al.*, *Phytochemistry*, 1976, **15**, 822 (4,5-Dihydropiperlonguminine)

Olsen, R.A. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 942 (*synth*, *uv*, *ir*, *pmr*)

Soares Costa, S. *et al.*, *Phytochemistry*, 1981, **20**, 1305-1307 (*Piperlonguminine*, *Isopiperlonguminine*, *Corcovadine*, *Isocorcovadine*)

Kubo, I. *et al.*, *Experientia*, 1984, **40**, 340 (4,5-Dihydropiperlonguminine)

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (6'-methoxy)

Kaga, H. *et al.*, *Synlett*, 1994, 607 (*synth*)


Anuradha, V. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 245-251 (*Isodihydropiperlonguminine*)

Min, K.R. *et al.*, *Planta Med.*, 2004, **70**, 1115-1118 (*isol*, *pmr*, *cmr*)

Marques, J.V. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2036-2039 (*Scutifoliamides A-B*, *Hoffmannseggiamides A-B*)

Pipermethystine P-463

5-Acetoxy-5,6-dihydro-1-(3-phenylprop-1-enyl)-2-(1H)-pyridinone [71627-22-0]



C₁₆H₁₇NO₄ 287.315

Alkaloid from the leaves of *Piper methysticum* (kava). Oil. [α]_D²³ -176.4 (c, 0.5 in Me₂CO).

3,4-Epoxy: 3,4-Epoxy pipermethystine

C₁₆H₁₇NO₅ 303.314

Alkaloid from the aerial parts of *Piper methysticum*. Needles (hexane). Mp 59°. [α]_D²² -98.8 (c, 0.5 in Me₂CO). Possesses 3α,4α,5β-config.

Deacetoxy: 5,6-Dihydro-1-(3-phenylprop-1-enyl)-2-(1H)-pyridinone. **Piperchabamide A**

C₁₄H₁₅NO₂ 229.278

Alkaloid from the fruit of *Piper chaba* (Javanese long pepper). Oil. λ_{max} 243 (log ε 3.72) (EtOH).

Smith, R.M. *et al.*, *Tetrahedron*, 1979, **35**, 437-439 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)


Gomez, R. *et al.*, *Org. Lett.*, 2001, **3**, 3381-3383 (*synth*)

Dragull, K. *et al.*, *Phytochemistry*, 2003, **63**, 193-198 (3,4-Epoxy pipermethystine)

Morikawa, T. *et al.*, *Planta Med.*, 2004, **70**, 152-159 (*Piperchabamide A*)

Pipernonaline P-464

1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,8-nonadienyl]piperidine, 9CI. 9-(3,4-Methylenedioxyphenyl)-2,8-nonadienoic acid piperidide [88660-10-0]



C₂₁H₂₇NO₃ 341.449

Alkaloid from the fruits of *Piper longum* (long pepper). Cryst. (EtOAc/hexane). Mp 54-55.5°.

2,3-Dihydro- Piperolein B. 9-(3,4-Methylenedioxyphenyl)-8-nonenic acid piperidide
[30505-89-6]
C₂₁H₂₉NO₃ 343.465
Alkaloid from *Piper nigrum* (black pepper). Oil. λ_{\max} 214 (ϵ 28000); 260 (ϵ 16600); 268 (ϵ 16100); 305 (ϵ 10800) (MeOH) (Derep).


4,5-Didehydro: 9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienoic acid piperidide. **Dehydropiperonaline**
[107584-38-3]
C₂₁H₂₅NO₃ 339.433
Alkaloid from the fruits of *Piper longum* (long pepper). Possesses coronary vasodilating activity. Plates (hexane/Me₂CO). Mp 98-99°.

2,3-Dihydro, 4,5,6,7-tetrahydro (E,E)-: 9-(3,4-Methylenedioxyphenyl)-4,6,8-nonatrienoic acid piperidide. **Piptigrine**
C₂₁H₂₅NO₃ 339.433
Alkaloid from the seeds of *Piper nigrum* (black pepper). Insecticidal agent. Dark yellow rods (CHCl₃/MeOH). Mp 159-161°. λ_{\max} 260 (ϵ 10100); 313 (ϵ 19500); 331 (ϵ 35200); 363 (ϵ 42500) (MeOH).

Grewe, R. et al., *Chem. Ber.*, 1970, **103**, 3752 (*isol, uv, ir, pmr, ms, synth, Piperolein B*)
Tabuneng, W. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3562 (*isol, pmr, cmr, ir, uv, ms, struct*)
Shoji, N. et al., *J. Pharm. Sci.*, 1986, **75**, 1188-1189 (*Dehydropiperonaline*)
Kaga, H. et al., *Synlett*, 1994, 607 (*Piperonaline, Dehydropiperonaline, synth*)
Strunz, G.M. et al., *Can. J. Chem.*, 1996, **74**, 419 (*Dehydropiperonaline, synth*)
Yang, Y.-C. et al., *J. Agric. Food Chem.*, 2002, **50**, 3765-3767 (*isol, pmr, cmr*)
Siddiqui, B.S. et al., *Nat. Prod. Res.*, 2004, **18**, 473-477 (*Piptigrine*)

Piperolein A P-465

1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-6-heptenyl]piperidine, 9CI. 1-[7-(3,4-Methylenedioxyphenyl)-6-heptenyl]piperidine



C₁₉H₂₅NO₃ 315.411
 λ_{\max} 214 (ϵ 28000); 260 (ϵ 16600); 268 (ϵ 16100); 305 (ϵ 10800) (MeOH) (Derep).

(E)-form [30505-92-1]

Minor constit. of *Piper nigrum* (pepper) (Piperaceae). Pale yellow oil.

2,3-Didehydro, 6,7-dihydro: Piperine S. 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2-heptenyl]piperidine. 1-[7-(3,4-Methylenedioxyphenyl)-2-heptenyl]piperidine
[156970-11-5]
C₁₉H₂₅NO₃ 315.411


Isol. from stems and leaves of *Piper puberulum* (Piperaceae). Yellow nee-

dles. Mp 52-54°. λ_{\max} 202 ; 232 (sh) ; 280 (MeOH).

Grewe, R. et al., *Chem. Ber.*, 1970, **103**, 3752 (*isol, synth, struct*)
Wu, Q. L. et al., *Phytochemistry*, 1997, **44**, 727; *CA*, **121**, 104122z (*Piperine S*)
Chaves, M.C. de O. et al., *Fitoterapia*, 2003, **74**, 181-183 (*Piperine S*)

Piperovatine

6-(4-Methoxyphenyl)-N-(2-methylpropyl)-2,4-hexadienamide, 9CI. N-Isobutyl-6-p-methoxyphenylsorbamide. 6-(4-Methoxyphenyl)-2,4-hexadienoic acid isobutylamide



C₁₇H₂₃NO₂ 273.374

(E,E)-form [25090-18-0]

Alkaloid from *Ottonia vahlii* (preferred genus name *Piper*) and *Ottonia corcovadensis* (Piperaceae). Cryst. (Et₂O). Mp 120-121° (115-119°).

Demethoxy: [99083-23-5]

C₁₆H₂₁NO 243.348

Isol. from *Dinosperma erythrocca*. Shows activity against house fly (*Musca domestica*). Needles. Mp 116°. λ_{\max} 258 (MeOH).

2'-Hydroxy, demethoxy: [120262-71-7]

C₁₆H₂₁NO₂ 259.347

Isol. from *Dinosperma erythrocca*. Shows lethal activity against tobacco budworm (*Heliothis virescens*). Yellow waxy solid. Mp 77°. λ_{\max} 257 (MeOH).

Price, S.J. et al., *J.O.C.*, 1970, **35**, 2568-2571 (*isol, uv, ir, pmr, ms, synth*)

Vig, B. et al., *J. Indian Chem. Soc.*, 1974, **51**, 817-818 (*synth, uv, ir, pmr*)


Soares Costa, S. et al., *Phytochemistry*, 1981, **20**, 1305-1307 (*isol, uv, ir, pmr, ms*)

Latif, Z. et al., *J. Nat. Prod.*, 1998, **61**, 614-619 (*demethoxy derivs*)

Rendon, W. et al., *Spectroscopy (Amsterdam)*, 1998, **14**, 35-40 (*pmr, cmr*)

1-Piperoylpiperidine

1-[5-(1,3-Benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine, 9CI. 1-[5-(3,4-Methylenedioxyphenyl)-2,4-pentadienyl]piperidine. 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid piperidide



C₁₇H₁₉NO₃ 285.342

(2E,4E)-form

Piperine. FEMA 2909
[94-62-2]

Constit. of pepper (*Piper nigrum*) and many other *Piper* spp. (Piperaceae). Responsible for the hot taste of pepper. Flavouring agent, insecticide. Analeptic,

bactericidal agent. Sol. EtOH; poorly sol. H₂O. Mp 129°. pK_a 1.98 (15°). Log P 2.7 (calc).

▶LD₅₀ (rat, orl) 514 mg/kg. Exp. reprod. and teratogenic effects. TN2321500

4,5-Dihydro: 1-[5-(1,3-Benzodioxol-5-yl)-1-oxo-2-pentenyl]piperidine. 5-(3,4-Methylenedioxyphenyl)-2-pentenoic acid piperidide. **Piperanine.** β -Hydropiperylpiperidine. $\Delta^{\alpha,\beta}$ -Dihydro-piperine
[23512-46-1]
C₁₇H₂₁NO₃ 287.358

Alkaloid from *Piper nigrum*, *Piper guineense* and *Piper novae-hollandiae* (Piperaceae). Cryst. (EtOAc or petrol). Mp 79-80°. λ_{\max} 232 (ϵ 11800); 285 (ϵ 4350) (MeOH).

2,3,4,5-Tetrahydro: 1-[5-(3,4-Methylenedioxyphenyl)pentanoyl]piperidine. **Tetrahydropiperine**

C₁₇H₂₃NO₃ 289.374

Alkaloid from the fruit of *Piper longum* (long pepper).

(2E,4Z)-form

Isochavicine

[30511-77-4]

Constit. of pepper (*Piper nigrum*) (Piperaceae). Cryst. (CHCl₃/hexane). Mp 110° (103°).

(2Z,4E)-form

Isopiperine

[30511-76-3]

Constit. of pepper (*Piper nigrum*) (Piperaceae). Cryst. (CHCl₃/hexane). Mp 89° (86°).

(2Z,4Z)-form

Chavicine

[495-91-0]

Constit. of pepper (*Piper nigrum*) (Piperaceae). Oil. Bp_{0.0001} 110°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 362C (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1379B (*nmr*)

Rügheimer, L. et al., *Ber.*, 1882, **15**, 1390 (*synth, Piperine*)

Loder, J.W. et al., *Aust. J. Chem.*, 1969, **22**, 1531 (*Piperanine*)

Grewe, R. et al., *Chem. Ber.*, 1970, **103**, 3752 (*isol, uv, ir, pmr, synth*)

Traxler, J.T. et al., *J. Agric. Food Chem.*, 1971, **19**, 1135 (*Piperanine*)

Wenkert, E. et al., *J.A.C.S.*, 1971, **93**, 6271 (*pmr, cmr, Piperine*)

Grynpas, M. et al., *Acta Cryst. B*, 1975, **31**, 2663 (*cryst struct, Piperine*)

DeCleyne, R. et al., *Chromatographia*, 1975, **8**, 342 (*isol, uv*)

Addae-Mensah, I. et al., *Phytochemistry*, 1977, **16**, 483 (*Piperanine*)

Tsuboi, S. et al., *Tet. Lett.*, 1979, 1043 (*synth, Piperine, Isochavicine*)

Schulze, A. et al., *Annalen*, 1981, 1725 (*synth*)

Kaga, H. et al., *Synlett*, 1994, 607 (*synth, Piperine*)

De Araujo-Junior, J.X. et al., *Phytochemistry*, 1997, **44**, 559 (*pmr, cmr*)


Madhusudhan, P. et al., *Biochem. Syst. Ecol.*, 2001, **29**, 537-538

(*Tetrahydropiperine*)


Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, PIV600

Piperpense


[193225-26-2]

C₁₆H₁₇NO₄ 287.315Alkaloid from *Piper ponapense* (Piperaceae). Cryst. (EtOAc/hexane).Patil, A.D. et al., *Nat. Prod. Lett.*, 1997, **9**, 217-223 (isol, uv, pmr, cmr, cryst struct)**Piperstachine****P-469**


11-(1,3-Benzodioxol-5-yl)-N-(2-methylpropyl)-2,8,10-undecatrienamide, 9CI. 11-(3,4-Methylenedioxyphenyl)-2,8,10-undecatrienoic acid isobutylamide

C₂₂H₂₉NO₃ 355.476**(E,E,E)-form** [57155-80-3]Alkaloid from the stems of *Piper trichostachyon* (Piperaceae). Cryst. (CH₂Cl₂/hexane). Mp 153°. λ_{max} 216 (ε 34700); 283 (ε 21900); 293 (ε 24500); 315 (ε 18200) (EtOH).8,9-Dihydro: 11-(3,4-Methylenedioxyphenyl)-2,10-undecadienoic acid isobutylamide. **Piperchabamide D**C₂₂H₃₁NO₃ 357.492Alkaloid from the fruit of *Piper chaba* (Javanese long pepper). Oil. λ_{max} 261 (log ε 4.12) (EtOH).Joshi, B.S. et al., *Helv. Chim. Acta*, 1975, **58**, 1551 (isol, uv, ir, pmr, cmr, ms, struct)
Viswanathan, N. et al., *Helv. Chim. Acta*, 1975, **58**, 2026 (synth, uv, ir, ms)
Vig, O.P. et al., *Indian J. Chem., Sect. B*, 1980, **19**, 276 (synth, uv, ir, pmr)
Morikawa, T. et al., *Planta Med.*, 2004, **70**, 152-159 (Piperchabamide D)**Pipertipine****P-470**

1-[8-(1,3-Benzodioxol-5-yl)-1-oxo-7-octadecenyl]piperidine, 9CI. 1-[8-(3,4-Methylenedioxyphenyl)-7-octadecenyl]piperidine [485794-76-1]

C₂₀H₂₇NO₃ 329.438Alkaloid from the dried seeds of pepper *Piper nigrum*. Yellow wax. λ_{max} 263 (ε 13400); 270 (ε 12800); 310 (ε 5850); 322 (sh) (MeOH).Siddiqui, B.S. et al., *Heterocycles*, 2002, **57**, 1653-1658 (isol, pmr, cmr, ms)**Piperundecalidine****P-471**



1-[11-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,10-undecatrienyl]piperidine, 9CI. 11-(3,4-Methylenedioxyphenyl)-2,4,10-undecatrienoic acid piperidide [88660-11-1]

C₂₃H₂₉NO₃ 367.487Alkaloid from the fruits of *Piper longum* (long pepper) (Piperaceae). Cryst. (EtOAc/hexane). Mp 64.5-65.5°.

4,5-Dihydro: 11-(3,4-Methylenedioxyphenyl)-2,10-undecadienoic acid piperidide.

Piperchabamide BC₂₃H₃₁NO₃ 369.503Alkaloid from the fruit of *Piper chaba* (Javanese long pepper). Oil. λ_{max} 260 (log ε 4.24) (EtOH).Tabuneng, W. et al., *Chem. Pharm. Bull.*, 1983, **31**, 3562 (isol, uv, ir, pmr, cmr, ms, struct)
Morikawa, T. et al., *Planta Med.*, 2004, **70**, 152-159 (Piperchabamide B)**Pipgularine****P-472**

[622405-48-5]

C₂₃H₃₃NO₃ 371.519Alkaloid from the dried seeds of *Piper nigrum* (pepper). Yellow needles (CHCl₃/MeOH). Mp 98-100°. λ_{max} 263 (ε 5450); 313 (ε 14200) (MeOH).Siddiqui, B.S. et al., *Helv. Chim. Acta*, 2003, **86**, 2760-2767 (isol, pmr, cmr, ms)**Piplartine****P-473**5,6-Dihydro-1-[1-oxo-3-(3,4,5-trimethoxyphenyl)-2-propenyl]-2(1H)-pyridinone, 9CI. N-(3,4,5-Trimethoxycinnamoyl)-Δ³-piperidin-2-one. 5,6-Dihydro-1-(3,4,5-trimethoxycinnamoyl)-2(1H)-pyridinone. **Piperlongumine** [20069-09-4]C₁₇H₁₉NO₅ 317.341Alkaloid from the roots of *Piper longum* (long pepper) and *Piper sylvaticum* (Piperaceae). Also isol. from leaves of *Piper tuberculatum*. Effective drug for treatment of asthma and chronic bronchitis. Needles (EtOH/Et₂O). Mp 124°. Iso-meric Δ⁵ struct. originally proposed. λ_{max} 222; 240; 325 (MeOH) (Berdy). λ_{max} 225 (ε 11000); 324 (ε 10000) (EtOH) (Berdy).**4'-O-De-Me: 4'-O-Demethylpiplartine**C₁₆H₁₇NO₅ 303.314Alkaloid from *Piper cenocladum*. Yellow oil. λ_{max} 220 (sh) (log ε 4.74); 270 (log ε 4.62); 345 (log ε 4.5) (MeOH).**8,9-Dihydro: 8,9-Dihydropiplartine**

[137760-69-1]

C₁₇H₂₁NO₅ 319.357Alkaloid from the aerial parts of *Piper rugosum* (Piperaceae). Semi-solid.**8,9-Dihydro, 3,4-epoxide: 3,4-Epoxy-8,9-dihydropiplartine**

[174544-87-7]

C₁₇H₂₁NO₆ 335.356Alkaloid from leaves and twigs of *Piper verrucosum*. Cryst. (Me₂CO/petrol). Mp 85-87°. [α]_D²⁰ -121.6 (c, 0.020 in CHCl₃).**Tetrahydro:**Prisms (Et₂O). Mp 83-84°.**3'-Demethoxy: N-(3,4-Dimethoxycinnamoyl)-Δ³-piperidin-2-one. 3'-Demethoxypiplartine**

[130263-10-4]

C₁₆H₁₇NO₄ 287.315Alkaloid from the stems of *Piper arboreoscens* (Piperaceae) and leaves of *Piper tuberculatum*. Displays significant activity as a repellent to the leafcutter ant *Atta cephalotes*. Pale yellow needles (MeOH). Mp 116-117°. λ_{max} 222 (ε 45710); 338 (ε 53700) (MeOH) (Berdy).**3'-Demethoxy, 8,9-dihydro: 1-[3-(3,4-Dimethoxyphenyl)propanoyl]-5,6-dihydro-2(1H)-pyridinone. 8,9-Dihydro-3'-demethoxypiplartine. Sintenpyridone**C₁₆H₁₉NO₄ 289.33Alkaloid from the whole plant of *Piper sintenense*. Cytotoxic. Needles (CHCl₃/MeOH). Mp 125-127°. λ_{max} 220 (log ε 4.68); 276 (log ε 4.03) (EtOH).**3'-Demethoxy, 8,9-dihydro, 3,4-epoxide: Piplaroxide**


[179382-48-0]

C₁₆H₁₉NO₅ 305.33Alkaloid from leaves of *Piper tuberculatum*. Displays significant activity as a repellent to the leafcutter ant *Atta cephalotes*. Small needles (MeOH/CHCl₃). Mp 91-92°. [α]_D²⁵ +67.7 (c, 0.8 in CHCl₃).**(Z)-Isomer: cis-Piplartine**C₁₇H₁₉NO₅ 317.341Alkaloid from *Piper tuberculatum*. Amorph. solid. λ_{max} 332 (CHCl₃).Chatterjee, A. et al., *Tetrahedron*, 1967, **23**, 1769 (isol, uv, ir, pmr)Joshi, B.S. et al., *Tet. Lett.*, 1968, 2395 (pmr, struct)Banerji, J. et al., *Phytochemistry*, 1974, **13**, 2327 (isol)Banerjee, T. et al., *Can. J. Chem.*, 1986, **64**, 876 (cryst struct)Duh, C.-Y. et al., *Phytochemistry*, 1990, **29**, 2689 (3'-Demethoxypiplartine)Maxwell, A. et al., *J. Nat. Prod.*, 1991, **54**, 1150 (8,9-Dihydropiplartine)Seeram, N.P. et al., *J. Nat. Prod.*, 1996, **59**, 436 (3,4-Epoxy-8,9-dihydropiplartine)

- Capron, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 794 (*Piplaroxide*)
 Dodson, C.D. *et al.*, *Phytochemistry*, 2000, **53**, 51-54 (*4'-O-Demethylpiplartine*)
 Navickiene, H.M.D. *et al.*, *Phytochemistry*, 2000, **55**, 621-626 (*cis-Piplartine*)
 Chen, J.-J. *et al.*, *Planta Med.*, 2002, **68**, 980-985 (*Sintenpyridone*)

Piplartine dimer A P-474

1,1'-[2,4-Bis(3,4,5-trimethoxyphenyl)-1,3-cyclobutanediyl]dicarbonyl[bis[5,6-dihydro-2-(1H)-pyridone]] [80248-71-1]



$C_{34}H_{38}N_2O_{10}$ 634.682
 Alkaloid from *Piper arborescens*, *Piper rugosum* and *Piper tuberculatum*. Cytotoxic. Cryst. (MeOH). Mp 269-270° (262-264°). λ_{max} 214 (log ϵ 4.53); 240 (sh) (log ϵ 4.25) (MeOH).

3'-Demethoxy: Piperarborescine B

$C_{33}H_{36}N_2O_9$ 604.655
 Alkaloid from the stems of *Piper arborescens*. Needles (MeOH). Mp 158-162°. λ_{max} 214 (log ϵ 4.52); 240 (sh) (log ϵ 4.31); 280 (sh) (log ϵ 3.67) (MeOH).


3',3''-Bis(demethoxy): Piperarborescine A

$C_{32}H_{34}N_2O_8$ 574.629
 Alkaloid from the stems of *Piper arborescens*. Needles (MeOH). Mp 220-222°. $[\alpha]_D^{25} +5.5$ (c, 0.05 in $CHCl_3$). λ_{max} 212 (log ϵ 4.53); 240 (sh) (log ϵ 4.34); 280 (sh) (log ϵ 3.92) (MeOH).

1,2-Diepimer, 3'-demethoxy: Piperarborescine D


$C_{33}H_{36}N_2O_9$ 604.655
 Alkaloid from the stems of *Piper arborescens*. Brownish oil. λ_{max} 217 (log ϵ 4.55); 240 (sh) (log ϵ 4.33); 280 (sh) (log ϵ 3.73) (MeOH).

- Filho, R.B. *et al.*, *Phytochemistry*, 1981, **20**, 345-346 (*isol, synth, ir, pmr, cmr, ms*)
 Maxwell, A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1150-1152 (*isol*)
 Lee, F.-P. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 463-468 (*Piperarborescines A-B*)
 Tsai, I.-L. *et al.*, *Planta Med.*, 2005, **71**, 535-542 (*Piperarborescine D*)

Pipsaecedine P-475

$C_{34}H_{38}N_2O_6$ 570.684
 Alkaloid from the fruit of *Piper nigrum* (pepper). Amorph. powder. λ_{max} 254 (ϵ 11900); 264 (ϵ 13000); 301 (ϵ 25000); 314 (ϵ 24200); 321 (ϵ 35000); 357 (ϵ 41000); 373 (ϵ 45200) (MeOH).

Siddiqui, B.S. *et al.*, *Helv. Chim. Acta*, 2004, **87**, 660-666 (*isol, pmr, cmr, ms*)

Piptanthine P-476

(-)-form

$C_{20}H_{35}N_3$ 317.517
 Stereoisomeric with Ormosanine, O-115, Dasycarpine, D-85 and Templettine, T-68.


(-)-form [7344-67-4]
 Alkaloid from *Piptanthus nanus* (Fabaceae). Mp 142-146° (136-137°). $[\alpha]_D -24$.

N,N'-Di-Ac:
 Cryst. by subl. Mp 218-221°.
 N-Me: Mp 114-115°.

(+)-form
 Obt. by reduct. of Panamine, P-53. Mp 136-137°. $[\alpha]_D +23$.

(±)-form
 Minor alkaloid from leaves and stems of *Hovea linearis* and from *Templetonia retusa*, *Ormosia semicastrata* and *Amopiptanthus mongolicus* (Fabaceae). Cryst. (MeOH). Mp 220-222°.

- Konovalova, R.A. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1951, **78**, 705; **81**, 1069 (*isol*)
 Eisner, U. *et al.*, *Coll. Czech. Chem. Comm.*, 1959, **24**, 2348-2355 (*purifn*)
 Deslongchamps, P. *et al.*, *Tet. Lett.*, 1964, 3893-3898 (*struct, abs config*)
 McLean, S. *et al.*, *Can. J. Chem.*, 1971, **49**, 1976-1978; 1972, **50**, 1639-1641 (*isol*)
 Cannon, J.R. *et al.*, *Tet. Lett.*, 1974, 1683-1686 (*isol, abs config*)
 Liu, H.-J. *et al.*, *Can. J. Chem.*, 1976, **54**, 97-109 (*synth*)
 Lamberton, J.A. *et al.*, *Aust. J. Chem.*, 1982, **35**, 2577-2582 (*isol*)
 Le, P.M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 283-293 (*pmr, cmr*)

Pipyahyine P-477

$C_{24}H_{33}NO_3$ 383.53
 Alkaloid from the dried fruits of *Piper nigrum* (pepper). Needles (petrol/EtOAc). Mp 109-110.5°. λ_{max} 265 (ϵ 38500); 315 (ϵ 3500) (MeOH).

Siddiqui, B.S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1349-1352 (*isol, pmr, cmr, ms*)

Piricularin P-478


Toxin B. *Pyricularin* [56283-47-7]
 $C_{18}H_{30}N_2O_5$ 354.445
 Phenazine antibiotic. Struct. unknown. Isol. from *Piricularia oryzae*. Active against gram-positive bacteria and fungi. Needles. Sol. MeOH, EtOAc; fairly sol. H_2O ; poorly sol. Et_2O , hexane. Mp 73°. $[\alpha]_D -19$ (H_2O). λ_{max} 240 (E1%/1cm 2824); 308 (E1%/1cm 350) (MeOH) (Berdy).

Tamari, K. *et al.*, *Nippon Nogei Kagaku Kaishi*, 1954, **28**, 254-258; *CA*, **50**, 41086 (*isol*)

Umetsu, N. *et al.*, *Agric. Biol. Chem.*, 1974, **38**, 1867-1874 (*isol, uv, ir, ms*)

Piriferine P-479


2-Methyl-N-[1-(1-oxo-3-phenyl-2-propenyl)-2-pyrrolidinyl]propanamide, 9CI. N-Cinnamoyl-2-(2-methylpropanoylamino)-pyrrolidine [113689-36-4]



$C_{17}H_{22}N_2O_2$ 286.373
 Abs. config. unknown. (R)-form ill. by analogy with Odorine, O-67. Major alkaloid from the leaves of *Aglaia pirifera* (Meliaceae). Cryst. ($Et_2O/CHCl_3$). Mp 164-165.5°. $[\alpha]_D^{28} +30$ (c, 0.01 in EtOH).

2'-Hydroxy: **Piriferinol**
 $C_{17}H_{22}N_2O_3$ 302.372
 Alkaloid from *Aglaia elaeagnoides*. $[\alpha]_D^{20} +8$ (c, 0.1 in $CHCl_3$). λ_{max} 284 (log ϵ 4.16) (MeOH).

- Saifah, E. *et al.*, *J. Nat. Prod.*, 1988, **51**, 80-82; 1993, **56**, 473-477 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Brader, G. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1482-1490 (*Piriferinol*)

Piscarinine A P-480


$C_{26}H_{29}N_3O_3$ 431.533

(S)-form [312956-60-8]
 Prod. by *Penicillium piscarium* NKM F-691. Waxy solid. $[\alpha]_D^{25} +14$ (c, 1 in MeOH). λ_{max} 216 ; 244 ; 262 (MeOH).

Kozlovsky, A. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 333-340 (*isol, uv, ms, pmr, cmr*)

Piscarinine B

P-481

C₂₆H₂₇N₃O₃ 429.518**(S)-form** [312956-61-9]


Prod. by *Penicillium piscarium*. Waxy solid. $[\alpha]_D^{25} +132$ (c, 1 in MeOH). λ_{\max} 218 ; 240 ; 265 (MeOH).

Kozlovsky, A. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 333-340 (*isol, uv, ms, pmr, cmr*)

Piscerythoxazole

P-482

[151590-43-1]

C₂₂H₁₉NO₅ 377.396


Constit. of the root bark of *Piscidia erythrina*. λ_{\max} 203 ; 218 ; 242 (sh) ; 250 ; 295 ; 305 (sh) (MeOH).

Moriyama, M. *et al.*, *Phytochemistry*, 1993, **32**, 1317 (*isol, uv, pmr, cmr, ms*)

Piscerythramine

P-483

2-[4-Amino-3-hydroxy-5-methoxy-2,6-bis(3-methyl-2-butenyl)phenyl]-5,7-dihydroxy-4H-1-benzopyran-4-one, 9CI. 4'-Amino-3',5',7-trihydroxy-5'-methoxy-2',6'-diprenylisoflavone [132923-36-5]

C₂₆H₂₉NO₆ 451.518


The first naturally occurring aminoisoflavone. Isol. from the root bark of *Piscidia erythrina*. Pale yellow gum. λ_{\max} 213 ; 260 ; 294 (broad maximum) (MeOH).

Moriyama, M. *et al.*, *Phytochemistry*, 1993, **32**, 1317 (*isol, uv, pmr, cmr*)

Pisopowine

P-484

[107913-08-6]



Absolute Configuration

C₄₀H₄₈N₂O₆ 652.829

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D -152$ (c, 0.4 in MeOH).

6-O-De-Me: Pisopowidine

[107882-04-2]

C₃₉H₄₆N₂O₆ 638.802

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D -137$ (c, 1.2 in MeOH).

6,6'-Di-O-de-Me: Pisopowiarine

[107882-05-3]

C₃₈H₄₄N₂O₆ 624.775

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Cryst. (Me₂CO). Mp 187-188°. $[\alpha]_D -112$ (c, 1 in MeOH).

12,12'-Di-O-de-Me: Pisopowetene

[107882-06-4]

C₃₈H₄₄N₂O₆ 624.775

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D -80$ (c, 0.2 in MeOH).

12,12'-Di-O-de-Me, 2'-N-de-Me: Pisopowamine

[107882-08-6]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D -68$ (c, 0.15 in MeOH).

6,6',12'-Tri-O-de-Me: Pisopowiaridine

[107882-07-5]

C₃₇H₄₂N₂O₆ 610.749

Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Cryst. (MeOH). Mp 184-185°. $[\alpha]_D -78$ (c, 1 in MeOH).

6,6',12'-Tri-O-de-Me, 2'-N-de-Me: 2'-Norpisopowiaridine

[107882-09-7]

C₃₆H₄₀N₂O₆ 596.722


Alkaloid from the bark of *Popowia pisocarpa* (Annonaceae). Amorph. $[\alpha]_D -46$ (c, 0.33 in MeOH).

Jossang, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1018 (*isol, uv, pmr, cmr, ms, cd, struct*)

Pistacidelorel

P-485

[935854-05-0]



Absolute Configuration

C₂₄H₃₈N₂O 370.577


Alkaloid from the leaves of *Pistacia atlantica* ssp. *mutica*.

Meshkatsadat, M.H. *et al.*, *Asian J. Chem.*, 2007, **19**, 1599-1601 (*isol, struct*)

Pithecolobine

P-486

[22368-82-7]



A mixt. of homologues and analogues. Estimated percentages of the 3 main components are: m = 3, n = 6: 24-30%; m = 1, n = 8: 40-49%; m = 1, n = 10: 13.5-16.5%. Closely related to the Budmunchiamines. Alkaloid from *Samanea saman* (*Pithecolobium saman*) (Fabaceae). Pale-yellow oil which slowly crystallises but cannot be recryst. Taxonomy of the source plant is complex. Authors give alt. genera *Samanea* or *Pithecolobium* (*Pithecolobium*). Kew system refers *Samanea* to *Albizzia*.

Piconolate salt:


Cryst. (Me₂CO/MeOH). Mp 134-135°.

Wiesner, K. *et al.*, *Can. J. Chem.*, 1968, **46**, 1881; 3617 (*struct, bibl*)

Pitiamide A

P-487

[198416-38-5]



Absolute Configuration

C₂₂H₃₆ClNO₂ 381.985

Isol. from a mixt. of *Lynghya majuscula* and *Microcoleus* sp. growing on coral. $[\alpha]_D -10.3$ (c, 3 in CHCl₃). λ_{\max} 223 (MeOH).


Nagle, D.G. *et al.*, *Tet. Lett.*, 1997, **38**, 6969-6972 (*isol, uv, ir, pmr, cmr*)

Ribe, S. *et al.*, *J.A.C.S.*, 2000, **122**, 4608-4617 (*synth, cd, ir, pmr, abs config*)

Pityriacitrin

P-488

1-(1H-Indole-3-carbonyl)-9H-pyrido[3,4-b]indole. 1-(1H-Indole-3-carbonyl)-β-carboline [244295-64-5]

C₂₀H₁₃N₃O 311.342

Prod. by the yeast *Malassezia furfur* and

the marine bacterium *Paracoccus* sp. Uv protectant. Mp 227-230°. λ_{\max} 193 (log ϵ 4.24); 216 (log ϵ 4.47); 262 (sh) (log ϵ 3.91); 281 (sh) (log ϵ 3.95); 289 (log ϵ 3.99); 310 (log ϵ 3.86); 389 (log ϵ 3.88) (MeCN).

5',6-Dihydroxy-1-(5-Hydroxy-1H-indole-3-carbonyl)-9H-pyrido[3,4-b]indol-6-ol. Hyrtiosulawesine

[452067-33-3]

$C_{20}H_{13}N_3O_3$ 343.341

Isol. from the sponge *Hyrtios erectus*. Phospholipase A_2 inhibitor. λ_{\max} 217 (ϵ 145000); 260 (sh); 305 (ϵ 53000); 325 (sh); 403 (ϵ 22700) (MeOH).

Japan. Pat., 1999, 99 269 175; CA, 131, 242087z (isol, activity)


Mayer, P. et al., Arch. Dermatol. Res., 2002, 294, 131-134 (isol, ir, pmr, cmr, ms)

Salmoun, M. et al., J. Nat. Prod., 2002, 65, 1173-1176 (Hyrtiosulawesine)

Sauleau, P. et al., J. Nat. Prod., 2006, 69, 1676-1679 (Hyrtiosulawesine)

Pityriacitrin B P-489

1-(1H-Indol-3-ylcarbonyl)-9H-pyrido[3,4-b]indole-3-carboxylic acid. 1-(1H-Indole-3-carbonyl)- β -carboline-3-carboxylic acid
[863766-95-4]




$C_{21}H_{13}N_3O_3$ 355.352

Alkaloid from the yeast *Malassezia furfur*. Yellow solid. Mp > 300° dec. λ_{\max} 214 (log ϵ 4.49); 266 (log ϵ 4.18); 280 (log ϵ 4.17); 289 (log ϵ 4.17); 391 (log ϵ 3.74) (MeOH).

Irlinger, B. et al., Helv. Chim. Acta, 2005, 88, 1472-1485 (isol, ir, pmr, cmr)

Pityrialactone P-490

4,5-Di-1H-indol-3-yl-2,3-furandione




$C_{20}H_{12}N_2O_3$ 328.326

Isol. from the yeast *Malassezia furfur*. Pale yellow powder. Mp > 250°. λ_{\max} 224 (log ϵ 4.07); 292 (sh) (log ϵ 3.55); 356 (log ϵ 3.55) (MeOH).

Irlinger, B. et al., Angew. Chem., Int. Ed., 2004, 43, 1098-1100 (isol, pmr, cmr, ms)

Pityrianhidride P-491

3,4-Di-1H-indol-3-yl-2,5-furandione, 9CI. 3,4-Bis(1H-indol-3-yl)maleic anhydride
[115684-57-6]




$C_{20}H_{12}N_2O_3$ 328.326

Isol. from the yeast *Malassezia furfur*. Orange solid. Mp 227-230°.

Brenner, M. et al., Tetrahedron, 1988, 44, 2887-2892 (synth, pmr)

Irlinger, B. et al., Angew. Chem., Int. Ed., 2004, 43, 1098-1100 (isol)

Pityriarubin A P-492




$C_{32}H_{22}N_4O_4$ 526.55

Isol. from the yeast *Malassezia furfur*. Orange-red powder. $[\alpha]_D^{25}$ -80 (c, 0.05 in MeCN). Mp > 200° dec. λ_{\max} 198 (log ϵ 4.46); 219 (log ϵ 4.43); 277 (log ϵ 4.04); 385 (log ϵ 3.63); 436 (log ϵ 3.68) (MeCN).

Irlinger, B. et al., Angew. Chem., Int. Ed., 2004, 43, 1098-1100 (isol, pmr, cmr, cd)

Pityriarubin B P-493



X = NH

$C_{32}H_{20}N_4O_4$ 524.534

Isol. from the yeast *Malassezia furfur*. Orange powder. Mp 180-183° dec. λ_{\max} 196 (log ϵ 4.38); 218 (log ϵ 4.34); 277 (log ϵ 4.05); 382 (log ϵ 3.55); 446 (log ϵ 3.61) (MeCN).

Irlinger, B. et al., Angew. Chem., Int. Ed., 2004, 43, 1098-1100 (isol, pmr, cmr, ms)

Pityriarubin C P-494

As Pityriarubin B, P-493 with X = O


$C_{32}H_{19}N_3O_5$ 525.519

Isol. from the yeast *Malassezia furfur*. Red solid. Mp 125-130° dec. λ_{\max} 195 (log ϵ 4.66); 218 (log ϵ 4.6); 277 (log ϵ 4.37); 323 (sh) (log ϵ 4.08); 394 (log ϵ 3.74); 462 (log ϵ 3.83) (MeCN).

Irlinger, B. et al., Angew. Chem., Int. Ed., 2004, 43, 1098-1100 (isol, pmr, cmr, ms)

Pityriazole P-495

2-Hydroxy-1-(1H-indol-3-yl)-9H-carbazole-3-carboxylic acid
[454222-42-5]



$C_{21}H_{14}N_2O_3$ 342.353


Alkaloid from the yeast *Malassezia furfur*. Amorph. solid. Mp > 250°. λ_{\max} 221 (log ϵ 3.77); 279 (log ϵ 3.52); 336 (sh) (log ϵ 2.95) (MeOH).

Irlinger, B. et al., Helv. Chim. Acta, 2005, 88, 1472-1485 (isol, pmr, cmr, ms)

Forke, R. et al., Org. Biomol. Chem., 2008, 6, 2481-2483 (synth)

Plakinamine A P-496

[93474-13-6]



$C_{29}H_{46}N_2$ 422.696

Alkaloid from a sponge *Plakina* sp. Active against *Staphylococcus aureus* and *Candida albicans*. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 120-130° dec. $[\alpha]_D^{+16}$ (c, 1.02 in $CHCl_3$). First report of a steroidal alkaloid from a marine organism. Acetylation causes rearrangement and ring opening of the pyrroline system. λ_{\max} 246 (ϵ 7100) (MeOH) (Derep).

24,25-Dihydro-24,25-Dihydroplakinamine A

[224576-02-7]

$C_{29}H_{48}N_2$ 424.712

Alkaloid from a *Corticium* sp. Cytotoxic agent. $[\alpha]_D^{+7.4}$ (c, 0.01 in $CHCl_3/MeOH$). C-24 config. not de-

terminated.

23ξ,24ξ,25,30-Tetrahydro: **Tetrahydro-plakinamine A**

[454476-89-2]

C₂₉H₅₀N₂ 426.727

Alkaloid from a *Corticium* sp. Gum. [α]_D²² +23.2 (c, 0.19 in MeOH).

23ξ,24ξ,25,30-Tetrahydro, N³⁰-Me:

[224576-05-0]

C₃₀H₅₂N₂ 440.754

Alkaloid from a *Corticium* sp. Cytotoxic agent. [α]_D +23 (c, 0.02 in CHCl₃/MeOH).

4β-Acetoxy, 23ξ,30-dihydro, N³-Me:

Plakinamine K

[64734-88-9]

C₃₂H₅₂N₂O₂ 496.775

Alkaloid from the sponge *Corticium niger*. Cytotoxic. Oil. [α]_D +38.4 (c, 0.25 in MeOH). λ_{max} 203 (ε 59200); 246 (ε 1200) (MeOH).

4β-Acetoxy, 23ξ,24ξ,25,30-tetrahydro,

N³-Me: **Dihydroplakinamine K**

[640734-89-0]

C₃₂H₅₄N₂O₂ 498.791

Alkaloid from the sponge *Corticium niger*. Cytotoxic. Oil (as dihydrochloride). [α]_D +5.7 (c, 0.05 in MeOH) (dihydrochloride). λ_{max} 203 (ε 54800); 247 (ε 1350) (MeOH) (dihydrochloride).

3-Epimer, 4-oxo, N³,N³-di-Me: **Plakinamine F**

[383859-60-7]

C₃₁H₄₈N₂O 464.733

Alkaloid from an undescribed sponge of the genus *Corticium*. Gum. [α]_D²⁵ +8.4 (c, 0.1 in MeOH). λ_{max} 245 (log ε 3.97) (MeOH).

3-Epimer, 4β-hydroxy, N³-Ac: **Lokysterolamine B**

[159934-15-3]

C₃₁H₄₈N₂O₂ 480.732

Alkaloid from the sponge *Corticium* sp. Immunosuppressive agent. Semicryst. solid. [α]_D²⁶ -3.1 (c, 1.6 in CHCl₃). λ_{max} 247 (ε 8750) (MeOH) (Berdy).

3-Epimer, 4β-hydroxy, N³,N³-di-Me: **Lokysterolamine A**

[159934-14-2]

C₃₁H₅₀N₂O 466.749

Alkaloid from an undescribed sponge of the genus *Corticium*. Immunosuppressant. Oil. [α]_D²⁵ +17.7 (c, 0.1 in MeOH). λ_{max} 247 (log ε 3.91) (MeOH).

3-Epimer, 4β-hydroxy, N³,N³-di-Me, N³⁰-oxide: **Plakinamine E**. Lokysterolamine A N³⁰-oxide

[383859-59-4]

C₃₁H₅₀N₂O₂ 482.748

Alkaloid from an undescribed sponge of the genus *Corticium*. Gum. [α]_D²⁵ +9.3 (c, 0.2 in MeOH). λ_{max} 249 (log ε 3.85); 279 (sh) (log ε 3.56) (MeOH).

Rosser, R.M. *et al.*, *J.O.C.*, 1984, **49**, 5157-5160 (*isol, ir, uv, pmr, cmr, struct*)

Jurek, J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1004-1007 (*Lokysterolamines*)

De Marino, S. *et al.*, *Eur. J. Org. Chem.*, 1999, 697-701 (*derivs*)


Lee, H.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1474-1476 (*Plakinamine E, Plakinamine F*)

Borbone, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1206-1209 (*Tetrahydroplakinamine A*)

Ridley, C.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1536-1539 (*Plakinamine K, Dihydroplakinamine K*)

Plakinamine B

[93474-14-7]



C₃₁H₅₀N₂ 450.749

Alkaloid from a marine sponge *Plakina* sp. Active against *Staphylococcus aureus* and *Candida albicans*. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 180-200° dec. (as hydrochloride). [α]_D +29 (c, 1.19 in MeOH). λ_{max} 241 (ε 2700) (MeOH) (Derep).

4α-Hydroxy, N³-de-Me: **4α-Hydroxy-N³-demethylplakinamine B**

C₃₀H₄₈N₂O 452.722

Alkaloid from the sponge *Corticium* sp. Gum. [α]_D²² +6.7 (c, 0.09 in MeOH). λ_{max} 241 (log ε 3.8) (MeOH).

3-Epimer, 4-oxo, N³-Me: **N³-Methyl-4-oxo-3-epiplakinamine B**

C₃₂H₅₀N₂O 478.76

Alkaloid from the sponge *Corticium* sp. Cytotoxic agent. [α]_D +35.4 (c, 0.01 in CHCl₃/MeOH).

3-Epimer, 4-oxo, N²⁷-de-Me, N³-Me:

Plakinamine H

C₃₁H₄₈N₂O 464.733

Alkaloid from a *Corticium* sp. Gum. [α]_D²² +29 (c, 0.1 in MeOH). λ_{max} 250 (log ε 3.77) (MeOH).


Rosser, R.M. *et al.*, *J.O.C.*, 1984, **49**, 5157-5160 (*isol, ir, uv, pmr, cmr, struct*)

De Marino, S. *et al.*, *Eur. J. Org. Chem.*, 1999, 697-701 (*deriv, pmr, cmr*)

Borbone, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1206-1209 (*4-Hydroxy-N³-demethylplakinamine B, Plakinamine H*)

Plakinamine C

[224575-95-5]



C₃₃H₅₄N₂O₂ 510.802

Alkaloid from the sponge *Corticium* sp. Cytotoxic agent. [α]_D +29.4 (c, 0.02 in CHCl₃/MeOH).

Δ⁸-Isomer: **Plakinamine D**

[224575-99-9]


C₃₃H₅₄N₂O₂ 510.802

Alkaloid from *Corticium* sp. Cytotoxic agent. [α]_D +25.2 (c, 0.01 in CHCl₃/MeOH).

De Marino, S. *et al.*, *Eur. J. Org. Chem.*, 1999, 697-701 (*isol, pmr, cmr*)

Plakinamine G

[454476-87-0]




C₂₉H₄₄N₂O 436.679

Alkaloid from the sponge *Corticium* sp. Gum. [α]_D²² -24.4 (c, 0.09 in MeOH). λ_{max} 276 (log ε 4.11) (MeOH).

Borbone, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1206-1209 (*isol, pmr, cmr*)

Plakinamine I†

P-500



C₃₂H₅₆N₂O₂ 500.807


Alkaloid from the sponge *Corticium* sp. Gum. [α]_D²⁵ -18.3 (c, 0.03 in MeOH). Rel. config. of C-23/C-24 is *trans*.

Zampella, A. *et al.*, *Eur. J. Org. Chem.*, 2005, 4359-4363 (*isol, pmr, cmr*)

Plakinamine I†

P-501

[640734-86-7]




C₃₁H₅₀N₂ 450.749

Alkaloid from the sponge *Corticium niger*. Cytotoxic. Oil (as dihydrochloride). [α]_D +45.2 (c, 0.32 in MeOH) (dihydrochloride). λ_{max} 202 (ε 49800); 247 (ε 1500) (MeOH) (dihydrochloride).

Ridley, C.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1536-1539 (*isol, pmr, cmr*)

Plakinamine J

[640734-87-8]


C₃₀H₅₀N₂ 438.738

Alkaloid from the sponge *Corticium niger*. Cytotoxic. Oil (as dihydrochloride). $[\alpha]_D^{25}$ +25 (c, 0.1 in MeOH) (dihydrochloride). λ_{\max} 203 (€ 56800); 249 (€ 2030) (MeOH) (dihydrochloride).

Ridley, C.P. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1536-1539 (*isol*, *pmr*, *cmr*)

Plakinidine A**P-503**

10,11-Dihydro-7-(methylamino)benzo[b]pyrrolo[4,3,2-de][1,10]phenanthroline-8(9H)-one, 9CI
[124512-44-3]

C₁₈H₁₄N₄O 302.335

First reported natural pyrrolo[2,3,4-*k*]acridines. Alkaloid from the sponge *Plakortis* sp. Exhibits *in vitro* activity against the parasite *Hippostrongylus brasiliensis* and shows weak activity against reverse transcriptase. Flat purple plates (CHCl₃). Fairly sol. AcOH; poorly sol. MeOH, hexane. Mp 248-250°. λ_{\max} 226 (€ 15800); 247 (€ 20400); 279 (€ 11600); 336 (€ 22700); 378 (€ 6200); 537 (€ 3600) (MeOH) (Derep).

N⁷-Me: Plakinidine B

[124512-45-4]

C₁₉H₁₆N₄O 316.362

Alkaloid from the sponge *Plakortis* sp. Exhibits *in vitro* activity against the parasite *Nocardia brasiliensis*. Reverse transcriptase inhibitor. Purple oil. Fairly sol. AcOH; poorly sol. MeOH, hexane. λ_{\max} 226 (€ 13800); 248 (€ 17500); 283 (€ 9100); 347 (€ 19700); 383 (€ 4700); 544 (€ 4000) (MeOH) (Derep).

9,10-Didehydro: Plakinidine C

[129744-15-6]

C₁₈H₁₂N₄O 300.319

Alkaloid from the sponge *Plakortis* sp. Cytotoxic, anthelmintic. Reverse transcriptase inhibitor. λ_{\max} 244 (€ 42800); 254 (€ 39100); 323 (€ 25200); 370 (€

10900); 508 (€ 13800); 541 (€ 11400) (MeOH) (Derep).

N⁷-De-Me: Plakinidine D

[197234-19-8]

C₁₇H₁₂N₄O 288.308

Alkaloid from *Didemnum rubeum* and an unidentified *Didemnum* sp. Dark red solid. λ_{\max} 250 (€ 4770); 282 (€ 3500); 328 (€ 2980); 388 (€ 1940); 436 (€ 820); 514 (€ 926) (CHCl₃/MeOH).

Inman, W.D. *et al.*, *J.A.C.S.*, 1990, **112**, 1 (*isol*, *pmr*, *cmr*, *ms*, *struct*)


West, R.R. *et al.*, *Tet. Lett.*, 1990, **31**, 3271 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Smith, C.J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1048-1050 (*Plakinidine D*)

Ford, P.W. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1051-1053 (*Plakinidine D*)

Plakinidine E

[928673-36-3]


P-504C₁₇H₁₁N₃O₂ 289.293

Alkaloid from *Plakortis quasiamphiaster*. Dark red-purple solid.

Ralifo, P. *et al.*, *J. Nat. Prod.*, 2007, **70**, 95-99 (*isol*, *pmr*, *cmr*)

Plakoridine A

[155944-28-8]

P-505

Relative Configuration

R = -CH₂CH₂CH₃C₃₅H₅₇NO₅ 571.839

Alkaloid from the Okinawan marine sponge *Plakortis* sp. Weakly cytotoxic against murine lymphoma L1210 cells *in vitro*. Oil. $[\alpha]_D^{19}$ -0.4 (c, 0.5 in CHCl₃). λ_{\max} 317 (€ 22000) (MeOH) (Derep).

Takeuchi, S. *et al.*, *J.O.C.*, 1994, **59**, 3712-3713 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Ma, D. *et al.*, *Tet. Lett.*, 2000, **41**, 1947-1950 (*synth*)

Etchells, L.L. *et al.*, *Tetrahedron*, 2006, **62**, 10914-10927 (*synth*)

Plakoridine B

[164415-61-6]

P-506

As Plakoridine A, P-505 with

R = -(CH₂)₁₄CH₃C₄₇H₈₁NO₅ 740.161

Alkaloid from an Okinawan sponge *Plakortis* sp. Oil. λ_{\max} 224 (€ 12200); 318 (€ 16000) (MeOH).


Takeuchi, S. *et al.*, *Tetrahedron*, 1995, **51**,

5979-5986 (*isol*, *pmr*, *cmr*)

Etchells, L.L. *et al.*, *Tetrahedron*, 2006, **62**, 10914-10927 (*synth*)

Plakortamine A**P-507**

7-Bromo-N,N-dimethyl-9H-pyrido[3,4-b]indole-1-ethanamine. 7-Bromo-1-[2-(dimethylamino)ethyl]-β-carboline
[467419-71-2]


C₁₅H₁₆BrN₃ 318.216

Alkaloid from the sponge *Plakortis nigra*. Cytotoxic. Pale yellow oil. λ_{\max} 243 (€ 20000); 295 (€ 11700) (MeOH).

Sandler, J.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1258-1261 (*isol*, *pmr*, *cmr*)

Plakortamine C

[467419-73-4]


P-508C₂₇H₂₃Br₂N₅ 577.32

Alkaloid from the sponge *Plakortis nigra*. Cytotoxic. Pale yellow gum. λ_{\max} 243 (€ 57400); 296 (€ 29000) (MeOH).

Sandler, J.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1258-1261 (*isol*, *pmr*, *cmr*)

Plakortamine D


[467419-74-5]

P-509C₁₅H₁₄BrN₃O 332.199**(±)-form**

[467419-74-5]

Alkaloid from the sponge *Plakortis nigra*. Cytotoxic agent. Pale yellow oil. $[\alpha]_D^{25}$ -2.1 (c, 0.6 in MeOH). λ_{\max} 243 (€ 26600); 296 (€ 14700) (MeOH).

Sandler, J.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1258-1261 (*isol*, *pmr*, *cmr*)

Planchonelline**P-510**C₁₂H₁₉NO₂S 241.354

Ester of laburnine with 3-(Methylthio)-2-propenoic acid, M-577. A major alkaloid from the leaves of *Planchonella thyrsoidea*

(preferred genus name *Pouteria*) and *Planchonella anteridifera* (Sapotaceae).
Oil. $[\alpha]_D^{25} +9$ (c, 0.85 in EtOH).

8-Epimer: Isoetronocyl trans-β-methylthioacrylate

$C_{12}H_{19}NO_2S$ 241.354

Alkaloid from leaves of an unidentified *Planchonella* sp. (Sapotaceae). Pale yellow oil. $[\alpha]_D^{25} -57.2$ (c, 1.52 in EtOH). Ester of isetronocanol with 3-(Methylthio)-2-propenoic acid, M-577.

8-Epimer: picrate:


Yellow needles (EtOH). Mp 111-113°.

Hart, N.K. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1259; 1968, **21**, 1393 (*isol, uv, ir, struct, synth*)

Platencin

P-511

[869898-86-2]



$C_{24}H_{27}NO_6$ 425.48

Closely related to Platensimycin, P-513. The abs. config. is prob. as shown. Prod. by *Streptomyces platensis* MA 7339. FabH and FabF dual inhibitor. Broad spectrum antibiotic. Amorph. powder. $[\alpha]_D^{23} -7$ (c, 0.85 in MeOH). λ_{max} 226 (ε 16840); 296 (ε 2660) (MeOH).

Jayasuriya, H. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 4684-4688 (*isol, pmr, cmr, ms*)

Wang, J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, **104**, 7612-7616 (*isol, activity*)

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1780-1783 (*synth*)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 2008, **130**, 11292-11293; 14016 (*synth*)


Austin, K.A.B. *et al.*, *Org. Lett.*, 2008, **10**, 4465-4468 (*synth*)

Herath, K. *et al.*, *Tet. Lett.*, 2008, **49**, 5755-5758 (*biosynth*)

Platensimide A

P-512

[1022904-20-6]



$C_{23}H_{32}N_2O_6$ 432.516

Prod. by *Streptomyces platensis* MA7327. Gum. $[\alpha]_D^{23} +29.2$ (c, 1.3 in MeOH). λ_{max} 230 (ε 4867) (MeOH).

1-Parent acid: Platensic acid

[1023284-62-9]

$C_{17}H_{22}O_4$ 290.358

Prod. by *Streptomyces platensis* MA7327. Amorph. powder. $[\alpha]_D^{23} -30$ (c, 1.5 in MeOH). λ_{max} 234 (ε 8405) (MeOH).

1-Parent acid, Me ester: Methyl platensinoate

[1022904-19-3]

$C_{18}H_{24}O_4$ 304.385


Prod. by *Streptomyces platensis* MA7327. Gum. $[\alpha]_D^{23} -26.6$ (c, 3.5 in MeOH). λ_{max} 234 (ε 10078) (MeOH).

Herath, K.B. *et al.*, *Org. Lett.*, 2008, **10**, 1699-1702 (*isol, pmr, cmr*)

Platensimycin

P-513

[835876-32-9]



Absolute Configuration

$C_{24}H_{27}NO_7$ 441.48

Prod. by *Streptomyces platensis* isol. from soil. Inhibitor of cellular lipid biosynth. Active against gram-positive bacteria incl. MRSA. Cryst. (MeNO₂). Mp 220-222°. $[\alpha]_D^{23} -51.1$ (c, 0.135 in MeOH). λ_{max} 227 (ε 28167); 240 (sh); 296 (ε 4825) (MeOH).

Singh, S.B. *et al.*, *J.A.C.S.*, 2006, **128**, 11916-11920; 15547 (*isol, struct, abs config*)

Wang, J. *et al.*, *Nature (London)*, 2006, **441**, 358-361 (*isol, pmr, cmr, cryst struct*)

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 3942-3945; 2008, **47**, 944-946 (*synth*)

Nicolaou, K.C. *et al.*, *Chem. Comm.*, 2007, 1922-1923 (*synth*)


Zou, Y. *et al.*, *Org. Lett.*, 2007, **9**, 1825-1828 (*synth*)

Matsuo, J. *et al.*, *Org. Lett.*, 2008, **10**, 4049-4052 (*synth*)

Platylerine

P-514

5,6,11,12-Tetrahydro-3,8,9-trimethoxy-13-methylidibenzo[a,e]cycloocten-5,11-imin-4-ol, 9CI. 1-Hydroxy-2,8,9-trimethoxypavanine



(-)-form

$C_{20}H_{23}NO_4$ 341.406

(-)-form [18826-68-1]

Alkaloid from the above-ground parts of *Argemone gracilentia*, the roots and above-ground parts of *Argemone platyceras*, and the tops of *Thalictrum revolutum* (Papaveraceae, Ranunculaceae). Cryst. (Et₂O). $[\alpha]_D^{21} -267$ (c, 0.24 in CHCl₃). $[\alpha]_D^{25} -305$ (c, 0.2 in MeOH).

Picrate: Mp 155-160°.

N-Me: Platylerine N-methosalt

$C_{21}H_{26}NO_4^{\oplus}$ 356.441

Quaternary alkaloid from the aerial parts of *Argemone platyceras* (Papaveraceae). Cryst. (MeOH)(as chloride). Mp 165° (chloride). $[\alpha]_D^{25} -258$ (c, 0.47 in MeOH).

Me ether: O-Methylplatylerine. O,O-Dimethylmunitagine

[7688-85-9]

$C_{21}H_{25}NO_4$ 355.433

Alkaloid from *Argemone platyceras*.

Cryst. (Et₂O/pentane). Mp 124-125.5°. $[\alpha]_D^{27} -292$ (c, 0.35 in CHCl₃).

O⁸-De-Me: 1,8-Dihydroxy-2,9-dimethoxypavanine. Munitagine

[7691-07-8]

Alkaloid from the above-ground parts of *Argemone gracilentia* and *Argemone munita* var. *rotundata* (Papaveraceae). Needles (MeOH/Et₂O). Mp 167-169°. $[\alpha]_D^{27} -239$ (c, 0.5 in CHCl₃). Has also been named as 2,7-dihydroxy-3,8-dimethoxy.

(±)-form [38863-79-5]

Synthetic. Noncryst.

O⁸-De-Me: [55903-28-1]

Synthetic. Powder (MeOH/Et₂O). Mp not recorded.

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 1728 (*isol, uv, ir*)

Stermitz, F.R. *et al.*, *J.O.C.*, 1966, **31**, 2925; 1969, **34**, 555; 1973, **38**, 1761 (*Munitagine, Platylerine, isol, pmr, ms, synth, struct*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 2513; 1976, **41**, 285 (*Platylerine, Platylerine N-methosalt, isol, uv, ir, ms, synth, struct*)

Stermitz, F.R. *et al.*, *Indian J. Chem.*, 1974, **12**, 1249 (*Munitagine, synth, uv*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 593; 1980, **43**, 270 (*Platylerine, isol, uv, cd, pmr*)


Israilov, I.A. *et al.*, *Khim. Prirod. Soedin.*, 1986, **22**, 204-206; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 189-192 (*O-Methylplatylerine*)

Lee, S.S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 803-810 (*Munitagine, ed, pmr, cmr*)

Platydesmine

P-515

2,3-Dihydro-4-methoxy-α,α-dimethylfuro[2,3-b]quinoline-2-methanol, 9CI



(R)-form

$C_{15}H_{17}NO_3$ 259.304

Absolute configuration revised in 2000.

(R)-form [318465-89-3]

Alkaloid from the leaves of *Choisya ternata*. Mp 137-138°. $[\alpha]_D -47$ (c, 0.76 in MeOH).

N-Me:

$C_{16}H_{20}NO_3^{\oplus}$ 274.339

Alkaloid from the leaves, roots and stems of *Choisya ternata*. Mp 180° (as perchlorate). $[\alpha]_D -24.3$ (c, 0.11 in MeOH) (perchlorate).

(S)-form [2824-86-4]

Alkaloid from *Platydesma campanulata*, *Geijera salicifolia* and *Zanthoxylum belizense* (Rutaceae). Needles (C₆H₆/petrol). Mp 137-138°. $[\alpha]_D +47$ (c, 0.7 in MeOH).

Ac: O-Acetylplatydesmine

$C_{17}H_{19}NO_4$ 301.341

Minor alkaloid from the leaves of *Geijera salicifolia* (Rutaceae). Prisms (Me₂CO/Et₂O). Mp 126-127°. $[\alpha]_D^{23} +23$ (c, 1.8 in CHCl₃).

N-Me: Platydesminium

$C_{16}H_{20}NO_3^{\oplus}$ 274.339

Alkaloid from *Skimmia japonica*, *Ruta*

graveolens (rue) and many other Rutaceae spp. Mp 192-196° (as chloride). $[\alpha]_D^{20} +31$ (c, 0.74 in MeOH) (as iodide).

(±)-**form** [7764-73-0]

Alkaloid from the leaves of *Melicope perspicuineria* (Rutaceae). Plates (C₆H₆/petrol). Mp 137-138°.

Ac:

Synthetic. Needles (C₆H₆). Mp 126-127°.

N-Me:

Prisms (MeOH/diisopropyl ether). Mp 156-158° dec.

Wernhey, F. *et al.*, *Tetrahedron*, 1963, **19**, 1293 (*isol. struct. ir, uv, ms*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1991 (*isol, pmr*)

Bowman, R.M. *et al.*, *J.C.S. (C)*, 1966, 1504 (*synth, pmr*)

Murphy, S.T. *et al.*, *Aust. J. Chem.*, 1974, **27**, 187 (*isol*)

Rideau, M. *et al.*, *Phytochemistry*, 1979, **18**, 155 (*isol, ms*)

Ramesh, M. *et al.*, *Heterocycles*, 1984, **22**, 125 (*synth, ir, pmr*)


Boyd, D.R. *et al.*, *J.C.S. Perkin 1*, 2000, 3397-3405 (*synth, abs config*)

Boyd, D.R. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 2983-2991 (*isol*)

Platyphylline

P-516

1,2-Dihydro-12-hydroxysenecionan-11,16-dione, 9CI. Platifillin
[480-78-4]



Absolute Configuration

C₁₈H₂₇NO₅ 337.415

Alkaloid from *Senecio platyphyllus*, *Senecio adnatus*, *Senecio hygrophyllus*, other *Senecio* spp., *Adenostyles* spp. (preferred genus name *Cacalia*) and *Petasites laevigatus* (Asteraceae). Shows atropine-like activity. Antispasmodic, mydriatic agent. Has been used in the USSR for treatment of gastrointestinal hypermotility and peptic ulceration. Used as tartrate. Needles (H₂O). Spar. sol. hot H₂O, petrol. Mp 129° (124-125°). $[\alpha]_D^{20} -56$ (CHCl₃). $[\alpha]_D^{20} -59$ (EtOH). Log P 0.45 (uncertain value) (calc). Rel. nontoxic; lacks hepatotoxicity of other pyrrolizidine alkaloids.

▶ LD₅₀ (rat, ipr) 252 mg/kg. TP2518000

Methodide:

Needles (EtOH). Mp 216-217°. $[\alpha]_D^{20} -31.3$.

N-Oxide: Platyphylline N-oxide

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio* spp. The form in which Platyphylline is present in *Senecio hygrophyllus*, and probably in other spp. (Asteraceae). Mp 180-181°. $[\alpha]_D^{20} -44.6$ (CHCl₃). $[\alpha]_D^{20} -59$ (H₂O).

1 α -Hydroxy: Hadiensine

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio hadiensis* (As-

teraceae). Gum; cryst. (MeOH/EtO-H) (as perchlorate). Mp 279-280° (perchlorate). $[\alpha]_D^{20} -83$ (c, 1.6 in MeOH).

1 α -Hydroxy, 12-Ac: 12-O-Acetylhadiensine

C₂₀H₂₉NO₇ 395.452

Alkaloid from *Senecio hadiensis* (Asteraceae). Gum.

14 α -Hydroxy: Hygrophylline

[3573-82-8]

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio hygrophyllus* (Asteraceae). Prisms (Me₂CO). Mp 176° (173-174°). $[\alpha]_D^{20} -67.3$ (c, 2.9 in EtOH). Hydrol. gives platynecine and hygrophyllineic acid.

18-Hydroxy: Dihydroretorsine

C₁₈H₂₇NO₆ 353.414

Alkaloid from the roots of *Senecio subulatus* var. *erectus* (Asteraceae). Needles (CHCl₃/MeOH). Mp 62-63°. $[\alpha]_D^{20} -2.68$ (c, 0.078 in CHCl₃).

2 α ,18-Dihydroxy: Pettitanine

[137760-49-7]

C₁₈H₂₇NO₇ 369.414

Alkaloid from *Senecio hadiensis* (Asteraceae). Powder (MeOH). Mp 207-209°. $[\alpha]_D^{20} -60$ (c, 0.5 in MeOH).

(15E)-Isomer: Neoplatyphylline

[20361-76-6]

C₁₈H₂₇NO₅ 337.415

Alkaloid from *Senecio platyphyllus* and *Senecio rhombifolius* (Asteraceae). Mp 131-133°. $[\alpha]_D^{20} +1.95$.

(15E)-Isomer, N-oxide: Neoplatyphylline N-oxide

[153323-56-9]

C₁₈H₂₇NO₆ 353.414

Alkaloid from roots of *Cacalia hupehensis* (Asteraceae).

(15E)-Isomer, 1 α -hydroxy, 12-Ac: 12-O-Acetylneohadiensine

C₂₀H₂₉NO₇ 395.452

Alkaloid from *Senecio hadiensis* (Asteraceae). Pale yellow oil.

12-Epimer: Ligularinine

[90364-90-2]

C₁₈H₂₇NO₅ 337.415

Alkaloid from the roots of *Ligularia dentata* (Asteraceae). Prisms (Et₂O). Mp 103-104°. $[\alpha]_D^{24} -88$ (c, 0.82 in CHCl₃).

[1257-59-6]

Richardson, M.F. *et al.*, *J.C.S.*, 1943, 452 (*isol, Hygrophylline*)

Koekemoer, M.J. *et al.*, *J.C.S.*, 1951, 66; 1955, 63 (*isol, oxide, struct*)

Warren, F.L. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1955, **12**, 198 (*bibl, Platyphylline*)

Schlosser, F.D. *et al.*, *J.C.S.*, 1965, 5707 (*isol, struct, Hygrophylline*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1968, **21**, 1671 (*Neoplatyphylline*)

Aasen, A.J. *et al.*, *J.O.C.*, 1969, **34**, 4137 (*struct, pmr*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (*cd*)

Abdullaev, U.A. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 538 (*ms*)

Drewes, S.E. *et al.*, *J.C.S. Perkin 1*, 1981, 287 (*cmr*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)

Molyneux, R.J. *et al.*, *Phytochemistry*, 1982, **21**, 439 (*cmr*)

Asada, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 475 (*Ligularinine*)

Mackay, M.F. *et al.*, *Acta Cryst. C*, 1985, **41**, 395 (*cryst struct, Hygrophylline*)

Pestchanker, M.J. *et al.*, *Planta Med.*, 1985, 165 (*Dihydroretorsine*)

Mattocks, A.R. *et al.*, *Chemistry and Toxicology of Pyrrolizidine Alkaloids*, Academic Press, 1986, 318 (*pharmacol*)

Were, O. *et al.*, *J. Nat. Prod.*, 1991, **54**, 491; 1415 (*Hadiensine, Pettitanine*)


Miao, Z. *et al.*, *Bopuxue Zazhi*, 1993, **10**, 55; *CA*, **120**, 164605y (*Neoplatyphylline N-oxide*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 431

Plectrodorine

P-517

[123085-04-1]



Relative configuration

C₁₁H₁₃NO₄ 223.228

Alkaloid from aerial parts of *Plectronia odorata* (preferred genus name *Olinia*) (Oliniaceae). Noncryst. $[\alpha]_D^{20} 0$ (c, 1 in MeOH).

5-Epimer: Isopectrodorine

[123164-26-1]

C₁₁H₁₃NO₄ 223.228

Alkaloid from aerial parts of *Plectronia odorata* (Oliniaceae). Noncryst. $[\alpha]_D^{20} 0$ (c, 1 in MeOH). Rel. config.


Gournelis, D. *et al.*, *J. Nat. Prod.*, 1989, **52**, 306-316 (*isol, uv, ir, pmr, ms*)

Ohba, M. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 63-67 (*synth, bibl*)

Pleiocarpamine

P-518

Methyl 19,20-didehydro-1,16-cyclocorynan-17-oate, 9CI
[6393-66-4]



C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Pleiocarpa mutica*, *Pleiocarpa pycnantha*, *Pleiocarpa tubicina*, *Kopsia dasyrachis* some *Alstonia* spp. and some other spp. in Apocynaceae. Also from *Hunteria eburnea* bark. Cryst. (Et₂O). Mp 164-165°. $[\alpha]_D^{20} +152.5$ (c, 0.04 in CHCl₃). pK_a 7.3. λ_{max} 230 (ε 29600); 285 (ε 8200); 294 (sh) (ε 6700) (EtOH).

N^d-Me: N-Methylpleiocarpamine

[2574-48-3]

C₂₁H₂₅N₂O₂ 337.441

Alkaloid from *Hunteria eburnea* (Apocynaceae). Mp 242-243° dec. (as

chloride). $[\alpha]_D^{25} +165$ (c, 0.5 in MeOH aq.). λ_{\max} 223 (log ϵ 4.47); 274 (log ϵ 3.92); 283 (sh) (log ϵ 3.87); 292 (sh) (log ϵ 3.75) (EtOH).

11-Hydroxy: 11-Hydroxypleiocarpamine

[32101-16-9]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Vinca erecta* (Apocynaceae). Cryst. (Me₂CO). Mp 228-229°. $[\alpha]_D^{30} +71$ (c, 0.7 in MeOH). λ_{\max} 233 (log ϵ 4.25); 285 (log ϵ 3.64) (no solvent reported).

16-Epimer, N^d-oxide: Epipleiocarpamine N^d-oxide

[69930-64-9]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the leaves of *Vinca minor* (Apocynaceae). Amorph.; cryst. (EtOH) (as tartrate salt). Mp 150-153° (tartrate). $[\alpha]_D^{23} +4.5$ (c, 1 in MeOH) (tartrate).

16-Hydroxymethyl: 16-Hydroxymethylpleiocarpamine

[180869-66-3]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from stem bark of *Kopsia deverrei*. $[\alpha]_D^{20} +115$ (c, 0.35 in CHCl₃). λ_{\max} 227 (log ϵ 4.29); 284 (log ϵ 3.79) (EtOH).

Kump, W.G. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1503-1516 (*isol, uv*)

Bartlett, M.F. *et al.*, *J.O.C.*, 1963, **28**, 2197-2199 (*isol, uv*)

Hesse, M. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 878-911 (*uv, ir, pmr, ms, struct*)

Il'yasova, Kh.T. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 717-719; *Chem. Nat. Compd. (Engl. Transl.)*, 728-729 (*11-Hydroxypleiocarpamine*)

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1974, **52**, 2327-2330 (*N-Methylpleiocarpamine*)

Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 123-127 (*Epipleiocarpamine N-oxide*)


Kan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 275-281 (*16-Hydroxymethylpleiocarpamine*)

Keawpradab, N. *et al.*, *Planta Med.*, 1997, **63**, 97-101 (*isol, pmr, cmr*)

Pleiocorine, 9CI

[55732-60-0]

P-519

C₄₁H₄₆N₄O₅ 674.838

Alkaloid from the stems and leaves of *Alstonia deplanchei* and from the leaves of *Alstonia odontophora* (Apocynaceae). $[\alpha]_D^{25} +142$ (c, 1.04 in CHCl₃). Dec. slowly above 300° without melting.

N-De-Me: N-Demethylpleiocorine

[74045-65-1]

C₄₀H₄₄N₄O₅ 660.811

Isol. from the leaves of *Alstonia odontophora* (Apocynaceae). Amorph. $[\alpha]_D^{25} +80$ (c, 1.0 in CHCl₃).


Das, B.C. *et al.*, *Tet. Lett.*, 1974, 4299 (*uv, ir, pmr, cmr, ms, struct*)

Vercauteren, J. *et al.*, *Phytochemistry*, 1979, **18**, 1729 (*deriv*)

Pleioacraline, 9CI

[62509-84-6]

P-520

C₄₁H₄₆N₄O₅ 674.838

Alkaloid from *Alstonia deplanchei* and the leaves of *Alstonia odontophora* (Apocynaceae). Plates (MeOH). $[\alpha]_D^{20} +124$ (c, 1 in CHCl₃). Dec. >300°. Stereochem. of interunit link not proven.

Das, B.C. *et al.*, *J.O.C.*, 1977, **42**, 2785-2786 (*ir, uv, pmr, cmr, ms, struct*)


Vercauteren, J. *et al.*, *Phytochemistry*, 1979, **18**, 1729 (*isol*)

Pleiomutine

Methyl 15-(14,15-dihydroeburnamenin-14-yl)-1-methylaspidofractinine-3-carboxylate, 9CI

[5263-34-3]

P-521



Absolute Configuration

C₄₁H₅₀N₄O₂ 630.872

Minor alkaloid from *Pleiocarpa mutica* and from *Kopsia dasyrachis*. Cryst. (MeOH/CHCl₃). Mp 225° dec. $[\alpha]_D^{26} -111$ (c, 1.93 in CHCl₃). Thomas and Lavaud show the wrong abs. config. of the pleiocarpine half of the molecule. λ_{\max} 230 (log ϵ 4.5); 263 (log ϵ 4.2); 285 (log ϵ 3.97); 292 (log ϵ 3.97) (MeOH).

Dipicrate:

Cryst. (2-propanol/Me₂CO or Me₂CO/Et₂O). Dec. >230° without melting.

N-De-Me: Norpleiomutine. Kopsoffine

[82529-52-0]

[89783-67-5]

C₄₀H₄₈N₄O₂ 616.845

Alkaloid from the stem bark of

Hunteria zeylanica and from *Kopsia dasyrachis* and *Kopsia officinalis* (Apocynaceae). Amorph. Mp 240°. $[\alpha]_D^{25} -65$ (c, 0.5 in CHCl₃). $[\alpha]_D^{25} +4$ (c, 1.1 in CHCl₃). MP. and lower opt. rotn. refer to Kopsoffine. Higher opt. rotn. refers to Norpleiomutine, descr. as amorph.

N-De-Me, parent acid: [96935-24-9]C₃₉H₄₆N₄O₂ 602.818

Alkaloid from the stem bark of *Kopsia pauciflora* (Apocynaceae). Cryst. (Me₂CO). Mp 260°. $[\alpha]_D^{20} -130$ (c, 0.54 in CHCl₃).

19'ξ-Hydroxy, N-de-Me: Kopsoffinol

[96935-25-0]

C₄₀H₄₈N₄O₃ 632.844

Alkaloid from the stem-bark of *Kopsia pauciflora* and *Kopsia dasyrachis* (Apocynaceae). Cryst. (Me₂CO). Mp 250°. $[\alpha]_D^{20} +21.1$ (c, 1 in CHCl₃).

Kump, W.G. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1503-1516 (*isol, uv*)

Hesse, M. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 964-974 (*ms, struct, synth*)

Thomas, D.W. *et al.*, *J.A.C.S.*, 1966, **88**, 1537-1544 (*isol, uv, ir, pmr, ms, struct, synth*)

Lavaud, C. *et al.*, *Phytochemistry*, 1982, **21**, 445-447 (*isol, uv, ir, pmr, ms, struct, Norpleiomutine*)

Feng, X.Z. *et al.*, *J. Nat. Prod.*, 1984, **47**, 117-122 (*Kopsoffine*)

Magnus, P. *et al.*, *Chem. Comm.*, 1985, 184-188 (*Norpleiomutine*)


Kan-Fan, C. *et al.*, *J. Nat. Prod.*, 1985, **48**, 124-127 (*Kopsoffinol, N-de-Me parent acid*)

Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 159-169 (*Kopsoffine, isol, pmr, cmr*)

Pleurospermine

P-522

1-(4-Hydroxy-3-methoxyphenyl)-2-(2-piperidinyl)ethanone, 9CI. 4'-Hydroxy-3'-methoxy-2-(2-piperidinyl)acetophenone, 8CI. 2-(4-Hydroxy-3-methoxyphenacyl)-piperidine
[7363-20-4]

C₁₄H₁₉NO₃ 249.309

Alkaloid from leaves of *Cryptocarya pleurosperma* (Lauraceae). Prob. a biosynth. precursor of Cryptopleurine, the bark alkaloid. Cryst. (H₂O). Mp 177° dec. $[\alpha]_D^{25} +44$. pK_a 8 (50% EtOH aq.). The positive opt. rotn. reported could not be substantiated. On reexamination, the original isolate was found to be racemic, and only racemic Pleurospermine could be obt. from fresh plant. This could however, be a case of v. ready racemisation (cf. Pelletierine, P-173).

Hydrochloride:

Cryst. (EtOH). Mp 219-220°.

Picrate:

Cryst. (Me₂CO). Mp 166° dec. Dec. from 150°.

Me ether: 3,4-Dimethoxy-ω-(2-piperidyl)acetophenone

[23837-54-9]

C₁₅H₂₁NO₃ 263.336

Alkaloid from *Boehmeria cylindrica* var. *drummondii*, *Boehmeria platyphyllo* and *Boehmeria cylindrica* (Urticaceae). Shows strong activity against *Candida albicans*. Needles (Me₂CO). Poorly sol. hexane. Mp 81-82°. Opt. inactive.

2'-Methoxy: 4'-Hydroxy-2',3'-dimethoxy-2-(2-piperidyl)acetophenone. **Caulophyllumine A**

C₁₅H₂₁NO₄ 279.335

Alkaloid from the roots of *Caulophyllum thalictroides* (blue cohosh). Brown powder. [α]_D²⁸ -11.6 (c, 0.27 in MeOH). Possesses (S)-config. λ_{max} 280 (log ε 3.04) (MeOH).

Demethoxy, N-Me: 4'-Hydroxy-2-(1-methylpiperidin-2-yl)acetophenone.

Thalictroidine

[248259-06-5]

C₁₄H₁₉NO₂ 233.31

Alkaloid from the rhizomes of *Caulophyllum thalictroides* (blue cohosh). Glassy solid.

Gellert, E. et al., *Aust. J. Chem.*, 1959, **12**, 90-96 (*isol, ir*)

Loder, J.W. et al., *Aust. J. Chem.*, 1962, **15**, 296-300 (*synth, struct*)

Hart, N.K. et al., *Aust. J. Chem.*, 1968, **21**, 1397-1398; 1969, **22**, 1805-1807 (*Me ether, isol, pmr, struct*)


Al-Shamma, A. et al., *Phytochemistry*, 1982, **21**, 485-487 (*Me ether, isol, pharmacol*)

Kennelly, E.J. et al., *J. Nat. Prod.*, 1999, **62**, 1385-1389 (*Thalictroidine*)

Ali, Z. et al., *Phytochemistry*, 2008, **69**, 1037-1042 (*Caulophyllumine A*)

Pleurostyline P-523

2,3,6,7,8,9,10,11-Octahydro-6-phenyl-1H-[1,5,9]triazacyclotridecino[2,1-a][2]benzazepine-4,13(5H,19bH)-dione, 9CI [67257-79-8]



Probable Absolute Configuration

C₂₅H₂₉N₃O₂ 403.523

Alkaloid from *Pleurostyliya africana* (Celastraceae). Mp 246-247°. [α]_D²⁵ -194 (c, 0.825 in CHCl₃).

7R-Hydroxy: 7-Hydroxypleurostyline

[145199-53-7]

C₂₅H₂₉N₃O₃ 419.522

Minor alkaloid from leaves of *Pleurostyliya opposita* (Celastraceae). Amorph. [α]_D -110 (c, 1 in CHCl₃).

7',8'-Dihydro, 7'-hydroxy: 7'-Hydroxy-

7',8'-dihydropleurostyline

[145199-55-9]

C₂₅H₃₁N₃O₃ 421.538

Minor alkaloid from leaves of *Pleurostyliya opposita* (Celastraceae). Amorph. [α]_D 0 (c, 1 in CHCl₃).


Wagner, H. et al., *Helv. Chim. Acta*, 1981, **64**, 283 (*uv, ir, pmr, cmr, ms, struct*)

Séguineau, C. et al., *Helv. Chim. Acta*, 1992, **75**, 2283 (7-Hydroxypleurostyline, 7'-Hydroxy-7',8'-dihydropleurostyline)

Khanjin, N.A. et al., *Helv. Chim. Acta*, 2001, **84**, 1253 (*abs config*)

Plicamine

[232597-76-1]

C₂₆H₂₆N₂O₆ 462.501


Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. [α]_D +74.4 (c, 0.12 in MeOH). λ_{max} 226 (log ε 4.41); 265 (sh) (log ε 3.71); 299 (log ε 3.49) (MeOH).

Unver, N. et al., *Phytochemistry*, 1999, **50**, 1255-1261 (*isol, uv, ir, cd, pmr, cmr*)

Baxendale, I.R. et al., *Tetrahedron*, 2002, **58**, 6285-6304 (*synth*)

Plicane

[343983-02-8]


C₁₈H₁₈N₂O₄ 326.351

Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. [α]_D +205.9 (c, 0.12 in MeOH). λ_{max} 231 (log ε 4.3); 285 (log ε 3.35); 319 (log ε 3.18) (MeOH).

Unver, N. et al., *Heterocycles*, 2001, **55**, 641-652 (*isol, pmr, cmr, uv*)

Plumerinine

Octahydro-6,9-dimethyl-4-(1-methylethyl)-2H-quinolizin-2-ol, 9CI. Octahydro-8-hydroxy-6-isopropyl-1,4-dimethyl-quinolizine [125797-64-0]

C₁₄H₂₇NO 225.373

Struct. in doubt after 2002 synthesis. Alkaloid from the stems of *Plumeria rubra* (Apocynaceae). Light brown vis-


cous oil. [α]_D²⁰ +14.4 (c, 0.31 in MeOH).

Najam-ul-Hussain Kazmi, S. et al., *Heterocycles*, 1989, **29**, 1901 (*isol, uv, ir, pmr, cmr, ms, struct*)

Comins, D.L. et al., *Org. Lett.*, 2002, **4**, 1611-1613 (*synth*)

Plumocraline

[79123-75-4]

C₄₂H₅₀N₄O₅ 690.881


Alkaloid from the root bark of *Alstonia plumosa* (Apocynaceae). Cryst. (CHCl₃). [α]_D +20 (c, 1 in CHCl₃). Dec. above 300° without melting.

Massiot, G. et al., *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1981, **292**, 191 (*uv, ir, pmr, cmr, ms, struct*)

Jacquier, M.J. et al., *Phytochemistry*, 1982, **21**, 2973 (*isol, uv, ir, pmr, ms, struct*)

Pluviine

[548-11-8]



Absolute Configuration

C₁₇H₂₁NO₃ 287.358

Alkaloid from *Lycoris radiata*, several *Narcissus* spp. and some other spp. in the Amaryllidaceae. Mp 225-227°. [α]_D -171 (c, 0.18 in EtOH).

Perchlorate: Mp 260° dec.

Ac: Mp 184°.

O⁹-De-Me: 9-Norpluviine

[517-99-7]

C₁₆H₁₉NO₃ 273.331

Isol. from *Lycoris radiata*, "Texas" daffodils and *Pancreatium maritimum* (Amaryllidaceae). Plates (MeOH). Mp 239-241° (274-275°) dec. [α]_D -232 (c, 0.056 in MeOH). [α]_D -160 (c, 0.15 in MeOH).

O⁹-De-Me, O¹-Ac: 1-O-Acetyl-9-norpluviineC₁₈H₂₁NO₄ 315.368

Alkaloid from the bulbs of *Brunsvigia radulosa* and *Ammocharis coranica*. Amorph. powder (Me₂CO/hexane). Mp 185-187° (173°). [α]_D²² -67 (c, 0.25 in EtOH).

O⁹-De-Me, di-Ac:

Cubes (petrol). Mp 150-151°.

O¹⁰-De-Me: 10-Norpluviine

[162411-72-5]

C₁₆H₁₉NO₃ 273.331

Alkaloid from bulbs of *Narcissus pseudonarcissus* ssp. *pseudonarcissus* cv. Carlton. Mp 148° (dec. >136°). [α]_D²⁵

+61 (c, 2.4 in MeOH).

O¹⁰-De-Me, O¹-Ac: 1-O-Acetyl-10-norpluviine
[162411-71-4]
C₁₈H₂₁NO₄ 315.368
From bulbs of *Narcissus pseudonarcissus* ssp. *pseudonarcissus* cv. Carlton. Mp 203° (dec. >180°). [α]_D²⁵ +38 (c, 1.0 in MeOH).

O¹⁰-De-Me, di-Ac: 1,10-Diacetyl-10-norpluviine
[170172-02-8]
C₂₀H₂₃NO₅ 357.405
From bulbs of *Narcissus pseudonarcissus* ssp. *pseudonarcissus* cv. Carlton. Cryst. (EtOH). Mp 147°. [α]_D²⁵ +8.3 (c, 1.1 in MeOH).

16-Epimer, O⁹-de-Me: Fortucine
[115730-50-2]
C₁₆H₁₉NO₃ 273.331
Alkaloid from the leaves of *Narcissus* sp. Mp 160-162°. [α]_D +66.7 (c, 0.23 in EtOH).

16-Epimer, O⁹-de-Me; hydrochloride: Mp 208-209°.

A^{4,12}-Isomer, O⁹-de-Me: Kirkine
C₁₆H₁₉NO₃ 273.331
Alkaloid from the bulbs of *Crinum kirkii*. Mp 170-172°. [α]_D²⁵ +59.6 (c, 0.57 in MeOH). Struct. revised in 2008, originally assigned same struct. as Fortucine.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 363 (uv, ir, struct)

Fales, H.M. *et al.*, *J.A.C.S.*, 1958, 4395 (struct)

Uyeo, S. *et al.*, *J.C.S.*, 1959, 172 (deriv)

Kirby, G.W. *et al.*, *J.C.S.(C)*, 1966, 676 (deriv, biosynth, uv, pmr)

Sandberg, F. *et al.*, *Acta Pharm. Suec.*, 1968, **5**, 61; *CA*, **69**, 25043d (deriv)

Bruce, I.T. *et al.*, *Chem. Comm.*, 1968, 207 (deriv, biosynth)

Razakov, R. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 19; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 15 (ms)

Harken, R.D. *et al.*, *J.O.C.*, 1976, **41**, 2450 (biosynth)

Tokhtabaeva, G.M. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 872; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 727 (Fortucine)

Bastida, J. *et al.*, *Phytochemistry*, 1995, **40**, 1291-1293 (Kirkine)

Kreh, M. *et al.*, *Phytochemistry*, 1995, **40**, 1303 (10-Norpluviine, 1-O-Acetyl-10-norpluviine, 1,10-Diacetyl-10-norpluviine)


Campbell, W.E. *et al.*, *Phytochemistry*, 2000, **53**, 587-591 (1-O-Acetyl-9-norpluviine)

Koorbanally, N. *et al.*, *Phytochemistry*, 2000, **54**, 93-97 (1-Acetyl-9-norpluviine)

Biechy, A. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 1436-1438 (Fortucine, Kirkine, synth)

Podopetaline P-529

16,17-Didehydroormosanine, 9CI. Ormo-castrine
[38966-20-0]



Absolute configuration

C₂₀H₃₃N₃ 315.501

Ormo-castrine was actually identical with Podopetaline hydrochloride. Major alkaloid from *Podopetalum ormondii* (preferred genus name *Ormosia*), also isol. from *Ormosia semicastrata* (Fabaceae). Mp 77.5-79°. [α]_D -48 (c, 0.95 in MeOH).

Hydrobromide: Mp 253-256°.

6-Epimer: see Epipodopetaline, E-99

Hart, N.K. *et al.*, *Tet. Lett.*, 1972, 5333 (isol)

McLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**,

1639; 1974, **52**, 1907 (isol, struct)

Mackay, M.F. *et al.*, *Tetrahedron*, 1975, **31**,

1295 (cryst struct)

Poecillanosine P-530

2-Acetoxy-1-(hydroxynitrosoamino)heptadecane
[200564-72-3]

H₃C(CH₂)₁₄CH(OAc)CH₂N(OH)NO

C₁₉H₃₈N₂O₄ 358.52

Isol. from the sponge *Poecillastra* spec. aff. *tenuilaminaris*. Cytotoxic agent and free radical scavenger. Solid. [α]_D²⁵ -20.2 (c, 0.1 in MeOH). Ref. states abs. config. as (R)- in text but shows (S)- in diag. λ_{max} 232 (ε 5700) (MeOH).

Natori, T. *et al.*, *Tet. Lett.*, 1997, **38**, 8349-8350 (isol, uv, ir, pmr, cmr)

Xian, M. *et al.*, *Tet. Lett.*, 2007, **48**, 1209-1212 (synth)

Poetamine P-531

[1360-49-2]

C₂₉H₃₂N₂O₇ 520.581

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Narcissus poeticus* var. *ornatus* (Amaryllidaceae). Prisms (MeOH). Mp 258-260° dec. [α]_D²⁴ -160 (c, 0.2 in EtOH).

Perchlorate: Mp 248° dec.

Picrate: Mp 246° dec.

Methiodide: Mp 191° dec.

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 354 (isol, ir)

Poetaricine P-532

C₁₆H₁₇NO₄ 287.315

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Narcissus poeticus* var. *ornatus* (Amaryllidaceae). Prisms (MeOH). Mp 273° dec. [α]_D²³ -60 (c, 0.2 in EtOH).

Perchlorate:

Prisms (H₂O). Mp 228-230° dec.

Picrate:

Fine needles (MeOH). Mp 190° dec.

Ac:

Prisms (MeOH). Mp 165°.

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 595 (isol, ir)

Poeticine P-533

C₂₀H₂₃NO₆ 373.405

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Narcissus poeticus* (Amaryllidaceae). Needles (Me₂CO). Mp 209-210°. [α]_D²⁵ -89 (c, 0.3

in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 2462-2465 (isol)

Pogonopamine P-534

Struct. unknown. No analytical data recorded. Alkaloid from the bark of *Pogonopus tubulosis* (Rubiaceae).

Dalma, G. *et al.*, *CA*, 1949, **43**, 5548i

Pogonopeine P-535

Struct. unknown. No analytical data recorded. Alkaloid from the bark of *Pogonopus tubulosis* (Rubiaceae).

Dalma, G. *et al.*, *CA*, 1949, **43**, 5548i

Pogonopidine P-536

Struct. unknown. No analytical data recorded. Alkaloid from the bark of *Pogonopus tubulosis* (Rubiaceae).

Dalma, G. *et al.*, *CA*, 1949, **43**, 5548i


Pogonopine P-537

Struct. unknown. No analytical data recorded. Alkaloid from the bark of *Pogonopus tubulosis* (Rubiaceae).

Dalma, G. *et al.*, *CA*, 1949, **43**, 5548i

Pohakuline P-538

[60394-95-8]



Relative configuration


C₁₅H₂₆N₂O₂ 266.383

Trace alkaloid from the bark of *Sophora chrysophylla* (Fabaceae). Cryst. (Me₂CO). Mp 170-171°. [α]_D²⁵ -17.2 (c, 0.82 in EtOH).

Kadooka, M.M. *et al.*, *Tetrahedron*, 1976, **32**, 919 (isol, ir, pmr, ms, cryst struct)

Polonicumtoxin C P-539

[120693-76-7]



C₁₁H₁₉NO 181.277

Alkaloid from cultured cells of the freshwater dinoflagellate *Peridinium polonicum*. Ichthyotoxin. Ichthyotoxin.

▶ LD₅₀ (mus, ipr) 2 mg/kg.

O-Ac: Polonicumtoxin B

[120693-75-6]

C₁₃H₂₁NO₂ 223.314

From *Peridinium polonicum*. Ichthyotoxin.

O-(3-Butenoyl): Polonicumtoxin A

[120693-74-5]

C₁₅H₂₃NO₂ 249.352


From *Peridinium polonicum*. Ichthyotoxin.

▶ LD₅₀ (mus, ipr) 1.5 mg/kg.

Oshima, Y. *et al.*, *CA*, 1989, **110**, 209003n
 Yotsu-Yamashita, M. *et al.*, *Heterocycles*,
 1998, **48**, 79-93 (*synth*, *pmr*, *cmr*)
 Van, T.N. *et al.*, *Tetrahedron*, 2000, **56**, 7969-
 7973 (*synth*)

Polyalthenol**P-540**

[62121-09-9]


C₂₃H₃₁NO 337.504

Alkaloid from *Greenwayodendron oliveri*
 (*Polyalthia oliveri*) (Annonaceae). Mp
 149-150°. [α]_D +50 (c, 0.4 in EtOH). λ_{\max}
 226 (ε 33100); 285 (ε 5750); 292 (ε 5370)
 (EtOH) (Derep).

Leboeuf, M. *et al.*, *Tet. Lett.*, 1976, 3559 (*uv*,
pmr, *cmr*, *ms*, *struct*)

Polyandrocarpamide A**P-541**

N-[2-(3-Bromo-4-hydroxyphenyl)ethyl]-
 α -oxo-1H-indole-3-acetamide, 9CI
 [129369-41-1]

C₁₈H₁₅BrN₂O₃ 387.232

Alkaloid from the marine ascidian
Polyandrocarpa sp. Needles. Mp 178-
 179°. λ_{\max} 207 (ε 58500); 247 (ε 1600);
 266 (ε 9900); 272 (ε 8600); 309 (ε 9800)
 (MeOH/NaOH) (Derep). λ_{\max} 205 (ε
 41400); 255 (ε 10300); 267 (ε 9900); 274
 (ε 9600); 290 (ε 5000); 324 (ε 8400)
 (MeOH) (Derep).

Debromo: Polyandrocarpamide C

[107610-00-4]

C₁₈H₁₆N₂O₃ 308.336

Alkaloid from *Polyandrocarpa* sp.
 Amorph. solid. λ_{\max} 205 (ε 21400);
 246 (ε 8300); 266 (ε 6100); 272 (ε
 5300); 318 (ε 4600) (MeOH/NaOH)
 (Derep). λ_{\max} 205 (ε 19200); 254
 (ε 6400); 266 (ε 6100); 272 (ε 5700);
 285 (sh); 324 (ε 4600) (MeOH)
 (Derep).

3'-Debromo, 3'-iodo: Polyandrocarpamide B

[129369-42-2]

C₁₈H₁₅IN₂O₃ 434.233


Alkaloid from *Polyandrocarpa* sp. λ_{\max}
 203 (ε 79100); 247 (ε 13200); 267 (sh);
 274 (ε 7600); 311 (ε 8200) (MeOH/
 NaOH) (Derep). λ_{\max} 205 (ε 50000);
 230 (sh); 254 (ε 9100); 267 (sh); 274
 (ε 8200); 325 (ε 6600) (MeOH)
 (Derep).

Da Settimo, A. *et al.*, *Farmaco, Ed. Sci.*, 1987,
42, 17-26 (*synth*)

Lindquist, N. *et al.*, *Tet. Lett.*, 1990, **31**, 2521-
 2524 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Polyandrocarpamide D**P-542**

5-(1H-Indol-3-yl)-5-methoxy-2,4-imida-
 zolidinedione, 9CI

C₁₂H₁₁N₃O₃ 245.237**(±)-form [129369-43-3]**


Alkaloid from the marine ascidian
Polyandrocarpa sp. and from the
 sponge *Zyza massalis*. Exhibits anti-
 bacterial and antifungal activity. Yellow
 amorph. solid. Artifact. λ_{\max} 212 (ε
 27500); 268 (ε 5000); 276 (ε 400); 287 (ε
 3800) (MeOH) (Derep). λ_{\max} 204 (ε
 44800); 217 (sh); 268 (ε 4600); 277 (ε
 4600); 289 (ε 3700) (MeOH/NaOH)
 (Derep).

Lindquist, N. *et al.*, *Tet. Lett.*, 1990, **31**, 2521-
 2524 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Mancini, I. *et al.*, *Helv. Chim. Acta*, 1994, **77**,
 1886-1894 (*isol*, *pmr*, *cmr*)

Polyandrocarpamine A**P-543**

[444106-74-5]

C₁₁H₁₁N₃O₃ 233.226

Isol. from the Fijian ascidian *Polyandro-*
carpa sp. Yellow solid. λ_{\max} 246 (ε 2000);
 352 (ε 2000) (MeOH).

O-De-Me: Polyandrocarpamine B

[444106-75-6]


C₁₀H₉N₃O₃ 219.199

Isol. from *Polyandrocarpa* sp. Yellow
 solid. λ_{\max} 248 (ε 2000); 352 (ε 2000)
 (MeOH).

Davis, R.A. *et al.*, *Tetrahedron*, 2002, **58**, 3263-
 3269 (*isol*, *synth*, *uv*, *ir*, *pmr*, *cmr*)

Polyandrocarpidine A**P-544**

[5-[2-(2,5-Octadienylidene)-5-oxo-1-pyr-
 rolidinyl]pentyl]guanidine, 9CI
 [68838-36-8]



n = 5

C₁₈H₃₀N₄O 318.461

The polyandrocarpidines constitute a
 9:1 mixt. of homologues, each
 homologue being a mixt. of isomers
 with isomer ratios varying from 4:1
 to 1:1. Polyandrocarpidines A and B
 correspond to the former
 Polyandrocarpidine I, and the minor
 polyandrocarpidines C and D to the
 former Polyandrocarpidine II. Isol.
 from a marine tunicate *Polyandrocarpa*
 sp. Shows antimicrobial and
 cytotoxic props. λ_{\max} 275 (MeOH)
 (Berdy).

(2'E)-Isomer: Polyandrocarpidine B

[84453-26-9]

C₁₈H₃₀N₄O 318.461

From *Polyandrocarpa* sp. Shows anti-
 microbial and cytotoxic props. Sol.
 MeOH, CHCl₃; poorly sol. H₂O,
 hexane. λ_{\max} 276 (ε 1000) (MeOH)
 (Berdy).

Cheng, M.T. *et al.*, *J.A.C.S.*, 1978, **100**, 7409-
 7411 (*isol*)

Carté, B. *et al.*, *Tet. Lett.*, 1982, **23**, 3863-3866
 (*isol*, *uv*, *pmr*, *cmr*, *struct*)

Rinehart, K.L. *et al.*, *Tet. Lett.*, 1983, **24**,
 1593-1596 (*struct*)

Polyandrocarpidine C**P-545**

[4-[2-(2,5-Octadienylidene)-5-oxo-1-pyr-
 rolidinyl]butyl]guanidine, 9CI
 [68838-37-9]

As Polyandrocarpidine A, P-544 with
 n = 4

C₁₇H₂₈N₄O 304.434

See note under Polyandrocarpidine A,
 P-544. Constit. of a marine tunicate
Polyandrocarpa sp. Shows
 antimicrobial and cytotoxic props.
 Sol. MeOH, CHCl₃; poorly sol.
 H₂O, hexane. λ_{\max} 275 (MeOH) (Ber-
 dy).

(2'E)-Isomer: Polyandrocarpidine D

[84453-27-0]

C₁₇H₂₈N₄O 304.434

From *Polyandrocarpa* sp. Shows anti-
 microbial and cytotoxic props. Sol.
 MeOH, CHCl₃; poorly sol.
 H₂O, hexane. λ_{\max} 275 (MeOH)
 (Berdy).


Cheng, M.T. *et al.*, *J.A.C.S.*, 1978, **100**, 7409-
 7411 (*isol*)

Carté, B. *et al.*, *Tet. Lett.*, 1982, **23**, 3863-3866
 (*isol*, *uv*, *pmr*, *cmr*, *struct*)

Rinehart, K.L. *et al.*, *Tet. Lett.*, 1983, **24**,
 1593-1596 (*struct*)

Polyavolinamide**P-546**

[81525-54-4]

C₂₅H₃₅NO₂ 381.557Relative
Configuration


Alkaloid from the root bark and stem of *Greenwayodendron suaveolens* (*Polyalthia suaveolens*) (Annonaceae). Cryst. (CHCl₃/MeOH). Mp 249-251°. λ_{max} 217 (log ε 4.1); 255 (log ε 4.02); 285 (log ε 3.42); 293 (log ε 3.34) (MeOH). λ_{max} 262 (log ε 3.89); 285 (log ε 3.48); 293 (log ε 3.4) (MeOH/NaOH).

Ac:

Cryst. (CHCl₃/MeOH). Mp 246-248°. λ_{max} 222 (log ε 3.66); 261 (log ε 3.94); 282 (log ε 3.76); 290 (log ε 3.68) (MeOH).

Okorie, D.A. *et al.*, *Phytochemistry*, 1981, **20**, 2575-2578 (*isol, uv, ir, pmr, ms, struct*)

***Epilachna borealis* Polyazamacrolides P-547**



m, n, p, = 5, 6 or 7


A large family of lactonic ring structures derived from a small group of (2-hydroxyethylamino)alkanoic acids. Example shown contains 3 acids. Constit. of the pupal defensive secretion of the coccinellid beetle *Epilachna borealis*. Cited as a case of natural combinatorial chemistry.

Schroeder, F.C. *et al.*, *Science (Washington, D.C.)*, 1998, **281**, 428-431 (*isol, struct*)

Schroeder, F.C. *et al.*, *Tet. Lett.*, 1998, **39**, 6625-6628 (*abs config*)

Garcia-Rubio, S. *et al.*, *J.O.C.*, 2001, **66**, 1082-1096 (*synth*)

Polyberbine P-548
[73777-77-2]




C₂₀H₁₉NO₆ 369.373

Alkaloid from *Berberis valdiviana* (Berberidaceae). Amorph.

Murugesan, N. *et al.*, *Tet. Lett.*, 1979, 4521 (*synth, uv, ir, pmr, ms*)

Firdous, S. *et al.*, *J.A.C.S.*, 1984, **106**, 6099 (*occur, uv, ir, pmr, ms*)

Polycanthidine P-549
[155416-31-2]




C₂₃H₂₇NO₄ 381.471

Alkaloid from *Astragalus polycanthus* (Fabaceae).

Gupta, R.K. *et al.*, *Indian Drugs*, 1993, **30**, 595; *CA*, **120**, 319334k (*isol, struct*)

Polycanthisine P-550
Alkaloid APAL-VII
[161068-62-8]




C₁₃H₂₁NO 207.315

Alkaloid from aerial parts of *Astragalus polycanthus* (Fabaceae). Cryst. Mp 160-161°. [α]_D³⁰ +16 (c, 0.75 in MeOH). Enol, with OH group strongly H-bonded to N. Is not acetylated under usual conditions.

Gupta, R.K. *et al.*, *Indian Drugs*, 1993, **30**, 651 (*isol*)

Gupta, R.K. *et al.*, *Indian J. Chem., Sect. B*, 1995, **34**, 76 (*uv, ir, pmr, ms, struct*)

Polycarpaurine B P-551
[921987-38-4]



R = -SO₃H

C₁₁H₁₃N₃O₄S 283.307

Related to Polycarpine, P-553. Isolated from *Polycarpa aurata*. Powder. λ_{max} 222 (log ε 4.51); 261 (log ε 4.38) (MeOH).

Wang, W. *et al.*, *Tetrahedron*, 2007, **63**, 409-412 (*isol, pmr, cmr, ms*)

Polycarpaurine C P-552
[921987-39-5]

As Polycarpaurine B, P-551 with


R = -S-SO₃H

C₁₁H₁₃N₃O₄S₂ 315.373

Isolated from *Polycarpa aurata*. Pale yellow powder. λ_{max} 242 (log ε 4.27); 300 (log ε 4.12) (MeOH).

Wang, W. *et al.*, *Tetrahedron*, 2007, **63**, 409-412 (*isol, pmr, cmr, ms*)

Polycarpine† P-553
[175669-17-7]



C₂₂H₂₄N₆O₂S₂ 468.603

Neutral form illus. The nat. prod. appears to be the dication. Alkaloid from the ascidians *Polycarpa clavata* and *Polycarpa aurata*. Cytotoxic against the human colon tumour cell line HCT-116. Red rods (MeOH/CH₂Cl₂) or orange amorph. solid. Mp 201-204°. λ_{max} 226 (ε 21820); 264 (ε 23708); 364 (ε 9323) (no solvent reported).

Hydrochloride (1:2): [175669-29-1]

Orange rods. Mp 201-203°. Delocalised charges within the two iminoimidazole residues. λ_{max} 202 (log ε 4.45); 259 (log ε 4.28); 394 (log ε 3.58) (MeOH).

Monosulfide analogue: Polycarpaurine A

C₂₂H₂₄N₆O₂S 436.537

Isolated from *Polycarpa aurata*. Amorph. yellow powder (as bistrifluoroacetate salt). λ_{max} 228 (log ε 4.59); 281 (log ε 4.52) (MeOH) (bistrifluoroacetate).


Abbas, S.A. *et al.*, *J.O.C.*, 1996, **61**, 2709-2712 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Kang, H. *et al.*, *Tet. Lett.*, 1996, **37**, 2369-2372 (*isol, uv, ir, pmr, cmr, struct*)

Radchenko, O.S. *et al.*, *Tet. Lett.*, 1997, **38**, 3581-3584 (*synth*)

Wang, W. *et al.*, *Tetrahedron*, 2007, **63**, 409-412 (*Polycarpaurine A*)

Polycarpine† P-554
3,4-Dihydro-1-[(2-hydroxy-3,4-dimethoxyphenyl)methylene]-6,7-dimethoxy-2(1H)-isoquinolinecarboxaldehyde, 9CI
[63490-92-6]



C₂₁H₂₃NO₆ 385.416


Alkaloid from *Enantia polycarpa* trunk bark (Annonaceae). Cryst. (MeOH). Sol. MeOH, H₂O. Mp 179-180°. Unique alkaloid which prob. arises in the plant by hydroperoxide attack on a protoberberinium precursor. λ_{max} 202 (ε 28183); 259 (ε 19054); 394 (ε 8710) (MeOH) (Berdy). λ_{max} 203 (ε 42660); 278 (ε 12300) (MeOH-NaOH) (Berdy).

Jössang, A. *et al.*, *C. R. Hebd. Seances Acad.*

Sci. Ser. C, 1977, **284**, 467 (isol, cryst struct, pmr)
Murugesan, N. *et al.*, *Tet. Lett.*, 1979, 4521
(synth, biosynth)

Polycephalin C

[220422-37-7]

C₃₂H₃₆N₂O₈ 576.645

Tetramic acid deriv. Information on abs. config. is confused. That shown is assigned by Blumenthal *et al.* Longbottom *et al.* have assigned the opposite abs. configs. at the cyclohexane ring positions. Both sets of authors call their configs. *R,R*-; Longbottom *et al.* appear to be correct but assignment of *R,S*-priorities is difficult. Isol. from plasmodia of the slime mold *Physarum polycephalum*. Orange-red powder. λ_{\max} 253 (log ϵ 4.15); 390 (log ϵ 4.74) (MeOH).

1-N-De-Me: Polycephalin B

[220422-47-9]


C₃₁H₃₄N₂O₈ 562.618

Isol. from *Physarum polycephalum*. Orange-red powder. λ_{\max} 247 (log ϵ 4.14); 386 (log ϵ 4.74); 406 (sh) (log ϵ 4.69) (MeOH).

Nowak, A. *et al.*, *Angew. Chem., Int. Ed.*, 1998, **37**, 3139-3141 (isol, ir, cd, pmr, cmr)
Blumenthal, F. *et al.*, *Tetrahedron*, 2002, **58**, 8433-8437 (cd, abs config)
Longbottom, D.A. *et al.*, *Tetrahedron*, 2003, **59**, 6955-6966 (synth, abs config)

Polycitone A

[153212-84-1]



P-556

C₃₈H₂₁Br₈NO₇ 1242.818

Alkaloid from the marine ascidian *Polycitor* sp. Inhibits retroviral reverse transcriptase and DNA polymerase. Fibroblast inhibitor. Yellowish needles + 1 Me₂CO (Me₂CO). Mp 285°. λ_{\max} 285 (ϵ 13500) (MeOH) (Berdy).

N-Dealkyl: Polycitone B

[272118-06-6]

C₃₀H₁₃Br₈NO₆ 1122.668Alkaloid from *Polycitor africanus*.

Yellow oil.

Rudi, A. *et al.*, *J.O.C.*, 1994, **59**, 999-1003 (isol, uv, ir, pmr, cmr, cryst struct)

Loya, S. *et al.*, *Biochem. J.*, 1999, **344**, 85-92 (activity)


Rudi, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 832-833 (*Polycitone B*)

Kreipl, A.T. *et al.*, *Org. Lett.*, 2002, **4**, 3287-3288 (synth)

Gupton, J.T. *et al.*, *Tetrahedron*, 2008, **64**, 5246-5253 (synth)

Polycitorol A

P-557



Relative Configuration


C₁₇H₃₁NO 265.438

Alkaloid from an ascidian of the family Polycitoridae. Oil. $[\alpha]_{\text{D}}^{25}$ +15.3 (c, 0.2 in CHCl₃).

Issa, H.H. *et al.*, *Mar. Drugs*, 2005, **3**, 78-85 (isol, pmr, cmr)

Polycitorol B

P-558



Relative Configuration


C₁₇H₃₁NO 265.438

Alkaloid from an ascidian of the family Polycitoridae. Oil. $[\alpha]_{\text{D}}^{25}$ -10.3 (c, 0.6 in CHCl₃).

Issa, H.H. *et al.*, *Mar. Drugs*, 2005, **3**, 78-85 (isol, pmr, cmr)

Polycitrin A

[153212-85-2]



P-559

C₂₄H₁₅Br₄NO₅ 717.002

Alkaloid from the marine ascidian *Polycitor* sp. Yellowish oil or red cryst. Mp 180-181°.

17-Me ether: Polycitrin B

[153212-86-3]

C₂₅H₁₇Br₄NO₅ 731.029

From *Polycitor* sp. Cryst. (CH₂Cl₂/pentane) (synthetic). Mp 140-142° (synthetic).

Rudi, A. *et al.*, *J.O.C.*, 1994, **59**, 999 (isol, uv, ir, pmr, cmr, struct)

Terpin, A. *et al.*, *Tetrahedron*, 1995, **51**, 9941-9946 (synth)


Beccalli, E.M. *et al.*, *Tetrahedron*, 2000, **56**, 2699-2702 (synth, Polycitrin B)

Winkhofer, C. *et al.*, *Synthesis*, 2006, 3043-3047 (synth)

Polyervinine

P-560

[169971-14-6]

C₄₃H₄₆N₄O₁₀ 778.857


Zwitterionic. Alkaloid from leaves of *Ervatamia polyneura*. Air-sensitive amorph. purple powder. λ_{\max} 203 (log ϵ 4.49); 244 (sh) (log ϵ 4.11); 316 (log ϵ 4.18); 393 (sh) (log ϵ 3.35); 583 (log ϵ 3.33) (MeOH).

Clivio, P. *et al.*, *Phytochemistry*, 1995, **40**, 953 (isol, uv, ir, pmr, cmr, struct)

Polygonapholine

P-561

[188302-30-9]


C₃₄H₂₉NO₇ 563.606

Alkaloid from rhizomes of *Polygonatum alte-lobatum*. Pale yellow powder. Mp 238-240°. $[\alpha]_{\text{D}}^{25}$ -76 (c, 0.5 in MeOH). λ_{\max} 225 ; 306 ; 330 (sh) ; 429 (sh) (MeOH).

Lin, C.-N. *et al.*, *Tetrahedron*, 1997, **53**, 2025 (isol, uv, ir, pmr, cmr, ms, struct)

Polyneuridine**P-562**

Methyl 17-hydroxysarpagan-16-carboxylate, 9CI†. *Normacusine A* [6872-44-2]



Absolute Configuration

$C_{21}H_{24}N_2O_3$ 352.432

Epimeric with Akuammidine, A-229. Alkaloid from *Aspidosperma polyneuron*, *Rauwolfia suaveolens*, *Rauwolfia volkensii*, *Strychnos potatorum* and *Hazunia* spp. Cryst. (EtOH or C_6H_6). Mp 245-247.5° (solvate). $[\alpha]_D^{25}$ -73 (c, 0.82 in Py). $[\alpha]_D^{25}$ +1 (c, 1.03 in $CHCl_3$).

N^4 -Oxide: **Polyneuridine N-oxide**

$C_{21}H_{24}N_2O_4$ 368.432

Alkaloid from the roots of *Ochrosia acuminata*. Light yellow powder. $[\alpha]_D^{25}$ -19.5 (c, 0.1 in MeOH).

O-Ac:

Cryst. (MeOH). Mp 270-272°. $[\alpha]_D$ -92 (c, 0.71 in Py).

N^1 -Me: **Voachalotine**

[664-25-5]

$C_{22}H_{26}N_2O_3$ 366.459

Alkaloid from *Voacanga chalongiana*, *Tabernaemontana fuchsiaefolia* and *Alstonia legouixiae* (Apocynaceae). Shows cardiotoxic props. Prisms (C_6H_6). Mp 223-224°. $[\alpha]_D^{22}$ -2.8 (c, 6 in $CHCl_3$).

N^4 -Me: **Macusine A**

[6801-39-4]

$C_{22}H_{27}N_2O_3^{\oplus}$ 367.467

Quaternary alkaloid from the bark of *Strychnos toxifera* (Loganiaceae). Cryst. (EtOH/Et₂O) (as chloride). Mp 252° (darkens at 240°, sinters at 248°) (chloride). $[\alpha]_D^{25}$ -57.5 (c, 1.46 in H₂O).

N^1, N^4 -Di-Me: **N^4 -Methylvoachalotine**

[109269-77-4]

$C_{23}H_{29}N_2O_3^{\oplus}$ 381.494

Quaternary alkaloid from the roots of *Tabernaemontana laeta*. Cryst. (MeOH). Mp 259-260°. $[\alpha]_D^{25}$ -21.3 (c, 6.3 in DMSO). λ_{max} 223 (ε 42570); 282 (ε 6290) (MeOH).

N^1 -Me, O-Ac:

Needles (MeOH aq.). Mp 180-181.5°.

$[\alpha]_D^{20}$ -34.8 (c, 1 in $CHCl_3$).

17-Aldehyde: Voachalotinal. Polyneuridine aldehyde

[92138-23-3]

$C_{21}H_{22}N_2O_3$ 350.416

Alkaloid from the leaves of *Alstonia undulata* and bark of *Aspidosperma dasycarpon*. Cryst. (Et₂O). Mp 231°. The *A. dasycarpon* isolate was obtained in v. small amt. and C-17 config. was not assigned. It could therefore have been Akuammidine aldehyde. Identity with Voachalotinal is assigned here based solely on the identical reported Mps. The uv spectrum is that of this isolate. λ_{max} 226 (log ε 4.5); 282

(log ε 3.67); 290 (log ε 3.65) (EtOH).

19,20ξ-Dihydro: 19,20-Dihydropolyneuridine

[85799-35-5]

$C_{21}H_{26}N_2O_3$ 354.448

Minor alkaloid from the leaves of *Alstonia venenata* (Apocynaceae). Amorph.

19S,20-Dihydro, N¹-Me, O-Ac: 17-O-Acetyl-19,20-dihydrovoachalotine

[62404-92-6]

$C_{24}H_{30}N_2O_4$ 410.512

Alkaloid from the root bark of *Voacanga chalongiana* (Apocynaceae). Cryst. (EtOH). Mp 201°. $[\alpha]_D^{20}$ -54 ($CHCl_3$).

11-Hydroxy, N¹, N⁴-di-Me: N¹-Methyl-11-hydroxymacusine A

[163131-24-6]

[163131-24-6 (chloride)]

$C_{23}H_{29}N_2O_4^{\oplus}$ 397.493

Quaternary alkaloid from stem bark of *Stemmadenia obovata* (Apocynaceae). Needles (MeOH/ $CHCl_3$) (as chloride). Mp 300° (chloride). $[\alpha]_D^{25}$ -76 (c, 0.167 in MeOH) (chloride).

11-Methoxy, N⁴-Me: 11-Methoxymacusine A

[87340-28-1]

$C_{23}H_{29}N_2O_4^{\oplus}$ 397.493

Alkaloid from the root and stem bark of *Strychnos angolensis* (Loganiaceae). Muscular relaxant. Mp 224-226° (chloride). λ_{max} 262; 266 (sh); 293; 300 (sh) (MeOH).

►VQ5800000

12-Methoxy, N¹, N⁴-di-Me: 12-Methoxy-4-methylvoachalotine

[109304-75-8]

$C_{24}H_{31}N_2O_4^{\oplus}$ 411.52

Quaternary alkaloid from stem bark of *Peschiera fuchsiaefolia* (Apocynaceae). Amorph. solid. Mp 221-223°. $[\alpha]_D^{25}$ -106.19 (c, 1 in MeOH). No counterion mentioned.

12-Methoxy, N¹, N⁴-di-Me, Et ester: 12-Methoxy-4-methylvoachalotine ethyl ester

[109269-78-5]

$C_{25}H_{33}N_2O_4^{\oplus}$ 425.547

Quaternary alkaloid from the stem bark of *Peschiera fuchsiaefolia* (Apocynaceae). Amorph. solid. Mp 211-214°. $[\alpha]_D^{25}$ -55.94 (c, 1 in MeOH). No counterion mentioned.

16-Epimer: see Akuammidine, A-229

Pecher, J. et al., *Chem. Ind. (London)*, 1960,

1481 (*Voachalotine, isol, struct*)

Battersby, A.R. et al., *J.C.S.*, 1960, 1848-1854 (*Macusine A*)

Pecher, J. et al., *Tet. Lett.*, 1961, 270-273 (*Voachalotine, struct*)

Janot, M.-M. et al., *Bull. Soc. Chim. Fr.*, 1962,

1079-1081 (*abs config*)

Antonaccio, L.D. et al., *J.A.C.S.*, 1962, **84**,

2161-2169 (*isol, uv, ir, ms, struct*)

Iball, J. et al., *Acta Cryst.*, 1963, **16**, 434 (*cryst struct, Voachalotine*)

McPhail, A.T. et al., *J.C.S.*, 1963, 1832-1840

(*cryst struct, Macusine A*)

Ohashi, M. et al., *Tetrahedron*, 1963, **19**, 2241-2246 (*ms*)

Battersby, A.R. et al., *J.C.S.*, 1964, 4419-4427 (*Macusine A, struct, abs config*)

Joule, J.A. et al., *Tetrahedron*, 1965, **21**, 1717-1734 (*Polyneuridine aldehyde*)

Achenbach, H. et al., *Tet. Lett.*, 1966, 4405-4407 (*Voachalotine*)

Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1974, **39**, 3168-3176 (*uv, cd, abs config, Voachalotine, synth*)

Bombardelli, E. et al., *Phytochemistry*, 1976, **15**, 2021-2022

(*Acetyldihydrovoachalotine*)

Majumder, P. et al., *Phytochemistry*, 1982, **21**, 2389-2392 (*19,20-Dihydropolyneuridine*)

Verpoorte, R. et al., *J. Nat. Prod.*, 1983, **46**, 572-575 (*11-Methoxymacusine A*)

Braga, R.M. et al., *Phytochemistry*, 1987,

26, 833-836 (*12-Methoxy-4-methylvoachalotine*)

Pinchon, T.-M. et al., *Phytochemistry*, 1990,

29, 3341-3344 (*Voachalotinal*)

Madinaveitia, A. et al., *J. Nat. Prod.*, 1995, **58**,

250 (*N¹-Methyl-11-hydroxymacusine A*)

Medeiros, W.L.B. et al., *J. Braz. Chem. Soc.*, 2001, **12**, 368-372 (*N-Methylvoachalotine*)

Salim, A.A. et al., *J. Nat. Prod.*, 2004, **67**,


1719-1721 (*N-oxide*)

Yin, W. et al., *Org. Lett.*, 2007, **9**, 295-298

(*synth*)

Polysignine**P-563**

[102759-47-7]



$C_{21}H_{29}NO_3$ 343.465

A secobenzyloquinoline alkaloid. Alkaloid from bark of *Polyalthia insignis*.

3'-Methoxy: Methoxypolysignine

[66920-24-9]

$C_{22}H_{31}NO_4$ 373.491

Alkaloid from bark of *Polyalthia insignis*.


Lee, K.-H. et al., *Tet. Lett.*, 1997, **38**, 1253

(*isol, pmr, cmr, ms, struct*)

Nimgirawath, S. et al., *Aust. J. Chem.*, 2000, **53**, 523-525 (*synth*)

Polyveoline**P-564**

[67701-60-4]




$C_{23}H_{33}NO$ 339.52

Alkaloid from the stem bark of *Greenwayodendron suaveolens* (*Polyalthia suaveolens*) (Annonaceae). Mp 172°. $[\alpha]_D$ -23 (c, 1.1 in $CHCl_3$). λ_{max} 249 (ε 5750); 301 (ε 2340) (EtOH) (Derep).

Hocquemiller, R. et al., *Tet. Lett.*, 1981, **22**, 5057 (*uv, pmr, ms, cryst struct, abs config*)

Ponapensin

[922169-95-7]



Relative Configuration

C₃₂H₃₅NO₈ 561.63


Alkaloid from *Aglaia ponapensis*. NF-κB inhibitor. Yellow oil. [α]_D²² -167 (c, 0.4 in CHCl₃). λ_{max} 224 (log ε 3.84) (MeOH).

Salim, A.A. *et al.*, *Bioorg. Med. Chem. Lett.*, 2007, 17, 109-112 (*isol, pmr, cmr*)

Pontevedrine

P-566

1,2,9,10-Tetramethoxy-6-methyl-4H-dibenzo[de,g]quinoline-4,5(6H)-dione, 9CI [34647-65-9]



C₂₁H₁₉NO₆ 381.384

Alkaloid from *Glaucium flavum* var. *vestitum*. Also obt. by treatment of Cataline, C-190 with I₂ or DDQ (Papaveraceae). Red needles (EtOH/CHCl₃). Mp 269-271°.

5-Alcohol: **Dihydropontevedrine**

[72498-24-9]

C₂₁H₂₁NO₆ 383.4

Minor alkaloid from *Glaucium flavum* (Papaveraceae). Red needles (EtOH). Mp 251-253°.

Ribas, I. *et al.*, *Tet. Lett.*, 1971, 3093 (*uv, ir, pmr*)

Castedo, L. *et al.*, *Tet. Lett.*, 1976, 501; 1978, 2179 (*struct, synth*)

Daskalova, E. *et al.*, *Phytochemistry*, 1988, 27, 953 (*Dihydropontevedrine*)

Atanes, N. *et al.*, *J.O.C.*, 1991, 56, 2984 (*synth*)


Estévez, J.C. *et al.*, *Heterocycles*, 1994, 38, 1 (*synth*)

Estévez, J.C. *et al.*, *Tetrahedron*, 1994, 50, 2107 (*synth*)

Popolohuanone A

P-567

[133056-07-2]



Absolute Configuration

C₄₂H₅₇NO₃ 623.917

Terpenoid (clerodane) numbering shown; different numbering given in the lit. Isol. from the sponge *Dysidea* sp. Intense purple-blue powder. λ_{max} 222 (log ε 4.28); 266 (log ε 4.21); 512 (log ε 3.58) (hexane).

Δ^{3,4}-Isomer: **Popolohuanone B**

[133056-09-4]

C₄₂H₅₇NO₃ 623.917

Isol. from *Dysidea* sp. Purple-blue powder. λ_{max} 222 (log ε 4.08); 266 (log ε 3.98); 512 (log ε 3.36) (hexane).

8,8',9,9',10,10'-Hexaepimer, Δ^{3,4:3',4'}-isomer: **Popolohuanone C**

[144587-57-5]

C₄₂H₅₇NO₃ 623.917

Isol. from *Dysidea avara*. Dark violet amorph. solid. Mp 125-135°. [α]_D²⁰ -9 (c, 0.08 in CH₂Cl₂). λ_{max} 232 ; 275 ; 532 (MeOH).


Rodriguez, A.D. *et al.*, *Tetrahedron*, 1990, 46, 8025-8030 (*Popolohuanones A-B*)

Alvi, K.A. *et al.*, *J.O.C.*, 1992, 57, 6604-6607 (*Popolohuanone C*)

Popolohuanone D

P-568

[144587-58-6]



C₄₂H₅₇NO₃ 623.917


Isol. from the sponge *Dysidea avara*. Dark violet amorph. solid. Mp 113-120°. [α]_D -15 (c, 0.06 in CHCl₃). λ_{max} 234 ; 275 ; 533 (MeOH).

Alvi, K.A. *et al.*, *J.O.C.*, 1992, 57, 6604-6607 (*isol, uv, pmr, cmr, ms*)

Poranthericine

P-569

1-Ethyl-dodecahydro-9a-methylpyrido[2,1,6-de]quinolizin-2-ol, 9CI [39027-76-4]



Absolute configuration

C₁₅H₂₇NO 237.384

Alkaloid from the shrub *Poranthera corymbosa* (Euphorbiaceae). Gum. [α]_D -20 (c, 1.57 in CHCl₃).

Hydrobromide:

Cryst. (EtOH). Mp 308° dec. (subl. >260°). [α]_D -8 (c, 0.51 in MeOH).

Ac: **O-Acetylporanthericine**

[38976-55-5]

C₁₇H₂₉NO₂ 279.422

Alkaloid from *Poranthera corymbosa* (Euphorbiaceae). Gum. [α]_D +2 (c, 0.76 in CHCl₃).

1-Epimer:

Needles (Me₂CO). Mp 124-126°. [α]_D +31 (c, 0.73 in CHCl₃).

1-Epimer, hydrobromide:

Cryst. (Me₂CO). Mp 278° (with subl. >250°). [α]_D +36 (c, 0.50 in MeOH).

1-Epimer, Ac: **Porantheriline**

[54352-65-7]

C₁₇H₂₉NO₂ 279.422

Alkaloid from the shrub *Poranthera corymbosa* (Euphorbiaceae). Prisms (Me₂CO). Mp 76-77°. [α]_D +87 (c, 0.71 in CHCl₃).

Denne, W.A. *et al.*, *Tet. Lett.*, 1972, 1767 (*cryst struct, abs config*)


Denne, W.A. *et al.*, *J. Cryst. Mol. Struct.*, 1973, 3, 139 (*cryst struct, abs config*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1974, 27, 2025 (*isol, ir, pmr, ms, synth*)

Porantheridine

P-570

Decahydro-9a-methyl-2-propyl-2H-[1,3]oxazino[2,3,4-de]quinolizine, 9CI



Absolute configuration

C₁₅H₂₇NO 237.384

(-)-form [39027-77-5]

Alkaloid from the shrub *Poranthera corymbosa* (Euphorbiaceae). Oil. [α]_D -26 (c, 0.57 in EtOH).

Hydrobromide:

Prisms (Me₂CO). Mp 165-166°. [α]_D -19 (c, 0.89 in MeOH).

(±)-form

Synthetic. Oil.

Denne, W.A. *et al.*, *J. Cryst. Mol. Struct.*, 1973, 3, 87 (*cryst struct, abs config*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1974, 27, 2025 (*isol, ms*)

Gössinger, E. *et al.*, *Monatsh. Chem.*, 1980, 111, 783 (*synth, ir, pmr, cmr, ms*)

Gössinger, E. *et al.*, *Tet. Lett.*, 1980, 21, 2229 (*synth, pmr, cmr, ms*)


Comins, D.L. *et al.*, *J.A.C.S.*, 1993, 115, 8851 (*synth*)

David, M. *et al.*, *J.O.C.*, 1999, 64, 8402-8405 (*synth*)

Porantherilidine

P-571

Octahydro-6-methyl-α-propyl-2H-quinolizine-4-ethanol benzoate, 9CI



Absolute configuration

C₂₂H₃₃NO₂ 343.508

(-)-form [51105-10-3]

Alkaloid from the shrub *Poranthera corymbosa* (Euphorbiaceae). Gum. $[\alpha]_D^{25}$ -47 (c, 0.75 in CHCl_3).

Hydrobromide:

Needles (Me_2CO). Mp 251-252°. $[\alpha]_D^{25}$ -27 (c, 0.46 in MeOH).

(±)-form [74164-01-5]

Synthetic. Noncryst.

Denne, W.A. *et al.*, *J. Cryst. Mol. Struct.*, 1973, 3, 367 (*cryst struct, abs config*)


Johns, S.R. *et al.*, *Aust. J. Chem.*, 1974, 27, 2025 (*isol, ir, pmr, ms*)

Gössinger, E. *et al.*, *Monatsh. Chem.*, 1980, 111, 143; 783 (*synth, ir, pmr, ms*)

Porantherine

P-572

3a,4,5,6,7,8,9,9a-Octahydro-9a-methyl-1H-1,6a-ethanopyrido[2,1,6-de]quinolizine, 9CI



Absolute configuration

$\text{C}_{15}\text{H}_{23}\text{N}$ 217.353

(+)-form [33529-61-2]

Major alkaloid from the shrub *Poranthera corymbosa* (Euphorbiaceae). Mp 36-40°. $[\alpha]_D^{25}$ +29 (c, 0.35 in CHCl_3).

Hydrochloride:

Prisms (EtOH). Mp 307° dec.

Hydrobromide:

Prisms (EtOH). $[\alpha]_D^{25}$ -13 (c, 0.35 in EtOH). Dec. >340° without melting.

Picrate:

Yellow needles (EtOH). Mp 295° (dec.).

Dihydro: Mp 81-82°. $[\alpha]_D^{25}$ +15 (c, 0.55 in CHCl_3).

(±)-form [54382-19-3]

Synthetic. Oil.

Denne, W.A. *et al.*, *Tet. Lett.*, 1971, 3107 (*cryst struct, abs config*)

Denne, W.A. *et al.*, *J. Cryst. Mol. Struct.*, 1973, 3, 79 (*cryst struct, abs config*)


Johns, S.R. *et al.*, *Aust. J. Chem.*, 1974, 27, 2025 (*isol, pmr, ms*)

Corey, E.J. *et al.*, *J.A.C.S.*, 1974, 96, 6516 (*synth*)

Porothramycin A

P-573

BMV 28121. BU 2916T. Antibiotic BMV 28121. Antibiotic BU 2916T [110652-73-8]



$\text{C}_{18}\text{H}_{21}\text{N}_5\text{O}_4$ 343.382

Anthramycin-type antibiotic. Prod. by a *Streptomyces albus* sp. Active against gram-positive bacteria and tumours. Pale

yellow powder. Sol. MeOH, CHCl_3 , Py, EtOH, DMSO; fairly sol. EtOAc, Et_2O ; poorly sol. H_2O , hexane. Mp 140-150° dec. $[\alpha]_D^{25}$ +432 (c, 0.46 in CHCl_3). Porothramycin A or B obt. depending on isol. process used. λ_{max} 214 (ε 22000); 235 (ε 20300); 336 (ε 45500) (MeOH) (Derep). λ_{max} 214 (ε 12000); 235 (ε 20300); 335 (ε 45500) (MeOH) (Berdy). λ_{max} 214; 235; 335 (MeOH-HCl) (Berdy).

► UC6360000

Me ether: Porothramycin B

[110652-72-7]

$\text{C}_{19}\text{H}_{23}\text{N}_3\text{O}_4$ 357.408

From *Streptomyces albus*. Similar biol. props. as parent. Yellow needles (EtOAc/MeOH). Sol. MeOH, CHCl_3 , Py, DMSO, EtOH; fairly sol. EtOAc, Et_2O ; poorly sol. H_2O , hexane. Mp 164-166°. $[\alpha]_D^{25}$ +669 (c, 0.62 in CHCl_3). λ_{max} 214 (ε 22000); 235 (ε 20300); 336 (ε 45500) (MeOH) (Derep). λ_{max} 213 (ε 27800); 236 (ε 27100); 338 (ε 57500) (MeOH) (Berdy). λ_{max} 213; 236; 338 (MeOH-HCl) (Berdy).


► LD₅₀ (mus, ipr) 981 mg/kg. UC6350000

Tsunakawa, M. *et al.*, *J. Antibiot.*, 1988, 41, 1366 (*isol, pmr, cmr, struct*)

Porphobilinogen

P-574

5-(Aminomethyl)-4-(carboxymethyl)-1H-pyrrole-3-propanoic acid, 9CI [487-90-1]



$\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_4$ 226.232

Isol. from urine of patients suffering from certain types of porphyria. Key monopyrrolic intermed. in porphyrin, chlorophyll and vitamin B₁₂ biosynth. Cryst. + 1H₂O (H₂O). Mp 167° Mp 175-180° dec. Unstable in air and in aq. soln.; turns pink. On heating in acid (in air) forms a mixt. of Uroporphyrins I, II, III and IV.

Lactam: [3468-98-2]

$\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_3$ 208.216

Cryst. (AcOH aq.). Dec. >325°.

Rimington, C. *et al.*, *Nature (London)*, 1955, 175, 630 (*isol*)

Jackson, A.H. *et al.*, *Can. J. Chem.*, 1957, 35, 717 (*synth*)

Bezold, G. *et al.*, *Angew. Chem., Int. Ed.*, 1967, 6, 883 (*rev*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1973, 1546

Jones, M.I. *et al.*, *Chem. Comm.*, 1976, 472 (*synth*)

Kenner, G.W. *et al.*, *J.C.S. Perkin I*, 1977, 332 (*synth*)

Gibbs, P.N.B. *et al.*, *Biochem. J.*, 1986, 236, 447 (*biosynth*)

Blanche, F. *et al.*, *Angew. Chem., Int. Ed.*, 1995, 34, 383 (*rev*)

Bobal, P. *et al.*, *Trends Org. Chem.*, 1997, 6, 125-144 (*rev, synth*)


Jacobi, P.A. *et al.*, *J. Appl. Chem.*, 2001, 123, 9307-9312 (*synth, pmr, cmr*)

Portulacaxanthin I

P-575

Portulacaxanthine

[11042-69-6]



$\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_7$ 324.29


Betalaine pigment. Alkaloid from the flowers *Portulaca grandiflora* (Portulacaceae).

Piattelli, M. *et al.*, *CA*, 1967, 66, 112959p (*isol*)
Trezza, G.F. *et al.*, *Phytochemistry*, 1991, 30, 1897

Portulacaxanthin II

P-576

4-[[[Carboxy-(4-hydroxyphenyl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid. Tyrosinebetaxanthin [135545-98-1]



$\text{C}_{18}\text{H}_{18}\text{N}_2\text{O}_7$ 374.349


Betalaine pigment. Alkaloid from the flowers of *Portulaca grandiflora* (Portulacaceae).

Trezza, G.F. *et al.*, *Phytochemistry*, 1991, 30, 1897 (*isol, struct*)

Portulacaxanthin III

P-577

4-[[[(Carboxymethyl)imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid. Glycinebetaxanthin [135545-99-2]




$\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_6$ 268.226

Betalaine pigment. Alkaloid from the flowers of *Portulaca grandiflora* (Portulacaceae). λ_{max} 470 nm.

Trezza, G.F. *et al.*, *Phytochemistry*, 1991, 30, 1897 (*isol, struct*)


Potanisine G

[664304-12-5]

P-578C₃₈H₅₅N₃O₁₀ 713.867

Closely related to Delsemine, D-196. Alkaloid from the roots of *Delphinium potaninii*. Amorph. powder. $[\alpha]_D^{10}$ -24.5 (c, 1 in CHCl₃). Possible artifact.

Chen, D.-L. *et al.*, *J. Asian Nat. Prod. Res.*, 2003, **5**, 209-213 (*isol*, *pmr*, *cmr*)

Powellamine*Cripaline***P-579**

Relative configuration

C₁₆H₁₇NO₃ 271.315

May be the C7-C8 methylenedioxy isomer. Historically Powellamine was the (-)-form and Cripaline the (+)-form.

(+)-form

Alkaloid from *Crinum powellii* var. *harlemense* (Amaryllidaceae). Cryst. (Me₂CO). Mp 198-199°. $[\alpha]_D^{24}$ +50 (c, 0.2 in CHCl₃).

Perchlorate:

Cryst. (MeOH). Mp 257-258°.

Picrate:

Cryst. (MeOH). Mp 185-186°.

(-)-form

Alkaloid from the bulbs of *Crinum powellii* and from the *Crinum* hybrid "Ellen Bosanquet" (Amaryllidaceae). Mp 198-200°. $[\alpha]_D^{25}$ -49 (c, 0.25 in CHCl₃).

Perchlorate: Mp 258°.**Picrate:** Mp 186°.**(±)-form**

Obt. from equimolar amts. of the enantiomers. Mp 201°.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1959, **46**, 475; 1961, **48**, 406 (*isol*, *struct*)

Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1962, **295**, 868; *C.A.*, **58**, 11415g (*Cripaline*)

Powellidine

[1360-53-8]

P-580C₁₆H₁₇NO₃ 271.315


Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Crinum powellii* (Amaryllidaceae). Mp 207-209°.

 $[\alpha]_D^{26}$ +100 (c, 0.2 in CHCl₃).**Perchlorate:** Mp 177°.**Picrate:** Mp 198°.

Boit, H.-G. *et al.*, *Naturwissenschaften*, 1959, **46**, 475 (*isol*)

Powelline

[7363-25-9]

P-581

Absolute Configuration

C₁₇H₁₉NO₄ 301.341

Alkaloid from *Crinum powellii* and several other spp. in the Amaryllidaceae. Mp 197-198°. $[\alpha]_D^0$ (CHCl₃). λ_{\max} 288 (log ϵ 3.23) (EtOH).

Picrate: Mp 228-231° dec.**Me ether:** *Buphanidrine*. *Distichine*

[1162-10-3]

Alkaloid from *Boophone fischeri*, *Boophone disticha* and some other spp. in the Amaryllidaceae. Analgesic. Mp 90-92° Mp 144° (hydrate). $[\alpha]_D^{23}$ -6.9 (c, 1 in CHCl₃). $[\alpha]_D^{27}$ +4.2 (c, 0.54 in EtOH).

► Toxic, LD₅₀ (mus, scn) 8.9 mg/kg.

Me ether, hydrobromide: Mp 195-197°. $[\alpha]_D^{20}$ +5.6 (95% EtOH).**Me ether, picrate:** Mp 238-239°.**1β,2β-Epoxyde:** see Crinamidine, C-743**6α-Hydroxy:** **6α-Hydroxypowelline**

[31128-98-0]

C₁₇H₁₉NO₅ 317.341

Isol. from the bulbs of *Nerine bowdenii* and from an unidentified *Crinum* sp. Prisms (Me₂CO). Mp 233-235°. $[\alpha]_D^{24}$ -36 (c, 0.19 in MeOH). λ_{\max} 218 (ϵ 17500); 235 (ϵ 6800); 286 (ϵ 2150) (95%EtOH).

6α-Hydroxy, di-Ac:Cryst. (Me₂CO). Mp 114-117°. $[\alpha]_D^{24}$ -26 (c, 0.20 in MeOH).**6α-Hydroxy, O³-Me:** **6α-Hydroxybuphanidrine**

[31128-91-3]

C₁₈H₂₁NO₅ 331.368

Isol. from the bulbs of *Nerine bowdenii* and *Nerine filifolia*. Cytotoxic agent active against Rauscher leukaemia virus. Prisms (CHCl₃). Mp 95-96°. $[\alpha]_D^{24}$ -64 (c, 0.19 in MeOH). $[\alpha]_D^{20}$ +19 (c, 0.8 in MeOH). λ_{\max} 215 (ϵ 18250); 238 (ϵ 7750); 285 (ϵ 1880) (95%EtOH).

6α-Hydroxy, O³-Me, hydrochloride: Amorph. $[\alpha]_D^{24}$ -22 (c, 0.16 in MeOH).**6α-Hydroxy, O³-Me, Ac:**Cryst. (C₆H₆). Mp 165-168°. $[\alpha]_D^{24}$ -42 (c, 0.23 in MeOH).**6α-Methoxy, O³-Me:** **6α-Methoxybuphanidrine**C₁₉H₂₃NO₅ 345.394

Alkaloid from the bulbs of *Nerine filifolia*. $[\alpha]_D^{20}$ +34.6 (c, 0.13 in MeOH).

Wildman, W.C. *et al.*, *J.A.C.S.*, 1958, **80**, 2567-2575 (*ir*, *struct*, *Powelline*, *Buphanidrine*)

Goosen, A. *et al.*, *J.C.S.*, 1960, 1094-1096 (*isol*, *Buphanidrine*)

Haugwitz, R.D. *et al.*, *J.C.S.*, 1965, 2001-2009 (*pmr*, *Buphanidrine*)

Clardy, J.C. *et al.*, *J.A.C.S.*, 1970, **92**, 6337-6339 (*6-Hydroxybuphanidrine*, *cryst struct*)

Slabaugh, M.R. *et al.*, *J.O.C.*, 1971, **36**, 3202-3207 (*6-Hydroxypowelline*, *6-Hydroxybuphanidrine*)

Zetta, L. *et al.*, *J.C.S. Perkin 2*, 1973, 1180-1184 (*cmr*)

Longévialle, P. *et al.*, *Org. Mass Spectrom.*, 1973, **7**, 401-415; 417-430 (*ms*)

Viladomat, F. *et al.*, *Phytochemistry*, 1995, **40**, 961-965 (*Buphanidrine*)

Nair, J.I. *et al.*, *Phytochemistry*, 2005, **66**, 373-382 (*6α-Hydroxybuphanidrine*, *6α-Methoxybuphanidrine*)

Poweramine**P-582**

[1360-54-9]

C₂₃H₃₀N₂O₄ 398.501

Tentative mol. formula. Struct. unknown. Alkaloid from leaves of *Ochrosia poweri* (Apocynaceae). Prisms (MeOH). Mp 241-242° dec. Conts. a 6-methoxyindole chromophore.

Picrate:

Red needles (MeOH), turning brown on drying. Mp 156°.

Doy, F.A. *et al.*, *Aust. J. Chem.*, 1962, **15**, 548-554 (*isol*, *uv*)

Powerchrine**P-583**

[1360-55-0]

C₂₂H₂₆N₂O₃ 366.459


Struct. unknown. Alkaloid from *Ochrosia poweri* (Apocynaceae). Needles (MeOH). Mp 213°. $[\alpha]_D^{25}$ -80.8 (c, 1.1 in Me₂CO).

Douglas, B. *et al.*, *Aust. J. Chem.*, 1964, **17**, 246-255 (*isol*, *uv*)

Pramanicin**P-584**

3,4-Dihydroxy-5-(hydroxymethyl)-3-[3-(3-nonyloxiranyl)-1-oxo-2-propenyl]-2-pyrrolidinone, 9CI. Antibiotic L 746628. L 746628

[154445-05-3]



Absolute Configuration

C₁₉H₃₁NO₆ 369.457

Tetramic acid deriv. Prod. by *Stagonospora* sp. ATCC 74235. Antimicrobial agent. Cryst. (MeOH aq.). Mp 110-113°. $[\alpha]_D^{25}$ -35 (c, 0.2 in MeOH). λ_{\max} 245 (ϵ 14430) (MeOH).

Schwartz, R.E. *et al.*, *Tetrahedron*, 1994, **50**, 1675-1686 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)


Barrett, A.G.M. *et al.*, *J.O.C.*, 1999, **64**, 6005-6018 (*synth*, *abs config*)

Harrison, P.M. *et al.*, *J.C.S. Perkin 1*, 2000, 4390-4402 (*biosynth*)

Aoki, S. *et al.*, *Heterocycles*, 2006, **69**, 49-54 (*synth*)

Prangosine**P-585**

2-(1-Amino-1-methylethyl)-7H-furo[3,2-g][1]benzopyran-7-one, 9CI
[15399-67-4]



C₁₄H₁₃NO₃ 243.262

Alkaloid from *Prangos pabularia* (Apiaceae). Cryst. Mp 131-132°.

Mukhamedova, K.S. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 117; 357; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 97; 303 (*struct.*, *ms*)


Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 287; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 243 (*pmr*)

Schneiders, G.E. *et al.*, *J. Chem. Res., Synop.*, 1982, 182 (*synth*)

Zubia, E. *et al.*, *Tetrahedron*, 1992, **48**, 4239 (*synth*)

Pratorinine**P-586**

9-Hydroxy-10-methoxy-7H-pyrrolo[3,2,1-de]phenanthridin-7-one [88660-12-2]



C₁₆H₁₁NO₃ 265.268

There is considerable confusion regarding the structs. of Pratorimine and Pratorinine due to labelling errors. In the paper by Maddry *et al.*, the names are reversed. The alkaloid synthesised by Joshi *et al.*, and named Pratorimine now corresponds to Pratorinine. Alkaloid from the bulbs of *Crinum latifolium* and *Crinum bulbispermum*, also from *Crinum asiaticum* and *Crinum augustum* (Amaryllidaceae). Needles (MeOH). Mp 224-225°. The earlier reported Mp 263-265° is erroneous.

Me ether: Pratosine. O-Methylpratorimine. O-Methylpratorinine. Criasiaticidine A
[88720-89-2]

C₁₇H₁₃NO₃ 279.295

Alkaloid from the bulbs of *Crinum latifolium* and *Crinum asiaticum*. Fine needles (Et₂O/MeOH). Mp 277-279°.

O-De-Me: Hippacine[†]

C₁₅H₉NO₃ 251.241

Alkaloid from the bulbs of *Crinum bulbispermum*. Amorph. powder.

O-De-Me, O^o-Me: Pratorimine

[80787-57-1]

C₁₆H₁₁NO₃ 265.268

Alkaloid from the bulbs of *Crinum pratense*, *Crinum latifolium* and *Crinum bulbispermum*. Also from *Crinum asiaticum* (Amaryllidaceae). Flakes (CHCl₃/EtOH). Mp 265-267°.

Ghosal, S. *et al.*, *Phytochemistry*, 1981, **20**, 2003; 1983, **22**, 2305; 1985, **24**, 2141 (*isol.*, *uv*, *ir*, *pmr*, *ms*, *struct.*, *synth.*, *occur.*, *rev*)

Maddry, J.A. *et al.*, *Tet. Lett.*, 1985, **26**, 4301 (*struct*)

Joshi, B.S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 445 (*Pratorinine, synth*)

Iwao, M. *et al.*, *Heterocycles*, 1994, **38**, 1717 (*synth.*, *Pratosine*)

Hutchings, R.H. *et al.*, *J.O.C.*, 1996, **61**, 1004 (*synth.*, *Pratosine*)

Miki, Y. *et al.*, *Tet. Lett.*, 1999, **40**, 4347-4348 (*Pratosine, synth*)

Ramadan, M.A. *et al.*, *Phytochemistry*, 2000, **54**, 891-896 (*Hippacine*)


Min, B.S. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 1217-1219 (*Criasiaticidine A*)

Ganton, M.D. *et al.*, *Org. Lett.*, 2005, **7**, 4777-4779 (*synth*)

Mentzel, U.V. *et al.*, *J.O.C.*, 2006, **71**, 5807-5810 (*Pratosine, synth*)

Prekuammicine**P-587**

Methyl 1,2,19,20-tetrahydro-17-hydroxycycuran-16-carboxylate, 9CI
[23924-90-5]



Absolute Configuration


C₂₁H₂₄N₂O₃ 352.432

The C-16 config. was not assigned, but its ready conversion to Akuammicine, A-228 and Stemmadenine, S-510 (abs. configs. now fully known) indicates it to be as shown. Alkaloid from *Catharanthus roseus* seedlings (Apocynaceae). Non-cryst. λ_{max} 262 (ε 6000) (EtOH).

Scott, A.I. *et al.*, *J.A.C.S.*, 1969, **91**, 4932-4933; 5874-5876 (*isol.*, *uv*, *pmr*, *ms*, *struct.*, *biosynth*)

Prealmazole C**P-588**

[161068-70-8]



C₂₁H₂₃N₃O₂ 349.432


Alkaloid from an unidentified red Senegalese seaweed of the family Delesseriaceae. [α]_D²⁰ +38 (c, 0.25 in MeOH).

Guella, G. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1999-2006 (*isol.*, *pmr*, *cmr*, *ms*, *struct.*, *synth*)

Preapiodionene**P-589**

3-(3,5-Dihydroxy-4-methyl-1-oxo-6-octenyl)-2,4-pyrrolidinedione, 9CI
[142808-39-7]

[147920-90-9, 147782-11-4]




C₁₃H₁₉NO₅ 269.297

Tetramic acid deriv. Isol. from *Apiosordaria effusa*. Precursor of Apiodionene, A-1339.

Japan. Pat., 1992, 92 49 289; *CA*, **117**, 88730f (*isol*)

Takahashi, S. *et al.*, *CA*, 1993, **119**, 24279z (*isol.*, *pmr*, *cmr*)

Preaxinellamine**P-590**


C₂₂H₂₃Br₄ClN₁₀O₄ 846.557

There is indirect evidence for the occurrence of this alkaloid. Probably occurs in sponges. Precursor of pyrrole-imidazole dimers.

Grube, A. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 6721-6724; 8107 (*occur*)

Prebalamide**P-591**

[190905-97-6]



Relative Configuration

C₁₉H₁₈N₂O₂ 306.363

Alkaloid from leaves of *Clausena indica*. Cryst. (Et₂O). Mp 125-127°. [α]_D²⁰ +30 (c, 0.5 in CHCl₃). λ_{max} 224; 274 (sh); 283; 292 (Et₂O).

Riemer, B. *et al.*, *Phytochemistry*, 1997, **45**, 337-341 (*isol.*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)


Precasine**P-592**

Struct. unknown. Alkaloid from the roots of *Abrus precatorius* (Fabaceae). Mp 234-236° (as picrate). Mp given as 234-326° in CAS, presumably due to a misprint.

Khaleque, A. *et al.*, *Sci. Res. (Dacca)*, 1966, **3**, 203; *CA*, **67**, 90987n

Precatorine**P-593**

[36675-57-7]




C₁₄H₁₁NO₆ 289.244

Alkaloid from the seeds, leaves, stems and roots of *Abrus precatorius* (Fabaceae). Needles (MeOH). Mp 218-220°.

Ghosal, S. *et al.*, *Phytochemistry*, 1971, **10**, 195
(*isol, uv, ir, struct*)

Preclathridine A P-594

4-(1,3-Benzodioxol-5-ylmethyl)-1-methyl-1H-imidazol-2-amine, 9CI. 2-Amino-1-methyl-4-(3,4-methylenedioxybenzyl)-1H-imidazole
[146556-27-6]



C₁₂H₁₃N₃O₂ 231.254

Metab. from the nudibranch *Notodoris gardineri*. Anthelminthic. Yellow oil. λ_{max} 208 ; 285 (MeOH).


Alvi, K.A. *et al.*, *Tetrahedron*, 1993, **49**, 329-336 (*isol, uv, pmr, cmr, ms, struct*)

Kawasaki, I. *et al.*, *Heterocycles*, 1996, **43**, 1375 (*synth*)

Molina, P. *et al.*, *J.O.C.*, 1999, **64**, 2540-2544 (*synth, pmr, cmr*)

Precondylocarpine P-595

Methyl 1,2,14,19-tetrahydro-16-(hydroxymethyl)-condyfolan-16-carboxylate
[2779-17-1]



Absolute Configuration

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the leaves of *Vallesia dichotoma* (Apocynaceae). Cryst. (C₆H₆/hexane). Mp 143-146°.

Walser, A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 391 (*isol, uv, ir, pmr, ms, struct*)

Precurarine P-596


[1360-58-3]

Struct. unknown. Alkaloid from *Strychnos solimoesana* (Loganiaceae). Blue col. with Ce(IV).

Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148

Predicentrine P-597

5,6,6a,7-Tetrahydro-1,9,10-trimethoxy-6-methyl-4H-dibenzo[de,g]quinolin-2-ol, 9CI. 2-Hydroxy-1,9,10-trimethoxyaporphine



(S)-form

C₂₀H₂₃NO₄ 341.406

(S)-form [517-65-7]

Alkaloid from *Beilschmiedia podagrifera*, *Litsea triflora*, *Ocotea minarum*, *Ocotea brachybotra*, *Liriodendron tulipifera*, *Glaucium oxylobum*, *Corydalis cava*, *Corydalis bulbosa* and *Corydalis slivenensis* (Lauraceae, Magnoliaceae, Papaveraceae).

Hydrochloride:

Cryst. (MeOH/EtOAc). Mp 214-217°.

Hydrobromide:

Needles (MeOH). Mp 200-205° dec. [α]_D +97 (c, 0.2 in EtOH).

N-De-Me: 2-Hydroxy-1,9,10-trimethoxy-noraporphine. *Norpredicentrine*
[23768-71-0]

C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Beilschmiedia podagrifera* and *Guatteria scandens* (Lauraceae, Annonaceae). Noncryst.

N-De-Me, N-Ac:

Needles + 1MeOH (MeOH). Mp 133-135°. [α]_D +406 (c, 0.10 in CHCl₃).

(±)-form [20051-87-0]

Synthetic. Oil.

Hydrochloride:

Cryst. (MeOH/Et₂O). Mp 215-217° dec.

Hydrobromide: Mp 209-210°.

N-De-Me: [56533-87-0]

Synthetic. Cryst. (Me₂CO). Mp 204°.

N-De-Me, N-Ac:

Cryst. (CHCl₃/MeOH). Mp 234-236°.

Charubala, R. *et al.*, *Chem. Ber.*, 1968, **101**, 2665 (*synth, uv, pmr*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**,

1277 (*isol, uv, pmr, ms, struct, deriv*)

Kametani, T. *et al.*, *J.C.S. (C)*, 1971, 1032

(*synth*)

Premila, M.S. *et al.*, *Indian J. Chem.*, 1975, **13**,

13 (*deriv, synth, uv, pmr, ms*)

Preininger, V. *et al.*, *J. Pharm. Sci.*, 1976, **65**,

294 (*isol*)

Chen, C.-L. *et al.*, *Phytochemistry*, 1976, **15**,

1161 (*isol, uv, pmr, ms*)

Vecchiotti, V. *et al.*, *Farmacol. Ed. Sci.*, 1977,

32, 767; 1979, **34**, 829; *CA*, **88**, 133240s; **92**,

18789y (*isol*)

Shafiee, A. *et al.*, *J. Pharm. Sci.*, 1977, **66**, 593

(*isol, uv, pmr, ms*)

Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1978,

26, 3920 (*synth*)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1979, **42**,

325; 329 (*rev, cmr*)

Castedo, L. *et al.*, *An. Quim., Ser. C*, 1980, **76**,

171; *CA*, **94**, 61694s (*isol*)

Kiryakov, H.G. *et al.*, *Planta Med.*, 1981, **43**,

51; 1982, **44**, 168 (*isol*)

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1983, **46**,


335 (*deriv*)

Rasoanaivo, P. *et al.*, *Planta Med.*, 1998, **64**,

58-62 (*cmr, Norpredicentrine*)

Pregna-4,14-diene-3,20-diamine P-598

3,20-Diaminopregna-4,14-diene



C₂₁H₃₄N₂ 314.513

(3β,20S)-form

N²⁰,N²⁰-Di-Me, N³-tigloyl: *Salonine C*
[657411-75-1]

C₂₈H₄₄N₂O 424.668


Alkaloid from *Sarcococca saligna*.

Cholinesterase inhibitor. Amorph. solid. [α]_D²⁰ -120 (c, 0.05 in MeOH). λ_{max} 206 (log ε 2.23) (MeOH).

Atta-ur-Rahman, *et al.*, *Helv. Chim. Acta*, 2004, **87**, 439-448 (*Salonine C*)

Pregna-5,16-diene-3,20-diamine P-599

3,20-Diaminopregna-5,16-diene



C₂₁H₃₄N₂ 314.513

(3α,20S)-form

N³,N³,N²⁰-Tri-Me, N²⁰-Ac: *Saracocinaene*. *Saracocinaene*

[182815-72-1]

C₂₆H₄₂N₂O 398.631

Alkaloid from aerial parts of *Sarcococca saligna*. Solid. Mp 129-130°. [α]_D²⁵ +170 (c, 0.05 in CHCl₃).

(3β,20S)-form

N³,N²⁰,N²⁰-Tri-Me: *Sarconidine*

[193629-88-8]

C₂₄H₄₀N₂ 356.593


Alkaloid from aerial parts of *Sarcococca saligna*. Yellow amorph. solid. [α]_D²⁰ -66 (c, 0.02 in CHCl₃).

Naeem, I. *et al.*, *Phytochemistry*, 1996, **43**, 903-906 (*Saracocinaene*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1997, **45**, 861-864 (*Sarconidine*)

Pregna-14,16-diene-3,20-diamine P-600

3,20-Diaminopregna-14,16-diene



C₂₁H₃₄N₂ 314.513

(3α,5α,20S)-form

N³,N³,N²⁰-Tri-Me, N²⁰-formyl: *Sarcocinine A*. N-[3-(Dimethylamino)pregna-14,16-dien-20-yl]-N-methylformamide, 9CI
[129707-39-7]

C₂₅H₄₀N₂O 384.604

Alkaloid from the aerial parts of


Sarcococca ruscifolia (Buxaceae).

14,15-Dihydro, N³,N³,N²⁰-tri-Me, N²⁰-formyl: **Sarcocurinine A1**
[143086-41-3]
C₂₅H₄₂N₂O 386.62
From aerial parts of *Sarcococca ruscifolia*.

Qiu, M. et al., *Yunnan Zhiwu Yanjiu*, 1990, **12**, 111; 1991, **13**, 445; *CA*, **113**, 148873j; **117**, 108079f (isol)

Pregnane-3,20-diamine 3,20-Diaminopregnane

P-601



(3α,5α,20S)-form

C₂₁H₃₈N₂ 318.545

Epipachysamines are not epimers of the Pachysamines.

(3α,5α,20S)-form

N³,N²⁰-Di-Me, N²⁰-Ac: N³-**Demethylsaracodine**
[4947-49-3]
C₂₅H₄₄N₂O 388.635
Constit. of *Sarcococca saligna*.
Amorph. solid. λ_{max} 203 (log ε 2.12)
(no solvent reported).

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenyl): **Pachysamine E**. 3-(β,β-Dimethylacryloylamino)-20-dimethylaminopregnane
[6579-73-3]
C₂₈H₄₈N₂O 428.7
Alkaloid from *Sarcococca pruniformis* and *Pachysandra terminalis*. Cytotoxic. Powder. Mp 148-150° Mp 172-178°. [α]_D²⁵ +46 (CHCl₃). C-3 epimer of Epipachysamine E.

N³,N³,N²⁰-Tri-Me, N²⁰-Ac: **Saracodine**.
Saracodine
[15437-92-0]
C₂₆H₄₆N₂O 402.662
Alkaloid from *Sarcococca pruniformis* and *Sarcococca saligna*. Shows monoamine oxidase inhibitor-like activity. Cholinesterase inhibitor. Cryst. (MeOH). Mp 247°. [α]_D²⁵ -11.9 (CHCl₃).

N³,N²⁰,N²⁰-Tri-Me: 20-Dimethylamino-3-methylaminopregnane. **Pachysamine A**
[6801-29-2]
C₂₄H₄₄N₂ 360.625
Alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO/CH₂Cl₂). Mp 171-173° (167-168°). [α]_D¹⁰ +20 (CHCl₃).

▶ TU4157095

N³,N²⁰,N²⁰-Tri-Me, N³-Ac: N³-**Acetyl-pachysamine A**. **Isosaracodine**. *Isosaracodine*
[15027-96-0]
C₂₆H₄₆N₂O 402.662
Alkaloid from *Pachysandra* spp. and *Sarcococca saligna*. Cholinesterase inhibitor. Mp 150-152°.

N³,N²⁰,N²⁰-Tri-Me, N³-(3-methyl-2-butenyl): **Pachysamine B**
[6792-14-9]
C₂₉H₅₀N₂O 442.727
Alkaloid from *Pachysandra terminalis* (Buxaceae). Mp 171-173°. [α]_D¹⁰ +67 (CHCl₃).

N³,N²⁰,N²⁰-Tri-Me, N³-benzoyl: **Pachysamine H**
[128255-13-0]
C₃₁H₄₈N₂O 464.733
Alkaloid from *Pachysandra axillaris*. Mp 150-153°. [α]_D²² +57.4 (c, 0.5 in CHCl₃).

N³,N³,N²⁰,N²⁰-Tetra-Me: 3,20-Bis(dimethylamino)pregnane. **N-Methylpachysamine A**
[6584-68-5]
C₂₅H₄₆N₂ 374.652
Alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO/CH₂Cl₂). Mp 165-166°. [α]_D +16 (CHCl₃).

(3β,5α,20R)-form [95272-07-4]
[96391-90-1 (dihydrochloride)]
Noncryst. Mp 300° (as hydrochloride).

N³,N²⁰-Di-Ac: [95869-16-2]
Mp 284-288°. [α]_D +13 (MeOH).

N-Tetra-Me: [1248-06-2]
Cryst. (Me₂CO). Mp 120-121°.

(3β,5α,20S)-form [7050-25-1]

N³,N³-Di-Me: **Funtudiamine A**. *Epipachysamine F*
[5141-99-1]
C₂₃H₄₂N₂ 346.598
Alkaloid from *Funtumia latifolia*. Minor alkaloid from *Pachysandra terminalis* (Apocynaceae, Buxaceae). Mp 80°. [α]_D +6 (CHCl₃).

N³,N³-Di-Me, N²⁰-formyl: N²⁰-**Formylepipachysamine F**. *Iso-N-formylchone-morphine*
[657403-04-8]
C₂₄H₄₂N₂O 374.609
Alkaloid from *Sarcococca brevifolia*. Amorph. solid. Mp 220-222°. [α]_D²⁵ -14 (c, 0.2 in CHCl₃).

N³,N³-Di-Me, N²⁰-Ac:
Cryst. (Me₂CO). Mp 250-253°. [α]_D +6 (CHCl₃).

N³,N²⁰-Di-Me: **Epipachysamine C**. *Dictyodiamine*
[4215-74-1]
C₂₃H₄₂N₂ 346.598
Alkaloid from *Pachysandra terminalis* and *Dictyophleba lucida* roots (Buxaceae, Apocynaceae). Mp 116°. [α]_D +22 (CHCl₃).

N³,N²⁰-Di-Me, N³,N²⁰-di-Ac: **N,N-Diacetylepachysamine C**
[4947-48-2]
C₂₇H₄₆N₂O₂ 430.673
Alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO). Mp 242-243°. [α]_D -16 (CHCl₃).

N³,N²⁰-Di-Me, N³-benzoyl: **Hookerianamide I**
[957466-16-9]
C₃₀H₄₆N₂O 450.706
Alkaloid from *Sarcococca hookeriana*. Amorph. solid. [α]_D²⁵ +186 (c, 0.3 in

MeOH). λ_{max} 220 (log ε 2.1); 266 (log ε 3) (MeOH).

N²⁰,N²⁰-Di-Me: 3-Amino-20-dimethylaminopregnane. **Chonemorphine**
[4282-07-9]
C₂₃H₄₂N₂ 346.598
Main alkaloid of *Chonemorpha macrophylla* (*Chonemorpha fragrans*), isol. from *Chonemorpha penangensis*, *Malouetia bequaertiana*, *Dictyophleba lucida* and *Funtumia latifolia* (Apocynaceae). Mp 144-146°. [α]_D²² +25 (CHCl₃).

N²⁰,N²⁰-Di-Me, hydrochloride: Mp 318-320°.

N²⁰,N²⁰-Di-Me, N³-formyl: **N-Formylchonemorphine**
[51873-91-7]
C₂₄H₄₂N₂O 374.609
Alkaloid from *Chonemorpha macrophylla* (Apocynaceae) and *Sarcococca saligna* (Buxaceae). Mp 289-291°. [α]_D²⁵ +25 (c, 0.05 in MeOH).

N²⁰,N²⁰-Di-Me, N³-Ac: **Saracarine**. *N-Acetylchonemorphine*. *Saracarine*
[3514-38-3]
C₂₅H₄₄N₂O 388.635

Alkaloid possibly from *Malouetia bequaertiana* and from aerial parts of *Sarcococca saligna*. Cryst. (Me₂CO) (also descr. as an oil). Mp 272°. [α]_D +14 (c, 1 in CHCl₃). [α]_D²⁵ +49 (c, 0.815 in CHCl₃) (*Saracarine*). *Saracarine* and the previously tentatively identified *N-Acetylchonemorphine* are prob. identical but have not been compared.

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenyl): **Epipachysamine E**
[3697-79-8]
C₂₈H₄₈N₂O 428.7

Alkaloid from *Pachysandra terminalis* (Buxaceae). Cytotoxic to P388 cells. Mp 210-212°. [α]_D +20 (CHCl₃).

N²⁰,N²⁰-Di-Me, N³-tigloyl: **Pachysamine G**
[128255-14-1]
C₂₈H₄₈N₂O 428.7
Alkaloid from *Pachysandra axillaris*. Mp 205-206°. [α]_D²² +11.4 (CHCl₃).

N²⁰,N²⁰-Di-Me, N³-(3,4-dimethyl-2E-pentenyl): **Saligenamide A**
[208835-28-3]
C₃₀H₅₂N₂O 456.754
Alkaloid from *Sarcococca saligna*. [α]_D³⁰ +52 (c, 0.02 in MeOH). λ_{max} 219 (log ε 3.68); 268 (log ε 3.52) (MeOH).

N²⁰,N²⁰-Di-Me, N³-benzoyl: **Epipachysamine D**
[3697-77-6]
C₃₀H₄₆N₂O 450.706
Alkaloid from *Pachysandra terminalis* and *Sarcococca saligna*. Mp 245-248°. [α]_D +13 (CHCl₃). λ_{max} 223 (MeOH) (Berdy).

N²⁰,N²⁰-Di-Me, N³-(3-pyridinecarbonyl): **Epipachysamine B**
[2552-06-9]
C₂₉H₄₅N₃O 451.694
Alkaloid from *Pachysandra terminalis* (Buxaceae). Cytotoxic to P388 cells. Mp 260-262°. [α]_D +16 (CHCl₃).

N³,N³,N²⁰-Tri-Me: **Funtudiamine B**.

Deacetylepachysamine A

[15112-47-7]

C₂₄H₄₄N₂ 360.625Alkaloid from *Funtumia latifolia* and *Pachysandra terminalis* (Apocynaceae, Buxaceae). Mp 96-98° (90-100°). [α]_D²⁰ +23 (+20) (CHCl₃).N³,N³,N²⁰-Tri-Me, N²⁰-Ac: *Epipachysamine A*

[2309-42-4]

C₂₆H₄₆N₂O 402.662Alkaloid from *Pachysandra terminalis* (Buxaceae). Cryst. (Me₂CO). Mp 201-203°. [α]_D²⁰ -17 (-10) (CHCl₃).

▶ AB7910500

N³,N²⁰,N²⁰-Tri-Me: *Dictyophlebin*

[4947-43-7]

C₂₄H₄₄N₂ 360.625Alkaloid from the roots of *Dictyophleba lucida* and from *Funtumia latifolia* (Apocynaceae). Mp 148-151°. [α]_D³⁰ +32 (c, 1.0 in CHCl₃) (+20).N³,N²⁰,N²⁰-Tri-Me, N³-(3,4-dimethyl-2E-pentenyl): *Salignenamide F*

[426815-01-2]

C₃₁H₅₄N₂O 470.78Alkaloid from *Sarcococca saligna*. Powder. λ_{max} 266 (log ε 2.7); 279 (log ε 3.78) (MeOH).N³,N²⁰,N²⁰-Tri-Me, N³-benzoyl: *3-Epipachysamine H*. N³-Methylepipachysamine D

[202583-17-3]

C₃₁H₄₈N₂O 464.733Alkaloid from *Sarcococca saligna* (Buxaceae). [α]_D²⁵ +66 (c, 0.04 in CHCl₃). λ_{max} 232 (log ε 1.62) (MeOH).N³,N²⁰,N²⁰-Tri-Me, N³-E-cinnamoyl:*Salgicinnamide*

[202597-28-2]

C₃₃H₅₀N₂O 490.771Alkaloid from *Sarcococca saligna* (Buxaceae). Powder. Mp 156-160°. [α]_D²⁵ +96 (c, 0.02 in CHCl₃). λ_{max} 279 (log ε 3.78) (MeOH). λ_{max} 226 (ε 10300); 274 (ε 6000) (MeOH) (Berdy).N³,N³,N²⁰,N²⁰-Tetra-Me: *Dihydrokurchessine*

[1172-08-3]

C₂₅H₄₆N₂ 374.652Alkaloid from *Malouetia tamaquarina* (Apocynaceae). Mp 106-108°. [α]_D²⁰ +28 (CHCl₃).N-Hexa-Me: *Malouetine*

[14486-61-4]

[10308-44-8]

C₂₇H₅₂N₂²⁺ 404.721Quaternary alkaloid from *Malouetia bequaertiana* (Apocynaceae).Curarising agent. Noncryst. (as dichloride). [α]_D²⁰ +3 (H₂O) (chloride).**(3β,5β,20S)-form**Bp_{0.05} 220-230°. [α]_D³¹ +13 (c, 1.3 in EtOH).

[152985-06-3, 33467-60-6, 24663-41-0, 85719-26-2]

Crabbé, P. et al., *Bull. Soc. Chim. Belg.*, 1962, **71**, 203-216 (3β,5α,20R-form, di-Ac, tetra-Me)Janot, M.M. et al., *Bull. Soc. Chim. Fr.*, 1962, 111-118 (*Malouetine*)Schmitt, J. et al., *Bull. Soc. Chim. Fr.*, 1964, 761-771 (*synth*)Chien, P.L. et al., *J.O.C.*, 1964, **29**, 315-318


(Chonemorphine, synth, bibl)

Tomita, M. et al., *Tet. Lett.*, 1964, **5**, 1641-1644 (*Pachysamines A, B, isol, struct*)Kikuchi, T. et al., *Tet. Lett.*, 1964, **5**, 1817-1823; 1965, **6**, 1993-1999 (*Pachysandra terminalis constits*)Khuong-Huu, Q. et al., *Bull. Soc. Chim. Fr.*, 1965, 3035-3040 (*occur, bibl*)Chatterjee, A. et al., *Chem. Ind. (London)*,1966, 769-770 (*Sarcococca pruniformis derivs*)Kikuchi, T. et al., *Chem. Pharm. Bull.*, 1967,**15**, 307-316 (*Epipachysamines*)Chatterjee, A. et al., *Indian J. Chem.*, 1967, **5**, 146-150 (*Chonemorphine, isol, ir, pmr, synth, struct*)Sóti, F. et al., *Tet. Lett.*, 1967, 1437-1441

(Dihydrokurchessine)

Kohli, J.M. et al., *Tetrahedron*, 1967, **23**, 3829-3835 (*Saracodine*)Tomita, M. et al., *Yakugaku Zasshi*, 1967, **87**,215-227 (*Pachysamines A, B, N-Methylpachysamine A, Epipachysamines A, B, D, E, N, N-Diacetylepachysamine C, N²⁰-Acetylepachysamine F*)Goutarel, R. et al., *Tetrahedron*, 1968, **24**,7013-7026 (*Saracodine, synth*)Banerji, J. et al., *Indian J. Chem., Sect. B*, 1978,**16**, 346-349 (*Formylchonemorphine, Sarcorine*)Qiu, M. et al., *Zhivuo Xuebao (Acta Bot. Sin.)*,1990, **32**, 626-630; *CA*, **114**, 118538g

(Pachysamines G-H)

Naem, I. et al., *Phytochemistry*, 1996, **43**, 903-906 (*Saracodine*)Atta-ur-Rahman, et al., *Phytochemistry*, 1997,**45**, 861-864; **46**, 771-775 (*Saracodine, N-Formylchonemorphine*)Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1998, **61**,202-206 (*Salgicinnamide, N³-Methylepipachysamine D, Epipachysamine D*)Atta-ur-Rahman, et al., *Nat. Prod. Lett.*, 1998,**11**, 81-91; 297-304 (*N³-Demethylsaracodine, Salgicinnamide A*)Jayasinghe, U.L.B. et al., *Nat. Prod. Lett.*,1998, **12**, 103-109; 2000, **14**, 293-298 (*N²⁰-Formylepipachysamine F*)Funayama, S. et al., *Biol. Pharm. Bull.*, 2000,**23**, 262-264 (*Pachysamine E, Epipachysamines, isol, activity*)Atta-ur-Rahman, et al., *Helv. Chim. Acta*,2002, **85**, 678-688 (*Salgicinnamide F*)Gilani, A.-U.H. et al., *Planta Med.*, 2005, **71**,120-125 (*Saracodine, Sarcorine*)Devkota, K.P. et al., *Chem. Pharm. Bull.*, 2007,**55**, 1397-1401 (*Hookerianamide I*)**Pregnane-3,8,12,14,17,20-hexol** P-602

(3β,5α,8β,12β,14β,17S,20S)-form

C₂₁H₃₆O₆ 384.512**(3β,5α,8β,12β,14β,17S,20S)-form***Dihydrosarcostin. Deacyldrevogenin*

[3080-22-6]

Cryst. (EtOAc). Mp 240-245°. [α]_D²⁰ +41.2 (c, 1.03 in MeOH).12-O-(2-Methylaminobenzoyl), O²⁰-Ac: *Stephanthraniline B*

[69476-88-6]

C₃₁H₄₅NO₈ 559.698Alkaloid from *Stephanotis japonica* aerial parts (Asclepiadaceae). Needles (Me₂CO/diisopropyl ether). Mp 165-168°. [α]_D²⁰ -24.6 (c, 1.14 in CHCl₃). Posn. of ester groups not definitely establ., probable by analogy with Stephanthraniline A (see Pregn-5-ene-3,8,12,14,17,20-hexol, P-610).

20-O-(2-Methylaminobenzoyl), 12-Ac:

Cryst. Mp 157-159°. [α]_D²⁰ -11 (c, 0.1 in MeOH).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-

[β-D-glucopyranosyl-(1→4)-6-

deoxy-3-O-methyl-β-D-allopyranosyl-

(1→4)-β-D-cymaropyranosyl-(1→4)-

β-D-cymaropyranoside]: [906366-98-1]

C₅₈H₉₁NO₂₃ 1170.35Constit. of *Stephanotis mucronata*.Amorph. powder. Mp 183-185°. [α]_D²⁰ -50.1 (c, 0.1 in MeOH).

20-O-(3-Pyridinecarbonyl), 12-cinnamoyl:

Dihydrogagamimine


[72537-21-4]

C₃₆H₄₅NO₈ 619.753Alkaloid from *Stephanotis japonica* aerial parts (Asclepiadaceae).Amorph. [α]_D²⁰ +105 (c, 0.5 in CHCl₃).

Posn. of ester residues not rigorously detd.; probable by analogy with Gagamimine (see Pregn-5-ene-3,8,12,14,17,20-hexol, P-610).

(3β,5ξ,8β,12β,14β,17S,20S)-form20-O-(3-Pyridinecarbonyl), 12-Ac: *Dihydrodrostratine*

[39012-47-0]

C₂₉H₄₁NO₈ 531.645Constit. of *Marsdenia rostrata* (Asclepiadaceae). Cryst. (MeOH/Et₂O). Mp 257-258°. [α]_D²² +55 (c, 0.15 in MeOH).Jaeggi, K.A. et al., *Helv. Chim. Acta*, 1963, **46**, 694-700 (*Dihydrosarcostin*)Gellert, E. et al., *Aust. J. Chem.*, 1974, **27**, 919 (*Dihydrodrostratine*)Terada, S. et al., *Chem. Pharm. Bull.*, 1979, **27**, 2304 (*Dihydrogagamimine, Stephanthraniline B*)Ye, Y. et al., *Bioorg. Med. Chem. Lett.*, 2006,**16**, 4586-4591 (*Stephanotis mucronata saponin*)**Pregnane-3,11,12,14,20-pentol** P-603C₂₁H₃₆O₅ 368.512**(3β,5α,11α,12β,14β,20ξ)-form***Desacylkondurangenin C*

[13872-82-7]

Constit. of *Marsdenia roylei*. Small cryst. (Me₂CO). Mp 225-229°. [α]_D²⁰ +135.6 (c, 0.14 in CHCl₃).11-O-(3-Pyridinecarbonyl), 12-Ac: *Con-*

durangamine A. Kondurangamine A


[55785-23-4]

C₂₉H₄₁NO₇ 515.645Alkaloid from the bark of *Marsdenia condurango* (Asclepiadaceae).

Amorph.

20-O-(3-Pyridinecarbonyl): Condurangamine B. Kondurangamine B

[55785-27-8]

C₂₇H₃₉NO₆ 473.608Alkaloid from *Marsdenia condurango* (Asclepiadaceae). Mp 119-121° Mp 163-165° (double Mp).Berger, S. et al., *Phytochemistry*, 1988, **27**, 1451-1458 (*Condurangoglycosides*)Umehara, K. et al., *Chem. Pharm. Bull.*, 1994, **42**, 611 (*Condurangosides*)**Pregnane-3,12,14,17,20-pentol P-604**C₂₁H₃₆O₅ 368.512**(3β,5α,12β,14β,17βOH,20S)-form****Tomentogenin**

[3513-00-6]

Constit. of *Marsdenia tomentosa*. Cryst. (Me₂CO). Mp 264-267°. [α]_D¹⁶ +36 (c, 0.95 in MeOH).**20-O-(2-Aminobenzoyl), 12-Ac, 3-O-[6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-01-8]C₅₁H₇₉NO₁₇ 978.182Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +10 (c, 0.1 in MeOH). λ_{max} 330 (log ε 3.8) (MeOH).**20-O-(2-Aminobenzoyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871736-97-9]C₅₇H₈₉NO₂₂ 1140.324Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +22 (c, 0.1 in MeOH). λ_{max} 330 (log ε 3) (MeOH).**20-O-(2-Aminobenzoyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-05-2]C₅₆H₈₇NO₂₃ 1142.296Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +20 (c, 0.1 in MeOH). λ_{max} 323 (log ε 2.9) (MeOH).**20-O-(2-Aminobenzoyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-oleandro-****pyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-18-7]C₆₄H₁₀₁NO₂₅ 1284.494Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +28 (c, 0.1 in MeOH). λ_{max} 327 (log ε 2.9) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-02-9]C₅₀H₇₇NO₁₇ 964.155Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +29 (c, 0.1 in MeOH). λ_{max} 265 (log ε 2.7) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-11-0]C₅₀H₇₇NO₁₇ 964.155Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +20 (c, 0.1 in MeOH). λ_{max} 268 (log ε 3.5) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871736-98-0]C₅₆H₈₇NO₂₂ 1126.297Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +11 (c, 0.1 in MeOH). λ_{max} 263 (log ε 4.2) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-10-9]C₅₆H₈₇NO₂₂ 1126.297Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +15 (c, 0.1 in MeOH). λ_{max} 270 (log ε 4) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranosyl-(1→4)-β-D-cymaropyranoside]**: [871737-06-3]C₅₅H₈₅NO₂₃ 1128.269Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +49 (c, 0.1 in MeOH). λ_{max} 263 (log ε 3) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranosyl-(1→4)-β-D-oleandropyranoside]**: [871737-16-5]C₅₅H₈₅NO₂₃ 1128.269Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +48 (c, 0.1 in MeOH). λ_{max} 262 (log ε 3) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-Ac, 3-O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranosyl-(1→4)-β-D-oleandropyranoside]**: [871737-13-2]C₅₅H₈₅NO₂₄ 1144.269Constit. of *Caralluma dalzielii*.Amorph. powder. [α]_D²⁵ +20 (c, 0.1 in MeOH). λ_{max} 270 (log ε 3) (MeOH).**20-O-(3-Pyridinecarbonyl), 12-cinnamoyl: Tomentomin**

[63524-08-3]

C₃₆H₄₅NO₇ 603.754Alkaloid from *Marsdenia tomentosa* (Asclepiadaceae). Plates (EtOAc/hexane). Mp 155-157°. [α]_D²⁰ +137 (c, 0.4 in CHCl₃).**20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: Martsomentoside C**


[238398-64-6]

C₅₇H₈₁NO₁₇ 1052.263Constit. of *Marsdenia tomentosa*. Solid. [α]_D²⁶ +159.2 (c, 0.24 in MeOH).**20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1→4)-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: Marstomentoside T**

[260791-48-8]

C₆₃H₉₁NO₂₂ 1214.405Constit. of *Marsdenia tomentosa*. Solid. [α]_D²² +107.5 (c, 0.25 in MeOH).**20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1→4)-β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: Marstomentoside D**

[238398-66-8]


C₆₃H₉₁NO₂₂ 1214.405Constit. of *Marsdenia tomentosa*. Solid. [α]_D³⁰ +98.9 (c, 0.96 in MeOH).Fukuoka, M. et al., *Chem. Pharm. Bull.*, 1968, **16**, 1634 (*struct*)Seto, H. et al., *Chem. Pharm. Bull.*, 1975, **23**, 1552-1554; 2397-2400; 1976, **24**, 443-449; 1552-1554 (*Tomentosin, Tomentin, Tomentoin, Tomentodin, Deacetyltomentosin, Tomentidin*)Seto, H. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2457-2460; 1977, **25**, 611-616 (*stereochem, biosynth*)Seto, H. et al., *Chem. Pharm. Bull.*, 1977, **25**, 876-878 (*Tomentomin*)Abe, F. et al., *Chem. Pharm. Bull.*, 1999, **47**, 869-875; 2000, **48**, 154-156 (*Isotomentosin, Marstomentosides*)De Leo, M. et al., *Steroids*, 2005, **70**, 573-585 (*Caralluma dalzielii* saponins)**Pregn-4-ene-3,20-diamine P-605****3,20-Diaminopregn-4-ene**C₂₁H₃₆N₂ 316.529**(3β,20S)-form**N²⁰,N²⁰-Di-Me, N³-tigloyl: **Salignarine E**

[299203-57-9]

C₂₈H₄₆N₂O 426.684

Alkaloid from *Sarcococca saligna*.
Brown gum. $[\alpha]_D^{25} +53$ (c, 6.1 in
CHCl₃). λ_{\max} 204 (log ϵ 1.6) (MeOH).
Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000,
63, 1364-1368

Pregn-5-ene-3,20-diamine P-606
3,20-Diaminopregn-5-ene



C₂₁H₃₆N₂ 316.529

(3α,20S)-form

Mp 151°. $[\alpha]_D -70.5$ (CHCl₃).

N³,N²⁰,N²⁰-Tri-Me: Croomionidine

[150036-82-1]

C₂₄H₄₂N₂ 358.609

Alkaloid from roots of *Croomia japonica* (Stemonaceae). Cryst. Mp 150-152°. $[\alpha]_D^{25} -120$ (c, 0.15 in MeOH).

N³,N³,N²⁰,N²⁰-Tetra-Me: α-Kurchessine.

Epiheteroconessine

C₂₅H₄₄N₂ 372.636

Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 148-150°. $[\alpha]_D -28$ (CHCl₃).

(3β,20S)-form

Irehdiamine A

[3614-57-1]

Alkaloid from leaves of *Funtumia elastica* (Apocynaceae). Shows antibacterial activity. Mp 148°. $[\alpha]_D -47$ (c, 1 in 1:1 MeOH/CHCl₃).

▶ TU4812000

Hydrochloride (1:2): Mp 300°. $[\alpha]_D -44$ (c, 1 in MeOH).

N³-Ac, N²⁰-formyl: Holarrifine

[125187-53-3]

C₂₄H₃₈N₂O₂ 386.576

Alkaloid from bark of *Holarrhena antidysenterica*. The 20-config. is not ill. in the abstract and it is assumed here to be the usual 20S-.

N³,N²⁰-Di-Ac:

C₂₅H₄₀N₂O₂ 400.603

Mp 270°. $[\alpha]_D -49$ (c, 1 in MeOH).

N³-Me: Irehdiamine B. Kurchamine

[3734-04-1]

C₂₂H₃₈N₂ 330.556

Alkaloid from *Holarrhena antidysenterica* and *Funtumia elastica* (Apocynaceae). Mp 115-117°. $[\alpha]_D -56$ (c, 1 in CHCl₃).

N³-Me, N³,N²⁰-di-Ac:

C₂₆H₄₂N₂O₂ 414.63

Mp 231°. $[\alpha]_D -48$ (c, 1 in CHCl₃).

N²⁰-Me: Kurchimine

C₂₂H₃₈N₂ 330.556

Alkaloid from *Holarrhena antidysenterica* (Apocynaceae). Mp 104-106°. $[\alpha]_D -21$ (c, 1.29 in CHCl₃).

N³,N³-Di-Me: Irehdiamine C

[52617-38-6]

C₂₃H₄₀N₂ 344.582

Alkaloid from *Funtumia latifolia* (Apocynaceae). Mp 105-106°. $[\alpha]_D -47$ (CHCl₃). Incorr. descr. as an N-monomethyl compd. in CA.

N³,N³-Di-Me, N²⁰-formyl: N²⁰-Formylirehdiamine C

[124162-11-4]

C₂₄H₄₀N₂O 372.593

Alkaloid from *Sarcococca zeylanica*.

Mp 220-222°. $[\alpha]_D^{22} +2.1$ (c, 2.9 in CHCl₃).

N²⁰,N²⁰-Di-Me: Irehdiamine F

[52617-39-7]

C₂₃H₄₀N₂ 344.582

Alkaloid from *Vahadenia laurentii* root cortex (Apocynaceae).

N²⁰,N²⁰-Di-Me, N³-Ac: N³-Acetylirehdiamine F

[52342-67-3]

C₂₅H₄₂N₂O 386.62

Alkaloid from stem bark of *Didymeleis* cf. *madagascariensis* (Didymelaceae).

N³,N³,N²⁰-Tri-Me, N²⁰-Ac: Saracocine.

Alkaloid C†

[15437-93-1]

C₂₆H₄₄N₂O 400.646

Alkaloid from *Sarcococca pruniformis* and *Sarcococca saligna*. Cryst.

(Me₂CO). Mp 231°. $[\alpha]_D^{20} -58$

(CHCl₃). λ_{\max} 202 (log ϵ 3.2)

(MeOH).

N³,N³,N²⁰,N²⁰-Tetra-Me: Kurchessine.

Irehdiamine I. Saracodinine

[6869-45-0]

C₂₅H₄₄N₂ 372.636

Alkaloid from *Holarrhena antidysenterica*, *Pachysandra terminalis*, *Sarcococca pruniformis* and *Malouetia tamaquarina* (Apocynaceae, Buxaceae). Mp 140-141°. $[\alpha]_D^{30} -37$ (c, 1.9 in CHCl₃).

(3α,20R)-form [17934-39-3]

Mp 158°. $[\alpha]_D -66$ (CHCl₃).


(3β,20R)-form [17934-37-1]

Mp 140°. $[\alpha]_D -64$ (CHCl₃).

Truong-Ho, M. et al., *Bull. Soc. Chim. Fr.*, 1963, 594 (*Irehdiamine A, Irehdiamine B*)
Läbler, L. et al., *Coll. Czech. Chem. Comm.*, 1963, 28, 2345 (*isol. struct*)
Khuong-Huu, Q. et al., *Bull. Soc. Chim. Fr.*, 1965, 3035 (*isol. struct*)
Goutarel, R. et al., *Bull. Soc. Chim. Fr.*, 1967, 4575-4582 (*synth*)
Kohli, J.M. et al., *Tetrahedron*, 1967, 23, 3829 (*Saracocine*)
Goutarel, R. et al., *Tetrahedron*, 1968, 24, 7013 (*synth*)
Longeviall, P. et al., *Org. Mass Spectrom.*, 1970, 3, 803 (*ms*)
Lamotte, J. et al., *Acta Cryst. B*, 1977, 33, 2392 (*cryst struct*)
Sánchez, V. et al., *Bull. Soc. Chim. Fr.*, Part II, 1984, 71 (*N-Acetylirehdiamine F*)
Siddiqui, S. et al., *Pak. J. Sci. Ind. Res.*, 1989, 32, 1; *CA*, 112, 95430h (*Holarrifine*)
Lin, W.H. et al., *Yaoxue Xuebao*, 1993, 28, 202; *CA*, 119, 156210e (*Croomionidine*)
Atta-ur-Rahman, et al., *Phytochemistry*, 1997, 45, 861-864 (*Saracocine*)
Jayasinghe, U.L.B. et al., *Nat. Prod. Lett.*, 2001, 15, 151-155 (*N²⁰-Formylirehdiamine C*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, IGH700

Pregn-14-ene-3,20-diamine P-607
3,20-Diaminopregn-14-ene



C₂₁H₃₆N₂ 316.529

(3β,5α,20S)-form

N³,N²⁰,N²⁰-Tri-Me: 20-(Dimethylamino)-3-(methylamino)pregn-14-ene.

Hookerianamine A


[771533-18-7]

C₂₄H₄₂N₂ 358.609

Alkaloid from *Sarcococca hookeriana*. Cholinesterase inhibitor. Powder. $[\alpha]_D^{24} -60$ (c, 0.04 in MeOH). λ_{\max} 211 (log ϵ 3.6) (MeOH).

Choudhary, M.I. et al., *Helv. Chim. Acta*, 2004, 87, 1099-1108 (*Hookerianamine A*)

Pregn-16-ene-3,20-diamine P-608
3,20-Diaminopregn-16-ene



C₂₁H₃₆N₂ 316.529

(3β,5α,20S)-form

N²⁰,N²⁰-Di-Me, N³-(3-methyl-2-butenoyl): Salignarine D

[299203-56-8]

C₂₈H₄₆N₂O 426.684

Alkaloid from *Sarcococca saligna*. Powder. $[\alpha]_D^{23} +66$ (c, 0.2 in CHCl₃). λ_{\max} 210 (log ϵ 1.7) (MeOH).

N³,N²⁰,N²⁰-Tri-Me: 5,6-Dihydrosarconidine

C₂₄H₄₂N₂ 358.609

Alkaloid from *Sarcococca saligna*.

Cholinesterase inhibitor. Powder. $[\alpha]_D^{20} -60$ (c, 0.2 in CHCl₃). λ_{\max} 212 (log ϵ 2.66) (MeOH).

N³,N²⁰,N²⁰-Tri-Me, N³-(3,4-dimethyl-2E-pentenoyl): Salignenamide E

[426815-00-1]

C₃₁H₅₂N₂O 468.765

Alkaloid from *Sarcococca saligna*. Amorph. solid. λ_{\max} 219 (log ϵ 3.7); 270 (log ϵ 3.52) (MeOH).

Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000,


63, 1364-1368 (*Salignarine D*)

Atta-ur-Rahman, et al., *Helv. Chim. Acta*,

2002, 85, 678-688 (*Salignenamide E*)

Atta-ur-Rahman, et al., *Helv. Chim. Acta*,
2004, **87**, 439-448 (5,6-Dihydrosarconidine)

Pregn-6-ene-3,5,8,12,14,17,20-heptol P-609



C₂₁H₃₄O₇ 398.495

(3β,5α,8β,12β,14β,17β,20S)-form


20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside L** [185507-42-0]
C₅₇H₇₉NO₁₉ 1082.246
Constit. of *Stephanotis lutchuensis*. Cryst. Mp 142-144°. [α]_D²⁰ +80.2 (c, 3.8 in CHCl₃). λ_{max} 206 (log ε 4.29); 218 (log ε 4.34); 263 (log ε 4.2); 280 (log ε 4.2) (EtOH).

20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside N** [185507-44-2]
C₅₇H₇₉NO₁₉ 1082.246
Constit. of *Stephanotis lutchuensis*. Cryst. Mp 148-150°. [α]_D²⁰ +64.2 (c, 0.4 in CHCl₃). λ_{max} 207 (log ε 4.21); 218 (log ε 4.24); 280 (log ε 4.13) (EtOH).

20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside P** [185507-46-4]
C₆₄H₉₁NO₂₂ 1226.416
Constit. of *Stephanotis lutchuensis*. Cryst. Mp 152-154°. [α]_D²⁰ +96 (c, 2.3 in CHCl₃). λ_{max} 206 (log ε 4.34); 218 (log ε 4.37); 281 (log ε 4.26) (EtOH).

Yoshikawa, K. et al., *Chem. Pharm. Bull.*,
1996, **44**, 2243-2248 (*Stephanosides*)

Pregn-5-ene-3,8,12,14,17,20-hexol P-610



C₂₁H₃₄O₆ 382.496

(3β,12β,14β,17βOH,20S)-form

Sarcostin
[18607-76-6]

Found in *Asclepias glaucophylla*, *Sarcos-*

temma viminale, *Pachycarpus lineolatus*, *Cynanchum caudatum*, *Cynanchum wilfordii*, *Cynanchum paniculatum*, *Marsdenia tomentosa*, *Metaplexis japonica* and other plants in the Asclepiadaceae. Mp 150° (170°) Mp 250-255° (266-267°) (double Mp). [α]_D²³ -67 (c, 1 in MeOH).

12-O-(2-Methylaminobenzoyl), 20-Ac:

Stephanthraneline A
[65429-40-5]

C₃₁H₄₃NO₈ 557.683

Alkaloid from stems of *Stephanotis japonica* (Asclepiadaceae). Cryst. (Me₂CO/hexane). Mp 170-173°. [α]_D +17.9 (c, 0.089 in CHCl₃).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-[6-deoxy-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside B** [183116-98-5]
C₅₂H₇₉NO₁₈ 1006.192

Constit. of *Stephanotis lutchuensis*. Cryst. Mp 140-142°. [α]_D²⁰ +2 (c, 3.3 in CHCl₃). λ_{max} 221 (log ε 5.32); 255 (log ε 4.96); 280 (log ε 4.17); 350 (log ε 4.65) (EtOH).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside D** [183117-00-2]
C₅₂H₇₉NO₁₇ 990.193

Constit. of *Stephanotis lutchuensis*. Cryst. Mp 148-150°. [α]_D²⁰ -48.8 (c, 1 in CHCl₃). λ_{max} 221 (log ε 5.2); 256 (log ε 4.75); 280 (log ε 4.2); 352 (log ε 4.77) (EtOH).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside E** [183117-02-4]
C₅₂H₇₉NO₁₈ 1006.192

Constit. of *Stephanotis lutchuensis*. Cryst. Mp 153-155°. [α]_D²⁰ -4.8 (c, 3.8 in CHCl₃). λ_{max} 220 (log ε 5.32); 256 (log ε 4.76); 280 (log ε 4.25); 350 (log ε 4.75) (EtOH).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside C** [183116-99-6]
C₅₂H₇₉NO₁₈ 1006.192

Constit. of *Stephanotis lutchuensis*. Cryst. Mp 151-153°. [α]_D²⁰ +5.9 (c, 1.5 in CHCl₃). λ_{max} 220 (log ε 5.33); 255 (log ε 4.8); 281 (log ε 4.2); 351 (log ε 4.68) (EtOH).

20-O-(2-Methylaminobenzoyl), 12-Ac, 3-O-[β-D-thevetopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Stephanoside A** [183116-97-4]
C₅₉H₉₁NO₂₁ 1150.362

Constit. of *Stephanotis lutchuensis*. Cryst. Mp 143-145°. [α]_D²⁰ +12 (c, 1.9 in CHCl₃). λ_{max} 206 (log ε 4.05); 222 (log ε 4.17); 257 (log ε 3.74); 279 (log ε 3.61); 354 (log ε 3.38) (EtOH).

12-O-(3-Pyridinecarbonyl): **Gagamine**

[289634-18-0]

C₂₇H₃₇NO₇ 487.592

Constit. of *Cynanchum caudatum*. Amorph. solid. Mp 257-258° dec. [α]_D²⁵ -12.7 (c, 1 in CHCl₃). λ_{max} 221 (log ε 3.61); 263 (log ε 4.13) (MeOH).

12-O-(3-Pyridinecarbonyl), 3-O-[α-L-cymaropyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]:
C₅₅H₈₅NO₁₉ 1064.272

Constit. of *Cynanchum caudatum*. Amorph. powder. [α]_D²⁰ -13.9 (c, 0.61 in MeOH).

12-O-(3-Pyridinecarbonyl), 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: **Hainaneoside A** [156891-23-5]
C₆₁H₉₅NO₂₅ 1242.413

Constit. of *Marsdenia hainanensis*. Amorph. powder (MeOH). Mp 166-167°. [α]_D²⁵ +17.6 (c, 0.6 in Me₂CO).

12-O-(3-Pyridinecarbonyl), 20-Ac: [165746-37-2]
C₂₉H₃₉NO₈ 529.629
Constit. of *Leptadenia hastata*. Cryst. Mp 155-160°. [α]_D²⁵ +145.5 (c, 1 in MeOH).

12-O-(3-Pyridinecarbonyl), 20-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: [175667-74-0]
C₆₃H₉₇NO₂₆ 1284.45

Constit. of *Leptadenia hastata*. Amorph. powder. [α]_D²⁵ +100.5 (c, 1 in MeOH).

12-O-(3-Pyridinecarbonyl), 20-Ac, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranoside]: [175667-89-7]
C₆₃H₉₇NO₂₆ 1284.45

Constit. of *Leptadenia hastata*. Amorph. powder. [α]_D²⁵ +160 (c, 1 in MeOH).

12-O-(3-Pyridinecarbonyl), 20-E-cinnamoyl: **Isogagamine** [175702-35-9]
C₃₆H₄₃NO₈ 617.738

Constit. of *Leptadenia hastata*. Cryst. Mp 155-160°. [α]_D²⁵ +135.5 (c, 1 in MeOH).

12-O-(3-Pyridinecarbonyl), 20-E-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: [906366-99-2]
C₆₃H₈₉NO₂₃ 1228.389


Constit. of *Stephanotis mucronata*. Amorph. powder. Mp 178-180°. [α]_D²⁰ -23.1 (c, 0.1 in MeOH).

12-O-(3-Pyridinecarbonyl), 20-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1→4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1→4)-β-D-oleandropyranosyl-

- (1 → 4)-β-D-cymaropyranoside]: **Hainaneoside B**
[185217-99-6]
C₆₃H₈₉NO₂₃ 1228.389
Constit. of *Marsdenia hainanensis*.
Amorph. powder (MeOH). Mp 174-175°. [α]_D²⁵ +86.2 (c, 2.7 in Me₂CO).
- 12-O-(3-Pyridinecarbonyl), 20-cinnamoyl, 17-Ac, 3-O-[β-D-glucopyranosyl-(1 → 4)-6-deoxy-3-O-methyl-β-D-allopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]: [175667-94-4]
C₇₂H₁₀₃NO₂₇ 1414.596
Constit. of *Leptadenia hastata*.
Amorph. powder. [α]_D²⁵ +135.5 (c, 1 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-Ac: **Rostratine**
[39007-91-5]
C₂₉H₃₉NO₈ 529.629
Alkaloid from *Marsdenia rostrata* (Asclepiadaceae). Cryst. (MeOH/Et₂O). Mp 259-260°. [α]_D²⁵ +49 (c, 1.0 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-E-cinnamoyl: **Gagaminine**
[41060-37-1]
C₃₆H₄₃NO₈ 617.738
Alkaloid from *Metaplexis japonica*, *Marsdenia condurango*, *Marsdenia tomentosa*, *Stephanotis japonica* and *Cynanchum caudatum* (Asclepiadaceae). Mp 160-170°. [α]_D²⁰ +143.8 (c, 0.7 in CHCl₃).
- 20-O-(3-Pyridinecarbonyl), 12-E-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1 → 4)-β-D-thevetopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Marstomentoside O**
[260789-62-6]
C₆₃H₈₉NO₂₃ 1228.389
Constit. of *Marsdenia tomentosa*. Solid. [α]_D²⁶ +43.6 (c, 0.7 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-Z-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1 → 4)-β-D-thevetopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Marstomentoside P**
[260789-63-7]
C₆₃H₈₉NO₂₃ 1228.389
Constit. of *Marsdenia tomentosa*. Solid. [α]_D²⁶ +44.7 (c, 0.6 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[α-L-cymaropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₅₇H₇₉NO₁₇ 1050.248
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +111 (c, 0.41 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[α-L-cymaropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₅₇H₇₉NO₁₇ 1050.248
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +86.8 (c, 0.82 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]: [174097-88-2]
C₅₇H₇₉NO₁₇ 1050.248
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +127 (c, 2 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₅₇H₇₉NO₁₇ 1050.248
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +96 (c, 0.75 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Stephanoside H**
[183117-05-7]
C₅₇H₇₉NO₁₈ 1066.247
Constit. of *Stephanotis lutchuensis*.
Cryst. Mp 160-162°. [α]_D²⁰ +92 (c, 1.1 in CHCl₃). λ_{max} 206 (log ε 4.18); 217 (log ε 4.3); 280 (log ε 4.23) (EtOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Stephanoside G**
[183117-04-6]
C₅₇H₇₉NO₁₈ 1066.247
Constit. of *Stephanotis lutchuensis*.
Cryst. Mp 150-152°. [α]_D²⁰ +60.6 (c, 1.1 in CHCl₃). λ_{max} 206 (log ε 4.18); 218 (log ε 4.23); 280 (log ε 4.11) (EtOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-cymaropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₆₄H₉₁NO₂₀ 1194.418
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +130 (c, 0.34 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[6-deoxy-3-O-methyl-β-D-allopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranoside]: [175667-83-1]
C₆₄H₉₁NO₂₁ 1210.417
Constit. of *Leptadenia hastata*.
Amorph. powder. [α]_D²⁵ +121 (c, 1 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₆₄H₉₁NO₂₀ 1194.418
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁰ +101 (c, 0.7 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Stephanoside F**
[183117-03-5]
C₆₄H₉₁NO₂₁ 1210.417
Constit. of *Stephanotis lutchuensis*.
Cryst. Mp 152-154°. [α]_D²⁰ +90.8 (c, 2.5 in CHCl₃). λ_{max} 217 (log ε 4.3); 281 (log ε 4.22) (EtOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[α-L-cymaropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₇₁H₁₀₃NO₂₃ 1338.588
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²² +74.1 (c, 0.74 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1 → 4)-α-L-cymaropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-α-L-diginyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Wilfoside GIG**
[100776-59-8]
C₇₀H₁₀₁NO₂₅ 1356.56
Constit. of *Cynanchum wilfordii*.
Amorph. powder. Mp 164-167°. [α]_D¹⁷ +28.2 (c, 0.98 in CHCl₃).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-glucopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]: [188826-02-0]
C₇₀H₁₀₁NO₂₅ 1356.56
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²⁴ +91.8 (c, 0.73 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]:
C₇₁H₁₀₃NO₂₃ 1338.588
Constit. of *Cynanchum caudatum*.
Amorph. powder. [α]_D²² +87.7 (c, 0.59 in MeOH).
- 20-O-(3-Pyridinecarbonyl), 12-cinnamoyl, 3-O-[β-D-thevetopyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-oleandropyranosyl-(1 → 4)-β-D-cymaropyranosyl-(1 → 4)-β-D-cymaropyranoside]: **Marstomentoside A**
[238398-33-9]
C₇₁H₁₀₃NO₂₄ 1354.587
Constit. of *Marsdenia tomentosa*. Solid. [α]_D³⁰ +74.1 (c, 0.32 in MeOH).
- [1832-88-8, 3513-04-0]
Yamagishi, T. et al., *Chem. Pharm. Bull.*, 1972, **20**, 2289-2292 (Gagaminin)
Farnsworth, N.R. et al., *J. Pharm. Sci.*, 1973, **62**, 1028 (nomencl. Wilforine)
Gellert, E. et al., *Aust. J. Chem.*, 1974, **27**, 919-921 (Rostratine)
Terada, S. et al., *Chem. Pharm. Bull.*, 1977, **25**, 2802-2805 (*Stephanotis japonica constits*)
Tsukamoto, S. et al., *Chem. Pharm. Bull.*, 1985, **33**, 2294-2304 (*Wilfosides, cmr*)
Kaur, K.J. et al., *Indian J. Chem., Sect. B*, 1985, **24**, 1053-1056 (*Orine*)
Ma, B.X. et al., *Chin. Chem. Lett.*, 1994, **5**, 209-210 (*Hainaneoside A*)
Warashina, T. et al., *Chem. Pharm. Bull.*, 1995, **43**, 977-982; 1996, **44**, 358-363 (*Cynanchum caudatum saponins*)
Aquino, R. et al., *J. Nat. Prod.*, 1995, **58**, 672-679; 1996, **59**, 555-564 (*Leptadenia hastata constits*)


Yoshikawa, K. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1790-1796; 1998, **46**, 1239-1243 (*Stephanosides*, *Gymnepregoside F*)
 Ma, B. *et al.*, *J. Nat. Prod.*, 1997, **60**, 134-138 (*Hainaneosides*)
 Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 869-875; 2000, **48**, 154-156 (*Isokidjolidinin*, *Marstomentosides*)
 Lee, D.-U. *et al.*, *Planta Med.*, 2000, **66**, 480-482 (*Gagamine*, *Gagaminine*)
 Ye, Y. *et al.*, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 4586-4591 (*Stephanotis mucronata saponin*)

Preliisoclinamide 2 P-611
 [104928-31-6]



$C_{33}H_{43}N_7O_6S_2$ 697.878
 Isol. from the tunicate *Lissoclinum patella*. $[\alpha]_D +18.1$ (c, 2.24 in $CHCl_3$).
 Sesin, D.F. *et al.*, *Bull. Soc. Chim. Belg.*, 1986, **95**, 853

Diaphus Preluciferin P-612
 [107503-09-3]



$C_{32}H_{29}N_3O_9$ 599.596
 Bioluminescent subst. from the liver of the myctophina fish *Diaphus elucens*. Also found in *Diaphus coeruleus*, *Diaphus suborbitalis*, *Benthosema fibulata* and *Myctophum asperum*.
 Inoue, S. *et al.*, *Chem. Lett.*, 1987, 417 (*isol, uv, pmr, struct*)
 Teranishi, K. *et al.*, *Bioorg. Chem.*, 2007, **35**, 82-111 (*rev*)

Premavacurine P-613
 [1360-59-4]


Struct. unknown. Alkaloid from *Strychnos solimoesana* (Loganiaceae). Rose-violet col. with $Ce(IV)$.
 Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148

Premine P-614
 $C_{14}H_{15}NO$ 213.279

Struct. unknown. Alkaloid from the stem bark of *Premna integrifolia*. Amorph. Mp

82° Mp 211-213° (as hydrochloride).
 Basu, N.K. *et al.*, *J. Am. Pharm. Assoc.*, 1947, **36**, 389-391; *CA*, **42**, 3535i (*isol*)


Prenostodione P-615



$C_{19}H_{15}NO_5$ 337.331
 Precursor of Nostodione A, N-323. Isol. from a *Nostoc* sp. (TAU strain IL-235). Uv absorbant. Yellow oil. λ_{max} 217 (ϵ 29200); 230 (ϵ 24150); 287 (ϵ 17000); 318 (ϵ 15640) (MeOH).
 Ploutno, A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 544-545

Prenylcitpressine P-616

1,3,6-Trihydroxy-5-methoxy-10-methyl-4-(3-methyl-2-butenyl)-9-(10H)-acridinone, 9CI. 1,3,6-Trihydroxy-5-methoxy-10-methyl-4-C-prenylacridone [81525-60-2]




$C_{20}H_{21}NO_5$ 355.39
 Alkaloid from the root bark of *Citrus depressa* and *Citrus grandis* f. *hakunikuyu* (Rutaceae). Yellow plates + $\frac{1}{2}$ H_2O (Et_2O). Mp 160-162°.
3-Me ether: Grandisinine. 1,6-Dihydroxy-3,5-dimethoxy-10-methyl-4-(3-methyl-2-butenyl)-9-(10H)-acridinone [87959-95-3]
 $C_{21}H_{23}NO_5$ 369.416
 Alkaloid from the root bark of *Citrus grandis* f. *hakunikuyu* (Rutaceae). Yellow plates (Me_2CO). Mp 194-196°.
3,6-Di-Me ether: Baiyumine B. 1-Hydroxy-3,5,6-trimethoxy-10-methyl-4-(3-methyl-2-butenyl)-9-(10H)-acridinone, 9CI. 1-Hydroxy-3,5,6-trimethoxy-10-methyl-4-prenylacridone [109030-95-7]
 $C_{22}H_{25}NO_5$ 383.443
 Alkaloid from the root of *Citrus grandis* f. *hakunikuyu* (Rutaceae). Yellow plates (Et_2O). Mp 145-147°.
Tri-Me ether:
 Yellow syrup.

A^{3'}-Isomer, 2'-hydroxy, 3-Me ether: 1,6-Dihydroxy-3,5-dimethoxy-4-(2-hydroxy-3-methyl-3-butenyl)-10-methylacri-


done. Buntanmine A [126622-49-9]
 $C_{21}H_{23}NO_6$ 385.416
 Alkaloid from the stem bark of *Citrus grandis* f. *buntan* (Rutaceae). Yellowish granules (Me_2CO). Mp 201-202°. $[\alpha]_D -116.7$ (c, 0.12 in MeOH).
 Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 895 (*isol, uv, ir, pmr, ms, struct*)
 Wu, T.-S. *et al.*, *Phytochemistry*, 1983, **22**, 1493; 1987, **26**, 871 (*isol, uv, ir, pmr, cmr, ms, struct, derivs*)
 Huang, S.C. *et al.*, *Phytochemistry*, 1989, **28**, 3574 (*Buntanmine A*)

6-Prenyl-1H-indole-3-carboxylic acid P-617
6-(3-Methyl-2-butenyl)-1H-indole-3-carboxylic acid




$C_{14}H_{15}NO_2$ 229.278
 Alkaloid prod. by *Colletotrichum* sp. isol. from *Artemisia annua*. Needles. Mp 54-55°.
 Lu, H. *et al.*, *Plant Sci. (Limerick, Irel.)*, 2000, **151**, 67-73 (*isol, pmr, cmr*)

9-Prenylpaxilline P-618
9-Isopentenylpaxilline [163019-15-6]



$C_{32}H_{41}NO_4$ 503.68
 Numbering systems vary; also called 21-isopentenyl. Metab. of the ascostromata of *Eupenicillium shearii* NRRL3324. Insecticide. Mycotoxin. Oil. $[\alpha]_D -12$ (c, 0.30 in $CHCl_3$). λ_{max} 236 (ϵ 9800); 283 (ϵ 2500) (MeOH) (Berdy).
 Belofsky, G.N. *et al.*, *Tetrahedron*, 1995, **51**, 3959 (*isol, uv, pmr, cmr, ms*)
 Smith, A.B. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 3908-3938 (*synth, abs config*)


Preoxazin 7 P-619
 [1001439-33-3]




$C_{22}H_{21}N_3O_5$ 407.425
 Isol. from toxic mussels, *Mytilus gallo-provincialis*.
 Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2007, 5434-5439 (*isol, synth, pmr, cmr, ms*)

Prepatellamide A

[350593-23-6]

Absolute
ConfigurationC₃₅H₅₂N₈O₇S₂ 760.977Isol. from the tunicate *Lissoclinum patella*. Amorph. solid. [α]_D²⁰ +80.1 (c, 0.1 in MeOH).Fu, X. *et al.*, *Sci. China, Ser. B*, 2000, **43**, 643-648; *CA*, **135**, 105198e (isol, pmr, cmr)**Prepatellamide B formate**

[104928-32-7]


C₃₉H₅₀N₈O₈S₂ 823.005Cyclic peptide antibiotic. Isol. from the tunicate *Lissoclinum patella*. [α]_D²⁰ +36.5 (c, 0.3 in CHCl₃).Sesin, D.F. *et al.*, *Bull. Soc. Chim. Belg.*, 1986, **95**, 853 (isol, pmr, cmr)**Prepsammaplin A**

P-622

Methyl 3-oxo-2-oxa-7,8-dithia-4,11-diazadodecan-12-oate, 9CI. Dimethyl (dithiodiethylene)dicarbamate, 8CI [28138-84-3]

MeOOCNHCH₂CH₂-S-S-CH₂CH₂NHCOOMeC₈H₁₆N₂O₄S₂ 268.357Isol. from the marine sponge *Psammoplysilla purpurea*. Oil.Jimenez, C. *et al.*, *Tetrahedron*, 1991, **47**, 2097 (isol, pmr, cmr, ms)**Prepseudopalmanine**


P-623

C₂₁H₂₁NO₇ 399.399**(±)-form [75091-35-9]**Alkaloid from the stems of *Berberis darwinii* (Berberidaceae). Cryst. (Et₂O) or amorph. Mp 160-161°.

[95530-37-3]

Manikumar, G. *et al.*, *Heterocycles*, 1980, **14**, 827 (synth, ir, pmr)Valencia, E. *et al.*, *Tetrahedron*, 1984, **40**, 3957 (isol, uv, ir, pmr, ms)**Presecamine**


[29089-93-8]

Probable
Relative
ConfigurationC₄₂H₅₂N₄O₄ 676.897Minor alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Noncryst. λ_{max} 227 (log ε 4.3); 288 (sh) (log ε 4.07); 295 (log ε 4.14); 329 (log ε 4.17) (EtOH).**19,20- Andlor 19',20'-dihydro: Dihydro-presecamine**C₄₂H₅₄N₄O₄ 678.913Minor alkaloid from leaves of *Rhazya stricta* (Apocynaceae). Noncryst. Possibly a mixt. of 2 components.**19,19',20,20'-Tetrahydro: Tetrahydro-presecamine**

[56875-50-4]

C₄₂H₅₆N₄O₄ 680.929Alkaloid from *Pandaca minutiflora*, the leaves of *Hunteria elliotii* and *Rhazya stricta*, and from the roots of *Rhazya orientalis* and *Amsonia tabernaemontana* (Apocynaceae). Noncryst. [α]_D²⁰ -1.9 (95% EtOH). [α]_D²⁰ -10 (95% EtOH). [α]_D²⁰ +57 (EtOH). Stereochem. undefined, isolates have various opt. rotns., the (-)-form being probably largely racemic. Rearranges quantitatively to Secamine, S-174 in dil. HCl at r.t. λ_{max} 227 (log ε 4.3); 288 (sh) (log ε 4.07); 295 ; 329 (log ε 4.17) (EtOH).Cordell, G.A. *et al.*, *Chem. Comm.*, 1970, 191-192 (uv, ir, pmr, ms, struct)Zsaron, B. *et al.*, *Phytochemistry*, 1975, **14**, 1438-1439 (tetrahydro, uv, ir, ms)Petitfrère, N. *et al.*, *Phytochemistry*, 1975, **14**, 1648-1649 (tetrahydro, isol, uv, ir, pmr, ms)Cordell, G.A. *et al.*, *J. Indian Chem. Soc.*, 1978, **55**, 1083-1091 (config)**Preskimmianine**

P-625

3-(3,3-Dimethylallyl)-4,7,8-trimethoxy-2(1H)-quinolinone. 4,7,8-Trimethoxy-3-prenyl-2(1H)-quinolinone. *Dasyrcarpamine* [38695-41-9]C₁₇H₂₁NO₄ 303.357Alkaloid from root of *Dictamnus albus* and *Dictamnus dasycarpus* (Rutaceae). Cryst. (C₆H₆). Mp 151-152°.**N-Me: N-Methylpreskimmianine**

[76525-28-5]

C₁₈H₂₃NO₄ 317.384Alkaloid from the stem bark of *Vepris louisii* (Rutaceae). Plates (hexane). Mp 88-89°.**Δ^{3',4'}-Isomer, 2'-ξ-hydroxy: 3-(2-Hydroxy-3-methyl-3-butenyl)-4,7,8-trimethoxy-2(1H)-quinolinone. Isoptelefoline**

[27745-39-7]

C₁₈H₂₃NO₅ 333.383Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (Me₂CO/hexane). Mp 145-146°.**3',4'-Dihydro, 2'-oxo, 3'-hydroxy: Veprisilone**


[82848-01-9]

C₁₈H₂₃NO₆ 349.383Alkaloid from the trunk bark of *Vepris louisii* (Rutaceae). Prisms (EtOAc). Mp 135-136°.Reisch, J. *et al.*, *Tet. Lett.*, 1970, 1945 (Isoptelefoline)Storer, R. *et al.*, *Tetrahedron*, 1973, **29**, 1217 (Preskimmianine)Ayafor, J.F. *et al.*, *Phytochemistry*, 1982, **21**, 955; 2733 (Veprisilone, N-Methylpreskimmianine)Ramesh, M. *et al.*, *Tetrahedron*, 1984, **40**, 4041-4049 (synth, pmr)**Pretazettine**

P-626

Isotazettine. PTZ

[17322-84-8]

Absolute
ConfigurationC₁₈H₂₁NO₅ 331.368Alkaloid from *Sprekelia formosissima*, *Sprekelia formosissima*, *Ruta graveolens*, *Narcissus bicolor*, *Hymenocallis littoralis* and *Ismene calithina* (preferred genus name *Hymenocallis*). Prob. very widely distributed in the Amaryllidaceae. Antineoplastic agent active against Rauscher leukaemia virus and others. Noncryst. [α]_D²⁴ +180 (c, 0.2 in CHCl₃). Log P -0.65 (uncertain value) (calc). Easily rearr. to Tazettine, T-53. λ_{max} 238 (ε 5060); 291 (ε 4300) (EtOH) (Berdy).Hydrochloride: Mp 224-225°. [α]_D²⁴ +30.3 (c, 0.15 in H₂O).**O⁸-Me: 8-O-Methylpretazettine**C₁₉H₂₃NO₅ 345.394Alkaloid from *Eucharis amazonica*. Amorph. solid.**O⁸-Et, N-de-Me: 8β-Ethoxy-N-demethyl-pretazettine**

[250253-94-2]

C₁₉H₂₃NO₅ 345.394Alkaloid from *Crinum bulbispermum*.

- Amorph. $[\alpha]_D^{28} +34$ (c, 0.14 in CHCl_3).
- 8-Ketone (lactone): 3-Epimacronine**
[17322-73-5]
 $\text{C}_{18}\text{H}_{19}\text{NO}_5$ 329.352
Isol. from *Sprekelia formosissima* and bulbs of *Narcissus bicolor* (Amaryllidaceae). Prisms (Me_2CO). Mp 129-131°. $[\alpha]_D^{24} +276$ (c, 0.95 in CHCl_3).
- 8-Ketone (lactone), N-de-Me: Ungspirolidone. N-Demethyl-3-epimacronine**
[17245-18-0]
 $\text{C}_{17}\text{H}_{17}\text{NO}_5$ 315.325
Alkaloid from *Ungernia spiralis* (Amaryllidaceae). Cryst. ($\text{Et}_2\text{O}/\text{Me}_2\text{CO}$) (synthetic). Mp 148-149° (nat.) Mp 154-155° (synthetic). $[\alpha]_D^{20} +105$ (c, 0.6 in CHCl_3) (nat.). $[\alpha]_D^{24} +207$ (c, 0.36 in CHCl_3) (synthetic). Abs. config. incorrectly shown in the 1977 paper. λ_{max} 228 (ϵ 31400); 268 (ϵ 6000); 308 (ϵ 6800) (95% EtOH).
- 8-Ketone (lactone), O-de-Me: O-Demethyl-3-epimacronine**
[233767-23-2]
 $\text{C}_{17}\text{H}_{17}\text{NO}_5$ 315.325
Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. powder. $[\alpha]_D +120$ (c, 0.2 in MeOH). λ_{max} 228 (log ϵ 4.34); 262 (sh) (log ϵ 3.66); 309 (log ϵ 3.42) (MeOH).
- 1,2-Dihydro, 8-ketone (lactone): 1,2-Dihydro-3-epimacronine**
[64507-57-9]
 $\text{C}_{18}\text{H}_{21}\text{NO}_5$ 331.368
Alkaloid from *Ungernia spiralis* (Amaryllidaceae). Cryst. Mp 98-99°. $[\alpha]_D +11$ (CHCl_3). λ_{max} 227 (log ϵ 4.36); 272 (log ϵ 3.85); 314 (log ϵ 3.73) (EtOH).
- 1,2-Dihydro, 8-ketone (lactone), N-de-Me: Ungspirolidine**
[66408-49-9]
 $\text{C}_{17}\text{H}_{19}\text{NO}_5$ 317.341
Alkaloid from *Ungernia spiralis* (Amaryllidaceae). Mp 142-143°. $[\alpha]_D^{20} +11$ (c, 0.45 in CHCl_3). λ_{max} 230 (log ϵ 4.3); 270 (log ϵ 3.7); 309 (log ϵ 3.63) (EtOH).
- 3-Epimer: Precriwelline**
[17245-16-8]
 $\text{C}_{18}\text{H}_{21}\text{NO}_5$ 331.368
Alkaloid from bulbs of *Crinum powellii* var. *album*, *Narcissus tazetta*, *Sprekelia formosissima* (Amaryllidaceae). Active against Rauscher leukaemia virus. Amorph. $[\alpha]_D^{24} +228$ (c, 0.18 in CHCl_3). λ_{max} 242 (ϵ 3010); 291 (ϵ 3200) (EtOH) (Berdy).
- 3-Epimer, hydrochloride:**
Needles + $\frac{1}{2}$ H_2O (H_2O). Mp 199-201°. $[\alpha]_D^{24} +82$ (c, 0.2 in H_2O).
- 3-Epimer, 8-ketone (lactone): Macronine**
[2124-70-1]
 $\text{C}_{18}\text{H}_{19}\text{NO}_5$ 329.352
Alkaloid from the leaf sheaths of *Crinum macrantherum* and the bulbs of *Crinum erubescens* (Amaryllidaceae). Prisms (Me_2CO). Mp 203-205°. $[\alpha]_D^{22} +413$ (c, 0.228 in CHCl_3). $[\alpha]_D^{22} +380$ (c, 0.225 in EtOH). λ_{max} 229 (log ϵ 4.49); 268 (log ϵ 3.74); 308 (log ϵ 3.78) (EtOH).


- 3-Epimer, 8-ketone (lactone), N-de-Me: N-Demethylmacronine**
 $\text{C}_{17}\text{H}_{17}\text{NO}_5$ 315.325
Isol. from the bulbs of *Crinum erubescens* (Amaryllidaceae). Cryst. (Me_2CO). Mp 176-177°. Not indexed by CAS to 14CI.
- 3-Epimer, 8-ketone (lactone), N-de-Me, N-ethoxycarbonyl: N-Demethyl-N-ethoxycarbonylmacronine. N-Demethylcarboethoxymacronine**
[17245-22-6]
 $\text{C}_{20}\text{H}_{21}\text{NO}_7$ 387.388
Isol. from the bulbs of *Crinum erubescens* (Amaryllidaceae). Amorph. $[\alpha]_D^{24} +313$ (c, 0.17 in CHCl_3).
- 3-Epimer, 8-ketone (lactone), O-de-Me: O-Demethylmacronine**
[233767-22-1]
 $\text{C}_{17}\text{H}_{17}\text{NO}_5$ 315.325
Alkaloid from *Galanthus gracilis*. Amorph. powder. $[\alpha]_D +227$ (c, 0.16 in MeOH). λ_{max} 228 (log ϵ 4.53); 264 (log ϵ 3.82); 309 (log ϵ 3.66) (MeOH).
- 8-Epimer, O⁸-Et, N-de-Me: 8 α -Ethoxy-N-demethylpretazettine**
[250253-93-1]
 $\text{C}_{19}\text{H}_{23}\text{NO}_5$ 345.394
Alkaloid from *Crinum bulbispermum*. Amorph. $[\alpha]_D^{28} +160.6$ (c, 0.09 in CHCl_3).
- 3,8-Diepimer, O⁸-Et: 8 α -Ethoxyprecirwelline**
[250282-46-3]
 $\text{C}_{20}\text{H}_{25}\text{NO}_5$ 359.421
Alkaloid from *Crinum bulbispermum*. Amorph. $[\alpha]_D^{28} +116.6$ (c, 0.06 in CHCl_3).
- [61045-12-3]
- Hauth, H. et al., *Helv. Chim. Acta*, 1964, **47**, 185-194 (Macronine, isol, uv, ir, pmr)
- Murphy, C.F. et al., *Tet. Lett.*, 1964, 3857-3861 (Macronine, uv, ir, pmr, synth)
- Wildman, W.C. et al., *J.A.C.S.*, 1967, **89**, 5514-5515; 1969, **91**, 150-157 (isol, struct, synth)
- Wildman, W.C. et al., *J.O.C.*, 1968, **33**, 3749-3753 (Precirwelline, 3-Epimacronine, N-Demethylmacronine, N-Demethyl-3-epimacronine, N-Demethylcarboethoxymacronine, Ungspirolidone, Ungspirolidine)
- Döpke, W. et al., *Tet. Lett.*, 1968, 1307-1310 (pmr, cd, struct)
- Isobe, K. et al., *Tet. Lett.*, 1976, 2331-2334 (Macronine, synth)
- Kadyrov, K.A. et al., *Khim. Prir. Soedin.*, 1977, **13**, 426-427; 719; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 364; 606 (Dihydroepimacronine, Ungspirolidone, Ungspirolidine)
- Martin, S.F. et al., *J.A.C.S.*, 1984, **106**, 6431-6433 (synth)
- Viladomat, F. et al., *Phytochemistry*, 1990, **29**, 1307-1310 (3-Epimacronine)
- Ishibashi, H. et al., *J.O.C.*, 1993, **58**, 2360-2368 (synth)
- Weniger, B. et al., *Planta Med.*, 1995, **61**, 77-79 (activity)
- Wagner, J. et al., *Tetrahedron*, 1996, **52**, 6591-6600 (cd)
- Linden, A. et al., *Acta Cryst. C*, 1998, **54**, 1653-1659 (3-Epimacronine, cryst struct)
- Seaforth, C.E. et al., *Fitoterapia*, 1998, **69**, 79 (cmr)
- Nishimata, T. et al., *J.O.C.*, 1998, **63**, 7586-7587 (synth)

- Jones, K. et al., *Tetrahedron*, 1998, **54**, 2275-2280 (synth)
- Elgorashi, E.E. et al., *Phytochemistry*, 1999, **52**, 533-536 (*Crinum bulbispermum* derivis)
- Unver, N. et al., *Planta Med.*, 1999, **65**, 347-350 (*O*-Demethylmacronine, *O*-Demethyl-epimacronine)
- Baldwin, S.W. et al., *Org. Lett.*, 2000, **2**, 99-102 (synth)
- Cabezas, F. et al., *Chem. Pharm. Bull.*, 2003, **51**, 315-317 (*8*-*O*-Methylpretazettine)
- Zhang, F.-M. et al., *Tetrahedron*, 2006, **62**, 9445-9455 (synth)

Preulicyclamide

P-627

[104928-30-5]




$\text{C}_{33}\text{H}_{41}\text{N}_7\text{O}_6\text{S}_2$ 695.862
Isol. from the tunicate *Lissoclinum patella*. $[\alpha]_D +5.4$ (c, 0.24 in CHCl_3).

- Sesin, D.F. et al., *Bull. Soc. Chim. Belg.*, 1986, **95**, 853 (isol, pmr, cmr)
- Sugiura, T. et al., *Tet. Lett.*, 1987, **28**, 2251 (synth)

Preulithiacyclamide

P-628

[105637-44-3]




$\text{C}_{32}\text{H}_{46}\text{N}_8\text{O}_8\text{S}_4$ 799.028
Isol. from the ascidian *Lissoclinum patella*. Amorph. powder. Mp 178-181° Mp 260-263° (synthetic). $[\alpha]_D^{22} +65$ (c, 0.5 in DMF) (synthetic). λ_{max} 211 ; 241 (no solvent reported).

- Kato, S. et al., *Tet. Lett.*, 1986, **27**, 2653-2656 (synth)
- Schmidt, U. et al., *Tet. Lett.*, 1986, **27**, 3495-3496 (synth)
- Patil, A.D. et al., *Nat. Prod. Lett.*, 1997, **9**, 181-187 (isol, ir, pmr, cmr)

Prianosin C

[116302-36-4]



C₁₈H₁₅N₃O₃S 353.401

Alkaloid from the Okinawan marine sponge *Prianos melanos*. Exhibits potent antineoplastic activity. Green solid. Fairly sol. DMF; poorly sol. MeOH, hexane. Mp 300°. [α]_D²⁵ +358 (c, 0.01 in MeOH). Unstable in soln. λ_{max} 231 (ε 12300); 263 (ε 3900); 292 (ε 2100); 370 (ε 1280) (MeOH) (Derep).

O²,N,N-Tri-Ac:

Yellow cryst. Mp 200-201° dec. [α]_D²³ +384 (c, 0.1 in CHCl₃).

Cheng, J.F. *et al.*, *J.O.C.*, 1988, **53**, 4621 (*isol, uv, ir, pmr, cmr, ms, cd*)

Kobayashi, J. *et al.*, *Tet. Lett.*, 1991, **32**, 1227 (*struct*)

Primocarcin

P-630

5-(Acetylamino)-4-oxo-5-hexenamamide, 9CI. 5-Acetamido-4-oxo-5-hexenamamide, 8CI

[3750-26-3]



C₈H₁₂N₂O₃ 184.194

Isol. from *Nocardia fukaya*. Antitumour and antimicrobial antibiotic. Needles (MeOH). Sol. H₂O, MeOH, butanol, Me₂CO, Py, EtOH; fairly sol. EtOAc; poorly sol. C₆H₆, Et₂O, hexane. Mp 130-131°. λ_{max} 215 (ε 3230); 347 (ε 304) (0.1M NaOH) (Derep). λ_{max} 254 (ε 3500) (H₂O) (Derep). λ_{max} 215 ; 347 (NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 55.5 mg/kg; LD₅₀ (mus, ivn) 300 mg/kg. MP6475000

Nagatsu, J. *et al.*, *J. Antibiot.*, Ser. A, 1962, **15**, 75; 77; 80 (*isol, struct*)


Bowman, R.E. *et al.*, *J.C.S.*, 1965, 470 (*synth, ir, uv*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PMC750

Prioline

P-631

8-Hydroxy-3-methyl-7-(1-methylethyl)-benz[cd]indol-2(1H)-one, 9CI. 8-Hydroxy-7-isopropyl-3-methylbenz[cd]indol-2(1H)-one



C₁₅H₁₅NO₂ 241.289


P-629

Alkaloid from the roots of *Salvia prionitis*. Yellow needles (EtOAc). Mp 210-212°. λ_{max} 215 (log ε 4.43); 256 (log ε 4.39); 354 (log ε 3.71) (MeOH).

Li, M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 139-141

Procerine†

[68622-81-1]



C₁₃H₂₁NO₅ 271.313


Alkaloid from *Senecio procerus* var. *procerus* (Asteraceae). Mp 238-290° dec. Tentative struct.

Jovčeva, R.J. *et al.*, *Coll. Czech. Chem. Comm.*, 1978, **43**, 2312 (*isol, ms*)

Procevine

P-633

18,28-Cyclo-16,28-secosolanid-5-en-3-ol, 9CI. 5-Cevaniden-3-ol. Pseudosolanidine [468-24-6]



Absolute Configuration

C₂₇H₄₃NO 397.643

Alkaloid from *Veratrum grandiflorum* (Liliaceae) seedlings grown under controlled conditions. Mp 235-237°. [α]_D -12.2 (c, 0.33 in CHCl₃). This alkaloid probably represents a biosynthetic intermed. between solanidanes and cevanines.

O-Ac: Mp 166-168°.

Pelletier, S.W. *et al.*, *J.A.C.S.*, 1952, **74**, 4218-4219; 1953, **75**, 4442-4446 (*synth*)


Weisenborn, F.L. *et al.*, *J.A.C.S.*, 1953, **75**, 259-262 (*synth*)

Sheehan, J.C. *et al.*, *J.A.C.S.*, 1960, **82**, 6147-6153 (*synth, struct*)

Kaneko, K. *et al.*, *Tet. Lett.*, 1978, 4801-4804 (*isol, ir, ms, pmr, struct*)

Procumbine

[109389-82-4]



C₂₀H₁₇NO₇ 383.357

Alkaloid from *Hypecoum procumbens* and *Hypecoum leptocarpum*

(Hypecoaceae). Orange-red needles. Mp 191-192°.

Táborská, E. *et al.*, *Coll. Czech. Chem. Comm.*, 1987, **52**, 508 (*isol, uv, ir*)

Táborská, E. *et al.*, *Heterocycles*, 1988, **27**, 39 (*uv, ir, pmr, cmr, ms, struct*)


Prodigosene

P-635

5-(2H-Pyrrol-2-ylidenemethyl)-2,2'-bi-1H-pyrrole, 9CI

[22187-69-5]

[22187-70-8 (hydrobromide)]



C₁₃H₁₁N₃ 209.25

Pigment isol. from *Serratia marcescens*.

Long metallic blue needles (as hydrobromide). Mp 360° (hydrobromide). λ_{max} 546 (EtOH/HCl) (Berdy).


Hearn, W.R. *et al.*, *J.O.C.*, 1970, **35**, 142

Boger, D.L. *et al.*, *J.O.C.*, 1988, **53**, 1405 (*synth, bibl*)

Prodigosin

P-636

4-Methoxy-5-[(5-methyl-4-pentyl-2H-pyrrol-2-ylidene)methyl]-2,2'-bi-1H-pyrrole, 9CI. 2-Methyl-3-pentylprodiginine [82-89-3]



C₂₀H₂₅N₃O 323.437

Pyrrole antibiotic. Pigment from *Serratia marcescens*, *Serratia polymuthica*, *Actinomadura pelletieri*, *Hahella chejuensis*, *Streptomyces variegatus*, *Nocardia* sp., *Pseudomonas magnesorubra*, *Vibrio psychroerythreus*, *Vibrio gazogenes*, *Alteromonas rubra*, *Cladophora* sp., *Beneckea gazogenes* and *Cladophora* spp. Shows antileukaemic and antifungal props. Active against plant-pathogenic fungi; insecticide enhancer. Shows antimalarial activity but too toxic for clinical use. Shows strong immunosuppressive activity. Lustrous dark red square pyramids. Sol. MeOH, acids, Et₂O; poorly sol. H₂O. Mp 151-152°. Sinters at 70-80°. λ_{max} 216 (ε 8660); 275 (sh) (ε 6440); 296 (ε 12200); 371 (ε 7750); 382 (sh) (ε 7240); 510 (sh) (ε 54400); 541 (ε 132000) (EtOH at pH 2.9) (Derep). λ_{max} 257 (ε 7950); 281 (ε 9520); 336 (ε 7860); 469 (ε 42400) (EtOH at pH 11) (Derep). λ_{max} 226 (ε 10800); 289 (ε 9820); 337 (ε 7590); 471 (ε 41600); 539 (ε 17200) (EtOH at pH 7.4) (Derep).

▶ LD₅₀ (mus, ipr) 18 mg/kg; LD₅₀ (mus, ivn) 10 mg/kg. DW2977000

Hydrochloride:

Magenta cryst. (C₆H₆/petrol). Mp 148.5-150° dec.

Perchlorate:

Deep red cryst. (EtOH). Mp 228°.

O-De-Me: De-O-methylprodigiosin. Norprodigiosin

[27005-43-2]

C₂₀H₂₅N₃O 323.437

Pigment prod. by *Hahella chejuensis* and a mutant of *Serratia marcescens*. Orange-brown cryst. (CH₂Cl₂/petrol). Sol. Me₂CO; poorly sol. H₂O. Mp 184-185°. λ_{max} 490 (EtOH) (Berdy).

Demethoxy: Demethoxyprodigiosin

[112373-41-8]

C₁₉H₂₃N₃ 293.411

From *Serratia marcescens*. Purple solid (as hydrobromide). λ_{max} 568 (EtOH/HCl) (Berdy).

Lower homologue (n = 2): 2-Methyl-3-propylprodigiosin

[955127-08-9]

C₁₈H₂₁N₃O 295.383

Prod. by the marine bacterium *Hahella chejuensis* KCTC 2396.

Lower homologue (n = 3): 3-Butyl-2-methylprodiginine

[955127-09-0]

C₁₉H₂₃N₃O 309.41

Prod. by *Hahella chejuensis* KCTC 2396.

Homologue (n = 5): 3-Hexyl-2-methylprodiginine

[955127-10-3]

C₂₁H₂₇N₃O 337.464

Prod. by *Hahella chejuensis* KCTC 2396.

Homologue (n = 6): 3-Heptyl-2-methylprodiginine. Methylosin B

[89014-12-0]

C₂₂H₂₉N₃O 351.491

Prod. by *Hahella chejuensis* KCTC 2396 and earlier isol. from *Methylosinus trichosporium*. The struct. of the accompanying Methylosin A has not been established.

[89072-96-8 (Methylosin A)]

Wrede, F. *et al.*, *Ber.*, 1929, **62**, 2678

Rapoport, H. *et al.*, *J.A.C.S.*, 1962, **84**, 635 (synth)

Hearn, W.R. *et al.*, *Biochem. Biophys. Res. Commun.*, 1964, **17**, 517 (Norprodigiosin)

Williams, R.P. *et al.*, *Antibiotics (N.Y.)*, 1967, **2**, 410; 449 (rev)

Jackson, A.H. *et al.*, *Tetrahedron*, 1967, **23**, 603 (ms)

Hearn, W.R. *et al.*, *J.O.C.*, 1970, **35**, 142 (synth)

Deol, B.S. *et al.*, *Aust. J. Chem.*, 1974, **27**, 2657 (Norprodigiosin)

Lim, D.V. *et al.*, *J. Bacteriol.*, 1977, **129**, 124 (biosynth)

Sveshnikova, M.A. *et al.*, *Antibiotiki (Moscow)*, 1983, **28**, 723 (isol)

Strauss, D.G. *et al.*, *Z. Allg. Mikrobiol.*, 1983, **23**, 661-668 (Methylosins A-B)

Boger, D.L. *et al.*, *J.O.C.*, 1988, **53**, 1405-1415 (synth, biosynth)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**, 1725; 1999, **40**, 7587-7589 (synth)

Okamoto, H. *et al.*, *CA*, 1999, **130**, 207057x; 2000, **132**, 47317 (activity)


Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (rev)

Kim, D. *et al.*, *J. Appl. Microbiol.*, 2007, **102**, 937-944 (*Hahella* homologues)

Proferrosamine B P-637

2-Carboxy-3,4-dihydro-α-hydroxy-5-(2-pyridinyl)-2H-pyrrole-4-acetic acid, 9CI

5-Carboxy-2-(2-pyridyl)-1-pyrroline-3-glycolic acid, 8CI
[21045-59-0]



C₁₂H₁₂N₂O₅ 264.237

Prod. by *Bacillus roseus fluorescens*. Siderophore.

Fe complex (2:1): [21711-47-7]


C₂₄H₂₂FeN₄O₁₀ 582.306

Isol. from *Bacillus roseus fluorescens*. Red pigment.

Pouteau-Thouvenot, M. *et al.*, *Bull. Soc. Chim. Biol.*, 1968, **50**, 222 (isol)

Prolifine P-638

8,9-Dihydroxy-4-methyl-10H-[1]benzopyrano[3,2-b]pyridin-10-one, 9CI
[263871-95-0]




C₁₃H₉NO₄ 243.218

Alkaloid from the roots of *Saccopetalum proliferum* (preferred genus name *Milium*). Yellow needles. Mp 243-245°. λ_{max} 205 (log ε 4.43); 222 (log ε 4.27); 253 (log ε 4.02); 269 (log ε 4.06); 286 (log ε 4.04); 352 (log ε 3.97) (MeOH).

Wang, M.L. *et al.*, *Chin. Chem. Lett.*, 2000, **11**, 129-130

Prometaphanine P-639

6,7-Didehydro-8,10-epoxy-3,4,7-trimethoxy-17-methylhasubanan-8-ol, 9CI
[6858-85-1]



C₂₀H₂₅NO₅ 359.421

Alkaloid from *Stephania japonica* (Menispermaceae). Amorph. Mp 207° (as methiodide). [α]_D¹⁰ -32 (MeOH) ((methiodide)).

16-Oxo: 16-Oxoprometaphanine
[58738-31-1]

C₂₀H₂₃NO₆ 373.405

Alkaloid from the leaves, stems and rhizomes of *Stephania japonica* (Menispermaceae). Prisms (MeOH). Mp

195°. [α]_D²⁰ -52.3 (c, 0.5 in CHCl₃).


Tomita, M. *et al.*, *Tet. Lett.*, 1964, 3617 (ir, pmr, struct)

Matsui, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 1323; 1982, **45**, 497 (Prometaphanine, 16-Oxoprometaphanine, isol, uv, ir, pmr, ms, struct)

Watanabe, Y. *et al.*, *Phytochemistry*, 1975, **14**, 2695 (uv, ir, pmr, ms, struct, deriv)

Promoinducin P-640

58-(1-Carboxyethenyl)sulfomycin I
[164802-54-4]




C₅₇H₅₄N₁₆O₁₈S₂ 1315.282

Prod. by *Streptomyces* sp. SF2741. tipA promotor. Powder. Mp 290-300° (dec.). [α]_D²⁰ -15.6 (c, 0.1 in CHCl₃/MeOH). λ_{max} 204 (ε 87600); 251 (ε 86800); 315 (sh) (ε 15500) (EtOH).

Yun, B.-S. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 876-880 (isol, struct)

Promothiocin A P-641

SF 2741A. Antibiotic SF 2741A. 10381W. Antibiotic 10381W. pre-W
[156737-05-2]
[164455-23-6]



R = NH₂

C₃₆H₃₇N₁₁O₈S₂ 815.889

Thiopeptide antibiotic. Prod. by *Streptomyces* sp. SF2741. Powder. Sol. Py, DMF, DMSO; poorly sol. H₂O, MeOH. Mp 268-272° dec. [α]_D²¹ +79.2 (c, 0.69 in CHCl₃/MeOH). λ_{max} 224 (ε 34000); 313 (sh) (ε 9000) (MeOH) (Derep).

Yun, B.-S. *et al.*, *J. Antibiot.*, 1994, **47**, 510-514 (isol, uv, ir, pmr, cmr)

- Pat. Coop. Treaty (WIPO)*, 1995, 95 07 292; CA, **123**, 54272s (10381W)
 Bagley, M.C. et al., *J.A.C.S.*, 2000, **122**, 3301-3313 (synth)
 Yun, B.-S. et al., *Tetrahedron*, 2001, **57**, 9683-9687 (abs config, conformn)

Promothiocin B P-642


SF 2741B. Antibiotic SF 2741B. 10381X. Antibiotic 10381X. pre-X
 [156737-06-3]
 [164455-24-7]
 As Promothiocin A, P-641 with
 R = -NHC(=CH₂)CONHC(=CH₂)CONH₂

C₄₂H₄₃N₁₃O₁₀S₂ 954.015

Thiopeptide antibiotic. Prod. by *Streptomyces* sp. SF2741. Powder. Sol. Py, DMF, DMSO; poorly sol. H₂O, MeOH. Mp 250-255° dec. [α]_D²¹ -62.5 (c, 0.3 in CHCl₃/MeOH). λ_{max} 225 (ε 42000); 313 (sh) (ε 10000) (MeOH) (Derep). λ_{max} 225 (ε 41000); 313 (ε 10000) (MeOH) (Berdy). Yun, B.-S. et al., *J. Antibiot.*, 1994, **47**, 510 (isol, uv, ir, pmr, cmr)
Pat. Coop. Treaty (WIPO), 1995, 95 07 292; CA, **123**, 54272s (10381X)
 Yun, B.-S. et al., *Tetrahedron*, 2001, **57**, 9683-9687 (abs config, conformn)

Pronuciferine P-643

Miltanthin. N,O-Dimethylcrotsparine



(R)-form

C₁₉H₂₁NO₃ 311.38

(R)-form [2128-60-1]

Alkaloid from *Nelumbo nucifera* (East India lotus), *Stephania glabra*, *Papaver caucasicum*, *Meconopsis cambrica* and *Isolona pilosa* (Nelumbonaceae, Menispermaceae, Papaveraceae, Annonaceae). Shows strong local anaesthetic activity. Mp 127-129°. [α]_D²⁰ +99 (c, 0.2 in CHCl₃). [α]_D²⁰ +106 (c, 0.41 in EtOH).

▶ WH0167500

Picrate: Mp 119-123°.

N-De-Me: **Stepharine**. *Stephaglabrine*

[2810-21-1]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Stephania glabra* (Menispermaceae). Mp 179-181°. [α]_D²⁰ +143 (c, 1.88 in CHCl₃).

N-De-Me, N-formyl: **N-Formylstepharine**

[104012-91-1]

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Caryomene olivascens*. Negative opt. rotn.

N-De-Me, N-Ac: **N-Acetylstepharine**

[4880-87-9]

[75023-53-9]

C₂₀H₂₁NO₄ 339.39

Alkaloid from *Stephania sasakii*. Mp 234-235°. [α]_D²⁴ -80 (c, 1.6 in CHCl₃).

N-De-Me, N-carbamoyl: **N-Carboxami-**

dostepharine. N-Carbamoylstepharine

[113145-62-3]

C₁₉H₂₀N₂O₄ 340.378

Alkaloid from the rhizomes of *Stephania venosa* (Menispermaceae). Amorph. [α]_D²⁵ +22 (c, 0.2 in CHCl₃).

N-De-Me, N-methoxycarbonyl: **Promucosine**

[275355-87-8]

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Annona purpurea* (socoaya). Brown powder. Mp 125-127°. [α]_D²⁵ +147 (c, 0.07 in CHCl₃). λ_{max} 217 (log ε 3.78); 267 (log ε 4.45); 303 (log ε 3.87) (MeOH).

O¹-De-Me: **6-Methylcrotsparine**. *Suavedol*. *Glazen*. *PM 297*

[6808-72-6]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Ocotea glaziovii*, *Annona purpurea*, *Meconopsis cambrica* and *Litsea laurifolia* (Lauraceae, Papaveraceae). Claimed antidepressant activity. Weak tumour inhibitor. Active against gram-positive bacteria. Mp 235-237° dec. [α]_D²⁰ +7 (c, 1 in CHCl₃). The natural alkaloid is largely racemic. In some plants there is an excess of the (-)-form.

O²-De-Me: Mp 216-218°. [α]_D²⁰ +122 (c, 0.45 in MeOH).

O²-De-Me, N-de-Me: **Crotonosine**

[2241-43-2]

Alkaloid from *Croton linearis* (Euphorbiaceae). Shows strong local anaesthetic activity. Depolarising neuromuscular blocker. Mp 300° (begins to dec. at 165°, softens at 197°). [α]_D²⁸ +180 (c, 0.83 in MeOH).

O²-De-Me, N-de-Me, N,O-di-Ac: Mp 203-205°.

(S)-form [16654-37-8]

Alkaloid from *Ocotea glaziovii* (Lauraceae) and *Croton sparsiflorus* (Euphorbiaceae). Needles (EtOAc). Mp 125-127°.

O¹-De-Me: **N-Methylcrotsparine**

[18058-57-6]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Croton sparsiflorus* (Euphorbiaceae). Cryst. (EtOAc). Mp 223-225°. [α]_D²⁰ -113 (c, 1.52 in CHCl₃).

O¹-De-Me, N-de-Me: **Crotsparine**. *Nor-glaziovine*. *Crotoflorine*

[18373-79-0]

C₁₇H₁₇NO₃ 283.326

Alkaloid from *Croton sparsiflorus* and *Uvaria klaineana*. Cryst. (CHCl₃/EtOAc). Mp 193-195°. [α]_D²⁰ -64 (c, 0.73 in MeOH). λ_{max} 235 (log ε 3.37) (no solvent reported).

O¹-De-Me, N-de-Me, hydrochloride: Mp 278° dec.

O¹-De-Me, N-de-Me, N,O-di-Ac: Mp 185-186°.

O²-De-Me: **Homolinearisine**

[10214-68-3]

C₁₈H₁₉NO₃ 297.353

Alkaloid from *Caryomene linearis* (Euphorbiaceae). Mp 218-220° dec.

O²-De-Me, perchlorate: Mp 184-186°.

2',3',5',6'-Tetrahydro: **Tetrahydropronuciferine**

[16562-08-6]

C₁₉H₂₅NO₃ 315.411

Alkaloid from the leaves of *Phoebe scortechinii*. Amorph. gum. [α]_D²⁷ -50 (c, 1 in CHCl₃). λ_{max} 249 (log ε 2.3); 285 (log ε 1.93); 306 (log ε 3.51) (MeOH).

(±)-form [17236-29-2]

Synthetic. Mp 148-151°.

O¹-De-Me: **Glaziovine**, *INN*

[17127-48-9]

From *Ocotea glaziovii*. Neuropharmacological agent. Antitumorogenic agent. Mp 227-228° dec.

O²-De-Me:

Synthetic. Noncryst.

Tetrahydro, O¹-de-Me: **Tetrahydroglaziovine**

[50300-14-6]

C₁₈H₂₃NO₃ 301.385

Alkaloid from *Croton sparsiflorus* (Euphorbiaceae). Mp 163-164°.

[32410-24-5]

Bernauer, K. et al., *Helv. Chim. Acta*, 1963, **46**, 1783-1785 (*R*-form, isol, uv, ir, pmr, struct)

Gilbert, B. et al., *J.A.C.S.*, 1964, **86**, 694-696 (*Glaziovine*, isol, uv, ir, pmr, struct)

Haynes, L.J. et al., *J.C.S.(C)*, 1966, 1676-1679 (*Crotonosine*, ir, pmr, struct)

Baldwin, M. et al., *J.C.S.(C)*, 1967, 154-161 (*ms*)

Barton, D.H.R. et al., *J.C.S.(C)*, 1967, 1295-1298 (*Crotonosine*, biosynth)

Kametani, T. et al., *J.C.S.(C)*, 1967, 2182-2184 (*Glaziovine*, synth, pmr)

Bhakuni, D.S. et al., *Experientia*, 1968, **24**, 10 (*Crotsparine*, uv, ir, pmr, ms, struct)

Ishiwata, S. et al., *Chem. Pharm. Bull.*, 1970, **18**, 1841-1845 (*Homolinearisine*, synth)

Ferrari, G. et al., *Farmaco, Ed. Sci.*, 1970, **25**, 449 (config)

Bhakuni, D.S. et al., *Phytochemistry*, 1970, **9**, 2573-2580; 1974, **13**, 2767-2769 (*Crotsparine*, isol, pmr, struct, derivs, biosynth)

Kametani, T. et al., *Tet. Lett.*, 1973, 4219-4220 (*Glaziovine*, synth)

Casagrande, C. et al., *Farmaco, Ed. Sci.*, 1975, **30**, 479-490 (*Glaziovine*, *Crotsparine*, isol)

Casagrande, C. et al., *J.C.S. Perkin 1*, 1975, 1659-1663 (*Tetrahydroglaziovine*)

Marzo, A. et al., *Arch. Int. Pharmacodyn. Ther.*, 1977, **227**, 254-262 (*Glaziovine*, metab)

De Angelis, L. et al., *Drugs of Today (Barcelona)*, 1977, **13**, 22 (rev, pharmacol)

Bindra, J.S. et al., *J.O.C.*, 1977, **42**, 910-911 (*Glaziovine*, synth)

Ricca, G.S. et al., *Org. Magn. Reson.*, 1977, **9**, 8 (cmr)

Chaumontet, M. et al., *Arzneim.-Forsch.*, 1978, **28**, 2119-2121 (*Glaziovine*, pharmacol)

Horii, Z. et al., *Chem. Pharm. Bull.*, 1978, **28**, 481-483 (synth, uv, ir, pmr)

Marzo, A. et al., *Eur. J. Clin. Pharmacol.*, 1978, **13**, 219-221 (*Glaziovine*, pharmacokinetic)

Kunitomo, J. et al., *Yakugaku Zasshi*, 1981, **101**, 431-436 (*N-Acetylstepharine*)

Fajardo, V. et al., *Chem. Comm.*, 1982, **23**, 1350-1352 (pmr)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 12785

Madhusudanan, K.P. et al., *Indian J. Chem., Sect. B*, 1985, **24**, 523-526 (ms)


Lavault, M. et al., *Chem. Pharm. Bull.*, 1986, **34**, 1148-1152 (*N-Formylstepharine*)

Charles, B. et al., *J. Nat. Prod.*, 1987, **50**, 1113-1117 (*N-Carboxamidostepharine*)

- Simeon, S. *et al.*, *Pharmazie*, 1989, **44**, 593-597 (Glazioline, activity)
 Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 746-748 (*Promucosine*)
 Akendengue, B. *et al.*, *Planta Med.*, 2002, **68**, 167-169 (*Crosparine*)
 Honda, T. *et al.*, *Org. Lett.*, 2006, **8**, 657-659 (synth)
 Awang, K. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 704-709 (*Tetrahydropromuciferine*)

Prooxocryptochine

P-644

C₁₇H₁₅NO₄ 297.31


Alkaloid from the wood of *Cryptocarya chinensis*. Yellow syrup. $[\alpha]_D^{25}$ -17.2 (c, 0.02 in MeOH). λ_{max} 229 ; 254 ; 331 ; 379 (MeOH).

Wu, T.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1404-1407

2-Propanoylpyridine

P-645

1-(2-Pyridinyl)-1-propanone, 9CI. Ethyl 2-pyridyl ketone [3238-55-9]

C₈H₉NO 135.165

Alkaloid from the leaves *Semnostachya menglaensis* (preferred genus name *Strobilanthes*). Oil. Bp 206-207° Bp₉ 81-85°.

Phenylhydrazone: Mp 139-140° (128-129°).

Wibaut, J.P. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1951, **70**, 1054-1066 (synth)

Teague, P.C. *et al.*, *J.A.C.S.*, 1953, **75**, 3429-3430 (synth)

Prasad, K.B. *et al.*, *Chem. Ber.*, 1965, **98**, 2822-2824 (synth)

Berg-Nielsen, K. *et al.*, *Acta Chem. Scand., Ser. B*, 1975, **29**, 985-989 (synth)

Knebel, N.G. *et al.*, *J. Med. Chem.*, 1991, **34**, 2145-2152 (synth, pmr)

Sato, N. *et al.*, *Synthesis*, 2001, 1551-1555 (synth, ir, pmr, cmr)

Naef, R. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 9161-9164 (isol)

2-Propenoic acid, 9CI

P-646

Acrylic acid

[79-10-7]

H₂C=CHCOOHC₃H₄O₂ 72.063

Manuf. by oxidn. of 1-Propene or by reaction of Ni(CO)₄ with Acetylene and H₂O. Found in green algae. Prod. by *Phaeocystis* sp., *Phaeocystis pouchetii*,

Enteromorpha sp., *Ulva* sp., *Codium* sp., *Patinopecten yessoensis*, *Protogonyaulax* sp. Undergoes free-radical polym. in aq. soln. to give polyacrylates, acrylic resins etc. USA production in 1999 1.39 million tons. Corrosive liq. with acrid odour and fumes. Misc. H₂O, EtOH, Et₂O; sol. Me₂CO. d₄¹⁶ 1.06. Mp 13°. Bp 141° (polymerises). n_D²⁰ 1.4424. pK_a 4.25 (25°). Vp 4 mmHg (25°). Polym. readily in the presence of O₂. Acrylic acid and its esters usually contain hydroquinone monomethyl ether as polym. inhibitor. Q/e values for copolym., Q 0.83, e +0.88. Copolym. with vinyl chloride: r_{AA} 6.8; r_{VC} 0.107 (typical values, 60°); Q 1.15; e 0.77.

► Flammable, fl. p. 52°, autoignition temp. 438°. May polym. exothermically (possibly violently). Irritant to all tissues. LD₅₀ (rat, orl) 33.5 mg/kg. LD₅₀ (rbt, skn) 280 mg/kg. Exp. teratogen (ipr route). OES: long-term 10 ppm; short-term 20 ppm (Sk). AS4375000

Amide: Acrylamide. 2-Propenamide, 9CI

[79-06-1]

C₃H₅NO 71.079

Constit. of *Pongamia glabra* leaf galls. Manuf. by hydration of the nitrile. Undergoes free radical polym. in aq. soln. to give polyacrylamide. Copolym. easily with many vinyl monomers. Leaflets (C₆H₆). Sol. H₂O, EtOH, MeOH, Et₂O, Me₂CO, DMF, DMSO. Mp 85°. Bp 192.6°. Polym. on heating or under uv light. Aq. solns. are stabilised by O₂ and Cu²⁺ ions. Q/e values for copolym., Q 0.23, e +0.54.

► Polym. can be violent at Mp. Symptoms of exposure (dermal, inhalation or ingestion) include cold sweating hands and peripheral neuropathies with numbness, paraesthesias and weakness. Dermal absorption can also cause erythema and peeling of the palms. Probable human carcinogen (IARC 2A). LD₅₀ (rat, orl) 124 mg/kg. LD₅₀ (rat, skn) 400 mg/kg. Exp. carcinogen and reprod. effects. Adverse exp. CNS effects. MEL: long-term 0.3 mg m⁻³ (Sk). AS3325000

[6292-01-9 , 10344-93-1 , 9003-05-8 , 7446-81-3 , 10192-85-5 , 9003-06-9 , 14643-87-9]

Tareke, E. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 4998-5006 (amide, occur; bibl)

Mottram, D.S. *et al.*, *Nature (London)*, 2002, **419**, 448 (amide, occur)


Friedman, M. *et al.*, *J. Agric. Food Chem.*, 2003, **51**, 4504-4526 (acrylamide, rev, chem, biochem, safety)

2-(1-Propenyl)quinoline

P-647

Chimanine B

[57078-89-4]

C₁₂H₁₁N 169.226

Alkaloid from leaves of *Galipea longiflora*

(Rutaceae). Oil. Bp 249-253° Bp_{0.01} 80°. n_D²⁵ 1.6359.

1',2'-Epoxide: 2-(3-Methyloxiranyl)quinoline, 9CI. 1-Methyl-2-(2-quinolinyl)oxirane. 2-(1,2-Epoxypropyl)quinoline. Chimanine D C₁₂H₁₁NO 185.225

Alkaloid from leaves of *Galipea longiflora* (Rutaceae). Oil.

4-Methoxy: 4-Methoxy-2-(1-propenyl)quinoline. Chimanine C

[150044-85-2]

C₁₃H₁₃NO 199.252

Alkaloid from leaves of *Galipea longiflora* (Rutaceae). Oil.

Borsche, W. *et al.*, *Chem. Ber.*, 1950, **83**, 78 (synth)

Gilman, H. *et al.*, *J.A.C.S.*, 1959, **81**, 4000 (synth)

Ishikura, M. *et al.*, *Heterocycles*, 1985, **23**, 2375 (synth, pmr)

Fourmet, A. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1547 (isol, uv, pmr, cmr, ms, struct)

Munos, M.-H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1914 (synth, Chimanine D)

1-Propylamine, 8CI

P-648

Propanamine, 9CI. 1-Aminopropane

[107-10-8]

H₃CCH₂CH₂NH₂C₃H₉N 59.111

Isol. from ergot, other fungi and other vegetable sources. Liq. d₄²⁰ 0.72. Fp -83. Bp 49°. n_D²⁰ 1.3910. pK_a 10.4 (30°).

► Highly flammable, fl. p. -37°, autoignition temp. 318°. Respiratory tract irritant. Corrosive and irritating to skin and eyes. LD₅₀ (rbt, skn) 560 mg/kg. UH9100000 Hydrochloride: [556-53-6] Deliquescent cryst. Mp 157-158°.

► UI3057000

N-Ac: N-Propylacetamide, 9CI

[5331-48-6]

C₅H₁₁NO 101.148Oil. Bp 222-225° Bp₂ 60-65°.

N-Ac; hydrochloride:

Needles. Mp 47°.

N-Benzoyl: N-Propylbenzamide

[10546-70-0]

C₁₀H₁₃NO 163.219

Cryst. (EtOH or C₆H₆). Mp 84.5°. Bp 294-295° sl. dec.

► CV5800000

N-Me: [627-35-0]

C₄H₁₁N 73.138

Liq. with fishy odour. Bp 62-64°.

N,N-Di-Me: [926-63-6]

C₅H₁₃N 87.164

Liq. Bp 65-67°.

N-Hydroxy: N-Hydroxypropylamine. N-Propylhydroxylamine, 8CI. 1-Hydroxylaminopropane

[627-38-3]

C₃H₉NO 75.11

Needles (Et₂O). Mp 45.5-46°.

N-Hydroxy, oxalate: Mp 154-155°.

[17033-39-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 280C; 280D (ir)


Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 450A; 450B; 473C (nmr)

- Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 364C; 379C (*ir*)
- Aldrich Library of NMR Spectra*, **2**, A Hofmann, A.W. *et al.*, *Ber.*, 1882, **15**, 762 (*synth*)
- Graymore, J. *et al.*, *J.C.S.*, 1931, 1490 (*N-Me*)
- Beregi, L. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1947, **224**, 1508
- Ryer, A.I. *et al.*, *J.A.C.S.*, 1951, **73**, 5675 (*N-hydroxy*)
- Iffland, D.C. *et al.*, *J.A.C.S.*, 1952, **74**, 6284 (*synth*)
- Sasa, T. *et al.*, *CA*, 1957, **51**, 2780 (*synth*)
- Feuer, H. *et al.*, *J.O.C.*, 1965, **30**, 2877; 2880 (*N-hydroxy*)
- Iverson, P.E. *et al.*, *Acta Chem. Scand.*, 1966, **19**, 2303 (*N-hydroxy*)
- Nagasawa, A. *et al.*, *CA*, 1966, **65**, 15213 (*synth*)
- Faure, R. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 20 (*cmr*)
- Colosimo, M. *et al.*, *Org. Mass Spectrom.*, 1982, **17**, 286 (*ms*)
- Watanabe, Y. *et al.*, *J.O.C.*, 1984, **49**, 4451 (*N-Ac, synth, pmr, cmr, ir*)
- Durig, J.R. *et al.*, *J. Raman Spectrosc.*, 1989, **20**, 311 (*ir, Raman*)
- Kirk-Othmer Encycl. Chem. Technol.*, 4th edn., Wiley, 1991, **2**, 369 (*rev*)
- Ethel Browning's Toxicity and Metabolism of Industrial Solvents*, 2nd edn., (ed. Snyder, R.), Elsevier, Volume 2, 1990, 120
- Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 1072
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PND250

Propyleine

P-649

1,2,4,5,6,6a,7,8,9,9a-Decahydro-2-methylpyrido[2,1,6-de]quinolizine, 9CI [38211-32-4]



Relative configuration

$C_{13}H_{21}N$ 191.316


Alkaloid from the defence secretion of the beetle *Propylaea quatuordecimpunctata*, the Australian soldier-beetle *Chauliognathus pulchellus* and the seven-spot ladybug *Coccinella septempunctata*. Amorph. Undergoes rapid degradn. when kept as the free base.

- Tursch, B. *et al.*, *Chimia*, 1972, **26**, 74 (*uv, ir, pmr, struct*)
- Mueller, R.M. *et al.*, *J.O.C.*, 1984, **49**, 2217 (*synth, pmr, cmr*)

N,N'-Propylenebis(cantharimide)

P-650

[936025-61-5]



$C_{23}H_{30}N_2O_6$ 430.5

Alkaloid from the Chinese blister beetle, *Mylabris phalerata*. Powder. Mp 156-157°.

Homologue (n = 4): N,N'-Butylenebis(cantharimide)

[89068-48-4]

$C_{24}H_{32}N_2O_6$ 444.527

Alkaloid from *Mylabris phalerata*. Powder. Mp 246-247°.

Homologue (n = 5): N,N'-Pentylenebis(cantharimide)

[936025-62-6]

$C_{25}H_{34}N_2O_6$ 458.553


Alkaloid from *Mylabris phalerata*. Powder. Mp 148-149°.

- Nakatani, T. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 92-94 (*isol, pmr, cmr*)

5-Propyl-1-oxa-4-azacyclo-tridecan-13-one, 9CI

P-651

9-Propyl-10-azacyclododecan-12-olide [147363-85-7]



$C_{14}H_{27}NO_2$ 241.373


Related to Epilachnene, E-93. Trace alkaloid from the pupal defensive secretion of the Mexican bean beetle *Epilachna varivestis*.

- Attygalle, A.B. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 5204 (*isol, ms, struct*)
- King, A.G. *et al.*, *Tet. Lett.*, 1996, **37**, 2141 (*synth*)
- Gribble, G.W. *et al.*, *Tet. Lett.*, 1996, **37**, 2145 (*synth*)
- Rao, B.V. *et al.*, *Tet. Lett.*, 1996, **37**, 8613 (*synth*)

2-Propylpiperidine

P-652

Coniine. Cicutine. Conicine



(*R*)-form

$C_8H_{17}N$ 127.229

The first alkaloid to be synthesised. The alkaloid (and its *N-Me* deriv.) from *Conium maculatum* are largely racemic but opt. active forms can be obt. from residues.

(*R*)-form [5985-99-9]

Bp 166.5°. $[\alpha]_D$ -15.6.

Hydrochloride: [88057-03-8]

Mp 220-221° (214-215°). $[\alpha]_D^{20}$ -6.8 (c, 3.5 in EtOH).

Hydrobromide: Mp 205°.

N-Me: N-Methyl(-)-coniine

$C_9H_{19}N$ 141.256

Alkaloid from *Conium maculatum* (hemlock) (Apiaceae). Oil. Mp 175-176°. $[\alpha]_D^{20}$ -81.9.

N-Me; hydrochloride: Mp 189-190°.

(*S*)-form [458-88-8]

Alkaloid of *Conium maculatum*, also prob. in *Arum maculatum*, *Aethusa cynapium*, *Arisarum vulgare*, *Amorphophallus rivieri* (devil's tongue) and *Caladium bulbosum* (stereochem. not detd.) (Apiaceae, Araceae). d_4^{23} 0.84. Mp -2°. Bp 166-167°. $[\alpha]_D$ +15.7 (neat). $[\alpha]_D$ +8 (c, 4 in $CHCl_3$). n_D^{23} 1.4537.

► Exp. teratogen. LD₅₀ (mus, orl) 100 mg/kg. TN3730000

Hydrochloride:

Rhomboids. Mp 220°.

Hydrobromide: Mp 207°.

Picrate: Mp 75°.

N-Me: N-Methyl(+)-coniine

[35305-13-6]

$C_9H_{19}N$ 141.256

Alkaloid from *Caladium maculatum* and *Pimpinella acuminata* (Araceae, Apiaceae). Bp₇₅₇ 173-174°. $[\alpha]_D^{23}$ +82.4.

N-Me; hydrochloride: Mp 192-193° (188-189°). $[\alpha]_D$ +27.8.

N-Di-Me: N,N-Dimethyl(+)-coniine

$C_{10}H_{22}N^{\oplus}$ 156.291

Quaternary alkaloid from *Aloe sabaea*. $[\alpha]_D^{24}$ +7.5 (c, 0.1 in MeOH).

(±)-form [3238-60-6]

Alkaloid from *Caladium maculatum* (Araceae). Bp 166° Bp₇₀ 93°. pK_a 16 (25°NH, aq. KOH).

Hydrochloride: [15991-59-0]
Mp 216-217° (211°).

N-Me: Bp 175.5° Bp₁₃ 61-63°.

N-Me; hydrochloride:

Needles (Me₂CO). Mp 168-168.5°.

N-Me; picrate: Mp 112-114°.

[553-75-3]

Giesecke, *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1827, **20**, 97 (*isol*)

Hofmann, A.W. *et al.*, *Ber.*, 1884, **17**, 825 (*struct*)

Ladenburg, A. *et al.*, *Ber.*, 1886, **19**, 2578; 1894, **27**, 3062 (*synth, resoln*)

Wolfenstein, R. *et al.*, *Ber.*, 1894, **27**, 2611 (*isol, deriv*)

Ahrens, F.B. *et al.*, *Ber.*, 1902, **35**, 1330 (*isol, deriv*)

Chemnitius, F. *et al.*, *J. Prakt. Chem.*, 1928, **118**, 25 (*isol*)

Fairbairn, J.W. *et al.*, *Phytochemistry*, 1961, **1**, 38

Weitkamp, H. *et al.*, *Chem. Ber.*, 1962, **95**, 2896 (*pmr*)

Leete, E. *et al.*, *J.A.C.S.*, 1970, **92**, 3835 (*biosynth*)

Craig, J.C. *et al.*, *J.O.C.*, 1971, **36**, 3648 (*resoln*)

Roberts, M.F. *et al.*, *Phytochemistry*, 1974, **13**, 1841 (*biosynth*)

Aketa, K. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 621 (*synth*)

Kondrat, R.W. *et al.*, *Science (Washington, D.C.)*, 1978, **199**, 978 (*ms*)

Astier, A. *et al.*, *Tet. Lett.*, 1978, 2051 (*synth*)

Craig, J.C. *et al.*, *Tetrahedron*, 1978, **34**, 501 (*abs config*)

Guerrier, L. *et al.*, *J.A.C.S.*, 1983, **105**, 7754 (*synth*)


Husson, H.P. *et al.*, *Youji Huaxue*, 1987, 314; *CA*, **109**, 73693g (*synth*)

Nagasaka, T. *et al.*, *Heterocycles*, 1988, **27**, 1685 (*synth, ir, pmr*)

- Teng, T.-F. *et al.*, *Heterocycles*, 1990, **31**, 1201 (synth)
- Kiguchi, T. *et al.*, *Heterocycles*, 1990, **31**, 1525 (synth)
- Higashiyama, K. *et al.*, *Heterocycles*, 1992, **33**, 17 (synth)
- Ito, M. *et al.*, *Tet. Lett.*, 1992, **33**, 3765 (synth)
- Enders, D. *et al.*, *Annalen*, 1993, 173 (synth)
- Al-awar, R.S. *et al.*, *J.O.C.*, 1993, **58**, 7732 (synth)
- Hattori, K. *et al.*, *Tetrahedron*, 1993, **49**, 1749 (synth)
- Hirai, Y. *et al.*, *Chem. Lett.*, 1994, 21 (synth)
- Amat, M. *et al.*, *Tet. Lett.*, 1994, **35**, 2223 (synth)
- Kim, Y.H. *et al.*, *Tet. Lett.*, 1996, **37**, 5543 (synth)
- Weymann, M. *et al.*, *Synthesis*, 1997, 1151-1160 (synth)
- Reding, M.T. *et al.*, *J.O.C.*, 1998, **63**, 6344-6347 (synth)
- Jo, E. *et al.*, *Tet. Lett.*, 1999, **40**, 5581-5582 (synth, *N-Boc-coniine*)
- Wilkinson, T.J. *et al.*, *Org. Lett.*, 2000, **2**, 155-158 (synth)
- Blitzke, T. *et al.*, *Phytochemistry*, 2000, **55**, 979-982 (*N-Dimethylconiine*)
- Bois, F. *et al.*, *Tet. Lett.*, 2000, **41**, 8769-8772 (synth)
- Pachamuthu, K. *et al.*, *J. Organomet. Chem.*, 2001, **624**, 359-363 (synth)
- Burke, A.J. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 1387-1394 (synth)
- Nagata, K. *et al.*, *Heterocycles*, 2006, **70**, 335-344 (synth)
- Castro, A. *et al.*, *Heterocycles*, 2007, **71**, 2699-2708 (synth)
- Lebrun, S. *et al.*, *Org. Lett.*, 2007, **9**, 2473-2476 (synth)
- Etzebarria, J. *et al.*, *Tetrahedron*, 2007, **63**, 11421-11428 (synth)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PNT000

6-Propyl-3-piperidinol, 9CI P-653

5-Hydroxy-2-propylpiperidine



C₈H₁₇NO 143.228

(3*S*,6*S*)-form

(+)-trans-form. *Pseudoconiine*. *ψ*-*Conhydrine*
[140-55-6]
Alkaloid from hemlock (*Conium maculatum*) (Apiaceae). Mp 105-106° (93-94°). [α]_D¹⁵ +11 (EtOH).

► UL5300000

Hydrochloride: [99545-16-1]
Cryst. (MeOH/Et₂O). Mp 220° (212-213°). [α]_D²⁰ +7.6 (c, 1 in MeOH).
N-Benzoyl: Mp 132-133°. [α]_D¹⁷ +23.4.

N-Me: *N-Methylpseudoconiine*

[78962-69-3]
C₉H₁₉NO 157.255
Alkaloid from *Conium maculatum* (Apiaceae). Hexagonal rods (as hydrochloride). Mp 157° (hydrochloride). [α]_D²⁵ +25 (MeOH).

(3*S*,6*R*)-form

Epipseudoconiine

Hydrochloride: [165883-07-8]
Yellow solid (CH₂Cl₂/Et₂O). Mp 148°. [α]_D²⁰ -9.1 (c, 1 in EtOH).

(3*R*,6*R*)-form [5457-27-2]

Mp 90-92°.

Hydrochloride: [41221-92-5]

Mp 135-140°.

N-Me; *hydrochloride*: Mp 171-173°.

[40336-06-9]

Ladenburg, A. *et al.*, *Ber.*, 1891, **24**, 1671 (*isol*)
Späth, E. *et al.*, *Ber.*, 1933, **66**, 591 (*struct*)
Sorm, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1949, **14**, 331 (*synth*)

Marion, L. *et al.*, *J.A.C.S.*, 1949, **71**, 3402

(*synth*)

Hill, R.K. *et al.*, *J.A.C.S.*, 1958, **80**, 1611 (*abs config*)

Yanai, H.S. *et al.*, *Tetrahedron*, 1959, **6**, 103

(*cryst struct*)

Brown, E. *et al.*, *Tetrahedron*, 1973, **29**, 455

(*synth, pmr*)

Roberts, M.F. *et al.*, *Phytochemistry*, 1981, **20**, 447 (*deriv*)

Shono, T. *et al.*, *Chem. Lett.*, 1984, 1101 (*synth*)

Harding, K.E. *et al.*, *J.O.C.*, 1984, **49**, 40

(*synth*)

Tadano, K. *et al.*, *J. Carbohydr. Chem.*, 1985, **4**, 129-139 (*synth, abs config*)

Shono, T. *et al.*, *J.O.C.*, 1988, **53**, 4118 (*N-Methylpseudoconiine*)

Takahata, H. *et al.*, *Heterocycles*, 1994, **38**, 269

(*synth, Pseudoconiine, N-Methylpseudoconiine*)

Herdeis, C. *et al.*, *Annalen*, 1995, 1295 (*synth, pmr, cmr, ir*)

Sakagami, H. *et al.*, *Chem. Comm.*, 1996, 1433

(*synth*)

Dockner, M. *et al.*, *Liebigs Ann./Recl.*, 1997, 1267 (*synth, pmr, cmr, cryst struct*)

Cossy, J. *et al.*, *Synlett*, 1997, 905-906 (*synth*)

Herdeis, C. *et al.*, *Synthesis*, 1997, 1405-1410 (*Epipseudoconiine*)


Agami, C. *et al.*, *Tetrahedron*, 1998, **54**, 8783-8796 (*synth*)

Lofstedt, J. *et al.*, *Tetrahedron*, 2000, **56**, 2225-2230 (*synth*)

Liu, G. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 1297-1303 (*synth*)

2-Propylquinoline, 9CI P-654

[1613-32-7]



C₁₂H₁₃N 171.241

Alkaloid from the stems of *Galipea bracteata* (Rutaceae). Shows an inhibitory effect on shoot and root growth of lettuce. Pale yellow-green oil. d₄¹⁷ 1.04. Bp₁₃ 142-145° Bp₁₀ 130-131°. n_D²³ 1.5886.

Picrate: [1613-40-7]

Yellow needles or leaflets (EtOH). Mp 163-164°.

Methiodide: [55602-26-1]

Mp 184°.

Decahydro: see *Dendrobates* Alkaloid 181B, A-401

Meisenheimer, J. *et al.*, *Ber.*, 1923, **56**, 1353-1362 (*synth*)

Delaby, R. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1930, **191**, 845-847 (*synth*)

Kudo, T. *et al.*, *Yakugaku Zasshi*, 1975, **95**, 521-527; *CA*, **83**, 146716y (*synth*)

Vieira, P.C. *et al.*, *Phytochemistry*, 1990, **29**, 813-815 (*isol, ir, pmr, cmr, ms*)


Wiebe, J.M. *et al.*, *Tetrahedron*, 1996, **52**, 11705-11724 (*synth, pmr*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 332-334 (*synth, ir, pmr*)

Prosafrine

P-655

α-Ethyl-5-hydroxy-6-methyl-2-piperidine-decanol, 9CI. 3-Hydroxy-6-(10-hydroxy-dodecyl)-2-methylpiperidine



C₁₈H₃₇NO₂ 299.496

(-)-form [38764-78-2]

Alkaloid from the leaves of *Prosopis africana* (Fabaceae). Cryst. (Me₂CO). Mp 68°. [α]_D -8.5 (CHCl₃).

N,O,O-Tri-Ac:

Oil.

10'-Ketone: *Prosafrine*. 12-(5-Hydroxy-6-methyl-2-piperidinyl)-3-dodecanone, 9CI. 3-Hydroxy-2-methyl-6-(10-oxododecyl)piperidine [38764-77-1]

C₁₈H₃₅NO₂ 297.48

Alkaloid from leaves of *Prosopis africana* (Fabaceae). Cryst. (Me₂CO). Mp 56°. [α]_D -8 (CHCl₃). [α]_D -65 (c, 0.9 in EtOH).

(±)-form

10'-Ketone: [74843-66-6]

Mp 78.5-79°.

Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Belg.*, 1972, **81**, 425; 443 (*isol, ms, ir, pmr, struct, abs config*)


Natsume, M. *et al.*, *Heterocycles*, 1980, **14**, 615

(*synth, deriv*)

Prosopine†

P-656

5-Hydroxy-6-(hydroxymethyl)-*α*-methyl-2-piperidineundecanol, 9CI. 3-Hydroxy-6-(11-hydroxydodecyl)-2-hydroxymethylpiperidine [14058-38-9]



C₁₈H₃₇NO₃ 315.495

Alkaloid from leaves of *Prosopis africana* (Fabaceae). Cryst. (Me₂CO). Mp 126°. [α]_D +25 (CHCl₃).

N,O,O,O-Tetra-Ac:

Oil. [α]_D -20 (CHCl₃).

N-Me:

Cryst. (EtOAc). Mp 66°. [α]_D +6.6 (CHCl₃).

11'-Ketone: *Prosopinone*. 12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-2-dodecanone, 9CI. 3-Hydroxy-2-hydroxymethyl-6-(11-oxododecyl)piperidine [14058-55-0]


C₁₈H₃₅NO₃ 313.479

Alkaloid from *Cassia carnaval* (Fabaceae). Mp 90° (synthetic). $[\alpha]_D^{25} +15$.

- Ratle, G. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 2945 (*isol, ir, struct*)
 Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Belg.*, 1972, **81**, 425; 443 (*ir, uv, pmr, ms, abs config*)
 Lythgoe, D. *et al.*, *An. Asoc. Quim. Argent.*, 1972, **60**, 317; *CA*, **77**, 164901 (*Prosopinine*)
 Khuong-Huu, Q. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 267 (*cmr*)

Prosopinine P-657

12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-3-dodecanone, 9CI. 3-Hydroxy-2-hydroxymethyl-6-(10-oxododecyl)piperidine



(+)-form

$C_{18}H_{35}NO_3$ 313.479

(+)-form [14058-39-0]

Alkaloid from leaves of *Prosopis africana* (Fabaceae). Cryst. (EtOAc). Mp 94.5°. $[\alpha]_D^{25} +12$ (CHCl₃).

Hydrochloride: Mp 87°. $[\alpha]_D^{25} 0$ (CHCl₃).

N,O,O-Tri-Ac:

Oil. $[\alpha]_D^{25} -13$ (CHCl₃).

N-Me:

Cryst. (EtOAc). Mp 71°. $[\alpha]_D^{25} +2$ (CHCl₃).

(±)-form

6-Epimer: *Prosophylline*

[38764-76-0]

$C_{18}H_{35}NO_3$ 313.479

Alkaloid from the leaves of *Prosopis africana* (Fabaceae). Cryst. (Me₂CO). Mp 82-83° (79°). Racemic.

6-Epimer, N,O,O-tri-Ac:

Oil.

Ratle, G. *et al.*, *Bull. Soc. Chim. Fr.*, 1966, 2945 (*isol*)

Khuong-Huu, Q. *et al.*, *Bull. Soc. Chim. Belg.*, 1972, **81**, 425; 443 (*ir, uv, pmr, ms, abs config*)
 Natsume, M. *et al.*, *Heterocycles*, 1981, **16**, 973 (*synth, deriv*)

Khuong-Huu, Q. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 267 (*cmr*)

Cook, G.R. *et al.*, *Tet. Lett.*, 1994, **35**, 1669 (*synth*)

Hirai, Y. *et al.*, *J.O.C.*, 1997, **62**, 776 (*synth*)

Kadota, I. *et al.*, *Tet. Lett.*, 1997, **38**, 7469-7470 (*synth, deoxo deriv*)


Koulochesi, S.D. *et al.*, *Tet. Lett.*, 1999, **40**, 6869-6870 (*Prosophylline, synth*)

Cossy, J. *et al.*, *J.O.C.*, 2002, **67**, 1982-1992 (*Prosophylline, synth*)

Couladouros, E.A. *et al.*, *Tet. Lett.*, 2007, **48**, 8227-8229 (*Prosophylline, synth*)

Prostaphabysine P-658

6,7-Didehydro-8,10-epoxy-3,7-dimethoxy-17-methylhasubanan-4,8-diol, 9CI [36871-88-2]



$C_{19}H_{23}NO_5$ 345.394

Alkaloid from the roots and rhizomes of *Stephania abyssinica* and the roots and stems of *Stephania longa* (Menispermaceae). Pale-yellow glass.

Methiodide:

Prisms (MeOH/C₆H₆). Mp 196-198° dec. $[\alpha]_D^{25} -105$ (c, 1.98 in MeOH).

N-De-Me: *Norprostaphabysine*

[863712-30-5]

$C_{18}H_{21}NO_5$ 331.368

Alkaloid from *Stephania longa*. Powder. $[\alpha]_D^{20} -80.4$ (c, 0.52 in CHCl₃). λ_{max} 276 (log ϵ 3.45) (MeOH).

10-Epimer: *Isoprostaphabysine*

[863712-32-7]

$C_{19}H_{23}NO_5$ 345.394


Alkaloid from *Stephania longa*. Powder. $[\alpha]_D^{20} -242.4$ (c, 0.32 in CHCl₃). λ_{max} 276 (log ϵ 3.68) (MeOH).

Kupchan, S.M. *et al.*, *J.O.C.*, 1973, **38**, 151-153 (*Prostaphabysine*)

Zhang, H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1201-1207 (*Norprostaphabysine, isoprostaphabysine*)

Prostaphanaberrine

[105608-27-3]



$C_{19}H_{21}NO_5$ 343.379

Alkaloid from the fresh fruits of *Stephania japonica* (Menispermaceae). Light-yellow prisms (MeOH). Mp 225° dec. $[\alpha]_D^{25} -219.1$ (c, 0.25 in CHCl₃).

Ac:


Light-yellow prisms (MeOH). Mp 169.5°.

Matsui, M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 588 (*isol, uv, ir, pmr, cmr, ms, struct*)

Prothracarcin

P-660

2-Ethylidene-1,2,3,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI. DC 62. Antibiotic DC 62 [81542-99-6]



$C_{14}H_{14}N_2O$ 226.277

Anthramycin antibiotic. Prod. by *Streptomyces umbrosus* ssp. *raffinophilus* NRRL12143. Active against gram-positive and -negative bacteria, murine tumours and leukaemia cells. Analgesic. Sedative. Antitumour and antispasmodic agent. Cryst. (EtOAc). Sol. MeOH; fairly

sol. Et₂O; poorly sol. H₂O, hexane. Mp 87° dec. $[\alpha]_D^{22} +17.1$ (c, 0.1 in EtOAc). λ_{max} 218 (ϵ 21000); 239 (sh) (ϵ); 255 (sh) (ϵ); 316 (ϵ 4000) (MeOH) (Derep). λ_{max} 217 (ϵ 21000); 271; 316 (MeOH) (Berdy). λ_{max} 218 (ϵ 21000); 316 (ϵ 4000) (H₂O) (Berdy).

▶LD₅₀ (mus, ipr) 42 mg/kg. UY8533500

Japan. Pat., 1981, 56 158 785; *CA*, **96**, 179428 (*isol*)

Shimizu, K. *et al.*, *J. Antibiot.*, 1982, **35**, 972

Mori, M. *et al.*, *Tetrahedron*, 1986, **42**, 3793


(*synth, config*)

Weidner-Wells, M.A. *et al.*, *J.O.C.*, 1989, **54**, 5746 (*synth*)

Protophelin

P-661

[131610-88-3]



$C_{31}H_{36}N_4O_{10}$ 624.646

Isol. from a bacterium (DMS No. 5746). Also from *Azotobacter vinelandii*. Siderophore. λ_{max} 248 (ϵ 25120); 312 (ϵ 10000) (MeOH) (Berdy). λ_{max} 550 (ϵ 1995) (pH 1.5 buffer) (Berdy). λ_{max} 480 (ϵ 3980) (pH 13 buffer) (Berdy).

Taraz, K. *et al.*, *Z. Naturforsch., B*, 1990, **45**, 1327 (*isol, pmr, cmr*)

Cornish, A.S. *et al.*, *BioMetals*, 1995, **8**, 332 (*isol, ms*)

Protochondocurarine


P-662

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from Curare. Mp 265° (as nitrate). $[\alpha]_D^{20} +175$ (c, 2 in H₂O) (nitrate).

Bodendorf, K. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1954, **287**, 555; *CA*, **52**, 10119h (*isol*)

Protopuridine

P-663



$C_{36}H_{38}N_2O_6$ 594.706

Probable struct. Alkaloid from curare (Menispermaceae). Plates + 0.5 Py (Py aq.). Mp 295°. Stereochem. undetermined but the two (non-identical) chiral centres have the same chirality since methylation to a symmetrical product does not give a meso-form.

Hydrochloride (1:2): [66723-63-5]

Cryst. + 6H₂O. Mp 295° (efferv.). $[\alpha]_{H_g}^{20} +7.6$ (c, 0.3 in H₂O).

Isomer: *Neoprotocuridine*


$C_{36}H_{38}N_2O_6$ 594.706

Alkaloid from curare (Menispermaceae). Leaflets + 8H₂O (H₂O). Mp 232°. This is either the 6,7 or 6',7' positional isomer of Protopuridine; stereochem. undetermined.

King, H. *et al.*, *J.C.S.*, 1937, 1472-1482 (*isol*)
Bick, I.R.C. *et al.*, *J.C.S.*, 1960, 2402-2407
(*struct*)

Protoemetine

P-664



(-)-form

C₁₉H₂₇NO₃ 317.427

(-)-form [549-91-7]

Alkaloid from *Cephaelis (Psychotria) ipecacuanha* root (Rubiaceae). Noncryst. [α]_D²⁸ -14.38 (c, 0.57 in EtOH) (synthetic).
Perchlorate: Mp 140-142° (hydrate) Mp 193-195° (anhyd.).

Semicarbazone: Mp 168-169°.

Alcohol: **Protoemetinol**. Dihydroprotoemetine

[7344-78-7]

C₁₉H₂₉NO₃ 319.443

Alkaloid detected by mass spectrometry in *Alangium lamarkii* (Alangiaceae) and identified by tlc comparison with an authentic specimen. Prisms (EtOH aq. or H₂O) (as perchlorate). Mp 199-200° (perchlorate). [α]_D²⁴ -15.3 (c, 0.26 in MeOH) (synthetic, perchlorate).

Alcohol, O⁹-de-Me: **9-Demethylprotoemetinol**

[84590-09-0]

C₁₈H₂₇NO₃ 305.416

Alkaloid from the seeds of *Alangium lamarkii* (Alangiaceae). Mp 157-158.5° (synthetic). [α]_D²⁵ -61 (c, 0.50 in EtOH). A direct comparison of the synthetic prod. and natural alkaloid was not possible.

Alcohol, O⁹-de-Me, di-O-Ac:Oil. [α]_D²⁵ -34.3 (c, 0.40 in CHCl₃).Alcohol, O¹⁰-de-Me: **10-Demethylprotoemetinol**

[84590-10-3]

C₁₈H₂₇NO₃ 305.416

Alkaloid from the seeds of *Alangium lamarkii* (Alangiaceae). Glass or amorph. powder. [α]_D¹⁷ -35.2 (c, 0.50 in EtOH). [α]_D -11.9 (CHCl₃).

Alcohol, O¹⁰-de-Me, di-O-Ac:Oil. [α]_D²⁵ -29.7 (c, 0.38 in CHCl₃). [α]_D -15.8 (CHCl₃).

(±)-form [17132-38-6]

Synthetic. Noncryst.

Semicarbazone: Mp 185-186°.

Alcohol: [65982-36-7]

Synthetic. Syrup; prisms (EtOH/EtOAc) (as perchlorate). Mp 178-181° (perchlorate).


Alcohol, O⁹-de-Me:Synthetic. Prisms (Me₂CO). Mp 171-171.5°.Alcohol, O¹⁰-de-Me: [84985-54-6]Synthetic. Prisms (hexane/CHCl₃). Mp 151-152°.

Alcohol, O¹⁰-de-Me, di-O-Ac: Mp 97-98°.
Battersby, A.R. *et al.*, *J.C.S.*, 1959, 1744-1748; 1748-1753 (*isol, struct, synth*)
Albright, J.D. *et al.*, *J. Nat. Prod.*, 1965, 28, 212-217 (*Protoemetinol, occur*)
Szántay, C. *et al.*, *J.O.C.*, 1966, 31, 1447-1451 (*synth*)
Kametani, T. *et al.*, *Heterocycles*, 1977, 8, 119-124 (*synth*)
Takano, S. *et al.*, *Heterocycles*, 1979, 12, 765-770 (*synth, Protoemetinol*)
Kametani, T. *et al.*, *J.C.S. Perkin I*, 1979, 1211-1217 (*synth, ir, pmr*)
Ali, E. *et al.*, *Heterocycles*, 1982, 19, 2301-2304 (*Demethylprotoemetinols*)
Brown, R.T. *et al.*, *Tet. Lett.*, 1984, 25, 3127-3130 (*synth*)
Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1987, 35, 2755-2760 (*Demethylprotoemetinols*)
Hirai, Y. *et al.*, *Chem. Pharm. Bull.*, 1988, 36, 1343-1350 (*synth*)
Ihara, M. *et al.*, *J.C.S. Perkin I*, 1990, 1469-1476 (*synth*)
Huang, C.G. *et al.*, *Tet. Lett.*, 2002, 43, 2721-2723 (*synth*)
Chang, J.-K. *et al.*, *Tetrahedron*, 2008, 64, 9685-9688 (*Protoemetinol, synth, pmr, cmr*)

Protomethine

P-665

2-Phenyl-1,5,9,14-tetraazobicyclo[12.3.1]octadecan-4-one, 9CI
[216855-67-3]

C₂₀H₃₂N₄O 344.499

Related to Protoverbine, P-673. Alkaloid from *Verbascum pseudonobile*. Solid. [α]_D +4.9 (c, 1.4 in CHCl₃).

N⁹-E-Cinnamoyl: **Verbamethine**. Incasine C

[162465-73-8]

C₂₉H₃₈N₄O₂ 474.645

Alkaloid from *Incarvillea sinensis* and *Verbascum pseudonobile*. Off-white powder. [α]_D²⁵ +13.2 (c, 1.3 in CHCl₃) (synthetic). λ _{max} 280 (log ϵ 4.33) (EtOH).

N⁹-Z-Cinnamoyl: **Isoverbamethine**. Incasine C

[162709-45-7]

C₂₉H₃₈N₄O₂ 474.645

Alkaloid from *Incarvillea sinensis* and *Verbascum pseudonobile*. Off-white powder. [α]_D²⁴ +4.2 (c, 0.4 in CHCl₃). Props. refer to Incasine C. Incorr. struct. of Incasine C shown in the ref.

N⁹-(4-Methoxy-E-cinnamoyl): **Verbamekrine**

[249739-46-6]

C₃₀H₄₀N₄O₃ 504.671Alkaloid from *Verbascum pseudonobile*.N⁹-(4-Methoxy-Z-cinnamoyl): **Isoverbamekrine**

[249739-47-7]

C₃₀H₄₀N₄O₃ 504.671Alkaloid from *Verbascum pseudono-*

bile.

N⁹-(3,4-Dimethoxy-E-cinnamoyl): **Verbametrine**

[223519-65-1]

C₃₁H₄₂N₄O₄ 534.697

Alkaloid from *Verbascum pseudonobile*. Glass. [α]_D²⁸ -8 (c, 2 in CHCl₃). λ _{max} 232 (sh) (log ϵ 3.44); 296 (sh) (log ϵ 3.44); 320 (log ϵ 3.5) (EtOH).

N⁹-(3,4-Dimethoxy-Z-cinnamoyl): **Isoverbametrine**

[223519-66-2]

C₃₁H₄₂N₄O₄ 534.697Alkaloid from *Verbascum pseudonobile*.18-Oxo, N⁹-E-cinnamoyl: **Verbascine**.*Verbaskine*

[12651-38-6]

C₂₉H₃₆N₄O₃ 488.628

Alkaloid from *Verbascum pseudonobile* (Scrophulariaceae). Cryst. (Me₂CO). Mp 120-123°. [α]_D²⁰ -26 (CHCl₃). Artifact. λ _{max} 219 (ϵ 14500); 225 (ϵ 11200); 284 (ϵ 15800) (EtOH) (Derep).

 $\Delta^{1,18}$ ($\Delta^{14,18}$)-Dehydro, N⁹-E-cinnamoyl:**Incasine B'**

[185139-47-3]

C₂₉H₃₇N₄O₂[⊕] 473.637

Alkaloid from *Incarvillea sinensis* (Bignoniaceae). Off-white powder (as chloride). [α]_D³¹ +49.2 (c, 1.3 in CHCl₃) (chloride).

 $\Delta^{1,18}$ ($\Delta^{14,18}$)-Dehydro, N⁹-Z-cinnamoyl:**Incasine B**

[185139-27-9]

C₂₉H₃₇N₄O₂[⊕] 473.637


Alkaloid from *Incarvillea sinensis*. Off-white powder (as chloride). [α]_D³² +62.5 (c, 0.5 in CHCl₃) (chloride). +ve Charge delocalised over the two ring N atoms.

Koblicová, Z. *et al.*, *Tet. Lett.*, 1983, 24, 4381 (*Verbascine*)Ishihara, K. *et al.*, *J.A.C.S.*, 1996, 118, 1569 (*Verbascine, synth*)Chi, Y.M. *et al.*, *Tetrahedron*, 1997, 38, 2713-2716 (*Incasines*)Drandarov, K. *et al.*, *Helv. Chim. Acta*, 1999, 82, 229-237; 1185-1194 (*Verbascum alkaloids*)Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 2000, 83, 3035-3042 (*Protomethine*)Drandarov, K. *et al.*, *Helv. Chim. Acta*, 2002, 85, 979-989 (*synth*)Chi, Y.-M. *et al.*, *J. Asian Nat. Prod. Res.*, 2007, 9, 115-118 (*Incasine C'*)

Protophomin

P-666

[52212-93-8]

C₂₉H₃₅NO₃ 445.6

Constit. of cultures of *Phoma* spp. Cryst. (Me₂CO). Mp 252-256°. [α]_D²⁴ -112 (c, 0.042 in CHCl₃). λ_{\max} 206 (ε 23100); 217 (ε 12300); 243 (ε 10700) (EtOH) (Derep). λ_{\max} 248 (ε 13200) (EtOH) (Berdy).


Binder, M. et al., *Helv. Chim. Acta*, 1973, **56**, 2387 (isol)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 337

Protopine

P-667

4,6,7,14-Tetrahydro-5-methylbis[1,3]benzodioxolo[4,5-c:5',6'-g]azecin-13(5H)-one, 9CI. Fumarine. *Corydalis* C. *Corydine*. *Biflorine*†. *Macleyine* [130-86-9]



C₂₀H₁₉NO₅ 353.374

The N-protonated form adopts a tetracyclic struct. in soln. or in the solid state. Alkaloid from a wide variety of genera in the Papaveraceae (*Argemone*, *Bocconia*, *Chelidonium*, *Eschscholtzia*, *Glaucium*, *Hunnemannia*, *Hylomecon*, *Macleaya*, *Meconella*, *Meconopsis*, *Papaver*, *Roe-meria*, *Romneya*, *Sanguinaria*, *Stylomecon*, *Corydalis*, *Dactylicapnos*, *Dicentra*, *Fumaria*), Berberidaceae (*Berberis*) and Sapindaceae (*Pteridophyllum*). Shows weak spasmolytic, smooth muscle stimulant and weak antineoplastic activity. Exp. coronary vessel dilator. Shows bactericidal activity. Sedative. Antihypertensive and anti-amnesic agent. Cryst. (CHCl₃/EtOH). Mp 207°. pK_a 5.99. Log P 2.63 (uncertain value) (calc). λ_{\max} 288 (ε 4000) (MeOH) (Berdy). λ_{\max} 293 (ε 8500) (EtOH) (Berdy). λ_{\max} 240 ; 290 (MeOH/HCl) (Berdy).

► LD₅₀ (gpg, orl) 237 mg/kg. VS2800000

Hydrochloride: [6164-47-2]

Mp 261-264° dec.

Perchlorate: Mp 321-322°.

Methiodide: Mp 217°.

N-Oxide: Protopine N-oxide

[87264-51-5]

C₂₀H₁₉NO₆ 369.373

Minor alkaloid from *Bocconia cordata* (Papaveraceae). Cryst. + 1H₂O (MeOH/Me₂CO). Mp 145-146° dec.

14-Alcohol: *Dihydroprotopine*

C₂₀H₂₁NO₅ 355.39

Alkaloid from *Papaver somniferum* (Papaveraceae).

8-Oxo: *Leptocarpine*†. Alkaloid HLI. 8-Oxoprotopine

[164176-15-2]

C₂₀H₁₇NO₆ 367.357

Alkaloid from whole plants of *Hypecoum leptocarpum* (Papaveraceae). Yellow needles (as hydroiodide). Mp 246-248° (hydroiodide).

13-Oxo: *13-Oxoprotopine*. 14-Oxoprotopine. *Oxyprotopine*

[15211-02-6]

C₂₀H₁₇NO₆ 367.357

Alkaloid from the seeds of *Papaver atlanticum* (Papaveraceae). Also from *Hypecoum procumbens* (Hypecoaceae). Mp 222-225°.

1-Methoxy: *Coulteropine*

[6014-62-6]

[26012-86-2]

C₂₁H₂₁NO₆ 383.4

Alkaloid from the roots of *Romneya coulteri* var. *trichocalyx* (Papaveraceae). Cryst. (C₆H₆). Mp 168-170°.

1-Methoxy, hydrochloride: Mp 149-150°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 616C (nmr)

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 1513F (ir)

Haworth, R.D. et al., *J.C.S.*, 1926, 1769 (synth)

Dolejš, L. et al., *Coll. Czech. Chem. Comm.*, 1964, **29**, 2479 (ms)

Ma, J.C.N. et al., *Can. J. Chem.*, 1965, **43**, 1849 (pmr)

Stermitz, F.R. et al., *Tetrahedron*, 1966, **22**, 1095 (*Coulteropine*)

Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1967, **32**, 3414 (uv)

Anet, F.A.L. et al., *Tet. Lett.*, 1967, 4881 (pmr)

Stermitz, F.R. et al., *Tet. Lett.*, 1968, 3915

(*Coulteropine, cryst struct*)

Preininger, V. et al., *Pharmazie*, 1970, **25**, 356 (isol, 13-Oxoprotopine)

Battersby, A.R. et al., *J.C.S. Perkin 1*, 1975, 1147 (biosynth)

Takao, N. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2859 (biosynth)

Holland, H.L. et al., *Can. J. Chem.*, 1979, **57**, 1588-1597 (biosynth)

Šantavý, F. et al., *Alkaloids (Academic Press)*, 1981, **19**, 461 (*Dihydroprotopine*)

Iwasa, K. et al., *Phytochemistry*, 1983, **22**, 627 (isol, uv, ir, pmr, ms, struct, synth, oxide)

Castedo, L. et al., *Heterocycles*, 1986, **24**, 5 (synth, 13-Oxoprotopine)

Táborská, E. et al., *Coll. Czech. Chem. Comm.*, 1987, **52**, 508 (*Alkaloid HLI*)

Kulkarni, B.K. et al., *J. Het. Chem.*, 1990, **27**, 623 (synth)

Táborská, E. et al., *Heterocycles*, 1995, **41**, 799 (*Leptocarpine*)

Marek, R. et al., *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Dostál, J. et al., *Acta Cryst. C*, 2001, **57**, 651-652 (*hydrochloride, cryst struct*)

Seger, C. et al., *Magn. Reson. Chem.*, 2004, **42**, 882-886 (pmr, cmr)

Toušek, J. et al., *Magn. Reson. Chem.*, 2005, **43**, 578-581 (*HCl salts, pmr, cmr*)


Wada, Y. et al., *J.O.C.*, 2007, **72**, 7301-7306 (synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FOW000

Protosinomenine

P-668

[30883-59-1]



C₁₉H₂₃NO₄ 329.395

Alkaloid from the pods of *Erythrina*

lithosperma (Fabaceae). Gum.

Picolonate:

Yellow needles (EtOH). Mp 172-174°.

N-De-Me: *Norprotosinomenine*

[26153-70-8]

C₁₈H₂₁NO₄ 315.368

Alkaloid from the pods of *Erythrina lithosperma* (Fabaceae). Mp 242-244° (as hydrochloride). [α]_D²⁵ +18 (c, 0.88 in H₂O).

Di-Me ether: see Laudanosine, L-69

Ghosal, S. et al., *Aust. J. Chem.*, 1971, **24**,


2733-2735 (isol, uv, ir, pmr, ms, struct)

Maier, U.H. et al., *Phytochemistry*, 1999, **52**, 373-382 (biosynth, pmr, cmr)

Protostemodiol

P-669

[1002109-45-6]



C₂₃H₃₃NO₇ 435.516

Alkaloid from *Stemona japonica*.


Amorph. yellow powder. [α]_D²³ +124.7 (c, 0.15 in MeOH). λ_{\max} 305 (log ε 4.25) (MeOH).

Tang, C.-P. et al., *J. Nat. Prod.*, 2008, **71**, 112-116 (isol, pmr, cmr, ms)

Protostemonine

P-670

[27495-40-5]



Absolute Configuration

C₂₃H₃₁NO₆ 417.501

Alkaloid from the roots of *Stemona*

cochinchinensis and *Stemona japonica*.

Cryst. + 1MeOH (MeOH). Mp 172-173° (sintering at 90-100°). [α]_D +147.8.

Picolonate: Mp 237°.

1 α ,8 α -Epoxide: *Oxyprotostemonine*

C₂₃H₂₉NO₇ 431.485

Alkaloid from the roots of *Stemona*

curtisii and *Stemona kerrii*. Amorph.

[α]_D²⁰ +142 (c, 0.2 in MeOH). λ_{\max} 298 (MeOH aq.).

7,8-Didehydro: *Dehydroprotostemonine*

C₂₃H₂₉NO₆ 415.485

Alkaloid from the roots of *Stemona*

curtisii and *Stemona kerrii*. Amorph.

[α]_D²⁰ +72 (c, 0.3 in MeOH). λ_{\max} 312 (MeOH aq.).

1,2,3,9a-Tetrahydro: *Bisdehydroprotostemonine*. *Tetrahydroprotostemonine*

[145701-15-1]

C₂₃H₂₇NO₆ 413.469

Alkaloid from roots of *Stemona japonica*. Mp 192-194°. [α]_D +169 (c, 0.81 in EtOH).

Demethoxy, 11S,12R-dihydro: 12-Epistemochinine

$C_{22}H_{31}NO_5$ 389.491

Alkaloid from *Stemona japonica*. Yellow oil. $[\alpha]_D^{23} +53.3$ (c, 0.04 in MeOH). λ_{max} 294 (log ϵ 3.49) (MeOH).

Demethoxy, 11S,12S-dihydro: Stemocochinine

$C_{22}H_{31}NO_5$ 389.491

Alkaloid from the roots of *Stemona cochinchinensis*, *Stemona curtisii* and *Stemona kerrii*. Amorph. $[\alpha]_D^{20} -52$ (c, 0.2 in MeOH).

Demethoxy, 1,2,3,9a-tetrahydro, 11S,12R-dihydro: Isobisdehydrostemocochinine

[1003567-57-4]

$C_{22}H_{27}NO_5$ 385.459

Alkaloid from the roots of *Stemona cochinchinensis*. Amorph. yellow powder. $[\alpha]_D^{20} -55$ (c, 0.17 in $CHCl_3$). λ_{max} 240 (MeOH).

Demethoxy, 1,2,3,9a-tetrahydro,

11S,12S-dihydro: Bisdehydrostemocochinine

[1003567-56-3]

$C_{22}H_{27}NO_5$ 385.459

Alkaloid from the roots of *Stemona cochinchinensis*. Amorph. yellow powder. $[\alpha]_D^{20} -9$ (c, 0.15 in $CHCl_3$). λ_{max} 239 (MeOH).

(E)-Isomer: *Isoprotostemonine*

[145841-08-3]

$C_{23}H_{31}NO_6$ 417.501

Alkaloid from roots of *Stemona japonica*. Light yellow cryst. Mp 165-167°. $[\alpha]_D -23.6$ (c, 0.47 in EtOH).

Irie, H. *et al.*, *Chem. Comm.*, 1970, 268-269 (uv, ir, pmr, struct)

Irie, H. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 451-452 (cryst struct, abs config)

Ye, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1205-1208 (*Isoprotostemonine*, *Bisdehydroprotostemonine*)

Kaltenegger, E. *et al.*, *Phytochemistry*, 2003, **63**, 803-816 (isol, pmr, cmr, ms)

Greger, H. *et al.*, *Planta Med.*, 2006, **72**, 99-113 (rev, occur, activity)


Lin, L.-G. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2167-2175 (*Bisdehydrostemocochinine*, *Isobisdehydrostemocochinine*)

Wang, C.-P. *et al.*, *J. Nat. Prod.*, 2008, **71**, 112-116 (*12-Epistemocochinine*, *Stemocochinine*)

Protostephanine

P-671

6,7,8,9-Tetrahydro-2,3,10,12-tetramethoxy-7-methyl-5H-dibenz[d,f]azocine, 9CI
[549-28-0]



$C_{21}H_{27}NO_4$ 357.449

Alkaloid from *Stephania japonica* (Menispermaceae). Antihypertensive agent. Cryst. (MeOH). Mp 73-74° (MeOH solvate) Mp 90-95° (solvent-free). Log P 3.68 (calc).

Hydrochloride: Mp 150° dec.

Picrate: Mp 207-209°.

Barton, D.H.R. *et al.*, *J.C.S. (C)*, 1966, 2313 (isol, pmr)

Pecherer, B. *et al.*, *J.O.C.*, 1967, **32**, 1053 (synth)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1968, 1214; 1974, 773

Pecherer, B. *et al.*, *J. Med. Chem.*, 1969, **12**, 149 (synth)

Battersby, A.R. *et al.*, *Tet. Lett.*, 1977, 1321 (synth, biosynth)


Matsui, M. *et al.*, *Phytochemistry*, 1979, **18**, 1087 (isol)

Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1981, 2002 (synth, uv, pmr, ms)

Protostrychnine

P-672

12-Hydroxy-12,24-secostrychnidin-10-one, 9CI. 12,13-Dihydro-12-hydroxyisostrychnine
[71610-48-5]



Absolute Configuration

$C_{21}H_{24}N_2O_3$ 352.432

Minor alkaloid from the root bark of *Strychnos nux-vomica* and root of *Strychnos icaja* (Loganiaceae). Mp 201° dec. $[\alpha]_D^{20} +5$ (c, 0.39 in MeOH).

17-Deoxy, 17,18-didehydro: *Isostrychnine*.

Isostrychnine I

[467-16-3]

$C_{21}H_{22}N_2O_2$ 334.417

Alkaloid from root bark seeds and leaves of *Strychnos nux-vomica* (Loganiaceae). Needles (EtOAc). Mp 225-228°. $[\alpha]_D +25$ (EtOH). $[\alpha]_D -41.9$ (AcOH). λ_{max} 256 (log ϵ 4.05); 285 (log ϵ 3.6); 294 (log ϵ 3.5) (EtOH).

17-Deoxy, 17,18-didehydro, N⁴-oxide:

Isostrychnine N^b-oxide

[130641-44-0]

$C_{21}H_{22}N_2O_3$ 350.416

Alkaloid from the seeds of *Strychnos nux-vomica*. Component of Ma Qian Zi.

17-Deoxy, 17,18-didehydro, 19,20-dihydro: *19,20-Dihydroisostrychnine*

[14839-10-2]

$C_{21}H_{24}N_2O_2$ 336.433

Alkaloid from leaves of *Strychnos nux-vomica* (Loganiaceae). $[\alpha]_D^{21} +55$ (c, 0.044 in MeOH). λ_{max} 254 ; 282 ; 292 (MeOH).

11-Methoxy, 10-hydroxy, 17-deoxy, 17,18-didehydro: *O-Demethylisobrucine*

[130641-45-1]

$C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from the processed seeds of *Strychnos nux-vomica*. Possible artifact.

10,11-Dimethoxy, 17-deoxy, 17,18-didehydro: *Isobrucine*

[129724-78-3]

$C_{23}H_{26}N_2O_4$ 394.469

Alkaloid from the processed seeds of *Strychnos nux-vomica*. Possible artifact.

10,11-Dimethoxy, 17-deoxy, 17,18-didehydro, N⁴-oxide: *Isobrucine N-oxide*
[130641-43-9]

$C_{23}H_{26}N_2O_5$ 410.469

Alkaloid from the processed seeds of *Strychnos nux-vomica*. Possible artifact.

Leuchs, H. *et al.*, *Ber.*, 1942, **75**, 573-579 (*Isostrychnine*, synth)

Heimberger, S.I. *et al.*, *Chem. Comm.*, 1973, 217-218 (*Isostrychnine*, isol)

Galeffi, C. *et al.*, *J. Chromatogr.*, 1974, **88**, 416-418 (*Isostrychnine*, uv, ms, pmr)

Leung, J. *et al.*, *Org. Magn. Reson.*, 1977, **9**, 333-337 (cmr)

Baser, K.H.C. *et al.*, *Phytochemistry*, 1979, **18**, 512-514; 1982, **21**, 1423-1429 (*Isostrychnine*, *Dihydroisostrychnine*, isol, uv, ir, pmr, cmr, ms, struct)

Cai, B.C. *et al.*, *CA*, 1990, **113**, 237637c (isol, struct)

Cai, B.C. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1295-1298 (*Strychnos nux-vomica* alkaloids)


Cai, B.C. *et al.*, *Yaoxue Xuebao*, 1994, **29**, 44-48 (*Isostrychnine N-oxide*)

Philippe, G. *et al.*, *Phytochemistry*, 2003, **62**, 623-629 (isol, pmr, cd)

Protoberbine

P-673

8-Phenyl-1,5,9,13-tetraazaheptadecan-6-one, 9CI



$C_{19}H_{32}N_4O$ 332.488

Numbering shown here as given in the lit. by analogy with related alkaloid types, e.g. Protomethine, P-665. CAS numbering differs. $[\alpha]_D -28.6$ (c, 1.5 in $CHCl_3$).

(S)-form [216855-63-9]

Alkaloid from *Verbascum pseudonobile*. Solid.

N⁹-E-Cinnamoyl: *Verbacine*

[164230-52-8]

[12651-39-7 Verbasine]

$C_{28}H_{38}N_4O_2$ 462.634

Alkaloid from leaves of *Verbascum pseudonobile*. Glass. $[\alpha]_D^{26} -14.4$ (c, 3.7 in $CHCl_3$). The earlier isolate Verbasine was a mixt. of Verbacine with other alkaloids. λ_{max} 280 (ε 20400) (EtOH) (Derep).

N⁹-E-Cinnamoyl, N¹⁴-formyl: *Verbamedine. Incasine A'*

[459867-41-5]

[185135-06-2]

$C_{29}H_{38}N_4O_3$ 490.644

Alkaloid from *Incarvillea sinensis* and *Verbascum pseudonobile*. Glassy solid. $[\alpha]_D -31$ (c, 1.6 in $CHCl_3$). Incasine A' shown to be identical to Verbamedine and prev. published struct. incorrect.

N⁹-E-Cinnamoyl, N¹⁴-Ac: *Verbascenine*
[85559-65-5]

$C_{30}H_{40}N_4O_3$ 504.671


Alkaloid from the aerial parts of *Verbascum phoenicum* and *Verbascum*

- nigrum*. Amorph. $[\alpha]_D^{22}$ -15 (c, 0.43 in MeOH). λ_{\max} 217 (sh) (ϵ 21400); 223 (sh) (ϵ 15100); 280 (ϵ 18600); 299 (sh) (ϵ 12300) (no solvent reported).
- N⁹-Z-Cinnamoyl: Verballocline**
[164176-10-7]
C₂₈H₃₈N₄O₂ 462.634
Alkaloid from leaves of *Verbasicum pseudonobile*. Glass. $[\alpha]_D^{26}$ -18 (c, 2.37 in CHCl₃). λ_{\max} 256 (ϵ 16600) (EtOH) (Derep).
- N⁹-Z-Cinnamoyl, N¹⁴-formyl: Isoverbamedine**
Incasine A
[459867-42-6]
[185054-72-2]
C₂₉H₃₈N₄O₃ 490.644
Alkaloid from *Incarvillea sinensis* and *Verbasicum pseudonobile*. Glassy solid. $[\alpha]_D$ -18 (c, 0.9 in CHCl₃). Incasine A shown to be identical to Isoverbamedine and prev. published struct. incorrect.
- N⁹-Z-Cinnamoyl, N¹⁴-Ac: Verballoscennine**
[188838-50-8]
C₃₀H₄₀N₄O₃ 504.671
Alkaloid from leaves of *Verbasicum phoenicum*. Glass. $[\alpha]_D^{25}$ -19 (c, 2.6 in CHCl₃), $[\alpha]_D^{25}$ -10.7 (c, 2.38 in MeOH). λ_{\max} 256 (log ϵ 4.03) (EtOH).
- N⁹-(4-Methoxy-E-cinnamoyl): Verbasikrine**
[249739-44-4]
[247079-00-1]
C₂₉H₄₀N₄O₃ 492.66
Alkaloid from *Verbasicum pseudonobile*.
- N⁹-(4-Methoxy-Z-cinnamoyl): Isoverbasisikrine**
[249739-45-5]
[247079-01-2]
C₂₉H₄₀N₄O₃ 492.66
Alkaloid from *Verbasicum pseudonobile*.
- N⁹-(3,4-Dimethoxy-E-cinnamoyl): Verbasitrine**
[223519-63-9]
C₃₀H₄₂N₄O₄ 522.686
Minor alkaloid from leaves of *Verbasicum pseudonobile* (Scrophulariaceae). $[\alpha]_D^{25}$ -14 (c, 1.85 in CHCl₃). λ_{\max} 294 (sh) (log ϵ 3.66); 320 (log ϵ 3.71) (EtOH).
- N⁹-(3,4-Dimethoxy-Z-cinnamoyl): Isoverbasisitrine**
[223519-64-0]
C₃₀H₄₂N₄O₄ 522.686
Minor alkaloid from leaves of *Verbasicum pseudonobile*. $[\alpha]_D^{28}$ -15 (c, 2 in CHCl₃). λ_{\max} 230 (sh) (log ϵ 4.19); 272 (log ϵ 4.08); 296 (sh) (log ϵ 3.97) (EtOH).
- 4-Hydroxy: Prelandrine**
[312934-15-9]
C₁₉H₃₂N₄O₂ 348.487
Alkaloid from the roots of *Aphelandra squarrosa*. Solid (as trihydrochloride). $[\alpha]_D^{20}$ -9.3 (c, 0.76 in MeOH) (trihydrochloride).
- 4-Methoxy: Buchnerine**
[151813-62-6]
C₂₀H₃₄N₄O₂ 362.514

- Trace alkaloid from leaves of *Clerodendrum buchneri*. Oil. $[\alpha]_D^{22}$ -26 (c, 0.5 in MeOH).
- 4'-Methoxy, N⁹-(4-methoxy-Z-cinnamoyl): N¹-(Z)-p-Methoxycinnamoylbuchnerine**
[151194-27-3]
C₃₀H₄₂N₄O₄ 522.686
Trace alkaloid from leaves of *Clerodendrum buchneri* (Verbenaceae). Oil. $[\alpha]_D^{22}$ -28 (c, 1.2 in MeOH).
- Seifert, K. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2540-2547 (*Verbasennine*)
Lumbu, S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1418-1420 (*Buchnerine*)
Drandarov, K. *et al.*, *Tet. Lett.*, 1995, **36**, 617-620 (*Verbasine, Verballocline, isol, struct*)
Ishihara, K. *et al.*, *J.A.C.S.*, 1996, **118**, 1569-1570 (*Verbasennine, Verbasine, Buchnerine, synth*)
Drandarov, K. *et al.*, *Phytochemistry*, 1997, **44**, 971 (*Verballoscennine*)
Chi, Y.M. *et al.*, *Tet. Lett.*, 1997, **38**, 2713-2716 (*Incasines*)
Drandarov, K. *et al.*, *Helv. Chim. Acta*, 1998, **81**, 1773-1791; 1999, **82**, 229-237; 1185-1194 (*Verbasitrine, Isoverbasisitrine, Verbasikrine, Isoverbasisikrine*)
Guggisberg, A. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 3035-3042 (*Protoverbine, isol, biosynth*)
Nezbedova, L. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 172-179 (*Prelandrine*)
Drandarov, K. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 979-989 (*synth*)
Drandarov, K. *et al.*, *Tet. Lett.*, 2002, **43**, 5025-5027 (*Verbamedine, Isoverbamedine, Incasines*)

Protoverine P-674

4,9-Epoxy-3,4,6,7,14,15,16,20-cevaneocitol, 9Cl. 6-Hydroxygermine [76-45-9]



- C₂₇H₄₃NO₉ 525.638
Hydrol. prod. of Protoveratrine and other ester alkaloids. Veratrum extracts have been used medicinally to reduce blood pressure. Mp 210-216° dec. $[\alpha]_D^{25}$ -12 (c, 0.96 in Py). Log P -4.7 (uncertain value) (calc).
- ▶ LD₅₀ (mus, scu) 520 mg/kg. FL5700000
14,15-O-Isopropylidene: Mp 253-256°.
15-O-(2-Methylbutanoyl), 6,7-di-Ac: *Desatrine* [6746-01-6]
Obt. semisynthetically by oxidn. of Protoveratrine B. Patent claims use as insecticide, but doubtless an antihypertensive agent. Needles (Me₂CO/petrol). Mp 232-233°.
- 3-Angeloyl, 15-O-(2R-methylbutanoyl), 6,7-di-Ac: *Escholerine* [663-93-4]
Alkaloid from *Veratrum eschscholtzii*. Antihypertensive agent. Mp 235°. $[\alpha]_D$


- 30 (c, 1.0 in Py).
- 3-O-(2-Hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl): *Didesacetylprotoveratrine A* [41535-99-3]
C₃₇H₅₉NO₁₂ 709.873
Alkaloid from *Veratrum nigrum* roots. Cryst. (C₆H₆). Mp 202-204°. $[\alpha]_D^{22}$ -18.7 (c, 0.95 in Py). Shown not to be an artifact.
- 3-O-(2R-Hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 6-Ac: *Desacetylprotoveratrine A* [67375-42-2]
C₃₉H₆₁NO₁₃ 751.91
Alkaloid from *Veratrum album* (Liliaceae). Mp 191-192°. $[\alpha]_D$ -15 (Py).
- 3-O-(2R-Hydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 6,7-di-Ac: *Protoveratrine A. Protoveratrine. Pro-Amid. Protalba. Puroverin* [143-57-7]
C₄₁H₆₃NO₁₄ 793.947
Alkaloid from *Veratrum album, Veratrum viride* and *Veratrum nigrum* (Liliaceae). Shows antifungal activity. Strong antihypertensive agent showing high toxicity in use due to powerful emetic action. Mp 270-271°. $[\alpha]_D$ -40 (Py). Log P 0.59 (uncertain value) (calc). Also used as mixt. with Protoveratrine B (Ebrant, Provell, Tensatrin, Veralba, PVS 295).
- ▶ LD₅₀ (rat, orl) 5 mg/kg. V. toxic if swallowed. FL5750100
- 3-O-(2R,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 6-Ac: *Desacetylprotoveratrine B. Desacetylneoprotoveratrine* [67370-03-0]
C₃₉H₆₁NO₁₄ 767.909
Alkaloid from *Veratrum album* and *Veratrum viride*. Weak antihypertensive agent. Mp 182-183°. $[\alpha]_D$ -9 (Py).
- 3-O-(2R,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 6,7-di-Ac: *Protoveratrine B. Neoprotoveratrine. Veratrine* [124-97-0]
C₄₁H₆₃NO₁₅ 809.946
Alkaloid from *Veratrum album* and *Veratrum viride* (Liliaceae). Shows antifungal activity. Strong antihypertensive agent showing high toxicity in use due to powerful emetic action. Mp 269-270°. $[\alpha]_D$ -39 (Py).
- ▶ LD₅₀ (mus, scu) 0.21 mg/kg. V. toxic by subcutaneous and other routes. FL5750000
- 3-O-(2S,3R-Dihydroxy-2-methylbutanoyl), 15-O-(2R-methylbutanoyl), 6,7-di-Ac: *Protoveratrine C* [82535-71-5]
C₄₁H₆₃NO₁₅ 809.946
Alkaloid from *Veratrum album*. Cryst. solid. Mp 258-260° dec. $[\alpha]_D^{26}$ -6.6 (CHCl₃).
- 3-O-(3,4-Dimethoxybenzoyl), 15-O-(2 ζ -methylbutanoyl): *15-O-(2-Methylbutanoyl)-3-O-veratroylprotoverine* C₄₁H₅₉NO₁₃ 773.916
Alkaloid from *Veratrum nigrum*. Amorph. powder. $[\alpha]_D^{25}$ +12.5 (c, 0.02

in CHCl_3).

- Jacobs, W.A. *et al.*, *J. Biol. Chem.*, 1943, **149**, 271-279 (*isol*)
 Klohs, M.W. *et al.*, *J.A.C.S.*, 1954, **76**, 1152-1153 (*Escholerine*)
U.S. Pat., 1959, 3 009 917; *CA*, **56**, 4810b (*Desatrine*)
 Kupchan, S.M. *et al.*, *J.A.C.S.*, 1960, **82**, 2242-2251; 2252-2258 (*struct, Protoveratrines A,B*)
 Tomko, J. *et al.*, *Planta Med.*, 1962, **10**, 138-142 (*isol*)
 Wolters, B. *et al.*, *Planta Med.*, 1970, **19**, 189-193 (*activity*)
 Bondarenko, N.V. *et al.*, *Khim. Prir. Soedin.*, 1972, 810-811; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, 800-801 (*Didesacetylprotoveratine A*)
 Carey, F.A. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 141-144 (*cmr*)
 Saksena, A.K. *et al.*, *Tet. Lett.*, 1982, **23**, 811-814 (*Protoveratine C*)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., *Pharmaceutical Press*, 1993, 391
 Wang, B. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 244-248 (*15-Methylbutanoyl-3-veratroylprotoverine*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., *J. Wiley*, 2000, NCI600; POF000

Proviolacein P-675


5-(5-Hydroxy-1H-indol-3-yl)-3-(1H-indol-3-yl)-2H-pyrrol-2-one, 9CI
 [165133-94-8]



- $\text{C}_{20}\text{H}_{13}\text{N}_3\text{O}_2$ 327.342
 Prod. by *Chromobacterium violaceum*.
 Red cryst. (MeOH). Mp >240°. λ_{max} 271 (ε 12070); 290 (ε 13000); 385 (ε 8050); 412 (ε 8500); 492 (ε 16700) (EtOH aq.).
 5-Deoxy: 3,5-Di-1H-indol-3-yl-2H-pyrrol-2-one, 9CI. **Prodeoxyviolacein**
 [158099-50-4]
 $\text{C}_{20}\text{H}_{13}\text{N}_3\text{O}$ 311.342
 Prod. by *Chromobacterium violaceum*.
 Dark red cryst. (MeOH). Mp >275°.
 Hoshino, T. *et al.*, *J.C.S. Perkin 1*, 1995, 1565-1571 (*isol, pmr, cmr, ir, 5-deoxy*)

Proximicin A P-676

[1029061-04-8]




- $\text{C}_{12}\text{H}_{11}\text{N}_3\text{O}_6$ 293.235
 Analogue of Netropsin, N-180. *Isol.* from a marine-derived *Verrucosipora maris* AB-18-032. Cytotoxic.
 N^1 -[2-(4-Hydroxyphenyl)ethyl]: **Proxi-**

micin B

- [1029061-05-9]
 $\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_7$ 413.386
Isol. from *Verrucosipora maris* AB-18-032.
 N^1 -[2-(1H-Indol-3-yl)ethyl]: **Proximicin C**
 [1029061-06-0]
 $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_6$ 436.423
Isol. from *Verrucosipora maris* AB-18-032.
 Schneider, K. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 3258-3261 (*ms, struct*)
 Fiedler, H.-P. *et al.*, *J. Antibiot.*, 2008, **61**, 158-163 (*isol, pmr, cmr*)

Proxiphomin P-677

16-Methyl-10-phenylmethyl[13]cytochalasa-6,13,21-triene-1,23-dione, 9CI
 [52212-92-7]



- $\text{C}_{29}\text{H}_{37}\text{NO}_2$ 431.617
 Constit. of cultures of *Phoma* spp. Has antibiotic props. Amorph. solid. $[\alpha]_{\text{D}}^{24}$ -140 (c, 0.156 in CHCl_3). λ_{max} 206 (ε 23100); 217 (ε 12300); 243 (ε 10700) (EtOH) (Derep). λ_{max} 217 (ε 12300); 243 (ε 10700) (EtOH) (Berdy).
 Binder, M. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 2387 (*isol, struct, ir, uv, ms, pmr*)
 Schmidlin, T. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 2096 (*synth*)
 Tapoleczay, D.J. *et al.*, *Chem. Comm.*, 1985, 143 (*synth*)
 Thomas, E.J. *et al.*, *J.C.S. Perkin 1*, 1989, 499 (*synth*)
 Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 336

Prurienidine P-678

- Struct. unknown. Possibly contains a pyridine nucleus. Alkaloid from *Mucuna pruriens* (Fabaceae). Light-buff micro-cryst. solid. Mp 287-288°. *Hydrochloride*: Mp 282-283°. *Sulfate*: Mp 175-177°. *Picrate*: Mp 269-270°. *Chloroaurate*: Mp 300° (blackens at 200°).
 Majumdar, D.N. *et al.*, *Indian Pharm.*, 1954, **10**, 79-84; *CA*, **49**, 9881a (*isol*)

Prurienine P-679

- $\text{C}_6\text{H}_{12}\text{N}_2\text{O}_2$ 144.173
 Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Vasodilator, hypotensive agent, peristaltic stimulant. Fine needles (Me_2CO). Mp 263-264° dec. *Sulfate*: Mp 226°. *Picrate*: Yellowish cryst. Mp 246° dec.


Chloroaurate: Mp 300°.

- Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1953, **15**, 62-65; *CA*, **48**, 8794b (*isol*)

Prurieninine P-680

- $\text{C}_8\text{H}_{16}\text{N}_2\text{O}_2$ 172.227
 Struct. unknown. Alkaloid from *Mucuna pruriens* (Fabaceae). Clusters of needles (EtOH). Mp 213-214° (209-210° dec.). *Sulfate*: Mp 202° (197-198°). *Picrate*: Mp 276-278°. *Chloroaurate*: Mp 199-200°.
 Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1953, **15**, 62-65; *CA*, **48**, 8794b (*isol*)
 Majumdar, D.N. *et al.*, *Indian Pharm.*, 1954, **10**, 79-84; *CA*, **49**, 9881a (*isol*)


Przewalidine P-681



- $\text{C}_{20}\text{H}_{24}\text{NO}_4^{\oplus}$ 342.414
 Quaternary alkaloid from the roots of *Thalictrum przewalskii*. Powder (MeOH) (as chloride). Mp 170° dec. (chloride). $[\alpha]_{\text{D}}^{25}$ -141.7 (c, 1.8 in MeOH). λ_{max} 260 (log ε 3.74); 286 (log ε 4.37) (MeOH).
 Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**, 165-171 (*isol, uv, ir, pmr, cmr, ms*)

Przewaline P-682


[205520-57-6]



- $\text{C}_{41}\text{H}_{42}\text{N}_2\text{O}_{10}$ 722.79
 Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl_3). Mp 180-181°. $[\alpha]_{\text{D}}^{20}$ +70.1 (c, 0.5 in CHCl_3). λ_{max} 203 (log ε 4.5); 220 (sh) (log ε 4.41); 258 (sh) (log ε 4.34); 280 (sh) (log ε 4.09); 300 (sh) (log ε 3.96); 315 (sh) (log ε 3.88); 340 (sh) (log ε 3.35) (MeOH).
 Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**, 165-171 (*isol, uv, ir, pmr, cmr, ms*)

Przewalskine P-683

[205520-59-8]



C₄₂H₄₈N₂O₁₀ 740.849

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl₃). Mp 71-75°. [α]_D²⁰ +98.1 (c, 0.1 in CHCl₃). λ_{max} 291 (log ε 4.5); 302 (log ε 4.42); 322 (sh) (log ε 4.37) (MeOH).

10'-Demethoxy, 13'-methoxy: Thalmelatidine

[31199-55-0]

C₄₂H₄₈N₂O₁₀ 740.849

Alkaloid from the roots of *Thalictrum minus* ssp. *elatum* (Ranunculaceae). Needles (EtOH). Mp 120-122°. [α]_D¹⁹ +47 (c, 1.0 in CHCl₃). Struct. revised in 1989. Isomeric 5-methoxy-6,7-methylenedioxy struct. originally proposed.

10'-Demethoxy, 14'-methoxy: Przewalskinine


[205520-58-7]

C₄₂H₄₈N₂O₁₀ 740.849

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl₃). Mp 88-90°. [α]_D²⁰ +60 (c, 0.3 in CHCl₃). λ_{max} 282 (log ε 4.56); 300 (log ε 4.48); 317 (sh) (log ε 4.43) (MeOH).

El Sheikh, M.O.A. *et al.*, *Phytochemistry*, 1989, **28**, 1967 (*Thalmelatidine*)

Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**, 165-171 (*Przewalskine, Przewalskinine*)

Przewalstine**P-684**C₄₂H₅₀N₂O₉ 726.865

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl₃). Mp 96-98°. [α]_D²⁰ +47.1 (c, 0.3 in CHCl₃). λ_{max} 281 (log ε 4.56); 302 (log ε 4.44); 315 (sh) (log ε 4.39) (MeOH).

5'-Hydroxy: Przewalstinine

[205520-62-3]

C₄₂H₅₀N₂O₁₀ 742.864

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl₃). Mp 107-111°. [α]_D²⁰ +44.4 (c, 0.4 in CHCl₃). λ_{max} 281 (log ε 4.41); 302 (log ε 4.31); 315 (sh) (log ε 4.26) (MeOH).

10'-Demethoxy, O¹²-de-Me: Thalibulamine

[105437-17-0]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D +63 (c, 0.2 in MeOH).

10'-Demethoxy, O⁷, O¹²-di-de-Me: Istanbulamine

[82953-24-0]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from the roots and rhizomes of *Thalictrum minus* var. *microphyllum*. [α]_D²⁵ +60 (c, 0.09 in MeOH).

10'-Demethoxy, 14'-methoxy: Przewalstinine

[205520-60-1]

C₄₂H₅₀N₂O₉ 726.865

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder. Mp 91-93°. [α]_D +36.4 (c, 0.3 in CHCl₃). [α]_D +65.5 (c, 0.1 in MeOH). λ_{max} 281 (log ε 4.56); 300 (log ε 4.51); 317 (log ε 4.45) (MeOH).

10'-Demethoxy, 14'-methoxy, 5'-hydroxy: Przewalstinidine

[205520-63-4]

C₄₂H₅₀N₂O₁₀ 742.864

Alkaloid from the roots of *Thalictrum przewalskii*. Yellow powder (CHCl₃). Mp 116-120°. [α]_D +31 (c, 0.2 in CHCl₃). [α]_D +61.3 (c, 0.9 in MeOH). λ_{max} 281 (log ε 4.5); 300 (log ε 4.42); 315 (sh) (log ε 4.35) (MeOH).

Guinaudeau, H. *et al.*, *Tet. Lett.*, 1982, **23**, 2523 (*Istanbulamine*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1986, **49**, 494 (*Thalibulamine*)

Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**, 165-171 (*Przewalstine, Przewalstinine, Przewalstinidine, Przewalstinidine, isol, uv, ir, pmr, cmr, ms*)

Psammaplina I**P-685**

[540466-77-1]

As Psammaplina B, P-687 with

R = -SO₂MeC₁₂H₁₅BrN₂O₅S 379.231

Alkaloid from the sponge *Pseudoceratina purpurea*.


Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)

Psammaplina A**P-686**

N,N'-Bis[3-(3-bromo-4-hydroxyphenyl)-2-oximidopropanoyl]cystamine

[110659-91-1]

[109152-27-4, 109152-30-9]

C₂₂H₂₄Br₂N₄O₆S₂ 664.395

Both the (*E,E*)- (Mp 172-174°) and (*E,Z*)-isomers have been *isol.* and characterised. See Park *et al.* (2003). Major metab. from a sponge tentatively identified as *Thorectopsamma xana*. Also from the sponge *Pseudoceratina purpurea* and *isol.* from another unidentified sponge. Inhibitor of chitinase, histone deacetylase and DNA methyltransferase. Antifungal agent. Exhibits farnesyl protein transferase and leucine aminopeptidase inhibitory activity. Shows cytotoxic and antimicrobial activity. Semicryst. solid. Mp 67-75°. λ_{max} 219 (ε 64600); 277 (ε 4875); 286 (sh); 290 (ε 10000) (MeOH) (*Derep.*)

4'-O-Sulfate: Psammaplina A₁. Psammaplina A 11'-sulfate

[263009-15-0]

[325155-70-2]

C₂₂H₂₄Br₂N₄O₉S₃ 744.459

Isol. from the sponge *Aplysinella rhax*. Oil; amorph. pale yellow solid (as *N,N*-dimethylguanidine salt). Mp 76-80° (*N,N*-dimethylguanidine salt). λ_{max} 206 (ε 33000); 283 (ε 3000); 324 (ε 750) (MeOH). λ_{max} 204 (log ε 4.73); 280 (log ε 3.88) (MeOH) (*N,N*-dimethylguanidine).

4',4''-Di-O-sulfate: Psammaplina A₂

[325155-74-6]

C₂₂H₂₄Br₂N₄O₁₂S₄ 824.523

Isol. from the sponge *Aplysinella rhax*. Amorph. pale yellow solid (as *N,N*-dimethylguanidine salt). Mp 155-160° (*N,N*-dimethylguanidine salt). λ_{max} 204 (log ε 4.6); 281 (log ε 3.98) (MeOH) (*N,N*-dimethylguanidine).

S-Oxide: Psammaplina J

[540466-78-2]

C₂₂H₂₄Br₂N₄O₇S₂ 680.394

Alkaloid from the sponge *Pseudoceratina purpurea*.

5'-Hydroxy: Psammaplina K

[439089-58-4]

C₂₂H₂₄Br₂N₄O₇S₂ 680.394

Isol. from the sponge *Aplysinella rhax*. Light pinkish oil. λ_{max} 290 (ε 3438) (MeOH).

5',5''-Dihydroxy: Psammaplina L

[439089-59-5]

C₂₂H₂₄Br₂N₄O₈S₂ 696.394

Isol. from the sponge *Aplysinella rhax*. Light pinkish oil. λ_{max} 291 (ε 3139) (MeOH).

5',5''-Dimer: Bisaprasin

[112514-43-9]

C₄₄H₄₆Br₄N₈O₁₂S₄ 1326.774

Minor metab. of a sponge tentatively identified as *Thorectopsamma xana*. Also *isol.* from *Pseudoceratina purpurea* and *Aplysinella rhax*. Inhibitor of DNA methyltransferase and histone deacetylase. Exhibits farnesyl protein transferase and leucine aminopeptidase inhibitory activity. Shows cytotoxic and antimicrobial activity. Foam. Sol. EtOH, butanol; poorly sol. H₂O, EtOAc. λ_{max} 204 (ε 77000); 284 (ε 16000) (MeOH) (*Berdy*).

5',5''-Dimer, 4'-O-sulfate: Bisaprasin 11'-sulfate

[263009-16-1]

C₄₄H₄₆Br₄N₈O₁₃S₅ 1406.838

Isol. from *Aplysinella rhax*. Oil. λ_{max} 203 (ε 83000); 283 (ε 7900); 324 (ε 5300) (MeOH).

5'-Bromo: 5-Bromopsammaplina A

[612089-85-7]

C₂₂H₂₃Br₃N₄O₆S₂ 743.291

Isol. from an association of the sponges *Jaspis wondoensis* and *Poecillastra wondoensis*. Amorph. solid. Possesses (*E,E*)-config.

Trisulfide: Psammaplina A trisulfide

[263009-15-0]

C₂₂H₂₄Br₂N₄O₆S₃ 696.461


Isol. from an association of the sponges *Jaspis wondoensis* and *Poecillastra wondoensis*. Amorph. solid. Possesses (*E,E*)-config.

Arabshahi, L. *et al.*, *J.O.C.*, 1987, **52**, 3584-3586 (*isol, struct*)

- Rodriguez, A.D. *et al.*, *Tet. Lett.*, 1987, **28**, 4989-4992 (*isol, uv, ir, pmr, cmr, activity*)
- Jimenez, C. *et al.*, *Tetrahedron*, 1991, **47**, 2097-2102 (*activity*)
- Hoshino, O. *et al.*, *Bioorg. Med. Chem. Lett.*, 1992, **2**, 1561-1562 (*synth*)
- Pham, N.B. *et al.*, *J. Nat. Prod.*, 2000, **63**, 393-395 (*11'-sulfates*)
- Shin, J. *et al.*, *Tetrahedron*, 2000, **56**, 9071-9077 (*Psammaplins A₁, A₂, activity*)
- Tabudravu, J.N. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 1123-1128 (*pmr, cmr, Psammaplins K, L*)
- Park, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1495-1498 (*isol, pmr, cmr*)
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)
- Godert, A.M. *et al.*, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 3330-3333 (*synth*)
- Mora, F.D. *et al.*, *J. Nat. Prod.*, 2006, **69**, 547-552 (*Psammaplin A, activity*)

Psammaplin B P-687

2-[[3-(3-Bromo-4-hydroxyphenyl)-2-(hydroxyimino)-1-oxopropyl]amino]ethyl thiocyanate, 9CI
[133991-67-0]



R = SCN

- C₁₂H₁₂BrN₃O₃S 358.215
Isol. from the marine sponge *Pseudoceratina purpurea*. Oil. λ_{max} 280 (ε 5450) (MeOH) (Berdy).
- Jimenez, C. *et al.*, *Tetrahedron*, 1991, **47**, 2097-2102 (*isol, pmr, cmr*)
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, activity*)

Psammaplin C P-688

- N-[2-(Aminosulfonyl)ethyl]-3-bromo-4-hydroxy-α-(hydroxyimino)benzenepropanamide, 9CI
[133991-68-1]
As Psammaplin B, P-687 with R = -SO₂NH₂
C₁₁H₁₄BrN₃O₅S 380.219
Isol. from the marine sponge *Pseudoceratina purpurea*. Oil. λ_{max} 282 (ε 2200) (MeOH) (Berdy).
- Jimenez, C. *et al.*, *Tetrahedron*, 1991, **47**, 2097-2102 (*isol, pmr, cmr*)
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, activity*)

Psammaplin D P-689

- [133991-69-2]
As Psammaplin B, P-687 with R = -SSCH₂CH₂NHCOOMe
C₁₅H₂₀BrN₃O₅S₂ 466.376
Isol. from the marine sponge *Pseudoceratina purpurea*. Exhibits antimicrobial and mild tyrosine kinase inhibitory activities. Oil. λ_{max} 280 (ε 17733) (MeOH) (Berdy).
- Jimenez, C. *et al.*, *Tetrahedron*, 1991, **47**, 2097-2102 (*isol, pmr, cmr*)
- Park, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1495-1498 (*isol, pmr, cmr*)

- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, cmr, activity*)

Psammaplin E P-690

- [540466-73-7]
As Psammaplin B, P-687 with R = -S-S-CH₂CH₂NHCOCONH₂
C₁₅H₁₉BrN₄O₅S₂ 479.375
Alkaloid from the sponge *Pseudoceratina purpurea*. λ_{max} 206 (ε 18070); 284 (ε 1860) (MeOH).
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)

Psammaplin F P-691

- [540466-74-8]
As Psammaplin B, P-687 with R = -S-S-CH₂CH₂NHCOCOOH
C₁₅H₁₈BrN₃O₅S₂ 480.36
Alkaloid from the sponge *Pseudoceratina purpurea*. Potent histone deacetylase inhibitor. λ_{max} 206 (ε 25935); 280 (ε 2060) (MeOH).
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)

Psammaplin G P-692


- [540466-75-9]
As Psammaplin B, P-687 with R = -S-S-CH₂CH₂NHC(COOH)=NNH₂
C₁₅H₂₀BrN₅O₅S₂ 494.39
Alkaloid from the sponge *Pseudoceratina purpurea*. Potent DNA methyltransferase inhibitor.
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)

Psammaplin H P-693

- [540466-76-0]
As Psammaplin B, P-687 with R = -S-S-CH₂CH₂NHCOOEt
C₁₆H₂₂BrN₃O₅S₂ 480.403
Alkaloid from the sponge *Pseudoceratina purpurea*.
- Piña, I.C. *et al.*, *J.O.C.*, 2003, **68**, 3866-3873 (*isol, pmr, cmr*)

Psammaplysene A P-694

- [850013-02-4]
C₂₇H₃₅Br₄N₃O₃ 769.208
Isol. from the sponge *Psammaplysin* sp. Specific inhibitor of FOXO1a nuclear export.

**N¹⁵-Me: Psammaplysene D**

- [1000582-60-4]
C₂₈H₃₇Br₄N₃O₃ 783.235
Isol. from a *Psammoclemma* sp. Amorph. solid. Isol. as a 4:3 mixt. of *E/Z*-isomers. λ_{max} 206 (log ε 4.83); 219 (sh) (log ε 4.59); 286 (log ε 4.37) (MeOH).


N¹-De-Me: Psammaplysene B

- [865722-85-6]
C₂₆H₃₃Br₄N₃O₃ 755.181
Isol. from *Psammaplysin* sp.

7-Debromo, N¹⁵-Me: Psammaplysene C

- [1000582-18-2]
C₂₈H₃₈Br₃N₃O₃ 704.339
Isol. from a *Psammoclemma* sp. Cytotoxic. Amorph. solid. Isol. as a 4:3 mixt. of *E/Z*-isomers. λ_{max} 206 (log ε 4.87); 218 (sh) (log ε 4.61); 294 (log ε 4.36); 312 (sh) (log ε 4.3) (MeOH).

- Schroeder, F.C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 574-576 (*isol, pmr, cmr, ms*)
- Georgiades, S.N. *et al.*, *Org. Lett.*, 2005, **7**, 4091-4094 (*synth*)
- Buchanan, M.S. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1827-1829 (*Psammaplysenes C, D*)

Psammaplysin A P-695

- C₂₁H₂₃Br₄N₃O₆ 733.045
Isol. from Red Sea sponge *Psammaplysin* sp. Shows anti-biotic props.; active against gram-positive bacteria. Exhibits moderate *in vitro* cytotoxicity towards the human colon tumour cell-line HCT116. Also shows antifouling activity. Foam. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁵ -65.2 (c, 0.52 in MeOH). λ_{max} 229 (ε 12700); 262 (sh) (ε 7100); 276 (ε 2500); 283 (sh) (ε 3200) (MeOH) (Derep). λ_{max} 228 (ε 9000); 256 (ε 4000); 272 (ε 2800); 276 (ε 2500) (MeOH) (Berdy). λ_{max} 229 (ε 12500); 276 (ε 7100); 283 (ε 3200) (MeOH/HCl) (Berdy). λ_{max} 229 (ε 12500) (EtOH) (Berdy).

Di-Ac: [95739-54-1]
Needles (MeOH). Mp 140°.

N-Formyl: Ceratinamide A. N-Formylpsammaplysin A

- [178214-21-6]
C₂₂H₂₃Br₄N₃O₇ 761.056
Metab. from the marine sponge *Pseudoceratina purpurea*. Shows potent antifouling activity. Solid. Sol. MeOH. [α]_D²⁴ -89.7 (c, 0.146 in MeOH). λ_{max} 207 (ε 64200) (MeOH) (Berdy).

N-(13-Methyltetradecanoyl): Ceratinamide B. 1'-Deoxypsammaplysin D

[178214-22-7]

C₃₆H₅₁Br₄N₃O₇ 957.431
 Metab. from the marine sponge
Pseudoceratina purpurea. Shows
 antifouling activity. Solid. Sol.
 MeOH. [α]_D²⁴ -53.5 (c, 0.263 in
 Me₂CO). λ_{max} 207 (ε 51100) (MeOH)
 (Berdy).

N-Me: Psammaplysin F

[190510-86-2]
 C₂₂H₂₅Br₄N₃O₆ 747.072
 Alkaloid from *Aplysinella* sp. [α]_D -62.3
 (c, 1.2 in MeOH/CH₂Cl₂).

I'-Hydroxy: Psammaplysin B

[85819-67-6]
 C₂₁H₂₃Br₄N₃O₇ 749.045
 Isol. from *Psammaplysinilla purpurea*.
 Active against gram-positive
 bacteria. Shows moderate *in vivo*
 cytotoxicity vs human colon tumour
 cells. Foam. Sol. MeOH, CHCl₃;
 poorly sol. H₂O, hexane. [α]_D²⁵ -60.2
 (c, 0.632 in MeOH). λ_{max} 218 (ε
 28600); 256 (sh) (ε 7000); 262 (sh) (ε
 6400); 277 (sh) (ε 4000); 282 (ε 3000)
 (MeOH) (Derep). λ_{max} 218 (ε
 28600); 282 (ε 3000) (MeOH/HCl)
 (Berdy).

I'-Hydroxy, N-Me: Psammaplysin C

[142449-78-3]
 C₂₂H₂₅Br₄N₃O₇ 763.071
 Metab. from the marine sponge
Druinella purpurea (*Psammaplysinilla*
purpurea). Exhibits moderate *in vitro*
 cytotoxicity towards the human
 colon tumour cell-line HCT116.
 Glass. Sol. MeOH, CHCl₃; poorly
 sol. H₂O. [α]_D²³ -57.1 (c, 0.014 in
 MeOH). λ_{max} 218 (ε 17600); 255 (ε
 6700); 279 (ε 3000) (MeOH)
 (Berdy).

I'-Hydroxy, N-(13-methyltetradecanoyl):


Psammaplysin D
 [149415-75-8]
 C₃₆H₅₁Br₄N₃O₈ 973.43
 Alkaloid from the sponge
Aplysinella sp. Immunosuppressant.
 Oil. [α]_D¹⁸ -71.4 (c, 2.8 in Me₂CO).
 λ_{max} 210 (ε 50900); 224 ; 298
 (MeOH) (Berdy).

Rotem, M. *et al.*, *Tetrahedron*, 1983, **39**, 667
 (isol, uv, ir, pmr, cmr, ms, struct)
 Roll, D.M. *et al.*, *J.A.C.S.*, 1985, **107**, 2916 (uv,
 ir, pmr, cmr, ms, struct)
 Copp, B.R. *et al.*, *J. Nat. Prod.*, 1992, **55**, 822
 (*Psammaplysin C*)
 Ichiba, T. *et al.*, *J.O.C.*, 1993, **58**, 4149
 (*Psammaplysin D*)
 Tsukamoto, S. *et al.*, *Tetrahedron*, 1996, **52**,
 8181 (*Ceratinamide A*, *Ceratinamide B*)
 Liu, S. *et al.*, *J. Nat. Prod.*, 1997, **60**, 614
 (*Psammaplysin F*)

Psammaplysin E

P-696

[149415-76-9]



C₂₇H₂₅Br₄N₃O₈ 839.126


Alkaloid from the sponge
Aplysinella sp. and *Pseudoceratina*
purpurea. Immunosuppressive,
 antifouling agent. Bright yellow oil.
 [α]_D¹⁸ -80.3 (c, 0.3 in Me₂CO). λ_{max} 208
 (ε 53400); 224 ; 298 (ε 25000) (MeOH)
 (Berdy).

Ichiba, T. *et al.*, *J.O.C.*, 1993, **58**, 4149 (isol, uv,
 ir, pmr, cmr, struct)

Psammopemmin A

P-697

[145940-63-2]



C₁₂H₉BrN₄O 305.133

Free base unstable, isol. as salt (prob.
 hydrochloride). Alkaloid from an Ant-
 arctic sponge *Psammopemina* sp. Yellow
 solid (as salt). λ_{max} 220 (ε 24000); 287 (ε
 4300); 362 (ε 9800) (EtOH) (salt) (Ber-
 dy).

6-Bromo: Psammopemmin C

[145940-65-4]
 C₁₂H₈Br₂N₄O 384.029
 From *Psammopemina* sp. Yellow solid
 (as salt). λ_{max} 228 (ε 33000); 292 (ε
 2800); 362 (ε 15400) (EtOH) (salt)
 (Berdy).

7-Bromo: Psammopemmin B


[145940-64-3]
 C₁₂H₈Br₂N₄O 384.029
 From *Psammopemina* sp. λ_{max} 227 (ε
 34500); 292 (ε 2800); 361 (ε 15200)
 (EtOH) (as salt) (Berdy).

Butler, M.S. *et al.*, *Aust. J. Chem.*, 1992, **45**,
 1871 (isol, uv, pmr, cmr, ms, struct)

Pseudoanchynazine A

P-698

[445430-63-7]




C₃₅H₃₈N₁₀O₈ 726.747
 Alkaloid from the sponge *Clathria*
 sp. Yellow powder. [α]_D²⁵ -60.7 (c,
 0.83 in MeOH). λ_{max} 248 (ε 9700)
 (MeOH).

Zuleta, I.A. *et al.*, *Tetrahedron*, 2002, **58**, 4481-
 4486 (isol, pmr, cmr)

Pseudoanchynazine B

P-699

[445430-64-8]




C₃₅H₃₈N₁₀O₈ 726.747
 Alkaloid from the sponge *Clathria* sp.
 Yellow powder. [α]_D²⁵ +81.2 (c, 0.42 in
 MeOH). λ_{max} 242 (ε 11300) (MeOH).

Zuleta, I.A. *et al.*, *Tetrahedron*, 2002, **58**, 4481-
 4486 (isol, pmr, cmr)

Pseudoanchynazine C

P-700

[445430-65-9]




C₃₅H₃₈N₁₀O₈ 726.747
 Alkaloid from the sponge *Clathria* sp.
 Yellow powder. [α]_D²⁵ +96.6 (c, 2.21 in
 MeOH). λ_{max} 253 (ε 10900) (MeOH).

Zuleta, I.A. *et al.*, *Tetrahedron*, 2002, **58**, 4481-
 4486 (isol, pmr, cmr)

Pseudoargiopinin III

P-701

[117233-46-2]




C₁₉H₂₇N₅O₃ 373.454
 Constit. of *Argiope lobata* venom.

Grishin, E.V. *et al.*, *Bioorg. Khim.*, 1988, **14**,
 883 (isol)
 Grishin, E.V. *et al.*, *Toxicol.*, 1989, **27**, 541 (isol)

Pseudobactin B

P-702

[138145-39-8]



R = -COCH₂CH₂⁵COOH

C₄₂H₅₉N₁₁O₁₇ 989.992

Pyoverdinin chromopeptin occurring as Fe complex. Des-ferri form shown. Shows delocalisation of charge over the two naphthopyrimidine Ns. Prod. by *Pseudomonas fluorescens*. Siderophore.

5'-Amide: Pseudobactin

[76975-04-7]

C₄₂H₆₀N₁₂O₁₆ 989.007

Prod. by *Pseudomonas fluorescens*. Siderophore. Plant growth promotor, showing some control of *Fusarium* sp. Antifungal. Fluorescent yellow-green cryst. (as Fe complex). Poorly sol. hexane. λ_{\max} 400 (ϵ 15000) (H₂O) (Berdy). λ_{\max} 307; 362; 376 (pH 4.4 buffer) (Berdy). λ_{\max} 401 (pH 7.8 buffer) (Berdy).

5,6-Dihydro, 5'-amide: Pseudobactin A

[79438-64-5]

C₄₂H₆₂N₁₂O₁₆ 991.023

Prod. by *Pseudomonas* sp. B₁₀. Siderophore, active against *Erwinia carotovora*. Siderophore, active against *Erwinia carotovora*. λ_{\max} 226; 405 (pH 7.8 buffer) (Berdy). λ_{\max} 220; 360; 380 (pH 4.4 buffer) (Berdy).

7-Sulfo, 5,6-dihydro, 5'-amide: 7-Sulfopseudobactin A

[218433-00-2]

C₄₂H₆₂N₁₂O₁₉S 1071.087

Prod. by *Pseudomonas* sp. 267. Siderophore.

Analogue (R = COCH₂CH₂CH₂NH-COOH), 5,6-dihydro: [138145-41-2]

C₄₃H₆₄N₁₂O₁₇ 1021.049

Prod. by *Pseudomonas fluorescens*.

Analogue (R = COCH₂CH₂CH₂NH-COOH), 7-sulfo, 5,6-dihydro: [218432-99-6]

C₄₃H₆₄N₁₂O₂₀S 1101.113

Prod. by *Pseudomonas fluorescens*. Siderophore.

Teintze, M. *et al.*, *Biochemistry*, 1981, **20**, 6446; 6457 (*isol, cryst struct*)

Kolasa, T. *et al.*, *J.O.C.*, 1990, **55**, 4246 (*synth, bibl*)

Taraz, K. *et al.*, *Z. Naturforsch., C*, 1991, **46**, 522-526 (*Pseudobactin B*)

Andriollo, N. *et al.*, *J. Agric. Food Chem.*, 1992, **40**, 1245-1248 (*Pseudobactin B*)

Okonya, J.F. *et al.*, *J.O.C.*, 1995, **60**, 1932-1935 (*synth, ir, pmr, cmr*)


Buzikiewicz, H. *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 125-130 (*7-sulfo derivs*)

Pseudobactin 7SR1

P-703

Pyoverdinin cyclo-Pf P19

[90295-72-0]

C₄₆H₆₃N₁₃O₂₃ 1166.077

Struct. revised in 2000. Prod. by *Pseudomonas* sp. 7SR1 and *Pseudomonas fluorescens* P19. Siderophore. Sol. H₂O; poorly sol. Me₂CO, hexane. λ_{\max} 460 (pH 5 buffer) (Berdy). λ_{\max} 385; 401 (ϵ 21000) (H₂O) (Berdy).

3'-Deoxy: [280142-33-8]

C₄₆H₆₃N₁₃O₂₂ 1150.078

Prod. by *Pseudomonas fluorescens* P19.

Yang, C.C. *et al.*, *Biochemistry*, 1984, **23**, 3534-3540 (*isol, struct*)

Vossen, W. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 153-164 (*isol, struct*)

Pseudobaleabuxine F

P-704

Pseudobaleabuxine, 8CI

[31768-83-9]

C₃₀H₅₀N₂O₂ 470.737


Struct. unknown. Alkaloid from the leaves of *Buxus balearica* (Buxaceae). Mp 236-240°. [α]_D +120.7 (CHCl₃). Not identical with N³-Isobutyrylcyclobuxine F, in C-920. Suggested to be epimeric with it.

Kurakina, I.O. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 406-409; 1970, **6**, 231-235; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 337-339; 1970, **6**, 225-227 (*isol, ms, pmr*)

Pseudoberberine

P-705

[19716-72-4]

C₂₀H₁₈NO₄⁺ 336.367


Quaternary alkaloid from the leaves of *Isopyrum thalictroides* (Ranunculaceae). Pale yellow needles (as chloride). Mp 300° dec. (chloride).

Moullis, C. *et al.*, *Phytochemistry*, 1977, **16**, 1283 (*isol, uv, ms, struct*)

Pseudoceratinine

P-706

[174545-78-9]

C₁₇H₂₁Br₄N₅O₂ 647.001

Metab. from the marine sponge *Pseudoceratina purpurea*. Antifouling agent; larval settlement and development inhibitor. Solid. λ_{\max} 275 (ϵ 18400) (MeOH) (Berdy).

Tsukamoto, S. *et al.*, *Tet. Lett.*, 1996, **37**, 1439 (*isol, uv, ir, pmr, cmr, struct*)


Behrens, C. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 321 (*synth*)

Ponasiak, J.A. *et al.*, *Tetrahedron*, 1998, **54**, 6977-6986 (*synth*)

Pseudoceratin A

P-707

[929624-97-5]

C₃₅H₃₆Br₄N₆O₁₄ 1084.317

Isol. from *Pseudoceratina purpurea*. Antifungal agent. Amorph. solid. [α]_D²⁸ +11.7 (c, 0.1 in MeOH). λ_{\max} 207 (log ϵ 5.64) (MeOH).

21-Epimer: Pseudoceratin B

[929624-98-6]

C₃₅H₃₆Br₄N₆O₁₄ 1084.317


Isol. from *Pseudoceratina purpurea*. Antifungal agent. Amorph. solid. [α]_D²⁸ -11.6 (c, 0.1 in MeOH). λ_{\max} 207 (log ϵ 5.64) (MeOH).

Jang, J.-H. *et al.*, *J.O.C.*, 2007, **72**, 1211-1217 (*isol, pmr, cmr*)

Pseudoceratinine B

P-708

[172670-15-4]

C₁₈H₂₆Br₂N₅O₂⁺ 504.244


Alkaloid from the sponge *Pseudoceratina verrucosa*. Sol. MeOH, CH₂Cl₂; poorly sol. H₂O. [α]_D +17 (c, 1 in MeOH) (as hydrochloride). λ_{\max} 285 (ϵ 1480); 330 (ϵ 490) (MeOH) (Berdy).

Benharref, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 177 (*isol, uv, ir, pmr, cmr, cd, struct*)

Pseudoceratinine C

P-709

[172617-97-9]

C₂₇H₃₁Br₄N₇O₆ 869.201


Alkaloid from the sponge *Pseudoceratina verrucosa*. Sol. MeOH, CH₂Cl₂; poorly sol. H₂O. [α]_D -10 (c, 1 in

MeOH) (as hydrochloride). Closely related to Purealine, P-816. λ_{\max} 220 (ϵ 34670); 228 (ϵ 12880) (MeOH) (Berdy).

Benharref, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 177 (isol, uv, ir, cd, pmr, cmr, struct)

Pseudocoptisine

[19716-67-7]



$C_{19}H_{14}NO_4^{\oplus}$ 320.324

Quaternary alkaloid from the leaves of *Isopyrum thalictroides* (Ranunculaceae). Yellow cryst. (as chloride). Mp 300° (dec.)(chloride).

8-Hydroxy: 8-Hydroxypseudocoptisine

$C_{19}H_{14}NO_5^{\oplus}$ 336.323

Quaternary alkaloid from the roots of *Thalictrum przewalskii*. Powder (MeOH) (as chloride). Mp 270° (chloride). λ_{\max} 237 (log ϵ 3.97); 265 (log ϵ 4.08); 287 (log ϵ 4.24); 312 (log ϵ 4.06); 341 (log ϵ 3.9); 380 (sh) (log ϵ 3.53) (MeOH).


Moullis, C. *et al.*, *Phytochemistry*, 1977, **16**, 1283 (isol, uv, pmr, ms, struct)

Zhang, G.-L. *et al.*, *Planta Med.*, 1998, **64**, 165-171 (8-Hydroxypseudocoptisine)

Pseudodeoxyviolacein

P-711

3-[1,5-Dihydro-4-(1H-indol-3-yl)-5-oxo-2H-pyrrol-2-ylidene]-1,3-dihydro-2H-indol-2-one, 9CI
[155210-54-1]



$C_{20}H_{13}N_3O_2$ 327.342

Prod. by *Chromobacterium violaceum*. Orange-red powder. Mp >280° dec. λ_{\max} 270 (ϵ 9200); 285 (ϵ 9100); 383 (sh) (ϵ 6500); 483 (ϵ 15100) (MeOH).

5-Hydroxy: Pseudoviolacein

$C_{20}H_{13}N_3O_3$ 343.341

Prod. by *Chromobacterium violaceum*. Red cryst. (MeOH). Mp >240°. λ_{\max} 275 (ϵ 9100); 290 (ϵ 9500); 394 (ϵ 8500); 413 (ϵ 8900); 472 (ϵ 11100) (MeOH).


Hoshino, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 1994, **58**, 279-282 (Pseudodeoxyviolacein)

Hoshino, T. *et al.*, *J.C.S. Perkin 1*, 1995, 1565-1571 (Pseudoviolacein)

Pseudodistomin B

P-712

5-Amino-2-(6,8-tridecadienyl)-4-piperidinol, 9CI
[106293-83-8]



$C_{18}H_{34}N_2O$ 294.479

Struct. revised in 1992. Originally assigned as the 3',5'-diene. Metab. from the Okinawan tunicate *Pseudodistoma kanoko* and *Pseudodistoma megalarva*. Exhibits antineoplastic activity; phosphodiesterase inhibitor; calmodulin antagonist. Oil (as *N,N'*-di-Ac). Sol. $CHCl_3$, MeOH; poorly sol. H_2O , toluene. $[\alpha]_D^{24}$ +35 (c, 1 in MeOH) (di-Ac). λ_{\max} 233 (ϵ 17500) (MeOH) (Derep).

8'Z-Isomer: Pseudodistomin A

[106231-30-5]

Metab. from *Pseudodistoma kanoko*. Exhibits antineoplastic activity; phosphodiesterase inhibitor; calmodulin antagonist. Oil (as *N,N'*-di-Ac). Sol. $CHCl_3$, MeOH; poorly sol. H_2O , toluene. $[\alpha]_D^{24}$ +36 (c, 1 in MeOH) (di-Ac). λ_{\max} 233 (ϵ 17500) (MeOH) (Derep).

4-Epimer: Pseudodistomin D

[196099-42-0]

$C_{18}H_{34}N_2O$ 294.479

Isol. from *Pseudodistoma megalarva*. Gum. $[\alpha]_D^{25}$ +5 (c, 0.3 in MeOH). λ_{\max} 240 (MeOH).

4,5-Diepimer: Pseudodistomin E

[196099-44-2]

$C_{18}H_{34}N_2O$ 294.479

Isol. from *Pseudodistoma megalarva*. Gum. $[\alpha]_D^{25}$ -20.8 (c, 0.4 in MeOH). λ_{\max} 239 (MeOH).

Ishibashi, M. *et al.*, *J.O.C.*, 1987, **52**, 450-453 (Pseudodistomin A,B, isol, uv, ir, pmr, cmr, ms)

Kiguchi, T. *et al.*, *Tet. Lett.*, 1992, **33**, 7389-7390 (synth, struct)

Ishibashi, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 804-806 (struct)

Naito, T. *et al.*, *J.C.S. Perkin 1*, 1996, 281-288 (synth, struct, bibl)

Kiguchi, T. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1212-1215 (synth)

Freyer, A.J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 986-990 (Pseudodistomin D,E, isol, uv, ir, pmr, cmr)


Ninomiya, I. *et al.*, *Alkaloids (Academic Press)*, 1998, **50**, 317-342 (rev)

Trost, B.M. *et al.*, *Org. Lett.*, 2005, **7**, 823-826 (synth)

Pseudodistomin F

P-713

5-Amino-2-(1,3,8,10-pentadecatetraenyl)-4-piperidinol
[196099-46-4]



$C_{20}H_{34}N_2O$ 318.501

Isol. from the ascidian *Pseudodistoma megalarva*. Gum. $[\alpha]_D^{25}$ -13.9 (c, 0.4 in MeOH). λ_{\max} 233 (MeOH).

4,5-Diepimer: Pseudodistomin C

[169626-36-2]

$C_{20}H_{34}N_2O$ 318.501

Isol. from *Pseudodistoma kanoko* and *Pseudodistoma megalarva*. Cytotoxic agent. Oil (as *N^1,N^5,O*-tri-Ac). $[\alpha]_D^{22}$ +85 (c, 1 in $CHCl_3$) (tri-Ac). λ_{\max} 235 (ϵ 37000) (MeOH) (tri-Ac).

Kobayashi, J. *et al.*, *J.O.C.*, 1995, **60**, 6941-6945 (Pseudodistomin C)

Doi, Y. *et al.*, *Tetrahedron*, 1996, **52**, 4573-4580 (Pseudodistomin C, synth)

Freyer, A.J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 986-990 (Pseudodistomin F)


Langlois, N. *et al.*, *Org. Lett.*, 2002, **4**, 185-187 (Pseudodistomin C, synth)

Tanaka, K. *et al.*, *Heterocycles*, 2006, **68**, 183-192 (Pseudodistomin C, synth)

Pseudoepiisolaecarpiline

P-714

13,16-Dihydroisolaecarpine, 9CI
[28230-67-3]



$C_{16}H_{21}NO_2$ 259.347


Alkaloid from the leaves of *Elaeocarpus sphaericus* (Elaeocarpaceae). Gum. Mp 230-235° (as picrate). $[\alpha]_D$ +222 (c, 0.04 in $CHCl_3$).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1971, **24**, 1679 (isol, uv, ir, pmr, ms, struct, abs config)

Pseudolaudanine

P-715

1-[(3,4-Dimethoxyphenyl)methyl]-1,2,3,4-tetrahydro-7-methoxy-2-methyl-6-isoquinolinol, 9CI. 1-(3,4-Dimethoxybenzyl)-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline
[85-65-4]



(S)-form

$C_{20}H_{25}NO_4$ 343.422

(S)-form [13452-22-7]

Alkaloid from *Roemeria refracta* (Papaveraceae). Cryst. (MeOH/Et₂O). Mp 117-119°. $[\alpha]_D$ +73 (c, 0.4 in MeOH).

N-Me: Pseudorine. N-Methylpseudolaudanine

[87332-68-1]

$C_{21}H_{28}NO_4^{\oplus}$ 358.456

Alkaloid from the bark of *Fagara mayu* (Rutaceae) and from roots of *Papaver pseudo-orientale* (Papaveraceae). Prisms (EtOH) (as iodide). Mp 204-

206° (iodide). $[\alpha]_D^{20} +107$ (c, 0.5 in MeOH). Abs. config. not detd. It is possible that Pseudorine and *N*-Methylpseudolaudanine could have opposite abs. configs. Phys. props. refer to Pseudorine iodide.

Me ether: see Laudanosine, L-69

(±)-*form* [5876-49-3]

Cryst. (CHCl₃/petrol). Mp 110°.

Picrate:

Yellow cryst. (EtOH). Mp 162°.

Me ether: see Laudanosine, L-69

[105660-68-2, 13240-65-8]

Chatterjee, A. *et al.*, *Chem. Ber.*, 1966, **99**, 1764-1766 (*synth*)

Cassels, B.K. *et al.*, *Chem. Ind. (London)*, 1966, 1635-1636 (*resoln*)

Miller, L.L. *et al.*, *J.A.C.S.*, 1973, **95**, 2651-2656 (*synth*)

Assem, E.M. *et al.*, *Planta Med.*, 1983, **48**, 77 (*N-Methylpseudolaudanine*)


Věžník, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1986, **57**, 1752-1763 (*Pseudorine*)

Gözler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 666-668 (*isol, pmr, struct*)

Pseudolycorine

P-716

[29429-03-6]



C₁₆H₁₉NO₄ 289.33

Alkaloid from *Lycoris radiata* and several other spp. in the Amaryllidaceae. Antineoplastic agent active against Rauscher viral leukaemia. Mp 247-248° (hydrate) Mp 240-242° (anhyd.). $[\alpha]_D -62$ (c, 1.2 in EtOH). Log P -1.07 (uncertain value) (calc).

▶ VS3200000

Hydrochloride: Mp 266°.

1-O-β-D-Glucopyranoside: **Pseudolycorine-1-glucoside**

[93710-69-1]

C₂₂H₂₉NO₉ 451.472

Alkaloid from the roots of *Pancreatium biflorum*. Also detected in a number of *Crinum* spp. (Amaryllidaceae). Amorph. solid + 1H₂O. $[\alpha]_D^{20} -105.5$ (c, 0.3 in MeOH).

O¹, O²-Di-O-β-D-glucopyranoside: **Pseudolycorine 1,2-diglucoside**

[132472-47-0]

C₂₈H₃₉NO₁₄ 613.614

Alkaloid from *Hymenocallis caymanensis* (Amaryllidaceae). Mp 265°. $[\alpha]_{H_2}^{22} +68$ (c, 0.2 in CHCl₃).

1-Ac: **1-O-Acetylpsudolycorine**

[104386-94-9]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the aerial parts of *Narcissus requienii* (Amaryllidaceae). Mp 248-250°.

2-Ac: **2-O-Acetylpsudolycorine**

[104386-93-8]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the aerial parts of *Narcissus requienii* (Amaryllidaceae). Cryst. + ½ H₂O. Mp 168-170°.

O,O,O-Tri-Ac: Mp 205°.

O¹-Hexadecanoyl, O²-(1-O-hexadecanoyl-2-O-octadecanoyl)glycerophosphoryl: **1-O-Palmitoyl-2-O-(1-O-palmitoyl-2-O-stearoyl)glycerophosphorylpsudolycorine**

[128701-14-4]

C₆₉H₁₂₀NO₁₂P 1186.68

From fruits of *Crinum asiaticum*. Light brown amorph. solid. Obt. as a mixt. containing some octadecenoyl (oleoyl) residues in place of stearoyl.

O²-Phosphatidyl: **Phosphatidylpsudolycorine**

[109269-86-5]

[109269-88-7]

Alkaloid from the flowers of *Zephyranthes flava* (Amaryllidaceae). Light brown amorph. solid. Mixt. contg. stearoyl and oleoyl alkyl residues.

O²-Me: **Carinatine**

[64937-89-9]

C₁₇H₂₁NO₄ 303.357

Alkaloid from the bulbs of *Zephyranthes carinata* (Amaryllidaceae). Amorph. $[\alpha]_D^{27} -68.97$ (c, 0.696 in CHCl₃).

O²-Me, picrate: Mp 195-197° dec.

O⁹-De-Me, O¹⁰-Me: **Golceptine**

C₁₆H₁₉NO₄ 289.33

Alkaloid from *Narcissus jonquilla* (Amaryllidaceae). Cryst. (Me₂CO). Mp 146-148°. $[\alpha]_D^{24} -156$ (c, 0.2 in CHCl₃). Probable struct. Formulated as *O², O⁹*- or *O², O¹⁰*-didemethylgalanthine. The 2,10-isomer would correspond to Pseudolycorine, P-716 to which the props. clearly do not correspond.

O⁹-De-Me, O¹⁰-Me, 1-Ac: **Sternbergine**

[96253-76-8]

C₁₈H₂₁NO₅ 331.368

Alkaloid from the bulbs of *Sternbergia lutea* and *Brunsvigia radulosa*. Cytotoxic on P388 and BL6 melanoma. Antimalarial agent. Needles (CHCl₃). Mp 105-112° Mp 197-202°. Not correlated with Golceptine.

O¹⁰-Me: **Methylpsudolycorine**

[11042-05-0]

C₁₇H₂₁NO₄ 303.357

Alkaloid from *Narcissus pseudonarcissus* and several other spp. in the Amaryllidaceae. Mp 228-233°. $[\alpha]_D -40$ (DMF).

O¹, O¹⁰-Di-Me, stereoisomer: **Zaidine**

[63125-82-6]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Hymenocallis arenicola* (Amaryllidaceae). Prisms (Me₂CO). Mp 192-193°. $[\alpha]_D^{24} -4$ (c, 0.2 in CHCl₃). 1- And 2-configs. unknown.

O², O¹⁰-Di-Me: **Galanthine**

[517-78-2]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Galanthus voronovii* and a large number of other spp. in the

Amaryllidaceae. Hypotensive agent.

Mp 132-134° (hydrate) Mp 162-164° (anhyd.). $[\alpha]_D^{22} -85$ (c, 0.23 in CHCl₃).

O², O¹⁰-Di-Me, hydrochloride: Mp 198-199°.

O², O¹⁰-Di-Me, perchlorate: Mp 218° dec. (199-201°).

O², O¹⁰-Di-Me, 3,3αα-epoxide: **Incartine**

[145237-00-9]

C₁₈H₂₃NO₅ 333.383

Alkaloid from *Galanthus elwesii* and *Lycoris incarnata*. A biosynthetic intermediate from Galanthine to Narcissidine, N-43. Mp 183-185°.

O², O¹⁰-Di-Me, O⁹-de-Me: **Goleptine**

C₁₇H₂₁NO₄ 303.357

Alkaloid from the *Narcissus jonquilla* hybrid "Golden Sceptre" (Amaryllidaceae). Fine needles (Me₂CO). Mp 141°. $[\alpha]_D^{24} -99$ (c, 0.2 in CHCl₃). Probable struct. Shown to be *O⁹*- or *O¹⁰*-demethylgalanthine; clearly not the same as Carinatine.

O², O¹⁰-Di-Me, O⁹-de-Me, perchlorate:

Cryst. (MeOH). Mp 211° dec.

Boit, H.G. *et al.*, *Chem. Ber.*, 1956, **89**, 163 (*Galanthine, isol*)

Fales, H.M. *et al.*, *J.A.C.S.*, 1956, **78**, 4145; 1958, **80**, 4395 (*derivs, struct*)

Uyeo, S. *et al.*, *J.C.S.*, 1959, 172 (*struct*)

Döpke, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1964, **297**, 39 (*Goleptine, Golceptine*)

Kinstle, T.H. *et al.*, *Tet. Lett.*, 1966, 4659 (*Pseudolycorine, Galanthine, ms*)

Harken, R.D. *et al.*, *J.O.C.*, 1976, **41**, 2450 (*biosynth*)

Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1977, **25**, 2244 (*Carinatine*)

Döpke, W. *et al.*, *Z. Chem.*, 1977, **17**, 101 (*Zaidine*)

Evidente, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 1003 (*Sternbergine*)

Ghosal, S. *et al.*, *Phytochemistry*, 1984, **23**, 1167 (*glucoside*)

Llabrés, J.M. *et al.*, *Phytochemistry*, 1986, **25**, 1453 (*acetates*)

Ghosal, S. *et al.*, *Phytochemistry*, 1987, **26**, 823 (*Phosphatidylpsudolycorine*)

Ghosal, S. *et al.*, *Phytochemistry*, 1990, **29**, 805 (*glycerophosphorylpsudolycorine*)

Doepke, W. *et al.*, *Z. Chem.*, 1990, **30**, 375 (*1,2-diglucoside*)

Kihara, M. *et al.*, *Heterocycles*, 1992, **34**, 1299 (*Incartine*)


Seaforth, C.E. *et al.*, *Fitoterapia*, 1998, **69**, 79 (*Pseudolycorine, pmr, cmr*)

Campbell, W.E. *et al.*, *Phytochemistry*, 2000, **53**, 587 (*Sternbergine, isol, activity*)

Berkov, S. *et al.*, *Molecules*, 2007, **12**, 1430-1435 (*Incartine, pmr, cmr*)

Pseudomanibacanine

P-717



C₁₉H₂₁NO₃ 311.38

(*R*)-*form* [151851-39-7]

Alkaloid from stem bark of *Aniba*

canellilla (Lauraceae). $[\alpha]_D +182$ (c, 0.22 in MeOH).

(S)-form**O²-De-Me: Pseudoanibacanine**

[151851-38-6]


C₁₈H₁₉NO₃ 297.353

Alkaloid from stem bark of *Aniba canellilla* (Lauraceae). $[\alpha]_D -31$ (c, 2 in MeOH).

Oger, J.-M. *et al.*, *Can. J. Chem.*, 1993, **71**, 1128 (isol, uv, pmr, cmr, ms, struct)

Pseudomonine**P-718**

[172923-94-3]

C₁₆H₁₈N₄O₄ 330.343

Alkaloid from cultures of *Pseudomonas fluorescens* AH2 isol. from spoiled Nile perch from Lake Victoria. Siderophore. Amorph. brownish solid. Sol. H₂O. $[\alpha]_D -80$ (c, 0.041 in H₂O). λ_{max} 203 (ε 25118); 237 (ε 8912); 298 (ε 2291) (H₂O) (Berdy).


Anthoni, U. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1786 (isol, uv, pmr, cmr, struct)

Pseudopalmatine**P-719**

5,6-Dihydro-2,3,10,11-tetramethoxydibenzo[a,g]quinolininium(1+), 9CI

[19716-66-6]

[30045-17-1]

C₂₁H₂₂NO₄[⊕] 352.409

Alkaloid from *Stephania suberosa*, *Enantia chlorantha* and *Xylophia vieillardii*. Yellow needles (CHCl₃) (as chloride). Mp 212-213° (207-208°) (chloride). λ_{max} 265 (log ε 4.16); 288 (log ε 4.48); 310 (sh) (log ε 4.34); 339 (log ε 4.1); 379 (log ε 3.66) (CHCl₃) (as chloride).

O²-De-Me: Pseudocolumbamine

[64191-04-4]

C₂₀H₂₀NO₄[⊕] 338.382

Quaternary alkaloid from the leaves of *Isopyrum thalictroides* (Ranunculaceae). Yellow needles + 2H₂O (as chloride). Mp 300° dec. (chloride). λ_{max} 241 (log ε 4.42); 266 (log ε 4.46); 289 (log ε 4.79); 315 (log ε 4.52); 342 (log ε 4.35) (EtOH) (as chloride).

O³-De-Me: Dehydrodiscretine. Pseudojatrotrrhizine

[68947-61-5]

[75491-93-9]

C₂₀H₂₀NO₄[⊕] 338.382

Quaternary alkaloid from *Thalictrum fauriei*, *Fibraurea chloroleuca* and *Heptacyclum zenkeri* (preferred genus name *Penianthus*) (Ranunculaceae, Menispermaceae). Shows strong antiplasmodial activity. Fine reddish-brown needles (MeOH) (as chloride); orange needles (MeOH) (as iodide). Mp 230-234° (chloride) Mp 243-245° dec. (iodide). Identity of Pseudojatrotrrhizine with

Dehydrodiscretine is based on CA information only. λ_{max} 242 (log ε 4.24); 265 (log ε 4.13); 289 (log ε 4.32); 315 (sh) (log ε 4.15); 342 (log ε 3.97); 381 (log ε 3.97) (MeOH) (iodide). λ_{max} 254 (log ε 4.46); 309 (log ε 4.24); 379 (log ε 4.31) (MeOH/OH) (iodide).

O¹¹-De-Me: DehydrocorytenchineC₂₀H₁₉NO₄ 337.374

Alkaloid from *Xylophia vieillardii* (Annonaceae). Cryst. (MeOH). Mp 256-258°. Exists as zwitterion with negative charge on O-10. λ_{max} 208 (log ε 3.91); 233 (log ε 3.91); 290 (log ε 4.27); 318 (log ε 4.22); 344 (sh) (log ε 3.81); 383 (log ε 3.76) (MeOH). λ_{max} 240 (log ε 3.94); 266 (sh) (log ε 3.93); 288 (log ε 4.29); 313 (log ε 4.13); 340 (sh) (log ε 3.85); 376 (log ε 3.48) (MeOH/HCl).

O², O¹¹-Di-de-Me: DehydrocoreximineC₁₉H₁₈NO₄[⊕] 324.356

Quaternary alkaloid from the roots of *Xylophia parviflora*. Pale yellow crust. (as perchlorate). Mp 243-247° (perchlorate). λ_{max} 222 (log ε 4.23); 292 (log ε 4.54); 320 (log ε 4.48); 380 (log ε 4.11) (MeOH).

O³, O¹¹-Di-de-Me: 3,11-Dihydroxy-2,10-dimethoxy-5,6-dihydroprotoberberineC₁₉H₁₇NO₄ 323.348

Alkaloid from the stems and leaves of *Miliusa cuneata*. Yellow powder (MeOH). Mp 217-219°. Zwitterionic. λ_{max} 208 ; 239 (sh) ; 293 ; 323 ; 384 (MeOH).

Kametani, T. *et al.*, *Tetrahedron*, 1973, **29**, 73-76 (synth, pmr)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1974, 1712-1714 (synth)

Moulis, C. *et al.*, *Phytochemistry*, 1977, **16**, 1283-1287 (Pseudocolumbamine)

Siwon, J. *et al.*, *Pharm. Weekbl.*, 1978, **113**, 1153-1156 (Pseudocolumbamine)

Chen, C.H. *et al.*, *CA*, 1983, **99**, 212774z (Dehydrodiscretine, synth)

Duah, F.K. *et al.*, *Phytochemistry*, 1983, **22**, 321-322 (Dehydrodiscretine)

Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 547-549 (isol, uv)

Jossang, A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 466-472 (Dehydrocorytenchine)

Wright, C.W. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1638-1640 (activity)


Chen, B. *et al.*, *Nat. Prod. Res.*, 2003, **17**, 397-402 (3,11-Dihydroxy-2,10-dimethoxydihydroprotoberberine)

Nishiyama, Y. *et al.*, *Phytochemistry*, 2004, **65**, 939-944 (Dehydrocoreximine)

Grycová, L. *et al.*, *Phytochemistry*, 2007, **68**, 150-175 (rev)

Pseudophrynamine 258A**P-720**

Pseudophrynaminol. Pseudophryne Alkaloid 258A. PS 258A

C₁₆H₂₂N₂O 258.363**(+)-form**

Synthetic. Cryst. (MeOH). Mp 146°. $[\alpha]_D +87.8$ (c, 0.72 in CHCl₃) (+82.1).

(-)-form [113056-55-6]

Alkaloid from skin extracts of the Australian frog *Pseudophryne coriacea* (Myobatrachidae). Also present in *Pseudophryne guentheri*, *Pseudophryne occidentalis*, *Pseudophryne australis*, *Pseudophryne semimarmorata*, *Pseudophryne bibronii* and *Pseudophryne corroboree*. $[\alpha]_D^{29} -82.8$ (c, 0.98 in CHCl₃). λ_{max} 242 ; 295 (MeOH) (Berdy). λ_{max} 210 (ε 11000); 243 (ε 4800); 297 (ε 1800) (EtOH) (Berdy).

N⁸-Me: Pseudophrynamine 272A. N⁸-Methylpseudophrynaminol.

Pseudophryne Alkaloid 272A. PS 272A [128855-20-9]

C₁₇H₂₄N₂O 272.389

Trace constit. in one population of *Pseudophryne coriacea*. Tentative struct.

4'-Aldehyde: Pseudophrynamine 256.

Pseudophryne Alkaloid 256. PS 256 [128855-19-6]

C₁₆H₂₀N₂O 256.347

Trace alkaloid from skin extracts of the Australian frog *Pseudophryne coriacea*.

4'-Aldehyde, N⁸-Me: Pseudophrynamine 270.

Pseudophryne Alkaloid 270. PS 270

[433711-65-0]

C₁₇H₂₂N₂O 270.374

Alkaloid from skin extracts of *Pseudophryne semimarmorata*. Tentative struct.

4'-Carboxylic acid, Me ester: Pseudophrynamine 286A.

Pseudophryne Alkaloid 286A. PS 286A [113056-57-8]

C₁₇H₂₂N₂O₂ 286.373

Trace alkaloid from skin extracts of the Australian frog *Pseudophryne coriacea* (Myobatrachidae), also present in *Pseudophryne corroboree*, *Pseudophryne semimarmorata* and *Pseudophryne bibronii*. Probably an artifact.

Deoxy: Pseudophrynamine 242.

Pseudophryne Alkaloid 242. PS 242 [433711-61-6]

C₁₆H₂₂N₂ 242.363

Alkaloid from skin extracts of *Pseu-*

dophryne semimarmorata.

5-Hydroxy, 4'-carboxylic acid, Me ester: **Pseudophrynamine 302**. Pseudophryne Alkaloid 302. PS 302

[128646-02-6]

C₁₇H₂₂N₂O₃ 302.372

Alkaloid from skin extracts of *Pseudophryne australis*, *Pseudophryne coriacea*, *Pseudophryne semimarmorata* and *Pseudophryne corroboree*.

5-Methoxy, 4'-carboxylic acid, Me ester: **Pseudophrynamine 316**. Pseudophryne Alkaloid 316. PS 316

[128646-01-5]

C₁₈H₂₄N₂O₃ 316.399

Minor or trace alkaloid in skin extracts of *Pseudophryne coriacea*, *Pseudophryne australis* and *Pseudophryne occidentalis*.

6-Methoxy, 5-hydroxy, 4'-carboxylic acid, Me ester: **Pseudophrynamine 332**.

Pseudophryne Alkaloid 332. PS 332

[128646-03-7]

C₁₈H₂₄N₂O₄ 332.399

Minor or trace alkaloid from *Pseudophryne coriacea*, *Pseudophryne australis*, *Pseudophryne occidentalis* and one population of *Pseudophryne corroboree*.

5,6-Dimethoxy, 4'-carboxylic acid, Me ester: **Pseudophrynamine 346A**. Pseudophryne Alkaloid 346A. PS 346A

[128895-97-6]

C₁₉H₂₆N₂O₄ 346.425

Minor/trace alkaloid in skin of *Pseudophryne australis* and *Pseudophryne coriacea*.

Δ²-Isomer: **Pseudophrynamine 258B**.

Pseudophryne Alkaloid 258B. PS 258B

[433711-63-8]

C₁₆H₂₂N₂O 258.363

Alkaloid from skin extracts of *Pseudophryne semimarmorata*. Tentative struct.

Δ²-Isomer, 4'-carboxylic acid, Me ester: **Pseudophrynamine 286B**.

Pseudophryne Alkaloid 286B. PS 286B

[128946-40-7]

C₁₇H₂₂N₂O₂ 286.373

Trace alkaloid in *Pseudophryne coriacea* and *Pseudophryne semimarmorata*.

Deoxy, 6-bromo: see Flustramine E, F-108

Spande, T.F. *et al.*, *J.O.C.*, 1988, **53**, 1222-1226 (*isol, uv, ir, pmr, cmr, ms, cd, struct, PS 286A*)

Daly, J.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 407-421 (*occur, derivs, Alkaloid 256*)

Mitchell, M.O. *et al.*, *Tet. Lett.*, 1990, **31**, 2681-2684 (*synth*)

Sun, W.Y. *et al.*, *Synlett*, 1993, 337-338 (*synth*)

Fuji, K. *et al.*, *Synlett*, 1995, 367-368 (*synth*)

Crich, D. *et al.*, *Tetrahedron*, 1995, **51**, 6379-6384 (*synth, cd, abs config*)

Smith, B.P. *et al.*, *J. Nat. Prod.*, 2002, **65**, 439-447 (*biosynth, occur*)


Kawasaki, T. *et al.*, *Tet. Lett.*, 2003, **44**, 1591-1593 (*synth*)

Daly, J.W. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1556-1575 (*rev*)

Pseudophrynamine 512A P-721

Pseudophryne Alkaloid 512. Pseudophrynamine A. PS 512

[113056-56-7]



C₃₂H₄₀N₄O₂ 512.694

Alkaloid from skin extracts of the Australian frog *Pseudophryne coriacea* (Myobatrachidae).

Didehydro, N⁸-Me: **Pseudophrynamine 524**. Pseudophryne Alkaloid 524. PS 524

[129726-96-1]

C₃₃H₄₀N₄O₂ 524.705

Trace alkaloid in skin extracts of *Pseudophryne coriacea*. Tentative struct.

1''ξ-Hydroxy: **Pseudophrynamine 528**.

Pseudophryne Alkaloid 528. PS 528

[128855-18-5]

C₃₂H₄₀N₄O₃ 528.693

Minor or trace alkaloid in skin extracts of *Pseudophryne coriacea* and *Pseudophryne corroboree*. Minor constit. in *Pseudophryne semimarmorata*.

Spande, T.F. *et al.*, *J.O.C.*, 1988, **53**, 1222-1226 (*isol, ir, pmr, cmr, ms, struct*)


Daly, J.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 407-421 (*derivs*)

Smith, B.P. *et al.*, *J. Nat. Prod.*, 2002, **65**, 439-447 (*occur, biosynth*)

Pseudophrynamine 272B P-722

Pseudophryne Alkaloid 272B. PS 272B

[128646-04-8]



C₁₆H₂₀N₂O₂ 272.346

Tentative struct. Trace constit. in two populations of *Pseudophryne coriacea* and one population each of *Pseudophryne corroboree*, *Pseudophryne occidentalis* and *Pseudophryne semimarmorata*.

4'-Carboxylic acid, Me ester: **Pseudophrynamine 300**. Pseudophryne Alkaloid 300. PS 300

[128855-15-2]

C₁₇H₂₀N₂O₃ 300.357

Minor alkaloid from skin extracts of *Pseudophryne australis* and *Pseudophryne semimarmorata*.

6-Methoxy, 4'-carboxylic acid, Me ester: **Pseudophrynamine 330**. Pseudophryne Alkaloid 330. PS 330

[128855-16-3]

C₁₈H₂₂N₂O₄ 330.383

Minor alkaloid in skin extracts of the Australian frogs *Pseudophryne coriacea* and *Pseudophryne semimarmorata*.

Trace constit. in *Pseudophryne australis*, *Pseudophryne corroboree* and *Pseudophryne occidentalis*. Tentative struct. More than one isomer (*E,Z*-?) detected.


Daly, J.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 407-421 (*isol, ms*)

Smith, B.P. *et al.*, *J. Nat. Prod.*, 2002, **65**, 439-447 (*occur, biosynth*)

Pseudoptoxine P-723

5,7,8,15-Tetrahydro-6-methylbis[1,3]benzodioxolo[5,6-c,5',6'-g]azecin-14(6H)-one, 9CI

[24240-05-9]



C₂₀H₁₉NO₅ 353.374

Alkaloid from the bark of *Zanthoxylum conspersipunctatum* and the root and stem bark of *Fagara vitiensis* (Rutaceae). Cryst. (MeOH). Mp 205° (200-202°).

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2233 (*isol, uv, pmr, ms, struct*)


Fish, F. *et al.*, *Phytochemistry*, 1972, **11**, 1528 (*isol, uv*)

Sotelo, R.M. *et al.*, *Aust. J. Chem.*, 1972, **25**, 385 (*synth, uv*)

Orito, K. *et al.*, *Heterocycles*, 1980, **14**, 11 (*synth*)

Pseudopteroxazole P-724

[242150-01-2]



Absolute Configuration

C₂₁H₂₇NO 309.45

Struct. revised in 2001. Isol. from *Pseudopteroxgorgia elisabethae*. Active against *Mycobacterium tuberculosis*. Yellowish oil. [α]_D²⁵ +101 (c, 1 in CHCl₃). λ_{max} 220 (ε 15400); 250 (ε 5000); 284 (ε 2000) (MeOH).

Rodriguez, A.D. *et al.*, *Org. Lett.*, 1999, **1**, 527-530 (*isol*)

Johnson, T.W. *et al.*, *J.A.C.S.*, 2001, **123**, 4475-4479 (*struct*)


Davidson, J.P. *et al.*, *J.A.C.S.*, 2003, **125**, 13486-13489 (*synth, abs config*)

Harmata, M. *et al.*, *Org. Lett.*, 2005, **7**, 3581-3583 (*synth*)

Pseudorepanduline

P-725

[57821-67-7]

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from *Daphnandra dielsii* and from an unnamed *Daphnandra* sp. (Monimiaceae). Mp 168-173° dec. [α]_D¹⁸ +229 (CHCl₃).


Bick, I.R.C. *et al.*, *Tet. Lett.*, 1975, 2219-2220 (*ir, pmr, ms, struct*)

Koike, L. *et al.*, *J. Nat. Prod.*, 1992, **55**, 455-460 (*pmr, cmr, abs config*)

Pseudostrychnine

P-726

16-Hydroxystrychnidin-10-one, 9CI. 3-Hydroxystrychnine. ψ -Strychnine [465-62-3]



Absolute Configuration

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from *Strychnos nux-vomica*, *Strychnos wallichiana*, *Strychnos ligustrina* and bark of *Strychnos gaultheriana* (Loganiaceae). Cryst. (EtOAc), powder (CHCl₃). Mp 266-268° (248-251°) dec. (dimorph.). [α]_D -85.9 (CHCl₃). [α]_D -43.8 (EtOH). Is formed as an artifact when CHCl₃ or CH₂Cl₂ solns. of Strychnine are left to stand. Pseudostrychnine is prob. a genuine alkaloid, however. λ_{\max} 255 (log ϵ 4.13); 279 (sh) (log ϵ 3.7); 280 (log ϵ 3.58) (EtOH).

▶ UL7450000

Hydrochloride:

Cryst. + 2H₂O (H₂O). [α]_D¹⁹ +3.9 (H₂O).

Perchlorate:

Needles (H₂O). Mp 300°.

Me ether: 3-Methoxystrychnine. 16-Methoxystrychnine

[5096-72-0]

C₂₂H₂₄N₂O₃ 364.443

Alkaloid from *Strychnos icaja* leaves, prob. as methylation artifact (Loganiaceae). Feathery needles (MeOH). Mp 195-200°. [α]_D -66 (c, 0.67 in CHCl₃).

▶ WL2420000

Et ether: 16-Ethoxystrychnine

Alkaloid from *Strychnos ignatii* as ethylation artifact.

12-Hydroxy: 3,12-Dihydroxystrychnine.

12,16-Dihydroxystrychnine

[75326-94-2]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from leaves of *Strychnos nux-vomica* (Sri Lankan, poss. hybrid) (Loganiaceae). [α]_D²¹ +55 (c, 0.21 in MeOH). λ_{\max} 260 ; 280 (sh) (CHCl₃).

10-Methoxy: Pseudo- β -colubrine. 3-Hydroxy- β -colubrine

[29079-34-3]

C₂₂H₂₄N₂O₄ 380.443

Minor alkaloid from *Strychnos nux-vomica* seeds and *Strychnos ligustrina* bark (Loganiaceae). Cryst. (EtOAc/hexane). Mp 140-143°. [α]_D²⁰ -56 (c, 1 in CHCl₃). λ_{\max} 263 (log ϵ 4.06); 296 (log ϵ 3.56); 310 (sh) (90% EtOH).

11-Methoxy: Pseudo- α -colubrine. 3-Hydroxy- α -colubrine

[34479-58-8]

C₂₂H₂₄N₂O₄ 380.443

Minor alkaloid from seeds of *Strychnos nux-vomica* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 225-227°. [α]_D²⁰ -54 (c, 1 in CHCl₃). λ_{\max} 257 (sh) (log ϵ 3.94); 293 (log ϵ 3.68); 300 (log ϵ 3.68) (90% EtOH).

11-Methoxy, 12-hydroxy: 3,12-Dihydroxy-11-methoxystrychnine

[83905-58-2]

C₂₂H₂₄N₂O₅ 396.442

Alkaloid from leaves of *Strychnos nux-vomica* (Sri Lankan, poss. hybrid) (Loganiaceae). Prisms (EtOAc). [α]_D²¹ +38 (c, 0.25 in MeOH). λ_{\max} 240 ; 270 (MeOH).

10,11-Dimethoxy: Pseudobrucine

[560-30-5]

C₂₃H₂₆N₂O₅ 410.469

Alkaloid from *Strychnos wallichiana* leaves, from *Strychnos nux-vomica*, *Strychnos gaultheriana* and *Strychnos ligustrina* (Loganiaceae). Cryst. (EtOAc). Mp 259-260°. [α]_D -70 (-55) (CHCl₃). λ_{\max} 265 (log ϵ 4.21); 299 (sh) (log ϵ 6.03) (no solvent reported) (unlikely extinction value).

Warnat, K. *et al.*, *Helv. Chim. Acta*, 1931, **14**, 997-1007 (*isol*)

Bailey, A.S. *et al.*, *J.C.S.*, 1948, 703-705 (*synth*)

Hendrickson, J.B. *et al.*, *Alkaloids (Academic Press)*, 1960, **6**, 179-217 (*rev, bibl*)

Boit, H.G. *et al.*, *Naturwissenschaften*, 1960, **47**, 136 (*Pseudobrucine, isol*)

Marini-Bettolo, G.B. *et al.*, *J. Assoc. Off. Anal. Chem.*, 1968, **51**, 185-191 (*isol, props, Pseudostrychnine, Pseudobrucine*)

delle Monache, F. *et al.*, *J. Chromatogr.*, 1968, **32**, 178-179 (*isol*)

Galeffi, C. *et al.*, *Ann. Chim. (Rome)*, 1970, **60**, 444-453 (*Pseudocolubrines*)

Bisset, N.G. *et al.*, *Tetrahedron*, 1973, **29**, 4137-4148 (*Strychnos icaja alkaloids*)

Bisset, N.G. *et al.*, *Phytochemistry*, 1974, **13**, 259-263 (*isol*)

Verpoorte, R. *et al.*, *Org. Magn. Reson.*, 1977, **9**, 567-571 (*cmr*)

Baser, K.H.C. *et al.*, *Phytochemistry*, 1982, **21**, 1423-1429 (*3,12-Dihydroxystrychnine, 3,12-Dihydroxy-11-methoxystrychnine*)


Asai, F. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 690-694 (*Strychnos ligustrina alkaloids*)

Philippe, G. *et al.*, *Phytochemistry*, 2003, **62**, 623-629 (*isol, pmr*)

Pseudotheonamide A₁

P-727

[224577-31-5]

C₃₆H₄₅N₉O₈ 731.807

The pseudotheonamides were initially named as Pseudocolubrines. Isol. from the sponge *Theonella swinhoei*. Serine protease inhibitor. Amorph. solid. [α]_D²⁹ -28 (c, 0.08 in MeOH). λ_{\max} 227 (sh) (ϵ 14000); 243 (sh) (ϵ 4500); 278 (sh) (ϵ 2000) (MeOH).

2-Epimer: Pseudotheonamide A₂

[224577-33-7]

C₃₆H₄₅N₉O₈ 731.807


Isol. from *Theonella swinhoei*. Serine protease inhibitor. Amorph. solid. [α]_D²⁹ -34 (c, 0.06 in MeOH). λ_{\max} 227 (sh) (ϵ 17000); 243 (sh) (ϵ 7100); 278 (sh) (ϵ 2300) (MeOH).

Nakao, Y. *et al.*, *J.A.C.S.*, 1999, **121**, 2425-2431

Pseudotheonamide B₂

P-728

[224577-34-8]

C₃₇H₄₅N₉O₈ 743.818


Isol. from the sponge *Theonella swinhoei*. Serine protease inhibitor. Amorph. solid. [α]_D²⁹ -17 (c, 0.05 in MeOH). Pseudotheonamide B not isolated. λ_{\max} 226 (sh) (ϵ 15000); 243 (sh) (ϵ 4500); 278 (sh) (ϵ 1800) (MeOH).

Nakao, Y. *et al.*, *J.A.C.S.*, 1999, **121**, 2425-2431

Pseudothiobinupharidine P-729C₃₀H₄₀N₂O₂S 492.724

Nuphar alkaloid. Struct. unknown. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Prisms + 2H₂O (as diperchlorate). Mp 173-175° (diperchlorate). [α]_D +186.2 (EtOH) (diperchlorate).

Achmatowicz, O. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1962, **36**, 1815-1825 (*isol, ir*)

Pseudotrienoic acid A P-730

Absolute Configuration

C₃₁H₅₀N₂O₆ 546.746

Prod. by *Pseudomonas* sp. MF381-IODS. Antibacterial agent. Oil.

Pohanka, A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1380-1385 (*isol, pmr, cmr*)

Pseudotrienoic acid B P-731

As Pseudotrienoic acid A, P-730 with R = H

C₂₉H₄₆N₂O₆ 518.692


Prod. by *Pseudomonas* sp. MF381-IODS. Antibacterial agent. Oil. [α]_D²⁰ +1.9 (c, 0.4 in MeOH).

Pohanka, A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1380-1385 (*isol, pmr, cmr*)

Amans, D. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 5870-5874 (*synth*)

Pseudovinblastinediol P-732

Pseudovincalaukoblastine diol [58511-80-1]

C₄₄H₅₆N₄O₈ 768.948

Proposed struct. The alternative, carbimolamine struct. with the hydroxyl group located at C-3' instead of at C-15' cannot be conclusively excluded. Alkaloid from *Catharanthus roseus* (Apocynaceae).


Tafur, S. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 1953-1957 (*isol, pmr, ms, struct*)

Pseudovincadiformine P-733

Methyl 2,3-didehydro-7-ethyl-20,21-dinoraspido-permidine-3-carboxylate, 9CI. ψ-Vincadiformine

[26251-92-3]

[56815-95-3 (+)-form, 73837-57-7 (±)-form]



Absolute Configuration

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Pandaca caducifolia* and *Melodinus polyadenus* (Apocynaceae). Noncryst. [α]_D +430 (MeOH).

15,20-Didehydro: Pseudotabersonine.

Δ^{6,7}-ψ-Vincadiformine. ψ-Catharanthine. ψ-Tabersonine. Pseudocatharanthine

[54244-12-1]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from *Pandaca caducifolia* (Apocynaceae). Noncryst. [α]_D +320 (MeOH).

20-Hydroxy: 20-Epipandoline

[56698-80-7]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Pandaca caducifolia*, *Ervatamia lifuana* and *Ervatamia obtusiuscula* (Apocynaceae). Noncryst. [α]_D +462 (MeOH).

18,19ξ-Dihydroxy: 18,19-Dihydroxy-ψ-vincadiformineC₂₁H₂₆N₂O₄ 370.447

Alkaloid from the stem bark of *Tabernaemontana albiflora* (Apocynaceae). Cryst. (MeOH). Mp 190°. [α]_D²⁰ +264 (c, 0.8 in CHCl₃).

19ξ,20-Dihydroxy: 19,20-Dihydroxy-ψ-vincadiformine. 19-Hydroxy-20-epipandolineC₂₁H₂₆N₂O₄ 370.447

Alkaloid from stem bark of *Tabernaemontana albiflora* (Apocynaceae). Cryst. (MeOH). Mp 204°. [α]_D²⁰ +511 (c, 1 in CHCl₃).

20-Epimer, 20-hydroxy: Pandoline

[53507-78-1]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Pandaca caducifolia* and other *Pandaca* spp., *Ervatamia orientalis*, *Ervatamia lifuana* and *Melodinus polyadenus* (Apocynaceae). Noncryst. [α]_D +417 (MeOH).

Kutney, J.P. *et al.*, *J.A.C.S.*, 1970, **92**, 1708

(*Pseudovincadiformine, Pseudotabersonine, synth, uv, ir, pmr*)

Hoizey, M.-J. *et al.*, *Phytochemistry*, 1974, **13**, 1995 (*Pandoline, isol, ms, uv, ir, pmr*)

Hoizey, M.-J. *et al.*, *Tet. Lett.*, 1974, 1601 (*Pandoline, struct*)

Zeche, M. *et al.*, *Phytochemistry*, 1975, **14**, 1122

(*Pseudotabersonine, Pandoline, Epipandoline, isol, uv, ir, pmr, ms, struct*)

LeMen, J. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1976, **283**, 759 (*Pandoline, struct, abs config*)

Bruneton, J. *et al.*, *Tet. Lett.*, 1976, 3567

(*Pandoline, pmr, cmr*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1980, **45**, 3259; 1982, **47**, 1335 (*Pandoline, Pseudovincadiformine, synth, uv, ir, pmr, ms*)

Kan, C. *et al.*, *Planta Med.*, 1981, **41**, 195

(*19,20-Dihydroxy-ψ-vincadiformine*)


Bornmann, W.G. *et al.*, *J.O.C.*, 1992, **57**, 1752 (*synth*)

Carroll, W.A. *et al.*, *J.A.C.S.*, 1993, **115**, 1164 (*Pseudotabersonine, synth*)

Kalaus, G. *et al.*, *J.O.C.*, 1993, **58**, 6076 (*synth*)

Pseudovobparicine P-734

[99023-48-0]

C₃₉H₄₄N₄O₂ 600.802

Alkaloid from the root bark of *Tabernaemontana divaricata* (Apocynaceae).

Van Beek, T.A. *et al.*, *Planta Med.*, 1985, **277-279** (*uv, pmr, ms, cd, struct*)


Pseudowithanine P-735

Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Mp 155-156° dec.

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f

Pseudoxandrine P-736

[104818-08-8]

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from the bark of *Pseudoxandra* aff. *lucida* (Annonaceae). Amorph. [α]_D +23 (c, 1.13 in CHCl₃). [α]_D +60 (c, 0.1 in MeOH).

7-Me ether: Pseudoxandrinine

[104778-23-6]


C₃₈H₄₀N₂O₇ 636.743

Alkaloid from *Pseudoxandra* aff. *lucida* (Annonaceae). Amorph. [α]_D +7 (c, 0.7 in CHCl₃).

Cortes, D. *et al.*, *Can. J. Chem.*, 1986, **64**, 1390 (*isol, uv, ir, pmr, cmr, ms, cd, struct, deriv*)

Pseudozoanthoxanthin A P-737

4-Methyl-1H-cyclohepta[1,2-d:3,4-d']diimidazole-2,8-diamine, 9CI. 2,8-Diamino-4-methyl-1H-cyclohepta[1,2-d:3,4-d']diimidazole [60049-47-0]



C₁₀H₁₀N₆ 214.229

Tautomeric with 9H-form. Note that when the 1, 8 or 9-posn. is substd., CA numbering is reversed and derivs. become 6-methyl instead of 4-methyl. Isol. from *Parazoanthus axinellae* and other Zoanthidae. Cryst. (EtOH). Mp 280° dec.

1-Me: 3-Norpseudozoanthoxanthin

[94080-60-1]

C₁₁H₁₂N₆ 228.256

From *Parazoanthus gracilis* and *Epizoanthus arenaceus*. Yellow needles (MeOH). Mp 300°.

9-Me: Norpseudozoanthoxanthin

[55827-12-8]

C₁₁H₁₂N₆ 228.256

Pigment from *Epizoanthus arenaceus*. Yellow prisms (MeOH). Darkens >230°.

N⁸,N⁸-Di-Me: N⁸,N⁸-Dimethylpseudozoanthoxanthin A

[94080-62-3]

C₁₂H₁₄N₆ 242.283

From *Parazoanthus gracilis*. Yellow needles (MeOH). Mp 300°.

1,N⁸-Di-Me: 1,N⁸-Dimethylpseudozoanthoxanthin A

[71337-50-3]

C₁₂H₁₄N₆ 242.283

Isol. from the coral *Gerardia* sp. Amorph. yellow powder. Dec. at ca. 180°.

N²,N⁸-Di-Me: N²,N⁸-Dimethylpseudozoanthoxanthin A. Norparagraine

[94080-61-2]

[71827-21-9 (tautomer)]

C₁₂H₁₄N₆ 242.283

From *Parazoanthus gracilis* and *Epizoanthus arenaceus*. Yellow needles (MeOH/CH₂Cl₂). Mp 190-192° dec.

7,9-Di-Me: Pseudozoanthoxanthin

[55827-11-7]

C₁₂H₁₄N₆ 242.283

Pigment from *Parazoanthus axinellae*, *Parazoanthus gracilis*, *Epizoanthus arenaceus*, *Zoanthus sociatus* and *Palythoa mammilosa*. Yellow prisms (MeOH). Mp 310°. Imino-tautomer on C-2. λ_{max} 290 (ε 75900); 335 (ε 5890); 348 (ε 6030); 399 (ε 21400) (MeOH/HCl) (Derep). λ_{max} 281 (ε 30200); 307 (ε 46800); 367 (ε 11700); 421 (ε 8510) (MeOH) (Derep).

1,N⁸,N⁸-Tri-Me: 1,N⁸,N⁸-Trimethylpseudozoanthoxanthin A

[68395-65-3]

C₁₃H₁₆N₆ 256.31

From *Gerardia* sp., *Palythoa* sp. and *Parazoanthus* sp. Cryst. (EtOH aq.) or amorph. yellow powder. Dec. at ca. 200°. Imino-tautomer on C-2.

N²,N⁸,N⁸-Tri-Me: Paragraine

[57695-32-6]

C₁₃H₁₆N₆ 256.31

Isol. from *Parazoanthus gracilis* together with *Dentitheca habererii* (the former is parasitic on the latter). Also

from *Epizoanthus* sp. Shows papaverine-like activity and antihistamine activity. Selective Na channel blocker in squid axon membranes. Yellow needles (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 258-262° dec. Strong yellow-green fluor. in soln. λ_{max} 230 (sh) (ε 15000); 252 (ε 13000); 305 (sh) (ε 47900); 316 (ε 62200); 373 (ε 15000); 409 (ε 14400); 428 (sh) (ε 3800) (EtOH) (Derep). λ_{max} 252 (ε 11900); 314 (ε 52000); 372 (ε 12600); 409 (ε 13100) (MeOH) (Berdy). λ_{max} 245 (ε 13500); 301 (ε 54800); 363 (ε 6900); 410 (ε 21300) (MeOH/HCl) (Berdy).

N²,N⁸,N⁸-Tri-Me, dihydrobromide:

Yellow needles + 3H₂O. Mp 280-282° dec.

7,9,N²-Tri-Me: Homopseudozoanthoxanthin

[71827-22-0]

C₁₃H₁₆N₆ 256.31

Trace pigment from *Palythoa* sp. and *Zoanthus sociatus*. Yellow prisms (MeOH). Mp 256-259°. Imino-tautomer on C-2. λ_{max} 226 (ε 10500); 261 (ε 10500); 306 (ε 38900); 317 (sh) (ε 30900); 377 (ε 10000); 416 (ε 15100) (MeOH) (Derep).

1,N²,N⁸,N⁸-Tetra-Me: 9-Methylparagraine

[81904-42-9]

C₁₄H₁₈N₆ 270.336

Isol. from *Parazoanthus gracilis*. Yellow prisms. Mp 269-271° dec. λ_{max} 251 (ε 1875); 309 (ε 8250); 369 (ε 1625); 409 (ε 2125) (MeOH).

7,9,N²,N²-Tetra-Me: Dimethylpseudozoanthoxanthin

[55827-18-4]

C₁₄H₁₈N₆ 270.336

Pigment from *Palythoa* sp. Yellow needles (MeOH). Mp 310°. Imino-tautomer on C-2.

1,N²,N²,N⁸,N⁸-Penta-Me:

N²,N²,N⁸,N⁸,1,4-Hexamethyl-1H-cyclohepta[1,2-d:3,4-d']diimidazole-2,8-diamine, 9CI

[71337-51-4]

C₁₅H₂₀N₆ 284.363

Minor constit. of *Gerardia* sp. Oily yellow solid. Mp 115° dec. approx.

Cariello, L. et al., *Tetrahedron*, 1974, **30**, 4191-4196 (*Epizoanthus arenaceus* constits)

Komoda, Y. et al., *Chem. Pharm. Bull.*, 1975, **23**, 2464 (uv, pmr, cryst struct)

Braun, M. et al., *J.A.C.S.*, 1978, **100**, 4208-4213 (synth)

Schwartz, R.E. et al., *Tet. Lett.*, 1978, 2235-2238 (*1,N⁸,N⁸-tri-Me, isol, cryst struct*)

Schwartz, R.E. et al., *Can. J. Chem.*, 1979, **57**, 1707-1711 (*Gerardia* constits, isol, cryst struct)


Cariello, L. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1979, **63**, 77-82 (*Zoanthus* constits, *Palythoa* constits)

Komoda, Y. et al., *Chem. Pharm. Bull.*, 1982, **30**, 502; 1984, **32**, 3873-3879 (*Parazoanthus gracilis* constits, uv, pmr, cmr, ms)

Jiménez, C. et al., *J. Nat. Prod.*, 1993, **56**, 9-14 (*9-Methylparagraine, isol*)

Pseurotins

P-738



Pseurotin A, R = -CH^Z=CHCH₂CH₃

Pseurotin B, R = -CH^Z=CHCH(OH)CH₃

Pseurotin C, R = -CH^E=CHCH(OH)CH₃

Pseurotin E, R = -CH^E=CHCOCH₃

Pseurotin A [58523-30-1]

C₂₂H₂₅NO₈ 431.441

Prod. by *Pseudeurotium ovalis*. Also from terrestrial and marine-derived *Aspergillus fumigatus*. Monoamine oxidase and chitin synthase inhibitor, apomorphine antagonist, nerve growth factor. Rhombic cryst. (CH₂Cl₂/hexane). Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 162-163.5°. [α]_D²⁵ -5 (MeOH). λ_{max} 202 (ε 23400); 251 (ε 14700); 284 (ε 10300) (EtOH) (Derep). λ_{max} 253 (ε 16600); 279 (ε 11220) (EtOH) (Berdy).

Tri-O-Ac: Mp 126-128°. [α]_D²⁰ -23 (c, 0.23 in MeOH).

11-Me ether: 11-O-Methylpseurotin A

[956904-34-0]

C₂₃H₂₇NO₈ 445.468

Prod. by a marine-derived *Aspergillus fumigatus*. Pale yellow solid. [α]_D²⁸ -1.8 (c, 0.2 in MeOH).

O-De-Me: Pseurotin F₂. De-O-methylpseurotin A

[149883-42-1]

C₂₁H₂₃NO₈ 417.415

Prod. by *Aspergillus fumigatus*. Oil.

8-Epimer, O-de-Me: Pseurotin F₁

[149792-62-1]

C₂₁H₂₃NO₈ 417.415

Prod. by *Aspergillus fumigatus*.

Pseurotin B [77409-67-7]

C₂₂H₂₅NO₉ 447.441

Prod. by *Pseudeurotium ovalis*. Cryst. (THF/Et₂O). Mp 204-206°.

Pseurotin C [77449-67-3]

C₂₂H₂₅NO₉ 447.441

Prod. by *Pseudeurotium ovalis*. Monoamine oxidase inhibitor. Foam. Rather unstable.

Pseurotin D [77409-68-8]

C₂₂H₂₅NO₈ 431.441

Prod. by *Pseudeurotium ovalis* and *Aspergillus fumigatus*. Neuroleptic agent, apomorphine antagonist. Unstable yellowish oil. Sol. CH₂Cl₂, EtOH, CHCl₃, 2-propanol, EtOAc. Allylically rearranged isomer of Pseurotin A with -CH(OH)CH=CHCH(OH)CH₂CH₃ side-chain.

Pseurotin E [77409-69-9]

C₂₂H₂₃NO₉ 445.425
 Prod. by *Pseudeurotium ovalis*. Oil.
 Rather unstable.
 Bloch, P. *et al.*, *Helv. Chim. Acta*, 1981, **64**,
 304-315 (*isol, pmr, cmr*)
 Breitenstein, W. *et al.*, *Helv. Chim. Acta*, 1981,
64, 379-388 (*isol, struct*)
 Mohr, P. *et al.*, *Tetrahedron, Suppl.*, No. 9,
 1981, **37**, 201-212 (*cmr, biosynth*)
 Wenke, J. *et al.*, *Biosci., Biotechnol., Biochem.*,
 1993, **57**, 961-964 (O-Demethylpseurotin A,
Pseurotin A)
Eur. Pat., 1993, 546 474; *CA*, **119**, 137528y
 (*Pseurotins F₁, F₂*)
 Hayashi, Y. *et al.*, *Org. Lett.*, 2003, **5**, 2287-
 2280 (*synth*)
 Aoki, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 2004, **77**,
 1703-1716 (*synth*)
 Boot, C.M. *et al.*, *J. Nat. Prod.*, 2007, **70**,
 1672-1675 (*marine, isol*)

Psilocauline


P-739

C₁₅H₃₁N₃O 269.429
 Struct. unknown. MF not certain.
 Alkaloid from *Psilocaulon ab simile*. Mp
 162°.
 Rimington, C. *et al.*, *S. Afr. J. Sci.*, 1933, **30**,
 503

Psycholeine

P-740

[144424-79-3]




Absolute
 Configuration

C₄₄H₅₀N₈ 690.933
 Alkaloid from *Psychotria oleoides* (Ru-
 biaceae). [α]_D²⁰ -150 (c, 0.4 in EtOH).
 λ_{max} 240 (ε 18600); 310 (ε 5300)
 (EtOH).

Guéritte-Voegelein, F. *et al.*, *J. Nat. Prod.*,
 1992, **55**, 923-930 (*isol, uv, ir, pmr, cmr, ms*,
cd, struct)
 Lebsack, A.D. *et al.*, *J.A.C.S.*, 2002, **124**, 9008-
 9009 (*synth*)

Psychopentamine

P-741



C₅₅H₆₂N₁₀ 863.162


Alkaloid from the leaves of *Psychotria
 rostrata*. Amorph. powder. [α]_D¹⁸ +42 (c,
 0.2 in CHCl₃). λ_{max} 210 (log ε 4.81);
 250 (log ε 4.37); 304 (log ε 4.1)
 (MeOH).

Takayama, H. *et al.*, *Org. Lett.*, 2004, **6**, 2945-
 2948 (*isol, cd, pmr, cmr, ms*)

Psychotridine

P-742

[52617-25-1]




C₅₅H₆₂N₁₀ 863.162
 Struct. revised in 1999. Alkaloid from
Psychotria beccaroides and *Calycoden-
 dron milneri* (Rubiaceae). Platelet ag-
 gregation inhibitor; analgesic, sedative.
 Sol. MeOH, CHCl₃; poorly sol. H₂O.
 Mp 180-181°. [α]_D -38 (c, 0.9 in
 CHCl₃).

Stereoisomer: Alkaloid E†
 C₅₅H₆₂N₁₀ 863.162
 Alkaloid from leaves of
Psychotria forsteriana. Exhibits
 potent cytotoxicity. [α]_D²⁰ +153
 (c, 0.35 in EtOH). [α]_D²⁰ +49
 (c, 0.2 in CHCl₃).
 Hart, N.K. *et al.*, *Aust. J. Chem.*, 1974, **27**,
 639-646 (*isol, uv, ir, pmr, ms*)
 Roth, A. *et al.*, *Planta Med.*, 1986, 450
 (*Alkaloid E*)
 Jannic, V. *et al.*, *J. Nat. Prod.*, 1999, **62**, 838-
 843 (*cd, struct*)

Psychotridine C

P-743


[129651-30-5]



C₅₅H₆₂N₁₀ 863.162
 Alkaloid from *Calycodendron milnei*.
 Adjibade, Y. *et al.*, *Planta Med.*, 1990, **56**, 212-
 215
 Saad, H.-E.A. *et al.*, *Planta Med.*, 1995, **61**,
 313-316

Psychotrimine

P-744




C₃₃H₃₈N₆ 518.703
 Alkaloid from the leaves of *Psychotria
 rostrata*. Amorph. powder. [α]_D¹⁸ +179 (c,
 0.2 in CHCl₃). λ_{max} 206 (log ε 4.55); 222
 (log ε 4.57); 296 (log ε 3.99) (MeOH).

Takayama, H. *et al.*, *Org. Lett.*, 2004, **6**, 2945-
 2948 (*isol, cd, pmr, cmr, ms*)
 Newhouse, T. *et al.*, *J.A.C.S.*, 2008, **130**,
 10886-10887 (*synth*)
 Matsuda, Y. *et al.*, *Org. Lett.*, 2008, **10**, 125-
 128 (*synth*)

Psychrophilin A

P-745




C₂₃H₂₀N₄O₅ 432.435
 Prod. by *Penicillium ribeum* IBT 16537.
 Solid. Mp 164-167°. [α]_D²⁵ +43.7 (MeCN).
 λ_{max} 217 (log ε 4.37); 244 (log ε 4.15); 304
 (log ε 3.74) (MeCN).

Dalsgaard, P.W. *et al.*, *J. Nat. Prod.*, 2004, **67**,
 878-881 (*isol, pmr, cmr*)

Psychrophilin B

P-746



R = -CH(CH₃)₂

C₂₃H₂₂N₄O₅ 434.451
 Prod. by *Penicillium rivulum* IBT 24420.
 Amorph. powder (MeCN aq.). Mp 186-
 190°. [α]_D²⁵ +15.2 (c, 0.03 in MeCN). λ_{max}
 214 (log ε 4.27); 244 (log ε 4.09); 302 (log
 ε 3.65) (MeCN).

Dalsgaard, P.W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1950-1952 (*isol, cd, pmr, cmr*)

Psychrophilin C P-747

As Psychrophilin B, P-746 with
R = CH₃

C₂₁H₁₈N₄O₅ 406.397

Prod. by *Penicillium rivulum* IBT 24420. Amorph. powder (MeCN aq.). Mp 167-171°. [α]_D²⁵ +16.7 (c, 0.02 in MeCN).

Dalsgaard, P.W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1950-1952 (*isol, cd, pmr, cmr*)

Psychrophilin D P-748

As Psychrophilin B, P-746 with
R = -CH₂CH(CH₃)₂


C₂₄H₂₄N₄O₅ 448.477

Prod. by *Penicillium algidum* IBT 22067. Powder (MeCN aq.). Mp 93-95°. [α]_D²⁵ +24.4 (c, 0.18 in MeCN). λ_{\max} 194 (log ϵ 4.52); 244 (log ϵ 4.13); 303 (log ϵ 3.76) (MeCN).

Dalsgaard, P.W. *et al.*, *J. Antibiot.*, 2005, **58**, 141-144 (*isol, cd, pmr, cmr*)

Psylloborine A P-749

[215596-11-5]



C₂₆H₄₀N₂ 380.615

Alkaloid from the ladybird beetle *Psyllobora vigintiduopunctata* (*Psyllobora 22-punctata*). Oil.

 $\Delta^{3a,4}$ -Isomer: **Isopsylloborine A**

C₂₆H₄₀N₂ 380.615


Alkaloid from the ladybird beetles *Halyzia 16-guttata* and *Vibidia 12-guttata*. Oil. [α]₅₇₉ +17 (c, 0.53 in CHCl₃). Possesses 3 β H config.

Schroeder, F.C. *et al.*, *Tetrahedron*, 1998, **54**, 12243-12248 (*isol, pmr, cmr*)

Lebrun, B. *et al.*, *Tet. Lett.*, 1999, **40**, 8115-8116 (*Isopsylloborine A*)

Ptelecortine P-750

4,8-Dimethoxy-1-methyl-6,7-methylene-dioxy-3-prenyl-2(1H)-quinolinone [36017-59-1]



C₁₈H₂₁NO₅ 331.368

Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (hexane). Mp 126-128°.

$\Delta^{3',4'}$ -Isomer, 2'- ζ -hydroxy: **Ptelefructine** [27741-35-1]

C₁₈H₂₁NO₆ 347.367

Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (EtOAc/hexane). Mp 146-149°.

Reisch, J. *et al.*, *Tet. Lett.*, 1970, 1945; 1972, 449 (*uv, ir, pmr, ms, struct*)

Pteledimericine P-751

[72751-06-5]


C₃₀H₃₀N₂O₅ 498.577

Struct. unknown. Dimeric quinolinone alkaloid. Alkaloid from the root bark of *Ptelea trifoliata* (Rutaceae). Needles (EtOH/Me₂CO). Mp 275-277°.

Mester, I. *et al.*, *Annalen*, 1979, 1785-1788 (*isol, uv*)

Pteledimeridine P-752

[72681-97-1]




C₃₀H₃₀N₂O₄ 482.578

Alkaloid from the root bark of *Ptelea trifoliata* (Rutaceae). Needles (Me₂CO). Mp 340-342°.

Mester, I. *et al.*, *Annalen*, 1979, 1785 (*isol, uv, ir, pmr, ms, struct*)

Pteledimerine P-753

[70509-83-0]




C₃₀H₃₀N₂O₄ 482.578

Alkaloid from the root bark of *Ptelea trifoliata* (Rutaceae). Cryst. (Me₂CO). Mp 319-321°.

Reisch, J. *et al.*, *Tet. Lett.*, 1978, 3681 (*ir, uv, pmr, ms, struct*)

Pteleflorine P-754

[58880-20-9]



C₁₆H₁₇NO₅ 303.314


Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (C₆H₆/hexane). Mp 93-96°.

Reisch, J. *et al.*, *Phytochemistry*, 1975, **14**, 2722 (*isol, uv, ir, ms, struct*)

Neville, C.F. *et al.*, *J.C.S. Perkin 1*, 1991, 259 (*synth, pmr*)

Ptelefoline P-755

3-(2-Hydroxy-3-methyl-3-butenyl)-4,6,8-trimethoxy-1-methyl-2(1H)-quinolinone [254888-61-3]



C₁₈H₂₃NO₅ 333.383

Alkaloid from leaves of *Ptelea trifoliata* (Rutaceae). Mp 91-93°. Opt. inactive.

Me ether: **O-Methylptelefoline**

C₁₉H₂₅NO₅ 347.41

Alkaloid from *Ptelea trifoliata* (Rutaceae). Oil.

3',4'-Dihydro, 3'-hydroxy: 3-(2,3-Dihydroxy-3-methylbutyl)-4,6,8-trimethoxy-1-methyl-2(1H)-quinolinone, **9CI. Ptelefolinol, 9CI** [59719-25-4]


C₁₈H₂₅NO₆ 351.399

Alkaloid from *Ptelea trifoliata* (Rutaceae).

Reisch, J. *et al.*, *Tet. Lett.*, 1969, 3803; 1970, 1945 (*Ptelefoline, O-Methylptelefoline*)
Korosi, J. *et al.*, *Herba Hung.*, 1976, **15**, 9; *CA*, **85**, 30608b (*Ptelefolinol*)

Ptelefolone P-756

3,9-Dihydro-6,8-dimethoxy-9-methyl-2-(1-methylethenyl)furo[2,3-b]quinolin-4(2H)-one, **9CI. 3,9-Dihydro-6,8-dimethoxy-2-isopropenyl-9-methylfuro[2,3-b]quinolin-4(H)-one** [28385-67-3]



C₁₇H₁₉NO₄ 301.341


Alkaloid from the leaves and flowers of *Ptelea trifoliata* (Rutaceae). Cryst. (hexane/Me₂CO). Mp 70-71°. [α]_D²⁸ +1.9 (c, 1.5 in CHCl₃).

Reisch, J. *et al.*, *Tet. Lett.*, 1970, 3365 (*uv, ir, pmr, ms, struct*)

Gaston, J.L. *et al.*, *Tet. Lett.*, 1978, 2629 (*synth*)

Ptelefolonium P-757

2,3-Dihydro-4,6,8-trimethoxy-9-methyl-2-(1-methylethenyl)furo[2,3-b]quinolinium(1+), **9CI. O⁴-Methylptelefolonium** [52768-97-5]




C₁₈H₂₂NO₄[⊕] 316.376Alkaloid from the leaves and callus strains of *Ptelea trifoliata* (Rutaceae).Chloride: Mp 135° (94-96°). [α]_D²⁰ +15.28 (c, 0.006 in MeOH).


Perchlorate: Mp 254-256° (236-238°).

Reisch, J. et al., *Phytochemistry*, 1973, **12**, 2552 (isol, uv, pmr, struct)Mitscher, L.A. et al., *J. Nat. Prod.*, 1975, **38**, 109 (isol, ir, pmr)Gaston, J.L. et al., *Tet. Lett.*, 1978, 2629 (synth)Rideau, M. et al., *Phytochemistry*, 1979, **18**, 155 (isol, pmr, ms)Petit-Paly, G. et al., *Plant Cell Rep.*, 1987, **6**, 309 (isol, uv, pmr, ms)**Pteleoline****P-758**

[36017-61-5]

C₁₈H₂₁NO₆ 347.367Alkaloid from *Ptelea trifoliata* (Rutaceae). Oil. [α]_D²⁵ -1.8 (c, 1.12 in CHCl₃).Reisch, J. et al., *Tet. Lett.*, 1972, 449 (uv, ir, pmr, ms, struct)**Pteleprenine****P-759**

6-Methoxy-9-methyl-7-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-h]quinolin-8(9H)-one, 9CI. 4-Methoxy-1-methyl-7,8-methylenedioxy-3-prenyl-2-quinolone [17232-50-7]

C₁₇H₁₉NO₄ 301.341Alkaloid from leafless shoots of *Ptelea trifoliata* (Rutaceae). Needles (petrol). Mp 76-77° (synthetic).3',4'-Dihydro, 2'ξ,3'-dihydroxy: **Hydroxylumidine**

[518-59-2]

C₁₇H₂₁NO₆ 335.356Minor alkaloid from *Lunasia amara* (Rutaceae). Rosettes (EtOAc/cyclohexane). Mp 124-125°. [α]_D²³ +27.6 (c, 0.820 in EtOH).Δ^{3',4'}-Isomer, 2'ξ-hydroxy: **Ptelefolidine**

[27745-40-0]

C₁₇H₁₉NO₅ 317.341Alkaloid from *Ptelea trifoliata* (Rutaceae). Cryst. (EtOAc/hexane). Mp 118-119°.Δ^{3',4'}-Isomer, 2'ξ-methoxy: **O-Methylptelefolidine**

[27745-41-1]

C₁₈H₂₁NO₅ 331.368Alkaloid from *Ptelea trifoliata* (Ruta-

ceae). Cryst. (hexane). Mp 133-135°.


3',4'-Dihydro, 2'-oxo: see Lunidonine, L-295

Goodwin, S. et al., *J.A.C.S.*, 1959, **81**, 6209 (Hydroxylumidine)Bowman, R.M. et al., *J.C.S. (C)*, 1967, 2368 (Pteleprenine)Novak, I. et al., *Herba Hung.*, 1970, **9**, 23; *CA*, **75**, 77090k (Hydroxylumidine, isol)Reisch, J. et al., *Tet. Lett.*, 1970, 1945

(Ptelefolidine, O-Methylptelefolidine)

Korosi, J. et al., *Herba Hung.*, 1974, **13**, 23; *CA*, **83**, 93836d (Pteleprenine, isol)**Pteriatoxin A****P-760**

[352275-99-1]

R = -³⁴CH(OH)CH₂CH₂CH(NH₂)COOH (34S,2'R-)C₄₅H₇₀N₂O₁₀S 831.121Exists as a zwitterion. Neutral form shown. Related to Pinnatoxin A, P-437. Toxin from the bivalve *Ptereria penguin*.

▶ Acutely toxic to mice.

Takada, N. et al., *Tet. Lett.*, 2001, **42**, 3495-3497 (isol)Kuramoto, M. et al., *Mar. Drugs*, 2004, **2**, 39-54 (rev, tox)Matsuura, F. et al., *J.A.C.S.*, 2006, **128**, 7463-7465; 7742-7743 (synth, abs config)**Pteriatoxin B****P-761**

[352276-00-7]

As Pteriatoxin A, P-760 with

R = ³⁴CH(CH₂OH)SCH₂²CH(NH₂)-COOH (34R,2'R-)C₄₅H₇₀N₂O₁₀S 831.121Exists as a zwitterion. Neutral form shown. Toxin from the bivalve *Ptereria penguin*.


▶ Acutely toxic.

34-Epimer: **Pteriatoxin C**

[352276-01-8]

C₄₅H₇₀N₂O₁₀S 831.121Toxin from *Ptereria penguin*.

▶ Acutely toxic.

Takada, N. et al., *Tet. Lett.*, 2001, **42**, 3495-3497 (isol)Matsuura, F. et al., *J.A.C.S.*, 2006, **128**, 7463-7465; 7742-7743 (synth, abs config)**2,4(1H,3H)-Pteridinedione, 9CI****P-762***Lumazine*, 8CI. 2,4-Pteridinediol. 2,4-DihydroxypteridineC₆H₄N₄O₂ 164.123

Di-NH-form occurs in aq. soln.

Intermediate tautomers also possible.

Yellow cryst. (H₂O). Mp 350°. Subl. 0.05 145. pK_{a1} 7.95 (20°). Bluish-green fluor. in aq. soln., green in alkali, blue in acid.1-Me: **1-Methyl-2,4-pteridinedione** [50256-18-3]C₇H₆N₄O₂ 178.15Isol. from the New Caledonian sponge *Corallistes fulvodesmus*. Needles (H₂O). Mp 290-291° (271°).

3-Me: [50256-19-4]

C₇H₆N₄O₂ 178.15Cryst. (H₂O). Mp 332°.

1-Ph: [32433-26-4]

C₁₂H₈N₄O₂ 240.221

Cryst. Mp 307-308°.

3-Ph:

C₁₂H₈N₄O₂ 240.221

Cryst. (MeOH). Mp 359-360°.

Di-NH-form [487-21-8]

▶ UO3416000

1,3-Di-Me: [13401-18-8]

C₈H₈N₄O₂ 192.177Cryst. (H₂O). Mp 200°.

1,3-Di-Et: [94591-34-1]

C₁₀H₁₂N₄O₂ 220.23Cryst. (Et₂O/petrol). Mp 87-88°.

1,3-Dibenzyl: [94591-35-2]

C₂₀H₁₆N₄O₂ 344.372

Needles (EtOH aq.). Mp 140°.

Aldrich Library of NMR Spectra, 2nd edn.,1983, **2**, 768A (nmr)*Aldrich Library of FT-IR Spectra*, 1st edn.,1985, **2**, 894B (ir)Pfleiderer, W. et al., *Chem. Ber.*, 1957, **90**, 2582

(1-Me, 3-Me, 1,3-di-Me, uv)

Brown, D.J. et al., *J.C.S.*, 1961, 4413

(synth)

Pfleiderer, W. et al., *Chem. Ber.*, 1973, **106**,

3149 (synth)

Ewers, J. et al., *Chem. Ber.*, 1974, **107**, 3275

(cmr)

Pfleiderer, W. et al., *Annalen*, 1984, 1798

(derivs, synth)

Tsuzuki, K. et al., *J. Het. Chem.*, 1986, **23**,


1299 (1-Me)

Klein, R. et al., *Photochem. Photobiol.*, 1987,**45**, 55 (tautom, uv)Debitus, C. et al., *Tet. Lett.*, 1989, **30**, 1534-

1538 (1-Me, isol)

Goswami, S. et al., *Eur. J. Org. Chem.*, 2007,

4056-4064 (synth, ir, pmr)

2,4,7(1H,3H,8H)-Pteridine-trione, 9CI, 8CI**P-763***Isoxantholumazine*. *Violapterin*. 7-Oxolumazine. 2,4,7-Trihydroxypteridine [2577-38-0]C₆H₄N₄O₃ 180.123

Prod. by *Physarum polycephalum* and the ant *Formica polyctena*. Metab. of 2-Amino-4,7-dihydroxypteridine, A-751. Cryst. (H₂O). Mp 350°.

1-Me: [2614-44-0]
C₇H₆N₄O₃ 194.149
Mp > 340.

3-Me: [2622-65-3]
C₇H₆N₄O₃ 194.149
Cryst. (H₂O). Mp > 340°.

6-Me: [31053-46-0]
C₇H₆N₄O₃ 194.149
Cryst. + 1H₂O (H₂O). Mp 340°.

8-Me: 8-Methyl-2,4,7-(1H,3H,8H)-pteridinetriene, 9CI, 8CI. 1,2,3,4,7,8-Hexahydro-8-methyl-2,4,7-trioxopteridine.

Luciopterin

[19845-00-2]
C₇H₆N₄O₃ 194.149

Constit. of the Japanese firefly *Luciola cruciata*. Strongly fluorescent, pale yellow microcryst. Mp > 340°. pK_{a1} 3.69; pK_{a2} 0 (H₂O).

1,3-Di-Me: [2614-43-9]
C₈H₈N₄O₃ 208.176
Mp 264°.

1,6-Di-Me:
C₈H₈N₄O₃ 208.176
Cryst. (H₂O). Mp 330° dec.

1,8-Di-Me: [70916-42-6]
C₈H₈N₄O₃ 208.176
Cryst. (EtOH). Mp 310° dec.

3,6-Di-Me:
C₈H₈N₄O₃ 208.176
Cryst. (H₂O). Mp 340°.

3,8-Di-Me: [70916-39-1]
C₈H₈N₄O₃ 208.176
Cryst. Mp 350°. Darkens at 300°.

6,8-Di-Me: [6743-25-5]
C₈H₈N₄O₃ 208.176
Cryst. Mp 350°.

1,3,6-Tri-Me: [2625-21-0]
C₉H₁₀N₄O₃ 222.203
Needles (H₂O). Mp 308°.

1,3,8-Tri-Me: [70674-02-1]
C₉H₁₀N₄O₃ 222.203
Pale yellow cryst. (EtOH). Mp 220°.

1,6,8-Tri-Me: [70916-41-5]
C₉H₁₀N₄O₃ 222.203
Cryst. (EtOH). Dec. at 317°.

3,6,8-Tri-Me: [6743-26-6]
C₉H₁₀N₄O₃ 222.203
Cryst. (EtOH). Mp 331-334°.

1,3,6,8-Tetra-Me:
C₁₀H₁₂N₄O₃ 236.23
Pale yellow needles (EtOH aq.). Mp 253°.

Pfleiderer, W. et al., *Chem. Ber.*, 1957, **90**, 2588-2603; 1958, **91**, 1671-1680 (*synth*, *N-Me* derivs)

McNutt, W.S. et al., *J. Biol. Chem.*, 1963, **238**, 1116-1121 (7-Oxolumazine, *metab*)


Kishi, Y. et al., *Tet. Lett.*, 1968, **9**, 2847-2850 (*Luciopterin*, *isol*)

Schmidt, G.H. et al., *Z. Naturforsch., B*, 1969, **24**, 1153-1169 (*isol*)

Bergmann, F. et al., *J.C.S. Perkin 2*, 1979, 35-39 (1,8-di-Me, *synth*)

Loidl, P. et al., *CA*, 1984, **100**, 20127 (*isol*)

Pterocellin A
[593235-02-0]



R¹ = CH₂CH(CH₃)₂, R² = H

C₁₆H₁₆N₂O₃ 284.314
Isol. from the marine bryozoan *Pterocella vesiculosa*. Cytotoxic. Dark red needles (toluene). Mp 172-173°. λ_{max} 202 (log ε 3.83); 256 (log ε 3.7); 280 (log ε 3.87); 483 (log ε 2.93) (MeOH).

6,6'-Dimer: Pterocellin E
C₃₂H₃₀N₄O₆ 566.612
Isol. from *Pterocella vesiculosa*. Rose-pink solid. λ_{max} 201 (log ε 4.65); 230 (sh) (log ε 4.48); 258 (sh) (log ε 4.48); 286 (log ε 4.57); 480 (log ε 3.61) (MeOH).

Yao, B. et al., *J. Nat. Prod.*, 2003, **66**, 1074-1077 (*isol*, *pmr*, *cmr*, *N-15 nmr*, *cryst struct*)

O'Malley, M.M. et al., *Org. Lett.*, 2006, **8**, 2651-2652 (*synth*)

Prinsep, M.R. et al., *J. Nat. Prod.*, 2008, **71**, 134-136 (*Pterocellin E*)

Pterocellin B

[593235-03-1]
As Pterocellin A, P-764 with
R¹ = -CH₂Ph, R² = H

C₁₉H₁₄N₂O₃ 318.331
Isol. from the marine bryozoan *Pterocella vesiculosa*. Cytotoxic. Amorph. red solid. λ_{max} 202 (log ε 4.08); 287 (log ε 3.63); 485 (log ε 2.63) (MeOH).

Yao, B. et al., *J. Nat. Prod.*, 2003, **66**, 1074-1077 (*isol*, *pmr*, *cmr*, *N-15 nmr*)

Pterocellin C

[1000598-23-1]
As Pterocellin A, P-764 with
R¹ = CH₂CH(CH₃)₂, R² = -
(CH₂)₄COCH₃

C₂₂H₂₆N₂O₄ 382.458
Isol. from *Pterocella vesiculosa*. Orange-red solid. λ_{max} 203 (log ε 4.87); 224 (sh) (log ε 4.76); 257 (log ε 4.74); 290 (log ε 5.02); 489 (log ε 4.06) (MeOH).

Prinsep, M.R. et al., *J. Nat. Prod.*, 2008, **71**, 134-136 (*isol*, *pmr*, *cmr*, *ms*)


Pterocellin D

[1000598-24-2]
As Pterocellin A, P-764 with
R¹ = -CH₂CH(CH₃)₂, R² = -CH(CH₃)₂
C₁₉H₂₂N₂O₃ 326.394
Isol. from *Pterocella vesiculosa*. Red-orange solid. λ_{max} 203 (log ε 4.29); 255 (log ε 4.01); 290 (log ε 4.11); 490 (log ε 3.08) (MeOH).

Prinsep, M.R. et al., *J. Nat. Prod.*, 2008, **71**, 134-136 (*isol*, *pmr*, *cmr*, *ms*)

P-764

Pterocellin F
[1000598-26-4]



C₃₃H₃₂N₄O₆ 580.639
Isol. from *Pterocella vesiculosa*. Dark red solid. λ_{max} 202 (log ε 3.8); 286 (log ε 3.36); 494 (log ε 2.2) (MeOH).
Prinsep, M.R. et al., *J. Nat. Prod.*, 2008, **71**, 134-136 (*isol*, *pmr*, *cmr*, *ms*)

Pterogynidine

P-769
N,N'-Bis(3-methyl-2-butenyl)guanidine, 9CI, N,N'-Diisopentenylguanidine, N,N'-Diprenylguanidine
[25713-89-7]
HN=C[NHCH₂CH=C(CH₃)₂]₂


C₁₁H₂₁N₃ 195.307
Alkaloid from *Alchornea cordifolia* and *Pterogyne nitens*. Mp 105-106° (as picrate). λ_{max} 203 ; 273 (MeOH).

Corral, R.A. et al., *Chem. Comm.*, 1970, 556 (*isol*, *synth*, *pmr*)

Mavar-Manga, H. et al., *Nat. Prod. Commun.*, 2006, **1**, 1097-1100 (*isol*, *pmr*, *cmr*)

Pterophorine

P-770
12-(Acetyloxy)-3,8-didehydro-14-methyl-21-norsenecionan-5,11,16-trione, 9CI
[62786-99-6]



C₂₀H₂₃NO₇ 389.404
Alkaloid from seeds of *Senecio pterophorus* and *Senecio inaequidens* (Asteraceae). Oil. [α]_D +27 (c, 4.6 in CHCl₃).

Stereoisomer: Isopterophorine
[71075-40-6]


C₂₀H₂₃NO₇ 389.404
Alkaloid from roots of *Senecio pulviniformis*. Oil. [α]_D -25 (CHCl₃). Partial stereochemistries are shown for Pterophorine and Isopterophorine but these do not appear to be well established.

Bohlmann, F. et al., *Chem. Ber.*, 1977, **110**, 474 (*isol*, *ord*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1977, **16**, 965; 1979, **18**, 125 (*isol, ir, pmr, struct*)

Pterorhodine, 8CI**P-771**

2-Amino-7-[(2-amino-1,4,5,6-tetrahydro-4,6-dioxo-7-pteridiny)lmethylene]-1,5,7,8-tetrahydro-4,6-pteridinedione, 9CI. Rhodomelanochrome. Lepidoporphyrin. Rhosoptein. Platynerepterin [6538-79-0]




C₁₃H₁₀N₁₀O₄ 370.287

Isol. from wings of butterflies, e.g. *Appias nero*, and from the skin of the leaf frog *Agalychnis danielcolor*. Red-violet cryst.

Purrmann, R. *et al.*, *Annalen*, 1944, **556**, 189 (*isol*)
 Russell, P.B. *et al.*, *J.A.C.S.*, 1949, **71**, 3412
 Pfeleiderer, W. *et al.*, *Z. Naturforsch.*, B, 1963, **18**, 420 (*isol*)
 Viscontini, M. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1207-1209 (*isol, struct*)
 Misuraca, G. *et al.*, *Comp. Biochem. Physiol.*, B: *Comp. Biochem.*, 1977, **57**, 41 (*isol*)
 Pfeleiderer, W. *et al.*, *J. Het. Chem.*, 1992, **29**, 583 (*rev*)

Pteroylglutamic acid**P-772**

N-[4-[[(2-Amino-1,4-dihydro-4-oxo-6-pteridiny)lmethyl]amino]benzoyl]glutamic acid, 9CI. Folic acid, BAN, INN. Vitamin B_c. Vitamin M. Coenzyme F. Leucal. CoF. Citrovorin. Folacin. Folvite. Foliamin. NSC 3073 [59-30-3]



C₁₉H₁₉N₇O₆ 441.402

Other tautomers also possible. Natural isolates normally contain homologues having n > 1. Present in liver and kidney tissue, fungi, green plants, etc. Essential cofactor. Used in the treatment of megaloblastic anaemias to stimulate erythrocyte, lymphocyte and platelet production. Recommended as dietary supplement during pregnancy as preventative against spina bifida. Pale lemon-yellow to light-orange needles (H₂O at pH 3). Almost insol. H₂O; sl. sol. MeOH; mod. sol. AcOH; insol. Et₂O. [α]_D²⁵ +23 (c, 0.5 in 1M NaOH). pK_{a1} 4.65; pK_{a2} 6.75; pK_{a3} 9 (30°, 0.01M KNO₃). Log P -1.02 (calc). Chars above 250°. Component of numerous preparations. Also used as calcium salt (Calcium folate, BAN, INN).

► LD₅₀ (mus, orl) 10,000 mg/kg. Exp. teratogen. LP5425000

7,8-Dihydro: 7,8-Dihydrofolic acid. 7,8-Dihydropteroylglutamic acid

[4033-27-6]

C₁₉H₂₁N₇O₆ 443.418

Metabolic product of reduction of folic acid by dihydrofolate reductase. Yellow cryst. (H₂O). pK_{a1} 0.34; pK_{a2} 4.2; pK_{a3} 10.3.

5,6,7,8-Tetrahydro: see 5,6,7,8-Tetrahydrofolic acid, T-172

Homologue (n = 2): Pteroyldiglutamic acid. Pteroylglutamylglutamic acid.

Diopterin

[6807-82-5]

C₂₄H₂₆N₈O₉ 570.518

Prod. by microorganisms.

Homologue (n = 2), 7,8-dihydro, N⁸-Me: 7-Hydro-8-methylpteroylglutamylglutamic acid

[108402-49-9]

C₂₅H₃₀N₈O₉ 586.56

Isol. from actinomycete strain SK 2049, identified as *Promicromonospora sukumoe*. Antifolate agent, thymidilate synthase inhibitor. Shows antimalarial props. Yellowish powder + 5H₂O. Sol. bases; fairly sol. H₂O; poorly sol. EtOH, hexane, CHCl₃. Mp 196° dec. [α]_D²⁰ +24 (c, 0.25 in H₂O). λ_{max} 200 (ε 13000); 219 (ε 11000); 292 (ε 17500) (0.01M HCl) (Derep). λ_{max} 203 (ε 22000); 215 (sh) (ε 13500); 282 (ε 17000) (0.01N NH₄OH) (Derep). λ_{max} 216 (sh) (ε 13500); 290 (ε 17500) (H₂O) (Derep). λ_{max} 217 (ε 13800); 299 (ε 18200) (H₂O) (Berdy). λ_{max} 220 (ε 12000); 302 (ε 19000) (HCl) (Berdy).

Homologue (n = 3): Pteroyltriglutamic acid. Pteroylglutamylglutamylglutamic acid. Dripterin. Pteropterin. Terapterin. Lactobacillus casei Factor. Teropterin [89-38-3]
 C₂₉H₃₃N₉O₁₂ 699.633
 Prod. by *Lactobacillus casei*, *Corynebacterium* spp. and others. Antineoplastic agent. Amorph. Log P -3.65 (calc).

Homologue (n = 5): Pteroylpentaglutamic acid

[33611-85-7]

C₃₉H₄₇N₁₁O₁₈ 957.863

Found in yeast etc.

Homologue (n = 7): Pteroylheptaglutamic acid. Pteroylhexaglutamylglutamic acid. Folic acid conjugate. Vitamin B_c conjugate

[6484-74-8]

C₄₉H₆₁N₁₃O₂₄ 1216.094

Found in yeast, pollen etc. Microcryst. (H₂O).

[1492-18-8, 6484-89-5, 75708-92-8, 6713-32-2]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 898A (*ir*)

Pfiffner, J.J. *et al.*, *J.A.C.S.*, 1946, **68**, 1392 (*Pteroylheptaglutamic acid*)

Boothe, J.H. *et al.*, *J.A.C.S.*, 1948, **70**, 1099; 1949, **71**, 2304

Weisblat, D.I. *et al.*, *J.A.C.S.*, 1953, **75**, 3625 (*synth*)

Blakley, R.L. *et al.*, *Nature (London)*, 1960, **188**, 231 (*uv*)

Jaenicke, L. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1963, **21**, 183 (*rev*)

Pastore, E.J. *et al.*, *Ann. N.Y. Acad. Sci.*, 1971, **186**, 43 (*pmr*)

Shiota, T. *et al.*, *Compr. Biochem.*, Elsevier, Amsterdam, 1971, **21**, 111 (*rev, biosynth*)

Godwin, H.A. *et al.*, *J. Biol. Chem.*, 1972, **247**, 2266 (*Pteroyltriglutamic acid*)

Bieri, J. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 2905 (*synth*)

Frick, W. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 2658 (*cmr*)

Drey, C.N.C. *et al.*, *Chem. Comm.*, 1977, 144 (*Pteroyltriglutamic acid*)

Pont, L.O. *et al.*, *Anal. Profiles Drug Subst.*, 1979, **8**, 315 (*rev*)

Murphy, M. *et al.*, *Biochim. Biophys. Acta*, 1979, **583**, 535 (*metab*)

Nixon, P.F. *et al.*, *Clin. Exp. Pharmacol. Physiol.*, Suppl. 1, 1979, **5**, 35 (*rev, pharmacol*)

Poe, M. *et al.*, *Methods Enzymol.*, 1980, **66**, 483 (*pmr*)

Mastro Paolo, D. *et al.*, *Science (Washington, D.C.)*, 1980, **210**, 334 (*cryst struct*)

Straw, J.E. *et al.*, *Cancer Res.*, 1984, **44**, 3114 (*metab*)

Cooper, B.A. *et al.*, *Clin. Biochem.*, 1984, **17**, 95 (*metab*)

Blakeley, R.L. *et al.*, *Folates Pterins*, (Eds.), J. Wiley & Sons, NY, 1984, **1**, (*chem, biochem*)

Brody, T. *et al.*, *Food Sci. Technol.*, 1984, **13**, 459 (*biol studies*)

Charlton, P.A. *et al.*, *J.C.S. Perkin 1*, 1985, 1349 (*7,8-dihydro, synth*)

D'Ari, L. *et al.*, *Methods Enzymol.*, 1985, **113**, 169 (*Pteroyltriglutamic acid*)

Shane, B. *et al.*, *Methods Enzymol.*, 1986, **122**, 269; 323 (*detn, synth*)

Moran, R.G. *et al.*, *Methods Enzymol.*, 1986, **122**, 309 (*synth*)

Young, D.W. *et al.*, *Nat. Prod. Rep.*, 1986, **3**, 395 (*biosynth*)

Rees, L. *et al.*, *Tetrahedron*, 1986, **42**, 117 (*7,8-dihydro, synth*)

Rees, L. *et al.*, *Chem. Comm.*, 1987, 470 (*synth*)

Murata, M. *et al.*, *J. Antibiot.*, 1987, **40**, 251 (*7-Hydro-8-methylpteroylglutamylglutamic acid*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 5190

Al-Shammary, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1990, **19**, 221 (*rev*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1039

Rossi, C. *et al.*, *Spectrosc. Lett.*, 1993, **26**, 1603 (*pmr, conform*)


Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 1122-1123

Roje, S. *et al.*, *Phytochemistry*, 2007, **68**, 1904-1921 (*biosynth, rev*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FMT000

Ptilocaulin**P-773**

[78777-02-3]



(+)-form

C₁₅H₂₅N₃ 247.383

Isol. from the Caribbean sponge *Ptilocaulis* aff. *Ptilocaulis spiculifer*, may be a symbiont metab. Also from *Batzella* sp., *Monanchora arbuscula* and *Hemimycala* sp. Active against gram-positive bacteria and leukaemia cells. Cryst. (as nitrate). Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 183-185° (nitrate). [α]_D +74.4 (MeOH). λ_{max} 225 (ε 10000) (MeOH).

8b-Hydroxy: **8b-Hydroxyptilocaulin**
[167499-84-5]

C₁₅H₂₅N₃O 263.382

Alkaloid from the marine sponge *Monanchora arbuscula*. Oil (as hydrochloride). [α]_D +77.5 (c, 0.12 in MeOH) (hydrochloride).

7-Epimer, Δ^{8a,8b}-isomer (8αH): **7-Epineoptilocaulin**. 8a,8b-Dehydroptilocaulin

C₁₅H₂₅N₃ 247.383

Renamed in 2008. Isol. from the sponge *Batzella* sp. Oil. [α]_D +13.3 (c, 1.2 in MeOH). λ_{max} 228 ; 242 ; 306 (MeOH).

7-Epimer, Δ^{8a,8b}-isomer, 8α-hydroxy: **8α-Hydroxy-7-epineoptilocaulin**. 8a,8b-Dehydro-8-hydroxyptilocaulin

C₁₅H₂₅N₃O 263.382

Renamed in 2008. Isol. from the sponge *Batzella* sp. Gum. [α]_D +24.2 (c, 0.33 in MeOH). λ_{max} 237 ; 299 (MeOH).

[86594-30-1 , 88154-76-1]

Harbour, G.C. et al., *J.A.C.S.*, 1981, **103**, 5604 (isol, uv, pmr, cmr, ms, struct)

Roush, M.R. et al., *J.A.C.S.*, 1984, **106**, 721 (synth)

Snider, B.B. et al., *J.A.C.S.*, 1984, **106**, 1443 (synth, ir, pmr, cmr, ms, cd)

Watts, A.E. et al., *Tetrahedron*, 1985, **41**, 3463 (synth)

Uyehara, T. et al., *Chem. Comm.*, 1986, 539 (synth)

Murthy, K.S.K. et al., *Isr. J. Chem.*, 1991, **31**, 239 (synth)

Snider, B.B. et al., *J.O.C.*, 1993, **58**, 3828-3839 (biosynth)

Tavares, R. et al., *J. Nat. Prod.*, 1995, **58**, 1139 (8b-Hydroxyptilocaulin)


Patil, A.D. et al., *J. Nat. Prod.*, 1997, **60**, 704-707 (isol, uv, pmr, cmr, derivs)

Yu, M. et al., *J.O.C.*, 2008, **73**, 9065-9074 (synth, pmr, cmr)

Ptilomycalin A

P-774

[124512-47-6]

C₄₅H₈₀N₆O₅ 785.164

Alkaloid from the Caribbean sponge *Ptilocaulis spiculifer*, the Red Sea sponge *Hemimycala* sp., the New Caledonian starfish *Celerina heffernani* and the Caribbean sponge *Batzella* sp. Exhibits remarkable antifungal, antiviral and antitumour activities. [α]_D²⁵ -2.5 (c, 0.70 in

CHCl₃).36R-Hydroxy: **Celeromycalin**

[163597-72-6]

C₄₅H₈₀N₆O₆ 801.164

Alkaloid from the starfish *Celerina heffernani*. Highly cytotoxic. [α]_D -4.5.

38-Parent acid: **Crambescidic acid**C₃₈H₆₃N₃O₆ 657.932

Constit. of *Monanchora dianchora* and *Monanchora unguifera*. [α]_D²⁵ +4.4 (c, 0.2 in MeOH).

Kashman, Y. et al., *J.A.C.S.*, 1989, **111**, 8925-8926 (isol, ir, pmr, cmr, struct)

Ohtani, I. et al., *J.A.C.S.*, 1992, **114**, 8472-8479 (isol, ir, pmr, cmr, struct)

Overman, L.E. et al., *J.A.C.S.*, 1995, **117**, 2657-2658 (synth)

Patil, A.D. et al., *J.O.C.*, 1995, **60**, 1182-1188 (isol)

Palagiano, E. et al., *Tetrahedron*, 1995, **51**, 3675-3682 (*Celeromycalin*)

Coffey, D.S. et al., *J.A.C.S.*, 2000, **122**, 4893-4903 (synth)


Gallimore, W.A. et al., *J. Nat. Prod.*, 2005, **68**, 1420-1423 (*Crambescidic acid*)

Bensemhoun, J. et al., *J. Nat. Prod.*, 2007, **70**, 2033-2035 (*Crambescidic acid*)

Ptilomycalin D

P-775

[1002730-82-6]

C₃₈H₆₅N₃O₄ 627.949


Alkaloid from *Monanchora dianchora*. [α]_D²⁵ -7.2 (c, 0.26 in CHCl₃).

Bensemhoun, J. et al., *J. Nat. Prod.*, 2007, **70**, 2033-2035 (isol, pmr, cmr)

Puberaconitine

P-776

[69787-06-0]

C₃₆H₅₀N₂O₁₁ 686.798

Alkaloid from the roots of *Aconitum barbatum* var. *puberulum* (Ranunculaceae). Amorph. [α]_D²⁰ +34 (c, 0.5 in CHCl₃).

Me ester: Septentriodine. Cashmiradelphine

[69787-05-9]

C₃₇H₅₂N₂O₁₁ 700.825

Alkaloid from the roots of *Delphinium cashmirianum*, *Aconitum septentrionale* and *Aconitum barbatum* var. *puberulum*, and from *Aconitum gigas* (Ranunculaceae). Cryst. (MeOH/Et₂O). Mp 130-135°. [α]_D²⁷ +56 (c, 0.39 in EtOH). λ_{max} 222 (log ε 3.06); 227 (sh) (log ε 2.97); 252 (log ε 2.71); 260 (sh) (log ε 2.61); 310 (log ε 2.28) (MeOH).

Amide: **Avadharidine**

[509-16-0]

C₃₆H₅₁N₃O₁₀ 685.813

Alkaloid from the roots of *Aconitum orientale*, *Aconitum finetianum* and *Delphinium cashmirianum* (Ranunculaceae). Amorph. [α]_D²⁷ +40 (c, 0.37 in EtOH).

O⁸-Me: **Puberaconitidine**

[83685-21-6]

C₃₇H₅₂N₂O₁₁ 700.825

Alkaloid from the roots of *Aconitum barbatum* var. *puberulum* (Ranunculaceae). [α]_D²⁰ +22.4 (c, 0.9 in CHCl₃).

O⁸-Me, Me ester: **Septentriodine**

[70553-63-8]

C₃₈H₅₄N₂O₁₁ 714.851

Alkaloid from the roots of *Aconitum septentrionale* and *Aconitum barbatum* var. *puberulum* (Ranunculaceae). Mp 123-125°. [α]_D²⁶ +21.2 (c, 1.0 in CHCl₃).

O¹⁴-De-Me, Me ester: **Vulparine**

[935754-38-4]

C₃₆H₅₀N₂O₁₁ 686.798

Alkaloid from *Aconitum vulparia*. Amorph. solid. [α]_D²⁷ +47 (c, 0.1 in CHCl₃). λ_{max} 223 (log ε 3.04); 227 (sh) (log ε 3.01); 252 (log ε 2.75); 260 (sh) (log ε 2.65); 311 (log ε 2.32) (MeOH).

O¹⁴-De-Me, 14-Ac, Me ester: **Finetiadine**

[178494-88-7]

C₃₈H₅₂N₂O₁₂ 728.835

Minor alkaloid from roots of *Aconitum finetianum*. Amorph. powder (Me₂CO). Mp 116-117°. [α]_D¹⁸ +31.63 (c, 0.1 in CHCl₃).

Kuzovkov, A.D. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1955, **25**, 161-163; 1959, **29**, 2746-2749 (*Avadharidine*)

Sakai, S. et al., *Yakugaku Zasshi*, 1978, **98**, 1376-1384 (*Septentriodine*)

Pelletier, S.W. et al., *Heterocycles*, 1979, **12**, 377-381 (*Septentriodine*)

Shamma, M. et al., *J. Nat. Prod.*, 1979, **42**, 615-623 (*Avadharidine*, uv, ir, pmr, ms, struct)

Pelletier, S.W. et al., *J.A.C.S.*, 1981, **103**, 6536-6538 (config)

Edwards, O.E. et al., *Can. J. Chem.*, 1982, **60**, 2661-2667 (config)

Jiang, S. et al., *Zhongcaoyao*, 1982, **13**, 5; *CA*, **97**, 107027u (*Avadharidine*, isol)

Yu, D. et al., *Planta Med.*, 1983, **49**, 85-89 (*Puberaconitidine*, *Septentriodine*)


Joshi, B.S. et al., *J. Nat. Prod.*, 1990, **53**, 1028-1030 (*Puberaconitine*, *Puberaconitidine*, config)

Wu, G. et al., *Phytochemistry*, 1996, **42**, 1253-1255 (*Finetiadine*)

Csupor, D. et al., *Z. Naturforsch., B*, 2007, **62**, 135-141 (*Vulparine*, *Septentriodine*)

Pubesamide B

P-777

C₂₅H₂₉NO₃ 391.509

Constit. of the seeds of *Casimiroa pubescens*. Yellow solid. Mp 89-92°. Isol. as a mixt. with Pubesamide A to which data refers.

(3'Z)-Isomer: **Pubesamide A**C₂₅H₂₉NO₃ 391.509

Constit. of the seeds of *Casimiroa pubescens*. Yellow solid. Mp 88-91°. λ_{max} 269 (log ε 4.38) (CHCl₃).

5'-Deoxo, Δ^{4'}-isomer (E-), *3'ξ*-hydroxy:**Pubesamide C**C₂₅H₃₁NO₃ 393.525


Constit. of the seeds of *Casimiroa pubescens*. Yellowish solid. Mp 83-84°. λ_{max} 202 (log ε 4.37); 230 (log ε 4.52) (MeOH).

García-Argáez, A.N. *et al.*, *Z. Naturforsch., B*, 2004, **59**, 245-248 (*isol, pmr, cmr, ms*)

Pubeschimine

P-778

[159249-81-7]

C₂₄H₄₀N₂O 372.593


Alkaloid from bark of *Holarrhena pubescens* (Apocynaceae). Irregular plates (Me₂CO). Mp 220-222°.

Begum, S. *et al.*, *Phytochemistry*, 1994, **36**, 1537-1541 (*isol, uv, ir, pmr, cmr, ms, struct*)

Pubeschine†

P-779

[159194-89-5]

C₂₁H₂₉NO 311.466


Alkaloid from bark of *Holarrhena pubescens* (Apocynaceae). Irregular plates (Me₂CO). Mp 224-226°.

Begum, S. *et al.*, *Phytochemistry*, 1994, **36**, 1537-1541 (*isol, uv, ir, pmr, cmr, ms, struct*)

Puhinamide

P-780

[204641-53-2]

C₂₃H₂₉NO₄ 383.486

Alkaloid from *Glycosmis cf. parva*. Cryst. (Et₂O). Mp 88-90°. λ_{max} 227; 296 (MeOH).

2'',3''-Dihydro: **Khaochamide**

[204641-52-1]

C₂₃H₃₁NO₄ 385.502


Alkaloid from *Glycosmis parva*. Cryst. (Et₂O). Mp 114-115°. λ_{max} 227; 241 (sh); 291 (MeOH).

Hofer, O. *et al.*, *Monatsh. Chem.*, 1998, **129**, 213-219

Pukateine

P-781

11-Hydroxy-1,2-methylenedioxyaporphine. N-Methylbovanine

C₁₈H₁₇NO₃ 295.337*(R)*-form [81-67-4]

Alkaloid from the bark of *Laurelia novae-zelandiae* (Monimiaceae). Cryst. (EtOH). Mp 210-211°. [α]_D¹⁵ -257 (c, 1.0 in EtOH).

α-N-Oxide:

Synthetic. Plates. Mp 160-162° (MeOH solvate) Mp 176-178°.

β-N-Oxide: **Laurepukine**

[34029-94-2]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the bark of *Laurelia novae-zelandiae* (Monimiaceae). Pale beige needles (MeOH). Mp 220-222°. [α]_D²⁵ -257 (c, 0.11 in CHCl₃/MeOH 1:1).

N-De-Me-11-Hydroxy-1,2-methylenedioxyaporphine. **Obovanine**

[53254-79-8]

C₁₇H₁₅NO₃ 281.31

Alkaloid from *Magnolia obovata*, the stem bark of *Duguetia calycina*, and the bark of *Laurelia novae-zelandiae* (Magnoliaceae, Annonaceae, Monimiaceae). Mp 260-262° (250°) dec. (as hydrochloride). [α]_D¹² -263 (c, 0.11 in MeOH) (-164) (hydrochloride).

Me ether-11-Methoxy-1,2-methylenedioxyaporphine. **O-Methylpukateine**

[16625-66-4]

C₁₉H₁₉NO₃ 309.364

Alkaloid from *Laurelia novae-zelandiae* and the stem bark of *Duguetia calycina* (Monimiaceae, Annonaceae). Mp 136-138°. [α]_D²⁵ -293.4 (c, 0.199 in CHCl₃). [α]_D²⁵ -271 (c, 0.105 in EtOH).

Me ether, N-de-Me-11-Methoxy-1,2-methylenedioxyaporphine. **Puterine**

[65012-38-6]

C₁₈H₁₇NO₃ 295.337

Alkaloid from stem bark of *Guatteria elata* and *Duguetia calycina* (Annonaceae). Shows antimicrobial activity.

*Me ether, β-N-oxide:*Cryst. (MeOH/Et₂O). Mp 146-149°

Mp 169-175° (double Mp).

Me ether, N-de-Me, N-formyl: N-Formylputerine

[89368-30-9]

C₁₉H₁₇NO₄ 323.348

Alkaloid from *Duguetia calycina* (Annonaceae). Noncryst. [α]_D -467 (c, 0.42 in CHCl₃).

*Me ether, N-de-Me, Ac:*Amorph. powder (Me₂CO). Mp 178-182°.[α]_D²⁵ -352 (c, 0.25 in MeOH).*7S-Hydroxy: Duguxine*

[112494-62-9]

C₁₈H₁₇NO₄ 311.337

Alkaloid from *Duguetia spixiana* (Annonaceae). Obt. only in admixture with Spixianine, S-469.

7S-Hydroxy, N-oxide: Duguxine N-oxide

[112494-65-2]

C₁₈H₁₇NO₅ 327.336

Alkaloid from *Duguetia spixiana* (Annonaceae).

(±)-form

Synthetic. Cryst. (EtOH). Mp 232-233° (213-215°).

Barger, G. *et al.*, *Helv. Chim. Acta*, 1931, **14**, 481 (*Pukateine, isol, struct*)

Girardet, A. *et al.*, *Helv. Chim. Acta*, 1931, **14**, 504 (*Laurepukine, isol*)

Bernhauer, K. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 1583 (*O-Methylpukateine*)

Zymalkowsky, F. *et al.*, *Chem. Ber.*, 1969, **102**, 2959 (*synth, pmr*)

Weiss, E. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 1342 (*Laurepukine, synth, ms, cd, ord, struct*)

Oberhänsli, W.E. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 1389 (*Laurepukine, cryst struct*)

Kametani, T. *et al.*, *J.C.S. Perkin 1*, 1972, 1435 (*synth, pmr*)

Ito, K. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 729 (*Obovanine, isol, uv, ir, pmr, ms, struct*)

Hsu, C.C. *et al.*, *J. Nat. Prod.*, 1977, **40**, 505 (*Puterine*)

Roblot, F. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 259; *CA*, **91**, 2517b (*Obovanine, O-Methylpukateine*)

Roblot, F. *et al.*, *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1981, **293**, 373 (*O-Methylpukateine, cmr*)

Urzúa, A. *et al.*, *Phytochemistry*, 1982, **21**, 773 (*pmr, ms*)


Roblot, F. *et al.*, *J. Nat. Prod.*, 1983, **46**, 862 (*N-Formylputerine*)

Villar, A. *et al.*, *Farm. Tijdschr. Belg.*, 1984, **61**, 300 (*Puterine, activity*)


Debourges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664 (*Duguxine, Duguxine N-oxide*)

Pukeensine

[144442-84-2]

C₄₄H₆₄N₂O₃ 669.001Alkaloid from the whole plant of *Aconitum pukeense* (Ranunculaceae).Ding, L.S. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 394; *CA*, **117**, 230089z (*isol, struct*)**Pukeleimide**

1-/[1,5-Dihydro-4-(hydroxymethyl)-1-methyl-5-oxo-2H-pyrrol-2-ylidene]acetyl]-1,5-dihydro-4-methoxy-2H-pyrrol-2-one, 9CI



(E)-form

C₁₃H₁₄N₂O₅ 278.264**(E)-form****Pukeleimide A**

[72362-15-3]

Metab. from the marine cyanophyte *Lyngbya majuscula*. Solid. λ_{max} 224 (ε 11200); 287 (ε 25200) (MeOH) (Derep).**Me ether: Pukeleimide G**

[72362-23-3]

C₁₄H₁₆N₂O₅ 292.291Metab. from *Lyngbya majuscula*. Solid.9,10-Dihydro, 10-hydroxy: **Pukeleimide D**

[72362-20-0]

C₁₃H₁₆N₂O₆ 296.279Metab. from *Lyngbya majuscula*. Solid.9,10-Dihydro, 10-methoxy: **Pukeleimide C**

[72362-19-7]

C₁₄H₁₈N₂O₆ 310.306Metab. from *Lyngbya majuscula*. Cryst.(CHCl₃/hexane or Me₂CO/hexane).Mp 186° (part. subl.). Racemic. λ_{max} 224 (ε 11200); 287 (ε 25200) (MeOH)

(Derep).

Δ^{10,13}-Isomer, 13-deoxy: **Pukeleimide E**

[72362-21-1]

C₁₃H₁₄N₂O₄ 262.265Metab. from *Lyngbya majuscula*. Solid.**(Z)-form****Pukeleimide B**


[72362-17-5]

Metab. from *Lyngbya majuscula*. Solid.**Me ether: Pukeleimide F**


[72362-22-2]

C₁₄H₁₆N₂O₅ 292.291Metab. from *Lyngbya majuscula*. Solid.Cardellina, J.H. *et al.*, *Tet. Lett.*, 1979, 2003;2007 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)James, G.D. *et al.*, *Tet. Lett.*, 1985, **26**, 3617(*synth, Pukeleimide A*)**P-782****Pulchellalactam**

1,5-Dihydro-4-methyl-5-(2-methylpropylidene)-2H-pyrrol-2-one

C₉H₁₃NO 151.208Isol. from the marine-derived fungus *Corollospora pulchella*. Inhibitor of CD45 protein tyrosine phosphatase. Oil. λ_{max} 276 (no solvent reported). λ_{max} 275 (MeOH) (Berdy).Alvi, K.A. *et al.*, *J. Antibiot.*, 1998, **51**, 515-517 (*isol, pmr, cmr*)Li, W.-R. *et al.*, *J.O.C.*, 2002, **67**, 4702-4706(*synth*)Bessho, J. *et al.*, *Heterocycles*, 2004, **63**, 1013-1016 (*synth*)Chavan, S.P. *et al.*, *Synth. Commun.*, 2007, **37**,1503-1510 (*synth, ir, pmr, cmr*)Felluga, F. *et al.*, *Synthesis*, 2007, 1882-1886(*synth*)**Pulchellidine†**

[23367-86-4]

C₂₀H₃₃NO₄ 351.485Sesquiterpene alkaloid from *Gaillardia pulchella* (Asteraceae). Cryst. (Me₂CO/Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 185-186°. [α]_D²⁵ -22.5 (c, 1.33 in EtOH).*Hydrobromide*: Mp 208-210°.*Di-Ac*: Mp 135-136°. [α]_D²⁷ -24 (c, 1 in CHCl₃).8,11-Diepimer: **Neopulchellidine**


[28513-27-1]

C₂₀H₃₃NO₄ 351.485Alkaloid from *Gaillardia pulchella* (Asteraceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 131-134°. [α]_D²¹ -13 (c, 1 in CHCl₃). The 11-config. is not clearly shown.Yanagita, M. *et al.*, *Tet. Lett.*, 1969, 2073;1970, 131 (*ir, pmr, isol, stereochem*)Sekita, T. *et al.*, *Tet. Lett.*, 1970, 135 (*cryst struct*)Yanagita, M. *et al.*, *Tet. Lett.*, 1970, 3007(*Neopulchellidine*)Inayama, S. *et al.*, *Heterocycles*, 1985, **23**, 377(*synth*)**Pulchelline†**

[1360-68-5]

C₁₈H₂₃NO₃ 301.385Struct. unknown. Alkaloid from leaves of *Neolitsea pulchella* (Lauraceae). Cryst. (Me₂O). Mp 189°. [α]_D +80.1 (c, 0.81 in**P-784**CHCl₃).Hui, W.H. *et al.*, *J.C.S.*, 1965, 2285-2286**Pulcherrimine****P-787**


4-[(2-Carboxy-2-hydroxyethyl)thio]-2-piperidinecarboxylic acid [309946-45-0]



Absolute Configuration

C₉H₁₅NO₅ 249.287Abs. config. revised in 2002. Isol. from the ovaries of the sea urchin, *Hemicentrotus pulcherrimus*. Amorph. powder. [α]_D²⁴ -16.5 (c, 0.2 in H₂O). All 8 stereoisomers have been synthesised.Murata, Y. *et al.*, *J. Agric. Food Chem.*, 2000,**48**, 5557-5560 (*isol, pmr, cmr, ms*)Sata, N.U. *et al.*, *Tet. Lett.*, 2002, **43**, 115-118(*synth, abs config*)**Pulcherrimic acid, 8CI****P-788**


3,6-Bis(2-methylpropyl)-2,5-pyrazinediol 1,4-dioxide, 9CI. 2,5-Dihydroxy-3,6-diisobutylpyrazine N,N-dioxide [957-86-8]

C₁₂H₂₀N₂O₄ 256.301Microbial metab. of *Bacillus*, *Micrococcus*, *Candida*, etc. spp. Found in soil, manure etc., usually as Pulcherrimine. Active against gram-positive bacteria. Pale-yellow cryst. or solid. Poorly sol. H₂O. Mp ca. 156-157° dec. (Mp varies with recryst. solv.). λ_{max} 243 ; 282 ; 410 (NaOH) (Berdy).*Fe complex: Pulcherrimine†*C₃₆H₆₀FeN₆O₁₂ 824.75

Formed by soil bacteria. Active against mycobacteria and fungi. Red solid. Dec. >200°.

MacDonald, J.C. *et al.*, *Can. J. Chem.*, 1963,**41**, 165 (*isol, struct, bibl, ir, pmr*)Ohta, A. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**,125 (*synth, struct*)Uffen, R.L. *et al.*, *J. Bacteriol.*, 1972, **111**, 86(*biosynth*)**Puloupone****P-789**

[97190-30-2]



C₂₁H₂₇NO 309.45

Probable abs. config. shown based on the tentative assignment to Isopuloupone. A minor metab. of the cephalaspidean mollusc *Philinopsis speciosa*. Oil. Sol. MeOH, hexane; poorly sol. H₂O. [α]_D²⁵ -10 (c, 0.20 in hexane). λ_{max} 206 (ε 10700); 257 (ε 3870); 263 (ε 4200); 269 (ε 3130) (EtOH) (Derep).

A^{1',2'}-Isomer: Isopuloupone

[149155-21-5]

C₂₁H₂₇NO 309.45

Isol. from the mollusc *Navanax inermis* and its prey *Bulla gouldiana*. Ichthyotoxin. [α]_D²⁵ -119 (c, 0.4 in hexane). λ_{max} 242 (ε 11400); 281 (ε 5400) (MeOH) (Berdy).

Coval, S.J. *et al.*, *J.O.C.*, 1985, **50**, 3024-3025 (isol, uv, ir, pmr, ms, struct)


Sugahara, T. *et al.*, *Tet. Lett.*, 1989, **30**, 1821 (synth, abs config)

Spinella, A. *et al.*, *Tetrahedron*, 1993, **49**, 3203 (*Isopulo'upone*)

Pulvinic acid**P-790**

α-(3-Hydroxy-5-oxo-4-phenyl-2(5H)-furan-2-ylidene)benzeneacetic acid, 9CI. Pulvic acid

[26548-70-9]

C₁₈H₁₂O₅ 308.29

Numbering systems vary. Pigment from various lichen spp. Orange powder (Et₂O or CHCl₃), orange prisms (C₆H₆). V. sol. EtOH; sol. H₂O; spar. sol. Et₂O, CHCl₃. Mp 216-217°. Forms solvates with MeOH, EtOH.

▶ HF3325000

Amide: Pulvinamide

[31673-63-9]


C₁₈H₁₃NO₄ 307.305

Isol. from *Pseudocyphellaria crocata*. Cryst. (CHCl₃). Mp 220-221° dec.

Maass, W.S.G. *et al.*, *Phytochemistry*, 1970, **9**, 2477 (*Pulvinamide*)

Pumiliotoxin A**P-791**

Octahydro-6-(6-hydroxy-2,5-dimethyl-4-octenylidene)-8-methyl-8-indolizidinol, 9CI. 8-Hydroxy-6-(6-hydroxy-2,5-dimethyl-4-octenylidene)-8-methylindolizidine. *Dendrobates Alkaloid 307A*. PTX-A [67054-00-6]



(6'R)-form

C₁₉H₃₃NO₂ 307.475

Consists of two isomers which appear to be C-6' epimers. A major alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates pumilio*; minor or trace constit. in *Dendrobates auratus*, *Dendrobates granuliferus*, *Dendrobates lehmanni*, *Dendrobates minutus*, *Dendrobates occul-tator*, *Dendrobates viridis* and *Dendro-bates* sp. (Colombia, Dendrobatidae). Potentiator of striated muscle contraction, affects Ca transport across plasma membranes. [α]_D²⁵ +22.7 (c, 1.0 in MeOH). [α]_D²⁵ +14.2 (c, 0.51 in CHCl₃) (2:1 mixt. of epimers).

▶ Highly toxic, LD₅₀ = 1 mg/kg (mice). NM2411000

(6'R)-form**Pumiliotoxin 307A'**

[α]_D²⁵ +14.3 (c, 0.74 in CHCl₃). Major natural isomer.

6'-Me ether: Pumiliotoxin 321C₂₀H₃₅NO₂ 321.502

Alkaloid from skin extracts of *Dendrobates pumilio* (Dendrobatidae). [α]_D²⁵ +21.3 (c, 1.0 in MeOH). A mixt. of (15R)- and (15S)-isomers. Prob. artifact.

4',5'-Dihydro, 6'-ketone: Pumiliotoxin 307FC₁₉H₃₃NO₂ 307.475

Alkaloid from skin extracts of *Dendrobates pumilio* (Dendrobatidae). [α]_D²⁵ -8.5 (c, 1.0 in MeOH). Possibly an artifact.

7β-Hydroxy: Allopumiliotoxin 323B'

[109279-26-7]

C₁₉H₃₃NO₃ 323.475

Minor or trace alkaloid in *Dendrobates abditus*, *Dendrobates leucomelas*, *Dendrobates pumilio*, *Melanophryniscus moreirae*, the Madagascan frogs *Mantella aurantiaca* and *Mantella madagascariensis* (Ranidae, subfamily Mantellinae) and the Australian frog *Pseudophryne semimarmorata* (Myobatrachidae). [α]_D²⁵ +22.3 (c, 1.0 in MeOH). Config. at C-15 unknown, but presumed to be (R-). An epimeric mixt. of Allopumiliotoxin 323B' and Allopumiliotoxin 323B'' was formerly called Allopumiliotoxin B.

7'S-Hydroxy: Pumiliotoxin B. Dendrobates Alkaloid 323A. PTX-B

[67016-65-3]

C₁₉H₃₃NO₃ 323.475

A major alkaloid in skin extracts of the neotropical poison-frogs *Dendrobates auratus*, *Dendrobates granuliferus*, *Dendrobates lehmanni*, *Dendrobates leucomelas*, *Dendrobates occul-tator*, *Dendrobates pumilio*, *Dendrobates viridis*; minor or trace constit. in *Dendrobates abditus*, *Dendrobates fulguritus*, *Dendrobates histrionicus*, and *Dendrobates minutus*, also from the Madagascan frog *Mantella aurantiaca* (Ranidae, subfamily Mantellinae). [α]_D²⁵ +19.8 (c, 2.28 in MeOH).

▶ RH2241000

7α, 7'S-Dihydroxy: Allopumiliotoxin 339AC₁₉H₃₃NO₄ 339.474

Minor alkaloid from *Dendrobates auratus* (Dendrobatidae), also from skin extracts of *Mantella aurantiaca* (Ranidae, subfamily Mantellinae). [α]_D²⁵ +29.4 (c, 1.0 in MeOH).

7β, 7'S-Dihydroxy: Allopumiliotoxin 339BC₁₉H₃₃NO₄ 339.474

Minor alkaloid from *Dendrobates auratus* (Dendrobatidae). [α]_D²⁵ +4.5 (c, 0.5 in MeOH).

(6'S)-form**Pumiliotoxin 307A''**

Alkaloid from *Dendrobates* spp. (Dendrobatidae): see under Pumiliotoxin 307A' above.

7β-Hydroxy: Allopumiliotoxin 323B''C₁₉H₃₃NO₃ 323.475

Source as cited for Allopumiliotoxin 323B'. [α]_D²⁵ +55 (c, 0.1 in MeOH).

Daly, J.W. *et al.*, *Science (Washington, D.C.)*, 1967, **156**, 970

Daly, J.W. *et al.*, *J.A.C.S.*, 1980, **102**, 830 (pmr, cmr, struct)

Tokuyama, T. *et al.*, *Tet. Lett.*, 1982, **23**, 2121 (config)

Overman, L.E. *et al.*, *Tet. Lett.*, 1982, **23**, 2355 (config)

Uemura, M. *et al.*, *Tet. Lett.*, 1982, **23**, 4369 (abs config, Pumiliotoxin B)

Overman, L.E. *et al.*, *J.A.C.S.*, 1984, **106**, 4192 (synth, ir, pmr, cmr, ms)

Tokuyama, T. *et al.*, *Tetrahedron*, 1984, **40**, 1183 (*Allopumiliotoxin 323B*)

Daly, J.W. *et al.*, *Toxicol.*, 1984, **22**, 905 (*Allopumiliotoxin 323B*)

Overman, L.E. *et al.*, *J.O.C.*, 1985, **50**, 3669 (synth, pmr)

Daly, J.W. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1986, **4**, 142 (rev, pharmacol)

Tokuyama, T. *et al.*, *Tetrahedron*, 1987, **43**, 643 (*Pumiliotoxins 307F and 321*)

Overman, L.E. *et al.*, *Tet. Lett.*, 1988, **29**, 901 (synth)

Trost, B.M. *et al.*, *J.A.C.S.*, 1989, **111**, 4988 (synth, *Allopumiliotoxin 339B*)

Overman, L.E. *et al.*, *J.A.C.S.*, 1992, **114**, 368 (synth, *Allopumiliotoxin 339A*)

Goldstein, S.W. *et al.*, *J.O.C.*, 1992, **57**, 1179 (synth, *Allopumiliotoxin 339B*)

Franklin, A.S. *et al.*, *Chem. Rev.*, 1996, **96**, 505-522 (rev, synth)

Hirashima, S. *et al.*, *J.A.C.S.*, 1999, **121**, 9873-9874 (synth)

Tang, X.-Q. *et al.*, *J.A.C.S.*, 2000, **122**, 6950-6954 (synth, *Allopumiliotoxin 339A*, *Allopumiliotoxin 339B*)


Kibayashi, C. *et al.*, *J. Organomet. Chem.*, 2002, **653**, 229-233 (synth)

Aoyagi, S. *et al.*, *J.O.C.*, 2002, **67**, 5517-5526 (synth)

Daly, J.W. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1556-1575 (rev)

Pumiliotoxin 341A**P-792**

Allopumiliotoxin 341A. *Dendrobates Alkaloid 341A* [227606-38-4]



Probable Absolute Configuration

C₁₉H₃₅NO₄ 341.49

Alkaloid from skin extracts of the Ecuadorian poison frog *Epipedobates tricolor*, also from *Dendrobates auratus*, *Dendrobates granuliferus* and *Dendrobates viridis*. Oil.

Hydroxy (?): **Dendrobates Alkaloid 357** [164324-99-6]

C₁₉H₃₅NO₅ 357.489

Minor alkaloid from skin extracts of *Dendrobates auratus* (Dendrobatidae). m/e 357 (3), 340 (8), 324 (8), 272 (4), 182 (20), 138 (10), 110 (20), 84 (20), 70 (100).

Stereoisomer (?): **Dendrobates Alkaloid 341B**

C₁₉H₃₅NO₄ 341.49

Trace alkaloid from skin extracts of *Dendrobates lehmanni* (Dendrobatidae). m/e 341 (1), 324 (4), 182 (60), 114 (20), 112 (20), 70 (100). CAS No. not found to 2008.

Daly, J.W. *et al.*, *Toxicol.*, 1978, **16**, 163-188 (isol)


Daly, J.W. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 205-340 (ms, rev)

Jain, P. *et al.*, *Heterocycles*, 1999, **50**, 903-912 (isol, ir, pmr, ms, struct)

Michael, J.P. *et al.*, *Nat. Prod. Rep.*, 2000, **17**, 579-602 (rev)

Pumiliotoxin 307B**P-793**

Octahydro-6-(4-hydroxy-2,5-dimethyl-5-octenyldene)-8-methyl-8-indolizininol, 9CI [151805-33-3]


C₁₉H₃₃NO₂ 307.475

Consists of two diastereomers, poss. allylic rearr. prods. of Pumiliotoxin A, P-791. Trace alkaloid from skin extracts of the Madagascan frogs *Mantella auran-tiaca* and *Mantella crocea*. Minor constit. of *Mantella viridis*.

Garraffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (isol, ir, ms, struct)

Pumiliotoxin C**P-794**

Decahydro-5-methyl-2-propylquinoline, 9CI. **Dendrobates Alkaloid 195A. PTX-C** [27766-71-8] [55785-29-0 ((±)-form)]



Absolute Configuration

C₁₃H₂₅N 195.347

A major alkaloid in skin extracts of the poison-frogs *Dendrobates pumilio* and *Dendrobates auratus*; trace constit. in *Dendrobates lehmanni* and *Dendrobates minutus* (Dendrobatidae). Liq. Mp 230-240° (as hydrochloride) (natural) Mp 277-278° (hydrochloride) (synthetic). Opt. rotn. not reported. The hydrochloride of the unnatural enantiomer had [α]_D +16.4 (MeOH). Rel. nontoxic, causes convulsions in v. large doses. Many syntheses reported. For a review of synthetic work to 1996, see Kibayashi *et al.*, 1997.

6β-Hydroxy: **Dendrobates Alkaloid 211A** [109175-45-3]

C₁₃H₂₅NO 211.347

Trace alkaloid from *Dendrobates pumilio* (Dendrobatidae). [α]_D -11.7 (c, 1.0 in CHCl₃).

Daly, J.W. *et al.*, *Annalen*, 1969, **729**, 198-204 (isol, cryst struct, ms, pmr)

Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 593; 1977, **60**, 48; 204-207 (synth, ir, pmr, ms, abs config)

Habermehl, G. *et al.*, *Annalen*, 1976, 1577-1583; 1977, 800-805 (synth, ir, pmr, ms, abs config)

Overman, L.E. *et al.*, *J.A.C.S.*, 1978, **100**, 5179-5185 (synth, ir, pmr, cmr, ms)

Tokuyama, T. *et al.*, *Tetrahedron*, 1987, **43**, 643-652 (6-hydroxy)

Kibayashi, C. *et al.*, *Stud. Nat. Prod. Chem.*, 1997, **19**, 3-88 (rev, synth)

Riechers, T. *et al.*, *Eur. J. Org. Chem.*, 1998, 2641-2646 (synth)

Back, T.G. *et al.*, *J.O.C.*, 1998, **63**, 6566-6571 (synth, bibl)

Weymann, M. *et al.*, *Tet. Lett.*, 1998, **39**, 7835-7838 (synth)

Spande, T.F. *et al.*, *J. Nat. Prod.*, 1999, **62**, 5-21 (pmr, cmr, occur)

Shieh, Y.-S. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2000, **47**, 283-284 (synth)

Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 2001, **84**, 141-145 (synth)

Akashi, M. *et al.*, *J.O.C.*, 2001, **66**, 7873-7874 (synth)

Toyooka, N. *et al.*, *J.O.C.*, 2002, **67**, 6078-6081 (synth)


Dijk, E.W. *et al.*, *Tetrahedron*, 2004, **60**, 9687-9693 (synth)

Girard, N. *et al.*, *Eur. J. Org. Chem.*, 2005, 2269-2280 (synth)

Holub, N. *et al.*, *Org. Lett.*, 2005, **7**, 1227-1229 (trans-form, synth)

Pumiliotoxin 209F**P-795**

Octahydro-8-methyl-6-(2-methylpropyldene)-8-indolizininol, 9CI [103190-29-0]

C₁₃H₂₃NO 209.331

Alkaloid from skin extracts of the Panamanian poison frog *Dendrobates pumilio*. [α]_D -11.6 (c, 0.10 in CHCl₃).

7β-Hydroxy: Octahydro-8-methyl-6-(2-methylpropyldene)-7,8-indolizinediol,

9CI. **Allopumiliotoxin 225E**

[141643-32-5]

C₁₃H₂₃NO₂ 225.33

From skin extracts of *Dendrobates pumilio*.

12-Hydroxy: Octahydro-6-(3-hydroxy-2-methylpropyldene)-8-methyl-8-indolizininol, 9CI. **Pumiliotoxin 225F**

[141671-16-1]

C₁₃H₂₃NO₂ 225.33

From skin extracts of *Dendrobates pumilio*. [α]_D -87.4 (c, 0.23 in CHCl₃).

Tokuyama, T. *et al.*, *Tetrahedron*, 1991, **47**, 5415 (isol, ir, cmr, ms, struct)

Hirashima, S. *et al.*, *J.A.C.S.*, 1999, **121**, 9873-9874 (*Pumiliotoxin 225F, synth*)


Aoyagi, S. *et al.*, *J.O.C.*, 2002, **67**, 5517-5526 (*Pumiliotoxin 225F, synth*)

Woodin, K.S. *et al.*, *J.O.C.*, 2007, **72**, 7451-7454 (synth)

Pumiloside**P-796**

[126722-26-7]

[126624-21-3]

C₂₆H₂₈N₂O₉ 512.515

Tautomeric with 4-quinolone form. Alkaloid from the whole plant of *Ophiorrhiza pumila* and the bark of *Nauclea orientalis*. Intermed. in biosynth. of Camptothecin, C-70. Prisms (MeOH). Mp 307-308°. [α]_D²² -39.8 (c, 0.15 in MeOH). λ_{max} 211; 241 (sh); 244; 303 (sh); 314; 317 (MeOH) (Derep).

7-Deoxy: **(3S)-Deoxypumiloside**

[126624-20-2]

C₂₆H₂₈N₂O₈ 496.516

Alkaloid from whole plant of *Ophiorrhiza pumila* (Rubiaceae). Plausible intermed. for biosynth. of Camptothecin, C-70. Amorph. powder. λ_{max} 206; 235; 293; 300; 306; 312; 320 (MeOH) (Derep).

7-Deoxy, 3-epimer: **(3R)-Deoxypumiloside**

[185020-37-5]

C₂₆H₂₈N₂O₈ 496.516

Alkaloid from *Ophiorrhiza pumila*.

Aimi, N. *et al.*, *Tet. Lett.*, 1989, **30**, 4991-4994 (isol, uv, pmr, synth, struct)

Kitajima, M. *et al.*, *Tet. Lett.*, 1997, **38**, 4255 (isol, struct, cd, deriv)

Zhang, Z. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1001-1005 (isol, pmr, cmr)

Kitajima, M. *et al.*, *Tetrahedron*, 2002, **58**, 9169-9178 (*Deoxypumilosides*)

Punarnavine**P-797**C₁₇H₂₂N₂O 270.374

Struct. unknown. Alkaloid from *Boer-*

haavia diffusa (Nyctaginaceae). Cryst. (C₆H₆). Mp 236-237° dec.

Sulfate: Mp 204-205°.


Picrate: Mp 114-115° dec.

Basu, N.K. *et al.*, *Q. J. Pharm. Pharmacol.*, 1947, **20**, 38-42 (*isol, bibl*)

Punctaneine

P-798

[145204-91-7]



C₂₀H₃₃NO₆ 383.484

Abs. stereochem. is tentative. Alkaloid from *Liatris punctata* (Asteraceae).

Yellow oil. $[\alpha]_D^{23}$ -2.5 (c, 0.4 in CHCl₃).

[145307-23-9]

Mead, E.W. *et al.*, *Phytochemistry*, 1992, **31**, 3255 (*isol, pmr, cmr, ms, struct*)

Punikathine

P-799

C₁₉H₂₃NO₅ 345.394


Struct. unknown. Amaryllidaceae alkaloid. Isol. from the *Haemanthus* hybrid "King Albert" (Amaryllidaceae). Cryst. (Me₂CO or EtOAc/Et₂O). Mp 177-178°. $[\alpha]_D^{22}$ -46 (c, 0.25 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129-1134 (*isol*)

Punjabine

P-800

[84435-36-9]



C₃₅H₃₂N₂O₇ 592.647

Minor alkaloid from the roots of *Berberis lycium* (Berberidaceae). Needles. $[\alpha]_D^{25}$ -40 (c, 0.48 in MeOH).

Me ether: O-Methylpunjabine

[58194-22-2]

C₃₆H₃₄N₂O₇ 606.674

Trace alkaloid from *Stephania sasaki* (Menispermaceae). Also isol. from leaves of *Anisocyclus jollyana*. Needles (Me₂CO/MeOH). Mp 144-146°. $[\alpha]_D$ -4.12 (c, 0.19 in CHCl₃).

α'-Carboxylic acid, Me ester: Gilgitine

[84435-37-0]

C₃₆H₃₄N₂O₈ 622.673

Minor alkaloid from the roots of *Berberis lycium* (Berberidaceae). Specific rotn. could not be detd. CD curve resembles that of Punjabine.

α'-Alcohol, O¹²-Me: Secojollyanine

[175097-23-1]

C₃₆H₃₆N₂O₇ 608.69

Alkaloid from leaves of *Anisocyclus*

jollyana. $[\alpha]_D^{20}$ -8.4 (c, 2.0 in CHCl₃).

The name Secojollyanine is misleading (Jollyanine is an iboga alkaloid).

α'-Deoxy, Me ether: O-Methyldeoxopunjabine

[89503-80-0]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from *Stephania sasaki* (Menispermaceae). Prisms + 1.5MeOH (MeOH or Me₂CO). Mp 183-185°.

Leet, J.E. *et al.*, *Heterocycles*, 1982, **19**, 2355 (*isol, uv, cd, ir, pmr, ms, struct*)


Kunimoto, J. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 135 (*O-Methylpunjabine, O-Methyldeoxopunjabine*)

Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1996, **59**, 498 (*Secojollyanine, O-Methylpunjabine*)

Puntarenine

P-801

5,8,9,15a-Tetrahydro-3,4-dimethoxy[1,3]-dioxolo[4,5-h]isoquino[1,2-b][3]benzazepine-6,15-dione, 9CI. 3,4-Dimethoxy-11,12-methylenedioxy-8,14-dioxohexahydroisohomoprotoberberine



C₂₁H₁₉NO₆ 381.384

(±)-*form* [85769-44-4]

Alkaloid from the stems and above-ground wood of *Berberis empetrifolia* and from *Berberis actinacantha* (Berberidaceae). Cryst. (MeOH). Mp 214-216° Mp 250-255°. Prob. an artifact.

Sepulveda-Boza, S. *et al.*, *Planta Med.*, 1983, **49**, 32 (*isol, ir, pmr, cmr, ms, cryst struct*)


Fajardo, V. *et al.*, *Tet. Lett.*, 1983, **24**, 155 (*uv, ir, pmr, cmr, ms, struct*)

Shamma, M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 398

Puqienine A

P-802

[848644-14-4]



C₂₈H₄₇NO₃ 445.684

Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles (Me₂CO). Mp 260-261°. $[\alpha]_D^{20}$ -31.1 (c, 0.2 in MeOH).

6-Ketone: Puqienine B

[848644-15-5]

C₂₈H₄₅NO₃ 443.668


Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles (Me₂CO). Mp 226-227°. $[\alpha]_D^{20}$ -31.2 (c, 0.1 in MeOH).

Jiang, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 264-267 (*isol, pmr, cmr, cryst struct*)

Puqienine C

P-803

[924889-52-1]



C₂₈H₄₅NO₄ 459.668


Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles (MeOH). Mp 291-293°. $[\alpha]_D^{20}$ +10.9 (c, 0.1 in MeOH). λ_{max} 250 (MeOH).

Jiang, Y. *et al.*, *Steroids*, 2006, **71**, 843-848 (*isol, pmr, cmr*)

Puqienine D

P-804

[924889-53-2]



C₂₈H₄₅NO₄ 459.668


Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles. Mp 275-277°. $[\alpha]_D^{20}$ -73.1 (c, 0.1 in MeOH).

Jiang, Y. *et al.*, *Steroids*, 2006, **71**, 843-848 (*isol, pmr, cmr*)

Puqienine E

P-805

[924889-54-3]



C₂₈H₄₃NO₃ 441.653


Alkaloid from the bulbs of *Fritillaria puqiensis*. Cubes (MeOH). Mp 235-237°. $[\alpha]_D^{20}$ -141.1 (c, 0.1 in MeOH). λ_{max} 305 (MeOH).


Jiang, Y. *et al.*, *Steroids*, 2006, **71**, 843-848 (*isol, pmr, cmr*)

Puqienine F

P-806

[906089-20-1]



C₂₈H₄₅NO₅ 475.667Alkaloid from the bulbs of *Fritillaria puqiensis*. Cryst. (MeOH). Mp 216-218°. [α]_D²⁰ -65.5 (c, 0.1 in MeOH).Li, H.-J. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 722-724 (*isol, pmr, cmr, cryst struct*)**Puqietinone****P-807****3-Hydroxy-28-methyl-16,28-secosolanidan-6-one, 9CI**
[133362-87-5]C₂₈H₄₇NO₂ 429.685Stereochem. revised in 1995. Struct. incorrect in abstract. Alkaloid from the bulbs of *Fritillaria puqiensis* (Liliaceae). Exhibits antitussive and antitumour activity. Mp 240-245°. [α]_D²⁰ +29.4 (c, 0.64 in CHCl₃).**3-O-β-D-Glucopyranoside: Puqietinonoside**

[848644-16-6]

C₃₄H₅₇NO₇ 591.827Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles (Me₂CO). Mp 327-328°. [α]_D²⁰ -22.7 (c, 0.1 in MeOH).**N-De-Me: N-Demethylpuqietinone**


[848616-39-7]

C₂₇H₄₅NO₂ 415.658Alkaloid from the bulbs of *Fritillaria puqiensis*. Needles (Me₂CO). Mp 285-286°. [α]_D²⁰ -10.4 (c, 0.1 in MeOH).**N-De-Me, 3-O-[β-D-glucopyranosyl-(1→4)-β-D-galactopyranoside]: Yibeinoside C**


[157536-48-6]

C₃₉H₆₅NO₁₂ 739.942Alkaloid from *Fritillaria pallidiflora*. Cryst. Mp 209-211°.**3-Ketone: Puqietinedione**


[924889-56-5]

C₂₈H₄₅NO₂ 427.669Alkaloid from the bulbs of *Fritillaria puqiensis* (Liliaceae). Needles (MeOH). Mp 281-283°. [α]_D²⁰ +30.1 (c, 0.1 in MeOH).Li, P. *et al.*, *Zhongguo Yaoke Daxue Xuebao*, 1990, **21**, 198-200; *CA*, **114**, 182054y (*isol*)Xu, Y. *et al.*, *Yaaxue Xuebao*, 1994, **29**, 200-203 (*Yibeinoside C*)Lin, G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1662-1667 (*config, cryst struct*)Jiang, Y. *et al.*, *J. Nat. Prod.*, 2005, **68**, 264-267 (*N-de-Me, Puqietinonoside*)Jiang, Y. *et al.*, *Steroids*, 2006, **71**, 843-848 (*Puqietinedione*)**PUR 1****P-808**C₂₅H₃₄N₂O₄ 426.555Constit. of mixed purees of onion and garlic (*Allium cepa* and *Allium sativum*). Purple powder. Mp 161-164°. λ_{max} 570 (ε 94000) (MeOH). λ_{max} 570 (ε 148000) (MeOH/HCl). λ_{max} 300 (ε 20000) (MeOH/NaOH).Imai, S. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 843-847 (*isol, uv, pmr, cmr*)**Purealidin C****P-809**


[138590-56-4]

C₂₃H₂₈Br₄N₄O₄ 744.115Alkaloid from the marine sponge *Psammmaplysilla purea*. Antifungal agent. Cytotoxic. Amorph. λ_{max} 210 (ε 23000); 285 (ε 1600) (MeOH) (Berdy).Kobayashi, J. *et al.*, *Tetrahedron*, 1991, **47**, 6617 (*isol, uv, ir, pmr, cmr, ms, struct*)**Purealidin D****P-810**

[143052-17-9]

C₂₂H₂₅Br₂N₆O₃[⊕] 581.286Alkaloid from the sponge *Psammmaplysilla purea*. Na[⊕] and K[⊕]-ATPase inhibitor. *Isol.* as the bis(trifluoroacetate) salt. λ_{max} 217 (ε 15300); 260 (ε 2900); 269 (sh); 286 (ε 800) (MeOH) (Derep).Tsuda, M. *et al.*, *Tet. Lett.*, 1992, **33**, 2597 (*isol, pmr, cmr*)**Purealidin H****P-811**

[164301-30-8]


C₁₄H₁₅Br₂N₅O₃ 461.112Alkaloid from the Okinawan sponge *Psammmaplysilla purea*. Oil. λ_{max} 277 (ε 1700); 284 (ε 1400) (MeOH).O⁴-(3-Aminopropyl): **Purealidin A**

[134850-51-4]

C₁₇H₂₂Br₂N₆O₃ 518.207Alkaloid from *Psammmaplysilla purea* and *Pseudoceratina verrucosa*. Cytotoxic, inhibitor of Na[⊕]/K[⊕] ATPase. Ichthyotoxic agent. Amorph. solid.O⁴-(3-Trimethylaminopropyl): **Purealidin E**

[145205-26-1]

[145205-28-3]

C₂₀H₂₉Br₂N₆O₃[⊕] 561.296Alkaloid from *Psammmaplysilla purea*. Na[⊕]/K[⊕] ATPase inhibitor. Amorph. solid (as bis(trifluoroacetate)). Sol. MeOH. λ_{max} 218 (ε 13600); 287 (ε 1000) (MeOH).Ishibashi, M. *et al.*, *Experientia*, 1991, **47**, 299 (*Purealidin A*)Tsuda, M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1325-1327 (*Purealidin E*)Kobayashi, J. *et al.*, *J. Nat. Prod.*, 1995, **58**, 467 (*Purealidin H*)**Purealidin J****P-812****Pseudoceratinine A**C₁₅H₁₇Br₂N₅O₄ 491.138

Pseudoceratinine A was the (-)-form.


(+)-formAlkaloid from the Okinawan marine sponge *Psammmaplysilla purea*. Shows moderate inhibitory activity against epidermal growth factor receptor kinase. Oil (as trifluoroacetate). Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²¹ +24 (c, 0.98 in MeOH) (trifluoroacetate). λ_{max} 277 (ε 1700); 284 (ε 1400) (MeOH) (Berdy).**12,13-Dihydro, 13-oxo (12RS-): Purealidin K**

[167394-76-5]

C₁₅H₁₇Br₂N₅O₅ 507.138Alkaloid from *Psammmaplysilla purea*. Shows mod. inhibition of epidermal growth factor receptor kinase. Oil (as trifluoroacetate). Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁴ +26 (c, 0.38 in MeOH) (trifluoroacetate). λ_{max} 231 (ε 6300); 284 (ε 2400) (MeOH) (Berdy).*Deamino*: see Aerophobin 1, A-154**(-)-form** [172723-35-2]Alkaloid from the sponge *Pseudoceratina verrucosa*. [α]_D -158 (c, 1 in MeOH) (as hydrochloride). Incorrect struct. assigned in CA. λ_{max} 220 (log ε 4.47); 284 (log ε 4.08) (MeOH).Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403 (*isol, uv, ir, pmr, cmr, cd, struct*)Benharref, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 177-180 (*Pseudoceratinine A*)

Puralidin L

[167394-77-6]

C₁₅H₂₁Br₂N₅O₄ 495.17

Alkaloid from the Okinawan marine sponge *Psammaphysilla purea* and from *Ailochroia crassa*. Oil (as trifluoroacetate). $[\alpha]_D^{24} +27$ (c, 0.18 in MeOH) (trifluoroacetate). Genus name sometimes misspelt Ailochroia.

11ξ-Hydroxy-: Caissarine A

[439132-04-4]


C₁₅H₂₁Br₂N₅O₅ 511.169

Isol. from the Brazilian sponge *Aplysina caissara*. Glassy solid. λ_{\max} 232 (ε 9100); 283 (ε 4250) (MeOH).

Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403-407 (*isol, uv, ir, pmr, cmr, cd, struct*)
 Saeki, B.M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 796-799; 2003, **66**, 1039 (*Caissarine A*)

Puralidin M**P-814**

N-[2-(2-Amino-1H-imidazol-4-yl)ethyl]-3,5-dibromo-2-hydroxy-α-(hydroxyimino)-4-methoxybenzenepropanamide, 9CI [167394-78-7]

C₁₅H₁₇Br₂N₅O₄ 491.138

Alkaloid from the Okinawan marine sponge *Psammaphysilla purea*. Oil (as trifluoroacetate).

Deamino: Puralidin N

[167394-79-8]


C₁₅H₁₆Br₂N₄O₄ 476.124

From *Psammaphysilla purea*. Exhibits cytotoxicity against murine lymphoma L1210 cells and human epidermoid carcinoma KB cells *in vitro*. Oil. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 235 (ε 2600); 290 (ε 2800) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403 (*isol, uv, ir, pmr, cmr, struct*)
 Boehlow, T.R. *et al.*, *J.O.C.*, 2001, **66**, 3111-3118 (*Puralidin N, synth*)

Puralidin O**P-815**

N-[4-[(Aminoiminomethyl)amino]butyl]-3,5-dibromo-2-hydroxy-α-(hydroxyimino)-4-methoxybenzenepropanamide [167394-80-1]


C₁₅H₂₁Br₂N₅O₄ 495.17

Alkaloid from the Okinawan marine sponge *Psammaphysilla purea*. Oil (as trifluoroacetate).

Kobayashi, J. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 403 (*isol, uv, ir, pmr, cmr, struct*)

Puraline

[100101-28-8]

C₂₇H₃₀Br₄N₇O₇[⊕] 884.193

Isol. from the Okinawan marine sponge *Psammaphysilla purea*. Na/K-ATPase inhibitor. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{\max} 276 (sh); 284; 292 (sh) (MeOH) (Derep). λ_{\max} 276 (ε 8); 284; 292 (MeOH) (Berdy).


Chloride:C₂₇H₃₀Br₄ClN₇O₇ 919.645

Amorph. solid. Mp 142-145°. $[\alpha]_D$ -85 (c, 2.1 in MeOH).

Nakamura, H. *et al.*, *Tet. Lett.*, 1985, **26**, 4517 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

2,6-Purinediamine, 9CI**P-817**

2,6-Diaminopurine, 8CI. 2-Aminoadenine [1904-98-9]

C₅H₆N₆ 150.143

Prod. by *Bacillus subtilis*. Biochemistry much studied. Mp 302° Mp 270°. λ_{\max} 220 (ε 24500); 246 (ε 7400); 280 (ε 9100) (H₂O).

► LD₅₀ (mus, ipr) 202 mg/kg. UO7524000

N¹-Oxide:C₅H₆N₆O 166.142

Cryst.

N²-Me: [7390-65-0]C₆H₈N₆ 164.169

Mp 300°.

N⁶-Me: [4421-04-9]C₆H₈N₆ 164.169

Mp 300°.

N², N²-Di-Me:C₇H₁₀N₆ 178.196

Mp 300°.

N², N⁶-Di-Me: [6333-66-0]C₇H₁₀N₆ 178.196

Mp 300°.

N⁶, N⁶-Di-Me: [5437-49-0]C₇H₁₀N₆ 178.196

Mp 300°.

N², N², N⁶-Tri-Me: [5426-14-2]C₈H₁₂N₆ 192.223

Mp 234° (block).

N², N⁶, N⁶-Tri-Me:C₈H₁₂N₆ 192.223

Mp 300°.

[7280-83-3]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 714B (*ir*)

Cavalieri, L.F. *et al.*, *J.A.C.S.*, 1948, **70**, 3875 (*uv*)

Robins, R.K. *et al.*, *J.A.C.S.*, 1953, **75**, 263 (*synth*)

Montgomery, J.A. *et al.*, *J.A.C.S.*, 1958, **80**, 404 (*derivs*)

Stevens, M.A. *et al.*, *J.A.C.S.*, 1958, **80**, 2755; 2759 (*synth, uv, cryst struct, oxide*)

Taylor, E.C. *et al.*, *J.A.C.S.*, 1959, **81**, 2442 (*synth, uv*)

Coburn, W.C. *et al.*, *J.O.C.*, 1965, **30**, 1110 (*pmr*)

Thorpe, M.C. *et al.*, *J. Magn. Reson.*, 1974, **15**, 98 (*cmr*)

Marumoto, R. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 759-774 (*CV 1808*)

Ger. Pat., 1975, ((*Abbott Lab*))2 460 553; *CA*, **83**1, 47702 (*synth*)

Chollet, A. *et al.*, *Chem. Scr.*, 1986, **26**, 37

Gereneer, R.Z. *et al.*, *Br. J. Pharmacol.*, 1992, **107**, 1048-1056 (*CV 1808, pharmacol*)

Florian, J. *et al.*, *J. Phys. Chem.*, 1992, **96**, 9278-9282 (*Raman, ir*)


Wang, Z. *et al.*, *Org. Lett.*, 2003, **5**, 2067-2070 (*uv*)

Virta, P. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 2924-2929 (*fluorescence*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, POJ500

6,8-Purinediol**P-818**

6,8-Dihydroxypurine. 7,9-Dihydro-1H-purine-6,8-dione



9H-form

7,9-Dihydro-1H-form

C₅H₄N₄O₂ 152.112

Complex tautom. 7,9-Dihydro-1H-form appears to predominate for parent compd. Formed by electrochemical oxidn. of 6-Purinol, P-820. Component of hen egg yolk. Non-species specific mammalian cell growth promoting agent. Needles (H₂O). Mp 350°. pK_{a1} 7.65; pK_{a2} 9.87 (20°, H₂O).

1H-7,9-Dihydro-form [13231-00-0]*1-Me*: [54638-26-5]C₆H₆N₄O₂ 166.139

Rectangular plates (H₂O). Mp 300° dec. pK_{a1} 1.8; pK_{a2} 8.2; pK_{a3} 13 (7-anion). Violet fluor. Other tautomers possible.

7-Me: [54638-28-7]

Rectangular plates (H₂O). Mp 300° dec. pK_{a1} 2; pK_{a2} 7.8; pK_{a3} 10.5 (1-anion). Violet fluor. Other tautomers possible.

9-Me: [54638-29-8]C₆H₆N₄O₂ 166.139

Solid. Mp 390° dec. (sealed tube). pK_{a1} 2; pK_{a2} 8.4; pK_{a3} 13. Violet fluor. Other tautomers possible.

1,7-Di-Me: [54638-36-7]

C₇H₈N₄O₂ 180.166

Rods (H₂O). Mp 300° dec. pK_{a1} 2;
pK_{a2} 8.5 (9-anion). Violet fluor.

1,9-Di-Me: [54638-30-1]

Needles (H₂O). Mp 300° dec. pK_{a1} 1.5;
pK_{a2} 8.5 (7-anion). Violet fluor.

7,9-Di-Me: [54638-31-2]

Needles (H₂O). Mp 300° dec. pK_{a1} 1.5;
pK_{a2} 9.2 (1-anion). Violet fluor.

1,7,9-Tri-Me: [54638-34-5]

C₈H₁₀N₄O₂ 194.193

Solid. Mp 229-230° (sealed tube). pK_a
0 (cation). Violet fluor.

1-(3-Oxobutyl): 7,9-Dihydro-1-(3-oxobu-
tyl)-1H-purine-6,8-dione. 1-(3-Oxobu-
tyl)-6,8-purinediol

C₉H₁₀N₄O₃ 222.203

Alkaloid from *Subergorgia suberosa*.
Powder. λ_{max} 212 ; 264 (MeOH).

9-(3-Oxobutyl): 7,9-Dihydro-9-(3-oxobu-
tyl)-1H-purine-6,8-dione. 9-(3-Oxobu-
tyl)-6,8-purinediol

C₉H₁₀N₄O₃ 222.203

Alkaloid from *Subergorgia suberosa*.
Powder. λ_{max} 212 ; 264 (MeOH).

3H-form [51953-35-6]

3-Me: [54638-27-6]

C₆H₆N₄O₂ 166.139

Plates (H₂O). Mp 300° dec. pK_{a1} 1.3;
pK_{a2} 9 (9-anion). Violet fluor.

3,7-Di-Me: [54638-37-8]

C₇H₈N₄O₂ 180.166

Prisms (H₂O). Mp 300° dec. pK_{a1} 1.5;
pK_{a2} 6 (9-anion). Violet fluor.

3,9-Di-Me: [54638-38-9]

Prisms (H₂O). Mp 300° dec. pK_{a1} 1.3;
pK_{a2} 9 (9-anion). Violet fluor.

3,7,9-Tri-Me: [54638-41-4]

C₈H₁₀N₄O₂ 194.193

Prisms (DMF). Mp 300° dec. pK_a 1
(cation).

Fischer, E. et al., *Ber.*, 1899, **32**, 250; 1897, **30**,
1846 (*synth, derivs*)

Albert, A. et al., *J.C.S.*, 1954, 2060 (*synth*)

Mason, S.F. et al., *J.C.S.*, 1954, 2071 (*uv*)

Robins, R.K. et al., *J.A.C.S.*, 1958, **80**, 6671

(*synth, uv*)

Rahat, M. et al., *J.C.S. Perkin 1*, 1974, 2229

(*synth, pmr, uv*)


Ogawa, K. et al., *Chem. Pharm. Bull.*, 1992,

40, 612 (*synth, uv, pmr*)

Qi, S. et al., *J. Nat. Prod.*, 2008, **71**, 716-718 (3-
oxobutylpurinediols)

Subergorgia suberosa Purine- imidazole alkaloid P-819

[1016169-23-5]



C₁₃H₁₃N₇O₃ 315.291


Alkaloid from *Subergorgia suberosa*.

Powder. λ_{max} 213 ; 264 ; 310 (MeOH).

Qi, S.-H. et al., *J. Nat. Prod.*, 2008, **71**, 716-718
(*isol, pmr, cmr*)

6-Purinol

1,7-Dihydro-6H-purin-6-one, 9CI. **Hy-
poxanthine**. *Sarcine*. *Sarkine*. 6-Hydroxy-
purine. 6(1H)-Purinone
[68-94-0]



C₅H₄N₄O 136.113

Many alternative tautomeric forms possible. Occurs widely in plant and animal tissue. Used as a 0.8% soln. in DMF for pptn. detn. of Ag, Bi, Pd. Needles. Mp 360°. pK_{a1} 1.91; pK_{a2} 8.96; pK_{a3} 12.18 (20°).

► Exp. teratogen (large doses). LD₅₀ (mus, ipr) 750 mg/kg. UP0791000

1-Amino: 1-Aminohypoxanthine. 6-Amino-6,7-dihydroimidazo[4,5-d]pyrimidin-7-one

C₅H₅N₅O 151.127

Cryst. (DMF). Mp 314°.

1-(4-Nitrobenzylideneamino):

Pale yellow solid. Mp 311-312° dec.

1,7-Dihydro-form

1-Oxide:

C₅H₄N₄O₂ 152.112

► Exp. carcinogen.

1-Me: 1,7-Dihydro-1-methyl-6H-purin-6-one, 9CI. 1-Methylhypoxanthine, 8CI

[1125-39-9]

C₆H₆N₄O 150.14

Cryst. (MeOH aq.). Mp 311-312°.

pK_{a1} 8.8.

7-Me: 1,7-Dihydro-7-methyl-6H-purin-6-one, 9CI. 7-Methylhypoxanthine

[1006-08-2]

C₆H₆N₄O 150.14

Needles (EtOH). Mp 355° dec Mp 356-357° dec. pK_{a1} 9.4.

1,7-Di-Me: 1,7-Dihydro-1,7-dimethyl-6H-purin-6-one, 9CI. 1,7-Dimethylhypoxanthine

[33155-83-8]

C₇H₈N₄O 164.166

Cryst. Mp 245-246° (243°).

1,9-Dihydro-form [51953-23-2]

7-Oxide: [118974-81-5]

C₅H₄N₄O₂ 152.112

Needles. Mp 300°.

9-β-D-Ribopyranosyl: 1,9-Dihydro-9-β-D-
ribofuranosyl-6H-purin-6-one, 9CI. 9-
β-D-Ribopyranosylhypoxanthine

[52678-79-2]

C₁₀H₁₂N₄O₅ 268.229

Prod. by *Streptomyces antibioticus*.

Cryst. (MeOH aq.). Mp 245-246° dec.

[α]_D²³ -37 (c, 1 in 0.1M NaOH). λ_{max}
252 (ε 13150) (NaOH aq.).

9-Me: 1,9-Dihydro-9-methyl-6H-purin-6-one. 9-Methylhypoxanthine

[875-31-0]

C₆H₆N₄O 150.14

Plates (H₂O). Mp 390° dec. pK_{a1} 10.3.

1,9-Di-Me: 1,9-Dihydro-1,9-dimethyl-6H-purin-6-one, 9CI. 1,9-Dimethylhypoxanthine

[20535-82-4]

C₇H₈N₄O 164.166

Isol. from the sponge *Spongosorites* sp.

Cryst. (EtOH). Mp 255-256°. λ_{max} 209
(log ε 3.7); 260 (log ε 3.4) (H₂O).

3,7-Dihydro-form Minor tautomer.

3-Me: 3,7-Dihydro-3-methyl-6H-purin-6-one, 9CI. 3-Methylhypoxanthine

[1006-11-7]

C₆H₆N₄O 150.14

Cryst. (H₂O). Mp 307-309° dec. pK_{a1}
8.4.

3,7-Di-Me: 3,7-Dihydro-3,7-dimethyl-6H-purin-6-one, 9CI. 3,7-Dimethylhypoxanthine

[19143-59-0]

C₇H₈N₄O 164.166

Cryst. Mp 242-245°.

3,9-Dihydro-form Minor tautomer.

3,9-Di-Me: 3,9-Dihydro-3,9-dimethyl-6H-purin-6-one, 9CI. 3,9-Dimethylhypoxanthine

C₇H₈N₄O 164.166

Cryst. (AcOH/C₆H₆). Mp 276-278°.

7,9-Dihydro-form

Imidazolium betaine form

7,9-Di-Me: 6-Hydroxy-7,9-dimethyl-7H-purinium hydroxide inner salt, 9CI

[5752-16-9]

C₇H₈N₄O 164.166

Cryst. (MeOH). Mp 330-332°.

OH,3H-form

3-Me: [3324-66-1]

C₆H₆N₄O 150.14

Needles (H₂O). Mp 162-163°. pK_a 5.7.

OH,7H-form

Me ether: 6-Methoxyhypurine

[1074-89-1]

C₆H₆N₄O 150.14

Needles (H₂O). Mp 262-263°. pK_{a1}
2.17; pK_{a2} 9.18 (20°).

Me ether, 7-Me: 6-Methoxy-7-methyl-7H-purine

[38917-24-7]

C₇H₈N₄O 164.166

Needles (C₆H₆/heptane). Mp 200°.

Benzyl ether: 6-Benzoyloxypurine

[57500-07-9]

C₁₂H₁₀N₄O 226.237

Cryst. Mp 171-172°.

OH,9H-form

Me ether, 9-Me: 6-Methoxy-7-methyl-9H-purine
[38917-25-8]
C₇H₈N₄O 164.166
Needles (C₆H₆/heptane). Mp 152-153°.

Ph ether: 6-Phenoxyurine
[66085-17-4]

C₁₁H₈N₄O 212.21
Cryst. (2-propanol). Mp 217-218°.

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 584A (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 709A (ir)

Fischer, E. *et al.*, *Ber.*, 1897, **30**, 2400-2415; 1898, **31**, 104-122; 431-446 (7-Me, 9-Me, synth)

Traube, W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1906, **244**, 11 (3-Me)

Krebs, E.G. *et al.*, *Arch. Biochem. Biophys.*, 1949, **24**, 49 (1,9-di-Me)

Robins, R.K. *et al.*, *J.A.C.S.*, 1957, **79**, 490 (synth, 9-Me)

Prasad, R.N. *et al.*, *J.A.C.S.*, 1957, **79**, 6401 (synth, 7-Me)

Jones, J.W. *et al.*, *J.A.C.S.*, 1962, **84**, 1914-1919 (7,9-dihydro 7,9-di-Me, 3,7-dihydro 3-Me, synth)

Elion, G.B. *et al.*, *J.O.C.*, 1962, **27**, 2478-2491 (3,7-dihydro 3-Me, 1,7-dihydro 1-Me, 1,9-dihydro 9-Me, 1,9-dihydro 1,9-di-Me, synth)

Bullock, F.I. *et al.*, *J.O.C.*, 1964, **29**, 1988 (pmr)
Montgomery, J.A. *et al.*, *J.O.C.*, 1965, **30**, 3235 (1-Me)

Neiman, Z. *et al.*, *Isr. J. Chem.*, 1966, **3**, 161 (3,9-di-Me)

Bergmann, F. *et al.*, *J.C.S. (C)*, 1966, 10 (synth)

Ovcharova, I.M. *et al.*, *CA*, 1968, **69**, 52099 (3,7-di-Me)

Deutsch, J. *et al.*, *Org. Mass Spectrom.*, 1970, **3**, 1219 (7-Me)

Lister, J.H. *et al.*, *Chem. Heterocycl. Compd.*, (Weissberger, A. *et al.*, Ed.), Part II, 1971, (rev)

Sekiya, M. *et al.*, *Chem. Pharm. Bull.*, 1972, **20**, 209; 1975, **23**, 2401 (synth, 1-Me, uv, pmr)

Lichtenberg, D. *et al.*, *Isr. J. Chem.*, 1972, **10**, 805 (nmr, uv, tautomer)

Karrer, W. *et al.*, *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 2553 (occur)

Marzilli, L.G. *et al.*, *J.A.C.S.*, 1975, **97**, 3351 (7-Me)

Chenon, M.T. *et al.*, *J.A.C.S.*, 1975, **97**, 4627; 4636 (cmr, tautomer)

Raurez, G. *et al.*, *Quim. Anal. (Madrid)*, 1975, **29**, 216 (use)

Keck, J.H. *et al.*, *J.O.C.*, 1978, **43**, 2587-2590 (Ph ether)

Brown, T. *et al.*, *J.C.S. Perkin 1*, 1979, 3107 (3-Me)

Kern, D.L. *et al.*, *J. Het. Chem.*, 1980, **17**, 461-463 (9-ribofuranosyl)

Scheiner, P. *et al.*, *J. Het. Chem.*, 1985, **22**, 1435 (synth, pmr, uv)

Schmalle, H.W. *et al.*, *Acta Cryst. C*, 1988, **44**, 732 (cryst struct)

Ogawa, K. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 612 (9-Me, 7-oxide)

Capon, R.J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 261-262 (isol, 1,9-Dimethylhypoxanthine)

Fridman, N. *et al.*, *Acta Cryst. C*, 2004, **60**, o44-o46 (Me ether, cryst struct)

Lagoja, I.M. *et al.*, *Chem. Biodiversity*, 2004, **1**, 106-111 (synth)

Wanner, M.J. *et al.*, *J. Med. Chem.*, 2004, **47**, 6875-6883 (benzyl ether)


Lauffer, S.A. *et al.*, *J. Med. Chem.*, 2005, **48**, 710-722 (Ph ether)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DMC000

Purpuramine A

P-821

N-[3-[4-(2-Aminoethyl)-2,6-dibromophenoxy]propyl]- α -(hydroxyimino)benzene-propanamide, 9CI
[149636-92-0]



C₂₀H₂₃Br₂N₃O₃ 513.228

Isol. from the sponge *Psammaphysilla purpurea*. λ_{\max} 280 (€ 834) (MeOH).

2-Bromo, 4-methoxy, N¹⁰-Me: **Purpuramine L**

C₂₂H₂₆Br₃N₃O₄ 636.177

Isol. from *Psammaphysilla purpurea*.

Amorph. solid. Mp 175-178°. λ_{\max} 214 (log € 5.12); 280 (log € 4.02) (MeOH).

14-Debromo: **Purpuramine B**

[149636-93-1]

C₂₀H₂₄BrN₃O₃ 434.332

Isol. from *Psammaphysilla purpurea*. λ_{\max} 280 (€ 641) (MeOH).

14-Debromo, 2,4-dibromo, 3-methoxy: **14-Debromopurpuraraplysillin I**

[136685-30-8]

C₂₁H₂₄Br₃N₃O₄ 622.15

Alkaloid from the sponge *Psammaphysilla purpurea*. Glass.

James, D.M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1137 (14-Debromopurpuraraplysillin I)


Yagi, H. *et al.*, *Tetrahedron*, 1993, **49**, 3749-3754 (Purpuramines A,B)

Goud, T.V. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 990-993 (Purpuramine L)

Purpuramine C

P-822

[149636-94-2]



C₂₅H₂₆Br₂N₃O₃[⊕] 576.307


Isol. from the sponge *Psammaphysilla purpurea*. λ_{\max} 250 (€ 5600) (MeOH).

Yagi, H. *et al.*, *Tetrahedron*, 1993, **49**, 3749-3754 (isol)

Purpuramine H

P-823

Aplysamine 3
[149636-99-7]



C₂₁H₂₄Br₃N₃O₄ 622.15

Metab. from the sponge *Psammaphysilla purpurea*. Cytotoxic agent. Amorph. or semicryst. solid. Isol. as a salt, counterion not specified. λ_{\max} 214 (€ 17000); 222 (€ 16700); 280 (€ 3100) (MeOH).

N²⁰-(13-Methyltetradecanoyl): **Aplysamine 5**

[152606-62-7]

C₃₆H₅₂Br₃N₃O₅ 846.536

Isol. from sponge *Psammaphysilla purpurea*. Cytotoxic agent. Gum. λ_{\max} 214 (€ 18500); 222 (€ 17200); 282 (€ 2700) (MeOH).

N²⁰-Me: **Purpuramine I**

[149637-00-3]

C₂₂H₂₆Br₃N₃O₄ 636.177

Isol. from *Psammaphysilla purpurea* and *Suberea* sp. λ_{\max} 280 (€ 1800) (MeOH).

N²⁰-Me, N²⁰-(methoxycarbonyl): **Purpuramine K**

[627544-34-7]

C₂₄H₂₈Br₃N₃O₆ 694.214

Isol. from *Psammaphysilla purpurea*. Amorph. solid. Mp 190-195°. λ_{\max} 218 (log € 3.65); 280 (log € 4.78) (MeOH).

N²⁰, N²⁰-Di-Me: **Aplysamine 2**

[172486-24-7]

[125547-39-9]

C₂₃H₂₈Br₃N₃O₄ 650.204

Isol. from *Aplysina* sp., *Psammaphysilla purpurea* and *Pseudoceratina purpurea*. Pale tan semicryst. solid (as hydrochloride). Mp 87-88.5° (hydrochloride). λ_{\max} 214 (€ 17000); 222 (€ 16700); 280 (€ 3100) (MeOH).

N²⁰, N²⁰-Di-Me, N²⁰-oxide: **Purpuramine J**

[478487-88-6]

C₂₃H₂₈Br₃N₃O₅ 666.203

Isol. from the Fijian sponge *Druinella* sp. Cytotoxic. Oil. λ_{\max} 280 (log € 3.26) (MeOH).

O-De-Me: **Purpuramine F**

[149636-97-5]

C₂₀H₂₂Br₃N₃O₄ 608.124

Isol. from *Psammaphysilla purpurea*. λ_{\max} 282 (€ 2600) (MeOH).

O-De-Me, N²⁰-Me: **Purpuramine G**

[149636-98-6]

C₂₁H₂₄Br₃N₃O₄ 622.15

Isol. from *Psammaphysilla purpurea*. λ_{\max} 282 (€ 2800) (MeOH).

4-Bromo: **Aplysamine 4**

[152606-61-6]

C₂₁H₂₃Br₄N₃O₄ 701.046

From *Psammaphysilla purpurea*. Cytotoxic agent. Semicryst. solid. Isol. as a salt, counterion not specified. λ_{\max} 218 (€ 17600); 222 (€ 17400); 274 (€ 2100) (MeOH).

2-Debromo, 3-demethoxy: **Purpuramine D**

[149636-95-3]

C₂₀H₂₃Br₂N₃O₃ 513.228

Isol. from *Psammaphysilla purpurea*. λ_{\max} 282 (€ 512) (MeOH).

2-Debromo, 3-demethoxy, N²⁰-Me: **Purpuramine E**

[149636-96-4]

C₂₁H₂₅Br₂N₃O₃ 527.255

Isol. from *Psammaphysilla purpurea*.

λ_{\max} 282 (ε 600) (MeOH).

14-Debromo: [159157-29-6]

$C_{21}H_{25}Br_2N_3O_4$ 543.254

Isol. from *Psammaphysilla purpurea*.

14-Debromo, 4-bromo: **14-Debromoaplysamine 4**

[159157-30-9]

$C_{21}H_{24}Br_3N_3O_4$ 622.15

Isol. from *Psammaphysilla purpurea*.

14-Debromo, 4-bromo, N²⁰-Me: **Purpurealidin H**

[799246-92-7]

$C_{22}H_{26}Br_3N_3O_4$ 636.177

Isol. from *Psammaphysilla purpurea*.

Amorph. solid (MeOH). λ_{\max} 218 (ε 12675); 280 (ε 2675) (MeOH).

Kynas, R. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1427-1433 (*Aplysamine 2*)

Yagi, H. *et al.*, *Tetrahedron*, 1989, **49**, 3749-3754 (*Purpuramines*)

Pakrashi, S.C. *et al.*, *Tetrahedron*, 1990, **50**, 12009-12014; 12783 (*isol*)

Jurek, J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1609-1612 (*Aplysamines 3-5*)

Rao, M.R. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 1301-1303 (*Aplysamine 2*)

Tabudravu, J.N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1798-1801 (*Purpuramine J*)

Goud, T.V. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 990-993 (*Purpuramine K*)


Tilvi, S. *et al.*, *Tetrahedron*, 2004, **60**, 10207-10215 (*Purpurealidin H*, 14-*Debromoaplysamine 4*)

Kijjoa, A. *et al.*, *Z. Naturforsch., B*, 2005, **60**, 904-908 (*Aplysamine 2*)

Purpurealidin B

P-824

[799246-87-0]



$C_{22}H_{24}Br_3N_3O_4$ 634.161


Isol. from the sponge *Psammaphysilla purpurea*. Amorph. solid. Mp 175.8°. λ_{\max} 283 (ε 1320) (MeOH).

Tilvi, S. *et al.*, *Tetrahedron*, 2004, **60**, 10207-10215 (*isol*, *pmr*, *cmr*, *ms*)

Purpurone

P-825

[147362-37-6]



$C_{40}H_{27}NO_{11}$ 697.653

Alkaloid from the marine sponge *Iotrochota* sp. Exhibits ATP-citrate lyase inhibitory activity. Lipogenesis inhibitor. Purple glass. λ_{\max} 215 (ε 30400); 315 (ε 16400); 511 (ε 8500) (MeOH/KOH) (Derep). λ_{\max} 216 (ε 28100); 296 (ε 15100); 500 (ε 11400) (MeOH) (Derep).

3''-Hydroxy: **Ningalin D**

[188111-70-8]

$C_{40}H_{27}NO_{12}$ 713.653

Alkaloid from the ascidian *Didemnum* sp. Dark red solid. λ_{\max} 206 (ε 41000); 276 (ε 18000); 294 (ε 17000); 404 (sh); 508 (ε 9900) (MeOH).

Chan, G.W. *et al.*, *J.O.C.*, 1993, **58**, 2544-2546 (*isol*, *uv*, *ir*, *pmr*, *cmr*)

Kang, H. *et al.*, *J.O.C.*, 1997, **62**, 3254-3262 (*Ningalin D*)

Peschko, C. *et al.*, *Tet. Lett.*, 2000, **41**, 9477-9481 (*synth*)

Hamasaki, A. *et al.*, *J.A.C.S.*, 2005, **127**, 10767-10770 (*Ningalin D*, *synth*)

Putreanine

P-826

N-(4-Aminobutyl)-β-alanine, 9CI. N-(4-Aminobutylamino)propanoic acid

[25887-39-2]

$H_2NCH_2CH_2CH_2CH_2NHCH_2CH_2CO-OH$

$C_7H_{16}N_2O_2$ 160.216

Amino acid present in mammalian brain. Cryst. + ½ H₂O (as sulfate salt). Mp 250-251° (sulfate).


Shiba, T. *et al.*, *Tetrahedron*, 1970, **26**, 4307 (*struct*, *synth*)

Anselini, C. *et al.*, *CA*, 1979, **90**, 39227n (*synth*)

Pycnanthine†

P-827

[19553-44-7]



Absolute configuration

$C_{40}H_{44}N_4O_2$ 612.813

Alkaloid from *Pleiocarpa pycnantha* var. *pycnantha* (Apocynaceae). Mp 250°. $[\alpha]_D^{25}$ +321 (c, 1 in CHCl₃).

14',15'-Dihydro: **Pleiomutinine**

[21400-49-7]

$C_{40}H_{46}N_4O_2$ 614.829

Alkaloid from *Pleiocarpa mutica*, *Hunteria congolana* and *Gonioma malagasy* (Apocynaceae). Mp 250°. $[\alpha]_D^{25}$ +274 (c, 0.44 in CHCl₃).

19'-Epimer, 14',15'-dihydro: **19'-Epipleiomutinine**

$C_{40}H_{46}N_4O_2$ 614.829

Isol. from trunkwood of *Hunteria congolana* (Apocynaceae). Amorph. $[\alpha]_D$ +188 (c, 1 in CHCl₃).

Gorman, A.A. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 33 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)


Rasoanaivo, P. *et al.*, *J.O.C.*, 1976, **41**, 376 (*cmr*, *struct*)

Vercauteren, J. *et al.*, *Bull. Soc. Chim. Fr., Part 2*, 1982, 291 (*Epipleiomutinine*)

Pycnanthinine, 9CI

P-828

[25480-38-0]



$C_{40}H_{46}N_4O_2$ 614.829

Minor alkaloid from the root bark of *Pleiocarpa pycnantha* (Apocynaceae). Noncryst. $[\alpha]_D^{25}$ +183 (c, 0.567 in CHCl₃).

Gorman, A.A. *et al.*, *Monatsh. Chem.*, 1967, **98**, 1554 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

Pycnarrhenamine

P-829

$C_{35}H_{40}N_2O_9$ 632.709

Bisbenzylisoquinoline alkaloid. Struct. unknown. Phenolic. Alkaloid from the roots of *Pycnarrhena manillensis* (Menispermaceae). Mp 203°. Phenolic.

v. Bruchhausen, F. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1960, **293**, 454-461 (*isol*)

Pycnarrhenine

P-830

$C_{36}H_{42}N_2O_9$ 646.736


Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the roots of *Pycnarrhena manillensis* (Menispermaceae). Mp 193°.

v. Bruchhausen, F. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1960, **293**, 454-461 (*isol*)

Pycnosanguin

P-831

[133056-31-2]




$C_{19}H_{20}N_2O_6$ 372.377

Isol. from *Pycnoporus sanguineus*. Yellow needles. Mp 246-249°.

Achenbach, H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 3 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *synth*, *ms*, *struct*)

Pyochelin I **P-832**

2-[4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolyl]-3-methyl-4-thiazolidinecarboxylic acid
[164104-31-8]
[79236-62-7, 164104-33-0, 164104-34-1]



$C_{14}H_{16}N_2O_3S_2$ 324.424

Prod. by *Pseudomonas aeruginosa* and *Burkholderia cepacia* (*Pseudomonas cepacia*). Siderophore. $[\alpha]_D^{25} +54.8$ (c, 0.4 in $CHCl_3$) (as Me ester).

2-Epimer: Pyochelin II

[164104-32-9]

$C_{14}H_{16}N_2O_3S_2$ 324.424

Prod. by *Pseudomonas aeruginosa*.

$[\alpha]_D^{25} -51.2$ (c, 1.2 in $CHCl_3$) (as Me ester).

[163581-05-3, 163581-04-2, 163581-03-1, 163581-02-0]

Rinehart, K.L. *et al.*, *J.O.C.*, 1995, **60**, 2786-2791 (*isol, synth, pmr, cmr, cryst struct, abs config*)


Zamri, A. *et al.*, *Tetrahedron*, 2000, **56**, 249-256 (*synth, bibl*)

Rivault, F. *et al.*, *Tetrahedron*, 2006, **62**, 2247-2254 (*synth*)

Schlegel, K. *et al.*, *Z. Naturforsch., C*, 2006, **61**, 263-266 (*Fe complex, cryst struct*)

Pyocyanine **P-833**

1-Hydroxy-5-methylphenazinium hydroxide inner salt, 9CI. 1,5-Dihydro-5-methyl-1-phenazinone. Cyanomycin. Sanasin. Antibiotic 4738 B
[85-66-5]



$C_{13}H_{10}N_2O$ 210.235

Phenazine antibiotic. Prod. by *Bacillus pyocyanus*, *Streptomyces cyanoflavus* and *Pseudomonas aeruginosa*. Increases respiration of living cells. Active against gram-positive and -negative bacteria. Blue needles (H_2O or $CHCl_3$ /petrol). Sol. $CHCl_3$, Me_2CO , hot H_2O , insol. Et_2O , C_6H_6 , petrol, CCl_4 , CS_2 . Mp 133°. pK_a 4.85 (30°). Can be kept. for several weeks *in vacuo* in the dark, but on longer standing dec. to 1-Phenazinol, P-327. λ_{max} 240 (€ 11600); 278 (€ 44900); 384 (€ 16400); 520 (€ 2460) (0.1N HCl) (Derep). λ_{max} 239 (€ 23500); 293 (€ 27300); 370 (€ 5570); 690 (0.1N NaOH) (Derep). λ_{max} 239 (€ 16000); 278 (€ 24800); 310 (€ 15300); 384 (€ 10500); 690 (€ 4300) (H_2O at pH 5) (Derep).

Perchlorate: [55171-01-2]

Dark-red needles (H_2O contg. $HClO_4$). Mp 221-223° dec.

Picrate:

Very dark-red leaflets with violet sheen + 2MeOH (MeOH), violet leaflets + 1EtOH (EtOH). Mp 194-195° dec.

Mellwain, H. *et al.*, *J.C.S.*, 1937, 1704 (*synth Org. Synth., Coll. Vol.*, 3, 1955, 753 (*synth*))

Swan, G.A. *et al.*, *Phenazines*, Interscience, N.Y., London, 1957, 174

Funaki, M. *et al.*, *J. Antibiot., Ser. A*, 1958, **11**, 143 (*isol*)

MacDonald, J.C. *et al.*, *Can. J. Microbiol.*,

1963, **9**, 809; 1966, **12**, 771 (*biosynth*)

Von Saltza, M. *et al.*, *J. Antibiot.*, 1969, **22**, 49 (*isol*)

Flood, M.E. *et al.*, *J.C.S. Perkin 1*, 1972, 622 (*biosynth*)

Knight, M. *et al.*, *Anal. Biochem.*, 1979, **95**, 19 (*synth*)

Baron, S.S. *et al.*, *Antimicrob. Agents*

Chemother., 1981, **20**, 814 (*props*)

Muller, M. *et al.*, *Free Radical Biol. Med.*,

2002, **33**, 1527-1533; 2006, **41**, 1670-1677 (*pharmacol*)

Allen, L. *et al.*, *J. Immunol.*, 2005, **174**, 3643-3649 (*pharmacol*)

Angell, S. *et al.*, *Chem. Biol.*, 2006, **13**, 1349-1359 (*marine, isol, pmr, cmr*)


Sax, N.I. *et al.*, *Dangerous Properties of*

Industrial Materials, 7th edn., Van Nostrand

Reinhold, 1989, 2322

Pyoluteorin **P-834**

(4,5-Dichloro-1H-pyrrol-2-yl) (2,6-dihydroxyphenyl)methanone, 9CI. 2,3-Dichloro-5-(2,6-dihydroxybenzoyl)pyrrole
[25683-07-2]



$C_{11}H_7Cl_2NO_3$ 272.087

Pyrrole antibiotic. Isol. from *Pseudomonas aeruginosa*, *Pseudomonas fluorescens* and a bacterial symbiont of *Sogatella furcifera*. Shows antibacterial, antifungal and herbicidal props. Yellow needles. Sol. MeOH, Et_2O ; fairly sol. acids; poorly sol. H_2O , hexane. Mp 174-175° dec. λ_{max} 257 (€ 12600); 309 (€ 39100) (MeOH/HCl) (Derep). λ_{max} 288 (€ 12000); 368 (€ 35500) (MeOH/NaOH) (Derep). λ_{max} 256 (€ 12600); 308 (€ 37400) (MeOH) (Derep). λ_{max} 255 (€ 4200); 310 (€ 13000) (EtOH) (Berdy). λ_{max} 282; 369 (MeOH-NaOH) (Berdy). λ_{max} 338 (€ 18400) (NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 45 mg/kg, LD₅₀ (mus, orl) 125 mg/kg. OB1804400

Di-O-Ac: Mp 204-206°.

N,O,O-Tri-Me:

Prisms (Et_2O). Mp 136-137°.

3'-Nitro: 3'-Nitropyloluteorin

[69093-95-4]

$C_{11}H_6Cl_2N_2O_5$ 317.084

Prod. by *Pseudomonas aeruginosa*.

Shows antibacterial, antifungal and herbicidal props. Yellow needles. Mp 228-230°. λ_{max} 313 (€ 33700) (MeOH/HCl) (Derep). λ_{max} 310 (€ 30100) (MeOH) (Derep). λ_{max} 330 (€ 28700);

400 (€ 34500) (MeOH/NaOH) (Derep).

Takeda, R. *et al.*, *J.A.C.S.*, 1958, **80**, 4749 (*isol, uv*)

Birch, A.J. *et al.*, *J.C.S.*, 1964, 2641 (*struct*)
Utsumi, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1970, **43**, 2640 (*cryst struct*)

Birchall, G.R. *et al.*, *Tet. Lett.*, 1970, 4879 (*synth, bibl*)

Bailey, D.M. *et al.*, *J. Med. Chem.*, 1973, **16**,

1298 (*synth, analogues*)

Ohmori, T. *et al.*, *Agric. Biol. Chem.*, 1978, **43**,

2031 (*isol, derivs*)


Cuppels, D.A. *et al.*, *Z. Naturforsch., C*, 1986,

41, 532 (*biosynth*)

Kenny, P.T.M. *et al.*, *Pestic. Sci.*, 1989, **27**, 117 (*isol*)

Pyoverdin I, 9CI

[114616-35-2]

P-835

$C_{56}H_{82}N_{16}O_{21}$ 1315.358

Cyclic depsipeptide antibiotic complex.

Isol. from *Pseudomonas fluorescens*.

Siderophore.

Poppe, K. *et al.*, *Tetrahedron*, 1987, **43**, 2261-2272 (*struct, bibl*)


Pyoverdin 2112**P-836**

Contains glycine/threonine/serine/glutamic acid/alanine/lysine in ratio 3:2:1:1:1:1, DL-threo-β-hydroxyaspartic acid and N^δ-hydroxyornithine. Prod. by *Pseudomonas fluorescens* 2112. Siderophore. Inhibits *Phytophthora capsici*, a red pepper blight-causing fungus.

Lee, E.-T. *et al.*, *J. Microb. Biotechnol.*, 2003, **13**, 415-421 (*isol, uv, activity*)

Pyoverdin 11370

[264189-47-1]

P-837

$C_{45}H_{62}N_{12}O_{22}$ 1123.052

Isol. from *Pseudomonas putida*.

Budzikiewicz, H. *et al.*, *Z. Naturforsch., C*, 1999, **54**, 1021-1026

Pyoverdin A1**P-838**


Reported aminoacid analysis cannot be correct. Prod. by *Pseudomonas putida* A1 isol. from a chickpea rhizosphere. Siderophore. Shows antifungal activity against phytopathogens. CAS no. not found to 2008.

Boopathi, E. et al., *Biochim. Biophys. Acta*, 1999, **1435**, 30-40 (*isol, cd, activity*)

Pyoverdins BTP 2

P-839

[367252-76-4]




C₄₃H₆₀N₁₂O₁₉ 1049.016
Isol. from *Pseudomonas fluorescens* BTP2. Siderophore.

Ongena, M. et al., *Tet. Lett.*, 2001, **42**, 5849-5851

Pyoverdins BTP7

P-840

[205181-21-1]



C₅₆H₈₄N₁₆O₂₄ 1365.372
Prod. by *Pseudomonas fluorescens* BTP7. Siderophore.

Amide:

C₅₆H₈₅N₁₇O₂₃ 1364.388
Prod. by *Pseudomonas fluorescens* BTP7. Siderophore.


Homologue (R = COOH):

C₅₇H₈₄N₁₆O₂₅ 1393.383
Prod. by *Pseudomonas fluorescens* BTP7.

Ongena, M. et al., *J. Mass Spectrom. Soc. Jpn.*, 1998, **46**, 53-56 (*isol, ms, struct*)

Pyoverdins BTP7G

P-841




C₅₇H₈₇N₁₇O₂₄ 1394.414
Stereochem. of Ser residues not determined. Prod. by *Pseudomonas fluorescens* BTP7. Siderophore.

Ongena, M. et al., *J. Mass Spectrom. Soc. Jpn.*, 1998, **46**, 53-56 (*isol, ms, struct*)

Pyoverdins BTP16G

P-842

[431982-43-3]



C₅₅H₈₁N₁₅O₂₈ 1400.329
Prod. by *Pseudomonas putida* BTP16. Siderophore.


Ongena, M. et al., *J. Mass Spectrom. Soc. Jpn.*, 1998, **46**, 53-56 (*isol, ms*)

Ongena, M. et al., *Lett. Pept. Sci.*, 2002, **8**, 21-27 (*isol*)

Pyoverdins C

P-843

Pyoverdins PaB
[104022-78-8]



C₅₆H₈₃N₁₇O₂₃ 1362.372
Prod. by *Pseudomonas aeruginosa* and *Pseudomonas fluorescens*. Siderophore. No phys. props. reported. Sol. H₂O; poorly sol. Me₂CO, hexane. λ_{max} 230 (ε 36300); 400 (ε 16980); 470 (ε 5495); 555 (ε 2512) (MeOH) (Berdy).

Wendenbaum, S. et al., *Tet. Lett.*, 1983, **24**, 4877-4880 (*isol*)

Hider, R.C. et al., *Struct. Bonding (Berlin)*, 1984, **58**, 25-87 (*rev*)


Briskot, G. et al., *Annalen*, 1989, 375-384 (*isol, struct*)

Budzikiewicz, A. et al., *Z. Naturforsch., C*, 1997, **52**, 713-720 (*rev*)

Pyoverdins CFBP 2392

P-844

[273378-29-3]




C₅₅H₈₂N₁₆O₂₂ 1319.347
Prod. by *Pseudomonas fluorescens* CFBP 2392. Siderophore.

Beiderbeck, H. et al., *BioMetals*, 1999, **12**, 331-338 (*isol, struct*)

Pyoverdins CFBP 2461

P-845

Pyoverdins WCS358
[701285-46-3]
[273378-27-1]




C₅₆H₈₄N₁₅O₂₄ 1351.366
Struct. revised in 2003. Metab. of *Pseudomonas putida* CFBP 2461. Siderophore. 3'-Hydroxy: [273378-28-2]
C₅₆H₈₄N₁₅O₂₃ 1367.365
Metab. of *Pseudomonas putida* CFBP 2461. Siderophore.

Beiderbeck, H. et al., *BioMetals*, 1999, **12**, 331-338 (*isol, pmr, cmr, ms*)
Fernandez, D.U. et al., *Monatsh. Chem.*, 2003, **134**, 1421-1431 (*struct*)

Pyoverdins CHAO

P-846

[178497-43-3]



C₅₅H₈₁N₁₅O₂₁ 1288.333
Cyclic depsipeptide. Prod. by *Pseudomonas fluorescens*. Siderophore.

4'-Amide: **Pyoverdins CHAO-A**


[178497-44-4]
C₅₅H₈₂N₁₆O₂₀ 1287.348
Prod. by *Pseudomonas fluorescens*. Siderophore.

Wong-Lung-Sang, S. et al., *Tet. Lett.*, 1996, **37**, 3329-3332 (*isol, pmr, cmr*)

Pyoverdins C₄₄H₆₁N₁₃O₁₇

P-847

[203261-53-4]



C₄₄H₆₁N₁₃O₁₇ 1044.043
Prod. by *Pseudomonas fluorescens* and *Pseudomonas putida*. Siderophore. λ_{max} 227 (ε 30771); 399 (ε 15551) (H₂O) (Berdy). λ_{max} 218 (ε 19904); 245 (ε 8526); 362 (ε 6516); 374 (ε 6244) (HCl) (Berdy).


4'-Amide: **Pyoverdins C₄₄H₆₂N₁₄O₁₆**

[203261-55-6]
C₄₄H₆₂N₁₄O₁₆ 1043.058
Prod. by *Pseudomonas fluorescens* and *Pseudomonas putida*. Siderophore. λ_{max} 227 (ε 30771); 399 (ε 15551)

(H₂O) (Berdy). λ_{\max} 218 (ϵ 19904); 245 (ϵ 8526); 362 (ϵ 6516); 374 (ϵ 6244) (HCl) (Berdy).

Budzikiewicz, H. *et al.*, *Z. Naturforsch., C*, 1997, **52**, 721-728 (*isol, pmr, cmr, ms*)

Pyoverdin C₄₅H₆₃N₁₁O₂₄ **P-848**
[393802-69-2]



C₄₅H₆₃N₁₁O₂₄ 1142.052

Prod. by *Pseudomonas syringae* and *Pseudomonas viridiflava*.

4'-Amide: Pyoverdin C₄₅H₆₄N₁₂O₂₃
[393802-71-6]

C₄₅H₆₄N₁₂O₂₃ 1141.067

Prod. by *Pseudomonas syringae*, *Pseudomonas cichorii* and *Pseudomonas viridiflava*.

(1⁷→3⁴)-Lactone, 4'-amide: [848934-20-3]

C₄₅H₆₂N₁₂O₂₂ 1123.052

Metab. of *Pseudomonas syringae*. Siderophore.

5-Gly analogue, (1⁷→3⁴)-lactone, 4'-amide: Pyoverdin
[848934-21-4]


C₄₄H₆₀N₁₂O₂₁ 1093.026

Metab. of *Pseudomonas syringae*. Stereochem. not fully shown. Has serine at residue 5 replaced by glycine.

Jülich, M. *et al.*, *Z. Naturforsch., C*, 2001, **56**, 687-694 (*isol, pmr, cmr, amide*)

Bultreys, A. *et al.*, *Z. Naturforsch., C*, 2004, **59**, 613-618 (*Pseudomonas syringae* constits)

Pyoverdin C₅₂H₇₅N₁₅O₂₃ **P-849**
[222841-35-2]



C₅₂H₇₅N₁₅O₂₃ 1278.251

Prod. by *Pseudomonas aureofaciens*. Siderophore. Prob. artifact. λ_{\max} 399 (pH 7.1). λ_{\max} 363 (pH 3.2 buffer).

4'-Amide: Pyoverdin C₅₂H₇₆N₁₆O₂₂
[222841-34-1]


C₅₂H₇₆N₁₆O₂₂ 1277.266

Prod. by *Pseudomonas aureofaciens*. Siderophore.

Beiderbeck, H. *et al.*, *Z. Naturforsch., C*, 1999,

54, 1-5

Pyoverdin C₅₅H₈₃N₁₇O₂₁ **P-850**
[278618-18-1]




C₅₅H₈₂N₁₇O₂₁⁺ 1317.354

Succinopyoverdin *isol.* from *Pseudomonas aeruginosa* ATCC 15692. Siderophore. λ_{\max} 414 (log ϵ 4.29) (pH 7.0). λ_{\max} 376 (log ϵ 4.18) (pH 3.0).

Lenz, C. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 146-152 (*isol, uv, pmr, cmr*)

Pyoverdin C₅₇H₈₅N₁₇O₂₄ **P-851**
[278618-20-5]



C₅₆H₈₃N₁₇O₂₄ 1378.371

Isol. from *Pseudomonas fluorescens*. Siderophore.

(4'→4)-Lactam: **Pyoverdin**

C₅₇H₈₅N₁₇O₂₃

[278618-19-2]

C₅₆H₈₂N₁₇O₂₃[±] 1361.364

Isol. from *Pseudomonas fluorescens*. Siderophore. λ_{\max} 400 (pH 7). λ_{\max} 374 (pH 3).

Lenz, C. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 146-152 (*Pseudomonas fluorescens* constits)

Pyoverdin D **P-852**
Pyoverdin PaA. Pyoverdin PaO1

[104022-79-9]

As Pyoverdin C, P-843 with

R = -COOH

C₅₅H₈₃N₁₇O₂₂ 1334.361

Prod. by *Pseudomonas aeruginosa* and *Pseudomonas fluorescens*. Siderophore. Sol. H₂O; poorly sol. Me₂CO, hexane. λ_{\max} 385; 400 (pH 7 buffer) (Berdy). λ_{\max} 365 (ϵ 14000); 380 (ϵ 14000) (pH 4 buffer) (Berdy).

Amide: Pyoverdin E. Pyoverdin Pa

[88966-86-3]

C₅₅H₈₄N₁₈O₂₁ 1333.377

From *Pseudomonas aeruginosa* and *Pseudomonas fluorescens*. Siderophore. Sol. H₂O; poorly sol. Me₂CO, hexane. λ_{\max} 385; 400 (pH 7 buffer) (Berdy). λ_{\max} 403 (ϵ 19000); 460; 540 (pH 4


buffer) (Berdy).

Wendenbaum, S. *et al.*, *Tet. Lett.*, 1983, **24**, 4877-4880 (*Pyoverdin Pa*)

Hider, R.C. *et al.*, *Struct. Bonding (Berlin)*, 1984, **58**, 25-87 (*rev*)

Briskot, G. *et al.*, *Annalen*, 1989, 375-384 (*isol, struct*)

Pyoverdin G 173 **P-853**
Pyoverdin Pf G 173
[535994-93-5]



C₄₉H₇₀N₁₄O₂₀ 1175.174

Isol. from *Pseudomonas fluorescens* G173. Siderophore.


3''-Hydroxy:

C₄₉H₇₀N₁₄O₂₁ 1191.173

Isol. from *Pseudomonas fluorescens* G173. Siderophore. CAS no. not found to 2004.

Fernández, D.U. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 1-10 (*isol, pmr, cmr, ms*)

Pyoverdin GM II **P-854**
[131647-40-0]



C₅₉H₈₆N₁₈O₂₄ 1431.435

Isol. from *Pseudomonas fluorescens*. Siderophore. Anthelmintic agent. Acaricide. Sol. H₂O. λ_{\max} 400 (ϵ 11750) (pH 7 buffer) (Berdy).

3'-Amide: Pyoverdin GM I


[131647-39-7]

C₅₉H₈₇N₁₉O₂₃ 1430.45

Isol. from *Pseudomonas fluorescens*. Siderophore. Anthelmintic agent. Acaricide. Sol. H₂O. λ_{\max} 400 (ϵ 11220) (pH 7 buffer) (Berdy).

Mohn, G. *et al.*, *Z. Naturforsch., B*, 1990, **45**, 1437-1450 (*isol, pmr, cmr, ms*)

Pyoverdin G4RA **P-855**
[197846-88-1]



C₄₄H₅₉N₁₃O₁₉ 1074.026
 Prod. by *Pseudomonas putida*. Siderophore. λ_{max} 365 ; 380 (pH 5 buffer).

4'-Amide: **Pyoverdinin G4R**
 [197846-90-5]

C₄₄H₆₀N₁₄O₁₈ 1073.041
 Prod. by *Pseudomonas putida*. Siderophore.

Salah el Din, A.L.M. *et al.*, *Tetrahedron*, 1997, **53**, 12539-12552 (*isol, uv, pmr, cmr, N-15 nmr, ms*)

Pyoverdinin II P-856

[114587-58-5]
 As Pyoverdinin I, P-835 with
 R = -COOH

C₅₅H₈₂N₁₆O₂₀ 1287.348
 Isol. from *Pseudomonas fluorescens*. Siderophore.


Amide: **Pyoverdinin III**

[114587-59-6]
 C₅₅H₈₃N₁₇O₁₉ 1286.363
 From *Pseudomonas fluorescens*. Siderophore.

Poppe, K. *et al.*, *Tetrahedron*, 1987, **43**, 2261-2272 (*struct, bibl*)

Pyoverdinin Pa II P-857

[137341-42-5]



C₄₈H₆₇N₁₅O₂₀ 1174.146
 Prod. by *Pseudomonas aeruginosa* strain R. Siderophore.

3'-Amide: **Pyoverdinin Pa I**


[137341-41-4]
 C₄₈H₆₈N₁₆O₁₉ 1173.161

Prod. by *Pseudomonas aeruginosa* strain R. Siderophore.

Gipp, S. *et al.*, *Z. Naturforsch., C*, 1991, **46**, 534-541

Pyoverdinin Pa TII P-858

[148337-19-3]



C₄₅H₆₅N₁₃O₁₉ 1092.084
 Prod. by *Pseudomonas aeruginosa*. Siderophore.

Tappe, R. *et al.*, *J. Prakt. Chem.*, 1993, **335**, 83-87 (*struct*)


Kinzel, O. *et al.*, *J. Antibiot.*, 1998, **51**, 499-507 (*synth*)

Pyoverdinin PaC P-859

Contains alanine, 2 β-hydroxyaspartic acids, 2 glycines, 2 threonines and an unknown amino acid. Prod. by *Pseudomonas asplenii* LMG2137. Siderophore. Bultreys, A. *et al.*, *Appl. Environ. Microbiol.*, 2003, **69**, 1143-1153 (*isol, props*)

Pyoverdinin Pf 1.3 P-860


[244021-59-8]



C₅₃H₇₅N₁₇O₂₁ 1286.277
 Isol. from *Pseudomonas fluorescens* 1.3. Siderophore. λ_{max} 410 (pH 6.8). λ_{max} 368 ; 379 (pH 3.0).

Georgias, H. *et al.*, *Z. Naturforsch., C*, 1999, **54**, 301-308


Pyoverdinin Pf/3 17400 P-861



C₅₂H₇₅N₁₇O₁₉ 1242.267
 Metab. of *Pseudomonas fluorescens* ATCC 17400. Siderophore. CAS no. not found to 2008.

Demange, P. *et al.*, *Tet. Lett.*, 1990, **31**, 7611-7614 (*isol, pmr, cmr, ms*)

Pyoverdinin Pf12 P-862



Illus. in ref. as the cations. Isol. from cultures of *Pseudomonas fluorescens*. Siderophore.

Pyoverdinin Pf12-IA [141073-88-3]

C₆₂H₉₂N₁₈O₂₇ 1521.513
 Sol. H₂O. λ_{max} 400 (H₂O) (Berdy). λ_{max}

470 ; 550 (H₂O/FeCl₃) (Berdy).

Pyoverdinin Pf12-IB [141073-89-4]

C₆₃H₉₂N₁₈O₂₈ 1549.524
 Sol. H₂O. λ_{max} 400 (H₂O) (Berdy). λ_{max} 470 ; 550 (H₂O/FeCl₃) (Berdy).

Pyoverdinin Pf12-IIA [141095-99-0]

C₆₂H₉₃N₁₉O₂₆ 1520.528
 Sol. H₂O. λ_{max} 400 (H₂O) (Berdy). λ_{max} 470 ; 550 (H₂O/FeCl₃) (Berdy).


Pyoverdinin Pf12-IIB [141073-90-7]

C₆₃H₉₅N₁₉O₂₇ 1550.555
 Sol. H₂O. λ_{max} 400 (H₂O) (Berdy). λ_{max} 470 ; 550 (H₂O/FeCl₃) (Berdy).

Geisen, K. *et al.*, *Monatsh. Chem.*, 1992, **123**, 151-178 (*isol, pmr, cmr, ms, struct*)

Pyoverdinin Pf 1547 P-863

[292840-54-1]




C₆₄H₉₇N₁₉O₂₆ 1548.582
 Prod. by *Pseudomonas* spp. Siderophore. λ_{max} 399 (pH 7). λ_{max} 363 ; 375 (pH 3 buffer).

Ruangviriyachai, C. *et al.*, *Z. Naturforsch., C*, 2000, **55**, 323-327

Pyoverdinin Pf CCM 2798 P-864

[133174-10-4]




C₄₈H₆₆N₁₆O₂₁ 1203.144
 Metab. of *Pseudomonas fluorescens* CCM 2798. Siderophore.

Demange, P. *et al.*, *Tet. Lett.*, 1990, **31**, 7611-7614 (*isol, pmr, cmr, ms*)

Pyoverdinin Pf P19 P-865

[280577-27-7]



C₄₆H₆₄N₁₂O₂₅ 1185.077

Prod. by *Pseudomonas fluorescens* P19. Siderophore.

3'-Deoxy: [280142-32-7]

C₄₆H₆₄N₁₂O₂₄ 1169.078


Prod. by *Pseudomonas fluorescens* P19. Siderophore.

Vossen, W. et al., *Z. Naturforsch., C*, 2000, **55**, 153-164

Pyoverdinin PL7

P-866

[422563-91-5]



C₄₄H₆₂N₁₂O₁₈ 1047.043

Prod. by *Pseudomonas fluorescens* PL7. Siderophore.

Barelmann, I. et al., *Z. Naturforsch., C*, 2002, **57**, 9-16 (isol, pmr, cmr)

Pyoverdinin PL8

P-867

Pyoverdinin C₄₇H₆₉N₁₃O₁₈

[422563-93-7]

As Pyoverdinin PL7, P-866 with

R¹ = CH₂OH, R² = -(CH₂)₃CH₂NH₂

C₄₇H₆₉N₁₃O₁₈ 1104.138

Prod. by *Pseudomonas fluorescens* PL8. Siderophore.

Barelmann, I. et al., *Z. Naturforsch., C*, 2002, **57**, 9-16 (isol, pmr, cmr)

Pyoverdinin Pm

P-868

[149316-40-5]


Contains β-hydroxyaspartic acid, lysine, N⁶-hydroxyornithine, 2 serines and 3 threonines. Prod. by *Pseudomonas putida* M (VKPMB-4308). Siderophore.

Maksimova, N.P. et al., *Mikrobiologiya*, 1994, **63**, 1038-1044; *Microbiology (Engl. Transl.)*, 1994, **63**, 587-590 (isol, uv)

Pyoverdinin Pp 1

P-869

[144447-59-6]



C₅₇H₈₃N₁₇O₂₅ 1406.382

Isol. from *Pseudomonas putida*. Siderophore. Sol. H₂O. λ_{max} 225 ; 297 ; 398 (ε 16600) (H₂O) (Berdy).

4'-Amide: Pyoverdinin Pp 2

[144447-60-9]

C₅₇H₈₄N₁₈O₂₄ 1405.397

Isol. from *Pseudomonas putida*. Siderophore. Sol. H₂O. λ_{max} 225 ; 297 ; 398 (ε 16600) (H₂O) (Berdy).

5,6-Dihydro, 4'-amide: Dihydropyoverdinin

Pp 2

[144447-61-0]

C₅₇H₈₆N₁₈O₂₄ 1407.413

Isol. from *Pseudomonas putida*. Siderophore. Sol. H₂O. λ_{max} 225 ; 297 ; 398 (ε 16600) (H₂O) (Berdy).


Gwose, I. et al., *Z. Naturforsch., C*, 1992, **47**, 487-502 (isol, pmr, cmr, ms)

Pyoverdinin PpC 3B

P-870

Pyoverdinin C 3B

[148714-91-4]



C₅₄H₇₈N₁₄O₂₈ 1371.287

Prod. by *Pseudomonas putida* C and *Pseudomonas putida* BTB16. Siderophore. Sol. H₂O, MeOH. λ_{max} 401 (pH 7 buffer) (Berdy). λ_{max} 363 ; 370 (pH 3 buffer) (Berdy).

4'-Amide: Pyoverdinin PpC 2B. Pyoverdinin C 2B. Pyoverdinin BTB 16

[148714-89-0]

C₅₄H₇₉N₁₅O₂₇ 1370.302

Prod. by *Pseudomonas putida* C and *Pseudomonas putida* BTB16. Siderophore. Sol. H₂O, MeOH. λ_{max} 401 (pH 7 buffer) (Berdy). λ_{max} 363 ; 370 (pH 3 buffer) (Berdy).

3'-Hydroxy: Pyoverdinin PpC 3A. Pyoverdinin C 3A

[148714-90-3]

C₅₄H₇₈N₁₄O₂₉ 1387.287

Prod. by *Pseudomonas putida* C. Siderophore. Sol. H₂O, MeOH. λ_{max} 401 (pH 7 buffer) (Berdy). λ_{max} 363 ; 370 (pH 3 buffer) (Berdy).

3'-Hydroxy, 4'-amide: Pyoverdinin PpC 2A. Pyoverdinin C 2A

[148714-83-4]

C₅₄H₇₉N₁₅O₂₈ 1386.302

Prod. by *Pseudomonas putida* C. Siderophore. Sol. H₂O, MeOH. λ_{max} 401 (pH 7 buffer) (Berdy). λ_{max} 363 ; 370 (pH 3 buffer) (Berdy).

Seinsche, D. et al., *J. Prakt. Chem.*, 1993, **335**, 157-168 (*Pseudomonas putida* C constits, isol, struct)

Ongena, M. et al., *J. Mass Spectrom. Soc. Jpn.*, 1998, **46**, 53-56 (isol, amide, ms)

Ongena, M. et al., *Lett. Pept. Sci.*, 2002, **8**, 21-27 (*Pseudomonas putida* BTB16 constits)

Pyoverdinin PS

P-871


Contains 3 threonines, 3 serines, 1 lysine and δ-N-hydroxyornithine. Prod. by *Pseudomonas syringae*. CAS no. not found to 2008.

Torres, L. et al., *Appl. Environ. Microbiol.*, 1986, **52**, 157-160 (isol)

Pyoverdinin PS 6.10

P-872

[948556-39-6]



C₄₆H₆₉N₁₃O₁₈ 1092.127

Prod. by *Pseudomonas* sp. PS 6.10.

4'-Amide: [948556-38-5]

C₄₆H₇₀N₁₄O₁₇ 1091.143

Prod. by *Pseudomonas* PS 6.10.

Budzikiewicz, H. et al., *Z. Naturforsch., C*, 2006, **61**, 815-820 (isol, pmr, cmr, ms)

Pyoverdinin PSS

P-873

[110737-08-1]


Contains 2 β-hydroxyaspartic acids, 1 lysine, 2 serines and 2 threonines. Prod. by *Pseudomonas syringae* pv *syringae*. Siderophore.

Cody, Y.S. et al., *Appl. Environ. Microbiol.*, 1987, **53**, 928-934 (isol, props)

Pyoverdinin R'

P-874

[403806-52-0]



C₄₃H₅₉N₁₃O₁₈ 1046.015

Prod. by *Pseudomonas aeruginosa* strain R'.

4'-Amide: Pyoverdinin C₄₃H₆₀N₁₄O₁₇

C₄₃H₆₀N₁₄O₁₇ 1045.031

Prod. by *Pseudomonas aeruginosa* strain R'. CAS no. not found to 2008.

Homologue (R = COOH): Pyoverdinin

C₄₄H₅₉N₁₃O₁₉

C₄₄H₅₉N₁₃O₁₉ 1074.026

Prod. by *Pseudomonas aeruginosa* strain R'. CAS no. not found to 2008.


Ruangviriyachai, C. et al., *Z. Naturforsch., C*, 2001, **56**, 933-938

Pyoverdinin 51WA

P-875

Pyoverdinin 51W-Suc

[235101-94-7]



C₅₆H₈₁N₁₇O₂₄ 1376.355
 Isol. from *Pseudomonas fluorescens* strain 51W. Siderophore.

3'-Amide: **Pyoverdin 51W**. Pyoverdin 51W-Suca
 [235101-93-6]


C₅₆H₈₂N₁₈O₂₃ 1375.371
 Isol. from *Pseudomonas fluorescens* strain 51W. Siderophore.

Dallakian, P. et al., *Chirality*, 1999, **11**, 381-386 (abs config)

Voss, J. et al., *Z. Naturforsch., C*, 1999, **54**, 156-162 (isol, struct)

Pyracrimycin A P-876

3-(3,4-Dihydro-2H-pyrrol-5-yl)-2-propenamamide, 9CI. 3-(1-Pyrrolin-2-yl)acrylamide. Cyclamidomycin. Desdanine. U 22662. Antibiotic U 22662



C₇H₁₀N₂O 138.169
 λ_{max} 237 (ε 16900) (0.1N HCl) (Derep).
 λ_{max} 235 (ε 20100) (0.1N NaOH) (Derep).
 λ_{max} 235 (ε 24000) (MeOH) (Derep).

(E)-form [35663-85-5]

Isol. from *Streptomyces* spp. Shows antibiotic props. Cryst. (MeOH). Mp 215-217° dec.

► LD₅₀ (mus, orl) 240 mg/kg. UC6320000

1,2-Dihydro: 3-(2-Pyrrolinyl)-2-propenamamide, 9CI. **Dihydropyracrimycin A**. 2-Pyrrolidine-2-acrylamide
 [43043-23-8]

C₇H₁₂N₂O 140.185
 From *Streptomyces eridani*. Sol. H₂O, DMSO, DMF; fairly sol. MeOH, acids; poorly sol. butanol, hexane.

Coronelli, C. et al., *J. Antibiot.*, 1971, **24**, 491; 497 (isol, struct)


Takahashi, S. et al., *J. Antibiot.*, 1971, **24**, 902 (isol)

Argoudelis, A.D. et al., *J. Antibiot.*, 1972, **25**, 432 (uv, pmr, ms, struct, bibl)

Martin, W. et al., *Tet. Lett.*, 1973, 1459 (synth)
 Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, COV125

Pyracrimycin B P-877

3-(3,4-Dihydro-2H-pyrrol-5-yl)-2-hydroxy-2-propenamamide, 9CI



C₇H₁₀N₂O₂ 154.168
 Exists as zwitterion protonated on the pyrrole N.

(Z)-form [12688-72-1]


Isol. from *Streptomyces eridani*. Shows antibiotic props. Pale-yellow cryst. (CHCl₃). Mp 222-224°. λ_{max} 322 (ε 16000) (MeOH) (Berdy). λ_{max} 292 (ε

7000) (MeOH-HCl) (Berdy).

Coronelli, C. et al., *J. Antibiot.*, 1971, **24**, 491 (isol)

Gallo, G.G. et al., *Gazz. Chim. Ital.*, 1975, **105**, 51 (struct)

Pyramidaglaine A P-878



C₄₀H₄₂N₂O₉ 694.78

Flavaglaine compd. Alkaloid from *Aglaia andamanica*. Needles (MeOH). Mp 135-137°. [α]_D²⁰ -117 (c, 0.1 in MeOH). λ_{max} 215 (ε 12850); 270 (ε 420) (MeOH).

O-De-Ac: Isofoveoglin

[948312-50-3]
 C₃₈H₄₀N₂O₈ 652.743
 Alkaloid from the leaves of *Aglaia foveolata*. Amorph. powder. [α]_D²⁰ +10 (c, 0.3 in CHCl₃). λ_{max} 203 (log ε 4.72); 214 (sh) (log ε 4.63) (MeOH).


3,10-Diepimer: Pyramidaglaine B

C₄₀H₄₂N₂O₉ 694.78
 Alkaloid from *Aglaia andamanica*. Needles (MeOH). Mp 146-148°. [α]_D²⁰ +22.8 (c, 0.1 in MeOH). λ_{max} 215 (ε 11110); 270 (ε 1740) (MeOH).

Puripattanavong, J. et al., *Planta Med.*, 2000, **66**, 740-745 (isol, pmr, cmr)
 Salim, A.A. et al., *Tetrahedron*, 2007, **63**, 7926-7934 (Isofoveoglin)

Pyranofoline P-879

2,11-Dihydro-5,10-dihydroxy-12-methoxy-2,2,11-trimethyl-6H-pyrano[3,2-b]acridin-6-one
 [82644-79-9]




C₂₀H₁₉NO₅ 353.374
 Alkaloid from the root and stem bark of *Glycosmis citrifolia* (Rutaceae). Orange needles (Me₂CO). Mp 212-214°.

10-Methoxymethyl ether:
 Orange needles (Et₂O). Mp 112-114°.

Wu, T.-S. et al., *Heterocycles*, 1982, **19**, 1227; *J.C.S. Perkin 1*, 1983, 1681 (isol, uv, ir, ms, pmr, cmr, struct)

Pyranonigrin A P-880



Absolute Configuration


C₁₀H₉NO₅ 223.185
 Struct. revised in 2007. Prod. by the marine-derived *Aspergillus niger* LL-LV3020 and the terrestrial *Aspergillus niger* LL-RB17. Amorph. solid. [α]_D²⁵ +84.4 (c, 1 in DMSO). [α]_D²⁵ +38.2 (c, 1 in MeOH). λ_{max} 210 (ε 19400); 250 (ε 9900); 314 (ε 13200) (MeOH). λ_{max} 208; 250; 312 (MeCN aq.).

1-Deoxy: Pyranonigrin S

C₁₀H₉NO₄ 207.185
 Prod. by *Aspergillus niger* LL-LV3020. λ_{max} 210 (ε 18000); 250 (ε 8100); 312 (ε 13300) (MeCN aq.).

Hiort, J. et al., *J. Nat. Prod.*, 2004, **67**, 1532-1543 (isol, pmr, cmr)
 Schlingmann, G. et al., *J. Nat. Prod.*, 2007, **70**, 1180-1187 (isol, cd, uv, pmr, cmr, ms)

Pyranonigrin B P-881




C₁₁H₁₁NO₆ 253.211
 Prod. by *Aspergillus niger* isol. from the sponge *Axinella damicornis*. Amorph. solid.

O-De-Me, 2-Me ether: Pyranonigrin C

C₁₁H₁₁NO₆ 253.211
 Prod. by *Aspergillus niger* from *Axinella damicornis*. Amorph. solid.

Hiort, J. et al., *J. Nat. Prod.*, 2004, **67**, 1532-1543 (isol, pmr, cmr)

Pyranonigrin D P-882




C₁₁H₉NO₅ 235.196
 Prod. by *Aspergillus niger* isol. from the sponge *Axinella damicornis*. Amorph. solid.

Hiort, J. et al., *J. Nat. Prod.*, 2004, **67**, 1532-1543 (isol, pmr, cmr)

Pyrayafoline A P-883

3,11-Dihydro-9-methoxy-3,3,8-trimethylpyrano[3,2-a]carbazole, 9CI
 [104778-00-9]



C₁₉H₁₉NO₂ 293.365
 Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Plates (Et₂O). Mp 228-231°. λ_{max} 222; 239; 286 (sh);

295 ; 334 (MeOH).

O-De-Me: Pyrayafoline C

[138822-92-1]

C₁₈H₁₇NO₂ 279.338

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Pale yellow oil. λ_{max} 227 ; 238 ; 286 (sh) ; 296 ; 323 ; 351 (MeOH).

Demethoxy, 4-methoxy: 3,11-Dihydro-5-methoxy-3,3,8-trimethylpyrano[3,2-a]carbazole, 9CI. Heptazolidine
[53424-58-1]

C₁₉H₁₉NO₂ 293.365

Alkaloid from *Clausena heptaphylla* (Rutaceae). Cryst. (MeOH). Mp 145°. λ_{max} 230 (log ε 4.68); 238 (log ε 4.86); 278 (log ε 4.29); 282 (log ε 4.49); 335 (log ε 3.7) (EtOH).

Chakraborty, D.P. *et al.*, *Chem. Ind. (London)*, 1974, 303 (Heptazolidine)


Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1986, 34, 2672-2675 (synth, uv, ir, pmr, ms, struct)

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, 39, 1668-1671 (Pyrayafoline C)

Pyrayafoline B

P-884

[138822-91-0]



C₁₈H₁₇NO₂ 279.338

Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Shows cytotoxic activity. Brown powder.


Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, 39, 1668 (isol, uv, ir, pmr, ms, synth, struct)

Itoigawa, M. *et al.*, *J. Nat. Prod.*, 2000, 63, 893-897 (activity)

Pyrayafoline D

P-885

[138822-93-2]



C₂₄H₂₇NO₂ 361.483


Alkaloid from the stem bark of *Murraya euchrestifolia*. Pale brown powder. λ_{max} 222 ; 238 ; 288 (sh) ; 296 ; 331 ; 342 (MeOH).

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1991, 39, 1668-1671

Pyrayaquinone A

P-886

[96861-90-4]



C₁₈H₁₅NO₃ 293.321

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Mp 222° dec.


Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, 33, 1320 (isol, uv, ir, pmr, ms, struct, synth)

Yogo, M. *et al.*, *Chem. Pharm. Bull.*, 1991, 39, 328 (synth)

Pyrayaquinone B

P-887

[96861-91-5]



C₁₈H₁₅NO₃ 293.321

Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Mp 244° dec.

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1985, 33, 1320 (isol, uv, ir, pmr, ms, struct, synth)


Ramesh, K. *et al.*, *Chem. Ind. (London)*, 1986, 614 (synth)

Yogo, M. *et al.*, *Chem. Pharm. Bull.*, 1991, 39, 328 (synth)

Pyrayaquinone C

P-888

3,11-Dihydro-3,8-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-a]carbazole-7,10-dione, 9CI
[117591-99-8]



C₂₃H₂₃NO₃ 361.44


Alkaloid from the root bark of *Murraya euchrestifolia* (Rutaceae). Dark-violet prisms (Me₂CO). Mp 223° dec.

Ito, C. *et al.*, *Chem. Pharm. Bull.*, 1988, 36, 2377 (isol, uv, ir, pmr, ms, struct)

Pyrazinol, 8CI

P-889

2-Hydroxypyrazine. 2(1H)-Pyrazinone



C₄H₄N₂O 96.088

NH-form predominates. Cryst. (C₆H₆). Mp 185-186°. pK_{a1} -0.1; pK_{a2} 8.25 (20°, H₂SO₄ aq.).

***OH*-form**

Me ether: Methoxypyrazine. FEMA 3302 [3149-28-8]

C₅H₆N₂O 110.115

Flavouring agent. Bp₂₉ 60-61° (Bp₂₈ 68°). pK_a 0.75. Odour threshold 700 ppb in H₂O.

►Skin and eye irritant.

Me ether, 1-oxide: [32046-05-2]

C₅H₆N₂O₂ 126.115

Mp 159-160.5°.

Me ether, 4-oxide: [23902-69-4]

C₅H₆N₂O₂ 126.115

Mp 80-81.5°.

Et ether: Ethoxypyrazine

[38028-67-0]

C₆H₈N₂O 124.142

Bp₉₀ 90-92°.

Ph ether: 2-Phenoxypyrazine

[107697-82-5]

C₁₀H₈N₂O 172.186

Mp 50-51°.

***NH*-form** [6270-63-9]

1-N-Me:

C₃H₆N₂O 110.115

Cryst. (petrol). Mp 83-84°.

4-Oxide: Emimycin

[3735-46-4]

C₄H₄N₂O₂ 112.088

Prod. by *Streptomyces griseochromogenes*, *Streptomyces antibioticus* 2020i,

Bacillus sp. Nk84-0218 and *Bacillus megatherium*. Weakly active against

gram-positive and -negative bacteria

and fungi. Anticoccidial agent and

acaricide. Nucleic acid antimetabolite.

Cryst. (EtOH/Et₂O). Sol. H₂O, AcOH,

MeOH; fairly sol. EtOH, Me₂CO;

poorly sol. EtOAc, hexane. Mp 245-

250° dec. λ_{max} 222 (E1%/1cm 1545);

276 (E1%/1cm 610); 331 (E1%/1cm

393) (MeOH) (Berdy). λ_{max} 220 (E1%/

1cm 1955); 272 (E1%/1cm 800); 328

(E1%/1cm 460) (HCl) (Berdy). λ_{max}

229 (E1%/1cm 2410); 256 (E1%/1cm

831); 330 (E1%/1cm 569) (NaOH)

(Berdy).

►LD₅₀ (mus, ipr) 1000-2000 mg/kg.

UQ4385000

Aldrich Library of 13C and 1H FT NMR

Spectra, 1992, 3, 403B (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase,

1989, 3, 1559D (ir)

Erickson, A.E. *et al.*, *J.A.C.S.*, 1946, 68, 400

(synth)

Karmas, G. *et al.*, *J.A.C.S.*, 1952, 74, 1580

(synth)

Albert, A. *et al.*, *J.C.S.*, 1956, 1294

Mason, S.F. *et al.*, *J.C.S.*, 1957, 4874; 5010;

1959, 1253 (uv, tautom)

Cheeseman, G.W.H. *et al.*, *J.C.S.*, 1960, 242

(struct)

Tamura, C. *et al.*, *Bull. Chem. Soc. Jpn.*, 1963,

36, 1187 (Emimycin)

Terao, M. *et al.*, *J. Antibiot., Ser. A*, 1963, 16,

182 (deriv)

Okada, S. *et al.*, *Chem. Pharm. Bull.*, 1971, 19,

1344

Uchimani, F. *et al.*, *J. Het. Chem.*, 1971, 8, 99

(ms)

Bloch, A. *et al.*, *J. Med. Chem.*, 1972, 15, 164;

1973, 16, 183 (Emimycin)

Marx, G.S. *et al.*, *J.O.C.*, 1972, 37, 111 (pmr)

MacDonald, J.C. *et al.*, *Tetrahedron*, 1976, 32,

655 (nmr)

Tsujimoto, T. *et al.*, *Chem. Pharm. Bull.*, 1979,

27, 1169 (pmr)

Mano, M. *et al.*, *Chem. Pharm. Bull.*, 1980, 28,

2720 (Emimycin)

Isono, K. *et al.*, *J. Antibiot.*, 1988, 41, 1711

(rev)


Lewis, R.J. *et al.*, *Food Additives Handbook*,

Van Nostrand Reinhold International, New

York, 1989, MFN285

- Peters, D.A. *et al.*, *Acta Cryst. C*, 1992, **48**, 307 (cryst struct, Me ether oxide)
 Plé, N. *et al.*, *Synthesis*, 1996, 838 (Me ether)
 Cherng, Y.-J. *et al.*, *Tetrahedron*, 2002, **58**, 887-890 (Et ether, Ph ether)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MFN285

3(5)-Pyrazolecarboxylic acid, P-890 9CI



$C_4H_4N_2O_2$ 112.088
 Prisms (H_2O). Sol. EtOH; mod. sol. Et_2O , AcOH; insol. $CHCl_3$, C_6H_6 , Mp 293° dec. A claimed isolation in 1997 from a marine sponge was shown to be incorr. (2003).

Me ester: [15366-34-4]
 $C_5H_6N_2O_2$ 126.115
 Cryst. (C_6H_6). Sol. MeOH; spar. sol. petrol. Mp 139-140°.

Et ester: [5932-27-4]
 $C_6H_8N_2O_2$ 140.141
 Cryst. (hexane/ $EtOAc$). Mp 158-159°.

Amide: [33064-36-7]
 $C_4H_5N_3O$ 111.103
 Mp 203-204°.

Nitrile: 3-Cyanopyrazole
 [33064-34-5]
 $C_4H_3N_3$ 93.088
 Cryst. (C_6H_6). Mp 76-77°.

1H-form [1621-91-6]

N-Me: 1-Methyl-1H-pyrazole-3-carboxylic acid, 9CI
 [25016-20-0]
 $C_5H_6N_2O_2$ 126.115
 Cryst. Mod. sol. hot H_2O . Mp 155-155.5° (146°) Mp 222°.

N-Me, Me ester: [17827-61-1]
 $C_6H_8N_2O_2$ 140.141
 Liq. Bp₉ 120°.

N-Me, Et ester:
 $C_7H_{10}N_2O_2$ 154.168
 Liq. Mp 9-11°. Bp₁₃ 140-143°.

N-Ph: 1-Phenyl-1H-pyrazole-3-carboxylic acid, 9CI
 [4747-46-0]
 $C_{10}H_8N_2O_2$ 188.185
 Needles (H_2O). Mp 146°. p*K*_{a1} 3.6 (20°).

N-Ph, Me ester:
 $C_{11}H_{10}N_2O_2$ 202.212
 Needles. Mp 77°.

N-Ph, nitrile: 3-Cyano-1-phenyl-1H-pyrazole
 [18093-85-1]
 $C_{10}H_7N_3$ 169.185
 Cryst. (Et_2O /petrol).

2H-form

N-Me: 1-Methyl-1H-pyrazole-5-carboxylic acid, 9CI
 [16034-46-1]
 [25832-41-1, 17827-60-0]

$C_5H_6N_2O_2$ 126.115
 Isol. from the gorgonian *Echinomuraceae splendens*. Cryst. (H_2O). Mp 223-224° (219-220°).

N-Me, Et ester:
 Liq. Bp₁₈ 95°.

N-Ph: 1-Phenyl-1H-pyrazole-5-carboxylic acid, 9CI
 [1133-77-3]
 $C_{10}H_8N_2O_2$ 188.185
 Needles (H_2O). Mp 183°. p*K*_{a1} 2.7 (20°).

N-Ph, Me ester: [55115-07-6]
 Needles. Mp 67°.

N-Ph, nitrile: 5-Cyano-1-phenyl-1H-pyrazole
 $C_{10}H_7N_3$ 169.185
 Liq. Bp_{0.625} 95-100° (lit. gives a pressure range).

Claisen, L. *et al.*, *Annalen*, 1894, **278**, 292 (*N-Ph*)
 v. Pechmann, H. *et al.*, *Ber.*, 1900, **33**, 3594 (*synth*)

Buchner, E. *et al.*, *Ber.*, 1902, **35**, 37 (*synth*)
 Rojahn, C.A. *et al.*, *Annalen*, 1924, **437**, 303 (*N-Ph*)

v. Auwers, K. *et al.*, *J. Prakt. Chem.*, 1935, **143**, 259 (*ester*)

Cusmano, S. *et al.*, *Gazz. Chim. Ital.*, 1940, **70**, 227 (*N-Ph*)

Panizzi, L. *et al.*, *Gazz. Chim. Ital.*, 1947, **77**, 556 (*N-Ph*)

Roedig, A. *et al.*, *Annalen*, 1955, **597**, 214 (*N-Ph*)

Hüttel, R. *et al.*, *Annalen*, 1959, **625**, 55 (*synth*)
 Gundermann, K. *et al.*, *Chem. Ber.*, 1960, **93**, 883 (*synth*)

Wijnberger, C. *et al.*, *J. Het. Chem.*, 1969, **6**, 545 (*N-Me*)

Manecke, G. *et al.*, *Tet. Lett.*, 1969, 617 (*synth*)
 Sasaki, T. *et al.*, *J.C.S. (C)*, 1971, 2147 (*derivs*)

Akhrem, A.A. *et al.*, *Tet. Lett.*, 1973, 2655 (*pmr, ir, uv*)

Scherowsky, G. *et al.*, *Annalen*, 1977, 1235 (*N-Ph*)

Beck, J.R. *et al.*, *J. Het. Chem.*, 1988, **25**, 555 (*N-Ph*)

Pavlik, J.W. *et al.*, *J. Het. Chem.*, 1992, **29**, 1357 (*N-Me*)

Martins, M.A.P. *et al.*, *Synthesis*, 1995, 1491 (*Et ester*)


Parameswaran, P.S. *et al.*, *Indian J. Chem., Sect. B*, 2002, **41**, 1093-1096 (*isol, N-Me*)

Rao, K.V. *et al.*, *J. Nat. Prod.*, 2003, **66**, 823-828 (*claimed isol*)

Pyrazomycin

P-891

4-Hydroxy-3-ribofuranosyl-1H-pyrazole-5-carboxamide, 9CI. A 23813. Antibiotic A 23813



$C_9H_{13}N_3O_6$ 259.218
 Nucleoside antibiotic. Sol. H_2O , EtOH, butanol, MeOH; fairly sol. $EtOAc$; poorly sol. $CHCl_3$, hexane. Log P -2.91

(calc). λ_{max} 232 (ϵ 7400); 263 (ϵ 6200) ($EtOH/HCl$) (Derep). λ_{max} 235 (ϵ 5100); 307 (ϵ 5100) ($EtOH/NaOH$) (Derep). λ_{max} 232 (ϵ 6200); 263 (ϵ 6210) (pH 7 H_2O) (Derep). λ_{max} 226 (ϵ 7600); 267 (ϵ 6000) (MeOH) (Berdy). λ_{max} 232 (ϵ 7400); 263 (ϵ 6200) ($EtOH$) (Berdy). λ_{max} 235 (ϵ 5100); 307 (ϵ 5100) ($EtOH-NaOH$) (Berdy). λ_{max} 232 (ϵ 6000) ($EtOH-HCl$) (Berdy). λ_{max} 263 (ϵ 6200) (pH 7 buffer) (Berdy). λ_{max} 307 (ϵ 8100) (pH 12 buffer) (Berdy).

α -D-form

Pyrazofurin B. Pyrazomycin B

[41855-21-4]
 Isol. from *Streptomyces candidus*. Dihydrate. Sol. H_2O , MeOH; poorly sol. Et_2O , hexane. Mp 69-70°. λ_{max} 232 (ϵ 7400); 263 (ϵ 6200) ($EtOH/HCl$) (Derep). λ_{max} 235 (ϵ 5100); 307 (ϵ 5100) ($EtOH/NaOH$) (Derep). λ_{max} 232 (ϵ 6200); 263 (ϵ 6210) (pH 7 H_2O) (Derep). λ_{max} 225 (ϵ 8000); 276 (ϵ 6700) ($EtOH$) (Berdy).

β -D-form

Pyrazofurin, USAN. Pirazofurin, INN.

Antibiotic A 23812. NSC 143095. A 23812
 [30868-30-5]
 From *Streptomyces candidus*. Shows antibiotic and antineoplastic props. Cryst. (H_2O). Mp 112-115°. [α]_D²⁵ -49.6 (c, 0.80 in H_2O).

► Gastrointestinal disturbances and adverse skin effects reported when used therapeutically. UQ6360000

Ger. Pat., 1971, 2 019 838; *CA*, **74**, 41133s (*isol*)
 Farkas, J. *et al.*, *Tet. Lett.*, 1972, 2279 (*synth*)
 Wenkert, E. *et al.*, *Biochem. Biophys. Res. Commun.*, 1973, **51**, 318 (*cmr*)
 Sweeney, M.J. *et al.*, *Cancer Res.*, 1973, **33**, 2619

Crain, P.F. *et al.*, *J. Het. Chem.*, 1973, **10**, 843 (*ms*)

de Bernardo, S. *et al.*, *J.O.C.*, 1976, **41**, 287 (*synth*)

U.S. Pat., 1976, 3 998 999; *CA*, **86**, 155899z (*synth*)

Neumann, J.M. *et al.*, *Biochim. Biophys. Acta*, 1977, **479**, 427 (*conformn, nmr*)

Miles, D.W. *et al.*, *J. Phys. Chem.*, 1983, **87**, 2444 (*conformn*)

Karagiri, N. *et al.*, *J.C.S. Perkin 1*, 1984, 553 (*synth*)

Lopez Herrera, F.J. *et al.*, *Carbohydr. Res.*, 1985, **139**, 95 (*analogues*)

Petrie, C.R. *et al.*, *J. Med. Chem.*, 1986, **29**, 268 (*derivs*)


Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 1300

Isono, K. *et al.*, *J. Antibiot.*, 1988, **41**, 1711 (*rev*)

Pyrenoline A†

P-892

7,8-Dihydro-6,10-dihydroxy-7-methoxy-3-methylbenz[*g*]isoquinolin-9(6H)-one, 9CI. Chrysanthone†
 [129134-97-0]



C₁₅H₁₅NO₄ 273.288

Related to Bostroycoidin, B-255. Isol. from *Ascochyta chrysanthemi*. Prod. by the fungus *Pyrenophora teres*. Phytotoxin, shows weak antifungal and antibacterial activity. Orange cryst. Mp 193° (80° dec.). [α]_D -37.2 (c, 0.1 in Py).

Demethoxy: Pyrenoline B

[129134-98-1]


C₁₄H₁₃NO₃ 243.262

From *Pyrenophora teres*. Phytotoxin. Phytotoxin. Light orange oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D -22 (c, 0.0014 in MeOH). λ_{max} 244 (ε 8300); 270 (ε 7300); 350 (ε 7400) (MeOH) (Berdy).

Albinati, A. et al., *Phytochemistry*, 1989, **28**, 923 (isol, uv, ir, pmr, cmr, ms, cryst struct)
Coval, S.J. et al., *Tet. Lett.*, 1990, **31**, 2117 (isol, struct)

Pyridindolol P-893

α'-(Hydroxymethyl)-9H-pyrido[3,4-b]indole-1,3-dimethanol, 9CI. 1-(1,2-Dihydroxyethyl)-3-hydroxymethyl-β-carbazole. Tu 2480F₁. Antibiotic Tu 2480F₁



C₁₄H₁₄N₂O₃ 258.276

λ_{max} 256 (ε 38000); 306 (ε 20000); 380 (ε 5620) (MeOH/HCl) (Derep). λ_{max} 216 (ε 22900); 238 (ε 41690); 245 (sh) (ε 33880); 255 (sh) (ε 28840); 285 (sh); 345 (ε 5130); 350 (ε 5130) (MeOH) (Derep). λ_{max} 216 ; 237 ; 291 ; 343 ; 355 (EtOH) (Berdy).

(R)-form [55812-46-9]

Alkaloid from *Streptomyces alboverticillatus* and *Streptomyces parvulus*. Shows antibiotic props. β-Galactosidase inhibitor. Needles. Sol. MeOH, butanol; poorly sol. CHCl₃, hexane. Mp 167-168°. [α]_D²⁰ -49 (c, 1 in MeOH).

▶UU9810000

14-O-β-D-Glucopyranoside: Antibiotic Tu 2480F₂, Tu 2480F₂

[115178-47-7]

C₂₀H₂₄N₂O₈ 420.418

From *Streptomyces parvulus*. Glycosidase inhibitor. Needles + 1/8H₂O (MeOH/EtOAc). Mp 228-231° dec. [α]_D²⁰ -100 (c, 0.5 in MeOH). λ_{max} 256 (ε 30000); 306 (ε 20000); 380 (ε 5620) (EtOH/HCl) (Derep). λ_{max} 215 (ε 11200); 237 (ε 32400); 290 (ε 14100); 343 (ε 4570); 354 (ε 4570) (EtOH) (Derep).

15-O-β-D-Glucopyranoside: Antibiotic Tu 2480F₃, Tu 2480F₃

[115178-45-5]

C₂₀H₂₄N₂O₈ 420.418

From *Streptomyces parvulus*. Glycosidase inhibitor. Powder. [α]_D²⁰ -90 (c, 0.5 in MeOH). λ_{max} 256 (ε 30000); 306 (ε 20000); 380 (ε 5620) (EtOH/HCl)

(Derep). λ_{max} 215 (ε 11200); 237 (ε 32400); 290 (ε 14100); 343 (ε 4570); 354 (ε 4570) (EtOH) (Derep).

16-O-β-D-Glucopyranoside: Antibiotic Tu 2480F₄, Tu 2480F₄

[115178-46-6]

C₂₀H₂₄N₂O₈ 420.418

From *Streptomyces parvulus*. Glycosidase inhibitor. Powder. [α]_D²² -80 (c, 0.5 in MeOH). λ_{max} 256 (ε 30000); 306 (ε 20000); 380 (ε 5620) (EtOH/HCl) (Derep). λ_{max} 215 (ε 11200); 237 (ε 32400); 290 (ε 14100); 343 (ε 4570); 354 (ε 4570) (EtOH) (Derep). λ_{max} 214 (ε 11750); 237 (ε 35480); 239 (ε 214800); 340 (ε 1445); 351 (ε 4570) (EtOH) (Berdy).

16-Ac: Pyridindolol K2

[189756-51-2]

C₁₆H₁₆N₂O₄ 300.313

Alkaloid from *Streptomyces* sp. K93-0711. Needles. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 123-124°. [α]_D²⁰ -35 (c, 0.4 in MeOH). λ_{max} 215 (log ε 4.23); 238 (log ε 4.42); 245 (sh) (log ε 4.39); 254 (sh) (log ε 4.32); 262 (sh) (log ε 4.12); 282 (sh) (log ε 3.84); 291 (log ε 4.02); 343 (log ε 3.53); 356 (sh) (log ε 3.48) (MeOH).

15,16-Di-Ac: Pyridindolol K1

[189756-49-8]

C₁₈H₁₈N₂O₅ 342.351

Alkaloid from *Streptomyces* sp. K93-0711. Oil. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁰ -16 (c, 0.2 in MeOH). λ_{max} 205 (sh) (log ε 4.02); 215 (log ε 4.04); 238 (log ε 4.14); 247 (sh) (log ε 4.08); 254 (sh) (log ε 4.03); 262 (log ε 3.87); 282 (sh) (log ε 3.62); 290 (log ε 3.75); 346 (log ε 3.35); 356 (sh) (log ε 3.28) (MeOH).

l'-Carboxylic acid: see 1-(1,2-Dihydroxyethyl)-β-carboline-3-carboxylic acid, D-579

(±)-form [66183-62-8]

Cryst. (EtOAc/MeOH). Mp 169-170°.

Aoyagi, T. et al., *J. Antibiot.*, 1975, **28**, 555-557 (isol, ir, uv, struct)

Naganawa, H. et al., *J. Antibiot.*, 1975, **28**, 876-880 (struct, nmr, cryst struct)

Soerens, D. et al., *J.O.C.*, 1979, **44**, 535-545 (synth)

Bieraugel, H. et al., *Tetrahedron*, 1983, **39**, 3987-3990 (synth)

Hagmann, L. et al., *J. Antibiot.*, 1988, **41**, 289-295 (derivs)

Kim, Y.-P. et al., *J. Antibiot.*, 1997, **50**, 189-193 (*Pyridindolol K1/K2*)

Kanekiyo, N. et al., *Heterocycles*, 2000, **53**, 1877-1880 (*Pyridindolol K2*, synth)

Kanekiyo, N. et al., *J.O.C.*, 2002, **66**, 8793-8798 (synth)

Pyridine, 9CI, 8CI

Azabenzene. Azine. FEMA 2966

[110-86-1]



C₅H₅N 79.101

Extracted in quantity from coal tar.

P-894

Found in traces in some plants, e.g. *Nicotiana paniculata*, *Atropa belladonna* (Solanaceae). Powerful solvent; dissolves inorg. salts. Org. intermed. Solvent in Karl Fischer reagent for detn. of H₂O, denaturant for industrial ethanol. Important analytical solvent; often used in separation and detn. of elements. Liq. with characteristic unpleasant odour and sharp taste. Misc. H₂O, petrol, Et₂O, etc. d₄²⁵ 0.98. Fp -42. Bp 115.5°. n_D²¹ 1.5092. pK_a 5.58 (25°). pK_a 5.23 (25°). Forms azeotropic mixt. with 3 mols. H₂O, Bp 92-3°. Hygroscopic. Steam-volatile.

▶ Flammable, fl. p. 17°, autoignition temp. 482°. Eye and skin irritant. May cause dermatitis; symptoms include headache, vertigo, insomnia, nausea. Prolonged or repeated exposure injures kidney and liver. LD₅₀ (rat, orl) 891 mg/kg. LD₅₀ (rbt, skn) 1121 mg/kg. OES: long-term 5 ppm; short-term 10 ppm. UR8400000 [16969-45-2, 7291-22-7]

Luckner, M. et al., *Pharmazie*, 1964, **19**, 1 (biosynth)

Boodman, N.S. et al., *Chem. Heterocycl. Compd.*, Suppl. 1, 1974, **14**, 183 (synth)

Kirk-Othmer Encycl. Chem. Technol., 3rd edn., Wiley, 1978, **19**, 454 (rev)

3-Pyridinecarboxamide, 9CI P-895


Nicotinamide, 8CI, 1N. Niacinamide.

Nicotinic acid amide. *Antipellagra factor*.

Nicosan 2. *PP-Factor*. *Vitamin B₃*. *Many other names*

[98-92-0]

[119047-33-5 (perchlorate)]



C₆H₆N₂O 122.126

Widespread in plants, e.g. rice, yeast and fungi. Prod. by the marine bacterium strain He159b. Enzyme cofactor, used in treatment of pellagra, low mammalian toxicity. Needles (C₆H₆). Mp 129-130°. Bp_{0.0005} 150-160°. pK_a 3.33 (20°). Log P -0.21 (calc).

▶QS3675000

N'-Me: *N*-Methyl-3-pyridinecarboxamide, 9CI. *N*-Methylnicotinamide, 8CI. 3-(Methylcarbamoyl)pyridine [114-33-0]

C₇H₈N₂O 136.153

Metabolised from Tryptophan, T-640 and 3-Pyridinecarboxylic acid, P-897; principal urinary excretion product of 3-Pyridinecarboxylic acid, P-897 in man, dogs, pigs, rats, etc. Mp 105°.

[6485-44-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 796B; 917A (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 338B; 342B (nmr)

La Forge, F.B. et al., *J.A.C.S.*, 1928, **50**, 2477 (synth)


Georg, A. et al., *Helv. Chim. Acta*, 1943, **26**, 358 (synth)

Long, K.R. et al., *J. Magn. Reson.*, 1972, **8**, 207-216 (*N*'-Me, cmr)

Bernardi, L. *et al.*, *Chem. Comm.*, 1975, 320-321 (N'-Me)
 Auerbach, J. *et al.*, *J.O.C.*, 1976, **41**, 725 (synth)
 Kirk-Othmer *Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **24**, 59 (rev)
 Oppenheimer, N.J. *et al.*, *Org. Magn. Reson.*, 1980, **13**, 14 (N-15 nmr)
 Shabaan, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (marine, isol)
 Roje, S. *et al.*, *Phytochemistry*, 2007, **68**, 1904-1921 (biosynth, rev)

2-Pyridinecarboxylic acid, 9CI P-896

Picolinic acid, 8CI
 [98-98-6]



C₆H₅NO₂ 123.111
 Isol. from culture medium of *Piricularia oryzae*. Catalyst for CrO₃ oxidns. Used as 8% aq. soln. for extraction separation of V(III). Needles (H₂O, EtOH or C₆H₆). Sol. MeOH, EtOAc; fairly sol. H₂O, C₆H₆, Et₂O; poorly sol. hexane. Mp 136-138° (melts with sublimation). pK_{a1} 0.99; pK_{a2} 5.39 (25°). λ_{max} 265 (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 360 mg/kg. TJ7344000
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 784C; 792C; 800A; 916B (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 316C; 328C; 347A (nmr)
 Singer, A.W. *et al.*, *Org. Synth.*, Coll. Vol., **3**, 1955, 740-741 (synth, bibl)
 Tamari, K. *et al.*, *CA*, 1965, **63**, 6049 (isol)
 Pollak, P.I. *et al.*, *Chem. Heterocycl. Compd.*, Suppl. 3, 1974, **14**, (bibl)
 Cook, I.B. *et al.*, *Aust. J. Chem.*, 1989, **42**, 1493 (cmr)
 Csanády, G. *et al.*, *Org. Prep. Proced. Int.*, 1990, **22**, 507 (synth)
 de Oliveira, S.M. *et al.*, *Org. Mass Spectrom.*, 1991, **26**, 855 (ms)

3-Pyridinecarboxylic acid P-897

Nicotinic acid, INN. Niacin, USAN. Nicamin. Nicobid. Nyclin. Pelonin. Wampocap. Many other names [59-67-6]

C₆H₅NO₂ 123.111
 Present in fruits and other plant materials. Used as 2.5% aq. soln. for extraction-photometric detn. of U(VI) (λ_{max} 556 nm, ε 120000). Vitamin, enzyme cofactor, vasodilator. Antihyperglycaemic agent. Used to treat lipid disorders. Needles (H₂O or EtOH). Sol. hot H₂O; spar. sol. Et₂O. Mp 236° (225-227°). pK_{a1} 2; pK_{a2} 4.82 (25°). Log P 0.8 (calc). Sublimes.

▶ Can cause dermal and gastrointestinal reactions and hepatic effects at high therapeutic doses. Low acute mammalian toxicity. QT0525000

β-D-Glucopyranosyl ester: 1-O-Nicotinoyl-β-D-glucopyranose [35214-85-8]
 C₁₂H₁₅NO₇ 285.253
 Present in tobacco (*Nicotiana tabacum*)

cells grown in a high conc. of niacin. Prob. a detoxification form of niacin in the plant. λ_{max} 263 (log ε 3.22) (H₂O).
 N-(2-Hydroxyethyl), betaine: 3-Carboxy-1-(2-hydroxyethyl)pyridinium betaine. **Pyridinebetaine A** [52578-48-0]
 C₈H₉NO₃ 167.164
 Isol. from the sponge *Agelas dispar*. Amorph. solid. λ_{max} 220 (ε 3000); 266 (ε 2500) (MeOH).

[1976-28-9]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 785B; 793D; 794A; 797C; 800C; 916C (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 317B; 330B; 331A; 331B; 332B; 336C; 340B; 347C (nmr)
Org. Synth., Coll. Vol., **1**, 1932, 385 (synth)
 Pollak, P.I. *et al.*, *Chem. Heterocycl. Compd.*, Suppl. 3, 1974, **14**, (bibl)
 Kirk-Othmer *Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **24**, 59 (rev)
 Domisse, R. *et al.*, *Heterocycles*, 1981, **16**, 1893-1897 (cmr)
 Staedeli, W. *et al.*, *Org. Magn. Reson.*, 1981, **15**, 106-109 (N-15 nmr)
 Ohshima, N. *et al.*, *Phytochemistry*, 1997, **44**, 1483-1484 (glucosyl ester)
 Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1171-1173 (Pyridinebetaine A)
 McCormack, P.L. *et al.*, *Drugs*, 2005, **65**, 2719-2740 (rev)

4-Pyridinecarboxylic acid, P-898


9CI
 Isonicotinic acid, 8CI
 [55-22-1]

C₆H₅NO₂ 123.111
 Needles (H₂O). Subl. 315. pK_{a1} 1.77; pK_{a2} 4.84 (25°).

▶ LD₅₀ (rat, orl) 5000 mg/kg. NS1103000
 2R-Methylheptyl ester: 2-Methylheptyl isonicotinate
 C₁₄H₂₁NO₂ 235.325
 Prod. by *Streptomyces* sp. 201. Antibacterial and antifungal agent. Brown solid. [α]_D²⁰ +8.3 (c, 0.6 in CHCl₃). λ_{max} 220 (CHCl₃).
 Bordoloi, G.N. *et al.*, *Biosci., Biotechnol., Biochem.*, 2001, **65**, 1856-1858 (2-methylheptyl ester, isol)

2,6-Pyridinedicarbothioic acid, 9CI P-899

Pyridine-2,6-di(monothiocarboxylic acid) [69945-42-2]




C₇H₅NO₂S₂ 199.254
 Prod. by *Pseudomonas putida* and *Pseudomonas stutzeri*. Siderophore. λ_{max} 267; 340 (MeOH) (Berdy).
Fe complex: [97253-20-8] [95787-42-1]
 C₁₄H₆FeN₂O₄S₄²⁻ 450.323
 Prod. by *Pseudomonas putida*. Blue powder.

Di-OH-form

Di-Me ester: [84877-69-0]
 C₉H₉NO₂S₂ 227.308
 Orange cryst. (MeOH). Mp 100-100.5°
 Ockels, W. *et al.*, *Tet. Lett.*, 1978, 3341-3342 (isol)
 Jones, B.A. *et al.*, *J.O.C.*, 1983, **48**, 2635-2639 (synth, ester)
 Hildebrand, U. *et al.*, *Phosphorus Sulfur Relat. Elem.*, 1983, **16**, 361-364 (cryst struct)
 Hildebrand, U. *et al.*, *Z. Naturforsch., C*, 1985, **40**, 201-207 (isol)
 Lee, C.-H. *et al.*, *Biochem. Biophys. Res. Commun.*, 1999, **261**, 562-566 (isol)
 Zawadzka, A.M. *et al.*, *Appl. Environ. Microbiol.*, 2006, **72**, 3119-3129 (isol, activity)

2,3-Pyridinedicarboxylic acid, 9CI, 8CI P-900

Quinolinic acid
 [89-00-9]



C₇H₅NO₄ 167.121
 Intermed. in the biosynth. of nicotinic acid dinucleotide in microorganisms. Convulsive agent. Prisms (H₂O). Spar. sol. H₂O; prac. insol. EtOH, Et₂O. Mp 188° Mp 190-195° dec. pK_{a1} 2.43; pK_{a2} 4.78 (25°, 0.1M NaClO₄).

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 326A; 336A (nmr)
Aldrich Library of NMR Spectra, **9**, 44D (pmr)
 Oliveto, E.P. *et al.*, *Chem. Heterocycl. Compd.*, (Part 3), 1962, **14**, 179 (bibl)
 Caswell, L.R. *et al.*, *J. Het. Chem.*, 1972, **9**, 551 (rev)
 Sutter, P. *et al.*, *J. Het. Chem.*, 1986, **23**, 29 (synth, bibl)

2,6-Pyridinedicarboxylic acid, 9CI, 8CI P-901

Dipicolinic acid
 [499-83-2]

C₇H₅NO₄ 167.121
 Occurs in bacteria and fungi, e.g. *Verticillium*, *Pseudomonas*, *Penicillium*, *Bacillus* spp. Used as 0.1M aq. soln. for photometric detn. of Ag, Cr, V and Mn; gravimetric detn. of U. Polyamide fibres have been synth. by reaction of the di-Ph ester or dichloride with diamines. Coml. antibacterial agent. Also shows mycotoxic and insecticidal props. Cryst. + 1½H₂O (H₂O). Mp 252° dec. (anhyd.). pK_{a1} 2.24; pK_{a2} 4.67 (25°). λ_{max} 270; 278 (MeOH) (Berdy).

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 791D; 795D
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 327B; 334A; 337B (nmr)
 Biemann, K. *et al.*, *J.A.C.S.*, 1957, **79**, 5558-5564 (synth, ir)
 Oliveto, E.P. *et al.*, *Chem. Heterocycl. Compd.*, (Part 3), 1962, **14**, 179-346 (bibl)
 Hodson, P.H. *et al.*, *J. Bacteriol.*, 1966, **91**, 562-569 (occur)
 Brown, E.V. *et al.*, *J. Het. Chem.*, 1971, **8**, 189-192 (ms)

3,5-Pyridinedicarboxylic acid, 9CI, 8CI P-902*Dimicotinic acid*


[499-81-0]

C₇H₅NO₄ 167.121Cryst. Prac. insol. H₂O. Mp 323° dec.pK_{a1} 1.1; pK_{a2} 2.72; pK_{a3} 4.62 (25°).


Several polyamides have been synth.

Diamide: 3,5-Pyridinedicarboxamide. Dinicotinic amide

[4663-99-4]

C₇H₇N₃O₂ 165.151Alkaloid from *Aristolochia zollingeriana* and *Peristrophe japonica*. Cryst. + ½H₂O (DMF). Mp 318-319° dec. (310-311°).Kuthan, J. et al., *Coll. Czech. Chem. Comm.*, 1966, **31**, 2618 (esters, diamide)Chiang, C.-Y. et al., *J. Chin. Chem. Soc. (Taipei)*, 1998, **45**, 93-97 (isol, diamide)Carelli, V. et al., *J.C.S. Perkin 1*, 2002, 542-547 (diamide, bismethylamide, synth, pmr)**3-Pyridinedodecanal** P-903*12-(3-Pyridinyl)dodecanal*C₁₇H₂₇NO 261.406*Oxime:* [291775-86-5]C₁₇H₂₈N₂O 276.421Isol. from *Amphimedon* sp. Config. not determined. λ_{max} 264 (ε 3200) (MeOH).Hirano, K. et al., *Chem. Pharm. Bull.*, 2000, **48**, 974-977 (isol, pmr, cmr, uv)**3-Pyridinemethanol, 9CI, 8CI** P-904*3-Hydroxymethylpyridine. Nicotinyl alcohol, BAN. NSC 526046*


[100-55-0]

C₆H₇NO 109.127Vasodilator. Bp₁₅ 142-143° Bp_{0.1} 110°.pK_a 4.93 (25°). Log P -0.39 (calc).


▶ UT4690000

O-β-D-Glucopyranoside: Nicoloside

[151870-75-6]

C₁₂H₁₇NO₆ 271.269Alkaloid from whole plants of *Anoectochilus kosshumensis* (Orchidaceae).Oil. [α]_D²⁴ -37.7 (c, 2.60 in MeOH).Ito, A. et al., *Phytochemistry*, 1993, **33**, 1133 (*Nicoloside*)**3-Pyridinetricanal** P-905*13-(3-Pyridinyl)tridecanal*C₁₈H₂₉NO 275.433*Oxime:* [291775-85-4]C₁₈H₃₀N₂O 290.448Isol. from *Amphimedon* sp. Config. not determined. λ_{max} 264 (ε 3100) (MeOH).Hirano, K. et al., *Chem. Pharm. Bull.*, 2000, **48**, 974-977 (isol, pmr, cmr, uv)**3-Pyridinol** P-906*3-Hydroxypyridine. 3-Pyridol*


[109-00-2]

C₅H₅NO 95.101Tautomerises ca. 50% to zwitterion in H₂O at r.t. Reagent used in peptide synth. Needles. Mp 129°. pK_{a1} 4.79; pK_{a2} 8.75 (20°).▶ LD₅₀ (mus, ipr) 1822 mg/kg. UU7701400**OH-form***Me ether: 3-Methoxypyridine*

[7295-76-3]


C₆H₇NO 109.127Isol. from *Thermopsis rhombifolia* and *Equisetum arvense*. Bp 178-179°. n_D²¹ 1.5165. n_D²¹ 1.5202.Manske, F. et al., *Can. J. Res., Sect. B*, 1942, **20**, 265 (*Me ether, isol*)**Pyridinoline** P-907*4-(2-Amino-2-carboxyethyl)-1-(5-amino-5-carboxy-2-hydroxypentyl)-3-(3-amino-3-carboxypropyl)-5-hydroxypyridinium(1+), 9CI. Hydroxylysylpyridinoline* [77464-35-8]

[79646-35-8, 128112-67-4, 220243-86-7, 63800-01-1, 125659-26-9]

C₁₈H₂₈N₄O₈[⊕] 428.441Several diastereomers possible. -ve charge also shown on the phenolic O. Derived from collagen cross-links formed between adjacent hydroxylysine residues in type I and type II collagen of bone and cartilage respectively. Released into serum and urine during bone resorption. Clinical marker. Used to assess bone resorption levels in metabolic bone diseases. Solid. [α]_D²⁰ +39.17 (c, 0.24 in MeOH) (as trifluoroacetate salt). CAS no. refers to hydroxide inner salt.*2'-Deoxy: Deoxyypyridinoline. Lysylpyridinoline*

[82637-02-3]

[119241-22-4, 200405-37-4, 220243-98-1, 83462-55-9, 90032-33-0]

C₁₈H₂₈N₄O₇[⊕] 412.442Derived from collagen cross-links formed between adjacent lysine residues in type I collagen of bone and dentine. Released into serum and urine during bone resorption. Marker for bone metabolism. Clinical marker of bone turnover in osteoporosis. Pale yellow solid (as trifluoroacetate salt). [α]_D²⁰ +31.6 (c, 0.25 in MeOH) (as trifluoroacetate).*2'-Epimer:* [220348-30-1]C₁₈H₂₈N₄O₈[⊕] 428.441Synthetic. [α]_D²⁰ +43.23 (c, 0.26 in MeOH) (as trifluoroacetate salt). CAS no. refers to trifluoroacetate.Fujimoto, D. et al., *Biochem. Biophys. Res. Commun.*, 1977, **76**, 1124-1129; 1978, **84**, 52-57 (isol, uv, pmr, cmr, nmr, ms, fluorescence)Ogawa, T. et al., *Biochem. Biophys. Res. Commun.*, 1982, **107**, 1252-1257 (*Deoxyypyridinoline, isol, pmr*)Barber, M. et al., *Biochem. Biophys. Res. Commun.*, 1982, **109**, 1041-1046 (*struct, ms, uv*)Black, D. et al., *Anal. Biochem.*, 1988, **169**, 197-203 (*deoxyypyridinoline, isol, chromatog*)Acil, Y. et al., *J. Chromatogr., A*, 1994, **664**, 183-188 (*isol*)Arbault, P. et al., *J. Liq. Chromatogr.*, 1994, **17**, 1981-1993 (*isol, chromatog*)McGuinness, B.J. et al., *Biochem. Soc. Trans.*, 1995, **23**, 3865 (*deoxyypyridinoline, pmr*)Waelchli, R. et al., *Bioorg. Med. Chem. Lett.*, 1997, **7**, 2831-2836 (*synth*)Lamprecht, G. et al., *J. Chromatogr., B*, 1997, **691**, 297-304 (*deoxyypyridinoline, isol, chromatog*)Knott, L. et al., *Bone*, 1998, **22**, 181-187 (*rev*)
St Clair, E.W. et al., *J. Rheumatol.*, 1998, **25**, 1472-1479 (*detn, clinical*)U.S. Pat., 1998, 5 723 619; CA, **128**, 205137s (*synth*)Yilmaz, N. et al., *Clin. Chem. Lab. Med.*, 1999, **37**, 137-143 (*detn, Deoxyypyridinoline*)Adamczyk, M. et al., *Tetrahedron*, 1999, **55**, 63-88; 2000, **56**, 2379-2390 (*synth, pyridinoline, deoxyypyridinoline, pmr, cmr, bibl*)Adamczyk, M. et al., *Tetrahedron: Asymmetry*, 2000, **11**, 2289-2298; 5017 (*synth, pyridinoline, 2'-deoxy, pmr, erratum*)Allevi, P. et al., *Tetrahedron: Asymmetry*, 2002, **13**, 1901-1910; 2004, **14**, 2005-2012 (*synth, pyridinoline, 2'-deoxy*)Allevi, P. et al., *J.O.C.*, 2007, **72**, 3478-3483 (*synth*)**10-(3-Pyridinyl)-3,5,7,9-decatetraen-2-one, 9CI** P-908*3-(9-Oxo-1,3,5,7-decatetraenyl)pyridine*



(all-E)-form

C₁₅H₁₅NO 225.29**(all-E)-form***Navenone A*


[62695-67-4]

Major constit. of the alarm pheromone of the marine opisthobranch mollusc *Navanax inermis* (*Chelidonura inermis*). Yellow cryst. (C₆H₆). Mp 144-145°. λ_{max}

399 (ε 14400) (MeOH) (Berdy).

3-Methyl: 3-Methylnavenone A
[73414-53-6]C₁₆H₁₇NO 239.316Minor constit. of the alarm pheromone of *Navanax inermis*.**(3Z,5E,7E,9E)-form** [73465-89-1]Minor constit. of the alarm pheromone of *Navanax inermis*.**(3Z,5Z,7E,9E)-form** [73952-89-3]Minor constit. of the alarm pheromone of *Navanax inermis*.Sleeper, H.L. *et al.*, *J.A.C.S.*, 1977, **99**, 2367-2368 (*isol, uv, ir, pmr, cmr, ms, struct*)Sakakibara, M. *et al.*, *Agric. Biol. Chem.*, 1979, **43**, 117 (*synth*)Fenical, W. *et al.*, *Pure Appl. Chem.*, 1979, **51**, 1865 (*derivs*)Sleeper, H.L. *et al.*, *J. Chem. Ecol.*, 1980, **6**, 57 (*isol*)Soulez, D. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 203 (*synth*)**12-(2-Pyridinyl)-6,8-dodecadien-2-ol** P-909C₁₇H₂₅NO 259.391**(2R,6E,8E)-form****Phormidinine A**Alkaloid from the cyanobacterium *Phormidium* sp. Oil. [α]_D²⁰ -4 (c, 0.09 in CHCl₃). λ_{max} 232 (ε 17100); 262 (ε 3900); 268 (ε 1700) (MeOH).**Ac: Phormidinine B**C₁₉H₂₇NO₂ 301.428Isol. from *Phormidium* sp. Oil.Teruya, T. *et al.*, *Tet. Lett.*, 2005, **46**, 4001-4003 (*isol, pmr, cmr*)**12-(3-Pyridinyl)-2,4,6-dodecatrien-1-ol, 9CI** P-910C₁₇H₂₃NO 257.375**(2E,4E,6E)-form****Ac: Haminol C**

[147362-36-5]

C₁₉H₂₅NO₂ 299.412Alkaloid from the mollusc *Haminoea orsteai*. Alarm pheromone.Spinella, A. *et al.*, *Tetrahedron*, 1993, **49**, 1307-1314 (*isol, struct*)**12-(3-Pyridinyl)-4,6,8-dodecatrien-2-ol, 9CI** P-911C₁₇H₂₃NO 257.375**(2S,4E,6E,8E)-form****Haminol 1**

[147391-85-3]

Alkaloid from the molluscs *Haminoea orbignyana* and *Haminoea fusari*. Alarm pheromone.**Ac: Haminol 2**


[147391-84-2]

C₁₉H₂₅NO₂ 299.412From *Haminoea orbignyana* and *Haminoea fusari*. Alarm pheromone. [α]_D²⁰ -19 (c, 1.3 in MeOH).**8,9-Dihydro: 12-(3-Pyridinyl)-4,6-dodecadien-2-ol, 9CI. Haminol 3**

[147362-30-9]

C₁₇H₂₅NO 259.391From *Haminoea fusari*. Alarm pheromone.**8,9-Dihydro, Ac: Haminol 4**


[147362-25-2]

C₁₉H₂₇NO₂ 301.428From *Haminoea fusari*. Alarm pheromone. [α]_D²⁰ -12.5 (c, 0.7 in MeOH).Spinella, A. *et al.*, *Tetrahedron*, 1993, **49**, 1307 (*isol, uv, ir, pmr, cmr, ms, struct*)Solladie, G. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 801-810 (*synth, Haminol 1*)**12-(3-Pyridinyl)-4,6,11-dodecatrien-2-ol** P-912C₁₇H₂₃NO 257.375**(2S,4E,6E,11E)-form****Haminol A**

[133412-13-2]

Alkaloid from the molluscs *Haminoea navicula* and *Haminoea orsteai*. Alarm pheromone. Oil. [α]_D +5 (c, 0.3 in MeOH). λ_{max} 232 (ε 30200); 282 (ε 4340) (MeOH) (Derep).**Ac: Haminol B**

[133412-14-3]

C₁₉H₂₅NO₂ 299.412From *Haminoea navicula*. Alarm pheromone. Oil. [α]_D -24 (c, 0.4 in MeOH). λ_{max} 230 (ε 23400); 282 (ε 3440) (MeOH) (Derep).Cimino, G. *et al.*, *Experientia*, 1991, **47**, 61-63 (*isol, uv, ir, pmr, cmr, ms, struct*)Spinella, A. *et al.*, *Tetrahedron*, 1993, **49**, 1307Matikainen, J. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1622 (*synth*)Alvarez, R. *et al.*, *Tetrahedron: Asymmetry*, 1998, **9**, 3065-3072 (*Haminol B*)**12-(3-Pyridinyl)-5,8,10-dodecatrien-2-ol, 9CI** P-913C₁₇H₂₃NO 257.375**(2S,5E,8E,10E)-form****Haminol 5**

[147391-86-4]

Alkaloid from the mollusc *Haminoea fusari*. Alarm pheromone.**Ac: Haminol 6**

[147362-35-4]

C₁₉H₂₅NO₂ 299.412From *Haminoea fusari*. Alarm pheromone. [α]_D²⁰ -4.2 (c, 0.2 in MeOH).Spinella, A. *et al.*, *Tetrahedron*, 1993, **49**, 1307-1314 (*isol, uv, ir, pmr, cmr, ms, struct*)**9H-Pyrido[3,4-b]indol-1-ol** P-914**2,9-Dihydro-1H-pyrido[3,4-b]indol-1-one. 1-Hydroxy-β-carboline**

[19839-52-2]

C₁₁H₈N₂O 184.197

Solid. Mp 255-257° (245-247°).

Me ether: 1-Methoxy-9H-pyrido[3,4-b]indole. 1-Methoxy-β-carboline. Taraxacine AC₁₂H₁₀N₂O 198.224Alkaloid from the aerial parts of *Taraxacum formosanum*. Pale yellow syrup. λ_{max} 212 (log ε 3.9); 246 (log ε 3.6); 257 (log ε 3.6); 273 (log ε 3.5); 300 (log ε 3.4); 367 (log ε 3.2) (MeOH).Bracher, F. *et al.*, *Annalen*, 1992, 1315 (*synth, ir, pmr, cmr*)Leu, Y.-L. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 599-601 (*Taraxacine A*)**9H-Pyrido[3,4-b]indol-6-ol** P-915**6-Hydroxy-β-carboline**

[58982-28-8]

C₁₁H₈N₂O 184.197Cryst. (toluene or Me₂CO). Mp 189-190°.**Me ether: 6-Methoxy-9H-pyrido[3,4-b]indole, 9CI**

[30684-42-5]

C₁₂H₁₀N₂O 198.224Leaflets (C₆H₆). Mp 202°.**Me ether, N²-Me: 6-Methoxy-2-methyl-β-carbolinium(1+)**

[22022-18-0]

C₁₃H₁₃N₂O⁺ 213.258Quaternary alkaloid from *Desmodium gangeticum* and *Desmodium pulchellum* (Fabaceae). Mp 155-157° dec. (as reineckate). λ_{max} 240 ; 250 ; 273 (sh) ; 290 ; 342 (EtOH).Bannerjee, P.K. *et al.*, *Aust. J. Chem.*, 1969, **22**, 275-277 (*N²-Me Me ether, isol, uv, struct*)Ghosal, S. *et al.*, *Phytochemistry*, 1970, **9**, 429-433 (*N²-Me Me ether, occur*)Périn-Roussel, O. *et al.*, *J.C.S. Perkin 1*, 1972, 531-532 (*Me ether, synth*)Rocca, P. *et al.*, *J. Het. Chem.*, 1995, **32**, 1171-1175 (*synth, pmr, ir*)**9H-Pyrido[3,4-b]indol-7-ol, 9CI** P-916**7-Hydroxy-β-carboline**

[99896-91-0]

C₁₁H₈N₂O 184.197Cryst. (toluene or Me₂CO). Mp >260°.**Me ether: 7-Methoxy-9H-pyrido[3,4-**

b]indole. 7-Methoxy- β -carboline. **Norharmine**

[6253-19-6]
C₁₂H₁₀N₂O 198.224

Alkaloid from the seeds of *Passiflora harmala* (Zygophyllaceae). Needles (C₆H₆). Mp 213-215°. Sublimes. λ_{\max} 225 ; 256 ; 285 ; 325 ; 370 (MeOH).

1,2,3,4-Tetrahydro, Me ether, N⁹-formyl: 1,2,3,4-Tetrahydro-7-methoxy-9H-pyrido[3,4-b]indole-9-carboxaldehyde. 9-Formyl-1,2,3,4-tetrahydro-7-methoxy- β -carboline. **Harmaline**

[110064-66-9]
C₁₃H₁₄N₂O₂ 230.266

Alkaloid from the seeds of *Peganum harmala* (Zygophyllaceae). Fine needles (MeOH). Mp 251-252°. λ_{\max} 205 ; 235 (sh) ; 300 ; 330 (MeOH).

1,2,3,4-Tetrahydro, Me ether, N⁹-formyl, Ac: [110064-67-0]
Fine needles. Mp 235-236°.

Siddiqui, S. et al., *Heterocycles*, 1987, **26**, 1563-1567; 1989, **29**, 521 (*Harmaline*, *Norharmine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)
Rocco, P. et al., *J. Het. Chem.*, 1995, **32**, 1171-1175 (*synth*, *pmr*, *ir*)

9H-Pyrido[3,4-b]indol-8-ol P-917

8-Hydroxy- β -carboline

[334709-41-0]

C₁₁H₈N₂O 184.197

Cryst. (toluene or Me₂CO). Mp 250°.

N²-Me: 8-Hydroxy-2-methyl- β -carbolineum

[356073-89-7]
C₁₂H₁₁N₂O⁺ 199.232

Quaternary alkaloid from a *Pseudodistoma* sp. Gum. Counterion not specified. λ_{\max} 228 (log ϵ 3.68); 268 (log ϵ 3.56); 317 (log ϵ 3.23); 389 (log ϵ 2.7) (MeOH).

Me ether: 8-Methoxy-9H-pyrido[3,4-b]indole, 9CI. 8-Methoxy- β -carboline

[30684-48-1]
C₁₂H₁₀N₂O 198.224


Cryst. (xylene). Mp 204-205°. λ_{\max} 216 ; 243 ; 268 ; 278 ; 287 ; 343 ; 354 (sh) (EtOH).

Ho, B.T. et al., *J. Pharm. Sci.*, 1970, **59**, 1445-1448 (*Me ether*, *synth*)

Rocco, P. et al., *J. Het. Chem.*, 1995, **32**, 1171-1175 (*synth*, *ir*, *pmr*)

Rashid, M.A. et al., *Tetrahedron*, 2001, **57**, 5751-5755 (*N²-Me*, *isol*)

Pyridomacrolidin P-918



C₃₁H₃₅NO₁₀ 581.618

Related to Tenellin, T-72. Prod. by *Beauveria bassiana*. Sol. MeOH, EtOAc, DMSO, Me₂CO; poorly sol. H₂O, hexane. Mp 192-194° dec. $[\alpha]_{\text{D}}^{15}$ +19.4 (c, 0.1


in MeOH). λ_{\max} 238 (ϵ 14500); 342 (ϵ 14600) (MeOH).

Takahashi, S. et al., *J. Antibiot.*, 1998, **51**, 596-598; 1051-1054 (*isol*, *uv*, *pmr*, *cmr*)

Pyridomycin P-919

3-Hydroxy-N-[10-hydroxy-5,11-dimethyl-2-(1-methylpropylidene)-3,7,12-trioxo-9-(3-pyridylmethyl)-1,4-dioxo-8-azacyclododec-6-yl]-2-pyridinecarboxamide, 9CI. U 24544. Antibiotic U 24544. Erizomycin

[18791-21-4]
[11052-02-1]



C₂₇H₃₂N₄O₈ 540.572

Macrolide antibiotic. Prod. by *Streptomyces pyridomyceticus*, *Streptomyces griseus* var. *erizensis* and *Streptomyces albidofuscus*. Cytotoxic and antibacterial agent. Needles (EtOH). Sol. MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 221-222°. $[\alpha]_{\text{D}}^{25}$ -82 (c, 1.35 in EtOH). λ_{\max} 227 (ϵ 35000); 303 (ϵ 13500) (EtOH/HCl) (Derep). λ_{\max} 226 (ϵ 36000); 303 (ϵ 17900); 331 (ϵ 8960) (EtOH/NaOH) (Derep). λ_{\max} 303 (ϵ 11300) (EtOH) (Derep).

► LD₅₀ (mus, ivn) 300 mg/kg, LD₅₀ (mus, orl) 1000 mg/kg. UV1130000

Hydrochloride:

Cryst. Mp 194-196°. $[\alpha]_{\text{D}}^{16}$ -53.2 (H₂O).

Hydrobromide (1:2):

Cryst. + H₂O (EtOAc). Mp 203-205° dec.

2',3'-Dihydro, 3' ζ -hydroxy: **Pyridomycinol**

C₂₇H₃₄N₄O₉ 558.587

Prod. by *Streptomyces* sp. Stamm Ku2. Mp 108°. $[\alpha]_{\text{D}}^{20}$ -52 (c, 1 in MeOH).

λ_{\max} 256 (log ϵ 0.14); 262 (log ϵ 0.15); 303 (log ϵ 0.25) (MeOH). λ_{\max} 261 (log ϵ 0.18); 268 (log ϵ 0.12); 335 (log ϵ 0.27) (MeOH/NaOH).

Maeda, K. et al., *J. Antibiot.*, Ser. A, 1957, **10**, 94 (*isol*)

Herr, R.R. et al., *Appl. Microbiol.*, 1967, **15**, 1142 (*isol*)

Reusser, F. et al., *J. Bacteriol.*, 1967, **94**, 1040 (*props*)

Koyama, G. et al., *Tet. Lett.*, 1967, 3587 (*struct*, *cryst struct*, *abs config*)

Ogawara, H. et al., *Biochemistry*, 1968, **7**, 3296-3302 (*biosynth*)

U.S. Pat., 1968, 3 367 833; CA, **71**, 122302 (*Erizomycin*)

Kinoshita, M. et al., *Bull. Chem. Soc. Jpn.*, 1975, **48**, 2081 (*synth*)

Kinoshita, M. et al., *Tet. Lett.*, 1989, **30**, 7419 (*synth*)


Schulze, A. et al., *Dissertation*, Univ. of Göttingen, 2003, (*Pyridomycinol*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PPI775

9H-Pyrido[4,3,2-mn]thiazolo[4,5-b]acridin-9-one, 9CI P-920

Kuanoniamine A

[133401-10-2]



C₁₆H₇N₃OS 289.317


Alkaloid from a tunicate and its predator mollusc *Chelynotus semperi*. Also from *Oceanapia sagittaria*. Cytotoxic. Yellow needles (CHCl₃). Mp 255-258° dec. λ_{\max} 214 (ϵ 18200); 224 (ϵ 19100); 250 (ϵ 15800); 258 (ϵ 15500); 295 (ϵ 6310); 354 (ϵ 5750); 394 (ϵ 4070) (MeOH) (Derep). λ_{\max} 208 (ϵ 17200); 226 (ϵ 17780); 292 (ϵ 15000); 382 (ϵ 5150); 430 (MeOH/NaOH) (Berdy). λ_{\max} 480 (MeOH/HCl) (Berdy).

Carroll, A.R. et al., *J.O.C.*, 1990, **55**, 4426-4431 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Kitahara, Y. et al., *Tetrahedron*, 1997, **53**, 17029-17038 (*synth*)

Kijjoo, A. et al., *Mar. Drugs*, 2007, **5**, 6-22 (*isol*, *activity*)

Pyridovericin P-921



C₂₁H₂₃NO₅ 369.416

Related to Tenellin, T-72. Prod. by the fungus *Beauveria bassiana*. Pale yellow powder. Sol. MeOH, DMSO, EtOAc, Me₂CO; poorly sol. H₂O, hexane. Mp 203-206° dec. $[\alpha]_{\text{D}}^{15}$ -20.3 (c, 0.1 in MeOH). λ_{\max} 248 (ϵ 16400); 336 (ϵ 15900) (MeOH). λ_{\max} 248 (ϵ 16400); 336 (ϵ 15900) (MeOH) (Berdy).

Takahashi, S. et al., *J. Antibiot.*, 1998, **51**, 596-598; 1051-1054 (*isol*, *pmr*, *cmr*, *uv*)

Zhang, Q. et al., *J.A.C.S.*, 2002, **124**, 5774-5781 (*synth*)

Irlapati, N.R. et al., *Tetrahedron*, 2004, **60**, 9307-9317 (*synth*)


Pyridoxatin P-922

3-(2-Ethenyl-4,6-dimethylcyclohexyl)-1,4-dihydroxy-2(1H)-pyridinone, 9CI.

Tolypocin

[135529-30-5]

[149196-98-5]



C₁₅H₂₁NO₃ 263.336
 Prod. by the fungus *Acremonium* sp. BX86 and *Tolypocladium geodes*. Free radical scavenger. Inhibitor of gelatinase A. Cytotoxic agent. Needles (Me₂CO/EtOAc). Sol. MeOH, EtOAc, Me₂CO; poorly sol. H₂O. Mp 195-197° dec. [α]_D²³ -23 (c, 0.09 in MeOH). λ_{max} 216 (ε 10200); 246 (ε 3900); 267 (ε 3600) (MeOH/HClaq) (Derep). λ_{max} 230 (ε 15900); 262 (ε 4700); 295 (ε 3600) (MeOH/NaOH) (Derep). λ_{max} 218 (ε 11600); 288 (ε 4700) (MeOH) (Derep).

Fe complex: Terricolin

[149153-43-5]
 [149250-21-5]
 C₄₅H₆₀FeN₃O₉ 842.831
 Prod. by *Tolypocladium terricola*. Red plates + 2MeOH (MeOH). Dec. at 220°. λ_{max} 221 (ε 94500); 250 (sh) (ε 20000); 288 (ε 11100); 390 (ε 5100) (MeOH/HCl aq.).


Teshima, Y. *et al.*, *J. Antibiot.*, 1991, **44**, 685 (isol, pmr, cmr)

Jegorov, A. *et al.*, *J.C.S. Dalton*, 1993, 1287-1293 (*Tolypocin*, *Terricolin*, *cryst struct*)

Lee, H.-J. *et al.*, *J. Membr. Biol.*, 1996, **6**, 445-450 (isol, props)

Pyridoxine, BAN, INN P-923

5-Hydroxy-6-methyl-3,4-pyridinedimethanol, 9CI. Vitamin B₆, 8CI. 3-Hydroxy-4,5-bis(hydroxymethyl)-2-methylpyridine. Adermin. Pyridoxol. Many other names [65-23-6]



C₈H₁₁NO₃ 169.18
 Found in rice husks, cane molasses, yeast, wheat germ and cod liver oils. Vitamin, dietary supplement, nutrient. Needles (Me₂CO). Mp 160°. pK_{a1} 5; pK_{a2} 8.96 (25°, 0.15M NaCl). Log P -0.7 (calc). Readily sublimates.

- ▶ LD₅₀ (rat, orl) 4000 mg/kg. UV1300000
Hydrochloride: Pyridoxine hydrochloride, U.S.A.N. Aderoxin. Becilan. Gravidox [58-56-0]
 Rhombic cryst. (EtOH/Me₂CO). Mp 206-208°. Component of Bendectin and Debendox.
- ▶ Long-term treatment with large doses associated with development of peripheral neuropathies. Human and exp. reprod. effects. LD₅₀ (rat, orl) 4000 mg/kg. UV1350000
Picrate: Mp 156°.

O-(7-Theophyllinylethyl)sulfate (1:1):

Pyridofylline, INN. Atherophylline. Theodoxine
 [53403-97-7]
 Respiratory stimulant, coronary vasodilator. Cryst. (EtOH). Mp 144-146°.

- ▶ LD₅₀ (mus, orl) 1600 mg/kg. UO8430000
O^{1'}-Phosphate: Pyridoxine 5'-phosphate. Pyridoxol dihydrogen phosphate
 [447-05-2]

C₈H₁₂NO₆P 249.16
 Prod. by fungi and bacteria from pyridoxine. Cryst. (EtOH aq.). Mp 208-210°.

Tri-Ac: [10030-93-0]

C₁₄H₁₇NO₆ 295.291
 Viscous oil. Bp_{0.2} 145-150°.

Tribenzoyl:

C₂₉H₂₃NO₆ 481.504
 Mp 121-122°.

5-Me ether: [633-72-7]

C₉H₁₃NO₃ 183.207
 Needles (CHCl₃/petrol). Mp 89.5-90°.

5-Me ether, hydrochloride: [62511-98-2]
 Mp 150°.

O^{1''}-Me: 5-Hydroxy-4-(methoxymethyl)-6-methyl-3-pyridinemethanol, 9CI. 3-Hydroxy-5-hydroxymethyl-4-methoxymethyl-2-methylpyridine. 4'-Methoxy-pyridoxine. Ginkgotoxin
 [1464-33-1]

C₉H₁₃NO₃ 183.207
 Isol. from seeds of maidenhair tree *Ginkgo biloba*. Sitotoxic (convulsive) agent with antivitamin B₆ activity. Mp 181° (as hydrochloride).

- ▶ LD₅₀ (mus, orl) 32 mg/kg. UT5009000

O^{1''}-Me, 5-O-β-D-glucopyranoside:

[121531-33-7]
 C₁₅H₂₃NO₈ 345.349
 Isol. from the seeds of *Albizia lucida*. Neurotoxin.

O^{1''}-Me, O^{2''}-[β-D-glucuronopyranosyl-(1→2)-β-D-glucopyranoside]: Julibrine I

C₂₁H₃₁NO₁₄ 521.474
 Isol. from the stem bark of *Albizia julibrissin*. Powder. [α]_D -20 (MeOH).

O^{1''}-Me, O^{2''}-[β-D-apiofuranosyl-(1→2)-β-D-glucopyranoside]: Julibrine II

C₂₀H₃₁NO₁₂ 477.464
 From the stem bark of *Albizia julibrissin*. Shows arrhythmic-inducing activity. Powder. [α]_D -65.2 (MeOH).

O^{1''}-Me, O^{2''}-Ac: [82470-47-1]

C₁₁H₁₅NO₄ 225.244
 Constit. of *Albizia tanganyicensis*.

Dihydroxyacetic acid monoether: Piridoxilate, BAN, INN. Pyridoxylate. Nargin. Venartan

[24340-35-0] Antianginal agent. Mp 167°. Log P -1.16 (calc). A 1:1 complex of the 3 and 1' positional isomers. Exists as zwitterions.

[79636-52-5, 63797-00-2, 8064-77-5, 70829-29-7, 62154-08-9, 74536-44-0, 62602-94-2, 3131-27-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 759D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 282B; 283A (nmr)

Harris, S.A. *et al.*, *J.A.C.S.*, 1939, **61**, 1245; 3307 (synth)

Mowatt, J.H. *et al.*, *J.A.C.S.*, 1943, **65**, 954 (synth)

Fr. Pat., 1961, M828; *CA*, **58**, 9098g (Pyridofylline)

Ogata, K. *et al.*, *Agric. Biol. Chem.*, 1966, **30**, 829 (1''-phosphate)

De Jongh, D.C. *et al.*, *J.A.C.S.*, 1966, **88**, 1233 (ms)

Firestone, R.A. *et al.*, *Tetrahedron*, 1967, **23**, 943 (synth, ir, uv)

Efimovsky, O. *et al.*, *Eur. J. Med. Chem. (Chim. Ther.)*, 1969, **4**, 89 (*Piridoxilate, synth, struct*)

Fourneau, J.P. *et al.*, *Boll. Chim. Farm.*, 1970, **109**, 702 (rev)

Witherup, T.H. *et al.*, *J.O.C.*, 1975, **40**, 2229 (cmr)

Hill, R.E. *et al.*, *J.A.C.S.*, 1977, **99**, 4179 (*biosynth, cmr*)

Coffen, D.L. *et al.*, *Kirk-Othmer Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **24**, 94 (rev)

Longo, J. *et al.*, *Acta Cryst. B*, 1982, **38**, 2721 (*cryst struct*)

Aboul-Enain, H.Y. *et al.*, *Anal. Profiles Drug Subst.*, 1984, **13**, 447 (rev, synth, anal, pharmacol)

Shimada, S. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 38 (synth)

Wada, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3555; 1988, **36**, 1779 (*Ginkgotoxin*)

Parnell, C.A. *et al.*, *Tetrahedron*, 1985, **41**, 5791 (synth)

Likos, J.J. *et al.*, *Vitamin B6 Pyridoxal Phosphate: Chemical, Biochemical and Medical Aspects, Part A*, (ed. Dolphin, D. *et al.*), Wiley-Interscience, 1986, 13 (rev)

Korpela, T. *et al.*, *Biochemistry of Vitamin B6*, Birkhäuser Verlag, 1987, (book)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 951

Steyn, P.S. *et al.*, *S. Afr. J. Chem.*, 1987, **40**, 191 (*deriv*)

Lewis, R.J. *et al.*, *Food Additives Handbook*, Van Nostrand Reinhold International, New York, 1989, PPK500

Orsini, F. *et al.*, *Gazz. Chim. Ital.*, 1989, **119**, 63 (5-glucoside)

Dakshinamurti, K. *et al.*, *Ann. N.Y. Acad. Sci.*, 1990, **585**, 1 (revs)

Lycka, A. *et al.*, *Pharmazie*, 1990, **45**, 371 (N-15 nmr)

Higuchi, H. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 829 (*Julibrine*)

Martindale. The Extra Pharmacopoeia, 30th edn., Pharmaceutical Press, 1993, 1054; 1318; 1403

Spenser, I.D. *et al.*, *Nat. Prod. Rep.*, 1995, **12**, 555 (rev, biosynth)

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2389; 2391 (rev)

Fiehe, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 185-189 (*Ginkgotoxin, biosynth*)

Kall, M.A. *et al.*, *Food Chem.*, 2003, **82**, 315-327 (*detn, occur*)

Zeidler, J. *et al.*, *J.O.C.*, 2003, **68**, 3486-3493 (*biosynth*)


Dumond, Y.R. *et al.*, *Molecules*, 2003, **8**, 873-881 (synth)

Roje, S. *et al.*, *Phytochemistry*, 2007, **68**, 1904-1921 (*biosynth, rev*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MEY000; PPK250; PPK500; MFN600

Pyrifolidine P-924

1-Acetyl-16,17-dimethoxyaspidospermidine, 9CI
 [639-26-9]



C₂₃H₃₂N₂O₃ 384.517

(+)-form

Alkaloid from *Aspidosperma pyrifolium* (Apocynaceae). Cryst. (MeOH aq.). Mp 147.5-150°. $[\alpha]_D^{25} +90$ (c, 1.04 in CHCl₃). λ_{\max} 223 (log ϵ 4.5); 252 (log ϵ 3.94); 286 (log ϵ 3.32) (EtOH).

Perchlorate:

Cryst. (MeOH aq.). Mp 255-260.5°.

N-De-Ac: Deacetylpyrifolidine. O-

Methyl-N-depropionylaspidolimine

[25139-33-7]

C₂₁H₃₀N₂O₂ 342.48

Minor alkaloid from *Aspidosperma neblinae* (Apocynaceae). Mp 147-149°. $[\alpha]_D^{25} +6.7$ (c, 0.89 in CHCl₃). λ_{\max} 215 (log ϵ 4.4); 293 (log ϵ 3.5) (EtOH).

(-)-form

Alkaloid from *Aspidosperma quebrachoblanco* (quebracho) (Apocynaceae). Mp 148-150°. $[\alpha]_D^{28} -93$ (c, 0.9 in CHCl₃).

O¹²-De-Me: Aspidocarpine

[466-45-5]

C₂₂H₃₀N₂O₃ 370.491

Alkaloid from *Geissospermum argenteum*, *Aspidosperma meglacarpum*, *Aspidosperma formasanum*, *Aspidosperma marcgravianum*, *Aspidosperma neblinae* and *Aspidosperma album* (Apocynaceae). Prisms (MeOH or heptane). Mp 168.5-169.5°. $[\alpha]_D^{25} +140$ (c, 2.3 in CHCl₃). pK_a 6.55.

O¹²-De-Me, hydrochloride:

Needles (EtOH/EtOAc). Mp 224-230° dec.

O¹²-De-Me, N-de-Ac, N-propanoyl: Aspidolimine

[5516-66-5]

C₂₃H₃₂N₂O₃ 384.517

Alkaloid from *Aspidosperma limae* and *Aspidosperma obscurinervium* (Apocynaceae). Cryst. (MeOH). Mp 150-152°. $[\alpha]_D^{21} +133$ (c, 1.156 in CHCl₃).

O¹²-De-Me, N-de-Ac, N-propanoyl, hydrochloride:

Cryst. (EtOAc/Et₂O). Mp 200-205° dec.

Di-O-de-Me: O-Demethylaspidocarpine

[21401-34-3]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Aspidosperma album*, *Aspidosperma cuspa* and *Aspidosperma melanocalyx* (Apocynaceae). Cryst. (C₆H₆). Mp 130° Mp 156-158°.

Di-O-de-Me; hydrochloride:

Needles (EtOH/Et₂O). Mp 283° dec.

McLean, S. *et al.*, *Can. J. Chem.*, 1960, **38**, 1547 (*Aspidocarpine, O-Demethylaspidocarpine*)

Gilbert, B. *et al.*, *Experientia*, 1960, **16**, 61 (*isol, uv, ir*)

Djerassi, C. *et al.*, *Experientia*, 1961, **17**, 162 (*Deacetylpyrifolidine*)

Djerassi, C. *et al.*, *Tetrahedron*, 1961, **16**, 212 (*isol, uv, ord*)

Gilbert, B. *et al.*, *Chem. Ind. (London)*, 1962, 1949 (*isol*)

Pinar, M. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 1283 (*isol, uv, ir, struct*)

Ferrari, C. *et al.*, *Can. J. Chem.*, 1963, **41**, 1531 (*isol, struct, ir, Aspidocarpine, O-Demethylaspidocarpine*)

Biemann, K. *et al.*, *J.A.C.S.*, 1963, **85**, 631 (*isol, uv, ms*)

Brown, K.S. *et al.*, *J.A.C.S.*, 1964, **86**, 2451 (*isol*)

Klyne, W. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 833 (*ord, abs config*)


Burnell, R.H. *et al.*, *Phytochemistry*, 1968, **7**, 2045 (*Pyrifolidine, O-Demethylaspidocarpine, isol, synth, uv, ir, pmr*)

Miranda, E.C. *et al.*, *Experientia*, 1969, **25**, 575 (*O-Demethylaspidocarpine*)

Lyon, R.L. *et al.*, *J. Pharm. Sci.*, 1973, **62**, 218 (*isol, uv, ir, pmr*)

Garcia, M. *et al.*, *Phytochemistry*, 1976, **15**, 1093 (*isol, uv, ir, pmr, ms*)

Paccioni, J.-P. *et al.*, *Phytochemistry*, 1978, **17**, 2146 (*isol*)

Pyrinadine A**P-925**

X = $-\text{CH}_2\text{CH}_2\text{CH}=\overset{Z}{\text{C}}\text{H}(\text{CH}_2)_{10}-$

Y = $-(\text{CH}_2)_{10}\overset{Z}{\text{C}}\text{H}=\text{CHCH}_2\text{CH}_2-$

C₃₈H₆₀N₄O 588.918

Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. λ_{\max} 213 (€ 10600); 251 (€ 3600); 257 (€ 4200); 262 (€ 4700); 269 (€ 3700) (MeOH).

Kariya, Y. *et al.*, *Tet. Lett.*, 2006, **47**, 997-998 (*isol, pmr, cmr*)

Pyrinadine B**P-926**

[918302-92-8]

As Pyrinadine A, P-925 with

X = Y = $-(\text{CH}_2)_{13}-$

C₃₆H₆₀N₄O 564.896

Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. λ_{\max} 214 (€ 11700); 250 (€ 4500); 257 (€ 4700); 262 (€ 4900); 269 (€ 3500) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyrinadine C**P-927**

[918302-80-4]

As Pyrinadine A, P-925 with

X = $-(\text{CH}_2)_{13}-$, Y = $-(\text{CH}_2)_{12}-$

C₃₅H₅₈N₄O 550.869

N-Oxide functionality may be on the other azine N. Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. λ_{\max} 214 (€ 9200); 250 (€ 4900); 256 (€ 4700); 262 (€ 4700); 269 (€ 3400) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyrinadine D**P-928**

[918302-85-9]

As Pyrinadine A, P-925 with

X = $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_2)_{10}-$, Y = $-(\text{CH}_2)_{13}-$

C₃₇H₆₀N₄O 576.907

N-Oxide functionality may be on the other azine N. Alkaloid from

Cribrachalina sp. (SS-1115). Cytotoxic. Oil. X-linker chain has Z-configuration. λ_{\max} 213 (€ 9900); 251 (€ 3600); 257 (€ 4200); 262 (€ 4600); 269 (€ 3400) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyrinadine E**P-929**

[918302-88-2]

As Pyrinadine A, P-925 with

X = $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_2)_{10}-$, Y = $-(\text{CH}_2)_{12}-$

C₃₆H₅₈N₄O 562.88

N-Oxide functionality may be on the other azine N. Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. X-linker chain has Z-configuration. λ_{\max} 213 (€ 10100); 250 (€ 4100); 257 (€ 4400); 262 (€ 4600); 269 (€ 3300) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyrinadine F**P-930**

[918302-90-6]

As Pyrinadine A, P-925 with

X = $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_2)_{10}-$, Y = $-(\text{CH}_2)_3\text{CH}=\text{CH}(\text{CH}_2)_6-$

C₃₈H₆₀N₄O 588.918

N-Oxide functionality may be on the other azine N. Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. X- and Y-linker chains have Z-configuration. λ_{\max} 215 (€ 9300); 251 (€ 3700); 257 (€ 4400); 263 (€ 4800); 269 (€ 3500) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyrinadine G**P-931**

[918302-91-7]

As Pyrinadine A, P-925 with

X = $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_2)_{10}-$, Y = $-(\text{CH}_2)_8\text{CH}=\text{CH}(\text{CH}_2)_4-$


C₃₈H₆₀N₄O 588.918

N-Oxide functionality may be on the other azine N. Alkaloid from *Cribrachalina* sp. (SS-1115). Cytotoxic. Oil. X- and Y-linker chains have Z-configuration. λ_{\max} 213 (€ 10300); 251 (€ 4600); 257 (€ 4600); 262 (€ 4800); 269 (€ 3700) (MeOH).

Kariya, Y. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8415-8419 (*isol, pmr, cmr, ms*)

Pyridicine**P-932**

5-(1-Butenyl)-3,4,6,7-tetrahydro-3-methyl-2H-cyclopenta[b]pyridine, 9CI [50719-83-0]



C₁₃H₁₉N 189.3

Metab. of *Streptomyces griseofulvus* var. *pyrindicus*. Weak antibiotic, active against gram-positive and -negative bacteria. Pale yellow solid. [α]_D²¹ -38.8 (c, 0.05 in H₂O). λ_{\max} 311 (ε 41500) (MeOH) (Derep).

Hydrochloride: Mp 145°. [α]_D²⁰ +22.9 (c, 0.7 in H₂O).


Picrate: Mp 174-176° dec.

Methiodide: Mp 187-188°.

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1973, **21**, 2048 (*uv, pmr, ms, struct*)

Iwai, Y. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 736 (*cmr, biosynth*)

Hegde, V.R. *et al.*, *J. Antibiot.*, 1994, **47**, 110 (*isol, pmr, cmr*)

Pyrinodemin A**P-933**C₃₈H₅₉N₃O 573.904

Isol. from the sponge *Amphimedon* sp. Cytotoxic agent. Oil. [α]_D²⁵ -9 (c, 1 in CHCl₃). λ_{\max} 264 (ε 6300) (MeOH).

Tsuda, M. *et al.*, *Tet. Lett.*, 1999, **40**, 4819-4820 (*isol, uv, ir, pmr, cmr, ms*)

Snider, B.B. *et al.*, *Tet. Lett.*, 2001, **42**, 1639-1642 (*synth*)

Romeril, S.P. *et al.*, *Tet. Lett.*, 2002, **43**, 327-329; 2003, **44**, 7757-7761 (*synth, abs config*)


Mormoto, Y. *et al.*, *Org. Lett.*, 2003, **5**, 2611-2614 (*synth, abs config*)

Ishiyama, H. *et al.*, *Molecules*, 2005, **10**, 312-316 (*synth, struct*)

Pouilhès, A. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 1502-1510 (*synth*)

Pyrinodemin B**P-934**

[292041-81-7]

C₃₇H₅₉N₃O 561.893

Isol. from *Amphimedon* sp. Cytotoxic agent. λ_{\max} 264 (ε 6000) (MeOH).

4',5'-Didehydro (Z-): Pyrinodemin C
[292041-82-8]

C₃₇H₅₇N₃O 559.877

Isol. from *Amphimedon* sp. Cytotoxic agent. λ_{\max} 263 (ε 5800) (MeOH).

Lower homologue (n = 7): Pyrinodemin D
[292041-83-9]

C₃₆H₅₇N₃O 547.866

Isol. from *Amphimedon* sp. Cytotoxic agent. λ_{\max} 264 (ε 3300) (MeOH).

Hirano, K. *et al.*, *Chem. Pharm. Bull.*, 2000,


48, 974-977 (*Pyrinodemins B-D, isol, pmr, cmr, uv*)

Snider, B.B. *et al.*, *Tet. Lett.*, 2001, **42**, 1639-1642 (*synth*)

Romeril, S.P. *et al.*, *Tet. Lett.*, 2003, **44**, 7757-7761 (*config*)

Pyrizinostatin**P-935**

2,4,4a,8-Tetrahydro-2,6,8-trimethyl-4a-(2-oxo propyl)pyrimido[5,4-e]-1,2,4-triazine-3,5,7(6H)-trione, 9CI
[146406-84-0]



Relative configuration

C₁₁H₁₅N₅O₄ 281.271

Prod. by *Streptomyces* sp. Pyroglutamyl peptidase inhibitor. Cryst (MeOH). Sol. H₂O, CHCl₃, DMSO, MeOH; poorly sol. EtOAc, hexane, Et₂O. Mp 188-190°. [α]_D²⁴ -15.6 (c, 1 on MeOH). λ_{\max} 280 (ε 4600) (MeOH) (Derep).


Aoyagi, T. *et al.*, *J. Antibiot.*, 1992, **45**, 1795; 1961 (*isol, struct, props*)

Kitagawa, M. *et al.*, *Heterocycles*, 1994, **38**, 1747 (*synth*)

Tatsuka, K. *et al.*, *J. Antibiot.*, 1994, **47**, 389 (*synth*)

Pyrodisynoic acid**P-936**

[385801-57-0]



Relative Configuration


C₁₇H₂₃NO₃ 289.374

Isol. from a *Dysidea* sp. Pale yellow oil. [α]_D²³ +9.1 (c, 0.07 in MeOH). λ_{\max} 204 (log ε 1.46) (MeOH).

Goetz, G.H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1486-1488 (*isol, pmr, cmr*)

Pyroglutamylglucosamine**P-937**

2-Deoxy-2-(5-oxo-2-pyrrolidinecarboxamido)glucopyranose, 9CI
[29227-87-0]


C₁₁H₁₈N₂O₇ 290.272

Isol. from the field mushroom *Agaricus campestris*. Tentative identification.

Altamura, M.R. *et al.*, *J. Food Sci.*, 1970, **35**, 134-139 (*isol*)

Pyronaamidine**P-938**

[124535-77-9]

C₂₅H₂₇N₅O₆ 493.518

Alkaloid from a Micronesian sponge, *Leucetta* sp. Cytotoxic against KB cells (MIC 5 μg/mL). Yellow feathery cryst. Mp 185-187°. λ_{\max} 208 (ε 49000); 224 (ε 34700); 276 (ε 6030); 388 (ε 16600) (MeOH) (Derep).

19-Hydroxy: Naamidine E

[152273-84-2]

C₂₅H₂₇N₅O₇ 509.518

Alkaloid from *Leucetta* sp. and *Notodoris gardineri*. Yellow amorph. solid. Mp 115-118°.

9-(N-Methylimide): Pyronaamidine 9-N-methylimide

[191362-11-5]

C₂₆H₃₀N₆O₅ 506.56

Alkaloid from *Leucetta* sp. cf. *chagosensis*. Yellow cryst. (CH₂Cl₂/MeOH). Mp 222-225°. λ_{\max} 211 (5.14); 274 (4.78); 388 (4.96) (MeOH).

Akee, R.K. *et al.*, *J.O.C.*, 1990, **55**, 1944 (*isol, uv, ir, pmr, cmr, ms, struct*)

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1993, **46**, 1229 (*Naamidine E*)


Plubrukarn, A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 712 (*9-methylimide*)

Nakamura, S. *et al.*, *J.C.S. Perkin 1*, 2002, 1061-1066 (*synth, pmr*)

Pyronopaline**P-939**

[86630-17-3]

[141032-89-5]

C₁₁H₁₈N₄O₅ 286.287


Opine. Isol. from crown gall tumour. Cryst. (EtOH aq.). Mp 245° (dec.). [α]_D²² -40 (c, 0.5 in H₂O). Possible artifact.

Hall, L.M. *et al.*, *J. Biol. Chem.*, 1983, **258**, 7276-7279 (*isol, pmr, struct*)

U.S. Pat., 1992, 5 081 289; *CA*, **116**, 210751k (*synth*)

Pyrrocidine B**P-940**

[428439-25-2]



C₃₁H₃₉NO₄ 489.633
 Prod. by a fungus, LL-Cyan 426. Active against gram-positive bacteria. Yellowish amorph. powder. [α]_D +88.2 (c, 0.22 in MeOH).

17,18-Didehydro: **Pyrrocidine A**

[428439-24-1]

C₃₁H₃₇NO₄ 487.638


Prod. by a fungus, LL-Cyan426. Active against gram-positive bacteria. Yellowish amorph. powder. [α]_D +91.2 (c, 0.58 in MeOH).

He, H. et al., *Tet. Lett.*, 2002, **43**, 1633-1636 (isol, pmr, cmr)

Pyrroindomycin A

P-941

[160472-94-6]



C₆₀H₈₀N₄O₁₅ 1097.31

Prod. by a strain of *Streptomyces rugosporus*. Active against gram-positive bacteria. λ_{max} 226 (ε 26100); 273 (ε 18500); 335 (ε 26200) (MeOH) (Derep).

5'-Chloro: **Pyrroindomycin B**

[160472-95-7]

C₆₀H₇₉ClN₄O₁₅ 1131.755


Prod. by a strain of *Streptomyces rugosporus*. Active against gram-positive bacteria. λ_{max} 229 (ε 26100); 281 (ε 23200); 335 (ε 25600) (MeOH) (Derep).

Ding, W. et al., *J. Antibiot.*, 1994, **47**, 1249-1257; 1258-1265 (isol, pmr, cmr, ir, props)
 Abbanat, D. et al., *J. Antibiot.*, 1999, **52**, 117-126 (biosynth)

1H-Pyrrole-2-carboxaldehyde, 9CI

P-942

Pyrrole-2-aldehyde. 2-Formylpyrrole
 [1003-29-8]



C₅H₅NO 95.101

Constit. of numerous plant spp. incl. tea, coffee and various legumes. Prisms (petrol). Mp 50-51°. Bp 217-219° Bp₂ 78°.

Oxime: [32597-34-5]

C₅H₆N₂O 110.115

Needles (CHCl₃ or C₆H₆). Mp 165-166°.

Phenylhydrazone: [81280-72-0]

Needles (petrol). Mp 139-139.5°.

4-Nitrophenylhydrazone:

Red needles (xylene). Mp 182.5-183°.

Ethylene dithioketal: 2-(2-Pyrrolyl)-1,3-dithiolane. 2-(1,3-Dithiolan-2-yl)-1H-pyrrole, 9CI

[70582-57-9]

C₇H₉NS₂ 171.287

Light yellow oil. Mp 27-30°.

N-Ac: [30186-49-3]

C₇H₇NO₂ 137.138

Plates (hexane) cryst. (Et₂O/petrol). Mp 78-78.5°.

N-Benzoyl:

C₁₂H₉NO₂ 199.209

Cryst. (petrol). Mp 90°.

N-Me: 1-Methyl-2-pyrrolicarboxaldehyde

[1192-58-1]

C₆H₇NO 109.127

Oil with odour resembling benzaldehyde. Bp₃₂ 94-96° Bp₁₁ 75-76°. On boiling rapidly turns yellow, then purple, finally deep red. Stable in alkali resinifies with acids.

► UX9350000

N-Me, oxime:

C₆H₈N₂O 124.142

Mp 152°.

N-Me, semicarbazone:

Cryst. (EtOH). Mp 207-208°.

N-Me, phenylhydrazone: Mp 127-128°.

N-Et: 1-Ethyl-1H-pyrrole-2-carboxaldehyde

[2167-14-8]

C₇H₉NO 123.154

Light yellow oil. Bp₇ 73-75°.

N-Et, phenylhydrazone: Mp 75°.

N-Butyl: [17164-95-3]

C₉H₁₃NO 151.208

Oil. Bp₁₁ 148-150°.

N-Benzyl: [18159-24-5]

C₁₂H₁₁NO 185.225

Oil. Bp_{0,1} 96-98°.

N-Ph: 1-Phenyl-1H-pyrrole-2-carboxaldehyde

C₁₁H₉NO 171.198

Yellow cryst. Mp 98°.

N-Hydroxymethyl: 1-Hydroxymethyl-2-pyrrolicarboxaldehyde

[116967-16-9]

C₆H₇NO₂ 125.127

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 568B; 568C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 6C; 7A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1448A; 1448B (ir)

Smith, G.F. et al., *J.C.S.*, 1954, 3842 (synth)
 Silverstein, R.M. et al., *J.O.C.*, 1955, **20**, 668 (synth)

Fournari, P. et al., *Bull. Soc. Chim. Fr.*, 1963, 488 (I-Me)

Org. Synth., Coll. Vol., **4**, 1963, 831 (synth)
 Khan, M.K.A. et al., *J.C.S.*, 1964, 2579 (ir)
 Jones, R.F. et al., *J.C.S.(C)*, 1970, 2563 (synth)

Matsuo, T. et al., *Bull. Chem. Soc. Jpn.*, 1972, **45**, 1349 (uv)

Chadwick, D.J. et al., *Tet. Lett.*, 1974, 3183 (nmr)

Cushley, R.J. et al., *Can. J. Chem.*, 1975, **53**, 148 (cmr)

Severin, T. et al., *Chem. Ber.*, 1975, **108**, 1768 (I-Me)

Limage, M.H. et al., *J. Chim. Phys. Phys.-Chim. Biol.*, 1975, **72**, 914 (ir, Raman)

Nikisch, K. et al., *Chem. Ber.*, 1980, **113**, 2036 (synth, ir, pmr)

Comins, D.L. et al., *J.O.C.*, 1987, **52**, 104 (Et)
 Boykin, D.W. et al., *J. Het. Chem.*, 1988, **25**, 643 (O-17 nmr)

Katritzky, A.R. et al., *Magn. Reson. Chem.*, 1988, **26**, 129-133 (N-hydroxymethyl, synth, pmr, cmr)

Katritzky, A.R. et al., *Org. Prep. Proced. Int.*, 1988, **20**, 585 (synth)

Pindur, U. et al., *J. Het. Chem.*, 1989, **26**, 1563 (synth, ir, ms, pmr)

Wallace, D.M. et al., *J.O.C.*, 1993, **58**, 7245 (synth, deriv, pmr)

Dumoulin, H. et al., *J. Het. Chem.*, 1995, **32**, 1703 (N-Ph)


1H-Pyrrole-3-carboximide, 9CI

P-943

Pyrrole-3-carbamidine. 3-Amidinopyrrole.

Brunfelsamine. Hopamidine

[97744-98-4]



C₅H₇N₃ 109.13

Alkaloid from the root bark of *Brunfelsia grandiflora* ssp. *schultesii* and from *Leptonychia pubescens* and *Nierembergia hippomanica*. Shows convulsant props., toxic to livestock. Hygroscopic solid. Sol. MeOH, H₂O; fairly sol. butanol; poorly sol. CHCl₃, hexane. λ_{max} 232 ; 262 (MeOH) (Berdy).

► Toxic.

Hydrochloride: Mp 127-128°.

Picrate:

Cryst. (EtOH). Mp 247-249°.

1-β-D-Ribofuranosyl: 1-β-D-Ribofuranosylbrunfelsamine

[117654-70-3 (HCl salt)]

C₁₀H₁₅N₃O₄ 241.246

Alkaloid from *Leptonychia pubescens*. Amorph. solid. [α]_D³⁰ -1 (c, 0.5 in H₂O). λ_{max} 199 (log ε 4.05); 236 (log ε 3.18) (H₂O).

Lloyd, H.A. et al., *Tet. Lett.*, 1985, **26**, 2623-2624 (uv, pmr, cmr, ms)

Buschi, C.A. et al., *Phytochemistry*, 1987, **26**, 863-865 (isol, synth, ir, pmr, cmr, ms)


Yang, L.K. et al., *Planta Med.*, 2005, **71**, 1071-1072 (ribofuranosyl)

1H-Pyrrole-2-carboxylic acid, 9CI

P-944

Minaline

[634-97-9]



C₅H₅NO₂ 111.1

Occurs in various microorganisms, e.g. *Streptomyces griseoflavus*. Imino-acid occurring in diastase, prob. resulting

from dehydration of hydroxyprolines. Esters possess local anaesthetic props. Protein racemase inhibitor. Platelet aggregation inhibitor. Leaflets (H₂O). Sol. H₂O, EtOH, Et₂O. Mp 208.5° dec. pK_{a1} 1.48; pK_{a2} 4.5 (25°). Decarboxylates at Mp. Isoelectric point 2.9. λ_{max} 263 (MeOH) (Berdy).

α-L-Rhamnopyranosyl ester: 1H-Pyrrol-2-ylcarbonyl α-L-rhamnopyranoside
C₁₁H₁₅NO₆ 257.243
Prod. by *Streptomyces* sp. Mp 89° dec. [α]_D²⁰ -39 (c, 0.56 in MeOH). λ_{max} 210 (log ε 3.37); 267 (log ε 4.23) (MeOH).

(6-Deoxy-α-L-talopyranosyl) ester:
C₁₁H₁₅NO₆ 257.243
Prod. by *Streptomyces* sp. strain G6M1. Amorph. solid. [α]_D²⁰ -60 (c, 1 in MeOH). λ_{max} 231 (log ε 3.49); 267 (log ε 3.96) (MeOH).

Me ester: [1193-62-0]
C₆H₇NO₂ 125.127
Isol. from an *Agelas* sp. Needles (petrol). Sol. EtOH, Et₂O. Mp 73°. Bp₇₄₀ 220-223° Bp₁₄ 115-120° (lit. gives a pressure range).

Benzyl ester: [35889-87-3]
C₁₂H₁₁NO₂ 201.224
Pale yellow solid. Mp 55°.

Et ester: [2199-43-1]
C₇H₉NO₂ 139.154
Cryst. V. sol. Et₂O, EtOH, C₆H₆, petrol. Mp 41-42°. Bp 230-232°.

2-Phenylethyl ester: Phenethyl 2-pyrrole-carboxylate
[168324-00-3]
C₁₃H₁₃NO₂ 215.251
Alkaloid from the rhizome of *Sparganium stoloniferum*.

Chloride: [5427-82-7]
C₅H₄ClNO 129.545
Cryst. (Et₂O/petrol). V. sol. Et₂O, CHCl₃. Sinters at 110°, dec. at higher temps. Dec. rapidly in moist air.

Amide: 1H-Pyrrole-2-carboxamide
[4551-72-8]
C₅H₆N₂O 110.115
Alkaloid from the sponge *Agelas oroides*. Plates (H₂O or EtOH). Mp 176°. Sweet taste. λ_{max} 262 (ε 3700) (MeOH) (Berdy).

N-Formylamide: N-Formyl-1H-pyrrole-2-carboxamide, 9CI
[160156-23-0]
C₆H₆N₂O₂ 138.126
Alkaloid from the sponge *Agelas oroides*. Powder. λ_{max} 284 (ε 8140) (MeOH).

Hydrazide:
C₅H₇N₃O 125.13
Cryst. (EtOH aq.). Mp 231-232° dec.

Azide: [67307-19-1]
C₅H₄N₄O 136.113
Cryst. Mp 105° dec.

Nitrile: 1H-Pyrrole-2-carbonitrile, 9CI. 2-Cyanopyrrole
[4513-94-4]
C₅H₄N₂ 92.1
Oil. Bp₁₄ 118-120° Bp_{0.05} 50°.

N-Methoxycarbonyl, Me ester:
C₈H₉NO₄ 183.163

Cryst. (hexane). Mp 48-49.5° (37-40°). Bp₂₈ 145°.

N-Ethoxycarbonyl, Et ester: [66202-48-0]
C₁₀H₁₃NO₄ 211.217
Bp₈ 132-134° Bp₁ 86-87°.

N-tosyl, nitrile: [220105-73-7]
C₁₂H₁₀N₂O₂S 246.289
Cryst. Mp 114-115°.

N-Me: 1-Methyl-1H-pyrrole-2-carboxylic acid
[6973-60-0]
C₆H₇NO₂ 125.127
Needles (H₂O). Mp 135°.

N-Me, Me ester: [37619-24-2]
C₇H₉NO₂ 139.154
Bp₁₃ 86.5-87°.

N-Me, Et ester: [23466-27-5]
C₈H₁₁NO₂ 153.18
Bp_{0.5} 68°. n_D²⁰ 1.5043.

N-Me, nitrile: 2-Cyano-1-methyl-1H-pyrrole
[34884-10-1]
C₆H₆N₂ 106.127
Yellow oil. Bp₂₀ 60-62°.

N-Et: [4778-76-1]
C₇H₉NO₂ 139.154
Needles (H₂O). Mp 78°.

N-Ph: 1-Phenyl-1H-pyrrole-2-carboxylic acid, 9CI
[78540-03-1]
C₁₁H₉NO₂ 187.198
Cryst. (EtOH or C₆H₆). Mp 166° dec. Decarboxylates in boiling H₂O.

N-Ph, Me ester: [35524-54-0]
C₁₂H₁₁NO₂ 201.224
Cryst. (EtOH aq.). Mp 88°. Bp 282°.

N-Ph, Et ester: [24601-00-1]
C₁₃H₁₃NO₂ 215.251
Cryst. (EtOH aq.). Mp 88-89°. Bp 289°.

N-Ph, amide: [4778-75-0]
C₁₁H₁₀N₂O 186.213
Cryst. (C₆H₆). Mp 152-155°.

N-Ph, nitrile: 2-Cyano-1-phenyl-1H-pyrrole
C₁₁H₈N₂ 168.198
Solid (heptane). Mp 58-59°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 568D; 569A (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 7C; 8A (nmr)
Sadtler Standard C-13 NMR Spectra, 14815 (cmr)
Schwanert, H. et al., *Annalen*, 1860, 116, 257 (synth)

Pictet, A. et al., *Ber.*, 1902, 35, 2529 (N-Ph)
Blicke, F.F. et al., *J.A.C.S.*, 1930, 52, 238 (esters)

Treibs, A. et al., *Annalen*, 1958, 619, 80 (Et ester)
Doyle, F.P. et al., *J.C.S.*, 1958, 4458 (synth)
Fournari, P. et al., *Bull. Soc. Chim. Fr.*, 1963, 488 (N-Me)

Hodge, P. et al., *J.C.S.*, 1963, 2543 (synth)
Org. Synth., 1971, 51, 100 (Et ester)
Streith, J. et al., *Helv. Chim. Acta*, 1976, 59, 2786 (nitrile)

Wang, N.-C. et al., *Can. J. Chem.*, 1977, 55, 4103 (N-carboxy)
Lauti, A. et al., *Spectrochim. Acta A*, 1977, 33, 121 (ir)

Barnett, G.H. et al., *Can. J. Chem.*, 1980, 58, 409 (amide, nitrile)
Höfle, G. et al., *Annalen*, 1983, 835 (isol, pmr)

Katritzky, A.R. et al., *Org. Prep. Proced. Int.*, 1988, 20, 585 (synth, pmr)

Costa, C. et al., *Org. Mass Spectrom.*, 1991, 26, 972 (ms)

Wallace, D.M. et al., *J.O.C.*, 1993, 58, 7245 (Et ester)

Koenig, G.M. et al., *Nat. Prod. Lett.*, 1994, 5, 141-146 (isol, amides)

Niclas, H.-J. et al., *Synth. Commun.*, 1994, 24, 3241 (nitrile)

Miyaichi, Y. et al., *Nat. Med. (Tokyo)*, 1995, 49, 24-28 (phenethyl ester)

Hughes, T.V. et al., *J.O.C.*, 1999, 64, 313-315 (tosyl nitrile)

Grond, S. et al., *Eur. J. Org. Chem.*, 2000, 929-937; 1875-1881 (rhamnosyl ester)

Assman, M. et al., *Marine Ecol.: Progr. Ser.*, 2000, 207, 255-262 (*Agelas* constit)

Smaliy, R.V. et al., *Synthesis*, 2002, 2416-2420 (N-Me nitrile, N-Ph nitrile)

Dieter, A. et al., *J. Antibiot.*, 2003, 56, 639-646 (isol, pmr, cmr)


Bitzer, J. et al., *Eur. J. Org. Chem.*, 2006, 3661-3666 (6-deoxytalopyranosyl ester)

La Regina, G. et al., *J. Med. Chem.*, 2007, 50, 922-931 (N-Ph amide)

Dohi, T. et al., *J.O.C.*, 2007, 72, 109-116 (tosyl nitrile)

Schmuck, C. et al., *Eur. J. Org. Chem.*, 2008, 324-329 (Me ester, benzyl ester)

1H-Pyrrole-2,3-dicarboxylic acid, 9CI P-945 [1125-32-2]



C₆H₅NO₄ 155.11

Alkaloid from the leaves and flowers of *Parietaria officinalis* (Urticaceae). Needles. Mp 224° dec Mp 242-243° dec.

3-Me ester:
C₇H₇NO₄ 169.137
Needles (EtOH). Mp 200°.

3-Et ester:
C₈H₉NO₄ 183.163
Mp 145°.

Di-Me ester: [2818-08-8]
C₈H₉NO₄ 183.163
Mp 70°.

Di-Et ester: [25472-60-0]
C₁₀H₁₃NO₄ 211.217
Bp_{0.08} 110°.

N-Me:
C₇H₇NO₄ 169.137
Solid.

N-Me, di-Me ester: [25472-58-6]
C₉H₁₁NO₄ 197.19
Cryst. (Et₂O).

Nicolaus, R.A. et al., *Gazz. Chim. Ital.*, 1956, 86, 757 (synth)

Swan, G.A. et al., *J.C.S. (C)*, 1970, 285 (synth, derivs)

Budzianowski, J. et al., *Phytochemistry*, 1990, 29, 3299 (isol)

Woo, J. et al., *J.A.C.S.*, 1993, 115, 3407 (deriv, synth, pmr, ms)

1H-Pyrrole-2,5-dicarboxylic acid, 9CI P-946 [937-27-9]

C₆H₅NO₄ 155.11
Alkaloid from leaves of *Berberis koreana* (Berberidaceae). Needles (EtOH aq.). Sol. Et₂O, Me₂CO; insol. CHCl₃, EtOAc, C₆H₆, petrol. Mp 260° dec. (245°).

Mono-Me ester: [1199-64-0]

C₇H₇NO₄ 169.137
Alkaloid from leaves of *Berberis koreana* (Berberidaceae). Mp 241-242° (204-206°).

Di-Me ester: [1757-29-5]

C₈H₉NO₄ 183.163
Needles (H₂O). Sol. EtOH, Et₂O, C₆H₆. Mp 132° (126-127°).

Di-Et ester:

C₁₀H₁₃NO₄ 211.217
Needles. Mp 82°.

Dinitrile: 2,5-Dicyanopyrrole

[59215-66-6]
C₆H₃N₃ 117.11
Yellow solid.

1-Ph:

C₁₂H₉NO₄ 231.207
Light yellow cryst. Mp 140-144°. Various subst. 1-Ph derivs have also been prepd. and cryst. structs. determined.

Nicolaus, R.A. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 358 (*synth*)

U.S. Pat., 1959, 2 900 386; *CA*, **54**, 1557h (*synth*)


Barker, P. *et al.*, *J.O.C.*, 1978, **43**, 4849 (*derivs*)
Košťálová, D. *et al.*, *Phytochemistry*, 1992, **31**, 3669 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lin, Q. *et al.*, *J.C.S. Perkin 2*, 1998, 2109-2116 (*1-Ph, synth, cryst struct, pmr, cmr*)

Gade, L.H. *et al.*, *Chem. Eur. J.*, 2002, **8**, 4308-4318 (*dinitrile, synth, ir, pmr, cmr, ms*)

1H-Pyrrole-2,5-dione, 9CI P-947

Maleimide, 8CI
[541-59-3]



C₄H₃NO₂ 97.073
Maleimide and its *N*-substituted derivs. are used in polym. compositions. Plates. Mp 93°. Subl. readily.

► LD₅₀ (mus, orl) 80 mg/kg. Exp. reprod. and teratogenic effects. ON4800000

Monooxime: Maleimide monooxime

[74230-05-0]
C₄H₄N₂O₂ 112.088
Isol. from *Haliclona baeri*, *Haliclona cymaeformis* and *Cliona patera*. Yellow gum.

Wattanadilok, R. *et al.*, *Mar. Drugs*, 2007, **5**, 40-51 (*monooxime, isol*)

1H-Pyrrole-3-propanoic acid P-948

3-(3-Pyrrolyl)propionic acid
[134448-22-9]

C₇H₉NO₂ 139.154

Amide: 1H-Pyrrole-3-propanamide. Cystaminin A

[152509-75-6]
C₇H₁₀N₂O 138.169

Prod. by *Streptomyces* sp. KP-1241. Calpain inhibitor. Powder. Sol. H₂O, C₆H₆, MeOH, Me₂CO, CHCl₃. Mp 146-148°. λ_{max} 203 (ε 4400); 213; 260 (MeOH) (Berdy).

[150985-69-6, 107748-37-8]

Eur. Pat., 1993, 569 122; *CA*, **120**, 105131 (*Cystaminin A, isol, pmr, cmr, uv, ir*)
Hodges, L.M. *et al.*, *J.O.C.*, 1993, **58**, 4788 (*Me ester*)

Pyrrolidine, 9CI, 8CI P-949

Tetrahydropyrrole. Tetramethyleneimine. Azacyclopentane. FEMA 3523
[123-75-1]



C₄H₉N 71.122

More complex acyl derivatives not given here may be listed as pyrrolidides under the parent carboxylic acids. Manuf. by catalytic reaction of NH₃ with tetrahydrofuran or butanediol. Present in tobacco and carrot leaves (*Daucus carota*) (Apiaceae). Widely distributed in trace amts., presumably as bacterial decarboxylation prod. of proline. Widely used org. base, used in the prep. of enamines. Reagent used in the ms detn. of the position of double bonds and of methyl branching in fatty acids. Flavouring ingredient. Liq. with odour resembling Piperidine, P-454. Misc. H₂O. d₄²⁰ 0.86. Fp -63. Bp 88.5-89°. pK_a 11.31 (25°). Fumes in air. Strongly alkaline.

► Highly flammable, fl. p. 3°. Skin, eye and respiratory tract irritant. LD₅₀ (rat, orl) 300 mg/kg. UX9650000

Hempicrate:

Dark red cryst. Mp 163-164°.

Monopicrate: [1689-56-1]

Yellow needles (EtOH). Mp 112°.

N-Ac: [4030-18-6]

C₆H₁₁NO 113.159
Liq. Bp 224-225°.

N-Hexadecanoyl: N-Hexadecanoylpyrrolidine. Hexadecanoic acid pyrrolidide.

N-Palmitoylpyrrolidine. Alkaloid MQ-B₂

[70974-48-0]

C₂₀H₃₉NO 309.534

Alkaloid from *Piper amalago*, *Ipomoea aquatica* (water spinach) and *Merremia quinquefolia*.

N-(2-Hexadecenoyl) (E-): N-(2-Hexadecenoyl)pyrrolidine. 2-Hexadecenoic acid pyrrolidide

[102934-30-5]

C₂₀H₃₇NO 307.518

Alkaloid from *Piper amalago*.

N-(3,6-Hexadecadienoyl): N-(3,6-Hexadecadienoyl)pyrrolidine. 3,6-Hexadecadienoic acid pyrrolidide

[102819-18-1]

C₂₀H₃₅NO 305.503

Alkaloid from *Piper amalago*. Struct. not certain, may be the 3,7-dienoyl

isomer.

N-Octadecanoyl: N-Octadecanoylpyrrolidine. N-Stearoylpyrrolidine. Octadecanoic acid pyrrolidide. Alkaloid MQ-B₄

[33707-76-5]

C₂₂H₄₃NO 337.588

Alkaloid from *Piper amalago* and *Merremia quinquefolia*.

N-(2-Octadecenoyl) (E-): N-(2-Octadecenoyl)pyrrolidine. 2-Octadecenoic acid pyrrolidide

[102934-31-6]

C₂₂H₄₁NO 335.572

Alkaloid from *Piper amalago*.

N-(9-Octadecenoyl): N-(9-Octadecenoyl)pyrrolidine. 9-Octadecenoic acid pyrrolidide

[52380-36-6]

[4637-54-1]

C₂₂H₄₁NO 335.572

Alkaloid from *Piper amalago*.

N-(3,6-Octadecadienoyl): N-(3,6-Octadecadienoyl)pyrrolidine. 3,6-Octadecadienoic acid pyrrolidide

[102819-20-5]

C₂₂H₃₉NO 333.556

Alkaloid from *Piper amalago*. Struct. uncertain, may be the 3,7-dienoyl isomer.

N-(2,4-Eicosadienoyl): see Trichonine, T-491

N-(2,11-Eicosadienoyl) (E,E-): N-(2,11-Eicosadienoyl)pyrrolidine. 2,11-Eicosadienoic acid pyrrolidide

[102934-34-9]

C₂₄H₄₃NO 361.61

Alkaloid from *Piper amalago*.

N-(3,6-Eicosadienoyl): N-(3,6-Eicosadienoyl)pyrrolidine. 3,6-Eicosadienoic acid pyrrolidide

[102819-22-7]

C₂₄H₄₃NO 361.61

Alkaloid from *Piper amalago*. Struct. not certain, may be the 3,7-dienoyl isomer.

N-(14S-Methylhexadecanoyl): N-(14-Methylhexadecanoyl)pyrrolidine. 14-Methylhexadecanoic acid pyrrolidide.

Alkaloid MQ-A₃

[260058-83-1]

C₂₁H₄₁NO 323.561

Alkaloid from *Ipomoea aquatica* (water spinach) and *Merremia quinquefolia*. [α]_D²⁰ +5 (c, 0.4 in CHCl₃).

N-Benzoyl: [3389-54-6]

C₁₁H₁₃NO 175.23

Liq. Bp₂₀ 187°.

N-Nitroso: [930-55-2]

C₄H₈N₂O 100.12

Found in fried bacon, tobacco smoke. Oil. Bp 214°. n_D²⁰ 1.4900.

► Exp. hepatocarcinogen. Possible human carcinogen (IARC 2B). UY1575000

N-Me: see 1-Methylpyrrolidine, M-543

N-Et: [7335-06-0]

C₆H₁₃N 99.175

Liq. Bp₁₀₀ 50°.

N-Propyl: [7335-07-1]

C₇H₁₅N 113.202

Bp 130-135°.

N-Butyl: [767-10-2]

C₈H₁₇N 127.229
Liq. Bp 155-157° Bp₂₂ 55-56°. n_D²⁰
1.4400.

▶ UX9800000

N-(2-Chloroethyl): [5050-41-9]

[7250-67-1]

C₆H₁₂ClN 133.62

Mp 169.5-173° (as hydrochloride). CAS
no. refers to hydrochloride.

▶ UY0703150

N-Benzyl: [29897-82-3]

C₁₁H₁₅N 161.246

Liq. Sol. EtOH, Et₂O. Bp 237°. Ab-
sorbs CO₂ and H₂O from the air.

N-Triphenylmethyl: 1-Tritylpyrrolidine,
8CI

[18265-12-8]

C₂₃H₂₃N 313.441

Mp 124-125°.

N-(2,4-Dinitrophenyl): [14552-00-2]

Cryst. (cyclohexane/EtOAc). Mp
101.5-102.5°.

N-Amino: 1-Aminopyrrolidine. 1-Pyrroli-
dinamine, 9CI

[16596-41-1]

C₄H₁₀N₂ 86.136

Bp₁₃ 28-30°.

N-Amino; hydrochloride: [63234-71-9]

Cryst. (2-propanol). Mp 117-119°.

N-Amino, picrate: Mp 165° dec.

Aldrich Library of FT-IR Spectra, 1st edn.,
1985, **1**, 353C; 355A; 398B (*ir*)

*Aldrich Library of 13C and 1H FT NMR
Spectra, 1992, 1*, 557C; 623B (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase,
1989, **3**, 442A; 443B; 481B (*ir*)

Aldrich Library of NMR Spectra, 2, 58D
(*pmr*)

De Jong, M. *et al.*, *Rec. Trav. Chim. (J. R.
Neth. Chem. Soc.)*, 1930, **49**, 237 (*synth*)

Ochiai, E. *et al.*, *Ber.*, 1934, **67**, 1017 (*derivs*)

Rink, M. *et al.*, *Naturwissenschaften*, 1961, **48**,
51 (*N-amino*)

Wright, J.B. *et al.*, *J. Med. Pharm. Chem.*,
1962, **5**, 815 (*N-amino, synth*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 2920
(*ms*)

Wilshire, J.F.K. *et al.*, *Aust. J. Chem.*, 1966, **19**,
1935 (*deriv*)

Andersson, B.A. *et al.*, *Lipids*, 1974, **9**, 185;
443; 1975, **10**, 215; 716 (*use*)

Hawthorne, D.G. *et al.*, *Aust. J. Chem.*, 1976,
29, 315 (*cmr*)

Ahmed, M.G. *et al.*, *J.C.S. Perkin 2*, 1978, 372
(*cmr, derivs*)

*Fieser and Fieser's Reagents for Organic
Synthesis*, Wiley, 1980, **8**, 428 (*use*)

Ullmann's Encycl. Ind. Chem., 5th Ed., VCH,
Weinheim, 1985, **A2**, 13 (*rev*)

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**,
12-18 (*isol, pyrrolidides*)

Barbry, D. *et al.*, *Magn. Reson. Chem.*, 1990,
28, 560 (*cmr, N-15 nmr*)

Hunt, E.J. *et al.*, *Carcinogenesis (London)*,
1991, **12**, 571 (*N-nitroso, bibl*)

*Encyclopaedia of Reagents for Organic
Synthesis*, (ed. Paquette, L.A.), Wiley, 1995,
6, 4381-4384 (*use*)

Encyclopedia of Food and Color Additives, (ed.
Burdock, G.A.), CRC Press, 1997, 2398-
2399 (*occur*)

Akula, M.R. *et al.*, *Org. Prep. Proced. Int.*,
1999, **31**, 214-215 (*N-trityl*)

Tofern, B. *et al.*, *Phytochemistry*, 1999, **52**,
1437-1441 (*MQ alkaloids*)

Yajima, A. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 2001, **65**, 463-465 (*MQ-A₃, synth,
abs config*)

IARC Monog. (Web), (*N-nitroso*)

Luxon, S.G. *et al.*, *Hazards in the Chemical
Laboratory, 5th edn.*, Royal Society of
Chemistry, 1992, 1090

Lewis, R.J. *et al.*, *Sax's Dangerous Properties
of Industrial Materials, 8th edn.*, Van
Nostrand Reinhold, 1992, BSK250; NLP500;
PPS500

2-Pyrrolidineacetic acid, 9CI P-950

β-Homoprolin†



C₆H₁₁NO₂ 129.158

Present in flowerheads of *Arnica montana*,
Arnica chamissonis ssp. *foliosa*, *Arnica
amplexicaulis* and *Arnica sachalinensis*.
Also occurs in *Tussilago farfara* (colts-
foot).

(R)-form [61350-65-0]

Mp 188-190°. [α]_D²⁵ -44.8 (c, 1 in 3.41M
HCl).

(S)-form [56633-75-1]

Present in cured tobacco leaves. Needles
(propanol). Mp 189-191° (185°) (dec.).
[α]_D²⁵ +4 (c, 1 in H₂O). [α]_D +45 (c, 1 in
3.41M HCl).

Me ester: [53912-83-7]

C₇H₁₃NO₂ 143.185

Oil. Bp₁ 135-145°. [α]_D²⁰ +7 (c, 2.6 in
CHCl₃).

N-tert-Butyloxycarbonyl: [56502-01-3]

C₁₁H₁₉NO₄ 229.275

Cryst. (hexane). Mp 99-101°. [α]_D -39.5
(c, 1.9 in DMF) (-41.6).

N-Benzoyloxycarbonyl: [56633-73-9]

C₁₄H₁₇NO₄ 263.293

Cryst. (diisopropyl ether). Mp 74.5-
75.5°. [α]_D -35.3 (c, 1 in AcOH).

N-Benzoyloxycarbonyl, *Me ester*: [56633-
72-8]

C₁₅H₁₉NO₄ 277.319

Oil. [α]_D -35.1 (c, 1 in CHCl₃).

N-[(9H-Fluoren-9-ylmethoxy)carbonyl]:
[193693-60-6]

C₂₁H₂₁NO₄ 351.401

Cryst. (MeOH). Mp 191-192°. [α]_D²⁰ -
33.6 (c, 0.9 in DMF).

N-Me: 1-Methyl-2-pyrrolidineacetic acid.
Homohygrinic acid

[137693-28-8]

C₇H₁₃NO₂ 143.185

Plates (C₆H₆). Mp 120-122°. [α]_D -31.3
(c, 1.0 in EtOH).

N-Me, *Me ester*: [51856-76-9]

C₈H₁₅NO₂ 157.212

Constit. of *Solanum sturtianum*. Oil.
Bp₇ 80-85°.

N-Me, nitrile: 2-Cyanomethyl-1-methyl-
pyrrolidine

[67824-39-9]

C₇H₁₂N₂ 124.185

Oil. Bp_{0.5} 45-46°. [α]_D -36.6 (c, 3 in
MeOH).

N-Amidino: N-(Aminoiminomethyl)-2-
pyrrolidineacetic acid. **Cimipronidine**

C₇H₁₃N₃O₂ 171.199

Alkaloid from the roots of *Cimicifuga
racemosa*. Off-white powder. [α]_D²⁰
+36.2 (c, 0.1 in H₂O). Stereochem.
not confirmed. Zwitterionic. λ_{max}
285 (log ε 1.04); 333 (log ε 1.62)
(H₂O).

[52484-80-7]

Bremner, J.B. *et al.*, *Aust. J. Chem.*, 1973, **26**,
2559-2561 (*N-Me Me ester, isol, synth*)

Cassal, J.-M. *et al.*, *Helv. Chim. Acta*, 1976, **59**,
1917-1924 (*synth*)

Wakabayashi, T. *et al.*, *Synth. Commun.*, 1977,
7, 239-244 (*synth*)

Leete, E. *et al.*, *J.A.C.S.*, 1991, **113**, 9286-9292
(*1-Me, nitrile*)

Passreiter, C.M. *et al.*, *Phytochemistry*, 1992,
31, 4135-4137 (*isol*)

Ellmerer-Müller, E.P. *et al.*, *Helv. Chim. Acta*,
1998, **81**, 59-65 (*Fmoc deriv, synth, ir, pmr,
cmr, ms*)

Harrison, J.R. *et al.*, *J.C.S. Perkin 1*, 1999,
3623-3631 (*Me ester, synth, ir, pmr, cmr*)

Chippindale, A.M. *et al.*, *Tetrahedron*, 2003,
59, 3253-3265 (*S-form, synth*)


Calvet, S. *et al.*, *Tetrahedron*, 2003, **59**, 6333-
6339 (*S-form, Me ester*)

Fabricant, D.S. *et al.*, *J. Nat. Prod.*, 2005, **68**,
1266-1270 (*Cimipronidine*)

1-Pyrrolidinecarboxamide, P-951

9CI

N-Carbamoylpyrrolidine. 1-(Aminocarbo-
nyl)pyrrolidine
[4736-71-4]



C₅H₁₀N₂O 114.147

Metab. of the Mediterranean sponge
Aplysina (Verongia) cavernicola. Cryst.
(H₂O). Mp 148-152° (nat.) Mp 218°
(synth.).


Reppe, W. *et al.*, *Annalen*, 1955, **596**, 80 (*synth*)

D'Ambrosio, M. *et al.*, *Comp. Biochem.
Physiol., B: Comp. Biochem.*, 1986, **83**, 309
(*isol, ir, pmr, cmr, ms*)

3-Pyrrolidinecarboxylic acid P-952

β-Proline

[59378-87-9]



C₅H₉NO₂ 115.132

N-Me: **Achyranthine**

C₆H₁₁NO₂ 129.158

Alkaloid from *Achyranthes aspera*.
Vasodilator, diuretic, purgative, anti-
hypertensive agent. Monohydrate. Mp
292-293° dec. [α]_D²⁵ -6.5. Log P -2.23
(calc). Struct. proof not convincing.
Abs. config. not detd.

N-Me; hydroiodide: Mp 204-205°.

Dimethylbetaine: 3-Carboxy-1,1-di-

methylpyrrolidinium hydroxide inner salt, 9CI. β -Stachydrine
[86849-51-6]

$C_7H_{13}NO_2$ 143.185
Alkaloid from the marine red alga *Griffithsia flosculosa*. Mp 238°. Natural β -Stachydrine was characterised as the hydrochloride. The Mp given here is for semisynthetic material obt. by methylation of Achyranthine. Abs. config. not detd.

Dimethylbetaine; hydrochloride: Mp 236-238° dec. $[\alpha]_D^{23}$ -5.5 (c. 0.27 in MeOH). Deriv. of β -Stachydrine.

Dimethylbetaine; hydroiodide: Mp 267°. Derived from Achyranthine.

(R)-form [72580-54-2]

Synthetic. Cryst. (EtOH). Mp 188-190°. $[\alpha]_D^{25}$ -18.9 (c, 2.0 in H_2O).

(S)-form [72580-53-1]

Mp 186-188.5°. $[\alpha]_D^{25}$ +19.1 (c, 2 in H_2O). Struct. incorr. drawn by Fleš \acute{e} t *et al* (1964).

Me ester:

$C_6H_{11}NO_2$ 129.158
Volatile oil. $[\alpha]_D^{23}$ +7.2 (c, 1.0 in $CHCl_3$).

(\pm)-form [68464-02-8]

Plates. Mp 186°.

Et ester: [72925-15-6]

$C_7H_{13}NO_2$ 143.185
Bp₄ 70-74°.

Basu, N.K. *et al.*, *J. Proc. Inst. Chem. (India)*, 1957, **29**, 55; 73 (*Achyranthine*)

Miyamoto, M. *et al.*, *Yakugaku Zasshi*, 1957, **77**, 568; *CA*, **51**, 16422h (*synth*)

Fleš \acute{e} , D. *et al.*, *Croat. Chem. Acta*, 1964, **36**, 27-32; *CA*, **61**, 10644b (*S-form, synth, abs config*)

Yuki, H. *et al.*, *J. Polym. Sci., Polym. Chem. Ed.*, 1979, **17**, 3867-3878 (*R-form, synth, resoln, pmr*)

Blunden, G. *et al.*, *Phytochemistry*, 1983, **22**, 293 (*β -Stachydrine*)


Mazzini, C. *et al.*, *J.O.C.*, 1997, **62**, 5215-5218 (*R-form, S-form, synth, pmr, cmr*)

Thomas, C. *et al.*, *Synthesis*, 1998, 1491-1496 (*R-form, S-form, synth, ir, pmr*)

Blanchet, J. *et al.*, *Tet. Lett.*, 2007, **48**, 5727-5730 (*synth*)

2,4-Pyrrolidinedicarboxylic acid P-953

4-Carboxypyrrolidine



(2R,4R)-form

$C_6H_9NO_4$ 159.141

(2R,4R)-form

D-cis-form

[130830-78-3]

Prisms. Mp 224-225°. $[\alpha]_D^{21}$ +41.5 (c, 1.01 in H_2O) (+37).

N-Benzoyl: [188345-79-1]

$C_{13}H_{13}NO_5$ 263.249
Mp 164-165°. $[\alpha]_D^{20}$ +82 (c, 1.04 in

EtOH).

(2R,4S)-form

D-trans-form

[130830-77-2]

Prisms. Mp 219-220°. $[\alpha]_D^{25}$ +51 (c, 1.04 in H_2O).

(2S,4R)-form

L-trans-form

[64769-66-0]

Constit. of the Mediterranean red algae *Chondria coerulescens*, *Chondria dasiphylla* and *Ceramium rubrum*; found in the seeds of *Azelia bella* (Fabaceae). Cryst. (MeOH aq.). Mp 223-225°. $[\alpha]_D$ -54 (c, 1.04 in H_2O).

4-Me ester:

$C_7H_{11}NO_4$ 173.168

No phys. props. reported. $[\alpha]_D$ -11 (c, 1.5 in $CHCl_3$).

N-Benzyl: [627100-87-2]

$C_{13}H_{15}NO_4$ 249.266

Mp 60-62° (as hydrochloride). $[\alpha]_D^{20}$ +29.4 (c, 0.78 in MeOH) (hydrochloride). CAS no. refers to the hydrochloride.

2-Et ester, 4-nitrile, N-benzyl: [627100-83-8]

$C_{15}H_{18}N_2O_2$ 258.319

Oil. $[\alpha]_D^{20}$ +64.8 (c, 1 in $CHCl_3$).

(2S,4S)-form

L-cis-form

[64927-38-4]

Prisms (EtOH aq./Me₂CO). Mp 225-256°. $[\alpha]_D^{25}$ -40 (c, 1.02 in H_2O).

(2RS,4RS)-form

(\pm)-*cis-form*

[136033-11-9]

Prisms (EtOH). Mp 226-230°.

Impellizzeri, G. *et al.*, *Phytochemistry*, 1977, **16**, 1601-1602 (*isol, pmr, ms, struct*)

Dupont, L. *et al.*, *Acta Cryst. B*, 1978, **34**, 850-854 (*cryst struct*)

Welter, A. *et al.*, *Phytochemistry*, 1978, **17**, 131-134 (*isol, ir, pmr, cmr, ms, cd, struct*)

Bridges, R.J. *et al.*, *J. Med. Chem.*, 1991, **34**, 717-725 (*synth, pmr, cmr, ir*)

Arakawa, Y. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 255-259 (*2R,3R-form, 2S,3S-form, 2RS,3RS-form, synth, pmr, cmr*)

Wang, Q. *et al.*, *Tetrahedron*, 1998, **54**, 15759-15780 (*2S,4R-form, synth, ir, nmr, 4-Me ester*)


Heindl, C. *et al.*, *Tetrahedron: Asymmetry*, 2003, **14**, 3141-3152 (*2S,4R-form, benzyl derivs*)

2,4-Pyrrolidinedione, 9CI P-954

1,5-Dihydro-4-hydroxy-2H-pyrrol-2-one.

Tetramic acid

[37772-90-0]



$C_4H_5NO_2$ 99.089

Pale yellow solid. Mp 360°.

N-Ac: [182352-40-5]

$C_6H_7NO_3$ 141.126

Cryst. solid (MeOH). Mp 165-168°.

N-Benzoyl: [68661-20-1]

$C_{11}H_9NO_3$ 203.197

Solid. Mp 180° dec.

N-Ethoxycarbonyl:

$C_7H_9NO_4$ 171.152

Solid. Mp 105-108° (96-102°).

N-tert-Butyloxycarbonyl:

$C_9H_{13}NO_4$ 199.206

Cryst. solid (EtOAc). Mp 130-132° dec.

N-Benzylloxycarbonyl:

$C_{12}H_{11}NO_4$ 233.223

Cryst. solid. Mp 151-153° dec.

Enol-form

Me ether, 1,5-Dihydro-4-methoxy-2H-pyrrol-2-one, 9CI

[69778-83-2]

$C_5H_7NO_2$ 113.116

Constit. of the marine cyanophyte *Lyngbya majuscula*. Hydrol. prod. of Malyngamide A. Needles (C_6H_6). Mp 128-129°.

Me ether, N-Ac: 1-Acetyl-1,5-dihydro-4-methoxy-2H-pyrrol-2-one

$C_7H_9NO_3$ 155.153

Minor constit. of *Lyngbya majuscula*. Mp 89-90°.

Benzyl ether, 4-Benzylloxy-1,5-dihydro-2H-pyrrol-2-one

[113896-95-0]

$C_{11}H_{11}NO_2$ 189.213

Mp 150-153°.

N^d, 4-Di-Me: 1,5-Dihydro-4-methoxy-1-methyl-2H-pyrrol-2-one, 9CI

[90968-33-5]

$C_6H_9NO_2$ 127.143

Prisms (C_6H_6 /hexane). Mp 86-87°.

Cram, D.J. *et al.*, *J.A.C.S.*, 1963, **85**, 1430 (*deriv, synth*)

Mulholland, T.P.C. *et al.*, *J.C.S. Perkin 1*, 1972, 2121 (*synth*)

Lowe, G. *et al.*, *J.C.S. Perkin 1*, 1973, 2907 (*synth, nmr*)

Cardellina, J.H. *et al.*, *J.A.C.S.*, 1979, **101**, 240 (*occur, derivs*)

Kochhar, K.S. *et al.*, *Tet. Lett.*, 1984, **25**, 1871 (*synth*)

Gill, G.B. *et al.*, *J.C.S. Perkin 1*, 1993, 2567 (*deriv, synth, ir, pmr, cmr*)

James, G.D. *et al.*, *J.C.S. Perkin 1*, 1993, 2581 (*deriv, synth, ir, pmr, cmr*)

Hamilakis, S. *et al.*, *J. Het. Chem.*, 1996, **33**, 825 (*acyl derivs*)


Parsons, A.F. *et al.*, *Tet. Lett.*, 1996, **37**, 1667 (*benzyl ether, synth*)

Lu, Z. *et al.*, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 2505-2507 (*synth*)

2-Pyrrolidinol

2-Hydroxypyrrolidine

P-955



C_4H_9NO 87.121

(\pm)-form

N-Formyl: 2-Hydroxy-1-pyrrolidinecarboxaldehyde, 9CI

[69352-24-5]

C₅H₉NO₂ 115.132

Liq. Bp_{1.5} 122-125°.

Me ether, N-formyl: [61020-06-2]

C₆H₁₁NO₂ 129.158

Liq. Bp₄₈ 135-137° Bp_{1.5} 77-80°. n_D²⁵ 1.4700.

Me ether, N-E-cinnamoyl: **Aglamide D**

[922149-10-8]

C₁₄H₁₇NO₂ 231.294

Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. λ_{max} 206 (log ε 4.2); 218 (log ε 4.22); 280 (log ε 4.35) (MeOH) (206).

(ξ)-form

Me ether, N-carbamoyl: N-Carbamoyl-2-methoxy-pyrrolidine. 2-Methoxy-1-pyrrolidinecarboxamide, 9CI

[83459-48-7]

C₆H₁₂N₂O₂ 144.173

Alkaloid from the stem bark of *Hexalobus crispiflorus*, twigs of *Oxymitra velutina* (preferred genus name *Friesodielsia*) and roots and stem bark of *Piptostigma fugax* (Annonaceae). Needles (petrol/Me₂CO). Mp 96-98°. [α]_D 0 (c, 0.66 in MeOH).

Me ether, N-cinnamoyl: N-Cinnamoyl-2-methoxy-pyrrolidine. **Piperlotine I**

C₁₄H₁₇NO₂ 231.294

Alkaloid from the leaves of *Piper lolot*. Syrup. [α]_D +23.2 (c, 0.08 in MeOH). λ_{max} 210 ; 216 ; 222 ; 274 (MeOH).

Nyberg, K. et al., *Acta Chem. Scand., Ser. B*, 1976, **30**, 640-642 (*synth*)

Mitzlaff, M. et al., *Annalen*, 1978, 1713-1733 (*Me ether N-formyl*)

Iwasaki, T. et al., *J.O.C.*, 1979, **44**, 1552-1554 (*synth*)

Achenbach, H. et al., *Annalen*, 1982, 1623-1633 (*Me ether N-carbamoyl*)


Achenbach, H. et al., *Phytochemistry*, 1991, **30**, 1265-1267; 1995, **38**, 1037-1048 (*Me ether N-carbamoyl*)

Kim, S. et al., *J. Nat. Prod.*, 2006, **69**, 1769-1775 (*Aglamide D*)

Li, C.-Y. et al., *J. Agric. Food Chem.*, 2007, **55**, 9436-9442 (*Piperlotine I*)

2-Pyrrolidinone, 9CI, 8CI P-956

2-Oxopyrrolidine. Butyrolactam. 2-Pyrrolidone. Piperidinic lactam. 4-Aminobutanoic acid lactam [616-45-5]



C₄H₇NO 85.105

Manuf. from γ-butyrolactone and NH₃. Constit. of *Artemisia asiatica nakai*, *Solanum vestissimum* and red ginseng. Isol. from the rump glands of the antelope *Antilocapra americana* as a possible alerting pheromone. Reagent used in the ms detn. of the branching points in long-chain primary alcohols, low acute tox. Plasticiser/emulsifier. Sol-

vent. Anionic polym. gives nylon-4. Cryst. (petrol). V. sol. H₂O, most org. solvs.; spar. sol. petrol. Mp 24.6°. Bp 245° Bp₁₂ 133°. pK_a -0.94 (25°). Spar. steam-volatile. Forms a monohydrate in moist air, Mp 35° (30°). Ceiling temp. for polym., T_c 76°.

► Eye, skin and mucous membrane irritant. UY5715000

N-E-Cinnamoyl: 1-(1-Oxo-3-phenyl-2-propenyl)-2-pyrrolidinone. N-Cinnamoyl-2-pyrrolidinone. **Piperlotine F**

[141236-49-9]

C₁₃H₁₃NO₂ 215.251

Alkaloid from the leaves of *Piper lolot*. Needles. Mp 101-102°. λ_{max} 211 ; 273 (MeOH).

N-(4-Methoxy-E-cinnamoyl): N-(4-Methoxycinnamoyl)-2-pyrrolidinone.

Piperlotine G

[141236-52-4]

C₁₄H₁₅NO₃ 245.277

Alkaloid from the leaves of *Piper lolot*. Needles. Mp 140-142°. λ_{max} 222 ; 284 (MeOH).

N-(4-Methoxy-Z-cinnamoyl): **Piperlotine H**

[958296-15-6]

C₁₄H₁₅NO₃ 245.277

Alkaloid from the leaves of *Piper lolot*. Syrup. λ_{max} 224 ; 274 (MeOH).

N-(2,3,6-Tribromo-4,5-dihydroxybenzyl): N-(2,3,6-Tribromo-4,5-dihydroxybenzyl)-2-pyrrolidinone

[946585-49-5]

C₁₁H₁₀Br₃NO₃ 443.917

Isol. from red alga *Symphyocladia latiuscula*. Amorph. yellow solid. Mp 166-168°. λ_{max} 216 (log ε 3.86); 259 (log ε 2.9); 295 (log ε 2.7) (MeOH).

Suarez, M. et al., *Flavour Fragrance J.*, 1993, **8**, 215-220 (*isol*)

Wood, W.F. et al., *Biochem. Syst. Ecol.*, 2002, **30**, 361-363 (*isol*)

Li, C.-Y. et al., *J. Agric. Food Chem.*, 2007, **55**, 9436-9442 (*Piperlotines F,G,H*)

Duan, X.-J. et al., *J. Nat. Prod.*, 2007, **70**, 1210-1213


(*Tribromodihydroxybenzylpyrrolidinone*)

3-(2-Pyrrolidinyl)pyridine, P-957

9CI

2-(3-Pyridyl)pyrrolidine. **Nornicotine**

[13450-58-3]



C₉H₁₂N₂ 148.207

(**R**)-form [7076-23-5]

Alkaloid from *Duboisia hopwoodii* (Solanaceae). d₄¹⁰ 1.08. Bp_{3.6} 117°. [α]_D²⁰ +86.3 (60% o.p.). n_D¹⁸ 1.5490.

► RC8050000

Dipicrate: Mp 191-192°.

N-*Me*: see Nicotine, N-190

(**S**)-form [494-97-3]

Constit. of tobacco, *Nicotiana tabacum*

(Solanaceae). Principal alkaloid of some tobacco strains. Insecticide. Now superseded. d_{19.5}^{19.5} 1.07. Bp₁₁ 130-131° Bp₁ 120°. n_D^{18.5} 1.5378.

► QS6405000

N-Formyl: **N-Formylnornicotine**. 2-(3-Pyridinyl)-1-pyrrolidinecarboxaldehyde, 9CI

[38840-03-8]

C₁₀H₁₂N₂O 176.218

Alkaloid from *Nicotiana tabacum* and *Duboisia hopwoodii* (Solanaceae). No opt. rotn. reported for the *D. hopwoodii* isolate, could be the enantiomer (cf. Nornicotine).

N-*Ac*: **N-Acetylnornicotine**. 1-Acetyl-2-(3-pyridinyl)pyrrolidine, 9CI

[5979-94-2]

C₁₁H₁₄N₂O 190.244

Alkaloid from *Nicotiana tabacum* and *Duboisia hopwoodii* (Solanaceae). Bp₁₂ 212-214°. [α]_D²⁰ -3.24 (C₆H₆). No opt. rotn. recorded for the *D. hopwoodii* isolate (Solanaceae).

► UX9682000

N-Hexanoyl: **N-Hexanoylnornicotine**. 1-(1-Oxohexyl)-2-(3-pyridinyl)pyrrolidine, 9CI

[38854-09-0]

C₁₅H₂₂N₂O 246.352

Minor alkaloid from cured tobacco (*Nicotiana tabacum*) (Solanaceae).

N-Octanoyl: **N-Octanoylnornicotine**. 1-(1-Oxooctyl)-2-(3-pyridinyl)pyrrolidine, 9CI

[38854-10-3]

C₁₇H₂₆N₂O 274.405

Minor alkaloid from cured tobacco (*Nicotiana tabacum*) (Solanaceae).

N-*Me*: see Nicotine, N-190

N-*Et*: 3-(1-Ethyl-2-pyrrolidinyl)pyridine, 9CI. **N-Ethylnornicotine**

[5979-92-0]

C₁₁H₁₆N₂ 176.261

Alkaloid from leaves of *Nicotiana tabacum* (Solanaceae). Oil.

► UT3770000

N-Isopropyl: 3-[1-(1-Methylethyl)-2-pyrrolidinyl]pyridine, 9CI. **N-Isopropyl-nornicotine**

[72461-69-9]

C₁₂H₁₈N₂ 190.288

Alkaloid from *Nicotiana tabacum* (Solanaceae).

N-(3-Hydroxy-12-methyltridecanoyl): 1-(3-Hydroxy-12-methyl-1-oxotridecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. **N-(3-Hydroxy-12-methyltridecanoyl)nornicotine**

[115849-84-8]

C₂₃H₃₈N₂O₂ 374.565

Alkaloid from trichomes of *Nicotiana stocktonii*.

N-(β-D-Fructopyranos-1-yl): 1-Deoxy-1-[2-(3-pyridinyl)-1-pyrrolidinyl]-β-D-fructopyranose

C₁₅H₂₂N₂O₅ 310.349

Amadori compd. isol. from flue-cured Delhi tobacco. Needles. Mp 67-70°. [α]_D²⁵ -94 (c, 1 in H₂O).

N'-Nitroso: [16543-55-8]
[84237-38-7, 80508-23-2, 61601-78-3]
C₉H₁₂N₃O₂[⊕] 194.213
Carcinogen in tobacco and tobacco
smoke. Bp_{0.2} 150-160°.

▶ Possible human carcinogen (IARC 2B).

(±)-form [5746-86-1]

Bp_{0.5} 94°.

▶ RC8400000

N-Et: [86500-39-2]

Bp₁₃ 128-130°.

(ξ)-form

N-(3-Hydroxy-10-methylundecanoyl): 1-(3-Hydroxy-10-methyl-1-oxoundecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-10-methylundecanoyl)nornicotine

[115849-79-1]

C₂₁H₃₄N₂O₂ 346.512

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp. Plant growth inhibitor.

N-(3-Hydroxydodecanoyl): 1-(3-Hydroxy-1-oxododecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxydodecanoyl)nornicotine

[115849-80-4]

C₂₁H₃₄N₂O₂ 346.512

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp. Plant growth inhibitor.

N-(3-Hydroxy-10-methyldodecanoyl): 1-(3-Hydroxy-10-methyl-1-oxododecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-10-methyldodecanoyl)nornicotine

[115849-82-6]

C₂₂H₃₆N₂O₂ 360.539

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp.

N-(3-Hydroxy-11-methyldodecanoyl): 1-(3-Hydroxy-11-methyl-1-oxododecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-11-methyldodecanoyl)nornicotine

[115849-81-5]

C₂₂H₃₆N₂O₂ 360.539

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp.

N-(3-Hydroxy-12-methyltetradecanoyl): 1-(3-Hydroxy-12-methyl-1-oxotetradecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-12-methyltetradecanoyl)nornicotine

C₂₄H₄₀N₂O₂ 388.592

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp.

N-(3-Hydroxy-13-methyltetradecanoyl): 1-(3-Hydroxy-13-methyl-1-oxotetradecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-13-methyltetradecanoyl)nornicotine

[116353-94-7]

C₂₄H₄₀N₂O₂ 388.592

Alkaloid from *Nicotiana repanda* and other *Nicotiana* spp.

N-(3-Hydroxy-14-methylpentadecanoyl): 1-(3-Hydroxy-14-methyl-1-oxopentadecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxy-14-methylpentadecanoyl)nornicotine

[120042-33-3]

C₂₅H₄₂N₂O₂ 402.619

Alkaloid from *Nicotiana repanda*.

N-(3-Hydroxyhexadecanoyl): 1-(3-Hydroxy-1-oxohexadecyl)-2-(3-pyridinyl)pyrrolidine, 9CI. N-(3-Hydroxyhexadecanoyl)nornicotine

[120042-32-2]

C₂₅H₄₂N₂O₂ 402.619

Alkaloid from *Nicotiana repanda*.

[86900-40-5, 91429-66-2, 123348-72-1, 123348-68-5]

Späth, E. et al., *Ber.*, 1935, **68**, 1388; 1667 (isol)

Swain, M.L. et al., *J.A.C.S.*, 1949, **71**, 1342 (uv)

Erdtman, H. et al., *Acta Chem. Scand.*, 1963, **17**, 1717 (synth)

Duffield, A.M. et al., *J.A.C.S.*, 1965, **87**, 2926 (ms)

Leete, E. et al., *J.O.C.*, 1972, **37**, 4465 (synth)

Bolt, A.J.N. et al., *Phytochemistry*, 1972, **11**, 2341 (N-Hexanoynornicotine, N-Octanoynornicotine)

Warfield, A.H. et al., *Phytochemistry*, 1972, **11**, 3371 (N-Acetylnornicotine, N-Formylnornicotine, synth)

Seeman, J.I. et al., *Synthesis*, 1977, 498 (synth, nm, ms)

Hecht, S.S. et al., *J.O.C.*, 1978, **43**, 72-76 (N'-nitroso, synth, bibl)

Nakane, M. et al., *J.O.C.*, 1978, **43**, 3922 (synth)

Heck, R.F. et al., *Acc. Chem. Res.*, 1979, **12**, 146 (synth)

Miyano, M. et al., *Agric. Biol. Chem.*, 1979, **43**, 2205 (N-Isopropynornicotine)

Siddiqui, I.R. et al., *Carbohydr. Res.*, 1981, **98**, 57 (fructosyl)

Leete, E. et al., *Phytochemistry*, 1981, **20**, 1037 (N-Isopropynornicotine)

Jacob, P. et al., *J.O.C.*, 1982, **47**, 4165 (synth, resoln)

Castonguay, A. et al., *Methods Enzymol.*, 1982, **84**, 641-650 (N'-nitroso, rev)

Luanratana, O. et al., *Phytochemistry*, 1982, **21**, 449 (N-Acetylnornicotine, N-Formylnornicotine)

Langhals, E. et al., *Annalen*, 1983, 330 (synth)

Seeman, J.J. et al., *J.O.C.*, 1985, **50**, 5419 (synth)

Zador, E. et al., *Plant Physiol.*, 1986, **82**, 479 (N-(3-Hydroxy-12-methyltridecanoyl), isol, biosynth)

Severson, R.F. et al., *ACS Symp. Ser.*, 1988, **380**, 335-362 (isol, N-acyl derivs)

Mahboobi, S. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 175 (synth)

Kuno, H. et al., *Agric. Biol. Chem.*, 1989, **53**, 535 (N-(3-Hydroxy-12-methyltridecanoyl), synth)

Braumann, T. et al., *Phytochemistry*, 1990, **29**, 3693 (N-Ethynornicotine)

Pesticide Manual, 9th edn., 1991, No. 9050

Aislaitner, G. et al., *Med. Sci. Res.*, 1992, **20**, 897 (metab)

Deo, N.M. et al., *Tet. Lett.*, 1996, **37**, 1137 (synth)

Isomura, S. et al., *J.O.C.*, 2001, **66**, 4115-4121 (R-form, S-form, synth, pmr, cmr)

Felplin, F.-X. et al., *J.O.C.*, 2001, **66**, 6305-6312 (synth)


Baxendale, I.R. et al., *J.C.S. Perkin 1*, 2002, 143-154 (synth)

IARC Monog. (Web),

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NNR500; NNS500

11-(2-Pyrrolidinyl)-2-undecylamine P-958

α-Methyl-2-pyrrolidinedecanamine, 2-(10-Aminoundecyl)pyrrolidine



C₁₅H₃₂N₂ 240.431

N¹-(2-Hydroxyethyl): 2-(10-Aminoundecyl)-1-pyrrolidineethanol, 9CI

[192213-03-9]


C₁₇H₃₆N₂O 284.484

Alkaloid from the ladybird *Epilachna borealis*.

Radford, P. et al., *J. Nat. Prod.*, 1997, **60**, 755-759 (isol, ms)

Asteronotus cespitosus Pyrrolidone P-959

[489469-96-7]



C₁₇H₂₂Cl₆N₂O₄ 531.088


Related to Dysideapyrrolidone, D-969.

Alkaloid from the dorid nudibranch *Asteronotus cespitosus*. [α]_D +4.8 (c, 0.0004 in CHCl₃).

Fahey, S.J. et al., *J. Chem. Ecol.*, 2002, **28**, 1773-1785 (isol, pmr, cmr)

1-Pyrrolizidinecarboxylic acid P-960

Hexahydro-1H-pyrrolizine-1-carboxylic acid, 9CI



(1R,7aR)-form

C₈H₁₃NO₂ 155.196

All 4 stereoisomers occur as necine base components of pyrrolizidine alkaloids.

(1R,7aR)-form [22269-14-3]

Cryst. by subl. Mp 244-245°. [α]_D²⁵ +82 (c, 0.7 in H₂O).

Picrate:

Yellow prisms (EtOH). Mp 218-219°.

Methodide:

Cryst. (Me₂CO). Mp 141-142°.

Me ester: *Chysin A*. *Chysine*. 1-Methoxycarbonylpyrrolizidine [22269-11-0]

C₉H₁₅NO₂ 169.223

Alkaloid from *Chysis bractescens*

(Orchidaceae). Oil. [α]_D²⁵ +64 (c, 1.1 in CHCl₃).

Et ester: Chysis BC₁₀H₁₇NO₂ 183.25Alkaloid from *Chysis bractescens* (Orchidaceae). No phys. props. reported.**(1R,7aS)-form***Trachelanthamidinic acid*

[526-66-9]

Mp 215-216°. [α]_D -43.4 (H₂O). [α]_D -32.5 (EtOH).*Picrate*: Mp 178-179°.**(1S,7aR)-form***Laburnoic acid*

[488-07-3]

Cryst. + 1H₂O. Mp 215-216°. [α]_D +44.2 (H₂O).*Picrate*: Mp 175-176°.**(1S,7aS)-form***Isotronecanolic acid*

[526-65-8]

Mp 228-229° dec. [α]_D -71.4 (EtOH).*Picrate*: Mp 219-220°.*Me ester*: [63527-06-0]C₉H₁₅NO₂ 169.223Oil. [α]_D²⁰ -65.1 (c, 2.16 in CHCl₃).**(1RS,7aSR)-form**

(±)-cis-form

Et ester, picrate:


Cryst. (EtOH). Mp 149-151°.

(1RS,7aRS)-form

(±)-trans-form

Et ester: [34951-60-5]Bp_{0.1} 113-115°.*Et ester, picrate*:

Cryst. (EtOH). Mp 120-121°.

Adams, R. *et al.*, *J.A.C.S.*, 1942, **64**, 2597 (synth)Galinsky, F. *et al.*, *Monatsh. Chem.*, 1954, **85**, 913 (synth)Lüning, B. *et al.*, *Tet. Lett.*, 1965, 921 (Chysins)Lüning, B. *et al.*, *Acta Chem. Scand.*, 1968, **22**, 2324 (isol, synth)Schneckenburger, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1977, **310**, 177 (synth)Borch, R.F. *et al.*, *J.O.C.*, 1977, **42**, 1225 (synth)Pinnick, H.W. *et al.*, *J.O.C.*, 1978, **43**, 4662 (synth)Terao, Y. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 3167 (synth)David, O. *et al.*, *J.O.C.*, 1999, **64**, 3122-3131 (Me ester, synth, pmr, cmr, ir)**Pyrrolnitrin, INN, USAN P-961**3-Chloro-4-(3-chloro-2-nitrophenyl)-1H-pyrrole, 9CI. *Icar. Micutrin. Pyro-Ace. Rugosin H. A 10338. FR 005759. Lilly 52230. NSC 107654. Antibiotic A 10338 [1018-71-9]*C₁₀H₆Cl₂N₂O₂ 257.075Pyrrole antibiotic. Produced by *Pseudomonas pyrrocinia*, *Pseudomonas aureofaciens*, *Pseudomonas fluorescens*, other *Pseudomonas* spp. and *Myxococcus fulvus*. Antifungal agent. Pale-yellow cryst. (C₆H₆). Mp 125°. Log P 3.86 (calc). Dec. on exp. to air and light. λ_{max} 252 (ε 7500) (EtOH) (Derep).▶ LD₅₀ (mus, orl) 1000 mg/kg. UX9450000**2-Chloro-2,3-Dichloro-4-(3-chloro-2-nitrophenyl)-1H-pyrrole, 8CI. Chloropyrrolnitrin**

[21390-86-3]

C₁₀H₅Cl₃N₂O₂ 291.52Isol. from *Pseudomonas aureofaciens* and *Pseudomonas cepacia*. Active against *Neurospora* sp. Yellow cryst. (C₆H₆/hexane). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 140-141°. λ_{max} 212 (ε 29000); 240; 280 (MeOH) (Berdy). λ_{max} 252 (EtOH) (Berdy).**4'-Fluoro-3-Chloro-4-(3-chloro-4-fluoro-2-nitrophenyl)-1H-pyrrole. 4-Fluoropyrrolnitrin**

[17307-19-6]

C₁₀H₅Cl₂FN₂O₂ 275.065Prod. by *Pseudomonas aureofaciens* in the presence of 6-Fluorotryptophan. Active against *Neurospora* sp. and fungi. Yellow plates (CHCl₃). Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{max} 248 (ε 6550) (EtOH) (Berdy).**6'-Hydroxy-4-Chloro-2-(4-chloro-1H-pyrrol-3-yl)-3-nitrophenol, 9CI. 3-Chloro-4-(3-chloro-6-hydroxy-2-nitrophenyl)-1H-pyrrole. Oxypyrrrolnitrin**

[15345-51-4]

C₁₀H₆Cl₂N₂O₃ 273.074Produced by *Pseudomonas pyrrolnitrinica*. Pale-yellow cryst. (Et₂O/hexane). Sol. MeOH, EtOAc; fairly sol. CHCl₃, Et₂O, C₆H₆; poorly sol. H₂O, hexane. Mp 215-216° dec. λ_{max} 290 (ε 3470) (EtOH) (Derep).**3-Dechloro, bromo analogue: 3-(3-Bromo-2-nitrophenyl)-1H-pyrrole, 9CI**

[88899-31-4]

C₁₀H₇BrN₂O₂ 267.082Prod. by *Pseudomonas aureofaciens*. Active against *Neurospora crassa*.**3'-Dechloro-3-Chloro-4-(2-nitrophenyl)-pyrrole**


[7123-69-5]

C₁₀H₇ClN₂O₂ 222.63Metab. of *Pseudomonas pyrrolnitrinica*. Light-yellow needles (C₆H₆/petrol). Mp 111°.**3'-Dechloro, 2-chloro-2,3-Dichloro-4-(2-nitrophenyl)-1H-pyrrole, 9CI. Isopyrrolnitrin**

[6991-29-3]


C₁₀H₆Cl₂N₂O₂ 257.075Prod. by *Pseudomonas* sp. Yellowish cryst. Mp 105-108°.Arima, K. *et al.*, *Agric. Biol. Chem.*, 1964, **28**, 575-576 (isol, ir, nmr)Imanaka, H. *et al.*, *J. Antibiot., Ser. A*, 1965, **18**, 207-210 (struct, ir, uv, nmr)Hashimoto, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1966, **39**, 410; *Chem. Pharm. Bull.*, 1966, **14**, 1314-1316; 1968, **16**, 1144 (Pyrrolnitrin,*Isopyrrolnitrin, Oxypyrrrolnitrin, 3'-dechloro deriv*)Hamill, R.L. *et al.*, *Antimicrob. Agents Chemother.*, 1967, 388-396 (Chloropyrrolnitrin, isol, props)Gorman, M. *et al.*, *Biochem. Biophys. Res. Commun.*, 1968, **31**, 294 (4-Fluoropyrrolnitrin)Jones, N. *et al.*, *J. Antibiot.*, 1968, **21**, 451 (4-Fluoropyrrolnitrin, cryst struct)Japan. Pat., 1968, 68 16 135; CA, **70**, 57621 (Chloropyrrolnitrin, synth)Morimoto, Y. *et al.*, *Tet. Lett.*, 1968, 209-211 (cryst struct)Kumada, S. *et al.*, CA, 1969, **70**, 36371 (pharmacol)Umio, S. *et al.*, *Chem. Pharm. Bull.*, 1969, **17**, 559-566; 588-595 (Pyrrolnitrin, 3'-chloro deriv, synth, ir, uv, nmr)Hamill, R.L. *et al.*, *Appl. Microbiol.*, 1970, **19**, 721 (4-Fluoropyrrolnitrin)Japan. Pat., 1971, 70 29 432; CA, **74**, 52144 (Oxypyrrrolnitrin, isol)Gosteli, J. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 451-460 (synth, ir, uv, nmr)Martin, L.L. *et al.*, *J.A.C.S.*, 1972, **94**, 8942-8944 (cmr)Salcher, O. *et al.*, *Tet. Lett.*, 1978, 3097-3100 (isol)Howell, C.R. *et al.*, *Phytopathology*, 1979, **69**, 480-482 (isol)Chang, C.J. *et al.*, *J. Antibiot.*, 1981, **34**, 555-566 (biosynth)Gerth, K. *et al.*, *J. Antibiot.*, 1982, **35**, 1101-1103 (isol, bibl)Van Pée, K.-H. *et al.*, *J. Antibiot.*, 1983, **36**, 1735 (3-(3-Bromo-2-nitrophenyl)pyrrole)Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 387Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CFC000**Pyrrolo[3,4-d]imidazole-4,6(1H,5H)-dione**

P-962

C₅H₃N₃O₂ 137.098N⁵-(3-Methyl-2-butenyl): **Bredilonine**C₁₀H₁₁N₃O₂ 205.216Constit. of the leaves of *Bridelia balansae*. Powder. Mp > 300°. λ_{max} 209 (log ε 4.13); 274 (log ε 4.04) (MeOH).Tsai, Y.-H. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 2452-2457 (isol, pmr, cmr, ms)**Pyrrolomycin B**

P-963

2,4-Dichloro-6-[4,5-dichloro-3-nitro-1H-pyrrol-2-yl)methyl]phenol, 9CI. 2,3-Dichloro-5-(3,5-dichloro-2-hydroxybenzyl)-4-nitropyrrrole. SF 2080B. Antibiotic SF 2080B [79763-00-1]




C₁₁H₆Cl₄N₂O₃ 355.991
 prod. by *Streptomyces* strain SF 2080 (*Actinosporangium vitaminophilum*). Active against gram-positive and -negative bacteria and some fungi. Immunopotentiator by acting on the membrane of lymphocytes. Yellow cryst. Sol. MeOH, Me₂CO, dioxan; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 222-225°. [α]_D²⁵ 0 (c, 1 in MeOH). λ_{max} 277 (ε 7080); 332 (ε 3720) (MeOH/HCl) (Derep). λ_{max} 245 (ε 12000); 312 (ε 12100) (MeOH/NaOH) (Derep). λ_{max} 277 (ε 6440); 330 (ε 3720) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 100 mg/kg. SK9200000
 O-Ac:

Needles (C₆H₆). Mp 180-181°.
 Ezaki, N. *et al.*, *J. Antibiot.*, 1981, **34**, 1363-1365; 1366-1368; 1483-1489 (*isol, uv, ir, pmr, cmr, cryst struct*)
 Umezawa, K. *et al.*, *J. Antibiot.*, 1984, **37**, 1253-1256 (*props*)
 Ratnayake, A.S. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1923-1926 (*biosynth*)

Pyrrolomycin C P-964

(3,5-Dichloro-2-hydroxyphenyl)(4,5-dichloro-1H-pyrrol-2-yl)methanone, 9CI. 2,3-Dichloro-5-(3,5-dichloro-2-hydroxybenzoyl)pyrrole. SF 2080C. Antibiotic SF 2080C [81910-06-7]



C₁₁H₅Cl₄NO₂ 324.977
 Pyrrole antibiotic. Prod. by *Actinosporangium vitaminophilum* SF-2080 and *Streptomyces fumanus* LL-F42248. Active against gram-positive bacteria. Yellow needles (C₆H₆). Sol. bases, EtOAc, Me₂CO; fairly sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 220-221°. λ_{max} 220 (sh) (ε 21300); 260 (ε 8180); 327 (ε 14900); 340 (sh) (ε 14700) (MeOH) (Derep).

► LD₅₀ (mus, ipr) 50 mg/kg. PC4948000

Me ether: Pyrrolomycin I
 C₁₂H₇Cl₄NO₂ 339.004
 Prod. by *Streptomyces fumanus* LL-F42248.

4-Chloro-2,3,4-Trichloro-5-(3,5-dichloro-2-hydroxybenzoyl)pyrrole. **Pyrrolomycin D**. SF 2080D. Antibiotic SF 2080D [81910-07-8] [82987-07-3]
 C₁₁H₄Cl₅NO₂ 359.422
 Prod. by *Actinosporangium vitaminophilum* SF-2080. Active against gram-positive and -negative bacteria and fungi. Yellow needles (C₆H₆). Sol. Me₂CO, bases, EtOAc; fairly sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 195-198°. Pyrrole

antibiotic. λ_{max} 220 (sh) (ε 26900); 336 (ε 13800); 370 (ε 14500) (MeOH) (Derep). λ_{max} 310 (MeOH/HCl) (Berdy). λ_{max} 331 (MeOH/NaOH) (Berdy).

► LD₅₀ (mus, ipr) 20 mg/kg. PC4949000

4-Chloro, *Me ether: Pyrrolomycin J*
 C₁₂H₆Cl₅NO₂ 373.448


Prod. by *Streptomyces fumanus* LL-F42248. Yellow solid.

[82987-08-4]

Ezaki, N. *et al.*, *J. Antibiot.*, 1983, **36**, 1263-1267; 1483-1489 (*isol, uv, ir, pmr, cmr*)
 Charan, R.D. *et al.*, *J. Nat. Prod.*, 2005, **68**, 277-279 (*Pyrrolomycins I-J*)


Pyrrolomycin E P-965

2,4-Dichloro-6-(5-chloro-3-nitro-1H-pyrrol-2-yl)phenol, 9CI. 5-Chloro-2-(3,5-dichloro-2-hydroxyphenyl)-3-nitropyrrole. SF 2080E. Antibiotic SF 2080E [87376-16-7]



C₁₀H₅Cl₃N₂O₃ 307.519
 Pyrrole antibiotic. Isolated from *Actinosporangium vitaminophilum*. Active against gram-positive bacteria. Yellow needles (C₆H₆). Sol. MeOH, bases; fairly sol. EtOAc, CHCl₃; poorly sol. H₂O, hexane. Mp 250°. λ_{max} 221 (ε 39400); 245 (sh) (ε 15600); 330 (ε 5590); 340 (ε 6170) (MeOH) (Derep). λ_{max} 312 (MeOH/NaOH) (Berdy).
 Ezaki, N. *et al.*, *J. Antibiot.*, 1983, **36**, 1263; 1483 (*isol, struct*)

Pyrrolomycin F P-966



	R	R'
Pyrrolomycin F ₁	Br	Br
Pyrrolomycin F _{2a}	Cl	Br
Pyrrolomycin F _{2b}	Br	Cl
Pyrrolomycin F ₃	Cl	Cl

Pyrrole antibiotic complex. Consists of 4 components.

Pyrrolomycin F₁

2,3,4-Tribromo-5-(5-bromo-2-hydroxybenzoyl)pyrrole. SF 2080F₁. Antibiotic SF 2080F₁ [88217-19-0]
 C₁₁H₅Br₄NO₂ 502.782
 From *Actinosporangium vitaminophilum*. Active against gram-positive bacteria and fungi. Yellow needles (C₆H₆/hexane). Mp 188-190°.

► PC4937680

Pyrrolomycin F_{2a}

2,4-Dibromo-5-(5-bromo-2-hydroxybenzoyl)-3-chloropyrrole. SF 2080F_{2a}. Antibiotic SF 2080F_{2a} [88477-78-5]

C₁₁H₅Br₃ClNO₂ 458.331
 From *Actinosporangium vitaminophilum* ATCC31673. Active against gram-positive bacteria and fungi. Yellow needles (C₆H₆/hexane). Sol. MeOH, bases, EtOAc; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 182°. λ_{max} 267 ; 315 (MeOH) (Berdy).

Pyrrolomycin F_{2b}

3,4-Dibromo-2-(5-bromo-2-hydroxybenzoyl)-5-chloropyrrole. SF 2080F_{2b}. Antibiotic SF 2080F_{2b} [88477-79-6]

C₁₁H₅Br₃ClNO₂ 458.331
 From *Actinosporangium vitaminophilum* ATCC31673. Active against gram-positive bacteria and fungi. Yellow needles (C₆H₆/hexane). Sol. MeOH, bases, EtOAc; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 206-208°. λ_{max} 267 ; 315 (MeOH) (Berdy).


Pyrrolomycin F₃

3-Bromo-2-(5-bromo-2-hydroxyphenyl)-4,5-dichloropyrrole. SF 2080F₃. Antibiotic SF 2080F₃ [88228-92-6]

C₁₁H₅Br₂Cl₂NO₂ 413.879
 From *Actinosporangium vitaminophilum* ATCC31673. Active against gram-positive bacteria and fungi. Yellow needles (C₆H₆/hexane). Sol. MeOH, EtOAc, bases; fairly sol. CHCl₃; poorly sol. H₂O, hexane. Mp 192-193°. λ_{max} 268 ; 314 (MeOH) (Berdy).

Shomura, T. *et al.*, *Int. J. Syst. Bacteriol.*, 1983, **33**, 557 (*isol*)
 Ezaki, N. *et al.*, *J. Antibiot.*, 1983, **36**, 1431 (*isol, uv, ir, pmr, ms*)
 Japan. Pat., 1983, 83 126 863; CA, **100**, 21555 (*isol*)

Pyrrolomycin G P-967



Absolute Configuration


C₁₁H₆Cl₄N₂O₄ 371.99
 Prod. by *Streptomyces fumanus* (culture LL-F42248). Yellow solid. [α]_D -15.6 (c, 0.28 in MeOH).

2'-Me ether: Pyrrolomycin H

C₁₂H₈Cl₄N₂O₄ 386.017
 Prod. by *Streptomyces fumanus* (strain LL-F42248). Yellow solid. [α]_D -91 (c, 0.3 in MeOH).

Charan, R.D. *et al.*, *J. Nat. Prod.*, 2005, **68**, 277-279 (*isol, cd, pmr, cmr*)

Pyrrolo[1,2-*a*]pyrazine-3,6(4*H*,7*H*)-dione, 9CI
[870136-68-8]




C₇H₆N₂O₂ 150.137
Prod. by a marine-derived *Bacillus* sp.

Zhang, H.-L. *et al.*, *Zhongguo Yaowu Huaxue Zazhi*, 2003, **13**, 294-296; *CA*, **144**, 2926x (isol)

Pyrrolosporin A

BMY 42448. Antibiotic BMY 42448
[140395-71-7]




C₄₄H₅₄Cl₂N₂O₁₀ 841.823
Prod. by *Micromonospora narashinoensis*. Antitumour agent. Cryst. Mp 235°. [α]_D²⁶ +2.8 (c, 0.2 in MeOH). λ_{max} 268 ; 341 (MeOH).

U.S. Pat., 1992, 5 082 933; *CA*, **116**, 190938w (ir, uv, pmr, cmr)
Lam, K.S. *et al.*, *J. Antibiot.*, 1996, **49**, 860 (props)
Schroeder, D.R. *et al.*, *J. Antibiot.*, 1996, **49**, 865 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Pyrrolosedamine

2-(2-Hydroxy-2-phenylethyl)-1-methylpyrrolidine. 2-(1-Methyl-2-pyrrolidinyl)-1-phenylethanol. 1-Methyl-α-phenyl-2-pyrrolidineethanol, 9CI
[175447-14-0]




C₁₃H₁₉NO 205.299
Alkaloid from leafy parts of *Sedum oryzifolium*.

Stereoisomer: *Pyrrolallosedamine*
[175447-15-1]
C₁₃H₁₉NO 205.299

P-968

From leafy parts of *Sedum oryzifolium*.
Kim, J.H. *et al.*, *Phytochemistry*, 1996, **41**, 1319 (*Pyrrolosedamine*, *Pyrrolallosedamine*)

3-(1*H*-Pyrrol-3-yl)-2-propenoic acid, 9CI
3-Pyrroleacrylic acid
[856597-79-0]



C₇H₇NO₂ 137.138

(*E*)-form [85310-57-2]

Isol. from *Streptomyces parvulus*. No antimicrobial activity. Cryst. (EtOAc). Mp 182-184°. λ_{max} 300 (MeOH) (Berdy).

Me ester: [97055-89-5]
C₈H₉NO₂ 151.165
Cryst. (Et₂O/petrol). Mp 82-83°.

Amide: [97055-88-4]
C₇H₈N₂O 136.153
From *Streptomyces parvulus*. Cryst. (EtOAc). Mp 142-143°. λ_{max} 300 (MeOH) (Berdy).

Dimethylamide: [897921-42-5]
C₉H₁₂N₂O 164.207
Pale yellow cryst. (EtOAc/hexane). Mp 171-173°.


Nitrile: [897921-49-2]
C₇H₆N₂ 118.138
Light brown solid. Mp 106-108°.

N-4-Methylbenzenesulfonyl: [864815-12-3]
C₁₄H₁₃NO₄S 291.327
Yellow cryst. (EtOAc/hexane). Mp 184-185°.

Labouta, I.M. *et al.*, *Acta Chem. Scand., Ser. B*, 1982, **36**, 669-674 (synth, ir, uv, pmr)
Keller-Schierlein, W. *et al.*, *Helv. Chim. Acta*, 1985, **68**, 559-562 (isol, ir, uv, pmr, ms, *Me ester*, *amide*)
Nishikawa, Y. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 684-687 (synth, pmr)
Karousis, N. *et al.*, *Synthesis*, 2006, 1494-1498 (synth, *amide*, *dimethylamide*, *nitrile*)

3-(2-Pyrrolyl)pyridine

2-(3-Pyridinyl)pyrrole. β-Nornicotyrine.
Nornicotyrine
[494-98-4]



C₉H₈N₂ 144.176
Needles (petrol/C₆H₆). V. sol. EtOH, Et₂O, CHCl₃, C₆H₆; spar. sol. petrol, H₂O, EtOH. Mp 102°. pK_a 4.35 (24°). Et₂O solns. show blue fluor. Gives cryst. K salt.

► Exp. tumourigenic data. UT7875000
Methodide:

P-971

Pale yellow needles (EtOH). Mp 170-171°.

Picrate:

Yellow prisms (H₂O or EtOH). Mp 203-205°.

N-Me: 3-(1-Methyl-1*H*-pyrrol-2-yl)pyridine, 9CI. *Nicotyrine*. β-*Nicotyrine*
[487-19-4]
C₁₀H₁₀N₂ 158.202

Alkaloid from cured tobacco (*Nicotiana tabacum*) (Solanaceae). Shows insecticidal props. Liq. Bp 281°. pK_a 4.76 (24°).

► LD₅₀ (mus, orl) 575 mg/kg. UT6130000
N-Me, picrate: Mp 170-171°.

Späth, E. *et al.*, *Ber.*, 1937, **70**, 2450 (*Nicotyrine*)
Swain, M.L. *et al.*, *J.A.C.S.*, 1949, **71**, 1341 (*Nicotyrine*)

Ryang, H. *et al.*, *Chem. Comm.*, 1972, 594-595 (synth)

Shibagaki, M. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 2651 (synth)

Mullen, G.B. *et al.*, *J.O.C.*, 1989, **54**, 2476 (synth, ir, pmr, bibl)

Savoia, D. *et al.*, *J.O.C.*, 1991, **56**, 1822 (synth, ir, pmr, ms)


Noland, W.E. *et al.*, *Acta Cryst. C*, 2003, **59**, o263-o267 (cryst struct)

Wei, X. *et al.*, *Synth. Commun.*, 2003, **33**, 3305-3315 (*Nicotyrine*, synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, NDX300; PQB500

Pyrromycin A

P-973



C₂₃H₂₉N₁₁O₅ 539.553

Pyrrole antibiotic. Prod. by *Streptomyces* sp. KY11768. Active against gram-positive and -negative bacteria and tumours.

7'-*N*-Hydroxy: *Pyrromycin B*

C₂₃H₂₉N₁₁O₆ 555.552


Prod. by *Streptomyces* sp. KY11768. Active against gram-positive and -negative bacteria and tumours.

Asai, A. *et al.*, *J. Antibiot.*, 2000, **53**, 66-69 (isol, activity)

Pyrroxamycin

P-974

Dioxapyrrolomycin. Al-R 2081. LL-F42248a. SS 46506A. Antibiotic Al-R 2081. Antibiotic LL-F42248a. Antibiotic SS 46506A. Antibiotic HS 3. HS 3
[105888-54-8]



C₁₂H₆Cl₄N₂O₄ 384.001

Pyrrole antibiotic. Prod. by *Streptomyces*

spp. Active against gram-positive bacteria and dermatophytes. Pale-yellow cryst. (EtOH) or (C₆H₆). Mp 216-220° dec. (200-207°). $[\alpha]_D^{25}$ -110 (c, 0.5 in EtOH). $[\alpha]_D^{25}$ -37.2 (c, 1 in Me₂CO). λ_{\max} 211 (ε 27500); 273 (ε 7080); 321 (ε 3470) (MeOH/HCl) (Derep). λ_{\max} 212 (ε 30200); 296 (ε 6920); 319 (ε 8910) (MeOH/NaOH) (Derep). λ_{\max} 211 (ε

27500); 278 (ε 6030); 320 (ε 4470) (MeOH) (Derep). λ_{\max} 220 (sh) (ε); 276 (ε 6800); 285 (ε 6950); 294 (ε 6500); 322 (ε 5300) (MeOH) (Derep).

► UX9458500

[118187-02-3]

Rengaraju, S. *et al.*, *CA*, 1987, **106**, 152653 (*isol*)

Carter, G.T. *et al.*, *J. Antibiot.*, 1987, **40**, 233-236 (*isol, struct, props*)

Nakamura, H. *et al.*, *J. Antibiot.*, 1987, **40**, 899-903 (*isol, struct, props*)


Yano, K. *et al.*, *J. Antibiot.*, 1987, **40**, 961-969 (*isol, struct, props*)

Masuda, K. *et al.*, *J. Antibiot.*, 1991, **44**, 533-540 (*isol*)

Charan, R.D. *et al.*, *J. Nat. Prod.*, 2006, **69**, 29-33 (*biosynth*)

Quadrigemine I

[235099-49-7]



Absolute Configuration

C₄₄H₅₀N₈ 690.933

Alkaloid from *Psychotria oleoides*. Amorph. powder. [α]_D²⁵ +16 (c, 0.34 in CHCl₃). [α]_D²⁰ +199 (c, 0.34 in EtOH). λ_{max} 244 (log ε 4.45); 302 (log ε 4.13) (EtOH).

Homologue (n = 4): Oleoidine

C₅₅H₆₂N₁₀ 863.162
Alkaloid from *Psychotria oleoides*. Amorph. yellow powder. [α]_D²⁵ +89 (c, 0.4 in CHCl₃). [α]_D²⁰ +371 (c, 0.2 in EtOH). λ_{max} 244 (log ε 4.55); 302 (log ε 4.26) (EtOH).


Homologue (n = 5): Caledonine

[235099-23-7]
C₆₆H₇₄N₁₂ 1035.391
Alkaloid from *Psychotria oleoides*. Amorph. yellow powder. [α]_D²⁵ +125 (c, 0.4 in CHCl₃). [α]_D²⁰ +462 (c, 0.2 in EtOH). λ_{max} 243 (log ε 4.62); 303 (log ε 4.35) (EtOH).

Jannic, V. et al., *J. Nat. Prod.*, 1999, **62**, 838-843 (isol, uv, ir, cd, pmr, cmr, ms)

Quadrigemine A

[69937-02-6]



C₄₄H₅₀N₈ 690.933

Minor alkaloid from *Hodgkinsonia frutescens* and *Psychotria forsteriana* (Rubiaceae). Cytotoxic. Amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²³ +32 (EtOH). Probably a mixt. of diastereoisomers. λ_{max} 243 (ε 17380); 302 (ε 9120) (EtOH) (Berdy).

Stereoisomer: Quadrigemine C

[112295-93-9]
C₄₄H₅₀N₈ 690.933
Alkaloid from *Psychotria oleoides* and *Calycodendron milnei* (Rubiaceae). Cytotoxic. Shows analgesic props. Amorph. [α]_D -69 (c, 1 in CHCl₃). Probable stereochem. has been suggested.

Q-1

Stereoisomer (2): Alkaloid C†

C₄₄H₅₀N₈ 690.933
Alkaloid from leaves of *Psychotria forsteriana*. Exhibits potent cytotoxicity. [α]_D²⁰ +32.5 (c, 0.2 in EtOH). [α]_D²⁰ +115.7 (c, 0.2 in CHCl₃).

Parry, K.P. et al., *J.C.S. Perkin 1*, 1978, 1671 (isol, uv, ir, ms, pmr, struct)

Roth, A. et al., *Planta Med.*, 1986, **52**, 450 (Alkaloid C)

Libot, F. et al., *J. Nat. Prod.*, 1987, **50**, 468 (Quadrigemine C)


Adjibade, Y. et al., *Planta Med.*, 1989, **55**, 567 (isol, props)

Jannic, V. et al., *J. Nat. Prod.*, 1999, **62**, 838-843 (Quadrigemine C, stereochem)

Lebsack, A.D. et al., *J.A.C.S.*, 2002, **124**, 9008-9009 (Quadrigemine C, synth)

Quadrigemine B

[69937-10-6]



C₄₄H₅₀N₈ 690.933

Minor alkaloid from *Hodgkinsonia frutescens* (Rubiaceae), *Psychotria forsteriana* and *Psychotria rostrata*. Cytotoxic. Active against gram-positive bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 229-243°. [α]_D²³ +263 (EtOH). Probably a mixt. of diastereoisomers. λ_{max} 244 (ε 17780); 302 (ε 6040) (EtOH) (Berdy).

Stereoisomer: Psychotria Alkaloid B

C₄₄H₅₀N₈ 690.933
Alkaloid from leaves of *Psychotria forsteriana*. Exhibits potent cytotoxicity. [α]_D²⁰ +215 (EtOH). [α]_D²⁰ -10 (CHCl₃).


Parry, K.P. et al., *J.C.S. Perkin 1*, 1978, 1671 (isol, uv, ir, pmr, ms, struct)

Roth, A. et al., *Planta Med.*, 1986, 450 (Alkaloid B)

Mahmud, Z. et al., *Int. J. Pharmacogn.*, 1993, **31**, 142-146 (isol, activity)

Quadrigemine H

[168611-79-8]



Q-4


C₄₄H₅₀N₈ 690.933

Alkaloid from *Calycodendron milnei*.

Saad, H.-E.A. et al., *Planta Med.*, 1995, **61**, 313-316

3,2':3',4'':2'',3'''-Quaterpyridine, 9CI Nemertelline

[59697-14-2]



Q-5

C₂₀H₁₄N₄ 310.357

Struct. revised in 1995. Alkaloid from the marine hoplonemertine *Amphiporus angulatus*. Neurotoxin. Cryst. (Et₂O). Mp 154-156°.

Kem, W.R. et al., *Experientia*, 1976, **32**, 684-686 (isol)

Cruskie, M.P. et al., *J.O.C.*, 1995, **60**, 7491-7495 (synth, pmr, cmr, cryst struct)


Zoltewicz, J.A. et al., *Tetrahedron*, 1995, **51**, 11401-11410 (synth)

Bouillon, A. et al., *J.O.C.*, 2003, **68**, 10178-10180 (synth, pmr, cmr)

Quebrachamine

Q-6

7-Ethyl-1,4,5,6,7,8,9,10-octahydro-2H-3,7-methanoazacycloundecino[5,4-b]indole, 9CI. Kamassine



(R)-form

C₁₉H₂₆N₂ 282.428

(R)-form [4850-21-9]

Alkaloid from *Aspidosperma quebrachoblanco* (quebracho), *Aspidosperma chakensis* and several other *Aspidosperma* spp., *Gonioma kamassi*, *Hunteria elliptii* and *Rhazya stricta* (Apocynaceae). Shows DNA-binding properties, α₂ receptor antagonist. Cryst. (C₆H₆, hexane or EtOH). Mp 147-149°. [α]_D -100 (CHCl₃). [α]_D -110 (c, 0.496 in Me₂CO).

Picrate:

Scarlet needles (EtOH). Mp 195-196°.

N-Me: N-Methylquebrachamine

[2671-48-9]

C₂₀H₂₈N₂ 296.455

Alkaloid from *Vallesia dichotoma* (Apocynaceae). Cryst. (EtOH). Mp 82-84° (nat. prod.) Mp 89° (synthetic). [α]_D -109 (c, 0.5 in EtOH) (synthetic).

11-Methoxy: 11-Methoxyquebrachamine

C₂₀H₂₈N₂O 312.454

Alkaloid from the stem bark of *Melodinus australis* (Apocynaceae). Prisms (Et₂O/pentane). Mp 157-160°. [α]_D +68 (c, 1.3 in CHCl₃). Dec. in CHCl₃ soln. Struct. uncertain, may be the 12-methoxy isomer.

(S)-form [14430-17-2]

Alkaloid from *Vinca erecta*, the leaves of *Pleiocarpa tubicina*, the root bark of *Pleiocarpa pycnantha* var. *pycnantha*, and the bark of *Stemmadenia donnell-smithii* (Apocynaceae). Cryst. (MeOH or Et₂O/petrol). Mp 147-149°. [α]_D +98 (CHCl₃). [α]_D +108.9 (Me₂CO).

Picrate:

Bright red cryst. Mp 190-193°.

N-Me: Synthetic. Cryst. (MeOH). Mp 86-87°. [α]_D +110 (CHCl₃).

14β-Hydroxy: 14β-Hydroxyquebrachamine

C₁₉H₂₆N₂O 298.427

Alkaloid from the leaves of *Stemmadenia grandiflora* (Apocynaceae). Mp 123-126°. [α]_D²¹ +86 (c, 0.14 in CHCl₃).

(±)-form [7689-01-2]

Synthetic. Needles (MeOH). Mp 113-116°.

Picrate:

Cryst. (MeOH). Mp 176-179°.

N-Me: Alkaloid from *Vinca minor* (Apocynaceae). Cryst. (EtOH). Mp 70-72°.

Hesse, O. *et al.*, *Annalen*, 1882, **211**, 249 (*isol*)

Field, E. *et al.*, *J.C.S.*, 1924, **125**, 1444 (*derivs*)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1951, **34**, 920 (*isol, uv*)

Gellért, E. *et al.*, *Helv. Chim. Acta*, 1952, **35**, 114 (*struct*)

Orazi, O.O. *et al.*, *J.O.C.*, 1956, **21**, 978 (*isol, ord*)

Witkop, B. *et al.*, *J.A.C.S.*, 1957, **79**, 3193 (*ir, degradn*)

Walls, F. *et al.*, *Tetrahedron*, 1958, **2**, 173 (*isol, uv*)

Cohen, L.A. *et al.*, *J.A.C.S.*, 1960, **82**, 2184 (*pmr*)

Biemann, K. *et al.*, *J.A.C.S.*, 1962, **84**, 4578 (*ms*)

Mokry, J. *et al.*, *Tet. Lett.*, 1962, 1185 (*N-Me*)

Stork, G. *et al.*, *J.A.C.S.*, 1963, **85**, 2872 (*synth*)

Trojánek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1964, **29**, 1904 (*synth, ms, N-Me*)

Bycroft, B.W. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1147 (*isol*)

Walsler, A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 391 (*N-Me, isol, uv, ir, pmr, ms, synth*)

Linde, H.H.A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 1822 (*11-Methoxyquebrachamine*)

Kutney, J.P. *et al.*, *J.A.C.S.*, 1966, **88**, 3656; 1968, **90**, 3891 (*synth*)

Dugan, J.J. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 60 (*synth*)

Gorman, A.A. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 33 (*isol, uv*)

Barth, G. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 2168 (*N-Me, cd, mcd*)

Zsádon, B. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1973, **78**, 207 (*synth, N-Me*)

Bruneton, J. *et al.*, *Phytochemistry*, 1974, **13**, 1963 (*11-Methoxyquebrachamine*)

Puglisi, C. *et al.*, *Acta Cryst. B*, 1976, **32**, 1900 (*cryst struct*)

Wenkert, E. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2711 (*cmr*)

Takano, S. *et al.*, *J.A.C.S.*, 1979, **101**, 6414 (*synth*)

Takano, S. *et al.*, *Chem. Comm.*, 1980, 616; 1981, 1153 (*synth*)

Giri, V.S. *et al.*, *J. Het. Chem.*, 1980, **17**, 1133 (*synth, uv, ir, pmr, ms*)

Wenkert, E. *et al.*, *Tetrahedron*, 1981, **37**, 4017 (*synth*)

Ban, Y. *et al.*, *Tetrahedron*, 1983, **39**, 3657 (*synth*)

Mandal, S.B. *et al.*, *J.O.C.*, 1988, **53**, 4236 (*synth*)

Torrenegra, R. *et al.*, *Phytochemistry*, 1988, **27**, 1843 (*14β-Hydroxyquebrachamine*)

Asaoka, M. *et al.*, *Heterocycles*, 1989, **29**, 243 (*synth*)

Node, M. *et al.*, *J.O.C.*, 1990, **55**, 517 (*synth*)

Deutsch, H.F. *et al.*, *J. Pharm. Biomed. Anal.*, 1994, **12**, 1253-1287 (*isol, activity*)

Temme, O. *et al.*, *J.O.C.*, 1998, **63**, 6007-6015 (*synth*)

Wee, A.G.H. *et al.*, *Tetrahedron*, 1998, **54**, 13435-13448 (*synth*)

Meier, A. *et al.*, *J. Antibiot.*, 1999, **52**, 957-959 (*activity*)


Eles, J. *et al.*, *J. Het. Chem.*, 2002, **39**, 767-771 (*synth*)

Kozmin, S.A. *et al.*, *J.A.C.S.*, 2002, **124**, 4628-4641 (*synth*)

Fujimura, T. *et al.*, *Tet. Lett.*, 2002, **43**, 97-99 (*synth*)

***Aesculus hippocastanum* Quercetin indoxylacetylglucoside**

Q-7



C₄₂H₄₅NO₂₄ 947.81

Not named in the lit. Constit. of *Aesculus hippocastanum*. [α]_D²⁰ +52.8 (c, 0.2 in MeOH).


Huebner, G. *et al.*, *Planta Med.*, 1999, **65**, 636-642

Quettamine

[77844-42-9]

[82373-00-0]

Q-8



C₁₉H₂₂NO₃⁺ 312.388

Alkaloid from *Berberis baluchistanica* (Berberidaceae). Noncryst. (as chloride). Opt. inactive.


Zarga, M.H.A. *et al.*, *Tet. Lett.*, 1981, **22**, 541 (*uv, pmr, ms, struct*)

Chattopadhyay, S. *et al.*, *Heterocycles*, 1982, **19**, 697 (*synth*)

Queuine

Q-9

2-Amino-5-[[[4,5-dihydroxy-2-cyclopenten-1-yl]amino]methyl]-1,7-dihydro-4H-pyrrolo[2,3-d]pyrimidin-4-one, 9CI. 2-Amino-5-(4,5-dihydroxycyclopent-1-en-3-ylaminomethyl)pyrrolo[2,3-d]pyrimidine. 7-(4,5-Dihydroxy-2-cyclopent-1-en-3-ylaminomethyl)-7-deazaguanine. *Q* base [72496-59-4] [86496-18-6]



Absolute configuration

C₁₂H₁₅N₅O₃ 277.282

Prisms (as dihydrochloride). Mp 230-235° dec. (hydrochloride). [α]_D²⁶ +113 (c, 0.3 in H₂O).

7-β-D-Ribofuranosyl: **Queuosine**. Nucleoside *Q* [57072-36-3] [71050-05-0]

C₁₇H₂₃N₅O₇ 409.398

Hypermodified nucleoside occurring in the first anticodon posn. in tRNAs of plants and animals. Pale yellow cryst. (H₂O); glassy solid (as hydrochloride). Mp 225-230° dec. [α]_D +43.4 (c, 0.1 in H₂O).

7-O-β-D-Ribofuranosyl, 4'-O-β-D-galactopyranoside: **Galactosylqueuosine** [60426-58-6]

C₂₃H₃₃N₅O₁₂ 571.54

Modified nucleoside found in tRNAs.

7-O-β-D-Ribofuranosyl, 4'-O-β-D-mannopyranoside: **Mannosylqueuosine** [60398-20-1]

C₂₃H₃₃N₅O₁₂ 571.54

Modified nucleoside found in tRNAs.

Kasai, H. *et al.*, *J.A.C.S.*, 1976, **98**, 5044-5046 (*Galactosylqueuosine, Mannosylqueuosine*)


Ohgi, T. *et al.*, *J.A.C.S.*, 1979, **101**, 3629 (*deriv, synth, abs config, bibl*)

Akimoto, H. *et al.*, *J. Med. Chem.*, 1986, **29**, 1749 (*biochem, ir, pmr, bibl*)

Kondo, T. *et al.*, *Tetrahedron*, 1986, **42**, 207
(*synth, uv, pmr, ir*)
Akimoto, H. *et al.*, *J.C.S. Perkin 1*, 1988, 1637
(*synth, ir, pmr*)
Barnett, C.J. *et al.*, *Tetrahedron*, 2000, **56**,
9221-9225 (*synth, cmr*)

Quimbeline
[55786-21-5]

Q-10




C₄₃H₄₈N₄O₆ 716.875
Alkaloid from the root bark of *Voacanga chaloniana* (Apocynaceae). Mp 270°. [α]_D -195 (c, 1 in CHCl₃).

Bombardelli, E. *et al.*, *Experientia*, 1975, **31**,
139 (*uv, ir, pmr, cmr, ms, struct*)

Quinaldopeptin

Q-11

BMV 28662. BU 3845T. Antibiotic BMY 28662. Antibiotic BU 3845T
[130743-07-6]




C₆₂H₇₈N₁₄O₁₄ 1243.384
Cyclic peptide antibiotic. Related to Luzopeptin and Sandramycin, S-50. Prod. by *Streptovercillium album* and an *Amycolata* spp. Active against gram-positive bacteria and anaerobes. Exhibits cytotoxic and anthelmintic activity. Rods (CHCl₃/Et₂O). Sol. Py; fairly sol. DMSO; poorly sol. H₂O, Me₂CO, hexane. Mp 300°. [α]_D²⁴ +51 (c, 0.5 in Py). λ_{max} 211 (sh); 222; 251; 307 (sh); 396 (MeOH/NaOH) (Derep). λ_{max} 214 (ε 78000); 230 (ε 73000); 299 (ε 9700); 359 (ε 9300) (MeOH) (Derep). λ_{max} 214 ; 230 ; 299 ; 359 (MeOH/HCl) (Berdy).

Toda, S. *et al.*, *J. Antibiot.*, 1990, **43**, 796 (*isol, struct*)
Rickards, R.W. *et al.*, *J. Antibiot.*, 1998, **51**,
1093-1095 (*isol, activity*)

Quinamine
[464-85-7]

Q-12



Absolute Configuration

C₁₉H₂₄N₂O₂ 312.411
Stereochem. not clear. CAS assigns the abs. config. as shown, but proof for this is not clear. Minor alkaloid from *Cinchona ledgeriana* and many other *Cinchona* spp. (Rubiaceae). No detectable antimalarial props. Mp 185-186° (172°). [α]_D +104.5 (c, 2 in EtOH).

Hydrochloride:
Cryst. + 1H₂O. Mp 166-167°. [α]_D +103 (H₂O).

Picrate: Mp 175-176°. [α]_D +90 (Me₂CO).

9,10-Dihydro: Dihydroquinamine

[22660-97-5]
C₁₉H₂₆N₂O₂ 314.427
Main alkaloid of *Iserertia hypoleuca* (Rubiaceae). Silky needles (Me₂CO aq.). Mp 154-156°. [α]_D +36 (c, 0.113 in CHCl₃).

3-Epimer: Conquinamine. Epiquinamine

[464-86-8]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from *Cinchona succirubra*, *Cinchona rosulenta* and *Cinchona ledgeriana* (Rubiaceae). No detectable antimalarial props. Large prisms. Mp 123° Mp 115-118°. [α]_D +144 (c, 0.4 in CHCl₃). [α]_D +164 (c, 0.4 in EtOH). Conquinamine (1877), semi-synthetic epiquinamine (1950) and natural 3-epiquinamine (1980) have not been compared, but they are prob. identical. λ_{max} 242 ; 300 (MeOH).

3-Epimer, 9,10-dihydro: 3-Epidihydroquinamine

[89708-22-5]
C₁₉H₂₆N₂O₂ 314.427
Alkaloid from *Iserertia hypoleuca* (Rubiaceae). Mp 181-185°. [α]_D²⁰ -57 (c, 0.71 in CHCl₃).

Stereoisomer: Cinchona ledgeriana Alkaloid C

[77549-88-3]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from *Cinchona ledgeriana* leaves (Rubiaceae). Noncryst. [α]_D +35 (CHCl₃). Stereochem. unknown.


Hesse, O. *et al.*, *Ber.*, 1877, **10**, 2152-2162 (*Quinamine, Conquinamine, isol*)
Goutarel, R. *et al.*, *Helv. Chim. Acta*, 1950, **33**,
150-164 (*uv, ir, struct*)
Witkop, B. *et al.*, *J.A.C.S.*, 1950, **72**, 2311-2312 (*struct*)
Culvenor, C.C.J. *et al.*, *J.C.S.*, 1950, 1485-1493 (*uv, struct*)

Bohlmann, H. *et al.*, *Phytochemistry*, 1969, **8**,
645-652 (*Dihydroquinamine*)
Zeches, M. *et al.*, *Phytochemistry*, 1980, **19**,
2451-2454 (*Epiquinamine, Alkaloid C*)
Mulder-Krieger, T. *et al.*, *Pharm. Weekbl., Sci. Ed.*, 1982, 91-92 (*cmr*)
Esterl, A. *et al.*, *Planta Med.*, 1983, **49**, 244-
245 (*Dihydroquinamine, Epidihydroquinamine*)

2,4(1H,3H)-Quinazolidione, 9CI

Q-13

2,4-Quinazolidinol. 2,4-Dihydroxyquinazoline. *Benzoyleneurea*
[86-96-4]



C₈H₆N₂O₂ 162.148
Di-NH-form predominates for parent compd. Constit. of *Strobilanthes cusia*. Prisms (DMF/CHCl₃). Mp 358°.
▶ LD₅₀ (rat, ipr) 1200 mg/kg. Exp. reprod. effects. VA1390000

Diol-form

Di-Me ether: 2,4-Dimethoxyquinazoline
C₁₀H₁₀N₂O₂ 190.201
Mp 66-67°.

Di-NH-form

1-Me: 1-Methyl-2,4(1H,3H)-quinazolidione, 9CI. Glycosmicine
[604-50-2]
C₉H₈N₂O₂ 176.174
Minor alkaloid from the leaves of *Glycosmis arborea* (Rutaceae). Mp 270-271°.
▶ VA1411500

3-Me: 3-Methyl-2,4(1H,3H)-quinazolidione
[607-19-2]
C₉H₈N₂O₂ 176.174
Solid. Mp 242-243.5°.

1,3-Di-Me: 1,3-Dimethyl-2,4(1H,3H)-quinazolidione

[1013-01-0]
C₁₀H₁₀N₂O₂ 190.201
Sex pheromone of the pale brown chafer *Phyllopertha diversa*. Shows antiinflammatory, analgesic and anticonvulsant activity. Cryst. (H₂O). Mp 165-166° (170°). λ_{max} 220 (log ε 4.7); 312 (log ε 3.63) (no solvent reported).

3-Ph: 3-Phenyl-2,4(1H,3H)-quinazolidione, 9CI

[603-23-6]
C₁₄H₁₀N₂O₂ 238.245
Cryst. (MeOH). Mp 282°.

3-Ph, 1-Ac: [35118-04-8]

C₁₆H₁₂N₂O₃ 280.282
Cryst. (EtOH). Mp 202°.

3-Amino: 3-Amino-2-hydroxy-4(3H)-quinazolinone. 3-Amino-2,4(1H,3H)-quinazolidione

[30386-01-7]
C₈H₇N₃O₂ 177.162
Cryst. (H₂O). Mp 291-293°.

3-Benzyl: 3-Benzyl-2,4(1H,3H)-quinazo-

linedione

[1932-42-9]

C₁₅H₁₂N₂O₂ 252.272Solid (Et₂O). Mp 226-226.5°.*Aldrich Library of NMR Spectra*, 2nd edn., 1983, **2**, 359C (nmr)*Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 882B (ir)Kunckell, F. *et al.*, *Ber.*, 1910, **43**, 1021 (3-amino)*Org. Synth.*, *Coll. Vol.*, **2**, 1943, 79 (synth)Pakrashi, S.C. *et al.*, *Tetrahedron*, 1963, **19**, 1011 (uv, ir, pmr, ms, struct, Glycosmicine)Spence, T.W.M. *et al.*, *J.C.S. Perkin I*, 1972, 97 (3-Ph)Fahmy, A.F.M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1978, **51**, 2148 (synth)Yamamoto, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1633 (synth)Michman, M. *et al.*, *Org. Prep. Proced. Int.*, 1978, **10**, 13 (synth, ms)Petridou-Fischer, J. *et al.*, *J. Het. Chem.*, 1982, **19**, 123 (cmr)Papadopoulos, E.P. *et al.*, *J. Het. Chem.*, 1982, **19**, 123; 269 (synth, pmr, cmr)Lalezari, I. *et al.*, *J. Het. Chem.*, 1984, **21**, 5Yoshida, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1987, **60**, 1793 (synth)Beckwith, A.L.J. *et al.*, *Aust. J. Chem.*, 1990, **43**, 451 (synth)Gineinah, M.M. *et al.*, *J. Het. Chem.*, 1990, **27**, 723 (synth)Miyata, T. *et al.*, *Heteroat. Chem.*, 1991, **2**, 473 (synth)Li, L. *et al.*, *Yaouxue Xuebao*, 1993, **28**, 238 (isol)Leal, W.S. *et al.*, *Nature (London)*, 1997, **385**, 213 (occur, 1,3-di-Me)Pfeiffer, W.-D. *et al.*, *J. Het. Chem.*, 1999, **36**, 1327-1336 (3-amino)Choo, H.Y.P. *et al.*, *Bioorg. Med. Chem.*, 2002, **10**, 517-523 (3-Ph, synth, ir, pmr)Mizuno, T. *et al.*, *Tetrahedron*, 2002, **58**, 3155-3158 (synth)Wu, Y.-Q. *et al.*, *J. Het. Chem.*, 2003, **40**, 191-193 (synth, pmr)Petrov, J.S. *et al.*, *Org. Prep. Proced. Int.*, 2005, **37**, 560-565 (synth, ir, pmr, ms, 3-Ph)Huang, Y. *et al.*, *J. Het. Chem.*, 2007, **44**, 1421-1424 (3-Me, 3-benzyl)Mizuno, T. *et al.*, *Synthesis*, 2007, 2524-2528 (synth, ir, pmr, cmr, ms)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, QEJ800C₈H₆N₂O 146.1483H-form predominates. Isol. from the Chinese drug Ch'ang Shan (roots of *Dichroa febrifuga*) (Hydrangeaceae). Needles (EtOH). Mp 215-216°. pK_{a1} 2.06; pK_{a2} 9.87 (20°). Prob. an artifact.▶ LD₅₀ (mus, orl) 609 mg/kg. VA2300000*Hydrochloride*: [33238-79-8]

Mp 212-215° Mp 247°.

1H-form

N-Me: see 1-Methyl-4(1H)-quinazolinone, M-556

N-Et: 1-Ethyl-4(1H)-quinazolinone [16347-92-5]

C₁₀H₁₀N₂O 174.202

Mp 127-129°.

N-Ph: 1-Phenyl-4(1H)-quinazolinone [35242-42-3]

C₁₄H₁₀N₂O 222.246

Cryst. (EtOH or EtOAc). Mp 184° (180-181°).

3H-form [132305-20-5]

N-Me: 3-Methyl-4(3H)-quinazolinone [2436-66-0]

C₉H₈N₂O 160.175

Mp 104-106°.

N-Et: 3-Ethyl-4(3H)-quinazolinone [3476-65-1]

C₁₀H₁₀N₂O 174.202

Mp 101-102°.

N-Ph: 3-Phenyl-4(3H)-quinazolinone, 9CI

[16347-60-7]

C₁₄H₁₀N₂O 222.246

Cryst. Mp 138-140°.

N-Ph; hydrochloride: Mp 222-223°.

N-(2-Hydroxyethyl): 3-(2-Hydroxyethyl)-4(3H)-quinazolinone [2981-86-4]

C₁₀H₁₀N₂O₂ 190.201

Needles. Mp 152-153°. This struct. was erroneously assigned to Echinazolinone, an alkaloid, in 1987.

N-(2-Phenylethenyl) (E-): (E)-Bogorin [220457-08-9]

C₁₆H₁₂N₂O 248.284Trace constit. of *Glycosmis* cf. *chlorosperma*. Cryst. (petrol). Mp 154-157°.

N-(2-Phenylethenyl) (Z-): 3-(2-Phenylethenyl)-4(3H)-quinazolinone. N-Styryl-4(3H)-quinazolinone. (Z)-Bogorin [220457-05-6]

C₁₆H₁₂N₂O 248.284Alkaloid from *Glycosmis* cf. *chlorosperma*. Cryst. Mp 73-76°. λ_{max} 227 (log ε 3.81); 269 (log ε 3.45); 310 (log ε 3.12) (EtOH). λ_{max} 227 (ε 6456); 269 (ε 2818); 310 (ε 1318) (MeOH) (Berdy).

N-(2-Carboxyphenyl): 3-(2-Carboxyphenyl)-4(3H)-quinazolinone [25380-15-8]

C₁₅H₁₀N₂O₃ 266.256Alkaloid from the roots of *Isatis indigotica* (Brassicaceae). Mp 279-281°.

N-(2-Hydroxyphenyl): 3-(2-Hydroxyphenyl)-4(3H)-quinazolinone [3977-51-3]

C₁₄H₁₀N₂O₂ 238.245Alkaloid from *Isatis indigotica* (Bras-

sicaceae). Mp 183-184° (synthetic).

OH-form [134434-33-6]

[17227-47-3]

Me ether: 4-Methoxyquinazolinone

[16347-95-8]

C₉H₈N₂O 160.175

Cryst. Mp 37.5°.

▶ VA1465000

Et ether: 4-Ethoxyquinazolinone


[16347-96-9]

C₁₀H₁₀N₂O 174.202

Cryst. Mp 46-48°.


Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 881D (ir)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 466C (nmr)Chou, T.Q. *et al.*, *J.A.C.S.*, 1948, **70**, 1765-1767 (isol, synth)Breukink, K.W. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1957, **76**, 401-414 (Me ether, synth, deriv)Armarego, W.F.L. *et al.*, *J. Appl. Chem.*, 1961, **11**, 70-72 (synth)Batterham, T.J. *et al.*, *J.C.S. (B)*, 1967, 892-897 (ms)Irwin, W.J. *et al.*, *J.C.S. Perkin I*, 1972, 353-355 (1-Ph, synth, uv, ms)Peet, N.P. *et al.*, *J.O.C.*, 1974, **39**, 1931-1935 (synth, N-carboxyphenyl)Kametani, T. *et al.*, *Heterocycles*, 1978, **9**, 1585-1591; 1980, **14**, 1469-1473 (synth, cmr)Bhattacharya, J. *et al.*, *Heterocycles*, 1980, **14**, 1469-1473 (cmr)Baig, G.U. *et al.*, *J.C.S. Perkin I*, 1984, 2765-2766 (aryl derivs)Reisch, J. *et al.*, *J. Nat. Prod.*, 1989, **52**, 404-407 (2-hydroxyethyl, synth, ir, pmr, ms)Chakrabarty, M. *et al.*, *Monatsh. Chem.*, 1995, **126**, 789-794 (pmr, cmr)Deng, K.M. *et al.*, *Chin. Chem. Lett.*, 1997, **8**, 237-238 (2-hydroxyphenyl)Wu, X. *et al.*, *Planta Med.*, 1997, **63**, 55-57 (isol, N-carboxyphenyl)Seger, C. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1926-1928 (synth, Bogorin)Wang, L. *et al.*, *Synthesis*, 2003, 1241-1247 (3-Ph)Wiklund, P. *et al.*, *J.O.C.*, 2004, **69**, 6371-6376 (3-N-Et)Shemchuk, L.A. *et al.*, *Russ. J. Org. Chem. (Engl. Transl.)*, 2007, **43**, 719-722 (synth, pmr)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, QFA000**4-Quinazolinol, 8CI****Q-14**

4-Hydroxyquinazolinone. 4(1H)-Quinazolinone, 9CI. Quinazolone [491-36-1]



OH-form


3H-form



1H-form


Quinazomycin**Q-15**

[28171-80-4]

C₅₁H₆₂N₁₂O₁₃S₂ 1115.255Depsipeptide antibiotic. Molecular formula and structure in original paper disagree. Prod. by *Streptomyces* sp. in the presence of 4-Oxo-3-quinazolineacetic acid. Active against *Staphylococcus aureus*. Cryst. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{max} 225 ; 244 ; 290 ; 328 (MeOH) (Berdy).Khan, A.W. *et al.*, *Indian J. Biochem.*, 1969, **6**, 220

Dhar, M.M. et al., *Pure Appl. Chem.*, 1971, **28**, 469


Quindolinocryptotackeine Q-16
[625830-78-6]



C₃₁H₂₀N₄ 448.526
Alkaloid from *Cryptolepis sanguinolenta*.

Blinov, K. et al., *Magn. Reson. Chem.*, 2003, **41**, 577-584 (isol, pmr, cmr)

Quinine Q-17
3-(3-Ethenyl-4-piperidinyl)-1-(6-methoxy-4-quinolinyl)-1-propanone, 9CI. Me-quiniverine. Quinotoxine. **Viquidil**, INN [84-55-9]



C₂₀H₂₄N₂O₂ 324.422
Alkaloid from many *Cinchona* spp. (Rubiaceae), poss. an artifact from rearr. of Quinine, Q-20. Readily obt. by thermal rearr. of quinine or quinidine. Spasmodic agent, cerebral vasodilator. Antiarrhythmic agent. Noncryst. Sol. Et₂O, C₆H₆, Et₂O, CHCl₃; poorly sol. H₂O, hexane. Mp 60°. [α]_D +39 (CHCl₃). Log P 3.03 (calc).

Hydrochloride: Desclidium. Permiran. Vasexeten [36342-92-4]
Mp 180-182°. [α]_D¹⁴ +16.3 (H₂O).

Oxime: Mp 112-116°.

9S-Alcohol, N-Me: **N-Methylquinicinalol** [83255-68-9]
C₂₁H₂₈N₂O₂ 340.464
Alkaloid from the bark of *Guettarda noumeana* (Rubiaceae). Gum. [α]_D -70 (c, 0.5 in CHCl₃). λ_{max} 224 (log ε 4.55); 277 (log ε 3.97); 308 (sh) (log ε 3.51) (MeOH).

10,11-Dihydro, N-Me: **N-Methylhydroquinicinalol** [99964-74-6]
C₂₁H₂₈N₂O₂ 340.464
Alkaloid from the trunk bark of *Guettarda trimera* (Rubiaceae). Amorph. [α]_D²⁰ -3 (c, 1 in CHCl₃). λ_{max} 222 ; 245 ; 262 ; 300 ; 356 (EtOH).

10,11-Dihydro, 9R-alcohol, N-Me: **9-Epi-N-methylhydroquinicinalol** [99964-75-7]
C₂₁H₃₀N₂O₂ 342.48
Alkaloid from the trunk bark of

Guettarda trimera (Rubiaceae). Cryst. (Me₂CO). Mp 164°. [α]_D²⁰ +58 (c, 0.3 in CHCl₃).

10,11-Dihydro, 9S-alcohol, N-Me: **N-Methylhydroquinicinalol** [99964-76-8]
C₂₁H₃₀N₂O₂ 342.48
Alkaloid from the trunk bark of *Guettarda trimera* (Rubiaceae). Cryst. (Me₂CO). Mp 156°. [α]_D²⁰ -84 (c, 1.2 in CHCl₃). λ_{max} 233 ; 271 ; 281 ; 291 ; 323 ; 333 (EtOH).

Demethoxy: **Cinchonine. Cinchotoxine** [69-24-9]
C₁₉H₂₂N₂O 294.396
Poss. minor congener of Cinchonine, C-447, occurrence not well documented. Readily obt. by thermal rearr. of cinchonine or cinchonidine. Long thin prisms (Et₂O). Mp 58-59°. [α]_D +46.5 (c, 2 in CHCl₃). [α]_D +57.7 (EtOH).

Demethoxy; nitrate: Mp 160°.
Demethoxy, 9R-alcohol: **Epicinchoninol** [120949-74-8]
C₁₉H₂₄N₂O 296.411
Alkaloid from the bark of *Ladenbergia oblongifolia* and cortex of *Cinchona succirubra*. Brownish-yellow amorph. powder. [α]_D²⁵ -29 (c, 0.1 in EtOH).

Demethoxy, 9R-alcohol, N-Ac: **N-Acetylcinchoninol**
C₂₁H₂₆N₂O₂ 338.449
Alkaloid from the leaves of *Remijia peruviana*. Amorph. [α]_D²⁵ +111.5 (c, 0.1 in EtOH). λ_{max} 206 (log ε 4.68); 225 (log ε 4.5); 284 (log ε 3.8); 300 (log ε 3.78); 314 (log ε 3.7) (EtOH).

Demethoxy, 9S-alcohol: **Cinchoninol. Cinchonidinol. 1,8-Secocinchonine** [27875-43-0]
C₁₉H₂₄N₂O 296.411
Alkaloid from *Cinchona succirubra* and the bark of *Ladenbergia oblongifolia*. Monoamine oxidase inhibitor. Pale yellow amorph. powder. [α]_D²⁵ -15 (c, 0.1 in EtOH). [α]_D²¹ +5.6 (c, 0.09 in EtOH). λ_{max} 224 (log ε 4.1); 281 (log ε 2.92); 300 (log ε 2.79); 314 (log ε 2.72) (EtOH).


Demethoxy, 10,11-dihydro, 9-alcohol: **Di-hydrocinchoninol** [27875-44-1]
C₁₉H₂₆N₂O 298.427
Alkaloid from the bark of *Ladenbergia oblongifolia*. Mixt. of C-9 epimers.

Demethoxy, deoxo, N-Ac: **N-Acetyldeoxycinchoninol**
C₂₁H₂₆N₂O 322.449
Alkaloid from the leaves of *Remijia peruviana*. Amorph. [α]_D²⁵ +84 (c, 0.79 in EtOH). λ_{max} 206 (log ε 4.8); 224 (log ε 4.8); 282 (log ε 3.96); 300 (log ε 3.82); 316 (log ε 3.7) (EtOH).

[52211-63-9]
Howard, D. et al., *J.C.S.*, 1871, **24**, 61-64; 1872, **25**, 101-106 (isol, synth)
Proštenik, M. et al., *Helv. Chim. Acta*, 1943, **26**, 1965-1971 (synth)
Quevauviller, A. et al., *Ann. Pharm. Fr.*, 1966, **24**, 39-50 (rev. pharmacol, synth)
Guerey, C. et al., *Arzneim.-Forsch.*, 1972, **22**, 1336-1340; 1341-1346 (metab)

Grethe, G. et al., *Helv. Chim. Acta*, 1973, **56**, 1485-1494 (synth)
Smith, G. et al., *J. Chromatogr.*, 1984, **299**, 233-244 (hplc)
Kan-Fan, C. et al., *Phytochemistry*, 1985, **24**, 2773-2775 (*N-Methylhydroquinicinalol, N-Methylhydroquinicinalol, 9-Epidihydroquinicinalol*)
Mitsui, N. et al., *Chem. Pharm. Bull.*, 1989, **37**, 363-366 (*Cinchoninol, Cinchonine*)
Montagnac, A. et al., *Phytochemistry*, 1997, **46**, 973-975 (*N-Methylquinicinalol*)
Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 16442
Okunade, A.L. et al., *Fitoterapia*, 2001, **72**, 717-719 (*Cinchoninol, Epicinchoninol, Dihydrocinchoninols*)
Diaz, J.G. et al., *J. Nat. Prod.*, 2004, **67**, 1667-1671 (*N-Acetyldeoxycinchoninol, N-Acetylcinchoninol*)

Quinidine, BAN Q-18
6'-Methoxycinchonan-9-ol, 9CF. Conquinine. Conchinine. β-Quinine. Pitayine. Many other names [56-54-2]



C₂₀H₂₄N₂O₂ 324.422
Alkaloid from *Cinchona officinalis* and very many other *Cinchona* spp., and from *Remijia pedunculata* (Rubiaceae). Potassium channel blocker. Antiarrhythmic agent, antimalarial. Cardiac depressant, has been used as clinical antifibrillatory agent. Mp 174-175° (anhyd.). [α]_D¹⁵ +262 (EtOH). pK_{a1} 8.54; pK_{a2} 4. Log P 2.93 (calc). λ_{max} 279 (ε 3990); 332 (ε 5280) (MeOH).

▶ Adverse human effects incl. gastrointestinal and cardiac. Eye irritant. LD₅₀ (rat, orl) 263 mg/kg. VA4725000
Hydrochloride: Mp 258-259° dec. (anhyd.). [α]_D²⁰ +200 (H₂O).
Sulfate (2:1): **Quinidine sulfate, USAN. CinQuin. Quinidex** [50-54-4]
▶ Adverse human systemic effects. LD₅₀ (rat, orl) 456 mg/kg. VA5951000
Picrate: Mp 243°.

D-Gluconate salt: **Quinidine gluconate, USAN. Duraquin. Quinaglute** [7054-25-3]

▶ Harmful in contact with skin (allergic dermatitis). LZ5250000
Ketone: 6'-Methoxycinchonan-9-one, 9CI. **Quinidinone** [14528-53-1]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from *Cinchona* sp. Cryst. (Et₂O). Sol. CHCl₃, C₆H₆, Et₂O; poorly sol. H₂O, hexane. Mp 99-100.5°. [α]_D +73.8 (c, 1 in EtOH).
10,11-Dihydro: **Hydroquinidine. Hydroconquinine. Hydroconchinene. Hydroquinine. Lentoquinine. Quinotidine. Serecor** [1435-55-8]
C₂₀H₂₆N₂O₂ 326.438

Minor congener of quinidine from *Cinchona* bark, also obt. from *Remijia pedunculata* and *Aspidosperma marcgravianum* bark and leaves (Rubiaceae, Apocynaceae). Antimalarial. Antiarhythmic. Needles (EtOH). Mp 168-169°. $[\alpha]_D^{18} +237$ (EtOH). Log P 3.42 (calc).

▶ LD₅₀ (rat, orl) 369 mg/kg. MX3016000
10,11-Dihydro; hydrochloride: [1476-98-8]
Cryst. (H₂O). Mp 273-274°. $[\alpha]_D^{26} +183.9$.

▶ MX3018000

10,11-Dihydro, O-(4-chlorobenzoyl):

DHQD-CLB

[113162-02-0]

C₂₇H₂₉ClN₂O₃ 464.99

Chiral ligand for asymmetric dihydroxylation of alkenes. Pale yellow solid. Mp 102-105°. $[\alpha]_D -73$ (c, 1 in EtOH).

9-Epimer: **Epiquinidine**

[572-59-8]

Occurs in *Cinchona* spp., poss. as artifact. Cryst. (EtOAc). Mp 113°. $[\alpha]_D^{20} +103.7$ (EtOH).

9-Epimer; hydrochloride (1:2):

Cryst. (EtOH). Mp 195.7° dec. $[\alpha]_D^{20} +45.5$ (EtOH).

[6591-63-5]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 865D; 866B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 441A; 441B; 442A (nmr)

Van Hejningen, et al., *Annalen*, 1849, **72**, 301-305 (isol)

Dirscherl, W. et al., *Annalen*, 1935, **521**, 48-72 (Epiquinidine, isol)

Lyle, G.G. et al., *Tetrahedron*, 1967, **23**, 51-63; 3253-3263 (uv, ord, config)

Battersby, A.R. et al., *Chem. Comm.*, 1971, 30-31; 31-32 (biosynth)

Gutzwiller, J. et al., *Helv. Chim. Acta*, 1973, **56**, 1494-1503 (Quinidinone, synth)

Moreland, C.G. et al., *J.O.C.*, 1974, **39**, 2413-2416 (cmr)

Grethe, G. et al., *J.A.C.S.*, 1978, **100**, 589-593 (synth)

Soeterboek, A.M. et al., *Int. Congr. Ser. Excerpta Med.*, 1980, **501**, 180-192 (rev)

Kashino, S. et al., *Acta Cryst. C*, 1983, **39**, 310-312 (cryst struct)

Loutfy, M.A. et al., *Anal. Profiles Drug Subst.*, 1983, **12**, 483-546 (rev, uv, ir, pmr, cmr, ms, anal)

Roden, D.M. et al., *Pharmacol. Ther.*, 1983, **23**, 179-191 (rev)

Smith, E. et al., *J. Chromatogr.*, 1984, **299**, 233-244 (hplc)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 5748

Robins, R.J. et al., *Phytochemistry*, 1987, **26**, 551-556 (pmr, cmr)

Prakash, O. et al., *Indian J. Chem., Sect. B*, 1988, **27**, 950-952 (pmr)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **3**, 1887-1888 (dihydro chlorobenzoyl, use)

Silva, T.H.A. et al., *Bioorg. Med. Chem.*, 1997, **5**, 353-361 (conform)

Reichardt, C. et al., *Liebigs Ann./Recl.*, 1997, 707-720 (dihydro chlorobenzoyl, synth)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 938

Raheem, I.T. et al., *J.A.C.S.*, 2004, **126**, 706-707 (synth)



Igarashi, J. et al., *Tet. Lett.*, 2004, **45**, 3783-3786 (synth)

Guo, M.-J. et al., *Acta Cryst. E*, 2007, **63**, o1483-o1484 (cryst struct)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, HIG500; 1992, QFS000; QHA000; QDS225

Quinindoline, 9CI

Indolo[2,3-b]quinoline. 6-Azabenzob[6]-carbazole
[243-38-9]



6H-form

C₁₅H₁₀N₂ 218.257

Free compd. is in the 6H-form. Constit. of anthracite pitch. Degradn. prod. of β-Isoidingo. Yellow needles or leaflets (PhNO₂). Sol. hot PhNO₂, hot PhNH₂; insol. most solvs. Mp 346°.

Hydrochloride:

Yellow needles. Mp 280° dec.

5H-form

N-Me: 5-Methyl-5H-indolo[2,3-b]quinoline. *Cryptotackieine*. **Neocryptolepine**
[114414-78-7]

C₁₆H₁₂N₂ 232.284

Alkaloid from root bark of *Cryptolepis sanguinolenta*. Shows antibacterial and antifungal activities. Yellowish amorph. powder or cryst. (hexane). Mp 107-109°.

6H-form

Ac:

C₁₇H₁₂N₂O 260.295

Needles. Mp 189° (185°).

Gabriel, S. et al., *Ber.*, 1897, **30**, 3017-3022

Lawson, G. et al., *J.C.S.*, 1924, **125**, 626-657

Pailer, M. et al., *Monatsh. Chem.*, 1965, **96**, 1695 (synth)

Sagitullin, R.S. et al., *CA*, 1974, **81**, 13408m (synth)

Kaczmarek, L. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1988, **321**, 463-467

Marais, F. et al., *J. Het. Chem.*, 1989, **26**, 1589 (synth, ir, pmr)

Reczynska-Czoch, W. et al., *J. Med. Chem.*, 1994, **37**, 3503 (synth)

Sharaf, E.H.M. et al., *J. Het. Chem.*, 1996, **33**, 239 (Cryptotackieine)

Cimanga, K. et al., *Tet. Lett.*, 1996, **37**, 1703 (Neocryptolepine, isol, uv, pmr, cmr, ms, struct)

Alajarin, M. et al., *J. Nat. Prod.*, 1997, **60**, 747 (synth, cmr)

Timari, G. et al., *Synlett*, 1997, 1067-1068 (Neocryptolepine, synth)

Cimanga, K. et al., *Phytomedicine*, 1998, **5**, 209-214 (Neocryptolepine, activity)

Molina, P. et al., *Synthesis*, 1999, 326-329 (Neocryptolepine, synth, pmr, cmr, ms)

Fresneda, P.M. et al., *Tetrahedron*, 2001, **57**, 6197-6202 (Neocryptolepine, synth)

Ho, T.L. et al., *Helv. Chim. Acta*, 2002, **85**, 3823-3827 (Neocryptolepine, synth, ir, pmr, cmr)

Sundaram, G.S.M. et al., *J.O.C.*, 2004, **69**, 5760-5762 (Neocryptolepine, synth)

Engqvist, R. et al., *Synth. Commun.*, 2004, **34**, 386-390 (synth, pmr, Neocryptolepine)


Dhanabal, T. et al., *Tetrahedron*, 2006, **62**, 6258-6263 (Neocryptolepine, synth)

Parvatkar, P.T. et al., *Tet. Lett.*, 2007, **48**, 7870-7872 (Neocryptolepine, synth)

Quinine, BAN

6'-Methoxycinchonan-9-ol, 9CI†
[130-95-0]

Q-20



Absolute Configuration

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Cinchona officinalis* and most other *Cinchona* spp., also in *Remijia pedunculata*. Occurs mainly in the bark (cinchona bark) but has also been isol. from trunkwood (Rubiaceae). Used as 1% aq. soln. (as sulfate) for photometric detn. of Bi, W, As(I), P(I). Resolving agent for organic acids, used in asymmetric syntheses. Used in tonics and bitter drinks. Potassium channel blocker. Traditional antimalarial drug esp. important in treating *Plasmodium falciparum* which is resistant to other antimalarial drugs. Activity is stereochemistry-independent, i.e. racemates and stereoisomers show similar activity. Illicit abortifacient in large doses. Stimulant for horses, has been used in horse doping. Weak cardiac depressant, antipyretic and antifibrillatory agent. Cryst. with v. bitter taste. Sol. H₂O, EtOH, C₆H₆, CHCl₃. Mp 177° (anhyd.). $[\alpha]_D^{25} -158.7$ (Et₂O). $[\alpha]_D -158$ (-145) (EtOH). pK_{a1} 8.34; pK_{a2} 4.21 (20°). Log P 2.93 (calc). Forms a dihydrate and a trihydrate, Mp 57° (efflorescent). Triboluminescent. Blue fluor. in soln. λ_{max} 279 (ε 3870); 332 (ε 514) (MeOH).

▶ Adverse effects from therapeutic or larger doses of quinine or its salts (cinchonism). Irritant to mucous membranes. Human and exp. teratogenic effects. LD₅₀ (mus, ipr) 115 mg/kg. VA6020000

Hydrochloride (1:1): Quinine hydrochloride, JAN. FEMA 2976

[130-89-2]

[7549-43-1]

Needles + 2H₂O, dehydrating at 100°. Mp 158-160° (anhyd.). $[\alpha]_D^{15} -145.5$ (EtOH).

▶ VA7700000

Hydrochloride (1:2): [60-93-5]

[7549-43-1]

Needles. Mp 180-185°. $[\alpha]_D^{18} -233$ (H₂O).

▶ LD₅₀ (rat, orl) 1393 mg/kg. VA6478000

Hydrobromide: [549-49-5]

Hygroscopic needles + H₂O. Mp 200° (softens from 152°).

▶ VA7650000

Hydrobromide (1:2):

Yellow prisms + 3H₂O. Mp 81-82°.


Sulfate salt (1:1): Quinine bisulfate.

- FEMA 2975**
[549-56-4] Flavouring agent with bitter taste.
- Sulfate salt (2:1): Quinine sulfate, JAN, USAN. Quinanium. Quine. Quinite. FEMA 2977**
[804-63-7]
[549-56-4, 6119-70-6] Flavouring agent with bitter taste. Mp 205°. $[\alpha]_D^{15}$ -235 (H₂SO₄ aq.). Forms a dihydrate and an octahydrate. Component of Quinamin.
- ▶ **VA8440000**
Gluconate salt (1:1): [4325-25-1]
Mp 159-161° (anhyd.).
- Compd. with ascorbic acid (1:2): Quinine ascorbate, USAN. Quinine biascorbate**
[146-40-7] Smoking deterrent.
- Ethylcarbonate: Quinine ethylcarbonate, JAN. Euquinine. Tasteless Quinine**
[83-75-0]
C₂₃H₂₈N₂O₄ 396.485
Antimalarial. Fine needles. Mp 90-92°. $[\alpha]_D^{20}$ -43.7. Log P 4.22 (uncertain value) (calc).
- Methiodide (1:2):**
Yellow plates + 3H₂O. Mp 167-168° dec. $[\alpha]_D^{18}$ -151 (2M H₂SO₄).
- N-Benzyl:** [67174-25-8]
C₂₇H₃₁N₂O₂[⊕] 415.554
Chiral phase-transfer catalyst. Mp 200-205° dec. (as chloride). $[\alpha]_D^{20}$ -235 (c, 1.5 in H₂O). CAS no. refers to chloride.
- Ac:** [18797-86-9]
Cryst. (petrol). Mp 116-117°. $[\alpha]_D^{15}$ -54.3 (EtOH).
- Benzoyl:** [69758-70-9]
Prisms (Et₂O). Mp 139°. $[\alpha]_D^{17}$ +121.6 (EtOH). Log P 5.48 (uncertain value) (calc).
- O-(2-Hydroxybenzoyl): Saloquinine. Salochimin**
C₂₇H₂₈N₂O₄ 444.529
Analgesic, antipyretic. Cryst. (EtOH). Mp 140°. Log P 5.81 (uncertain value) (calc).
- N¹-Et: N¹-Ethylquinine**
C₂₂H₂₉N₂O₂[⊕] 353.483
Alkaloid from the bark of *Remijia peruviana*. Amorph. $[\alpha]_D^{25}$ +128 (c, 0.12 in EtOH). Counterion not specified. λ_{\max} 230 (log ϵ 3.02); 279 (sh) (log ϵ 2.14); 333 (sh) (log ϵ 2.17) (EtOH).
- O-De-Me: Cupreine. Ultraquinine**
[524-63-0]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from *Remijia pedunculata*, poss. also present in *Picrolemma pseudocoffea* (Rubiaceae, Simaroubaceae). Mp 198°. $[\alpha]_D$ -175 (EtOH).
- O-De-Me, sulfate salt:** Mp 257°.
- O-De-Me, 6'-Ac: 6'-O-Acetylcupreine**
C₂₁H₂₄N₂O₃ 352.432
Alkaloid from the bark of *Remijia peruviana*. $[\alpha]_D^{25}$ -116.6 (c, 0.32 in EtOH). λ_{\max} 232 (log ϵ 3.25); 286 (sh) (log ϵ 2.41); 336 (sh) (log ϵ 2.48) (EtOH).
- O-De-Me, di-Ac:** Mp 88°.
- O-De-Me, O⁶-Et: Quinethyline**
[83348-45-2]
C₂₁H₂₆N₂O₂ 338.449
Febrifuge. Mp 160°. $[\alpha]_D$ -169.4 (EtOH).
- 10,11-Dihydro: Hydroquinine**
[522-66-7]
C₂₀H₂₆N₂O₂ 326.438
Minor congener of Quinine found in *Cinchona* spp., also from *Remijia pedunculata* (Rubiaceae). Antimalarial. Needles (Et₂O or CHCl₃). Mp 172° (169°). $[\alpha]_D$ -142.2 (EtOH).
- 10,11-Dihydro; hydrochloride:** Mp 235-240° (anhyd.).
- 10,11-Dihydro, Me ether:** Mp 93°.
- 10,11-Dihydro, O-de-Me: Dihydrocupreine. Hydrocupreine**
[5962-19-6]
C₁₉H₂₄N₂O₂ 312.411
Alkaloid from the bark of *Timonius kaniensis* (Rubiaceae). Glass or foam. $[\alpha]_D$ -143 (c, 0.7 in EtOH).
- 10,11-Dihydro, O-de-Me, picrate:** Mp 250-252°.
- 10,11-Dihydro, O-(4-chlorobenzoyl):**
[113216-88-9]
C₂₇H₂₉ClN₂O₃ 464.99
Chiral ligand used in asymmetric dihydroxylation of alkenes. Pale yellow solid. Mp 130-133°. $[\alpha]_D$ +150 (c, 1 in EtOH).
- 10,11-Dihydro, O-de-Me, O⁶-Et: Ethyldihydrocupreine. Neumolisina. Neumolysin. Numoquin. Optochin. Optoquine**
[522-60-1]
C₂₁H₂₈N₂O₂ 340.464
Antibacterial (ophthalmologic) agent. Bitter cryst. Mp 123-128°. $[\alpha]_D^{25}$ -136.2 (EtOH). Log P 3.94 (calc).
- ▶ Eye irritant (rbt). LD_{Lo} (mus, orl) 400 mg/kg. MW5950000
- 10,11-Dihydro, O-de-Me, O⁶-Et, hydrochloride:** [3413-58-9]
Rhombic cryst. (Me₂CO/Et₂O). Mp 252-254°. $[\alpha]_D^{25}$ -123.6 (H₂O).
- ▶ MW6300000
- 3S-Hydroxy: 3-Hydroxyquinine**
C₂₀H₂₄N₂O₃ 340.421
Major metab. of quinine in humans. Mp 148-150°. $[\alpha]_D^{25}$ -196.8 (c, 0.5 in MeOH).
- 9-Epimer: Epiquinine**
[572-60-1]
C₂₀H₂₄N₂O₂ 324.422
Minor constit. of *Cinchona* spp. (Rubiaceae). Oil. $[\alpha]_D^{20}$ +43.3 (EtOH).
- 9-Epimer; hydrochloride (1:2):**
Cryst. (Me₂CO). Mp 196° dec. $[\alpha]_D^{21}$ +33.3 (EtOH).
- [6119-47-7]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 864D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 440B; 441C (nmr)
Paul, et al., *Pharm. J.*, 1881, 12, 497 (Cupreine, isol)
U.S. Pat., 1913, 1 062 203 (Ethyldihydrocupreine)
Giemsa, G. et al., *Ber.*, 1918, 51, 1325-1333 (Cupreine Me ether, Ethyldihydrocupreine)
Woodward, R.B. et al., *J.A.C.S.*, 1945, 67, 860-874 (synth)
- Muir, K. et al., *CA*, 1947, 41, 4582 (Ethyldihydrocupreine, synth, pharmacol)
Welcher, F.J. et al., *Organic Analytical Reagents*, Van Nostrand, NY, 1948, 4, 251 (use)
Altman, R.F.A. et al., *CA*, 1958, 52, 506a (Cupreine, occur)
Willemot, J. et al., *Ann. Pharm. Fr.*, 1965, 23, 203-209 (gluconate)
Lyle, G.G. et al., *Tetrahedron*, 1967, 23, 51-63; 3253-3263 (config, uv, ord, pmr)
Johns, S.R. et al., *Aust. J. Chem.*, 1970, 23, 211-212 (Dihydrocupreine)
Fales, H.M. et al., *J.A.C.S.*, 1970, 92, 1590-1597 (ms)
Battersby, A.R. et al., *Chem. Comm.*, 1971, 30-31; 31-32 (biosynth)
Chao, J.M. et al., *Anal. Chem.*, 1974, 46, 296-298 (ms)
Moreland, C.G. et al., *J.O.C.*, 1974, 39, 2413-2416 (cmr)
Grethe, G. et al., *J.A.C.S.*, 1978, 100, 589-593 (synth)
Hahn, F.E. et al., *Antibiotics (N.Y.)*, 1979, 5, 353-362 (rev, pharmacol, tox)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1980, 8, 430 (use)
Muhtadi, F.J. et al., *Anal. Profiles Drug Subst.*, 1983, 12, 547-621 (rev, uv, ir, pmr, cmr, ms, anal)
Imanishi, T. et al., *Chem. Pharm. Bull.*, 1983, 31, 1551-1560 (synth)
Hofheinz, W. et al., *Handb. Exp. Pharmacol.*, 1984, 68, 61-81 (rev)
Smith, E. et al., *J. Chromatogr.*, 1984, 299, 233-244 (hplc)
Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 5747 (synonyms)
Prakash, O. et al., *Indian J. Chem., Sect. B*, 1988, 27, 950-952 (pmr)
Christie, D.J. et al., *J. Lab. Clin. Med.*, 1988, 112, 92-98 (Cupreine, activity)
Diaz-Araujo, H. et al., *J. Nat. Prod.*, 1990, 53, 112-124 (synth)
Lander, V. et al., *Z. Lebensm.-Unters.-Forsch.*, 1990, 190, 410-413 (hplc)
Karle, J.M. et al., *Acta Cryst. C*, 1992, 48, 1975-1980 (Epiquinine, cryst struct)
Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 3, 1887-1888 (dihydro chlorobenzoyl, use)
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2407 (rev)
Reichardt, C. et al., *Liebigs Ann./Recl.*, 1997, 707-720 (dihydro chlorobenzoyl, synth)
Montagnac, A. et al., *Phytochemistry*, 1997, 46, 973-975 (Cupreine, isol, pmr, cmr)
Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 439
Fenaroli's Handbook of Flavor Ingredients, 4th edn., (ed. Burdock, G.A.), CRC Press, 2001, 1574-1577 (use, salts)
Stork, G. et al., *J.A.C.S.*, 2001, 123, 3239-3242 (synth, bibl)
Raheem, I.T. et al., *J.A.C.S.*, 2004, 126, 706-707 (synth)
Igarashi, J. et al., *Tet. Lett.*, 2004, 45, 3783-3786 (synth)
Kaufman, T.S. et al., *Angew. Chem., Int. Ed.*, 2005, 44, 854-885 (rev)
Ruiz-Mesia, L. et al., *J. Agric. Food Chem.*, 2005, 53, 1921-1926 (N¹-Ethylquinine, O-Acetylcupreine)
Sarma, P.V.V.S. et al., *J. Nat. Prod.*, 2005, 68, 942-944 (3-Hydroxyquinine)
Seeman, J.I. et al., *Angew. Chem., Int. Ed.*, 2007, 46, 1378-1413 (rev, synth)
Webber, P. et al., *J.O.C.*, 2008, 73, 9379-9387 (synth)
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley,

2000, QHJ000; 1992, QIJ000; QJS000;
QMA000; QJJ100; ELC500; HHR700

Quinocarcin

DC 52. Antibiotic DC 52
[84573-33-1]



Absolute
Configuration

$C_{18}H_{22}N_2O_4$ 330.383

Isol. from *Streptomyces melanovinaceus*. Active against gram-positive bacteria and tumours. Powder. Sol. H_2O , butanol, EtOH, MeOH; fairly sol. EtOAc, Me_2CO ; poorly sol. $CHCl_3$, Et_2O , hexane. $[\alpha]_D^{22}$ -32 (c, 0.5 in H_2O). Browns at 170° , indefinite Mp. Fairly unstable, forms a more stable HCN adduct. λ_{max} 271 ; 277 (MeOH) (Derep).

► LD₅₀ (mus, ipr) 27 mg/kg. Mutagenic. NJ8620820

[159571-56-9 , 101311-75-5]

Tomita, F. *et al.*, *J. Antibiot.*, 1983, **36**, 463; 468; 1984, **37**, 1268 (isol, uv, pmr, cmr, struct) *Eur. Pat.*, 1985, ((*Kyowa Hakko Kogyo*))157 126; *CA*, **104**, 155970f (synth, pharmacol) Danishefsky, S.J. *et al.*, *J.A.C.S.*, 1985, **107**, 1421

Fujimoto, K. *et al.*, *Cancer Res.*, 1987, **47**, 1516

Saito, H. *et al.*, *Tet. Lett.*, 1987, **28**, 4065 (synth)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1988, **110**, 5196 (synth, bibl)

Garner, P. *et al.*, *J.A.C.S.*, 1993, **115**, 10742-10753 (synth, abs config)

Katoh, T. *et al.*, *Tetrahedron*, 1994, **50**, 6193; 6209; 6221; 6239 (synth, bibl, ir, pmr)

Plowman, J. *et al.*, *Cancer Res.*, 1995, **55**, 862
Katoh, T. *et al.*, *Stud. Nat. Prod. Chem.*, 1997, **19**, 289-350 (rev, synth)


Kwon, S. *et al.*, *J.A.C.S.*, 2005, **127**, 16796-16797 (synth)

Wu, Y.-C. *et al.*, *J.A.C.S.*, 2008, **130**, 7148-7152 (synth)

Allan, K.M. *et al.*, *J.A.C.S.*, 2008, **130**, 17270-17271 (synth)

Quinocarcinol

DC 52d. Antibiotic DC 52d
[84573-32-0]



Absolute
Configuration

$C_{18}H_{24}N_2O_4$ 332.399

See also Quinocarcin, Q-21. Isol. from *Streptomyces melanovinaceus* NRRL12388. Active against gram-positive bacteria and shows marginal antitumour activity. Powder. Sol. H_2O , butanol, MeOH; fairly sol. Me_2CO , EtOAc; poorly sol. $CHCl_3$, hexane. $[\alpha]_D^{22}$ -8 (c, 0.5 in H_2O). Browns at 188° , dec. at

Q-21

230° . λ_{max} 271 ; 277 (MeOH) (Derep).


Tomita, F. *et al.*, *J. Antibiot.*, 1983, **36**, 463; 468 (isol, uv, pmr, cmr, struct)

Hirayama, N. *et al.*, *J.C.S. Perkin 2*, 1983, 1705 (cryst struct)

Danishefsky, S.J. *et al.*, *J.A.C.S.*, 1985, **107**, 1421 (synth)

Quinocitrinine A

[630119-02-7]



Relative
Configuration

$C_{16}H_{19}N_2O_2^{\oplus}$ 271.338

Prod. by *Penicillium citrinum* VKM FW-800. Sl. brownish solid. Mp $142-143^\circ$. $[\alpha]_D^{22}$ +25.3 (c, 0.32 in MeOH). Counterion not specified. λ_{max} 216 (€ 8108); 248 (€ 5285); 256 (€ 5430); 300 (€ 2388); 314 (€ 2785); 328 (€ 2499) (MeOH).

Diastereoisomer: Quinocitrinine B

[630119-04-9]


$C_{16}H_{19}N_2O_2^{\oplus}$ 271.338

Prod. by *Penicillium citrinum* VKM FW-800. Sl. brownish solid. Mp $152-153^\circ$. $[\alpha]_D^{22}$ -13.2 (c, 0.62 in MeOH). Counterion not specified. λ_{max} 216 (€ 8108); 248 (€ 5772); 256 (€ 5831); 300 (€ 2542); 314 (€ 3074); 328 (€ 1973) (MeOH).

Kozlovsky, A.G. *et al.*, *J. Antibiot.*, 2003, **56**, 488-491 (isol, pmr, cmr)

Quinolactacide

10H-Pyrrolizino[1,2-b]quinoline-9,10(4H)-dione, 9CI
[856216-29-0]




$C_{14}H_8N_2O_2$ 236.229

Isol. from *Penicillium citrinum* F 1539. Insecticide. Yellow cryst. ($CHCl_3$ /MeOH). Mp $373-374^\circ$ dec. λ_{max} 241 (€ 12300); 288 (€ 32200) ($CHCl_3$ /MeOH).

Abe, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2005, **69**, 1202-1205; 2006, **70**, 303-306 (isol, synth, pmr, cmr, ms)

Quinolactacin A₁

[386211-68-3]



Absolute
Configuration

$C_{16}H_{18}N_2O_2$ 270.33

Q-23

Stereochem. revised in 2006. It has been demonstrated that only Quinolactacins A₂ and B₂ are genuine nat. prods.; the remainder are dec. artifacts. Prod. by *Penicillium citrinum* sp. 90648. Acetylcholinesterase inhibitor. Powder. $[\alpha]_D^{25}$ +30.3 (c, 0.16 in DMSO). λ_{max} 215 (€ 31300); 248 (€ 20150); 256 (€ 20250); 315 (€ 11150); 327 (€ 10150) (MeOH).

3-Hydroxy: Quinolactacin C₁

[888728-74-3]

$C_{16}H_{18}N_2O_3$ 286.33

Prod. by *Penicillium citrinum* (MST-F10130).

3-Epimer: Quinolactacin A₂. Quinolactacin A (obsol.)

[386211-69-4]

$C_{16}H_{18}N_2O_2$ 270.33

Prod. by *Penicillium citrinum* sp. 90648 and a *Penicillium* sp. EPF-6. Acetylcholinesterase inhibitor; shows antitumour props. Powder. Mp $262-265^\circ$. $[\alpha]_D^{25}$ +17.9 (c, 0.13 in DMSO).

3-Epimer, 3-hydroxy: Quinolactacin C₂

[888728-73-2]

$C_{16}H_{18}N_2O_3$ 286.33

Prod. by *Penicillium citrinum* and *Penicillium* sp. EPF-6. Powder. Mp $180-185^\circ$ dec. $[\alpha]_D^{25}$ +5.9 (c, 0.19 in DMSO). λ_{max} 215 (€ 31050); 250 (€ 18900); 258 (€ 17850); 315 (€ 12200); 328 (€ 11750) (MeOH).

Kakinuma, N. *et al.*, *J. Antibiot.*, 2000, **53**, 1247-1251; 1252-1256 (isol, uv, pmr, cmr)

Kim, W.G. *et al.*, *J. Antibiot.*, 2001, **54**, 831-835 (isol, uv, cd, ir, pmr, cmr)

Zhang, X. *et al.*, *J.O.C.*, 2003, **68**, 4523-4526 (synth)


Park, S.-J. *et al.*, *Tet. Lett.*, 2004, **45**, 8793-8795 (Quinolactacin A₂, synth)

Sasaki, T. *et al.*, *J. Antibiot.*, 2006, **59**, 418-427 (biosynth)

Clark, B. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1512-1519 (isol, uv, abs config)

Quinolactacin B

Q-26



(R)-form

$C_{15}H_{16}N_2O_2$ 256.304

Abs. configs. not clear. Quinolactacin B has only one chiral centre. It appears that the R- config. refers to the laevorotatory enantiomer (known as Quinolactacin B1) but Shankaraiah *et al* give the opposite assignment. Opt.rotns. are small however.

(R)-form**Quinolactacin B₁**

[549500-83-6]


Prod. by *Penicillium citrinum* (MST-F10130) and *Penicillium* sp. EPF-6. Powder. Mp $260-263^\circ$ dec. $[\alpha]_D^{25}$ -3.3 (c, 0.15 in DMSO). λ_{max} 214 (€ 24800); 248 (€ 15650); 255 (€ 15600); 314 (€ 8500); 327 (€ 7850) (MeOH).

(S)-form**Quinolactacin B₂**

[319917-26-5]

Prod. by *Penicillium citrinum* (MST-F10130).Kakinuma, N. *et al.*, *J. Antibiot.*, 2000, **53**, 1247-1251; 1252-1256 (*isol, struct*)Tatsuta, K. *et al.*, *J. Antibiot.*, 2001, **54**, 109-112 (*synth*)Zhang, X. *et al.*, *J.O.C.*, 2003, **68**, 4523-4526 (*synth*)Clark, B. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1512-1519 (*isol, uv, abs config*)Shankaraiah, N. *et al.*, *Tet. Lett.*, 2008, **49**, 4289-4291 (*synth*)**Quinolactacin D₁**


[888728-69-6]



Absolute Configuration

C₁₆H₁₈N₂O₃ 286.33Prod. by *Penicillium citrinum* (MST-F10130). *Isol.* as a mixt. with D₂. λ_{max} 216 (ε 23100); 250 (ε 12600); 259 (ε 12100); 315 (ε 8200) (MeOH) (mixt. of D₁/D₂).**3-Epimer: Quinolactacin D₂**

[888728-68-5]

C₁₆H₁₈N₂O₃ 286.33Prod. by *Penicillium citrinum* (MST-F10130). *Isol.* as a mixt. with D₁.Clark, B. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1512-1519 (*isol, pmr, cmr, ms*)**2-Quinolinamine, 9CI****2-Aminoquinoline, 8CI. α-Aminoquinoline**
[580-22-3]C₉H₈N₂ 144.176Alkaloid from the mushroom *Leucopaxillus aebissimus* var. *paradoxus* form *albiformis* (Agaricaceae). Possesses antibacterial, protease inhibitory, mutagenic, antitumour and anthelmintic activity. Cryst. (H₂O). Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 131.5-132.5° (129°). pK_a 7.3 (20°).

▶ VA9621800

Picrate: Mp 266-267°.**Methochloride:**Cryst. + 1H₂O. Mp 268°.**Methiodide:**Cryst. (H₂O). Mp 247°.**N-Ac:**C₁₁H₁₀N₂O 186.213
Mp 88-90°.**N-Ph: 2-Anilinoquinoline**

[5468-85-9]

C₁₅H₁₂N₂ 220.273

Cryst. (EtOH). Mp 98°.

N-Ph, 1-oxide:C₁₅H₁₂N₂O 236.273

Mp 211-212°.


N-Nitro: 2-Nitraminoquinoline. 2-QuinolylnitramineC₉H₇N₃O₂ 189.173

Yellow needles (AcOH). Mp 223-225°.

Turns red in air.

Tschitschibabin, A.E. *et al.*, *Ber.*, 1925, **58**, 803-807 (*synth*)Steck, E.A. *et al.*, *J.A.C.S.*, 1948, **70**, 3397 (*synth*)Dymek, W. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1964, **38**, 925 (2-Anilinoquinoline)Kametani, T. *et al.*, *J. Het. Chem.*, 1965, **2**, 330 (*synth*)Cook, M.J. *et al.*, *J.C.S. Perkin 2*, 1973, 1080 (*tautom*)Tondys, H. *et al.*, *J. Het. Chem.*, 1985, **22**, 353 (*synth*)Pfister, J.R. *et al.*, *J. Nat. Prod.*, 1988, **51**, 969 (*isol, cmr, pmr*)Yamada, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1991, **64**, 1821 (*uv*)Strekowski, L. *et al.*, *Synth. Commun.*, 1994, **24**, 2387 (*N-Ac*)Campagnone, R.S. *et al.*, *Synth. Commun.*, 1997, **27**, 1631-1641 (*synth, ir, pmr, cmr*)**Quinoline****Q-29****1-Benzazine. 1-Azanaphthalene. FEMA**
3470

[91-22-5]

C₉H₇N 129.161Occurs in cocoa, black tea, Scotch whisky, coal tar and in 'stupp' fat. A fairly major constit. of cigarette smoke. Alkaloid from various plant spp. incl. *Mentha* spp., *Peganum harmala* and *Rubus laciniata*. Solv. for decarboxylations. Dehydrohalogenation reagent. Used for photometric detn. of Mo (λ_{max} 470 nm, ε 18000), gravimetric detn. of P(V); as a synergistic agent in extraction of Co. Flavouring ingredient. Antimalarial agent. Liq. Mod. sol. H₂O. d₄²⁰ 1.09. Fp -15.6. Bp 238.05° Bp₁₇ 114°. n_D²⁰ 1.6268. pK_a 4.9 (20°). Hygroscopic, steam-volatile.▶ Fl. p. 99°, autoignition temp. 480°. Often violent synth. (Skraup reaction). Skin and severe eye irritant. Contact with liquid can damage eyes. LD₅₀ (rat, orl) 331 mg/kg. LD₅₀ (rbt, skn) 540 mg/kg. Reported to be mutagenic and carcinogenic. VA9275000**Hydrochloride:** [530-64-3]Deliquescent prisms + ½ H₂O. Mp 94°
Mp 134.5° (anhyd.).**Sulfate (2:1):** [54957-90-3]

Cryst. (EtOH or AcOH). Mp 163.5-164.5°.

Picrate: [14876-03-0]Yellow needles (C₆H₆). Mp 207-208°.**N-Oxide:** [1613-37-2]C₉H₇NO 145.16Needles + 2H₂O. Mp 62°.

▶ VC2335000

N-Oxide, picrate:

Cryst. Mp 143°.

Methochloride:C₁₀H₁₀CIN 179.648Cryst. + 1H₂O (EtOH). Mp 126°.**Methobromide:** [2516-72-5]C₁₀H₁₀BrN 224.1

Needles. Mp 70°.

Methiodide: [3947-76-0]C₁₀H₁₀IN 271.1

Orange-red cryst. (EtOH). Mp 133°.

▶ VC3955000

Ethobromide: [26670-42-8]C₁₁H₁₂BrN 238.127Plates + 1H₂O (H₂O or EtOH). Mp 80°.**Ethiodide:** [634-35-5]C₁₁H₁₂IN 285.127

Yellow prisms (EtOH). Mp 158°.

▶ VC3677000

N-Benzyl: [19801-99-1]C₁₆H₁₄N[⊕] 220.293Used as a 0.8% soln. in HNO₃ for gravimetric detn. of Ce, Th. Cryst. (HNO₃) (as nitrate). Mp 73-74° (nitrate).


[5436-28-2, 1701-63-9, 27926-80-3, 26323-01-3]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 853A (*ir*)*Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 421A (*nmr*)*Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 1565A (*ir*)Cole, W.G. *et al.*, *J.C.S. (B)*, 1968, 1284 (*ms*)
Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1969, **2**, 356 (*use*)Lienhard, G.E. *et al.*, *Biochemistry*, 1970, **9**, 3011 (*uv*)Markov, G.S. *et al.*, *Zh. Anal. Khim.*, 1970, **25**, 278; 1971, **26**, 1215 (*N-benzyl, synth, detn, Th, Ce*)Hideo, A. *et al.*, *Nippon Kagaku Zasshi*, 1971, **92**, 1156; *CA*, **76**, 104508g (*detn, Co*)Zharekeev, B.K. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 264-265; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 282-283 (*isol*)Rao, V.P. *et al.*, *Mikrochim. Acta*, 1975, 265 (*detn, Mo*)Johns, S.R. *et al.*, *Aust. J. Chem.*, 1976, **29**, 1617 (*cmr*)Hirano, K. *et al.*, *Cancer Res.*, 1976, **36**, 329-335 (*tox*)Jones, G. *et al.*, *Chem. Heterocycl. Compd.*, 1977, **32**, 1 (*rev*)*Kirk-Othmer Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **19**, 532 (*rev*)Attimonelli, M. *et al.*, *Org. Magn. Reson.*, 1979, **12**, 17 (*pmr*)Horino, H. *et al.*, *Tet. Lett.*, 1979, 2403 (*synth*)Boyd, D.R. *et al.*, *Tet. Lett.*, 1991, **32**, 2963-2966 (*bibl, tox, metab*)*Encyclopaedia of Reagents for Organic Synthesis*, (ed. Paquette, L.A.), Wiley, 1995, **6**, 4397-4399 (*use*)*Fenaroli's Handbook of Flavor Ingredients, 3rd edn.*, (ed. Burdock, G.A.), CRC Press, 1995, **II**, 714 (*occur, use*)Asseline, U. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 1949-1957 (*methiodide*)Kalinin, A.I. *et al.*, *CA*, **80**, 127816c (*detn, P*)Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards, 4th edn.*, Butterworths, 1990, 2923Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory, 5th edn.*, Royal Society of Chemistry, 1992, 1091

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, QMJ000

4-Quinolinecarboxaldehyde, 9CI Q-30

4-Formylquinoline. Cinchoninaldehyde [4363-93-3]



C₁₀H₇NO 157.171

Alkaloid prod. by *Archangium gephyra*. Orange needles, plates + 1H₂O. Sol. H₂O, EtOH, Et₂O, toluene. Mp 51-53° (monohydrate 84-84.5°). Bp₄ 122-123°.

Picrate:

Cryst. (EtOH). Mp 179° (sinters at 170°).

Oxime: [39977-74-7]

C₁₀H₈N₂O 172.186

Alkaloid prod. by *Archangium gephyra*. Needles (MeOH). Mp 181-182°.

Phenylhydrazone:

Yellow cryst. Mp 175-176°.

Semicarbazone: Mp 244-245° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 869A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 446C (*nmr*)

Clemon, G.R. *et al.*, *J.C.S.*, 1939, 1241-1244 (*synth*)

Schultz, O.E. *et al.*, *Annalen*, 1970, 740, 192-195 (*synth*)


Forbis, R.M. *et al.*, *J.A.C.S.*, 1973, 95, 5003-5013 (*synth*)

Rodriguez, J.G. *et al.*, *J. Het. Chem.*, 1988, 25, 819-821 (*synth*)

Böhlendorf, B. *et al.*, *Annalen*, 1996, 49-53 (*pmr, ms*)

2-Quinolinecarboxylic acid, 9CI Q-31

Quinaldic acid. Quinaldinic acid [93-10-7]



C₁₀H₇NO₂ 173.171

Used for gravimetric detn. of Cd, Cu, Zn, Te. Needles + 2H₂O (H₂O). Sol. hot H₂O, C₆H₆; mod. sol. cold H₂O. Mp 157° (anhyd.). pK_{a1} 1.45; pK_{a2} 4.91 (25°, 0.1M NaClO₄). Heat → quinoline.

►UZ9100000

N-Me, zwitterion: 2-Carboxy-1-methylisoquinolinium hydroxide inner salt, 9CI.

2-Carboxy-1-methylquinolinium betaine. **N-Methylquinaldic acid**

[27104-82-1]

C₁₁H₉NO₂ 187.198

Alkaloid from the firefly *Photuris versicolor*. Defensive agent. Mp 110° dec. (natural) Mp 166-171° dec. (synthetic). λ_{max} 235 (log ε 4.35); 320 (sh) (log ε 3.97); 326 (log ε 3.99) (MeOH).

Quast, H. *et al.*, *Annalen*, 1970, 732, 64-69 (*synth, N-Me*)

Gonzalez, A. *et al.*, *J. Nat. Prod.*, 1999, 62, 378-380 (*N-Me, isol, spectra*)

4-Quinolinecarboxylic acid, 9CI, 8CI Q-32

Cinchoninic acid

[486-74-8]

C₁₀H₇NO₂ 173.171

Metab. of *Archangium gephyra*. Needles + 1 or 2H₂O. Insol. Et₂O; spar. sol. H₂O, EtOH. Mp 253-254°. Possible artifact.

►GD3850000

Methochloride:

Prisms (H₂O). Mp 243° dec.

Methobromide: [17323-98-7]

Needles (H₂O). Mp 262°.

Methiodide:

Orange-red needles (EtOH). Mp 224° dec.

Me ester: [21233-61-4]

C₁₁H₉NO₂ 187.198

Mp 24°.

Et ester: [10447-29-7]

C₁₂H₁₁NO₂ 201.224

Mp 13°. Bp₁₅ 173°.

Chloride: [50821-72-2]

C₁₀H₆ClNO 191.616

Yellow needles. Mp 68°. Bp₁₂ 148°.

Amide: [4363-95-5]

C₁₀H₈N₂O 172.186

Needles. Sol. hot H₂O, hot EtOH. Mp 181°.

Nitrile: 4-Cyanoquinoline. Cinchoninonitrile

[2973-27-5]

C₁₀H₆N₂ 154.171

Mp 102°. Steam-volatile.

N-Oxide: [40614-43-5]

C₁₀H₇NO₃ 189.17

Prisms (AcOH). Mp 256° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 869D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 447C (*nmr*)

Elderfield, R.C. *et al.*, *J.A.C.S.*, 1951, 73, 5622 (*synth*)

Schultz, O.-E. *et al.*, *Annalen*, 1970, 740, 192 (*synth*)


Zalis, B. *et al.*, *Talanta*, 1973, 20, 33 (*uv*)

Böhlendorf, B. *et al.*, *Annalen*, 1996, 49-53 (*isol, pmr, cmr, ms*)

Dobson, A.J. *et al.*, *Acta Cryst. C*, 1998, 54, 1883-1885; 1999, 55, 935-937 (*cryst struct*)

2,4-Quinolinediol Q-33

4-Hydroxy-2(1H)-quinolinone, 9CI. 4-Hydroxycarboxystyryl, 8CI. 2,4-Dihydroxyquinoline. 2-Hydroxy-4(1H)-quinolinone [70254-43-2] [52851-41-9]



C₉H₇NO₂ 161.16

4-One-form predominates. Mp 300°.

pK_{a1} 0.76; pK_{a2} 5.9 (20°).

2-One-form [86-95-3]

►VB7877500

4-Ac: 4-Acetoxy-2(1H)-quinolinone

C₁₁H₉NO₃ 203.197

Needles. Mp 214-216°.

4-Benzoyl:

C₁₆H₁₁NO₃ 265.268

Needles (C₆H₆). Mp 220°.

N-Me: [1677-46-9]

C₁₀H₉NO₂ 175.187

Mp 274-275°.

N-Me, O⁴-benzoyl:

C₁₇H₁₃NO₃ 279.295

Cryst. (EtOH). Mp 173-174°.

N-Benzyl, O⁴-benzoyl:

C₂₃H₁₇NO₃ 355.392

Cryst. (EtOH). Mp 150-152°.

N-Ph: [14994-75-3]

C₁₅H₁₁NO₂ 237.257

Light yellow cryst. (EtOH aq.). Mp 291°.

N-Ph, O⁴-benzoyl: [685895-32-3]

C₂₂H₁₅NO₃ 341.365

Yellow prisms (EtOH). Mp 150-151°.

4-Me ether: 2-Hydroxy-4-methoxyquinoline. 4-Methoxy-2(1H)-quinolinone

[27667-34-1]

C₁₀H₉NO₂ 175.187

Alkaloid from *Haplophyllum bungei*.

Cryst. (hexane). Mp 254-255°.

Me ether, N-Me: 4-Methoxy-1-methyl-2(1H)-quinolinone, 9CI

[32262-18-3]

C₁₁H₁₁NO₂ 189.213

Alkaloid from *Evodia lunu-ankenda*, *Xanthoxylum inerme*, *Hesperethusa crenulata* (preferred genus name *Nar- ingi*), *Fagara boninensis*, *Hortia longifolia*, *Myrtopsis sellingsii*, *Zanthoxylum decaryi*, *Adiscanthus fusciflorus*, and others (Rutaceae). Cryst. (C₆H₆/petrol). Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 100-100.5° Mp 68-69° (hydrate). λ_{max} 229 (ε 48000); 269 (ε 5120); 278 (ε 5130); 318 (ε 4470) (EtOH) (Berdy).

4-Me ether, N-methoxy: 1,4-Dimethoxy-2(1H)-quinolinone, 9CI. **Haplotusine**

[27667-33-0]

C₁₁H₁₁NO₃ 205.213

Alkaloid from the aerial parts of *Haplophyllum obtusifolium*. Mp 118-119°.

4-One-form [70254-44-3]

2-Me ether, N-Me: 2-Methoxy-1-methyl-4(1H)-quinolinone

C₁₁H₁₁NO₂ 189.213

Mp 195°.

2-Et ether: 2-Ethoxy-4-hydroxyquinoline.

2-Ethoxy-4(1H)-quinolinone

C₁₁H₁₁NO₂ 189.213

Mp 228°.

Diol-form

Di-Me ether: 2,4-Dimethoxyquinoline.

Montrutanine. *Montanine*†

[40335-00-0]

C₁₁H₁₁NO₂ 189.213

Alkaloid from the fruit of *Evodia hupehensis* and the aerial parts of *Ruta montana* (Rutaceae). Cryst. (MeOH).

Mp 82°. Doubts have been expressed

by Osborne *et al* as to the veracity of this struct.

Di-Et ether: 2,4-Diethoxyquinoline

[57839-56-2]

C₁₃H₁₅NO₂ 217.267

Cryst. (EtOH) (as picrate). Mp 55-56° (EtOH aq.) Mp 192-193° (picrate).

[4510-76-3]

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 736D (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 859A (ir)

Arndt, F. *et al.*, *Chem. Ind. (London)*, 1950, 465 (Me ethers)

Ziegler, E. *et al.*, *Monatsh. Chem.*, 1965, **96**, 418 (synth)

Paul, B.D. *et al.*, *J. Indian Chem. Soc.*, 1968, **45**, 552-553 (N-Me)

Omori, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1970, **43**, 1135-1138 (synth)

Nayar, M.N.S. *et al.*, *Phytochemistry*, 1971, **10**, 2843 (4-Methoxy-1-methylquinolinone)

Razakova, D.M. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 635; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 599 (Haplotusine)

Reisch, J. *et al.*, *Pharmazie*, 1985, **40**, 812-814 (di-Me ether, isol)

Ulubelen, A. *et al.*, *J. Nat. Prod.*, 1990, **53**, 207 (Montrutanine)

Osborne, A.G. *et al.*, *J. Nat. Prod.*, 1992, **55**, 589-595 (di-Me ether, di-Et ether, pmr, cmr)

Kappe, T. *et al.*, *J. Het. Chem.*, 1996, **33**, 663-670 (O⁴-benzoyl derivs)

Jung, J.-C. *et al.*, *Synth. Commun.*, 2001, **31**, 1195-1200 (synth, ir, pmr, cmr)

Jones, K. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 4380-4383 (4-Me ether, 2,4-di-Me ether, synth, ir, pmr, cmr)

Stadlbauer, W. *et al.*, *J. Het. Chem.*, 2004, **41**, 681-690 (N-Ph O⁴-benzoyl)

Nadaraj, V. *et al.*, *ARKIVOC*, 2006, **x**, 82-89 (synth, 4-Me ether N-Me)

Sheibani, H. *et al.*, *Synthesis*, 2006, 435-438 (N-Ph)

Ambrozin, A.R.P. *et al.*, *Quim. Nova*, 2008, **31**, 740-743 (Haplotusine)

Selig, P. *et al.*, *Synthesis*, 2008, 2177-2182 (4-Ac)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, QNA000

3,8-Quinolinediol

Q-34

3,8-Dihydroxyquinoline. Jineol

[178762-28-2]

C₉H₇NO₂ 161.16

Alkaloid from the centipede *Scolopendra subspinipes*. Exhibits modest cytotoxicity *in vitro* against the growth of several human tumour cell lines. Orange amorph. solid. Mp 139-141°. λ_{max} 254 ; 267 (MeOH).

Di-Ac:

C₁₃H₁₁NO₄ 245.234

Mp 101-102°.

Moon, S.-S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 777-779 (isol, uv, ir, pmr, cmr, ms, struct)

4,8-Quinolinediol

Q-35

8-Hydroxy-4(1H)-quinolinone, 9CI. 4,8-Dihydroxyquinoline

[53846-46-1]

C₉H₇NO₂ 161.16

Alkaloid from octopus ink (*Octopus dofleini martini*) and the urine of silk-worm pupae. Cryst. (CH₂Cl₂). Mp 305-308° dec.

Hydrochloride:

Needles + ½H₂O (Me₂CO/EtOH). Mp 295-298° dec.

8-(4-Methylbenzenesulfonyl): [18996-01-5]

C₁₆H₁₃NO₄S 315.349

Cryst. (EtOH). Mp 237°.

Inagami, K. *et al.*, *CA*, 1956, **50**, 8077a (isol)
Matsuura, Y. *et al.*, *CA*, 1959, **53**, 8373b (synth)
Siuda, J.F. *et al.*, *J. Nat. Prod.*, 1974, **37**, 501-503 (isol, synth, pmr)

Grandjean, D. *et al.*, *Bull. Soc. Chim. Fr.*, 1989, 657-660 (synth, pmr)


Heiskanen, J.P. *et al.*, *J.O.C.*, 2007, **72**, 920-922 (synth, tosyl)

2-Quinolinemethanethiol, 9CI

Q-36

2-(Thiomethyl)quinoline. 2-(Mercapto-methyl)quinoline

[83492-32-4]



C₁₀H₉NS 175.254

Constit. of the defensive secretions of the skunks *Conepatus mesoleucus*, *Mephitis mephitis* and *Spilogale putorius*. Prisms (EtOH/petrol) (as hydrobromide). Mp 150-151° dec. (hydrobromide).

S-Ac: *S*-(2-Quinolinylmethyl) thioacetate. 2-(Acetylthiomethyl)quinoline [129120-37-2]

C₁₂H₁₁NOS 217.291

Isol. from the anal sac of *Mephitis mephitis*.

[99409-21-9 ; 83492-33-5 ; 83492-25-5]

Yousif, M.M. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 1974-1979 (synth)

Sturis, A. *et al.*, *CA*, 1986, **104**, 33980p (synth)


Wood, W.F. *et al.*, *J. Chem. Ecol.*, 1990, **16**, 2057-2065; 1991, **17**, 1415-1420; 1993, **19**, 837-841 (occur)

2-Quinolinemethanol, 9CI

Q-37

2-(Hydroxymethyl)quinoline. α-Hydroxy-quinaldine. α-Quinolylcarbinol. α-Quinaldinol. Antibiotic PO 1227B. PO 1227B

[1780-17-2]



C₁₀H₉NO 159.187

Prod. by *Kitasatoa griseophaeus*. Exhibits hypoglycaemic activity. Needles (EtOH or petrol). Mp 66-67° (64°). Bp 145-160°. Steam-volatile. λ_{max} 230 (ε 47700); 233 (ε 44500); 277 (ε 4000); 284 (ε 3800); 290 (ε 3700); 296 (ε 3300); 303 (ε 4000) (MeOH) (Berdy).

Ac: 2-(Acetoxy-methyl)quinoline. α-Acetoxyquinaldine. Quinaldyl acetate. Antibiotic PO 1227A. PO 1227A [60483-07-0]

C₁₂H₁₁NO₂ 201.224

Prod. by *Kitasatoa griseophaeus* PO-1227. Cryst. (hexane).

Benzoyl:

C₁₇H₁₃NO₂ 263.295

Mp 52-53°.

Hammick, D.L. *et al.*, *J.C.S.*, 1926, 1302-1304 (synth)

Kaslow, C.E. *et al.*, *J.O.C.*, 1953, **18**, 55-58 (synth)

Kato, T. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 1649-1652; *CA*, **59**, 2765 (synth)

Omura, S. *et al.*, *J. Antibiot.*, 1976, **29**, 797-803 (isol)

4-Quinolinemethanol, 9CI

Q-38

4-(Hydroxymethyl)quinoline

[6281-32-9]

C₁₀H₉NO 159.187

Alkaloid from the fungi *Polyporus sanguineus* and *Polyporus versicolor*. Cryst. (MeOH). Mp 96-97° (92° dec.).

Ac: [35982-82-2]

C₁₂H₁₁NO₂ 201.224

Mp 55-57°.

Palmer, M.H. *et al.*, *Tet. Lett.*, 1968, 2147 (synth)

Forbis, R.M. *et al.*, *J.A.C.S.*, 1973, **95**, 5003 (synth)

Takadate, A. *et al.*, *Synthesis*, 1983, 806 (synth, ir, pmr)

Abraham, W.-R. *et al.*, *Phytochemistry*, 1991, **30**, 371 (isol)

Wender, P.A. *et al.*, *Synthesis*, 1994, 1278 (synth, ir, pmr, cmr)


Böhendorf, B. *et al.*, *Annalen*, 1996, 49 (isol, ir, pmr, cmr, ms)

Mikolajczyk, M. *et al.*, *J.O.C.*, 2006, **71**, 8818-8823 (synth)

2,3,4,7,8-Quinolinetetrol

Q-39

2,3,4,7,8-Pentahydroxyquinoline. 3,4,7,8-Tetrahydroxy-2(1H)-quinolinone



C₉H₇NO₅ 209.158

2,3-Methylene, 4,7,8-tri-Me ether: 4,7,8-Trimethoxy-2,3-methylenedioxyquinoline

[200337-59-3]

C₁₃H₁₃NO₅ 263.249

Alkaloid from the leaves of *Acanthosyrhis paulo-alvini* (Santalaceae). Needles. Mp 112-114°. λ_{max} 243 (log ε 4.01); 272 (log ε 3.33) (no solvent reported).


Chavez, J.P. *et al.*, *Phytochemistry*, 1997, **46**, 967-968 (isol, uv, ir, pmr, cmr, ms)

2,3,4,7-Quinolinetetrol

Q-40

3,4,7-Trihydroxy-2(1H)-quinolinone.

2,3,4,7-Tetrahydroxyquinoline



C₉H₇NO₄ 193.159

2,3-Methylene, 4,7-di-Me ether: 4,7-Di-

methoxy-2,3-methylenedioxyquinolineC₁₂H₁₁NO₄ 233.223Alkaloid from the roots of *Acronychia laurifolia*. Plates (MeOH). Mp 176-178°. λ_{max} 242 (log ε 1.5); 255 (log ε 2.8); 308 (log ε 3.5); 335 (log ε 3.8) (MeOH).Cui, B. et al., *Phytochemistry*, 1999, **52**, 95-98 (isol, uv, ir, pmr, cmr, ms)**2,3,4,8-Quinolinetretol** Q-41

3,4,8-Trihydroxy-2(1H)-quinolinone.

2,3,4,8-Tetrahydroxyquinoline

C₉H₇NO₄ 193.159**NH-form**3,4,8-Tri-Me ether: **3,4,8-Trimethoxy-2(1H)-quinolinone**

[164163-88-6]

C₁₂H₁₃NO₄ 235.239Alkaloid from aerial parts of *Eriostemon gardneri* (Rutaceae). Amorph. solid.N,O³,O⁸-Tri-Me: 4-Hydroxy-3,8-dimethoxy-1-methyl-2(1H)-quinolinone.**Swietenidin A**

[64595-74-0]

C₁₂H₁₃NO₄ 235.239Alkaloid from the bark of *Chloroxylon swietenia* (Flindersiaceae). Cryst. (EtOAc). Mp 164-166° (158-159°).Bhide, K.S. et al., *Indian J. Chem., Sect. B*, 1977, **15**, 440 (isol, uv, ir, pmr, struct, Swietenidin A)Coppola, G.M. et al., *J. Het. Chem.*, 1985, **22**, 1087 (synth, ir, pmr, cmr, Swietenidin A)Sarkar, S.D. et al., *J. Nat. Prod.*, 1995, **58**, 574 (isol, uv, ir, pmr, cmr, ms, struct, deriv)Fischer, M. et al., *Sci. Pharm.*, 1996, **64**, 353 (synth, tri-Me ether)**2,4,6,8-Quinolinetretol** Q-42

4,6,8-Trihydroxy-2(1H)-quinolinone,

9CI. 2,4,6,8-Tetrahydroxyquinoline

C₉H₇NO₄ 193.159

6,8-Di-Me ether: 6,8-Dimethoxy-2,4-dihydroxyquinoline. 4-Hydroxy-6,8-dimethoxy-2(1H)-quinolinone. 6,8-Dimethoxy-2,4-quinolinediol

[39022-15-6]

C₁₁H₁₁NO₄ 221.212

Mp 290°.

4,6,8-Tri-Me ether: 4,6,8-Trimethoxy-2(1H)-quinolinone. 2-Hydroxy-4,6,8-trimethoxyquinoline. **Halfordamine**

[21144-33-2]

C₁₂H₁₃NO₄ 235.239Alkaloid from *Halfordia scleroxyla* (Rutaceae). Cryst. (C₆H₆/petrol). Mp 240-241°.

4,6,8-Tri-Me ether, N-Me: 4,6,8-Trimethoxy-1-methyl-2(1H)-quinolinone

[38989-33-2]

C₁₃H₁₅NO₄ 249.266

Mp 163°.

Crow, W.D. et al., *Aust. J. Chem.*, 1968, **21**, 3075 (isol, uv, ir, pmr, ms)Storer, R. et al., *Tetrahedron*, 1973, **29**, 1215 (synth, uv, ir, pmr, struct)Venturella, P. et al., *Gazz. Chim. Ital.*, 1974, **104**, 297 (synth, uv, ir, pmr, struct)Mali, R.S. et al., *Indian J. Chem., Sect. B*, 1977, **15**, 865 (synth)Nadaraj, V. et al., *ARKIVOC*, 2006, x, 82-89 (6,8-di-Me ether, 4,6,8-tri-Me ether N-Me)**2,4,7,8-Quinolinetretol, 9CI** Q-43

4,7,8-Trihydroxy-2(1H)-quinolinone.

2,4,7,8-Tetrahydroxyquinoline

C₉H₇NO₄ 193.159

7,8-Di-Me ether: 2,4-Dihydroxy-7,8-dimethoxyquinoline. 7,8-Dimethoxy-2,4-quinolinediol. 7,8-Dimethoxy-4-hydroxy-2(1H)-quinolinone

[43215-49-2]

C₁₁H₁₁NO₄ 221.212

Mp 246°.

4,7,8-Tri-Me ether: 4,7,8-Trimethoxy-2(1H)-quinolinone. 2-Hydroxy-4,7,8-trimethoxyquinoline. **Haplobungine**

[121949-99-3]

C₁₂H₁₃NO₄ 235.239Alkaloid from aerial parts of flowering *Haplophyllum bungei* (Rutaceae). Mp 174-175°.**NH-form**

4,7,8-Tri-Me, N-Me: 4,7,8-Trimethoxy-1-methyl-2(1H)-quinolinone

[43215-43-6]

C₁₃H₁₅NO₄ 249.266Minor alkaloid from the roots and stems of *Spathelia sorbifolia* (Rutaceae). Cryst. (petrol). Mp 143-148°.Storer, R. et al., *Tetrahedron*, 1973, **29**, 1721 (Trimethoxymethylquinolinone)Bessonova, I.A. et al., *Khim. Prir. Soedin.*, 1989, **25**, 23; *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 18 (Haplobungine)Nadaraj, V. et al., *ARKIVOC*, 2006, x, 82-89 (7,8-di-Me ether, tri-Me ether N-Me)**3,4,5,8-Quinolinetretol** Q-44


3,5,8-Trihydroxy-4(1H)-quinolinone,

9CI. 3,4,5,8-Tetrahydroxyquinoline. **Uranidine**

[92264-09-0]

C₉H₇NO₄ 193.159Yellow zoochore of the sponge *Verngeria aerophoba*. Also from *Tylodina perversa*. HIV reverse transcriptase (HIV-rt) inhibitor. Very unstable; rapidly turns blue, then purple, and finally black on exposure to air. λ_{max} 240 (ε 15000); 321 (ε 2600); 368 (ε 2900) (pH 1 H₂O) (Derep). λ_{max} 236 (ε 14700); 315 (ε 2900); 604 (ε 3000); 641 (ε 3100) (pH 13 H₂O) (Derep). λ_{max} 239 (ε 15000); 319 (ε 2600); 365 (ε 2900) (H₂O pH 7) (Derep).Cimino, G. et al., *Tet. Lett.*, 1984, **25**, 2925-2928 (isol, uv, pmr, cmr)Loya, S. et al., *Arch. Biochem. Biophys.*, 1994, **309**, 315-322 (activity)**2,3,4-Quinolinetriol, 9CI** Q-45

3,4-Dihydroxy-2(1H)-quinolinone. 2,3,4-Trihydroxyquinoline

C₉H₇NO₃ 177.159C₉H₇NO₃ 177.1593,4-Di-Me ether: 3,4-Dimethoxy-2(1H)-quinolinone. 2-Hydroxy-3,4-dimethoxyquinoline. **Swietenidin B**

[2721-56-4]

C₁₁H₁₁NO₃ 205.213Alkaloid from the bark of *Chloroxylon swietenia* (Flindersiaceae). Needles (C₆H₆/Me₂CO or EtOAc). Mp 182°.**NH-form**

3,4-Di-Me ether, N-Me: 3,4-Dimethoxy-1-methyl-2(1H)-quinolinone, 9CI. N-

Methylswietenidin B

[123497-83-6]

C₁₂H₁₃NO₃ 219.24Alkaloid from combined stem bark and root extracts of *Clausena anisata* (Rutaceae). Plates (CH₂Cl₂). Mp 74-75°.Ziegler, E. et al., *Monatsh. Chem.*, 1965, **96**, 889 (synth)Bhide, K.S. et al., *Indian J. Chem., Sect. B*, 1977, **15**, 440 (Swietenidin B)Ngadjui, B.T. et al., *Phytochemistry*, 1989, **28**, 1517 (N-Methylswietenidin B)**2,4,6-Quinolinetriol** Q-46

4,6-Dihydroxy-2(1H)-quinolinone. 2,4,6-Trihydroxyquinoline

C₉H₇NO₃ 177.159

6-Me ether: 2,4-Dihydroxy-6-methoxyquinoline. 6-Methoxy-2,4-quinolinediol. 4-Hydroxy-6-methoxy-2(1H)-quinolinone

[14300-45-9]

C₁₀H₉NO₃ 191.186

Mp 300°.

NH-form

4-Me ether, N-Me: 6-Hydroxy-4-methoxy-1-methyl-2(1H)-quinolinone, 9CI.

Integrinquinone

[81943-13-7]

C₁₁H₁₁NO₃ 205.213Alkaloid from roots of *Zanthoxylum integrifolium* (Rutaceae). Needles (MeOH). Mp 257-260°.

4,6-Di-Me ether, N-Me: 4,6-Dimethoxy-1-methyl-2(1H)-quinolinone, 9CI

[52345-94-5]

C₁₂H₁₃NO₃ 219.24Alkaloid from aerial parts of an *Agathosma* sp. (Rutaceae). Needles (C₆H₆/petrol). Mp 143-144°.Ishii, H. et al., *Chem. Pharm. Bull.*, 1982, **30**, 1992 (uv, ir, pmr, struct, Integrinquinone)Campbell, W.E. et al., *Phytochemistry*, 1990, **29**, 1303 (isol, uv, ir, pmr, cmr, ms)Nadaraj, V. et al., *ARKIVOC*, 2006, x, 82-89 (6-Me ether, 4,6-di-Me ether N-Me)**2,4,8-Quinolinetriol** Q-47

2,4,8-Trihydroxyquinoline. 4,8-Dihydroxy-2(1H)-quinolinone

C₉H₇NO₃ 177.159

Other tautomers possible.

8-Me ether: 2,4-Dihydroxy-8-methoxyquinoline. 8-Methoxy-2,4-quinolinediol. 4-Hydroxy-8-methoxy-2(1H)-quinolinone

[7224-68-2]

C₁₀H₉NO₃ 191.186
Mp 248°.

4,8-Di-Me ether: 4,8-Dimethoxy-2(1H)-quinolinone. **Edulitine**. Robustinine [15272-24-9]

C₁₁H₁₁NO₃ 205.213

Alkaloid from bark of *Casimiroa edulis* (Mexican apple) (Rutaceae). Needles (EtOAc). Mp 235-236°.

4,8-Di-Me ether, picrate:

Yellow cryst. (MeOH). Mp 189-191°.

NH-form

O⁴,N-Di-Me, 8-O-β-D-glucopyranoside:

C₁₆H₁₉NO₈ 353.328

Alkaloid from the aerial parts of *Echinops gmelinii*. Needles (MeOH). Mp 242° dec. [α]_D²⁰ -4.3 (c, 0.23 in Py).

O⁴,N-Di-Me: 8-Hydroxy-4-methoxy-1-methyl-2(1H)-quinolinone. **Folifidine** [3148-23-0]

C₁₁H₁₁NO₃ 205.213

Alkaloid from *Haplophyllum foliosum* (Rutaceae). Cryst. (EtOH). Mp 226-227°.

O⁴,N-Di-Me; hydrochloride:

Cryst. (EtOH). Mp 232° dec.

O⁴,N-Di-Me, picrate:

Cryst. (EtOH). Mp 218°.

O⁴,N-Di-Me, 8-O-β-D-glucopyranoside:

C₁₇H₂₁NO₈ 367.355

Alkaloid from the aerial parts of *Echinops gmelinii*. Needles (MeOH). Mp > 300°. [α]_D²⁰ -20.4 (c, 0.54 in Py).

O⁴,O⁸,N-Tri-Me: 4,8-Dimethoxy-1-methyl-2(1H)-quinolinone. **Folimine** [3148-24-1]

C₁₂H₁₃NO₃ 219.24

Alkaloid from *Haplophyllum foliosum* and *Haplophyllum perforatum* (Rutaceae). Mp 139-140°.

O⁴,O⁸,N-Tri-Me; hydrochloride: Mp 171-172°.

O⁴,O⁸,N-Tri-Me, picrate: Mp 193-194°.

8-O-(3-Methyl-2-butenyl), O⁴,N-di-Me: 4-Methoxy-1-methyl-8-[(3-methyl-2-butenyl)oxy]-2(1H)-quinolinone. 4-Methoxy-1-methyl-8-prenyloxyquinolone. **Daurine**

[54357-79-8]

C₁₆H₁₉NO₃ 273.331

Alkaloid from roots of *Haplophyllum dauricum* (Rutaceae). Needles (CH₂Cl₂/petrol). Mp 118-119°.

Iriarte, J. et al., *J.C.S.*, 1956, 4170 (*Edulitine, isol, uv, ir*)

Faizutdinova, Z.Sh. et al., *Khim. Prir. Soedin.*, 1967, **3**, 257; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 215 (*Folifidine*)

Toube, T.P. et al., *Tetrahedron*, 1967, **23**, 2061 (*Edulitine, pmr, ms*)

Narasimhan, N.S. et al., *Tetrahedron*, 1971, **27**, 1351 (*synth*)

Razzakova, I.A. et al., *Khim. Prir. Soedin.*, 1972, **8**, 133; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 139 (*Folimine*)

Bessonova, I.A. et al., *Khim. Prir. Soedin.*, 1983, **19**, 116; 1984, **20**, 73; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 117; 1984, **20**, 68 (*Daurine*)

Reisch, J. et al., *Monatsh. Chem.*, 1988, **119**, 1169 (*Daurine, synth, pmr, ms*)

Su, Y.-F. et al., *J. Asian Nat. Prod. Res.*, 2004, **6**, 223-227 (*Echinops gmelinii glucosides*)

Nadaraj, V. et al., *ARKIVOC*, 2006, **x**, 82-89 (8-Me ether, folimine)

4,7,8-Quinolinetriol

Q-48

4,7,8-Trihydroxyquinoline. 7,8-Dihydroxy-4(1H)-quinolinone

C₉H₇NO₃ 177.159

8-Me ether: 4,7-Dihydroxy-8-methoxyquinoline. 8-Methoxy-4,7-quinolinediol. 7-Hydroxy-8-methoxy-4(1H)-quinolinone

[98267-21-1]

C₁₀H₉NO₃ 191.186

Isol. from the soft corals *Simularia polydactyla* and *Simularia microclavata*. Cardiovascular agent.

7,8-Di-Me ether: 4-Hydroxy-7,8-dimethoxyquinoline. 7,8-Dimethoxy-4(1H)-quinolinone. 7,8-Dimethoxy-4-quinolinol

[99878-76-9]

C₁₁H₁₁NO₃ 205.213

Cryst. (EtOAc/MeOH). Mp 187-190°.

[108915-16-8]

Long, K. et al., *CA*, 1985, **103**, 128867; 1990, **112**, 118619; 1991, **115**, 71358 (*synth, isol, pharmacol*)

Kelly, T.R. et al., *Tetrahedron*, 1985, **41**, 3033 (*deriv*)


Huang, S. et al., *Jiegou Huaxue*, 1986, **5**, 1; *CA*, **107**, 39495 (*cryst struct*)

3-Quinolinol, 9CI

Q-49

3-Hydroxyquinoline

[580-18-7]



C₉H₇NO 145.16

Alkaloid from aerial parts of *Ruta montana*. Cryst. (MeOH). Mp 201-202°. pK_{a1} 4.28; pK_{a2} 8.08 (20°).

▶VC4050000

N-Oxide:

C₉H₇NO₂ 161.16

Solid (EtOH/CHCl₃). Mp 218°.

O-Ac:

C₁₁H₉NO₂ 187.198

Needles (petrol). Mp 37.5°.

Me ether: 3-Methoxyquinoline

[6931-17-5]

C₁₀H₉NO 159.187

Pale brown oil. Mp 222° (as picrate).

Et ether: 3-Ethoxyquinoline

C₁₁H₁₁NO 173.214

Mp 224-228° (as picrate).

[1321-40-0]

Org. Synth., 1960, **40**, 54 (*synth*)

Kaneko, C. et al., *Chem. Pharm. Bull.*, 1967, **15**, 663 (*synth, uv*)

Naumann, C. et al., *Synthesis*, 1990, 279 (*synth, ir, pmr*)


Ulubelen, A. et al., *Fitoterapia*, 1991, **62**, 279 (*isol, pmr, ms*)

Bergeron, R.J. et al., *J. Med. Chem.*, 1999, **42**, 95-108 (*synth, oxide, pmr*)

2(1H)-Quinolinone, 9CI

Q-50

Carbostyryl, 8CI. 2-Hydroxyquinoline. 2-Quinololinol [59-31-4]



C₉H₇NO 145.16

NH-Tautomer predominates. Cryst. + 1H₂O. Mp 199-200° (anhyd.). pK_{a1} -0.31; pK_{a2} 11.74.

▶LD₅₀ (mus, ipr) 150 mg/kg. FG7175000

NH-form

N-Me: 1-Methyl-2(1H)-quinolinone

[606-43-9]

C₁₀H₉NO 159.187

Alkaloid from the bark of *Galipea officinalis*. Needles (petrol). Mp 74°.

N-Et: [53761-50-5]

C₁₁H₁₁NO 173.214

Mp 54°. Bp 316-318°.

N-Ph: [67176-94-7]

C₁₅H₁₁NO 221.258

Needles (EtOH aq.). Mp 138°.

I-Amino: [26539-38-8]

C₉H₈N₂O 160.175

Plates (EtOH). Mp 107-108°.

OH-form [70254-42-1]

Me ether: 2-Methoxyquinoline

C₁₀H₉NO 159.187

Bp 246-247°.

Me ether, N-oxide:

C₁₀H₉NO₂ 175.187

Mp 70-72°.

Et ether: 2-Ethoxyquinoline

C₁₁H₁₁NO 173.214

Bp 266°.

Et ether, N-oxide:

C₁₁H₁₁NO₂ 189.213

Mp 75-77°.

[1321-40-0]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 431C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 1570A (*ir*)

Bamberger, E. et al., *Ber.*, 1894, **27**, 203

Dieckmann, W. et al., *Ber.*, 1908, **41**, 3268

Chichibabin, A.E. et al., *CA*, 1925, **19**, 1572 (*synth*)

Effenberger, F. et al., *Angew. Chem.*, 1964, **76**, 188 (*synth, ethers*)

Brasiunas, V. et al., *CA*, 1964, **61**, 5608 (*synth*)

Mack, W. et al., *Angew. Chem.*, *Int. Ed.*, 1966, **5**, 896 (*synth, ir, uv, pmr, ms*)

Hampson, P. et al., *Chem. Comm.*, 1967, 371 (*N-14 nmr, ethers*)

Katritzky, A.R. et al., *An. Quim.*, 1974, **70**, 994 (*I-amino*)

Claret, P.A. et al., *Spectrosc. Lett.*, 1976, **9**, 157; 167 (*pmr, cmr*)


Ribar, B. et al., *Cryst. Struct. Commun.*, 1977, **6**, 677 (*cryst struct*)

Kido, M. et al., *Chem. Pharm. Bull.*, 1982, **30**, 1488 (*cryst struct*)

Bird, C.W. et al., *Tetrahedron*, 1985, **41**, 1409 (*props*)

- Gronowitz, S. *et al.*, *Heteroat. Chem.*, 1986, **5**, 73 (rev)
 Wawzonek, S. *et al.*, *J. Het. Chem.*, 1988, **25**, 381 (pmr, deriv)
 Grignon-Dubois, M. *et al.*, *Synth. Commun.*, 1995, **25**, 2999 (N-Me)
 Houghton, P.J. *et al.*, *Planta Med.*, 1999, **65**, 250-254 (isol, N-Me)
 Horaguchi, T. *et al.*, *J. Het. Chem.*, 2002, **39**, 61-67 (synth, ir, pmr, cmr)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CCC250

1-(2-Quinoliny)-β-carboline Q-51
 1-(2-Quinoliny)-9H-pyrido[3,4-b]indole, 9CI. **Nitramarine**
 [95360-17-1]



- $C_{20}H_{13}N_3$ 295.343
 Alkaloid from the aerial parts of *Nitraria komarovii* (Zygophyllaceae). Promotes sleep in expl. animals. Needles. Mp 180.5-182° (172-173°).
 1,2,3,4-Tetrahydro: 1,2,3,4-Tetrahydro-1-(2-quinoliny)-β-carboline. **Tetrahydro-nitramarine**
 [95360-18-2]
 $C_{20}H_{17}N_3$ 299.374
 Alkaloid from aerial parts of *Nitraria komarovii*. Mp 193-194°. Opt. inactive.
 Tulyaganov, T.S. *et al.*, *Khim.-Farm. Zh.*, 1984, **18**, 1474; *CA*, **102**, 128811g
 Hibino, S. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2261 (synth, uv, pmr, pharmacol)
 Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1990, **25**, 61; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **25**, 49 (Tetrahydro-nitramarine)
 Rocca, P. *et al.*, *Tetrahedron*, 1993, **49**, 3325 (synth, ir, pmr, cmr)
 Molina, P. *et al.*, *Tet. Lett.*, 1994, **35**, 8851 (synth)
 Bracher, F. *et al.*, *Tetrahedron*, 1994, **50**, 12329 (synth)

1-(4-Quinoliny)-β-carboline Q-52
 1-(4-Quinoliny)-9H-pyrido[3,4-b]indole. **Nitraridine**
 [915708-51-9]

- $C_{20}H_{13}N_3$ 295.343
 Alkaloid from the aerial parts of *Nitraria komarovii*. Cryst. (CH₂Cl₂). Mp 272-273°. λ_{max} 211 (log ε 4.65); 233 (log ε 4.65); 252 (sh) (log ε 4.32); 294 (log ε 4.18); 317 (sh) (log ε 3.93); 360 (log ε 3.77) (EtOH).

- 3,4-Dihydro: **Dihydroneitraridine**
 [915708-52-0]
 $C_{20}H_{15}N_3$ 297.359
 Alkaloid from *Nitraria komarovii*. Mp 262-263°. λ_{max} 218 (log ε 4.51); 244 (sh) (log ε 4.07); 307 (log ε 3.93); 312

- (log ε 3.98) (EtOH).
 1,2,3,4-Tetrahydro: **Tetrahydroneitraridine**
 [6649-85-0]
 $C_{20}H_{17}N_3$ 299.374
 Alkaloid from *Nitraria komarovii*. Mp 176-177°. Racemate. λ_{max} 227 (log ε 4.52); 280 (log ε 4.01); 304 (log ε 3.84); 316 (log ε 3.72) (EtOH).
 Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2006, **42**, 459-461 (isol, synth, pmr)

1-(5-Quinoliny)-β-carboline Q-53
 1-(5-Quinoliny)-9H-pyrido[3,4-b]indole, 9CI. **Isokomarovine**
 [85403-68-5]

- $C_{20}H_{13}N_3$ 295.343
 Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Mp 321-322°. λ_{max} 220 (log ε 4.65); 292 (log ε 4.18); 358 (log ε 3.69) (no solvent reported).

- 3,4-Dihydro: **Dihydroisokomarovine**
 [85403-69-6]
 $C_{20}H_{15}N_3$ 297.359
 Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (CHCl₃/MeOH). Mp 252-253°. λ_{max} 220 (log ε 4.6); 246 (sh) (log ε 4.66); 293 (log ε 4.1); 320 (log ε 3.87) (no solvent reported).

- 1,2,3,4-Tetrahydro: 2,3,4,9-Tetrahydro-1-(5-quinoliny)-1H-pyrido[3,4-b]indole. **Tetrahydroisokomarovine**
 [127498-35-5]

- $C_{20}H_{17}N_3$ 299.374
 Alkaloid from *Nitraria komarovii*. Cryst. (CH₂Cl₂). Mp 274-275°.
 Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 635-638; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 601-603 (isol, struct)
 Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1990, **26**, 61-67; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 49-53 (Dihydroisokomarovine, Tetrahydroisokomarovine)

1-(6-Quinoliny)-β-carboline Q-54
 1-(6-Quinoliny)-9H-pyrido[3,4-b]indole. **Komarovine**
 [85412-79-9]

- $C_{20}H_{13}N_3$ 295.343
 Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (MeOH/CHCl₃). Mp 239-240°.

- 1,2,3,4-Tetrahydro: **Tetrahydrokomarovine**
 [85403-70-9]
 [112583-69-4]
 $C_{20}H_{17}N_3$ 299.374
 Alkaloid from *Nitraria komarovii*. Cryst. (C₆H₆/CH₂Cl₂). Mp 252-253°.

- Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 638-640; 1990, **26**, 61-67; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 604-605; 1990, **26**, 49-53 (isol, ir, pmr, ms)

1-(8-Quinoliny)-β-carboline Q-55
 1-(8-Quinoliny)-9H-pyrido[3,4-b]indole, 9CI. **Komarovine**
 [62209-25-0]

- $C_{20}H_{13}N_3$ 295.343


- Alkaloid from the above-ground parts of *Nitraria komarovii* (Zygophyllaceae). Cryst. (MeOH/CHCl₃). Mp 229-230°.

- 3,4-Dihydro: **Komarovidine**
 [76971-22-7]
 $C_{20}H_{15}N_3$ 297.359
 Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (MeOH/CHCl₃). Mp 219-220°.

- 1,2,3,4-Tetrahydro: **Komarovicine**
 [85403-71-0]
 $C_{20}H_{17}N_3$ 299.374
 Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (MeOH/CHCl₃). Mp 209-210°. Racemate.

- Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1980, **15**, 654 (Komarovicine)
 Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 598-600 (Komarovicine, Komarovidine)


N²-[(2-Quinoliny)carbonyl]arginine, 9CI Q-56
 N²-Quinaldylarginine



- $C_{16}H_{19}N_5O_3$ 329.358

- (S)-form
 L-form
 [180082-49-9]
 Isol. from the ladybird *Subcoccinella 24-punctata*. Insect defence substance, feeding deterrent. Amorph. solid. (as hydrochloride). Sol. MeOH. [α]_D +26.8 (c, 0.47 in MeOH) (monohydrochloride). CAS no. refers to hydrochloride. λ_{max} 238 (ε 22535); 315 (ε 2066); 352 (ε 1485) (MeOH) (monohydrochloride). λ_{max} 238 (ε 22535); 315 (ε 2066); 352 (ε 1485) (MeOH) (Berdy).
 Wang, S.F. *et al.*, *Experientia*, 1996, **52**, 628-630 (isol, synth, pmr, cmr)

***Oceanapia* Quinolone alkaloid** Q-57




Absolute Configuration

- $C_{24}H_{22}Br_2N_6O_8$ 682.281
 Isol. from the sponge *Oceanapia* sp. Inhibitor of mycotoxin S-conjugate amidase. [α]_D²⁰ -150 (c, 0.19 in MeOH).

- Nicholas, G.M. *et al.*, *Org. Lett.*, 2001, **3**, 1543-1545 (isol, cd, pmr, cmr)

Quinomycin A, 9CI **Q-58**

Echinomycin. Actinoleukin. X 948. X 53III. Antibiotic 1491. Antibiotic 59266. Antibiotic X 948. Antibiotic X 53III [512-64-1]



R = R' = -CH(CH₃)₂
Absolute Configuration

C₅₁H₆₄N₁₂O₁₂S₂ 1101.272

Depsideptide antibiotic. Produced by *Streptomyces echinatus* and *Streptomyces lasaliensis*. Active against gram-positive and negative bacteria and some viruses. Under investigation as an antineoplastic agent. Cryst. Mp 221-223° (217-218°). [α]_D²⁰ -310 (c, 0.86 in CHCl₃). Intercalates with DNA. λ_{max} 242 (ε 63300); 322 (ε 11560) (MeOH) (Derep).

► Adverse effects reported when used therapeutically. LD₅₀ (mus, ipr) 0.4 mg/kg. LD₅₀ (mus, scu) 3.8 mg/kg. JW5250000

7-S-Oxide: **Antibiotic FD 991. FD 991** [160918-29-6]

C₅₁H₆₄N₁₂O₁₃S₂ 1117.271
Prod. by *Streptomyces echinatus*. Cytotoxic agent. λ_{max} 243 (ε 66000); 320 (ε 12000) (MeOH) (Berdy).

Corbaz, R. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 199 (*isol*)

Japan. Pat., 1970, 70 17 592; *CA*, **73**, 75659 (*deriv*)

Martin, G.D. *et al.*, *J. Antibiot.*, 1975, **28**, 332 (*struct*)

Dell, A. *et al.*, *J.A.C.S.*, 1975, **97**, 2497 (*struct*)
Cheung, H.T. *et al.*, *J.A.C.S.*, 1978, **100**, 46 (*conform*)

Waring, M.J. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 173 (*rev*)

Steinerova, N. *et al.*, *Folia Microbiol. (Prague)*, 1987, **32**, 1 (*isol*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 477

Japan. Pat., 1994, 94 316 595; *CA*, **122**, 131143j (*FD 991*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, EAD500

Quinomycin B₀, 8CI **Q-59**

[13602-50-1]

As Quinomycin A, Q-58 with

R = -CH(CH₃)₂, R' = -CH(CH₃)CH(CH₃)₂

C₅₃H₆₈N₁₂O₁₂S₂ 1129.325

Depsideptide antibiotic. Isol. from *Streptomyces aureus* with added leucine. Active against gram-positive bacteria, shows some antitumour activity. Powder. Sol. MeOH, C₆H₆; poorly sol. H₂O, hexane. Mp 215-219° dec. λ_{max} 243 (ε 66070); 320 (ε 12600) (MeOH) (Berdy). Yoshida, T. *et al.*, *J. Antibiot., Ser. A*, 1961, **14**, 330 (*isol, ir, uv*)

Ōtsuka, M. *et al.*, *Tetrahedron*, 1967, **23**, 1535 (*isol, ir, uv*)

Dell, A. *et al.*, *J.A.C.S.*, 1975, **97**, 2497 (*struct, ir, ms, pmr, cmr*)

Quinomycin B **Q-60**

A528. *Antibiotic A528*

[13602-52-3]

As Quinomycin A, Q-58 with
R = R' = -CH(CH₃)CH₂CH₃ (R-)

C₅₃H₆₈N₁₂O₁₂S₂ 1129.325

Depsideptide antibiotic. Prod. by *Streptomyces aureus* in a medium supplemented with DL-isoleucine. Active against gram-positive bacteria, shows some antitumour activity. Needles. Sol. Me₂CO, EtOAc, dioxan, acids, HCl; fairly sol. MeOH, C₆H₆, EtOH; poorly sol. H₂O, hexane. Mp 221° dec. [α]_D²⁶ -300 (c, 1.308 in CHCl₃). λ_{max} 242 (E1%/1cm 678); 322 (E1%/1cm 118) (MeOH) (Berdy).

► LD₅₀ (mus, ivn) .054 mg/kg, LD₅₀ (mus, ipr) .04 mg/kg. VC8935000

[11001-72-2]

Yoshida, T. *et al.*, *J. Antibiot., Ser. A*, 1962, **15**, 272 (*isol, ir, uv*)

Ōtsuka, H. *et al.*, *Tetrahedron*, 1967, **23**, 1535 (*isol, ir, uv*)

Martin, G.D. *et al.*, *J. Antibiot.*, 1975, **28**, 332-336 (*struct*)

Dell, A. *et al.*, *J.A.C.S.*, 1975, **97**, 2497 (*struct, ir, ms, pmr, cmr*)

Quinomycin C **Q-61**

Antibiotic 6270†. U 48160. Antibiotic U 48160

[11001-74-4]

As Quinomycin A, Q-58 with

R = R' = -CH(CH₃)CH(CH₃)₂

C₅₅H₇₂N₁₂O₁₂S₂ 1157.379

Depsideptide antibiotic. Prod. by *Streptomyces aureus* and *Streptomyces flavochromogenes*. Cytotoxic. Needles (CHCl₃). Mp 218-220° dec. [α]_D²⁵ -250 (c, 1 in CHCl₃). λ_{max} 243 ; 322 (MeOH). Brazhnikova, M.G. *et al.*, *CA*, 1960, **54**, 25028eg; 1961, **55**, 7664b (*isol*)

Yoshida, T. *et al.*, *J. Antibiot.*, 1961, **14**, 324-329; 330-334 (*isol*)

Otsuka, H. *et al.*, *J. Antibiot.*, 1966, **19**, 128-131 (*isol*)

Martin, G.D. *et al.*, *J. Antibiot.*, 1975, **28**, 332-336 (*struct*)

Quinomycin D, 8CI **Q-62**

[11016-62-9]

As Quinomycin A, Q-58 with

R = -CH(CH₃)₂, R' = CH(CH₃)CH₂CH₃ (R-)

C₅₂H₆₆N₁₂O₁₂S₂ 1115.299

Depsideptide antibiotic. Isol. from

Streptomyces aureus as trace component. Shows antibacterial activity. Sol. MeOH, C₆H₆; poorly sol. hexane, H₂O. λ_{max} 243 (ε 63100); 320 (ε 12300) (MeOH) (Berdy). Otsuka, M. *et al.*, *Tetrahedron*, 1967, **23**, 1535 (*isol, ir, uv*)

Dell, A. *et al.*, *J.A.C.S.*, 1975, **97**, 2497 (*struct, ir, ms, pmr, cmr*)

Quinomycin E, 8CI **Q-63**

[14222-15-2]

As Quinomycin A, Q-58 with

R = -CH(CH₃)CH₂CH₃, R' = CH(CH₃)CH(CH₃)₂

C₅₄H₇₀N₁₂O₁₂S₂ 1143.352


Depsideptide antibiotic. Mixt. of two components in 4:1 ratio. Struct. of major component shown; minor component has R and R' interchanged. Isol. from *Streptomyces aureus*. Powder. Sol. Me₂CO, CHCl₃; poorly sol. MeOH, H₂O, hexane. Mp 215-219° dec. λ_{max} 243 (ε 66000); 320 (ε 12500) (MeOH) (Berdy). Otsuka, M. *et al.*, *Tetrahedron*, 1967, **23**, 1535 (*isol, ir, uv*)

Dell, A. *et al.*, *J.A.C.S.*, 1975, **97**, 2497 (*struct, ir, ms, pmr, cmr*)

Shoji, J. *et al.*, *J. Antibiot.*, 1976, **29**, 1246 (*struct, cmr*)

Quinoxapeptin A **Q-64**

[175484-07-8]



Absolute Configuration

C₆₈H₈₄N₁₆O₂₂ 1477.506

Depsideptide antibiotic. Prod. by actinomycete sp. MA7095 (ATCC 55599). Inhibitor of HIV-1 and -2 reverse transcriptase. Cytotoxic agent. Amorph. solid. [α]_D²⁰ +11 (c, 0.09 in CH₂Cl₂). Similar to Luzopeptin C, L-312.

4-Deacyl, 4-Ac: **Quinoxapeptin B**

[175484-08-9]

C₆₅H₈₀N₁₆O₂₂ 1437.441

Prod. by actinomycete sp. MA7095 (ATCC 55599). Inhibitor of HIV-1 and -2 reverse transcriptase. Cytotoxic agent. Amorph. solid. [α]_D²⁰ +36 (c, 0.01 in CH₂Cl₂).


Lingham, R.S. *et al.*, *J. Antibiot.*, 1996, **49**, 253-259 (*isol, props*)

Boger, D.L. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 2424-2426 (*synth, abs config*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 11375-11383 (*synth*)

Racemigerine

Methyl 7-methyl-5H-2-pyridine-4-carboxylate, 9CI
[76655-39-5]




C₁₁H₁₁NO₂ 189.213
Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae). Prisms (CH₂Cl₂). Mp 108°.

6,7-Epoxyde: 6,7-Epoxyracemigerine
[99339-51-2]

C₁₁H₁₁NO₃ 205.213
Alkaloid from *Scaevola racemigera* (Goodeniaceae). Noncryst. Opt. inactive (racemic).

Skaltsounis, A.-L. et al., *Helv. Chim. Acta*, 1985, **68**, 1679 (isol, uv, ir, pmr, ms, struct, epoxide)


Racemulosine



C₂₂H₃₂N₂O₃ 372.506
Alkaloid from *Aconitum racemulosum* Franch var. *pengzhouense*. Cryst. (CHCl₃/MeOH). Mp 228-230°. [α]_D¹⁷ -19.2 (c, 0.5 in CHCl₃/MeOH).

Wang, F.-P. et al., *Tetrahedron*, 2000, **56**, 7443-7446

Racemulotine




C₃₅H₃₇NO₇ 583.68
Alkaloid from the whole plant of *Aconitum racemulosum* var. *pengzhouense*. Amorph. powder. [α]_D¹⁷ -34.4 (c, 0.2 in CHCl₃). Not in CA.

Peng, C.-S. et al., *J. Asian Nat. Prod. Res.*, 2001, **3**, 49-54 (isol, pmr, cmr)

Rachelmycin

CC 1065. ML 1065. NSC 298223. Antibiotic CC 1065
[69866-21-3]
[110352-07-3 (enantiomer)]

R-1



C₃₇H₃₃N₇O₈ 703.71
Obt. from terrestrial *Streptomyces zelessis* and *Streptomyces canulus*. Antineoplastic antibiotic. One of the most potent antineoplastic agents known. Alkylates B-DNA minor groove sites. Clustered needles or amorph. amber-coloured foam. [α]_D²⁵ +98 (c, 0.2 in DMF). λ_{max} 233 (ε 23100); 257 (ε 19500); 300; 362 (MeOH/HCl) (Derep). λ_{max} 230 (sh); 278; 375 (H₂O/pH 8) (Derep). λ_{max} 232 (sh) (ε 21400); 258 (sh) (ε 18700); 371 (ε 27800) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 0.009 mg/kg. LD₅₀ (mus, ipr) 0.007 mg/kg. Very toxic by intravenous and intraperitoneal route. Mutagenic props. DF4936780

N-De(aminocarbonyl), N-Ac: *Gilvusmycin*

[195052-09-6]
C₃₈H₃₄N₆O₈ 702.722
Prod. by *Streptomyces* sp. QM16. Antitumour agent. Yellow powder. Mp 160-162° dec. [α]_D²⁰ +85 (c, 0.1 in DMF). Stereochem. not confirmed. λ_{max} 276 (ε 27500); 370 (ε 33300) (MeOH).

Hanka, L.J. et al., *J. Antibiot.*, 1978, **31**, 1211-1217 (isol)

Martin, D.G. et al., *J. Antibiot.*, 1980, **33**, 902-903; 1981, **34**, 1119-1125 (isol, struct)

Chidester, C.G. et al., *J.A.C.S.*, 1981, **103**, 7629-7635 (cryst struct)

Frolova, V.I. et al., *Antibiotiki (Moscow)*, 1982, **27**, 483-487; *CA*, **97**, 123511 (CC 1065, isol, uv, pmr, cmr)

Sundberg, R.I. et al., *J.O.C.*, 1985, **50**, 425-432 (bibl)

Wierenga, W. et al., *Adv. Enzyme Regul.*, 1986, **25**, 141-155 (rev, pharmacol)

Reynolds, V.L. et al., *J. Antibiot.*, 1986, **39**, 319-334 (rev, pharmacol)

Rawal, V.H. et al., *Heterocycles*, 1987, **25**, 701-728 (rev, synth)

Kelly, R.C. et al., *J.A.C.S.*, 1987, **109**, 6837-6838 (synth)

Watt, W. et al., *Acta Cryst. C*, 1988, **44**, 1675-1677 (cryst struct)

Bolton, R.E. et al., *J.C.S. Perkin 1*, 1988, 2491-2499 (synth, bibl)

Lee, C.-S. et al., *Chem. Res. Toxicol.*, 1991, **4**, 203-213 (pharmacol)

Hurley, L.H. et al., *J. Mol. Recognit.*, 1994, **7**, 123-132 (rev)

Tietze, L. et al., *J.O.C.*, 1994, **59**, 192-196 (synth, bibl)

Linseman, D.A. et al., *Nat. Toxins*, 1995, **3**, 32-40 (pharmacol, enantiomers)

Boger, D.L. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1995, **92**, 3642-3649 (pharmacol, rev)


Boger, D.L. et al., *Chem. Rev.*, 1997, **97**, 787-828 (rev, synth)

Tokoro, Y. et al., *J. Antibiot.*, 1999, **52**, 263-268 (*Gilvusmycin*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, APT375

Radamycin

R-5




C₄₈H₄₇N₁₅O₁₁S₃ 1106.191
Related to Sulfomycin, S-638. Prod. by *Streptomyces* sp. RSP9. Pale yellow powder. [α]_D²⁵ -274.2 (c, 0.58 in CHCl₃). λ_{max} 250 (MeOH).

Holgado, G.G. et al., *J. Antibiot.*, 2002, **55**, 383-390; 391-395 (isol, pmr, cmr)

Radarin B

R-6

[138629-07-9]



C₂₈H₄₁NO₂ 423.637
Indole diterpenoid antibiotic. Prod. by *Aspergillus sulphureus*. Possesses anti-insectant and cytotoxic props. Yellow solid. Mp 115-118° dec. [α]_D +39.4 (c, 0.003 in CHCl₃). λ_{max} 210 (ε 6220); 226 (ε 6990); 290 (ε 1510) (MeOH) (Derep).

6-Deoxy: *Radarin D*

[138606-37-8]
C₂₈H₄₁NO 407.638
Prod. by *Aspergillus sulphureus*. Yellow oil. [α]_D +31.8 (c, 0.003 in CHCl₃). λ_{max} 210 (ε 6220); 226 (ε 6990); 290 (ε 1510) (MeOH) (Derep).

7'-Ketone: *Radarin A*

[138606-35-6]
C₂₈H₃₉NO₂ 421.622
Prod. by *Aspergillus sulphureus*. Pink oil. [α]_D +11.1 (c, 0.005 in CHCl₃). λ_{max} 210 (ε 6220); 226 (ε 6990); 290 (ε 1510) (MeOH) (Derep).

6-Deoxy, 7'-ketone: *Radarin C*

[138606-36-7]
C₂₈H₃₉NO 405.622
Prod. by *Aspergillus sulphureus*. Yellow oil. [α]_D +6.7 (c, 0.002 in CHCl₃). λ_{max} 210 (ε 6220); 226 (ε 6990); 290 (ε 1510) (MeOH) (Derep).

Laakso, J.A. et al., *J.O.C.*, 1992, **57**, 138 (isol, pmr, cmr, struct)

Raddeamine

R-7

C₂₃H₃₇NO₂ 359.551

Prob. a steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria raddeana* (Liliaceae). Cryst. Mp 271-272°. Occurs with Alvanidine, A-665 and Alvanine, A-666.


Hydrochloride: Mp 235-236°.

Aslanov, Kh.A. et al., *Zh. Obshch. Khim.*, 1956, **26**, 579-583; *J. Gen. Chem. USSR (Engl. Transl.)*, 1956, **26**, 623-627 (isol)

Raddeanamine

R-8

3',4',6,8-Tetrahydro-6',7'-dimethoxy-2',6-dimethylspiro[7H-indeno[4,5-d]-1,3-dioxole-7,1'(2'H)-isoquinolin]-6-ol, 9CI [59614-35-6]



Absolute Configuration

C₂₂H₂₅NO₅ 383.443


Alkaloid from *Corydalis ochotensis* var. *raddeana* (Papaveraceae). Gum. [α]_D²⁰ +166 (c, 0.68 in MeOH).

Kametani, T. et al., *J.C.S. Perkin 1*, 1977, 390 (isol, ir, pmr, ms, struct)

Raddeanine†

R-9

3',4',6,8-Tetrahydro-6',7'-dimethoxy-2'-methylspiro[7H-indeno[4,5-d]-1,3-dioxole-7,1'(2'H)-isoquinoline]-6,8-diol, 9CI. 13-Epiyenusomine [59654-07-8]



(+)-form

C₂₁H₂₃NO₆ 385.416**(+)-form**

Alkaloid from *Corydalis ochotensis* var. *raddeana* and *Corydalis govaniana* (Papaveraceae). Prisms (Me₂CO). Mp 208-209° (200-202°). [α]_D +79.4 (c, 0.11 in MeOH).

8-Ac: Raddeanidine

[59614-36-7]

C₂₃H₂₅NO₇ 427.453

Alkaloid from *Corydalis ochotensis* var. *raddeana* (Papaveraceae). Gum. [α]_D²⁰ +82.7 (c, 0.52 in MeOH).

2-O-De-Me: Ledeboridine. Ledebouridine [64191-01-1]C₂₀H₂₁NO₆ 371.389

Alkaloid from *Corydalis ledebouriana* (Papaveraceae). Mp 140-141°. [α]_D +114 (c, 0.28 in MeOH).

13-Ketone: Raddeanone

[59654-08-9]

C₂₁H₂₁NO₆ 383.4

Alkaloid from *Corydalis ochotensis* var. *raddeana* (Papaveraceae). Prisms (Me₂CO). Mp 132-133° Mp 168-170° Mp 181-182°.

13-Ketone, 2-O-de-Me: Ledeborine. Ledebourine. Ledebourine

[56816-35-4]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Corydalis ledebouriana* (Papaveraceae). Cryst. (CHCl₃). Mp 184-186°. Opt. rotn. not reported. Struct. may be erroneous: see *Alkaloids (N.Y.)*, 1979, **17**, 502.

8-Epimer, 13-ketone: Yenusomidine. O-Methylcorpaine

[60102-30-9]

C₂₁H₂₁NO₆ 383.4

Alkaloid from *Corydalis vaginans* (Papaveraceae). Cryst. (MeOH). Mp 220-221°. [α]_D²² -36.7 (c, 0.44 in CHCl₃).

8-Epimer, 13-ketone, 2-O-de-Me: Corpaine

[31002-20-7]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Corydalis paczoskii* (Papaveraceae). Cryst. (EtOH). Mp 204°. No opt. rotn. reported. A mixt. containing Corpaine was known as Coryline.

13-Epimer: Yenusomine

[56435-41-7]

C₂₁H₂₃NO₆ 385.416

Alkaloid from *Corydalis ochotensis* (Papaveraceae). Prisms + 1H₂O (Me₂CO). Mp 127-128° dec. [α]_D¹⁹ +48 (c, 1.0 in MeOH).

(±)-form [64234-40-8]

Alkaloid from *Corydalis ledebouriana* (Papaveraceae). Cryst. (MeOH/CHCl₃). Mp 219-220° Mp 201-202° (synthetic). O⁸-Ac: Synthetic. Mp 169-171°.

13-Ketone, 2-O-de-Me: Corysolidine

[105926-86-1]

C₂₀H₁₉NO₆ 369.373

Alkaloid from the whole plants of *Corydalis ochotensis* and *Corydalis solida*. Amorph. Racemate.

13-Ketone, 2-O-de-Me, 8-Ac: 8-O-Acetylcorpaine

[314729-31-2]

C₂₂H₂₁NO₇ 411.41

Alkaloid from the aerial parts of *Corydalis ochotensis*.

8-Epimer, 13-ketone: (±)-YenusomidineC₂₁H₂₁NO₆ 383.4

Alkaloid from *Corydalis ochotensis* (Papaveraceae). Prisms (Me₂CO). Mp 242-243° Mp 228-229° (synthetic).

8-Epimer, 13-ketone, picrate:

Yellow prisms (Me₂CO). Mp 214-215° dec.

8-Epimer, 13-ketone, Ac:

Yellow-brown prisms (MeOH). Mp 173-174°.

13-Epimer: [60761-13-9]

Synthetic. Cryst. (MeOH). Mp 223-225°.

Baishева, Kh.S. et al., *Khim. Prir. Soedin.*,

1970, **6**, 574; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 590 (Corpaine)

Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 268; 1977, **13**, 428; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 284; 1977, **13**, 366 (Raddeamine, Ledeborine, Ledeboridine)

Irie, H. et al., *Heterocycles*, 1976, **4**, 1083 (synth)

Lu, S.T. et al., *J.C.S. Perkin 1*, 1976, 63 (isol, cd, uv, ir, pmr, ms, struct)

Margelashvili, N.N. et al., *Khim. Prir. Soedin.*, 1976, **12**, 123 (O-Methylcorpaine)

Nalliah, B.C. et al., *Can. J. Chem.*, 1977, **55**, 922; 1979, **57**, 1545 (synth)

McLean, S. et al., *Can. J. Chem.*, 1977, **55**, 924 (synth)

Hughes, D.W. et al., *Can. J. Chem.*, 1977, **55**, 3304 (nmr)

Kametani, T. et al., *J.C.S. Perkin 1*, 1977, 390 (Raddeanine, Raddeanone, isol, ir, pmr, ms, struct)

Dime, D. et al., *Can. J. Chem.*, 1979, **57**, 1569 (Corpaine, synth)

Rahimizadeh, M. et al., *Phytochemistry*, 1986, **25**, 2245-2246 (Corysolidine)

Mukhopadhyay, S. et al., *J. Nat. Prod.*, 1987, **50**, 270 (isol, uv, ir, pmr, cmr, ms)

Hanaoka, M. et al., *Chem. Pharm. Bull.*, 1988, **36**, 4248 (Raddeanine, Raddeanidine, Yenusomine, Yenusomidine, synth, ir, pmr)


Kessar, S.V. et al., *Chem. Comm.*, 1994, 1327 (synth)

Kim, D.K. et al., *Arch. Pharmacol. Res.*, 2000, **23**, 459-460 (8-O-Acetylcorysolidine)

Radicamine A

R-10

3,4-Dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-5-hydroxymethylpyrrolidine



(+)-form

C₁₂H₁₇NO₅ 255.27**(+)-form [431981-74-7]**

Alkaloid from *Lobelia sinensis*. Pale yellow oil. [α]_D²⁵ +43.7 (c, 0.13 in H₂O). Related to Codonopsine, C-557. λ_{max} 228 (log ε 3.64); 281 (log ε 3.94) (no solvent reported).

3'-Deoxy, O-de-Me: Radicamine B

[431981-75-8]

C₁₁H₁₅NO₄ 225.244

Alkaloid from *Lobelia sinensis*. Pale yellow oil. [α]_D²⁵ +72 (c, 0.1 in H₂O). λ_{max} 227 (log ε 4.19); 277 (log ε 3.6) (no solvent reported).

(-)-form**O^{3'},N-Di-Me: Codonopsinol**C₁₄H₂₁NO₅ 283.324

Alkaloid from *Codonopsis clematidea*. Amorph. powder. [α]_D -3.5 (c, 0.2 in MeOH).

Shibano, M. et al., *Chem. Pharm. Bull.*, 2001, **49**, 1362-1365 (isol)

Shibano, M. et al., *Heterocycles*, 2002, **57**, 1539-1553 (biosynth)


Yu, C.-Y. et al., *Org. Lett.*, 2006, **8**, 3021-3024 (synth, abs config)

Gurjar, M.K. et al., *Tet. Lett.*, 2006, **47**, 6979-6981 (Radicamine B, synth, abs config)

Liu, C. *et al.*, *Lett. Org. Chem.*, 2007, **4**, 556-558 (*Radicalamine B, synth*)
 Merino, P. *et al.*, *Eur. J. Org. Chem.*, 2008, 2929-2947 (*Radicalamine B, synth*)
 Ishida, S. *et al.*, *J. Nat. Med. (Tokyo)*, 2008, **62**, 236-238 (*Codonopsinol*)
 Ribes, C. *et al.*, *J.O.C.*, 2008, **73**, 7779-7782 (*Radicalamine B, synth*)

Radixisatidis A
Banlangenjiasu
 [221181-27-7]

R-11




$C_{21}H_{23}NO_6$ 385.416
 Constit. of *Radix Strobilanthes (Strobilanthes sp.)* Component of Ban Lan Gen.

Wang, S. *et al.*, *CA*, 1999, **130**, 220357q

Radpetine

[139751-06-7]

R-12



$C_{29}H_{45}NO_3$ 455.679
 Alkaloid from epigeal parts of *Petilium raddeana* (Liliaceae). Cryst. (Me_2CO /petrol). Mp 229-231°.

Ac:


Cryst. Mp 100-103°.

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1991, 384; *Chem. Nat. Compd. (Engl. Transl.)*, 332 (*isol, ir, pmr, ms, cryst struct*)

Rakicidin C

[310440-15-4]

R-13

 $C_{26}H_{42}N_4O_6$ 506.641


Depsideptide antibiotic. Prod. by *Streptomyces sp.* GT61042. Powder. $[\alpha]_D^{25} +90.9$ (c, 0.24 in DMSO). λ_{max} 201 (log ϵ 3.73); 243 (log ϵ 1.71) (MeOH).

Hu, J.-F. *et al.*, *Eur. J. Org. Chem.*, 2000, 3353-3356 (*isol, pmr, cmr*)

Ramosine A

[530092-39-8]

R-14

 $C_{27}H_{40}N_4O_4$ 484.637


Alkaloid from the roots of *Paliurus ramosissimus*. Amorph. solid. Mp 55-56°. $[\alpha]_D^{26} -125$ (c, 0.76 in MeOH). λ_{max} 221 (log ϵ 4.52) (MeOH).

Lin, H.-Y. *et al.*, *Helv. Chim. Acta*, 2003, **86**, 127-138 (*isol, cd, pmr, cmr*)

Rankiniridine

[1036975-00-4]

R-15

 $C_{30}H_{36}N_2O_8$ 552.623

Related to Humantenirine, H-367. Alkaloid from *Gelsemium rankinii*. Amorph. powder. $[\alpha]_D^{21} -270.5$ (c, 0.14 in MeOH). λ_{max} 208 (log ϵ 4.18); 296 (log ϵ 4.25) (MeOH).

11-Methoxy, 7'-deoxy: Humanteniridine
 [1036975-02-6]

 $C_{31}H_{38}N_2O_8$ 566.65


Alkaloid from the leaves of *Gelsemium elegans*. Amorph. yellow powder. $[\alpha]_D^{15} -409.4$ (c, 0.26 in MeOH). λ_{max} 215 (log ϵ 4.51); 293 (log ϵ 4.35) (MeOH).

Kogure, N. *et al.*, *Tet. Lett.*, 2008, **49**, 3638-3642 (*isol, cd, pmr, cmr*)

Raocyclamide A

[174846-65-2]

R-16



Absolute
 Configuration

 $C_{27}H_{30}N_6O_5S$ 550.637

Stereochem. revised in 1998. Cyclic hexapeptide from the terrestrial cyanobacterium *Oscillatoria raoi*. Exhibits moderate cytotoxicity against sea urchin embryos. Glassy solid. $[\alpha]_D^{25} +25.5$ (c, 4.4 in MeOH). λ_{max} 242 (ϵ 17750) (MeOH).


Admi, V. *et al.*, *J. Nat. Prod.*, 1996, **59**, 396 (*isol, uv, ir, pmr, cmr, ms, struct*)

Freeman, D.J. *et al.*, *Tet. Lett.*, 1998, **39**, 3251-3254 (*synth, abs config*)

Raocyclamide B

[174846-66-3]

R-17



Absolute
 Configuration

 $C_{27}H_{32}N_6O_6S$ 568.652

Stereochem. revised in 1998. Cyclic hexapeptide from the terrestrial cyanobacterium *Oscillatoria raoi*. Glassy solid. $[\alpha]_D^{25} +5.1$ (c, 4.1 in MeOH). λ_{max} 241 (ϵ 17166) (MeOH).


Admi, V. *et al.*, *J. Nat. Prod.*, 1996, **59**, 396 (*isol, uv, ir, pmr, cmr, ms, struct*)

Freeman, D.J. *et al.*, *Tet. Lett.*, 1998, **39**, 3251-3254 (*synth, abs config*)

Raphanusanin

3-[(Methylthio)methylene]-2-pyrrolidinethione, 9CI. Raphantin

R-18



(E) - form

C₆H₉NS₂ 159.276

There have been various struct. revisions. The most recent incorrect struct. proposed was a MeOH covalent solvate of the struct. shown. The names Raphanusanin A and Raphanusanin B were erroneously applied to stereoisomers of this solvate.

(E)-form [128463-44-5]
[104760-73-8]

Isol. from *Raphanus sativus* var. *hortensis* f. *gigantissimus* (Brassicaceae). Growth inhibitor involved in phototropism of hypocotyls. Needles (MeOH).

(Z)-form [148225-27-8]
[104760-72-7]

Isol. from *Raphanus sativus* var. *hortensis* f. *gigantissimus* (Brassicaceae). Growth inhibitor involved in phototropism of hypocotyls. Amorph. pale yellow powder. Sakoda, M. *et al.*, *Phytochemistry*, 1990, **29**, 1031 (*isol*)
Harada, N. *et al.*, *Tet. Lett.*, 1991, **32**, 6757 (*uv, ir, pmr, cmr, ms, bibl*)
Kosemura, S. *et al.*, *Tet. Lett.*, 1993, **34**, 481-484 (*struct*)
Matsuoka, H. *et al.*, *Phytochemistry*, 1998, **47**, 975-977 (*isol*)
Hasegawa, T. *et al.*, *Phytochemistry*, 2000, **54**, 275-279 (*biosynth*)

Raucaffridine

R-19

[107527-78-6]


C₂₁H₂₄N₂O₃ 352.432

Struct. unknown. Alkaloid from *Rauwolfia caffra* (Apocynaceae). Mp 221° dec.

Khan, N.H. *et al.*, *Pak. J. Sci. Ind. Res.*, 1965, **8**, 23-27

Raucanine

R-20



Absolute Configuration

C₂₂H₂₈N₂O₄ 384.474


The struct. is badly drawn in the paper but this appears to be the relative stereochemistry. Alkaloid from *Rauwolfia canescens*. Mp 183-185°. [α]_D⁻²⁷ (CHCl₃). CAS no. not found 8-14CI. λ_{max} 251 ; 295 (no solvent reported).

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 761-863; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 737-858 (*rev*)

Raucubaine

R-21

1,2,19,20-Tetrahydro-19,20-dihydroxy-1-methylakuummilian-17-oic acid γ-lactone, 9CI. *Quaternoline* [75418-95-0]



Absolute Configuration

C₂₀H₂₄N₂O₃ 340.421

Identity of Raucubaine and Quaternoline not certain, may be stereoisomers. Alkaloid from the leaves of *Rauwolfia salicifolia* (Apocynaceae). Cryst. (MeOH). Mp 224°. [α]_D²⁰ -18 (CHCl₃). Poss. artifact, also obt. by acid treatment of Cathafoline, C-191.

11-Methoxy: **Caberoline**

[56283-51-3]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Cabucala* spp. and *Alstonia plumosa* (Apocynaceae). Poss. artifact; also obt. by acid treatment of Caberine.

[57498-99-4]

Mamatas-Kalamaras, S. *et al.*, *Phytochemistry*, 1975, **14**, 1849 (*Quaternoline*)

Kutney, J.P. *et al.*, *Heterocycles*, 1980, **14**, 1309 (*uv, ir, pmr, cd, ms, cryst struct*)

Paupit, R.A. *et al.*, *Can. J. Chem.*, 1981, **59**, 1007 (*cryst struct*)

Jacquier, M.J. *et al.*, *Phytochemistry*, 1982, **21**, 2973 (*isol*)


Sierra, P. *et al.*, *Coll. Czech. Chem. Comm.*, 1982, **47**, 2912 (*isol, spectra, abs config*)

Raumacine

R-22

4-Demethyl-20,21-dihydroalstophyllan-17-ol, 9CI

[132923-02-5]



C₂₀H₂₆N₂O₂ 326.438

Alkaloid from cell cultures of *Rauwolfia serpentina* (Apocynaceae). Mp 197-200°. [α]_D²⁶ -26.43 (c, 0.28 in CHCl₃) (*synthetic*).

N^b-Me: **N^b-Methylraumacline**

[132943-65-8]

C₂₁H₂₈N₂O₂ 340.464

Alkaloid from cell cultures of *Rauwolfia serpentina* (Apocynaceae). Amorph. solid. [α]_D²⁵ -67.5 (c, 0.16 in CHCl₃) (*synthetic*).

6α-Hydroxy: **6α-Hydroxyraumacline**

[145051-93-0]

C₂₀H₂₆N₂O₃ 342.437

Alkaloid from cell cultures of *Rauwolfia serpentina* (Apocynaceae). Needles (Me₂CO/Et₂O). Mp 199-200°.

10-Hydroxy, 19,20-didehydro, N-de-Me: **10-Hydroxy-N-demethyl-19,20-dehydro-raumacline**

C₁₉H₂₂N₂O₃ 326.394

Alkaloid from hairy root cultures of *Rauwolfia serpentina*. Powder. [α]_D²⁵ +29.6 (c, 0.5 in MeOH). Mp >300°. Isol. as a mixt. of *E*- and *Z*-isomers. λ_{max} 226 (log ε 4.12); 279 (log ε 3.65) (MeOH).

19*S*-Hydroxy, N^b-Me: **19(S)-Hydroxy-**

N^b-methylraumacline

[141810-20-0]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from cell suspension cultures of *Rauwolfia serpentina* following addition of Ajmaline, A-216. Amorph. powder. [α]_D²⁵ -86.7 (c, 0.18 in CHCl₃).

20-Epimer: **Isoramacline**

[149507-97-1]

C₂₀H₂₆N₂O₂ 326.438

Alkaloid from cell cultures of *Rauwolfia serpentina*. [α]_D²⁰ +15.6 (c, 0.39 in CHCl₃). λ_{max} 228 ; 284 ; 293 (MeOH).

Polz, L. *et al.*, *Tet. Lett.*, 1990, **31**, 6693 (*isol, uv, cd, pmr, cmr, struct*)

Endress, S. *et al.*, *Planta Med.*, 1992, **58**, 410 (*6α-Hydroxyraumacline*)

Takayama, H. *et al.*, *Tetrahedron*, 1992, **48**, 2627 (*19(S)-Hydroxy-N^b-methylraumacline*)

Endress, S. *et al.*, *Helv. Chim. Acta*, 1993, **76**, 2544 (*synth*)

Fu, X. *et al.*, *J.O.C.*, 1993, **58**, 661 (*synth*)

Endress, S. *et al.*, *Phytochemistry*, 1993, **32**, 725-730 (*Isoramacline*)

Bailey, P.D. *et al.*, *J.C.S. Perkin 1*, 1997, 1209 (*synth*)

Sheludko, Y. *et al.*, *Planta Med.*, 2002, **68**, 435-438 (*19,20-Didehydro-N-demethyl-10-hydroxyraumacline*)

Bailey, P.D. *et al.*, *Tet. Lett.*, 2008, **49**, 2150-2153 (*synth*)

Raunamine

R-23

[1360-77-6]

C₂₃H₃₀N₂O₄ 398.501

Struct. unknown. Alkaloid from the root bark of *Rauwolfia micrantha* (Apocynaceae). Needles (EtOH). Mp 206-207°. [α]_D¹⁷ +60.1 (c, 0.189 in EtOH).

Hydrochloride:

Needles (EtOH). Mp 236-238°.

Picrate:


Yellow microcryst. needles (EtOH aq.). Mp 190-192° dec.

Pillay, P.P. *et al.*, *J. Sci. Ind. Res., Sect. B*, 1960, **19**, 135-137 (*isol, uv, ir*)

Raureflexine

R-24

[226994-19-0]



Absolute Configuration


C₄₃H₄₈N₄O₄ 684.877

Alkaloid from *Rauwolfia reflexa*. Amorph. solid. [α]_D²⁵ +328.2 (c, 1 in CHCl₃). Dec. at 217-218°. λ_{max} 213 (log ε 4.76); 256 (log ε 4.19); 300 (log ε 4) (EtOH).


Chatterjee, A. *et al.*, *J. Indian Chem. Soc.*, 1998, **75**, 695-697

Rausutranine


[161068-56-0]

C₄₁H₄₆N₄O₇ 706.837Alkaloid from leaves of *Rauwolfia sumatrana* (Apocynaceae). Dark red amorph. powder.Subhadhirasakul, S. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1427-1431 (*isol*, *uv*, *cd*, *pmr*, *cmr*, *ms*, *struct*)**Rausutrine**

[161068-57-1]




Absolute Configuration

C₄₂H₄₈N₄O₇ 720.864Alkaloid from leaves of *Rauwolfia sumatrana* (Apocynaceae). Dark red amorph. powder.Subhadhirasakul, S. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1427-1431 (*isol*, *uv*, *cd*, *pmr*, *cmr*, *ms*, *struct*)**Rauviridine***Vobasine methine*
[57567-12-1]C₂₂H₂₆N₂O₃ 366.459Alkaloid from the stem bark of *Rauwolfia viridis* (Apocynaceae). Needles (Et₂O/petrol). Mp 145-146°. [α]_D -103.7 (c, 0.9 in CHCl₃). Data given relates to synthetic prod.*Epimer: Epirauviridine*
[128112-82-3]

R-25

C₂₂H₂₆N₂O₃ 366.459Alkaloid from the stem bark of *Rauwolfia viridis* (Apocynaceae). Config. not determined.Renner, U. *et al.*, *Helv. Chim. Acta*, 1963, **46**,2186-2208 (*synth*)Knox, J.R. *et al.*, *Aust. J. Chem.*, 1975, **28**,1843 (*synth*)Martinez, P.J.A. *et al.*, *Rev. Cubana Farm.*,1989, **23**, 117; 129; *CA*, **113**, 37702; 41096 (*isol*)**Rauvolcinine**

[78853-47-1]

C₂₂H₂₈N₂O₅ 400.474Alkaloid from the roots of *Rauwolfia volkensii* (Apocynaceae).*Stereoisomer: Cabucinine*

[38839-94-0]

C₂₂H₂₈N₂O₅ 400.474Alkaloid from *Cabucala madagascariensis*, *Cabucala erythrocarpa*, *Cabucala fasciculata* and *Cabucala striolata* (Apocynaceae). Mp 170°. [α]_D 0 (CHCl₃). Epimeric with Rauvolcinine at C-20; C-16 and C-17 configs. undetd.Douzoua, L.L. *et al.*, *Ann. Pharm. Fr.*,1972, **30**, 199-204; *CA*, **77**, 72560v

(Cabucinine)


Akinloye, B.A. *et al.*, *J. Ethnopharmacol.*,1981, **4**, 99-109; *CA*, **95**, 111700j

(Rauvolcinine)

R-28


Ravencic acid

R-30


3-(1-Hydroxy-4-methyl-2,4,6,8-decateranylidene)-2,4-pyrrolidinedione, 9CIC₁₅H₁₇NO₃ 259.304Tetramic acid. Prod. by *Penicillium* sp. MINAP9902. Orange-yellow solid.λ_{max} 240 (ε 9900); 366 (ε 11000) (EtOH).Michael, A.P. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1360-1362 (*isol*, *uv*, *pmr*, *cmr*, *ms*)**Ravenoline**

R-31

[20105-23-1]

C₁₅H₁₇NO₂ 243.305**(+)-form**Alkaloid from *Ravenia spectabilis* (Rutaceae). Mp 144°. [α]_D²⁵ +6 (CHCl₃).**(-)-form**From *Ravenia spectabilis*. [α]_D -5 (CHCl₃). The question of the *isol.* of 2 enants. from the same sp. has not been resolved.**(±)-form***2',3'-Dihydro, 2'-hydroxy: 4-Hydroxy-3-(2-hydroxy-1,2-dimethylpropyl)-1-methyl-2(1H)quinolinone. Paraensine†*
[33172-19-9]C₁₅H₁₉NO₃ 261.32Alkaloid from the heartwood of *Euxylophora paraënsis* (Rutaceae). Needles (CHCl₃ or Me₂CO/MeOH). Mp 217-218°. [α]_D²⁵ -0.54 (c, 1.0 in CHCl₃).Paul, B.D. *et al.*, *Indian J. Chem.*, 1969, **7**, 678 (*ir*, *uv*, *pmr*, *struct*, *synth*)Talapatra, S.K. *et al.*, *Tet. Lett.*, 1971, 2683-6 (*isol*)Jurd, L. *et al.*, *Aust. J. Chem.*, 1983, **36**, 759 (*Paraensine*)**Rauwolfinine***Perakenine*

R-29



Probable Structure


C₂₀H₂₆N₂O₂ 326.438Alkaloid from the root bark of *Rauwolfia serpentina* (Apocynaceae). Plates (EtOAc). Mp 235-236° dec. [α]_D²² -34.7 (EtOH).*Hydrochloride*: Mp 195° (dec.).*Hydrobromide*:

Cryst. (EtOH). Mp 280-281° (dec.).

Bose, S. *et al.*, *J. Indian Chem. Soc.*, 1954, **31**, 47; 311; 691; 1958, **35**, 72 (*isol*, *uv*, *ir*, *derivs*, *struct*)

Rebeccamycin**R-32**

1,11-Dichloro-12,13-dihydro-12-(4-O-methyl-β-D-glucopyranosyl)-5H-indolo[2,3-a]pyrrolo[3,4-c]carbazole-5,7(6H)-dione, 9CI
[93908-02-2]



$C_{27}H_{21}Cl_2N_3O_7$ 570.384

Isol. from *Saccharothrix aerocolonigenes* ATCC 39243. Antineoplastic agent. Yellow solid. Sol. THF, DMSO; fairly sol. Me_2CO , $CHCl_3/MeOH$; poorly sol. H_2O , hexane, MeOH. Mp 326-330° dec. $[\alpha]_D^{21} +131$ (THF). $[\alpha]_D^{20} +143$ (c, 1 in THF) (synthetic). Log P 3.69 (calc). λ_{max} 238 (€ 43130); 256 (sh) (€); 293 (€ 31466); 314 (€ 51594); 362 (€ 4780); 390 (€ 4495) (MeOH) (Derep). λ_{max} 237; 285; 314 (MeOH/NaOH) (Berdy).

▶ **NM3455000**

6-N-[2-(Diethylamino)ethyl]: **Becatecarin**, INN, USAN. BMY 27557. BMS 181176. NSC 655649. XL 119 [119673-08-4]

[127099-93-8, 119673-09-5]

$C_{33}H_{34}Cl_2N_4O_7$ 669.56

Antineoplastic agent. Water-soluble Rebeccamycin deriv. active against P388 leukaemia and B16 melanoma. Sol. H_2O .

Aglycone: [118458-56-3]

Golden-yellow needles (DMF). Mp 360°.

Dibromo analogue: **Bromorebeccamycin**

[137605-02-8]

$C_{27}H_{21}Br_2N_3O_7$ 659.287

Prod. by *Saccharothrix aerocolonigenes* in a medium containing 0.5% KBr. Antineoplastic agent. Bright yellow amorph. solid. Sol. DMSO, DMF, THF; fairly sol. Me_2CO ; poorly sol. H_2O . λ_{max} 206 (€ 22500); 238 (€ 27400); 294 (€ 20500); 316 (€ 34500); 390 (€ 3240) (MeOH) (Derep).

11-Dechloro: **11-Deschlororebeccamycin**

[97938-09-5]

$C_{27}H_{22}ClN_3O_7$ 535.939

From *Saccharothrix aerocolonigenes* ATCC 39243. Active against gram-positive and -negative bacteria and tumours. Yellow solid. $[\alpha]_D^{20} +128.1$ (c, 1 in THF) (synthetic). λ_{max} 235; 257 (sh); 290; 315; 400 (MeOH).

Bis(dechloro), 2,10-difluoro: **Fluoroindolocarbazole A**

[138829-50-2]

$C_{27}H_{21}F_2N_3O_7$ 537.476

Prod. by *Saccharothrix aerocolonigenes* ATCC 39243 in a medium containing 6-fluorotryptophan. Antitumour agent. Bright yellow amorph. solid. λ_{max} 228 (€ 780); 256 (€ 454); 280 (€ 387); 316 (€ 1050); 398 (€ 103) (MeOH).

Bis(dechloro), 2,10-difluoro, O-de-Me:

Fluoroindolocarbazole B

[138829-47-7]

$C_{26}H_{19}F_2N_3O_7$ 523.449

Prod. by *Saccharothrix aerocolonigenes* ATCC 39243 in a medium containing 6-fluorotryptophan. Antitumour agent. Bright yellow amorph. solid. λ_{max} 226 (€ 526); 256 (€ 313); 280 (€ 259); 316 (€ 640); 398 (€ 60) (MeOH).

Bis(dechloro), 3,9-difluoro, O-de-Me:

Fluoroindolocarbazole C

[138829-46-6]

$C_{26}H_{19}F_2N_3O_7$ 523.449

Prod. by *Saccharothrix aerocolonigenes* in a medium containing 5-fluorotryptophan. Antitumour agent. Bright yellow amorph. solid. λ_{max} 230 (€ 348); 259; 277; 288; 322 (€ 457); 405 (MeOH).

Nettleton, D.E. *et al.*, *Tet. Lett.*, 1985, **26**,

4011-4014 (*isol, struct, props*)

Kaneko, T. *et al.*, *Tet. Lett.*, 1985, **26**, 4015-

4018 (*synth*)

U.S. Pat., 1985, ((*Bristol-Myers*))4 524 145;

CA, **103**, 159104d (*11-dechloro*)

Bush, J.A. *et al.*, *J. Antibiot.*, 1987, **40**, 668-678

(*manuf, props, bibl*)

Eur. Pat., 1988, ((*Bristol-Myers*))269 025; CA,

110, 135647b (*becatecarin, synth, pharmacol*)

Golik, J. *et al.*, *J. Antibiot.*, 1989, **42**, 1784-

1789 (*cmr*)

Bergman, J. *et al.*, *J.O.C.*, 1989, **54**, 824-828

(*aglycone, synth, pmr*)

Kaneko, T. *et al.*, *J. Antibiot.*, 1990, **43**, 125-

127 (*becatecarin, synth, pharmacol*)

Lam, K.S. *et al.*, *J. Antibiot.*, 1991, **44**, 934-939

(*Bromorebeccamycin*)

Fabre, S. *et al.*, *Bioorg. Med. Chem.*, 1994, **2**,

73-77 (*sar*)

Saulnier, M.G. *et al.*, *Tet. Lett.*, 1995, **36**, 7841-

7844 (*synth*)

Faul, M.M. *et al.*, *J.O.C.*, 1999, **64**, 2465-2470

(*synth, pmr, cmr*)

Lam, K.S. *et al.*, *J. Antibiot.*, 2001, **54**, 1-9

(*Fluoroindolocarbazoles*)

Wang, J. *et al.*, *Tet. Lett.*, 2001, **42**, 8935-8937

(*aglycone, synth*)


Rewcastle, G.W. *et al.*, *IDrugs*, 2005, **8**, 838-

847 (*becatecarin, rev*)

Reductiline

AIS 2M3. Antibiotic AIS 2M3

[85145-25-1]

**R-33**

$C_{16}H_{20}N_2O_3S$ 320.412

Prod. by *Streptomyces orientalis* in the presence of methionine. Shows anticancer, antibacterial and antiviral props. Yellow cryst. (MeOH). Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. Mp 203-204°. λ_{max} 273 (€ 16000); 327 (€ 28700) (MeOH) (Derep). λ_{max} 260 (€ 13300); 305 (€ 12800) (MeOH/NaOH) (Berdy). λ_{max} 330 (€ 13700) (MeOH/HCl) (Berdy).


Japan. Pat., 1982, 82 57 206 676; CA, **99**, 4062

Ojika, M. *et al.*, *Tet. Lett.*, 1982, **23**, 4977

(*struct, synth*)

Reductiomycin**R-34**

3-[5-(Acetyloxy)-4,5-dihydro-3-furanyl]-N-(2-hydroxy-5-oxo-1-cyclopenten-1-yl)-2-propenamide. AM 6201. Antibiotic AM 6201. S 5511I. Antibiotic S 5511I [68748-55-0]



$C_{14}H_{15}NO_6$ 293.276

Identity of Reductiomycin and AM6201 not yet conclusively proven.

Isol. from *Streptomyces orientalis*, *Streptomyces xanthochromogenus* and *Streptomyces griseorubiginosus*. Active against gram-positive bacteria, fungi and Newcastle disease virus. Yellow needles (MeOH). Mp 215° dec Mp 232-233°. λ_{max} 290 (€ 26000) (MeOH/HCl) (Derep). λ_{max} 265 (sh) (€ 25900); 282 (€ 28700); 326 (sh) (€ 15500) (MeOH) (Derep).

▶ **UC6220000**

[79303-68-7, 83377-76-8, 85439-32-3]

Kondo, Y. *et al.*, *J. Antibiot.*, 1981, **34**, 1222

(*isol*)

Shimizu, K. *et al.*, *J. Antibiot.*, 1981, **34**, 649;

654 (*isol*)

Shizuri, Y. *et al.*, *Tet. Lett.*, 1981, **22**, 4291 (*uv, ir, pmr, cmr, ms, struct*)

Ojika, M. *et al.*, *Chem. Comm.*, 1982, 628

(*synth*)

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**,

1210 (*isol, uv, cd, ir, pmr, cmr, struct*)

Donohue, J. *et al.*, *J. Crystallogr. Spectrosc.*

Res., 1984, **14**, 35 (*cryst struct*)

Floss, H.G. *et al.*, *Pure Appl. Chem.*, 1989, **61**,


485 (*rev, biosynth*)

Cho, H. *et al.*, *J.A.C.S.*, 1993, **115**, 12296

(*biosynth*)

Reframidine**R-35**

11,12-Dihydro-14-methyl-11,5-(imino-methano)-5H-cyclohepta[1,2-f:4,5-f']bis[1,3]benzodioxole, 9CI



(-)-form

$C_{19}H_{17}NO_4$ 323.348

(-)-form [21051-82-1]

Alkaloid from *Papaver anomalum* and *Roemeria refracta* (*Roemeria rheadiflora*) (Papaveraceae). Amorph. $[\alpha]_D^{23}$ -123 (c, 0.4 in MeOH).

N-Oxide (R-): (R)-Reframidine N-oxide

[143051-75-6]
C₁₉H₁₇NO₅ 339.347

Alkaloid from *Roemeria refracta* (Papaveraceae). $[\alpha]_D$ -146.1 (c, 0.167 in MeOH).

N-Oxide (S-): (S)-Reframidine N-oxide

[143120-44-9]
C₁₉H₁₇NO₅ 339.347

Alkaloid from *Roemeria refracta* (Papaveraceae). $[\alpha]_D$ -128.2 (c, 0.117 in MeOH).

N-Me: Mp 223-225° (as iodide).

N-De-Me: Norreframidine

[119181-90-7]
C₁₈H₁₅NO₄ 309.321

Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D$ -98 (c, 0.11 in MeOH). First known naturally occurring N-norisopavine.

15-Oxo: Roelactamine. 15-Oxoreframidine

[143070-35-3]
C₁₉H₁₅NO₅ 337.331

Alkaloid from *Roemeria refracta* (Papaveraceae). $[\alpha]_D$ -22 (c, 0.123 in MeOH). First example of a natural isopavine lactam.

15-(2-Oxopropyl): 15-(2-Oxopropyl)reframidine

[143120-82-5]
C₂₂H₂₁NO₅ 379.412

Alkaloid from *Roemeria refracta* (Papaveraceae). $[\alpha]_D$ -100.5 (c, 0.19 in MeOH). Isol. as an inseparable mixt. of 2 epimers which are possibly artifacts.

15-(Carboethoxy)methyl: Ethyl (raframidin-15-yl)acetate

[143120-45-0]
C₂₃H₂₃NO₆ 409.438

Alkaloid from *Roemeria refracta* (Papaveraceae). $[\alpha]_D$ -104.2 (c, 0.19 in MeOH). Isol. as an inseparable mixt. of 2 epimers which are possibly artifacts.

(±)-form [33535-61-4]

Synthetic. Golden oil.

N-Me:

Cryst. (EtOH). Mp 267-269° (as iodide).

[143051-76-7, 143070-36-4]

Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 3917 (ms)

Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 4066 (isol, uv, ir, struct)

Dyke, S.F. *et al.*, *Tetrahedron*, 1971, **27**, 3803 (synth, pmr, ms)

Pfeifer, S. *et al.*, *Pharmazie*, 1972, **27**, 48 (isol, ms)


Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 1614 (synth)

Gözler, B. *et al.*, *J. Nat. Prod.*, 1988, **51**, 760 (Norreframidine)

Gözler, B. *et al.*, *Helv. Chim. Acta*, 1992, **75**, 260 (oxides, Roelactamine)

Reframine

R-36
10,11-Dihydro-7,8-dimethoxy-13-methyl-10,5-(iminomethano)-5H-benzo[4,5]cyclohepta[1,2-f]-1,3-benzodioxole, 9CI [21305-35-1]



C₂₀H₂₁NO₄ 339.39

Biogenetic (benzylisoquinoline-type) numbering shown. Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D^{24}$ -146 (c, 0.5 in MeOH).

N-Me: Remrefine. Roemrefine. Reframine

N-methosalt
[26012-99-7]

[21059-77-8, 21059-78-9]

C₂₁H₂₄NO₄[±] 354.425

Quaternary alkaloid from *Roemeria refracta* (Papaveraceae). Cryst. (EtOH/Me₂CO or MeOH)(as chloride). Mp 258-259° (241-242°)(chloride). $[\alpha]_D$ -188 (c, 0.33 in H₂O) (-147).

O⁶-De-Me: Reframoline

[21238-92-6]

Alkaloid from *Roemeria refracta* (*Roemeria rheadiflora*) (Papaveraceae). Cryst. (Et₂O). Mp 160°. $[\alpha]_D^{20}$ -144 (c, 0.37 in EtOH).

O⁷-De-Me: Refractamine

C₁₉H₁₉NO₄ 325.363

Alkaloid from *Roemeria refracta* (Papaveraceae). Amorph. $[\alpha]_D$ -114 (c, 0.11 in MeOH).

(±)-form [33535-62-5]

Synthetic. Red-brown gum.

N-Me:

Cryst. (EtOH). Mp 272-274°.

Yunusov, M.S. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 68; 1968, **4**, 225; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 58; 1968, **4**, 193 (isol, uv, pmr, struct)

Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 3917 (ms)

Slavík, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 4066 (isol, uv, ir, struct)


Hoshino, O. *et al.*, *Heterocycles*, 1973, **1**, 223 (synth)

Dyke, S.F. *et al.*, *Tetrahedron*, 1978, **34**, 241; 3803 (synth, uv, cd, pmr, ms)

Gözler, B. *et al.*, *J. Nat. Prod.*, 1988, **51**, 760 (Refractamine)

Regecoline

[97682-67-2]



C₁₉H₂₃NO₄ 329.395

Alkaloid from the aerial parts of *Colchicum kesselringii* (Liliaceae). Mp 312-314° dec.

11-Epimer: Isoregecoline

[104972-88-5]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Colchicum kesselringii* (Liliaceae). Mp 284-286°.


Usmanov, A.M. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 248; *Chem. Nat. Compd. (Engl. Transl.)*, 233 (isol, pmr, cmr, ms, struct, synth)

Chommadov, B. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 808; *Chem. Nat. Compd. (Engl. Transl.)*, 767 (Isoregecoline)

Alikulov, R.V. *et al.*, *Khim. Prir. Soedin.*, 1986, **22**, 464; *Chem. Nat. Compd. (Engl. Transl.)*, 433 (synth)

Regelinone

[66855-44-5]



C₁₉H₁₉NO₅ 341.363

Struct. revised in 1985. Alkaloid from *Colchicum kesselringii* (Liliaceae). Mp 316-317°. λ_{max} 248; 300; 395 (no solvent reported).

11-Epimer: Isoregelinone

[79027-71-7]

C₁₉H₁₉NO₅ 341.363

Alkaloid from *Colchicum kesselringii* (Liliaceae). Cryst. (Me₂CO/MeOH). Mp 321-323°. Struct. revised in 1985.

Yusupov, M.K. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 867-868; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 736-737 (isol, uv, pmr, ms)


Usmanov, A.M. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 195-199; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, **17**, 152-156 (Isoregelinone)

Usmanov, A.M. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 81-84; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 76-79 (pmr, struct)

Regholarrhennine C

R-39

N-Methyl-23-norconca-3,5-dienin-3-amine, 9CI. 3-Methylamino-N-norconca-3,5-diene [114687-92-2]



C₂₂H₃₄N₂ 326.524


Alkaloid from the stem bark of *Holarhena antidyserterica* collected at flowering (Apocynaceae). Cryst. (MeOH/Me₂CO). Mp 302-304°. $[\alpha]_D^{33}$ -80 (c, 0.2 in MeOH).

N,N'-Di-Ac: Mp 191-192°.

Bhutani, K.K. *et al.*, *Phytochemistry*, 1988, **27**, 925 (isol, uv, ir, pmr, ms, cd, ord, struct)

Remijimine

R-40

Relative
ConfigurationC₁₉H₂₄N₂O₂ 312.411

Alkaloid from the leaves of *Remijia peruviana*. Prisms (MeOH/EtOAc). Mp 214-216°. [α]_D²⁵ -21.9 (c, 0.57 in MeOH). λ_{max} 207 (log ε 4.6); 251 (log ε 3.97); 282 (log ε 3.57) (EtOH).

3-Epimer: *Epiremijimine*C₁₉H₂₄N₂O₂ 312.411


Alkaloid from the leaves of *Remijia peruviana*. Amorph. [α]_D²⁵ +41.6 (c, 0.13 in MeOH). λ_{max} 206 (log ε 4.05); 251 (log ε 3.14); 287 (log ε 2.85) (EtOH).

Díaz, J.G. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1667-1671 (*isol, pmr, cmr, cryst struct*)

Renieramycin A

R-41

[79664-60-1]

C₃₀H₃₄N₂O₉ 566.607

Metab. from the marine sponge *Reniera* sp. Shows antimicrobial and antitumour props.; active against *Staphylococcus aureus* and *Bacillus subtilis*. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁰ -36.3 (c, 0.16 in MeOH). λ_{max} 267 (ε 15800); 363 (ε 1400) (MeOH) (Derep). λ_{max} 268 (ε 15800); 365 (ε 1370) (MeOH) (Berdy).

O-*Detigloyl*, 22-*Ac*: *Xestomycin*C₂₇H₃₀N₂O₉ 526.542

Isol. from *Xestospongia* sp. Antibacterial agent. Yellow powder. [α]_D -56 (MeOH). Poorly documented. λ_{max} 212 ; 267 ; 347 (MeOH).

Et ether: *Renieramycin B*

[79664-61-2]

C₃₂H₃₈N₂O₉ 594.66

Prod. by *Reniera* sp. Shows antimicrobial and antitumour props. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁰ -32.2 (c, 0.15 in MeOH). Prob. an artifact. λ_{max} 265 (ε 16100); 370 (ε 1450) (MeOH) (Derep). λ_{max} 268 (ε 17400); 365 (ε 1460) (MeOH) (Berdy).

21α-*Hydroxy*, *Me ether*: *Renieramycin F*

[123641-96-3]

C₃₁H₃₆N₂O₁₀ 596.633

From *Reniera* sp. Yellow powder. λ_{max} 265 (ε 16100); 370 (ε 1450) (MeOH) (Derep). λ_{max} 265 (ε 12000) (MeOH) (Berdy).

21α-*Hydroxy*, 14-*deoxy*: *Renieramycin E*

[123641-95-2]

C₃₀H₃₄N₂O₉ 566.607

From *Reniera* sp. Yellow powder. λ_{max} 267 (ε 15800); 363 (ε 1400) (MeOH) (Derep). λ_{max} 266 (ε 17000) (MeOH) (Berdy).

21-*Oxo*: *Renieramycin C*

[79664-62-3]

C₃₀H₃₂N₂O₁₀ 580.59

Prod. by *Reniera* sp. Shows antimicrobial and antitumour props. Poorly sol. hexane. [α]_D²⁰ -89.2 (c, 0.065 in MeOH). λ_{max} 267 (ε 15800); 363 (ε 1400) (MeOH) (Derep). λ_{max} 266 (ε 14900); 360 (ε 21600) (MeOH) (Berdy).

21-*Oxo*, *Et ether*: *Renieramycin D*

[79664-63-4]

C₃₂H₃₆N₂O₁₀ 608.644

Prod. by *Reniera* sp. Shows antimicrobial and antitumour props. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²⁰ -100.7 (c, 0.092 in MeOH). λ_{max} 265 (ε 16100); 370 (ε 1450) (MeOH) (Derep).

21-*Oxo*, 8α,9-*didehydro*, *Me ether*: *Renieramycin I*

[224566-79-4]

C₃₁H₃₂N₂O₁₀ 592.601

Isol. from the sponge *Haliclona cribri-cutis*. λ_{max} 220 (log ε 4.4); 260 (log ε 4.1); 340 (log ε 3.5); 520 (log ε 3.1) (MeOH).

21-*Oxo*, 14-*deoxy*: *Renieramycin G*

[143458-75-7]

C₃₀H₃₂N₂O₉ 564.591

Major metab. from the Fijian sponge *Xestospongia caycedoi*. Exhibits moderate cytotoxicity. Unstable. λ_{max} 267 (ε 15800); 363 (ε 1400) (MeOH) (Derep). λ_{max} 269 (ε 20000) (MeOH) (Berdy).

21α-*Cyano*: *Renieramycin O*

[631913-68-3]

C₃₁H₃₃N₃O₉ 591.616

Isol. from a *Xestospongia* sp. pre-treated with KCN. Oxidn. prod. of Renieramycin N. Pale yellow powder. [α]_D²⁰ -134.4 (c, 0.7 in CHCl₃). λ_{max} 269 (log ε 4.62); 370 (log ε 3.1) (no solvent reported).

21α-*Cyano*, *Me ether*: *Renieramycin R*

[724707-54-4]

C₃₂H₃₅N₃O₉ 605.643

Isol. from a *Xestospongia* sp. pre-treated with KCN. Pale yellow solid. [α]_D¹⁸ -17.6 (c, 0.1 in CHCl₃). λ_{max} 269 (log ε 4.62); 370 (log ε 3.12) (no solvent reported).

21α-*Cyano*, 14-*deoxy*: *Renieramycin M*

[631913-64-9]

C₃₁H₃₃N₃O₈ 575.617

Isol. from a *Xestospongia* sp. pre-treated with KCN. Cytotoxic. Dark yellow prisms (EtOAc). Mp 194.5-197°. [α]_D²⁰ -49.5 (c, 1 in CHCl₃). λ_{max}

269 (log ε 4.63); 370 (log ε 3.11) (no solvent reported).

21α-*Cyano*, 14-*deoxy*, O-*deacyl*: *Jorunnamycin A*C₂₆H₂₇N₃O₇ 493.515

Isol. from *Jorunna funebris* pre-treated with KCN. Pale yellow solid. [α]_D²⁵ -270.6 (c, 1 in CHCl₃). λ_{max} 269 (log ε 4.61); 370 (log ε 3.11) (no solvent reported).

21α-*Cyano*, 14-*deoxy*, O-*deacyl*, O-*propa-nyl*: *Jorunnamycin C*C₂₉H₃₁N₃O₈ 549.579

Isol. from *Jorunna funebris* pre-treated with KCN. Yellow solid. [α]_D²⁰ -91.6 (c, 0.1 in CHCl₃). λ_{max} 268 (log ε 4.25); 370 (log ε 3.14) (no solvent reported).

21α-*Cyano*, 14-*deoxy*, 7-O-*de-Me*: *Renieramycin S*C₃₀H₃₁N₃O₈ 561.59

Isol. from a *Xestospongia* sp. pre-treated with KCN. Pale yellow needles (EtOAc/petrol). Mp 179-180°. [α]_D²⁰ -38.8 (c, 0.1 in CHCl₃). λ_{max} 269 (log ε 4.62); 370 (log ε 3.12) (no solvent reported).

21α-*Cyano*, 15,18-*hydroquinone*: *Renieramycin N*

[631913-65-0]

C₃₁H₃₅N₃O₉ 593.632

Isol. from a *Xestospongia* sp. pre-treated with KCN. Cytotoxic. Pale yellow prisms (EtOH). Mp 162.5-164°. [α]_D²⁰ -24.7 (c, 0.01 in MeOH). λ_{max} 273 (log ε 3.99); 370 (log ε 2.7) (no solvent reported).

21α-*Cyano*, 15,18-*hydroquinone*, 14-*ketone*: *Renieramycin Q*

[724707-53-3]

C₃₁H₃₃N₃O₉ 591.616

Isol. from a *Xestospongia* sp. pre-treated with KCN. Pale yellow solid. [α]_D¹⁸ -69.8 (c, 0.1 in CHCl₃). λ_{max} 224 (log ε 4.02); 276 (log ε 4.07); 374 (log ε 3.61) (no solvent reported).

21α-*Cyano*, 15,18-*hydroquinone*, 14-*ketone*, O-*deacyl*: *Jorunnamycin B*C₂₆H₂₇N₃O₈ 509.515

Isol. from *Jorunna funebris* pre-treated with KCN. Orange-yellow solid. [α]_D²⁰ +117.6 (c, 0.11 in CHCl₃). λ_{max} 254 (log ε 3.96); 275 (log ε 4.15); 371 (log ε 3.64) (no solvent reported).

Frincke, J.M. *et al.*, *J.A.C.S.*, 1982, **104**, 265-269 (*Renieramycins A-D, isol, uv, ir, pmr, cmr, ms, struct*)

He, H. *et al.*, *J.O.C.*, 1989, **54**, 5822-5824 (*Renieramycins E,F*)

Fukuyama, T. *et al.*, *Tet. Lett.*, 1990, **31**, 5989-5992 (*synth, struct*)

Gulavita, N.K. *et al.*, *CA*, 1992, **117**, 230454q (*Xestomycin*)

Davidson, B.S. *et al.*, *Tet. Lett.*, 1992, **33**, 3721-3724 (*Renieramycin G*)

Parameswaran, P.S. *et al.*, *Indian J. Chem., Sect. B*, 1998, **37**, 1258-1263 (*Renieramycin I*)

U.S. Pat., 2001, 7 241 892; *CA*, **134**, 17616 (*Xestomycin*)

Suwanborirux, K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1441-1446; 2004, **67**, 1023-1028 (*Renieramycins M,N,O,Q,R,S*)


Magnus, P. *et al.*, *J.A.C.S.*, 2005, **127**, 12476-12477 (*Renieramycin G, synth*)

Lane, J.W. *et al.*, *J.A.C.S.*, 2005, **127**, 12684-12690 (*Renieramycin G, synth*)
 Charupant, K. *et al.*, *Chem. Pharm. Bull.*, 2007, **55**, 81-86 (*Jorunnamycins A-C*)

Renieramycin H

R-42

Cribrostatin 4
 [276682-85-0]
 [224566-78-3]



$C_{30}H_{30}N_2O_{10}$ 578.574


Struct. revised in 2001. Isol. from sponges *Cribrachalina* sp. and *Haliclona cribicutis*. Cytotoxic agent. Red prisms (MeOH). Mp 190-192° dec. λ_{max} 209 (€ 40770); 274 (€ 13680); 359 (€ 8070); 507 (€ 3420) (MeOH).

Pasameswaran, P.S. *et al.*, *Indian J. Chem., Sect. B*, 1998, **37**, 1258-1263 (*isol*)
 Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 793-798 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)
 Saito, N. *et al.*, *Heterocycles*, 2001, **55**, 21-28 (*pmr, cmr, struct*)
 Chan, C. *et al.*, *J.A.C.S.*, 2005, **127**, 4596-4598; 7262 (*synth*)
 Vincent, G. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 1517-1520 (*synth*)
 Chen, X. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 3962-3965 (*synth*)

Renieramycin K

R-43

[631913-66-1]



$C_{34}H_{40}N_2O_{10}$ 636.697

Isol. from a Thai sponge *Xestospongia* sp. Amorph. yellow powder. λ_{max} 271 (log € 4.4); 372 (log € 3.11) (no solvent reported).

3-Demethoxy: Renieramycin J

[500111-59-1]

$C_{33}H_{38}N_2O_9$ 606.671

Isol. from *Xestospongia* sp. Amorph. yellow powder. λ_{max} 273 (log € 4.32); 370 (log € 3) (no solvent reported).

4,7-Hydroquinone, O³-de-Me, 3-ketone:**Renieramycin L**

[631913-67-2]

$C_{33}H_{38}N_2O_{10}$ 622.671

Isol. from a *Xestospongia* sp. Pale yellow powder. λ_{max} 244 (log € 4.02); 276 (log € 4.07); 375 (log € 3.75) (no solvent reported).


Suwanborirux, K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1441-1446 (*isol, pmr, cmr*)

Renieramycin P

R-44

Renieramycin J (obsol.)†

[593280-18-3]



Absolute Configuration

$C_{30}H_{36}N_2O_{10}$ 584.622


Originally called Renieramycin J; renamed to avoid duplication. Isol. from the sponge *Neopetrosia* sp. Brownish solid. $[\alpha]_D^{21} +1.2$ (c, 0.02 in MeOH). λ_{max} 203 (€ 39000); 273 (€ 12000) (MeOH).

Oku, N. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1136-1139 (*isol, pmr, cmr*)
 Suwanborirux, K. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1441-1446 (*nomencl*)

Renierol†

R-45

1-(Hydroxymethyl)-7-methoxy-6-methyl-5,8-isoquinolinedione, 9CI
 [77640-19-8]



$C_{12}H_{11}NO_4$ 233.223

Alkaloid from the Fijian sponge *Xestospongia caycedoi*; also from *Oceanapia* sp. Shows antibiotic activity. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . λ_{max} 205 (€ 15000); 241 (€ 13000); 314 (€ 4100) (MeOH) (Berdy).

O-Ac: Renierol acetate

[97603-18-4]

$C_{14}H_{13}NO_5$ 275.26

Metab. from the marine sponge *Xestospongia* sp. and its associated nudibranch *Jorunna funebris*. Cryst. (MeOH). Mp 118-119°. λ_{max} 246 (€ 17780); 318 (€ 5011) (MeOH) (Berdy). λ_{max} 206 ; 237 ; 296 ; 315 (EtOH) (Berdy).

O-Propanoyl: Renierol propionate

[124909-65-5]

$C_{15}H_{15}NO_5$ 289.287

Isol. from the sponge *Xestospongia* sp. and its associated nudibranch *Jorunna funebris*. Yellow needles (MeOH). Mp 89-90°.

O-Angeloyl: Renierone

[73777-65-8]

$C_{17}H_{17}NO_5$ 315.325

Major metab. from an intense blue sponge *Reniera* sp., possibly prod. by symbiotic microorganism. Active against gram-positive bacteria, fungi and tumours. Mp 91.5-92.5°. Related to Mimocin, M-622. λ_{max} 214 (€ 20000); 312 (€ 4000) (MeOH) (Derep).

O⁷-De-Me, O¹-Ac: O-Demethylrenierol acetate

[149022-06-0]

$C_{13}H_{11}NO_5$ 261.234

Isol. from a bright blue sponge *Petrosia* sp. Shows antibacterial activity. Yellow oil. λ_{max} 226 (€ 19400); 300 (€ 4094) (MeOH).

O⁷-De-Me, O-angeloyl: O-Demethylrenierone

[77515-81-2]

$C_{16}H_{15}NO_5$ 301.298

Isol. from an unidentified *Reniera* sp. and a *Cribrachalina* sp. Shows antibacterial props. Orange solid. Mp 135-136°. λ_{max} 208 (€ 12500); 248 (€ 7640); 290 (€ 3260); 316 (€ 1610) (MeOH) (Derep). λ_{max} 228 (log € 3.92); 298 (log € 3.31) (MeOH).

1,2-Dihydro, O-Ac, N-formyl: N-Formyl-1,2-dihydrorenierol acetate

[124909-71-3]

$C_{15}H_{15}NO_6$ 305.287

Isol. from the sponge *Xestospongia* sp. and its associated nudibranch *Jorunna funebris*. Dark red oil. Sol. MeOH. λ_{max} 269 (€ 9772); 340 (€ 3467); 500 (€ 1550) (MeOH) (Berdy). λ_{max} 204 ; 217 ; 269 ; 344 (EtOH) (Berdy).

1,2-Dihydro, O-propanoyl, N-formyl: N-Formyl-1,2-dihydrorenierol propionate

[124909-72-4]

$C_{16}H_{17}NO_6$ 319.313

Isol. from the sponge *Xestospongia* sp. and its associated nudibranch *Jorunna funebris*. Dark red oil. λ_{max} 268 (€ 10000); 340 (€ 3715); 500 (€ 1737) (MeOH) (Berdy).

1,2R-Dihydro, O-angeloyl, N-formyl: (-)-N-Formyl-1,2-dihydrorenierone

[79664-57-6]

$C_{18}H_{19}NO_6$ 345.351

Isol. from a *Reniera* sp. Active against tumours. Red non-cryst. solid. Sol. MeOH, $CHCl_3$; poorly sol. H_2O , hexane. $[\alpha]_D^{20} -227$ (c, 0.023 in MeOH). Shown to be a 2:1 mixt. of two rotamers by pmr and cmr. λ_{max} 216 (€ 31000); 265 (€ 16500); 340 (€ 6000); 515 (€ 3500) (MeOH) (Derep).

1,2S-Dihydro, O-angeloyl, N-formyl: (+)-N-Formyl-1,2-dihydrorenierone

[80108-30-1]

$C_{18}H_{19}NO_6$ 345.351

Isol. from a *Cribrachalina* sp. and a *Reniera* sp. Dark red solid. $[\alpha]_D +230$ (c, 0.04 in MeOH). λ_{max} 218 (log € 4.35); 268 (log € 3.96); 344 (log € 3.53) (MeOH).

1,2-Dihydro, O-angeloyl, N-(3-oxo-1-butenyl): N-(3-Oxo-1-butenyl)-1,2-dihydrorenierone

[182293-46-5]

C₂₁H₂₃NO₆ 385.416

Isol. from the Philippine marine sponge *Xestospongia* sp. Active against gram-positive bacteria. Shows weak insecticidal activity. Blue gum. λ_{\max} 223 (ε 27000); 312 (ε 23000); 330 (ε 21000) (MeOH).

McIntyre, D.E. *et al.*, *Tet. Lett.*, 1979, 4163-4166 (*Renierone*, *isol*, *uv*, *ir*, *cmr*, *pmr*, *cryst struct*)

Danishefsky, S. *et al.*, *Tet. Lett.*, 1980, 4819-4822 (*Renierone*, *synth*)

Kubo, A. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 595-596; 1983, **31**, 341-343; 1986, **34**, 4056-4068; 1989, **37**, 1384-1386 (*synth*, *ir*, *derivs*)

Frincke, J.M. *et al.*, *J.A.C.S.*, 1982, **104**, 265-269 (*Renierone*, *N-formyl-1,2-dihydrorenierone*, *O-Demethylrenierone*)

Berman, E. *et al.*, *Diss. Abstr. Int.*, **B**, 1984, **45**, 556 (*synth*)

Kitahara, Y. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 823-830 (*cmr*)

McKee, T.C. *et al.*, *J. Nat. Prod.*, 1987, **50**, 754-756 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Saito, N. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1493-1499 (*Renierone*, *synth*)

Venkateswarlu, Y. *et al.*, *Indian J. Chem., Sect. B*, 1993, **32**, 704 (*O-Demethylrenierol acetate*)

Edrada, R.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 973-976 (*N-(3-Oxo-1-butenyl)-1,2-dihydrorenierone*)

Ramesh, P. *et al.*, *J. Nat. Prod.*, 1999, **62**, 780-781 (*activity*, *isol*)

Fontana, A. *et al.*, *Tetrahedron*, 2000, **56**, 7305-7308 (*isol*)


Plubrukarn, A. *et al.*, *Pharm. Biol.*, 2003, **41**, 439-442 ((+)-*N-Formyl-1,2-dihydrorenierone*)

Kuwabara, N. *et al.*, *Tetrahedron*, 2004, **60**, 2943-2952 (*synth*)

Repandine† R-46

6,6',7'-Trimethoxy-2,2'-dimethyloxycanthan-12'-ol, 9CI

[518-92-3]

C₃₇H₄₀N₂O₆ 608.733

Alkaloids covered by this entry (*S,S*-config.) are enantiomeric with respect to those in Limacusine, L-175 and diastereomeric with those in Oxyacanthine, O-1 (1*R*,1'*S*). Alkaloid from the bark of *Daphnandra repandula*, and the leaves, stem and bark of *Daphnandra johnsonii* (Monimiaceae). Shows selective anti-plasmodial activity. Needles (EtOH). Mp 255°. $[\alpha]_{\text{D}}^{15}$ -106 (c, 1.2 in CHCl₃). Epimer of Oxyacanthine in O-1.

N²-De-Me: Demerarine

[15353-21-6]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the bark of *Ocotea rodiaei* (Lauraceae). Needles (MeOH). Mp 222-223°. $[\alpha]_{\text{D}}^{25}$ -162 (c, 0.35 in CHCl₃/MeOH 5:95). Epimer of Seperine in O-1.

N²-De-Me; hydrochloride (1:2):Cryst. + 1H₂O. Mp 278°. $[\alpha]_{\text{D}}$ -181 (c,1.0 in H₂O).**Me ether: O-Methylrepandine**

[4021-17-4]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from the bark of *Daphnandra dielsii* and *Daphnandra repandula*, the leaves, stem and bark of *Daphnandra johnsonii*, and from *Isopyrum thalictroides* (Monimiaceae, Ranunculaceae). Needles (MeOH). Mp 211°. $[\alpha]_{\text{D}}^{13}$ -73 (c, 0.3 in CHCl₃). $[\alpha]_{\text{D}}^{20}$ -108 (c, 1.7 in 0.1M HCl).

Me ether, N²,N²-di-Me:

Needles + 2.5H₂O (H₂O)(as diiodide). Mp 255-260° dec. (diiodide). $[\alpha]_{\text{D}}^{20}$ -93 (c, 1.6 in 50% EtOH aq.).

O⁶-De-Me, O¹²-Me: Johnsonine

[69064-34-2]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the leaves, stems and bark of *Daphnandra johnsonii* (Monimiaceae). Pale-yellow cryst. + 0.5 MeOH. Mp 225° dec. (sinters at 150-152°). $[\alpha]_{\text{D}}^{19}$ -86 (CHCl₃).

O⁷-De-Me, O¹²-Me: Cycleapeltine. Faralaotrine

[38849-80-8]

Alkaloid from the roots of *Cyclea peltata* and the leaves, stem bark and root bark of *Colubrina faralaotra* ssp. *faralaotra* (Menispermaceae, Rhamnaceae). Cryst. (EtOH/CHCl₃). Mp 232-234°. $[\alpha]_{\text{D}}^{25}$ -106 (c, 1.0 in CHCl₃). Enantiomer of Limacusine, L-175. λ_{\max} 282 (ε 5200) (MeOH) (Berdy).

Bick, I.R.C. *et al.*, *J.C.S.*, 1948, 2170; 1949, 2767; 1953, 695 (*Repandine*, *O-Methylrepandine*, *isol*, *struct*, *uv*)

Fujita, E. *et al.*, *Yakugaku Zasshi*, 1952, **72**, 1232; *CA*, **47**, 12409a (*struct*, *synth*)

Hearst, P.J. *et al.*, *J.O.C.*, 1964, **29**, 466; 1968, **33**, 1229 (*Demerarine*)

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 2239 (*uv*, *ord*)

Falco, M.R. *et al.*, *Tet. Lett.*, 1968, 1953 (*pmr*)

Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592 (*ms*)

Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1978, **31**, 2539 (*Johnsonine*)

Moullis, C. *et al.*, *J. Nat. Prod.*, 1981, **44**, 101 (*O-Methylrepandine*, *isol*, *uv*, *ir*, *pmr*, *ms*)


Koike, L. *et al.*, *J.O.C.*, 1982, **47**, 4351 (*cmr*)

Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*Cycleapeltine*)

Angerhoffer, C.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 59-66 (*activity*)

Repanduline R-47

[20398-02-1]

C₃₇H₃₆N₂O₇ 620.701Alkaloid from the bark of *Daphnandra*

repandula, *Daphnandra dielsii* and *Daphnandra tenuipes* (Monimiaceae). Yellow needles. Mp 215-232° dec. (shrinks at 180-185°). $[\alpha]_{\text{D}}^{17}$ +473 (c, 0.98 in CHCl₃) (c, 0.2 in MeOH).

Hydrochloride (1:2): Dec. slowly >100°.

Bick, I.R.C. *et al.*, *J.C.S.*, 1953, 692-695; 695-700 (*isol*, *uv*)

Harley-Mason, J. *et al.*, *J.C.S. (C)*, 1967, 1948-1951 (*ir*, *struct*)

Bick, I.R.C. *et al.*, *J.C.S. (C)*, 1967, 1951-1957 (*pmr*, *ms*, *struct*)

Shamma, M. *et al.*, *Heterocycles*, 1975, **3**, 297-300 (*biosynth*)


Neuhaus, D. *et al.*, *J.A.C.S.*, 1983, **105**, 5996-6002 (*pmr*, *abs config*)

Koike, L. *et al.*, *J. Nat. Prod.*, 1992, **55**, 455-460 (*cmr*)

Rescinnaminol

R-48

[109028-40-2]

C₃₂H₄₂N₂O₆ 550.694


The proposed hemiacetal struct. would not be expected to be stable or isolable. Alkaloid from roots of *Rauwolfia serpentina*. Mp 241-243°.

Siddiqui, S. *et al.*, *Pak. J. Sci. Ind. Res.*, 1986, **29**, 401-403; *CA*, **107**, 36566f

Reserpenediol

R-49

18-Hydroxy-11,17-dimethoxyyohimban-16-methanol, 9CI. *Reserpine alcohol* [482-98-4]



Absolute Configuration

C₂₂H₃₀N₂O₄ 386.49

Alkaloid from *Rauwolfia serpentina* root cultures (Apocynaceae), well known prev. as semisynthetic prod. Mp 217-218°. λ_{\max} 228 ; 273 ; 298 (MeOH).


[75331-13-4 , 75331-14-5]

Uddin, A. *et al.*, *Bangladesh Pharm. J.*, 1978, **7**(3), 15-16; *CA*, **91**, 35721s (*isol*, *uv*)

Reserpine acid

R-50

18-Hydroxy-11,17-dimethoxyyohimban-16-carboxylic acid, 9CI. *Reserpinolic acid* [83-60-3]



Absolute configuration

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Rauwolfia vomitoria*, hydrolyt. prod. of several ester alkaloids. Only recently isol. in unesterified form (Apocynaceae). Mp 241-243°.

Hydrochloride: [1910-70-9]

Mp 257-263°. [α]_D²⁰ -80 (CHCl₃).

Me ester: Methyl reserpate

[2901-66-8]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from *Rauwolfia serpentina*, *Rauwolfia amsoniaefolia*, *Rauwolfia macrophylla* and *Rauwolfia vomitoria* (Apocynaceae). Mp 244-245°. [α]_D²⁰ -101 (c, 0.5 in CHCl₃). λ_{max} 228 (log ε 4.49); 271 (log ε 3.67); 298 (log ε 3.78) (EtOH).

▶LD₅₀ (rat, orl) 479 mg/kg. ZF9890000

O-(3,4-Dimethoxybenzoyl), *Me ester: 3,4-Dimethoxybenzoylreserpine acid methyl ester*

[2901-64-6]

C₃₂H₃₈N₂O₈ 578.661

Alkaloid from root bark of *Rauwolfia vomitoria* (Apocynaceae). Mp 225-230° dec. Mp 230° (synthetic). [α]_D²⁰ -125.5 (c, 1 in CHCl₃). λ_{max} 223 (log ε 4.74); 263 (log ε 4.27); 296 (log ε 4.13) (MeOH).

O-(3,4,5-Trimethoxybenzoyl), *Me ester*: see Reserpine, R-52

O-(3-Methoxy-4,5-methylenedioxybenzoyl), *Me ester: Veneserpine*

[23924-95-0]

C₃₂H₃₆N₂O₉ 592.644

Alkaloid from *Alstonia venenata* stem bark (Apocynaceae). Mp 249-250° dec. [α]_D²⁰ -100 (CHCl₃). pK_a 6.1. λ_{max} 225 (log ε 4.65); 274 (log ε 4.18); 292 (log ε 4.08) (EtOH).

O-(2-Acetamidobenzoyl), *Me ester: Anthraserpine*

[123690-03-9]

C₃₂H₃₇N₃O₇ 575.66

Alkaloid from *in vitro* hairy-root cultures of *Catharanthus trichophyllus*. Highly unstable. Opt. rotn. not recorded. λ_{max} 220 (log ε 4.74); 251 (log ε 4.48); 310 (log ε 4.02) (EtOH).

O-(2-Acetamido-4,5-dimethoxybenzoyl), *Me ester: Dimethoxyanthraserpine*

[123690-04-0]

C₃₄H₄₁N₃O₉ 635.713

Major alkaloid from *in vitro* hairy-root cultures of *Catharanthus trichophyllus*. Amorph. solid. [α]_D²⁰ -73 (c, 0.6 in CHCl₃). λ_{max} 229 (log ε 4.58); 272 (log ε 4.18); 307 (log ε 3.96) (EtOH).

O-(3,4,5-Trimethoxycinnamoyl), *Me ester: Rescinnamine, BAN, INN, JAN. Reserpiline†. Anaprel. Apoterin. Cinnaloid. Rescaloid. Moderil. Scinnamina. Many other names*

[84-34-4]

[24815-24-5]

C₃₅H₄₂N₂O₉ 634.725

Alkaloid from *Rauwolfia vomitoria*, *Rauwolfia caffra*, *Rauwolfia cuminsii*, *Rauwolfia macrophylla*,

Rauwolfia oreogiton, *Rauwolfia volkensii*, *Rauwolfia mombasiana* and many other *Rauwolfia* spp.

(Apocynaceae). Antihypertensive agent and tranquilliser agent. Sedative. Cryst. (Me₂CO aq.). Mp 238-239° (in vacuo). [α]_D¹⁷ -99 (c, 1.3 in CHCl₃). Log P 3.6 (calc). Not the same as Reserpiline, R-53. Shows similar pharmacol. to Reserpine but is less potent.

▶LD₅₀ (rat, orl) 1000 mg/kg.

O-(3,4,5-Trimethoxycinnamoyl), O¹⁷-*De-Me, Me ester: Rescidine*

[3314-86-1]

C₃₄H₄₀N₂O₉ 620.698

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae). Needles (MeOH). Mp 183-186° dec. [α]_D²⁰ -63.4 (c, 1 in CHCl₃). λ_{max} 229 (log ε 4.76); 302 (log ε 4.42) (no solvent reported).

O-[3-(3,4,5-Trimethoxyphenyl)propionyl], *Me ester: Rescinnamidine*

[110222-52-1]

C₃₅H₄₄N₂O₉ 636.741

Alkaloid from roots of *Rauwolfia serpentina*. Yellow needles (CHCl₃/Et₂O). Mp 260-261°. [α]_D²⁰ +207 (CHCl₃). λ_{max} 210 ; 225 ; 270 ; 295 (MeOH).

O¹⁷-*De-Me, 17-O-(3,4,5-trimethoxybenzoyl), Me ester: Raugustine*

[522-79-2]

C₃₂H₃₈N₂O₉ 594.66

Alkaloid from *Rauwolfia ligustrina* (Apocynaceae). Cryst. + 1H₂O. Mp 160-170° dec. (in vacuo). [α]_D²⁴ -50 (c, 0.609 in CHCl₃). λ_{max} 217 (ε 54600); 268 (ε 15400); 291 (ε 9300) (EtOH).

O¹⁷-*De-Me, 18-O-(2-acetamidobenzoyl), Me ester: Pseudoanthraserpine*

[123690-05-1]

C₃₁H₃₅N₃O₇ 561.633

Alkaloid from *in vitro* hairy-root cultures of *Catharanthus trichophyllus*. Amorph. Tentative struct. No stereochem. implied.

3-*Epimer, O-(3,4,5-trimethoxycinnamoyl), Me ester: 3-Epirescinnamine*

[187082-77-5]

C₃₅H₄₂N₂O₉ 634.725

Alkaloid from root bark of *Rauwolfia vomitoria* (Apocynaceae). Mp 238°. [α]_D²⁰ -137.2 (c, 1 in CHCl₃). λ_{max} 230 (log ε 4.57); 300 (log ε 4.2) (MeOH).

Furlenmeyer, A. et al., *Experientia*, 1953, 9, 331-333 (*Me ester, uv, ir*)

Klohs, M.W. et al., *J.A.C.S.*, 1955, 77, 2241-2243 (*Rescinnamine, isol*)

Kroneberg, G. et al., *Arzneim.-Forsch.*, 1956, 6, 579-583 (*Rescinnamine, pharmacol*)

Müller, J.M. et al., *Experientia*, 1957, 13, 479-481 (*Raugustine, isol*)

Zoha, M.S. et al., *J. Pharm. Pharmacol., Suppl.*, 1958, 10, 231T-240T (*Rescinnamine, pharmacol*)

Woodward, R.B. et al., *Tetrahedron*, 1958, 2, 1-57 (*synth, Me ester*)

Rosen, W.E. et al., *J.A.C.S.*, 1961, 83, 4816-4819 (*pmr, Me ester*)

Popelak, A. et al., *Naturwissenschaften*, 1961, 48, 73-74 (*Rescidine, isol*)

Popelak, A. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1963, 296, 261-265 (*Rescidine, struct*)

Chatterjee, A. et al., *Chem. Ind. (London)*, 1969, 1388-1389 (*Veneserpine, isol, struct*)

Pantarotto, C. et al., *Adv. Mass Spectrom. Biochem. Med.*, 1977, 2, 351-364 (*Rescinnamine, detn, glc, ms*)

Suzuki, Y. et al., *CA*, 1982, 96, 115441 (*pharmacol*)

Malik, A. et al., *J. Nat. Prod.*, 1983, 46, 939 (*isol*)

Lounasmaa, M. et al., *Heterocycles*, 1985, 23, 371-375 (*pmr*)

Siddiqui, S. et al., *J. Nat. Prod.*, 1987, 50, 238-240 (*Rescinnamidine*)

Davioud, E. et al., *Phytochemistry*, 1989, 28, 1383-1387 (*Anthraserpine, Dimethoxyanthraserpine, Pseudoanthraserpine*)

Awang, D.V.C. et al., *J. Pharm. Biomed. Anal.*, 1990, 8, 91; *CA*, 113, 115663n (*cmr, Rescinnamine*)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 11465 (*synonyms*)

Petati, M. et al., *Fitoterapia*, 1996, 67, 422-426 (*3-Epirescinnamine, 3,4-Dimethoxybenzoylreserpine acid methyl ester*)


Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 941

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, MPH300; TLN500

Reserpiline

Elliptamine

[131-02-2]



Absolute Configuration

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Rauwolfia canescens* and very many other *Rauwolfia* spp., *Aspidosperma discolor* and some *Ochrosia* spp. (Apocynaceae). Noncryst. [α]_D²⁰ -12 (CHCl₃). λ_{max} 229 (log ε 4.57); 250 (log ε 4.07); 300 (log ε 4.03); 304 (log ε 4.03); 307 (sh) (log ε 4) (EtOH).

▶RP5776550

Hydrochloride: Mp 205-207°. [α]_D²⁴ -40 (c, 0.44 in EtOH).

(4R)-*N-Oxide: Reserpiline (4R)-N-oxide*

[77795-01-8]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{max} 224 ; 274 ; 291 ; 296 ; 302 ; 308 (EtOH).

(4S)-*N-Oxide: Reserpiline (4S)-N-oxide*

[77795-02-9]

C₂₃H₂₈N₂O₆ 428.484

Alkaloid from the trunk bark of *Ochrosia moorei* (Apocynaceae). λ_{max} 225 ; 276 ; 290 ; 297 ; 302 ; 307 (EtOH).

N²-*Me: 4-Methylreserpiline*

[49685-85-0 (bromide)]

C₂₄H₃₁N₂O₅[⊕] 427.519

Quaternary alkaloid from the root bark of *Rauwolfia confertiflora* (Apocynaceae).

19,20-Didehydro: 19,20-Dehydroreserpiline

[70509-80-7]
C₂₃H₂₆N₂O₅ 410.469

Alkaloid from the stem bark of *Rauwolfia vomitoria* (Apocynaceae).

3-Epimer: Isoreserpiline

[572-67-8]

C₂₃H₂₈N₂O₅ 412.485

From *Rauwolfia* spp., *Aspidosperma discolor*, *Vinca* and *Ochrosia* spp. (Apocynaceae). Mp 211-212° dec. [α]_D²⁰ -82 (Py). λ_{\max} 220 (log ϵ 4.58); 250 (log ϵ 4.16); 300 (log ϵ 4.01); 304 (log ϵ 4.02); 311 (sh) (log ϵ 3.98) (EtOH or MeOH).

3-Epimer, N⁴-Me: Holecimine. N^b-Methylisoreserpilinium

[6879-77-2]

C₂₄H₃₁N₂O₅[⊕] 427.519

Quaternary alkaloid from the bark of *Ochrosia sandwicensis* (Apocynaceae). Needles (MeOH/EtOAc) (as chloride). Mp 283-285° dec. (chloride). [α]_D²³ -134.5 (c, 2.97 in MeOH) (chloride).

3-Epimer, 18,19-didehydro: 18,19-Dehydro-3-isoreserpiline. Darcyribeirine

[432031-09-9]

Alkaloid from the root bark of *Rauwolfia grandiflora*. Amorph. solid. [α]_D²⁰ -36 (c, 0.05 in CHCl₃). λ_{\max} 226 (log ϵ 4.53); 299 (log ϵ 4.01) (no solvent reported).

3,20-Diepimer: 10,11-Dimethoxyajmalicine

[55529-56-1]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Cabucala striolata*, *Rauwolfia cummingsii* and *Neisosperma glomerata* (Apocynaceae).

19,20-Diepimer: 3-Epirauvanine

[6835-93-4]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from the bark of *Neisosperma oppositifolia* (Apocynaceae). Amorph. [α]_D +111 (CHCl₃).

3,19,20-Triepimer: Rauvanine

[3148-42-3]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae). Leaflets + 0.5H₂O (MeOH). Mp 129-135° dec. [α]_D +32.5 (c, 1 in CHCl₃). λ_{\max} 228 (log ϵ 4.54); 298 (log ϵ 3.98); 315 (log ϵ 3.99) (no solvent reported).

▶ RP5776500

3,19,20-Triepimer, hydrochloride: Mp 280-282° dec.

Stereoisomer: Neoreserpiline

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from leaves and stems of *Rauwolfia perakensis* (Apocynaceae). Mp 129-131°. [α]_D²³ -78 (Py). Config. undetd. No CAS no. reported.

Stereoisomer, nitrate: Mp 252-254° dec.

Stoll, A. et al., *Helv. Chim. Acta*, 1955, **38**, 270-283 (isol, uv, ir, struct)

Goutarel, R. et al., *C. R. Hebd. Seances Acad. Sci.*, 1961, **253**, 2589-2591 (*Rauvanine*, uv, ir, pmr, struct)

Scheuer, P.J. et al., *J.O.C.*, 1961, **26**, 3069-3071 (*Holecimine*)

Shamma, M. et al., *J.A.C.S.*, 1962, **84**, 1739-1740; 1963, **85**, 2507-2512 (pmr, config)

Spiteller, G. et al., *Monatsh. Chem.*, 1963, **94**, 779-785 (ms)

Poison, J. et al., *Bull. Soc. Chim. Fr.*, 1964, 2853-2856 (*3-Epirauvanine*)

Kiang, A.K. et al., *J. Nat. Prod.*, 1964, **27**, 220-225 (*Neoreserpiline*)

Finch, N. et al., *Tetrahedron*, 1966, **22**, 1327-1333 (uv, ord, config)

Bombardelli, E. et al., *Fitoterapia*, 1973, **43**, 67-70; 1974, **45**, 183-187; *CA*, **79**, 137335p; **83**, 25051u (*4-Methylreserpiline*, *10,11-Dimethoxyajmalicine*)

Sabri, N.N. et al., *Phytochemistry*, 1978, **17**, 2023-2026 (*19,20-Dehydroreserpiline*)

Rüffer, M. et al., *Tet. Lett.*, 1978, 1593-1596 (*biosynth*)

Ahond, A. et al., *J. Nat. Prod.*, 1981, **44**, 193-199 (*oxides*)

Seguin, E. et al., *J. Nat. Prod.*, 1982, **45**, 738-744; 1984, **47**, 687-691 (*10,11-Dimethoxyajmalicine*)

Seguin, E. et al., *Helv. Chim. Acta*, 1983, **66**, 2059-2067 (*cd*)

Amarasekara, A.S. et al., *Fitoterapia*, 1986, **57**, 55-57 (*3-Epirauvanine*, *isol*)


De Bruyn, A. et al., *Magn. Reson. Chem.*, 1989, **27**, 935-940 (pmr, cmr)

Cancelieri, N.M. et al., *Tet. Lett.*, 2002, **43**, 1783-1787 (*Darcyribeirine*, *Isoreserpiline*)

Reserpine, BAN, INN, JAN, USAN

Serpasil. Sermix. Reserpoid. Sandril.

Apoplone. Carpacil. Many other names



C₃₃H₄₀N₂O₉ 608.687

Antiadrenergic, anticonvulsant agent. Antineoplastic agent. Possesses sedative tranquilliser props. Antihypertensive agent acting by depleting stores of noradrenaline in sympathetic neurones. Many semisynthetic analogues have been investigated. Log P 3.37 (calc).

(-)-form [50-55-5]

Alkaloid from *Rauwolfia serpentina*, *Rauwolfia vomitoria*, *Rauwolfia macrophylla*, very many other *Rauwolfia* spp., *Vinca minor*, *Alstonia constricta* and many other spp. in the Apocynaceae, e.g. *Vallesia dichotoma* and *Excavatia coccinea* (preferred genus name *Ochrosia*). Active against fungi. Long prisms (Me₂CO aq.). Sol. CHCl₃, AcOH; fairly sol. H₂O. Mp 262-266° Mp 284-285°. λ_{\max} 216 (ϵ 61700); 267 (ϵ 1700); 295 (ϵ 12200) (MeOH) (Berdy).

▶ Adverse effects reported when used therapeutically incl. mental depression and gastrointestinal disturbances. Human and exp. reprod. effects. Exp. teratogen.

Exp. neoplastic agent. LD₅₀ (rat, orl) 420 mg/kg. ZG0350000

Hydrochloride: Mp 224°.

Perchlorate: Mp 238-239°.

Picrate: Mp 183-186°.

N^b-Oxide: Renoxydine. Reserpoxidine.

Renoxidine

[474-48-6]

C₃₃H₄₀N₂O₁₀ 624.686

Alkaloid from *Rauwolfia vomitoria*, *Rauwolfia serpentina*, *Rauwolfia canescens*, several other *Rauwolfia* spp. and *Melodinus balansae* (Apocynaceae). Shows similar pharmacol. props. to Reserpine but with about half the activity. Cryst. + ½ H₂O (Me₂CO/hexane). Mp 238-241° dec. [α]_D -100 (CHCl₃). λ_{\max} 218 (ϵ 64300); 267 (ϵ 16600); 296 (ϵ 11000) (EtOH).

1-N-(Diethylaminoethyl): Bietaserpine, INN

[53-18-9]

C₃₉H₅₃N₃O₉ 707.862

Antihypertensive agent. [α]_D¹⁷ -121 (c, 2 in CHCl₃). Log P 4.88 (calc).

▶ Exp. teratogen. ZF9380000

1-N-(Diethylaminoethyl), bitartrate:

Pleiatensin. Pleiatensin simplex. Tensibar. DL 152. S 1210

[1111-44-0]

Mp 145-150° dec.

▶ Exp. reprod. and teratogenic effects (large dose). LD₅₀ (rat, orl) 4885 mg/kg. ZF9385000

O¹⁷-De-Me: Pseudoreserpine. Norreserpine.

ψ-Reserpine

[522-80-5]

C₃₂H₃₈N₂O₉ 594.66

Alkaloid from *Rauwolfia canescens*, *Rauwolfia nitida* and *Rauwolfia ligustrina* (Apocynaceae). Shows antihypertensive and sedative activity, weaker than that of Reserpine. Cryst. (Me₂CO aq.). Mp 257-258°. [α]_D²⁴ -65 (c, 1 in CHCl₃). Log P 2.7 (calc). λ_{\max} 218 (log ϵ 4.76); 268 (log ϵ 4.22); 296 (log ϵ 4.02) (EtOH).

O¹⁷-De-Me; nitrate:

Fine needles. Mp 253-255°.

3-Epimer: 3-Isoreserpine

[482-85-9]

[72173-53-6]

C₃₃H₄₀N₂O₉ 608.687

Alkaloid from *Rauwolfia vomitoria*. Cryst. (MeOH). Mp 152-156° (150-151°). [α]_D²³ -165.1 (c, 1 in CHCl₃).

16,17-Diepimer, O¹⁷-de-Me: 16,17-Diepipseudoreserpine

[82262-76-8]

C₃₂H₃₈N₂O₉ 594.66

Alkaloid from *Rauwolfia yunnanensis* (Apocynaceae). Alkaloid not named in the abstract.

Stereoisomer, O¹⁷-de-Me: Neonorreserpine

[74958-26-2]

Alkaloid from *Rauwolfia vomitoria* (Apocynaceae). Silky prismatic rods (MeOH/C₆H₆). Mp 291-292°. Config. unknown.

Stereoisomer, O¹⁷-de-Me, hydrochloride:

Mp 230-231°.

(±)-form [75331-15-6]


Mp 260-262° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 1069C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 615A (nmr)

Furlenmeyer, A. et al., *Experientia*, 1953, 9, 331-333 (uv, ir)Ulshafer, P.R. et al., *C. R. Hebd. Seances Acad. Sci.*, 1957, 244, 2989-2991 (oxide)Klohs, M.W. et al., *J.A.C.S.*, 1957, 79, 3763-3766 (*Pseudoreserpine*)Velluz, L. et al., *Bull. Soc. Chim. Fr.*, 1958, 145-146; 673-677 (synth)Woodward, R.B. et al., *Tetrahedron*, 1958, 2, 1-57 (synth)Rosen, W.E. et al., *J.A.C.S.*, 1961, 83, 4816-4819 (pmr)U.K. Pat., 1961, 894 866; *CA*, 57, 9910e (*Bietaserpine*)Berthaux, P. et al., *Arzneim.-Forsch.*, 1964, 14, 1040-1045 (*Bietaserpine*, pharmacol, clin trials)Ban, Y. et al., *Tetrahedron*, 1964, 20, 2877-2884 (abs config)Schlittler, E. et al., *Alkaloids (Academic Press)*, 1965, 8, 287-334 (rev)Levin, R.H. et al., *J.O.C.*, 1973, 38, 1983-1986 (cmr)Taylor, W.I. et al., *The Vinca Alkaloids*, Marcel Dekker, 1973, (rev, props)Becker, O. et al., *Org. Mass Spectrom.*, 1977, 12, 461-464 (ms)Mehri, M.H. et al., *Phytochemistry*, 1978, 17, 1451-1452 (oxide)Fanso-Free, S.N.Y. et al., *J.A.C.S.*, 1979, 101, 1549-1553 (*N-15 nmr*)Pearlman, B.A. et al., *J.A.C.S.*, 1979, 101, 6404-6408 (synth)Bein, H.J. et al., *Handb. Exp. Pharmacol.*, (Part 1), 1980, 55, 43-58 (rev, pharmacol)*IARC Monog.*, 1980, 24, 211; *Suppl.*, 6, 485; *Suppl.*, 7, 330 (rev, tox)Wender, P.A. et al., *J.A.C.S.*, 1980, 102, 6157-6159 (synth)Voelter, W. et al., *Z. Naturforsch., B*, 1980, 35, 920-923 (*Neonoreserpine*, isol, cmr)Feng, X. et al., *Yaoxue Xuebao*, 1981, 16, 510-518; *CA*, 97, 3593j (*Diepseudoreserpine*, isol)Muhtadi, F.J. et al., *Anal. Profiles Drug Subst.*, 1984, 13, 737-765 (rev, ir, pmr, cmr, ms, anal)Lounasmaa, M. et al., *Heterocycles*, 1985, 23, 371-375 (pmr)Szántay, C. et al., *Alkaloids (Academic Press)*, 1986, 27, 253-268 (pharmacol, bibl)Martín, S.F. et al., *J.A.C.S.*, 1987, 109, 6124-6134 (synth)Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 7993 (synonyms)Zhang, L.-H. et al., *J.O.C.*, 1989, 54, 4708-4712 (*3-Isoserpine*)Chu, C.-S. et al., *Chem. Comm.*, 1996, 1537-1538 (synth)Nicolaou, K.C. et al., *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, 55 (bibl, synth)Hanessian, S. et al., *J.O.C.*, 1997, 62, 465-473 (synth)Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 942Mehta, G. et al., *J.C.S. Perkin 1*, 2000, 1399-1404 (synth)Sparks, S.M. et al., *J.O.C.*, 2003, 68, 5274-5285 (synth)Stork, G. et al., *J.A.C.S.*, 2005, 127, 16255-16262 (synth)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn.,*J. Wiley*, 2000, DIG800; RDK000; DIH000**Reserpine†**

Methyl 16,17-didehydro-11-methoxy-19-methyloxayohimban-17-carboxylate, 9CI. Raubasinine. Pubescine† [482-96-2]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from *Rauwolfia serpentina*, many other *Rauwolfia* spp., *Vinca major* and several other *Vinca* spp., also in *Ochrosia oppositifolia* (Apocynaceae). Cryst. (Me₂CO). Mp 243-244° dec. (229°). [α]_D²⁰ -131 (c, 1.18 in CHCl₃) (-120). λ_{max} 228 (log ε 4.65); 250 (sh) (log ε 4.12); 296 (log ε 3.81) (no solvent reported).

Hydrochloride: Mp 244-246° dec.

3-Epimer: 3-Isoserpine

[482-95-1]

C₂₂H₂₆N₂O₄ 382.458From *Rauwolfia canescens*, *Ochrosia oppositifolia*, *Vinca major* and *Vinca herbacea* (Apocynaceae). Cryst. (Et₂O/EtOH). Mp 225-226° dec. [α]_D²⁰ -5 (c, 0.7 in Py).**19-Epimer: Raunitidine**

[14883-83-1]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from leaves of *Rauwolfia nitida* (Apocynaceae). Cryst. (EtOH). Mp 276-278°. [α]_D -30.6 (Py). [α]_D -69.5 (CHCl₃). λ_{max} 229 (log ε 4.63); 250 (log ε 4.06); 298 (log ε 3.8) (no solvent reported).▶ LD₅₀ (mus, ivn) 56 mg/kg. NM3400000**19-Epimer, hydrochloride:**Cryst. + 1EtOH (EtOH). Mp 230° dec. [α]_D -108 (MeOH).**20-Epimer: Tetraphylline**

[482-94-0]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from *Rauwolfia* spp. and leaves of *Ochrosia glomerata* (Apocynaceae). Mp 220-223° dec. [α]_D²⁸ -73 (CHCl₃). [α]_D -35 (Py). λ_{max} 229 (log ε 4.65); 298 (log ε 3.81) (EtOH).**20-Epimer, nitrate:** Mp 264-266°.**3,19-Diepimer: Isoraunitidine**

[13555-43-6]

Synthetic. Cryst. (MeOH). Mp 259-261°. [α]_D +131 (Py). λ_{max} 229 (log ε 4.66); 249 (log ε 4.11); 298 (log ε 3.86) (EtOH).**3,20-Diepimer: Raufloridine**


[36063-55-5]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from root bark of *Rauwolfia confertiflora* and bark of *Neisosperma glomerata* (Apocynaceae). Amorph. powder. [α]_D²⁰ +5.95 (c, 0.25 in Me₂CO).

3,20-Diepimer, hydrochloride: Mp 183°.

R-53Weisenborn, F.L. et al., *Chem. Ind. (London)*, 1954, 375-376 (isol, uv, ir, struct)Djerassi, C. et al., *J.A.C.S.*, 1957, 79, 1217-1222 (*Tetraphylline*)Salkin, R. et al., *J. Pharm. Sci.*, 1961, 50, 1038-1041 (*Raunitidine*)Shamma, M. et al., *J.A.C.S.*, 1963, 85, 2507-2512 (pmr, config)Göggel, H. et al., *Chem. Comm.*, 1965, 538-539 (*biosynth*)Finch, N. et al., *Tetrahedron*, 1966, 22, 1327-1333 (uv, ord, config)Danieli, B. et al., *Chim. Ind. (Milan)*, 1971, 53, 1042-1043 (*Raufloridine*)Levin, R.H. et al., *J.O.C.*, 1973, 38, 1983-1986 (cmr)Taylor, W.I. et al., *The Vinca Alkaloids*, Marcel Dekker, 1973, (rev, props)Sakai, S. et al., *Yakugaku Zasshi*, 1978, 98, 850; *CA*, 79, 18923w (synth)Seguin, E. et al., *J. Nat. Prod.*, 1982, 45, 738-744; 1984, 47, 687-691 (*Tetraphylline*, *Raufloridine*, isol)Seguin, E. et al., *Helv. Chim. Acta*, 1983, 66, 2059-2067 (cd)De Bruyn, A. et al., *Magn. Reson. Chem.*, 1989, 27, 935-940 (*Tetraphylline*, pmr)Mukhopadhyay, G. et al., *Phytochemistry*, 1991, 30, 2447-2449 (3-epimer, cryst struct)**Resorcinomycin A****R-54**

N-[N-(Aminoiminomethyl)-2-[3,5-dihydroxy-4-(1-methylethyl)phenyl]glycyl]glycine, 9CI. DO 248A. Antibiotic DO 248A [100234-70-6]

R = -CH(CH₃)₂C₁₄H₂₀N₄O₅ 324.336Peptide antibiotic. Isol. from *Streptovorticillum roseovorticillatum*. Active against acid-fast bacteria. Powder + 2H₂O. Sol. H₂O, DMF, EtOH, MeOH; fairly sol. Me₂CO, EtOAc, CHCl₃; poorly sol. C₆H₆, hexane, Et₂O. Mp 194-200°. λ_{max} 276 (ε 1170); 283 (ε 1170) (MeOH/HCl) (Derep). λ_{max} 297 (ε 2400) (MeOH/NaOH) (Derep). λ_{max} 276 (ε 1170); 283 (ε 1170) (MeOH) (Derep).*Eur. Pat.*, 1985, 159 004; *CA*, 104, 67502 (isol) Kondo, E. et al., *J. Antibiot.*, 1989, 42, 1; 463 (isol, struct, props)**Resorcinomycin B****R-55**

N-[N-(Aminoiminomethyl)-2-(4-ethyl-3,5-dihydroxyphenyl)glycyl]glycine, 9CI. DO 248B. Antibiotic DO 248B [100234-69-3]

As Resorcinomycin A, R-54 with

R = -CH₂CH₃C₁₃H₁₈N₄O₅ 310.309Peptide antibiotic. Isol. from *Streptovorticillum roseovorticillatum*. Active against acid-fast bacteria. Powder + 2H₂O. Sol. H₂O, MeOH, DMF, EtOH; fairly sol. Me₂CO, EtOAc, CHCl₃; poorly sol.


C₆H₆, Et₂O, hexane. Mp 189-192°. λ_{max} 276 (ε 1170); 283 (ε 1170) (MeOH/HCl) (Derep). λ_{max} 297 (ε 2400) (MeOH/NaOH) (Derep). λ_{max} 276 (ε 1170); 283 (ε 1170) (MeOH) (Derep).

Ger. Pat., 1985, 159 004; CA, 104, 67502 (isol)

Kondo, E. et al., J. Antibiot., 1989, 42, 1 (isol, struct, props)

Retamine

12α-Hydroxysparteine
[2122-29-4]



Absolute configuration

C₁₅H₂₆N₂O 250.383

The 12-config. does not appear to be unequivocally established. Originally thought to be 7-, 8-, or 9-hydroxysparteine. Alkaloid from *Genista sphaerocarpa*, *Genista junceum* and most *Genista* spp. (Fabaceae). Powerful hypotensive agent. Mp 166°. [α]_D²⁰ +46.2 (c, 1.02 in EtOH).

▶ VH5373000

Battandier, J. et al., C. R. Hebd. Seances Acad. Sci., 1897, 125, 360; 450 (isol)

Fraga, F. et al., Tetrahedron, 1960, 11, 78 (isol, ir)

Bohlmann, F. et al., Chem. Ber., 1965, 98, 659 (struct)


Shin, K.H. et al., Can. J. Chem., 1965, 43, 2012 (struct, synth, abs config)

Neuner-Jehle, N. et al., Monatsh. Chem., 1967, 98, 836 (ms)

Bohlmann, F. et al., Chem. Ber., 1975, 108, 1043 (cmr)

Reticulatae

[552846-52-3]



C₁₈H₁₂N₂O₂ 288.305

Alkaloid from the sponges *Fascaplysinopsis reticulata* and *Thorectandra* sp. Light yellow solid. λ_{max} 215 (log ε 3.72); 260 (log ε 3.71); 312 (log ε 3.6); 388 (log ε 2.94) (MeOH).

Me ester: **Reticulatine A**. 1-Deoxysecofascaplysin A. Methyl reticulatae [552846-51-2]

C₁₉H₁₅N₂O₂[⊕] 303.34

Alkaloid from the sponges *Fascaplysinopsis reticulata* and a *Thorectandra* sp. Light yellow solid. Counterion not specified. λ_{max} 218 (log ε 3.89); 261 (log ε 3.85); 313 (log ε 3.71); 348 (log ε 3.18); 394 (log ε 3.14) (MeOH).

Me ester, salt with Dehydrolyffariellolide diacid: **Reticulatine B** [135091-12-2]

C₄₄H₅₂N₂O₆ 704.905

Isol. from *Fascaplysinopsis reticulata*. Red amorph. solid. Consists of alkaloid cation with sesterterpenoid anion. λ_{max} 250; 308; 340 (sh); 382 (MeOH) (Derep).

Me ester, salt with 16-Oxodehydrolyffariellolide diacid: **Reticulatine A** [135091-11-1]

C₄₄H₅₀N₂O₇ 718.888

Alkaloid from the sponge *Fascaplysinopsis reticulata*. Red amorph. solid. Consists of alkaloid cation with sesterterpenoid anion. λ_{max} 250; 308; 340 (sh); 382 (MeOH) (Derep).

4'-Bromo: **14-Bromoreticulatae**

[548773-66-6]

C₁₈H₁₁BrN₂O₂ 367.201

Alkaloid from the sponge *Fascaplysinopsis reticulata*. Yellow solid.

4'-Bromo, Me ester: **14-Bromoreticulatine**

[693790-76-0]

C₁₉H₁₄BrN₂O₂[⊕] 382.236

Alkaloid from the Indo-Pacific sponge *Fascaplysinopsis reticulata*. Yellow solid (as chloride).

7-Bromo, Me ester: **Methyl 7-bromoreticulatae**

[1001202-15-8]

C₁₉H₁₄BrN₂O₂[⊕] 382.236

Alkaloid from an undescribed thorectid sponge.

4',7-Dibromo, Me ester: **Methyl 7,14-dibromoreticulatae**

[693790-77-1]

C₁₈H₁₀Br₂N₂O₂ 446.097

Alkaloid from *Fascaplysinopsis reticulata*. Yellow solid. Different numbering systems used.

Jiménez, C. et al., Tet. Lett., 1991, 32, 1843-1846 (*Reticulatines A,B*)

Segraves, N.L. et al., Tet. Lett., 2003, 44, 3471-3475 (isol, pmr, cmr)


Segraves, N.L. et al., J. Nat. Prod., 2004, 67, 783-792 (isol, activity)

Charan, R.D. et al., Nat. Prod. Res., 2004, 18, 225-229 (*Thorectandra* alkaloids)

Ankietty, S. et al., Nat. Prod. Commun., 2007, 2, 1145-1148 (*Methyl 7-bromoreticulatae*)

Reticulatul

[694436-62-9]



C₁₇H₁₃N₂O[⊕] 261.303

Alkaloid from the sponge *Fascaplysinopsis reticulata*. Yellow solid (as chloride).

4'-Bromo: **14-Bromoreticulatul**

[693790-79-3]


C₁₇H₁₂BrN₂O[⊕] 340.199

Alkaloid from *Fascaplysinopsis reticulata*. Yellow solid (as chloride).

Segraves, N.L. et al., J. Nat. Prod., 2004, 67, 783-792 (isol, pmr, cmr, 4'-bromo)

Reticuline

Laudanosine 4',6-dimethyl ether. Coclanoline



(R)-form

C₁₉H₂₃NO₄ 329.395

(R)-form

Alkaloid from *Romneya coulteri* var. *trichocalyx* (Papaveraceae). Intermediate in the biosynth. of Morphine, M-704. Noncryst. [α]_D²⁸ -55 (c, 0.44 in EtOH).

(S)-form [485-19-8]

Alkaloid from *Ammona reticulata* (custard apple), *Cananga odorata* (ylang ylang), *Phyllica rogersii* and *Papaver somniferum* and many spp. over many different plant families (Annonaceae, Rhamnaceae, Papaveraceae). Shows antiplatelet aggregation activity. Needles. Mp 200-202°. [α]_D¹⁵ +132 (MeOH). λ_{max} 258 (log ε 4.5); 285 (log ε 4.39); 334 (log ε 2.17) (MeOH).

▶ LD₅₀ (mus, ipr) 56 mg/kg. NX5993000

Perchlorate: Mp 203-204° (211°). [α]_D³¹ +88 (EtOH).

N-Oxide (R-): **Reticuline N^β-oxide**

[71657-63-1]

C₁₉H₂₃NO₅ 345.394

Alkaloid from the leaves of *Neolitsea sericea* var. *aurata* and *Pachygone ovata*.

N-Oxide (S-): **Reticuline N^α-oxide**

[937018-75-2]

C₁₉H₂₃NO₅ 345.394

Alkaloid from the leaves of *Neolitsea sericea* var. *aurata*.

N-Me: **Tembetarine**

[18446-73-6]

C₂₀H₂₆NO₄[⊕] 344.43

Alkaloid from the bark of *Fagara naranjillo* and several other *Fagara* spp. and from *Zanthoxylum* spp. (Rutaceae). Convulsive agent producing transient hypertension and brachycardia in exp. animals. Mp 236-237° (as chloride). [α]_D²⁹ +123 (c, 0.9 in EtOH).

▶ Toxic, LD₅₀ 63mg/kg (rat).

Di-Me ether: see Laudanosine, L-69

(±)-form [1699-46-3]

Alkaloid from *Pachygone somniferum* (Papaveraceae). Noncryst. Mp 190-192° (as picrate).

Billek, G. et al., Monatsh. Chem., 1956, 87, 106-112 (synth, deriv)

Gopinath, K.W. et al., Chem. Ber., 1959, 92, 1657-1661 (synth)

Arndt, R.R. et al., J.C.S., 1964, 2244-2248 (isol, uv)

Albonico, S.M. et al., Annalen, 1965, 685, 200-206 (*Tembetarine*)


Battersby, A.R. et al., J.C.S., 1965, 3323-3332 (synth)

Albonico, S.M. et al., J.C.S.(C), 1966, 1340-1342 (*Tembetarine*, uv, ord, abs config)

- Konda, M. et al., *Chem. Pharm. Bull.*, 1975, **23**, 1063-1076 (*pmr, synth, ms*)
 Bhakuni, D.S. et al., *J.C.S. Perkin 1*, 1977, 1662-1666 (*biosynth*)
 Dasgupta, S. et al., *J. Nat. Prod.*, 1979, **42**, 399-406 (*oxide*)
 Hirsenkorn, R. et al., *Tet. Lett.*, 1990, **31**, 7591-7594 (*synth*)
 Chen, J. et al., *Planta Med.*, 2000, **66**, 251-256 (*activity*)
 Chang, Y.-C. et al., *Z. Naturforsch., C*, 2003, **58**, 521-526 (*isol, pmr, ms*)
 Lee, S.-S. et al., *J. Nat. Prod.*, 2007, **70**, 637-642 (*N-oxides*)
 Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SLX500

Retrofractamides

R-60



Retrofractamide A: n = 1
 B: n = 3
 D: n = 2

Historically, Pipericide was the first alkaloid characterised but they are described here as retrofractamides for ease of presentation.

Retrofractamide A [94079-67-1]

$C_{20}H_{25}NO_3$ 327.422
 Alkaloid from the above-ground parts of *Piper retrofractum* (Javanese long pepper) and the fruits of *Piper nigrum* (pepper). Cryst. (MeOH). Mp 129° (125°).

4,5-Dihydro: Retrofractamide C

[96386-33-3]

 $C_{20}H_{27}NO_3$ 329.438

Alkaloid from *Piper retrofractum* (Javanese long pepper) (Piperaceae). Mp 120°.

2,3,4,5-Tetrahydro: Dihydroretrofractamide C. Dehydroretrofractamide C (*in-corr.*) $C_{20}H_{29}NO_3$ 331.454

Alkaloid from the fruit of *Piper nigrum* (pepper). Powder. λ_{max} 216 ; 261 ; 306 (MeOH).

Hexahydro:

Cryst. (MeOH). Mp 65°. Obt. from either Retrofractamide A or C.

Retrofractamide B

Pipericide

[54794-74-0]

 $C_{22}H_{29}NO_3$ 355.476

Alkaloid from the aerial parts of *Piper retrofractum* (Javanese long pepper) and the fruits of *Piper nigrum* (pepper) (Piperaceae). Insecticidal agent. Cryst. (EtOAc). Mp 114-115°.

10,11-Dihydro: 10,11-Dihydropipericide.

Dihydroretrofractamide B

[75022-26-3]

[85383-60-4]

 $C_{22}H_{31}NO_3$ 357.492

Alkaloid from the fruit of *Piper nigrum* (pepper). Insecticide. Needles (petrol/EtOAc/hexane). Mp 94-95° (89.5-91.5°).

Retrofractamide D [96386-34-4]

 $C_{21}H_{27}NO_3$ 341.449

Alkaloid from *Piper brachystachyum* and *Piper retrofractum* (Javanese long pepper). Amorph. solid. Mp 118-120°. λ_{max} 211 (log ϵ 4.42); 262 (log ϵ 4.5) (EtOH).

Miyakado, M. et al., *Agric. Biol. Chem.*, 1979, **43**, 1609; 1980, **44**, 1701 (*Pipericide, Dihydropipericide*)

Crombie, L. et al., *Tet. Lett.*, 1984, **25**, 4267-4270 (*Pipericide, synth*)

Banerji, A. et al., *Phytochemistry*, 1985, **24**, 279; 1987, **26**, 3345 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Sabharwal, A. et al., *J. Indian Chem. Soc.*, 1990, **67**, 318-320 (*Pipericide, synth*)

Strunz, G.M. et al., *Can. J. Chem.*, 1996, **74**, 419 (*synth, Retrofractamide A*)

Rotherham, L.W. et al., *J.O.C.*, 1998, **63**, 6667-6672 (*Pipericide, Dihydropipericide, synth*)


Park, I.-K. et al., *J. Agric. Food Chem.*, 2002, **50**, 1866-1870 (*isol, pmr, cmr*)

Banerji, A. et al., *Phytochemistry*, 2002, **59**, 897-901 (*Retrofractamide D*)

Rho, M.-C. et al., *Phytochemistry*, 2007, **68**, 899-903 (*Dihydroretrofractamide C*)

Retronecine 9-(2,3-dihydroxy-2-hydroxymethylbutanoate) 7-senecioate

R-61

 $C_{18}H_{27}NO_7$ 369.414

Not named in the lit. Alkaloid from *Senecio caudatus* (Asteraceae).

1'''-Deoxy: Retronecine 9-(2,3-dihydroxy-2-methylbutanoate) 7-senecioate

 $C_{18}H_{27}NO_6$ 353.414

Alkaloid from *Senecio caudatus* (Asteraceae). Not named in the lit.

1'''-Deoxy, N-oxide: Retronecine 9-(2,3-dihydroxy-2-methylbutanoate) 7-senecioate N-oxide

 $C_{18}H_{27}NO_7$ 369.414

Alkaloid from *Senecio caudatus* (Asteraceae).

1'''-Deoxy, O^{3''}-Ac: Retronecine 9-(3-acetoxy-2-hydroxy-2-methylbutanoate) 7-senecioate $C_{20}H_{29}NO_7$ 395.452

Alkaloid from *Senecio caudatus* (Asteraceae). Not named in the lit.

1'''-Deoxy, O^{3''}-Ac, N-oxide: Retronecine 9-(3-acetoxy-2-hydroxy-2-methylbutanoate) 7-senecioate N-oxide $C_{20}H_{29}NO_8$ 411.451

Alkaloid from *Senecio caudatus* (Asteraceae).

2'',3''-Dideoxy, 2'',3''-didehydro (Z-):

Retronecine 9-sarracinate 7-senecioate

 $C_{18}H_{25}NO_5$ 335.399

Alkaloid from *Senecio variabilis*, *Senecio caudatus* and *Senecio triangularis* (Asteraceae). Gum. $[\alpha]_D^{25}$ +11 (c, 1.0 in MeOH). Not named in the lit.

2'',3''-Dideoxy, 2'',3''-didehydro, N-oxide: Retronecine 9-sarracinate 7-senecioate

N-oxide

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Senecio caudatus* and *Senecio umgeniensis* (Asteraceae).

Rüeger, H. et al., *Can. J. Chem.*, 1983, **61**, 2526 (*isol, ir, pmr, cmr, cd, struct*)


Bohlmann, F. et al., *Phytochemistry*, 1986, **25**, 1151 (*isol, ir, ms, struct*)

Retrorsine

R-62

12,18-Dihydroxysenecionan-11,15-dione, 9CI. β -Longilobine

[480-54-6]

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Senecio* spp., *Crotalaria* spp. and *Erechtites quadridentata* (Asteraceae, Fabaceae). Cryst. (EtOH). Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 217° (207-208°). $[\alpha]_D^{20}$ -62.4 (c, 1.0 in $CHCl_3$). $[\alpha]_D$ -17.6 (c, 1.99 in EtOH). λ_{max} 217 (ϵ 7100) (MeOH) (Berdy).

► Hepatotoxin, causes liver and kidney neoplasms. Highly toxic. VH7525000

Nitrate:

Prisms (EtOH/Et₂O). Mp 145°.

Methiodide:

Prisms (MeOH). Mp 266° (260°).

N-Oxide: Isatidine. Retrorsine N-oxide

[15503-86-3]

 $C_{18}H_{25}NO_7$ 367.398

Alkaloid from *Senecio* spp., *Erechtites quadridentata* and *Werneria decora*. Mp 145°. $[\alpha]_D$ -8.2 (H_2O).

► VH7700000

15E-Isomer: Usaramine. Usaramine.

Mucronatine†

[15503-87-4]

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Crotalaria brevifolia*, *Crotalaria incana*, *Crotalaria mucronata* and *Crotalaria usaramoensis* (Fabaceae). Poorly sol. hexane. Mp 182.5-183.5°. $[\alpha]_D^{20}$ +7.1 (c, 1.83 in EtOH). Uv light → Retrorsine.

► Hepato- and pneumotoxin. VT5707200

Stereoisomer: Mucronatinine

[20824-37-7]

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Crotalaria mucronata* (Fabaceae). Cryst. Mp 161-163°. Of undetd. config.

Stereoisomer, picrate: [20824-38-8]

Mp 228-229°.

Aldrich Library of ¹³C and ¹H FT NMR Spectra, 1992, **1**, 1164B (*nmr*)

Manske, R.H.F. et al., *Can. J. Res.*, 1931, **5**, 651 (*isol*)

Barger, G. et al., *J.C.S.*, 1935, **11** (*isol*)

Leisegang, E.C. et al., *J.C.S.*, 1950, 702 (*struct*)


Culvenor, C.C.J. et al., *Aust. J. Chem.*, 1966, **19**, 2127 (*Usaramine*)

Bhacca, N.S. et al., *Tetrahedron*, 1968, **24**, 6319 (*Mucronatine*)

- Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (cd)
- Stoeckli-Evans, H. *et al.*, *Acta Cryst. B*, 1979, **35**, 2798 (cryst struct)
- Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (cmr)
- Molyneux, R.J. *et al.*, *Phytochemistry*, 1982, **21**, 439 (cmr)
- Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1203 (isol)
- Segall, H.J. *et al.*, *Phytochemistry*, 1983, **22**, 1271-1273 (pmr)
- White, J.D. *et al.*, *J.O.C.*, 1989, **54**, 4268 (synth, Usaramine)
- Lock de Ugaz, O. *et al.*, *Phytochemistry*, 1990, **29**, 2373-2375 (N-Oxide, cmr)
- White, J.D. *et al.*, *J.O.C.*, 1992, **57**, 2270 (synth, Usaramine)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, RFP000; RFU000

Retuline**R-63**

1-Acetyl-19,20-didehydrocuran-17-ol, 9CI
[2616-16-2]



Absolute
Configuration

$C_{21}H_{26}N_2O_2$ 338.449
Alkaloid from *Strychnos holstii* var. *reticulata* f. *condensata*, *Strychnos henningsii*, *Strychnos kasengaensis* and some other *Strychnos* spp. (Loganiaceae). Mp 167-173° (170°). $[\alpha]_D^{25} +7$ (+23) (MeOH). $[\alpha]_D^{25} -13.5$ (c, 0.73 in $CHCl_3$).

▶ GS3310000

N^b-Oxide: Retuline N^b-oxide. Oxyretuline
[38769-04-9]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Strychnos camptoneura* and *Strychnos henningsii* (Loganiaceae). Prisms (Et₂O/MeOH). Mp 178-180°.

N-De-Ac: Desacetylretuline. 2β,16β-Dihydroakuammicinol. N-Deacetylretuline
[18397-04-1]
 $C_{19}H_{24}N_2O$ 296.411
Alkaloid from *Strychnos variabilis*, *Strychnos kasengaensis* and other *Strychnos* spp. (Loganiaceae). Cryst. (MeOH/Et₂O). Mp 177° (165°). $[\alpha]_D -7$ (MeOH). $[\alpha]_D -42$ ($CHCl_3$).

O-Ac: O-Acetylretuline

[18397-05-2]
 $C_{23}H_{28}N_2O_3$ 380.486
Alkaloid from bark and leaves of *Strychnos henningsii*, *Strychnos pungens* and *Strychnos kasengaensis* (Loganiaceae). Amorph. powder. Mp 120°.

1,2-Didehydro. N-de-Ac: 1,2-Dehydrodesacetylretuline
[88721-04-4]
 $C_{19}H_{22}N_2O$ 294.396
Structure erroneously assigned to an alkaloid from *Strychnos longicaudata*. For revised struct. see Normavacurine in M-130.

17-Aldehyde: Retulinal

[77096-54-9]
 $C_{21}H_{24}N_2O_2$ 336.433
In tautomeric equilib. with Isoretulinal. Major alkaloid from dried root of *Strychnos variabilis* (Loganiaceae). λ_{max} 212 (log ε 4.4); 251 (log ε 4.07); 282 (log ε 3.46) (MeOH).

17-Carboxylic acid, N-de-Ac, N¹,N⁴-di-Me, Me ester: N^a-Methyl-2,16-dihydroakuammicine N^b-methosalt
[69754-34-3]
 $C_{22}H_{29}N_2O_2^{\oplus}$ 353.483
Quaternary alkaloid from leaves of *Vinca minor* (Apocynaceae). Cryst. (Me₂CO)(as iodide). Mp 310° (iodide).

12-Hydroxy, 17-aldehyde: 12-Hydroxyretulinal
[77053-11-3]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from dried root of *Strychnos variabilis* (Loganiaceae). In tautomeric equilib. with 12-Hydroxyisoretulinal. λ_{max} 217 (log ε 4.05); 252 (log ε 3.45); 288 (log ε 3.08) (MeOH).

11-Methoxy: 11-Methoxyretuline

[94943-26-7]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from *Strychnos kasengaensis*.

16-Epimer: Isoretuline

[10388-62-2]
 $C_{21}H_{26}N_2O_2$ 338.449
Trace alkaloid from *Strychnos variabilis* and *Strychnos kasengaensis* (Loganiaceae).

▶ GS3320000

16-Epimer, N-de-Ac: N^a-Desacetylisoretuline. Isodeacetylretuline. Desacetylisoretuline
[13941-27-0]
 $C_{19}H_{24}N_2O$ 296.411
Alkaloid from *Strychnos variabilis*, *Strychnos henningsii*, *Strychnos floribunda*, *Strychnos kasengaensis*, *Strychnos scheffleri* and other *Strychnos* spp. (Loganiaceae). Prisms (Me₂CO). Mp 177-178°.

16-Epimer, O-Ac: O-Acetylisoretuline

[62860-81-5]
 $C_{23}H_{28}N_2O_3$ 380.486
Alkaloid from *Strychnos variabilis* and *Strychnos kasengaensis*.

16-Epimer, 17-aldehyde: Isoretulinal. N-Acetyl-2,16-dihydroakuammicinol. 18-Deoxydiaboline
[62835-85-2]
 $C_{21}H_{24}N_2O_2$ 336.433
Alkaloid from dried root of *Strychnos variabilis* and from *Strychnos kasengaensis* (Loganiaceae). Mp 192-197° (188-191°). $[\alpha]_D +157$ (MeOH). In tautomeric equilib. with Retulinal.

16-Epimer, 17-aldehyde, N-de-Ac: Desacetylisoretulinal. Dihydrocheminortoxiferine. 16-Deoxy-Wieland-Gumlich aldehyde
[6879-70-5]
 $C_{19}H_{22}N_2O$ 294.396

Alkaloid from *Strychnos variabilis*, *Strychnos dolichothyrsa*, *Strychnos kasangaensis*, *Strychnos matopensis* and *Strychnos minfiensis* (Loganiaceae).

16-Epimer, 12-hydroxy, 17-aldehyde: 12-Hydroxyisoretulinal

[77096-55-0]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from dried root of *Strychnos variabilis* (Loganiaceae). In tautomeric equilib. with 12-hydroxyretulinal.

16-Epimer, 16-hydroxy, 17-aldehyde: 16-Hydroxyisoretulinal

[76177-21-4]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from the root bark of *Strychnos variabilis* (Loganiaceae). Powder.

16-Epimer, 18-hydroxy: 18-Hydroxyisoretuline

$C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from leaves and stems of *Strychnos henningsii* (Loganiaceae). Small prisms (Me₂CO). Mp 245-248°.

16-Epimer, 18-hydroxy, N-de-Ac: N^a-Desacetyl-18-hydroxyisoretuline. Wieland-Gumlich diol

[900-98-1]
 $C_{19}H_{24}N_2O_2$ 312.411
Alkaloid from *Strychnos henningsii*, *Strychnos longicaudata* and *Strychnos kasengaensis* root bark (Loganiaceae). Needles (Me₂CO). Mp 247-250°.

16-Epimer, 18-hydroxy, N-de-Ac, O¹⁷-Ac: 17-O-Acetyl-N^a-desacetyl-18-hydroxyisoretuline

$C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Strychnos henningsii* (Loganiaceae). Noncryst.

16-Epimer, 18-acetoxy, N-de-Ac: 18-Acetoxy-N^a-desacetylisoretuline

[88721-02-2]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from the stem bark of *Strychnos longicaudata* and from *Strychnos kasengaensis* (Loganiaceae). $[\alpha]_D -31$ (c, 0.55 in $CHCl_3$). λ_{max} 218; 246; 300 (EtOH).

16-Epimer, 10-methoxy, N-de-Ac: Tsilanimbine

[59630-35-2]
 $C_{20}H_{26}N_2O_2$ 326.438
Alkaloid from *Strychnos henningsii* leaves and stems (Loganiaceae). Noncryst.

16-Epimer, 11-methoxy: 11-Methoxyisoretuline

[94898-70-1]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from the stem bark of *Strychnos kasengaensis* and from *Strychnos variabilis* (Loganiaceae). $[\alpha]_D +145$ (c, 0.2 in MeOH). λ_{max} 223 (log ε 4.32); 250 (log ε 4.05); 292 (log ε 3.69) (MeOH).


16-Epimer, 11-methoxy, O-Ac: O-Acetyl-11-methoxyisoretuline. 11-Methoxy-O-acetylisoretuline

[127160-69-4]
 $C_{24}H_{30}N_2O_4$ 410.512
 Alkaloid from seeds of *Strychnos variabilis* (Loganiaceae). $[\alpha]_D^{20} +62$ (c, 0.6 in $CHCl_3$).

- Bisset, N.G. *et al.*, *Chem. Ind. (London)*, 1965, 1036-1037 (*uv, ir, pmr, ms, struct*)
 Hymon, J.R. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 2067-2071 (*synth, pmr, ir*)
 Wenkert, E. *et al.*, *J.O.C.*, 1966, **31**, 2689-2691; 1978, **43**, 1099-1105 (*synth, cmr, pmr*)
 Koch, M. *et al.*, *Ann. Pharm. Fr.*, 1972, **30**, 299-306 (*oxide*)
 Angenot, L. *et al.*, *Phytochemistry*, 1975, **14**, 2519-2520 (*Desacetylretuline*)
 Koch, M. *et al.*, *Phytochemistry*, 1976, **15**, 321-324 (*18-Hydroxyisoretuline: 17-Acetyl-N-desacetyl-18-hydroxyisoretuline*)
 Richard, C. *et al.*, *Phytochemistry*, 1976, **15**, 1805-1806 (*Isoretuline, Isoretulinal*)
 Tavernier, D. *et al.*, *Bull. Soc. Chim. Belg.*, 1978, **87**, 595-607 (*pmr*)
 Tits, M. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 92-95 (*18-Deoxydiaboline*)
 Votický, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 123-127 (*N-Methylidihydroakummicine N-methosalt*)
 Tits, M. *et al.*, *Phytochemistry*, 1980, **19**, 1531-1534 (*16-Hydroxyisoretulinal*)
 Tits, M. *et al.*, *Tet. Lett.*, 1980, **21**, 2439-2442 (*Retulinal, 12-Hydroxyretulinal, isol, ms, uv, ir, pmr, tautom, struct*)
 Angenot, L. *et al.*, *Planta Med.*, 1981, **41**, 240-243 (*O-Acetylretuline*)
 Massiot, G. *et al.*, *Tetrahedron*, 1983, **39**, 3645-3656 (*Wieland-Gumlich diol, 18-Acetoxy-N²-desacetylretuline*)
 Thépenier, P. *et al.*, *Phytochemistry*, 1984, **23**, 2659-2663; 1990, **29**, 686-687 (*11-Methoxyretuline, 11-Methoxyisoretuline, O-Acetylretuline, O-Acetyl-11-methoxyisoretuline, Desacetylretulinal, Isoretulinal*)
 Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304 (*cmr*)
 Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, derivs, rev*)

Retusamine, 9CI**R-64**

[6883-16-5]



Absolute configuration

 $C_{19}H_{25}NO_7$ 379.409

Cyclic diester of Otonecine, O-136 with retusaminic acid. Alkaloid from *Crotalaria retusa*, *Crotalaria crassipes*, *Crotalaria mitchellii* and *Crotalaria novae-hollandiae* (Fabaceae). Needles (C_6H_6 /petrol), prisms (EtOH). Mp 174.5° (164-164.5°). $[\alpha]_D^{20} +12$ (c, 2.25 in EtOH).


▶ VH7980000

- Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1957, **10**, 464; 1965, **18**, 1625; 1967, **20**, 801 (*isol, struct, ir, nmr*)
 Wunderlich, J.A. *et al.*, *Acta Cryst. B*, 1967, **23**, 846 (*cryst struct, abs config*)

Kiyooka, S. *et al.*, *Chem. Lett.*, 1973, 963 (*synth*)

Retusine†**R-65**

1,2,14,19-Tetrahydro-12-hydroxy-20-nor-crotalanan-11,15-dione, 9CI
 [480-86-4]


 $C_{16}H_{25}NO_5$ 311.377

Alkaloid from *Crotalaria retusa* and *Crotalaria spectabilis* (Fabaceae). Needles (EtOH or Me_2CO). Mp 174-175°. $[\alpha]_D^{20} +16$ (c, 1 in $CHCl_3$). Hydrol. gives Hexahydro-7-hydroxy-1H-pyrrolizine-1-methanol, H-233 and a mixt. of epimeric α - and β -retusanecic acids (lactone acids derived by cyclisation of the necic acid portion).

- Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1957, **10**, 464 (*isol*)
 Aasen, A.J. *et al.*, *J.O.C.*, 1969, **34**, 4137 (*pmr, stereochem*)
 Kiyooka, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 3609 (*config*)
 Smith, L.W. *et al.*, *Aust. J. Chem.*, 1988, **41**, 429

Reumycin**R-66**

6-Methylpyrimido[5,4-e]-1,2,4-triazine-5,7(1H,6H)-dione, 9CI. 1-Demethyltoxoflavin. Reumitsin
 [5016-18-2]

 $C_8H_5N_5O_2$ 179.138

Isol. from *Actinomyces rectus bruneus*. Antineoplastic agent. Yellow powder or cryst. Sol. MeOH, H_2O , EtOAc, DMF; fairly sol. Me_2CO , C_6H_6 ; poorly sol. hexane, Et_2O . Mp 210° dec
 Mp 244-245°. λ_{max} 235 (ε 17800); 265 (sh) (ε 3090); 340 (ε 5248); 400 (ε 741) (EtOH) (Derep). λ_{max} 258 (ε 15130); 400 (ε 2630) (EtOH/NaOH) (Berdy).

▶ UW7930000

1H-form

1-N-Me: 1,6-Dimethylpyrimido[5,4-e]-1,2,4-triazine-5,7(1H,6H)-dione. 1,5,6,7-Tetrahydro-1,6-dimethyl-5,7-dioxopyrimido[5,4-e]-as-triazine. Toxoflavin. Xanthothricin. 1-Methylreumycin
 [84-82-2]
 $C_7H_7N_5O_2$ 193.165
 Prod. by *Pseudomonas cocovenenans*

and *Burkholderia gladioli*. Shows broad-spectrum antibiotic activity but is highly toxic. Shows phytotoxic activity. Yellow needles (EtOH or MeOH). Sol. H_2O , EtOH. Mp 172°. λ_{max} 258 (ε 18400); 397 (ε 4500) (pH 1 H_2O) (Derep). λ_{max} 322 (ε 6400) (pH 11) (Derep). λ_{max} 259 (ε 22500); 330 (sh) (ε 1950); 395 (ε 5500) (EtOH) (Derep).

▶ UW7875000

1-N-Me, 4-oxide: [32496-16-5]

Yellow needles (EtOH or dioxan). Mp 189° (215°).

7OH-form

7-Me ether: 7-Methoxy-6-methylpyrimido[5,4-e]-1,2,4-triazin-5(6H)-one, 9CI. 7-Methoxyreumycin
 [51590-95-5]

 $C_7H_7N_5O_2$ 193.165

From *Actinomyces rectus bruneus*. Antineoplastic agent. Sol. MeOH, EtOAc, DMF, H_2O ; fairly sol. Me_2CO , C_6H_6 ; poorly sol. hexane, Et_2O .

Van Damme, P.A. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1960, **79**, 255 (*Toxoflavin, struct, uv, ir*)

Latuasan, H.E. *et al.*, *Biochim. Biophys. Acta*, 1961, **52**, 502 (*Toxoflavin, tox*)

Hellendoorn, A.J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1961, **80**, 307 (*Toxoflavin, cryst struct*)

Daves, G.D. *et al.*, *J.A.C.S.*, 1962, **84**, 1724 (*synth*)

Levenberg, P. *et al.*, *J. Biol. Chem.*, 1966, **241**, 846 (*biosynth*)

Esipov, S.E. *et al.*, *J. Antibiot.*, 1973, **26**, 537 (*struct*)

Yoneda, F. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2001-2009 (*synth, ir, uv, nmr*)

Yoneda, F. *et al.*, *J.C.S. Perkin 1*, 1976, 713


Adanin, V.M. *et al.*, *Khim. Geterotsikl. Soedin.*, 1979, 1270 (*ms*)

Voragen, A.G.J. *et al.*, *Food Chem.*, 1982, **9**, 167 (*Toxoflavin, hplc*)

Esipov, S.E. *et al.*, *Antibiot. Med. Biotechnol.*, 1987, **32**, 111; 116 (*ir, pmr, struct*)

Furuya, N. *et al.*, *J. Fac. Agric., Kyushu Univ.*, 1997, **22**, 33-37; *CA*, **128**, 214320q (*Toxoflavin, isol, activity*)

Koshino, H. *et al.*, *Heterocycles*, 2000, **52**, 811-817 (*Toxoflavin, N-15 nmr*)

Reutericyclin**R-67** $C_{20}H_{31}NO_4$ 349.469

Tetramic acid. Tautomeric β -triketonic system; one tautomer shown.

(R)-form [303957-69-9]


Prod. by *Lactobacillus reuteri* LTH2584. Active against gram-positive bacteria. Mp 161-164° (synthetic). $[\alpha]_D^{20} +13$ (c, 0.29 in EtOH) (synthetic).

λ_{\max} 238 ; 286 (MeCN). λ_{\max} 228 (€ 17870); 266 (€ 23875); 278 (€ 25165) (EtOH).

Höltzel, A. et al., *Angew. Chem., Int. Ed.*, 2000, **39**, 2766-2768 (*pmr, cmr, struct*)
 Gänzle, M.G. et al., *Appl. Environ. Microbiol.*, 2000, **66**, 4325-4333 (*isol*)
 Marquardt, U. et al., *Synlett*, 2000, 1131-1132 (*synth*)
 Böhme, R. et al., *Helv. Chim. Acta*, 2005, **88**, 2837-2841 (*synth*)
 Schobert, R. et al., *Synthesis*, 2006, 3902-3914 (*synth, ir, pmr, cmr, ms*)

Revolutinone
R-68

[74046-19-8]



 $C_{38}H_{40}N_2O_8$ 652.743

Trace alkaloid from the fruit of *Thalictrum revolutum* (Ranunculaceae), also obt. by oxidn. of *O*-Methylthalicberine in T-313. Amorph. $[\alpha]_D^{25}$ -10 (c, 0.5 in MeOH).

Wu, J. et al., *J. Nat. Prod.*, 1980, **43**, 270 (*isol, uv, cd, ir, pmr, ms, struct, synth*)

Rexostatine
R-69

3-[[[1-[[[4-[(Aminoiminomethyl)amino]butyl]amino]carbonyl]-3-methylbutyl]amino]carbonyl]oxiranecarboxylic acid, 9CI. Proteinase inhibitor E 64. E 64 [66701-25-5]


 $C_{15}H_{27}N_5O_5$ 357.409

Isol. from *Aspergillus japonicus*. Thiol proteinase inhibitor. Sensitizes human lung and colon carcinomas to antineoplastic chemotherapy. Needles (Me₂CO aq.). Sol. MeOH aq., AcOH, DMSO, Py; fairly sol. EtOH; poorly sol. Me₂CO, hexane. $[\alpha]_D^{25}$ +24.4 (c, 1 in 0.1M HCl). Log P -1.71 (calc). Dec. at 231-234°.


► Exp. teratogen. RR0390000

[66701-24-4]

Hanada, K. et al., *Agric. Biol. Chem.*, 1978, **42**, 523; 529; 537 (*isol, synth, struct, props*)
 Hanada, K. et al., *Pept. Chem.*, 1979, **17**, 31 (*synth, props*)
 Barrett, A.J. et al., *Acta Biol. Med. Ger.*, 1981, **40**, 1513 (*rev. pharmacol*)
 Barrett, A.J. et al., *Biochem. J.*, 1982, **201**, 189 (*props*)
 Yamamoto, D. et al., *Chem. Pharm. Bull.*, 1989, **37**, 2577; 1990, **38**, 2339 (*cryst struct, conformn*)
 Tachikura, T. et al., *Acta Paediatr. Jpn.*, 1990, **32**, 495 (*tox*)
 Jani, J.P. et al., *Cancer Res.*, 1992, **52**, 2931
 Sarabia, F. et al., *Bioorg. Med. Chem.*, 2005, **13**, 1691-1705 (*synth*)

Rhappallin A
R-70

5-Methyl-1,3,7-triazecine-2,6,8(1H,3H,7H)-trione, 9CI. 5-Methyl-1,3,7-triazacyclodeca-4,9-diene-2,6,8-trione



(E,E)-form

 $C_8H_9N_3O_3$ 195.177

(E,E)-form

 Isol. from the sponge *Rhaphisia pallida*.

(Z,Z)-form [193224-03-2]

 Alkaloid from the sponge *Rhaphisia pallida*.

*N*⁷-Bromo-7-Bromo-5-methyl-1,3,7-triazecine-2,6,8(1H,3H,7H)-trione [193224-05-4]

 $C_8H_8BrN_3O_3$ 274.074


 Alkaloid from the sponge *Rhaphisia pallida*.

Cen, Y.Z. et al., *CA*, 1997, **127**, 147229f (*isol, uv, pmr, cmr, ms*)

Zeng, L. et al., *Pure Appl. Chem.*, 1999, **71**, 1147-1151

Rhopsamine
R-71

1,28-Bis[(3-amino-2-hydroxypropyl)amino]-3,6,9,12,15,18,25-octacosaeptaene [198826-15-2]


 $C_{34}H_{60}N_4O_2$ 556.874

Alkaloid from the sponge *Leucetta leptorhopsis*. Amorph. solid. Racemic. λ_{\max} 232 (€ 6340) (MeOH).


Jayatilake, G.S. et al., *Tet. Lett.*, 1997, **38**, 7507-7510 (*isol, uv, pmr, cmr*)

Rhazidigenine
R-72

7-Ethyl-1,2,4,5,6,7,8,9-octahydro-14bH-3,7-methanoazacycloundecino[5,4-b]indol-14b-ol, 9CI. Rhazidine base. Strictanol

[3384-38-1]

[109063-87-8]



Absolute configuration

 $C_{19}H_{26}N_2O$ 298.427

Strictanol appears to be identical with the prev. descr. Rhazidine. The opt. rotn. reported is similar to that of the hydrochloride. Alkaloid from *Rhazya stricta*, *Aspidosperma quebracho-blanco*, *Amsonia angustifolia* and *Gonioma kamassi*. Mp 187°. $[\alpha]_D$ -612 (EtOH/NaOH). λ_{\max} 210 (log € 4.33); 281 (log € 3.66); 292 (log € 3.62); 307 (log € 3.58) (heptane).

Hydrochloride: Rhazidine. Rhazidine salt [15381-61-0]

Mp 285° dec. $[\alpha]_D$ -40 (neutral or acidic EtOH) ((-37)). Quaternary pentacyclic structure. λ_{\max} 236 (log € 3.91); 293 (log € 3.38) (EtOH).

N^b-Oxide: Rhazidigenine *N*^b-oxide

[26066-43-3]

 $C_{19}H_{26}N_2O_2$ 314.427

Alkaloid from the seeds of *Rhazya stricta* and the bark of *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Noncryst. Mp 240-245°. λ_{\max} 228 ; 285 ; 293 ; 308 (no solvent reported).

N^b-Oxide; hydrochloride: Mp 325-327°.

λ_{\max} 237 (log € 3.89); 293 (log € 3.29) (no solvent reported).

Spiteller-Friedmann, M. et al., *Monatsh. Chem.*, 1964, **95**, 1228 (*uv, ir, ms, struct*)

Markey, S. et al., *Tet. Lett.*, 1967, 157 (*uv, struct*)


Tunmann, P. et al., *Z. Naturforsch., B*, 1969, **24**, 1665 (*isol, uv, ir, ms, struct, oxide*)

Miana, G.A. et al., *J. Chem. Soc. Pak.*, 1982, **4**, 119; *CA*, **97**, 159507v (*isol*)

Atta-ur-Rahman, et al., *Phytochemistry*, 1987, **26**, 589-591 (*Strictanol*)

Rhazimine
R-73

[93772-08-8]



Absolute Configuration

 $C_{21}H_{22}N_2O_3$ 350.416

Struct. revised in 1987. Rearranged indole alkaloids which are remotely related to the Akuammiline type. Alkaloid from the leaves of *Rhazya stricta* and *Melodinus acutiflorus* and from *Amsonia sinensis*. Inhibitor of arachidonic acid metabolism and of platelet activating factor. Mp 170.5° (168°) dec. $[\alpha]_D$ +131 (CHCl₃). $[\alpha]_D^{22}$ +252 (c, 0.23 in CHCl₃).

1,2-Dihydro, 2-ξ-methoxy: 1,2-Dihydro-2-methoxyrhazimine. 2-Methoxy-1,2-dihydro-rhazimine

[98264-40-5]

 $C_{22}H_{26}N_2O_4$ 382.458

Alkaloid from leaves of *Rhazya stricta* (Apocynaceae). Hygroscopic cryst. $[\alpha]_D$ +85 (CHCl₃). Extremely labile, transformed to Rhazimine after 2-3 hr. at 30° in CHCl₃ soln. The struct. of this compd. is presumed to be as shown following the struct. revision of Rhazimine.

19*Z*-Isomer, 5α-methoxy: (Z)-5α-Methoxyrhazimine

[214263-41-9]

 $C_{22}H_{24}N_2O_4$ 380.443

Alkaloid from *Alstonia villosa*. Prisms. Mp 260-263°. $[\alpha]_D^{25}$ +366.6 (c, 0.06 in CHCl₃). λ_{\max} 224 (log € 4.24); 270 (log € 3.79); 290 (sh) (log € 3.72) (MeOH).


Atta-ur-Rahman, et al., *Tet. Lett.*, 1984, **25**, 3913-3916 (*isol, uv, ir, pmr, cmr, ms*)

- Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1985, **24**, 1625-1626 (1,2-dihydro 2-methoxy)
 Hu, W.-L. *et al.*, *Phytochemistry*, 1987, **26**, 2625-2630 (isol, uv, ir, pmr, cmr, ms, struct)
 Saeed, S.A. *et al.*, *Planta Med.*, 1993, **59**, 566-568 (activity)
 Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1235-1238 (5-Methoxyrhazimine)

Rhazinal

R-74

[197141-77-8]



Absolute Configuration


$C_{20}H_{22}N_2O_2$ 322.406
 Alkaloid from *Kopsia singapurensis*.
 Light yellow oil. $[\alpha]_D^{25}$ -187 (c, 0.19 in $CHCl_3$). λ_{max} 203 ; 234 ; 302 (no solvent reported).

- Kam, T.-S. *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 307-310 (isol, pmr, cmr)
 Banwell, M.G. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 296-305 (synth)
 Banwell, M.G. *et al.*, *ARKIVOC*, 2006, **iii**, 163-174 (synth)
 Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1783-1789 (isol)

Rhazinaline

R-75

[60706-90-3]



Absolute Configuration

$C_{21}H_{22}N_2O_3$ 350.416
 Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Stout needles (C_6H_6 /petrol). Mp 137°. $[\alpha]_D^{25}$ +61 (EtOH).

N^b-Oxide: Rhazinaline N⁴-oxide

[154879-27-3]

 $C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from leaves of *Leuconotis eugenifolius* (Apocynaceae). Solid. $[\alpha]_D^{27}$ -87.3 (c, 0.70 in MeOH).

16-Epimer: 16-Formylstrictamine. Rhazimal. 16-Epirhazinaline

[73466-11-2]

 $C_{21}H_{22}N_2O_3$ 350.416

Alkaloid from *Rhazya stricta* (Apocynaceae). Minor alkaloid from leaves of *Melodinus acutiflorus*. Mp 101-102°. $[\alpha]_D$ +57 (MeOH). $[\alpha]_D^{20}$ +16.19 ($CHCl_3$). λ_{max} 222 (ϵ 14100); 265 (ϵ 3660) (MeOH).

16-Epimer; (Z)-isomer, 5 α -methoxy: (Z)-16-Formyl-5 α -methoxystrictamine $C_{22}H_{24}N_2O_4$ 380.443


Alkaloid from *Alstonia villosa*. Solid. $[\alpha]_D^{25}$ -18.4 (c, 0.06 in MeOH). λ_{max} 217 (log ϵ 4.18); 230 (sh) (log ϵ 4.09); 297

(log ϵ 3.65) (MeOH).

- Chatterjee, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 2000 (isol, uv, ir, pmr, ms, struct)
 Ahmad, Y. *et al.*, *Phytochemistry*, 1983, **22**, 1017-1019 (Rhazimal)
 Atta-ur-Rahman, *et al.*, *Planta Med.*, 1986, 230-231 (pmr, cmr, Rhazimal)
 Hu, W.-L. *et al.*, *Planta Med.*, 1988, **54**, 235-236 (epimer)
 Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 169 (oxide)
 Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1235-1238 (16-Formyl-5-methoxystrictamine)

Rhazinilam

R-76



(R)-form

 $C_{19}H_{22}N_2O$ 294.396**(R)-form [36193-36-9]**

Isol. from *Rhazya stricta*, *Aspidosperma quebracho-blanco* (quebracho), *Leuconotis eugenifolia* and *Melodinus australis* (Apocynaceae). Antimitotic, spindle poison, antitubulin. Mp 176° Mp 214-216°. Artifact.

5,21-Dihydro: 5,21-Dihydrorhazinilam

[109305-80-8]

 $C_{19}H_{24}N_2O$ 296.411

Alkaloid from *Leuconotis eugenifolia*.
 Converted to Rhazinilam on prolonged exp. to air.

3-Oxo: Rhazinicine. 3-Oxorhazinilam

[197141-93-8]

 $C_{19}H_{20}N_2O_2$ 308.379

Alkaloid from *Kopsia dasyrachis* and a *Rauwolfia serpentina* x *Rhazya stricta* hybrid. Powder. Mp 201-204°. $[\alpha]_D^{25}$ -247.2 (c, 0.3 in $CHCl_3$). λ_{max} 207 (log ϵ 4.37); 229 (sh) (log ϵ 4.19); 275 (sh) (log ϵ 3.61) (EtOH).

3-Oxo, 14,15-didehydro: 14,15-Didehydro-3-oxorhazinilam. 3-Oxo-14,15-dehydrorhazinilam. AQC-3

[139955-87-6]

 $C_{19}H_{18}N_2O_2$ 306.363

Alkaloid from *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Amorph.

(±)-form [52745-52-5]

Synthetic. Mp 222-223°.


- Banerji, A. *et al.*, *Phytochemistry*, 1970, **9**, 1491-1493 (isol, uv, ir, pmr, ms)
 Abraham, D.J. *et al.*, *Tet. Lett.*, 1972, 909-912 (cryst struct)
 De Silva, K.T. *et al.*, *Tet. Lett.*, 1972, 913-916 (pmr, struct)
 Ratcliffe, A.H. *et al.*, *Tet. Lett.*, 1973, 5179-5184 (synth)
 Goh, S.H. *et al.*, *Tet. Lett.*, 1986, **27**, 2501-2504 (5,21-Dihydrorhazinilam)
 Aimi, N. *et al.*, *Tet. Lett.*, 1991, **32**, 4949-4952 (3-Oxo-14,15-dehydrorhazinilam)
 Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 159-169 (Rhazinicine)
 Gerasimenko, I. *et al.*, *J. Nat. Prod.*, 2001, **64**, 114-116 (Rhazinicine)

- Magnus, P. *et al.*, *Tetrahedron*, 2001, **57**, 8647-8651 (synth)
 Johnson, J.A. *et al.*, *J.A.C.S.*, 2002, **124**, 6900-6903 (synth)
 Bowie, A.L. *et al.*, *Org. Lett.*, 2005, **7**, 5207-5209 (synth)
 Banwell, M.G. *et al.*, *ARKIVOC*, 2006, **iii**, 163-174 (synth)

Rhazizine

R-77

[123871-29-4]

 $C_{21}H_{24}N_2O_4$ 368.432


Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). $[\alpha]_D$ +84 (MeOH).

Atta-ur-Rahman, *et al.*, *Tetrahedron*, 1989, **45**, 3507 (isol, uv, ir, pmr, cmr, ms, struct)

Rhexifoline

R-78

[93915-32-3]

 $C_{11}H_{13}NO_3$ 207.229


Alkaloid from *Castilleja rhexifolia* and *Castilleja rhexifolia* aff. *miniata* (Scrophulariaceae). $[\alpha]_D^{23}$ +18 (c, 0.82 in $CHCl_3$). λ_{max} 225 ; 273 (EtOH).

Roby, M.R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 846-853 (isol, uv, ir, pmr, cmr, ms, struct)

Rhinolidine

R-79

20-(Hexahydro-1-methyl-1H-azepin-2-yl)pregn-5-ene-3,15-diol, 9CI
 [70474-40-7]

 $C_{28}H_{47}NO_2$ 429.685

Struct. dubious. More likely to be a piperidine related to Sevcordinine, S-275 or a solanidane. No further reports to 2006. Alkaloid from *Fritillaria valujevii* (Liliaceae). Mp 199-201°. $[\alpha]_D$ -52.9 (EtOH).

3-O-β-D-Glucopyranoside: Rhinoline

[70474-39-4]

 $C_{34}H_{57}NO_7$ 591.827

Alkaloid from *Rhinopetalum bucharicum*, *Rhinopetalum karelini* and *Fritillaria valujevii* (Liliaceae). Mp 255-257°.

$[\alpha]_D^{20}$ -53.2 (EtOH).

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside]: **Rhinoline**
[72060-81-2]


C₄₀H₆₇NO₁₂ 753.969

Alkaloid from *Rhinopetalum bucharicum* and *Fritillaria valujevii* (Liliaceae).
Mp 301-302°. $[\alpha]_D$ -36.5.

Samikov, K. et al., *Khim. Prir. Soedin.*, 1978, **14**, 815-817; 1979, **15**, 350-353; 1981, **17**, 530; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 701-702; 1979, **15**, 303-305 (*Rhinoline, Rhinoline*)

Rhizoferrin R-80

2,2'-[1,4-Butanediy]bis[imino(2-oxo-2,1-ethanediy)]bis[2-hydroxybutanedioic acid]. N¹,N⁴-Bis(4,4-dicarboxy-3-hydroxy-1-oxobutyl)-1,4-butanediamine
[138846-62-5]



C₁₆H₂₄N₂O₁₂ 436.372
 λ_{\max} 200 ; 280 (H₂O) (Berdy).

(R,R)-form

Prod. by *Rhizopus microsporus* var. *rhizopodiformis*, *Rhizopus arrhizus* and *Cunninghamella elegans*. Siderophore, growth promotor. Sol. H₂O. $[\alpha]_D^{26}$ -16.7 (c, 0.16 in MeOH).

(S,S)-form

Enantiomorph

[251366-27-5]

Prod. by *Ralstonia pickettii* DSM 6297. Siderophore. Glass.

Drechsel, H. et al., *Biol. Met.*, 1991, **4**, 238;

1992, **5**, 141 (*isol, uv, ir, pmr, cmr, ms*)

Thieken, A. et al., *FEMS Microbiol. Lett.*, 1992, **94**, 37 (*occur*)

Shenker, M. et al., *Soil Sci. Soc. Am. Proc.*,

1995, **59**, 837 (*isol, pmr, cmr, cd*)

Bergeron, R.J. et al., *Tetrahedron*, 1997, **53**,


427-434 (*synth, pmr, cd, abs config*)

Munzinger, M. et al., *BioMetals*, 1999, **12**,

189-193 (*Enantiomorph*)

Rhizopodin R-81

[150346-23-9]



C₇₈H₁₂₄N₄O₂₂ 1469.851

Major struct. revision in 2008. Prod. by *Mycrococcus stipitatus* strain Mxf 164. Antitumour agent. Protein phosphorylation inhibitor. Amorph. solid. Sol. MeOH, EtOH, Me₂CO; fairly sol. CHCl₃, Et₂O; poorly sol. H₂O, hexane.

$[\alpha]_D^{20}$ -53.4 (c, 1 in MeOH). λ_{\max} 231 (log ϵ 4.65); 241 (sh) (log ϵ 4.49) (MeOH).

Sasse, F. et al., *J. Antibiot.*, 1993, **46**, 741-748


(*isol, uv, cmr, props*)

Jansen, R. et al., *Tet. Lett.*, 2008, **49**, 5796-

5799 (*isol, pmr, cmr*)

Rhizopterin R-82

4-[(2-Amino-1,4-dihydro-4-oxo-6-pteridinyl)methyl]formylaminobenzoic acid, 9CI. 12-Formylpteroic acid
[119-20-0]



C₁₅H₁₂N₆O₄ 340.298

Occurs in *Rhizopus nigricans*. Folic acid growth factor. Pale yellow needles (AcOH aq.). Insol. H₂O, org. solvs.; sol. acids and alkalis. Mp 300°.

Benzoyl:

Cryst. (AcOH aq.). Mp 250° dec.

Rickes, E.L. et al., *J.A.C.S.*, 1947, **69**, 2749;

2751 (*isol, uv*)


Wolf, D.E. et al., *J.A.C.S.*, 1947, **69**, 2753

Rhizoxin R-83

WF 1360. Antibiotic WF 1360. NSC

332598

[90996-54-6]



C₃₅H₄₇NO₉ 625.758

Macrolide antibiotic. Originally thought to be prod. by *Rhizopus microsporus* but it has been shown to be prod. by its bacterial endosymbiont *Burkholderia rhizoxina*. Also prod. by *Pseudomonas fluorescens* Pf-5. Antifungal agent. Phytotoxic to rice seedlings. Inhibits microtubule assembly. Pale-yellow powder. Mp 131-135°. $[\alpha]_D^{24}$ +155 (c, 0.8 in MeOH). λ_{\max} 230 (ϵ 9000); 240 (sh) (ϵ 8000); 295 (ϵ 42300); 308 (ϵ 54000); 325 (ϵ 39000) (MeOH) (Derep).

► VI1800000

O-De-Me: 17-De-O-methylrhizoxin

[103528-74-1]

C₃₄H₄₅NO₉ 611.731

Prod. by *Burkholderia rhizoxina* and *Rhizopus chinensis*. Active against fungi and tumours. Plant growth inhibitor. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{24}$ +116.3 (c, 0.11 in MeOH). λ_{\max} 297 (ϵ 32300); 309 (ϵ 41000); 323 (ϵ 30180) (MeOH) (Berdy).

2,3-Deepoxy, 2,3-didehydro: **Antibiotic**

WF 1360F. WF 1360F. FR 900216F.

Antibiotic FR 900216F

[103528-75-2]

C₃₅H₄₇NO₈ 609.758

Prod. by *Burkholderia rhizoxina* and *Pseudomonas fluorescens* Pf-5. Active against fungi and tumours. Powder. $[\alpha]_D^{24}$ +136.8 (c, 0.71 in MeOH). $[\alpha]_D^{23}$ +97 (c, 1.9 in CHCl₃).

2,3-Deepoxy, 2,3-didehydro, O-de-Me: **Antibiotic WF 1360B. WF 1360B. FR 900216B. Antibiotic FR 900216B**
[103528-76-3]

C₃₄H₄₅NO₈ 595.731

Prod. by *Burkholderia rhizoxina* and *Pseudomonas fluorescens* Pf-5. Active against fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{24}$ +99.3 (c, 0.61 in MeOH). $[\alpha]_D^{23}$ +110 (c, 1.9 in CHCl₃). λ_{\max} 297 (ϵ 39100); 308 (ϵ 50300); 323 (ϵ 36600) (MeOH) (Berdy).

2,3:11,12-Bis(deepoxy), 2,3,11,12-tetra-dehydro: **Rhizoxin D**
[103528-77-4]

C₃₅H₄₇NO₇ 593.759

Prod. by *Burkholderia rhizoxina* and *Pseudomonas fluorescens* Pf-5. Active against fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{24}$ +287.1 (c, 0.48 in MeOH). λ_{\max} 216 (ϵ 25900); 235 (ϵ 24700); 238 (ϵ 24500); 297 (ϵ 38700); 309 (ϵ 49900); 323 (ϵ 36500) (MeOH) (Berdy).

2,3:11,12-Bis(deepoxy), 2,3,11,12-tetra-dehydro, O-de-Me: **Rhizoxin D₁. Antibiotic WF 1360C. WF 1360C. FR 900216C. Antibiotic FR 900216C**
[103528-78-5]

C₃₄H₄₅NO₇ 579.732

Prod. by *Burkholderia rhizoxina* and *Pseudomonas fluorescens* Pf-5. Active against fungi and tumours. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{24}$ +246 (c, 0.66 in MeOH). $[\alpha]_D^{23}$ +197 (c, 0.2 in CHCl₃). λ_{\max} 215 (ϵ 28400); 232 (ϵ 26200); 238 (ϵ 25700); 297 (ϵ 42500); 308 (ϵ 55100); 323 (ϵ 40300) (MeOH) (Berdy).

Iwasaki, S. et al., *J. Antibiot.*, 1984, **37**, 354-362; 1986, **39**, 424-429; 1987, **40**, 66-72 (*isol, uv, ir, pmr, cmr, bibl*)

Tsuruo, T. et al., *Cancer Res.*, 1986, **46**, 381-385 (*props*)

Iwasaki, S. et al., *Chem. Comm.*, 1986, 1702-1703 (*biosynth*)

Iwasaki, S. et al., *Chem. Pharm. Bull.*, 1986, **34**, 1387-1390 (*derivs*)

Kiyoto, S. et al., *J. Antibiot.*, 1986, **39**, 762-772 (*WF 1360*)

Li, Y. et al., *Biochem. Biophys. Res. Commun.*, 1992, **187**, 722-729 (*Rhizoxin, activity*)

Boger, D.L. et al., *J.O.C.*, 1992, **57**, 2235-2244 (*synth, bibl*)

Nakada, M. et al., *Tet. Lett.*, 1993, **34**, 1035-1038; 1039-1042 (*synth*)

Keck, G.E. et al., *Angew. Chem., Int. Ed.*,

2001, **40**, 231-234 (*Rhizoxin D, synth*)

White, J.D. et al., *J.O.C.*, 2002, **67**, 7750-7760 (*Rhizoxin D, synth*)

Lafontaine, J.A. et al., *J.O.C.*, 2003, **68**, 4215-4234 (*Rhizoxin D, synth*)

Jiang, Y. et al., *Org. Lett.*, 2004, **6**, 1445-1448 (*Rhizoxin D, synth*)


Hong, J. et al., *Tetrahedron*, 2004, **60**, 5653-5681 (*rev*)

Mitchell, I.S. et al., *Org. Biomol. Chem.*, 2005, **3**, 4412-4431 (*Rhizoxin D, synth*)

- Scherlach, K. *et al.*, *J.A.C.S.*, 2006, **128**, 11529-11536 (*Burkholderia*, *isol*, *pmr*, *cmr*)
 Partida-Martinez, L.P. *et al.*, *ChemBioChem*, 2007, **8**, 41-45 (*biosynth*)
 Loper, J.E. *et al.*, *Appl. Environ. Microbiol.*, 2008, **74**, 3085-3093 (*Pseudomonas*, *isol*)

Rhizoxin S₁

[911676-00-1]

C₃₄H₄₇NO₉ 613.747

Closely related to Rhizoxin, R-83. Prod. by *Burkholderia rhizoxina*, the bacterial endosymbiont of *Rhizopus microsporus*. Antitumour agent. Amorph. powder. $[\alpha]_D^{25} +31$ (c, 0.08 in MeOH). λ_{max} 299 (log ϵ 4.29); 310 (log ϵ 4.4); 324 (log ϵ 4.3) (MeOH).

Me ester: Rhizoxin M₁

[911676-02-3]

C₃₅H₄₉NO₉ 627.773

Potent antitumour agent. Amorph. powder. $[\alpha]_D^{25} +32$ (c, 0.1 in MeOH).

17-Me ether: Rhizoxin S₂

[308285-11-2]

C₃₅H₄₉NO₉ 627.773

Prod. by *Burkholderia rhizoxina*, the bacterial endosymbiont of *Rhizopus microsporus*. Amorph. powder. $[\alpha]_D^{25} +41$ (c, 0.13 in MeOH). λ_{max} 298 (log ϵ 4.42); 310 (log ϵ 4.52); 324 (log ϵ 4.39) (MeOH).

17-Me ether, Me ester: Rhizoxin M₂

[911676-03-4]

C₃₆H₅₁NO₉ 641.8

Potent antitumour agent. Amorph. powder. $[\alpha]_D^{25} +36$ (c, 0.3 in MeOH).

Deepoxy, 11,12-didehydro(E-): Rhizoxin D₃

[911676-06-7]

C₃₄H₄₇NO₈ 597.747

Prod. by *Burkholderia rhizoxina*, the bacterial endosymbiont of *Rhizopus microsporus*. Amorph. powder.

22Z-Isomer, 17-Me ether: Rhizoxin Z₁

[911676-04-5]

C₃₅H₄₉NO₉ 627.773


Prod. by *Burkholderia rhizoxina*, the bacterial endosymbiont of *Rhizopus microsporus*. Amorph. powder. $[\alpha]_D^{25} +116$ (c, 0.07 in MeOH). λ_{max} 298 (sh); 310 (log ϵ 4.55); 324 (log ϵ 4.43) (MeOH).

22Z-Isomer, 17-Me ether, Me ester: Rhizoxin Z₂

[911676-05-6]

C₃₆H₅₁NO₉ 641.8

Potent antitumour agent. Amorph. powder. $[\alpha]_D^{25} +138$ (c, 0.15 in MeOH).


Scherlach, K. *et al.*, *J.A.C.S.*, 2006, **128**,11529-11536 (*isol*, *pmr*, *cmr*)**Rhodesiacridone****R-85**C₂₀H₁₉NO₇ 385.373**(-)-form**

Alkaloid from the roots of *Thamnosma rhodesica*. Amorph. yellow powder. $[\alpha]_D -47.7$ (c, 0.1 in MeOH). λ_{max} 249 (log ϵ 3.96); 265 (sh) (log ϵ 4.01); 272 (log ϵ 4.09); 299 (log ϵ 3.73); 330 (log ϵ 3.44); 397 (log ϵ 3.18) (MeOH).

Ahua, K.M. *et al.*, *Phytochemistry*, 2004, **65**, 963-968 (*isol*, *pmr*, *cmr*)

Rhoeadine**R-86**

8-Methoxy-16-methyl-2,3:10,11-bis[methylenebis(oxy)]rheadan, 9CI. Rhoeadine. O-Methylrhoeagenine. N-Methylpapaverrubine E

**(+)-form**C₂₁H₂₁NO₆ 383.4

Various numbering schemes in use.

(+)-form [2718-25-4]

Alkaloid from *Bocconia frutescens*, *Mecconopsis betonicifolia* and other *Mecconopsis* spp., *Papaver albiflorum* and other *Papaver* spp. (Papaveraceae). Increases intraocular pressure in expt. animals, causes mydriasis. *P. rhoeas* extracts have been used in medicine as sedatives and expectorants but this does not appear to be caused by Rhoeadine. Needles (MeOH or CHCl₃/MeOH). Mp 253-254°. $[\alpha]_D^{22} +228.5$ (c, 1.008 in CHCl₃).

► Toxic, LD₅₀ (rat, ipr) 530 mg/kg. VI0650000

N-De-Me: Papaverrubine E. N-De-methylrhoeagenine

[6807-95-0]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Bocconia frutescens*, *Mecconopsis* spp. and many *Papaver* spp. (Papaveraceae). Mp 230-231°. $[\alpha]_D +331$ (CHCl₃).

O-De-Me: Rhoegenine. Rheagenine

[5574-77-6]

C₂₀H₁₉NO₆ 369.373

In some solvents the alkaloid exists as a mixt. of C-14 epimers (hemiacetal equilibrium). Alkaloid from *Papaver arenarium*, *Papaver argemone*, *Papaver*

armeniaceum, *Papaver atlanticum*, *Papaver californicum*, *Papaver commutatum*, *Papaver dubium*, *Papaver hispidum*, *Papaver latericum*, *Papaver nudicaule*, *Papaver oreophilum*, *Papaver pilosum*, *Papaver rhoeas* (corn poppy), *Papaver rupifragum*, *Papaver strigosum*, *Papaver syriacum* and *Papaver tauricola*. Mp 236-238°. $[\alpha]_D^{22} +134$ (c, 0.96 in Py).

O-De-Me, N-Me:

Needles (MeOH) (as iodide). Mp 255-258° dec. (iodide). $[\alpha]_D^{22} +154$ (c, 0.95 in CHCl₃) (iodide).

O-De-Me, Et ether: Dubirheine. O-*Ethylrhoeagenine*

[2650-36-4]

C₂₂H₂₃NO₆ 397.427

Isol. from *Papaver armeniaceum*, *Papaver dubium* and *Papaver tauricola* (Papaveraceae). Needles (CHCl₃/EtOH). Mp 236-237°. $[\alpha]_D^{24} +236$ (c, 0.06 in CHCl₃). Prob. an artifact.

14-Epimer: Isorhoeadine. N-Methylpapaverrubine A. Alkaloid RA

[4046-21-3]

C₂₁H₂₁NO₆ 383.4

Alkaloid from *Mecconopsis betonicifolia*, *Papaver arenarium*, *Papaver argemone*, *Papaver commutatum* and other *Papaver* spp. (Papaveraceae). Mp 159-161°. $[\alpha]_D +314$ (c, 0.576 in CHCl₃).

14-Epimer, N-de-Me: Papaverrubine A. N-Demethylisorhoeadine. Alkaloid RS

[6807-93-8]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Mecconopsis betonicifolia* and many *Papaver* spp. (Papaveraceae). Mp 223-225°. $[\alpha]_D^{22} +406$ (c, 0.978 in CHCl₃).

8,14-Diepimer, O-de-Me: Isorhoegenine. Isorheagenine

[17948-35-5]

C₂₀H₁₉NO₆ 369.373

Alkaloid from *Papaver commutatum* and *Papaver rhoeas* (corn poppy) (Papaveraceae). Needles (MeOH/Me₂CO). Mp 210-215°. $[\alpha]_D^{22} +153$ (c, 0.65 in 1:1 CHCl₃/MeOH).

8,14-Diepimer, O-de-Me, O-α-D-glucopyranoside: Alkaloid RC

[17948-36-6]

C₂₆H₂₉NO₁₁ 531.515

Alkaloid from *Papaver commutatum* and *Papaver rhoeas* (corn poppy) (Papaveraceae). Mp 240-242°. $[\alpha]_D^{22} +250$ (c, 0.40 in CHCl₃/MeOH 1:1).

(-)-form [35321-09-6]

Synthetic. Mp 253-254°. $[\alpha]_D^{22} -228.2$ (c, 1 in CHCl₃).

(±)-form [30634-53-8]

Synthetic. Mp 222-224°.

N-De-Me: [70681-69-5]

Cryst. (CH₂Cl₂/petrol). Mp 216-218°.


Pfeifer, S. *et al.*, *Pharmazie*, 1964, **19**, 286;1972, **27**, 48 (*Papaverrubines*)Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*,1965, **30**, 2864; 1980, **45**, 2706; 1981, **46**,2587 (*Rhoeadine, Dubirheine*)Šantavý, F. *et al.*, *Coll. Czech. Chem. Comm.*,1965, **30**, 3479; 1967, **32**, 461 (*Rhoeadine, Isorhoegenine, uv, pmr, struct, synth*)

- Němečková, A. et al., *Naturwissenschaften*, 1967, **54**, 45 (Alkaloid RC)
- Dolejš, L. et al., *Tetrahedron*, 1967, **23**, 2997 (ms)
- Shamma, M. et al., *Chem. Comm.*, 1968, 212 (stereochem)
- Huber, C.S. et al., *Acta Cryst. B*, 1970, **26**, 373 (Rhoegenine, *cryst struct, abs config*)
- Klötzer, W. et al., *Helv. Chim. Acta*, 1972, **55**, 2228 (*synth*)
- Hohlbrugger, R. et al., *Chem. Ber.*, 1979, **112**, 849 (*synth*)
- Slavková, L. et al., *Coll. Czech. Chem. Comm.*, 1980, **45**, 761 (*isol*)
- Sariyar, G. et al., *Phytochemistry*, 1980, **19**, 2189 (*Dubirheine, Rhoeadine, isol, pmr, ms*)
- Věžník, F. et al., *Coll. Czech. Chem. Comm.*, 1981, **46**, 926 (*isol*)
- Phillipson, J.D. et al., *Planta Med.*, 1981, **41**, 105 (*Dubirheine, Rhoeadine*)
- Sariyar, G. et al., *Planta Med.*, 1982, **46**, 175; 1983, **49**, 43 (*isol*)
- Rönsch, H. et al., *Alkaloids (Academic Press)*, 1986, **28**, 78 (*pharmacol*)
- Marek, R. et al., *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr*)

Rhoifoline B

R-87

2,3,13-Trimethoxy[1,3]benzodioxolo[5,6-c]phenanthridine




C₂₁H₁₇NO₅ 363.369
Alkaloid from the stem bark of *Zanthoxylum rhoifolium*. Powder. Mp 258-259°.

Gonzaga, W. de A. et al., *Planta Med.*, 2003, **69**, 371-374 (*isol, pmr, cmr*)

Rhoifolinic acid

R-88

5-(2-Carboxy-1-ethenyl)-4-methoxyfuro[2,3-b]pyridine-6-carboxylic acid



(E)-form

C₁₂H₉NO₆ 263.206
Secofuroquinoline.

(E)-form

Di-Me ester: **Dimethyl (E)-rhoifolinate** [142543-00-8]

C₁₄H₁₃NO₆ 291.26
Alkaloid from *Glycosmis citrifolia*, *Metrodorea nigra* and *Zanthoxylum rhoifolium*.

9-Methoxy(E-), di-Me ester: **Fuomegistine I** [352320-25-3]

C₁₅H₁₅NO₇ 321.286
Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow solid. λ_{max} 298 (sh) ; 345 (sh) (log ε 2.64) (MeOH).

(Z)-form

Di-Me ester: **Dimethyl (Z)-rhoifolinate** [142542-99-2]


C₁₄H₁₃NO₆ 291.26
Alkaloid from *Glycosmis citrifolia*, *Metrodorea nigra* and *Zanthoxylum rhoifolium*.

Arruda, M.S.P. et al., *Biochem. Syst. Ecol.*, 1992, **20**, 173-178 (*isol*)
Mueller, A.H. et al., *Phytochemistry*, 1995, **40**, 1797-1800 (*isol*)
Fokialakis, N. et al., *Phytochemistry*, 2001, **57**, 593-596 (*Fuomegistine I*)

Rhopaladin A

R-89

[212069-48-2]



C₂₁H₁₃BrN₄O₃ 449.263
Alkaloid from the marine tunicate *Rhopalaea* sp. Amorph. red solid. λ_{max} 218 (ε 29000); 268 (ε 10000); 280 (ε 9700); 291 (ε 9600); 344 (ε 6700); 488 (ε 13000) (MeOH).

6-Deoxy: **Rhopaladin C**

[212069-50-6]
C₂₁H₁₃BrN₄O₂ 433.263
Alkaloid from *Rhopalaea* sp. Amorph. red solid. λ_{max} 212 (ε 56000); 258 (sh) ; 282 (ε 4900); 291 (sh) ; 471 (ε 5600) (MeOH).

Debromo: **Rhopaladin B**

[212069-49-3]
C₂₁H₁₄N₄O₃ 370.367
Alkaloid from *Rhopalaea* sp. Amorph. red solid. λ_{max} 213 (ε 19000); 267 (ε 6900); 281 (ε 5800); 293 (ε 5100); 350 (ε 4400); 485 (ε 8900) (MeOH).

Debromo, 6-deoxy: **Rhopaladin D**

[212069-51-7]
C₂₁H₁₄N₄O₂ 354.367
Alkaloid from *Rhopalaea* sp. Amorph. red solid. λ_{max} 211 (ε 22000); 259 (sh) ; 276 (ε 5400); 281 (sh) ; 468 (ε 4800) (MeOH).


Sato, H. et al., *Tetrahedron*, 1998, **54**, 8687-8690 (*Rhopaladins A-D, isol, uv, ir, pmr, cmr, ms*)

Janosik, T. et al., *Tetrahedron*, 2002, **58**, 2813-2819 (*synth*)

Rhyncophylline

R-90

Mitrinermine. Rhyncophylline [76-66-4]



Absolute Configuration

C₂₂H₂₈N₂O₄ 384.474
Alkaloid from *Uncaria rhyncophylla*, some other *Uncaria* spp., several *Mitragyna* spp. and *Cephalanthus occidentalis*

(Rubiaceae). Antipyretic, hypotensive agent. Mp 214° (208-209°). [α]_D -24 (CHCl₃) (-15). λ_{max} 245 (log ε 4.24); 280 (log ε 3.15) (no solvent reported).

N-Oxide: **Rhyncophylline N-oxide**.

Rhyncophylline N-oxide [35128-90-6]
C₂₂H₂₈N₂O₅ 400.474
Alkaloid from *Mitragyna inermis* and *Cephalanthus occidentalis* (Rubiaceae). Noncryst.

18,19-Didehydro: **Corynoxine**

[630-94-4]
C₂₂H₂₆N₂O₄ 382.458
Alkaloid from *Pseudocinchona africana* (preferred genus name *Corynanthe*), *Mitragyna rotundifolia*, *Mitragyna speciosa* and *Uncaria attenuata* (Rubiaceae). Cryst. (Me₂CO). Mp 212-214°. [α]_D -21 (CHCl₃). [α]_D +23 (c, 1 in Py). λ_{max} 244 (log ε 4.27) (EtOH).

Parent acid: **Rhyncophyllinic acid**. Rhyncophyllinic acid

[151124-30-0]
C₂₁H₂₆N₂O₄ 370.447
Alkaloid from stems of *Uncaria sinensis* (Rubiaceae). Amorph. powder. λ_{max} 237 (log ε 4.2); 280 (log ε 3.36) (EtOH).

7-Epimer: **Isorhyncophylline**. *Isorhyncophylline*

[6859-01-4]
C₂₂H₂₈N₂O₄ 384.474
Alkaloid from *Uncaria* and *Mitragyna* spp. (Rubiaceae). Mp 144°. [α]_D +13 (+8.3) (CHCl₃). λ_{max} 243 (log ε 4.27) (EtOH).

▶ GN1605000

7-Epimer, β-N-oxide: **Isorhyncophylline N-oxide**. *Isorhyncophylline N-oxide*

[35129-00-1]
C₂₂H₂₈N₂O₅ 400.474
Alkaloid from *Mitragyna inermis*, *Mitragyna rotundifolia* and *Cephalanthus occidentalis* (Rubiaceae). Mp 242-243°. N-oxide config. *anti*.

7-Epimer, 18,19-didehydro: **Isocorynoxine**

[51014-29-0]
C₂₂H₂₆N₂O₄ 382.458
Alkaloid from *Uncaria attenuata attenuata* and *Mitragyna rotundifolia* (Rubiaceae). Noncryst.

7-Epimer, parent acid: **Isorhyncophyllinic acid**. *Isorhyncophyllinic acid*

[144525-05-3]
C₂₁H₂₆N₂O₄ 370.447
Alkaloid from stems of *Uncaria sinensis* (Rubiaceae). Mp 175-176.5°. [α]_D +5.2 (c, 0.08 in MeOH). λ_{max} 242 (log ε 4.27); 280 (log ε 3.5) (EtOH).

7-Epimer, demethoxy: **17-Demethoxyisorhyncophylline**

[94482-54-9]
C₂₁H₂₆N₂O₃ 354.448
Alkaloid from the whole plant of *Amsonia brevifolia* (Apocynaceae). Cytotoxic agent. Amorph. gum. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D 0 (c, 0.2 in MeOH). λ_{max} 210 ; 252 ; 282 (MeOH) (Berdy).

7-*Epimer, demethoxy, 16,17-dihydro:*
16,17-Dihydro-17-demethoxyisorhynchophylline
 [259540-05-1]
 $C_{21}H_{28}N_2O_3$ 356.464
 Alkaloid from *Ervatamia yunnanensis*.
 Gum.

7-*Epimer, demethoxy, 16,17-dihydro, N-oxide:* **16,17-Dihydro-17-demethoxyisorhynchophylline N-oxide**
 [259540-06-2]
 $C_{21}H_{28}N_2O_4$ 372.463
 Alkaloid from *Ervatamia yunnanensis*.
 Gum.

20-*Epimer: Corynoxine B. Isocorynoxine*
 [17391-18-3]
 $C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from *Uncaria macrophylla*
 (Rubiaceae). Noncryst.

20-*Epimer, demethoxy: 17-Demethoxycorynoxine B*
 [119365-65-0]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the whole plant of
Amsonia brevifolia (Apocynaceae).
 Cytotoxic agent. Fine needles. Sol.
 MeOH, $CHCl_3$; poorly sol. H_2O .
 Mp 178-180°. $[\alpha]_D^{25} +61$ (c, 0.2 in
 MeOH). λ_{max} 214 ; 253 ; 286 (MeOH)
 (Berdy).

3,15-*Diepimer, demethoxy: Catharinensine*
 [94444-28-7]
 $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the bark of *Peschiera*
catharinensis (Apocynaceae). $[\alpha]_D^{25} -194$
 (c, 0.5 in $CHCl_3$). Unusual 15-epi
 config. CD of opposite sign to that of
 Corynoxine.

7,20-*Diepimer: Corynoxine. Corynoxine A*
 [6877-32-3]
 $C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from *Pseudocinchona africa-*
na, *Uncaria macrophylla* and *Mitragyna*
speciosa (Rubiaceae). Mp 166-168°.
 $[\alpha]_D -3$ ($CHCl_3$). $[\alpha]_D -14$ (c, 1 in Py).

(±)-*form* [39032-62-7]
 Synthetic. Mp 197-199°.

7-*Epimer:* Mp 225-227°.


Cu, N.A. et al., *Bull. Soc. Chim. Fr.*, 1957,
 1292-1294 (*Corynoxine, Corynoxine*)
 Seaton, J.C. et al., *Can. J. Chem.*, 1957, **35**,
 1102-1108; 1960, **38**, 1035-1042
 (*Rhynchophylline, Isorhynchophylline*)
 Finch, N. et al., *J.A.C.S.*, 1962, **84**, 3871-3877
 (*Corynoxine, config*)
 Gilbert, B. et al., *J.A.C.S.*, 1963, **85**, 1523-1528
 (*ms*)
 Pousset, J.L. et al., *Tet. Lett.*, 1966, **7**, 6283-
 6289; 1967, **8**, 952; 1919-1927 (*Corynoxine,*
cd, config, synth)
 Pousset, J.-L. et al., *Bull. Soc. Chim. Fr.*, 1967,
 2766-2779 (*pmr, config*)
 Becket, A.H. et al., *Tetrahedron*, 1969, **25**,
 5961-5969 (*ms*)
 Shellard, E.J. et al., *Phytochemistry*, 1971, **10**,
 2505-2511 (*oxides*)
 Phillipson, J.D. et al., *Phytochemistry*, 1973,
12, 2795-2798; 1974, **13**, 2621-2622
 (*Corynoxine B, N-oxides*)
 Hough, P.J. et al., *Planta Med.*, 1974, **26**, 104
 (*Isocorynoxine*)
 Ban, Y. et al., *Chem. Pharm. Bull.*, 1975, **23**,
 2605-2613 (*Isorhynchophylline, synth, ir, ms*)

Phillipson, J.D. et al., *J. Chromatogr.*, 1975,
105, 163-178 (*tlc, gc, ms*)
 Brown, R.T. et al., *Tet. Lett.*, 1976, 1401-1402
 (*Isorhynchophylline, synth*)
 Araujo, A.R. et al., *Phytochemistry*, 1984, **23**,
 2359 (*Catharinensine*)
 Sharma, P. et al., *Phytochemistry*, 1988, **27**,
 3649-3652 (*17-Demethoxycorynoxine B, 17-*
Demethoxyisorhynchophylline)
 Liu, H.-M. et al., *Phytochemistry*, 1993, **33**,
 707-710 (*Rhynchophyllic acid,*
Isorhynchophyllic acid)
 Yu, Y. et al., *Chin. Chem. Lett.*, 1999, **10**, 575-
 578 (*Dihydrodemethoxyisorhynchophylline*)
 Laus, G. et al., *Helv. Chim. Acta*, 2003, **86**,
 181-187 (*cryst struct*)
 Deiters, A. et al., *J.O.C.*, 2006, **71**, 6547-6561
 (*synth*)

Ribaline

R-91

3,9-*Dihydro-6-hydroxy-2-(1-hydroxy-1-*
methylethyl)-9-methylfuro[2,3-b]quinolin-
4(2H)-one, 9CI. Rutalinidine
 [50894-68-3]



(R)-form

 $C_{15}H_{17}NO_4$ 275.304

(R)-*form* [6872-52-2]

Alkaloid from the trunk bark and
 heartwood of *Balfourodendron riedelianum*
 (Rutaceae). Cryst. (MeOH). Mp
 210-211° (indefinite). $[\alpha]_D^{25} +86$ (c, 0.44 in
 MeOH) (+72.1).

O⁶-*Ac:*

Glistening plates (Me₂CO). Mp 222-
 223°. $[\alpha]_D^{25} +82.1$ (c, 0.6 in MeOH).

(±)-*form* [41234-32-6]

Minor alkaloid from *Balfourodendron*
riedelianum and *Ruta graveolens* (rue)
 (Rutaceae). Needles (MeOH). Mp 268-
 269° (259-260° dec.).

Picrate: Mp 223-224° dec.

O⁶-*Ac:*


Needles (MeOH). Mp 240°.

Szendrei, K. et al., *Herba Hung.*, 1971, **10**, 131-
 139 (*isol*)
 Corral, R.A. et al., *Tetrahedron*, 1973, **29**, 205
 (*isol, uv, ir, struct, synth*)
 Hammerum, S. et al., *Acta Chem. Scand., Ser.*
B, 1977, **31**, 31 (*ms*)
 Jurd, L. et al., *Aust. J. Chem.*, 1983, **36**, 1615
 (*isol, ir, pmr, cmr, ms*)
 Corral, R.A. et al., *Tet. Lett.*, 1983, **24**, 2359
 (*synth*)

Ribalinine

R-92

2,3,4,10-*Tetrahydro-3-hydroxy-2,2,10-tri-*
methyl-5H-pyrano[2,3-b]quinolin-5-one,
9CI. Folifine
 [16117-68-3]



(R)-form

 $C_{15}H_{17}NO_3$ 259.304

(R)-*form* [62928-56-7]

Alkaloid from *Haplophyllum bucharicum*,
Haplophyllum foliosum and *Araliopsis*
tabouensis (Rutaceae). Mp 222°. $[\alpha]_D$
 +14.1 (c, 0.6 in MeOH).

7-*Hydroxy: Ribalinidine*

[87936-14-9]

$C_{15}H_{17}NO_4$ 275.304

Minor alkaloid from the heartwood of
Balfourodendron riedelianum (Ruta-
 ceae). Needles (MeOH). Mp 260-261°.
 $[\alpha]_D^{25} +15.3$ (c, 0.5 in MeOH).

9-*Methoxy: Isobalfourodine*

[478-38-6]

$C_{16}H_{19}NO_4$ 289.33

Alkaloid from *Balfourodendron riedelianum*
 (Rutaceae). Mp 204-205°. $[\alpha]_D$
 +15 (c, 1 in EtOH).

9-(3-*Methyl-2-butenyloxy*): **Ribaliprenylene**

$C_{20}H_{25}NO_4$ 343.422

Alkaloid from *Skimmia laureola*. Pale
 yellow gum. $[\alpha]_D^{25} +10$ (c, 1 in $CHCl_3$).
 λ_{max} 265 (log ε 2.67); 304 (log ε 3.84);
 318 (log ε 3.86); 325 (log ε 3.74); 330
 (log ε 3.76) (MeOH).

(S)-*form* [7688-58-6]

Alkaloid from *Araliopsis soyauxii*. Mp
 222°. $[\alpha]_D -14$ (c, 0.5 in MeOH).

7-*Hydroxy:* [19783-80-3]

Alkaloid from *Balfourodendron*
riedelianum. Cryst. (EtOH). Mp
 257-258° dec. $[\alpha]_D^{20} -15$ (c, 1.0 in
 MeOH).

9-*Methoxy: Lunasia II. Lunacrinol*

$C_{16}H_{19}NO_4$ 289.33

Alkaloid from *Lunasia amara* (Ruta-
 ceae). Mp 201-203°. $[\alpha]_D -14$ (EtOH).

(±)-*form*

Alkaloid from the bark of *Balfourodendron*
riedelianum (Rutaceae). Mp 233-
 234°.

7-*Hydroxy:* [86851-77-6]

Synthetic. Cryst. (EtOH). Mp 256-
 257°.

9-*Methoxy:* Mp 184-186°.

Rapoport, H. et al., *J.A.C.S.*, 1960, **82**, 4395
 (*Isobalfourodine*)

Bowman, R.M. et al., *J.C.S.(C)*, 1966, 1504
 (*synth, pmr*)

Faizutdinova, Z.Ch. et al., *Khim. Prir. Soedin.*,
 1967, **3**, 257; *Chem. Nat. Compd. (Engl.*
Transl.), 1967, **3**, 215 (*Ribalinine*)

Collins, J.F. et al., *Chem. Comm.*, 1969, 1078
 (*config*)

Corral, R.A. et al., *Tetrahedron*, 1973, **29**, 205
 (*Ribalinine, Ribalinidine, isol, ir, uv, pmr,*
struct, synth)

Vaquette, J. et al., *Phytochemistry*, 1976, **15**,
 743 (*Ribalinine, isol, ir, uv, ms, pmr, struct*)
 Hammerum, S. et al., *Acta Chem. Scand., Ser.*
B, 1977, **31**, 31 (*ms*)

Grundon, M.F. et al., *Chem. Comm.*, 1978,
 624 (*config*)

Jurd, L. et al., *Aust. J. Chem.*, 1983, **36**, 1615
 (*Ribalinidine, isol, ir, pmr, cmr, ms, struct*)

Corral, R.A. et al., *Tet. Lett.*, 1983, **24**, 2359
 (*synth*)


Ramesh, M. et al., *Heterocycles*, 1984, **22**, 125
 (*synth, ir, pmr*)

Ramesh, M. et al., *Indian J. Chem., Sect. B*,
 1985, **24**, 602 (*Isobalfourodine, synth*)

- Subramanian, M. *et al.*, *Z. Naturforsch., B*, 1992, **47**, 1016 (*synth*)
 Barr, S.A. *et al.*, *Chem. Comm.*, 1994, 153 (*synth*)
 Boyd, D.R. *et al.*, *J.C.S. Perkin 1*, 2000, 3397-3405 (*synth, abs config*)
 Sultana, N. *et al.*, *Z. Naturforsch., B*, 2006, **60**, 1186-1191 (*Ribaliprenylene*)

Ribalinium**R-93**

2,3-Dihydro-6-hydroxy-2-(1-hydroxy-1-methylethyl)-4-methoxy-9-methylfuro[2,3-b]quinolinium(1+), 9CI [6883-22-3]



$C_{16}H_{20}NO_4^{\oplus}$ 290.338

Alkaloid from the trunk bark of *Balfourodendron riedelianum* and the leaves of *Ruta graveolens* (rue) (Rutaceae).

Chloride:

Cryst. (MeOH/EtOAc or MeOH). Mp 188-190° (gas evolv.). $[\alpha]_D^{24} +40$ (c, 1.25 in MeOH).

Perchlorate: Mp 205° (218-220° dec.). $[\alpha]_D^{20} +23.6$ (c, 0.006 in MeOH).

Picrate:


Cryst. (Me₂CO/EtOH). Mp 192-193° (darkening, gas evolv.).

Corral, R.A. *et al.*, *Tetrahedron*, 1965, **21**, 909 (*isol, uv, ir, pmr, struct*)

Rideau, M. *et al.*, *Phytochemistry*, 1979, **18**, 155 (*isol, pmr, ms, struct*)

Ribasine**R-94**

Limogine. Grandiflorine† [87099-54-5]



$C_{20}H_{17}NO_5$ 351.358

Alkaloid from *Sarcocapnos crassifolia*, *Corydalis claviculata* and *Argemone grandiflora* subsp. *grandiflora* (Fumariaceae, Papaveraceae). Cryst. (EtOH). Mp 202.5-203.5° (194-195°). $[\alpha]_D +125$ (c, 1 in CHCl₃).

N-De-Me: Norribasine

[133084-01-2]

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. $[\alpha]_D^{25} +72$ (c, 0.36 in MeOH).

13β-Hydroxy: **Ribasidine**. 13β-Hydroxyribasine

[88151-44-4]

$C_{20}H_{17}NO_6$ 367.357

Alkaloid from *Sarcocapnos emneaphyl-*

la and *Corydalis claviculata* (Papaveraceae). Needles (CHCl₃). Mp 230°. $[\alpha]_D^{25} +120$ (c, 0.04 in EtOH).

13β-Hydroxy, O-Ac:

Cryst. (MeOH). Mp 198°.

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2029 (*uv, pmr, ms, cryst struct*)

Allais, D.P. *et al.*, *Tet. Lett.*, 1983, **24**, 2445 (*uv, ir, pmr, cmr, ms, cd, struct*)

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 4481 (*uv, ir, pmr, cmr, ms, struct, Ribasidine*)

Mitchell, R.E. *et al.*, *Can. J. Chem.*, 1984, **62**, 258 (*isol, uv, ir, pmr, ms, struct*)

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1280 (*Norribasine*)


Ollero, L. *et al.*, *Tet. Lett.*, 1998, **39**, 1413-1416 (*synth*)

Padwa, A. *et al.*, *J.O.C.*, 1999, **64**, 4079-4088 (*synth*)

Ollero, L. *et al.*, *Tetrahedron*, 1999, **55**, 4445-4456 (*synth, pmr, cmr*)

Ricinidine**R-95**

1,2-Dihydro-1-methyl-2-oxo-3-pyridine-carbonitrile, 9CI. 3-Cyano-1-methyl-2-pyridone [767-88-4]



$C_7H_6N_2O$ 134.137

Alkaloid from *Trewia nudiflora* (Euphorbiaceae), long known previously as synthetic compd. Needles (CHCl₃/Et₂O). Mp 146°. Bp₂₈ 243°.


Späth, E. *et al.*, *Ber.*, 1923, **56**, 880 (*bibl*)

Robinson, W.G. *et al.*, *J. Biol. Chem.*, 1964, **239**, 4257 (*synth*)

Ganguly, S.N. *et al.*, *Phytochemistry*, 1970, **9**, 1667 (*isol, uv*)

Ricinine**R-96**

1,2-Dihydro-4-methoxy-1-methyl-2-oxo-3-pyridinecarbonitrile, 9CI. 4-Methoxy-2-oxo-1-methyl-1,2-dihydro nicotinonitrile. 3-Cyano-4-methoxy-1-methyl-2-pyridone. *Ricinine* [524-40-3]



$C_8H_8N_2O_2$ 164.163

Alkaloid from *Ricinus communis* (castor-oil) seed (Euphorbiaceae). Leaflets or prisms (H₂O). Sol. hot H₂O. Mp 201°.

► Human systemic effects by ingestion, hepatotoxic, can cause death. QT3100000

HgCl₂ compd.: Mp 204°.

Tuson, R.V. *et al.*, *J.C.S.*, 1864, **17**, 195 (*isol*)

Späth, E. *et al.*, *Ber.*, 1925, **58**, 2124 (*synth, bibl*)

Robinson, W.E. *et al.*, *J. Biol. Chem.*, 1964, **239**, 4257 (*ir, uv, synth*)

Walker, G.R. *et al.*, *Anal. Biochem.*, 1966, **16**, 277 (*ms*)

Sugasawa, T. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 763 (*synth, ir, pmr*)


Robinson, T. *et al.*, *Phytochemistry*, 1978, **17**, 1903 (*biosynth*)

Soriano-Garcia, M. *et al.*, *Acta Cryst. C*, 1989, **45**, 957 (*cryst struct*)

Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979,

Riedelianine**R-97**

2-(1-Hydroxy-1-methylethyl)-2,3-dihydrofuro[2,3-b]quinolin-6-ol. 2,3-Dihydro-6-hydroxy- α, α -dimethylfuro[2,3-b]quinoline-2-methanol [87893-11-6]




$C_{14}H_{15}NO_3$ 245.277

Alkaloid from *Balfourodendron riedelianum* heartwood (Rutaceae). Thick needles (MeOH). Mp 257°. $[\alpha]_D^{25} +25.3$ (c, 0.6 in MeOH).

Jurd, L. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1615 (*isol, cryst struct, ms, ir, cmr, pmr*)

Rigidin A**R-98**

6-(4-Hydroxybenzoyl)-5-(4-hydroxyphenyl)-1H-pyrrolo[2,3-d]pyrimidine-2,4-(3H,7H)-dione, 9CI. *Rigidin* [132160-44-2]



$C_{19}H_{13}N_5O_5$ 363.329

Alkaloid from the marine tunicate *Eudistoma cf. rigida* and *Cystodytes* sp. Calmodulin antagonist. Purple solid. Mp 300°. λ_{max} 232 (sh); 276 (€ 4500); 356 (€ 2500); 552 (€ 200) (MeOH/HCl) (Derep). λ_{max} 240 (sh); 285 (€ 4300); 346 (€ 4500); 401 (€ 2500) (MeOH/KOH) (Derep). λ_{max} 240 (sh); 285 (€ 4300); 346 (€ 2500); 401 (€ 2500) (MeOH) (Derep). λ_{max} 206 (€ 39780); 278 (€ 21070); 285 (€ 4300); 308 (€ 14860); 346 (€ 2500); 356 (€ 14110); 401 (€ 2500) (MeOH) (Berdy).

3-N-Me: **Rigidin E**

$C_{20}H_{15}N_3O_5$ 377.356

Alkaloid from a *Eudistoma* sp. Dark green film. λ_{max} 276 (€ 7000); 362 (€ 4000) (MeOH).

3'-Methoxy: **Rigidin B**

$C_{20}H_{15}N_3O_6$ 393.355

Alkaloid from the marine tunicate *Cystodytes* sp. Amorph. yellow

solid. λ_{\max} 236 (sh) ; 278 (ϵ 13200); 304 (sh) ; 319 (ϵ 9600); 364 (ϵ 9400) (MeOH).

3''-Methoxy: Rigidin C

$C_{20}H_{15}N_3O_6$ 393.355
Alkaloid from *Cystodytes* sp.
Amorph. yellow solid. λ_{\max} 236 (sh) ; 272 (ϵ 14400); 304 (sh) ; 317 (ϵ 11600); 352 (ϵ 9400) (MeOH).

3',3''-Dimethoxy: Rigidin D

$C_{21}H_{17}N_3O_7$ 423.381
Alkaloid from *Cystodytes* sp.
Amorph. yellow solid. λ_{\max} 238 (sh) ; 270 (ϵ 12000); 304 (sh) ; 322 (ϵ 11000); 352 (ϵ 9300) (MeOH).

Kobayashi, J. *et al.*, *Tet. Lett.*, 1990, **31**, 4617-4620 (*isol, uv, ir, pmr, cmr, struct*)

Edstrom, E.D. *et al.*, *J.O.C.*, 1993, **58**, 403-407 (*synth*)

Sakamoto, T. *et al.*, *J.C.S. Perkin 1*, 1996, 459-464 (*synth*)


Tsuda, M. *et al.*, *J. Nat. Prod.*, 2003, **66**, 292-294 (*Rigidins B,C,D*)

Davis, R.A. *et al.*, *Mar. Drugs*, 2003, **1**, 27-33 (*Rigidin E*)

Gupton, J.T. *et al.*, *Tetrahedron*, 2006, **62**, 8243-8255 (*synth*)

Ripromycin**R-99**

[617711-42-9 ((+)-form)]




$C_{30}H_{40}N_2O_6$ 524.656

Prod. by *Streptomyces* sp. Tü 6239.
Exhibits antibacterial and cytostatic activities. Yellow-beige powder. Mp 190°. $[\alpha]_D^{20}$ +114 (c, 0.77 in MeOH). λ_{\max} 204 (log ϵ 4.4); 238 (sh) ; 290 (sh) ; 314 (log ϵ 4.05) (MeOH).

Bertasso, M. *et al.*, *J. Antibiot.*, 2003, **56**, 364-371 (*isol, activity*)

Ritterazine V**R-100**

[184770-01-2]




$C_{54}H_{76}N_2O_9$ 897.202
Alkaloid from the tunicate *Ritterella tokioka*. $[\alpha]_D$ +109.2 (c, 0.05 in $CHCl_3$). λ_{\max} 290 (ϵ 9209); 308 (sh) ($CHCl_3$).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

Ritterazine A**R-101**

[160391-62-8]



$C_{54}H_{76}N_2O_{10}$ 913.202

Alkaloid from the tunicate *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +112 (c, 0.1 in MeOH). λ_{\max} 287 (ϵ 8580); 308 (sh) (ϵ) (MeOH) (Derep).

22-Epimer: Ritterazine D

[165172-44-1]

$C_{54}H_{76}N_2O_{10}$ 913.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +81.4 (c, 0.1 in MeOH). λ_{\max} 286 (ϵ 9200) (MeOH) (Berdy).

7',17'-Dideoxy: Ritterazine T

[184768-89-6]

$C_{54}H_{76}N_2O_8$ 881.203


Alkaloid from *Ritterella tokioka*. $[\alpha]_D$ +106.6 (c, 0.1 in $CHCl_3$). λ_{\max} 290 (ϵ 9870); 308 (sh) ($CHCl_3$).

Fukuzawa, S. *et al.*, *J.O.C.*, 1994, **59**, 6164-6166; 1997, **62**, 4484-4491 (*isol, uv, ir, pmr, cmr, struct*)

Fukuzawa, S. *et al.*, *Tetrahedron*, 1995, **51**, 6707-6716 (*Ritterazine D*)

Ritterazine B**R-102**

[160568-10-5]



$C_{54}H_{78}N_2O_9$ 899.218

22-Config. revised in 2007. C-27 is subject to epimerisation, and the stereochem. of some of the Ritterazines is subject to revision. Alkaloid from the tunicate *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +43 (c, 0.1 in MeOH). λ_{\max} 287 (ϵ 6880); 308 (sh) (ϵ) (MeOH) (Derep).

7',17'-Dideoxy: Ritterazine Y

[184770-06-7]

$C_{54}H_{78}N_2O_7$ 867.219

Alkaloid from *Ritterella tokioka*. Cytotoxic agent. $[\alpha]_D$ +57.4 (c, 0.1 in $CHCl_3$). λ_{\max} 289 (ϵ 9156); 308 (sh) ($CHCl_3$).

22-Epimer: Ritterazine F

[165172-45-2]

$C_{54}H_{78}N_2O_9$ 899.218

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leu-

kaemia cells. Glassy solid. $[\alpha]_D$ +59 (c, 0.1 in MeOH). λ_{\max} 288 (ϵ 7900) (MeOH) (Berdy).

22-Epimer, 14,15-didehydro: Ritterazine G

[164991-70-2]

$C_{54}H_{76}N_2O_9$ 897.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +91.4 (c, 0.1 in MeOH). λ_{\max} 288 (ϵ 11200) (MeOH) (Berdy).

22-Epimer, 12-ketone: Ritterazine H

[165074-71-5]

$C_{54}H_{76}N_2O_9$ 897.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +96 (c, 0.1 in MeOH). λ_{\max} 287 (ϵ 8920) (MeOH) (Berdy).

22-Epimer, 14-hydroxy, 12-ketone: Ritterazine I

[164991-71-3]

$C_{54}H_{76}N_2O_{10}$ 913.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +74.5 (c, 0.1 in MeOH). λ_{\max} 286 (ϵ 9120) (MeOH) (Berdy).


Fukuzawa, S. *et al.*, *J.O.C.*, 1995, **60**, 608-614; 1997, **62**, 4484-4491 (*isol, uv, ir, pmr, cmr, struct*)

Fukuzawa, S. *et al.*, *Tetrahedron*, 1995, **51**, 6707-6716 (*Ritterazines F-I*)

Phillips, S.T. *et al.*, *J.A.C.S.*, 2007, **129**, 6589-6598 (*stereochem, bibl*)

Ritterazine C**R-103**

[160604-68-2]



$C_{54}H_{78}N_2O_9$ 899.218

Alkaloid from the tunicate *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +72 (c, 0.1 in MeOH). λ_{\max} 287 (ϵ 6880); 308 (sh) (MeOH) (Derep). λ_{\max} 285 (ϵ 8720); 303 (MeOH) (Berdy).

Fukuzawa, S. *et al.*, *J.O.C.*, 1995, **60**, 608-614 (*isol, uv, ir, pmr, cmr, struct*)

Ritterazine E**R-104****24-Methylritterazine D**

[164991-69-9]

As Ritterazine A, R-101 with R = CH_3 , 22-epimer


$C_{55}H_{78}N_2O_{10}$ 927.229

Alkaloid from the tunicate *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. $[\alpha]_D$ +70.8 (c, 0.1 in MeOH). λ_{\max} 288 (ϵ 10100) (MeOH) (Berdy).

Fukuzawa, S. *et al.*, *Tetrahedron*, 1995, **51**, 6707-6716 (*isol, uv, ir, pmr, cmr, struct*)

Ritterazine J

[164991-72-4]

C₅₄H₇₆N₂O₁₁ 929.201

Stereochem. revised in 2001. Alkaloid from the tunicate *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. [α]_D +66.1 (c, 0.1 in MeOH). λ_{max} 289 (ε 8420) (MeOH) (Berdy).

7'-Deoxy: Ritterazine K

[164991-73-5]

C₅₄H₇₆N₂O₁₀ 913.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. [α]_D +74 (c, 0.1 in MeOH). λ_{max} 288 (ε 7100) (MeOH) (Berdy).

7,17-Dideoxy: Ritterazine L

[164991-74-6]

C₅₄H₇₆N₂O₉ 897.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. [α]_D +85.5 (c, 0.1 in MeOH). λ_{max} 288 (ε 11000) (MeOH) (Berdy).

7,17-Dideoxy, 25-epimer: Ritterazine M

[165172-46-3]

C₅₄H₇₆N₂O₉ 897.202

From *Ritterella tokioka*. Shows potent cytotoxicity against P388 murine leukaemia cells. Glassy solid. [α]_D +95.1 (c, 0.1 in MeOH). Struct. revised in 2002. λ_{max} 289 (ε 11900) (MeOH) (Berdy).


Fukuzawa, S. *et al.*, *Tetrahedron*, 1995, **51**, 6707-6716 (*isol, uv, ir, pmr, cmr, struct*)

Jeong, J.U. *et al.*, *J.A.C.S.*, 1999, **121**, 2071-2084 (*synth, Ritterazine K*)

Lee, S. *et al.*, *Org. Lett.*, 2002, **4**, 313-316; 317-318 (*Ritterazine M, synth, struct*)

Ritterazine N

[184764-19-0]

C₅₄H₇₆N₂O₈ 881.203

Alkaloid from the tunicate *Ritterella tokioka*. Cytotoxic agent. [α]_D +121.7 (c, 0.05 in CHCl₃). λ_{max} 289 (ε 9388); 309 (sh) (CHCl₃).

22'-Epimer: Ritterazine O

[184971-68-4]

C₅₄H₇₆N₂O₈ 881.203


From *Ritterella tokioka*. [α]_D +108.6 (c, 0.1 in CHCl₃). λ_{max} 289 (ε 9379); 308 (sh) (CHCl₃).

R-105

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

Ritterazine P

[184765-85-3]

C₅₄H₇₈N₂O₇ 867.219

Alkaloid from the tunicate *Ritterella tokioka*. Cytotoxic agent. [α]_D +42.5 (c, 0.05 in CHCl₃). λ_{max} 289 (ε 10618); 309 (sh) (CHCl₃).

22'-Epimer: Ritterazine Q

[184971-69-5]


C₅₄H₇₈N₂O₇ 867.219

From *Ritterella tokioka*. Cytotoxic agent. [α]_D +57.8 (c, 0.05 in CHCl₃). λ_{max} 289 (11225); 309 (sh) (CHCl₃).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

Ritterazine R

[184766-99-2]

C₅₄H₈₀N₂O₆ 853.236

Alkaloid from the tunicate *Ritterella tokioka*. [α]_D +26.3 (c, 0.05 in CHCl₃). λ_{max} 288 (9557); 309 (sh) (CHCl₃).

22'-Epimer: Ritterazine S


[184971-73-1]

From *Ritterella tokioka*. Cytotoxic agent. [α]_D +43.3 (c, 0.05 in CHCl₃). λ_{max} 290 (ε 10480); 308 (sh) (CHCl₃).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

Ritterazine U

[184769-28-6]


C₅₄H₇₆N₂O₉ 897.202

Alkaloid from the tunicate *Ritterella tokioka*. [α]_D +89 (c, 0.1 in CHCl₃). λ_{max} 290 (9424); 308 (sh) (CHCl₃).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

R-107**Ritterazine W**

[184770-05-6]

C₅₄H₇₆N₂O₈ 881.203

Alkaloid from the tunicate *Ritterella tokioka*. [α]_D +120.4 (c, 0.05 in CHCl₃). λ_{max} 290 (ε 8155); 308 (sh) (CHCl₃).

22-Epimer: Ritterazine X

[184971-98-0]


C₅₄H₇₆N₂O₈ 881.203

From *Ritterella tokioka*. [α]_D +108 (c, 0.05 in CHCl₃). λ_{max} 290 (ε 8816); 308 (sh) (CHCl₃).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

Ritterazine Z

[184770-08-9]

C₅₅H₇₈N₂O₉ 911.229

Alkaloid from the tunicate *Ritterella tokioka*. [α]_D +105.8 (c, 0.1 in CHCl₃). λ_{max} 289 (8958); 308 (sh) (CHCl₃).

Fukuzawa, S. *et al.*, *J.O.C.*, 1997, **62**, 4484-4491 (*isol, uv, pmr, cmr*)

R-110**R-111****Robecine**

[1360-89-0]

C₁₇H₂₁NO₃ 287.358

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Narcissus* sp. (Amaryllidaceae).


Hydroiodide: Mp 240-241° dec. [α]_D²⁵ -95 (c, 0.2 in DMF).

Methiodide: Mp 248-249° dec.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 2197-2202 (*isol*)

R-112**Robustamine**

[174847-42-8]



Absolute Configuration

C₂₀H₂₇NO₄ 345.438

Alkaloid from aerial parts of *Merendera robusta*. Cryst. (Me₂CO). Mp 251-252°. [α]_D -40 (c, 0.50 in CHCl₃).

R-113


α -N-Oxide: **Robustamine cis-N-oxide**
[191356-25-9]
C₂₀H₂₇NO₅ 361.437
Alkaloid from *Merendera robusta*.
Cryst. (Me₂CO). Mp 140-142°. [α]_D²² -24 (c, 0.4 in CHCl₃).

Yunusov, M.K. *et al.*, *Khim. Prir. Soedin.*, 1995, **31**, 109; 1996, **32**, 734-738; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 87; 1996, **32**, 716-719

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 761-863; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 737-858 (N-oxide)

Rodiasine, 9CI**R-114**

6'-O-Methylphlebicine
[6391-64-6]



C₃₈H₄₂N₂O₆ 622.76

Alkaloids covered by this entry (1*S*,1'*R*-) are enantiomeric with those in Cordobine, C-643 and diastereomeric with those in Guattaguianine, G-212 (S,S-). Alkaloid from the bark and seeds of *Ocotea rodiaei* (Lauraceae). Needles (MeOH). Mp 203-204°. [α]_D²⁴ +157 (c, 1.0 in CHCl₃). [α]_D +168 (c, 0.3 in CHCl₃). Hydrochloride (1:2): Mp 247-249°.

N²-De-Me: **Ocotine**. 2'-Demethylrodiasine
[18529-55-0]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the bark and seeds of *Ocotea rodiaei* (Lauraceae). Needles (MeOH). Mp 165°. [α]_D²⁴ +40 (c, 1.0 in CHCl₃).

N²-De-Me, hydrochloride (1:2):
Needles + 2H₂O (dil. HCl). Mp 250-260° dec. [α]_D²⁹ +84 (c, 2.6 in MeOH).

N-De-Me: **Norrodiasine**

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the bark of *Ocotea rodiaei* (Lauraceae). Cryst. + 2H₂O (as dihydrochloride). Mp 282° (dihydrochloride). [α]_D +74 (c, 1.0 in H₂O). May be identical with Ocotine or may be the N²-de-Me isomer.

Me ether: Obt. by methylation of Rodiasine, Tiliageine or Antioquine. Cryst. (Et₂O/hexane or EtOH) or amorph. solid. Mp 170-172° (163-165°, 168-169°). [α]_D²⁴ +86.9 (c, 0.45 in MeOH). [α]_D +78 (c, 0.9 in CHCl₃).

Me ether, N²,N²-di-Me:
Prisms (H₂O). Mp 270-280° dec.

O⁶-De-Me: **Phlebicine**

[52674-06-3]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the bark of *Crematosperma polyphlebium* (Annonaceae). Cryst. + 1.5H₂O (CHCl₃/MeOH). Mp 195° (sinters at 180°). [α]_D +182.5 (c, 1.0 in CHCl₃).

O⁶-De-Me, di-Ac:

Yellow powder. Mp 200-203° (sinters at 190°).

O⁷-De-Me: **(+)-Antioquine**

[93767-27-2]

Alkaloid from *Pseudoxandra* aff. *lucida* (Annonaceae). Needles (MeOH). Mp 197°. [α]_D +214 (c, 0.9 in CHCl₃).

O⁷-De-Me, di-Ac:

Needles (MeOH). Mp 164-166°. [α]_D +150 (c, 0.2 in CHCl₃).

O⁷-De-Me, N²-de-Me: **Tilitriandrine**

[120139-68-6]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from aerial parts of *Tiliacora triandra* (Menispermaceae). Cryst. + 1CH₂Cl₂ (MeOH/CH₂Cl₂). Mp 192°. [α]_D +197.9 (c, 1.1 in CHCl₃).

O¹²-De-Me, O¹²-Me: **Funiferine**. O¹²-Demethyl-O¹²-methylrodiasine

[1394-44-1]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from roots of *Tiliacora funifera* and *Tiliacora dinklagei* (Menispermaceae). Exhibits antiparasitic and cytotoxic activities. Active against gram-positive bacteria. Cryst. (EtOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 232-234° (218-220°). [α]_D³² +184.3 (c, 2.0 in CHCl₃).

O¹²-De-Me, O¹²-Me, N²-oxide: **Funiferine N-oxide**

[61912-73-0]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from the roots of *Tiliacora funifera* (Menispermaceae). Powder. Mp 207-209° dec. [α]_D²⁵ +44 (c, 0.1 in MeOH).

O¹²-De-Me, O¹²-Me, N,N-di-Me: **Funiferine N,N'-dimethosalt**

[1394-46-3]

C₄₀H₄₈N₂O₆[⊕] 652.829

Alkaloid from the roots of *Tiliacora funifera* (Menispermaceae). Cryst. (Me₂CO)(as diiodide). Mp 268° (diiodide). [α]_D²⁹ +13.8 (c, 0.65 in MeOH). CAS no. refers to diiodide.

O¹²-De-Me, O¹²-Me, N²-de-Me: **2'-Norfuniferine**

[116064-70-1]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). [α]_D +196 (c, 0.17 in CHCl₃).

O⁷,O¹²-Di-de-Me, O¹²-Me: **Tiliageine**

[53755-51-4]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the roots of *Tiliacora dinklagei* (Menispermaceae). Exhibits antiparasitic and cytotoxic activities. Needles (Et₂O/EtOAc). Mp 270°. [α]_D²⁵ +132.6 (c, 1.43 in Py). Abs. config. determined by tritium labelling studies.

O⁷,O¹²-Di-de-Me, O¹²-Me, N²-de-Me: **2'-Nortiliageine**

[116084-92-5]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the stem bark of *Guatteria guianensis* (Annonaceae). [α]_D +203 (c, 0.18 in CHCl₃).

Grundon, M.F. *et al.*, *J.C.S.*, 1960, 2739-2745 (*Rodiasine, Ocotine, isol, uv, ir*)

Hearst, P.J. *et al.*, *J.O.C.*, 1964, **29**, 466-470 (*Norrodiasine*)

Grundon, M.F. *et al.*, *J.C.S.(C)*, 1966, 1082-1084 (*Rodiasine, pmr, struct*)

Chan, K.C. *et al.*, *J.C.S.(C)*, 1967, 2479-2488 (*Ocotine, isol, uv, ord, pmr, ms*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 157-176 (*Funiferine, Tiliageine, activity*)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1973, **36**, 66-71 (*Tiliageine, Funiferine, isol, uv, ir, pmr, ms, struct*)

Tackie, A.N. *et al.*, *Experientia*, 1974, **30**, 847-848 (*Tiliageine*)

Cava, M.P. *et al.*, *J.O.C.*, 1974, **39**, 3588-3591 (*Phlebicine*)

Dwuma-Badu, D. *et al.*, *J. Pharm. Sci.*, 1977, **66**, 1242-1244 (*Funiferine N-oxide*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1978, **34**, 1409-1410 (*Tiliageine, biosynth, abs config*)

Cortes, D. *et al.*, *J. Nat. Prod.*, 1985, **48**, 76-85 (*Antioquine, isol, uv, ir, pmr, ms, cd, struct*)


Pachaly, P. *et al.*, *Planta Med.*, 1988, **54**, 516-519 (*Tilitriandrine*)

Chiaroni, P.A. *et al.*, *Acta Cryst. C*, 1990, **46**, 825-828 (*Antioquine, cryst struct*)

Roumy, V. *et al.*, *Planta Med.*, 2006, **72**, 894-898 (*isol, pmr, cmr*)

Roecarboline**R-115**

[131984-68-4]



C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Roemeria hybrida* (Papaveraceae). [α]_D -27 (c, 0.14 in CHCl₃). [α]_D -5 (c, 0.13 in MeOH).

N²-De-Me: **Norroecarboline**

[131984-69-5]


C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Roemeria hybrida* (Papaveraceae). [α]_D +35 (c, 0.07 in CHCl₃). [α]_D +4 (c, 0.21 in MeOH).

Gözler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 740 (*isol, uv, cd, ms, pmr, struct*)

Roehybridine**R-116**

[123000-13-5]



C₃₁H₃₉N₃O₅ 533.666

Alkaloid from *Roemeria hybrida* (Papaveraceae). Cryst. (MeOH). Mp 210–211°. [α]_D -16 (c, 0.056 in MeOH).

N⁶-Oxide (α -): **Roehybridine α -N-oxide** [409120-36-1]

C₃₁H₃₉N₃O₆ 549.666

Alkaloid from *Roemeria hybrida*. Amorph. solid. [α]_D -15 (c, 0.13 in MeOH). λ_{\max} 208 (log ϵ 4.89); 224 (sh) (log ϵ 4.67); 289 (log ϵ 3.91) (MeOH).

N-Oxide (β -): **Roehybridine β -N-oxide** [131669-93-7]

C₃₁H₃₉N₃O₆ 549.666

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -36 (c, 0.056 in MeOH).

1-Me ether: **O-Methylroehybridine** [131669-94-8]

C₃₂H₄₁N₃O₅ 547.693

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -17 (c, 0.187 in MeOH).

N-De-Me: **Norroehybridine**

[131690-43-2]

C₃₀H₃₇N₃O₅ 519.639

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -29 (c, 0.149 in MeOH).

4'-Demethoxy: **Roehybramine**

[131669-92-6]

C₃₀H₃₇N₃O₄ 503.64

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -13 (c, 0.185 in MeOH).

4'-Demethoxy, N⁶-oxide (β -): **Roehybramine β -N-oxide**

[409105-30-2]

C₃₀H₃₇N₃O₅ 519.639

Alkaloid from *Roemeria hybrida*. Amorph. solid. [α]_D -26.6 (c, 0.15 in MeOH). λ_{\max} 208 (log ϵ 4.83); 226 (log ϵ 4.65); 289 (log ϵ 3.81) (MeOH).

3',9-Bis(demethoxy): **Phoebegrandine A** [194737-72-9]

C₂₉H₃₅N₃O₃ 473.614

Alkaloid from *Phoebe grandis* (Lauraceae). Gum. Racemic. λ_{\max} 224 (log ϵ 4.55); 277 (log ϵ 3.97); 308 (sh) (log ϵ 3.51) (MeOH).

3',9-Bis(demethoxy), 1-Me ether: **Phoebescortechiniine**

[1007220-51-0]

C₃₀H₃₇N₃O₃ 487.641

Alkaloid from the leaves of *Phoebe scortechinii*. Amorph. gum. [α]_D²⁷ -3.3 (c, 1 in MeOH). λ_{\max} 215 (log ϵ 4.55); 272 (log ϵ 3.97) (MeOH).

3',9-Bis(demethoxy), 2-O-de-Me, 1-Me ether: **Phoebegrandine B**

[194737-77-4]

C₂₉H₃₅N₃O₃ 473.614

Alkaloid from *Phoebe grandis* (Lauraceae). Gum. Racemic. λ_{\max} 225 (log ϵ 4.55); 278 (log ϵ 3.97); 308 (sh) (log ϵ 3.51) (MeOH).

3',4',9-Tris(demethoxy), 8 α -hydroxy:

Phoebegrandine D

[909118-75-8]

C₂₈H₃₃N₃O₃ 459.587

Alkaloid from the leaves of *Phoebe grandis*. Amorph. brown solid. [α]_D²³ -22.2 (c, 0.16 in MeOH). λ_{\max} 233; 300 (MeOH).

3',4',9-Tris(demethoxy), 8 β -hydroxy:

Phoebegrandine C

[690957-78-9]

C₂₈H₃₃N₃O₃ 459.587

Alkaloid from the leaves of *Phoebe grandis*. Amorph. brown solid. [α]_D²³ +3.8 (c, 0.27 in MeOH). λ_{\max} 238; 301; 376 (MeOH).

9-Epimer: **Roehyimine. Misrhybridine**

[131724-52-2]

C₃₁H₃₉N₃O₅ 533.666

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D +29 (c, 0.065 in MeOH). Identity of Misrhybridine with Roehyimine not certain.

9-Epimer, 4'-demethoxy: **Roebromine**

[132075-36-6]

C₃₀H₃₇N₃O₄ 503.64

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D +7 (c, 0.092 in MeOH).

7a,10-Diepimer: **Roemeridine. Remeridine**

[122890-33-9]

C₃₁H₃₉N₃O₅ 533.666

Alkaloid from *Roemeria hybrida* and *Papaver pavoninum* (Papaveraceae). Cryst. (MeOH) or amorph. solid. Mp 236–237°. [α]_D -21 (c, 0.075 in MeOH).

7a,10-Diepimer, 1-Me ether: **O-Methylroemeridine**

C₃₂H₄₁N₃O₅ 547.693

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -18 (c, 0.317 in MeOH).

7a,10-Diepimer, 4'-demethoxy: **Roemebromine**

C₃₀H₃₇N₃O₄ 503.64

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. solid. [α]_D -21 (c, 0.193 in MeOH).

Platonova, T.F. *et al.*, *Zh. Obshch. Khim.*,

1956, **26**, 173-180; *CA*, **50**, 13960

(Roemeridine)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*,

1974, **39**, 888-894 (*Roehybridine, isol*)

Podlaha, J. *et al.*, *Phytochemistry*, 1989, **28**, 1779-1781 (*Roemeridine, pmr, cmr, ms, cryst struct*)

Gözler, B. *et al.*, *Tet. Lett.*, 1989, **30**, 1165-1168

(Roemeridine, Roehybridine, pmr, cmr, ms, struct)

El-Masry, S. *et al.*, *Alexandria J. Pharm. Sci.*,

1990, **4**, 90-93; *CA*, **114**, 160639

(Misrhybridine)

Gözler, B. *et al.*, *J. Nat. Prod.*, 1990, **53**, 675-

685 (*isol, uv, cd, pmr, ms, Roemeridine, β -oxide*)

Mukhtar, M.R. *et al.*, *Phytochemistry*, 1997,

45, 1543-1546 (*Phoebegrandine A,B*)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999,

37, 195-202 (*N-15 nmr*)

Gunes, H.S. *et al.*, *Fitoterapia*, 2001, **72**, 875-

887 (*Roehybridine α -N-oxide, Roehybramine β -N-oxide*)

Mukhtar, M.R. *et al.*, *Nat. Prod. Res.*, 2004,

18, 163-167; 2006, **20**, 567-572


(Phoebegrandine C,D)

Awang, K. *et al.*, *Nat. Prod. Res.*, 2007, **21**,

704-709 (*Phoebescortechiniine*)**Roehybrine***Rehybrine*

[52705-95-0]

R-117



Absolute Configuration

C₁₉H₂₅NO₄ 331.411

Alkaloid from *Roemeria hybrida* (Papaveraceae). Needles (Et₂O/hexane). Mp 191-192°. [α]_D²⁴ -67 (c, 0.29 in MeOH).

Perchlorate:

Long needles (H₂O or MeOH). Mp 290-291°.

10S-Alcohol: **α -Roemehybrine**

[112515-39-6]


C₁₉H₂₇NO₄ 333.427

Alkaloid from *Roemeria hybrida* (Papaveraceae). Amorph. [α]_D -49 (c, 0.15 in MeOH).

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 888 (*isol, uv, ir, ms, struct*)
Gözler, B. *et al.*, *Tetrahedron*, 1987, **43**, 1765 (*isol, uv, ir, pmr, ms, cd, struct, abs config*)

Roemecarine

R-118



(+)-form

C₂₀H₂₅NO₅ 359.421

The first naturally occurring C-4 hydroxylated tetrahydrobenzylisoquinoline alkaloid. Config. revised in 1987.

(+)-form

Alkaloid from *Roemeria carica* (Papaveraceae). Amorph. [α]_D +9 (c, 0.07 in MeOH).

2 α -N-Oxide: **Roemecarine 2 α -N-oxide**

C₂₀H₂₅NO₆ 375.421

Alkaloid from *Roemeria carica*. [α]_D +18 (c, 0.09 in MeOH).

(-)-form

Synthetic. [α]_D²⁷ -9.18 (c, 2.1 in MeOH).

(\pm)-form

Synthetic. Cryst. (C₆H₆/hexane). Mp 115.5-117°.

4-Epimer: Synthetic. Cryst. (C₆H₆/hexane). Mp 140-141.5°.

Gözler, T. *et al.*, *Heterocycles*, 1986, **24**, 1227


(isol, uv, pmr, ms, cd, struct, oxide)

Hoshino, O. *et al.*, *Chem. Pharm. Bull.*, 1990,

38, 3277 (*synth, abs config*)

Roemeramine

[21059-81-4]

Absolute
configuration $C_{18}H_{21}NO_3$ 299.369

Alkaloid from *Roemeria refracta* (Papaveraceae). Cryst., fine needles + 1Et₂O (Et₂O). Mp 66-67° (59-62° solvate). $[\alpha]_D^{22}$ -93 (c, 0.30 in MeOH). Browns rapidly in air, satisfactory anal. could not be obt. Chemically correlated with (+)-Mecambrine, the enantiomer of natural (-)-Mecambrine.

Hydrochloride:

Cryst. (HCl aq.). Mp 284-285°. Rel. stable in air.

10-Ketone: Roemeroinine

[21059-83-6]


 $C_{18}H_{19}NO_3$ 297.353

Alkaloid from *Roemeria refracta* (Papaveraceae). Prisms (Et₂O). Mp 130-135°. $[\alpha]_D^{23}$ -167 (c, 0.10 in MeOH). Extremely unstable in air.

Slávik, J. et al., *Coll. Czech. Chem. Comm.*, 1968, **33**, 4066

Roemerine**R-120**

6,7,7a,8-Tetrahydro-7-methyl-5H-benzoz[*g*]-1,3-benzodioxolo[6,5,4-*de*]/quinoline. 1,2-Methylenedioxyaporphine. N-Methylanonaine. Remerine†. Aporheine. Isoeroemerine. Isoeroemerine



(R)-form

 $C_{18}H_{17}NO_2$ 279.338

Aporheine and Isoeroemerine were names applied historically to the S-enantiomer.

(R)-form [548-08-3]

Alkaloid from *Roemeria refracta*, *Annona senegalensis* and *Nelumbo nucifera* (Nelumbonaceae). Present in the Annonaceae, Magnoliaceae, Menispermaceae and Rhamnaceae. Mp 102-103° (85-86°). $[\alpha]_D^{25}$ -97 (c, 0.20 in EtOH). λ_{max} 234 (log ϵ 4.17); 272 (log ϵ 4.21); 312 (log ϵ 4.21); 372 (log ϵ 3.52) (EtOH).

▶ **Mutagenic.** CE0908000**N-Oxide: Roemerine N-oxide**

[75088-35-6]

 $C_{18}H_{17}NO_3$ 295.337

Alkaloid from *Papaver* sp. (Papaveraceae) and *Liriodendron tulipifera*.

N-Me: (-)-Roemrefidine. Remrefidine. N-Methylroemerinium $C_{19}H_{20}NO_2^{\oplus}$ 294.372

Quaternary alkaloid from *Roemeria refracta* (Papaveraceae). Cryst. (MeOH aq.) (as hydroxide). Mp 223-224° (hydroxide).

N-De-Me: 1,2-Methylenedioxyaporphine. Anonaine

[1862-41-5]

 $C_{17}H_{15}NO_2$ 265.311

Alkaloid from *Annona reticulata*, *Annona muricata* (soursop) (Annonaceae) and *Nelumbo nucifera* (East India lotus) (Nelumbonaceae), also from the Lauraceae, Magnoliaceae, Monimiaceae, Papaveraceae, Rhamnaceae and Menispermaceae. Serotonin receptor antagonist. Shows cytotoxic, antimicrobial and insecticidal activity. Mp 122-123°. $[\alpha]_D^{21}$ -68 (c, 0.77 in EtOH). λ_{max} 233 (log ϵ 4.12); 273 (log ϵ 4.2); 314 (log ϵ 3.53) (EtOH).

N-De-Me, N-formyl: N-Formylanonaine

[83459-45-4]

 $C_{18}H_{15}NO_3$ 293.321

Alkaloid from the stem bark and root bark of *Hexalobus crispiflorus* (Annonaceae). Needles (MeOH). Mp 249-250°. $[\alpha]_D^{20}$ -319 (c, 0.1 in CHCl₃). λ_{max} 233 (sh); 274 (log ϵ 4.32); 292 (sh); 315 (log ϵ 3.73) (EtOH).

N-De-Me, N-Ac: N-Acetylanonaine

[5894-74-6]

 $C_{19}H_{17}NO_3$ 307.348

Alkaloid from the heartwood of *Magnolia obovata* and *Liriodendron tulipifera*, the roots of *Zanthoxylum bungeanum* and the root bark of *Zanthoxylum simulans* (Szechuan pepper) (Magnoliaceae, Rutaceae). Cryst. (EtOH). Mp 229-230°. $[\alpha]_D^{25}$ -356 (c, 0.49 in CHCl₃).

N-De-Me, N-methoxycarbonyl: Romucosine. N-Methoxycarbonylanonaine

[181718-71-8]

 $C_{19}H_{17}NO_4$ 323.348

Alkaloid from fresh unripe fruits of *Rollinia mucosa* (biriba). Needles (CHCl₃). Mp 152-153°. $[\alpha]_D^{25}$ -106.5 (c, 0.04 in CHCl₃).

N-De-Me, N-carbamoyl: N-Carbamoylanonaine

[83459-46-5]

 $C_{18}H_{16}N_2O_3$ 308.336

Alkaloid from the stem bark and root bark of *Hexalobus crispiflorus* (Annonaceae). Amorph. (diisopropyl ether). Mp 172° dec. $[\alpha]_D^{20}$ -336 (c, 0.49 in MeOH). λ_{max} 232 (log ϵ 4.22); 275 (log ϵ 4.15); 313 (log ϵ 3.53) (MeOH).

N-De-Me, N-nitroso: N-Nitrosoanonaine $C_{17}H_{14}N_2O_3$ 294.309

Alkaloid from *Duguetia furfuracea*. Cryst. (MeOH/CHCl₃). Mp 174-175°. $[\alpha]_D^{25}$ -298 (c, 0.5 in CHCl₃).

6a,7-Didehydro: see Dehydroroemerine, D-166**(S)-form** [2030-53-7]

Alkaloid from *Liriodendron tulipifera* (Magnoliaceae) and from the Papaveraceae. Low doses are hypotensive, high doses hypertensive. Strychnine-like convulsive agent in high doses. $[\alpha]_D^{25}$ +68 (c,

0.27 in EtOH).

N-Oxide: Mp 176-177° dec.

N-Me: Aporheine N-methosalt. (+)-Roemrefidine

[21153-67-3]

 $C_{19}H_{20}NO_2^{\oplus}$ 294.372

Quaternary alkaloid from *Papaver caucasicum*, *Papaver dubium*, *Papaver litwinowii*, *Papaver albiflorum* ssp. *astrumoravicum*, *Papaver fugax* and *Roemeria refracta* (Papaveraceae). Curare-like neuromuscular blocking agent. Small needles (H₂O or MeOH) (as iodide). Mp 236-237° (223-224°) (iodide). $[\alpha]_D^{25}$ +49 (c, 0.23 in MeOH). λ_{max} 270 (log ϵ 4.26); 315 (log ϵ 3.73) (no solvent reported).

N-De-Me: [58846-21-2] $C_{17}H_{15}NO_2$ 265.311

Alkaloid from *Neolitsea aurata* (Annonaceae).

(±)-form

Mp 85-87°.

N-De-Me: [7241-00-1] $C_{17}H_{15}NO_2$ 265.311

Alkaloid from *Xylopiya pancheri* (Annonaceae). Mp 116.5°.

N-Me: Mp 221°.

Yunousoff, S. et al., *Bull. Soc. Chim. Fr.*, 1939, **6**, 811-817 (*Roemerine*)

Marion, L. et al., *J.A.C.S.*, 1944, **66**, 1290-1292 (*Roemerine*)

Marion, L. et al., *Can. J. Res., Sect. B*, 1950, **28**, 21 (*Anonaine*)

Gopinath, K.W. et al., *Chem. Ber.*, 1959, **92**, 776-779 (*Anonaine, isol*)

Barton, D.H.R. et al., *Chem. Comm.*, 1966, 259-260 (*Roemerine*)

Cava, M.P. et al., *J.O.C.*, 1966, **31**, 1281-1283 (*Anonaine*)

Johns, S.R. et al., *Aust. J. Chem.*, 1968, **21**, 1383-1386 (*Anonaine, pmr, isol*)

Akramov, S.T. et al., *Khim. Prir. Soedin.*, 1968, **4**, 199; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 173 (*Roemrefidine*)

Bhakuni, D.S. et al., *Phytochemistry*, 1972, **11**, 1819-1822 (*Roemerine, Anonaine, ms, pmr, uv, isol*)

Hufford, C.D. et al., *Phytochemistry*, 1976, **15**, 1169-1171 (*N-Acetylanonaine*)

Nieto, M. et al., *Planta Med.*, 1976, **30**, 48-58 (*±-form, N-de-Me*)

Sashida, Y. et al., *Yakugaku Zasshi*, 1976, **96**, 659; *CA*, **88**, 3043u (*N-Acetylanonaine*)

Manushkayan, M.A. et al., *Khim. Prir. Soedin.*, 1977, **13**, 713; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 599 (*N-metho salt*)

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1980, **45**, 914-920; 1981, **46**, 1534-1538; 2587-2593 (*Roemrefidine, Aporheine*)

Gray, A.I. et al., *Planta Med.*, 1980, **39**, 209 (*N-Acetylanonaine*)

Phillipson, J.D. et al., *J. Nat. Prod.*, 1981, **44**, 296-307 (*Roemerine oxide*)

Ren, L. et al., *Yaoxue Xuebao*, 1981, **16**, 672-677; *CA*, **96**, 48974e (*N-Acetylanonaine*)


Achenbach, H. et al., *Annalen*, 1982, 1623-1633 (*N-Formylanonaine, N-Carbamoylanonaine*)

Chen, Y.-Y. et al., *J. Nat. Prod.*, 1996, **59**, 904-906 (*Romucosine*)

Hasrat, J.A. et al., *Phytomedicine*, 1997, **4**, 133-140 (*Anonaine, activity*)

Chen, C.-Y. et al., *J. Nat. Prod.*, 2000, **63**, 1475-1478 (*Anonaine, activity*)

Carollo, C.A. et al., *J. Nat. Prod.*, 2006, **69**, 1222-1224 (*N-Nitrosoanonaine*)

Rohitukine†[71294-60-5]
[97907-01-2]Absolute
ConfigurationC₁₆H₁₉NO₅ 305.33

Alkaloid from the leaves and stems of *Amoora rohituka*, the stem bark of *Dysoxylum binectariferum* and from *Schumannioophyton magnificum*. Shows antiinflammatory and immunomodulatory activity. Yellow cryst. Mp 218-219° Mp 227-232°. [α]_D²⁰ +44.3 (MeOH). Log P -0.49 (calc). λ_{\max} 208 (ϵ 23442); 262 (ϵ 12600); 330 (ϵ 4786) (MeOH) (Berdy).

Hydrochloride: Mp 242-243°. [α]_D -27.2 (MeOH).

N-Oxide: Rohitukine N-oxideC₁₆H₁₉NO₆ 321.329

Alkaloid from the stem bark of *Dysoxylum binectariferum*. Amorph. powder (MeOH). λ_{\max} 227; 252; 257; 297; 320 (sh) (no solvent reported).

N-De-Me, O^{3'}-Ac: N-Demethylrohitukine 3'-acetateC₁₇H₁₉NO₆ 333.34

Alkaloid from the stem bark of *Schumannioophyton magnificum*. Cream amorph. solid. Isol. as a mixt. of the 3' α - and 3' β - isomers in an approximately 2:1 ratio.

Harmon, A.D. *et al.*, *Tet. Lett.*, 1979, 721-724 (*isol, ir, ms, pmr, cmr, cryst struct*)

Houghton, P.J. *et al.*, *Planta Med.*, 1988, **54**, 239 (*N-Demethylrohitukine 3'-acetate*)

Naik, R.G. *et al.*, *Tetrahedron*, 1988, **44**, 2081-2086 (*isol, ir, pmr, synth*)


De Souza, N.J. *et al.*, *ACS Symp. Ser.*, 1993, **534**, 331 (*rev*)

Yang, D.H. *et al.*, *Chin. Chem. Lett.*, 2003, **14**, 720-723 (*cryst struct, abs config*)

Yang, D.-H. *et al.*, *J. Asian Nat. Prod. Res.*, 2004, **6**, 233-236 (*N-oxide*)

Rollipyrrole

R-122

C₁₆H₂₀N₂O₃ 288.346**(±)-form**

Constit. of the leaves of *Rollinia mucosa* (biriba). Amorph. yellow powder. Possible artifact. λ_{\max} 266 (no solvent reported).


Kuo, R.-Y. *et al.*, *Tet. Lett.*, 2001, **42**, 7907-7909

R-121

Romneine

R-123

5-[(3,4-Dimethoxyphenyl)methyl]-5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-g]isoquinoline, 9CI. 1-(3,4-Dimethoxybenzyl)-1,2,3,4-tetrahydro-6,7-methylenedioxy-2-methylisoquinoline

**(S)-form**C₂₀H₂₃NO₄ 341.406**(R)-form**

Alkaloid from the bark of *Laurelia novae-zelandiae* (Monimiaceae). Amorph. Has the same registry number as the (S)-form.

Hydrobromide: Mp 223-225°. [α]_D¹⁵ -75 (c, 0.2 in EtOH).

(S)-form [5544-49-0]

Alkaloid from the roots of *Romneya coulteri* var. *trichocalyx* (Papaveraceae). Oil. [α]_D²⁷ +37 (c, 0.11 in EtOH).

Hydrobromide: Mp 226-227°. [α]_D²⁸ +40 (c, 0.26 in EtOH). The stereochemical purity of this sample has been questioned.

N-Me: Escholinine

[51550-01-7]

C₂₁H₂₆NO₄[⊕] 356.441

Quaternary alkaloid from *Eschscholtzia californica* (Papaveraceae).

N-Me, iodide:C₂₁H₂₆INO₄ 483.345

Cryst. (MeOH). Mp 197-198°.

N-Me, perchlorate:C₂₁H₂₆ClNO₈ 455.891Prisms (MeOH). Mp 209-210°. [α]_D²⁵ +74 (c, 0.31 in MeOH).**O^{4'}-De-Me, N-Me: Phyllocryptine**

[126298-57-5]

[126262-25-7]

C₂₀H₂₄NO₄[⊕] 342.414

Quaternary alkaloid from various parts of *Cryptocarya phyllostemon* (Lauraceae). Yellow amorph. powder (as chloride). Positive opt. rotn. which could not be measured owing to the dark-coloured soln.

(±)-form [5544-50-3]

Alkaloid from leaves of *Cryptocarya chinensis* (Lauraceae). Pale yellow oil or amorph. solid.

Hydrobromide: Mp 194-196°.

Stermitz, F.R. *et al.*, *Tetrahedron*, 1966, **22**,

1095 (*isol, uv, ir, pmr, ms, struct, synth*)

Stermitz, F.R. *et al.*, *Tet. Lett.*, 1967, 1601 (*abs config*)

Slavk, J. *et al.*, *Coll. Czech. Chem. Comm.*,

1970, **35**, 2597; 1973, **38**, 3514*(Escholinine)*

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1974,

22, 966 (*synth, pmr*)

Urzúa, A. *et al.*, *Phytochemistry*, 1982, **21**, 773

(isol, pmr)


Cavé, A. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2243

(Phyllocryptine)

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1993, **56**, 227
(*isol, uv, ir, pmr, cmr, ms*)

Romucosine I

R-124

C₂₀H₂₁NO₅ 355.39


Alkaloid from the stems of *Rollinia mucosa* (biriba). Brown amorph. powder. λ_{\max} 251; 259; 304; 316 (MeOH).

Kuo, R.-Y. *et al.*, *Z. Naturforsch., B*, 2004, **59**, 334-336 (*isol, pmr*)

Roquefortine C

R-125

[58735-64-1]

Absolute
ConfigurationC₂₂H₂₃N₅O₂ 389.456

CAS numbering shown. The abs. config. ref. is inaccessible but the results are given in the 1998 Synthesis ref. Metab. of *Penicillium roquefortii* ATCC 10110, *Penicillium commune*, *Penicillium cyclopium* NRRL 6093, *Penicillium farinosum* and many other *Penicillium* spp. incl. a marine-derived sp. Common constit. of blue cheese. Neurotoxin. Needles + 1MeOH (MeOH aq.). Mp 195-200° dec Mp 225-228° dec. [α]_D²² -703 (c, 1 in CHCl₃). [α]_D¹⁵ -764 (c, 0.5 in Py). λ_{\max} 210 (ϵ 31620); 243 (ϵ 18200); 325 (ϵ 28800) (EtOH).

▶ LD₅₀ (mus, ipr) 15 mg/kg. UQ4730500**N⁶-Formyl: N⁶-Formylroquefortine C**

[158200-10-3]

C₂₃H₂₃N₅O₃ 417.466

Metab. of *Penicillium verrucosum* var. *cyclopium* AV 67718. Mycotoxin. Solid (MeOH aq.). Mp 216-219° dec. λ_{\max} 207; 242; 326 (MeOH) (Berdy).

N¹⁶-Et: N¹⁶-Ethylroquefortine C

[174792-02-0]

C₂₄H₂₇N₅O₂ 417.51Prod. by *Penicillium janczewskii*.**3 α ,12-Dihydro: 3,12-Dihydroroquefortine.****Roquefortine D. Alkaloid Z**


[58735-66-3]

C₂₂H₂₅N₅O₂ 391.472

Prod. by *Penicillium* sp. incl. *Penicillium roquefortii* ATCC 10110. Neurotoxin. Prisms (MeCN). Mp 153-154°. [α]_D¹⁵ -370 (c, 0.23 in Py). λ_{\max} 208 (ϵ 195000); 244 (ϵ 35480); 298 (ϵ 15140) (EtOH).

▶ Toxic.


3 α ,12-Dihydro, N¹⁶-ethoxycarbonyl: N¹⁶-Ethoxycarbonylroquefortine D. N¹⁶-

Carboxyethylidihydroroquefortine CC₂₅H₂₉N₅O₄ 463.535Prod. by *Penicillium aureovirens* VKM FW-766. Wax. $[\alpha]_D^{22}$ -66.8 (c, 10.1 in MeOH). λ_{\max} 210; 239; 300 (MeOH).Ohmomo, S. *et al.*, *Agric. Biol. Chem.*, 1975, **39**, 1333-1334; 1977, **41**, 2097-2098; 1978, **42**, 2387-2389; 1979, **43**, 2035-2038; 1980, **44**, 1929-1930 (*isol, uv, ir, pmr, ms, biosynth, Roquefortine D*)Scott, P.M. *et al.*, *Experientia*, 1976, **32**, 140-142 (*isol, uv, ir, pmr, cmr, struct*)Barrow, K.D. *et al.*, *Chem. Comm.*, 1979, 225-226 (*biosynth*)Scott, P.M. *et al.*, *J. Agric. Food Chem.*, 1979, **27**, 201-202 (*config*)Engel, G. *et al.*, *J. Chromatogr.*, 1979, **170**, 288-291 (*tlc, uv*)Wagner, R.E. *et al.*, *Appl. Environ. Microbiol.*, 1980, **39**, 882-887 (*isol, uv, ir, ms*)Vesonder, R.F. *et al.*, *Experientia*, 1980, **36**, 1308 (*isol, uv, ir*)Kozlovsky, A.G. *et al.*, *Experientia*, 1981, **37**, 472-473 (*Roquefortine D, isol, biosynth*)Gorst-Allman, C.P. *et al.*, *Chem. Comm.*, 1982, 652-653 (*biosynth, cmr*)Mantle, P.G. *et al.*, *Appl. Environ. Microbiol.*, 1983, **45**, 1486-1490 (*biosynth*)Scott, P.M. *et al.*, *Dev. Food Sci.*, 1984, **8**, 463-468 (*rev*)Yamaguchi, T. *et al.*, *Proc. Jpn. Assoc. Mycotoxicol.*, 1991, **34**, 29-32 (*abs config*)Bhat, B. *et al.*, *Tetrahedron*, 1993, **49**, 10663-10669 (*biosynth*)Musuku, A. *et al.*, *J. Nat. Prod.*, 1994, **57**, 983-987 (*N⁶-Formylroquefortine C*)Kozlovsky, A.G. *et al.*, *Prikl. Biokhim. Mikrobiol.*, 1997, **33**, 70-74 (*N-Ethylroquefortine*)Chen, W.-C. *et al.*, *Tet. Lett.*, 1998, **39**, 8401-8404 (*Roquefortine D, synth*)Kozlovsky, A.G. *et al.*, *Heterocycles*, 2003, **60**, 1639-1644 (*Ethoxycarbonylroquefortine D*)Bringmann, G. *et al.*, *Tetrahedron*, 2005, **61**, 7252-7265 (*marine isol*)Shangguan, N. *et al.*, *J.A.C.S.*, 2008, **130**, 6281-6287 (*synth, props*)**Roquefortine E****R-126**


Absolute Configuration

C₂₇H₃₁N₅O₂ 457.574Prod. by *Gymnoascus reessii* (MST-F9977). Amorph. solid. $[\alpha]_D$ -223 (c, 0.06 in CHCl₃). λ_{\max} 228 (ε 13600); 342 (ε 11100) (MeOH).Clark, B. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1661-1664 (*isol, pmr, cmr*)**Roseadine****R-127**


[58005-25-7]

C₄₆H₅₆N₄O₉ 808.97Alkaloid from *Catharanthus roseus* (Apocynaceae). Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{\max} 214 (ε 49000); 225 (ε 147000); 286 (ε 12000); 294 (ε 12400); 326 (ε 8500) (EtOH) (Berdy).El-Sayed, A. *et al.*, *J. Nat. Prod.*, 1983, **46**, 517 (*isol, uv, ir, pmr, cmr, ms, struct*)**Rosellichalasin****R-128**

[123452-64-2]

C₂₈H₃₃NO₅ 463.572Stereochem. of the compds. in this entry not all determined. Metab. of *Rosellinia necatrix*. Plant growth inhibitor. Needles (Me₂CO/hexane). Mp 121-123°. $[\alpha]_D^{20}$ -2.7 (c, 1 in MeOH). λ_{\max} 245 (ε 12500) (EtOH) (Derep).*16-Hydroxy, 18,19-dihydro:*C₂₈H₃₅NO₆ 481.588Prod. by *Rhinochadiella* sp. Powder.*22-Hydroxy, 18,19-dihydro:*C₂₈H₃₅NO₆ 481.588Prod. by *Rhinochadiella* sp. Powder.*A¹⁹-Isomer, 17-alcohol:*C₂₈H₃₅NO₅ 465.588Prod. by *Rhinochadiella* sp. Powder.Kimura, Y. *et al.*, *Agric. Biol. Chem.*, 1989, **53**, 1699-1701 (*isol, pmr, cmr*)Wagenaar, M.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1692-1695 (*Rhinochadiella* constits)**Roseophilin****R-129**

[142386-38-7]




Absolute Configuration

C₂₇H₃₃ClN₂O₂ 453.023Prod. by *Streptomyces griseoviridis*. Cytotoxic agent. Mp 137-178° dec. λ_{\max} 296 (ε 24800); 477 (ε 29800); 572 (ε 14400) (MeOH/NaOH) (Derep). λ_{\max} 306 (ε 15000); 364 (ε 8800); 378 (ε 8600); 533 (ε 112400) (MeOH) (Derep).*Hydrochloride:* [142435-87-8]


Reddish-purple powder.

Hayakawa, Y. *et al.*, *Tet. Lett.*, 1992, **33**, 2701-2704 (*isol*)Fürstner, A. *et al.*, *J.A.C.S.*, 1998, **120**, 2817-2825 (*synth, pmr, cmr*)Trost, B.M. *et al.*, *J.A.C.S.*, 2000, **122**, 3801-3810 (*synth*)Bamford, S.J. *et al.*, *Org. Lett.*, 2000, **2**, 1157-1160 (*synth*)Harrington, P.E. *et al.*, *J.A.C.S.*, 2001, **123**, 8509-8514 (*synth, abs config*)Boger, D.L. *et al.*, *J.A.C.S.*, 2001, **123**, 8515-8519 (*synth, cd, abs config*)Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (*rev, synth, chem, biol, history*)**Roserine****R-130***1,2,3,3a,4,5-Hexahydro-9,10,11-trimethoxyppyrrrolo[3,2,1-de]phenanthridinium(1+)*, 9C1

[139955-89-8]

C₁₈H₂₂NO₃⁺ 300.377Alkaloid from aerial parts and bulbs of *Narcissus pallidulus* (Amaryllidaceae). Counterion not specified.Bastida, J. *et al.*, *J. Nat. Prod.*, 1992, **55**, 134 (*isol, pmr, cmr, ms, struct*)Bunnell, A.E. *et al.*, *J.O.C.*, 1997, **62**, 9305-9313 (*synth*)**Rosibiline****R-131**

[76247-84-2]




Absolute Configuration

C₂₀H₂₄N₂O 308.422Alkaloid from the root bark of *Strychnos variabilis* (Loganiaceae). Powder.*16-Epimer: Isorosibiline*

[55969-04-5]


C₂₀H₂₄N₂O 308.422Alkaloid from *Strychnos floribunda* stem bark and from *Strychnos mato-pensis* (Loganiaceae).Tits, M. *et al.*, *Phytochemistry*, 1980, **19**, 1531-1534 (*Rosibiline, isol, uv, cd, ms, pmr, struct*)Verpoorte, R. *et al.*, *Planta Med.*, 1981, **42**, 32-36 (*Isorosibiline, isol, ms, pmr, synth*)Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304 (*Isorosibiline, isol*)**Rosicine****R-132**

[95690-65-6]



Absolute Configuration

C₁₉H₂₀N₂O₃ 324.379A dinoraspidosperma alkaloid. Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Pale-yellow gum. $[\alpha]_D$

+188.67 (CHCl₃).Atta-ur-Rahman, *et al.*, *Tet. Lett.*, 1984, **25**, 6051 (*isol, uv, ir, pmr, cmr, ms, struct*)**Rosmarinine****R-133**1,2-Dihydro-2,12-dihydroxysenecionan-11,16-dione, 9CI
[520-65-0]

Absolute configuration

C₁₈H₂₇NO₆ 353.414

Cyclic diester of Rosmarinecine in D-597 with Senecic acid. Alkaloid from *Senecio rosmarinifolius*, *Senecio taiwanensis*, *Senecio brachyphodoides*, *Senecio triangularis*, *Senecio pauciligulatus*, *Senecio hygrophyllus*, *Senecio angulatus* and *Senecio halimifolius* (Asteraceae). Laminae (Me₂CO or EtOH). Mp 209° (205-208°). [α]_D -120 (CHCl₃). [α]_D -94 (EtOH). Not hepatotoxic. Dehydration gives Senecionine, S-238.

12-Ac: 12-O-Acetytrosmarinine

[137760-53-3]

C₂₀H₂₉NO₇ 395.452

Alkaloid from *Senecio hadiensis* (Asteraceae). Pale amber oil. [α]_D -36 (c, 1.8 in EtOH).

13,19-Didehydro: Angularine†

C₁₈H₂₅NO₆ 351.399

Alkaloid from *Senecio angulatus* (Asteraceae). Cryst. (EtOAc). Mp 200-201°. [α]_D²⁵ -98 (c, 0.022 in EtOH). Cyclic diester of Rosmarinecine in D-597 with Seneciphillic acid.

(15E)-Isomer: Neorosmarinine

[137821-11-5]

C₁₈H₂₇NO₆ 353.414

Alkaloid from *Senecio hadiensis* (Asteraceae). Gum; cryst. (EtOH)(as perchlorate). Mp 205-207° (perchlorate). [α]_D -12 (c, 0.6 in EtOH).

Stereoisomer: Isorosmarinine

[155322-44-4]


C₁₈H₂₇NO₆ 353.414

Trace alkaloid from *Senecio pterophorus* (Asteraceae). Mp 137-142°.

Richardson, M.F. *et al.*, *J.C.S.*, 1943, 452 (*isol, bibl, struct*)Porter, L.A. *et al.*, *J.O.C.*, 1962, **27**, 4132 (*Angularine*)Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918 (*cd*)Lu, S.T. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1972, **19**, 127 (*isol*)Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1203 (*isol, cmr*)Freer, A.A. *et al.*, *Acta Cryst. C*, 1986, **42**, 1348 (*cryst struct*)Kunec, E.K. *et al.*, *Chem. Comm.*, 1986, 250 (*biosynth*)Were, O. *et al.*, *J. Nat. Prod.*, 1991, **54**, 491 (*12-O-Acetytrosmarinine, Neorosmarinine*)Liddell, J.R. *et al.*, *Phytochemistry*, 1993, **34**, 1629 (*Isorosmarinine*)**Rotundifoline**

Stipulatine

[6883-25-6]



Absolute Configuration

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Mitragyna rotundifolia* and several other *Mitragyna* spp. (Rubiaceae). Mp 238-240° (248-249°). [α]_D¹⁵ +124 (CHCl₃). λ_{max} 223 (log ε 4.36); 242 (sh) (log ε 4.15); 292 (log ε 3.42) (EtOH).

Hydrochloride: Mp 222°.

β-N^b-Oxide: Rotundifoline N-oxide

[64470-93-5]

C₂₂H₂₈N₂O₆ 416.473

Alkaloid from *Mitragyna rubrostipulata* (Rubiaceae). Noncryst. λ_{max} 222; 242 (sh); 292 (EtOH).

Me ether: Rhynchociline. Rhynchociline

[18002-97-6]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from *Mitragyna ciliata*, *Mitragyna inermis*, *Mitragyna tubulosa*, *Mitragyna parvifolia*, *Mitragyna rotundifolia* and *Uncaria* spp. (Rubiaceae). Mp 178-180°. [α]_D²⁰ +6.2 (c, 2 in CHCl₃).

Me ether; perchlorate: Mp 221-222°.

18,19-Didehydro: Rotundifoleine

[55872-09-8]

C₂₂H₂₆N₂O₅ 398.458

Alkaloid from *Mitragyna parvifolia* (Rubiaceae). λ_{max} 223; 243 (sh); 286; 295 (EtOH).

18,19-Didehydro, Me ether: Isospecionoxeine

[18003-00-4]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Mitragyna speciosa* (Rubiaceae). Cryst. (Et₂O/hexane). Mp 179°. λ_{max} 223 (log ε 4.46); 244 (log ε 4.26); 288 (log ε 3.52) (EtOH).

18,19-Didehydro, Me ether, perchlorate:

Cryst. (EtOH/Et₂O). Mp 230-231°.

7-Epimer: Isorotundifoline. Mitragynol

[6884-20-4]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Mitragyna ciliata* and some other *Mitragyna* spp. (Rubiaceae). Mp 131-132° Mp 200° (dimorph.). λ_{max} 222 (log ε 4.43); 242 (sh) (log ε 4.13); 290 (log ε 3.49) (EtOH).

7-Epimer, Me ether: Ciliaphylline

[18002-98-7]

C₂₃H₃₀N₂O₅ 414.5

Alkaloid from *Mitragyna ciliata*, *Mitragyna inermis*, *Mitragyna tubulosa*, *Mitragyna parvifolia*, *Mitragyna rotundifolia* and *Uncaria* spp. (Rubiaceae). Mp 222-223°. [α]_D^{25.5} -89.5 (c, 0.65 in CHCl₃). λ_{max} 222 (log ε 4.44); 244 (log ε 4.24); 287 (log ε 3.46) (EtOH).

7-Epimer, Me ether, picrate: Mp 130-131°.

7-Epimer, Me ether, N-oxide: Ciliaphyl-

R-134**line N-oxide**

[50506-22-4]

C₂₃H₃₀N₂O₆ 430.5

Alkaloid from *Mitragyna tubulosa* (Rubiaceae). λ_{max} 221; 243; 297 (no solvent reported).

7-Epimer, 18,19-didehydro: Isorotundifoleine

[55923-84-7]

C₂₂H₂₆N₂O₅ 398.458

Alkaloid from *Mitragyna parvifolia* (Rubiaceae). λ_{max} 222; 224 (sh); 278; 294 (EtOH).

7-Epimer, 18,19-didehydro, Me ether: Specionoxeine

[18002-99-8]

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Mitragyna speciosa* (Rubiaceae). Cryst. (Me₂CO). Mp 225°. λ_{max} 223 (log ε 4.49); 245 (log ε 4.18); 288 (log ε 3.29) (EtOH).

20-Epimer: Mitrafoline

[55903-79-2]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae). λ_{max} 224; 243 (sh); 292 (EtOH).

3,20-Diepimer: Isospeciofoline

[56687-49-1]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae).

7,20-Diepimer: Isomitrafoline

[55903-80-5]

C₂₂H₂₈N₂O₅ 400.474


Alkaloid from the leaves of *Mitragyna speciosa* (Rubiaceae). λ_{max} 225; 237 (sh); 290 (EtOH).

3,7,20-Triepimer: Speciofoline

[5171-42-6]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from leaves of *Mitragyna speciosa* (Rubiaceae). Needles. Mp 202-204°. [α]_D²² -103 (c, 2 in CHCl₃). pK_a 6.3 (H₂O). λ_{max} 223 (log ε 4.47); 242 (sh) (log ε 4.27); 290 (log ε 3.49) (EtOH).

Beckett, A.H. *et al.*, *Tet. Lett.*, 1963, 1709-1714 (*Rotundifoline, Speciofoline*)Trager, W.F. *et al.*, *Tetrahedron*, 1968, **24**, 523-543 (*Ciliaphylline, Specionoxeine, Isospecionoxeine*)Beckett, A.H. *et al.*, *Tetrahedron*, 1969, **25**, 5961-5969 (*ms*)Phillipson, J.D. *et al.*, *Phytochemistry*, 1973, **12**, 2043-2048 (*Ciliaphylline oxide*)Hemingway, S.R. *et al.*, *Phytochemistry*, 1975, **14**, 557-563 (*Mitrafoline, Isomitrafoline, Rotundifoleine, Isospeciofoline*)Shellard, E.J. *et al.*, *Phytochemistry*, 1977, **16**, 1427-1429 (*Rotundifoline oxide*)**Rotundine B****R-135**

C₁₅H₂₃NO 233.353

Alkaloid from the rhizomes of *Cyperus rotundus* (nutgrass). Oil. $[\alpha]_D^{25}$ -14.7 (c, 0.15 in CHCl₃). λ_{\max} 264 (log ϵ 3.4) (MeOH).

13-Ketone: Rotundine AC₁₅H₂₁NO 231.337

Alkaloid from the rhizomes of *Cyperus rotundus* (nutgrass). Oil. $[\alpha]_D^{27}$ -12.3 (c, 0.31 in CHCl₃). λ_{\max} 260 (log ϵ 3.15) (MeOH).


6-Epimer: Rotundine CC₁₅H₂₃NO 233.353

Alkaloid from the rhizomes of *Cyperus rotundus* (nutgrass). Oil. $[\alpha]_D^{25}$ -10.2 (c, 0.11 in CHCl₃). λ_{\max} 262 (log ϵ 3.48) (MeOH).

Jeong, S.-J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 673-675

Routiennocin**R-136**

Antibiotic CP 61405. CP 61405 [99623-84-4]

C₂₆H₃₀N₂O₇ 482.532

Polyether antibiotic related to Calcimycin, C-26. Prod. by *Streptomyces routienii*. Active against gram-positive bacteria. Controls swine dysentery. Sol. MeOH, C₆H₆; poorly sol. H₂O. λ_{\max} 258 (E₁%/1cm 287); 308 (E₁%/1cm 400) (MeOH) (Berdy). λ_{\max} 258 (E₁%/1cm 277); 269 (E₁%/1cm 247); 290 (E₁%/1cm 331) (MeOH/HCl) (Berdy). λ_{\max} 254 (E₁%/1cm 247); 310 (E₁%/1cm 356) (MeOH/NaOH) (Berdy).

U.S. Pat., 1985, 4 547 523; CA, **104**, 49844
Kotecha, N.R. *et al.*, *Synlett*, 1992, 395; 399 (synth)

Diez-Martin, D. *et al.*, *Tetrahedron*, 1992, **48**, 7899 (synth)

Rovidine**R-137**


[1360-83-4]

Bisindole alkaloid. Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Thin blades (EtOH) (as sulfate). Mp 320° dec. (sulfate). Mol. formula not reported. λ_{\max} 214 (E₁%/1cm 513); 265 (E₁%/1cm 153); 296 (E₁%/1cm 112) (EtOH) (Berdy).

Sulfate:

Thin blades (EtOH). Mp 320° dec.

Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1964, **53**, 1227-1231 (isol, uv, ir)

Roxburghines**R-138**C₃₁H₃₂N₄O₂ 492.619**(3R,15S,19R,20R)-form****Roxburghine B**

[27932-46-3]

Isol. from the leaves of *Uncaria gambir* (Rubiaceae). Mp 255°. $[\alpha]_D^{20}$ -350 (MeOH).

(3R,15S,19R,20S)-form**Roxburghine E**

[27773-33-7]

Isol. from the leaves and stems of *Uncaria gambir* (Rubiaceae). Mp 234-236°. $[\alpha]_D^{20}$ -105 (MeOH).

(3R,15S,19S,20S)-form**Roxburghine D. Roxburghine**

[27802-03-5]

Isol. from the leaves and stems of *Uncaria gambir* and from the bark of *Uncaria elliptica* (Rubiaceae). Faint-yellow cryst. (EtOH). Mp 197-200°. $[\alpha]_D^{20}$ +160 (MeOH).

(3S,15S,19S,20S)-form**Roxburghine C**

[27773-32-6]

Isol. from the leaves of *Uncaria gambir* (Rubiaceae). Mp 245-250°. $[\alpha]_D^{20}$ -221 (MeOH).

(3ξ,15S,19ξ,20ξ)-form**Roxburghine X**

[73651-11-3]

Isol. from the bark of *Uncaria elliptica* (Rubiaceae). Faint-pink cryst. (EtOH). Mp 215°. $[\alpha]_D^{27}$ -29 (MeOH).

(3ξ,15ξ,19ξ,20ξ)-form**Roxburghine A**

[30908-03-3]

Alkaloid from the leaves of *Uncaria gambir* (Rubiaceae). Mp 290-295°. $[\alpha]_D^{20}$ -264 (MeOH).

Merlini, L. *et al.*, *Tetrahedron*, 1970, **26**, 2259 (isol, uv, ir, pmr, ms, ord, struct)

Riesner, H. *et al.*, *Chem. Comm.*, 1972, 786 (synth)


Cistaro, C. *et al.*, *Gazz. Chim. Ital.*, 1973, **103**, 153 (pmr)

Benz, G. *et al.*, *Chem. Ber.*, 1975, **108**, 248 (synth)

Cistaro, C. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2249 (pmr, config)

Merlini, L. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2254 (cmr)

Herath, W.H.M.W. *et al.*, *Phytochemistry*, 1979, **18**, 1385 (Roxburghine X)

Roxiamine A**R-139**

R = COOMe

C₁₈H₁₉NO₅ 329.352**(S)-form** [175097-25-3]

Alkaloid from *Evodia roxburghiana*. Yellow oil. $[\alpha]_D$ +2 (c, 1.0 in CHCl₃). 2',3'-Didehydro (E-): **Roxiamine B**

[175097-26-4]

C₁₈H₁₇NO₅ 327.336From *Evodia roxburghiana*. Solid.

McCormick, J.L. *et al.*, *J. Nat. Prod.*, 1996, **59**, 469 (isol, uv, ir, pmr, cmr, ms, struct)

Roxiamine C**R-140**

As Roxiamine A, R-139 with

R = OH


C₁₆H₁₇NO₄ 287.315**(S)-form** [175097-27-5]

Alkaloid from *Evodia roxburghiana*. Solid. $[\alpha]_D$ +4 (c, 1.0 in CHCl₃).

McCormick, J.L. *et al.*, *J. Nat. Prod.*, 1996, **59**, 469 (isol, uv, ir, pmr, cmr, ms, struct)

Rubenine**R-141**

[52467-52-4]

C₂₈H₃₂N₂O₁₁ 572.568


Alkaloid from the heartwood of *Adina rubescens* (Nucleaceae). $[\alpha]_D^{25}$ -42 (MeOH).

Tetra-O-Ac: Mp 154-156° (as picrolonate). $[\alpha]_D$ -46 (CHCl₃).

Brown, R.T. *et al.*, *Chem. Comm.*, 1973, 765 (pmr, ms, cd, struct)

Rubesamide**R-142**

N-[2-(1,3-Benzodioxol-5-yl)ethyl]cyclopropanecarboxamide, 9CI. N-(3,4-Methylenedioxyphenethyl)cyclopropanecarboxamide [58738-58-2]


C₁₃H₁₅NO₃ 233.266

Alkaloid from *Fagara rubescens* root bark (Rutaceae). Needles (C₆H₆/petrol). Mp 131.5-132.5°.

Dadson, B.A. *et al.*, *J.C.S. Perkin 1*, 1976, 146 (isol, uv, ir, pmr, ms, synth)

Rubescenamine**R-143**

[125187-30-6]


C₂₀H₂₁NO₅ 355.39

Alkaloid from the stem bark of *Zanthoxylum rubescens* (Rutaceae). Mp

145-147°.

N-Me: Rubescenamide

[125187-29-3]

C₂₁H₂₃NO₅ 369.416Alkaloid from the stem bark of *Zanthoxylum rubescens* (Rutaceae). Amorph. solid.Adesina, S.K. *et al.*, *Planta Med.*, 1989, **55**, 324 (*isol, uv, ir, pmr, cmr, ms, synth, struct*)**Rubijervine****R-144**22 α H,25 β H-Solanid-5-ene-3 β ,12 α -diol
[79-58-3]

Absolute configuration

C₂₇H₄₃NO₂ 413.642Alkaloid from *Veratrum album*, *Veratrum californicum*, *Veratrum eschscholtzii*, *Veratrum nigrum*, *Veratrum viride*, *Veratrum lobelianum* and *Veratrum oxyssepalum* (Liliaceae). Shows antifungal activity. Mp 240-242°. [α]_D +19 (EtOH).

▶ WF0300000

Hydrobromide: Mp 265-270°.**Hydroiodide:** Mp 293-296°.**Di-Ac:** Mp 160-163°.**Dibenzoyl:** Mp 186-187.5°.**12-Epimer: Epirubijervine**


[472-00-4]

C₂₇H₄₃NO₂ 413.642Alkaloid from illuminated *Veratrum grandiflorum* (Liliaceae). Mp 231-234°. [α]_D -32.3 (c, 0.22 in CHCl₃).**12-Epimer, dibenzoyl:** Mp 264-267°.Jacobs, W.A. *et al.*, *J. Biol. Chem.*, 1943, **148**, 41 (*isol, synth, Epirubijervine*)Sato, Y. *et al.*, *J. Biol. Chem.*, 1949, **179**, 623 (*struct*)Höhne, E. *et al.*, *Tetrahedron*, 1966, **22**, 673 (*config*)Wolters, B. *et al.*, *Planta Med.*, 1970, **19**, 189-193 (*activity*)Kaneko, K. *et al.*, *Phytochemistry*, 1975, **14**, 1295 (*biosynth*)Kaneko, K. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 2534 (*Epirubijervine*)Machaidze, N.L. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 659; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 628 (*isol*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SKR000**Rubiverine****R-145**

[1360-91-4]

C₂₅H₃₉NO₂ 385.589Steroidal alkaloid. Struct. unknown. Alkamine from *Veratrum album* (Liliaceae). Mp 245-247°. [α]_D²⁰ 0 (c, 0.5 in CHCl₃).Cionga, E. *et al.*, *Acta Pol. Pharm.*, 1957, **14**, 73-76; *CA*, **52**, 12882c**Rubradirin B****R-146**


[68833-11-4]

C₄₀H₃₃N₃O₁₅ 795.712Isol. from *Streptomyces achromogenes rubradiris*. Active against gram-positive bacteria. Red cryst. (CHCl₃). Mp 300° dec. λ_{\max} 235 (sh) (ϵ 34000); 280 (ϵ 25000); 312 (ϵ 33000); 515 (ϵ 1580) (85% MeOH/pH 2) (Derep). λ_{\max} 300 (ϵ 42500); 345 (sh) (ϵ 18000); 412 (ϵ 11000) (85% MeOH/pH 11) (Derep). λ_{\max} 235 (sh) (ϵ 34000); 280 (ϵ 25000); 312 (ϵ 33000); 515 (ϵ 1580) (acidic solvent) (Derep). λ_{\max} 222 (ϵ 47900); 270 (ϵ 19960); 312 (ϵ 24950); 510 (ϵ 1711) (EtOH-HCl) (Berdy). λ_{\max} 285 (ϵ 25000); 315 (ϵ 32500); 512 (MeOH-HCl) (Berdy).**3''-Deoxy: Rubradirin C**

[69279-50-1]

C₄₀H₃₃N₃O₁₄ 779.712From *Streptomyces achromogenes rubradiris*. Less biol. active than other rubradirins. Cryst. (DMSO). Sol. bases, DMSO, DMF; fairly sol. MeOH, THF, EtOH, CHCl₃; poorly sol. H₂O, hexane, C₆H₆, toluene. Mp 300° dec. λ_{\max} 235 (sh) (ϵ 34000); 280 (ϵ 25000); 312 (ϵ 33000); 515 (ϵ 1580) (85% MeOH/pH 2) (Derep). λ_{\max} 300 (ϵ 42500); 345 (sh) (ϵ 18000); 412 (ϵ 11000) (85% MeOH/pH 11) (Derep). λ_{\max} 235 (sh) (ϵ 34000); 280 (ϵ 25000); 312 (ϵ 33000); 515 (ϵ 1580) (acidic solvent) (Derep). λ_{\max} 203 (ϵ 86205); 220 (ϵ 41644); 271 (ϵ 26239); 320 (ϵ 30400); 530 (MeOH-HCl) (Berdy). λ_{\max} 302 (ϵ 35818); 410 (ϵ 14100) (MeOH-NAOH) (Berdy).Bhuyan, B.K. *et al.*, *Antimicrob. Agents Chemother.*, 1964, **91**; 97 (*isol*)Hoeksema, H. *et al.*, *J. Antibiot.*, 1978, **31**, 945 (*isol, uv, ir, pmr, cmr*)Hoeksema, H. *et al.*, *J.A.C.S.*, 1982, **104**, 5173 (*isol, struct, pmr, cmr*)**Rubrifacine****R-147**


[91403-60-0]



1680

C₁₃H₁₀N₂O₉ 338.23Red pigment from *Erwinia rubrifaciens*. λ_{\max} 295 ; 350 ; 490 (pH 2 buffer) (Berdy). λ_{\max} 350 ; 522 (pH 9 buffer) (Berdy).Feistner, G. *et al.*, *Can. J. Chem.*, 1985, **63**, 495 (*cmr, ms, struct*)**Rubrobramide****R-148**

[205386-57-8]




Relative Configuration

C₁₂H₁₇NO₅ 255.27Prod. by *Cladobotryum rubrobrunnescens*. Phytotoxic and cytotoxic agent. Cryst. (EtOH). Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 131-133°. [α]_D +177 (c, 1 in CHCl₃).Wagner, C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 501-502 (*isol, ir, pmr, cmr, ms*)**Rubrocurarines****R-149**

[1360-95-8 (Rubrocurarine IV), 1360-92-5 (Rubrocurarine I), 1360-94-7 (Rubrocurarine III), 1360-93-6 (Rubrocurarine II)]

Struct. unknown. Alkaloids from *Strychnos solimoesana* (Loganiaceae). Rubrocurarines I, II, III and IV identified chromatographically, all giving red spots. Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148**Rubrolone****R-150**

[65445-21-8]



Absolute Configuration

C₂₃H₂₃NO₈ 441.437The opposite abs. config. is given in CAS. Prod. by *Streptomyces echinoruber* NRRL8144. Red pigment. Sol. H₂O, MeOH; fairly sol. EtOH; poorly sol. butanol, hexane. [α]_D²⁵ -937.5 (c, 0.2 in H₂O). A second pigment, prob. isomeric, was also isol. λ_{\max} 216 (ϵ 18800); 275 (ϵ 31500); 421 (ϵ 9500); 523 (ϵ 8600) (H₂O) (Berdy).

▶ UQ0350000


Palleroni, N.J. *et al.*, *J. Antibiot.*, 1978, **31**, 1218-1225 (*isol*)Schuop, W. *et al.*, *J. Antibiot.*, 1978, **31**, 1226-1232 (*cryst struct*)Sweeny, J.G. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 1292-1293 (*props*)Kelly, T.R. *et al.*, *Tet. Lett.*, 1986, **27**, 6049-6050 (*synth*)Boger, D.L. *et al.*, *J.A.C.S.*, 2000, **122**, 12169-12173 (*synth*)

Rubrolone B

R-151
Antibiotic Ro 21-4958. Ro 21-4958
 $C_{23}H_{23}NO_8$ 441.437
 Struct. closely related to Rubrolone, R-150. Prod. by *Streptomyces echinoruber*. Red powder. Sol. H_2O , MeOH; fairly sol. EtOH; poorly sol. butanol, hexane. λ_{max} 218 (ϵ 18000); 273 (ϵ 4300); 333 (ϵ 8600); 412 (ϵ 6700); 520 (ϵ 5200) (H_2O).
 Palleroni, N.J. *et al.*, *J. Antibiot.*, 1978, **31**, 1218-1225
 Schuep, W. *et al.*, *J. Antibiot.*, 1978, **31**, 1226-1232

Rubropunctamine

R-152
9a-Methyl-3-(1-oxohexyl)-6-(1-propenyl)furo[3,2-g]isoquinoline-2,9(7H,9aH)-dione
 [514-66-9]



Absolute Configuration

$C_{21}H_{23}NO_4$ 353.417
 Homologue of Monascorubramine, M-686. Prod. by *Monascus anka* and *Monascus purpureus*. Mycotoxin. Purple needles (EtOH). Mp 217-218° dec. λ_{max} 252 (log ϵ 4.16); 305 (log ϵ 4.43); 382 (log ϵ 3.93); 426 (log ϵ 4.17); 542 (log ϵ 4.24) (no solvent reported).

N-(*IR-Carboxyethyl*): [186649-33-2]
 $C_{24}H_{27}NO_6$ 425.48
 Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20}$ -2800 (c, 0.003 in MeOH). λ_{max} 202 ; 248 ; 292 ; 420 ; 500 (MeOH).

N-(*IS-Carboxyethyl*): [186649-32-1]
 $C_{24}H_{27}NO_6$ 425.48
 Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20}$ -2500 (c, 0.002 in MeOH). λ_{max} 202 ; 248 ; 292 ; 420 ; 498 (MeOH).

N-(*IR,2-Dicarboxyethyl*): [186649-37-6]
 $C_{25}H_{27}NO_8$ 469.49
 Prod. by *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20}$ -1400 (c, 0.002 in MeOH).

N-(*IS,2-Dicarboxyethyl*): [186649-36-5]
 $C_{25}H_{27}NO_8$ 469.49
 Prod. by a *Monascus* sp. Amorph. red solid. $[\alpha]_D^{20}$ -2200 (c, 0.003 in MeOH). λ_{max} 204 ; 248 ; 294 ; 418 ; 500 (MeOH).

Haws, E.J. *et al.*, *J.C.S.*, 1959, 3598-3610 (*synth, struct*)
 Hadfield, J.R. *et al.*, *J.C.S.(C)*, 1967, 751-755 (*isol*)
 Sweeny, J. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 1189-1193 (*isol, pmr*)
 Sato, K. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 227-229 (*N-carboxyethyl derivis*)
 Akihisa, T. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 562-565 (*isol, pmr*)

Rubroside G

R-153
 [227597-42-4]

 $C_{33}H_{38}Cl_2N_2O_9$ 677.577
 Similar to Aurantoside A, A-1550. Isol. from the marine sponge *Siliquariaspongia japonica*. Antifungal agent. Amorph. red solid. $[\alpha]_D^{24}$ -1212 (c, 0.001 in MeOH). λ_{max} 248 (ϵ 32400); 435 (ϵ 154400); 460 (ϵ 175500) (MeOH).
 2'-O-[5-Deoxy- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside B**
 [227597-37-7]
 $C_{43}H_{54}Cl_2N_2O_{16}$ 925.809
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -1460 (c, 0.001 in MeOH). λ_{max} 250 (ϵ 18900); 432 (ϵ 71300); 457 (ϵ 69700) (MeOH).
 2'-O-[5-Deoxy-2-O-methyl- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside D**
 [227597-39-9]
 $C_{44}H_{56}Cl_2N_2O_{16}$ 939.836
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -1024 (c, 0.001 in MeOH). λ_{max} 251 (ϵ 16400); 432 (ϵ 109100) (MeOH).
 Sata, N.U. *et al.*, *J.O.C.*, 1999, **64**, 2331-2339 (*Rubrosides B,D,G*)

Rubroside H

R-154
 [227597-43-5]

 $C_{31}H_{35}Cl_3N_2O_9$ 685.984
 Similar to Aurantoside A, A-1550. Isol. from the marine sponge *Siliquariaspongia japonica*. Antifungal agent. Amorph. red solid. $[\alpha]_D^{24}$ -1308 (c, 0.001 in MeOH). λ_{max} 245 (ϵ 13700); 440 (ϵ 44700) (MeOH).
 2'-O-[5-Deoxy- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside E**
 [227597-40-2]
 $C_{41}H_{51}Cl_3N_2O_{16}$ 934.216
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -1424 (c, 0.001 in MeOH). λ_{max} 243 (ϵ 17000); 450 (ϵ 61700) (MeOH).

2'-O-[5-Deoxy-2-O-methyl- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside F**
 [227597-41-3]
 $C_{42}H_{53}Cl_3N_2O_{16}$ 948.243
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -490 (c, 0.001 in MeOH). λ_{max} 250 (ϵ 9100); 425 (ϵ 80000) (MeOH).
 21-Dechloro, 2'-O-[5-deoxy-2-O-methyl- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside C**
 [227597-38-8]
 $C_{42}H_{54}Cl_2N_2O_{16}$ 913.798
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -1088 (c, 0.001 in MeOH). λ_{max} 240 (ϵ 18600); 450 (ϵ 60600) (MeOH).
 21-Dechloro, 20*Z*-isomer, 22-epimer, 2'-O-[5-deoxy-2-O-methyl- β -D-arabinofuranosyl-(1 \rightarrow 4)- β -D-xylopyranosyl]: **Rubroside A**
 [227597-36-6]
 $C_{42}H_{54}Cl_2N_2O_{16}$ 913.798
 Isol. from *Siliquariaspongia japonica*. Amorph. red solid. $[\alpha]_D^{24}$ -1824 (c, 0.001 in MeOH). λ_{max} 246 (ϵ 29100); 418 (ϵ 110400); 430 (ϵ 110200) (MeOH).
 Sata, N.U. *et al.*, *J.O.C.*, 1999, **64**, 2331-2339 (*Rubrosides A,C,E,F,H*)

Rugosanine B

R-155
 [123519-68-6]

 $C_{36}H_{39}N_5O_5$ 621.735
 Alkaloid from the stem bark of *Zizyphus rugosa* (Rhamnaceae). Shows antibacterial and antifungal props. Granules (MeOH). Mp 216-218° dec.
 Tripathi, Y.C. *et al.*, *Phytochemistry*, 1989, **28**, 1563 (*isol, uv, ir, ms, struct*)

Rugosinone


R-156
Linaresine
 [73609-04-8]

 $C_{19}H_{15}NO_6$ 353.331

Linaresine and Dihydrolinaresine originally assigned α -hydroxycularine structs. Identity with Rugosinone and Dihydrosugosinone established in 1996. Minor alkaloid from the roots of *Thalictrum rugosum* (Ranunculaceae). Also isol. (as Linaresine) from *Berberis valdiviana* (Berberidaceae). Yellow needles (EtOAc). Mp 223-224° (220-221°).

3,4-Dihydro: Dihydrosugosinone. Dihydrolinaresine
[74909-04-9]
C₁₉H₁₇NO₆ 355.346
Alkaloid from stems of *Berberis darwinii* and twigs of *Berberis actinocantha* (Berberidaceae). Also isol. (as Dihydrolinaresine) from *Berberis valdiviana*. Yellow amorph. solid (MeOH). Mp 172-174° Mp 220-223°.
[92219-94-8, 92240-99-8]

Murugesan, N. et al., *Heterocycles*, 1980, **14**, 585 (synth)
Wu, W.-N. et al., *J. Nat. Prod.*, 1980, **43**, 143 (isol, uv, ir, pmr, ms, struct)
Cheng, H.-Y. et al., *J. Nat. Prod.*, 1980, **43**, 151 (synth)
Valencia, E. et al., *J. Nat. Prod.*, 1984, **47**, 1050 (Dihydrosugosinone)
Firdous, S. et al., *J.A.C.S.*, 1984, **106**, 6099 (isol, Linaresine, Dihydrolinaresine)
Garcia, A. et al., *Tetrahedron*, 1996, **52**, 5929 (struct)

Rugulovasine A**R-157**C₁₆H₁₆N₂O₂ 268.315

(+)-form [91465-62-2]
Synthetic. Mp 120°. [α]_D²⁵ +43 (c, 0.5 in MeOH).

Hydrochloride: Mp 224°. [α]_D²⁰ -142 (c, 1 in MeOH). [α]_D +30 (c, 0.5 in DMSO).

(±)-form [26909-33-1]

Alkaloid from *Penicillium concavo-rugulosum* and *Penicillium islandicum*. Shows hypotensive and bradycardiac props. Mycotoxic. Prisms or needles (C₆H₆, CHCl₃ or MeCN). Sol. MeOH, EtOAc; fairly sol. CHCl₃, C₆H₆; poorly sol. H₂O, hexane. Mp 138° dec. λ_{\max} 224 (€ 30200); 288 (€ 6600); 295 (€ 6300) (EtOH) (Berdy).

► LD₅₀ (mus, ivn) 80 mg/kg.

Hydrochloride:
Prisms (H₂O). Mp 225° dec.

Picrate:
Yellow prisms (MeOH). Mp 229° dec.

N-De-Me: N-Demethylrugulovasine A. RUG 13A. Antibiotic RUG 13A
[176901-08-9]
[177019-39-5]

C₁₅H₁₄N₂O₂ 254.288

Prod. by *Saccharothrix* sp. Rug-13.

Substance P antagonist. Analgesic. Powder (as hydrochloride). Sol. MeOH; poorly sol. CHCl₃. λ_{\max} 220 (€ 40600); 284 (€ 7600); 293 (€ 7600) (MeOH) (hydrochloride). λ_{\max} 220 (€ 40600); 284 (€ 7600); 293 (€ 7600) (MeOH) (Berdy).

8-Chloro: 8-Chlororugulovasine A. Toxin B
[59787-45-0]
C₁₆H₁₅ClN₂O₂ 302.759
Alkaloid from cultures of *Penicillium islandicum*. Noncryst. Isol. as the racemate, poss. because of racemisation during isol. Interconverts with 8-chlororugulovasine B below in hot MeOH.

► WG9667000

5-Epimer: Rugulovasine B
[26909-34-2]

Alkaloid from cultures of *Penicillium concavo-rugulosum* and *Penicillium islandicum*. Also prod. by *Lenzites trabea*, *Pellicularia filamentosa*, *Corticium caeruleum*, *Penicillium Coryophyloides*. Hypotensive agent, mycotoxin. Amorph. solid; prisms (as hydrochloride). Sol. MeOH, EtOAc, bases, Py; fairly sol. H₂O, CHCl₃, Et₂O, C₆H₆; poorly sol. hexane. Mp 187° dec. (as hydrochloride). Rugulovasines A and B are readily interconverted. λ_{\max} 227 (€ 14450); 228 (€ 4790); 288 (€ 5370); 295 (€ 5250) (EtOH) (Berdy).

5-Epimer, N-de-Me: N-Demethylrugulovasine B. RUG 13B. Antibiotic RUG 13B

[177019-41-9]
[177185-32-9]
C₁₅H₁₄N₂O₂ 254.288

Prod. by *Saccharothrix* sp. Rug-13. Substance P antagonist. Analgesic. Powder (as hydrochloride). Sol. MeOH; poorly sol. CHCl₃. λ_{\max} 220 (€ 39500); 284 (€ 6800); 293 (€ 6600) (MeOH) (hydrochloride). λ_{\max} 220 (€ 39500); 284 (€ 6800); 293 (€ 6600) (MeOH) (Berdy).

5-Epimer, 8-chloro: 8-Chlororugulovasine B. Toxin A

[59787-46-1]
C₁₆H₁₅ClN₂O₂ 302.759

Alkaloid from cultures of *Penicillium islandicum*. Noncryst.

Abe, M. et al., *Agric. Biol. Chem.*, 1969, **33**, 469 (isol, uv, ir)

Yamatodani, S. et al., *Agric. Biol. Chem.*, 1970, **34**, 485 (uv, pmr, struct)

Nagaoka, A. et al., *Arzneim.-Forsch.*, 1972, **22**, 137; 143 (props)

Cole, R.J. et al., *Can. J. Microbiol.*, 1976, **22**, 741 (8-Chlororugulovasines)

Cole, R.J. et al., *Tet. Lett.*, 1976, 3849 (cryst struct)

Ohmomo, S. et al., *Agric. Biol. Chem.*, 1977, **41**, 1707 (biosynth)

Dorner, J.W. et al., *Appl. Environ. Microbiol.*, 1980, **40**, 685 (isol)

Rebek, J. et al., *J.O.C.*, 1984, **49**, 3540 (synth, conformn)

Martin, S.F. et al., *J.A.C.S.*, 1993, **115**, 10450 (synth)

Japan. Pat., 1996, 96 81470; *CA*, **125**, 8685x (N-demethylrugulovasines)

Liras, S. et al., *J.A.C.S.*, 2001, **123**, 5918-5924 (synth)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 537

Rulodine**R-158**C₁₇H₂₁NO₄ 303.357

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Narcissus odoros* var. *rugulosus* (Amaryllidaceae). Cryst. (Me₂CO). Mp 193°. [α]_D²⁵ -15 (c, 0.2 in CHCl₃).


Hydroiodide:

Cryst. (Me₂CO). Mp 135-136° dec.

Boit, H.-G. et al., *Naturwissenschaften*, 1961, **48**, 603 (isol)

Rumberine**R-159**

[73354-04-8]

C₂₁H₂₄N₂O₅ 384.431

Alkaloid from the aerial parts of *Hamelia patens* (Rubiaceae). Cryst. (cyclohexane). Mp 188-189°. [α]_D²⁰ -48. λ_{\max} 227 (€ 12600); 255 (€ 18100); 312 (€ 2900) (EtOH).

Me ether: Palmirine

[73367-02-9]

C₂₂H₂₆N₂O₅ 398.458


Alkaloid from the aerial parts of *Hamelia patens* (Rubiaceae). Cryst. (MeOH aq.). Mp 101-102°. [α]_D²⁰ -56.5. λ_{\max} 255 (€ 24900); 312 (€ 2990) (EtOH).

Borges, J. et al., *Tet. Lett.*, 1979, **20**, 3197-3200 (*Rumberine, Palmirine, uv, ir, pmr, ms, struct*)

Rumbrin**R-160**

N 13. Antibiotic N 13

[150206-14-7]

C₂₀H₂₀ClNO₃ 357.836

Related to Auxarconjugatin A, A-1559. Isol. from the fungi *Auxarthron umbrium* and *Gymnoascus reessii*. Cytoprotective agent. Calcium accumulation inhibitor and active oxygen remover. Lipid peroxidation inhibitor. Red needles (MeOH). Sol. MeOH, EtOAc, CHCl₃, DMSO; poorly sol. H₂O, hexane. Mp 170-171° dec. λ_{\max} 269 (€ 12800); 341 (€ 17400); 442 (€ 33800) (MeOH).

12E-Isomer: 12E-Rumbrin. 12E-Isorumbrin

C₂₀H₂₀ClNO₃ 357.836

Isol. from *Gymnoascus reessii* strain MST-F9977. Cytotoxic. Red solid. λ_{max} 225 (ϵ 7700); 269 (ϵ 9900); 330 (ϵ 10600); 340 (ϵ 12000); 439 (ϵ 35000) (MeOH).


[148528-19-2]

Matsuoka, M. *et al.*, *J. Antibiot.*, 1993, **46**, 884-887; 888-891 (*isol, pmr, cmr, struct, props*)

Clark, B.R. *et al.*, *Org. Lett.*, 2006, **8**, 701-704 (*12E-Rumbrin*)

Runanine

[100485-12-9]



$\text{C}_{21}\text{H}_{27}\text{NO}_5$ 373.448

Alkaloid from the roots of *Stephania sinica*. Mp 100-102°. $[\alpha]_{\text{D}}^{18}$ -400 (c, 0.8 in CHCl_3).

N-De-Me: 7,8-Didehydro-2,3,7,8-tetramethoxyhasubanan-6-one. **Stephanubine**

[111509-16-1]

$\text{C}_{20}\text{H}_{25}\text{NO}_5$ 359.421

Alkaloid from the tuberous roots of *Stephania suberosa*. Amorph. $[\alpha]_{\text{D}}^{25}$ -170 (c, 0.22 in CHCl_3).

O²-De-Me: **Cephatonine**

[175862-78-9]

$\text{C}_{20}\text{H}_{25}\text{NO}_5$ 359.421

Alkaloid from tubers of *Stephania cepharantha*. Amorph. powder. $[\alpha]_{\text{D}}^{28}$ -264 (c, 0.13 in CHCl_3).

8-Demethoxy: **8-Demethoxyrunanine**

[1017277-31-4]

$\text{C}_{20}\text{H}_{25}\text{NO}_4$ 343.422

Alkaloid from the rhizomes of *Sinomenium acutum*. Mp 169.5-170.5°. $[\alpha]_{\text{D}}^{20}$ -244.1 (c, 0.48 in CHCl_3).

Zhi-Da, M. *et al.*, *Phytochemistry*, 1985, **24**, 3084-3085 (*Runanine*)


Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 2391-2395 (*Stephanubine*)

Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 1996, **59**, 476-480 (*Cephatonine*)

Wang, X. *et al.*, *Fitoterapia*, 2007, **78**, 593-595 (*8-Demethoxyrunanine*)

Rupancamine

[96203-73-5]



R-162

$\text{C}_{37}\text{H}_{40}\text{N}_2\text{O}_6$ 608.733


Alkaloid from *Berberis actinacantha* (Berberidaceae). Amorph. $[\alpha]_{\text{D}}^{25}$ +116.9 (c, 0.12 in CHCl_3).

Weiss, I. *et al.*, *Heterocycles*, 1984, **22**, 2231 (*uv, ir, pmr, ms, struct*)

Ruprechstylil

R-163

6-Methoxy-3-pentyl-1,8-isoquinolinediol. 2,8-Dihydroxy-6-methoxy-2-pentylisoquinoline. 8-Hydroxy-6-methoxy-3-pentyl-1(2H)-isoquinolinone. 8-Hydroxy-6-methoxy-3-pentylisocarbostylil



$\text{C}_{15}\text{H}_{19}\text{NO}_3$ 261.32


Tautomeric with pyridone form. Alkaloid from the stem bark and twigs of *Ruprechtia tangarana*. Needles (Me₂CO). Mp 139-141°. λ_{max} 244 (log ϵ 4.7) (MeOH).

Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1065-1069 (*isol, pmr, cmr, cryst struct*)

Rurrebanidine

R-164

[112494-67-4]



$\text{C}_{18}\text{H}_{19}\text{NO}_4$ 313.352

Alkaloid from the bark of Bolivian *Duguetia spixiana* (Annonaceae). Negative opt. rotn. in EtOH.

3-Me ether: **Rurrebanine**

[112494-68-5]

$\text{C}_{19}\text{H}_{21}\text{NO}_4$ 327.379


Alkaloid from *Duguetia spixiana* (Annonaceae). $[\alpha]_{\text{D}}$ -43 (c, 0.72 in EtOH).

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1987, **50**, 674 (*isol, uv, pmr, ms, cd, struct*)

Ruspolinone

R-165

1-(3,4-Dimethoxyphenyl)-2-(2-pyrrolidinyl)ethanone, 9CI. 2-(3,4-Dimethoxyphenacyl)pyrrolidine. 2-(2-Pyrrolidinyl)-3',4'-dimethoxyacetophenone. 3',4'-Dimethoxy-2-pyrrolidin-2-yl-acetophenone



$\text{C}_{14}\text{H}_{19}\text{NO}_3$ 249.309

(±)-form [60890-27-9]

Alkaloid from seeds of *Ruspolia hypercrateriformis* (Acanthaceae). Intermed. in

Tylophorinine biosynth. in *Tylophora asthmatica* (Asclepiadaceae). Oil. Unstable, intractable.

O⁴-De-Me: **Norruspolinone**. 1-(4-Hydroxy-3-methoxyphenyl)-2-(2-pyrrolidinyl)ethanone, 9CI. 2-(2-Pyrrolidinyl)-4'-hydroxy-3'-methoxyacetophenone [67257-66-3]

$\text{C}_{13}\text{H}_{17}\text{NO}_3$ 235.282

Alkaloid from seeds of *Ruspolia hypercrateriformis* (Acanthaceae). Mp 175° dec.

O⁴-De-Me, N-Me: **Phyllosterone**

[126262-24-6]

$\text{C}_{14}\text{H}_{19}\text{NO}_3$ 249.309

Alkaloid from *Cryptocarya phyllostemon* (Lauraceae). Gum. $[\alpha]_{\text{D}}$ -5 (c, 0.69 in EtOH). Apparently not completely racemic. No information on opt. purity or abs. config.

(S)-form

Synthetic. Pale yellow powder (CH_2Cl_2 /petrol). Mp 114-115°. $[\alpha]_{\text{D}}$ -29.73 (c, 0.74 in CH_2Cl_2).

[62024-33-3]

Roessler, F. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 1200 (*isol, uv, ir, pmr, ms, struct, deriv*)

Mangla, V.K. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 748 (*synth*)

Cragg, J.E. *et al.*, *J.C.S. Perkin I*, 1982, 2477 (*synth*)

Herbert, R.B. *et al.*, *J.C.S. Perkin I*, 1984, 825

Cavé, A. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2243 (*Phyllosterone*)

Brown, D.S. *et al.*, *Tetrahedron*, 1991, **47**, 1311 (*synth*)


Jones, K. *et al.*, *Tetrahedron*, 1991, **47**, 7179 (*synth*)

Courture, A. *et al.*, *Tet. Lett.*, 1996, **37**, 7749 (*synth*)

Russupteridine yellow IV

R-166

1-Deoxy-1-(1,2,3,6,7,8-hexahydro-2,6,8-trioxo-4H-imidazo[4,5-g]pteridin-4-yl)-D-ribitol, 9CI [70858-25-2]



$\text{C}_{12}\text{H}_{14}\text{N}_6\text{O}_7$ 354.279

Isol. from *Russula sardonia* (Basidiomycetes). Pale yellow powder. Mp 300° dec. pK_{a} 2.97.

2-Alcohol: **Russupteridine yellow I**

[70858-26-3]


$\text{C}_{12}\text{H}_{16}\text{N}_6\text{O}_7$ 356.294

Isol. from *Russula* spp. Yellowish-brown needles (AcOH aq.). Mp 300°. $\text{pK}_{\text{a}1}$ 0.25; $\text{pK}_{\text{a}2}$ 4.8; $\text{pK}_{\text{a}3}$ 12.1. pH > 5 is in equilibrium with an open formylimino form.

Iten, P.X. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 550 (isol, pmr, cmr, uv, ir, ms)

Rutacridone R-167

1,11-Dihydro-5-hydroxy-11-methyl-2-(1-methylethenyl)furo[2,3-c]acridin-6(2H)-one
[17948-33-3]



C₁₉H₁₇NO₃ 307.348

Natural Rutacridone has recently (2005) been shown to have the *R*-config. illus. Related alkaloids included in this entry may not necessarily be configurationally related. Alkaloid from *Ruta graveolens* (rue) and *Ruta chalepensis* (Rutaceae). Mp 161-162°. [α]_D²⁵ -44.9 (c, 0.1 in MeOH). λ_{max} 212 (log ε 4.35); 227 (log ε 4.3); 249 (log ε 4.56); 273 (log ε 4.72); 300 (log ε 4.35); 332 (log ε 4); 398 (log ε 3.83) (MeOH).

▶ VM2050000

1'*R*,2'-Epoxide: **Rutacridone epoxide**

[77996-03-3]
C₁₉H₁₇NO₄ 323.348

Alkaloid from roots and callus tissue cultures of *Ruta graveolens* (rue). Shows antimicrobial activity. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Abs. config. determined in 2005. λ_{max} 226 (ε 3710); 263 (ε 6760); 300 (ε 3710); 390 (ε 1200) (MeOH) (Berdy).

▶ LV0850000

3'-Chloro: **Alkaloid A6**

[142741-38-6]
C₁₉H₁₆ClNO₃ 341.793

Alkaloid from callus cultures of several *Ruta* spp.

1ξ-Hydroxy, 1'ξ,2'-epoxide: **1-Hydroxyrutacridone epoxide**

[101330-59-0]
C₁₉H₁₇NO₅ 339.347

Alkaloid from the callus tissue of *Ruta graveolens* (rue). Shows antimicrobial activity.

3'-Hydroxy: **Gravacridonol**

[81545-69-9]
C₁₉H₁₇NO₄ 323.348

Alkaloid from *Ruta graveolens* (rue). Mp 155-156° (153°). λ_{max} 230 (log ε 4.28); 248 (log ε 4.33); 280 (log ε 4.37); 338 (log ε 3.9); 424 (log ε 3.67) (MeOH).

3'-Hydroxy, O-β-D-glucopyranoside: **Gravacridonol monoglucoside**

[155861-84-0]
C₂₅H₂₇NO₉ 485.49

Alkaloid from cell suspension cultures of *Thamnosma montana*.

3'-Hydroxy, 1',2'-epoxide: **20-Hydroxyrutacridone epoxide**

[80565-10-2]

C₁₉H₁₇NO₅ 339.347

Alkaloid from the roots and callus tissue cultures of *Ruta graveolens* (rue). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. 20-Posn. in the authors' numbering scheme called here 3'. λ_{max} 227 (ε 8910); 250 (ε 9770); 264 (ε 10470); 274 (ε 13800); 301 (ε 5370); 330 (ε 2450); 392 (ε 1740) (MeOH) (Berdy).

1',2'-Dihydro, 1'-chloro, 2'-hydroxy: **Gravacridonechlorine**

[38494-84-7]
C₁₉H₁₈ClNO₄ 359.808

Alkaloid from *Ruta graveolens* (rue). Mp 254-257°.

1',2'-Dihydro, 2'-chloro, 1'-hydroxy: **Iso-gravacridonechlorine**

[62512-94-1]
C₁₉H₁₈ClNO₄ 359.808

Alkaloid from *Ruta graveolens* (rue). Shows weak antibacterial activity. Yellow powder (Me₂CO), Mp 224-226° dec Mp 247-250°. [α]_D²⁵ -9.2 (c, 0.08 in CHCl₃).

▶ Mutagen.

1',2'-Dihydro, 1',2'-dihydroxy: **Gravacridonediol**

[37551-75-0]
C₁₉H₁₉NO₅ 341.363

Alkaloid from the root tissue cultures of *Ruta graveolens* (rue). Cryst. (Me₂CO or Me₂CO/C₆H₆). Mp 224-227° dec Mp 227-233° Mp 246-250°. [α]_D²⁵ -111 (c, 0.1 in MeOH). λ_{max} 214 (ε 14517); 227 (ε 14517); 249 (ε 26127); 273 (ε 3731); 300 (ε 15690); 331 (ε 7310); 400 (ε 4845) (MeOH).

1',2'-Dihydro, 1',2'-dihydroxy, 2'-O-β-D-glucopyranoside: **Gravacridonediol glucoside**

[59086-97-4]
C₂₅H₂₉NO₁₀ 503.505

Alkaloid from *Ruta graveolens* (rue). Isol. as a mixt. with Gravacridonetriol glucoside.

1',2'-Dihydro, 1',2'-dihydroxy, 2'-Ac:

Gravacridonediol acetate
[136591-55-4]

C₂₁H₂₁NO₆ 383.4

Alkaloid from *Ruta graveolens* (rue). Yellow needles (CHCl₃). Mp 221-223°. [α]_D²⁵ +59.8 (c, 0.4 in CHCl₃).

1',2'-Dihydro, 1'-hydroxy, 2'-methoxy:

Gravacridonediol methyl ether
[37551-76-1]

C₂₀H₂₁NO₅ 355.39

Alkaloid from the root of *Ruta graveolens* (rue). Cryst. (EtOAc). Mp 219-221°.

1',2'-Dihydro, 1',3'-dihydroxy, 2'-chloro:

2-[1-Chloro-2-hydroxy-1-(hydroxymethyl)ethyl]-1,11-dihydro-5-hydroxy-11-methylfuro[2,3-c]acridin-6(2H)-one. Gravacridonolchlorine
[38494-85-8]

C₁₉H₁₈ClNO₅ 375.808

Alkaloid from *Ruta graveolens* (rue). Mp 223-227°.

1',2'-Dihydro, 1',2',3'-trihydroxy: **Gravacridonetriol**

[59086-94-1]
C₁₉H₁₉NO₆ 357.362

Alkaloid from *Ruta graveolens* (rue). Cryst. (MeOH). Mp 230-232°.

1',2'-Dihydro, 1',2',3'-trihydroxy, 2'-O-β-D-glucopyranoside: **Gravacridonetriol glucoside**

[59086-96-3]
C₂₅H₂₉NO₁₁ 519.504

Alkaloid from *Ruta graveolens* (rue). Isol. as a mixt. with Gravacridonediol glucoside.

10-Hydroxy, N-de-Me: **Oriaciadone C**

C₁₈H₁₅NO₄ 309.321

Alkaloid from the stem bark of *Oriaciopsis glaberrima*. α-Glucosidase inhibitor. Yellow needles (MeOH). Mp 253-254°. [α]_D²⁵ +21.8 (c, 0.39 in MeOH). Possesses *R*-config. λ_{max} 253 (log ε 6.09); 277 (log ε 5.76); 279 (log ε 5.77); 303 (log ε 5.66); 346 (log ε 4.76) (MeOH).

Reisch, J. *et al.*, *Phytochemistry*, 1972, **11**,

2121-2122; 2359; 1976, **15**, 240-241; 1977,

16, 151 (*Gravacridonediol derivs.*,

Gravacridonechlorine, *Gravacridonolchlorine*,

Gravacridonetriol, *Isogravacridonechlorine*)

Rózsa, Zs. *et al.*, *Fitoterapia*, 1976, **47**, 147-149

(*Gravacridonol*, *isol*)

Reisch, J. *et al.*, *Z. Naturforsch.*, **B**, 1978, **33**,

957 (*rev. pmr, struct*)

Bergenthal, D. *et al.*, *Phytochemistry*, 1979, **18**,

161-163 (*Gravacridonediol*, *cmr*)

Wolters, B. *et al.*, *Planta Med.*, 1981, **43**, 166-

174 (*epoxides*, *isol*, *activity*)

Nahrstedt, A. *et al.*, *Z. Naturforsch.*, **C**, 1981,

36, 200-203 (*Rutacridone epoxide*)

Zschunke, A. *et al.*, *Chem. Comm.*, 1982, 1263-

1265 (*biosynth*)

Eilert, U. *et al.*, *Z. Naturforsch.*, **C**, 1982, **37**,

132-133 (*20-Hydroxyrutacridone epoxide*)

Nahrstedt, A. *et al.*, *Planta Med.*, 1985, 517-

518 (*1-Hydroxyrutacridone epoxide*)

Reisch, J. *et al.*, *Monatsh. Chem.*, 1990, **121**,

829-835 (*synth*)

Reisch, J. *et al.*, *Annalen*, 1991, 299

(*Gravacridonol*, *synth*, *ir*, *uv*, *pmr*, *cmr*)

Paulini, H. *et al.*, *Planta Med.*, 1991, **57**, 59;

82-83 (*Isogravacridonechlorine*,

Gravacridonediolacetate)

Baumert, A. *et al.*, *Plant Cell, Tissue Organ*

Cult., 1992, **28**, 159-162; *C.A.*, **117**, 66678t

(*Alkaloid A6*)

Reisch, J. *et al.*, *J. Chem. Res., Synop.*, 1994,

157; *J. Chem. Res., Miniprint*, 1994, 911-927

(*Gravacridonediol*, *synth*, *ir*, *pmr*, *cmr*)

Baumert, A. *et al.*, *Planta Med.*, 1994, **60**, 143-

145 (*Gravacridonol monoglucoside*)

El Sayed, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 995-

997 (*Isogravacridonechlorine*, *activity*)

Meepagala, K.M. *et al.*, *Phytochemistry*, 2005,

66, 2689-2695 (*cd*, *abs config*)

Wansi, J.D. *et al.*, *Chem. Pharm. Bull.*, 2006,


54, 292-296 (*Oriaciadone C*)

Rutaecarpine**R-168**

8,13-Dihydroindolo[2',3':3,4]pyrido[2,1-b]quinazolin-5(7H)-one, 9CI. *Rhetine*.

Rutecarpine

[84-26-4]



$C_{18}H_{13}N_3O$ 287.32

A major component of the Chinese drug Wou-chou-yu (the dried fruit of *Evodia rutaecarpa*). Also isol. from *Hortia arborea*, *Hortia badinii*, *Zanthoxylum rhetsa*, *Zanthoxylum ovalifolium* and *Zanthoxylum integrifolium*. Antihypertensive agent., vasodilator. *Evodia rutaecarpa* is used in Chinese medicine against headache, abdominal pain, dysentery and cholera. Vasodilator. Cryst. (EtOAc, C_6H_6 , EtOH or AcOH). Mp 259.5-260°. Log P 2.57 (calc). λ_{max} 278; 290; 332; 345; 364 (MeOH) (Berdy).

7,8-Didehydro: 7,8-Dehydrorutaecarpine
[55786-24-8]

$C_{18}H_{11}N_3O$ 285.304

Alkaloid from *Phellodendron amurense*. Pale yellow powder. λ_{max} 217 (log ϵ 4.52); 250 (log ϵ 4.49); 281 (log ϵ 4.31); 330 (log ϵ 4.42); 348 (log ϵ 4.38); 368 (log ϵ 4.43); 390 (log ϵ 4.42) (MeOH).

1-Hydroxy: 1-Hydroxyrutaecarpine
[53600-24-1]

$C_{18}H_{13}N_3O_2$ 303.32

Minor alkaloid from the bark of *Euxylophora paraensis* and the stem bark of *Vepris louisii* (Rutaceae). Pale yellow needles (CHCl₃/MeOH). Mp 318-320°.

1-Hydroxy, hydrochloride:

Yellow needles (MeOH/CHCl₃). Mp 326-328°.

1-Hydroxy, 7,8-didehydro: 7,8-Dehydro-1-hydroxyrutaecarpine
[85769-53-5]

$C_{18}H_{11}N_3O_2$ 301.304

Minor alkaloid from the stem bark of *Vepris louisii* (Rutaceae). Yellow powder. Mp 340°.

3-Hydroxy: 3-Hydroxyrutaecarpine
[381723-48-4]

$C_{18}H_{13}N_3O_2$ 303.32

Alkaloid from *Leptothrysa sprucei*. Needles (MeOH). Mp 280-282°. λ_{max} 220 (log ϵ 4.62); 230 (log ϵ 4.61); 340 (log ϵ 4.66) (MeOH).

7 β -Hydroxy: 7-Hydroxyrutaecarpine
[163815-35-8]

$C_{18}H_{13}N_3O_2$ 303.32

Alkaloid from heartwood of *Tetradium glabrifolium* (*Evodia meliaefolia*) and fruit of *Tetradium ruticarpum* (Rutaceae). Needles. Mp 228-230°. [α]_D +187.8 (c, 0.0017 in MeOH).

11-Hydroxy: 11-Hydroxyrutaecarpine

11-Hydroxy, O- β -D-glucopyranoside: Ternatoside C

$C_{24}H_{23}N_3O_7$ 465.462

Alkaloid from the roots of *Ranunculus ternatus*. Brown powder (MeOH). Mp 185-187°. λ_{max} 222 (log ϵ 4.61); 353 (log ϵ 4.86) (MeOH).

11-Hydroxy, O-[α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside]: Ternatoside D

$C_{30}H_{33}N_3O_{11}$ 611.604

Alkaloid from the roots of *Ranunculus ternatus*. Brown powder (MeOH). Mp 172-175°. λ_{max} 220 (log ϵ 4.51); 351 (log ϵ 4.88) (MeOH).

7 β ,8 α -Dihydroxy: 7,8-Dihydroxyrutaecarpine
[212138-47-1]

$C_{18}H_{13}N_3O_3$ 319.319

Alkaloid from *Phellodendron amurense*. Needles (EtOAc/EtOH). Mp 247-250°. [α]_D²⁵ -38 (c, 0.0016 in MeOH). Incorrect name assigned in CA. λ_{max} 215 (log ϵ 4.77); 240 (sh) (log ϵ 4.63); 275 (log ϵ 4.19); 288 (log ϵ 4.16); 332 (log ϵ 4.64); 345 (log ϵ 4.7); 362 (log ϵ 4.6) (EtOH).

1,2-Dihydroxy: 1,2-Dihydroxyrutaecarpine
[622842-36-8]

$C_{18}H_{13}N_3O_3$ 319.319

Alkaloid from the aerial parts of *Bouchardatia neurococca*. Amorph. yellow solid. λ_{max} 256 (log ϵ 3.6); 343 (log ϵ 3.5); 360 (log ϵ 3.5); 377 (sh) (no solvent reported).

1-Methoxy: 1-Methoxyrutaecarpine
[53600-26-3]

$C_{19}H_{15}N_3O_2$ 317.346

Alkaloid from fruits of *Zanthoxylum integrifolium*. Shows antiplatelet aggregation activity. Yellowish prisms (MeOH). Mp 253-255°.

1-Methoxy, 7,8-didehydro: 7,8-Dehydro-1-methoxyrutaecarpine
[864179-28-2]

$C_{19}H_{13}N_3O_2$ 315.331

Alkaloid from the root bark of *Zanthoxylum integrifolium*. Yellowish needles (CHCl₃/MeOH). Mp 287-289°. λ_{max} 210 (log ϵ 4.64); 248 (sh) (log ϵ 4.48); 285 (log ϵ 4.27); 326 (log ϵ 4.42); 345 (log ϵ 4.39); 364 (log ϵ 4.46); 385 (log ϵ 4.42) (EtOH).

2-Methoxy: 2-Methoxyrutaecarpine
[53600-27-4]

$C_{19}H_{15}N_3O_2$ 317.346

Alkaloid from the stem bark of *Araliopsis tabouensis*. Amorph. yellow powder. λ_{max} 214 (log ϵ 1.95); 242 (log ϵ 1.11); 324 (log ϵ 0.7) (MeOH).

2-Methoxy, N¹³-Me: 2-Methoxy-N¹³-methylrutaecarpine
[53600-30-9]

$C_{20}H_{17}N_3O_2$ 331.373

Alkaloid from the stem bark of *Araliopsis tabouensis*. Amorph. yellow powder (EtOAc). Mp 164° (synthetic). λ_{max} 212 (log ϵ 1.8); 326 (log ϵ 0.9); 372 (log ϵ 1.52) (MeOH).

Asahina, Y. et al., *J.C.S.*, 1927, 1708-1710 (synth)

Pachter, I.J. et al., *J.A.C.S.*, 1960, **82**, 5187-5193 (isol)

Yamazaki, M. et al., *Tet. Lett.*, 1966, **7**, 3221-3224; 1967, **8**, 3317-3320 (biosynth)

Danieli, B. et al., *Phytochemistry*, 1974, **13**, 1603-1606 (isol, uv, ir, pmr, ms, struct, synth, 1-Hydroxyrutaecarpine)

Danieli, B. et al., *Gazz. Chim. Ital.*, 1975, **105**, 45-49 (2-Methoxy-N¹³-methylrutaecarpine, synth)

De Barros Corrêa, D. et al., *Phytochemistry*, 1975, **14**, 2059-2060 (isol)

Tamás, J. et al., *Acta Chim. Acad. Sci. Hung.*, 1976, **89**, 85-89 (ms)

Kametani, T. et al., *J.A.C.S.*, 1976, **98**, 6186-6188; 1977, **99**, 2306-2309 (synth, uv, ir, pmr, ms)

Toth, G. et al., *Annalen*, 1977, 529-536 (cmr)

Kametani, T. et al., *Chem. Pharm. Bull.*, 1978, **26**, 1922-1926 (synth)

Danieli, B. et al., *Heterocycles*, 1978, **9**, 803-806 (synth)

Kökösi, J. et al., *Tet. Lett.*, 1981, **22**, 4861-4862; 1992, **33**, 2995-2998 (synth)

Ayafor, J.F. et al., *Phytochemistry*, 1982, **21**, 2733-2736 (7,8-Dehydro-1-hydroxyrutaecarpine)

Kaneko, C. et al., *Heterocycles*, 1985, **23**, 1385-1390 (synth)

Bergman, J. et al., *J.O.C.*, 1985, **50**, 1246-1255 (synth)

Eguchi, S. et al., *Heterocycles*, 1992, **33**, 153-156 (synth)

Chiou, W.-F. et al., *Eur. J. Pharmacol.*, 1994, **257**, 59-66 (pharmacol)

Ko, H.-C. et al., *J. Chromatogr. B*, 1994, **655**, 27-31 (hplc)

Wu, T.-S. et al., *Heterocycles*, 1995, **41**, 1071-1076 (7-Hydroxyrutaecarpine)

Sheen, W.-S. et al., *Planta Med.*, 1996, **62**, 175-176 (1-Methoxyrutaecarpine)

Ikuta, A. et al., *J. Nat. Prod.*, 1998, **61**, 1012-1014 (7,8-Dihydroxyrutaecarpine, 7-Hydroxyrutaecarpine)

Ikuta, A. et al., *Phytochemistry*, 1998, **48**, 285-291 (pmr, cmr, 7,8-Dehydrorutaecarpine)

Sheu, J.-R. et al., *Cardiovasc. Drug Rev.*, 1999, **17**, 237-245 (rev)

Lee, S.L. et al., *Heterocycles*, 2001, **55**, 1555-1559 (synth)

Li, X.-C. et al., *Phytochemistry*, 2001, **58**, 627-629 (3-Hydroxyrutaecarpine)

Christopher, E. et al., *Helv. Chim. Acta*, 2003, **86**, 2914-2918 (2-Methoxyrutaecarpines)

Wattanapiromsakul, C. et al., *Phytochemistry*, 2003, **64**, 609-615 (1,2-Dihydroxyrutaecarpine)

Chavan, S.P. et al., *Tet. Lett.*, 2004, **45**, 997-999 (synth)

Harayama, T. et al., *Tetrahedron*, 2004, **60**, 10645-10649 (synth)

Chen, J.-J. et al., *Planta Med.*, 2005, **71**, 470-475 (7,8-Dehydro-1-methoxyrutaecarpine)

Hamid, A. et al., *Tet. Lett.*, 2006, **47**, 1777-1781 (synth)

Mhaske, S.B. et al., *Tetrahedron*, 2006, **62**, 9787-9826 (rev, synth)

Zhang, L. et al., *Chem. Pharm. Bull.*, 2007, **55**, 1267-1269 (11-Hydroxyrutaecarpine derivs)


Bowman, W.R. et al., *Org. Biomol. Chem.*, 2007, **5**, 103-113 (synth)

Lee, S.H. et al., *Molecules*, 2008, **13**, 272-300 (rev)

Rutagravine

[101330-60-3]

R-169



$C_{19}H_{17}NO_5$ 339.347


Alkaloid from the callus tissue of *Ruta graveolens* (rue) (Rutaceae). Cryst. (MeOH). Mp 209-212°. [α]_D²⁵ -152 (CHCl₃). λ_{max} 212 (ϵ 14414); 226 (ϵ 17057); 251 (ϵ 28602); 272 (ϵ 48289); 325 (ϵ 7207); 393 (ϵ 5700) (MeOH).

Nahrstedt, A. et al., *Planta Med.*, 1985, 517-519 (isol, pmr, cmr, struct)

Reisch, J. *et al.*, *J. Chem. Res., Synop.*, 1994, 157; *J. Chem. Res., Miniprint*, 1994, 911-927 (synth, ir, pmr, cmr, struct, abs config)

Rutalexin R-170

3-Methyl-1,3-thiazino[6,5-b]indole-2,4-(3H,9H)-dione




C₁₁H₈N₂O₂S 232.262

Alkaloid from the tubers of *Brassica napus* ssp. *rapifera* (swede, rutabaga). Phytoalexin. Solid. Mp 313-315°. λ_{max} 213 (log ε 4); 242 (log ε 3.7); 275 (log ε 3.6) (no solvent reported).

Pedras, M.S.C. *et al.*, *J.O.C.*, 2004, **69**, 4471-4476 (isol, synth, pmr, cmr, ms)

Rutalinium R-171

[27539-40-8]



C₁₆H₂₀NO₄[⊕] 290.338

Alkaloid from *Ruta graveolens* (rue), *Ruta graveolens* ssp. *hortensis* and *Balfourendendron riedelianum* (Rutaceae).

Szendrei, K. *et al.*, *Herba Hung.*, 1971, **10**, 131; *CA*, **79**, 15853k (isol, pmr, struct)

Varga, E. *et al.*, *Acta Pharm. Hung.*, 1978, **48**, 193; *CA*, **89**, 176400g (isol)

Ruticin R-172

[1393-17-5]

Phenazine antibiotic. Struct. unknown. Prod. by *Streptomyces rutgersensis*. Antibacterial agent. Orange cryst. Sol. MeOH, EtOAc; fairly sol. H₂O. λ_{max} 262; 364 (MeOH).

▶ LD₅₀ (mus, ivn) 20 mg/kg.

Fisher, W.P. *et al.*, *Antibiot. Annu.*, 1953, 174-176

Ruzorine R-173

C₁₈H₂₇NO₈ 385.413

Pyrrrolizidine alkaloid. Struct. unknown. Possibly identical with Bisline, B-194. Alkaloid from *Senecio ruwenzoriensis* (= *Senecio othonniformis*) (Asteraceae). Prisms or needles. V. sol. H₂O. Mp 161-163° (dec. from 134°).

Hydrochloride:

V. unstable cryst. Mp 148° dec. approx.

Picrate: Mp 133-137° (dec. from 125°).


Sapiro, M.L. *et al.*, *J.C.S.*, 1953, 1942-1943 (isol)

Benn, M. *et al.*, *Phytochemistry*, 1992, **31**, 3295-3296

Susag, L. *et al.*, *Phytochemistry*, 2000, **54**, 933-935

Ryania diterpene ester C₂ R-174

[99964-43-9]




C₂₅H₃₃NO₉ 491.537

Alkaloid from *Ryania speciosa* (Flacourtiaceae). Cryst. (CHCl₃). Mp 274-275°.

Ruest, L. *et al.*, *Can. J. Chem.*, 1985, **63**, 2840-2843 (isol, ir, pmr, struct)

Ryanodine R-175

Ryanex. Ryanicide. Ryno-tox [15662-33-6]



C₂₅H₃₅NO₉ 493.553

Alkaloid from *Ryania speciosa* (Flacourtiaceae). Calcium transport blocker. Insecticide, now superseded. Tan solid. Mp 219-220°. [α]_D²⁵ +26 (c, 1.02 in MeOH). Log P -1.22 (uncertain value) (calc). Hydrol. yields 2-Pyrrolicarboxylic acid and Ryanodol. For diterpene ester C₂ see Ryania diterpene ester C₂, R-174.

▶ Human systemic effects by ingestion (gastrointestinal and CNS). Harmful in contact with skin. LD₅₀ (rat, orl) 750 mg/kg. LD₅₀ (rat, skn) 750 mg/kg. VM4025000

3-O-(3-Pyridinecarbonyl) analogue: **Ryanodyl 3-nicotinate** [137441-82-8]

C₂₆H₃₅NO₉ 505.564

Alkaloid from *Ryania speciosa* (Flacourtiaceae). Prisms (Me₂CO/hexane). Mp 200-205°.

8,9-Didehydro: **8,9-Didehydoryanodine** [99964-45-1]

C₂₅H₃₃NO₉ 491.537

Constit. of *Ryania speciosa*.

9,21-Didehydro: **Dehydoryanodine**. 9,21-Didehydoryanodine [94513-55-0]

C₂₅H₃₃NO₉ 491.537

Second most abundant ryanodine-type alkaloid from *Ryania speciosa* (Flacourtiaceae). Mp 175-179°.

▶ VM4155000

4-Deoxy, 20-hydroxy: **Spiganthine**. 4-Deoxy-20-hydroxyryanodine

[168034-68-2]

C₂₅H₃₅NO₉ 493.553

Alkaloid from *Spigelia anthelmia*.

Cardiotonic agent. Cryst. (CHCl₃/Me₂CO). Mp 158°. [α]_D²³ +30 (c, 0.1 in MeOH). λ_{max} 227 (log ε 3.86); 267 (log ε 4.1) (MeOH).

8-Oxo, 10-deoxy, 9,10-didehydro: **Ryania diterpene ester B** [99964-44-0]

C₂₅H₃₁NO₉ 489.521

Alkaloid from *Ryania speciosa* (Flacourtiaceae).

3-Epimer: 3-Epiryanodine

[150736-85-9]

Cryst. (CH₂Cl₂/MeOH). Mp 228-230°.

10-Epimer, 8β-hydroxy, O⁴-Me: **Ryania diterpene ester A** [100082-08-4]

C₂₆H₃₇NO₁₀ 523.579

Alkaloid from *Ryania speciosa* (Flacourtiaceae).

10-Epimer, 8β-hydroxy, 19,21-didehydro, O⁴-Me: **Ryania diterpene ester D** [100082-09-5]

C₂₆H₃₅NO₁₀ 521.563

Alkaloid from *Ryania speciosa* (Flacourtiaceae).

10-Epimer, 8β-hydroxy, 4-deoxy, O¹⁰-Me: **4-Deoxyester A** [226923-09-7]

C₂₆H₃₇NO₉ 507.58

Constit. of *Ryania speciosa*. Amorph. solid.

10-Epimer, 9α-hydroxy: **Ryania diterpene ester C₁** [137844-84-9]

C₂₅H₃₅NO₁₀ 509.552

Alkaloid from *Ryania speciosa* (Flacourtiaceae). Cryst. (CHCl₃). Mp 188-190°.

[100427-92-7]

Wiesner, K. *et al.*, *Tet. Lett.*, 1967, 221-225 (struct)

Wiesner, K. *et al.*, *Adv. Org. Chem.*, 1972, **8**, 295-316 (pmr, rev, struct)

Bélanger, A. *et al.*, *Can. J. Chem.*, 1979, **57**, 3348-3354 (synth)

Jones, L.R. *et al.*, *J. Biol. Chem.*, 1981, **256**, 11809-11818 (pharmacol)

Waterhouse, A.L. *et al.*, *Chem. Comm.*, 1984, 1265-1268 (Dehydoryanodine)

Ruest, L. *et al.*, *Can. J. Chem.*, 1985, **63**, 2840-2843 (*Ryania diterpene esters*)

Waterhouse, A.L. *et al.*, *J.C.S. Perkin 2*, 1985, 1011-1016 (cmr, pmr)

Deslongchamps, P. *et al.*, *Can. J. Chem.*, 1990, **68**, 115-126; 127-152; 153-185; 186-192 (synth)

Jefferies, P.R. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1147-1149 (*Ryanodyl 3-nicotinate*)

Pesticide Manual, 9th edn., 1991, No. 10620

Ruest, L. *et al.*, *Can. J. Chem.*, 1993, **71**, 634-638 (3-Epiryanodine)

Achenbach, H. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1092-1096 (*Spiganthine*)

Ruest, L. *et al.*, *Can. J. Chem.*, 1999, **77**, 12-15 (8,9-Didehydoryanodine, 4-Deoxyester A)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, RSZ000

Sabadillines

S-1


Veragenine†

Alkaloids of unknown struct. Sabadillines I, II and III, which may not be pure, were separated. Isol. from *Schoenocaulon sabadilla* (Sabadilla). Amorph. solids. Elemental analyses were inconclusive. Svoboda, G.R. *et al.*, *J. Pharm. Sci.*, 1963, **52**, 772-776 (isol, ir, uv)

Safracin A

S-2

Quinonamine B. Y 16601. Y 16482β. Antibiotic Y 16601. Antibiotic Y 16482β [87578-98-1]



Absolute Configuration

C₂₈H₃₆N₄O₆ 524.616

Quinone antibiotic. CAS numbering shown. Isol. from *Pseudomonas fluorescens*. Active against gram-positive and -negative bacteria *in vitro*. Shows antitumour activity. Inhibits primary RNA synthesis. Pale yellow needles + 1H₂O (as dihydrochloride). Mp 300° dec. (dihydrochloride). [α]_D²⁰ -144 (c, 0.5 in MeOH). λ_{max} 292 (MeOH/NaOH) (Derep). λ_{max} 271 (ε 8510) (MeOH) (Derep).

▶ TX1404580

7α-Hydroxy: Safracin B. Quinonamine A. EM 5519. Y 16760. Y 16482α. Antibiotic EM 5519. Antibiotic Y 16760. Antibiotic Y 16482α [87578-99-2]

C₂₈H₃₆N₄O₇ 540.615

From *Pseudomonas fluorescens*. Active against gram-positive and -negative bacteria *in vitro*. Shows antitumour activity. Pale yellow needles + 1H₂O (as dihydrochloride). Mp 260° dec. (dihydrochloride). [α]_D²⁰ -106 (c, 0.5 in MeOH). λ_{max} 292 (MeOH/NaOH) (Derep). λ_{max} 271 (ε 8510) (MeOH) (Derep).

▶ TX1404550

7-Cyano: Cyanoquinonamine [96996-50-8]

C₂₉H₃₅N₅O₆ 549.625

Semisynthetic. Active against gram-positive bacteria and tumours.

[97059-07-9, 82029-27-4, 82029-28-5]

Pat. Coop. Treaty (WIPO), 1982, 00 146; *CA*, **97**, 4669 (isol)

Ikeda, Y. *et al.*, *J. Antibiot.*, 1983, **36**, 1279; 1284; 1290 (isol, uv, ir, pmr, cmr, ms)

Meyers, E. *et al.*, *J. Antibiot.*, 1983, **36**, 190

Cooper, R. *et al.*, *J. Antibiot.*, 1985, **38**, 24 (uv, ir, pmr, cmr, ms, struct)

Saframycin A_{D-1}

S-3

[107140-33-0]

As Saframycin A, S-4 with

R = -COCOCH₂CH₃, R' = CNC₃₀H₃₂N₄O₈ 576.605Prod. by *Streptomyces lavendulae*

NRRL11002 by directed biosynth.

Antitumour agent. Yellow powder. Sol.

MeOH, EtOAc; poorly sol. H₂O. Mp124-128°. λ_{max} 268 (ε 18600); 340 (sh)(MeOH) (Derep). λ_{max} 268 (ε 19950)

(MeOH) (Berdy).

Yazawa, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1639-

1650 (isol, uv, ir, pmr, cmr, ms, struct)


Kaneda, S. *et al.*, *J. Antibiot.*, 1987, **40**, 1640-

1642 (activity)

Saframycin A

S-4

[66082-27-7]

R = -CO²⁵COCH₃, R' = CNC₂₉H₃₀N₄O₈ 562.578

CAS numbering shown. Isol. from *Streptomyces lavendulae*. Shows antibiotic and antineoplastic props. Yellow powder. Sol. MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. hexane, H₂O. Mp 122-126°. [α]_D²⁰ +18.2 (c, 0.9 in MeOH). λ_{max} 268 (ε 21880); 368 (ε 1450) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 4.9 mg/kg; LD₅₀ (mus, ivn) 3.3 mg/kg. VN2265000

25-Alcohol (1): Saframycin AR₁. 25-Dihydrosafamycin A. Saframycin AH₂ [81382-09-4]

C₂₉H₃₂N₄O₈ 564.594

Prod. by microbial conversion of Saframycin A. Active against gram-positive bacteria, weakly active against gram-negative bacteria. Weak antineoplastic agent. Yellow powder + 1/3 H₂O. Sol. EtOAc. Mp 142-145°. [α]_D²⁵ +21.4 (MeOH). Log P -0.42 (calc). λ_{max} 268 (ε 10500) (MeOH) (Berdy).

25-Alcohol (2): Saframycin AH₁ [92621-76-6]

C₂₉H₃₂N₄O₈ 564.594

Semisynthetic, prod. by redn. of Saframycin A. Potent antineoplastic agent. Log P -0.42 (calc). C-25 Epimer of Saframycin AR₁.

5α-Hydroxy: Saframycin G

[92569-02-3]

C₂₉H₃₀N₄O₉ 578.577From *Streptomyces lavendulae*.

Yellow powder. Sol. MeOH; poorly

sol. H₂O. Mp 134-136°. [α]_D²² -28(c, 1 in MeOH). λ_{max} 268 (ε 21880);368 (ε 1450) (MeOH) (Derep). λ_{max}

264 (ε 17500); 370 (ε 891) (MeOH)

(Berdy).

14-Oxo, 1,4-hydroquinone: Saframycin F [92569-03-4]

C₂₉H₃₀N₄O₉ 578.577Prod. by *Streptomyces lavendulae*.

Antineoplastic agent. Light yellow

powder. Sol. MeOH; poorly sol. H₂O.Mp 134-136° dec. [α]_D²² +28.4 (c, 0.1

in MeOH). Planar struct. only currently

known. λ_{max} 243 (ε 13800); 274

(ε 17400); 369 (ε 5620) (MeOH)

(Derep). λ_{max} 231 (ε 9550); 277 (ε

12880); 375 (ε 4466) (MeOH)

(Berdy).

Decyano: Saframycin B

[66082-28-8]

C₂₈H₃₁N₃O₈ 537.568From *Streptomyces lavendulae*. Or-ange-yellow prisms (Et₂O). Sol.MeOH, Py, CHCl₃, C₆H₆; fairly sol.Et₂O; poorly sol. hexane, H₂O. Mp108-109°. [α]_D²⁰ -54.4 (c, 1 in MeOH).λ_{max} 268 (ε 21880); 368 (ε 1450)

(MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 250 mg/kg. TX1418200

Decyano, 25-alcohol: Saframycin AR₃. 25-Dihydrosafamycin B. 21-Decyano-25-dihydrosafamycin A

[81853-83-0]

C₂₈H₃₃N₃O₈ 539.584

Prod. by microbial conversion of

Saframycin B. Active against gram-

positive and -negative bacteria. Yellow

powder + 1/2 H₂O. Sol. EtOAc. Mp 123-126°. [α]_D²⁵ -76.7 (MeOH). Log P 0.12(calc). λ_{max} 269 (ε 13100) (MeOH)

(Berdy).

Decyano, 5α-methoxy: Saframycin C

[66082-29-9]

C₂₉H₃₃N₃O₉ 567.594From *Streptomyces lavendulae*. Orangeneedles (Et₂O). Sol. MeOH, Py, C₆H₆,CHCl₃; fairly sol. Et₂O; poorly sol.H₂O, hexane. Mp 143-146°. [α]_D²⁰ -20.8(c, 1 in MeOH). λ_{max} 268 (ε 21880);

368 (ε 1450) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 250 mg/kg. TX1418300

Decyano, 14-oxo, 1,4-hydroquinone: Saframycin D

[66082-30-2]

C₂₈H₃₁N₃O₉ 553.568Prod. by *Streptomyces lavendulae*.

Shows antibiotic and antineoplastic

props. Yellow needles. Sol. MeOH,

CHCl₃, Py, C₆H₆; fairly sol. Et₂O,EtOAc, Me₂CO; poorly sol. hexane,H₂O. Mp 150-154°. [α]_D²⁰ +141 (c, 1 inMeOH). λ_{max} 243 (ε 13800); 274 (ε

17400); 369 (ε 5620) (MeOH) (De-

rep).

▶ LD₅₀ (mus, ipr) 250 mg/kg. TX1486750

Decyano, 5α-hydroxy, 1,4-hydroquinone: Saframycin E

[66082-31-3]

C₂₈H₃₃N₃O₉ 555.583Isol. from *Streptomyces lavendulae*.Yellow powder. Sol. MeOH, CHCl₃,Py, C₆H₆; fairly sol. Et₂O, Me₂CO,EtOAc, CHCl₃; poorly sol. hexane,H₂O. Mp 146-148°. [α]_D²⁰ -37.3 (c, 0.53in MeOH). λ_{max} 243 (ε 13800); 274 (ε

17400); 369 (ε 5620) (MeOH) (Derep).

λ_{max} 272 (ε 10700); 368 (ε 8500)

(MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 250 mg/kg. VN2265100

(±)-form

Decyano: [82660-65-9]

C₂₈H₃₁N₃O₈ 537.568

Synthetic. Mp 175-180° dec.

Arai, T. et al., *J. Antibiot.*, 1977, **30**, 1015-1018
(*Saframycins A,B,C,D,E, isol, ir, uv, ms, pmr*)

Ger. Pat., 1979, 2 839 668; *CA*, **90**, 166572
(*isol*)

Arai, T. et al., *Tet. Lett.*, 1979, 2355-2358
(*Saframycin C, cryst struct, Saframycin B*)

Arai, T. et al., *Experientia*, 1980, **36**, 1025-1027
(*uv, ir, pmr, cmr, ms, cd, struct*)

Lown, J.W. et al., *Can. J. Chem.*, 1981, **59**,
2945-2952 (*pmr, bibl*)

Takahashi, K. et al., *J. Antibiot.*, 1982, **35**, 196-
202 (25-alcohol, decyano 25-alcohol)

Yazawa, K. et al., *J. Antibiot.*, 1982, **35**, 915-
917 (*biosynth*)

Fukuyama, T. et al., *J.A.C.S.*, 1982, **104**, 4957-
4958; 1990, **112**, 3712-3713 (*Saframycin B, synth*)

Arai, T. et al., *Alkaloids (Academic Press)*,
1983, **21**, 89-95 (*biol props*)

Kishi, K. et al., *J. Antibiot.*, 1984, **37**, 847-852
(*struct, props, Saframycins D,F,G*)

Haruyama, H. et al., *Chem. Pharm. Bull.*,
1985, **33**, 905-915 (*pmr, abs config*)

Cooper, R. et al., *J. Antibiot.*, 1985, **38**, 24-30
(*ms*)

Mikami, Y. et al., *J. Biol. Chem.*, 1985, **260**,
344-348 (*biosynth*)

Sachleben, R.A. et al., *Diss. Abstr. Int.*, **B**,
1986, **46**, 1429 (*synth*)

Kubo, A. et al., *Chem. Pharm. Bull.*, 1987, **35**,
440-442 (*Saframycin D, uv, ir, pmr, cmr, ms, struct*)

Mikami, Y. et al., *J. Antibiot.*, 1988, **41**, 734-
740 (*Saframycins F,G*)

Kubo, A. et al., *J.O.C.*, 1988, **53**, 4295-4310
(*Saframycin B, synth*)

Saito, N. et al., *Tetrahedron*, 1990, **46**, 7711-
7728 (*Saframycins C,D, synth*)

Shawe, T.T. et al., *Tetrahedron*, 1991, **47**, 5643-
5666 (*Saframycin B, synth*)

Saito, N. et al., *Chem. Pharm. Bull.*, 1995, **43**,
777-782 (*Saframycin E, synth*)

Myers, A.G. et al., *J.A.C.S.*, 1999, **121**, 10828-
10829 (*synth*)

Martinez, E.J. et al., *Org. Lett.*, 1999, **1**, 75-77
(*Saframycin A, synth*)

Saito, N. et al., *Heterocycles*, 2001, **55**, 21-28
(*Saframycin D, pmr, cmr*)

U.S. Pat., 2001, ((*Pharma Mar S.A.*))7 241
892; *CA*, **134**, 17616 (*biol*)

Saframycin H

S-5

[92569-01-2]

As Saframycin A, S-4 with

R = -COC(OH)(CH₃)CH₂COCH₃, R' = CN

C₃₂H₃₆N₄O₉ 620.658

Prod. by *Streptomyces lavendulae*

NRRL11002. Antitumour agent. Light yellow needles. Sol. MeOH; poorly sol. H₂O. Mp 184-186° dec. [α]_D²² -7 (c, 0.1 in MeOH). λ_{max} 268 (ε 15500) (MeOH) (Derep).

Takahashi, K. et al., *J. Antibiot.*, 1982, **35**, 196-
202 (*isol*)


Yazawa, K. et al., *J. Antibiot.*, 1982, **35**, 915-
917 (*isol*)

Kishi, K. et al., *J. Antibiot.*, 1984, **37**, 847-852
(*struct*)

Mikami, Y. et al., *J. Antibiot.*, 1988, **41**, 734-
740 (*struct*)

Saframycin Mx 1

[113036-78-5]



C₂₉H₃₈N₄O₉ 586.641

Quinone antibiotic. CAS numbering shown. Isol. from *Myxococcus xanthus*. Sol. MeOH, EtOAc; poorly sol. H₂O, C₆H₆, hexane. [α]_D²⁰ -70.7 (c, 0.5 in MeOH). λ_{max} 273 (ε 8910) (MeOH) (Derep). λ_{max} 290 (pH 3.5 buffer) (Berdy).

▶ TX1404590

1,4-Quinone: Saframycin Mx 1BC

[116925-60-1]

C₂₉H₃₆N₄O₉ 584.625

Prod. by *Myxococcus xanthus*. Orange-yellow solid.

7-Deoxy: Saframycin Mx 2

[113036-79-6]

C₂₉H₃₈N₄O₈ 570.641

Isol. from *Myxococcus xanthus*. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁰ -119.8 (c, 0.5 in MeOH). λ_{max} 273 (ε 8910) (MeOH) (Derep). λ_{max} 290 (pH 3.5 buffer) (Berdy).

7-Deoxy, 1,4-quinone: Saframycin Mx 2BC. Mx 2BC

[116925-61-2]

C₂₉H₃₆N₄O₈ 568.625

Prod. by *Myxococcus xanthus*. Orange-yellow solid.

Trowitzsch-Kienast, W. et al., *Annalen*, 1988,
475-481 (*isol, pmr, cmr, struct*)


Irschik, H. et al., *J. Antibiot.*, 1988, **41**, 993-
998 (*isol, ir, pmr, cmr, struct*)

Japan. Pat., 1988, 88 49 092; *CA*, **110**, 55972p
(*Mx 1BC, Mx 2BC*)

Saito, N. et al., *Tetrahedron*, 1995, **51**, 8213-
8230; 8231-8246 (*synth, abs config*)

Saframycin R

[85411-35-4]



C₃₁H₃₄N₄O₁₀ 622.63

Quinone antibiotic. Struct. revised in 2000. Prod. by *Streptomyces lavendulae*.

S-6

Shows antitumour props. and activity against gram-positive and -negative bacteria. Pale yellow needles. Sol. DMSO, MeOH, EtOAc; fairly sol. CHCl₃, C₆H₆; poorly sol. H₂O, hexane. Mp 184-186° dec. [α]_D²⁰ -79.2 (c, 0.6 in CHCl₃). λ_{max} 270 (ε 8910) (MeOH) (Derep).

▶ LD₅₀ (mus, ivn) 29.5 mg/kg, LD₅₀ (mus, ipr) 26 mg/kg.

Asaoka, T. et al., *J. Antibiot.*, 1982, **35**, 1708-
1710 (*isol, pmr, cmr*)

Lown, J.W. et al., *J. Antibiot.*, 1983, **36**, 1184-
1195 (*struct, nmr*)

Saito, N. et al., *Tetrahedron*, 2000, **56**, 9937-
9944 (*pmr, cmr, struct*)

Saframycin S

S-8

21-Hydroxysaframycin B

[75425-66-0]

As Saframycin A, S-4 with R = -COC(OH)₃, R' = OH

C₂₈H₃₁N₃O₉ 553.568

Isol. from *Streptomyces lavendulae*.

Shows antineoplastic and antibiotic props. Dark yellow powder. Sol.

MeOH, CHCl₃; fairly sol. Et₂O; poorly sol. hexane, H₂O. Mp 107-115° dec.

[α]_D²⁵ +32.5 (c, 0.5 in MeOH). λ_{max} 268 (ε 21880); 368 (ε 1450) (MeOH) (Derep).

λ_{max} 268 (ε 15200) (MeOH) (Berdy).

▶ VN2265300

Arai, T. et al., *J. Antibiot.*, 1980, **33**, 951-960
(*isol, uv, ir, pmr, cmr, struct*)


Yazawa, K. et al., *J. Antibiot.*, 1982, **35**, 915-
917 (*biosynth*)

Cooper, R. et al., *J. Antibiot.*, 1985, **38**, 24-30
(*cmr*)

Saframycin Y_{2b}

S-9

[107140-34-1]



C₅₈H₆₄N₁₀O₁₄ 1125.202

Prod. by *Streptomyces lavendulae*

NRRL11002 by directed biosynth.

Antitumour agent. Orange-yellow powder + 1H₂O. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 200° dec. [α]_D²² -42.2 (c, 0.1 in MeOH). λ_{max} 268 (ε 45700) (MeOH) (Derep).

Yazawa, K. et al., *J. Antibiot.*, 1986, **39**,
1639-1650 (*isol, uv, ir, pmr, cmr, ms, struct*)

Kaneda, S. et al., *J. Antibiot.*, 1987, **40**, 1640-
1642 (*activity*)

Saframycin Y_{2b-D}

S-10

[107163-30-4]

As Saframycin Y_{2b}, S-9 with

R = CH₂CH₃

C₆₀H₆₈N₁₀O₁₄ 1153.256

Prod. by *Streptomyces lavendulae* NRRL11002 by directed biosynth. Antitumour agent. Orange-yellow powder + 1H₂O. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{25}$ -41.9 (c, 0.1 in MeOH). Dec. >200°. λ_{\max} 268 (ε 45700) (MeOH) (Derep). λ_{\max} 269 (ε 43650) (MeOH) (Berdy).

Yazawa, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1639-1650 (*isol, uv, ir, pmr, cmr, ms, struct*)
Kameda, S. *et al.*, *J. Antibiot.*, 1987, **40**, 1640-1642 (*activity*)

Saframycin Y₃ S-11

25-Deoxy-25-aminosaframycin A [98205-62-0]

As Saframycin A, S-4 with R = -COCH(NH₂)CH₃, R' = CN
C₂₉H₃₃N₅O₇ 563.609

Quinone antibiotic. Prod. by *Streptomyces lavendulae* by directed biosynth. Shows antibacterial and antitumour props. Yellow amorph. powder. Sol. MeOH, C₆H₆, CHCl₃, EtOAc; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 143-146° dec. $[\alpha]_D^{25}$ -46.1 (c, 0.7 in MeOH). λ_{\max} 268 (ε 18600); 340 (sh) (MeOH) (Derep).

Yazawa, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1639-1650 (*isol, uv, ir, pmr, cmr, struct*)
Kameda, S. *et al.*, *J. Antibiot.*, 1987, **40**, 1640-1642 (*activity*)

Saframycin Y_{D-1} S-12

26-Homosafamycin Y₃ [98205-63-1]

As Saframycin A, S-4 with R = -COCH(NH₂)CH₂CH₃, R' = CN
C₃₀H₃₅N₅O₇ 577.636
Isol. from *Streptomyces lavendulae* by directed biosynth. Antitumour agent. Yellow amorph. powder + 1H₂O. Sol. MeOH, CHCl₃, C₆H₆, EtOAc; fairly sol. Et₂O; poorly sol. H₂O, hexane. Mp 124-127° dec. $[\alpha]_D^{25}$ -43.5 (c, 1 in MeOH). λ_{\max} 268 (ε 18600); 340 (sh) (MeOH) (Derep).

Yazawa, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1639-1650 (*isol, uv, ir, pmr, cmr, struct*)
Kameda, S. *et al.*, *J. Antibiot.*, 1987, **40**, 1640-1642 (*activity*)

Saframycin Y_{D-2} S-13


26-Demethylsaframycin Y₃ [98205-61-9]

As Saframycin A, S-4 with R = -COCH₂NH₂, R' = CN
C₂₈H₃₁N₅O₇ 549.582
Prod. by *Streptomyces lavendulae* NRRL11002 by directed biosynth. Antitumour agent. Dark yellow powder + 1/2 H₂O. Sol. CHCl₃, C₆H₆; fairly sol. Et₂O; poorly sol. hexane. Mp 144-148° dec. λ_{\max} 268 (ε 18600); 340 (sh) (MeOH) (Derep). λ_{\max} 268 (ε 17100) (MeOH) (Berdy).

Yazawa, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1639-1650 (*isol, uv, ir, pmr, struct*)
Kameda, S. *et al.*, *J. Antibiot.*, 1987, **40**, 1640-1642 (*activity*)

Sagitol

[185543-92-4]




C₂₁H₁₈N₄O₂S 390.465
Minor alkaloid from the sponge *Oceanapia sagittaria*. Yellow glass. Closely related to Kuanoniamine C in D-93. λ_{\max} 260 (ε 22810); 318 (ε 3430); 335 (ε 3920); 351 (ε 3430); 396 (ε 1470) (MeOH).

Salomon, C.E. *et al.*, *Tet. Lett.*, 1996, **37**, 9147 (*isol, uv, ir, pmr, cmr, struct*)

Sagittamide A

S-15



C₄₈H₈₁N₃O₁₈ 988.177
Absolute configuration confirmed in 2007. Isol. from an unidentified didemnid ascidian from Micronesia. Glass. $[\alpha]_D^{25}$ -22 (c, 0.6 in MeOH). λ_{\max} 214; 228; 276 (MeOH).

7-O-De-Ac: Sagittamide B


C₄₆H₇₉N₃O₁₇ 946.14
Isol. from an unidentified didemnid ascidian from Micronesia. Glass. $[\alpha]_D^{25}$ -19.7 (c, 0.17 in MeOH). λ_{\max} 214 (ε 1534); 256 (ε 702) (MeOH).

Lievens, S.C. *et al.*, *Org. Lett.*, 2005, **7**, 2281-2284 (*isol, pmr, cmr*)
Lievens, S.C. *et al.*, *J.A.C.S.*, 2006, **128**, 11764-11765 (*abs config*)
Seike, H. *et al.*, *Org. Lett.*, 2006, **8**, 3865-3868 (*synth, abs config*)
Schuetz, A. *et al.*, *J.A.C.S.*, 2007, **129**, 15114-15115 (*nmr, abs config*)

Saifine

S-16

[161099-48-5]



C₁₉H₂₆N₂O₃ 330.426
Indole alkaloid possibly derived from an Eburna type with loss of C-22, but the details are not yet known. Alkaloid from roots of *Rhazya stricta* (Apocynaceae).

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1995, **5**, 245 (*isol*)

S-14

Saieimine


S-17

C₂₇H₄₃NO₄ 445.641
Steroidal alkaloid. Struct. unknown. Alkaloid from the Chinese drug Bei-son-pei-mu. Mp 256-258°. λ_{\max} 288 nm. Chu, T.T. *et al.*, *CA*, 1951, **51**, 445

Salacin†

S-18

[128638-14-2]



Relative Configuration

C₁₇H₂₀N₂O₃ 300.357

Related to the Corynanthe alkaloids. CAS indexes under the abs. config. as shown, but there does not appear to be any firm evidence for this. Alkaloid from the leaves of *Uncaria salaccensis* (*Uncaria attenuata*) (Rubiaceae). Amorph. λ_{\max} 208; 250; 282 (MeOH).


[128706-55-8]

Ponglux, D. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 573-575 (*isol, synth, struct*)

Salarin A

S-19

[1005147-97-6]




C₃₅H₄₆N₂O₁₂ 686.755
Isol. from a *Fascaplysinopsis* sp. Cytotoxic. Pale yellow oil. $[\alpha]_D^{23}$ -57 (c, 0.37 in CHCl₃).

Bishara, A. *et al.*, *Org. Lett.*, 2008, **10**, 153-156 (*isol, pmr, cmr*)

Salarin B

S-20

[1005147-98-7]



C₃₆H₅₂N₂O₁₃ 720.812
Isol. from a *Fascaplysinopsis* sp. Oil. $[\alpha]_D^{23}$


-130 (c, 0.12 in CHCl₃).

Bishara, A. *et al.*, *Org. Lett.*, 2008, **10**, 153-156
(*isol*, *pmr*, *cmr*)

Salfredin C₁

S-21

[139542-56-6]



C₁₃H₁₁NO₆ 277.233

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Pale yellow powder. Sol. EtOAc, Me₂CO, MeOH, EtOH. λ_{max} 234 (ε 21600); 247 (sh) (ε 16100); 393 (ε 4160) (MeOH/NaOH) (Derep). λ_{max} 223 (ε 18600); 248 (ε 23300); 335 (ε 3050) (MeOH) (Derep). λ_{max} 223 (E1%/1cm 380); 248 (E1%/1cm 770); 335 (E1%/1cm 110) (MeOH/HCl) (Berdy).

N-(Carboxymethyl): **Salfredin C₂**

[139542-57-7]

C₁₅H₁₃NO₈ 335.27

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Prisms. Sol. EtOAc, EtOH, Me₂CO, MeOH. Mp 246-247°. λ_{max} 223 (E1%/1cm 600); 247 (E1%/1cm 900); 335 (E1%/1cm 110) (MeOH) (Berdy). λ_{max} 223 (E1%/1cm 520); 247 (E1%/1cm 840); 335 (E1%/1cm 110) (MeOH/HCl) (Berdy). λ_{max} 241 (E1%/1cm 900); 395 (E1%/1cm 150) (MeOH/NaOH) (Berdy).

N-(1-Carboxyethyl): **Salfredin C₃**

[139542-58-8]

C₁₆H₁₅NO₈ 349.296

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Pale yellow oil. Sol. EtOAc, EtOH, MeOH, Me₂CO. λ_{max} 223; 247; 335 (MeOH) (Berdy). λ_{max} 223; 247; 335 (MeOH/HCl) (Berdy). λ_{max} 241; 395 (MeOH/NaOH) (Berdy).

5-Deoxo, N-(carboxymethyl): **Salfredin A₄**

[139542-54-4]

C₁₅H₁₅NO₇ 321.286

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Prisms. Sol. EtOAc, MeOH, Me₂CO, EtOH. Mp >300°. λ_{max} 220 (sh) (ε 20200); 275 (ε 4820); 316 (ε 1930) (MeOH/NaOH) (Derep). λ_{max} 215 (ε 26000); 258 (ε 6100); 303 (ε 1930) (MeOH) (Derep). λ_{max} 215 (E1%/1cm 710); 258 (E1%/1cm 190); 303 (E1%/1cm 60) (MeOH/HCl) (Berdy).

5-Deoxo, N-(1-carboxyethyl): **Salfredin A₇**

[139542-55-5]

C₁₆H₁₇NO₇ 335.313

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Prisms. Sol. EtOAc, EtOH, MeOH, Me₂CO. Mp 293-296° dec. λ_{max} 220 (sh) (ε 20200); 275 (ε 4820); 316 (ε 1930)

(MeOH/NaOH) (Derep). λ_{max} 215 (ε 26000); 258 (ε 6100); 303 (ε 1930) (MeOH) (Derep).

5-Deoxo, N-(1,3-dicarboxypropyl): **Salfredin A₃**

[139542-53-3]

C₁₈H₁₉NO₉ 393.349

Prod. by *Crucibulum* sp. RF-3817. Aldose reductase inhibitor. Powder. Sol. EtOAc, Me₂CO, EtOH, MeOH. λ_{max} 220 (sh) (ε 24800); 275 (ε 5900); 316 (ε 2360) (MeOH/NaOH) (Derep). λ_{max} 215 (ε 31800); 258 (ε 7470); 303 (ε 2360) (MeOH) (Derep). λ_{max} 217 (E1%/1cm 770); 260 (E1%/1cm 230); 303 (E1%/1cm 70) (MeOH/HCl) (Berdy).

Matsumoto, K. *et al.*, *J. Antibiot.*, 1995, **48**, 439 (*isol*, *pmr*, *cmr*)

Salicilobine

S-22

Struct. unknown. Alkaloid from *Lobelia salicifolia*. Mp 280-285°.

Steinegger, I.E. *et al.*, *Pharm. Acta Helv.*, 1956, **31**, 89-96 (*isol*)

Salicornine

S-23

[1361-03-1]

C₃H₁₁NO₂ 117.147

Struct. unknown. Alkaloid from *Salicornia herbacea* (Chenopodiaceae).

Hydrochloride: Mp 253-256°.

Picrate: Mp 210-211°.

Borkowski, B. *et al.*, *Pharmazie*, 1965, **20**, 390-393; *C.A.*, **63**, 15223c

Saliherbine

S-24

[1361-04-2]

C₉H₂₁N₅O₄ 263.296

Struct. unknown. Alkaloid from *Salicornia herbacea* (Chenopodiaceae).


Hydrochloride: Mp 318-321°.

Picrate: Mp 261-262°.

Borkowski, B. *et al.*, *Pharmazie*, 1965, **20**, 390-393; *C.A.*, **63**, 15223c

Salimine

S-25



C₂₂H₂₅NO₇ 415.442

Struct. revised in 1994.

(S)-form [138704-12-8]

Alkaloid from *Colchicum decaisnei* (Liliaceae). Amorph. solid. [α]_D -17 (c, 0.11 in MeOH).


Abu Zarga, M.H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 936 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Banwell, M.G. *et al.*, *Chem. Comm.*, 1994, 2647 (*struct*)

Saliniketol A

S-26

[923952-66-3]



C₂₂H₃₇NO₅ 395.538

Prod. by the marine-derived *Streptomyces arenicola* CNR-005. Inhibitor of ornithine decarboxylase induction. Amorph. powder. [α]_D -13.7 (c, 0.13 in MeOH). λ_{max} 240 (log ε 3.1) (MeOH).

18-Hydroxy: **Saliniketol B**

[923952-68-5]

C₂₂H₃₇NO₆ 411.537

Prod. by a marine-derived *Streptomyces arenicola* CNR-059. Amorph. powder. [α]_D -22.4 (c, 0.11 in MeOH). λ_{max} 242 (log ε 4) (MeOH).

Williams, P.G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 83-88 (*isol*, *pmr*, *cmr*)

Paterson, I. *et al.*, *Org. Lett.*, 2008, **10**, 3295-3298 (*synth*)

Salinosporamide I

S-27

[872360-12-8]

As Salinosporamide A, S-28 with R¹ = -CH₂CH₂Cl, R² = -CH₂CH₃

C₁₆H₂₂ClNO₄ 327.807

Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²⁵ -20 (c, 0.9 in MeCN). Full stereochem. not confirmed. λ_{max} 205 (log ε 3.19); 220 (sh) (log ε 2.99) (MeCN).


Reed, K.A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 269-276 (*isol*, *pmr*, *cmr*, *ms*)

Salinosporamide A

S-28

NPI 0052. Antibiotic NPI 0052

[437742-34-2]



R¹ = -CH₂CH₂Cl, R² = CH₃

C₁₅H₂₀ClNO₄ 313.78

Prod. by the marine bacterium *Salinospora tropica* (strain CNB-392). Proteasome inhibitor. Cytotoxic. Needles (EtOAc/2,3,3-trimethylpentane). Mp 169-171°. [α]_D²⁵ -72.9 (c, 0.55 in MeOH). λ_{max} 205 (log ε 4.03); 225 (log ε 3.3) (MeOH).

5-Deoxy: **Salinosporamide J**

[932739-03-2]

C₁₅H₂₀ClNO₃ 297.781

Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²² -120 (c, 0.3 in MeCN). λ_{max} 217 (log ε 3.21); 230 (sh) (log ε 3.19) (MeCN).

Dechloro: **Salinosporamide B**. NPI 0047.

Antibiotic NPI 0047

[863126-95-8]

C₁₅H₂₁NO₄ 279.335

Prod. by *Salinospora tropica* (strain CNB-392). Cryst. (EtOAc). Mp 143-145°. [α]_D²⁵ -54.5 (c, 0.29 in MeOH). λ_{max} 256 (log ε 3.7) (MeOH).

Bromo analogue: Bromosalinosporamide

[872360-15-1]

C₁₅H₂₀BrNO₄ 358.231

Semisynthetic. Prod. by *Salinospora tropica* in NaBr. Amorph. solid. [α]_D²² -51.6 (c, 0.95 in MeCN).

2-Epimer: Salinosporamide F

[872360-11-7]

C₁₅H₂₀ClNO₄ 313.78

Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²² -108.9 (c, 8.78 in MeCN). λ_{max} 206 (log ε 3.16); 223 (sh) (log ε 3.05) (MeCN).

2-Epimer, dechloro: Salinosporamide H

[872360-16-2]

C₁₅H₂₁NO₄ 279.335

Prod. by *Salinospora tropica* in NaBr. Amorph. solid. λ_{max} 221 (sh) (MeCN aq.).

Feling, R.H. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 355-357 (*isol, pmr, cmr, cryst struct*)

Macherla, V.R. *et al.*, *J. Med. Chem.*, 2005, **48**, 3684-3687 (*sar*)

Endo, A. *et al.*, *J.A.C.S.*, 2005, **127**, 8298-8299 (*synth*)

Williams, P.G. *et al.*, *J.O.C.*, 2005, **70**, 6196-6203 (*Salinosporamide B, isol, pmr, cmr*)

Reddy, L.R. *et al.*, *Org. Lett.*, 2005, **7**, 2699-2701 (*synth*)

Groll, M. *et al.*, *J.A.C.S.*, 2006, **128**, 5136-5141 (*proteasome complex, struct, cryst struct*)

Ruiz, S. *et al.*, *Mol. Cancer Ther.*, 2006, **5**, 1836-1843 (*pharmacol*)

Shibasaki, M. *et al.*, *Chem. Asian J.*, 2007, **2**, 20-38 (*rev, synth*)

Lam, K.S. *et al.*, *J. Antibiot.*, 2007, **60**, 13-19 (*isol, props*)

Reed, K.A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 269-276 (*isol, pmr, cmr, cryst struct*)

Beer, L.L. *et al.*, *Org. Lett.*, 2007, **9**, 845-848 (*biosynth*)

Ma, G. *et al.*, *Org. Lett.*, 2007, **9**, 2143-2146 (*synth*)

Ling, T. *et al.*, *Org. Lett.*, 2007, **9**, 2289-2292 (*synth*)

Caubert, V. *et al.*, *Tet. Lett.*, 2007, **48**, 381-384 (*synth*)

Mulholland, N.P. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 2782-2789 (*synth*)


Fukuda, T. *et al.*, *Org. Lett.*, 2008, **10**, 4239-4242 (*synth*)

Margalef, I.V. *et al.*, *Tetrahedron*, 2008, **64**, 7896-7901 (*synth*)

Salinosporamide C

S-29

[863126-96-9]



Relative Configuration

C₁₄H₁₈ClNO₃ 283.754

Prod. by the marine bacterium *Salinospora tropica* (strain CNB-392). Cytotoxic. Oil. [α]_D²⁵ -33.6 (c, 0.27 in MeOH). Possible artifact. λ_{max} 222 (log ε 3.9)

(MeOH).

Williams, P.G. *et al.*, *J.O.C.*, 2005, **70**, 6196-6203 (*isol, pmr, cmr*)

Salinosporamide D

S-30

[823229-26-1]

As Salinosporamide A, S-28 with

R¹ = R² = CH₃C₁₄H₁₉NO₄ 265.308

Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²² -63.8 (c, 4.71 in MeCN). λ_{max} 208 (log ε 3.06); 223 (sh) (log ε 3.06) (MeCN).

2-Epimer: Salinosporamide G

[872360-13-9]

C₁₄H₁₉NO₄ 265.308

Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²² -54.5 (c, 1.1 in MeCN). λ_{max} 209 (log ε 3.09); 218 (sh) (log ε 3.08) (MeCN).

Reed, K.A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 269-276 (*isol, pmr, cmr*)

Salinosporamide E

S-31

[872360-24-2]

As Salinosporamide A, S-28 with

R¹ = CH₂CH₂CH₃, R² = CH₃C₁₆H₂₃NO₄ 293.362


Prod. by *Salinospora tropica* NPS000465. Amorph. solid. [α]_D²² -60 (c, 0.1 in MeCN).

Reed, K.A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 269-276 (*isol, pmr, cmr*)

Salmoachelin 4

S-32

[768386-89-6]

C₄₂H₄₇N₃O₂₅ 993.838


Related to Enterobactin, E-86. Isol. from *Salmonella enterica*. Catecholate siderophore.

Bister, B. *et al.*, *BioMetals*, 2004, **17**, 471-481 (*isol, pmr, cmr, ms*)

Salmoachelin X

S-33

N-(5-β-D-Glucopyranosyl-2,3-dihydroxybenzoyl)serine. Pacifarinic acid
[768386-88-5]

C₁₆H₂₁NO₁₁ 403.342


Isol. from *Salmonella enterica*. Catecholate siderophore.

Bister, B. *et al.*, *BioMetals*, 2004, **17**, 471-481

Salmycin B

S-34

[163596-95-0]



n = 4

C₄₁H₆₉FeN₆O₂₁ 1037.871

Prod. by *Streptomyces violaceus* DSM 8286. Siderophore. Antibacterial agent. Sol. H₂O, MeOH; poorly sol. EtOH, hexane. Possesses same aglycone as Danomycin, D-16. λ_{max} 427 (ε 1995) (MeOH) (Berdy).

2'-Oxime: Salmycin A

[163596-94-9]

C₄₁H₇₀FeN₇O₂₁ 1052.885

Prod. by *Streptomyces violaceus* DSM 8286. Siderophore. Sol. H₂O, MeOH; poorly sol. EtOH, hexane. λ_{max} 430 (ε 1995) (MeOH) (Berdy).

Vertesy, L. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 46-60 (*isol, ir, pmr, cmr, struct*)

Dong, L. *et al.*, *J.A.C.S.*, 2002, **124**, 15001-15005 (*synth*)

Salmycin C

S-35

[163596-96-1]

As Salmycin B, S-34 with

n = 3

C₄₀H₆₇FeN₆O₂₁ 1023.844

Prod. by *Streptomyces violaceus* DSM 8286. Siderophore. Antibacterial agent. Sol. H₂O, MeOH; poorly sol. EtOH, hexane. λ_{max} 430 (ε 2000) (MeOH) (Berdy).

2'-Oxime: Salmycin D

[163596-97-2]


C₄₀H₆₈FeN₇O₂₁ 1038.858

Prod. by *Streptomyces violaceus* DSM 8286. Siderophore. Sol. H₂O, MeOH; poorly sol. EtOH, hexane. λ_{max} 427 (ε 200) (MeOH) (Berdy).

Vertesy, L. *et al.*, *Helv. Chim. Acta*, 1995, **78**, 46-60 (*isol, ir, pmr, cmr, struct*)

Salsamine

S-36

C₂₆H₃₆N₂O₄ 440.581

Defective diag. in *CA*. Alkaloid from


Salsola richteri (Chenopodiaceae). Mp 165-167°.

Proskurnina, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1937, **4**, 1265 (*isol*)

Proskurnina, N.F. *et al.*, *Zh. Obshch. Khim.*, 1958, **28**, 256; *CA*, **52**, 12879f (*struct*)

Saludimerine A

S-37



Absolute Configuration

C₃₈H₄₀N₂O₈ 652.743

Dimer of Salutaridine, S-39. Alkaloid from the leaves of *Croton flavens*. Mp 220° dec. [α]_D²⁰ -219 (c, 1.7 in MeOH). λ_{max} 241 (log ε 4.56); 283 (log ε 4.08) (MeOH).

Atropisomer: **Saludimerine B**

C₃₈H₄₀N₂O₈ 652.743


Alkaloid from the leaves of *Croton flavens*. Mp 225° dec. [α]_D²⁰ -154 (c, 1.4 in MeOH). λ_{max} 244 (log ε 4.54); 283 (log ε 4.07) (MeOH).

Bracher, F. *et al.*, *J.O.C.*, 2004, **69**, 8602-8608 (*isol, cd, pmr, cmr*)

Salutadimerine

S-38

[131690-44-3]



C₃₈H₄₀N₂O₈ 652.743


The first reported dimeric morphinan-dienone alkaloid. Alkaloid from *Papaver lasiothrix* and *Papaver pseudo-orientale* (Papaveraceae). Amorph. [α]_D +36 (c, 0.1 in MeOH).

Sariyar, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1383 (*isol, uv, ir, pmr, cmr, ms, struct*)

Salutaridine

S-39

Floripavine. Sinoacutine. Salutarine [1936-18-1]



(+)-form

C₁₉H₂₁NO₄ 327.379

Salutaridine was historically the (+)-form, Sinoacutine the (-)-form and Salutarine the (±)-form. However Salutaridine occurs in at least some sources as a partial racemate.

(+)-form

Alkaloid from *Croton salutaris*, *Croton balsamifera*, *Croton stenophylus*, *Papaver* spp. and *Glaucium* spp. (Euphorbiaceae, Papaveraceae). Intermediate in biosynth. of morphine alkaloids. Shows antitumour activity vs. Walker 256 carcinosarcoma. Rods (EtOAc). Mp 197-198°. [α]_D +111 (c, 1.69 in EtOH). λ_{max} 236 ; 279 (MeOH) (Berdy).

N-Oxide: **Salutaridine N-oxide**

[144860-15-1]

C₁₉H₂₁NO₅ 343.379

Alkaloid from capsules of *Papaver bracteatum* (Papaveraceae). Amorph. [α]_D +52 (c, 0.146 in MeOH).

N-De-Me: **Norsalutaridine**

[55781-26-5]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Croton salutaris* (Euphorbiaceae). [α]_D²⁴ +38.6 (c, 3.4 in EtOH).

Ac: Mp 171°. [α]_D +120 (c, 1.26 in EtOH).

5,6ξ-Dihydro: **Alkaloid Or₁**

[57157-67-2]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Papaver orientale* and *Papaver pseudo-orientale* (Papaveraceae). Cryst. (petrol). Mp 109°. [α]_D²⁰ +93 (c, 0.45 in CHCl₃). Proposed struct. Cryst. with difficulty, not obt. completely pure.

5,6-Dihydro, 2,4-dinitrophenylhydrazone: Mp 110-115°.

5,6ξ-Dihydro; methiodide: Mp 186-188°.

5,6-Dihydro, N-de-Me: **N-Noralkaloid Or₁**

[199527-84-9]

C₁₈H₂₁NO₄ 315.368

Alkaloid from *Papaver pseudo-orientale*.

(-)-form [4090-18-0]

Alkaloid from *Sinomenium acutum*, *Corydalis* spp., *Nandina* spp. and others in the Menispermaceae, Papaveraceae and Lauraceae. Mp 198°. [α]_D¹⁶ -112 (EtOH).

Perchlorate: Mp 117°.

Methiodide: Mp 216°.

N-De-Me: **Norsinoacutine**

[13186-21-5]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Croton* spp. (Euphorbiaceae). Cryst. + EtOAc (MeOH/EtOAc). Mp 113-115°. [α]_D¹⁶ -107 (c, 1.14 in MeOH).

Me ether; methiodide:

Cryst. + Me₂CO. Mp 163-165°.

O³-De-Me, O⁴-Me: **Isosinoacutine**

[76202-24-9]

C₁₉H₂₁NO₄ 327.379

Alkaloid from the roots of *Stephania elegans* (Menispermaceae). Mp 120°. [α]_D -41 (CHCl₃). [α]_D -82 (90% EtOH).

O⁶-De-Me: **Mocrispatine**

[84622-66-2]

[27669-33-6]

C₁₈H₁₉NO₄ 313.352

Alkaloid from *Monodora crispata* (Annonaceae).

O⁶-De-Me, N-de-Me: **3-Methoxy-4,6-dihydroxymorphinandien-7-one**

[80248-95-9]

C₁₇H₁₇NO₄ 299.326

Alkaloid from *Croton bonplandianum* (Euphorbiaceae). Cryst. + 1EtOAc (MeOH/EtOAc). Mp 143°.

(±)-form [23979-21-7]

Alkaloid from *Croton salutaris* (Euphorbiaceae). Mp 222-224°. Co-occurs with the (+)-form.

Picrate: Mp 219-222°.

Barton, D.H.R. *et al.*, *Chem. Comm.*, 1965, 52 (*struct*)

Barton, D.H.R. *et al.*, *J.C.S.*, 1965, 2423 (*pmr, uv*)

Chambers, C. *et al.*, *Chem. Comm.*, 1966, 449

Wheeler, D.M.S. *et al.*, *J.A.C.S.*, 1967, **89**, 4494 (*ms*)

Stuart, K.L. *et al.*, *J.C.S.(C)*, 1969, 1681 (*isol, uv*)

Kametani, T. *et al.*, *J.C.S.(C)*, 1969, 2030 (*synth*)

Stuart, K.L. *et al.*, *Phytochemistry*, 1973, **12**, 1967 (*biosynth*)

Shafiee, A. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 1570 (*Alkaloid Or₁*)

Khosa, R.L. *et al.*, *Chem. Ind. (London)*, 1980, 662 (*Isosinoacutine*)

Barnes, R.A. *et al.*, *Phytochemistry*, 1981, **20**, 543 (*N-Norsalutaridine*)

Djakoure, A.L. *et al.*, *CA*, 1983, **98**, 107592v (*Mocrispatine*)

Ludwig, W. *et al.*, *Angew. Chem., Int. Ed.*, 1986, **25**, 1025 (*synth*)

Sariyar, G. *et al.*, *Planta Med.*, 1992, **58**, 368 (*Salutaridine N-oxide*)

Ribár, B. *et al.*, *Acta Cryst. C*, 1993, **49**, 978 (*cryst struct*)


Shafiee, A. *et al.*, *J. Sci., Islamic Repub. Iran*, 1997, **8**, 105-107 (*N-Noralkaloid Or₁*)

Bracher, F. *et al.*, *J.O.C.*, 2004, **69**, 8602-8608 (*isol, pmr, cmr*)

Salviadione

S-40

[862832-47-1]




C₁₉H₁₉NO₂ 293.365

Alkaloid from the roots of *Salvia miltiorrhiza*. Orange solid (EtOAc/hexane). Mp 212-214°.

Don, M.-J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1066-1070 (*isol, pmr, cmr*)

Salviमितamide S-41

1,6,7,8-Tetrahydro-2,10,11-trihydroxy-8-methoxy-2H-benzo[e]azecine-3,5-dione




C₁₄H₁₇NO₆ 295.291

Alkaloid from the rhizomes of *Salvia miltiorrhiza*. Grey powder. [α]_D²⁰ +7.3 (c, 0.1 in MeOH). λ_{max} 230 (log ϵ 3.62); 283 (log ϵ 3.45) (MeOH).

Choi, J.S. *et al.*, *Fitoterapia*, 2001, **72**, 30-34

Salvianene S-42

[862832-45-9]



C₂₁H₂₁NO₂ 319.402

Alkaloid from the roots of *Salvia miltiorrhiza*. Cytotoxic. Brown needles (EtOAc/hexane). Mp 90-92°.

15,16-Dihydro: **Salvianane**

[862832-46-0]


C₂₁H₂₃NO₂ 321.418

Alkaloid from the roots of *Salvia miltiorrhiza*. Pale green plates (EtOAc/hexane).

Don, M.-J. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1066-1070 (*isol, pmr, cmr, ms*)

Samandaridine S-43

[6384-73-2]



C₂₁H₃₁NO₃ 345.481

Present in salamanders (*Salamandra* spp.). Also occurs as glycosides in *Torulopsis magnoliae* and prod. by various *Acetobacter* spp. Cryst. (EtOH). Mp 287-288°. [α]_D²¹ +29.5 (2N AcOH).

► CL8019000

Hydrochloride:

Cryst. (H₂O). Mp 343°. [α]_D²⁴ +14.1 (H₂O).

Walker, T.K. *et al.*, *Arch. Biochem. Biophys.*, 1959, **83**, 161 (*isol*)

Gorin, P.A.J. *et al.*, *Can. J. Chem.*, 1961, **39**, 846 (*isol*)

Habermehl, G. *et al.*, *Chem. Ber.*, 1963, **96**, 840

Habermehl, G. *et al.*, *Annalen*, 1967, **706**, 213 (*ms*)


Habermehl, G. *et al.*, *Chem. Ber.*, 1968, **101**, 198 (*biosynth*)

Shimizu, Y. *et al.*, *J.O.C.*, 1976, **41**, 1930 (*synth*)

Oka, K. *et al.*, *J.A.C.S.*, 1977, **99**, 3859 (*struct*)

Samandarine S-44

1,4-Epoxy-3-aza-A-homoandrostan-16-ol [467-51-6]



Absolute configuration

C₁₉H₃₁NO₂ 305.459

Secretion of *Salamandra maculosa* and *Salamandra atra*. Toxin causing respiratory paralysis, hypertension and cardiac arrhythmia. Potent local anaesthetic not used because of its toxicity. Cryst. (EtOH aq. or Me₂CO). Mp 187-188°. [α]_D¹⁷ +43.7.

► LD₅₀ (mus, ipr) 0.3 mg/kg; Highly toxic, LD₅₀ (mus, scu) 1.5 mg/kg; LD₅₀ (rbt, ivn) 1 mg/kg. VP2405000

Hydrochloride: Mp 321-322°.

O-Ac: O-Acetylsamandarine

[1857-07-4]

C₂₁H₃₃NO₃ 347.497

Alkaloid from *Salamandra maculosa*.

Mp 159°.

Di-Ac:

Needles (EtOH aq.). Mp 167-168°.

16-Ketone: Samandarone

[467-52-7]

C₁₉H₂₉NO₂ 303.444

Main alkaloid of *Salamandra maculosa*, minor alkaloid in other *Salamandra* spp. Also from *Leptodactylus pentadactylus*. Paralytic toxin. Mp 190°.

► Toxic, LD₅₀ (mus, scu) 1.2 mg/kg.

CL8017800

Wolfel, E. *et al.*, *Chem. Ber.*, 1961, **94**, 2361-2373 (*struct, ir, ord, abs config*)

Habermehl, G. *et al.*, *Alkaloids (Academic Press)*, 1967, **9**, 427-439 (*rev, tox*)

Habermehl, G. *et al.*, *Annalen*, 1967, **706**, 213-222 (*ms*)

Habermehl, G. *et al.*, *Chem. Ber.*, 1968, **101**, 198-200 (*biosynth*)


Altona, C. *et al.*, *Tetrahedron*, 1968, **24**, 13-32 (*abs config*)

Shizimu, Y. *et al.*, *J.O.C.*, 1976, **41**, 1930-1934 (*synth*)

Samandenone S-45

1,4-Epoxy-20-methyl-3-aza-A-homo-pregn-14-en-16-one

[6400-81-3]



C₂₂H₃₃NO₂ 343.508

Alkaloid from the skin of *Salamandra maculosa*. Platelets (EtOH aq.). Mp 189-191°.

► VP2405500

Hydroiodide: Mp 310-311°.


Habermehl, G. *et al.*, *Chem. Ber.*, 1966, **99**,

1439 (*isol, struct, ir, ms, pmr*)

Habermehl, G. *et al.*, *Chem. Ber.*, 1968, **101**, 198 (*biosynth*)

Samandinine S-46

[24206-15-3]



C₂₄H₃₉NO₃ 389.577

Prob. struct. Minor alkaloid from the skin of *Salamandra maculosa*. Mp 170°.


► CL8019200

Habermehl, G. *et al.*, *Toxicol.*, 1969, **7**, 163; *CA*, **71**, 113121z (*isol, ms, ir*)

Samanine S-47

3-Aza-A-homoandrostan-16-ol

[22614-24-0]



C₁₉H₃₃NO 291.476

Alkaloid from the toxic secretion of *Salamandra maculosa taeniata*. Cryst. (EtOH). Mp 197°.

Hydrochloride: Mp 304-306°.

O,N-Di-Ac:

Cryst. (MeOH aq.). Mp 70-74°.

Habermehl, G. *et al.*, *Prog. Org. Chem.*, 1968, **7**, 35 (*isol*)


Habermehl, G. *et al.*, *Annalen*, 1969, **722**, 155 (*synth, ms*)

Oka, K. *et al.*, *Tet. Lett.*, 1969, 1193 (*synth*)

Rao, R.B. *et al.*, *Tet. Lett.*, 1973, 4971

Sambutoxin S-48

[160047-56-3]



C₂₈H₃₉NO₄ 453.62

Related to Funiculosin, F-197 and Oxysporidinone, O-226. Prod. by *Fusarium sambucinum* PZF4 and *Fusarium oxysporum*. Mycotoxin. Shows respiratory chain inhibiting activity. Prisms (MeOH). Sol. MeOH, toluene, C₆H₆, EtOH, butanol, EtOAc; poorly sol. H₂O, hexane. Mp 196.5-197.5°. [α]_D²⁵ -200 (c, 0.1 in MeOH). λ_{\max} 213 (ε 38000); 233 (ε 17000); 254 (ε 29000) (MeOH) (Berdy).

N-De-Me: N-Demethylsambutoxin

C₂₇H₃₇NO₄ 439.594

Prod. by *Fusarium oxysporum* strain N17B. Amorph. solid. [α]_D²⁶ -98.6 (c, 0.1 in CHCl₃). λ_{\max} 212 (log ε 4.32); 252 (log ε 4.03) (CHCl₃).

Kim, J.-C. *et al.*, *Tet. Lett.*, 1995, **36**, 1047-1050 (*isol, uv, ir, pmr, cmr*)


Kawaik, S. *et al.*, *Cereal Res. Commun.*, 1997, **25**, 325-326 (*activity*)

Jayasinghe, L. *et al.*, *J. Nat. Prod.*, 2006, **69**, 439-442 (*N-Demethylsambutoxin*)

Sampangine

S-49

7H-Naphtho[1,2,3-ij][2,7]naphthyridin-7-one, 9CI
[116664-93-8]



C₁₅H₈N₂O 232.241

A drawing error in the original CAS abstract incorrectly depicted Sampangine as Indeno[1,2,3-ij][2,7]naphthyridine, I-65 which caused errors in registration. Alkaloid from the stem bark of *Cananga odorata* (ylang ylang) (Annonaceae). Shows antifungal activity. Bright yellow needles (C₆H₆). Mp 210° dec. λ_{\max} 220 ; 252 ; 312 ; 326 ; 392 (MeOH) (Berdy).

3-Methoxy: 3-Methoxysampangine

[128129-42-0]

C₁₆H₁₀N₂O₂ 262.267

Alkaloid from the root bark of *Cleistopholis patens* (Annonaceae). Exhibits antifungal activity. Yellow needles (EtOAc/hexane). Mp 225-227° (213-215°).

4-Methoxy: Eupomatidine 2. 4-Methoxysampangine

[139253-60-4]

C₁₆H₁₀N₂O₂ 262.267

Alkaloid from bark of *Eupomatia laurina* (Eupomatiaceae). Fine lemon needles (MeOH/CH₂Cl₂). Mp 262-265° dec.

8-Hydroxy: 8-Hydroxysampangine

[225938-87-4]

C₁₅H₈N₂O₂ 248.24

Alkaloid from *Cleistopholis patens*. Orange powder. Mp 223-226°. λ_{\max} 233 ; 315 ; 337 ; 447 (MeOH).

9-Methoxy: Eupomatidine 1. 9-Methoxysampangine

[139220-15-8]

C₁₆H₁₀N₂O₂ 262.267

Alkaloid from aerial parts of *Eupomatia bennettii* (Eupomatiaceae). Yellow prisms (MeOH). Mp 195-197° (natural) Mp 228-231° (synthetic).

4,9-Dimethoxy: Eupomatidine 3. 4,9-Dimethoxysampangine

[139220-16-9]

C₁₇H₁₂N₂O₃ 292.293

Alkaloid from bark of *Eupomatia laurina* (Eupomatiaceae). Yellow needles (MeOH/CH₂Cl₂). Mp 245-248° dec. (natural) Mp 278-281° dec. (synthetic).

Rao, J.U.M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 346 (*isol, uv, ir, pmr, ms, struct, activity*)

Bracher, F. *et al.*, *Annalen*, 1989, 87 (*synth, uv, ir, pmr, cmr, ms*)

Liu, S. *et al.*, *Antimicrob. Agents Chemother.*, 1990, **34**, 529 (*3-Methoxysampangine*)

Carroll, A.R. *et al.*, *Aust. J. Chem.*, 1991, **44**, 1615 (*Eupomatidines, isol*)

Zjawiony, J.K. *et al.*, *J. Het. Chem.*, 1997, **34**, 1233-1237 (*3-Methoxysampangine, synth, ir, pmr, cmr*)


Kitahara, Y. *et al.*, *Tetrahedron*, 1997, **53**, 6001 (*Eupomatidines, synth*)

Akendengue, B. *et al.*, *Nat. Prod. Lett.*, 1999, **13**, 147-150 (*8-Hydroxysampangine*)

Sandramycin, 9CI

S-50

[100940-65-6]



C₆₀H₇₆N₁₂O₁₆ 1221.331

Depsipeptide antibiotic. Prod. by *Nocardioides* sp. ATCC 39419. Cytotoxic agent; moderately active against leukaemia, gram-positive bacteria and tumours; inhibits HIV-1 reverse transcriptase. Cryst. or powder + 8H₂O (MeOH/CHCl₃). Sol. EtOAc, C₆H₆; fairly sol. toluene, MeOH; poorly sol. H₂O, hexane. Mp 208-212°. λ_{\max} 210 (ε 76000); 228 (ε 71200); 306 (ε 9880); 356 (ε 10100) (MeOH/HCl) (Derep). λ_{\max} 246 (ε 73000); 301 (ε 8780); 395 (ε 11200) (MeOH/NaOH) (Derep). λ_{\max} 217 (ε 77700); 229 (ε 76600); 356 (ε 9880) (MeOH) (Derep).

Ger. Pat., 1985, 3 521 436; *CA*, **104**, 128220 (*isol, struct*)


Matson, J.A. *et al.*, *J. Antibiot.*, 1989, **42**, 1763-1767; 1993, **46**, 162-166 (*pmr, struct*)

Boger, D.L. *et al.*, *J.A.C.S.*, 1993, **115**, 11624; 1996, **118**, 1629; 2001, **123**, 561-568 (*synth, activity*)

Sandwicolidine

S-51

[99612-65-4]



Relative Configuration

C₂₁H₂₈N₂O₂ 340.464

Struct. is peculiarly shown and described. It appears to be an abeoajmaline. Alkaloid from the undried roots of *Rauwolfia serpentina* (Apocynaceae). Plates (EtOH/EtOAc). Mp 213-214°. [α]_D²⁰ +227 (CHCl₃).

Ac: Mp 241-242°.


Siddiqui, S. *et al.*, *Tetrahedron*, 1985, **41**, 4577 (*isol, uv, ir, pmr, cmr, ms, struct*)

Sandwicoline

S-52

4-Methyl-4,21-secosandwicine

[95753-42-7]



Absolute Configuration

C₂₁H₃₀N₂O₂ 342.48

Alkaloid from the undried winter roots of *Rauwolfia serpentina* (Apocynaceae). Elongated rods (MeOH). Mp 177-178°. [α]_D²⁰ +202 (CHCl₃).

Di-Ac:

Powder. Mp 203-204°.

Siddiqui, S. *et al.*, *Heterocycles*, 1985, **23**, 617 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)

Haider, S.I. *et al.*, *Indian J. Chem., Sect. B*, 1987, **26**, 1065 (*synth*)


Ahmad, S.S. *et al.*, *Synthesis*, 1988, 478 (*synth, uv, ir, pmr, ms*)

Sangivamycic acid

S-53

4-Amino-7-β-D-ribofuranosyl-7H-pyrrolo[2,3-d]pyrimidine-5-carboxylic acid, 8CI

[18418-00-3]



C₁₂H₁₄N₄O₆ 310.266

Cryst. (H₂O). Mp 238° dec.

Hydrochloride: [21090-38-0]

Plates (MeOH). Mp 236-238°.

Me ester: 5-(Methoxycarbonyl)tubercidin
[18440-68-1]
C₁₃H₁₆N₄O₆ 324.293
Isol. from the sponge *Jaspis johnstoni*. Cytotoxic agent. Cryst. (MeOH/CHCl₃). Sol. MeOH, CHCl₃; poorly sol. hexane. Mp 216-218°. λ_{max} 211; 232 (sh); 281 (MeOH).

Me ester, hydrochloride:
Prisms (MeOH). Mp 216-218°.

Amide: Sangivamycin. B 14437. BA 90912. NSC 65346. OS 1998. Antibiotic B 14437. Antibiotic BA 90912
[18417-89-5]
C₁₂H₁₅N₃O₅ 309.281
Produced by a strain of *Streptomyces rimosus*. Possesses cytotoxicity against HeLa cells and exhibits significant activity against leukaemia 1210 in mice. Mp 260°. [α]_D²⁶ -45.7 (c, 1 in 0.1M HCl). λ_{max} 229 (ε 8200); 278 (ε 15100) (EtOH).

► **LD₅₀** (mus, orl) 11 mg/kg. UY9355000
Nitrile: 4-Amino-7-β-D-ribofuranosyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile, 9CI. Toyocamycin. Cyanotubercidin. Niamycin B. Vengicide. E 212. Antibiotic 1037. Antibiotic E 212. Anhygroscopin B. Naritheracin
[606-58-6]
C₁₂H₁₃N₅O₄ 291.266
Prod. by *Streptomyces toyocaensis* and *Streptomyces fungicidicus*. Isol. from the sponge *Jaspis johnstoni*. Active against gram-positive organisms; shows some exp. antineoplastic activity. Needles or prisms + 1H₂O. Mp 243°. [α]_D²⁶ -55.6 (c, 1 in 0.1M HCl). λ_{max} 235 (ε 22100); 273 (ε 16300) (0.1N HCl) (Derep). λ_{max} 235 (ε 11600); 280 (ε 16600) (0.1N NaOH) (Derep). λ_{max} 207; 232 (ε 11060); 274 (sh); 278 (ε 15700); 289 (sh) (MeOH) (Derep).

► **LD₅₀** (mus, orl) 8 mg/kg. UY9100000
Nitrile, 5'-O-α-D-glucopyranoside: Toyocamycin 5'-α-D-glucopyranoside
[117456-79-8]
C₁₈H₂₃N₅O₉ 453.408
Isol. from *Tolypothrix tenuis* and *Plectonema radiosum*. Cytotoxic and antifungal agent. Amorph. Sol. H₂O. [α]_D²⁵ +2.1 (c, 0.03 in H₂O). λ_{max} 230 (ε 7400); 278 (ε 10000) (H₂O) (Derep).


5'-Deoxy, nitrile: 5'-Deoxytoyocamycin
[65562-55-2]
C₁₂H₁₃N₅O₃ 275.266
Prod. by *Streptomyces* sp. A14345. Cryst. + ½H₂O (H₂O). Mp 187-188°. λ_{max} 235 (ε 22100); 273 (ε 16300) (0.1N HCl) (Derep). λ_{max} 235 (ε 11600); 280 (ε 16600) (0.1N NaOH) (Derep). λ_{max} 207; 232 (ε 11060); 274 (sh); 278 (ε 15700); 289 (sh) (MeOH) (Derep).

Suzuki, S. et al., *J. Antibiot.*, Ser. A, 1961, **14**, 343
Aszalos, A. et al., *J. Antibiot.*, 1966, **19**, 285
Rao, K.V. et al., *J. Med. Chem.*, 1968, **11**, 939 (*Sangivamycin, isol. struct*)
Tolman, R.L. et al., *J.A.C.S.*, 1968, **90**, 524; 1969, **91**, 2102 (*Sangivamycin acid, Sangivamycin, Toyocamycin, struct, synth*)
U.S. Pat., 1969, 3 423 398; CA, **70**, 86268y (*Sangivamycin*)

Uematsu, T. et al., *J. Biol. Chem.*, 1970, **245**, 4365 (*biosynth*)
Japan. Pat., 1970, 70 19 638; CA, **73**, 108253f (*isol*)
Nichol, C.A. et al., *Handb. Exp. Pharmacol.*, 1975, **38**, 434 (*rev. pharmacol*)
Chenon, M.-T. et al., *J.A.C.S.*, 1975, **97**, 4627 (*nmr*)
Wang, Y. et al., *Carbohydr. Res.*, 1977, **59**, 449 (*5'-Deoxytoyocamycin*)
Prusiner, P. et al., *Acta Cryst. B*, 1978, **34**, 517 (*cryst struct, deriv*)
Bergstrom, D.E. et al., *J.O.C.*, 1981, **46**, 1423-1431 (*Methoxycarbonyltubercidin, synth*)
Stewart, J.B. et al., *J. Antibiot.*, 1988, **41**, 1048 (*deriv*)
Isono, K. et al., *J. Antibiot.*, 1988, **41**, 1711 (*rev*)
Osada, H. et al., *J. Antibiot.*, 1989, **42**, 102 (*props*)
Zabriskie, T.M. et al., *J. Nat. Prod.*, 1989, **52**, 1353-1356 (*Jaspis johnstoni constits*)
Isaac, B.G. et al., *J. Antibiot.*, 1991, **44**, 729 (*5'-Deoxytoyocamycin*)
Iimori, T. et al., *Tet. Lett.*, 1991, **32**, 7273 (*pmr, conformn*)
Sharma, M. et al., *Nucleosides Nucleotides*, 1993, **12**, 643 (*Toyocamycin, synth*)
Porcari, A.R. et al., *Nucleosides Nucleotides*, 1999, **18**, 153-159 (*Toyocamycin, synth*)
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SAU000; VGZ000

Sanguidimerine S-54

1,3-Bis(13,14-dihydro-13-methyl[1,3]-benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridin-14-yl)-2-propanone, 9CI. 1,3-Bis(8-hydrosanguinarinyl)acetone. 1,3-Bis(11-hydrosanguinarinyl)acetone



C₄₃H₃₂N₂O₉ 720.734
Antineoplastic agent.

(8R*,8'R*)-form
(+)-form. Alkaloid SC2
[41758-45-6]

Alkaloid from the rhizomes of *Sanguinaria canadensis*. Also obt. by base-catalysed condensation of Sanguinarine, S-56 with acetonedicarboxylic acid (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 174°. [α]_D²⁶ +21.2 (c, 0.5 in Py).


(8RS,8'SR)-form
meso-form. **Chelidimerine. Alkaloid CMI**
[39110-99-1]
Alkaloid from the rhizomes and roots of *Chelidonium majus*. Also obt. by base-cat. condensation of Sanguinarine, S-56 with acetonedicarboxylic acid (Papaveraceae). Cryst. (CHCl₃/MeOH). Mp 258-260°. Pharmacol. active isomer. Opt. inactive.

The N-oxide shows higher cytotoxic activity.

Kim, H.K. et al., *J. Pharm. Sci.*, 1969, **58**, 372 (*isol, uv, ir*)
Tin-Wa, M. et al., *J. Nat. Prod.*, 1970, **33**, 267; 1972, **35**, 87 (*isol, uv, ir, pmr, ms, cryst struct*)
Tin-Wa, M. et al., *J. Pharm. Sci.*, 1972, **61**, 1846 (*synth, uv, ms, struct*)

Sanguilutine S-55

2,3,7,8,10-Pentamethoxy-5-methylbenzo[c]phenanthridinium(1+), 9CI
[55950-34-0]



C₂₃H₂₄NO[⊕] 394.446
Struct. revised in 1978. Alkaloid from *Sanguinaria canadensis* (Papaveraceae). Sol. MeOH, CHCl₃; poorly sol. H₂O.


Chloride:
Golden-yellow needles (CHCl₃/EtOH). Mp 163-164° (137-138°, 140-145° dec.).

Pseudocyanide:
Prisms (CHCl₃/EtOH). Mp 232-233° dec.

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1960, **25**, 1667; 1968, **33**, 1619 (*isol, pmr*)
Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 3420 (*uv*)
Kessar, S.Y. et al., *Tet. Lett.*, 1977, 1459 (*synth*)
Ishii, H. et al., *Chem. Pharm. Bull.*, 1978, **26**, 864; 3252 (*struct, synth*)
Ishii, H. et al., *J.C.S. Perkin I*, 1984, 2283 (*synth, pmr*)
Hanaoka, M. et al., *J.C.S. Perkin I*, 1987, 677 (*synth, uv, pmr*)
Hanaoka, M. et al., *Chem. Pharm. Bull.*, 1989, **37**, 857 (*synth*)
Dostal, J. et al., *Phytochemistry*, 1998, **47**, 879-885 (*isol, struct*)
Marek, R. et al., *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)
Seckárová, P. et al., *Magn. Reson. Chem.*, 2002, **40**, 147-152 (*pmr, cmr*)

Sanguinarine S-56

13-Methyl[1,3]benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridinium(1+), 9CI. 5-Methyl-2,3,7,8-bis(methylenedioxy)benzo[c]phenanthridinium(1+). Pseudocheletrythrine. ψ-Cheletrythrine
[2447-54-3]



C₂₀H₁₄NO[⊕] 332.335
Several numbering systems have been used. Alkaloid from a wide variety of genera in the Papaveraceae (*Argemone, Boconia, Chelidonium, Dicterostigma, Eschscholtzia, Glaucium, Hunnemannia, Hylomecon, Hypocum, Macleaya, Me-*

conopsis, *Papaver*, *Platystemon*, *Romneya*, *Sanguinaria*, *Stylomecon*, *Stylophorum*, *Papaveraceae* (*Corydalis*, *Dicentra*, *Fumaria*), *Rutaceae* (*Zanthoxylum*), and *Sapindaceae* (*Pteridophyllum*). Potent cytotoxic agent functioning by DNA intercalation and uncoupling of oxidative phosphorylation. Inhibitor of alanine aminotransferase, acetylcholine transferase and of human plasma diamine oxidase. Adrenergic receptor antagonist. Antimicrobial and antiinflammatory agent showing antiplaque activity. Anaesthetic, antifungal and hepatotoxic agent. Has been used in toothpastes and oral rinses. Causes temporary change in intraocular pressure. Consumption of Sanguinarine, present in poppy seeds and in the oil of *Argemone mexicana* which has been used as an adulterant for mustard oil in India, has been linked to development of glaucoma. Banned by FDA. Sol. MeOH, CHCl₃. Log P 0.01 (uncertain value) (calc). λ_{max} 234; 284; 324 (MeOH) (Berdy). λ_{max} 234 (ε 31500); 283 (ε 33100); 325 (ε 14100) (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 22 mg/kg; LD₅₀ (mus, ipr) 18 mg/kg; LD₅₀ (mus, scu) 112 mg/kg. VP5200000

Chloride: *Sanguinarium chloride*, INN, *USAN*

[5578-73-4]

C₂₀H₁₄ClNO₄ 367.788

Orange needles (dil. HCl). Mp 277-280° dec. (273-274°, 284-289° dec.).

▶ VP5200000

Pseudocyanide:

C₂₁H₁₄N₂O₄ 358.353

Mp 244-245° (236-237°).

N-De-Me: [1,3]Benzodioxolo[5,6-c]-1,3-dioxolo[4,5-i]phenanthridine. 2,3:7,8-Bis(methylenedioxy)benzo[c]phenanthridine. *Norsanguinarine*

[522-30-5]

C₁₉H₁₁NO₄ 317.3

Alkaloid from *Corydalis incisa*, *Corydalis pallida*, *Dicentra peregrina*, *Argemone mexicana*, *Argemone albiflora*, *Chelidonium japonicum*, *Eschscholtzia californica*, *Glauclium flavum* var. *vestitum*, *Macleaya cordata*, *Papaver somniferum*, *Papaver setigerum*, *Papaver bracteatum*, *Papaver orientale*, *Papaver rhoeas* (corn poppy), *Pteridophyllum racemosum* and *Fumaria vaillantii* (Papaveraceae, Sapindaceae). Pale yellow needles (Me₂CO, CHCl₃/MeOH or CHCl₃/EtOH). Mp 285-287° (278-280° dec.).

12-Methoxy: see Chelirubine, C-360

8-Methoxy (?): see Pancorine, P-55

Späth, E. et al., *Ber.*, 1931, **64**, 370 (struct)

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1968, **33**, 1619 (ms, pmr)

Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 3420 (uv)

Sainsbury, M. et al., *J.C.S. (C)*, 1970, 1797 (synth, uv, ir)

Onda, M. et al., *Chem. Pharm. Bull.*, 1971, **19**, 31 (synth)

Furuya, T. et al., *Phytochemistry*, 1972, **11**, 3401 (*Norsanguinarine*)

Haisová, K. et al., *Coll. Czech. Chem. Comm.*, 1973, **38**, 3312; 1975, **40**, 1576

(*Norsanguinarine*)

Ikuta, A. et al., *Phytochemistry*, 1974, **13**, 2175; 1976, **15**, 577 (*Norsanguinarine*)

Battersby, A.R. et al., *J.C.S. Perkin 1*, 1975, 1140; 1147 (*isol, biosynth*)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1978, **26**, 864 (pmr)

Lenfeld, J. et al., *Planta Med.*, 1981, **43**, 161; 1983, **48**, 111 (*pharmacol*)

Maiti, M. et al., *FEBS Lett.*, 1982, **142**, 280 (*pharmacol*)

Takao, N. et al., *Helv. Chim. Acta*, 1983, **66**, 473 (*biosynth*)

Šmidrkal, J. et al., *Coll. Czech. Chem. Comm.*, 1984, **49**, 1412 (*synth*)

Šimánek, V. et al., *Alkaloids (Academic Press)*, 1985, **26**, 185 (rev)

Nagao, Y. et al., *Nucleic Acids Symp. Ser.*, 1985, **16**, 37 (*pharmacol*)

Hanaoka, M. et al., *Chem. Lett.*, 1986, 739 (*synth*)

Blaskó, G. et al., *Heterocycles*, 1988, **27**, 911 (*cmr*)

Martindale, The Extra Pharmacopoeia, 30th edn., *Pharmaceutical Press*, 1993, 1410

Marek, R. et al., *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Seckárová, P. et al., *Magn. Reson. Chem.*, 2002, **40**, 147-152 (pmr, cmr)

Dvorák, Z. et al., *Heterocycles*, 2006, **68**, 2403-2422 (rev)


Mackraj, I. et al., *Cardiovasc. Ther.*, 2008, **26**, 75-83 (rev)

Sax, N.I. et al., *Dangerous Properties of Industrial Materials*, 5th edn., *Van Nostrand Reinhold*, 1979, 964

Sanguinolentaquinone

S-57

[950842-11-2]



C₁₃H₁₆N₂O₄ 264.28

Isol. from the mushrooms *Mycena haematopus* and *Mycena sanguinolenta*. Red solid. λ_{max} 241 (log ε 3.95); 341 (log ε 3.81); 531 (log ε 2.79) (H₂O).


Peters, S. et al., *J. Nat. Prod.*, 2007, **70**, 1274-1277 (*isol, pmr, cmr, ms*)

Peters, S. et al., *Eur. J. Org. Chem.*, 2008, 319-323 (*isol*)

Sanguinone A†

S-58

[950842-09-8]



R = H

Absolute Configuration

C₁₅H₁₄N₄O₃ 298.301

Isol. from mushroom *Mycena sanguinolenta*. Blue solid. [α]_D²⁵ -45 (c, 0.01 in H₂O). λ_{max} 251 (log ε 3.9); 376 (log ε 3.73); 578 (log ε 2.64) (H₂O).

4-Decarboxy, 3,4-didehydro: Decarboxy-didehydrosanguinone A. Decarboxydehydrosanguinone A

[949005-10-1]

C₁₄H₁₂N₄O 252.275

Yellow solid. Artifact. Oxidative decarboxylation product. λ_{max} 231 (log ε 3.32); 252 (log ε 3.19); 428 (log ε 3.05); 452 (log ε 3.25) (H₂O).

Peters, S. et al., *J. Nat. Prod.*, 2007, **70**, 1274-1277 (*isol, cd, pmr, cmr, ms*)

Sanguinone B

S-59

[950842-10-1]

As Sanguinone A, S-58 with

R = CH₃

C₁₆H₁₆N₄O₃ 312.327

Isol. from mushroom *Mycena sanguinolenta*. Blue solid. λ_{max} 247 (log ε 3.86); 374 (log ε 3.75); 571 (log ε 2.59) (H₂O).


Peters, S. et al., *J. Nat. Prod.*, 2007, **70**, 1274-1277 (*isol, cd, pmr, cmr, ms*)

Sanguirubine

S-60

5,9,10-Trimethoxy-12-methylbenzo[c]-1,3-dioxolo[4,5-i]phenanthridinium(1+), 9Cl. 2,3,10-Trimethoxy-5-methyl-7,8-methylenedioxybenzo[c]phenanthridinium(1+)

[55950-33-9]



C₂₂H₂₀NO₅⁺ 378.404

Alkaloid from *Sanguinaria canadensis* (Papaveraceae). Sol. MeOH, CHCl₃; poorly sol. H₂O.

Chloride:

C₂₂H₂₀ClNO₅ 413.856

Red needles (CHCl₃/EtOH or H₂O). Mp 281-284° Mp 275-276°.

Pseudocyanide:

C₂₃H₂₀N₂O₅ 404.421

Prisms (CHCl₃/EtOH). Mp 237-238°.

Slavk, J. et al., *Coll. Czech. Chem. Comm.*, 1960, **25**, 1667; 1968, **33**, 1619 (*isol, pmr*)

Hruban, L. et al., *Coll. Czech. Chem. Comm.*, 1970, **35**, 3420 (uv)


Ishii, H. et al., *Chem. Pharm. Bull.*, 1978, **26**, 864; 3252 (*struct, synth*)

Ishii, H. et al., *J.C.S. Perkin 1*, 1984, 2283 (*synth, pmr*)

Seckárová, P. et al., *Magn. Reson. Chem.*, 2002, **40**, 147-152 (pmr, cmr)

Sanjoinine G₁

[107462-36-2]

Absolute
ConfigurationC₃₁H₄₄N₄O₅ 552.712

Isol. from the seeds of *Zizyphus vulgaris* var. *spinosus*. Mp 236-238°. $[\alpha]_D^{20}$ -68.6 (c, 0.175 in CHCl₃). λ_{\max} 232 (log ϵ 3.97); 278 (log ϵ 3.3) (No solvent reported).

Me ether: Sanjoinine D


[107462-34-0]

C₃₂H₄₆N₄O₅ 566.739

Isol. from the seeds of *Zizyphus vulgaris* var. *spinosus*. Needles (CHCl₃/MeOH). Mp 256-258°. $[\alpha]_D^{26}$ -53.6 (c, 0.25 in CHCl₃). λ_{\max} 231 (log ϵ 3.82); 279 (log ϵ 2.82) (MeOH).

Han, B.H. *et al.*, *Phytochemistry*, 1990, **29**,3315-3319 (*isol, pmr, cmr, ms*)Han, B.H. *et al.*, *Heterocycles*, 1995, **41**, 1909(*synth*)Park, M.H. *et al.*, *Phytochemistry*, 1996, **43**,701-704 (*isol, uv, cd, pmr, abs config*)East, S.P. *et al.*, *Tetrahedron*, 1998, **54**, 13371-13390 (*synth*)Temal-Laib, T. *et al.*, *J.A.C.S.*, 2002, **124**, 583-590 (*synth*)Sanjoinine G₂

[107494-21-3]

C₃₀H₄₂N₄O₅ 538.686

A seco-ansapeptide alkaloid. Alkaloid from Sanjoin (the seeds of *Zizyphus vulgaris* var. *spinosus* (Rhamnaceae)). Needles (Me₂CO aq.). Mp 182°. $[\alpha]_D$ -79.2. λ_{\max} 284 (log ϵ 3.81) (no solvent reported).

Han, B.H. *et al.*, *Pure Appl. Chem.*, 1989, **61**,443-448 (*isol, struct*)Han, B.H. *et al.*, *Phytochemistry*, 1990, **29**,3315-3319 (*isol, ir, uv, pmr, cmr*)

Sankhpuspine

C₁₇H₂₃NO₃ 289.374


Struct. unknown. Alkaloid from *Convolvulus pluricaulis* (Convolvulaceae). Cryst. (EtOH). Mp 162-164°.

Picrate: Mp 76° dec.*Aurichloride*: Mp 208° dec.*Chloroplatinate*: Mp 180° dec.Basu, N.K. *et al.*, *J. Am. Pharm. Assoc.*, 1948, **37**, 27-28 (*isol*)

S-61

Santiagonamine

[92664-91-0]


C₁₉H₁₈N₂O₃ 322.363

Alkaloid from the stems and branches of *Berberis darwinii* (Berberidaceae). The first known phenanthridine alkaloid.

Valencia, E. *et al.*, *Tet. Lett.*, 1984, **25**, 3163(*uv, ir, pmr, ms, struct*)Markey, M.D. *et al.*, *Org. Lett.*, 2007, **9**, 3255-3257 (*synth*)

Santiaguine

[528-31-4]

(2*R*,2'*R*)-formC₃₈H₄₈N₄O₂ 592.823

α -Truxillate dimer of Adenocarpine, A-138. The truxillate residue is *meso*-, so there are four possible isomers; 2 enantiomers, one (\pm)-form and one *meso*-. All have been found in nature.

(2*R*,2'*R*)-form [11016-82-3]

Alkaloid from *Adenocarpus commutatus* and *Adenocarpus grandiflorus* (Fabaceae). Mp 240-245°. $[\alpha]_D^{20}$ -18.42 (c, 9.6 in 85% HCOOH aq.).

Hydrochloride (1:2): [11016-90-3] $[\alpha]_D^{18}$ -16.66 (c, 3.96 in 85% HCOOH).(2*S*,2'*S*)-form [11016-83-4]

Alkaloid from *Adenocarpus complicatus*, *Adenocarpus intermedius*, *Adenocarpus foliosus* and *Adenocarpus viscosus* (Fabaceae). Cryst. (EtOH). Mp 235-236°. $[\alpha]_D^{17}$ +3.3 (c, 3.68 in EtOH).

Hydrochloride (1:2): [11016-92-5]Cryst. + 1H₂O (H₂O); cryst. (EtOH). Mp 238° (anhyd. 241°).*Perchlorate* (1:2): [11017-00-8]

Mp 244-245° dec.

Picrate: Mp 285° (explodes, prior blackening).(2*R**S*,2'*R**S*)-form(\pm)-form

S-64

[11016-84-5]

Alkaloid from *Adenocarpus commutatus*, *Adenocarpus grandiflorus*, *Adenocarpus foliosus* and *Adenocarpus viscosus* (Fabaceae). Cryst. + 1H₂O. Mp 237-239° dec.

Hydrochloride (1:2): [11016-93-6]Cryst. + 2H₂O. Mp 245-250° dec.*Perchlorate* (1:2):Cryst. + 1½ H₂O. Mp 252-254° dec.(2*R**S*,2'*S*R)-form*meso*-form

[11016-85-6]


Alkaloid from *Adenocarpus commutatus* (Fabaceae). Cryst. + 3H₂O (MeOH aq.). Mp 238-240° dec.

Hydrochloride (1:2): [11016-91-4]Cryst. + 3½ H₂O (EtOH aq.). Mp 245-247° dec.*Perchlorate* (1:2): [11017-01-9]Cryst. + 1½ H₂O. Mp 245-248° dec.

[11016-88-9, 11016-89-0, 11016-86-7, 11016-96-9, 11016-95-8, 11016-87-8, 11016-97-0, 11016-94-7]

Ribas, I. *et al.*, *An. R. Soc. Esp. Fis. Quim.*,*Ser. B*, 1950, **46**, 489; 1951, **47**, 730;1953, **49**, 783; 1956, **52**, 133; 1958, **54**, 157;215; 1966, **62**, 845 (*isol, struct, bibl, abs config*)Ribas, I. *et al.*, *Ann. Pharm. Fr.*, 1952, **10**, 54(*isol*)Bernasconi, R. *et al.*, *Pharm. Acta Helv.*, 1970,**45**, 421 (*isol*)O'Donovan, D.G. *et al.*, *J.C.S. Perkin I*, 1974,2524 (*biosynth*)Nehme, M. *et al.*, *An. Quim.*, 1977, **73**, 307(*isol*)

Sarachine


22,26-Iminocholest-5-en-3-amine
[214706-01-1]C₂₇H₄₆N₂ 398.674

Alkaloid from *Saracha punctata*. Anti-protozoal agent. Greenish powder. $[\alpha]_D^{25}$ -7 (c, 0.3 in CHCl₃). $[\alpha]_D^{25}$ -13 (c, 0.3 in MeOH). Only relative config. of C-22:C-25 is known.

Moretti, C. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1390-1393 (*isol, pmr, cmr, ms*)

Sarain A

[123117-93-1]

Absolute
Configuration

n = 1

S-67

$C_{32}H_{51}N_2O_3^{\oplus}$ 511.767

Alkaloid from the sponge *Reniera sarai*. Vasodilator, insecticide. Amorph. powder. $[\alpha]_D^{20} +66$ (c, 1.3 in $CHCl_3$). The title of the 2007 paper was 'Synthesis of (-)-Sarain A'. This was an error and should be (+)-Sarain A. A correction is in press March 2008. λ_{max} 238 (ε 14939) (MeOH).

13,14-Didehydro(Z-), n = 2: **Sarain B**

[123391-00-4]

$C_{33}H_{51}N_2O_3^{\oplus}$ 523.778

Alkaloid from *Reniera sarai*.

Insecticide, acaricide. Toxic to sea urchins. Powder. $[\alpha]_D^{20} +76.3$ (c, 1.2 in $CHCl_3$). λ_{max} 238 (ε 13415) (MeOH).

[123117-96-4, 123391-01-5, 123117-94-2]

Cimino, G. *et al.*, *Tetrahedron*, 1989, **45**, 3863-3872 (*isol, uv, ir, pmr, cmr, cryst struct*)

Cimino, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1519-1525 (*uv, ir, pmr, cmr, ms*)

Guo, Y. *et al.*, *Tetrahedron*, 1996, **52**, 8341-8348 (*pmr, cmr, abs config*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (*rev*)

Becker, M.H. *et al.*, *J.A.C.S.*, 2007, **129**, 11987-12002 (*synth, bibl*)

Sarain C

S-68

[123089-18-9]

As Sarain A, S-67 with

n = 2

$C_{33}H_{53}N_2O_3^{\oplus}$ 525.793

Alkaloid from the sponge *Reniera sarai*. Insecticide, acaricide, toxic to sea urchins. Amorph. powder. $[\alpha]_D^{20} +67.4$ (c, 1 in $CHCl_3$). λ_{max} 238 (ε 13582) (MeOH).

[123089-19-0]

Cimino, G. *et al.*, *Tetrahedron*, 1989, **45**, 3863 (*isol, uv, ir, pmr, cmr*)


Caprioli, V. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1992, **103**, 293 (*activity*)

Guo, Y. *et al.*, *Tetrahedron*, 1996, **52**, 8341 (*abs config*)

Saraine 1

S-69

[105418-77-7]



Absolute Configuration

X = $-\text{CH}=\text{CH}-\text{Z}$

$C_{31}H_{50}N_2O$ 466.749

Alkaloid from the marine sponge *Reniera sarai*. Antiarrhythmic, ion chelator, insecticide. Amorph. powder. $[\alpha]_D^{20} -47.8$ (c, 1.2 in $CHCl_3$).

► LD₅₀ (mus, ipr) 200 mg/kg.

1,2,9-Triepimer: **Isosaraine 1**

[119766-95-9]

$C_{31}H_{50}N_2O$ 466.749

Minor alkaloid from the Mediterranean sponge *Reniera sarai*. Shows insecticidal props. Amorph. solid. $[\alpha]_D^{20} -23.1$ (c, 1.2 in $CHCl_3$).

Cimino, G. *et al.*, *Bull. Soc. Chim. Belg.*, 1986, **95**, 783-800 (*isol, pmr, cmr, biosynth*)

Cimino, G. *et al.*, *Tet. Lett.*, 1989, **30**, 133-136 (*isol, pmr, cmr, ms, struct*)

Caprioli, V. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1992, **103**, 293-296 (*biochem*)

Guo, Y. *et al.*, *Tetrahedron*, 1996, **52**, 14961-14974 (*cmr*)

Guo, Y. *et al.*, *Tet. Lett.*, 1998, **39**, 463-466 (*abs config*)

Saraine 2

S-70

[105418-80-2]

As Saraine 1, S-69 with

X = $-\text{CH}_2-$

$C_{30}H_{50}N_2O$ 454.738

Alkaloid from the marine sponge *Reniera sarai*. Antiarrhythmic, ion-chelator, insecticide. Amorph. powder. $[\alpha]_D^{20} -117$ (c, 1.1 in $CHCl_3$).

1,2,9-Triepimer: **Isosaraine 2**

[135091-28-0]

$C_{30}H_{50}N_2O$ 454.738

Alkaloid from the marine sponge *Reniera sarai*. Oil. $[\alpha]_D^{20} -34.6$ (c, 2 in $CHCl_3$).

Cimino, G. *et al.*, *Bull. Soc. Chim. Belg.*, 1986, 783-800 (*isol, pmr, cmr, biosynth*)

Cimino, G. *et al.*, *Tet. Lett.*, 1989, **30**, 133-136 (*struct*)

Cimino, G. *et al.*, *Magn. Reson. Chem.*, 1991, **29**, 327-332 (*isol, pmr, cmr, struct*)

Caprioli, V. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1992, **103**, 293-296 (*biochem*)

Guo, Y. *et al.*, *Tetrahedron*, 1996, **52**, 14961-14974 (*abs config, cmr*)

Guo, Y. *et al.*, *Tet. Lett.*, 1998, **39**, 463-466 (*config*)

Saraine 3

S-71

[105305-54-2]

As Saraine 1, S-69 with

X = $-\text{CH}=\text{CH}-\text{CH}_2-$

$C_{32}H_{52}N_2O$ 480.776

Alkaloid from the marine sponge *Reniera sarai*. Antiarrhythmic, ion-chelator, insecticide. Amorph. powder. $[\alpha]_D^{20} -27.4$ (c, 0.8 in $CHCl_3$).

1,2,9-Triepimer: **Isosaraine 3**

[184429-62-7]

$C_{32}H_{52}N_2O$ 480.776

Alkaloid from the sponge *Reniera sarai*. Amorph. powder. $[\alpha]_D^{20} -16.3$ (c, 1.34 in $CHCl_3$).

Cimino, G. *et al.*, *Bull. Soc. Chim. Belg.*, 1986, **95**, 783-800 (*isol, pmr, cmr, biosynth*)

Cimino, G. *et al.*, *Tet. Lett.*, 1989, **30**, 133-136 (*struct*)

Caprioli, V. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1992, **103**, 293-296 (*biochem*)


Guo, Y. *et al.*, *Tetrahedron*, 1996, **52**, 14961-14974 (*pmr, cmr, struct*)

Guo, Y. *et al.*, *Tet. Lett.*, 1998, **39**, 463-466 (*abs config*)

Sarcinapterin

S-72

[89157-12-0]



$C_{35}H_{48}N_7O_{19}P$ 901.773

Isol. from *Methanosarcina barkeri*. Co-factor in methanogenic bacteria. Solid.


Van Beelen, P. *et al.*, *Eur. J. Biochem.*, 1984, **139**, 359 (*isol, uv, pmr*)

Gorris, L.G. *et al.*, *Prog. Biotechnol.*, 1986, **2**, 144; *CA*, **106**, 210634t (*props*)

Sarcocapnidine

S-73

2,3,12,12a-Tetrahydro-6,9-dimethoxy-1-methyl-1H-[1]-benzoxepino[2,3,4-ij]isoquinolin-8-ol, 9CI
[87069-33-8]



$C_{19}H_{21}NO_4$ 327.379

Biogenetic (benzylisoquinoline-type) numbering shown. Alkaloid from *Sarcocapnos crassifolia* and *Corydalis claviculata* (Papaveraceae). Cryst. (EtOH). Mp 126-127°. $[\alpha]_D^{20} +385.4$ (c, 0.07 in MeOH).

N- α -Oxide: **cis-Sarcocapnidine N-oxide**
[122621-72-1]

$C_{19}H_{21}NO_5$ 343.379

Alkaloid from aerial parts of *Sarcocapnos baetica* subsp. *integrifolia* (Papaveraceae). Amorph. powder. $[\alpha]_D^{20} +430$ (c, 0.20 in MeOH).

N-De-Me: **Norsarcocapnidine**

$C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Sarcocapnos crassifolia*. Amorph. solid. $[\alpha]_D^{25} +347.5$ (c, 0.8 in EtOH).

Me ether: **Sarcocapnine**

[81874-23-9]

$C_{20}H_{23}NO_4$ 341.406

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Cryst. (EtOH/Et₂O) (as hydrochloride). Mp 213-215° (hydrochloride). $[\alpha]_D^{25} +218$ (c, 0.3 in EtOH).

Me ether, N- α -oxide: **cis-Sarcocapnine N-**

oxide

[122555-13-9]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Ceratocarpus heterocarpa*. Amorph. powder. Mp 110-114°. [α]_D²⁵ +174 (c, 0.065 in MeOH). λ_{\max} 230 (sh) (log ϵ 4.08); 284 (log ϵ 3.4) (MeOH).

Me ether, N-de-Me: Norsarcocapnine

[81855-38-1]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Ceratocarpus heterocarpa*. Pale yellowish cryst. Mp 110°. [α]_D²⁵ +165 (c, 0.06 in MeOH). λ_{\max} 206 (log ϵ 4.68); 228 (sh) (log ϵ 4.26); 280 (log ϵ 3.61) (MeOH).

4R-Hydroxy: 4-Hydroxysarcocapnine

[120186-23-4]

C₁₉H₂₁NO₅ 343.379

Alkaloid from aerial parts of *Sarcocapnos baetica* subsp. *integrifolia* (Papaveraceae). Amorph. solid. [α]_D²⁵ +157 (c, 0.13 in CHCl₃).

4R-Hydroxy, Me ether: 4-Hydroxysarcocapnine

[95377-94-9]

C₂₀H₂₃NO₅ 357.405

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Cryst. (EtOH). Mp 145-146°. [α]_D²⁵ +311 (c, 0.11 in CHCl₃). λ_{\max} 218 (log ϵ 4.26); 230 (sh) (log ϵ 4.09); 282 (log ϵ 3.46) (EtOH).

 α -Oxo, 1,2,3,4-tetrahydro, N-de-Me: 8-Hydroxy-6,9-dimethoxy-12H-[1]benzoxepino[2,3,4-ij]isoquinolin-12-one, 9CI. Oxosarcocapnine

[87035-68-5]

C₁₈H₁₃NO₅ 323.304

Alkaloid from *Sarcocapnos crassifolia* (Papaveraceae). Yellow cryst. (MeOH). Mp 231-232°.

 α -Oxo, 1,2,3,4-tetrahydro, Me ether, N-de-Me: 6,8,9-Trimethoxy-12H-[1]benzoxepino[2,3,4-ij]isoquinolin-12-one, 9CI. Oxosarcocapnine

[81855-39-2]

C₁₉H₁₅NO₅ 337.331

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Yellow cryst. (EtOH). Mp 202-203°.

Campello, M.J. *et al.*, *Tet. Lett.*, 1982, **23**, 239 (isol, uv, pmr, ms, struct, abs config, Sarcocapnine, Oxosarcocapnine)

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2303 (uv, ir, pmr, cmr, ms, struct, Sarcocapnine, Oxosarcocapnine)

Blaschke, G. *et al.*, *Z. Naturforsch., C*, 1983, **38**, 670

Castedo, L. *et al.*, *Tet. Lett.*, 1984, 4573-4576 (4-Hydroxysarcocapnine)

Castedo, L. *et al.*, *Heterocycles*, 1987, **26**, 29; 1988, **27**, 2783 (Norsarcocapnine, 4-Hydroxysarcocapnine, Sarcocapnine N-oxide)

Suau, R. *et al.*, *Phytochemistry*, 1989, **28**, 3511 (Norsarcocapnine)


de Sousa, J.D.F. *et al.*, *J.A.C.S.*, 1994, **116**, 9745 (synth, Sarcocapnine, Sarcocapnine)

Suau, R. *et al.*, *Phytochemistry*, 1996, **43**, 1389 (Sarcocapnine N-oxide, Sarcocapnine N-oxide)

Rodrigues, J.A.R. *et al.*, *J.O.C.*, 2004, **69**, 2920-2928 (synth)

Sarcodifurine A

S-74



Relative Configuration


C₂₃H₂₇NO₅ 397.47

Alkaloid from the leaves of *Sarcomelicope follicularis*. Yellow needles (MeOH). Mp 155-157°. [α]_D²⁵ -27 (c, 0.6 in MeOH). λ_{\max} 244; 255 (sh); 272; 320 (MeOH).

Chosson, E. *et al.*, *Heterocycles*, 2004, **63**, 2043-2050 (isol, pmr, cmr, cryst struct)

Sarcodifurine B

S-75



Relative Configuration

C₂₃H₂₇NO₅ 397.47


Alkaloid from the leaves of *Sarcomelicope follicularis*. Yellow needles (MeOH). Mp 166-167°. [α]_D²⁵ +99 (c, 0.6 in MeOH). λ_{\max} 258; 267; 276; 354 (MeOH).

Chosson, E. *et al.*, *Heterocycles*, 2004, **63**, 2043-2050 (isol, pmr, cmr, cryst struct)

Sarcodonin

S-76

[263846-30-6]



Absolute Configuration

C₃₆H₃₈N₂O₁₄ 722.701

Isol. from the fruiting bodies of the basidiomycete *Sarcodon leucopus*. Moderate cytotoxic agent. Amorph. powder. [α]_D²⁵ -20.4 (c, 1.6 in CHCl₃) (as tetra-Ac). λ_{\max} 202 (ϵ 40450); 256 (ϵ 15230) (EtOH).

7-Ac: Sarcodonin γ

[679001-56-0]

C₃₈H₄₀N₂O₁₅ 764.738

Isol. from *Sarcodon leucopus*. Amorph. solid. [α]_D²⁵ -11.7 (c, 0.24 in CHCl₃).

4''-O-De-Ac: Sarcodonin α

[679001-53-7]

C₃₄H₃₆N₂O₁₃ 680.664

Isol. from *Sarcodon leucopus*. Amorph. powder. [α]_D²⁵ -9.9 (c, 1.2 in CHCl₃). λ_{\max} 210 (log ϵ 4.66); 270 (log ϵ 4.24) (EtOH).

4''-O-De-Ac, 5',6'-di-Ac: Sarcodonin β

[679001-55-9]

C₃₈H₄₀N₂O₁₅ 764.738

Isol. from *Sarcodon leucopus*. Amorph. powder. [α]_D²⁵ -41.7 (c, 0.12 in CHCl₃).

N-Deoxy, N-methoxy, 4''-O-de-Ac: Sarcodonin δ

[870773-03-8]

C₃₅H₃₈N₂O₁₃ 694.691

Isol. from *Hydnellum geogerirum*, *Hydnellum suaveolens* and *Sarcodon scabrosus*. Amorph. red powder. [α]_D²⁰ -2.7 (c, 0.1 in MeOH). λ_{\max} 205 (log ϵ 4.26); 260 (log ϵ 4.79) (MeOH).

5',6'-Quinone, tri-de-Ac: Sarcoviolin α

[679001-64-0]

C₃₀H₃₀N₂O₁₁ 594.574

Isol. from the fruiting bodies of the basidiomycete *Sarcodon leucopus*. Cytotoxic. Amorph. violet solid. Obt. as a 2:1 mixt. with Episcarvicolin α to which the data refers. λ_{\max} 290 (log ϵ 4.08); 350 (log ϵ 3.52); 540 (log ϵ 2.71) (MeOH).

8-Epimer: Episcarvicolin

[679001-58-2]

C₃₆H₃₈N₂O₁₄ 722.701

Isol. from *Sarcodon leucopus*. Amorph. solid. [α]_D²⁵ -5.4 (c, 0.4 in MeOH).

8-Epimer, 4''-O-de-Ac: Episcarvicolin α

[679001-59-3]

C₃₄H₃₆N₂O₁₃ 680.664

Isol. from *Sarcodon leucopus*. Amorph. solid. [α]_D²⁵ -1.8 (c, 0.55 in CHCl₃). λ_{\max} 210 (log ϵ 4.71); 270 (log ϵ 4.27) (EtOH).

8-Epimer, 4''-O-de-Ac, 5',6'-di-Ac: Episcarvicolin β

[679001-61-7]

C₃₈H₄₀N₂O₁₅ 764.738

Isol. from *Sarcodon leucopus*. Amorph. powder. [α]_D²⁵ -28 (c, 0.23 in CHCl₃).

8-Epimer, N-methoxy, 4''-O-de-Ac: Hydrellin A

[910099-73-9]

C₃₅H₃₈N₂O₁₃ 694.691

Isol. from *Hydnellum suaveolens*. [α]_D²⁰ -125.1 (c, 1 in CHCl₃). λ_{\max} 257 (log ϵ 4.2) (EtOH).

8-Epimer, 5',6'-quinone, tri-de-Ac: Episcarvicolin α

[679001-65-1]

C₃₀H₃₀N₂O₁₁ 594.574

Isol. from *Sarcodon leucopus*.

Geraci, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 347-351 (Sarcodonin, isol, pmr, cmr)


Cali, V. *et al.*, *Eur. J. Org. Chem.*, 2004, 592-599 (Sarcodonins α, β, γ , Episcarvicolin, Sarcoviolin α , Episcarvicolin α)

Ma, B.-J. *et al.*, *Z. Naturforsch., B*, 2005, **60**, 565-568 (Sarcodonin δ)

Hashimoto, T. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 912-914 (Hydnellin A)

Sarcomejine

S-77

C₁₆H₁₇NO₆ 319.313


Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow powder. $[\alpha]_D^{25} +3$ (c, 0.1 in CH_2Cl_2). λ_{max} 290 (sh); 327 (log ϵ 3.67); 341 (log ϵ 3.73) (MeOH).

Fokialakis, N. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1004-1005

Sarcomejine B

S-78

[939403-86-8]



$\text{C}_{18}\text{H}_{19}\text{NO}_7$ 361.351


Alkaloid from the bark of *Sarcomelicope megistophylla*. λ_{max} 297 (sh); 314 (sh); 326 (sh); 337 (log ϵ 3.32) (MeOH).

Mitaku, S. *et al.*, *Fitoterapia*, 2007, **78**, 169-170 (isol, pmr, cmr, ms)

Sarcopharyngine

S-79

Methyl 19-(hydroxymethyl)-12-methoxy-yibogamine-18-carboxylate, 9CI [110011-72-8]




$\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_4$ 398.501

Alkaloid from the seeds of *Sarcopharyngia crassa* (preferred genus name *Tabernaemontana*) (Apocynaceae). $[\alpha]_D^{22} -38$.

Batchily, F. *et al.*, *Ann. Pharm. Fr.*, 1986, **44**, 449

Sarcophylline

S-80



$\text{C}_{19}\text{H}_{21}\text{NO}_4$ 327.379

Biogenetic (benzylisoquinoline-type) numbering shown.

(S)-form

Alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Amorph. $[\alpha]_D^{25} +200$ (c, 1.65 in CHCl_3).

O¹¹-De-Me: Clavicine

[87035-67-4]

Alkaloid from *Sarcocapnos crassifolia* (Papaveraceae). Cryst. (EtOH). Mp 126° (112-113°). $[\alpha]_D +443$ (c, 0.41 in

(MeOH).

Boente, J.M. *et al.*, *Tet. Lett.*, 1983, **24**, 2303 (Clavicine)


Blaschke, G. *et al.*, *Phytochemistry*, 1985, **24**, 585-588 (Clavicine)

Castedo, L. *et al.*, *Heterocycles*, 1987, **26**, 29-33 (uv, ir, pmr, ms, struct)

Sarcotragin A

S-81

[346583-99-1]



$\text{C}_{31}\text{H}_{43}\text{NO}_4$ 493.685

Alkaloid from the sponge *Sarcotragus* sp. Gum. $[\alpha]_D^{25} +16$ (c, 0.11 in MeOH). λ_{max} 204 (log ϵ 4.1); 236 (log ϵ 4.07) (MeOH).

N-De-(2-phenylethyl), N-(carboxymethyl): **Sarcotragin B**

[346584-00-7]

 $\text{C}_{25}\text{H}_{37}\text{NO}_6$ 447.57


Alkaloid from *Sarcotragus* sp. Gum (as Na salt). $[\alpha]_D^{25} +17.5$ (c, 0.12 in MeOH) (Na salt). λ_{max} 209 (log ϵ 4.12); 232 (log ϵ 4.15) (MeOH) (Na salt).

Shin, J. *et al.*, *Tet. Lett.*, 2001, **42**, 3005-3007 (isol, pmr, cmr)

Sarcotrine A

S-82

[472974-46-2]



$\text{C}_{30}\text{H}_{45}\text{NO}_4$ 483.69

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +36.1$ (c, 0.18 in MeOH). λ_{max} 240 (log ϵ 4.81) (MeOH).

N-Dealkyl, N-(2-methylbutyl): **Sarcotrine C**

[472974-48-4]

 $\text{C}_{30}\text{H}_{45}\text{NO}_4$ 483.69

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +19.1$ (c, 0.18 in MeOH). λ_{max} 245 (log ϵ 4.28) (MeOH).

N-Dealkyl, N-(2-phenylethyl): **Sarcotrine B**

[472974-47-3]

 $\text{C}_{33}\text{H}_{43}\text{NO}_4$ 517.707

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +36.9$ (c, 0.07 in MeOH).

N-Dealkyl, N-(carboxymethyl): **Sarcotrine E**

[627895-36-7]

 $\text{C}_{27}\text{H}_{37}\text{NO}_6$ 471.592

Constit. of *Sarcotragus* sp. Oil. $[\alpha]_D^{21} +38.8$ (c, 0.18 in MeOH).

17-Epimer: **Episarcotrine A**

[471278-72-5]

 $\text{C}_{30}\text{H}_{45}\text{NO}_4$ 483.69

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +42.8$ (c, 0.05 in MeOH). λ_{max} 242 (log ϵ 4.56) (MeOH).

17-Epimer, N-dealkyl, N-(2-methylbu-

tyl): **Episarcotrine C**

[472974-54-2]

 $\text{C}_{30}\text{H}_{45}\text{NO}_4$ 483.69

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D +24$ (c, 0.06 in MeOH).

17-Epimer, N-dealkyl, N-(2-phenylethyl):

Episarcotrine B

[471278-73-6]

 $\text{C}_{33}\text{H}_{43}\text{NO}_4$ 517.707

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +42.1$ (c, 0.04 in MeOH).


Liu, Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1307-1314 (isol, pmr, cmr)

Liu, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1451-1456 (*Sarcotrine E*)

Sarcotrine D

S-83

[472974-56-4]



$\text{C}_{30}\text{H}_{45}\text{NO}_4$ 483.69

Constit. of a *Sarcotragus* sponge. Oil. $[\alpha]_D^{21} +65.2$ (c, 0.01 in MeOH). λ_{max} 239 (log ϵ 4.34) (MeOH).

N-Dealkyl, N-(carboxymethyl): **Isosarcotrine E**

[627895-37-8]

 $\text{C}_{27}\text{H}_{37}\text{NO}_6$ 471.592

Constit. of a *Sarcotragus* sp. Oil. $[\alpha]_D^{21} +33.3$ (c, 0.09 in MeOH).


Liu, Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1307-1314 (*Sarcotrine D*)

Liu, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1451-1456 (*Isosarcotrine E*)

Sarcotrine F

S-84

[627895-38-9]



$\text{C}_{26}\text{H}_{39}\text{NO}_6$ 461.597

Constit. of a *Sarcotragus* sp. Oil. Isol. as a mixt. with Isosarcotrine F.

3-Deoxo, 5-oxo: **Isosarcotrine F**

[627895-39-0]

 $\text{C}_{26}\text{H}_{39}\text{NO}_6$ 461.597


Constit. of a *Sarcotragus* sp. Oil. Stereochem. need not necessarily correspond.

Liu, Y. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1451-1456 (isol, pmr, cmr)

Sargassumlactam

S-85

[75956-49-9]



$\text{C}_9\text{H}_{11}\text{NO}_6$ 229.189


Isol. from the marine brown alga *Sar-*

gassum kjellmanianum. Mp 140-141.5°. $[\alpha]_D^0$ 0. λ_{\max} 208 (ϵ 6100); 293 (ϵ 3400) (EtOH) (Derep).

Nozaki, H. *et al.*, *Chem. Lett.*, 1980, 1453 (*uv, ir, pmr, cmr, cryst struct*)

Sarmetamide A S-86

1,5-Dihydro-1-(1-oxo-3-phenylpropyl)-2H-pyrrol-2-one. N-(3-Phenylpropanoyl)-A³-pyrrolidone




C₁₃H₁₃NO₂ 215.251

Constit. of the roots of *Piper sarmentosum*. Oil. λ_{\max} 209 (log ϵ 4.23); 230 (sh) (log ϵ 3.89) (MeOH).

Tuntiwachwuttikul, P. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 149-151 (*isol, pmr, cmr*)

Sarmetamide B S-87

3,4-Diacetoxy-N-cinnamoylpyrrolidine



Relative Configuration

C₁₇H₁₉NO₅ 317.341

Constit. of the roots of *Piper sarmentosum*. Wax. $[\alpha]_D^{25}$ +68.3 (c, 0.12 in MeOH). λ_{\max} 206 (log ϵ 4.27); 217 (log ϵ 4.21); 233 (sh) (log ϵ 4.08); 258 (log ϵ 3.78); 283 (log ϵ 4.25) (MeOH).

Tuntiwachwuttikul, P. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 149-151 (*isol, pmr, cmr*)

Sarothamine S-88

[1361-10-0]

C₃₀H₅₀N₄ 466.752

Struct. unknown. Prob. a Sparteine-type dimer. Alkaloid from *Cytisus scoparius* (Fabaceae). Mp 173-174°. $[\alpha]_D$ -48.4 (CHCl₃). Ir spectrum resembles that of sparteine; indicates no CH₃ groups or sites of unsatn.


Valeur, A. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1918, **167**, 26 (*isol*)

Delaby, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1949, **16**, 152-154

Paris, R.R. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 1597-1599 (*chromatog*)

Sarpagine S-89

Sarpagan-10,17-diol, 9CI. Raupine [482-68-8]



C₁₉H₂₂N₂O₂ 310.395

Alkaloid from many *Rauwolfia* spp., *Alstonia yunnanensis*, *Vinca major* and *Vinca difformis* (Apocynaceae). Needles (EtOH or MeOH). $[\alpha]_D^{20}$ +54 (Py). Dec. at 320° without melting.

Hydrochloride:

Needles (EtOH). Dec. at 220° without melting.

Hydrobromide:

Prisms (EtOH). Dec. at 220° without melting.

O¹⁷-Ac: 17-Acetylsarpagine

[102358-21-4]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from roots of *Alstonia yunnanensis*.

Di-O-Ac:

Cryst. (C₆H₆). Mp 244-245°. $[\alpha]_D^{20}$ -23 (c, 0.46 in Py). $[\alpha]_D$ +24 (c, 0.45 in EtOH).

N¹-Me: N-Methylsarpagine

[17801-05-7]

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from the bark of *Alstonia spectabilis* (Apocynaceae). Prisms (MeOH). Mp 300° dec.

N¹,N⁴-Di-Me: N^{1,4}-Methylsarpagine

methosalt

[4849-01-8 (chloride)]

C₂₁H₂₇N₂O₂⁺ 339.456

Alkaloid from the roots of *Pleiocarpa mutica* and *Pleiocarpa tubicina* (Apocynaceae). Cryst. (MeOH) (as chloride). Mp 275-280° (dec.) (as chloride). $[\alpha]_D$ +56 (c, 0.174 in MeOH aq.).

10-Me ether: Lochnerine. Lochnerine.

10-Methoxysarpagan-17-ol, 9CI. Alkaloid C†. C-Alkaloid T

[522-47-4]

C₂₀H₂₄N₂O₂ 324.422

Alkaloid present in curare, *Lochnera rosea* (preferred genus name *Catharanthus*), *Vinca major*, *Rauwolfia suaveolens*, *Rauwolfia macrophylla* and *Rauwolfia nitida* (Apocynaceae). Strong hypoglycaemic agent. Cryst. (MeOH aq.). Mp 202.5-203.5° (195-197° dec.). $[\alpha]_D^{25}$ +72 (c, 0.624 in EtOH). $[\alpha]_D^{24}$ +52 (c, 0.385 in Py).

10-Me ether, 17-Ac:

Needles (CH₂Cl₂/Et₂O). Mp 243°. $[\alpha]_D^{24}$ -5.6 (c, 0.358 in Py).

10-Me ether, N¹-Me: 10-Methoxyaffinisine

[7689-00-1]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from the stem bark of *Alstonia macrophylla*. Needles (MeOH). Mp 205-206°. $[\alpha]_D$ +75 (c, 0.62 in CHCl₃). λ_{\max} 223 (log ϵ 4.25); 283 (log ϵ 4.02); 300 (log ϵ 3.91) (EtOH).

10-Me ether, N⁴-Me: Lochneram. 17-

Hydroxy-10-methoxy-4-methylsarpagine

nium

[6901-26-4]

C₂₁H₂₇N₂O₂⁺ 339.456

Alkaloid from curare (Loganiceae).

Cryst. (Me₂CO) (as iodide). Mp 235-238° (iodide). $[\alpha]_D^{23}$ +41 (c, 0.48 in 96% EtOH) (iodide).

17-Aldehyde, Me ether: 10-Methoxyvellonimine

[2149-40-8]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Vinca major* (Apocynaceae). Needles (C₆H₆/EtOAc or MeOH). Mp 224-226°.

17-Aldehyde, Me ether, N¹-Me: Majvinine

[63959-49-9]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from the aerial parts of *Vinca major* (Apocynaceae). Needles (C₆H₆). Mp 195-197° dec.

18-Hydroxy, 10-Me ether: 18-Hydroxylochnerine

[107603-56-5]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the leaves, stem and root bark of *Rauwolfia biauiculata* (Apocynaceae). Amorph. solid.

16-Epimer: 16-Episarpagine

[102490-01-7]

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from roots of *Alstonia yunnanensis*. Cryst. Mp 300° dec. $[\alpha]_D^{30}$ +34.7 (c, 0.08 in EtOH).

16-Epimer, 17-carboxylic acid, Me ester: 10-Hydroxypericyclivine

[132242-29-6]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the leaves of *Alstonia undulata* (Apocynaceae).

16-Epimer, 17-carboxylic acid, N¹-Me, Me ester: 10-Hydroxy-N-methylpericyclivine

[132242-28-5]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the leaves of *Alstonia undulata* (Apocynaceae).

16-Epimer, 17-carboxylic acid, Me ether, Me ester: 10-Methoxypericyclivine

[132242-30-9]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the leaves of *Alstonia undulata* (Apocynaceae). Cryst. (Me₂CO). Mp 235-238°. $[\alpha]_D$ +1.7 (c, 1 in CHCl₃).

16-Epimer, 17-carboxylic acid, Me ether, N¹-Me, Me ester: 10-Methoxy-N-methylpericyclivine

[132242-31-0]

C₂₂H₂₆N₂O₃ 366.459

Alkaloid from the leaves of *Alstonia undulata* (Apocynaceae).

Also isol. from *Haplophyton crooksii* (Apocynaceae).

Cryst. (Me₂CO). Mp 230-231° (213-215°). $[\alpha]_D$ +23 (c, 0.5 in CHCl₃). Erroneously claimed to be novel by Mroue *et al* (1996).

Stoll, A. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 1143-1147 (*isol, uv*)

Janot, M.M. *et al.*, *C. R. Hebd. Seances Acad. Sci.*, 1956, **243**, 1789 (*uv, ir*)

Mors, W.B. *et al.*, *Chem. Ind. (London)*, 1956, 173-174 (*Lochnerine, isol, uv, ir*)


Stauffer, D. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 508-513 (*synth, Lochnerine*)

Arnold, W. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 705-716 (*Lochnerine, Lochneram*)

- Biemann, K. *et al.*, *J.A.C.S.*, 1961, **83**, 4801-4805 (*ms. synth. Lochnerine*)
- Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 2497-2501 (*10-Methoxyvellosimine*)
- Khan, Z.M. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 1957-1965; 1967, **50**, 625-627 (*N^o-Methylsarpagine methochloride*)
- Hart, N.K. *et al.*, *Aust. J. Chem.*, 1972, **25**, 2739-2741 (*N-Methylsarpagine*)
- Blaha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 3168-3176 (*uv, cd, ord*)
- Banerji, A. *et al.*, *Phytochemistry*, 1974, **13**, 2309-2312; 1977, **16**, 1124-1125 (*Sarpagine, Lochnerine, 10-Methoxyvellosimine, Majvinine, isol, uv, ir, pmr, ms, struct*)
- Chen, W. *et al.*, *Yaoxue Xuebao*, 1985, **20**, 906; *CA*, **105**, 3520s (*epimer, 17-Acetylsarpagine*)
- Abaul, J. *et al.*, *J. Nat. Prod.*, 1986, **49**, 829-832 (*18-Hydroxylochnerine*)
- Pinchon, T.-M. *et al.*, *Phytochemistry*, 1990, **29**, 3341-3344 (*Alstonia undulata constits*)
- Mroue, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 890-893 (*10-Methoxy-N-methylpericyclivine*)
- Kam, T.-S. *et al.*, *Phytochemistry*, 1999, **51**, 839-844 (*10-Methoxyaffinisine*)
- Zhao, S. *et al.*, *J.O.C.*, 2003, **68**, 6279-6295 (*N-Methylsarpagine, Majvinine, 10-Methoxyaffinisine, synth*)
- Sarma, P.V.V.S. *et al.*, *Org. Lett.*, 2006, **8**, 1017-1020 (*10-Hydroxy-N-methylpericyclivine, synth*)
- Lewis, S.E. *et al.*, *Tetrahedron*, 2006, **62**, 8655-8681 (*rev, chem*)

Sarracine

Mikanoidine
[2492-09-3]



$C_{18}H_{27}NO_5$ 337.415

Double ester of Platynecine, H-233 with Sarracinic acid, H-574 and Angelic acid. Alkaloid from *Senecio sarraceni*, *Senecio mikanoides*, *Senecio sylvaticus*, *Senecio rhombifolius*, *Senecio franchetii* and *Adenostyles alliariae* (preferred genus name *Cacalia*) (Asteraceae). Was used in the USSR for treatment of gastrointestinal hypermotility and peptic ulceration. Spasmolytic agent. Cryst. (EtOH). Mp 51-52° (45-46°). $[\alpha]_D^{20}$ -129.7 (EtOH). Log P 1.81 (calc).

► LD₅₀ (rat, ipr) 218 mg/kg. EM9251400

Picrate:

Cryst. (EtOH). Mp 140-141°.

N-Oxide: Sarracine N-oxide

[19038-27-8]

$C_{18}H_{27}NO_6$ 353.414

Alkaloid from *Senecio sarraceni*, *Senecio franchetii* and *Senecio mikanoides* (Asteraceae). Cryst. + 1H₂O (Me₂CO). Mp 123-124°. $[\alpha]_D^{20}$ -81.65 (H₂O). $[\alpha]_D^{20}$ -94 (EtOH). Becomes

anhyd. at 100°, rehydrates on standing in air.

N-Oxide, picrate:

Cryst. (EtOH). Mp 107.5-108.5°.

O-(3-Hydroxy-3-methylbutanoyl): Racemodine

[147554-28-7]

$C_{23}H_{35}NO_7$ 437.532

Alkaloid from *Senecio racemosus* (Asteraceae). Oil. $[\alpha]_D^{20}$ -135 (c, 0.35 in EtOH).

(2'E,2''E)-isomer: Neosarracine

[136173-27-8]

$C_{18}H_{27}NO_5$ 337.415

Alkaloid from *Senecio serra*, *Senecio hydrophyllus* and *Senecio mikanoides* (Asteraceae).

(2'E,2''Z)-isomer: Sarracine

[136173-25-6]

$C_{18}H_{27}NO_5$ 337.415

Alkaloid from *Senecio serra*, *Senecio hydrophyllus* and *Senecio mikanoides* (Asteraceae).

(2'Z,2''E)-isomer: Neosarracine

[136173-26-7]

$C_{18}H_{27}NO_5$ 337.415

Alkaloid from *Senecio chrysocoma* and *Senecio kaschkarovii*. Tentatively identified also in a GC-MS study in *Senecio hydrophyllus* and *Senecio mikanoides* (Asteraceae). Oil. $[\alpha]_D^{20}$ +60.5 (c, 0.5 in EtOH).

7a-Epimer: 8-Episarracine

$C_{18}H_{27}NO_5$ 337.415

Alkaloid from the roots of *Senecio macedonicus*.

7a-Epimer, N-oxide: 8-Episarracine N-oxide

$C_{18}H_{27}NO_6$ 353.414

Alkaloid from the roots of *Senecio macedonicus*. Oil.

7a-Epimer, (2'Z,2''E)-isomer: 8-Epineosarracine

$C_{18}H_{27}NO_5$ 337.415

Alkaloid present in the roots of *Senecio macedonicum*. Identified by gc-ms.

Danilova, A.V. *et al.*, *Zh. Obshch. Khim.*, 1953, **23**, 1417; 1597; *CA*, **47**, 12759 (*isol, Sarracine, oxide*)

Culvenor, C.C.J. *et al.*, *J.O.C.*, 1961, **26**, 3045 (*isol, pmr, struct*)

Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 351; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 296 (*isol*)

White, E.P. *et al.*, *An. Quim.*, 1972, **68**, 723 (*isol*)

Dauksha, V.E. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 831; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 753 (*isol*)

Stelljes, M.E. *et al.*, *J. Nat. Prod.*, 1991, **54**, 759 (*Neosarracine, Neosarracine, Sarracine*)

Cheng, D.L. *et al.*, *Phytochemistry*, 1992, **31**, 3671 (*Neosarracine*)


Ahmed, W. *et al.*, *Phytochemistry*, 1993, **32**, 224 (*Racemodine*)

Grue, M.R. *et al.*, *Phytochemistry*, 1993, **33**, 1517 (*Neosarracine*)

Christov, V. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 780-784 (*8-Episarracine, 8-Episarracine N-oxide, 8-Epineosarracine*)

Sarusubine A

[956298-05-8]



Relative Configuration

$C_{56}H_{70}N_2O_{12}$ 963.175

Alkaloid from the leaves of *Lagerstroemia subcostata*. Pale yellow solid. $[\alpha]_D^{23}$ +40.1 (c, 1 in MeOH).


Watanabe, K. *et al.*, *Tet. Lett.*, 2007, **48**, 7502-7504 (*isol, pmr, cmr, ms*)

Sativanine A

S-92

2-(Dimethylamino)-3-methyl-N-[7-(1-methylethyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]pentanamide, 9CI

[72361-59-2]



$C_{30}H_{40}N_4O_4$ 520.67

Alkaloid from the bark of *Zizyphus sativa* and from *Zizyphus spina-christi* (Rhamnaceae). Mp 80°. Opt. rotn. not recorded.


Tschesche, R. *et al.*, *Phytochemistry*, 1979, **18**, 702 (*isol, uv, ms, struct*)

Sativanine B

S-93

4-(4-Butyl-3-methyl-5-oxo-1-imidazolidinyl)-7-(1-methylethyl)-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraene-5,8-dione, 9CI

[72361-60-5]




$C_{30}H_{38}N_4O_4$ 518.655

Alkaloid from the bark of *Zizyphus sativa* (Rhamnaceae). Amorph. Opt. rotn. not recorded. Related to Nummularine G, N-333.

Tschesche, R. *et al.*, *Phytochemistry*, 1979, **18**, 702 (*isol, uv, ms, struct*)

Sativanine C

[92664-86-3]



$C_{29}H_{43}N_5O_6$ 557.689
Alkaloid from the bark of *Zizyphus sativa*. Mp 113-114°.

N²-Formyl: Sativanine M


$C_{30}H_{43}N_5O_7$ 585.699
Alkaloid from the bark of *Zizyphus sativa*. Cryst. (MeOH). Mp 210-212°. $[\alpha]_D^{20}$ -215 (MeOH). λ_{max} 260 (log ϵ 3.7); 320 (log ϵ 3.6) (MeOH).

Shah, A.H. *et al.*, *Phytochemistry*, 1984, **23**, 931-933 (*Sativanine C*)

Pandey, M.B. *et al.*, *Nat. Prod. Res.*, 2008, **22**, 219-221 (*Sativanine M*)

Sativanine D

[99694-93-6]




$C_{30}H_{43}N_5O_6$ 569.7
Alkaloid from the bark of *Zizyphus sativa* (Rhamnaceae). Mp 119-121°.

Shah, A.H. *et al.*, *Phytochemistry*, 1985, **24**, 2765 (*isol, uv, ir, ms, struct*)

Sativanine E

[98618-05-4]




S-94

$C_{33}H_{41}N_5O_5$ 587.717
Alkaloid from the bark of *Zizyphus sativa* (Rhamnaceae). Mp 127-128°. $[\alpha]_D^{20}$ -99 (c, 0.2 in $CHCl_3$).

Shah, A.H. *et al.*, *J. Nat. Prod.*, 1985, **48**, 555 (*isol, uv, ir, pmr, ms, struct*)

Sativanine F

[99694-92-5]




S-97

$C_{34}H_{43}N_5O_7$ 633.743
Alkaloid from the bark of *Zizyphus sativa* (Rhamnaceae). Mp 139-141°.

Shah, A.H. *et al.*, *Phytochemistry*, 1985, **24**, 2768 (*isol, uv, ir, ms, struct*)

Sativanine G

[93754-84-8]



Absolute Configuration

$C_{28}H_{42}N_4O_5$ 514.664
Alkaloid from *Paliurus ramossissimus* and *Zizyphus sativa* (Rhamnaceae). Amorph. solid. Mp 93-94°. $[\alpha]_D^{26}$ -327 (c, 0.85 in MeOH). λ_{max} 217 (log ϵ 4.66); 230 (log ϵ 4.51); 269 (log ϵ 4.65) (MeOH).

N,N-Di-de-Me: Sativanine N

$C_{26}H_{38}N_4O_5$ 486.61
Alkaloid from the bark of *Zizyphus sativa*. Cryst. (MeOH). Mp 176-178°. $[\alpha]_D^{20}$ -158 (c, 0.18 in $CHCl_3$). λ_{max} 270 (log ϵ 2.78); 322 (log ϵ 2.58) (MeOH).

N,N-Di-de-Me, 2'-N-formyl: Sativanine K

[108906-92-9]
 $C_{27}H_{38}N_4O_6$ 514.62
Alkaloid from the bark of *Zizyphus sativa*. Mp 160-162°. λ_{max} 260; 320 (MeOH).

Shah, A.H. *et al.*, *Phytochemistry*, 1984, **23**, 2120-2121; 1987, **26**, 1230 (*isol, uv, ir, ms, struct*)


Lin, H.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1338-1343 (*isol, uv, cd, pmr, cmr*)

Singh, S. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 733-737 (*Sativanine N*)

Sativanine H

[107603-59-8]

S-99




$C_{29}H_{43}N_5O_6$ 557.689
Alkaloid from the stem bark of *Zizyphus sativa* and *Zizyphus rugosa*. Mp 191-192°.

Shah, A.H. *et al.*, *Planta Med.*, 1986, 500-501 (*isol*)

Tripathi, Y.C. *et al.*, *Phytochemistry*, 1989, **28**, 1563-1565 (*occur*)

Sativanine O

S-100




$C_{32}H_{34}N_4O_5$ 554.644
Alkaloid from the bark of *Zizyphus sativa*. Cryst. (MeOH). Mp 224-226°. $[\alpha]_D^{20}$ -195 (c, 0.15 in $CHCl_3$). λ_{max} 270 (log ϵ 2.8); 320 (log ϵ 2.4) (MeOH).

Singh, S. *et al.*, *J. Asian Nat. Prod. Res.*, 2006, **8**, 733-737 (*isol, ms*)

Sattazolol

2-Hydroxy-1-(1H-indol-3-yl)-5-methyl-3-hexanone

S-101


 $C_{15}H_{19}NO_2$ 245.321**(+)-form** [169272-85-9]

Prod. by *Bacillus* sp. B-60. Antiviral agent. Sol. MeOH, $CHCl_3$. $[\alpha]_D$ +58 (c, 1.5 in $CHCl_3$).

Me ether: 1-(1H-Indol-3-yl)-2-methoxy-5-methyl-3-hexanone. **O-Methylsattazolol** [168916-24-3]

$C_{16}H_{21}NO_2$ 259.347
Prod. by *Bacillus* sp. B-60. Antiviral agent. Sol. MeOH, $CHCl_3$. $[\alpha]_D$ -22 (c, 0.5 in $CHCl_3$).
Lampis, G. *et al.*, *J. Antibiot.*, 1995, **48**, 967 (isol, pmr, activity)

Saulatine S-102



$C_{22}H_{23}NO_6$ 397.427

(±)-**form** [91897-61-9]
Alkaloid from the roots of *Abuta bullata* (Menispermaceae). Cryst. (MeOH). Mp 226-228°. Prob. an artifact.

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 539 (uv, ir, pmr, cmr, ms, struct)
Shamma, M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 398
Kim, D.C. *et al.*, *J. Het. Chem.*, 1993, **30**, 1431 (synth)


Saururidine S-103

$C_{17}H_{25}NO_3$ 291.389
Struct. unknown. Alkaloid from *Lycopodium saururus* (Lycopodiaceae).
MacLean, D.B. *et al.*, *Alkaloids* (Academic Press), 1985, **26**, 241

Saururine S-104

$C_{10}H_{19}N$ 153.267
Struct. unknown. Alkaloid from the leaves of *Lycopodium saururus* (Lycopodiaceae). Syrup.
Picrate:
Yellow prisms (Me₂CO aq.). Mp 202°.
Methiodide:
Fine needles (H₂O). Mp 242-244°.
Deulofeu, V. *et al.*, *J.A.C.S.*, 1942, **64**, 968-969 (isol)


Saussureamine A S-105
[148245-82-3]



$C_{20}H_{29}NO_4$ 347.453
Constit. of *Saussurea lappa* root. Prisms (MeOH aq.). Mp 115-117°. $[\alpha]_D$ +36.7 (c, 0.4 in MeOH).


Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 214-216 (isol, pmr, cmr)
Matsuda, H. *et al.*, *Tetrahedron*, 2000, **56**, 7763-7777 (synth, abs config)

Saussureamine C S-106
[148245-83-4]



$C_{19}H_{26}N_2O_5$ 362.425
Constit. of *Saussurea lappa* root. Powder. $[\alpha]_D$ -17.2 (c, 0.5 in MeOH).
Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 214-216 (isol, pmr, cmr)
Matsuda, H. *et al.*, *Tetrahedron*, 2000, **56**, 7763-7777 (synth, abs config)


Saussureamine D S-107
[148225-51-8]



$C_{20}H_{29}NO_5$ 363.453
Constit. of *Saussurea lappa* root. Needles (MeOH aq.). Mp 235-237°. $[\alpha]_D$ +13.3 (c, 0.3 in MeOH).

$\Delta^4(15)$ -**Isomer: Saussureamine E**
[148225-52-9]
 $C_{20}H_{29}NO_5$ 363.453
Constit. of *Saussurea lappa* root. Needles (MeOH aq.). Mp 150-153°. $[\alpha]_D$ +33.7 (c, 0.9 in MeOH).
Yoshikawa, M. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 214-216 (isol, pmr, cmr)
Matsuda, H. *et al.*, *Tetrahedron*, 2000, **56**, 7763-7777 (synth, abs config)

Sauvagine S-108
[133084-03-4]




$C_{18}H_{11}NO_6$ 337.288
Struct. revised in 1996. Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph. Mp 215-217°.

3,4-Dihydro: Dihydrosauvagine
[133083-99-5]
 $C_{18}H_{13}NO_6$ 339.304
Alkaloid from *Corydalis claviculata* (Papaveraceae). Amorph.

Allais, D.P. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1280-1286 (isol, pmr)
Garcia, A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 806-807 (synth, ir, struct)

Saxitoxin S-109
STX
[35523-89-8]



$C_{10}H_{17}N_7O_4$ 299.289
Isol. from Alaska butter clams (*Saxidomus giganteus*), toxic mussels (*Mytilus californianus*), the plankton *Gonyaulax catenella*, *Protogonyaulax tamarensis* and other marine organisms. Sodium channel (I, II, III, h1) blocker. Causal agent of paralytic shellfish poisoning. Noncryst. Sol. H₂O, MeOH; fairly sol. EtOH, AcOH; poorly sol. butanol, hexane. $[\alpha]_D$ +130. See also 11-Hydroxysaxitoxin, H-731.

► Extremely toxic, among the most toxic substances known. LD₅₀ (rat, orl) 0.192 mg/kg. LD₅₀ (mus, ipr) 0.005 mg/kg. LD₅₀ (mus, ivn) 0.008 mg/kg. UY8708500
N¹⁴-Sulfonic acid: Gonyautoxin V. GTX5.
Toxin B₁
[64296-25-9]
 $C_{10}H_{17}N_7O_7S$ 379.353
Present in *Gonyaulax* and *Protogonyaulax* spp. and other marine organisms. Neurotoxin.
► Toxic.

N¹⁴-Me: N'-Methylsaxitoxin
[209856-64-4]
 $C_{11}H_{19}N_7O_4$ 313.316
Isol. from the freshwater puffer *Tetraodon cutcutia*.

Decarbamoyl: Decarbamoylsaxitoxin
[58911-04-9]
 $C_9H_{16}N_6O_3$ 256.264
Isol. from various shellfish.

Decarbamoyl, 13-Ac: LWTX5
[133144-32-8]
 $C_{11}H_{18}N_6O_4$ 298.301
Isol. from *Lyngbya wollei*.

Decarbamoyl, 13-O-(4-hydroxybenzoyl): GC 3
[603125-82-2]
 $C_{16}H_{20}N_6O_5$ 376.371
Isol. from *Gymnodinium catenatum*.

De(carbamoyloxy): Decarbamoyloxysaxitoxin
[143084-69-9]
[164905-60-6]
 $C_9H_{16}N_6O_2$ 240.264
Isol. from *Gymnodinium catenatum*.

N¹⁴-Hydroxy: Neosaxitoxin
[64296-20-4]
 $C_{10}H_{17}N_7O_5$ 315.288

Prod. by *Protogonyaulax* and found in shellfish, marine algae and other organisms incl. *Gonyaulax tamarensis*, *Aphanizomenon flos-aquae*, *Saxidomus giganteus*, *Vibrio* sp., *Pyrodinium* sp. Neurotoxic neuromuscular blocker.

- Potent neurotoxin; LD₅₀ (mus, ipr) 5 mg/kg.

N¹-Hydroxy, N¹⁴-sulfonic acid: **Gonyautoxin VI. GTX6. Toxin B₂**
[82810-44-4]

C₁₀H₁₇N₇O₈S 395.352

Prod. by *Gonyaulax* and *Protogonyaulax* spp., isol. from shellfish. Neurotoxin; causal agent of shellfish poisoning.

- Toxic.

N¹-Hydroxy, decarbamoyl: **Decarbamoylneosaxitoxin**

[68683-58-9]

C₉H₁₆N₆O₄ 272.263

Isol. from the crab *Zosimus aeneus*.

N¹⁴-Hydroxy: **N'-Hydroxysaxitoxin**

[153856-77-0]

C₁₀H₁₇N₇O₅ 315.288

Isol. from the crab *Zosimus aeneus*.

Contains an N-hydroxycarbamoyl group.

N¹⁴,1-Dihydroxy: **N'-Hydroxyneosaxitoxin**

[153856-78-1]

C₁₀H₁₇N₇O₆ 331.288

Isol. from the crab *Zosimus aeneus*.

Toxic paralytic agent.

12-Deoxy(12S-), decarbamoyl: **Decarbamoyl-12-deoxysaxitoxin. LWTX4**

[75352-31-7]

[80844-68-4]

C₉H₁₆N₆O₂ 240.264

Isol. from *Lyngbya wollei*. Neurotoxin.

12-Deoxy(12S-), decarbamoyl, 13-O-Ac: **LWTX6**

[200816-98-4]

C₁₁H₁₈N₆O₃ 282.302

Isol. from *Lyngbya wollei*.

11-Hydroxy: see 11-Hydroxysaxitoxin, H-731

[35554-08-6]

Schantz, E.J. *et al.*, *Biochemistry*, 1966, **5**, 1191-1195 (*isol, bibl*)

Bordner, J. *et al.*, *J.A.C.S.*, 1975, **97**, 6008-6012 (*cryst struct, pmr, cmr*)

Tanino, H. *et al.*, *J.A.C.S.*, 1977, **99**, 2818-2819 (*synth*)

Koehn, F.E. *et al.*, *Bioorg. Chem.*, 1981, **10**, 412-428 (*decarbamoyl, synth*)

Fix Wichman, C. *et al.*, *Tet. Lett.*, 1981, **22**, 1941-1944 (*isol*)

Harada, T. *et al.*, *Agric. Biol. Chem.*, 1982, **46**, 1861-1864 (*Gonyautoxins V, VI*)

Onoue, Y. *et al.*, *J. Agric. Food Chem.*, 1983, **31**, 420-423 (*props*)

Oshima, Y. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 1707-1711 (*chromatog, struct, bibl*)

Maruyama, J. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 2783-2788 (*ms*)

Shimizu, Y. *et al.*, *Tetrahedron*, 1984, **40**, 539-544 (*props*)

Kodama, M. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 1075-1077 (*isol*)

Hall, S. *et al.*, *ACS Symp. Ser.*, 1990, **418**, (*book*)

Arakawa, O. *et al.*, *Toxicon*, 1994, **32**, 175-183 (*Zosimus aeneus constits*)

Onodera, H. *et al.*, *Nat. Toxins*, 1997, **5**, 175-186 (*LWTX toxins*)

Zaman, L. *et al.*, *Toxicon*, 1998, **36**, 627-630 (*N'-Methylsaxitoxin*)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., *Pharmaceutical Press*, 1999, 16273

Food Sci. Technol., Seafood and Freshwater Toxins, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (revs)

Negri, A. *et al.*, *Chem. Res. Toxicol.*, 2003, **16**, 1029-1033 (*GC3*)

Ciminiello, P. *et al.*, *Eur. J. Org. Chem.*, 2004, 2533-2551 (*rev*)

Llewellyn, L.E. *et al.*, *Nat. Prod. Rep.*, 2006, **23**, 200-222 (*rev*)

Jang, J. *et al.*, *Toxicon*, 2006, **48**, 980-987 (*occur*)

Iwamoto, O. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 8625-8628 (*Decarbamoylsaxitoxin, synth*)

Fleming, J.J. *et al.*, *J.A.C.S.*, 2007, **129**, 9964-9975 (*synth*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., *J. Wiley*, 2000, SBA500

11-Saxitoxinacetic acid

S-110

11-Saxitoxinethanoic acid

[172888-62-9]



Absolute Configuration

C₁₂H₁₉N₇O₆ 357.325


Assumed to exist as a mixt. of 3 tautomers. Isol. from the xanthid crab *Atergatis floridus*.

Arakawa, O. *et al.*, *Toxicon*, 1995, **33**, 1577-1584

Saxoguttine

S-111

[91174-13-9]



C₂₁H₂₅N₆O₆ 387.432

Alkaloid from the stem bark of *Guatteria discolor* (Annonaceae). Noncryst.

O^{4'}, O⁶-Di-de-Me, O^{3'}-Me: **Pseudoronine**
C₂₀H₂₃N₆O₆ 373.405

Alkaloid from the roots of *Papaver pseudo-orientale* (Papaveraceae). Yellow needles (MeOH). Mp 249-252°.

Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1984, **47**, 353 (*isol, uv, ir, pmr, cmr, ms, struct*)

Věžník, F. *et al.*, *Coll. Czech. Chem. Comm.*, 1986, **51**, 1752 (*Pseudoronine*)


Rozwadowska, M.D. *et al.*, *Pol. J. Chem. (Rocz. Chem.)*, 1991, **65**, 1287 (*synth*)

Bentley, K.W. *et al.*, *The Isoquinoline Alkaloids*, 1998, 41 (*Pseudoronine, struct*)

Saxorumamide

S-112

[949092-28-8]



Relative Configuration

C₁₇H₂₃NO₄ 305.373

Alkaloid from the roots of *Stemona saxorum*. Light yellow powder. [α]_D²⁰ -15.4 (c, 0.35 in MeOH). λ_{max} 204 (log ε 4.04); 300 (log ε 2.22) (MeOH).

12-Epimer: **Isosaxorumamide**

[949092-29-9]

C₁₇H₂₃NO₄ 305.373

Alkaloid from the roots of *Stemona saxorum*. Light yellow powder. [α]_D²⁰ -15.2 (c, 0.13 in MeOH). λ_{max} 205 (log ε 4.2); 299 (log ε 2.36) (MeOH).

Wang, Y.-Z. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1356-1359 (*isol, pmr, cmr*)

Scabrine

S-113

[111-04-6]

C₂₂H₃₅NO 329.525

Struct. unknown. An isobutylamide of an acid C₁₈H₂₆O₂ which may be an octadecapentaenoic acid or may be acetylenic. Isol. from seeds of *Heliopsis scabra* (Asteraceae). Yellow oil.


Jacobson, M. *et al.*, *J.A.C.S.*, 1951, **73**, 100-103

Bohlmann, F. *et al.*, *Annalen*, 1983, 1202-1206

Scabrosin

S-114

[67866-98-2]



Absolute Configuration

C₁₈H₁₆N₂O₆S₂ 420.466

Epithiodioxopiperazine antibiotics. Struct. revised in 1999.

Di-Ac: **Scabrosin 4,4'-diacetate**

[67705-18-4]

C₂₂H₂₀N₂O₈S₂ 504.541

Isol. from lichen *Xanthoparmelia scabrosa*. Mp 262°.

4'-Butanoyl, 4-Ac: **Scabrosin 4-acetate-4'-butyrate**

[67705-19-5]

C₂₄H₂₄N₂O₈S₂ 532.594

Isol. from *Xanthoparmelia scabrosa*. Mp 216.5-217.5°. [α]_D²⁵ -9600.

Dibutanoyl: **Scabrosin 4,4'-dibutyrate**

[67705-20-8]

C₂₆H₂₈N₂O₈S₂ 560.648

Isol. from *Xanthoparmelia scabrosa*. Mp 197-199° (196-197°).

4'-Hexanoyl, 4-Ac: **Scabrosin 4-acetate-4'-hexanoate**

[67705-21-9]

$C_{26}H_{28}N_2O_8S_2$ 560.648
Isol. from *Xanthoparmelia scabrosa*.
Mp 213° (209°).

4'-Hexanoyl, 4-butanoyl: Scabrosin 4-butanoate-4'-hexanoate

$C_{28}H_{32}N_2O_8S_2$ 588.701
Isol. from *Xanthoparmelia scabrosa*.
Cytotoxic agent. Amorph. solid. Mp 139-141°.


Begg, W.R. *et al.*, *Tet. Lett.*, 1978, 1047 (*ir, ms, cmr, struct*)

Ernst-Russell, M.A. *et al.*, *Aust. J. Chem.*, 1999, **52**, 279-283 (*isol, pmr, cmr, ms, cryst struct*)

Scaevodimerine A

S-115

[112579-70-1]



$C_{21}H_{25}N_3O_4$ 383.446

Scaevodimerines are the first bis-monoterpene alkaloids. Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae) and from *Strychnos spinosa*. Amorph. solid. $[\alpha]_D^{20} +31$ (c, 1 in MeOH).

1',2'-Didehydro: Scaevodimerine B

[112561-59-8]

$C_{21}H_{23}N_3O_4$ 381.43

Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae) and from *Strychnos minfiensis*. Amorph. solid. $[\alpha]_D^{20} +39$ (c, 1 in MeOH).

1,2,4a,7a-Tetrahydro: Scaevodimerine C

[112561-60-1]

$C_{21}H_{21}N_3O_4$ 379.415

Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae). $[\alpha]_D^{20} -18$ (c, 1 in MeOH).

1',2'-Didehydro, 1-oxo: Scaevodimerine D

[112579-71-2]

$C_{21}H_{21}N_3O_5$ 395.414

Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae) and from *Strychnos pungens*. $[\alpha]_D^{20} +48$ (c, 0.2 in MeOH).

Skaltsounis, A.-L. *et al.*, *Heterocycles*, 1987, **26**, 599 (*uv, ir, pmr, struct, derivs*)


Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1992, **61**, 429-440 (*isol, Strychnos*)

Thépenier, P. *et al.*, *Bull. Soc. R. Sci. Liege*, 1996, **65**, 379-382 (*isol, Strychnos*)

Scaevoline

S-116

[99339-50-1]




$C_{11}H_{11}NO_3$ 205.213

Alkaloid from the aerial parts of *Scaevola racemigera* (Goodeniaceae). Prisms (CHCl₃). Mp 119-121°. $[\alpha]_D^{20} 0$ (c, 1 in MeOH). λ_{max} 235 (log ϵ 3.88); 268 (log ϵ 3.62); 298 (log ϵ 3.24); 310 (log ϵ 3.12) (EtOH).

Skaltsounis, A.-L. *et al.*, *Helv. Chim. Acta*, 1985, **68**, 1679-1685 (*isol, uv, ir, pmr, ms, struct*)

Scalusamide A

S-117



$C_{16}H_{27}NO_3$ 281.394

Prod. by marine-derived *Penicillium citrinum* strain N 055. Antifungal agent. Amorph. solid. $[\alpha]_D^{22} -28$ (c, 1 in CHCl₃).

8',9'-Dihydro: Scalusamide C

$C_{16}H_{29}NO_3$ 283.41

Prod. by *Penicillium citrinum* strain N 055. Amorph. solid. $[\alpha]_D^{26} -13$ (c, 0.2 in CHCl₃).

6',7'-Didehydro(E-), 8',9'-dihydro: Scalusamide B

$C_{16}H_{27}NO_3$ 281.394


Prod. by *Penicillium citrinum* strain N 055. Amorph. solid. $[\alpha]_D^{26} -13$ (c, 0.2 in CHCl₃).

Tsuda, M. *et al.*, *J. Nat. Prod.*, 2005, **68**, 273-276 (*isol, pmr, cmr*)

Scandine

S-118

3-(Methoxycarbonyl)meloscine, 9CI
[24314-59-8]



Absolute Configuration

$C_{21}H_{22}N_2O_3$ 350.416

Alkaloid from *Melodinus scandens* (Apocynaceae). Prisms (MeOH/EtOH). Mp 188-192°. $[\alpha]_D^{25} +254$ (c, 0.2 in EtOH).

N^b-Oxide: Scandine N^b-oxide

[140701-69-5]

$C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from *Melodinus fusiformis* (Apocynaceae).

14 ξ ,15 ξ -Epoxide: 14,15-Epoxyscandine

$C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from aerial parts of *Melodinus hemsleyanus* (Apocynaceae). Prisms (Me₂CO). Mp 116-118°.

10-Hydroxy: 10-Hydroxyscandine

[119188-47-5]

$C_{21}H_{22}N_2O_4$ 366.416

Alkaloid from the stem bark of *Melodinus tenuicaudatus* (Apocynaceae).

Mp 180°. $[\alpha]_D +228$ (CHCl₃).

Bernauer, B.K. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 1886 (*uv, ms, pmr*)

Daudon, M. *et al.*, *J.O.C.*, 1975, **40**, 2838 (*cmr*)

Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1655 (*cryst struct, abs config*)

Zhou, Y.L. *et al.*, *Planta Med.*, 1988, **54**, 315 (*10-Hydroxyscandine*)

He, X. *et al.*, *Huaxue Xuebao*, 1992, **50**, 96; *CA*, **116**, 191092c (*oxide*)


Guo, L.-W. *et al.*, *Phytochemistry*, 1993, **34**, 563 (*14,15-Epoxyscandine*)

Gu, J.-M. *et al.*, *Acta Cryst. E*, 2006, **62**, o4775-o4777 (*cryst struct*)

Scandomelidine

S-119

[139682-13-6]



$C_{42}H_{46}N_4O_5$ 686.849


Alkaloid from stems and leaves of *Melodinus scandens* (Apocynaceae). Amorph. powder. $[\alpha]_D^{20} -160$ (c, 1 in CHCl₃).

Mehri, H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 241 (*isol, uv, ir, pmr, cmr, ms, struct*)

Scandomeline

S-120

[59813-32-0]



Absolute Configuration

$C_{42}H_{46}N_4O_6$ 702.849

Alkaloid from the leaves and stems of *Melodinus scandens* (Apocynaceae). Cryst. (Me₂CO). Mp 300° dec. $[\alpha]_D^{22} -170$ (c, 1 in CHCl₃). λ_{max} 232 (sh) (log ϵ 3.93); 259 (log ϵ 3.94); 302 (log ϵ 3.92); 328 (log ϵ 4.01) (EtOH).

19-Epimer: Episcandomeline

[59830-07-8]

$C_{42}H_{46}N_4O_6$ 702.849


Isol. from *Melodinus scandens* (Apocynaceae). Cryst. (Me₂CO). Mp 300° dec. $[\alpha]_D^{22} -112$ (c, 0.5 in CHCl₃). λ_{max} 232 (sh) (log ϵ 3.93); 259 (log ϵ 3.94); 302 (log ϵ 3.92); 328 (log ϵ 4.01) (EtOH).

Daudon, M. *et al.*, *J.O.C.*, 1976, **41**, 3275 (*isol, uv, ir, pmr, cmr, ms, struct, epimer*)

Scandomelonine

[59813-31-9]

S-121

C₄₁H₄₂N₄O₅ 670.807

Alkaloid from the stems and leaves of *Melodinus scandens* (Apocynaceae). Cryst. (Me₂CO). Mp 300° dec. [α]_D²² -25 (c, 1.0 in CHCl₃).

19-Epimer: Episcandomelonine

[59830-06-7]

C₄₁H₄₂N₄O₅ 670.807


Alkaloid from *Melodinus scandens* (Apocynaceae). Amorph. [α]_D²² +25 (c, 1.0 in CHCl₃).

Daudon, M. *et al.*, *J.O.C.*, 1976, **41**, 3275 (*isol, uv, ir, pmr, cmr, ms, struct, epimer*)

Scleratine

S-122

15,20-Dihydro-12,15,18-trihydroxy-14-methyl-21-norsenecanone-11,16-dione [6190-25-6]

C₁₈H₂₇NO₇ 369.414

Cyclic diester of Retronecine with Scleraneic acid. Struct. revised in 1985. Constit. of *Senecio sceleratus* and *Senecio latifolius* (Asteraceae). Cryst. (EtOH). Mp 176.5-177°. [α]_D²⁰ +55 (c, 1.09 in EtOH).

▶ VR1225000

N-Oxide: Scleratine N-oxide

[103184-92-5]

C₁₈H₂₇NO₈ 385.413

Alkaloid from *Senecio latifolius*. Cryst. (EtOH). Mp 152° dec. [α]_D²⁰ +48.4 (c, 1.27 in MeOH).

18-Deoxy, 18-chloro, N-oxide: Merenskine N-oxide

[96657-95-3]

C₁₈H₂₆ClNO₇ 403.859

Alkaloid from *Senecio latifolius* (Asteraceae). Solid + 1EtOH (EtOH). Mp 146° dec. [α]_D²⁰ +26.1 (c, 0.33 in EtOH).

18-Deoxy, 18-chloro, picrate: Mp 213-215°.

18-Deoxy, 18-chloro: Merenskine. Scleratinyl chloride. Chlorodeoxyscleratine

[96657-94-2]

[15215-30-2]

C₁₈H₂₆ClNO₆ 387.859

Alkaloid from *Senecio sceleratus* (now regrouped under *Senecio latifolius*) (Asteraceae). Mp 196° (194.5-195°). [α]_D^{20,4} +32.6 (c, 0.24 in EtOH). Revised struct.

de Waal, H.L. *et al.*, *CA*, 1944, **38**, 833 (*isol*)

de Waal, H.L. *et al.*, *J.C.S.*, 1963, 953

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1965,

18, 1625 (*pmr*)

Gordon-Gray, C.G. *et al.*, *J.C.S. (C)*, 1967,

781 (*Merenskine, isol*)

Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653


(cd)

Bredenkamp, M.W. *et al.*, *Tet. Lett.*, 1985, **26**,

929; 5721 (*Scleratine, Merenskine, ir, cmr,*
ms, cryst struct, oxides)**Sceletone**

S-123

1,2,3,3a,7,7a-Hexahydro-3a-(4-hydroxyphenyl)-1-methyl-6H-indol-6-one, 9CI [51934-31-7]

C₁₅H₁₇NO₂ 243.305

Minor alkaloid from *Sceletium namaquense* (Aizoaceae). Cardiotonic antiphogistic. Oil, purified by hplc. Opt. inactive.

Jeffs, P.W. *et al.*, *J.O.C.*, 1974, **39**, 2703 (*isol, uv, ir, pmr, cmr, ms, struct*)


Martin, N.H. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 1 (*ms*)

Sánchez, I.H. *et al.*, *Synth. Commun.*, 1983, **13**, 35 (*synth, deriv*)

Sceptrine

S-124

[79703-25-6]

C₂₂H₂₄Br₂N₁₀O₂ 620.305

Conts. 2 tautomeric aminoimidazole fragments. *Isol.* from the sponge *Agelas sceptrum*, *Agelas confifera*, *Agelas schmidtii*, *Agelas dispar*, *Agelas nakamurai*, *Agelas novaecaledoniae*, *Agelas longissima*, *Agelas clathrodes*, *Axinella* sp., *Hymeniacidon* sp. and from an unidentified Micronesian sponge. Exhibits antimicrobial, antiserotonergic, prophage induction, ichthyotoxic, adrenergic antagonist, serotonin antagonist and anti-muscarinic activities; somatostatin inhibitor, vasoactive intestinal peptide inhibitor and antihistamine. Cryst. + H₂O (H₂O) (as hydrochloride). Sol. MeOH; poorly sol.

Me₂CO, hexane, H₂O, EtOAc. Mp 215-225° dec. (hydrochloride). [α]_D -7.4 (c, 1.2 in MeOH). The elemental anal. of a dried sample of the hydrochloride required 1H₂O but the X-ray study indicated 3H₂O. λ_{max} 265 (ε 20900) (MeOH) (Derep). λ_{max} 203 (ε 27300); 214 (ε 28300); 267 (ε 25200) (MeOH) (Berdy).

2-Bromo: Bromosceptrine

[335214-63-6]

C₂₂H₂₃Br₃N₁₀O₂ 699.202

Isol. from *Agelas confifera*.

2,2'-Dibromo: Dibromosceptrine

[117417-71-7]

C₂₂H₂₂Br₄N₁₀O₂ 778.098

Isol. from *Agelas confifera* and *Agelas clathrodes*. Antifouling agent, neurotoxin. Sol. MeOH, CHCl₃, EtOAc; poorly sol. H₂O. [α]_D²⁵ -44 (c, 0.108 in MeOH) (as di-Ac). λ_{max} 202 (ε 27300); 215 (ε 24800); 273 (ε 26300) (MeOH) (Derep).

3'-Debromo: Debromosceptrine

[117442-38-3]

C₂₂H₂₅BrN₁₀O₂ 541.409

Isol. from *Agelas confifera*. [α]_D²⁵ -30 (c, 1.03 in MeOH) (as di-Ac). λ_{max} 203 (ε 25700); 212 (ε 26700); 265 (ε 25600) (MeOH) (Derep).

Bis(debromo): Didebromosceptrine

[217807-68-6]

[217633-87-9 (dihydrochloride)]

C₂₂H₂₆N₁₀O₂ 462.513

Isol. from *Agelas confifera*. Amorph. pale yellow powder (as dihydrochloride). [α]_D +53.8 (c, 0.36 in EtOH) (dihydrochloride). λ_{max} 267 (log ε 3.5) (EtOH) (dihydrochloride).

Walker, R.P. *et al.*, *J.A.C.S.*, 1981, **103**, 6772-6773 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Albizzati, K.F. *et al.*, *J.O.C.*, 1985, **50**, 4163-4164 (*isol*)

Keifer, P.A. *et al.*, *J.O.C.*, 1991, **56**, 2965-2975; 6728 (*isol, pmr, cmr, struct, Dibromosceptrine*)

Shen, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1302-1303 (*Debromosceptrine, Didebromosceptrine*)

Assmann, M. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 157-160 (*Bromosceptrine*)

Birman, V.B. *et al.*, *Org. Lett.*, 2004, **6**, 2369-2371 (*synth*)


O'Malley, D.P. *et al.*, *J.A.C.S.*, 2007, **129**, 4762-4775; 7702 (*synth*)

Schelhammeridine

S-125

Alkaloid C†

[22373-00-8]

C₁₉H₂₁NO₃ 311.38

A major alkaloid from *Schelhammera pedunculata* (Liliaceae). Needles (Me₂CO). Mp 118°. [α]_D -108 (c, 0.53 in CHCl₃).

Picrate:

Yellow needles (EtOH). Mp 202-207° dec.

Methiodide: Mp 215-216°.

6ξ,7ξ-Epoxyde: Cycloxyisoschelhammericine

[66322-14-3]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Cephalotaxus hainanensis* and *Cephalotaxus sinensis* (Cephalotaxaceae).

8-Oxo: 8-Oxoschelhammeridine. Alkaloid K

[24204-40-8]

C₁₉H₁₉NO₄ 325.363

Minor alkaloid from *Schelhammera pedunculata* (Liliaceae). Needles (Me₂CO). Mp 170-171°. [α]_D +35 (c, 0.23 in CHCl₃).

11a-Oxo: 11a-Oxoschelhammeridine. 12-Oxoschelhammeridine, 8Cl. Alkaloid J†

[24204-39-5]

C₁₉H₁₉NO₄ 325.363

Minor alkaloid from *Schelhammera pedunculata* (Liliaceae). Cryst. (Me₂CO). Mp 151-153°. [α]_D -47 (c, 0.15 in CHCl₃).

3-Epimer: 3-Epischelhammeridine. Alkaloid G†

[24204-35-1]

C₁₉H₂₁NO₃ 311.38

Minor alkaloid from *Schelhammera pedunculata* (Liliaceae). Needles (Me₂CO). Mp 131-133°. [α]_D +24 (c, 0.04 in CHCl₃).

3-Epimer, 6S,7R-epoxyde: Fortunine

[196796-76-6]

Alkaloid from *Cephalotaxus fortunei*. May be identical with Comosimine above.

3-Epimer; 6ξ,7ξ-epoxyde: Comosimine

[31690-01-4]

C₁₉H₂₁NO₄ 327.379

Alkaloid from leaves of *Phelline comosa* (Phellinaceae). Mp 126°. [α]_D +63 (c, 1.8 in CHCl₃).

3-Epimer, 6ξ,7ξ-epoxyde; hydrochloride: Cryst. (MeOH/Et₂O). Mp 218-219°. [α]_D +71 (c, 1.6 in EtOH).

Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2187 (*isol, uv, pmr, ms, struct*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2219 (*oxo derivs*)

Langlois, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 3535 (*epoxyde*)

Battersby, A.R. *et al.*, *Tet. Lett.*, 1975, 3419 (*biosynth*)

Ren, L. *et al.*, *Zhongcaoyao*, 1981, **12**, 1; *CA*, **96**, 3672p (*epoxyde*)


Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 515 (*synth, Schelhammeridine, 3-Epischelhammeridine*)

Qiu, M. *et al.*, *CA*, 1997, **127**, 275355f (*Fortunine*)

Schelhammerine

S-126

Alkaloid D†
[21030-71-7]



Absolute
Configuration

C₁₉H₂₃NO₄ 329.395

Major alkaloid from *Schelhammera pedunculata* (Liliaceae). Prisms (Me₂CO).

Mp 173-174°. [α]_D +186 (c, 0.5 in CHCl₃).

Hydrobromide: Mp 276°.

Methiodide:

Prisms (MeOH). Mp 210-212°.

Ac:

Needles (MeOH). Mp 143-144°. [α]_D +242 (c, 0.15 in CHCl₃).

2-Deoxy: Schellhammericine. Alkaloid F†
[21030-78-4]

C₁₉H₂₃NO₃ 313.396

Minor alkaloid from *Schelhammera pedunculata* (Liliaceae). Needles (petrol). Mp 76-77°. [α]_D +122 (c, 0.08 in CHCl₃).

2-Deoxy, picrate:

Yellow cryst. (MeOH). Mp 169-172°.

3-Epimer: 3-Epischelhammerine. Alkaloid H†. Alkaloid 3. 2-Epihomoerythratine

[24204-38-4]

C₁₉H₂₃NO₄ 329.395

Minor alkaloid from *Schelhammera pedunculata*. Also isol. from leaves of *Phelline comosa* and from the twigs and foliage of *Athrotaxis cupressoides* (Liliaceae, Phellinaceae, Taxodiaceae). Cryst. (Me₂CO). Mp 184-185°. [α]_D¹⁹ +170 (c, 1.4 in CHCl₃).

3-Epimer, O-de-Me: Robustidine

[128508-52-1]

C₁₈H₂₁NO₄ 315.368

Minor alkaloid from the leaves of *Phelline comosa* var. *robusta* (Phellinaceae). Mp 200-202°. [α]_D +141 (c, 0.83 in CHCl₃).

3-Epimer, 2-deoxy: 3-Epischelhammericine. Alkaloid E†

[24204-36-2]

C₁₉H₂₃NO₃ 313.396

Major alkaloid of *Schelhammera multiflora*; a constit. of *Schelhammera pedunculata*, *Schelhammera undulata*, *Phelline comosa*, *Phelline* sp. aff. *Phelline lucida*, *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus hainanensis*, *Cephalotaxus wilsoniana*, *Cephalotaxus drupacea* and *Dysoxylum lenticellare* (Liliaceae, Phellinaceae, Cephalotaxaceae, Meliaceae). Noncryst. [α]_D +123 (c, 0.3 in CHCl₃).

3-Epimer, 2-deoxy, 1α,6α-epoxyde: O-Methylrobustimine

[128508-53-2]

C₁₉H₂₃NO₄ 329.395

Minor alkaloid from leaves of *Phelline comosa* var. *robusta* and *Phelline brachyphylla* (Phellinaceae). Mp 146-148°. [α]_D +113 (c, 0.66 in CHCl₃).

3-Epimer, 2-deoxy, 1α,6α-epoxyde, O-de-Me: Robustimine

[128533-18-6]

C₁₈H₂₁NO₄ 315.368

Minor alkaloid from leaves of *Phelline comosa* var. *robusta* and *Phelline brachyphylla* (Phellinaceae).

Mp 134-136°. [α]_D +94 (c, 0.76 in CHCl₃).

3-Epimer, 2-deoxy, 12ξ-hydroxy: 3-Epi-12-hydroxyschelhammericine

[91897-63-1]

C₁₉H₂₃NO₄ 329.395

Alkaloid from the leaves of *Dysoxylum lenticellare* (Meliaceae) and *Lagarostrobos colensoi* (*Dacrydium colensoi*). Cryst. (EtOAc). Mp 235-236°.

2,3-Diepimer: Homoerythratine
[84471-41-0]

C₁₉H₂₃NO₄ 329.395

Alkaloid from *Phelline brachyphylla*, *Athrotaxis cupressoides*, *Athrotaxis selaginoides* and *Athrotaxis laxifolia* (Phellinaceae). Cryst. (Et₂O). Mp 176°. [α]_D +75 (c, 0.65 in CHCl₃).

2,3-Diepimer, 1α,6α-epoxyde, Me ether: O-Methylphellinine

[82482-18-6]

C₂₀H₂₅NO₅ 359.421

Alkaloid from leaves of *Phelline brachyphylla* (Phellinaceae). Mp 202°. [α]_D +45.4 (c, 0.86 in CHCl₃).

2,3-Diepimer, 1α,6α-epoxyde, O³-de-Me, O²-Me: Phellinine

[82463-44-3]

C₁₉H₂₃NO₅ 345.394

Alkaloid from the leaves of *Phelline brachyphylla* (Phellinaceae). Cryst. (MeOH). Mp 230°. [α]_D +40.2 (c, 1.25 in CHCl₃).

Johns, S.R. *et al.*, *Chem. Comm.*, 1968, 1102-1104 (*cryst struct, pmr*)

Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2187-2201 (*Schelhammerine, Schelhammericine, isol, uv, ir, pmr, ms, struct*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2219-2231 (*Schelhammerine, 3-Epischelhammerine, Schelhammericine, uv, pmr, ms, struct, synth*)

Langlois, N. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1969, **269**, 639-641 (*Schelhammerine, 3-Epischelhammerine, Schelhammericine, occur, uv, pmr, ms, struct*)

Kowala, C. *et al.*, *Z. Kristallogr., Kristallgeom., Kristallphys., Kristallchem.*, 1969, **130**, 121-138; *CA*, **72**, 94254 (*cryst struct, abs config*)

Powell, R.G. *et al.*, *Phytochemistry*, 1972, **11**, 1467-1472 (*Schelhammericine, isol, uv, pmr, ms*)

Debourges, D. *et al.*, *J. Nat. Prod.*, 1982, **45**, 163-167 (*Homoerythratine, Phellinine, O-Methylphellinine*)

Aledesamni, A.J. *et al.*, *J. Chem. Res., Synop.*, 1984, 108 (*3-Epi-12-hydroxyschelhammericine*)

Panichanun, S. *et al.*, *Tetrahedron*, 1984, **40**, 2677-2684; 2685-2689 (*isol, uv, ir, pmr, ms, epimers*)

Tsuda, Y. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3574; 1996, **44**, 500


(*Schelhammericine, 3-Epischelhammericine, synth*)

Langlois, N. *et al.*, *Heterocycles*, 1990, **30**, 659-664 (*Robustidine, Robustimine, O-Methylrobustimine*)


Bloor, S.J. *et al.*, *Phytochemistry*, 1996, **41**, 801-802 (*3-Epi-12-hydroxyschelhammericine, isol*)

Schischkiniin

[865866-93-9]

Relative
ConfigurationC₂₆H₂₄N₆O₂ 452.515Alkaloid from the seeds of *Centaurea schischkini*. Cytotoxic and antioxidant. Gum. λ_{max} 220 ; 280 (MeOH).Shoeb, M. *et al.*, *Tetrahedron*, 2005, **61**, 9001-9006 (*isol, pmr, cmr*)**Schizanthine X**

[136945-84-1]

C₃₁H₄₄N₂O₈ 572.697Alkaloid from *Schizanthus grahamii* (Solanaceae). Oil. [α]_D²⁵ -20 (c, 0.92 in EtOH).6-O-Deangeloyl: **Schizanthine Y**

[222716-55-4]

C₂₆H₃₈N₂O₇ 490.595Alkaloid from *Schizanthus porrigens*. Oil. [α]_D²⁵ +8.1 (c, 0.43 in EtOH).6-O-Deangeloyl, 6-O-(3-methyl-2-butenoyl): **Schizanthine B**

[70474-25-8]

C₃₁H₄₄N₂O₈ 572.697Alkaloid from the leaves and stems of *Schizanthus pinnatus* (Solanaceae). Amorph. powder. [α]_D²⁵ -64.7 (c, 0.71 in CHCl₃).6-De(angeloyloxy): **Schizanthine C**


[109031-02-9]

C₂₆H₃₈N₂O₆ 474.596Alkaloid from *Schizanthus grahamii* (Solanaceae). Oil. [α]_D²⁵ -17.2 (c, 0.32 in EtOH).Δ^{2'',3''}-Isomer, 6-O-deangeloyl: **Schizanthine Z**

[222716-56-5]

C₂₆H₃₈N₂O₇ 490.595Alkaloid from *Schizanthus porrigens*.Oil. [α]_D²⁵ +11.1 (c, 0.24 in EtOH).Ripperger, H. *et al.*, *Phytochemistry*, 1979, **18**, 171 (*Schizanthine B*)San-Martin, A. *et al.*, *Phytochemistry*, 1987, **26**, 819 (*Schizanthine C*)Muñoz, O. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1094 (*Schizanthine X*)Muñoz, O. *et al.*, *Pharm. Biol.*, 1998, **36**, 162-166 (*Schizanthine Y*, *Schizanthine Z*)**Schizanthine A**


[70474-24-7]

Absolute
configuration

S-127

C₂₀H₂₉NO₆ 379.452Alkaloid from the leaves and stems of *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D²⁵ -24.6 (c, 0.92 in CHCl₃).Ripperger, H. *et al.*, *Phytochemistry*, 1979, **18**, 171 (*isol, uv, ir, pmr, ms, struct*)**Schizanthine E**


[109031-04-1]

C₂₆H₃₈N₂O₇ 490.595Alkaloid from *Schizanthus grahamii* (Solanaceae). Oil. [α]_D²⁵ -2.3 (c, 0.086 in EtOH).Δ^{2,3}-Isomer: **Schizanthine D**

[109031-03-0]

C₂₆H₃₈N₂O₇ 490.595Alkaloid from *Schizanthus grahamii* (Solanaceae). [α]_D²⁵ -9.7 (c, 0.584 in EtOH).San-Martín, A. *et al.*, *Phytochemistry*, 1987, **26**, 819 (*isol, ir, pmr, cmr, ms, struct*)**Schizanthine G**

[119736-74-2]

C₁₉H₂₇NO₆ 365.425Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -11.4 (c, 0.23 in EtOH).Et ester analogue: **Schizanthine M**

[119736-78-6]

C₂₀H₂₉NO₆ 379.452Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -8.5 (c, 0.06 in EtOH). Poss. artifact.2'Z-Isomer: **Schizanthine H**

[119736-75-3]

C₁₉H₂₇NO₆ 365.425Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -12.5 (c, 0.1 in EtOH).2'Z-Isomer, Et ester analogue: **Schizanthine L**

[119736-77-5]

C₂₀H₂₉NO₆ 379.452Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -7.7 (c, 0.1 in EtOH).Δ^{2'',3''}-Isomer: **Schizanthine F**

[119736-73-1]

C₁₉H₂₇NO₆ 365.425Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -13.3 (c, 0.16 in EtOH).Δ^{2'',3''}-Isomer, 2'Z-isomer: **Schizanthine I**

[119763-87-0]

C₁₉H₂₇NO₆ 365.425Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -13.2 (c, 0.18 in


EtOH).

Δ^{2'',3''}-Isomer, Et ester analogue: **Schizanthine K**

[119736-76-4]


C₂₀H₂₉NO₆ 379.452Alkaloid from *Schizanthus pinnatus* (Solanaceae). Oil. [α]_D -12.1 (c, 0.24 in EtOH). Poss. artifact.de la Fuente, G. *et al.*, *Heterocycles*, 1988, **27**, 1887 (*isol, ir, pmr, cmr, ms, struct*)**Schizocommunine**

S-132

1,3-Dihydro-3-[(4-hydroxy-3-cinnolinyl)-methylene]-2H-indol-2-one, 9CI
[272119-70-7]C₁₇H₁₁N₃O₂ 289.293Prod. by *Schizophyllum commune* *isol.* from a human with allergic bronchopulmonary mycosis. Cytotoxic.Hosoe, T. *et al.*, *Mycopathologia*, 1999, **146**, 9-12**Schizogaline**

S-133


[2671-28-5]

Absolute
ConfigurationC₂₀H₂₂N₂O₂ 322.406Alkaloid from *Schizogygia coffaeoides* (Apocynaceae). Cryst. (Et₂O). Mp 156-157°. [α]_D²⁵ +28.2 (c, 1 in CHCl₃).10-Methoxy: **Schizogamine**

[2671-27-4]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from *Schizogygia coffaeoides* (Apocynaceae). Cryst. (petrol). Mp 123-125°. [α]_D²⁵ -7.9 (c, 1 in CHCl₃).Renner, U. *et al.*, *Experientia*, 1963, **19**, 244-246 (*cd, ir, uv, pmr*)Renner, U. *et al.*, *J. Nat. Prod.*, 1964, **27**, 406-415 (*struct*)Stephens, P.I. *et al.*, *J.O.C.*, 2007, **72**, 2508-2524 (*cd, abs config*)**Schizokinen A†**

S-134

N,1-Bis[3-(acetylhydroxyamino)propyl]-3-hydroxy-2,5-dioxo-3-pyrrolidineacetamide, 9CI
[83948-77-0]

$C_{16}H_{26}N_4O_8$ 402.403
Isol. from *Bacillus megaterium* ATCC 19213. Siderophore. Powder. Not the same as Schizokinen.


Mullis, K.B. *et al.*, *Biochemistry*, 1971, **10**, 4894-4898 (*isol, struct*)
Lee, B.H. *et al.*, *J.O.C.*, 1983, **48**, 24-31 (*synth, struct*)
Budzikiewicz, H. *et al.*, *Z. Naturforsch.*, **C**, 1997, **52**, 496-503 (*isol, synth, pmr, cmr*)

Schizoluteine S-135

Struct. unknown. Mol. formula not recorded. Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Mp 210-212°. Renner, U. *et al.*, *Experientia*, 1963, **19**, 244-246 (*isol, uv, ir*)

Schizophylline S-136

[2447-60-1]




Relative Configuration

$C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Cryst. (MeOH). Mp 129-130°. $[\alpha]_D^{24}$ -64 (c, 1 in $CHCl_3$).

Renner, U. *et al.*, *Experientia*, 1963, **19**, 244 (*cd, ir, uv*)
Renner, U. *et al.*, *J. Nat. Prod.*, 1964, **27**, 406 (*synth, struct*)

Schizozygine S-137

[2047-63-4]



Absolute Configuration

$C_{20}H_{20}N_2O_3$ 336.39
Alkaloid from the leaves and roots of *Schizozygia coffaeoides* (Apocynaceae). Cryst. (Et₂O). Mp 192-194°. $[\alpha]_D^{23}$ +15.5 (c, 1 in $CHCl_3$).

19β-Hydroxy, 6,7-didehydro: 6,7-Didehydro-19-hydroxyschizozygine. 6,7-Dehydro-19-hydroxyschizozygine
 $C_{20}H_{18}N_2O_4$ 350.373
Alkaloid from the leaves of *Schizozygia coffaeoides*. Amorph. powder. Mp 212-214°. λ_{max} 250 (log ϵ 3.78); 254 (log ϵ 4.03); 260 (log ϵ 4.12); 263 (log ϵ 4.12); 272 (log ϵ 3.81); 320 (log ϵ 3.03) ($CHCl_3$).

14,15-Dihydro, 15ξ-hydroxy: α-Schizozygol

[2772-65-8]
 $C_{20}H_{22}N_2O_4$ 354.405
Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Cryst. (MeOH). Mp 210-211°. $[\alpha]_D^{24}$ +51.3 (c, 1 in $CHCl_3$). pK_a 4.8.

Renner, U. *et al.*, *Experientia*, 1963, **19**, 244-246 (*cd, uv, ir*)

Renner, U. *et al.*, *J. Nat. Prod.*, 1964, **27**, 406-415 (*α-Schizozygol*)
Renner, U. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 308-317 (*struct, pmr*)
Hesse, M. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 1875-1899 (*ms*)
Kariba, R.M. *et al.*, *J. Nat. Prod.*, 2002, **65**, 566-569 (*Dehydrohydroxyschizozygine*)
Stephens, P.J. *et al.*, *J.O.C.*, 2007, **72**, 2508-2524 (*cd, abs config*)


β-Schizozygol S-138

$C_{20}H_{22}N_2O_4$ 354.405
Struct. unknown. Alkaloid from *Schizozygia coffaeoides* (Apocynaceae). Mp 247-250°.

Renner, U. *et al.*, *Experientia*, 1963, **19**, 244-246 (*isol, uv, ir*)

Schoberidine, 9CI S-139

[56775-83-8]




Probable Relative configuration

$C_{20}H_{21}N_3$ 303.406
Alkaloid from *Nitraria schoberi* (Zygophyllaceae). Yellow cryst. Mp 204-205°.
Hydrochloride: Mp 267°.

Ibragimov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 275; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 297 (*isol, uv, ms, struct*)

Schoberine S-140

Dodecahydro-1H,4H,9H-dipyrido[2,1-b:3',2',1'-ij]quinazoline, 9CI
[63653-27-0]




$C_{15}H_{26}N_2$ 234.384
Alkaloid from *Myrioneuron nutans*, *Nitraria schoberi* and *Nitraria komarovii*. Cryst. (MeOH). Mp 62-63°. $[\alpha]_D^{20}$ -12.2 (c, 1 in MeOH). Racemates isol. from *Nitraria* spp.

1,2-Didehydro: Dehydroschoberine

[169564-18-5]
 $C_{15}H_{24}N_2$ 232.368
Alkaloid from *Nitraria komarovii*. Oil.
Ibragimov, A.A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 71-74 (*isol*)
Tashkhodzhaev, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 24-28 (*cryst struct*)
Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 31-34 (*Dehydroschoberine*)
Pham, V.C. *et al.*, *J.O.C.*, 2008, **73**, 7565-7573 (*isol, synth, pmr, cmr, ms, abs config*)

Scholarisine A S-141

[1002321-69-8]




Absolute Configuration

$C_{19}H_{18}N_2O_2$ 306.363
Alkaloid from the leaves of *Alstonia scholaris*. Powder. $[\alpha]_D^{20}$ +188 (c, 0.55 in MeOH). λ_{max} 220 (ϵ 2370); 268 (ϵ 604) (MeOH).

Cai, X.-H. *et al.*, *Org. Lett.*, 2008, **10**, 577-580 (*isol, pmr, cmr*)

Schulzeine A S-142



R = CH₃
Absolute Configuration

$C_{42}H_{72}N_2O_{16}S_3$ 957.233
Alkaloid from the marine sponge *Penares schulzei*. α -Glucosidase inhibitor. Powder (as tri-Na salt). $[\alpha]_D^{22}$ +40 (c, 0.1 in MeOH) (tri-Na salt). λ_{max} 209 (ϵ 27500); 284 (ϵ 1390) (MeOH) (tri-Na salt).

Takada, K. *et al.*, *J.A.C.S.*, 2004, **126**, 187-193 (*isol, cd, pmr, cmr, ms*)

Schulzeine C S-143

As Schulzeine A, S-142 with R = H

$C_{41}H_{70}N_2O_{16}S_3$ 943.206
Alkaloid from the marine sponge *Penares schulzei*. α -Glucosidase inhibitor. Powder (as tri-Na salt). $[\alpha]_D^{22}$ +33 (c, 0.1 in MeOH) (tri-Na salt). λ_{max} 209 (ϵ 27800); 284 (ϵ 1500) (MeOH) (tri-Na salt).

11b-Epimer: Schulzeine B


$C_{41}H_{70}N_2O_{16}S_3$ 943.206
Alkaloid from *Penares schulzei*. α -Glucosidase inhibitor. Powder (as tri-Na salt). $[\alpha]_D^{22}$ -23 (c, 0.1 in MeOH) (tri-Na salt). λ_{max} 209 (ϵ 20800); 281 (ϵ 1500) (MeOH) (tri-Na salt).

Takada, K. *et al.*, *J.A.C.S.*, 2004, **126**, 187-193 (*isol, pmr, cmr, ms*)

Gurjar, M.K. *et al.*, *J.O.C.*, 2007, **72**, 6591-6594 (*synth*)

Schumagnine S-144


[112448-53-0]



$C_{15}H_{15}NO_4$ 273.288
Alkaloid from *Schumanniphyton magnificum* (Rubiaceae). Cream amorph. solid.

N-Me: N-MethylschumannifineC₁₆H₁₇NO₄ 287.315Alkaloid from *Schumanniohyton magnificum* (Rubiaceae). Cream amorph. solid.Houghton, P.J. *et al.*, *Planta Med.*, 1987, **53**, 262 (*isol, uv, ir, pmr, ms, struct, deriv*)**Schumannifine***Schumannioficine*


[88720-91-6]

S-145C₁₆H₁₅NO₆ 317.298Struct. revised in 1987 and again in 1995. Alkaloid from the root bark of *Schumanniohyton magnificum* (Rubiaceae). Anti-HIV activity. Mp 234°. λ_{max} 220; 255; 260; 280; 290; 296; 310; 320; 333 (MeOH) (Berdy).**N-Me: N-Methylschumannifine**

[88720-92-7]

C₁₇H₁₇NO₆ 331.324Alkaloid from the root bark of *Schumanniohyton magnificum* (Rubiaceae). Cryst. + 1MeOH. Mp 208-209°. λ_{max} 220 (ε 13182); 225 (ε 13182); 253 (ε 7762); 260 (ε 7940); 277 (ε 7800); 290 (ε 7940); 310 (ε 8912); 320 (ε 9200) (MeOH) (Berdy).**7'-Deoxy, 3',7'-didehydro: Anhydroschumannifine**C₁₆H₁₃NO₅ 299.282Alkaloid from the root bark of *Schumanniohyton magnificum* (Rubiaceae). Yellowish amorph. solid. Struct. revised in 1987 and again in 1995. λ_{max} 224 (ε 12020); 253 (ε 20900); 310 (ε 250) (MeOH) (Berdy).**7'-Deoxy, 3',7'-didehydro, N-Me: N-Methylanhydroschumannifine**C₁₇H₁₅NO₅ 313.309Alkaloid from the root bark of *Schumanniohyton magnificum* (Rubiaceae). Yellow cryst. (CHCl₃). λ_{max} 224 (ε 15848); 246 (ε 12600); 256 (ε 12600); 310 (ε 200) (MeOH) (Berdy).Okogun, J.I. *et al.*, *Planta Med.*, 1983, **49**, 95 (*isol, uv, ir, pmr, ms*)Houghton, P.J. *et al.*, *Planta Med.*, 1995, **61**, 154 (*struct*)**Schumanniohytine**


[69735-24-6]

S-146C₁₆H₉NO₅ 295.251Alkaloid from the root bark of *Schumanniohyton problematicum* and *Schumanniohyton magnificum* (Rubiaceae). Anti-HIV agent. Cryst. (AcOH). Mp 284-286°. λ_{max} 225 (ε 25118); 237 (ε 25700); 251 (ε 28840); 256 (ε 28180); 292 (ε 11748); 318 (ε 12880) (MeOH) (Berdy).


Ac: Mp 243-245°.

N-Me: N-MethylschumanniohytineC₁₇H₁₂NO₅[⊕] 310.285Quaternary alkaloid from *Schumanniohyton magnificum* (Rubiaceae). Anti-HIV agent. Yellow prismatic cryst. (as chloride); cryst. + 1H₂O (as iodide). Mp 286° (iodide). Confusingly, the struct. of *N*-Methylschumanniohytine is shown by Houghton *et al.* as the *N*-methyl deriv. of the now discarded structure for Schumanniohytine which is now allocated to Isoschumanniohytine, 1-314.Schlittler, E. *et al.*, *Tet. Lett.*, 1978, 2911 (*uv, ir, pmr, ms, struct*)Houghton, P.J. *et al.*, *Planta Med.*, 1985, **51**, 23; 1987, **53**, 262 (*isol, uv, ir, pmr, ms, struct, deriv*)Kelly, T.R. *et al.*, *J.O.C.*, 1992, **57**, 1593 (*synth, pmr, ms, struct*)Macklin, T.K. *et al.*, *Eur. J. Org. Chem.*, 2008, 1507-1509 (*synth*)**Schweinine**

[133084-04-5]

S-147C₃₀H₄₀N₄O₅ 536.67Alkaloid from the whole plant of *Schweinfurthia papilionacea* (Scrophulariaceae). Amorph. Mp 263° dec. [α]_D +61.7 (c, 0.65 in MeOH). Related to Aphelandrine, A-1334.Ahmad, V.U. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1162-1167 (*isol, pmr, cmr, struct*)**Sciadenine**


[59043-23-1]

S-148C₃₇H₄₀N₂O₆ 608.733Alkaloid from *Sciadotenia toxifera* (Menispermaceae). Needles (CHCl₃/Me₂CO). Mp 254-256° dec. [α]_D³⁰ -43 (c, 1.22 in Py). [α]_D²⁵ +15 (c, 1.185 in CHCl₃).**Me ether: Isocycleanine. O-Methylsciadenine**

[18210-67-8]

C₃₈H₄₂N₂O₆ 622.76Alkaloid from the roots of *Cyclea sutchuenensis*. Powder. Mp 262-263° dec. (semisynthetic). Opt. inactive (meso-compd.).*Et ether*: Mp 233-234° dec.Takahashi, K. *et al.*, *Heterocycles*, 1976, **4**, 471 (*Sciadenine, O-Methylsciadenine*)Lai, S. *et al.*, *Yaoxue Xuebao*, 1993, **28**, 599-603 (*Isocycleanine*)**Sciadoline**

[62404-95-9]

S-149C₃₆H₃₄N₂O₆ 590.674Alkaloid from *Sciadotenia toxifera* (Menispermaceae). Needles (CHCl₃/Me₂CO). Mp 225-228° dec. [α]_D²² +46 (c, 0.46 in CHCl₃).


Ac: Mp 236-237°.

Et ether: Mp 192-194° dec.**3,4-Dihydro: Sciadoferine**

[68676-59-5]

C₃₆H₃₆N₂O₆ 592.69Alkaloid from the stem wood of *Sciadotenia toxifera* (Menispermaceae). Cryst. (EtOAc). Mp 188-189°. [α]_D²⁰ +84.7 (c, 1.2 in CHCl₃).**α-Hydroxy: Cavanine**C₃₆H₃₄N₂O₇ 606.674Alkaloid from *Sciadotenia toxifera*. Needles (MeOH/CH₂Cl₂). λ_{max} 242 (log ε 4.1); 259 (sh) (log ε 4.03); 276 (log ε 3.77); 290 (sh) (log ε 3.09); 328 (sh) (log ε 3.5); 336 (log ε 3.58) (EtOH aq.).Takahashi, K. *et al.*, *Heterocycles*, 1976, **5**, 367 (*uv, ms, pmr, struct*)Galeffi, C. *et al.*, *Gazz. Chim. Ital.*, 1978, **108**, 97 (*Sciadoferine*)Menachery, M.D. *et al.*, *Tet. Lett.*, 2000, **41**, 2843-2846 (*Cavanine*)**Sciadole**


[157536-57-7]

S-150C₂₂H₂₆N₂O₂ 350.46

Alkaloid from fruit bodies of *Tricholoma sciodes*. Solid. Mp 70–75°. [α]_D +44 (c, 1.5 in CDCl₃). λ_{\max} 222 (ϵ 36000); 275 (ϵ 16500) (EtOH) (Derep).

Sterner, O. *et al.*, *Nat. Prod. Lett.*, 1994, **4**, 9–14 (*isol, uv, ir, pmr, cmr, ms*)

Scleritodermin A S-151
[663597-92-0]



C₄₂H₅₅N₇O₁₃S₂ 930.068


Stereochem. revised in 2008. Isol. from the sponge *Scleritoderma nodosum*. Cytotoxic. Off-yellow powder (as Na salt). [α]_D -41 (c, 0.1 in MeOH) (Na salt). λ_{\max} 200 (ϵ 26500); 305 (ϵ 11300) (MeOH) (Na salt).

Schmidt, E.W. *et al.*, *J. Nat. Prod.*, 2004, **67**, 475–478 (*isol, pmr, cmr*)

Liu, S. *et al.*, *Org. Lett.*, 2008, **10**, 3765–3768 (*synth, abs config*)

Sclerominol S-152

5-Benzyl-3-benzylidene-1-hydroxy-2,6-(1H,3H)-pyrazinedione



C₁₈H₁₄N₂O₃ 306.32

Isol. from hypovirulent isolates of *Sclerotinia minor*. Also obt. from debilitated isolates of *Sclerotinia sclerotiorum*. Cryst. (MeOH). λ_{\max} 246 (sh); 251; 260 (sh); 372 (CHCl₃).


α -Hydroxy(E-): 5-Benzyl-1-hydroxy-3-(hydroxybenzylidene)-2,6-(1H,3H)-pyrazinedione [915972-36-0]

C₁₈H₁₄N₂O₄ 322.32

Prod. by *Menisporopsis theobromae* BCC 3975. Antimalarial agent; cytotoxic. Yellow needles (MeOH). Mp 168–171°. λ_{\max} 204 (log ϵ 4.6); 251 (log ϵ 4.2); 365 (log ϵ 4.3) (MeOH).

Savard, M.E. *et al.*, *J. Nat. Prod.*, 2003, **66**, 306–309 (*isol, pmr, cmr, ms, cryst struct*)
Chinworrungsee, M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1404–1410 (α -hydroxy)

Sclerotiamide S-153



(-)-form

C₂₆H₂₉N₃O₅ 463.532

(-)-form [184025-39-6]

Isol. from the sclerotia of *Aspergillus sclerotiorum* NRRL 5167. Exhibits potent antiinsectan activity. Larvicide. Feeding deterrent. Mp 239–242° dec. [α]_D -55.1 (c, 0.1 in MeOH). λ_{\max} 215 (ϵ 5100); 250 (ϵ 8700); 280 (ϵ 3700) (MeOH).

10-Deoxy: **Notoamide B**

[937791-52-1]

C₂₆H₂₉N₃O₄ 447.533

Isol. from a marine-derived *Aspergillus* sp. [α]_D -118 (c, 0.06 in MeOH). λ_{\max} 246 (log ϵ 4.3); 283 (sh) (log ϵ 3.8); 294 (sh) (log ϵ 3.6); 330 (sh) (log ϵ 3.1) (MeOH).

10-Deoxy, N¹-hydroxy: **Notoamide A**

[937810-92-9]

C₂₆H₂₉N₃O₅ 463.532

Prod. by a marine-derived *Aspergillus* sp. [α]_D -112 (c, 0.08 in MeOH). λ_{\max} 247 (log ϵ 4.3); 283 (sh) (log ϵ 3.9); 294 (sh) (log ϵ 3.7); 330 (sh) (log ϵ 3.2) (MeOH).

3,19-Diepimer, 10-deoxy: **Versicolamide B**

C₂₆H₂₉N₃O₄ 447.533

Prod. by *Aspergillus versicolor* NRRL 35600. [α]_D +26 (c, 0.1 in Me₂CO).

(+)-form

10-Deoxy: (+)-**Notoamide B**

C₂₆H₂₉N₃O₄ 447.533

Prod. by *Aspergillus versicolor* NRRL 35600. [α]_D +102 (c, 0.05 in MeOH).

Whyte, A.C. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1093–1095 (*isol, uv, ir, pmr, cmr, ms, struct*)


Kato, H. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2254–2256 (*Notoamides A,B*)

Grubbs, A.W. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 2257–2261; 2262–2265 (*Notoamide B, synth*)

Greshock, T.J. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 3573–3577 (*Versicolamide B, (+)-Notoamide B*)

Sclerotigenin S-154

6,7-Dihydroquinolino[3,2-a][1,4]benzodiazepine-5,13-dione, 9CI [65641-84-1]




C₁₆H₁₁N₃O₂ 277.282

Isol. from the sclerotia of *Penicillium sclerotigenum*. Antiinsectan. Solid. Mp 235–238° (natural) Mp 277–280° (synthetic). λ_{\max} 235 (ϵ 7800); 271 (ϵ 3700); 311 (ϵ 1600) (MeOH).

Joshi, B.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 650–652 (*isol, uv, ir, pmr, cmr, ms*)

Snider, B.B. *et al.*, *Tetrahedron*, 2001, **57**, 3301–3307 (*synth, pmr, cmr*)

Sclerotiorinamine S-155



C₂₁H₂₄ClNO₄ 389.878

The parent (not isol.) was represented as the 8-oxo-NH-tautomer.


Me ether: **8-O-Methylsclerotiorinamine**

C₂₂H₂₆ClNO₄ 403.904

Prod. by *Penicillium multicolor*. Antagonist of the Grb2-SH2 domain. Orange gum. [α]_D +102 (c, 0.15 in MeOH). Related to Isochromophilone VI, I-205. λ_{\max} 204 (ϵ 5600); 271 (ϵ 16500); 286 (ϵ 17800); 364 (ϵ 22500) (MeOH).

Nam, J.-Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1303–1305

Scolopendrine S-156




C₁₈H₁₇NO₈S 407.4

Isol. from the centipede *Scolopendra subspinipes*. Yellow powder. Mp 192–194°.

Noda, N. *et al.*, *Chem. Pharm. Bull.*, 2001, **49**, 930–931

Scopadonnine S-157

1,2,3,4-Tetrahydro-1-phenyl-1,4-naphthalenedicarboxylic acid bis(9-methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl) ester, 9CI, 1-Phenyl-1,4-tetraindicarboxylic acid discopine ester



C₃₄H₃₈N₂O₆ 570.684

Dimerisation prod. of Apohyoscine, A-1364. Prob. artifact. May be a minor congener of Apohyoscine, A-1364 in various plants.

(1*R*,4*R*)-form

cis-form. **β -Scopadonnine**

[6882-53-7]

Minor alkaloid from leaves of *Duboisia leichhardtii* (Solanaceae). Also isol. from seeds of *Datura innoxia*. Needles (EtOH), cryst. + $\frac{1}{2}$ H₂O. Mp 128° (hemihydrate) Mp 130–133°.

Perchlorate (1:2):

Cryst. + $\frac{1}{2}$ H₂O. Mp 210–220° dec.

Dipicrate: Mp 210–220° dec.

(1*R*,4*S*)-formtrans-form. **α -Scopadonnine**

[545-85-7]

Minor alkaloid from leaves of *Duboisia leichhardtii* (Solanaceae). Also isol from seeds of *Datura innoxia*. Prisms (EtOH). Mp 195-196° (191°). $[\alpha]_D^{20}$ 0 (c, 0.38 in EtOH).

Perchlorate (1:2):

Cryst. + 1½ H₂O. Mp 229-230° dec.

Dipicrate: Mp 325°.

Voigtländer, H.W. *et al.*, *Annalen*, 1959, **625**, 196 (struct)Voigtländer, H.W. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1959, **262**, 632Kagei, K. *et al.*, *Yakugaku Zasshi*, 1980, **100**, 216 (isol, uv, pmr, ms, cmr)Aripova, S.F. *et al.*, *Khim. Prir. Soedin.*, 1991, 532-537; *Chem. Nat. Compd. (Engl. Transl.)*, 1991, 464-468 (isol, cryst struct)**Scopine****S-158**

9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]-nonan-7-ol, 9*CI*. 6,7-Epoxytropan-3-ol, 8*CI*

[498-45-3]

C₈H₁₃NO₂ 155.196

Tropane numbering shown. Base obt. by mild hydrol. of Scopolamine, S-159, prob. present in scopolamine-yielding plants in traces. Needles (petrol). Mp 76°. Opt. inactive (*meso*-).

Hydrochloride: [85700-55-6]

Platelets (EtOH).

Picrate:

Leaflets. Mp 231°.

N-De-Me, O-(3-acetoxy-2-methylpropanoyl): [855784-82-6]

C₁₃H₁₉NO₅ 269.297Alkaloid from *Datura stramonium*.

O-(2-Hydroxy-3-phenylpropanoyl): 6,7-Epoxytrotine

C₁₇H₂₁NO₄ 303.357

Alkaloid from leaves of a *Datura candida* cultivar (Solanaceae). Waxy semicryst. Tentative struct.

O-Phenylacetyl: **Phenylacetylscopine**

[130288-54-9]

C₁₆H₁₉NO₃ 273.331Alkaloid from the seeds of *Datura ferox*.3-Ketone: **Scopinone**

[498-47-5]

C₈H₁₁NO₂ 153.18Needles and plates (Et₂O/petrol). Mp 65-67°.3-Epimer: **Pseudoscopine**. ψ -Scopine

[498-46-4]

C₈H₁₃NO₂ 155.196

Long needles. Mp 122-123°.

Willstätter, R. *et al.*, *Ber.*, 1923, **56**, 1079

(synth)

Fodor, G. *et al.*, *Chem. Ind. (London)*, 1956, 764 (synth)Heusner, A. *et al.*, *Chem. Ber.*, 1958, **91**, 2399-2405 (synth, Scopine, Pseudoscopine, Scopinone)Werner, G. *et al.*, *Tet. Lett.*, 1967, 1283 (synth)Mandava, N. *et al.*, *Can. J. Chem.*, 1968, **46**, 2761 (pmr)Schink, H.E. *et al.*, *J.O.C.*, 1991, **56**, 2769


(synth)

Griffin, W.J. *et al.*, *Phytochemistry*, 1992, **31**, 367 (6,7-Epoxytrotine)Vitale, A.A. *et al.*, *J. Ethnopharmacol.*, 1995, **49**, 81-89 (Phenylacetylscopine)Justice, D.E. *et al.*, *Tetrahedron*, 1996, **52**,

11977 (synth, Scopine, Pseudoscopine)

Doncheva, T. *et al.*, *CA*, 2005, **143**, 74802f (*Datura stramonium consti*)**Scopolamine****S-159**

9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]-non-7-yl α -(hydroxymethyl)benzeneacetate, 9*CI*. **Hyoscine**, **BAN**. **Scopolamine tropate**. **Oscine tropate**. **Escopolamina**. **Joscina**. **Transcop**. **Scopoderm**

C₁₇H₂₁NO₄ 303.357

Antimuscarinic agent. Analgesic. Anti-inflammatory, antispasmodic agent.

Smooth muscle relaxant. Used as a cycloplegic and mydriatic, antinauseant, e.g. in travel sickness, sedative, e.g. in preoperative medication. Banned for use in cosmetics in UK. Log P 0.26 (calc).

(-)-form [51-34-3]

Alkaloid from *Atropa*, *Datura*, *Hyoscyamus* and *Scopolia* spp. and several other genera in the Solanaceae. Coml. sources are *Datura metel*, *Datura meteloides* and *Datura fastuosa* var *alba*. Noncryst. $[\alpha]_D^{20}$ -18 (EtOH). $[\alpha]_D$ -20 (H₂O). Forms a hydrate, Mp 59°; racemises in alkaline soln. Pharmacol. active isomer.

► Adverse ocular effects. Absorbed through skin. Human CNS effects when used therapeutically (esp. drowsiness). Psychotic effects when administered transdermally. LD₅₀ (mus, orl) 1275 mg/kg. Exp. reprod. and teratogenic effects. VR3675000

Hydrochloride: [55-16-3]

Mp 200°.

► VR4290000

Hydrobromide: **Scopolamine hydrobromide**, **USAN**. **Hyosol**. **Scopamin**. **Scopos**. **Tranaxine**

[114-49-8]

Cycloplegic and mydriatic. Mp 193-194° (anhyd.). $[\alpha]_D$ -25.9 (H₂O). $[\alpha]_D$ -15.72 (EtOH). Component of Benacine, Donnagel, Donnatal, Donnazyme and Kinesed.

► LD₅₀ (rat, orl) 1270 mg/kg. Exp. reprod. and teratogenic effects. YM4550000

Picrate:

Yellow needles. Mp 187-188°.

Methobromide: **Hyoscine methobromide**,**BAN**. **Methscopolamine bromide**,**USAN**. **Pamine**

[155-41-9]

[13265-10-6]

Parasympatholytic and spasmolytic agent.

► LD₅₀ (rat, orl) 3400 mg/kg. YM3675000

N-Oxide: Hyoscine N-oxide 1C₁₇H₂₁NO₅ 319.357

Alkaloid from *Atropa*, *Datura*, *Hyoscyamus* and *Scopolia* spp. (Solanaceae). Mp 125-130° dec. (as hydrochloride). Equatorial O. The axial-O-isomer (obt. synthetically) was not detected.

N-De-Me: Norhyoscine. **Norscopolamine**

[4684-28-0]

C₁₆H₁₉NO₄ 289.33

Principal alkaloid of the corollas of *Datura suaveolens* (Solanaceae). Mp 205° (as hydrochloride).

N-De-Me; picrate:

Serrated needles (EtOH). Mp 232-233°.

N-Butyl, bromide: **Butylscopolammonium bromide**. **Hyoscine butyl bromide**. **Scopolamine butyl bromide**. **Buscopan**.**Scobutil**. **Scopolan**. Many other names

[149-64-4]

Anticholinergic, spasmolytic agent.

Mp 142-144°.

► LD₅₀ (mus, orl) 1170 mg/kg. Exp. por-

phyrinogen. Exp. reprod. effects.

YM3500000

(±)-form**Atrosine**

[138-12-5]

Alkaloid from *Scopolia atropoides* and *Duboisia leichhardtii* (Solanaceae). Noncryst. Forms a monohydrate, Mp 56-7°, and a dihydrate, Mp 37-9°.

Picrate: Mp 173.5-174.5°.

Hydrobromide: pK_a 7.55 (23°).

[92714-23-3, 6533-68-2]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 286C (ir)Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 1231A (nmr)Ladenburg, A. *et al.*, *Annalen*, 1880, **206**, 299 (isol)Dobó, P. *et al.*, *J.C.S.*, 1959, 3461 (synth)Johns, S.R. *et al.*, *Chem. Comm.*, 1965, 458 (pmr)Ivanov, V. *et al.*, *Farmatsiya (Sofia)*, 1967, **17**,18; *CA*, **68**, 29912p (synth, N-butyl)Joyce, C.R. *et al.*, *Life Sci.*, 1968, **7**, 533

(pharmacol)

Pauling, P. *et al.*, *Chem. Comm.*, 1969, 1001 (cryst struct)Fales, H.M. *et al.*, *Arch. Mass Spectral Data*, 1971, **2**, 654 (ms)Evans, W.C. *et al.*, *Phytochemistry*, 1972, **11**,

3293 (isol, Norhyoscine)

Greenblat, D.J. *et al.*, *N. Engl. J. Med.*, 1973, **288**, 1215 (rev, pharmacol)Simeral, L. *et al.*, *Org. Magn. Reson.*, 1974, **6**,

226 (cmr)

Innes, I.R. *et al.*, *Pharmacol. Basis Ther.*, 5th edn., 1975, 514 (rev, pharmacol)Phillipson, J.D. *et al.*, *Phytochemistry*, 1975,**14**, 999 (oxide)Shutt, L.E. *et al.*, *Anaesthesia*, 1979, **34**, 476 (rev, pharmacol)Clissold, S.P. *et al.*, *Drugs*, 1985, **29**, 189 (rev, activity)Seeger, R. *et al.*, *Dtsch. Apoth. -Ztg.*, 1986,**126**, 1930 (rev, chem, props, occur, tox,

pharmacol)


Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 4459; 4970; 6354

(synonyms)

Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1990, **19**, 477 (rev)

- Sarazin, C. *et al.*, *Magn. Reson. Chem.*, 1991, **29**, 291 (*pmr, cmr*)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 425
 Glaser, R. *et al.*, *J.O.C.*, 1999, **64**, 9217-9224 (*conformn*)
 Glaser, R. *et al.*, *Can. J. Chem.*, 2000, **78**, 212-223 (*cryst struct*)
 Tytgat, G.N. *et al.*, *Drugs*, 2007, **67**, 1343-1357 (*hyoscine butyl bromide, rev*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, HOT500; SBG000; SBG500; SBH500

Scopoline† **S-160**
 Hexahydro-4-methyl-2,5-methano-2H-furo[3,2-b]pyrrol-6-ol, 9CI. *Oscine* [487-27-4]




$C_8H_{13}NO_2$ 155.196
 Alkaloid from several *Datura* spp. and *Scopolia tangutica* (Solanaceae). Prod. in the hydrol. of Scopolamine, S-159 by rearr. Hygroscopic prisms (Et₂O). Mp 108-109°. Bp 248°. p*K*_a 8.2 (as nitrate). Racemic; has been resolved.

Hydrochloride: [21030-89-7]
 Plates (hydrate), prisms (anhyd.). Mp 273-274° (244-250° dec.).

- Hesse, O. *et al.*, *Annalen*, 1892, **271**, 100
 King, H. *et al.*, *J.C.S.*, 1919, **115**, 476 (*resoln*)
 Heusner, A. *et al.*, *Chem. Ber.*, 1954, **87**, 1063 (*struct*)
 Zeile, K. *et al.*, *Chem. Ber.*, 1957, **90**, 2809 (*synth*)
 Mandava, N. *et al.*, *Can. J. Chem.*, 1968, **46**, 2761 (*pmr*)
 Evans, W.C. *et al.*, *J.C.S.(C)*, 1968, 2775 (*ms, ir*)
 Bristol, M.L. *et al.*, *J. Nat. Prod.*, 1969, **32**, 123 (*isol*)
 Scheiber, P. *et al.*, *Magn. Reson. Chem.*, 1989, **27**, 1097 (*pmr, cmr*)
 Mann, J. *et al.*, *J.C.S. Perkin 1*, 1992, 787 (*synth*)


Scorazanone **S-161**
 4-Hydroxy-2,3-dimethoxybenzo[g]quinoline-5,10-dione, 9CI [126657-23-6]



$C_{15}H_{11}NO_5$ 285.256
 Alkaloid from the roots of *Goniothalamus scortechinii* (Annonaceae). Yellow needles (Et₂O). Mp 205.5-206.5°.

Din, L.B. *et al.*, *Phytochemistry*, 1990, **29**, 346 (*isol, ir, pmr, cmr, ms, struct*)


Scorodophlone A **S-162**
 Dihydro-6-(methylsulfonyl)-1,2,3-dithiazin-4(3H)-one [933770-81-1]



$C_4H_7NO_3S_3$ 213.302
 Constit. of the seeds of *Scorodophloeus zenkeri*.

Songue, J.L. *et al.*, *Bull. Chem. Soc. Ethiop.*, 2006, **20**, 173-176 (*isol, struct*)

Scorpioidine† **S-163**
 [80405-18-1]



$C_{20}H_{31}NO_6$ 381.468
 Alkaloid from *Myosotis scorpioides* (Boraginaceae). Oil. $[\alpha]_D^{24}$ -5.06 (c, 1.72 in MeOH). λ_{max} 218 (ε 15200) (MeOH) (Berdy).

*O*⁷-Ac: 7-Acetylscorpioidine [80405-17-0]


$C_{22}H_{33}NO_7$ 423.505
 Alkaloid from *Myosotis scorpioides* (Boraginaceae). Oil. $[\alpha]_D^{24.5}$ -8.17 (c, 1.18 in MeOH).

Diastereoisomer, N-oxide: Anadoline N-oxide. Anadoline [28513-29-3]

$C_{20}H_{31}NO_7$ 397.467
 Alkaloid from *Symphytum orientale* (Boraginaceae). Mp 186° dec. $[\alpha]_D^{22}$ +9.2 (c, 0.7 in CHCl₃). Epimeric with Scorpioidine at C2' or C3', i.e. a tiglate ester of a retronecine trachelanthate N-oxide (abs. config. of necic ester portion not known). Thus a deriv. of either Indicine or Intermedine.

- Ulubelen, A. *et al.*, *Tet. Lett.*, 1970, 2583 (*isol, Anadoline*)
 Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1975, **28**, 173 (*struct, Anadoline*)
 Resch, J.F. *et al.*, *J. Nat. Prod.*, 1982, **45**, 358 (*isol, uv, ir, pmr, cmr, ms*)

Scoulerine **S-164**
 Alkaloid HFI. Aurotensine



(*R*)-form

$C_{19}H_{21}NO_4$ 327.379
 Aurotensine was a partial racemate with lower Mp. Active against gram-positive and gram-negative bacteria.

(*R*)-form [6507-34-2]
 Alkaloid from *Corydalis tuberosa* and *Corydalis cava* (Papaveraceae). Mp 195° (vac.). $[\alpha]_D^{22}$ +284 (c, 0.82 in MeOH).

(*S*)-form [6451-73-6]
 Alkaloid from many *Corydalis* spp., *Erythrina orientalis*, *Bocconia frutescens*, *Glaucium* spp., *Eschscholtzia lobbii* and *Fumaria officinalis* (Fabaceae, Papaveraceae). Shows antiemetic and antitussive activities. Active against gram-positive and gram-negative bacteria. Fine grey needles (MeOH). Mp 204°. $[\alpha]_D^{28}$ -289 (c, 0.62 in MeOH). $[\alpha]_D^{26}$ -318 (c, 0.41 in MeOH).

► DR9808600

Hydrochloride: Mp 268-269°.

N-Me: see Cyclanoline, C-833

Di-Me ether: see Tetrahydropalmatine, T-212

(±)-form [6451-72-5]
 Alkaloid from *Corydalis* and *Fumaria* spp. Plates (MeOH), cryst. (EtOH). Mp 198-200° (183-185°).

Picrate:

Cryst. powder (EtOH). Mp 206° (200°) dec.

Gadamar, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1927, **265**, 675

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1936, **16**, 153; 1939, **17**, 57; 1940, **18**, 414; 1942, **20**, 49

Battersby, A.R. *et al.*, *J.C.S.(C)*, 1966, 1052 (*synth*)


Brochmann-Hanssen, G. *et al.*, *Tet. Lett.*, 1966, 2261 (*cd, ms, pmr*)

Kametani, T. *et al.*, *J.C.S.(C)*, 1967, 530 (*ir, synth*)

Abbasoglu, U. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 379-380 (*activity*)

Cheng, X. *et al.*, *Chem. Biodiversity*, 2008, **5**, 1335-1344 (*isol, pmr, cmr*)

Scutianine A **S-165**
Scutianine [144051-13-9]



Absolute Configuration

$C_{39}H_{47}N_5O_5$ 665.831
 Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Needles (EtOH aq. or CHCl₃/petrol). Mp 186-187°. $[\alpha]_D^{20}$ -399 (c, 0.15 in CHCl₃).

N-De-Me: Scutianine F N-Demethylscutianine A

[64309-19-9]


C₃₈H₄₅N₅O₅ 651.804Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Needles (EtOH aq.). Mp 208°. [α]_D²⁰ -132 (c, 0.02 in MeOH).**N^{pro}-Deacyl, N^{pro}-Me: Ceanothine B**

[19471-43-3]

C₂₉H₃₆N₄O₄ 504.628Alkaloid from the root bark of *Ceanothus americanus* (New Jersey tea) and *Ceanothus sanguineus* (Rhamnaceae). Needles (CHCl₃/Et₂O). Mp 238.5-240.5°. [α]_D²⁵ -293 (c, 0.68 in CHCl₃).Warnhoff, E.W. *et al.*, *Can. J. Chem.*, 1965, **43**, 2594 (*Ceanothine B, isol, ms, pmr*)Tschesche, R. *et al.*, *Chem. Ber.*, 1967, **100**, 323-334 (*Scutianine A*)Klein, F.K. *et al.*, *J.A.C.S.*, 1968, **90**, 3576 (*Ceanothine B, ms, struct*)Servis, R.E. *et al.*, *J.A.C.S.*, 1968, **90**, 4179 (*Ceanothine B, struct*)Tschesche, R. *et al.*, *Phytochemistry*, 1977, **16**, 1025-1028 (*Scutianine F*)Lagarias, J.C. *et al.*, *J. Nat. Prod.*, 1979, **42**, 663 (*Ceanothine B, isol, ms*)**Scutianine C**

[53766-26-0]

S-166



Absolute Configuration

C₃₁H₄₂N₄O₄ 534.697Various nomenclature and numbering schemes used. Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Needles (MeOH/cyclohexane). Mp 267°. [α]_D²⁰ -231 (c, 0.1 in CHCl₃).**10,11-Dihydro:**Needles (CHCl₃/hexane). Mp 297-299°. [α]_D²⁰ -102 (c, 0.1 in CHCl₃).**10,11-Dihydro, 11 ξ -hydroxy: Pandamine**

[10233-81-5]


C₃₁H₄₄N₄O₅ 552.712Alkaloid from *Panda oleosa* (Pandaecae). Cryst. (EtOAc). Mp 256°. [α]_D -103 (c, 0.5 in CHCl₃). Gross struct. only, no stereochem. Not interrelated with Scutianine C.**10,11-Dihydro, 11-hydroxy, hydrochloride:**Cryst. + 1H₂O. Mp 235° dec. [α]_D -77 (c, 1 in MeOH).**1' ξ -Hydroxy: Scutianine H**

[72047-69-9]

C₃₁H₄₂N₄O₅ 550.697Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Cryst. (CHCl₃/Et₂O). Mp 242-243°. [α]_D -233 (c, 0.1 in CHCl₃). Gross struct. only, no stereochem. Not interrelated with Scutianine C.Pais, M. *et al.*, *Ann. Chim. (Paris)*, 1966, **11**, 83 (*Pandamine*)Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 2274-2283 (*Scutianine C*)Morel, A.F. *et al.*, *Phytochemistry*, 1979, **18**, 473-477 (*Scutianines C,H*)Pais, M. *et al.*, *Phytochemistry*, 1979, **18**, 1869-1872 (*Pandamine, cmr*)Da Silva, U.F. *et al.*, *Phytochem. Anal.*, 1996, **7**, 20-23 (*abs config*)**Scutianine D**

[53766-27-1]

S-167



Absolute Configuration

C₃₄H₄₀N₄O₅ 584.714

Various changes in CAS nomenclature and numbering. That shown here is in accordance with one CAS scheme.

Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae) and from *Condalia buxifolia*, *Zizyphus jujuba* and *Zizyphus nummularia*. Needles (EtOH aq.). Mp 219-220°. [α]_D²⁰ -196 (c, 0.1 in CHCl₃).**O-Ac:**Cryst. (EtOH aq.). Mp 236°. [α]_D -163 (c, 0.1 in CHCl₃).**10,11-Dihydro:**Needles (EtOH aq.). Mp 256-258°. [α]_D²⁰ -74.5 (c, 0.1 in CHCl₃).**4' ξ -Hydroxy: Scutianine J**

[163136-11-6]

C₃₄H₄₀N₄O₆ 600.713Alkaloid from bark of *Scutia buxifolia* (Rhamnaceae). Amorph. solid. Stereochem. not determined.**7'-Deoxy: Scutianine B**

[35467-41-5]

C₃₄H₄₀N₄O₄ 568.714Alkaloid from the bark and roots of *Scutia buxifolia* and from *Condalia buxifolia*, *Discaria febrifuga*, *Melochia tomentosa* and *Waltheria douradahina*. Antibacterial. Needles (CH₂Cl₂/cyclohexane). Mp 248-250°. [α]_D²⁰ -296 (c, 0.1 in CHCl₃).**7-Epimer: Scutianine G**

[65494-36-2]

C₃₄H₄₀N₄O₅ 584.714Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Needles (EtOH aq.). Mp 162°. [α]_D²⁰ -112 (c, 0.02 in MeOH).**7,7'-Diepimer: Scutianine K**


[202394-59-0]

C₃₄H₄₀N₄O₅ 584.714Alkaloid from *Scutia buxifolia*. Cryst. (MeOH/Et₂O). Mp 215-217°. [α]_D²⁵ -20.9 (c, 0.1 in CHCl₃).**3,4,7,7'-Tetraepimer: Scutianine E**

[53797-27-6]


C₃₄H₄₀N₄O₅ 584.714Alkaloid from the bark of *Scutia buxifolia* (Rhamnaceae). Needles (EtOH aq.). Mp 121°. [α]_D²⁰ -22.2 (c, 0.1 in CHCl₃). Contains two D-amino acid residues.Tschesche, R. *et al.*, *Tet. Lett.*, 1971, 4405-4408 (*Scutianine B*)Tschesche, R. *et al.*, *Chem. Ber.*, 1974, **107**, 2274-2283 (*Scutianine D*)Merkuza, V.M. *et al.*, *Phytochemistry*, 1974, **13**, 1279-1282 (*Scutianine B*)Tschesche, R. *et al.*, *Phytochemistry*, 1977, **16**, 1817-1818 (*Scutianine G*)Morel, A.F. *et al.*, *Phytochemistry*, 1979, **18**, 473-477; 1998, **47**, 125-129 (*Scutianines D,K*)Menezes, A.S. *et al.*, *Phytochemistry*, 1995, **38**, 783-786 (*Scutianine J*)Da Silva, U.F. *et al.*, *Phytochem. Anal.*, 1996, **7**, 20-23 (*abs config*)Tan, N.H. *et al.*, *Chem. Rev.*, 2006, **106**, 840-895 (*bibl*)**Scytonemin†**

S-168

3,3'-Bis[(4-hydroxyphenyl)methylene]-[1,1'-bicyclopent[b]indole]-2,2'-(3H,3'H)-dione, 9CI
[152075-98-4]C₃₆H₂₀N₂O₄ 544.565Pigment from the sheaths of cyanobacteria *Calothrix* spp. and *Stigonema* sp. UV sunscreen. Antiproliferative agent. Yellow-green cryst. Mp > 325°.Proteau, P.J. *et al.*, *Experientia*, 1993, **49**, 825-829 (*isol, pmr, cmr*)Edwards, H.G. *et al.*, *Spectrochim. Acta A*, 2000, **56**, 193-200 (*Raman*)Stevenson, C.S. *et al.*, *J. Pharmacol. Exp. Ther.*, 2002, **303**, 858-866 (*activity*)**Scytonine**

[683771-38-2]


S-169

C₃₁H₂₂N₂O₆ 518.525Isol. from a terrestrial cyanobacterium *Scytonema* sp. Amorph. brown solid. λ_{\max} 207 (ε 38950); 225 (ε 37050); 270 (ε 22480) (MeOH).Bultel-Poncé, V. *et al.*, *J. Nat. Prod.*, 2004, **67**, 678-681 (*isol, pmr, cmr*)

Secukidine

S-170

2,15-Dihydroxy-6,21-secohetisan-6,13-dione, 9CI
[135626-91-4]



C₂₁H₂₇NO₄ 357.449
Alkaloid from *Aconitum szukini* (Ranunculaceae).

2-Ac: **Secukinine**

[135626-90-3]
C₂₃H₂₉NO₅ 399.486

Alkaloid from *Aconitum szukini* (Ranunculaceae).

2-O-(2-Methylbutanoyl), 15-Ac: **Secukitine**

[135626-92-5]
C₂₈H₃₇NO₆ 483.603

Alkaloid from *Aconitum szukini* (Ranunculaceae).

6-Deoxo, 2-ketone: **Delcarduchol**

[240121-83-9]
C₂₁H₂₇NO₃ 341.449

Alkaloid from *Delphinium carduchorum*.


Chen, D. et al., *CA*, 1991, **115**, 110581r
(*Secukinine, Secukitine*)

Mericli, A.H. et al., *Phytochemistry*, 1999, **51**, 337-340 (*Delcarduchol*)

Sebastianine A

S-171

[448264-64-0]



C₁₇H₉N₃O 271.278
Isol. from the Brazilian ascidian *Cystodytes dellechiaiei*. Cytotoxic. Amorph. yellow solid. λ_{max} 223 (log ε 4.56); 238 (log ε 4.49); 265 (log ε 4.36); 301 (log ε 4.11); 390 (log ε 3.85) (MeOH).


Torres, Y.R. et al., *J.O.C.*, 2002, **67**, 5429-5432
(*isol, pmr, cmr*)

Legentil, L. et al., *Tet. Lett.*, 2003, **44**, 2473-2475 (*synth*)

Sebastianine B

S-172

[448264-65-1]



Relative Configuration

C₂₂H₁₉N₃O₃ 373.41


Isol. from the Brazilian ascidian *Cystodytes dellechiaiei*. Cytotoxic. Pale yellow solid. λ_{max} 206 (log ε 4.57); 252 (log ε 4.58); 292 (log ε 4.23); 366 (log ε 4.19) (MeOH).

Torres, Y.R. et al., *J.O.C.*, 2002, **67**, 5429-5432
(*isol, cd, pmr, cmr*)

Secaloid A

S-173

[188834-49-3]



Relative Configuration

C₄₆H₅₁NO₂₄ 1001.901
Constit. of the pollen of rye (*Secale cereale*). Antitumour agent. Amorph. powder. Mp 178-186° dec. [α]_D +4.6 (c, 0.6 in MeOH). λ_{max} 280; 320 (sh) (H₂O).

6'''-O-β-D-Glucopyranoside: **Secaloid C**

[188834-50-6]

C₅₂H₆₁NO₂₉ 1164.043
Constit. of rye pollen. Antitumour agent. Tentative struct. assigned.

3-Epimer: **Secaloid B**

[188855-24-5]

C₄₆H₅₁NO₂₄ 1001.901
Constit. of the pollen of rye. Antitumour agent. Amorph. powder. Mp 192-201° dec. [α]_D +6.8 (c, 0.4 in MeOH). λ_{max} 280; 320 (sh) (H₂O).

3-Epimer, 6'''-O-β-D-glucopyranoside: **Secaloid D**

[188899-19-6]

C₅₂H₆₁NO₂₉ 1164.043
Constit. of rye pollen. Antitumour agent. Tentative struct. assigned.


[187988-51-8, 187988-50-7]

Jaton, J.-C. et al., *J. Nat. Prod.*, 1997, **60**, 356-360; 361-365

Secamine

S-174

[34304-16-0]



Probable Relative Configuration

C₄₂H₅₂N₄O₄ 676.897
The *R,R*- absolute config. is given in CAS but the evidence supporting this cannot be located. The evidence summarised here supports the *RS,SR*-config., i.e. racemic with *R,S* relative config. Alkaloid from the roots of *Rhazya stricta* (Apocynaceae). Noncryst. Opt. inactive.

15,20/15',20'-Dihydro: **Dihydrosecamine**

C₄₂H₅₄N₄O₄ 678.913
Alkaloid from *Rhazya stricta* (Apocynaceae), obt. only as a mixt. Noncryst.

15,15',20,20'-Tetrahydro: **Tetrahydrosecamine**

[33633-19-1]
C₄₂H₅₆N₄O₄ 680.929

Alkaloid from *Rhazya stricta*, *Amsonia elliptica*, *Amsonia tabernaemontana*, *Aspidosperma excelsum*, *Aspidosperma maregravianum* and *Rhazya orientalis* (Apocynaceae). Shows cytotoxic and antimicrobial activity. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{max} 220; 283; 292 (MeOH) (Berdy). λ_{max} 225; 272; 285; 294 (EtOH) (Berdy).

16-Hydroxy, 15,15',20,20'-tetrahydro: **16-Hydroxytetrahydrosecamine**

[88607-52-7]
C₄₂H₅₆N₄O₅ 696.928

Alkaloid from root bark of *Aspidosperma excelsum*. Shows antimicrobial activity. Sol. MeOH, Me₂CO; poorly sol. H₂O. λ_{max} 220; 276 (sh); 282; 291 (MeOH). λ_{max} 220; 282; 291 (MeOH) (Berdy).

16-De(methoxycarbonyl), 16-hydroxy, 15,15',20,20'-tetrahydro: **16-Hydroxy-16-decarbomethoxytetrahydrosecamine**

[88607-51-6]
C₄₀H₅₄N₄O₃ 638.892

From root bark of *Aspidosperma excelsum*. Shows antimicrobial activity. Sol. MeOH, Me₂CO; poorly sol. H₂O. λ_{max} 220; 276 (sh); 285; 292 (MeOH).

16'-De(methoxycarbonyl), 15,15',20,20'-tetrahydro: **Decarbomethoxytetrahydrosecamine**

[56842-20-7]
C₄₀H₅₄N₄O₂ 622.892

Alkaloid from *Amsonia tabernaemontana*, *Rhazya stricta*, *Aspidosperma maregravianum* and *Aspidosperma excelsum*. Antiseptic. Noncryst. Sol. Me₂CO; poorly sol. H₂O. [α]_D²⁰ 0 (EtOH). Log P 8.76 (uncertain value) (calc). λ_{max} 220; 283; 292 (MeOH) (Berdy).

Bis(demethoxycarbonyl), 15,15',20,20'-tetrahydro: **Didemethoxycarbonyltetrahydrosecamine**

[57905-97-2]
C₃₈H₅₂N₄ 564.856

Alkaloid from roots of *Rhazya stricta*. Amorph. [α]_D²⁰ +30 (CHCl₃). λ_{max} 224; 284; 290 (sh) (MeOH).

[88728-16-9]

Evans, D.A. et al., *Chem. Comm.*, 1968, 859-861 (*isol, ms, struct*)

Zsardon, B. et al., *Phytochemistry*, 1975, **14**, 1438-1439 (*Tetrahydrosecamine, Decarbomethoxytetrahydrosecamine*)

Cordell, G.A. et al., *J. Indian Chem. Soc.*, 1978, **55**, 1083-1091 (*config*)

Mukhopadhyay, S. et al., *J. Nat. Prod.*, 1981, **44**, 696-700 (*Tetrahydrosecamine, isol, uv, ir, pmr, ms*)


Verpoorte, R. et al., *Planta Med.*, 1983, **48**, 283-289 (*16-hydroxy derivis*)

Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1986, **49**, 1138-1139 (*Didemethoxycarbonyltetrahydrosecamine*)

Secantioquine

S-175

[93767-29-4]




C₃₇H₃₈N₂O₈ 638.716

Alkaloid from the bark of a *Pseudoxandra* sp. (Annonaceae). Amorph. [α]_D -15 (c, 1 in CHCl₃). First example of a secobisbenzylisoquinoline alkaloid to contain a biphenyl linkage.

Cortes, D. et al., *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1984, **298**, 591 (uv, ir, pmr, ms, cd, struct)

6,7-Secoagroclavine

S-176



C₁₆H₂₀N₂ 240.347

Natural-form [68156-97-8]

Minor alkaloid from *Claviceps purpurea*. Mp 132° (126-129°). Opt. rotn. not recorded.

N-Me: Mp 136° (132-134°).

(±)-form

Synthetic. Mp 202-205°.

Horwell, D.C. et al., *Phytochemistry*, 1979, **18**, 519 (isol, uv, ir, pmr, ms, struct)

Natsume, M. et al., *Heterocycles*, 1980, **14**,

1101 (synth, uv, ir, pmr, ms)

Porter, J.K. et al., *J. Agric. Food Chem.*, 1981, **29**, 653 (isol, synth)

Oppolzer, W. et al., *Tetrahedron*, 1983, **39**,

3695 (synth, uv, ir, pmr, ms)

Somei, M. et al., *Chem. Pharm. Bull.*, 1984, **32**,

5064; 1986, **34**, 677 (synth)

Hatanaka, N. et al., *Tet. Lett.*, 1986, **27**, 3169

(synth)

Somei, M. et al., *Heterocycles*, 1987, **26**, 2823

(synth)

Nakagawa, K. et al., *Heterocycles*, 1991, **32**,

873 (synth)


Yamada, F. et al., *Heterocycles*, 2007, **72**, 599-

620 (synth)

6,7-Secoangustilobine A

S-177

[123064-77-7]



C₂₀H₂₄N₂O₃ 340.421

Alkaloid from leaves of *Alstonia congenisis* (Apocynaceae). [α]_D +78 (c, 0.3 in CHCl₃).

N-De-Me: Nor-6,7-secoangustilobine A [112448-49-4]

C₁₉H₂₂N₂O₃ 326.394

Alkaloid from *Alstonia angustiloba* and *Alstonia pneumatophora* (Apocynaceae). [α]_D +83 (c, 0.3 in MeOH).

7-Formyl, N-de-Me: 4,6-Secoangustilobinal [112464-22-9]

C₂₀H₂₂N₂O₄ 354.405

Alkaloid from *Alstonia angustiloba* and *Alstonia pneumatophora* (Apocynaceae). [α]_D +28 (c, 0.23 in MeOH). Rel. config. only detd.


Zeches, M. et al., *J. Nat. Prod.*, 1987, **50**, 714 (4,6-Secoangustilobinal, Nor-6,7-secoangustilobine A)

Caron, C. et al., *Phytochemistry*, 1989, **28**, 1241 (isol, uv, ir, pmr, cmr, ms, struct)

Secobatzelline A

S-178

[247590-59-6]



C₁₀H₁₀ClN₃O₃ 255.66

Alkaloid from the sponge *Batzella* sp. Mp 169-170° (as di-Ac) Mp > 300° (blackens at 165-170°). [α]_D²⁴ -135 (c, 0.01 in MeOH). λ_{max} 203 (log ε 4.19); 233 (log ε 4.13); 322 (log ε 4) (MeOH).

4-Deimino, 4-oxo: Secobatzelline B

[247590-60-9]

C₁₀H₉ClN₂O₄ 256.645

Alkaloid from *Batzella* sp. Mp 170-171° (as di-Ac) Mp > 300° (blackens at 180-182°). [α]_D²⁴ -18 (c, 0.01 in MeOH). Prob. artifact. λ_{max} 203 (log ε 4.32); 245 (log ε 4); 335 (log ε 4.32) (MeOH).

Gunasekera, S.P. et al., *J. Nat. Prod.*, 1999, **62**,

1208-1211 (Secobatzellines A,B)


Shinkre, B.A. et al., *Synth. Commun.*, 2007, **37**,

2399-2409 (Secobatzelline B, synth)

Secocepharanthine

S-179

[89503-78-6]



C₃₇H₃₆N₂O₈ 636.7

Trace alkaloid from *Stephania sasakii* (Menispermaceae). Amorph. [α]_D -7.2 (c,

2.08 in CHCl₃).

α'-Alcohol: **Dihydrosecocepharanthine**

[89503-79-7]


C₃₇H₃₈N₂O₈ 638.716

Alkaloid from *Stephania sasakii*. Plates (MeOH). Mp 192-194°. [α]_D -1.67 (c, 0.12 in CHCl₃).

Kunitomo, J. et al., *Chem. Pharm. Bull.*, 1985, **33**, 135 (uv, ir, pmr, ms, cd, ord, struct, synth)

Secocularidine

S-180



C₂₀H₂₃NO₄ 341.406

Alkaloid from *Corydalis claviculata* (Papaveraceae). Prisms (MeOH). Mp 189-190°.

N-De-Me: Norssecocularidine

[110883-32-4]

C₁₉H₂₁NO₄ 327.379

Trace alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Noncryst.

Me ether: Secocularine

C₂₁H₂₅NO₄ 355.433

Alkaloid from *Sarcocapnos crassifolia* (Papaveraceae). Amorph.

Me ether, perchlorate:

Cryst. (EtOH). Mp 194-196°.

Me ether, N-de-Me: Norssecocularine

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Corydalis claviculata* (Papaveraceae).

Boente, J.M. et al., *Chem. Biol. Isoquinoline*

Alkaloids, Int. Symp., Phytochem. Soc.

Eur., Abstr. Pap., 1984, **7**

(Norssecocularine)

Boente, J.M. et al., *Tet. Lett.*, 1984, **25**, 889 (uv,

ir, pmr, ms, struct, synth)

Castedo, L. et al., *Heterocycles*, 1987, **26**, 591

(Norssecocularidine)

Tojo, E. et al., *Heterocycles*, 1988, **27**,


2367 (synth, Norssecocularidine,

Norssecocularine)

Secodaphniphylline

S-181

[28606-61-3]



Absolute Configuration

C₃₀H₄₇NO₃ 469.706

Alkaloid from the leaves and bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Cryst. (hexane). Mp 129-130° Mp 124-125° (synthetic). [α]_D²⁵ -50 (c, 0.35 in CHCl₃) (synthetic).

22S-Acetoxy: **Daphniteijsmine**. Acetoxy-secodaphniphylline

[55855-00-0]

C₃₂H₄₉NO₅ 527.743

Minor alkaloid from the fruits of *Daphniphyllum teijsmannii* (Daphniphyllaceae). Cryst. (hexane/EtOAc). Mp 228-232° (187-188°).

Sasaki, K. *et al.*, *J.C.S. (B)*, 1971, 1565 (*abs config*)

Toda, M. *et al.*, *Tetrahedron*, 1972, **28**, 1477 (*isol, ir, pmr, ms, struct*)


Yamamura, S. *et al.*, *Tet. Lett.*, 1974, **15**, 2849-2852; 3673-3676 (*isol, ir, ms, struct, Daphniteijsmine*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1992, **57**, 2566 (*synth, ir, pmr, cmr, ms*)

Morita, H. *et al.*, *Tetrahedron*, 1999, **55**, 12549-12556 (*isol, cmr*)

3,4-Seco-3,14-dehydrocubucraline S-182


[87596-50-7]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from the root bark of *Alstonia plumosa* (Apocynaceae). [α]_D²⁵ -21 (c, 1.0 in CHCl₃).

Jacquier, M.J. *et al.*, *Phytochemistry*, 1982, **21**, 2973 (*isol, uv, ir, pmr, ms, struct*)

12-Seco-14-deoxyparsonsia-nine S-183

C₂₁H₃₃NO₉ 443.493

Me ester: [136997-63-2]


C₂₂H₃₅NO₉ 457.52

Alkaloid from leaves of *Parsonsia laevigata* (Apocynaceae). Solid. [α]_D³¹ +31.6 (c, 0.70 in MeOH). Possibly an artifact.

Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1576 (*isol, pmr, cmr, struct*)

Secodine S-184

Methyl 3-[2-(3-ethyl-5,6-dihydro-1(2H)-pyridinyl)ethyl]-α-methylene-1H-indole-2-acetate, 9CI
[27825-42-9]

C₂₁H₂₆N₂O₂ 338.449

Presumed precursor of Presecamine, P-624. Noncryst. Reactive, readily dimerises.

15,20-Dihydro: 15,20-Dihydrosecodine [27570-39-4]

Formed by the pyrolysis of Tetrahydropresecamine in P-624. Noncryst. [α]_D²⁰ -2.5 (MeOH). Dimerises back to Tetrahydropresecamine on standing.

16,17-Dihydro: 16,17-Dihydrosecodine [27825-43-0]

C₂₁H₂₈N₂O₂ 340.464

Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Gum.

16,17-Dihydro, 17-hydroxy: 16,17-Dihydrosecodin-17-ol [27825-45-2]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Rhazya orientalis* (Apocynaceae). Cryst. (CH₂Cl₂). Mp 131.5-132° Mp 150-151°. λ_{max} 222 (log ε 4.49); 274 (log ε 3.87); 284 (log ε 3.93); 292 (log ε 3.86) (EtOH).

15,16,17,20-Tetrahydro: Tetrahydrosecodine [27498-98-2]

C₂₁H₃₀N₂O₂ 342.48

Alkaloid from *Rhazya orientalis* and the leaves of *Rhazya stricta* (Apocynaceae). Gum. Mp 223-225° (216-218°) dec. (as picrolonate). [α]_D²⁵ -6.5 (CHCl₃). [α]_D²⁵ 0 (EtOH). Probably a mixt. of diastereoisomers.

15,16,17,20-Tetrahydro, 17-hydroxy: Tetrahydrosecodinol [28423-88-3]

C₂₁H₃₀N₂O₃ 358.48

Alkaloid from the roots of *Rhazya orientalis* (Apocynaceae). Noncryst.

Cordell, G.A. *et al.*, *Chem. Comm.*, 1970, 189-190; 191-192 (*isol, uv, ir, pmr, ms, struct, 16,17-dihydro, tetrahydro*)

Brown, R.T. *et al.*, *Chem. Comm.*, 1970, 190-191 (*Tetrahydrosecodine, synth, biosynth*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1970, 193-194 (*synth, struct*)


Marazano, C. *et al.*, *Chem. Comm.*, 1977, 742-743 (*synth*)

Kutney, J.P. *et al.*, *Can. J. Chem.*, 1979, **57**, 289-299 (*synth, uv, ir, pmr, ms*)

Raucher, S. *et al.*, *J.A.C.S.*, 1981, **103**, 2419-2421 (*synth, pmr*)

Kalaus, G. *et al.*, *Heterocycles*, 1985, **23**, 2783 (*synth, pmr, cmr, ms, Tetrahydrosecodine*)

1,2-Secodysidamide S-185

C₁₆H₂₅Cl₆NO₄ 508.094

Constit. of the Red Sea sponge *Lamellosidea herbacea*. Needles. Mp 173-174°. [α]_D²⁵ -41.8 (c, 1.1 in CH₂Cl₂).

3-Ketone: 1,2-Secodysidamide H


C₁₆H₂₃Cl₆NO₄ 506.078

Constit. of *Lamellosidea herbacea*. Needles. Mp 95-96°. [α]_D²⁵ -29.6 (c, 0.44 in CH₂Cl₂).

Sauleau, P. *et al.*, *Tetrahedron*, 2005, **61**, 955-963 (*isol, pmr, cmr*)

Secofascaplysin A S-186

[132911-53-6]

C₁₉H₁₄N₂O₃ 318.331

Alkaloid from the sponge *Fascaplysinopsis reticulata*. Pale yellow cryst.; originally isol. as a red oil. Mp 105-107°. λ_{max} 238 (ε 22500); 286; 296 (ε 4780); 334 (ε 3600); 350 (sh) (ε 3000) (MeOH) (Derep).

4'-Bromo: 3-Bromosecofascaplysin A [693790-80-6]

C₁₉H₁₃BrN₂O₃ 397.227

Alkaloid from *Fascaplysinopsis reticulata*. Yellow solid.

4'-Bromo, parent acid: 3-Bromosecofascaplysin B [693790-81-7]

C₁₈H₁₁BrN₂O₃ 383.201

Alkaloid from *Fascaplysinopsis reticulata*. Yellow solid.


Jimenez, C. *et al.*, *J.O.C.*, 1991, **56**, 3403-3410 (*isol, uv, ir, pmr, cmr, struct*)

Fürster, A. *et al.*, *Tetrahedron*, 1996, **52**, 7329-7344 (*synth, ir, pmr, cmr*)

Segraves, N.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 783-792 (*3-Bromosecofascaplysin*)

Secojesaconitine S-187

[119212-27-0]

C₃₃H₄₅NO₁₀ 615.719

Alkaloid from the roots of *Aconitum japonicum* (Ranunculaceae). Prisms + 1CHCl_3 ($\text{CHCl}_3/\text{Me}_2\text{CO}$). Mp 175-180°. $[\alpha]_{\text{D}}^{20} +12.5$ (c, 1.0 in MeOH).


15-Deoxy: Secoyunaconitine

$\text{C}_{33}\text{H}_{45}\text{NO}_9$ 599.72

Alkaloid from the roots of *Aconitum episcopale*. Needles ($\text{CHCl}_3/\text{Me}_2\text{CO}$). Mp 180-182°. $[\alpha]_{\text{D}}^{14} +56.1$ (c, 0.77 in CHCl_3).

Bando, H. et al., *Chem. Pharm. Bull.*, 1988, **36**, 1604 (*Secoyunaconitine, cryst struct*)

Li, Z.Y. et al., *Helv. Chim. Acta*, 2004, **87**, 2085-2087 (*Secoyunaconitine*)

Secojuliprosopinal S-188


$\text{C}_{36}\text{H}_{68}\text{N}_2\text{O}_3$ 576.945

Alkaloid from *Prosopis juliflora*. Gum. $[\alpha]_{\text{D}}^{28} +5$ (c, 1 in MeOH). λ_{max} 224 (ε 4700) (MeOH).

Nakano, H. et al., *Phytochemistry*, 2004, **65**, 587-591 (*isol, pmr, cmr*)

Secokaraconitine S-189

[475978-90-6]




$\text{C}_{30}\text{H}_{39}\text{NO}_9$ 557.639

Alkaloid from the tubers of *Aconitum karacolicum*. Cryst. + $\frac{1}{2}\text{H}_2\text{O}$ (Me_2CO). Mp 230-232°.

Sultankhodzhaev, M.N. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2002, **38**, 78-81; 2003, **39**, 512 (*isol, pmr, cmr, cryst struct*)

Secolucidine S-190


[107739-86-6]



$\text{C}_6\text{H}_{34}\text{N}_2\text{O}_7$ 606.674

Minor alkaloid from the bark of *Pseudoxandra sclerocarpa* (Annonaceae). $[\alpha]_{\text{D}}$ +82 (c, 0.5 in CHCl_3).

Cortes, D. et al., *J. Nat. Prod.*, 1986, **49**, 854 (*isol, uv, ir, pmr, ms, cd, struct*)

Secomartinelline S-191


$\text{C}_{33}\text{H}_{54}\text{N}_{10}\text{O}_2$ 622.855

Alkaloid from the root of *Martinella iquitosensis*. Bradykinin receptor antagonist. Muscarinic antagonist. Adrenergic and antihistaminic agents. $[\alpha]_{\text{D}}^{25} +9.4$. λ_{max} 314 (ε 12460) (H_2O).

U.S. Pat., 1994, 5 288 725; *CA*, **121**, 91779s (*isol, pmr, cmr*)

Secoobaberine S-192

Auroramine



$\text{C}_{38}\text{H}_{40}\text{N}_2\text{O}_8$ 652.743

Secoobaberine as isolated was the *S*-form and Auroramine the *R*-form.

(R)-form [115653-32-2]

Alkaloid from the leaves of *Gyrocarpus americanus* (Hernandiaceae).

O⁶-De-Me: Maroumine

[115610-42-9]

$\text{C}_{37}\text{H}_{38}\text{N}_2\text{O}_8$ 638.716

Alkaloid from the leaves of *Gyrocarpus americanus* (Hernandiaceae).

(S)-form [55085-45-5]

Minor alkaloid from *Pseudoxandra aff. lucida*. Amorph. $[\alpha]_{\text{D}} -5$ (c, 0.2 in CHCl_3).

O⁷-De-Me: Secohomoaromaline

[175097-24-2]

$\text{C}_{37}\text{H}_{38}\text{N}_2\text{O}_8$ 638.716

Obt. by the oxidn. of Homoaromaline in O-1. Alkaloid from leaves of *Anisocycla jollyana*. Needles. $[\alpha]_{\text{D}}^{20} -5.4$ (c, 5.3 in CHCl_3).

O^{12'}-De-Me: Baluchistanamine

[55085-44-4]

$\text{C}_{37}\text{H}_{38}\text{N}_2\text{O}_8$ 638.716

Obt. in low yield by the oxidn. of Oxyacanthine in O-1. Alkaloid from *Berberis baluchistanica* and *Berberis lycium*. Cryst. (cyclohexane/ C_6H_6).

1^{1'}-Carboxylic acid, O^{12'}-de-Me, Me ester: Tejedine

[221001-23-6]

$\text{C}_{38}\text{H}_{40}\text{N}_2\text{O}_9$ 668.742

Alkaloid from *Berberis vulgaris* ssp. *australis* (barberry). Amorph. powder. Mp 132-134°. $[\alpha]_{\text{D}}$ -40.6 (c, 0.06 in MeOH). λ_{max} 206 (log ε 4.83); 224 (log ε 4.79); 260 (log ε 4.36); 292 (log ε 3.99) (MeOH).

Shamma, M. et al., *J.A.C.S.*, 1974, **96**, 7809 (*isol, ir, uv, ms, pmr, cd, struct, Baluchistanamine*)

Leete, J.E. et al., *Heterocycles*, 1982, **19**, 2355 (*isol, Baluchistanamine*)

Cortes, D. et al., *J. Nat. Prod.*, 1985, **48**, 76 (*Secoobaberine*)

Dutè, P. et al., *Phytochemistry*, 1988, **27**, 655 (*Auroramine, Maroumine*)


Kanyinda, B. et al., *J. Nat. Prod.*, 1996, **59**, 498 (*Secohomoaromaline*)

Suau, R. et al., *Phytochemistry*, 1998, **49**, 2545-2549 (*Tejedine*)

Wang, Y.-C. et al., *Org. Lett.*, 2002, **4**, 2675-2678 (*Tejedine, synth*)

14,15-Seco-3-oxokopsinal S-193

[90373-27-6]



Absolute Configuration


$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from the stem bark of *Melodinus guillauminii* (Apocynaceae). $[\alpha]_{\text{D}}$ -41 (c, 0.4 in CHCl_3).

Zeches, M. et al., *Phytochemistry*, 1984, **23**, 171 (*isol, uv, ir, pmr, ms, struct*)

Secopenitrem B S-194

[139975-54-5]



$\text{C}_{37}\text{H}_{47}\text{NO}_5$ 585.782

Metab. of *Aspergillus sulphureus*. Shows antiinsect props. Orange-brown solid. Mp 185-189° dec. $[\alpha]_{\text{D}}$ -5.2 (CHCl_3). λ_{max} 220 (ε 31900); 286 (ε 11900) (MeOH) (Derep).

15-Hydroxy, 18,19-didehydro: Thomitrem E

[502759-47-9]

$\text{C}_{37}\text{H}_{45}\text{NO}_6$ 599.766

Prod. by *Penicillium crustosum* grown on rice. Named as Thomitrem E because it is structurally related to Penitrem E. Thomitrems B-D have not been isol. λ_{max} 228 ; 249 ; 284 (no solvent reported).

15-Hydroxy, 6-chloro, 18,19-didehydro: Thomitrem A

[502759-46-8]


$\text{C}_{37}\text{H}_{44}\text{ClNO}_6$ 634.21

Prod. by *Penicillium crustosum* grown on rice. λ_{max} 236 ; 254 ; 282 (no solvent reported).

Laakso, J.A. et al., *J.O.C.*, 1992, **57**, 2066-2071 (*isol, pmr, cmr*)

Rundberget, T. et al., *Phytochemistry*, 2002, **61**, 979-985 (*Thomitrems*)


Secoplicamine S-195
[232597-77-2]



$C_{26}H_{28}N_2O_6$ 464.517
Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. $[\alpha]_D^{25}$ -16.9 (c, 0.14 in MeOH). λ_{max} 221 (sh) (log ϵ 4.41); 243 (sh) (log ϵ 4.02); 287 (log ϵ 3.67) (MeOH).

Unver, N. et al., *Phytochemistry*, 1999, **50**, 1255-1261 (*isol, uv, ir, cd, pmr, cmr*)


Secopseudopteraxazole S-196
[242150-02-3]



$C_{21}H_{29}NO$ 311.466
Isol. from *Pseudopterogorgia elisabethae*. Active against *Mycobacterium tuberculosis*. Yellowish oil. $[\alpha]_D^{25}$ +28.2 (c, 0.85 in $CHCl_3$). Isolate was contaminated with traces of Elisabethin A.

Rodriguez, A.D. et al., *Org. Lett.*, 1999, **1**, 527-530

Secoquettamine S-197
4-[4-[2-(Dimethylamino)ethyl]-7-methoxy-2-benzofuranyl]phenol, 9CI. 4-(2-Dimethylaminoethyl)-2-(4-hydroxyphenyl)-7-methoxybenzofuran
[77765-52-7]



$C_{19}H_{21}NO_3$ 311.38
Alkaloid from *Berberis baluchistanica* (Berberidaceae). Needles (MeOH). Mp 171-172°.


Dihydro: **Dihydrosecoquettamine**
[77765-54-9]

$C_{19}H_{23}NO_3$ 313.396
Alkaloid from *Berberis baluchistanica* (Berberidaceae). Noncryst. Opt. inactive.

Zarga, M.H.A. et al., *Tet. Lett.*, 1981, **22**, 541 (*uv, pmr, ms, struct*)

Chattopadhyay, S. et al., *Heterocycles*, 1982, **19**, 697 (*synth*)
Biftu, T. et al., *J. Chem. Res., Synop.*, 1982, 270 (*synth*)

Secosarcocapnidine S-198
9-[2-(Dimethylamino)ethyl]-3,6-dimethoxydibenz[b,f]oxepin-4-ol, 9CI
[110883-33-5]



$C_{20}H_{23}NO_4$ 341.406
Trace alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Noncryst.

Hydrochloride:

Cryst. (EtOH/Et₂O). Mp 240-241°.

N-De-Me: **Norsecosarcocapnidine**. 3,6-Dimethoxy-9-[2-(methylamino)ethyl]-dibenz[b,f]oxepin-4-ol, 9CI
[110883-30-2]

$C_{19}H_{21}NO_4$ 327.379
Trace alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Noncryst.

Me ether: **Secosarcocapnine**. 4,6,7-Trimethoxy-N,N-dimethyldibenz[b,f]oxepin-1-ethanamine, 9CI
[110883-34-6]

$C_{21}H_{25}NO_4$ 355.433
Trace alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Noncryst.

Me ether hydrochloride:

Cryst. (EtOH/Et₂O). Mp 168-188°.

Me ether, N-De-Me: **Norsecosarcocapnine**. 4,6,7-Trimethoxy-N-methyldibenz[b,f]oxepin-1-ethanamine, 9CI
[110883-31-3]


$C_{20}H_{23}NO_4$ 341.406
Trace alkaloid from *Sarcocapnos crassifolia* and *Sarcocapnos enneaphylla* (Papaveraceae). Noncryst.

Castedo, L. et al., *Heterocycles*, 1987, **26**, 591 (*isol, uv, ir, pmr, ms, struct*)

Tojo, E. et al., *Heterocycles*, 1988, **27**, 2367 (*synth, Norsecosarcocapnidine, Norsecosarcocapnine*)

16,28-Secosolanida-5,22(28)-diene-3,16-diol, 9CI S-199

20-(3,4,5,6-Tetrahydro-5-methyl-2-pyridinyl)-pregn-5-ene-3,16-diol, 13CI. 22,26-Epiminocholesta-5,22(N)-diene-3,16-diol, 9CI



(3β,16α,20S,25α)-form

$C_{27}H_{43}NO_2$ 413.642

(3β,16α,20S,25α)-form
Etioline. *Veralosidine*. *Veralozidine*. 16-Hydroxyverazine
[29271-49-6]

Alkaloid from *Veratrum album* ssp. *lobelianum*, *Veratrum grandiflorum* and *Solanum havanense* (Liliaceae, Solanaceae). Hepatoprotective agent. Mp 153-155°. $[\alpha]_D^{26}$ -92.2 (c, 0.47 in EtOH). Identity of Etioline and Veralosidine apparently not conclusively establ.

3-O-β-D-Glucopyranoside: **Deacetylveralosine**. *Etioline* 3-glucoside
[54557-67-4]

$C_{33}H_{53}NO_7$ 575.784
Alkaloid from epigeal parts of *Veratrum lobelianum* and from roots of *Solanum spirale*. Cryst. (MeOH). Mp 238-240° (225-227° dec.). $[\alpha]_D$ -30.3 (c, 0.5 in $CHCl_3$). $[\alpha]_D^{26}$ -78.8 (c, 0.72 in Py).

3-O-[β-D-Glucopyranosyl-(1→4)-β-D-glucopyranoside]: **Etiolinine**
[107783-36-8]

$C_{39}H_{63}NO_{12}$ 737.926
Alkaloid from *Solanum havanense* (Solanaceae).

16-Ac: **Veralosinine**. 16-O-Acetyletioline. *Veralozinine*
[36506-65-7]

$C_{29}H_{45}NO_3$ 455.679
Alkaloid from *Veratrum lobelianum* (Liliaceae). Mp 161-163°. $[\alpha]_D^{25}$ -186.2 (c, 0.92 in $CHCl_3$).

16-Ac, 3-O-β-D-glucopyranoside: **Veralosine**. *Veralozine*. *Havanine*†
[30511-97-8]

$C_{35}H_{55}NO_8$ 617.821
Alkaloid from *Veratrum album* ssp. *lobelianum* and from leaves of *Solanum havanense* (Liliaceae, Solanaceae). Mp 213-215° (186-187°). $[\alpha]_D$ -147.7 (MeOH). $[\alpha]_D^{25}$ -110.8 (c, 1 in MeOH). Struct. of Veralosine not certain and identity of Veralosine with Havanine not establ.

(3β,16α,20S,25β)-form
25-Isoetioline. 25-Epietioline
[129938-55-2]

Alkaloid from the leaves and stems of *Solanum canense* and *Solanum fraxinifolium* (Solanaceae). Mp 141-143°. $[\alpha]_D^{20}$ +73.6 (c, 0.4 in $CHCl_3$).

(3β,16α,20R,25β)-form
20,25-Bisoietioline
[175777-78-3]

Alkaloid from leaves and stems of *Solanum canense* and *Solanum fraxinifolium*. Cryst. (MeOH). Mp 158-161°. $[\alpha]_D^{20}$ -11.8 (c, 0.69 in $CHCl_3$).

Khashimov, A.M. et al., *Khim. Prir. Soedin.*, 1970, **6**, 339; 1971, **7**, 779; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 338; 1971, **7**, 751 (*isol, pmr*)

Kaneko, K. et al., *Tet. Lett.*, 1971, 4251 (*isol, spectra*)


Abaidullaev, K.A. et al., *Khim. Prir. Soedin.*, 1974, **10**, 678; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 703 (*Deacetylveralosine*)

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1976, **12**, 623; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 557 (*cd*)

- Ripperger, H. *et al.*, *Pharmazie*, 1977, **32**, 537 (isol)
- Basterrechea, M. *et al.*, *Phytochemistry*, 1984, **23**, 2057-2058 (*Havanine*)
- Basterrechea, M.J. *et al.*, *Rev. Cubana Quim.*, 1986, **2**, 71; *CA*, **106**, 153103t (*Etiolinine*)
- Ripperger, H. *et al.*, *Phytochemistry*, 1990, **29**, 3375-3376; 1996, **41**, 1629-1631 (25-*Isoetioline*, 20,25-*Bisioetioline*)
- Le Thi Quyen, *et al.*, *Annalen*, 1991, 143 (synth)
- Ripperger, H. *et al.*, *Phytochemistry*, 1996, **43**, 705-707 (*Etioline 3-O-glucoside*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, MRV750

16,28-Secosolanid-5-ene-3,16-diol S-200

20-(5-Methyl-2-piperidinyl)-pregn-5-ene-3,16-diol, 13CI. 22,26-Epiminocholest-5-ene-3,16-diol



(3β,16α,22R,25R)-form

C₂₇H₄₅NO₂ 415.658

(3β,16α,22R,25R)-form Capsimine

[123123-41-1]

Constit. of *Solanum capsicastrum*. Antihepatotoxic agent. Prisms (MeOH) + 2H₂O. Mp 264-267°. [α]_D²⁵ -104 (c, 0.5 in MeOH).

3-O-β-D-Glucopyranoside: *Capsimine 3-glucoside*

[145681-73-8]

C₃₃H₅₅NO₇ 577.8

Alkaloid from root bark of *Solanum capsicastrum* (Solanaceae). Antihepatotoxic agent. Powder (MeOH). Mp 214-216°. [α]_D²⁰ -68.8 (c, 0.05 in cyclohexane/EtOAc/MeOH).

(3β,16α,22R,25S)-form

Teinimine

[65027-01-2]

Isol. from *Veratrum grandiflorum* terrestrial parts (Liliaceae). Plates (Me₂CO). Mp 204-209°. [α]_D²⁰ -35.8 (c, 1.19 in CHCl₃).

► QG1360000

3-O-β-D-Glucopyranoside: *Isocapsicarine. Tienimine 3-glucoside*

[123164-25-0]

C₃₃H₅₅NO₇ 577.8

Isol. from *Solanum capsicastrum*. Pale yellow powder (MeOH/CHCl₃). [α]_D²⁵ +202 (c, 0.25 in Py).

(3β,16α,22S,25R)-form

Schlechtendamine. 22-Epicapsimine

[36456-16-3]

Alkaloid from *Solanum schlechtendalia-num*. Cryst. Mp 252-253° (synthetic). [α]_D²³ -33 (c, 0.4 in CHCl₃) (synthetic).

(3β,16α,22S,25S)-form

Isoleinimine

[65027-00-1]

Isol. from *Veratrum grandiflorum* terrestrial parts (Liliaceae). Flakes (Me₂CO). Mp 217-220°.

3-O-β-D-Galactopyranoside: *Capsicarine. Isoleinimine galactoside*

[107585-56-8]

C₃₃H₅₅NO₇ 577.8

Alkaloid from root bark of *Solanum capsicastrum* (Solanaceae). Needles (Me₂CO). Mp 220-221°. [α]_D²¹ -25.5 (c, 0.1 in CHCl₃).

3-O-[α-L-Rhamnopyranosyl-(1→4)-[α-L-rhamnopyranosyl-(1→4)-β-D-glucopyranosyl-(1→2)]]-β-D-glucopyranoside]: *Solateinimine*

[211683-81-7]

C₅₁H₈₅NO₂₀ 1032.227

Alkaloid from the root bark of *Solanum pseudocapsicastrum*.

16-Ac: *Muldamine. Alkaloid Q*

[36069-45-1]

C₂₉H₄₇NO₃ 457.695

Alkaloid from *Veratrum californicum* (Liliaceae). Mp 210-211°. [α]_D²⁵ -95 (c, 1 in EtOH/CHCl₃ 3:1). c-Nor-D-homosteroid struct. originally assigned.

► Exp. reprod. and teratogenic effects. QG1320000

(3β,16β,22S,25R)-form

Solaverbascine. Dihydrosolasodine A

[6785-55-3]

Alkaloid from leaves of *Solanum verbascifolium* (Solanaceae). Abortifacient responsible for the folk medicine use of *S. verbascifolium*. Cryst. (MeOH aq.). Mp 263-265°. [α]_D -67.9 (c, 0.3 in CHCl₃).

► Poss. teratogen.

Tri-Ac: Mp 171°. [α]_D -16.8 (CHCl₃).

(3β,16β,22S,25S)-form

N-Me: 28-Methyl-16,28-secosolanid-5-ene-3,16-diol. *Hapepunine*

[68422-01-5]

C₂₈H₄₇NO₂ 429.685

Alkaloid from the aerial parts of *Fritillaria camtschatcensis* (Liliaceae). Needles (EtOH). Mp 201-202° (196.5-198.5°). [α]_D -72.6.

N-Me, 3-O-[β-D-glucopyranosyl-(1→4)-β-D-glucopyranoside]: *Hapepunine 3-β-cellobioside*

[169786-66-7]

C₄₀H₆₇NO₁₂ 753.969

Alkaloid from bulbs of *Fritillaria maximowiczii* (Liliaceae). Amorph. [α]_D²⁷ -29.1 (c, 0.43 in MeOH).

N-Me, 3-O-[α-L-rhamnopyranosyl-(1→2)-β-D-glucopyranoside]: *Hapepunine 3-neohesperidoside*

C₄₀H₆₇NO₁₁ 737.969

Alkaloid from aerial parts of *Fritillaria thunbergii* (Liliaceae). Needles + 1H₂O (MeOH aq.). Mp 269-274° dec. [α]_D -67.2 (Py).

N-Me, di-Ac: Mp 207-212°.

Kaneko, K. *et al.*, *Phytochemistry*, 1977, **16**, 1620-1622; 1981, **20**, 157-160 (*Teinimine, Isoleinimine, Hapepunine*)

Adam, G. *et al.*, *Phytochemistry*, 1980, **19**, 1002-1003 (*Solaverbascine*)

Katajima, J. *et al.*, *Phytochemistry*, 1982, **21**, 187-192 (*Hapepunine rhamnosylglucoside*)

Gaffield, W. *et al.*, *Phytochemistry*, 1982, **21**, 2397-2400 (*Muldamine*)

Lin, C.N. *et al.*, *Phytochemistry*, 1987, **26**, 305-307 (*Capsicarine*)

Lin, C.N. *et al.*, *Planta Med.*, 1989, **55**, 48-50 (*Capsimine*)

Gan, K.H. *et al.*, *J. Nat. Prod.*, 1993, **56**, 15-21 (*Capsimine 3-glucoside*)

Qian, Z.Z. *et al.*, *Phytochemistry*, 1995, **40**, 979-981 (*Hapepunine cellobioside*)


Gan, K.-H. *et al.*, *Chin. Pharm. J. (Taipei)*, 1997, **49**, 315-320 (*Solateinimine*)

Ferrer, A. *et al.*, *Rev. Cubana Quim.*, 1998, **10**, 3 (*Schlechtendamine*)

Weissenberg, M. *et al.*, *Phytochemistry*, 2001, **58**, 501-508 (*Solaverbascine glycosides*)

16,28-Secosolanid-22(28)-ene-3,6-diol S-201

22,26-Imincholest-22(N)-ene-3,6-diol



(3β,5α,6β,25R)-form

C₂₇H₄₅NO₂ 415.658

(3β,5α,6β,25R)-form

O³-β-D-Glucopyranoside: *Cordatine A*

[107585-57-9]

C₃₃H₅₅NO₇ 577.8

Alkaloid from the petals of *Lilium cordatum* (Liliaceae). Pale-red powder. [α]_D²⁹ -3.5 (c, 0.85 in MeOH).

(3β,5α,6β,25S)-form

O³-β-D-Glucopyranoside: *Cordatine B*

[107566-79-1]


C₃₃H₅₅NO₇ 577.8

Alkaloid from the petals of *Lilium cordatum* (Liliaceae). Pale-red needles. Mp 187-190°. [α]_D²⁹ -11.2 (c, 1.07 in MeOH).

Nakano, K. *et al.*, *Phytochemistry*, 1987, **26**, 301 (isol, ir, pmr, cmr, ms, struct)

Secuamamine A

S-202



Absolute Configuration

C₁₃H₁₅NO₃ 233.266

Alkaloid from the leaves and twigs of *Securinega suffruticosa* var. *amamiensis*. Oil. [α]_D²³ -479 (c, 0.15 in CHCl₃). λ_{max} 254 (log ε 4.06) (MeOH).


Ohsaki, A. *et al.*, *Tet. Lett.*, 2003, **44**, 3097-3099 (isol, pmr, cmr)

Magnus, P. *et al.*, *Org. Lett.*, 2006, **8**, 3569-3571 (biosynth)

Liu, P. *et al.*, *J.A.C.S.*, 2008, **130**, 7562-7563 (synth)

Securamine A

[173220-55-8]

C₂₀H₂₀BrClN₄O 447.761

Exists in equilib. with Securine A.
Alkaloid from the marine bryozoan
Securiflustra securifrons. Cryst. Mp 200°
dec. $[\alpha]_D^{20}$ -87.5 (c, 0.064 in CHCl₃).

16-Bromo: Securamine B

[173220-56-9]


C₂₀H₁₉Br₂ClN₄O 526.657

From *Securiflustra securifrons*. Cryst.
 $[\alpha]_D^{20}$ -316.7 (c, 0.030 in CHCl₃). Exists
in equilib. with Securine B.

Rahbaek, L. *et al.*, *J.O.C.*, 1996, **61**, 887-889
(*isol, uv, ir, cd, pmr, cmr, struct*)

Securamine C

[173220-57-0]

Relative
ConfigurationC₂₀H₁₈BrClN₄O₂ 461.744

Alkaloid from the marine bryozoan
Securiflustra securifrons. Amorph.
yellow solid. $[\alpha]_D^{20}$ -433.5 (c, 0.033 in
CHCl₃).

Debromo: Securamine D

[173220-58-1]

C₂₀H₁₉ClN₄O₂ 382.848

Isol. from *Securiflustra securifrons*.
Amorph. green solid. $[\alpha]_D^{20}$ -320 (c,
0.069 in CHCl₃).

18-Bromo: Securamine E

[185616-44-8]

C₂₀H₁₇Br₂ClN₄O₂ 540.641

Isol. from *Securiflustra securifrons*.
Amorph. yellow solid. $[\alpha]_D^{20}$ -115 (c,
0.0078 in CHCl₃). λ_{max} 303 (log ϵ
3.82); 327 (log ϵ 3.83) (CHCl₃).

10-Epimer: Securamine F

[185616-45-9]

C₂₀H₁₈BrClN₄O₂ 461.745

Isol. from *Securiflustra securifrons*.
Amorph. orange solid. $[\alpha]_D^{20}$ -200 (c,
0.0045 in CHCl₃). λ_{max} 299 (log ϵ
3.86); 328 (log ϵ 3.88) (CHCl₃).

2,3,4,5-Tetrahydro, 3,4-didehydro: Securamine G

[185616-47-1]

C₂₀H₂₀BrClN₄O₂ 463.76

Isol. from *Securiflustra securifrons*.
Amorph. brown solid. $[\alpha]_D^{20}$ -15.6 (c,

S-203


0.032 in CHCl₃). λ_{max} 298 (log ϵ 2.81)
(CHCl₃).

Rahbaek, L. *et al.*, *J.O.C.*, 1996, **61**, 887-889
(*isol, uv, ir, cd, pmr, cmr, struct*)

Rahbaek, L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 175-
177 (*Securamines E-G*)

Securinegine†

[72741-90-3]

C₁₃H₁₅NO₂ 217.267


Alkaloid from the leaves of *Securinega
suffruticosa*. Yellow cryst. (EtOH).

Shabana, M.M. *et al.*, *CA*, 1980, **92**, 72692z
(*isol, struct*)

Securinine, INN

S-206

**9,10,11,11a-Tetrahydro-8H-6,11b-metha-
nofuro[2,3-c]pyrido[1,2-a]pyridoazepin-
2(6H)-one, 9Cl. Virosecurinine**



(-)-form

C₁₃H₁₅NO₂ 217.267

Virosecurinine was originally the (+)-
enantiomer. Log P 0.21 (uncertain value)
(calc).

(-)-form [5610-40-2]

Alkaloid from the leaves, roots and stems
of *Securinega suffruticosa* and *Phyl-
lanthus discoides*, also isol. from *Secur-
inega durissima*, *Securinega fluggeoides*
and the bark of *Securidaca longepedun-
culata* (Euphorbiaceae, Fabaceae).
Stereospecific GABA_A receptor antago-
nist. CNS stimulant with strychnine-like
activity but lower toxicity, causes res-
piratory stimulation, increased muscle
tone and hypotension. Yellow cryst.
(EtOH). Mp 142-143°. $[\alpha]_D^{20}$ -1042 (c, 1.0
in EtOH). pK_a 7.17.

▶ LD₅₀ (mus, orl) 270 mg/kg. VS4115000**Hydrobromide:**

Pale yellow needles + 2H₂O (Et₂O).
Mp 250° dec. (dehydrating range 110-
120°).

Picolonate: Mp 194° (208°).**14,15-Dihydro: Dihydrosecurinine**

[1878-04-2]

Alkaloid from the leaves of *Securinega
suffruticosa* (Euphorbiaceae). Flakes
(petrol). Mp 58-60°. $[\alpha]_D^{20}$ 0 (c, 1 in
EtOH). pK_a 8.35.

14,15-Dihydro, hydrochloride:

Needles (MeOH/Et₂O). Mp 256-258°.
 $[\alpha]_D^{22}$ -71.5 (c, 1 in EtOH).

14,15-Dihydro, picrate: Mp 244-246°.

14,15-Dihydro, 15 α -methoxy: Securamine C

[1004529-77-4]

C₁₄H₁₉NO₃ 249.309

Alkaloid from the wood of *Securinega
suffruticosa* var. *amamiensis*. Amorph.
solid. $[\alpha]_D^{25}$ +82.1 (c, 0.15 in CHCl₃).
Only relative config. determined. λ_{max}
232 (log ϵ 3.41); 283 (log ϵ 2.53)
(MeOH).

2-Epimer: see Allosecurinine, A-630

(+)-form [6704-68-3]

Alkaloid from the leaves of *Securidaca
virosa* (Euphorbiaceae). Yellow cryst.
(EtOH). Mp 141-142°. $[\alpha]_D^{20}$ +1035 (c,
0.43 in EtOH). $[\alpha]_D^{20}$ +1148 (c, 0.4 in
CHCl₃). It is remarkable that while
Securinine and Allosecurinine (2-*epi*-
Securinine) co-occur in *Securinega suffru-
ticosa*, their optical antipodes,
Virosecurinine and Viroallosecurinine,
co-occur in *S. virosa*.

▶ ZA5500000**Hydrochloride:**

Cryst. (Me₂CO). Mp 252-253° (sealed
tube). $[\alpha]_D^{21}$ +344 (c, 0.52 in EtOH).

Picrate:

Cryst. (Me₂CO). Mp 218°.

(±)-form [5588-52-3]

Synthetic. Yellow needles (hexane). Mp
109-111°.

Parello, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 898
(*uv, ir, pmr*)

Nakano, T. *et al.*, *J.O.C.*, 1963, **28**, 2619; 1964,
29, 3441 (*uv, ord, cd, config, Virosecurinine*)
Mukherjee, R. *et al.*, *Naturwissenschaften*,
1963, **50**, 155 (*isol, uv, ir, ms*)

Nakano, T. *et al.*, *Tetrahedron*, 1963, **19**, 609
(*isol, uv, ir, struct, Virosecurinine*)

Saito, S. *et al.*, *Tetrahedron*, 1963, **19**, 2085 (*uv,
ir, struct, synth*)

Horii, Z. *et al.*, *Tetrahedron*, 1963, **19**, 2085;
2101; 1967, **23**, 1165 (*uv, ir, pmr, struct, abs
config, synth*)

Saito, S. *et al.*, *Yakugaku Zasshi*, 1963, **83**, 800;
CA, **59**, 15535h (*Dihydrosecurinine, isol*)

Bevan, C.W.L. *et al.*, *Chem. Ind. (London)*,
1964, 838 (*occur*)

Imado, S. *et al.*, *Chem. Ind. (London)*, 1964,
1691 (*cryst struct*)

Chatterjee, A. *et al.*, *J. Indian Chem. Soc.*,
1964, **41**, 163 (*ir*)

Clauder, O. *et al.*, *Acta Pharm. Hung.*, 1968,
38, 126; *CA*, **69**, 54266j (*isol*)

Audier, H.E. *et al.*, *Bull. Soc. Chim. Fr.*, 1968,
1552 (*pmr, ms*)

Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**,
1967 (*synth, ir*)

Snieckus, V. *et al.*, *Alkaloids (London)*, 1971,
1, 455 (*rev*)

Snieckus, V. *et al.*, *Alkaloids (Academic
Press)*, 1973, **14**, 425 (*rev*)

Parry, R.J. *et al.*, *Tet. Lett.*, 1974, 307; *Chem.
Comm.*, 1975, 144 (*biosynth*)

Sankawa, U. *et al.*, *Tet. Lett.*, 1974, 1867
(*biosynth*)

Golebiewski, W.M. *et al.*, *Chem. Comm.*, 1976,
217 (*biosynth*)

Sankawa, U. *et al.*, *Phytochemistry*, 1977, **16**,
561 (*biosynth*)

Parry, R.J. *et al.*, *Bioorg. Chem.*, 1978, **7**, 277
(*biosynth*)

Beutler, J.A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 677
(*cmr*)


Niu, C.S. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 347;
CA, **110**, 154639s (*synth*)

- Luger, P. *et al.*, *Acta Cryst. C*, 1995, **51**, 127 (cryst struct)
 Liras, S. *et al.*, *Org. Lett.*, 2001, **3**, 703-706 (synth)
 Honda, T. *et al.*, *Heterocycles*, 2002, **59**, 169-187 (synth)
 Honda, T. *et al.*, *Org. Lett.*, 2004, **6**, 87-89 (synth)
 Alibés, R. *et al.*, *Org. Lett.*, 2004, **6**, 1813-1816 (synth)
 Ohsaki, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 2003-2005 (Secuamamine C)
 Yuan, W. *et al.*, *Nat. Prod. Res.*, 2007, **21**, 234-242 (isol, pmr, cmr)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SBN350

Securinol A

S-207

[5008-48-0]



Absolute Configuration

C₁₃H₁₇NO₃ 235.282

Struct. revised in 1991. Alkaloid from *Securinea suffruticosa* and *Margaritaria indica*. Cryst. (C₆H₆). Mp 135-136°. [α]_D¹⁷ +58.2 (c, 0.14 in CHCl₃).

Picrate:

Cryst. + 1H₂O (EtOH). Mp 186-187° dec.

15-Epimer: Securinol B. Virosine A

[30155-10-3 (Securinol B)]

C₁₃H₁₇NO₃ 235.282

Alkaloid from *Flueggea virosa* and *Securinea suffruticosa*. Needles (petrol). Mp 159-160°. [α]_D +120 (c, 0.15 in EtOH) (Securinol B). [α]_D +92 (c, 0.1 in MeOH) (Virosine B).

Probable struct. of Securinol B based on revised struct. for Securinol A.

Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 1307-1311; 1970, **18**, 2009-2012 (isol, uv, ord, ir, pmr, ms, Securinol B)

Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1991, 1863-1869 (isol, pmr, cmr, cryst struct, abs config)


Wang, G.-C. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 1124-1129 (Virosine A, B)

Securinol C

S-208

14,15-Dihydro-14-hydroxysecurinane-11-one, 9CI

[30155-11-4]



Tentative structure

C₁₃H₁₇NO₃ 235.282

Struct. doubtful. Alkaloid from the leaves of *Securinea suffruticosa* (Euphorbiaceae). Needles (C₆H₆/petrol). Mp 114-115°. [α]_D -81.9 (c, 0.58 in EtOH).

Picrate: Mp 194° dec.


Horii, Z. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 1307; 1970, **18**, 2009 (isol, ir, uv, pmr, ms,

struct)
 Arbain, D. *et al.*, *J.C.S. Perkin 1*, 1991, 1863

Sedacryptine

S-209

Octahydro-2-hydroxy-2,4-dimethyl-α-phenylfuro[3,2-b]pyridine-5-ethanol, 9CI



(-)-form

C₁₇H₂₅NO₃ 291.389**(-)-form** [77784-07-7]

Minor alkaloid from *Sedum acre* (Crassulaceae). Cryst. (cyclohexane). Mp 125-127°.

Hydrochloride: [77844-22-5]

Mp 141-143°. [α]_D²² -49 (c, 1.6 in MeOH).

(±)-form

Mp 99-101°.

Kooy, J.H. *et al.*, *Planta Med.*, 1976, **30**, 295 (isol)

Hootelé, C. *et al.*, *Tet. Lett.*, 1980, **21**, 5061 (isol, cryst struct, ms)

Colau, B. *et al.*, *Can. J. Chem.*, 1983, **61**, 470 (abs config)

Natsume, M. *et al.*, *Heterocycles*, 1983, **20**, 601 (synth)


Akiyama, E. *et al.*, *Synlett*, 1996, 100 (synth, abs config)

Wee, A.G.H. *et al.*, *Org. Lett.*, 2008, **10**, 3869-3872 (synth)

Sedamine

S-210

1-Methyl-α-phenyl-2-piperidineethanol, 9CI. 2-(2-Hydroxy-2-phenylethyl)-1-methylpiperidine. 2-(1-Methyl-2-piperidinyl)-1-phenylethanol. 8-Phenyllobelol



(2R,2'R)-form

C₁₄H₂₁NO 219.326**(2R,2'R)-form** [67008-23-5]

Synthetic. Cryst. (pentane). Mp 61-62°. [α]_D¹⁸ +91.5 (c, 3.29 in EtOH). [α]_D²² +65 (c, 0.1 in EtOH).

4R-Hydroxy: 4-Hydroxy-1-methyl-α-phenylpiperidineethanol. 4-Hydroxy-2-(2-hydroxy-2-phenylethyl)piperidine. 4-Hydroxysedamine

[61535-08-8]

[103823-30-9]

C₁₄H₂₁NO₂ 235.325

Alkaloid from *Sedum acre* (Crassulaceae). Amorph. Mp 130-131° (as picrate). [α]_D²² +58 (c, 1.6 in MeOH). A hydroxy deriv. of the enantiomer of natural Sedamine present in *S. acre*.

(2R,2'S)-form**(+)-Allosedamine**

Synthetic. Cryst. (pentane). Mp 79-80°. [α]_D²⁰ +18.6 (c, 2.42 in EtOH).

Picrate: Mp 129-130°.**4R-Hydroxy: 4-Hydroxyallosedamine**

[103805-62-5]

C₁₄H₂₁NO₂ 235.325

Alkaloid from *Sedum acre* (Crassulaceae). Mp 91-92°. [α]_D²² +20.6 (c, 1.3 in MeOH). A hydroxy deriv. of the enantiomer of natural allosedamine present in *S. acre*.

(2S,2'R)-form**Allosedamine. 8-Phenyllobelol I**

[497-89-2]

Alkaloid from *Lobelia inflata* and *Sedum acre* (Campanulaceae). Mp 81-82°. [α]_D^{19.5} -18.6 (EtOH). [α]_D²² -32.2 (MeOH).

Hydrochloride: Mp 118-119°.**N-De-Me: Norallosedamine. 8-Phenyl-norlobelol I**C₁₃H₁₉NO 205.299

Alkaloid from *Lobelia inflata* (Campanulaceae). Mp 102-103°. [α]_D²³ +49.3 (EtOH). [α]_D +37.2 (MeOH).

N-De-Me, hydrochloride: Mp 133-134°.**3R-Hydroxy: 3-Hydroxyallosedamine**C₁₄H₂₁NO₂ 235.325

Minor alkaloid from *Sedum acre* (Crassulaceae). Amorph. [α]_D²² -17 (c, 1.3 in MeOH). Racemate synthesised.

3R-Hydroxy, perchlorate: Mp 162-163°. [α]_D²² -40 (c, 2.4 in MeOH).

3R-Hydroxy, N-de-Me: 3-HydroxynorallosedamineC₁₃H₁₉NO₂ 221.299

Minor alkaloid from *Sedum acre* (Crassulaceae). Mp 155-156°. [α]_D²² -66 (c, 1.6 in MeOH). Racemate synthesised and resolved.

(2S,2'S)-form [497-88-1]

Alkaloid from *Sedum acre*, *Sedum lydium* and other *Sedum* spp. (Crassulaceae). Mp 75-76° (89°). [α]_D -88.4 (c, 4.65 in EtOH).

Hydrochloride: Mp 210-211°.**Picolonate:** Mp 159-160°.**N-De-Me: Norsedamine**

Semisynthetic. Mp 108-109°. [α]_D²⁰ -32.5 (MeOH).

2'-Ketone: 2-(1-Methyl-2-piperidinyl)-1-phenylethanol, 9CI. Sedaminone

[103771-49-9]

C₁₄H₁₉NO 217.31

Alkaloid from spp. in the Crassulaceae.

N-De-Me; hydrochloride: Mp 169-171°.**5S-Hydroxy: 5-Hydroxysedamine**C₁₄H₂₁NO₂ 235.325

Minor alkaloid from *Sedum acre* (Crassulaceae). Noncryst. [α]_D²⁰ -50 (c, 0.3 in MeOH).

5S-Hydroxy, di-Ac: [α]_D²⁰ -66 (c, 0.6 in MeOH).

(2RS,2'RS)-form**(±)-Sedamine**

[1630-44-0]

Alkaloid from *Sedum acre*, *Sedum aizoon*, *Sedum purpureum*, *Sedum hybridum* and *Sedum ewersii* (Crassulaceae). Mp 88°.

▶ TM9314000

Hydrochloride: Mp 185°.

2'-Ketone: [121961-27-1]

Oil. Bp_{0.15} 92-93°.

N-De-Me: Mp 98-99°.

(2RS,2'SR)-form

(±)-Allosedamine

Mp 67-68°.

N-De-Me: Mp 112-113°.

[67008-24-6, 67008-23-5]

Wieland, C.H. *et al.*, *Annalen*, 1939, **540**, 103 (isol, deriv)

Marion, L. *et al.*, *Can. J. Res., Sect. B*, 1945, **23**, 165 (isol)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1956, **75**, 82; 1957, **76**, 415; 1959, **78**, 43 (synth)

Schöpf, C. *et al.*, *Annalen*, 1959, **626**, 134; **628**, 101 (synth, abs config, derivs, bibl)

Spiteller-Friedmann, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 104 (ms)

Gupta, R.N. *et al.*, *Phytochemistry*, 1970, **9**, 2329 (biosynth)

Leistner, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4715 (biosynth)

Wakabayashi, T. *et al.*, *Chem. Lett.*, 1977, 223 (synth)

Krasnov, E.A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 585; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 492 (isol)

Tufariello, J. *et al.*, *Tet. Lett.*, 1978, 4647 (synth, pmr)

Gerdes, H.J. *et al.*, *Phytochemistry*, 1979, **18**, 771 (biosynth)

Irie, K. *et al.*, *Chem. Comm.*, 1985, 633 (synth)

Halin, F. *et al.*, *Tetrahedron*, 1985, **41**, 2891 (4-Hydroxysedamine, 4-Hydroxyallosedamine)

Hootele, C. *et al.*, *Tetrahedron*, 1985, **41**, 5563 (pmr, cmr, conformn, occur)

Ibebeke-Bomangwa, W. *et al.*, *Tetrahedron*, 1987, **43**, 935 (5-Hydroxysedamine, 3-Hydroxynorallosedamine)

Liguori, A. *et al.*, *Chem. Ber.*, 1989, **122**, 2019 (synth, ir, pmr, ms)

Takahata, H. *et al.*, *J.C.S. Perkin 1*, 1989, 1211 (synth, ir, pmr, bibl)

Pyne, S.G. *et al.*, *J.O.C.*, 1990, **55**, 1086 (synth)

Ozawa, N. *et al.*, *Heterocycles*, 1991, **32**, 889 (Sedamine, Allosedamine, synth)

Pilli, R.A. *et al.*, *Synth. Commun.*, 1991, **21**, 2213 (Sedamine, Allosedamine, synth)

Stevens, J.F. *et al.*, *Phytochemistry*, 1992, **31**, 3917 (isol, Sedamone)

Comins, D.L. *et al.*, *J.O.C.*, 1993, **58**, 5035 (synth)

Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 554 (Allosedamine, synth)

Herdeis, C. *et al.*, *Annalen*, 1995, 1295 (5-Hydroxysedamine)

Ghiaci, M. *et al.*, *Org. Prep. Proced. Int.*, 1996, **28**, 474 (synth)

Cossy, J. *et al.*, *J.O.C.*, 2002, **67**, 1982-1992 (synth)

Felplin, F.-X. *et al.*, *J.O.C.*, 2002, **67**, 9192-9199 (synth)

Bates, R.W. *et al.*, *Tetrahedron*, 2002, **58**, 5957-5978 (synth)

Angoli, M. *et al.*, *J.O.C.*, 2003, **68**, 9525-9527 (synth)

Raghavan, S. *et al.*, *Tetrahedron*, 2004, **60**, 5059-5067 (Allosedamine, synth)

Kang, B. *et al.*, *Tetrahedron*, 2004, **60**, 7353-7359 (Allosedamine, synth)

Yadav, J.S. *et al.*, *Synthesis*, 2006, 4005-4012 (synth)


Bates, R.W. *et al.*, *Synthesis*, 2008, 1033-1038 (synth, pmr, cmr)

Liu, G. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 1297-1303 (5-Hydroxysedamine, synth)

Sederine

S-211

Methyl 6-(2-hydroxy-2-phenylethyl)-1-methyl-2-piperidinecarboxylate, 9CI [70786-43-5]



C₁₇H₂₅NO₃ 291.389


Minor alkaloid from *Sedum acre* (Crassulaceae). Mp 199-201° (as hydrochloride). [α]_D²⁰ -82 (c, 1.4 in MeOH).

Hootele, C. *et al.*, *Bull. Soc. Chim. Belg.*, 1979, **88**, 111; 1983, **92**, 21 (isol, pmr, ms, cryst struct, abs config)

Sediendione

S-212

1-(6-Benzoyl-1,6-dihydro-2-pyridinyl)-2-propanone, 9CI. 2-Benzoyl-1,2-dihydro-6-acetylpyridine [133056-36-7]



C₁₅H₁₅NO₂ 241.289

(ξ)-form

Alkaloid from *Sedum acre* (Crassulaceae).


Maksimovic, M. *et al.*, *Rapid Commun. Mass Spectrom.*, 1990, **4**, 503; *CA*, **114**, 160721m (Sediendione, ms)

Maksimovic, M. *et al.*, *CA*, 2005, **143**, 474940y (Sediendione)

Sediene

S-213

1-[1,2,5,6-Tetrahydro-1-methyl-6-(2-oxopropyl)-2-pyridinyl]-3-buten-2-one, 9CI [133056-35-6]



C₁₃H₁₉NO₂ 221.299

Alkaloid from *Sedum acre* (Crassulaceae). Shows antimicrobial activity. Pale yellow powder. Mp 94°. λ_{max} 210; 325 (Me₂CO) (Berdy).


Nikolin, B. *et al.*, *Acta Pharm. Jugosl.*, 1990, **40**, 555 (isol)

Maksimovic, M. *et al.*, *Rapid Commun. Mass Spectrom.*, 1990, **4**, 503; *CA*, **114**, 160721m (isol, ms)

Sedinine

S-214

1,2,3,6-Tetrahydro-α²,1-dimethyl-α²-phenyl-2,6-pyridinediethanol, 9CI. 8-Methyl-10-phenyl-3,4-dehydrolobelidol [492-49-9]



C₁₇H₂₅NO₂ 275.39

The double bond was originally considered to be in the alternative 4,5-position. Alkaloid from *Sedum acre* (Crassulaceae). Prisms (cyclohexane). Mp 121°. [α]_D²⁰ -105 (c, 0.46 in MeOH).

Hydrochloride: [86161-46-8] Mp 172°. [α]_D²⁰ -145 (c, 1.04 in MeOH).

8-Ketone: **Sedacrine**. 1-[1,2,5,6-Tetrahydro-6-(2-hydroxy-2-phenylethyl)-1-methyl-2-pyridinyl]-2-propanone, 9CI [61550-92-3]

C₁₇H₂₃NO₂ 273.374

Major alkaloid from *Sedum acre* (Crassulaceae). Oil. [α]_D²² -156 (c, 1.7 in CHCl₃). Thermolabile, dec. on extraction with hot MeOH.

8-Ketone; perchlorate:

Cryst. (butanone). Mp 169°. [α]_D²⁰ -115 (c, 2 in MeOH).

3,4-Dihydro: **Dihydrosedinine**

[86108-22-7]

C₁₇H₂₇NO₂ 277.406

Alkaloid from *Sedum acre* (Crassulaceae). Cryst. (butanone) (as hydrochloride). Mp 165° (hydrochloride). [α]_D²⁰ -64 (c, 0.4 in MeOH).

3,4-Dihydro, diastereoisomer: **8-Methyl-10-phenyllobelidol**

C₁₇H₂₇NO₂ 277.406

Alkaloid from *Lobelia inflata* (Campanulaceae). Mp 91-92° (as hydrochloride). [α]_D²⁰ +3 (EtOH). Of unknown stereochem.

Schöpf, C. *et al.*, *Annalen*, 1957, **608**, 88 (8-Methyl-10-phenyllobelidol)

Franck, B. *et al.*, *Chem. Ber.*, 1958, **91**, 2803; 1959, **92**, 1001; 1960, **93**, 2360 (isol, ir, uv, biosynth)

Hootele, C. *et al.*, *Tet. Lett.*, 1980, **21**, 5063 (cryst struct, abs config)


Colau, B. *et al.*, *Can. J. Chem.*, 1983, **61**, 470 (isol, pmr, ms, struct, abs config, Sedacrine, Dihydrosedinine)

Driessens, F. *et al.*, *Can. J. Chem.*, 1991, **69**, 211 (synth)

Sedinone

S-215

1-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-2-propanone, 9CI. 8-Methyl-10-phenyllobelionol [67010-71-3]



C₁₇H₂₅NO₂ 275.39

Alkaloid from *Sedum acre* (Crassulaceae). Cryst. (petrol). Mp 93°.

Hydrochloride:

Needles (butanone). Mp 176-178° dec. (175°). [α]_D²² -79.4 (c, 1.02 in MeOH).

Hydroxy: **Hydroxysedinone**

[61565-33-1]

C₁₇H₂₅NO₃ 291.389


Alkaloid from *Sedum acre* (Crassulaceae). Posn. of hydroxylation unknown.

Hydroxy, 2''-alcohol: **Hydroxysedinol**

[66921-64-0]
 $C_{17}H_{27}NO_3$ 293.405
 Alkaloid from *Sedum acre* (Crassulaceae). Posn. of OH-group and stereochem. undetd.
 Franck, B. *et al.*, *Chem. Ber.*, 1958, **91**, 2803-2818; 1959, **92**, 1001 (*isol, uv, struct*)
 Kooy, J.H. *et al.*, *Planta Med.*, 1976, **30**, 295-296 (*Hydroxysedinine*)
 Francis, L.P.S. *et al.*, *Planta Med.*, 1977, **32**, 268-274 (*isol, Sedinine, Hydroxysedininol*)
 Colau, B. *et al.*, *Can. J. Chem.*, 1983, **61**, 470-472 (*struct, abs config, pmr, ms*)
 Driessens, F. *et al.*, *Can. J. Chem.*, 1991, **69**, 211-217 (*synth*)

Sedridine S-216

α -Methyl-2-piperidineethanol, 9CI. 2-(2-Hydroxypropyl)piperidine. 1-(2-Piperidino)-2-propanol. 8-Methylnorlobelol. Dihydroisopelletierine

(2*S*,2'*S*)-form $C_8H_{17}NO$ 143.228

Note potential confusion in stereochem. of allosedridines depending on whether the side-chain chiral centre is numbered 2 or 2'. Both have been used. Here it is numbered 2'.

(2*R*,2'*R*)-form

Synthetic. Mp 83-84°. $[\alpha]_D^{21}$ -28.1 (c, 2.31 in EtOH).

(2*S*,2'*S*)-form [501-83-7]

Alkaloid from *Sedum acre* and *Sedum maximum* (Crassulaceae). Mp 83-84°. $[\alpha]_D^{23}$ +29.3 (c, 2.28 in EtOH).

O,N-Dibenzoyl: Mp 101-103°. $[\alpha]_D$ -13.6 (c, 1.5 in $CHCl_3$).

(2*R*,2'*S*)-form*(+)*-Allosedridine

[26171-47-1]

Synthetic. Mp 62-63°. $[\alpha]_D^{29}$ +16.2 (c, 4 in MeOH).

O,N-Dibenzoyl: Mp 97-100°. $[\alpha]_D^{20}$ +75.6 (c, 1.8 in EtOH).

N-Me: *N*-Methylallosedridine $C_9H_{19}NO$ 157.255

Alkaloid from *Sedum sarmentosum* (Crassulaceae). Mp 125° (as picrate). $[\alpha]_D^{25}$ +67 (c, 0.9 in EtOH) (synthetic). $[\alpha]_D$ +25 (c, 2.48 in EtOH) (natural).

(2*S*,2'*R*)-form*(-)*-Allosedridine

Synthetic. Mp 62°. $[\alpha]_D^{29}$ -16.4 (c, 4 in MeOH).

(2*R*,2'*R*)-form*(±)*-Sedridine

Alkaloid from *Sedum acre* (Crassulaceae). Mp 75°.

Picrate: Mp 135-137°.*N*-Me: $C_9H_{19}NO$ 157.255

Mp 113-114° (as picrate). Bp₂₄ 103-105°.

(2*R*,2'*S*)-form*(±)*-Allosedridine

Mp 71-72°.

N-Me: Mp 126-127° (as picrate). Bp₂₄ 118-120°.

Marion, L. *et al.*, *Can. J. Res., Sect. B*, 1949, **27**, 215 (*isol, deriv*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1955, **74**, 1568; 1965, **84**, 1367; 1969, **88**, 1012 (*isol, synth, abs config, ir, ord*)

Fodor, G. *et al.*, *Tetrahedron, Suppl.*, No. 8, 1966, 113 (*synth*)

Schöpf, C. *et al.*, *Annalen*, 1970, **732**, 181

(synth)

Butruille, D. *et al.*, *Tetrahedron*, 1971, **27**, 2055 (*abs config, ir, pmr, ms*)

Beyerman, H.C. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1972, **91**, 1441 (*N-Methylallosedridine*)

Tufariello, J.J. *et al.*, *Tet. Lett.*, 1978, 4647

(synth)

Uyehara, T. *et al.*, *Tet. Lett.*, 1991, **32**, 4371

(synth)

Littler, B.J. *et al.*, *Synlett*, 1997, 22 (*synth*)


Nagata, K. *et al.*, *Heterocycles*, 2006, **70**, 335-344 (*synth*)

Chen, L.-J. *et al.*, *Tetrahedron: Asymmetry*, 2008, **19**, 715-720 (*Allosedridine, synth*)

Segoline A

S-217

[117694-96-9]



Absolute Configuration

 $C_{23}H_{19}N_3O_3$ 385.421

Alkaloid from the Red Sea tunicate *Eudistoma* sp. Cell proliferation inhibitor. Amorph. powder. Sol. MeOH. Mp 276°. $[\alpha]_D^{24}$ -322 (c, 1 in $CHCl_3$). λ_{max} 245 (ε 7400); 278 (sh) (ε 15600); 298 (ε 21300); 366 (ε 1400); 382 (ε 2500); 545 (ε 2500) (MeOH/HCl) (Derep). λ_{max} 236 (sh) (ε 9000); 270 (ε 16200); 308 (ε 5100); 320 (ε 5600); 368 (ε 1600); 383 (ε 2600); 460 (ε 3100) (MeOH) (Derep).

16-Epimer: Segoline C

[340162-11-0]

 $C_{23}H_{19}N_3O_3$ 385.421

Alkaloid from *Eudistoma bituminis*. $[\alpha]_D$ -374 ($CHCl_3$).

9,13-Diepimer: Segoline B

[122795-54-4]

 $C_{23}H_{19}N_3O_3$ 385.421

Alkaloid from the Red Sea tunicate *Eudistoma* sp. Amorph. powder. $[\alpha]_D^{24}$ +355 (c, 1 in $CHCl_3$). λ_{max} 245 (ε 7400); 278 (sh) (ε 15600); 298 (ε 21300); 366 (ε 1400); 382 (ε 2500); 545 (ε 2500) (MeOH/HCl) (Derep). λ_{max} 236 (sh) (ε 9000); 270 (ε 16200); 308 (ε 5100); 320 (ε 5600); 368 (ε 1600); 383 (ε 2600); 460 (ε 3100) (MeOH) (Derep).


Rudi, A. *et al.*, *J.O.C.*, 1989, **54**, 5331-5337 (*isol, uv, ir, pmr, cmr, cd, struct*)

Viracaoundin, I. *et al.*, *Tet. Lett.*, 2001, **42**, 2669-2671 (*isol, cd, pmr, cmr, abs config*)

Selaginoidine

S-218

[93888-55-2]

 $C_{18}H_{23}NO_4$ 317.384

Alkaloid from the twigs and foliage of *Athrotaxis selaginoides* and *Athrotaxis laxifolia* (Taxodiaceae). Mp 62-63°. $[\alpha]_D^{19}$ +166.7 (c, 0.25 in $CHCl_3$).


Panichanun, S. *et al.*, *Tetrahedron*, 1984, **40**, 2685 (*isol, uv, ir, pmr, cmr, ms, struct*)

Cassidy, M.P. *et al.*, *Org. Lett.*, 2005, **7**, 1339-1342 (*synth*)

Selenobiotin

S-219

Hexahydro-2-oxo-1*H*-selenolo[3,4-*d*]imidazole-4-pentanoic acid, 9CI
 [57956-29-3]

 $C_{10}H_{16}N_2O_3Se$ 291.208

Identified as an excretion prod. of *Phycomyces blakesleeanus*. Supports the growth of biotin-requiring microorganisms. Mp 234°. $[\alpha]_D^{25}$ +80.5 (c, 1 in 0.1*M* NaOH).

Bury, S. *et al.*, *Tet. Lett.*, 1976, 2033 (*synth*)


Lindblow-Kull, C. *et al.*, *Biochem. Biophys. Res. Commun.*, 1980, **93**, 572 (*isol*)

De Titta, G.T. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1980, **77**, 333 (*cryst struct*)

Sellowiine

S-220

N-Demethyl-20-deethylsuaveoline
 [173693-51-1]

 $C_{17}H_{15}N_3$ 261.326


Struct. reported has been shown by synth. to be incorrect in 2004. Alkaloid from leaves of *Rauwolfia sellowii*. Cryst. (MeOH). Mp 202-206°. $[\alpha]_D$ -135 (c, 0.54 in $CHCl_3$). λ_{max} 222 ; 268 ; 282 ; 290 (MeOH).

Batista, C.V.F. *et al.*, *Phytochemistry*, 1996, **41**, 969-973 (*isol, pmr, cmr, ms*)

Ohba, M. *et al.*, *Heterocycles*, 2004, **63**, 2845-2850 (*synth*)

Semecarpifoline S-221

4,7-Dimethoxy-3-(methoxymethyl)-2(1H)-quinolinone




$C_{13}H_{15}NO_4$ 249.266
Alkaloid from the root bark of *Melicope semecarpifolia*. Yellow needles (MeOH). Mp 123-127°. λ_{max} 218 (log ϵ 4.57); 284 (log ϵ 3.83); 325 (log ϵ 4.06); 337 (log ϵ 3.97) (MeOH).

Chen, I.S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1143-1147

Semecarpine S-222

[632335-16-1]




$C_{17}H_{17}NO_3$ 283.326
Alkaloid from the leaves of *Melicope semecarpifolia*. Pale yellow needles ($CHCl_3/Me_2CO$). Mp 145-147°. λ_{max} 227 (log ϵ 4.61); 262 (log ϵ 4.51); 279 (log ϵ 4.44); 325 (log ϵ 4.05); 411 (log ϵ 3.78) (EtOH).

Chen, J.-J. *et al.*, *Planta Med.*, 2003, **69**, 542-546 (*isol*, *pmr*, *cmr*, *ms*)

Semiplenamidine C S-223

[630100-43-5]



Absolute Configuration

$C_{22}H_{41}NO_3$ 367.571
Alkaloid from *Lyngbya semiplena*. Amorph. solid. $[\alpha]_D^{26}$ -5 (c, 0.3 in $CHCl_3$). λ_{max} 213 (ϵ 6500) (MeOH).

Han, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1364-1368 (*isol*, *pmr*, *cmr*)

Davies, I.R. *et al.*, *Tet. Lett.*, 2005, **46**, 5547-5549 (*synth*)

Semiplenamidine D S-224

[630100-44-6]

As Semiplenamidine C, S-223 with $n = 16$

$C_{26}H_{49}NO_3$ 423.678
Alkaloid from *Lyngbya semiplena*. Amorph. solid. $[\alpha]_D^{26}$ -10.6 (c, 0.15 in

$CHCl_3$). λ_{max} 207 (ϵ 5200) (MeOH).

Han, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1364-1368 (*isol*, *pmr*, *cmr*)

Semiplenamidine E S-225

[630100-45-7]

As Semiplenamidine C, S-223 with $n = 14$

$C_{24}H_{45}NO_3$ 395.624
Alkaloid from *Lyngbya semiplena*. Amorph. solid. $[\alpha]_D^{26}$ -7.1 (c, 0.28 in $CHCl_3$). λ_{max} 207 (ϵ 4300) (MeOH).

2R*,3S*-Epoxide: **Semiplenamidine G**

[630100-47-9]

$C_{24}H_{45}NO_4$ 411.624
Alkaloid from *Lyngbya semiplena*. Amorph. solid. $[\alpha]_D^{26}$ -3 (c, 0.6 in $CHCl_3$). 1'-Config. not determined. λ_{max} 207 (ϵ 6100) (MeOH).

2R*,3S*-Epoxide, O-de-Ac: **Semiplenamidine F**


[630100-46-8]

$C_{22}H_{43}NO_3$ 369.587
Alkaloid from *Lyngbya semiplena*. Amorph. solid. $[\alpha]_D^{26}$ -5 (c, 0.3 in $CHCl_3$). 1'-Config. not determined. λ_{max} 205 (ϵ 5500) (MeOH).

Han, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1364-1368 (*Semiplenamides E-G*)

Semperviraminol S-226

[231628-55-0]



$C_{35}H_{52}N_2O_4$ 564.807
Alkaloid from the roots of *Buxus sempervirens*. Amorph. solid. $[\alpha]_D^{20}$ +97 (c, 0.6 in $CHCl_3$). λ_{max} 226 (log ϵ 2.3) (MeOH).

9,11-Didehydro: **Hyrscanol**


$C_{35}H_{50}N_2O_4$ 562.791
Alkaloid from the leaves of *Buxus hyrcana*. Amorph. solid. $[\alpha]_D^{24}$ -4.8 (c, 0.08 in $CHCl_3$). λ_{max} 226 (log ϵ 2.3) (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1999, **62**, 665-669 (*Semperviraminol*)

Choudhary, M.I. *et al.*, *Chem. Biodiversity*, 2006, **3**, 1039-1042 (*Hyrscanol*)

Semperviraminone S-227

[191288-30-9]



$C_{26}H_{39}NO$ 381.6

Alkaloid from the roots of *Buxus sempervirens* (Buxaceae). Gum. $[\alpha]_D^{20}$ +0.5 (c, 1.2 in $CHCl_3$). λ_{max} 240 (MeOH).

N-De-Me: **N-Demethylsemperviraminone**

[191101-05-0]

$C_{25}H_{37}NO$ 367.573


Alkaloid from the roots of *Buxus sempervirens* (Buxaceae). Gum. $[\alpha]_D^{20}$ +0.5 (c, 0.8 in $CHCl_3$). λ_{max} 240 (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1997, **60**, 770-774 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Sempervirine† S-228

2,3,4,13-Tetrahydro-1H-benz[*g*]indolo[2,3-*a*]quinolizin-6-ium, 9CI. *Sempervirine*

[6882-99-1 (cation)]



$C_{19}H_{16}N_2$ 272.349

Alkaloid from *Gelsemium sempervirens* and *Rauwolfia* sp. Shows antitumour props. Intercalating agent. Red-orange powder. Sol. EtOH, $CHCl_3$, Py; poorly sol. C_6H_6 , Et_2O . Mp 258-260° (223°, 254°). λ_{max} 242 ; 249 ; 294 ; 342 ; 385 (95% EtOH).

Hydrochloride: pK_a 10.6 (18°).

Hydrobromide: Mp 323-325° dec.

Nitrate: Mp 267° dec.

N¹-Me: [6484-78-2]

[47197-61-5 (cation)]

$C_{20}H_{19}N_2^+$ 287.384

Mp 330-332° (as chloride).

Stevenson, A.E. *et al.*, *J. Am. Pharm. Assoc.*, 1916, **4**, 1458-1463; *CA*, **10**, 804 (*isol*)

Prelog, V. *et al.*, *Helv. Chim. Acta*, 1948, **31**, 588-593 (*uv*)

Woodward, R.B. *et al.*, *J.A.C.S.*, 1949, **71**, 379-380 (*struct*, *synth*, *methochloride*)

Potts, K.T. *et al.*, *J.O.C.*, 1968, **33**, 3985-3987 (*synth*, *ir*, *uv*)


Beljanski, M. *et al.*, *Oncology*, 1986, **43**, 198-203; *CA*, **105**, 54186v (*pharmacol*)

Gribble, G.W. *et al.*, *Tetrahedron*, 1988, **44**, 3195-3202 (*synth*, *uv*, *ir*, *pmr*, *cmr*)

Chatterjee, A. *et al.*, *Monatsh. Chem.*, 1996, **127**, 1259-1262 (*synth*, *uv*)

Sempervirone S-229

[116537-79-2]



$C_{26}H_{39}NO_2$ 397.6


Stereochem. revised in 1998.
Alkaloid from the leaves of *Buxus sempervirens* (Buxaceae). Amorph. solid. $[\alpha]_D^{20}$ -65 (c, 0.5 in CHCl_3). λ_{max} 243 (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1988, **51**, 783-786 (*isol, uv, ir, pmr, ms, struct*)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1998, **48**, 519-528 (*isol, pmr, ms, config*)

Semperviroxazolidine S-230

[220705-50-0]




$\text{C}_{26}\text{H}_{39}\text{NO}_2$ 397.6

Alkaloid from the roots of *Buxus sempervirens*. Gum. $[\alpha]_D^{20}$ -45 (c, 0.38 in CHCl_3). λ_{max} 231 (sh); 238; 245; 254 (sh) (MeOH).

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 299-306 (*isol, pmr, cmr*)

Senacarcin A S-231

DC 59A. Antibiotic DC 59A [82800-43-9]



$\text{C}_{25}\text{H}_{26}\text{N}_4\text{O}_7$ 494.503

Phenazine antibiotic. Deriv. of Griseoliteic acid in H-514. Prod. by *Streptomyces endus* subsp. *aureus*. Active against gram-positive bacteria and tumours and weakly active against gram-negative organisms. Yellow needles. Sol. MeOH, H_2O , CHCl_3 , DMSO; poorly sol. hexane, H_2O . Mp 190°. λ_{max} 210 (ϵ 3950); 263 (ϵ 4200); 350 (sh) (ϵ 500); 366 (ϵ 700) (95% EtOH) (Derep).

O-Deacyl: **Senacarcin H**. DC 59H. Antibiotic DC 59H

[81782-37-8]
 $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_5$ 379.371


Prod. by *Streptomyces endus aureus*. Orange cryst. Sol. MeOH, H_2O , CHCl_3 , DMSO; poorly sol. hexane, H_2O . λ_{max} 215; 263; 365; 445 (EtOH).

Nakano, H. *et al.*, *J. Antibiot.*, 1982, **35**, 760 (*isol, uv, ir, pmr*)

Japan. Pat., 1982, 82 4 975; CA, **96**, 197887f (*isol, pharmacol*)

Senaetnine S-232

12-(Acetyloxy)-3,8-didehydro-senecionan-5,11,16-trione, 9CI [64191-69-1]



$\text{C}_{20}\text{H}_{23}\text{NO}_7$ 389.404

Alkaloid from *Senecio* spp. (Asteraceae). Cryst. (Et_2O /petrol). Mp 183.5°. $[\alpha]_D^{24}$ +10.6 (c, 2.5 in CHCl_3).

15E-Isomer: **14,15-trans-Senaetnine**

[71075-41-7]

$\text{C}_{20}\text{H}_{23}\text{NO}_7$ 389.404

Alkaloid from *Senecio aucheri* (Asteraceae). Oil. Not obt. free of Senaetnine. The 14,15-numbering arises from use of a nonstandard numbering system. No stereochem. detd.

13,19-Didehydro: **Dehydro-senaetnine**.

(+)-Dehydroisosenaetnine

[68353-30-0]

$\text{C}_{20}\text{H}_{21}\text{NO}_7$ 387.388

Alkaloid from *Senecio barbertonicus* (Asteraceae). Oil. Referred to by the authors as (+)-Dehydroisosenaetnine, which seems misleading as it is apparently shown to be dehydro-senaetnine.

7-Epimer: **Isosenaetnine**

[67920-39-2]

$\text{C}_{20}\text{H}_{23}\text{NO}_7$ 389.404

Alkaloid from above-ground parts of *Kleinia kleinioides* (Asteraceae). Cryst. (Et_2O /petrol). Mp 198.5°. $[\alpha]_D^{24}$ -34.8 (c, 1.0 in CHCl_3). Stereochem. undefined.

7-Epimer, 13,19-didehydro: **Dehydroisosenaetnine**

[67951-60-4]

$\text{C}_{20}\text{H}_{21}\text{NO}_7$ 387.388


Alkaloid from *Kleinia kleinioides* (Asteraceae). Viscous oil. $[\alpha]_D^{24}$ -45.6 (c, 0.5 in CHCl_3).

Bohlmann, F. *et al.*, *Phytochemistry*, 1977, **16**, 965; 1978, **17**, 599; 1979, **18**, 79 (*isol, ir, pmr, ms, struct*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1978, **111**, 3009 (*cd, stereochem*)

Klose, W. *et al.*, *Chem. Ber.*, 1980, **113**, 2694

Senampelines S-233



Senampeline	R	R'	12-configuration
A	Tigloyl	(H,C) ₂ C=CHCO	(E-)
B	Tigloyl	Tigloyl	(E-)
C	Tigloyl	(H,C) ₂ C=CHCO	(Z-)
D	Tigloyl	Tigloyl	(Z-)
E	Tigloyl	Angeloyl	(E-)
F	(H,C) ₂ C=CHCO	Angeloyl	(E-)
G	Angeloyl	Angeloyl	(E-)

A family of isomeric alkaloids from *Senecio cissampelinus*, *Senecio mikanioides* and other *Senecio* spp. (Asteraceae).

Senampeline A [62787-00-2]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Oil. Not separated from Senampeline B.

Senampeline B [62860-52-0]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Oil. Not sepd. from Senampeline A.

Senampeline C [62841-11-6]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Oil. Not sepd. from Senampeline D.

Senampeline D [62787-01-3]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Oil. Not sepd. from Senampeline C.

Senampeline E [71075-42-8]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Not sepd. from Senampelines F and G.

Senampeline F [71075-43-9]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Not sepd. from Senampelines E and G.

Senampeline G [71075-44-0]

$\text{C}_{25}\text{H}_{31}\text{NO}_8$ 473.522

Not sepd. from Senampelines E and F.

5-Deacetoxy, 5-oxo: **5-Deacetoxy-5-oxo-senampeline G**

$\text{C}_{23}\text{H}_{27}\text{NO}_7$ 429.469


Alkaloid from *Senecio stapeliaeformis* (Asteraceae). Alkaloid not named in the lit.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1977, **110**, 474 (*isol, uv, ir, pmr, ms, struct*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1979, **18**, 79; 1986, **25**, 1151 (*isol, struct, deriv*)

Sendaverine S-234

1,2,3,4-Tetrahydro-6-methoxy-2-[(4-methoxyphenyl)methyl]-7-isoquinolinol, 9CI. 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-(4-methoxybenzyl)isoquinoline. Alkaloid F28 [5056-80-4]



$\text{C}_{18}\text{H}_{21}\text{NO}_3$ 299.369

Alkaloid from *Corydalis aurea*, *Corydalis gortschakovii* and *Corydalis tashiroi* (Papaveraceae). Fine needles (Et_2O), cubes (hexane). Mp 140-141.5°.

Hydrochloride:

Needles (MeOH/ Et_2O). Mp 225-230° dec.

N-Oxide: **Sendaverine N-oxide**

[65907-09-7]

$\text{C}_{18}\text{H}_{21}\text{NO}_4$ 315.368

Alkaloid from the aerial parts of *Corydalis gortschakovii* (Papaveraceae).

O^4 -*De-Me*: 1,2,3,4-Tetrahydro-2-[(4-hydroxyphenyl)methyl]-6-methoxy-7-isquinolinol. 1,2,3,4-Tetrahydro-7-hydroxy-2-(4-hydroxybenzyl)-6-methoxyisoquinoline. **Corgoine** [15778-86-6]
 $C_{17}H_{19}NO_3$ 285.342
 Alkaloid from *Corydalis gortschakovii* (Papaveraceae). Needles (MeOH/Et₂O), cryst. (CHCl₃/MeOH). Mp 198° (190-191°).

O^4 -*De-Me*, hydrochloride:
 Cryst. (MeOH/Et₂O). Mp 239-240°.

O^6 -*De-Me*, O^7 -*Me*: **Isosendaverine** [156401-02-4]
 $C_{18}H_{21}NO_3$ 299.369
 Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Pale yellow cryst. (MeOH). Mp 181-182°.

3'-Hydroxy: **Capnosine** [148234-97-3]
 $C_{18}H_{21}NO_4$ 315.368
 Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Yellowish powder. Mp 105-106°.

3'-Hydroxy, O^7 -*Me*: **Capnosinine** [156401-03-5]
 $C_{19}H_{23}NO_4$ 329.395
 Alkaloid from *Ceratocarpus heterocarpa* (Papaveraceae). Cryst. (EtOH). Mp 172-173°.

3'-Methoxy, O^7 -*Me*: **Intebrimine** $C_{20}H_{25}NO_4$ 343.422
 Alkaloid from leaves of *Berberis integerrima*. Cryst. (as hydrochloride). Mp 169-170° (hydrochloride).

Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1936, **16**, 81 (isol)

Kametani, T. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 608 (uv, ir, pmr, ms, struct, synth)

Ibragimova, M.U. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 438; 638; 1971, **7**, 211; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 447; 660; 1971, **7**, 209 (Sendaverine, Corgoine, isol, synth, pmr)

Suguna, H. *et al.*, *Indian J. Chem.*, 1974, **12**, 1141 (Corgoine, synth, uv, ir, pmr)

Kametani, T. *et al.*, *Tetrahedron*, 1975, **31**, 235 (Corgoine, synth, ir, pmr)

Mori, M. *et al.*, *Heterocycles*, 1977, **6**, 1841 (synth, ir, pmr, ms)

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 834; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 702 (isol, oxide)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1979, 2836 (synth)

Pandey, G.D. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 160 (synth)

Pandey, G.D. *et al.*, *Pol. J. Chem. (Roc. Chem.)*, 1980, **54**, 763 (Corgoine, synth)

Masood, M. *et al.*, *Synth. Commun.*, 1980, **10**, 541 (synth, ir, pmr, ms)

Tani, C. *et al.*, *Planta Med.*, 1981, **41**, 403 (isol)

Otomasu, H. *et al.*, *J.C.S. Perkin I*, 1982, 2399 (Sendaverine, Corgoine, synth, pmr)


Takano, S. *et al.*, *Heterocycles*, 1993, **35**, 47 (synth)

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 70; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 57 (Intebrimine)

Suau, R. *et al.*, *Phytochemistry*, 1994, **36**, 241 (Isosendaverine, Capnosine, Capnosinine)

Senecicaudatin

[106283-48-1]

 $C_{13}H_{21}NO_5$ 271.313

O-(3-Methylbutanoyl): **Senecicaudatin O-isopentanoate**

[106283-50-5]

 $C_{18}H_{29}NO_6$ 355.43

Alkaloid from *Senecio caudatus* (Asteraceae). Alkaloid not named in the paper.

O-(3-Methyl-2-butanoyl): **Senecicaudatin O-senecioate**

[106283-49-2]


 $C_{18}H_{27}NO_6$ 353.414

Alkaloid from *Senecio caudatus* (Asteraceae). Not named in the paper.

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151-1159 (isol, ir, ms, struct)

Senecicaudatinal hemiacetal

S-236

 $C_{13}H_{19}NO_4$ 253.297


Alkaloid from *Senecio caudatus* (Asteraceae).

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151 (isol, ir, ms, struct)

Seneciocannabine

S-237

[81855-31-4]

 $C_{18}H_{23}NO_7$ 365.382

Diepoxide of Seneciphylline, S-240. Alkaloid from roots and aerial parts of *Senecio cannabifolius* (Asteraceae). Prisms. Mp 198°. $[\alpha]_D^{25}$ -8.9 (CHCl₃).

► Poss. carcinogen. VT5704550

Asada, Y. *et al.*, *Tet. Lett.*, 1982, 189 (isol, cmr, cryst struct)

Asada, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 4616 (synth)

Mori, H. *et al.*, *Cancer Res.*, 1985, **45**, 3125 (tox)

Absolute configuration


Senecionine

S-238

12-Hydroxysenecionan-11,16-dione, 9CI.

Aureine†

[130-01-8]



Absolute configuration

 $C_{18}H_{25}NO_5$ 335.399

Cyclic ester of 2,3,5,7a-Tetrahydro-1-hydroxy-1H-pyrrolizine-7-methanol, T-188 with 5-Ethylidene-2-hydroxy-2,3-dimethylhexanedioic acid. Alkaloid from many *Senecio* spp., *Brachyglottis repanda*, *Emilia sonchifolia*, *Erechtites hieracifolia*, *Petasites hybridus* and other Asteraceae. Large amts. obtainable from *Senecio triangularis*; also obt. from *Castilleja rhexifolia* (Scrophulariaceae). Shows antineoplastic activity vs. Walker 256 carcinosarcoma. Mp 232-233°. $[\alpha]_D$ -56 (CHCl₃). Log P -0.04 (uncertain value) (calc). Hieracifoline and Petenephine were mixts. of Senecionine and Seneciphylline, S-240.

► LD₅₀ (rat, ivn) 41 mg/kg. LD₅₀ (mus, ivn) 64 mg/kg. Exp. reprod. and teratogenic effects. Hepatotoxic, also causes lung lesions. VT5710000

N-Oxide: **Senecionine N-oxide**

[13268-67-2]

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from seeds of *Crotalaria anagyroides* (Fabaceae). Antineoplastic agent. Hygroscopic solid. Mp 140-141° dec. (synthetic). Log P -2.01 (uncertain value) (calc).

► LD₅₀ (mus, ipr) 75 mg/kg.

15E-Isomer: **Integerrimine. Squalidine.**

Alkaloid SD

[480-79-5]

 $C_{18}H_{25}NO_5$ 335.399

Alkaloid from *Senecio* spp., *Crotalaria incana* and other *Crotalaria* spp., *Cacalia hastata* and others (Fabaceae, Asteraceae). Strong insect antifeedant agent. Mp 172°. $[\alpha]_D$ -29 (-18.3) (CHCl₃). $[\alpha]_D$ +4 (MeOH). λ_{max} 210 (ε 10620) (MeOH) (Berdy).

► NM9080000

15E-Isomer, N-oxide: **Integerrimine N-oxide**

[85955-28-8]

 $C_{18}H_{25}NO_6$ 351.399

Alkaloid from *Senecio erucifolius*, *Senecio nebrodensis* and *Senecio vulgaris*. Strong insect antifeedant agent. Oil. $[\alpha]_D^{25}$ +8.5 (c,0.02 in EtOH).

15E-Isomer, O-Ac: **Acetylintegerrimine**

[119328-82-4]

 $C_{20}H_{27}NO_6$ 377.436

Alkaloid from the seeds of *Crotalaria naragutensis* (Fabaceae).

15E-Isomer, 12-epimer: **Usaramoensine.**

Usaramensine, 8CI

[480-80-8]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio usaramoensis*. Prisms (Me₂CO). Mp 221° dec. (sealed tube). [α]_D²⁵ -25.2 (CHCl₃). Struct. not certain. Considered by Adams *et al.* to be the 12-epimer of Senecionine, reassigned by Gonzalez *et al.* as the 12-epimer, 15*E*-isomer but the evidence is not conclusive.

15*E*-Isomer, 12-epimer, picrate:

Cryst. (MeOH). Mp 235° dec.

15*E*-Isomer, 21-hydroxy: **21-Hydroxyintegerrimine**. *Eruciflorine*

[135626-66-3]

C₁₈H₂₅NO₆ 351.399

Alkaloid from *Senecio argunensis*, *Senecio erucifolius* and *Senecio jacobaea* (Asteraceae). Gum. [α]_D²⁵ -2 (c, 1.5 in EtOH).

18-Hydroxy: see Retrorsine, R-62

Barger, G. *et al.*, *J.C.S.*, 1936, 743 (*isol*)Kropman, M. *et al.*, *J.C.S.*, 1950, 700

(Integerrimine)

Adams, R. *et al.*, *J.A.C.S.*, 1953, 75, 4631

(Integerrimine, Usaramoensine)

Warren, F.L. *et al.*, *Prog. Chem. Org. Nat.**Prod.*, 1955, 12, 198; 1966, 24, 329 (*occur*,*bibl*)Gonzalez Gonzalez, A. *et al.*, *An. R. Soc. Esp.**Fis. Quim.*, Ser. B, 1958, 54, 223

(Usaramoensine)

Šantavý, F. *et al.*, *Coll. Czech. Chem. Comm.*,1962, 27, 1666 (*props*, Integerrimine)Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*,1969, 34, 1832 (*uv*)Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653

(cd)

Crout, D.H.G. *et al.*, *J.C.S. Perkin I*, 1972, 671

(biosynth)

Abdullaev, U.A. *et al.*, *Khim. Prir. Soedin.*,1974, 10, 538; *Chem. Nat. Compd. (Engl.**Transl.)*, 1974, 10, 556 (*ms*)Bale, N.M. *et al.*, *J.C.S. Perkin I*, 1978, 101

(biosynth)

Mackay, M.F. *et al.*, *Acta Cryst. B*, 1982, 38,2754 (*cryst struct*)Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, 35, 1173

(cmr)

Rastogi, R. *et al.*, *Pharmazie*, 1982, 37, 75 (*N-**oxide*)Molyneux, R.J. *et al.*, *Phytochemistry*, 1982,21, 439 (*cmr*)Robins, D.J. *et al.*, *Prog. Chem. Org. Nat.**Prod.*, 1982, 41, 115 (*occur*, *bibl*)Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1983, 36,1203 (*isol*)Narasaka, K. *et al.*, *J.A.C.S.*, 1984, 106, 2954

(synth, Integerrimine)

Mattocks, A.R. *et al.*, *Chemistry and**Toxicology of Pyrrolizidine Alkaloids*,Academic Press, 1986, (*tox*, *rev*)Hanson, B.A. *et al.*, *J.A.C.S.*, 1988, 110, 6314

(conformn)

Mattocks, A.R. *et al.*, *Phytochemistry*, 1988,27, 3289 (*Acetylintegerrimine*)Barrero, A.F. *et al.*, *An. Quim.*, 1991, 87, 386-390 (*Integerrimine N-oxide*)Liu, K. *et al.*, *Phytochemistry*, 1991, 30, 1303

(21-Hydroxyintegerrimine)

White, J.D. *et al.*, *J.O.C.*, 1992, 57, 2270 (*synth*,*Integerrimine*)Witte, L. *et al.*, *Phytochemistry*, 1992, 31, 559

(Eruciflorine)

Niwa, H. *et al.*, *Tetrahedron*, 1992, 48, 393


(synth, Integerrimine)

Niwa, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1994,67, 1990 (*synth*)Reina, M. *et al.*, *J. Nat. Prod.*, 2001, 64, 6-11(Integerrimine *N-oxide*, activity)Lewis, R.J. *et al.*, *Sax's Dangerous Properties**of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, ARS500; IDG000

Senecioracene

[146573-88-8]


C₁₉H₂₇NO₇ 381.425

Alkaloid from *Senecio racemosus* (Asteraceae). Oil. [α]_D²⁵ +50 (c, 0.2 in CHCl₃).

Ahmed, W. *et al.*, *J. Nat. Prod.*, 1992, 55, 1764-1767 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)**Seneciphylline**

S-240

13,19-Didehydro-12-hydroxysenecionan-11,16-dione, 9CI. α-Longilobine. Jacodine [480-81-9]



Absolute configuration

C₁₈H₂₃NO₅ 333.383

Alkaloid from numerous *Senecio* and *Crotalaria* spp., also *Erechtites hieracifolia* and *Erechtites quadridentata* (Asteraceae, Fabaceae). Cause of grazing toxicity in animals. Plates (EtOAc). Mp 217-218° dec. [α]_D²⁵ -139 (CHCl₃).

► LD₅₀ (rat, ipr) 77 mg/kg, hepatotoxic, other exp. toxic effects. VT5977000

N-Oxide: Seneciphylline N-oxide

[38710-26-8]

C₁₈H₂₃NO₆ 349.383

Alkaloid from whole plants of *Senecio persoonii* (Asteraceae). [α]_D²⁰ -56 (c, 1 in EtOH).

O-Ac: Seneciphyllinine. 12-Acetyl-seneciphylline

[90341-45-0]

C₂₀H₂₅NO₆ 375.421

Alkaloid from *Gynura segetum* and *Senecio pterophorus* (Asteraceae). Cryst. Mp 82-83°.

13ξ,19-Epoxide: Seneciphylline epoxide

[106356-85-8]

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Senecio megaphyllus* and *Senecio usgorensis* (Asteraceae). Alkaloid not named in the lit.

18-Hydroxy: Riddelline. Riddelliine

[23246-96-0]

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Senecio riddellii*, *Senecio longiflorus*, *Senecio eremophilus*, *Crotalaria juncea* etc. (Asteraceae, Fabaceae). Mp 197-198° dec. [α]_D²⁵ -109.5 (CHCl₃).

► LD₅₀ (mus, ivn) 105 mg/kg, hepatotoxic. Possible human carcinogen (IARC 2B). VJ3850000

15*E*-Isomer: (E)-Seneciphylline

[129029-37-4]

C₁₈H₂₃NO₅ 333.383

Alkaloid from *Gynura segetum* (Asteraceae).

15*E*-Isomer, 13ξ,19-epoxide: (E)-Seneciphylline epoxide

[106356-86-9]

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Senecio megaphyllus* (Asteraceae). Not named in the lit.

12-Epimer, 15*E*-isomer: Spartioidine

[520-59-2]

C₁₈H₂₃NO₅ 333.383

Alkaloid from *Senecio spartioides* (Asteraceae). Mp 178°. [α]_D²⁵ -83.7 (c, 1.73 in 95% EtOH).

► WG6475000

12-Epimer, 15*E*-isomer, N-oxide: Spartioidine N-oxide

[121123-61-3]

C₁₈H₂₃NO₆ 349.383

Alkaloid from *Senecio vulgaris*.

Adams, R. *et al.*, *J.A.C.S.*, 1942, 64, 2760-2763; 1957, 79, 174-177 (*Riddelline*,*Spartioidine*)Warren, F.L. *et al.*, *Prog. Chem. Org. Nat.**Prod.*, 1955, 12, 201 (*occur*)Masamune, S. *et al.*, *Chem. Ind. (London)*,1959, 21 (*struct*)Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1965,18, 1625 (*pmr*)Cervinka, O. *et al.*, *Chem. Comm.*, 1968, 261

(abs config)

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*,1969, 34, 1832 (*uv*)Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*,1972, 37, 3918 (*cd*)Abdullaev, U.A. *et al.*, *Khim. Prir. Soedin.*,1974, 10, 538; *Chem. Nat. Compd. (Engl.**Transl.)*, 1974, 10, 556 (*ms*)Edgar, J.A. *et al.*, *Experientia*, 1976, 32, 1535

(Acetyl-seneciphylline)

Molyneux, R.J. *et al.*, *Phytochemistry*, 1982,21, 439 (*cmr*)Robins, D.J. *et al.*, *Prog. Chem. Org. Nat.**Prod.*, 1982, 41, 115-203 (*occur*)Wiedenfeld, H. *et al.*, *Arch. Pharm.*(Weinheim, Ger.), 1984, 317, 97 (*cryst**struct*)Rana, J. *et al.*, *Chem. Comm.*, 1985, 1742

(biosynth)


Mattocks, A.R. *et al.*, *Chemistry and**Toxicology of Pyrrolizidine Alkaloids*,Academic Press, 1986, (*tox*, *rev*)Bohlmann, F. *et al.*, *Phytochemistry*, 1986, 25,1151 (*epoxides*)Yuan, S.Q. *et al.*, *Yaoxue Xuebao*, 1990, 25,191-197 (*Seneciphyllinine*, (E)-*Seneciphylline*)Roeder, E. *et al.*, *Phytochemistry*, 1993, 32,1051 (*Seneciphylline N-oxide*)Liddell, J.R. *et al.*, *Phytochemistry*, 1993, 34,1629 (*Acetyl-seneciphylline*)IARC Monog. (Web), (*tox*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties**of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, RJZ000; SBX500

Senecivernine

S-241

12-Hydroxy-14-methyl-21-norsenecionan-11,16-dione, 9CI
[72755-25-0]



Absolute
Configuration

C₁₈H₂₅NO₅ 335.399

Cyclic diester of retronecine with seneci-verninc acid (unique to this alkaloid). Alkaloid from *Senecio vernalis* above-ground parts (Asteraceae). Mp 105-107° dec. [α]_D²⁰ -34.9 (EtOH).

N-Oxide: Senecivernine N-oxide
[101687-28-9]

C₁₈H₂₅NO₆ 351.399

Alkaloid from *Senecio vernalis* and *Senecio procumbens*. Brown oil.

Röder, E. et al., *Planta Med.*, 1979, **37**, 131 (isol, ms, pmr, cmr, ir, struct)


Parvez, M. et al., *Acta Cryst. C*, 1995, **51**, 1202-1204 (cryst struct, abs config)

Liu, Z.-Y. et al., *Tet. Lett.*, 1999, **40**, 5593-5596 (synth)

Kostova, N. et al., *J. Serb. Chem. Soc.*, 2006, **71**, 1275-1280 (N-oxide)

Senepodine A

S-242



Absolute
Configuration

C₂₃H₄₀N₂ 344.582

Alkaloid from *Lycopodium chinense*. Cytotoxic agent. Solid. [α]_D -33 (c, 0.6 in MeOH).

N-De-Me: Senepodine C

C₂₂H₃₈N₂ 330.556

Alkaloid from *Lycopodium chinense*. Solid. [α]_D²⁵ -15 (c, 1 in MeOH).

N-De-Me, N-formyl: Senepodine D

C₂₃H₃₈N₂O 358.566

Alkaloid from *Lycopodium chinense*. Solid. [α]_D²⁵ -33 (c, 0.5 in MeOH).

N-De-Me, N-Ac: Senepodine E

C₂₄H₄₀N₂O 372.593

Alkaloid from *Lycopodium chinense*. Solid. [α]_D²² -106 (c, 1 in MeOH).

4αβ,5'α-Dihydro, N-de-Me, N-Ac: Senepodine F

C₂₄H₄₂N₂O 374.609

Alkaloid from *Lycopodium chinense*. Amorph. solid. [α]_D²³ -35 (c, 0.5 in MeOH).

6(N)-Dehydro: Senepodine B

C₂₃H₃₉N₂[⊕] 343.575

Quaternary alkaloid from *Lycopodium chinense*. Solid. [α]_D²⁷ -38 (c, 0.2 in

MeOH). Counterion not specified.

Morita, H. et al., *Tet. Lett.*, 2001, **42**, 4199-4201 (isol, pmr, cmr)


Hirasawa, Y. et al., *Tetrahedron*, 2003, **59**, 3567-3573 (isol, pmr, cmr, abs config)

Hirasawa, Y. et al., *Heterocycles*, 2004, **64**, 515-521 (Senepodine F)

Senepodine G

S-243

1,2,3,6,7,8,9,9a-Octahydro-2,4-dimethyl-quinolinizinium(1+)
[760947-76-0]



Relative
Configuration

C₁₁H₂₀N[⊕] 166.286

Quaternary alkaloid from *Lycopodium chinense*. Amorph. solid (as trifluoroacetate). [α]_D²⁶ -35 (c, 0.3 in MeOH) (trifluoroacetate).


Morita, H. et al., *Tetrahedron*, 2004, **60**, 7015-7023 (isol, pmr, cmr)

Snider, B.B. et al., *J.O.C.*, 2007, **72**, 1039-1042 (synth)

Senepodine H

S-244

[760947-77-1]



Absolute
Configuration

C₁₄H₂₆NO[⊕] 224.366


Quaternary alkaloid from *Lycopodium chinense*. Amorph. solid (as trifluoroacetate salt). [α]_D²⁷ -35 (c, 0.2 in MeOH) (trifluoroacetate salt).

Morita, H. et al., *Tetrahedron*, 2004, **60**, 7015-7023 (isol, pmr, cmr)

Senkirkine

S-245

12-Hydroxy-4-methyl-4,8-secosenecionan-8,11,16-trione, 9CI. Renardine
[2318-18-5]



C₁₉H₂₇NO₆ 365.425

Cyclic otonecine diester. Alkaloid from *Nardosmia laevigata*, (preferred genus name *Petasites*), *Farfugium japonicum*, *Crotalaria laburnifolia*, *Senecio anonyms*, *Senecio kirkii* and others (Asteraceae, Fabaceae). Bevelled plates (EtOAc or Me₂CO). Sol. MeOH, CHCl₃; poorly

sol. H₂O. Mp 196.5-197.5°. [α]_D²⁵ -16 (c, 1.89 in MeOH). pK_a 6.4. λ_{max} 219 (ε 10500) (MeOH) (Berdy).

▶ LD₅₀ (rat, ipr) 220 mg/kg. Exp. neoplastic agent, hepatotoxic. VT5960000

Picrate:

Yellow rods (EtOH aq.). Mp 225-225.5° (219-220°).

Ac: O-Acetylsenkirkine

[53092-44-7]

C₂₁H₂₉NO₇ 407.463

Alkaloid from *Senecio kirkii* (Asteraceae). Needles (EtOAc/Me₂CO). Mp 195-196°. [α]_D²⁵ -34 (c, 0.44 in MeOH).

1R,2-Dihydro: Dihydrosenkirkine

[135574-55-9]

C₁₉H₂₉NO₆ 367.441

Alkaloid from *Senecio integrifolius* var. *fauriri* (Asteraceae). Exists in equilib. as two ring structs.

18-Hydroxy: Hydroxysenkirkine

[37819-28-6]

C₁₉H₂₇NO₇ 381.425

Alkaloid from *Crotalaria laburnifolia* (Fabaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 124-125°. [α]_D²⁶ +5.3 (c, 0.682 in EtOH).

▶ Hepatotoxic.

19-Hydroxy: 18-Hydroxysenkirkine

[106312-35-0]

C₁₉H₂₇NO₇ 381.425

Alkaloid from *Senecio laricifolius* (Asteraceae). The trivial name 18-Hydroxysenkirkine results from use of a nonstandard numbering system.

19-Acetoxy: 18-Acetoxyenkirkine

[106283-54-9]

C₂₁H₂₉NO₈ 423.462

Alkaloid from *Senecio laricifolius* (Asteraceae). Trivial name results from a nonstandard numbering system.

12-Epimer, 12-Ac: Neogularidine

[90364-91-3]

C₂₁H₂₉NO₇ 407.463

Alkaloid from the aerial parts of *Ligularia dentata* (Asteraceae). Needles (hexane). Mp 117-119°. [α]_D²⁷ -58 (c, 0.13 in CHCl₃).

▶ VS3592000

(15E)-Isomer: Neosenkirkine

[57194-70-4]

C₁₉H₂₇NO₆ 365.425

Alkaloid from *Senecio auricola* (Asteraceae). Cryst. (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 175°. [α]_D²³ -21 (c, 1.08 in CHCl₃).

15E-Isomer, 18-hydroxy: Hydroxyneosenkirkine

C₁₉H₂₇NO₇ 381.425

Alkaloid from *Senecio anonyms*. Sol. MeOH, CHCl₃; poorly sol. H₂O.

15E-Isomer, 21-hydroxy: Anonamine

[111566-66-6]

C₁₉H₂₇NO₇ 381.425

Alkaloid from *Senecio anonyms* (Asteraceae). Mp 202°. [α]_D +33.5 (c, 1 in CHCl₃).

15E-Isomer, 21-acetoxy: Acetylanonamine

[138079-62-6]

C₂₁H₂₉NO₈ 423.462

Alkaloid from *Senecio anonyms*

(Asteraceae). Mp 124-125°. [α]_D +17.99 (c, 0.91 in CHCl₃).

15E-Isomer, 12-epimer, 12-Ac: Ligularidine

[60872-63-1]
C₂₁H₂₉NO₇ 407.463

Alkaloid from the aerial parts and roots of *Ligularia dentata* (Asteraceae). Needles (Et₂O). Mp 196° Mp 210°. [α]_D²⁸ -49.8 (c, 1.02 in EtOH).

▶ VS3591000

Stereoisomer: Isosenkirkine

C₁₉H₂₇NO₆ 365.425

Alkaloid from *Crotalaria walkeri* (Fabaceae). Mp 195-196°. [α]_D +32 (MeOH). No details in accessible lit. Of undetd. stereochem. For a further stereoisomer see Crotaverrine in S-245.

15E-Isomer, stereoisomer: Crotaverrine

[60827-69-2]

C₁₉H₂₇NO₆ 365.425

Alkaloid from *Crotalaria verrucosa* and *Crotalaria walkeri* (Fabaceae). Hygroscopic cryst. (EtOAc). Mp 142-144°. [α]_D²⁵ +32.7 (c, 1.04 in MeOH). Stereochem. incorrectly given in CAS (indexed as *O*-deacetyliligularidine). The stereochem. is poorly shown in the original paper which appears to show that the acid obt. from Crotaverrine is Usaramenic acid. This would make it identical with Neosenkirkine, but the props. are completely different.

15E-Isomer, 12-Ac, stereoisomer: O-Acetylcrotaverrine

C₂₁H₂₉NO₇ 407.463

Alkaloid from *Crotalaria verrucosa* (Fabaceae). Noncryst. [α]_D²⁵ +45.54 (c, 1.01 in MeOH).

[6882-01-5]

Chalmers, A.H. *et al.*, *J. Chromatogr.*, 1965, **20**, 270-277 (*tlc*)

Briggs, L.H. *et al.*, *J.C.S.*, 1965, 2492-2498 (*isol, bibl, pmr, Senkirkine, Acetylsenkirkine*)

Šimánek, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832-1837 (*uv*)

Hrbek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1972, **37**, 3918-3935 (*cd*)

Crout, D.H.G. *et al.*, *J.C.S. Perkin 1*, 1972, 1602-1607 (*isol, pmr, Hydroxysenkirkine*)

Atal, C.K. *et al.*, *Indian J. Pharm.*, 1973, **35**, 1-12 (*Isosenkirkine*)

Panizo, F.M. *et al.*, *An. Quim.*, 1974, **70**, 1043-1048 (*Neosenkirkine*)

Birnbaum, G.I. *et al.*, *J.A.C.S.*, 1974, **96**, 6165-6168 (*cryst struct*)

Abdullaev, U.A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 66-71; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 55-59 (*ms*)

Suri, O.P. *et al.*, *Phytochemistry*, 1976, **15**, 1061-1063 (*Crotaverrine, O-Acetylcrotaverrine*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173-1184 (*cmr*)

Molyneux, R.J. *et al.*, *Phytochemistry*, 1982, **21**, 439-443 (*cmr*)

IARC Monog., 1983, **31**, 231; *Suppl.* 7, 71 (*rev. tox*)

Asada, Y. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 475-482 (*Ligularidine, Neoligularidine*)

Mattocks, A.R. *et al.*, *Chemistry and Toxicology of Pyrrolizidine Alkaloids*, Academic Press, 1986, (*tox, rev*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151-1159 (*isol, 19-hydroxy, 19-acetoxy*)

Glinski, J.A. *et al.*, *Acta Cryst. C*, 1988, **44**, 1593-1598 (*cryst struct, Anonamine, Neosenkirkine, Hydroxysenkirkine*)

Zalkow, L.H. *et al.*, *J. Nat. Prod.*, 1988, **51**, 690-702 (*Anonamine, Hydroxysenkirkine*)

Asibal, C.F. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1425-1426 (*Acetylanonamine*)


Roeder, E. *et al.*, *Phytochemistry*, 1991, **30**, 1734-1737 (*Dihydroxysenkirkine*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DMX200

Sepiapterin, 8CI

S-246

2-Amino-7,8-dihydro-6-(2-hydroxy-1-oxo-propyl)-4(1H)-pteridinone, 9CI. 6-Lactoyl-7,8-dihydropterin



C₉H₁₁N₅O₃ 237.218

(S)-form

L-form

[17094-01-8]

Eye pigment from *Drosophila melanogaster* mutant *sepia*. Yellow powder or cryst.

Deoxy: **Deoxysepiapterin**. *Iosepiapterin* [1797-87-1]

C₉H₁₁N₅O₂ 221.218

Eye pigment from *Drosophila melanogaster*, more readily obt. from the blue-green alga *Anacystis nidulans*. Yellow cryst. (EtOH aq.). pK_{a1} 1.35; pK_{a2} 10.05 (cation).

7,8-Didehydro, 2'-ketone: Ramiopterin

[88848-62-8]

C₉H₇N₅O₃ 233.186

Present in *Drosophila* sp. Precursor of Sepiapterin, S-246 and Drosopterin, D-941.

[35101-75-8]

Forrest, H.S. *et al.*, *J.A.C.S.*, 1954, **76**, 5656; 5658 (*isol, uv*)

Forrest, H.S. *et al.*, *Arch. Biochem. Biophys.*, 1959, **83**, 508 (*isol*)

Nawa, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1960, **33**, 1555 (*struct, uv*)

Schircks, B. *et al.*, *Helv. Chim. Acta*, 1978, **61**, 2731 (*synth, uv, cd, bibl*)

Pfleiderer, W. *et al.*, *Chem. Ber.*, 1979, **112**, 2750 (*synth, abs config*)

Hearl, W.G. *et al.*, *C.A.*, 1984, **100**, 134386x (*Ramiopterin*)

Baur, R. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 531 (*synth, uv, pmr, deriv*)


Primus, J.P. *et al.*, *Insect Biochem.*, 1994, **24**, 907 (*biosynth*)

Septatisine

Septedinine

[156400-93-0]

S-247



C₂₂H₃₁NO₃ 357.492

Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Mp 129-130°. [α]_D¹⁹ +30.55 (c, 0.37 in CHCl₃).

16β,17-Dihydro, 15-ketone: Septedine

[176181-91-2]

C₂₂H₃₁NO₃ 357.492

Alkaloid from roots of *Aconitum septentrionale* (Ranunculaceae). Cryst. (Me₂CO). Mp 160-161°.


Joshi, B.S. *et al.*, *Can. J. Chem.*, 1994, **72**, 100 (*isol, pmr, cmr, ms, struct*)

Usmanova, S.K. *et al.*, *Khim. Prir. Soedin.*, 1995, **31**, 104; 1996, **32**, 77-81; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, **31**, 83; 1996, **32**, 62-65 (*Septedine, Septedinine*)

Septicine

S-248

6,7-Bis(3,4-dimethoxyphenyl)-1,2,3,5,8,8a-hexahydroindolizine, 9CI



C₂₄H₂₉NO₄ 395.497

Phenanthroindolizidine numbering shown.

(R)-form [42922-10-1]

Minor alkaloid from *Tylophora asthma-tica* (Asclepiadaceae). Needles (Et₂O). Mp 136°. [α]_D +38.8 (c, 1 in MeOH). λ _{max} 238 (sh); 287 (log ϵ 4.2) (EtOH).

7-Demethoxy: Secoantofine

[126262-27-9]

C₂₃H₂₇NO₃ 365.471

Alkaloid from *Cynanchum vincetoxicum*. Gum. [α]_D -81.9 (c, 0.4 in CHCl₃).

7-Demethoxy, N-oxide(β-): Di-O-methylphyllostemine β-N-oxide. Secoantofine β-N-oxide

[314744-21-3]

C₂₃H₂₇NO₄ 381.471

Alkaloid from *Cynanchum vincetoxicum*. Gum. [α]_D²¹ -83.4 (c, 0.15 in CHCl₃). λ _{max} 260 (ϵ 7100); 280 (ϵ 300) (MeOH).

7-Demethoxy, O³,O⁶-di-de-Me: Phyllostemine

[126262-22-4]

C₂₁H₂₃NO₃ 337.418

Alkaloid from the leaves, twigs, roots and fruit of *Cryptocarya phyllostemon* (Lauraceae). Pale yellow cryst. (MeOH/CHCl₃ or Py). Mp 205-207° (200-202°) dec. [α]_D -8 (c, 0.28 in EtOH).

7-Demethoxy, O⁶-de-Me: 6-O-Demethyl-secoantofine

C₂₂H₂₅NO₃ 351.444

Alkaloid from *Cynanchum vincetoxicum*. Gum. [α]_D²⁵ -92.6 (c, 0.2 in CHCl₃).

(S)-form [24316-18-5]

Alkaloid from *Tylophora crebriflora* and *Ficus septica* (Asclepiadaceae, Moraceae). Cryst. (MeOH). Mp 135-136°. $[\alpha]_D^{22}$ -16.2 (c, 1 in MeOH). λ_{max} 240 (sh) (log ϵ 4.2); 288 (log ϵ 4.03) (no solvent reported).

2-Demethoxy: Hispidine†

[82958-04-1]
 $C_{23}H_{27}NO_3$ 365.471
 Alkaloid from the leaves of *Ficus hispida* (Moraceae). Mp 124-125°.

(±)-form [26503-67-3]

Synthetic. Cryst. (MeOH or MeOH, aq.). Mp 135-136°.

(ξ)-form

13a-Hydroxy: 13a-Hydroxysepticine


[95066-47-0]
 $C_{24}H_{29}NO_5$ 411.497
 Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Cryst. (C_6H_6). Mp 290-292° dec. $[\alpha]_D^{25}$ +100 (c, 0.4 in MeOH).

- Russel, J.H. *et al.*, *Naturwissenschaften*, 1963, **50**, 443-444 (*occur, struct*)
 Russel, J.H. *et al.*, *Tet. Lett.*, 1969, 4035-4036 (*synth*)
 Rao, K.V. *et al.*, *J. Pharm. Sci.*, 1970, **59**, 1608-1611 (*uv, ms, pmr, struct*)
 Govindachari, T.R. *et al.*, *Tetrahedron*, 1970, **26**, 715-719; 1973, **29**, 891-897 (*isol, synth, uv, ir, pmr, ms*)
 Herbert, R.B. *et al.*, *Chem. Comm.*, 1976, 450-451 (*synth*)
 Stevens, R.V. *et al.*, *Tet. Lett.*, 1977, 979-982 (*synth*)
 Iwashita, T. *et al.*, *Chem. Lett.*, 1980, 383-386 (*synth*)
 Mangla, V.K. *et al.*, *Indian J. Chem., Sect. B*, 1980, **19**, 748-749 (*synth, uv, ir, pmr, ms*)
 Cragg, J.E. *et al.*, *J.C.S. Perkin I*, 1982, 2477-2485 (*synth*)
 Venkatachalam, S.R. *et al.*, *Naturwissenschaften*, 1982, **69**, 287-288 (*Hispidine*)
 Iida, H. *et al.*, *J.O.C.*, 1984, **49**, 2412-2418 (*synth, pmr, ms*)
 Bhutani, K.K. *et al.*, *Phytochemistry*, 1984, **23**, 1765-1769 (*13a-Hydroxysepticine*)
 Cavé, A. *et al.*, *Aust. J. Chem.*, 1989, **42**, 2243-2263 (*Phyllostemine*)
 Yerxa, B.R. *et al.*, *Tetrahedron*, 1994, **50**, 6173-6180 (*synth*)
 Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1996, **118**, 12082-12089 (*synth*)
 Comins, D.L. *et al.*, *J.O.C.*, 1997, **62**, 7435-7438 (*synth*)
 Staerk, D. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1584-1586; 2002, **65**, 1299-1302 (*Secoantofine, Secoantofine β-N-oxide*)

Septorine

S-249

[67332-36-9]



Absolute Configuration

$C_{16}H_{18}N_2O_4$ 302.329

Alkaloid from the fungus *Septoria nodorum*. Respiration inhibitor. Amorph. $[\alpha]_D^{20}$ +19 (MeOH). λ_{max} 230 (ϵ 6500); 300 (ϵ 9000); 370 (ϵ 4000) (MeOH) (Derep).

N-Methoxy: N-Methoxyseptorine

[85657-19-8]
 $C_{17}H_{20}N_2O_5$ 332.355
 Alkaloid from *Septoria nodorum*. Cryst. (hexane). Mp 167°. $[\alpha]_D$ +30 ($CHCl_3$). λ_{max} 230 (ϵ 6500); 300 (ϵ 9000); 370 (ϵ 4000) (MeOH) (Derep).

N-Methoxy, 1'-alcohol: N-Methoxyseptorinol


[146428-58-2]
 $C_{17}H_{22}N_2O_5$ 334.371
 Metab. of *Septoria nodorum*. Phytotoxin. Yellow amorph. powder. $[\alpha]_D$ +0.5 ($CHCl_3$). λ_{max} 223 (ϵ 7980); 282 (ϵ 2300); 360 (ϵ 4700) (MeOH) (Berdy).

- Devys, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1978, **286**, 457-458 (*ms, uv, pmr, struct*)
 Devys, M. *et al.*, *Tet. Lett.*, 1982, **23**, 5409-5412 (*uv, pmr, cmr, ms, struct*)
 Ohta, A. *et al.*, *Heterocycles*, 1991, **32**, 923-936 (*synth*)
 Devys, M. *et al.*, *Phytochemistry*, 1992, **31**, 4393-4394 (*N-Methoxyseptorinol*)
 Devys, M. *et al.*, *Org. Prep. Proced. Int.*, 1993, **25**, 696-698 (*synth*)
 Barbier, M. *et al.*, *Phytochemistry*, 1994, **35**, 955-957 (*abs config, N-Methoxyseptorinol*)

Sequoiamonascin D

S-250

[561305-73-5]




$C_{26}H_{33}NO_7$ 471.549

Azaphilone. Similar to Monascorubramine, M-686. Prod. by *Aspergillus parasiticus* isol. from *Sequoia sempervirens*. Red oil. $[\alpha]_D^{25}$ +1.7 (c, 0.09 in MeOH). λ_{max} 202 (log ϵ 3.94); 260 (log ϵ 3.74); 402 (log ϵ 3.5); 481 (log ϵ 3.54) (MeOH). Stierle, D.B. *et al.*, *J.O.C.*, 2003, **68**, 4966-4969 (*isol, pmr, cmr, ms*)

Seredine

S-251

Methyl 17-hydroxy-10,11-dimethoxy-yohimban-16-carboxylate, 9CI, 10,11-Dimethoxy-α-yohimbine [522-77-0]



Absolute Configuration

$C_{23}H_{30}N_2O_5$ 414.5

Alkaloid from roots of a variety of

Rauwolfia vomitoria. Also detected in *Rauwolfia serpentina* (Apocynaceae). Small prisms (Me_2CO). Mp 291° Mp 308° (sealed tube). $[\alpha]_D$ -1 (c, 0.43 in $CHCl_3$).

17-Epimer: 10,11-Dimethoxy-17-epi-α-yohimbine

[84667-06-1]
 $C_{23}H_{30}N_2O_5$ 414.5
 Alkaloid from leaves of *Neisosperma glomerata* (Apocynaceae). Cryst. (MeOH). Mp 259°. $[\alpha]_D^{20}$ -50 (c, 0.1 in EtOH).


- Poisson, J. *et al.*, *Bull. Soc. Chim. Fr.*, 1958, 1195-1200 (*isol, uv, ir, ord, struct*)
 Kaiser, F. *et al.*, *Chem. Ber.*, 1959, **92**, 278-287 (*occur*)
 Seguin, E. *et al.*, *J. Nat. Prod.*, 1982, **45**, 738-744 (*epimer*)

Serotobenine

S-252

Moschamindole

[99615-94-8]



$C_{20}H_{18}N_2O_4$ 350.373

Isol. from safflower meal (*Carthamus tinctorius*), *Centaurea moschata* and *Rhaphidophora decursiva*. Cryst. (MeOH). Mp 282-284.5° dec.

Ac:


Needles. Mp 211-214°. $[\alpha]_D^{25}$ 0 (c, 0.15 in Me_2CO).

- Sato, H. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 2969 (*isol, ir, pmr, cmr, ms, cryst struct*)
 Sarker, S.D. *et al.*, *Nat. Prod. Lett.*, 1997, **9**, 189-199 (*Moschamindole*)
 Zhang, H. *et al.*, *Pharm. Biol.*, 2002, **40**, 221-224 (*isol, pmr, cmr*)
 Koizumi, Y. *et al.*, *J.A.C.S.*, 2008, **130**, 16854-16855 (*synth*)

Serpentinine

S-253

[36519-42-3]



Absolute Configuration

$C_{42}H_{44}N_4O_5$ 684.833

Alkaloid from several *Rauwolfia* spp. (Apocynaceae). Mp 265-270° dec. pK_{a1} 6; pK_{a2} 10.6 (66% DMF aq.).

Hydrochloride (1:2): Mp 250°. $[\alpha]_D^{25}$ +189 (c, 0.45 in H₂O).

Dipicrate: Mp 271-273°.

Schlitter, E. et al., *Helv. Chim. Acta*, 1954, **37**, 1912 (uv, ir)

Kaneko, H. et al., *Yakugaku Zasshi*, 1960, **80**, 1357; *CA*, **55**, 6511


Gorman, A.A. et al., *Alkaloids* (London), 1971, **1**, 287 (uv, ms, pmr)

Irie, H. et al., *Chem. Comm.*, 1972, 871 (cryst struct)

Serratezomine A

S-254

[301679-55-0]



Absolute Configuration

C₁₆H₂₅NO₃ 279.378


Alkaloid from *Lycopodium serratum* var. *serratum*. Amorph. solid. $[\alpha]_D$ +13 (c, 0.5 in MeOH).

Morita, H. et al., *J.O.C.*, 2000, **65**, 6241-6245; 2002, **67**, 5378-5381 (isol, synth, abs config)

Serratine

S-255

[15252-93-4]



R = OH, R' = H

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium serratum* var. *serratum* form. *serratum* (*Lycopodium serratum* var. *thunbergii*). Pillars (Me₂CO or EtOAc). Mp 253°. $[\alpha]_D^{22}$ -15 (c, 1.02 in EtOH).

Di-Ac:

Cryst. (hexane). Mp 212-214°.

8α-Hydroxy: Serratimidine

[19637-58-2]

C₁₆H₂₅NO₄ 295.378

Alkaloid from *Lycopodium serratum* var. *serratum* form. *serratum*. Mp 210-211°. $[\alpha]_D^{12.5}$ -52 (c, 1.01 in EtOH).

Inubushi, Y. et al., *Chem. Pharm. Bull.*, 1967, **15**, 250-252; 1968, **16**, 2463-2470 (ir, pmr, struct)

Inubushi, Y. et al., *Chem. Pharm. Bull.*, 1968, **16**, 561-563 (*Serratimidine*)


Nishio, K. et al., *Tet. Lett.*, 1969, **10**, 861-864 (config)

Katakawa, K. et al., *J. Nat. Prod.*, 2007, **70**, 1024-1028 (isol, pmr, cmr, ms, cryst struct)

Serratimidine

S-256

[7689-04-5]

C₁₈H₂₈N₂O₂ 304.431

Alkaloid from *Lycopodium serratum* (Lycopodiaceae). Mp 232-234°. $[\alpha]_D^{10}$ +224.2 (c, 1.08 in EtOH).

O-Ac:Cryst. + 2H₂O. Mp 105-108°. $[\alpha]_D^{16}$ +172 (c, 1.19 in EtOH).*Deoxy: 8-Deoxyserratimidine*

[82841-99-4]

C₁₈H₂₈N₂O 288.432

Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Mp 198-199°.

Yasui, B. et al., *Tet. Lett.*, 1966, 3967 (ir, pmr, struct)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1970, **18**, 1880 (struct)

Inubushi, Y. et al., *Yakugaku Zasshi*, 1982, **102**, 434; *CA*, **97**, 107030q (8-Deoxyserratimidine)

Harayama, T. et al., *Heterocycles*, 1984, **22**, 1327 (config)

Serratine

S-257

[5545-99-3]

As Serratine, S-255 with

R = H, R' = OH

C₁₆H₂₅NO₃ 279.378

Alkaloid from *Lycopodium serratum* (Lycopodiaceae). Prisms (Me₂CO). Mp 244-245°. $[\alpha]_D^8$ -27.8 (c, 1.44 in EtOH).

*N-Oxide: Serratezomine B. Serratine**N-oxide*

[301679-57-2]

C₁₆H₂₅NO₄ 295.378

Alkaloid from *Lycopodium serratum* var. *serratum*. Amorph. solid. $[\alpha]_D$ +7 (c, 1.3 in MeOH).

Di-Ac: Mp 157-158°.*O⁸-(4-Methoxyphenylacetyl): Huperserratine*

[158182-11-7]

C₂₅H₃₃NO₅ 427.539

Alkaloid from the whole plant of *Huperzia serrata* (Lycopodiaceae). Mp 217-218°. $[\alpha]_D$ -3.5 (c, 0.2 in EtOH).

8-Deoxy: 8-Deoxyserratine

[18331-31-2]

C₁₆H₂₅NO₂ 263.379

Alkaloid from *Lycopodium serratum* (Lycopodiaceae). Fine columns (EtOAc or Me₂CO). Mp 223-224°. $[\alpha]_D^{17}$ -15.6 (c, 1.00 in EtOH).

8-Deoxy, 13-ketone: 8-Deoxy-13-dehydroserratine

[14478-54-7]

C₁₆H₂₃NO₂ 261.363

Alkaloid from *Lycopodium phlegmaria* (Lycopodiaceae). Fine needles (hexane). Mp 110-112°.

Inubushi, Y. et al., *Tet. Lett.*, 1966, **6**, 1537-1549 (ir, struct)

Inubushi, Y. et al., *Yakugaku Zasshi*, 1967, **87**, 1394-1404; *CA*, **68**, 47027h (*Serratine*, 8-Deoxyserratine, isol)

Inubushi, Y. et al., *Tetrahedron*, 1968, **24**, 3541-3556 (ms)

Ishii, H. et al., *Chem. Pharm. Bull.*, 1970, **18**, 1880-1885 (synth, 8-Deoxyserratine, 8-Deoxy-13-dehydroserratine)

Harayama, T. et al., *Chem. Pharm. Bull.*, 1975, **23**, 1511-1515; 1980, **28**, 2394-2402 (*Serratine*, 8-Deoxyserratine, synth)

Inubushi, Y. et al., *Yakugaku Zasshi*, 1982, **102**, 434-439; *CA*, **97**, 107030q (8-Deoxy-13-dehydroserratine)

Zhou, B.-N. et al., *Phytochemistry*, 1993, **34**, 1425-1428 (pmr, cmr)


Zhu, D.-Y. et al., *Phytochemistry*, 1994, **36**, 1069-1072 (*Huperserratine*)

Morita, H. et al., *J.O.C.*, 2000, **65**, 6241-6245 (*Serratezomine B*)

Serratiochelin

S-258

[155070-21-6]

C₂₁H₂₃N₃O₇ 429.429


Isol. from *Serratia marcescens*. Siderophore. λ_{max} 254 (ε 2400); 316 (ε 20000) (MeOH) (Berdy).

Ehlert, G. et al., *Z. Naturforsch., C*, 1994, **49**, 11 (isol, synth, pmr, cmr)

Serratoline

S-259

[74260-86-9]



Absolute Configuration

C₂₀H₂₆N₂O 310.438

Minor alkaloid from *Aristotelia serrata* (Elaeocarpaceae). Rhombs (MeOH). Mp 157-160° Mp 167-170° Mp 178° dec. (synthetic). $[\alpha]_D^{19}$ -68.25 (CHCl₃).

13-Oxo: Triabunnine

[90930-66-8]

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Aristotelia peduncularis* (Elaeocarpaceae). Amorph. λ_{max} 219 (sh) (log ε 4.31); 223 (log ε 4.35); 266 (log ε 3.74) (EtOH).

15α-Hydroxy: Aristotelinine

[66833-16-7]

C₂₀H₂₆N₂O₂ 326.438


Trace alkaloid from *Aristotelia chilensis* (Elaeocarpaceae). Mp 246-250°. λ_{max} 220 (ε 18000); 263 (ε 6000)

(MeOH).

- Gopalakrishna, E.M. *et al.*, *Acta Cryst. B*, 1978, **34**, 3778-3780 (*cryst struct*, *Aristolelinine*)
 Bittner, M. *et al.*, *Chem. Comm.*, 1978, 79-80 (*Aristolelinine*, *uv, struct*)
 Bick, I.R.C. *et al.*, *Tet. Lett.*, 1980, **21**, 545-546 (*uv, pmr*)
 Bick, I.R.C. *et al.*, *Heterocycles*, 1983, **20**, 667-669 (*pmr, cmr, struct*)
 Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1984, **67**, 804-814 (*Triabunnine*)
 Stoermer, D. *et al.*, *J.O.C.*, 1993, **58**, 564-568 (*synth*)

Sesbanimide A
Sesbanimide

S-260

C₁₅H₂₁NO₇ 327.333**(+)-form** [85719-78-4]

Alkaloid from the seeds of *Sesbania drummondii* and *Sesbania punicea* (Fabaceae). Prod. by a marine *Agrobacterium* strain PH-103. Shows notable cytotoxicity against KB cells *in vitro* and potent inhibitory activity against P338 murine leukaemia *in vivo*. Cryst. (Et₂O/CH₂Cl₂ or MeOH/CH₂Cl₂). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 158-159° (155-156°). [α]_D²⁰ +54.7 (c, 0.17 in CHCl₃). [α]_D²⁰ -3.8 (c, 0.28 in MeOH) (-5.6).

(-)-form

Synthetic. Solid (CH₂Cl₂/Et₂O). Mp 154.5-155.5°. [α]_D +6 (c, 0.27 in MeOH). [α]_D -56.9 (c, 0.21 in CHCl₃).

11-Epimer: Sesbanimide BC₁₅H₂₁NO₇ 327.333

Alkaloid from seeds of *Sesbania drummondii* (Fabaceae). Glass. Sol. MeOH, CHCl₃. [α]_D²⁰ -22.4 (c, 0.17 in CHCl₃) (synthetic). Isol. as a mixt. of 2 components, B1 and B2, which are C-10 epimers (assignments of configs. not yet made).

[92282-10-5, 95839-03-5]


- Powell, R.G. *et al.*, *J.A.C.S.*, 1983, **105**, 3739-3741 (*isol, uv, ir, pmr, ms, cryst struct*)
 Gorst-Allman, C.P. *et al.*, *J.C.S. Perkin 1*, 1984, 1311-1314 (*isol, ir, pmr, cmr*)
 Powell, R.G. *et al.*, *Phytochemistry*, 1984, **23**, 2789-2796 (*Sesbanimide B, isol, uv, ir, pmr, cmr, ms, struct*)
U.S. Pat., 1985, 4 532 327; *CA*, **103**, 166147 (*activity*)
 Schlessinger, R.H. *et al.*, *J.O.C.*, 1986, **51**, 2621-2623 (*synth*)
 Wanner, M.J. *et al.*, *Tetrahedron*, 1987, **43**, 2549-2556 (*synth, ir, pmr*)
 Matsuda, F. *et al.*, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 2123 (*synth, props*)
 Tomioka, K. *et al.*, *Tet. Lett.*, 1988, **29**, 3095-3096 (*synth*)
 Matsuda, F. *et al.*, *Tetrahedron*, 1988, **44**, 4721-4736 (*synth, ir, pmr, abs config*)

- Vloon, W.J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1991, **110**, 414-419 (*synth*)
 Grieco, P.A. *et al.*, *Chem. Comm.*, 1992, 368-370 (*synth*)
 Cirillo, P.F. *et al.*, *J.O.C.*, 1994, **59**, 3055-3063 (*synth*)
 Honda, T. *et al.*, *Tetrahedron: Asymmetry*, 1994, **5**, 247-254 (*synth*)
 Acebal, C. *et al.*, *J. Antibiot.*, 1998, **51**, 64-67 (*marine isol, pmr, cmr*)

Sesbanimide C

S-261

[95599-43-2]


C₁₅H₂₃NO₆ 313.35

Alkaloid from seeds of *Sesbania drummondii* (Fabaceae). Prod. by a marine *Agrobacterium* strain PH-A034C. Powder. Sol. MeOH. [α]_D²⁵ +9.52 (c, 0.21 in CHCl₃).

- Powell, R.G. *et al.*, *Phytochemistry*, 1984, **23**, 2789-2796 (*isol, pmr, ms, struct*)
 Acebal, C. *et al.*, *J. Antibiot.*, 1998, **51**, 64-67 (*isol, pmr, cmr*)

Sesbanine

S-262



Absolute configuration

C₁₂H₁₂N₂O₃ 232.238

Shows weak antineoplastic, cytotoxic activity. Log P -1.77 (uncertain value) (calc).

(+)-form [70521-94-7]

Alkaloid from the seeds of *Sesbania drummondii* (Fabaceae). Cryst. (MeOH). Mp 240-243°. [α]_D²³ +14.6 (c, 0.56 in MeOH) (natural). [α]_D +43.1 (c, 0.52 in MeOH) (synthetic). Originally reported to exhibit potent antineoplastic activity. However, subsequent studies revealed that Sesbanine is only marginally cytotoxic and that the major antineoplastic agent isolated from the plant is Sesbanimide A, S-260.


(±)-form

Synthetic. Cryst. (MeOH/EtOAc). Mp 241-243° (238-240°).
 10-Epimer: Synthetic. Mp 240-242°.

- Powell, R.G. *et al.*, *J.A.C.S.*, 1979, **101**, 2784 (*pmr, cmr, ir, uv, cryst struct*)
 Bottaro, J.C. *et al.*, *J.O.C.*, 1980, **45**, 1176 (*synth*)
 Kende, A.S. *et al.*, *Tet. Lett.*, 1980, **21**, 715 (*synth, ir, pmr, cmr, ms*)
 Wanner, M.J. *et al.*, *Tetrahedron*, 1982, **38**, 2741 (*synth, ir, pmr*)
 Iwao, M. *et al.*, *Tet. Lett.*, 1983, **24**, 2649 (*synth, ir, pmr*)
 Wada, M. *et al.*, *Tet. Lett.*, 1985, **26**, 3267 (*synth, ir, pmr*)
 Tomioka, K. *et al.*, *Tetrahedron*, 1988, **44**, 4351 (*synth, ir, pmr, cmr, abs config*)

Sespendole

S-263



Absolute Configuration

C₃₃H₄₅NO₄ 519.723


Prod. by *Pseudobotrytis terrestris* FKA-25. Inhibitor of lipid droplet synthesis in macrophages. Amorph. solid. [α]_D²³ -18 (c, 0.1 in MeOH). λ_{max} 239 (ε 37900); 288 (ε 8800) (MeOH).

- Uchida, R. *et al.*, *J. Antibiot.*, 2006, **59**, 93-97; 298-302; 338-344 (*isol, pmr, cmr, biosynth, activity*)

Sessiflorine

S-264

2,3-Dihydro-5-methoxy-3-[4-(methylamino)phenyl]pyrrolo[2,1-b]quinazolin-9(1H)-one, 9CI
 [35226-54-1]


C₁₉H₁₉N₃O₂ 321.378

Minor alkaloid from the leaves and branches of *Anisotes sessiliflorus* (Acanthaceae). Cryst. (MeOH). Mp 195-197°. Opt. inactive.

- Arndt, R.R. *et al.*, *Tetrahedron*, 1967, **23**, 3521 (*isol, uv, ir, pmr, ms*)
 Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4387 (*struct, synth, ir, pmr, ms*)

Sessilifoliamide B

S-265

C₁₇H₂₇NO₃ 293.405Alkaloid from the roots of *Stemona*

sessilifolia. Oil. $[\alpha]_D^{24}$ -43 (c, 0.1 in CHCl_3).


12,13-Didehydro: Sessilifoliamide C

$\text{C}_{17}\text{H}_{25}\text{NO}_3$ 291.389

Alkaloid from the roots of *Stemona sessilifolia*. Oil. $[\alpha]_D^{26}$ -140 (c, 0.17 in CHCl_3).

Kakuta, D. *et al.*, *Tetrahedron*, 2003, **59**, 7779-7786 (*isol, pmr, cmr*)

Sessilifoliamide D S-266




$\text{C}_{18}\text{H}_{29}\text{NO}_4$ 323.431

Alkaloid from the roots of *Stemona sessilifolia*. Oil. $[\alpha]_D^{26}$ -94 (c, 0.16 in CHCl_3).

Kakuta, D. *et al.*, *Tetrahedron*, 2003, **59**, 7779-7786 (*isol, pmr, cmr*)

Sessilifoliamide E S-267

[922500-42-3]



Relative Configuration

$\text{C}_{22}\text{H}_{31}\text{NO}_6$ 405.49

Alkaloid from the roots of *Stemona sessilifolia*. Prisms. Mp 197-200°. $[\alpha]_D$ -18 (c, 0.1 in CHCl_3).

1-Hydroxy: Sessilifoliamide F

[922500-43-4]


$\text{C}_{22}\text{H}_{31}\text{NO}_7$ 421.489

Alkaloid from the roots of *Stemona sessilifolia*. Prisms. Mp 200-202°. $[\alpha]_D$ -42 (c, 0.8 in CHCl_3).

Hitotsuyanagi, Y. *et al.*, *Tetrahedron*, 2007, **63**, 1008-1013 (*isol, pmr, cmr, cryst struct*)

Sessilifoliamide G S-268

[922500-44-5]




$\text{C}_{22}\text{H}_{31}\text{NO}_7$ 421.489

Alkaloid from the roots of *Stemona sessilifolia*. Prisms. Mp 213-215°. $[\alpha]_D$ -85 (c, 0.06 in MeOH).

Hitotsuyanagi, Y. *et al.*, *Tetrahedron*, 2007, **63**, 1008-1013 (*isol, pmr, cmr, cryst struct*)

Sessiline S-269

5-[(5-Oxo-2-pyrrolidinyl)oxy]methyl]-2-furancarboxaldehyde




$\text{C}_{10}\text{H}_{11}\text{NO}_4$ 209.201

Alkaloid from the fruit of *Acanthopanax sessiliflorus*. Mp 171-172°. Racemate. λ_{max} 223 (log ϵ 3.58); 278 (log ϵ 4.17) (MeOH).

Lee, S. *et al.*, *Planta Med.*, 2002, **68**, 936-939 (*isol, pmr, cmr, ms*)

Sessilistemonamine A S-270



Relative Configuration

$\text{C}_{22}\text{H}_{33}\text{NO}_5$ 391.506

Alkaloid from the roots of *Stemona sessilifolia*. Prisms (EtOAc/petrol). Mp 193-194°. $[\alpha]_D^{20}$ +117 (c, 0.07 in MeOH).

10-Epimer: Sessilistemonamine C

$\text{C}_{22}\text{H}_{33}\text{NO}_5$ 391.506

Alkaloid from the roots of *Stemona sessilifolia*. Prisms (EtOAc/petrol). Mp 221-222°. $[\alpha]_D^{20}$ -157 (c, 0.05 in MeOH).

9a,10,11,12-Tetraepimer: Sessilistemonamine B


$\text{C}_{22}\text{H}_{33}\text{NO}_5$ 391.506

Alkaloid from the roots of *Stemona sessilifolia*. Plates (EtOAc/petrol). Mp 252-254°. $[\alpha]_D^{20}$ -29 (c, 0.05 in MeOH).

Wang, P. *et al.*, *Chem. Biodiversity*, 2007, **4**, 523-530 (*isol, pmr, cmr, ms*)

Sessilistemonamine D S-271

[941271-34-7]



$\text{C}_{22}\text{H}_{33}\text{NO}_4$ 375.507

Alkaloid from the roots of *Stemona sessilifolia*. Prisms. Dec. at 208°.

Wang, P. *et al.*, *Chin. Chem. Lett.*, 2007, **18**, 152-154 (*isol, pmr, cmr*)

Sessilistemonine S-272

$\text{C}_{25}\text{H}_{35}\text{NO}_7$ 461.554

Struct. unknown. Alkaloid from *Stemona sessilifolia* (Stemonaceae). Mp 172°. $[\alpha]_D$ +112.5 (c, 1 in MeOH).

Perchlorate:

Needles. Mp 201-205°.

Picrate:

Yellow needles. Mp 163-165°.

Picrolonate:


Yellow needles. Mp 236°.

Methiodide:

Needles. Mp 231-232°.

Chu, J.-H. *et al.*, *Chem. Zentralbl.*, 1960, **131**, 1858 (*isol*)

Setoclavine S-273



(+)-form

$\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}$ 254.331

(+)-form [519-12-0]

Alkaloid from ergot found on *Pennisetum typhoideum* (pearl millet) and *Elymus mollis*. Also occurs in *Argyrea cuneata*, *Argyrea nervosa* and *Rivea corymbosa* (Convolvulaceae). Prisms (MeOH or Me_2CO). Mp 229-234° dec. $[\alpha]_D^{20}$ +174 (c, 1.1 in Py). $[\alpha]_D^{20}$ +165 (c, 0.3 in EtOH). λ_{max} 242 (ϵ 26300); 317 (ϵ 12600) (EtOH).

Hydrochloride:

Needles (H_2O or EtOH). Mp 300° (darkens above 200°).

Nitrate:

Needles (H_2O). Mp 300° (darkens above 125°). $[\alpha]_D^{20}$ +190 (c, 0.5 in 50% EtOH aq.).

N-De-Me: 6-Norsetoclavine

[19641-11-3]

$\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}$ 240.304

Isol. from the culture filtrates of *Pennisetum ergot* strains 47A and 231. Mp 163-165°. Prob. a secondary prod. arising in the fungus by demethylation of Setoclavine. λ_{max} 225 (log ϵ 4.28); 241 (log ϵ 4.23); 314 (log ϵ 3.9) (EtOH).

18-Hydroxy: Penniclavine

[519-13-1]

$\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}_2$ 270.33

Alkaloid from ergot (and several other *Claviceps* spp.), *Ipomoea muelleri*, *Argyrea* spp. and several other spp. (Convolvulaceae). Mp 222-225° dec. $[\alpha]_D^{20}$ +151 (c, 0.5 in Py). λ_{max} 242 (ϵ 26300); 317 (ϵ 12600) (EtOH) (Derep).

8-Epimer: Isosetoclavine

[519-11-9]

$\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}$ 254.331

Alkaloid from *Claviceps purpurea*, *Argyrea nervosa*, *Argyrea cuneata*, *Rivea corymbosa* and *Ipomoea muelleri* (Convolvulaceae). Cryst. (MeOH). Mp 234-237° dec. $[\alpha]_D^{20}$ +107 (c, 0.5 in Py). $[\alpha]_D^{20}$ +129 (c, 0.4 in EtOH).

8-Epimer, picrate:

Orange-yellow needles (MeOH). Mp 150-155°.

8-Epimer, 9,10 α -dihydro: **Dihydroisose-to-clavine**

[52146-68-6]
 $C_{16}H_{20}N_2O$ 256.347
 Stereochem. revised in 2006 by Moldvai *et al.* Formerly known as Dihydrosetoclavine. Prod. by *Claviceps paspali*. Cryst. (MeOH or Me₂CO). Mp 276-278° dec. $[\alpha]_D^{20}$ -51 (c, 0.2 in Py). λ_{max} 223 (log ϵ 4.52); 281 (log ϵ 3.83); 292 (log ϵ 3.74) (MeOH).

8-Epimer, 18-hydroxy: **Isopenniclavine**
 [478-92-2]
 $C_{16}H_{18}N_2O_2$ 270.33
 Alkaloid from ergot *Claviceps purpurea* and from the seeds of *Ipomoea muelleri* and *Stictocardia campanulata* (Convolvulaceae). Cryst. (H₂O). Mp 163-165° dec. $[\alpha]_D^{20}$ +146 (c, 0.7 in Py). $[\alpha]_D^{20}$ +140 (c, 0.9 in EtOH). λ_{max} 242 (ϵ 26300); 317 (ϵ 12600) (EtOH) (Derep).

(±)-**form** [86194-81-2]
 Synthetic. Mp 206°.


9,10 α -Dihydro:
 Synthetic. Mp 252-256°.

8-Epimer: [34964-90-4]
 Synthetic. Mp 203-205°.

Stoll, A. *et al.*, *Helv. Chim. Acta*, 1954, **37**, 1815 (*Penniclavine, isol, uv*)
 Hofmann, A. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 1358-1373 (*Isosetoclavine, Penniclavine, Isopenniclavine, isol, uv, ir, struct*)
 Ramstad, E. *et al.*, *J. Nat. Prod.*, 1967, **30**, 441-444 (*6-Norsetoclavine*)
 Mrtek, R.G. *et al.*, *Phytochemistry*, 1968, **7**, 1535-1541 (*pmr*)
 Kornfeld, E.C. *et al.*, *Chem. Ind. (London)*, 1971, 1233-1234 (*Isosetoclavine, synth*)
 Chao, J.M. *et al.*, *J. Pharm. Sci.*, 1973, **62**, 588-591 (*Isosetoclavine, Penniclavine, occur*)
 Chao, J.-M. *et al.*, *Phytochemistry*, 1973, **12**, 2435-2440 (*Isosetoclavine, occur*)
 Tschertner, H. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 113-121 (*Dihydrosetoclavine*)
 Der Marderosian, A. *et al.*, *Planta Med.*, 1974, **25**, 6-16 (*Penniclavine, Isopenniclavine, occur*)
 Bach, N.J. *et al.*, *Tet. Lett.*, 1974, 3225-3228 (*Penniclavine, synth*)
 Bernardi, L. *et al.*, *Tetrahedron*, 1974, **30**, 3447-3450 (*Isosetoclavine, synth*)
 Wurst, M. *et al.*, *J. Chromatogr.*, 1978, **150**, 477-483 (*Isosetoclavine, hplc*)
 Lee, T.M. *et al.*, *Planta Med.*, 1979, **35**, 247-252 (*Penniclavine, occur*)
 Natsume, M. *et al.*, *Heterocycles*, 1981, **16**, 375-379 (*Dihydrosetoclavine*)
 Rebek, J. *et al.*, *J.A.C.S.*, 1984, **106**, 1813-1819 (*synth, ir, pmr, ms*)
 Liras, S. *et al.*, *J.A.C.S.*, 2001, **123**, 5918-5924 (*synth*)
 Moldvai, I. *et al.*, *Heterocycles*, 2006, **67**, 291-298 (*synth, pmr, cmr*)

Sevanine S-274

1-(1,3-Benzodioxol-5-ylmethyl)-6-methoxy-7-isoquinolinol, 9CI. 7-Hydroxy-6-methoxy-1-(3,4-methylenedioxybenzyl)isoquinoline
 [54293-58-2]



$C_{18}H_{15}NO_4$ 309.321
 Alkaloid from *Papaver macrostomum* (Papaveraceae). Mp 218°.


O- β -D-Glucopyranoside: **Glycomarine**.
Glicomarine
 [77396-64-6]
 $C_{24}H_{25}NO_9$ 471.463
 Alkaloid from *Papaver arenarium* (Papaveraceae). Mp 205-206°. $[\alpha]_D$ -51 (CHCl₃/MeOH).

Me ether: Mp 106-108°.

α -Methoxy, Me ether: **Setigeridine**
 [180386-77-0]
 $C_{20}H_{19}NO_5$ 353.374
 Minor alkaloid from *Papaver setigerum*.
 Mnatsakanyan, V.A. *et al.*, *Tet. Lett.*, 1974, 851 (*ms, uv, pmr, struct*)
 Simanek, V. *et al.*, *Heterocycles*, 1976, **4**, 1263 (*synth*)
 Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1980, **16**, 852 (*Glycomarine*)
 Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1996, **61**, 1047-1052 (*Setigeridine*)

Sevcoridine S-275

21-(1,5-Dimethyl-2-piperidinyl)pregn-20-ene-3,6-diol, 9CI
 [60269-68-3]



$C_{28}H_{47}NO_2$ 429.685
 Revised structs. Originally assigned with azepine side-chains. Alkaloid from the aerial parts of *Korolkowia sewerzowii*. Cryst. (Me₂CO). Mp 241-243°. $[\alpha]_D$ -43.2 (c, 0.11 in EtOH).

3-O- β -D-Glucopyranoside: **Sevcorine**
 [19625-06-0]
 $C_{34}H_{57}NO_7$ 591.827
 Alkaloid from *Korolkowia sewerzowii*. Mp 236-238°. $[\alpha]_D$ -41.1 (c, 1.01 in MeOH).

6-Deoxy, 5,6-didehydro: **Sevcoridine**
 $C_{28}H_{45}NO$ 411.67
 Aglycone obt. by hydrol. of Sevcorine. Cryst. (Me₂CO). Mp 179-180°. $[\alpha]_D$ -78 (c, 0.846 in MeOH).

22,25-Diepimer: **Edpetilidine**
 [27509-73-5]
 $C_{28}H_{47}NO_2$ 429.685
 Alkaloid from the epigeal parts of *Petilium eduardii*. Cryst. (MeOH). Mp 269-271°. $[\alpha]_D$ +42.48 (c, 0.306 in EtOH).

22,25-Diepimer, hydrochloride: Mp 283° dec.


22,25-Diepimer, 3-O- β -D-xylopyranoside: **Edpetilinine**
 [28440-32-6]
 $C_{33}H_{55}NO_6$ 561.801

Alkaloid from aerial parts of *Petilium eduardii* (Liliaceae). Mp 262-264°. $[\alpha]_D^{26}$ -10.86 (Py).

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 384-392; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 302-307 (*Edpetilidine, isol*)
 Nuriddinov, R.N. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 605-606; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 527 (*Edpetilinine*)
 Samikov, S.M. *et al.*, *Khim. Prir. Soedin.*, 1976, **13**, 269-270; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **13**, 244-245 (*Sevcorine*)
 Abdullaeva, D.U. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 817; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 702-703 (*Sevcoridine, Sevcorine*)
 Nasirov, S.M. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 864-869; *Chem. Nat. Compd. (Engl. Transl.)*, 1988, **23**, 721-723 (*Edpetilidine, Sevcoridine, cryst struct*)

Severibuxine S-276

6-Hydroxy-3,3-bis(3,7-dimethyl-2,6-octadienyl)-2,4(1H,3H)-quinolinedione
 [219998-24-0]




$C_{29}H_{39}NO_3$ 449.632
 Alkaloid from the root bark of *Severinia buxifolia*. Yellowish needles (hexane). Mp 113-115°. λ_{max} 209 ; 242 ; 265 ; 382 (MeOH).

Wu, T.-S. *et al.*, *Phytochemistry*, 1998, **49**, 1467-1470 (*isol, uv, ir, pmr, cmr*)

Severidine S-277

1,3,13-Trihydroxy-11,12-secocevane-6,12-dione, 9CI. **Severidine**
 [80248-74-4]



$C_{27}H_{43}NO_5$ 461.64
 A C-seco-cevane alkaloid. Alkaloid from *Korolkowia sewerzowii* (*Fritillaria sewerzowii*) (Liliaceae). Cryst. (Me₂CO). Mp 200-202°. $[\alpha]_D$ +97.7 (c, 1.504 in EtOH).

3-Ac: **Severidinine**. **Seweridinine**
 [88755-00-4]
 [88823-94-3]
 $C_{29}H_{45}NO_6$ 503.678


Alkaloid from *Korolkowia sewerzovi*. Cryst. (Me₂CO/petrol). Mp 145-147°. [α]_D -50.2 (c, 0.78 in CHCl₃).

Nasirov, S.M. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 342-349; *Chem. Nat. Compd. (Engl. Transl.)*, 1981, 265-272 (isol, cryst struct, abs config)

Abdullaeva, D.U. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 488-490; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 459-461 (Severidinine)

Severifoline**S-278**

3,12-Dihydro-6-hydroxy-3,3-dimethyl-5-(3-methyl-2-butenyl)-7H-pyrano[2,3-c]acridin-7-one, 9CI
[84306-81-0]



C₂₃H₂₃NO₃ 361.44

Alkaloid of *Severinia buxifolia* (Rutaceae). Yellow needles (Me₂CO). Mp 253-254°. λ_{\max} 224 ; 240 ; 253 ; 278 ; 296 ; 338 ; 410 (MeOH).

N-Me: N-Methylseverifoline

[84306-82-1]

C₂₄H₂₅NO₃ 375.466

Alkaloid from *Severinia buxifolia* (Rutaceae). Orange-yellow cryst. (Me₂CO). Mp 152-154°. λ_{\max} 228 ; 254 ; 288 ; 295 ; 343 ; 417 (MeOH).

11-Hydroxy: Atalaphyllinine

[56296-87-8]

C₂₃H₂₃NO₄ 377.439

Alkaloid from *Severinia buxifolia* and *Atalantia monophylla* (Rutaceae). Yellow needles (C₆H₆/EtOAc). Mp 205-207°. λ_{\max} 264 (log ϵ 3.85); 293 (log ϵ 4.05); 305 (log ϵ 4.12); 347 (log ϵ 4.34) (EtOH).

11-Hydroxy, N-Me: N-Methylatalaphyllinine

[49620-08-8]

C₂₄H₂₅NO₄ 391.466

Alkaloid from *Severinia buxifolia*, *Atalantia monophylla* and *Atalantia ceylanica* (Rutaceae). Red needles (Et₂O/petrol). Mp 192-193°.

11-Methoxy, N-Me:

Yellow needles (Et₂O/petrol). Mp 126.5°.

Fraser, A.W. *et al.*, *J.C.S. Perkin I*, 1973, 1173-1175 (isol, uv, ir, nmr, ms, N-Methylatalaphyllinine)

Basa, S.C. *et al.*, *Phytochemistry*, 1975, **14**, 835-836 (Atalaphyllinine)


Banerji, J. *et al.*, *Indian J. Chem., Sect. B*, 1981, **20**, 835-838 (cmr, N-Methylatalaphyllinine)

Wu, T.-S. *et al.*, *Phytochemistry*, 1982, **21**, 1771-1773 (*Severinia buxifolia* constits)

Chukaew, A. *et al.*, *Acta Cryst. E*, 2008, **63**, o3723-o3724 (N-Methylatalaphyllinine, cryst struct)

Severzinine

Sewercinine. Sewerzinine
[68736-95-8]



Absolute Configuration

C₂₀H₁₉NO₆ 369.373


Diastereoisomer of Ochrobirine, O-20.

CAS name is defective (spiro posn. numbered 7' instead of 5). Alkaloid from *Corydalis severtzovii* (Papaveraceae). Cryst. (EtOH). Mp 90-91° dec. [α]_D +109 (c, 1.2 in CHCl₃).

Irgashev, T. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 536; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 464 (isol, uv, ir, pmr, ms, struct)

Shearinine I**S-280**

[925427-39-0]



C₃₇H₄₅NO₇ 615.765

Prod. by a mangrove-derived *Penicillium* sp. HKI0459. Amorph. powder. [α]_D +6.5 (c, 0.3 in CHCl₃). λ_{\max} 245 (MeOH).

$\Delta^{23,28}$ -Isomer: Shearinine H

[925427-38-9]

C₃₇H₄₅NO₇ 615.765


Prod. by *Penicillium* sp. HKI0459.

Amorph. powder. [α]_D +20.2 (c, 0.3 in CHCl₃). λ_{\max} 246 ; 308 (MeOH).

Xu, M. *et al.*, *Tetrahedron*, 2007, **63**, 435-444 (isol, cd, pmr, cmr, ms)

Shearinine A**S-281**

[163136-24-1]



Absolute Configuration

C₃₇H₄₅NO₅ 583.766

Shearinines have been isol. by two groups of workers simultaneously, and various Shearinine code letters have been given to different but closely similar compds.

Prod. by *Eupenicillium shearii*, a marine-derived *Penicillium janthinellum* and *Penicillium* sp. HKI0459. Apoptosis inducer. Exhibits antiinsectan activity. Mp 250° dec. [α]_D +16 (c, 0.20 in CHCl₃).

22 α -Hydroxy (1): Shearinine D \dagger

[925427-34-5]

C₃₇H₄₅NO₆ 599.766

S-279

Prod. by *Penicillium* sp. HKI0459 and marine-derived *Penicillium janthinellum*. Amorph. powder. [α]_D +67.7 (c, 0.3 in CHCl₃). λ_{\max} 210 ; 236 ; 260 ; 330 (MeOH).

22 α -Hydroxy (2): Shearinine E \dagger

C₃₇H₄₅NO₆ 599.766

Prod. by a marine-derived *Penicillium janthinellum*. Cryst. (hexane/EtOAc). Mp > 300° dec. [α]_D +3.5 (c, 0.17 in CHCl₃). C-22:C-23 relative config. only. May be identical with Shearinine D above. λ_{\max} 231 (log ϵ 4.2); 256 (sh) (log ϵ 4.36); 264 (log ϵ 4.38); 286 (log ϵ 4.03); 330 (log ϵ 4.04) (CHCl₃).

22 β -Hydroxy: Shearinine D \ddagger

C₃₇H₄₅NO₆ 599.766

Prod. by a marine-derived *Penicillium janthinellum*. Cryst. (hexane/EtOAc). Mp 180-185° dec. [α]_D +21.8 (c, 0.13 in CHCl₃). C-22/C-23 relative config. only determined. λ_{\max} 233 (log ϵ 4.13); 256 (sh) (log ϵ 4.36); 263 (log ϵ 4.34); 330 (log ϵ 4.08) (CHCl₃).

22 α -Methoxy: Shearinine E \ddagger

[925427-35-6]

C₃₈H₄₇NO₆ 613.792

Prod. by *Penicillium* sp. HKI0459. Amorph. powder. [α]_D +8.9 (c, 0.2 in CHCl₃). λ_{\max} 204 ; 234 ; 264 ; 332 (MeOH).

$\Delta^{23,28}$ -Isomer: Shearinine F \ddagger

[925427-36-7]

C₃₇H₄₅NO₅ 583.766

Prod. by *Penicillium* sp. HKI0459 isol. from a mangrove. Amorph. powder. [α]_D +26.4 (c, 0.3 in CHCl₃). λ_{\max} 204 ; 240 ; 306 (MeOH).

$\Delta^{23,28}$ -Isomer, 22-oxo: Shearinine G

[925427-37-8]

C₃₇H₄₃NO₆ 597.75

Prod. by *Penicillium* sp. HKI0459. Amorph. powder. [α]_D +35.7 (c, 0.1 in CHCl₃). λ_{\max} 280 (MeOH).


Belofsky, G.N. *et al.*, *Tetrahedron*, 1995, **51**, 3959-3968 (Shearinine A)

Smetanina, O.F. *et al.*, *J. Nat. Prod.*, 2007, **70**, 906-909; 2054 (Shearinines D,E, isol, pmr, cmr)

Xu, M. *et al.*, *Tetrahedron*, 2007, **63**, 435-444 (Shearinines D-G)

Shearinine C**S-282**

[163136-26-3]



C₃₇H₄₇NO₇ 617.781

Metab. from the ascostromata of *Eupenicillium shearii*. Exhibits antiinsectan activity. Pale yellow solid. Mp 180-190° dec. [α]_D -146 (c, 0.20 in CHCl₃).

1'-Deoxy, 1',2'-didehydro, 3 β -alcohol:

Antibiotic PF 1101B. PF 1101B

[157408-11-2]

C₃₇H₄₇NO₆ 601.781

Prod. by *Penicillium* sp. Anthelminthic,

shows insecticidal props. Cryst.


Japan. Pat., 1994, 94 65 246; CA, 121, 155915t (PF 1101B)

Belofsky, G.N. et al., *Tetrahedron*, 1995, 51, 3959 (*Shearinine C*)

Shearinine J

S-283

[925427-40-3]



C₃₇H₄₇NO₆ 601.781

Prod. by a mangrove-derived *Penicillium* sp. HKI0459. Amorph. powder. [α]_D +48.5 (c, 0.3 in CHCl₃). λ_{max} 206 ; 232 (MeOH).


Xu, M. et al., *Tetrahedron*, 2007, 63, 435-444 (isol, pmr, cmr, ms)

Shearinine K

S-284

Shearinine F†

[925427-41-4]



C₃₇H₄₇NO₄ 569.783

See note under Shearinine A, S-281.

Prod. by a mangrove-derived *Penicillium* sp. HKI0459 and *Penicillium janthinellum*. Amorph. powder. [α]_D +26.7 (c, 0.2 in CHCl₃). [α]_D +76 (c, 0.25 in CHCl₃). λ_{max} 238 ; 280 (MeOH). λ_{max} 234 (log ε 3.9); 300 (log ε 3.28) (CHCl₃).


Smetanina, O.F. et al., *J. Nat. Prod.*, 2007, 70, 906-909; 2054 (isol, pmr, cmr)

Xu, M. et al., *Tetrahedron*, 2007, 63, 435-444 (isol, pmr, cmr, ms)

Shepherdine

S-285

2,3,4,9-Tetrahydro-1-methyl-1H-pyrido[3,4-b]indol-6-ol, 9CI. 1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-β-carboline [3000-36-0] [28090-87-1]



(R)-form

C₁₂H₁₄N₂O 202.255

(R)-form [104485-07-6]

Occurs in urine of man and other mammals. No phys. props. reported.

(S)-form [104885-06-5]

Occurs in urine of man and other mammals. No phys. props. reported.

(±)-form

Cryst. (MeOH). Mp 225-230° Mp 257-258°. λ_{max} 276 ; 298 (sh) ; 308 (sh) (EtOH) (HCl salt, pH7). λ_{max} 222 (log ε 4.3); 274 (log ε 3.9); 297 (sh) (log ε 3.72) (no solvent reported) (phosphate buffer, pH7).

Me ether: Mp 224-226° (156°, 196-196.5°).

Me ether, picrate: Mp 223-226° dec.

Me ether, N^b-Me: 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl-β-carboline

C₁₄H₁₈N₂O 230.309

Constit. of *Piptadenia peregrina*. Cryst. (CHCl₃/hexane). Mp 155-157°.

(ξ)-form

Alkaloid from the roots of *Shepherdia canadensis* and from the sponge *Hyrtios* sp. Under certain conditions, a metab. of Serotonin in animals. Mp 266° dec. (as hydrochloride). [α]_D 0.

N₂O-Di-Ac:

Cryst. (MeOH). Mp 192-194°.

N^b-Me: 1,2,3,4-Tetrahydro-6-hydroxy-1,2-dimethyl-β-carboline. 1,2,3,4-Tetrahydro-6-hydroxy-2-methylharman

C₁₃H₁₆N₂O 216.282

Alkaloid from *Croton moritibensis*.

Me ether: 1,2,3,4-Tetrahydro-6-methoxy-1-methyl-β-carboline. Adrenoglomerulotropin. 6-Methoxytetrahydroharman [1210-56-6] [58911-03-8]

C₁₃H₁₆N₂O 216.282

Alkaloid from *Desmodium pulchellum* and *Virola cuspidata* (Fabaceae, Myristicaceae). Also isol. from bovine pineal gland and human urine. Shows some psychotomimetic activity.

► UV0740000

Me ether, N^b-Me: [21890-60-8]

Alkaloid from *Anadenanthera peregrina* (Fabaceae). No opt. rotn. reported.

Akabori, S. et al., *Ber.*, 1930, 63, 2245-2248 (Me ether, synth)

Farrell, G.L. et al., *Arch. Biochem. Biophys.*,

1961, 94, 543-544 (Me ether, isol, struct)

Taboisky, R.G. et al., *J. Med. Chem.*, 1964, 7, 135-141 (synth, uv)

Agurell, S. et al., *Acta Chem. Scand.*, 1969, 23, 903-916 (Me ether N^b-Me, occur, synth)

Ayer, W.A. et al., *Can. J. Chem.*, 1970, 48, 1980-1984 (isol, uv, ir, pmr, synth)

Cassady, J.M. et al., *CA*, 1972, 77, 98688 (Me ether, isol)

Gynther, J. et al., *Acta Chem. Scand., Ser. B*, 1985, 39, 849-859 (Me ether, synth, ms)

Matsubara, K. et al., *Alcohol Alcohol*,

1986, 21, 339-345 (Me ether, hplc, ms, occur)

Beck, O. et al., *Biochem. Pharmacol.*, 1986, 35, 2636-2639 (R-form, S-form, hplc, occur, bibl)

Zhang, F. et al., *J. Med. Chem.*, 1992, 35, 82-93 (synth, hplc, pmr, cmr)

Callaway, J.C. et al., *J. Het. Chem.*, 1994, 31, 431-435 (synth, bibl)

Herraiz, T. et al., *J. Agric. Food Chem.*, 2000, 48, 4900-4904 (hplc, occur)

Somei, M. et al., *Chem. Pharm. Bull.*, 2001, 49, 1159-1165 ((±)-form, synth, Me ether, Me ether N^b-Me)


Salmoun, M. et al., *J. Nat. Prod.*, 2002, 65, 1173-1176 (isol, pmr, cmr)

De Araújo-Júnior, V.T. et al., *Pharm. Biol.*, 2004, 42, 62-67 (N^b-Me)

Shermilamine A

S-286

[116302-28-4]



C₂₁H₁₇BrN₄O₂S 469.361

Alkaloid from a tunicate *Trididemnum* sp. collected from Guam and from *Chelynotus semperi*. Cytotoxic agent. Orange prisms (CHCl₃/MeOH). Mp 300°. Erroneous spectroscopic data quoted in the original paper. λ_{max} 242 (ε 25100); 286 (ε 22900); 312 (ε 35500); 320 (ε 34700); 364 (ε 5500); 382 (ε 5620); 536 (ε 5890) (MeOH/HCl) (Derep). λ_{max} 238 (ε 33100); 282 (ε 28200); 298 (ε 24500); 350 (ε 7940); 392 (ε 5130); 470 (ε 5750) (MeOH) (Derep).

Debromo: Shermilamine B. Debromoshermilamine A

[122271-41-4]

C₂₁H₁₈N₄O₂S 390.465

Alkaloid from a tunicate *Trididemnum* sp., from the Red Sea tunicate *Eudistoma* sp. and from *Chelynotus semperi*. Fine orange prisms (MeOH). Mp 254° dec. λ_{max} 232 (ε 27500); 282 (ε 25100); 302 (ε 33900); 318 (ε 49000); 364 (ε 6030); 382 (ε 6170); 536 (ε 6760) (MeOH/HCl) (Derep). λ_{max} 234 (ε 34700); 282 (ε 28200); 298 (ε 24500); 348 (ε 9330); 390 (ε 5750); 468 (ε 6460) (MeOH) (Derep).

Debromo, N-de-Ac, N^o,N^o-di-Me: Shermilamine D

[211311-08-9]

C₂₁H₂₀N₄O₂S 376.481

Alkaloid from the tunicate *Cystodytes violatinctus*. Amorph. orange powder. λ_{max} 232 (log ε 4.35); 281 (log ε 4.25); 297 (log ε 4.18); 348 (log ε 3.74); 392 (log ε 3.51); 468 (log ε 3.58) (MeOH). λ_{max} 230 ; 281 ; 302 ; 316 ; 363 ; 382 ; 530 (MeOH/HCl) (Berdy).

Debromo, N-de-Ac, N^o,N^o-di-Me, N-oxide: Shermilamine E

[211311-09-0]

C₂₁H₂₀N₄O₂S 392.481

Alkaloid from the tunicate *Cystodytes violatinctus*. Amorph. brown powder. λ_{max} 232 (log ε 4.29); 296 (log ε 4.1); 349 (log ε 3.7); 391 (log ε 3.51); 464 (log ε 3.49) (MeOH). λ_{max} 230 ; 281 ; 300 ; 315 ; 361 ; 382 ; 526 (MeOH/HCl) (Berdy).


Debromo, N-de-Ac, N^o-(3-methyl-2-butenoyl): Shermilamine C

[158758-41-9]
 $C_{24}H_{22}N_4O_2S$ 430.529
 Isol. from *Cystodytes* sp. Antitumour agent. Topoisomerase II inhibitor. DNA intercalator. Orange solid. λ_{max} 231 (ϵ 9185); 279 (ϵ 6670); 296 (ϵ 5690); 350 (ϵ 2680); 390 (ϵ 1896); 463 (ϵ 1664) (MeOH).

Cooray, N.M. *et al.*, *J.O.C.*, 1988, **53**, 4619-4620 (*isol, cryst struct*)
 Carroll, A.R. *et al.*, *J.O.C.*, 1989, **54**, 4231-4232 (*isol, uv, ir, pmr, cmr, ms, struct, Shermilamine B*)
 Rudi, A. *et al.*, *J.O.C.*, 1989, **54**, 5331 (*isol, uv, ir, pmr, cmr, Shermilamine B*)
 Steffan, B. *et al.*, *Tetrahedron*, 1993, **49**, 6223 (*biosynth, Shermilamine B*)
 Eldredge, G.S. *et al.*, *J. Med. Chem.*, 1994, **37**, 3819-3827 (*Shermilamine C*)
 Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1995, **117**, 12460-12469 (*synth*)
 Koren-Goldshlager, G. *et al.*, *J.O.C.*, 1998, **63**, 4601-4603 (*Shermilamines D,E, isol, uv, ir, pmr, cmr*)

Shihumidine S-287

1'-Methylspiro[1H-isoindole-1,2'-pyrrolidin]-3(2H)-one, 9CI
 [135626-84-5]




$C_{12}H_{14}N_2O$ 202.255
 Artifact from *Dendrobium loddigesii*. Inhibitor of Na and K ATPase of rat kidney. Cryst. (C_6H_6). Mp 173-174°.

Li, M.F. *et al.*, *Yaoxue Xuebao*, 1991, **26**, 307-310 (*isol, struct*)

Shihunine S-288

1'-Methylspiro[isobenzofuran-1(3H),2'-pyrrolidin]-3-one, 9CI
 [4031-12-3]



$C_{12}H_{13}NO_2$ 203.24
 pK_a 3.6 (50% EtOH). Log P 2.46 (uncertain value) (calc).

(±)-form

Alkaloid from the orchid *Dendrobium lohohense* (component of the Chinese drug Shi-hu) and from *Dendrobium pierardii* (Orchidaceae). Na/K-ATP-ase inhibitor. Needles (Et₂O/hexane). Mp 78-79°. Log P 2.46 (uncertain value) (calc).


Picrate:

Yellow prisms (MeOH). Mp 154-155.5° Mp 163-164°.

Inubushi, Y. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 749; 1968, **16**, 1014 (*isol, uv, ir, pmr, struct*)
 Elander, M. *et al.*, *Acta Chem. Scand.*, 1971, **25**, 721 (*isol, uv, ir, pmr*)
 Breuer, E. *et al.*, *Tetrahedron*, 1975, **31**, 499 (*synth, ir, pmr*)
 Leete, E. *et al.*, *J.A.C.S.*, 1976, **98**, 6321 (*biosynth, cmr*)
 Bodem, G.B. *et al.*, *J.O.C.*, 1979, **44**, 4696 (*synth*)
 Gore, V.G. *et al.*, *Tetrahedron*, 1990, **46**, 2483 (*synth*)
 Li, M.F. *et al.*, *Yaoxue Xuebao*, 1991, **26**, 307-310 (*isol, pharmacol*)

Shishididemiol A S-289

[929641-64-5]



Absolute Configuration

$C_{45}H_{81}N_3O_{11}$ 840.148
 Isol. from a tunicate of the family Didemnidae. Antibacterial agent. Oil. $[\alpha]_D^{20}$ -33.7 (c, 1 in MeOH). λ_{max} 275 (ϵ 1195); 281 (ϵ 1020) (MeOH).

O²-Ac: Shishididemiol E

[945421-79-4]
 $C_{47}H_{83}N_3O_{12}$ 882.186
 Isol. from a tunicate of the family Didemnidae. Oil. $[\alpha]_D^{19}$ -30.7 (c, 0.5 in MeOH). λ_{max} 275 (ϵ 760); 281 (ϵ 650) (MeOH).

N^{10'}-Depropanoyl, N^{10'}-Ac: Shishididemiol C

[945421-77-2]
 $C_{44}H_{79}N_3O_{11}$ 826.122
 Isol. from a tunicate of the family Didemnidae. Oil. $[\alpha]_D^{18}$ -19 (c, 1 in MeOH). λ_{max} 275 (ϵ 870); 281 (ϵ 770) (MeOH).

Deepoxy, 7R-hydroxy, 8R-chloro: Shishididemiol B

[929641-65-6]
 $C_{45}H_{82}ClN_3O_{11}$ 876.609
 Isol. from a tunicate of the family Didemnidae. Antibacterial agent. Oil. $[\alpha]_D^{20}$ -19.4 (c, 1 in MeOH). λ_{max} 275 (ϵ 1220); 281 (ϵ 1040) (MeOH).

Deepoxy, 7R-hydroxy, 8R-chloro, N^{10'}-depropanoyl, N^{10'}-Ac: Shishididemiol D


[945421-78-3]
 $C_{44}H_{80}ClN_3O_{11}$ 862.582
 Isol. from a tunicate of the family Didemnidae. Oil. $[\alpha]_D^{19}$ -16.6 (c, 1 in MeOH). λ_{max} 275 (ϵ 1100); 281 (ϵ 900) (MeOH).

Kobayashi, H. *et al.*, *J.O.C.*, 2007, **72**, 1218-1225 (*Shishididemniols A,B*)

Kobayashi, H. *et al.*, *Tetrahedron*, 2007, **63**, 6748-6754 (*Shishididemniols C,D,E*)

Shishijimicin A S-290

[503860-50-2]



$C_{46}H_{52}N_4O_{12}S_4$ 981.2
 Isol. from the ascidian *Didemnum proliferum*. Cytotoxic.

4'-N-Deisopropyl, 4'-N-Et: Shishijimicin C

[503860-52-4]
 $C_{45}H_{50}N_4O_{12}S_4$ 967.174
 Isol. from *Didemnum proliferum*. Cytotoxic.


4'-De(methylthio): Shishijimicin B

[503860-51-3]
 $C_{45}H_{50}N_4O_{12}S_3$ 935.108
 Isol. from *Didemnum proliferum*. Cytotoxic.

Oku, N. *et al.*, *J.A.C.S.*, 2003, **125**, 2044-2045 (*isol*)

Siamenol S-291

6-Methyl-3-(3-methyl-2-butenyl)-9H-carbazol-2-ol. 2-Hydroxy-6-methyl-3-prenyl-carbazole
 [264622-44-8]




$C_{18}H_{19}NO$ 265.354
 Alkaloid from *Murraya siamensis*. HIV inhibitor. Pale yellow solid. λ_{max} 216 (log ϵ 4.03); 238 (log ϵ 4.11); 261 (log ϵ 3.73); 308 (log ϵ 3.71); 328 (sh) (log ϵ 3.4) (MeOH).

Meragelman, K.M. *et al.*, *J. Nat. Prod.*, 2000, **63**, 427-428

Sibirinine S-292

Hexahydro-2-methyl-4H,7H-methano-2H-1,3-benzoxazocine 3-oxide, 9CI
 [113866-79-8]



Absolute Configuration

$C_{12}H_{21}NO_2$ 211.303
 Alkaloid from *Nitraria sibirica* (Zygophyllaceae). Cryst. Mp 40°. $[\alpha]_D^{20}$ -9.4 (c,

0.53 in CHCl_3).


[120095-27-4]

Ibragimov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 685-689; *Chem. Nat. Compd. (Engl. Transl.)*, 1987, **23**, 569-573 (*isol, pmr, abs config, synth*)

Ibragimov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1988, **24**, 82-91; *Chem. Nat. Compd. (Engl. Transl.)*, 1988, **24**, 71-78 (*pmr, cmr*)

Sibyllimycin S-293

6,7-Dihydro-3-methylimidazo[1,2-a]pyridin-8(5H)-one, 9CI
[176520-65-3]



$\text{C}_8\text{H}_{10}\text{N}_2\text{O}$ 150.18

Metab. of *Thermoactinomyces* sp. Cryst. (toluene). Mp 122-123°. λ_{max} 240 (ϵ 9379); 297 (ϵ 24481) (CHCl_3).

Hafenbradl, D. *et al.*, *Angew. Chem., Int. Ed.*, 1996, **35**, 545 (*isol, pmr, cmr, synth, struct*)

Sickenbergine S-294

[1361-17-7]

$\text{C}_{16}\text{H}_{19}\text{NO}_5$ 305.33

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Pancreatium sickenbergeri*, *Pancreatium arabicum*, *Pancreatium maritimum* and *Pancreatium tortuosum* (Amaryllidaceae). Mp 110° dec.

Hydrochloride: Mp 175° dec.


Fahmy, I.R. *et al.*, *J. Pharm. Sci. U.A.R.*, 1960, **1**, 133-137; *CA*, **56**, 10280i (*isol*)

Ahmed, Z.F. *et al.*, *J. Nat. Prod.*, 1964, **27**, 115-134; *CA*, **61**, 13630d (*occur*)

Siderochelin A S-295

3,4-Dihydro-4-hydroxy-5-(3-hydroxy-2-pyridinyl)-4-methyl-2H-pyrrole-2-carboxamide, 9CI. EM 4940A. Antibiotic EM 4940A

[77550-87-9]



R = CH_3

$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_3$ 235.242

Isol. from the fermentation broths of *Nocardia* sp. SC 11,340 and from an unusual *Actinomycetes* sp. Weakly active against gram-positive, gram-negative and anaerobic bacteria. Sol. MeOH, CHCl_3 ; fairly sol. MeCN; poorly sol. H_2O , hexane. Mp 165-168° (161-163°). $[\alpha]_{\text{D}}^{22}$ +1.7 (c, 0.30 in MeOH). λ_{max} 315 (ϵ 8350) (MeOH) (Derep). λ_{max} 206 (E1%/1cm 480); 315 (E1%/1cm 220) (MeOH) (Berdy). λ_{max} 316 (ϵ 8350) (EtOH)

(Berdy).

4-Epimer: Siderochelin B

[77518-26-4]

$\text{C}_{11}\text{H}_{13}\text{N}_3\text{O}_3$ 235.242

Isol. from *Nocardia* sp. SC 11,340 and from an unusual *Actinomycetes* sp.

Weakly active against gram-positive, gram-negative and anaerobic bacteria. Sol. MeOH, CHCl_3 , MeCN; poorly sol. H_2O , hexane. Mp 200-201° (186-186.5°). $[\alpha]_{\text{D}}^{22}$ -1.9 (c, 0.21 in MeOH). λ_{max} 315 (ϵ 8350) (MeOH) (Derep).

Liu, W.-C. *et al.*, *J. Antibiot.*, 1981, **34**, 791 (*isol, uv, ir, pmr, cryst struct*)

Okuyama, D. *et al.*, *J. Antibiot.*, 1982, **35**, 1240 (*abs config*)

Mitscher, L.A. *et al.*, *J. Antibiot.*, 1984, **37**, 1260 (*isol, uv, ir, pmr, cd*)

Siderochelin C S-296

[93973-61-6]

As Siderochelin A, S-295 with

R = $-\text{CH}_2\text{CH}_3$


$\text{C}_{12}\text{H}_{15}\text{N}_3\text{O}_3$ 249.269

Fermentation product of *Streptoalloteichus* sp. 69-209B. Weakly active against *E. coli*. Cryst. Sol. MeOH, EtOAc; poorly sol. H_2O , hexane. Mp 168-171°. λ_{max} 315 (ϵ 8350) (MeOH) (Derep). λ_{max} 316 (ϵ 7340) (MeOH) (Berdy).

Mitscher, L.A. *et al.*, *J. Antibiot.*, 1984, **37**, 1260 (*isol, uv, ir, pmr, cd, struct*)

Sieboldine A S-297

[619326-44-2]



Relative Configuration


$\text{C}_{16}\text{H}_{23}\text{NO}_4$ 293.362

Alkaloid from *Lycopodium sieboldii*. ACE inhibitor. Needles (MeOH). $[\alpha]_{\text{D}}$ +139 (c, 0.3 in MeOH). Dec. at 160°.

Hirasawa, Y. *et al.*, *Org. Lett.*, 2003, **5**, 3991-3993 (*isol, pmr, cmr*)

Siechuantane S-298

[162616-59-3]



$\text{C}_{27}\text{H}_{43}\text{NO}_3$ 429.642

Alkaloid from bulbs of *Fritillaria siechuanica* (Liliaceae).

Wang, F. *et al.*, *CA*, 1995, **122**, 261018h (*isol, struct*)

Signatipennine S-299

[183163-68-0]

$\text{HOCH}_2\text{CH}_2\text{NH}(\text{CH}_2)_{15}\text{CH}(\text{CH}_3)\text{NHC}-\text{H}_2\text{CH}_2\text{O}_2$


$\text{C}_{21}\text{H}_{46}\text{N}_2\text{O}_2$ 358.607

Alkaloid from the beetle *Epilachna signatipennis* (Coccinellidae). Isol. and characterised as the tetra-Ac.

Wang, S.F. *et al.*, *Bull. Soc. Chim. Belg.*, 1996, **105**, 483 (*isol*)

Silvaticamide S-300

[73606-98-1]



$\text{C}_{25}\text{H}_{29}\text{NO}_5$ 423.508

Secoanthraquinone. Prod. by *Aspergillus silvaticus*. Mycotoxin. Sol. MeOH, C_6H_6 ; poorly sol. H_2O . Mp 191-192° dec. Unstable to light. λ_{max} 290 (ϵ 4400); 310 (ϵ 5300) (MeOH) (Berdy).


Yamazaki, M. *et al.*, *Heterocycles*, 1981, **15**, 889 (*isol, struct*)

Maebayashi, Y. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 4296 (*biosynth*)

Simaomicin β S-301

7a,8,11,12,13,14-Hexahydro-11,14,16,17-tetrahydroxy-9-methoxy-3-methyl-2H-xantho[2',3':6,7][1,3]benzodioxino[4,5-fg]isoquinoline-1,15-dione, 9CI. LL-D 42067 β

[100111-39-5]



Relative Configuration


$\text{C}_{27}\text{H}_{23}\text{NO}_{10}$ 521.479

Prod. by *Actinomadura madura* ssp. *simaoensis*. Anticoccidial agent. Yellow cryst. Similar to Actinoplanone A in A-128. λ_{max} 253 (ϵ 35500); 320 (ϵ 12900); 379 (sh) (ϵ 22900); 395 (ϵ 25100) (MeOH) (Derep).

- N-Me: **Simaomicin α** . LL-D 42067 α
[100157-22-0]
C₂₈H₂₅NO₁₀ 535.506
From *Actinomadura madura* ssp. *si-maensis* NRRL15734 and *Micromonospora purpurochromogenes* wuxiensis NRRL18075. Anticoccidial agent. Yellow cryst. Sol. MeOH, CH₂Cl₂, EtOAc; poorly sol. H₂O. $[\alpha]_D^{26}$ +836 (c, 0.3 in DMF). λ_{\max} 253 (ε 35500); 320 (ε 12900); 379 (sh) (ε 22900); 395 (ε 25100) (MeOH) (Dererp). λ_{\max} 213 (ε 27000); 253 (ε 34500); 321 (ε 12200); 374 (ε 21100); 389 (ε 8290) (MeOH/HCl) (Berdy). λ_{\max} 217 (ε 42100); 253 (ε 13900); 312 (ε 5700); 395 (ε 10700) (MeOH/NaOH) (Berdy).
- U.S. Pat., 1985, 4551533; CA, 104, 62068 (LL-D 42067 α)
- Lee, T.M. et al., *Chem. Comm.*, 1989, 1771-1772 (isol, struct)
- Carter, G.T. et al., *J.O.C.*, 1989, 54, 4321-4323 (biosynth)
- Maiese, W.M. et al., *J. Antibiot.*, 1990, 1059-1063 (isol)

Simmondsin S-302

[6-(β -D-Glucopyranosyloxy)-2-hydroxy-3,4-dimethoxycyclohexylidene]acetone-trile, 9CI. 2-Cyanomethylene-1-glucosyloxy-3-hydroxy-4,5-dimethoxycyclohexane
[51771-52-9]



C₁₆H₂₅NO₉ 375.375
Constit. of *Simmondsia californica* and *Simmondsia chinensis* (jojoba). Appetite suppressant. Waxy solid. Mp 95-100°. $[\alpha]_D^{25}$ -78 (c, 1 in MeOH).

▶ AM0250000

Penta-Ac: Mp 165-166°.

2'-O-(4-Hydroxy-3-methoxy-E-cinnamoyl): **2'-O-Feruloylsimmondsin**
[179466-18-3]
[67411-22-7, 179466-19-4]

C₂₆H₃₃NO₁₂ 551.546
Constit. of the seeds of jojoba (*Simmondsia chinensis*).

3-O-De-Me, 2'-(4-hydroxy-3-methoxy-Z-cinnamoyl): [162290-39-3]
[179466-20-7]

C₂₅H₃₁NO₁₂ 537.519
Constit. of jojoba meal (*Simmondsia chinensis*).

4-O-De-Me: **4-O-Demethylsimmondsin**
[135105-75-8]

C₁₅H₂₃NO₉ 361.348
Constit. of jojoba meal (*Simmondsia chinensis*).

4-O-De-Me, 2'-(4-hydroxy-3-methoxy-E-cinnamoyl): [162290-38-2]
[179601-21-9]
C₂₅H₃₁NO₁₂ 537.519

Constit. of jojoba meal (*Simmondsia chinensis*).

Di-O-de-Me: **Dide-O-methylsimmondsin**
[135074-86-1]

C₁₄H₂₁NO₉ 347.321
Constit. of jojoba meal (*Simmondsia chinensis*). Amorph. $[\alpha]_D^{20}$ -50 (c, 0.52 in H₂O). λ_{\max} 215 (log ε 3.97) (H₂O).

Di-O-de-Me, 4-O- α -D-glucopyranoside: **Dide-O-methyl-4-O- α -D-glucopyranosylsimmondsin**
[370068-00-1]

C₂₀H₃₁NO₁₄ 509.463
Constit. of jojoba meal (*Simmondsia chinensis*).

Di-O-de-Me, 2-O-(3-methyl-2-butenoyl): **Ehretioside A₃**
[156368-87-5]

C₁₉H₂₇NO₁₀ 429.423
Constit. of *Ehretia philippinensis*. Waxy solid. $[\alpha]_D^{25}$ -53 (c, 1 in MeOH). Has (E)-config. (change of Cahn-Ingold-Prelog priorities).

Di-O-de-Me, 3-O-(3-methyl-2-butenoyl): **Ehretioside A₁**
[156368-85-3]

C₁₉H₂₇NO₁₀ 429.423
Constit. of *Ehretia philippinensis*. Amorph. $[\alpha]_D^{25}$ +39 (c, 1 in Py).

Di-O-de-Me, 4-O-(3-methyl-2-butenoyl): **Ehretioside A₂**
[156368-86-4]

C₁₉H₂₇NO₁₀ 429.423
Constit. of *Ehretia philippinensis*. Waxy solid. $[\alpha]_D^{25}$ -53 (c, 1 in MeOH).

Di-O-de-Me, 2-O-benzoyl: **Lanceolin A**
[158022-45-8]

C₂₁H₂₅NO₁₀ 451.429
Constit. of *Lophira lanceolata*. Amorph. solid. Has (E)-config. (change of Cahn-Ingold priorities).

Di-O-de-Me, 3-O-benzoyl: **Lanceolin C**
[221149-42-4]

C₂₁H₂₅NO₁₀ 451.429
Constit. of *Lophira alata*. Amorph. solid.

Di-O-de-Me, 4-O-benzoyl: **Lanceolin B**
[151197-19-2]

C₂₁H₂₅NO₁₀ 451.429
Constit. of *Lophira lanceolata*. Amorph. solid.

Di-O-de-Me, 2,4-di-O-benzoyl: **Lophiroside A₂**
[151171-81-2]

C₂₈H₂₉NO₁₁ 555.537
Constit. of the bark of *Lophira alata*. Amorph. $[\alpha]_D^{20}$ -22.6 (c, 0.93 in MeOH). λ_{\max} 229 (log ε 4.54); 274 (log ε 3.26) (MeOH).

Di-O-de-Me, 3,4-di-O-benzoyl: **Lophiroside A₁**
[151171-80-1]

C₂₈H₂₉NO₁₁ 555.537
Constit. of the bark of *Lophira alata*. Amorph. $[\alpha]_D^{20}$ +11.5 (c, 0.87 in MeOH). λ_{\max} 229 (log ε 4.54); 274 (log ε 3.26) (MeOH).

Di-O-de-Me, 2-O-E-cinnamoyl, 4-O-benzoyl: **Lophiroside B₂**
[151171-83-4]

C₃₀H₃₁NO₁₁ 581.575

Constit. of the bark of *Lophira alata*. Amorph. $[\alpha]_D^{20}$ -22.9 (c, 0.83 in MeOH). Has (E)-config. (change of Cahn-Ingold-Prelog priorities). λ_{\max} 222 (log ε 4.25); 279 (log ε 4.14) (MeOH).

Di-O-de-Me, 3-O-E-cinnamoyl, 4-O-benzoyl: **Lophiroside B₁**
[151171-82-3]

C₃₀H₃₁NO₁₁ 581.575
Constit. of the bark of *Lophira alata*. Amorph. $[\alpha]_D^{20}$ +63 (c, 0.54 in MeOH). λ_{\max} 222 (log ε 4.25); 279 (log ε 4.14) (MeOH).

Elliger, C.A. et al., *J.O.C.*, 1974, 39, 2930 (struct)

Elliger, C.A. et al., *Phytochemistry*, 1974, 13, 2319 (isol)

Verbiscar, A.J. et al., *J. Agric. Food Chem.*, 1978, 26, 1456-1459 (2-ferulate)

Chida, N. et al., *J.C.S. Perkin 1*, 1992, 1131 (synth, abs config)

Van Boven, M. et al., *J. Agric. Food Chem.*, 1993, 41, 1605; 1994, 42, 2684; 1996, 44, 2239-2243 (isol, cryst struct, derivs)

Murakami, A. et al., *Phytochemistry*, 1993, 32, 1461-1466 (*Lophirosides*)

Tih, A.E. et al., *J. Nat. Prod.*, 1994, 57, 971-974 (*Lanceolin A*, *Lanceolin B*)


Simpol, L.R. et al., *Phytochemistry*, 1994, 36, 91-95 (*Ehretiosides*)

Messanga, B.B. et al., *Fitoterapia*, 1998, 79, 439-442 (*Lanceolin C*)

Van Boven, M. et al., *J. Agric. Food Chem.*, 2001, 49, 4278-4283 (*Didemethyl-4-glucosylsimmondsin*)

Simplakidine A

S-303



Absolute Configuration

C₂₄H₃₇NO₆ 435.559


Alkaloid from the sponge *Plakortis simplex*. Weakly cytotoxic. Amorph. solid. $[\alpha]_D^{25}$ -21.7 (c, 1.2 in MeOH). λ_{\max} 272 (log ε 3.55) (MeOH).

Campagnuolo, C. et al., *Org. Lett.*, 2003, 5, 673-676 (isol, pmr, cmr, ms)

Simplexine†

S-304

[118636-89-8]



Absolute Configuration


C₁₃H₁₇NO₄ 251.282

Alkaloid from the whole plant of *Phyllanthus simplex* (Euphorbiaceae). $[\alpha]_D^{20}$ -20.4 (c, 0.2 in EtOH).



Negi, R.S. et al., *Phytochemistry*, 1988, 27, 3027-3028 (isol, uv, ir, pmr, ms, struct)

Simulanoquinoline S-305

[155416-22-1]

C₃₇H₃₄N₂O₇ 618.685Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Prisms (CHCl₃/MeOH). Mp 240-243°.Wu, S.-J. *et al.*, *Phytochemistry*, 1993, **34**, 1659 (isol, uv, ir, pmr, ms, struct)**Simulansamide** S-306

[176713-29-4]

C₂₂H₂₃NO₆ 397.427A secobenzo[*c*]phenanthridine alkaloid. Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper). Platelet aggregation inhibitor. Cryst. (MeOH/CHCl₃). Mp 257-259°.Wu, S.-J. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1996, **43**, 195-198; *CA*, **124**, 337827k (synth, uv, ir, pmr, ms)**Sinactine** S-307*Tetrahydroepiberberine*

(R)-form

C₂₀H₂₁NO₄ 339.39**(R)-form** [82051-55-6]Alkaloid from *Corydalis meifolia* (Papaveraceae). Cryst. (MeOH). Mp 177-178°. [α]_D +298 (c, 1.04 in CHCl₃).**(S)-form** [22554-51-4]Alkaloid from *Fumaria officinalis* and *Sinomenium acutum* (Papaveraceae, Menispermaceae). Mp 175°. [α]_D -312 (CHCl₃). λ_{max} 232 (ε 13600); 287 (ε 7200) (EtOH).

N-Me: N-Methylsinactine


[82667-90-1]

C₂₁H₂₄NO₄[⊕] 354.425Quaternary alkaloid from *Fumaria officinalis* (Papaveraceae). Needles (EtOH/CHCl₃) (as hydroxide). Mp 318-320° (hydroxide). [α]_D²⁰ -260 (c, 0.11 in H₂O) (hydroxide).**O²-De-Me: Cheilanthifoline**

[483-44-3]

C₁₉H₁₉NO₄ 325.363Alkaloid from *Corydalis cheilanthifolia*, *Corydalis scouleri*, *Corydalis sibirica* and some *Argemone* spp. (Papaveraceae). Mp 184°. [α]_D²⁰ -311 (MeOH).**(±)-form** [38853-67-7]Alkaloid from *Fumaria officinalis* (Papaveraceae). Mp 169-170°.**O²-De-Me:** [41431-80-5]C₁₉H₁₉NO₄ 325.363Alkaloid from whole plants of *Eschscholtzia californica*. Mp 176-178° (synthetic) Mp 190° (natural).Goto, K. *et al.*, *J.C.S.*, 1930, 1234-1237 (*S-form*, isol, struct)Manske, R.H.F. *et al.*, *Can. J. Res.*, 1940, **18**, 100-102 (*isol*)Cooper, S.F. *et al.*, *Planta Med.*, 1972, **21**, 313 (*synth*, pmr, ir)Ninomiya, I. *et al.*, *J.C.S. Perkin 1*, 1975, 1720-1724 (*synth*)Pavelka, S. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3157-3169 (*uv*)Furuya, T. *et al.*, *Phytochemistry*, 1978, **17**, 891-893 (*biosynth*)Pai, B.R. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 607-611 (*synth*)Mardirossian, Z.H. *et al.*, *Phytochemistry*,1983, **22**, 759-761 (N-Methylsinactine)Bhakuni, D.S. *et al.*, *Tetrahedron*, 1983, **39**, 455-459 (*biosynth*, abs config)Narasimhan, N.S. *et al.*, *Tetrahedron*, 1983, **39**, 1975-1982 (*synth*)Jain, L. *et al.*, *Planta Med.*, 1996, **62**, 188 ((±)-form O²-de-Me, isol)Seger, C. *et al.*, *Magn. Reson. Chem.*, 2004, **42**, 882-886 (pmr, cmr)Wang, X.-L. *et al.*, *Acta Cryst. E*, 2006, **62**, 81-82; 1271-1272 (*Sinactine*, *Cheilanthifoline*, *cryst structs*)**Sinapine** S-308

2-[[3-(4-Hydroxy-3,5-dimethoxyphenyl)-1-oxo-2-propenyl]oxy]-N,N,N-trimethylthalaminium(1+), 9CI. Choline 4-hydroxy-3,5-dimethoxycinnamate, 8CI [18696-26-9]

C₁₆H₂₄NO₅[⊕] 310.369Alkaloid from seeds of *Brassica nigra*, *Brassica alba* (*Sinapis alba*) and many other spp. in the Brassicaceae. Also from the Chinese drug Ting Li (*Draba nemorosa*) (Brassicaceae). Astringent antinutrient of practical importance, esp. in cattle feed, e.g. rape meal. Reported to have radioprotective effects in some cruciferous plants and in *Drosophila*.

Iodide: [5655-06-1]

C₁₆H₂₄INO₅ 437.274

Mp 185-186° (178-179°).

Sulfate: [6509-38-2]C₁₆H₂₅NO₈S 391.441

Mp 186-188° (anhyd.) Mp 126.5-127.5° dec. (dihydrate).

O-β-D-Glucopyranoside: [21848-40-8]


[87206-57-3, 87206-59-5]

C₂₂H₃₄NO₁₀[⊕] 472.511Alkaloid from the seeds of *Alliaria officinalis*. Semicryst. (as acetate salt).Obt. as a mix. of *E*- and *Z*-isomers. λ_{max} 315 (H₂O).

[84123-22-8, 84123-21-7]

Gadamer, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1897, **235**, 44 (*isol*)Späth, E. *et al.*, *Monatsh. Chem.*, 1920, **41**, 271 (*synth*)Kung, H.P. *et al.*, *J.A.C.S.*, 1949, **71**, 1836 (*isol*, uv)Schwartz, P. *et al.*, *Naturwissenschaften*, 1949, **36**, 88 (*isol*)Tzagoloff, A. *et al.*, *Plant Physiol.*, 1963, **38**, 202 (*metab*)Durkee, A.B. *et al.*, *J. Food Sci.*, 1975, **40**, 820 (*occur*)Sosulski, F.W. *et al.*, *Dev. Food Proteins*, 1983, **2**, 109 (*rev*)Larsen, L.M. *et al.*, *Phytochemistry*, 1983, **22**, 219-222 (*glucoside*)Regenbrecht, J. *et al.*, *Phytochemistry*, 1985, **24**, 407**Singapurensine A** S-309

[158182-15-1]



Absolute Configuration

C₂₃H₂₆N₂O₇ 442.468Alkaloid from *Kopsia singapurensis* (Apocynaceae).**Me ether: Singapurensine B**

[158182-16-2]

C₂₄H₂₈N₂O₇ 456.494From *Kopsia singapurensis* (Apocynaceae).**12-Methoxy: Kopsidine C**

[164081-05-4]

C₂₄H₂₈N₂O₈ 472.494Alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D +12 (c, 0.208 in CHCl₃). λ_{max} 217 (log ε 4.5); 254 (log ε 3.99); 282 (log ε 3.28); 289 (log ε 3.25) (EtOH).**12-Methoxy, O¹⁵-Me: Kopsidine A**

[149355-64-6]

C₂₅H₃₀N₂O₈ 486.521Novel heptacyclic skeleton. Alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D +16 (c, 0.21 in CHCl₃).**12-Methoxy, O¹⁵-Et: Kopsidine B**

[149355-65-7]

C₂₆H₃₂N₂O₈ 500.547Alkaloid from leaves of *Kopsia teoi* (Apocynaceae). [α]_D +11 (c, 0.073 in CHCl₃).**11,12-Methylenedioxy: Singapurensine C**

[158182-33-3]
 $C_{24}H_{26}N_2O_9$ 486.477
 From *Kopsia singaporensis* (Apocynaceae).

11,12-Methylenedioxy, Me ether: **Singaporensine D**
 [158182-17-3]
 $C_{25}H_{28}N_2O_9$ 500.504
 From *Kopsia singaporensis* (Apocynaceae).

15-Epimer, 12-methoxy: **Kopsidine D**
 [192995-27-0]
 $C_{24}H_{28}N_2O_8$ 472.494
 Alkaloid from leaves of *Kopsia teoi*.
 $[\alpha]_D^{25}$ -16 (c, 0.058 in $CHCl_3$). λ_{max} 217 (log ϵ 4.36); 254 (log ϵ 3.9); 282 (log ϵ 3.33); 289 (log ϵ 3.3) (EtOH).

Awang, K. et al., *Nat. Prod. Lett.*, 1993, **3**, 283-289 (*Singaporensines*)
 Kan-Fan, C. et al., *Nat. Prod. Lett.*, 1993, **3**, 291-298 (*Kopsidines, synth*)
 Kam, T.-S. et al., *Tet. Lett.*, 1993, **34**, 1819-1822 (*Kopsidines, isol, uv, pmr, cmr, ms, struct*)
 Tan, G.-H. et al., *Tet. Lett.*, 1995, **36**, 1327-1330 (*Kopsidines, synth*)
 Kam, T.-S. et al., *Phytochemistry*, 1997, **45**, 623-625 (*Kopsidine D*)

Sinharine S-310

$PhCH_2CH_2NHCOCH=CHSM_e$
 $C_{12}H_{15}NOS$ 221.323
 Struct. revised in 1994. λ_{max} 263 (Et₂O) (Berdy).


(E)-form [142717-65-5]
 Isol. from *Glycosmis cyanocarpa* (Rutaceae). Antifungal agent. Cryst. Mp 74-75°.

N-Me: **Methylsinharine**
 [142717-66-6]
 $C_{13}H_{17}NOS$ 235.349
 From *Glycosmis cyanocarpa*. Antifungal agent. Oil. λ_{max} 269 (Et₂O) (Berdy).

Greger, H. et al., *Tetrahedron*, 1992, **48**, 1209 (*isol, pmr, cmr*)
 Johnson, W.M. et al., *Aust. J. Chem.*, 1994, **47**, 751-756 (*synth*)
 Hinterberger, S. et al., *Tetrahedron*, 1994, **50**, 6279 (*synth, struct*)

Sinococuline S-311

8,14-Didehydro-3,8-dimethoxymorphinan-4,6,7-triol, 9CI. Alkaloid FK 1000. FK 1000
 [109351-36-2]



$C_{18}H_{23}NO_5$ 333.383
 Alkaloid from the stems and rhizomes of *Cocculus trilobus* (Menispermaceae). Possesses antitumour activity. Amorph.

powder. $[\alpha]_D^{25}$ -137.4 (c, 0.12 in MeOH). The previously reported specific rotn. (-77) is erroneous. λ_{max} 234 (ϵ 8000); 283 (ϵ 2700) (EtOH).

6,7-Di-Ac: **Alkaloid FK 3000. FK 3000.**
 6,7-Di-O-acetylsinococuline
 [115439-63-9]
 $C_{22}H_{27}NO_7$ 417.458
 Alkaloid from roots of *Stephania cepharantha* (Menispermaceae). Needles (EtOAc). Mp 185-186°. $[\alpha]_D^{25}$ -163 (c, 0.73 in MeOH). The CA name is erroneous: 4,5,6-triol instead of 4,6,7-triol. λ_{max} 258 (ϵ 3720); 284 (ϵ 1660) (MeOH) (Berdy). λ_{max} 217 (log ϵ 4.28); 286 (log ϵ 3.4) (no solvent reported).

N-Me: N-Methylsinococuline. **Cephasugine**
 [189390-47-4]
 $C_{19}H_{25}NO_5$ 347.41
 Alkaloid from *Stephania cepharantha*. Amorph. powder. $[\alpha]_D^{25}$ -98 (c, 0.3 in $CHCl_3$). λ_{max} 283 (log ϵ 3.22) (MeOH).

6,7-Di-Ac, N-Me: **Cephakicine**
 [175617-22-8]
 $C_{23}H_{29}NO_7$ 431.485
 Alkaloid from tubers of *Stephania cepharantha*. Amorph. powder. $[\alpha]_D^{28}$ -161 (c, 0.22 in $CHCl_3$). CAS name for Cephakicine is defective (the methyl group at N-17 is omitted).

16,17-Didehydro: **Cepharmorphanine**
 [142905-23-5]
 $C_{18}H_{21}NO_5$ 331.368
 Alkaloid from roots of *Stephania cepharantha* (Menispermaceae). Needles (Me₂CO/MeOH). Mp 284-285°. $[\alpha]_D^{25}$ -118 (c, 0.07 in MeOH). λ_{max} 215 (log ϵ 4.23); 235 (sh) (log ϵ 4); 270 (log ϵ 3.25) (no solvent reported).


6-Epimer, 10 β -hydroxy, O⁶, O⁷-di-Ac: **Fenfangine G**
 [205533-81-9]
 $C_{22}H_{27}NO_8$ 433.457
 Alkaloid from the roots of *Stephania tetrandra*. Needles (EtOAc). Mp 203-205°. $[\alpha]_D^{24}$ -63.7 (c, 0.8 in $CHCl_3$).

2-Methoxy, 4-deoxy, O³-de-Me: **Isosinococuline**
 [164176-42-5]
 $C_{18}H_{23}NO_5$ 333.383
 Alkaloid from rhizomes of *Cocculus trilobus*. Antitumour agent. Cryst. powder. Mp 149-151° dec. $[\alpha]_D^{26}$ -66 (c, 0.31 in MeOH). λ_{max} 208 (log ϵ 4.5); 287 (log ϵ 3.68); 297 (sh) (log ϵ 3.62) (MeOH).

Itokawa, H. et al., *Chem. Pharm. Bull.*, 1987, **35**, 1660-1662 (*isol, uv, pmr, cmr, cd, struct*)
 Deng, J.-Z. et al., *Phytochemistry*, 1992, **31**, 1448-1450 (FK-3000, *Cepharmorphanine*)
 Itokawa, H. et al., *Bioorg. Med. Chem. Lett.*, 1995, **5**, 821 (*Isosinococuline*)
 Hitotsuyanagi, Y. et al., *J. O. C.*, 1995, **60**, 4549-4558 (*synth, Sinococuline, epimers*)
 Kashiwaba, N. et al., *J. Nat. Prod.*, 1996, **59**, 476-480 (*Cephakicine*)
 Kashiwaba, N. et al., *Chem. Pharm. Bull.*, 1997, **45**, 545-548 (*Cephasugine*)
 Ogino, T. et al., *Heterocycles*, 1998, **48**, 311-317 (*Fenfangine G*)

Sinomendine S-312

[143601-13-2]




$C_{20}H_{19}NO_4$ 337.374
 Minor alkaloid from rhizomes of *Sinomenium acutum* (Menispermaceae).

Chen, Y. et al., *Beijing Yike Daxue Xuebao*, 1991, **23**, 235; *CA*, **117**, 167656q

Sinomenine S-313

7,8-Didehydro-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one, 9CI. **Cucoline. Kukoline**
 [115-53-7]



Absolute configuration

$C_{19}H_{23}NO_4$ 329.395
 Alkaloid from *Sinomenium acutum* and *Stephania cepharantha* (Menispermaceae). Starting material for synth. of enantiomeric morphine derivs. Abortifacient in large doses. Immunosuppressant. Analgesic, antiinflammatory, antirheumatic props. Mp 161° Mp 182° (double Mp). $[\alpha]_D^{25}$ -71 (c, 2.1 in EtOH). Log P 0.46 (uncertain value) (calc).

▶ LD₅₀ (mus, orl) 580 mg/kg. LD₅₀ (mus, ipr) 285 mg/kg. QD2170000

Hydrochloride:

Cryst. + 2H₂O. Mp 231° dec. $[\alpha]_D^{17}$ -82 (c, 4.4 in H₂O).

N-Oxide: **Sinomenine N-oxide**

$C_{19}H_{23}NO_5$ 345.394
 Alkaloid from the stems of *Sinomenium acutum*. Powder (MeOH). Mp 155-156°. $[\alpha]_D^{20}$ -79 (c, 0.5 in MeOH).

N-De-Me: **N-Demethylsinomenine**
 $C_{18}H_{21}NO_4$ 315.368
 Alkaloid from the stems of *Sinomenium acutum*. Cryst. ($CHCl_3$). Mp 222-223°. $[\alpha]_D^{20}$ -108 (c, 0.5 in MeOH).

8-Methoxy: **Cephamonine**

[162857-99-0]
 $C_{20}H_{25}NO_5$ 359.421
 Alkaloid from tubers of *Stephania cepharantha* (Menispermaceae). Amorph. powder; prisms (MeOH/Et₂O) as hydrochloride. Mp 183-186° (hydrochloride). $[\alpha]_D^{22}$ -36 (c, 0.11 in $CHCl_3$).

14-Epimer: **14-Episinomenine**

[60761-53-7]
 $C_{19}H_{23}NO_4$ 329.395
 Alkaloid from *Ocotea brachybotra* (Lauraceae). Cryst. + 1/2 C₆H₆ (C₆H₆).

Mp 118-120°. $[\alpha]_D^{20}$ -40 (c, 1 in CHCl_3).

14-Epimer; hydrochloride:

Cryst. + $2\text{H}_2\text{O}$ (MeOH). Mp 200-210°.

14-Epimer, 8-methoxy: **Cephamuline**

[162990-77-4]

$\text{C}_{20}\text{H}_{25}\text{NO}_5$ 359.421

Alkaloid from tubers of *Stephania cepharantha* (Menispermaceae).

Amorph. powder. $[\alpha]_D^{25}$ -63 (c, 0.18 in CHCl_3).

1,1'-Dimer: **1,1'-Bisinomenine**. Dehydrosinomenine. 1,1'-Disinomenine

[596-58-7]

$\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_8$ 656.774

Alkaloid from seeds and stems of *Sinomenium acutum*. Also obt. by oxidn. of Sinomenine. Tablets (MeOH), needles (C_6H_6). Mp 222°. $[\alpha]_D^{25}$ +97.4 (c, 0.4 in MeOH). Possible artifact of oxidn.

1,1'-Dimer, 7S,8-dihydro: **7,8-Dihydro-1,1'-bisinomenine**. 7,8-Dihydro-1,1'-disinomenine

$\text{C}_{38}\text{H}_{46}\text{N}_2\text{O}_8$ 658.79

Alkaloid from the stems of *Sinomenium acutum*. Amorph. powder. Mp 220-221°. $[\alpha]_D^{25}$ +79 (c, 0.06 in MeOH). Possible artifact of oxidn. λ_{max} 211; 290 (MeOH).

2,2'-Dimer: **2,2'-Bisinomenine**. 2,2'-Disinomenine

$\text{C}_{38}\text{H}_{44}\text{N}_2\text{O}_8$ 656.774

Alkaloid from the stems of *Sinomenium acutum*. Needles ($\text{CHCl}_3/\text{MeOH}$). Mp 214-215°. $[\alpha]_D^{25}$ +48 (c, 0.16 in CHCl_3). Possible artifact of oxidn. λ_{max} 208; 294 (MeOH).

Goto, K. *et al.*, *Annalen*, 1931, **485**, 247 (struct)

Bentley, K.W. *et al.*, *J.C.S.*, 1955, 3252

Goto, K. *et al.*, *Sinomenine*, Kitasoto Institute, Tokyo, 1964, (book)

Vecchietti, V. *et al.*, *Tet. Lett.*, 1976, 1631 (epimer)

Bjorevåg, S.V. *et al.*, *Acta Cryst. C*, 1983, **39**, 1066 (cryst struct, bibl)

Hojo, H. *et al.*, *J. Immunopharmacol.*, 1985, **7**, 33 (pharmacol)

Liu, L. *et al.*, *Arzneim.-Forsch.*, 1994, **44**, 1223 (pharmacol)

Kashiwaba, N. *et al.*, *Chem. Pharm. Bull.*, 1994, **47**, 2452 (Cephamonine, Cephamuline)

Tai, Z. *et al.*, *Drugs of the Future*, 1998, **23**, 45-49 (rev)


Bao, G.-H. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1128-1130 (pmr, cmr, N-oxide, N-demethyl)

Jin, H.-Z. *et al.*, *J. Nat. Prod.*, 2008, **71**, 127-129 (1,1'-Bisinomenine, 2,2'-Bisinomenine)

Sinomontanine I

S-314

[872977-67-8]



$\text{C}_{34}\text{H}_{44}\text{N}_2\text{O}_{10}$ 640.729


Alkaloid from the roots of *Aconitum sinomontanum*.

Peng, C.-S. *et al.*, *Youji Huaxue*, 2005, **25**, 1235-1239; *CA*, **144**, 103949n

Sinomontanine C

S-315

[433717-27-2]



$\text{C}_{34}\text{H}_{44}\text{N}_2\text{O}_9$ 624.73

Alkaloid from the roots of *Aconitum sinomontanum*. Amorph. powder. $[\alpha]_D$ +21.8 (c, 0.5 in CHCl_3).

7-Hydroxy: **Sinomontanine I**

$\text{C}_{34}\text{H}_{44}\text{N}_2\text{O}_{10}$ 640.729

Alkaloid from *Aconitum sinomontanum*. $[\alpha]_D^{17}$ -9.8 (c, 0.5 in CHCl_3).


Wang, F.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 15-22 (*Sinomontanine C*)

Peng, C.-S. *et al.*, *Youji Huaxue*, 2005, **25**, 1235-1239 (*Sinomontanine I*)

Sinomontanitine A

S-316

[433717-24-9]



$\text{C}_{35}\text{H}_{44}\text{N}_2\text{O}_9$ 636.741

Alkaloid from the roots of *Aconitum sinomontanum*. Foam. $[\alpha]_D$ +24.2 (c, 1 in CHCl_3).

1-Me ether: **Sinomontanitine B**

[433717-25-0]

$\text{C}_{36}\text{H}_{46}\text{N}_2\text{O}_9$ 650.767

Alkaloid from the roots of *Aconitum sinomontanum*. Amorph. powder. $[\alpha]_D$ +1 (c, 0.5 in CHCl_3).

Wang, F.-P. *et al.*, *J. Asian Nat. Prod. Res.*, 2001, **3**, 15-22 (isol, pmr, cmr)

Sinostemonine

S-317

$\text{C}_{21}\text{H}_{36}\text{NO}_5$ 382.519

Struct. unknown. Constit. of the Chinese drug pai pu, stated to be derived from an

unidentified *Stemona* sp. (Stemonaceae). Needles (H_2O). Mp 138-138.5°. $[\alpha]_D^{25}$ -37 (H_2O).


Lee, H.M. *et al.*, *J. Am. Pharm. Assoc.*, 1940, **29**, 391-394 (isol, pharmacol)

Sinulamide

S-318

[122027-56-9]

[221278-53-1, 122027-57-0]



$\text{C}_{34}\text{H}_{67}\text{N}_4\text{O}^{2\oplus}$ 547.93

Alkaloid from a soft coral *Sinularia* sp. ATP-ase inhibitor. Cytotoxic agent. Powder (as dichloride, hydrochloride).


Japan. Pat., 1988, 88 307 849; *CA*, **111**, 102689j

Sata, N.U. *et al.*, *Tet. Lett.*, 1999, **40**, 719-722 (isol, synth, pmr, cmr)

Sinulamine II

S-319

[139579-28-5]



$\text{C}_{24}\text{H}_{37}\text{NO}_5$ 419.56

Constit. of a *Sinularia* coral. Needles. Mp 147-148°. $[\alpha]_D$ +40 (c, 0.1 in CHCl_3).

O^{13} -De-Ac, 13-ketone: **Sinulamine I**

[139579-27-4]

$\text{C}_{22}\text{H}_{33}\text{NO}_4$ 375.507

Constit. of a *Sinularia* coral. Needles. Mp 139-140°. $[\alpha]_D$ +34 (c, 0.4 in CHCl_3).


Iguchi, K. *et al.*, *Chem. Lett.*, 1992, 127 (isol, pmr, cmr)

Kobayashi, M. *et al.*, *J. Chem. Res., Synop.*, 1992, 340 (struct)

Sinulamine III

S-320

[139579-29-6]



$\text{C}_{24}\text{H}_{37}\text{NO}_5$ 419.56

Constit. of a *Sinularia* coral. Pale yellow viscous oil. $[\alpha]_D$ -57.7 (c, 1.02 in CHCl_3).


Iguchi, K. *et al.*, *Chem. Lett.*, 1992, 127 (isol, pmr, cmr)

Kobayashi, M. *et al.*, *J. Chem. Res., Synop.*, 1992, 340 (struct)

Siomycin A, 8CI

S-321

1-Valine-2-(2,3-didehydroalanine)thio-strepton, 9CI. Mutabilysin. *Sporangiomyces*. Antibiotic 6741-21. Mutabilicin [12656-09-6]



$C_{71}H_{81}N_{19}O_{18}S_5$ 1648.867

Depsideptide antibiotic. Close homologue of Thiostrepton, T-392. Metab. of *Streptomyces sioyaensis* and *Streptomyces mutabilis-thiostreptoni*. Active against gram-positive bacteria. Cryst. (MeOH/dioxan). Mp 255-260° dec. $[\alpha]_D^{23}$ -90.9 (c, 1 in dioxan). λ_{max} 240 (sh); 245 (ε 50000); 280 (sh); 294 (ε 18000); 305 (sh); 306 (ε 14900) (MeOH) (Derep).

7'-Parent amide: Siomycin B

[61419-19-0]

 $C_{65}H_{75}N_{17}O_{16}S_5$ 1510.742

Cyclic depsipeptide antibiotic. Metab. of *Streptomyces sioyaensis*. Active against gram-positive bacteria. Cryst. (MeOH/CHCl₃). Mp 255-260° dec. $[\alpha]_D^{23}$ -102.9 (c, 0.5 in dioxan). Probably derived from Siomycin A, S-321. λ_{max} 245 (sh) (ε 60000); 290 (ε 20000) (MeOH) (Derep).

1'-Parent acid, Me ester: Siomycin C

[75524-62-8]

 $C_{72}H_{82}N_{18}O_{19}S_5$ 1663.879

Metab. of *Streptomyces sioyaensis*. Active against gram-positive organisms and mycobacteria. Cryst. (MeOH/CHCl₃). Sol. CHCl₃, dioxan, DMF; fairly sol. Me₂CO, MeOH; poorly sol. H₂O, EtOAc, hexane, bases, acids. Mp 255-260° dec. $[\alpha]_D^{23}$ -84.5 (c, 0.5 in dioxan). λ_{max} 241; 255 (MeOH) (Berdy).

49S,49'-Dihydro, 26-thione, 4'-parent**amide: Thiopeptin A_{4b}**

[70591-35-4]

 $C_{68}H_{80}N_{18}O_{16}S_6$ 1597.887

From *Streptomyces tateyamensis*. Active against gram-positive bacteria. Sol. CHCl₃; poorly sol. H₂O.

49S,49'-Dihydro, 26-thione, 7'-parent**amide: Thiopeptin A_{3b}**

[70788-67-9]

 $C_{65}H_{77}N_{17}O_{15}S_6$ 1528.824

From *Streptomyces tateyamensis*. Active against gram-positive bacteria. Sol. CHCl₃; poorly sol. H₂O.

49S,49'-Dihydro, 26-thione, 1'-parent acid:**Thiopeptin B_b**

[70591-33-2]

 $C_{71}H_{82}N_{18}O_{18}S_6$ 1667.935

Prod. by *Streptomyces tateyamensis* ATCC21389. Active against gram-

positive bacteria.

49S,49'-Dihydro, 26-thione, 1'-parent acid,**Me ester: Thiopeptin A_{1b}**

[70591-34-3]

 $C_{72}H_{84}N_{18}O_{18}S_6$ 1681.961

Prod. by *Streptomyces tateyamensis* ATCC21389. Active against gram-positive bacteria. Sol. CHCl₃; poorly sol. H₂O. λ_{max} 250 (E1%/1cm 335) (MeOH) (Berdy).

1,2R,49S,49'-Tetrahydro, 26-thione, 4'-parent**amide: Thiopeptin A_{4a}**

[70591-37-6]

[11075-34-6]

 $C_{68}H_{82}N_{18}O_{16}S_6$ 1599.903

Depsideptide antibiotic. Isol. from *Streptomyces tateyamensis*. Active against gram-positive organisms. Cryst. Sol. CHCl₃, DMSO, DMF, dioxan; fairly sol. MeOH, EtOAc, Me₂CO; poorly sol. C₆H₆, H₂O, hexane. Mp ca. 200° (dec. browns). $[\alpha]_D^{23}$ -81.5 (c, 1 in CHCl₃). Data refers to mixt. of A_{4a} and A_{4b}. λ_{max} 242; 290; 305 (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 1000 mg/kg.**1,2R,49S,49'-Tetrahydro, 26-thione, 7'-parent****amide: Thiopeptin A_{3a}**

[70591-38-7]

 $C_{65}H_{79}N_{17}O_{15}S_6$ 1530.84

Isol. from *Streptomyces tateyamensis* ATCC21389. Active against gram-positive organisms. Fine needles. Sol. DMSO, CHCl₃, dioxan, THF; fairly sol. EtOAc, Me₂CO; poorly sol. C₆H₆, H₂O, hexane. Mp 200° dec. approx. $[\alpha]_D^{23}$ -10.8 (c, 1 in CHCl₃). Data refers to a mixt. of A_{3a} and A_{3b}. λ_{max} 245 (E1%/1cm 410); 292 (E1%/1cm 160); 306 (E1%/1cm 155) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 250 mg/kg.**1,2R,49S,49'-Tetrahydro, 26-thione, 1'-parent****acid: Thiopeptin B_a. Thiofeed**

[70606-90-5]

[12789-58-1]

 $C_{71}H_{84}N_{18}O_{18}S_6$ 1669.95

Cyclic depsipeptide antibiotic. Isol. from *Streptomyces tateyamensis* ATCC21389. Used as a feed additive. Active against gram-positive bacteria. Faintly yellow cryst. Sol. DMF, dioxan, Py, THF, CHCl₃, DMSO; fairly sol. MeOH, Me₂CO, EtOAc; poorly sol. C₆H₆, H₂O, hexane. Mp 219-222°. $[\alpha]_D^{23}$ -80 (c, 1 in CHCl₃). λ_{max} 248 (E1%/1cm 430); 250 (E1%/1cm 327) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 40-80 mg/kg, LD₅₀ (mus, scu) 3600 mg/kg.**1,2R,49S,49'-Tetrahydro, 26-thione, 1'-parent****acid, Me ester: Thiopeptin A_{1a}**

[70591-36-5]

 $C_{72}H_{86}N_{18}O_{18}S_6$ 1683.977

Prod. by *Streptomyces tateyamensis* ATCC21389. Active against gram-positive bacteria. Yellow plates (EtOAc). Sol. DMSO, DMF, CHCl₃, dioxan, Py; fairly sol. MeOH, Me₂CO, EtOAc; poorly sol. C₆H₆, hexane, H₂O. Mp ca. 200° (browns). $[\alpha]_D$ -71 (CHCl₃). Data refers to inseparable mixt. of Thiopeptins A_{1a} and A₁. λ_{max} 295; 305

(MeOH) (Berdy).

► LD₅₀ (mus, ipr) 500 mg/kg.

Ebata, M. et al., *J. Antibiot., Ser. A*, 1969, **22**, 364-368; 423-433; 434-441 (isol, uv)
Japan. Pat., 1969, 70 05 034; *CA*, **72**, 109825k (isol, manu)
Miyairi, N. et al., *J. Antibiot.*, 1970, **23**, 113-119 (*Thiopeptins*, isol)
Miyairi, N. et al., *Antimicrob. Agents Chemother.*, 1972, **1**, 192-196 (*Thiopeptins*)
Oleskar, A. et al., *Chem. Comm.*, 1978, 577-578 (struct, nmr)
Motoki, Y. et al., *Pept. Chem.*, 1979, **17**, 13; 19
Tokura, K. et al., *J. Antibiot.*, 1980, **33**, 1563-1567 (pmr, cmr, struct)
Tori, K. et al., *J. Antibiot.*, 1981, **34**, 124-129 (pmr, cmr, struct)
Hensens, O.D. et al., *J. Antibiot.*, 1983, **36**, 799-813; 814-831; 832-845 (*Thiopeptins*)
Mori, T. et al., *Chem. Asian J.*, 2008, **3**, 984-1012; 1013-1025 (synth)

Siomycin D₁

S-322

[61419-61-2]

As Siomycin A, S-321 with

R = H


 $C_{70}H_{79}N_{19}O_{18}S_5$ 1634.841

Cyclic depsipeptide antibiotic. Isol. from *Streptomyces sioyaensis*. Active against gram-positive bacteria. Cryst. Sol. CHCl₃, DMF. Mp ca. 260° (dec.). $[\alpha]_D$ -69.9 (dioxan). λ_{max} 205; 250; 285 (EtOH).

Tokura, K. et al., *J. Antibiot.*, 1980, **33**, 1563-1567 (struct)

Siphonazole

S-323

 $C_{25}H_{25}N_3O_6$ 463.489

Isol. from *Herpetosiphon* sp. Amorph. solid. λ_{max} 225 (log ε 4.54); 334 (log ε 4.11) (MeOH).


Nett, M. et al., *Angew. Chem., Int. Ed.*, 2006, **45**, 3863-3867 (isol, pmr, cmr, ms)

Linder, J. et al., *Org. Biomol. Chem.*, 2008, **6**, 3908-3916 (synth)

Siphonodictidine

S-324

[88316-91-0]

 $C_{16}H_{25}N_3O$ 275.393

Constit. of the sponge *Siphonodictyon* sp. Sol. MeOH, EtOAc.

Sullivan, B. et al., *Science (Washington, D.C.)*, 1983, **221**, 1175-1176 (isol, pmr, cmr, ms)


Jefford, C.W. et al., *Synlett*, 1990, 745-746 (synth)

Jefford, C.W. et al., *Gazz. Chim. Ital.*, 1993, **123**, 317-320 (synth)

Jefford, C.W. et al., *J. Nat. Prod.*, 2004, **67**, 1383-1386 (synth)

Sitsirikine

[1245-00-7]

Absolute
ConfigurationC₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Catharanthus roseus*, the seeds of *Aspidosperma oblongum*, leaves of *Rauwolfia caffra* and from *Strychnos pungens*. Mp 206–209° (218–220°). [α]_D²⁵ -58 (MeOH) (synthetic). [α]_D +23 (MeOH) (natural). λ_{max} 226 (log ε 4.56); 282 (log ε 3.9); 290 (log ε 3.84) (MeOH).

Picrate: Mp 226–228° dec.

Ac: Mp 198°. [α]_D²⁶ -26 (MeOH).

18,19-Dihydro: (16R)-Dihydrositsirikine

[6519-26-2]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from *Catharanthus roseus*, *Pausinystalia yohimbe* and *Aspidosperma maregravianum* (Rubiaceae, Apocynaceae). Cryst. (MeOH aq.). Mp 215°. [α]_D²⁶ -55 (MeOH). Forms solvated cryst. from Me₂CO, Mp 180°. λ_{max} 226 (log ε 4.61); 282 (log ε 3.95); 290 (log ε 3.87) (MeOH).

18,19-Dihydro, picrate:

Amber prisms (MeOH). Mp 228–230° dec.

18,19-Dihydro, Ac:

Needles (Me₂CO/petrol). Mp 187°. [α]_D²⁶ -31 (MeOH).

3,4,5,6-Tetrahydro: 3,4,5,6-Tetrahydrositsirikine

[88607-63-0]

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). [α]_D -167 (c, 0.56 in MeOH). Anhydronium base.

A^{19,20}-Isomer: see Isositsirikine, I-319

10-Methoxy: 10-Methoxysitsirikine

[88607-59-4]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. [α]_D -9 (c, 0.54 in MeOH).

16-Epimer: 16-Epitsirikine. (16S)-Sitsirikine

[68346-08-7]

Alkaloid from the seeds of *Aspidosperma album* (Apocynaceae). Mp 205–207°. [α]_D²⁵ -22 (MeOH). [α]_D +218 (MeOH).

16-Epimer, 18,19-dihydro: (16S)-Dihydrositsirikine

[72401-66-2]

C₂₁H₂₈N₂O₃ 356.464

Alkaloid from the root bark of *Aspidosperma maregravianum* (Apocynaceae). Amorph. [α]_D -14 (c, 0.35 in MeOH). λ_{max} 226 (log ε 4.82); 274 (log ε 3.89); 284 (log ε 3.91); 291 (log ε 3.65) (EtOH).

S-325

16-Epimer, 10,11-dimethoxy, 18,19-dihydro: 10,11-Dimethoxy-18,19-dihydro-16S-sitsirikine

[77795-00-7]

C₂₃H₃₂N₂O₅ 416.516

Alkaloid from the trunk bark of *Ochrosia moorei* and *Ochrosia glomerata* (Apocynaceae). λ_{max} 227; 297 (EtOH).

Van der Meulen, Th.H. et al., *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1964, **83**, 148–153; 154–166 (*Dihydrositsirikine*)

Kutney, J.P. et al., *Tetrahedron*, 1966, **22**, 321–336 (*Dihydrositsirikine, isol, uv, ir, pmr, ms, struct*)

Fujii, T. et al., *Heterocycles*, 1977, **7**, 149–154 (*Dihydrositsirikine, synth*)

Stöckigt, J. et al., *Planta Med.*, 1978, **33**, 188–192 (*config, uv, ir, pmr, ms*)

Brown, R.T. et al., *Tet. Lett.*, 1979, 1805–1808 (*config*)

Ahond, A. et al., *J. Nat. Prod.*, 1981, **44**, 193–199 (*10,11-Dimethoxy-16S-dihydrositsirikine*)

Kohl, W. et al., *Z. Naturforsch., B*, 1982, **37**, 1346–1351 (*Dihydrositsirikine*)

Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 694–707; 708–722 (*16R- and 16S-Dihydrositsirikine, 10-Methoxysitsirikine, Tetrahydrositsirikine*)


Seguin, E. et al., *J. Nat. Prod.*, 1984, **47**, 687–691 (*10,11-Dimethoxy-16S-dihydrositsirikine*)

Thépenier, P. et al., *Phytochemistry*, 1990, **29**, 2384–2386 (*isol*)

Skytanthine

S-326

Octahydro-2,4,7-trimethyl-1H-cyclopenta[c]pyridine



(4R,4aR,7S,7aR)-form

C₁₁H₂₁N 167.294

►Tremorigenic agent of low toxicity.

(4R,4aR,7S,7aR)-form

α-Skytanthine

[2065-32-9]

Minor alkaloid of *Skytanthus acutus* (Apocynaceae). Liq. Mp 120° (as picrate). [α]_D +79.

4,4a-Didehydro: Δ⁴-Dehydroskytanthine.

4,4a-Didehydroskytanthine

C₁₁H₁₉N 165.278

Alkaloid from *Skytanthus acutus* (Apocynaceae). Also obt. by treating 4-Hydroxyskytanthine, H-735 with SOCl₂. Mp 127° (as picrate).

(4S,4aR,7S,7aS)-form

β-Skytanthine

[24282-31-3]

Major component of the skytanthine mixt. from *Skytanthus acutus* (Apocynaceae). Liq. Mp 135° (as picrate). Bp₁₀ 88°. [α]_D +16.4.

N-Oxide: β-Skytanthine N-oxide

[25499-86-9]

C₁₁H₂₁NO 183.293

Alkaloid from *Skytanthus acutus* (Apocynaceae). Cryst. + 2H₂O (C₆H₆/

petrol). Mp 218–222° dec. [α]_D 0 (c, 1 in MeOH).

(4R,4aR,7S,7aS)-form

γ-Skytanthine

[6545-32-0]

Synthetic. Liq. Mp 162° (as picrate). [α]_D +59.

(4S,4aR,7S,7aR)-form

δ-Skytanthine

[2883-89-8]

Minor alkaloid of *Skytanthus acutus* (Apocynaceae). Also isol. from *Tecoma stans* (Bignoniaceae). Mp 144–146° (139°) (as picrate). [α]_D +9.

4a,5-Didehydro: Δ^{4a}-Dehydroskytanthine.

4a,5-Didehydroskytanthine

C₁₁H₁₉N 165.278

Alkaloid from *Tecoma stans* (Bignoniaceae). Mp 167° (as picrate). [α]_D²⁶ -89 (c, 0.72 in CHCl₃). Catalytic hydrogenation yields δ-skytanthine.

(4ξ,4aξ,7ξ,7aξ)-form

N-De-Me: N-Normethylskytanthine

[22324-93-2]

C₁₀H₁₉N 153.267

Alkaloid from *Tecoma stans* (Bignoniaceae). Mp 179–180° (as picrate). Bp₁₃ 125–130°. [α]_D²² +35.

Eisenbraun, E.J. et al., *Chem. Ind. (London)*, 1962, 1242 (*isol, struct*)

Casinovi, C.G. et al., *Gazz. Chim. Ital.*, 1962, **92**, 479 (*synth*)

Marini-Bettolo, G.B. et al., *Gazz. Chim. Ital.*, 1963, **93**, 1367 (*Δ⁴-Dehydroskytanthine*)

Adolphsen, G. et al., *Tetrahedron*, 1967, **23**, 3147 (*Δ⁴-Dehydroskytanthine*)

Dickinson, E.M. et al., *Tetrahedron*, 1969, **25**, 1523 (*N-Normethylskytanthine*)

Eisenbraun, E.J. et al., *J.O.C.*, 1970, **35**, 1364 (*ms*)

Gross, D. et al., *Phytochemistry*, 1973, **12**, 201 (*Skytanthine, Δ^{4a}-Dehydroskytanthine*)

Oppolzer, W. et al., *Tet. Lett.*, 1986, **27**, 1141 (*synth*)

Tsunoda, T. et al., *Tet. Lett.*, 1996, **37**, 2463 (*synth*)

Ernst, M. et al., *Synthesis*, 2002, 1953–1955 (*α-Skytanthine, synth*)


Schöllhorn, B. et al., *Eur. J. Org. Chem.*, 2006, 901–908 (*synth*)

Kaneda, K. et al., *Tetrahedron*, 2008, **64**, 11589–11593 (*α-Skytanthine, synth*)

Slaframine

S-327

6-Aminoctahydro-1-indolizolinol acetate, 9CI. Alkaloid RL-A



(–)-form

C₁₀H₁₈N₂O₂ 198.264

Sol. MeOH, EtOAc.

►Toxic.

(–)-form [20084-93-9]

Alkaloid toxin from the fungus *Rhizoctonia leguminicola*. Strong parasymptomatic agent. Cause of 'slobbers' disease (excessive salivation) in cattle feeding on *Rhizoctonia leguminicola*. Toxicity requires metabolic activation.

Has been suggested as therapeutic agent for cystic fibrosis and similar diseases. Mp 183-184° (as dipicrate). $[\alpha]_D^{25}$ -38 (c, 0.16 in CHCl_3) (synthetic).

N-Ac:

Mp 140-142°. $[\alpha]_D^{25}$ -15.9 (c, 5 in EtOH).

6-Epimer:

Synthetic. $[\alpha]_D^{22}$ -11.6 (c, 0.37 in EtOH).

1,8a-Diepimer:

Synthetic. $[\alpha]_D^{25}$ +20 (c, 0.5 in EtOH).

(±)-form [30591-15-2]

Synthetic. Oil; extremely hygroscopic solid as dihydrochloride.

Dipicrate:

Yellow needles (20% EtOH aq.). Mp 215-221° dec.

N-Ac:

Cryst. (EtOH/hexane). Mp 143-146°.

6-Epimer: [114817-63-9]

Synthetic. Oil.

6-Epimer, N-Ac: [114817-64-0]

Cryst. by subl. Mp 203°.

Rainey, D.P. et al., *Nature (London)*, 1965, **205**, 203 (isol)

Aust, S.D. et al., *Nature (London)*, 1965, **205**, 204 (isol)

Whitlock, B.J. et al., *Tet. Lett.*, 1966, 3819 (isol, ms)

Gardiner, R.A. et al., *J.A.C.S.*, 1968, **90**, 5639 (ms, struct, abs config)

Cartwright, D. et al., *J.A.C.S.*, 1970, **92**, 7615 (synth)

Gensler, W.J. et al., *J.O.C.*, 1973, **38**, 3848 (synth)

Clevenstine, E.C. et al., *Biochemistry*, 1979, **18**, 3658; 3663 (biosynth)

Gobao, R.A. et al., *J.A.C.S.*, 1982, **104**, 7065 (synth, ir, pmr, ms)

Schneider, M.J. et al., *J.O.C.*, 1984, **49**, 3681 (synth, ir, pmr, cmr, ms)

Howard, A.S. et al., *Alkaloids (Academic Press)*, 1986, **28**, 269 (pharmacol)

Elbein, A.D. et al., *Alkaloids: Chem. Biol. Perspect.*, 1987, **5**, 1 (rev)

Dartmann, M. et al., *Annalen*, 1988, 695 (synth, ir, pmr, cmr, ms)

Pearson, W.H. et al., *J.O.C.*, 1991, **56**, 1976; 1992, **57**, 3977 (synth, abs config)

Choi, J.-R. et al., *Tet. Lett.*, 1991, **32**, 6469 (synth)

Sibi, M.P. et al., *J.O.C.*, 1992, **57**, 4329 (synth)

Knapp, S. et al., *J.O.C.*, 1992, **57**, 4802 (synth)

Hua, D.H. et al., *J.O.C.*, 1993, **58**, 2144 (synth)

Knight, D.W. et al., *J.C.S. Perkin 1*, 1997, 2179-2187 (synth)

Kang, S.H. et al., *Tet. Lett.*, 1998, **39**, 9047-9050 (synth)

Sibi, M.P. et al., *J.O.C.*, 1999, **64**, 6434-6443 (synth)

Comins, D.L. et al., *Org. Lett.*, 1999, **1**, 1941-1943 (synth)

Pourashraf, M. et al., *J.O.C.*, 2000, **65**, 6966-6972 (synth)


Cossy, J. et al., *Synthesis*, 2002, 951-957 (synth, ir, pmr, cmr, ms)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 912

Slagenine A

S-328

[246136-98-1]



Absolute Configuration

$\text{C}_{11}\text{H}_{13}\text{BrN}_4\text{O}_4$ 345.152

Alkaloid from the sponge *Agelas nakamurai*. Cytotoxic. Amorph. solid. $[\alpha]_D^{27}$ +11 (c, 1.2 in MeOH). λ_{max} 270 (ε 10500) (MeOH).

11-Me ether: **Slagenine B**

[246137-00-8]

$\text{C}_{12}\text{H}_{15}\text{BrN}_4\text{O}_4$ 359.179

Alkaloid from *Agelas nakamurai*. Cytotoxic agent. Amorph. solid. $[\alpha]_D^{26}$ +33 (c, 0.2 in MeOH). λ_{max} 269 (ε 9000) (MeOH).

11,15-Diepimer, 11-Me ether: **Slagenine C**

[246137-03-1]

$\text{C}_{12}\text{H}_{15}\text{BrN}_4\text{O}_4$ 359.179

Alkaloid from *Agelas nakamurai*. Cytotoxic agent. Amorph. solid. $[\alpha]_D^{25}$ -35 (c, 0.2 in MeOH). λ_{max} 269 (ε 9000) (MeOH).

Tsuda, M. et al., *Tet. Lett.*, 1999, **40**, 5709-

5712 (isol, ir, uv, pmr, cmr)


Gurjar, M.K. et al., *Org. Lett.*, 2002, **4**, 3569-3570 (synth)

Jiang, B. et al., *J.O.C.*, 2003, **68**, 2376-2384 (synth, abs config)

Smenospongine

S-329

[113021-53-7]



Absolute Configuration

$\text{C}_{21}\text{H}_{29}\text{NO}_3$ 343.465

Quinoterpenoid antibiotic. Prod. by the sponges *Smenospongia* sp., *Dactylospongia elegans* and *Petrosaspongia metachromia*. Cytotoxic and antimicrobial agent. Red cryst. Sol. MeOH, CHCl_3 ; poorly sol. H_2O . Mp 153-155°. λ_{max} 209 (ε 16200); 317 (ε 15600) (EtOH) (Derrep).

N-(2-Methylpropyl): **Smenospongiorine**

[121994-53-4]

$\text{C}_{25}\text{H}_{37}\text{NO}_3$ 399.572

Constit. of *Smenospongia* sp. and *Dactylospongia elegans*. Antimicrobial and cytotoxic agent. Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. λ_{max} 210 (ε 14000); 329 (ε 20200) (EtOH) (Derrep).

N-Pentyl: **Pentylsmenospongine. Sesquiterpenylamylaminohydroxyquinone**

[127524-59-8]

$\text{C}_{26}\text{H}_{39}\text{NO}_3$ 413.599

Isol. from a *Smenospongia* sp. Red needles (EtOH aq.). Mp 131-134°. No stereochem. indicated. Presumed here to be a Smenospongine deriv. Named by the authors as Sesquiterpenylamylaminohydroxyquinone.

N-(3-Methylbutyl): **Smenospongiarine**

[121994-52-3]

$\text{C}_{26}\text{H}_{39}\text{NO}_3$ 413.599

Constit. of *Smenospongia* sp. and *Dactylospongia elegans*. Antimicrobial and cytotoxic agent. Cryst. Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. Mp 170-172°. λ_{max} 210 (ε 14000); 329 (ε 20200) (EtOH) (Derrep). λ_{max} 210 ; 325 (MeOH) (Berdy). λ_{max} 204 (ε 27230); 324 (ε 14700) (EtOH) (Berdy).

N-(2-Phenylethyl): **Smenospongidine**

[121994-51-2]

$\text{C}_{29}\text{H}_{37}\text{NO}_3$ 447.616

Constit. of *Smenospongia* sp. and *Dactylospongia elegans*. Antimicrobial and cytotoxic agent. Cryst. (MeOH). Sol. MeOH, CHCl_3 ; poorly sol. H_2O , hexane. Mp 168-170°. λ_{max} 207 (ε 25000); 327 (ε 17000) (EtOH) (Derrep). λ_{max} 210 ; 330 (MeOH) (Berdy).

N-(Carboxymethyl): **Glycinyllimaquinone**

[159240-55-8]

$\text{C}_{23}\text{H}_{31}\text{NO}_5$ 401.502

Constit. of a *Fasciospongia* sp. Amorph. red powder. λ_{max} 203 (ε 23000); 320 (ε 11000); 454 (ε 1900); 492 (ε 2000) (MeOH) (Berdy).

5-Epimer: **5-Epismenospongine**

[287967-28-6]

$\text{C}_{21}\text{H}_{29}\text{NO}_3$ 343.465

Constit. of *Petrosaspongia metachromia*. Purple powder. $[\alpha]_D^{25}$ +73.1 (c, 0.03 in CHCl_3).

5-Epimer, N-(3-methylbutyl): **5-Epismenospongiarine**

$\text{C}_{26}\text{H}_{39}\text{NO}_3$ 413.599

Constit. of *Dactylospongia elegans*. Oil. $[\alpha]_D^{25}$ +96.7 (c, 0.12 in CHCl_3). CAS no. not found 9-14Cl.

5-Epimer, N-(2-phenylethyl): **5-Epismenospongidine**

[144335-13-7]

$\text{C}_{29}\text{H}_{37}\text{NO}_3$ 447.616

Constit. of *Dactylospongia elegans*. Oil. $[\alpha]_D^{25}$ +37.5 (c, 0.16 in CHCl_3).

5-Epimer, N-(2-methylpropyl): **5-Epismenospongiorine**

[764648-37-5]

$\text{C}_{25}\text{H}_{37}\text{NO}_3$ 399.572

Constit. of *Dactylospongia elegans*. $[\alpha]_D^{25}$ +23 (c, 0.06 in CHCl_3). λ_{max} 205 (ε 20500); 320 (ε 9800) (no solvent reported).

Kondracki, M.-L. et al., *Tetrahedron*, 1989, **45**, 1995-2004 (isol, ir, uv, pmr, cmr)

Utkina, N.K. et al., *Khim. Prir. Soedin.*, 1990, **26**, 47-51; *Chem. Nat. Compd. (Engl. Transl.)*, 1990, **26**, 37-40

(Pentylsmenospongine)

Rodriguez, J. et al., *Tetrahedron*, 1992, **48**, 6667-6680 (isol, pmr, cmr, abs config, bibl)

Evans, T.P. et al., *Nat. Prod. Lett.*, 1994, **4**, 287-291 (Glycinyllimaquinone, isol, pmr, cmr, uv, ir, activity)

Kwak, J.H. et al., *J. Nat. Prod.*, 2000, **63**, 1153-1156 (5-Epismenospongine)


Ling, T. et al., *J.A.C.S.*, 2002, **124**, 12261-12267 (synth)

Aoki, S. et al., *Chem. Pharm. Bull.*, 2004, **52**, 935-937 (*Dactylospongia elegans* constits, bibl)

Smipine

S-330

2-(3,4-Dihydro-2H-pyrrol-5-yl)-1-piperidinecarboxaldehyde, 9CI. 2-(1-Formyl-2-piperidyl)-1-pyrroline [52196-11-9]



C₁₀H₁₆N₂O 180.249


Alkaloid from *Lupinus formosus* (Fabaceae). Lack of opt. activity prob. due to tautomerism with Δ^{2,2'}-form.

Fitch, W.L. *et al.*, *J.O.C.*, 1974, **39**, 2974-2979 (isol, uv, ir, pmr, ms, synth)

SMTP 3

S-331

[221161-84-8]



R = CH₂OH

C₂₆H₃₅NO₇ 473.565

Analogue of Staplabin in S-490. Abs. config. of C-27 as shown; rel. config. only of other centres known. Prod. by *Stachybotrys microspora*. Promotes plasminogen activation and plasminogen binding to fibrin. Light brown oil. [α]_D²⁰ +19 (c, 0.32 in Me₂CO). λ_{max} 214 (ε 43000); 256 (ε 9500); 298 (ε 3400) (MeOH).

Hasumi, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1059-1068 (isol, uv, ir, pmr, cmr)

SMTP 4

S-332

[221161-86-0]

As SMTP 3, S-331 with

R = CH₂Ph

C₃₂H₃₉NO₆ 533.663

Analogue of Staplabin in S-490. Prod. by *Stachybotrys microspora*. Promotes plasminogen activation and plasminogen binding to fibrin. Light brown oil. [α]_D²⁰ -79 (c, 0.9 in MeOH). λ_{max} 214 (ε 40700); 259 (ε 8400); 302 (ε 2500) (MeOH).

27-Epimer: SMTP 4D. Antibiotic SMTP 4D

Prod. by *Stachybotrys microspora* fed with D-Phe. Light brown oil. [α]_D²⁰ +51.4 (c, 0.9 in MeOH).

Hasumi, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1059-1068 (isol, uv, ir, pmr, cmr)

Hu, W. *et al.*, *J. Antibiot.*, 2003, **56**, 832-837 (SMTP 4D)

SMTP 5

S-333

[221161-89-3]

As SMTP 3, S-331 with

R = CH₂CH(CH₃)₂

C₂₉H₄₁NO₆ 499.646

Analogue of Staplabin in S-490. Prod. by *Stachybotrys microspora*. Promotes plasminogen activation and plasminogen binding to fibrin. Light brown oil. [α]_D²⁰ -12.3 (c, 0.5 in MeOH). λ_{max} 215 (ε 36700); 259 (ε 7600); 300 (ε 2300) (MeOH).

27-Epimer: SMTP 5D. Antibiotic SMTP 5D

Prod. by *Stachybotrys microspora* fed with D-Leu. Light brown oil. [α]_D²⁰ -9.8 (c, 0.5 in MeOH).

Hasumi, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1059-1068 (isol, uv, ir, pmr, cmr)

Hu, W. *et al.*, *J. Antibiot.*, 2003, **56**, 832-837 (SMTP 5D)

SMTP 6

S-334

[221161-92-8]

As SMTP 3, S-331 with

R = 3-indolylmethyl

C₃₄H₄₀N₂O₆ 572.7

Analogue of Staplabin in S-490. Prod. by *Stachybotrys microspora*. Promotes plasminogen activation and plasminogen binding to fibrin. Light brown oil. [α]_D²⁰ -44.6 (c, 0.5 in MeOH). λ_{max} 216 (ε 50300); 260 (ε 10100); 282 (sh) (ε 6200); 290 (sh) (ε 5400); 308 (sh) (ε 2500) (MeOH).

27-Epimer: SMTP 6D. Antibiotic SMTP 6D

Prod. by *Stachybotrys microspora* fed with D-Trp. Light brown oil. [α]_D²⁰ +19.2 (c, 0.5 in MeOH).


Hasumi, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1059-1068 (isol, uv, ir, pmr, cmr)

Hu, W. *et al.*, *J. Antibiot.*, 2003, **56**, 832-837 (SMTP 6D)

Solacallinidine

S-335

3-Amino-16,28-secosolanid-22(28)-en-16-ol, 9CI. 3-Amino-22,26-epimino-5α-cholest-22(N)-en-16-ol [61899-01-2]



C₂₇H₄₆N₂O 414.673

Isol. from dried leaf and stem of *Solanum callium* (Solanaceae). Noncryst. Mp 175-178°. [α]_D³⁵ +51.3 (c, 0.9 in CHCl₃).

Hydrochloride (1:2): [71609-90-0]

Cryst. + H₂O. Mp 310° dec. Strongly retains H₂O.

Picrate: [71609-94-4]

Minute yellow cryst. (EtOH aq.). Mp 210° dec.

O-Ac:


Pale-yellow solid. Mp 150-152°. [α]_D³⁵ -115.6 (c, 1.23 in CHCl₃).

Bird, G.J. *et al.*, *Aust. J. Chem.*, 1979, **32**, 597; 797 (isol, struct, ir, uv, pmr, cmr)

Solacapine

S-336

3-Amino-22,26-iminocholestane-16,23-diol [63785-22-8]



C₂₇H₄₈N₂O₂ 432.688

Alkaloid from *Solanum pseudocapsicum* (Solanaceae). Fine needles (MeOH/CHCl₃). Mp 286-288°. [α]_D +47.1 (c, 0.68 in CHCl₃/MeOH 1:1).

N-Tri-Me: Mp 228-230°.

22-Epimer: **Isosolacapine** [90762-21-3]

C₂₇H₄₈N₂O₂ 432.688

Alkaloid from *Solanum pseudocapsicum* (Solanaceae). Fine needles (MeOH/MeCN). Mp 238-240°. [α]_D -12.3 (c, 0.73 in CHCl₃/MeOH 1:1).

22-Epimer, N-tri-Me: Mp 200-201°.

23-Epimer: **Episolacapine** [90762-20-2]

C₂₇H₄₈N₂O₂ 432.688

Alkaloid from *Solanum pseudocapsicum* (Solanaceae). Prisms (MeOH/CHCl₃). Mp 256-258°. [α]_D -41.5 (c, 0.41 in CHCl₃/MeOH 1:1).


23-Epimer, N³,N³-di-Me: Mp 242-244°.

Chakravarty, A.K. *et al.*, *J.C.S. Perkin I*, 1984, 467 (isol, ir, pmr, struct)

Solacongostidine

S-337

16,28-Secosolanid-22(28)-en-3-ol, 9CI. 22,26-Epiminocholest-22(28)-en-3-ol [984-82-7]



C₂₇H₄₅NO 399.659

Alkaloid from *Solanum congestiflorum*, occurring as glycosides (Solanaceae). Shows antifungal activity. Rod-shaped cryst. (MeOH aq., Me₂CO or Et₂O) Mp

169-174°. $[\alpha]_D^{20} +35.6$ (c, 1.6 in CHCl_3).

Glycoside (1): Solacongестine

[57495-66-6]

$\text{C}_{39}\text{H}_{65}\text{NO}_{11}$ 723.943

Alkaloid from *Solanum congestiflorum* (Solanaceae). Conts. 1 D-glucose and 1 D-galactose.

Glycoside (2): α -Solacongестinine

[57572-75-5]

$\text{C}_{44}\text{H}_{73}\text{NO}_{14}$ 840.059

Alkaloid from *Solanum congestiflorum* (Solanaceae). Conts. 1 D-glucose, 1 D-xylose and 1 D-rhamnose.

Glycoside (3): β -Solacongестinine

[57512-69-3]

$\text{C}_{38}\text{H}_{63}\text{NO}_{10}$ 693.916

Alkaloid from *Solanum congestiflorum* (Solanaceae). Conts. 1 D-glucose and 1 D-xylose.

Ac:

Cryst. (MeOH). Mp 156-158°.

23-Oxo: 23-Oxosolacongестidine

[19374-60-8]

$\text{C}_{27}\text{H}_{43}\text{NO}_2$ 413.642

Alkaloid from *Solanum congestiflorum* (Solanaceae). Pale yellow needles. Mp 213-223° (unsharp) and >300° (part dec.).

24-Oxo: 24-Oxosolacongестidine

[19374-61-9]

$\text{C}_{27}\text{H}_{43}\text{NO}_2$ 413.642

Alkaloid from *Solanum congestiflorum* (Solanaceae). Pale yellow plates + 1 MeOH (MeOH). Mp 158-162° (anhyd.). $[\alpha]_D^{23} +40.9$ (c, 0.8 in CHCl_3).

Sato, Y. *et al.*, *J.O.C.*, 1969, **34**, 1577-1582

(*isol, ms, ir, uv, pmr, oxo-derivs*)

Kusano, G. *et al.*, *J.O.C.*, 1970, **35**, 2624-2626

(*synth*)

Katz, R. *et al.*, *CA*, 1975, **83**, 193635f

(*glycosides*)

Bird, G.J. *et al.*, *Aust. J. Chem.*, 1979, **32**, 797-

816 (*cmr*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1987,


35, 4862-4867 (*activity*)

Solafloridine

S-338

16,28-Secosolanid-22(28)-ene-3,16-diol, 9CI. 22,26-Epiminocholest-22(N)-ene-3,16-diol

[2385-18-4]



$\text{C}_{27}\text{H}_{41}\text{NO}_2$ 415.658

Alkaloid from *Solanum congestiflorum*, *Solanum umbellatum* and *Solanum verbascifolium*, occurring as glycosides (Solanaceae). Shows antifungal activity. Needles (Me_2CO , rapidly), prisms (Me_2CO , slowly). Mp 162-165° Mp 172-175°. $[\alpha]_D^{20} +123$ (c, 1.15 in CHCl_3).

Hydrochloride:

Cryst. + $1\text{H}_2\text{O}$. Mp 280-288°.

20-Epimer: 20-Isosolafloridine

[175777-77-2]

$\text{C}_{27}\text{H}_{45}\text{NO}_2$ 415.658

Alkaloid from roots of *Solanum abutiloides*. Needles (Me_2CO). Mp 136-137°. $[\alpha]_D^{25} +45.1$ (c, 0.89 in CHCl_3).

25-Epimer: 25-Isosolafloridine

[36297-66-2]

$\text{C}_{27}\text{H}_{45}\text{NO}_2$ 415.658

Alkaloid from *Solanum callium* and *Solanum aculeatum* (Solanaceae). Mp 164.5-166.5°. $[\alpha]_D^{25} +44.8$ (c, 0.85 in CHCl_3).

► VS3600000

25-Epimer, hydrochloride: Mp 325-330° dec.

Sato, Y. *et al.*, *J.O.C.*, 1969, **34**, 1577 (*ir, uv, pmr, ms*)

Kusano, G. *et al.*, *J.O.C.*, 1970, **35**, 2624

(*synth*)

Bird, G.J. *et al.*, *Tet. Lett.*, 1976, 3653 (*epimer*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1987,

35, 4862-4867 (*activity*)


Ripperger, H. *et al.*, *Phytochemistry*, 1996, **41**,

1629 (*20-Isosolafloridine*)

Solamaladine

S-339

21-(3,4-Dihydro-3-methyl-2H-pyrrol-5-yl)-3-hydroxy-20-methylpregnane-4,21-dione, 9CI. 23,26-Epimino-3-hydroxycholest-23(N)-ene-4,22-dione [51059-59-7]



$\text{C}_{27}\text{H}_{41}\text{NO}_3$ 427.626

Alkaloid *isol.* from green fruits of *Solanum hypomalacophyllum* (Solanaceae).

Mp 178-180°. $[\alpha]_D +39.5$ (EtOH).

Ac: Mp 202-208°. $[\alpha]_D -69$ (EtOH).

Usubillaga, A. *et al.*, *Rev. Latinoam. Quim.*,

1973, **4**, 32; *CA*, **79**, 123632w (*isol, ms, pmr, ir*)

Usubillaga, A. *et al.*, *Acta Cryst. B*, 1982, **38**,

966 (*cryst struct*)

Usubillaga, A. *et al.*, *J. Nat. Prod.*, 1984, **47**,

52 (*isol*)

Solamine

S-340

N'-[4-(Dimethylamino)butyl]-N,N-dimethyl-1,4-butanediamine, 9CI. 4,4'-Bis(dimethylamino)dibutylamine, 8CI. N^o-Tetramethylhomospermine [17232-87-0]

$\text{Me}_2\text{N}(\text{CH}_2)_4\text{NH}(\text{CH}_2)_4\text{NMe}_2$

$\text{C}_{12}\text{H}_{29}\text{N}_3$ 215.381

Constit. of the roots of *Solanum carolinense* and *Cyphomandra betacea* (Solanaceae). Bp₁₀ 80-90° (bath).

Trihydrochloride:

Small prisms (EtOH/Et₂O). Mp 262°

dec.

Tripicrate:

Shining plates (EtOH aq.). Mp 113-114°.

N-Ac:

Oil; needles (as methiodide). Mp 241-243° (methiodide).

N-Ethoxycarbonyl: Solaurethine

[65427-96-5]

$\text{C}_{15}\text{H}_{33}\text{N}_3\text{O}_2$ 287.445

Alkaloid from the roots of *Solanum carolinense* (Solanaceae). Rosettes (EtOH aq.) (as picrate). Mp 122° (picrate).

N-Hexanoyl: Solacaproine. N,N-Bis[4-(dimethylamino)butyl]hexanamide. N-Hexanoylsolamine

[35771-90-5]

$\text{C}_{18}\text{H}_{39}\text{N}_3\text{O}$ 313.526

Alkaloid from the roots of *Cyphomandra betacea* (tree tomato) (Solanaceae). Gum; fine needles (EtOH) (as picrate). Mp 149-150° (picrate).

N-Heptanoyl: N-Heptanoylsolamine

$\text{C}_{19}\text{H}_{41}\text{N}_3\text{O}$ 327.552

Tentatively identified as a constit. of *Cyphomandra betacea* (tree tomato) (Solanaceae).

N-Hexadecanoyl: Solapalmitine. Hexadecanoylsolamine. NSC 123124

[17232-85-8]

$\text{C}_{28}\text{H}_{59}\text{N}_3\text{O}$ 453.794

Alkaloid from *Solanum tripartitum* (Solanaceae). Shows significant anti-neoplastic activity. Liq. Bp_{0,05} 150° (bath). Log P 7.11 (uncertain value) (calc). A mixt. of Solapalmitine and Solapalmitenine was called Solapartine.

N-(2E-Hexadecenyl): Solapalmitenine.

N-(2-Hexadecenyl)solamine. Solpalmitene. NSC 123125

[17232-86-9]

$\text{C}_{28}\text{H}_{57}\text{N}_3\text{O}$ 451.778

Alkaloid from *Solanum tripartitum* (Solanaceae). Shows significant anti-neoplastic activity. Liq. Bp_{0,08} 153° (bath). Log P 6.81 (uncertain value) (calc).

Kupchan, S.M. *et al.*, *J.O.C.*, 1969, **34**, 3888

(*Solamine, Solapalmitine, Solapalmitenine, synth, ir, pmr, ms*)

Evans, W.C. *et al.*, *J.C.S. Perkin I*, 1972, 2017

(*Solamine, Solacaproine, N-Heptanoylsolamine*)


Evans, W.C. *et al.*, *Phytochemistry*, 1977, **16**,

1859 (*Solaurethine*)

Solanapyrone L

S-341

[950731-13-2]



Relative Configuration

C₁₉H₂₅NO₃ 315.411
Prod. by an unidentified fungus. Oil. [α]_D²⁵
-53 (c, 0.04 in MeOH). Isol. as a mixt. of
E/Z-isomers.

N-(2-Hydroxyethyl): **Solanapyrone K**
[950731-12-1]

C₂₁H₂₉NO₄ 359.464

Prod. by an unidentified fungus. Oil.
[α]_D²⁵ -45 (c, 0.12 in MeOH).

Schmidt, L.E. *et al.*, *J. Nat. Prod.*, 2007, **70**,
1317-1320 (*isol*, *pmr*, *cmr*)

Solangustidine

S-342

[138464-51-4]

C₂₇H₄₃NO₂ 413.642

Not well characterised. Struct. unknown,
poss. a steroidal alkaloid such as a
stereoisomer of Spirosol-5-en-3-ol, S-456.
Plates (as hydrochloride). Mp 325° (hy-
drochloride).

Glucoside: **Solangustine**

[1401-35-0]

C₃₃H₅₃NO₇ 575.784

Isol. from *Solanum angustifolium*
(Solanaceae). Cryst. + 1H₂O. Mp 235°
dec.

Ac:

Needles. Mp 256°.


Tutin, F. *et al.*, *J.C.S.*, 1914, **105**, 559-576

3,5-Solanidadiene, 9CI

S-343

Solanthrene

[26516-51-8]




C₂₇H₄₁N 379.628

Alkaloid from *Fritillaria*
camtschaticensis, *Notholirion*
bulbuliferum (Liliaceae) and *Solanum* sp.
(Solanaceae). Mp 162-165°. Possible
artifact.

Soltys, A. *et al.*, *Ber.*, 1933, **66**, 762
Mitsuhashi, H. *et al.*, *Chem. Pharm. Bull.*,
1969, **17**, 2370 (*isol*)

3,5,6-Solanidanetriol

S-344



C₂₇H₄₅NO₃ 431.657

(3 β ,5 α ,6 β)-form [119766-93-7]

Alkaloid from the Chinese herbal drug
"Bei-mu" (dried bulbs of *Fritillaria dela-*

vayi) (Liliaceae).


Di-Ac: Mp 100-104°.

Kaneko, K. *et al.*, *Chem. Pharm. Bull.*, 1988,
36, 4700 (*isol*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Solanidine

S-345

Solanid-5-en-3-ol, 9CI. *Solatubine*. *Sola-*
nidine T
[80-78-4]



C₂₇H₄₃NO 397.643

Alkaloid from *Solanum tuberosum*,
Cestrum purpureum, *Fritillaria*
camtschaticensis, *Rhinopetalum buchari-*
cum and *Rhinopetalum stantherum*
(Solanaceae, Liliaceae). Glycosides (esp.
Solanines and Chaconine) are trace
toxic constits. of potato tubers, and
interbreeding of potatoes with wild
strains may increase their concn. or
introduce other more toxic solanidine
glycosides. Mp 219°. [α]_D -27 (CHCl₃)
(Py).

► WF0500000

Hydrochloride: Mp 335° dec.

O- β -D-Galactopyranoside: γ -**Solanine**
[511-37-5]

C₃₃H₅₃NO₆ 559.785

Alkaloid from potato spp. (*Solanum*
tuberosum and *Solanum chacoense*) and
Veratrum album ssp. *lobelianum* (Sola-
naceae, Liliaceae). Antiasthmatic,
antibronchitic, antiepileptic agent. Mp
ca. 250°. [α]_D -26 (MeOH). Log P 4.15
(uncertain value) (calc).

O- β -D-Glucopyranoside: γ -**Chaconine**
[511-36-4]

C₃₃H₅₃NO₆ 559.785

Alkaloid from potato spp. *Solanum*
tuberosum and *Solanum chacoense*
(Solanaceae). Mp 243-244°. [α]_D -40
(Py).

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 4)- β -D-
glucopyranoside]: β ₂-**Chaconine**
[469-14-7]

C₃₉H₆₃NO₁₀ 705.927

Alkaloid from *Solanum tuberosum*
(potato) and other *Solanum* spp.
Amorph. powder.

O-[β -D-Glucopyranosyl-(1 \rightarrow 3)- β -D-
glucopyranoside]: β -**Solanine**
[61877-94-9]

C₃₉H₆₃NO₁₁ 721.927

Alkaloid from potato spp. *Solanum*
wrightii, *Solanum tuberosum* and *Sola-*
num chacoense (Solanaceae). Mp 290°.
[α]_D -31 (MeOH).

O-[β -D-Glucopyranosyl-(1 \rightarrow 6)- β -D-
glucopyranoside]: **Stenantinid**
[80248-81-3]

C₃₉H₆₃NO₁₁ 721.927

Alkaloid from aerial parts of *Rhino-*
petalum stantherum (Liliaceae). Mp

269-271°. [α]_D -47.5.

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 2)- β -D-
glucopyranoside]: β ₁-**Chaconine**
[472-51-5]

C₃₉H₆₃NO₁₀ 705.927

Alkaloid from potato spp. (*Solanum*
chacoense, *Solanum commersonii*, *Sola-*
num crispum, *Solanum tuberosum*,
Solanum ajanhuiri and *Solanum steno-*
tomum, Solanaceae). Mp 287-292° dec.
[α]_D -52.5 (c, 0.9 in Py).

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 4)- β -D-
glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyra-
noside]: **Stenantine**
[80248-79-9]

C₄₅H₇₃NO₁₅ 868.069

Alkaloid from aerial parts of *Rhino-*
petalum stantherum (Liliaceae). Mp
262-264°. [α]_D +46.5. Partial hydrol \rightarrow
 β ₂ and γ -Chaconines.

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 2)-[β -D-
glucopyranosyl-(1 \rightarrow 4)]- β -D-glucopyra-
noside]: [81942-09-8]

C₄₅H₇₃NO₁₅ 868.069

Alkaloid from aerial parts of *Fritillaria*
thunbergii. Needles + 2H₂O (MeOH).
Mp 278-283° dec. [α]_D -58.4 (c, 1 in
Py).

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 2)-O-[β -
D-glucopyranosyl-(1 \rightarrow 3)]- β -D-galacto-
pyranoside]: α -**Solanine**
[20562-02-1]

C₄₅H₇₃NO₁₅ 868.069

Alkaloid from potato (*Solanum tu-*
berosum) and very many other *Sola-*
num spp. (Solanaceae). Shows
cytotoxic activity against mouse as-
cites tumour cells. Cholinesterase in-
hibitor. Responsible for the
teratogenicity of sprouting potatoes.
Poorly sol. H₂O, Et₂O, CHCl₃. Mp
286°. [α]_D -59 (Py). Log P 0.96
(uncertain value) (calc).

► Adverse human CNS and gastrointestinal
effects by ingestion (associated with
consumption of damaged or rotten po-
tatoes). LD₅₀ (rat, orl) 590 mg/kg. LD₅₀
(mus, ipr) 42 mg/kg. Exp. reprod. and
teratogenic effects. WF0250000

O-[α -L-Rhamnopyranosyl-(1 \rightarrow 2)-O-[α -
L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-
glucopyranoside]: α -**Chaconine**
[20562-03-2]

C₄₅H₇₃NO₁₄ 852.07

Alkaloid from *Solanum chacoense* and
very many other *Solanum* spp. (Sola-
naceae). Mp 243°. [α]_D -85 (Py).

► Adverse human CNS and gastrointestinal
effects by ingestion (associated with
consumption of damaged or rotten po-
tatoes). LD₅₀ (rat, ipr) 84 mg/kg. Exp.
reprod. and teratogenic effects.
FL6700000

O-[β -D-Glucopyranosyl-(1 \rightarrow 4)- β -D-xy-
lopyranosyl-(1 \rightarrow 4)]- β -D-xylopyrano-
side]: **Solacauline**

C₄₃H₆₉NO₁₄ 824.016

Alkaloid from potato spp. *Solanum*
acaule, *Solanum punae* and *Solanum*
schreiteri (Solanaceae). Mp 260-265°.
[α]_D -30 (Py).

O-[β -D-Glucopyranosyl-(1 \rightarrow 3)- β -D-
glucopyranosyl-(1 \rightarrow 4)]- α -L-rhamnopyra-

nosyl-(1 → 2)-β-D-glucopyranoside]:

Hyacinthoside

[102728-60-9]

C₅₁H₈₃NO₂₀ 1030.211

Alkaloid from the bulbs of *Notholirion hyacinthinum* (Liliaceae).

O-[β-D-Glucopyranosyl-(1 → 2)-[β-D-xylopyranosyl-(1 → 3)]-β-D-glucopyranosyl-(1 → 4)-β-D-galactopyranoside]:

Δ⁵-Demissine, 5,6-Dehydrodemissine

[195433-57-9]

C₅₀H₈₁NO₂₀ 1016.185

Alkaloid from potato (*Solanum comersonii*).

O-[β-D-Glucopyranosyl-(1 → 6)-[α-L-rhamnopyranosyl-(1 → 2)]-β-D-glucopyranosyl-(1 → 3)-β-D-glucopyranosyl-(1 → 4)-β-D-glucopyranoside]:

Neohyacinthoside

[115491-59-3]

C₅₇H₉₃NO₂₅ 1192.353

Isol. from the bulbs of *Notholirion hyacinthinum*. Powder. Mp 262-265°.

[α]_D²⁰ -25.9 (c, 0.29 in Py).

O-Ac: Mp 208°.

5,6-Dihydro: see Demissidine, D-205

27-Hydroxy: **Camtschatcanidine**

[78719-99-0]

C₂₇H₄₃NO₂ 413.642

Isol. from *Fritillaria camtschatcensis* (Liliaceae). Needles (Me₂CO). Mp 261-265°. [α]_D -19.4 (c, 0.1 in MeOH).

Clemono, G.R. *et al.*, *J.C.S.*, 1936, 1299 (*isol*)
Prelog, V. *et al.*, *Helv. Chim. Acta*, 1942, **25**, 1306; 1944, **27**, 390 (*struct*)

Schreiber, K. *et al.*, *Chem. Ber.*, 1954, **87**, 1007 (*Solacauline*)

Correia Alves, L. *et al.*, *Garcia de Orta*, 1961, **9**, 713; *CA*, **61**, 12326d (*trioside*)

Budzikiewicz, H. *et al.*, *Tetrahedron*, 1964, **20**, 2267 (*ms*)

Höhne, E. *et al.*, *Tetrahedron*, 1966, **22**, 673 (*cryst struct*)

Schreiber, K. *et al.*, *Alkaloids (Academic Press)*, 1968, **10**, 1 (*rev*)

Kessar, S.V. *et al.*, *Tetrahedron*, 1971, **27**, 2153 (*synth*)

Radeglia, R. *et al.*, *Tet. Lett.*, 1977, 903 (*cmr*)

Ripperger, H. *et al.*, *Alkaloids (Academic Press)*, 1981, **19**, 81 (*rev*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 349; *Chem. Nat. Compd. (Engl. Transl.)*, 273 (*Stenantine, Stenantidine*)

Kaneko, K. *et al.*, *Phytochemistry*, 1981, **20**, 327 (*Camtschatcanidine*)

Katajima, J. *et al.*, *Phytochemistry*, 1982, **21**, 187 (*glycosides*)

Qiu, F. *et al.*, *CA*, 1983, **98**, 104286n (*glycoside*)

Keeler, R.F. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1984, **4**, 389 (*rev, tox*)

Xu, W.H. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 177; *CA*, **105**, 21663 (*Hyacinthoside*)

Xu, W.H. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 61-63 (*Neohyacinthoside*)

Vazquez, A. *et al.*, *Euphytica*, 1997, **95**, 195-201 (*Δ⁵-Demissine*)

Lawson, D.R. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 4122-4126 (*pmr, cmr, ms*)


Abouzeid, S. *et al.*, *Nat. Prod. Res.*, 2008, **22**, 147-153 (*α-Chaconine, β-Chaconine*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, CDG500; SKS000

Solanocapsine

S-346

3-Amino-16,23-epoxy-16,28-secosolanidan-23-ol, 9CI. 3-Amino-22,26-epimino-16,23-epoxycholestan-23-ol [639-86-1]



Absolute Configuration

C₂₇H₄₆N₂O₂ 430.673

Alkaloid from *Solanum pseudocapsicum*, *Solanum capsicastrum*, *Solanum henderstonii*, *Solanum seafortianum* and *Solanum tucumanense* (Solanaceae). Shows antifungal and antiyeast activity. Antibacterial agent. On injection produces local irritation and cardiac disorganisation. Mp 222° dec. (213-215° dec.) (variable). [α]_D +25 (MeOH). Log P 4.7 (uncertain value) (calc). λ_{max} 286; 293 (EtOH) (Berdy).

▶ LD₅₀ (mus, ipr) 100 mg/kg, LD₅₀ (mus, scu) 300 mg/kg. WF0539000

Sulfate: Mp 324°.

Picrate: Mp 194° (200-201°).

Me ether: **O-Methylsolanocapsine**

[90819-92-4]

C₂₈H₄₈N₂O₂ 444.699

Alkaloid from *Solanum pseudocapsicum* (Solanaceae). Fine needles (MeOH/MeCN). [α]_D +44.4 (c, 0.63 in CHCl₃).

5,6-Didehydro: **Solanoforthine**

[64919-50-2]

C₂₇H₄₄N₂O₂ 428.657

Isol. from *Solanum seafortianum* (Solanaceae). Cryst. (C₆H₆). Mp 208-210°. [α]_D -26.6 (c, 0.6 in CHCl₃).

22,N-Didehydro, Me ether: **Solacassine**

[59957-97-0]

C₂₈H₄₆N₂O₂ 442.684

Isol. from *Solanum pseudocapsicum* flower tops (Solanaceae). Antibacterial agent. Shows antifungal and antiyeast activity. Mp 215-220° dec. [α]_D +29 (MeOH). Probable struct. OMe group may be an artifact of extraction. Drawing error in the paper.

7β-Hydroxy, Me ether: **7-Hydroxy-O-methylsolanocapsine**

[114847-23-3]

C₂₈H₄₈N₂O₃ 460.699

Minor alkaloid from leaves of *Solanum capsicastrum* (Solanaceae). Amorph. powder. [α]_D²⁴ +65.2 (c, 0.46 in CHCl₃).

Barger, G. *et al.*, *J.C.S.*, 1936, 1537 (*isol*)

Boll, P.M. *et al.*, *Antibiot. Annu.*, 1955, 255-259 (*activity*)

Budzikiewicz, H. *et al.*, *Tetrahedron*, 1964, **20**, 2267 (*ms*)

Ripperger, H. *et al.*, *Annalen*, 1969, **723**, 159 (*pmr, ir, struct*)

Höhne, E. *et al.*, *Tetrahedron*, 1970, **26**, 3569 (*cryst struct*)

Ripperger, H. *et al.*, *Tetrahedron*, 1972, **28**, 1629 (*synth*)

Mitscher, L. *et al.*, *Experientia*, 1976, **32**, 415 (*Solacassine, activity*)

Ali, E. *et al.*, *Tetrahedron*, 1977, **33**, 1371 (*isol, ms, Solanoforthine*)

Bird, G.J. *et al.*, *Aust. J. Chem.*, 1979, **32**, 797 (*cmr*)

Ripperger, H. *et al.*, *Pharmazie*, 1982, **37**, 870 (*isol*)


Chakravarty, A.K. *et al.*, *J.C.S. Perkin I*, 1984, 467 (*O-Methylsolanocapsine*)

Chakravarty, A.K. *et al.*, *Phytochemistry*, 1988, **27**, 956 (*7β-Hydroxy-O-methylsolanocapsine*)

Solanocastrine

S-347

[112156-49-7]



C₂₇H₄₂N₂O 410.642


Very minor alkaloid from the leaves of *Solanum capsicastrum* (Solanaceae). Mp 258-260° dec. [α]_D 0 (CHCl₃).

Chakravarty, A.K. *et al.*, *Tet. Lett.*, 1987, **28**, 4753 (*uv, ir, pmr, cmr, ms, struct*)

Solanudine

S-348

[118655-44-0]



C₂₇H₄₃NO₃ 429.642

Alkaloid from the green berries of *Solanum nudum* (Solanaceae). Needles. Mp 225°. [α]_D²³ +352 (c, 1.02 in dioxan).


Tri-Ac: Mp 162-165°.

Usubillaga, A. *et al.*, *Phytochemistry*, 1988, **27**, 3031-3032 (*isol, uv, ir, pmr, cmr, ms, struct*)

Solaphyllidine

S-349

16-Acetoxy-3,23-dihydroxy-16,28-secosolanidan-4-one, 9CI. Solafilidine [26520-58-1]



C₂₉H₄₇NO₅ 489.694

Alkaloid from leaves and berries of *Solanum hypomalacophyllum* and from *Solanum ecuadorensis* (Solanaceae). Cryst. (MeOH). Mp 165-170°. [α]_D²⁰ -25.4 (c, 0.887 in MeOH).

▶ TU4382400

O-De-Ac: **Deacetylolaquidine**.

3,16,23-Trihydroxy-16,28-secosolanidan-4-one, 9CI

[26520-60-5]

C₂₇H₄₅NO₄ 447.657

Alkaloid from *Solanum ecuadorensis* and *Solanum hypomalacophyllum* (Solanaceae). Mp 272-276°. [α]_D²⁰ +48.3 (c, 0.258 in MeOH).

▶ TU4381800

O,O,N-Tri-Ac:

Cryst. (propan-2-ol). Mp 204-206°. [α]_D²⁰ -25.4 (c, 0.75 in MeOH).

Usubillaga, A. *et al.*, *J.A.C.S.*, 1970, **92**, 700-701 (*isol, ms, cryst struct*)


Usubillaga, A. *et al.*, *CA*, 1973, **79**, 113186 (*isol*)

Usubillaga, A. *et al.*, *Planta Med.*, 1973, **23**, 286-289 (*isol*)

Solaquidine

S-350

3,3-Dimethoxy-16,28-secosolanidan, 9CI. 3,3-Dimethoxy-22,26-epiminocholestane [65180-55-4]



Absolute Configuration

C₂₉H₅₁NO₂ 445.727

Alkaloid from *Solanum pseudoquina* (Solanaceae). Needles (MeOH). Mp 278-281°.

Ac:

Cryst. (Me₂CO/hexane). Mp 129-134°.


Usubillaga, A. *et al.*, *Phytochemistry*, 1977, **16**, 1861-1862 (*ir, ms, pmr, struct*)

Meccia, G. *et al.*, *J. Nat. Prod.*, 1987, **50**, 642-645 (*abs config*)

Solaseaforthine

S-351

3-Amino-16,28-secosolanidan-22,24,26(28)-triene-16,23-diol [70191-12-7]



C₂₇H₄₂N₂O₂ 426.641

Alkaloid from *Solanum seaforthianum* and *Solanum giganteum* (Solanaceae).

Mp 172-178°. [α]_D +22 (MeOH). Prob. artifact.

N,N-Di-Me, O,O-di-Ac:

Glass.

20-Epimer: **Isosolaseaforthine**

[70139-40-1]

From *Solanum seaforthianum* (Solanaceae). Amorph. Mp 172-180°. [α]_D +26 (MeOH).


20-Epimer, N,N-di-Me, O,O-di-Ac: Mp 202-204°.

Ali, E. *et al.*, *Tet. Lett.*, 1978, **19**, 3871-3874 (*isol, uv, cd, pmr, struct*)

Solaspinalidine

S-352

[181884-05-9]



C₂₇H₄₁NO₃ 427.626


Alkaloid from roots of *Solanum spirale*. Needles (Me₂CO). Mp 212-214°. [α]_D²⁶ -63.1 (c, 0.86 in CHCl₃).

Ripperger, H. *et al.*, *Phytochemistry*, 1996, **43**, 705-707 (*isol, pmr, cmr, ms, cd, struct*)

Solidaline

S-353

[67257-83-4]



Relative Configuration

C₂₃H₂₇NO₆ 413.469

Alkaloid from *Corydalis solida*. Cryst. (CHCl₃/MeOH). Mp 198°. λ _{max} 233 (log ϵ 4.03); 281 (log ϵ 3.51); 316 (log ϵ 3.42); 364 (log ϵ 3.33) (MeOH). λ _{max} 248 (log ϵ 4.02); 318 (log ϵ 3.92); 367 (log ϵ 3.9) (MeOH/HCl aq.).

Manske, R.H.F. *et al.*, *Can. J. Chem.*, 1978, **56**, 383-386 (*isol, uv, pmr, cmr, ms*)

Suau, R. *et al.*, *Tetrahedron*, 1999, **55**, 4019-4028 (*synth*)

Solimocurarine

S-354

[1361-30-4]

Struct. unknown. Alkaloid from *Strychnos solimoesana* (Loganiaceae). Blue col. with Ce(IV).

Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148

Solimoesines

S-355

[1361-29-1 (Solimoesine III), 1361-28-0 (Solimoesine II), 1361-27-9 (Solimoesine I)]


Struct. unknown. Alkaloids from *Strychnos solimoesana* (Loganiaceae). Solimoesines I, II and III identified, giving different colour reacts.

Marini-Bettolo, G.B. *et al.*, *Gazz. Chim. Ital.*, 1956, **86**, 1148

Sollasin B

S-356

[149297-98-3]



C₂₀H₃₂N₂O₂ 332.485


Constit. of *Poecillastra sollasi*. Solid. Sol. MeOH, C₆H₆; poorly sol. H₂O. [α]_D²⁴ +29.8 (c, 2.14 in CHCl₃). λ _{max} 215 (ϵ 18900) (heptane).

Killday, K.B. *et al.*, *J. Nat. Prod.*, 1993, **56**, 500-507 (*isol, pmr, cmr*)

Sollasin C

S-357

[149297-99-4]




C₂₁H₃₄N₂O₂ 346.512

Constit. of *Poecillastra sollasi*. Oil. Sol. MeOH, C₆H₆; poorly sol. H₂O. [α]_D²⁴ +10 (c, 1.26 in CHCl₃). λ _{max} 222 (ϵ 21772) (heptane) (Berdy).

Killday, K.B. *et al.*, *J. Nat. Prod.*, 1993, **56**, 500 (*isol, pmr, cmr*)

Solsodamine A

S-358




C₁₀H₁₂N₄O 204.231

Alkaloid from the fruit of *Solanum sodomaemum*. Yellow needles (EtOH). Mp 118-119°. λ _{max} 280 (log ϵ 3.08); 320 (log ϵ 3.06); 348 (log ϵ 2.93) (MeOH). λ _{max} 280 (ϵ 1200); 320 (ϵ 1150); 348 (ϵ 850) (MeOH) (Berdy).

El Sayed, K.A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 848-850 (*isol, uv, ir, pmr, cmr, ms*)

Solsodamine B

S-359




C₈H₁₃N₃O 167.21

Constit. of the fruit of *Solanum sodomaeum*. Needles (EtOH). Mp 112-113°. λ_{max} 271 (log ε 3.1); 317 (log ε 3.01); 342 (log ε 2.03) (MeOH). λ_{max} 271 (ε 1280); 317 (ε 1030); 342 (ε 550) (MeOH) (Berdy).

El Sayed, K.A. *et al.*, *J. Nat. Prod.*, 1998, **61**, 848-850 (*isol, uv, ir, pmr, cmr, ms*)

Somamide A**S-360**

[350811-51-7]

R¹ = -CH₂CH₂SOMe, R² = -(CH₂)₄CH₃

Absolute Configuration

C₄₈H₆₇N₇O₁₂S 966.163

Depsipeptide antibiotic. Related to Dolastatin 13. *Isol.* from an assemblage of *Lyngbya majuscula* and *Schizothrix* sp. Oil. [α]_D²⁵ -2.5 (c, 0.08 in MeOH).

Nogle, L.M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 716-719 (*isol*)

Yokokawa, F. *et al.*, *Tet. Lett.*, 2002, **43**, 8673-8677 (*synth*)

Somniferine†**S-361**


Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Mp 185-187° dec.

Majumdar, D.N. *et al.*, *Curr. Sci.*, 1952, **21**, 46; *CA*, **47**, 2184b

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f

Somniferine†**S-362**

[117611-63-9]

C₃₆H₃₆N₂O₇ 608.69

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). Hypnotic agent. Amorph. [α]_D²⁰ -297 (c, 1 in

CHCl₃). Log P 0.28 (uncertain value) (calc).

3-Me ether: O-Methylsomniferine

[117611-62-8]

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from *Papaver somniferum* (opium poppy) (Papaveraceae). No phys. props. reported.

Dragar, C. *et al.*, *Tet. Lett.*, 1988, **29**, 3115 (*struct*)

Somniferine**S-363**

Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Mp 120° dec.

Majumdar, D.N. *et al.*, *Curr. Sci.*, 1952, **21**, 46; *CA*, **47**, 2184b

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f


Somnine**S-364**

Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Mp 300°.

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f

Somocystinamide A**S-365**

[422312-35-4]



Absolute Configuration

C₄₂H₇₀N₄O₄S₂ 759.171

Isol. from a cyanobacterial assemblage of *Lyngbya majuscula* and *Schizothrix* sp. Cytotoxic. Amorph. solid. [α]_D²² +13.5 (c, 0.75 in CHCl₃). λ_{max} 210 (ε 17700); 241 (ε 9900) (MeOH).

Nogle, L.M. *et al.*, *Org. Lett.*, 2002, **4**, 1095-1098 (*isol, pmr, cmr*)

Suyama, T.L. *et al.*, *Org. Lett.*, 2008, **10**, 4449-4452 (*synth*)


Songorinine**S-366**C₂₇H₃₃NO₉ 515.559

Struct. unknown. Alkaloid from the upper parts of *Aconitum soongoricum* (Ranunculaceae). Cryst. (MeOH). Mp 203-204° dec.

Samatov, A.S. *et al.*, *Dokl. Akad. Nauk UzSSR*, 1965, **22**, 21-22; *CA*, **63**, 12000f

Sonodione**S-367**

Isocorydione
[75652-90-3]

C₂₀H₁₉NO₅ 353.374

Alkaloid from stem bark of *Hernandia sonora* and *Dehaasia triandra*. Violet prisms (MeOH). Mp 207-209°. λ_{max} 224 (log ε 4.43); 277 (log ε 4.16); 323 (log ε 4.22); 565 (log ε 3.49) (EtOH).

N-De-Me: Norsonodione. *N*-Demethylsonodione. *Norisocorydione.* *N*-Demethylisocorydione
[173220-46-7]

C₁₉H₁₇NO₅ 339.347

From stem bark of *Hernandia sonora* and *Dehaasia triandra*. Violet prisms (CHCl₃/MeOH). Mp 216-219°. λ_{max} 223 (log ε 4.44); 276 (log ε 4.13); 327 (log ε 4.23); 564 (log ε 3.58) (EtOH).

O¹-De-Me: 1-Demethylsonodione

[174581-72-7]

C₁₉H₁₇NO₅ 339.347

From stem bark of *Hernandia sonora*. Greenish prisms (MeOH). Mp 230-233°. λ_{max} 219 (log ε 4.44); 283 (log ε 4.09); 336 (log ε 4.28); 492 (sh) (log ε 3.36); 642 (log ε 3.67) (EtOH).

4,5-Didehydro: Dehydroisocorydione

[176901-23-8]

C₂₀H₁₇NO₅ 351.358

Alkaloid from leaves of *Dehaasia triandra*. Needle-like cryst. (Me₂CO). Mp 252.5°.

Chen, I.-S. *et al.*, *Planta Med.*, 1995, **61**, 537 (*isol, uv, ir, pmr, ms, struct*)

Lee, S.-S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 55 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lee, S.S. *et al.*, *Tetrahedron*, 1996, **52**, 6561 (*Dehydroisocorydione*)


Sonpeimine**S-368**C₂₇H₄₃NO₄ 445.641

Struct. unknown. Steroidal alkaloid from "Bei-son-pei-mu" (*Fritillaria* spp.) (Liliaceae). Mp 256-258°.

Chu, T.-T. *et al.*, *CA*, 1957, **51**, 446a

Sophazrine**S-369**

[137760-66-8]


C₁₉H₂₅N₃O₂ 327.425

Alkaloid from *Sophora griffithii* (Fabaceae). Amorph. solid. [α]_D²³ +213 (c, 0.02 in MeOH). New type of tetracyclic quinolizidine alkaloid. λ_{max} 233 (log ε 3.81); 309 (log ε 3.87) (MeOH).

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1991, **54**, 929-935 (*isol, uv, ir, pmr, cmr, ms, struct*)

Sophoattarine


S-370

C₂₂H₂₅N₃O₄ 395.457Alkaloid from *Sophora griffithii*. Gum. Opt. inactive. λ_{max} 234 (log ε 3.71); 310 (log ε 3.77) (MeOH).

Ajaz, M.S. et al., Thesis, Univ. of Karachi, 1993, (isol, pmr, cmr, ms, struct)

Sophocarpine

S-371

13,14-Didehydromatridin-15-one, 9CI
[6483-15-4]Absolute
ConfigurationC₁₅H₂₂N₂O 246.352Alkaloid from *Sophora alopecuroides*, *Sophora pachycarpa*, *Sophora griffithii*, *Sophora flavescens*, *Sophora prodani* and other *Sophora* spp., *Ammothamnus lehmannii* (preferred genus name *Sophora*), *Goebelia pachycarpa* (preferred genus name *Sophora*), and *Leontice smirnovii* (Fabaceae, Leonticaceae). Antineoplastic agent, inhibitor of Walker 256 carcinosarcoma and other tumours. Antiarrhythmic agent, nematocide. Cryst. + 1H₂O. Poorly sol. hexane. Mp 53-54° (81-82°). [α]_D -29.4 (EtOH). Log P 0.28 (uncertain value) (calc). λ_{max} 260 (log ε 3.6) (EtOH). λ_{max} 260 (ε 2500) (EtOH).▶ LD₅₀ (rat, orl) 196 mg/kg. LD₅₀ (rat, ipr) 124 mg/kg. WG1136000

Hydrochloride: Mp 300°.

Picrate:

Yellow cryst. (H₂O). Mp 155-157°.1-Oxide: **Sophocarpidine**. *Sophocarpine* N-oxide. *Oxysophocarpine*
[26904-64-3]C₁₅H₂₂N₂O₂ 262.351Alkaloid from *Sophora pachycarpa*, *Sophora alopecuroides* and *Sophora flavescens* (Fabaceae). Mp 202-204° (200° dec.). [α]_D +32.4 (c, 2.0 in EtOH). [α]_D¹⁸ +21 (c, 2.0 in EtOH).1-Oxide, picrate: [26904-65-4]
Mp 200-202°.5α-Hydroxy: **5-Hydroxysophocarpine**
[131871-12-0]C₁₅H₂₂N₂O₂ 262.351Alkaloid from *Sophora flavescens* var. *angustifolia* (Fabaceae). Amorph. solid. [α]_D -36 (c, 0.06 in MeOH).9-Hydroxy: **9α-Hydroxysophocarpine**

[220961-52-4]

C₁₅H₂₂N₂O₂ 262.351Alkaloid from *Sophora viciifolia* (Fabaceae). Cryst. (C₆H₆). Mp 120°. [α]_D²⁵ -44.2 (c, 0.36 in EtOH).9α-Hydroxy, N-oxide: **9-Hydroxysophocarpine N-oxide**C₁₅H₂₂N₂O₃ 278.35Alkaloid reported from *Sophora flavescens*. Details not accessible. Cited by Murakoshi et al. *Phytochemistry*, 1982, **21**, 2379-2384. CAS no. not found 8-14CI.12α-Hydroxy: **12α-Hydroxysophocarpine**C₁₅H₂₂N₂O₂ 262.351Alkaloid from the roots of *Sophora flavescens*. Plates (petrol/Me₂CO). Mp 90-92°. [α]_D²⁴ +137.3 (c, 0.1 in MeOH). λ_{max} 206 (log ε 3.86); 258 (log ε 3.37) (MeOH).12β-Hydroxy: **12β-Hydroxysophocarpine**
[259860-47-4]C₁₅H₂₂N₂O₂ 262.351Alkaloid from the seeds of *Sophora viciifolia*. Cryst. (C₆H₆). Mp 146°. [α]_D²⁵ -215.1 (c, 0.22 in EtOH).5-Epimer: **5-Episophocarpine**. 13,14-Dehydrosophoridine

[68398-59-4]

C₁₅H₂₂N₂O 246.352Alkaloid from epigeal parts of *Sophora flavescens* and from *Sophora alopecuroides*. Prisms + 1H₂O (hexane). Mp 83-84°. [α]_D²⁵ +76.8 (c, 0.266 in EtOH) (synthetic).5-Epimer, 1-oxide: **13,14-Dehydrosophoridine N-oxide**

[64838-17-1]

C₁₅H₂₂N₂O₂ 262.351Alkaloid from *Sophora alopecuroides* (Fabaceae). Cryst. (Et₂O). Mp 68-70°. [α]_D²⁵ +27.5 (c, 0.75 in H₂O). C-7 config. wrongly shown, apparently a drawing error.

[78003-71-1, 68365-91-3]

Orekhov, A.P. et al., *Ber.*, 1934, **67**, 77-83 (isol)Okuda, S. et al., *Chem. Ind. (London)*, 1962, 1326-1327 (synth, struct)Iskandarov, S. et al., *Khim. Prir. Soedin.*, 1968, **4**, 106-109; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **4**, 90-92 (ms)Zainutdinov, U.N. et al., *CA*, 1970, **72**, 121754u (oxide)Yunusov, T.K. et al., *Khim. Prir. Soedin.*, 1972, **8**, 200-207; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 201-206 (ir)Kushmuradov, J.K. et al., *Phytochemistry*, 1972, **11**, 3441-3445 (isol, oxide)Karakozova, S.A. et al., *Khim. Prir. Soedin.*, 1975, **11**, 664-665; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 703 (isol)Kuchkarov, S. et al., *Khim. Prir. Soedin.*, 1977, **13**, 541-544; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 451-454 (epimer oxide)Abdusalimov, B.A. et al., *Khim. Prir. Soedin.*, 1977, **13**, 549-552; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 458-460 (biosynth)Ueno, A. et al., *Chem. Pharm. Bull.*, 1978, **26**, 1832-1836 (isol, struct, oxide)Morinaga, K. et al., *Chem. Pharm. Bull.*, 1978, **26**, 2483-2488 (isol, synth, cryst struct, epimer)Kushmuradov, Yu.K. et al., *Khim. Prir. Soedin.*, 1978, **14**, 231-233; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 190-191 (epimer)Wang, Y.L. et al., *J. Tradit. Chin. Med.*, 1987, **7**, 35 (pharmacol)Saito, K. et al., *J. Chromatogr.*, 1989, **462**, 333-340 (hplc)Saito, K. et al., *Planta Med.*, 1990, **56**, 487 (5-Hydroxysophocarpine)Matsuda, K. et al., *J. Agric. Food Chem.*, 1991, **39**, 189-191 (isol, activity)Xiao, P. et al., *Phytochemistry*, 1999, **50**, 189-193 (12β-Hydroxysophocarpine, 9α-Hydroxysophocarpine, cmr)Atta-ur-Rahman, et al., *J. Nat. Prod.*, 2000, **63**, 190-192 (12β-Hydroxysophocarpine, cryst struct)Ding, P.-L. et al., *Bioorg. Med. Chem. Lett.*, 2006, **16**, 1231-1235 (12α-Hydroxysophocarpine)

Sophochrysin

S-372

[1361-32-6]

C₁₅H₁₉N₃O₂ 273.334Tentative mol. formula. Struct. unknown. Alkaloid from *Sophora chrysophylla*, *Sophora microphylla*, *Sophora tetraptera* and bark of *Genista monosperma* (Fabaceae). Amorph. Mp 284-287°. [α]_D -113.2 (EtOH).


Picrate:

Short yellow needles. Mp 360° (darkens at 250°).

Briggs, L.H. et al., *J.C.S.*, 1937, 1795; 1938, 1206; 1942, 507-509; 555-556; 1960, 1955-1956González, A. et al., *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1959, **55**, 607-616; *CA*, **54**, 8874c (isol)Ahmed, Z.F. et al., *J. Chem. U.A.R.*, 1963, **6**, 227-242; *CA*, **63**, 3312a (isol)Morales Mendez, A. et al., *CA*, 1975, **82**, 121629p (isol)

Sophohejrine


S-373

Relative
ConfigurationC₃₀H₄₄N₄O₂ 492.703Alkaloid from *Sophora griffithii*. Yellow gum. [α]_D²⁵ -250 (c, 0.204 in MeOH).

Ajaz, M.S. et al., Thesis, Univ. of Karachi, 1993, (isol, pmr, cmr, ms, struct.)

Sopholupanizidone

S-374

C₂₅H₃₅N₃O₂ 409.57


Picrate: Mp 122-123°.

Abdusalamov, B.A. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 71

Abdusalamov, B.A. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 549; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 458 (*isol, uv, ir, ms, cmr, struct, biosynth*)

Sophoridine**S-380**

(5β)-*Matridin-15-one*, 9CI
[6882-68-4]



(-)-form

C₁₅H₂₄N₂O 248.367

Stereoisomer of Allomatrine, A-628, Matrine, M-121, Isomatrine, I-244 and Darvasamine, D-79.

▶ WG1250000

(+)-form

Alkaloid from epigeal parts of *Leontice albertii* (Leonticaceae). Cryst. (petrol). Mp 108-109°. [α]_D²⁵ +59.3 (c, 1.30 in EtOH).

(-)-form

Alkaloid from *Sophora alopecuroides* and *Sophora prodani* (Fabaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 109-110°. [α]_D¹⁸ -63.8 (c, 0.53 in H₂O).

▶ LD₅₀ (mus, ipr) 60 mg/kg.

1α-Oxide:

Semisynthetic. Cryst. [α]_D²⁵ -42.9 (c, 0.17 in EtOH).

1β-Oxide: Sophoridine N-oxide

[54809-74-4]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Euchresta japonica* (Fabaceae). Strongly hygroscopic needles. [α]_D²⁸ -5.6 (c, 0.14 in EtOH). Sophoridine N-oxide of undetd. config. was also isol. from *S. alopecuroides*.

3α-Hydroxy: 3α-Hydroxysophoridine

[41645-69-6]

C₁₅H₂₄N₂O₂ 264.367

Minor alkaloid from epigeal parts of *Sophora alopecuroides* (Fabaceae). Mp 162-164°. [α]_D²⁰ -50.6 (c, 0.555 in EtOH).

14β-Hydroxy: 14β-Hydroxysophoridine

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from the seeds of *Sophora viciifolia*. Needles (CH₂Cl₂/hexane). Mp 90°. [α]_D²⁵ -94.8 (c, 0.47 in EtOH).

Kamalitinov, D. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 352; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 297 (*isol*)

Yunusov, T.K. *et al.*, *Khim. Prir. Soedin.*, 1972, **8**, 200; *Chem. Nat. Compd. (Engl. Transl.)*, 1972, **8**, 201 (*ir*)

Monakhova, T.E. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 59; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 52 (*deriv*)

Vulfson, N.S. *et al.*, *Khim. Geterotsikl. Soedin.*, 1974, **10**, 251; *Chem. Heterocycl. Compd. (Engl. Transl.)*, 1974, **10**, 221 (*ms*)

Kamaev, F.G. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 744; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 765 (*pmr, struct*)

Morinaga, K. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 2483 (*pmr*)


Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 355; 1981, **17**, 588; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 308; 1981, **17**, 424 (*cryst struct, Sophoridine, Sophoridine N-oxide*)

Kuchkarov, S. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 364; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 364 (*isol*)

Ohmiya, S. *et al.*, *Chem. Pharm. Bull.*, 1980, **28**, 546 (*isol, struct, oxide*)

Ibragimov, B.T. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 71; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 66 (*rev, stereochem*)

Xiao, P. *et al.*, *Phytochemistry*, 1999, **50**, 189-193 (*14β-Hydroxysophoridine*)

Sophosalimine**S-381**


C₂₇H₃₂N₄O₃ 460.575

Alkaloid from *Sophora griffithii*. Readily dec. Opt. inactive. λ_{max} 234 (log ε 3.81); 310 (log ε 3.85) (MeOH).

Ajaz, M.S. *et al.*, *Thesis*, Univ. of Karachi, 1993, (*isol, pmr, cmr, ms, struct*)

Sorbicillactone A**S-382**

[664987-12-6]



Absolute Configuration

C₂₁H₂₃NO₈ 417.415

Prod. by *Penicillium chrysogenum* isol. from the sponge *Ircinia fasciculata*. Cytotoxic. Anti-HIV agent. Yellow needles (MeOH aq.). Mp 205° dec. [α]_D²⁰ -939 (c, 0.2 in MeOH).

2',3'-Dihydro: Sorbicillactone B

[861434-14-2]

C₂₁H₂₅NO₈ 419.43


Prod. by *Penicillium chrysogenum* isol. from *Ircinia fasciculata*. Light brown solid (MeOH aq.). Mp 109-115°. [α]_D²⁰ -327 (c, 0.2 in MeOH).

Bringmann, G. *et al.*, *Tetrahedron*, 2005, **61**, 7252-7265 (*isol, cd, pmr, cmr*)

Bringmann, G. *et al.*, *Mar. Drugs*, 2007, **5**, 23-30 (*prodn*)

Sorelline**S-383**

[73004-62-3]



Absolute Configuration

C₂₀H₂₄N₂ 292.423

Alkaloid from *Aristotelia peduncularis* (Elaeocarpaceae). Mp 165-168°. [α]_D²² +157 (c, 1.067 in CHCl₃).

9-Epimer: Isosorelline

C₂₀H₂₄N₂ 292.423

Alkaloid from *Aristotelia serrata* and *Aristotelia fruticosa* (Elaeocarpaceae). Cryst. (MeOH). Mp 160-162°. [α]_D^{21.5} +120 (CHCl₃). CAS no. not found 8-14CI.

(±)-form

Synthetic. Cryst. (CHCl₃). Mp 167-168°.

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 2539-2546 (*isol, uv, ir, pmr, cmr, ms, struct*)


Bick, I.R.C. *et al.*, *Alkaloids (Academic Press)*, 1985, **24**, 113-151 (*rev, Isosorelline*)

Burkard, S. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 275-289 (*synth, abs config*)

Sparsomycin, INN, USAN**S-384**

Sparsogenin. B 79B. NSC 59729. U 19183. Antibiotic B 79B. Antibiotic U 19183

[1404-64-4]



C₁₃H₁₉N₃O₅S₂ 361.442

Prod. by *Streptomyces sparsogenes* and *Streptomyces cuspidosporus*. Antineoplastic antibiotic. Inhibits protein biosynth. Sol. H₂O, MeOH, EtOH; poorly sol. CHCl₃, hexane. Mp 208-209° dec. [α]_D²⁵ +69 (c, 0.5 in H₂O). Log P -1.92 (calc). λ_{max} 284 (sh) (ε 10500); 328 (ε 25600) (0.01N KOH) (Derep). λ_{max} 270 (sh) (ε 15500); 302 (ε 23500) (H₂O) (Derep). λ_{max}. λ_{max} 328 (E1%/1cm 710) (H₂SO₄) (Berdy).

▶ LD₅₀ (mus, orl) 22 mg/kg. LD₅₀ (mus, ipr) 2.4 mg/kg. WG5850000

(Z)-Isomer: Isosparsomycin

[58462-97-8]

Cryst. (H₂O). Mp 158-164°.

S¹³-Epimer: S-Episparsomycin

[77880-77-4]

[α]_D²⁵ +48 (c, 0.175 in H₂O). Other diastereoisomers also prepd.

S¹⁵-Oxide (R-): Sparoxomycin A2

[174390-02-4]

C₁₃H₁₉N₃O₆S₂ 377.442

Prod. by *Streptomyces sparsogenes* SN2325. Cell proliferation modulator. Powder. Sol. MeOH, H₂O. Mp 170-174° dec. [α]_D²⁸ +3.6 (c, 0.33 in H₂O). λ_{max} 203 (ε 8800); 303 (ε 13600) (MeOH) (Berdy).

S¹⁵-Oxide (S-): Sparoxomycin A1


[174390-01-3]

C₁₃H₁₉N₃O₆S₂ 377.442

Prod. by *Streptomyces sparsogenes* SN2325. Cell proliferation modulator. Powder. Sol. MeOH, H₂O. Mp 169-173° dec. [α]_D²⁸ +47.8 (c, 0.13 in H₂O). λ_{max} 203 (ε 9300); 303 (ε 15200)

(MeOH) (Berdy).

[77880-75-2, 77880-76-3, 63222-12-8]

Wiley, P.F. *et al.*, *J.O.C.*, 1976, **41**, 1858 (*struct. uv, ir, nmr*)Lin, C.-C.L. *et al.*, *J. Med. Chem.*, 1977, **20**, 337 (*synth*)Goldberg, I.H. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 264 (*rev*)Ottenheim, H.C.J. *et al.*, *J.A.C.S.*, 1981, **103**, 1720-1723 (*abs config*)Ottenheim, H.C.J. *et al.*, *J.O.C.*, 1981, **46**, 3273 (*synth*)Ash, R.J. *et al.*, *Biochem. Biophys. Res. Commun.*, 1984, **125**, 784 (*stereochem, props*)Liskamp, R.M.J. *et al.*, *J. Med. Chem.*, 1984, **27**, 301 (*sar*)Hwang, D.R. *et al.*, *J.O.C.*, 1985, **50**, 1264 (*synth*)Ottenheim, H.C.J. *et al.*, *Prog. Med. Chem.*, 1986, **23**, 219-268 (*rev*)Parry, R.J. *et al.*, *J.A.C.S.*, 1992, **114**, 5946 (*biosynth*)Parry, R.J. *et al.*, *Tet. Lett.*, 1994, **35**, 7497 (*biosynth*)Ubukata, M. *et al.*, *J. Antibiot.*, 1996, **49**, 65-70; 1096-1100 (*Sparoxomicin*)Nakajima, N. *et al.*, *Biosci. Biotechnol. Biochem.*, 2003, **67**, 2556-2566 (*synth, pmr, cmr, ms*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SKX000**Sparteine, INN** **S-385***Dodecahydro-7,14-methano-2H,6H-dipyr-ido[1,2-a:1',2'-e]diazocine, 9CI. Lupinidine (obsol.). Pachycarpine (obsol.)*

(-)-form

C₁₅H₂₆N₂ 234.384Diastereoisomeric with α -Isosparteine, I-321 and β -Isosparteine, I-322. Anti-inflammatory, cathartic, diuretic agent. Shows oxytocic props. Shows antimicrobial activity. Log P 2.15 (calc).**(+)-form** [492-08-0]Alkaloid from several genera of the family Fabaceae, subfamily Faboideae, notably *Lupinus pusillus*, *Lupinus niger*, *Cytisus caucasicus*, *Baptisia* spp., *Cytisus scoparius*, *Genista monosperma*, *Pelargonium acutifolia*, *Pelargonium longifolia*, *Sophora pachycarpa* and *Ammodendron* spp. Used as 1% aq. soln. of iodide for extraction-photometric detn. of Ti (λ_{\max} 390 nm, ϵ 10300, CHCl₃). Antiinflammatory, cathartic, diuretic agent. Shows oxytocic props. Shows antimicrobial activity. Bp₈ 173-174°. [α]_D +17.1 (EtOH). n_D^{20} 1.5312. λ_{\max} 268 (ϵ 16000) (HCl) (Berdy).▶ LD₅₀ (mus, ivn) 26 mg/kg. RT0620000*Perchlorate:*

Plates (EtOH). Mp 171-172°.

*Dipicrate:*Cryst. (EtOH/Me₂CO). Mp 205-206°.N¹⁶-Oxide: *Pachycarpine N¹⁶-oxide*

[30301-23-6]

C₁₅H₂₆N₂O 250.383Alkaloid from *Ammodendron karelinii* (Fabaceae). Cryst. (Me₂CO).

Mp 153-154°.

(-)-form [90-39-1]Alkaloid from several genera of the family Fabaceae, subfamily Faboideae, notably many *Lupinus* and *Adenocarpus* spp., also *Piptanthus nanus*, *Sarothamnus* spp. *Chamaecytisus* spp. Used mainly as sulfate used in chiral catalytic systems. Useful complexing agent for metal ions. Antiarrhythmic agent. Has been used in treatment of cardiac insufficiency. Shows curare-like properties. Liq. Bp₁₈ 181°. [α]_D -17 (EtOH). n_D^{19} 1.5289. pK_{a1} 11.96; pK_{a2} 4.8 (15°). Log P 2.15 (calc). This is the commonest form, frequently referred to as just Sparteine.▶ LD₅₀ (rat, orl) 960 mg/kg. LD₅₀ (rat, ipr) 42 mg/kg. WG5950000*Hydrobromide (1:2):* Mp 194-195°.*Hydroiodide (1:2):* Mp 257-258°.*Sulfate: Sparteine sulfate, USAN. Many other names*


[299-39-8]

Cryst. + 5H₂O. Dec. at 136°.▶ Human reprod. effects. LD₅₀ (mus, orl) 350 mg/kg. LD₅₀ (mus, ivn) 15 mg/kg. WG6430000*Dipicrate:* Mp 208° (205-206°).*Methiodide:* Mp 239-240°.**(±)-form** [4985-24-4]Alkaloid from *Chamaecytisus proliferus* and *Adenocarpus hispanicus* (Fabaceae). Liq. Bp₁ 119-121°. n_D^{20} 1.5223.*Perchlorate:* Mp 131-133°.*Dipicrate:* Mp 208°.

[6160-12-9]

Stenhouse, J. *et al.*, *Annalen*, 1858, **78**, 1 (*isol*)Clemo, G.R. *et al.*, *J.C.S.*, 1933, 644; 1949, 663 (*struct*)Jaretsky, R. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1934, **272**, 152 (*occur*)Leonard, N.J. *et al.*, *J.A.C.S.*, 1950, **72**, 1316 (*ir, synth, rev, resolu*)Anet, E. *et al.*, *Nature (London)*, 1950, **165**, 35 (*synth*)Okuda, S. *et al.*, *Chem. Ind. (London)*, 1961, 1116 (*abs config*)van Tاملen, E.E. *et al.*, *J.A.C.S.*, 1969, **91**, 7372 (*synth*)Vard, A.G. *et al.*, *CA*, 1970, **76**, 121173m (*use*)Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1971, **24**, 1537 (*isol*)Binnig, F. *et al.*, *Arzneim.-Forsch.*, 1974, **24**, 752; 753 (*pharmacol*)Klyne, W. *et al.*, *J.C.S. Perkin 1*, 1974, 2565 (*cd*)Nakano, T. *et al.*, *J.O.C.*, 1974, **39**, 3584 (*ms*)Bohlmann, F. *et al.*, *Chem. Ber.*, 1975, **108**, 1043 (*cmr*)Kushmuradov, Yu.K. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 717; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 604 (*oxide*)Fanso-Free, S.N.Y. *et al.*, *J.A.C.S.*, 1979, **101**, 1549 (*N nmr, conform*)Golebiewski, W.M. *et al.*, *Chem. Comm.*, 1983, 1509 (*biosynth*)Wink, M. *et al.*, *Z. Naturforsch., C*, 1984, **39**, 548-552 (*activity*)Wippich, C. *et al.*, *Experientia*, 1985, **41**, 1477-1479 (*activity*)Eichelbaum, M. *et al.*, *Xenobiotica*, 1986, **16**, 465 (*rev, metab*)Takatsu, N. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 4990 (*synth*)Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 3779Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1416Bour, P. *et al.*, *J. Phys. Chem. A*, 1997, **101**, 9783-9790 (*vibration cd, uv, ir*)Lee, Y.-M. *et al.*, *Acta Cryst. C*, 2002, **58**, 6733-6734 (*perchlorate, cryst struct*)Smith, B.T. *et al.*, *Org. Lett.*, 2002, **4**, 2577-2579 (*synth*)Iyengar, R. *et al.*, *Chemtracts: Org. Chem.*, 2004, **17**, 92-96 (*rev, synth*)He, H. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 1557-1567 (*synth*)Norcross, N.R. *et al.*, *J.O.C.*, 2008, **73**, 7939-7951 (*synth*)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, PAB250; SKX500; SKX750**Spartocytisine****S-386**


[1361-33-7]

C₁₂H₁₇NO₃ 223.271Struct. not fully established. Alkaloid from the seeds of *Spartocytisus filipes* (preferred genus name *Cytisus*) (Fabaceae). Cryst. (heptane/C₆H₆). Mp 158-159°. [α]_D +11.4.*Picrate:*Cryst. (H₂O). Mp 174-175°.*Styphnate:*

Cryst. (EtOH). Mp 212°.

N,O-Di-Ac:

Cryst. (petrol). Mp 125-126°.

González, A.G. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1962, **58**, 483-490; *CA*, **59**, 2869d (*isol*)**Spesciosine****S-387**C₂₈H₃₁NO₆ 477.556

▶ DF9485000

(S)-form [16892-03-8]Alkaloid from the tubers of *Colchicum speciosum* (Liliaceae). Yellow prisms (C₆H₆/Me₂CO). Mp 209-211°. [α]_D²⁰ -21.2 (c, 1.75 in CHCl₃).*Me ether: O-Methylspesciosine*

[172685-48-2]

C₂₉H₃₃NO₆ 491.583Alkaloid from *Colchicum ritchii*.Amorph. [α]_D²⁵ -17 (c, 0.20 in MeOH).

O²-De-Me: Specicolchine

[111509-13-8]

C₂₇H₂₉NO₆ 463.529Alkaloid from *Colchicum ritchii* (Liliaceae). Amorph. [α]_D²⁰ -37 (c, 0.13 in MeOH).**O³-De-Me: Specioritchine**


[111509-12-7]

C₂₇H₂₉NO₆ 463.529Alkaloid from *Colchicum ritchii* (Liliaceae). Amorph. [α]_D²⁰ -57 (c, 0.11 in MeOH).**O¹⁰-De-Me: Specioseine**

[142735-47-5]

C₂₇H₂₉NO₆ 463.529Minor alkaloid from *Colchicum speciosum* (Liliaceae). Cryst. (Me₂CO). Mp 169-171°. [α]_D²⁰ -78 (c, 1.01 in CHCl₃).**Deoxy: Speciosamine**

[97763-01-4]

C₂₈H₃₁NO₅ 461.557Alkaloid from *Colchicum speciosum* (Liliaceae). Mp 192-194°. [α]_D²⁰ -42 (c, 0.26 in CHCl₃).Kiselev, V.V. et al., *Zh. Obshch. Khim.*, 1956, **26**, 3218; *CA*, **51**, 8119i (isol)Ramage, R. et al., *Tetrahedron*, 1971, **27**, 1499 (ir, uv, ms, pmr, struct, synth)Hrbek, J. et al., *Coll. Czech. Chem. Comm.*, 1982, **47**, 2258 (cd)Chommadov, B. et al., *Khim. Prir. Soedin.*, 1985, **21**, 417; 1991, **27**, 253; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 218; 1991, **27**, 394 (Speciosamine, Specioseine)Freyer, A.J. et al., *J. Nat. Prod.*, 1987, **50**, 684 (Specicolchine, Specioritchine)Abu Zarga, M.H. et al., *Z. Naturforsch., B*, 1995, **50**, 1424 (O-Methylspecioseine)Izotova, L.Y. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1997, **33**, 469-472 (cryst struct)**Spectraline S-388****14-(5-Hydroxy-6-methyl-2-piperidinyl)-2-tetradecanone. 3-Hydroxy-2-methyl-6-(13-oxotetradecyl)piperidine. Cassinicine**(2*R*,5*R*,6*R*)-formC₂₀H₃₉NO₂ 325.534

The more recently isol. (-)-spectaline may or may not be identical to the prev. isol. and poorly documented Cassinicine, Mp 170°.

(2*R*,5*R*,6*R*)-form**6-Isospectraline**

[709046-69-5]

Alkaloid from the flowers of *Cassia spectabilis*. Amorph. yellow solid. Mp 116.7-119.6°. [α]_D²⁰ -7 (c, 0.31 in CH₂Cl₂).**(2*R*,5*S*,6*S*)-form [64384-95-8]**Alkaloid from *Cassia spectabilis*, *Cassia leptophylla* and *Prosopis africana* (Fabaceae). Oil; cryst. (EtOAc) (as hydrochloride). Mp 155° (hydrochloride). [α]_D²⁵ +8 (c, 0.27 in CHCl₃).**(2*S*,5*R*,6*R*)-form [65560-25-0]**Alkaloid from leaves of *Cassia leptophylla* and *Cassia spectabilis* (Fabaceae).Antinociceptive agent. Shows DNA-damaging props. Amorph. solid. Mp 118° (48.9-52.5°). [α]_D²⁰ -12 (c, 0.14 in CH₂Cl₂).**Ac: 3-O-Acetyl(-)-spectaline**

[709046-67-3]

C₂₂H₄₁NO₃ 367.571Alkaloid from the flowers of *Cassia spectabilis*.**7-Hydroxy: 7-Hydroxy(-)-spectaline**

[709046-68-4]

C₂₀H₃₉NO₃ 341.533Alkaloid from the flowers of *Cassia spectabilis*. Oil. [α]_D²⁰ -7 (c, 0.04 in CH₂Cl₂).**(2ξ,5ξ,6ξ)-form****13'ξ-Alcohol: 5-Hydroxy-α,6-dimethyl-2-piperidinetridecanol. 6-(13-Hydroxytridecyl)-2-methyl-3-piperidinol. Spectalinine**

[66408-16-0]


C₂₀H₄₁NO₂ 327.549Alkaloid from leaves and seeds of *Cassia spectabilis* (Fabaceae). Oil; cryst. (as hydrochloride). Mp 120° (hydrochloride). [α]_D²⁵ -8.4 (c, 0.12 in CHCl₃).

[96614-54-9 ((±)-form)]

Mulchandani, N.B. et al., *Planta Med.*, 1977, **32**, 357-361 (Cassinicine)Christofidis, I. et al., *Tetrahedron*, 1977, **33**, 977; 3005 (isol, ir, pmr, ms, cmr, struct)Khuong-Huu, Q. et al., *J. Chem. Soc. Pak.*, 1982, **4**, 267 (cmr, struct)Paterne, M. et al., *J. Chem. Res., Synop.*, 1985, 278 (synth, Spectaline)Hasseberg, H.-A. et al., *Annalen*, 1989, 255 (synth, ir, pmr, cmr, Spectaline)Momose, T. et al., *Tet. Lett.*, 1993, **34**, 5785 (synth, Spectaline)Bolzani, V. da S. et al., *Tetrahedron*, 1995, **51**, 5929-5934 ((-)-Spectraline)Bolzani, V. da S. et al., *Ann. Acad. Bras. Cinc.*, 1999, **71**, 181 ((-)-Spectraline, activity)Viegas, C. et al., *J. Nat. Prod.*, 2004, **67**, 908-910 ((-)-Spectraline, 6-Isospectraline, 7-Hydroxyspectraline, Ac)**Sperabillin C S-389**

TAN 749C. Antibiotic TAN 749C

[111337-84-9]



R = H

C₁₅H₂₇N₅O₃ 325.41Peptide antibiotic. Prod. by *Pseudomonas fluorescens*. Active against gram-positive and -negative bacteria. Powder + ½H₂O (as hydrochloride). Sol. H₂O, DMSO, MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁰ -11 (c, 0.68 in H₂O). λ_{max} 264 (ε 28400) (H₂O) (Derep).**4'Z-Isomer: Sperabillin A. TAN 749A.**

Antibiotic TAN 749A


[111337-86-1]

C₁₅H₂₇N₅O₃ 325.41From *Pseudomonas fluorescens*. Active against gram-positive and -negative bacteria. Powder + 1H₂O (as hydro-chloride). Sol. H₂O, DMSO, MeOH; poorly sol. Me₂CO, hexane. [α]_D²⁰ -11 (c, 1.06 in H₂O). λ_{max} 264 (ε 28400) (H₂O) (Derep).▶ LD₅₀ (mus, scu) 500 mg/kg, LD₅₀ (mus, orl) 2000-4000 mg/kg, LD₅₀ (mus, ipr) 200 mg/kg.

[111337-88-3, 111337-87-2]


Hashiguchi, S. et al., *J.C.S. Perkin 1*, 1991, 2435 (synth)Katayama, N. et al., *J. Antibiot.*, 1992, **45**, 10 (isol, pmr, cmr, props)Hida, T. et al., *Bull. Chem. Soc. Jpn.*, 1993, **66**, 863 (isol, struct)**Speranberculatine A**

S-390

C₁₂H₁₄N₂O₆ 282.252Alkaloid from *Speranskia tuberculata*. Pale yellow gum. [α]_D¹⁸ -14.8 (c, 0.5 in MeOH). λ_{max} 204 (log ε 3.79); 230 (log ε 2.44); 271 (sh) (log ε 2.04) (MeOH).Shi, J.G. et al., *Chin. Chem. Lett.*, 2000, **11**, 785-788Shi, J.G. et al., *J. Nat. Prod.*, 2000, **63**, 782-786**Speranculatine A**

S-391

[268728-59-2]




Relative Configuration

C₂₀H₂₄N₂O₁₀ 452.417Alkaloid from *Speranskia tuberculata*. Prisms (EtOAc). Mp 192-194°. [α]_D¹⁸ +21.9 (c, 0.54 in MeOH). λ_{max} 225 (log ε 4.87); 267 (log ε 1.62) (MeOH).**5-Epimer: Speranculatine B**

[268728-60-5]

C₂₀H₂₄N₂O₁₀ 452.417Alkaloid from *Speranskia tuberculata*. Prisms (EtOAc). Mp 147-149°. [α]_D¹⁸ +6 (c, 0.3 in MeOH). λ_{max} 224 (log ε 4.96); 267 (log ε 1.66) (MeOH).**5-Epimer, N^{1'}-de-Me: Speranculatine C**C₁₉H₂₂N₂O₁₀ 438.39Alkaloid from *Speranskia tuberculata*. Cryst. (EtOAc). Mp 183-185°. [α]_D¹⁸ +9.6 (c, 0.43 in MeOH). λ_{max} 221 (log ε 4.57); 262 (log ε 1.38) (MeOH).Shi, J.G. et al., *Chin. Chem. Lett.*, 2000, **11**, 225-228Shi, J.G. et al., *J. Nat. Prod.*, 2000, **63**, 782-786**Speranskatine A**

S-392



C₁₀H₁₃NO₅ 227.216

Abs configs. not yet determined, but Speranskatines A and B belong to opposite enantiomeric series.

(+)-form [171090-84-9]

Alkaloid from whole plants of *Speranskia tuberculata* (Euphorbiaceae). Prisms (EtOAc). Mp 158-160°. [α]_D²⁵ +14.8 (c, 0.54 in MeOH).

(-)-form

N-De-Me: Speranskatine B

[171090-85-0]

C₉H₁₁NO₅ 213.19


From *Speranskia tuberculata* (Euphorbiaceae). Prisms (EtOAc). Mp 171-172°. [α]_D²⁵ -30 (c, 0.30 in MeOH).

Shi, J.-G. *et al.*, *Phytochemistry*, 1995, **40**, 1299-1302 (isol, uv, ir, pmr, cmr, ms, cryst struct)

Zhu, Y. *et al.*, *Acta Cryst. C*, 1996, **52**, 2791-2792 (cryst struct)

Speranskilatine A

S-393



C₁₆H₁₈N₂O₈ 366.327


Alkaloid from *Speranskia tuberculata*. Prisms (CHCl₃). Mp 166-167°. [α]_D¹⁸ +3.8 (c, 1.3 in MeOH). λ_{\max} 229 (log ϵ 4.34); 268 (log ϵ 1.89) (MeOH).

Shi, J.-G. *et al.*, *J. Nat. Prod.*, 2000, **63**, 782-786

Spergualin

S-394

1-Amino-19-guanidino-11,15-dihydroxy-4,9,12-triazanonadecane-10,13-dione. BMG 162 aF₂. Antibiotic BMG 162 aF₂ [80902-43-8] [80952-47-2 (trihydrochloride)]



C₁₇H₃₇N₇O₄ 403.524

Prod. by *Bacillus laterosporus*. Active against gram-negative and -positive bacteria and tumours. Immunosuppressant, angiogenesis inhibitor. Sol. H₂O, MeOH; fairly sol. EtOH; poorly sol. Me₂CO, hexane. Mp 62-78° (as tripicrate). [α]_D²⁵ -11 (c, 1 in H₂O) (as trihydrochloride). Log P -4.07 (calc).

► LD₅₀ (mus, ipr) 150 mg/kg; LD₅₀ (mus, ivn) 80-120 mg/kg. MI7601060

15-Deoxy: 7-[(Aminoiminomethyl)amino]-N-[2-[4-[(3-aminopropyl)ami-

no]butyl]amino]-1-hydroxy-2-oxoethyl]heptanamide. N-[[[4-[(3-Aminopropyl)amino]butyl]carbamoyl]hydroxymethyl]-7-guanidinoheptanamide. 1-Amino-19-guanidino-11-hydroxy-4,9,12-triazanonadecane-10,13-dione. 15-Deoxyspergualin. Gusperimus, INN. Spanidin [89149-10-0] [104317-84-2 (racemate)]

C₁₇H₃₇N₇O₃ 387.524

Isol. from *Bacillus laterosporus*.

Immunosuppressant used clinically in kidney transplantation, antineoplastic agent. Angiogenesis inhibitor. [α]_D²⁵ -14.3 (c, 1 in H₂O) (synthetic). Log P -3 (calc). Commercial product is racemic.

15-Deoxy, trihydrochloride: Gusperimus trihydrochloride. BMS 181173. NKT 01. BMY 42215-1. NSC 356894

[84937-45-1]

[85468-01-5 (racemate)]

Syrup. [α]_D²⁵ -7.3 (c, 1 in H₂O). The pharmaceutical prod. appears to be the racemate.

[114760-38-2]

Takeuchi, T. *et al.*, *J. Antibiot.*, 1981, **34**, 1619-1621 (isol)

Umezawa, H. *et al.*, *J. Antibiot.*, 1981, **34**, 1622-1624 (struct)

Kondo, S. *et al.*, *J. Antibiot.*, 1981, **34**, 1625-1627 (synth)

Belg. Pat., 1982, ((Microbiochemical Research Foundation))890 229; CA, **97**, 90399s (isol)

Iwasawa, H. *et al.*, *J. Antibiot.*, 1982, **35**, 1665-1669; 1985, **38**, 886-898; 1987, **40**, 561-562; 1316-1324 (Gusperimus, synth, ir, pharmacol, pmr)

Eur. Pat., 1983, ((Microbiochemical Research Foundation))94 632; CA, **100**, 120781k (Gusperimus, synth, pharmacol)

Umezawa, K. *et al.*, *Biomed. Pharmacother.*, 1987, **41**, 227 (rev, pharmacol, props)

Bergeron, R.J. *et al.*, *J.O.C.*, 1987, **52**, 1700-1703 (Gusperimus, synth, bibl, pmr)

Nishimura, K. *et al.*, *Cell Sci.*, 1990, **6**, 639 (Gusperimus, rev, pharmacol)

Nishikawa, K. *et al.*, *J. Antibiot.*, 1991, **44**, 1101-1109 (Gusperimus, activity)

Oikawa, T. *et al.*, *Anti-Cancer Drugs*, 1992, **3**, 293-300 (Gusperimus, pharmacol)

Maeda, K. *et al.*, *Ann. N.Y. Acad. Sci.*, 1993, **685**, 123-135 (Gusperimus, rev, pharmacol)

Schorlemmer, H.U. *et al.*, *Ann. N.Y. Acad. Sci.*, 1993, **685**, 155-174 (Gusperimus, rev, pharmacol)

Tepper, M.A. *et al.*, *Ann. N.Y. Acad. Sci.*, 1993, **685**, 175-192; **696**, 123-132; 270-280 (Spergualin, Gusperimus, rev, pharmacol)

Suzuki, S. *et al.*, *Ann. N.Y. Acad. Sci.*, 1993, **696**, 263-269 (Gusperimus, rev)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1360 (Gusperimus)

Yuh, D.D. *et al.*, *Transplantation*, 1993, **55**, 578-591 (Gusperimus, pharmacol)

Tanabe, K. *et al.*, *J. Urol. (Baltimore)*, 1994, **152**, 562 (Gusperimus, pharmacol)

Mayo, D.J. *et al.*, *J. Pharm. Biomed. Anal.*, 1996, **14**, 457-463 (chiral hplc)


Durand, P. *et al.*, *J.O.C.*, 1998, **63**, 9723-9727 (synth, abs config)

Durand, P. *et al.*, *Tetrahedron*, 2001, **57**, 2757-2760 (synth)

Spermacoecine

S-395

2-[2,4a,10,10a-Tetrahydro-1,1,3,13-tetramethyl-1H,5H-4b,9b-(iminoethano)indenol[1,2-b]indol-10-yl]-1H-indole-3-ethanol, 9CI [135091-05-3]



C₃₁H₃₇N₃O 467.653

Related to Borreverine, B-253. Alkaloid from *Borreria verticillata* (Rubiaceae). λ_{\max} 226 ; 285 ; 295 (EtOH) (Berdy). Baldé, A.M. *et al.*, *Phytochemistry*, 1991, **30**, 997 (isol, pmr, cmr, struct)

Spermatherine

S-396

Struct. unknown. Mol. formula not reported. Trace alkaloid from *Atherosperma moschatum* (Monimiaceae). Needles (Me₂CO/petrol). Mp 124-125°. Dec. to a brown mass on exposure to light. Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1956, **9**, 111-119 (isol)

Spermic acid

S-397

N,N'-1,4-Butanediybis- β -alanine, 9CI. 4,9-Diazadodecanedioic acid. N¹,N⁴-Bis(2-carboxyethyl)-1,4-butanediamine. N,N'-Tetramethylenedi- β -alanine. 1,4-Diaminobutane-N,N'-dipropanoic acid [14209-33-7] HOOCCH₂CH₂NH(CH₂)₄NHCH₂CH₂COOH

C₁₀H₂₀N₂O₄ 232.279

Isol. from bovine brain.

Di-Et ester: [87952-71-4]

Powder (as hydrochloride). Mp 237-238° dec. (hydrochloride). CAS no. refers to hydrochloride.

[61345-82-2]

Imaoka, N. *et al.*, *J. Neurochem.*, 1974, **22**, 859 (isol)

Tabor, H. *et al.*, *Methods Enzymol.*, 1983, **94**, 418 (rev, synth)

Wasserman, H.H. *et al.*, *Tetrahedron*, 1983, **39**, 2459 (synth, ester)

Ioannou, P.V. *et al.*, *Chim. Chron.*, 1991, **20**, 85 (synth)

Spermidine

S-398

N-(3-Aminopropyl)-1,4-butanediamine, 9CI. 1,5,10-Triazadecane. 1,8-Diamino-4-azaotane [124-20-9] H₂NCH₂CH₂CH₂CH₂N'HCH₂CH₂CH₂CH₂N'H₂

C₇H₁₉N₃ 145.247

Prod. by many microorganisms including

- the thermophilic eubacteria, *Thermoleophilum album*, *Thermoleophilum minutum*, *Hydrogenobacter thermophilus* and *Bacillus schlegelii*. Isol. from the edible shaggy ink cap mushroom (*Coprinus comatus*) and from *Stichopus japonicus*, *Halocynthia roretzi* and *Crassostrea gigas*. Bp₁₄ 128-130°.
- ▶ LD₅₀ (mus, ivn) 78 mg/kg. EJ7000000
- Hydrochloride (1:3)**: [334-50-9]
Mp 256-258°.
- ▶ EJ7023000
- Tripicrate**:
Yellow needles (H₂O). Mp 211°.
- N-Ac**: N-[3-[(4-Aminobutyl)amino]propyl]acetamide, 9CI. N¹-Acetylspermidine
[14278-49-0]
[34450-16-3]
C₉H₂₁N₃O 187.284
Oil. Mp 191-193° (as dihydrochloride).
- N'-Ac**: N-(4-Aminobutyl)-N-(3-aminopropyl)acetamide. N⁵-Acetylspermidine
[66039-56-3]
C₉H₂₁N₃O 187.284
Mp 187.5-188.5° (as oxalate).
- N''-Ac**: N-[4-[(3-Aminopropyl)amino]butyl]acetamide, 9CI. N¹-(3-Aminopropyl)-N⁴-acetyl-1,4-butanediamine. N⁸-Acetylspermidine
[13431-24-8]
[34450-15-2]
C₉H₂₁N₃O 187.284
Inhibits spermidine-stimulated polyphenylalanine synth. in an *E. coli* cell-free system. Oil or cryst. (as dihydrochloride). Mp 202-203° (EtOH) (dihydrochloride).
- N'-Hexanoyl**: N⁵-Hexanoylspermidine
[97141-37-2]
C₁₃H₂₉N₃O 243.392
Alkaloid from senescing ovaries of pea (*Pisum sativum*).
- N'-Benzoyl**: N-(4-Aminobutyl)-N-(3-aminopropyl)benzamide. N⁵-Benzoylspermidine
[83392-14-7]
C₁₄H₂₃N₃O 249.355
Isol. from leaves of *Oncinotis tenuiloba* (Apocynaceae). Viscous oil.
- N,N''-Dibenzoyl**: N¹,N¹⁰-Dibenzoylspermidine
[73038-07-0]
C₂₁H₂₇N₃O₂ 353.463
Alkaloid from leaves of *Cassia floribunda* (Fabaceae). Needles (EtOH/EtOAc). Mp 133-134°.
- N,N''-Bis(2,3-dihydroxybenzoyl)**: N,N''-Bis(2,3-dihydroxybenzoyl)spermidine
[54135-84-1]
[73038-10-5]
C₂₁H₂₇N₃O₆ 417.461
Alkaloid from *Paracoccus denitrificans* (*Micrococcus denitrificans*). Siderophore (microbial iron-transport agent). Fragment present in Agrobactin, A-203. Cryst. (H₂O)(as hydrochloride). Mp 205-207° (hydrochloride).
- N,N''-Bis(3,4-dihydroxybenzoyl)**: N¹,N¹⁰-Bis(3,4-dihydroxybenzoyl)spermidine. **Pistillarine**
[89647-69-8]
C₂₁H₂₇N₃O₆ 417.461
Alkaloid from *Clavariadelphus pistillar* and several *Ramaria* spp. Also prod. by the marine-derived *Penicillium bilaii*.
- N,N''-Bis(3,4-dihydroxyphenylprop-1-en-1-yl)**: N¹,N¹⁰-Bis(dihydrocaffeoyl)spermidine
[132194-39-9]
C₂₅H₃₃N₃O₆ 473.568
Alkaloid from *Ichroma cyaneum* (Solanaceae).
- Tris-(3-nitrobenzoyl)**: Mp 148-150° (anhyd.).
- N²-(4-Hydroxycinnamoyl)**: **p-Coumaroylspermidine**
[70185-59-0]
C₁₆H₂₅N₃O₂ 291.392
Alkaloid from *Salix* sp. (Salicaceae). Position of coumaroyl group not determined.
- N²-(3,4-Dihydroxycinnamoyl)**: **Caffeoylspermidine**
[58994-15-3]
C₁₆H₂₅N₃O₃ 307.392
Alkaloid from *Nicotiana tabacum* and *Pennisetum americanum* (Solanaceae, Poaceae). Position of caffeoyl group not determined.
- N²-(4-Hydroxy-3-methoxycinnamoyl)**: **Feruloylspermidine**
[70185-60-3]
C₁₇H₂₇N₃O₃ 321.419
Alkaloid from *Ananas comosus* (pineapple) and *Lycopersicon esculentum* (tomato) (Bromeliaceae, Solanaceae). Position of feruloyl group not determined.
- N²-(4-Hydroxy-3,5-dimethoxycinnamoyl)**: **Sinapoylspermidine**
[70185-62-5]
C₁₈H₂₉N₃O₄ 351.445
Alkaloid from *Lilium* sp. (Liliaceae). Position of sinapoyl group not determined.
- N,N''-Di-E-cinnamoyl**: N¹,N⁸-Dicinnamoylspermidine. **Maytenine**
[41590-65-2]
C₂₅H₃₁N₃O₂ 405.539
Alkaloid from *Maytenus chuchuhua*. Isol. from *Pholiota spumosa*. Cryst. (Me₂CO). Mp 161-162° (158°).
- N,N''-Bis(4-hydroxycinnamoyl)**: N¹,N⁵-Di-p-coumaroylspermidine
[114916-04-0]
C₂₅H₃₁N₃O₄ 437.538
Alkaloid from the pollen of *Pterocarya fraxinifolia* (Juglandaceae). A dicoumaroylspermidine of undetd. regiochemistry was isol. from *Alnus glutinosa* (Betulaceae).
- N,N''-Bis(4-hydroxycinnamoyl)**: N¹,N¹⁰-Bis(4-hydroxycinnamoyl)spermidine. N¹,N¹⁰-Dicoumaroylspermidine
[65715-79-9]
[101330-61-4]
C₂₅H₃₁N₃O₄ 437.538
Alkaloid from *Dianthus caryophyllus*, *Helianthus annuus* (sunflower), *Aesculus hippocastanum*, *Vicia faba*, *Pyrus communis* (pear)
- (Caryophyllaceae, Hippocastanaceae, Fabaceae, Rosaceae, Solanaceae).
- N,N''-Bis(3,4-dihydroxycinnamoyl)**: **Dicaffeoylspermidine**
[59001-33-1]
C₂₅H₃₁N₃O₆ 469.536
Alkaloid from *Aesculus hippocastanum*, *Salix* spp. and *Nicotiana tabacum* (Hippocastanaceae, Salicaceae, Solanaceae). Present in fruits of *Solanum melongena* (egg plant).
- N-or N''-(4-Hydroxy-3-methoxycinnamoyl)**, N'' or N-(3,4-dihydroxycinnamoyl): **Caffeoylferuloylspermidine**
[124935-81-5]
C₂₆H₃₃N₃O₆ 483.563
Alkaloid from the pollen of *Corylus avellana* (filbert) (Betulaceae).
- N,N''-Bis(4-hydroxy-3-methoxycinnamoyl)**: N¹,N¹⁰-Diferuloylspermidine
[70185-61-4]
C₂₇H₃₅N₃O₆ 497.59
Alkaloid from the pollen of *Corylus avellana* (filbert) (Betulaceae), also from *Ananas comosus* (pineapple), *Dianthus caryophyllus*, *Vicia faba* and *Lunaria esculentum* (Betulaceae, Bromeliaceae, Caryophyllaceae, Fabaceae, Solanaceae). Only the alkaloid from *C. avellana* has been shown to have the exact struct. shown. The other isolates are diferuloylspermidine with undefined regioisomerism (i.e. could be N⁵-substituted).
- N',N''-Bis(4-hydroxy-3-methoxycinnamoyl)**: N⁵,N¹⁰-Diferuloylspermidine
[115532-15-5]
C₂₇H₃₅N₃O₆ 497.59
Alkaloid from the pollen of *Betula verrucosa* (Betulaceae).
- N,N',N''-Tris(4-hydroxy-E-cinnamoyl)**: N,N',N''-Tris(4-hydroxycinnamoyl)spermidine. N,N',N''-Tris(E-p-coumaroyl)spermidine. N,N',N''-Tricoumaroylspermidine
[131086-78-7]
C₃₄H₃₇N₃O₆ 583.683
Constit. of *Artemisia caruifolia*, *Carthamus tinctorius* (safflower), *Crataegus* spp. and the flowers of *Arachis hypogaea* (peanut). Cryst. (MeOH).
- N''-(3,4-Dihydroxycinnamoyl)**, N,N''-bis(4-hydroxycinnamoyl): N⁵-Caffeoyl-N¹,N¹⁰-di-p-coumaroylspermidine
[167868-57-7]
C₃₄H₃₇N₃O₇ 599.682
Isol. from pollen of *Quercus dentata* (Fagaceae).
- N''-(3,4-Dihydroxycinnamoyl)**, N,N''-bis(4-hydroxycinnamoyl): N¹⁰-Caffeoyl-N¹,N⁵-di-p-coumaroylspermidine
[181264-26-6]
C₃₄H₃₇N₃O₇ 599.682
Minor constit. of *Quercus dentata* pollen.
- N',N''-Bis(3,4-dihydroxycinnamoyl)**, N-(4-hydroxycinnamoyl): N⁵,N¹⁰-Dicaffeoyl-N¹-p-coumaroylspermidine
[167868-56-6]
C₃₄H₃₇N₃O₈ 615.682
Isol. from pollen of *Quercus dentata* (Fagaceae).

- N,N',N'' -Tris(3,4-dihydroxycinnamoyl): N^1,N^5,N^{10} -Tricaffeoylspermidine [167868-55-5]
 $C_{34}H_{37}N_3O_9$ 631.681
 Isol. from pollen of *Quercus dentata* (Fagaceae).
- N'' -(4-Hydroxy-3-methoxycinnamoyl), N -(3,4-dihydroxycinnamoyl): N^1 -Caffeoyl- N^{10} -feruloylspermidine [101164-74-3]
 $C_{26}H_{33}N_3O_6$ 483.563
 Alkaloid from *Corylus avellana* (filbert) (Betulaceae) pollen.
- N -(4-Hydroxy-3-methoxycinnamoyl), N',N'' -bis(3,4-dihydroxycinnamoyl): N^5,N^{10} -Dicaffeoyl- N^1 -feruloylspermidine
 $C_{35}H_{39}N_3O_9$ 645.708
 Minor constit. of *Quercus dentata* pollen.
- N -(4-Hydroxy-3-methoxy-E-cinnamoyl), N',N'' -bis(4-hydroxy-E-cinnamoyl): N^5,N^{10} -Di-p-coumaroyl- N^1 -feruloylspermidine. Keayanidine A [927818-88-0]
 $C_{35}H_{39}N_3O_7$ 613.709
 Constit. of the roots of *Microdesmis keayana*. Yellow powder. λ_{max} 219 (log ϵ 4.66); 232 (log ϵ 4.65); 294 (log ϵ 4.73); 317 (log ϵ 4.82) (MeOH).
- N'' -(4-Hydroxy-3-methoxycinnamoyl), N' -(3,4-dihydroxycinnamoyl), N -(4-hydroxycinnamoyl): N^5 -Caffeoyl- N^1 -p-coumaroyl- N^{10} -feruloylspermidine [181264-10-8]
 $C_{35}H_{39}N_3O_8$ 629.708
 Minor constit. of *Quercus dentata* pollen.
- N,N'' -Bis(4-hydroxy-3,5-dimethoxycinnamoyl): Disinapoylspermidine [70185-63-6]
 $C_{29}H_{39}N_3O_8$ 557.642
 Alkaloid from *Lilium* sp. (Liliaceae). Regioisomerism not rigorously established.
- N,N'' -Bis(4-hydroxy-3-methoxy-E-cinnamoyl), N' -(4-hydroxy-E-cinnamoyl): N^5 -p-Coumaroyl- N^1,N^{10} -diferuloylspermidine. Keayanidine B [927818-89-1]
 $C_{36}H_{41}N_3O_8$ 643.735
 Constit. of the roots of *Microdesmis keayana*. Yellow powder. λ_{max} 220 (log ϵ 4.71); 232 (log ϵ 4.7); 295 (log ϵ 4.75); 319 (log ϵ 4.81) (MeOH).
- N,N',N'' -Tris(4-hydroxy-3-methoxy-E-cinnamoyl): N^1,N^5,N^{10} -Triferuloylspermidine. Keayanidine C [364368-18-3]
 $C_{37}H_{43}N_3O_9$ 673.761
 Constit. of *Hippeastrum x hortorum* and *Microdesmis keayana*. Uv absorbant. Yellow powder. λ_{max} 320 (no solvent reported).
- N'' -(3,5-Dibromo-4-methoxy-E-cinnamoyl): N^{10} -(3,5-Dibromo-4-methoxycinnamoyl)spermidine. Tokaradine C [368422-40-6]
 $C_{17}H_{25}Br_2N_3O_2$ 463.211
 Isol. from the sponge *Pseudoceratina purpurea*. Amorph. yellow solid. λ_{max} 229 (ϵ 12200); 279 (ϵ 9900) (MeOH).
- N' -Me: N -(3-Aminopropyl)- N -methyl-1,4-butanediimine, 9CI. N^5 -Methylspermidine [51460-23-2]
 $C_8H_{21}N_3$ 159.274
 Alkaloid from *Trapa natas*. Oil. Mp 205° dec. (as tripicrate). Bp₁₀ 119-121°.
- N,N -Di-Me, N' -E-cinnamoyl, N'' -benzoyl: Dovyacin F
 $C_{25}H_{33}N_3O_2$ 407.555
 Alkaloid from *Dovyalis abyssinica*. Gum. [49721-50-8]
 Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 309D; 310A (ir)
 Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 501B (nmr)
 v. Braun, J. et al., Ber., 1937, 70, 1230-1240 (synth)
 Danzig, M. et al., J.A.C.S., 1952, 74, 1836-1837 (synth)
 Herbst, E.J. et al., Arch. Biochem. Biophys., 1958, 75, 171-177 (isol)
 Raina, A. et al., Acta Chem. Scand., 1970, 24, 3061-3064 (biosynth)
 Husson, H.-P. et al., C. R. Hebd. Seances Acad. Sci. Ser. C, 1973, 276, 1039 (Maytenine, synth)
 Englert, G. et al., Helv. Chim. Acta, 1973, 56, 474-478 (Maytenine)
 Schlittler, E. et al., Helv. Chim. Acta, 1973, 56, 1097-1099 (N^5 -Methylspermidine, Maytenine, synth)
 Delétang, J. et al., Ann. Tab., Sect. 2, 1974, 11, 123-130; CA, 84, 147656m (Caffeoylspermidine, Dicaffeoylspermidine, Dicomaroylspermidine)
 Williams-Ashman, H.G. et al., CA, 1974, 81, 23332w (rev)
 Tait, G.H. et al., Biochem. J., 1975, 146, 191-204 (bis(2,3-dihydroxybenzoyl))
 Tait, G.H. et al., Biochem. Soc. Trans., 1976, 4, 610-612 (biosynth)
 Cabanne, F. et al., Physiol. Veg., 1977, 15, 429-443; CA, 88, 86095m (Dicoumaroylspermidine)
 Martin-Tanguy, J. et al., Phytochemistry, 1978, 17, 1927-1928 (Coumaroylspermidine, Feruloylspermidine, Sinapoylspermidine, Dicomaroylspermidine)
 Bhargava, K.K. et al., J. Pharm. Sci., 1980, 69, 986-989 (bis(2,3-dihydroxybenzoyl), synth, pmr)
 Bergeron, R.J. et al., J.O.C., 1980, 45, 1589-1592; 1981, 46, 3712-3718; 4524-4529 (bis(2,3-dihydroxybenzoyl), synth, pmr)
 Nagao, Y. et al., Tet. Lett., 1980, 21, 841-844 (Maytenine, synth)
 Tice, C.M. et al., J.O.C., 1983, 48, 2106-2108 (N-Ac, synth, ir, pmr)
 Joshua, A.V. et al., Tet. Lett., 1984, 25, 5725-5728 (Dibenzoylspermidine, Bis(2,3-dihydroxybenzoyl)spermidine, synth)
 Steglich, W. et al., Z. Naturforsch., C, 1984, 39, 10-12 (Pistillarine)
 Meurer, B. et al., Phytochemistry, 1986, 25, 433-435; 1988, 27, 839 (Dicoumaroylspermidine, Diferuloylspermidine, Caffeoylferuloylspermidine)
 Murahashi, S. et al., Chem. Lett., 1987, 879-882 (bis(2,3-dihydroxybenzoyl), synth)
 Miyasaka, T. et al., J.C.S. Perkin 2, 1987, 1543-1549 (bis(2,3-dihydroxybenzoyl), synth)
 Alemayehu, G. et al., Phytochemistry, 1988, 27, 3255-3258 (N^5,N^{10} -Dibenzoylspermidine)
 Almeida, M.L.S. et al., Acta Chem. Scand., 1989, 43, 990-994 (N'-Ac, synth, pmr, cmr)
- Meurer-Grimes, B. et al., Z. Naturforsch., C, 1989, 44, 635-640 (Caffeoylferuloylspermidine)
 Koumoto, Y. et al., Chem. Pharm. Bull., 1990, 38, 1648-1652 (N'-Ac, activity)
 Strack, D. et al., Phytochemistry, 1990, 29, 2893-2896 (iris-4-hydroxycinnamoyl)
 Sattar, E.A. et al., Phytochemistry, 1990, 29, 3931-3933 (N^1,N^{10} -Bis(dihydrocaffeoyl)spermidine)
 Hamana, K. et al., Comp. Biochem. Physiol., B: Comp. Biochem., 1991, 100, 59-62 (occur, hplc)
 Hamana, K. et al., Biochem. J., 1992, 284, 741-747 (thermophilic eubacteria constits)
 Polyamines in the Gastrointestinal Tract, (eds. Dowling, R.H. et al), Kluwer Academic Publishers, Dordrecht, 1992, (book)
 Doll, M.K.-H. et al., Helv. Chim. Acta, 1994, 77, 1229-1235 (N^5 -Benzoylspermidine)
 Bokern, M. et al., Phytochemistry, 1995, 39, 1371-1375 (*Quercus dentata* constits)
 Nimtz, M. et al., Phytochemistry, 1996, 43, 487-489 (*Quercus dentata* constits)
 Perez-Amador, M.A. et al., Plant Physiol., 1996, 110, 1177-1186 (Hexanoylspermidine)
 Hernandez-Jover, T. et al., J. Agric. Food Chem., 1997, 45, 2098-2102 (occur)
 Hamana, K. et al., Can. J. Bot., 1998, 76, 130-133 (N^5 -Methylspermidine, isol)
 Kalač, P. et al., Biologically-active Phytochemicals in Food, (eds. Pfannhauser, W. et al), Royal Society of Chemistry, 2001, 217-220 (sauerkraut)
 Ma, C.-M. et al., Chem. Pharm. Bull., 2001, 49, 915-917 (tri-p-coumaroyl)
 Fusetani, N. et al., Tetrahedron, 2001, 57, 7507-7511 (Tokaradine C)
 Youhnovski, N. et al., Z. Naturforsch., C, 2001, 56, 526-530 (Triferuloylspermidine)
 Whitaker, B.D. et al., J. Agric. Food Chem., 2003, 51, 3448-3454 (Dicaffeoylspermidine, occur, pmr)
 Zamble, A. et al., Chem. Biodiversity, 2006, 3, 982-989 (Keayanidines A-C)
 Rasmussen, B. et al., J. Nat. Prod., 2006, 69, 1300-1304 (Dovyacin F)
 Clericuzio, M. et al., Eur. J. Org. Chem., 2007, 5551-5559 (Maytenine)
 Capon, R.J. et al., J. Nat. Prod., 2007, 70, 1764-1752 (Pistillarine)
 Sobolev, V.S. et al., J. Agric. Food Chem., 2008, 56, 2960-2969 (peanut coumarates)
 Jiang, J.-S. et al., J. Asian Nat. Prod. Res., 2008, 10, 447-451 (safflower coumarates)
 Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 10th edn., J. Wiley, 2000, SLA000

Spermine

S-399

N,N' -Bis(3-aminopropyl)-1,4-butanediimine, 9CI. Bis(3-aminopropyl)tetramethylenediamine. 1,12-Diamino-4,9-diazadodecane. 1,5,10,14-Tetraazatetradecane. Gerontine. Neuridine. Musculamine [71-44-3]

$H_2NCH_2CH_2CH_2^{\oplus}NH(CH_2)_4NHCH_2CH_2CH_2NH_2$

$C_{10}H_{26}N_4$ 202.342

Occurs as phosphate in semen, ox pancreas, yeast and meat prods. Isol. from *Anthocidaris crassispina*, *Stichopus japonicus*, *Halocynthia roretzi*, *Tapes philippinarum* and venom of the spider *Agelenopsis aperta*. Deliquescent cryst. Bp_{0.5} 141-142°. Absorbs CO₂ from the air.

▶ LD₅₀ (mus, ivn) 56 mg/kg. EJ7175000 Hydrochloride (1:4): [306-67-2]

- Cryst. (EtOH). Mp 312-314.5°.
- EJ7230000
- Phosphate:** [3891-79-0] Sol. hot H₂O, insol. EtOH, Et₂O. Mp 230-234°.
- N¹-Ac:** *N*-Acetylspermine. *Monoacetylspermine* [25593-72-0] C₁₂H₂₈N₄O 244.379 Inhibits oat and maize polyamine oxidase. No phys. props. reported.
- Tetrabenzoyl:** Needles (Me₂CO aq.). Mp 155°.
- N¹,N¹⁴-Bis(4-hydroxy-3-methoxyphenylacetyl): 1,14-Bis(homovanillyl)spermine. Orthidine F** C₂₈H₄₂N₄O₆ 530.663 Alkaloid from *Aplidium orthium*. Pale yellow gum. λ_{max} 206 (log ε 4.1); 230 (log ε 3.46); 285 (log ε 3.1) (MeOH).
- N¹,N¹⁰-Bis[3-(3,4-dihydroxyphenyl)propionyl]: 1,10-Bis(dihydrocaffeoyl)spermine. Kukoamine B** [164991-67-7] C₂₈H₄₂N₄O₆ 530.663 Alkaloid from root bark of *Lycium chinense* (Chinese boxthorn) (Solanaceae). Pale yellow powder.
- N¹,N¹⁴-Bis[3-(3,4-dihydroxyphenyl)propionyl]: 1,14-Bis(dihydrocaffeoyl)spermine. Kukoamine A** [75288-96-9] C₂₈H₄₂N₄O₆ 530.663 Alkaloid from the root bark of *Lycium chinense* (Chinese boxthorn) and *Solanum tuberosum* (potatoes). Shows hypotensive activity. Noncryst. λ_{max} 284 (possibly MeOH, not reported) (De-rep).
- N¹,N⁵,N¹⁰-Tris(4-hydroxy-E-cinnamoyl): N¹,N⁵,N¹⁰-Tris-trans-p-coumaroylspermine** [402598-51-0] C₃₇H₄₄N₄O₆ 640.778 Constit. of *Matricaria chamomilla* (German chamomile). Yellow powder (as trifluoroacetate). Mp 109-111° dec. (trifluoroacetate).
- N¹,N⁵,N¹⁰,N¹⁴-Tetakis(4-hydroxy-E-cinnamoyl): N¹,N⁵,N¹⁰,N¹⁴-Tetra-trans-p-coumaroylspermine** [364048-95-3] C₄₆H₅₀N₄O₈ 786.923 Constit. of *Matricaria chamomilla* (German chamomile) and other species of Asteraceae. Tachykinin NK₁ receptor antagonist. Cryst. (MeOH aq.). Mp 139-141° dec. λ_{max} 227 (log ε 4.7); 299 (log ε 4.93); 311 (log ε 4.94) (MeOH).
- N¹,N¹⁴-Bis(4-hydroxy-3-methoxy-E-cinnamoyl), N⁵-(4-hydroxy-E-cinnamoyl): N⁵-trans-p-Coumaroyl-N¹,N¹⁴-di-trans-feruloylspermine. Keayanine A** [942914-01-4] C₃₉H₄₈N₄O₈ 700.83 Alkaloid from the roots of *Microdesmis keayana*. Yellow powder. λ_{max} 218 (log ε 4.78); 232 (log ε 4.72); 293 (log ε 4.64); 315 (log ε 4.68) (MeOH).
- Bis(4-hydroxy-3-methoxycinnamoyl): Di-feruloylspermine** [70208-06-9]


- C₃₀H₄₂N₄O₆ 554.685 Alkaloid from *Ananas comosus* (pine-apple) and *Gomphrena globosa*. Substituted on two of the N atoms. Exact struct. unknown.
- N-(4-Hydroxy-3,5-dimethoxycinnamoyl): Sinapoyspermine** [70185-65-8] C₂₁H₃₆N₄O₄ 408.54 Alkaloid from *Brassica oleracea* var. *botrytis* (cauliflower) (Brassicaceae). Mixt. of N¹ and/or N⁵-substitution.
- Bis(4-hydroxy-3,5-dimethoxycinnamoyl): Disinapoyspermine** [70185-66-9] C₃₂H₄₆N₄O₈ 614.737 Alkaloid from *Lilium* sp. Substituted on two of the N atoms. Exact struct. unknown.
- N⁵-Hydroxy:** [864812-09-9] C₁₀H₂₆N₄O 218.342 Isol. from venom of the spider *Hololena curta*.
- Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 312B (ir)*
- Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 503C (nmr)*
- Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 399A (ir)*
- Dudley, H.W. et al., *Biochem. J.*, 1924, **18**, 1263-1272; 1926, **20**, 1082-1094; 1927, **21**, 97-103 (*isol. struct*)
- Dubin, D.T. et al., *J. Biol. Chem.*, 1960, **235**, 776-782 (*Monoacetylspermine*)
- Israel, M. et al., *J. Med. Chem.*, 1964, **7**, 710-716 (*synth*)
- Raina, A. et al., *Acta Chem. Scand.*, 1970, **24**, 3061-3064 (*biosynth*)
- Williams-Ashman, M.G. et al., *Adv. Enzyme Regul.*, 1972, **10**, 225-245; *CA*, **81**, 23332w (*rev*)
- Martin-Tanguy, J. et al., *Phytochemistry*, 1978, **17**, 1927-1928 (*Sinapoyspermine, Diferuloylspermine, Disinapoyspermine*)
- Funayama, S. et al., *Tet. Lett.*, 1980, **21**, 1355 (*Kukoamine A*)
- Chantrapromma, K. et al., *Tet. Lett.*, 1981, **22**, 23 (*Kukoamine A, synth*)
- Federico, R. et al., *Phytochemistry*, 1989, **28**, 2045-2046; 1990, **29**, 2411-2414 (*Monoacetylspermine, activity*)
- Hamana, K. et al., *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 59-61 (*occur, hplc, gc*)
- Polyamines in the Gastrointestinal Tract*, (eds. Dowling, R.H. et al), Kluwer Academic Publishers, Dordrecht, 1992, (*book*)
- Funayama, S. et al., *Phytochemistry*, 1995, **38**, 1529 (*Kukoamine B*)
- Hernandez-Jover, T. et al., *J. Agric. Food Chem.*, 1997, **45**, 2098-2102 (*occur*)
- Karigiannis, G. et al., *Tet. Lett.*, 1998, **39**, 5117-5120 (*Kukoamine A, synth*)
- Yamamoto, A. et al., *Chem. Pharm. Bull.*, 2002, **50**, 47-52 (*Tricoumaroylspermine, Tetraoumaroylspermine*)
- Rubio, M. et al., *Synth. Commun.*, 2002, **32**, 2441-2452 (*synth, pmr, cmr*)
- Garnelis, T. et al., *Chem. Lett.*, 2005, **34**, 264-265 (*Kukoamine A, synth*)
- Parr, A.J. et al., *J. Agric. Food Chem.*, 2005, **53**, 5461-5466 (*Kukoamines, isol*)
- Tzouros, M. et al., *Toxicol.*, 2005, **46**, 350-354 (*N-hydroxy*)
- Zamble, A. et al., *Chem. Pharm. Bull.*, 2007, **55**, 643-645 (*Keayanine*)
- Pearce, A.N. et al., *Tetrahedron*, 2008, **64**, 5748-5755 (*Orthidine F*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, DCC400; GEK000

Spermostrychnine

S-400

1-Acetyl-17,19-epoxycuran, 9CI [639-34-9]



Absolute Configuration

C₂₁H₂₆N₂O₂ 338.449 Alkaloid from *Strychnos psilosperma*, *Strychnos aculeata* and *Strychnos brasiliensis* (Loganiaceae). Needles (Me₂CO). Mp 208-209°. [α]_D²² +88 (c, 2.2 in CHCl₃).

► WG8700000

Picrate:

Small yellow hexagonal plates. Mp 173-175° dec.

Methiodide: Mp 301-303°.

N-De-Ac: N-Deacetylspermostrychnine.

Desacetylspermostrychnine

[22153-12-4]

C₁₉H₂₄N₂O 296.411

Alkaloid from root bark of *Strychnos panganensis*. [α]_D -120 (c, 0.48 in CHCl₃). λ_{max} 210; 245; 301 (MeOH).

N-De-Ac, N-hydroxyacetyl: 23-Hydroxy-spermostrychnine

[141544-25-4]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Strychnos henningsii* (Loganiaceae). [α]_D +70 (c, 0.95 in CHCl₃). λ_{max} 212; 250; 280; 290 (MeOH).

N-De-Ac, N-hydroxyacetyl, N⁴-oxide: 23-Hydroxyspermostrychnine N⁴-oxide

[141544-32-3]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Strychnos henningsii*.

17-Hydroxy, N-de-Ac, N-hydroxyacetyl:

17,23-Dihydroxyspermostrychnine

[141544-33-4]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from *Strychnos henningsii* (Loganiaceae). λ_{max} 211; 250; 281; 290 (sh) (MeOH).

11-Methoxy: Strychnospermine

[509-45-5]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Strychnos psilosperma* leaves (Loganiaceae). Needles (MeOH aq.). Mp 208-209°. [α]_D¹⁶ +60.3 (c, 1.37 in CHCl₃).

11-Methoxy, hydrochloride:

Plates (H₂O). Mp 330-332° dec. (chars from 300°). [α]_D¹⁵ +66.6 (c, 1.43 in H₂O).

11-Methoxy, picrate:

Yellow needles (EtOH). Mp 254° dec.

11-Methoxy, 12-hydroxy: 12-Hydroxy-11-methoxyspermostrychnine

[34327-10-1]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from trunk bark of *Strychnos brasiliensis* (Loganiaceae). Small plates

(EtOH). Mp 272-273°. $[\alpha]_D^{25}$ -217 (c, 0.58 in CHCl₃). λ_{\max} 225 (log ϵ 4.36); 255 (log ϵ 3.77); 290 (log ϵ 2.88) (EtOH).

11-Methoxy, 12-hydroxy; methiodide: Cryst. (Me₂CO). Mp 289-291°.

19-Epimer: N²-Acetyl-3-deoxystrychnosplendine

[22153-15-7]
C₂₁H₂₆N₂O₂ 338.449

Minor alkaloid detected in the bark of *Strychnos splendens* (Loganiaceae). Cryst. (CHCl₃). Mp 280-283°. $[\alpha]_D^{20}$ +98 (c, 1 in EtOH). λ_{\max} 218 (log ϵ 4.41); 250 (log ϵ 4.05); 277 (sh) (log ϵ 3.38) (no solvent reported).

19-Epimer, N-de-Ac, N-hydroxyacetyl: 19-Epi-23-hydroxyspermostrychnine

[141544-26-5]
C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Strychnos hemingsii* (Loganiaceae). λ_{\max} 212; 250; 280; 290 (MeOH).

Anet, F.A.L. *et al.*, *Aust. J. Chem.*, 1953, **6**, 58-65 (*Spermostrychnine, Strychnospermine, isol*)

Anet, F.A.L. *et al.*, *J.C.S.*, 1955, 2253-2262 (*Spermostrychnine, Strychnospermine, uv, ir, struct*)

Biemann, K. *et al.*, *Mass Spectrom. [SPR]*, McGraw-Hill, 1962, 331 (*Strychnospermine, ms*)

Anet, F.A.L. *et al.*, *Can. J. Chem.*, 1963, **41**, 883-888 (*Spermostrychnine, Strychnospermine, pmr*)

Plat, M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1968, **267**, 1419-1422 (*pmr, config*)

Koch, M. *et al.*, *Ann. Pharm. Fr.*, 1969, **27**, 229-238 (*N²-Acetyl-3-deoxystrychnosplendine*)

Koch, M. *et al.*, *Tetrahedron*, 1969, **25**, 3377-3382 (*config*)

Iwataki, I. *et al.*, *Tetrahedron*, 1971, **27**, 2541-2552 (*12-Hydroxy-11-methoxyspermostrychnine*)

Wenkert, E. *et al.*, *J.O.C.*, 1978, **43**, 1099-1105 (*Strychnospermine, pmr, cmr*)

Massiot, G. *et al.*, *Phytochemistry*, 1991, **30**, 3449-3456 (*23-Hydroxyspermostrychnine, 19-Epi-23-hydroxyspermostrychnine, 1,23-Dihydroxyspermostrychnine, 23-Hydroxyspermostrychnine N-oxide, 17,23-Dihydroxyspermostrychnine*)

Nuzillard, J.-M. *et al.*, *Phytochemistry*, 1996, **43**, 897-902 (*N-Deacetylspermostrychnine*)

Sphaeranthine† S-401

C₁₃H₁₉NO₅ 269.297

Struct. unknown. Alkaloid from *Sphaeranthus indicus* (Asteraceae). Mp 166-168° dec.

Hydrochloride: Mp 158-160° dec.

Sulfate: Mp 185° dec.

Oxalate: Mp 110-111° dec.

Picrate: Mp 154-156° dec.

Chloroplatinate: Mp 196-198° dec.

Basu, N.K. *et al.*, *J. Am. Pharm. Assoc.*, 1946, **35**, 274 (*isol*)

Sphaeranthine I† S-402

C₁₄H₈N₂O₄ 268.228

Struct. unknown. Alkaloid from the flowers of *Sphaeranthus indicus*. Yellowish needles. Mp 190°.

Naqvi, S.B.S. *et al.*, *Thesis*, Univ. of Karachi, 1997, (*isol, uv, ir, pmr, cmr, ms*)

Sphaeranthine IV S-403

C₁₃H₁₇NO₆ 283.28

Struct. unknown. Alkaloid from flowers of *Sphaeranthus indicus*. Mp 125°.

Naqvi, S.B.S. *et al.*, *Thesis*, Univ. of Karachi, 1997, (*isol, uv, ir, pmr, cmr, ms*)

Sphaeranthine VI S-404

C₁₇H₁₈N₂O 266.342

Struct. unknown. Alkaloid from flowers of *Sphaeranthus indicus*. Cryst. Mp 135°.

Naqvi, S.B.S. *et al.*, *Thesis*, Univ. of Karachi, 1997, (*isol, uv, ir, pmr, cmr, ms*)

Sphaeranthine VII S-405


C₁₂H₈N₂O₅ 260.206

Struct. unknown. Alkaloid from flowers of *Sphaeranthus indicus*. Mp 170°. λ_{\max} 220; 280 (no solvent reported).

Naqvi, S.B.S. *et al.*, *Thesis*, Univ. of Karachi, 1997, (*isol, uv, ir, pmr, cmr, ms*)

Sphaerophysine S-406

N-(4-Aminobutyl)-N-(3-methyl-2-butenyl)guanidine, 9CI. N-(4-Aminobutyl)-N-prenylguanidine. *Sphaerophysine* [25978-54-5]



C₁₀H₂₂N₄ 198.311

Two incorrect structs. were previously proposed. Alkaloid from the leaves of *Sphaerophysa salsula* and from *Galega officinalis*, *Eremosparton flocciodum* and *Swainsona galegifolia* (Fabaceae). Ganglion blocker. V. hygroscopic solid.

Dipicrate: Mp 154-155°.

N⁴-Ac: **Smirnovine**, *Smyrnovine* [2758-69-2]

C₁₂H₂₄N₄O 240.348

Alkaloid from roots of *Smirnowia turkestanica* and from *Astragalus* sp. and *Galega officinale* (Fabaceae). Golden needles (as picrate). Mp 154° (picrate).

N⁴-(Carboxyacetyl): **Smirnovinine**.

Smyrnovinine

[27586-70-5]

C₁₃H₂₄N₄O₃ 284.358

Alkaloid from *Smirnowia turkestanica* and *Astragalus* sp. (Fabaceae). Mp 145° dec. (as picrate).

4-Hydroxy (2Z-), N⁴-Ac: **4-Hydroxysmirnovine**

[180156-26-7]

C₁₂H₂₄N₄O₂ 256.347

Alkaloid from *Galega orientalis*. Oil. [70293-13-9]

Rubinschtein, M.M. *et al.*, *CA*, 1945, **39**, 2291⁴ (*isol*)

Rjabinin, A.A. *et al.*, *CA*, 1948, **42**, 4718f; 1951, **45**, 8458 (*derivis*)

Reuter, G. *et al.*, *Planta Med.*, 1965, **13**, 494 (*isol*)

Heesing, A. *et al.*, *Chem. Ber.*, 1970, **103**, 534 (*struct, ms*)


Steinger, J.R.G. *et al.*, *Pharmazie*, 1973, **28**, 682; 1974, **29**, 422 (*isol, biosynth*)

Benn, M.H. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 2286-2288 (*4-Hydroxysmirnovine, isol*)

Yadav, V.K. *et al.*, *Can. J. Chem.*, 2000, **78**, 542-545 (*4-Hydroxysmirnovine, synth*)

Spicaledonine S-407

[126617-13-8]



C₃₇H₄₉NO₁₀ 667.795

Alkaloid from the bark of *Austrotaxus spicata* (Taxaceae). Amorph. $[\alpha]_D$ +29 (c, 0.3 in CHCl₃).

2'-Deoxy: Taxuspine H

[164991-81-5]

C₃₇H₄₉NO₉ 651.795


Isol. from *Taxus cuspidata* and *Taxus brevifolia*. Microtubule depolymerisation inhibitor. Amorph. solid. $[\alpha]_D^{23}$ +6.8 (c, 0.29 in CHCl₃). λ_{\max} 230 (ϵ 14000) (MeOH) (Berdy).

Ettouati, L. *et al.*, *Bull. Soc. Chim. Fr.*, 1989, 687-694 (*isol, pmr, cmr, struct*)

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**, 5971-5978 (*Taxuspine H*)

Spicatine† S-408

4-Ethenyl-3-ethoxy-1,3,4,5,6,7-hexahydro-8H-pyrano[3,4-c]pyridin-8-one, 9CI [11053-08-0]



C₁₂H₁₇NO₃ 223.271

Monoterpene alkaloid. Alkaloid from *Centaurium spicatum* (Gentianaceae). Mp 182-183°.

$\Delta^{1,8a}$ -Isomer: [70474-61-2]

C₁₂H₁₇NO₃ 223.271

Alkaloid from *Centaurium spicatum* (Gentianaceae). Mp 149-150°.

Khafagy, S.M. *et al.*, *Acta Pharm. Suec.*, 1968, **5**, 135; *CA*, **69**, 80134j


Bishay, D. *et al.*, *J. Pharm. Pharmacol.*, 1978, **30**, Suppl.; 80P (*struct*)

Bishay, D. *et al.*, *Planta Med.*, 1979, **37**, 253 (*deriv*)

Spicigerine

S-409

5-Hydroxy-6-methyl-2-piperidinedodecanoic acid, 9CI
[52998-86-4]



Absolute Configuration

$C_{18}H_{35}NO_3$ 313.479
Alkaloid from *Prosopis spicigera* (Fabaceae). Mp 196°.

N,O-di-Ac, Me ester:
Oil. $[\alpha]_D^{26}$ -17.2 (c, 1.0 in $CDCl_3$).

(±)-form


Me ester: [87243-58-1]
Mp 69-71.2°. Bp_{0.001} 160°.

Jewers, K. *et al.*, *Phytochemistry*, 1976, **15**, 238-240 (*isol*)
Paterne, M. *et al.*, *J. Chem. Res., Synop.*, 1985, 278 (*synth, ms, ir, pmr*)
Hasseberg, H.-A. *et al.*, *Annalen*, 1989, 255 (*synth, ir, pmr, cmr*)
Momose, T. *et al.*, *Heterocycles*, 1995, **40**, 137 (*abs config*)
Leverett, C.A. *et al.*, *J.O.C.*, 2006, **71**, 8591-8601 (*synth*)

Spiduxine

S-410

[112494-59-4]

 $C_{21}H_{23}NO_5$ 369.416**(ξ)-form**


Alkaloid from the trunk bark of Colombian *Duguetia spixiana* (Annonaceae). Amorph.

Debourges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664 (*isol, uv, ir, pmr, ms, struct*)

Spiguetidine

S-411

5,6,7,7a-Tetrahydro-7-methylbenzo[d]-1,3-dioxolo[4,5-g]pyrido[4,3,2-jk][2]benzazepin-11-ol, 9CI
[112494-57-2]



$C_{18}H_{16}N_2O_3$ 308.336
Rearranged aporphine (azahomoaporphine).

(ξ)-form

Alkaloid from the bark of Bolivian *Duguetia spixiana* (Annonaceae).

Me ether: Spiguetine

[112494-56-1]
 $C_{19}H_{18}N_2O_3$ 322.363
Alkaloid from *Duguetia spixiana* (Annonaceae). $[\alpha]_D$ -2 (c, 0.15 in EtOH).

Deoxy: Dragabine

[107030-45-5]
 $C_{18}H_{16}N_2O_2$ 292.337
Alkaloid from the bark and leaves of *Guatteria sagotiana* (Annonaceae). Noncryst. Insignificant opt. rotn.

Deoxy, N-de-Me: Nordragabine


[107030-44-4]
 $C_{17}H_{14}N_2O_2$ 278.31
Trace alkaloid from the trunk bark of *Meiogyne virgata* (Annonaceae). Noncryst. Undergoes considerable dec. on standing.

Cassels, B.K. *et al.*, *Chem. Comm.*, 1986, 1481 (*Dragabine, Nordragabine*)
Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1987, **50**, 674 (*Spiguetidine, Spiguetine*)

Spinanine A

S-412

[135091-07-5]



Absolute Configuration


$C_{24}H_{32}N_4O_4$ 440.541
Alkaloid from the stem bark of *Zizyphus spinachristi* (Rhamnaceae). Cryst. Mp 175-176°. $[\alpha]_D$ -121 (c, 0.1 in MeOH).

Abdel-Galil, F.M. *et al.*, *Phytochemistry*, 1991, **30**, 1348 (*isol, uv, ir, ms, struct*)

Spiraeaine A

S-413

[473714-84-0]



$C_{22}H_{33}NO_3$ 359.508
Alkaloid from the stems of *Spiraea formosana*. Oil. $[\alpha]_D$ -7.8 (c, 0.7 in $CHCl_3$).

19-Oxo, 6-Ac: Spiramine Y

$C_{24}H_{33}NO_5$ 415.528
Alkaloid from *Spiraea japonica*. Needles (Me_2CO). Mp 144-145°. $[\alpha]_D^{27}$ -152 (c, 0.75 in $CHCl_3$).

19-Oxo, di-Ac: Spiramine X

$C_{26}H_{35}NO_6$ 457.566
Alkaloid from *Spiraea japonica*. Needles (Me_2CO). Mp 125-127°. $[\alpha]_D^{25}$ -139.7 (c, 0.23 in $CHCl_3$).


Wang, B.-G. *et al.*, *Heterocycles*, 2000, **53**, 1343-1350 (*Spiramines X and Y*)

Wu, T.S. *et al.*, *Heterocycles*, 2002, **57**, 1495-1500 (*Spiraeaine A*)

Spirafine II

S-414

[20071-95-8]



$C_{22}H_{31}NO_2$ 341.492
Alkaloid from *Spiraea fritschiana* var. *parvifolia*. Needles. Mp 155-156°. $[\alpha]_D^{22}$ -33.2 (c, 2 in Py).

5-Epimer, Δ¹⁷-isomer: Spirafine III


[20072-00-8]
 $C_{22}H_{31}NO_2$ 341.492
Alkaloid from *Spiraea fritschiana* var. *parvifolia*. Needles. Mp 192-193°. $[\alpha]_D^{22}$ -46.1 (c, 2 in Py).

Li, M. *et al.*, *Chin. Chem. Lett.*, 1999, **10**, 827-830

Spirajine

S-415

17-Hydroxy-20-(methylamino)-19,21-dinorchole-4,11-diene-3,22-dione, 9CI
[21728-69-8]




$C_{23}H_{33}NO_3$ 371.519
No details in abstract; struct. is taken from CA details. Unprecedented struct. Alkaloid from leaves of *Spiraea koreana* (Rosaceae). Mp 182-184°. $[\alpha]_D^{19}$ +3.4 ($CHCl_3$).

Kab Duck Jin, *et al.*, *CA*, 1969, **70**, 26359

Spiramine C

S-416

[114485-97-1]



$C_{22}H_{31}NO_3$ 357.492
Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae). Cryst. (Et_2O). Mp 167-169°. $[\alpha]_D^{25}$ -149.9 (c, 1.0 in C_6H_6).

15-Ac: Spiramine A

[114531-28-1]
 $C_{24}H_{33}NO_4$ 399.529

Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae). Shows antiplatelet aggregation activity. Cryst. (hexane). Mp 137.5-139°. $[\alpha]_D^{25}$ -103.1 (c, 0.9 in C₆H₆).

19-Epimer: Spiramine D

[114485-98-2]

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae). Cryst. (Et₂O). Mp 160-162°. $[\alpha]_D^{25}$ -169 (c, 0.7 in C₆H₆).

19-Epimer, 15-Ac: Spiramine B

[114485-96-0]

C₂₄H₃₃NO₄ 399.529

Alkaloid from *Spiraea japonica* var. *acuminata* (Rosaceae). Shows antiplatelet aggregation activity. Cryst. (hexane). Mp 129-131°. $[\alpha]_D^{25}$ -159.5 (c, 0.9 in C₆H₆).

Hao, X. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 1670 (*ir, pmr, cmr, cryst struct*)


Node, M. *et al.*, *Heterocycles*, 1990, **30**, 635 (*isol, ir, pmr, cmr, ms, cryst struct*)

Wang, B.-G. *et al.*, *Heterocycles*, 2000, **53**, 1343-1350 (*activity*)

Spiramine J

S-417

[147395-98-0]



C₂₃H₃₃NO₃ 371.519

C-19 has (S)-config. Alkaloid from the roots of *Spiraea japonica* var. *acuminata* (Rosaceae).

7-Ac: Spiramine M

[137760-59-9]

C₂₅H₃₅NO₄ 413.556

Alkaloid from the roots of *Spiraea japonica* var. *acuminata* (Rosaceae).

15-Ac: Spiramine L

[137760-58-8]

C₂₅H₃₅NO₄ 413.556

Alkaloid from the roots of *Spiraea japonica* var. *acuminata* (Rosaceae).

19-Epimer: Spiramine K

[137760-57-7]

C₂₃H₃₃NO₃ 371.519


Alkaloid from the roots of *Spiraea japonica* var. *acuminata* (Rosaceae).

Hao, X. *et al.*, *CA*, 1993, **118**, 209386

Spiramine P

S-418

[162901-90-8]



C₂₂H₃₃NO₄ 375.507

Alkaloid from roots of *Spiraea japonica* var. *incisa* (Rosaceae). Needles (MeOH). Mp 237-239°. $[\alpha]_D$ -49 (c, 0.81 in CHCl₃). Struct. revised in 2000.

6-Ac: Thalicsiline. Spiramine U

[119420-10-9]

[205871-37-0]

C₂₄H₃₅NO₅ 417.544

Alkaloid from the roots of *Thalictrum sessile* and from roots of *Spiraea japonica* var. *acuta*. Demonstrates antiinflammatory activity in rats. Prisms + 1/2H₂O (MeOH). Mp 183-186°. $[\alpha]_D^{20}$ -11.4 (c, 0.1 in MeOH). Struct. of Spiramine U revised in 2000. Props. refer to Thalicsiline. Exists in soln. as a mixt. of C-19 epimers.

16-Deoxy, 16,17-didehydro: Spiradine G

[21040-66-4]

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 168-170°. $[\alpha]_D$ -137 (c, 1.16 in MeOH). pK_a 6.9 (50% MeOH aq.).

16-Deoxy, 16,17-didehydro, 6-Ac: Spiradine F

[21040-64-2]

C₂₄H₃₃NO₄ 399.529

Alkaloid from *Spiraea japonica* and *Spiraea japonica* var. *acuta* (Rosaceae). Shows antiplatelet aggregation activity. Mp 114-117° (as hydrochloride).

16-Epimer: Spiramine Q

[162993-29-5]

C₂₂H₃₃NO₄ 375.507

Alkaloid from roots of *Spiraea japonica* var. *incisa* (Rosaceae). Platelet aggregation inhibitor, decreases serotonin secretion. Needles. Mp 197-199°. $[\alpha]_D$ -70 (c, 0.84 in CHCl₃). Struct. revised in 2000.

16,19-Diepimer: Spiramine W

[304645-14-5]

C₂₂H₃₃NO₄ 375.507

Constit. of *Spiraea japonica* var. *acuta*.

16,19-Diepimer, 6-Ac: Spiramine T

[205871-36-9]

C₂₄H₃₅NO₅ 417.544

Alkaloid from roots of *Spiraea japonica* var. *acuta*. Needles. Mp 183-185°. $[\alpha]_D^{25}$ -151.6 (c, 0.67 in CHCl₃). Struct. revised in 2000.

Toda, M. *et al.*, *Tet. Lett.*, 1968, **9**, 5565-5568 (*Spiradines F,G*)

Wu, Y.-C. *et al.*, *Phytochemistry*, 1988, **27**, 3949-3953 (*Thalicsiline*)

Hao, X.-J. *et al.*, *Phytochemistry*, 1995, **38**, 545-547 (*Spiramines P,Q, isol, ir, pmr, cmr, ms*)

Nie, J.L. *et al.*, *CA*, 1998, **128**, 280808f

(*Spiramines T,U*)

Wang, B.G. *et al.*, *CA*, 2000, **133**, 332129k

(*Spiramine W*)

Wang, B.-G. *et al.*, *Heterocycles*, 2000, **53**, 1343-1350 (*Spiradines F,G, activity*)

Wang, B.G. *et al.*, *J. Asian Nat. Prod. Res.*, 2000, **2**, 271-281 (*Spiramines P,Q,T,U, Spiradine F, struct*)


Shen, Z. *et al.*, *Planta Med.*, 2000, **66**, 287-289 (*Spiramine Q, activity*)

Li, L. *et al.*, *Planta Med.*, 2001, **67**, 142-145 (*Spiramine T*)

Spiramine Z₂

S-419

Spiramidine A



C₂₂H₃₁NO₃ 357.492

Two different sets of names have been assigned by the same authors. Alkaloid from *Spiraea japonica* var. *ovalifolia*. Isol. as a mixture with Spiramine Z₃.

19-Epimer: Spiramine Z₃, Spiramidine B

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Spiraea japonica* var. *ovalifolia*. Isol. as a mixture with Spiramine Z₂. The 19-configs. are ambiguously descr. as 19 α - and 19 β -.

Zuo, G.Y. *et al.*, *Chin. Chem. Lett.*, 2001, **12**, 147-150


Zuo, G.Y. *et al.*, *Heterocycles*, 2001, **55**, 487-493

Spirasine X

S-420

13-Hydroxyhetisan-11-one, 9CI

[112543-29-0]



C₂₀H₂₅NO₂ 311.423


Alkaloid from *Spiraea japonica* (Rosaceae). Needles (Me₂CO). Mp 224-227°. $[\alpha]_D^{17}$ +51 (c, 1.5 in CHCl₃).

Sun, F. *et al.*, *J. Nat. Prod.*, 1987, **50**, 923 (*isol, ir, pmr, cmr, ms, cd, struct*)

Spirasine III

S-421

[112496-48-7]



C₂₂H₂₇NO₄ 369.46

Exists in soln. as a mixt. of C-19 epimers. Alkaloid from *Spiraea japonica* (Rosaceae). Mp 210-212°. $[\alpha]_D^{21}$ -9 (c, 1.1 in CHCl₃).

9-Deoxy: Spiredine. 11-Oxospiradine D, 9CI

[60062-45-5]

C₂₂H₂₇NO₃ 353.46

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 163°.

11-Deoxo: Spirasine II

[106777-12-2]

C₂₂H₂₉NO₃ 355.476

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 208-209°.

11-Deoxo, 9-deoxy: Spiradine D, 9CI

[20071-93-6]

C₂₂H₂₉NO₂ 339.477

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 134-135°. pK_a 9.35 (50% MeOH).

11-Deoxo, Δ¹⁵-isomer: Spirasine I

[106777-13-3]

C₂₂H₂₉NO₃ 355.476

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 244-246°. [α]_D²¹ -131 (c, 1.1 in CHCl₃).

11-Deoxo, 16,17-dihydro, 16α-hydroxy: Spirasine VII

[106798-64-5]

C₂₂H₃₁NO₄ 373.491

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 191-193°. [α]_D²⁸ -78 (c, 1.3 in CHCl₃).

11-Deoxo, 16,17-dihydro, 16β-hydroxy: Spirasine VIII

[106861-54-5]

C₂₂H₃₁NO₄ 373.491

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 207-209°. [α]_D²⁸ -57 (c, 1.0 in CHCl₃).

11-Deoxo, 9-deoxy, 16,17-dihydro, 16α-hydroxy: Spirasine V

[106798-63-4]

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 177-179°. [α]_D²⁸ -47 (c, 0.77 in CHCl₃).

11-Deoxo, 9-deoxy, 16,17-dihydro, 16β-hydroxy: Spirasine VI

[106861-53-4]

C₂₂H₃₁NO₃ 357.492

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 202-203°. [α]_D²¹ -107 (c, 1.0 in CHCl₃).

Goto, G. *et al.*, *Tet. Lett.*, 1968, 2989

(Spiradine D)

Gorbunov, V.D. *et al.*, *Chem. Nat. Compd.*

(*Engl. Transl.*), 1976, **12**, 119-120

(Spiradine)

Sun, F. *et al.*, *Heterocycles*, 1986, **24**, 2105-

2108 (Spirasines I,II,VII,VIII)

Sun, F. *et al.*, *Tet. Lett.*, 1986, **27**, 275-278

(Spirasines V,VI, *ir, cmr, cd, cryst struct*)

Sun, F. *et al.*, *Heterocycles*, 1987, **26**, 19-22


(Spirasine III)

Fan, L.-M. *et al.*, *J. Integ. Plant Biol.*, 2005, **47**,

120-123 (Spiradine, Spirasine V,VI)

Spiratine B

S-422



C₂₄H₃₃NO₅ 415.528

Alkaloid from the roots of *Spiraea japonica*. Amorph. powder. [α]_D²⁵ +129.5 (c, 5 in CHCl₃).

O¹⁹-Et: Spiramine Z

C₂₆H₃₇NO₅ 443.582

Alkaloid from *Spiraea japonica*. Waxy solid. [α]_D²⁷ +81.7 (c, 0.35 in CHCl₃). Possible artifact.


Wang, B.G. *et al.*, *Heterocycles*, 2000, **53**,

1343-1350 (Spiramine Z)

He, H.-P. *et al.*, *J. Nat. Prod.*, 2001, **64**, 379-380 (Spiratine B)

Spireine

S-423



C₂₂H₂₇NO₄ 369.46

Alkaloid from *Spiraea japonica* (Rosaceae). Mp 230°.


Gorbunov, V.D. *et al.*, *Khim. Prir. Soedin.*,

1969, **5**, 454; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 379-380 (*ir, pmr*)

Spiroarogenic acid

S-424

8-Hydroxy-1-oxo-2-azaspiro[4.5]deca-6,9-diene-3-carboxylic acid, 9CI
[86334-99-8]



C₁₀H₁₁NO₄ 209.201

Prod. by *Neurospora crassa*. No phys. props. reported.

[87726-54-3]


Zamir, L.O. *et al.*, *J. Biol. Chem.*, 1983, **258**,

6486 (*isol, pmr, cmr, strut*)

1-Spiroaxen-6-amine

S-425

6-Amino-1-spiroaxene



C₁₅H₂₇N 221.385

(6β,10α)-form

N-Formyl: Axamide 3

[59633-82-8]

C₁₆H₂₇NO 249.395

Constit. of sponges *Axinella cannabina* and *Axinella cavernosa*. Oil. [α]_D -6.86 (c, 1 in CHCl₃).

Isocyanide: Axisonitrile 3

[59633-83-9]

C₁₆H₂₅N 231.38

Isol. from *Axinella cannabina*, *Topsentiasp.*, *Ciocalypta sp.*, *Acanthella sp.*, *Axinyssa aplysinoides* and *Phyllidia pustulosa*. Shows strong antimycobacterial activity. Anthelmintic, antifouling agent, larval settlement inhibitor, spongicide. Cryst. (petrol). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 101-103°. [α]_D +68.44 (c, 1 in CHCl₃). Has -NCS replacing -NH₂.

Isothiocyanate: Axisothiocyanate 3

[59633-81-7]

C₁₆H₂₅NS 263.446

Isol. from *Axinella cannabina* and *Axinella cavernosa*. Oil. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D +165.2 (c, 1 in CHCl₃). Has -NCS replacing -NH₂.

3-Oxo, isocyanide: 3-Oxoaxisonitrile 3

[922518-37-4]

C₁₆H₂₃NO 245.364

Constit. of an *Acanthella sp.* Oil. [α]_D²⁰ +70 (c, 0.13 in CHCl₃).

(6β,10β)-form [509078-99-3]

N-Formyl: Exiguamide

[481054-36-8]

C₁₆H₂₇NO 249.395

Constit. of *Geodia exigua*. Cryst.

(MeOH aq.). Mp 139-140°. [α]_D²⁵ +31.7 (c, 0.08 in CHCl₃).

N-Methoxycarbonyl: Exicarbamate

[509095-16-3]

C₁₇H₂₉NO₂ 279.422

Constit. of *Geodia exigua*. Oil. [α]_D²⁵ +28 (c, 0.02 in CHCl₃).

N-[(N-Methoxycarbonylmethyl)-N-methylamino]acetyl: Exigurin

[509095-20-9]

C₂₁H₃₆N₂O₃ 364.527

Constit. of *Geodia exigua*. Oil. [α]_D²⁵ -32 (c, 0.03 in CHCl₃).

Isocyanide: 10-Epiaxisonitrile 3

[178963-62-7]

C₁₆H₂₅N 231.38

Constit. of *Geodia exigua*, *Phyllidia pustulosa* and *Phyllidia acelata*. Antifouling agent; larval settlement inhibitor. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²³ -5.6 (c, 0.025 in CHCl₃).

Blasio, B.Di. *et al.*, *Tetrahedron*, 1976, **32**, 473-

478 (*Axisonitrile 3, Axisothiocyanate 3, Axamide 3*)

Fusetani, N. *et al.*, *Tet. Lett.*, 1992, **33**, 6823-

6826 (*Axisonitrile 3*)

Hirota, H. *et al.*, *Tetrahedron*, 1996, **52**, 2359-

2368 (*Axisothiocyanate 3, Axamide 3*)

Okino, T. *et al.*, *Tetrahedron*, 1996, **52**, 9447-

9454 (*10-Epiaxisonitrile 3*)

Koenig, G.M. *et al.*, *Planta Med.*, 2000, **66**,

337-342 (*Axisonitrile 3, activity*)

Uy, M.M. *et al.*, *Bioorg. Med. Chem. Lett.*,

2002, **12**, 3037-3038 (*Exiguamide, cryst struct*)

Uy, M.M. *et al.*, *Tetrahedron*, 2003, **59**, 731-

736 (*Exiguamide, Exicarbamate, Exigurin, cryst struct*)


Yan, X.-H. *et al.*, *J. Asian Nat. Prod. Res.*,

2006, **8**, 579-584 (*3-Oxoaxisonitrile 3*)

Spirobrassinin

S-426

2'-(Methylthio)spiro[3H-indole-3,5'-(4'H)-thiazol]-2(1H)-one, 9CI
[294857-91-3]



(R)-form

C₁₁H₁₀N₂O₂S₂ 250.345

(R)-form [294864-10-1]

N¹-Methoxy: **N-Methoxyspirobrassinin**

[156499-63-7]

C₁₂H₁₂N₂O₂S₂ 280.371

Phytoalexin from *Brassica oleracea* var. *gongyloides* (kohlrabi). [α]_D +61.1 (c, 1 in MeOH). Abs. config. shown to be opposite of parent Spirobrassinin in 2005. λ_{max} 220 (ε 23400); 257 (sh) (ε 6300); 290 (ε 1300) (MeOH).

(S)-form [113866-40-3]


Alkaloid from *Brassica oleracea* var. *botrytis* (cauliflower), *Raphanus sativus* var. *hortensis* (daikon) and other crucifers as response to fungal infection. Phytoalexin. Cryst. [α]_D -69.5 (c, 1.14 in CHCl₃) (natural). [α]_D -143.6 (opt. pure). The nat. prod. is a partial racemate. Undergoes strong enantiomeric enrichment in achiral chromatographic phase. λ_{max} 215 (ε 23500); 250 (ε 8200); 300 (ε 1900) (MeOH).

6-Methoxy: **6-Methoxyspirobrassinin**.**Capparine A**C₁₂H₁₂N₂O₂S₂ 280.371

Alkaloid from *Capparis himalayensis*. Pale yellow plates (CHCl₃/MeOH). Mp 170-173°. [α]_D -9.9 (c, 0.16 in MeOH). Partial racemate. λ_{max} 198 (log ε 2.86); 224 (log ε 3.34) (MeOH).

Takasugi, M. *et al.*, *Chem. Lett.*, 1987, 1631-1632 (*isol. struct*)Monde, K. *et al.*, *J.A.C.S.*, 1994, **116**, 6650-6657 (*synth*)Gross, D. *et al.*, *Z. Naturforsch., C*, 1994, **49**, 281-285 (*1-Methoxyspirobrassinin*)Monde, K. *et al.*, *Chem. Lett.*, 2000, 886-887 (*abs config*)Monde, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1312-1314 (*opt purity*)Pedras, M.S.C. *et al.*, *Phytochemistry*, 2000, **53**, 161-176 (*rev*)Suchy, M. *et al.*, *J.O.C.*, 2001, **66**, 3940-3947 (*synth. cd, abs config*)Monde, K. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 5206-5212 (*1-Methoxyspirobrassinin, abs config*)Pedras, M.S.C. *et al.*, *Chem. Comm.*, 2006, 1848-1850 (*biosynth*)Pedras, M.S.C. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 691-701 (*synth*)Curilova, Z. *et al.*, *Tet. Lett.*, 2007, **48**, 8200-8204 (*1-Methoxyspirobrassinin, synth*)Li, Y.-Q. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 189-191 (*Capparine A*)**Spirobrassinol**

S-427



Absolute Configuration

C₁₁H₁₂N₂O₂S₂ 252.361N-Methoxy: **N-Methoxyspirobrassinol**

[166020-09-3, 165746-57-6]

C₁₂H₁₄N₂O₂S₂ 282.387

Stress metab. from the Japanese radish daikon (*Raphanus sativus* var. *hortensis*) inoculated with *Pseudomonas cichorii*. Gum. [α]_D²⁰ 0 (c, 0.52 in CHCl₃). Unusual hemiaminal struct. Occurs as a mixt. of diastereoisomers, ratio α-OH:β-OH estimated as 2.5:1. λ_{max} 211 (ε 27300); 246 (ε 15500); 292 (ε 3630) (EtOH).

Me ether, N-methoxy: **N-Methoxyspirobrassinol methyl ether**

[113866-41-4]


C₁₃H₁₆N₂O₂S₂ 296.414

Stress metab. from *Raphanus sativus* var. *hortensis* inoculated with *Pseudomonas cichorii*. Gum. [α]_D¹⁸ -1.9 (c, 1.57 in CHCl₃). 2R,3R-config. determined in 2005. λ_{max} 210 (ε 26400); 291 (ε 2880) (EtOH).

Monde, K. *et al.*, *Phytochemistry*, 1995, **39**, 581-586 (*isol, pmr, cmr, ms*)Monde, K. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 5206-5212 (*Me ether, abs config*)Curilova, Z. *et al.*, *Tet. Lett.*, 2007, **48**, 8200-8204 (*synth*)**Spirocalcaridine A**

S-428

[568561-18-2]



Relative Configuration

C₁₉H₂₁N₃O₄ 355.393

Isol. from the calcareous sponge *Leucetta* sp. Yellow oil. [α]_D -59.3 (c, 0.11 in MeOH).

4-Me ether: **Spirocalcaridine B**

[568561-22-8]


C₂₀H₂₃N₃O₄ 369.419

Isol. from *Leucetta* sp. Yellow oil. [α]_D -50.8 (c, 0.13 in MeOH).

Edrada, R.A. *et al.*, *J. Nat. Prod.*, 2003, **66**, 939-942 (*isol, pmr, cmr*)**Spiroformabuxine**

S-429

[190720-50-4]



Absolute Configuration


C₂₅H₃₅NO 365.558

Highly modified steroidal alkaloid. Al-

kaloid from the leaves of *Buxus sempervirens* (Buxaceae).Fourneau, C. *et al.*, *Tet. Lett.*, 1997, **38**, 2965-2968 (*isol, pmr, cmr, abs config*)**Spiroleucettadine**

S-430

[813460-81-0]



Absolute Configuration


C₂₀H₂₃N₃O₄ 369.419

Structure revised in 2008. Alkaloid from a *Leucetta* sp. Yellow powder. [α]_D -27.1 (c, 0.38 in MeOH). λ_{max} 227 (ε 2410) (MeOH).

Ralifo, P. *et al.*, *J.O.C.*, 2004, **69**, 9025-9029 (*isol, pmr, cmr*)White, K.N. *et al.*, *J.O.C.*, 2008, **73**, 8719-8722 (*pmr, cmr, cryst struct*)**Spirolide B**

S-431

[170713-72-1]



Absolute Configuration

R = H

C₄₂H₆₃NO₇ 693.962

Isol. from *Alexandrium ostenfeldii* found in various contaminated shellfish such as *Mytilus edulis*, *Placopecten magellanicus*. Calcium channel activating agent, phycotoxin.

2,3-Didehydro: **Spirolide A**

[318996-47-3]

C₄₂H₆₁NO₇ 691.946

Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.

Hu, T. *et al.*, *Chem. Comm.*, 1995, 2159-2161 (*isol, pmr, cmr, ms*)Hu, T. *et al.*, *J. Nat. Prod.*, 2001, **64**, 308-312 (*Spirolide A*)Falk, M. *et al.*, *Tetrahedron*, 2001, **57**, 8659-8665 (*config*)Sleno, L. *et al.*, *Anal. Bioanal. Chem.*, 2004, **378**, 969-976; 977-986 (*isol, ms*)Meilert, K. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 2164-2192 (*synth, abs config*)**Spirolide D**

S-432

[170713-73-2]

As Spirolide B, S-431 with

R = CH₃C₄₃H₆₅NO₇ 707.989

Isol. from *Alexandrium ostenfeldii* found in contaminated shellfish. Ca channel activator; phycotoxin.

2,3-Didehydro: **Spirolide C**

[318996-49-5]
 $C_{43}H_{63}NO_7$ 705.973
 Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.

13-Demethyl-13-Demethylspirolide D
 $C_{42}H_{63}NO_7$ 693.962
 Isol. from *Alexandrium ostenfeldii*.

13-Demethyl, 2,3-didehydro-13-De-methylspirolide C. 13-Desmethylspirolide C
 [334974-07-1]
 $C_{42}H_{61}NO_7$ 691.946
 Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.

13,19-Didemethyl, 2,3-didehydro-13,19-Didemethylspirolide C
 $C_{41}H_{59}NO_7$ 677.92
 Isol. from *Alexandrium ostenfeldii*.

13,19-Didemethyl, 27 ξ -hydroxy, 2,3-didehydro-27-Hydroxy-13,19-didemethylspirolide C. 13,19-Didemethyl-27-hydroxyspirolide C
 [1000865-10-0]
 $C_{41}H_{59}NO_8$ 693.919
 Isol. from *Alexandrium ostenfeldii*.

Hu, T. et al., *Chem. Comm.*, 1995, 2159-2161 (isol, pmr, cmr, ms)

Hu, T. et al., *J. Nat. Prod.*, 2001, **64**, 308-312 (Spirolide C, 13-Demethylspirolide C)

Falk, M. et al., *Tetrahedron*, 2001, **57**, 8659-8665 (config)

Sleno, L. et al., *Anal. Bioanal. Chem.*, 2004, **378**, 969-976; 977-986 (isol, ms)

MacKinnon, S.L. et al., *J. Nat. Prod.*, 2006, **69**, 983-987 (13,19-Didemethylspirolide C)

MacKinnon, S.L. et al., *J.O.C.*, 2006, **71**, 8724-8731 (13-Demethylspirolide C, biosynth)


Meilert, K. et al., *Org. Biomol. Chem.*, 2006, **4**, 2184-2192 (synth, abs config)

Ciminiello, P. et al., *J. Nat. Prod.*, 2007, **70**, 1878-1883 (27-Hydroxy-13,19-didemethylspirolide C)

Spirolide E

S-433

[183159-71-9]



$C_{42}H_{63}NO_8$ 709.962
 Isol. from *Alexandrium ostenfeldii* found in contaminated shellfish.

2,3-Dihydro- Spirolide F
 [183159-73-1]

$C_{42}H_{65}NO_8$ 711.977
 Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.


Hu, T. et al., *Tet. Lett.*, 1996, **37**, 7671-7674 (isol, pmr, cmr, ms)

Sleno, L. et al., *Anal. Bioanal. Chem.*, 2004, **378**, 969-976; 977-986 (isol, ms)

Spirolide G

S-434

[849215-96-9]



$C_{42}H_{61}NO_7$ 691.946
 Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.

20-Methyl-20-Methylspirolide G
 [849215-95-8]

$C_{43}H_{63}NO_7$ 705.973
 Isol. from *Alexandrium ostenfeldii* in contaminated shellfish.

20-Methyl, 17-O-hexadecanoyl-20-Methyl-17-O-palmitoylspirolide G
 [901124-53-6]

$C_{59}H_{93}NO_8$ 944.386
 Isol. from *Alexandrium ostenfeldii* in contaminated mussels. Most abundant member of a large group of fatty acid esters.

Aasen, J. et al., *Chem. Res. Toxicol.*, 2005, **18**, 509-515 (isol, pmr, cmr, ms)


MacKinnon, S.L. et al., *J. Nat. Prod.*, 2006, **69**, 983-987 (isol, pmr, cmr, ms)

Aasen, J. et al., *Rapid Commun. Mass Spectrom.*, 2006, **20**, 1531-1537 (20-Methylspirolide G esters)

Spirolucidine

S-435

[89647-79-0]




$C_{30}H_{49}N_3O_2$ 483.736
 Alkaloid from *Lycopodium lucidulum* (Lycopodiaceae). Viscous oil.

Ayer, W.A. et al., *Can. J. Chem.*, 1984, **62**, 298 (isol, ir, pmr, ms, struct)

Spiropachysine

S-436

[19587-41-8]



$C_{31}H_{46}N_2O$ 462.717

Absolute configuration

Alkaloid from leaves of *Pachysandra terminalis* (Buxaceae). Needles. Mp 290-292°. $[\alpha]_D^{25} +35$ (CHCl₃).

▶WH1314450

N²-De-Me:

Leaves (Me₂CO). Mp 283-285°. $[\alpha]_D^{18} +54$ (c, 1.0 in CHCl₃).

N²⁰-De-Me:

Prisms (Me₂CO). Mp 287-290°.

3-Epimer: Isospiropachysine


$C_{31}H_{46}N_2O$ 462.717
 Alkaloid from *Pachysandra axillaris* (Buxaceae). Needles. Mp 297-298°. $[\alpha]_D^{22} +39.7$ (CHCl₃).

Kikuchi, T. et al., *Tet. Lett.*, 1968, 2077 (isol)
 Kikuchi, T. et al., *Chem. Pharm. Bull.*, 1975, **23**, 416 (struct, ms, uv, ir, pmr, cd, ord)

Minghua, C. et al., *Phytochemistry*, 1990, **29**, 3927 (Isospiropachysine)

Spiropyrrolizidine 252

S-437



$C_{14}H_{24}N_2O_2$ 252.356

Struct. revised in 1992. Alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates pumilio* (Dendrobatidae). $[\alpha]_D +18.4$ (c, 0.47 in MeOH). $[\alpha]_D -4.3$ (c, 0.47 in CHCl₃).

Deoxy- Spiropyrrolizidine 236. Dendrobates Alkaloid 236

$C_{14}H_{24}N_2O$ 236.356
 Alkaloid from skin extracts of *Dendrobates pumilio* and from the millipede *Rhinotus purpureus*. $[\alpha]_D +55.6$ (c, 1.0 in MeOH).

Deoxy, O-de-Me: Spiropyrrolizidine 222. Dendrobates Alkaloid 222

$C_{13}H_{22}N_2O$ 222.33
 Struct. revised in 1992. Alkaloid from skin extracts of the Panamanian poison-frog *Dendrobates pumilio* (Dendrobatidae).

Tokuyama, T. et al., *Tetrahedron*, 1987, **43**, 643; 1992, **48**, 4247 (isol, ir, pmr, cmr, ms, struct)


Hutchinson, K.D. et al., *Tetrahedron*, 1994, **50**, 6129 (synth)

Saporito, R.A. et al., *J. Chem. Ecol.*, 2003, **29**, 2781-2786 (Spiropyrrolizidine 236, isol)

Spiroquinazoline

S-438

[157075-58-6]



Absolute Configuration


$C_{23}H_{19}N_5O_3$ 413.435

Isol. from *Aspergillus flavipes*. Substance P inhibitor. Yellow solid. Mp 166-168°. $[\alpha]_D^{25}$ -2.5 (c, 0.7 in MeOH). λ_{\max} 212 (ε 22500); 225 (ε 20600); 265 (ε 7010); 277 (ε 6350); 303 (ε 2440); 315 (ε 1980) (MeOH) (Derep).

Barrow, C.J. *et al.*, *J. Nat. Prod.*, 1994, **57**, 471-476 (isol, uv, ir, pmr, cmr, cd, struct)

Spirosolan-3-amine, 9CI S-439

3-Aminospirosolane
[125520-81-2]



C₂₇H₄₆N₂O 414.673

(3α,5α,22S,25S)-form [182689-75-4]
Alkaloid from *Solanum arboreum*.
Amorph. solid.

(3β,5α,22R,25R)-form [76932-66-6]
Alkaloid from *Solanum triste*. Cryst. (MeOH aq.). Mp 178-182°. $[\alpha]_D^{26}$ -57.3 (c, 0.26 in CHCl₃).

(3β,5α,22S,25S)-form

Soladunalinidine

[66934-59-6]

Alkaloid from *Solanum arboreum* and *Solanum dumaliatum*. Amorph. solid. Mp 145-153°. $[\alpha]_D^{22}$ +13.3 (c, 1.5 in MeOH) (natural). $[\alpha]_D$ +3.7 (c, 0.25 in CHCl₃) (synthetic).

Picrate: Mp 235-255° dec.


Bird, G.J. *et al.*, *Aust. J. Chem.*, 1979, **32**, 611-617; 797-816 (isol, ir, pmr, ms, struct, cmr)

Quyen, L. T. *et al.*, *Annalen*, 1990, 519-524 (synth)

Maxwell, A. *et al.*, *Phytochemistry*, 1995, **58**, 625-628 (isol)

Maxwell, A. *et al.*, *Phytochemistry*, 1996, **43**, 913 (isol, pmr, cmr, ms)

Spirosolane-2,3-diol S-440




C₂₇H₄₅NO₃ 431.657

(2α,3β,5α,22R,25R)-form
2-Hydroxysoladulcinine

Alkaloid from roots of *Lycianthes biflora* (Solanaceae). Needles (MeOH aq.). Mp 252-255° dec. $[\alpha]_D^{22}$ -62.5 (c, 0.45 in CHCl₃).

Ripperger, H. *et al.*, *Phytochemistry*, 1992, **31**, 725-726 (isol)

Spirosolane-3,15-diol S-441



C₂₇H₄₅NO₃ 431.657

(3β,5α,15α,22R,25R)-form

15α-Hydroxysoladulcinine

[16137-74-9]

Isol. from *Solanum dulcamara* (Solanaceae). Prisms + 1H₂O (MeOH aq.). Mp 167-168° Mp 209-212° (double Mp, rapid htg.). On slow htg. the prisms turn to needles at 150-170°. Anhyd. material (by subl.) has Mp 213-215°.

15-Ketone: 15-Oxosoladulcinine

C₂₇H₄₃NO₃ 429.642

May occur in *Solanum dulcamara* (Solanaceae).

(3β,5α,15α,22S,25S)-form

15α-Hydroxytomatidine

[4828-39-1]

Alkaloid from *Solanum dulcamara* (Solanaceae). Needles (MeOH). Mp 150-155°. $[\alpha]_D^{18}$ +17.8 (c, 0.8 in CHCl₃).

Tri-Ac: Mp 203.5-204.5°. $[\alpha]_D^{18}$ -3.1 (c, 0.8 in CHCl₃).

(3β,5α,15β,22R,25R)-form

15β-Hydroxysoladulcinine

[16137-76-1]

Isol. from *Solanum dulcamara* (Solanaceae).


Ronsch, H. *et al.*, *Tet. Lett.*, 1965, **6**, 1947-1952 (*15α-Hydroxysoladulcinine*, *15α-Hydroxytomatidine*)

Ronsch, H. *et al.*, *Annalen*, 1966, **694**, 169-182 (*15α-Hydroxysoladulcinine*, *15α-Hydroxytomatidine*)

Ronsch, H. *et al.*, *J. Chromatogr.*, 1967, **30**, 149-154 (*15β-Hydroxysoladulcinine*)

Willuhn, G. *et al.*, *Planta Med.*, 1970, **18**, 354-360 (*15α-Hydroxytomatidine*)

Spirosolane-3,23-diol S-442



C₂₇H₄₅NO₃ 431.657

The 23-OH group changes the R,S-priorities compared with other spirosolanes.

(3β,5α,22R,23R,25R)-form

23R-Hydroxysoladulcinine

Alkaloid from roots of *Solanum pandur-aeforme* (Solanaceae). Mp 192-193°. $[\alpha]_D^{22}$ -65.4 (c, 0.44 in MeOH).

(3β,5α,22R,23S,25R)-form

23-Ac: 23-Acetoxy-soladulcinine

[152128-85-3]

C₂₉H₄₇NO₄ 473.695

Alkaloid from the roots of a *Lycopersicon esculentum-Lycopersicon hirsutum* hybrid. Amorph. $[\alpha]_D^{20}$ -46 (c, 0.37 in CHCl₃).

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: Lycoperside B

[176300-86-0]

C₅₂H₈₅NO₂₃ 1092.236

Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. $[\alpha]_D^{27}$ -40.3 (c, 0.56 in MeOH).

(3β,5α,22R,23S,25S)-form

23-Ac: 23-Acetoxy-25-episoladulcinine

[152128-86-4]

C₂₉H₄₇NO₄ 473.695

Alkaloid from roots of a *Lycopersicon esculentum-Lycopersicon hirsutum* hybrid. Amorph. $[\alpha]_D^{20}$ -36.7 (c, 1.2 in CHCl₃).

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: Lycoperside C

[176300-87-1]

C₅₂H₈₅NO₂₃ 1092.236

Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. $[\alpha]_D^{27}$ -26.3 (c, 0.63 in MeOH).

(3β,5α,22S,23R,25S)-form

23-Ac: 23R-Acetoxytomatidine

[152128-84-2]

C₂₉H₄₇NO₄ 473.695

Alkaloid from roots of a *Lycopersicon esculentum/Lycopersicon hirsutum* hybrid. Amorph. $[\alpha]_D^{20}$ -2.8 (c, 0.42 in CHCl₃).

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: 23R-Acetoxytomatidine.

Lycoperside A

[176181-33-2]

C₅₂H₈₅NO₂₃ 1092.236


Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. $[\alpha]_D^{27}$ -31.6 (c, 0.8 in MeOH).

Ripperger, H. *et al.*, *Phytochemistry*, 1993, **32**, 1607-1609 (*23-Hydroxysoladulcinine*)

Nagaoka, T. *et al.*, *Phytochemistry*, 1993, **34**, 1153-1157 (*23-Acetoxy-soladulcinine*, *23-Acetoxy-25-episoladulcinine*, *23-Acetoxytomatidine*)

Yahara, S. *et al.*, *Phytochemistry*, 1996, **42**, 169-172 (*Lycopersides*)

Spirosolane-3,23,24-triol S-443



C₂₇H₄₅NO₄ 447.657

(3β,5α,22S,23S,24R,25S)-form

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside], 24-O-β-D-glucopyranoside:


Lycoperside G

[675828-27-0]

C₅₈H₉₅NO₂₉ 1270.378

Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. [α]_D²⁰ -44.1 (c, 0.68 in MeOH).

Yahara, S. et al., *J. Nat. Prod.*, 2004, **67**, 500-502 (isol, pmr, cmr)

Spirosolan-3,23,27-triol**S-444**C₂₇H₄₅NO₄ 447.657**(3β,5α,22R,23R,25S)-form****Isoesculeogenin A**

[854381-39-8]

Needles. Mp 206-213°. [α]_D -87.2 (c, 0.64 in Py).

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside], 27-O-β-D-glucopyranoside:

Lycoperside F

[675828-26-9]

C₅₈H₉₅NO₂₉ 1270.378

Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. [α]_D²⁹ -38.5 (c, 0.8 in MeOH).

(3β,5α,22S,23S,25S)-form**Esculeogenin A**

[854381-37-6]

Needles. Mp 215-220°. [α]_D²⁶ -99.6 (c, 0.7 in Py).

23-Ac, 3-O-[[β-D-glucopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside], 27-O-β-D-glucopyranoside:

Esculeoside A

[532387-86-3]


C₅₈H₉₅NO₂₉ 1270.378

Alkaloid from ripe tomatoes (*Lycopersicon esculentum* var. *cerasiforme*). Needles (MeOH). Mp 225° dec. [α]_D²⁶ -52.5 (c, 0.6 in MeOH).

Yahara, S. et al., *J. Nat. Prod.*, 2004, **67**, 500-502 (*Lycoperside F*)

Fujiwara, Y. et al., *Tetrahedron*, 2004, **60**, 4915-4920 (*Esculeoside A*)

Yoshizaki, M. et al., *Chem. Pharm. Bull.*, 2005, **53**, 839-840 (*Isoesculeogenin A*)

Spirosolan-3-ol, 9CI**S-445**C₂₇H₄₅NO₂ 415.658

Log P 6.08 (uncertain value) (calc).

(3β,5α,22R,25R)-form

Soladulcidine. *Megacarpidine*. *Dihydrosolasodine*. *Solasodanol*. 5α-Solasodan-3β-ol (obsol.)

[511-98-8]

Aglycone of several very widespread glycosides found in *Solanum* spp. Said to occur free in *Lycopersicon pimpinellifolium* (currant tomato). Shows antifungal activity. Mp 209-211°. [α]_D -53 (c, 0.64 in Py). [α]_D -50 (c, 0.4 in CHCl₃).

Hydrochloride: Mp 298-299°.

Picrate: Mp 152-154°.

3-O-[[β-D-Xylopyranosyl-(1→4)]-β-D-galactopyranoside]: **Solalyratine A**

C₃₈H₆₃NO₁₁ 709.916

Alkaloid from *Solanum lyratum*. Needles + 3H₂O. Mp 252-254°. [α]_D -89.2 (c, 0.5 in Py).

3-O-[[β-D-Xylopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: **Solalyratine B**

C₄₄H₇₃NO₁₆ 872.058

Alkaloid from *Solanum lyratum*. Needles + 2H₂O. Mp 274-279°. [α]_D -17.9 (c, 0.5 in Py).

3-O-[[α-L-Rhamnopyranosyl-(1→2)]-[[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Soladulcine A**. *Soladulcidine 3-chacotrioxide*

[156555-52-1]

C₄₅H₇₅NO₁₅ 870.085

Alkaloid from *Solanum dulcamara*. Needles + 1H₂O. Mp 256-258°. [α]_D²⁴ -91.8 (c, 0.33 in Py).

3-O-[[β-D-Glucopyranosyl-(1→2)]-[[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: **Soladulcine B**. *Soladulcidine 3-lycote-triaxide*

[90366-11-3]

C₅₀H₈₃NO₂₁ 1034.2

Alkaloid from *Solanum dulcamara* and *Solanum japonense*. Needles + 2H₂O. Mp 264-266°. [α]_D²⁴ -58.4 (c, 0.5 in Py). Possible identity with α-Soladulcine.

3-O-[[β-D-Glucopyranosyl-(1→2)]-β-D-glucopyranosyl-(1→4)]-[[α-L-rhamnopyranosyl-(1→2)]-β-D-galactopyranoside]: **Dihydrosolasuaveoline**

C₅₁H₈₅NO₂₁ 1048.227

Alkaloid from *Solanum suaveolens* (Solanaceae). Cryst. (MeOH aq.). Mp 275-277° dec. [α]_D²⁴ -43.8 (c, 0.5 in Py).

Glycoside(1): **Megacarpine**

C₄₄H₇₃NO₁₆ 872.058

Alkaloid from *Solanum megacarpum* (Solanaceae). Mp 259-260°. [α]_D -61.6 (Py). Hydrol. gives 1 Glu, 1 Gal and 2 Xyl.

Glycoside(2): **Soladulcidine tetraoxide**

C₄₅H₇₅NO₁₇ 902.084

Alkaloid from *Solanum dulcamara* (Solanaceae). Shows antifungal activity. Mp 268-270°. [α]_D -53 (Py). Hydrol. gives 2 Glu, 1 Gal and 1-Xyl.

Glycoside(3): **α-Soladulcine**

[37337-73-8]

C₅₀H₈₃NO₂₁ 1034.2

Alkaloid from *Solanum dulcamara* and *Solanum pseudopersicum* (Solanaceae). Mp 265-269°. [α]_D -56 (MeOH). Hydrol. gives 2 Glu, 1 Gal and 1 D-Xyl.

Glycoside(4): **β-Soladulcine**

[11093-43-9]

C₄₅H₇₅NO₁₆ 886.085

Alkaloid from *Solanum dulcamara* (Solanaceae). Mp 246-248° dec. [α]_D -39 (c, 0.38 in MeOH). Hydrol. gives 1 Glu, 1 Gal and 1 Rha.

Glycoside(5): **γ-Soladulcine**

Alkaloid from *Solanum dulcamara*

(Solanaceae). Not well characterised. Isol. only (Schreiber) as a mixt. with α- and β-Soladulcines.

(3β,5α,22S,25S)-form

Tomatidine. 5α-Tomatidan-3β-ol (obsol.)

[77-59-8]

Alkaloid from *Lycopersicon esculentum* (tomato) and *Solanum demissum*, found as glycosides in many other *Lycopersicon* and *Solanum* spp. (Solanaceae). Shows antifungal and cytostatic activity, has been used clinically against dermatitis. Cholinesterase inhibitor, antihistamine. Repellent to the Colorado beetle *Leptinotarsa decemlineata*. Mp 210-211°. [α]_D²⁰ +5 (MeOH). Log P 6.08 (uncertain value) (calc).

Hydrochloride: Mp 275-280° dec. [α]_D²⁵ -9 (MeOH).

Picrate: Mp 131-133°.

O-β-D-Glucopyranoside: [177354-90-4]

C₃₃H₅₅NO₇ 577.8

Alkaloid from aerial parts of *Solanum arboreum*. Cryst. Mp 174-177° (as penta-Ac). [α]_D²³ -13.6 (c, 0.44 in CHCl₃).

O-β-D-Galactopyranoside: **δ-Tomatine**

[99304-79-7]

C₃₃H₅₅NO₇ 577.8

Metab. of α-Tomatine.

O-[[β-D-Xylopyranosyl-(1→6)]-β-D-glucopyranoside]: [177354-91-5]

C₃₈H₆₃NO₁₁ 709.916

Alkaloid from aerial parts of *Solanum arboreum*. Needles. Mp 238-240° (as hepta-Ac). [α]_D²³ -7.3 (c, 1.64 in CHCl₃).

3-O-[[β-D-Glucopyranosyl-(1→4)]-β-D-galactopyranoside]: **γ-Tomatine**. *Lycoperside D*

[81037-16-3]

C₃₉H₆₅NO₁₂ 739.942

Alkaloid from *Lycopersicon esculentum* (tomato). Amorph. powder. [α]_D²⁷ -10.8 (c, 0.6 in MeOH).

O-[[β-D-Xylopyranosyl-(1→2)]-[[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranoside]: **Polyanine**

C₄₃H₇₁NO₁₅ 842.032

Alkaloid from *Solanum polyadenium* (Solanaceae). Needles (MeOH aq.). Mp 263-270° dec. [α]_D²¹ -36.3 (c, 1.05 in Py).

3-O-[[β-D-Glucopyranosyl-(1→2)]-[[β-D-glucopyranosyl-(1→3)]-β-D-glucopyranosyl-(1→4)]-β-D-galactopyranoside]: **Sisunine**. *Neotomatine*

[85547-35-9]

C₅₁H₈₅NO₂₂ 1064.226

Alkaloid from clones of hybrids of

Solanum acaule and *Solanum ajanhuiri*.

3-O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-galactopyranoside]: **β -Tomatine**

[17406-46-1]

C₄₅H₇₅NO₁₇ 902.084

Alkaloid from *Lycopersicon esculentum* and *Lycopersicon pimpinellifolium*, also obt. by partial hydrolysis of Tomatine (Solanaceae). Amorph. solid + 1H₂O. Mp 240-260° dec. $[\alpha]_D^{22}$ -28 (c, 1 in Py). Not obt. completely pure. Other Tomatines (β -, γ - and δ -) are other nonalkaloidal partial hydrolysis prods. of Tomatine.

O- $[\beta$ -D-Glucopyranosyl-(1 \rightarrow 2)]- $[\beta$ -D-xylopyranosyl-(1 \rightarrow 3)]- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-galactopyranoside]:

Tomatine. α -Tomatine. *Lycopersicin*

[17406-45-0]
C₅₀H₈₃NO₂₁ 1034.2

Alkaloid from *Lycopersicon esculentum*, other *Lycopersicon* spp., *Solanum polyadenium*, *Solanum tomatillo*, *Solanum boerhaavii*, *Solanum demissum* and other *Solanum* spp. Antiinflammatory agent. Antibacterial, antifungal agent. Acetylcholine esterase inhibitor, immunostimulant. Shows molluscicidal propls. Sol. MeOH, EtOH, dioxan; fairly sol. H₂O, Et₂O, acids, hexane; poorly sol. CHCl₃, EtOAc, C₆H₆. Mp 263-267° dec. $[\alpha]_D$ -19 (Py). Log P -3.27 (uncertain value) (calc).

► LD₅₀ (rat, orl) 900 mg/kg; LD₅₀ (mus, ipr) 25 mg/kg; LD₅₀ (mus, scu) 1000 mg/kg; LD₅₀ (mus, orl) 500 mg/kg. XW1050000

N-Nitroso: **N-Nitrosotomatidine**

[4847-05-6]
C₂₇H₄₄N₂O₃ 444.656

Alkaloid from leaves and stems of *Lycopersicon esculentum* (Solanaceae). Hexagonal cryst. changing to flat rods on heating. Mp 234-237°. Mp. quoted is for synthetic material.

3-Ketone: **5 α -Tomatidan-3-one**

[6870-84-4]
C₂₇H₄₃NO₂ 413.642

Alkaloid from roots of a *Lycopersicon esculentum*/*Lycopersicon hirsutum* hybrid. Needles (MeOH). Mp 195-198°. $[\alpha]_D^{20}$ +19 (c, 0.38 in CHCl₃).

Brink, N.G. et al., *J.A.C.S.*, 1951, **73**, 4018 (isol, Tomatidine)

Kuhn, R. et al., *Chem. Ber.*, 1957, **90**, 203 (Tomatines)

Sato, Y. et al., *J.A.C.S.*, 1957, **79**, 6089 (Tomatidine, struct)

Schreiber, K. et al., *Planta Med.*, 1958, **6**, 94 (Soladulcines)

Uhle, F.C. et al., *J.O.C.*, 1962, **27**, 656 (synth, Soladulcine)

Schreiber, K. et al., *Annalen*, 1964, **674**, 168 (Polyanine)

Budzikiewicz, H. et al., *Tetrahedron*, 1964, **20**, 2265 (Tomatidine, ms)

Boll, P.M. et al., *Acta Chem. Scand.*, 1965, **19**, 1365 (pmr, stereochem)

Schreiber, K. et al., *Annalen*, 1965, **682**, 219 (config)

Schreiber, K. et al., *Phytochemistry*, 1966, **5**, 707 (isol)

Adam, G. et al., *Tetrahedron*, 1966, **22**, 3591 (synth)

Kennard, O. et al., *J.C.S.(C)*, 1967, 956 (cryst struct, Tomatidine)

Wolters, B. et al., *Planta Med.*, 1970, **19**, 189-193 (Soladulcine, activity)

Tukalo, E.A. et al., *Khim. Prir. Soedin.*, 1971, **7**, 207; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 203 (Soladulcines)

Roddick, J.G. et al., *Phytochemistry*, 1974, **13**, 1459 (biosynth)

Weston, R.J. et al., *Aust. J. Chem.*, 1977, **30**, 917 (Tomatidine, cmr)

Garnero, J. et al., *CA*, 1982, **96**, 148962t (N-Nitrosotomatidine)

Osman, S.F. et al., *Phytochemistry*, 1986, **25**, 967 (Sisunine)

Willker, W. et al., *Magn. Reson. Chem.*, 1992, **30**, 645-650 (pmr, cmr)

Nagaoka, T. et al., *Phytochemistry*, 1993, **34**, 1153 (5 α -Tomatidan-3-one)

Lee, Y.Y. et al., *Chem. Pharm. Bull.*, 1994, **42**, 707-709 (Soladulcines)

Yahara, S. et al., *Phytochemistry*, 1996, **42**, 169-172 (Lycopersosides)

Maxwell, A. et al., *Phytochemistry*, 1996, **42**, 543 (derivs)


Friedman, M. et al., *J. Agric. Food Chem.*, 2002, **50**, 5751-5780 (Tomatines, rev)

Zha, X. et al., *Chem. Biodiversity*, 2007, **4**, 25-31 (Soladulcine, synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, THG250

Spirosol-5-en-3-amine, 9CI S-446

3-Aminospirosol-5-ene



C₂₇H₄₄N₂O 412.657

(3 α ,22R,25R)-form [173074-72-1]

Alkaloid from *Solanum triste*. Amorph. solid. Mp 166-170°.

(3 α ,22S,25S)-form [173074-71-0]

Alkaloid from *Solanum triste*. Amorph. solid. Mp 147-154°.

(3 β ,22R,25R)-form [164117-53-7]

Alkaloid from *Solanum triste*. Needles. Mp 165-167°. $[\alpha]_D^{27}$ -52 (c, 1 in MeOH).


(3 β ,22S,25S)-form [182689-76-5]

Alkaloid from *Solanum arboreum*. Amorph. solid.

Maxwell, A. et al., *J. Nat. Prod.*, 1995, **58**, 625-628; 1996, **59**, 200-201 (*Solanum triste* constits)

Maxwell, A. et al., *Phytochemistry*, 1996, **43**, 913-915 (*Solanum arboreum* constit)

Spirosol-5-ene-3,12-diol S-447



C₂₇H₄₃NO₃ 429.642

(3 β ,12 α ,22R,25R)-form

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside]: **Robeneoside A**. 12 α -Hydroxysolamargine

C₄₅H₇₃NO₁₆ 884.069

Alkaloid from the dried fruit of *Solanum lycocarpum*. Powder (CHCl₃/MeOH). Mp 205-208°. $[\alpha]_D^{29}$ -112.2 (c, 1 in Py).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 3)]- β -D-galactopyranoside]: **Robeneoside B**. 12 α -Hydroxysolasonine

C₄₅H₇₃NO₁₇ 900.068

Alkaloid from the dried fruit of *Solanum lycocarpum*. Powder (CHCl₃/MeOH). Mp 206-210°. $[\alpha]_D^{29}$ -22.2 (c, 1 in Py).

(3 β ,12 β ,22R,25R)-form

Solanaviol. Alkaloid SN-b

[74131-93-4]

Alkaloid from the leaves of *Solanum aviculare*. Mp 229-232.5°. $[\alpha]_D$ -113 (c, 0.97 in CHCl₃).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\alpha$ -L-rhamnopyranosyl-(1 \rightarrow 4)]- β -D-glucopyranoside]: **Lobofrutoside**. 12 β -Hydroxysolamargine

C₄₅H₇₃NO₁₆ 884.069

Alkaloid from the fruit of *Solanum lycocarpum*. Powder. $[\alpha]_D^{27}$ -92.3 (c, 1 in MeOH).

3-O- $[\alpha$ -L-Rhamnopyranosyl-(1 \rightarrow 2)]- $[\beta$ -D-glucopyranosyl-(1 \rightarrow 3)]- β -D-galactopyranoside]: Alkaloid SN-e. 12 β -Hydroxysolasonine

[107484-57-1]

C₄₅H₇₃NO₁₇ 900.068

Alkaloid from the fruits of *Solanum lycocarpum* and *Solanum nigrum*. Amorph. powder. $[\alpha]_D$ -53.3 (Py).


Kaneko, K. et al., *Phytochemistry*, 1980, **19**, 299-302 (*Solanaviol*)

Yoshida, X. et al., *Chem. Pharm. Bull.*, 1987, **35**, 1645-1648 (Alkaloid SN-e)

Yoshikawa, M. et al., *J. Nat. Prod.*, 2007, **70**, 210-214 (*Robeneosides A,B*)

Nakamura, S. et al., *Phytochemistry*, 2008, **69**, 1565-1572 (*Lobofrutoside*)

Spirosol-5-ene-3,15-diol S-448



C₂₇H₄₃NO₃ 429.642


(3 β ,15 α ,22R,25R)-form

15 α -Hydroxysolasonine




Alkaloid from *Solanum dulcamara* (Solanaceae). Prisms (MeOH aq.). Mp 212-216°. $[\alpha]_D^{21}$ -84.5 (c, 0.68 in CHCl₃).

(3β,15α,22S,25S)-form**15α-Hydroxytomatidenol**


[7755-31-9]

Isol. from *Solanum dulcamara* (Solanaceae).**(3β,15β,22R,25R)-form****15β-Hydroxysolasodine**Said to occur in *Solanum dulcamara* (Solanaceae). Details unpubl. (K. Schreiber).Rönsch, H. *et al.*, *Annalen*, 1966, **694**, 169-182 (*15α-Hydroxysolasodine*, *15α-Hydroxytomatidenol*)Willuhn, G. *et al.*, *Planta Med.*, 1970, **18**, 354-360 (*15α-Hydroxysolasodine*, *15α-Hydroxytomatidenol*)**Spirosol-5-ene-3,21-diol S-449**C₂₇H₄₃NO₃ 429.642**(3β,22R,25R)-form**Aglycone from *Solanum sycophanta*.3-O-[β-D-Xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-[α-L-rhamnopyranosyl-(1→2)]-β-D-glucopyranoside]: **21-Hydroxysycophantine** [188342-01-0]C₅₀H₈₁NO₂₀ 1016.185Alkaloid from *Solanum sycophanta*.


Powder (MeOH). Mp 256-258° dec.

[α]_D²¹ -78 (c, 0.9 in MeOH).Usubillaga, A. *et al.*, *Phytochemistry*, 1997, **44**, 537-543 (*isol*, *pmr*, *cmr*, *ms*)**Spirosol-5-ene-3,23-diol, 9CI S-450**C₂₇H₄₃NO₃ 429.642The 23-OH group changes the *R,S*-priority at C-22 compared with other spirosolanines.**(3β,22S,23S,25R)-form****Solaverol A**. 23S-Hydroxysolasodine [129273-42-3]Alkaloid from the roots of *Solanum canense* and *Solanum fraxinifolium*; hydrolyt. prod. of Solaverines I and II. Pale yellow powder or cryst. (Me₂CO aq.). [α]_D²⁰ -67.1 (CHCl₃). [α]_D²² -93.2 (c, 0.5 in CHCl₃).3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-galactopyranoside]: **23-Hydroxyisoanguivine** [188483-45-6]C₄₄H₇₁NO₁₆ 870.042Alkaloid from *Solanum uporo*. Needles (MeOH aq.). Mp 225-230° dec. [α]_D²⁶ -90.1 (c, 0.56 in Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranoside]: **23-Hydroxyanguivine** [188483-43-4]C₄₄H₇₁NO₁₆ 870.042Alkaloid from *Solanum uporo*. Mp 215-222° dec. [α]_D²² -94 (c, 0.71 in Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-glucopyranosyl-(1→3)]-β-D-galactopyranoside]: **Solaverine II** [128397-86-4]C₄₅H₇₃NO₁₇ 900.068Alkaloid from the aerial parts of *Solanum toxicarium* and *Solanum verbascifolium* (Solanaceae). Pale yellow powder. [α]_D -74.6 (Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Solaverine I** [128397-85-3]C₄₅H₇₃NO₁₆ 884.069Alkaloid from the aerial parts of *Solanum toxicarium* and *Solanum verbascifolium* (Solanaceae). Pale yellow powder. [α]_D -75.7 (Py).Yamashita, T. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 827-829 (*Solaverines*, *Solaverol A*)Ripperger, H. *et al.*, *Phytochemistry*, 1991, **30**, 1299-1301; 1997, **44**, 731-734 (*23-Hydroxyanguivine*, *23-Hydroxyisoanguivine*, *23-Hydroxysolasodine*)**Spirosol-5-ene-3,25-diol S-451**C₂₇H₄₃NO₃ 429.642**(3β,25S)-form**25-Ac, 3-O-[α-L-arabinopyranosyl-(1→3)-[α-L-rhamnopyranosyl-(1→2)]-β-D-glucopyranoside]: **25-Acetoxyrobustine** [156122-57-5]C₅₂H₈₃NO₂₁ 1058.222Alkaloid from the roots of *Solanum robustum*. Amorph. [α]_D²⁰ -71.4 (c, 0.85 in Py).Ripperger, H. *et al.*, *Annalen*, 1994, 517-520 (*isol*, *pmr*, *cmr*, *ms*)**Spirosol-5-ene-3,27-diol S-452**C₂₇H₄₃NO₃ 429.642**(3β,22R,25R)-form****Solaparnaine**

[103239-11-8]

Alkaloid from green berries of *Solanum asperum* (Solanaceae). [α]_D²³ -77.8 (MeOH).Bhattacharyya, J. *et al.*, *Heterocycles*, 1985, **23**, 3111-3112 (*isol*, *cmr*, *ms*)**Spirosol-5-ene-3,12,23-triol S-453**C₂₇H₄₃NO₄ 445.641**(3β,12β,22S,23ξ,25R)-form**23-Ac: **23-Acetoxy-12-hydroxysolasodine**. 23-O-Acetyl-12-hydroxysolasodine (*incorr.*)

[117803-97-1]

C₂₉H₄₅NO₅ 487.678Alkaloid from *Solanum nigrum* (Solanaceae). Mp 145°.Döpke, W. *et al.*, *Z. Chem.*, 1988, **28**, 185**Spirosol-5-ene-3,12,27-triol, 9CI S-454**C₂₇H₄₃NO₄ 445.641**(3β,12β,22R,25R)-form****Alkaloid SN-c**. 12,17-Dihydroxysolasodine


[107484-55-9]

Alkaloid from the berries of *Solanum nigrum* (Solanaceae). Mp 233-237°. [α]_D -54.7 (MeOH).3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Alkaloid SN-f**C₄₅H₇₃NO₁₇ 900.068Alkaloid from *Solanum nigrum* berries (Solanaceae). Amorph. powder. [α]_D -64.1 (Py).27-Carboxylic acid: **3,12-Dihydroxyspirosol-5-en-27-oic acid, 9CI. Alkaloid SN-d**

[107484-56-0]

C₂₇H₄₁NO₅ 459.625Alkaloid from the berries of *Solanum nigrum* (Solanaceae). Mp 276-278°. [α]_D -71.2 (Py).Yoshida, K. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 1645-1648 (*pmr*, *cmr*, *ms*, *struct*)

Spirosol-5-ene-3,23,27-triol, 9CI S-455

C₂₇H₄₃NO₄ 445.641

(3β,22S,23S,25R)-form

Solaverol B


[129273-43-4]

Hydrol. prod. of Solaverine III.

3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Solaverine III** [128397-87-5]C₄₅H₇₃NO₁₇ 900.068Alkaloid from the aerial parts of *Solanum toxicarium* and *Solanum verbascifolium* (Solanaceae). Pale yellow powder. [α]_D²⁰ -66 (Py).Yamashita, T. et al., *Chem. Pharm. Bull.*, 1990, **38**, 827-829 (isol, struct)

Spirosol-5-en-3-ol

S-456

C₂₇H₄₃NO₂ 413.642

▶ WF1300000

(3β,22R,25R)-form

Solasodine. Solanocarpidine. Purapuridine. Solanidine S. Solanacarpine [126-17-0]Isol. from *Solanum aculeatissimum*, *Solanum canense*, *Solanum cyananthum*, *Solanum fraxinifolium*, *Cestrum purpureum* and other plants, v. widely distributed in the Solanaceae as the alkaline constit. of glycoside alkaloids. Antineoplastic agent. Also reputed to have antiandrogenic, antiinflammatory and antispermatogenic props. Much investigated as potential starting material for coml. steroid synth, as alternative to Diosgenin. Shows strong antifungal and antiyeast activity. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 200-201°. [α]_D²⁰ -98 (MeOH). Log P 5.59 (uncertain value) (calc). Dimorph., both forms with same Mp.▶ LD₅₀ (rat, orl) 4978 mg/kg. Exp. reprod. and teratogenic effects.

Hydrochloride: Mp 313-314°.

Picrate: Mp 144-145°.

3-O-β-D-Glucopyranoside: **Solasodine 3-O-β-D-glucopyranoside**

[14197-65-0]

C₃₃H₅₃NO₇ 575.784Alkaloid from whole plants of *Solanum umbelliferum*. Amorph. powder. Mp 246-248°. [α]_D²⁰ -81.1 (c, 0.24 in

MeOH).

3-O-[α-L-Rhamnopyranosyl-(1→2)-β-D-glucopyranoside]: **β₁-Solamargine** [73069-20-2]C₃₉H₆₃NO₁₁ 721.927Alkaloid from *Solanum marginatum* (Solanaceae). Antifungal and molluscicidal agent. Mp 261°. [α]_D²⁰ -89. Log P 2.31 (uncertain value) (calc). β-Solamargine, which has been isol. from several *S. spp.*, was prob. a mixt. of β₁ and β₂-Solamargines. The identity of pure β₁-Solamargine is not conclusively established.3-O-[α-L-Rhamnopyranosyl-(1→4)-β-D-glucopyranoside]: **Khasianine**. β₂-Solamargine [32449-98-2]C₃₉H₆₃NO₁₁ 721.927Alkaloid from *Solanum khasianum* and *Solanum incanum* (Solanaceae). Cytotoxic agent. Microneedles (MeOH). Mp 226-228°. [α]_D²⁰ -95 (MeOH). [α]_D²⁴ -84 (Py). Also a partial hydrol. prod. of Solamargine.3-O-[β-D-Glucopyranosyl-(1→1)-β-L-rhamnopyranos-4-yl]: **Solaplumbine** [54302-48-6]C₃₉H₆₃NO₁₁ 721.927Alkaloid from *Nicotiana plumbaginifolia* (Solanaceae). Shows antineoplastic activity. Pale yellow needles (EtOH/C₆H₆). Sol. EtOH, butanol; poorly sol. CHCl₃, C₆H₆. Mp 180-181°. [α]_D²⁰ -90 (c, 1 in MeOH). Log P 2.31 (uncertain value) (calc).3-O-[D-Glucopyranosyl-(1→?)-L-mannopyranoside]: **Solasurine** [27028-76-8]C₃₉H₆₃NO₁₁ 721.927Alkaloid from *Solanum surattense*, *Solanum brusquense*, *Solanum xanthocarpum*, *Solanum elaeagnifolium* and *Solanum viarum* (Solanaceae). Mp 228-230°.3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-galactopyranoside]: **Isoanguivine**

[163252-96-8]

C₄₄H₇₁NO₁₅ 854.043From roots of *Solanum anguivi* (Solanaceae). Cryst. (MeOH aq.). Mp 286-291° dec. [α]_D²⁰ -88.8 (c, 0.64 in Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranoside]: **Anguivine**

[163135-92-0]

C₄₄H₇₁NO₁₅ 854.043From roots of *Solanum anguivi* (Solanaceae). Amorph. [α]_D²² -83.2 (c, 0.87 in Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-xylopyranoside]: **Ravifoline**

[157382-29-1]

C₄₄H₇₁NO₁₄ 838.043Alkaloid from *Solanum plataniifolium*. Cryst. (MeOH aq.). Mp 280-282°. [α]_D²⁵ -104 (Py).3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Solamargine**

[20311-51-7]

C₄₅H₇₃NO₁₅ 868.069Alkaloid from *Solanum marginatum* and a very large number of *Solanum spp.* (Solanaceae). Shows strong antifungal and antiyeast activity. Cytotoxic. Mp 301° dec. [α]_D²⁰ -105 (MeOH). [α]_D²⁰ -102 (Py).3-O-[α-L-Rhamnopyranosyl-(1→3)-[α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Solanelagnine** [181023-53-0]C₄₅H₇₃NO₁₅ 868.069Alkaloid from *Solanum elaeagnifolium*.3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-glucopyranosyl-(1→3)]-β-D-galactopyranoside]: **Solasonine**. Solasodamine [19121-58-5]C₄₅H₇₃NO₁₆ 884.069Alkaloid from *Solanum sodomeum* and a very large number of *Solanum spp.* (Solanaceae). Molluscicidal agent. Sol. MeOH; poorly sol. H₂O, hexane. Mp 300-301°. [α]_D²⁰ -74 (MeOH). Log P 0.12 (uncertain value) (calc).▶ LD₅₀ (mus, ipr) 30 mg/kg. WF1450000

3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-D-glucopyranosyl-(1→4)]-β-D-glucopyranoside]: [132922-48-6]

C₄₅H₇₃NO₁₆ 884.069Alkaloid from the bulbs of *Lilium brownii* var. *colchesteri* (Liliaceae). Amorph. powder. [α]_D²⁸ -77.3 (c, 0.11 in MeOH).3-O-[α-L-Arabinopyranosyl-(1→3)-[α-L-rhamnopyranosyl-(1→2)]-α-L-rhamnopyranosyl-(1→4)]-β-D-glucopyranoside]: **Robustine**

[156431-09-3]

C₅₀H₈₁NO₁₉ 1000.185Alkaloid from roots of *Solanum robustum* (Solanaceae). Needles (MeOH aq.). Mp 217-223°. [α]_D²⁰ -73.7 (c, 0.66 in Py).3-O-[β-D-Xylopyranosyl-(1→4)-α-L-rhamnopyranosyl-(1→4)-[β-D-xylopyranosyl-(1→3)]-β-D-glucopyranoside]: **Incanumine**

[128585-03-5]

C₄₉H₇₉NO₁₉ 986.158Alkaloid from the fruit of *Solanum incanum*. Powder (MeOH). Mp > 300. [α]_D²⁰ +78 (c, 0.1 in MeOH).3-O-[β-D-Xylopyranosyl-(1→2)-α-L-rhamnopyranosyl-(1→4)-[α-L-rhamnopyranosyl-(1→2)]-β-D-glucopyranoside]: **Sycophantine**

[175701-97-0]

C₅₀H₈₁NO₁₉ 1000.185Constit. of *Solanum sycophanta* and *Solanum coccineum*. Cryst. (MeOH). Mp 269-271° dec. [α]_D²⁰ -90 (c, 0.6 in MeOH).3-O-[α-L-Rhamnopyranosyl-(1→2)-[α-L-rhamnopyranosyl-(1→4)]-β-D-xylopyranosyl-(1→3)]-β-D-glucopyranoside]: **Arudonine**

[737763-55-2]

C₅₀H₈₁NO₁₉ 1000.185Alkaloid from the root bark of *Solanum arundo*. Allelopathic agent.

3-O-[α-L-Rhamnopyranosyl-(1→2)-[β-

- d*-glucopyranosyl-(1→2)-O-β-*D*-glucopyranosyl-(1→3)]-β-*D*-galactopyranoside]: **Solaradixine**
[28512-69-8]
C₅₁H₈₃NO₂₁ 1046.211
Alkaloid from roots of *Solanum laciniatum* and fruits of *Solanum mammosum* (Solanaceae). Mp 275-278°. [α]_D²⁰ -69.7 (Py). [α]_D²⁰ -51.9 (MeOH). Partial enzymic hydrol. gives Solasoline.
- 3-O-[β-*D*-Glucopyranosyl-(1→2)-β-*D*-glucopyranosyl-(1→4)]-α-*L*-rhamnopyranosyl-(1→2)]-β-*D*-galactopyranoside]: **Solasuaveoline**
[201805-24-5]
C₅₁H₈₃NO₂₁ 1046.211
Alkaloid from *Solanum suaveolens* (Solanaceae). Cryst. (MeOH aq.). Mp 277-278° dec. [α]_D²² -64.6 (c, 1 in Py).
- 3-O-[β-*D*-Glucopyranosyl-(1→6)-β-*D*-glucopyranosyl-(1→3)]-α-*L*-rhamnopyranosyl-(1→2)]-β-*D*-galactopyranoside]: **Isosolasuaveoline**. *Solashabanine*
[37449-88-0]
[201805-28-9]
C₅₁H₈₃NO₂₁ 1046.211
Isol. from *Solanum aviculare*, *Solanum laciniatum* and *Solanum suaveolens* (Solanaceae). Needles (MeOH). Mp 277-279° dec. [α]_D²³ -71.4 (c, 1 in Py). Struct. of *Solashabanine* not fully substantiated.
- 3-O-[β-*D*-Glucopyranosyl-(1→6)-β-*D*-glucopyranosyl-(1→2)-β-*D*-glucopyranosyl-(1→3)]-α-*L*-rhamnopyranosyl-(1→2)]-β-*D*-galactopyranoside]: **Solaradinine**
[37647-41-9]
C₅₇H₉₃NO₂₆ 1208.353
Isol. from *Solanum laciniatum* and *Solanum aviculare* (Solanaceae). Mp 227-230°. [α]_D²⁰ -45.8 (50% EtOH aq.).
- Glycoside (1): β-Solanigrine**
Alkaloid from *Solanum americanum*, *Solanum atropurpureum*, *Solanum boerhaaviaefolium*, *Solanum curtipes*, *Solanum gracile*, *Solanum humile*, *Solanum integrifolium*, *Solanum melano-cerasum* (garden huckleberry), *Solanum miniatum*, *Solanum nigrum*, *Solanum paraense* and *Solanum villosum* (Solanaceae). Struct. unknown. Hydrol. gives 2 L-rhamnose, 1 D-glucose and 1 D-galactose. No CAS no., isol. pre 1967.
- Glycoside (2): Isosolasonine**
C₄₅H₇₃NO₁₆ 884.069
Alkaloid from *Solanum tomatillo* (Solanaceae). Mp 224-229° dec. [α]_D²⁰ -75. Struct. unknown. Hydrol. gives glucose, rhamnose and galactose. No CAS no., isol. pre 1967.
- Glycoside (3): Solapersine**
[58037-78-8]
C₄₉H₇₉NO₂₀ 1002.158
Alkaloid from *Solanum persicum* (Solanaceae). Cryst. (MeOH). Mp 282-284°. [α]_D²⁰ -48.1 (c, 0.5 in MeOH). Struct. unknown. Hydrol. gives 1 galactose, 1 glucose and 2 xylose.
- Glycoside (4): Solavilline**
Alkaloid from *Solanum villosum* (Solanaceae). Struct. unknown. Not v. well-characterised. Gives L-rhamnose, D-xylose, D-glucose and D-galactose on hydrol. No CAS no. in 8CI.
- Glycoside (5): α-Solanigrine**
Alkaloid from *Solanum americanum*, *Solanum melano-cerasum* (garden huckleberry) and *Solanum nigrum* (Solanaceae). Sugar residues not identified. No CAS no., isol. pre 1967.
- Glycoside (6): Solakhasianine**
Glycoalkaloid from *Solanum khasianum* (Solanaceae). Needles (EtOH). Mp 251-256° dec. Struct. unknown. Hydrol. gives Solasodine, galactose and rhamnose. No CAS no. in 8CI.
- Glycoside (7): Solatifoline**
[57903-98-7]
Alkaloid from *Solanum plataniifolium* (Solanaceae). Mp 292-295°. [α]_D²⁵ -119. Struct. unknown. Hydrol. gives Solasodine, glucose, galactose and rhamnose.
- O-Ac: O-Acetylsolasodine**
[6159-99-5]
C₂₉H₄₅NO₃ 455.679
Alkaloid from whole plants of *Solanum umbelliferum*. Amorph. powder. Mp 192-194°. [α]_D²⁶ -108.4 (c, 0.62 in CHCl₃).
- N-Hydroxy: N-Hydroxysolasodine**
[142182-57-8]
C₂₇H₄₃NO₃ 429.642
Alkaloid from roots of *Solanum robustum* (Solanaceae). Cryst. (Me₂CO). Mp 206-209° dec. [α]_D²⁰ -119.5 (c, 1.00 in CHCl₃).
- N-Hydroxy, 3-O-[α-L-arabinopyranosyl-(1→3)]-α-L-rhamnopyranosyl-(1→2)]-β-D-glucopyranoside]: N-Hydroxyrobustine**
[156122-58-6]
C₅₀H₈₁NO₂₀ 1016.185
From roots of *Solanum robustum* (Solanaceae). Amorph. [α]_D²⁰ -63.4 (c, 0.68 in Py).
- (3β,22S,25R)-form**
3-O-[β-*D*-Xylopyranosyl-(1→2)-α-*L*-rhamnopyranosyl-(1→4)]-α-*L*-rhamnopyranosyl-(1→2)]-β-*D*-glucopyranoside]: **22-Episcyphantine**
C₅₀H₈₁NO₁₉ 1000.185
Constit. of *Solanum coccineum* and *Solanum sycophanta*. Cryst. (MeOH). Mp 268-272° dec. [α]_D²¹ -52 (c, 0.6 in MeOH).
- (3β,22S,25S)-form**
Tomatidenol. *Soladulcamaridine*. *Tomatid-5-en-3-ol*. δ⁵-*Tomatidenol*
[546-40-7]
Aglycone from many *Solanum* spp. and from *Fritillaria camtschatscensis*. Occurs free in *Solanum trilobatum* and *Solanum dulcamara* (Liliaceae, Solanaceae). Shows antifungal activity. Mp 235-238° (206°, 226-228°). [α]_D²⁰ -45 (CHCl₃). *Soladulcamaridine* was shown to be impure *Tomatidenol*.
- 3-O-[α-*L*-Rhamnopyranosyl-(1→2)-β-*D*-glucopyranoside]: **γ₁-Solamarine**
[11040-30-5]
C₃₉H₆₃NO₁₁ 721.927
Alkaloid from *Solanum dulcamara* (Solanaceae). Small needles + 1H₂O (MeOH aq.). Mp 268-271° dec. [α]_D²² -60.3 (c, 1 in Py).
- 3-O-[α-*L*-Rhamnopyranosyl-(1→4)-β-*D*-glucopyranoside]: **γ₂-Solamarine**. **γ-Solamarine**
[11034-34-7]
C₃₉H₆₃NO₁₁ 721.927
Alkaloid from *Solanum dulcamara*, *Solanum dasyphyllum*, *Solanum juzepczukii* (bitter potato) and *Solanum schimperianum* (Solanaceae). Mp 243-248° dec. [α]_D²⁰ -86 (c, 0.36 in Py).
- 3-O-[β-*D*-Glucopyranosyl-(1→3)-β-*D*-galactopyranoside]: **δ-Solamarine**
[15299-07-7]
C₃₉H₆₃NO₁₂ 737.926
Alkaloid from *Solanum dulcamara* (Solanaceae). Small needles (MeOH aq.). Mp 265-269° dec. [α]_D²¹ -39.6 (c, 0.92 in Py).
- 3-O-[α-*L*-Rhamnopyranosyl-(1→2)-α-*L*-rhamnopyranosyl-(1→4)]-β-*D*-glucopyranoside]: **β-Solamarine**
[3671-38-3]
C₄₅H₇₃NO₁₅ 868.069
Alkaloid from *Solanum dulcamara*, *Solanum dasyphyllum*, *Solanum juzepczukii* (bitter potato), *Solanum macrocarpum* and *Solanum schimperianum* (Solanaceae). Antineoplastic agent. Shows activity against mouse 180 sarcoma. Mp 275-277° dec. [α]_D²⁰ -85.6 (c, 0.4 in Py). Log P 1.31 (uncertain value) (calc).
- 3-O-[α-*L*-Rhamnopyranosyl-(1→2)-β-*D*-glucopyranosyl-(1→3)-β-*D*-galactopyranoside]: **α-Solamarine**
[20318-30-3]
C₄₅H₇₃NO₁₆ 884.069
Alkaloid from *Solanum dulcamara*, *Solanum macrocarpum*, *Solanum dasyphyllum* and *Solanum juzepczukii* (bitter potato) (Solanaceae). Mp 278-281° dec. [α]_D²⁰ -45 (c, 0.65 in Py).
- 3-O-[β-*D*-Xylopyranosyl-(1→2)-α-*L*-rhamnopyranosyl-(1→4)]-α-*L*-rhamnopyranosyl-(1→2)]-β-*D*-glucopyranoside]: **22,25-Diepiscyphantine**
[175779-31-4]
C₅₀H₈₁NO₁₉ 1000.185
Alkaloid from *Solanum sycophanta*. Cryst. (MeOH). Mp 268-272° dec. [α]_D²⁰ -52 (c, 0.6 in MeOH).
- O-[α-*L*-Rhamnopyranosyl-(1→2)]-β-*D*-xylopyranosyl-(1→2)-α-*L*-rhamnopyranosyl-(1→4)]-β-*D*-galactopyranoside]: **Solaculine A**
[408345-39-1]
C₅₀H₈₁NO₁₉ 1000.185
Alkaloid from the root bark of *Solanum aculeastrum*. Cryst. Mp 220-224°. [α]_D²⁴ -52 (c, 0.2 in MeOH/CHCl₃).
- 3-O-[β-*D*-Glucopyranosyl-(1→2)]-β-*D*-xylopyranosyl-(1→3)]-β-*D*-glucopyranosyl-(1→4)-β-*D*-galactopyranoside]: **Dehydrotomatine**. *Tomatidenol*

3-lycotetraoside
[157604-98-3]
C₅₀H₈₁NO₂₁ 1032.184
Alkaloid from *Lycopersicon esculentum* (tomato) (Solanaceae).

O,N-Di-Ac: Mp 160-162°. [α]_D -39.6 (c, 0.36 in MeOH).

Uhle, F.C. *et al.*, *J.A.C.S.*, 1954, **76**, 4245-4246 (synth, struct)

Schreiber, K. *et al.*, *Planta Med.*, 1958, **6**, 435-439 (*Solanigrines*)

Bianchi, E. *et al.*, *CA*, 1961, **55**, 26002g (*Isosolasonine*)

Boll, P.M. *et al.*, *Acta Chem. Scand.*, 1962, **16**, 1819-1830 (*Solanum dulcamara glycosides*)

Wolters, B. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1964, **297**, 748-754 (*Tomatidenol, activity*)

Schreiber, K. *et al.*, *Tetrahedron*, 1964, **20**, 1939-1945; 1966, **22**, 3591-3595 (struct, synth)

Budzikiewicz, H. *et al.*, *Tetrahedron*, 1964, **20**, 2267-2278 (ms)

Boll, P.M. *et al.*, *Acta Chem. Scand.*, 1965, **19**, 1365-1370 (pmr, config)

Bognár, R. *et al.*, *Pharmazie*, 1965, **20**, 40-42 (*Solavilline*)

Rönsch, H. *et al.*, *Phytochemistry*, 1966, **5**, 1227-1233 (*Solamarines*)

Wolters, B. *et al.*, *Planta Med.*, 1966, **14**, 392-401 (*activity*)

Maitli, P.C. *et al.*, *Indian J. Chem.*, 1968, **6**, 547-548 (*Solakhastanine*)

Puroshathaman, K.K. *et al.*, *Aust. J. Chem.*, 1969, **22**, 1569-1570 (*isol*)

Kessar, S.V. *et al.*, *Tetrahedron*, 1971, **27**, 2153-2159 (*synth*)

Bite, P. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1972, **73**, 361-362 (*Solaradixine, Solashabanine, Solaradinine*)

Singh, S. *et al.*, *Phytochemistry*, 1974, **13**, 2020-2022 (*Solaphumbine*)

Novruzov, E.N. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 453 (*Solapersine*)

Briggs, L.H. *et al.*, *J.C.S. Perkin I*, 1975, 2455-2457 (*Solamargine*)

Puri, R.K. *et al.*, *Phytochemistry*, 1975, **14**, 2096 (*Solatifoline*)

Weston, R.J. *et al.*, *Aust. J. Chem.*, 1977, **30**, 917-921 (cmr)

Dnyansagar, V.R. *et al.*, *Planta Med.*, 1977, **31**, 21-25 (*Solasurine*)

Bird, G.J. *et al.*, *Aust. J. Chem.*, 1979, **32**, 783-796 (cmr)

Mahato, S.B. *et al.*, *Phytochemistry*, 1980, **19**, 2017-2020 (*Solamargines*)

Ripperger, H. *et al.*, *Alkaloids (Academic Press)*, 1981, **19**, 81-192 (rev)

Neszmelyi, A. *et al.*, *Phytochemistry*, 1988, **27**, 603-605 (*Solaradinine, Solashabanine*)

Mimaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 3055-3059 (*rhamnosylglucosylglucoside*)

Lin, C.-N. *et al.*, *J. Nat. Prod.*, 1990, **53**, 513-516 (*Incanumine*)

Ripperger, H. *et al.*, *Phytochemistry*, 1991, **30**, 1299-1301 (*Solasodine, Tomatidenol*)

Ripperger, H. *et al.*, *Phytochemistry*, 1992, **31**, 1837-1839 (*N-Hydroxysolasodine*)

Ripperger, H. *et al.*, *Annalen*, 1994, 517-520 (*Robustine, N-Hydroxyrobustine*)

Puri, R. *et al.*, *J. Nat. Prod.*, 1994, **57**, 587-596 (*Ravifoline*)

Ripperger, H. *et al.*, *Phytochemistry*, 1994, **37**, 1725-1727 (*Anguivine, Isoanguivine*)

Hanna, A.G. *et al.*, *Fitoterapia*, 1996, **67**, 223-226 (*Solanelagnine*)

Kim, Y.C. *et al.*, *J. Nat. Prod.*, 1996, **59**, 283-285 (*Ac. glucoside*)

Lorey, S. *et al.*, *Phytochemistry*, 1996, **41**, 1633-1635 (*Sycophantine, 22-Episcyphantine*)

Friedman, M. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 1541-1547 (*Dehydrotomatine*)

Usubillaga, A. *et al.*, *Phytochemistry*, 1997, **44**, 537-543 (*Sycophantine, 22-Episcyphantine, Diepiscyphantine*)

Ripperger, H. *et al.*, *Phytochemistry*, 1997, **46**, 1279-1282 (*Solasuaveoline, Isosolasuaveoline*)

Hu, K. *et al.*, *Planta Med.*, 1999, **65**, 35-38 (*isol, activity*)

Weissenberg, M. *et al.*, *Phytochemistry*, 2001, **58**, 501-508 (*β-Solamarine, cmr, ms*)

Wanyonyi, A.W. *et al.*, *Phytochemistry*, 2002, **59**, 79-84 (*Solaculine A, Solamargine, α-Solamarine*)


Fukuhara, K. *et al.*, *Phytochemistry*, 2004, **65**, 1283-1286 (*Arudonine*)

Vega-Baez, J.L. *et al.*, *Acta Cryst. E*, 2006, **62**, 4741-4743 (*Solasodine, cryst struct*)

Zha, X. *et al.*, *Chem. Biodiversity*, 2007, **4**, 25-31 (*Solasodine, synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, POJ000 (*Solasodine*)

Spirosol-4-en-3-one, 9CI S-457

C₂₇H₄₁NO₂ 411.626

(22R,25R)-form

Solasodenone

[17094-86-9]

Alkaloid from *Solanum hainanense* and *Solanum aviculare* spp. Needles (Me₂CO aq.). Poorly sol. H₂O. Mp 183-184°. [α]_D²⁶ +28 (c, 0.4 in CHCl₃).

N-Ac:

Cryst. (Et₂O). Mp 124°. [α]_D²⁵ +40.1 (c, 0.513 in CHCl₃).

4,5β-Dihydro: Solasodan-3-one

C₂₇H₄₃NO₂ 413.642

Alkaloid from leaves of *Solanum aviculare* (Solanaceae). Needles (Me₂CO aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 169-170° (164-165°). [α]_D²⁰ -49 (c, 2.8 in CHCl₃).

Tolstikov, G.A. *et al.*, *Khim. Prir. Soedin.*, 1968, **3**, 286; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **3**, 242 (*synth*)

Radeglia, R. *et al.*, *Tet. Lett.*, 1977, **18**, 903-906 (cmr)


Adam, G. *et al.*, *Phytochemistry*, 1978, **17**, 1070-1071 (*Solasodenone, isol, synth, ms, pmr, uv*)

Adam, G. *et al.*, *Tet. Lett.*, 1980, **21**, 1931-1932 (*synth*)

Rowan, D.D. *et al.*, *Phytochemistry*, 1983, **22**, 2102-2104 (*Solasodenone, Solasodan-3-one*)

Spirostan-3-amine S-458

3-Aminospirostane



(3β,5α,25R)-form

C₂₇H₄₅NO₂ 415.658

(3β,5α,25R)-form

Isojurubidine

[32562-75-7]

Alkaloid from the roots of *Solanum paniculatum* (Solanaceae). Mp 185-187°. [α]_D -63 (c, 0.032 in CHCl₃).

N-Ac: Mp 262-265°.

5,6-Didehydro: Antillaridine. Δ⁵-Isojurubidine

[155416-19-6]

C₂₇H₄₃NO₂ 413.642

Alkaloid from *Solanum antillarum* (Solanaceae).

(3β,5α,25S)-form

Jurubidine

[6084-44-2]

Cyclised aglycone from Jurubine, J-60. Mp 182-186°. [α]_D¹⁸ -78.7 (c, 1.8 in CHCl₃).

(3α,25R)-form

5,6-Didehydro: Antillidine. 3α-Δ⁵-Isojurubidine

[155416-18-5]

C₂₇H₄₃NO₂ 413.642

From *Solanum antillarum* (Solanaceae).


Ripperger, H. *et al.*, *Chem. Ber.*, 1967, **100**, 1725

Irismetov, M.P. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1980, 81 (*synth*)

Coll, P.F. *et al.*, *Rev. Cubana Farm.*, 1992, **6**, 66; *CA*, **121**, 5053w (*Antillaridine, Antillidine*)

Spirostaphylotrichin E S-459

[123548-68-5]

C₁₄H₁₇NO₅ 279.292

Metab. of *Staphylotrichum coccosporum*. Cryst. (CH₂Cl₂/Et₂O). Mp 107-109°. λ_{max} 225 (ε 12100) (EtOH) (Derep).

6-Epimer: Spirostaphylotrichin F

[123548-49-2]

C₁₄H₁₇NO₅ 279.292

From *Staphylotrichum coccosporum*.

Cryst. (CH₂Cl₂/Et₂O). Mp 114-116°. λ_{max} 225 (ε 12100) (EtOH) (Derep). λ_{max} 225 (ε 12200) (MeOH) (Berdy).

7-Alcohol: Spirostaphylotrichin L

[123573-21-7]


C₁₄H₁₉NO₅ 281.308

Metab. of *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 131-137°. Configs. of the 6- and 7-OH groups not certain.

Sandmeier, P. *et al.*, *Helv. Chim. Acta*, 1989, **72**, 784; 1107 (*isol, pmr, cmr, ms, struct*)

Spirostaphylotrichin G

[123548-69-6]

C₁₄H₁₅NO₅ 277.276

Metab. of *Staphylotrichum coccosporum*. Cryst. Mp 160-176°. λ_{max} 202 (ε 10800); 232 (ε 10000); 279 (ε 3200) (EtOH) (Derep).

6-Epimer: Spirostaphylotrichin H

[124441-13-0]


C₁₄H₁₅NO₅ 277.276

From *Staphylotrichum coccosporum*. λ_{max} 202 (ε 10800); 232 (ε 10000); 279 (ε 3200) (EtOH) (Derep).

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 1107 (*isol, struct*)

Spirostaphylotrichin K

[123548-71-0]


(4*S*,6*R*)-formC₁₄H₁₇NO₅ 279.292

Isol. from *Staphylotrichum coccosporum*. Mp 200-205°. λ_{max} 215 (sh) (ε 7000); 235 (sh) (ε 5000) (EtOH) (Derep).

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 1107 (*isol, pmr, cmr, struct*)

Spirostaphylotrichin M

[123548-72-1]

C₁₄H₁₇NO₅ 279.292


Metab. of *Staphylotrichum coccosporum*. Cryst. Mp 187-190°. λ_{max} 226 (ε 15100) (EtOH) (Derep).

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 1107 (*isol, pmr, cmr, struct*)

Spirostaphylotrichin O

S-463

4,9,10-Trihydroxy-2-methoxy-3-methylene-6-(1-propenyl)-2-azaspiro[4.5]dec-7-en-1-one, 9CI
[129666-00-8]

C₁₄H₁₉NO₅ 281.308

Isol. from a blocked mutant of *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 147-151°.

4-Ketone: Spirostaphylotrichin N

[129665-99-2]

C₁₄H₁₇NO₅ 279.292

From *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 92-95°.

4-Ketone, N-demethoxy: Spirostaphylotrichin P

[129666-01-9]


C₁₃H₁₅NO₄ 249.266

From *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 141-144°.

Sandmeier, P. et al., *Helv. Chim. Acta*, 1990, **73**, 975 (*isol, pmr, cmr, struct*)

Spirostaphylotrichin Q

S-464

C₁₄H₁₇NO₅ 279.292


Metab. of *Staphylotrichum coccosporum* and *Curvularia pallescens*. Cryst. (CH₂Cl₂/Et₂O). Mp 163-169°. λ_{max} 205 (ε 8900); 291 (ε 14200) (EtOH) (Berdy).

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 784 (*isol, pmr, cmr, struct*)

Abraham, W.-R. et al., *Tetrahedron*, 1995, **51**, 4947 (*isol*)

Spirostaphylotrichin R

S-465

C₁₄H₁₉NO₆ 297.307

Metab. of *Staphylotrichum coccosporum* and *Curvularia pallescens*. Cryst. Mp 157-166°. λ_{max} 204 (ε 9100); 291 (ε 14100) (EtOH) (Berdy).

3-Me ether: Spirostaphylotrichin VC₁₅H₂₁NO₆ 311.334

From *Curvularia pallescens*. Glassy, yellowish cryst. [α]_D²⁷ -147.6 (c, 0.50 in MeOH).

3-Epimer: Spirostaphylotrichin UC₁₄H₁₉NO₆ 297.307

From *Curvularia pallescens*. Not completely separated from Spirostaphylotrichin R.


Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 784 (*isol, pmr, cmr, struct*)

Abraham, W.-R. et al., *Tetrahedron*, 1995, **51**, 4947 (*isol*)

Spirostaphylotrichin S

S-466

[123620-40-6]

C₁₄H₁₉NO₆ 297.307

Metab. of *Staphylotrichum coccosporum*. Cryst. Mp 166-173°. λ_{max} 215 (sh) (ε 7000); 235 (sh) (ε 5000) (EtOH) (Derep).

Stereoisomer: Spirostaphylotrichin I

[123548-70-9]

C₁₄H₁₉NO₆ 297.307


Metab. of *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 167-176°.

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 1107 (*isol, pmr, cmr, struct*)

Spirostaphylotrichin T

S-467

[129681-31-8]

C₁₄H₁₉NO₅ 281.308


Tentative config. shown. Isol. from blocked mutant of *Staphylotrichum coccosporum*. Cryst. (Et₂O). Mp 152-159°.

Sandmeier, P. et al., *Helv. Chim. Acta*, 1990, **73**, 975 (*isol, pmr, cmr, struct*)

Spirotryprostatin A

S-468

[182234-25-9]



Relative Configuration

$C_{22}H_{25}N_3O_4$ 395.457
Metab. from *Aspergillus fumigatus* BM939 and the marine-derived *Aspergillus sydowi* PFW1-13. Mammalian cell cycle inhibitor. Pale yellow amorph. powder. $[\alpha]_D^{26}$ -34 (c, 0.10 in $CHCl_3$). λ_{max} 219 (ε 15920); 245 (sh) (ε 4400); 270 (ε 3020); 280 (sh) (ε 2960) (MeOH).

Demethoxy, 8,9-didehydro: **Spirotryprostatin B**

[182234-26-0]
 $C_{21}H_{21}N_3O_3$ 363.415
Prod. by *Aspergillus fumigatus* BM939. Mammalian cell cycle inhibitor. Pale yellow cryst. Mp 137-138°. $[\alpha]_D^{22}$ -162.1 (c, 0.92 in $CHCl_3$). λ_{max} 212 (ε 36660); 227 (sh) (ε 28820); 242 (sh) (ε 24830); 272 (sh) (ε 17350); 286 (sh) (ε 14810) (MeOH).

Cui, C.-B. *et al.*, *Tetrahedron*, 1996, **52**, 12651-12666 (*isol, uv, ir, pmr, cmr, ms*)
Edmondson, S. *et al.*, *J.A.C.S.*, 1999, **121**, 2147-2155 (*synth*)

von Nussbaum, F. *et al.*, *Angew. Chem., Int. Ed.*, 2000, **39**, 2175-2178 (*Spirotryprostatin B, synth*)

Wang, H. *et al.*, *J.O.C.*, 2000, **65**, 4685-4693 (*Spirotryprostatin B, synth*)

Bagul, T.D. *et al.*, *Org. Lett.*, 2002, **4**, 249-251 (*Spirotryprostatin B, synth*)

Sebahar, P.R. *et al.*, *Tetrahedron*, 2002, **58**, 6311-6322 (*Spirotryprostatin B, synth*)
Meyers, C. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 694-696 (*Spirotryprostatin B, synth*)

Miyake, F.Y. *et al.*, *Org. Lett.*, 2004, **6**, 4249-4251 (*synth*)


Onishi, T. *et al.*, *Tetrahedron*, 2004, **60**, 9503-9515 (*synth*)

Marti, C. *et al.*, *J.A.C.S.*, 2005, **127**, 11505-11515 (*Spirotryprostatin B, synth*)

Trost, B.M. *et al.*, *Org. Lett.*, 2007, **9**, 2763-2766 (*synth*)

Spixianine

[112494-63-0]



$C_{19}H_{19}NO_5$ 341.363
Alkaloid from the trunk bark of Colombian *Duguetia spixiana* (Annonaceae). Not isol. pure; always mixed with Duguexine in P-781. Pure material obt. by redn. of the N-oxide.

N-Oxide: **Spixianine N-oxide**


[112494-64-1]
 $C_{19}H_{19}NO_6$ 357.362
Alkaloid from *Duguetia spixiana* (Annonaceae). Amorph. $[\alpha]_D$ -84 (c, 0.68 in $CHCl_3$).

Debourges, D. *et al.*, *J. Nat. Prod.*, 1987, **50**, 664 (*isol, uv, pmr, ms, struct*)

Splendidine

S-470

1,2,4-Trimethoxy-7H-dibenzo[de,g]quinolin-7-one, 9CI
[68353-25-3]




$C_{19}H_{15}NO_4$ 321.332

Alkaloid from *Abuta rufescens* (*Abuta splendida*) (Menispermaceae). Bright-yellow needles (MeOH/ $CHCl_3$). Mp 235°.

Skiles, J.W. *et al.*, *Can. J. Chem.*, 1979, **57**, 1642 (*isol, ir, uv, ms, pmr, synth, struct*)
Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1997, **60**, 1328-1330 (*isol, cmr*)

Spongiacysteine

S-471



Absolute Configuration

$C_{17}H_{33}NO_5S$ 363.517


Isol. from the Japanese sponge *Spongia* sp. Oil. $[\alpha]_D^{20}$ -238 (c, 0.02 in MeOH) (natural). $[\alpha]_D^{24}$ -115 (c, 0.02 in MeOH) (synthetic).

Kobayashi, K. *et al.*, *Chem. Lett.*, 2004, **33**, 1262-1263 (*isol, synth, pmr, cmr*)

Spondidine A

S-472

[263143-53-9]



R = -CH₂COOH

$C_{23}H_{34}NO_2^{\oplus}$ 356.527

Alkaloid from a *Spongia* sp. Inhibitor of human phospholipase A₂. $[\alpha]_D$ -16.2 (c, 0.01 in MeOH). Counterion not specified.

17-Acetoxy: **Spondidine B**

[263143-54-0]

$C_{25}H_{36}NO_4^{\oplus}$ 414.564

Alkaloid from a *Spongia* sp. Inhibitor of human phospholipase A₂. $[\alpha]_D$ +7.8 (c, 0.01 in MeOH). Counterion not specified.

De Marino, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 323-326

Spondidine D

S-473

[263143-56-2]

As Spondidine A, S-472 with

R = -CH₂CH₂SO₃H

$C_{23}H_{36}NO_5^{\oplus}$ 406.608

Alkaloid from a *Spongia* sp. Inhibitor of human phospholipase A₂. $[\alpha]_D$ -6 (c, 0.02

in MeOH). Counterion not specified.


De Marino, S. *et al.*, *J. Nat. Prod.*, 2000, **63**, 323-326

Sporidesmin

S-474

Sporidesmin A

[1456-55-9]



Absolute configuration

n = 2

$C_{18}H_{20}ClN_3O_6S_2$ 473.957

Alkaloid from *Pithomyces chartarum*. Causes facial eczema in sheep. Shows antitumour props. Sol. MeOH, Et₂O, bases; poorly sol. H₂O, hexane. λ_{max} 219 (ε 31600); 254 (ε 12000); 304 (ε 2570) (Et₂O) (Derep). λ_{max} 218 (ε 39820); 254 (ε 13200); 302 (ε 2820) (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 1 mg/kg. KB4740000

CCl₄ complex:

Cryst. Mp 130-134° (sintering from 109°). $[\alpha]_D^{24}$ -23 (c, 0.96 in MeOH).

Di-Ac:

Needles. Mp 170-171°. $[\alpha]_D$ -25 (c, 1.2 in MeOH).

8-N-De-Me: **Sporidesmin J**

[69799-25-3]

$C_{17}H_{18}ClN_3O_6S_2$ 459.93

Metab. of *Pithomyces chartarum*. Cryst. (Et₂O). Mp 168-169°. $[\alpha]_D^{18}$ +43 (c, 0.25 in $CHCl_3$). λ_{max} 219 (ε 31600); 254 (ε 12000); 304 (ε 2570) (Et₂O) (Derep). λ_{max} 216 (ε 23400); 252 (ε 11300); 290 (ε 8000) (MeOH) (Berdy).

11-Deoxy: **Sporidesmin B**

[3351-96-0]

$C_{18}H_{20}ClN_3O_5S_2$ 457.958

Constit. of *Pithomyces chartarum*. Needles (Me₂CO aq.). Sol. MeOH, $CHCl_3$; fairly sol. Et₂O, hexane, C₆H₆; poorly sol. H₂O. Mp 183°. $[\alpha]_D^{21}$ -27 (c, 1 in MeOH). λ_{max} 218 (ε 31620); 256 (ε 12000); 307 (ε 2570) (EtOH) (Berdy).

▶ Toxic.

Hodges, R. *et al.*, *Chem. Ind. (London)*, 1963, 42 (*isol, ir, uv, pmr*)

Shannon, J.S. *et al.*, *Tet. Lett.*, 1963, 801 (*ms*)

Herrmann, H. *et al.*, *J.C.S.*, 1964, 4315 (*ord, cd*)

Beecham, A.F. *et al.*, *Tet. Lett.*, 1966, 3131 (*cryst struct*)

Kishi, Y. *et al.*, *J.A.C.S.*, 1973, **95**, 6439 (*synth*)

Nagarajan, R. *et al.*, *J.A.C.S.*, 1973, **95**, 7212 (*cd, ord*)

Nakatsuka, S. *et al.*, *Tet. Lett.*, 1974, 1549 (*synth*)

Ronaldson, J.W. *et al.*, *Aust. J. Chem.*, 1976, **29**, 2307; 1981, **34**, 1215 (*cmr, ir*)

White, E.P. *et al.*, *Mycotoxic Fungi, Mycotoxins, Mycotoxicoses*, 1977, **1**, 427 (*rev*)

Rahman, R. *et al.*, *J.C.S. Perkin 1*, 1978, 1476 (*Sporidesmin J*)


Wróbel, J.T. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 70 (*rev*)

Gallagher, R.T. *et al.*, *J. Agric. Food Chem.*, 1992, **40**, 701 (*synth*)

Miles, C.O. *et al.*, *J. Agric. Food Chem.*, 1992, **40**, 2458 (*pmr*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 584

Sporidesmin C S-475
[7051-84-5]



Absolute Configuration

$C_{18}H_{20}ClN_3O_6S_3$ 506.023
Metab. of *Pithomyces chartarum*. Toxin. Possesses antitumour props. Cryst. λ_{max} 217 (ϵ 33100); 252 (ϵ 16600); 295 (ϵ 3160) (MeOH) (Derep).


Di-Ac:

Needles (2-propanol). Mp 195-200° (resolidifies and remelts at 230-240° dec.). $[\alpha]_D^{25}$ -215 (c, 0.46 in $CHCl_3$).

Hodges, R. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1059 (*isol*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 590

Sporidesmin D S-476
[18180-71-7]



Absolute Configuration

$C_{20}H_{26}ClN_3O_6S_2$ 504.027
Metab. of *Pithomyces chartarum*. Toxin. Shows antibacterial activity. Cryst. + Et_2O . Sol. MeOH, $CHCl_3$; fairly sol. Et_2O ; poorly sol. H_2O , hexane. Mp 105-107°. $[\alpha]_D^{20}$ +58 (c, 0.1 in $CHCl_3$). λ_{max} 219 (ϵ 31600); 254 (ϵ 12000); 304 (ϵ 2570) (Et_2O) (Derep). λ_{max} 216 (ϵ 28200); 252 (ϵ 10000); 300 (ϵ 1900) (EtOH) (Berdy).

Di-Ac: Mp 202-204°. $[\alpha]_D^{20}$ +67 (c, 0.1 in $CHCl_3$).

Rahman, R. *et al.*, *Chem. Comm.*, 1967, 1032 (*isol*)

Jamieson, W.D. *et al.*, *J.C.S. (C)*, 1969, 1564 (*isol, struct*)

Ronaldson, J.W. *et al.*, *Aust. J. Chem.*, 1976, **29**, 2307; 1981, **34**, 1215

White, E.P. *et al.*, *CA*, 1978, **88**, 84087m (*rev*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 591

Sporidesmin E S-477
[22327-77-1]

As Sporidesmin, S-474 with $n = 3$

$C_{18}H_{20}ClN_3O_6S_3$ 506.023
Toxic metab. of *Pithomyces chartarum*. Shows antibacterial and antitumour props. Cryst. + Et_2O . Mp 180-187° (155-165°). $[\alpha]_D^{20}$ -131 (c, 0.065 in $CHCl_3$). λ_{max} 217 (ϵ 33100); 252 (ϵ 16600); 295 (ϵ 3160) (MeOH) (Derep).

Di-Ac: [24569-98-0]


Mp 112-115°.

Rahman, R. *et al.*, *J.C.S. (C)*, 1969, 1665 (*isol*)

White, E.P. *et al.*, *CA*, 1978, **88**, 84087m (*rev*)
Ronaldson, J.W. *et al.*, *Aust. J. Chem.*, 1981, **34**, 1215 (*ir*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 594

Sporidesmin F S-478
[23044-40-8]



Absolute Configuration

$C_{19}H_{22}ClN_3O_6S$ 455.918
Metab. of *Pithomyces chartarum*. Fungal toxin. Amorph. solid. Mp 65-67°. λ_{max} 217 (ϵ 33100); 252 (ϵ 16600); 295 (ϵ 3160) (MeOH) (Derep). λ_{max} 216 (ϵ 28840); 250 (ϵ 13800); 298 (ϵ 2000) (EtOH) (Berdy).

Jamieson, W.D. *et al.*, *J.C.S. (C)*, 1969, 1564 (*isol*)

White, E.P. *et al.*, *CA*, 1978, **88**, 84087m (*rev*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 595

Sporidesmin G S-479
[32608-68-7]

As Sporidesmin, S-474 with $n = 4$

$C_{18}H_{20}ClN_3O_6S_4$ 538.089
Metab. of *Pithomyces chartarum*. Toxin. Shows antitumour props. Cryst. (Et_2O at 4°). Mp 148-153°. $[\alpha]_D^{20}$ -217 (c, 0.023 in $CHCl_3$). λ_{max} 217 (ϵ 33100); 252 (ϵ 16600); 295 (ϵ 3160) (MeOH) (Derep). λ_{max} 216 (ϵ 44700); 250 (ϵ 11500); 298 (ϵ 2950) (EtOH) (Berdy).


Francis, E. *et al.*, *J.C.S. Perkin I*, 1972, 470 (*isol, struct*)

Przybylska, M. *et al.*, *Acta Cryst. B*, 1974, **30**, 597 (*cryst struct*)

White, E.P. *et al.*, *CA*, 1978, **88**, 84087m (*rev*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 596

Sporidesmin H S-480
[65492-17-3]




$C_{18}H_{20}ClN_3O_4S_2$ 441.958
Constit. of *Pithomyces chartarum*. Shows antitumour props. Amorph. solid (Et_2O /petrol). Mp 150-152°. λ_{max} 216 (ϵ 23400); 252 (ϵ 11200); 290 (ϵ 6460) (MeOH) (Derep).

Rahman, R. *et al.*, *J.C.S. Perkin I*, 1978, 1476 (*isol*)

White, E.P. *et al.*, *CA*, 1978, **88**, 84087m (*rev*)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 597

Sporovexin A S-481



$C_{12}H_{14}O_5$ 238.24
Prod. by *Sporormiella vexans*. Pale yellow solid. Mp 177-179°. $[\alpha]_D^{25}$ -22 (c, 0.4 in MeOH). λ_{max} 251 (ϵ 3400) (MeOH).

2-Carboxylethylamide: Sporovexin C

$C_{15}H_{19}NO_6$ 309.318
Prod. by *Sporormiella vexans*. Oil. $[\alpha]_D^{26}$ -11 (c, 0.15 in MeOH). λ_{max} 250 (ϵ 4900) (MeOH).

Soman, A.G. *et al.*, *J. Nat. Prod.*, 1999, **62**, 659-661 (*isol, Sporovexin C*)

Sprintillamine S-482
[1393-32-4]

$C_{28}H_{45}NO_4$ 459.668

Struct. unknown. Alkaloid from *Helleborus viridus*. Respiratory stimulant, convulsant. Small needles ($EtOH$ aq.). Mp 228-229°. Contains one N-Me group.

Hydrochloride: Mp 287-288° (prev. browns and sinters).

Keller, O. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1928, **266**, 545-572 (*isol*)

Franzen, G. *et al.*, *CA*, 1931, **25**, 4052 (*pharmacol*)

Sprintilline S-483
[1401-42-9]

$C_{25}H_{41}NO_3$ 403.604


Struct. unknown. Alkaloid from roots of *Helleborus viridus*. Respiratory stimulant, convulsant. Mp 141-142° (sinters at 132°).

Hydrochloride: Mp 278-279°.

Keller, O. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1928, **266**, 545-572 (*isol*)

Franzen, G. *et al.*, *CA*, 1931, **25**, 4052 (*pharmacol*)

Spylidone S-484




Relative Configuration

$C_{26}H_{39}NO_4$ 429.598
Tetramic acid deriv. Prod. by *Phoma* sp. FKI-1840. Inhibitor of lipid droplet accumulation in macrophages. Oil. $[\alpha]_D^{26}$ +5.6 (c, 0.1 in MeOH). λ_{max} 203 (ϵ 17300) (MeOH).

Koyama, N. *et al.*, *J. Antibiot.*, 2005, **58**, 338-345 (*isol, pmr, cmr*)

Squalamine**S-485**

[148717-90-2]



$C_{34}H_{65}N_3O_5S$ 627.971

Constit. of the shark *Squalus acanthias*. Active against gram-positive and -negative bacteria. Broad-spectrum steroidal antibiotic. Inhibits angiogenesis and solid tumour growth. Antineoplastic agent, haemolytic, laxative, gastric secretion inhibitor, diuretic. ATP-ase inhibitor. Sol. H_2O , MeOH.

Lactate: **Squalamine lactate**, *USAN*. *Evi-zon*. *MSI 1256F* [320725-47-1]

Angiogenesis inhibitor. Antineoplastic used in the treatment of advanced malignancies.

N^{ω} -(3-Aminopropyl): **Trodusquamine**, *INN, USAN*. 7'-N-(3-Aminopropyl)-squalamine. *MSI 1436* [186139-09-3]
 $C_{37}H_{72}N_4O_5S$ 685.066
Constit. of liver of *Squalus acanthias*. Used for treatment of medically significant obesity. Appetite suppressant. Powder.

24-O-Desulfo, 26-sulfooxy: [259173-54-1]
 $C_{34}H_{65}N_3O_6S$ 643.97
Constit. of liver of *Squalus acanthias*. Powder.

24-Desulfo, 24-ketone, 26-sulfooxy: [232613-79-5]
 $C_{34}H_{63}N_3O_6S$ 641.954
Constit. of liver of *Squalus acanthias*. Powder.

24-Desulfo, 24-ketone, 26-[(2-amino-2-carboxyethyl)thio]: [259173-51-8]
 $C_{37}H_{68}N_4O_4S$ 665.035
Constit. of the liver of the dogfish shark *Squalus acanthias*. Powder.

25,26-Didehydro, 24-desulfo, 24-ketone: 3-[[3-[(4-Aminobutyl)amino]propyl]amino]-7-hydroxycholest-25-en-24-one [186139-08-2]
 $C_{34}H_{61}N_3O_2$ 543.875
Constit. of liver of *Squalus acanthias*. Powder.

12 α -Hydroxy: **12-Hydroxysqualamine** [268733-21-7]
 $C_{34}H_{65}N_3O_6S$ 643.97
Constit. of liver of *Squalus acanthias*. Powder.

24-Hydroxymethyl: [268733-22-8]
 $C_{35}H_{67}N_3O_6S$ 657.997
Constit. of liver of *Squalus acanthias*. Powder.

- Moore, K.S. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1993, **90**, 1354-1358 (*isol*)
Wehrli, S.L. *et al.*, *Steroids*, 1993, **58**, 370-378 (*pmr, cmr*)
Moriarty, R.M. *et al.*, *Tet. Lett.*, 1994, **35**, 8103; 1995, **36**, 5139-5142 (*synth*)
Sills, A.K. *et al.*, *Cancer Res.*, 1998, **58**, 2784-2792 (*pharmacol*)
Jones, S.R. *et al.*, *J.O.C.*, 1998, **63**, 3786-3789 (*synth*)
Zhang, X. *et al.*, *J.O.C.*, 1998, **63**, 8599-8603 (*synth*)
Selinsky, B.S. *et al.*, *Biochim. Biophys. Acta*, 2000, **1464**, 135-141 (*pharmacol*)
Rao, M.N. *et al.*, *J. Nat. Prod.*, 2000, **63**, 631-635 (*isol, pmr, cmr, config, derivs*)
Zhang, X. *et al.*, *Org. Lett.*, 2000, **2**, 2921-2922 (*synth*)
Bhargava, P. *et al.*, *Clin. Cancer Res.*, 2001, **7**, 3912-3919 (*Squalamine lactate, pharmacol*)
Ahima, R.S. *et al.*, *Diabetes*, 2002, **51**, 2099-2104 (*MSI 1436, pharmacol*)
Okamura, K. *et al.*, *Chem. Pharm. Bull.*, 2003, **51**, 117-1182 (*synth*)
Hao, D. *et al.*, *Clin. Cancer Res.*, 2003, **9**, 2465-2471 (*clin trial*)
Zhang, D.-H. *et al.*, *Org. Lett.*, 2003, **5**, 3257-3259 (*synth*)
Ciulla, T.A. *et al.*, *Retina*, 2003, **23**, 808-814 (*pharmacol*)
Zhang, D.-H. *et al.*, *Chin. J. Chem.*, 2005, **23**, 176-181 (*synth*)
Zhang, D.-H. *et al.*, *Chin. J. Chem.*, 2005, **23**, 176-181 (*synth*)

Squamigerine**S-486**

[1361-35-9]


 $C_{18}H_{21}NO_5$ 331.368

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Lycoris squamigera* (Amaryllidaceae). Mp 260°. $[\alpha]_D^{25} +165$ ($CHCl_3$).

Hung, S.-H. *et al.*, *Yaoxue Xuebao*, 1964, **11**, 1-14; *CA*, **61**, 3154f (*isol*)

Squamolone**S-487**

2-Oxo-1-pyrrolidinecarboxamide, 9CI. 1-Carbamoyl-2-pyrrolidone [40451-67-0]

 $C_5H_8N_2O_2$ 128.13

An incorrect diazepinedione struct. was originally assigned. Constit. of *Annona squamosa* (sugar apple) and *Hexalobus crispiflorus* (Annonaceae). Also from *Asimina triloba* (pawpaw). Prisms (C_6H_6). Mp 148°.

Yang, T.H. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1972, **19**, 149; *CA*, **78**, 16151w (*isol, uv, pmr, ms*)

Marquez, V.E. *et al.*, *J.O.C.*, 1980, **45**, 5308 (*struct, synth, ir, pmr, ms*)


Achenbach, H. *et al.*, *Annalen*, 1982, 1623 (*isol, cmr*)

Zhao, G.-X. *et al.*, *Phytochemistry*, 1993, **33**, 1065 (*isol, synth*)

Sahai, M. *et al.*, *Indian J. Chem., Sect. B*, 1996, **35**, 510 (*isol, synth, ir, pmr, cmr, ms*)

Squamosamide**S-488**

4-Hydroxy- α -[(4-hydroxy-3-methoxyphenyl)methylene]-N-[2-(4-hydroxyphenyl)ethyl]-3,5-dimethoxybenzeneacetamide, 9CI

 $C_{26}H_{27}NO_7$ 465.502**(E)-form** [142750-35-4]

Isol. from branches of *Annona squamosa* (sugar apple) (Annonaceae). Mp 206-207°.


Yang, X.J. *et al.*, *Yaoxue Xuebao*, 1992, **27**, 185-190; *CA*, **117**, 86699d (*isol, struct*)

Ji, X. *et al.*, *Chin. Chem. Lett.*, 1993, **4**, 297-298 (*synth, pmr, cmr*)

Stachylocin A**S-489****Spirodihydrobenzofuranlactam VI**

[158827-60-2]

[173220-05-8]

 $C_{52}H_{70}N_2O_{10}$ 883.132

Probable struct. Struct. of Spirodihydrobenzofuranlactam VI revised in 2003. Presumed synthetic material showed some differences from the nat. product (2003). Prod. by *Stachybotrys* sp. M6222. Endothelin receptor antagonist. Powder. Sol. MeOH, DMSO, bases; poorly sol. H_2O , C_6H_6 , hexane. $[\alpha]_D^{23} -57$ (c, 0.4 in MeOH). λ_{max} 218 (ϵ 70500); 266 (ϵ 15400); 300 (ϵ 5300) (MeOH) (Berdy). λ_{max} 218 (ϵ 70600); 266 (ϵ 15400); 300 (ϵ 5300) (MeOH/HCl) (Berdy). λ_{max} 222 (ϵ 61600); 274 (ϵ 13200); 300 (ϵ 5300) (MeOH/NaOH) (Berdy).

2 α -Hydroxy: **Stachylocin B**

[158827-61-3]

 $C_{52}H_{70}N_2O_{11}$ 899.132

Prod. by *Stachybotrys* sp. M6222. Endothelin receptor antagonist. Powder. Sol. MeOH, bases, DMSO; poorly sol. H_2O , C_6H_6 , hexane. $[\alpha]_D^{23} -64$ (c, 0.4 in MeOH). λ_{max} 218 (ϵ 70500); 265 (ϵ 15700); 300 (ϵ 5400) (MeOH) (Berdy). λ_{max} 218 (ϵ 70500); 265 (ϵ 15700); 300 (ϵ 5400) (MeOH/HCl) (Berdy). λ_{max} 222 (ϵ 62000); 274 (ϵ 13500); 310 (ϵ 5400) (MeOH/NaOH) (Berdy).

2' β -Hydroxy: **Stachylocin C**

[158827-62-4]

 $C_{52}H_{70}N_2O_{11}$ 899.132

Prod. by *Stachybotrys* sp. M6222. Endothelin receptor antagonist. Powder. Sol. MeOH, DMSO, bases; poorly sol. H_2O , hexane, C_6H_6 . $[\alpha]_D^{23} -65$ (c,

0.4 in MeOH). λ_{\max} 218 (ϵ 69100); 265 (ϵ 15300); 300 (ϵ 5400) (MeOH) (Berdy). λ_{\max} 222 (ϵ 59100); 275 (ϵ 12800); 310 (ϵ 5200) (MeOH/HCl) (Berdy).


2 α ,2'-*Dihydroxy*: **Stachybotrin D**

[174423-35-9]
C₅₂H₇₀N₂O₁₂ 915.131
Prod. by *Stachybotrys* sp. M6222.
Endothelin antagonist.

Nakamura, M. *et al.*, *J. Antibiot.*, 1995, **48**, 1389-1395; 1396-1400 (*Stachybotrins B-C, isol, pmr, cmr, ir, uv, props*)
Japan. Pat., 1995, 95 330622; *CA*, **124**, 200280 (*Stachybotrin D*)
Roggo, B.E. *et al.*, *J. Antibiot.*, 1996, **49**, 13-19; 374-379 (*Spirodihydrobenzofuranlactam V*)
Deng, W.P. *et al.*, *J.O.C.*, 2003, **68**, 7422-7427 (*synth, struct*)

Stachybotrin B

S-490



C₂₃H₃₁NO₄ 385.502

(+)-form [144385-02-4]

From *Stachybotrys* sp. Exhibits antibacterial and antifungal activity. Sol. EtOAc, CHCl₃, MeOH; poorly sol. H₂O. Mp 178-180° dec. $[\alpha]_D^{25} +39.1$ (c, 0.11 in MeOH). λ_{\max} 220 (ϵ 17000); 254 (ϵ 6400); 302 (ϵ 2900) (MeOH) (Derep). λ_{\max} 254; 270; 302 (MeOH) (Berdy).

N-[2-(4-Hydroxyphenyl)ethyl]: **Stachybotrin C**. NG 245. Antibiotic NG 245

[150351-23-8]
C₃₁H₃₉NO₅ 505.653
Isol. from *Stachybotrys parvispora*.
Neuritogenic agent. Yellowish powder.
Mp 89-92°. $[\alpha]_D^{25} -28.2$ (c, 0.1 in MeOH). λ_{\max} 216 (ϵ 52800); 258 (ϵ 12600); 301 (ϵ 4000) (MeOH).

24-Hydroxy: **Stachybotrin A†**

[144373-26-2]
C₂₃H₃₁NO₅ 401.502
Alkaloid from an aquatic isolate of the fungus *Stachybotrys* sp. Exhibits antibacterial and antifungal activity. Sol. EtOAc, CHCl₃, MeOH; poorly sol. H₂O. $[\alpha]_D^{25} +8.8$ (c, 0.61 in MeOH). λ_{\max} 220 (ϵ 17000); 254 (ϵ 6400); 302 (ϵ 2900) (MeOH) (Derep).

(-)-form

Antibiotic SMTP O. SMTP O

Prod. by *Stachybotrys microspora*.
Amorph. solld. $[\alpha]_D^{25} -7.4$ (c, 0.5 in CHCl₃). λ_{\max} 215 (ϵ 55770); 251 (ϵ 10620); 300 (ϵ 4000) (MeOH).

N-(2-Hydroxyethyl): **Antibiotic SMTP 1**. SMTP 1. NG 243. Antibiotic NG 243

[189819-17-8]
[161128-86-5]
C₂₅H₃₅NO₅ 429.555
Prod. by *Stachybotrys microspora* and *Stachybotrys parvispora*. Light brown oil. $[\alpha]_D^{20} -7.1$ (c, 2 in Me₂CO). λ_{\max} 214 (ϵ 39700); 254 (ϵ 7200); 299 (ϵ 2800) (MeOH).

20,21-Dihydro, 20,21-dihydroxy, N-(2-hydroxyethyl): **Antibiotic SMTP 2**.

SMTP 2
[189819-19-0]
C₂₅H₃₇NO₇ 463.57
Prod. by *Stachybotrys microspora*.
Plasminogen activator. Light brown oil. Sol. CHCl₃, MeOH, Me₂CO, DMSO; poorly sol. H₂O. $[\alpha]_D^{20} -2.7$ (c, 0.2 in Me₂CO). λ_{\max} 212 (ϵ 34000); 252 (ϵ 8000); 294 (ϵ 3000) (MeOH).

N-(4-Carboxybutyl): **Staplabin**

[183006-79-3]
C₂₈H₃₉NO₆ 485.619
Prod. by *Stachybotrys microspora*.
Plasminogen activator. Fibrin binding enhancer. Light brown oil. Sol. MeOH, butanol; poorly sol. EtOAc, hexane. $[\alpha]_D^{20} -11$ (c, 0.1 in CHCl₃). λ_{\max} 216 (ϵ 33500); 258 (ϵ 7300); 300 (ϵ 2500) (MeOH).

(ξ)-form

N-(2-Oxopropyl): **Antibiotic NG 129**.

Antibiotic F 4708. NG 129. F 4708
[157758-74-2]
C₂₆H₃₅NO₅ 441.566
Prod. by *Stachybotrys parvispora*.
Nerve growth factor-like substance.
Powder. Stereochem. not determined.
 λ_{\max} 243 (ϵ 12000) (MeOH) (Berdy).

N-(4-Hydroxy-2-oxobutyl): **Antibiotic NG 242**. NG 242

[160632-28-0]
C₂₇H₃₇NO₆ 471.592
Prod. by *Stachybotrys parvispora*.
Nerve growth factor-like substance.
Powder. Stereochem. not detd.

Xu, X. *et al.*, *J.O.C.*, 1992, **57**, 6700-6703 (*isol, uv, ir, pmr, cmr, ms, struct*)

Japan. Pat., 1994, 94 128 264; 94 256 350; 94 239 869; *CA*, **121**, 177851w; **122**, 29880k; 142492y (NG 129,242,243,245)

Shinohara, C. *et al.*, *J. Antibiot.*, 1996, **49**, 961-966 (*Staplabin*)

Takayasu, R. *et al.*, *FEBS Lett.*, 1997, **418**, 58-62 (*activity*)

Kohyama, T. *et al.*, *J. Antibiot.*, 1997, **50**, 172-174 (*SMTP1, SMTP2*)

Nozawa, Y. *et al.*, *J. Antibiot.*, 1997, **50**, 635-640; 641-645 (*Stachybotrin C*)


Hasumi, K. *et al.*, *J. Antibiot.*, 1998, **51**, 1059-1068 (*activity*)

Hasumi, K. *et al.*, *J. Antibiot.*, 2007, **60**, 463-468 (*SMTP O*)

Stachydrine

S-491

2-Carboxy-1,1-dimethylpyrrolidinium hydroxide inner salt, 9CI. Hygic acid methylbetaine, 8CI. N-Methylproline methylbetaine. Cadabine. Chrysanthemine [32039-73-9] [167936-48-3]



C₇H₁₃NO₂ 143.185

Chrysanthemine was a mixt. of stachydrine with Choline, C-419. Cardiotoxic agent.

(S)-form

L-form

[471-87-4]

Alkaloid from *Capparis tomentosa*, *Chrysanthemum* spp., *Citrus* spp., *Galeopsis grandiflora*, *Lagochilus hirtus*, *Medicago sativa* and *Stachys* spp. (Lamiaceae, Capparidaceae, Asteraceae, Rutaceae, Fabaceae). Also from *Aspergillus oryzae*. Systolic depressant. Stachydrine-contg. spp. e.g. *Capparis* spp. are widely used against rheumatism and many other diseases. Mp 116-118° (monohydrate) Mp 235° dec. (anhyd.). $[\alpha]_D -40.25$ (c, 4 in H₂O). Fairly readily racemised, (±)-form often obt.

Hydrochloride:

Cryst. (EtOH). Mp 222°. $[\alpha]_D^{20} -28.1$ (c, 4.83 in H₂O).

Et ester: Stachydrine ethyl ester

C₉H₁₈NO₃[±] 172.247

Present in roots of *Citrus virgata* (Capparidaceae). $[\alpha]_D -4$ (H₂O). Isol. as periodide.

Et ester, picrate: Mp 101°.

(±)-form [50298-93-6]

Mp 235°.

Hydrochloride: Mp 195-196°.

Picrate:

Cryst. (EtOH). Mp 195-196°.

King, H. *et al.*, *J.C.S.*, 1950, 2866, (*isol, struct*)
Cornforth, J.W. *et al.*, *J.C.S.*, 1952, 597; 601 (*isol, props*)

Marian, L. *et al.*, *Phytochemistry*, 1962, **1**, 209 (*biosynth*)

Paudler, W.W. *et al.*, *Chem. Ind. (London)*, 1963, 1693 (*pmr*)

Mandava, N. *et al.*, *Annalen*, 1970, **741**, 167 (*pmr*)

Musich, J.A. *et al.*, *J.O.C.*, 1977, **42**, 139 (*synth*)


Massiot, G. *et al.*, *Alkaloids (Academic Press)*, 1986, **27**, 310 (*bibl, pharmacol*)

Jones, G.P. *et al.*, *Acta Cryst. C*, 1988, **44**, 1669 (*cryst struct*)

Stachyflin

S-492

[189185-60-2]



Absolute Configuration

C₂₃H₃₁NO₄ 385.502

Prod. by *Stachybotrys* sp. RF-7260.
Active against influenza virus A. Needles. $[\alpha]_D +138.7$ (c, 1 in MeOH). Mp >300°. λ_{\max} 216 (ϵ 42700); 257 (ϵ 6600); 302 (ϵ 3000) (MeOH).

3-Ac: **3-Acetylstachyflin**

[189185-59-9]

C₂₅H₃₃NO₅ 427.539

Prod. by *Stachybotrys* sp. RF-7260.
Active against influenza virus A.

Needles. $[\alpha]_D^{24} +136.4$ (c, 1 in MeOH).
Mp $>300^\circ$. λ_{\max} 215 (ϵ 45700); 256 (ϵ 7000); 301 (ϵ 3200) (MeOH).

N-(1-Carboxy-2-methylpropyl), 3-Ac:
Antibiotic SQ 02-S-VI. SQ 02-S-V1
C₃₀H₄₁NO₇ 527.656

Prod. by *Stachybotrys* sp. RF-7260.
Powder. $[\alpha]_D^{24} +113.5$ (c, 0.34 in MeOH). λ_{\max} 218 (ϵ 38500); 262 (ϵ 9200); 300 (ϵ 2900) (MeOH).

N-(5-Amino-5-carboxypentyl), 3-Ac:
Antibiotic SQ 02-S-L2. SQ 02-S-L2
C₃₁H₄₄N₂O₇ 556.698
Prod. by *Stachybotrys* sp. RF-7260.
Powder. $[\alpha]_D^{24} +96.6$ (c, 0.56 in MeOH).
 λ_{\max} 218 (ϵ 36000); 261 (ϵ 7600); 301 (ϵ 2500) (MeOH).

[189185-71-5]


Taishi, T. *et al.*, *Tet. Lett.*, 1998, **39**, 4347-4350 (synth)

Minagawa, K. *et al.*, *J. Antibiot.*, 2002, **55**, 155-164; 165-171; 239-248 (isol, pmr, cmr, activity)

Staphinine

S-493

[104736-03-0]



Relative Configuration

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from the leaves of *Ervatamia coronaria* (Apocynaceae). Amorph. $[\alpha]_D +25$ (CHCl₃). λ_{\max} 222 (log ϵ 4.58); 275 (log ϵ 3.84); 292 (log ϵ 3.83) (no solvent reported).

11-Hydroxy, 5-ketone: **Ervatinine**

[99877-65-3]

C₁₉H₂₂N₂O₃ 326.394


Alkaloid from the leaves of *Ervatamia coronaria* (Apocynaceae). Amorph. $[\alpha]_D +74$ (CHCl₃). No stereochem. given.

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1985, **24**, 2473; 1986, **25**, 1781-1782 (isol, uv, ir, pmr, cmr, ms, struct)

Staphinine

S-494

13-Methoxystaphimine, 9CI
[59588-19-1]



C₄₂H₅₆N₂O₂ 620.917

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Amorph. $[\alpha]_D^{24} -57.5$ (c, 1.0 in C₆H₆). Extremely sensitive to heat and light in comparison with Staphisine and Staphidine.

Demethoxy: **Staphimine**

[59588-18-0]

C₄₁H₅₄N₂O 590.89


Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Amorph. $[\alpha]_D -58.5$ (c, 1.0 in C₆H₆). Extremely sensitive to heat and light.

Pelletier, S.W. *et al.*, *Tet. Lett.*, 1976, 1055 (uv, ir, pmr, cmr, struct)

Staphisagnine

S-495

13-Methoxystaphisagnine, 9CI
[60723-40-2]



Absolute Configuration

C₄₄H₆₂N₂O₃ 666.985

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Resin. $[\alpha]_D^{25} -104.5$ (c, 2.0 in C₆H₆).

Demethoxy: **Staphisagnine**

[1415-71-0]

C₄₃H₆₀N₂O₂ 636.959


Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Mp 229-231°. $[\alpha]_D^{25} -105.6$ (c, 1.4 in C₆H₆).

Pelletier, S.W. *et al.*, *Tet. Lett.*, 1976, 1749 (isol, ir, pmr, cmr, struct)

Staphisine

S-496

[36576-56-4]



C₄₃H₆₀N₂O₂ 636.959

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Stout needles (Me₂CO). Mp 205° (sinters above 170° and gradually softens to a resin at about 195°). The Staphisine originally isol. (Jacobs, 1941) was shown to be a mixt. of Staphisine and Staphidine.

Methodide:

Thin plates (EtOH). Mp 285-290° dec.

Demethoxy: **Staphidine**

[59588-15-7]

C₄₂H₅₈N₂O 606.933

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Mp 213-216°. $[\alpha]_D^{24} -160$ (c, 2.0 in C₆H₆).

19-Oxo: **Staphigine**

[59588-13-5]

C₄₃H₅₈N₂O₃ 650.943

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Mp 225-227°. $[\alpha]_D^{25} -116$ (c, 2.0 in C₆H₆).

19-Oxo, demethoxy: **Staphirine**

[59588-14-6]

C₄₂H₅₆N₂O₂ 620.917

Alkaloid from *Delphinium staphisagria* seeds (Ranunculaceae). Mp 222-225°. $[\alpha]_D^{25} -126$ (c, 0.5 in C₆H₆).

Jacobs, W.A. *et al.*, *J. Biol. Chem.*, 1941, **141**, 67 (isol, uv)

Craig, L.C. *et al.*, *J. Biol. Chem.*, 1944, **152**, 645 (struct)

Huebner, C.H. *et al.*, *J. Biol. Chem.*, 1947, **169**, 211 (struct)

Pelletier, S.W. *et al.*, *J.A.C.S.*, 1972, **94**, 1754 (uv, ir, pmr, ms, cryst struct)

Pelletier, S.W. *et al.*, *J.O.C.*, 1976, **41**, 3042 (uv, ir, pmr, cmr, struct, derivs)


Pelletier, S.W. *et al.*, *Tet. Lett.*, 1976, 1055 (uv, ir, pmr, cmr, ms, struct, Staphidine)

Pelletier, S.W. *et al.*, *Acta Cryst. B*, 1980, **36**, 3040 (cryst struct)

Staudine

S-497

[67701-63-7]



Relative configuration

C₃₁H₃₃NO₇ 531.604


Alkaloid from *Pachypodanthium staudtii* (Annonaceae). Yellow cryst. Mp 205-206°. Opt. inactive.

Dihydro: Mp 195-196°.

Cavé, A. *et al.*, *J. Nat. Prod.*, 1980, **43**, 103 (isol, uv, ir, ms, pmr, cmr, cryst struct)

Staurosporine**S-498**

AM 2282. CGP 39360. M 193. Antibiotic 230. Antibiotic AM 2282. Antibiotic CGP 39360. Antibiotic M 193 [62996-74-1]



Absolute
Configuration

$C_{28}H_{26}N_4O_3$ 466.538

Alkaloid from *Streptomyces staurosporeus*, *Streptomyces* sp. M 193 and the marine actinomycete strain N96C-47. Isol. from *Eudistoma toعالensis* and its predator *Pseudoceros* sp. Active against fungi and yeasts. Possesses high hypotensive activity and is a potent platelet aggregation inhibitor. Antiparasitic, nematocidal, vasorelaxant, antiarrhythmic, smooth muscle relaxant, neurotropic, antihyperplastic, endothelin agonist and cell cycle progression inhibitor. Pale yellow plates. Sol. Py; fairly sol. $CHCl_3$, MeOH; poorly sol. butanol, C_6H_6 , Me_2CO -EtOAc, hexane. Mp 270° dec. $[\alpha]_D^{20} +35$ (c, 1 in MeOH). λ_{max} 207 (€ 25000); 243 (€ 25000); 267 (sh) (€ 31800); 292 (€ 57200); 322 (sh) (€ 14200); 335 (€ 14600); 356 (€ 10300); 372 (€ 11400) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 6.6 mg/kg (as hydrochloride). KD5084000

4'-N-Formyl: N-Formylstaurosporine

[161973-03-1]

$C_{29}H_{26}N_4O_4$ 494.549

From *Streptomyces longisporoflavus* and the marine *Streptomyces* sp. QD518. Cryst. (EtOAc/ CH_2Cl_2). Mp 221-226°.

4'-N-Ac: [120685-33-8]

$C_{30}H_{28}N_4O_4$ 508.576

Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856. Protein kinase C inhibitor; antihypertensive; diuretic. Yellow powder. λ_{max} 206 (€ 34000); 243 (€ 22000); 292 (€ 49000); 320 (€ 9000); 334 (€ 12000); 355 (€ 8000); 372 (€ 9000) (MeOH).

4'-N-Benzoyl: Midostaurin, INN, USAN. N-Benzoylstaurosporine. CGP 41251. PKC 412

[120685-11-2]

$C_{35}H_{30}N_4O_4$ 570.646

Selective protein kinase C inhibitor. Antineoplastic agent.

4'-N-Carbamoyl: N-Carboxamidostaurosporine. N-Carbamoylstaurosporine

$C_{29}H_{27}N_5O_4$ 509.563

Prod. by the marine *Streptomyces* sp. QD518. Pale yellow solid. λ_{max} 243 (log € 3.67); 292 (log € 4.01); 318 (log € 3.43); 333 (log € 3.43); 354 (log € 3.25); 372 (log € 3.3) (MeOH).

4'-N-Me: N-Methylstaurosporine

[129623-30-9]

$C_{29}H_{28}N_4O_3$ 480.565

Prod. by *Streptomyces longisporoflavus*. Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp. Cryst. (CH_2Cl_2 /2-propanol). Mp 133-137°.

4'-N-De-Me: 4'-N-Demethylstaurosporine

[126221-77-0]

$C_{27}H_{24}N_4O_3$ 452.512

Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp.

O-De-Me: De-O-methylstaurosporine.

CGP 58546. Antibiotic CGP 58546

[161743-35-7]

$C_{27}H_{24}N_4O_3$ 452.512

Prod. by a blocked mutant of *Streptomyces longisporoflavus* and the marine actinomycete strain N96C-47. Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp. Inhibitor of protein kinase C. Pale yellow cryst. (EtOAc). Sol. MeOH, $CHCl_3$. Mp 220° dec. $[\alpha]_D^{20} +82$ (c, 1 in DMSO). λ_{max} 207 (€ 25000); 243 (€ 25000); 267 (sh) (€ 31800); 292 (€ 57200); 322 (sh) (€ 14200); 335 (€ 14600); 356 (€ 10300); 372 (€ 11400) (MeOH) (Derep).

O-De-Me, 4'-N-de-Me: 4'-N,O-Dide-methylstaurosporine

[406703-32-0]

$C_{26}H_{22}N_4O_3$ 438.485

Isol. from the flatworm *Pseudoceros* sp. Yellowish amorph. powder. λ_{max} 206 (€ 2190); 291 (€ 3440); 322 (€ 690); 334 (€ 760); 355 (€ 510); 372 (€ 550) (MeOH).

4'-N-Hydroxy: [161973-04-2]

[137888-67-6]

$C_{28}H_{26}N_4O_4$ 482.538

Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856.

4'-N-Hydroxy, 4'-N-de-Me, 4'-N-formyl:

[161927-20-4]

$C_{28}H_{24}N_4O_5$ 496.521

From *Streptomyces longisporoflavus*. Cryst. (MeCN aq.). Mp 220° dec.

4'-N-Hydroxy, O-de-Me: [137888-68-7]

$C_{27}H_{24}N_4O_4$ 468.511

Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856.

4'-N-(Acetoxymethoxy): 4'-N-(Acetoxy-methoxy)staurosporine

[161927-21-5]

$C_{31}H_{30}N_4O_6$ 554.601

From *Streptomyces longisporoflavus*. Cryst. (CH_2Cl_2 /propanol). Mp 140-143°.

4'-De(methylamino), 4'α-nitro: 4'-De-methylamino-4'-nitrostaurosporine

[161927-19-1]

$C_{27}H_{22}N_4O_5$ 482.495

From *Streptomyces longisporoflavus*. Enzyme inhibitor. Powder (CH_2Cl_2 /2-propanol). Mp 237-242°. λ_{max} 233 (€ 29400); 244 (sh) (€ 28000); 263 (sh) (€ 31300); 275 (sh) (€ 42000); 289 (€ 71000); 319 (sh) (€ 13400); 333 (€ 17700); 352 (€ 12100); 369 (€ 13400) (MeOH) (Derep). λ_{max} 209 (€ 23000); 230 (€ 7800); 263; 295 (€ 16000); 336; 364 (EtOH) (Berdy).

4'-De(methylamino), 4'-hydroxy: 4'-De-methylamino-4'-hydroxystaurosporine.

Antibiotic RK 286C. RK 286C

[126572-73-4]

$C_{27}H_{23}N_3O_4$ 453.496

Prod. by *Streptomyces* sp. RK 286C. Protein kinase C inhibitor; platelet aggregation inhibitor; antiinflammatory; cell cycle progression inhibitor; bleb formn. inhibitor. Pale yellow powder. Sol. EtOAc, DMSO; fairly sol. MeOH; poorly sol. H_2O . Mp 265° dec. $[\alpha]_D^{20} +45.3$ (c, 0.22 in EtOAc). λ_{max} 207 (€ 25000); 243 (€ 25000); 267 (sh) (€ 31800); 292 (€ 57200); 322 (sh) (€ 14200); 335 (€ 14600); 356 (€ 10300); 372 (€ 11400) (MeOH) (Derep). λ_{max} 237 (€ 30450); 245 (€ 30120); 292 (€ 69500); 335 (€ 16550); 356 (€ 13240); 372 (€ 14890) (MeOH) (Berdy). λ_{max} 237; 245; 292; 335; 356; 372 (MeOH/HCl) (Berdy). λ_{max} 237; 245; 292; 335; 356; 372 (MeOH/NaOH) (Berdy).

4'-De(methylamino), 4'α,5'α-dihydroxy: 4'-Demethylamino-4',5'-dihydroxystaurosporine. Antibiotic MLR 52. MLR 52

[155416-34-5]

$C_{27}H_{23}N_3O_5$ 469.496

Prod. by a *Streptomyces* sp. Inhibitor of protein kinase C. Sol. MeOH. Mp 263-268°. $[\alpha]_D +68$ (c, 0.09 in MeOH). λ_{max} 206 (€ 48500); 234 (€ 48500); 286 (€ 72100); 317 (€ 25000); 332 (€ 30000); 351 (€ 21500); 368 (€ 22600) (MeOH) (Derep).

4'-De(methylamino), 4',7-dihydroxy:

[137888-70-1]

$C_{27}H_{23}N_3O_5$ 469.496

Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856. Enzyme inhibitor. λ_{max} 207; 239; 292; 300; 358; 375 (MeOH) (Berdy).

4'-De(methylamino), 4',7-dihydroxy, O-de-Me: [137888-71-2]

$C_{26}H_{21}N_3O_5$ 455.469

Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856. Enzyme inhibitor. λ_{max} 208 (€ 24000); 239 (€ 15000); 301 (€ 27000); 358 (€ 4000); 376 (€ 4000) (MeOH) (Berdy).

7-Oxo: 7-Oxostaurosporine. BMY 41950.

RK 1409. Antibiotic BMY 41950.

Antibiotic RK 1409

[125035-83-8]

$C_{28}H_{24}N_4O_4$ 480.522

Prod. by *Streptomyces platensis* ssp. *malvinus* and *Streptomyces staurosporeus* ATCC55006. Inhibitor of protein kinase C. Yellow powder. Mp 235° dec. $[\alpha]_D^{20} +38.3$ (c, 0.06 in $CHCl_3$). λ_{max} 208 (€ 19600); 238 (€ 26400); 260 (€ 13600); 287 (€ 16800); 305 (sh) (€ 20800); 317 (€ 33600); 340 (sh) (€ 8800); 410 (€ 2400) (MeOH) (Derep).

3-Hydroxy: 3-Hydroxystaurosporine

[320384-51-8]

$C_{28}H_{26}N_4O_4$ 482.538

Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp.


3-Hydroxy, 4'-N-Me: 3-Hydroxy-4'-N-methylstaurosporine

- [399031-46-0]
C₂₉H₂₈N₄O₄ 496.565
Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp.
- 3-*Hydroxy*, 4'-*N*-*de*-*Me*: **3-Hydroxy-4'-N-demethylstauroporine**
[406703-29-5]
C₂₇H₂₄N₄O₄ 468.511
Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp. Yellowish amorph. powder. λ_{max} 204 (ε 2710); 287 (ε 3100); 297 (ε 4080); 342 (ε 1150); 366 (ε 700); 384 (ε 700) (MeOH).
- 3-*Hydroxy*, *O*-*de*-*Me*: **3-Hydroxy-O-demethylstauroporine**
[399031-45-9]
C₂₇H₂₄N₄O₄ 468.511
Isol. from *Eudistoma toعالensis* and *Pseudoceros* sp. Amorph. yellow powder. λ_{max} 237 (ε 8350); 251 (ε 6600); 270 (ε 7120); 297 (ε 13380); 342 (ε 3790); 366 (ε 2400); 384 (ε 2420) (MeOH).
- 5'α-*Hydroxy*: **5'-Hydroxystauroporine**
[308847-74-7]
C₂₈H₂₆N₄O₄ 482.538
Prod. by a marine *Micromonospora* sp. Pale yellow powder. Mp > 220° dec. [α]_D²⁵ +53 (c, 0.1 in MeOH). λ_{max} 206; 242; 291; 320; 334; 354; 370 (MeOH).
- 5'α-*Hydroxy*, *N*-*Me*: **5'-Hydroxy-4'-N-methylstauroporine**
[308847-73-6]
C₂₉H₂₈N₄O₄ 496.565
Prod. by a marine *Micromonospora* sp. Pale yellow powder. Mp > 220° dec. [α]_D²⁵ +30 (c, 0.11 in CHCl₃). λ_{max} 206; 242; 291; 320; 334; 354; 370 (MeOH).
- 7ξ-*Hydroxy*: **7-Hydroxystauroporine**. UCN 01. Antibiotic UCN 01. NSC 638850
[112953-11-4]
C₂₈H₂₆N₄O₄ 482.538
From *Streptomyces* sp. N-126 and *Streptomyces longisporoflavus* R19. Selective inhibitor of protein kinase C. Cytotoxic. Pale yellow needles. Sol. MeOH, CHCl₃, DMSO; poorly sol. H₂O, hexane. Mp 245-250° dec. [α]_D²² +132 (c, 0.3 in MeOH). λ_{max} 240 (ε 29000); 264 (sh) (ε 20000); 274 (sh) (ε 21000); 300 (ε 55000); 326 (sh) (ε 9600); 338 (sh) (ε 8300); 358 (ε 7500); 374 (ε 8500) (MeOH) (Derep). λ_{max} 241; 266; 280; 298; 325; 356; 375 (EtOH) (Berdy).
- LD₅₀ (mus, ipr) 30 mg/kg. KC6600010
- 11-*Hydroxy*: **11-Hydroxystauroporine**
[143682-17-1]
C₂₈H₂₆N₄O₄ 482.538
Isol. from the marine tunicate *Eudistoma* sp. Cytotoxic. Inhibitor of protein kinase C. Amorph. [α]_D +10.3 (c, 0.3 in MeOH).
- 11-*Hydroxy*, 4'-*N*-*de*-*Me*: **11-Hydroxy-4'-N-demethylstauroporine**
[272766-45-7]
C₂₇H₂₄N₄O₄ 468.511
Isol. from *Eudistoma toعالensis*, *Pseudoceros* sp., and from the mollusc *Coriocola nigra*. Amorph. yellow powder. λ_{max} 290 (ε 1040); 356 (ε 180); 373 (ε 180) (MeOH).
- 3,11-*Dihydroxy*: **3,11-Dihydroxystauroporine**
[143682-18-2]
C₂₈H₂₆N₄O₅ 498.537
Isol. from *Eudistoma* sp. and the mollusc *Coriocola nigra*. Cytotoxic. Amorph.
- 7ξ-*Methoxy*, 4'-*N*-*hydroxy*: [137888-69-8]
C₂₉H₂₈N₄O₅ 512.564
Prod. by *Saccharothrix aerocolonigenes copiosa* ATCC53856.
- 10-*Methoxy*: **10-Methoxystauroporine**. Antibiotic TAN 999. TAN 999
[124843-68-1]
C₂₉H₂₈N₄O₄ 496.565
From *Nocardioopsis dassonvillei*. Possesses macrophage-activating props. Pale yellow cryst. + ½H₂O. Sol. MeOH; poorly sol. H₂O. Mp 221° dec. [α]_D²⁴ +42 (c, 0.5 in DMF). λ_{max} 245 (ε 31200); 296 (ε 60300); 341 (ε 19500); 352 (sh) (ε 16200); 368 (ε 11000) (MeOH) (Derep).
- 3'-*Epimer*, 4'-*de* (methylamino), 4'-*hydroxy*: **4'-Demethylamino-4'-hydroxy-3'-epistauroporine**. Antibiotic RK 1409B. RK 1409B
[145212-39-1]
C₂₇H₂₃N₃O₄ 453.496
Prod. by *Streptomyces platensis* ssp. *malvinus*. Inhibitor of protein kinase C; antiinflammatory; cell cycle progression inhibitor. Pale yellow powder. Sol. DMSO; fairly sol. MeOH, EtOAc, CHCl₃, Me₂CO; poorly sol. H₂O, hexane. Mp 260° dec. [α]_D²² +147 (c, 0.2 in DMSO). λ_{max} 203 (ε 37870); 245 (ε 25370); 293 (ε 58710); 336 (ε 13140); 357 (ε 9970); 374 (ε 10870) (MeOH) (Berdy).
- 7-*Epimer*, 7-*hydroxy*: **Antibiotic UCN 02**. UCN 02
[121569-61-7]
C₂₈H₂₆N₄O₄ 482.538
From *Streptomyces* sp. N-126. Cytotoxic, protein kinase C inhibitor. Pale yellow needles. Sol. CHCl₃, DMSO, MeOH; poorly sol. H₂O. Mp 245-250° dec. [α]_D²⁵ -38.6 (c, 0.35 in MeOH). λ_{max} 240 (ε 29000); 264 (sh) (ε 20000); 274 (sh) (ε 21000); 300 (ε 55000); 326 (sh) (ε 9600); 338 (sh) (ε 8300); 358 (ε 7500); 374 (ε 8500) (MeOH) (Derep).
- Omura, S. et al., *J. Antibiot.*, 1977, **30**, 275-282 (isol, ms, uv, ir, pmr)
Cordell, G.A. et al., *Alkaloids (Academic Press)*, 1981, **20**, 4-5 (rev, tox)
Furusaki, A. et al., *Bull. Chem. Soc. Jpn.*, 1982, **55**, 3681-3685 (cryst struct)
Weinreb, S.M. et al., *Heterocycles*, 1984, **21**, 309-324 (synth)
Oka, S. et al., *Agric. Biol. Chem.*, 1986, **50**, 2723-2727 (isol, props)
Eur. Pat., 1988, ((*Ciba-Geigy*))296 110; CA, **111**, 96977h (synth, pharmacol)
Meyer, T. et al., *Int. J. Cancer*, 1989, **43**, 851-856 (*Midostaurin*, pharmacol)
Takahashi, I. et al., *J. Antibiot.*, 1989, **42**, 564-570; 571-576 (UCN01, 02)
Tanida, S. et al., *J. Antibiot.*, 1989, **42**, 1619-1630 (TAN 999)
Osada, H. et al., *J. Antibiot.*, 1990, **43**, 168-173 (RK 286C)
- Pat. Coop. Treaty (WIPO)*, 1991, 9 034; CA, **116**, 39768v (*Saccharothrix aerocolonigenes copiosa* constits)
Tsuboi, S. et al., *Tetrahedron*, 1991, **47**, 3565-3574 (TAN 999, cmr)
Krause, K.H. et al., *Eur. J. Pharmacol., Mol. Pharmacol.*, 1992, **227**, 221-224 (*Midostaurin*, pharmacol)
Osada, H. et al., *J. Antibiot.*, 1992, **45**, 189-194; 195-198 (7-*Oxostauroporine*)
Koshino, H. et al., *J. Antibiot.*, 1992, **45**, 1428-1432 (RK 1409B)
Kinnel, R.B. et al., *J.O.C.*, 1992, **57**, 6327-6329 (11-*Hydroxystauroporine*, 3,11-*Dihydroxystauroporine*)
Caravatti, G. et al., *Bioorg. Med. Chem. Lett.*, 1994, **4**, 399-404 (*Midostaurin*, synth, pharmacol)
McAlpine, J.B. et al., *J. Antibiot.*, 1994, **47**, 281-288 (MLR 52)
Funato, N. et al., *Tet. Lett.*, 1994, **35**, 1251-1254 (cryst struct, abs config)
Sedlak, J. et al., *Anti-Cancer Drugs*, 1995, **6**, 70-76 (*Midostaurin*, pharmacol)
Cai, Y. et al., *J. Antibiot.*, 1995, **48**, 143-148 (*Demethylaminonitrostauroporine*, *Acetoxymethoxystauroporine*, *N-Formylstauroporine*)
Hoehn, P. et al., *J. Antibiot.*, 1995, **48**, 300-305 (*De-O-methylstauroporine*)
Omura, S. et al., *J. Antibiot.*, 1995, **48**, 535-548 (rev)
Link, J.T. et al., *J.A.C.S.*, 1995, **117**, 552-553 (synth)
Begemann, M. et al., *Clin. Cancer Res.*, 1996, **2**, 1017-1030 (*Midostaurin*, pharmacol)
Cai, Y. et al., *J. Antibiot.*, 1996, **49**, 1060-1062 (*N-Methylstauroporine*)
Yang, S.W. et al., *J. Nat. Prod.*, 1996, **59**, 828-833; 1997, **60**, 236-241 (biosynth, ms)
Wood, J.L. et al., *J.A.C.S.*, 1996, **118**, 10656-10657; 1997, **119**, 9652-9661 (synth)
Wood, J.L. et al., *Tet. Lett.*, 1996, **37**, 7335-7338 (synth, RK 286C)
Beltran, P.J. et al., *Biochem. Pharmacol.*, 1997, **53**, 245-247 (*Midostaurin*, pharmacol)
Weidner, S. et al., *J. Antibiot.*, 1998, **51**, 679-682 (biosynth)
Schupp, P. et al., *J. Nat. Prod.*, 1999, **62**, 959-962 (isol, pmr, cmr)
Contrell, C.L. et al., *Nat. Prod. Lett.*, 1999, **14**, 39-46 (*Coriocola nigra* constits)
Williams, D.E. et al., *Tet. Lett.*, 1999, **40**, 7171-7174 (marine strain N96C-47, isol)
Hernandez, L.M.C. et al., *J. Antibiot.*, 2000, **53**, 895-902 (5'-*Hydroxystauroporine*, 5'-*Hydroxy-N-methylstauroporine*)
Cantrell, C.L. et al., *Nat. Prod. Lett.*, 2000, **14**, 39-46 (11-*Hydroxy-4'-N-demethylstauroporine*)
Schupp, P. et al., *Cancer Lett. (Shannon, Irel.)*, 2001, **174**, 165-172; 2003, **190**, 119 (*Eudistoma*, *Pseudoceros*, isol)
Zaugg, K. et al., *Cancer Res.*, 2001, **61**, 732-738 (*Midostaurin*, pharmacol)
Schupp, P. et al., *J. Nat. Prod.*, 2002, **65**, 295-298 (*N,O*-*Didemethylstauroporine*, 3-*Hydroxy-N-demethylstauroporine*)
Wu, S.J. et al., *J. Antibiot.*, 2006, **59**, 331-337 (marine strain QD518, isol)

Stawamycin

[167568-90-3 (Na salt)]

S-499




C₂₉H₃₉NO₅ 481.631

Prod. by *Streptomyces* sp. Inhibitor of Epstein-Barr virus transcription factor. Amorph. powder (as Na salt). Sol. MeOH; poorly sol. H₂O. [α]_D²⁵ +179 (c, 0.3 in MeOH) (Na salt). Related to Indanomycin, I-64. λ_{max} 206 (ε 13000); 244 (ε 12000); 289 (ε 25000) (MeOH).

Miao, S. *et al.*, *Tet. Lett.*, 1995, **36**, 5699-5702 (isol, uv, pmr, cmr)

Stealthin A S-500

11-Amino-4,5,9-trihydroxy-2-(hydroxy-methyl)-10H-benzo[b]fluoren-10-one, 9CI. Antibiotic CA 39A. CA 39A [146474-93-3]

C₁₈H₁₃NO₅ 323.304

Prod. by *Streptomyces viridochromogenes*. Free radical scavenger. Haemolysis inhibitor. Antioxidant. Red powder. Unstable in light. λ_{max} 274 (ε 37600); 470 (sh) (ε 7500); 504 (ε 11600); 538 (ε 13900) (MeOH). λ_{max} 269 (ε 36400); 275 (ε 37600); 330 (ε 7000); 430 (ε 5800); 505 (ε 8700); 538 (ε 11000) (MeOH/HCl) (Berdy). λ_{max} 268 (ε 29500); 275 (ε 30100); 357 (ε 11600); 583 (ε 12100) (MeOH/NaOH) (Berdy).

12-Aldehyde: **Stealthin B**. Antibiotic CA 39B. CA 39B [146474-94-4]

C₁₈H₁₁NO₅ 321.289


Prod. by *Streptomyces viridochromogenes*. Free radical scavenger. Haemolysis inhibitor. Antioxidant. Purple powder. Unstable in light. λ_{max} 286 (ε 21900); 315 (ε 19200); 415 (ε 6300); 550 (sh) (ε 11600); 585 (ε 13900) (MeOH). λ_{max} 268 (ε 34000); 275 (ε 35300); 330 (ε 5500); 345 (ε 8100); 415 (ε 4900); 437 (ε 6700); 503 (ε 9800); 535 (ε 12500) (MeOH/HCl) (Berdy).

Shin-ya, K. *et al.*, *Tet. Lett.*, 1992, **33**, 7025-7028 (isol, uv, ir, pmr, cmr)

Koyama, H. *et al.*, *J.C.S. Perkin 1*, 1998, 203-210 (synth)

Stealthin C S-501

11-Amino-4,5,9-trihydroxy-2-methyl-10H-benzo[b]fluoren-10-one, 9CI [185253-25-2]

C₁₈H₁₃NO₄ 307.305

Prod. by *Streptomyces murayamaensis*. Biosynthetic intermed. of Kinamycin D. λ_{max} 214 ; 276 ; 340 ; 506 ; 540 (MeCN/AcOH aq.).


Gould, S.J. *et al.*, *J.O.C.*, 1997, **62**, 320-324 (isol, uv, ir, struct)

Koyama, H. *et al.*, *J.C.S. Perkin 1*, 1998, 203-210 (synth)

Stebisimine S-502

N-Nor-1,2-dehydroepistephanine

[5692-04-6]

C₃₆H₃₄N₂O₆ 590.674

Alkaloid from *Stephania japonica*, *Anisocycla grandidieri* and *Triclisia gillettii* (Menispermaceae). Prisms (MeOH). Mp 233-235° (227-229°). [α]_D 0 (c, 1.26 in CHCl₃). λ_{max} 238 (ε 51900); 279 (ε 24200); 308 (ε 12500) (EtOH) (Berdy).

Hydrochloride (1:2): Mp 290° dec.

Dipicrate: Mp 254-256°.

Methiodide (1:2): Mp 290° (gradual dec.).

7'-O-De-Me: 7'-O-Demethylstebisimine [916480-88-1]

C₃₅H₃₂N₂O₆ 576.648

Alkaloid from the roots of *Cocculus laurifolius*. Amorph. solid.

12-O-De-Me: **Puertogaline B**C₃₅H₃₂N₂O₆ 576.648

Alkaloid from the stem bark of *Guatteria boliviana*. Amorph. λ_{max} 205 (log ε 4.64); 235 (log ε 4.5); 278 (log ε 4.17) (EtOH).

7',12-Di-O-de-Me: **Puertogaline A**C₃₄H₃₀N₂O₆ 562.621

Alkaloid from the stem bark of *Guatteria boliviana*. Amorph. λ_{max} 204 (log ε 4.64); 235 (sh) (log ε 4.45); 278 (log ε 4.09) (EtOH).

1',2'-Dihydro: see Coclobine, C-551

α,α'-Dioxo, 7'-O-de-Me: 7'-O-Demethyl-α,α'-dioxostebisimine [916480-87-0]

C₃₅H₂₈N₂O₈ 604.615

Alkaloid from the roots of *Cocculus laurifolius*. Amorph. solid.

Barton, D.H.R. *et al.*, *J.C.S. (C)*, 1966, 2313

(isol, ms, ir, uv, pmr, struct)

Kametani, T. *et al.*, *Chem. Comm.*, 1967, 1212 (synth, uv, ir, pmr)

Schlittler, E. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 2061 (isol)

Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592 (ms)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 1 (isol, uv, ir, pmr, ms)

Owusu, P.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 61 (isol)

Matsui, M. *et al.*, *J. Nat. Prod.*, 1982, **45**, 497 (isol, uv, ir, pmr, ms)


Nakova, E. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 83-87; 88-91 (synth)

Mahiou, V. *et al.*, *Phytochemistry*, 2000, **54**, 709-716 (*Puertogalines A, B*)

Zhang, H. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 553-557 (7'-Demethylstebisimines)

Stelletadine A

S-503



Absolute Configuration

C₂₀H₃₆N₆O 376.544**(R)-form** [179732-83-3]

Alkaloid from the marine sponge *Stelletta* sp. Induces larval metamorphosis in ascidians, RNA cleaving agent. [α]_D²⁴ -42.7 (c, 0.09 in MeOH). [α]_D²⁴ -32.8 (c, 1.00 in MeOH). λ_{max} 285 (ε 11100) (MeOH).

Tsukamoto, S. *et al.*, *Tet. Lett.*, 1996, **37**, 5555-5556 (isol, uv, ir, pmr, cmr, struct)


Nozawa, D. *et al.*, *Bioorg. Med. Chem. Lett.*, 2001, **11**, 1481-1483 (synth, abs config)

Takikawa, H. *et al.*, *J.C.S. Perkin 1*, 2001, 657-661 (synth, abs config)

Stelletamide A

S-504

Octahydro-4-methyl-1-[[[4,8,12-trimethyl-1-oxo-2,7,11-tridecatrieny]amino]methyl]indolizinium(1+), 9CI [129744-24-7] [129744-25-8]

C₂₆H₄₅N₂O[⊕] 401.654

Alkaloid from the marine sponge *Stelletta* sp. Antifungal agent. Cytotoxic. Semisolid (as phosphate). [α]_D²⁵ +23.1 (c, 0.3 in EtOH) (phosphate).

Δ³-Isomer: **Stelletamide C**C₂₆H₄₅N₂O[⊕] 401.654

Alkaloid from *Stelletta* sp. Antibacterial agent. Amorph. solid. Counterion not specified.

Hirota, H. *et al.*, *Tet. Lett.*, 1990, **31**, 4163-4164 (isol, ir, pmr, cmr)

Whitlock, G.A. *et al.*, *J.O.C.*, 1997, **62**, 7916-7917 (synth, enantiomer)

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1202-1204 (*Stelletamide C*)


Whitlock, G.A. *et al.*, *Helv. Chim. Acta*, 2000, **83**, 2007-2022 (synth)

Yamazaki, N. *et al.*, *Heterocycles*, 2008, **75**, 285-290 (synth)

Stelletamide B

S-505

[189580-08-3]


C₂₄H₄₁N₂O[⊕] 373.601

Alkaloid from the sponge *Stelletta* sp. Antifungal agent. Yellow gum (as chloride). $[\alpha]_D^{25}$ -24.2 (c, 0.5 in CHCl_3) (chloride). CAS no. refers to chloride.

Shin, J. *et al.*, *J. Nat. Prod.*, 1997, **60**, 611-613 (*isol, uv, ir, pmr, cmr*)

Yamazaki, N. *et al.*, *Org. Lett.*, 2001, **3**, 193-196 (*synth*)

Stelletamine S-506
[139427-07-9]




$\text{C}_{20}\text{H}_{14}\text{N}_4\text{S}$ 342.423

Alkaloid from the marine sponge *Stelletta* sp. Dark brown cryst. (CHCl_3). Mp 280-282°. λ_{max} 205 (ϵ 15700); 260 (ϵ 9400); 309 (ϵ 10800); 364 (ϵ 3760); 391 (ϵ 3420); 460 (ϵ 1370) (MeOH) (Derep).

Gunawardana, G.P. *et al.*, *J.O.C.*, 1992, **57**, 1523 (*isol, ir, uv, pmr, cmr, cryst struct*)

Stelletazole A S-507




$\text{C}_{26}\text{H}_{46}\text{N}_6\text{O}^{2+}$ 458.689

Alkaloid from a marine sponge, *Stelletta* sp. Antibacterial agent. Also inhibits Ca/calmodulin-dependent phosphodiesterase. $[\alpha]_D^{24}$ -35 (c, 0.08 in MeOH). Counterion not specified.

Tsakamoto, S. *et al.*, *Tet. Lett.*, 1999, **40**, 737-738 (*isol, pmr, cmr*)

Stelletazole B S-508
[247086-88-0]



$\text{C}_{30}\text{H}_{51}\text{N}_6\text{O}^+$ 511.773

The +ve charge is delocalised over the imidazole N's. Alkaloid from the sponge *Stelletta* sp. Antibacterial agent.

Amorph. solid. Counterion not specified. Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1202-1204 (*isol, pmr, cmr*)

Stelletazole C S-509

[247086-89-1]
As Stelletazole B, S-508 with
R = H


$\text{C}_{29}\text{H}_{49}\text{N}_4\text{O}^+$ 469.732

Alkaloid from the sponge *Stelletta* sp. Antibacterial agent. Amorph. solid. Counterion not specified.

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1202-1204 (*isol, pmr, cmr*)

Stemmadenine S-510

Methyl 2,7,19,20-tetrahydro-17-hydroxy-3,7-secocuran-16-carboxylate, 9CI
[10012-73-4]



Absolute Configuration

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3$ 354.448

Alkaloid from *Stemmadenia donnell-smithii*, *Pandaca minutiflora* leaves, *Diplorhynchus condylocarpon*, *Craspidospermum verticillatum*, *Rhazya stricta* and *Tabernaemontana dichotoma* (Apocynaceae). Shows hypotensive and weak muscle relaxant activity. Active against gram-positive bacteria and fungi. Mp 199-200° (189-191°) dec. $[\alpha]_D^{25}$ +324 (Py).

► W1810000

Hydrochloride: Mp 215-216° dec.

15-Epimer: **15-Epistemmadenine**. 15 β -Stemmadenine

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3$ 354.448

Alkaloid from the fruit of *Tabernaemontana heyneana*. Yellow powder. Mp 160°. $[\alpha]_D^{25}$ +247 (c, 0.09 in EtOH). λ_{max} 220 (log ϵ 1.01); 280 (log ϵ 0.24) (MeOH).

De(hydroxymethyl), stereoisomer: **Desformylstemmadenine**. Deformylstemmadenine

[55785-32-5]

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_2$ 324.422

Alkaloid from *Craspidospermum verticillatum* var. *petiolare* (Apocynaceae). Cryst. (Me_2CO). Mp 218°. $[\alpha]_D^{20}$ +112 (c, 0.5 in Py). The trivial name is misleading. Stereochem. not detd.

Walls, F. *et al.*, *Tetrahedron*, 1958, **2**, 173 (*isol, uv, ir*)

Stauffer, D. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 2006 (*isol, ir, uv*)

Sandoval, A. *et al.*, *Tet. Lett.*, 1962, 409, (*pmr*)

Kan-Fan, C. *et al.*, *Bull. Soc. Chim. Fr.*, 1974, **12**, 2839 (*isol, ms, deriv*)

Kutney, J.P. *et al.*, *Heterocycles*, 1975, **3**, 197 (*abs config, pmr*)

Petitfrere, N. *et al.*, *Phytochemistry*, 1975, **14**, 1648 (*isol*)

Ohiri, F.C. *et al.*, *Planta Med.*, 1983, **49**, 17 (*isol*)

Perera, P. *et al.*, *J. Ethnopharmacol.*, 1985, **13**, 165 (*pharmacol*)


Mariee, N.K. *et al.*, *J. Nat. Prod.*, 1988, **51**, 186-187 (*isol, activity*)

Achenbach, H. *et al.*, *Phytochemistry*, 1997, **45**, 325 (*cmr*)

Grover, R.K. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 474-476 (15-Epistemmadenine)

Stemoamide S-511

[142905-25-7]



Absolute Configuration

$\text{C}_{12}\text{H}_{17}\text{NO}_3$ 223.271

Alkaloid from roots of *Stemona tuberosa* (Stemonaceae). Insecticidal agent. Amorph. (natural); Cryst. (EtOAc) (synthetic). Mp 190-191° dec Mp 187-188° (synthetic). $[\alpha]_D^{21.6}$ -28.1 (c, 0.125 in MeOH) (natural). $[\alpha]_D^{30}$ -219.3 (c, 0.5 in MeOH) (synthetic) (-181).

Lin, W.-H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 571-576 (*isol, pmr, cmr, ms*)

Williams, D.R. *et al.*, *Tet. Lett.*, 1994, **35**, 6417 (*synth, abs config*)

Kohno, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 1996, **69**, 2063 (*synth*)


Kinoshita, A. *et al.*, *Heterocycles*, 1997, **46**, 287-300 (*synth*)

Jacobi, P.A. *et al.*, *J.A.C.S.*, 2000, **122**, 4295-4303 (*synth*)

Olivo, H.F. *et al.*, *J.O.C.*, 2006, **71**, 3287-3290 (*synth*)

Torrsell, S. *et al.*, *J.O.C.*, 2007, **72**, 4246-4249 (*synth*)

Stemoburkilline S-512



Absolute Configuration


$\text{C}_{22}\text{H}_{31}\text{NO}_5$ 389.491

Alkaloid from the roots of *Stemona burkillii*. Yellow-brown gum. $[\alpha]_D^{26}$ +37.5 (c, 0.28 in CHCl_3).

Mungkornasawakul, P. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1740-1743 (*isol, pmr, cmr*)

Stemochinamine S-513

[1003567-55-2]




Absolute Configuration

$\text{C}_{16}\text{H}_{28}\text{N}_2\text{O}_3$ 296.409

Alkaloid from the roots of *Stemona cochinchinensis*. Needles (hexane/ Me_2CO). Racemic.

Lin, L.-G. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 2167-2175 (*isol, pmr, cmr, ms*)

Stemocurtisine S-514
Pyridostemine



Relative Configuration

C₁₉H₂₅NO₅ 347.41

Alkaloid from the roots of *Stemona curtisii* and an unidentified *Stemona* sp.²⁵ Prisms (EtOAc/Et₂O). Mp 149–151°. [α]_D²⁵ +334 (c, 0.66 in CHCl₃). [α]_D²⁰ +473 (c, 0.4 in MeOH).

Mungkornasawakul, P. et al., *J. Nat. Prod.*, 2003, **66**, 980–982 (isol, pmr, cmr, cryst struct)


Kaltenegger, E. et al., *Phytochemistry*, 2003, **63**, 803–816 (isol, pmr, cmr)

Pyne, S.G. et al., *Maejo Int. J. Sci. Technol.*, 2007, **1**, 157–165 (rev, biosynth)

Stemofoline

S-515

[29881-57-0]



C₂₂H₂₉NO₅ 387.475

Alkaloid from the stems and leaves of *Stemona japonica* (Stemonaceae). Has insecticidal props. Mp 87–89°. [α]_D +273 (MeOH). λ_{max} 295 (MeOH).

Hydrobromide:

Cryst. + 1H₂O. Mp 224° dec.

11β,12α-Dihydro: **11,12-Dihydrostemofoline**

C₂₂H₃₁NO₅ 389.491

Alkaloid from the roots of *Stemona burkillii*. Yellow-brown gum. [α]_D²⁶ +38.9 (c, 0.35 in CHCl₃).

1',2'-Didehydro(E-): **1',2'-Didehydrostemofoline. Asparagamine A** [156798-15-1]

C₂₂H₂₇NO₅ 385.459

Alkaloid from the roots of *Stemona collinsae* and *Asparagus racemosus*. Insecticide and antitumour agent. Cryst. (MeOH). Mp 180° (172–174°). [α]_D¹⁸ +230 (c, 0.74 in MeOH) (+202.5). Original source of *A. racemosus* now suspected of being *S. collinsae* (2004). λ_{max} 294 (log ε 4.41) (MeOH).

1',2'-Didehydro(E-), N-oxide: **1',2'-Didehydrostemofoline N-oxide**

C₂₂H₂₇NO₆ 401.458

Alkaloid from an unidentified *Stemona* sp. Yellow-brown gum. [α]_D²⁵ +99 (c, 0.28 in CHCl₃).

2'R-Hydroxy: **2'R-Hydroxystemofoline**

C₂₂H₂₉NO₆ 403.474

Alkaloid from an unidentified *Stemona* sp. Yellow-brown gum. [α]_D²² +168 (c, 0.29 in CHCl₃).

2'S-Hydroxy: **2'S-Hydroxystemofoline**

C₂₂H₂₉NO₆ 403.474

Alkaloid from *Stemona collinsae*. [α]_D²¹ +249 (c, 0.33 in CHCl₃).

3'R-Hydroxy, 1',2'-didehydro: **(3'R)-Stemofolenol**

C₂₂H₂₇NO₆ 401.458

Alkaloid from an unidentified *Stemona* sp.

3'S-Hydroxy: **3'S-Hydroxystemofoline**

[947697-38-3]

C₂₂H₂₉NO₆ 403.474

Alkaloid from a *Stemona* sp.

3'S-Hydroxy, 1',2'-didehydro: **(3'S)-Stemofolenol**

C₂₂H₂₇NO₆ 401.458

Alkaloid from an unidentified *Stemona* sp.

4'-Hydroxy: **4'-Hydroxystemofoline.**

Oxystemofoline

[139750-70-2]

C₂₂H₂₉NO₆ 403.474

Minor alkaloid from roots of *Stemona parviflora*. Drawing error in Chem. Abs.

6S-Hydroxy: **6S-Hydroxystemofoline**

C₂₂H₂₉NO₆ 403.474

Alkaloid from *Stemona japonica*.

Amorph. yellow powder. [α]_D +490 (c, 0.05 in MeOH). λ_{max} 295 (log ε 4.7) (MeOH).

16-Hydroxy: **16-Hydroxystemofoline**

C₂₂H₂₉NO₆ 403.474

Alkaloid from *Stemona japonica*.

Amorph. yellow powder. [α]_D²⁰ +286.7 (c, 0.14 in MeOH). λ_{max} 296 (log ε 4.39) (MeOH).

4'-Methoxy: **4'-Methoxystemofoline**

[139768-07-3]

C₂₃H₃₁NO₆ 417.501

Minor alkaloid from roots of *Stemona parviflora*.

5-O-β-D-Glucopyranosyloxy, 1',2'-didehydro(E-): **Stemofolinolide**

C₂₈H₃₇NO₁₁ 563.6

Alkaloid from an unidentified *Stemona* sp. Yellow-brown solid. [α]_D²² +138 (c, 0.16 in MeOH).

(E)-Isomer: **Isostemofoline**

[247089-64-1]

C₂₂H₂₉NO₅ 387.475

Alkaloid from *Stemona* spp.

(E)-Isomer, 1',2'-didehydro(E-): **1',2'-Didehydroisostemofoline**

C₂₂H₂₇NO₅ 385.459

Alkaloid from *Stemona collinsae*. Insecticide. Glassy resin. [α]_D¹⁸ +130 (c, 0.01 in MeOH). λ_{max} 295 (log ε 4.34) (MeOH).

Irie, H. et al., *Chem. Comm.*, 1970, 1066 (uv, ir, pmr, cryst struct, abs config)

Lin, W. et al., *Huaxue Xuebao*, 1991, **49**, 1034; *CA*, **116**, 148189c (4'-Hydroxystemofoline, 4'-Methoxystemofoline)

Sekine, T. et al., *J.C.S. Perkin 1*, 1995, 391 (*Asparagamine A*, cryst struct)

Kende, A.S. et al., *J.A.C.S.*, 1999, **121**, 7431–7432 (*Isostemofoline*, synth)

Jiwajinda, S. et al., *Phytochemistry*, 2001, **56**, 693–695 (*Didehydrostemofoline*, *Didehydroisostemofoline*)

Brem, B. et al., *J. Agric. Food Chem.*, 2002, **50**, 6383–6388 (cmr, 2'S-Hydroxystemofoline)

Seger, C. et al., *Chem. Biodiversity*, 2004, **1**, 265–279 (isol, pmr, cmr, biosynth, cryst struct)

Mungkornasawakul, P. et al., *J. Nat. Prod.*, 2004, **67**, 1740–1743 (11,12-Dihydrostemofoline)

Sastrarajji, T. et al., *J. Nat. Prod.*, 2005, **68**, 1763–1767 (2'-Hydroxystemofolines,

Stemofolenols, *Stemofolinolide*, 1',2'-Didehydrostemofoline N-oxide)


Sastrarajji, T. et al., *Nat. Prod. Commun.*, 2006, **1**, 813–818 (3'S-Hydroxystemofoline)

Greger, H. et al., *Planta Med.*, 2006, **72**, 99–113 (rev, occur, activity)

Tang, C.-P. et al., *J. Nat. Prod.*, 2008, **71**, 112–116 (6-Hydroxystemofoline, 16-Hydroxystemofoline)

Stemokerrin

S-516



C₂₂H₃₁NO₅ 389.491

Alkaloid from the roots of *Stemona kerrii*. Plates (CH₂Cl₂/hexane). Mp 138–141°. [α]_D²⁰ +136 (c, 0.3 in MeOH). λ_{max} 216 ; 306 (MeOH aq.).

N-Oxide: **Stemokerrin N-oxide**

[948994-13-6]

C₂₂H₃₁NO₆ 405.49

Alkaloid from the roots of *Stemona saxorum*. Pale yellow powder. [α]_D²² +118 (c, 0.24 in MeOH). λ_{max} 212 (log ε 3.89); 306 (log ε 4.3) (MeOH).

1'-Me ether, N-oxide: **1'-O-Methylstemokerrin N-oxide. Methoxystemokerrin N-oxide**

C₂₃H₃₃NO₆ 419.517

Alkaloid from the roots of *Stemona kerrii*. Amorph. [α]_D²⁰ +255 (c, 0.2 in MeOH). λ_{max} 216 ; 308 (MeOH aq.).

8,9-Dihydro, 1α,9α-epoxide: **Oxystemokerrin**

C₂₂H₃₁NO₆ 405.49

Alkaloid from the roots of *Stemona curtisii* and *Stemona kerrii*. Amorph. [α]_D²⁰ +289 (c, 0.4 in MeOH). λ_{max} 298 (MeOH aq.).

8,9-Dihydro, 1α,9α-epoxide, N-oxide: **Oxystemokerrin N-oxide**

C₂₂H₃₁NO₇ 421.489

Alkaloid from the roots of *Stemona kerrii*. Amorph. [α]_D²⁰ +247 (c, 0.3 in MeOH). λ_{max} 296 (MeOH aq.).

1',4-Diepimer, 8,9-dihydro, 1α,9α-epoxide: **Stemocurtisinol**

C₂₂H₃₁NO₆ 405.49

Alkaloid from the roots of *Stemona curtisii*. Pale yellow needles (EtOAc). Mp 209–211°. [α]_D²⁵ +233 (c, 0.33 in CHCl₃).

Kaltenegger, E. et al., *Phytochemistry*, 2003, **63**, 803–816 (isol, pmr, cmr, ms, cryst struct)

Mungkornasawakul, P. et al., *J. Nat. Prod.*, 2004, **67**, 675–677 (*Stemocurtisinol*)

Mungkornasawakul, P. et al., *ACGC Chem. Res. Commun.*, 2005, **19**, 30–33 (*Oxystemokerrin*, cryst struct)


Greger, H. et al., *Planta Med.*, 2006, **72**, 99–113 (rev)

Wang, Y.-Z. et al., *J. Nat. Prod.*, 2007, **70**, 1356–1359 (*N-oxide*)

Pyne, S.G. et al., *Maejo Int. J. Sci. Technol.*, 2007, **1**, 157–165 (rev, biosynth)

Stemonamide

[157009-09-1]

C₁₈H₂₁NO₅ 331.368

Alkaloid from roots of *Stemona japonica* (Stemonaceae). Mp 182.5-184°. [α]_D -120 (c, 0.79 in EtOH). Amide of Stemonamine, S-518.

12-Epimer: Isostemonamide

[156953-97-8]

C₁₈H₂₁NO₅ 331.368

From roots of *Stemona japonica* (Stemonaceae). Mp 234-236°. [α]_D -177 (c, 0.37 in EtOH).

Ye, Y. *et al.*, *J. Nat. Prod.*, 1994, **57**, 665-669 (*isol, ir, pmr, cmr, ms*)


Kende, A.S. *et al.*, *Tetrahedron*, 2002, **58**, 61-74 (*synth*)

Taniguchi, T. *et al.*, *Tetrahedron*, 2008, **64**, 8773-8779 (*synth*)

Stemonamine

S-518

2,3,5,6,7,8-Hexahydro-3'-methoxy-4',9-dimethylspiro[1H-cyclopenta[b]pyrrolo[1,2-a]azepine-11(10H),2'(5'H)-furan]-5',10-dione, 9CI



Relative configuration

C₁₈H₂₃NO₄ 317.384**(±)-form [41758-66-1]**

Alkaloid from the roots of *Stemona japonica* (Stemonaceae). Mp 172-174°. Stemonamine racemises and epimerises via a tricyclic achiral dipolar intermediate.

Hydrochloride:

Cryst. + 2H₂O (Me₂CO). Mp 148-151°.

Epimer: Isostemonamine

[41758-67-2]

C₁₈H₂₃NO₄ 317.384

Alkaloid from the roots of *Stemona japonica* (Stemonaceae). Mp 165-169°. Racemic. Epimeric with Stemonamine at the lactone spiro carbon.

Iizuka, H. *et al.*, *Chem. Comm.*, 1973, 125 (*uv, ir, cryst struct*)

Iizuka, H. *et al.*, *Cryst. Struct. Commun.*, 1973, **2**, 375 (*cryst struct*)

Zhao, Y.-M. *et al.*, *Org. Lett.*, 2008, **10**, 1763-1766 (*synth*)


Taniguchi, T. *et al.*, *Tetrahedron*, 2008, **64**, 8773-8779 (*synth*)

S-517

Stemonine†

S-519

Decahydro-1-methyl-8-(tetrahydro-4-methyl-5-oxo-2-furanyl)-2H-furo[3,2-c]pyrrolo[1,2-a]azepin-2-one, 9CI [27498-90-4]



Absolute Configuration

C₁₇H₂₅NO₄ 307.389

See also Stemonine, S-520. Alkaloid from *Stemona japonica* and *Stemona pierrei*. Also a degrad. prod. of Protostemonine, P-670. Mp 151°. [α]_D¹⁶ -113.8.

Hydrobromide:

Cryst. + ½ H₂O. Mp 275°.

1,2,3,9a-Tetrahydro: **Bisdehydrostemonine**. Tetrahydrostemonine. Didehydrostemonine [27498-91-5]

C₁₇H₂₁NO₄ 303.357

Alkaloid from *Stemona japonica*. Mp 172-175°.

Suzuki, K. *et al.*, *Yakugaku Zasshi*, 1929, **49**, 78; 1931, **51**, 44; 1934, **54**, 101 (*isol*)

Irie, H. *et al.*, *Chem. Comm.*, 1970, 268-269 (*uv, ir, pmr, ms, struct*)

Koyama, H. *et al.*, *J.C.S.(B)*, 1970, 1330 (*cryst struct, abs config*)

Zou, C.-Y. *et al.*, *J. Chin. Pharm. Sci.*, 2000, **9**, 113-115 (*Bisdehydrostemonine*)

Williams, D.R. *et al.*, *Org. Lett.*, 2003, **5**, 3361-3364 (*synth*)

Stemonine†

S-520

[20460-41-7]

C₂₂H₃₃NO₄ 375.507

Struct. unknown. Not the same as the alkaloid *isol.* from *Stemona ovata*. Probably a stereoisomer of Tuberostemonine, T-664. Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Cryst. (MeOH). Mp 162°. [α]_D²⁰ +76 (c, 1.0 in abs. EtOH).

Hydroiodide: Mp 247-248° dec.

Perchlorate: Mp 274-276° dec.


Methiodide: Mp 258-261° dec.

Pfeifer, S. *et al.*, *Pharmazie*, 1968, **23**, 342; *CA*, **69**, 74480c (*isol, uv, ir*)

Stemonidine

S-521

[69772-72-1]

C₂₂H₃₁NO₅ 389.491

Alkaloid from the roots of *Stemona sessilifolia*. Mp 113-115°. [α]_D¹⁴ -110 (c, 1.02 in EtOH).

12,13β-Dihydro: **Dihydrostemonidine**

C₂₂H₃₃NO₅ 391.506

Alkaloid from the roots of *Stemona*

sessilifolia. Prisms (EtOAc/petrol).

1,2,3,9a-Tetrahydro: **Bisdehydrostemonine**

C₂₂H₂₇NO₅ 385.459

Alkaloid from the roots of *Stemona sessilifolia* and *Stemona tuberosa*.

Cryst. (EtOAc). Mp 180-181° (*S. sessilifolia*) Mp 138-140° (*S. tuberosa*).

[α]_D²⁰ -57 (c, 0.13 in CHCl₃) (*S. tuberosa*). [α]_D²⁰ -125 (c, 0.11 in MeOH) (*S. sessilifolia*).

11-Epimer, 1,2,3,9a-tetrahydro: **Isobisdehydrostemonidine**

C₂₂H₂₇NO₅ 385.459

Alkaloid from the roots of *Stemona tuberosa*. Needles (EtOAc). Mp 178-182°.

Kuo, C. *et al.*, *Huaxue Xuebao*, 1978, **36**, 291; *CA*, **90**, 164717y (*isol*)

Cheng, D. *et al.*, *J. Nat. Prod.*, 1988, **51**, 202-211 (*pmr, cmr, struct*)


Lin, L.-G. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1051-1054 (*Bisdehydrostemonidine, Isobisdehydrostemonidine*)

Wang, P. *et al.*, *Chem. Biodiversity*, 2007, **4**, 523-530 (*Dihydrostemonidine*)

Stemoninoamide

S-522

[158340-34-2]



Absolute Configuration

C₁₇H₂₃NO₄ 305.373

Minor alkaloid from roots of *Stemona tuberosa* (Stemonaceae). Mp 155-157°. [α]_D -94 (c, 0.06 in MeOH).

12,13β-Dihydro: **Sessilifoliamide A**

C₁₇H₂₅NO₄ 307.389

Alkaloid from the roots of *Stemona sessilifolia*. Prisms (EtOAc). Mp 166-168°. [α]_D²⁷ -128 (c, 0.35 in CHCl₃).

[156280-89-6]

Lin, W.H. *et al.*, *Phytochemistry*, 1994, **36**,

1333-1335 (*isol, ir, pmr, cmr, ms, struct*)


Kakuta, D. *et al.*, *Tetrahedron*, 2003, **59**, 7779-7786 (*abs config, Sessilifoliamide A*)

Stemospironine

S-523

Decahydro-8'-methoxy-4-methyl-3'-(tetrahydro-4-methyl-5-oxo-2-furanyl)spiro[furan-2(5H),9'-[9H]pyrrolo[1,2-a]azepin]-5-one, 9CI. Stemonidine [66267-46-7]

[85700-47-6 (Stemonidine)]



Absolute Configuration

C₁₉H₂₉NO₅ 351.442

Alkaloid from *Stemona japonica* and *Stemona ovata*. Insecticidal agent. Non-cryst. Mp 283-284° dec. (as hydrobromide). [α]_D²⁷ -8.2 (c, 0.92 in CHCl₃).

Identity of the struct. of Stemonidine with Stemospironine demonstrated in 2007.

Demethoxy: Croomine

[71239-66-2]

C₁₈H₂₇NO₄ 321.416Alkaloid from *Croomia heterosepala* and *Stemona tuberosa*. Bp_{0.002} 210-215°. [α]_D¹⁸ +9.8 (c, 0.11 in CHCl₃).**Demethoxy; picrate:**

Cryst. (EtOH). Mp 125°.

Demethoxy, 6 α -hydroxy: 6 α -Hydroxy-croomineC₁₈H₂₇NO₅ 337.415Alkaloid from the roots of *Stemona tuberosa*. Powder. [α]_D²⁰ +24.6 (c, 0.5 in MeOH).**9-Epimer, demethoxy, 10R-hydroxy: Tuberospiroline**C₁₈H₂₇NO₅ 337.415Alkaloid from the roots of *Stemona tuberosa*. Powder. [α]_D²⁰ -14.4 (c, 0.5 in MeOH).Sakata, K. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 457; *CA*, **88**, 166716e (*isol. ir. pmr. cmr. ms. cryst. struct. abs. config.*)Noro, T. *et al.*, *Chem. Pharm. Bull.*, 1979, **27**, 1495 (*uv. ir. pmr. cmr. ms. cryst. struct. Croomine*)Xu, R.S. *et al.*, *Tetrahedron*, 1982, **38**, 2667 (*Croomine, pmr*)Williams, D.R. *et al.*, *J.A.C.S.*, 1989, **111**, 1923 (*synth. Croomine*)Martin, S.F. *et al.*, *J.A.C.S.*, 1999, **121**, 6990-6997 (*Croomine, synth*)Williams, D.R. *et al.*, *Org. Lett.*, 2001, **3**, 2721-2724 (*synth*)Schinnerl, J. *et al.*, *Monatsh. Chem.*, 2005, **136**, 1671-1680 (*6-Hydroxycroomine*)Jiang, R.-W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 749-754 (*Tuberospiroline*)Jiang, R.-W. *et al.*, *Phytochemistry*, 2006, **67**, 52-57 (*Croomine, 6-Hydroxycroomine*)Pyne, S.G. *et al.*, *Maejo Int. J. Sci. Technol.*, 2007, **1**, 157-165 (*rev. biosynth*)Sánchez-Izquierdo, F. *et al.*, *Org. Lett.*, 2007, **9**, 1769-1772 (*Stemomidine*)


ceae). Mp 77-82° dec.

Pfeifer, S. *et al.*, *Pharmazie*, 1968, **23**, 342-343 (*isol. uv*)**Stenanzine**

S-526

3,23-Dihydroxyveratraman-6(5H)-one, 9CI


[83133-08-8]

C₂₇H₄₃NO₃ 429.642Alkaloid from *Rhinopetalum stenantherrum* (Liliaceae). Mp 203-205°. [α]_D -44 (c, 0.5 in CHCl₃).Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 340; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 314 (*isol. struct*)**Stendomycidine**

S-527

 α -Amino-3,4,5,6-tetrahydro-3-methyl-2-(methylamino)-4-pyrimidineacetic acid, 9CI

[21948-17-4]


C₈H₁₆N₄O₂ 200.24

Component of Stendomycin. No antibiotic activity.

Bodanszky, M. *et al.*, *J. Antibiot.*, 1968, **21**, 77; 1969, **22**, 40 (*isol. struct*)Marconi, G.G. *et al.*, *J. Antibiot.*, 1970, **23**, 120 (*abs. config.*)**Stemotinine**

S-524

[85644-15-1]

C₁₈H₂₅NO₅ 335.399Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Prisms (Me₂CO/Et₂O). Mp 207-208°. [α]_D²² +91.7 (c, 1.1 in MeOH).**Epimer: Isostemotinine**

[85644-16-2]


C₁₈H₂₅NO₅ 335.399Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Prisms (Me₂CO/Et₂O). Mp 245-246°. [α]_D²² +47.5 (c, 0.6 in MeOH). Epimeric at the lactone spiro carbon atom.Xu, R.-S. *et al.*, *Tetrahedron*, 1982, **38**, 2667 (*isol. ir. pmr. cmr. ms. cd. struct*)**Stemotuberine**

S-525

Struct. unknown. Alkaloid from the roots of *Stemona tuberosa* (Stemona-**Stenine**

S-528

[16625-37-9]

[130760-62-2 (\pm)-form]C₁₇H₂₇NO₂ 277.406Numbering systems vary. Alkaloid from the roots of *Stemona tuberosa*. Prisms (hexane). Mp 65-67°. [α]_D -30.2 (MeOH).

3-Oxo: 3-Oxostenine. 2-Oxostenine

[18058-98-5]

C₁₇H₂₅NO₃ 291.389Alkaloid from *Stemona sessilifolia*. Mp 135-138°.

Absolute Configuration

13-Epimer: 15-Epistenine. Sessilifoline B

[929637-36-5]


C₁₇H₂₇NO₂ 277.406Alkaloid from the stems of *Stemona sessilifolia*. Amorph. powder. [α]_D²⁰ +34 (c, 0.2 in CHCl₃).**1,11,15-Triepimer: Neostenine**

[477953-07-4]

[1023306-02-6 (\pm)-form]C₁₇H₂₇NO₂ 277.406Alkaloid from the roots of *Stemona tuberosa*. Antitussive agent. Prisms (hexane/EtOAc). Mp 90-92°. [α]_D²⁰ +73.6 (c, 0.1 in MeOH). Both Neostenine and Isostenine below have been assigned the same stereochem., 1,11,15-triépistenine, although they have different props.**Stereoisomer: Isostenine**C₁₇H₂₇NO₂ 277.406Alkaloid from the roots of *Stemona collinsae*. Cryst. Mp 213-215°. [α]_D +92 (c, 0.6 in CHCl₃). See note under Neostenine above.Harada, H. *et al.*, *Chem. Comm.*, 1967, 460 (*abs. config.*)Uyeo, S. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 768 (*isol. ir. pmr. ms. struct.*)Chen, C.-Y. *et al.*, *J.O.C.*, 1993, **58**, 3840 (*synth.*)Wipf, P. *et al.*, *J.A.C.S.*, 1995, **117**, 11106 (*synth.*)Morimoto, Y. *et al.*, *Chem. Eur. J.*, 2001, **7**, 4107-4116 (*synth.*)Golden, J.E. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 4316-4318 (*synth.*)Pham, H.-D. *et al.*, *J. Asian Nat. Prod. Res.*, 2002, **4**, 81-85 (*Isostenine*)Ginn, J.D. *et al.*, *Org. Lett.*, 2002, **4**, 1515-1517 (*synth.*)Chung, H.-S. *et al.*, *Planta Med.*, 2003, **69**, 914-920 (*Neostenine*)Kakuta, D. *et al.*, *Tetrahedron*, 2003, **59**, 7779-7786 (*2-Oxostenine*)Zeng, Y. *et al.*, *J.A.C.S.*, 2005, **127**, 15712-15713 (*synth.*)Qian, J. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 326-331 (*Sessilifoline B*)Frankowski, K.J. *et al.*, *J.A.C.S.*, 2008, **130**, 6018-6024 (*synth.*)Lainchbury, M.D. *et al.*, *J.O.C.*, 2008, **73**, 6497-6505 (*Neostenine, synth.*)**Stephabenine**

S-529

[87853-56-3]

C₂₇H₂₉NO₇ 479.529Alkaloid from the fresh fruits of *Stephania japonica* (Menispermaceae). Prisms (EtOH). Mp 170°. [α]_D¹⁶ -15.24 (c, 1.65 in CHCl₃).**16-Oxo: Oxostephabenine**

[99964-46-2]


C₂₇H₂₇NO₈ 493.512

Alkaloid from the fruits of *Stephania japonica* (Menispermaceae). Needles (MeOH). Mp 272-273° dec. $[\alpha]_D^{24} +65$ (c, 1.04 in CHCl₃).

Kondo, S. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2574 (*isol, uv, ir, pmr, cmr, ms, struct*)
Yamamura, Y. *et al.*, *J. Nat. Prod.*, 1985, **48**, 746 (*Oxostephabenine*)

Stephabinamine S-530

5,8,13,13a-Tetrahydro-2,3,10-trimethoxy-6H-dibenzo[a,g]quinolizine-1,11-diol, 9CI



C₂₀H₂₃NO₅ 357.405

(S)-form [109028-34-4]

Alkaloid from the roots of *Stephania suberosa* (Menispermaceae). $[\alpha]_D^{25} -212$ (c, 0.13 in CHCl₃).

O¹¹-De-Me: **Tetrahydrostephabinine**

[109063-88-9]

C₂₁H₂₅NO₅ 371.432

Alkaloid from the roots of *Stephania suberosa* (Menispermaceae). $[\alpha]_D^{25} -277$ (c, 0.59 in CHCl₃).


[109028-33-3]

Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 547 (*isol, pmr, ms, cd, struct*)

Stephazine S-531

5,6-Dihydro-1-hydroxy-2,3,10,11-tetramethoxydibenzo[a,g]quinolizinium(1+), 9CI

[109028-32-2]




C₂₁H₂₂NO₅⁺ 368.409

Alkaloid from the roots of *Stephania suberosa* (Menispermaceae).

Patra, A. *et al.*, *Phytochemistry*, 1987, **26**, 547 (*isol, uv, pmr, ms, struct*)

Stephacidin B S-532

[360765-75-9]



Absolute Configuration

C₅₂H₅₄N₆O₈ 891.034

Prod. by *Aspergillus ochraceus* WC76466. Cytotoxic. Off-white amorph. solid. λ_{max} 240 ; 268 (sh) ; 301 ; 346 (sh) (MeOH).

Qian-Cutrone, J. *et al.*, *J.A.C.S.*, 2002, **124**, 14556-14557 (*isol, struct*)

Von Nussbaum, F. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3068-3071 (*rev, biosynth*)

Herzon, S.B. *et al.*, *J.A.C.S.*, 2005, **127**, 5342-5344 (*synth, abs config*)


Baran, P.S. *et al.*, *J.A.C.S.*, 2006, **128**, 8678-8693 (*synth, abs config*)

Artman, G.D. *et al.*, *J.A.C.S.*, 2007, **129**, 6336-6342 (*synth*)

Stephadiamine

[94443-36-4]

S-533



C₁₉H₂₄N₂O₄ 344.41

Alkaloid from *Stephania japonica* (Menispermaceae). Prisms (Me₂CO). Mp 180°. $[\alpha]_D^{20} +51.8$ (c, 0.54 in CHCl₃). First example of a C-norhasubanan alkaloid.


N-Ac: Mp 248°.

Taga, T. *et al.*, *Chem. Pharm. Bull.*, 1984, **32**, 4223 (*ir, pmr, cryst struct*)

Stephadiamine

S-534

4,7-Dihydroxy-1,2-methylenedioxyaporphine [115044-74-1]



C₁₈H₁₇NO₄ 311.337

The abs. configs. of the alkaloids in this entry may not be well established.

N-Oxide (β -): **Stephadiamine β -N-oxide**

[113145-66-7]

C₁₈H₁₇NO₅ 327.336

Alkaloid from the rhizomes of *Stephania venosa* (Menispermaceae). Amorph. $[\alpha]_D -37$ (c, 0.06 in MeOH).

4,7-Diepimer, O⁷-Me: **Pachystaudine**

[67627-76-3]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the bark of *Pachypodanthium staudtii* (Annonaceae). Mp 157°. $[\alpha]_D +34$ (c, 0.5 in CHCl₃).

4,7-Diepimer, O⁷-Me, N-de-Me: **Norpachystaudine**

[67627-78-5]

C₁₈H₁₇NO₄ 311.337


Alkaloid from the bark of *Pachypodanthium staudtii* (Annonaceae). Mp 214-216°. $[\alpha]_D +5$ (c, 0.5 in CHCl₃).

Bevalot, F. *et al.*, *Plant. Med. Phytother.*, 1977, **11**, 315 (*Pachystaudine, Norpachystaudine*)
Charles, B. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1113 (*Stephadiamine N-oxide*)

Stephadione

S-535

[142905-22-4]



C₁₉H₁₁NO₆ 349.299


Alkaloid from aerial parts of *Stephania tetrandra* (Menispermaceae). Amorph. red powder. Mp 300°.

Si, D.Y. *et al.*, *J. Nat. Prod.*, 1992, **55**, 828 (*isol, uv, ir, pmr, ms, struct*)

Stephalagine

S-536

3-Methoxy-1,2-methylenedioxyaporphine [52617-22-8]



C₁₉H₁₉NO₃ 309.364

Alkaloid from the stems of *Stephania dinklagei* (Menispermaceae). Needles (Et₂O). Mp 128-130° dec. $[\alpha]_D^{28} -30$ (c, 1.0 in CHCl₃).

Hydrochloride: Mp 263-264° dec.

Hydroiodide:

Cryst. (MeOH). Mp 218° dec.

N-De-Me: **Norstephalagine. 3-Methoxy-1,2-methylenedioxyaporphine** [80151-82-2]

C₁₈H₁₇NO₃ 295.337

Alkaloid from *Xylopi buxifolia* and from the stem and root barks of *Hexalobus crispiflorus* (Annonaceae). Cryst. (diisopropyl ether). Mp 94-95°. $[\alpha]_D^{20} -35$ (c, 0.98 in EtOH).

6a,7-Didehydro: **Dehydrostephalagine**

[107882-28-0]

C₁₉H₁₇NO₃ 307.348

Alkaloid from the leaves of *Gutteria sagotiana* (Annonaceae). Amorph.

7-Hydroxy: see Guatterine, G-213

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 6 (*isol, uv, pmr, ms, struct*)


Hocquemiller, R. *et al.*, *J. Nat. Prod.*, 1981, **44**, 551 (*Norstephalagine*)

Achenbach, H. *et al.*, *Annalen*, 1982, 1132; 1623 (*Norstephalagine*)

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1078 (*Dehydrostephalagine*)

Stephamiersine S-537

8,10-Epoxy-3,4,7,8-tetramethoxy-17-methylhasubanan-6-one, 9CI
[52309-76-9]



C₂₁H₂₇NO₆ 389.447
Alkaloid from the stems and rhizomes of *Stephania japonica* (Menispermaceae). Prisms (MeOH). Mp 165°. [α]_D²⁰ +33 (c, 1.28 in CHCl₃).

Hydroiodide:

Cryst. + ½ H₂O (MeOH). Mp 253° dec.

16-Oxo: **Oxostephamiersine**. 8,10-Epoxy-3,4,7,8-tetramethoxy-17-methylhasubanan-6,16-dione, 9CI
[52466-83-8]

C₂₁H₂₅NO₇ 403.431

Alkaloid from leaves, stems and rhizomes of *Stephania japonica*, and from *Stephania japonica* var. *australis* (Menispermaceae). Prisms (MeOH). Mp 256°. Mp 290° (dimorph.). [α]_D²⁰ +88.25 (c, 1.87 in CHCl₃).

7-Epimer: **Epistephamiersine**

[52389-15-8]

C₂₁H₂₇NO₆ 389.447

Alkaloid from stems and rhizomes of *Stephania japonica* (Menispermaceae). Prisms (MeOH). Mp 98°. [α]_D²⁰ +64.1 (c, 1.12 in CHCl₃).

7-Epimer; hydroiodide:

Cryst. (MeOH). Mp 217° dec.

7-Epimer, 16-oxo: **Oxoepistephamiersine**

[51804-68-3]

C₂₁H₂₅NO₇ 403.431

Alkaloid from roots of *Stephania japonica* (Menispermaceae). Light-yellow prisms (MeOH). Mp 228°. [α]_D¹³ +104.88 (c, 1.0 in CHCl₃).

Matsui, M. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 1323 (*isol, uv, ir, pmr, ms, struct*)

Matsui, M. *et al.*, *Phytochemistry*, 1979, **18**, 1087 (*Oxostephamiersine*)

Matsui, M. *et al.*, *J. Nat. Prod.*, 1982, **45**, 247 (*cmr*)

Matsui, M. *et al.*, *J. Nat. Prod.*, 1982, **45**, 497; 1984, **47**, 858 (*Oxostephamiersine, Oxoepistephamiersine*)


Stephanoline S-538

C₃₁H₄₂N₂O₇ 554.682

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from *Stephania japonica* (Menispermaceae). Mp 186°. [α]_D¹⁷ -255.

Kondo, H. *et al.*, *Yakugaku Zasshi*, 1928, **48**, 1141-1156; *CA*, **23**, 2978

Stephanosporin S-539




C₁₀H₁₀ClN₃O₆ 303.658

Prod. by the carrot truffle, *Stephanospora caroticolor*. Yellow solid. Mp 191-194° dec. λ _{max} 226 (ε 11925); 258 (ε 12580); 312 (ε 4740); 351 (sh) (ε 3750) (MeOH).

Lang, M. *et al.*, *Angew. Chem., Int. Ed.*, 2001, **40**, 1704-1705 (*isol, synth, uv, ir, pmr, cmr*)

Stephanotic acid S-540

[268541-40-8]



C₃₃H₄₆N₆O₇ 638.762


Constit. of *Stephanotis floribunda*. Mp 250-252°. [α]_D²⁵ -143 (c, 1.16 in MeOH). λ _{max} 231 (log ε 4.31); 275 (log ε 3.76); 280 (log ε 3.79); 292 (log ε 3.72) (MeOH).

Yoshikawa, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 540-542 (*Stephanotic acid*)

Bentley, D.J. *et al.*, *Org. Lett.*, 2006, **8**, 1975-1978 (*synth*)

Stephaoxocanine S-541

[179458-25-4]



Absolute Configuration

C₁₈H₂₁NO₄ 315.368

Alkaloid from tubers of *Stephania cepharantha*. Fine needles (Me₂CO). Mp 160-162°. [α]_D²⁴ +60 (c, 0.67 in CHCl₃). λ _{max} 255 (log ε 4.48); 286 (log ε 4.15); 330 (sh) (log ε 3.65) (MeOH).

3,4,14,15-Tetrahydro: **Stephaoxocandine**

[189393-37-1]

C₁₈H₁₇NO₄ 311.337

Alkaloid from the tubers of *Stephania cepharantha* (Menispermaceae). Pale yellow needles (Et₂O). Mp 188-191°. [α]_D²⁵ +30 (c, 0.9 in CHCl₃). λ _{max} 235

(sh) (log ε 4.44); 247 (log ε 4.51); 340 (log ε 3.89) (MeOH).

3,4,14,15-Tetrahydro, 5-methoxy: **Eletefine**

[216165-78-5]

C₁₉H₁₉NO₅ 341.363

Alkaloid from the roots of *Cissampelos glaberrima*. Red-brown wax.


Kashiwaba, N. *et al.*, *J. Nat. Prod.*, 1996, **59**, 803-805 (*Stephaoxocanine*)

Kashiwaba, N. *et al.*, *Nat. Prod. Lett.*, 1997, **9**, 177-180 (*Stephaoxocandine*)

Da-Cunha, E.V.L. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1140-1142 (*Eletefine*)

Stepharinosine S-542

[113145-63-4]



C₁₈H₂₁NO₄ 315.368

Alkaloid from the rhizomes of *Stephania venosa* (Menispermaceae). Amorph. Unstable and tends to dec. in soln.

Me ether: **O-Methylstepharinosine**

[113145-64-5]


C₁₉H₂₃NO₄ 329.395

Alkaloid from rhizomes of *Stephania venosa* (Menispermaceae). Amorph. [α]_D -26 (c, 0.11 in CHCl₃). Unstable and tends to dec. in soln.

Charles, B. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1113 (*isol, ir, pmr, ms, struct*)

Stepharotine S-543

5,8,13,13a-Tetrahydro-2,3,9,10-tetramethoxy-6H-dibenzo[a,g]quinolizin-11-ol, 9CI



C₂₁H₂₅NO₅ 371.432

Struct. still doubtful despite the synth. of the racemate.

(S)-form [13588-89-1]

[7668-83-9]

Alkaloid from root tubers of *Stephania rotunda* (Menispermaceae). Mp 227-229° (as hydrobromide). [α]_D -303 (MeOH).

Me ether:

Cryst. (H₂O). Mp 133-135°.

8-Oxo, O²,O⁹-di-O-de-Me: **8-Oxopolyalthiane**

C₁₉H₁₉NO₆ 357.362

Alkaloid from *Polyalthia longifolia* var.

pendula. Amorph. yellow powder. Mp 95-97°. $[\alpha]_D^{25}$ -264.8 (c, 0.02 in CHCl_3). λ_{max} 238 (log ϵ 4.3); 270 (log ϵ 3.9); 353 (log ϵ 3.5) (MeOH).

(±)-form [19598-17-5]

V. pale-yellow prisms (MeOH). Mp 192-193° dec.

Hydrobromide: [19598-18-6]

Needles (MeOH). Mp 258-260° dec.

Tomita, M. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 460 (*isol*)


Augstein, W. *et al.*, *J.O.C.*, 1969, **34**, 1349 (*synth, struct, uv, pmr*)

Ringdahl, B. *et al.*, *J. Nat. Prod.*, 1981, **44**, 75 (*cd, abs config*)

Chen, C.Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1475-1478 (*8-Oxopolyalthiane*)

Stephasubine

S-544



$\text{C}_{36}\text{H}_{34}\text{N}_2\text{O}_6$ 590.674

(R)-form [104386-89-2]

Alkaloid from the tuberous roots of *Stephania suberosa* (Menispermaceae) and stems of *Stephania hernandifolia*. $[\alpha]_D^{25}$ +339 (c, 0.09 in MeOH). λ_{max} 240 (log ϵ 4.56); 287 (log ϵ 3.61); 337 (log ϵ 3.43) (MeOH).

N²-Me: Colorflamine

[80550-39-6]

$\text{C}_{37}\text{H}_{37}\text{N}_2\text{O}_6$ 605.709

Alkaloid from the stem and roots of *Pycnarrhena longifolia* (Menispermaceae). $[\alpha]_D^{20}$ +1050 (c, 0.06 in CHCl_3). No counterion indicated, prob. chloride. λ_{max} 233; 292; 333; 439 (MeOH).

N-De-Me: Norstephasubine

$\text{C}_{35}\text{H}_{32}\text{N}_2\text{O}_6$ 576.648

Isol. from *Stephania suberosa*. $[\alpha]_D^{25}$ +309 (c, 0.09 in MeOH). λ_{max} 240 (log ϵ 4.53); 286 (log ϵ 3.62); 338 (log ϵ 3.42) (MeOH).

O¹²-De-Me, N-de-Me: Pycnazanthine

[11481-94-8]

$\text{C}_{34}\text{H}_{30}\text{N}_2\text{O}_6$ 562.621

Alkaloid from the stems of *Pycnarrhena ozantha* (Menispermaceae). $[\alpha]_D^{25}$ +186 (c, 0.29 in MeOH). Tentative struct. May be the 6-OH, 12-OMe isomer. λ_{max} 230 (log ϵ 4.8); 284 (sh) (log ϵ 4.12); 318 (sh) (log ϵ 3.8) (MeOH).

1,2-Didehydro, N-de-Me: Stephasubimine

[104386-88-1]

$\text{C}_{35}\text{H}_{30}\text{N}_2\text{O}_6$ 574.632

Alkaloid from tuberous roots of *Stephania suberosa* (Menispermaceae). λ_{max} 242 (log ϵ 4.64); 281 (log ϵ 3.91); 323 (log ϵ 3.7) (MeOH).

Van Beek, T.A. *et al.*, *J.O.C.*, 1982, **47**, 898-900 (*Colorflamine*)

Patra, A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 424-427 (*Stephasubimine, Stephasubine, Norstephasubine*)


Abouchacra, M.L. *et al.*, *J. Nat. Prod.*, 1987, **50**, 375-380 (*Pycnazanthine*)

Chen, C.-K. *et al.*, *Chin. Pharm. J. (Taipei)*, 2003, **55**, 35-47 (*Colorflamine*)

Stephavanine

[33116-33-5]

S-545



Absolute configuration

$\text{C}_{26}\text{H}_{27}\text{NO}_9$ 497.501

Alkaloid from rhizomes of *Stephania abyssinica* (Menispermaceae). Antineoplastic agent. Mp 229-230° dec. $[\alpha]_D^{25}$ +30 (c, 0.90 in Py). Log P 0.61 (uncertain value) (calc).

Hydrochloride: Mp 217-218° dec. $[\alpha]_D^{22}$ +16 (c, 0.73 in MeOH).

Hydrobromide: Mp 191-192° dec.

N,O,O-Tri-Ac: Mp 189-190°.

4'-Me ether: 4'-O-Methylstephavanine

[152013-83-7]

$\text{C}_{27}\text{H}_{29}\text{NO}_9$ 511.527

Alkaloid from roots of *Stephania abyssinica* (Menispermaceae).

Amorph. solid. Mp 187-190° dec. $[\alpha]_D^{20}$ -4 (c, 0.01 in CHCl_3).

N,O,O-Tri-Me: N,O,O-Trimethylstephavanine

[53111-19-6]

$\text{C}_{29}\text{H}_{33}\text{NO}_9$ 539.581

Alkaloid from *Stephania abyssinica* (Menispermaceae).

Kupchan, S.M. *et al.*, *J.A.C.S.*, 1970, **92**, 5756 (*ms, pmr, struct, cryst struct*)


Van Wyk, A.J. *et al.*, *J. S. Afr. Chem. Inst.*, 1974, **27**, 95 (*deriv*)

Dagne, E. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2022 (*4'-O-Methylstephavanine*)

Stephananthrine

S-546

N,N-Dimethylphenanthro[3,4-d]-1,3-dioxole-5-ethanamine, *9CI*. 1-(2-Dimethylaminoethyl)-3,4-methylenedioxyphenanthrene. *De-N-methyl-1-isoremerine* [22108-99-2]



$\text{C}_{19}\text{H}_{19}\text{NO}_2$ 293.365

Aporphine-derived numbering shown. Alkaloid from the roots of *Stephania tetrandra* (Menispermaceae). Needles (CH_2Cl_2 /hexane). Mp 92-93°. λ_{max} 213 (log ϵ 0.81); 239 (log ϵ 1.07); 249 (log ϵ 1.31); 284 (log ϵ 0.43); 320 (log ϵ 0.25); 351 (log ϵ 0.04); 369 (log ϵ 0.03) (MeOH).

N-Oxide: Stephenanthrine N-oxide

[138706-50-0]

$\text{C}_{19}\text{H}_{19}\text{NO}_3$ 309.364

Alkaloid from *Monocyclanthus vignei* (Annonaceae). Oil. Possible artifact.

1'-ξ-Hydroxy: Fenfangjine F

$\text{C}_{19}\text{H}_{19}\text{NO}_3$ 309.364

Alkaloid from the roots of *Stephania tetrandra*. Oil.

8-Hydroxy: 8-Hydroxystephananthrine

[138690-44-5]

$\text{C}_{19}\text{H}_{19}\text{NO}_3$ 309.364

Alkaloid from *Monocyclanthus vignei* (Annonaceae). Oil.

8-Hydroxy, N-oxide: 8-Hydroxystephananthrine N-oxide

[138690-46-7]

$\text{C}_{19}\text{H}_{19}\text{NO}_4$ 325.363

Alkaloid from *Monocyclanthus vignei* (Annonaceae). Oil. Possible artifact.

Hu, T. *et al.*, *Yaoxue Xuebao*, 1986, **21**, 29; *CA*, **104**, 221975q (*Stephananthrine*)

Achenbach, H. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1331 (*Stephananthrine N-oxide, 8-Hydroxystephananthrene, 8-Hydroxystephananthrine N-oxide*)


Ogino, T. *et al.*, *Heterocycles*, 1998, **48**, 311-317 (*Fenfangjine F*)

Lopez-Martin, J. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1613-1615 (*isol, pmr, cmr*)

Stepholidine

S-547

5,8,13,13a-Tetrahydro-3,9-dimethoxy-6H-dibenzo[a,g]quinolizine-2,10-diol, *9CI*



(S)-form

$\text{C}_{19}\text{H}_{21}\text{NO}_4$ 327.379

(S)-form [16562-13-3]

Alkaloid from *Stephania glabra* (Menispermaceae). Needles (Me_2CO). Mp 126-138°. $[\alpha]_D^{18}$ -311 (c, 0.523 in EtOH). λ_{max} 214 (log ϵ 4.4); 287 (log ϵ 3.79) (EtOH).

►DR9823500

Di-Me ether: see Tetrahydropalmatine, T-212

(±)-form [16562-14-4]

Mp 136-140° dec.

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2785-2789 (*uv, ir, ms*)


Chiang, H.C. *et al.*, *J.O.C.*, 1977, **42**, 3190-3194 (*ms, pmr, synth*)

Wu, S. *et al.*, *Acta Cryst. C*, 1987, **43**, 2126-2128 (*cryst struct*)

Stepinonine

S-548

[38835-79-9]



Absolute configuration

C₃₆H₃₄N₂O₇ 606.674

Alkaloid from *Stephania japonica* (Menispermaceae). Yellow prisms. Mp 244-245° Mp 280° (dimorph.). [α]_D³⁰ -28 (c, 1 in Py).

Oxalate: Mp 208-210° dec.


Ac: Mp 157°. [α]_D²⁰ -12 (c, 0.5 in CHCl₃).

Ibukawa, T. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 114 (*isol, uv, ir, ms, pmr, struct*)

Steporphine

S-549

4-Hydroxy-1,2-methylenedioxyaporphine.
4-Hydroxyroemerine

C₁₈H₁₇NO₃ 295.337(-)-*form* [24191-98-8]

Alkaloid from *Stephania sasakii* and from the stems of *Stephania dinklagei* (Menispermaceae). Pale-yellow needles (Me₂CO/Et₂O). Mp 177-179°. [α]_D -90.6 (MeOH).

▶ CE1055500

4-Epimer: *Episteporphine*

[56688-64-3]

C₁₈H₁₇NO₃ 295.337

Alkaloid from the leaves of *Colubrina faralaoetra* subsp. *faralaoetra* (Rhamnaceae). Cryst. (MeOH). Mp 198°. [α]_D²⁰ -35 (c, 1.1 in CHCl₃).

4-Epimer, N-de-Me: 4-Hydroxyanonaine

[35597-00-3]

C₁₇H₁₅NO₃ 281.31

Alkaloid from the bark and leaves of *Laurelia philippiana* (Monimiaceae). Amorph. Isol. in admixture with 4-Hydroxynornantenine in N-29.

(±)-*form*

Synthetic. Needles (Me₂CO/Et₂O). Mp 172-173°.

Kunitomo, J. *et al.*, *Tet. Lett.*, 1969, 3287 (*uv, ir, pmr, ms, struct*)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 6 (*isol, uv, pmr, ms*)


Guinaudeau, H. *et al.*, *Planta Med.*, 1975, **27**, 304 (*epimer*)

Kunitomo, J. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 4283 (*synth, uv, ir, pmr, ms*)

Urzúa, A. *et al.*, *Phytochemistry*, 1982, **21**, 773 (*4-Hydroxyanonaine*)

Sterculinine I

S-550

C₁₈H₂₀N₂O₆ 360.366

Alkaloid from the seeds of *Sterculia lychnophora* (Pangdahai). Powder. Mp 184-186°. [α]_D²⁰ +24.2 (c, 0.5 in H₂O). λ _{max} 271 (log ϵ 3.75); 331 (log ϵ 3.12) (H₂O).

Me ester analogue: *Sterculinine II*C₁₅H₁₄N₂O₆ 318.285

Alkaloid from the seeds of *Sterculia lychnophora*. Powder. Mp 176-178°. [α]_D²⁰ +25 (c, 0.5 in H₂O). λ _{max} 275 (log ϵ 3.76); 333 (log ϵ 3.15) (H₂O).

Wang, R.-F. *et al.*, *Phytochemistry*, 2003, **63**, 475-478 (*isol, pmr, cmr, ms*)

Sternine

S-551

C₁₈H₂₁NO₃ 299.369


Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Sternbergia fischeriana* (Amaryllidaceae). Mp 231-232°. [α]_D +12.5 (EtOH).

Hydrochloride: Mp 181-182°.*Hydrobromide*:Cryst. + 1H₂O (H₂O). Mp 226-226.5°.*Picrate*: Mp 227-228°.*Methiodide*: Mp 202-203°.Zh. *Obshch. Khim.*, 1953, **23**, 2056-2059

Proskurnina, N.F. *et al.*, *Chem. Zentralbl.*, 1954, **125**, 8353 (*isol*)

Stichoneurine A

S-552

C₂₂H₃₅NO₄ 377.523

Alkaloid from the roots and rhizomes of *Stichoneuron caudatum*. Isol. as an inseparable mixt. with Stichoneurine B.

10-Epimer: *Stichoneurine B*C₂₂H₃₅NO₄ 377.523

Alkaloid from the roots and rhizomes of *Stichoneuron caudatum*.


Schinnerl, J. *et al.*, *Monatsh. Chem.*, 2005, **136**, 1671-1680 (*isol, pmr, cmr*)

Stizolamine

S-553

[3,4-Dihydro-5-(hydroxymethyl)-4-methyl-3-oxopyrazinyl]guanidine, 9CI. 3-Guanidino-6-hydroxymethyl-1-methylpyrazin-2-one

[61481-34-3]

C₇H₁₁N₅O₂ 197.196

Alkaloid from seeds of *Stizolobium hassjoo* (Fabaceae). Mp 208° dec.

Hydrochloride: Mp 294-296° dec.

Yoshida, T. *et al.*, *Phytochemistry*, 1976, **15**, 1723; 1977, **16**, 1824 (*isol, uv, ir, ms, pmr, struct, biosynth*)

Stizolophine

S-554

C₁₅H₂₃NO₅ 297.35


Struct. unknown. Alkaloid from the aerial parts of *Stizolophus balsamita*. Cryst. (CHCl₃). Mp 122-123°. [α]_D +24.6 (EtOH).

Kuzovkov, A.D. *et al.*, *Zh. Obshch. Khim.*, 1953, **23**, 157-158; *CA*, **48**, 696i

Stockerine

S-555

[136771-39-6]

C₂₄H₄₆N₂O₂ 394.64


Metab. from the alga *Stockeyia indica*. Amorph.

Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1991, **62**, 77 (*isol, ir, pmr, cmr, ms, struct*)

Storniamide D

S-556

[174285-74-6]

C₄₂H₃₅N₃O₁₃ 789.751

Alkaloid from a Patagonian sponge *Cliona* sp. Shows antibiotic activity against gram-positive bacteria. Yellow oil. λ _{max} 212 (€ 29500); 284 (€ 11500); 338 (€ 12400) (MeOH) (Berdy).

3'-Deoxy: *Storniamide B*

[174232-39-4]

C₄₂H₃₅N₃O₁₂ 773.751

From *Cliona* sp. Active against gram-positive bacteria. Yellow oil. λ _{max} 210 (€ 24500); 282 (€ 10200); 334 (€ 10800) (MeOH) (Berdy).

3'''-Deoxy: *Storniamide C*

[174232-40-7]

C₄₂H₃₅N₃O₁₂ 773.751

From *Cliona* sp. Active against gram-positive bacteria. Yellow oil. λ _{max} 214 (€ 27000); 284 (€ 11100); 342 (€ 14300) (MeOH) (Berdy).

3',3''-Dideoxy: *Storniamide A*

[174232-38-3]

C₄₂H₃₅N₃O₁₁ 757.752

From *Cliona* sp. Active against gram-positive bacteria. Yellow oil. λ _{max} 208 (€ 22500); 286 (€ 11300); 334 (€ 11500) (MeOH) (Berdy).

Palermo, J.A. *et al.*, *Tetrahedron*, 1996, **52**, 2727 (*isol, uv, ir, pmr, cmr, struct*)

Ebel, H. *et al.*, *Tet. Lett.*, 1998, **39**, 9165-9166 (*synth*)


Boger, D.L. *et al.*, *J.A.C.S.*, 1999, **121**, 54-62 (*Storniamide A, synth*)

Iwao, M. *et al.*, *Tet. Lett.*, 2003, **44**, 4443-4446 (*Storniamide A, synth*)

Gupton, J.T. *et al.*, *Tetrahedron*, 2008, **64**, 5246-5253 (*synth*)

Strellidimine

[112547-87-2]


C₃₄H₂₈N₄O 508.621

Alkaloid from the bark of *Strychnos dinklagei* (Loganiaceae). Yellow amorph. solid. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁰ 0 (c. 0.1 in MeOH). λ_{max} 244 (ε 24550); 253 (ε 26300); 265 (ε 21380); 279 (ε 19500); 340 (ε 4265); 410 (ε 3830) (EtOH) (Berdy).

Michel, S. *et al.*, *Chem. Comm.*, 1987, 229-230 (uv, ir, pmr, ms, struct)

Strepeliopidine

[96861-87-9]


C₃₈H₄₈N₄ 560.824

Alkaloid from the roots of *Strepeliopsis strepelioides* (Apocynaceae). Mp 133-135°. [α]_D²⁰ +100 (CHCl₃).

Laguna, A. *et al.*, *Planta Med.*, 1984, **50**, 285-288 (isol, uv, ir, pmr, ms, struct)

Strepeliopine

[79808-95-0]



Absolute Configuration

C₁₉H₂₂N₂O 294.396

Alkaloid from *Strepeliopsis strepelioides* (Apocynaceae). Cryst. (MeOH). Mp 152-154°. [α]_D -25.4 (c. 1.8 in MeOH). A prev. reported opt. rotn. of -120° was erroneous.


Laguna, A. *et al.*, *Planta Med.*, 1984, **50**, 285 (isol, uv, ir, pmr, ms, struct)

S-557

Hájček, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1986, **51**, 1731 (synth, uv, ir, pmr, ms, abs config)

Streptazoline

4-Ethylidene-2a,3,4,6,7,7b-hexahydro-3-hydroxy-1H-2-oxa-7a-azacyclopent[*c*-*d*]inden-1-one, 9CI
[80152-07-4]



Absolute Configuration

C₁₁H₁₃NO₃ 207.229

Authors' numbering shown (Puder *et al.*). Isol. from *Streptomyces viridochromogenes* and the marine-derived *Streptomyces* sp. B5525. [α]_D²⁵ +22 (c. 2.8 in CHCl₃). Readily polymerises. λ_{max} 256 (ε 8510) (EtOH) (Derep). λ_{max} 256 (ε 9000) (EtOH) (Berdy).

5-O-β-D-Xylopyranoside: 5-O-β-D-Xylopyranosylstreptazolin

C₁₆H₂₁NO₇ 339.344

Prod. by *Streptomyces* sp. strain A1. Solid. Mp 118°. [α]_D²² +1.3 (c. 0.8 in MeOH). λ_{max} 252 (log ε 4.23) (MeOH).

9R-Hydroxy: 9-Hydroxystreptazolin

C₁₁H₁₃NO₄ 223.228

Prod. by *Streptomyces* sp. strain A1. Oil. [α]_D²² +38 (c. 1.1 in MeOH). λ_{max} 256 (log ε 3.84) (MeOH).

13-Hydroxy: 13-Hydroxystreptazolin

C₁₁H₁₃NO₄ 223.228

Prod. by *Streptomyces* sp. strain A1. Oil. [α]_D²² +39 (c. 1.1 in MeOH). λ_{max} 256 (log ε 4.04); 315 (log ε 2.7) (MeOH).

Drautz, H. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 1752-1765 (isol)

Karrer, A. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 1432 (cryst struct)

Flann, C.J. *et al.*, *J.A.C.S.*, 1987, **109**, 6115 (synth)

Kozikowski, A.P. *et al.*, *J.O.C.*, 1990, **55**, 4668 (synth)

Mayer, M. *et al.*, *J.O.C.*, 1993, **58**, 3486 (biosynth, bibl)

Huang, S. *et al.*, *Chem. Comm.*, 2000, 569-570 (synth)

Cosy, J. *et al.*, *Eur. J. Org. Chem.*, 2001, 2841-2850 (Dihydrostreptazolin, synth)

Puder, C. *et al.*, *J. Nat. Prod.*, 2001, **64**, 42-45 (9-hydroxy, 13-hydroxy, 5-xylopyl)


Maskey, R.P. *et al.*, *Z. Naturforsch.*, **B**, 2002, **57**, 823-829 (marine, isol)

Trost, B.M. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 4327-4329 (synth)

Li, F. *et al.*, *J.O.C.*, 2006, **71**, 5221-5227 (synth)

Streptazone B₁

S-561

C₁₀H₁₁NO 161.203

Prod. by *Streptomyces* sp. FORM5. Oil. Related to Abikoviromycin, A-14. λ_{max} 262 (log ε 4.14); 315 (log ε 3.99) (MeOH).

4ξ,4αξ-Epoxyde: Streptazone A

C₁₀H₁₁NO₂ 177.202

Prod. by *Streptomyces* sp. FORM5. Mp 133°. [α]_D²² +118 (c. 0.34 in MeOH). Config. not determined. λ_{max} 239 (log ε 4.19); 310 (log ε 4.37) (MeOH).

(E)-Isomer: Streptazone B₂


C₁₀H₁₁NO 161.203

Prod. by *Streptomyces* sp. FORM5. Oil. λ_{max} 257 (log ε 4.1); 319 (log ε 3.91) (MeOH).

Puder, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1258-1260

Streptazone C

S-562

C₁₀H₁₁NO 161.203


Prod. by *Streptomyces* sp. FORM5. Yellow oil. λ_{max} 286 (log ε 3.76); 368 (log ε 3.39) (MeOH).

Puder, C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1258-1260

Streptimidone

S-563

4-(2-Hydroxy-5,7-dimethyl-4-oxo-6,8-nonadienyl)-2,6-piperidinedione, 9CI. Protomycin. S 632A₁. Antibiotic S 632A₁ [738-72-7]

C₁₆H₂₃NO₄ 293.362

Glutarimide-type antibiotic. Isol. from *Streptomyces* spp. and *Micromonospora coerulea*. Shows antibiotic props. Cryst. (Me₂CO/diisopropyl ether). Mp 72-73°. [α]_D²⁵ +238 (c. 0.5 in CHCl₃). λ_{max} 232 (ε 23100); 291 (ε 790) (EtOH) (Derep).

▶ LD₅₀ (mus, ipr) 280 mg/kg. TM7525000

Oxime:

Cryst. (MeOH aq.). Mp 141-146°. [α]_D²⁷ +65 (c. 0.5 in MeOH).

Ac:

Cryst. (Me₂CO/diisopropyl ether). Mp 108-110°. [α]_D +232 (c. 0.5 in MeOH).

Frohardt, R.P. *et al.*, *J.A.C.S.*, 1959, **81**, 5500-5506 (isol)

Van Tاملen, E.E. *et al.*, *J.A.C.S.*, 1960, **82**, 2974 (struct)

Woo, P.W.K. *et al.*, *J.A.C.S.*, 1961, **83**, 3085 (struct)


Sugawara, R. *et al.*, *J. Antibiot., Ser. A*, 1963, **16**, 111; 115 (isol)

Allen, M.S. *et al.*, *Aust. J. Chem.*, 1976, **29**, 673-679 (biosynth)

- Becker, A.M. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 2393-2401 (*abs config*)
 Becker, A.M. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1405
 Otani, T. *et al.*, *J. Antibiot.*, 1989, **42**, 647; 654 (*S 632*)
 Kim, B.S. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 3372-3380 (*isol*)
 Kondo, H. *et al.*, *Eur. J. Org. Chem.*, 2000, 3459-3462 (*synth, pmr, cmr*)

Streptindole S-564

β-1*H*-Indol-3-yl-1*H*-indole-3-ethyl acetate, 9*CI*. 2,2-Di(3-indolyl)ethyl acetate [88321-08-8]




$C_{20}H_{18}N_2O_2$ 318.374
 Metab. from intestinal bacteria (*Streptococcus faecium* IB 37). Sol. MeOH, EtOAc; poorly sol. H_2O . Uncertain low mp. λ_{max} 273 (ϵ 6100); 283 (ϵ 6500); 290 (ϵ 5800) (EtOH) (Derep).

▶ Genotoxic agent.

- Deacetoxy: Vibrindole A**, 1,1-Di(3-indolyl)ethane [5030-91-1]
 $C_{18}H_{16}N_2$ 260.338
 Metab. from the marine bacterium *Vibrio parahaemolyticus*, *isol.* from the toxic mucus of the boxfish *Ostracion cubicus*. Exhibits antimicrobial activity. Red powder. Mp 92°. Also descr. as oil, prob. impure. λ_{max} 241 (ϵ 12200); 275 (ϵ 8100); 282 (ϵ 8160) ($CHCl_3$) (Berdy).
 Osawa, T. *et al.*, *Tet. Lett.*, 1983, **24**, 4719-4722 (*isol, uv, ir, pmr, ms, struct*)
 Hogan, I.T. *et al.*, *Synthesis*, 1984, 872 (*synth, uv, ir, pmr*)
 Singh, H. *et al.*, *Tetrahedron*, 1988, **44**, 5897-5904 (*synth, ir, pmr*)
 Bell, R. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1587-1590 (*Vibrindole A*)
 Chakrabarty, M. *et al.*, *Tet. Lett.*, 2002, **43**, 4075-4078 (*Vibrindole A, synth*)
 Bartoli, G. *et al.*, *Synthesis*, 2004, 895-900 (*synth*)
 Bandagar, B.P. *et al.*, *ARKIVOC*, 2007, **xvi**, 252-259 (*Vibrindole A, synth, ir, pmr*)

Streptolidine S-565

2-Amino-5-(2-amino-1-hydroxyethyl)-4,5-dihydro-1*H*-imidazole-4-carboxylic acid, 9*CI*. Geamine. Roseonine [29307-61-7]



$C_6H_{12}N_4O_3$ 188.186
 Hydrol. prod. of many *Streptomyces* antibiotics, e.g. Geomycin. pK_{a1} 2.5; pK_{a2} 8.72; pK_{a3} 11.3 (H_2O). pK_{a1} 3.95; pK_{a2} 9.1; pK_{a3} 12.65 (66% DMF).


Hydrochloride (1:2): Mp 175-180° dec
 Mp 208-215° dec
 Mp 220°
 Mp 239° dec. $[\alpha]_D^{25} +56$.

Dipicrate:

- Deep yellow prisms (H_2O). Mp 237° dec.
 Carter, H.E. *et al.*, *J.A.C.S.*, 1961, **83**, 4296 (*isol, struct*)
 Borders, D.B. *et al.*, *Tet. Lett.*, 1967, 4187 (*pmr, ir, struct, derivs*)
 Bycroft, B.W. *et al.*, *Chem. Comm.*, 1972, 652 (*cryst struct*)
 Goto, T. *et al.*, *Tet. Lett.*, 1974, 1413 (*synth, ir, ms, nmr*)
 Kusumoto, S. *et al.*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3611 (*pmr, synth*)
 Kinoshita, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1977, **50**, 2375 (*synth*)

Streptolydigin, 9CI S-566

Portamycin. *Afragilimycin A* [7229-50-7]



$C_{32}H_{44}N_2O_9$ 600.708
 Nucleoside antibiotic. Prod. by *Streptomyces lydicus* and *Streptomyces neohydroscopicus afragilimyceticus*. Active against gram-positive bacteria. RNA-polymerase inhibitor. Cryst. + H_2O (Me_2CO). Mp 147-148° Mp 190-195°. $[\alpha]_D^{25} +93$ (c, 1.6 in $CHCl_3$). $[\alpha]_D^{25} -65.7$ (c, 2.28 in 0.005*M* NaOH). λ_{max} 240 (ϵ 8000); 357 (ϵ 35500); 370 (ϵ 33700) (EtOH/HCl) (Derep). λ_{max} 262 (ϵ 13600); 291 (ϵ 16400); 335 (ϵ 20000) (EtOH/NaOH) (Derep). λ_{max} .


▶ WK4250000

Na salt: Mp 225°. $[\alpha]_D^{25} +153$ (c, 1.35 in $CHCl_3$). λ_{max} 240 (ϵ 7280); 360 (ϵ 29400); 373 (ϵ 29500) (50% MeOH/HCl) (Derep). λ_{max} 257 (ϵ 10800); 296 (sh) (ϵ 13100); 310 (sh) (ϵ 13600); 336 (ϵ 17900) (50% MeOH) (Derep). λ_{max} 259 (ϵ 11200); 296 (sh) (ϵ 13700); 336 (ϵ 18500) (MeOH aq./NaOH) (Derep).

[76559-69-8]

- Rinehart, K.L. *et al.*, *J.A.C.S.*, 1963, **85**, 4035; 4037; 4038 (*struct, uv, ir, ms, pmr*)
 Stevens, C.L. *et al.*, *J.A.C.S.*, 1964, **86**, 3592 (*isol*)
 Duchamp, D.J. *et al.*, *J.A.C.S.*, 1973, **95**, 4077 (*cryst struct, stereochem*)
 Von Meyenburg, K. *et al.*, *Antimicrob. Agents Chemother.*, 1978, **13**, 324


- Lee, V.J. *et al.*, *J. Antibiot.*, 1980, **33**, 408 (*cmr*)
 McClure, W.R. *et al.*, *J. Biol. Chem.*, 1980, **255**, 1610 (*props*)
 Japan. Pat., 1980, 80 127 398; CA, **94**, 101308 (*Afragilimycin A*)
 Pearce, C.J. *et al.*, *J. Antibiot.*, 1983, **36**, 1536 (*biosynth*)
 Perez, J.P. *et al.*, *Diss. Abstr. Int.*, B, 1984, **45**, 1781 (*synth*)
 Ireland, R.E. *et al.*, *J.A.C.S.*, 1988, **110**, 854 (*synth*)
 Bols, M. *et al.*, *J. Antibiot.*, 1991, **44**, 678 (*Afragilimycin A*)
 Chen, H. *et al.*, *Org. Lett.*, 2004, **6**, 4033-4036 (*biosynth*)
 Chen, H. *et al.*, *Org. Lett.*, 2006, **8**, 5329-5332 (*biosynth*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, SLW475

Streptomyceamide A S-567

Relative Configuration

$C_{13}H_{16}N_2O_6$ 296.279
 Prod. by *Streptomyces* sp. LR4612. Yellowish powder. $[\alpha]_D^{26} +77$ (c, 1.1 in $CHCl_3$). λ_{max} 242 ($\log \epsilon$ 4.07); 322 ($\log \epsilon$ 3.52); 378 ($\log \epsilon$ 2.4) ($CHCl_3$).

Zhao, P.-J. *et al.*, *Chem. Biodiversity*, 2006, **3**, 337-342 (*isol, pmr, cmr*)

Streptomyceamide B S-568


Relative Configuration

$C_{12}H_{12}N_2O_5$ 264.237
 Prod. by *Streptomyces* sp. LR4612. Amorph. yellow solid. $[\alpha]_D^{25} +45$ (c, 0.2 in MeOH). λ_{max} 224 ($\log \epsilon$ 4.13); 333 ($\log \epsilon$ 3.38) (MeOH).

Zhao, P.-J. *et al.*, *Chem. Biodiversity*, 2006, **3**, 337-342 (*isol, pmr, cmr*)

Streptonigrin, USAN S-569

Rufocromomycin, BAN, INN. Bruneomycin. Nigrin. Valacidin. NSC 45383. RP 5278. Antibiotic RP 5278 [3930-19-6]



Absolute Configuration

$C_{25}H_{22}N_4O_8$ 506.471
 Benzoquinone antibiotic. Absolute configuration conclusively assigned in 2008. *Isol.* from *Streptomyces flocculus* and

other *Streptomyces* spp. Shows immunosuppressant, antineoplastic, anti-HIV and antileukaemic activities. Coffee-brown to almost black rectangular plates (Me₂CO or dioxan) or brown needles. Mp 262–263° (needles) Mp 275° dec. (plates). pK_a 6.2–6.4 (50% dioxan aq.). Log P 1.12 (uncertain value) (calc). λ_{max} 247 (ε 46000); 295 (sh) (ε 17000); 386 (ε 19700) (MeOH) (Derep).

▶ LD₅₀ (dog, orl) 0.5 mg/kg. LD₅₀ (mus, orl) 2.3 mg/kg. Exp. reprod. and teratogenic effects. TJ7350000

N⁷-(1-Methyl-2-oxopropyl): N⁷-(1-Methyl-2-oxopropyl)streptonigrin

[436142-66-4]

C₂₉H₂₈N₄O₉ 576.562

Prod. by *Micromonospora* sp. IM 2670. Red solid. λ_{max} 250 ; 300 (sh) ; 377 (no solvent reported).

6-O-De-Me: 6-Demethylstreptonigrin.

MG 883-12F₂. Antibiotic MG 883-12F₂ [104462-58-0]

C₂₄H₂₀N₄O₈ 492.444

From a *Streptomyces* sp. Active against mouse leukaemia.

10'-O-De-Me: 10'-Demethylstreptonigrin

[105933-56-0]

C₂₄H₂₀N₄O₈ 492.444

From *Streptomyces albus*. Shows antineoplastic and weak antibacterial props. Brown powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 177–181°. λ_{max} 247 (ε 46000); 295 (sh) (ε 17000); 386 (ε 19700) (MeOH) (Derep). λ_{max} 245 (ε 32800); 385 (ε 12300) (MeOH) (Berdy).

10'-Demethoxy: 10'-Demethoxystreptonigrin

[136803-89-9]

C₂₄H₂₀N₄O₇ 476.445

Prod. by *Streptomyces albus* and *Streptomyces otagonensis*. Shows broad spectrum of antibacterial activity. Cytotoxic. Dark red needles (Me₂CO). λ_{max} 247 (ε 41600); 379 (ε 17400) (MeOH/HCl) (Derep). λ_{max} 246 (ε 43600); 380 (ε 16100) (MeOH/NaOH) (Derep). λ_{max} 247 (ε 34800); 379 (ε 14900) (MeOH) (Derep).

▶ US5177500

Rao, K.V. *et al.*, *Antibiot. Annu.*, 1959, 950-953; *CA*, **54**, 17723y (isol)

Rao, K.V. *et al.*, *J. Het. Chem.*, 1975, **12**, 725-730; 731-735 (synth, ir, uv, nmr)

Chiu, Y.Y. *et al.*, *J.A.C.S.*, 1975, **97**, 2525-2530 (cryst struct)

Hibino, S. *et al.*, *Heterocycles*, 1977, **6**, 1485-1507 (rev, synth)

Gould, S.J. *et al.*, *J.A.C.S.*, 1977, **99**, 5496-5497 (biosynth)

Kim, D. *et al.*, *J.O.C.*, 1978, **43**, 121-125 (synth)

Mizuno, N.S. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 372-384 (rev, props)

Gould, S.J. *et al.*, *J.A.C.S.*, 1982, **104**, 343-346 (biosynth)

Weinreb, S.M. *et al.*, *J.A.C.S.*, 1982, **104**, 536-544 (synth)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 1862

Gould, S.J. *et al.*, *Prog. Chem. Org. Nat. Prod.*, 1982, **41**, 77-114 (rev)

Herlt, A.J. *et al.*, *J. Antibiot.*, 1985, **38**, 516-518 (isol, biosynth)

Boger, D.L. *et al.*, *J.A.C.S.*, 1985, **107**, 5745-5754 (synth, bibl)

Eur. Pat., 1986, 185 979; *CA*, **105**, 151549 (MG 883-12F₂)

Isshiki, K. *et al.*, *J. Antibiot.*, 1986, **39**, 1013-1015 (10'-Demethylstreptonigrin)

Si, S. *et al.*, *CA*, 1991, **115**, 202849 (10'-Demethoxystreptonigrin)

Japan. Pat., 1991, 91 83 980; *CA*, **115**, 230526 (10'-Demethoxystreptonigrin)

Liu, W.C. *et al.*, *J. Antibiot.*, 1992, **45**, 454-457 (10'-Demethoxystreptonigrin)

Wang, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 721-724 (N-Methylxopropylstreptonigrin, pmr, cmr)

Anderberg, P.I. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 313-315 (cmr, N-15 nmr)

Bringmann, G. *et al.*, *Tetrahedron*, 2004, **60**, 3539-3574 (rev, synth)

Bringmann, G. *et al.*, *Tetrahedron*, 2008, **64**, 515-521 (cd, abs config)


Bringmann, G. *et al.*, *Tetrahedron*, 2008, **64**, 515-521 (cd, uv, abs config)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, SMA000

Streptonigrone

S-570

[96684-38-7]



C₂₄H₂₂N₄O₇ 478.46

Quinone antibiotic. Opt. inactive, poss. racemic. Prod. by *Streptomyces flocculus* and *Streptomyces albus-bruneomycini*.

Antineoplastic agent. Less active than Streptonigrin, S-569. Rods (CH₂Cl₂/petrol). Mp 268-269°. λ_{max} 425 (ε 12500) (MeOH) (Berdy). λ_{max} 342 (ε 15000) (MeOH/HCl) (Berdy).

Herlt, A.J. *et al.*, *J. Antibiot.*, 1985, **38**, 516 (isol)

Kozlova, N.V. *et al.*, *Antibiot. Khimioter.*, 1990, **35**, 13 (isol)

Boger, D.L. *et al.*, *J.O.C.*, 1991, **56**, 880 (synth)

Tennant, S. *et al.*, *Tetrahedron*, 1997, **53**, 15101-15114 (config)

Chan, B.K. *et al.*, *J.O.C.*, 2007, **72**, 8489-8495 (synth)


Streptopyrrole

S-571

2-Chloro-6,8-dihydroxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI.

Antibiotic XR 587. XR 587

[192819-12-8]



C₁₄H₁₂ClNO₄ 293.706

Prod. by *Streptomyces armeniacus* and *Streptomyces rimosus*. Inhibitor of NR11c protein histidine kinase. Shows

strong antimicrobial activity against gram positive bacteria and weak activity against yeasts. Needles (MeCN). Sol. MeOH, DMSO, CHCl₃; poorly sol. H₂O. Mp 181-183° dec. λ_{max} 241 (log ε 4.51); 294 (log ε 4.3); 335 (log ε 3.65) (MeOH). λ_{max} 254 ; 293 ; 336 (MeOH) (Berdy).

6-Me ether: 2-Chloro-8-hydroxy-6-methoxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI

C₁₅H₁₄ClNO₄ 307.733

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 248 ; 291 ; 341 (MeCN aq.).

1-Chloro: 1,2-Dichloro-6,8-dihydroxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI

[208707-46-4]

C₁₄H₁₁Cl₂NO₄ 328.151

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 247 ; 295 ; 342 (MeCN aq.).

1-Chloro, 6-Me ether: 1,2-Dichloro-8-hydroxy-6-methoxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI

C₁₅H₁₃Cl₂NO₄ 342.177

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 252 ; 291 ; 348 (MeCN aq.).

Bromo analogue: 2-Bromo-6,8-dihydroxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, 9CI

[208707-48-6]

C₁₄H₁₂BrNO₄ 338.157

Prod. by *Streptomyces rimosus*. Powder. λ_{max} 243 ; 295 ; 333 (MeCN aq.).

Breinholt, J. *et al.*, *Acta Chem. Scand.*, 1998, **52**, 1040-1044 (isol, uv, pmr, cmr, ms)

Pat. Coop. Treaty (WIPO), 1998, 98 25 931; *CA*, **129**, 52003e (isol)


Raggatt, M.E. *et al.*, *Chem. Comm.*, 1999, 1039-1040 (biosynth)

Trew, S.I. *et al.*, *J. Antibiot.*, 2000, **53**, 1-11 (isol, uv, ir, pmr, cmr, cryst struct, activity)

Streptorubin B

S-572

[56208-07-2]



C₂₅H₃₃N₃O 391.555

Pyrrole antibiotic. Structure finally shown to be distinctly different to Butylcycloheptylprodigiosine, B-427 in 2007. Prod. by *Streptomyces coelicolor*. λ_{max} 293 (ε 6920); 361 (ε 5250); 510 (ε 37200); 537 (ε 83200) (EtOH/HCl) (Derep).

Tsao, S.-W. *et al.*, *J. Antibiot.*, 1985, **38**, 128-131 (isol)


Laatsch, H. *et al.*, *J. Antibiot.*, 1991, **44**, 187-191 (pmr, cmr)

Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (rev)

Fürstner, A. *et al.*, *Chem. Eur. J.*, 2007, **13**, 1929-1945 (struct)

Streptovercillin S-573

1-(2-Hydroxypropyl)-3,4-dimethoxy-2-methyl-9H-carbazole
[941687-52-1]



Absolute
Configuration

C₁₈H₂₁NO₃ 299.369

Prod. by *Streptomyces morookaense* strain SC1169. Antifungal agent. Amorph. yellow powder. [α]_D²⁰ +18.4 (c, 0.18 in MeOH). λ_{max} 220 (log ε 4.41); 242 (log ε 4.48); 293 (log ε 4.12); 328 (log ε 3.61) (MeOH).

Feng, N. *et al.*, *J. Antibiot.*, 2007, **60**, 179-183 (isol, pmr, cmr, ms)

Streptovitacin D, 8CI S-574


[11035-06-6]

C₁₅H₂₃NO₅ 297.35

Closely related isomer of Streptovitacin A in D-765, full struct. not known. Produced by *Streptomyces griseus*. Shows antitumour activity, phytotoxin. Cryst. Sol. H₂O, EtOAc, MeOH; fairly sol. CHCl₃, Et₂O; poorly sol. hexane. Mp 67-69°. λ_{max} 234 ; 240 (H₂O) (Berdy).

Herr, R.R. *et al.*, *J.A.C.S.*, 1959, **81**, 2595-2596 (struct)

U.S. Pat., 1967, 3 305 554; *CA*, **66**, 98487z (manuf)

Stresgenin B S-575

Relative
Configuration


C₁₁H₁₃NO₅ 239.227

Prod. by *Streptomyces* sp. AS-9. Inhibitor of heat-induced heat shock protein gene expression. Powder. Mp 183-184°. [α]_D²⁵ -12.6 (c, 0.16 in MeOH). λ_{max} 275 (ε 25800) (MeOH).

Akagawa, H. *et al.*, *J. Antibiot.*, 1999, **52**, 960-970

Strictamine S-576

Methyl akuammilan-17-oate, 9CI. *Vinca-midine*. *Desacetyldesfermoakuammiline* [6475-05-4]



Absolute
Configuration

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Vinca minor*, *Vinca libanotica*, *Alstonia scholaris*, *Rhazya stricta* and *Rauwolfia vomitoria* (Apocynaceae). Cryst. (Me₂CO or Et₂O). Mp 80-83° (hydrate) Mp 114-115° (anhyd.) Mp 135°. λ_{max} 213 (log ε 4.37); 262 (log ε 3.8) (EtOH).

▶ LD₅₀ (mus, ipr) 162 mg/kg. WK7150000

N⁴-Oxide: **Strictamine N-oxide**

[90850-95-6]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from the leaves of *Rauwolfia stricta* (Apocynaceae). Amorph. [α]_D -75 (CHCl₃). λ_{max} 213 (log ε 3.15); 262 (log ε 2.66) (MeOH).

N⁴-Me: **N^b-Methylstrictamine**

[123789-23-1]

C₂₁H₂₅N₂O₂⁺ 337.441

Quaternary alkaloid from the leaves of *Rauwolfia stricta* and *Vinca minor* (Apocynaceae). Mp 176-178°. [α]_D -65 (CHCl₃). Counterion not specified. CAS no. refers to chloride. λ_{max} 242 (MeOH).

17-Aldehyde: **Strictalamine**

[1897-36-5]

C₁₉H₂₀N₂O 292.38

Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Needles. Mp 152-154°.

10-Hydroxy: **10-Hydroxystriactamine**

[88754-97-6]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Alstonia macrophylla* and *Vinca erecta*. Amorph. [α]_D +81.6 (c, 0.02 in CHCl₃). λ_{max} 230 (log ε 3.6); 280 (log ε 3.4) (MeOH).

11-Hydroxy: **11-Hydroxystriactamine**

[88754-98-7]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from *Vinca erecta* and *Alstonia angustifolia* (Apocynaceae). Mp 228-229°. λ_{max} 233 (log ε 4.25); 285 (log ε 3.64) (EtOH).

12-Hydroxy: **12-Hydroxystriactamine**

[136952-70-0]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from leaves of *Rauwolfia sumatrana* (Apocynaceae). Feathery needles (MeOH). Mp 282-286° dec. λ_{max} 259 (ε 5300); 297 (sh) (ε 2500) (MeOH).

12-Hydroxy, Ac:

Stout rods (EtOAc/petrol). Mp 177-181°.

18-Hydroxy: **Gomaline**. 18-Hydroxystriactamine

[89384-07-6]

C₂₀H₂₂N₂O₃ 338.405

Alkaloid from leaves of *Catharanthus roseus* (Apocynaceae). [α]_D +94 (CHCl₃). λ_{max} 210 ; 262 (MeOH).

5β-Methoxy: **5-Methoxystriactamine**

[870995-64-5]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the leaves of

Alstonia scholaris. Powder. Mp 232-235°. [α]_D¹⁴ -119.3 (c, 0.39 in CHCl₃).

11-Methoxy: **11-Methoxystriactamine**

[160920-59-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from leaves of *Rauwolfia sumatrana* (Apocynaceae). Amorph. powder. λ_{max} 229 (4.08); 279 (3.4) (MeOH).

5α,10-Dimethoxy: **5,10-Dimethoxystriactamine**

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from *Alstonia villosa*. Solid. [α]_D²⁵ -71.8 (c, 1.23 in MeOH). Incorrectly indexed in CAS. λ_{max} 207 (log ε 4.31); 220 (sh) (log ε 4.24); 285 (log ε 3.87) (MeOH).

5α,10,11-Trimethoxy: **5,10,11-Trimethoxystriactamine**

C₂₃H₂₈N₂O₅ 412.485

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. [α]_D²² -73.4 (c, 1.30 in MeOH).

16-Epimer, 17-alcohol: **Rhazinol**

[72778-18-8]

C₁₉H₂₂N₂O 294.396

Alkaloid from *Rhazya stricta* (Apocynaceae). Glass. λ_{max} 225 (ε 13610); 266 (ε 5440) (MeOH).

16-Epimer, 17-aldehyde: **Rhazinolone**

[949932-16-5]

C₁₉H₂₀N₂O 292.38

Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D +136 (c, 0.07 in CHCl₃). λ_{max} 211 (log ε 4.27); 260 (log ε 3.7) (EtOH).

Trojánek, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1960, **25**, 2045-2048 (isol, uv, ir)

Schnoes, H.K. *et al.*, *J.O.C.*, 1966, **31**, 1641-1642 (uv, ir, pmr, ms, struct)

Taylor, W.I. *et al.*, *The Vinca Alkaloids*, Marcel Dekker, 1973, (props)

Ahmad, Y. *et al.*, *J.A.C.S.*, 1977, **99**, 1943-1946 (uv, ir, pmr, ms, cryst struct, abs config)

Yagudaev, M.R. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 454-458 (10-Hydroxystriactamine, 11-Hydroxystriactamine)

Ahmad, Y. *et al.*, *Phytochemistry*, 1983, **22**, 1017-1019 (Rhazinol)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1984, **22**, 85-86 (Gomaline)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1984, **23**, 709-710 (Strictamine N-oxide)

Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 1987, **42**, 91-93 (N^b-Methylstrictamine)

Ghedira, K. *et al.*, *Phytochemistry*, 1988, **27**, 3955-3962 (11-Hydroxystriactamine)

Uhrin, D. *et al.*, *J. Nat. Prod.*, 1989, **52**, 637-639 (N^b-Methylstrictamine)

Proksa, B. *et al.*, *Planta Med.*, 1989, **55**, 188-190 (N^b-Methylstrictamine)

Arbain, D. *et al.*, *Aust. J. Chem.*, 1991, **44**, 1007-1011 (12-Hydroxystriactamine)


Subhadhirasakul, S. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 1427-1431 (11-Methoxystriactamine)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1994, **5**, 201-209 (10-Hydroxystriactamine)

Abe, F. *et al.*, *Phytochemistry*, 1994, **35**, 253-257 (5,10,11-Trimethoxystriactamine)

- Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1235-1238 (5,10-Dimethoxystrictamine)
 Zhou, H. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 2508-2512 (5-Methoxystrictamine)
 Lim, K.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*Rhazinoline*)


Strictaminolamine S-577
 [119328-84-6]



Absolute Configuration

- $C_{21}H_{26}N_2O_3$ 354.448
 Alkaloid from the leaves of *Alstonia macrophylla* (Apocynaceae). Amorph. $[\alpha]_D -30$ (CHCl₃).
16-Epimer, 10-methoxy, N⁴-oxide: Alstozine N-oxide
 [134328-12-4]
 $C_{22}H_{28}N_2O_5$ 400.474
 Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Amorph. solid. $[\alpha]_D +107$ (CHCl₃).
 Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1988, **27**, 3653 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Atta-ur-Rahman, *et al.*, *Fitoterapia*, 1990, **61**, 230 (*Alstozine N-oxide*)


Strictanine S-578
 [109028-31-1]



Absolute Configuration


- $C_{20}H_{26}N_2O_2$ 326.438
 Trace alkaloid from the fruits of *Rhazya stricta* (Apocynaceae). λ_{max} 212 ; 253 ; 260 (sh) ; 280 ; 290 (MeOH).
11,12-Dihydroxy, N-deformyl, N-Ac: Spegazzinidine
 [6899-72-5]
 $C_{21}H_{28}N_2O_4$ 372.463
 Alkaloid from *Aspidosperma chakensis* (Apocynaceae). Cryst. Mp 237-238°. $[\alpha]_D^{24} +186$ (CHCl₃).
12-Hydroxy, N-deformyl, N-Ac: Spegazzinine
 [6882-72-0]
 $C_{21}H_{28}N_2O_3$ 356.464
 Alkaloid from *Aspidosperma chakensis* (Apocynaceae). Cryst. (MeOH aq.). Mp 104.5-106°. $[\alpha]_D +175.6$ (CHCl₃).
 Djerassi, C. *et al.*, *J.A.C.S.*, 1962, **84**, 3480 (*Spegazzinine, Spigazzinidine*)
 Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1987, **26**, 589-591 (*Strictanine*)
 Belferdi, F. *et al.*, *Molecules*, 2001, **6**, 803-814 (*synth*)

Strictibine S-579
 [183803-82-9]




- $C_{13}H_{11}NO_2$ 213.235
 Alkaloid from leaves of *Rhazya stricta*. λ_{max} 208 (log ϵ 3.9); 237 (log ϵ 4.13); 286 (log ϵ 2.74); 375 (log ϵ 2.25) (MeOH).
 Habib-ur-Rehman, *et al.*, *Fitoterapia*, 1996, **67**, 145 (*isol, uv, ir, pmr, ms, struct*)

Stricticine S-580
 [72847-49-5]




- $C_{20}H_{22}N_2O_3$ 338.405
 Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Amorph. $[\alpha]_D +515$ (CHCl₃). Related to 20-Epilochneridine; see L-224. λ_{max} 208; 228; 292; 327 (MeOH) (Derep).
 Ahmad, Y. *et al.*, *Phytochemistry*, 1983, **22**, 1017-1019 (*isol*)
 Atta-ur-Rahman, *et al.*, *Tet. Lett.*, 1987, **28**, 3609 (*uv, ir, pmr, cmr, ms, struct*)

Strictimine S-581
1,1'-Carbonylbis[3-ethylpiperidine], 9CI
 [92631-62-4]



- $C_{15}H_{28}N_2O$ 252.399
 Alkaloid from the roots of *Rhazya stricta*. Racemic.
 Atta-ur-Rahman, *et al.*, *Heterocycles*, 1984, **22**, 2023-2027


Strictine S-582
 [72847-50-8]



Absolute Configuration

- $C_{20}H_{20}N_2O_3$ 336.39
 Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). Amorph.
 Atta-ur-Rahman, *et al.*, *Heterocycles*, 1987, **26**, 2125 (*isol, uv, ir, pmr, cmr, ms, struct*)

Strictosidinic acid S-583
 [150148-81-5]



Absolute Configuration

- $C_{26}H_{32}N_2O_9$ 516.547
 Most refs. refer to Strictosidine. Alkaloid from the roots of *Rhazya orientalis* (Apocynaceae) and leaves of *Ophiorrhiza filistipula* (Rubiaceae). Feathery needles (MeOH/EtOAc). Mp 253-255° dec. $[\alpha]_D -223$ (c, 0.19 in H₂O).
3'- α -D-Glucopyranosyl: Hunterioside B
 [168434-16-0]
 $C_{32}H_{42}N_2O_{14}$ 678.689
 Alkaloid from the stem bark of *Hunteria zeylanica* (Apocynaceae). Amorph. solid. λ_{max} 221 ; 272 ; 289 (EtOH).
6'- α -D-Glucopyranosyl: Hunterioside. Strictosidinic acid 6'- α -D-glucoiside
 [156431-08-2]
 $C_{32}H_{42}N_2O_{14}$ 678.689
 Alkaloid from stem bark of *Hunteria zeylanica* (Apocynaceae). Amorph. powder. The first biose bound mono-terpenoid indole alkaloid.
Me ester: Strictosidine. Isovincoside
 [20824-29-7]
 $C_{27}H_{34}N_2O_9$ 530.574
 Alkaloid from the leaves of *Rhazya stricta* and *Anthocephalus cadamba*, *Catharanthus roseus* (*Vinca rosea*) and *Strychnos mellodora* (Apocynaceae, Rubiaceae). Key intermediate in the biosynth. of terpenoid indole alkaloids. Noncryst. $[\alpha]_D -143$ (MeOH) (as hydrochloride). Very reactive compound; hydrol. to Vallesiachotamine, V-10.
N⁴-Ac, Me ester:
 Needles (MeOH aq.). Mp 172-173°. $[\alpha]_D -107$ (MeOH).
N⁴-Me: Palicoside
 [123828-68-2]
 $C_{27}H_{34}N_2O_9$ 530.574
 Alkaloid from leaves of *Palicourea marcgravi* (Rubiaceae) and from *Strychnos mellodora*. Yellowish needles. Mp 206-208°. $[\alpha]_D -61.6$ (c, 0.5 in H₂O).
N⁴-Me, Me ester: Dolichantoside
 [68727-52-6]
 $C_{28}H_{36}N_2O_9$ 544.6
 Alkaloid from roots of *Strychnos gossweileri* and stem bark of *Strychnos tricalysoides* and from *Strychnos mellodora* (Loganiaceae). Noncryst. C-3 config. revised to R- in 1993 but reversed again in 1997.
3,4-Didehydro: 3,4-Dehydrostrictosidinic acid
 $C_{26}H_{30}N_2O_9$ 514.531
 Alkaloid from *Chimarrhis turbinata*. Amorph. yellow powder. λ_{max} 240 (log

- ϵ 3); 356 (log ϵ 3.5) (MeOH).
3,4-Didehydro, Me ester: 3,4-Dehydros-strictosidine
 $C_{27}H_{32}N_2O_9$ 528.558
 Alkaloid from *Chimarrhis turbinata*.
 Amorph. yellow powder. λ_{\max} 239 (log ϵ 3); 358 (log ϵ 3.9) (MeOH).
5-Oxo, Me ester: 5-Oxostrictosidine
 [56470-42-9]
 $C_{27}H_{32}N_2O_{10}$ 544.557
 Alkaloid from the heartwood of *Adina rubescens* (Rubiaceae). Amorph. solid (as N^4, O, O, O, O -penta-Ac). $[\alpha]_D^{25}$ -20 (CHCl₃) (penta-Ac).
3-Epimer: 3-Epistrictosidinic acid. Vinso-sidinic acid
 $C_{26}H_{32}N_2O_9$ 516.547
 Alkaloid from the leaves of *Palicourea coriacea*. Brown cryst. Mp 190-194°. $[\alpha]_D$ -72 (c, 8.2 in MeOH).
3-Epimer, Me ester: Vincoside
 [19456-89-4]
 $C_{27}H_{34}N_2O_9$ 530.574
 Alkaloid from *Catharanthus roseus* (Apocynaceae). Noncryst. $[\alpha]_D$ -112 (MeOH) (as hydrochloride). Unstable.
3-Epimer, N⁴-Ac, Me ester: N-Acetylvincoside
 $C_{29}H_{36}N_2O_{10}$ 572.611
 Alkaloid from *Catharanthus roseus* (Apocynaceae). Needles (EtOH aq.). Mp 180-181°. $[\alpha]_D$ -170 (MeOH).
3-Epimer, N⁴-Me, Me ester: Isodolichantoxide
 [75363-97-2]
 $C_{28}H_{36}N_2O_9$ 544.6
 Minor alkaloid from root bark of *Strychnos gossweileri* (Loganiaceae). C-3 config. revised to *S*- in 1993 but reversed again in 1997.
 Smith, G.N. *et al.*, *Chem. Comm.*, 1968, 912 (*isol, uv, ir, pmr, ms*)
 Battersby, A.R. *et al.*, *J.C.S.(C)*, 1969, 1193 (*isol, uv, ir, pmr*)
 De Silva, K.T.D. *et al.*, *Chem. Comm.*, 1971, 905; 908 (*Strictosidinic acid, config*)
 Blackstock, W.P. *et al.*, *Chem. Comm.*, 1971, 910 (*abs config*)
 Brown, R.T. *et al.*, *Experientia*, 1975, **31**, 505 (*5-Oxostrictosidine*)
 Mattes, K.C. *et al.*, *J.A.C.S.*, 1975, **97**, 6270 (*abs config*)
 Coune, C. *et al.*, *Planta Med.*, 1978, **34**, 53 (*Dolichantoxide*)
 Ruffer, M. *et al.*, *Tet. Lett.*, 1978, 1593 (*biosynth*)
 Stöckigt, J. *et al.*, *Phytochemistry*, 1979, **18**, 965 (*biosynth*)
 Coune, C.A. *et al.*, *Herba Hung.*, 1980, **19**, 189; *CA*, **93**, 217906e (*Isodolichantoxide*)
 Leclercq, J. *et al.*, *Planta Med.*, 1984, **50**, 457 (*Dolichantoxide*)
 Morita, H. *et al.*, *Planta Med.*, 1989, **55**, 288 (*Palicoside*)
 Subhadhirasakul, S. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 991 (*Hunterioside*)
 Tits, M. *et al.*, *Planta Med.*, 1996, **62**, 73-74 (*isol, derivs, Strychnos*)
 Pathy-Lukáts, A. *et al.*, *J. Nat. Prod.*, 1997, **60**, 69 (*pmr, cmr, stereochem*)
 Achenbach, H. *et al.*, *Phytochemistry*, 1997, **44**, 1387-1390 (*Dolichantoxide, Isodolichantoxide, abs config*)
 Takayama, H. *et al.*, *Heterocycles*, 1998, **47**, 87-90 (*Hunteriosides*)


- Ohmori, O. *et al.*, *Tet. Lett.*, 1998, **39**, 7737-7740 (*Dolichantoxide, Isodolichantoxide*)
 Pathy-Lukáts, A. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1492-1499 (*conformn, config*)
 Ohmori, O. *et al.*, *Tet. Lett.*, 1999, **40**, 5039-5042 (*Hunterioside, synth*)
 Cardoso, C.L. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1882-1885 (*Dehydrostrictosidine, Dehydrostrictosidinic acid*)
 Do Nascimento, C.A. *et al.*, *Z. Naturforsch., B*, 2006, **61**, 1443-1446 (*Epistrictosidinic acid*)
 Szabó, L.F. *et al.*, *Molecules*, 2008, **13**, 1875-1896 (*rev*)

Strictosidinic ketone S-584

- [929686-15-7]
 As Strictosidinic acid, S-583 with
 $R = COCH_3$
 $C_{27}H_{34}N_2O_8$ 514.574
 Alkaloid from the leaves of *Palicourea coriacea*. Yellowish cryst. $[\alpha]_D$ -60 (c, 10 in MeOH).
 Do Nascimento, C.A. *et al.*, *Z. Naturforsch., B*, 2006, **61**, 1443-1446

Strigosine S-585

- Trachelanthamidine strigosate*
 [33981-76-9]




Absolute configuration

- $C_{14}H_{25}NO_4$ 271.356
 Ester of Trachelanthamidine in H-629 with Strigosic acid. Major alkaloid from *Heliotropium strigosum* (Boraginaceae). Gum. $[\alpha]_D^{20}$ -19.3 (c, 6.56 in EtOH).
Hydrochloride:
 Needles (EtOH/Et₂O). Mp 137.5°.
- Picrate:*
 Yellow leaflets (C₆H₆/petrol). Mp 141°.
- Methiodide:*
 Leaflets (EtOH/Et₂O). Mp 135-136°. $[\alpha]_D$ -15.5 (c, 2.84 in EtOH).
 Mattocks, A.R. *et al.*, *J.C.S.*, 1964, 1974 (*isol, struct*)
 Crout, D.H.G. *et al.*, *J.C.S. Perkin 1*, 1977, 544 (*abs config*)

Strobamine S-586

- 2,3,6,7,8,9-Hexahydro-10-methyl-2-phenylcyclohepta[b]pyran-5,8-imin-4(5H)-one, 9CI**
 [75656-91-6]




- $C_{17}H_{19}NO_2$ 269.343
 Rel. config. only detd. Alkaloid from the leaves of *Knightsia strobilina* (Proteaceae). Viscous oil. $[\alpha]_D^{20}$ +36 (CHCl₃).
7-Hydroxy: Strobolamine
 [75638-69-6]
 $C_{17}H_{19}NO_3$ 285.342

Alkaloid from the leaves of *Knightsia strobilina* (Proteaceae). Cryst. (Me₂CO). Mp 159-161°. $[\alpha]_D^{20}$ +34 (CHCl₃).

- Lounasmaa, M. *et al.*, *Phytochemistry*, 1980, **19**, 953 (*uv, ir, pmr, ms, struct, deriv*)
 Lounasmaa, M. *et al.*, *J. Nat. Prod.*, 1983, **46**, 429 (*synth*)

Strobiline S-587

6,7,8,9-Tetrahydro-10-methylcyclohepta[b]pyran-5,8-imin-4(5H)-one, 9CI
 [77053-09-9]



- $C_{11}H_{13}NO_2$ 191.229
 Rel. config. only established. Alkaloid from the leaves of *Knightsia strobilina* (Proteaceae). Cryst. (petrol). Mp 108-109°. $[\alpha]_D^{20}$ +60 (CHCl₃).

10,11-Dihydro: Dihydrostrobiline
 $C_{11}H_{15}NO_2$ 193.245
 Alkaloid from the leaves of *Knightsia strobilina* (Proteaceae). Viscous oil. $[\alpha]_D^{20}$ -30 (CHCl₃). Incorrectly labelled 9,10-dihydro in the lit.


- Lounasmaa, M. *et al.*, *Phytochemistry*, 1980, **19**, 949 (*isol, uv, ir, pmr, cmr, ms, struct, deriv*)

Strychnambarine S-588

- [39453-52-6]
 Struct. unknown. Minor alkaloid from *Strychnos usambarensis* (Loganiaceae). Conts. 3N. λ_{\max} 227; 268; 283; 290 (no solvent reported).
 Koch, M. *et al.*, *Ann. Pharm. Fr.*, 1973, **31**, 45-48 (*isol, ms*)

Strychnine, 8CI, BSI, ISO S-589

Strychnidin-10-one, 9CI
 [57-24-9]



Absolute configuration

- $C_{21}H_{22}N_2O_2$ 334.417
 Alkaloid from the seeds of *Strychnos nux-vomica*, from *Strychnos wallichiana* and from very many other *Strychnos* spp. Bark of *Strychnos icaja* is richest known source (6.6%) (Loganiaceae). Violent tetanic poison, employed comly. in vermin killers. Small doses formerly used in tonics. Use as animal poison prohibited in UK except for destruction of moles and other burrowing animals, and is being phased out in most countries. Pharmaceutical use essentially obsolete although still features in homeopathic medicines. Important tool in neurophysiological research. CNS stimulant.

- Convulsant. Cholinesterase inhibitor. Used as 80% aq. soln. for photometric detn. of PO_4^{3-} , NO_2^- , NO_3^- . Potent CNS glycine antagonist. Mp 270-271° (slow heat) Mp 275-285° Mp 286-288°. $[\alpha]_D^{18}$ -139 (CHCl₃). $\text{p}K_{a1}$ 8; $\text{p}K_{a2}$ 2.3 (20°). Log P -0.31 (uncertain value) (calc). Readily forms crystalline chloromethochloride artifacts when CHCl₃ or CH₂Cl₂ is used in isolan.
- ▶ Very toxic by ingestion and other routes of administration. Lethal human dose 30-60 mg. OES: long-term 0.15 mg m⁻³; short-term 0.45 mg m⁻³. LD₅₀ (rat, orl) 2.35 mg/kg. Exp. reprod. effects. WL2275000
- Sulfate (2:1): Strynervene. Strychnine sulfate. Vampirol*
[60-41-3]
Cryst. + 5H₂O. Mp 200° (anhyd.).
- ▶ LD₅₀ (rat, orl) 5 mg/kg. Very toxic by ingestion and other routes of administration. WL2550000
- Nitrate:* [66-32-0]
Needles.
- ▶ LD₅₀ (rat, orl) 16.2 mg/kg. WL2450000
- Picrate:* Mp 270° dec.
- N⁴-Oxide: Strychnine N-oxide. Genos-trychnine. Movellan. Stricnogen. Z 203**
[7248-28-4]
C₂₁H₂₂N₂O₃ 350.416
Alkaloid from *Strychnos wallichiana* and *Strychnos nux-vomica* leaves (Loganiaceae). Log P -2.62 (uncertain value) (calc). Is formed as an artifact when CHCl₃ or CH₂Cl₂ solns. of strychnine are allowed to stand. Appears to be a genuine alkaloid, however.
 - ▶ LD₅₀ (mus, scu) 24.5 mg/kg. WL2460300

N⁴-Me: N⁴-Methylstrychninium
[47466-17-1]
C₂₂H₂₅N₂O₂[⊕] 349.452
Quaternary alkaloid from *Strychnos icaja*. λ_{max} 253 (log ϵ 4.31); 280 (log ϵ 3.89); 289 (log ϵ 3.76) (no solvent reported).

Deoxo: Strychnidine
[2054-81-1]
Needles (EtOH). Mp 258° (246-248°). Subl. 150. $[\alpha]_D$ -8.28 (CHCl₃). $[\alpha]_D^{19}$ -63 (c, 1.5 in CHCl₃). Not reported as a natural alkaloid.

3-Hydroxy: see Pseudostrychnine, P-726

10-Hydroxy: 10-Hydroxystrychnine
[19894-24-7]
C₂₁H₂₂N₂O₃ 350.416
Alkaloid in root bark of *Strychnos nux-vomica* (Sri Lankan, poss. hybrid). Metab. of strychnine *in vivo* (Loganiaceae). Identified spectroscopically. λ_{max} 260 ; 299 ; 310 (sh) (MeOH).

 - ▶ WL2330000

12-Hydroxy: 12-Hydroxystrychnine
[22595-26-2]
C₂₁H₂₂N₂O₃ 350.416
Alkaloid from *Strychnos icaja*, seeds of *Strychnos wallichiana* and fruits of *Strychnos nux-vomica* (Loganiaceae). Convulsant. Cryst. (EtOH). $[\alpha]_D^{20}$ -8 (c, 0.7 in CHCl₃).
 - ▶ WL2340000

12-Hydroxy, N⁴-oxide: 12-Hydroxy-strychnine N^b-oxide
[83905-59-3]
C₂₁H₂₂N₂O₄ 366.416
Alkaloid from leaves of *Strychnos nux-vomica* (Sri Lankan, poss. hybrid) (Loganiaceae). λ_{max} 264 ; 295 (MeOH).

15-Hydroxy: 15-Hydroxystrychnine
[72994-79-7]
C₂₁H₂₂N₂O₃ 350.416
Alkaloid from seeds of *Strychnos nux-vomica* (Loganiaceae). Cryst. (EtOAc). Mp 204-206°. $[\alpha]_D^{20}$ -192.7 (c, 0.4 in CHCl₃). λ_{max} 255 (log ϵ 4.08); 280 (log ϵ 3.6); 291 (log ϵ 3.49) (EtOH).

10-Methoxy: β -Colubrine. 10-Methoxystrychnine
[509-36-4]
C₂₂H₂₄N₂O₃ 364.443
Alkaloid from *Strychnos nux-vomica* and *Strychnos ligustrina* (Loganiaceae). Cryst. (EtOH aq.). Mp 222°. $[\alpha]_D^{21}$ -156 (c, 0.042 in CHCl₃). $[\alpha]_D^{19}$ -107.7 (c, 2.47 in EtOH).
 - ▶ GJ9990000

10-Methoxy; hydrochloride:
Plates + 1H₂O. $[\alpha]_D^{18}$ -32.7 (H₂O).

10-Methoxy, N⁴-oxide: β -Colubrine N-oxide
[19442-97-8]
C₂₂H₂₄N₂O₄ 380.443
Alkaloid from *Strychnos ligustrina* (Loganiaceae). Mp 205°. $[\alpha]_D^{17}$ +6.6 (c, 0.2 in EtOH).

11-Methoxy: α -Colubrine. 11-Methoxystrychnine
[509-44-4]
[27137-63-9 , 29135-26-0]
C₂₂H₂₄N₂O₃ 364.443
Alkaloid from *Strychnos nux-vomica* (Loganiaceae). Cryst. + 4H₂O (EtOH aq.). Mp 184°. $[\alpha]_D^{19}$ -76.5 (c, 2.025 in EtOH).
 - ▶ GJ9980000

11-Methoxy; hydrochloride:
Plates + 3H₂O (H₂O). $[\alpha]_D^{18}$ -3.1 (c, 1.943 in EtOH).

11-Methoxy, 10-hydroxy: 10-O-De-methylbrucine. 10-Hydroxy-11-methoxystrychnine. 10-Hydroxy- α -colubrine
[1103-85-1]
C₂₂H₂₄N₂O₄ 380.443
Alkaloid from the seeds of *Strychnos nux-vomica*. Yellow-brown cryst. (EtOH aq.). Dec. at 245°. λ_{max} 208 ; 264 ; 303 (MeOH).

11-Methoxy, 12-hydroxy: 12-Hydroxy-11-methoxystrychnine
[50837-32-6]
C₂₂H₂₄N₂O₄ 380.443
Alkaloid from seeds of *Strychnos wallichiana* (*Strychnos cinnamomifolia*) also detected in root bark and leaves of *Strychnos nux-vomica* (Loganiaceae). Mp 263-264° dec. $[\alpha]_D$ -18 (c, 0.22 in CHCl₃). λ_{max} 231 ; 270 ; 301 (EtOH).

11-Methoxy, 12-hydroxy, N⁴-oxide: 12-Hydroxy-11-methoxystrychnine N^b-oxide
[83905-57-1]
C₂₂H₂₄N₂O₅ 396.442
Alkaloid from leaves of *Strychnos nux-vomica* (Sri Lankan, poss. hybrid) (Loganiaceae). Plates (MeOH). $[\alpha]_D^{21}$ +91 (c, 0.175 in MeOH). λ_{max} 237 ; 268 (MeOH).

10,11-Dimethoxy: Brucine. 10,11-Dimethoxystrychnine
[357-57-3]
[5892-11-5]
C₂₃H₂₆N₂O₄ 394.469
Alkaloid from *Strychnos nux-vomica* and many other *Strychnos* spp. (Loganiaceae). Resolving agent. Used in photometric detn. of ClO_3^- , NO_3^- , NO_2^- , NO_2 . Antineoplastic agent. A const. of the herbal and homoeopathic medicine nux vomica. Cryst. powder. Sol. EtOH, CHCl₃; sl. sol. C₆H₆, Et₂O. Mp 105° (hydrate) Mp 78° (anhyd.). $[\alpha]_D$ -127 (CHCl₃). $[\alpha]_D$ -85 (EtOH). $\text{p}K_{a1}$ 8.16; $\text{p}K_{a2}$ 2.5 (15°). Log P -0.85 (uncertain value) (calc).
 - ▶ Reported to have one-sixth toxicity of strychnine. LD₅₀ (rat, ipr) 91 mg/kg. EH8925000

10,11-Dimethoxy; sulfate: [4845-99-2]
[5787-00-8 , 60583-39-3]
Mp 186° dec. (as monohydrate). $[\alpha]_D^{22}$ -24 (c, 1 in H₂O).

10,11-Dimethoxy; perchlorate: Mp 208-210°.

10,11-Dimethoxy, N⁴-oxide: Brucine N-oxide
[17301-81-4]
C₂₃H₂₆N₂O₅ 410.469
Alkaloid from *Strychnos wallichiana* and *Strychnos ligustrina*, also detected in *Strychnos nux-vomica* leaves (Loganiaceae).

10,11-Dimethoxy, N⁴-Me: Mp 295° dec. Δ^{20} -Isomer: Neostrychnine
[466-69-3]
C₂₁H₂₂N₂O₂ 334.417
Semisynthetic. Mp 227-229.5°.

[116708-31-7 , 126590-09-8 , 116728-44-0]
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 1066B (ir)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 1067C; 1067D (ir)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 613A (nmr)
Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 614A; 614B (nmr)
Pelletier, J. et al., *Ann. Chim. (Paris)*, 1819, **10**, 142-177; 1820, **12**, 113-133 (*Strychnine, Brucine, isol*)
Warnat, K. et al., *Helv. Chim. Acta*, 1931, **14**, 997-1007 (α -Colubrine, β -Colubrine)
Woodward, R.B. et al., *J.A.C.S.*, 1948, **70**, 2107-2115 (*struct, bibl*)
Welcher, F.J. et al., *Organic Analytical Reagents*, Van Nostrand, NY, 1948, **4**, 259 (*use*)
Holmes, H.L. et al., *Alkaloids (Academic Press)*, 1949, **1**, 375-500; 1952, **2**, 513-551 (*rev, bibl*)
Kenner, G.W. et al., *J.C.S.*, 1950, 406 (*Strychnidine, synth*)
Robinson, R. et al., *Prog. Org. Chem.*, 1952, **1**, 1-21 (*rev, struct*)
Eger, C. et al., *Anal. Chem.*, 1955, **27**, 1199-1200 (*detn, ClO₃⁻*)

Peerdeman, A.F. *et al.*, *Acta Cryst.*, 1956, **9**, 824 (*cryst struct*)

Hendrickson, J.B. *et al.*, *Alkaloids (Academic Press)*, 1960, **6**, 179-217 (*rev, bibl*)

Rosenmund, P. *et al.*, *Chem. Ber.*, 1962, **95**, 2639-2644 (*struct, synth, biosynth*)

Woodward, R.B. *et al.*, *Tetrahedron*, 1963, **19**, 247-288 (*synth*)

Jenkins, D. *et al.*, *Anal. Chem.*, 1964, **36**, 610-612 (*detn, NO₂[±]*)

Rosenmund, P. *et al.*, *Chem. Ber.*, 1964, **97**, 1677-1683 (*α-Colubrine, synth*)

Szabo, L. *et al.*, *CA*, 1965, **63**, 7064a (*Strychnidine, synth*)

Hesse, M. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 674-689 (*ms*)

Smith, G.A. *et al.*, *Analyst (London)*, 1967, **92**, 456-462 (*detn, NO₂*)

Sandberg, F. *et al.*, *Tet. Lett.*, 1968, 6217-6218 (*12-Hydroxystrychnine*)

Tedeschi, E. *et al.*, *Tetrahedron*, 1968, **24**, 4573-4580 (*Brucine, β-Colubrine, synth*)

Fadrus, H. *et al.*, *Fresenius' Z. Anal. Chem.*, 1969, **246**, 239-241 (*detn, NO₂[±]*)

Phillipson, J.D. *et al.*, *Phytochemistry*, 1972, **11**, 2547-2553 (*chloromethochlorides*)

Heimberger, S.I. *et al.*, *Chem. Comm.*, 1973, 217-218 (*biosynth*)

Bisset, N.G. *et al.*, *J. Pharm. Pharmacol.*, 1973, **25**, 563-569 (*12-Hydroxy-11-methoxystrychnine*)

Bisset, N.G. *et al.*, *Phytochemistry*, 1974, **13**, 259-263 (*Strychnine N-oxide*)

Bose, A.K. *et al.*, *Anal. Biochem.*, 1978, **89**, 284-291 (*ms*)

Snow, J.W. *et al.*, *Can. J. Chem.*, 1978, **56**, 1222-1230 (*uv, cd*)

Wenkert, E. *et al.*, *J.O.C.S.*, 1978, **43**, 1099-1105 (*pmr, cmr*)

Kambu, K. *et al.*, *Planta Med.*, 1979, **37**, 161-164 (*N⁴-Methylstrychninium*)

Galeffi, C. *et al.*, *Tetrahedron*, 1979, **35**, 2545-2549 (*15-Hydroxystrychnine*)

Baser, K.H.C. *et al.*, *Phytochemistry*, 1982, **21**, 1423-1429 (*10-Hydroxystrychnine, 12-Hydroxystrychnine N-oxide, 12-Hydroxy-11-methoxystrychnine N-oxide*)

Asai, F. *et al.*, *Yakugaku Zasshi*, 1982, **102**, 690-694 (*Strychnine N-oxide*)

Chazin, W.J. *et al.*, *Can. J. Chem.*, 1983, **61**, 1749-1755 (*pmr*)

Goldinger, A. *et al.*, *Med. Monatsschr.*, 1983, **6**, 141-146 (*rev, pharmacol*)

Fieser and Fieser's Reagents for Organic Synthesis, Wiley, 1984, **11**, 83-87 (*Brucine, use*)

Verpoorte, R. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 345-348 (*cmr, 10-O-Demethylbrucine*)

Mostad, A. *et al.*, *Acta Chem. Scand., Ser. B*, 1985, **39**, 705-716 (*cryst struct*)

Glover, S.S.C. *et al.*, *Acta Cryst. C*, 1985, **41**, 990-994 (*Brucine, cryst struct*)

Bernstein, M.A. *et al.*, *Can. J. Chem.*, 1985, **63**, 483-490 (*pmr*)

Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1986, **15**, 563-646 (*rev, bibl*)

Gould, R.O. *et al.*, *Acta Cryst. C*, 1987, **43**, 2405-2410 (*cryst struct*)

Dangerous Prop. Ind. Mater. Rep., 1988, **8**, 78-83 (*tox, rev*)

Ghosh, R. *et al.*, *Acta Cryst. C*, 1989, **45**, 1794-1797 (*cryst struct*)

Glycine Neurotransm. 1990, (Eds. Ottersen, O.P. *et al.*), J. Wiley, Chichester, 1990, (*book*)

Smith, B.A. *et al.*, *J. Emergency Med.*, 1990, **8**, 321 (*tox, rev*)

Agrochemicals Handbook, 3rd edn., Royal Society of Chemistry, 1992, A729

Nicholson, J.W. *et al.*, *Educ. Chem.*, 1993, **30**, 46-47 (*rev, bibl*)

Magnus, P. *et al.*, *J.A.C.S.*, 1993, **115**, 8116-8129 (*synth*)

Knight, S.D. *et al.*, *J.A.C.S.*, 1993, **115**, 9293-9294 (*synth*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1993, **58**, 7490-7497 (*synth*)

Beddoes, R.L. *et al.*, *Acta Cryst. C*, 1994, **50**, 447-450 (*Neostrychnine, cryst struct*)

Beifuss, U. *et al.*, *Angew. Chem., Int. Ed.*, 1994, **33**, 1144-1149 (*synth*)

Kuijpers, G.A. *et al.*, *Br. J. Pharmacol.*, 1994, **113**, 471-478 (*pharmacol*)

Rawal, V.H. *et al.*, *J.O.C.*, 1994, **59**, 2685-2686 (*synth*)

Cai, B.C. *et al.*, *Yaoxue Xuebao*, 1994, **29**, 44-48 (*10-O-Demethylbrucine*)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **1**, 797-799 (*use*)

Aprison, M.H. *et al.*, *J. Neurosci. Res.*, 1995, **40**, 396-400 (*pharmacol, conformn*)

Nicholaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, **21**; 641 (*bibl, synth*)

Dijksma, F.J.J. *et al.*, *Acta Cryst. C*, 1998, **54**, 1948-1951 (*cryst struct*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1998, **63**, 9427-9433 (*synth*)

Solé, D. *et al.*, *Angew. Chem., Int. Ed.*, 1999, **38**, 395-397 (*synth*)

Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 1633.1; 1609.3

Martin, G.E. *et al.*, *J. Nat. Prod.*, 2000, **63**, 543-585 (*Strychnine, Brucine, N-15 nmr*)

Pesticide Manual, 12th edn., 2000, 708

Nakanishi, M. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **114**, 2014-2016 (*synth*)

Mori, M. *et al.*, *J.A.C.S.*, 2003, **125**, 9801-9807 (*synth*)

Bialonska, A. *et al.*, *Acta Cryst. C*, 2004, **60**, o853-o855 (*Brucine, cryst struct*)

Ohshima, T. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 1031-1052 (*rev, synth*)

Shibasaki, M. *et al.*, *Alkaloids (Academic Press)*, 2007, **64**, 103-138 (*rev, synth*)

Zhang, H. *et al.*, *Org. Lett.*, 2007, **9**, 279-282 (*synth*)

Buckingham, J. *et al.*, *Bitter Nemesis: The Intimate History of Strychnine*, CRC Press, 2008, (*book*)


Chemical Hazards of the Workplace, 3rd edn., (eds. Proctor, N.H. *et al.*), Van Nostrand Reinhold, 1991, 517

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 10th edn.*, J. Wiley, 2000, SMN5000; SMO000; SMP000; BOL750

Strychnobaridine

S-590

[67650-02-6]



Proposed structure

C₃₀H₃₄N₄O₂ 482.624

Conceivably a mixt. Alkaloid from the leaves of *Strychnos usambarensis* (Loganiaceae).


Angenot, L. *et al.*, *J. Pharm. Belg.*, 1978, **33**, 11-23; *CA*, **89**, 129770t

Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 49 (*rev*)

Strychnobiline

S-591

1-Acetyl-19,20-didehydro-16-(23-deoxy-11-oxa-12,24-secostrychnidin-10-yl)-17-norcuran, 9CI
[68727-53-7]



Absolute Configuration

C₄₀H₄₆N₄O₂ 614.829

Alkaloid from *Strychnos variabilis* (Loganiaceae).

12'-Hydroxy: 12'-Hydroxystrychnobiline
[88668-50-2]

C₄₀H₄₆N₄O₃ 630.828

Alkaloid from *Strychnos variabilis* root bark (Loganiaceae).

16',17'-Diepimer: Isostrychnobiline

[68780-12-1]

C₄₀H₄₆N₄O₂ 614.829

Alkaloid from *Strychnos variabilis* (Loganiaceae). Microcryst. powder. Violet col. with Ce(IV).

16',17'-Diepimer, 16,17-didehydro: 16,17-Dehydroisostrychnobiline

[90332-58-4]

C₄₀H₄₄N₄O₂ 612.813

Alkaloid from root bark of *Strychnos kasengaensis* (Loganiaceae). [α]_D +49 (c, 0.64 in MeOH).

16',17'-Diepimer, 12'-hydroxy: 12'-Hydroxyisostrychnobiline

[71610-50-9]

C₄₀H₄₆N₄O₃ 630.828

Alkaloid from *Strychnos variabilis* (Loganiaceae). Powder. Violet col. with Ce(IV).

Tits, M.J.G. *et al.*, *Planta Med.*, 1978, **34**, 57-61 (*isol*)

Tits, M.J.G. *et al.*, *Phytochemistry*, 1979, **18**,

515-516 (*12'-Hydroxyisostrychnobiline*)

Tits, M.J.G. *et al.*, *J. Nat. Prod.*, 1983, **46**, 638-


645 (*isol, struct, pmr, ms*)

Thepenier, P. *et al.*, *Phytochemistry*, 1984, **23**,

2659 (*16,17-Dehydroisostrychnobiline*)**Strychnochromine**

S-592

[69345-20-6]

C₁₈H₂₂N₂O₂ 298.384


Related to the Condylocarpan alkaloids. Alkaloid from *Strychnos gossweileri* (Loganiaceae). Gives intense colour reagents with numerous reagents, e.g. H₂SO₄

(stable violet), Ce(SO₄)₂ (red). Mp 75°.
 Coune, C. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 106 (*isol, uv, ir, ms*)
 Quetin-Leclercq, J. *et al.*, *Tet. Lett.*, 1991, **32**, 4295 (*pmr, cmr, cryst struct*)

Strychnochrysin

S-593

[201009-99-6]

Absolute
ConfigurationC₃₈H₃₉N₄O₂⁺ 583.752

Related to Longicaudatine Z, L-250.
 Quaternary alkaloid from the roots of *Strychnos nux-vomica* (Loganiaceae).
 Orange amorph. powder. Counterion not specified. λ_{max} 210 (log ε 4.25); 252 (log ε 3.8); 275 (log ε 3.76); 311 (log ε 3.88); 428 (log ε 3.75) (MeOH).

9'-Methoxy, N⁴-Me: Guiachrysin[284675-26-9]
C₄₀H₄₄N₄O₃⁺ 628.813

Quaternary alkaloid from *Strychnos guianensis*. Orange amorph. powder.
 Counterion not specified. λ_{max} 205 (log ε 4.26); 254 (log ε 3.96); 320 (log ε 3.87); 426 (log ε 3.86) (MeOH).

9'-Methoxy, 5',6'-didehydro, N⁴-Me:**5',6'-Dehydroguaiachrysin**[381724-72-7]
C₄₀H₄₂N₄O₃⁺ 626.797

Quaternary alkaloid from *Strychnos guianensis*. Orange-brown powder.
 Counterion not specified. λ_{max} 205 (log ε 4.38); 258 (log ε 4.29); 279 (log ε 4.07); 328 (log ε 3.87); 421 (log ε 3.98) (MeOH).


Biala, R.G. *et al.*, *J. Nat. Prod.*, 1998, **61**, 139-141 (*Strychnochrysin*)

Penelle, J. *et al.*, *Phytochemistry*, 2000, **53**, 1057-1066; 2001, **58**, 619-626 (*Guiachrysin, Dehydroguaiachrysin*)

Strychnofendlerine

S-594

l-Acetyl-17,19-epoxy-4-methyl-3,4-seco-curaran-3-one, 9CI
 [62278-92-6]

Absolute
ConfigurationC₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Strychnos fendleri*, *Strychnos scheffleri*, *Strychnos soubrensis* and *Strychnos aculeata* (Loganiaceae).
 Cryst. (EtOAc). Mp 245-247°. [α]_D²⁰ +97 (c, 1.4 in CHCl₃). [α]_D²⁰ +264 (c, 1 in CHCl₃). λ_{max} 248 (log ε 4.16); 278 (log ε

3.53); 286 (sh) (log ε 3.45) (EtOH).

20,21-Didehydro: Strychnobrasiline[34227-00-4]
C₂₂H₂₆N₂O₃ 366.459

Alkaloid from *Strychnos brasiliensis*, *Strychnos tabascana*, *Strychnos soubrensis*, *Strychnos alvimiana*, *Strychnos myrtilloides* and *Strychnos scheffleri* (Loganiaceae). Shows significant chloroquine potentiating action against resistant *Plasmodium* spp.
 Cryst. (Et₂O or EtOAc). Mp 218-222° (205-208°). [α]_D²⁵ +156 (c, 0.91 in CHCl₃). λ_{max} 247 (log ε 4.15); 277 (sh) (log ε 3.5); 285 (sh) (log ε 3.43) (EtOH).

20,21-Didehydro, N-de-Ac: Deacetylstrychnobrasiline[139694-99-8]
C₂₀H₂₄N₂O₂ 324.422

Alkaloid from root bark of *Strychnos mostueoides*.

14β-Hydroxy, 20,21-didehydro: 14-Hydroxystrychnobrasiline[92202-53-4]
C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the stem bark of *Strychnos soubrensis* (Loganiaceae).

10-Methoxy, 20,21-didehydro: 10-Methoxystrychnobrasiline[35611-65-5]
C₂₃H₂₈N₂O₄ 396.485

Alkaloid from *Strychnos tabascana* (Loganiaceae). Cryst. (EtOH). Mp 255-257°.

11-Methoxy: 11-Methoxystrychnofendlerine[72994-78-6]
C₂₃H₃₀N₂O₄ 398.501

Alkaloid from *Strychnos fendleri* stem bark (Loganiaceae). Cryst. (EtOAc/hexane). Mp 242-244°. λ_{max} 251 (log ε 4.03); 290 (log ε 3.72); 296 (sh) (log ε 3.69) (EtOH).

11-Methoxy, 12-hydroxy: 12-Hydroxy-11-methoxystrychnofendlerine[62258-28-0]
C₂₃H₃₀N₂O₅ 414.5

Alkaloid from *Strychnos fendleri* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 262-263°. [α]_D²⁰ -3.8 (c, 0.85 in CHCl₃). λ_{max} 226 (log ε 4.41); 254 (log ε 3.9); 290 (sh) (log ε 3.28) (EtOH).

11-Methoxy, 12-hydroxy, 20,21-didehydro: 12-Hydroxy-11-methoxystrychnobrasiline[34227-01-5]
C₂₃H₂₈N₂O₅ 412.485

Alkaloid from *Strychnos brasiliensis* trunk bark (Loganiaceae). Needles (MeOH). Mp 229-230°. [α]_D¹⁸ -31 (c, 0.57 in CHCl₃). λ_{max} 227 (log ε 4.65); 263 (sh) (log ε 3.5); 290 (sh) (log ε 3.4) (EtOH).

10,11-Dimethoxy, 20,21-didehydro: 10,11-Dimethoxystrychnobrasiline[34327-12-3]
C₂₄H₃₀N₂O₅ 426.511

Alkaloid from *Strychnos brasiliensis* trunk bark (Loganiaceae). Small plates (EtOH). Mp 168-175°. [α]_D²⁰ +148 (c, 0.61 in CHCl₃). λ_{max} 217 (log ε 4.47); 258 (log ε 4.1); 297 (log ε 3.67) (EtOH).

10,11-Dimethoxy, 12-hydroxy, 20,21-didehydro: 12-Hydroxy-10,11-dimethoxystrychnobrasilineC₂₄H₃₀N₂O₆ 442.511

Alkaloid from the stem bark of *Strychnos mattogrossensis*. Powder (EtOAc). Mp 260-262°. No CAS no. found CA 136-138. λ_{max} 222 (log ε 4.29); 255 (log ε 3.94); 294 (log ε 3.14) (MeOH).

19-Epimer: Isosplendine[23943-38-6]
C₂₂H₂₈N₂O₃ 368.475

Alkaloid from fruits of *Strychnos splendens* and from *Strychnos aculeata* and *Strychnos soubrensis* (Loganiaceae). Prisms (Et₂O/EtOH). Mp 232-233° (block). [α]_D²⁰ +206 (c, 1 in EtOH).

Koch, M. *et al.*, *Ann. Pharm. Fr.*, 1969, **27**, 229-238 (*Isosplendine, isol, ms, struct*)

Koch, M. *et al.*, *Tetrahedron*, 1969, **25**, 3377-3382 (*Isosplendine, synth*)

Galeffi, C. *et al.*, *Farmacol. Ed. Sci.*, 1971, **26**, 1100-1114; *CA*, **76**, 59837a (*Strychnobrasiline, 10-*

Methoxystrychnobrasiline, isol, uv, pmr, ms)

Iwataki, I. *et al.*, *Tetrahedron*, 1971, **27**, 2541-2552 (*Strychnos brasiliensis constits*)

Galeffi, C. *et al.*, *Gazz. Chim. Ital.*, 1976, **106**, 773-777 (*Strychnos fendleri constits*)

Mirand, C. *et al.*, *Plant. Med. Phytother.*, 1979, **13**, 84-86 (*isol*)

Galeffi, C. *et al.*, *Gazz. Chim. Ital.*, 1980, **110**, 81-85 (*11-Methoxystrychnofendlerine, 12-*

Hydroxy-11-methoxystrychnofendlerine)

Caprasse, M. *et al.*, *Planta Med.*, 1981, **42**, 364-370 (*isol*)

Marini-Bettolo, G.B. *et al.*, *An. Asoc. Quim. Argent.*, 1982, **70**, 263-270 (*isol*)

Ohiri, F.C. *et al.*, *J. Nat. Prod.*, 1983, **46**, 369-373 (*Strychnobrasiline, Isosplendine, 14β-*

Hydroxystrychnobrasiline)

Mostad, A. *et al.*, *Acta Chem. Scand., Ser. B*, 1984, **38**, 381-386 (*cryst struct, 14β-*

Hydroxystrychnobrasiline)

Angenot, L. *et al.*, *Phytochemistry*, 1990, **29**, 2746-2749 (*Strychnobrasiline, 12-Hydroxy-*

10,11-dimethoxystrychnobrasiline)

Rasoanivo, P. *et al.*, *Rev. Latinoam. Quim.*, 1991, **22**, 32-34; *CA*, **116**, 148160m (*Deacetylstrychnobrasiline*)

Rasoanivo, P. *et al.*, *Planta Med.*, 1994, 13-16 (*isol, activity*)

Martin, M.-T. *et al.*, *Magn. Reson. Chem.*, 1996, **34**, 489-491 (*pmr, cmr*)


Trigalo, F. *et al.*, *Tetrahedron*, 1999, **55**, 6139-6146 (*Deacetylstrychnobrasiline, synth*)

Belem-Pinheiro, M.L. *et al.*, *Nat. Prod. Lett.*, 2002, **16**, 229-233 (*12-Hydroxy-10,11-*

dimethoxystrychnobrasiline)**Strychnofoline**

S-595

[63408-17-3]

C₃₀H₃₄N₄O₂ 482.624

Principal alkaloid of leaves of *Strychnos usambarensis* (Loganiaceae). Antimitotic

agent. Powder. Mp 175°.

Deoxy: Barterine

$C_{30}H_{34}N_4O$ 466.625

Alkaloid from leaves of *Strychnos barteri*. Cryst. (EtOAc/hexane). Mp 166-168°. $[\alpha]_D^{20} +58.1$ (c, 0.7 in EtOH). λ_{max} 218 (log ϵ 4.7); 269 (log ϵ 4.22); 288 (log ϵ 4.09) (EtOH).

Deoxy, 10-hydroxy: 10-Hydroxybarterine

$C_{30}H_{34}N_4O_2$ 482.624

Alkaloid from leaves of *Strychnos barteri*. Cryst. (EtOAc/hexane). Mp 189-192°. $[\alpha]_D^{20} -9.5$ (c, 1 in EtOH). λ_{max} 218 (log ϵ 4.71); 270 (log ϵ 4.24); 289 (log ϵ 4.09) (EtOH).

7-Epimer: Isostrychnofoline

[69351-04-8]

$C_{30}H_{34}N_4O_2$ 482.624

Alkaloid from leaves of *Strychnos usambarensis* (Loganiaceae). Powder. Mp 182°.

Dideberg, O. *et al.*, *Acta Cryst. B*, 1977, **33**, 1796-1801 (*cryst struct*)

Angenot, L. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 123-129 (*isol, uv, ir, pmr, ms, struct*)

Oguakwa, J.U. *et al.*, *Rend. Accad. Naz. Quaranta*, 1979, **4(2)**, 9-14 (*Barterine, 10-Hydroxybarterine*)


Coune, C. *et al.*, *J. Pharm. Belg.*, 1982, **37**, 189 (*cmr, struct*)

Lerchner, A. *et al.*, *Chem. Eur. J.*, 2006, **12**, 8208-8219 (*synth*)

Strychnofuranine

S-596

[119308-20-2]



$C_{38}H_{40}N_4O$ 568.76


Alkaloid from the root bark of *Strychnos matopensis* (Loganiaceae). $[\alpha]_D +14$ (c, 0.47 in $CHCl_3$).

Massiot, G. *et al.*, *Phytochemistry*, 1988, **27**, 3293-3304 (*isol, uv, ir, pmr, cmr, ms, struct*)

Strychnogucine A

S-597

[324519-12-2]



$C_{42}H_{42}N_4O_3$ 650.819

Alkaloid from the roots of *Strychnos icaja*. Antiplasmodial agent. Amorph. powder. λ_{max} 214 (log ϵ 3.22); 255 (log ϵ 2.98); 281 (log ϵ 2.64); 289 (log ϵ 2.58) (MeOH).

 $\Delta^{17,23}$ -Isomer: Strychnogucine C

[532971-12-3]

$C_{42}H_{42}N_4O_3$ 650.819

Alkaloid from the roots of *Strychnos icaja*. Yellowish amorph. powder. λ_{max} 208 (log ϵ 4.63); 255 (log ϵ 4.14); 283 (log ϵ 3.93) (MeOH).

► Possesses 16'βH-config.


Frederich, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 12-16 (*Strychnogucine A*)

Philippe, G. *et al.*, *Phytochemistry*, 2003, **62**, 623-629 (*Strychnogucine C*)

Strychnogucine B

S-598

[324519-13-3]



Absolute
Configuration

$C_{42}H_{42}N_4O_4$ 666.818


Alkaloid from the roots of *Strychnos icaja*. Antiplasmodial agent. Amorph. powder. λ_{max} 222 (log ϵ 4.45); 252 (log ϵ 4.12); 283 (log ϵ 3.91); 292 (log ϵ 3.86); 304 (log ϵ 3.81) (MeOH).

Frederich, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 12-16 (*isol, pmr, cmr, uv, cd*)

Strychnohexamine

S-599

[442123-70-8]



Absolute
Configuration

$C_{59}H_{60}N_6O$ 869.163

Alkaloid from the roots of *Strychnos icaja*. Antiplasmodial agent. Yellowish powder. λ_{max} 210 (log ϵ 3.55); 293 (log ϵ 3.25); 320 (log ϵ 2.99) (MeOH).


Philippe, G. *et al.*, *Tet. Lett.*, 2002, **43**, 3387-3390 (*isol, pmr, cmr*)

Philippe, G. *et al.*, *Phytochemistry*, 2003, **62**, 623-629 (*cd, ms*)

Strychnohirsutine

S-600

16-De(methoxycarbonyl)-17-deoxy-19,20-dihydro-19-hydroxy-21-methyl-18-nortaltbotine, 9CI
[73061-90-2]



$C_{19}H_{24}N_2O_2$ 312.411

Alkaloid from *Strychnos hirsuta* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 141-143°. $[\alpha]_D^{20} -6.3$ (c, 0.5 in $CHCl_3$).

3,4,5,6-Tetrahydro: Tetrahydrostrychnohirsutine

[73061-91-3]

$C_{19}H_{20}N_2O_2$ 308.379


Alkaloid from *Strychnos hirsuta* (Loganiaceae). Cryst. (Et_2O). Mp 221-223°. Fluorescent. Cryst. with difficulty.

Galeffi, C. *et al.*, *Tetrahedron*, 1981, **37**, 3167 (*isol, struct, ir, uv*)

Strychnopentamine

S-601

[63209-34-7]



Absolute
configuration

$C_{35}H_{43}N_5O$ 549.758

Alkaloid from the leaves of *Strychnos usambarensis* (Loganiaceae). Shows strong antimetabolic activity. Shows anti-tumour activity *in vitro* against Ehrlich ascites, HW165 hepatoma and B16 melanoma test systems. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . λ_{max} 226 (ϵ 23000); 275 (ϵ 6650); 291 (ϵ 5500) (MeOH) (Berdy).

2''-Epimer: Isostrychnopentamine A

$C_{35}H_{43}N_5O$ 549.758

Alkaloid from the leaves of *Strychnos usambarensis* (Loganiaceae). Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Epimeric at the asymmetric carbon atom of the *N*-methylpyrrolidin-2-yl group.

Dupont, L. *et al.*, *Acta Cryst. B*, 1977, **33**, 1801-1807 (*cryst struct*)


Angenot, L. *et al.*, *J. Pharm. Belg.*, 1978, **33**, 11-23 (*isol, uv, ir, pmr, ms, cd, struct*)

Tavernier, D. *et al.*, *Phytochemistry*, 1987, **26**, 557-560 (*Isostrychnopentamine A*)

Strychnophylline

S-602

[69306-88-3]

C₃₅H₄₃N₅O₂ 565.757

Probable abs. config. shown. Alkaloid from *Strychnos usambarensis* (Loganiaceae). Shows cytotoxic activity. Powder. Mp 216°.

7-Epimer: Isostrychnophylline

[69351-03-7]

C₃₅H₄₃N₅O₂ 565.757

Trace alkaloid from leaves of *Strychnos usambarensis* (Loganiaceae).


Angenot, L. *et al.*, *Plant. Med. Phytother.*, 1978, **12**, 123 (*isol, uv, ir, pmr, ms, struct*)

Strychnopivotine

S-603

1-Acetyl-19,20-didehydro-17-norcuran-16-one, 9CI

[76177-20-3]



Absolute configuration

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Strychnos variabilis* (Loganiaceae). Powder.


Tits, M. *et al.*, *Phytochemistry*, 1980, **19**, 1531-1534 (*isol, uv, ir, pmr, ms, struct*)

Strychnosilidine

S-604

Acetyltabascanine

[34327-13-4]



Absolute Configuration

C₂₆H₃₄N₂O₆ 470.564

A 3,4-secocuran. Alkaloid from the bark of *Strychnos brasiliensis*, *Strychnos tabascanina* and *Strychnos alvimiana* (Loganiaceae). Needles (EtOH). Mp 236-237°. [α]_D +233 (c, 0.63 in CHCl₃). λ_{\max} 214 (log ϵ 4.5); 260 (log ϵ 4.14); 290 (log ϵ 3.92) (EtOH).

O-De-Ac: Tabascanine

[35611-61-1]

C₂₄H₃₂N₂O₅ 428.527

Alkaloid from *Strychnos tabascanina* (Loganiaceae). Cryst. (EtOAc). Mp

210-213°.

19,20-Dihydro, 18,19,20,21-tetradecydro:

Strychnosiline

[34436-25-4]

C₂₆H₃₂N₂O₆ 468.549

Alkaloid from the bark of *Strychnos brasiliensis* and from *Strychnos alvimiana* (Loganiaceae). Powder. [α]_D²⁰ +35 (c, 0.82 in CHCl₃). λ_{\max} 217 (log ϵ 4.47); 256 (log ϵ 3.82); 299 (log ϵ 3.37) (EtOH).

Bis(demethoxy), O-de-Ac: Malagashine

[139682-32-9]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from root bark of *Strychnos mostueoides* (Loganiaceae). Probable stereochem.

Galeffi, C. *et al.*, *Farmaco, Ed. Sci.*, 1971, **26**, 1100-1114; *CA*, **76**, 59837a

(*Tabascanine, Strychnosilidine, isol, ir, uv, ir, pmr, ms, ord, struct*)

Iwataki, I. *et al.*, *Tetrahedron*, 1971, **27**, 2541 (*Strychnosilidine, Strychnosiline, isol, uv, cd, ms, struct*)

Marini-Bettolo, G.B. *et al.*, *An. Asoc. Quim. Argent.*, 1982, **70**, 263-270; *CA*, **97**, 3556 (*Strychnosilidine, isol*)


Rasonaivo, P. *et al.*, *Rev. Latinoam. Quim.*, 1991, **22**, 32-34; *C.A.*, **116**, 148160m (*Malagashine*)

Strychnosplendine

S-605

17,19-Epoxycuran-3-ol, 9CI

[6516-49-0]



Absolute Configuration

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from leaves and fruit of *Strychnos splendens* (Loganiaceae). Cryst. Mp 203-204° (block). [α]_D²⁰ -82 (c, 1 in EtOH). pK_a 8.35 (DMF). λ_{\max} 245 (log ϵ 3.7); 297 (log ϵ 3.29) (no solvent reported).

N¹-Ac: N^a-Acetylstrychnosplendine

[14013-80-0]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from *Strychnos fendleri* stem bark, *Strychnos aculeata*, *Strychnos scheffleri* and *Strychnos henningsii* root bark (Loganiaceae). Cryst. (EtOAc). Mp 155-157°. [α]_D²⁰ +121 (c, 0.75 in CHCl₃).

N¹-Hydroxyacetyl: Splendoline

[22152-58-5]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from leaves of *Strychnos splendens* and from *Strychnos henningsii* (Loganiaceae). Cryst. (C₆H₆). Mp 250° dec. [α]_D²² +122 (c, 1 in CHCl₃). pK_a 7.55 (DMF). λ_{\max} 250 (log ϵ 4.12); 277 (sh) (log ϵ 3.48) (no solvent reported).

Me ether, N¹-Ac: N^a-Acetyl-O-methylstrychnosplendine

[35611-63-3]

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Strychnos tabascanina* after MeOH extraction, also from *Strychnos aculeata* and *Strychnos scheffleri* (Loganiaceae). Cryst. (EtOAc/hexane). Mp 151-153°. Prob. artifact, also obt. by heating N^a-Acetylstrychnosplendine with MeOH.

11-Methoxy, N¹-Ac: N^a-Acetyl-11-methoxystrychnosplendine

[90890-73-6]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from *Strychnos henningsii* root bark (Loganiaceae). Cryst. (EtOAc). Mp 200-202°. [α]_D²⁰ -101.1 (c, 1 in CHCl₃). λ_{\max} 224 (log ϵ 4.32); 251 (log ϵ 4.03); 290 (log ϵ 3.73); 296 (sh) (log ϵ 3.7) (EtOH).

11-Methoxy, 12-hydroxy, N¹-Ac: 12-Hydroxy-11-methoxy-N^a-acetylstrychnosplendine

[62230-08-4]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from *Strychnos fendleri* (Loganiaceae). Cryst. (EtOAc). Mp 242-244°. [α]_D²⁰ -141 (c, 0.75 in CHCl₃). λ_{\max} 226 (log ϵ 4.42); 254 (log ϵ 3.87); 296 (sh) (log ϵ 3.3) (EtOH).

19-Epimer: Isostrychnosplendine

[23943-35-3]

C₁₉H₂₄N₂O₂ 312.411

Alkaloid from leaves, fruit and trunkwood of *Strychnos splendens* (Loganiaceae). Amorph.; cryst. (as hydrochloride). Mp 330° (hydrochloride). [α]_D²⁰ -59 (c, 1 in H₂O) (hydrochloride). λ_{\max} 244 (log ϵ 3.7); 297 (log ϵ 3.34) (no solvent reported).

19-Epimer, N¹-Ac: N^a-Acetylisostrychnosplendine

[22152-57-4]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from leaves, trunk and fruit of *Strychnos splendens* (Loganiaceae). Mp 190-192°. [α]_D²⁰ +125 (c, 1 in EtOH). λ_{\max} 250 (log ϵ 3.89); 277 (sh) (log ϵ 3.22) (no solvent reported).

19-Epimer, N¹-hydroxyacetyl: Isosplendoline

[22153-10-2]

C₂₁H₂₆N₂O₄ 370.447

Alkaloid from leaves, trunk and fruit of *Strychnos splendens* (Loganiaceae). Mp 250° dec. [α]_D²⁰ +110 (c, 1 in CHCl₃).

Koch, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1968, 3250-3252 (*isol, struct, config, uv, ir, pmr, ms, Strychnos splendens constits*)

Koch, M. *et al.*, *Tetrahedron*, 1969, **25**, 3377-3382 (*isol, struct, config, uv, ir, pmr, ms*)

Galeffi, C. *et al.*, *Gazz. Chim. Ital.*, 1976, **106**, 773-777; 1983, **113**, 773-775 (*Strychnos fendleri constits, Strychnos henningsii constits*)


Caprasse, M. *et al.*, *Planta Med.*, 1981, **42**, 364-370 (*Strychnos scheffleri constits*)

Massiot, G. *et al.*, *Phytochemistry*, 1991, **30**, 3449-3456 (*Splendoline, isol*)

Strychnoxanthine

S-606

[69345-02-4]

C₁₉H₁₉N₂O₂⁺ 307.371

An isoyohimban. Alkaloid from the root bark of *Strychnos gossweileri* (Loganiaceae).

Deoxo: Melinonine E

[483-11-4]

C₁₉H₂₁N₂O⁺ 293.388

Quaternary alkaloid from the bark of *Strychnos melinoniana* (Loganiaceae). Fine brown cryst. (MeOH) (as chloride). Mp 283.6–283.8° (chloride). [α]_D²² -13.9 (c, 1.02 in MeOH) (chloride). Relative config. only appears to be known.

Bächli, E. et al., *Helv. Chim. Acta*, 1957, **40**, 1167 (*Melinonine E*, isol, uv, ir)

Borris, R.P. et al., *Helv. Chim. Acta*, 1984, **67**, 455 (*Melinonine E*, uv, ir, pmr, cmr, ms, struct)


Coune, C. et al., *Planta Med.*, 1984, **50**, 93-95 (*Strychnoxanthine*, isol, uv, ir, pmr, cmr, ms, cd, struct)

Quirante, J. et al., *J.O.C.*, 1998, **63**, 968-976 (*synth*)

Strychnozairine

S-607

1-Acetyl-20,21-didehydro-19-oxocuran-17-al, 9CI
[95360-06-8]



Absolute Configuration

C₂₁H₂₂N₂O₃ 350.416


Alkaloid from root bark of *Strychnos variabilis* (Loganiaceae).

Tits, M. et al., *Phytochemistry*, 1985, **24**, 205-207 (*isol, uv, ir, ms, pmr, cmr, cd, struct*)

Styelsamine C

S-608

[216446-11-6]



R = CHO

C₁₆H₁₀N₂O₂ 262.267

Isol. from the ascidian *Eusynstyela latericius*. Cytotoxic. Orange solid (as TFA salt). λ_{max} 248 (log ε 4.9); 284 (log ε 4.82); 334 (log ε 4.22); 350 (log ε 4.27); 418 (log ε 4.75); 480 (log ε 4.41) (CHCl₃).

Copp, B.R. et al., *J.O.C.*, 1998, **63**, 8024-8026 (*isol, pmr, cmr*)

Nakahara, S. et al., *Heterocycles*, 2003, **60**, 2017-2022; 2005, **65**, 1925-1929 (*synth*)

Styelsamine D

S-609

[216365-44-5]

As Styelsamine C, S-608 with

R = -¹³CH₂¹⁴CH₂NH₂C₁₇H₁₅N₃O 277.325

Closely related to Deacylcystodytin, D-97. Isol. from the ascidian *Eusynstyela latericius*. Purple solid (as bistrifluoroacetate salt). λ_{max} 222 (log ε 4.3); 238 (log ε 4.17); 276 (log ε 4.37); 294 (log ε 4.38); 320 (log ε 3.8); 372 (log ε 3.54); 386 (log ε 3.6); 560 (log ε 3.43) (MeOH/trifluoroacetate). λ_{max} 270 (log ε 4.26); 376 (log ε 3.95) (MeOH/KOH).

N¹⁴-Ac: Styelsamine B

[216387-14-3]

C₁₉H₁₇N₃O₂ 319.362

Isol. from *Eusynstyela latericius*. Purple solid (as trifluoroacetate salt). λ_{max} 222 (sh) (log ε 4.43); 244 (log ε 4.28); 278 (log ε 4.55); 294 (log ε 4.6); 372 (sh) (log ε 3.57); 388 (log ε 3.7); 572 (log ε 3.72) (MeOH/trifluoroacetate). λ_{max} 262 (log ε 4.38); 378 (log ε 3.97) (KOH).

13-Hydroxy: Styelsamine A

[216446-06-9]

C₁₇H₁₅N₃O₂ 293.324

Isol. from *Eusynstyela latericius*. Purple solid (as bistrifluoroacetate salt). λ_{max} 222 (sh) (log ε 3.86); 244 (log ε 3.71); 276 (log ε 3.93); 294 (log ε 3.96); 386 (log ε 3.17); 554 (log ε 3.02) (MeOH/trifluoroacetate). λ_{max} 272 (log ε 3.85); 378 (log ε 3.39) (MeOH/KOH).

Copp, B.R. et al., *J.O.C.*, 1998, **63**, 8024-8026 (*Styelsamines A, B*, isol, ir, pmr, cmr, uv)


Skyler, D. et al., *Org. Lett.*, 2001, **3**, 4323-4324 (*Styelsamine B*, synth)

Stylessadine A

S-610

Flabellazole B

[914769-87-2]



Absolute Configuration

C₄₄H₄₆Br₈N₂₀O₉ 1638.208

Dimer of Massadine, M-112. Alkaloid from *Stylissa caribica* and *Stylissa flabellata*. Specific agonist of the P2X₇ receptor. [α]_D²² -15.2 (c, 1.21 in MeOH). λ_{max} 235; 275 (no solvent reported). λ_{max} 277 (log ε 4.49) (MeOH).

2-Epimer: Stylessadine B. Flabellazole A
[914769-88-3]

C₄₄H₄₆Br₈N₂₀O₉ 1638.208

Alkaloid from *Stylissa caribica* and *Stylissa flabellata*. [α]_D²² -20 (c, 0.62 in MeOH). λ_{max} 277 (log ε 4.56) (MeOH).

Grube, A. et al., *Org. Lett.*, 2006, **8**, 4675-4678 (*isol, pmr, cmr, ms*)


Buchanan, M.S. et al., *J.O.C.*, 2007, **72**, 2309-2317 (*isol, pmr, cmr, ms*)

Styloguanidine

S-611

Isopalauamine

[163089-72-3]



Absolute Configuration

C₁₇H₂₂ClN₉O₂ 419.873

Stereochem. revised in 2007. Alkaloid from the marine sponge *Stylorella aurantium*. Potent chitinase inhibitor. Antifouling agent. Amorph. solid. [α]_D +20.7 (c, 3.5 in MeOH). λ_{max} 224 (ε 7800); 272 (ε 7900) (MeOH) (Derep).

3-Bromo: 3-Bromostyloguanidine

[162339-62-0]

C₁₇H₂₁BrClN₉O₂ 498.769

From *Stylorella aurantium*. Potent chitinase inhibitor; antifouling agent. Tan solid. [α]_D +57.5 (c, 0.7 in MeOH). λ_{max} 277 (ε 5000) (MeOH) (Derep).

2,3-Dibromo: 2,3-Dibromostyloguanidine

[162339-63-1]

C₁₇H₂₀Br₂ClN₉O₂ 577.665

From *Stylorella aurantium*. Potent chitinase inhibitor; antifouling agent. Off-white cryst. (2-propanol/MeOH). [α]_D -70.8 (c, 0.6 in MeOH). λ_{max} 283 (ε 5800) (MeOH) (Derep).

2,3-Dibromo, N¹⁹-(3,4-dibromopyrrole-2-carbonyl): Carteramine A. Tetrabromostyloguanidine

[934986-32-0]

C₂₂H₂₁Br₄ClN₁₀O₃ 828.542

Alkaloid from *Stylissa carteri* and *Stylissa caribica*. Inhibitor of neutrophil chemotaxis. Light yellow powder. [α]_D¹⁷ -44 (c, 0.5 in MeOH). [α]_D²³ -42 (c, 1.26 in MeOH). Dimer of Oroidin, O-123. The name tetrabromostyloguanidine is misleading. λ_{max} 217 (ε 11700); 277 (ε 13700) (MeOH).

Kato, T. et al., *Tet. Lett.*, 1995, **36**, 2133-2136 (*isol, uv, ir, pmr, cmr*)

Kinnel, R.B. et al., *J.O.C.*, 1998, **63**, 3281-3286 (*isol, pmr, cmr*)

Grube, A. et al., *Angew. Chem., Int. Ed.*, 2007, **46**, 2320-2324 (*config, Tetrabromostyloguanidine*)


Buchanan, M.S. et al., *J.O.C.*, 2007, **72**, 2309-2317 (*config*)

Kobayashi, H. et al., *Tet. Lett.*, 2007, **48**, 2127-2129 (*Carteramine A*)

Stylopine

S-612

6,7,12b,13-Tetrahydro-4H-bis[1,3]benzodioxolo[5,6-a:4',5'-g]quinolizine, 9CI.
Chelidamine. Corydalis E. Fumjudaine. Stilopine. Tetrahydrocoptisine
 [7461-02-1]
 [11050-51-4 Fumjudaine]

*(R)*-form

C₁₉H₁₇NO₄ 323.348
 Shows antipsychotic and neuroleptic activity in mice and rats. Active against gram-positive and gram-negative bacteria.

(R)-form [2884-83-5]

Alkaloid from several *Corydalis* spp. (Papaveraceae). Mp 204°. [α]_D²⁰ +310 (c, 0.88 in CHCl₃).
Methiodide: Mp 248° dec.

(S)-form [84-39-9]

Alkaloid from *Corydalis* spp., *Chelidonium majus* (Papaveraceae) and several other spp. in Papaveraceae. Prisms. Mp 204°. [α]_D²³ -310 (c, 0.3 in CHCl₃).
 ▶ LD₅₀ (mus, ipr) 368 mg/kg. DR9808480
Hydrochloride: Mp 250°.

N-Me (trans-): [90457-61-7]
 [41431-71-4, 41431-72-5, 61989-70-6, 51795-13-2, 56711-20-7, 56245-69-3]
 C₂₀H₂₀NO₄[⊕] 338.382

Quaternary alkaloid isol. from *Argemone* spp., *Corydalis* spp., *Eschscholtzia* sp., *Stylophorum* spp. and *Glaucium* spp. Small prisms (MeOH)(as iodide). Mp 297-302° (iodide). [α]_D²³ -125 (c, 0.1 in MeOH) (iodide). CAS no. refers to iodide.

N-Me (cis-): **N-Methylstylopine. N-Methylstylopinium**
 [90457-62-8]
 [64474-10-8, 56245-68-2, 113349-14-7]
 C₂₀H₂₀NO₄ 338.382
 Quaternary alkaloid isol. from *Argemone* spp., *Corydalis* spp., *Eschscholtzia* sp. and *Stylophorum* spp. Small prisms (MeOH)(as iodide). Mp 279-281° (iodide). [α]_D²⁰ -120 (c, 0.13 in MeOH) (iodide). Normally obt. as a mixt. with its epimer at N. CAS no. refers to iodide. λ_{max} 209 (log ε 4.67); 232 (sh) (log ε 3.96); 289 (log ε 3.92) (MeOH).

13β-Hydroxy: 13-Hydroxystylopine

[53777-76-7]
 C₁₉H₁₇NO₅ 339.347

Alkaloid from *Corydalis ophiocarpa* (Papaveraceae). Prisms (EtOH). Mp 213-215°. [α]_D -259 (c, 0.1 in CHCl₃).

13β-Hydroxy, N-Me: 13β-Hydroxy-N-methylstylopinium

[108385-06-4]
 C₂₀H₂₀NO₅[⊕] 354.382

Quaternary alkaloid from aerial parts

and roots of *Papaver atlanticum* (Papaveraceae). Cryst. (MeOH) (as iodide). Mp 254-256° (iodide). [α]_D²⁰ -144 (c, 0.1 in MeOH) (iodide). CAS No. refers to iodide.

13β-Acetoxy:

Fine needles (CH₂Cl₂/Et₂O). Mp 237-239° (sealed tube).

(±)-form [4312-32-7]

Alkaloid from several *Corydalis* spp. (Papaveraceae). Needles (CHCl₃). Mp 222-223°. Bp_{0.01} 260°.

Hydrochloride: Mp 246° (266°).

13β-Hydroxy: [53833-90-2]

Synthetic. Fine needles (EtOH). Mp 219-221° dec.

13β-Acetoxy:

Small plates (CH₂Cl₂/petrol). Mp 242-243° dec.

(±)-form

Alkaloid from leaves and stems of *Fumaria judaica* (as Fumjudaine).

8ξ-Hydroxy, 8-O-β-D-glucopyranoside: 8-Glucopyranosyloxystylopine
 [188560-51-2]

C₂₅H₂₇NO₁₀ 501.489

Alkaloid from *Fumaria indica* (Papaveraceae). Mp 250°. λ_{max} 214 ; 240 ; 291 (MeOH).

Späth, E. *et al.*, *Ber.*, 1931, **64**, 1131-1137 (isol)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1973, **38**, 2513-2517; 1975, **40**, 1095-1102; 1991, **56**, 1116-1122 (*N-Methylstylopinium*)

Battersby, A.R. *et al.*, *J.C.S. Perkin I*, 1975, 1140-1147 (isol, biosynth)

Jeffs, P.W. *et al.*, *J.O.C.*, 1975, **40**, 644-647 (*13-Hydroxystylopine*)

Bhattacharya, S.K. *et al.*, *Arzneim.-Forsch.*, 1976, **26**, 2187 (pharmacol)

Takao, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 2859-2868; 3185; 1977, **25**, 1426-1435 (pmr, cmr, biosynth)

Abou-Donia, A.H.A. *et al.*, *Planta Med.*, 1980, **40**, 295-298 (*Fumjudaine*)

Narasimhan, N.S. *et al.*, *Tetrahedron*, 1983, **39**, 1975-1982 (synth)

Bhakuni, D.S. *et al.*, *Alkaloids (Academic Press)*, 1986, **28**, 95-181 (rev)

Táborská, E. *et al.*, *Coll. Czech. Chem. Comm.*, 1986, **51**, 2232-2239 (*13β-Hydroxy-N-methylstylopinium*)

Dai-Ho, G. *et al.*, *J.O.C.*, 1988, **53**, 5113-5127 (synth, uv, ir, pmr, cmr)

Abbasoglu, U. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1991, **324**, 379-380 (activity)

Chrzanowska, M. *et al.*, *J. Nat. Prod.*, 1995, **58**, 401-407 (synth)

Khan, S.A. *et al.*, *J. Indian Chem. Soc.*, 1997, **74**, 62-63 (*13-glucosyloxy*)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202; 2002, **40**, 687-692 (*N-15 nmr, cryst struct*)

Li, M. *et al.*, *Planta Med.*, 2001, **67**, 189-190 (*Thalictrum delavayi derivs*)

Stylorellane A

[199732-52-0]

S-613

C₁₆H₂₅Cl₂N 302.286Constit. of *Stylorella aurantium*.

Δ³⁽¹⁵⁾-Isomer, 2ξ-chloro: Stylorellane B
 [64789-89-5]

C₁₆H₂₄Cl₃N 336.73

Constit. of the sponge *Pseudaxinyssa pitys* and *Stylorella aurantium*.

Δ³⁽¹⁵⁾-Isomer, 1,2-didehydro(E-): Ulosin A

[321557-59-9]

C₁₆H₂₃Cl₂N 300.27

Constit. of *Stylorella aurantium* and *Ulosa spongia*. Oil. λ_{max} 284 (log ε 4.5) (MeOH).

Wratten, S.J. *et al.*, *J.A.C.S.*, 1977, **99**, 7367-7368 (isol, pmr, cmr)

Simpson, J.S. *et al.*, *Tet. Lett.*, 1997, **38**, 7947-7950 (isol, pmr, cmr, biosynth)

Musman, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 111-113 (*Ulosin A*)

Kehraus, S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 939-941 (isol, pmr, cmr)


Brust, A. *et al.*, *Tet. Lett.*, 2003, **44**, 327-330 (biosynth)

Simpson, J.S. *et al.*, *Org. Biomol. Chem.*, 2004, **2**, 949-956 (biosynth)

Stylorellin

S-614

[108648-45-9]

C₁₆H₂₅N 231.38

Constit. of sponge *Stylorella* sp. Oil. [α]_D -47 (c, 1.7 in CHCl₃).

Paś, M. *et al.*, *Tet. Lett.*, 1987, **28**, 1409

Suaedin

S-615

[12712-74-2]

C₅₅H₅₄N₂O₃₁ 1239.026


An O-citryl betacyanin of unknown struct. Pigment from *Suaeda fruticosa* (Chenopodiaceae). λ_{max} 294 ; 330 ; 546 (H₂O).

Piattelli, M. *et al.*, *Phytochemistry*, 1971, **10**, 3133-3134

Suaeolenine

S-616

[138642-02-1]


C₂₁H₂₂N₂O₄ 366.416

A secoaspidosperma alkaloid. Alkaloid from the trunk of *Melodinus suaveolens* (Apocynaceae). Mp 205°.

Ye, J.H. *et al.*, *Phytochemistry*, 1991, **30**, 3168 (isol, uv, ir, pmr, cmr, ms, struct)

Suaveolindole

S-617

C₂₃H₃₁NO₂ 353.503


Alkaloid from the fruit of *Greenwayodendron suaveolens*. Antibacterial agent. $[\alpha]_D^{25} +26$ (c, 1 in MeOH). λ_{\max} 230; 283; 292 (MeOH).

Yoo, H.-D. *et al.*, *J. Nat. Prod.*, 2005, **68**, 122-124 (*isol, pmr, cmr*)
Velthuisen, E.J. *et al.*, *J.A.C.S.*, 2007, **129**, 10640-10641 (*synth, abs config*)

Suaveoline[†]

S-618

[38993-99-6]



Absolute configuration

C₂₀H₂₁N₃ 303.406

Indole alkaloid related to the Macroline group with two cyclisations. Alkaloid from *Rauwolfia suaveolens* (Apocynaceae). Noncryst. $[\alpha]_D$ 0 (c, 1 in CHCl₃). $[\alpha]_D^{25} -9.33$ (c, 0.30 in CHCl₃) (*synthetic*). N-Ac: Mp 207-208°.

N-Me: Mp 194-195°. $[\alpha]_D -93$ (c, 0.89 in CHCl₃).

N-De-Me: Norsuaveoline

[93552-58-0]

C₁₉H₁₉N₃ 289.379Alkaloid from *Rauwolfia caffra*.

Majumdar, S.P. *et al.*, *Phytochemistry*, 1973, **12**, 1167 (*isol, synth, uv, pmr, ms*)

Nasser, A.M.A.G. *et al.*, *J. Ethnopharmacol.*, 1984, **11**, 99-117 (*Norsuaveoline*)

Trudell, M.L. *et al.*, *Tetrahedron*, 1992, **48**, 1805 (*synth*)

Bailey, P.D. *et al.*, *J.C.S. Perkin 1*, 1993, 441; 1997, 1209; 2000, 3566-3577 (*synth*)

Fu, X. *et al.*, *J.O.C.*, 1993, **58**, 661 (*synth, cmr*)

Wang, T. *et al.*, *Tet. Lett.*, 1998, **39**, 8009-8012 (*synth*)


Li, J. *et al.*, *J.A.C.S.*, 1999, **121**, 6998-7010 (*Norsuaveoline, synth*)

Ohba, M. *et al.*, *Tetrahedron*, 2007, **63**, 10337-10344 (*synth*)

Subarine

S-619

[445471-63-6]

C₁₉H₁₃N₃O₃ 331.33

Alkaloid from a Singaporean ascidian. Pale yellow gum. λ_{\max} 233 (log ϵ 4.45); 267 (log ϵ 4.14); 335 (log ϵ 3.72) (MeOH).


Nilar, *et al.*, *J. Nat. Prod.*, 2002, **65**, 1198-1200 (*isol, pmr, cmr, ms*)

Bijeire, L. *et al.*, *Eur. J. Org. Chem.*, 2004, 1891-1893 (*synth, pmr, cmr*)

Subereatensin

S-620

[474670-21-8]


C₁₂H₁₇NO₅ 255.27

Isol. from the sponge *Suberea* aff. *praetensa*. Gum.

Kijjoa, A. *et al.*, *Z. Naturforsch., C*, 2002, **57**, 732-738 (*isol, pmr, cmr, ms*)

Suberedamine A

S-621

C₂₃H₃₀Br₃N₃O₃ 636.22

Related to Purpuramine H, P-823.

(S)-form

Isol. from a sponge, *Suberea* sp. Cytotoxic. Amorph. solid. Mp 64-67°. $[\alpha]_D^{25} +21$ (c, 1 in MeOH). λ_{\max} 281 (ϵ 2400) (MeOH).

N⁸-Me: Suberedamine BC₂₄H₃₂Br₃N₃O₃ 650.247


Isol. from a *Suberea* sp. Cytotoxic. Amorph. solid. Mp 79-81°. $[\alpha]_D^{25} +16$ (c, 1 in MeOH). λ_{\max} 281 (ϵ 2900) (MeOH).

Tsuda, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 980-982

Subhirsine

S-622

3,4-Dimethoxybenzoic acid carbonylbis(8-azabicyclo[3.2.1]octane-8,3-diyl) ester; 9CI. N,N'-Carbonylbis-3-veratroyloxy-nortropine [85412-77-7]

C₃₃H₄₀N₂O₉ 608.687


Dimeric tropane alkaloid with unusual 3 β configs. Alkaloid from *Convolvulus subhirsutus* (Convolvulaceae). Cryst. (Me₂CO). Mp 190-191°.

Aripova, S.F. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 640; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **18**, 606 (*isol, ms, pmr, struct*)

Subincanadine A

S-623

[467420-43-5]



Absolute Configuration

C₁₉H₂₃N₂O[⊕] 295.404

Corynanthe-type alkaloid related to Hunteracine, H-368, biosynth. not yet fully worked out. Quaternary alkaloid from the bark of *Aspidosperma subincanum*. Amorph. solid. $[\alpha]_D^{23} -11$ (c, 1 in MeOH). Counterion not specified. λ_{\max} 225 (log ϵ 4.12); 271 (log ϵ 3.67); 279 (log ϵ 3.64); 289 (log ϵ 3.5) (MeOH).

Deoxy: Subincanadine C

[467420-45-7]

C₁₉H₂₃N₂O[⊕] 279.404

Alkaloid from the bark of *Aspidosperma subincanum*. Amorph. solid. $[\alpha]_D^{23} +5$ (c, 1 in MeOH). Counterion not specified. λ_{\max} 225 (log ϵ 4.09); 271 (log ϵ 3.63); 279 (log ϵ 3.6); 289 (log ϵ 3.45) (MeOH).

Deoxy, 10-hydroxy: Subincanadine G

C₁₉H₂₃N₂O[⊕] 295.404

Alkaloid from the bark of *Aspidosperma subincanum*. Amorph. solid. $[\alpha]_D^{23} -32$ (c, 0.4 in MeOH). λ_{\max} 273 (log ϵ 3.27); 315 (sh) (log ϵ 2.98) (MeOH).

16-Epimer: Subincanadine B

[467420-44-6]

C₁₉H₂₃N₂O[⊕] 295.404

Alkaloid from the bark of *Aspidosperma subincanum*. Amorph. solid. $[\alpha]_D^{23} +41$ (c, 1 in MeOH). Counterion not specified. λ_{\max} 223 (log ϵ 4.23); 273 (log ϵ 3.64); 279 (log ϵ 3.64); 289 (log ϵ 3.54) (MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 2002, **67**, 6449-6455 (*isol, pmr, cmr*)


Ishiyama, H. *et al.*, *Heterocycles*, 2005, **66**, 651-658 (*Subincanadine G*)

Suzuki, K. *et al.*, *Org. Lett.*, 2007, **9**, 4605-4608 (*synth*)

Subincanadine D

S-624

[467420-46-8]

C₁₉H₂₂N₂O 294.396

Rearranged indole alkaloid related to the Akuammicine type. Alkaloid from the bark of *Aspidosperma subincanum*. Amorph. solid. $[\alpha]_D^{23}$ -3.3 (c, 1 in MeOH). λ_{\max} 223 (log ϵ 4); 294 (log ϵ 3.58) (MeOH).

Deoxy: **Pericine**. *Subincanadine E*

[84638-28-8]
[467420-48-0]
 $C_{19}H_{22}N_2$ 278.396

Alkaloid from *Aspidosperma subincanum* and *Picalima nitida*. Amorph. solid. $[\alpha]_D^{23}$ +39 (c, 1 in MeOH). λ_{\max} 226 (log ϵ 4.01); 301 (log ϵ 3.8) (MeOH).

Deoxy, N-oxide: **Pericine N-oxide**

[906096-64-8]
 $C_{19}H_{22}N_2O$ 294.396

Alkaloid from the stem bark of *Kopsia arborea*. Cryst. Mp 205-208°. $[\alpha]_D$ +71 (c, 0.21 in $CHCl_3$). λ_{\max} 227 (log ϵ 4.41); 301 (log ϵ 4.23) (EtOH).


Arens, H. *et al.*, *Planta Med.*, 1982, **46**, 210-214 (*Pericine*)

Kobayashi, J. *et al.*, *J.O.C.*, 2002, **67**, 6449-6455 (*isol*, *pmr*, *cmr*)

Lim, K.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*Kopsia arborea* *constit*)

Subincanadine F S-625

[467420-48-0]



$C_{17}H_{18}N_2O$ 266.342


Truncated indole alkaloid most closely related to the Akuammicine type. Alkaloid from the bark of *Aspidosperma subincanum*. Amorph. yellow solid. $[\alpha]_D^{23}$ +17.8 (c, 1 in MeOH). λ_{\max} 223 (log ϵ 4.12); 274 (log ϵ 3.5); 356 (sh) (log ϵ 2.87) (MeOH).

Kobayashi, J. *et al.*, *J.O.C.*, 2002, **67**, 6449-6455 (*isol*, *pmr*, *cmr*)

Gao, P. *et al.*, *J.O.C.*, 2006, **71**, 9495-9498 (*synth*)

Subincanine S-626

[26237-70-7]



$C_{22}H_{24}N_2O_2$ 348.444

Two stereoisomers of the struct. depicted were synth., neither is the same as the natural alkaloid. It was argued that the other stereoisomers are also very unlikely to correspond to the alkaloid and that the struct. suggested is incorr. Minor alkaloid from *Aspidosperma subincanum* (Apocynaceae). Mp 160-163°. No further


reports to 2006.

Gaskell, A.J. *et al.*, *Tet. Lett.*, 1970, 77-82 (*uv*, *ir*, *pmr*, *ms*)

Cohylakis, D. *et al.*, *J.C.S. Perkin 1*, 1980, 1290-1293 (*synth*, *uv*, *ir*, *pmr*, *ms*)

Subsessiline† S-627

Splendaboline
[49620-02-2]



$C_{19}H_{15}NO_5$ 337.331

Alkaloid from the bark of *Guatteria subsessilis*, the stem bark of *Guatteria ourougou*, and from *Telitoxicum peruvianum* (Annonaceae, Menispermaceae). Shows antimicrobial activity. Deep red needles (Me₂CO). Mp 247-249° (241-243°).

O¹-De-Me, O⁹-Me ether: **Sinofranine**

$C_{19}H_{15}NO_5$ 337.331

Alkaloid from the dried stems of *Sinofranchetia chinensis*. Yellowish cryst. Mp 193-195°.

Hasegawa, M. *et al.*, *Acta Cient. Venez.*, 1972, **23**, 165; *CA*, **79**, 42716z (*isol*)

Skiles, J.W. *et al.*, *J.O.C.*, 1979, **44**, 409 (*synth*, *ir*, *pmr*, *ms*, *struct*)

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 320 (*isol*)


Leboeuf, M. *et al.*, *Planta Med.*, 1983, **48**, 234 (*isol*, *pmr*)

Villar, A. *et al.*, *Planta Med.*, 1986, **52**, 556-557 (*activity*)

Xu, C. *et al.*, *Fitoterapia*, 2004, **75**, 239-241 (*Sinofranine*)

Succinopyoverdin G173 S-628

[535994-95-7]



$C_{49}H_{68}N_{13}O_{20}^{\oplus}$ 1159.151

Prod. by *Pseudomonas fluorescens* G173. Siderophore.

Fernandez, D.U. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 1-10 (*isol*, *pmr*, *cmr*, *ms*)

Succirubine S-629

[1361-38-2]

$C_{19}H_{24}N_2O_2$ 312.411

Struct. unknown. Alkaloid from the bark of *Cinchona succirubra* (Rubiaceae). Mp

80-81°. $[\alpha]_D^{30}$ +1.63 (EtOH).

Quadrat-i-Khuda, M. *et al.*, *Sci. Res. (Dacca)*, 1965, **2**, 1-7; *CA*, **63**, 12004g

Suffruticodine S-630

[1361-40-6]

$C_{13}H_{15}NO_3$ 233.266

Struct. unknown. Alkaloid from *Securinega suffruticosa* (Euphorbiaceae). Mp 120-122°. Opt. inactive. Ir 3048, 1756 and 1636 cm^{-1} (OH or NH, C=O and C=C).

Murav'eva, V.I. *et al.*, *Zh. Obshch. Khim.*, 1963, **33**, 693-694; *CA*, **59**, 2884h (*isol*)

Suffruticonine S-631

[1361-41-7]


$C_{13}H_{17}NO_2$ 219.283

Struct. unknown. Alkaloid from *Securinega suffruticosa* (Euphorbiaceae). Mp 237-238°. Opt. inactive. Ir 3050 (OH or NH), 1773 (C=O), 1655 (C=C) cm^{-1} .

Murav'eva, V.I. *et al.*, *Zh. Obshch. Khim.*, 1963, **33**, 693-694; *CA*, **59**, 2884h (*isol*)

Suinin S-632

[175993-00-7]



$C_{31}H_{37}NO_5$ 503.637


Prod. by a mutant of *Streptomyces nasri*. Antibiotic. Reddish needles.

Gohar, Y.M. *et al.*, *CA*, 1996, **124**, 311910g (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Sulfinemycin S-633

14-Methyl-2-pentadecenethioamide S-oxide, 9CI

[162857-73-0]



$C_{16}H_{31}NOS$ 285.493


Prod. by *Streptomyces albus*. Anthelmintic agent. Yellow powder. Sol. MeOH, $CHCl_3$, CH_2Cl_2 ; poorly sol. Et₂O, hexane. Mp 84°. Dec. at 105°. λ_{\max} 210; 247; 343 (MeOH) (Derep).

Lee, T.M. *et al.*, *J. Antibiot.*, 1995, **48**, 282 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)

Sulfobacin A S-634

Flavocristamide B

[169217-35-0]



C₃₄H₆₉NO₆S 619.988

Prod. by *Chryseobacterium* sp. NR 2993 and from a *Flavobacterium* sp. in the marine bivalve *Cristaria plicata*. Von Willebrand factor receptor antagonist. Powder. $[\alpha]_D^{24}$ -35 (c, 0.1 in MeOH).

4,5-Didehydro(E-): Flavocristamide A
[169217-34-9]

C₃₄H₆₇NO₆S 617.972

Prod. by a *Flavobacterium* sp. in the marine bivalve *Cristaria plicata*. DNA polymerase inhibitor. Amorph. solid. $[\alpha]_D^{20}$ -17 (c, 0.3 in MeOH).

Kamiyama, T. *et al.*, *J. Antibiot.*, 1995, **48**, 924-928; 929-936 (*isol, ir, pmr, cmr*)

Kobayashi, J. *et al.*, *Tetrahedron*, 1995, **51**, 10487-10490 (*isol, ir, pmr, cmr, ms*)

Irako, N. *et al.*, *Tet. Lett.*, 1998, **39**, 5793-5796 (*synth*)

Takikawa, H. *et al.*, *J.C.S. Perkin 1*, 1999, 2467-2477 (*synth*)

Shioiri, T. *et al.*, *Tetrahedron*, 2000, **56**, 9129-9142 (*synth*)

Labeeuw, O. *et al.*, *Tet. Lett.*, 2003, **44**, 6383-6386 (*synth*)


Gupta, P. *et al.*, *Tet. Lett.*, 2004, **45**, 9641-9643 (*synth*)

Sharma, A. *et al.*, *Tet. Lett.*, 2007, **48**, 3705-3707 (*synth*)

Sulfobacin B

S-635

[170242-21-4]

C₃₂H₆₅NO₅S 575.935

Prod. by *Chryseobacterium* sp. NR 2993. von Willebrand factor receptor antagonist. Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. $[\alpha]_D^{24}$ -19 (c, 0.1 in MeOH). λ_{max} (MeOH) (Berdy).

Kamiyama, T. *et al.*, *J. Antibiot.*, 1995, **48**, 924; 929 (*isol, ir, pmr, cmr, ms, props*)

Takikawa, H. *et al.*, *J.C.S. Perkin 1*, 1999, 2467-2477 (*synth*)


Shioiri, T. *et al.*, *Tetrahedron*, 2000, **56**, 9129-9142 (*synth*)

N-(2-Sulfoethyl)pyridinium betaine

S-636

2-Pyridinioethanesulfonic acid. **Pyridinebetaine B**

[24020-66-4]

C₇H₉NO₃S 187.219

Isol. from the sponge *Agelas dispar*. Cryst. (EtOH). Mp 260°. λ_{max} 222 (ε 6500); 260 (ε 4500) (MeOH).

Le Berre, A. *et al.*, *Bull. Soc. Chim. Fr.*, 1970, 954 (*synth, pmr*)


King, J.F. *et al.*, *Can. J. Chem.*, 1984, **62**, 1977-1995 (*synth*)

Cafieri, F. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1171-1173 (*isol, uv, ir, pmr, cmr*)

Sulfohalopterin 2

S-637

6,6'-Oxybis[7-(2-amino-1,4-dihydro-4-oxo-6-pteridinyl)-6,7-dideoxy-D-glycero-L-galacto-heptonic acid] 3,3'-bis(hydrogen sulfate), 9CI. **SHP 2**
[109686-75-1]

C₂₆H₃₂N₁₀O₂₁S₂ 884.725


Isol. from *Halobacterium marismortui*. No phys. props. reported.

Lin, X. *et al.*, *Biochemistry*, 1987, **26**, 6211 (*isol, pmr, cmr, cd, struct*)

Sulfomycin

S-638

[65454-59-3]



Prod. by *Streptomyces viridochromogenes* ssp. *sulfomycini*. Active against gram-positive bacteria. Closely related to Berninamycin A, B-104. λ_{max} 255 (ε 82400); 330 (sh) (ε 14400) (MeOH/NaOH) (Derep). λ_{max} 252 (ε 93300); 325 (sh) (ε 14400) (MeOH) (Derep).
► WR0176000

Sulfomycin I [62339-99-5]C₅₄H₅₂N₁₆O₁₆S₂ 1245.234

Prisms. Mp 280°. $[\alpha]_D^{20}$ -16 (c, 2 in MeOH).

N⁴⁹-Parent amide: **Antibiotic 10381V**
[164366-78-3]

C₄₅H₄₃N₁₃O₁₃S₂ 1038.046

Prod. by *Streptomyces arginensis*. Yellow powder. Sol. MeOH, EtOAc.

N⁵²-Parent amide: **Antibiotic 10381Y**

[164300-77-0]

C₄₈H₄₆N₁₄O₁₄S₂ 1107.109

Prod. by *Streptomyces arginensis*. Yellow powder. Sol. MeOH, EtOAc.

21-Demethoxy: **21-Demethoxysulfomycin I. Pre-B**

[164366-77-2]

C₅₃H₅₀N₁₆O₁₅S₂ 1215.208

Prod. by *Streptomyces arginensis*. Yellow powder. Sol. MeOH, EtOAc.

5-Methyl: **Methylsulfomycin I**

[202606-10-8]

C₅₅H₅₄N₁₆O₁₆S₂ 1259.261

Prod. by *Streptomyces* sp. HIL Y-9420704. Powder. $[\alpha]_D^{25}$ -57.1 (c, 0.07 in MeOH). Mp >250° dec. λ_{max} 250; 320 (MeOH).

Sulfomycin II [102489-42-9]C₅₄H₅₂N₁₆O₁₅S₂ 1229.235

$[\alpha]_D^{25}$ -11.8 (MeOH). Dec. at 190°.

Sulfomycin III [102489-41-8]C₅₃H₅₀N₁₆O₁₆S₂ 1231.208

$[\alpha]_D^{25}$ +3.2 (MeOH). Dec. at 183°.

65-Deoxy, 21-O-de-Me: **Antibiotic 10381Z₁**

[164366-75-0]

C₅₂H₄₈N₁₆O₁₅S₂ 1201.181

Prod. by *Streptomyces arginensis*. Yellow powder. Sol. MeOH, EtOAc.

65-Deoxy, 21-demethoxy: **Antibiotic 10381Z₂**

[164366-76-1]

C₅₂H₄₈N₁₆O₁₄S₂ 1185.182

Prod. by *Streptomyces arginensis*. Yellow powder. Sol. MeOH, EtOAc.

Egawa, Y. *et al.*, *J. Antibiot.*, 1969, **22**, 12-17 (*isol, props*)

Abe, H. *et al.*, *Tet. Lett.*, 1977, 735-736; 1978, 2791-2794; 1988, **29**, 1401-1404 (*struct*)

Pat. Coop. Treaty (WIPO), 1995, 95 07 292; CA, **123**, 54272s (*Antibiotics 10381*)

Kohno, J. *et al.*, *J. Antibiot.*, 1996, **49**, 1063-1065 (*Sulfomycins II, III, pmr, cmr, struct*)

Martin, G.E. *et al.*, *Magn. Reson. Chem.*, 1998, **36**, 635-644 (*N-15 nmr*)

Kumar, E.K.S.V. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1562-1564 (*Methylsulfomycin I*)


Sulphostin

S-639

Sulphostin

[307345-51-3]

[307345-53-5]



Absolute Configuration

C₅H₁₃N₄O₅PS 272.221


Prod. by *Streptomyces* sp. MK251-43F3. Inhibitor of dipeptidyl peptidase IV. Cryst. (EtOH aq.) (synthetic). Mp 203-208° dec. (synthetic). $[\alpha]_D^{23}$ -21.5 (c, 1 in H₂O) (synthetic). Readily epimerises at C-3. Isol. as a mixt. with the C-3 epimer.

Akiyama, T. *et al.*, *J. Antibiot.*, 2001, **54**, 744-746 (*isol*)

Abe, M. *et al.*, *J. Nat. Prod.*, 2004, **67**, 999-1004 (*synth, pmr, cmr, P-31 nmr, cryst struct*)

Abe, M. *et al.*, *J. Antibiot.*, 2005, **58**, 111-117 (activity)

Sulpinine B S-640
[139975-52-3]




$C_{32}H_{41}NO_3$ 487.681
Metab. of *Aspergillus sulphureus*. Shows antiinsect props. Orange oil. $[\alpha]_D -43.2$ ($CHCl_3$). λ_{max} 235 (ϵ 22600); 283 (ϵ 5600) (MeOH) (Derep).

18 α ,19 α -Epoxide: Sulpinine A
[139975-51-2]
 $C_{32}H_{41}NO_4$ 503.68
Metab. of *Aspergillus sulphureus*. Yellow-orange solid. Mp 149-152° dec. $[\alpha]_D -29.6$ ($CHCl_3$). λ_{max} 235 (ϵ 22600); 283 (ϵ 5600) (MeOH) (Derep).

Laakso, J.A. *et al.*, *J.O.C.*, 1992, **57**, 2066 (*isol*, *pmr*, *cmr*)


Sulpinine C S-641
[139975-53-4]



$C_{32}H_{41}NO_6$ 535.679
Metab. of *Aspergillus sulphureus*. Shows antiinsect props. Yellow oil. $[\alpha]_D +17.1$ ($CHCl_3$). λ_{max} 229 (ϵ 25600); 262 (ϵ 18200) (MeOH) (Derep).

Laakso, J.A. *et al.*, *J.O.C.*, 1992, **57**, 2066 (*isol*, *pmr*, *cmr*)

Sungucine S-642
[73020-53-8]



Absolute Configuration

$C_{42}H_{42}N_4O_2$ 634.819
Alkaloid from the root bark of *Strychnos icaja* (Loganiaceae). Shows antiplasmodial, antimalarial and cytotoxic activities. Prisms ($Me_2CO/CHCl_3$). Mp 350°. λ_{max} 220 ($\log \epsilon$ 4.38); 265 ($\log \epsilon$ 3.88); 292 ($\log \epsilon$ 4.03); 302 ($\log \epsilon$ 4.06) (MeOH).


16,17-Didehydro, 17,23-dihydro: Isosungucine
[279684-12-7]
 $C_{42}H_{42}N_4O_2$ 634.819
Alkaloid from *Strychnos icaja*. Amorph. powder. C-23 config. not determined. λ_{max} 220 ($\log \epsilon$ 4.38); 230 ($\log \epsilon$ 4.24); 252 ($\log \epsilon$ 4.04); 292 ($\log \epsilon$ 3.86); 302 (3.74) (MeOH).

18-Hydroxy: 18-Hydroxysungucine
[279258-77-4]
 $C_{42}H_{42}N_4O_3$ 650.819
Alkaloid from *Strychnos icaja*. Amorph. powder. λ_{max} 220 ($\log \epsilon$ 4.38); 265 ($\log \epsilon$ 3.88); 292 ($\log \epsilon$ 4.03); 302 ($\log \epsilon$ 4.06) (MeOH).

18-Hydroxy, 16,17-didehydro, 17,23-dihydro: 18-Hydroxyisosungucine
[279258-78-5]
 $C_{42}H_{42}N_4O_3$ 650.819
Alkaloid from *Strychnos icaja*. Amorph. powder. λ_{max} 220 ($\log \epsilon$ 4.38); 230 ($\log \epsilon$ 4.24); 252 ($\log \epsilon$ 4.04); 292 ($\log \epsilon$ 3.86); 302 (3.74) (MeOH).

Lamotte, J. *et al.*, *Tet. Lett.*, 1979, 4227-4228 (*isol*, *uv*, *ir*, *pmr*, *ms*, *cryst struct*, *abs config*)
Frederich, M. *et al.*, *Planta Med.*, 2000, **66**, 262-269 (*isol*, *activity*, *uv*, *ir*, *cd*, *pmr*, *cmr*, *ms*)
Frédérich, M. *et al.*, *J. Nat. Prod.*, 2001, **64**, 12-16 (*activity*)


Suomilide S-643
[195256-59-8]



$C_{45}H_{73}N_7O_{19}S$ 1048.173
Isol. from the marine cyanobacterium *Nodularia spumigena* HKUV. Enzyme inhibitor. Amorph. powder. $[\alpha]_D^{26} +74.2$ (c, 0.1 in MeOH).

Sivonen, K. *et al.*, *Hydrobiologia*, 1989, **185**, 3-8 (*isol*)
Fujii, K. *et al.*, *Tet. Lett.*, 1997, **38**, 5529-5532 (*isol*, *pmr*, *cmr*, *ms*)

Supinidine S-644
2,3,5,7a-Tetrahydro-1H-pyrrolizine-7-methanol, 9Cl. 1-Hydroxymethyl-1,2-didehydropyrrolizidine



(R)-form

$C_8H_{13}NO$ 139.197
► Exp. carcinogen.

(R)-form
Necine component of Cynaustine in A-669. Rare in nature. Oil. Bp_{0.14} 85°

(bulb). $[\alpha]_D^{20} +9.2$ (c, 2.07 in EtOH).
Picrate:
Yellow needles (EtOH). Mp 144-144.5°.


(S)-form [551-59-7]
Obt. by hydrol. of Supinine in A-669. $[\alpha]_D -10.3$.
O-[2S-Hydroxy-2-(1S-hydroxyethyl)-4-methylpentanoyl], N-oxide:
 $C_{16}H_{27}NO_5$ 313.393
Alkaloid from the roots of *Anchusa strigosa*. Orange oil. $[\alpha]_D^{25} -4.3$ (c, 0.1 in MeOH). λ_{max} 230 (sh); 270 (MeOH).

Me ether: 1-Methoxymethyl-1,2-dehydropyrrolizidine
[6029-76-1]
 $C_9H_{15}NO$ 153.224
Alkaloid from *Crotalaria trifoliastrum* and *Crotalaria medicaginea* (Fabaceae). Oil. Bp₁₀ 100°. $[\alpha]_D^{20} -24$ (c, 4.88 in EtOH).

Me ether, picrate: Mp 153-154°.

(±)-form [23185-51-5]
Cryst. (MeOH) (as picrate). Mp 124-126° (picrate).
Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1962, **15**, 121 (*isol*, *deriv*)
Sawhney, R.S. *et al.*, *J. Indian Chem. Soc.*, 1970, **47**, 741 (*isol*, *deriv*)
Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653 (*cd*)
Tufariello, J.J. *et al.*, *J.O.C.*, 1975, **40**, 3866 (*synth*, *bibl*)
Mody, N.V. *et al.*, *J. Nat. Prod.*, 1979, **42**, 417 (*cmr*)
Chamberlin, A.R. *et al.*, *Tet. Lett.*, 1982, **23**, 2619 (*synth*)
Hart, D.J. *et al.*, *Tet. Lett.*, 1982, **23**, 2761 (*synth*)
Macdonald, T.L. *et al.*, *J.O.C.*, 1983, **48**, 1129 (*synth*)
Burnett, D.A. *et al.*, *J.A.C.S.*, 1984, **106**, 8201 (*synth*)
Boynton, C.M. *et al.*, *J.C.S. Perkin 1*, 2000, 3599-3602 (*synth*)
Braca, A. *et al.*, *Planta Med.*, 2003, **69**, 835-841 (*Anchusa strigosa ester*)
Sarkar, T.K. *et al.*, *Tetrahedron*, 2005, **61**, 1155-1165 (*synth*)
Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 1005

Surugatoxin S-645
[40957-92-4]



$C_{25}H_{26}BrN_5O_{13}$ 684.41
Unique skeleton containing both indole and pteridine subunits. Toxin from the mid-gut gland of the Japanese Ivory shell *Babylonia japonica* (harvested from

a particular area only). Prod. from cyanobacteria. Marine toxin, causing shellfish poisoning. Prisms + 7H₂O. Mp 300°. Artifact formed during isol. λ_{\max} 276 (ε 15000) (0.1N HCl) (Derep). λ_{\max} 279 (ε 19000) (0.1N NaOH) (Derep). λ_{\max} 276 (ε 15000) (H₂O) (Derep).

► Toxic.

Kosuge, T. *et al.*, *Tet. Lett.*, 1972, 2545-2548 (isol, uv, ir, cryst struct)

Inoue, S. *et al.*, *Tet. Lett.*, 1984, **25**, 4407 (synth)


Okado, K. *et al.*, *Yakugaku Zasshi*, 1985, **105**, 375; 381 (synth)

Inoue, S. *et al.*, *Tetrahedron*, 1994, **50**, 2729 (synth)

Sutchueneneoneine

S-646

[151271-92-0]



R¹ = H, R² = OH

C₃₆H₄₀N₂O₆ 596.722

Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae). Powder. $[\alpha]_D^{26}$ +6.7 (c, 0.422 in EtOH).

Wang, X.-K. *et al.*, *Phytochemistry*, 1993, **33**, 1253 (isol, uv, ir, pmr, cmr, ms, struct)

Sutchuenenine

S-647

[151271-90-8]

As Sutchueneneoneine, S-646 with

R¹ = OH, R² = H

C₃₆H₄₀N₂O₆ 596.722


Alkaloid from roots of *Cyclea sutchuenensis* (Menispermaceae). Yellowish powder. $[\alpha]_D^{26}$ -47.4 (c, 0.703 in CHCl₃). Suggested to have 1R,1'R config.

Wang, X.-K. *et al.*, *Phytochemistry*, 1993, **33**, 1253 (isol, uv, ir, pmr, cmr, ms, struct)

Sutchuenensine

S-648

[152406-26-3]



C₃₈H₄₂N₂O₆ 622.76

Alkaloid from the roots of *Cyclea sutchuenensis*. Powder. $[\alpha]_D^{27}$ -110 (c, 0.13 in EtOH).


Lai, S. *et al.*, *Yaouxue Xuebao*, 1993, **28**, 599-603

Suvanine

S-649

[94203-53-9]

[202072-67-1]



C₂₅H₃₈O₅S 450.638

Constit. of sponges of the genera *Coscinoderma* and *Hippospongia*. Powder (as N,N-dimethylguanidine salt). Mp 224° (dimethylguanidine salt). $[\alpha]_D$ +9.5 (MeOH). Also isol. as the trimethylguanidinium salt. λ_{\max} 210 (ε 6300) (MeOH) (Derep).

Manes, L.V. *et al.*, *J.O.C.*, 1985, **50**, 284; 1988, **53**, 570 (struct)

Kimura, J. *et al.*, *J. Nat. Prod.*, 1998, **61**, 248-250; 862 (isol)


Swainsonine

S-650

Octahydro-1,2,8-indolizinetriol, 9CI.

1,2,8-Trihydroxyoctahydroindolizine. **Tri-dolgosir, INN**

[72741-87-8]



C₈H₁₅NO₃ 173.211

Alkaloid from *Swainsona canescens* and *Astragalus lentiginosus*. Also isol. from seeds of *Astragalus oxyphythus* (Fabaceae) and prod. by the fungus *Rhizoctonia leguminicola*. Neurological α-mannosidase inhibitor. Potent toxin producing symptoms similar to the genetic disorder mannosidosis. Antineoplastic agent. Needles (CHCl₃). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 144-145° dec. $[\alpha]_D^{20}$ -85.1 (c, 0.6 in MeOH) (synthetic). pK_a 7.4. The *Rhizoctonia leguminicola* isol. was originally assigned an isomeric struct. (Guengerich *et al.*, 1973, revised by Schneider *et al.*, 1983).

Hydrochloride: Tri-dolgosir hydrochloride, USAN. GD 0039

[214462-68-7]

Cryst. (MeOH/Et₂O). Sol. H₂O (3g/ml). Mp 190-191°.

N-Oxide: Swainsonine N-oxide

[81759-44-6]

C₈H₁₅NO₄ 189.211

Alkaloid from above-ground parts of *Astragalus lentiginosus* (Fabaceae). Hygroscopic, pale yellow prisms (Me₂CO). Mp 161-163°.

Tri-Ac:

Pale yellow oil. Dec. rapidly on standing in air.

8-Deoxy: see Octahydro-1,2-indolizinediol, O-46

1-Epimer: Synthetic. Cryst. (MeCN/MeOH). Mp 111-113°. $[\alpha]_D^{20}$ -32.14 (c,

1.5 in MeOH).

8-Epimer: Synthetic. Mp 92-93°. $[\alpha]_D^{20}$ -22.1 (c, 0.25 in MeOH).

8a-Epimer: Synthetic. α-D-Mannosidase inhibitor. Cryst. (CHCl₃). Mp 122-124° dec. $[\alpha]_D^{19}$ -64.5 (c, 0.95 in MeOH).

2,8a-Diepimer: Synthetic. Mp 138-142° dec. $[\alpha]_D^{26}$ -24 (c, 1.14 in MeOH).

8,8a-Diepimer: Synthetic. Potent inhibitor of human lysosomal α-D-mannosidase. Cryst. (CHCl₃/hexane). Mp 130-131° dec. $[\alpha]_D^{22}$ -21.2 (c, 0.78 in MeOH).

2,8,8a-Triepimer: Synthetic. Glass. $[\alpha]_D^{20}$ +31.62 (c, 1.07 in MeOH).

[41431-34-9]

Colegate, S.M. *et al.*, *Aust. J. Chem.*, 1979, **32**, 2257 (isol, ir, pmr, cmr, ms, struct)

Skelton, B.W. *et al.*, *Aust. J. Chem.*, 1980, **33**, 435 (cryst struct)

Schneider, M.J. *et al.*, *J.A.C.S.*, 1982, **104**, 6863 (biosynth)

Molyneux, R.J. *et al.*, *Science (Washington, D.C.)*, 1982, **216**, 190 (oxide)

Schneider, M.J. *et al.*, *Tetrahedron*, 1983, **39**, 29 (struct)

Yasuda, N. *et al.*, *Chem. Lett.*, 1984, 1201 (synth)

Fleet, G.W.J. *et al.*, *Tet. Lett.*, 1984, **25**, 1853 (synth)

Suami, T. *et al.*, *Carbohydr. Res.*, 1985, **136**, 67 (synth)

Ali, M.H. *et al.*, *Carbohydr. Res.*, 1985, **136**, 225 (synth)

Adams, C.E. *et al.*, *J.O.C.*, 1985, **50**, 420 (synth)

Setoi, H. *et al.*, *J.O.C.*, 1985, **50**, 3948 (synth)

Howard, A.S. *et al.*, *Alkaloids (Academic Press)*, 1986, **28**, 287 (rev. pharmacol, tox)

Elbein, A.D. *et al.*, *Alkaloids: Chem. Biol. Perspect.*, 1987, **5**, 1 (rev)

Ikota, N. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2140; 1988, **36**, 1143 (synth)

Bashyal, B.P. *et al.*, *Tetrahedron*, 1987, **43**, 3083 (synth, ir, pmr, ms)

Tadano, K. *et al.*, *J.O.C.*, 1988, **53**, 5209 (synth, ir, pmr, cmr)

Dener, J.M. *et al.*, *J.O.C.*, 1988, **53**, 6022 (synth, ir, ms)

Bennett, R.B. *et al.*, *J.A.C.S.*, 1989, **111**, 2580 (synth, pmr, cmr)

Kim, Y.G. *et al.*, *Tet. Lett.*, 1989, **30**, 5721 (synth, epimers)

Carpenter, N.M. *et al.*, *Tet. Lett.*, 1989, **30**, 7261 (synth)

Ikota, N. *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 2712 (synth)

Miller, S.A. *et al.*, *J.A.C.S.*, 1990, **112**, 8100 (synth)

Pearson, W.H. *et al.*, *Tet. Lett.*, 1990, **31**, 7571 (synth)

Kardono, L.B.S. *et al.*, *Phytochem. Anal.*, 1991, **2**, 120 (isol, pmr, cmr)

Gonzalez, F.B. *et al.*, *Bull. Chem. Soc. Jpn.*, 1992, **65**, 567 (synth)

Casiraghi, G. *et al.*, *J.O.C.*, 1993, **58**, 3397 (synth, epimers)

Honda, T. *et al.*, *J.C.S. Perkin 1*, 1994, 2091 (synth)

Naruse, M. *et al.*, *J.O.C.*, 1994, **59**, 1358 (synth)

Zhou, W.-S. *et al.*, *Tet. Lett.*, 1995, **36**, 1291 (synth)

Mukai, C. *et al.*, *J.O.C.*, 1998, **63**, 6281-6287 (synth)

Pat. Coop. Treaty (WIPO), 1998,


((Glycodesign))98 46 602; *CA*, **129**, 306525d (tridolgosir hydrochloride)

Klein, J.L. *et al.*, *Br. J. Cancer*, 1999, **80**, 87-95 (pharmacol)

- Trost, B.M. *et al.*, *Chem. Eur. J.*, 1999, **5**, 3279-3284 (*synth*)
- Misago, M. *et al.*, *Biochem. Biophys. Res. Commun.*, 2000, **269**, 219-225 (*pharmacol*)
- Bohle, D.S. *et al.*, *J.O.C.*, 2000, **65**, 5685-5692 (*synth*)
- El Nemr, A. *et al.*, *Tetrahedron*, 2000, **56**, 8579-8629 (*rev. synth*)
- Zhao, H. *et al.*, *J.O.C.*, 2001, **66**, 1761-1767 (*synth, pmr, cmr*)
- Buschmann, N. *et al.*, *J.O.C.*, 2002, **67**, 4325-4329 (*synth*)
- Lindsay, K.B. *et al.*, *J.O.C.*, 2002, **67**, 4325-4329 (*synth*)
- Kumar, N.S. *et al.*, *J.O.C.*, 2006, **71**, 1262-1264 (*synth*)
- Au, C.W.G. *et al.*, *J.O.C.*, 2006, **71**, 7097-7099 (*synth*)
- Nath, M. *et al.*, *Org. Lett.*, 2006, **8**, 317-320 (*synth*)
- Déschamps, I. *et al.*, *Tetrahedron*, 2007, **63**, 9082-9091 (*synth*)
- Abrams, J.N. *et al.*, *J.O.C.*, 2008, **73**, 1935-1940 (*epimer, synth*)
- Håkansson, A.E. *et al.*, *Tet. Lett.*, 2008, **49**, 179-184 (*synth*)
- Guo, H. *et al.*, *Tetrahedron*, 2008, **64**, 304-313 (*synth*)
- Shi, G.-F. *et al.*, *Tetrahedron*, 2008, **64**, 5005-5012 (*synth*)

Swazine**S-651**

[38763-74-5]



Absolute configuration


C₁₈H₂₃NO₆ 349.383

Cyclic retronecine diester. Alkaloid from *Senecio swaziensis* (Asteraceae). Needles (Me₂CO). Mp 165°. [α]_D²⁰ -104 (c, 1.16 in EtOH).

- Laing, M. *et al.*, *Tet. Lett.*, 1972, 5183 (*cryst struct*)
- Gordon-Gray, C.G. *et al.*, *J.C.S. Perkin 1*, 1974, 1556 (*isol, struct*)
- Drewes, S.E. *et al.*, *J.C.S. Perkin 1*, 1981, 287 (*cmr*)
- White, J.D. *et al.*, *Chem. Comm.*, 1998, 603-604; 1237 (*abs config, cryst struct*)

Swinhoeamide A**S-652**

[209408-31-1]

C₄₀H₆₅N₂O₁₂P 796.934

Related to Calyculin A, C-55 and Clavosine A, C-516. Isol. from the sponge *Theonella swinhoei*. Insecticide and antifungal agent. Cell proliferation inhibitor. Powder. [α]_D²⁰ -21.6 (c, 0.35 in EtOH).

λ_{max} 223 (MeOH).**O²¹-De-Me: Geometricin A**

[443769-46-8]

C₃₉H₆₃N₂O₁₂P 782.907

Isol. from the sponge *Luffariella geometrica*. Cytotoxic. Amorph. solid. [α]_D²³ -36.3 (c, 0.29 in MeOH). λ_{max} 206 (ε 16280); 221 (ε 16370) (MeOH).

- Kehraus, S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1056-1058 (*Geometricin A*)
- Edrada, R.A. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1168-1172 (*Swinhoeamide A*)

Sylvatine**S-653**

5-(1,3-Benzodioxol-5-yl)-N-(10-methyl-5-undecenyl)-2,4-pentadienamide, 9CI


C₂₄H₃₃NO₃ 383.53**(E,E,E)-form [42438-80-2]**

Alkaloid from *Piper aurantiacum*, *Piper brachystachyum*, *Piper sylvaticum*, *Piper chaba* (Javanese long pepper) and several other *Piper* spp. (Piperaceae). Needles (C₆H₆/hexane). Mp 116-117° (111-112°).

- Banerji, A. *et al.*, *Tetrahedron*, 1973, **29**, 977 (*isol, uv, ir, pmr, ms, struct*)
- Patra, A. *et al.*, *Phytochemistry*, 1974, **13**, 2889 (*isol*)
- Rao, J.M. *et al.*, *Curr. Sci.*, 1974, **43**, 76 (*isol, uv, ir, pmr*)
- Dutta, C.P. *et al.*, *Phytochemistry*, 1975, **14**, 2090 (*isol*)
- Vig, O.P. *et al.*, *Indian J. Chem.*, 1975, **13**, 225 (*synth, ir, pmr*)

Symbioimine**S-654**

[862479-86-5]



Absolute Configuration

C₁₉H₂₃NO₅S 377.46

Alkaloid from the marine dinoflagellate *Symbiodinium* sp.; also from *Amphiscollops* sp. Inhibitor of cyclooxygenase-2 and osteoclast differentiation. Cryst. + 1H₂O (H₂O). Mp 214-215° dec. [α]_D²⁷ +245 (c, 0.1 in DMSO).

- Kita, M. *et al.*, *J.A.C.S.*, 2004, **126**, 4794-4795 (*isol*)
- Kita, M. *et al.*, *Bioorg. Med. Chem.*, 2005, **13**, 5253-5258 (*isol, pmr, cmr, activity*)
- Varseev, G.N. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 4767-4771 (*synth*)
- Zou, Y. *et al.*, *Org. Lett.*, 2006, **8**, 5605-5608 (*synth*)
- Kim, J. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 3104-3106 (*synth*)

Synaine**S-655**C₂₄H₃₉NO 357.578


Struct. unknown. Alkaloid from the

roots and rhizomes of *Veratrum album* (Liliaceae). Mp 217-218° dec. [α]_D²⁰ -31.68 (c, 1 in CHCl₃). Proposed mol. formula seems improbable as it is prob. a steroidal alkaloid.

- Cionga, E. *et al.*, *Acta Pol. Pharm.*, 1957, **14**, 73-76; *CA*, **52**, 12882c
- Cionga, E. *et al.*, *Ann. Pharm. Fr.*, 1958, **16**, 511; *CA*, **53**, 5590h


Syncarpurea**S-656**

2,3,7b,10-Tetrahydro-7,7,9,9-tetramethylpyrrolo[1,2-c]quinazoline-5,8,10(1H,6H,9H)-trione, 9CI [89759-17-1]

C₁₅H₂₀N₂O₃ 276.335

Metab. from *Uvaria afzelii* (Annonaceae). Cryst. (CHCl₃). Mp 239-240°. λ_{max} 236 (ε 4310); 306 (ε 5520); 362 (MeOH/NaOH) (Derep). λ_{max} 236 (ε 4310); 306 (ε 5520) (MeOH) (Derep).

- Hufford, C.D. *et al.*, *Tet. Lett.*, 1984, **25**, 371 (*isol, uv, ir, pmr, ms, cryst struct*)

Synechobactin A**S-657**C₂₆H₄₈N₄O₉ 560.687


Isol. from the marine cyanobacterium *Synechococcus* sp. PCC 7002. Siderophore. Related to Schizokinens.

N¹-Deacyl, N¹-octanoyl: Synechobactin CC₂₂H₄₀N₄O₉ 504.579Isol. from *Synechococcus* sp. PCC 7002.**N¹-Deacyl, N¹-decanoyl: Synechobactin B**C₂₄H₄₄N₄O₉ 532.633Isol. from *Synechococcus* sp. PCC 7002.

- Ito, Y. *et al.*, *Limnol. Oceanogr.*, 2005, **50**, 1918-1923 (*isol, struct*)

Syneilesine**S-658**

15,20-Dihydro-12,14-dihydroxy-4-methyl-4,8-secosenecionan-8,11,16-trione, 9CI [55652-65-8]



Absolute configuration

C₁₉H₂₉NO₇ 383.441

Cyclic otonecine diester. Alkaloid from *Syneilesis palmata* (Asteraceae). Needles (petrol). Mp 194.5-195°.

► Poss. hepatocarcinogen. VS3595500

O¹⁴-Ac: **Acetylsyneilesine**

[62387-18-2]

C₂₁H₃₁NO₈ 425.478

From *Syneilesis palmata* (Asteraceae). Oil.

Hikichi, M. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 3178 (*isol, cd, cmr, pmr, struct*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173 (*cmr*)


Mori, H. *et al.*, *Cancer Res.*, 1985, **45**, 3125 (*tox*)

Synephrine

S-659

4-Hydroxy- α -[(methylamino)methyl]-benzenemethanol, 9CI. 1-(p-Hydroxyphenyl)-2-methylaminoethanol. p-(α -Hydroxy- β -methylaminoethyl)phenol. *Analeptin*. *Ethaphene*. *Oxedrine*, *BAN*. *Simpalon*. *Sympathol*. Many other names

[94-07-5]



(R)-form

C₉H₁₃NO₂ 167.207

Sympathomimetic agent showing vasoconstrictor, hypertensive and bronchial muscle relaxant props. Possesses decongestant props. Log P -0.09 (calc). Was investigated as a synthetic drug before it was found naturally.

► LD₅₀ (mus, ipr) 1000 mg/kg. DO7350000

(R)-form [614-35-7]

A widely distributed alkaloid, occurs in *Citrus* (Rutaceae) and *Coryphantha* (Cactaceae) spp., in *Ficus bengalensis* and a no. of Amaryllidaceae spp., also in *Haloxylon salicornicum* (Chenopodiaceae) (opt. rotn. of isolates not reported in many cases). Plates (Me₂CO aq. or H₂O). Mp 162-164° dec. [α]_D²⁵ -55.6 (c, 0.5 in M HCl).

► DO7175003

4-Me ether: **Longimammine**. 4-O-Methylsynephrine

[57286-93-8]

C₁₀H₁₅NO₂ 181.234

Alkaloid from *Dolichothele longimamma* (Cactaceae). Mp 144-146° (hydrochloride). [α]_D²⁵ -36.

N,O-Isopropylidene: **Synephrine acetone**

[225928-85-8]

C₁₂H₁₇NO₂ 207.272

Alkaloid from the seeds of *Casimiroa edulis* (Mexican apple). Hypertensive.

(±)-form [582-84-3]

Alkaloid from a number of *Coryphantha* spp. and from *Dolichothele longimamma* (Cactaceae). Spar. sol. org. solvs. Mp 184-185° dec. Used mainly as tartrate.

Hydrochloride: [5985-28-4]

Cryst. powder. Mp 149.5-157°.

► DO7700000

Tartrate (2:1): [16589-24-5]

Cryst. Mp 188-190°.

Di-Ac: Mp 157°.

Dibenzoyl: Mp 176°.

β -Me ether: **β -O-Methylsynephrine**

[60094-92-0]

[25006-35-3]

C₁₀H₁₅NO₂ 181.234

Alkaloid from *Coryphantha greenwoodii*, *Coryphantha ramulosa* and *Coryphantha cornifera* (Cactaceae); also obt. from tangerine leaves, poss. as artifact. Mp 182-184° (175-176°) (as hydrochloride). One isolate was definitely racemic; opt. rotn. of others not recorded, poss. enantiomeric.

4-Me ether:

C₁₀H₁₅NO₂ 181.234

Synthetic. Mp 117-118° (as hydrochloride).

β -Et ether: **β -O-Ethylsynephrine**

C₁₁H₁₇NO₂ 195.261

Alkaloid from *Dolichothele sphaerica* (Cactaceae). Mp 169-171° (as hydrochloride). Artifact.

[67-04-9]

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 613C (*nmr*)

Priestley, H.M. *et al.*, *J.O.C.*, 1940, **5**, 355 (*synth*)

Corrigan, J.R. *et al.*, *J.A.C.S.*, 1945, **67**, 1894 (*synth*)

Stewart, I.S. *et al.*, *J. Biol. Chem.*, 1964, **239**, 930 (*isol*)

Kappe, T. *et al.*, *J. Med. Chem.*, 1965, **8**, 368 (*uv*)

Wheaton, T.A. *et al.*, *Phytochemistry*, 1969, **8**, 85 (*biosynth*)

McLaughlin, J.L. *et al.*, *J. Pharm. Sci.*, 1972, **61**, 41; 1973, **62**, 408; 411; 1663 (*isol, derivs*)

Ranieri, R.L. *et al.*, *J.O.C.*, 1976, **41**, 319 (*isol, derivs*)

Smith, T.A. *et al.*, *Phytochemistry*, 1977, **16**, 9 (*occur*)

Hengstmann, J.H. *et al.*, *Arzneim.-Forsch.*, 1978, **28**, 2326 (*metab*)

Dattagupta, J.K. *et al.*, *Acta Cryst. B*, 1982, **38**, 2830 (*cryst struct*)

Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., *Akademie-Verlag*, 1987, 1283

Midgley, J.M. *et al.*, *J.C.S. Perkin 2*, 1989, 963 (*cryst struct, abs config*)

Martindale, *The Extra Pharmacopoeia*, 30th edn., *Pharmaceutical Press*, 1993, 1250

Kusu, F. *et al.*, *Chem. Pharm. Bull.*, 1995, **43**, 1158 (*resoln*)


Magos, G.A. *et al.*, *J. Ethnopharmacol.*, 1998, **64**, 35-44; *CA*, **131**, 410 (*acetone*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., *Van Nostrand Reinhold*, 1992, HLV500; SPD000

Synerazol

S-660

8-Benzoyl-2-[3-(1-butenyl)oxiranyl]-9-hydroxy-8-methoxy-3-methyl-1-oxa-7-azaspiro[4.4]non-2-ene-4,6-dione, 9CI [127941-81-5]



Absolute Configuration

C₂₂H₂₃NO₇ 413.426

Prod. by *Aspergillus fumigatus* SANK 10588. Antifungal agent. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. [α]_D²² +22.9 (c, 0.55 in CHCl₃). Related to Pseurotin, P-738. λ_{\max} 202 (ε 23400); 251 (ε 14700); 284 (ε 10300) (EtOH) (Derep). λ_{\max} 251 (E1%/1cm 340); 285 (E1%/1cm 280) (MeOH) (Berdy).

► RN7067000

Japan. Pat., 1989, 89 277 492; *CA*, **113**, 38904 (*isol*)

Ando, O. *et al.*, *J. Antibiot.*, 1991, **44**, 382-389 (*isol, pmr, cmr, ms*)


Igarashi, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 537-540 (*abs config*)

Hayashi, Y. *et al.*, *J.O.C.*, 2005, **70**, 5643-5654 (*synth, abs config*)

Syphilobin A

S-661

[11031-99-5]



C₂₈H₂₂N₂O₂ 418.494


Alkaloid from *Lobelia syphilitica* (Campanulaceae). Needles (Et₂O). Mp 192-194°. λ_{\max} 221 (log ε 4.67); 254 (log ε 4.64); 275 (log ε 4.46); 321 (sh) (log ε 3.33); 337 (log ε 3.47); 353 (log ε 3.51) (no solvent reported).

Tschesche, R. *et al.*, *Tetrahedron*, 1964, **20**, 2885-2893 (*isol, ms, uv, ir*)

Syphilobin F

S-662

[11032-00-1]



C₂₈H₂₆N₂O₃ 438.525

Alkaloid from *Lobelia syphilitica* (Campanulaceae). Needles (C₆H₆). Mp 222-223°. [α]_D²⁰ -1.64 (c, 1.217 in Py). λ_{\max} 274 (log ε 4.14); 280 (log ε 4.14) (no solvent reported).

Ac:

Plates (Me₂CO). Mp 124-125°.

Me ether:


Plates (Me₂CO). Mp 179-180°.

Methiodide:


Needles. Mp 244-246°.

Tabernaevovine


[220327-17-3]

Absolute
Configuration $C_{40}H_{50}N_4O_2$ 618.861Alkaloid from *Tabernaemontana bovina*. Cryst. (Me₂CO). Mp 163-165°. $[\alpha]_D^{24} +133$ (c, 1 in CHCl₃).Lien, T.P. *et al.*, *Phytochemistry*, 1998, **49**, 1797-1799 (isol, pmr, cmr)**Tabernaegantidine**Bisindole alkaloid (Iboga-vobasine type). Struct. unknown. Alkaloid from *Tabernaemontana elegans* (Apocynaceae). Mol. formula not reported.Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 266 (rev)**Tabernaegantinine A**

[58917-03-6]

Bond to C-12'
Absolute
Configuration $C_{46}H_{58}N_4O_6$ 762.987Alkaloid from the root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (hexane). Mp 160°. $[\alpha]_D^{20} -53.7$ (CHCl₃). Possibly an artifact.*19,20Z-Didehydro-3-(2-Oxopropyl)conodurine*
[64192-91-2] $C_{46}H_{56}N_4O_6$ 760.972Alkaloid from the roots of *Tabernaemontana holstii* (Apocynaceae). Shows high cytotoxicity vs. mouse P-388 cells. Mp 203-204°.Bombardelli, E. *et al.*, *J.C.S. Perkin 1*, 1976, 1432 (*Tabernaegantinine A*)Kingston, D.G.I. *et al.*, *J. Pharm. Sci.*, 1977, **66**, 1135; 1978, **67**, 249 (3-(2-Oxopropyl)conodurine)**Tabernaegantinine B**

[58917-02-5]




T-4

T-1

 $C_{46}H_{58}N_4O_6$ 762.987Alkaloid from the root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (MeOH). Mp 215°. $[\alpha]_D^{20} +39.1$ (CHCl₃). Possibly an artifact.Bombardelli, E. *et al.*, *J.C.S. Perkin 1*, 1976, 1432 (uv, pmr, cmr, ms, struct)**Tabernaegantinine C**


[72542-42-8]



T-5

 $C_{44}H_{53}N_5O_5$ 731.933Alkaloid from the root bark of *Tabernaemontana elegans* (Apocynaceae). Cryst. (MeOH). Mp 177°.Danieli, B. *et al.*, *J.C.S. Perkin 1*, 1980, 601 (uv, ir, pmr, cmr, ms, cd, struct)**Tabernaemontabovine**

[243974-08-5]



R = H

T-6

 $C_{41}H_{50}N_4O_3$ 646.871Struct. revised in 2000. Alkaloid from leaves and stems of *Tabernaemontana bovina*. Oil. $[\alpha]_D^{25} -74.2$ (c, 0.50 in MeOH).Ripperger, H. *et al.*, *J. Prakt. Chem.*, 1999, **341**, 506-508; 2000, **342**, 725-727 (isol, pmr, cmr, ms)**Tabernaemontavine**


[243974-09-6]

As Tabernaemontabovine, T-6 with R = CH₂OH $C_{42}H_{52}N_4O_4$ 676.897Struct. revised in 2000. Alkaloid from leaves and stems of *Tabernaemontana bovina*. Oil. $[\alpha]_D^{25} -56.8$ (c, 0.50 in MeOH).Ripperger, H. *et al.*, *J. Prakt. Chem.*, 1999, **341**, 506-508; 2000, **342**, 725-727 (isol, pmr, cmr, ms)

T-7

Tabernamine

[59626-92-5]

Absolute
Configuration $C_{40}H_{48}N_4O_2$ 616.845Alkaloid from the stem bark of *Tabernaemontana johnstonii* and *Tabernaemontana dichotoma* (Apocynaceae). Shows strong cytotoxic activity vs. mouse P-388 carcinoma. Amorph. powder. $[\alpha]_D^{22} -51$ (c, 0.18 in MeOH). λ_{max} 236 (ε 33000); 287 (ε 10900); 295 (ε 10000) (MeOH) (Berdy). λ_{max} 235 (ε 34000); 287 (ε 10400); 295 (ε 10000) (EtOH) (Berdy).*N-De-Me: N⁴-Demethyltabernamine. 13'-Peribogamine*
[66302-66-7] $C_{39}H_{46}N_4O_2$ 602.818Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Shows strong activity vs. P-388 lymphocytic leukaemia.*19,20ξ-Dihydro: 19,20-Dihydrotabernamine*

[79543-26-3]

 $C_{40}H_{50}N_4O_2$ 618.861Alkaloid from the root bark of *Hazunta modesta* (Apocynaceae). Cryst. (MeOH). Mp 230°. $[\alpha]_D -27$ (c, 0.6 in CHCl₃).*3'-Hydroxy: 3'-Hydroxytabernamine*
[99257-60-0] $C_{40}H_{48}N_4O_3$ 632.844Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Active against gram-positive bacteria. Sol. MeOH, butanol, CHCl₃; poorly sol. H₂O, hexane. Epimeric mixt. at C-3'. λ_{max} 237 (ε 69200); 287 (ε 21900); 295 (ε 20400) (EtOH) (Berdy).*3'-Hydroxy, N-de-Me: 3'-Hydroxy-N⁴-demethyltabernamine*

[99257-59-7]

 $C_{39}H_{46}N_4O_3$ 618.817Alkaloid from the stem bark of *Tabernaemontana dichotoma* (Apocynaceae). Active against gram-positive bacteria. Sol. MeOH, CHCl₃, butanol; poorly sol. H₂O, hexane. Epimeric mixt. at C-3'. λ_{max} 236 (ε 74000); 256 (ε 20000); 296 (ε 18600) (EtOH) (Berdy).*19'R-Hydroxy: 19'R-Hydroxytabernamine*

[482662-87-3]

 $C_{40}H_{48}N_4O_3$ 632.844Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Light

T-8

yellow oil. $[\alpha]_D$ -139 (c, 0.13 in CHCl_3); λ_{max} 231 (log ϵ 4.59); 286 (log ϵ 4.07); 295 (log ϵ 4.02) (EtOH).

19'S-Hydroxy: 19'S-Hydroxytabernamine
[482662-86-2]

$\text{C}_{40}\text{H}_{48}\text{N}_4\text{O}_3$ 632.844
Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Light yellow oil. $[\alpha]_D$ -144 (c, 0.07 in CHCl_3); λ_{max} 230 (log ϵ 4.64); 286 (log ϵ 4.09); 295 (log ϵ 4.04) (EtOH).

19'-Oxo: 19'-Oxotabernamine

[482662-88-4]
 $\text{C}_{40}\text{H}_{46}\text{N}_4\text{O}_3$ 630.828
Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Oil. $[\alpha]_D$ -158 (c, 0.08 in CHCl_3); λ_{max} 231 (log ϵ 4.57); 284 (log ϵ 4.03); 296 (log ϵ 3.98) (EtOH).

10'-Methoxy, 19,20 α -dihydro: 16'-Decarbomethoxy-20-epidihydrovoacamine
[56725-98-5]

$\text{C}_{41}\text{H}_{52}\text{N}_4\text{O}_3$ 648.887
Alkaloid from the bark of *Ervatamia orientalis* (Apocynaceae). Powdery solid (petrol/ C_6H_6). $[\alpha]_D$ +18 (c, 0.75 in CHCl_3). Dec. at 180° without melting.

10'-Methoxy, 19,20 β -dihydro: 16'-Decarbomethoxy-19,20-dihydrovoacamine
[56691-97-5]

$\text{C}_{41}\text{H}_{52}\text{N}_4\text{O}_3$ 648.887
Alkaloid from the bark of *Ervatamia orientalis* (Apocynaceae). Powder (C_6H_6 /petrol). $[\alpha]_D$ +22 (c, 1.7 in CHCl_3). Dec. at 215° without melting.

Knox, J.R. *et al.*, *Aust. J. Chem.*, 1975, **28**, 1813 (Decarbomethoxydihydrovincaamines)

Kingston, D.G.I. *et al.*, *Tet. Lett.*, 1976, 649 (uv, ir, pmr, ms, struct)

Kingston, D.G.I. *et al.*, *J. Pharm. Sci.*, 1978, **67**, 249 (isol)

Urrea, M. *et al.*, *Bull. Soc. Chim. Fr., Part II*, 1981, 147 (isol, uv, ir, pmr, cmr, ms, struct, 19,20-Dihydrotabernamine)


Perera, P. *et al.*, *Phytochemistry*, 1985, **24**, 2097 (isol, uv, ir, pmr, ms, derivs, activity)

Kam, T.-S. *et al.*, *Heterocycles*, 2002, **57**, 2137-2143 (19'-Hydroxytabernamines, 19'-Oxotabernamine)

Tabernoxidine

T-9

[91423-26-6]



$\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_5$ 400.474


Minor alkaloid from the leaves of *Tabernaemontana heyneana* (Apocynaceae). Mp 291-292°.

Joshi, B.S. *et al.*, *Indian J. Chem., Sect. B*, 1984, **23**, 101 (uv, ir, pmr, ord, cryst struct)

Taberpsychine

T-10

3,17-Epoxyvobasan, 9CI. Anhydrovobasinediol
[19452-84-7]



Absolute Configuration

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}$ 308.422

Alkaloid from *Tabernaemontana psychotrifolia*, *Tabernaemontana brachyantha* and *Conopharyngia durissima* (Apocynaceae). Cryst. (MeOH). Mp 208° dec. $[\alpha]_D^{23}$ -284 (c, 0.87 in CHCl_3).

Methiodide:

Cryst. (Me_2CO). Mp 272-274° dec.

19Z-Isomer: (19Z)-Taberpsychine

[113973-32-3]

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}$ 308.422

Minor alkaloid from roots of *Gelsemium elegans* (Loganiaceae). Oil. $[\alpha]_D^{23}$ -180 (c, 0.4 in CHCl_3) (natural). $[\alpha]_D^{23}$ -151 (c, 0.3 in CHCl_3) (synth.).

19Z-Isomer, N-methoxy: N-Methoxyanhydrovobasinediol

[125180-42-9]

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_2$ 338.449

Alkaloid from *Gelsemium elegans* (Loganiaceae). Needles. Mp 75°. $[\alpha]_D$ -272.5 (c, 0.24 in MeOH).

Dugan, J.J. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 701 (uv, ir, pmr, ms, struct, synth)

Burnell, R.H. *et al.*, *Can. J. Chem.*, 1971, **49**, 307 (isol, uv, ir, pmr, ms, struct)

Patel, M.B. *et al.*, *Phytochemistry*, 1973, **12**, 451 (isol, uv, ir, pmr, ms)

Van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400 (isol, pmr, ms)

Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075 (19Z-Taberpsychine)

Takayama, H. *et al.*, *J.C.S. Perkin I*, 1989, 1075 (Taberpsychine, 19Z-Taberpsychine, synth, abs config)

Lin, L.-Z. *et al.*, *Phytochemistry*, 1989, **28**, 2827 (N-Methoxyanhydrovobasinediol)

Magnus, P. *et al.*, *J.A.C.S.*, 1990, **112**, 5220 (synth)


Kitajima, M. *et al.*, *J.C.S. Perkin I*, 1991, 1773 (synth)

Bailey, P.D. *et al.*, *J.C.S. Perkin I*, 1993, 441 (synth)

Tabersonine

T-11

Methyl 2,3,6,7-tetrahydroaspidospermidine-3-carboxylate, 9CI
[4429-63-4]



Absolute Configuration

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_2$ 336.433

Numbering systems vary. Alkaloid from *Amsonia tabernaemontana*, *Amsonia an-*

gustifolia, *Voacanga africana* and v. many other spp. in the Apocynaceae. Shows hypotensive activity (ca. 25% of Reserpine, R-52). Amorph.; cryst. (as hydrochloride). Mp 193-197° (hydrochloride). $[\alpha]_D$ -310 (c, 0.13 in MeOH) (hydrochloride). λ_{max} 230 (log ϵ 3.85); 299 (log ϵ 3.99); 328 (log ϵ 3.17) (EtOH).

N⁴-Oxide: Tabersonine N^b-oxide

[67249-34-7]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from the seeds of *Amsonia elliptica* (Apocynaceae). Amorph. λ_{max} 231; 298; 335 (EtOH).

14 α ,15 α -Epoxide: Lochnericine. 14 α ,15 α -Epoxytabersonine. Tabersonine α -epoxide

[72058-36-7]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from *Catharanthus roseus*, *Amsonia tabernaemontana*, *Amsonia angustifolia*, *Alstonia lanceolata*, *Catharanthus trichophyllus*, *Tabernaemontana* spp., *Voacanga* spp. and several other genera in Apocynaceae. Shows strong cytotoxicity vs. human carcinoma KB cells. Mp 188-191° dec. $[\alpha]_D^{25}$ +473 (c, 0.37 in EtOH). λ_{max} 299 (ϵ 18000); 328 (ϵ 3000) (MeOH) (Berdy). λ_{max} 225; 295; 325 (EtOH) (Berdy).

14 β ,15 β -Epoxide: Pachysiphine. Tabersonine β -epoxide. 14 β ,15 β -Epoxytabersonine

[2447-58-7]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from *Tabernaemontana pachysiphon* (Apocynaceae). Amorph.; cryst. (EtOH/ Me_2CO) (as hydrochloride). Mp 163° (hydrochloride). $[\alpha]_D$ -455 (c, 1.05 in MeOH) (hydrochloride).

11-Hydroxy: 11-Hydroxytabersonine. 16-Hydroxytabersonine

[22149-28-6]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from the leaves of *Tabernanthe pubescens*, *Melodinus balsansae* and *Craspidospermum verticillatum* (Apocynaceae). Mp 176°. $[\alpha]_D$ -332 (CHCl_3).

14,15-Dihydro: see Vincadifformine, V-111

11-Hydroxy, 14 α ,15 α -epoxide: 14,15-Epoxy-11-hydroxytabersonine

[140680-64-4]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Melodinus fusiformis* (Apocynaceae). Shows significant antitumour activity. Spermatocide.

11-Methoxy: Ervamycine. 11-Methoxytabersonine. Ervamycine

[27773-39-3]

[22149-29-7]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_3$ 366.459

Alkaloid from *Vinca herbacea*, *Vinca erecta*, *Melodinus polyadenus* and *Melodinus aeneus* (Apocynaceae). Amorph. Mp 186° dec. (as hydrochloride). $[\alpha]_D^{23}$ -310 (c, 0.32 in CHCl_3). $[\alpha]_D^{20}$ -264.4 (c, 1.4 in CHCl_3).

11-Methoxy, 14 α ,15 α -epoxide: Lochnericine. Hazuntine

[22255-04-5]
 $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from *Catharanthus roseus*,
Melodinus aeneus, *Vinca herbacea*,
Cabucala erythrocarpa and *Cabucala*
fasciculata (Apocynaceae). Shows
 antitumour activity. Cryst. (Me₂CO/
 MeOH). Mp 168-169°. $[\alpha]_D^{16}$ -424
 (CHCl₃). λ_{max} 247 (ε 15500); 326 (ε
 21350) (EtOH) (Berdy).

11-Methoxy, 14,15-epoxide, stereoisomer:
Ervincinine

[22223-16-1]
 $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from *Vinca erecta* (Apocyna-
 ceae). Cryst. (MeOH/Me₂CO). Mp
 247-248°. $[\alpha]_D^{22}$ -80.5 (c, 0.39 in CHCl₃).
 No config. information.

**10-Hydroxy, 11-methoxy: 10-Hydroxy-
 11-methoxytabersonine. 15-Hydroxy-
 16-methoxytabersonine**

[72713-28-1]
 $C_{22}H_{26}N_2O_4$ 382.458
 Alkaloid from *Hazunta* spp. (Apocy-
 naceae).

**10,11-Dimethoxy, 14β,15β-epoxide: Ha-
 zuntinine**

[22255-06-7]
 $C_{23}H_{28}N_2O_5$ 412.485
 Alkaloid from *Hazunta velutina*
 (Apocynaceae). Cryst. (MeOH). Mp
 132°. $[\alpha]_D$ -482 (c, 0.56 in CHCl₃).

**11,12-Dimethoxy, 10-hydroxy, 14β,15β-
 epoxide: Taberhanine**

$C_{23}H_{28}N_2O_6$ 428.484
 Alkaloid from the leaves of *Taber-
 naemontana divaricata*. Light yellow-
 ish oil. $[\alpha]_D$ -185 (c, 0.08 in CHCl₃).
 λ_{max} 202 (log ε 4.13); 238 (log ε 3.94);
 307 (log ε 3.96); 340 (log ε 3.93)
 (EtOH).

19-Hydroxy: see 19-Hydroxytabersonine,
 H-743

Dimer: Bistabersonine. Ditabersonine

[50645-77-7]
 $C_{42}H_{46}N_4O_4$ 670.85
 Alkaloid from the root bark of
Crioceras dipladeniiflorus (Apocyna-
 ceae). Noncryst. Oxidative dimer,
 point of dimerisation unknown.
 Possibly identical with Voafrine A,
 V-176.

[22260-37-3]

Janot, M.-M. *et al.*, *Bull. Soc. Chim. Fr.*, 1954,
 707-708 (*isol, ir, uv*)

Nair, C.P.N. *et al.*, *Tetrahedron*, 1959, **6**, 89
 (*Lochnericine, isol, uv, ir*)

Moza, B.K. *et al.*, *Chem. Ind. (London)*, 1962,
 1425 (*isol, uv, ir*)

Plat, M. *et al.*, *Tet. Lett.*, 1962, 271-276 (*ms,
 struct*)

Moza, B.K. *et al.*, *Tet. Lett.*, 1964, 2561
 (*Tabersonine, Lochnericine, pmr, ms, struct*)

Patel, M.B. *et al.*, *Bull. Soc. Chim. Fr.*, 1966,
 427 (*Pachysiphine, isol, uv, ir, pmr, ms*)

Pyuskyulev, B. *et al.*, *Coll. Czech. Chem.
 Comm.*, 1967, **32**, 1289-1294 (*Ervamycine*)

Kan-Fan, C. *et al.*, *Ann. Pharm. Fr.*, 1968, **26**,
 577-582 (*11-Hydroxytabersonine*)

Potier, P. *et al.*, *Ann. Pharm. Fr.*, 1968, **26**, 621
 (*Hazuntinine, isol, ir, pmr, ms, struct*)

Mehri, M. *et al.*, *Ann. Pharm. Fr.*, 1972, **30**,
 643-650 (*11-Hydroxytabersonine*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4990-
 4995 (*cmr*)

Ziegler, F.E. *et al.*, *J.A.C.S.*, 1973, **95**, 7458-
 7464 (*synth*)

Bruneton, J. *et al.*, *Phytochemistry*, 1973, **12**,
 1475-1480 (*Ditabersonine*)

Cavé, A. *et al.*, *Tet. Lett.*, 1973, 5081
 (*Hazuntinine, cmr, stereochem*)

Titeux, F. *et al.*, *Phytochemistry*, 1974, **13**, 1620
 (*isol*)

Douzoua, L. *et al.*, *Phytochemistry*, 1974, **13**,
 1994 (*isol*)

Takano, S. *et al.*, *J.A.C.S.*, 1976, **98**, 3022-3023
 (*synth*)

Baassou, S. *et al.*, *Phytochemistry*, 1978, **17**,
 1449-1450 (*Ervamycine*)

Bui, A.M. *et al.*, *Phytochemistry*, 1979, **18**,
 1329-1331 (*10-Hydroxy-11-
 methoxytabersonine*)

Kunesch, N. *et al.*, *Tet. Lett.*, 1980, **21**, 1727
 (*Lochnericine, cmr*)

Rabaron, A. *et al.*, *Ann. Pharm. Fr.*, 1981, **39**,
 369-373 (*16-Methoxytabersonine*)

Mulamba, T. *et al.*, *J. Nat. Prod.*, 1981, **44**,
 184-189 (*16-Hydroxytabersonine*)

Vercauteren, J. *et al.*, *Phytochemistry*, 1981, **20**,
 1411 (*Lochnericine, isol*)

Kunesch, N. *et al.*, *Bull. Soc. Chim. Fr., Part II*,
 1982, 285 (*Pachysiphine, struct, synth*)

He, X. *et al.*, *Huaxue Xuebao*, 1992, **50**, 96-
 101; *C.A.* **116**, 191092c (*14,15-Epoxy-11-
 hydroxytabersonine*)

Kobayashi, S. *et al.*, *Tet. Lett.*, 1999, **40**, 1519-
 1522 (*synth*)

Kozmin, S.A. *et al.*, *J.A.C.S.*, 2002, **124**, 4621-
 4641 (*synth*)


Éles, J. *et al.*, *J.O.C.*, 2002, **67**, 7255-7260
 (*Lochnericine, synth*)

Kam, T.-S. *et al.*, *Org. Biomol. Chem.*, 2003, **1**,
 1292-1297 (*Taberhanine*)

Tacamine

Pseudovincamine
 [56942-57-5]

T-12



$C_{21}H_{26}N_2O_3$ 354.448

A major alkaloid from the leaves and
 twigs of *Tabernaemontana eglandulosa*,
 also from *Tabernaemontana pandacaqui*
 (Apocynaceae). λ_{max} 227 ; 277 ; 282 ; 290
 (sh) (MeOH).

**16-Deoxy, 16,17-didehydro: 16,17-Anhy-
 drotacamine. Apotacamine**

[56942-59-7]

$C_{21}H_{24}N_2O_2$ 336.433

Minor alkaloid from *Tabernaemontana
 eglandulosa* (Apocynaceae). λ_{max} 228 ;
 273 ; 312 (MeOH).

19S-Hydroxy: 19-Hydroxytacamine

[90702-15-1]

$C_{21}H_{26}N_2O_4$ 370.447

Minor alkaloid from *Tabernaemontana
 eglandulosa* (Apocynaceae). λ_{max} 227 ;
 277 ; 280 ; 291 (MeOH).

16-Epimer: 16-Epitacamine

[56942-58-6]

$C_{21}H_{26}N_2O_3$ 354.448

A major alkaloid from *Tabernaemon-*

tana eglandulosa (Apocynaceae).

λ_{max} 225 ; 274 ; 280 ; 289 (sh)
 (MeOH).

van Beek, T.A. *et al.*, *Tetrahedron*, 1984, **40**,
 737-748 (*Tacamine, 19-Hydroxytacamine,
 16,17-Anhydrotacamine, uv, pmr, cmr, ms, cd,
 abs config*)

Taesotikul, T. *et al.*, *Planta Med.*, 1990, **56**,
 688 (*isol*)

Din Belle, D. *et al.*, *Tetrahedron*, 1996, **52**,
 11361-11378 (*synth, cmr*)

Lounasmaa, M. *et al.*, *Tetrahedron*, 1998,
54, 14845-14858 (*synth, 19-
 Hydroxytacamine*)


Tacamonine

Pseudovincamine I

[90761-95-8]

[83607-40-3 ((±)-form)]

T-13



$C_{19}H_{22}N_2O$ 294.396

A major alkaloid from the leaves and
 twigs of *Tabernaemontana eglandulosa*
 (Apocynaceae). Mp 180-181°. λ_{max} 242 ;
 265 ; 273 (sh) ; 294 ; 303 (MeOH).

**16α-Alcohol: 16S-Descarbomethoxytaca-
 mine**

[90702-14-0]

$C_{19}H_{24}N_2O$ 296.411

Minor alkaloid from *Tabernaemontana
 eglandulosa* (Apocynaceae). λ_{max} 228 ;
 277 ; 282 ; 290 (MeOH).

**16β-Alcohol: 16R-Descarbomethoxytaca-
 mine**

[90702-13-9]

$C_{19}H_{24}N_2O$ 296.411

Minor alkaloid from *Tabernaemontana
 eglandulosa* (Apocynaceae). λ_{max} 228 ;
 277 ; 282 ; 290 (MeOH).

17β-Hydroxy: 17β-Hydroxytacamonine

[90702-12-8]

$C_{19}H_{22}N_2O_2$ 310.395

Trace alkaloid from *Tabernaemontana
 eglandulosa* (Apocynaceae). Config.
 revised in 1995. Originally formulated
 as 17α-Hydroxytacamonine. λ_{max} 221 ;
 225 ; 242 ; 267 ; 290 ; 302 (sh)
 (MeOH).

Massiot, G. *et al.*, *Bull. Soc. Chim. Fr.*, 1982,
 185-190 (*synth*)

van Beek, T.A. *et al.*, *Tetrahedron*, 1984, **40**,
 737-748 (*isol, uv, pmr, ms, cd, struct,
 derivs*)

Wasserman, H.H. *et al.*, *Tet. Lett.*, 1989, **30**,
 873-876 (*synth*)

Ihara, M. *et al.*, *J.O.C.*, 1994, **59**, 5317-5323
 (*synth*)

Din Belle, D. *et al.*, *Tetrahedron*, 1996, **52**,
 11361-11378 (*17-Hydroxytacamonine, synth,
 config*)

Lounasmaa, M. *et al.*, *Tetrahedron*, 1998, **54**,
 157-164 (*synth*)


Suzuki, M. *et al.*, *Heterocycles*, 2000, **52**, 1083-
 1085 (*synth*)

Lavilla, R. *et al.*, *Eur. J. Org. Chem.*, 2001,
 3719-3729 (*synth*)

- Danieli, B. *et al.*, *Tet. Lett.*, 2001, **42**, 7237-7240 (*synth*)
 Deiters, A. *et al.*, *J.O.C.*, 2006, **71**, 6547-6561 (*synth*)
 Ho, T.-L. *et al.*, *Tetrahedron*, 2008, **64**, 10401-10405 (*synth*)

Tacraline

[98776-13-7]



Absolute Configuration

$C_{23}H_{27}N_3O_4$ 409.484
 Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). λ_{max} 205 ; 235 ; 293 (MeOH).


O-De-Ac: Desacetyltacraline

[98798-64-2]
 $C_{21}H_{25}N_3O_3$ 367.447
 Trace alkaloid from *Tabernaemontana chippii* (Apocynaceae). λ_{max} 207 ; 236 ; 295 (MeOH).

Van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400-423 (*isol, uv, pmr, ms, struct, de-Ac*)

Takatonine

[6882-14-0]
 [4668-06-8]



$C_{21}H_{24}NO_4^{\oplus}$ 354.425
 Quaternary alkaloid from the Japanese commercial crude drug “Takato-gusa”, the dried leaves and stems of *Thalictrum minus* (Ranunculaceae). Weak spasmolytic agent showing atropine-like activity. Yellow plates (MeOH) (as iodide). Mp 181-182° (iodide).

► LD₅₀ (mus, ipr) 167mg/kg.

O⁶-De-Me: 6-De-O-methyltakatonine

$C_{20}H_{22}NO_4^{\oplus}$ 340.398
 Quaternary alkaloid from the leaves of *Thalictrum delavayi*. Yellow oil. Counterion not specified.


[5083-11-4]

- Fujita, E. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 1082; *CA*, **54**, 4643a (*isol*)
 Kubota, S. *et al.*, *J.O.C.*, 1966, **31**, 516 (*uv, pmr, synth, struct*)
 Birch, A.J. *et al.*, *J.C.S. Perkin 1*, 1974, 2190 (*synth, pmr*)
 Baser, K.H.C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 704 (*isol, uv, ir, pmr, ms*)
 Wang, Y. *et al.*, *Acta Bot. Sin.*, 2003, **45**, 500-502 (*6-O-Demethyltakatonine*)

Talarotoxin

TB Toxin
 [155679-73-5]

T-17




$C_{28}H_{37}NO_3$ 435.605
 Prod. by *Talaromyces bacillisporus* IFO 8397 cultured on rice. Cytotoxic. Yellow needles (EtOH). Mp 137-138°.

Ishii, K. *et al.*, *Appl. Environ. Microbiol.*, 1995, **61**, 941-943 (*isol, pmr*)

Talassamine

[142861-01-6]

T-18



$C_{20}H_{27}NO_2$ 313.439
 Minor alkaloid from epigeal parts of *Aconitum talassicum* (Ranunculaceae). Cryst. (Me₂CO). Mp 208-210°.

7-Ac: Talassimine

[142861-10-7]
 $C_{22}H_{29}NO_3$ 355.476
 Minor alkaloid from epigeal parts of *Aconitum talassicum* (Ranunculaceae). Cryst. (Et₂O). Mp 242-245°.


15-Ac: Talassimidine

[142861-11-8]
 $C_{22}H_{29}NO_3$ 355.476
 Minor alkaloid from epigeal parts of *Aconitum talassicum* (Ranunculaceae). Cryst. (Me₂CO). Mp 263-265°.
 Nishanov, A.A. *et al.*, *Khim. Prir. Soedin.*, 1991, 93; *Chem. Nat. Compd. (Engl. Transl.)*, 82 (*isol, ir, pmr, ms, cryst struct*)

Talbotine, 9CI

[30809-15-5]

T-19



$C_{21}H_{24}N_2O_4$ 368.432
 Alkaloid from the leaves and stem bark of *Pleiocarpa talbotii* (Apocynaceae). Cryst. (MeOH). Mp 212°. $[\alpha]_D^{23}$ -200 (c, 0.528 in CHCl₃). $[\alpha]_D^{23}$ -289 (c, 0.995 in Py).

N,O-Di-Ac:

Cryst. (Et₂O/pentane). Mp 190-193°. $[\alpha]_D^{25}$ -35 (c, 0.975 in CHCl₃).

Me ether:

Cryst. (Me₂CO/hexane). Mp 155-156°. $[\alpha]_D^{20}$ -240 (c, 0.915 in MeOH).

3,4-Didehydro: 3,4-Didehydrotalbotine

[51856-92-9]
 $C_{21}H_{22}N_2O_4$ 366.416
 Alkaloid from the leaves of *Pleiocarpa talbotii* (Apocynaceae). Needles (MeOH). Mp 196°. $[\alpha]_D^{22}$ -182 (c, 0.612 in MeOH).

3,4,5,6-Tetrahydro: 3,4,5,6-Tetrahydrotalbotine

[30809-25-7]
 $C_{21}H_{20}N_2O_4$ 364.4
 Alkaloid from leaves of *Pleiocarpa talbotii* (Apocynaceae). Needles (Me₂CO). Mp 189°. $[\alpha]_D^{22}$ -211 (c, 0.294 in MeOH).

3,4,5,6-Tetrahydro, Me ether:

Cryst. (Me₂CO/hexane). Mp 212-214°. $[\alpha]_D^{25}$ -230 (c, 0.482 in MeOH).

Pinar, M. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 15 (*isol, uv, ord, ir, pmr, ms, struct*)


Narango, J. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 752 (*isol*)

Pinar, M. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 2719 (*isol, uv, ord, ir, struct, Didehydrotalbotine, Tetrahydrotalbotine*)

Talcarpine

T-20

20,21-Dihydro-21-methyl-18-noralstophyllan-19-al, 9CI
 [38990-08-8]



Absolute Configuration

$C_{21}H_{26}N_2O_2$ 338.449
 Alkaloid from *Alstonia macrophylla* and *Pleiocarpa talbotii*. Plates (Et₂O/Me₂CO/hexane). Mp 167-169° dec. $[\alpha]_D$ -26 (c, 0.12 in CHCl₃). λ_{max} 209 (log ϵ 3.86); 226 (log ϵ 4.01); 277 (log ϵ 2.65); 285 (log ϵ 2.91); 294 (log ϵ 2.65) (EtOH).

21-Alcohol: Macrocarpine A

[685508-24-1]
 $C_{21}H_{28}N_2O_2$ 340.464
 Alkaloid from the bark of *Alstonia macrophylla*. Light yellow oil. $[\alpha]_D$ +117 (c, 0.11 in CHCl₃). λ_{max} 230 (log ϵ 4.15); 286 (log ϵ 3.46) (EtOH).

19,20-Didehydro: 19,20-Didehydrotalcarpine. Alstonerinal

[158725-12-3]
 $C_{21}H_{24}N_2O_2$ 336.433
 Alkaloid from *Alstonia macrophylla*. $[\alpha]_D$ -32 (c, 0.03 in CHCl₃). λ_{max} 228 (log ϵ 4.3); 267 (log ϵ 4); 291 (log ϵ 3.8) (EtOH).

19,20-Didehydro, N¹-de-Me: N¹-De-methylalstonerinal

Alkaloid from the bark of *Alstonia macrophylla*.


19,20-Didehydro, N⁴-de-Me: N⁴-De-methylalstonerinal

[701304-80-5]
 $C_{20}H_{22}N_2O_2$ 322.406
 Alkaloid from the leaves of *Alstonia angustifolia* var. *latifolia*.
20 α -Hydroxy-, 21-alcohol: Alstohentine
 [685508-22-9]
 $C_{21}H_{28}N_2O_3$ 356.464
 Alkaloid from the leaves of *Alstonia macrophylla*. Light yellowish oil. $[\alpha]_D^{25}$ -58 (c, 0.22 in $CHCl_3$); λ_{max} 228 (log ϵ 3.91); 289 (log ϵ 3.26) (EtOH).
10-Methoxy-, 19,20-didehydro-: 19,20-Dehydro-10-methoxytalcarpine
 [120374-20-1]
 $C_{22}H_{26}N_2O_3$ 366.459
 Alkaloid from the leaves of *Alstonia angustifolia*. Mp 300°. $[\alpha]_D^{25}$ -140 (c, 0.5 in EtOH).
20-Epimer: 20-Epitalcarpine. N^b-Methyl-N^b,21-secotalpinine
 [38990-35-1]
 Alkaloid from *Alstonia macrophylla*. $[\alpha]_D^{25}$ +19 (c, 0.45 in $CHCl_3$). λ_{max} 205 (log ϵ 3.95); 228 (log ϵ 4.21); 280 (log ϵ 3); 285 (log ϵ 3.42); 300 (log ϵ 3.12) (EtOH).
20-Epimer, 21-alcohol: Macrocarpine B
 [685508-25-2]
 $C_{21}H_{28}N_2O_2$ 340.464
 Alkaloid from the bark of *Alstonia macrophylla*. Light yellow oil. $[\alpha]_D^{25}$ -51 (c, 0.34 in $CHCl_3$); λ_{max} 230 (log ϵ 4.34); 288 (log ϵ 3.64) (EtOH).
20-Epimer, 21-alcohol, 21-Ac: Macrocarpine C
 $C_{23}H_{30}N_2O_3$ 382.502
 Alkaloid from the bark of *Alstonia macrophylla*. $[\alpha]_D^{25}$ -35 (c, 1.5 in $CHCl_3$). λ_{max} 230 (log ϵ 4.3); 287 (log ϵ 3.62) (EtOH).
 Naranjo, J. et al., *Helv. Chim. Acta*, 1972, **55**, 752-771 (isol, uv, ir, pmr, ms, struct)
 Garnick, R.L. et al., *J.A.C.S.*, 1978, **100**, 4213-4219 (synth, ir, pmr, ms, epimer)
 Ghedira, K. et al., *Phytochemistry*, 1988, **27**, 3955-3962 (19,20-Dehydro-10-methoxytalcarpine)
 Takayama, H. et al., *Tetrahedron*, 1991, **47**, 1383-1392 (synth, uv, ir, pmr, cmr, ms, cd)
 Wong, W.-H. et al., *Phytochemistry*, 1996, **41**, 313-315 (isol, pmr, cmr)
 Yu, P. et al., *J.O.C.*, 1998, **63**, 9160-9161; 2000, **65**, 3173-3191 (synth)
 Kam, T.-S. et al., *Phytochemistry*, 1999, **51**, 839-844 (Alstonerinal)
 Kam, T.-S. et al., *J. Nat. Prod.*, 2004, **67**, 547-552 (Alstohentine)
 Kam, T.-S. et al., *Phytochemistry*, 2004, **65**, 603-608 (N^b-Demethylalstonerinal)
 Kam, T.-S. et al., *Tetrahedron*, 2004, **60**, 3957-3966 (isol, pmr, cmr, 20-Epitalcarpine, Macrocarpines)
 Liao, X. et al., *J.O.C.*, 2006, **71**, 8884-8890 (synth)

Talpinine

T-21

17,19-Epoxy-19,20-dihydro-1-methylsarpagan-21-ol, 9CI
 [38990-06-6]



Absolute Configuration

$C_{20}H_{24}N_2O_2$ 324.422
 Alkaloid from the root bark and stem bark of *Pleiocarpa talbotii* (Apocynaceae). Plates (Et₂O/pentane). Mp 153-154°. $[\alpha]_D^{25}$ -30 (c, 0.302 in $CHCl_3$). pK_a 6.51 (MCS).

Picrate:


Cryst. (MeOH). Mp 245° dec.

10-Methoxy-, N-de-Me: 21-Hydroxycyclochlornerine
 [96688-58-3]
 $C_{20}H_{24}N_2O_3$ 340.421
 Alkaloid from *Catharanthus roseus* cell suspension cultures, and from the leaves, stem, and root bark of *Rauwolfia biauriculata* (Apocynaceae). Mp 172°. Naranjo, J. et al., *Helv. Chim. Acta*, 1972, **55**, 752-771 (isol, uv, ir, pmr, ms, cd, struct)
 Garnick, R.L. et al., *J.A.C.S.*, 1978, **100**, 4213-4219 (synth, ir, ms)
 Kohl, W. et al., *Planta Med.*, 1984, **50**, 242-244 (21-Hydroxycyclochlornerine, isol, uv, pmr, cmr, ms, cd, cryst struct)
 Abaul, J. et al., *J. Nat. Prod.*, 1986, **49**, 829-832 (21-Hydroxycyclochlornerine)
 Yu, P. et al., *J.O.C.*, 1998, **63**, 9160-9161; 2000, **65**, 3173-3191 (synth)

Tambjamine A

T-22

[85850-00-6]



$C_{10}H_{11}N_3O$ 189.216
 Isol. from the nembrothid nudibranchs *Tambja abdere*, *Tambja eliora*, *Atapozoa* sp., *Nembrotha crista*, *Nembrotha kubaryana* and *Roboastra tigris*, also isol. from the dietary source *Sessibugula translucens*. DNA binding agent. Oil. Sol. MeOH, $CHCl_3$; poorly sol. H₂O. λ_{max} 255 (ε 4600); 397 (ε 20000) (MeOH) (Derep).

1''-N-Et: Tambjamine E

[126584-10-9]
 $C_{12}H_{15}N_3O$ 217.27
 Alkaloid from the marine ascidian *Atapozoa* sp. and from *Nembrotha crista* and *Nembrotha kubaryana*. DNA binding agent, feeding deterrent. Yellow cryst. ($CDCl_3$). Mp 68-70°. λ_{max} 204 (ε 26600); 252 (ε 6900); 366 (ε 17700) (MeOH/NaOH) (Derep). λ_{max} 205 (ε 5000); 255 (ε 5700); 280 (sh); 405 (ε 30500) (MeOH) (Derep).

1''-N-(2-Methylpropyl): Tambjamine C

[85850-02-8]
 $C_{14}H_{19}N_3O$ 245.324
 Isol. from *Tambja abdere*, *Tambja eliora*, *Roboastra tigris*, *Sessibugula* and *Atapozoa* spp. DNA binding agent. Oil. Sol. MeOH, $CHCl_3$; poorly sol. H₂O. λ_{max} 204 (ε 26600); 252 (ε 6900); 366 (ε 17700) (MeOH/NaOH) (Derep). λ_{max} 205 (ε 5000); 255 (ε 5700); 280 (sh); 405 (ε 30500) (MeOH) (Derep). λ_{max} 204 (ε 5300); 251 (ε 6600); 258 (ε 5700); 405 (ε 23000) (MeOH) (Berdy).

1''-N-(2-Phenylethyl): Tambjamine F

[126584-09-6]
 $C_{18}H_{19}N_3O$ 293.368
 Alkaloid from *Atapozoa* sp. and from *Nembrotha crista* and *Nembrotha kubaryana*. DNA binding agent, feeding deterrent. Brown oil. λ_{max} 205 (ε 10200); 259 (ε 7000); 407 (ε 32100) (MeOH) (Berdy). λ_{max} 203 (ε 32600); 251 (ε 7000); 365 (ε 19200) (MeOH/NaOH) (Berdy).

1''-N-Dodecyl: Antibiotic BE 18591. BE 18591

[147138-01-0]
 $C_{22}H_{35}N_3O$ 357.538
 Prod. by *Streptomyces* sp. BA18591. Antitumour agent. Yellowish-green amorph. solid. Mp 50-53°. λ_{max} 257 (28400); 325 (100000); 406 (8500) (MeOH).

1''-N-(3Z-Dodeceny): [882419-92-3]

$C_{22}H_{33}N_3O$ 355.522
 Prod. by the marine bacterium *Pseudoalteromonas tunicata*. Yellow solid.

3'-Bromo, 1''-N-(2-methylpropyl): Tambjamine D

[85850-03-9]
 $C_{14}H_{18}BrN_3O$ 324.22
 Isol. from *Tambja abdere*, *Tambja eliora*, *Sessibugula* sp. and *Roboastra tigris* spp. DNA binding agent. Oil. Sol. MeOH, $CHCl_3$; poorly sol. H₂O. λ_{max} 257 (ε 6200); 401 (ε 23000) (MeOH) (Derep).

5'-Bromo: Tambjamine B

[85850-01-7]
 $C_{10}H_{10}BrN_3O$ 268.112
 Isol. from *Tambja abdere*, *Tambja eliora*, *Roboastra tigris*, *Sessibugula* and *Roboastra* spp. DNA binding agent. Oil. Sol. MeOH, $CHCl_3$; poorly sol. H₂O. λ_{max} 255 (ε 4600); 397 (ε 20000) (MeOH) (Derep).

5'-Bromo, 1''-N-Et: Tambjamine G

[157536-53-3]
 $C_{12}H_{14}BrN_3O$ 296.166
 Alkaloid from the bryozoan *Bugula dentata*.

5'-Bromo, 1''-N-propyl: Tambjamine H

[157536-54-4]
 $C_{13}H_{16}BrN_3O$ 310.193
 Alkaloid from *Bugula dentata*.

5'-Bromo, 1''-N-(2-methylpropyl): Tambjamine I

[157536-55-5]
 $C_{14}H_{18}BrN_3O$ 324.22
 Alkaloid from *Bugula dentata*.

5'-Bromo, 1''-N-(2-methylbutyl): Tambjamine J

[157536-56-6]
 $C_{15}H_{20}BrN_3O$ 338.246
 Alkaloid from *Bugula dentata*.


[153231-89-1]

Carté, B. et al., *J.O.C.*, 1983, **48**, 2314-2318 (*Tambjamines A-D, struct*)
 Carté, B. et al., *J. Chem. Ecol.*, 1986, **12**, 795-804 (*Tambjamines A-D, defence pheromones*)
 Paul, V.J. et al., *Marine Ecol.: Progr. Ser.*, 1990, **59**, 109 (occur)
 Lindquist, N. et al., *Experientia*, 1991, **47**, 504-506 (*Tambjamines C,E,F*)
 Kojiri, K. et al., *J. Antibiot.*, 1993, **46**, 1799-1803; 1894-1896 (*BE 18591*)

- Blackman, A.J. *et al.*, *Aust. J. Chem.*, 1994, **47**, 1625-1629 (*Tambjamines G-J*)
 Franks, A. *et al.*, *Molecules*, 2005, **10**, 1286-1291 (*Pseudoalteromonas* alkaloid)
 Pinkerton, D.M. *et al.*, *Org. Lett.*, 2007, **9**, 5127-5130 (*synth*)

Tamynine

[159509-33-8]




$C_{14}H_{13}NO_4$ 259.261
 Alkaloid from leaves of *Murraya paniculata* (Rutaceae). Mp 113-114°. $[\alpha]_D^{26} +103$ (c, 0.23 in $CHCl_3$).

Khan, M.A. *et al.*, *Heterocycles*, 1994, **38**, 2005 (*isol, uv, ir, pmr, cmr, ms, struct*)

Tanakamine

2-(1*H*-Indol-3-yl)-3-methyl-1,3-butane-diol, 9*CI*. 3-(2-Hydroxy-1-hydroxy-methyl-2-methylpropyl)-1*H*-indole [106449-17-6]




$C_{13}H_{17}NO_2$ 219.283

(ξ)-form

Alkaloid from the stems of *Limonia acidissima* (wood apple) (Rutaceae). Amorph. $[\alpha]_D^{25} +7.6$ (c, 0.69 in MeOH). Zarga, M.H.A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 901 (*isol, uv, pmr, cmr, ms, struct*)

Tanakine

1-(1*H*-Indol-3-yl)-3-methyl-2,3-butane-diol, 9*CI* [106449-15-4]



$C_{13}H_{17}NO_2$ 219.283
 Alkaloid from the stems of *Limonia acidissima* (wood apple) (Rutaceae). Amorph. $[\alpha]_D^{25} +38$ (c, 0.51 in MeOH).

N-β-D-Glucopyranosyl: **Bruceolline F** [159903-55-6]


$C_{19}H_{27}NO_7$ 381.425
 Alkaloid from root wood of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Pale yellow powder (Me₂CO/MeOH). Mp 209-211°. $[\alpha]_D^{20} -55.8$ (c, 0.9 in Py). Stereochem. not detd. Not correlated

with Tanakine.

- Zarga, M.H.A. *et al.*, *J. Nat. Prod.*, 1986, **49**, 901 (*isol, uv, pmr, ms, struct*)
 Ouyang, Y. *et al.*, *Phytochemistry*, 1994, **37**, 575 (*Bruceolline F*)

Tangirine

[152606-86-5]




$C_{49}H_{62}N_2O_7$ 791.038
 Alkaloid from *Aconitum tanguticum* (Ranunculaceae). Mp 266-268° dec. $[\alpha]_D^{19} +84.2$ (c, 0.36 in $CHCl_3$).

Joshi, B.S. *et al.*, *Tet. Lett.*, 1993, **34**, 7525 (*isol, ir, pmr, cmr, ms, struct*)

Tangutorine

[224426-78-2]




$C_{20}H_{24}N_2O$ 308.422
 Alkaloid from *Nitraria tangutorum*. Cryst. (CH₂Cl₂/MeOH). Mp 276-278°. Racemic. λ_{max} 225 ; 281 ; 292 (sh) (no solvent reported).

- Duan, J.-A. *et al.*, *Tet. Lett.*, 1999, **40**, 2593-2596 (*isol, uv, ir, pmr, cmr, cryst struct*)
 Putkonen, T. *et al.*, *Tet. Lett.*, 2001, **42**, 6593-6594 (*synth*)
 Luo, S. *et al.*, *Org. Lett.*, 2003, **5**, 4709-4712 (*synth*)
 Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 122-126 (*synth, pmr, cmr*)

Tannagine

[123750-34-5]



$C_{21}H_{27}NO_5$ 373.448

Relative configuration

T-26

This struct. was originally assigned to Iso-stephodeline, I-325. Alkaloid from aerial parts of *Stephania zippeliana* (Menispermaceae). $[\alpha]_D +40$ (c, 0.80 in $CHCl_3$).

14-Epimer: Stephodeline

[56596-12-4]

 $C_{21}H_{27}NO_5$ 373.448

Alkaloid from *Stephania delavayi* and *Stephania zippeliana* (Menispermaceae). Amorph. Mp 198-200° (as hydrochloride). $[\alpha]_D +20$ (c, 0.6 in $CHCl_3$).

14-Epimer, O²-de-Me: Zippelianine. 2-O-Demethylstephodeline

[123688-34-6]

 $C_{20}H_{25}NO_5$ 359.421


Alkaloid from aerial parts of *Stephania zippeliana* (Menispermaceae). $[\alpha]_D +25$ (c, 0.70 in $CHCl_3$).

Perel'son, M.E. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 188; *Chem. Nat. Compd. (Engl. Transl.)*, 197 (*Stephodeline*)

Charles, B. *et al.*, *Can. J. Chem.*, 1989, **67**, 1257 (*Zippelianine*)

Tantazole A**T-29**

[129941-13-5]



$R^1 = CH_3, R^2 = H$

$C_{24}H_{32}N_6O_2S_4$ 564.82

Stereochem. revised in 1993. Isol. from the blue-green alga *Scytonema mirabile*. Cytotoxic. $[\alpha]_D -31.9$ ($CHCl_3$). λ_{max} 222 (ε 25000); 255 (ε 12800) (EtOH) (Derep).

4''-Epimer: Tantazole I

[129895-75-6]

 $C_{24}H_{32}N_6O_2S_4$ 564.82

From *Scytonema mirabile*. Cytotoxic. No phys. props. reported. λ_{max} 222 (ε 25000); 255 (ε 12800) (EtOH) (Derep).

Carmeli, S. *et al.*, *J.A.C.S.*, 1990, **112**, 8195 (*isol, struct*)

Carmeli, S. *et al.*, *Tet. Lett.*, 1993, **34**, 6681 (*config*)

Tantazole B**T-30**

[129895-76-7]

As Tantazole A, T-29 with

 $R^1 = R^2 = CH_3$ $C_{25}H_{34}N_6O_2S_4$ 578.847

Stereochem. revised in 1993. Isol. from *Scytonema mirabile*. Cytotoxic. $[\alpha]_D -94$ ($CHCl_3$). λ_{max} 222 (ε 25000); 255 (ε 12800) (EtOH) (Derep).

Carmeli, S. *et al.*, *J.A.C.S.*, 1990, **112**, 8195 (*isol, struct*)

Fukuyama, T. *et al.*, *J.A.C.S.*, 1993, **115**, 8449 (*synth*)

Carmeli, S. *et al.*, *Tet. Lett.*, 1993, **34**, 6681 (*config*)

Parsons, R.L. *et al.*, *Synlett*, 1996, 1168 (*synth*)

Tantazole F**T-31**

[129895-77-8]

As Tantazole A, T-29 with

 $R^1 = H, R^2 = CH_3$ $C_{24}H_{32}N_6O_2S_4$ 564.82


Isol. from *Scytonema mirabile*. Cytotoxic.

$[\alpha]_D^{25}$ -63.7 (CHCl₃). λ_{\max} 222 (ϵ 25000); 255 (ϵ 12800) (EtOH) (Derep).

Carmeli, S. *et al.*, *J.A.C.S.*, 1990, **112**, 8195 (isol, struct)

Tardioxopiperazine A

T-32



$R^1 = -CH_2CH=C(CH_3)_2$, $R^2 = H$

C₂₄H₃₁N₃O₂ 393.528

Prod. by *Microascus tardifaciens*. Moderate immunosuppressant. Amorph. pale yellow solid. $[\alpha]_D^{25}$ -30 (c, 0.2 in CHCl₃). λ_{\max} 229 (log ϵ 4.46); 281 (sh) (log ϵ 3.85); 287 (log ϵ 3.86); 296 (sh) (log ϵ 3.79) (MeOH).

Fujimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1426-1432

Tardioxopiperazine B

T-33

As Tardioxopiperazine A, T-32 with $R^1 = H$, $R^2 = -CH_2CH=C(CH_3)_2$

C₂₄H₃₁N₃O₂ 393.528


Prod. by *Microascus tardifaciens*. Amorph. pale yellow solid. $[\alpha]_D^{25}$ -16 (c, 0.2 in CHCl₃). λ_{\max} 219 (sh) (log ϵ 3.7); 226 (log ϵ 4.4); 273 (sh) (log ϵ 3.79); 280 (log ϵ 3.79) (MeOH).

Fujimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 1426-1432

Tarennine⁺

T-34

2,3,4,9-Tetrahydro-1-[2-(1-pyrrolidinylethyl)-1H-pyrido[3,4-b]indole]-9CI. 1,2,3,4-Tetrahydro-1-(2-pyrrolidylethyl)- β -carboline. Dihydroelaeocarpidine [20069-08-3]



C₁₇H₂₃N₃ 269.389

(\pm)-form [24298-76-8]

Alkaloid from *Tarenna bipindensis* (Rubiaceae). Mp 124-126°. Boissier *et al.* propose that the name Dihydroelaeocarpidine be used in preference to Tarennine. However, since hydrogenation of Elaeocarpidine involves ring-opening the trivial name Tarennine is preferred here.

Hydrochloride (1:2): Mp 294-295°.

Picrate: Mp 200-202°.

Harley-Mason, J. *et al.*, *Chem. Comm.*, 1969, 281 (synth)

Gribble, G.W. *et al.*, *J.O.C.*, 1970, **35**, 1944 (synth)


Boissier, J.R. *et al.*, *Experientia*, 1971, **27**, 677 (struct, ir, ms)

Gribble, G.W. *et al.*, *J.O.C.*, 1988, **53**, 3164 (synth)

Tasmanine

T-35

[80640-70-6]



Absolute Configuration

C₂₀H₂₆N₂O 310.438

Alkaloid from *Aristotelia peduncularis* (Elaeocarpaceae). Cryst. (MeOH). Mp 250° (subl. from 220°). $[\alpha]_D^{20}$ -132 (c, 0.453 in CHCl₃). λ_{\max} 213 (log ϵ 4.42); 250 (sh) (log ϵ 3.77); 255 (log ϵ 3.8); 262 (sh) (log ϵ 3.74); 286 (log ϵ 3.22); 291 (sh) (log ϵ 3.17) (EtOH).

Kyburz, R. *et al.*, *Helv. Chim. Acta*, 1981, **64**, 2555-2561 (isol, uv, ir, pmr, cmr, ms, cd, struct)


Hai, M.A. *et al.*, *Tetrahedron*, 1984, **40**, 4359-4361 (abs config)

Güller, R. *et al.*, *Helv. Chim. Acta*, 1993, **76**, 1847-1862 (synth)

Taspine

T-36

1-[2-(Dimethylamino)ethyl]-3,8-dimethoxy[1]benzopyrano[5,4,3-cde][1]-benzopyran-5,10-dione, 9CI. Thaspine [602-07-3]



C₂₀H₁₉NO₆ 369.373

Alkaloid from *Leontice eversmannii*, *Caulophyllum robustum*, *Magnolia liliiflora* and from the bark of *Croton lechleri* (Leonticaceae, Euphorbiaceae). Antiinflammatory agent. Cryst. (MeOH). Mp 225° dec. Mp 370° dec. $[\alpha]_D^{24}$ +7.6 (c, 0.64 in Py). Log P 1.64 (calc).

Hydrochloride:

Needles. Mp 252-253° dec.

Platonova, T.F. *et al.*, *Zh. Obshch. Khim.*, 1956, **26**, 2651-2656; *CA*, **51**, 5102b (*Leontice eversmannii* constiti)

Safronich, L.N. *et al.*, *CA*, 1961, **55**, 18892 (*Caulophyllum robustum* constiti)

Shamma, M. *et al.*, *Chem. Comm.*, 1971, 1065-1066 (synth, uv, pmr)

Perdue, G.P. *et al.*, *J. Pharm. Sci.*, 1979, **68**, 124-126 (*Croton lechleri* constiti)


Talapatra, B. *et al.*, *Phytochemistry*, 1982, **21**, 747-750 (*Magnolia liliiflora* constiti)

Kelly, T.R. *et al.*, *J.O.C.*, 1998, **63**, 8045-8048 (synth)

Tatiopterin

T-37

[126026-65-1]



Tatiopterin O R = OH
1 R = -Glu-OH

Tatiopterin O

C₃₃H₄₄N₇O₁₉P 873.72

Isol. from *Methanogenium tationis*. Isol. as Na salt.

2',5'-Dihydroxy: **Thermopterin**

[135745-46-9]

C₃₃H₄₄N₇O₂₁P 905.719

Isol. from *Methanoculleus thermophilicum*.

Tatiopterin 1

C₃₈H₅₁N₈O₂₂P 1002.835

Isol. from *Methanogenium tationis*. Isol. as Na salt.

Raemakers-Franken, P.C. *et al.*, *BioFactors*, 1991, **3**, 127 (isol)


Raemakers-Franken, P.C. *et al.*, *Eur. J. Biochem.*, 1991, **200**, 783-787 (isol, uv, pmr, ms)

Tauret

T-38

Retinylidenetaurine

[133867-05-7]



C₂₂H₃₃NO₃S 391.574


Endogenous constiti. of the retina and pigmented epithelium of the frog *Rana ridibunda*.

Petrosian, A.M. *et al.*, *Adv. Exp. Med. Biol.*, 1996, **403**, 333-342

Tauroacidin A

T-39

[200566-34-3]



C₁₃H₁₆Br₂N₆O₅S 528.181

Tautomerism possible in the iminoimidazole ring. Closely related to Dispacamide A, D-863. Alkaloid from the sponge *Hymeniacidon* sp. Inhibits tyro-

sine kinase. Amorph. solid. $[\alpha]_D^{28}$ -4.3 (c, 0.1 in MeOH). Isol. as a partial racemate. λ_{\max} 273 (ϵ 9700); 312 (ϵ 3800) (MeOH).

9-Deoxy: **Taurodispacamide A**

[323177-96-4]

C₁₃H₁₆Br₂N₆O₄S 512.181

Alkaloid from the sponge *Agelas oroides*. Antihistaminic agent. Pale yellow solid. λ_{\max} 230; 270 (no solvent reported).

2-Debromo: **Tauroacidin B**

[200566-35-4]

C₁₃H₁₇BrN₆O₅S 449.285

Alkaloid from *Hymeniacidon* sp. Inhibits tyrosine kinase. Amorph. solid. Isol. as a racemate.

2-Debromo, 9-deoxy: **9-Deoxytauroacidin B**

2-Debromotaurodispacamide A

C₁₃H₁₇BrN₆O₄S 433.285

Alkaloid from *Axinella verrucosa*.

Kobayashi, J. *et al.*, *Tetrahedron*, 1997, **53**,


16679-16682 (isol, uv, ir, pmr, cmr, ms)

Fattorusso, E. *et al.*, *Tet. Lett.*, 2000, **41**, 9917-9922 (*Taurodispacamide A*)

Aiello, A. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 17-24 (*Deoxytauroacidin B*)

Taveuniamide E

T-40



C₁₉H₂₂Cl₃NO₃ 418.746

Isol. from a mixed assemblage of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25}$ +2.1 (c, 0.3 in CHCl₃). λ_{\max} 242 (ϵ 7200) (EtOH).

12,12,13,13,14,15-Hexahydro: **Taveuniamide A**

C₁₉H₂₈Cl₃NO₃ 424.793

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25}$ +11.7 (c, 0.4 in CHCl₃). Stereochem. not determined. λ_{\max} 236 (ϵ 6700) (EtOH).

1-Chloro, 1,2-dihydro: **Taveuniamide D**

C₁₉H₂₃Cl₄NO₃ 455.206

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25}$ +5.3 (c, 0.09 in CHCl₃). Stereochem. not determined. λ_{\max} 244 (ϵ 16500) (EtOH).

1-Chloro, 1,2,3E,4-tetrahydro: **Taveuniamide C**

C₁₉H₂₅Cl₄NO₃ 457.222

Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D$ +24 (c, 0.03 in CHCl₃). Stereochem. not determined. λ_{\max} 244 (ϵ 15400) (EtOH).

1-Chloro, 12,12,13,13,14,15-hexahydro: **Taveuniamide B**

C₁₉H₂₇Cl₄NO₃ 459.238


Isol. from a mixt. of *Lyngbya majuscula* and *Schizothrix* sp. $[\alpha]_D^{25}$ +10.4 (c, 0.1 in CHCl₃). Stereochem. not determined. λ_{\max} 244 (ϵ 18300) (EtOH).

Williamson, R.T. *et al.*, *Tetrahedron*, 2004, **60**, 7025-7033 (isol, pmr, cmr)

Tawicyclamide A

[143007-28-7]

T-41



R = CH₂Ph

C₃₉H₅₀N₈O₅S₃ 807.073

Cyclic peptide antibiotic. Isol. from the ascidian *Lissoclinum patella*. Weakly cytotoxic. Clear solid. $[\alpha]_D^{25}$ -15 (c, 0.43 in CHCl₃). λ_{\max} 249 (ϵ 16530) (MeOH) (Berdy).

McDonald, L.A. *et al.*, *J.O.C.*, 1992, **57**, 4616-4624 (isol, pmr, cmr, cryst struct)

Tawicyclamide B

[143007-29-8]

T-42

As Tawicyclamide A, T-41 with

R = -CH₂CH(CH₃)₂


C₃₆H₅₂N₈O₅S₃ 773.055

Cyclic peptide antibiotic. Isol. from the ascidian *Lissoclinum patella*. Weakly cytotoxic. Clear solid. $[\alpha]_D^{25}$ +2.1 (c, 0.35 in CHCl₃). λ_{\max} 249 (ϵ 15029) (MeOH) (Berdy).

McDonald, L.A. *et al.*, *J.O.C.*, 1992, **57**, 4616-4624 (isol, pmr, cmr, cryst struct)

4(20),11-Taxadiene-1,2,5,7,9,10,13-heptol

T-43



(1 β ,2 α ,5 α ,7 β ,9 α ,10 α ,13 α)-form

C₂₀H₃₂O₇ 384.469

(1 β ,2 α ,5 α ,7 β ,9 α ,10 α ,13 α)-form

5-O-(3-Dimethylamino-2-hydroxy-3-phenylpropanoyl), 2,7,9,10,13-penta-Ac: **2-Acetoxy-2-deacetyl-1-hydroxytaustrospicatin**


C₄₁H₅₅NO₁₄ 785.884

Alkaloid from *Taxus baccata* (Taxaceae). Cryst. Mp 120°. $[\alpha]_D^{25}$ +42 (c, 1.1 in CHCl₃).

Barboni, L. *et al.*, *Annalen*, 1995, 345 (2-Acetoxy-2'-deacetyl-1-hydroxytaustrospicatin)

4(20),12-Taxadiene-2,5,7,9,10,11,13-heptol

T-44



C₂₀H₃₂O₇ 384.469

(2 α ,5 α ,7 β ,9 α ,10 β ,11 β)-form

5-(3-Dimethylamino-3-phenylpropanoyl)(R-), 2,7,9,10,13-penta-Ac: **Taxuspine P**

[175739-43-2]

C₄₁H₅₅NO₁₃ 769.884

Constit. of *Taxus cuspidata*. Microtubule depolymerisation inhibitor. Amorph. solid. $[\alpha]_D^{19}$ +32.7 (c, 1.3 in MeOH). λ_{\max} 207 (ϵ 14500); 261 (ϵ 960) (MeOH) (Berdy).


Kobayashi, J. *et al.*, *Tetrahedron*, 1996, **52**,

5391 (*Taxuspine P*)

Kobayashi, J. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 393-398 (*Taxuspines*, activity)

4(20),11-Taxadiene-1,2,5,9,10,13-hexol

T-45



C₂₀H₃₂O₆ 368.469

(1 β ,2 α ,5 α ,9 α ,10 β ,13 α)-form [138613-22-6]

Constit. of *Taxus chinensis*. Cryst. Mp 120-121°. $[\alpha]_D$ -5.6 (CHCl₃/Py).

5-O-(3-Dimethylamino-3-phenylpropanoyl)(S-), 10,13-di-Ac: **13-Deoxo-13-acetyloxytaxine B**

[151124-22-0]

C₃₅H₄₉NO₉ 627.773

Alkaloid from *Taxus baccata* (Taxaceae). Powder. Mp 119°. $[\alpha]_D^{25}$ +42 (c, 0.26 in CH₂Cl₂).

5-O-(3-Dimethylamino-3-phenylpropanoyl)(S-), 2,9,10,13-tetra-Ac: **13-Dihydro-2,9,13-triacetyltaxine B**

[155512-17-7]

C₃₉H₅₃NO₁₁ 711.848

Constit. of *Taxus baccata*. $[\alpha]_D^{22}$ +60 (c, 0.52 in CHCl₃).

13-Ketone, 5-O-(3-dimethylamino-3-phenylpropanoyl)(S-), 10-Ac: **Taxine B. Taxine I**

[1361-51-9]

C₃₃H₄₅NO₈ 583.72

Alkaloid from yew *Taxus baccata* (Taxaceae). Amorph. solid. Mp 109°. $[\alpha]_D$ +116.9 (c, 0.33 in CHCl₃). The older lit. on substances known as Taxine B and Taxine I is confused. Struct. revised in 1991.


▶ WX1275000

13-Ketone, 5-O-(3-dimethylamino-3-phenylpropanoyl) (S-), 2,9,10-tri-Ac: **Diacetyltaxine B**
[101981-13-9]
C₃₇H₄₉NO₁₀ 667.795
Constit. of *Taxus baccata*. Mp 115°
Mp 156-158°. [α]_D²⁵ +107 (c, 0.88 in MeOH).

Ettouati, L. et al., *J. Nat. Prod.*, 1991, **54**, 1455-1458 (*Taxine B*, ir, uv, pmr, struct)
Appendino, G. et al., *J. Nat. Prod.*, 1993, **56**, 514-520 (*Diacetyltaxine B*)
Appendino, G. et al., *Phytochemistry*, 1993, **33**, 1521-1523 (*13-Deoxo-13-acetyloxytaxine B*)
Hassanzadeh, M.K. et al., *Iran. J. Sci. Technol.*, 1999, **23**, 313-319; *CA*, **132**, 262625x (*13-Dihydro-2,9,13-triacetyltaxine B*)

4(20),11-Taxadiene-2,5,7,9,10,13-hexol

T-46



(2α,5α,7β,9α,10α,13α)-form

C₂₀H₃₂O₆ 368.469**(2α,5α,7β,9α,10α,13α)-form**

5-(3-Dimethylamino-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: **2-Hydroxy-2'-deacetoxyaustrospicatine**
[119777-83-2]
C₃₉H₅₃NO₁₁ 711.848
Alkaloid from leaves of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +50 (c, 2.75 in CHCl₃).

5-(3-Dimethylamino-2-hydroxy-3-phenylpropanoyl), 2,7,9,10,13-penta-Ac: **2-Acetoxy-2'-deacetylaustrospicatine**
[119777-82-1]
C₄₁H₅₅NO₁₃ 769.884
Alkaloid from leaves of *Austrotaxus spicata* (Taxaceae). Cryst. (CHCl₃/MeOH). Mp 113-115°. [α]_D +19 (c, 2 in CHCl₃).

5-(2-Acetoxy-3-dimethylamino-3-phenylpropanoyl), 2,7,9,10,13-penta-Ac: **2α-Acetoxyaustrospicatine**
[119777-81-0]
C₄₃H₅₇NO₁₄ 811.922
Alkaloid from leaves of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D -41 (c, 0.3 in CHCl₃).

(2α,5α,7β,9α,10β,13α)-form

5-(3S-Dimethylamino-2R-hydroxy-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: [824951-59-9]
C₃₉H₅₃NO₁₂ 727.847
Constit. of *Taxus canadensis*. Gum. [α]_D²⁵ +76 (c, 0.21 in CHCl₃).


4β,20-Epoxyde, 5-O-(3ξ-dimethylamino-3-phenylpropanoyl), 2,7,9,10,13-penta-Ac: **7-Acetoxy-9-acetylspicataxine**
[156576-68-0]
C₄₁H₅₅NO₁₃ 769.884

Alkaloid from roots of *Taxus x media* cv. *Hicksii* (Taxaceae). Powder. Mp 212-213°. [α]_D²⁵ +56 (c, 1.1 in CHCl₃).

Ettouati, L. et al., *Bull. Soc. Chim. Fr.*, 1988, 749-755; 1989, 687-694 (*Autotaxus spicata* constits)
Appendino, G. et al., *J. Nat. Prod.*, 1994, **57**, 607 (*7-Acetoxy-9-acetylspicataxine*)
Shi, Q.-W. et al., *Phytochemistry*, 2004, **65**, 3097-3106 (*Taxus canadensis* consti)

4(20),11-Taxadiene-2,5,9,10,13-pentol

T-47

C₂₀H₃₂O₅ 352.47**(2α,5α,9α,10β,13α)-form**

5-O-(3-Methylamino-3-phenylpropanoyl) (R-), 10,13-di-Ac: **13-Deoxy-13-acetyloxy-1-deoxynortaxine B**
[151124-24-2]
C₃₄H₄₇NO₈ 597.747
Alkaloid from *Taxus baccata* and *Taxus canadensis* (Taxaceae). Powder or gum. Mp 165°. [α]_D²⁵ +76 (c, 0.62 in CH₂Cl₂). [α]_D²² +47 (c, 0.21 in CHCl₃).

5-(3-Methylamino-3-phenylpropanoyl), 2,10,13-tri-Ac: [525601-68-7]
C₃₆H₄₉NO₉ 639.784
Constit. of *Taxus canadensis*. Gum. [α]_D²² +37 (c, 0.1 in CHCl₃).

5-(3-Methylamino-3-phenylpropanoyl), 2,9,10,13-tetra-Ac: [256942-45-7]
C₃₈H₅₁NO₁₀ 681.822
Constit. of *Taxus chinensis* var. *mairei*. Gum. [α]_D²⁴ +47 (c, 0.002 in CHCl₃).

5-(3-Dimethylamino-3-phenylpropanoyl) (R-), 2,9,13-tri-Ac: [640278-65-5]
C₃₇H₅₁NO₉ 653.811
Constit. of *Taxus canadensis*. Gum. [α]_D²² +37 (c, 0.1 in CHCl₃).

5-(3-Dimethylamino-3-phenylpropanoyl) (R-), 2,10,13-tri-Ac: [640278-66-6]
C₃₇H₅₁NO₉ 653.811
Constit. of *Taxus canadensis*. Amorph. powder or gum. [α]_D²² +49 (c, 0.1 in CHCl₃). [α]_D²² +53 (c, 0.21 in CHCl₃).

5-(3-Dimethylamino-3-phenylpropanoyl) (R-), 9,10,13-tri-Ac: **Taxuspine Z**
[194782-04-2]
C₃₇H₅₁NO₉ 653.811
Constit. of *Taxus cuspidata*. Amorph. solid. [α]_D²⁸ +31.2 (c, 0.08 in CHCl₃). λ_{max} 210 (ε 15400); 265 (ε 2330) (MeOH).

5-(3-Dimethylamino-3-phenylpropanoyl) (R-), 2,9,10,13-tetra-Ac: **2-Acetyltaxuspine Z**
[214151-14-1]
[256942-48-0]
C₃₉H₅₃NO₁₀ 695.848
Constit. of *Taxus chinensis* var. *mairei*. Amorph. powder. λ_{max} 216 ; 276

(MeOH).

5-O-(3R-Dimethylamino-3-phenylpropanoyl), 10,13-di-Ac: **13-Deoxo-13-acetyloxy-1-deoxytaxine B**
[151124-23-1]
C₃₅H₄₉NO₈ 611.774

Alkaloid from *Taxus baccata* (Taxaceae). Powder. Mp 192°. [α]_D²⁵ +25 (c, 0.79 in CH₂Cl₂).

4β,20-Epoxyde, 5-O-(3ξ-dimethylamino-3-phenylpropanoyl), 2,9,13-tri-Ac: **9-Acetyl-10-deacetylspicataxine**
[126585-91-9]
C₃₇H₅₁NO₁₀ 669.811

From the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +43 (c, 1.1 in CHCl₃).

4β,20-Epoxyde, 5-O-(3ξ-dimethylamino-3-phenylpropanoyl), 2,10,13-tri-Ac: **Spicataxine**
[126585-86-2]
C₃₇H₅₁NO₁₀ 669.811

Alkaloid from the bark of *Austrotaxus spicata* (Taxaceae).

4β,20-Epoxyde, 9-O-(3-pyridinecarbonyl), 5-O-(3ξ-methylamino-3-phenylpropanoyl), 2,10,13-tri-Ac: **N-Demethylnicaustrine**
[126617-16-1]
C₄₂H₅₂N₂O₁₁ 760.88
From the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +93 (c, 0.52 in CHCl₃).

4β,20-Epoxyde, 9-O-(3-pyridinecarbonyl), 5-O-(3ξ-dimethylamino-3-phenylpropanoyl), 2,10,13-tri-Ac: **Nicaustrine**
[127211-02-3]
C₄₃H₅₄N₂O₁₁ 774.906
Alkaloid from the bark of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +67 (c, 1.8 in CHCl₃).

Ettouati, L. et al., *Bull. Soc. Chim. Fr.*, 1989, 687-755 (*Austrotaxus spicata* constits)

Appendino, G. et al., *Phytochemistry*, 1993, **33**, 1521-1523 (*5-dimethylaminophenylpropanoyl 10,13-di-Ac, 5-methylaminophenylpropanoyl 10,13-di-Ac*)

Shigemori, H. et al., *Chem. Pharm. Bull.*, 1997, **45**, 1205-1208 (*Taxuspine Z*)

Chen, Y.-J. et al., *J. Nat. Prod.*, 1999, **62**, 149-151 (*Decinamoyltaxine E, 2-Acetyltaxuspine Z*)

Shi, Q.-W. et al., *Phytochemistry*, 1999, **52**, 1571-1575 (*Taxus chinensis* var. *mairei* consti)


Shi, Q.-W. et al., *Tohoku J. Agric. Res.*, 1999, **50**, 33-46; *CA*, **132**, 134755c (*Taxus chinensis* var. *mairei* consti)

Shi, Q.-W. et al., *J. Nat. Prod.*, 2003, **66**, 470-476; 1480-1485 (*Taxus canadensis* constits)

Shi, Q.-W. et al., *Phytochemistry*, 2004, **65**, 3097-3106 (*Taxus canadensis* constits)

4(20),11-Taxadiene-5,7,9,10,13-pentol

T-48



(5α,7β,9α,10β,13α)-form

C₂₀H₃₂O₅ 352.47**(5 α ,7 β ,9 α ,10 β ,13 α)-form** [27854-03-1]Cryst. Mp 194-195°. [α]_D +65.

5-(3-Methylamino-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: [256942-46-8]

C₃₈H₅₁NO₁₀ 681.822Constit. of *Taxus chinensis* var. *mairei*.Gum. [α]_D²⁵ +10.3 (c, 0.005 in CHCl₃).5-(3-Dimethylamino-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: **2'-Deacetoxyaustrospicatin**

[119777-80-9]

C₃₉H₅₃NO₁₀ 695.848From the leaves of *Austrotaxus spicata*, *Taxus cuspidata* and *Taxus wallichiana*. Amorph. [α]_D +71 (c, 0.85 in CH₂Cl₂). [α]_D +117 (c, 1 in CH₂Cl₂).5-(3-Dimethylamino-2-hydroxy-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: **2'-Deacetylaustrospicatin**

[119777-78-5]

C₃₉H₅₃NO₁₁ 711.848From the leaves of *Austrotaxus spicata* (Taxaceae). Cryst. (CHCl₃/MeOH).Mp 298-300°. [α]_D +56 (c, 0.7 in CH₂Cl₂).5-(2-Acetoxy-3-dimethylamino-3-phenylpropanoyl), 10-Ac: **2',3,9-Trideacetylaustrospicatin**

[119789-85-4]

C₃₅H₄₉NO₉ 627.773From the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +41 (c, 1.05 in CHCl₃).5-(2-Acetoxy-3-dimethylamino-3-phenylpropanoyl), 10,13-di-Ac: **7,9-Dideacetylaustrospicatin**

[119789-84-3]

C₃₇H₅₁NO₁₀ 669.811From the leaves of *Austrotaxus spicata* (Taxaceae). Amorph.5-(2-Acetoxy-3-dimethylamino-3-phenylpropanoyl), 9,10,13-tri-Ac: **7-Deacetylaustrospicatin**

[119777-79-6]

C₃₉H₅₃NO₁₁ 711.848From the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +41 (c, 0.5 in CHCl₃).5-(2-Acetoxy-3-dimethylamino-3-phenylpropanoyl), 7,9,10,13-tetra-Ac: **Austrospicatin**

[119777-76-3]

C₄₁H₅₅NO₁₂ 753.885Alkaloid from the leaves of *Austrotaxus spicata* (Taxaceae). Amorph. [α]_D +52 (c, 0.7 in CH₂Cl₂).

13-Ketone: 5,7,9,10-Tetrahydroxy-4(20),11-taxadien-13-one

C₂₀H₃₀O₅ 350.454

Aglycone of 2-Deacetoxytaxinine B.

13-Ketone, 5-(3-dimethylamino-2-hydroxy-3-phenylpropanoyl), 7,9,10-tri-Ac: **Comptonine**†


[126585-69-1]

C₃₇H₄₉NO₁₀ 667.795Taxane alkaloid from the bark of *Austrotaxus spicata* (Taxaceae).Amorph. [α]_D +85 (c, 0.82 in CHCl₃).Ettouati, L. et al., *Bull. Soc. Chim. Fr.*, 1988, 749-755; 1989, 687-694 (*Austrotaxus spicata*

constit)

Konda, Y. et al., *Chem. Pharm. Bull.*, 1994, **42**, 2621-2624 (2'-Deacetoxyaustrospicatin, *cryst struct*)Chattopadhyay, S.K. et al., *Indian J. Chem., Sect. B*, 1996, **35**, 508-509 (2'-Deacetoxyaustrospicatin)Shi, Q.-W. et al., *Phytochemistry*, 1999, **52**, 1571-1575 (*Taxus chinensis* var. *mairei* *constit*)Shi, Q.-W. et al., *Tohoku J. Agric. Res.*, 1999, **50**, 33-46; *CA*, **132**, 134755c (*Taxus chinensis* var. *mairei* *constit*)**4(20),11-Taxadiene-5,9,10,13-tetrol**

T-49

C₂₀H₃₂O₄ 336.47**(5 α ,9 α ,10 β ,13 α)-form** [15877-46-0]Constit. of the heartwood of *Taxus baccata*. Cryst. Mp 195-198°. [α]_D +134.

5-(3-Methylamino-3-phenylpropanoyl), 9,10,13-tri-Ac: [256942-47-9]


C₃₆H₄₉NO₈ 623.785Constit. of *Taxus chinensis* var. *mairei*.5-(3-Dimethylamino-3-phenylpropanoyl), 9,10,13-tri-Ac: **2',7-Bisdeacetoxyaustrospicatin**. 2',7-Dideacetoxyaustrospicatin

[159384-73-3]

C₃₇H₅₁NO₈ 637.812Constit. of *Taxus wallichiana* and *Taxus mairei*. Needles (Me₂CO). Mp 199-200°. [α]_D¹⁵ +112.9 (c, 0.95 in CHCl₃) (+14.5).Shi, Q.W. et al., *Nat. Prod. Lett.*, 1999, **13**, 179-186 (2',7-Bisdeacetoxyaustrospicatin)Shi, Q.-W. et al., *Tohoku J. Agric. Res.*, 1999, **50**, 33-46; *CA*, **132**, 134755c (*Taxus chinensis* var. *mairei* *constit*)**Taxilamine**

T-50

(6,7-Dimethoxy-1-isoquinoliny)l (2-hydroxy-3,4-dimethoxyphenyl)methanone, 9CI. 1-(2-Hydroxy-3,4-dimethoxybenzoyl)-6,7-dimethoxyisoquinoline [81525-57-7]

C₂₀H₁₉NO₆ 369.373Alkaloid from the root bark of *Berberis aristata* (Berberidaceae). Amorph.3,4-Dihydro: **Dihydrotaxilamine**


[105798-92-3]

C₂₀H₂₁NO₆ 371.389Alkaloid from roots of *Berberis actinacantha* (Berberidaceae). Amorph.Blaskó, G. et al., *Heterocycles*, 1982, **19**, 257 (*isol, uv, pmr, ms, struct*)Rahimizadeh, M. et al., *Planta Med.*, 1986, 339 (*Dihydrotaxilamine*)**Taxodine**

T-51

1,6-Didehydro-3,16-dimethoxy-C-homoerythrinan-15-ol, 9CI. Alkaloid B†. 3-Epischellhammericine B. Alkaloid V†. 2,7-Dihydrohomoerysoline

[24204-37-3]

C₁₉H₂₅NO₃ 315.411A major alkaloid from *Schelhammera undulata*. Also isol. from *Schelhammera pedunculata*, *Cephalotaxus harringtonia* var. *harringtonia*, *Cephalotaxus drupacea*, *Cephalotaxus oliveri*, *Athrotaxis cupressoides*, *Athrotaxis selaginoides* and *Athrotaxis laxifolia* (Liliaceae, Cephalotaxaceae, Taxodiaceae). Needles (Me₂CO). Mp 152-153°. [α]_D +111 (c, 0.09 in CHCl₃). λ_{\max} 235 (sh) (ϵ 8000); 283 (ϵ 3700); 289 (sh) (ϵ 3300) (no solvent reported).*Me ether*: 2,7-Dihydrohomoerysotrine. O-Methyltaxodine. O-Methyl-3-epischellhammericine B

[51095-85-3]

C₂₀H₂₇NO₃ 329.438Minor alkaloid from leaves, stems and roots of *Cephalotaxus harringtonia* var. *harringtonia* and the leaves of *Phelline* sp. aff. *Phelline lucida*, *Dysoxylum lenticellare* and *Cephalotaxus wilsoniana* (Cephalotaxaceae, Phellinaceae, Meliaceae). Noncryst. [α]_D +122 (c, 0.50 in CHCl₃).17-O-De-Me: **Cephalozomine M**. 17-O-Demethyltaxodine

[462075-60-1]

C₁₈H₂₃NO₃ 301.385Alkaloid from the leaves of *Cephalotaxus harringtonia* var. *nana*. Amorph. solid. [α]_D +64 (c, 0.8 in MeOH). λ_{\max} 217 (ϵ 4500); 233 (ϵ 3600); 286 (ϵ 1800) (MeOH).3-Epimer: **Schelhammericine B**

[30633-33-1]

C₁₉H₂₅NO₃ 315.411Minor alkaloid from *Cephalotaxus harringtonia* var. *harringtonia* (Cephalotaxaceae). [α]_D +76 (c, 0.40 in CHCl₃). Not obt. pure. The struct. of this alkaloid has not been proven and it could be the 16-hydroxy-15-methoxy isomer. It has, however, been registered by CAS with the 16-methoxy struct.3-Epimer, *Me ether*: 3-Epi-2,7-dihydrohomoerysotrine. O-Methylschelhammericine B

[38750-53-7]

C₂₀H₂₇NO₃ 329.438


Minor alkaloid from leaves, stems and roots of *Cephalotaxus harringtonia* var. *harringtonia* (Cephalotaxaceae). [α]_D²⁵ +118 (c, 0.58 in CHCl₃).

- Johns, S.R. *et al.*, *Aust. J. Chem.*, 1969, **22**, 2219-2231 (*uv, ir, pmr, ms, struct*)
 Sioumis, A.A. *et al.*, *Aust. J. Chem.*, 1971, **24**, 2737 (*isol*)
 Powell, R.G. *et al.*, *Phytochemistry*, 1972, **11**, 1467-1472 (*isol, uv, ir, pmr, ms, struct, Cephalotaxus harringtonia constits*)
 Aladesanmi, A.J. *et al.*, *J. Nat. Prod.*, 1983, **46**, 127-131 (*2,7-Dihydrohomoerysotrine*)
 Langlois, N. *et al.*, *Heterocycles*, 1984, **22**, 2453-2457 (*2,7-Dihydrohomoerysotrine*)
 Panichanun, S. *et al.*, *Tetrahedron*, 1984, **40**, 2677-2684; 2685-2689 (*isol, uv, ir, pmr, ms, synth*)
 Morita, H. *et al.*, *Tetrahedron*, 2002, **58**, 5489-5495 (*Cephalozomine M*)

Taxol

T-52

Paclitaxel, BAN, INN, USAN. *Taxol A. Anzntax. Yewtaxan. BMS 181339-01. NSC 125973 [33069-62-4]*



C₄₇H₅₁NO₁₄ 853.918

A coml. semisynthesis introduced in 1993 uses 10-Deacetylbaaccatin III in E-141. Isol. from the stem bark of *Taxus brevifolia*, *Taxus cuspidata* and *Taxus yunnanensis* (Taxaceae). Also reported in traces from the fungus *Taxomyces andreanae* and from *Taxus baccata*, *Cephalotaxus mannii*, *Taxus wallichiana*, *Taxomyces andreanae*, *Taxus sumatrana*, *Pestalotiopsis wallachiana*, *Pestalotiopsis microspora*, *Periconia* sp., *Pestalotia heterocornis* and *Trichothecium roseum*. Antileukaemic and antineoplastic agent, esp. against melanoma and ovarian tumours. Use limited by low solubility. Biochemical tool extensively used to study cellular shape and function. Also shown to be active against oomycete fungi. Needles (MeOH aq.). Mp 213-216°. [α]_D²⁰ -49 (MeOH). Log P 2.47 (uncertain value) (calc). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).

- ▶ LD₅₀ (mus, ipr) 128 mg/kg. LD₅₀ (mus, ivn) 12 mg/kg. WX1272100

Poly(L-glutamic acid) conjugate: Paclitaxel poliglumex, INN, USAN. CT 2103. PG-TXL. XYOTAX [263351-82-2] Biodegradable polymer. Delivers higher levels of active chemotherapeutic to the tumour. Phase III trials (2002) for the treatment of non-small cell lung cancer.

Polyethylene glycol conjugate: Peg-paclitaxel
 Prodrug of taxol.

7-O-β-D-Xylopyranoside: 7-Xylosyltaxol [90332-66-4]

C₅₂H₅₉NO₁₈ 986.034

Constit. of *Taxus baccata*. Cryst. (2-propanol). Mp 236-238°. [α]_D²⁵ -23 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).

N-Debenzoyl, N-Ac: N-Acetyl-N-debenzoyltaxol [173101-48-9]

C₄₂H₄₉NO₁₄ 791.847

Constit. of *Taxus canadensis*. Gum.

N-Debenzoyl, N-butanoyl: Taxol D. Taxcultine [153415-46-4]

C₄₄H₅₃NO₁₄ 819.901

Isol. from roots of *Taxus x media* cv Hicksii and cell cultures of *Taxus baccata* and *Taxus canadensis* (Taxaceae). Powder. Mp 155° Mp 206-208°. [α]_D²⁵ -7.54 (c, 0.106 in MeOH). [α]_D²⁵ -16 (c, 0.45 in CHCl₃). Identity of Taxol D with Taxcultine not confirmed. λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).

N-Debenzoyl, N-butanoyl, 7-O-β-D-xylopyranoside: 7-Xylosyltaxol D [334981-00-9]

C₄₉H₆₁NO₁₈ 952.017

Constit. of bark of *Taxus yunnanensis*. Powder. [α]_D²⁸ -25 (c, 0.25 in MeOH). λ_{\max} 202 (log € 4.3); 229 (log € 4.2) (MeOH).

N-Debenzoyl, N-(2-methylbutanoyl): N-Debenzoyl-N-(α-methylbutyryl)paclitaxel [159001-25-9]

C₄₅H₅₅NO₁₄ 833.928

Constit. of *Taxus x media* cv Hicksii. Powder. [α]_D²⁵ -48 (c, 0.1 in MeOH).

N-Debenzoyl, N-tigloyl: Cephalomannine. Taxol B [71610-00-9]

C₄₅H₅₃NO₁₄ 831.912

Alkaloid from *Taxus wallichiana*, the bark of *Taxus yunnanensis* and *Cephalotaxus mannii*. Antileukaemic agent. [α]_D²³ -41 (c, 0.39 in MeOH). Abs. config. of side-chain not certain. λ_{\max} 225 (€ 25000); 273 (€ 1700) (MeOH) (Derep).

N-Debenzoyl, N-tigloyl, 7-O-β-D-xylopyranoside: 7-Xylosylcephalomannine. 7-Xylosyltaxol B [90352-19-5]

C₅₀H₆₁NO₁₈ 964.028

Constit. of *Taxus baccata*. [α]_D²⁵ -26 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).

N-Debenzoyl, N-hexanoyl: Taxol C. Taxuyunnanine A [153415-45-3]

C₄₆H₅₇NO₁₄ 847.955

From roots of *Taxus x media* cv Hicksii, *Taxus yunnanensis* and cell cultures of *Taxus baccata*. Also from *Taxus brevifolia* and *Taxus media*. Shows potent selective cytotoxicity vs. NCI human cell line. Tubulin assembly promotor. Cryst. (MeOAc/hexane) or yellowish amorph. solid (MeOH). Mp 204-205° (150°). [α]_D²⁰ -64.7 (c, 1.2 in

CHCl₃). [α]_D²⁵ -107 (c, 0.054 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 227 (€ 15490) (MeOH) (Berdy).

N-Debenzoyl, N-hexanoyl, 7-O-β-D-xylopyranoside: 7-Xylosyltaxol C [90332-67-5]

C₅₁H₆₅NO₁₈ 980.07

Constit. of *Taxus baccata*. Tubulin assembly promotor. Cryst. (EtOH aq.). Mp 229-231°. [α]_D²⁵ -4 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).

N-Debenzoyl, N-cinnamoyl: N-Debenzoyl-N-cinnamoylpaclitaxel [219783-77-4]

C₄₉H₅₃NO₁₄ 879.956

Constit. of *Taxus x media* cv Hicksii. Powder. Mp 180° dec. [α]_D²⁵ -16.6 (c, 0.9 in MeOH).

N-Me: Taxuspinanane I [212071-25-5]

C₄₈H₅₃NO₁₄ 867.945

Constit. of *Taxus cuspidata* var. *nana*. Amorph. powder. [α]_D²⁵ -71 (c, 0.04 in CHCl₃). λ_{\max} 204 (log € 4.41); 218 (log € 4.32); 272 (log € 3.27); 282 (log € 3.22) (MeOH).

N-Me, N-debenzoyl, N-hexanoyl: N-Methyltaxol C [153083-53-5]

C₄₇H₅₉NO₁₄ 861.981

From roots of *Taxus x media* cv Hicksii and *Taxus baccata*. Tubulin assembly promotor. Cryst. (MeOAc/hexane). Mp 225-228° dec. [α]_D²⁰ -52.7 (c, 1.0 in CHCl₃). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 223 (€ 16980) (MeOH) (Berdy).

N-Me, N-debenzoyl, N-hexanoyl, 7-O-β-D-xylopyranoside: 7-Xylosyl-N-methyltaxol C [550373-59-6]

C₅₂H₆₇NO₁₈ 994.097

Constit. of *Taxus yunnanensis*. Cryst. Mp 220°. λ_{\max} 202 (log € 4.41); 214 (log € 4.33) (MeOH).

2',7-Di-Ac: 2',7-Diacetyltaxol [81924-74-5]

C₅₁H₅₅NO₁₆ 937.993

Constit. of *Taxus brevifolia*. Tubulin assembly promotor. Needles. Poorly sol. hexane. Mp 227-228°.

7-[(2,3-Dihydroxypropoxy) carbonyl]: Paclitaxel ceribate, INN. Protaxel. BP 179 [186040-50-6]

C₅₁H₅₇NO₁₈ 972.007

H₂O sol. prodrug.

4-De-Ac, 4-methoxycarbonyl: BMS 188797 [172481-83-3]

C₄₇H₅₁NO₁₅ 869.918

Antineoplastic agent. Amorph. solid.

10-De-Ac: 10-Deacetyltaxol [78432-77-6]

C₄₅H₄₉NO₁₃ 811.881

Constit. of *Taxus wallichiana*, *Taxus mairei* and *Cephalotaxus mannii*. Tubulin assembly promotor. Amorph. powder. λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 211 (€ 15500); 231 (€ 15500); 277 (€ 3160)


- (MeOH) (Berdy).
- 10-De-Ac, 7-O- β -D-xylopyranoside: 10-Deacetyl-7-xylosyltaxol**
[90332-63-1]
C₅₀H₅₇NO₁₇ 943.997
Constit. of *Taxus baccata*, *Taxus yunnanensis* and *Taxus canadensis*. Tubulin assembly promotor. Cryst. (EtOH aq.). Mp 246-248°. [α]_D -2 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 229 (€ 20700); 274 (€ 1400); 281 (€ 1100) (MeOH) (Berdy).
- 10-De-Ac, N-debenzoyl, N-propanoyl: N-Debenzoyl-N-propanoyl-10-deacetylpaclitaxel**
[173101-59-2]
C₄₁H₄₉NO₁₃ 763.837
Constit. of *Taxus baccata*. Cryst. Mp 245° dec.
- 10-De-Ac, N-debenzoyl, N-butanoyl: N-Debenzoyl-N-butanoyl-10-deacetylpaclitaxel**
[173101-47-8]
C₄₂H₅₁NO₁₃ 777.864
Constit. of *Taxus baccata*. Cryst. Mp 244° dec.
- 10-De-Ac, N-debenzoyl, N-butanoyl, 7-O- β -D-xylopyranoside: 7-Xylosyl-10-deacetyltaxol D**
[172486-23-6]
C₄₇H₅₉NO₁₇ 909.98
Constit. of *Taxus baccata*. Amorph. powder. [α]_D²⁰ -14.3 (c, 0.27 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 227 (€ 6166); 263 (€ 933); 267 (€ 933); 273 (€ 955); 280 (€ 813) (MeOH) (Berdy).
- 10-De-Ac, N-debenzoyl, N-tigloyl: 10-Deacetylcephalomannine**
[76429-85-1]
C₄₃H₅₁NO₁₃ 789.875
Isol. from *Taxus wallichiana* and *Cephalotaxus mannii*. Amorph. Sol. H₂O. λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 10-De-Ac, N-debenzoyl, N-tigloyl, 7-O- β -D-xylopyranoside: 10-Deacetyl-7-xylosylcephalomannine. 10-Deacetyl-7-xylosyltaxol B**
[90332-64-2]
C₄₈H₅₉NO₁₇ 921.991
Constit. of *Taxus baccata*. Cryst. (EtOH aq.). Mp 250-252°. [α]_D +4 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 10-De-Ac, N-debenzoyl, N-hexanoyl: 10-Deacetyltaxol C. 10-Deacetyltaxuyunnanine A**
[154677-95-9]
C₄₄H₅₅NO₁₃ 805.917
From cell cultures of *Taxus baccata* and from *Taxus mairei* and *Taxus yunnanensis* (Taxaceae). Tubulin assembly promotor. Amorph. solid (MeOH). Mp 157° dec. [α]_D -14.78 (c, 0.115 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep). λ_{\max} 236 ; 272 (MeOH) (Berdy).
- 10-De-Ac, N-debenzoyl, N-hexanoyl, 7-O- β -D-xylopyranoside: 10-Deacetyl-7-xylosyltaxol C**
[90332-65-3]
C₄₉H₆₃NO₁₇ 938.033
Constit. of *Taxus baccata*. Cryst. (EtOH aq.). Mp 215-217°. [α]_D +3 (Py). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 10-De-Ac, 7-Ac: 7-Acetyl-10-deacetyltaxol**
[115842-86-9]
C₄₇H₅₁NO₁₄ 853.918
Constit. of *Taxus canadensis*. Gum. [α]_D²² -44 (c, 0.05 in CHCl₃).
- 10-De-Ac, 10-(3-hydroxybutanoyl): 10-Deacetyl-10-(3-hydroxybutyryl)taxol**
[90332-68-6]
C₄₉H₅₅NO₁₅ 897.971
Constit. of *Taxus baccata*. λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 10-De-Ac, 10-(3-hydroxybutanoyl), N-debenzoyl, N-tigloyl: 10-Deacetyl-10-(β -hydroxybutyryl)cephalomannine. 10-Deacetyl-10-(β -hydroxybutyryl)-taxol B**
[90332-69-7]
C₄₇H₅₇NO₁₅ 875.965
Constit. of *Taxus baccata*. λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 2-O-Debenzoyl, 2-tigloyl: 2-Debenzoyl-2-tigloylpaclitaxel**
[173101-54-7]
C₄₅H₅₃NO₁₄ 831.912
Constit. of *Taxus x media* cv. Hicksii. Powder. Mp 232° dec. [α]_D²⁵ -44 (c, 0.7 in MeOH).
- 13-Deacyl, 13-cinnamoyl(E-): Taxuspinnanane J**
[212071-26-6]
C₄₀H₄₄O₁₂ 716.78
Constit. of *Taxus cuspidata* var. *nana*. Amorph. powder. [α]_D -43.8 (c, 0.14 in CHCl₃). λ_{\max} 205 (log € 4.05); 232 (log € 4.1); 274 (log € 3.02); 282 (log € 2.94) (MeOH).
- 10-Ketone, 10-de-Ac, 7-Ac: 10-Deacetyl-10-dehydro-7-acetyltaxol A**
[171864-27-0]
C₄₇H₄₉NO₁₄ 851.902
Constit. of *Taxus x media*. Cryst. Mp 226-230°. [α]_D²⁵ -57 (c, 2.8 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 9 α -Alcohol: Yunnanxamine. 9-Deoxo-9 α -hydroxytaxol. Dihydrotaxol**
[148584-53-6]
C₄₇H₅₃NO₁₄ 855.934
From *Taxus yunnanensis* (Taxaceae). Powder. Mp 174-176°. [α]_D²² -13.1 (c, 0.08 in CHCl₃). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 7-(Methylthio)methyl: BMS 184476**
[160237-25-2]
C₄₉H₅₅NO₁₄S 914.038
Antineoplastic agent. Characterised by pmr.
- 7-Epimer: 7-Epitaxol**
[105454-04-4]
C₄₇H₅₁NO₁₄ 853.918
From *Taxus brevifolia* (Taxaceae). Tubulin assembly promotor. Cryst. Mp 168-171°. [α]_D²³ -32.3 (c, 0.012 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 7-Epimer, 2'-Ac: 2'-Acetyl-7-epitaxol**
[172586-62-8]
C₄₉H₅₃NO₁₅ 895.955
Constit. of *Taxus canadensis*. [α]_D²² -30 (c, 0.05 in CHCl₃).
- 7-Epimer, 10-de-Ac: 10-Deacetyl-7-epitaxol. Ormosin VI**
[78454-17-8]
C₄₅H₄₉NO₁₃ 811.881
Isol. from callus cultures of *Taxus baccata*, the bark of *Taxus yunnanensis* and from *Taxus cuspidata*. Tubulin assembly promotor.
- 7-Epimer, N-debenzoyl, N-(2-methylbutanoyl): Taxoline**
[213539-42-5]
C₄₅H₅₅NO₁₄ 833.928
Constit. of *Taxus yunnanensis*. Cryst. Mp 184-186°. [α]_D²⁰ -79.5 (c, 0.515 in MeOH). λ_{\max} 230 (log € 4.21); 274 (log € 3.01) (MeOH).
- 7-Epimer, N-debenzoyl, N-tigloyl: 7-Epi-cephalomannine. 7-Epitaxol B**
[150547-36-7]
C₄₅H₅₃NO₁₄ 831.912
Constit. of *Taxus x media* cv. Hicksii and *Taxus yunnanensis*. Cryst. Mp 187-189° Mp 210° dec. [α]_D -7.3 (c, 0.2 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 7-Epimer, N-debenzoyl, N-hexanoyl: 7-Epitaxuyunnanine A**
[156130-25-5]
C₄₆H₅₇NO₁₄ 847.955
Alkaloid from *Taxus yunnanensis*. [α]_D -47.3 (CHCl₃).
- 7-Epimer, 2'-Ac, N-debenzoyl, N-tigloyl: 2'-Acetyl-7-epicephalomannine**
[340739-83-5]
C₄₇H₅₅NO₁₅ 873.949
Constit. of *Taxus canadensis*. Gum. [α]_D²⁴ +15 (c, 0.01 in CHCl₃).
- 7-Epimer, 10-de-Ac, N-debenzoyl, N-hexanoyl: 7-Epi-10-deacetyltaxuyunnanine A. Taxuspinnanane E**
[156130-26-6]
C₄₄H₅₅NO₁₃ 805.917
Alkaloid from *Taxus yunnanensis* and *Taxus cuspidata* var. *nana*. [α]_D -34.9 (CHCl₃). [α]_D -22.9 (c, 0.1 in MeOH). λ_{\max} 204 (log € 4.38); 218 (log € 4.25); 228 (log € 4.2); 272 (log € 3.18); 282 (log € 3.17) (MeOH).
- 7-Epimer, 10-de-Ac, 10-ketone: 10-Deacetyl-10-oxo-7-epi-taxol**
[105377-71-7]
C₄₅H₄₇NO₁₃ 809.865
From *Taxus brevifolia*. Tubulin assembly promotor. Oil. [α]_D²³ -60.4 (c, 0.002 in MeOH). λ_{\max} 227 (€ 29800); 273 (€ 1700) (MeOH) (Derep).
- 7-Epimer, 10-de-Ac, 10-ketone, N-debenzoyl, N-tigloyl: 10-Deacetyl-10-oxo-7-epicephalomannine**
[213276-08-5]
C₄₃H₄₉NO₁₃ 787.859
Constit. of *Taxus canadensis*. Gum. [α]_D²⁴ -1.2 (c, 0.05 in CHCl₃).
- 7-Epimer, 10-de-Ac, 10-ketone, N-debenzoyl, N-hexanoyl: 7-Epi-10-deacetyl-10-oxotaxuyunnanine A**

- [156053-35-9]
 $C_{44}H_{53}NO_{13}$ 803.902
 Constit. of *Taxus yunnanensis*. $[\alpha]_D -70.8$ (CHCl₃). The name assigned is not strictly correct.
- Docosahexaenoic acid conjugate: Docosahexaenoic acid-paclitaxel. DHA-paclitaxel*
 Inert prodrug composed of the natural fatty acid DHA covalently linked to the 2-position of paclitaxel.
- Wani, M.S. *et al.*, *J.A.C.S.*, 1971, **93**, 2325-2327 (*isol, nmr, ms, ir, uv*)
 McLaughlin, J.L. *et al.*, *J. Nat. Prod.*, 1981, **44**, 312-319 (*Cephalomannine, 10-Deacetylcephalomannine, 10-Deacetyltaxol*)
 Miller, R.W. *et al.*, *J.O.C.*, 1981, **46**, 1469-1474 (*Cephalomannine*)
 Kingston, D.G.I. *et al.*, *J. Nat. Prod.*, 1982, **45**, 466-470 (*2',7-di-Ac*)
 S nith, V. *et al.*, *J. Nat. Prod.*, 1984, **47**, 131-137 (*xylosides*)
 Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 1-18 (*rev, pharmacol*)
 Horwitz, S.B. *et al.*, *Ann. N.Y. Acad. Sci.*, 1986, **466**, 733-744 (*rev, pharmacol*)
 Huang, C.H.O. *et al.*, *J. Nat. Prod.*, 1986, **49**, 665-669 (*7-Epitaxol, 10-Deacetyl-10-oxo-7-epitaxol*)
 Blechert, S. *et al.*, *Alkaloids (Academic Press)*, 1990, **39**, 195-238 (*rev, pharmacol*)
 Kingston, D.G.I. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1-12; 802 (*bibl, rev*)
 Slichenmyer, W.J. *et al.*, *Anti-Cancer Drugs*, 1991, **2**, 519-530 (*rev*)
 Kingston, D.G.I. *et al.*, *Pharmacol. Ther.*, 1991, **52**, 1-34 (*rev, pharmacol*)
 Rose, W.C. *et al.*, *Anti-Cancer Drugs*, 1992, **3**, 311-321 (*rev*)
 Potier, P. *et al.*, *Chem. Soc. Rev.*, 1992, **21**, 113-119 (*rev*)
 Young, D.H. *et al.*, *Experientia*, 1992, **48**, 882-885 (*antifungal activity*)
 Chmurny, G.N. *et al.*, *J. Nat. Prod.*, 1992, **55**, 414-423 (*pmr, cmr*)
 Wheeler, N.C. *et al.*, *J. Nat. Prod.*, 1992, **55**, 432-440 (*occur*)
 Hilton, B.D. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1157-1161 (*pmr*)
 Falzone, C.J. *et al.*, *Tet. Lett.*, 1992, **33**, 1169-1172 (*pmr, cmr*)
 Winkler, J.D. *et al.*, *Tetrahedron*, 1992, **48**, 6953-6964 (*rev*)
 Ojima, I. *et al.*, *Tetrahedron*, 1992, **48**, 6985-7012 (*synth*)
 Horwitz, S.B. *et al.*, *Trends Pharmacol. Sci.*, 1992, **13**, 134-136 (*rev*)
 Chen, W.M. *et al.*, *Chin. Chem. Lett.*, 1993, **4**, 699-702 (*Yunnanxamine*)
 Wall, M.E. *et al.*, *Chron. Drug Discovery*, 1993, **3**, 327-348 (*rev*)
 Fleming, P.E. *et al.*, *J.A.C.S.*, 1993, **115**, 805-807 (*biosynth*)
 Stierle, A. *et al.*, *Science (Washington, D.C.)*, 1993, **260**, 214-217 (*isol, ms*)
 Williams, H.J. *et al.*, *Tetrahedron*, 1993, **49**, 6545-6560 (*pmr, conformn*)
 Zhang, H. *et al.*, *Yunnan Zhiwu Yanjiu*, 1993, **15**, 424-426; *CA*, **121**, 78282e
 Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 1994, **33**, 15-44 (*rev*)
 Joel, S.P. *et al.*, *Chem. Ind. (London)*, 1994, 172-175 (*rev*)
 Sonnichsen, D.S. *et al.*, *Clin. Pharmacokinet.*, 1994, **27**, 256-269 (*rev, pharmacokinet*)
Eur. Pat., 1994, ((*Bristol-Myers Squibb*))604 910; *CA*, **122**, 81691p (*BMS 184476, synth, pharmacol*)
 Hongjie, Z. *et al.*, *Heterocycles*, 1994, **38**, 975-980
 Ma, W. *et al.*, *J. Nat. Prod.*, 1994, **57**, 116-122 (*Taxucultine, Taxol C*)
 Hoke, S.H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 277-286 (*ms, hplc*)
 Appendino, G. *et al.*, *J. Nat. Prod.*, 1994, **57**, 607-613 (*Taxol D*)
 Kerns, E.H. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1391-1403 (*ms, hplc*)
 Gimon, M.E. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1404-1410 (*ms*)
 Holton, R.A. *et al.*, *J.A.C.S.*, 1994, **116**, 1597-1598; 1599-1600 (*synth*)
 Fleming, P.E. *et al.*, *J.A.C.S.*, 1994, **116**, 4137-4138 (*biosynth*)
 Nicolaou, K.C. *et al.*, *Nature (London)*, 1994, **367**, 630-634 (*synth*)
 Barboni, L. *et al.*, *Phytochemistry*, 1994, **36**, 987-990 (*Taxol C, N-Methyltaxol C*)
 Fleming, P.E. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2045-2048 (*biosynth*)
 Kingston, D.G.I. *et al.*, *Tet. Lett.*, 1994, **35**, 4483-4484 (*synth*)
 Chen, W.M. *et al.*, *Yaoxue Xuebao*, 1994, **29**, 751-757 (*Yunnanxamine*)
 Georg, G.I. *et al.*, *ACS Symp. Ser.*, 1995, **583**, (*book*)
 Gao, Q. *et al.*, *Acta Cryst. C*, 1995, **51**, 295-298 (*cryst struct*)
 Zamir, L.O. *et al.*, *Can. J. Chem.*, 1995, **73**, 655-665 (*pmr, cmr, 7-Epitaxol*)
 De Bellis, P. *et al.*, *Fitoterapia*, 1995, **66**, 521-524 (*7-Epicephalomannine*)
 Huizang, M.T. *et al.*, *J. Chromatogr. B*, 1995, **664**, 373-382 (*hplc*)
 Grothaus, P.G. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1003-1014 (*immunoassay*)
 Guo, Y. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1906-1912 (*7-Xylosyl-10-deacetyltaxol D*)
 Nicolaou, K.C. *et al.*, *J.A.C.S.*, 1995, **117**, 624-633; 634-644; 645-652; 653-659 (*synth*)
 Rowinsky, E.K. *et al.*, *N. Engl. J. Med.*, 1995, **332**, 1004-1014 (*use, tox, rev*)
 Nogales, E. *et al.*, *Nature (London)*, 1995, **375**, 424-427 (*taxol-tubulin heterodimer, cryst struct*)
 Gabetta, B. *et al.*, *Phytochemistry*, 1995, **40**, 1825-1828 (*10-Deacetyl-10-dehydro-7-acetyltaxol A*)
 Vyas, D.M. *et al.*, *Prog. Med. Chem.*, 1995, **32**, 289-337 (*rev*)
Taxol: Science and Applications, Ed., Suffness, M., CRC Press, Boca Raton, USA, 1995, (*book*)
The Chemistry and Pharmacology of Taxol and its Derivatives, (Ed. Farina, V.), Elsevier, 1995, (*book*)
 Li, C. *et al.*, *Anti-Cancer Drugs*, 1996, **7**, 642-648 (*peg-paclitaxel*)
 Nicolaou, K.C. *et al.*, *Classics in Total Synthesis, Targets, Strategies, Methods*, VCH, 1996, 655 (*bibl, synth*)
 Danishefsky, S.J. *et al.*, *J.A.C.S.*, 1996, **118**, 2843-2859 (*synth*)
 Heinstein, P. *et al.*, *J.C.S. Perkin 1*, 1996, 845-851 (*ms*)
 Strobel, G. *et al.*, *Microbiology (Reading, U.K.)*, 1996, **142**, 435-440 (*Taxol[®], isol, biosynth*)
Pat. Coop. Treaty (WIPO), 1996, ((*Biophysica Foundation*))96 38 138; *CA*, **126**, 113164v (*paclitaxel ceribate, synth*)
 Zamir, L.O. *et al.*, *Phytochemistry*, 1996, **41**, 803-805 (*Taxucultine*)
 Gao, Q. *et al.*, *Tetrahedron*, 1996, **52**, 2291-2300 (*cryst struct*)
 Chen, Z. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1996, **38**, 323-327; *CA*, **126**, 169062c (*7-Epitaxol B*)
 Rao, K.V. *et al.*, *J. Het. Chem.*, 1997, **34**, 675-680 (*synth*)
 Wender, P.A. *et al.*, *J.A.C.S.*, 1997, **119**, 2757-2758 (*synth*)
 Gennari, C. *et al.*, *J.O.C.*, 1997, **62**, 4746-4755 (*synth*)
 Hezari, M. *et al.*, *Planta Med.*, 1997, **63**, 291-295 (*biosynth, rev*)
 Li, C. *et al.*, *Cancer Res.*, 1998, **58**, 2404-2409 (*paclitaxel poliglumex*)
 Shiina, I. *et al.*, *Chem. Lett.*, 1998, 3-4 (*synth*)
 Eisenhauer, E.A. *et al.*, *Drugs*, 1998, **55**, 5-30 (*pharmacol, rev*)
 Gabetta, B. *et al.*, *Phytochemistry*, 1998, **47**, 1325-1329 (*N-Debenzoyl-N-butanoyl-10-deacetylpaclitaxel, N-Debenzoyl-N-propanoyl-10-deacetylpaclitaxel, 2-Debenzoyl-2-tigloylpaclitaxel*)
 Morita, H. *et al.*, *Phytochemistry*, 1998, **48**, 857-862 (*Taxuspinnananes I and J*)
 Morita, H. *et al.*, *Planta Med.*, 1998, **64**, 183-186 (*Taxuspinnanane E*)
 Cao, X. *et al.*, *Prep. Biochem. Biotechnol.*, 1998, **28**, 79-87 (*10-Deacetyl-7-epitaxol, Cephalomannine, isol*)
 Xu, X. *et al.*, *Zhongcaoyao*, 1998, **29**, 361-364; *CA*, **129**, 257658u (*Taxoline*)
 Mukaiyama, T. *et al.*, *Chem. Eur. J.*, 1999, **5**, 121-161 (*Taxol, synth, rev*)
 Li, C. *et al.*, *Clin. Cancer Res.*, 1999, **5**, 891-897; 2000, **6**, 2824-2834 (*paclitaxel poliglumex*)
 Ranson, M. *et al.*, *Expert Opin. Invest. Drugs*, 1999, **8**, 837-848
 Das, B. *et al.*, *Indian J. Chem., Sect. B*, 1999, **38**, 1018-1024 (*rev*)
 Gabetta, B. *et al.*, *J. Nat. Prod.*, 1999, **62**, 219-223 (*N-debenzoyl-N-cinnamoyl, N-debenzoyl-N-2-methylbutanoyl, 2-debenzoyl-2-tigloyl*)
 Baloglu, E. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1068-1071 (*synth*)
 Gunatilaka, A.A.L. *et al.*, *J.O.C.*, 1999, **64**, 2694-2703 (*BMS 188797, synth, pmr, cmr*)
Martindale, The Extra Pharmacopoeia, 32nd edn., Pharmaceutical Press, 1999, 556
 Zhang, J. *et al.*, *J. Nat. Prod.*, 2000, **63**, 929-933 (*2'-Acetyl-7-epitaxol*)
 Snyder, J.P. *et al.*, *J.A.C.S.*, 2000, **122**, 724-725 (*pmr, conformn*)
 Kusama, H. *et al.*, *J.A.C.S.*, 2000, **122**, 3811-3820 (*synth*)
 Zhang, J. *et al.*, *Phytochemistry*, 2000, **54**, 221-230 (*N-Acetyl-N-debenzoyltaxol, 2'-Acetyl-7-epicephalomannine*)
 Mandai, T. *et al.*, *Tet. Lett.*, 2000, **41**, 243-244 (*synth*)
 Seligson, A.L. *et al.*, *Anti-Cancer Drugs*, 2001, **12**, 305-313 (*paclitaxel ceribate, synth, pharmacol*)
 Niethammer, A. *et al.*, *Bioconj. Chem.*, 2001, **12**, 414-420 (*paclitaxel ceribate*)
 Damascelli, B. *et al.*, *Cancer (Philadelphia)*, 2001, **92**, 2592-2602 (*ABI 007*)
 Bradley, M.O. *et al.*, *Clin. Cancer Res.*, 2001, **7**, 3229-3238; 2003, **9**, 151-159; 3589-3597 (*DHA paclitaxel, pharmacol*)
 Richard, A. *et al.*, *Crit. Rev. Biotechnol.*, 2001, **21**, 219-232 (*paclitaxel poliglumex, rev*)
 Alstadt, T.J. *et al.*, *J. Med. Chem.*, 2001, **44**, 4577-4583 (*BMS 184476, synth, pmr, ir, pharmacol*)
 Zhang, J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 450-455 (*7-Acetyl-10-deacetyltaxol, 2'-Acetyl-7-epicephalomannine, 10-Deacetyl-10-oxo-7-epicephalomannine, 2'-Acetyl-7-epitaxol*)
 Walker, K. *et al.*, *Phytochemistry*, 2001, **58**, 1-7 (*biosynth, rev*)
 Li, S. *et al.*, *Phytochemistry*, 2001, **58**, 369-374 (*7-Xylosyltaxol D*)
 Ibrahim, N.K. *et al.*, *Clin. Cancer Res.*, 2002, **8**, 1038-1044 (*ABI 007, clinical trial*)
 Plummer, R. *et al.*, *Clin. Cancer Res.*, 2002, **8**, 2788-2797 (*BMS 184476, pharmacol*)
 Singer, J.W. *et al.*, *Adv. Exp. Med. Biol.*, 2003, **519**, 81-99 (*CT 2103, rev*)

- Advani, R. *et al.*, *Clin. Cancer Res.*, 2003, **9**, 5187-5194 (*BMS 188797, pharmacol*)
- Ojima, I. *et al.*, *Curr. Opin. Invest. Drugs*, 2003, **4**, 732-736 (*BMS 184476, rev*)
- Drugs RD*, 2003, **4**, 303-305 (*ABI 007, rev*)
- Li, S.-H. *et al.*, *Tetrahedron*, 2003, **59**, 37-45 (7-Xylosyl-N-methyltaxol C)
- Clarke's *Analysis of Drugs and Poisons*, 3rd edn., (eds. Moffat, A.C. *et al.*), Pharmaceutical Press, 2004, 1387 (*props, uv, ir, hplc*)
- Simpson, D. *et al.*, *Drugs*, 2004, **64**, 1839-1847 (*rev*)
- Oberlies, N.H. *et al.*, *J. Nat. Prod.*, 2004, **67**, 129-135 (*rev*)
- Horowitz, S.B. *et al.*, *J. Nat. Prod.*, 2004, **67**, 136-138 (*rev*)
- Man, M. *et al.*, *IDrugs*, 2005, **8**, 739-754 (*paclitaxel poliglumex, rev*)
- Doi, T. *et al.*, *Chem. Asian J.*, 2006, **1**, 370-383 (*synth*)
- Robinson, D.M. *et al.*, *Drugs*, 2006, **66**, 941-948 (*rev*)
- Ho, Y. *et al.*, *Magn. Reson. Chem.*, 2006, **44**, 581-585 (*solid-state nmr*)
- Boddy, A.V. *et al.*, *Drugs of the Future*, 2007, **32**, 776 (*paclitaxel poliglumex*)
- Lin, N.U. *et al.*, *Invest. New Drugs*, 2007, **25**, 369-375 (*paclitaxel poliglumex, pharmacol*)
- Kingston, D.G.I. *et al.*, *Phytochemistry*, 2007, **68**, 1844-1854 (*rev*)
- Hu, Z. *et al.*, *Chin. Chem. Lett.*, 2008, **19**, 130-132 (*synth*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, TAH775

Tazettine**T-53**

Ungernine. Sekisanine. Sekisanoline
[507-79-9]
[28405-99-4 ((±)-form)]



Absolute
Configuration

$C_{18}H_{21}NO_5$ 331.368
Isol. from *Narcissus tazetta*, *Hymenocallis expansa*, *Hippeastrum* spp., *Narcissus confusus* and a very large number of other spp. in the Amaryllidaceae. Weak hypotensive agent. Mp 210-211°. $[\alpha]_D^{16} +150$ ($CHCl_3$). Has been shown to be an artifact produced by base-catalysed rearr. of Pretazettine, P-626 in two cases studied, and is probably an artifact in all cases.

► LD₅₀ (dog, ivn) 71 mg/kg.

Picrate: Mp 205-208°.

Ac: Mp 125-126.5°.

O-De-Me: **Tazettinol**. O-Demethyltazettine
[560-31-6]

$C_{17}H_{19}NO_5$ 317.341

Alkaloid from bulbs of *Hippeastrum equestre*. Mp 179-181°. $[\alpha]_D^{22} +85$ (c, 1 in MeOH).

O-De-Me, O⁶-Me: **Zeylamine**. 6-O-Methyltazettinol
[110815-91-3]

Alkaloid from the rhizomes of *Crinum zeylanicum*. Cryst. (Me₂CO). Mp 138-140°. The stereochem. may not be certain.

O-De-Me, 3-O-(3-hydroxybutanoyl): **3-O-(3-Hydroxybutanoyl)tazettinol**
[343983-03-9]

$C_{21}H_{25}NO_7$ 403.431

Alkaloid from *Galanthus plicatus* ssp. *byzantinus*. Amorph. solid. $[\alpha]_D +63.5$ (c, 0.22 in MeOH). λ_{max} 215 (log ϵ 4.12); 234 (log ϵ 3.87); 292 (log ϵ 3.6) (MeOH).

1,2-Dihydro, Me ether: **Ungvedine**. 1,2-Dihydro-O-methyltazettine, 9CI
[73276-40-1]

$C_{19}H_{25}NO_5$ 347.41

Alkaloid from *Ungernia vvedenskyi* (Amaryllidaceae). Mp 148-150°. $[\alpha]_D +12.5$ (c, 0.528 in $CHCl_3$).

3-Epimer: **Crivelline**. 3-Epitazettine
[545-16-4]

[60944-17-4 ((±)-form)]

$C_{18}H_{21}NO_5$ 331.368

Alkaloid from *Crinum powellii*, *Crinum powellii* var. *album*, *Crinum macrantherum*, *Crinum erubescens* and *Galanthus nivalis* (Amaryllidaceae). Cryst. (Me₂CO). Mp 214° (205-206°). $[\alpha]_D^{25} +220$ (c, 0.15 in $CHCl_3$). $[\alpha]_D^{23} +288$ (c, 1.0 in $CHCl_3$).

3-Epimer, hydroiodide:

Prisms (H₂O). Mp 228-229° dec.

3-Epimer, picrate: Mp 233°.

3-Epimer, Me ether: Mp 127-128°. $[\alpha]_D^{25} +214$ (EtOH).

3-Epimer, O-de-Me: **Isotazettinol**. O-De-methylcrivelline

[465-63-4]

$C_{17}H_{19}NO_5$ 317.341

Alkaloid from *Galanthus gracilis*. Cryst. (Me₂CO). Mp 204°. $[\alpha]_D +194$ (c, 0.14 in MeOH) (natural). $[\alpha]_D +261.7$ (c, 0.5 in EtOH) (synthetic). λ_{max} 236 (sh) (log ϵ 4.07); 290 (log ϵ 3.65) (MeOH).

Boit, H.G. *et al.*, *Chem. Ber.*, 1956, **89**, 2093 (*Crivelline, isol*)

Ikeda, T. *et al.*, *J.C.S.*, 1956, 4749-4761 (*struct*)

Fales, M.M. *et al.*, *Chem. Ind. (London)*, 1959, 1415 (*Crivelline, struct*)

Jeffs, P.W. *et al.*, *J.C.S.*, 1960, 1090-1094 (*Crivelline, struct*)

Duffield, A.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4902-4912 (*ms*)

Haugwitz, R.D. *et al.*, *J.C.S.*, 1965, 2001-2009 (*pmr*)

Highet, R.J. *et al.*, *Tet. Lett.*, 1966, 4099-4101 (*config*)

Sato, T. *et al.*, *J.C.S. (B)*, 1971, 1070-1073 (*cryst struct*)

Hendrickson, J.B. *et al.*, *J.A.C.S.*, 1974, **96**, 7781-7789 (*synth*)

Isobe, K. *et al.*, *Tet. Lett.*, 1976, 2331-2334 (*Crivelline, synth*)

Kadyrov, K.A. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 585-586; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 513-514 (*Ungvedine*)

Danishefsky, S. *et al.*, *J.A.C.S.*, 1982, **104**, 7591-7599 (*synth, ir, pmr*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 203 (*occur*)

Döpke, W. *et al.*, *Z. Chem.*, 1986, **26**, 438 (*Zeylamine*)

Abelman, M.M. *et al.*, *J.A.C.S.*, 1990, **112**, 6959-6964 (*synth*)

Wagner, J. *et al.*, *Tetrahedron*, 1996, **52**, 6591-6600 (*cd*)

Ide, S. *et al.*, *Kristallografiya*, 1997, **42**, 88-92 (*cryst struct*)

Linden, A. *et al.*, *Acta Cryst. C*, 1998, **54**, 1653-1659 (*Tazettine, Isotazettinol, cryst struct*)

Rigby, J.H. *et al.*, *J.A.C.S.*, 1998, **120**, 3664-3670 (*synth*)

Pham, L.H. *et al.*, *Phytochemistry*, 1999, **51**, 327-332 (*pmr, cmr, Tazettinol*)

Unver, N. *et al.*, *Planta Med.*, 1999, **65**, 347-350 (*Isotazettinol*)

Baldwin, S.W. *et al.*, *Org. Lett.*, 2000, **2**, 99-102 (*synth*)

Unver, N. *et al.*, *Heterocycles*, 2001, **55**, 641-652 (*3-Hydroxybutanoyltazettinol*)

Zhang, F.-M. *et al.*, *Tetrahedron*, 2006, **62**, 9446-9455 (*synth*)

Tecleanine†**T-54**

$C_{28}H_{36}N_2O_4$ 464.603

Struct. unknown. Alkaloid from the bark of *Teclea natalensis* (Rutaceae). Needles (C₆H₆). Mp 204-205°. $[\alpha]_D^{24} -245$ (c, 0.45 in EtOH).


Hydrochloride (1:2):

Prisms (EtOH aq.). Mp 300° dec.

Wright, W.G. *et al.*, *J.C.S. (C)*, 1967, 2262

Tecleanone**T-55**

2-[(Methylamino)phenyl](2,4,6-trimethoxyphenyl)methanone, 9CI. 2'-Methylamino-2,4,6-trimethoxybenzophenone
[55950-37-3]



$C_{17}H_{19}NO_4$ 301.341

Alkaloid from *Teclea grandifolia*, *Teclea verdoorniana*, *Diphasia klaineana* and *Oricia suaveolens* (Rutaceae). Yellow solid. Mp 190-191°. Structurally related to an intermed. in the biosynth. of acridone alkaloids.


N-Me: Mp 128°.

Casey, A. *et al.*, *Tet. Lett.*, 1975, 401 (*ir, pmr, ms, isol, struct*)

Waterman, P. *et al.*, *Phytochemistry*, 1975, **14**, 2092 (*isol, uv, ir, pmr, ms*)

Tecleanthine**T-56**

6,11-Dimethoxy-5-methyl-1,3-dioxo[4,5-b]acridin-10(5H)-one, 9CI. 1,5-Dimethoxy-10-methyl-2,3-methylenedioxyacridone
[24966-05-0]



C₁₇H₁₅NO₅ 313.309

Alkaloid from the bark of *Teclea natalensis* and other *Teclea* spp. (Rutaceae). Yellow prisms (EtOH). Mp 158°. λ_{\max} 235 ; 268 ; 401 (MeOH) (Berdy). λ_{\max} 236 (ϵ 17000); 269 (ϵ 55000); 401 (ϵ 11200) (EtOH) (Berdy).

Picrate:

Needles (EtOH). Mp 171°.

O¹¹-De-Me: 1-Hydroxy-5-methoxy-10-methyl-2,3-methylenedioxyacridone. O-Nortecleanthine
Orange plates (EtOH). Mp 207°.


7-Methoxy: 1,5,6-Trimethoxy-10-methyl-2,3-methylenedioxyacridone. 6,7,11-Trimethoxy-5-methyl-1,3-dioxolo[4,5-b]acridin-10(5H)-one, 9CI
[52911-51-0]

C₁₈H₁₇NO₆ 343.335

Alkaloid from *Teclea boiviniana* (Rutaceae). Yellow cryst. (CHCl₃). Mp 168°. λ_{\max} 221 (ϵ 14000); 273 (ϵ 55000); 323 (ϵ 5600); 392 (ϵ 7600) (EtOH) (Berdy).

Pegel, K.H. *et al.*, *J.C.S. (C)*, 1969, 2327-2329 (*isol, uv, ir, pmr, ms, struct*)Ramachandran, V.N. *et al.*, *Indian J. Chem.*, 1972, **10**, 14-15 (*synth, uv, ir, pmr*)Laing, M. *et al.*, *J. S. Afr. Chem. Inst.*, 1974, **27**, 162-163; *CA*, **82**, 148605q (*cryst struct*)Vaquette, J. *et al.*, *Plant. Med. Phytother.*, 1974, **8**, 57-62 (7-methoxy, *isol, struct*)Rasoanaivo, P *et al.*, *Fitoterapia*, 1999, **70**, 625-627 (*isol, cmr*)**Tecleaverdoornine****T-57**

9-Methoxy-10-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-g]furo[2,3-b]quinolin-4-ol, 9CI. 8-Hydroxy-6,7-methylenedioxy-5-prenyldictamine
[65847-03-2]

C₁₈H₁₇NO₅ 327.336

Alkaloid from the stem bark of *Teclea verdoorniana* (Rutaceae). Long needles (Me₂CO). Mp 191°.

Ac:

Cryst. (MeOH). Mp 212°.


Me ether:

Cryst. (Me₂CO). Mp 139-140°.3',4'-Dihydro, 4'-hydroxy: **Tecleaverdine**
[82202-84-4]C₁₈H₁₉NO₆ 345.351

Alkaloid from stem bark of *Teclea verdoorniana* (Rutaceae). Cryst. (MeOH). Mp 218-219°.

Ayafor, J.F. *et al.*, *J.C.S. Perkin 1*, 1982, 309 (*isol, uv, ir, pmr, ms, struct*)**Tecomanine**

Tecomine[†]
[6878-83-7]



Absolute configuration

C₁₁H₁₇NO 179.261

Major alkaloid from *Tecoma stans* and *Tecoma fulva* (Bignoniaceae). Hypoglycaemic agent. *Tecoma* extracts have been used in Mexico for control of diabetes. Bp_{0.1} 125°. $[\alpha]_{\text{D}}^{24}$ -175 (c, 1.17 in CHCl₃). Log P 0.49 (calc).

▶WX8575000

Picrate: Mp 179.5-180.5°.

2,4-Dinitrophenylhydrazone: Mp 260°.


4-Hydroxy: 4-Hydroxytecomanine

C₁₁H₁₇NO₂ 195.261

Alkaloid from fruits of *Tecoma stans* (Bignoniaceae). Oil.

Dickinson, E.M. *et al.*, *Tetrahedron*, 1969, **25**, 1523 (*ir, uv, pmr, isol, struct*)Jones, G. *et al.*, *Chem. Comm.*, 1971, 994 (*cryst struct*)Berg, W. *et al.*, *Pharmazie*, 1977, **32**, 41 (*ms*)
Imanishi, T. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 1243 (*synth*)Miyashita, M. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 1614 (*synth*)Lins, A.P. *et al.*, *Phytochemistry*, 1993, **34**, 876-878 (4-Hydroxytecomanine)Kemp, M.I. *et al.*, *Synthesis*, 1998, 552-556 (*synth*)Ockey, D.A. *et al.*, *Tetrahedron*, 2003, **59**, 5377-5381 (*synth*)**Tecostanine****T-59**

[708-18-9]




Absolute Configuration

C₁₁H₂₁NO 183.293

Alkaloid from the leaves of *Tecoma stans* (Bignoniaceae). Mp 85°. $[\alpha]_{\text{D}}^{20}$ 0 (MeOH). $[\alpha]_{\text{D}}$ +6 (c, 2 in CHCl₃).

▶WX8690000

Methiodide:

Cryst. (MeOH/Et₂O). Mp 243-246°.Hammouda, Y. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2802 (*uv, ir, pmr, struct*)Costantino, L. *et al.*, *Pharmazie*, 2003, **58**, 140-142 (*isol, pmr, cmr, cryst struct*)**Tecostidine****T-60**

(R)-form

C₁₀H₁₃NO 163.219**(R)-form**

Alkaloid from the upper parts of *Pedicularis rhinanthoides* (Scrophulariaceae). Viscous oil. Bp₁ 150° (bath). $[\alpha]_{\text{D}}^{20}$ +5.9 (c, 0.508 in EtOH). $[\alpha]_{\text{D}}^{21}$ +3 (c, 1.1 in CHCl₃) (synthetic).

Picrate:

Yellow needles (EtOH/Et₂O). Mp 151-152°.**(S)-form**

Alkaloid from the leaves of *Tecoma stans* (Bignoniaceae). Viscous liq. $[\alpha]_{\text{D}}^{25}$ -4 (c, 1.221 in CHCl₃).

Picrate:

Cryst. (EtOH/Et₂O). Mp 152-153°.Me ether: **Valerianine**

[30634-66-3]

[30769-38-1]

Alkaloid from roots of *Valeriana officinalis* (valerian) (Valerianaceae). Mp 134° (as hydrochloride). $[\alpha]_{\text{D}}^{20}$ -10.5 (c, 0.37 in MeOH) (hydrochloride).

Hammouda, Y. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 2901 (*uv, ir, pmr, struct*)Cavill, G.W.K. *et al.*, *Aust. J. Chem.*, 1967, **20**, 349 (*synth, uv, ir, pmr*)Abdusamatov, A. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 334; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 285 (*isol*)Franck, B. *et al.*, *Angew. Chem., Int. Ed.*, 1970, **9**, 891 (*Valerianine*)Ranarivelo, Y. *et al.*, *Heterocycles*, 1990, **31**, 1727-1731 ((S)-form, *synth*)Robert, N. *et al.*, *Tetrahedron*, 2007, **63**, 3702-3706 (*synth*)**Tejeramycin****T-61**

[111643-99-3]

Nucleoside-peptide antibiotic complex. Struct. unknown. Prod. by *Streptomyces griseus* ATCC39208. Active against gram-positive and -negative bacteria.

Tejeramycin A [111644-00-9]Powder. Sol. H₂O; poorly sol. butanol, hexane.**Tejeramycin B** [111644-01-0]Powder. Sol. H₂O; poorly sol. butanol, hexane.**Tejeramycin C** [111644-02-1]Powder. Sol. H₂O; poorly sol. butanol, hexane.**Tejeramycin D** [111644-03-2]Powder. Sol. H₂O; poorly sol. butanol, hexane.**Tejeramycin E** [111644-05-4]Powder. Sol. H₂O; poorly sol. butanol, hexane.**Tejeramycin F** [111644-06-5]

[111644-07-6, 111644-08-7]


Powder. Sol. H₂O; poorly sol. butanol, hexane. λ_{\max} 215 ; 258 (H₂O). λ_{\max} 215 (E1%/1cm 340); 258 (E1%/1cm 206) (H₂O) (Berdy).

Tejeramycin G [111644-09-8]Powder. Sol. H₂O; poorly sol. butanol, hexane. λ_{\max} 215 (E1%/1cm 354); 258

(E1%/1cm 200) (H₂O) (Berdy).
U.S. Pat., 1987, 4 656 036; CA, 107, 234820x

Telazoline**T-62**

1-Amino-2-methoxy-7H-dibenzo[de,g]-quinolin-7-one, 9CI
[78432-75-4]



C₁₇H₁₂N₂O₂ 276.294

Struct. revised in 2003. Unusual aminoaporphine alkaloids. Alkaloid from the woody stems of *Telitoxicum peruvianum* (Menispermaceae). Reddish-brown prisms (MeOH/CH₂Cl₂). Mp 240-243°.

3-Methoxy: **Teladiazoline**. 1-Amino-2,3-dimethoxy-7H-dibenzo[de,g]quinolin-7-one

[119763-88-1]

C₁₈H₁₄N₂O₃ 306.32


Alkaloid from the woody stems of *Telitoxicum glaziovii* (Menispermaceae). Reddish-orange needles (CHCl₃/MeOH). Mp 197-199° dec. Struct. revised in 2003.

Menachery, M.D. et al., *J. Nat. Prod.*, 1981, 44, 320; 1988, 51, 1283 (*Telazoline*, *Teladiazoline*)

Killmer, L. et al., *J. Nat. Prod.*, 2003, 66, 115-118 (*pmr, cmr, ms, struct*)

Teleocidin B₁**T-63**

De-O-methylolivoretin B
[95044-71-6]



Absolute Configuration

C₂₈H₄₁N₃O₂ 451.651

Prod. by *Streptomyces mediocidicus*. Nematocide and acaricide. Prisms (diisopropyl ether). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 153-155°. [α]_D²⁰ -132 (MeOH). Closely related to Lyngbyatoxin A, L-367. λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

► Tumour promotor. Skin irritant.

14-Ac: 14-O-Acetylteleocidin B₃

C₃₀H₄₃N₃O₃ 493.688

Prod. by *Actinomyces* sp. Stamm K 17/7. Yellow oil. [α]_D²⁰ -76 (c, 0.1 in MeOH).

N-De-Me: **N-Demethylteleocidin B₁**

C₂₇H₃₉N₃O₂ 437.624

Prod. by *Streptomyces mediocidicus*. Gum.

Me ether: **Olivoretin B**. 16-Epiolivoretin A
[90599-27-2]

C₂₉H₄₃N₃O₂ 465.678

Prod. by *Streptovorticillium olivoreticuli*. Needles (MeOH). Sol. MeOH, CHCl₃. Mp 277.5-279°. [α]_D²⁰ -298.4 (c, 0.35 in CHCl₃). λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

► Strong vesicant, poss. tumour promotor.

16-Epimer: **Teleocidin B₄**. Olivoretin D
[11032-05-6]

C₂₈H₄₁N₃O₂ 451.651

Prod. by *Streptomyces mediocidicus* and *Streptovorticillium olivoreticuli*. Prisms (diisopropyl ether). Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O, bases, acids. Mp 230-232.5°. λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

► Very toxic by intraperitoneal and intravenous routes. LD₅₀ (mus, ivn) 0.22 mg/kg. WY1982000

16-Epimer, N-de-Me: **N-Demethylteleocidin B₄**

[110200-23-2]

C₂₇H₃₉N₃O₂ 437.624

Prod. by *Streptovorticillium mediocidicus*, *Streptovorticillium blastmyceticum* and *Streptovorticillium olivoreticuli*. Tumour promotor. Gummy solid.

16-Epimer, Me ether: **Olivoretin A**

[90297-52-2]

C₂₉H₄₃N₃O₂ 465.678

Prod. by *Streptovorticillium olivoreticuli* and *Streptovorticillium blastmyceticus*. Plates (MeOH). Sol. MeOH, CHCl₃; poorly sol. cyclohexane. Mp 251-253°. [α]_D²⁰ -314.9 (CHCl₃). λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

► Strong vesicant, tumour promotor.

19-Epimer: **Teleocidin B₃**

[95189-06-3]

C₂₈H₄₁N₃O₂ 451.651

Prod. by *Streptovorticillium mediocidicus*. Prisms (diisopropyl ether). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 160-162°. [α]_D²⁰ -163 (c, 0.1 in MeOH). λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

16,19-Diepimer: **Teleocidin B₂**

[95189-05-2]

C₂₈H₄₁N₃O₂ 451.651

Prod. by *Streptovorticillium mediocidicus*. Prisms (Me₂CO). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 203-204°. λ_{max} 233 (ε 34700); 287 (ε 9770); 298 (sh) (ε 8130) (MeOH) (Derep).

Nakata, H. et al., *Tet. Lett.*, 1966, 2515

(*dihydro, synth*)

Sakabe, N. et al., *Tet. Lett.*, 1966, 2523 (*cryst struct, dihydro*)

Sakai, S. et al., *Chem. Pharm. Bull.*, 1984, 32, 354 (*Olivoretin A, Teleocidin B₄*)

Hitotsuyanagi, Y. et al., *Chem. Pharm. Bull.*, 1984, 32, 3774; 4233 (*Olivoretin B, Teleocidins B₁-B₄*)

Irie, K. et al., *Agric. Biol. Chem.*, 1985, 49, 221; 1988, 52, 3193 (*isol, deriv*)

Sakai, S. et al., *Chem. Pharm. Bull.*, 1986, 34, 4883 (*De-N-methylteleocidin B₄*)

Nakatsuka, S. et al., *Tet. Lett.*, 1987, 28, 3671 (*synth*)

Muratake, H. et al., *Tet. Lett.*, 1988, 29, 6267 (*synth*)

Okabe, K. et al., *Chem. Pharm. Bull.*, 1989, 37, 563 (*synth*)

Irie, K. et al., *Tetrahedron*, 1990, 46, 2773 (*biosynth*)


Muratake, H. et al., *Tetrahedron*, 1991, 47, 8535; 8545; 8559 (*synth*)

Ströck, K. et al., *Dissertation*, Univ. of Göttingen, 2003, (14-O-Acetylteleocidin B₃)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TAK750

Telikovinone**T-64**

4-Amino-1,2-dimethoxy-3H-dibenzo[de,g]quinoline-3,5(6H)-dione, 9CI
[166196-18-5]



C₁₈H₁₄N₂O₄ 322.32

Unusual 4-aminoaporphine alkaloids. Alkaloid from woody stems of *Telitoxicum krukovii* (Menispermaceae). Blood red needles (MeOH/CHCl₃). Mp 275-277°. Artifact.

7-Methoxy: **Telitoxinone**

[174232-33-8]


C₁₉H₁₆N₂O₅ 352.346

Alkaloid from *Telitoxicum peruvianum*. Ruby-red needles (CH₂Cl₂/MeOH). Mp 247-248° dec.

Menachery, M.D. et al., *Heterocycles*, 1995, 41, 1425 (*isol, uv, ir, pmr, cmr, ms, struct*)
Menachery, M.D. et al., *J. Nat. Prod.*, 1995, 58, 1945 (*Telitoxinone*)

Telisatin A**T-65**

[64938-92-7]



C₂₀H₁₅NO₄ 333.343

Modified aporphine. Alkaloid from *Telitoxicum peruvianum*. Also obt. by reaction of Dehydronuciferine, D-164 with oxalyl chloride. Wine-red cryst. (CH₂Cl₂/MeOH). Mp 238-239° dec. (233-234°).

3-Methoxy: **Telisatin B**

[174232-32-7]

C₂₁H₁₇NO₅ 363.369


From *Telitoxicum peruvianum*. Reddish-purple needles (CH₂Cl₂/MeOH). Mp 221-222°.

Saá, J.M. et al., *J.O.C.*, 1978, 43, 1096 (*synth*)

Menachery, M.D. et al., *J. Nat. Prod.*, 1995, 58, 1945 (*isol, uv, ir, pmr, cmr, ms*)

Telomestatin

GM 95. Antibiotic GM 95

C₂₆H₁₄N₈O₇S 582.512

Prod. by *Streptomyces anulatus* 3533-SV4. Antitumour agent. Specific telomerase inhibitor. Yellowish powder. Mp 134-143° dec. $[\alpha]_D^{25}$ -9.4 (c, 0.13 in MeOH).

Pat. Coop. Treaty (WIPO), 2000, 24 747; CA, 132, 307353a (GM 95)


Shin-ya, K. et al., *J.A.C.S.*, 2001, 123, 1262-1263

Doi, T. et al., *Org. Lett.*, 2006, 8, 4165-4167 (synth)

Tembamide

T-67

N-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]benzamide, 9CI. N-Benzoyl-2-hydroxy-2-(4-methoxyphenyl)ethylamine [37791-56-3]



(R)-form

C₁₆H₁₇NO₃ 271.315**(R)-form** [15298-30-3]

Alkaloid from the stem bark of *Zanthoxylum hyemale*. Cryst. (EtOH). Mp 156-157° (145-147°). $[\alpha]_D^{25}$ -59.4 (c, 0.57 in CHCl₃). $[\alpha]_D^{21}$ +17.9 (c, 0.5 in EtOH).

Me ether:C₁₇H₁₉NO₃ 285.342

Alkaloid from the stem bark of *Zanthoxylum hyemale*. Needles. Mp 88-89°. $[\alpha]_D^{25}$ +17.7 (c, 0.1 in CHCl₃).

(S)-form [15779-24-5]

Synthetic. Cryst. (EtOH). Mp 156-157°. $[\alpha]_D^{21}$ +55.8 (c, 0.5 in CHCl₃). $[\alpha]_D^{20}$ -18.5 (c, 0.5 in EtOH).

(±)-form [15298-28-9]

Alkaloid from the bark of *Fagara heynealis*, *Zanthoxylum tingoassuiba*, *Zanthoxylum ocumarensis*, *Zanthoxylum conspersipunctatum*, *Zanthoxylum inermis*, the root of *Aegle marmelos* (bael fruit), and the leaves and branchlets of *Clausena brevistyla* (Rutaceae). Cryst. (EtOH, EtOAc, C₆H₆, CHCl₃/petrol or CH₂Cl₂/EtOH). Mp 156-157° (147-149°).

Ac: Tembamide acetate

[15298-31-4]

C₁₈H₁₉NO₄ 313.352

T-66

Alkaloid from aerial parts of *Piper guayranum* (Piperaceae). Needles (EtOAc or C₆H₆/petrol). Mp 159° (140-141°).

Me ether: O-MethyltembamideC₁₇H₁₉NO₃ 285.342

Alkaloid from the bark of *Zanthoxylum conspersipunctatum* (Rutaceae). Cryst. (C₆H₆/petrol). Mp 98-99°.

[50802-66-9]

Johns, S.R. et al., *Aust. J. Chem.*, 1967, 20, 2795 (isol, ir, pmr, ms)

Albónico, S.M. et al., *J.C.S. (C)*, 1967, 1327 (isol, uv, ir, pmr, synth)

Johns, S.R. et al., *Aust. J. Chem.*, 1969, 22, 2233 (isol)

Della Casa de Marcano, D.

et al., *Phytochemistry*, 1972, 11, 1531 (isol)

Krajniak, E.R. et al., *Aust. J. Chem.*, 1973, 26, 687 (isol, synth, deriv)

Shoeb, A. et al., *Phytochemistry*, 1973, 12, 2071 (isol, uv, ir, pmr, ms)

Bernhard, H.O. et al., *Helv. Chim. Acta*, 1978, 61, 2269 (isol, ir, pmr, ms)

Patra, A. et al., *Org. Magn. Reson.*, 1981, 16, 65 (cmr)

Maxwell, A. et al., *J. Nat. Prod.*, 1989, 52, 411 (isol, uv, ir, pmr, cmr, ms, struct, acetate)

Brown, R.F.C. et al., *Tetrahedron: Asymmetry*, 1993, 4, 205 (synth)


De Moura, N.F. et al., *Planta Med.*, 2002, 68, 534-538 (isol, cmr)

Lee, D.-M. et al., *Tetrahedron: Asymmetry*, 2007, 18, 2662-2667 (R-form, synth, pmr, cmr, ms)

Templetine

T-68

[54274-32-7]



Absolute configuration

C₂₀H₃₅N₃ 317.517

Stereoisomeric with Ormosanine, O-115, Dasycarpine, D-85 and Piptanthine, P-476. Alkaloid from the leaves of *Templetonia retusa* (Fabaceae). Prisms (Me₂CO). Mp 120.5-122°. $[\alpha]_D$ -52 (c, 1 in EtOH).

Hydrochloride (1:3): [54274-33-8]Cryst. + 2H₂O. Mp 295-300°.**Perchlorate** (1:3):Prisms + 1H₂O (EtOH). Mp 280-282° dec.

Cannon, J.R. et al., *Aust. J. Chem.*, 1991, 44, 509 (isol, ir, ms, cryst struct, abs config)

Tenacibactin B

T-69

[934349-30-1]

(H₃C)₂CHCH₂CON(OH)(CH₂)₅NH-COCH₂CH₂COOHC₁₄H₂₆N₂O₅ 302.37

Prod. by the marine-derived *Tenacibaculum* sp. A4K-17. Siderophore. Powder. λ_{\max} 205 (log ϵ 2.72) (MeOH).

Me ester: Tenacibactin A

[934490-72-9]

C₁₅H₂₈N₂O₅ 316.397


Prod. by *Tenacibaculum* sp. A4K-17. Siderophore. Powder. λ_{\max} 205 (log ϵ 2.69) (MeOH).

Jang, J.-H. et al., *J. Nat. Prod.*, 2007, 70, 563-566 (isol, pmr, cmr)

Tenacibactin C

T-70

[934490-73-0]



R = OH

C₂₃H₄₂N₄O₈ 502.607

Prod. by the marine-derived *Tenacibaculum* sp. A4K-17. Siderophore. Powder. λ_{\max} 214 (log ϵ 3.14) (MeOH).

Jang, J.-H. et al., *J. Nat. Prod.*, 2007, 70, 563-566 (isol, pmr, cmr)

Tenacibactin D

T-71

[934490-74-1]

As Tenacibactin C, T-70 with

R = -N(OH)(CH₂)₅NH₂C₂₈H₅₄N₆O₈ 602.77


Prod. by the marine-derived *Tenacibaculum* sp. A4K-17. Siderophore. Powder. λ_{\max} 212 (log ϵ 3.12) (MeOH).

Jang, J.-H. et al., *J. Nat. Prod.*, 2007, 70, 563-566 (isol, pmr, cmr)

Tenellin

T-72

3-(4,6-Dimethyl-1-oxo-2,4-octadienyl)-1,4-dihydroxy-5-(4-hydroxyphenyl)-2(1H)-pyridinone, 9CI

C₂₁H₂₃NO₅ 369.416**(-)-form** [53823-15-7]

Pigment from *Beauveria tenella* and *Beauveria bassiana*. Greenish-yellow. $[\alpha]_D^{24}$ -44 (c, 1 in Me₂CO). Similar to Bassianin, B-28.

(±)-form [81844-67-9]

Elongated bright-yellow platelets (CCl₄/CHCl₃). Mp 174-176°.

El Basyouni, S.H. et al., *Can. J. Bot.*, 1968, 46, 441 (isol)


Leete, E. et al., *Tet. Lett.*, 1975, 4103 (biosynth)

Wat, C.K. et al., *Can. J. Chem.*, 1977, 55, 4090 (struct, ms, cmr)

- Williams, D.R. *et al.*, *J.O.C.*, 1982, **47**, 2846 (synth)
 Rigby, J.H. *et al.*, *J.O.C.*, 1989, **54**, 5852 (synth)
 Cox, R.J. *et al.*, *J.C.S. Perkin 1*, 1991, 2537 (biosynth)
 Buck, J. *et al.*, *J.C.S. Perkin 1*, 1992, 67 (synth)
 Moore, M.C. *et al.*, *Tetrahedron*, 1998, **54**, 9195-9206 (biosynth)
 Eley, K.L. *et al.*, *ChemBioChem*, 2007, **8**, 289-297 (biosynth)

Tenuamine**T-73**

N-[*Hexahydro-8(5H)-indolizinyldene*]-methyl]acetamide. Norlusitanine



C₁₁H₁₈N₂O 194.276


(R)-form [256379-00-7]

Alkaloid from *Maackia tenuifolia*. Needles (CHCl₃). Mp 136-138°. [α]_D²³ -4.5 (c, 0.12 in EtOH).

Wang, Y.-H. *et al.*, *Heterocycles*, 1999, **51**, 3001-3004

Tenuazonic acid**T-74**

3-Acetyl-1,5-dihydro-4-hydroxy-5-(1-methylpropyl)-2H-pyrrol-2-one, 9CI. 3-Acetyl-5-sec-butyltetramic acid. Vivotoxin [610-88-8]



C₁₀H₁₅NO₃ 197.233

CA refers also to two tautomeric forms. Prod. by *Alternaria tenuis*, *Alternaria longipes*, *Alternaria linicola*, *Alternaria alternata*, *Alternaria tenuissima*, *Aspergillus* sp. F1404, *Pyricularia oryzae* T2, *Aspergillus tamarii*, *Penicillium* A374, and an imperfect fungus (order Sphaeropsidales). Causes rice leaf rot and brown spot disease of tobacco. Has insecticidal props. Tremorgenic toxin. Antineoplastic and antiviral agent. Inhibits MT1-matrix metalloproteinase. Viscous oil or pale brown cryst. Sol. MeOH, Et₂O, bases; poorly sol. H₂O, hexane. Mp 74-75.5°. Bp_{0.035} 117°. [α]_D²⁰ -136 (c, 0.2 in CHCl₃). λ_{max} 219 (ε 5600); 277 (ε 12700) (EtOH/HCl) (Derep). λ_{max} 240 (ε 11700); 279 (ε 14800) (dil. NaOH) (Derep). λ_{max} 240 (ε 10800); 280 (ε 13500) (H₂O pH 7) (Derep). λ_{max} 240 (ε 11480); 280 (ε 14450) (MeOH) (Berdy). λ_{max} 220 (ε 6500); 277 (ε 10000) (MeOH/HCl) (Berdy).
 ▶ LD₅₀ (mus, orl) 225 mg/kg; LD₅₀ (mus, ivn) 125 mg/kg; LD₅₀ (mus, ipr) 150 mg/kg; LD₅₀ (mus, scu) 145 mg/kg. UY7425000

Na salt: Sodium tenuazonate. NSC 525816

[1013-59-8]

[α]_D²⁰ -96.7 (c, 2 in MeOH).

▶ LD₅₀ (rat, orl) 168 mg/kg. UY7525000

Compd. with N,N-Dibenzylethylenediamine (1:1): Benzathine tenuazonate. NSC 82260

[21327-82-2]

Antineoplastic.

2,4-Dinitrophenylhydrazonate:

Needles (MeOH). Mp 199-200°.

Semicarbazone:

Cryst. Mp 187-189°.

[75652-74-3, 27778-66-1]

Rosett, T. *et al.*, *Biochem. J.*, 1957, **67**, 390 (isol)

Stickings, C.E. *et al.*, *Biochem. J.*, 1959, **72**, 332; 1961, **78**, 412 (struct, biosynth)

Harris, S.A. *et al.*, *J. Med. Chem.*, 1965, **8**, 478 (synth)

Yuki, H. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 1107; 1120 (synth, props)

Mikami, Y. *et al.*, *Agric. Biol. Chem.*, 1971, **35**, 611

Meronuck, R.A. *et al.*, *Appl. Microbiol.*, 1972, **23**, 613 (isol)

Gatenbeck, S. *et al.*, *Acta Chem. Scand.*, 1973, **27**, 1825 (biosynth)

Dippenaar, A. *et al.*, *J. Cryst. Mol. Struct.*, 1977, **7**, 189 (cryst struct)

Kono, Y. *et al.*, *Agric. Biol. Chem.*, 1986, **50**, 2401 (isol, pmr, ms)


Fujita, M. *et al.*, *Tetrahedron*, 2001, **57**, 1229-1234 (activity)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 488

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ADC250; VTA750

Tenucyclamide A**T-75**

[213539-46-9, 213539-45-8]



R¹ = R² = CH₃

Absolute Configuration

C₁₉H₂₀N₆O₄S₂ 460.537

Alkaloidal peptide from *Nostoc spongiaeforme* var. *tenu*. Amorph. solid. [α]_D²⁵ -8.8 (c, 2.5 in MeOH). λ_{max} 243 (ε 24500) (MeOH).

12-Epimer: **Tenucyclamide B**

C₁₉H₂₀N₆O₄S₂ 460.537

Alkaloidal peptide from *Nostoc spongiaeforme* var. *tenu*. Glassy solid. [α]_D²⁵ -36 (c, 0.5 in MeOH). CAS no. incorrectly assigned by CA. λ_{max} 244 (ε 16500) (MeOH).

Banker, R. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1248-1251 (isol, uv, ir, pmr, cmr, ms)

You, S.-L. *et al.*, *Org. Lett.*, 2004, **6**, 2627-2630 (synth, abs config)

Tenucyclamide C**T-76**

[213539-46-9]

As Tenucyclamide A, T-75 with

R¹ = -CH₂⁴CH₂SMe, R² = H

C₂₀H₂₂N₆O₄S₃ 506.63

Alkaloidal peptide from *Nostoc spongiaeforme* var. *tenu*. Glassy solid. [α]_D²⁵

+12 (c, 0.5 in MeOH). λ_{max} 244 (ε 17300) (MeOH).

14-S-Oxide: **Tenucyclamide D**

[213539-47-0]

C₂₀H₂₂N₆O₅S₃ 522.629


Alkaloidal peptide from *Nostoc spongiaeforme* var. *tenu*. Glassy solid. [α]_D²⁵ +44 (c, 0.5 in MeOH). λ_{max} 247 (ε 39170) (MeOH).

Banker, R. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1248-1251 (isol, uv, ir, pmr, cmr, ms)

You, S.-L. *et al.*, *Org. Lett.*, 2004, **6**, 2627-2630 (synth, abs config)

Tenucausine**T-77**

[119212-24-7]



C₄₁H₄₆N₄O₃ 642.839

Alkaloid from the stem bark of *Melodinus tenuicaudatus* (Apocynaceae). Mp 160°. [α]_D -27 (CHCl₃).

O-De-Me: **De-O-methyltenucausine**

[221640-45-5]

C₄₀H₄₄N₄O₃ 628.813

Alkaloid from *Melodinus hemsleyanus*. Component of Chuan Shan Cheng. Amorph. powder. Mp 190° dec. [α]_D¹³ -198.5 (c, 0.05 in CHCl₃). λ_{max} 201 (log ε 4.59); 228 (sh) (log ε 4.53); 260 (log ε 4.07); 290 (log ε 4.06); 330 (log ε 4.14) (EtOH).

Zhou, Y.L. *et al.*, *Planta Med.*, 1988, **54**, 315 (isol, uv, ir, pmr, cmr, ms, struct)

Yan, K. *et al.*, *Yaoxue Xuebao*, 1998, **33**, 597-599 (Demethyltenucausine)

Tenuidine†**T-78**

[11017-85-9]


C₂₁H₃₁N₃O₅ 405.493

Struct. unknown. Alkaloid from *Polygala tenuifolia* (Polygalaceae). Mp 256°. [α]_D^{18.5} +1200 (EtOH).

Kim, J.H. *et al.*, *Yakhak Hoeji*, 1964, **8**, 59-61; *CA*, **65**, 12248c



Tenuilobine**T-79**

N¹-(4-Aminobutyl)-N¹,N¹⁶-bis(3-aminopropyl)-N¹⁶-[4-[(3-aminopropyl)amino]butyl]hexadecanediamide [173429-85-1]



C₃₃H₇₁N₇O₂ 597.97Alkaloid from leaves of *Oncinotis tenuiloba*. Oil (as 5HCl).Doll, M.K.-H. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 541 (*synth, cmr*)Doll, M.K.-H. *et al.*, *Heterocycles*, 1996, **42**, 319 (*isol, ir, pmr, ms, struct*)Vassiss, S. *et al.*, *Tet. Lett.*, 2002, **43**, 2597-2600 (*synth*)**Tenuiphylline**

[197234-71-2]

C₄₂H₄₄N₄O₇ 716.832Alkaloid from the leaves of *Kopsia tenuis* (Apocynaceae). Amorph. [α]_D -22 (c, 0.05 in CHCl₃). λ_{\max} 202 (log ϵ 4.44); 212 (log ϵ 4.33); 249 (log ϵ 3.99); 301 (log ϵ 3.54); 318 (log ϵ 3.5) (EtOH).Kam, T.-S. *et al.*, *Tetrahedron*, 1997, **53**, 12661-12670 (*isol, uv, ir, pmr, cmr, ms*)**Tenuipine****T-81**C₃₈H₄₀N₂O₇ 636.743**(1R,1'R)-form**Alkaloid from the bark of *Daphnandra dielsii* and *Daphnandra tenuipes* (Monimiaceae). Prisms (MeOH or Me₂CO/MeOH). Mp 140-145°. [α]_D²⁰ -258 (c, 2.8 in CHCl₃).**N²,N^{2'}-Di-Me:**C₄₀H₄₆N₂O₇²⁺ 666.813Prisms + 3H₂O (50% EtOH aq.) (as diiodide). Mp 267-272° dec. (diiodide). [α]_D¹⁵ -165 (c, 1.0 in EtOH aq.).**O⁷-De-Me: (-)-Nortenuipine**C₃₇H₃₈N₂O₇ 622.716Alkaloid from the leaves of *Daphnandra tenuipes* (Monimiaceae). Needles (95% EtOH or CHCl₃/MeOH). Mp 211° dec. [α]_D²² -218 (c, 0.8 in CHCl₃). At first erroneously named *N*-Demethyltenuipine.**O⁷-De-Me, N²- β -oxide: Nortenuipine 2'- β -N-oxide**

[126624-25-7]

C₃₇H₃₈N₂O₈ 638.716Alkaloid from the bark of *Daphnandra dielsii* (Monimiaceae). Amorph. [α]_D²⁷ -71 (c, 0.24 in CHCl₃).**O⁷-De-Me, N²,N^{2'}-di-Me:**C₃₉H₄₄N₂O₇²⁺ 652.786Mp 261-265° dec. (as diiodide). [α]_D¹⁵ -154 (c, 0.4 in EtOH aq.).**(1R,1'S)-form****Isotenuipine**

[35306-97-9]

Alkaloid from the bark of a *Daphnandra* sp. of unknown origin (Monimiaceae).Mp 239-241°. [α]_D¹³ +129 (c, 1.2 in CHCl₃).**N²,N^{2'}-Di-Me:**Prisms + 3H₂O (30% EtOH aq.) (as diiodide). Mp 278° dec. (diiodide).[α]_D¹⁵ -50 (c, 0.48 in 50% EtOH aq.).**(1S,1'S)-form [1263-97-4]**Alkaloid from the leaves and terminal twigs of an unnamed *Daphnandra* sp. and from the bark of *Daphnandra tenuipes* (Monimiaceae). [α]_D²⁰ +224 (CHCl₃).**O⁷-De-Me: (+)-Nortenuipine**

[36067-01-3]

Alkaloid from the bark of *Daphnandra* sp. Dt-7 and *Daphnandra tenuipes*, and from the leaves, stems and bark of *Daphnandra johnsonii* (Monimiaceae). Mp 260°. [α]_D²⁰ +236.3 (CHCl₃).**(2RS,2'RS)-form****Repandinine**


[6883-11-0]

Alkaloid of the bark of *Daphnandra dielsii*, *Daphnandra repandula* and *Daphnandra tenuipes*, and from the leaves, stems and bark of *Daphnandra johnsonii* (Monimiaceae). Fine needles (Me₂CO/MeOH). Mp 243°.**N²,N^{2'}-Di-Me:**Needles + 2H₂O (H₂O) (as diiodide).


Mp 275° (diiodide).

Bick, I.R.C. *et al.*, *J.C.S.*, 1953, 695 (*isol, struct*)Bick, I.R.C. *et al.*, *CA*, 1964, **60**, 1811e (*isol*)Battersby, A.R. *et al.*, *J.C.S.*, 1965, 2239 (*ord*)Bick, I.R.C. *et al.*, *J.C.S. (C)*, 1971, 3779(Tenuipine, Isotenuipine, *config, pmr*)Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592 (*ms*)Bick, I.R.C. *et al.*, *Aust. J. Chem.*, 1978, **31**,2539 (*isol*)Koike, L. *et al.*, *Tet. Lett.*, 1979, 3765 (*cmr*)Galal, A.M. *et al.*, *Heterocycles*, 1989, **29**, 1689 (*Nortenuipine 2'- β -N-oxide*)**Terengganensine A****T-82**



[188549-37-3]



C₁₉H₂₂N₂O₃ 326.394Alkaloid from bark of *Kopsia terengganensis*. [α]_D -25. λ_{\max} 208 (log ϵ 4.24); 236 (log ϵ 3.89); 284 (log ϵ 3.34) (No solvent reported).Uzir, S. *et al.*, *Tet. Lett.*, 1997, **38**, 1571 (*isol, uv, pmr, cmr, struct*)**Terengganensine B****T-83**

[188549-41-9]


C₁₉H₂₂N₂O₂ 310.395Alkaloid from bark of *Kopsia terengganensis*. [α]_D -19. λ_{\max} 228 (log ϵ 3.74); 283 (log ϵ 4.13); 307 (log ϵ 3.6) (No solvent reported).Uzir, S. *et al.*, *Tet. Lett.*, 1997, **38**, 1571 (*isol, uv, pmr, cmr, struct*)**Terezine A****T-84**

[165133-88-0]

C₁₅H₁₈N₂O₃ 274.319Exists predominantly in the hydroxy-pyrazine tautomeric form rather than in the more common pyrazinone (amide) form. Metab. of the fungus *Sporormiella teretispora*. Antifungal agent. Yellow oil. [α]_D -20.2 (c, 0.21 in MeOH). λ_{\max} 204 (ϵ 15000); 230 (ϵ 8000); 325 (ϵ 7100) (MeOH) (Berdy).Wang, Y. *et al.*, *J. Nat. Prod.*, 1995, **58**, 93 (*isol, uv, ir, pmr, cmr, ms, struct*)**Terezine B****T-85****3,6-Dihydro-6-hydroxy-6-(hydroxyphenylmethyl)-3,5-dimethoxy-3-(1-methyl-ethyl)-2-(1H)-pyrazinone, 9CI**
[165133-89-1]C₁₆H₂₂N₂O₅ 322.36Abs. *config.* shown is tentative. Metab. of the fungus *Sporormiella teretispora*. Antifungal agent. Yellow oil. [α]_D -14.4 (c, 0.083 in MeOH). λ_{\max} 204 (ϵ 11000) (MeOH) (Berdy).


3-Epimer: Terezine C[165306-63-8]
C₁₆H₂₂N₂O₅ 322.36From *Sporormiella teretispora*. Antifungal agent. Yellow oil. [α]_D -63.1 (c, 0.59 in MeOH). λ_{max} 205 (ε 13000) (MeOH) (Berdy).Wang, Y. et al., *J. Nat. Prod.*, 1995, **58**, 93 (*isol, uv, ir, pmr, cmr, ms, struct*)**Terezine D****3-Methyl-6-[[7-(3-methyl-2-butenyl)-1H-indol-3-yl]methyl]-2,5-piperazine-dione, 9CI**
[165133-90-4]C₁₉H₂₃N₃O₂ 325.41Metab. of the fungus *Sporormiella teretispora*. Antifungal agent. Powder. Mp 192-194°. [α]_D +7 (c, 0.58 in MeOH). λ_{max} 204 (ε 24000); 222 (ε 24000); 279 (ε 3800) (MeOH) (Berdy).Wang, Y. et al., *J. Nat. Prod.*, 1995, **58**, 93 (*isol, uv, ir, pmr, cmr, ms, struct*)**[3,2':2'(3'H),3''-Ter-1H-indol]-3'-one, 9CI****2,2-Di-3-indolyl-3-indolone. 2,2-Bis(3-indolyl)indoxyl**
[17646-95-6]C₂₄H₁₇N₃O 363.418Metab. from the marine bacterium *Vibrio parahaemolyticus* isol. from the toxic mucus of the boxfish *Ostracion cubicus*. Also isol. as the product of indole oxidation by a strain of *Claviceps purpurea*. Exhibits antibacterial activity. Oil. λ_{max} 242 (ε 33600); 268 (ε 31800) (CHCl₃) (Berdy).Loo, Y.H. et al., *Chem. Ind. (London)*, 1957, 1123Bell, R. et al., *J. Nat. Prod.*, 1994, **57**, 1587 (*isol, uv, ir, pmr, cmr*)Stull, T.L. et al., *J. Biol. Chem.*, 1995, **270**, 5**[3,3':3'(2'H),3''-Ter-1H-indol]-2'-one**

T-88


3,3-Di-3-indolyl-2-indolone. TrisindolineC₂₄H₁₇N₃O 363.418Prod. by a bacterium of *Vibrio* sp. separated from the Okinawan marine sponge *Hyrtios altum* and *Vibrio parahaemolyticus* Bio249. Amorph. solid. Incorrectly named in CA; no CAS no. reported. λ_{max} 219 (ε 62000); 254 (ε 12000); 274 (ε 12000); 280 (ε 12000); 290 (ε 10000) (MeOH).Kobayashi, M. et al., *Chem. Pharm. Bull.*, 1994, **42**, 2449-2451 (*isol, synth, uv, ir, pmr, cmr*)Veluri, R. et al., *J. Nat. Prod.*, 2003, **66**, 1520-1523 (*isol*)**Termine**

T-89

[171090-90-7]

C₁₅H₂₂N₂O₂ 262.351Alkaloid from seeds of *Lupinus termis*.Mohamed, M.H. et al., *CA*, 1996, **124**, 4886v (*isol*)**Termisine**

T-90




Relative Configuration

C₁₆H₂₈N₂O₃ 296.409**(±)-form** [152606-58-1]Alkaloid from seeds of *Lupinus termis* (Fabaceae). Fine yellow cryst. Mp 98-99°. Zwitterionic.**(ξ)-form****14-Aldehyde, 10α-hydroxy: 14-Dehydro-10-hydroxytermisine**
[220672-12-8]C₁₆H₂₆N₂O₄ 310.392Alkaloid from *Lupinus albus*. Opt. active.Mohamed, M.H. et al., *J. Nat. Prod.*, 1993, **56**, 1999 (*isol, ir, pmr, cmr, ms, struct*)Mohamed, M.H. et al., *CA*, 1999, **130**, 179920c (*14-Dehydro-10-hydroxytermisine*)**Terpendole I**


T-91

[167612-17-1]

C₂₇H₃₅NO₅ 453.577Prod. by *Albophoma yamanashiensis*.Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, EtOH, C₆H₆, MeCN, EtOAc; poorly sol. H₂O. [α]_D²⁸ +7 (c, 1 in MeOH). λ_{max} 228 (ε 47400); 280 (ε 11200) (MeOH) (Berdy).**17-O-(3-Methyl-2-butenyl): Terpendole J**
[167427-26-1]C₃₂H₄₃NO₅ 521.695Prod. by *Albophoma yamanashiensis*.Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, EtOH, C₆H₆, MeCN, EtOAc; poorly sol. H₂O. Mp 248-250° dec. [α]_D²⁸ -30.3 (CHCl₃). λ_{max} 238 (ε 13100); 282 (ε 9400) (MeOH) (Berdy).Tomoda, H. et al., *J. Antibiot.*, 1995, **48**, 793 (*isol, pmr, cmr, struct*)**Terpendole B**

T-92

[156967-67-8]


C₂₇H₃₅NO₃ 421.578Full stereochem. not determined. Probable stereochem. shown here by analogy with related compounds. CAS numbering shown. Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, EtOAc, EtOH, C₆H₆, MeCN; poorly sol. H₂O. [α]_D²⁸ -3.6 (c, 1 in MeOH). λ_{max} 225; 275 (MeOH) (Derep).**3α-Hydroxy, 1'-O-(3-methyl-2-butenyl):****Terpendole D**

[156967-66-7]

C₃₂H₄₃NO₄ 505.696Prod. by *Albophoma yamanashiensis*.Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, MeCN, C₆H₆, EtOH, EtOAc; poorly sol. H₂O. [α]_D²⁸ -31.2 (c, 1 in MeOH). λ_{max} 225; 275 (MeOH) (Derep).Huang, X.-H. et al., *J. Antibiot.*, 1995, **48**, 1; 5 (*isol, pmr, cmr, struct*)

Terpendole C

[156967-65-6]

C₃₂H₄₁NO₅ 519.68

Prod. by *Albophoma yamanashiensis* FO 2546. Acyl-CoA:cholesterol acyl transferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆, EtOAc, EtOH, MeCN; poorly sol. H₂O. $[\alpha]_D^{28}$ -2.3 (c, 1 in MeOH). λ_{\max} 225 ; 275 (MeOH) (Derep).

33,34-Epoxyde: Terpendole A

[156967-64-5]

C₃₂H₄₁NO₆ 535.679

Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyl transferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆, MeCN, EtOAc, EtOH; poorly sol. H₂O. $[\alpha]_D^{28}$ +11.6 (c, 1 in MeOH). Only gross struct. is known. λ_{\max} 225 ; 275 (MeOH) (Derep). λ_{\max} 226 ; 282 (MeOH) (Berdy).

14 α -Hydroxy: Terpendole M

[222400-32-0]

C₃₂H₄₁NO₆ 535.679

Isol. from *Lolium perenne* infected with *Neotyphodium lolii*. Amorph. solid (MeCN).

6,7-Didehydro: Terpendole K

[167427-27-2]

C₃₂H₃₉NO₅ 517.664

Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, MeCN, EtOH, EtOAc, C₆H₆; poorly sol. H₂O. $[\alpha]_D^{28}$ -21.8 (c, 1 in MeOH). λ_{\max} 228 (ϵ 38600); 280 (ϵ 11200) (MeOH) (Berdy).


Huang, X.-H. *et al.*, *J. Antibiot.*, 1995, **48**, 1-4; 5-11 (*isol, pmr, cmr, struct*)

Tomoda, H. *et al.*, *J. Antibiot.*, 1995, **48**, 793-804 (*Terpendole K*)

Gatenby, W.A. *et al.*, *J. Agric. Food Chem.*, 1999, **47**, 1092-1097 (*Terpendole M*)

Terpendole H

[156967-69-0]


C₂₇H₃₃NO₅ 451.561

Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆; poorly sol. H₂O. $[\alpha]_D^{28}$ -47 (c, 1 in MeOH). λ_{\max} 227 (ϵ 35400); 279 (ϵ 7900) (MeOH) (Berdy).

Tomoda, H. *et al.*, *J. Antibiot.*, 1995, **48**, 793 (*isol, pmr, cmr, struct*)

T-93**Terpendole L**

[167612-18-2]

C₃₇H₄₉NO₅ 587.798


Prod. by *Albophoma yamanashiensis*. Acyl-CoA:cholesterol acyltransferase (ACAT) inhibitor. Powder. Sol. MeOH, C₆H₆; poorly sol. H₂O. Mp 148-150°. $[\alpha]_D^{28}$ +16.5 (c, 1 in MeOH). λ_{\max} 230 (ϵ 32500); 283 (ϵ 9500) (MeOH) (Berdy).

Tomoda, H. *et al.*, *J. Antibiot.*, 1995, **48**, 793 (*isol, pmr, cmr, struct*)

Terpeptin*Antibiotic RK-F 1010. RK-F 1010*

[188824-81-9]

[203200-22-0]

C₂₈H₄₀N₄O₃ 480.649

Peptide antibiotic. Prod. by *Aspergillus terreus* 95F-1. Mammalian cell cycle inhibitor. Yellow solid. Mp 92-95°. $[\alpha]_D$ -135.2 (c, 0.1 in CHCl₃). λ_{\max} 230 (ϵ 87000); 285 (ϵ 6700) (no solvent reported). λ_{\max} 230 (ϵ 87000); 280 (ϵ 6700) (MeOH) (Berdy).

Kagamizono, T. *et al.*, *Tet. Lett.*, 1997, **38**,

1223-1226 (*isol, uv, ir, pmr, cmr*)

Japan. Pat., 1998, 98 17 593; *CA*, **128**, 179450h (*isol*)


Su, S. *et al.*, *Tetrahedron*, 2003, **59**, 8931-8946 (*synth, pmr, cmr*)

T-95

Quang, D.N. *et al.*, *Phytochemistry*, 2003, **62**, 109-113; **63**, 919-924; 2004, **65**, 1179-1184 (*Thelephantins*)

3,2':4',3''-Terpyridine, 9CI, 8CI*Nicotelline*

[494-04-2]

C₁₅H₁₁N₃ 233.272

Alkaloid from tobacco (*Nicotiana tabacum*) (Solanaceae). Needles (petrol). Mp 147-148°. Originally considered to be C₁₀H₈N₂. Kuffner *et al.* (1956) initially proposed an isomeric struct.

Picrate: Mp 216-217°.

l-N-Oxide: [172424-37-2]

Mp 200° (min.).


Pictet, A. *et al.*, *Ber.*, 1901, **34**, 696 (*isol*)

Thesing, J. *et al.*, *Chem. Ber.*, 1957, **90**, 711 (*synth*)

Yamamoto, Y. *et al.*, *Synthesis*, 1986, 564 (*synth*)

Zoltewicz, J.A. *et al.*, *Tetrahedron*, 1995, **51**, 11393 (*synth, N-oxide, pmr*)

Yamamoto, Y. *et al.*, *Heterocycles*, 1996, **42**, 189 (*synth, ir, pmr*)


Terragine A**T-99**C₂₆H₃₈N₄O₇ 518.609

Prod. by *Streptomyces lividans* recombinant 436-s4-5b1. Siderophore.

Wang, G.Y.S. *et al.*, *Org. Lett.*, 2000, **2**, 2401-2404 (*isol, pmr, cmr*)

Terragine B**T-100**


N-[5-(2,5-Dioxo-1-pyrrolidinyl)pentyl]-*N*-hydroxybenzeneacetamide, 9CI [292862-75-0]

C₁₇H₂₂N₂O₄ 318.372

Prod. by *Streptomyces lividans* recombinant 436-s4-5b1. Siderophore.

T-94**[*p*-Terphenyl]-2',3',4,4'',5',6'-hexol****T-97**

*3,6-Bis(4-hydroxyphenyl)-1,2,4,5-benzenetetrol, 8CI. 2',3',4,4'',5',6'-Hexahydroxy-*p*-terphenyl. Leucoatromentin*

C₁₈H₁₄O₆ 326.305

2'-(3-Pyridinecarbonyl), 5'-benzoyl: Thelephantin N

[719279-84-2]

C₃₁H₂₁NO₈ 535.509

Isol. from the mushroom *Hydnellum caeruleum*. Greyish solid. λ_{\max} 227 (log ϵ 4.2); 264 (log ϵ 4.2) (MeOH).

Wang, G.Y.S. *et al.*, *Org. Lett.*, 2000, **2**, 2401-2404 (*isol*, *pmr*, *cmr*)

Terrigine C T-101

PhCH₂CON(OH)(CH₂)₅NHAc

C₁₅H₂₂N₂O₃ 278.35

Prod. by *Streptomyces lividans* recombinant 436-s4-5b1. Siderophore.

Wang, G.-Y.-S. *et al.*, *Org. Lett.*, 2000, **2**, 2401-2404 (*isol*, *pmr*, *cmr*)


Terrigine D T-102

As Terrigine A, T-99 with R = NHAc

C₂₄H₃₈N₄O₆ 478.587

Prod. by *Streptomyces lividans* recombinant 436-s4-5b1. Siderophore.

Wang, G.Y.S. *et al.*, *Org. Lett.*, 2000, **2**, 2401-2404 (*isol*, *pmr*, *cmr*)

Terresoxazine T-103


C₁₁H₁₁NO₃ 205.213

Alkaloid from the fruit of *Tribulus terrestris*. Prisms. Mp 194-196°. [α]_D²⁰ +0.7 (c, 0.38 in Me₂CO).

Huang, J.W. *et al.*, *Chin. Chem. Lett.*, 2004, **15**, 305-306 (*isol*, *pmr*, *cmr*)

Terrestrial acid† T-104

6-Amino-1,2,3,4-tetrahydro-4-oxo-1,3,5-triazine-2-carboxylic acid, 9CI [912850-67-0]




C₄H₆N₄O₃ 158.116

Constit. of *Tribulus terrestris*.

Chen, H. *et al.*, *CA*, 2006, **145**, 434756 (*isol*)

Terreusinone T-105

2,6-Bis(1-hydroxy-2-methylpropyl)-1H,5H-pyrrolo[2,3-b]indole-4,8-dione [637334-34-0]




Absolute Configuration

C₁₈H₂₂N₂O₄ 330.383

Prod. by the marine fungus *Aspergillus terreus* (MFA-460). Potent Uv-A protectant. Yellowish solid. Mp 230° dec. [α]_D +47 (c, 0.3 in MeOH). λ_{max} 204 (log ε 3.73); 247 (log ε 4.08); 277 (log ε 3.83);

355 (log ε 3.57) (MeOH).

Lee, S.M. *et al.*, *Tet. Lett.*, 2003, **44**, 7707-7710 (*isol*, *cd*, *pmr*, *cmr*, *ms*)

Terriquinone A T-106

C₃₂H₃₀N₂O₃ 490.601

Prod. by *Aspergillus terreus* found in *Ambrosia ambrosioides*. Cytotoxic. Purple powder. Mp 160-165°. λ_{max} 223 (log ε 5.63); 274 (log ε 5.18); 360 (log ε 4.24) (MeOH).

He, J. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1985-1991 (*isol*, *pmr*, *cmr*, *ms*)

4,8,12,16-Tetraazanonadecane-1,19-diamine, 9CI T-107

1,5,9,13,17,21-Hexaazaheneicosane. *Cal-dohexamine*

[63833-74-9]

[87265-97-2]

H₂N(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH₂

C₁₅H₃₈N₆ 302.505


Prod. by *Thermus thermophilus* and other thermophilic bacteria. Powder (as hexahydrochloride). Hexahydrochloride melts >250°.

Dietrich, B. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 1262-1278 (*synth*, *pmr*, *cmr*)

Hamana, K. *et al.*, *FEMS Microbiol. Lett.*, 1990, **68**, 31-34 (*isol*)

2,2',5,5'-Tetrabromo-3,3'-bi-1H-indole T-108

3,3'-Bi(2,5-dibromoindole). *Rivularin C* [81387-82-8]



C₁₆H₈Br₄N₂ 547.869


Isol. from the marine blue-green alga *Rivularia firma*. Antiinflammatory agent. Rosettes (CHCl₃). Mp 239-240°. λ_{max} 209 (ε 61400); 228 (ε 57400); 289 (ε 17600); 297 (ε 15500) (MeCN).

Norton, R.S. *et al.*, *J.A.C.S.*, 1982, **104**, 3628-3635 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)

Maehr, H. *et al.*, *J.O.C.*, 1984, **49**, 1549-1553 (*nomenclature*)

2,2',6,6'-Tetrabromo-3,3'-bi-1H-indole, 9CI T-109

3,3'-Bi(2,6-dibromoindole) [138779-89-2]




C₁₆H₈Br₄N₂ 547.869

Isol. from the cyanobacterium *Rivularia firma*. Viscous oil.

Hodder, A.R. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1661-1663 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

4,4',6,6'-Tetrabromo-2,2'-bis(methylthio)-3,3'-bi-1H-indole, 9CI T-110

3,3'-Bi[4,6-dibromo-2-methylthio-1H-indole] [128351-82-6]



C₁₈H₁₂Br₄N₂S₂ 640.054

Alkaloid from the Okinawan red alga *Laurencia brongniartii*. Cryst. (hexane/CHCl₃). Mp 186-188°. λ_{max} 236 (ε 50000); 306 (ε 20000) (EtOH).

2-S-Oxide: 4,4',6,6'-Tetrabromo-2-(methylsulfinyl)-2'-(methylthio)-3,3'-bi-1H-indole

C₁₈H₁₂Br₄N₂O₂S₂ 656.054

Isol. from *Laurencia brongniartii*. Mp 187-189°. [α]_D²⁵ +36 (c, 0.2 in CHCl₃). λ_{max} 238 (log ε 4.7); 312 (log ε 4.1) (MeOH).

2,2'-Di-S-oxide: 4,4',6,6'-Tetrabromo-2,2'-bis(methylsulfinyl)-3,3'-bi-1H-indole

C₁₈H₁₂Br₄N₂O₂S₂ 672.053

Isol. from *Laurencia brongniartii*. Mp 191-193°. λ_{max} 233 (log ε 4.5); 309 (log ε 3.08) (MeOH).

2-De(methylthio), 2-bromo: 2,4,4',6,6'-Pentabromo-2'-(methylthio)-3,3'-bi-1H-indole

[873202-17-6]

C₁₇H₉Br₅N₂S 672.858

Alkaloid from *Laurencia brongniartii*. λ_{max} 212 (sh) (ε 28390); 234 (ε 40925); 296 (ε 11930) (MeOH).

Tanaka, J. *et al.*, *Tetrahedron*, 1989, **45**, 7301-7310 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *cryst struct*)


Kubota, N.K. *et al.*, *Heterocycles*, 2005, **65**, 2675-2682 (*demethylthio-2-bromo*)

El-Gamal, A.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 815-817 (*oxides*)

3,3',4,4'-Tetrabromo-5,5'-dichloro-1,1'-dimethyl-2,2'-bi-1H-pyrrole


T-111

[253798-64-0]

C₁₀H₆Br₄Cl₂N₂ 544.692Isol. from the eggs of various seabirds. Cryst. (Me₂CO). Mp 221° dec.Gribble, G.W. *et al.*, *Chem. Comm.*, 1999, 2195-2196 (*synth*, *pmr*, *ms*, *cryst struct*)**5,5',7,7'-Tetrabromo-6,6'-dimethoxyindigotin**


T-112

[58933-45-2]

C₁₈H₁₀Br₄N₂O₄ 637.904Pigment from the marine invertebrate *Ptychodera flava laysanica*. Purple-blue powder. Mp 300°.Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 227-230 (*isol*, *struct*)**2,3,4,6-Tetrabromo-1H-indole, 9CI**

T-113

[128351-87-1]

C₈H₃Br₄N 432.734Alkaloid from the Okinawan red alga *Laurencia brongniartii*. Cryst. (hexane/CCl₄). Mp 137-141°.

N-Ac: [128351-91-7]

C₁₀H₅Br₄NO 474.772Cryst. (hexane/CCl₄). Mp 176-178°.

N-Me: 2,3,4,6-Tetrabromo-1-methyl-1H-indole

[957475-92-2]

C₉H₅Br₄N 446.761Alkaloid from *Laurencia decumbens*. Cryst.Tanaka, J. *et al.*, *Tetrahedron*, 1989, **45**, 7301-7310 (*isol*, *ir*, *pmr*, *ms*)Ji, N.-Y. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 1731-1736 (*N-Me*)**2,3,4,7-Tetrabromo-1H-indole, 9CI**

T-114

[68234-23-1]

C₈H₃Br₄N 432.734Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 144.5°.Brennan, M.R. *et al.*, *Tet. Lett.*, 1978, **19**,1637-1640 (*isol*, *pmr*, *struct*)Ohta, T. *et al.*, *Heterocycles*, 1989, **29**, 1663-1667 (*synth*)**2,3,5,6-Tetrabromo-1H-indole, 9CI**

T-115

[17826-06-1]


C₈H₃Br₄N 432.734Alkaloid from the algae *Laurencia brongniartii* and *Laurencia similis*. Mp 152.5-154° (149-151°). λ_{max} 230 (ε 47000); 294 (ε 9600); 301 (ε 9500) (EtOH) (Derep).

N-Me: 2,3,5,6-Tetrabromo-1-methyl-1H-indole

[25055-55-4]

C₉H₅Br₄N 446.761Alkaloid from *Laurencia brongniartii*. Cryst. (petrol). Mp 171.5-172° (168-170°). λ_{max} 233 (ε 49000); 296 (ε 10000); 303 (ε 10000) (EtOH) (Derep).Settimo, A.D. *et al.*, *Gazz. Chim. Ital.*, 1967, **97**, 1304-1316; 1977, **107**, 367-372 (*synth*)Settimo, A.D. *et al.*, *J.O.C.*, 1970, **35**, 2546-2551 (*N-Me*, *synth*)Carter, G.T. *et al.*, *Tet. Lett.*, 1978, 4479-4482 (*N-Me*, *isol*, *uv*, *pmr*, *ms*)Liu, Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 748-749 (*N-Me*, *synth*)Suárez-Castillo, O.R. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1596-1600 (*synth*)**2,3',5,5'-Tetrabromo-7'-methoxy-3,4'-bi-1H-indole Rivularin D₃**

T-116

C₁₇H₁₀Br₄N₂O 577.895**(+)-form** [81387-83-9]Isol. from the marine cyanobacterium *Rivularia firma*. Shows antiinflammatory, antiallergic, CNS depressant and tremogenic props. Prisms (CH₂Cl₂/hexane). Sol. MeOH, CHCl₃. Mp 178-179° dec. [α]_D²⁰ +71 (c, 1 in CHCl₃). Exhibits opt. activity owing to restricted rotn. λ_{max} 227 (ε 66000); 280 (ε 16000); 290 (ε 17000); 299 (ε 14000) (MeCN) (Berdy).

I-N-Ac:

Prisms (CH₂Cl₂/hexane). Mp 109-111°. [α]_D²⁰ +99 (c, 1 in CHCl₃).

Di-N-Ac:

Prisms (CH₂Cl₂/hexane). Mp 96-98°. [α]_D²⁰ +71 (c, 0.75 in CHCl₃).2-Debromo: 3',5,5'-Tribromo-7'-methoxy-3,4'-bi-1H-indole. **Rivularin D₁**

[81387-84-0]

C₁₇H₁₁Br₃N₂O 498.999Isol. from *Rivularia firma*. Antiinflammatory agent. Prisms (CHCl₃). Mp 220-223°. [α]_D²⁰ +8.5 (c, 1 in CHCl₃). Exhibits opt. activity owing to restricted rotn. λ_{max} 283 ; 288 ; 297 (MeOH) (Berdy). λ_{max} 225 (ε 73000);

283 (ε 12700); 288 (ε 13000); 297 (ε 11500) (MeCN).

3'-Debromo: 2,5,5'-Tribromo-7'-methoxy-3,4'-bi-1H-indole. **Rivularin D₂**

[81387-85-1]

C₁₇H₁₁Br₃N₂O 498.999Isol. from the marine blue-green alga *Rivularia firma*. Foam. [α]_D²⁰ +11.3 (c, 1 in CHCl₃). Exhibits opt. activity owing to restricted rotn. λ_{max} 227 (ε 69000); 279 (ε 17000); 289 (ε 16000); 298 (ε 13000) (MeCN).**(±)-form** [95739-89-2]Synthetic. Beige prisms (CH₂Cl₂/hexane). Mp 180°.

2-Debromo: [95739-88-1]

Synthetic. Cryst. (CHCl₃). Mp 220-223° dec.Norton, R.S. *et al.*, *J.A.C.S.*, 1982, **104**, 3628-3635 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)Maehr, H. *et al.*, *J.A.C.S.*, 1985, **107**, 2943-2945 (*synth*, *uv*, *ir*, *pmr*, *ms*)**3,3',5,5'-Tetrabromo-7'-methoxy-1,4'-bi-1H-indole Rivularin B**


T-117

[81387-86-2]

C₁₇H₁₀Br₄N₂O 577.895Isol. from the marine blue-green alga *Rivularia firma*. Poorly cryst. solid (CHCl₃/hexane). Mp 196-200°. [α]_D²⁰ -6 (c, 1 in MeCN). Exhibits opt. activity owing to restricted rotn.Norton, R.S. *et al.*, *J.A.C.S.*, 1982, **104**, 3628-3635 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *struct*)Maehr, H. *et al.*, *J.O.C.*, 1984, **49**, 1549-1553 (*nomenclature*)**2,3,4,5-Tetrabromo-1H-pyrrole, 9CI**

T-118

[54705-14-5]

C₄HBr₄N 382.675Metab. of marine *Chromobacterium* sp. Has antibiotic activity. Spar. sol. H₂O. Mp 250°. Unstable, v. light-sensitive.

N-Me: [56454-29-6]


C₅H₃Br₄N 396.701

Mp 154-155°. Blue melt.

De Varda, G. *et al.*, *Ber.*, 1888, **21**, 2871 (*synth*)Andersen, R.J. *et al.*, *Mar. Biol. (Berlin)*, 1974, **27**, 281 (*isol*, *ms*, *struct*)Gilow, H.M. *et al.*, *J.O.C.*, 1981, **46**, 2221 (*synth*, *glc*)**2,3,4,7-Tetrachloro-1H-indole, 9CI**

T-119

[68234-17-3]

C₈H₃Cl₄N 254.929

Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 119°.

Brennan, M.R. *et al.*, *Tet. Lett.*, 1978, **19**, 1637-1640 (*isol. struct*)

Ohta, T. *et al.*, *Heterocycles*, 1989, **29**, 1663-1667 (*synth*)

15-Tetracosenoic acid T-120
[26444-06-4]

$\text{H}_3\text{C}(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_{13}\text{COOH}$

$\text{C}_{24}\text{H}_{46}\text{O}_2$ 366.626

(E)-form [14490-79-0]

Cryst. (EtOH). Mp 66-67° (61°).

Me ester:

$\text{C}_{25}\text{H}_{48}\text{O}_2$ 380.653

Cryst. (Me_2CO). Mp 36.5-37.5°.

Bp_{0.00001} 180-190°. CAS no. not found to CA 130.

Amide:

$\text{C}_{24}\text{H}_{47}\text{NO}$ 365.641

Mp 97-98°. CAS no. not found to CA 130.

(Z)-form

Nervonic acid. Selacholeic acid

[506-37-6]

[31152-46-2]

Present in cerebrosides and in fish oils.

Isol. from the sponge *Pseudaxinella cf. lunaecharta*. Also in seed oils of *Lunaria biennis*, rape (*Brassica napus*), *Carthamus tinctorius* and *Tropaeolum speciosum*.

Cryst. (EtOH). Mp 40.5-41° Mp 44-45°.

Me ester: [2733-88-2]

Mp 14-15°. Bp_{0.02} 165° approx.

Amide: [208650-37-7]

Cryst. (EtOH). Mp 86.5-87.5°.

Benzylamide: N-Benzyl-15-tetracosenamide

[847361-94-8]

$\text{C}_{31}\text{H}_{53}\text{NO}$ 455.766

Alkaloid from the tubers of *Lepidium meyenii* (maca). Powder. λ_{max} 212 (log ϵ 3.86) (MeOH).

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 784B (*nmr*)

Hale, J.B. *et al.*, *J.A.C.S.*, 1930, **52**, 4536 (*synth*)

Bounds, D.G. *et al.*, *J.C.S.*, 1954, 448 (*synth*)

Litchfield, C. *et al.*, *Lipids*, 1970, **5**, 144-146 (*isol*)

Batchelor, J.G. *et al.*, *J.A.C.S.*, 1973, **95**, 6358 (*cmr*)

Lecerf, J. *et al.*, *J. Physiol. (Paris)*, 1975, **70**, 493

Morales, R.W. *et al.*, *Biochim. Biophys. Acta*, 1976, **431**, 206 (*isol. biosynth*)

Bohannon, M.B. *et al.*, *Lipids*, 1976, **11**, 157-159 (*isol*)

Barnathan, G. *et al.*, *Lipids*, 1996, **31**, 193-200 (*isol. Pseudaxinella*)

Zhao, J.-P. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 690-693 (*benzylamide*)

2,4-Tetradecadiene-8,10-diy- noic acid T-121

$\text{H}_3\text{CCH}_2\text{CH}_2\text{C}\equiv\text{CC}\equiv\text{CCH}_2\text{CH}_2\text{CH}=\text{CHCH}=\text{CHCOOH}$

$\text{C}_{14}\text{H}_{16}\text{O}_2$ 216.279

(2E,4E)-form

Cryst. (petrol/Et₂O). Mp 154-155° (58°).

2-Methylpropylamide: see Anacycline, A-968

3-Methylbutylamide: N-(3-Methylbutyl)-2,4-tetradecadiene-8,10-diy- noic acid

[113235-96-4]

$\text{C}_{19}\text{H}_{27}\text{NO}$ 285.428

Isol. from *Achillea wilhelmsii*. Cryst. Mp 95-97°. λ_{max} 250 nm (Et₂O).

Pyrrolidide: 2,4-Tetradecadiene-8,10-diy- noic acid pyrrolidide. 1-(1-Oxo-2,4-tetradecadiene-8,10-diy- nyl)pyrrolidine

[94413-14-6]

$\text{C}_{18}\text{H}_{23}\text{NO}$ 269.386

Isol. from underground parts of *Achillea nana*. Cryst. Mp 88-89°. λ_{max} 255nm (Et₂O).

(2E,4E)-form

Piperidide: 2,4-Tetradecadien-8,10-diy- noic acid piperidide. 1-(1-Oxo-2,4-tetradecadiene-8,10-diy- nyl)piperidine, 9CI

[43110-70-9]

$\text{C}_{19}\text{H}_{25}\text{NO}$ 283.413

Alkamide from *Achillea sudetica* (Asteraceae). Not obt. completely pure.

Crombie, L. *et al.*, *J.C.S.*, 1955, 999; 1957, 2767 (*isol. struct. synth. ir. uv*)

Jente, R. *et al.*, *Chem. Ber.*, 1972, **105**, 1694 (*isol. struct. uv. ms*)

Bohlmann, F. *et al.*, *Chem. Ber.*, 1973, **106**, 1328 (*isol. ms. struct*)

Greger, H. *et al.*, *Phytochemistry*, 1984, **23**, 1503; 1989, **28**, 2363 (*isol. struct. pmr. ir. uv. ms*)

Greger, H. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1100 (*isopentylamide*)

Kuroopka, G. *et al.*, *Planta Med.*, 1987, **53**, 440 (*isol*)

Crombie, L. *et al.*, *Tet. Lett.*, 1987, **28**, 4875 (*synth*)

10,12-Tetradecadiene-4,6- diynoic acid, 9CI T-122

$\text{H}_3\text{CCH}=\text{CHCH}=\text{CHCH}_2\text{CH}_2\text{C}\equiv\text{CC}\equiv\text{CCH}_2\text{CH}_2\text{COOH}$

$\text{C}_{14}\text{H}_{16}\text{O}_2$ 216.279

(10E,12E)-form

2-Methylpropylamide: N-(2-Methylpropyl)-10,12-tetradecadiene-4,6-diy- namide

[125085-13-4]

$\text{C}_{18}\text{H}_{25}\text{NO}$ 271.402

Isol. from *Achillea magnifica*.

Ulubelen, A. *et al.*, *Planta Med.*, 1989, **55**, 395 (*isol*)

2,4-Tetradecadiene-8,10,12- triynoic acid T-123

$\text{H}_3\text{CC}\equiv\text{CC}\equiv\text{CC}\equiv\text{CCH}_2\text{CH}_2\text{CH}=\text{CHCH}=\text{CHCOOH}$

$\text{C}_{14}\text{H}_{12}\text{O}_2$ 212.248

(E,E)-form

Piperidide: 2,4-Tetradeca-8,10,12-triynoic acid piperidide. 1-(1-Oxo-2,4-tetradecadiene-8,10,12-triynyl)piperidine, 9CI

[83182-43-8]

$\text{C}_{19}\text{H}_{21}\text{NO}$ 279.381

Alkamide from the roots of *Achillea lycanica* and *Achillea spinulifolia* (Asteraceae). Mp 115°.

Greger, H. *et al.*, *Phytochemistry*, 1982, **21**, 1071 (*isol. uv. ir. pmr. ms. struct*)

2,4-Tetradecadienoic acid, 9CI T-124

[75011-60-8]

$\text{H}_3\text{C}(\text{CH}_2)_8\text{CH}=\text{CHCH}=\text{CHCOOH}$

$\text{C}_{14}\text{H}_{24}\text{O}_2$ 224.342

(2E,4E)-form [24738-49-6]

Cryst. (petrol at low temp.). Mp 57-58°.

Me ester: [24738-47-4]

$\text{C}_{15}\text{H}_{26}\text{O}_2$ 238.369

Bp_{0.3} 117-118°.

2-Methylpropylamide: 2,4-Tetradecadienoic acid isobutylamide

[13891-73-1]

$\text{C}_{18}\text{H}_{33}\text{NO}$ 279.465

Alkaloid from *Chrysanthemum frutescens*, *Anacyclus pyrethrum* and *Leucocyclis formosus* (Asteraceae) (in admixture with homologues). Cryst. (petrol). Mp 88°.

2-Hydroxy-2-methylpropylamide: N-(2-Hydroxyisobutyl)-2,4-tetradecadienamide. **Tetrahydrobungeoil**

[200938-43-8]

$\text{C}_{18}\text{H}_{33}\text{NO}_2$ 295.464

Alkaloid from *Zanthoxylum bungeanum* and *Zanthoxylum integrifolium*. Unstable powder.

Pyrrolidide: N-(2,4-Tetradecadienoyl)-pyrrolidide. 2,4-Tetradecadienoic acid pyrrolidide

[88855-41-8]

[88855-38-3]

$\text{C}_{18}\text{H}_{31}\text{NO}$ 277.449

Alkaloid from *Achillea nana*. Mp 36-38° (synthetic).

Piperidide: 1-(1-Oxo-2,4-tetradecadienyl)piperidine, 9CI. 1-(2,4-Tetradecadienoyl)piperidine. 2,4-Tetradecadienoic acid piperidide

[52657-12-2]

$\text{C}_{19}\text{H}_{33}\text{NO}$ 291.476

Alkaloid from the roots of *Otanthus maritimus* (Asteraceae). Oil. Bp_{0.1} 150°.

4,5-Epoxyde: 3-(2-Nonyloxiranyl)-2-propenoic acid. 9CI. **4,5-Epoxy-2-tetradecenoic acid**

[154291-61-9]

$\text{C}_{14}\text{H}_{24}\text{O}_3$ 240.342

Constit. of a sterile fungus found growing in grass. Mp 81-81.5°.

(2E,4Z)-form

Me ester: [54977-81-0]

Oil.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1967, **100**, 104; 1974, **107**, 1038 (*isobutylamide, piperidide*)

Burden, R.S. *et al.*, *J.C.S. (C)*, 1969, 2477 (*synth*)

Greger, H. *et al.*, *Phytochemistry*, 1981, **20**, 2579; 1984, **23**, 1503-1505 (*isobutylamide, pyrrolidide*)

Tsuboi, S. *et al.*, *J.O.C.*, 1982, **47**, 4478-4482 (*2E,4Z-Me ester*)

Tsuboi, S. *et al.*, *J.O.C.*, 1984, **49**, 1204-1208 (*synth. amides*)

Schwartz, R.E. *et al.*, *Tetrahedron*, 1994, **50**, 1675 (*4,5-epoxide*)

Sisiri, W. *et al.*, *J.O.C.*, 1996, **61**, 5911 (*synth. ir. pmr*)

Cow, C. *et al.*, *Can. J. Chem.*, 1997, **75**, 884-889 (*synth. ir. pmr. cmr. ms. 4,5-epoxide*)

- Xiong, Q. *et al.*, *Phytochemistry*, 1997, **46**, 1123-1126 (*Tetrahydrobungeoanal*)
- 2,4-Tetradecadien-8-ynoic acid** T-125
 $\text{H}_3\text{C}(\text{CH}_2)_4\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}=\text{CHCH}=\text{CHCOOH}$
 $\text{C}_{14}\text{H}_{20}\text{O}_2$ 220.311
- (2E,4E)-form**
Pyrrrolidide: 2,4-Tetradecadien-8-ynoic acid pyrrrolidide. 1-(1-Oxo-2,4-tetradecadien-8-ynyl)pyrrrolidine, 9CI [111509-33-2]
 $\text{C}_{18}\text{H}_{27}\text{NO}$ 273.417
 Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.
- Greger, H. *et al.*, *Phytochemistry*, 1987, **26**, 2289 (*isol, uv, ir, pmr, ms, struct*)
- 2,4,6,8,10,12-Tetradecahexaenedioic acid** T-126
 $\text{HOOCCH}=\text{CHCH}=\text{CHCH}=\text{CHCH}=\text{CHCOOH}$
 $\text{C}_{14}\text{H}_{14}\text{O}_4$ 246.262
- (all-E)-form**
Corticocin
 [505-53-3]
 Pigment of the fungus *Corticium croceum*. Also from paprika (*Capsicum annuum*) and from *Encephalartos* spp. Orange-red needles. Insol. most solvs. Mp ca. 300° subl. dec.
- Di-Me ester:* [77186-00-6]
 $\text{C}_{16}\text{H}_{18}\text{O}_4$ 274.316
 Orange cryst. (CHCl_3 or AcOH). Mp 230-232°.
- Di-Et ester:* [1441-61-8]
 $\text{C}_{18}\text{H}_{22}\text{O}_4$ 302.369
 Mp 204-205°.
- l-L-Aspartic acid, 14-L-isoleucine diamide: Boletocrocin D*
 $\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_8$ 474.51
 Isol. from the fruit bodies of *Boletus laetissimus*.
- Erdtman, H. *et al.*, *Acta Chem. Scand.*, 1948, **2**, 209 (*isol, struct*)
 Shaw, B.L. *et al.*, *J.C.S.*, 1954, 3217 (*synth*)
 Kovalev, B.G. *et al.*, *CA*, 1963, **58**, 11209f (*synth*)
 Rives-Arnau, V. *et al.*, *Macromolecules*, 1985, **18**, 2088
 Kahner, L. *et al.*, *Phytochemistry*, 1998, **49**, 1693-1697 (*Boletocrocin D*)
 Schreiner, T. *et al.*, *Z. Naturforsch.*, C, 1998, **53**, 4-8 (*isol, synth*)
- 2,4,8,10,12-Tetradecapentaenoic acid** T-127
 $\text{H}_3\text{C}(\text{CH}=\text{CH})_3\text{CH}_2\text{CH}_2\text{CH}=\text{CHCH}=\text{CHCOOH}$
 $\text{C}_{14}\text{H}_{18}\text{O}_2$ 218.295
- (all-E)-form**
2-Hydroxy-2-methylpropylamide: Hydroxy-γ-isosanshool. N-(2-Hydroxyisobutyl)-2,4,8,10,12-tetradecapentaenamide [127514-62-9]
 $\text{C}_{18}\text{H}_{27}\text{NO}_2$ 289.417
 Constit. of *Zanthoxylum* spp. (Rutaceae). Powder. λ_{max} 258 (ε 26300); 268 (ε 28900); 279 (ε 26010) (EtOH).
- (2E,4E,8Z,10E,12E)-form**
2-Methylpropylamide: N-Isobutyl-2,4,8,10,12-tetradecapentaenamide. γ-Sanshool [78886-65-4]
 $\text{C}_{18}\text{H}_{27}\text{NO}$ 273.417
 Constit. of *Zanthoxylum ailanthoides*, *Zanthoxylum piperitum* (Japanese pepper tree) and other *Zanthoxylum* spp. (Rutaceae). Needles (hexane). Mp 88-89°. λ_{max} 260 (ε 48000); 272 (ε 57000); 280 (ε 45000) (EtOH).
- 2-Hydroxy-2-methylpropylamide: Hydroxy-γ-sanshool* [78886-66-5]
 $\text{C}_{18}\text{H}_{27}\text{NO}_2$ 289.417
 Constit. of *Zanthoxylum ailanthoides*, *Zanthoxylum bungeanum* and *Zanthoxylum piperitum* (Japanese pepper tree) (Rutaceae). Needles (CHCl_3). Mp 122-123°.
- 2-Methylenepropylamide: Dehydro-γ-sanshool*
 $\text{C}_{18}\text{H}_{25}\text{NO}$ 271.402
 Alkaloid from *Zanthoxylum bungeanum*. Unstable oil.
- (2E,4E,8Z,10Z,12E)-form**
Me ester: [114567-68-9]
 $\text{C}_{15}\text{H}_{20}\text{O}_2$ 232.322
 Constit. of *Sanvitalia oxymoides*. Gum. Incorrect name in ref.
- 2-Methylpropylamide:* [114612-73-6]
 $\text{C}_{18}\text{H}_{27}\text{NO}$ 273.417
 Constit. of *Sanvitalia oxymoides*. Gum. [98168-69-5, 98095-54-6, 117581-14-3]
 Yasuda, I. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1791-1793 (*γ-Sanshool, Hydroxy-γ-sanshool*)
 Crombie, L. *et al.*, *Tet. Lett.*, 1985, **26**, 2481 (*synth*)
 Dominguez, X.A. *et al.*, *Rev. Latinoam. Quim.*, 1987, **18**, 114 (*isol*)
 Mizutani, K. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**, 2362 (*isol, ir, pmr, cmr*)
 Xiong, Q. *et al.*, *Phytochemistry*, 1997, **46**, 1123-1126 (*isol, pmr, cmr, ms*)
- 2,4,6,12-Tetradecatetraene-8,10-diyenoic acid** T-128
 $\text{H}_3\text{CCH}=\text{CHC}\equiv\text{CC}\equiv\text{C}(\text{CH}=\text{CH})_3\text{COOH}$
 $\text{C}_{14}\text{H}_{12}\text{O}_2$ 212.248
- (all-E)-form**
 Isol. from the underground parts of *Achillea wilhelmsii*.
- Piperidide: 2,4,6,12-Tetradecatetraene-8,10-diyenoic acid piperidide. 1-(1-Oxo-2,4,6,12-tetradecatetraene-8,10-diyenyl)piperidine, 9CI* [83182-49-4]
 $\text{C}_{19}\text{H}_{21}\text{NO}$ 279.381
 Alkamide from the roots of *Achillea grandifolia* (Asteraceae). Unstable yellow foam.
- (2E,4E,6E,12Z)-form**
Me ester: [14109-34-3]
 $\text{C}_{15}\text{H}_{14}\text{O}_2$ 226.274
 Isol. from *Sanvitalia procumbens*. Cryst. (Et_2O /petrol). Mp 102°. λ_{max} 252; 275; 281; 348; 368 (no solvent reported).
- 2-Methylpropylamide: 2,4,6,12-Tetradecatetraene-8,10-diyenoic acid isobutylamide* [113235-98-6]
 $\text{C}_{18}\text{H}_{21}\text{NO}$ 267.37
 Isol. from underground parts of *Achillea wilhelmsii*. Oil. Contains 30% of the 6Z-isomer. λ_{max} 264; 279; 333 (sh); 348; 368 (Et_2O).
- 3-Methylbutylamide: 2,4,6,12-Tetradecatetraene-8,10-diyenoic acid isopentylamide* [113235-95-3]
 $\text{C}_{19}\text{H}_{23}\text{NO}$ 281.397
 Isol. from underground parts of *Achillea wilhelmsii*. Cryst. Mp 136-139°. λ_{max} 217; 279; 333; 347; 368 (Et_2O).
- (2E,4E,6Z,12Z)-form**
2-Methylpropylamide: [113235-97-5]
 Isol. from underground parts of *Achillea wilhelmsii*. Oil. Obt. as a mixt. containing 6E-isomer. λ_{max} 264; 277; 335 (sh); 348; 366 (sh) (Et_2O).
- 3-Methylbutylamide:* [113235-94-2]
 Isol. from underground parts of *Achillea wilhelmsii*. Oil. λ_{max} 265; 277; 332 (sh); 347; 365 (sh) (Et_2O).
- Bohlmann, F. *et al.*, *Chem. Ber.*, 1966, **99**, 3194 (*isol, struct, uv, ir, pmr*)
 Greger, H. *et al.*, *Phytochemistry*, 1982, **21**, 1071 (*piperidide*)
 Greger, H. *et al.*, *J. Nat. Prod.*, 1987, **50**, 1100 (*amides, isol, uv, ir, ms, pmr*)
- 2,4,8,11-Tetradecatetraenoic acid** T-129
 $\text{H}_3\text{CCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}_2\text{CH}=\text{CHCH}=\text{CHCOOH}$
 $\text{C}_{14}\text{H}_{20}\text{O}_2$ 220.311
- (2E,4E,8Z,11E)-form**
2-Methylpropylamide: Lanyuamide III
 $\text{C}_{18}\text{H}_{29}\text{NO}$ 275.433
 Alkaloid from the fruit of *Zanthoxylum integrifolium*. Oil. λ_{max} 259 (log ε 4.23) (EtOH).
- 2-Hydroxy-2-methylpropylamide: Isobungeoool. Hydroxylanyuamide III* [127460-60-0]
 [117568-41-9]
 $\text{C}_{18}\text{H}_{29}\text{NO}_2$ 291.433
 Alkaloid from *Zanthoxylum bungeanum*. Syrup. λ_{max} 268 (ε 27350) (EtOH).
- (2E,4E,8Z,11Z)-form**
2-Methylpropylamide: Hazaleamide [81427-15-8]
 $\text{C}_{18}\text{H}_{29}\text{NO}$ 275.433
 Alkaloid from *Leucocyclus formosus* (Asteraceae). Gum.
- 2-Hydroxy-2-methylpropylamide: Bungeoool* [117568-40-8]
 $\text{C}_{18}\text{H}_{29}\text{NO}_2$ 291.433
 Alkaloid from *Zanthoxylum bungeanum*. Syrup. λ_{max} 268 (ε 26190) (EtOH).
- Pyrrrolidide: 2,4,8,11-Tetradecatetraenoic acid pyrrrolidide. 1-(1-Oxo-2,4,8,11-tetradecatetraenyl)pyrrrolidine, 9CI*

[111509-32-1]
C₁₈H₂₇NO 273.417

Isol. from the underground parts of *Achillea ageratifolia* ssp. *serbica* (Asteraceae). Oil.

Piperidide: 1-(1-Oxo-2,4,8,11-tetradecatetraenyl)piperidine. (2,4,8,11-Tetradecatetraenyl)piperidine. 2,4,8,11-Tetradecatetraenoic piperidide
C₁₉H₂₉NO 287.444

Alkaloid from *Otanthus maritimus*. Oil.

Greger, H. et al., *Phytochemistry*, 1981, **20**, 2579; 1987, **26**, 2289 (isol, uv, ir, pmr, ms, struct)


Mizutani, K. et al., *Chem. Pharm. Bull.*, 1988, **36**, 2362-2365 (isol)

Shibuya, H. et al., *Chem. Pharm. Bull.*, 1992, **40**, 2325 (Hazaleamide)

Chen, I.-S. et al., *Phytochemistry*, 1999, **52**, 357-360 (isol, cmr)

Christodouloupoulou, L. et al., *J. Agric. Food Chem.*, 2005, **53**, 1435-1439 (piperidide)

2,4,10,12-Tetradecatetraen-8-yenoic acid T-130



C₁₄H₁₆O₂ 216.279


(2E,4E,10E,12Z)-form

2-Methylpropyl amide: N-(2-Methylpropyl)-2,4,10,12-tetradecatetraen-8-ynamide. N-Isobutyl-2,4,10,12-tetradecatetraen-8-ynamide
C₁₈H₂₅NO 271.402

Alkaloid from *Chrysanthemum morifolium*. Oil. λ_{max} 270 (MeCN aq.).

Tsao, R. et al., *J. Nat. Prod.*, 2003, **66**, 1229-1231 (isol, ir, pmr, cmr, ms)

3,5,13-Tetradecatrien-2-amine T-131



C₁₄H₂₅N 207.358

(2R,3E,5E)-form

13,14-Dihydro: 3,5-Tetradecadien-2-amine, 9CI. 2-Amino-3,5-tetradecadiene

[356073-87-5]
C₁₄H₂₇N 209.374

Isol. from a *Pseudodistoma* sp. Oil. [α]_D -1.2 (c, 0.25 in MeOH). λ_{max} 229 (log ε 4.24) (MeOH).

(2S,3E,5Z)-form

(+)-trans,cis-form
[151284-88-7]

Prod. by tunicate *Pseudodistoma novaezelandiae*. Viscous pale yellow oil. [α]_D²⁵ +17.8 (c, 1 in CHCl₃) (95% e.e.). λ_{max} 230 (ε 16600) (MeOH) (Berdy).

(2RS,3E,5E)-form

(±)-trans,trans-form
[134381-24-1]

Constit. of the ascidian *Pseudodistoma novaezelandiae*. λ_{max} 230 (ε 16600) (MeOH) (Berdy).

(2RS,3E,5Z)-form

(±)-trans,cis-form
[134381-23-0]

Constit. of *Pseudodistoma novaezelandiae*.

13,14-Dihydro: [134381-25-2]

Constit. of *Pseudodistoma novaezelandiae*.

Perry, N.B. et al., *Aust. J. Chem.*, 1991, **44**, 627-633 (isol, uv, pmr, cmr)

Enders, D. et al., *Annalen*, 1993, 551-555 (synth)

Rashid, M.A. et al., *Tetrahedron*, 2001, **57**, 5751-5755 (3R,3E,5E-form 13,14-dihydro)

2,4,12-Tetradecatriene-8,10-diynoic acid T-132



C₁₄H₁₄O₂ 214.263

(E,E,E)-form

2-Methylpropylamide: see Anacycline, A-968

Piperidide: 2,4,12-Tetradecatriene-8,10-diynoic acid piperidide. 1-(1-Oxo-2,4,12-tetradecatriene-8,10-diynyl)piperidine, 9CI

[83182-48-3]

C₁₉H₂₃NO 281.397

Alkamide from *Achillea biebersteinii* and *Achillea sudetica* (Asteraceae). Oil.

Pyrrolidide: 2,4,12-Tetradecatriene-8,10-diynoic acid pyrrolidide. 1-(1-Oxo-2,4,12-tetradecatriene-8,10-diynyl)pyrrolidine

[94413-15-7]

C₁₈H₂₁NO 267.37

Isol. from underground parts of *Achillea nana*. Cryst. Mp 107-110°. λ_{max} 282, 266, 253 nm (Et₂O).

(2E,4E,12Z)-form

2-Methylpropylamide: [112711-13-4]

Isol. from *Achillea ptarmica* and *Chrysanthemum morifolium*. Cryst. (petrol). Mp 100-101°. Rapidly becoming dull pink in light. λ_{max} 256 (35000), 267 (43000), 283 (19000) nm.

3-Methylbutylamide: N-(3-Methylbutyl)-2,4,12-tetradecatriene-8,10-diyname. 2,4,12-Tetradecatrien-8,10-diynoic acid isopentylamide

[113235-99-7]

C₁₉H₂₅NO 283.413

Isol. from *Achillea wilhelmsii*. Cryst. Mp 83-85°. λ_{max} 282, 266, 252 nm (Et₂O).

Crombie, L. et al., *J.C.S.*, 1957, 2767 (synth, uv, ir)


Bohlmann, F. et al., *Chem. Ber.*, 1973, **106**, 1328 (isol, ms, pmr, struct)

Greger, H. et al., *Phytochemistry*, 1981, **20**, 2579; 1984, **23**, 1503; 1989, **28**, 2363 (isol, struct, pmr, uv, ms)

Greger, H. et al., *J. Nat. Prod.*, 1987, **50**, 1100 (isol, struct, uv, ir, ms, pmr)

Kuropka, G. et al., *Planta Med.*, 1987, **53**, 440 (isol, pmr)

2,4,8-Tetradecatrienoic acid T-133



C₁₄H₂₂O₂ 222.327

(2E,4E,8Z)-form

2-Methylpropylamide: 2,4,8-Tetradeca-

trienoic acid isobutylamide. N-(2-Methylpropyl)-2,4,8-tetradecatrienamide, 9CI

[81427-14-7]

C₁₈H₃₁NO 277.449

Alkaloid from *Leucocyclyus formosus* (Asteraceae) and from fruits of *Phellodendron chinense* (Rutaceae). Gum.

2-Hydroxy-2-methylpropylamide: N-(2-Hydroxyisobutyl)-2,4,8-tetradecatrienamide. **Dihydrobungeanool**

[200938-41-6]

C₁₈H₃₁NO₂ 293.448

Alkaloid from *Zanthoxylum bungeanum*. Unstable oil.

Piperidide: 1-(1-Oxo-2,4,8-tetradecatrienyl)piperidine. N-(2,4,8-Tetradecatrienyl)piperidine. 2,4,8-Tetradecatrienoic piperidide

C₁₉H₃₁NO 289.46

Alkaloid from *Otanthus maritimus*. Oil.


Greger, H. et al., *Phytochemistry*, 1981, **20**, 2579-2581 (isol, ms, struct)

Su, R. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1994, **36**, 817-820; *CA*, **122**, 209774n

Xiong, Q. et al., *Phytochemistry*, 1997, **46**, 1123-1126 (*Dihydrobungeanool*)

Christodouloupoulou, L. et al., *J. Agric. Food Chem.*, 2005, **53**, 1435-1439 (piperidide)

2,4,10-Tetradecatrien-8-yenoic acid T-134



C₁₄H₁₈O₂ 218.295

(2E,4E,10Z)-form

N-(2-Methylpropyl)amide: 2,4,10-Tetradecatrien-8-yenoic acid isobutylamide.

N-(2-Methylpropyl)-2,4,10-tetradecatrien-8-ynamide

[83204-63-1]

C₁₈H₂₇NO 273.417

Alkaloid from the roots of *Achillea tomentosa* (Asteraceae). Oil.

Piperidide: 2,4,10-Tetradecatrien-8-yenoic acid piperidide. 1-(1-Oxo-2,4,10-tetradecatrien-8-ynyl)piperidine, 9CI

[83182-42-7]

C₁₉H₂₇NO 285.428

Alkaloid from the roots of *Achillea spinulifolia* (Asteraceae). Gum.

Pyrrolidide: 2,4,10-Tetradecatrien-8-yenoic acid pyrrolidide

[94413-13-5]

C₁₈H₂₅NO 271.402

Constit. of *Achillea nana* (Asteraceae).


Greger, H. et al., *Phytochemistry*, 1982, **21**, 1071 (isol, uv, ir, pmr, ms, struct)

Bohlmann, F. et al., *Tetrahedron*, 1983, **39**, 123 (synth)

Greger, H. et al., *Phytochemistry*, 1984, **23**, 1503-1505 (pyrrolidide)

Hofer, O. et al., *Tetrahedron*, 1986, **42**, 2707-2716 (pmr, cmr, alkamides)

2-Tetradecylpyrrolidine T-135



C₁₈H₃₇N 267.497**(R)-form**


N-Me: 1-Methyl-2-tetradecylpyrrolidine,

9Cl. Bgugaine

[155416-32-3]

C₁₉H₃₉N 281.524Alkaloid from tubers of *Arisarum vulgare*. Displays antifungal activity, active against gram-positive bacteria. Oil. [α]_D -48 (MeOH).Melhaoui, A. et al., *Nat. Prod. Lett.*, 1993, **2**, 237 (isol, struct)Jossang, A. et al., *Heterocycles*, 1996, **43**, 755 (synth)Takahata, H. et al., *Heterocycles*, 1997, **46**, 349-356 (synth, pmr, cmr)Majik, M.S. et al., *J. Chem. Res., Synop.*, 2008, 121-122 (synth, pmr, cmr)**3,4,5,6-Tetrahydrodihydro-corynantheol** T-136



[88644-42-2]

C₁₉H₂₂N₂O 294.396Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph. [α]_D -56 (c, 0.57 in MeOH).**19,20E-Didehydro: 3,4,5,6-Tetrahydrogeissoschizol**C₁₉H₂₀N₂O 292.38Alkaloid from *Rauwolfia serpentina*. Yellow powder. [α]_D²⁰ +30.9 (c, 0.3 in MeOH). λ_{max} 253 (log ε 4.09); 307 (log ε 3.92); 366 (log ε 3.28) (MeOH).**19,20Z-Didehydro:**C₁₉H₂₀N₂O 292.38Alkaloid from *Rauwolfia serpentina*. Yellow powder. [α]_D²⁰ +7 (c, 0.5 in MeOH). λ_{max} 253 (log ε 4.13); 308 (log ε 3.97); 366 (log ε 3.31) (MeOH).**10-Methoxy: 10-Methoxy-3,4,5,6-tetrahydro-18,19-dihydrocorynantheol** [92446-40-7]C₂₀H₂₄N₂O₂ 324.422Alkaloid from the bark of *Neisosperma glomerata* (Apocynaceae). Tentative struct. Isol. as an inseparable mixt. with 10-Methoxy-3,4,5,6-tetrahydrocorynantheol, 3,4,5,6-Tetrahydrochroprosinine and 3,4,5,6,18,19-Hexahydrochroprosinine. Anhydronium base.**10-Methoxy, 18,19-didehydro: 10-Methoxy-3,4,5,6-tetrahydrocorynantheol** [92446-39-4]C₂₀H₂₂N₂O₂ 322.406Alkaloid from the bark of *Neisosperma glomerata* (Apocynaceae). Tentative struct. Anhydronium base.**10,11-Dimethoxy: 3,4,5,6-Tetrahydrochroprosinine** [92446-41-8]C₂₁H₂₆N₂O₃ 354.448Alkaloid from the bark of *Neisosperma glomerata* (Apocynaceae). Tentative struct. Isol. as an inseparable mixt. with 10-Methoxy-3,4,5,6-tetrahydro-18,19-dihydrocorynantheol, 10-Methoxy-3,4,5,6-tetrahydrocorynantheol and 3,4,5,6,18,19-Hexahydrochroprosinine.**10,11-Dimethoxy, 18,19-didehydro:****3,4,5,6,18,19-Hexahydrochroprosinine** [92458-34-9]C₂₁H₂₄N₂O₃ 352.432Alkaloid from the bark of *Neisosperma glomerata* (Apocynaceae). Tentative struct. Anhydronium base.Seguin, E. et al., *J. Nat. Prod.*, 1982, **45**, 738; 1984, **47**, 687-691

(Tetrahydrochroprosinine, Hexahydrochroprosinine, Methoxytetrahydrodihydrocorynantheol, Methoxytetrahydrocorynantheol)

Fujii, T. et al., *Tetrahedron*, 1993, **49**, 1879-1890 (Tetrahydrodihydrocorynantheol)Wachsmuth, O. et al., *Phytochemistry*, 2002, **61**, 705-709 (Tetrahydrogeissoschizols)**Tetrahydroechinulin** T-137

Alkaloid E7 [64361-66-6]


C₂₉H₃₅N₃O₂ 457.614Alkaloid from *Aspergillus amstelodami*.Inoue, S. et al., *Yakugaku Zasshi*, 1977, **97**, 576 (isol)**3,4,5,6-Tetrahydrogeissoschizine** T-138

Absolute Configuration

C₂₁H₂₀N₂O₃ 348.401**O-β-D-Glucopyranoside:**C₂₇H₃₀N₂O₈ 510.543Alkaloid from *Rauwolfia serpentina*. Yellow powder. [α]_D²⁰ +118.3 (c, 0.1 in MeOH). CAS no. not found 8-14Cl. λ_{max} 252 (log ε 4.23); 307 (log ε 4.04); 366 (log ε 3.34) (MeOH).Wachsmuth, O. et al., *Phytochemistry*, 2002, **61**, 705-709 (isol, pmr, cmr, ms)**Tetrahydroisocalcathine** T-139

[202932-49-8]

[202932-50-1]



Absolute Configuration

C₂₂H₂₂N₄ 342.443The stereochem. of this group apart from Tetrahydroisocalcathine itself requires clarification. Alkaloid from the flowers of *Psychotria colorata*. Needles (diisopropyl ether/CHCl₃). Mp 253°. [α]_D²⁰ -449 (c, 1 in EtOH). λ_{max} 274 (log ε 4.38); 280 (log ε 4.37); 300 (sh) (log ε 4.03) (MeOH).**8,8a-Dihydro: Bhesine**

[155322-45-5]

C₂₂H₂₄N₄ 344.458Alkaloid from *Bhesa paniculata* (Celastraceae). No stereochem. determined.**8,8a,8',8'-a-Tetrahydro: Isocalcathine**

[1399-92-4]

C₂₂H₂₆N₄ 346.474Alkaloid from stem bark and fruits of *Psychotria forsteriana* (Rubiaceae). Mp 253°. [α]_D²⁰ -150 (c, 2.5 in EtOH). Stereochem. not determined.**8,8a,8',8'-a-Tetrahydro, stereoisomer: Alkaloid CPC 2**


[886752-87-0]

C₂₂H₂₆N₄ 346.474Alkaloid from *Chimonanthus praecox* f. *concolor*. Amorph. solid. [α]_D²⁶ +57 (c, 0.04 in EtOH). The enantiomeric 3*a*S,3'*a*S-config. was proposed on little evidence. λ_{max} 208 (log ε 4.53); 250 (log ε 4.12); 295 (log ε 3.59) (MeOH).**Stereoisomer (?): Dehydrobhesine**

[155210-51-8]

Alkaloid from *Bhesa paniculata*. No stereochem. determined.Adjibade, Y. et al., *Phytochemistry*, 1992, **31**, 317-319 (*Isocalcathine*)Balayer, A. et al., *Nat. Prod. Lett.*, 1993, **2**, 61-67 (*Bhesine, Dehydrobhesine*)Verotta, L. et al., *J. Nat. Prod.*, 1998, **61**, 392-396 (*Tetrahydroisocalcathine, crystal struct*)Kitajima, M. et al., *Tet. Lett.*, 2006, **47**, 3199-3202 (*CPC 2*)**3,4,5,6-Tetrahydroyohimbine** T-140

[181283-45-4]



C₂₁H₂₂N₂O₃ 350.416

Alkaloid from *Rauwolfia serpentina*. Yellow powder. Mp 260-264°. [α]_D²⁰ +175.4 (c, 0.3 in MeOH). λ_{\max} 252 (log ϵ 4.14); 306 (log ϵ 3.97); 365 (log ϵ 3.37) (MeOH).

17-Epimer: **3,4,5,6-Tetrahydro- β -yohimbinium**

[50834-28-1]

[50439-43-5, 50315-64-5]

C₂₁H₂₃N₂O₃[⊕] 351.424

Quaternary alkaloid from *Asonia elliptica* (Apocynaceae). Cryst. + 2H₂O (as hydrochloride). Mp 233-234° (hydrochloride). [α]_D -40.6 (c, 1.09 in H₂O) (hydrochloride).

Sakai, S. *et al.*, *Yakugaku Zasshi*, 1973, **93**, 483-489 (17-epimer)


Stahl, R. *et al.*, *Helv. Chim. Acta*, 1996, **79**, 1361-1378 (synth, pmr, cmr)

Wachsmuth, O. *et al.*, *Phytochemistry*, 2002, **61**, 705-709 (isol, pmr, cmr, ms)

Tetrahydroakagerine

T-141

[113122-66-0]

C₂₀H₂₈N₂O₂ 328.453

Alkaloid from the root bark and stem bark of *Strychnos johnsonii* (Loganiaceae). [α]_D +49 (c, 0.2 in MeOH).

1,2,3,4-Tetrahydro, N-de-Me: **Deppeaninol**

[180785-98-2]

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from leaves of *Deppea blumenaviensis* (Rubiaceae). Amorph. [α]_D²⁵ +54 (c, 0.41 in MeOH). R-Config. at C-15 assigned on biogenetic grounds. λ_{\max} 235 (log ϵ 4.36); 250 (sh) (log ϵ 3.93); 280 (sh) (log ϵ 3.61); 288 (log ϵ 4.07); 336 (log ϵ 3.54) (EtOH).

Massiot, G. *et al.*, *Phytochemistry*, 1987, **26**, 2839-2846 (isol, uv, ir, pmr, cmr, ms, struct)

Kan-Fan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 317-321 (Deppeaninol)


5,6,7,8-Tetrahydrobiopterin

T-142

2-Amino-6-(1,2-dihydroxypropyl)-5,6,7,8-tetrahydro-4(1H)-pteridinone, 9CI

[17528-72-2]

[90366-07-7]

C₉H₁₅N₅O₃ 241.249

► UO3516000

(1'R,2'S,6R)-form

L-erythro-form. **Sapropterin**, INN. Biop-

ten. Kuvan. *Phenoptin*. Sun 0588. 6R-BH4. T 1401

[99630-29-2]

Cofactor for phenylalanine, tyrosine and tryptophan hydroxylases which influences the biosynth. of neurotransmitting catecholamines. Used for treatment of hyperphenylalaninemia. pK_{a1} 1.29; pK_{a2} 5.02; pK_{a3} 10.77. Other stereoisomers known.

Hydrochloride (1:2): **Sapropterin dihydrochloride**, USAN

[69056-38-8]

Needles (HCl/EtOH). Mp 245-246°. [α]_D²⁰ -9.8 (c, 0.56 in 0.1M HCl). Also forms a 2:3 hydrochloride.

Tetra-Ac: [81873-18-9]

Cryst. (MeOH). Mp 292° dec. [α]_D²⁰ -144 (c, 0.5 in CHCl₃).

1',2'-Diketone: **Dyspropterin**. 6-Pyruvyl-tetrahydropterin. 6-Pyruvyl-5,6,7,8-tetrahydropterin

[89687-39-8]

[101383-42-0]

C₉H₁₁N₅O₃ 237.218

Intermed. in the biosynthetic pathway of Tetrahydrobiopterin.

[62961-57-3, 105119-63-9, 62989-33-7, 69056-39-9, 71031-45-3, 71074-53-8, 83023-72-7, 83709-59-5, 99630-30-5, 105119-62-8, 27070-47-9]

Brethauer, R. *et al.*, *Z. Naturforsch., B*, 1972, **27**, 580-581 (isol, tlc)

Ayling, J.E. *et al.*, *Biochemistry*, 1973, **12**, 2045-2051 (biochem)

Williams, V.P. *et al.*, *J. Het. Chem.*, 1973, **10**, 827-833 (ms)

Schircks, B. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 211-214; 1978, **61**, 2731-2738 (synth, pmr, cmr)

Bailey, S.W. *et al.*, *J. Biol. Chem.*, 1978, **253**, 1598-1605 (resoln, uv, cd)

Furrer, H.J. *et al.*, *Helv. Chim. Acta*, 1979, **62**, 2577 (synth, pmr, cmr)

Armarego, W.L.F. *et al.*, *Aust. J. Chem.*, 1982, **35**, 785-793; 1984, **37**, 355-366 (synth, pmr, cmr, abs config, bibl, tetra-Ac)

Whiteley, J.M. *et al.*, *Anal. Biochem.*, 1984, **137**, 394-396 (synth)

Masada, M. *et al.*, *Biochim. Biophys. Acta*, 1985, **840**, 235-244 (Dyspropterin)

Matsuura, S. *et al.*, *Heterocycles*, 1985, **23**, 3115-3120 (synth, uv, pmr, cmr, ms)

Matsuura, S. *et al.*, *J. Biochem. (Tokyo)*, 1985, **98**, 1341-1347; *CA*, **104**, 30828 (abs config, bibl, cryst struct)

Smith, G.K. *et al.*, *J. Biol. Chem.*, 1986, **261**, 2725-2737 (biosynth, uv)

Levine, R.A. *et al.*, *Ann. N.Y. Acad. Sci.*, 1988, **521**, 129-139 (rev, metab)

Hirotsu, I. *et al.*, *Oyo Yakuri*, 1989, **38**, 511-529 (pharmacol)

Hirotsu, I. *et al.*, *Eur. J. Pharmacol.*, 1990, **183**, 1461-1462 (pharmacol)

Duch, D.S. *et al.*, *J. Nutr. Biochem.*, 1991, **2**, 411-423 (rev, biosynth)

Werner, E.R. *et al.*, *Proc. Soc. Exp. Biol. Med.*, 1993, **203**, 1-12 (rev)

Steinfeld, R. *et al.*, *J. Inherited Metab. Dis.*, 2004, **27**, 449-453 (pharmacol)

Fiege, B. *et al.*, *Mol. Genet. Metab.*, 2004, **81**, 45-51 (pharmacokinetic)


Burnett, J.R. *et al.*, *IDrugs*, 2007, **10**, 805-813 (rev)

Michals-Matalon, K. *et al.*, *Expert Opin. Invest. Drugs*, 2008, **17**, 245-251 (pharmacol)

Burton, B.K. *et al.*, *Nat. Rev. Drug Discovery*, 2008, **7**, 199-200 (rev)

1,2,3,6-Tetrahydro-2,3'-bipyridine, 9CI

T-143

Anatabine

(S)-form

C₁₀H₁₂N₂ 160.218**(R)-form** [126454-22-6]

Synthetic.

Oil. [α]_D +156 (c, 1.6 in CHCl₃).

(S)-form [581-49-7]

Alkaloid from *Nicotiana tabacum*, *Nicotiana glutinosa* and *Duboisia hopwoodii*. Second most abundant alkaloid after Nicotine in most tobacco varieties (Solanaceae). Bp₁₀ 145-156°. [α]_D¹⁷ -177.8 (neat). n_D^{20} 1.5676.

► BV6495000

Picrate: Mp 191-193°.

N-Benzoyl: Bp_{0.01} 160-170°. [α]_D¹⁵ -15.4 (MeOH).

N-Me: N-Methylanatabine

[5953-51-5]

C₁₁H₁₄N₂ 174.245

Minor alkaloid from *Nicotiana tabacum* (Solanaceae). Oil. Bp₁ 120°. [α]_D¹⁸ -17.4 (MeOH).

N-Me, dipicrate: Mp 207-208°.

(±)-form [2743-90-0]

Alkaloid from tobacco (Solanaceae). Oil. Bp₁₀ 145-156°. n_D^{20} 1.5655.

Perchlorate (1:2): Mp 129-230°.

Dipicrate: Mp 201-201.5° (198-200°).

Späth, E. *et al.*, *Ber.*, 1936, **69**, 2448; 1937, **70**, 239; 704 (isol, struct, resoln)

Lukeš, R. *et al.*, *Coll. Czech. Chem. Comm.*, 1962, **27**, 751 (abs config)

Quan, P.M. *et al.*, *J.O.C.*, 1965, **30**, 2769 (synth, struct)

Leete, E. *et al.*, *Chem. Comm.*, 1975, 9 (biosynth)

Leete, E. *et al.*, *J.A.C.S.*, 1976, **98**, 6326 (biosynth)

Leete, E. *et al.*, *Symp. Pap. - IUPAC Int. Symp. Chem. Nat. Prod.*, 11th, 1978, 1978, **1**, 155 (rev)

Luanratana, O. *et al.*, *Phytochemistry*, 1982, **21**, 449 (isol)

Génisson, Y. *et al.*, *Heterocycles*, 1994, **39**, 811 (synth)

Deo, N.M. *et al.*, *Tet. Lett.*, 1996, **37**, 1137 (synth)

Yang, C.-M. *et al.*, *Can. J. Chem.*, 1997, **75**, 616-620 (synth, ir, pmr, cmr, ms)


Felpin, F.-X. *et al.*, *J.O.C.*, 2001, **66**, 6305-6312 (synth)

3,4,5,6-Tetrahydro-2,3'-bipyridine, 9CI

T-144

Anabaseine

[3471-05-4]



C₁₀H₁₂N₂ 160.218

Toxin of the marine hoplonemertean worms *Paranemertes peregrina* and *Amphiporus angulatus*. Also isol. from the ants *Aphaenogaster* spp. and *Messor* spp. Nicotinic receptor agonist with selectivity for $\alpha 7$ subtype. Oil. Bp₁ 110-120°.


▶ DW1890000

Dipicrate: Mp 174°.

- Späth, E. *et al.*, *Ber.*, 1936, **69**, 1082 (*synth*)
 Kem, W.R. *et al.*, *Toxicol.*, 1971, **9**, 15-22; 23-32 (*isol, occur*)
 Kem, W.R. *et al.*, *Experientia*, 1976, **32**, 684
 Wheeler, J.W. *et al.*, *Science (Washington, D.C.)*, 1981, **211**, 1051-1052 (*isol, ants*)
 Zoltewicz, J.A. *et al.*, *Heterocycles*, 1993, **35**, 171-179 (*pharmacol, derivs*)
 Kem, W.R. *et al.*, *J. Pharmacol. Exp. Ther.*, 1997, **283**, 979-992 (*pharmacol*)
 Leclercq, S. *et al.*, *J. Chem. Ecol.*, 2001, **27**, 945-952 (*ants, isol*)
 Kem, W.R. *et al.*, *Mar. Drugs*, 2006, **4**, 255-273 (*rev*)

Tetrahydrocantlyne**T-145**

[60022-24-4]



Absolute Configuration

C₁₁H₁₇NO₃ 211.26

Alkaloid from the trunk bark of *Lasianthera austrocaledonica* and the aerial parts of *Scaevola racemigera* (Icacinaeae, Goodeniaceae). Also from several *Strychnos* spp. Amorph. [α]_D²⁰ +108 (c, 1.01 in CHCl₃).

O-(3-Pyridinecarbonyl): 6-O-Nicotinoyl-tetrahydrocantlyne

[99339-53-4]
 C₁₇H₂₀N₂O₄ 316.356

Alkaloid from *Scaevola racemigera* (Goodeniaceae) and (?) from *Strychnos cocculoides*, *Strychnos minfiensis*, *Strychnos mitis* and *Strychnos spinosa*. Noncryst. [α]_D²⁰ +59 (c, 1 in MeOH). In the 1997 review an isomeric struct. with an extra O atom is drawn. This struct. does not appear elsewhere in the lit. and appears to be an error.

O-(5-Vinyl-3-pyridinecarbonyl): 6-O-(5-Vinylnicotinoyl)tetrahydrocantlyne

[99339-54-5]
 C₁₉H₂₂N₂O₄ 342.394

Alkaloid from *Scaevola racemigera* (Goodeniaceae). Noncryst. [α]_D²⁰ +36 (c, 0.2 in MeOH).

1-Oxo: Strychnovoline

[96861-97-1]
 C₁₁H₁₅NO₄ 225.244

Alkaloid from the leaves of *Strychnos dinklagei*, from *Strychnos cocculoides*, *Strychnos longicauda*, *Strychnos mitis*, *Strychnos variabilis* and the aerial parts of *Scaevola racemigera* (Loganiaceae, Goodeniaceae). Needles (CHCl₃). Mp 177-178°. [α]_D²⁰ +98 (c, 1 in MeOH).

1-Oxo, O-(3-pyridinecarbonyl): 6-O-Ni-**cotinoylstrychnovoline**


[99339-52-3]
 C₁₇H₁₈N₂O₅ 330.34

Alkaloid from *Scaevola racemigera* (Loganiaceae). Noncryst. [α]_D²⁰ +24 (c, 0.1 in MeOH).

- Sévenet, T. *et al.*, *Phytochemistry*, 1976, **15**, 576 (*isol, uv, ir, pmr, ms, struct, synth*)
 Skaltsounis, A.-L. *et al.*, *Helv. Chim. Acta*, 1985, **68**, 1679-1685 (*Scaevola racemigera constits, abs config, synth*)
 Michel, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 86 (*Strychnovoline, isol, uv, ir, pmr, cmr, ms, struct, synth*)
 Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, Strychnos*)

1,2,3,4-Tetrahydro- β -carboline**T-146**

1,2,3,4-Tetrahydro-9H-pyrido[3,4-b]indole
 [16502-01-5]



C₁₁H₁₂N₂ 172.229
 Cryst. (EtOH). Mp 204-205°.

N²-(2-Methylaminobenzoyl): Goshuyamide I. Gosuiamide I


[126223-62-9]
 C₁₉H₁₉N₃O 305.379

Alkaloid from the fruits of *Evodia rutaecarpa* (Rutaceae). Prisms (C₆H₆/Me₂CO). Mp 178-180°.

- Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 610C (*nmr*)
 Späth, E. *et al.*, *Ber.*, 1930, **63**, 2102-2111 (*synth*)
 Org. Synth., 1971, **51**, 136-138 (*synth*)
 Shoji, N. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1160-1162 (*Goshuyamide I*)
 Yu, H.T. *et al.*, *J.A.C.S.*, 1995, **117**, 348-357 (*cryst struct*)
 Herráiz, T. *et al.*, *J. Agric. Food Chem.*, 2004, **52**, 2652-2658 (*occur*)

1,2,3,4-Tetrahydro- β -carboline-1,3-dicarboxylic acid**T-147**

2,3,4,9-Tetrahydro-1H-pyrido[3,4-b]indole-1,3-dicarboxylic acid, 9CI
 [59132-30-8]
 [192180-81-7, 192180-80-6]

(1*R*,3*S*)-form

C₁₃H₁₂N₂O₄ 260.249

(1*R*,3*S*)-form

Mp 269-271°. Probable config. One set of authors gives the Mp but does not assign a config., the other authors show that the main prod. of condensation of Tryptophan, T-640 with Glyoxylic acid is the *cis*-isomer, but do not give a Mp.

(1*ξ*,3*ξ*)-form

Present in various foods as Maillard prod. Cryst. Mp 270°.


[123771-30-2, 123771-45-9]

Bobbitt, J.M. *et al.*, *J.O.C.*, 1980, **45**, 1978-1984 (*synth*)

Gutsche, B. *et al.*, *J. Agric. Food Chem.*, 1997, **45**, 2458-2462 (*isol, pmr, ms, synth, config*)

Tetrahydrocorysamine**T-148**

6,7,12b,13-Tetrahydro-13-methyl-4H-bis[1,3]benzodioxolo[5,6-a:4',5'-g]quinolizine, 9CI



(+)form

C₂₀H₁₉NO₄ 337.374**(+)-form [32043-26-8]**

Alkaloid from *Corydalis pallida* var. *tenuis* (Papaveraceae). Mp 180-181°. [α]_D +236.

 α -N-Me: Tetrahydrocorysamine N-methosalt

[72203-88-4]
 C₂₁H₂₂NO₄⁺ 352.409

Quaternary alkaloid from *Corydalis cava* (Papaveraceae). Cryst. (MeOH) (as iodide). Mp 285-293° (iodide). CAS no. refers to iodide.

(±)-form [19775-54-3]


Mp 207-209°.

[42511-94-4, 42511-93-3]

- Tani, C. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 748 (*synth*)
 Kaneko, H. *et al.*, *Yakugaku Zasshi*, 1971, **91**, 101 (*isol*)
 Slávik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261 (*deriv*)
 Iwasa, K. *et al.*, *J.O.C.*, 1981, **46**, 4744 (*abs config*)
 Hanaoka, M. *et al.*, *Chem. Comm.*, 1985, 1257 (*synth*)

Tetrahydrocytisine**T-149**

[18161-94-9]



Probable Relative Configuration

C₁₁H₁₈N₂O 194.276

Stereochem., and stereochem. of derivs., not firmly established. Alkaloid from *Thermopsis chinensis* (Fabaceae). Bp₀ 160° (bath). [α]_D³⁰ +81 (EtOH).

N-Me: N-MethyltetrahydrocytisineC₁₂H₂₀N₂O 208.303

Alkaloid from *Ormosia indurata* seeds (Fabaceae). Cryst. (Me₂CO). Mp 210°.

N-Et: N-Ethyltetrahydrocytisine. Alkaloid XIC₁₃H₂₂N₂O 222.33

Alkaloid from leaves of *Laburnocytisus adamii* (not a generally recognised genus; a hybrid of *Laburnum* and

Chamaecytisus). Tentative identification based on ms.

N-(3-Butenyl): Tetrahydrorhombifoline

[3382-84-1]
C₁₅H₂₄N₂O 248.367

Alkaloid from *Lupinus angustifolius*, *Lupinus truncatus*, *Lupinus mutabilis* seeds and *Lupinus oscar-haughtii* (Fabaceae). Oil. Bp_{0.001} 160°. [α]_D³⁰ +81 (EtOH).

N-(3-Butenyl), hydrochloride: Mp 189°.

Hydroxy, N-(3-butenyl): Hydroxytetrahydrorhombifoline

[89248-47-5]
C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Castilleja miniata* (Scrophulariaceae). Probable struct. Posn. of OH-group and abs. config. unknown. No phys. props. reported.

11-Oxo, N-(3-butenyl): 11-Oxotetrahydrorhombifoline

C₁₅H₂₂N₂O₂ 262.351

Alkaloid from *Ormosia coutinhoi* bark (Fabaceae). Cryst. (Me₂CO/hexane). [α]_D²¹ +18 (c, 1.1 in MeOH). Shown to belong to the opposite enantiomeric series to Rhombifoline, i.e. opposite of that illus.

Bohlmann, F. *et al.*, *Chem. Ber.*, 1963, **96**, 2254 (*Tetrahydrorhombifoline*)

McLean, S. *et al.*, *Can. J. Chem.*, 1967, **45**, 751; 1971, **49**, 1976 (*11-Oxotetrahydrorhombifoline, N-Methyltetrahydrocytisine*)

Balandrin, M.F. *et al.*, *J. Nat. Prod.*, 1981, **44**, 495 (*Tetrahydrorhombifoline*)


McCoy, J.W. *et al.*, *J. Nat. Prod.*, 1983, **46**, 902-907 (*Hydroxytetrahydrorhombifoline*)

Saito, K. *et al.*, *CA*, 1990, **114**, 203603 (*Tetrahydrocytisine*)

Greinwald, R. *et al.*, *Biochem. Physiol. Pflanz.*, 1991, **187**, 385 (*N-Ethyltetrahydrocytisine*)

Tetrahydrodeoxycordifoline lactam

T-150



C₂₇H₃₀N₂O₁₀ 542.541

3α-form

5α-Carboxystrictosamide
[38495-77-1]

Alkaloid from *Adina rubescens* and the bark of *Nauclea diderrichii* (Rubiaceae). [α]_D -84 (CHCl₃) (as tetra-*O*-Ac, Me ester).

22-Deoxo: Sickingine

[160927-80-0]
C₂₇H₃₂N₂O₉ 528.558

Alkaloid from bark of *Sickingia tinctoria* and *Sickingia williamsii* (preferred genus name *Simira*) (Rubiaceae). Pale yellow amorph. powder. Mp 136-138°. [α]_D²⁵ -250 (c, 1 in MeOH).

3β-form

5α-Carboxyvincosamide
[38495-79-3]

Alkaloid from *Adina rubescens* (Rubiaceae). [α]_D -79 (CHCl₃) (as tetra-*O*-Ac, Me ester).


Blackstock, W.P. *et al.*, *Chem. Comm.*, 1972, 1006 (*uv, ms, cd, struct*)

Aquino, R. *et al.*, *Phytochemistry*, 1994, **37**, 1471 (*Sickingine*)

Lamidi, M. *et al.*, *Magn. Reson. Chem.*, 2005, **43**, 427-429 (*Nauclea diderrichii constit*)

5,6,7,8-Tetrahydro-5,7-dihydroxy-1*H*-azocin-2-one

T-151



C₇H₁₁NO₃ 157.169


Prod. by the marine *Streptomyces* sp. strain QD518.

Fotso, S. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 9-13 (*isol*)

1,2,3,4-Tetrahydro-6,7-dihydroxy-1,1-dimethylisoquinoline

T-152

1,2,3,4-Tetrahydro-1,1-dimethyl-6,7-isoquinolinediol
[55661-23-9]



C₁₁H₁₅NO₂ 193.245

Alkaloid from the leaves of *Aristolochia arcuata*. Amorph. brown solid (MeOH). [α]_D²⁵ +13 (c, 0.61 in Me₂CO). Dec. at 250°. λ_{max} 289 (log ε 3) (MeOH).

N-(β-D-Fructopyranos-6-yl): [556104-22-4]

C₁₇H₂₅NO₇ 355.387

Alkaloid from the leaves of *Aristolochia arcuata*. Amorph. brown solid (MeOH). [α]_D²⁵ -44 (c, 0.26 in MeOH). Dec. at 105°. λ_{max} 287 (log ε 3.4) (MeOH).

N-Et: 2-Ethyl-1,2,3,4-tetrahydro-6,7-dihydroxy-1,1-dimethylisoquinoline. 2-Ethyl-1,2,3,4-tetrahydro-1,1-dimethyl-6,7-isoquinolinediol
[556104-21-3]

C₁₃H₁₉NO₂ 221.299

Alkaloid from the leaves of *Aristolochia arcuata*. Amorph. brown solid (MeOH). [α]_D²⁵ +13 (c, 0.7 in MeOH). Dec. at 184°. λ_{max} 285 (log ε 3.5) (MeOH).

N-(2-Hydroxy-1-hydroxymethylethyl): 1,2,3,4-Tetrahydro-6,7-dihydroxy-2-(2-hydroxy-1-hydroxymethylethyl)-1,1-dimethylisoquinoline

[556104-20-2]


C₁₄H₂₁NO₄ 267.324

Alkaloid from the leaves of *Aristolochia arcuata*. Amorph. brown solid (MeOH). [α]_D²⁵ -11 (c, 0.72 in MeOH). Dec. at 174°. λ_{max} 286 (log ε 3.1) (MeOH).

Francisco, M.C. *et al.*, *Phytochemistry*, 2003, **62**, 1265-1270 (*isol, pmr, cmr*)

1,2,3,4-Tetrahydro-6,7-dihydroxy-1,1-dimethyl-3-isoquinoline-carboxylic acid, 9CI

T-153



C₁₂H₁₅NO₄ 237.255

(S)-form

Alkaloid from the seeds of *Mucuna pruriens*. Cryst. [α]_D³⁵ -155.3 (c, 0.07 in MeOH).

Misra, L. *et al.*, *Phytochemistry*, 2004, **65**, 2565-2567 (*isol, pmr, cmr, ms*)

1,2,3,4-Tetrahydro-7,8-dihydroxy-1,1-dimethyl-3-isoquinoline-carboxylic acid, 9CI

T-154

C₁₂H₁₅NO₄ 237.255

(S)-form


Alkaloid from the stem bark of *Mucuna pruriens*. Cryst. [α]_D³⁵ -144.2 (c, 0.01 in MeOH).

Misra, L. *et al.*, *Phytochemistry*, 2004, **65**, 2565-2567 (*isol, pmr, cmr, ms*)

1,2,3,4-Tetrahydro-5,8-dihydroxy-1-(4-hydroxybenzyl)-2-methylisoquinoline

T-155

1,2,3,4-Tetrahydro-1-[(4-hydroxyphenyl)methyl]-2-methyl-5,8-isoquinolinediol



C₁₇H₁₉NO₃ 285.342


(E)-form

4',8-Di-Me ether: 1,2,3,4-Tetrahydro-5-hydroxy-8-methoxy-1-(4-methoxybenzyl)-2-methylisoquinoline
C₁₉H₂₃NO₃ 313.396

Alkaloid from the aerial parts of *Sabia parviflora*. Needles. Mp 87-89°. λ_{max} 275 (no solvent reported).

Chen, J. *et al.*, *Chin. Chem. Lett.*, 2002, **13**, 426-427 (*isol, pmr*)

4,5,6,7-Tetrahydro-4,5-dihydroxy-1*H*-indole-3-carboxaldehyde T-156
3-Formyl-4,5,6,7-tetrahydro-4,5-dihydroxy-1*H*-indole



C₉H₁₁NO₃ 181.191

(4*R,5*R**)-form**


5-*Ac*: [501373-69-9]

C₁₁H₁₃NO₄ 223.228

Isol. from a Puerto Rican *Lyngbya majuscula*. Amorph. solid. $[\alpha]_D^{25} +104.6$ (c, 0.5 in CHCl₃). λ_{\max} 214 (log ϵ 4.39); 254 (log ϵ 4.04); 287 (log ϵ 3.67) (MeOH).

Nogle, L.M. et al., *J. Nat. Prod.*, 2003, **66**, 217-220 (isol, pmr, cmr, ms)

4,5,6,7-Tetrahydro-6,7-dihydroxy-1*H*-indol-4-one T-157



Relative Configuration

C₈H₉NO₃ 167.164

(6*R,7*R**)-form**


(-)-*trans*-form

Prod. by *Nocardia* sp. FH-A 1527. Oil. $[\alpha]_D^{20} -185.6$ (c, 0.4 in MeOH). λ_{\max} 208 (ϵ 9500); 243 (ϵ 4400); 276 (ϵ 3600) (MeOH).

Henne, P. et al., *Nat. Prod. Lett.*, 1997, **10**, 43-47 (isol, uv, ir, cd, pmr, cmr)

1,2,3,4-Tetrahydro-6,7-dihydroxyisoquinoline T-158

1,2,3,4-Tetrahydro-6,7-isoquinolinediol, 9*CI*
[34827-33-3]



C₉H₁₁NO₂ 165.191

Direct and indirect acting sympathomimetic agent.

Hydrochloride: [42887-47-8]

Cryst. (HCl aq. or EtOH). Mp 262° (254-256°).

Hydrobromide: [52768-23-7]

Cryst. (EtOH/Et₂O aq.). Mp 267-268°.

N-Me: 1,2,3,4-Tetrahydro-2-methyl-6,7-isoquinolinediol, 9*CI*
[37491-98-8]

[52553-18-1 (hydrobromide)]

C₁₀H₁₃NO₂ 179.218

Occurs in cerebrospinal fluid of

humans. May be implicated in Parkinsons disease.

N-Me, hydrochloride: [63937-92-8]

Cryst. Mp 228-230°.

N,N-Di-Me:

[57553-19-2 (bromide)]

C₁₁H₁₆NO₂ 194.253

Cryst. (EtOH) (as bromide). Mp 285° (263-266°) (bromide).

6-*Me ether*: 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxyisoquinoline. 1,2,3,4-Tetrahydro-6-methoxy-7-isoquinolinol.

Norcorypalline

[1011-42-3]

C₁₀H₁₃NO₂ 179.218

Poss. alkaloid from the stem bark of *Zizyphus rugosa*. Cryst. (C₆H₆/EtOH)(synthetic); granules (MeOH) (nat.). Mp 197-199° (182-183°)(synth.) Mp 283-285° (nat.). Subl. 1.01 140. Tentative identification. The wide discrepancy in Mps. makes the struct. unlikely.

6-*Me ether, hydrochloride*: [1078-26-8]
Cryst. (EtOH). Mp 256-258° (246-248°).

6-*Me ether, picrate*:

Cryst. (EtOH aq.). Mp 252-254°.

6-*Me ether, N-Me*: 1,2,3,4-Tetrahydro-6-methoxy-2-methyl-7-isoquinolinol.

1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-methylisoquinoline. **Corypalline** [450-14-6]

C₁₁H₁₅NO₂ 193.245

Alkaloid from *Corydalis pallida*, *Corydalis aurea*, *Corydalis ophiocarpa*, *Thalictrum dasycarpum*, *Thalictrum rugosum*, *Thalictrum uchiyamai*, *Doryphora sassafras*, *Islaya minor* (preferred genus name *Neoporteria*) and *Papaver bracteatum*. Cryst. (CHCl₃, C₆H₆ or MeOH/Et₂O). Mp 171-173° (164-165°). λ_{\max} 227 (log ϵ 3.79); 286 (log ϵ 3.58) (EtOH). λ_{\max} 225 (log ϵ 3.79); 286 (log ϵ 3.56) (EtOH/HCl). λ_{\max} 247 (log ϵ 3.83); 302 (log ϵ 3.66) (EtOH/NaOH).

6-*Me ether, N-Me, picrate*: Mp 178°.

6-*Me ether, N,N-di-Me*: **N-Methylcorypalline**. **N-Methylcorypallinium**

[23594-91-4]

C₁₂H₁₈NO₂[⊕] 208.28

Quaternary alkaloid from *Corydalis stricta* (Papaveraceae). Mp 238-239° (as iodide).

7-*Me ether*: 1,2,3,4-Tetrahydro-6-hydroxy-7-methoxyisoquinoline. 1,2,3,4-Tetrahydro-7-methoxy-6-isoquinolinol

[1011-43-4]

[1078-27-9]

C₁₀H₁₃NO₂ 179.218

Cryst. (EtOH/Et₂O) (as hydrochloride). Mp 260-263° (247-249°) (hydrochloride).

7-*Me ether, N-Me*: 1,2,3,4-Tetrahydro-7-methoxy-2-methyl-6-isoquinolinol.

1,2,3,4-Tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline. **Isocorypalline**

[13871-59-5]

C₁₁H₁₅NO₂ 193.245

Alkaloid from *Berberis oblonga* and

Corydalis stricta. Cryst. (C₆H₆). Mp 163-164°. λ_{\max} 227 (log ϵ 3.55); 286 (log ϵ 3.44) (EtOH). λ_{\max} 227 (log ϵ 3.55); 286 (log ϵ 3.44) (EtOH/HCl). λ_{\max} 247 (log ϵ 3.69); 301 (log ϵ 3.55) (EtOH/NaOH).

7-*Me ether, N-Me, hydrochloride*: [30765-63-0]
Cryst. (EtOH). Mp 285-290° (260°-263°).

7-*Me ether, N-Me, picrate*:

Cryst. (H₂O). Mp 168°.

Di-Me ether: 1,2,3,4-Tetrahydro-6,7-dimethoxyisoquinoline, 9*CI*. **Heliamine**

[1745-07-9]

C₁₁H₁₅NO₂ 193.245

Alkaloid from *Pachycereus weberi*, *Pachycereus pringlei*, *Pachycereus pecten-aboriginum*, *Backebergia militaris* (preferred genus name *Pachycereus*) and *Carnegiea gigantea* (Cactaceae). Antineoplastic agent. Inhibitor of rat sarcoma 45. Mp 84-85°. Log P 0.98 (calc). Readily forms carbonate on exp. to air.

▶ NX5015000

Di-Me ether, hydrochloride: [2328-12-3]
Cryst. (EtOH/Et₂O). Mp 255° (248°, 252°).

Di-Me ether, picrate:

Cryst. (EtOH). Mp 223-225°.

Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methylisoquinoline. **O-Methylcorypalline**. **N-Methylheliamine**

[16620-96-5]

C₁₂H₁₇NO₂ 207.272

Alkaloid from *Pachycereus weberi*, *Thalictrum dioicum*, *Thalictrum polygamum*, *Backebergia militaris*, *Nehumbo nucifera* (East Indian lotus), *Pilosocereus guerreronis* and *Papaver bracteatum* (Cactaceae, Ranunculaceae, Nelumbonaceae, Papaveraceae). Cryst. (MeOH or petrol/cyclohexane), cryst. + 1/2 H₂O (Et₂O). Mp 69-70° Mp 82° (hemihydrate). λ_{\max} 290 (sh) (log ϵ 3.99); 295 (log ϵ 3.47) (no solvent reported).

▶ NX5018510

Di-Me ether, N-Me, hydrochloride:

[16135-43-6]

Cryst. (2-propanol). Mp 219-221° (215°, 210°).

Di-Me ether, N-Me, picrate: [28332-96-9]
Cryst. (C₆H₆/MeOH). Mp 160°.

Di-Me ether, N,N-di-Me: 1,2,3,4-Tetrahydro-6,7-dimethoxy-2,2-dimethylisoquinolinium (1+). **N-Dimethylheliamine**

[83527-64-4, 19626-05-2]

C₁₃H₂₀NO₂[⊕] 222.307

Quaternary alkaloid from *Berberis amurensis* (Berberidaceae). Mp 242-243° (as iodide).

Di-Et ether: 6,7-Diethoxy-1,2,3,4-tetrahydroisoquinoline

[52759-05-4]

[63905-65-7]

C₁₃H₁₉NO₂ 221.299

Flakes (as hydrochloride). Mp 268° (hydrochloride).


Buck, J.S. et al., *J.A.C.S.*, 1934, **56**, 1769-1771 (synth. *N-Me*, 6,7-*di-Me*, 6,7-*di-Me N-Me*)

- Manske, R.H.F. *et al.*, *Can. J. Res., Sect. B*, 1937, **15**, 159-167 (*Corypalline, occur, struct, synth*)
- Corrodi, H. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 911-918 (*synth*)
- Bobbitt, J.M. *et al.*, *J.O.C.*, 1965, **30**, 2247-2250; 1967, **32**, 2225-2227; 1968, **34**, 2001-2002 (*Corypalline, Me ethers, N-Me, synth*)
- Kupchan, S.M. *et al.*, *J.O.C.*, 1969, **34**, 1062-1065 (*Corypalline, isol*)
- Yang, T.-H. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1970, **17**, 54-56; 235-242 (*O-Methylcorypalline, isol, struct*)
- Chen, C.R. *et al.*, *J. Nat. Prod.*, 1974, **37**, 493-500 (*Corypalline, isol, uv*)
- Simpson, L.L. *et al.*, *J. Pharmacol. Exp. Ther.*, 1975, **192**, 365-371 (*activity*)
- Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 530-531; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 563 (*Isocorypalline*)
- Wood, G.W. *et al.*, *Anal. Chem.*, 1976, **48**(7), 981-984 (*ms*)
- Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252-2260; 1977, **55**, 3304-3311 (*cmr*)
- Von Weigrebe, W. *et al.*, *Helv. Chim. Acta*, 1976, **59**, 949-962 (*O-Methylcorypalline, synth*)
- Smissman, E.E. *et al.*, *J. Med. Chem.*, 1976, **19**, 127-131 (*N-Me, 6,7-di-Me ether, 6,7-di-Me ether N-Me, synth*)
- Lindgren, J.E. *et al.*, *J. Nat. Prod.*, 1976, **39**, 464-466 (*N-Me, occur, ms*)
- Mary, L.J.F. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 182-183 (*Corypalline, synth, uv*)
- Strömbom, J. *et al.*, *Acta Pharm. Suec.*, 1978, **15**, 127-132; *CA*, **89**, 126107w (*isol*)
- Strömbom, J. *et al.*, *J. Chromatogr.*, 1978, **147**, 513-515 (*O-Methylcorypalline, hplc*)
- Shamma, M. *et al.*, *J. Nat. Prod.*, 1978, **41**, 169-178 (*O-Methylcorypalline, isol, uv, ir, pmr, ms*)
- Miller, L.L. *et al.*, *J.O.C.*, 1978, **43**, 1580-1586 (*synth, pmr, ms*)
- Pandey, G.D. *et al.*, *Indian J. Chem., Sect. B*, 1979, **18**, 544-545 (*Corypalline, synth, pmr, ms*)
- Irie, H. *et al.*, *Chem. Lett.*, 1980, 875-878 (*Corypalline, synth*)
- Doetsch, P.W. *et al.*, *J. Chromatogr.*, 1980, **189**, 79-85 (*Corypalline, occur*)
- Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673-678; 1983, **22**, 1263-1270 (*Heliamine, N-Methylheliamine, isol, ir, uv, pmr, cmr, ms*)
- Pummangura, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 277-282 (*Heliamine, isol*)
- Irgashev, T. *et al.*, *Khim. Prir. Soedin.*, 1983, **19**, 490-493; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **19**, 461-463 (*N-Methylcorypalline, Isocorypalline*)
- Theuns, H.G. *et al.*, *Phytochemistry*, 1983, **22**, 247-250 (*Corypalline, Isocorypalline, O-Methylcorypalline, occur, synth, pmr, ms*)
- Ruchirawat, S. *et al.*, *Synth. Commun.*, 1984, **14**, 1221-1228 (*synth, pmr, ms*)
- Bates, H.A. *et al.*, *J.O.C.*, 1986, **51**, 3061-3063 (*N-Me, synth*)
- Pandey, V.B. *et al.*, *Phytochemistry*, 1988, **27**, 1915-1916 (*Norcorypalline, isol*)
- Niwa, T. *et al.*, *Biochem. Biophys. Res. Commun.*, 1991, **177**, 603-609 (*N-Me, isol*)
- Moser, A. *et al.*, *Life Sci.*, 1992, **50**, 1885-1891 (*N-Me, isol*)
- Takano, S. *et al.*, *Heterocycles*, 1993, **35**, 47-52 (*synth*)
- Walpole, C.S.J. *et al.*, *J. Med. Chem.*, 1994, **37**, 1942-1954 (*synth, pmr*)
- Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 410-512; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 386-512 (*N-Dimethylheliamine*)
- Shirahita, A. *et al.*, *Chem. Pharm. Bull.*, 1997, **45**, 1814-1819 (*synth, detm*)
- Cheng, X. *et al.*, *Chem. Biodiversity*, 2008, **5**, 1335-1344 (*Corypalline, pmr, cmr*)

1,2,3,4-Tetrahydro-7,8-dihydroxyisoquinoline

T-159

1,2,3,4-Tetrahydro-7,8-isoquinolinediol,
9CI
[102830-09-1]

C₉H₁₁NO₂ 165.191

7-Me ether, N-Me: 1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-2-methylisoquinoline. **Turcamine**

[28026-17-7]
C₁₁H₁₅NO₂ 193.245

Alkaloid from *Berberis turcomanica* (Berberidaceae) and *Ceratocarpus palaestinus* (Papaveraceae). Cryst. (Et₂O) petrol. Mp 102°.

7-Me ether, N-Me; hydrochloride: [19462-72-7]
Cryst. (EtOH). Mp 220°.

8-Me ether, N-Me: 1,2,3,4-Tetrahydro-7-hydroxy-8-methoxy-2-methylisoquinoline

[87339-94-4]
C₁₁H₁₅NO₂ 193.245
Cryst. (2-propanol). Mp 160°.

8-Me ether, N-Me; hydrochloride:
Cryst. (2-propanol/EtOH). Mp 265°.

Di-Me ether: 1,2,3,4-Tetrahydro-7,8-dimethoxyisoquinoline. **Lemaireocereine**

[52759-08-7]
[15365-56-7]
C₁₁H₁₅NO₂ 193.245

Alkaloid from *Pachycereus weberi* and *Backebergia militaris* (preferred genus name *Pachycereus*). Also identified in trace amount in *Pachycereus pringlei* (Cactaceae). Cryst. (EtOH/Et₂O)(as hydrochloride). Mp 198° (185°)(hydrochloride).

Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-7,8-dimethoxy-2-methylisoquinoline

[65644-75-9]
Mp 173-174° (as hydrochloride).

[102830-10-4, 27943-55-1]

Bobbitt, J.M. *et al.*, *J.O.C.*, 1969, **34**, 2001-2002 (*synth*)

Grethe, G. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 873-881 (*synth*)

Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673-678; 1983, **22**, 1263-1270 (*Lemaireocereine, isol, uv, ir, pmr, ms, cmr*)

Mata, R. *et al.*, *Planta Med.*, 1980, **38**, 180-182 (*Lemaireocereine, occur*)

Pummangura, S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 498-499 (*Lemaireocereine, isol*)

Dwuma-Badu, D. *et al.*, *J. Nat. Prod.*, 1983, **46**, 342-349 (*di-Me ether N-Me, synth*)

Theuns, H.G. *et al.*, *Phytochemistry*, 1983, **22**, 247-250 (*derivs*)

Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1990, **53**, 1006-1008 (*7-Me ether N-Me*)

Ishii, H. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 1993-1996 (*synth*)


Khamidov, I.I. *et al.*, *Khim. Prir. Soedin.*, 1996, **32**, 894-896; *Chem. Nat. Compd. (Engl. Transl.)*, 1996, **32**, 880-881 (*Turcamine*)

Shinohara, T. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 918-927 (*synth*)

1,2,3,4-Tetrahydro-6,7-dihydroxy-3-isoquinolinecarboxylic acid, 9CI

T-160

[41844-06-8]

C₁₀H₁₁NO₄ 209.201**(S)-form** [34312-81-7]

Alkaloid from the seeds of *Mucuna mutisiana* (Fabaceae). Also present in *Mucuna urens*, *Mucuna deeringiana*, *Mucuna andreana*, *Mucuna holtoni*, *Mucuna pruriens*, *Mucuna sloanei* and *Stizolobium hassjoo* (Fabaceae). Mp 286-288° (natural) Mp 293-294° dec. (synthetic). [α]_D²⁵ -114.9 (c, 1.65 in 20% HCl) (natural). [α]_D²⁵ -125.91 (c, 1 in 1M HCl) (synthetic).

(±)-form

Synthetic. Needles (H₂O). Mp 277°. No CAS No. found 8-14CI.

Shah, R.J. *et al.*, *J.O.C.*, 1961, **26**, 3533-3554 (*synth, ir*)

Bell, E.A. *et al.*, *Phytochemistry*, 1971, **10**, 2191-2194 (*isol, uv, ir, pmr, ms, synth*)

Brossi, A. *et al.*, *Helv. Chim. Acta*, 1972, **55**, 15-21 (*synth, pmr*)

Saito, K. *et al.*, *Phytochemistry*, 1982, **21**, 474-476 (*isol, synth*)


Misra, L. *et al.*, *Phytochemistry*, 2004, **65**, 2565-2567 (*isol, cmr*)

1,2,3,4-Tetrahydro-4,8-dihydroxy-2-methylisoquinoline

T-161

1,2,3,4-Tetrahydro-2-methyl-4,8-isoquinolinediol, 9CI. **Longimammamine**

[57236-57-4]

C₁₀H₁₃NO₂ 179.218**(±)-form**

Alkaloid from *Dolichothele longimamma* and *Dolichothele uberiformis* (Cactaceae). Cryst. (EtOH/Et₂O)(as hydrochloride). Mp 235-236.5° (224-228°)(as hydrochloride).

Ranieri, R.L. *et al.*, *J.O.C.*, 1976, **41**, 319 (*isol, uv, ir, pmr, ms, struct, synth*)


Ranieri, R.L. *et al.*, *J. Nat. Prod.*, 1977, **40**, 173 (*isol*)

1,2,3,4-Tetrahydro-6,7-dihydroxy-1-methylisoquinoline

T-162

1,2,3,4-Tetrahydro-1-methyl-6,7-isoquinolinediol, 9CI. **Salsolinol**

[525-72-4]

**(S)-form**

C₁₀H₁₃NO₂ 179.218**(R)-form** [53622-83-6]Synthetic. Mp 174-174° (as hydrobromide). [α]_D²⁰ +30 (MeOH) (hydrobromide).**6-Me ether:** see 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-1-methylisoquinoline, T-181**7-Me ether:** see 1,2,3,4-Tetrahydro-6-hydroxy-7-methoxy-1-methylisoquinoline, T-179**Di-Me ether:** see 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methylisoquinoline, T-167**(S)-form** [27740-96-1]Alkaloid from the leaves of *Aristolochia arcuata*. False neurotransmitter. Potent inhibitor of dopamine uptake into rat brain. Catecholamine O-methyltransferase inhibitor. Cryst. (EtOH/Et₂O) (as hydrobromide). Mp 174-175° (hydrobromide). [α]_D²⁰ -30.9 (MeOH).**N-(1-Deoxy-β-D-fructopyranos-1-yl):** [556799-99-6]C₁₆H₂₃NO₇ 341.36Alkaloid from the leaves of *Aristolochia arcuata*. Amorph. brown solid (MeOH). [α]_D²⁵ -45 (c, 1.9 in MeOH). Dec. at 105°. λ_{max} 290 (log ε 2.9) (MeOH).**(±)-form****Me ether, N-Me: O-Demethylcarnegine. O-Norcarnegine**

[176329-46-7]

C₁₂H₁₇NO₂ 207.272Alkaloid from *Echium humile*. Oil. Site of methylation not known. Identical with either (±)-N-Methylsalsoline or (±)-N-Methylisalsoline.**(±)-form**Alkaloid from *Annona reticulata* (custard apple), *Musa paradisiaca* (banana), *Theobroma cacao* (cocoa) and *Aconitum carmichaeli* (Annonaceae, Ranunculaceae, Musaceae, Sterculiaceae).Teitel, S. et al., *J. Med. Chem.*, 1972, **15**, 845 (synth, uv, cd, ord, pmr, abs config)Riggin, R.M. et al., *J. Agric. Food Chem.*, 1976, **24**, 189; 900 (occur)Forgacs, P. et al., *Plant. Med. Phytother.*, 1981, **15**, 10 (occur)Chen, D. et al., *Yaoxue Xuebao*, 1982, **17**, 792; *CA*, **98**, 40465kEl-Shazly, A. et al., *Phytochemistry*, 1996, **42**, 225 (O-Demethylcarnegine)Francisco, M. et al., *Phytochemistry*, 2003, **62**, 1265-1270 (isol, pmr, cmr)Zhao, S.-L. et al., *Chin. J. Chem.*, 2006, **24**, 439-441 (anal, resoln)**1,2,3,4-Tetrahydro-7,8-dihydroxy-1-methylisoquinoline** T-163**1,2,3,4-Tetrahydro-1-methyl-7,8-isoquinolinediol, 9CI. Isosalsolinol**

[53405-13-3]


[102917-28-2]

C₁₀H₁₃NO₂ 179.218**(±)-form**Alkaloid from the roots of *Mirabilis jalapana*.**7-Me ether:** 1,2,3,4-Tetrahydro-7-methoxy-1-methyl-8-isoquinolinol. 1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-1-methylisoquinoline. **Arizonine**

[60508-83-0]

C₁₁H₁₅NO₂ 193.245Alkaloid from *Carnegiea gigantea* and *Pachycereus pecten-aboriginum* (Cactaceae). Cryst. (MeOH/Et₂O) (as salicylate). Mp 207-209° (salicylate).Bruhn, J.G. et al., *J. Nat. Prod.*, 1976, **39**, 197 (isol, pmr, ms, struct, synth)Strömbom, J. et al., *Acta Pharm. Suec.*, 1978, **15**, 127-132 (Arizonine)Kaufman, T.S. et al., *Synth. Commun.*, 1992, **22**, 1913 (synth)Suau, R. et al., *Tet. Lett.*, 1995, **36**, 1315 (synth)Yi-Fen, W. et al., *Helv. Chim. Acta*, 2002, **85**, 2342-2348 (isol)**1,2,3,4-Tetrahydro-6,7-dihydroxy-1-methyl-3-isoquinolinecarboxylic acid, 9CI** T-164


[52618-26-5]

C₁₁H₁₃NO₄ 223.228**(1S,3S)-form** [35287-23-1]Alkaloid from the seeds of *Mucuna deeringiana* and from the seeds and callus of *Stizolobium hassjoo*. Antioxidant. Cryst. (20% AcOH). Mp 240-269° dec. (natural) Mp 288-291° dec. (synthetic). [α]_D²⁰ -157 (c, 1 in 1M HCl) (synthetic).**Et ester:** Mp 161-162°. [α]_D²⁰ -133.3 (c, 1 in MeOH).**(1RS,3RS)-form**



Cryst. Mp 270-275° dec.

Müller, P. et al., *Biochem. Physiol. Pflanz.*, 1971, **162**, 234 (biosynth)Brossi, A. et al., *Helv. Chim. Acta*, 1972, **55**, 15 (synth, pmr, ord)Daxenbichler, M.E. et al., *Tet. Lett.*, 1972, 1801 (isol, synth, ms, pmr, struct)Teitel, S. et al., *J. Nat. Prod.*, 1974, **37**, 196 (cd, pmr, cryst struct)Bruderer, H. et al., *Helv. Chim. Acta*, 1975, **58**, 795 (ester, synth, pmr)Saito, K. et al., *Phytochemistry*, 1982, **21**, 474 (isol, pmr, cmr, ms, synth)Misra, L. et al., *Phytochemistry*, 2004, **65**, 2565-2567 (isol, cmr)Shiraiwa, T. et al., *Chem. Pharm. Bull.*, 2005, **53**, 1197-1199 (synth, pmr, cmr, resoln)**5,5a,12,12a-Tetrahydro-7,14-dimethoxybenzo[1''',2''':4,5;4''',5''':4',5']difuro[2,3-b:2',3'-b']diindole-7b,14b-diol, 9CI** T-165

[78279-84-2]

C₂₄H₂₀N₂O₆ 432.432Metab. of *Aspergillus terreus* var. *africanus*. Needles (C₆H₆). Mp 221-223°. [α]_D¹⁹ +39 (c, 0.11 in dioxan).Arai, K. et al., *Chem. Pharm. Bull.*, 1981, **29**, 1005 (uv, pmr, struct)**1,2,3,4-Tetrahydro-7,8-dimethoxy-1,2-dimethylisoquinoline, 9CI** T-166**Tepepine**

[34319-92-1]

C₁₃H₁₉NO₂ 221.299Alkaloid from *Pachycereus tehuanepicanus* (Cactaceae). Details of isol. and struct. elucidation have not been published.Kapadia, G.J. et al., *Chem. Comm.*, 1970, 856 (occur, synth)**1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methylisoquinoline Salsolidine. O-Methylsalsoline. N-Norcarnegine** T-167C₁₂H₁₇NO₂ 207.272**(R)-form** [54193-08-7]Alkaloid from branches and fruits of *Genista purgans* (Fabaceae), also obt. by resoln. Mp 47.5-48.5° (71-73° from H₂O). [α]_D¹⁶ +59.9 (EtOH).**Hydrochloride:** Mp 235-236°. [α]_D¹⁸ +25.3 (H₂O).**Picrate:** Mp 193-194° dec.**N-Me:** [51745-28-9]Alkaloid from the callus of *Nandina domestica*. Oil. [α]_D¹⁹ +24.6 (c, 3 in EtOH).**(S)-form** [493-48-1]Alkaloid from *Salsola richteri* and *Salsola arbuscula* (Chenopodiaceae). Also obt. by resoln. Plates (H₂O). Mp 47.5-48.5° (anhyd.), 71-73° (dihydrate). [α]_D²⁸ -59 (c, 0.5 in EtOH).

▶ NX5018500

Hydrochloride:Plates (H₂O). Mp 239-241° (235-236°). [α]_D¹⁸ -24.8 (H₂O).**Picrate:**

Yellow cryst. (EtOH). Mp 193-194° dec.

N-Benzoyl: Mp 124-125°.**N-Me:** [38221-25-9]Synthetic. Oil. [α]_D¹⁸ -24.4 (c, 9 in EtOH).**N-Me, picrate:** Mp 233-234° (229-232°).**(±)-form** [38520-68-2]Alkaloid from *Genista purgans*,

Carnegiea gigantea, *Pachycereus pectenaboriginum*, *Bienertia cycloptera*, *Corispermum leptopyrum*, *Salsola arbuscula*, *Salsola kali*, *Salsola richteri*, *Salsola ruthenica*, *Salsola soda*, *Alhagi pseudalhagi*, *Calicotome spinosa*, *Desmodium tiliaefolium* and *Desmodium cephalotes* (Fabaceae, Cactaceae, Chenopodiaceae). Antihypertensive agent. Mp 53-53.5°. Bp₁ 140°.

Hydrochloride: Mp 196-197°.

Picrate: Mp 200-201°.

N-Benzoyl: Mp 127-128°.

N-Me: 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethylisoquinoline. **Carnegine**. *Pectenine*†. *N-Methylsalsolidine* [490-53-9]

Alkaloid from *Carnegiea gigantea*, *Haloxylon salicornicum*, *Haloxylon articulatum* and *Cereus pectenaboriginum* (*Pachycereus pectenaboriginum*) (Cactaceae, Chenopodiaceae). Convulsive agent. Oil. Bp₁ 170°.

► Toxic.

N-Me, hydrochloride: Mp 209-211°.

N-Me, hydrobromide: Mp 228°.

N-Me, picrate: Mp 213-215°.

Proskurnina, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1937, **4**, 1265-1271 (*isol*)

Späth, E. *et al.*, *Ber.*, 1938, **71**, 113-119 (*synth, ir, uv*)

Barca, R. *et al.*, *An. R. Soc. Esp. Fis. Quim.*, 1959, **55B**, 731; *CA*, **54**, 14289g (*isol*)

Battersby, A.R. *et al.*, *J.C.S.*, 1960, 1214-1221 (*config, bibl*)

Carling, C. *et al.*, *Acta Pharm. Suec.*, 1970, **7**, 285; *CA*, **73**, 63154f (*isol*)

Teitel, S. *et al.*, *J. Med. Chem.*, 1972, **15**, 845-846; 1974, **17**, 134-136 (*synth*)

Brown, S.D. *et al.*, *J.O.C.*, 1972, **37**, 1825 (*Carnegine, isol, ir, pmr, ms, synth*)

Ghosal, S. *et al.*, *Phytochemistry*, 1973, **12**, 193-197; 1974, **13**, 1628-1629 (*isol, uv, ms, pmr, occur*)

Kagan, H.B. *et al.*, *J. Organomet. Chem.*, 1975, **90**, 353 (*synth*)

Erhardt, P.W. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 53-62 (*synth, resoln, pharmacol, bibl*)

Middleditch, B.S. *et al.*, *J. Chromatogr.*, 1976, **126**, 581-589 (*ms*)

Bruhn, J.G. *et al.*, *J. Nat. Prod.*, 1976, **39**, 175-177; 197-203 (*Carnegine, isol*)

Drost-Karbowska, K. *et al.*, *Acta Pol. Pharm.*, 1977, **34**, 421; *CA*, **88**, 117814b (*isol*)

Cymerman Craig, J. *et al.*, *J.A.C.S.*, 1977, **99**, 7996-8002 (*uv, cd*)

Kametani, T. *et al.*, *J.C.S. Perkin I*, 1977, 579-581 (*synth, pmr, ir*)

Singh, S.P. *et al.*, *J. Het. Chem.*, 1978, **15**, 541-544 (*cmr*)

Pummangura, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 277-282 (*S-form, isol, ir, pmr, struct*)

Mata, R. *et al.*, *Photochemistry*, 1983, **22**, 1263-1270 (*cmr*)

Yamato, M. *et al.*, *Tet. Lett.*, 1988, **29**, 6949-6950 (*R-form, S-form, synth*)

Pyne, S.G. *et al.*, *J.O.C.*, 1990, **55**, 1086-1093 (*R-form, N-Me, synth*)

Comins, D.L. *et al.*, *Heterocycles*, 1991, **32**, 1869-1873 (*synth*)

Coppola, G.M. *et al.*, *J. Het. Chem.*, 1991, **28**, 1769-1772 (*synth, pmr, cmr*)

Org. Synth., *Coll. Vol.*, **8**, 1993, 573-577 (*S-form, synth*)

Lee, A.W.M. *et al.*, *J.C.S. Perkin I*, 1994, 477-481 (*R-form N-Me, synth*)

Chan, W. *et al.*, *Tet. Lett.*, 1995, **36**, 715-718 (*S-form, synth*)

Ponzo, V.L. *et al.*, *Tet. Lett.*, 1995, **36**, 9105-9108 (*S-form, synth*)

Wirth, T. *et al.*, *Synthesis*, 1998, 162-166 (*S-form, synth, pmr, cmr*)

Ukaji, Y. *et al.*, *Bull. Chem. Soc. Jpn.*, 2000, **73**, 447-452 (*S-form, synth, ir, pmr, ms*)

Taniyama, D. *et al.*, *Tet. Lett.*, 2000, **41**, 5533-5536 (*synth*)

Adam, S. *et al.*, *J.O.C.*, 2001, **66**, 8744-8750 (*R-form, synth, pmr, cmr*)

Bracca, A.B.J. *et al.*, *Tetrahedron*, 2004, **60**, 10575-10610 (*rev, Carnegine, synth*)

Werner, F. *et al.*, *Eur. J. Org. Chem.*, 2007, 3911-3915 (*R-form, synth, pmr, cmr*)


Grajewska, A. *et al.*, *Tetrahedron: Asymmetry*, 2007, **18**, 557-561 (*synth*)

Iwasa, K. *et al.*, *J. Nat. Prod.*, 2008, **71**, 1376-1385 (*Carnegine*)

1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-phenylisoquinoline, 9CI T-168

[4008-69-9]

[136904-97-7]



C₁₈H₂₁NO₂ 283.369

(±)-*form* [54329-12-3]

Cryst. (hexane). Mp 82-83°.

N-Oxide: [74675-30-2]

Cryst. (EtOAc). Mp 115-118°.

(ξ)-*form*

Alkaloid from *Adhatoda vasica*.

[16135-40-3]

Kametani, T. *et al.*, *J. Het. Chem.*, 1965, **2**, 222 (*synth*)

Bremner, J.B. *et al.*, *Aust. J. Chem.*, 1980, **33**, 833-841 (*synth*)


Barker, J.M. *et al.*, *J.C.S. Perkin I*, 1985, 275-281 (*synth*)

Minor, D.L. *et al.*, *J. Med. Chem.*, 1994, **37**, 4317-4328 (*synth*)

Hashem, F.A.E.-M. *et al.*, *Pharm. Pharmacol. Lett.*, 1998, **8**, 167-169 (*isol*)

1,3,4,9-Tetrahydro-1,1-dimethylpyrano[3,4-b]indole T-169

[42821-17-0]



C₁₃H₁₅NO 201.268

Alkaloid from *Monascus pilosus*. Cryst. (Et₂O/C₆H₆) (nat. alkaloid descr. as oil). Mp 142-144°. λ_{max} 288 (log ε 3.6) (MeOH).

Humber, L.G. *et al.*, *Eur. J. Med. Chem.*

(*Chim. Ther.*), 1975, **10**, 215-220 (*synth*)


Demerson, C.A. *et al.*, *Can. J. Chem.*, 1979, **57**, 3296-3300 (*synth*)

Zhang, X. *et al.*, *Org. Lett.*, 2005, **7**, 2043-2046 (*synth*)

Cheng, M.J. *et al.*, *Chem. Pharm. Bull.*, 2008, **56**, 394-397 (*isol, pmr, cmr, ms*)

5,6,7,8-Tetrahydro-2,4-dimethylquinoline T-170

[60169-66-6]



C₁₁H₁₅N 161.246

Alkaloid from the roots of *Glycyrrhiza uralensis* (Chinese licorice). Mp 19-20°. Bp 250-251° Bp₁₂ 122-123°.

Hydrochloride: Mp 195°.

Picrate: Mp 144°.

Methiodide: Mp 157°.

v. Braun, J. *et al.*, *Ber.*, 1923, **56**, 1346 (*synth*)
Yamaguchi, K. *et al.*, *Yakugaku Zasshi*, 1926, **533**, 53 (*synth*)

Shaw, J.E. *et al.*, *J. Het. Chem.*, 1987, **24**, 1477 (*synth*)

Han, Y.N. *et al.*, *Arch. Pharmacol. Res.*, 1990, **13**, 101; *CA*, **113**, 227984 (*isol*)

5,6,7,8-Tetrahydro-2,5-dimethylquinoline T-171

2,5-Dimethyl-5,6,7,8-tetrahydroquinoline [91245-80-6]

C₁₁H₁₅N 161.246

Alkaloid from *Fabiana imbricata* (Solanaceae). Bp_{0.04} 68-73°.

[94058-68-1]


Edwards, O.E. *et al.*, *Can. J. Chem.*, 1962, **40**, 256 (*isol, pmr, struct*)

Soccolini, F. *et al.*, *J. Het. Chem.*, 1984, **24**, 1001 (*synth*)

Sugita, T. *et al.*, *Heterocycles*, 1985, **23**, 2789 (*synth*)

5,6,7,8-Tetrahydrofolic acid T-172

N-[4-[[(2-Amino-1,4,5,6,7,8-hexahydro-4-oxo-6-pteridinyl)methyl]amino]benzoyl]glutamic acid, 9CI



C₁₉H₂₃N₇O₆ 445.434

All stereoisomers reported here have L-Glu config., therefore are diastereoisomers.

(6R)-*form* [74708-38-6]

5-*Formyl*: see Folic acid, F-121

(6S)-*form* [71963-69-4]

Essential coenzyme, product of reduction of folic acid by dihydrofolate reductase. Involved in C₁-transfer during synthesis of nucleoside bases. [α]_D -49.9 (c, 0.149 in 1.5M TRIS/HCl; 0.2M EtSH). Readily oxid. in air.

*N*¹⁰-*Formyl*: [2800-34-2]

[73650-39-2, 74644-66-9]

C₂₀H₂₃N₇O₇ 473.444

Product of formylation of parent tetrahydrofolic acid. Gives Folinic

acid, F-121 on heating.

5-Formyl: see Folinic acid, F-121

5-N-Me: 5-Methyltetrahydrofolic acid.
Levomefolic acid, USAN. Bodyfolin®.
Metafolin®. Nutrifolin®. LMSR
[31690-09-2]

C₂₀H₂₅N₇O₆ 459.461

Widely distributed in milk, fruit and vegetables. Folate cofactor. Source of Me groups in biosynth. of methionine from homocysteine. Used in the treatment and prevention of folate deficiency.

5-N-Me, Ca salt (1:1): **Levomefolate calcium**, USAN. LMCA
[151533-22-1]
Powder (H₂O).

(6RS)-form [135-16-0]

Powder; cryst. +1H₂O (as hydrochloride).
Sol. H₂O. pK_a -1.25; pK_a 1.24; pK_a 3.5;
pK_a 4.8; pK_a 4.82; pK_a 10.5
(2225°, CONH₂, 22).

► Exp. reprod. and teratogenic effects.
MA0600800

5-Formyl: see Folinic acid, F-121

5-N-Me: [134-35-0]


[26560-38-3] Has been used as a modulator of 5-fluorouracil toxicity in cancer patients, but effectiveness not supported by phase II clin. trials. Cryst. (H₂O + 2-mercaptoethanol). Mp >300° (as Ca salt). Predominant circulating folate monoglutamate in human plasma/serum. Most significant folate for diagnosing folate deficiency.

[31690-08-1]

- May, M. *et al.*, *J.A.C.S.*, 1951, **73**, 3067-3075 (formyl derivs)
Kallen, R.G. *et al.*, *J. Biol. Chem.*, 1966, **24**, 5845 (props)
Blair, J.A. *et al.*, *Anal. Biochem.*, 1970, **34**, 376-381 (synth, uv, pmr, N-Me)
Ruediger, H. *et al.*, *FEBS Lett.*, 1970, **11**, 265-267 (synth, biochem, N-Me)
Fontecilla-Camps, J.C. *et al.*, *J.A.C.S.*, 1978, **101**, 6114 (abs config)
Feeney, J. *et al.*, *J.C.S. Perkin 2*, 1980, 176 (derivs, pmr, cmr)
Kattchee, P.A. *et al.*, *Anal. Biochem.*, 1981, **118**, 85-90 (synth, deriv)
Charlton, P.A. *et al.*, *J.C.S. Perkin 1*, 1985, 1349 (synth, uv)
Sato, J.K. *et al.*, *Anal. Biochem.*, 1986, **104**, 516 (synth, derivs)
Selhub, J. *et al.*, *Anal. Biochem.*, 1989, **182**, 84-93 (anal, hplc, N-Me)
Brunner, H. *et al.*, *Chem. Ber.*, 1992, **125**, 2085 (synth, bibl)
Eur. Pat., 1993, ((*Applied Pharma Res.*))537 842; *CA*, **119**, 270915 (Ca salt, synth, pharmacol)
Hilhorst, E. *et al.*, *Tet. Lett.*, 1993, **34**, 4257-4260 (biochem)
Boll, E. *et al.*, *Cancer Chemother. Pharmacol.*, 1995, **35**, 339-342 (5-N-Me, clin trial)
Bailey, S.W. *et al.*, *Methods Enzymol.*, 1997, **281**, 3-16 (6S-form, synth)
Nelson, B.C. *et al.*, *J. Chromatogr. B*, 2001, **765**, 141-150 (5-N-Me, detn)
Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TCR400

1,2,3,4-Tetrahydro-6-hydroxy-β-carboline T-173

2,3,4,9-Tetrahydro-1H-pyrido[3,4-b]indol-6-ol, 9CI. **Plectocomine**
[23778-34-9]



C₁₁H₁₂N₂O 188.229

Alkaloid from the fruit of *Plectocomiopsis geminiflora* (Arecaceae). Mp 316° (evac. tube) (294°).

► UV1050000

Hydrochloride: Mp 300°.

Picrate: Mp 240°.

N,O-Di-Ac: [58100-30-4]
Mp 168-170°.

N²-Me: 1,2,3,4-Tetrahydro-6-hydroxy-2-methyl-β-carboline
[64226-26-2]
[171066-23-2 (monohydrochloride)]

C₁₂H₁₄N₂O 202.255

Alkaloid from *Evodia fargesii*.

N²-Me, N²-oxide, O-β-D-glucopyranoside: [890318-61-3]

C₁₈H₂₄N₂O₇ 380.397

Alkaloid from *Evodia fargesii*.
Amorph. brown powder. [α]_D²⁵ -21 (c, 0.19 in H₂O). λ_{max} 218 ; 274 (H₂O).

Me ether: 1,2,3,4-Tetrahydro-6-methoxy-β-carboline. 2,3,4,9-Tetrahydro-6-methoxy-1H-pyrido[3,4-b]indole. 6-Methoxytetrahydro-β-carboline. **Pinoline**. 6-Methoxytryptoline
[20315-68-8]

C₁₂H₁₄N₂O 202.255

Alkaloid from *Desmodium pulchellum* (Fabaceae). Monoamine oxidase inhibitor. Tan needles (MeOH aq.). Mp 223-224°.

Me ether, hydrochloride: [35764-54-6]
Needles (EtOH aq.). Mp 288-290° (268-269°).

Me ether, picrate:

Cryst. (EtOH). Mp 240-241°.

N²,O-Di-Me: 1,2,3,4-Tetrahydro-6-methoxy-2-methyl-β-carboline. 2,3,4,9-Tetrahydro-6-methoxy-2-methyl-1H-pyrido[3,4-b]indole, 9CI
[6582-80-5]

C₁₃H₁₆N₂O 216.282

Alkaloid from *Desmodium pulchellum*, *Nectandra megapotamica* and *Virola theiodora*. Fine needles (MeOH aq.). Mp 214° Mp 217-218° (183-184°). λ_{max} 227 (ε 27300); 283 (ε 8350) (EtOH).

N²,O-Di-Me, picrate:

Cryst. (EtOH). Mp 182-191°.

N²,N⁹,O-Tri-Me: 1,2,3,4-Tetrahydro-6-methoxy-2,9-dimethyl-β-carboline
[25968-13-2]

C₁₄H₁₈N₂O 230.309

Alkaloid from *Papaver* sp. (Papaveraceae). Pale yellow needles (MeOH). Mp 132-135°.

[17952-64-6 ; 58100-31-5]

Abramovitch, R.A. *et al.*, *J.C.S.*, 1956, 4589-

4592 (Me ether, synth)

Pachter, I.J. *et al.*, *J.A.C.S.*, 1961, **83**, 635 (Me ether, synth)

Kiang, A.K. *et al.*, *J. Nat. Prod.*, 1967, **30**, 189 (isol, uv, pmr, ms, struct, synth, N,O-di-Ac)
Agurell, S. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 903 (N,O-di-Me, isol, ms)

Audette, R.C.S. *et al.*, *Can. J. Chem.*, 1970, **48**, 149 (isol, uv, ir, pmr, ms)

Frahn, J.L. *et al.*, *Aust. J. Chem.*, 1971, **24**, 2189; 1974, **27**, 1367 (N,O-di-Me, isol, uv, synth)

Shannon, P.V.R. *et al.*, *J.C.S. (C)*, 1971, 2837-2839 (derivs, synth, pmr, uv, ms, struct)

Ghosal, S. *et al.*, *Planta Med.*, 1972, **21**, 200 (deriv)

Vijayanagar, H.M. *et al.*, *J. Nat. Prod.*, 1975, **38**, 442 (deriv, isol, synth, ms)

Filho, D. dos S. *et al.*, *Phytochemistry*, 1975, **14**, 821-822 (N,O-di-Me)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 3343; 1977, **42**, 132; 1978, **43**, 316 (isol, uv, ir)

Baudouin, G. *et al.*, *J. Nat. Prod.*, 1981, **44**, 546 (isol, deriv)

Jossang, A. *et al.*, *J.O.C.*, 1991, **56**, 6527 (N,O-di-Me, isol, ir, ms, pmr, cmr)


Dantale, S.W. *et al.*, *Tetrahedron*, 2003, **59**, 5507-5514 (N,O-di-Me, synth)

Qu, S.-J. *et al.*, *Planta Med.*, 2006, **72**, 264-266 (N-Me N-oxide glucoside)

Liu, Q.-W. *et al.*, *Zhongguo Tianran Yaowu*, 2006, **4**, 25-29; *CA*, **145**, 425429 (N²-Me)

1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinoline T-174

1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethyl-5-isoquinolinol, 9CI. **Gigantine**



(S)-form

C₁₃H₁₉NO₃ 237.298

(S)-form [32829-58-6]

Alkaloid from *Carnegiea gigantea* (Cactaceae). Poss. hallucinogen. Cryst. (Et₂O). Mp 151-152°. [α]_D²⁵ +27 (c, 1.99 in CHCl₃).

Hydrochloride:

Cryst. (EtOH). Mp 221.5-222.5°.

(±)-form

Synthetic. Cryst. (Et₂O). Mp 121-123°.

Me ether: 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1,2-dimethylisoquinoline
Mp 153-154° (as picrate).

I'-Hydroxy: see 1,2,3,4-Tetrahydro-5-hydroxy-1-hydroxymethyl-6,7-dimethoxy-2-methylisoquinoline, T-178


Kapadia, G.J. *et al.*, *Chem. Comm.*, 1970, 856 (synth)

Choudhury, A.M. *et al.*, *Chem. Ind. (London)*, 1971, 578 (synth)

Brown, S.D. *et al.*, *J.O.C.*, 1972, **37**, 1825 (isol, ir, pmr, ms, struct, abs config, synth)

Bruhn, J.G. *et al.*, *J. Nat. Prod.*, 1976, **39**, 197 (isol)

Cymerman Craig, J. *et al.*, *J.A.C.S.*, 1977, **99**, 7996 (uv, cd)

1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-isoquinolinecarboxylic acid**Peyoxylic acid**
[29193-99-5]C₁₂H₁₅NO₅ 253.254**(±)-form**


Mp 237-238° dec.

Me ether: [41303-73-5]




Synthetic. Mp 238-240° dec.

(ξ)-formTrace component of *Lophophora williamsii* (Peyote cactus) (Cactaceae). The nat. prod. was identified by glc comparison with synthetic racemic matl.*Me ether*: 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-isoquinolinecarboxylic acid, 9CI. **O-Methylpeyoxalic acid**C₁₃H₁₇NO₅ 267.281Trace component of *Lophophora williamsii* (Cactaceae). Mp 238-240° dec.Kapadia, G.J. *et al.*, *J.A.C.S.*, 1970, **92**, 6943 (*isol, ir, pmr, ms, synth, struct*)Kapadia, G.J. *et al.*, *J. Het. Chem.*, 1973, **10**, 135 (*deriv*)**6,7,9,10-Tetrahydro-5-(4-hydroxy-3,5-dimethoxyphenyl)-5H-benzol[1,3]dioxolo[4',5':4,5]benzo[de]pyrido[3,2,1-ij]-quinolin-7-one**

[779355-68-9]




Absolute Configuration


C₂₈H₂₃NO₆ 469.493Lignoaporphine. Alkaloid from the wood of *Annona dioica*. Amorph. yellow solid. Mp 211-213°. [α]_D²⁶ +326 (c, 2.5 in CHCl₃).Dos Santos, P.R.D. *et al.*, *J. Braz. Chem. Soc.*, 2003, **14**, 396-400 (*isol, pmr, cmr*)**1,2,3,4-Tetrahydro-8-hydroxy-1,3-dimethylisoquinoline**1,2,3,4-Tetrahydro-1,3-dimethyl-8-isoquinolinol. **Phylline**C₁₁H₁₅NO 177.246**(1R,3R)-form** [240133-91-9]Alkaloid from *Habropetalum dawei*.**N-Me: N-Methylphylline**C₁₂H₁₇NO 191.272Alkaloid from the root bark of *Ancistrocladus barteri*. Yellow oil. [α]_D²⁰ +22 (c, 0.4 in CHCl₃). λ_{max} 220 ; 277 (EtOH).**N,O-Di-Me: 1,2,3,4-Tetrahydro-8-methoxy-1,2,3-trimethylisoquinoline. N,O-Dimethylphylline**C₁₃H₁₉NO 205.299Alkaloid from the root bark *Ancistrocladus barteri*. Cryst. (Et₂O/CH₂Cl₂/petrol) (as hydrochloride). [α]_D²⁵ +39.1 (c, 0.54 in CHCl₃) (hydrochloride).Bringmann, G. *et al.*, *Anal. Chem.*, 1999, **71**, 2678-2686 (*isol, cd, pmr, ms*)Bringmann, G. *et al.*, *Z. Naturforsch., B*, 2003, **58**, 577-584 (*N-Methylphylline O,N-Dimethylphylline*)**1,2,3,4-Tetrahydro-5-hydroxy-1-hydroxymethyl-6,7-dimethoxy-2-methylisoquinoline**1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-2-methyl-1-isoquinolinemethanol, 9CI. **Deglucopteroceine**
[70475-63-7]C₁₃H₁₉NO₄ 253.297**(ξ)-form**Alkaloid from the cactus *Pterocereus gaumeri* (Cactaceae). Mp 247-248° (as hydrochloride). [α]_D²⁶ -1.04 (c, 2.2 in H₂O). Probably an artifact of extraction.**N-Oxide: Deglucopteroceine N-oxide**
[85769-27-3]C₁₃H₁₉NO₅ 269.297Alkaloid from *Pterocereus gaumeri* (Cactaceae). Cryst. (EtOH/Et₂O). Mp 210-213°.**5-O-β-D-Glucopyranoside: Pteroceine**
[70475-62-6]C₁₉H₂₉NO₉ 415.439Alkaloid from *Pterocereus gaumeri* (Cactaceae). Mp 198-199°. [α]_D²⁶ -4.5 (c, 1.35 in H₂O).Mohamed, Y.A.H. *et al.*, *J. Nat. Prod.*, 1979, **42**, 197 (*isol, uv, ir, pmr, cmr, ms, struct, Deglucopteroceine, Pteroceine*)Pummangura, S. *et al.*, *Phytochemistry*, 1982, **21**, 2375 (*oxide, isol, ir, pmr, cmr, synth*)**1,2,3,4-Tetrahydro-6-hydroxy-7-methoxy-1-methylisoquinoline**1,2,3,4-Tetrahydro-7-methoxy-1-methyl-6-isoquinolinol, 9CI. **Salsoline**
[63596-58-7]**(S)-form**C₁₁H₁₅NO₂ 193.245

Antihypertensive, has been used clinically in the USSR. Antihistamine. Plant growth inhibitor. Log P 1.23 (calc).

(R)-formAlkaloid from *Salsola richteri* and *Salsola arbuscula* (Chenopodiaceae). Monoamine oxidase inhibitor. Mp 215-216°. Log P 1.23 (calc).**Hydrochloride**: [51424-33-0]Mp 174-175° (171-172°). [α]_D +40.1(H₂O) (natural). [α]_D²⁵ +31 (c, 1 in

MeOH) (synthetic).

▶ LD₅₀ (mus, ivn) 140 mg/kg. NX6000000**(S)-form** [89-31-6]Alkaloid from *Salsola richteri* (Chenopodiaceae). Mp 218-221°.**Hydrochloride**: [881-26-5]Cryst. (H₂O). Mp 174-175°. [α]_D -39.2(H₂O). [α]_D²⁵ -31.5 (c, 1 in MeOH).*Me ether*: see 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methylisoquinoline, T-167**(±)-form**Alkaloid from *Alangium lamarckii*, *Echinocereus merkeri*, *Pachycereus pecten-aboriginum*, *Corispermum leptopyrum*, *Salsola arbuscula*, *Salsola kali*, *Salsola richteri*, *Salsola ruthenica*, *Salsola soda*, *Calycotome spinosa*, *Desmodium tiliaefolium* and *Genista purgans* (Cactaceae, Fabaceae, Alangiaceae, Chenopodiaceae). Mp 224-226°.**Hydrochloride**:Cryst. (EtOH/Et₂O). Mp 147-149°.**Picrate**:Yellow needles (H₂O). Mp 189-191°.Späth, E. *et al.*, *Ber.*, 1934, **67**, 1214 (*isol, struct*)Proskurnina, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1937, **4**, 1265 (*isol*)Kovács, O. *et al.*, *Chem. Ber.*, 1951, **84**, 795 (*synth*)Battersby, A.R. *et al.*, *J.C.S.*, 1960, 1214 (*abs config*)Ghosal, S. *et al.*, *Phytochemistry*, 1973, **12**, 193 (*isol, uv, ir, pmr, ms*)Teitel, S. *et al.*, *J. Med. Chem.*, 1974, **17**, 134 (*synth, pmr, uv*)Drost-Karbowska, K. *et al.*, *Acta Pol. Pharm.*, 1977, **34**, 421; *CA*, **88**, 117814b (*isol*)Cymerman Craig, J. *et al.*, *J.A.C.S.*, 1977, **99**, 7996 (*uv, cd*)Strömbom, J. *et al.*, *Acta Pharm. Suec.*, 1978, **15**, 127 (*occur*)Strömbom, J. *et al.*, *J. Chromatogr.*, 1978, **147**, 513 (*hplc*)Achari, B. *et al.*, *Planta Med. (Suppl.)*, 1980, 5 (*occur*)Bembenek, M.E. *et al.*, *J. Med. Chem.*, 1990, **33**, 147 (*pharmacol, enantiomers*)**1,2,3,4-Tetrahydro-7-hydroxy-5-methoxy-2-methylisoquinoline**1,2,3,4-Tetrahydro-5-methoxy-2-methyl-7-isoquinolinol, 9CI. **Uberine**
[63596-58-7]

C₁₁H₁₅NO₂ 193.245Alkaloid from *Dolichothele uberiformis* (Cactaceae). Mp 263-267° dec. (as hydrochloride).Ranieri, R.L. *et al.*, *J. Nat. Prod.*, 1977, **40**, 173 (isol, ir, pmr, ms, struct)**1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-1-methylisoquinoline** T-1811,2,3,4-Tetrahydro-6-methoxy-1-methyl-7-isoquinolinol, 9CI. *Isosalsoline* [4593-97-9]C₁₁H₁₅NO₂ 193.245**(R)-form**Synthetic. Mp 241-242° (as hydrochloride). [α]_D²⁰ +24.7 (c, 1 in MeOH) (hydrochloride).

N-Me: N-Methylisalsoline. 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl-7-isoquinolinol, 9CI. 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-1,2-dimethylisoquinoline. 1-Methylcorypalline [35048-35-2]

C₁₂H₁₇NO₂ 207.272Minor alkaloid from tubers of *Corydalis ambigua* (Papaveraceae). Also isol. from *Haloxylon articulatum* (Chenopodiaceae). Mp 156-158°. [α]_D²⁰ +33.5 (c, 0.23 in CHCl₃). [α]_D²⁰ +1 (c, 0.23 in EtOH).**(S)-form**Synthetic. Mp 241-241° (as hydrochloride). [α]_D²⁰ -26 (c, 1 in MeOH) (hydrochloride).N-Me: Alkaloid from aerial parts of *Arthrocnemum glaucum* (Chenopodiaceae). Cubes (MeOH). Mp 175-178°. [α]_D²⁰ -32 (c, 0.79 in MeOH).**(±)-form**


N-Me: [19641-12-4]

Synthetic. Mp 148-150°.

N-Me; hydrochloride: Mp 190-192°.

(ξ)-formAlkaloid from *Pachycereus pecten-aboriginum* (Cactaceae). Spec. rotn. and stereochem. unspecified.Strukov, I.T. *et al.*, *Zh. Obshch. Khim.*, 1959, **29**, 3831; 1961, **31**, 2709; *CA*, **54**, 19676a; **56**, 11567f (synth, deriv)Carling, C. *et al.*, *Acta Pharm. Suec.*, 1970, **7**, 285; *CA*, **73**, 63154f (isol, deriv)Naruto, S. *et al.*, *Phytochemistry*, 1973, **12**, 3008 (occur, deriv)Teitel, S. *et al.*, *J. Med. Chem.*, 1974, **17**, 134 (synth, uv, ord, cd, pmr)Strombom, J. *et al.*, *Acta Pharm. Suec.*, 1978, **15**, 127 (isol)Strombom, J. *et al.*, *J. Chromatogr.*, 1978, **147**, 513 (hplc)Khalil, A.T. *et al.*, *Phytochemistry*, 1992, **31**, 1023 (*Methylcorypalline*)**1,2,3,4-Tetrahydro-8-hydroxy-6-methoxy-1,2,3-trimethylisoquinoline** T-182

1,2,3,4-Tetrahydro-6-methoxy-1,2,3-trimethyl-8-isoquinolinol, 9CI

C₁₃H₁₉NO₂ 221.299**(1R,3S)-form****Gentrymine A**

[157740-47-1]

Alkaloid from leaves and twigs of *Ancistrocladus korupensis* (Ancistrocladaceae). Solid. [α]_D²⁰ +120 (c, 0.17 in MeOH).**(1S,3S)-form****N-Me: Gentrymine B**

[165689-38-3]

C₁₄H₂₂NO₂[⊕] 236.333Quaternary alkaloid from *Ancistrocladus korupensis* (Ancistrocladaceae).Cubes (Me₂CO)(as iodide). Mp 126° (iodide). [α]_D²⁵ -29.8 (c, 0.3 in MeOH).

[165330-33-6]

Haddock, Y.F. *et al.*, *J.O.C.*, 1994, **59**, 6349




(isol, uv, ir, pmr, cmr, struct)

Haddock, Y.F. *et al.*, *Tet. Lett.*, 1995, **36**, 4753


(Gentrymine B)

Bringmann, G. *et al.*, *Tetrahedron*, 2000, **56**, 581-585 (synth, abs config)**3,4,5,6-Tetrahydro-6-hydroxymethyl-3,6-dimethyl-4-pyrimidinecarboxylic acid, 9CI** T-183


[246137-84-8]

C₈H₁₄N₂O₃ 186.21Isol. from the Palauan sponge *Protophlitaspongia aga*. Antifouling agent against barnacles.Hattori, T. *et al.*, *Fish. Sci.*, 2001, **67**, 690-693 (isol)**1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-3-isoquinolinecarboxylic acid** T-184C₁₁H₁₃NO₃ 207.229**(1ξ,3ξ)-form** [19697-94-0]Alkaloid from the latex of *Euphorbia myrsinites* and *Euphorbia tirucalli*. Needles (propanol aq.). Mp 269-271° dec. λ_{max} 275 (H₂O).Müller, P. *et al.*, *Z. Naturforsch., B*, 1968, **23**, 491-493 (isol, struct)**1,4,5,6-Tetrahydro-5-hydroxy-2-methyl-4(6)-pyrimidine-carboxylic acid** T-185*Pyrostatin A*
[117229-60-4]C₆H₁₀N₂O₃ 158.157Prod. by *Streptomyces parvulus* and marine-derived *Streptomyces* sp. SA-3501 (Pyrostatin A). Powder. [α]_D²⁰ +125.4 (H₂O) (Pyrostatin A). Tentatively assigned struct. of Pyrostatin A after revision in 2006.Inbar, L. *et al.*, *J. Biol. Chem.*, 1988, **263**,16014-16022 (isol, cmr, N-15 nmr, biosynth)
Castellanos, L. *et al.*, *Org. Lett.*, 2006, **8**, 4967-4970 (Pyrostatin A)**2,7,19,20-Tetrahydro-3-hydroxy-1-methylvobasan-17-oic acid δ-lactone** T-186


[77485-23-5]

Absolute
ConfigurationC₂₁H₂₈N₂O₂ 340.464Alkaloid from the leaves of *Rauwolfia oreogiton* (Apocynaceae). Mp 180-181°.Akinloye, B.A. *et al.*, *Phytochemistry*, 1980, **19**, 2741 (isol, uv, ir, pmr, ms, struct)**1,2,3,6-Tetrahydro-3-hydroxy-2-pyridinecarboxylic acid, 9CI** T-187

1,2,3,6-Tetrahydro-3-hydroxycypolinic acid. 3-Hydroxybaikiain

C₆H₉NO₃ 143.142**(2S,3R)-form** [112269-85-9]Isol. from fruiting bodies of the toxic mushroom *Russula subnigricans*. Needles (MeOH aq.). Mp 300-302°. [α]_D²⁰ -332.7 (c, 0.3 in H₂O). [α]_D²⁰ -324 (c, 0.3 in 1M HCl).Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 3482 (isol, ir, pmr, cmr, ms, struct)

2,3,5,7a-Tetrahydro-1-hydroxy-1H-pyrrolizine-7-methanol, 9CI
7-Hydroxy-1-hydroxymethyl-1,2-didehydropyrrolizidine



(7R,8R)-form

C₈H₁₃NO₂ 155.196

Pyrrolizidine numbering shown. In CAS numbering 8 is 7a. Simple esters of these bases are given here. Many more complex esters have individual entries.

(7R,8R)-form**Retronecine**

[480-85-3]

Necine base from numerous pyrrolizidine alkaloids. Trace constit. in seedlings of *Crotalaria scassellatii*. Constit. of the pheromones of the flea beetles *Longitarsus* spp. Cryst. (Me₂CO or by subl.). Sol. H₂O, EtOH; spar. sol. Et₂O; fairly sol. Me₂CO, CHCl₃. Mp 121-122°. [α]_D²⁶ +50.2 (EtOH). pK_a 8.88 (26°). Does not form a picrate.

▶ LD₅₀ (mus, ivn) 634 mg/kg. VH7175000

Hydrochloride: Mp 162-163°. [α]_D¹⁵ -16 (EtOH).

N-Oxide: Retronecine N-oxide

[6870-33-3]

C₈H₁₃NO₃ 171.196

Alkaloid from *Senecio caudatus* and *Werneria decora*.

7-O-(3-Methylbutanoyl): O⁷-(3-Methylbutanoyl)retronecine

[63503-35-5]

C₁₃H₂₁NO₃ 239.314

Constit. of *Crotalaria scassellatii*.

7-O-(3-Methylbutanoyl), N-oxide: O⁷-(3-Methylbutanoyl)retronecine N-oxide

[121181-03-1]

C₁₃H₂₁NO₄ 255.313

Constit. of *Crotalaria scassellatii*.

7-O-Angeloyl: 7-Angeloylretronecine

[6029-82-9]

C₁₃H₁₉NO₃ 237.298

Constit. of *Cynoglossum latifolium* and *Senecio triangularis*. Constit. of the pheromones of *Longitarsus* spp. Plates (petrol). Mp 76-77° (74°). [α]_D²⁴ +49 (c, 1.38 in EtOH).

9-O-Angeloyl: 9-Angeloylretronecine

[6922-62-9]

C₁₃H₁₉NO₃ 237.298

Alkaloid from aerial parts of *Alkanna orientalis*. Constit. of the pheromone of *Longitarsus exoletus*. Active against gram-positive bacteria. Oil. [α]_D^{19.5} +2.01 (c, 0.6 in EtOH).

9-O-Angeloyl, N-oxide: 9-Angeloylretronecine N-oxide

[27773-86-0]

C₁₃H₁₉NO₄ 253.297

Alkaloid from the bark of *Bhesa archboldiana* (Celastraceae) and from *Heliotropium bursiferum*. Cryst. (Me₂CO). Mp 153-154°. [α]_D +30 (c,

0.98 in CHCl₃).**7-O-(3-Methyl-2-butenyl): O⁷-Seneciolyretronecine**

[63503-33-3]

C₁₃H₁₉NO₃ 237.298

Alkaloid from *Senecio variabilis* and *Senecio caudatus*. Constit. of the pheromones of *Longitarsus* spp.

7-O-(3-Methyl-2-butenyl), N-oxide: O⁷-Seneciolyretronecine N-oxide

[119590-86-2]

C₁₃H₁₉NO₄ 253.297

Constit. of *Crotalaria scassellatii*.

9-O-(3-Methyl-2-butenyl): 1,2-Dehydrofuchsisenecionine. O⁹-Seneciolyretronecine

[124262-88-0]

C₁₃H₁₉NO₃ 237.298

Alkaloid from *Senecio variabilis* and *Senecio nemorensis* var. *fuchsii*. Constit. of the pheromones of *Longitarsus* spp.

7-O-(2S-Hydroxy-2,3-dimethylbutanoyl): Heliohoustine

[366452-81-5]

C₁₄H₂₃NO₄ 269.34

Alkaloid from *Ageratum houstonianum*.

9-O-(2S-Hydroxy-2,3-dimethylbutanoyl): Retrohoustine

[366452-83-7]

C₁₄H₂₃NO₄ 269.34

Alkaloid from *Ageratum houstonianum*.

7-O-(4-Hydroxy-3-methyl-2Z-butenyl), 9-angeloyl: Sencalenine

[95066-37-8]

C₁₈H₂₅NO₅ 335.399

Diester of retronecine with angelic and hydroxyangelic acids. Alkaloid from *Senecio cacaliaster* (Asteraceae). Oil. [α]_D²⁰ -8 (EtOH).

9-O-(2-Hydroxymethyl-2-butenyl), 7-angeloyl: see Triangularine, T-454**7-O-(2S-Hydroxy-3S-methylpentanoyl), N-oxide: Isocreatonotine N-oxide**

[134306-31-3]

C₁₄H₂₃NO₅ 285.339

Alkaloid from the moth *Cretonotos transiens*.

9-O-(2S-Hydroxy-3S-methylpentanoyl), N-oxide: Creatonotine N-oxide

[134306-30-2]

C₁₄H₂₃NO₅ 285.339

Alkaloid from the moth *Cretonotos transiens*.

7-O-[2-Hydroxy-2-(1-hydroxyethyl)-3-methylpentanoyl]: Heliospathine. Retronecine 7-O-curassavate

[135683-57-7]

C₁₆H₂₇NO₅ 313.393

Alkaloid from *Heliotropium spathulum* (Boraginaceae). Pale yellow oil. [α]_D²⁰ +1.5 (EtOH).

9-O-[2-Hydroxy-2-(1-hydroxyethyl)-3-methylpentanoyl]: Retronecine 9-O-curassavate

[135626-74-3]

C₁₆H₂₇NO₅ 313.393

Alkaloid from *Heliotropium spathulum*. Yellow oil.

9-O-[2S-Hydroxy-2-(1R-hydroxyethyl)-4-methylpentanoyl], N-oxide:C₁₆H₂₇NO₆ 329.392

Alkaloid from the roots of *Anchusa strigosa*. Red oil. [α]_D²⁵ +2.3 (c, 0.1 in MeOH). CAS no. not found 8-14CI. λ_{max} 235 (sh); 282 (MeOH).

9-O-[2S-Hydroxy-2-(1S-hydroxyethyl)-4-methylpentanoyl]:C₁₆H₂₇NO₅ 313.393

Alkaloid from the roots of *Anchusa strigosa*. Orange-yellow oil. [α]_D²⁵ +2.2 (c, 0.05 in MeOH). CAS no. not found 8-14CI. λ_{max} 233 (sh); 285 (MeOH).

9-O-[2S-Hydroxy-2-(1S-hydroxyethyl)-4-methylpentanoyl], N-oxide:C₁₆H₂₇NO₆ 329.392

Alkaloid from the roots of *Anchusa strigosa*. Orange oil. [α]_D²⁵ +4 (c, 0.05 in MeOH). CAS no. not found 8-14CI. λ_{max} 230 (sh); 280 (MeOH).

7,9-Dibenzoyl: Alkaloid CG 1. Retronecine-7,9-dibenzoate

[25471-42-5]

C₂₂H₂₁NO₄ 363.412

Alkaloid from the flowers of *Caccinia glauca* (Boraginaceae). Noncryst. Mp 137-139° (as picrate).

9-Me ether: O⁹-Methylretronecine. 7-Hydroxy-1-methoxymethyl-1,2-dehydropyrrolizidine

[24999-22-2]

C₉H₁₅NO₂ 169.223

Constit. of *Crotalaria* spp. (Fabaceae). Mp 35-40°. Bp_{0.4} 77°. [α]_D +38 (EtOH).

9-Me ether, 7-Ac: 7-Acetoxy-1-methoxymethyl-1,2-dehydropyrrolizidine

[15211-06-0]

C₁₁H₁₇NO₃ 211.26

Alkaloid from *Crotalaria trifolium* (Fabaceae). Bp_{0.03} 61°. [α]_D +15.2 (EtOH).

(7S,8R)-form**Heliotridine**

[520-63-8]

Found in *Heliotropium ellipticum*, *Heliotropium eichwaldii* and *Heliotropium lasiocarpum*. Hydrol. prod. of many other pyrrolizidine alkaloids. Constit. of the pheromones of the flea beetles *Longitarsus* spp. Cryst. Mp 116-118°. [α]_D²⁰ +31.

▶ LD₅₀ (rat, ipr) 1500 mg/kg. UY8405000**7-O-Angeloyl: 7-Angeloylheliotridine.****Rivularine[†]**

[723-78-4]

C₁₃H₁₉NO₃ 237.298

Alkaloid from *Crotalaria officinale*, other *Crotalaria* spp., *Heliotropium eichwaldii*, *Senecio crispatis*, *Senecio rivularis* and other spp. Constit. of the pheromones of *Longitarsus* spp. Cryst. (EtOAc). Mp 116-117°. [α]_D²⁴ -18 (c, 0.78 in CHCl₃). [α]_D²⁴ +19 (c, 0.66 in EtOH). λ_{max} 218 (log ε 4.07) (EtOH).

▶ EM9257500

7-O-Angeloyl, N-oxide: Rivularine N-oxide

[64595-72-8]

C₁₃H₁₉NO₄ 253.297

Alkaloid from *Senecio integrifolius* var.

fauriri (Asteraceae). Fine needles. Mp 93-95°. $[\alpha]_D^{25} +8.3$ (c, 1 in EtOH).

7,9-Di-O-angeloyl: 7,9-Diangeloylheliotridine. **Asperumine**

[78513-20-9]

C₁₈H₂₅NO₄ 319.4

Alkaloid from *Symphytum asperum*. Mp 135-137° (as picrate).

7,9-Di-O-angeloyl, N-oxide: **Asperumine N-oxide**

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Echium vulgare*. Not indexed by CA to 14CI (2001).

9-O-(2S-Hydroxy-2,3-dimethylbutanoyl): **Isoretrohoustine**

[366452-82-6]

C₁₄H₂₃NO₄ 269.34

Alkaloid from *Ageratum houstonianum*.

9-O-[2S-Hydroxy-2-(1S-hydroxyethyl)-4-methylpentanoyl]:

C₁₆H₂₇NO₅ 313.393

Alkaloid from the leaves of *Anchusa strigosa*. Orange-yellow oil. $[\alpha]_D^{25} +14.3$ (c, 0.1 in MeOH). λ_{max} 233 (sh); 285 (MeOH).

9-Me ether: **O⁹-Methylheliotridine**

[15211-05-9]

C₉H₁₅NO₂ 169.223

Alkaloid from *Crotalaria trifoliastrum*, *Crotalaria aridicola* and *Crotalaria medicaginea* (Fabaceae). Mp 54°. Bp_{0.3} 90-94° (bath). $[\alpha]_D^{25} +25.2$ (c, 1.78 in EtOH).

[875-22-9, 6029-83-0, 133738-53-1, 133738-52-0, 17690-54-9]

Warren, F.L. *et al.*, *J.C.S.*, 1958, 4574-4575 (*abs config, bibl*)

Geissman, T.A. *et al.*, *J.O.C.*, 1962, **27**, 139-142 (*Retronecine, synth*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1965, **18**, 1605-1624; 1966, **19**, 1955-1964; 1967, **20**, 757-768; 1970, **23**, 1279-1282 (*isol, pmr, synth, esters*)

Klásek, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 2512-2522 (*Rivularine*)

Šimáček, V. *et al.*, *Coll. Czech. Chem. Comm.*, 1969, **34**, 1832-1837 (*uv*)

Man'ko, I.V. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1970, **40**, 2506 (*Asperumine*)

Sawhney, R.S. *et al.*, *J. Indian Chem. Soc.*, 1970, **47**, 741-742 (*O⁹-Methylheliotridine*)

Pedersen, E. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 249-256 (*ms*)

Culvenor, C.C.J. *et al.*, *J.C.S. (C)*, 1971, 3653-3664 (*cd*)

Karimov, A. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 452 (*Asperumine N-oxide*)

Bale, N.M. *et al.*, *Phytochemistry*, 1975, **14**, 2617-2622 (*biosynth*)

Siddiqui, M.A. *et al.*, *Phytochemistry*, 1978, **17**, 2049-2050 (*Alkaloid CG 1*)

Jones, A.J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 1173-1184 (*cmr*)

Grue-Sorensen, G. *et al.*, *Can. J. Chem.*, 1982, **60**, 643-662 (*biosynth*)

Niwa, H. *et al.*, *Chem. Lett.*, 1983, 125-126 (*Retronecine, synth*)

Rüeger, H. *et al.*, *Heterocycles*, 1983, **20**, 1331-1334 (*Retronecine, synth*)

Chamberlin, A.R. *et al.*, *J.A.C.S.*, 1983, **105**, 3653-3656 (*Heliotridine, synth*)

Ohsawa, T. *et al.*, *J.O.C.*, 1983, **48**, 3644-3648 (*Retronecine, synth*)

Röder, E. *et al.*, *Phytochemistry*, 1984, **23**, 1761-1763 (*Sencalenine*)

Hart, D.J. *et al.*, *J.O.C.*, 1985, **50**, 235-242 (*Heliotridine, synth*)

Vedejs, E. *et al.*, *J.O.C.*, 1985, **50**, 2170-2174 (*Retronecine, synth*)

Glinski, J.A. *et al.*, *Tet. Lett.*, 1985, **26**, 2857-2860 (*Heliotridine, Angeloylheliotridine, synth*)

Mattocks, A.R. *et al.*, *Chemistry and Toxicology of Pyrrolizidine Alkaloids*, Academic Press, 1986, (*tox, rev*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1986, **25**, 1151-1159 (*O⁷-angeloyl, O⁷-seneciyl, O⁹-seneciyl*)

Kano, S. *et al.*, *Heterocycles*, 1988, **27**, 253-256 (*Heliotridine, synth*)

Toppel, G. *et al.*, *Phytochemistry*, 1988, **27**, 3757-3760 (*Crotalaria scassellatii constits*)

Marquina, G. *et al.*, *Pharmazie*, 1989, **44**, 870-871 (*activity*)

Weidenfeld, H. *et al.*, *Sci. Pharm.*, 1989, **57**, 97; *CA*, **112**, 11824d (*1,2-Dehydrofuchsisenecionine*)

Hartmann, T. *et al.*, *Biochem. Syst. Ecol.*, 1990, **18**, 549-554 (*Creatonotine N-oxide, Isoreatonotine N-oxide*)

Roeder, E. *et al.*, *Phytochemistry*, 1990, **29**, 11-29 (*rev, cmr*)

Roeder, E. *et al.*, *Phytochemistry*, 1991, **30**, 1703-1706; 1734-1737 (*Heliospathine, Rivularine N-oxide*)

Roeder, E. *et al.*, *Fitoterapia*, 1992, **63**, 405-408 (*9-Angeloylretrohecine*)

Witte, L.R. *et al.*, *Phytochemistry*, 1992, **32**, 187-196 (*glc-ms*)

Logie, C.G. *et al.*, *Phytochemistry*, 1994, **37**, 43-109 (*rev, pmr*)

Dobler, S. *et al.*, *J. Chem. Ecol.*, 2000, **26**, 1291-1298 (*Longitarsus pheromones*)

Wiedefeld, H. *et al.*, *Phytochemistry*, 2001, **57**, 1269-1271 (*Heliohoustine, Retrohoustine, Isoretrohoustine*)

Braca, A. *et al.*, *Planta Med.*, 2003, **69**, 835-841 (*Anchusa strigosa esters*)

Huang, J.-M. *et al.*, *Tet. Lett.*, 2004, **45**, 3047-3050 (*synth*)

Roche, C. *et al.*, *J.O.C.*, 2005, **70**, 8352-8363 (*Retronecine, synth*)

Siciliano, T. *et al.*, *Phytochemistry*, 2005, **66**, 1593-1600 (*Anchusa strigosa esters*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, RfK000

5,6,7,7a-Tetrahydro-7a-hydroxy-3H-pyrrolizin-3-one, 9CI

T-189

5-Hydroxy-1-azabicyclo[3.3.0]oct-3-en-2-one

[119100-89-9]



C₇H₉NO₂ 139.154

(±)-**form**

Oil or pale yellow cryst. Mp 94-95°.

(ξ)-**form**

Prod. by *Streptomyces globisporus* and *Streptomyces griseus*.

[116515-49-2]


Guru, S. *et al.*, *Indian J. Chem., Sect. B*, 1987, **26**, 955 (*synth*)

Ikeda, M. *et al.*, *Heterocycles*, 1988, **27**, 943 (*synth, ir, pmr*)

Jizba, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1993, **58**, 1452-1456 (*isol, synth*)

1,2,3,4-Tetrahydro-4-hydroxy-4-quinolinecarboxylic acid Tuberosine B[†]

T-190



C₁₀H₁₁NO₃ 193.202

(ξ)-**form** [306955-19-1]


Alkaloid from the seeds of *Allium tuberosum* (Chinese chives).

Sang, S. *et al.*, *Tianran Chanwu Yanjiu Yu Kaifa*, 2000, **12**, 1; *CA*, **133**, 360801s (*isol*)

1,2,3,4-Tetrahydro-ar-hydroxy-ar-trimethoxy-2-methylisoquinoline

T-191

[93474-24-9]



C₁₃H₁₉NO₄ 253.297

Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).


Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109-114 (*occur*)

4,5,6,7-Tetrahydro-1H-imidazo[4,5-c]pyridine, 9CI

T-192

Spinaceamine

[6882-74-2]



C₆H₉N₃ 123.157

Isol. from skin extracts of the amphibians *Verongula* sp., *Leptodactylus pentadactylus labyrinthicus*, *Nyctimystes disrupta* and *Litoria moorei*. Shows antimicrobial activity. Genus name given as *Nyctimystes* in lit.

Hydrochloride (1:2): [62002-31-7] Mp 277-279° dec.

Dipicrate: Mp 224-225° dec.

N⁶-**Me**: **6-Methylspinaceamine**

[10517-40-5]

C₇H₁₁N₃ 137.184


Isol. from the skin of *Leptodactylus pentadactylus labyrinthicus* and *Nyctimystes disrupta*. Shows antimicrobial activity. Mp 272-274° (as dihydrochloride).

[64403-23-2]

Erspamer, V. *et al.*, *Experientia*, 1963, **19**, 346-347 (*occur*)

- Erspamer, V. *et al.*, *Arch. Biochem. Biophys.*, 1964, **105**, 620-629 (*occur*)
 Vitali, T. *et al.*, *Gazz. Chim. Ital.*, 1964, **94**, 296-305 (*pmr, struct, synth*)
 Nardelli, M. *et al.*, *Ric. Sci., Parte 2: Sez. A*, 1964, **7**, 718-719; *CA*, **63**, 12440g (*cryst struct*)
 Habermehl, G.G. *et al.*, *Heterocycles*, 1976, **5**, 127-134 (*synth*)
 Roseghini, M. *et al.*, *Z. Naturforsch., C*, 1976, **31**, 118-120 (*occur*)
 Ciminiello, P. *et al.*, *J. Nat. Prod.*, 1994, **59**, 1564-1569 (*occur*)

4,5,6,7-Tetrahydro-1H-imidazo[4,5-c]pyridine-6-carboxylic acid, 9CI **T-193**
Spinacine



C₇H₉N₃O₂ 167.167

(S)-form [59981-63-4]

Isol. from the crab *Crangon vulgaris*, from the liver of the shark *Acanthias vulgaris* and from spinach *Spinacia oleracea*. Also isol. from the roots of *Panax ginseng*. Mp 265°. [α]_D²⁰ -174.6 (H₂O). Forms a sesquihydrate.

Hydrochloride: Mp 286-287° (279-280°).

l-Me: [133807-71-3]

C₈H₁₁N₃O₂ 181.194

Cryst. (MeOH/Et₂O) (as dihydrochloride). Mp 287-290° dec. (dihydrochloride). [α]_D²³ -9597 (c, 1.11 in H₂O).

3-Me: [114787-98-3]

C₈H₁₁N₃O₂ 181.194

Cryst. (MeOH/Et₂O) (as dihydrochloride). Mp 262-265° dec. (dihydrochloride). [α]_D²³ -93.5 (c, 1.05 in H₂O).

(±)-form

Synthetic. Mp 265°.

Ackermann, D. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1941, **268**, 277-282; 1949, **284**, 129-131; 1962, **328**, 275-276 (*isol, struct, synth*)

Vitali, T. *et al.*, *Gazz. Chim. Ital.*, 1964, **94**, 296-305 (*synth*)

Nardelli, M. *et al.*, *Ric. Sci., Parte 2: Sez. A*, 1964, **7**, 718-719; *CA*, **63**, 12440g (*cryst struct*)


Andreotti, G.D. *et al.*, *Gazz. Chim. Ital.*, 1971, **101**, 625-634 (*cryst struct*)

Eilazyan, O.G. *et al.*, *CA*, 1982, **96**, 85467n (*synth*)

Han, Y.N. *et al.*, *Arch. Pharmacol. Res.*, 1987, **10**, 258-259; *CA*, **108**, 201742r (*isol*)

Klutchko, S. *et al.*, *J. Het. Chem.*, 1991, **28**, 97-108 (*synth, pmr, derivs*)

1,6,7,9-Tetrahydro-6-imino-1,9-dimethyl-8H-purin-8-one, 9CI **T-194**
1,9-Dimethyl-6-imino-8-oxapurine [98601-03-7]



C₇H₉N₅O 179.181

Isol. from the marine sponge *Hymeniacion sanguinea*. Mp 300° (synthetic). Isol. and struct. elucidated as the Ac deriv.


N-Ac: [98601-04-8]

Cryst. (CHCl₃). Mp 245-246°.

Cimino, G. *et al.*, *J. Nat. Prod.*, 1985, **48**, 523-528 (*isol, uv, ir, pmr, ms, cryst struct*)

Fujii, T. *et al.*, *Heterocycles*, 1988, **27**, 1145-1148 (*synth, ir, pmr*)

5,6,7,7a-Tetrahydro-1H-indol-2(3H)-one **T-195**
Thomandersine



C₈H₁₁NO 137.181

Alkaloid from leaves of *Thomandersia laurifolia* (Acanthaceae). Isol. and characterised as the acetate.

N-Ac: [167105-95-5]

C₁₀H₁₃NO₂ 179.218

Brown oil.

Δ^{3,3a}-Isomer: 4,5,6,7-Tetrahydro-1H-indol-2(3H)-one. Isothomandersine

C₈H₁₁NO 137.181

From leaves of *Thomandersia laurifolia* (Acanthaceae). Isol. and characterised as the acetate.

Δ^{3,3a}-Isomer, N-Ac: [167105-96-6]


C₁₀H₁₃NO₂ 179.218

Fine brown needles (hexane/EtOAc). Mp 92°.

Ngadjui, B.T. *et al.*, *Phytochemistry*, 1995, **39**, 1249 (*isol, uv, ir, pmr, cmr, ms, struct*)

1,2,3,4-Tetrahydro-6-isoquinolinol, 9CI, 8CI **T-196**

1,2,3,4-Tetrahydro-6-hydroxyisoquinoline. 6-Hydroxy-1,2,3,4-tetrahydroisoquinoline



C₉H₁₁NO 149.192

Cryst. (EtOH). Mp 192-193°.

Hydrochloride: [66393-01-9]

Mp 222-224° (233°).

Hydrobromide: [59839-23-5]

Cryst. (EtOH/Et₂O). Mp 196.7-197.8°.

N-Me: 1,2,3,4-Tetrahydro-6-hydroxy-2-methylisoquinoline. Longimammosine

[14097-39-3]

C₁₀H₁₃NO 163.219

Alkaloid from *Dolichothele longimamma* (Cactaceae). Cryst. (EtOH). Mp 180-182°.

▶NX6030700

N-Me; hydrochloride: [57196-60-8]

Cryst. (EtOH/Et₂O). Mp 236°.

Me ether: 1,2,3,4-Tetrahydro-6-methoxyisoquinoline, 9CI. Longimammatine

[42923-77-3]

[57196-62-0]

C₁₀H₁₃NO 163.219

Alkaloid from *Dolichothele*

longimamma (Cactaceae). Plates (EtOH/Et₂O)(as hydrochloride). Mp 244-245.5° (238-239°)(hydrochloride).

▶NX5082000

Buck, J.S. *et al.*, *J.A.C.S.*, 1934, **56**, 1769-1771 (*synth, Longimammosine*)

Schenker, F. *et al.*, *J. Het. Chem.*, 1971, **8**, 665-668 (*synth, pmr, Longimammatine*)

Ranieri, R.L. *et al.*, *J.O.C.*, 1976, **41**, 319-323 (*isol, uv, ir, pmr, ms, synth, Longimammatine, Longimammosine*)

Mata, R. *et al.*, *Phytochemistry*, 1983, **22**,

1263-1270 (*cmr, Longimammatine*)

Sall, D.J. *et al.*, *J. Med. Chem.*, 1987, **30**, 2208-2216 (*pmr, ir, cmr, ms*)

Shinohara, T. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 919-927 (*Me ether*)

Zhong, H.M. *et al.*, *Org. Process Res. Dev.*, 2007, **11**, 463-465 (*Me ether*)

1,2,3,4-Tetrahydro-7-isoquinolinol, 9CI **T-197**

1,2,3,4-Tetrahydro-7-hydroxyisoquinoline.

7-Hydroxy-1,2,3,4-tetrahydroisoquinoline [30798-64-2]

C₉H₁₁NO 149.192

Bp₁₈ 210-220°.

Hydrobromide: [110192-19-3]

Cryst. (EtOH). Mp 210.3-211.4°.

Me ether: 1,2,3,4-Tetrahydro-7-methoxyisoquinoline, 9CI. Weberidine

[43207-78-9]

C₁₀H₁₃NO 163.219

Alkaloid from *Pachycereus weberi*

(Cactaceae). Bp_{0.1} 115° (*synth.*).

Me ether; hydrochloride: [1745-05-7]

Cryst. (EtOH/Et₂O). Mp 233° (228°).

Pictet, A. *et al.*, *Ber.*, 1911, **44**, 2036-2045

(*synth*)

Schenker, F. *et al.*, *J. Het. Chem.*, 1971, **8**, 665-668 (*synth, pmr*)

Moniot, J.L. *et al.*, *Heterocycles*, 1978, **9**, 145-152 (*pmr*)

Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673-678 (*Weberidine, isol, uv, ir, pmr, ms, synth, deriv*)

Mata, R. *et al.*, *Phytochemistry*, 1983, **22**,

1263-1270 (*cmr, Weberidine*)

Sall, D.J. *et al.*, *J. Med. Chem.*, 1987, **30**, 2208-2216 (*synth, pmr, ir, cmr, ms*)

Shinohara, T. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 919-927 (*Me ether, synth*)

Kuczniarz, R. *et al.*, *J. Med. Chem.*, 1998, **41**, 4983-4994 (*synth, cmr*)

Ueno, H. *et al.*, *J. Med. Chem.*, 2005, **48**, 3586-3604 (*synth*)

1,2,3,4-Tetrahydro-8-isoquinolinol, 9CI, 8CI **T-198**

1,2,3,4-Tetrahydro-8-hydroxyisoquinoline.

8-Hydroxy-1,2,3,4-tetrahydroisoquinoline [32999-37-4]

C₉H₁₁NO 149.192

Cryst. by subl. Mp 181-181.5° (*in vacuo*).

Hydrochloride: [32999-38-5]

Cryst. (1M HCl). Mp 271-272° dec. (*in vacuo*).

Hydrobromide: [110192-20-6]

Cryst. (EtOH/Et₂O). Mp 257.5-258.7°.

N-Me: 1,2,3,4-Tetrahydro-8-hydroxy-2-methylisoquinoline, 9CI. Longimammidine

[14788-32-0]

C₁₀H₁₃NO 163.219

Alkaloid from *Dolichothele longimamma* (Cactaceae). Mp 177° (173-175°).

N-Me; hydrochloride: [34222-77-0]
Cryst. (EtOH/Et₂O). Mp 246-247°.


Me ether: 1,2,3,4-Tetrahydro-8-methoxy-isoquinoline
[34146-68-4]
C₁₀H₁₃NO 163.219
Mp 38-39°.

Me ether, hydrochloride: [24693-40-1]
Cryst. (EtOH/Et₂O). Mp 260-264°.

Okamoto, M. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 168-172 (*N-Me, synth*)
Schenker, F. *et al.*, *J. Het. Chem.*, 1971, **8**, 665-668 (*synth, uv, ir, pmr*)
Ranieri, R.L. *et al.*, *J.O.C.*, 1976, **41**, 319-323 (*N-Me, isol, uv, pmr, ms, struct, synth*)
Gray, R.W. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 315-319 (*N-Me, synth*)
Kashdan, D.S. *et al.*, *J.O.C.*, 1982, **47**, 2638-2643 (*Me ether, synth, pmr*)
Sall, D.J. *et al.*, *J. Med. Chem.*, 1987, **30**, 2208-2216 (*synth, pmr, ir, cmr, ms*)
Shinohara, T. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 919-927 (*Me ether, synth*)

Tetrahydroalathyrine T-199

$\alpha,2$ -Diamino-1,4,5,6-tetrahydro-4-pyrimidinepropanoic acid, 9CI
[72748-96-0]




C₇H₁₄N₄O₂ 186.213
Amino acid from seeds of *Lonchocarpus costaricensis*. Cryst. (Me₂CO aq.) (as hydrochloride). Mp 258-259° (hydrochloride). [α]_D²⁵ -18.9 (c, 0.175 in H₂O) (hydrochloride).

Fellows, L.E. *et al.*, *Phytochemistry*, 1979, **18**, 1333 (*isol*)

1,2,3,4-Tetrahydro-ar-methoxy-1-methylisoquinoline T-200

[93474-21-6]




C₁₁H₁₅NO 177.246
Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).

Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109-114 (*occur*)

1,2,3,4-Tetrahydro-4-methoxy-2-methyl-8-quinolinecarboxylic acid T-201

Helquinoline



C₁₂H₁₅NO₃ 221.255


(2*RS*,4*RS*)-form

Prod. by the marine-derived *Janibacter limosus* Hel 1. Antibacterial and antifungal agent. Oil. λ_{\max} 221 (log ϵ 4.05); 261 (log ϵ 3.67); 349 (log ϵ 3.49) (MeOH). λ_{\max} 201 (log ϵ 3.96); 223 (log ϵ 4.03); 261 (log ϵ 3.59); 358 (log ϵ 3.47) (MeOH/HCl).

Asolkar, R.N. *et al.*, *J. Antibiot.*, 2004, **57**, 17-23 (*isol, uv, pmr, cmr*)

1,2,3,4-Tetrahydro-1-methyl- β -carboline T-202

2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indole. **Tetrahydroharman**. *Calligonine*. *Eleagnine*



C₁₂H₁₄N₂ 186.256

(*R*)-form [2254-36-6]

Mp 144-148°. [α]_D²⁰ +82 (c, 1 in EtOH). *N*^b-*Me*: [148151-46-6] Synthetic. Mp 116-118°. [α]_D²⁵ -37 (c, 2 in CHCl₃).

5-Methoxy: 2,3,4,9-Tetrahydro-5-methoxy-1-methyl-1*H*-pyrido[3,4-*b*]indole.

5-Methoxytetrahydroharman
[32640-61-2]

C₁₃H₁₆N₂O 216.282
Alkaloid from leaves of *Banisteriopsis argentea* (Malphigiaceae). Pale brown gum. [α]_D²⁵ +34 (c, 0.83 in CHCl₃). λ_{\max} 216 (log ϵ 4.55); 290 (sh) (log ϵ 3.66) (EtOH).

7-Methoxy: 1,2,3,4-Tetrahydro-7-methoxy-1-methyl- β -carboline. **Tetrahydroharmine**. *Leptaflorine*
[17019-01-1]

[7671-30-9, 7759-46-8]
C₁₃H₁₆N₂O 216.282

Alkaloid from *Banisteriopsis caapi*, *Banisteriopsis argentea*, *Banisteriopsis inebrians*, *Leptactina densiflora*, *Peganum harmala* and snuff used by South American Indians (Malphigiaceae, Rubiaceae, Zygophyllaceae). Also obt. by resoln. of the racemate. Mp 198.4-199.8° Mp 210-211° (by resoln.). [α]_D²⁵ +32 (CHCl₃). [α]_D +70.9 (CHCl₃) (by resoln.). Physical constants refer to the alkaloid isol. from *B. caapi*. Spec. rotns. of isolates from other spp. not available.

(*S*)-form [23844-21-5]

Synthetic. Mp 146-148°. [α]_D²⁰ -81.6 (c, 1 in EtOH).

N^b-*Me*: 2,3,4,9-Tetrahydro-1,2-dimethyl-1*H*-pyrido[3,4-*b*]indole. **1,2,3,4-Tetrahydro-1,2-dimethyl- β -carboline**
[32640-60-1]

C₁₃H₁₆N₂ 200.283
Alkaloid from *Banisteriopsis argentea* (Malphigiaceae). Mp 118-120° synthetic. [α]_D²⁵ +36 (c, 2 in CHCl₃) (synthetic)(+26). [α]_D²⁵ +26 (c, 0.72 in

CHCl₃) (natural). The higher opt. rotn. refers to synthetic material. λ_{\max} 226 (log ϵ 4.5); 278 (log ϵ 3.83); 293 (log ϵ 3.81) (EtOH) (natural).

7-Methoxy:

Obt. by resoln. of the racemate. Mp 211-212°. [α]_D -69.6 (CHCl₃).

(\pm)-form [525-40-6]

[2506-10-7]

Alkaloid from *Leptactina densiflora* (Rubiaceae) and several spp. from Fabaceae and Chenopodiaceae. Mp 179-180°.

Hydrochloride: Mp 267° dec.

N-Benzoyl: Mp 168-169° Mp 194-195°.

N^b-*Me*: **Leptocladine**

[27297-47-8]

C₁₃H₁₆N₂ 200.283

Alkaloid from *Acacia complanata*, *Arthropytum leptocladum* and several other spp. in the Fabaceae and Chenopodiaceae. Mp 112°.

N^b-*Me, picrate*: Mp 182-184°.

5-Methoxy: [83788-97-0]

Cryst. (EtOAc/hexane). Mp 200-202°.

λ_{\max} 226 (log ϵ 4.25); 271 (log ϵ 3.59); 280 (log ϵ 3.52); 292 (log ϵ 3.53) (EtOH).

5-Methoxy, *N*^b-*Me*: [83789-00-8]

Cryst. (MeOH aq.). Mp 229-231°. λ_{\max} 226 (log ϵ 4.25); 271 (log ϵ 3.59); 280 (log ϵ 3.52); 292 (log ϵ 3.53) (EtOH).

7-Methoxy: [46501-01-3]

Synthetic. Needles (MeOH). Mp 198-199°.

(ξ)-form

N^b-*Oxide*: **Tetrahydroharman *N*^b-oxide**.

Calligonidine

[2288-25-7]

C₁₂H₁₄N₂O 202.255

Alkaloid from *Calligonum minimum* (Polygonaceae). Mp 132-133°.

N^b-*Oxide, hydrochloride*: Mp 215-215.5°.

N^b-*Ethoxycarbonyl*: 2-Ethoxycarbonyl-

1,2,3,4-tetrahydro-1-methyl- β -carboline.

2-Ethoxycarbonyltetrahydroharman

man

C₁₅H₁₈N₂O₂ 258.319

Alkaloid from *Croton moritibensis*.

Perkin, W.H. *et al.*, *J.C.S.*, 1919, **115**, 933-967 (*Tetrahydroharmine, synth*)

Paris, P.R. *et al.*, *Bull. Soc. Chim. Fr.*, 1957, 780-782 (*Tetrahydroharman, Tetrahydroharmine, isol, uv, ir*)

Hochstein, F.A. *et al.*, *J.A.C.S.*, 1957, **79**, 5735-5736 (*Tetrahydroharmine, isol*)

Absalamov, B.A. *et al.*, *CA*, 1963, **58**, 11412b (*oxide*)

Bernauer, K. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1075-1077 (*Tetrahydroharmine, isol*)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1966, **19**, 1539 (*Banisteriopsis argentea constits*)

Kobicova, Z. *et al.*, *Chem. Ind. (London)*, 1966, 1342 (*Tetrahydroharmine, abs config*)

O'Donovan, D.G. *et al.*, *J.C.S.(C)*, 1967, 1109-1110 (*biosynth*)

Ghosal, S. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1209-1212 (*Banisteriopsis argentea constits*)

Ghosal, S. *et al.*, *Planta Med.*, 1972, **21**, 200-203 (*Tetrahydroharmine, cmr*)


Bláha, K. *et al.*, *Coll. Czech. Chem. Comm.*, 1974, **39**, 3168-3176 (*resoln, config*)

Middleditch, B.S. *et al.*, *J. Chromatogr.*, 1976, **126**, 581-589 (*ms*)

Repke, D.B. *et al.*, *J. Het. Chem.*, 1982, **19**, 845-848 (5-methoxy, 5-methoxy-*N^b-Ne*, synth, bibl)
 Han, S.P. *et al.*, *Heterocycles*, 1985, **23**, 1671-1673 (*N^b-oxide*, synth)
 Chrisey, L.A. *et al.*, *Heterocycles*, 1990, **30**, 267-270 (Tetrahydroharmine, resoln)
 Qais, N. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 3338-3340 (synth)
 Peng Shiqi, *et al.*, *Annalen*, 1993, 137-140 (*N^b-Methyltetrahydroharman*, synth)
 Herraiz, T. *et al.*, *J. Agric. Food Chem.*, 2000, **48**, 4900-4904; 2004, **52**, 2652-2658 (hplc, occur, bibl)
 De Araújo-Júnior, V.T. *et al.*, *Pharm. Biol.*, 2004, **42**, 62-67 (2-Ethoxycarbonyl tetrahydroharman)

1,2,3,4-Tetrahydro-2-methyl-β-carboline T-203

2,3,4,9-Tetrahydro-2-methyl-1*H*-pyrido[3,4-*b*]indole, 9CI
 [13100-00-0]



C₁₂H₁₄N₂ 186.256
 Alkaloid from *Arthropytum leptocladum*, *Viola theiodora*, *Viola sebifera*, *Hammada leptoclada*, *Elaeagnus angustifolia*, *Phalaris aquatica*, *Phalaris arundinacea*, *Acacia simplicifolia*, *Anadenanthera peregrina*, *Banisteriopsis rusbyana*, *Testulea gabonensis*, *Gymnocranthera paniculata*, *Psychotria carthaginensis*, *Psychotria viridis* and *Flueggea microcarpa* (Chenopodiaceae, Myristicaceae, Eleagnaceae, Poaceae, Fabaceae, Malpighiaceae, Ochidaceae, Rubiaceae, Euphorbiaceae). Cryst. (MeOH). Mp 216-218°. λ_{max} 224 (log ε 3.5); 280 (log ε 3.9) (MeOH).

Hydrochloride:
 Cryst. (EtOH/Et₂O). Mp 245-247°.


Picrate:
 Cryst. (EtOH). Mp 195-197°.

Methiodide:
 Cryst. (EtOH). Mp 265-266°.

Bockelheide, V. *et al.*, *J.A.C.S.*, 1950, **72**, 2132 (synth)
 Johns, S.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1737 (isol, uv, ir, pmr, ms)
 Agurell, S. *et al.*, *Acta Chem. Scand.*, 1969, **23**, 903 (isol, ms)
 Allen, J.R.F. *et al.*, *Phytochemistry*, 1980, **19**, 1573 (rev, bibl)
 Kawanishi, K. *et al.*, *Phytochemistry*, 1985, **24**, 1373 (occur)
 Kumar, S. *et al.*, *Planta Med.*, 1985, 466 (isol, pmr, cmr, ms)
 Zhalolov, I.Z. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2003, **39**, 289-291 (cryst struct)

5,6,7,8-Tetrahydro-5-methyl-2-propylquinoline T-204

Mantella Alkaloid 189



C₁₃H₁₉N 189.3
 Trace alkaloid from skin extracts of the Madagascan frogs *Mantella* sp. cf. *madagascariensis*, *Mantella betsileo* and *Mantella laevigata*.

1,4,4a,8a-Tetrahydro: 1,4,4a,5,6,7,8,8a-Octahydro-5-methyl-2-propylquinoline.

Mantella Alkaloid 193D


C₁₃H₂₃N 193.331

Alkaloid from skin extracts of *Mantella* sp. cf. *madagascariensis*, *Mantella betsileo* and *Mantella laevigata*. Tentative struct.

Hexahydro: see Pumiliotoxin C, P-794
 Garaffo, H.M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1016 (isol, ir, ms)

1,2,5,6-Tetrahydro-1-methyl-3-pyridinecarboxylic acid, 9CI T-205

Arecaidine. *Arecaine*
 [499-04-7]



C₇H₁₁NO₂ 141.169
 Alkaloid from nuts of *Areca catechu* (betel nuts) (Arecaceae). Cholinergic and anthelmintic agent. Mp 223-224° dec. (anhyd. 232° dec.). pK_a 9.07 (25°). Log P -1.74 (calc). Strong base.

►QT2070800

Hydrochloride: Mp 262-263° (rapid heat) (250.5-262°).

Me ester: Arecoline

[63-75-2]

C₈H₁₃NO₂ 155.196

Alkaloid from *Areca catechu* (Arecaceae). Cathartic, vermifuge, hypotensive agent etc. Formerly used as a cholinergic agent; purgative and taenifuge in veterinary medicine. Liq. Bp₁₇ 94° Bp₇ 74°. pK_a 7.64 (25°). Log P 1.05 (calc).

►LD₅₀ (dog, scu) 5 mg/kg. LD₅₀ (rat, orl) 2500 mg/kg. QT2100000

Me ester, hydrochloride: [61-94-9]
 Mp 157-158°.

►QT2150000

Me ester, hydrobromide: [300-08-3]
 Mp 172°.

►QT2275000

Et ester: Homoarecoline

[28125-84-0]

[6027-10-7, 17210-50-3]

C₉H₁₅NO₂ 169.223

Constit. of *Areca catechu*. Oil.

Jahns, E. *et al.*, *Ber.*, 1888, **21**, 3404 (isol)

Wohl, A. *et al.*, *Ber.*, 1907, **40**, 4712 (struct, synth)

Chemnitzius, F. *et al.*, *J. Prakt. Chem.*, 1927, **117**, 147 (isol, deriv)

Mannich, C. *et al.*, *Ber.*, 1942, **75**, 1480 (synth)

Lyle, R.E. *et al.*, *J.O.C.*, 1955, **20**, 1761 (synth, uv)

Ma, J.C.N. *et al.*, *Can. J. Chem.*, 1965, **43**, 1849 (pmr)

Spiteller-Friedmann, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 104 (ms)

Nery, R. *et al.*, *Biochem. J.*, 1971, **122**, 503 (metab, tox, bibl)

Burrows, R.B. *et al.*, *Prog. Drug Res.*, 1973, **17**, 108 (rev)

Kozello, I.A. *et al.*, *Khim.-Farm. Zh.*, 1979, **13**, 61; *CA*, **92**, 110798y (synth)

Strunz, G.M. *et al.*, *Alkaloids* (Academic Press), 1985, **26**, 89 (rev)

Hollander, E. *et al.*, *Br. Med. Bull.*, 1986, **42**, 97 (pharmacol)


Kurihara, T. *et al.*, *Heterocycles*, 1990, **30**, 885-896 (Et ester, synth)

Martindale. *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 39

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, AQT625; AQU000; AQU250; AQT750

2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, 9CI T-206

1,2,3,4-Tetrahydro-1-methyl-β-carboline-3-carboxylic acid
 [5470-37-1]
 [143396-05-8 (cis-form)]



(1*R*,3*S*)-form

C₁₃H₁₄N₂O₂ 230.266

(1*R*,3*S*)-form [42438-72-2]

Constit. of cocoa and maca tubers (*Lepidium meyenii*). Cryst. (H₂O). Mp 242-244° (225-228° dec.). [α]_D²⁵ -69.1 (c, 1 in 1*M* HCl/MeOH 1:1).

N²-Me:

C₁₄H₁₆N₂O₂ 244.293

Mp 222-223° (as hydrochloride). [α]_D²⁵ -29.6 (c, 1 in 1*M* HCl MeOH/H₂O 1:1).

7-Hydroxy: 1,2,3,4-Tetrahydro-7-hydroxy-1-methyl-β-carboline-3-carboxylic acid. Brunnein B
 [955377-46-5]

C₁₃H₁₄N₂O₃ 246.265

Alkaloid from *Cortinarius brunneus*. Yellow solid. [α]_D²⁵ -35 (c, 0.03 in MeOH). λ_{max} 219 (log ε 4.4); 299 (log ε 3.4) (MeOH).

7-Hydroxy, 1,2-didehydro: 3,4-Dihydro-7-hydroxy-1-methyl-β-carboline-3-carboxylic acid. Brunnein A
 [955377-45-4]

C₁₃H₁₂N₂O₃ 244.249

Alkaloid from *Cortinarius brunneus*. Yellow-green solid. [α]_D²⁶ -22 (c, 0.02 in MeOH). λ_{max} 260 (log ε 3.3); 375 (log ε 3.6) (MeOH).

(1*S*,3*S*)-form [40678-46-4]

Alkaloid from the starfish *Lethasterias nanimensis chelifera*. Constit. of cocoa and garlic (*Allium sativum*). Cryst. (H₂O). Mp 293° (276-280° dec.). [α]_D²⁵ -106.6 (c, 1 in 1*M* HCl/MeOH 1:1).

(1*R**,3*R**)-form

6-Hydroxy: 1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-β-carboline-3-carboxylic acid. Hyrtioerectine B
 [42438-95-9 (1*S*,3*S*)-form]

C₁₃H₁₄N₂O₃ 246.265

Isol. from the sponge *Hyrtios erectus*. Amorph. solid. $[\alpha]_D^{25}$ -19.2 (c, 0.18 in MeOH). Stereochem. is 1*R*-3*R*-*rel* according to Youssef *et al.* (2005) but abstracted as 1*S*,3*S* by *CA*. λ_{\max} 231 (log ϵ 3.81); 276 (log ϵ 3.74); 357 (log ϵ 1.82) (MeOH).

(1*S*,3*S*)-form

Formed by Pictet-Spengler condensation between Acetaldehyde and Tryptophan, T-640 in nature or during food processing. Present in many foods. Alkaloid from the toxic fly agaric mushroom (*Amanita muscaria*). Shows little pharmacol. activity. Needles (H₂O). Mp 290° dec. $[\alpha]_D$ -142.8 (50 % Py aq.). λ_{\max} 219; 271; 277; 280 (sh); 287 (MeOH).

[22677-22-1]

Matsumoto, T. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 716-720 (*isol, uv, ir, struct*)

Brossi, A. *et al.*, *J. Med. Chem.*, 1973, **16**, 418-420 (1*R*,3*S*-*form*, 1*S*,3*S*-*form*, *synth, pharmacol, ord*)

Bobbitt, J.M. *et al.*, *J.O.C.*, 1980, **45**, 1978-1984 (*synth*)

Bosin, T.R. *et al.*, *J. Agric. Food Chem.*, 1986, **34**, 843-847 (*isol*)

Herraiz, T. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3057-3065; 1998, **46**, 3484-3490; 1999, **47**, 4883-4887; 2000, **48**, 4900-4904; 2004, **52**, 2652-2658 (*hplc, occur, bibl*)

Herraiz, T. *et al.*, *J. Chromatogr. A*, 1997, **765**, 265-277; 2000, **871**, 23-30 (*occur, gc, hplc*)

Piacenta, S. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 5621-5625 (1*R*,3*S*-*form, isol*)

Kicha, A.A. *et al.*, *Tet. Lett.*, 2003, **44**, 1935-1937 (*Lethasterias alkaloid, isol, pmr, cmr*)

Kuo, F.-M. *et al.*, *Tetrahedron*, 2004, **60**, 12075-12084 (*synth*)


Youssef, D.T.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1416-1419 (*Hyrtioerectine B*)

Teichert, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1529-1531 (*Brunneins A,B*)

1,4,5,6-Tetrahydro-2-methyl-4(6)-pyrimidinecarboxylic acid, 9CI

T-207

Ectoine. *Pyrostatin B*
[117305-84-7]



(S)-form

C₆H₁₀N₂O₂ 142.157

Exists as zwitterion in cryst. Pyrostatin B tentatively assigned this struct. after revision in 2006.

(S)-form [96702-03-3]

Isol. from halophilic bacteria such as *Ectothiorhodospira halochloris* and *Halomonas elongata*. Also from the marine-derived *Streptomyces* sp. SA-3501 (Pyrostatin B). Osmoprotective agent. Cryst. V. sol. H₂O. Mp 280°. $[\alpha]_D^{20}$ +140 (c, 1.0 in MeOH).

(±)-form

Cryst. (EtOH/Et₂O)(as hydrochloride). Mp 280° (hydrochloride). CAS no. refers to hydrochloride.

Galinski, E.W. *et al.*, *Eur. J. Biochem.*, 1985, **149**, 135-139 (*isol, cmr, ms, ir*)

Schuh, W. *et al.*, *Z. Naturforsch., C*, 1985, **40**, 780 (*cryst struct*)

Inbar, L. *et al.*, *Eur. J. Biochem.*, 1993, **214**, 897-906 (*isol, cryst struct, pmr*)

Filsak, G. *et al.*, *Z. Naturforsch., C*, 1994, **49**, 18 (*synth, pmr, cmr*)


Him-di-Kabbab, S. *et al.*, *Synth. Commun.*, 1995, **25**, 2223 (*synth, pmr, cmr*)

Castellanos, L. *et al.*, *Org. Lett.*, 2006, **8**, 4967-4970 (*Pyrostatin B*)

5,6,7,8-Tetrahydro-4-methylquinoline

T-208

5,6,7,8-Tetrahydrolepidine
[28971-03-1]

C₁₀H₁₃N 147.219

Alkaloid from the roots of *Glycyrrhiza uralensis* (Chinese licorice) (Fabaceae). Liq. Spar. sol. H₂O. Bp₁₁ 122°.

Hydrochloride: Mp 203-204°.

Picrate: [60166-61-2]

Mp 178-181° (170°).

Methiodide: Mp 183°.

v. Braun, J. *et al.*, *Ber.*, 1923, **56**, 1343 (*synth*)

Booth, H. *et al.*, *J.C.S. Perkin 2*, 1976, 759 (*pmr*)

Reimann, E. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1983, **316**, 210 (*synth*)

Koyama, J. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 2601 (*synth*)


Konno, K. *et al.*, *Tet. Lett.*, 1986, **27**, 3865 (*synth*)

Murahashi, S.-I. *et al.*, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 2968 (*synth*)

Han, Y.N. *et al.*, *Arch. Pharmacol. Res.*, 1990, **13**, 101; *CA*, **113**, 227984 (*isol*)

2,3,8*a*-Tetrahydro-3*a*-nitro-1*H*-pyrrolo[2,3-*b*]indole

T-209

C₁₀H₁₁N₃O₂ 205.216


Isol. from the marine-derived *Flavobacterium* sp. T436.

Schuhmann, I. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*isol*)

5,6,7,8-Tetrahydro-2-nonylquinoline, 9CI

T-210

[59652-32-3]

C₁₈H₂₉N 259.434


Isol. from egg tar. λ_{\max} 274 (MeOH) (Berdy). λ_{\max} 282 (MeOH-HCl) (Berdy).

Tsuji, K. *et al.*, *Yakugaku Zasshi*, 1976, **96**, 479-483 (*isol*)

Kosuge, T. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 611-619 (*isol*)

2,3,6,11-Tetrahydro-3-oxo-1*H*-indolizino[8,7-*b*]indole-5,11*b*(5*H*)-dicarboxylic acid, 9CI

T-211

(5*S*,11*bR*)-formC₁₆H₁₄N₂O₅ 314.297**(5*S*,11*bR*)-form** [82503-93-3]

Alkaloid from *Clerodendrum trichotomum* (Verbenaceae). Mp 189-190° (as di-Me ester). $[\alpha]_D$ +110 (c, 0.219 in MeOH) (di-Me ester). CAS no. refers to di-Me ester.

(5*S*,11*bS*)-form [82535-74-8]


Alkaloid from *Clerodendrum trichotomum* (Verbenaceae). $[\alpha]_D$ -107 (c, 0.205 in MeOH) (as di-Me ester). CAS no. refers to di-Me ester.

Toyoda, Y. *et al.*, *Chem. Lett.*, 1982, 903 (*ir, pmr, ms, struct, synth*)

Irikawa, H. *et al.*, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 880 (*isol, ir, pmr, cmr, ms, cd, struct*)

Tetrahydropalmatine T-212*Corydalis B. Gindarine. Caseanine.**Rotundine*†. *Hindarine*

[10097-84-4]



(R)-form

C₂₁H₂₅NO₄ 355.433

Hindarine is a transliteration error for Gindarine, from a Russian paper. Analgesic, sedative and hypnotic agent (hypnotic activity exceeds that of Morphine, M-704). Shows strong papaverine-like activity. Active against gram-positive bacteria. Log P 2.45 (uncertain value) (calc).

(R)-form [3520-14-7]

Alkaloid from several *Corydalis* spp., *Berberis tinctoria* and *Coptis tecta* (Papaveraceae, Menispermaceae, Berberidaceae, Ranunculaceae). Mp 143°. $[\alpha]_D^{20}$ +291 (EtOH).

Hydrochloride: Mp 266°.

(S)-form [483-14-7]

Alkaloid from *Corydalis* spp., *Pachypodanthium confine* (Papaveraceae, Annonaceae) and tubers of *Stephania glabra*. Active against gram-positive bacteria. Mp 141-142°. $[\alpha]_D^{21}$ -290.8 (EtOH).

▶HQ1792500

Hydrate: Mp 115-116°.

N-Oxide (R-): *Epicorynoxidine*
[58000-48-9]

C₂₁H₂₅NO₅ 371.432

From *Corydalis koidzumiana* (Papaveraceae). Cryst. (Me₂CO). Mp 193-195° dec. [α]_D -2.

N-Oxide(S-): Corynoxidine

[57906-85-1]

C₂₁H₂₅NO₅ 371.432

Alkaloid from *Corydalis koidzumiana* (Papaveraceae). Cryst. (Me₂CO). Mp 183.6-184° dec. [α]_D -57.

O²-De-Me: see Isocorypalmine, I-210

O³-De-Me: see Corypalmine, C-694

O¹⁰-De-Me: see Corydalmine, C-677

O²,O⁹-Di-de-Me: see Scoulerine, S-164

O²,O¹⁰-Di-de-Me: see Stepholidine, S-547

O³,O⁹-Di-de-Me: see Isoscoulerine, I-315

O³,O¹⁰-Di-de-Me: see Discretamine, D-856

O⁹,O¹⁰-Di-de-Me: see 9,10-Dihydroxy-2,3-dimethoxytetrahydroprotoberberine, D-570

(±)-form [2934-97-6]

Alkaloid from *Corydalis* spp. (Papaveraceae). Scales (MeOH). Mp 151° (147°).

Hydrochloride:

Needles (H₂O). Mp 215°.

Hydroiodide:

Orange-yellow needles. Mp 241° dec.

Späth, E. *et al.*, *Ber.*, 1923, **56**, 875-879 (*isol*)
Kametani, T. *et al.*, *J.C.S.(C)*, 1967, 530-532 (*synth, pmr, ir*)

Cava, M.P. *et al.*, *J.O.C.*, 1968, **33**, 2785-2789 (*S-form, isol*)

Tani, C. *et al.*, *Chem. Lett.*, 1975, 1081-1084 (*Corynoxidine, Epicorynoxidine*)

Hughes, D.W. *et al.*, *Can. J. Chem.*, 1976, **54**, 2252-2260 (*cmr*)

Tourwe, D. *et al.*, *Org. Magn. Reson.*, 1977, **9**, 341-346 (*pmr*)

Narasimhan, N.S. *et al.*, *Tetrahedron*, 1983, **39**, 1975-1982 (*synth*)

Wang, S. *et al.*, *Yiyao Gongye*, 1988, **19**, 444-446; *CA*, **110**, 173517s (*synth*)

Hussain, R.A. *et al.*, *Heterocycles*, 1989, **29**, 2257-2260 (*pmr, cmr*)

Orito, K. *et al.*, *Org. Prep. Proced. Int.*, 1989, **21**, 309-314 (*synth*)

Simeon, S. *et al.*, *Pharmazie*, 1989, **44**, 593-597 (*activity*)

Pyne, S.G. *et al.*, *J.O.C.*, 1990, **55**, 1932-1936 (*synth, pmr, cmr, ms*)

Kessar, S.V. *et al.*, *J.O.C.*, 1992, **57**, 6716-6720 (*synth*)

Ribár, B. *et al.*, *Acta Cryst. C*, 1993, **49**, 1691-1693 (*cryst struct*)

Lugar, P. *et al.*, *Acta Cryst. C*, 1998, **54**, 1977-1980 (*cryst struct*)


Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999, **37**, 195-202 (*N-15 nmr*)

Blanchfield, J.T. *et al.*, *Phytochemistry*, 2003, **63**, 711-720 (*S-form, isol, pmr, cmr*)

Boudou, M. *et al.*, *J.O.C.*, 2005, **70**, 9486-9494 (*synth*)

2,3,4,5-Tetrahydro-6-(1,3-pentadienyl)pyridine, 9CI T-213

1-[2-(3,4,5,6-Tetrahydropyridyl)]-1,3-pentadiene



C₁₀H₁₅N 149.235

(E,E)-form

Antibiotic NA 337. NA 337

[53696-65-4]

Alkaloid from *Streptomyces* sp. NA-337 and actinomyces strain MD736-C6. N-Methyltransferase inhibitor. Sol. H₂O, MeOH, butanol; poorly sol. Me₂CO, hexane. Originally considered to be a pyrrolidine. Erroneously reported to have positive opt. rotn. λ_{\max} 253 (ϵ 28000); 261 (ϵ 34200) (cyclohexane) (Berdy). λ_{\max} 303 (ϵ 33000) (HCl) (Berdy). λ_{\max} 267 (NaOH) (Berdy).

► LD₅₀ (mus, ivn) 5.9 mg/kg. UT8415000

Hydrochloride: [62928-54-5]

Plates. Mp 190-200° dec. Sinters at 155-160°, browns at 170°, dark-brown melt.

Picrate:

Yellow granules (EtOAc/MeOH).


[54583-43-6]

Onda, M. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 2916-2920; 1975, **23**, 2462-2463 (*isol, cmr, pmr, struct*)

Kumada, Y. *et al.*, *J. Antibiot.*, 1974, **27**, 726-728 (*isol, pmr, cmr*)

2,3,4,5-Tetrahydro-6-(4-pentenyl)pyridine, 9CI T-214

2-(4-Penten-1-yl)- Δ^1 -piperideine [83688-89-5]



C₁₀H₁₇N 151.251


Component of the venom of a *Solenopsis* sp. ant. Unique among the ant venom alkaloids in having an even no. of C atoms.

Jones, T.H. *et al.*, *Tetrahedron*, 1982, **13**, 1949 (*isol, ms, pmr, struct, synth*)

Tetrahydropentoxilyne T-215

1-C-(3-Carboxy-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)pentitol, 9CI. 1,2,3,4-Tetrahydro-1-(1,2,3,4,5-pentahydroxypentyl)- β -carboline-3-carboxylic acid

[154204-09-8]



C₁₇H₂₂N₂O₇ 366.37


Isol. from human urine and various fruit juices. The fruit juice isolate characterised as two stereoisomers having *cis*- and *trans*- stereochem. at the tetrahydrocarboline ring, but with undetermined pentitol config.

Horiuchi, K. *et al.*, *J. Biochem. (Tokyo)*, 1994, **115**, 362-366 (*isol, pmr, cmr*)

Herraiz, T. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 4690-4695 (*isol, pmr, cmr, ms*)

2,3,4,5-Tetrahydro-6-pentylpyridine, 9CI T-216

[5832-23-5]



C₁₀H₁₉N 153.267

Alkaloid from *Conium maculatum*. Mp 65-67° (as picrate). Bp₁₁ 90°.


Dietzsch, K. *et al.*, *Z. Chem.*, 1966, **6**, 98-102 (*synth*)

McGrane, P.L. *et al.*, *J.A.C.S.*, 1992, **114**, 5459-5460 (*synth*)

Lang, D.G. *et al.*, *CA*, 1999, **130**, 136554u (*isol*)

1,2,3,4-Tetrahydro-6-propionylpyridine T-217

1-(1,4,5,6-Tetrahydro-2-pyridinyl)-1-propanone, 9CI. 1-(3,4,5,6-Tetrahydro-2-pyridinyl)-1-propanone, 9CI. Ethyl 3,4,5,6-tetrahydro-2-pyridyl ketone, 8CI. 3,4,5,6-Tetrahydro-2-propionylpyridine. PTP [80933-75-1, 80933-74-0]



C₈H₁₃NO 139.197

Tautomeric; 1H-form predominates. Formed by thermal treatment of proline and glucose mixtures.

Constit. of the leaves of *Semnostachya menglaensis*. Liq. with intense roast-smelling odour. Bp₇ 71-76°.

De Kimpe, N. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 1515-1519 (*synth, pmr, cmr, ms*)

Hofmann, T. *et al.*, *J. Agric. Food Chem.*, 1998, **46**, 2721-2726 (*occur, ms*)


Naef, R. *et al.*, *J. Agric. Food Chem.*, 2005, **53**, 9161-9164 (*isol, synth*)

2,3,4,5-Tetrahydro-6-(1-propenyl)pyridine T-218

2-(1-Propenyl)- Δ^1 -piperideine

[16543-92-3]


[16543-93-4]



C₈H₁₃N 123.197

Incorr. named as 2-(2-propenyl)- Δ^1 -piperideine in the lit. Alkaloid from leaves of *Punica granatum* (pomegranate) (Punicaceae). Cryst. (as hydrochloride). Mp 112-114° (hydrochloride). Unstable, only obt. from fresh extracts. Hydrochloride dec. in moist air.

Roberts, M.F. *et al.*, *Phytochemistry*, 1967, **6**, 711 (*isol, ir, pmr, struct*)

1,2,3,4-Tetrahydro-1-propyl- β -carboline T-2192,3,4,9-Tetrahydro-1-propyl-1H-pyridolo[3,4-b]indole, 9CI. **Komaroidine**C₁₄H₁₈N₂ 214.31**(ξ)-form**Alkaloid from the aerial parts of *Nitraria komarovii* and *Nitraria schoberi*. Cryst. (EtOH/Me₂CO). Mp 207-208°.N²-Ac: 2-Acetyl-1,2,3,4-tetrahydro-1-propyl- β -carboline. N²-Acetylkomaroidine C₁₆H₂₀N₂O₂ 272.346Alkaloid from the aerial parts of *Nitraria komarovii* and *Nitraria schoberi*. Cryst. (EtOH/Me₂CO). Mp 229-230°.6-Hydroxy: 1,2,3,4-Tetrahydro-6-hydroxy-1-propyl- β -carboline. **Schobericine** C₁₄H₁₈N₂O 230.309Alkaloid from the aerial parts of *Nitraria schoberi*. Cryst. (EtOH). Mp 193-194°.Tulyaganov, T.S. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2006, **42**, 198-200 (*isol. synth., pmr*)**1,2,3,4-Tetrahydropyridine** T-220 Δ^2 -Piperidine
[37497-65-7]C₅H₉N 83.133


Liq. Dimerises on heating.

Hydrochloride: Mp 230°.**Hydrobromide**: Mp 178°.Ac: **1-Acetyl-1,2,3,4-tetrahydropyridine**C₇H₁₁NO 125.17Alkaloid from leaves and twigs of *Dichilus strictus*, *Dichilus reflexus*, *Dichilus lebeckioides*, *Dichilus pilosus* and *Dichilus gracilis* (Fabaceae). Bp 219.5-220.5°.Paal, C. *et al.*, *Ber.*, 1901, **34**, 2757 (*synth*)Van Wyk, B.-E. *et al.*, *Biochem. Syst. Ecol.*, 1988, **16**, 471 (*deriv. isol*)**1,2,3,6-Tetrahydro-2-pyridinecarboxylic acid, 9CI** T-221

1,2,3,6-Tetrahydropicolinic acid, 8CI.

Baikain

[498-98-6]

**(S)-form**C₆H₉NO₂ 127.143**(S)-form** [31456-71-0]Amino acid present in *Baikiaea plurijuga*, *Caesalpinia tinctoria*, red algae and red seaweeds (Fabaceae, Caesalpinaceae).Also in *Russula subnigricans*. Neurotransmission inhibitor. Prisms (MeOH). V. sol. H₂O. spar. sol. EtOH, insol. Me₂CO, EtOAc, Et₂O, C₆H₆. [α]_D²⁰ -182.6 (c, 0.3 in H₂O).**Hydrochloride**:

Prisms (MeOH or EtOH). Mp 264°.

[α]_D²⁰ -90.1 (H₂O).**Picrate**:Needles or plates (H₂O). Mp 172-173°.**Me ester**:C₇H₁₁NO₂ 141.169Bp₁₅ 110-112°.**Me ester; hydrochloride**:Prisms (MeOH/Me₂CO). Mp 164°.**N-Benzoyl**:C₁₃H₁₃NO₃ 231.251Prisms (EtOAc); needles (Et₂O or H₂O). Mp 178-179°. [α]_D²⁰ -91.9.**N-Di-Me, betaine: Baikain betaine**


[114622-09-2]

C₈H₁₃NO₂ 155.196Isol. from the red alga *Pterocladia capillacea*. Cryst. (2-propanol). Mp 184-186°. [α]_D²⁵ -103 (c, 1.6 in H₂O).**(±)-form** [7200-16-0]Small prisms (MeOH/Me₂CO). Mp 251-254° dec Mp 273-274° dec.**Hydrochloride**: [111257-58-0]

Cryst. (MeOH/EtOAc). Mp 264° dec.

King, T.J. *et al.*, *J.C.S.*, 1950, 3590 (*isol. struct. abs config*)Burgstahler, A.W. *et al.*, *J.O.C.*, 1960, **25**, 489 (*synth*)Watson, R. *et al.*, *Phytochemistry*, 1973, **12**, 617 (*isol*)Impellizzeri, G. *et al.*, *Phytochemistry*, 1975, **14**, 1549-1557 (*algae, isol*)Maeda, M. *et al.*, *Agric. Biol. Chem.*, 1980, **44**, 2725 (*isol. pmr, cmr, ir, occur, ms*)Miyazawa, K. *et al.*, *CA*, 1983, **98**, 194969q (*occur*)Sciuto, S. *et al.*, *J. Nat. Prod.*, 1988, **51**, 322-325 (*Baikain betaine*)Kinderman, S.S. *et al.*, *Adv. Synth. Catal.*, 2002, **344**, 736-748 (*S-form, synth, ir, pmr, cmr*)Ginesta, X. *et al.*, *Tet. Lett.*, 2002, **43**, 779-782 (*synth*)Chang, M.-Y. *et al.*, *Heterocycles*, 2006, **68**, 2365-2373 (*S-form, synth*)**1,2,3,6-Tetrahydro-4-pyridinecarboxylic acid, 9CI** T-222**Iso-guvacine**

[64603-90-3]

C₆H₉NO₂ 127.143Alkaloid from *Areca catechu* (betel nut). GABA_A-receptor agonist. Shows anxio-lytic-like activity in animal model. Pharmacol. tool. Mp 220°. pK_{a1} 3.6; pK_{a2} 9.8.**Hydrobromide**: [68947-42-2]Cryst. (2-propanol aq./Et₂O). Mp 284-287° dec.**Me ester**: [70684-82-1]C₇H₁₁NO₂ 141.169

Cryst. (MeCN) (as hydrochloride). Mp 159-161° (hydrochloride).

N-tert-Butyloxycarbonyl, Me ester:

[184368-74-9]

C₁₂H₁₉NO₄ 241.286

No phys. props. reported.


N-tert-Butyloxycarbonyl, Et ester:

[906663-30-7]

C₁₃H₂₁NO₄ 255.313

Characterised spectroscopically.

[68547-97-7]

Winterstein, E. *et al.*, *Hoppe-Seyler's Z.**Physiol. Chem.*, 1918, **104**, 48-53Krogsgaard-Larsen, P. *et al.*, *J. Neurochem.*, 1978, **30**, 1377 (*synth, sar, ir, pmr*)Krogsgaard-Larsen, P. *et al.*, *Eur. J. Med. Chem. (Chim. Ther.)*, 1979, **14**, 157; 1980, **15**, 529 (*synth, pharmacol*)Japan. Pat., 1979, 79 36 275; *CA*, **91**, 56827w (*synth*)Krogsgaard-Larsen, P. *et al.*, *Acta Chem. Scand., Ser. B*, 1981, **35**, 311 (*bibl, rev*)Falch, E. *et al.*, *J. Med. Chem.*, 1981, **24**, 285 (*sar*)Lipkowitz, K.B. *et al.*, *J. Mol. Struct.*, 1989, **195**, 65 (*cryst struct*)Yamasaki, K. *et al.*, *Jpn. J. Pharmacol.*, 1990, **52**, 255 (*pharmacol*)Corbett, R. *et al.*, *Psychopharmacology*, 1991, **104**, 312 (*pharmacol*)Chang, M.-Y. *et al.*, *Heterocycles*, 2002, **57**, 2321-2334 (*synth*)Hanessian, S. *et al.*, *J. Med. Chem.*, 2006, **49**, 4544-4567 (*N-Boc Me ester, N-Boc Et ester*)**1,2,5,6-Tetrahydro-3-pyridinecarboxylic acid, 9CI** T-223**Guvacine**. Δ^3 -Tetrahydronicotinic acid
[498-96-4]C₆H₉NO₂ 127.143Alkaloid from *Areca catechu* (betel nut) (Arecaceae). Potent inhibitor of 4-Aminobutanoic acid (GABA) uptake. Mp 293-295° dec. (271-272°).**Hydrochloride**: Mp 316° dec.**Me ester: Guvacoline. Norarecoline**


[495-19-2]

C₇H₁₁NO₂ 141.169Alkaloid from *Areca catechu* (Arecaceae). Mp 27°. Bp_{13.5} 114° (lit. gives a pressure range).**Me ester; hydrochloride**: Mp 121-122°.**N-Me**: see 1,2,5,6-Tetrahydro-1-methyl-3-pyridinecarboxylic acid, T-205Jahns, E. *et al.*, *Ber.*, 1891, **24**, 2615 (*isol*)Wohl, A. *et al.*, *Ber.*, 1907, **40**, 4698 (*struct*)Freudenberg, K. *et al.*, *Ber.*, 1918, **51**, 976; 1668 (*struct*)

- v. Euler, H. *et al.*, *Helv. Chim. Acta*, 1944, **27**, 382 (*isol*)
 McElvain, S.M. *et al.*, *J.A.C.S.*, 1946, **68**, 1049 (*synth*)
 Spittler-Friedmann, M. *et al.*, *Monatsh. Chem.*, 1965, **96**, 104 (*ms*)
 Morlacchi, F. *et al.*, *Ann. Chim. (Rome)*, 1967, **57**, 1456 (*synth*)
 Krosggaard-Larsen, P. *et al.*, *Acta Chem. Scand., Ser. B*, 1978, **32**, 327 (*synth, pharmacol*)

2,3,4,9-Tetrahydro-1H-pyrido[3,4-*b*]indole-3-carboxylic acid, 9CI T-224

1,2,3,4-Tetrahydro-β-carboline-3-carboxylic acid
 [6052-68-2]
 [72002-54-1 (*R*-form), 41509-88-0 (*±*-form)]



$C_{12}H_{12}N_2O_2$ 216.239

(S)-form

L-form. **Lycoperodine 1**
 [42438-90-4]

Formed by Pictet-Spengler condensation of Formaldehyde and Tryptophan, T-640 in nature or during food processing. Alkaloid from *Aleurites fordii* and *Allium tuberosum*. Isol. from starfish *Asterias rollestoni*. Constit. of cocoa and tomatoes. Cryst. (H_2O). Mp 282-284° Mp 315°. $[\alpha]_D^{20}$ -124.6 (c, 0.2 in 0.1M NaOH). $[\alpha]_D^{25}$ -49.6 (c, 1 in 1M HCl MeOH/ H_2O 1:1). λ_{max} 274 (3.72); 291 (sh) (3.5) (MeOH).

Me ester: [79815-18-2]

$C_{13}H_{14}N_2O_2$ 230.266

Solid. Mp 167°. $[\alpha]_D^{20}$ -73.1 (c, 1 in $CHCl_3$).

Me ester, hydrochloride: Mp 283° dec. $[\alpha]_D^{20}$ -74.7 (c, 1 in EtOH).

Et ester: [129848-93-7]

$C_{14}H_{16}N_2O_2$ 244.293

Fluffy needles (MeCN) (as hydrochloride). Mp 269-270° (hydrochloride). Unstable in air.

*N*²-*Me*: [42438-91-5]

Cryst. (MeOH/ Me_2CO). Mp 260-262°. $[\alpha]_D^{25}$ -2.99 (c, 1.0 in 1M HCl/MeOH).

*N*²-*Me, Me ester: Methyl 2-methyl-2,3,4,9-tetrahydro-β-carboline-3-carboxylate*
 [83159-20-0]

$C_{14}H_{16}N_2O_2$ 244.293

Alkaloid from the leaves of *Gastrolobium callistachys*. Needles (MeOH or C_6H_6) or prisms. Mp 196-197° (changes cryst. form at 175-180°) (153-154°). $[\alpha]_D$ +18 (c, 0.3 in EtOH). $[\alpha]_D$ -62.6 (c, 1.6 in MeOH).

*N*²-*Me, Me ester, hydrochloride*:

Needles. Mp 284-285°. $[\alpha]_D^{20}$ -62.5 (c, 0.4 in EtOH).

7-Hydroxy: 1,2,3,4-Tetrahydro-7-hydroxy-β-carboline-3-carboxylic acid.

Brunnein C

[955377-47-6]

$C_{12}H_{12}N_2O_3$ 232.238

Alkaloid from *Cortinarius brunneus*.

Yellowish solid. $[\alpha]_D^{22}$ -57 (c, 0.02 in MeOH). λ_{max} 222 (log ε 4.4); 267 (log ε 3.5); 296 (log ε 3.5) (MeOH).

[129848-94-8]

Harvey, D.G. *et al.*, *J.C.S.*, 1941, **63**, 153-159 (*synth*)

Brossi, A. *et al.*, *J. Med. Chem.*, 1973, **16**, 418-420 (*S*-form, *synth, pmr, ord, cd, N*²-*Me*)

Okuda, T. *et al.*, *Phytochemistry*, 1975, **14**,

2304-2305 (*ir, uv, ms, pmr, config*)

Cannon, J.R. *et al.*, *Aust. J. Chem.*, 1982, **35**,

1497-1500 (*N*²-*Me Me ester*)

Maclaren, J.A. *et al.*, *Aust. J. Chem.*, 1989, **42**,

813-821 (*synth, bibl*)

Tilstra, L. *et al.*, *J.A.C.S.*, 1990, **112**, 9176-

9182; 9182-9190 (*synth, pmr, ms, cryst struct*)

Herraiz, T. *et al.*, *J. Agric. Food Chem.*, 1996,

44, 3057-3065; 1998, **46**, 3484-3490; 2000,

48, 4900-4904; 2004, **52**, 2652-5658 (*hplc, occur, bibl*)

Herraiz, T. *et al.*, *J. Chromatogr., A*, 1997, **765**,

265-277; 2000, **871**, 23-30 (*occur, gc, hplc*)

Li, G.Q. *et al.*, *J. Chin. Pharm. Sci.*, 2004, **13**,

81-86 (*isol, Asterias*)

Yahara, S. *et al.*, *J. Nat. Prod.*, 2004, **67**, 500-


502 (*Lycoperodine 1, isol, pmr, cmr*)

Teichert, A. *et al.*, *J. Nat. Prod.*, 2007, **70**,

1529-1531 (*Brunnein C*)

2,3,4,9-Tetrahydro-1H-pyrido[3,4-*b*]indole-1,6-diol T-225

1,2,3,4-Tetrahydro-1,6-dihydroxy-β-carboline
 [452067-34-4]



$C_{11}H_{12}N_2O_2$ 204.228

Isol. from the sponge *Hyrtios reticulata*.

1-O-(4-Guanidinobutyl): 1-(4-Guanidinobutyl)-6-hydroxy-β-carboline.

PwTx II

$C_{16}H_{23}N_5O_2$ 317.39

Isol. from the venom of the spider

Parawixia bistriata.

Salmoun, M. *et al.*, *J. Nat. Prod.*, 2002, **65**,


1173-1176 (*isol*)

Cesar, L.M.M. *et al.*, *Toxicol.*, 2005, **46**, 786-

796 (*PwTx II*)

2,3,4,9-Tetrahydro-1H-pyrido[3,4-*b*]indol-1-one, 9CI T-226

1,2,3,4-Tetrahydro-1-oxo-β-carboline
 [17952-82-8]



$C_{11}H_{10}N_2O$ 186.213

Alkaloid from *Evodia rutaecarpa*. Also isol. from an Indonesian sponge. Needles (EtOH aq.). Mp 183-185°.

Picrate:

Orange-yellow plates (EtOH). Mp 195-197°.

*N*²-*(2-Methylaminobenzoyl): Rhetsinine.*

Hydroxyevodiamine

[526-43-2]

$C_{19}H_{17}N_3O_2$ 319.362

Alkaloid from the trunk bark of *Zanthoxylum rhetsa* and the fruits of *Evodia rutaecarpa* (Rutaceae). Also obt. by the mild oxidn. of Evodiamine. Yellow cryst. ($CHCl_3$ /EtOH). Mp 206-207° (192° dec.). λ_{max} 314 (log ε 4.15) (EtOH).

*N*²-*(2-Methylaminobenzoyl), hydrochloride*:

Yellow needles. Mp 228-229° dec.

*N*²-*(2-Methylaminobenzoyl), picrate*:

Slender orange-yellow needles. Mp 270-272° dec.

*N*²-*Me: 2,3,4,9-Tetrahydro-2-methyl-1H-*

*pyrido[3,4-*b*]indol-1-one. 1,2,3,4-Tetra-*

hydro-2-methyl-1-oxo-β-carboline.

Strychnocarpine

[59156-98-8]

$C_{12}H_{12}N_2O$ 200.24

Alkaloid from the stem bark of

Strychnos elaeocarpa and *Strychnos*

floribunda (Loganiaceae). Weak

muscular relaxant. Cryst. (EtOAc).

Mp 226° (198-200°). λ_{max} 227 (log ε 4.25); 242 (log ε 4.07); 305 (log ε 4.14)

(MeOH).

6-Hydroxy: 2,3,4,9-Tetrahydro-6-hydro-

*xy-1H-pyrido[3,4-*b*]indol-1-one.*

1,2,3,4-Tetrahydro-6-hydroxy-1-oxo-β-

carboline

[51085-95-1]

$C_{11}H_{10}N_2O_2$ 202.212

Isol. from the sponge *Hyrtios* sp.

7-Hydroxy: 2,3,4,9-Tetrahydro-7-hydro-

*xy-1H-pyrido[3,4-*b*]indol-1-one, 9CI*

[56409-32-6]

Mp 138°.

5-Methoxy: 2,3,4,9-Tetrahydro-6-meth-

*oxy-1H-pyrido[3,4-*b*]indol-1-one.*

1,2,3,4-Tetrahydro-5-methoxy-1-oxo-β-

carboline

[26579-73-7]

$C_{12}H_{12}N_2O_2$ 216.239

Alkaloid from the root bark of *Alstonia*

venenata (Apocynaceae). Prisms

(MeOH). Mp 216-217° (*synth.*) Mp

182-184° (*nat.*). The original nat. iso-

late was impure and the reported

spectroscopic data was subsequently

revised. λ_{max} 236 (log ε 4.46); 295 (log ε

4.16) (EtOH).

6-Methoxy: 2,3,4,9-Tetrahydro-6-meth-

*oxy-1H-pyrido[3,4-*b*]indol-1-one, 9CI.*

1,2,3,4-Tetrahydro-6-methoxy-1-oxo-β-

carboline

[17952-87-3]

Synthetic. Prisms (EtOH). Mp 275-

277° (265°) Mp 275-277°.

7-Methoxy: 2,3,4,9-Tetrahydro-7-meth-

*oxy-1H-pyrido[3,4-*b*]indol-1-one.*

1,2,3,4-Tetrahydro-7-methoxy-1-oxo-β-

carboline. Ketotetrahydronorharmine.

Harmalacidine

[26579-69-1]

$C_{12}H_{12}N_2O_2$ 216.239

Alkaloid from *Banisteriopsis caapi* (Malpighiaceae) and seeds of *Peganum harmala* (Zygophyllaceae). Needles (CHCl₃/MeOH). Mp 200-201° (197-198° dec.). λ_{max} 220 ; 250 ; 325 (MeOH).

8-Methoxy: 2,3,4,9-Tetrahydro-8-methoxy-1H-pyrido[3,4-b]indol-1-one, 9CI. 1,2,3,4-Tetrahydro-8-methoxy-1-oxo-β-carboline

[109021-64-9]


Synthetic. Needles (MeOH). Mp 244-246° (241-242°).

- Abramovitch, R.A. *et al.*, *J.C.S.*, 1956, 4589-4592 (6-methoxy, *synth*)
 Chatterjee, A. *et al.*, *Tetrahedron*, 1959, 7, 257-261 (*Rhetsinine*, *isol*, *uv*, *ir*)
 Pachter, I.J. *et al.*, *J.O.C.*, 1960, 25, 1680-1682 (*Rhetsinine*, *uv*, *struct*, *synth*)
 Hashimoto, Y. *et al.*, *Phytochemistry*, 1976, 15, 1559-1560 (*Ketotetrahydronorharmine*, *isol*, *uv*, *ir*, *pmr*, *ms*, *struct*, *synth*)
 Rolfsen, W. *et al.*, *Acta Pharm. Suec.*, 1980, 17, 333-340 (*Strychnocarpine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)
 Verpoorte, R. *et al.*, *Planta Med.*, 1981, 42, 32-36 (*Strychnocarpine*, *isol*, *uv*, *pmr*, *ms*)
 Banerji, J. *et al.*, *Phytochemistry*, 1982, 21, 2765-2767 (5-methoxy, *isol*, *struct*)
 Herdeis, C. *et al.*, *Heterocycles*, 1984, 22, 2277 (*Strychnocarpine*, *synth*, *uv*, *ir*, *pmr*, *cmr*)
 Yamada, F. *et al.*, *Heterocycles*, 1986, 24, 2619-2627 (5-methoxy, 8-methoxy, *synth*, *uv*, *ir*, *pmr*, *struct*)
 Lehmann, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, 321, 411-414 (*Strychnocarpine*, *synth*, *ir*, *pmr*)
 Siddiqui, S. *et al.*, *Heterocycles*, 1988, 27, 1401-1410 (*Harmalacidine*, *isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)
 Narayanan, K. *et al.*, *J.O.C.*, 1991, 56, 359-365 (*pmr*, *ir*, *ms*)
 Bracher, F. *et al.*, *Annalen*, 1992, 1315-1319 (*synth*)
 Abramovitch, R.A. *et al.*, *Synlett*, 1992, 795-796 (6-methoxy, *synth*)
 Bracher, F. *et al.*, *Pharmazie*, 1993, 48, 695 (*Harmalacidine*, *Strychnocarpine*, *synth*)
 Soti, F. *et al.*, *Synth. Commun.*, 1993, 23, 1689-1698 (8-methoxy, *synth*)
 Salmoun, M. *et al.*, *J. Nat. Prod.*, 2002, 65, 1173-1176 (6-hydroxy)
 Rao, K.V. *et al.*, *J. Nat. Prod.*, 2003, 66, 823-828 (*isol*, *pmr*, *cmr*)
 Lehmann, I. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1988, 321-411 (*synth*, *ir*, *pmr*)

6,8,9,11-Tetrahydro-7H-pyrido[2,1-b]quinazoline, 9CI

6,7,8,9-Tetrahydro-11H-pyrido[2,1-b]quinazoline

[261-08-5]



C₁₂H₁₄N₂ 186.256

Alkaloid from the leaves of *Macklinaya macroscladia* and *Macklinaya subulata* (Araliaceae). Plates (Et₂O or Et₂O/petrol). Mp 85-87°.

Hydroiodide:

Needles (EtOH). Mp 225-226°.

Sulfate:


Needles (EtOH). Mp 283-285°.

Picrate:

Cryst. (AcOH). Mp 216-217°.

- Späth, E. *et al.*, *Ber.*, 1935, 68, 2221 (*synth*)
 Munoz, G.G. *et al.*, *Chem. Ber.*, 1962, 95, 2182 (*synth*)
 Fitzgerald, J.S. *et al.*, *Aust. J. Chem.*, 1966, 19, 151 (*isol*, *pmr*, *ms*)

1,2,3,4-Tetrahydro-5-(2-pyrrolidinyl)pyridine



C₉H₁₆N₂ 152.239

N¹-Ac: *Maackiamine*

[54966-14-2]

C₁₁H₁₈N₂O 194.276


Alkaloid from the flowers of *Maackia amurensis* var. *buergeri* (Fabaceae). Oil. [α]_D²⁵ +110 (c, 0.01 in EtOH).

Saito, K. *et al.*, *Phytochemistry*, 1989, 28, 2533 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

5,6,7,7a-Tetrahydro-3H-pyrolizin-3-one, 9CI

1-Azabicyclo[3.3.0]oct-3-en-2-one. *Pyrrolam A*

[113727-94-9]



C₇H₉NO 123.154

(R)-form [126424-76-8]

Prod. by *Streptomyces olivaceus*. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 62°. [α]_D²⁰ -29.3 (c, 1 in CHCl₃). λ_{max} 209 (ε 6000); 240 (ε 2600) (MeOH) (Berdy).

(S)-form [161691-23-2]


Solid. [α]_D²⁷ +22.37 (c, 0.96 in CHCl₃).

- Flitsch, W. *et al.*, *Annalen*, 1988, 387 (*synth*, *pmr*)
 Grote, R. *et al.*, *Annalen*, 1990, 525 (*isol*, *pmr*, *cmr*, *struct*)
 Aoyagi, Y. *et al.*, *Tetrahedron*, 1996, 52, 869 (*synth*)
 Giovenzana, G.B. *et al.*, *Tetrahedron: Asymmetry*, 1997, 8, 515-518 (*synth*)
 Watson, R.T. *et al.*, *J.O.C.*, 2004, 69, 6105-6114 (*synth*, *bibl*)
 Majik, M.S. *et al.*, *Synthesis*, 2007, 663-665 (*S-form*, *synth*, *pmr*, *cmr*)
 Schobert, R. *et al.*, *Synthesis*, 2007, 1499-1502 (*R-form*, *synth*, *ir*, *pmr*, *cmr*, *ms*)

4,5,6,7-Tetrahydro-2H-pyrrolo[3,4-c]pyridine-6-carboxylic acid, 9CI

5H-Pyrrolo[3,4-d]piperidine-2-carboxylic acid

[134649-28-8]



C₈H₁₀N₂O₂ 166.179

(ξ)-form


Prod. by *Penicillium oxalicum*. Sol. H₂O; poorly sol. MeOH, EtOAc, DMSO. Mp 284°. λ_{max} 215 (H₂O).

Japan. Pat., 1990, 90 306 976; *CA*, 115, 27688k (*isol*)

1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazoline, 9CI

Deoxypeganine. *Deoxyvasicine*

[495-59-0]



C₁₁H₁₂N₂ 172.229

Alkaloid from the above-ground parts of *Peganum harmala* and the foliage of *Peganum nigellastrum* (Zygophyllaceae). Claimed to show cholinergic activity. Cryst. (Et₂O). Mp 99-100°.

Hydrochloride: Mp 260° (246°).

Picrate:

Cryst. (EtOH). Mp 205-206°.

N-Oxide: *Deoxypeganine N-oxide*

[168780-02-7]

C₁₁H₁₂N₂O 188.229

Alkaloid from *Nitraria komarovii* (Zygophyllaceae). Cryst. (EtOH/Me₂CO). Mp 243-244°. [α]_D 0. λ_{max} 214 (log ε 4.04); 217 (log ε 4.1); 223 (log ε 3.98); 285 (log ε 3.51) (no solvent reported).

1-Hydroxy: 1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazolin-1-ol, 9CI. 1-Hydroxydeoxypeganine

[119459-57-3]

C₁₁H₁₂N₂O 188.229

Alkaloid from aerial parts of *Galium aparine* collected at the flowering period (Rubiaceae).

3-Hydroxy: see *Peganine*, P-165

9-Oxo: see 2,3-Dihydropyrrolo[2,1-b]quinazolin-9(1H)-one, D-503

Hanford, W.E. *et al.*, *J.A.C.S.*, 1935, 57, 920 (*synth*)

Späth, E. *et al.*, *Ber.*, 1936, 69, 255 (*synth*)

Muñoz, G.G. *et al.*, *Chem. Ber.*, 1962, 95, 2182 (*synth*)

Timothy, J. *et al.*, *J. Med. Chem.*, 1973, 16, 633 (*synth*, *pmr*)

Batsuren, D. *et al.*, *Khim. Prir. Soedin.*, 1980, 16, 736 (*isol*)


Johns, S. *et al.*, *Alkaloids (Academic Press)*, 1986, 29, 129 (*rev*, *pharmacol*)

Sener, B. *et al.*, *Gazi Univ. Eczacilik Fak. Derg.*, 1988, 5, 33; *CA*, 110, 111673f (*1-Hydroxydeoxypeganine*)

Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin.*, 1993, 29, 87; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, 29, 73 (*Deoxypeganine N-oxide*)

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1995, 31, 421-425; *Chem. Nat. Compd. (Engl. Transl.)*, 1995, 31, 349-352 (*cryst struct*)

Tozhibaev, A.G. *et al.*, *Khim. Prir. Soedin.*, 2006, 42, 280-283; *Chem. Nat. Compd. (Engl. Transl.)*, 2006, 42, 340-344 (*cryst struct*)

1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-1-methylisoquinoline, 9CI**Pachycereine**
[82261-04-9]C₁₄H₂₁NO₄ 267.324Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).N-Me: 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-1,2-dimethylisoquinoline, 9CI. **N-Methylpachycereine**

[74046-25-6]

C₁₅H₂₃NO₄ 281.351Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).1,2-Didehydro: 3,4-Dihydro-5,6,7,8-tetramethoxy-1-methylisoquinoline, 9CI. **Dehydopachycereine**


[82261-02-7]

C₁₄H₁₉NO₄ 265.308Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).1,2,3,4-Tetrahydro: 5,6,7,8-Tetramethoxy-1-methylisoquinoline, 9CI. **Isopachycereine**

[93474-27-2]

C₁₄H₁₇NO₄ 263.293Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **77**, 109 (occur, derivs)**Tetrahydrothalifendine**


T-234



Absolute Configuration

C₁₉H₁₉NO₄ 325.363Shamma, M. *et al.*, *Tetrahedron*, 1971, **27**, 727 (uv, pmr, ir, ms, struct)Pinho, P.M.M. *et al.*, *Phytochemistry*, 1992, **31**, 1403 (8-Oxotetrahydrothalifendine)**1,2,3,4-Tetrahydro-1-(2-thio-3-pyrrolidinyl)-β-carboline-3-carboxylic acid**

T-235



(1R*,3R*,3'S*)-form

C₁₆H₁₇N₃O₂S 315.395

(1R,3R,3'S)-form [221657-55-2]

Alkaloid from fermented radish roots.


(1R,3S,3'R)-form [221657-57-4]

Alkaloid from fermented radish roots.

Ozawa, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 216-219**1,2,3,4-Tetrahydro-5,6,7-trihydroxyisoquinoline**

T-236

1,2,3,4-Tetrahydro-5,6,7-isoquinolinetriol

C₉H₁₁NO₃ 181.1916,7-Di-Me ether: 1,2,3,4-Tetrahydro-6,7-dimethoxy-8-isoquinolinol. 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxyisoquinoline. **Anhalamine**

[643-60-7]

C₁₁H₁₅NO₃ 209.244Alkaloid from *Anhalonium lewinii* (*Lophophora williamsii*), *Lophophora diffusa* and *Gymnocalycium gibbosum* (Cactaceae). Cryst. (EtOH). Mp 189-191°. pK_{a1} 8.8; pK_{a2} 11.3 (50% 2-propanol aq.).

▶ NX5948000

6,7-Di-Me ether, hydrochloride:

Cryst. (AcOH). Mp 277-278° dec Mp 257-258° (in vacuo).

6,7-Di-Me ether, picrate: Mp 273-274° Mp 249-249.5° (in vacuo).

6,7-Di-Me ether, N-formyl: **N-Formylanhalamine**C₁₂H₁₅NO₄ 237.255Constit. of peyote cactus, *Lophophora williamsii* (Cactaceae). Nat. prod. ident. by glc/ms in comparison with an authentic sample.6,7-Di-Me ether, N-Ac: **N-Acetylanhalamine**C₁₃H₁₇NO₄ 251.282Constit. of *Lophophora williamsii* (Cactaceae). Nat. prod. ident. by glc/ms.

6,7-Di-Me ether, N-benzoyl:

C₁₈H₁₉NO₄ 313.352

Prisms (EtOH). Mp 167.5°.

6,7-Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-8-isoquinolinol. 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-2-methylisoquinoline. **Anhalidine**. N-Methylanhalamine

[2245-94-5]

C₁₂H₁₇NO₃ 223.271Alkaloid from *Anhalonium lewinii* (*Lophophora williamsii*), *Pelecypora aselliformis* and *Stetsonia coryne* (Cactaceae). Cryst. by subl. Mp 131-133°. pK_{a1} 7.7; pK_{a2} 11.1 (50% 2-propanol aq.).

▶ NX5955350

6,7-Di-Me ether, N-Me, hydrochloride: Mp 243°.

6,7-Di-Me ether, N,N-di-Me: 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-2,2-dimethylisoquinolinium (1+). **Anhalotine**

[19267-93-7]

[19445-62-6]

C₁₃H₂₀NO₃⁺ 238.306Alkaloid from *Anhalonium lewinii* (Cactaceae). Cryst. (EtOH/EtOAc) (as iodide). Mp 219-220° (iodide).

7,8-Di-Me ether: 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxyisoquinoline.

1,2,3,4-Tetrahydro-7,8-dimethoxy-6-isoquinolinol. **Isoanhalamine**

[5308-58-7]

C₁₁H₁₅NO₃ 209.244Alkaloid from *Lophophora williamsii* (peyote cactus) (Cactaceae). Cryst. (CH₂Cl₂/Et₂O). Mp 172-174°.

7,8-Di-Me ether, hydrobromide: Mp 214.5-215.5°.

7,8-Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-7,8-dimethoxy-2-methyl-6-isoquinolinol. 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-2-methylisoquinoline. **Isoanhalidine**

[37484-64-3]

C₁₂H₁₇NO₃ 223.271Alkaloid from *Lophophora williamsii* (peyote cactus) (Cactaceae). Mp 215-218° (as hydrochloride).Tri-Me ether: 1,2,3,4-Tetrahydro-6,7,8-trimethoxyisoquinoline. **Anhalinine**. O-Methylanhalamine

[642-30-8]

C₁₂H₁₇NO₃ 223.271Alkaloid from *Anhalonium lewinii* (Cactaceae). Cryst. by subl. Mp 61-63°.

Tri-Me ether, hydrochloride: Mp 242-243°.

Tri-Me ether, picrate: Mp 184-185°.

Tri-Me ether, N-formyl: **N-Formylanhalidine**C₁₃H₁₇NO₄ 251.282Alkaloid from *Anhalonium lewinii* (Cactaceae). Identified by glc/ms comparison with an authentic specimen.

6,7-Methylene, 8-Me ether, N-Me: see Hydrocotarnine, H-396

Heffter, A. *et al.*, *Ber.*, 1901, **34**, 3004

(Anhalamine, isol)


Späth, E. *et al.*, *Ber.*, 1934, **67**, 2100; 1935, **68**, 501; 944 (Anhalamine, Anhalinine, isol, synth)Brossi, A. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 2089; 1966, **49**, 403 (synth, derivs)Kametani, T. *et al.*, *Yakugaku Zasshi*, 1966, **86**, 913; *CA*, **66**, 28631t (Anhalamine, Anhalidine, synth)Kapadia, G.J. *et al.*, *Chem. Comm.*, 1968, 1688 (N-Formylanhalinine, N-Formylanhalamine, N-Acetylanhalamine)Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 254 (Anhalotine)Lundström, J. *et al.*, *Tet. Lett.*, 1968, 4437 (biosynth)Todd, J.S. *et al.*, *J. Nat. Prod.*, 1969, **32**, 395 (Anhalamine, occur)

- Khanna, K.L. *et al.*, *Phytochemistry*, 1970, **9**, 1811 (*biosynth*)
 Lundström, J. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 1295 (*Isoanhalamine, Isoanhalidine*)
 Neal, J.M. *et al.*, *Science (Washington, D.C.)*, 1972, **176**, 1131 (*Anhalidine, isol*)

1,2,3,4-Tetrahydro-6,7,8-trihydroxy-1-methylisoquinoline

T-237

1,2,3,4-Tetrahydro-1-methyl-6,7,8-isoquinolinetriol



(R)-form

C₁₀H₁₃NO₃ 195.218

(R)-form

- 6,7-Di-Me ether: 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-methylisoquinoline. 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methyl-8-isoquinolinol. **Anhalonidine** [529-58-8]
 C₁₂H₁₇NO₃ 223.271
 Alkaloid from *Lophophora williamsii* (Cactaceae). [α]_D -21.2 (95% EtOH). Racemises readily.
 ▶ LD₅₀ (mus, orl) 700 mg/kg. NX5953000
 6,7-Di-Me ether, hydrochloride: [α]_D -0.7 (95% EtOH).

(S)-form

- 7,8-Di-Me ether: 1,2,3,4-Tetrahydro-7,8-dimethoxy-1-methyl-6-isoquinolinol. 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-1-methylisoquinoline. **Isoanhalonidine** [37484-65-4]
 C₁₂H₁₇NO₃ 223.271
 Alkaloid from *Lophophora williamsii* (peyote cactus) (Cactaceae). Cryst. (Et₂O/pentane). Mp 112-114°.
 7,8-Di-Me ether, hydrobromide: Mp 210.5-212°.
 7,8-Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-1,2-dimethylisoquinoline. 1,2,3,4-Tetrahydro-7,8-dimethoxy-1,2-dimethyl-6-isoquinolinol. **Isopellotine** [37484-66-5]
 C₁₃H₁₉NO₃ 237.298
 Alkaloid from *Lophophora williamsii* (Cactaceae). Mp 212-222° (as hydrochloride).
 Tri-Me ether: 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-methylisoquinoline. **O-Methylanhalonidine** [35646-08-3]
 C₁₃H₁₉NO₃ 237.298
 Alkaloid from *Lophophora williamsii* (peyote cactus) (Cactaceae). Plant growth inhibitor. Viscous oil. Bp_{0.05} 140°. [α]_D²⁵ +20.6 (c, 1 in MeOH) ((+11.5)). [α]_D²⁵ +19.3 (c, 1 in 1M HCl).

Tri-Me ether, hydrobromide:

Cryst. (H₂O). Mp 202-204°. [α]_D²⁵ +16.4 (c, 1 in MeOH).

(±)-form

- 6,7-Di-Me ether: [3851-33-0]
 Alkaloid from *Lophophora williamsii*, *Echinocactus lewinii* and *Lemaireocereus weberi* (Cactaceae). Narcotic and curarising agent to frogs but not to mammals. Cryst. (Me₂CO). Mp 161-161.5°.
 6,7-Di-Me ether, hydrochloride: Cryst. (EtOH/Et₂O). Mp 248.5-250° dec.
 6,7-Di-Me ether, picrate: Needles (EtOH). Mp 200.5-201.5° (201-208°).
 6,7-Di-Me ether, N,O-dibenzoyl: Mp 125-126°.
 6,7-Di-Me ether, N-Me: 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinoline. 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethyl-8-isoquinolinol. **Pellotine. Peyotiline** [83-14-7]
 C₁₃H₁₉NO₃ 237.298
 Alkaloid from *Anhalonium lewinii* (*Lophophora williamsii*), *Anhalonium williamsii*, *Pachycereus weberi* and *Pelecophora aselliformis*, also detected in *Lophophora diffusa* and *Islaya minor* (preferred genus name *Neoporteria*) (Cactaceae). Cryst. (EtOH or petrol). Mp 111-112°. [α]_D 0. pK_{a1} 8.3; pK_{a2} 11.6 (50% 2-propanol aq.). May be optically active in the plant, but racemises during extraction.
 ▶ RY0350000
 6,7-Di-Me ether, N-Me, hydrochloride: Cryst. (EtOH). Mp 243-244°.
 6,7-Di-Me ether, N-Me, picrate: Mp 172-173°.
 Tri-Me ether: [4838-93-1]
 Synthetic. Cryst. (EtOAc)(as hydrochloride). Mp 148-150° (hydrochloride).
 (±)-form
 6,7-Di-Me ether, N-formyl: **N-Formylanhalonidine**
 C₁₃H₁₇NO₄ 251.282
 Constit. of *Lemaireocereus williamsii* (Cactaceae). Abs. config. of the nat. alkaloid unknown; it was identified by glc/ms comparison with an authentic sample.
 6,7-Di-Me ether, N,N-di-Me: 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1,2,2-trimethylisoquinolinium(1+). **Peyotine** [25526-36-7]
 C₁₄H₂₂NO₃⁺ 252.333
 Alkaloid from *Lophophora williamsii* (Cactaceae). Cryst. (H₂O)(as iodide). Mp 185-186° (softens and resolidifies at 114°)(iodide).
 Tri-Me ether, N-formyl: **N-Formyl-O-methylanhalonidine** [84666-21-7]
 C₁₄H₁₉NO₄ 265.308

Constit. of *Lophophora williamsii* (Cactaceae). Abs. config. not detd., ident. by glc/ms.

Tri-Me ether, N-Me: 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1,2-dimethylisoquinoline. **O-Methylpellotine** [4973-61-9]
 C₁₄H₂₁NO₃ 251.325
 Constit. of *Lophophora diffusa* (Cactaceae). Nat. prod. identified by tlc and glc comparison with reference material.


- Heffter, A. *et al.*, *Ber.*, 1894, **27**, 2975-2979 (*Pellotine, isol*)
 Späth, E. *et al.*, *Monatsh. Chem.*, 1921, **42**, 97-115; 1922, **43**, 477-484 (*Anhalonidine, Pellotine, synth*)
 Späth, E. *et al.*, *Ber.*, 1932, **65**, 1778-1785; 1934, **67**, 266-268; 1936, **69**, 755-757 (*Pellotine, Anhalonidine, struct, synth*)
 Späth, E. *et al.*, *Ber.*, 1939, **72**, 334-338 (*O-Methylanhalonidine*)
 Djerassi, C. *et al.*, *J.A.C.S.*, 1954, **76**, 3215-3217 (*Anhalonidine, isol*)
 Brossi, A. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 2089-2098 (*Pellotine, Anhalonidine, synth, uv, ir, pmr*)
 Brossi, A. *et al.*, *Helv. Chim. Acta*, 1966, **49**, 403 (*Isoanhalonidine*)
 Leete, E. *et al.*, *J.A.C.S.*, 1966, **88**, 4218-4221 (*biosynth*)
 Battersby, A.R. *et al.*, *Tet. Lett.*, 1967, 563-565 (*biosynth*)
 Kapadia, G.J. *et al.*, *Chem. Comm.*, 1968, 1688-1689 (*N-Formylanhalonidine*)
 Kapadia, G.J. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 254-262 (*Peyotiline, isol, uv, ir*)
 Todd, J.S. *et al.*, *J. Nat. Prod.*, 1969, **32**, 395-398 (*tlc, occur*)
 Fales, H.M. *et al.*, *J.A.C.S.*, 1969, **91**, 3682-3685 (*O-Methylpellotine, ms*)
 Khanna, K.L. *et al.*, *Phytochemistry*, 1970, **9**, 1811-1815 (*biosynth*)
 Brossi, A. *et al.*, *J.A.C.S.*, 1971, **93**, 6248-6252 (*O-Methylanhalonidine, uv, cd, ord, pmr, config, cryst struct, resoln*)
 Lundström, J. *et al.*, *Acta Chem. Scand.*, 1972, **26**, 1295 (*Isopellotine*)
 Neal, J.M. *et al.*, *Science (Washington, D.C.)*, 1972, **176**, 1131-1133 (*Pellotine*)
 Kapadia, G. *et al.*, *J. Nat. Prod.*, 1973, **36**, 9-35 (*rev, ms*)
 Bruhn, J.G. *et al.*, *Phytochemistry*, 1975, **14**, 1442-1443 (*O-Methylpellotine, isol*)
 Cymerman Craig, J. *et al.*, *J.A.C.S.*, 1977, **99**, 7996-8002 (*uv, cd, abs config*)
 Stronböm, J. *et al.*, *J. Chromatogr.*, 1978, **147**, 513-515; 1980, **189**, 79-89 (*hplc, occur*)
 Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673 (*isol*)
 Sax, N.I. *et al.*, *Dangerous Properties of Industrial Materials*, 5th edn., Van Nostrand Reinhold, 1979, 379

1,2,3,4-Tetrahydro-5,6,7-trimethoxy-2-methylisoquinoline, 9CI

T-238

Tehaunine

[30147-93-4]



C₁₃H₁₉NO₃ 237.298

Alkaloid from *Pachycereus weberi*, *Pachycereus pringlei* and *Pachycereus tehuantepecanus* (Cactaceae). Mp 229-230° (219-221°) (as hydrochloride).

N-Oxide: Tehaunine N-oxide

[85769-25-1]

C₁₃H₁₉NO₄ 253.297

Alkaloid from *Pachycereus pringlei* (Cactaceae). Cryst. (EtOH/Et₂O) (as hydrochloride). Mp 186-187° (hydrochloride).

N-De-Me: Nortehaunine. 1,2,3,4-Tetrahydro-5,6,7-trimethoxyisoquinoline

[1745-06-8]

C₁₂H₁₇NO₃ 223.271

Alkaloid from *Pachycereus weberi* (Cactaceae). Mp 71-72°.

N-De-Me; hydrochloride: Mp 268-269° (260°).

N-De-Me, picrate: Mp 175-177°.

Bobbitt, J.M. et al., *J.O.C.*, 1965, **20**, 2247 (synth)

Mata, R. et al., *Phytochemistry*, 1980, **19**, 673 (isol, uv, ir, pmr, ms, struct, deriv)

Mata, R. et al., *Planta Med.*, 1980, **38**, 180 (isol)

Hara, H. et al., *Heterocycles*, 1982, **17**, 293 (synth)


Pummangura, S. et al., *Phytochemistry*, 1982, **21**, 2375 (oxide)

Mata, R. et al., *Phytochemistry*, 1983, **22**, 1263 (cmr, Tehaunine, Nortehaunine)

Takano, S. et al., *Heterocycles*, 1993, **35**, 47 (synth)

1,8,14,16-Tetrahydroxyaconitan-19-one

T-239



C₁₈H₂₅NO₅ 335.399

(1α,5β,14α,16β)-form

O',O¹⁶-Di-Me: Piepunendine A

[947321-58-6]


C₂₀H₂₉NO₅ 363.453

Alkaloid from the roots of *Aconitum piepunense*. Mp 107-109°. [α]_D²⁰ -14.3 (c, 0.5 in CHCl₃).

Cai, L. et al., *Nat. Prod. Commun.*, 2006, **1**, 191-194 (isol, pmr, cmr)

1,2,3,4-Tetrahydroxyacridone

T-240



C₁₃H₉NO₅ 259.218

2,3,4-Tri-Me ether: 1-Hydroxy-2,3,4-trimethoxyacridone

[114340-03-3]

C₁₆H₁₅NO₅ 301.298

Alkaloid from leaves of *Vepris fitoravina* and *Vepris macrophylla* (Rutaceae). Orange-yellow needles (EtOAc/hexane). Mp 208-209° (200°).

2,3,4-Tri-Me ether, N-Me: 1-Hydroxy-2,3,4-trimethoxy-10-methylacridone.

Normelicopine

[7008-68-6]

C₁₇H₁₇NO₅ 315.325

Alkaloid from *Acronychia baueri* and *Teclea trichocarpa*. Orange needles (EtOH aq.). Mp 125-127°. Prob. artifact.

Tetra-Me ether: 1,2,3,4-Tetramethoxyacridone

[75340-58-8]

C₁₇H₁₇NO₅ 315.325

Alkaloid from the leaves of *Bauerella simplicifolia* (preferred genus name *Acronychia*) (Rutaceae). Mp 238-240°.

Tetra-Me ether, N-Me: 1,2,3,4-Tetramethoxy-10-methylacridone. Melicopine

[517-73-7]

C₁₈H₁₉NO₅ 329.352

Alkaloid from *Acronychia baueri*, *Teclea boiviniana*, *Melicope fareana* and *Melicope leptococca* (Rutaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 133-134°. λ_{max} 218 ; 268 ; 397 (EtOH) (Berdy).

2,3-Methylene, 1,4-di-Me ether: see Xanthevodine, X-2

3,4-Methylene, 1,2-di-Me ether, N-Me: see Melicopine, M-169

Price, J.R. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 249-254; *CA*, **46**, 4010d (Melicopine)

Crow, W.D. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 282-300; *CA*, **46**, 4014a (Melicopine, struct)

Hughes, G.K. et al., *Aust. J. Sci. Res., Ser. A*, 1950, **3**, 497-503; *CA*, **46**, 4544a (2,3,4-tri-Me ether, tetra-Me ether, synth)

Brown, R.D. et al., *Aust. J. Sci. Res., Ser. A*, 1950, **3**, 593-614; *CA*, **45**, 9369a (uv)

Bowie, J.H. et al., *Aust. J. Chem.*, 1967, **20**, 1179-1193 (ms)

Mester, I. et al., *Z. Naturforsch., B*, 1979, **34**, 516-519 (cmr)

Tillequin, F. et al., *J. Nat. Prod.*, 1980, **43**, 498 (tetra-Me ether)

Skaltsounis, A.L. et al., *J. Nat. Prod.*, 1983, **46**, 732-735 (Melicopine)

Baudouin, G. et al., *J. Nat. Prod.*, 1985, **48**, 260-265 (2,3,4,10-tetra-Me)


Koffi, Y. et al., *Planta Med.*, 1987, **53**, 570-571

Rasoanaivo, P. et al., *Fitoterapia*, 1999, **70**, 625-627 (Melicopine, isol, cmr)

Muriithi, M.W. et al., *J. Nat. Prod.*, 2002, **65**, 956-959 (Normelicopine, isol, cryst struct)

1,2,3,5-Tetrahydroxyacridone

T-241



C₁₃H₉NO₅ 259.218

2,3-Di-Me ether, N-Me: 1,5-Dihydroxy-2,3-dimethoxy-10-methylacridone. 5-Hydroxyarborinine

[75821-37-3]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the stems and leaves of *Glycosmis bilocularis* (Rutaceae). Orange-yellow cryst. (Et₂O/petrol). Mp 206-207° (185-187°). Phys. props. differ from those of an alkaloid of the same assigned struct. (Mp 185-7°) isol. from *Atalantia monopylla*.

2,3-Di-Me ether, N-Me, 5-Ac:

C₁₈H₁₇NO₆ 343.335

Yellow prisms (Me₂CO/hexane). Mp 128-129°.

2,3-Di-Me ether, N-Me, 1,5-di-Ac:

C₂₀H₁₉NO₇ 385.373

Pale-yellow prisms (Me₂CO/hexane). Mp 192-193°.

2,3,5-Tri-Me ether, N-Me: 1-Hydroxy-2,3,5-trimethoxy-10-methylacridone. 5-Methoxyarborinine

[75821-42-0]

C₁₇H₁₇NO₅ 315.325

Alkaloid from stem bark of *Luvunga angustifolia* (Rutaceae). Yellow needles (EtOH). Mp 168-169° (135-136°).

Tetra-Me ether, N-Me: 1,2,3,5-Tetramethoxy-10-methylacridone

C₁₈H₁₉NO₅ 329.352

Yellow needles (EtOH). Mp 97-98°.

2,3-Methylene, 1,5-di-Me ether: see Tecleanthine, T-56


Bowen, I.H. et al., *Phytochemistry*, 1978, **17**, 2125; 1980, **19**, 1566 (isol, uv, ir, pmr, ms, synth)

Shah, J.S. et al., *Indian J. Chem., Sect. B*, 1982, **21**, 16 (isol, uv, ir, pmr, ms, struct)

Wijeratne, E.M.K. et al., *J. Nat. Prod.*, 1992, **55**, 1261 (5-Methoxyarborinine)

1,3,4,5-Tetrahydroxyacridone

T-242



C₁₃H₉NO₅ 259.218

4-Me ether, N-Me: 1,3,5-Trihydroxy-4-methoxy-10-methylacridone. Citrusinine II

[86680-33-3]

C₁₅H₁₃NO₅ 287.271

Alkaloid from the root bark of *Citrus sinensis* var. *brasiliensis* (navel orange) (Rutaceae). Yellow needles (Me₂CO). Mp 244-246°.

3,4-Di-Me ether, N-Me: 1,5-Dihydroxy-3,4-dimethoxy-10-methylacridone. Citrusinine I

[86680-32-2]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the root bark of *Citrus sinensis* var. *brasiliensis* (navel orange) and *Citrus grandis* (pummelo) (Rutaceae). Orange needles (Me₂CO). Mp

206-207°. λ_{\max} 222 ; 265 ; 319 ; 331 ; 416 (EtOH) (Berdy).

Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 901 (*isol, uv, ir, pmr, cmr, ms, struct*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1983, **22**, 1493 (*isol*)

Kato, N. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 445 (*synth, Citrusinine I*)

1,3,5,6-Tetrahydroxyacridone T-243

C₁₃H₉NO₅ 259.218

5-Me ether, N-Me: 1,3,6-Trihydroxy-5-methoxy-10-methylacridone. **Grandisine III**

C₁₅H₁₃NO₅ 287.271

Alkaloid from roots of several hybrid seedlings resulting from a cross of Pummelo (*Citrus grandis* cv. May Pummelo x Marsh grapefruit (*Citrus paradisi*)) (Rutaceae). Yellow amorph. powder.

3,5-Di-Me ether: 1,6-Dihydroxy-3,5-dimethoxyacridone. **Natsucitrine I. Des-N-methylcitpressine I** [96910-77-9]

C₁₅H₁₃NO₅ 287.271

Alkaloid from the root bark of *Citrus natsudaidai* (Rutaceae). Light yellow prisms (MeOH). Mp 292-293°.

3,5-Di-Me ether, N-Me: 1,6-Dihydroxy-3,5-dimethoxy-10-methylacridone. **Citpressine I** [81525-58-8]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the root of bark of *Citrus depressa* (Shekwasha mandarin) and *Citrus grandis*, f. *hakunikuyu* (Rutaceae). Yellow needles + ½ H₂O (Me₂CO). Mp 183-185°.

3,6-Di-Me ether, N-Me: 1,5-Dihydroxy-3,6-dimethoxy-10-methylacridone.

Grandisine I

[87959-97-5]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the root bark of *Citrus grandis* f. *hakunikuyu* (Rutaceae). Pale yellow needles (Me₂CO). Mp 262-264°.

5,6-Di-Me ether, N-Me: 1,3-Dihydroxy-5,6-dimethoxy-10-methylacridone.

Grandisine II

[87959-99-7]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the root bark of *Citrus grandis* f. *hakunikuyu* (Rutaceae). Yellow needles (Me₂CO). Mp 266-268°.

3,5,6-Tri-Me ether: 1-Hydroxy-3,5,6-trimethoxyacridone. **Natsucitrine II. Des-N-methylcitpressine II** [96910-78-0]

C₁₆H₁₅NO₅ 301.298

Alkaloid from the root bark of *Citrus natsudaidai* (Rutaceae). Light yellow needles (C₆H₆/Me₂CO). Mp 246-247°.

3,5,6-Tri-Me ether, N-Me: 1-Hydroxy-3,5,6-trimethoxy-10-methylacridone.

Citpressine II

[81525-59-9]

C₁₇H₁₇NO₅ 315.325

Alkaloid from the root of bark of *Citrus depressa* (Shekwasha mandarin) and *Citrus grandis* f. *hakunikuyu* (Ru-

taeae). Yellow needles (Et₂O). Mp 168-170°.

5,6-Methylene, 3-Me ether, N-Me: 1-Hydroxy-3-methoxy-10-methyl-5,6-methylenedioxyacridone. **Marshdine** [160927-87-7]

C₁₆H₁₃NO₅ 299.282

Alkaloid from roots of marsh grapefruit *Citrus paradisi* (Rutaceae). Yellow cubes. Mp 210-213°.


Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 1983, **31**, 895 (*isol, uv, ir, pmr, ms, struct, Citpressines*)

Wu, T.-S. *et al.*, *Phytochemistry*, 1983, **22**, 1493 (*isol, uv, ir, cmr, pmr, ms, struct, Grandisines*)

Ju-ichi, M. *et al.*, *Heterocycles*, 1985, **23**, 1131 (*isol, uv, ir, pmr, cmr, ms, struct, synth, Natsucitrines*)

Takemura, Y. *et al.*, *Heterocycles*, 1992, **34**, 2123; 1994, **39**, 315 (*Grandisine II, Marshdine*)

1,2,3,9-Tetrahydroxyaporphine T-244



C₁₇H₁₇NO₄ 299.326

(R)-form

2,3-Di-Me ether: 1,9-Dihydroxy-2,3-dimethoxyaporphine

[α]_D -88 (c, 0.07 in CHCl₃).

2,3-Di-Me ether, N-de-Me: 1,9-Dihydroxy-2,3-dimethoxynoraporphine.

Oureguattidine

[88607-28-7]

C₁₈H₁₉NO₄ 313.352

Alkaloid from the stem bark of *Guatteria ouregou* (Annonaceae). Cryst. (MeOH). Mp 140°. [α]_D -13 (c, 0.16 in EtOH).

2,3-Di-Me ether, N-de-Me, N,O,O-tri-Ac: Cryst. (Me₂CO). Mp 168-171°. [α]_D -211 (c, 0.13 in EtOH).

1,2,3-Tri-Me ether, N-de-Me: 9-Hydroxy-1,2,3-trimethoxynoraporphine. **Oureguattine**

[88607-30-1]

C₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae).

(S)-form

1,2-Di-Me ether, N-de-Me: 3,9-Dihydroxy-1,2-dimethoxynoraporphine.

Norguattevaline

[158182-27-5]

C₁₈H₁₉NO₄ 313.352

Alkaloid from stem bark of *Guatteria foliosa* (Annonaceae). Amorph. [α]_D +35 (EtOH). Guattevaline appears to be unknown.

Leboeuf, M. *et al.*, *Planta Med.*, 1983, **48**, 234

(*Oreguattidine*)


Cortes, D. *et al.*, *J. Nat. Prod.*, 1986, **49**, 878

(*Oureguattine*)

Mahiou, V. *et al.*, *J. Nat. Prod.*, 1994, **57**, 890

(*Norguattevaline*)

1,2,3,11-Tetrahydroxyaporphine T-245



C₁₇H₁₇NO₄ 299.326

(R)-form

1,2,3-Tri-Me ether: 11-Hydroxy-1,2,3-trimethoxyaporphine. **N-Methylstenantherine**

[119060-81-0]

C₂₀H₂₃NO₄ 341.406

Alkaloid from *Neostenanthera gabonensis* (Annonaceae). [α]_D -145 (c, 0.11 in EtOH).

1,2,3-Tri-Me ether, Ac:

Oil. [α]_D -60 (c, 0.10 in EtOH).

1,2,3-Tri-Me ether, N-de-Me: 11-Hydroxy-1,2,3-trimethoxynoraporphine. **Stenantherine**

[119089-37-1]

C₁₉H₂₁NO₄ 327.379

Alkaloid from *Neostenanthera gabonensis* (Annonaceae). Oil.

1,2-Methylene, 3,11-di-Me ether, N-de-Me: 3,11-Dimethoxy-1,2-methylenedioxyaporphine. **3-Methoxyputerine**

[158018-12-3]

C₁₉H₁₉NO₄ 325.363


Alkaloid from stem bark of *Guatteria foliosa* (Annonaceae). Amorph. [α]_D -75 (EtOH).

Renner, C. *et al.*, *J. Nat. Prod.*, 1988, **51**, 973 (*Stenantherine, N-Methylstenantherine*)

Mahiou, V. *et al.*, *J. Nat. Prod.*, 1994, **57**, 890 (*3-Methoxyputerine*)

1,2,9,10-Tetrahydroxyaporphine T-246

5,6,6a,7-Tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-1,2,9,10-tetrol, 9CI



C₁₇H₁₇NO₄ 299.326

Log P 2.97 (uncertain value) (calc).

(R)-form

Tetra-Me ether: [38325-02-9]

Mp 124-125°. $[\alpha]_D -115.4$ (EtOH aq.).

(S)-form

1,9-Di-Me ether: see Liriotulipiferine, L-195

1,10-Di-Me ether: see Boldine, B-243

2,9-Di-Me ether: see Bracteoline, B-276

2,10-Di-Me ether: see Isoboldine, I-196

9,10-Di-Me ether: see Lastourvilline, L-52

1,2,9-Tri-Me ether: see 10-Hydroxy-1,2,9-trimethoxyaporphine, H-752

1,2,10-Tri-Me ether: see Lauroschoztzine, L-75

1,9,10-Tri-Me ether: see Predicentrine, P-597

2,9,10-Tri-Me ether: see 1-Hydroxy-2,9,10-trimethoxyaporphine, H-751

Tetra-Me ether: *1,2,9,10-Tetramethoxyaporphine. Glaucine* [475-81-0]

$C_{21}H_{25}NO_4$ 355.433

Alkaloid from a wide variety of genera in the Annonaceae (*Alphonsea*, *Annona*, *Polyalthia*, *Schefferomitra*, *Uvaria*), Berberidaceae (*Berberis*, *Mahonia*), Euphorbiaceae (*Croton*), Papaveraceae (*Corydalis*, *Dicentra*), Lauraceae (*Beilschmiedia*, *Ocotea*, *Litsea*), Magnoliaceae (*Liriodendron*, *Magnolia*), Menispermaceae (*Chasmanthera*), Papaveraceae (*Glaucium*, *Papaver*), Ranunculaceae (*Thalictrum*, *Aconitum*), Rhamnaceae (*Colubrina*) and Monimiaceae (*Hedycarya*). Shows antithrombotic, analgesic and

antiinflammatory activity. Antifungal agent. Produces narcosis and convulsions in animals, also hypotension and respiratory depression. Antitussive agent with similar potency to Codeine. Mild muscle relaxant. Cryst. (EtOAc or petrol). Mp 120-121°. $[\alpha]_D +116$ (c, 0.75 in EtOH). Log P 2.97 (uncertain value) (calc). Pharmacol. active isomer.

► LD₅₀ (rat, orl) 545 mg/kg. LD₅₀ (rat, ipr) 143 mg/kg LD₅₀ (mus, ivn) 98 mg/kg. CE0925000

Tetra-Me ether; hydrobromide: Bromcholitit. Glauvent. Tussiglaucin [5996-06-5]
Cryst. Mp 235°.

Tetra-Me ether, N-Me: Mp 225-227° dec. (as iodide). $[\alpha]_D^{24} +80$ (EtOH). Prepd. as *O*-Methylxanthoplanine iodide.

Tetra-Me ether, N-de-Me: *1,2,9,10-Tetramethoxynoraporphine. Norglaucine* [21848-62-4]

$C_{20}H_{23}NO_4$ 341.406

Alkaloid from a variety of genera in the Annonaceae (*Alphonsea*, *Duguetia*, *Pseudoalaria*), Magnoliaceae (*Magnolia*, *Liriodendron*), Menispermaceae (*Chasmanthera*), Monimiaceae (*Monimia*) and Rhamnaceae (*Colubrina*). Noncryst.

► RB6000200

Tetra-Me ether, N-de-Me; hydrobromide: Mp 248-250°. $[\alpha]_D^{30} +102$ (c, 1 in MeOH).

Tetra-Me ether, N-de-Me, N-Ac: Mp

102-104°. $[\alpha]_D +333$ (c, 0.13 in MeOH).

1,2-Methylene, 9,10-di-Me ether: see Dicentrine, D-346

9,10-Methylene, 1,2-di-Me ether: see Nantenine, N-29

3-Chloro, tetra-Me ether, N-de-Me, N-methoxycarbonyl: Romucosine F $C_{22}H_{24}ClNO_6$ 433.887

Alkaloid from *Annona purpurea*. Needles. Mp 140-142°. $[\alpha]_D^{35} +134$ (c, 0.05 in CHCl₃). λ_{max} 222 (log ϵ 4.35); 280 (log ϵ 4.21); 305 (log ϵ 3.76) (MeOH).

(±)-form

Tetra-Me ether: [5630-11-5]
Mp 137-139°.

Tetra-Me ether, N-de-Me: [39945-40-9]
Synthetic. Gum.

Tetra-Me ether, N-de-Me, picrate: Yellow cryst. (MeOH/EtOH). Mp 149-150°.

6a,7-Didehydro, tetra-Me ether: see Dehydroglauconine, D-159

[73239-87-9]

Go, J. *et al.*, *Yakugaku Zasshi*, 1930, **50**, 933 (isol)

Corrodi, H. *et al.*, *Helv. Chim. Acta*, 1956, **39**, 889 (abs config)

Baarschers, W.H. *et al.*, *J.C.S.*, 1964, 4478 (pmr, ms)

Jackson, A.H. *et al.*, *J.C.S. (C)*, 1966, 2061 (synth, pmr, uv)

Shamma, M. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 262 (isol, ir)

Kametani, T. *et al.*, *Tetrahedron*, 1969, **25**, 3667 (synth, ir, uv)

Johns, S.R. *et al.*, *Aust. J. Chem.*, 1970, **23**, 423 (isol, pmr, Norglaucine)

Duchevska, K.H.B. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1973, **26**, 899; *CA*, **80**, 27410x (6,6a-Dehydronorglaucine)

Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1975, **38**, 275; 290; 1979, **42**, 325; 329 (cmr, uv, rev)

Premila, M.S. *et al.*, *Indian J. Chem., Sect. B*, 1976, **14**, 134 (synth)

Aleshinskaya, E.E. *et al.*, *Khim.-Farm. Zh.*, 1976, **10**, 144 (pharmacol)

Hartenstein, J. *et al.*, *Angew. Chem.*, 1977, **89**, 739 (synth)

Davis, P.J. *et al.*, *J.C.S. Perkin 1*, 1977, 1 (synth, ms)

Ricca, G. *et al.*, *Gazz. Chim. Ital.*, 1979, **109**, 1 (cmr)

Kase, Y. *et al.*, *Arzneim.-Forsch.*, 1983, **33**, 936; 947 (pharmacol)

Kerr, K.M. *et al.*, *J. Nat. Prod.*, 1986, **49**, 576 (pmr)

Czarnocki, Z. *et al.*, *Can. J. Chem.*, 1987, **65**, 2356 (synth)

Gupta, S. *et al.*, *Synth. Commun.*, 1989, **19**, 393 (synth)

Gottlieb, L. *et al.*, *J.O.C.*, 1990, **55**, 5659 (synth)

Ivorra, M.D. *et al.*, *Br. J. Pharmacol.*, 1992, **106**, 387 (pharmacol)

Oralla, F. *et al.*, *Br. J. Pharmacol.*, 1993, **110**, 943; 1995, **114**, 1419 (pharmacol)

Ozaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1993, **41**, 481 (synth)

Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 747

Estévez, J.C. *et al.*, *Tetrahedron*, 1994, **50**, 2107 (synth)

Comins, D.L. *et al.*, *Tetrahedron*, 1997, **48**, 16327-16340 (synth)


Chang, F.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 746-748 (*Romucosine F*)

Huang, W.-J. *et al.*, *Synth. Commun.*, 2002, **32**, 3681-3686 (synth, pmr, ms)

Huang, W.-J. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 167-174 (synth)

Gao, Y. *et al.*, *Synthesis*, 2004, 1093-1101 (*Glaucine*, synth, pmr, cmr, ms)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TD1475

1,2,9,11-Tetrahydroxyaporphine T-247

$C_{17}H_{17}NO_4$ 299.326


(S)-form

2,9-Di-Me ether, N-Me: Trilobinine [126595-92-4]

$C_{20}H_{24}NO_4^{\oplus}$ 342.414

Quaternary alkaloid from *Cissampelos glaberrima* and *Thalictrum acutifolium*. $[\alpha]_D +11.6$ (c, 0.06 in MeOH). Counterion not specified.

Lin, C.W. *et al.*, *Zhiwu Xuebao (Acta Bot. Sin.)*, 1989, **31**, 449-452; *CA*, **112**, 175576a

1,2,10,11-Tetrahydroxyaporphine T-248

(S)-form

$C_{17}H_{17}NO_4$ 299.326

(S)-form

1-Me ether: 2,10,11-Trihydroxy-1-methoxyaporphine. Glaufine [70474-42-9]

$C_{18}H_{19}NO_4$ 313.352

Alkaloid from *Glaucium fimbriigerum* (Papaveraceae). $[\alpha]_D +83$ (c, 0.2 in MeOH).

1,2-Di-Me ether: 10,11-Dihydroxy-1,2-dimethoxyaporphine. Suaveoline [99026-98-9]

$C_{19}H_{21}NO_4$ 327.379

Alkaloid from the bark of *Artabotrys suaveolens* (Annonaceae). Bundles of needles or prisms (MeOH). Mp 232°. $[\alpha]_D^{15} +164$ (c, 1.22 in CHCl₃).

1,2-Di-Me ether, N-Me: Fuzitine [142287-96-5]

$C_{20}H_{24}NO_4^{\oplus}$ 342.414

Quaternary alkaloid from tubers of *Aconitum carmichaeli* (Ranunculaceae). Mp 209-211° dec. $[\alpha]_D +258$ (c,

- 0.59 in MeOH). Counterion not specified.
- 1,2-Di-Me ether, N-de-Me: 10,11-Dihydroxy-1,2-dimethoxynoraporphine**
C₁₈H₁₉NO₄ 313.352
Alkaloid from *Xylopiya parviflora*. Amorph. powder. [α]_D²² +57 (c, 0.46 in MeOH). λ_{max} 266 (log ε 4.09); 273 (sh); 306 (log ε 3.85) (CHCl₃).
- 1,10-Di-Me ether, 2,11-Dihydroxy-1,10-dimethoxyaporphine. N-Methylindcarpine. Phoebe base**
[14028-97-8]
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Lindera pipericarpa* and *Phoebe clemensii*, also from *Magnolia*, *Beilschmiedia* and *Glaucium* spp. (Lauraceae, Magnoliaceae, Papaveraceae). Mp 198-200°. [α]_D²² +222 (c, 0.8 in MeOH). λ_{max} 270 (log ε 4.2); 303 (log ε 3.8) (no solvent reported).
- 1,10-Di-Me ether, N-Me: N,N-Dimethylindcarpine**
[5890-24-4 (iodide)]
C₂₀H₂₄NO₄⁺ 342.414
Quaternary alkaloid from *Menispermum canadense*, also from *Magnolia*, *Caltha*, *Coscinium*, *Fibraurea* and *Pycnarrhena* spp. (Menispermaceae, Magnoliaceae, Ranunculaceae). Mp 249-251° dec. (as iodide). [α]_D²⁷ +213 (c, 0.6 in MeOH) (iodide).
- 1,10-Di-Me ether, N-de-Me: 2,11-Dihydroxy-1,10-dimethoxynoraporphine. Lindcarpine**
[11010-63-2]
[14028-95-6]
C₁₈H₁₉NO₄ 313.352
Alkaloid from *Lindera pipericarpa* (Lauraceae). Mp 195° dec. [α]_D²² +166 (c, 1.35 in EtOH). λ_{max} 218 (log ε 4.57); 267 (log ε 4.14); 303 (log ε 3.82) (EtOH).
- 1,10-Di-Me ether, N-de-Me, hydrochloride: [14028-96-7]**
Mp 200° dec.
- 1,11-Di-Me ether, 2,10-Dihydroxy-1,11-dimethoxyaporphine. N-Methylhernovine**
[5550-65-2]
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Croton wilsonii* and *Hernandia peltata* (Euphorbiaceae, Hernandiaceae). Amorph. Mp 244-245° dec. (as hydrochloride). [α]_D²⁴ +209 (c, 0.55 in MeOH). λ_{max} 218 (4.41); 273 (4.02); 305 (3.63) (EtOH).
- 1,11-Di-Me ether, N-de-Me: 2,10-Dihydroxy-1,11-dimethoxynoraporphine. Hernovine†**
[5544-69-4]
C₁₈H₁₉NO₄ 313.352
Alkaloid from the bark of *Hernandia ovigera*, *Hernandia catalpifolia* and *Hernandia jamaicensis*. Also present in *Croton wilsonii* (Hernandiaceae, Euphorbiaceae). Plates (MeOH). Mp 235-237° dec. [α]_D¹⁸ +142 (c, 0.52 in Py). [α]_D +266 (EtOH). λ_{max} 221 (log ε 4.41); 272 (log ε 4.01); 306 (log ε 3.64) (EtOH).
- 2,10-Di-Me ether: see Corytuberine, C-702**
- 2,11-Di-Me ether: 1,10-Dihydroxy-2,11-dimethoxyaporphine. Isocorytuberine**
[6870-51-5]
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Glaucium fimbrilligerum*, *Glaucium oxylobum* and *Trivalvaria macrophylla* (Papaveraceae, Annonaceae). Mp 223° (as hydrochloride).
- 2,11-Di-Me ether, N-de-Me: 1,10-Dihydroxy-2,11-dimethoxynoraporphine. Norisocorytuberine**
[131989-91-8]
C₁₈H₁₉NO₄ 313.352
Alkaloid from *Trivalvaria macrophylla* (Annonaceae). [α]_D +170 (c, 0.03 in EtOH).
- 10,11-Di-Me ether, N-Me: Zizyphusine**
[107446-79-7]
C₂₀H₂₄NO₄ 342.414
Alkaloid from seeds of *Zizyphus vulgaris* var. *spinosa* and *Zizyphus jujuba* var. *inermis* (Rhamnaceae). Shows sedative props. Mp 214-216°. [α]_D +317. Defective diag. in paper. Counterion not specified.
- 1,2,10-Tri-Me ether: see Isocorydine, I-208**
- 1,2,11-Tri-Me ether: 10-Hydroxy-1,2,11-trimethoxyaporphine. Praecoxine. N-Methylhernagine**
[4668-25-1]
C₂₀H₂₃NO₄ 341.406
Alkaloid from the roots of *Parabenzoin praecox* (Lauraceae). Cryst. (Me₂CO). Mp 230° dec. [α]_D²⁷ +270 (c, 0.1 in MeOH). λ_{max} 271 (ε 12380); 302 (ε 5000) (95% EtOH).
- 1,2,11-Tri-Me ether, N-de-Me: 10-Hydroxy-1,2,11-trimethoxynoraporphine. Hernagine. Glaufinine**
[74133-19-0]
C₁₉H₂₁NO₄ 327.379
Alkaloid from the leaves of *Hernandia nymphaefolia* and *Hernandia peltata*, the stem bark of *Hernandia cordigera* and the aerial parts of *Glaucium fimbrilligerum* (Hernandiaceae). Cryst. (Me₂CO). Mp 222°. [α]_D +252 (c, 1.3 in CHCl₃). [α]_D +165 (c, 0.4 in MeOH). Hernagine and Glaufinine not compared, but same struct. assigned. λ_{max} 222 (log ε 4.35); 270 (log ε 3.82); 309 (log ε 3.43) (no solvent reported).
- 1,10,11-Tri-Me ether: 2-Hydroxy-1,10,11-trimethoxyaporphine. N,O-Dimethylhernovine. Litseglutine B**
[25368-01-8]
C₂₀H₂₃NO₄ 341.406
Alkaloid from *Croton wilsonii*, *Litsea glutinosa* and *Litsea laeta*. Amorph.; cryst. (as hydrochloride). Mp 218-219° dec. (as hydrochloride). [α]_D²³ +139 (c, 0.51 in MeOH). λ_{max} 220 (log ε 4.55); 273 (log ε 4.11); 304 (log ε 3.69) (EtOH).
- 1,10,11-Tri-Me ether, N-de-Me: 2-Hydroxy-1,10,11-trimethoxynoraporphine.**
- 10-O-Methylhernovine**
[17807-65-7]
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Croton wilsonii* (Euphorbiaceae). Mp 157-158° dec. [α]_D¹⁷ +188 (c, 1.0 in EtOH). λ_{max} 220 (log ε 4.56); 273 (log ε 4.11); 305 (log ε 3.7) (EtOH).
- 2,10,11-Tri-Me ether: see Corydine, C-682**
- Tetra-Me ether: 1,2,10,11-Tetramethoxyaporphine. O-Methylcorydine. O,O-Dimethylcorytuberine**
[6191-46-4]
C₂₁H₂₅NO₄ 355.433
Alkaloid from *Hernandia catalpifolia* and *Hernandia jamaicensis* and from stems of *Chasmanthera dependens* (Hernandiaceae, Menispermaceae). Also a methylation prod. of other aporphine alkaloids, e.g. Isocorydine, I-208. Mp 226-227° dec. (as tartrate). [α]_D²⁸ +147 (c, 0.7 in H₂O).
- Tetra-Me ether, N-oxide: O-Methylcorydine N-oxide**
[97457-23-3]
C₂₁H₂₅NO₅ 371.432
Alkaloid from whole plants of *Berberis chitria* (Berberidaceae). [α]_D³⁵ +193 (c, 1 in MeOH).
- Tetra-Me ether, N-Me: N,O-Dimethylisocorydine. O,O-Dimethylmagnoflorine**
[51827-27-1]
[58489-51-3, 5489-15-6]
C₂₂H₂₈NO₄⁺ 370.467
Quaternary alkaloid from leaves of *Cocculus laurifolius* and roots of *Pachygone ovata* (Menispermaceae). Cryst. (MeOH/Et₂O) (as chloride). Mp 228-230° dec. (chloride).
- Tetra-Me ether, N-de-Me: 1,2,10,11-Tetramethoxynoraporphine. Catalpifoline. Hemiargine A**
[10214-65-0]
C₂₀H₂₃NO₄ 341.406
Alkaloid from the bark of *Hernandia catalpifolia* and *Hernandia jamaicensis*. Also from *Croton hemiargyreus* var. *gymnodiscus*. Mp 174-175°. [α]_D²⁵ +220.5 (c, 0.10 in EtOH).
- 1,2-Methylene, 10-Me ether: see Bulbocapnine, B-397**
- 1,2-Methylene, 11-Me ether, N-de-Me: see Nandigerine, N-25**
- 10,11-Methylene, 1-Me ether, N-de-Me: see Laetine, L-10**
- 4-Oxo, 1,10-di-Me ether, N-Me: Saxicolaline A**
[1006029-76-0]
C₂₀H₂₂NO₅⁺ 356.398
Quaternary alkaloid from *Corydalis saxicola*. Yellow cryst. (MeOH). Mp 223-225°. [α]_D²⁷ +718.8 (c, 0.16 in Py). Counterion not specified. λ_{max} 205 (log ε 4.39); 231 (log ε 4.23); 289 (log ε 4.09); 358 (log ε 4.05) (MeOH).
- (±)-form**
2,11-Di-Me ether: [2273-24-7]

Synthetic. Mp 223° (as hydrochloride).
1,10,11-Tri-Me ether: [32563-01-2]
Synthetic. Yellow syrup; needles
(MeOH/Et₂O) (as hydrochloride). Mp
245-248° (hydrochloride).

(E)-form

5ξ-Hydroxy, tetra-Me ether, N-Me:

Magnoporphine

[188578-70-3]
C₂₂H₂₈NO₅[⊕] 386.467

Alkaloid from the leaves of *Magnolia sieboldii* (Magnoliaceae).

Barger, G. et al., *J.C.S.*, 1939, 991-997
(*Suaveoline*)

Jackson, A.H. et al., *J.C.S. (C)*, 1966, 2181-2183; 2222-2229 (*Isocorytuberine*, synth, pmr, ms)

Cava, M.P. et al., *Tet. Lett.*, 1966, 1577-1581 (*Hernovine*)

Johns, S.R. et al., *Aust. J. Chem.*, 1967, **20**, 1277-1281 (*N-Methylindocarpine*, isol, pmr)

Kiang, A.K. et al., *J.C.S. (C)*, 1967, 282-283 (*Lindocarpine*, isol, uv, pmr)

Stuart, K.L. et al., *Tet. Lett.*, 1967, 4135-4138 (*N-Methylhernovine*, 10-O-Methylhernovine, N,O-Dimethylhernovine)

Kametani, T. et al., *J.C.S. (C)*, 1971, 1923-1927 (*N,O-Dimethylhernovine*, synth)

Cava, M.P. et al., *Tetrahedron*, 1971, **27**, 2639-2643 (*Catalpifoline*)

Guinaudeau, H. et al., *J. Nat. Prod.*, 1975, **38**, 275-338; 1994, **57**, 1033-1035 (rev)

Karimova, S.U. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 699 (*Glaufine*)

Saxena, N.K. et al., *J. Indian Chem. Soc.*, 1979, **56**, 1020-1023 (*N,O-Dimethylisocorydine*)

Bruneton, J. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1980, **291**, 187-189 (*Hernagine*, uv, ms, struct)

Bhat, S.V. et al., *J. Nat. Prod.*, 1980, **43**, 588-591 (*O,O-Dimethylmagnoflorine*)

Yakushijin, K. et al., *Phytochemistry*, 1980, **19**, 161-162 (*Hernagine*, pmr, struct)

Rastogi, R.C. et al., *Phytochemistry*, 1980, **19**, 998-999 (*N,O-Dimethylhernovine*, isol, uv, ir, pmr, ms)

Ringdahl, B. et al., *J. Nat. Prod.*, 1981, **44**, 80-85 (*Hernovine*, cd)

Suess, T.R. et al., *J. Nat. Prod.*, 1981, **44**, 688-692 (synth)

Lavault, M. et al., *Planta Med.*, 1981, **42**, 50-54; 1982, **46**, 119-121 (*Hernagine*, N-Methylhernovine, isol, uv, pmr, ms)

Kozuka, M. et al., *Chem. Pharm. Bull.*, 1984, **32**, 5055-5058 (*Praecoxine*)

Hussaini, F.A. et al., *Phytochemistry*, 1985, **24**, 633 (*O-Methylcorydine N-oxide*)

Israilov, I.A. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **22**, 236 (*Glaufinine*)

Han, B.H. et al., *Arch. Pharmacol. Res.*, 1989, **12**, 263-268 (*Zizyphusine*, isol)

Han, B.H. et al., *Pure Appl. Chem.*, 1989, **61**, 443-448 (*Zizyphusine*, occur)

Cortes, D. et al., *J. Nat. Prod.*, 1990, **53**, 862-866 (*Norisocorytuberine*)

Chen, H. et al., *Chin. Chem. Lett.*, 1991, **2**, 787; *CA*, **117**, 44535s (*Fuzitine*)

Park, H.J. et al., *CA*, 1997, **126**, 235966d (*Magnoporphine*)

Rasoanaivo, P. et al., *Planta Med.*, 1998, **64**, 58-62 (cmr)


Yang, J.-H. et al., *Helv. Chim. Acta*, 2005, **88**, 2523-2526 (*Litsegultine B*)

Nishiyama, Y. et al., *Phytochemistry*, 2006, **67**, 2671-2675 (*10,11-Dihydroxy-1,2-dimethoxynoraporphine*)

Wu, Y.-R. et al., *Planta Med.*, 2007, **73**, 787-791 (*Saxicolaline A*)

3,3',4,4'-Tetrahydroxy-2,2'-bipyridine T-249

[2,2'-Bipyridine]-3,3',4,4'-tetrol, 9CI.
2,2'-Bi[3-hydroxy-4(1H)-pyridinone].
Orelline. *Orellanine II*
[72016-31-0]



C₁₀H₈N₂O₄ 220.184

Alkaloid detected in the toxic mushroom *Cortinarius orellanus*; formed by photochem. or thermal dec. of Orellanine.

Yellow solid. Sublimes readily. λ_{max} 243; 260; 285; 385; 407 (MeOH/NaOH) (Derep). λ_{max} 219 (ε 22500); 344 (ε 7900); 390 (ε 2400) (MeOH) (Derep).

Mono-N-oxide: Orellanine

[98726-96-6]
C₁₀H₈N₂O₅ 236.184

Alkaloid detected in *Cortinarius orellanus*, *Cortinarius scioosimus* and *Cortinarius rubellus*. Formed by photochem. or thermal dec. of Orellanine. Mycotoxin. Convulsive and nephrotoxic agent. Sol. DMSO, acids, bases; fairly sol. MeOH; poorly sol. butanol, H₂O, hexane. λ_{max} 213; 267; 342 (MeOH) (Berdy). λ_{max} 209; 261 (HCl) (Berdy). λ_{max} 223; 305 (NaOH) (Berdy).

► Toxic.

Di-N-oxide: Orellanine

[37338-80-0]
C₁₀H₈N₂O₆ 252.183

Alkaloid from *Cortinarius orellanus* and *Cortinarius speciosissimus*. Nephrotoxin, mycotoxin. Sol. bases, NH₄OH, DMSO, Py, THF; fairly sol. MeOH, acids; poorly sol. H₂O, butanol, hexane. Dec. slowly >150°, explosively at 267° to Orelline and O₂. λ_{max} 213; 265; 290 (MeOH/HCl) (Derep). λ_{max} 234; 292; 319 (MeOH/NaOH) (Derep). λ_{max} 219 (ε 21800); 248 (ε 13200); 282 (ε 8400); 352 (ε 7300) (MeOH) (Derep).

► Highly toxic, induces acute renal failure LD₅₀ = 5-8 mg/kg. Long latency period. RM0525000

Di-N-oxide, 4,4'-di-O-β-D-glucopyranoside: Orellanine 4,4'-diglucoside

C₂₂H₂₈N₂O₁₆ 576.467

Alkaloid from *Cortinarius orellanus*. Orellanine occurs mainly as this diglucoside. λ_{max} 237 (log ε 3.68); 262 (sh) (log ε 3.14); 318 (log ε 3.15) (H₂O).

3,3'-Di-Me ether: [405137-22-6]

C₁₂H₁₂N₂O₄ 248.238
Mp 151°.

4,4'-Di-Me ether: [104819-53-6]

C₁₂H₁₂N₂O₄ 248.238
Mp 252-254°.

3,3',4,4'-Tetra-Me ether: 3,3',4,4'-Tetramethoxy-2,2'-bipyridine

[101664-55-5]
C₁₄H₁₆N₂O₄ 276.291
Mp 186-187°.

Antkowiak, W.Z. et al., *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1975, **23**, 729 (isol)

Antkowiak, W.Z. et al., *Tet. Lett.*, 1979, 1931 (ir, uv, struct)

Tiecco, M. et al., *Tetrahedron*, 1986, **42**, 1475-1485 (synth, pmr, cmr, ms, uv)

Dehmlow, E.V. et al., *Annalen*, 1987, 857 (synth)

Tiecco, M. et al., *Experientia*, 1987, **43**, 462 (*Orellanine*, synth, uv, pmr, cmr)

Hasseberg, H.-A. et al., *Helv. Chim. Acta*, 1988, **71**, 957 (synth, ir, pmr, cmr)

Trécourt, F. et al., *Tetrahedron*, 1993, **49**, 8373 (synth)

Antkowiak, W.Z. et al., *Heterocycles*, 1994, **39**, 477; 2002, **58**, 137-146 (*Orellanine*, ms, pmr, atropisomerism, bibl)

Oubrahim, H. et al., *J. Chromatogr. A*, 1997, **758**, 145-157 (tlc)

Oubrahim, H. et al., *Free Radical Res.*, 1998, **28**, 497-505 (biochem)


Mongin, F. et al., *Tetrahedron*, 2002, **58**, 309-314 (synth, Me ethers)

Spiteller, P. et al., *Angew. Chem., Int. Ed.*, 2003, **42**, 2864-2867 (*Orellanine diglucoside*)

Michelot, D. et al., *Nat. Prod. Res.*, 2003, **17**, 41-46 (*Orellanine*, synth)

2-(1,2,3,4-Tetrahydroxybutyl)-5-(2,3,4-trihydroxybutyl)pyrazine T-250

1-[5-(2,3,4-Trihydroxybutyl)pyrazinyl]-1,2,3,4-butanetetrol, 9CI
[68510-02-1]



C₁₂H₂₀N₂O₇ 304.299

(1'R,2'R,2''R,3'R,3''R)-form

D-lyxo-*D*-threo-*form*

[220145-37-9]

Cryst. (MeOH). Mp 146°. [α]_D²⁰ -14.6 (c, 0.2 in H₂O).

(1'R,2'S,2''S,3'R,3''R)-form

D-arabino-*D*-erythro-*form*. *Deoxyfructosazine*

Exhibits DNA strand cleavage activity. Solid (MeOH/EtOH); cryst. (2-propanol aq.). Mp 161-162° (157-158°). [α]_D^{23.5} -79.8 (c, 0.2 in H₂O).

Hepta-Ac: [52057-08-6]

Oil. [α]_D²⁰ -18 (c, 1.0 in CHCl₃).

(1'R,2'S,2''S,3'S,3''S)-form

L-xylo-*L*-threo-*form*

[220145-38-0]

Beige cryst. (EtOH aq.). Mp 116°. [α]_D²⁰ -62.4 (c, 0.5 in H₂O).

(1'S,2'R,2''R,3'R,3''R)-form

D-xylo-*D*-threo-*form*

[220145-36-8]

Beige solid (EtOH/MeOH/Et₂O). Mp 90°. [α]_D²⁰ +71.3 (c, 0.5 in MeOH).

(1'S,2'S,2''S,3'R,3''R)-form

D-ribo-*D*-erythro-*form*


[220145-41-5]

Ochre cryst. (EtOH aq.). Mp 141°.

(1ξ,2ξ,2'ξ,3ξ,3'ξ)-form

Pedatisectine G

[206757-33-7]


Alkaloid from the rhizomes of *Pinellia pedatisecta*.Kuhn, R. *et al.*, *Annalen*, 1961, **644**, 122-127 (synth)Chilton, W.S. *et al.*, *J.A.C.S.*, 1967, **89**, 4129-4132 (ord, uv)Tsuchida, H. *et al.*, *Agric. Biol. Chem.*, 1973, **37**, 2571-2578; 1975, **39**, 1143-1148 (deoxyfructosazine, isol, pmr, uv)Tsuchida, H. *et al.*, *Carbohydr. Res.*, 1978, **67**, 549-563 (ms)Eitelman, S.J. *et al.*, *Carbohydr. Res.*, 1979, **77**, 205-211 (deoxyfructosazine, synth)Tsuchida, H. *et al.*, *Dev. Food Sci.*, 1986, **13**, 85-94 (occur)Tsuchida, H. *et al.*, *CA*, 1990, **113**, 229877r (occur, soy sauce)Sumoto, K. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 792-794 (deoxyfructosazine, synth)Magaletta, R.L. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 2629-2635 (anal, hplc)Wang, R. *et al.*, *Zhongguo Zhongyao Zazhi*, 1997, **22**, 421-423; *CA*, **128**, 306181y (Pedatisectine G)Pat. Coop. Treaty (WIPO), 1999, 9903842; *CA*, **130**, 139579s (stereoisomers, synth, pmr)Rohovec, J. *et al.*, *Eur. J. Org. Chem.*, 2001, 3899 (deoxyfructosazine, synth)**1,2,6,7-Tetrahydroxydibenz[cd,f]indol-4(5H)-one, 9CI** T-251C₁₅H₉NO₅ 283.24

CAS numbering shown. Alternative (phenanthrene) numbering frequently used, in which 1,2,6,7- are 4,3,9,8- respectively.


1,6,7-Tri-Me ether: **Enterocarpam I** [102719-94-8]C₁₈H₁₅NO₅ 325.32Alkaloid from the stem bark of *Ororopea enterocarpa* (Annonaceae). Shows platelet aggregation inhibitory activity. Cryst. (CHCl₃/MeOH). Mp 214°.

1,6,7-Tri-Me ether, N,O-di-Ac:


Pale yellow needles (MeOH). Mp 186°.

1,2-Methylene, 7-Me ether: **9-Hydroxyaristolactam I** [127172-86-5]C₁₇H₁₁NO₅ 309.278Alkaloid from rhizomes of *Aristolochia brevipes*, also prod. from 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262 by treatment with xanthine oxidase. Light yellow fluorescent cryst. (MeOH). Mp 290° dec.1,2-Methylene, 7-Me ether, 6-O-[β-D-glucopyranosyl-(1→6)-β-D-glucopyranoside]: **Triangularine B** [197006-83-0]C₂₉H₃₁NO₁₅ 633.562Alkaloid from the roots of *Aristolochia triangularis*.1,2-Methylene, 6,7-di-Me ether: **9-Methoxyaristolactam I** [133462-30-3]C₁₈H₁₃NO₅ 323.304Alkaloid from *Aristolochia auricularia*. Yellow solid.1,2-Methylene, 6-Et, 7-Me ether: **9-Ethoxyaristolactam I**C₁₉H₁₅NO₅ 337.331Alkaloid from *Aristolochia mollissima*. Component of Mian Mao Ma Dou Ling. Cryst. Mp 237-238°.Mahmood, K. *et al.*, *Phytochemistry*, 1986, **25**, 965 (*Enterocarpam I*)Lou, F. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 305-307 (*9-Ethoxyaristolactam I*)Houghton, P.J. *et al.*, *Phytochemistry*, 1991, **30**, 253 (*9-Methoxyaristolactam I*)Achenbach, H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 918-922 (*9-Hydroxyaristolactam I*)Lin, W. *et al.*, *J. Chin. Pharm. Sci.*, 1997, **6**, 8-13 (*Triangularine B*)Cao, S.-G. *et al.*, *Tetrahedron*, 1998, **54**, 2143-2148 (*Enterocarpam I*, activity)Chia, Y.C. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1160-1163 (*Enterocarpam I*, isol, activity)**1,2,7,8-Tetrahydroxydibenz[cd,f]indol-4(5H)-one** T-252C₁₅H₉NO₅ 283.24

Alternative (phenanthrene-type) numbering has also been used for this type of compd., in which posns. 1,2,8- become 4,3,7- respectively.

2,7-Di-Me ether: **Aristolactam E** [710319-83-8]C₁₇H₁₃NO₅ 311.293Alkaloid from *Aristolochia elegans*. Yellow syrup. λ_{max} 236 (log ε 4.22); 253 (log ε 4.12); 282 (log ε 3.65); 332 (log ε 3.52); 390 (log ε 3.44) (MeOH).1,2-Methylene, 8-Me ether: **Cepharanone C** [293314-08-6]C₁₇H₁₁NO₅ 309.278Alkaloid from *Aristolochia cucurbitifolia*. Yellowish powder. λ_{max} 236 (sh) (log ε 4.45); 241 (log ε 4.49); 276 (sh) (log ε 4.35); 288 (log ε 4.49); 357 (log ε 3.85); 373 (log ε 3.83); 395 (log ε 3.74) (MeOH).Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 1006-1009 (*Cepharanone C*)Shi, L.-S. *et al.*, *Bioorg. Med. Chem.*, 2004, **12**, 439-446 (*Aristolactam E*)**1,2,7,9-Tetrahydroxydibenz[cd,f]indol-4(5H)-one, 9CI** T-253C₁₅H₉NO₅ 283.24

CAS numbering shown. Alternative (phenanthrene) numbering frequently used in which 1,2,7,9- become 4,3,8,6- respectively.


1,2-Methylene, 7,9-di-Me ether: **Aristolactam IV** [17413-39-7]C₁₈H₁₃NO₅ 323.304Alkaloid from the roots of *Aristolochia indica* and the rhizomes of *Aristolochia argentina* (Aristolochiaceae). Yellow cryst. by subl. Mp 353° (345°).Kupchan, S.M. *et al.*, *J.O.C.*, 1968, **33**, 3735 (isol, uv, ir, pmr, struct, synth)Priestap, H.A. *et al.*, *Phytochemistry*, 1985, **24**, 849 (isol, uv, ir, pmr, ms)**1,2,3,7-Tetrahydroxydibenz[cd,f]indol-4(5H)-one, 9CI** T-254C₁₅H₉NO₅ 283.24

Numbering systems vary.

1,2,3-Tri-Me ether: **Oldhamlactam. Oldhamactam** [921936-15-4]C₁₈H₁₅NO₅ 325.32Alkaloid from the stems of *Fissistigma oldhamii*. Yellow-brown powder. Mp 224-226°. λ_{max} 252 (log ε 4.24); 296 (log ε 4.82); 397 (log ε 4.92) (CHCl₃).1,2-Methylene, 7-Me ether: **2-Hydroxy-8-methoxycepharanone A** [157196-46-8]C₁₇H₁₁NO₅ 309.278Alkaloid from roots of *Aristolochia cinnabarina* (Aristolochiaceae). Yellow amorph. powder. λ_{max} 230 (log ε 5.52); 250 (sh) (log ε 5.46); 290 (sh) (log ε 4.79); 425 (log ε 4.82) (MeOH).Hong, L. *et al.*, *Phytochemistry*, 1994, **37**, 237-239 (*Aristolochia cinnabarina* consti)Zhang, Y.-N. *et al.*, *Bioorg. Med. Chem.*, 2007, **15**, 988-996 (*Oldhamlactam*)

1,2,3,9-Tetrahydroxydibenz[cd,f]indol-4(5H)-one, 9CI

T-255


C₁₅H₉NO₅ 283.24

Numbering systems vary.

1,2,3-Tri-Me ether: 9-Hydroxy-1,2,3-trimethoxydibenz[cd,f]indol-4(5H)-one.

StigmatalactinC₁₈H₁₅NO₅ 325.32Alkaloid from *Fissistigma oldhamii*.Brownish-yellow needles. Mp 276-278°. λ_{max} 234 (log ε 4.58); 254 (log ε 4.56); 304 (log ε 4.3); 314 (log ε 4.32) (EtOH).Chia, Y.-C. et al., *J. Nat. Prod.*, 2000, **63**, 1160-1163 (isol, pmr, cmr)Rys, V. et al., *Eur. J. Org. Chem.*, 2003, 1231-1237 (synth)**1,2,6,9-Tetrahydroxydihydro-β-agarofuran**

T-256

C₁₅H₂₆O₅ 286.367**(1α,2α,6β,9β)-form**6-(3-Pyridinecarbonyl), 9-benzoyl, 1,2-di-Ac: **Triptogelin C2**

[135118-63-7]

C₃₂H₃₇NO₉ 579.646Alkaloid from *Tripterygium wilfordii*. Needles. Mp 191-194°. [α]_D²³ +43.2 (c, 0.22 in CHCl₃).6-(3-Pyridinecarbonyl), 9-benzoyl, 1-Ac: **Triptogelin C4**


[137740-10-4]

C₃₀H₃₅NO₈ 537.608Alkaloid from *Tripterygium wilfordii*. Amorph. powder. [α]_D²³ +193 (c, 0.46 in MeOH).2,6-Bis(3-pyridinecarbonyl), 9-benzoyl, 1-Ac: **Heterophylline**[†]

[379693-17-1]

C₃₆H₃₈N₂O₉ 642.704Constit. of *Maytenus heterophylla*.Cryst. Mp 132-135°. [α]_D +63.2 (c, 1 in CHCl₃). λ_{max} 234 (log ε 4.1); 258 (log ε 3.72); 264 (log ε 3.74) (CHCl₃).Takaishi, Y. et al., *Phytochemistry*, 1991, **30**, 1561-1566; 1567-1572; 3027-3031 (*Tripterygium wilfordii* constits)Orabi, K.Y. et al., *Phytochemistry*, 2001, **58**, 475-480 (*Heterophylline*)**1,4,6,9-Tetrahydroxydihydro-β-agarofuran**

T-257

C₁₅H₂₆O₅ 286.367**(1α,4β,6β,9β)-form**

6-(3-Pyridinecarbonyl), 1,9-dibenzoyl:

Regelidine

[114542-54-0]


C₃₅H₃₇NO₈ 599.679Alkaloid from the roots of *Tripterygium regelii* (Celastraceae). Prisms (CHCl₃). Mp 292-294°. [α]_D^{26.4} +50 (c, 6 in CHCl₃).

1,9-Bis(3-pyridinecarbonyl), 6-benzoyl:

[339151-93-8]

C₃₄H₃₆N₂O₈ 600.667Constit. of *Tripterygium wilfordii*.Amorph. powder. [α]_D²⁵ +67.7 (c, 0.3 in MeOH). λ_{max} 222 (log ε 4.4); 263 (log ε 3.7) (MeOH).1,6-Bis(3-pyridinecarbonyl), 9-cinnamoyl: **Wilfordicine**C₃₆H₃₈N₂O₈ 626.705Constit. of *Tripterygium wilfordii*.Needles (EtOH). Mp 189°. [α]_D²⁵ +17.4 (c, 0.5 in CHCl₃).Hori, H. et al., *Chem. Pharm. Bull.*, 1987, **35**, 4683 (*Regelidine*)He, Z.-S. et al., *J. Nat. Prod.*, 1994, **57**, 305 (*Wilfordicine*)Duan, H.-Q. et al., *Phytochemistry*, 2001, **56**, 341-346 (*Tripterygium wilfordii* constit)**1,6,8,9-Tetrahydroxydihydro-β-agarofuran**

T-258



(1α,6β,8α,9α)-form

C₁₅H₂₆O₅ 286.367**(1α,6β,8α,9α)-form**1-(3-Pyridinecarbonyl), 8,9-dibenzoyl, 6-Ac: **Triptogelin B2**

[135153-78-5]

C₃₇H₃₉NO₉ 641.716Alkaloid from *Tripterygium wilfordii*. Powder. [α]_D²³ +52.3 (c, 0.52 in MeOH).**(1α,6β,8α,9β)-form**9-(3-Pyridinecarbonyl), 6-benzoyl, 1,8-di-Ac: **Celapanigine**

[52691-07-3]

C₃₂H₃₇NO₉ 579.646Alkaloid from *Celastrus paniculatus* (Celastraceae). Cryst. (EtOH). Mp 184-185°.**(1α,6β,8β,9α)-form**

9-(3-Pyridinecarbonyl), 6-benzoyl, 1-Ac:

Celapagine

[58074-71-8]

C₃₀H₃₅NO₈ 537.608Alkaloid from *Celastrus paniculatus* (Celastraceae). Cryst. (MeOH). Mp 275-283°.9-(3-Pyridinecarbonyl), 6-(3-furancarboxonyl), 1,8-di-Ac: **Celapanine**


[52658-32-9]

C₃₀H₃₅NO₁₀ 569.607Alkaloid from *Celastrus paniculatus*(Celastraceae). Cryst. (MeOH). Mp 245-249°. [α]_D²⁵ -64.7 (c, 0.46 in CHCl₃).

4-Config. not certain.

Wagner, H. et al., *Tetrahedron*, 1975, **31**, 1949-1956 (*Celapanine*, *Celapagine*, *Celapanigine*)Takaishi, Y. et al., *Phytochemistry*, 1991, **30**, 1567-1572 (*Triptogelin B2*)**1,8,9,14-Tetrahydroxydihydro-β-agarofuran**

T-259



(1α,8α,9α)-form

C₁₅H₂₆O₅ 286.367**(1α,8α,9α)-form**

9-Benzoyl, 1,8,14-tri-Ac:

C₂₈H₃₆O₉ 516.587Constit. of *Celastrus angulatus*.Amorph. [α]_D²³ -77.6 (c, 0.515 in CHCl₃). [α]_D²³ -9.2 (c, 0.53 in CHCl₃).

9-(3-Pyridinecarbonyl), 14-benzoyl, 1,8-di-Ac: [209123-98-8]

C₃₂H₃₇NO₉ 579.646Constit. of *Celastrus angulatus*. Cryst.Mp 168-170°. λ_{max} 201 (log ε 3.23); 224 (log ε 3.28); 263 (log ε 2.56) (MeOH).

14-(3-Pyridinecarbonyl), 8,9-dibenzoyl, 1-Ac:

C₃₇H₃₉NO₉ 641.716Constit. of *Celastrus angulatus*.Amorph. powder. [α]_D²³ -102 (c, 0.53 in CHCl₃).**(1α,8β,9α)-form**1-Benzoyl, 8,9,14-tri-Ac: **Kupitengester 4**

[132185-62-7]

C₂₈H₃₆O₉ 516.587Constit. of *Celastrus angulatus*.9-Benzoyl, 1,8,14-tri-Ac: **Angulatueoid C**

[144789-24-2]

C₂₈H₃₆O₉ 516.587Constit. of *Celastrus angulatus*. Cryst.Mp 192-195°. [α]_D²³ -9.24 (c, 0.53 in CHCl₃).8-(3-Pyridinecarbonyl), 1-benzoyl, 9,14-di-Ac: **Kupitengester 2**

[149183-64-2]

C₃₂H₃₇NO₉ 579.646Constit. of *Celastrus angulatus*.

9-(3-Pyridinecarbonyl), 14-benzoyl, 1,8-di-Ac: [210763-52-3]

C₃₂H₃₇NO₉ 579.646

Constit. of *Celastrus angulatus*. Needles (Me₂CO). Mp 168-170°. [α]_D +25.4 (c, 0.52 in MeOH). λ_{\max} 201 (log ϵ 3.23); 224 (log ϵ 3.28); 263 (log ϵ 2.56) (MeOH).

14-(2-Pyridinecarbonyl), 9-benzoyl, 1,8-di-Ac: **Angulatueoid D** [145042-02-0]

C₃₂H₃₇NO₉ 579.646

Constit. of *Celastrus angulatus*. Cryst. Mp 184-185°. Unusual 2-pyridinecarbonyl residue.

14-(3-Pyridinecarbonyl), 9-benzoyl, 1-Ac: [145639-84-5]

C₃₀H₃₅NO₈ 537.608

Constit. of *Celastrus angulatus*. Amorph. powder. [α]_D²³ +47.6 (c, 0.5 in CHCl₃).

14-(3-Pyridinecarbonyl), 9-benzoyl, 1,8-di-Ac: [209123-95-5]

C₃₂H₃₇NO₉ 579.646

Constit. of *Celastrus angulatus*. Amorph. powder. [α]_D +48.6 (c, 0.35 in MeOH). λ_{\max} 201 (log ϵ 3.24); 224 (log ϵ 3.26); 263 (log ϵ 2.58) (MeOH).

14-(3-Pyridinecarbonyl), 9-benzoyl, 1-propanoyl, 8-Ac: **Angulatueoid F** [147029-06-9]

C₃₃H₃₉NO₉ 593.672

Constit. of *Celastrus angulatus*. Cryst. (Me₂CO). Mp 178.5-180.5°.

14-(3-Pyridinecarbonyl), 9-benzoyl, 1-(2-methylpropanoyl), 8-Ac: **Angulatueoid E** [147029-05-8]

C₃₄H₄₁NO₉ 607.699

Constit. of *Celastrus angulatus*. Cryst. (Me₂CO). Mp 175.5-177.5°.

14-(3-Pyridinecarbonyl), 9-benzoyl, 8-(2-methylpropanoyl), 1-Ac: [145613-76-9]

C₃₄H₄₁NO₉ 607.699

Constit. of *Celastrus angulatus*. Amorph. powder. [α]_D²³ +49.9 (c, 0.505 in CHCl₃).

14-(3-Pyridinecarbonyl), 9-benzoyl, 8-(2-methylbutanoyl), 1-Ac: [145613-75-8]

C₃₅H₄₃NO₉ 621.726

Constit. of *Celastrus angulatus*. Amorph. powder. [α]_D²³ +44.4 (c, 0.495 in CHCl₃).

9,14-Bis(3-pyridinecarbonyl), 8-benzoyl, 1-Ac: [145047-92-3]

C₃₆H₃₈N₂O₉ 642.704

Constit. of *Celastrus angulatus*. Amorph. [α]_D²³ +45.4 (c, 0.52 in CHCl₃).

(1 α ,8 β ,9 β)-form**Malkanguniol**

[42719-36-8]

Cryst. Mp 171-172°. [α]_D -32.94 (c, 3 in dioxan).

9-Benzoyl, 8-Ac: **Malkangunin**

[52691-06-2]

C₂₄H₃₂O₇ 432.513

Constit. of *Celastrus paniculatus*. Cryst. (Et₂O/petrol). Mp 240-245°. [α]_D -58.8 (c, 1 in CHCl₃).

9-(3-Furancarboxyl), 1,8,14-tri-Ac: [135153-81-0]

C₂₆H₃₄O₁₀ 506.549

Constit. of *Celastrus paniculatus*.

Amorph. powder. [α]_D²⁵ -8.6 (c, 0.32 in MeOH). λ_{\max} 212 (log ϵ 3.42); 232 (log ϵ 3.33); 274 (log ϵ 2.76) (MeOH).

Wagner, H. et al., *Tetrahedron*, 1975, **31**, 1949 (isol)

Lotter, H. et al., *Tet. Lett.*, 1978, 3243 (abs config)

Hong, S. et al., *Phytochemistry*, 1991, **30**, 1547 (deriv)

Tu, Y.Q. et al., *Chin. Chem. Lett.*, 1992, **3**, 625 (isol, pmr, cmr)

Chunquan, C. et al., *Phytochemistry*, 1992, **31**, 2777 (*Angulatueoids C and D*)

Tu, Y.Q. et al., *Phytochemistry*, 1992, **31**, 3633 (isol, pmr, cmr)

Wang, G. et al., *Zhìwù Xuebao (Acta Bot. Sin.)*, 1992, **34**, 777-780 (*Kupitengesters 2,4*)

Tu, Y.Q. et al., *J. Nat. Prod.*, 1993, **56**, 126 (isol, pmr, cmr)

Jikai, L. et al., *Phytochemistry*, 1993, **32**, 379 (*Angulatueoids E and F*)


Yong-Qiang, T. et al., *Phytochemistry*, 1993, **32**, 1339 (isol, pmr, cmr)

Li, Y. et al., *Zhongshan Daxue Xuebao Ziran Kexueban*, 1997, **36**, 123-124; *CA*, **129**, 65514x (*Celastrus angulatus esters*)

Wang, Y. et al., *J. Nat. Prod.*, 1998, **61**, 942-944 (*Celastrus angulatus esters*)

Borrelli, F. et al., *Planta Med.*, 2004, **70**, 652-656 (*Celastrus paniculatus constit*)

6,13,14,21-Tetrahydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-7,19-dien-1-one T-260

C₂₈H₃₉NO₅ 469.62

21-Ac:


C₃₀H₄₁NO₆ 511.657

Prod. by *Daldinia concentrica*. Needles. Mp 197-199°. [α]_D²⁴ +51.2 (c, 1.3 in MeOH). λ_{\max} 204 (MeOH).

Qin, X.-D. et al., *Helv. Chim. Acta*, 2006, **89**, 450-455 (isol, pmr, cmr)

1,2,6,7-Tetrahydroxy-3-(hydroxymethyl)-5-methyl-1H-pyrrolizidine T-261

Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizine-1,2,6,7-tetrol, 9CI



(1R*,2R*,3R*,5S*,6S*,7S*,7aR*)-form

C₉H₁₇NO₅ 219.237

(1R*,2R*,3R*,5S*,6S*,7S*,7aR*)-form Hyacinthacine C₅

[944408-28-0]

Alkaloid from the bulbs of *Scilla socialis*. Powder. [α]_D +1.5 (c, 0.22 in H₂O).

(1R*,2S*,3S*,5S*,6S*,7S*,7aS*)-form Hyacinthacine C₁

[240117-30-0]

Alkaloid from *Hyacinthoides non-scripta* and *Muscari armeniacum*. [α]_D +14.7 (c, 0.28 in H₂O).

(1S*,2R*,3R*,5R*,6R*,7R*,7aR*)-form Hyacinthacine C₄


[944408-27-9]

Alkaloid from the bulbs of *Scilla socialis*. Powder. [α]_D -37.9 (c, 0.44 in H₂O). Apparently the enantiomer of Hyacinthacine C₁.

Kato, A. et al., *Carbohydr. Res.*, 1999, **316**, 95-103 (*Hyacinthacine C₁*)

Kato, A. et al., *J. Nat. Prod.*, 2007, **70**, 993-997 (*Hyacinthacines C₄, C₅*)

1,2,6,7-Tetrahydroxy-3-(hydroxymethyl)-1H-pyrrolizidine T-262
Hexahydro-3-(hydroxymethyl)-1H-pyrrolizine-1,2,6,7-tetrol



(1R,2R,3R,6R,7S,7aR)-form

C₈H₁₅NO₅ 205.21**(1R,2R,3R,6R,7S,7aR)-form****6-Epicasuarine. Uniflorine A**

[194918-09-7]

[260247-75-4 (Uniflorine A)]

Struct. of Uniflorine A finally determined in 2008. Alkaloid from the leaves of *Eugenia uniflora* (Surinam cherry). α -Glucosidase inhibitor. Cryst. Mp 174-178°. [α]_D -4.4 (c, 1.2 in H₂O).

(1R,2R,3R,6S,7S,7aR)-form**Casuarine. Uniflorine B**

[159440-57-0]

[260247-76-5 (Uniflorine B)]

Structure of Uniflorine B was revised in 2008. Alkaloid from *Casuarina equisetifolia*, *Eugenia uniflora* and *Myrtus communis*. Cryst. (EtOH aq.). Mp 181-182°. [α]_D²⁴ +16.9 (c, 0.8 in H₂O).

6-O- α -D-Glucopyranoside: **Casuarine 6- α -D-glucoside**

[186795-20-0]

C₁₄H₂₅NO₁₀ 367.352

Alkaloid from *Casuarina equisetifolia* and *Eugenia jambolana* (jambolan).

(1R,2R,3S,6S,7S,7aR)-form**3-Epicasuarine**


[729593-71-9]

Alkaloid from *Myrtus communis* (myrtle). Cryst. (EtOH aq.). [α]_D²³ +5.7 (c, 0.5 in H₂O).

Nash, R.J. et al., *Tet. Lett.*, 1994, **35**, 7849-7852 (isol, pmr, cmr)

- Wormald, M.R. *et al.*, *Carbohydr. Lett.*, 1996, **2**, 169-174 (*Casuarine 6-glucoside*)
 Matsumura, T. *et al.*, *Pharm. Biol.*, 2000, **38**, 302-307 (*Uniflorines, isol*)
 Izquierdo, I. *et al.*, *Tetrahedron*, 2005, **61**, 6527-6533 (*synth*)
 Van Ameijde, J.V. *et al.*, *Tetrahedron: Asymmetry*, 2006, **17**, 2702-2712 (*Casuarine, 3-Epicasuarine, isol, synth, pmr, cmr*)
 Ritthiwigrom, T. *et al.*, *Org. Lett.*, 2008, **10**, 2769-2771 (*Uniflorine A, synth, struct*)
 Davies, A.S. *et al.*, *Tetrahedron*, 2008, **64**, 4868-4879 (*Uniflorines, synth, struct*)

3,5,7,15-Tetrahydroxy-6(17),12-lathyradien-14-one T-263



(2β,3β,5α,7β,12E,15β)-form

C₂₀H₃₀O₅ 350.454

(2β,3β,5α,7β,12E,15β)-form

7-(3-Pyridinecarbonyl), 3-benzoyl, 5,15-di-Ac: **L9**
 [129393-28-8]
 C₃₇H₄₁NO₉ 643.732
 Constit. of *Euphorbia lathyris*. Needles. Mp 119-122°. [α]_D²⁰ +130.7 (c, 0.76 in CHCl₃).


3,7-Dibenzoyl, 5-Ac: **Euphorbia factor L₁₁**
 [850560-45-1]
 C₃₆H₄₀O₈ 600.707
 Constit. of *Euphorbia lathyris*. Amorph. powder. Mp 117-118°. [α]_D²⁰ +26 (c, 0.94 in CHCl₃). λ_{max} 230 (log ε 3.51); 274 (log ε 3.13) (MeOH).

(2β,3β,5β,7β,12E,15β)-form

7-Hydroxylathyril
 [34208-98-5]
 Isol. from *Euphorbia lathyris*. Mp 222-224°.

Narayanan, P. *et al.*, *Tet. Lett.*, 1971, 1325-1328 (7-Hydroxylathyril)
 Itokawa, H. *et al.*, *Phytochemistry*, 1990, **29**, 2025-2026 (**L9**)
 Liao, S.-G. *et al.*, *Org. Lett.*, 2005, **7**, 1379-1382 (*Euphorbia factor L₁₁*)

2,3,4,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid T-264

C₁₅H₉NO₈ 331.238

3,4-Methylene, 8-Me ether: 4-Hydroxy-8-methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid. 2-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, 9CI. **Aristolochic acid VIa**
 [108766-27-4]
 C₁₇H₁₁NO₈ 357.276
 Minor alkaloid from the underground parts of *Aristolochia argentina* (Aris-

tolochiaceae). Mp 280-282°.

Priestap, H.A. *et al.*, *Phytochemistry*, 1987, **26**, 519 (*isol, uv, pmr, ms, struct*)

3,4,6,7-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid T-265

C₁₅H₉NO₈ 331.238

3,4-Methylene, 7-Me ether: 6-Hydroxy-7-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. 10-Hydroxy-9-methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. **Aristolochic acid Va**
 [108779-46-0]
 C₁₇H₁₁NO₈ 357.276

Minor alkaloid from the underground parts of *Aristolochia argentina* (Aristolochiaceae). Not obt. completely pure.

3,4-Methylene, 6,7-di-Me ether: 6,7-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. **Aristolochic acid V**
 [108766-28-5]
 C₁₈H₁₃NO₈ 371.303

Minor alkaloid from *Aristolochia argentina* (Aristolochiaceae).

3,4-Methylene, 6,7-di-Me ether, Me ester: [108766-30-9]
 Mp 290-300°.

[108779-47-1, 108766-32-1]

Priestap, H.A. *et al.*, *Phytochemistry*, 1987, **26**, 519 (*isol, uv, ir, pmr, ms, struct*)

3,4,6,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid T-266

C₁₅H₉NO₈ 331.238

3,4-Methylene, 8-Me ether: 10-Hydroxy-8-methoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid. 6-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. **Aristolochic acid D**. **Aristolochic acid IVa**
 [17413-38-6]
 C₁₇H₁₁NO₈ 357.276

Alkaloid from *Aristolochia acuminata*, *Aristolochia argentina*, *Aristolochia clematitis*, *Aristolochia esperanzae*, *Aristolochia indica*, *Aristolochia manshuriensis*, *Aristolochia multiflora* and *Aristolochia mollissima* (Aristolochiaceae). Wine-red cryst. (MeOH). Mp 254-259° Mp 269-271°. λ_{max} 220 (ε 29800); 242 (ε 37790); 252 (ε 37800); 292 (ε 13850); 325 (ε 11300) (EtOH) (Berdy).

3,4-Methylene, 8-Me ether, Me ester: Cryst. (EtOAc). Mp 243-250° dec.

3,4-Methylene, 8-Me ether, 6-O-β-D-glucopyranoside: **Aristoloside**. **Aristolochin†**
 [84014-70-0]
 C₂₃H₂₁NO₁₃ 519.418

Alkaloid from the stems of *Aristolochia manshuriensis* (Aristolochiaceae). Orange prisms (MeOH). Sol. MeOH, butanol; poorly sol. CHCl₃, hexane. Mp 193-196°. [α]_D¹⁵ -69.5 (c, 0.23 in MeOH). λ_{max} 222 (ε 25700); 243 (ε 32300); 252 (ε 33200); 318 (ε 10700); 392 (ε 8500) (MeOH) (Berdy).

3,4-Methylene, 8-Me ether, 6-O-β-D-glu-

copyranoside, Me ester:

Orange cryst. Mp 176-178°. [α]_D¹⁵ -59.4 (c, 0.32 in MeOH).

3,4-Methylene, 6,8-di-Me ether: 8,10-Dimethoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid. 6,8-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. **Aristolochic acid IV**
 [15918-62-4]
 C₁₈H₁₃NO₈ 371.303

Alkaloid from the roots of *Aristolochia argentina*, *Aristolochia esperanzae*, *Aristolochia clematitis*, *Aristolochia longa* and *Aristolochia pallida* and from the stems of *Aristolochia manshuriensis* (Aristolochiaceae). Deep-red prisms (DMF aq.). Mp 268-270° dec.

3,4-Methylene, 6,8-di-Me ether, Me ester: **Aristolochic acid IV methyl ester**
 [17448-02-1]

C₁₉H₁₅NO₈ 385.329
 Alkaloid from bulbs and roots of *Aristolochia kwangsiensis* and tubers of *Aristolochia championii* (Aristolochiaceae). Orange-brown cryst. (CHCl₃), orange rods (EtOAc). Mp 241-243° (238-240°).

Kupchan, S.M. *et al.*, *J.O.C.*, 1968, **33**, 3735-3738 (*Aristolochic acid D*)

Rüveda, E.A. *et al.*, *Monatsh. Chem.*, 1968, **99**, 2349-2358 (*Aristolochic acid IVa*)

Pakrashi, S.C. *et al.*, *Phytochemistry*, 1977, **16**, 1103-1104 (*Aristolochic acid D*)

Chou, F.H. *et al.*, *Yaoxue Tongbao*, 1981, **16**, 56; *CA*, **95**, 175618w (*Aristolochic acid IV Me ester*)

Nakanishi, T. *et al.*, *Phytochemistry*, 1982, **21**, 1759-1762 (*Aristoloside*)

De Pascual Teresa, J. *et al.*, *Phytochemistry*, 1983, **22**, 2745-2747 (*Aristolochic acid IV*)

3,4,7,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid T-267

C₁₅H₉NO₈ 331.238

3,4-Methylene, 7-Me ether: 8-Hydroxy-7-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. **Aristolochic acid E**
 [107259-48-3]
 C₁₇H₁₁NO₈ 357.276

Alkaloid from the roots of *Aristolochia contorta* (Aristolochiaceae).

3,4-Methylene, 8-Me ether: 7-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. 7-Hydroxyaristolochic acid A. **Aristolochic acid VIIa**
 [79185-75-4]
 C₁₇H₁₁NO₈ 357.276

Alkaloid from *Aristolochia debilis*. Also an active principle of Tong Cheng Hu Geng (roots of *Aristolochia tagala*) (Aristolochiaceae). Orange powder (CHCl₃/MeOH). Mp 267-269°. λ_{max} 226 ; 265 ; 313 ; 371 (MeOH).

3,4-Methylene, 7,8-di-Me ether: 7,8-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid. 8,9-Dimethoxy-6-nitrophenanthro[3,4-d]-1,3-dioxole-5-carboxylic acid, 9CI. 7-Methoxyaristolochic acid A. **Aristolochic acid VII**
 [79185-74-3]

C₁₈H₁₃NO₈ 371.303

Alkaloid from *Aristolochia argentina*, *Aristolochia contorta* and *Aristolochia debilis* (Aristolochiaceae). Yellowish needles (as Na salt). Mp >280° (Na salt). λ_{max} 222 ; 264 ; 310 ; 367 ; 372 (MeOH) (Na salt).

3,4-Methylene, 7,8-di-Me ether, Me ester: **Aristolochic acid VII methyl ester** [108766-31-0]

C₁₉H₁₅NO₈ 385.329

Alkaloid from *Aristolochia cucurbitifolia*. Yellow syrup. λ_{max} 227 ; 273 ; 313 ; 373 (MeOH).

Kunimoto, J.-I. et al., *Phytochemistry*, 1980, **19**, 2735 (*Aristolochic acid VII*)

Lou, F. et al., *Yaouxue Xuebao*, 1986, **21**, 702; *CA*, **106**, 135229j (*Aristolochic acid E*)

Priestap, H.A. et al., *Phytochemistry*, 1987, **26**, 519-529 (*Aristolochic acid VII*)


Leu, Y.-L. et al., *Phytochemistry*, 1998, **48**, 743-745 (*Aristolochic acid VIIa*)

Wu, T.S. et al., *Chem. Pharm. Bull.*, 1999, **47**, 571-573; 2000, **48**, 357-361 (*Aristolochic acid VII, Aristolochic acid VII Me ester*)

Nascimento, I.R. et al., *Phytochemistry*, 2003, **63**, 953-957 (*Aristolochic acid VIIa*)

3,8,12,14-Tetrahydroxy-pregn-5-en-20-one

T-268



(3β,12β,14β,17α)-form

C₂₁H₃₂O₅ 364.481

C-17 configs. in some of these derivs. may not be reliable owing to ready epimerisation.

(3β,12β,14β,17α)-form

Lineolone. Desacylcynanchogenin. Deacylcynanchogenin [6869-50-7]

Constit. of *Adonis amurensis* and *Cynanchum caudatum*. Cryst. (Me₂CO). Mp 238° (233-239°). [α]_D²⁴ +130 (c, 0.9 in MeOH). Equilibrates with isolineolone below under mild conditions.

12-(3-Pyridinecarbonyl), 3-O-[α-L-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: C₄₈H₇₁NO₁₅ 902.087

Constit. of *Cynanchum caudatum*. Amorph. powder.

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: C₄₈H₇₁NO₁₅ 902.087

Constit. of *Cynanchum caudatum*. Amorph. powder. [α]_D²⁰ -20.7 (c, 1.6 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[α-L-cymaropyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: C₅₅H₈₃NO₁₈ 1046.257

Constit. of *Cynanchum caudatum*. Amorph. powder. [α]_D²⁰ -51.8 (c, 0.9 in

MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-cymaropyranoside]: [267422-79-7]

C₅₃H₇₉NO₁₈ 1018.203

Constit. of *Asclepias incarnata*. Amorph. powder. [α]_D²¹ -14.4 (c, 0.33 in MeOH). λ_{max} 219 (log ε 4); 263 (log ε 3.49) (MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-digitoxopyranoside]: [267422-66-2]

C₅₂H₇₇NO₁₈ 1004.176

Constit. of *Asclepias incarnata*. Amorph. powder. [α]_D²⁴ -19.1 (c, 0.57 in MeOH). λ_{max} 219 (log ε 4); 258 (log ε 3.46); 263 (log ε 3.49); 269 (sh) (MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-oleandropyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranoside]: C₅₅H₈₃NO₁₈ 1046.257

Constit. of *Cynanchum caudatum*. Amorph. powder. [α]_D²⁰ -26.9 (c, 0.96 in MeOH).

12-(3-Pyridinecarbonyl), 3-O-[β-D-glucopyranosyl-(1→4)-β-D-oleandropyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-digitoxopyranosyl-(1→4)-β-D-cymaropyranoside]: [267422-86-6]

C₅₉H₈₉NO₂₃ 1180.345


Constit. of *Asclepias incarnata*. Amorph. powder. [α]_D²⁷ -12.1 (c, 0.45 in MeOH). λ_{max} 219 (log ε 4.07); 259 (log ε 3.45); 263 (log ε 3.49); 270 (sh) (MeOH).

Warashina, T. et al., *Chem. Pharm. Bull.*, 1994, **42**, 322; 1995, **43**, 977; 1734; 1996, **44**, 358; 2000, **48**, 99-107; 2003, **51**, 1036-1045 (*Cynanchum caudatum* saponins, *Asclepias saponins*, *Araujia sericifera* saponins)

Warashina, T. et al., *Phytochemistry*, 1994, **37**, 217; 1997, **44**, 917; 2000, **53**, 485-488 (*Cynanchum caudatum* saponins, *Asclepias saponins*)

3,11,12,14-Tetrahydroxy-pregn-5-en-20-one

T-269



(3β,11α,12β,14β)-form

C₂₁H₃₂O₅ 364.481

(3β,11ξ,12ξ,14β)-form

11-(3-Pyridinecarbonyl), 12-Ac: **Tinctoramine** [151756-68-2]

C₂₉H₃₇NO₇ 511.614


Alkaloid from *Marsdenia tinctoria*. Needles. Mp 246-255°.

Chowdhury, A.K.A. et al., *Pharmazie*, 1993, **48**, 628-629 (*Tinctoramine*)

1,3,4,5-Tetrahydroxy-2-prenylacridone

T-270

1,3,4,5-Tetrahydroxy-2-(3-methyl-2-butenyl)-9(10H)-acridinone, 9CI

C₁₈H₁₇NO₅ 327.336

4-Me ether, N-Me: 1,3,5-Trihydroxy-4-methoxy-10-methyl-2-prenylacridone.

Glycoctrine IV

C₂₀H₂₁NO₅ 355.39


Alkaloid from *Glycosmis citrifolia*. Yellow oil. λ_{max} 202 (log ε 4.44); 227 (sh) (log ε 4.21); 270 (log ε 4.5); 287 (sh) (log ε 4.36); 315 (log ε 4.09); 336 (sh) (log ε 4); 412 (log ε 3.63) (MeOH).

Ito, C. et al., *Chem. Pharm. Bull.*, 2000, **48**, 65-70

1,3,5,6-Tetrahydroxy-2-prenylacridone

T-271

1,3,5,6-Tetrahydroxy-2-(3-methyl-2-butenyl)-9H-acridin-9-one

C₁₈H₁₇NO₅ 327.336

5-Me ether, N-Me: 1,3,6-Trihydroxy-5-methoxy-10-methyl-2-prenylacridone.

Buntanine

[119116-85-7]

C₂₀H₂₁NO₅ 355.39


Alkaloid from the root bark of *Citrus grandis* (pummelo) (Rutaceae). Yellow plates (Me₂CO). Mp 247-249°.

Wu, T.-S. et al., *Phytochemistry*, 1988, **27**, 3717 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

3,4,7,8-Tetrahydroxy-2-quinolinecarboxylic acid

T-272

1,4-Dihydro-3,7,8-trihydroxy-4-oxo-2-quinolinecarboxylic acid

C₁₀H₇NO₆ 237.168

NH-form


O³,O⁸,N-Tri-Me, Me ester: **Megistonine II**

C₁₄H₁₅NO₆ 293.276

Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow powder. λ_{max} 241 (log ε 3.99); 268 (log ε 3.85); 337 (log ε 3.53) (MeOH).

Fokialakis, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 2009-2010

2,5,9,10-Tetrahydroxy-4(20),11-taxadien-13-one T-273
[5308-90-7]



$C_{20}H_{30}O_5$ 350.454

(2 α ,5 α ,9 α ,10 β)-form
Taxicin II

5-(3-Dimethylamino-3-phenylpropanoyl), 2,9-di-Ac: [313696-96-7]
 $C_{35}H_{47}NO_8$ 609.758
Constit. of *Taxus cuspidata*. Gum. $[\alpha]_D^{24}$ -10 (c, 0.01 in $CHCl_3$).

5-(3-Dimethylamino-3-phenylpropanoyl), 2,10-di-Ac: [313696-95-6]
 $C_{35}H_{47}NO_8$ 609.758
Constit. of *Taxus cuspidata*. Gum. $[\alpha]_D^{24}$ -19 (c, 0.01 in $CHCl_3$).

5-(3-Dimethylamino-3-phenylpropanoyl), 9,10-di-Ac: **2-O-Acetyltaxine II** [313696-97-8]
 $C_{35}H_{47}NO_8$ 609.758
Constit. of *Taxus cuspidata*. Gum. $[\alpha]_D^{24}$ -13 (c, 0.01 in $CHCl_3$).

5-(3-Dimethylamino-3-phenylpropanoyl), 2,9,10-tri-Ac: **Taxine II. 1-Deoxydiacetyltaxine B** [110042-00-7]
 $C_{37}H_{49}NO_9$ 651.795
Constit. of *Taxus baccata*. Gum. $[\alpha]_D^{25}$ +27 (c, 2.8 in CH_2Cl_2).


5-(3S-Dimethylamino-2R-hydroxy-3-phenylpropanoyl), 2,9,10-tri-Ac: **2'-Hydroxytaxine II** [189956-94-3]
[184291-02-9]
 $C_{37}H_{49}NO_{10}$ 667.795
Constit. of *Taxus cuspidata*. Amorph. powder. Mp 84-88°. $[\alpha]_D^{20}$ +43 (c, 1 in $CHCl_3$).

Appendino, G. *et al.*, *J. Nat. Prod.*, 1993, **56**, 514-520 (*Taxine II*)

Ando, M. *et al.*, *J. Nat. Prod.*, 1997, **60**, 499-501 (*2'-Hydroxytaxine II*)

Shi, Q.-W. *et al.*, *Nat. Prod. Lett.*, 2000, **14**, 265-272 (*Taxus cuspidata* constituents)

5,9,12,13-Tetrahydroxy-1,6-tigliadien-3-one T-274



$C_{20}H_{28}O_5$ 348.438

(4 β ,5 β ,9 α ,12 β ,13 α)-form

12-O-(2-Methylaminobenzoyl), 13-Ac:

Sapintoxin C


[81345-28-0]

$C_{30}H_{37}NO_7$ 523.625

Constit. of *Sapium indicum*.

Taylor, S.E. *et al.*, *Phytochemistry*, 1981, **20**, 2749-2751; 1982, **21**, 405-407 (*Sapintoxin C*, *Sapintoxin C*)

9,12,13,20-Tetrahydroxy-1,6-tigliadien-3-one T-275



$C_{20}H_{28}O_5$ 348.438

(4 α ,9 α ,12 β ,13 α)-form [37415-57-9]

12-O-(2-Methylaminobenzoyl): α -Sapintoxin

[80373-85-9]

$C_{28}H_{35}NO_6$ 481.588

Not a natural product.

12-O-(2-Methylaminobenzoyl), 13-Ac:

4 α -Sapinine

[64284-90-8]

$C_{30}H_{37}NO_7$ 523.625

Constit. of *Sapium indicum*. Mp 274-275°. λ_{max} 193 (ϵ 25000); 223 (ϵ 37500); 251 (ϵ 12500); 355 (ϵ 7600) (MeOH).

(4 β ,9 α ,12 β ,13 α)-form

4-Deoxyphorbol

[79083-67-3]

Exists as esters in *Euphorbia tirucalli*, *Euphorbia biglandulosa* and *Sapium indicum*.

12-O-(2-Methylaminobenzoyl), 13-Ac:

Sapintoxin A

[79083-69-5]

$C_{30}H_{37}NO_7$ 523.625

Isol. from unripe fruits of *Sapium indicum*.

► Irritant.

20-Aldehyde, 12-O-(2-methylaminobenzoyl), 13-Ac: [80321-12-6]

$C_{30}H_{35}NO_7$ 521.609

Constit. of *Sapium indicum*.

Fürstenberger, G. *et al.*, *Tet. Lett.*, 1977, **18**, 925-928 (*isol*)

Miana, G.A. *et al.*, *Z. Naturforsch., B*, 1977, **32**, 727 (*4 α -Sapinine*)


Taylor, S.E. *et al.*, *Experientia*, 1981, **37**, 681 (*Sapintoxin A*)

Taylor, S.E. *et al.*, *Phytochemistry*, 1982, **21**, 405-407; 1983, **22**, 1231-1233 (*Sapium esters*)

5,6,7,8-Tetramethoxyisoquinoline T-276

Isonorweberine

[93529-98-7]



$C_{13}H_{15}NO_4$ 249.266

The trivial name Isonorweberine is misleading. Alkaloid detected in *Pachycereus weberi* (Cactaceae) by tandem mass spectrom.

3,4-Dihydro-3,4-Dihydro-5,6,7,8-tetramethoxyisoquinoline. Dehydronorweberine

$C_{13}H_{17}NO_4$ 251.282

Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae). Not indexed in CAS.

1,2,3,4-Tetrahydro-1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxyisoquinoline. Norweberine

$C_{13}H_{19}NO_4$ 253.297

Alkaloid detected in *Pachycereus weberi* (Cactaceae) by tandem ms.

1,2,3,4-Tetrahydro, N-Me-1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-2-methylisoquinoline. Weberine

[74046-24-5]

$C_{14}H_{21}NO_4$ 267.324

Alkaloid from *Pachycereus weberi*. Also identified in trace amount in *Pachycereus pringlei* (Cactaceae). Cryst. (EtOAc)(as hydrochloride). Mp 165-166° (hydrochloride).

Mata, R. *et al.*, *Phytochemistry*, 1980, **19**, 673 (*Weberine, isol, uv, ir, pmr, ms, struct*)


Mata, R. *et al.*, *Planta Med.*, 1980, **38**, 180 (*Norweberine*)

Takahashi, K. *et al.*, *Heterocycles*, 1982, **19**, 691 (*Weberine, synth, ir, pmr, ms*)

Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109-114 (*Norweberine, Isonorweberine, Dehydronorweberine*)

5,8,9,10-Tetramethoxy-4-methylbenzo[g]quinolin-2(1H)-one, 9CI T-277

1-Aza-1,2-dihydro-7,8,9,10-tetramethoxy-4-methyl-2-oxoanthracene [925207-32-5]




$C_{18}H_{19}NO_5$ 329.352

Alkaloid from the bark of *Pseudoxandra cuspidata*. Amorph. yellow powder. $[\alpha]_D^{25}$ +3 (c, 1.7 in $CHCl_3$). λ_{max} 254 (log ϵ 5.13); 287 (log ϵ 5.18); 296 (log ϵ 5.16); 351 (log ϵ 4.25); 362 (log ϵ 4.3) ($CHCl_3$).

Roumy, V. *et al.*, *Planta Med.*, 2006, **72**, 894-898 (*isol, pmr, cmr*)

3,4,5,6-Tetramethoxyphenanthroindolizidine Tyloindicine A T-278



C₂₄H₂₇NO₄ 393.482**(E)-form** [126647-33-4]

Alkaloid from the aerial parts of *Tylophora indica* (Asclepiadaceae). Cryst. (Me₂CO). Mp 199-201°. [α]_D³⁵ +7.2 (c, 2.1 in MeOH).

7-Methoxy: 3,4,5,6,7-Pentamethoxyphenanthroindolizidine
Mp 217-219°.

7-Methoxy, O⁵-de-Me: **Tyloindicine D**
[126624-16-6]


C₂₄H₂₇NO₅ 409.481

Alkaloid from the aerial parts of *Tylophora indica* (Asclepiadaceae). Light yellow powder (Me₂CO). Mp 229-231° dec. [α]_D³⁵ +1.6 (c, 0.6 in MeOH).

Ali, M. et al., *Phytochemistry*, 1989, **28**, 3513-3517 (isol, uv, ir, pmr, ms, struct, Tyloindicine D)

Tetramethoxyscytonemin T-279

[683771-36-0]

C₄₀H₃₄N₂O₈ 670.717

Isol. from the terrestrial cyanobacterium *Scytonema* sp. Amorph. purple solid. λ_{\max} 212 (ε 35930); 562 (ε 5940) (MeOH).

3,9-Didemethoxy, 3,9-didehydro: **Dimethoxyscytonemin**
[683771-37-1]

C₃₈H₂₈N₂O₆ 608.649

Isol. from a terrestrial *Scytonema* sp. Dark red amorph. solid. λ_{\max} 215 (ε 60350); 316 (ε 18140); 422 (ε 23015) (MeOH).

Bultel-Poncé, V. et al., *J. Nat. Prod.*, 2004, **67**, 678-681 (isol, pmr, cmr)

Tetramethylammonium(1+), T-280 8CI

N,N,N-Trimethylmethanaminium(1+), 9CI

[51-92-3]

Me₄N⁺C₄H₁₂N⁺ 74.145

Occurs naturally in plant and animal tissues.

Chloride: **Tetramethylammonium chloride**
[75-57-0]

C₄H₁₂ClN 109.598

Cryst. (EtOH). Mp 230° dec.

▶ BS7700000

[15625-56-6, 80526-82-5, 79723-02-7]

Llinares, J. et al., *Org. Magn. Reson.*, 1980, **14**, 20 (cmr)

Neef, C. et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1984, **328**, 103; 111 (pharmacol)

Tsubaki, H. et al., *J. Pharmacobio-Dyn.*, 1986, **9**, 737 (pharmacol)

N,N,N',N'-Tetramethyl-1,4-butanediamine, 9CI T-281

1,4-Bis(dimethylamino)butane. **Tetramethylputrescine**

[111-51-3]

Me₂N(CH₂)₄NMe₂C₈H₂₀N₂ 144.259

Constit. of *Duboisia leichhardtii*, *Duboisia myoporoides*, *Hyoscyamus muticus*, *Hyoscyamus reticulatus*, *Hyoscyamus niger*, *Oldenlandia affinis* and *Ruellia rosea* (Solanaceae, Rubiaceae, Acanthaceae). Oil. Bp 169°.

▶ EJ7530000

Hydrochloride (1:2): Mp 280° (273°).

Dipicrate: Mp 197-200°.

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 498A (nmr)

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 307A (ir)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 394D (ir)

Willstätter, R. et al., *Ber.*, 1907, **40**, 3869 (struct)

Konowalowa, R.A. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1928, **266**, 449; *CA*, **23**, 472 (isol)

Griffin, W.J. et al., *Aust. J. Pharm.*, 1967, **48**, S20; *CA*, **67**, 79622u (isol)

Coulson, J.F. et al., *Planta Med.*, 1968, **16**, 174 (isol)


Veith, H.J. et al., *Helv. Chim. Acta*, 1971, **54**, 653 (ms)

Gran, L. et al., *J. Nat. Prod.*, 1973, **36**, 209 (isol, ms)

Johns, S. et al., *Phytochemistry*, 1975, **14**, 2635 (isol, uv, ir, cmr, ms)

2,2',4,4'-Tetramethyl-6,6'-di-nitroazobenzene T-282

Bis(2,4-dimethyl-6-nitrophenyl)diazene, 9CI

C₁₆H₁₆N₄O₄ 328.327**(E)-form** [811848-77-8]

Alkaloid from the leaves of *Aconitum sunpaganense*. Red cryst. Subl. 210.


Wang, X. et al., *Fitoterapia*, 2004, **75**, 789-791 (isol, pmr, cmr, cryst struct)

Wang, X.-L. et al., *Jieoug Huaxue*, 2004, **23**, 1005-1008 (cryst struct)

2,2,6,6-Tetramethyl-4-piperidinone, 9CI T-283

Triacetoneamine. Odoratin†. *Vincubine*
[826-36-8]

[956833-72-0 (nitrate salt)]

C₉H₁₇NO 155.239

Isol. from *Acalypha indica* and *Salsola tetrandra* (Euphorbiaceae, Chenopodiaceae), also from *Viola odorata* (sweet violet). Metab. of the soft coral *Lobophytum strictum* and sponge *Agelas ooides*. Plates + 1H₂O (Et₂O); anhyd. needles (dry Et₂O). Mp 34.9° (anhyd.). Bp 205°. The hydrochloride of the alkaloid from *Salsola tetrandra* was reported to have [α]_D³⁵ +14.3° (EtOH) which makes the struct. assignment highly dubious. Artifact.

▶ TO0127900

Monohydrate: [10581-38-1]

Mp 61-63° (58°). Bp₁₈ 102-105°.

Hydrochloride: [33973-59-0]

Mp 198° dec.

Oxime: [4168-79-0]

C₉H₁₈N₂O 170.254

Prisms (EtOH). Mp 153°.

▶ LD₅₀ (mus, ivn) 180 mg/kg. TO0128000

Semicarbazone:

Cryst. (EtOH). Mp 219-220°.

N,N-Di-Me: 1,1,2,2,6,6-Hexamethyl-4-oxopiperidinium(1+), 9CI. 1,1,2,2,6,6-Hexamethyl-4-piperidinone
[195864-37-0, 113308-50-2]

C₁₁H₂₂NO⁺ 184.301

Quaternary alkaloid from the alga *Laminaria japonica*.

N-Benzyl: [52981-86-9]

C₁₆H₂₃NO 245.364

Plates or prisms (EtOH/Et₂O) (as hydrochloride). Mp 137-138° (hydrochloride).

N-Bromo:

C₉H₁₆BrNO 234.135

Cryst. (petrol). Mp 44°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 461C; 461D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 720C; 721A (nmr)

Frenclowa, I. et al., *Acta Pol. Pharm.*, 1961, **18**, 187; 1964, **21**, 145; *CA*, **57**, 7381; **62**, 10817 (isol)

Rees, B. et al., *Acta Cryst. B*, 1971, **27**, 932 (cryst struct)

Karawya, M.S. et al., *Phytochemistry*, 1971, **10**, 3303 (isol)

Sosnovsky, G. et al., *Synthesis*, 1976, 735 (synth)

Ermakov, A.I. et al., *Khim. Geterotsikl. Soedin.*, 1979, 507 (ms)

Ozinskas, A.J. et al., *Helv. Chim. Acta*, 1980, **63**, 1407 (pmr)

Parameswaran, P.S. et al., *Indian J. Chem., Sect. B*, 1991, **30**, 449 (isol, pmr, cmr)

Shimoi, N. et al., *Fish. Sci.*, 1997, **63**, 650-651 (*Hexamethylpiperidone*)

Koenig, G.M. et al., *Planta Med.*, 1998, **64**, 88-89 (isol, pmr, cmr)

Bond, A.D. et al., *Acta Cryst. C*, 2002, **58**,

o115-o116 (*monohydrate, cryst struct*)

Rodrigues, V.H. et al., *Acta Cryst. C*, 2007, **63**,


o243-o245 (*nitrate, cryst struct*)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van

Nostrand Reinhold, 1992, TDU000

Tetramethylpyrazine**T-284**

Chuanxiongzine. Ligustrazine. Tetrapyrazine. BS Factor. FEMA 3237. NSC 36080. NSC 46451
[1124-11-4]



$C_8H_{12}N_2$ 136.196

Metab. from *Bacillus subtilis* and an arctic marine bacterium; isol. from Galbanum oil. Constit. of *Ligusticum chuanxiong* and *Ligusticum wallichii*. Fragrance and flavouring ingredient. Shows antiinflammatory activity. Bactericidal agent. Trihydrate; cryst. (H_2O). Mp $74-77^\circ$ (hyd.) Mp 86° (anhyd.). Bp 190° . pK_{a1} 3.61; pK_{a2} -2.73 (25° , H_2O). pK_{a1} 3.55; pK_{a2} -2.7 (27° , H_2SO_4 aq.). Log P 1.58 (calc). Odour threshold 10^4 ppb in H_2O . λ_{max} 294 (ϵ 8500) (MeOH) (Berdy).

► LD₅₀ (rat, orl) 1910 mg/kg. UQ3905000
Hydrochloride: Mp 91° (dihydrate) Mp 156° (anhyd.).

Dipicrate:

Needles. Mp $194-195^\circ$.

1,4-Dioxide:

$C_8H_{12}N_2O_2$ 168.195

Mp 224° .

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 841C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 402A (nmr)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, 3, 1559A (ir)

Piloty, O. *et al.*, *Ber.*, 1910, 43, 489

Kipping, F.B. *et al.*, *J.C.S.*, 1929, 2891

Gnitchel, H. *et al.*, *Chem. Ber.*, 1972, 105, 1865

Bus, J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1973, 92, 123 (ir)

Gumbley, S.J. *et al.*, *J. Het. Chem.*, 1985, 22, 1143 (props)

Eiermann, U. *et al.*, *Chem. Ber.*, 1990, 123, 523 (uv)


Ozaki, Y. *et al.*, *Chem. Pharm. Bull.*, 1992, 40, 954 (isol, pharmacol)

Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, 2, 318-353 (marine isol)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, TDV725

2,4,6,6-Tetramethyl-3(6H)-pyridinone**T-285**

[203524-64-5]




$C_9H_{13}NO$ 151.208

Isol. from the marine sponge *Agelas oroides*. Volatile oil/glass. Possible artifact. λ_{max} 221 (sh) (ϵ 9310); 245 (ϵ 7120) (MeOH).

Koenig, G.M. *et al.*, *Planta Med.*, 1998, 64, 88-89 (isol, uv, ir, pmr, cmr, ms)

Tetrandrine**T-286**

6,6',7,12-Tetramethoxy-2,2'-dimethylberbaman, 9CI. Menisine. NSC 77037
[518-34-3]



Absolute Configuration

$C_{38}H_{42}N_2O_6$ 622.76

Alkaloids covered by this entry (*S,S*-series) are enantiomeric with those covered by Phaeanthine, P-301 and diastereomeric with those included in Isotetrandrine, I-328 (1*R*, 1'*S*-) and Peinamine, P-171 (1*S*, 1'*R*-). Menisine, Mp 152° , was transformed into Tetrandrine on heating at 150° and was almost certainly a metastable modification of it. Obt. semisynthetically by methylation of Penduline. Alkaloid from the roots of *Cocculus sarmentosus*, *Cyclea barbata*, *Cyclea peltata*, *Cyclea burmanni*, *Stephania hernandifolia*, *Trichlisia subcordata* and *Isopyrum thalictroides*, (Menispermaceae). Menisine obt. from the Chinese drug Mu fang chi (tubers of *Stephania tetrandra*). Shows hypotensive and vasodilator props., gives exp. protection against myocardial infarction. Shows antineoplastic activity. Most effective antineoplastic agent of the bisbenzylisoquinoline alkaloids. Also shows antiinflammatory, antidiabetic, anaesthetic, antiarrhythmic, antimicrobial and antimotile props. Needles (Et_2O). Mp 218° ($215-217^\circ$). $[\alpha]_D^{24}$ +274 (c, 2.16 in $CHCl_3$) (+241.4). Log P 8.3 (calc). λ_{max} 282 (ϵ 8050) (EtOH).

► LD₅₀ (mus, ipr) 41.3 mg/kg. Chronic administration causes liver necrosis. Lymphotoxin.

N²-β-Oxide: Fenfangjine A. Tetrandrine 2β-N-oxide

[115556-32-6]

$C_{38}H_{42}N_2O_7$ 638.759

Alkaloid from "Fen-Fang-Ji" (roots of *Stephania tetrandra*) (Menispermaceae). Needles (Me_2CO). Mp $174-176^\circ$. $[\alpha]_D^{24}$ +328.2 (c, 1.043 in $CHCl_3$).

N²-β-Oxide: Tetrandrine 2'β-N-oxide

[62828-25-5]

$C_{38}H_{42}N_2O_7$ 638.759

Alkaloid from the Thai drug "Krung Kha Mao" (*Cyclea barbata*) (Menispermaceae). Mp $185-190^\circ$. $[\alpha]_D^{23}$ +198 ($CHCl_3$). $[\alpha]_D$ +157 (c, 0.24 in $CHCl_3$).

N²-De-Me: 2-Nortetrandrine

[19196-54-4]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the seeds of *Ocotea rodiaei* (Lauraceae). Rods (Me_2CO). Mp $222-226^\circ$. $[\alpha]_D^{24}$ +335 (c, 1.0 in $CHCl_3$). λ_{max} 227 (log ϵ 4.53); 282 (log ϵ 3.88) ($CHCl_3$).

N²-De-Me: perchlorate (1:2):

Cryst. + 4 H_2O . Mp $252-253^\circ$.

N²-De-Me, dipicrate: Mp $247-251^\circ$.

N²-De-Me: Cycleanorine. 2'-N-De-methyltetrandrine

[38769-08-3]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the roots of *Cyclea peltata* (Menispermaceae). Antineoplastic agent. Needles ($EtOH$ aq.). Mp $170-172^\circ$. $[\alpha]_D^{25}$ +308 (c, 0.52 in $CHCl_3$). Log P 7.6 (uncertain value) (calc). λ_{max} 252 (ϵ 10200) (EtOH).

N²-Me: 2-N-Methyltetrandrine. Cycleahomine

[41222-80-4]

[38849-82-0 (chloride salt)]

$C_{39}H_{45}N_2O_6^{\oplus}$ 637.794

Quaternary alkaloid from roots of *Stephania tetrandra* the source of the Chinese drug Fen-Fang-Ji. Also from *Cyclea peltata* roots. Needles (Me_2CO) or prisms ($CH_2Cl_2/EtOAc$) (as chloride). Mp $192-196^\circ$ (chloride). $[\alpha]_D$ +245 (c, 0.07 in $CHCl_3$) (chloride)(103). λ_{max} 280 (log ϵ 3.84) (EtOH) (as chloride).

N²-Me: 2'-N-Methyltetrandrine

[90308-88-6]

$C_{39}H_{45}N_2O_6^{\oplus}$ 637.794

Isol. from roots of *Stephania tetrandra*, the source of the Chinese drug Fen-Fang-Ji. Amorph. powder. Mp $208-211^\circ$ (chloride) (synthetic). $[\alpha]_D^{25}$ +166.5 (c, 0.2 in $CHCl_3$) (synthetic).

N-Me: Monomethyltetrandrinium

[53935-72-1 (chloride-hydrochloride)]

$C_{39}H_{45}N_2O_6^{\oplus}$ 637.794

Alkaloid from *Cyclea barbata* (Menispermaceae). Plates ($CHCl_3$ /petrol) (as chloride-hydrochloride). Mp 208° (chloride-hydrochloride). $[\alpha]_D^{20}$ +51.5 (MeOH) (as chloride-hydrochloride). Site of monomethylation unknown.

N²-Chloromethyl: N²-Chloromethyltetrandrinium

[38769-09-4 (chloride)]

$C_{39}H_{44}ClN_2O_6^{\oplus}$ 672.239

Artifact isol. from the roots of *Cyclea peltata* (Menispermaceae). Needles + $3\frac{1}{2}H_2O$ (Me_2CO) (as chloride). Mp $213-217^\circ$ dec. (chloride). $[\alpha]_D^{25}$ +156 (c, 0.41 in $CHCl_3$) (chloride).

N²,N²-Di-Me: N,N'-Dimethyltetrandrinium. Hanjisong

[26605-37-8]

[22445-73-4, 116171-98-3]

$C_{40}H_{48}N_2O_6^{\oplus}$ 652.829

Alkaloid from *Stephania tetrandra*. Needles (EtOH) (as diiodide). Mp 267.5° ($236-240^\circ$ dec.) (diiodide). $[\alpha]_D^{20}$ +234 (MeOH) (diiodide).

N²,N²-Bis(chloromethyl): N²,N²-Bis(chloromethyl)tetrandrinium

[136055-61-3 (dichloride salt)]

$C_{40}H_{46}Cl_2N_2O_6^{\oplus}$ 721.719

Artifact isol. from *Stephania tetrandra* (Menispermaceae).

O⁷-De-Me: Fangchinoline. Menisidine. 12-O-Methylatherospermoline

[436-77-1]

$C_{37}H_{40}N_2O_6$ 608.733

Alkaloid from the roots of *Cyclea peltata*, *Trichlisia subcordata* and *Stephania hernandifolia*, the tubers of *Stephania tetrandra*, and from the bark of *Daphnandra* sp. Dt-7 (Menispermaceae, Atherospermaceae).

Also, as Menisidine, isol. from the Chinese drug "Mu-fang-chi" (*Stephania tetrandra*) (Menispermaceae). Shows good *in vitro* antineoplastic activity against HeLa cells. Also anti-fibrogenic and antiinflammatory. Cryst. (Et₂O). Mp 240-242° dec. $[\alpha]_D^{24} +274$ (c, 0.55 in MeOH). Log P 8.12 (uncertain value) (calc). Menisidine, Mp 176°, was transformed into Fangchinoline on heating at 160-170° and was almost certainly a metastable modification of it. Opt. rotns. were similar.

► LD₅₀ (mus, ipr) 50 mg/kg. DR9280000
O⁷-De-Me, dipicrate: Mp 186° Mp 225°.

O⁷-De-Me, N²- α -oxide: **Fenfangjine B**.
Fangchinoline 2'- α -N-oxide
[115648-96-9]
C₃₇H₄₀N₂O₇ 624.732
Alkaloid from "Fen-Fang-Ji" (roots of *Stephania tetrandra*) (Menispermaceae). Prisms (EtOH). Mp 211-213°. $[\alpha]_D^{25} +242.5$ (c, 0.640 in CHCl₃/MeOH).

O⁷-De-Me, N²- β -oxide: **Fenfangjine C**.
Fangchinoline 2'- β -N-oxide
[115648-97-0]
C₃₇H₄₀N₂O₇ 624.732
Alkaloid from "Fen-Fang-Ji" (roots of *Stephania tetrandra*) (Menispermaceae). Needles (EtOH). Mp 165-166°. $[\alpha]_D^{25} +239.4$ (c, 0.630 in MeOH).

O⁷-De-Me, N²-Me: N²-Methylfangchinoline
[133744-51-1]
C₃₈H₄₃N₂O₆⁺ 623.767
Alkaloid from the roots of *Stephania tetrandra* (Menispermaceae). Needles (Me₂CO/MeOH) (as chloride). Mp 205-208° (chloride). $[\alpha]_D +256$ (c, 0.9 in MeOH). λ_{\max} 281 (log ϵ 3.83) (MeOH).

O¹²-De-Me: **Penduline**†. (+)-Pycnamine
[26137-45-1]
Alkaloid from the stems and leaves of *Cocculus pendulus*, and from the roots of *Cocculus laeabe*, *Berberis coriaria* and *Andrachne cordifolia* (Menispermaceae, Berberidaceae, Euphorbiaceae). Also obt. by epimerisation of Berbamine in I-328 with MeOH (40°/24h, then 2 days at r.t.). Cryst. (EtOAc). Mp 192-194°. $[\alpha]_D +265$ (c, 1.0 in CHCl₃). λ_{\max} 284 (log ϵ 3.84) (EtOH).

O¹²-De-Me, N²-de-Me: **Norpenduline**
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from the stems of *Cocculus pendulus*. $[\alpha]_D^{25} +260$ (c, 0.09 in MeOH).

O⁷,O¹²-Di-de-Me: **Atherospermoline**
[21008-67-3]
C₃₆H₃₈N₂O₆ 594.706
Alkaloid from the leaves of *Atherosperma moschatum* (Monimiaceae). Prisms + 1CHCl₃ (CHCl₃). Mp 183-188°. $[\alpha]_D^{18} +202$ (c, 0.15 in CHCl₃). λ_{\max} 284 (log ϵ 3.97) (EtOH).

(±)-form [23495-89-8]

Alkaloid from the roots of *Cyclea peltata*, *Cyclea barbata*, *Stephania hernandi-*

folia and *Isopyrum thalictroides* (Menispermaceae, Ranunculaceae). Needles (EtOH or CHCl₃/MeOH). Mp 257-258° (252-253°).

O⁷-De-Me: **Cycleadrine**
[38769-07-2]
Alkaloid from the roots of *Cyclea peltata*. Also isol. from the Thai drug Krung Kha Mao (*Cyclea barbata*) (Menispermaceae). Cryst. (Me₂CO/hexane). Mp 160-162°.

O⁷-De-Me, hydroiodide (1:2):
Cryst. (EtOH aq.). Mp 223-224°.
Kidd, D.A.A. *et al.*, *J.C.S.*, 1954, 669-677 (struct)

Tomita, M. *et al.*, *Pharm. Bull.*, 1954, **2**, 372-375 (struct)
Kupchan, S.M. *et al.*, *J. Pharm. Sci.*, 1961, **50**, 164-167; 819-822 (*Menisidine, isol*)
Bick, I.R.C. *et al.*, *J.C.S.*, 1961, 1896-2407 (*pmr*)
Bick, I.R.C. *et al.*, *Chem. Ind. (London)*, 1965, 694-695 (*Atherospermoline*)
Battersby, A.R. *et al.*, *J.C.S.*, 1965, 2239-2247 (*uv, ord*)
Chan, K.C. *et al.*, *J.C.S. (C)*, 1967, 2479-2488 (*ord, pmr, ms, Nortetrandrine*)
Tomita, M. *et al.*, *Yakugaku Zasshi*, 1967, **87**, 316; *CA*, **67**, 32855g (*Fangchinoline, isol*)

Gupta, N.C. *et al.*, *Experientia*, 1970, **26**, 12-13 (*Penduline*)
Baldas, J. *et al.*, *J.C.S. Perkin I*, 1972, 592-596 (*ms*)
Kupchan, S.M. *et al.*, *J.O.C.*, 1973, **38**, 1846-1852 (*Cycleanorine, Cycleahomine, Cycleadrine*)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 1-5 (*isol, uv, ir, pmr, ms*)
Hoffstadt, B. *et al.*, *Tetrahedron*, 1974, **30**, 307-310 (*Monomethyltetrandrinium*)
Dwuma-Badu, D. *et al.*, *Phytochemistry*, 1975, **14**, 2524-2525 (*isol, uv, ir, pmr, ms*)
Koike, L. *et al.*, *Tet. Lett.*, 1979, 3765-3768 (*cmr*)

Bhakuni, D.S. *et al.*, *Phytochemistry*, 1980, **19**, 2347-2350 (*biosynth*)
Moulis, C. *et al.*, *J. Nat. Prod.*, 1981, **44**, 101-103 (*isol, uv, ir, pmr, ms*)
Chung, B.H. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1984, **317**, 274-278 (*N-Me derivs, synth*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 165-171 (*rev, pharmacol*)
Fang, D. *et al.*, *Chin. Med. J. (Beijing, Engl. edn.)*, 1986, **99**, 638-644; *CA*, **106**, 43950j (*pharmacol*)

Ogino, T. *et al.*, *Heterocycles*, 1988, **27**, 1149-1154 (*Fenfangjines A-C, N-Me-Tetrandrines*)
Deng, J. *et al.*, *J. Nat. Prod.*, 1990, **53**, 993 (2-N-Methyltetrandrine, 2-N-Methylfangchinoline)

Deng, J. *et al.*, *Chin. Chem. Lett.*, 1991, **2**, 231-232; *CA*, **115**, 131982n (*N², N²-Bis(chloromethyl)tetrandrinium*)
Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1993, **56**, 1989-1992 (*Tetrandrine 2'- β -N-oxide*)


Wang, G. *et al.*, *Life Sci.*, 1994, **56**, 295-306 (*pharmacol, rev*)
Liu, Q.-Y. *et al.*, *J. Pharmacol. Exp. Ther.*, 1995, **273**, 32 (*pharmacol*)
Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (*pmr, cmr*)
Mousa, S.A. *et al.*, *Cardiovasc. Drug Rev.*, 1998, **16**, 48-61 (*rev*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, TDX830

Tetrapetalone C

[780781-76-2]

T-287

C₂₆H₃₃NO₈ 487.549

Structs. revised from those reported in isol. ref. Prod. by *Streptomyces* sp. USF-4727. Lipoygenase inhibitor. Amorph. powder. Mp 154-157°. λ_{\max} 206 (ϵ 10350); 250 (sh) (ϵ 5890); 311 (ϵ 3190); 350 (sh) (ϵ 2900) (MeOH).

2-Deoxy: **Tetrapetalone A**

[561056-66-4]

C₂₆H₃₃NO₇ 471.549

Prod. by *Streptomyces* sp. USF-4727. Lipoygenase inhibitor. Pale yellow powder. Mp 190°. Exists in 1-oxo-3-hydroxy enol form. λ_{\max} 240 (ϵ 13800); 385 (ϵ 10200) (MeOH).

17R-Acetoxy: **Tetrapetalone D**

[780781-77-3]

C₂₈H₃₅NO₁₀ 545.585

Prod. by *Streptomyces* sp. USF-4727. Lipoygenase inhibitor. Amorph. powder. Mp 148-151°. λ_{\max} 208 (sh) (ϵ 9920); 247 (sh) (ϵ 4470); 312 (ϵ 1720); 335 (sh) (ϵ 1440) (MeOH).

17R-Acetoxy, 2-deoxy: **Tetrapetalone B**

[780781-75-1]

C₂₈H₃₅NO₉ 529.586

Prod. by *Streptomyces* sp. USF-4727. Lipoygenase inhibitor. Pale yellow powder. Mp 191-193°. λ_{\max} 217 (ϵ 9310); 236 (ϵ 4190) (MeOH).


Komoda, T. *et al.*, *Biosci., Biotechnol., Biochem.*, 2004, **68**, 104-111; 903-908 (*Tetrapetalones A-D, isol, pmr, cmr, struct, abs config*)

Komoda, T. *et al.*, *Org. Biomol. Chem.*, 2007, **5**, 1615-1620 (*biosynth*)

Tetraphyllicine

T-288

19,20-Didehydroajmalan-17-ol, 9CI. Serpinine. **Semperflorine**
[509-38-6]

C₂₀H₂₄N₂O 308.422

Alkaloid from *Rauwolfia tetraphylla* and several other *Rauwolfia* spp. (Apocynaceae). Cryst. (Me₂CO). Mp 320-322°. $[\alpha]_D^{28} +21$ (Py). pK_a 8.5. λ_{\max} 250 (log ϵ 4.07); 294 (log ϵ 3.6) (EtOH).

Perchlorate:

Cryst. (EtOH). Mp 305-307° dec.

O-Ac:

Noncryst. Mp 60° (partial melt) Mp 154° (clear melt).

- O-(3,4,5-Trimethoxybenzoyl): **Rauvominitine**
[466-57-9]
C₃₀H₃₄N₂O₅ 502.609
Alkaloid from the roots and unripe fruits of *Rauwolfia vomitoria*, and the root of *Rauwolfia obscura* (Apocynaceae). Needles (EtOH aq.). Mp 115-117°. $[\alpha]_D^{20}$ -173.4 (c, 1 in CHCl₃). λ_{\max} 254 (log ϵ 4.18) (no solvent reported).
- O-(3,4,5-Trimethoxybenzoyl); hydrochloride:
Cryst. (H₂O). Mp 223° dec.
- N-De-Me: **Nortetraphyllicine**
[68160-76-9]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Rauwolfia cumminsii*, *Rauwolfia macrophylla*, *Rauwolfia mombasiana*, *Rauwolfia volkensii* and *Rauwolfia vomitoria* (Apocynaceae). Mp 280° dec. λ_{\max} 210 ; 240 ; 288 (MeOH).
- N-De-Me, O-Ac: **17-O-Acetylnortetraphyllicine**
[81525-52-2]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from the root bark of *Rauwolfia nitida* (Apocynaceae). Amorph. $[\alpha]_D^{22}$ +41 (c, 0.01 in MeOH). λ_{\max} 215 (log ϵ 4.31); 246 (log ϵ 4.5); 292 (log ϵ 3.63) (MeOH).
- N-De-Me, O-(3,4,5-trimethoxybenzoyl): **Norrauvomitine**
[64675-21-4]
C₂₉H₃₂N₂O₅ 488.582
Alkaloid from *Rauwolfia vomitoria* root and stem bark.
- 17-Ketone: **Demethoxypurpeline**
[457613-92-2]
C₂₀H₂₂N₂O 306.407
Alkaloid from the bark of *Rauwolfia bahiensis*. Amorph. yellow solid. $[\alpha]_D^{25}$ +22 (c, 0.57 in CHCl₃). λ_{\max} 246 (log ϵ 3.61); 280 (log ϵ 3.32); 294 (log ϵ 3.31) (MeOH).
- 17-Ketone, N-de-Me: **Rauflorine**
[36063-54-4]
C₁₉H₂₀N₂O 292.38
Alkaloid from the root bark of *Rauwolfia confertiflora* (Apocynaceae). Cryst. (MeOH). Mp 221°. $[\alpha]_D^{20}$ +312 (c, 1.54 in CHCl₃). λ_{\max} 215 (log ϵ 3.84); 240 (log ϵ 3.91); 292 (log ϵ 3.54) (MeOH).
- 10-Hydroxy, N-de-Me: **10-Hydroxynortetraphyllicine**
[70509-79-4]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from the stem bark of *Rauwolfia vomitoria* (Apocynaceae). Amorph. λ_{\max} 215 ; 222 (sh) ; 245 (sh) ; 270 ; 280 ; 290 (MeOH).
- 12-Hydroxy, 17-ketone: **Mitoridine**
[3911-19-1]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from *Rauwolfia vomitoria* and the stem bark of *Rauwolfia cumminsii* (Apocynaceae). Cryst. (MeOH). Mp 322°. $[\alpha]_D^{25}$ +175 (c, 1.04 in Py). pK_a 6.16 (methyl cellosolve). λ_{\max} 215 (log ϵ 4.55); 249 (log ϵ 3.89); 293 (log ϵ 3.42) (EtOH).


- 12-Hydroxy, 17-ketone, N-de-Me: **Nor-mitoridine**
[68160-75-8]
C₁₉H₂₀N₂O₂ 308.379
Alkaloid from the leaves of *Rauwolfia cumminsii* (Apocynaceae). λ_{\max} 249 (log ϵ 4.74); 290 (log ϵ 4.22) (MeOH).
- 10-Methoxy: **Vincamajoreine**
[3382-93-2]
C₂₁H₂₆N₂O₂ 338.449
Alkaloid from *Vinca major* (Apocynaceae). Needles (MeOH). Mp 266-267° (230°, 246-247°). λ_{\max} 246 (log ϵ 4.13); 310 (log ϵ 3.66) (EtOH).
- 10-Methoxy, O-Ac: **Majoridine. Majidine**
[6519-30-8]
C₂₃H₂₈N₂O₃ 380.486
Alkaloid from the aerial parts of *Vinca major* (Apocynaceae). Cryst. (EtOH). Mp 222-223° (210-213°). $[\alpha]_D^{22}$ -26 (c, 1.05 in CHCl₃). pK_a 6.95 (80% 2-methoxyethanol). λ_{\max} 247 (log ϵ 3.94); 310 (log ϵ 3.54) (MeOH).
- 10-Methoxy, 17-ketone, N-de-Me: **Endolobine**
[67627-71-8]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from the leaves and stem bark of *Rauwolfia cumminsii*, and the leaves of *Rauwolfia mombasiana* (Apocynaceae).
- 11-Methoxy, 17-ketone: **Rauflexine**
[70522-05-3]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from the leaves of *Rauwolfia reflexa* (Apocynaceae). Cryst. (petrol/EtOAc). Mp 154-155°. Originally confused with Purpeline.
- 12-Methoxy: **Seredamine**
[3911-20-4]
C₂₁H₂₆N₂O₂ 338.449
Alkaloid from the root and stem bark of *Rauwolfia cumminsii* and from *Rauwolfia vomitoria* (Apocynaceae). Cryst. (CHCl₃/Me₂CO). Mp 297° (280-283°). $[\alpha]_D^{25}$ +44 (c, 1.97 in CHCl₃). $[\alpha]_D^{18}$ +60 (c, 0.74 in CHCl₃/EtOH, 200:1). pK_a 7.79 (methyl cellosolve). λ_{\max} 210 (log ϵ 4.51); 254 (log ϵ 3.93); 290 (log ϵ 3.4) (EtOH).
- 12-Methoxy, O-Ac: Mp 260-262°.
- 12-Methoxy, O-(3,4,5-trimethoxybenzoyl): **3,4,5-Trimethoxybenzoylsередamine**
[68160-77-0]
C₃₁H₃₆N₂O₆ 532.635
Alkaloid from the leaves of *Rauwolfia cumminsii* (Apocynaceae). λ_{\max} 218 (log ϵ 4.62); 251 (log ϵ 4.2); 290 (log ϵ 3.85) (MeOH).
- 12-Methoxy, N-de-Me: **Norseredamine**
[30171-06-3]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from *Rauwolfia vomitoria*, *Rauwolfia sumatrana* and leaves of *Rauwolfia cumminsii*. Needles (CHCl₃). Mp 242-245°. $[\alpha]_D^{22}$ +32.6 (c, 3.65 in MeOH). λ_{\max} 246 (log ϵ 3.72); 287 (log ϵ 3.24) (EtOH).
- 12-Methoxy, N-de-Me, N,O-di-Ac: Cryst. (Et₂O/hexane). Mp 172-175°.

- 12-Methoxy, 17-ketone: **Purpeline**
[2246-33-5]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from the stem bark of *Rauwolfia cumminsii* and *Rauwolfia vomitoria*, and the leaves of *Rauwolfia mombasiana* and *Rauwolfia reflexa* (Apocynaceae). Cryst. (petrol/EtOAc, Et₂O/hexane or Me₂CO/hexane). Mp 155° Mp 164-165°. $[\alpha]_D^{22}$ +286 (c, 1.659 in CHCl₃). λ_{\max} 252 (log ϵ 3.93); 291 (log ϵ 3.42) (EtOH).
- 12-Methoxy, 17-ketone, N-de-Me: **Norpurpeline**
[65061-33-8]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from the stem bark of *Rauwolfia cumminsii*, *Rauwolfia mombasiana* and *Rauwolfia vomitoria*, and the leaves of *Rauwolfia mombasiana* (Apocynaceae). Yellow cryst. $[\alpha]_D$ 0. λ_{\max} 229 (log ϵ 4.5); 254 (log ϵ 3.92); 290 (log ϵ 3.45) (MeOH).
- 17-Epimer: **Mauisene. 17-Epitetraphyllicine**
[6883-73-4]
Alkaloid from the root bark of *Rauwolfia mauiensis* (Apocynaceae). Cryst. (Me₂CO). Mp 240-242° (235-240°). $[\alpha]_D$ +184 (MeOH). λ_{\max} 249 (log ϵ 3.92); 291 (log ϵ 3.35) (EtOH).
- 17-Epimer, hydrochloride: Needles (MeOH). Mp 295° dec.
- 17-Epimer, 11-methoxy: **Reflexine†**
[61091-18-7]
C₂₁H₂₆N₂O₂ 338.449
Alkaloid from the leaves of *Rauwolfia reflexa* (Apocynaceae). Needles (Me₂CO). Mp 260° dec. $[\alpha]_D$ +126 (CHCl₃). OMe group originally considered to be in 12-posn. λ_{\max} 250 (log ϵ 3.78); 295 (log ϵ 3.66) (EtOH).
- 17-Epimer, 12-hydroxy: **12-Hydroxymauisene**
[104748-99-4]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from root bark of *Rauwolfia media* (Apocynaceae). Mp 260°. $[\alpha]_D$ +100 (c, 1 in MeOH). λ_{\max} 255 (ε 7630); 295 (ε 2270) (EtOH).
- 17-Epimer, 12-methoxy, N-de-Me: **17-Epinorseredamine. Dihydronorpurpeline. N²-Demethyl-dihydropurpeline**
[65136-98-3]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from the stem bark of *Rauwolfia cumminsii* (Apocynaceae). Yellow amorph. powder. Erroneously indexed in CA.
- Poisson, J. et al., C. R. Hebd. Seances Acad. Sci., 1955, **241**, 1840-1842 (Rauvomitine)
- Haack, E. et al., Naturwissenschaften, 1955, **42**, 627 (Rauvomitine)
- Djerassi, C. et al., J.A.C.S., 1956, **78**, 1259-1260; 1957, **79**, 1217-1222 (Rauvomitine, Tetraphyllicine, isol, uv, ir, struct)
- Gorman, M. et al., Tetrahedron, 1957, **1**, 328-337 (Mauisene)
- Scheuer, P.J. et al., J.O.C., 1963, **28**, 2641-2643 (Mauisene)
- Poisson, J. et al., Bull. Soc. Chim. Fr., 1964, 2683-2689 (Purpeline, Mitoridine, Seredamine)

- Biemann, K. *et al.*, *J.A.C.S.*, 1964, **86**, 4624-4628 (*ms*)
- Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1965, 2497-2501 (*Vincamajoreine*)
- Kaul, J.L. *et al.*, *J. Nat. Prod.*, 1966, **29**, 26-34 (*Majoridine*)
- Kaul, J.L. *et al.*, *Coll. Czech. Chem. Comm.*, 1970, **35**, 116-123 (*Majoridine, struct, ms*)
- Hanaoka, M. *et al.*, *Helv. Chim. Acta*, 1970, **53**, 1723-1732 (*Purpeline, Sereadamine, Noreseredamine, uv, ord, ir, pmr, ms, abs config, synth*)
- Danieli, B. *et al.*, *Chim. Ind. (Milan)*, 1971, **53**, 1042-1043 (*Rauflorine*)
- Banerji, A. *et al.*, *Phytochemistry*, 1974, **13**, 2309-2312 (*Majoridine, Vincamajoreine*)
- Chatterjee, A. *et al.*, *Experientia*, 1976, **32**, 1236 (*Reflexine, Rauflexine, Purpeline*)
- Iwu, M.M. *et al.*, *Experientia*, 1977, **33**, 1268 (*Norpurpeline, Demethyldihydropurpeline*)
- Iwu, M.M. *et al.*, *Planta Med.*, 1977, **32**, 88-99; 1978, **33**, 232-236; 360-364 (*Norrauvomitine, Endolobine, Purpeline, Normitoridine, Trimethoxybenzoylsereadamine, Dihydronorpurpeline*)
- Iwu, M.M. *et al.*, *Phytochemistry*, 1978, **17**, 1651-1654 (*Endolobine*)
- Sabri, N.N. *et al.*, *Phytochemistry*, 1978, **17**, 2023-2026 (*Nortetraphylline, 10-Hydroxynortetraphylline*)
- Chatterjee, A. *et al.*, *Tet. Lett.*, 1978, 3879-3882 (*Reflexine, Rauflexine, struct, cmr*)
- Stöckigt, J. *et al.*, *Tet. Lett.*, 1979, 2615-2618 (*biosynth*)
- Amer, M.A. *et al.*, *Phytochemistry*, 1981, **20**, 2569-2573 (*Acetylnortetraphylline*)
- Kan, C. *et al.*, *Phytochemistry*, 1986, **25**, 1783-1784 (*12-Hydroxymauiensine*)
- Jokela, R. *et al.*, *Planta Med.*, 1996, **62**, 577-579 (*Rauflorine, pmr, cmr*)
- Kato, L. *et al.*, *Phytochemistry*, 2002, **60**, 315-320 (*Demethoxypurpeline*)

Tetraphylline pseudoindoxyl T-289

[92471-49-3]



Absolute Configuration

$C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from the bark of *Neisosperma glomerata* (*Ochrosia glomerata*) (Apocynaceae).

20-Epimer or 7,20-diepimer, demethoxy, 10-methoxy: **Aricine pseudoindoxyl** [1354-50-3]
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. $[\alpha]_D^{20}$ -341 (c, 0.91 in MeOH). C-7 config. not determined.

3,20-Diepimer or 3,7,20-triepimer, demethoxy: **Akuammigine pseudoindoxyl**
 $C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from leaves of *Uncaria elliptica* (Rubiaceae). C-7 config. not determined.

7,20-Diepimer, 10-methoxy: **Isoreserpiline pseudoindoxyl stereoisomer 2**
 $C_{23}H_{28}N_2O_6$ 428.484
Alkaloid from *Rauwolfia grandiflora*.

$[\alpha]_D^{20}$ -90 (c, 0.07 in $CHCl_3$). This stereochem. refers to the most recent isolate (2003) of the alkaloid called Isoreserpiline pseudoindoxyl. The earlier isolates were of a different stereoisomer of incompletely determined abs. config. (see below).

19,20-Diepimer or 7,19,20-triepimer, demethoxy: **Rauniticine pseudoindoxyl** [88375-63-7, 88335-35-7, 88335-34-6, 5802-00-6]

$C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from the leaves of *Uncaria elliptica* (Rubiaceae). C-7 config. not determined.

3,19,20-Trieppimer or 3,7,19,20-tetraepimer, demethoxy: **3-Isoarauniticine pseudoindoxyl**

$C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from leaves of *Uncaria elliptica* (Rubiaceae). C-7 config. not determined.

Stereoisomer, 10-methoxy: **Isoreserpiline pseudoindoxyl stereoisomer 1** [5802-11-9]

$C_{23}H_{28}N_2O_6$ 428.484
Alkaloid from *Rauwolfia vomitoria*, *Rauwolfia ligustrina*, *Rauwolfia volkensii*, *Aspidosperma discolor*, *Ochrosia moorei* and *Bleekeria viitensis* (preferred genus name *Ochrosia*) (Apocynaceae). Cryst. (Me_2CO /pentane). Mp 251-254° (250-253° dec.). $[\alpha]_D^{20}$ -280 (c, 0.108 in $CHCl_3$). This is the more frequently isol. stereoisomer.

Finch, N. *et al.*, *Experientia*, 1963, **19**, 296 (*Isoreserpiline pseudoindoxyl stereoisomer 1*)

Dastour, N.J. *et al.*, *Helv. Chim. Acta*, 1967, **50**, 213 (*Isoreserpiline pseudoindoxyl*)

Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 708 (*Aricine pseudoindoxyl*)


Phillipson, J.D. *et al.*, *Phytochemistry*, 1983, **22**, 1809 (*Rauniticine pseudoindoxyl, 3-Isoarauniticine pseudoindoxyl, Akuammigine pseudoindoxyl, uv, pmr, ms, struct, synth*)

Seguin, E. *et al.*, *J. Nat. Prod.*, 1984, **47**, 687 (*Tetraphylline pseudoindoxyl*)

Cancelieri, N.M. *et al.*, *Magn. Reson. Chem.*, 2003, **41**, 287-290 (*Isoreserpiline pseudoindoxyl stereoisomer 2*)

Tetraponerine 1 T-290

[171980-65-7]



Absolute Configuration

R = -CH₂CH₂CH₃

$C_{13}H_{24}N_2$ 208.346
Trace alkaloid from the poison gland of the New Guinean ant *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +11 (c, 0.14 in $CHCl_3$).

5-Epimer: **Tetraponerine 2** [172139-30-9]

$C_{13}H_{24}N_2$ 208.346
Trace alkaloid from *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +36 (c, 1.79 in $CHCl_3$).

Merlin, P. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 517

(isol)


Devijver, C. *et al.*, *Tetrahedron*, 1995, **51**, 10913 (*pmr, ms, struct*)

Yue, C. *et al.*, *J.O.C.*, 1996, **61**, 4949 (*synth, ir, pmr, cmr, ms*)

Bates, R.W. *et al.*, *Tetrahedron*, 2002, **58**, 5957-5978 (*synth*)

Tetraponerine 3 T-291

[114475-95-5]



Absolute Configuration

R = -CH₂CH₂CH₃ $C_{14}H_{26}N_2$ 222.373

Alkaloid from the poison gland of the New Guinean ant *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +27 (c, 0.07 in $CHCl_3$).

5-Epimer: **Tetraponerine 4**

[114530-37-9]

 $C_{14}H_{26}N_2$ 222.373

From *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +94 (c, 0.2 in $CHCl_3$).

Merlin, P. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 517 (*isol*)

Braekman, J.C. *et al.*, *Org. Mass Spectrom.*, 1989, **24**, 837 (*ms*)

Jones, T.H. *et al.*, *Tet. Lett.*, 1990, **31**, 1535; 4543 (*synth*)

Macours, P. *et al.*, *Tetrahedron*, 1995, **51**, 1415 (*pmr, cmr, cd, struct, abs config*)

Yue, C. *et al.*, *J.O.C.*, 1996, **61**, 4949 (*synth*)

Stragies, R. *et al.*, *J.A.C.S.*, 2000, **122**, 9584-9591 (*synth*)

Bates, R.W. *et al.*, *Tetrahedron*, 2002, **58**, 5957-5978 (*synth*)

Charette, A.B. *et al.*, *Org. Lett.*, 2005, **7**, 5401-5404 (*synth*)

Tetraponerine 5 T-292

[114475-96-6]

As Tetraponerine 1, T-290 with

R = CH₂CH₂CH₂CH₂CH₃ $C_{15}H_{28}N_2$ 236.4

Originally assigned an erroneous struct. Alkaloid from the poison gland of the New Guinean ant *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +10 (c, 0.2 in $CHCl_3$).

5-Epimer: **Tetraponerine 6**

[114530-38-0]

 $C_{15}H_{28}N_2$ 236.4

From *Tetraponera* sp. Oil. $[\alpha]_D^{20}$ +35 (c, 0.15 in $CHCl_3$). Incorrect struct. originally assigned.

Merlin, P. *et al.*, *J. Chem. Ecol.*, 1988, **14**, 517 (*isol*)

Braekman, J.C. *et al.*, *Org. Mass Spectrom.*, 1989, **24**, 837 (*ms*)

Devijver, C. *et al.*, *Tetrahedron*, 1995, **51**, 10913 (*pmr, cmr, cd, synth, struct, abs config*)

Yue, C. *et al.*, *J.O.C.*, 1996, **61**, 4949 (*synth, ir, pmr, cmr*)

Stragies, R. *et al.*, *J.A.C.S.*, 2000, **122**, 9584-9591 (*synth*)

Kim, J.T. *et al.*, *Org. Lett.*, 2002, **4**, 4697-4699 (*synth*)

Bates, R.W. *et al.*, *Tetrahedron*, 2002, **58**, 5957-5978 (*synth*)

Tetraponerine 7 **T-293**

Decahydro-5-pentyl-5H-pyrido[1,2-c]pyrrolo[1,2-a]pyrimidine, 9CI
[114530-39-1]

As Tetraponerine 3, T-291 with
R = (CH₂)₄CH₃

C₁₆H₃₀N₂ 250.426

Struct. revised in 1995. Alkaloid from the poison gland of the New Guinea ant *Tetraponera* sp. Oil. [α]_D +30 (c, 0.22 in CHCl₃).

5-Epimer: Tetraponerine 8

[109269-89-8]

C₁₆H₃₀N₂ 250.426

Major alkaloid from *Tetraponera* sp.

Mp 40°. [α]_D +102 (c, 0.15 in CHCl₃).

▶ Contact poison.

Braekman, J.C. et al., *Z. Naturforsch.*, C, 1987, **42**, 627 (*isol, ir, pmr, cmr, cryst struct*)

Merlin, P. et al., *J. Chem. Ecol.*, 1988, **14**, 517 (*isol*)

Braekman, J.C. et al., *Org. Mass Spectrom.*, 1989, **24**, 837 (*ms*)

Yue, C. et al., *J.O.C.*, 1990, **55**, 1140; 1996, **61**, 4949 (*synth*)

Jones, T.H. et al., *Tet. Lett.*, 1990, **31**, 4543 (*synth*)

Merlin, P. et al., *Tetrahedron*, 1991, **47**, 3805 (*synth*)

Renson, B. et al., *Can. J. Chem.*, 1994, **72**, 105 (*biosynth*)

Barluenga, J. et al., *J.O.C.*, 1994, **59**, 3699 (*synth*)


Macours, P. et al., *Tetrahedron*, 1995, **51**, 1415 (*pmr, cmr, cd, struct, synth, abs config*)

Stragies, R. et al., *J.A.C.S.*, 2000, **122**, 9584-9591 (*synth*)

Bates, R.W. et al., *Tetrahedron*, 2002, **58**, 5957-5978 (*synth*)

Tetrastachyne **T-294**

[79887-93-7]



C₄₂H₅₀N₄O₆ 706.88

Aspidosperma-Iboga dimer. Alkaloid from the leaves of *Bonafousia tetrastachya* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. [α]_D -248 (c, 0.5 in CHCl₃).

14,15-Didehydro: 14-Dehydrotetrastachyne

[91853-53-1]

C₄₂H₄₈N₄O₆ 704.864


Alkaloid from the leaves of *Tabernaemontana citrifolia* and *Peschiera echinata* (Apocynaceae). Amorph. [α]_D -164 (c, 0.31 in EtOH).

Damak, M. et al., *Bull. Soc. Chim. Fr., Ser. II*, 1981, 213 (*isol, uv, ir, pmr, cmr, ms, struct*)

Abaul, J. et al., *C. R. Hebd. Seances Acad. Sci., Ser. 2*, 1984, **298**, 627 (14-Dehydrotetrastachyne)

Tetrastachynine **T-295**

[79887-94-8]



C₄₂H₅₀N₄O₆ 706.88

Alkaloid from the leaves of *Bonafousia tetrastachya* (preferred genus name *Tabernaemontana*) (Apocynaceae). Amorph. [α]_D 0 (c, 1. in CHCl₃).

14,15-Didehydro: 14,15-Dehydrotetrastachynine

C₄₂H₄₈N₄O₆ 704.864

Alkaloid from the leaves of *Stemmadenia grandiflora* and from *Tabernaemontana citrifolia* (Apocynaceae). Oil or amorph. [α]_D²¹ -31 (c, 0.26 in CHCl₃). [α]_D -151 (c, 0.25 in EtOH). [α]_D -124 (c, 0.37 in CHCl₃). Dec. slowly in the presence of air and light. The low opt. rotn. refers to the *S. grandiflora* isolate.


Damak, M. et al., *Bull. Soc. Chim. Fr., Ser. II*, 1981, 213 (*isol, uv, ir, pmr, cmr, ms, struct*)

Torrenegra, R. et al., *Phytochemistry*, 1988, **27**, 1843 (14,15-Dehydrotetrastachynine)

Abaul, J. et al., *J. Nat. Prod.*, 1989, **52**, 1279 (14,15-Dehydrotetrastachynine)

Tetrazomine **T-296**

[132073-72-4]



C₂₄H₃₄N₄O₅ 458.556

Prod. by *Saccharothrix mutabilis* ssp. *chichijimaensis*. Active against gram-positive and -negative bacteria. Cytotoxic. Sol. H₂O, EtOH, MeOH; poorly sol. EtOAc, hexane. Mp 190° dec. [α]_D -62 (c, 1 in MeOH). λ_{max} 207; 224 (MeOH) (Derp). λ_{max} 209; 245 (MeOH) (Berdy). λ_{max} 207; 224 (EtOH) (Berdy).


Suzuki, K. et al., *J. Antibiot.*, 1991, **44**, 479 (*isol, props*)

Sato, T. et al., *J. Antibiot.*, 1991, **44**, 1367 (*pmr, cmr, struct*)

Scott, J.D. et al., *J.A.C.S.*, 2002, **124**, 2951-2956 (*synth, abs config*)

Tetrodonic acid **T-297**

[3270-35-7]



C₁₁H₁₇N₃O₈ 319.271

Isol. from the fish *Fugu pardalis*, *Fugu poccilonotus* and *Fugu rubripes*. Amorph.


Tsuda, K. et al., *Chem. Pharm. Bull.*, 1964, **12**, 634-642 (*struct*)

Tamura, C. et al., *Acta Cryst.*, 1966, **21**, 219-225 (*cryst struct*)

Nakamura, M. et al., *Toxicol.*, 1985, **23**, 271-276 (*isol*)

Tetrodotoxin **T-298**

Tarichatoxin. Spheroidine. Tetrodotoxin. *Fugu* poison. Maculotoxin. Araregai toxin. TTX
[4368-28-9]



C₁₁H₁₇N₃O₈ 319.271

Isol. from the ovaries and liver of Japanese puffer fish (*Sphoeroides rubripes*, *Sphoeroides vermicularis*, *Sphoeroides phyreus*), skin of Californian newt (*Taricha torosa*), from the Japanese ivory shell (*Babylonia japonica*), from the dinoflagellate *Alexandrium tamarense*, xanthid crab *Atergatis floridus* and *Cynops ensicauda*. Believed to be a metab. prod. of a *Pseudomonas* sp. Sodium channel (I, II, III, μ1, h1) blocker. Highly potent toxin. Of practical significance as a cause of accidental poisoning in Japan. Used in neurophysiological research. Anaesthetic synergist. Cryst. Sol. acids; fairly sol. H₂O; poorly sol. EtOAc, hexane. [α]_D²⁵ -8.64 (c, 8.55 in AcOH aq.). pK_a 8.76. Darkens at 225° but does not melt.

▶ LD₅₀ (mus, orl) 435 μg/kg; LD₅₀ (mus, ivn) 9 μg/kg; LD₅₀ (mus, ipr) 0.08 mg/kg. Highly toxic to mammals, birds, reptiles, amphibians and fish. IO1450000

Hydrobromide:

Prisms (EtOH aq.). No definite Mp.

Picrate:

Yellow needles + 1H₂O (H₂O). Darkens >200° without melting.

Hepta-Ac: Mp 188.5-191.5°. [α]_D³⁰ +15.8 (c, 0.39 in CHCl₃).

11-Deoxy: 11-Deoxytetrodotoxin

[112174-24-0]

C₁₁H₁₇N₃O₇ 303.271

Isol. from *Arothron nigropunctatus*, *Cynops ensicauda*, *Fugu* spp. and other marine spp. Paralytic poison, phycotoxin, Na ion blocker. Needles (AcOH aq.). Mp 202° (dec.). [α]_D²⁵ +5.4 (c. 0.3 in AcOH aq.).

▶ LD₅₀ (mus, ipr) 0.71 mg/kg.

11-Hydroxy: 11-Hydroxytetrototoxin. 11-Oxotetrototoxin

[123665-88-3]

C₁₁H₁₇N₃O₉ 335.27

Isol. from the puffer fish *Arothron nigropunctatus*, the frog *Brachycephalus ephippium* and the newt *Notophthalmus viridescens*. Also from the xanthid crab *Atergatis floridus*. Amorph. Exists as covalent hydrate.

4-Deoxy, 4R-[(2S-amino-2-carboxyethyl)thio]: 4-S-Cysteinyltetrototoxin

[853013-71-5]

C₁₄H₂₂N₄O₉S 422.415

Isol. from *Fugu pardalis*.

4-Epimer: 4-Epitetrototoxin

[98242-82-1]

C₁₁H₁₇N₃O₈ 319.271

Isol. from *Arothron nigropunctatus*, *Cynops ensicauda*, *Fugu* spp., *Octopus maculosus* and other marine spp. Phycotoxin, sodium channel blocker, paralytic poison. Amorph.

4-Epimer, 11-deoxy: 11-Deoxy-4-epitetrototoxin

[113564-23-1]

C₁₁H₁₇N₃O₇ 303.271

Isol. from the newt *Cynops ensicauda*. Amorph.

6-Epimer: 6-Epitetrototoxin

[112318-40-8]

C₁₁H₁₇N₃O₈ 319.271

Isol. from *Arothron nigropunctatus*, *Cynops ensicauda*, *Fugu* spp. and other marine spp. Phycotoxin. Amorph. [α]_D²¹ -4.8 (c. 0.3 in AcOH aq.).

▶ LD₅₀ (mus, ipr) 0.6 mg/kg.

6-De(hydroxymethyl): 11-Nortetrototoxin. 11-Nortetrototoxin-6S-ol

[156336-07-1]

C₁₀H₁₅N₃O₇ 289.244

Isol. from the fish *Arothron nigropunctatus*.

6-De(hydroxymethyl), 6-epimer: 6-Epi-11-nortetrototoxin. 6-De(hydroxymethyl)-6-epitetrototoxin. 11-Nortetrototoxin-6R-ol

[81520-41-4]

C₁₀H₁₅N₃O₇ 289.244

Isol. from *Arothron nigropunctatus*, *Atergatis floridus* and *Fugu* spp. incl. from the puffer fish *Fugu niphobles*. Amorph.

Yakoo, A. et al., *Nippon Kagaku Kaishi*, 1950, **71**, 590 (isol)

Tomiie, Y. et al., *Tet. Lett.*, 1963, 2101 (struct)

Woodward, R.B. et al., *Pure Appl. Chem.*, 1964, **9**, 49 (struct)

Goto, T. et al., *Tetrahedron*, 1965, **21**, 2059 (isol, ir, pmr, struct)

Furusaki, A. et al., *Bull. Chem. Soc. Jpn.*, 1970, **43**, 3332 (cryst struct)

U.S. Pat., 1971, 3 966 934; CA, **85**, 87545 (use)

Kishi, Y. et al., *J.A.C.S.*, 1972, **94**, 9219 (synth)

Yasumoto, T. et al., *Agric. Biol. Chem.*, 1986, **50**, 793-795 (isol)

Tetrototoxin, Saxitoxin and the Molecular Biology of the Sodium Channel, (eds. Kao, C.Y. et al), New York Academy of Sciences, 1986, (book)

Simidu, U. et al., *Appl. Environ. Microbiol.*, 1987, **53**, 1714-1715 (occur)

Yotsu, M. et al., *Toxicon*, 1987, **25**, 225-228 (isol)

Yasumoto, T. et al., *J.A.C.S.*, 1988, **110**, 2344-2345 (derivs, isol, pmr, cmr)

Endo, A. et al., *Tet. Lett.*, 1988, **29**, 4127-4128 (6-Epi-11-nortetrototoxin)

Khora, S.S. et al., *Tet. Lett.*, 1989, **30**, 4393-4394 (11-Oxotetrototoxin)

Yotsu, M. et al., *Biosci., Biotechnol., Biochem.*, 1992, **56**, 370-371 (11-Nortetrototoxin-6S-ol)

Arakawa, O. et al., *Fish. Sci.*, 1994, **60**, 769-771 (11-Oxotetrototoxin, 11-Nortetrototoxin-6R-ol)

Yasumoto, T. et al., *J. Toxicol., Toxin Rev.*, 1996, **15**, 81-90 (rev)

Kodama, M. et al., *Toxicon*, 1996, **34**, 1101-1105 (occur)

Food Sci. Technol., Seafood and Freshwater Toxins, (ed. Botana, L.M.), Marcel Dekker, 2000, **103**, (rev)

Miyazawa, K. et al., *J. Toxicol., Toxin Rev.*, 2001, **20**, 11-25 (rev)

Yotsu-Yamashita, M. et al., *J. Toxicol., Toxin Rev.*, 2001, **20**, 51-66 (rev)

Narahashi, T. et al., *J. Toxicol., Toxin Rev.*, 2001, **20**, 67-85 (rev, pharmacol)

Isobe, M. et al., *Tetrahedron*, 2001, **57**, 4543-4558 (5,11-Dideoxytetrototoxin, synth)

Ohyabu, N. et al., *J.A.C.S.*, 2003, **125**, 8798-8805 (synth)

Hinman, A. et al., *J.A.C.S.*, 2003, **125**, 11510-11511 (synth)

Nishikawa, T. et al., *Pure Appl. Chem.*, 2003, **75**, 251-257 (11-Deoxytetrototoxin, synth)

Yotsu-Yamashita, M. et al., *Toxicon*, 2003, **41**, 893-897 (11-Oxotetrototoxin)

Koert, U. et al., *Angew. Chem., Int. Ed.*, 2004, **43**, 5572-5576 (synth)

Daly, J.W. et al., *J. Nat. Prod.*, 2004, **67**, 1211-1215 (rev, occur)

Yotsu-Yamashita, M. et al., *Chem. Res. Toxicol.*, 2005, **18**, 865-871 (4-S-Cysteinyltetrototoxin)

Urabe, D. et al., *Chem. Asian J.*, 2006, **1**, 125-135 (synth)

Jang, J. et al., *Toxicon*, 2006, **48**, 980-987 (occur)

Sato, K. et al., *J.O.C.*, 2008, **73**, 1234-1242 (synth)


Noguchi, T. et al., *Mar. Drugs*, 2008, **6**, 220-242 (rev, occur, activity)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, FOQ000

Texaline

T-299

3-[5-(1,3-Benzodioxol-5-yl)-2-oxazolyl]-pyridine, 9CI. 5-(3,4-Methylenedioxyphenyl)-2-(3-pyridinyl)oxazole
[115070-72-9]



C₁₅H₁₀N₂O₃ 266.256

Alkaloid from the roots of *Amyris texana* (Rutaceae). Shows antimycobacterial activity. Cryst. (EtOAc/hexane). Mp 171-174°.

5'-Methoxy: 3-[5-(7-Methoxy-1,3-benzodioxol-5-yl)-2-oxazolyl]pyridine, 9CI. 2-(3-Methoxy-4,5-methylenedioxyphenyl)-5-(3-pyridinyl)oxazole
[89764-16-9]

C₁₆H₁₂N₂O₄ 296.282

Alkaloid from the aerial portions of *Amyris plumieri* (Rutaceae). Mp 188-189°.

Philip, S. et al., *Heterocycles*, 1984, **22**, 9 (5'-methoxy)


Dominguez, X.A. et al., *Heterocycles*, 1988, **27**, 35 (isol, uv, ir, pmr, cmr, ms, struct)

Rastogi, N. et al., *FEMS Immunol. Med. Microbiol.*, 1998, **20**, 267-273 (activity)

Texensine

T-300

2-(Dimethylamino)-N-[7-(1H-indol-3-ylmethyl)-3-(1-methylethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-4-methylpentanamide, 9CI
[52617-20-6]



C₃₃H₄₃N₅O₄ 573.734

Alkaloid from the aerial part of *Colubrina texensis* (Rhamnaceae). Cryst. (EtOAc). Mp 249-252°. [α]_D²⁵ -144 (c. 0.50 in CHCl₃).

N-De-Me (?): Homoamericine

[27710-05-0]

C₃₂H₄₁N₅O₄ 559.707


Alkaloid from *Ceanothus americanus* (Rhamnaceae). Isol. only as a minor contaminant of Americine, A-690. Struct. uncertain; the struct. of the terminal C5 chain was not determined. Could alternatively be identical with Discarine I in D-836.

Klein, F.K. et al., *J.A.C.S.*, 1968, **90**, 2398-2404 (Homoamericine)

Wani, M.C. et al., *Tet. Lett.*, 1973, 4675-4678 (Texensine)

Thaicanine

T-301



C₂₁H₂₅NO₅ 371.432

(S)-form [102828-06-8]

Alkaloid from the leaves of *Parabaena sagittata* (Menispermaceae). Cryst. (MeOH/CHCl₃). Mp 144-146°. [α]_D²⁰ -243 (CHCl₃).

N-Me: N-Methylthaicanine

[142934-42-7]

C₂₂H₂₈NO₅[⊕] 386.467Alkaloid from leaves of *Anisocycla cymosa* (Menispermaceae). Amorph. powder. [α]_D²⁰ -102.1 (c, 0.014 in MeOH).**Me ether: O-Methylthaicanine**

[102828-07-9]

C₂₂H₂₇NO₅ 385.459Isol. from the leaves of *Parabaena sagittata* (Menispermaceae). Cryst. (MeOH/CHCl₃). Mp 119-120°. [α]_D²⁰ -259 (CHCl₃).**Me ether, N-Me: N,O-Dimethylthaicanine**

[142934-43-8]

C₂₃H₃₀NO₅[⊕] 400.494Alkaloid from leaves of *Anisocycla cymosa* (Menispermaceae). Needles (CHCl₃). Mp 218°. [α]_D²⁰ -102.8 (c, 0.020 in CHCl₃).**O¹⁰-De-Me: Thaipetaline**

[132923-34-3]

C₂₀H₂₃NO₅ 357.405Alkaloid from *Polyalthia stenopetala* (Annonaceae). Amorph. [α]_D -130 (c, 1 in MeOH).**8-Oxo: 8-Oxothaicanine**C₂₁H₂₃NO₆ 385.416Alkaloid from stems of *Coscinium fenestratum* (Menispermaceae). Gum. [α]_D -667 (c, 0.003 in CHCl₃). Tentative struct.; the 3-OH, 4-OMe struct. cannot be excluded.**8-Oxo, Me ether: 4-Methoxy-8-oxotetrahydropalmatine**C₂₂H₂₅NO₆ 399.443Alkaloid from the stems of *Cocculus orbiculatus*. Amorph. yellow powder. [α]_D²⁵ -214.6 (c, 0.9 in MeOH). λ_{max} 224; 279; 308 (MeOH).**(±)-form [126640-91-3]****Me ether:**

Synthetic. Yellow solid. Mp 116-117°.

Ruangrunsi, N. *et al.*, *J. Nat. Prod.*, 1986, **49**, 253 (*isol, uv, ir, pmr, cmr, ms, struct*)Mali, R.S. *et al.*, *Indian J. Chem., Sect. B*, 1989, **28**, 107 (*synth, uv, ir, pmr, cmr, ms, O-Methylthaicanine*)Mali, R.S. *et al.*, *Synth. Commun.*, 1989, **19**, 2613 (*synth*)Janssen, R.H.A.M. *et al.*, *Phytochemistry*, 1990, **29**, 3331 (*cmr*)Lavault, M. *et al.*, *Phytochemistry*, 1990, **29**, 3845 (*Thaipetaline*)Kanyinda, B. *et al.*, *J. Nat. Prod.*, 1992, **55**, 607 (*N-Methylthaicanine, N,O-Dimethylthaicanine*)Pinho, P.M.M. *et al.*, *Phytochemistry*, 1992, **31**, 1403 (*8-Oxothaicanine*)Chang, F.-R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1056-1060 (*4-Methoxy-8-oxotetrahydropalmatine*)**Thaldimerine****T-302**


[11051-30-2]

C₄₀H₄₆N₂O₉ 698.811Struct. not fully known. Aporphine-benzylisoquinoline dimer alkaloid. Alkaloid from *Thalictrum fendleri* (Ranunculaceae). Light-brown oil. Phenolic,contains 6,7-dimethoxyisoquinoline residue. Pmr shows 2 *N*-Me and 6-*O*-Me groups.Shamma, M. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 262-268 (*isol, uv, pmr, ms*)**Thalebanin B****T-303***3-Methylbutanoic acid N-methyl-N-(2-phenylethenyl)amide*. *N*,*N*,*3*-Dimethyl-*N*-(2-phenylethenyl)butanamide, *9CI*(H₃C)₂CH₂CH₂CONMeCH=CHPhC₁₄H₁₉NO 217.31**(E)-form***2,3-Didehydro: Dehydrothalebanin A*. *3-Methyl-2-butenic acid N-methyl-N-(2-phenylethenyl)amide*


[184101-55-1]

C₁₄H₁₇NO 215.294Isol. from leaves of *Glycosmis crassifolia*. Exhibits antifungal activity. λ_{max} 221 (log ε 4.08); 296 (log ε 4.18); 305 (sh) (log ε 4.15) (Et₂O).**(Z)-form [184101-54-0]**Isol. from leaves of *Glycosmis crassifolia*. Exhibits antifungal activity. λ_{max} 216 (log ε 4.04); 270 (log ε 4.08) (Et₂O).*2,3-Didehydro: Dehydrothalebanin B*


[184101-52-8]

C₁₄H₁₇NO 215.294From leaves of *Glycosmis crassifolia* and *Glycosmis pentaphylla*. Exhibits antifungal activity. λ_{max} 221 (log ε 4.06); 281 (log ε 3.92) (Et₂O).Greger, H. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1163-1168 (*isol, uv, ir, pmr, cmr, ms*)Shapiro, J.P. *et al.*, *J. Agric. Food Chem.*, 2000, **48**, 4404-4409 (*isol*)**Thalfine****T-304***Thalphine*C₃₈H₃₆N₂O₈ 648.711**(S)-form [27764-05-2]**Alkaloid from the roots of *Thalictrum foetidum* and *Thalictrum minus* (Ranunculaceae). Antiseptic agent. Yellow prisms (MeOH). Mp 150-151° (141-142°). [α]_D²⁶ +85 (c, 0.35 in MeOH). Log P 8.57 uncertain value (calc).Abdizhabbarova, S. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 330; 1970, **6**, 279; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 281; 1970, **6**, 281 (*isol, uv, pmr, struct*)Shamma, M. *et al.*, *Heterocycles*, 1976, **4**, 1817 (*abs config*)Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257 (*isol, ms, cd, config*)Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 818; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 723 (*cd, abs config*)**Thalphine****T-305***Thalphine*

[27764-06-3]

C₃₉H₄₂N₂O₈ 666.769Alkaloid from the roots of *Thalictrum foetidum* and *Thalictrum minus* Race B, and from *Thalictrum faberi* (Ranunculaceae). Antiseptic. Amorph. Mp 117-118°. [α]_D¹⁶ +115 (c, 0.95 in EtOH) (c, 0.25 in MeOH). Log P 7.7 (uncertain value) (calc).**Hydrochloride (1:2):**

Prisms. Mp 223-226° dec.

Perchlorate (1:2):Needles. Mp 234-235° dec. [α]_D²¹ +135 (c, 1.16 in 2:1 EtOH/H₂O).Abdizhabbarova, S. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 330; 1970, **6**, 279; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 281; 1970, **6**, 281 (*isol, uv, ir, pmr, struct*)Shamma, M. *et al.*, *Heterocycles*, 1976, **4**, 1817 (*abs config*)Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257 (*isol, uv, pmr, ms, cd, config*)**Thalflavine****T-306***8,9-Dihydro-4-methoxy-7-methyl-1,3-dioxolo[4,5-f]isoquinolin-6(7H)-one, 9CI*. *3,4-Dihydro-7-methoxy-2-methyl-5,6-methylenedioxy-1(2H)-isoquinolinone* [125617-79-0]C₁₂H₁₃NO₄ 235.239Struct. revised in 1989. Isomeric 5-methoxy-6,7-methylenedioxy struct. originally proposed. Alkaloid from the roots of *Thalictrum flavum* (Ranunculaceae). Also obt. by the KMnO₄ oxidn. of the bisbenzylisoquinolines Thalracebine, Thalstyline and Thalmelatidine. Needles (MeOH or C₆H₆/petrol). Mp 140°.


[30278-27-4]

Umarov, Kh.S. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 444; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 452 (*isol, ir, pmr, ms*)Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1977, **40**, 281; 1980, **43**, 472 (*synth, uv, ir, pmr, ms*)

- Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257 (synth)
 Aly, Y. *et al.*, *Phytochemistry*, 1989, **28**, 1967 (struct)

Thaliatrine

T-307




$C_{39}H_{42}N_2O_8$ 666.769
 Alkaloid from the roots of *Thalictrum atriplex*. Amorph. powder. Mp 89-90°. $[\alpha]_D^{25} +42.2$ (c, 0.9 in $CHCl_3$). λ_{max} 210; 274 (MeOH).

Gao, G.-Y. *et al.*, *J. Asian Nat. Prod. Res.*, 2005, **7**, 805-809 (isol, cd, pmr, cmr, ms)

Thalibealine

T-308




$C_{42}H_{48}N_2O_9$ 724.849
 Alkaloid from the roots of *Thalictrum wangii*. Amorph. $[\alpha]_D^{24} -162$ (c, 0.5 in $CHCl_3$). λ_{max} 210 (log ϵ 4.8); 223 (sh) (log ϵ 4.69); 281 (log ϵ 4.37); 300 (log ϵ 4.21); 303 (sh) (log ϵ 4.19); 312 (log ϵ 4.17) (MeOH).

Al-Howiriny, T.A. *et al.*, *J. Nat. Prod.*, 2001, **64**, 819-822

Thalibrine

T-309

2'-N-Methylnorthalibrine
 [59614-34-5]



$C_{38}H_{44}N_2O_6$ 624.775
 Alkaloids covered by this entry are enantiomeric with those in Dauricine, D-88 (*R,R*-config.) and diastereomeric with those covered by Berbaminine, B-98 (1*R,1'S*-config.) and Grisabine, G-177 (1*S,1'R*-config.). Alkaloid from the roots

of *Thalictrum rochebrunianum* and *Thalictrum longistylum* (Ranunculaceae). Antiseptic. Amorph. powder. $[\alpha]_D +110$ (c, 0.135 in $CHCl_3$). Log P 6.67 (calc). λ_{max} 282 (ϵ 9000) (MeOH) (Berdy).

2'-N-De-Me: Northalibrine

[59614-33-4]
 $C_{37}H_{42}N_2O_6$ 610.749
 Alkaloid from the roots of *Thalictrum rochebrunianum* (Ranunculaceae). Amorph. powder. $[\alpha]_D +47$ (c, 0.2 in $CHCl_3$).

Me ether: O-Methylthalibrine

[59654-05-6]
 $C_{39}H_{46}N_2O_6$ 638.802
 Alkaloid from roots of *Thalictrum minus* Race B (Ranunculaceae). Shows antimicrobial activity. Antiseptic. Amorph. $[\alpha]_D +82$ (c, 0.36 in $CHCl_3$). Log P 6.95 (calc). Enantiomeric with O-Methylauricine. λ_{max} 280 (ϵ 10500); 285 (ϵ 10250) (MeOH) (Berdy).

O⁷-De-Me, O⁷-Me: Neothalibrine

[73609-03-7]
 $C_{38}H_{44}N_2O_6$ 624.775
 Alkaloid from the fruit of *Thalictrum revolutum* (Ranunculaceae). Amorph. $[\alpha]_D^{27} +155$ (c, 0.5 in MeOH).

O⁷-De-Me, O⁷-Me, 2'-N-Oxide (α):

Neothalibrine 2 α -N-oxide
 [112448-45-0]
 $C_{38}H_{44}N_2O_7$ 640.775
 Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. $[\alpha]_D +74$ (c, 0.2 in MeOH).


O⁷,O¹²-Di-de-Me, N^{2'}-de-Me: Northalibrine

O⁷,O¹²-Didemethylnorthalibrine
 [120091-14-7]
 $C_{35}H_{38}N_2O_6$ 582.695
 Alkaloid from the roots and rhizomes of *Thalictrum minus* var. *minus* (Ranunculaceae). Enantiomer of Pedroamine in D-88.

Saá, J.M. *et al.*, *Heterocycles*, 1976, **4**, 753 (Thalibrine, O-Methylthalibrine, Northalibrine, uv, ir, pmr, ms, struct, synth)
 Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1977, **40**, 281; 1980, **43**, 270; 472-481 (Thalibrine, Methylthalibrine, Neothalibrine, isol, uv, cd, ir, pmr, ms, struct, activity)
 Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 721 (Neothalibrine 2'-oxide)
 Başer, K.H.C. *et al.*, *Planta Med.*, 1988, **54**, 513 (Northalibrine)

Thalibrumine

T-310



$C_{38}H_{40}N_2O_8$ 652.743
 Struct. revised in 1980.

(S)-form [59553-87-6]

Alkaloid from the roots of *Thalictrum rochebrunianum* (Ranunculaceae). Needles (MeOH). Mp 198-200°. $[\alpha]_D +28$ (c,

0.19 in $CHCl_3$).

Me ether: O-Methylthalibrumine

[75956-50-2]
 $C_{39}H_{42}N_2O_8$ 666.769
 Alkaloid from *Thalictrum rochebrunianum* roots (Ranunculaceae). Cryst. ($CHCl_3$). Mp 183-185°. $[\alpha]_D -103.7$ (c, 0.5 in $CHCl_3$). Struct. assignment doubtful due to inconsistent publ. pmr and ms (only 5 OMe groups reported in pmr; parent m/e cited as 665).

 α '-Oxo: Oxothalibrumine

[72187-00-9]
 $C_{38}H_{38}N_2O_9$ 666.726
 Alkaloid from the roots of *Thalictrum rochebrunianum*, also obt. by air oxidn. of Thalibrumine, T-310 (Ranunculaceae). Mp 198-200° dec. $[\alpha]_D^{22} -70$ (c, 0.25 in MeOH).

 α '-Oxo, 3',4'-didehydro: Thalictrine

[72187-01-0]
 $C_{38}H_{36}N_2O_9$ 664.71
 Trace alkaloid from the roots of *Thalictrum rochebrunianum*, also obt. by Pd/C oxidn. of Thalibrumine, T-310 (Ranunculaceae). Shows strong antiinflammatory activity, weak antitumour agent. Cryst. (Me₂CO). Mp 199-201° dec. $[\alpha]_D^{22} -255$ (c, 0.24 in MeOH).

 α '-S-Hydroxy, 3',4'-didehydro: Dihydrothalictrine

[72187-02-1]
 $C_{38}H_{38}N_2O_9$ 666.726
 Minor alkaloid from the roots of *Thalictrum rochebrunianum* (Ranunculaceae). Mp 194-197°. $[\alpha]_D^{22} -125$ (c, 0.13 in MeOH).


Saá, J.M. *et al.*, *Tet. Lett.*, 1976, 513 (uv, pmr, ms)

Ahmad, R. *et al.*, *Islamabad J. Sci.*, 1978, **5**, 38; *CA*, **94**, 30976f (O-Methylthalibrumine)
 Wu, J. *et al.*, *J. O. C.*, 1980, **45**, 208; 213 (Thalictrine, Dihydrothalictrine, Thalibrumine, Oxothalibrumine, !struct, synth)

Thalibrumine

T-311

5,6,6',7,12-Pentamethoxy-2,2'-dimethylberbaman-14-ol, 9CI
 [11021-81-1]



$C_{39}H_{44}N_2O_8$ 668.785

Struct. revised in 1980. Alkaloid from the roots of *Thalictrum rochebrunianum* (Ranunculaceae). Mp 172-173°. $[\alpha]_D^{28} +160$ (c, 0.9 in MeOH).

Ac:

Needles (EtOAc). Mp 236-237°. $[\alpha]_D^{22} +161$ (c, 0.26 in MeCN).

N^{2'}-De-Me: N'-Northalibrumine. 2'-De-methylthalibrumine

[59553-88-7]
 $C_{38}H_{42}N_2O_8$ 654.758
 Alkaloid from roots of *Thalictrum rochebrunianum*, also obt. by redn. of Thalibrumine, T-310 (Ranunculaceae).


ceae). Needles (Me₂CO). Mp 158-161°. [α]_D²⁰ +79 (c, 0.16 in MeOH).

- Fong, H.H.S. *et al.*, *J. Nat. Prod.*, 1966, **29**, 94 (isol, uv)
Cava, M.P. *et al.*, *Tet. Lett.*, 1974, 4259 (cd, pmr, ms)
Wu, J. *et al.*, *J.O.C.*, 1980, **45**, 208; 213 (isol, struct, pmr, deriv, synth)

Thalicarpine

T-312

Thaliblastine. NSC 68075
[5373-42-2]



C₄₁H₄₈N₂O₈ 696.839

Alkaloid from *Thalictrum dasycarpum*, *Thalictrum minus*, several other *Thalictrum* spp. and *Hernandia ovigera* (Ranunculaceae, Hernandiaceae).

Antihypertensive agent acting via non-specific and myocardial depression. Shows antibacterial activity against *M. smegmatis*, but not against other bacterial species studied. Shows *in vitro* antineoplastic activity against HeLa-S₃ cells and Walker 256 carcinoma. Cryst. (EtOH aq. or Et₂O). Mp 108-110° Mp 160-161° Mp 155-157° (dimorph.). [α]_D +133 (MeOH). Log P 6.09 (uncertain value) (calc).

- LD₅₀ (rat, orl) 1500 mg/kg. LD₅₀ (rat, ipr) 140 mg/kg. HQ1775000

N²-Oxide: **Thalicarpine 2'-N-oxide**

C₄₁H₄₈N₂O₉ 712.838

Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). [α]_D +15 (c, 0.14 in CHCl₃).

N⁶-De-Me: **6-Northalicarpine**

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae).

N²-De-Me: **Northalicarpine**

[56021-86-4]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from the roots of *Thalictrum revolutum* (Ranunculaceae). Amorph. [α]_D²¹ +108 (c, 0.25 in MeOH).

O¹-De-Me: **Thalictropine**

[39032-60-5]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from *Thalictrum polygamum* and *Thalictrum dioicum*. Needles (MeOH). Mp 167°. [α]_D²⁵ +120 (c, 0.3 in MeOH). Extremely susceptible to aerial oxidn.

O¹-De-Me, Ac:

Cryst. (MeOH). Mp 182-183°.

O⁶-De-Me: **Thalilutidine**

[66408-23-9]

C₄₀H₄₆N₂O₈ 682.812

Minor alkaloid from the roots of *Thalictrum revolutum* (Ranunculaceae). Amorph. [α]_D²⁰ +74.2 (c, 0.11 in MeOH). λ _{max} 280 (ε 24000); 304 (ε 16000) (MeOH) (Berdy).

O⁷-De-Me: **Thalmelatine**

[5308-77-0]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from *Thalictrum dioicum*, *Thalictrum minus* ssp. *elatum*, and the roots, fruits and tops of *Thalictrum revolutum* (Ranunculaceae). Shows hypotensive activity. Shows antibacterial activity against *Mycobacterium smegmatis*. Cryst. (Et₂O). Mp 110-112° Mp 120-123°. Log P 5.82 (uncertain value) (calc). λ _{max} 287; 302 (MeOH) (Berdy).

O^{4'}-De-Me: **Thalidoxine**

[50802-24-9]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from *Thalictrum dioicum* (Ranunculaceae). Amorph. [α]_D²⁵ +113 (c, 0.2 in MeOH). Isomer of Pennsylvanine. In a subsequent reinvestigation of *T. dioicum*, no thalidoxine was obtd.

O^{4'}-De-Me, Ac:

Cryst. (CHCl₃). Mp 128-129°.

O^{5'}-De-Me: **Pennsylvanine**

[53466-31-2]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from *Thalictrum polygamum*, *Thalictrum dioicum* and the roots and tops of *Thalictrum revolutum* (Ranunculaceae). Shows antibacterial activity against *Mycobacterium smegmatis*. Cryst. (Et₂O). Mp 112-113°. [α]_D²⁴ +131 (c, 0.7 in MeOH). Isomer of Thalidoxine. λ _{max} 281; 303 (MeOH) (Berdy). λ _{max} 282; 305; 320 (MeOH-NaOH) (Berdy).

O^{5'}-De-Me, Ac:

Cryst. (Et₂O). Mp 137-138°.

O¹, O⁷-Di-de-Me: **Thalictrogamine**

[41928-76-1]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from *Thalictrum polygamum* and *Thalictrum dioicum*, and from the tops of *Thalictrum revolutum* (Ranunculaceae). Amorph. [α]_D²⁵ +135 (c, 0.2 in MeOH). Susceptible to aerial oxidn, turns green on standing.

O¹, O⁷-Di-de-Me, di-Ac:

Cryst. (Et₂O). Mp 147-148°.

O¹, O^{5'}-Di-de-Me: **Pennsylvanine**

[53416-85-6]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from *Thalictrum polygamum* (Ranunculaceae). Fine needles (Me₂CO/Et₂O), prisms (Et₂O). Mp 107-108° Mp 128-129°.

O¹, O^{5'}-Di-de-Me, di-Ac:

Cryst. (Et₂O). Mp 147-148°.

O⁷, O^{5'}-Di-de-Me: **Thalipine**

[62724-08-7]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from *Thalictrum polygamum*, *Thalictrum minus* and the fruits and tops of *Thalictrum revolutum* (Ranunculaceae). Amorph. [α]_D²⁵ +141 (c, 0.19 in MeOH). λ _{max} 282 (MeOH) (Berdy).

Didehydro: see Dehydrothalicarpine, D-170

Kupchan, S.M. *et al.*, *J. Polym. Sci.*, 1963, **52**, 985 (*Thalictropine*)

Mollov, N.M. *et al.*, *Tet. Lett.*, 1964, 2219 (*Thalmelatine*)

Tomita, M. *et al.*, *Tet. Lett.*, 1965, 4309 (*Thalicarpine*)

Kupchan, S.M. *et al.*, *J.A.C.S.*, 1973, **95**, 2995 (*synth, Thalictropine*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 508

(*Thalictrogamine, Thalmelatine, Thalipine, Pennsylvanine, Pennsylvanine*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 508; 593

(*Thalictrogamine, Thalmelatine, Thalipine, Pennsylvanine, Pennsylvanine*)

Wu, W.N. *et al.*, *Tetrahedron*, 1977, **33**, 2919 (*Thalilutidine*)

Shamma, M. *et al.*, *J. Nat. Prod.*, 1978, **41**, 169

(*Thalictrogamine, Thalictropine, Thalmelatine, Thalidoxine, Pennsylvanine, Pennsylvanine*)

Wu, W.N. *et al.*, *J. Nat. Prod.*, 1980, **43**, 567 (*Northalicarpine*)

Sidjimov, A. *et al.*, *Phytochemistry*, 1982, **21**, 871 (*biosynth*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1982, **38**, 729 (*biosynth*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 173 (rev. *pharmacol, Thalictropine*)

Chalandre, M.C. *et al.*, *Can. J. Chem.*, 1986, **64**, 123 (*synth, uv, Dehydrothalmelatine, 6-Northalicarpine, Thalictropine oxide*)

Chen, G. *et al.*, *Cancer Lett. (Shannon, Irel.)*, 1992, **62**, 173 (*pharmacol, Thalictropine*)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold*, 1992, (TEH250)

Thalicerberine

T-313

Dihydrothalmethine

[602-83-5]



C₃₇H₄₀N₂O₆ 608.733

Diastereoisomeric with Isothalicerberine, I-329. Alkaloid from the stems and leaves of *Thalictrum thunbergii* and the roots of *Thalictrum lucidum* and *Thalictrum minus* (Ranunculaceae). Shows weak *in vivo* antitubercular activity. Needles + 1H₂O. Mp 161°. [α]_D +231.2 (CHCl₃). [α]_D²⁵ +210 (c, 0.02 in MeOH). Log P 7.83 (uncertain value) (calc).

- LD₅₀ (mus, ipr) 125 mg/kg. XG2001000

Me ether: **O-Methylthalicerberine. Thalimidine**

[5096-71-9]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from the stems and leaves of *Thalictrum thunbergii* and other *Thalictrum* spp. (Ranunculaceae). Weak antineoplastic agent. Antiinflammatory, antihypertensive agent. Mp 186-187°. [α]_D¹⁹ +244.6. Log P 8.4 (uncertain value) (calc).

- LD₅₀ (mus, ipr) 125 mg/kg. XG2003000

Me ether, dipicrate: Mp 188-199° dec. [α]_D²⁵ +140 (c, 0.3 in CHCl₃).

O⁷-De-Me: **Thalivarmin. 7-O-De-methylthalicerberine**

[101312-86-1]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the above-ground parts of *Thalictrum minus* var. *minus* (Ranunculaceae).

O⁷-De-Me, O¹²-Me: **Thaliphylline**

[93780-79-1]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from the roots and rhizomes of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). $[\alpha]_D^{25} +198$ (c, 0.12 in MeOH).

**O⁷-De-Me, O¹²-Me, 2'-β-N-oxide: Thali-
phylline 2'β-N-oxide**
[112464-19-4]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. $[\alpha]_D +257$ (c, 0.7 in MeOH).

**O⁷-De-Me, O¹²-Me, N²-de-Me: 2'-
Northaliphylline**

[105418-70-0]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Also obt. by NaBH₄ reduct. of Thalsiviasine in T-339. Amorph. $[\alpha]_D +197$ (c, 0.15 in MeOH).

Yunusov, S. et al., *Zh. Obshch. Khim.*, 1950, **20**, 1151; *CA*, **45**, 1608c (isol, deriv)

Fujita, E. et al., *Yakugaku Zasshi*, 1959, **79**, 1256; 1260; 1963, **83**, 159; *CA*, **54**, 4643h; 4644b; **59**, 3971f (struct)

Tomimatsu, T. et al., *Yakugaku Zasshi*, 1963, **83**, 153; *CA*, **59**, 3971b (struct)

DeJongh, D.C. et al., *J.A.C.S.*, 1966, **88**, 1052 (ms, deriv)

Ismailov, Z.F. et al., *Khim. Prir. Soedin.*, 1968, **4**, 256; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 220 (ms, deriv)

Doskotch, R.W. et al., *J. Nat. Prod.*, 1969, **32**, 29 (isol, deriv)

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1970, **6**, 705; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 715 (ord, deriv)

Duchevska, Kh.B. et al., *Dokl. Bulg. Akad. Nauk*, 1971, **24**, 467; *CA*, **75**, 106101x (isol)

Fujita, E. et al., *Chem. Pharm. Bull.*, 1972, **20**, 368 (synth, pmr, ms, deriv)

Baldas, J. et al., *J.C.S. Perkin I*, 1972, 597 (ms)

Wu, W.-N. et al., *J. Nat. Prod.*, 1976, **39**, 204;

378; 1977, **40**, 508 (isol, uv, ir, pmr, cd)

Moiseeva, G.P. et al., *Khim. Prir. Soedin.*, 1979, **15**, 818; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 723 (cd, abs config)

Wu, J. et al., *J. Nat. Prod.*, 1980, **43**, 270 (isol, pmr, deriv)

Guinaudeau, H. et al., *Tetrahedron*, 1984, **40**, 1975 (Thaliphylline)

Baser, K.H.C. et al., *Planta Med.*, 1985, **448** (Thalivarmine)

Hussain, S.F. et al., *J. Nat. Prod.*, 1986, **49**, 488 (2'-Northaliphylline, synth)

Herath, W.H.M.W. et al., *J. Nat. Prod.*, 1987, **50**, 721 (2'-Northaliphylline, 2'-Northaliphylline N-oxide)

Thalicflavine T-314


[37274-00-3]

Struct. unknown. Alkaloid from *Thalictrum flavum*.

Vichkanova, S.A. et al., *CA*, 1973, **78**, 66905r

Thalicminine T-315

3,9,10-Trimethoxy-1,2-methylenedioxyoxoaporphine
[16408-77-8]



C₂₀H₁₅NO₆ 365.342

Alkaloid from *Thalictrum* spp. and *Ocotea puberula*. Respiratory stimulant, transient hypotensive agent. Golden-orange cryst. (MeOH/CHCl₃). Mp 263-265° (274-277°). λ_{\max} 254 (log ϵ 4.7); 280 (log ϵ 4.53); 357 (log ϵ 4.43); 452 (log ϵ 3.4) (EtOH).

O⁹-De-Me: Isofiliformine

C₁₉H₁₃NO₆ 351.315

Alkaloid from *Cassytha filiformis*. Amorph. red powder (CHCl₃/MeOH). Mp > 300°. λ_{\max} 250 (log ϵ 3.67); 281 (log ϵ 3.73); 356 (log ϵ 3.16); 461 (log ϵ 2.96) (MeOH). λ_{\max} 257 (log ϵ 3.9); 294 (log ϵ 4.03); 389 (log ϵ 3.32); 536 (log ϵ 3.04) (MeOH/NaOH).

O¹⁰-De-Me: Filiformine†

[208252-25-9]

C₁₉H₁₃NO₆ 351.315

Alkaloid from *Cassytha filiformis*. Amorph. red powder (CHCl₃). Mp 318-320°. λ_{\max} 250 (log ϵ 4.11); 279 (log ϵ 4.2); 320 (log ϵ 4.03); 406 (log ϵ 3.27) (MeOH).

Pulatova, K.G. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **2**, 349-350 (isol, ir, uv)


Baralle, F.N. et al., *Phytochemistry*, 1973, **12**, 948-949 (isol, ir, uv, pmr, ms)

Chang, F.R. et al., *J. Nat. Prod.*, 1998, **61**, 863-866 (Filiformine)

Tsai, T.-H. et al., *J. Nat. Prod.*, 2008, **71**, 289-291 (Isofiliformine)

Thalicipureine T-316

2,3,4,6,7-Pentamethoxy-N-methyl-1-phenanthrenemethanamine, 9CI. 2,3,4,6,7-Pentamethoxy-1-(2-methylaminoethyl)-phenanthrene
[218900-91-5]




C₂₂H₂₇NO₅ 385.459

Alkaloid from the leaves of *Annona purpurea* (soncoya). Antiplatelet aggregation agent. Yellow needles (CHCl₃). Mp 196-197°. λ_{\max} 220 ; 264 ; 308 ; 326 ; 345 ; 358 (sh) (EtOH).

Chang, F.-R. et al., *J. Nat. Prod.*, 1998, **61**, 1457-1461 (isol, uv, pmr, cmr, ms)

Thalicsessine T-317

[113807-86-6]



C₂₂H₂₇NO₄ 369.46

Alkaloid from the roots of *Thalictrum sessile* (Ranunculaceae). Prisms (MeOH). Mp 213-216°. $[\alpha]_D^{25} +113$ (c, 0.2 in CHCl₃).

**N-De(2-hydroxyethyl), N-Me: Cardu-
chorone**
[240121-82-8]

C₂₁H₂₅NO₃ 339.433


Alkaloid from *Delphinium carduchorum*.

Wu, Y.-C. et al., *Heterocycles*, 1987, **26**, 943-946 (Thalicsessine)

Merici, A.H. et al., *Phytochemistry*, 1999, **51**, 337-340 (Carduchorone)

Thalicthuberine T-318

1,2-Dimethoxy-N,N-dimethylphenanthro[2,3-d][1,3]dioxole-4-ethanamine, 9CI. 1-(2-Dimethylaminoethyl)-3,4-dimethoxy-6,7-methylenedioxyphenanthrene. Thalicthuberine
[477-35-0]



C₂₁H₂₃NO₄ 353.417

Aporphine-derived numbering shown. Alkaloid from the roots of *Thalictrum thunbergii* and *Thalictrum rugosum*, and from the roots and rhizomes of *Thalictrum strictum* (Ranunculaceae). Shows a wide range of *in vitro* antimicrobial activity. Needles (Et₂O). Sol. MeOH, H₂O; fairly sol. hexane. Mp 126-127°. λ_{\max} 216 (log ϵ 4.22); 261 (log ϵ 4.78); 288 (log ϵ 4.28); 310 (log ϵ 4.01); 322 (log ϵ 3.99); 347 (log ϵ 3.25); 364 (log ϵ 3.21) (MeOH).

Hydrochloride:

Needles + 0.5H₂O. Mp 209-210°.

N-Oxide: Thalicthuberine N-oxide

[149472-49-1]

C₂₁H₂₃NO₅ 369.416

Alkaloid from aerial parts of *Platycapnos spicata* (Papaveraceae). Prisms (EtOH). Mp 112-114°.

N-De-Me: Northalichthuberine

[160568-07-0]

C₂₀H₂₁NO₄ 339.39

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae).

**N-De-Me, N-hydroxy: N-Hydroxy-
northalichthuberine**


[177413-38-6]

C₂₀H₂₁NO₅ 355.39

Alkaloid from aerial parts of *Thalictrum simplex*.

**O²-De-Me: 2-O-Demethylthalicthuberine.
3-O-Demethylthalicthuberine**

[154814-42-3]

C₂₀H₂₁NO₄ 339.39Alkaloid from leaves of *Ocotea insularis* (Lauraceae). Mp 182-184°.Fujita, E. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 1252; *C.A.*, **54**, 4643f (*isol. uv. ir. struct*)Maekh, S.Kh. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 560; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 507 (*isol*)Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1980, **43**, 143 (*isol. pmr*)Hasbun, C. *et al.*, *Fitoterapia*, 1993, **64**, 440 (3-*O*-Demethylthalictuberine)Blanco, O.M. *et al.*, *Phytochemistry*, 1993, **32**, 1055 (*oxide*)Velcheva, M.P. *et al.*, *Planta Med.*, 1994, **60**, 485 (*Northalictuberine*)Velcheva, M.P. *et al.*, *Phytochemistry*, 1996, **42**, 535 (*N-Hydroxynorthalictuberine*)Ningirawath, S. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 2006, **53**, 443-447 (*synth. uv. pmr. cmr*)**Thalictricavine****T-319**

(+)-form


C₂₁H₂₃NO₄ 353.417**(+)-form** [29550-24-1]Alkaloid from *Corydalis tuberosa* and *Corydalis cava* (Papaveraceae). Mp 148-149° (darkens). [α]_D²⁵ +291.9 (c, 0.555 in CHCl₃).**O¹⁰-De-Me: Epiapocavidine**C₂₀H₂₁NO₄ 339.39Alkaloid from *Corydalis tuberosa* (Papaveraceae). Noncryst. No opt. rotn. reported. Not the epimer of Apocavidine.**(±)-form**Alkaloid from *Corydalis cava*, also obt. by racemisation of the (+)-form (Papaveraceae). Mp 209-210° (204-206°).**O¹⁰-De-Me:**

V. pale yellow cryst. Mp 230° (prior dec.).

Manske, R.H.F. *et al.*, *J.A.C.S.*, 1953, **75**, 4928 (*isol. struct*)Manske, R.H.F. *et al.*, *An. R. Soc. Esp. Fis. Quim., Ser. B*, 1972, **68**, 689 (*Epiapocavidine*)Taka, N. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 3185; 1977, **25**, 1426 (*ir. cmr. conformm*)Kametani, T. *et al.*, *J.C.S. Perkin I*, 1977, 1151 (*synth*)Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 2261 (*isol*)Cushman, M. *et al.*, *J.O.C.*, 1979, **44**, 407 (*synth*)Iwasa, K. *et al.*, *J.O.C.*, 1981, **46**, 4744 (*abs config. synth*)Hanaoka, M. *et al.*, *Chem. Comm.*, 1985, 1257 (*synth*)Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 399-404Marek, R. *et al.*, *Magn. Reson. Chem.*, 2002, **40**, 687-692 (*N-15 nmr*)**Thalidasine****T-320**

6,6',7,7',12-Pentamethoxy-2,2'-dimethylthalidasan, 9CI. 12-O-Methylthalfoetidine

[16623-56-6]

C₃₉H₄₄N₂O₇ 652.786Alkaloid from the roots of *Thalictrum dasycarpum*, *Thalictrum lucidum*, *Thalictrum revolutum*, *Thalictrum rugosum*, *Thalictrum alpinum*, *Thalictrum faberi* and *Thalictrum minus* Race B, and from the rhizomes of *Thalictrum foliolosum* (Ranunculaceae). Antineoplastic, antihypertensive and antibacterial agent. Shows *in vivo* activity against *Mycobacterium smegmatis*. Pale yellow amorph. powder. Mp 144-145° (105-107°). [α]_D²⁷ -70 (c, 0.89 in MeOH). [α]_D²⁵ -68 (c, 1.2 in CHCl₃). Log P 7.83 (uncertain value) (calc). λ_{max} 275 (ε 4560); 282 (ε 4530) (EtOH) (Derep). λ_{max} 275 (ε 4560); 285 (ε 4530) (MeOH) (Berdy).▶ LD₅₀ (mus, ivn) 120 mg/kg, LD₅₀ (mus, ipr) 520 mg/kg. XG2005000**Oxalate:**Cryst. + 2.5H₂O (EtOH). Mp 160-162°. [α]_D²⁶ -53 (c, 0.55 in MeOH).**Dipicrate:**

Cryst. (EtOH). Mp 175-177°.

N²-α-Oxide: Thalidasine 2α-N-oxideC₃₉H₄₄N₂O₈ 668.785Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D +6 (c, 0.15 in MeOH).**N²-De-Me: N²-Demethylthalidasine. 2-Northalidasine**

[78432-93-6]

C₃₈H₄₂N₂O₇ 638.759Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Antineoplastic agent. Pale yellow amorph. powder. Mp 137-139°. [α]_D -86.9 (c, 0.414 in MeOH). Log P 7.02 (uncertain value) (calc).**N²-De-Me, N²-formyl: Thalrugosinone**

[73609-02-6]

C₃₉H₄₂N₂O₈ 666.769Alkaloid from roots of *Thalictrum rugosum* and from *Thalictrum cultratum* (whole plant) (Ranunculaceae). Noncryst. [α]_D²² -42 (c, 0.30 in MeOH). Darkens on exp. to air.**O⁷-De-Me: Thalrugosidine**

[33954-34-6]

C₃₈H₄₂N₂O₇ 638.759Alkaloid from the roots of *Thalictrum rugosum*, *Thalictrum alpinum* and *Thalictrum foliolosum* (Ranunculaceae). Shows antibacterial activity. Needles (MeOH). Mp 175-176° (170-171°). [α]_D³⁰ -185 (MeOH). λ_{max} 275 (ε 9800); 282 (ε 9800) (MeOH) (Berdy).**O⁷-De-Me, N²-de-Me: N-Demethylthal-****rugosidine**

[74683-04-8]

C₃₇H₄₀N₂O₇ 624.732Alkaloid from roots of *Thalictrum alpinum* (Ranunculaceae). Needles (MeOH). Mp 205-206°. [α]_D²¹ -57 (c, 0.23 in MeOH).**O⁷-De-Me, N²-de-Me, N²-formyl: Thalpindione. 2-Formyl-2-northalidasine. 7-O-Demethylthalrugosinone**

[74690-97-4]

C₃₈H₄₀N₂O₈ 652.743Alkaloid from the roots of *Thalictrum alpinum* (Ranunculaceae). Pale yellow amorph. solid. [α]_D²¹ -41.5 (c, 0.29 in MeOH). Struct. revised in 1985 (formerly assigned as C₃₇H₃₆N₂O₉).**O¹²-De-Me: Thalfoetidine. Thaltrimine†**

[16687-93-7]

C₃₈H₄₂N₂O₇ 638.759Alkaloid from the aerial parts of *Thalictrum foetidum* (Ranunculaceae). Antiinflammatory agent. Prisms + 0.5 Et₂O (Et₂O). Mp 168-170°. [α]_D²¹ -88.6 (CHCl₃). Log P 7.25 (uncertain value) (calc). λ_{max} 275 (ε 4560); 282 (ε 4530) (EtOH) (Derep).**O⁷, O¹²-Di-de-Me: Thaligosidine**


[64252-82-0]

C₃₇H₄₀N₂O₇ 624.732Alkaloid from the roots of *Thalictrum rugosum* (Ranunculaceae). Antiseptic. Cryst. + 1.5H₂O (C₆H₆). Mp 175-177°. [α]_D²⁰ -45 (c, 0.26 in MeOH). Log P 7 (uncertain value) (calc). λ_{max} 275 (ε 5230); 283 (ε 5250) (MeOH) (Berdy). λ_{max} 275 (ε 5620); 284 (ε 5760); 310 (ε 1670) (NaOH) (Berdy).**5-Hydroxy: 5-Hydroxythalidasine**C₃₉H₄₄N₂O₈ 668.785Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D -51 (c, 0.1 in MeOH).**5-Hydroxy, N²-α-oxide: 5-Hydroxythalidasine 2α-N-oxide**C₃₉H₄₄N₂O₉ 684.785Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D -11 (c, 0.4 in MeOH).Mollov, N.M. *et al.*, *Chem. Ind. (London)*, 1966, 1178 (*Thalfoetidine, isol. uv. ir. pmr*)Kupchan, S.M. *et al.*, *J.O.C.*, 1968, **33**, 1052; 1969, **34**, 3884 (*Thalidasine, isol. uv. pmr, ms. struct, config, synth*)Ismailov, Z.F. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 256; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 220 (*Thalfoetidine, ms*)Kupchan, S.M. *et al.*, *J.O.C.*, 1969, **34**, 3884 (*Thalfoetidine, struct, config*)Moisseva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 705; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 715 (*ord*)Mitscher, L.A. *et al.*, *Dtsch. Apoth. -Ztg.*, 1971, **111**, 1704 (*activity*)Georgiev, V.S. *et al.*, *Phytochemistry*, 1971, **10**, 2161 (*Thalfoetidine, struct*)Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 167 (*Thalidasine, Thalrugosidine, isol. uv. pmr*)Baldas, J. *et al.*, *J.C.S. Perkin I*, 1972, 597 (*ms*)Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 204; 1977, **40**, 508; 1980, **43**, 143; 372 (*Thalidasine, N-Demethylthalrugosidine, Thalpindione*)

- Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257 (isol)
 Wu, W.-N. *et al.*, *J.O.C.*, 1978, **43**, 580
 (*Thalidasine, Thaligosidine, isol, uv, ir, ms, pmr, struct*)
 Lin, L.-Z. *et al.*, *CA*, 1981, **95**, 76882w; 86198s
 (*N-Demethylthalidasine*)
 Chattopadhyay, S.K. *et al.*, *J. Nat. Prod.*, 1981, **44**, 45 (*Thalrugosidine, isol, uv, cd, ir, pmr, ms*)
 Bhakuni, D.S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 252 (*Thalidasine, Thalrugosidine, isol, uv, pmr*)
 Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1985, **48**, 962; 1986, **49**, 488 (*Thalpidione, 5-Hydroxythalidasine, struct, uv, ir, pmr, ms, cd*)
 Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 721 (*oxides*)

Thalidastine

T-321



$C_{19}H_{16}NO_5^{\oplus}$ 338.339
 Darkens and dec. $>230^{\circ}$. Yellow-brown
 cryst. (as chloride). $[\alpha]_D^{25} +138$ (MeOH).

(R)-form [4839-14-9]

Quaternary alkaloid from *Thalictrum fendleri* and *Nandina domestica* (Ranunculaceae, Berberidaceae).

O¹⁰-Me: Berberastine

[2435-73-6]

 $C_{20}H_{18}NO_5^{\oplus}$ 352.366

Alkaloid from *Hydrastis canadensis* and *Coptis* sp. (Ranunculaceae). Yellow solid or cryst. (as iodide). Mp 300° (iodide). $[\alpha]_D +107$ (c, 0.06 in EtOH). No abs. config. correspondence with Thalidastine implied.

5-Deoxy, 5,6-didehydro: 10-Hydroxy-9-methoxybenzo[1,3-benzodioxolo][5,6-a]quinolinizinium(1+), 9CI.**Deoxythalidastine**

[4839-16-1]

 $C_{19}H_{14}NO_4^{\oplus}$ 320.324


Alkaloid from callus tissue of *Thalictrum minus*. Also obt. by dehydration of Thalidastine chloride. Yellow-brown solid (as chloride). Mp 210° dec. CAS No. refers to chloride.

- Nijland, M.M. *et al.*, *Pharm. Weekbl.*, 1961, **96**, 640; 1963, **98**, 301; *CA*, **55**, **59**; 26371h (isol, uv, ir, struct)
 Monković, I. *et al.*, *Can. J. Chem.*, 1965, **43**, 2017 (biosynth)
 Shamma, M. *et al.*, *Tet. Lett.*, 1965, 3825 (*Thalidastine, Deoxythalidastine, isol, struct*)
 Shamma, M. *et al.*, *J. Pharm. Sci.*, 1968, **57**, 262 (*Thalidastine, isol*)
 Dyke, S.F. *et al.*, *Tetrahedron*, 1975, **31**, 561 (synth, uv, ir, pmr)
 Zarga, M.H.A. *et al.*, *Tet. Lett.*, 1980, **21**, 3739 (abs config)
 Ikuta, A. *et al.*, *Phytochemistry*, 1982, **21**, 1419 (*Deoxythalidastine*)
 Ikuta, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 189 (isol)
 Napolitano, E. *et al.*, *J.C.S. Perkin I*, 1987, 2565 (synth)

Thalifaberine

O³-Methylthalifarapine

[88313-32-0]

 $C_{41}H_{48}N_2O_8$ 696.839

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Shows cytotoxic activity against several human cancer cell lines, as well as antimalarial activity. Amorph. yellowish solid. Sol. MeOH, $CHCl_3$; poorly sol. hexane. Mp $80-85^{\circ}$. $[\alpha]_D^{25} +94.6$ (c, 0.38 in MeOH). λ_{max} 282 (ε 23000) (MeOH) (Berdy).

O¹-De-Me: Thalifalandine

[107019-96-5]

 $C_{40}H_{46}N_2O_8$ 682.812

Minor alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Cytotoxic agent. Amorph. solid. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{14} +83.3$ (c, 0.375 in MeOH). λ_{max} 205 (ε 29500); 285 (ε 10715) (MeOH) (Berdy). λ_{max} 204 (ε 23500); 290 (ε 6760); 313 (ε 8910); 325 (ε 8900) (MeOH-NAOH) (Berdy).

O³-De-Me: Thalifarapine. Thalifaroline

[91926-02-2]

 $C_{40}H_{46}N_2O_8$ 682.812

Alkaloid from the roots of *Thalictrum faberi* and *Thalictrum cultratum* (Ranunculaceae). Amorph. $[\alpha]_D^{24} +98.6$ (c, 0.422 in MeOH) (+73).

O⁶-De-Me: Thalifarazine

[105418-68-6]

 $C_{40}H_{46}N_2O_8$ 682.812

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). $[\alpha]_D +72$ (c, 0.06 in MeOH).

O⁷-De-Me: Thalifaretine

[105418-69-7]

 $C_{40}H_{46}N_2O_8$ 682.812

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). $[\alpha]_D +61$ (c, 0.1 in MeOH).

O³,O⁶-Di-de-Me: 3-Hydroxy-6'-demethyl-9-O-methylthalifaboramine $C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +85.6$ (c, 0.005 in MeOH). λ_{max} 284 (log ε 4); 314 (sh) (log ε 3.85) (MeOH).

O³,O⁹-Di-de-Me: 3-Hydroxythalifaboramine $C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +92.3$ (c, 0.007 in MeOH). λ_{max} 282 (log ε 4.07); 314 (sh) (log ε 3.85) (MeOH).

O⁶,O⁹-Di-de-Me: Thalifaberidine $C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from roots of *Thalictrum faberi* (Ranunculaceae). Shows cyto-

T-322

toxic activity against several human cancer cell lines as well as antimalarial activity. Yellow powder. $[\alpha]_D +87$ (c, 0.1 in MeOH). λ_{max} 284 (ε 12600); 314 (ε 7950) (MeOH) (Berdy).

O⁷,O⁹-Di-de-Me: Thalifaricine

[106146-68-3]

 $C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). $[\alpha]_D +66$ (c, 0.1 in MeOH).

O³,O⁶,O⁹-Tri-de-Me: 3-Hydroxy-6'-odemethylthalifaboramine $C_{38}H_{42}N_2O_8$ 654.758

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +76.2$ (c, 0.002 in MeOH). λ_{max} 285 (log ε 4.01); 314 (sh) (log ε 3.78) (MeOH).

3-Demethoxy: Thalifarone

[105458-70-6]

 $C_{40}H_{46}N_2O_7$ 666.813

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). $[\alpha]_D +68$ (c, 0.1 in MeOH).

3-Demethoxy, O⁷-de-Me: Thalifaramine

[105437-16-9]

 $C_{39}H_{44}N_2O_7$ 652.786

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). $[\alpha]_D +76$ (c, 0.06 in MeOH).

3-Demethoxy, O⁹-de-Me: Thalifaboramine

[109028-30-0]

 $C_{39}H_{44}N_2O_7$ 652.786

Alkaloid from the roots of *Thalictrum faberi*. Yellow amorph. solid. $[\alpha]_D +107.4$ (c, 0.135 in MeOH).

3-Demethoxy, O⁶,O⁹-di-de-Me: 6'-O-De-methylthalifaboramine $C_{38}H_{42}N_2O_7$ 638.759

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +65.5$ (c, 0.005 in MeOH). λ_{max} 284 (log ε 4.05); 314 (sh) (log ε 3.8) (MeOH). λ_{max} 284 (ε 10700); 314 (MeOH) (Berdy).

5'-Hydroxy: Thalifabatine

[91926-03-3]

 $C_{41}H_{48}N_2O_9$ 712.838

Alkaloid from roots of *Thalictrum faberi* (Ranunculaceae). Yellow solid. $[\alpha]_D^{22} +60.5$ (c, 0.154 in MeOH).

5'-Hydroxy, O³-de-Me: Thalifasine

[91926-04-4]

 $C_{40}H_{46}N_2O_9$ 698.811

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Yellow solid. $[\alpha]_D^{25} +67.9$ (c, 0.80 in MeOH).

5'-Hydroxy, O³,O⁹-di-de-Me: 3,5'-Dihydroxythalifaboramine $C_{39}H_{44}N_2O_9$ 684.785

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +89.8$ (c, 0.008 in MeOH). λ_{max} 285 (log ε 4.08); 314 (sh) (log ε 3.88) (MeOH).

5'-Hydroxy, 3-demethoxy, O⁹-de-Me: 5'-Hydroxythalifaboramine $C_{39}H_{44}N_2O_8$ 668.785

Alkaloid from *Thalictrum faberi*. Yellow powder. $[\alpha]_D +97.3$ (c, 0.008 in MeOH). λ_{max} 283 (log ε 4.06); 314 (sh) (log ε 3.82) (MeOH).

6a,7-Didehydro: Dehydrothalifaberine

[91926-01-1]

C₄₁H₄₆N₂O₈ 694.823

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Amorph. yellow solid. [α]_D²⁴ +95.9 (c, 0.143 in MeOH).

Lin, L.-Z. *et al.*, *Planta Med.*, 1983, **49**, 55 (*uv, pmr, ms, cd, struct*)

Wagner, H. *et al.*, *Tetrahedron*, 1984, **40**, 2133 (*isol, uv, pmr, ms, cd, struct, derivs*)

Lin, L.-Z. *et al.*, *Heterocycles*, 1986, **24**, 2731 (*Thalifalandine*)


Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1986, **49**, 494 (*isol, uv, pmr, ms, cd, struct*)

Lin, L.Z. *et al.*, *Phytochemistry*, 1987, **26**, 583; 1999, **50**, 829-834 (*isol, uv, ir, pmr, cmr, ms, derivs*)

Lin, L.-Z. *et al.*, *J. Nat. Prod.*, 1994, **57**, 1430 (*Thalifaberidine*)

Thalifabine**T-323**

[88313-34-2]

C₄₁H₄₆N₂O₉ 710.822

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Yellow amorph. solid. [α]_D²⁵ +78.3 (c, 0.53 in MeOH).

Lin, L.-Z. *et al.*, *Planta Med.*, 1983, **49**, 55 (*uv, pmr, ms, cd, struct*)

Wagner, H. *et al.*, *Tetrahedron*, 1984, **40**, 2133 (*isol, uv, pmr, ms, cd, struct*)

Thalifabomine**T-324**

[92047-65-9]


C₃₉H₄₄N₂O₇ 652.786

Aporphine-benzylisoquinoline dimer of unknown struct. Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). [α]_D²⁶ +67.5 (c, 0.16 in MeOH).

Wagner, H. *et al.*, *Tetrahedron*, 1984, **40**, 2133-2139 (*isol, uv, pmr, ms, cd*)

Thalifaurine**T-325**

[75491-94-0]

C₁₉H₁₆NO₄[⊕] 322.34

Quaternary alkaloid from *Thalictrum fauriei* (whole plant) (Ranunculaceae). Fine yellow needles (MeOH)(as chloride). Mp 258-260° (chloride).

O-De-Me, O³-Me: *Dehydropseudocheilanthifoline*


[64191-03-3]

C₁₉H₁₆NO₄[⊕] 322.34

Quaternary alkaloid from the leaves of *Isopyrum thalictroides* (Ranunculaceae). Yellow cryst. + 1H₂O (as chloride). Mp >300° dec.

Moullis, C. *et al.*, *Phytochemistry*, 1977, **16**, 1283-1287 (*Dehydropseudocheilanthifoline*)

Chen, C.-H. *et al.*, *J. Pharm. Sci.*, 1980, **69**, 1061 (*Thalifaurine*)

Thalifenderine**T-326**C₂₀H₂₅NO₄ 343.422**(R)-form** [4668-18-2]

Alkaloid from *Thalictrum fendleri* (Ranunculaceae). Cryst. Mp 177-178°. [α]_D²⁵ -108 (MeOH).

4'-O- β -L-Mannopyranoside: *Veronamine*

[27894-67-3]

C₂₆H₃₅NO₉ 505.564


Alkaloid from *Thalictrum fendleri* (Ranunculaceae). Hypotensive agent (site of action unknown) prob. also a smooth muscle relaxant. Produces bradycardia. Noncryst. [α]_D -155 (MeOH).

Shamma, M. *et al.*, *Tet. Lett.*, 1965, 3595; 1969, 4951 (*uv, pmr, cmr, ms, struct, synth, Thalifenderine, Veronamine*)

Shamma, M. *et al.*, *Tetrahedron*, 1971, **27**, 727 (*Veronamine*)

Thaliglicinone**T-327**

3-[2-(Dimethylamino)ethyl]-1-methoxy-10H-[1,3]dioxolo[6,7]phenanthro[4,5-bcd]pyran-10-one, 9CI [35988-96-6]

C₂₁H₁₉NO₅ 365.385

Aporphine-derived numbering shown. Alkaloid from *Thalictrum polygamum*, *Thalictrum rugosum* and other *Thalictrum* spp. (Ranunculaceae). Shows antibacterial and hypotensive activity. Yellow cryst. (MeOH or Et₂O). Mp 197° (180-182°, 190-191°). A dimorphic form with Mp 115-116° also reported. λ_{\max} 238 ; 256 ; 264 ; 287 ; 322 ; 390 (MeOH) (Berdy).

N-Me: *Thaliglicinone N-methosalt*

[53416-40-3]

C₂₂H₂₂NO₅[⊕] 380.419

Alkaloid from *Thalictrum polygamum* (Ranunculaceae). Mp 274-274° (as

chloride). CAS no. refers to chloride.

Deoxo: *Thaliglicine*. 1-Methoxy-N,N-dimethyl-10H-[1,3]dioxolo[6,7]phenanthro[4,5-bcd]pyran-3-ethanamine, 9CI. *Thalphenine methine* [36018-39-0]

C₂₁H₂₁NO₄ 351.401

Alkaloid from *Thalictrum polygamum* and *Thalictrum rugosum* (Ranunculaceae). Cryst. (EtOH). Mp 122°.

Deoxo, N-Me: *Thaliglicine N-methosalt*

[53416-39-0]

C₂₂H₂₄NO₄[⊕] 366.436

Alkaloid from *Thalictrum polygamum* (Ranunculaceae). CAS no. refers to chloride.

3-Methoxy: *Thalflavidine*

[66834-83-1]

C₂₂H₂₁NO₆ 395.411

Alkaloid from *Thalictrum flavum*, *Thalictrum minus* and *Thalictrum revolutum* (Ranunculaceae). Cryst. (MeOH). Mp 229-230° (219-220°).

2-Demethoxy, 3-methoxy: *Thalicsine*.

Thalixine

[50657-28-8]

C₂₁H₁₉NO₅ 365.385

Alkaloid from the above-ground parts of *Thalictrum longipedunculatum* (Ranunculaceae). Mp 193-194°. Doubtful structural assignment; all natural aporphines are substituted at C-1 and C-2. λ_{\max} 237 (log ϵ 4.22); 265 (log ϵ 4.48); 313 (log ϵ 3.96); 390 (log ϵ 3.6) (EtOH).

Doskotch, R.W. *et al.*, *J. Nat. Prod.*, 1969, **32**, 29-35 (*isol, uv, ir*)

Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1971, **24**, 1047-1050; *CA*, **76**, 85970h

(*Thaliglicine, Thaliglicinone*)

Shamma, M. *et al.*, *Chem. Comm.*, 1972, 408-409 (*Thaliglicine, uv, pmr, ms, cryst struct*)

Mitscher, L.A. *et al.*, *J. Nat. Prod.*, 1972, **35**, 167-176 (*isol*)

Khodzhaev, V.G. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 441; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **9**, 421-422 (*Thalicsine*)

Shamma, M. *et al.*, *Heterocycles*, 1974, **2**, 427-431 (*methosalts*)


Shamma, M. *et al.*, *Tetrahedron*, 1974, **30**, 2279-2282 (*synth*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 65-75; 204-212; 1977, **40**, 384-394; 508-514; 1980, **43**, 567-570 (*Thaliglicinone, Thalflavidine*)

Umarov, Kh.S. *et al.*, *CA*, 1978, **89**, 39379 (*Thalflavidine*)

Thalihazine**T-328**

1,2,3-Trimethoxy-N,N-dimethylphenanthro[2,3-d][1,3]dioxole-4-ethanamine, 9CI. 1-(2-Dimethylaminoethyl)-2,3,4-trimethoxy-6,7-methylenedioxyphenanthrene [111537-36-1]



C₂₂H₂₅NO₅ 383.443

Alkaloid from the whole plant of *Thalictrum hazarica* (Ranunculaceae). λ_{\max} 261 (log ϵ 4.3); 283 (log ϵ 3.77); 315 (log ϵ 3.37); 344 (log ϵ 2.9) (MeOH).

N-Oxide: **Thalihazine N-oxide**

[177413-37-5]

C₂₂H₂₅NO₆ 399.443

Alkaloid from aerial parts of *Thalictrum simplex*.

N-De-Me: 1-(2-Methylaminoethyl)-2,3,4-trimethoxy-6,7-methylenedioxyphenanthrene. 1,2,3-Trimethoxy-N-methylphenanthro[2,3-d][1,3]dioxole-4-ethanamine. **Secophoebine**

[109175-38-4]

C₂₁H₂₃NO₅ 369.416

Alkaloid from the leaves of *Phoebe valeriana* (Lauraceae). Amorph. solid. λ_{\max} 234 (sh); 262; 284; 304; 317; 344; 362 (EtOH).


Castro, O. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1036-1040 (*N-de-Me*)

Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 757-758 (*isol, uv, pmr, ms, struct*)

Velcheva, M.P. *et al.*, *Phytochemistry*, 1996, **42**, 535-537 (*oxide*)

Thalimicrinone**T-329**

[84716-73-4]

C₂₀H₁₉NO₅ 353.374


Alkaloid from the leaves of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Needles (EtOH). Mp 130-132°.

Başer, K.H.C. *et al.*, *J. Nat. Prod.*, 1982, **45**, 704 (*isol, uv, ir, pmr, ms, struct*)

Al-Khalil, S. *et al.*, *J. Nat. Prod.*, 1985, **48**, 989 (*synth*)

Thalimonine**T-330**

2,8,9-Trimethoxy-3,4-methylenedioxy-pavine
[142735-70-4]

C₂₁H₂₃NO₅ 369.416

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae). $[\alpha]_D^{22}$ -118 (c, 0.20 in MeOH). λ_{\max} 287 (ϵ 6920) (MeOH) (Berdy).

N-Oxide: **Thalimonine N-oxide A**

[166266-74-6]

C₂₁H₂₃NO₆ 385.416

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae). $[\alpha]_D^{21}$ -53.3 (c, 0.05 in MeOH).

N-Oxide, epimer: **Thalimonine N-oxide B**
[166376-54-1]

C₂₁H₂₃NO₆ 385.416

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae). $[\alpha]_D^{21}$ -80 (c, 0.02 in MeOH).

O²-De-Me: 2-Hydroxy-8,9-dimethoxy-3,4-methylenedioxy-pavine. **2-De-methylthalimonine**

[150036-90-1]

C₂₀H₂₁NO₅ 355.39

Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae). $[\alpha]_D^{22}$ -43.8 (c, 0.02 in MeOH).

O⁹-De-Me: 9-Hydroxy-2,8-dimethoxy-3,4-methylenedioxy-pavine. **9-De-methylthalimonine**

[150036-91-2]

C₂₀H₂₁NO₅ 355.39


Alkaloid from aerial parts of *Thalictrum simplex* (Ranunculaceae). $[\alpha]_D^{22}$ -51.7 (c, 0.067 in MeOH).

Velcheva, M. *et al.*, *J. Nat. Prod.*, 1992, **55**, 679-680 (*isol, uv, ir, cd, pmr, cmr, ms, struct*)

Velcheva, M.P. *et al.*, *Planta Med.*, 1993, **59**, 262-263 (2-Demethylthalimonine, 9-Demethylthalimonine)

Velcheva, M.P. *et al.*, *Phytochemistry*, 1995, **39**, 683-687 (*Thalimonine N-oxides*)

Pabuccuoglu, V. *et al.*, *Heterocycles*, 1997, **45**, 1751-1758 (*synth*)

Thaliphine**T-331**C₄₁H₄₄N₂O₁₁ 740.805

Alkaloid from the roots and rhizomes of *Isopyrum thalictroides*. Amorph. solid. $[\alpha]_D$ -46.3 (c, 0.4 in CHCl₃). λ_{\max} 210 (log ϵ 8); 234 (sh) (log ϵ 7.61); 262 (log ϵ 7.02); 293 (log ϵ 7.32) (EtOH).


Diastereoisomer: **Isotaliphine**C₄₁H₄₄N₂O₁₁ 740.805

Alkaloid from the roots and rhizomes of *Isopyrum thalictroides*. Amorph. solid. $[\alpha]_D$ -51.8 (c, 0.3 in CHCl₃). λ_{\max} 210 (log ϵ 7.93); 234 (sh) (log ϵ 7.53); 262 (log ϵ 6.96); 293 (log ϵ 7.18) (EtOH).

Istatkova, R. *et al.*, *Nat. Prod. Res.*, 2004, **18**, 259-263 (*isol, cd, pmr, cmr, ms*)

Thaliracebine**T-332**

[67591-63-3]

C₃₉H₄₄N₂O₇ 652.786

Tentative struct. The 5-methoxy-6,7-methylenedioxy substitution pattern may

need revision to 5,6-methylenedioxy-7-methoxy following the structural revision of Thalflavine, T-306. Alkaloid from the roots of *Thalictrum minus* Race B, and from *Thalictrum faberi* (Ranunculaceae). Antiseptic, antihypertensive agent. Amorph. Mp 83-84°. $[\alpha]_D^{26}$ +121 (c, 0.28 in MeOH). Log P 6.83 (uncertain value) (calc). λ_{\max} 278 (MeOH) (Berdy).

*Methiodide (1:2):*Beige amorph. solid (MeOH/Et₂O).

Mp 203-205° dec.


Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257 (*isol, uv, cd, ir, pmr, ms, struct*)

Aly, Y. *et al.*, *Phytochemistry*, 1989, **28**, 1967

Thalirevolutine**T-333**

O-Methylfetidine

[62751-64-8]

C₄₁H₄₈N₂O₈ 696.839

Alkaloid from the roots of *Thalictrum revolutum* (Ranunculaceae). Hypotensive agent in rabbits. Cryst. + 1/2 H₂O (Et₂O). Mp 105-108°. $[\alpha]_D^{20}$ +134 (c, 0.1 in MeOH).

O¹-De-Me: **Fetidine. Foetidine†**

[7072-86-8]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from the aerial parts of *Thalictrum foetidum* (Ranunculaceae). Antihypertensive agent, nervous depressant in mice. Shows strong anti-inflammatory activity. Cryst. (EtOAc). Mp 132-135° (125-126°). $[\alpha]_D^{15}$ +121.4 (c, 2.57 in MeOH). Log P 5.82 (uncertain value) (calc).

O¹-De-Me, hydrochloride: Mp 228-230° dec. $[\alpha]_D^{20}$ -30.9 (c, 2.52 in H₂O).

O¹²-De-Me: **Thalirevoline**

[65853-12-5]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from the roots, tops and fruits of *Thalictrum revolutum* (Ranunculaceae). Hypotensive agent in rabbits. Shows antibiotic activity against *Mycobacterium smegmatis*. Cryst. + 1/2 H₂O (Et₂O). Mp 123-125°. $[\alpha]_D^{20}$ +95 (c, 0.1 in MeOH).

O⁷-De-Me: **Faberidine. 3-Demethoxyfaberonine**

[91925-99-4]

C₄₀H₄₆N₂O₈ 682.812

Alkaloid from the roots of *Thalictrum faberi* (Ranunculaceae). Amorph. solid. $[\alpha]_D^{22}$ +105.5 (c, 0.675 in MeOH).

O⁷, O¹²-Di-de-Me: **Revolutopine**

[62724-07-6]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from the tops of *Thalictrum revolutum* (Ranunculaceae). Powder (C₆H₆/hexane). $[\alpha]_D^{25}$ +126 (c, 0.1 in MeOH).


Sargazalov, D. *et al.*, *Dokl. Akad. Nauk SSSR*,

- Ser. Khim.*, 1963, **20**, 28; *CA*, **59**, 15336h (isol, Fetidine)
 Ismailov, Z.F. *et al.*, *Khim. Prir. Soedin.*, 1966, **2**, 43; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **2**, 35 (isol, Fetidine)
 Cava, M.P. *et al.*, *Tet. Lett.*, 1972, 2309 (pmr, struct, Fetidine)
 Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 593 (Revolutopine)
 Wu, W.N. *et al.*, *Tetrahedron*, 1977, **33**, 2919 (Thalirevoline, Thalirevolutine)
 Wagner, H. *et al.*, *Tetrahedron*, 1984, **40**, 2133 (Faberidine)

Thalirugidine

T-334

[64215-95-8]

 $C_{39}H_{46}N_2O_8$ 670.801

Alkaloid from the roots of *Thalictrum rugosum* and *Thalictrum foliolosum* (Ranunculaceae). Antiseptic. Amorph. powder. $[\alpha]_D^{20} +112$ (c, 0.19 in MeOH). Log P 5.34 (calc).

Di-Me ether:

Pale yellow amorph. solid. $[\alpha]_D^{20} +53$ (c, 0.21 in MeOH).

Di-Et ether:


Amorph. $[\alpha]_D^{20} +44$ (c, 0.17 in MeOH).

- Wu, W.-N. *et al.*, *J.O.C.*, 1978, **43**, 580 (isol, uv, cd, ir, pmr, ms, struct)
 Chattopadhyay, S.K. *et al.*, *J. Nat. Prod.*, 1981, **44**, 45 (isol, uv, ir, pmr, ms)

Thalirugine

T-335

[64235-41-2]

 $C_{38}H_{44}N_2O_7$ 640.775

Alkaloid from the roots of *Thalictrum rugosum* (Ranunculaceae). Antiseptic. Amorph. solid + $\frac{1}{2}$ H₂O. $[\alpha]_D^{20} +92$ (c, 0.25 in MeOH). Log P 5.87 (calc).

7-Me ether: Thaliruginine

[64215-93-6]

 $C_{39}H_{46}N_2O_7$ 654.802

Alkaloid from the roots of *Thalictrum rugosum* (Ranunculaceae). Amorph. solid + 1H₂O. $[\alpha]_D^{20} +104$ (c, 0.16 in MeOH).

Di-Me ether:


Pale-yellow amorph. solid. $[\alpha]_D^{21} +105$ (c, 0.2 in MeOH).

- Wu, W.-N. *et al.*, *J.O.C.*, 1978, **43**, 580 (isol, uv, cd, ir, pmr, ms, struct)

Thalisopavine

T-336

10,11-Dihydro-3,7,8-trimethoxy-12-methyl-10,5-(iminomethano)-5H-dibenzo[a,d]cyclohepten-2-ol, 9CI
 [18927-72-5]
 [18927-73-6 ((±)-form)]

 $C_{20}H_{23}NO_4$ 341.406

Biogenetic (benzylisoquinoline-type) numbering shown. Alkaloid from the roots of *Thalictrum dasycarpum* (Ranunculaceae). Needles (EtOH). Mp 211-212°. $[\alpha]_D^{25} -210$ (c, 0.21 in CHCl₃).

Me ether: O-Methylthalisopavine

[18944-92-8]

[33579-95-2 ((±)-form)]

 $C_{21}H_{25}NO_4$ 355.433

Alkaloid from *Papaver radicans* (*Papaver nudicaule* var. *radicans*) (Papaveraceae). Cryst. (diisopropyl ether); needles (EtOH/Et₂O). Mp 92-94°.

O⁷-De-Me: Thalidine

[63110-83-8]

[63162-85-6 ((±)-form)]

 $C_{19}H_{21}NO_4$ 327.379

Alkaloid from *Thalictrum dioicum* (Ranunculaceae). Cryst. (MeOH). Mp 205-207° (natural) Mp 222-223° (synthetic). $[\alpha]_D^{25} -172$ (c, 0.7 in MeOH) (natural).

O⁷, O¹²-Di-de-Me, O¹¹-Me: Thalidicine

[61774-85-4]

 $C_{19}H_{21}NO_4$ 327.379

Alkaloid from the roots of *Thalictrum dioicum* (Ranunculaceae). Cryst. (EtOH/Et₂O). Mp 200°. Structural determination incomplete. May be identical with Thalidine.

Kupchan, S.M. *et al.*, *J.O.C.*, 1969, **34**, 1062-1065 (*Thalisopavine*, isol, uv, ir, pmr, ms, struct, synth)

Dyke, S.F. *et al.*, *Tetrahedron*, 1971, **27**, 3803-3809 (*O-Methylthalisopavine*, synth, pmr)

Hoshino, O. *et al.*, *Heterocycles*, 1973, **1**, 223-226 ((±)-*O-Methylthalisopavine*, synth)

Böhm, V.H. *et al.*, *Planta Med.*, 1975, **28**, 210 (deriv, isol)

Ong, H. *et al.*, *Ann. Pharm. Fr.*, 1976, **34**, 223 (*Thalidicine*)

Shamma, M. *et al.*, *J. Nat. Prod.*, 1976, **39**, 395 (*Thalidine*)

Elliott, I.W. *et al.*, *J.O.C.*, 1979, **44**, 1162-1163 ((+)-*O-Methylthalisopavine*, synth, ms)

Maffrand, J.-P. *et al.*, *Heterocycles*, 1980, **14**, 325 (deriv, synth)


Rice, K.C. *et al.*, *J.O.C.*, 1980, **45**, 601 (*O-Methylthalisopavine*, *Thalidine*, synth, ms)

Carrillo, L. *et al.*, *J.O.C.*, 1997, **62**, 6716-6721 ((-)-*O-Methylthalisopavine*, synth)

Thalistine

T-337

[75352-25-9]

 $C_{39}H_{44}N_2O_8$ 668.785

Probable struct. The structs. of all the alkaloids covered by this entry were tentatively revised from 5-methoxy-6,7-methylenedioxy to 7-methoxy-5,6-methylenedioxy in 1989, and this has not subsequently been challenged. Alkaloid from the roots of *Thalictrum minus* Race B (Ranunculaceae). Antiseptic. Amorph. $[\alpha]_D^{20} +104$ (c, 0.35 in MeOH). Log P 6.03 (uncertain value) (calc). λ_{max} 278 (ε 8000) (MeOH) (Berdy).

N²-Me: Thalirabine. 5-O-Demethylthalistyline

[67624-63-9]

 $C_{40}H_{47}N_2O_8^{\oplus}$ 683.82

Alkaloid from the roots of *Thalictrum minus* Race B (Ranunculaceae). Antiseptic. λ_{max} 207; 276; 283 (MeOH) (Berdy). λ_{max} 276; 283 (HCl) (Berdy). λ_{max} 209; 273; 280 (CHCl₃) (Berdy).

N²-Me, hydroxide:

Beige-coloured amorph. solid. Mp 131-132° dec. $[\alpha]_D^{26} +142$ (c, 0.548 in MeOH).

N²-Me, iodide:

Tan-coloured rosettes (MeOH aq.). Mp 205-206° dec.

O⁵-Me: N-Demethylthalistyline

[62251-51-8]

 $C_{40}H_{46}N_2O_8$ 682.812

Alkaloid from the roots of *Thalictrum longistylum* and *Thalictrum podocarpum* (Ranunculaceae). Antiseptic, antihypertensive agent. Pale yellow amorph. powder. $[\alpha]_D^{25} +151$ (c, 0.2 in MeOH). Log P 6.26 (uncertain value) (calc). λ_{max} 282 (ε 8000) (MeOH) (Berdy).

O⁵-Me, N²-Me: Thalistyline. O-Methylthalirabine

[62251-53-0]

 $C_{41}H_{49}N_2O_8^{\oplus}$ 697.847

Alkaloid from roots of *Thalictrum longistylum* and *Thalictrum podocarpum* (Ranunculaceae). Antiseptic, antihypertensive agent. Shows antibacterial activity. λ_{max} 276 (ε 7300); 283 (ε 7000) (MeOH) (Berdy).

O⁵-Me, N²-Me, chloride: [62251-48-3]

Mp 150-153°. $[\alpha]_D^{25} +146$ (c, 0.1 in MeOH).

O⁵-Me, N²-Me, iodide: [62251-49-4]

Mp 220-223° dec.

O⁵-Me, N², N²-di-Me: Methothalistyline. 2'-N-Methylthalistyline

[65853-13-6]

 $C_{42}H_{52}N_2O_8^{\oplus}$ 712.881

Alkaloid from roots of *Thalictrum longistylum* and *Thalictrum podocarpum* (Ranunculaceae). Shows antibacterial activity. Antiseptic. Needles + 1H₂O (MeOH) (as diiodide). Mp 266-268° dec. (diiodide). $[\alpha]_D^{21} +125$ (c, 0.1 in MeOH).

Wu, W.-N. *et al.*, *Tet. Lett.*, 1976, 3687-3690 (*Thalistyline*)


Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1977, **40**, 281-289; 384-394 (*Demethylthalistyline*, *Thalistyline*, *Methothalistyline*, activity)

Liao, W.-T. *et al.*, *J. Nat. Prod.*, 1978, **41**, 257-270 (*Thalirabine*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1980, **43**, 472-481 (*isol, uv, cd, ir, pmr, ms, struct, synth*)
 Aly, Y. *et al.*, *Phytochemistry*, 1989, **28**, 1967-1971 (*struct*)

Thallusin**T-338**

[851369-86-3]



Relative Configuration


C₂₅H₃₁NO₇ 457.522

Isol. from a marine bacterium *Cytophaga* sp. YM2-23 obt. from a *Monostroma* sp. Morphogenesis inducer in algae. Amorph. powder.

Matsuo, Y. *et al.*, *Science (Washington, D.C.)*, 2005, **307**, 1598 (*isol, pmr, cmr, cryst struct*)
 Gao, X. *et al.*, *Org. Lett.*, 2006, **8**, 2123-2126; 2007, **9**, 379 (*synth*)
 Nishizawa, M. *et al.*, *Tet. Lett.*, 2007, **48**, 4229-4233 (*synth*)

Thalmethine**T-339**

[3729-83-7]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the above-ground parts of *Thalictrum minus* (Ranunculaceae). Cryst. (MeOH). Mp 275-277°. [α]_D²¹ +200 (c, 1 in CHCl₃).

Me ether: O-Methylthalmethine

[5979-99-7]

C₃₇H₃₈N₂O₆ 606.717

Alkaloid from the above-ground parts of *Thalictrum minus* and from the roots of *Thalictrum revolutum* (Ranunculaceae). Antiseptic. Needles (C₆H₆). Mp 245-246°. [α]_D²¹ +237 (c, 1 in CHCl₃). Log P 7.64 (uncertain value) (calc).

*O*⁷-*De-Me*, *O*¹²-*Me: Thalsivasine. Thalcultimine*

[101219-59-4]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the above-ground parts of *Thalictrum minus* var. *minus* (Ranunculaceae).

Mollov, N.M. *et al.*, *Chem. Ind. (London)*, 1965, 1595 (*struct*)

Duchevska, Kh.B. *et al.*, *Dokl. Bulg. Akad. Nauk*, 1971, **24**, 467; *CA*, **75**, 106101x, (*isol*)


Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 597 (*ms*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1977, **40**, 508

Başer, K.H.C. *et al.*, *Planta Med.*, 1985, 448 (*Thalsivasine*)

Thalmiculine**T-340**

[106146-69-4]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D -35 (c, 2.2 in MeOH).

*N*²-*De-Me*, *1',2'*-*didehydro: Thalmiculimine*

[105437-18-1]

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D -5 (c, 0.09 in MeOH).

*N*²-*De-Me*, *1',2'*-*didehydro, O*¹²-*de-Me: Cultithalminine*

[112448-46-1]

C₃₆H₃₆N₂O₇ 608.69

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D +7 (c, 0.17 in MeOH).

*N*²-*De-Me*, *1',2'*-*didehydro, 5-deoxy, O*¹²-*de-Me: Thalmiculatimine*

[105418-71-1]

C₃₆H₃₆N₂O₆ 592.69


Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D +7.5 (c, 0.093 in MeOH).

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1986, **49**, 488-493 (*Thalmiculine, Thalmiculimine, Thalmiculatimine, uv, pmr, ms, cd, struct*)

Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 721-725 (*Cultithalminine*)

Thalmine**T-341**

Talmine
 [7682-65-7]

C₃₇H₄₀N₂O₆ 608.733

These alkaloids (*S,S*-series) are diastereomeric with those covered by the entries Lauberine, L-67 (*2R,2'S*-) and Dryadodaphnine, D-944 (*2S,2'R*-). Alkaloid from *Thalictrum minus* and *Thalictrum kuhistanicum* (Ranunculaceae). Shows strong antiinflammatory activity. Significant activity against ascites lymphoma in mice and rats. Cryst. (EtOH/CHCl₃). Mp 253° dec. [α]_D -64.5. Log P 8.12 (uncertain value) (calc).

Hydrochloride:

Cryst. (EtOH). Mp 147-157°.

*N*²-*De-Me: 2-Northalmine*

[101488-79-3]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D²⁵ -31.8 (c, 0.43 in MeOH).

Me ether: O-Methylthalmine

[7682-67-9]

C₃₈H₄₂N₂O₆ 622.76

Alkaloid from *Thalictrum sultanabadense* and *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D -43 (MeOH).

*O*¹²-*De-Me: Thalbadenzine. Thalbadensine*

[66834-86-4]

C₃₆H₃₈N₂O₆ 594.706

Alkaloid from the aerial parts of *Thalictrum sultanabadense*, and from *Thalictrum minus* (Ranunculaceae). Amorph. No opt. rotn. reported.

*O*¹²-*De-Me, O*^{6'}-*Me: Thalicine*

[58092-24-3]

C₃₇H₄₀N₂O₆ 608.733

Alkaloid from stems and leaves of *Thalictrum thunbergii* (Ranunculaceae). Needles +1.5 H₂O (as nitrate salt). Mp 226-228° (nitrate). [α]_D²⁸ -15.8 (c, 1.203 in CHCl₃). [α]_D²⁸ -12.2 (c, 1.639 in MeOH). Rapidly oxid. in air.

5-Hydroxy: 5-Hydroxythalmine

[105418-73-3]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). [α]_D -69 (c, 0.08 in MeOH).

Yunusov, S. *et al.*, *Zh. Obshch. Khim.*, 1950, **20**, 1151; *CA*, **45**, 1608c (*isol*)

Telezhenetskaya, M.V. *et al.*, *Khim. Prir. Soedin.*, 1966, **2**, 107; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **2**, 83 (*struct*)

Baldas, J. *et al.*, *Tet. Lett.*, 1968, 6315 (*pmr, ms, struct*)

Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 705; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 715 (*ord*)

Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 597 (*ms*)
 Tomimatsu, T. *et al.*, *Chem. Pharm. Bull.*, 1975, **23**, 2279 (*Thalicine*)

Abdizhabbarova, S. *et al.*, *Khim. Prir. Soedin.*, 1978, **14**, 139; *Chem. Nat. Compd. (Engl. Transl.)*, 1978, **14**, 114 (*Thalbadenzine, isol, pmr, ms, struct*)


Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 818; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 723 (*cd, abs config*)

Mukhamedova, S. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 397; *CA*, **102**, 929037m (*O-Methylthalmine*)

Hussain, S.F. *et al.*, *J. Nat. Prod.*, 1985, **48**, 962; 1986, **49**, 488 (*Northalmine, 5-Hydroxythalmine*)

Thalmineline**T-342**

[28328-00-9]

C₄₂H₅₀N₂O₁₀ 742.864

Alkaloid from the roots of *Thalictrum minus* (Ranunculaceae). Cryst. (EtOH) or

Et₂O/heptane). Mp 108-110° (96-98°).

Reisch, J. *et al.*, *Tet. Lett.*, 1970, 2113 (*uv, ir, pmr, struct*)

Borkowski, B. *et al.*, *J. Chromatogr.*, 1971, **59**, 222 (*chromatog*)

Thalmirabine T-343

[75352-27-1]

As Thalfinine, T-305 with

R¹ = R² = Me, R³ = H


C₃₉H₄₄N₂O₈ 668.785

Alkaloid from the roots of *Thalictrum minus* (Ranunculaceae) Race B. Antiseptic; shows antibacterial activity. Amorph. solid + 1.5 H₂O. [α]_D +116 (c, 0.2 in MeOH). Log P 7.01 (uncertain value) (calc). λ_{max} 280 (ε 9000) (MeOH) (Berdy).

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1980, **43**, 472 (*uv, ir, pmr, ms, cd, struct, activity*)

Thalpetaline T-344

3,4-Dihydro-7-methoxy-2-methyl-5,6-methylenedioxyisoquinolinium(1+)




C₁₂H₁₄NO₃⁺ 220.248

Quaternary alkaloid from *Thalictrum petaloideum* var. *supradecompositum*. Amorph. (as iodide).

Al-Rehaily, A.J. *et al.*, *Planta Med.*, 1998, **64**, 681 (*isol, pmr*)

Thalphenine T-345

4,5,5a,11-Tetrahydro-1-methoxy-5,5-dimethyl-3H,6H-[1,3]benzodioxolo[4',5',6':4,5][2]benzopyrano[1,8,7-hi]isoquinolinium(1+), 9CI



(S)-form

C₂₁H₂₂NO₄⁺ 352.409

(S)-form [39027-72-0]

Quaternary alkaloid from *Thalictrum polygamum*, the roots of *Thalictrum minus* Race B, *Thalictrum rugosum* and *Thalictrum revolutum*, and from the stem bark of *Phellodendron wilsonii*. Shows hypotensive action in rabbits. Weak antibacterial activity vs. *Staphylococcus aureus*, *Mycobacterium smegmatis* and *Candida albicans*.

Chloride: [39027-97-9]

Needles (Me₂CO/MeOH). Mp 186-188°. [α]_D²⁵ +71 (c, 0.28 in MeOH).

Iodide: [39005-06-6]

Cryst. + 2H₂O (Me₂CO aq.). Mp 198-199°.

N-De-Me: N-Demethylthalphenine

[63640-46-0]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the tops and fruits of *Thalictrum revolutum* (Ranunculaceae). Shows *in vitro* antibacterial activity vs. *S. aureus*, *M. smegmatis* and *C. albicans*. Needles (MeOH or C₆H₆/petrol). Mp 179.5-180.5°. [α]_D²⁵ +104 (c, 0.18 in MeOH).

Di-N-de-Me: Bisnorthalphenine

[53416-38-9]

C₁₉H₁₇NO₄ 323.348

Alkaloid from the tops of *Thalictrum revolutum* and from *Thalictrum polygamum* (Ranunculaceae). Shows weak antibacterial activity vs. *S. aureus*, *M. smegmatis* and *C. albicans*. Mp 124-125°. [α]_D²⁵ +81 (c, 1.12 in MeOH).

(±)-form [51744-23-1]

Synthetic. Cryst. + 1MeOH (MeOH)(as iodide). Mp 193-194° (iodide). CAS no. refers to iodide.

N-De-Me: [51665-96-4]

Synthetic. Cryst. (MeOH). Mp 179-180°.

Shamma, M. *et al.*, *Chem. Comm.*, 1972, 408 (*uv, pmr, ms, cryst struct*)

Shamma, M. *et al.*, *Heterocycles*, 1974, **2**, 427 (*uv, pmr, ms, struct, Bisnorthalphenine*)


Shamma, M. *et al.*, *Tetrahedron*, 1974, **30**, 2279 (*synth*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 65; 249; 1977, **40**, 508; 1980, **43**, 472 (*isol, uv, ir, pmr, ms*)

Wu, J. *et al.*, *J. Nat. Prod.*, 1977, **40**, 294 (*isol, uv, pmr, derivs*)

Thalgugosaminine T-346

5,6,6',7,12'-Pentamethoxy-2,2'-dimethylloxycanthan, 9CI. 5-O-Methylthalisopine. 12'-O-Methylthalisopine. O,O-Dimethylthalisopidine [22226-73-9]



C₃₉H₄₄N₂O₇ 652.786

Alkaloid from the roots of *Thalictrum revolutum*, *Thalictrum rugosum*, *Thalictrum foliolosum*, *Thalictrum alpinum* and *Thalictrum minus* Race B. Also detected in *Thalictrum isopyroides* (Ranunculaceae). Antiseptic, antihypertensive agent. Shows *in vitro* activity against *Mycobacterium smegmatis*. Amorph. solid + 0.5H₂O. Mp 103-105°. [α]_D²⁵ -90.4 (c, 0.104 in MeOH). Log P 7.83 (uncertain value) (calc). λ_{max} 262 (ε 6310); 282 (ε 7600) (MeOH) (Berdy). λ_{max} 205; 282 (EtOH) (Berdy).

N²-β-Oxide: *Thalgugosaminine 2'β-N-*

oxide

[112448-43-8]

C₃₉H₄₄N₂O₈ 668.785

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D -33 (c, 0.2 in MeOH). Struct. in paper not drawn according to Shamma-Moniot convention; called in the paper the 2α-N-oxide.

O⁵-De-Me: *Thalisopine*. *Thaligosine*. 5-O-Demethylthalgugosaminine. *Talisopine* [22226-72-8]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from the seeds of *Thalictrum isopyroides*, the roots of *Thalictrum rugosum* and *Thalictrum foliolosum*, and from the subterranean parts of *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Antiseptic, anti-rhythmic agent. Nervous depressant in mice. Cryst. + 1H₂O (Et₂O). Mp 143-145°. [α]_D -109 (c, 0.17 in MeOH). Log P 7.54 (uncertain value) (calc). λ_{max} 282 (ε 7250) (MeOH) (Berdy). λ_{max} 282 (ε 7600); 305 (ε 4360) (NaOH) (Berdy).

O⁵-De-Me, N²-β-oxide: *Thaligosine 2'β-N-oxide*

C₃₈H₄₂N₂O₈ 654.758

Alkaloid from *Thalictrum cultratum* (Ranunculaceae). Amorph. [α]_D -59 (c, 0.13 in MeOH). Struct. in paper not drawn in standard orientation; alkaloid incorrectly designated as the 2α-N-oxide.

O¹²-De-Me: *Thaligosine*

[64235-38-7]

C₃₈H₄₂N₂O₇ 638.759

Alkaloid from the roots of *Thalictrum rugosum* (Ranunculaceae). Antiseptic. Cryst. (Et₂O). Mp 233-234.5°. [α]_D²¹ -58.5 (c, 0.316 in MeOH). Log P 7.25 (uncertain value) (calc).

O⁵, O¹²-Di-de-Me: *Thalisopidine*

[26989-49-1]

C₃₇H₄₀N₂O₇ 624.732

Alkaloid from the seeds of *Thalictrum isopyroides* (Ranunculaceae). Mp 215-216°. [α]_D¹⁹ -9 (EtOH).

Ismailov, Z.F. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1963, **20**, 21; *CA*, **61**, 4407g (*Thalisopine, struct*)

Pulatova, Kh.G. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 394; 1969, **5**, 609; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 336; 1969, **5**, 533 (*Thalgugosaminine, Thaliosopine, Thaliosopidine, isol, struct*)

Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 705; 1979, **15**, 818; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 715; 1979, **15**, 723 (*pmr, ms, cd, abs config*)

Shamma, M. *et al.*, *Heterocycles*, 1976, **4**, 1817 (*abs config*)

Wu, W.-N. *et al.*, *J. Nat. Prod.*, 1976, **39**, 65; 1977, **40**, 508 (*isol, uv, cd, pmr, ms, config, struct*)

Wu, W.N. *et al.*, *J.O.C.*, 1978, **43**, 580


(*Thalisopine, Thaliosopine, isol, uv, cd, ir, pmr, ms, struct*)

Chatopadhyay, S.K. *et al.*, *J. Nat. Prod.*, 1981, **44**, 45 (*Thalisopine, Thalgugosaminine, isol, uv, cd, ir, pmr, ms*)

Herath, W.H.M.W. *et al.*, *J. Nat. Prod.*, 1987, **50**, 721 (*Thaliosopine N-oxide, Thalgugosaminine N-oxide*)


ThalsimidineThalimidine
[22223-14-9]

T-347

C₃₇H₃₈N₂O₇ 622.716Alkaloid from the aerial parts of *Thalictrum simplex* (Ranunculaceae). Mp 195°. [α]_D²⁰ +48 (CHCl₃).*Me ether*: **Thalsimine**. Thalimine. O-Methylthalsimidine
[5525-36-0]C₃₈H₄₀N₂O₇ 636.743Alkaloid from the aerial parts of *Thalictrum rugosum* and roots of *Thalictrum rochebrunianum* (Ranunculaceae). Anti-inflammatory, antitussive agent. Shows weak antineoplastic activity. Inhibitor of conditioned avoidance reactions in expt. animals. Mp 149–150°. [α]_D²⁰ +22.6 (c, 0.7 in CHCl₃). Log P 7.18 (uncertain value) (calc). Exists in soln. as a 1:1 mixt. of conformers.Maekh, S.Kh. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 188; 1968, **4**, 138; 393; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 144; 1968, **4**, 119; 335 (*isol, uv, ms, struct*)Mollov, N.M. *et al.*, *Dokl. Bulg. Akad. Nauk.*, 1966, **19**, 491; 1967, **20**, 329; 1970, **23**, 181; *CA*, **65**, 13780d; **67**, 61598f; **73**, 32285v (*isol, deriv*)Ismailov, Z.F. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 256; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 220 (*ms, deriv*)Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 705; 1979, **15**, 818; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 715; 1979, **15**, 723 (*ord, cd, abs config*)Saá, J.M. *et al.*, *Tet. Lett.*, 1976, 513 (*pmr, deriv*)**Thapoxepine A**

[250161-96-7]

T-348

C₃₆H₃₈N₂O₁₀ 658.704Isol. as a C-13 epimeric mixt. Alkaloid from the roots of *Aglaia edulis*. [α]_D²⁰ +55 (c, 0.6 in CHCl₃). λ_{max} 231 (sh) (log ε 4.38); 249 (sh) (log ε 4.19); 280 (log ε 3.9); 297 (sh) (log ε 3.78); 335 (log ε 3.38)

(MeOH).

7,8-Diepimer: Thapoxepine B

[250278-72-9]


C₃₆H₃₈N₂O₁₀ 658.704Alkaloid from *Aglaia edulis*. Mp 260–262°. [α]_D²⁰ +33 (c, 0.5 in CHCl₃). Isol. as a C-13 epimeric mixt. λ_{max} 231 (sh) (log ε 4.23); 249 (sh) (log ε 4.04); 281 (log ε 3.81); 297 (sh) (log ε 3.73); 336 (log ε 3.43) (MeOH).*Homologue (R = CH₂CH₃): Homothapoxepine A*

[250161-98-9]

C₃₇H₄₀N₂O₁₀ 672.73Alkaloid from *Aglaia edulis*. [α]_D²⁰ +110 (c, 0.4 in CHCl₃). Possesses (13S)-config. λ_{max} 231 (sh) (log ε 4.42); 249 (sh) (log ε 4.19); 280 (log ε 3.9); 297 (sh) (log ε 3.78); 334 (log ε 3.48) (MeOH).Bacher, M. *et al.*, *Phytochemistry*, 1999, **52**, 253–263 (*isol, uv, ir, pmr, cmr, ms*)**Thapsakin B**

[250161-87-6]

T-349



Absolute Configuration

C₃₅H₃₈N₂O₉ 630.693Alkaloid from the roots of *Aglaia edulis*. [α]_D²⁰ +24 (c, 0.4 in CHCl₃). Isol. as a C-13 epimeric mixt. λ_{max} 252 (sh) (log ε 3.7); 271 (sh) (log ε 3.51); 282 (sh) (log ε 3.54); 298 (log ε 3.68) (MeOH).**10-Ketone: Thapsakone B**

[250161-93-4]

C₃₅H₃₆N₂O₉ 628.677Alkaloid from *Aglaia edulis*. [α]_D²⁰ +8 (c, 0.2 in CHCl₃). λ_{max} 279 (sh) (log ε 3.6); 304 (log ε 3.71) (MeOH).**6-Demethoxy, N¹³-deacyl, N¹³-(2ξ-methylbutanoyl), 10-Ac:** [251982-77-1]C₃₇H₄₀N₂O₉ 656.731Alkaloid from the stem bark of *Aglaia roxburghiana*.**6-Demethoxy, 19-hydroxy, N¹³-deacyl, N¹³-(2ξ-methylbutanoyl):** [251982-76-0]C₃₅H₃₈N₂O₉ 630.693Alkaloid from the stem bark of *Aglaia roxburghiana*. C-19 config. not determined.**10-Epimer: Isothapsakin B**

[250144-64-0]

C₃₅H₃₈N₂O₉ 630.693Alkaloid from *Aglaia edulis*. [α]_D²⁰ -51 (c, 0.2 in CHCl₃). λ_{max} 251 (sh) (log ε 3.68); 272 (sh) (log ε 3.45); 282 (sh) (log ε 3.53); 298 (log ε 3.64) (MeOH).**3,4-Diepimer, 10-Ac: 10-O-Acetylthapsakin A**

[250144-66-2]

C₃₇H₄₀N₂O₁₀ 672.73Alkaloid from *Aglaia edulis*. Mp 150–152°. [α]_D²⁰ -24 (c, 0.5 in CHCl₃). λ_{max} 251 (sh) (log ε 3.73); 272 (sh) (log ε 3.48); 282 (sh) (log ε 3.56); 298 (log ε 3.67) (MeOH).**3,4-Diepimer, 10-ketone: Thapsakone A**


[250161-91-2]

C₃₅H₃₆N₂O₉ 628.677Alkaloid from *Aglaia edulis*. [α]_D²⁰ +23 (c, 0.6 in CHCl₃). λ_{max} 280 (sh) (log ε 3.54); 304 (log ε 3.65) (MeOH).**3,4-Diepimer, N¹³-deacyl, N¹³-(2-methylbutanoyl): Homothapsakin A**

[250161-89-8]

C₃₆H₄₀N₂O₉ 644.72Alkaloid from *Aglaia edulis*. [α]_D²⁰ -135 (c, 0.3 in CHCl₃). λ_{max} 251 (sh) (log ε 3.76); 272 (sh) (log ε 3.51); 282 (sh) (log ε 3.58); 298 (log ε 3.69) (MeOH).**3,4-Diepimer, N¹³-deacyl, N¹³-(3-methylbutanoyl): Edulirin A**C₃₆H₄₀N₂O₉ 644.72Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. [α]_D²⁵ -148 (c, 0.2 in MeOH). C-13 config. not determined. λ_{max} 213 (log ε 4.61); 269 (log ε 3.27); 297 (log ε 3.52) (MeOH).**3,4-Diepimer, N¹³-deacyl, N¹³-(3-methylbutanoyl), 10-Ac: 10-O-Acetyledulirin A**C₃₈H₄₂N₂O₁₀ 686.757Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. [α]_D²² -71 (c, 0.1 in MeOH). C-13 config. not determined. λ_{max} 214 (log ε 4.59); 268 (log ε 3.31); 297 (log ε 3.55) (MeOH).**3,4-Diepimer, N¹³-deacyl, N¹³-(3-methyl-2-butenoyl): 19,20-Dehydroedulirin A**C₃₆H₃₈N₂O₉ 642.704Alkaloid from the bark of *Aglaia edulis*. Amorph. powder. [α]_D²² -46 (c, 0.1 in MeOH). C-13 config. not determined. λ_{max} 216 (log ε 4.57); 269 (log ε 3.39); 297 (log ε 3.53) (MeOH).Molloyres, L.-P. *et al.*, *Pestic. Sci.*, 1999, **55**, 486–503 (*Aglaia roxburghiana constits*)Bacher, M. *et al.*, *Phytochemistry*, 1999, **52**, 253–263 (*Thapsakins, Thapsakones, Isothapsakin B, Homothapsakin A*)Kim, S. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1769–1775; 2007, **70**, 714 (*Edulirin A, 10-Acetyledulirin A, 19,20-Dehydroedulirin A*)**Thebaine**

T-350

6,7,8,14-Tetrahydro-4,5-epoxy-3,6-dimethoxy-17-methylmorphinan, 9CI. Codeinone methyl enol ether. Paramorphine [115-37-7]

(-)-form

C₁₉H₂₁NO₃ 311.38

Log P 1.49 (uncertain value) (calc).

▶ QD2975000

(-)-form

Alkaloid from *Papaver* spp., esp. *Papaver bracteatum*; minor constit. of opium (Papaveraceae). Cholinesterase inhibitor. CNS stimulant. Narcotic showing higher narcotic activity and toxicity but weaker analgesic activity than Morphine, M-704. Causes histamine release from tissues. Rectangular plates (EtOH); needles by subl.; also obt. in cubes and prisms. Sol. C_6H_6 , $CHCl_3$, v. spar. sol. petrol. Mp 192.5°. $[\alpha]_D^{16}$ -221 (c, 0.35 in MeOH). Log P 1.49 (uncertain value) (calc). Pharmacol. active isomer.

► LD₅₀ (mus, ipr) 20 mg/kg.

N-Oxide: **Thebaine N-oxide**

$C_{19}H_{21}NO_4$ 327.379

Alkaloid from *Papaver bracteatum* (Papaveraceae).

N-Me: **Thebaine N-methosalt**. N-Methylthebaine

$C_{20}H_{24}NO_3^{\oplus}$ 326.414

Alkaloid from *Papaver bracteatum* (Papaveraceae). Needles (Me₂CO aq.) (as chloride). Mp 188-190° (chloride). $[\alpha]_D^{20}$ -125.8 (c, 0.81 in MeOH).

O³-De-Me: 6,7,8,14-Tetrahydro-4,5-epoxy-6-methoxy-17-methylmorphinan-3-ol. **Oripavine**

[467-04-9]

$C_{18}H_{19}NO_3$ 297.353

Alkaloid from *Papaver somniferum* (opium poppy), *Papaver orientale* and *Papaver bracteatum*. Metab. of Thebaine (Papaveraceae). Analgesic. Needles (EtOH). Mp 201-202°. $[\alpha]_D^{20}$ -216.9 (c, 3.44 in $CHCl_3$). Log P 0.91 (uncertain value) (calc).

► QD2097500

O³-De-Me, N-de-Me: **Oripavidine**. Nor-oripavine

[7168-66-3]

$C_{17}H_{17}NO_3$ 283.326

Alkaloid from *Papaver orientale* (Papaveraceae). Mp 250° dec. $[\alpha]_D$ -90 (c, 0.27 in MeOH).

16ξ-Hydroxy: **16-Hydroxythebaine**

[34388-67-5]

$C_{19}H_{21}NO_4$ 327.379

Constit. of opium. Pale yellow prisms (Me₂CO/petrol). Mp 126-128°.

(±)-form

Prisms (MeOH/Et₂O). Mp 184-186°. pK_a 8.15 (15°).

Kiselev, V.V. et al., *J. Gen. Chem. USSR (Engl. Transl.)*, 1948, **18**, 142; 855 (Oripavine)

Bentley, K.W. et al., *Chemistry of the Morphine Alkaloids*, Oxford Univ. Press, 1954, 184 (isol, props)

Okuda, S. et al., *Chem. Pharm. Bull.*, 1964, **12**, 104 (pmr)

Brochmann-Hanssen, E. et al., *J. Pharm. Sci.*, 1964, **53**, 1549 (glc, tlc)

Barton, D.H.R. et al., *J.C.S.*, 1965, 2423 (biosynth)

Rapoport, H. et al., *J.A.C.S.*, 1967, **89**, 1942 (synth)

Wheeler, D.M.S. et al., *J.A.C.S.*, 1967, **89**, 4494 (ms)

Kametani, T. et al., *J.C.S. (C)*, 1969, 2030 (synth, uv, ord, cd)

Brochmann-Hanssen, E. et al., *J.O.C.*, 1972, **37**, 1881 (16-Hydroxythebaine)

Barber, R.B. et al., *J. Med. Chem.*, 1975, **18**, 1074 (synth)

Terui, Y. et al., *Tet. Lett.*, 1975, 2853 (cmr)

Weller, D.D. et al., *J. Med. Chem.*, 1976, **19**, 1171 (synth)

Phillipson, J.D. et al., *Phytochemistry*, 1976, **15**, 1297 (oxide)

Shaffie, A. et al., *J. Polym. Sci.*, 1977, **66**, 1050 (Oripavine, isol, uv, ir)

Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1977, **13**, 714; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 600 (Oripavidine)

Rönsch, H. et al., *Phytochemistry*, 1979, **18**, 1089 (N-Metho salt)

Neilsen, B. et al., *Planta Med.*, 1983, **48**, 205 (Oripavine, isol, pmr, ms)

Theuns, H.G. et al., *J.C.S. Perkin 1*, 1984, 1701 (oxide)

Rice, K.C. et al., *Chem. Biol. Isoquinoline Alkaloids, Int. Symp., Phytochem. Soc. Eur., Abstr. Pap.*, 1985, 191 (rev, synth)

Mahler, C.H. et al., *Acta Cryst. C*, 1996, **52**, 3193-3195 (—)-form, cryst struct


Coop, A. et al., *Heterocycles*, 1998, **49**, 43-47 (synth)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TEN000

Thehaplosin

T-351

4,5-Dimethoxy-11-methylfuro[2,3-c]acridin-6(11H)-one, 9CI
[151870-95-0]



$C_{18}H_{15}NO_4$ 309.321


Alkaloid from aerial parts of *Haplophyl- lum thesioides*. Amorph. λ_{max} 242 (log ϵ 4.5); 270 (log ϵ 4.8); 322 (log ϵ 4.4); 396 (log ϵ 3.85) (MeOH).

Ulubelen, A. et al., *Nat. Prod. Lett.*, 1993, **1**, 269-272 (isol, uv, pmr, cmr, struct)

Agelas nakamurai Thelepo- gane-purine alkaloid

T-352

[213749-81-6]



$C_{26}H_{40}N_5$ 422.636

Alkaloid from the sponge *Agelas naka- murai*. Amorph. solid. $[\alpha]_D$ +12.9 (c, 0.18 in MeOH). Closely related to the Agela- sines. λ_{max} 215 (log ϵ 4.24); 272 (log ϵ 3.95) (MeOH).

Iwagawa, T. et al., *J. Nat. Prod.*, 1998, **61**, 1310-1312 (isol, uv, ir, pmr, cmr)

Thelepogidine

T-353

$C_{18}H_{27}NO_2$ 289.417

Struct. unknown. Alkaloid from *Thele-*

pogon elegans whole plant (Poaceae).


Needles (EtOH). Mp 254-255°. Conts. no C=O or OH groups.

Crow, W.D. et al., *Aust. J. Chem.*, 1962, **15**, 159-161

Thelepogine

T-354

[6899-83-8]



Absolute configuration

$C_{20}H_{31}NO$ 301.471

Alkaloid from the grass *Thelepogon elegans* (Poaceae). Mp 184-185°.

Methiodide: Mp 254-255°.


Crow, W.D. et al., *Aust. J. Chem.*, 1962, **15**, 159 (isol)

Fridrichsons, J. et al., *Acta Cryst.*, 1963, **16**, 206

Theobromine, BAN

T-355

3,7-Dihydro-3,7-dimethyl-1H-purine-2,6-dione, 9CI. 3,7-Dimethylxanthine. Diuro- bromine. Santheose. Thesal. FEMA 3591. Many other names [83-67-0]



$C_7H_8N_4O_2$ 180.166

Constit. of tea leaves (*Camellia thea*), *Theobroma cacao* (cocoa), *Cola acumi- nata*, *Paullinia cupana* and some *Ilex* and *Acer* spp. Adenosine receptor antagonist. Diuretic, cardiac stimulant, arterial dilator. Monoclinic needles. Mp 351°. Subl. 290. pK_{a1} 1; pK_{a2} 10 (25°). Log P -0.69 (calc).

► Human systemic effects by ingestion (CNS and gastrointestinal). LD₅₀ (rat, orl) 1265 mg/kg. Exp. reprod. and teratogenic effects (large dose). XH2275000

Perchlorate: Mp 271-273° dec.

1-Hexyl: **Pentifylline, BAN, INN. Cosal- don. Hexyltheobromine. SK 7** [1028-33-7]

$C_{13}H_{20}N_4O_2$ 264.327

Stabiliser for vitamin preparations. Vasodilator, diuretic agent. Mp 82-83°. Log P 2.59 (uncertain value) (calc).

► UO8450000

2-Hydroxy-form

O²-Me: 3,7-Dihydro-2-methoxy-3,7-di- methyl-6H-purin-6-one, 9CI. **2-O- Methyltheobromine** [19143-62-5]

$C_8H_{10}N_4O_2$ 194.193

Constit. of the gorgonian *Echinomur-*


aceae splendens. λ_{\max} 265 (ε 8555) (MeOH).

- Blout, E.R. *et al.*, *J.A.C.S.*, 1950, **72**, 479-484 (ir)
 Lockhart, E.L. *et al.*, *J.A.C.S.*, 1950, **72**, 5328-5329 (uv)
 Spitteller, G. *et al.*, *Monatsh. Chem.*, 1962, **93**, 632-638 (ms)
 Twanmoh, L.-M. *et al.*, *J. Het. Chem.*, 1973, **10**, 187-190 (pmr)
 Nicolau, C. *et al.*, *Z. Naturforsch., C*, 1974, **29**, 475-478 (cmr)
 Bergmann, F. *et al.*, *J.O.C.*, 1977, **42**, 2470-2473 (synth)
 Royer, R.F. *et al.*, *Actual. Pharmacol.*, 1982, **191**, 36-38 (rev. pharmacol)
 Ueda, T. *et al.*, *Heterocycles*, 1982, **19**, 2291-2294 (synth)
 Tarka, S.M. *et al.*, *Prog. Clin. Biol. Res.*, 1984, **158**, 9-16 (props. isol. biosynth)
 Shively, C.A. *et al.*, *Toxicol. Appl. Pharmacol.*, 1986, **84**, 593-598 (metabol. toxicol. rev)
 Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 688 (synonyms)
 Atta-ur-Rahman, *et al.*, *Alkaloids (Academic Press)*, 1990, **38**, 225-323 (spectral props)
 Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 2833-2862 (synth. uv, pmr)
IARC Monog., 1991, **51**, 421 (rev. tox)
Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, **2**, 751
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2771-2772
 Ford, K.A. *et al.*, *Acta Cryst. C*, 1998, **54**, 1980-1983 (cryst. struct)
 Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 765
 Parameswaran, P.S. *et al.*, *Indian J. Chem., Sect. B*, 2002, **41**, 1093-1096 (2-O-Methyltheobromine)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, TEO500; TEO750

Theoneberine

T-356

[145400-57-3]



$C_{27}H_{25}Br_4NO_6$ 779.114


Biogenetically unique isoquinoline alkaloid. Classified as a hybrid between 1-benzylisoquinolines and protoberberines. Alkaloid from the Okinawan marine sponge *Theonella* sp. Exhibits antimicrobial activity against gram-positive bacteria. Also shows cytotoxicity against murine lymphoma L1210 and human epidermoid carcinoma KB cells *in vitro*. Sol. MeOH, $CHCl_3$; poorly sol. H_2O . Mp 128°. $[\alpha]_D^{20}$ -53 (c, 0.6 in $CHCl_3$). λ_{\max} 286 (ε 5300) (MeOH/HCl) (Derep). λ_{\max} 300 (ε 6800) (MeOH/NaOH) (Derep). λ_{\max} 284 (ε 5300) (MeOH) (Derep).

Kobayashi, J. *et al.*, *J.O.C.*, 1992, **57**, 6680 (isol. uv, ir, pmr, cmr, ms, struct)

Theonelladine A

T-357

14-(3-Pyridinyl)-11-tetradecen-1-amine, 9CI. 3-(14-Amino-3-tetradecenyl)pyridine [125289-09-0]



$C_{19}H_{32}N_2$ 288.475

Mol. formula erroneously reported as $C_{20}H_{32}N_2$ in paper. Alkaloid from the Okinawan marine sponge *Theonella swinhoei*. Exhibits potent antineoplastic activity. Calcium release inducer. λ_{\max} 210 (ε 12300); 257 (ε 3300); 263 (ε 3600); 269 (ε 2700) (MeOH) (Derep).

1-Aldehyde, oxime (Z-): [291775-83-2]

$C_{19}H_{30}N_2O$ 302.459

Isol. from *Amphimedon* sp. Oxime config. undetermined. λ_{\max} 264 (ε 3300) (MeOH).

N-Me: *Theonelladine B*

[125289-10-3]

$C_{20}H_{34}N_2$ 302.502

Alkaloid from *Theonella swinhoei*. Exhibits potent antineoplastic activity. Calcium release inducer. Mol. formula erroneously reported as $C_{21}H_{34}N_2$ in paper. λ_{\max} 210 (ε 12300); 257 (ε 3300); 263 (ε 3600); 269 (ε 2700) (MeOH) (Derep).

3',4'-Dihydro, 9',10'-didehydro, 1-aldehyde, oxime (Z,Z-): [291775-84-3]

$C_{19}H_{30}N_2O$ 302.459

Isol. from *Amphimedon* sp. λ_{\max} 264 (ε 3200) (MeOH).

11,12-Didehydro: 14-(3-Pyridinyl)-11-tetradecyn-1-amine. 3-(11-Amino-3-tetradecynyl)pyridine. *Niphatesine A* [132923-10-5]

$C_{19}H_{30}N_2$ 286.459

Alkaloid from the marine sponge *Niphates* sp. Antineoplastic. Oil. λ_{\max} 207 (ε 5900); 258 (ε 2800); 263 (ε 3200); 268 (ε 2500) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *J.C.S. Perkin I*, 1990, 3301-3303 (isol. uv, pmr, cmr, ms, struct)

Rao, A.V.R. *et al.*, *J.O.C.*, 1991, **56**, 4545-4547 (synth)


Teubner, A. *et al.*, *Annalen*, 1993, 161-165 (synth)

Hirano, K. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 974-977 (oximes, isol)

Theonelladine C

T-358

3-Pyridinetridecanamine. 3-(13-Aminotridecyl)pyridine. 3-(3-Pyridyl)tridecylamine [125289-11-4]



$C_{18}H_{32}N_2$ 276.464

Alkaloid from the Okinawan marine sponge *Theonella swinhoei*. Exhibits potent antineoplastic activity. Calcium release inducer. Obt. mixed with traces of

analogues having branched alkyl chains. λ_{\max} 210 (ε 12300); 257 (ε 3300); 263 (ε 3600); 269 (ε 2700) (MeOH) (Derep).

N-Me: *Theonelladine D*

[125289-12-5]

$C_{19}H_{34}N_2$ 290.491

Alkaloid from *Theonella swinhoei*. Exhibits potent antineoplastic activity. Calcium release inducer. Not obt. completely pure. λ_{\max} 210 (ε 12300); 257 (ε 3300); 263 (ε 3600); 269 (ε 2700) (MeOH) (Derep).

N-Methoxy: N-Methoxy-3-pyridinetridecanamine. *Ikimine C*

[131479-32-8]

$C_{19}H_{34}N_2O$ 306.49

Alkaloid from *Niphates* sp. and other sponges. Cytotoxic. Oil. λ_{\max} 260 (ε 2000); 265 (ε 6610); 270 (ε 2880) (MeOH) (Derep). λ_{\max} 260 (ε 1995); 264 (ε 6166); 272 (ε 3630) (MeOH) (Berdy).

Kobayashi, J. *et al.*, *Tet. Lett.*, 1989, **30**, 4833 (isol. uv, pmr, cmr, ms, struct)

Carroll, A.R. *et al.*, *Tetrahedron*, 1990, **46**, 6637 (*Ikimine C*)

Rao, A.V.R. *et al.*, *J.O.C.*, 1991, **56**, 4545 (synth)


Teubner, A. *et al.*, *Annalen*, 1993, 161-165 (synth)

Wang, Y. *et al.*, *J.O.C.*, 2003, **68**, 3090-3098 (synth)

Shorey, B.J. *et al.*, *Tetrahedron*, 2007, **63**, 5587-5592 (synth)

Theonezolides

T-359



Macrolide antibiotics. n = 6,4,8 for Theonezolides A,B,C respectively. Partial stereochem. is known. Isol. from the marine sponge *Theonella* sp. Cytotoxic agents.

Theonezolid A [150243-49-5]

$C_{79}H_{140}N_4O_{22}S_2$ 1562.121

Needles + 3 H_2O . Mp 123°. $[\alpha]_D^{28}$ -8.1 (c, 1.5 in MeOH). λ_{\max} 210 (ε 22000) (MeOH).

Theonezolid B [157536-46-4]

$C_{77}H_{136}N_4O_{22}S_2$ 1534.067

Cryst. Mp 125°. $[\alpha]_D^{28}$ -8 (c, 1.5 in MeOH). λ_{\max} 211 (ε 22000) (MeOH).

Theonezolid C [157536-47-5]

$C_{81}H_{144}N_4O_{22}S_2$ 1590.174


Cryst. Mp 122°. $[\alpha]_D^{28}$ -7.5 (c, 1.5 in MeOH). λ_{\max} 211 (ε 25000) (MeOH).

Kobayashi, J. *et al.*, *J.A.C.S.*, 1993, **115**, 6661-6665 (isol. uv, ir, pmr, cmr, ms)

- Kondo, K. *et al.*, *Tetrahedron*, 1994, **50**, 8355-8362 (*isol, uv, ir, pmr, cmr, ms, struct*)
 Kobayashi, J. *et al.*, *Heterocycles*, 1998, **49**, 39-42 (*stereochem*)
 Sato, M. *et al.*, *Tetrahedron*, 1998, **54**, 4819-4826 (*stereochem*)

Theophylline, BAN, JAN, USAN **T-360**

3,7-Dihydro-1,3-dimethyl-1H-purine-2,6-dione, 9CI. 1,3-Dimethylxanthine. Austyn. Elan. Elixophyllin. Euphyllin. Nuclin. Spophyllin retard. Teonova. Theocin. Theodel. Theo-Dur. Theograd. Theolair. Theona P. Theotard. Uniphyllin. Many other names [58-55-9]



C₇H₈N₄O₂ 180.166
 Constit. of tea leaves (*Camellia thea*), *Ilex paraguariensis* and *Paullinia cupana*. Also *isol.* from the marine sponge *Haliclona* sp. Adenosine receptor antagonist. Smooth muscle relaxant of value in the treatment of bronchial conditions. Antiasthmatic agent. Synergises the antileukaemic activity of nitrosoureas. Diuretic and cardiac stimulant. Mp 268° Mp 264°. pK_{a1} 1; pK_{a2} 8.6 (25°). Log P -0.06 (uncertain value) (calc).

▶ Adverse effects reported when used therapeutically incl. gastrointestinal and CNS changes. Overdosage can cause death. LD₅₀ (rat, orl) 244 mg/kg. Exp. reprod. and teratogenic effects. XH3850000


N⁷-(2-Deoxy-β-D-erythro-pentofuranosyl): *Aplysidine* [144096-49-1]
 C₁₂H₁₆N₄O₅ 296.282
Isol. from the sponge *Aplysina* sp. Adenosine A1 receptor antagonist. Amorph. solid. Sol. H₂O, MeOH. Mp 162°. [α]_D²⁰ +17 (c, 0.4 in H₂O). λ_{max} 274 (ε 8500) (H₂O) (Berdy). λ_{max} 275 (ε 8400) (HCl) (Berdy). λ_{max} 274 (ε 8400) (NaOH) (Berdy).

[49746-06-7, 5967-84-0, 8002-89-9, 85531-28-8]

- Cohen, J.L. *et al.*, *Anal. Profiles Drug Subst.*, 1975, **4**, 466-493 (*rev, ir, pmr, uv, ms, anal*)
 Naqvi, A.A. *et al.*, *J. Appl. Crystallogr.*, 1981, **14**, 464 (*cryst struct*)
 Kondo, K. *et al.*, *Tetrahedron*, 1992, **48**, 7145-7148 (*Aplysidine*)
 Ebisuzaki, Y. *et al.*, *Acta Cryst. C*, 1997, **53**, 777-779 (*cryst struct*)
 Markham, A. *et al.*, *Drugs*, 1998, **56**, 1081-1091 (*rev*)

Thermopsamine **T-361**

13-Hydroxysparteine [14145-73-4]



Absolute Configuration

C₁₅H₂₆N₂O 250.383
 Alkaloid from *Thermopsis lanceolata* epigeal parts (Fabaceae). Also from *Virgilia divaricata* and *Virgilia oroboides* (no stereochem. indicated). Mp 154-155°. [α]_D +26.4 (EtOH).

Dipicrate: Mp 130-131°.
Methiodide: Mp 244-246° dec.

Ketone: 13-Oxosparteine [55869-87-9]
 C₁₅H₂₄N₂O 248.367
 Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae) (no stereochem. detd.).


4-Oxo: 4-Oxothermopsamine. 13α-Hydroxy-4-oxosparteine [162584-08-9]
 C₁₅H₂₄N₂O₂ 264.367
 Minor alkaloid from the seeds of *Lupinus varius*. Amorph. [α]_D²⁵ -131 (c, 0.1 in MeOH).

13-Epimer: 13-Epihydroxysparteine
 C₁₅H₂₆N₂O 250.383
 Alkaloid from *Thermopsis mongolica*. [72401-67-3, 72401-68-4, 10349-35-6]

- Neuner-Jehle, N. *et al.*, *Monatsh. Chem.*, 1964, **95**, 687-709 (*ms*)
 Vinogradova, V.I. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 463-466; *Chem. Nat. Compd. (Engl. Transl.)*, 1971, **7**, 440-442 (*isol, struct*)
 Christov, V. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1413-1415 (*13-epi-Hydroxysparteine*)
 Veen, G. *et al.*, *Phytochemistry*, 1991, **30**, 1891-1895 (*isol*)
 Mohamed, M.H. *et al.*, *Pharmazie*, 1999, **54**, 778-780 (*4-Oxothermopsamine*)

Thermopsidine **T-362**

Alternine. Dimethylenedicytisine




C₂₄H₃₀N₄O₂ 406.527
Isol. from *Thermopsis alterniflora* (Fabaceae). Mp 184-186°. [α]_D -216.3.
 Artifact arising from ClCH₂CH₂Cl used in *isol.*

Hydrate: Mp 70°.
Hydrochloride (1:2): Mp 270-271° dec.
 Padukina, Z.P. *et al.*, *CA*, 1960, **54**, 6783

Thermopsine **T-363**

Hexalupine. *Isoanagryrine*



C₁₅H₂₀N₂O 244.336

(+)-form
 Alkaloid from *Lupinus caudatus* and *Lupinus corymbosus* (Fabaceae). Mp 207°. [α]_D²⁵ +155 (EtOH).

Perchlorate: Mp 289°.


(-)-form [486-90-8]
 Alkaloid from *Thermopsis lanceolata*, *Thermopsis rhombifolia* and *Sophora secundiflora* seeds (Fabaceae). Mp 207°. [α]_D -159.6 (EtOH).
Hydrochloride: Mp 247-248° dec.

(±)-form
 Prisms (Me₂CO). Mp 171-172°.
Perchlorate: Mp 260-263°.

- 11-Epimer: see Anagryrine, A-970
 Orechov, A. *et al.*, *Ber.*, 1933, **66**, 625 (*isol*)
 Cockburn, W.F. *et al.*, *Can. J. Chem.*, 1951, **29**, 13 (*struct*)
 Marion, L. *et al.*, *Can. J. Chem.*, 1951, **29**, 22 (*isol, ir*)
 van Tamelen, E.E. *et al.*, *J.A.C.S.*, 1956, **78**, 2913 (*synth*)
 Bohlmann, F. *et al.*, *Chem. Ber.*, 1962, **95**, 944 (*synth*)
 Vul'fson, N.S. *et al.*, *Khim. Geterotsikl. Soedin.*, 1974, 251; *CA*, **81**, 13678z (*ms*)
 Robins, D.J. *et al.*, *Magn. Reson. Chem.*, 1992, **30**, 1125 (*config*)
 Gray, D. *et al.*, *Angew. Chem., Int. Ed.*, 2006, **45**, 2419-2423 (*synth*)

Thesine **T-364**

Diisoretronecanyl p,p'-dihydroxy-α-truxillate [528-37-0]



Absolute configuration

C₃₄H₄₂N₂O₆ 574.716
 Alkaloid from *Thesium minkwitzianum* whole plant (Santalaceae). Cryst. (EtOH). Mp 254-256°. Hydrol. gives (+)-isoretronecanol and thesinic acid (*p,p'*-dihydroxy-α-truxillic acid).

Sulfate: Mp 244-246°. [α]_D²⁰ +33.4 (c, 1.6 in H₂O). Forms a tetrahydrate.

Picrate:
 Fine-yellow cryst. (EtOH). Mp 224-226° dec.

Methiodide (1:2): Mp 140-150°. [α]_D²⁰ +33.24 (c, 2.2 in EtOH).

- Arendaruk, A.P. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 484; 489; 670; *J. Gen. Chem. USSR (Engl. Transl.)*, 507; 511; 693 (*isol, struct*)

Thesinicine **T-365**

C₁₀H₁₁NO₂ 177.202
 Prob. pyrrolizidine alkaloid. Struct. unknown. Alkaloid from epigeal parts of *Thesium minkwitzianum* (Santalaceae). Mp 124-125°.


Hydrochloride: Mp 197-199°.

Picrate: Mp 158-159° dec.

- Arendaruk, A.P. *et al.*, *Zh. Obshch. Khim.*, 1960, **30**, 670-676; *J. Gen. Chem. USSR (Engl. Transl.)*, 1960, **30**, 693

Thiamine, BAN, INN T-366

3-[(4-Amino-2-methyl-5-pyrimidinyl)-methyl]-5-(2-hydroxyethyl)-4-methylthiazolium(1+), 9CI, 8CI. Vitamin B₁. Torulin. Oryzantin. Aneurine. Vitaneurin. Many other names [70-16-6]



C₁₂H₁₇N₄OS[⊕] 265.358
Ubiquitous constit. of biol. materials. Produced by numerous bacterial spp. Used as aq. soln. for fluorimetric detn. of PO₄²⁻ (λ_{max} 440 nm, 5-100 ppb, via molybdatophosphate). Essential vitamin. Sol. H₂O; sl. sol. EtOH. pK_a 5.17. Log P -3.91 (uncertain value) (calc). Labile in acid or alkaline soln. or on heating. In alkaline soln. the thiazolium ring is reversibly opened to thiols. In therapy thiamine can be replaced by various ring-opened analogues, e.g. Benfotiamine.

Chloride: Thiamine monochloride. Thiacrat [59-43-8]
C₁₂H₁₇ClN₄OS 300.811
Cryst. + 1H₂O (H₂O). Mp 120-122° dec., 163-165° dec. (anhyd.). Infrequently encountered, hydrochloride more stable.

▶ XI6550000

Chloride; hydrochloride: Thiamine hydrochloride, USAN. Bewon. Betaxin. Betalin S. Vinothiam. FEMA 3322 [67-03-8]
C₁₂H₁₈Cl₂N₄OS 337.272
Clinically used vitamin source. Oral or parenterally administered doses. Well tolerated. Plates or cryst. (EtOH). V. sol. H₂O; spar. sol. EtOH; insol. Et₂O, C₆H₆. Mp 248° dec. Mp is not a good criterion of purity.

▶ XI7350000

Nitrate: Thiamine mononitrate, USAN [532-43-4]
C₁₂H₁₇N₅O₄S 327.363
Clinically used vitamin source. Mp 164-165° Mp 196-200°.

▶ XI7400000

O-Phosphate: Monophosphothiamine [532-40-1]
C₁₂H₁₈N₄O₄PS[⊕] 345.338
Vitamin B₁ deriv. used clinically. Mp 200° (as chloride).

O-Diphosphate: Thiamine diphosphate. Cocarboxylase. Diphosphothiamine. Pyruvodehydrase. Thiamine pyrophosphate. Aneurin diphosphate. Coxylase. Cozymase II. Interacton. Nutrase. Many other names [154-87-0]
C₁₂H₁₈N₄O₇P₂S 424.31
Coenzyme of the yeast enzyme carboxylase. Catalyses the decarboxylation of α-ketoacids. V. pale yellow needles (as hydrochloride). Mp 240°

(hydrochloride).

▶ XI7520000

O-Triphosphate: Thiamine triphosphate. Thiamine triphosphoric acid. TTP[†] [15666-52-1 (chloride)]
C₁₂H₂₀N₄O₁₃P₃S 553.296
Isol. from animal tissues. Cryst. (EtOH aq.).

[7019-71-8]


Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 515C (nmr)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 647A (ir)
Lohmann, K. et al., *Biochem. Z.*, 1937, 294, 188
Todd, A.R. et al., *J.C.S.*, 1937, 364; 1938, 26 (synth)
Wejllard, J. et al., *J.A.C.S.*, 1938, 60, 2263; 1941, 63-1160
Karrer, P. et al., *Helv. Chim. Acta*, 1946, 29, 711
Lenormant, H. et al., *Bull. Soc. Chim. Fr.*, 1954, 375 (ir, uv)
Greiling, H. et al., *Nature (London)*, 1958, 13, 251 (isol, triphosphate)
Kotera, K. et al., *Chem. Pharm. Bull.*, 1965, 13, 440 (pmr)
Hesse, M. et al., *Helv. Chim. Acta*, 1967, 50, 808 (ms)
Linnett, P.E. et al., *J.C.S.(C)*, 1967, 796 (biosynth)
Ullrich, et al., *Vitam. Horm. (N.Y.)*, 1970, 28, 365 (rev, biochem)
Florkin, M. et al., *Compr. Biochem.*, Elsevier, Amsterdam, Vol. 11, 1971, 3 (rev, biosynth)
Pletcher, J. et al., *Acta Cryst. B*, 1972, 28, 2928; 1997, 33, 3349 (cryst struct)
Karrer, W. et al., *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 2609
Holzbecher, J. et al., *Anal. Chim. Acta*, 1973, 64, 147 (detn, P)
Dwivedi, B.K. et al., *J. Agric. Food Chem.*, 1973, 21, 54 (props)
Gallo, A.A. et al., *J. Biol. Chem.*, 1974, 249, 1382 (cmr)
Thiamine, [Proc. Pap. Discuss. U.S.-Jpn. Semin.], 2nd, Wiley, N.Y., 1976, (book)
Chauvet-Monges, A. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1978, 286, 489 (nmr)
Oka, Y. et al., *Kirk-Othmer Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, 24, 124 (synth)
Penttinen, H.K. et al., *Methods Enzymol.*, 1979, 62, 112 (triphosphate)
Banan, E.J. et al., *Z. Naturforsch., B*, 1979, 34, 1615 (ir)
Rindi, G. et al., *Acta Vitaminol. Enzymol.*, 1982, 4, 59 (metab)
Brown, G.M. et al., *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1982, 53, 345 (biosynth)
Sable, H.Z. et al., *Ann. N.Y. Acad. Sci.*, 1982, 378, 78; 378; 454 (book, nmr, cryst struct, conformn)
Sanemori, H. et al., *Experientia*, 1982, 38, 1044 (metab)
Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 7836
Uray, G. et al., *Monatsh. Chem.*, 1982, 113, 1475 (synth)
Davis, R.E. et al., *Adv. Clin. Chem.*, 1983, 23, 93 (chemistry)
Ishida, T. et al., *Acta Cryst. C*, 1984, 40, 437 (cryst struct)
Gubler, C.J. et al., *Handbook of Vitamins*, (Machlin, L.J. Ed.), M. Dekker, New York, 1984, 245 (rev)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 2515; 2540; 2558; 4574
Thiamin Pyrophosphate Biochemistry, (eds. Schellenberger, A. et al), CRC Press, Boca Raton, Florida, 1988, (book)
Al-Rashood, K.A.M. et al., *Anal. Profiles Drug Subst.*, 1989, 18, 413 (rev)
Contant, P. et al., *Helv. Chim. Acta*, 1990, 73, 1300 (synth)
Hu, N. et al., *Acta Cryst. C*, 1992, 48, 1951 (cryst struct)
Bender, D.A. et al., *Nutritional Biochemistry of the Vitamins*, Cambridge University Press, Cambridge, 1992, 128 (rev)
Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1053
Casas, J.S. et al., *Acta Cryst. C*, 1994, 50, 1265 (cryst struct, acetate)
Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 2, 1797
Estramareix, B. et al., *New J. Chem.*, 1996, 20, 607 (rev, biosynth)
Spenser, I.D. et al., *Angew. Chem., Int. Ed.*, 1997, 36, 1033-1046 (rev, biosynth)
Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2772-2778 (rev, use, occur)
Roje, S. et al., *Phytochemistry*, 2007, 68, 1904-1921 (biosynth, rev)
Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TES750; TET300; TET500; TET750

Thiangazole

T-367

[138667-71-7]




C₂₆H₂₉N₅O₂S₃ 539.745
Metab. of *Polyangium* sp. Inhibitor of HIV-1. Anthelmintic. Mp 140°. [α]_D²² -287 (c, 0.1 in MeOH). Similar to Tantzazole B, T-30. λ_{max} 211 (sh); 218 (sh); 223 (ε 37800); 228 (sh); 288 (ε 36400); 300 (sh) (MeOH) (Derep). λ_{max} 223 (ε 38000); 288 (ε 36200) (MeOH) (Berdy). λ_{max} 223 (ε 49774); 289 (ε 41640) (EtOH) (Berdy).

Jansen, R. et al., *Annalen*, 1992, 357 (isol, uv, ir, pmr, cmr, ms, struct)
Boyce, R.J. et al., *Tet. Lett.*, 1994, 35, 5705 (synth)
Akaji, K. et al., *Tetrahedron*, 1999, 55, 10685-10694 (synth)

Thiaplidiaquinone A

T-368

[852872-92-5]



R¹ = -CH₂CH=C(CH₃)CH₂CH₂CH=C(CH₃)₂ (E), R² = H

C₃₄H₄₁NO₆S 591.767
Alkaloid from the ascidian *Aplidium*

conicum. Antitumour agent. λ_{\max} 354 (€ 9300) (MeOH).

Aiello, A. *et al.*, *J. Med. Chem.*, 2005, **48**, 3410-3416 (*isol*, *pmr*, *cmr*)

Thiaphlidiaquinone B T-369

[852872-93-6]

As Thiaphlidiaquinone A, T-368 with

$R^1 = H$, $R^2 = -CH_2CH=$

$C(CH_3)CH_2CH_2CH=C(CH_3)_2(E-)$

$C_{34}H_{41}NO_6S$ 591.767


Alkaloid from the ascidian *Aplidium conicum*. Antitumour agent. λ_{\max} 324 (€ 6300) (MeOH).

Aiello, A. *et al.*, *J. Med. Chem.*, 2005, **48**, 3410-3416 (*isol*, *pmr*, *cmr*)

Thiazohalostatin T-370

Antibiotic HQ 24A. HQ 24A

[152509-78-9]




$C_{20}H_{25}Cl_3N_2O_4S$ 495.853

Prod. by *Actinospira* sp. HQ24 and *Actinomadura* sp. Hq24-Bp4037. Cytoprotective agent. Antioxidant, anti-ischaemic agent, antiinflammatory. Powder. Sol. MeOH, EtOAc, EtOH, Me_2CO , $CHCl_3$. Mp 67-69°. $[\alpha]_D^{22}$ -122 (c, 1 in MeOH). λ_{\max} 252 (€ 7650); 275 (€ 5110); 287 (€ 4690); 345 (€ 33500); 360 (€ 28900) (MeOH) (Derep).

Yamagishi, Y. *et al.*, *J. Antibiot.*, 1993, **46**, 1633; 1638 (*isol*, *pmr*, *cmr*, *struct*, *props*)
Japan. Pat., 1994, 94 157 520; *CA*, **121**, 253888f

5-Thiazoleethanol, 9CI T-371

2-(5-Thiazolyl)ethanol. 5-(2-Hydroxyethyl)thiazole
[5664-55-1]



C_5H_7NOS 129.182


Prod. by an unidentified marine bacterium. Bp₁₀ 138-141°.

Ulf, H. *et al.*, *Acta Pharm. Suec.*, 1967, **4**, 269-280 (*synth*)

Laasch, H. *et al.*, *Dissertation*, Univ. of Göttingen, 2005, (*marine*, *isol*)

3-(2-Thiazolyl)-1H-indole, 9CI T-372

Camalexin. 2-(3-Indolyl)thiazole
[135531-86-1]



$C_{11}H_8N_2S$ 200.264

Alkaloid from the leaves of *Camelina sativa* (false flax) infected by the fungus *Alternaria brassicae*. Phytoalexin. Cryst. (MeOH). Mp 134-137°. λ_{\max} 214 (€ 22200); 274 (€ 7900); 318 (€ 13800) (MeOH) (Derep).

N^1 -Ac: [141690-88-2]

$C_{13}H_{10}N_2OS$ 242.301

Cryst. Mp 117.5°.

N^1 -Me: 1-Methyl-3-(2-thiazolyl)-1H-indole, 9CI. 1-Methylcamalexin

[194155-86-7]

$C_{12}H_{10}N_2S$ 214.29

From *Capsella bursa-pastoris* (shepherd's purse) infected with *Alternaria brassicae*. Phytoalexin. Brown solid (hexane). Mp 69-70°.

6-Methoxy: 6-Methoxy-3-(2-thiazolyl)-1H-indole. 6-Methoxycamalexin

[135531-87-2]

$C_{12}H_{10}N_2OS$ 230.29

From *Camelina sativa* (false flax) infected with *Alternaria brassicae*. Phytoalexin. Cryst. (MeOH/hexane). Mp 157-159°. λ_{\max} 218 (€ 21700); 296 (€ 11500); 324 (€ 11300) (MeOH) (Derep).

Browne, L.M. *et al.*, *Tetrahedron*, 1991, **47**,

3909 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Ayer, W.A. *et al.*, *Tetrahedron*, 1992, **48**, 2919

(*synth*)

Fürstner, A. *et al.*, *Tetrahedron*, 1995, **51**, 773

(*synth*)

Sakamoto, T. *et al.*, *J.C.S. Perkin 1*, 1996, 1927

(*synth*, *pmr*)

Moody, C.J. *et al.*, *Anti-Cancer Drugs*, 1997, **8**,

489-499 (*synth*, *ir*, *pmr*, *cmr*, *ms*, N^1 -Ac, N^1 -Me)

Pedras, M.S.C. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 2255-2260 (*metab*)

Jimenez, L.D. *et al.*, *Phytoprotection*, 1997, **78**,

99-103 (1-Methylcamalexin)

Pedras, M.S.C. *et al.*, *Phytochemistry*, 2000, **53**,

161-176 (*rev*)

Dzurilla, M. *et al.*, *Molecules*, 2001, **6**, 716-720

(*synth*)

Koradin, C. *et al.*, *Tetrahedron*, 2003, **59**, 1571-

1587 (*synth*)


Glawischig, E. *et al.*, *Phytochemistry*, 2007,

68, 401-406 (*rev*)

Thiazostatin A T-373

4-Methylpyochelin. AG 55. Antibiotic AG 55

[124508-94-7]



$C_{15}H_{18}N_2O_3S_2$ 338.451

Isol. from *Streptomyces toluosus*. Antioxidant. Pale yellow needles. Mp 69-72° dec. $[\alpha]_D^{25}$ +29.1 (c, 1 in MeOH). λ_{\max} 211 (€ 18700); 270 (€ 11800); 341 (€ 4700) (MeOH/HCl) (Derep). λ_{\max} 213 (€ 23800); 248 (€ 12200); 312 (€ 5400) (MeOH/NaOH)

(Derep). λ_{\max} 211 (€ 18700); 270 (€ 11800); 341 (€ 4700) (MeOH/HCl) (Derep).

2-Epimer: Thiazostatin B

[124509-83-7]

$C_{15}H_{18}N_2O_3S_2$ 338.451

From *Streptomyces toluosus*.

Antioxidant. Pale yellow needles. Sol. Me_2CO , MeOH, $CHCl_3$, EtOAc, DMSO; poorly sol. H_2O , hexane. Mp 75-78°. $[\alpha]_D^{25}$ -80.7 (c, 1 in MeOH). λ_{\max} 211 (€ 18700); 270 (€ 11800); 341 (€ 4700) (MeOH/HCl) (Derep). λ_{\max} 213 (€ 23800); 248 (€ 12200); 312 (€ 5400) (MeOH/NaOH) (Derep). λ_{\max} 211 (€ 18700); 270 (€ 11800); 341 (€ 4700) (MeOH/HCl) (Derep). λ_{\max} 211 (E1%/1cm 567); 270 (E1%/1cm 350); 341 (E1%/1cm 140) (MeOH/HCl) (Berdy). λ_{\max} 213 (E1%/1cm 717); 250 (E1%/1cm 363); 312 (E1%/1cm 157) (MeOH/NaOH) (Berdy).


[122054-40-4]

Shindo, K. *et al.*, *J. Antibiot.*, 1989, **42**, 1526-1529 (*isol*)

Thienodolin T-374

6-Chloro-8H-thieno[2,3-b]indole-2-carboxamide, 9CI. MJ 286A

[149127-27-5]



$C_{11}H_7ClN_2OS$ 250.708

Prod. by *Streptomyces albobriseolus*.

Plant growth regulator. Dark yellow prisms. Sol. MeOH, DMSO, Me_2CO ; poorly sol. H_2O . Mp 249-251°. λ_{\max} 239 (€ 44700); 271 (sh) (€ 12600); 288 (€ 18500); 327 (€ 18300) (MeOH). λ_{\max} 238; 288 (€ 18500); 327 (MeOH-HCl) (Berdy). λ_{\max} 238; 288; 327 (€ 18300) (MeOH-NaOH) (Berdy).

Kanbe, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 1993, **57**, 632-635; 636-637 (*isol*,


uv, *ir*, *pmr*, *cmr*, *cryst struct*)

Engqvist, R. *et al.*, *Eur. J. Org. Chem.*, 2004,

2589-2592 (*synth*)

Thiersindole A T-375

[610317-20-9]




$C_{28}H_{39}NO$ 405.622
Prod. by *Penicillium thiersii* MYC-500.
Amorph. solid. Mp 89-92°. $[\alpha]_D^{25} +26$ (c, 0.1 in CH_2Cl_2). λ_{max} 227 (log ϵ 4.26); 254 (log ϵ 3.43); 292 (log ϵ 3.48) (MeOH).

$\Delta^{2,3}$ -Isomer: **Thiersindole B**

[610317-21-0]
 $C_{28}H_{39}NO$ 405.622
Prod. by *Penicillium thiersii* MYC-500.
Amorph. solid. Mp 95-98°. $[\alpha]_D^{25} +21$ (c, 0.1 in CH_2Cl_2). λ_{max} 224 (log ϵ 4.28); 284 (log ϵ 3.49); 292 (log ϵ 3.45) (MeOH).

Li, C. et al., *J. Nat. Prod.*, 2003, **66**, 1232-1235 (isol, pmr, cmr, ms)


Thiersindole C
[610317-22-1]



$C_{28}H_{39}NO$ 405.622
Prod. by *Penicillium thiersii* MYC-500.
Oil. $[\alpha]_D^{25} -42$ (c, 0.1 in CH_2Cl_2). λ_{max} 220 (log ϵ 4.26); 288 (log ϵ 3.49); 292 (log ϵ 3.48) (MeOH).

Li, C. et al., *J. Nat. Prod.*, 2003, **66**, 1232-1235 (isol, pmr, cmr, ms)

Thiersinine A
[468733-26-8]



$C_{30}H_{37}NO_7$ 523.625
Prod. by *Penicillium thiersii* NRRL 28147. Antiinsectan. Solid. Mp 255-258°. $[\alpha]_D^{25} -80$ (c, 0.3 in CH_2Cl_2). λ_{max} 230 (ϵ 17000); 272 (ϵ 3000) (MeOH).


O-De-Me: **Thiersinine B**

[468733-27-9]
 $C_{29}H_{35}NO_7$ 509.598
Prod. by *Penicillium thiersii* NRRL 28147. Antiinsectan. Oil. $[\alpha]_D^{25} -93$ (c, 0.1 in CH_2Cl_2). A diketone. λ_{max} 230 (ϵ 20300); 284 (ϵ 3800) (MeOH).

Li, C. et al., *Org. Lett.*, 2002, **4**, 3095-3098 (isol, pmr, cmr, ms)

Thioactin

[161505-22-2]



$C_{43}H_{40}N_{14}O_{11}S_4$ 1057.14

Cyclic peptide antibiotic related to Thioxamycin. Prod. by *Streptomyces* sp. DP94. Powder. Sol. Me_2CO , $CHCl_3$; poorly sol. H_2O . Mp 250-255° dec. $[\alpha]_D^{25} -42.5$ (c, 0.1 in $CHCl_3/MeOH$). λ_{max} 215 (sh) (ϵ 40000); 236 (ϵ 42000) (EtOH).

Yun, B.-S. et al., *J. Antibiot.*, 1994, **47**, 1541 (isol, uv, ir, pmr, cmr)

Thiobideoxynupharidine

$C_{30}H_{40}N_2OS$ 476.725


Nuphar alkaloid. Struct. unknown. Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Prismatic needles (as diperchlorate). Mp 225-226° (diperchlorate). $[\alpha]_D^{25} +26.6$ (EtOH) (diperchlorate).

Achmatowicz, O. et al., *Pol. J. Chem. (Rocz. Chem.)*, 1962, **36**, 1815-1825 (isol, ir)

Thiobinupharidine

Thionupharlutine A

[30343-72-7]



Absolute Configuration

$C_{30}H_{42}N_2O_2S$ 494.74

Alkaloid from *Nuphar luteum* (Nymphaeaceae). Mp 129-130°. $[\alpha]_D^{25} +7.8$ (c, 0.18 in MeOH).

Hydrobromide (1:2):

Cryst. + 2 H_2O . Mp 245-247°.

Perchlorate (1:2): Mp 282-284°. $[\alpha]_D^{25} +49.8$ (H_2O).

S-Oxide (S-): **anti-Thiobinupharidine**

T-378

sulfoxide

$C_{30}H_{42}N_2O_3S$ 510.739

Alkaloid from *Nuphar luteum* (Nymphaeaceae). Glass. $[\alpha]_D^{25} -64$ (c, 0.73 in EtOH).

6 ξ -Hydroxy: **Thionupharoline. 6-Hydroxythiobinupharidine**

[50478-55-2]

$C_{30}H_{42}N_2O_3S$ 510.739

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Noncryst.; cryst. (as perchlorate). Mp 265-273° dec. (perchlorate). $[\alpha]_D^{25} +33$ (c, 3.6 in CH_2Cl_2). λ_{max} 204 (ϵ 35000) (EtOH).

6' ξ -Hydroxy: **6'-Hydroxythiobinupharidine**

[52002-85-4]

$C_{30}H_{42}N_2O_3S$ 510.739

Alkaloid from *Nuphar luteum* (Nymphaeaceae). Mp 216-220° (as perchlorate). $[\alpha]_D^{25} +34$ (c, 1 in 95% EtOH).

6R-Hydroxy, S-oxide (S-): **syn-6-Hydroxythiobinupharidine sulfoxide**

$C_{30}H_{42}N_2O_4S$ 526.739

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Amorph. $[\alpha]_D^{20} +3.04$ (c, 0.85 in $CHCl_3$).

6'R-Hydroxy, S-oxide (R-): **Nupharpumilamine A**

[205530-11-6]

$C_{30}H_{42}N_2O_4S$ 526.739

Alkaloid from the rhizomes of *Nuphar pumilum* (Nymphaeaceae). Oil. $[\alpha]_D^{25} -33.4$ (c, 0.6 in $CHCl_3$).

6'R-Hydroxy, S-oxide (S-): **syn-6'-Hydroxythiobinupharidine sulfoxide**

[105857-25-8]

$C_{30}H_{42}N_2O_4S$ 526.739

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Cryst. (C_6H_6). Mp 198-204°. $[\alpha]_D^{20} +9.7$ (c, 0.79 in $CHCl_3$).

6R,6'R-Dihydroxy: **Nuphleine. 6,6'-Dihydroxythiobinupharidine**

[30343-70-5]

$C_{30}H_{42}N_2O_4S$ 526.739

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Shows antifungal activity. Cryst. (MeOH) (as perchlorate). Mp 225-226° (perchlorate).

6S,6'S-Dihydroxy: **Thionupharodiolone**

[51153-11-8]

$C_{30}H_{42}N_2O_4S$ 526.739

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Mp 156-158°.

6R,6'R-Dihydroxy, S-oxide (S-): **syn-6,6'-Dihydroxythiobinupharidine sulfoxide**

$C_{30}H_{42}N_2O_5S$ 542.738

Alkaloid from the rhizomes of *Nuphar luteum* (Nymphaeaceae). Cryst. (C_6H_6). Mp 160-165°. $[\alpha]_D^{20} +39$ (c, 0.79 in $CHCl_3$). Incorrectly indexed in CAS.

6- or 6'-Ethoxy: **Ethoxythiobinupharidine**

[51141-41-4]

$C_{32}H_{46}N_2O_3S$ 538.793

Alkaloid from *Nuphar luteum* (Nymphaeaceae). Mp 270° (as perchlorate). Indexed in *CA* as 6-Ethoxythiobinupharidine, although the exact location of the ethoxy group appears to be unknown.

6ξ,6'ξ-Diethoxy: Diethoxythiobinupharidine

[50764-65-3]

C₃₄H₅₀N₂O₄S 582.846Alkaloid from *Nuphar luteum* (Nymphaeaceae). Mp 230° (as perchlorate).**1-Epimer: 1-Epithiobinupharidine**

[58845-61-7]

C₃₀H₄₂N₂O₂S 494.74Alkaloid from *Nuphar luteum* (Nymphaeaceae). [α]_D²⁵ +23.9 (EtOH) (for mixt. with 1'-epimer). Isol. in admixture with the 1'-epimer.**1'-Epimer: 1'-Epithiobinupharidine**

[58845-62-8]

C₃₀H₄₂N₂O₂S 494.74Alkaloid from *Nuphar luteum* (Nymphaeaceae). Isol. in admixture with the 1-epimer.**7-Epimer: Thionuphlutine B**

[30343-74-9]

Synthetic. Liq. [α]_D²⁵ -131 (c, 0.8 in MeOH).**7-Epimer, S-oxide(R-): Thionuphlutine B α-S-oxide**

[60252-64-4]

C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae).**7-Epimer, S-oxide(S-): Thionuphlutine B β-S-oxide**

[60252-63-3]

C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae).**7-Epimer, 6ξ-hydroxy: 6-Hydroxythionuphlutine B**

[52002-90-1]

C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae). λ_{max} 205 (ε 24400) (acidic 95% EtOH).**7-Epimer, 6'S-hydroxy, S-oxide(R-): Nupharpumilamine B**

[205530-12-7]

C₃₀H₄₂N₂O₄S 526.739Alkaloid from *Nuphar pumilum*. Oil. [α]_D²⁵ -60.1 (c, 1.6 in CHCl₃).**7-Epimer, 6ξ,6'ξ-dihydroxy: 6,6'-Dihydroxythionuphlutine B**

[30343-71-6]

C₃₀H₄₂N₂O₄S 526.739Alkaloid from *Nuphar luteum* (Nymphaeaceae). Glassy solid. [α]_D²⁵ -69 (c, 0.1 in CH₂Cl₂).**1,1'-Diepimer: 1-Epi-1'-epithiobinupharidine**

[58845-63-9]

C₃₀H₄₂N₂O₂S 494.74Alkaloid from *Nuphar luteum* (Nymphaeaceae). [α]_D²⁵ +30.9 (c, 0.9 in 95% EtOH).**1,1'-Diepimer, S-oxide(R-): Nupharpumilamine C**C₃₀H₄₂N₂O₃S 510.739Alkaloid from the rhizomes of *Nuphar pumilum* (Nymphaeaceae). Oil. [α]_D²⁵ -75.1 (c, 0.5 in CHCl₃).**1,1'-Diepimer, S-oxide(S-): Nupharpumilamine D**C₃₀H₄₂N₂O₃S 510.739Alkaloid from the rhizomes of *Nuphar**pumilum* (Nymphaeaceae). Oil. [α]_D²⁵ +20.8 (c, 0.9 in CHCl₃).**7,7'-Diepimer: Neothiobinupharidine**

[4850-09-3]

C₃₀H₄₂N₂O₂S 494.74Alkaloid from *Nuphar luteum* (Nymphaeaceae). Mp 159-160°.**7,7'-Diepimer, diperchlorate:** Mp 320° dec.**7,7'-Diepimer, S-oxide: Neothiobinupharidine sulfoxide**C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae). Cryst. (MeOH/Me₂CO). Mp 240-242°.**7,7'-Diepimer, 6ξ-hydroxy: 6-Hydroxyneothiobinupharidine**

[55869-57-3]

C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae).**7,7'-Diepimer, 6'ξ-hydroxy: 6'-Hydroxyneothiobinupharidine**




[55869-58-4]

C₃₀H₄₂N₂O₃S 510.739Alkaloid from *Nuphar luteum* (Nymphaeaceae).**7,7'-Diepimer, 6ξ,6'ξ-dihydroxy: 6,6'-Dihydroxyneothiobinupharidine**

[58846-67-6]

C₃₀H₄₂N₂O₄S 526.739Alkaloid from *Nuphar luteum* (Nymphaeaceae).

[205530-14-9, 205530-13-8]

Achmatowicz, O. *et al.*, *Tet. Lett.*, 1962, 1121-1124; 1964, 927-934 (*Thiobinupharidine, Neothiobinupharidine, isol, ir, pmr, ms*)Birnbaum, G.I. *et al.*, *Tet. Lett.*, 1965, 4149 (*Neothiobinupharidine, cryst struct*)I'inskaya, T.N. *et al.*, *Khim. Prir. Soedin.*, 1967, 3, 178; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, 3, 147-149 (*Nuphleine*)LaLonde, R.T. *et al.*, *J.O.C.*, 1971, 36, 3703-3706; 1974, 39, 2892-2897 (*6'-Hydroxythiobinupharidine, Thionuphlutine B, 6-Hydroxythionuphlutine B*)Wróbel, J.T. *et al.*, *Can. J. Chem.*, 1972, 50, 1968 (*Neothiobinupharidine sulfoxide*)Wróbel, J.T. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1973, 21, 551-554; 1976, 24, 99-100; *CA*, 79, 137334n; 85, 94579a (*Thionupharodioline, Ethoxythiobinupharidine, Diethoxythiobinupharidine, Thionuphlutine B oxide*)Wróbel, J.T. *et al.*, *Can. J. Chem.*, 1973, 51, 2810-2820 (*cryst struct, abs config, ms, pmr*)Cullen, W.P. *et al.*, *J. Pharm. Sci.*, 1973, 62, 826-827 (*activity*)LaLonde, R.T. *et al.*, *J.A.C.S.*, 1973, 95, 6342-6349 (*ms, pmr, Thionupharoline, Thionuphlutine B derivs*)Wong, C.F. *et al.*, *J.O.C.*, 1973, 38, 3225-3226 (*abs config, Thionuphlutine B*)Martin, T.I. *et al.*, *Can. J. Chem.*, 1974, 52, 2705-2713 (*Thionupharoline*)LaLonde, R.T. *et al.*, *Org. Mass Spectrom.*, 1974, 9, 714-725 (*ms*)LaLonde, R.T. *et al.*, *Can. J. Chem.*, 1975, 53, 1714-1715; 3545-3550 (*cmr, 1-Epithiobinupharidine, 1'-Epithiobinupharidine*)Perel'son, M.E. *et al.*, *Khim. Prir. Soedin.*, 1975, 11, 768-790; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, 11, 789 (*Nuphleine*)Wróbel, J. *et al.*, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1976, 24, 99; *CA*, 85, 94579a (*oxides*)LaLonde, R.T. *et al.*, *J.O.C.*, 1976, 41, 291 (*Hydroxyneothiobinupharidines*)Iwanow, A. *et al.*, *Phytochemistry*, 1986, 25, 2227-2231 (*sulfoxides*)Yoshikawa, M. *et al.*, *Heterocycles*, 1997, 46, 301-308 (*cmr, Nupharpumilamines*)**Thiobutacin****T-381****2-Amino-α-(methylthio)-γ-oxobenzenebutanoic acid. 4-(2-Aminophenyl)-2-(methylthio)-4-oxobutanoic acid**C₁₁H₁₃NO₃S 239.295Prod. by *Lechevalieria aerocolonigenes* strain VK-A9. Antifungal and antimycotic agent. Brown solid. λ_{max} 227; 260; 360 (MeOH).Lee, J.Y. *et al.*, *J. Nat. Prod.*, 2004, 67, 2076-2078 (*isol, pmr, cmr*)Chakor, N. *et al.*, *Tet. Lett.*, 2008, 49, 5056-5058 (*synth*)**Perophora viridis 1,2,3-Thiocane glycoside****T-382**C₃₀H₄₈N₂O₉S₄ 708.981Isol. from the Atlantic tunicate *Perophora viridis*. Pale yellow amorph. solid. [α]_D²⁴ +197.3 (c, 0.11 in EtOH). λ_{max} 257 (log ε 2.78) (EtOH).Rezanka, T. *et al.*, *Eur. J. Org. Chem.*, 2002, 2400-2404 (*isol, pmr, cmr, ms*)**Thiocillin I****T-383****13',19'-Didehydro-19'-deoxy-28,44-dihydro-41,44-dihydroxymicrococccin P, 9CI** [59979-01-0]

C₄₈H₄₉N₁₃O₁₀S₆ 1160.392

Thiopeptide antibiotic. Similar to Micrococins. Isol. from *Bacillus badius* and *Bacillus cereus*. Active against gram-positive bacteria. Amorph. powder. Sol. DMF, DMSO, THF, CHCl₃-MeOH; fairly sol. MeOH, CHCl₃, Me₂CO, EtOH; poorly sol. EtOAc, C₆H₆, Et₂O, H₂O, acids. [α]_D^{24.5} +97.7 (c, 2.028 in 90% EtOH aq.). λ_{max} 217 (E1%/1cm 734); 275 (E1%/1cm 326); 348 (E1%/1cm 199) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) 500 - 1500 mg/kg. XK6510000

26-Me ether: **Thiocillin II**

[59979-02-1]

C₄₉H₅₁N₁₃O₁₀S₆ 1174.419

From *Bacillus badius* and *Bacillus cereus*. Amorph. powder. Sol. DMSO, CHCl₃-MeOH, THF, DMF; fairly sol. MeOH, EtOH, CHCl₃, Me₂CO; poorly sol. EtOAc, H₂O, acids, Et₂O, C₆H₆. [α]_D^{24.5} +93.4 (c, 0.59 in 90% EtOH aq.). λ_{max} 217 (E1%/1cm 705); 278 (E1%/1cm 291); 348 (E1%/1cm 202) (MeOH) (Berdy).

► LD₅₀ (mus, ipr) -400 - 600 mg/kg. XK6520000

34-Me ether: **Thiocillin III**

[59979-03-2]

C₄₉H₅₁N₁₃O₁₀S₆ 1174.419

From *Bacillus badius*. Amorph. powder. Sol. DMF, THF, CHCl₃-MeOH, DMSO; fairly sol. MeOH, CHCl₃, EtOH, Me₂CO; poorly sol. EtOAc, Et₂O, H₂O, acids, C₆H₆. [α]_D^{24.5} +88 (c, 0.85 in 90% EtOH aq.). λ_{max} 217 (E1%/1cm 729); 275 (E1%/1cm 302); 347 (E1%/1cm 208) (MeOH) (Berdy).

2-Ketone: **Antibiotic YM 266183**. YM 266183

C₄₈H₄₇N₁₃O₁₀S₆ 1158.376

Prod. by the marine-derived *Bacillus cereus* QN03323. Active against gram-positive bacteria. Powder. [α]_D²⁵ +64.7 (c, 0.37 in MeOH). λ_{max} 213 (ε 79830); 290 (sh); 344 (ε 14460) (MeOH).

2-Ketone, 26-Me ether: **Antibiotic YM 266184**. YM 266184

C₄₉H₄₉N₁₃O₁₀S₆ 1172.403

Prod. by the marine-derived *Bacillus cereus* QN03323. Active against gram-positive bacteria. Powder. [α]_D²⁵ +60.9 (c, 0.15 in MeOH). λ_{max} 215 (ε 100960); 290 (sh); 346 (ε 29350) (MeOH).

Shoji, J. *et al.*, *J. Antibiot.*, 1976, **29**, 366-374;

1981, **34**, 1126-1136 (*isol, struct*);


Nagai, K. *et al.*, *J. Antibiot.*, 2003, **56**, 123-128; 129-134 (*YM 266183, YM 266184*)

Thiocoraline

T-384

PM 93135. *Antibiotic PM 93135*

[173046-02-1]



C₄₈H₅₆N₁₀O₁₂S₆ 1157.426

Depipeptide antibiotic. Prod. by a marine *Micromonospora* sp. L-13-ACM2-092. Cytotoxic agent. RNA synthesis inhibitor. Weakly inhibits HIV-1 reverse transcriptase. Pale yellow cryst. Mp 266-266.5°. [α]_D²⁵ -190.9 (c, 1 in CHCl₃). λ_{max} 218 (ε 76500); 230 (ε 76700); 298 (ε 9400); 360 (ε 8900) (MeOH).

Romero, F. *et al.*, *J. Antibiot.*, 1997, **50**, 734-737 (*isol, props*)

Baz, J.P. *et al.*, *J. Antibiot.*, 1997, **50**, 738-741 (*uv, ir, pmr, cmr*)


Boger, D.L. *et al.*, *J.A.C.S.*, 2001, **123**, 561-568 (*synth, abs config, activity*)

Lombó, F. *et al.*, *ChemBioChem*, 2006, **7**, 366-367 (*biosynth*)

Thiomarinol A

T-385

[146697-04-3]



C₃₀H₄₄N₂O₉S₂ 640.818

Prod. by the marine bacterium *Alteromonas rava*. Active against gram-positive and -negative bacteria. Inhibitor of isoleucyl transferase and RNA synthetase. Orange cryst. (MeOH). Sol. MeOH, DMSO, DMF, Et₂O; poorly sol. hexane, H₂O. Mp 106-110° dec. [α]_D²⁵ +4.3 (c, 1 in MeOH). Related to Pseudomononic acid A. λ_{max} 206 (ε 25000); 306 (ε 3200); 386 (ε 9600) (MeOH/NaOH) (Derep). λ_{max} 214 (ε 26000); 300 (ε 3500); 387 (ε 12000) (MeOH) (Derep). λ_{max} 214; 300; 387 (MeOH/HCl) (Berdy).

1'',1''-Dioxide: **Thiomarinol B**

[156098-42-9]

C₃₀H₄₄N₂O₁₁S₂ 672.816

Prod. by *Alteromonas rava*. Active against gram-positive and -negative bacteria. Inhibitor of isoleucyl transferase and RNA synthetase. Yellow cryst. (MeOH aq.). Sol. MeOH, DMSO, EtOAc, Et₂O, DMF, butanol, 1-propanol; poorly sol. H₂O, hexane. [α]_D²⁵ +7.7 (c, 1 in propanol). λ_{max} 215 (ε 21000); 301 (ε 13000); 377 (2900) (MeOH). λ_{max} 221 (ε 19000); 301 (ε 13000); 377 (ε 2900) (MeOH/NaOH) (Berdy).

13-Ketone: **Thiomarinol F**

[182155-01-7]

C₃₀H₄₂N₂O₉S₂ 638.802

Prod. by *Alteromonas rava*. Inhibitor of isoleucyl transferase and RNA synthetase. [α]_D²⁵ -1.66 (c, 0.8 in MeOH). λ_{max} 210 (ε 21200); 300 (ε 2600); 385 (ε 8500) (MeOH).

4-Deoxy: **Thiomarinol C**

[156343-39-4]

C₃₀H₄₄N₂O₈S₂ 624.818

Prod. by *Alteromonas rava*. Active

against gram-positive and -negative bacteria. Inhibitor of isoleucyl transferase and RNA synthetase. Yellow cryst. Sol. MeOH, DMF, Et₂O, EtOAc, DMSO, 1-propanol, butanol; poorly sol. H₂O, hexane. [α]_D²⁵ -1.4 (c, 1 in MeOH). λ_{max} 215 (ε 17000); 300 (ε 2700); 388 (9600) (MeOH). λ_{max} 205 (ε 49000); 336 (ε 8600) (MeOH/NaOH) (Berdy).

4,6-Dideoxy, 8-hydroxy: **Thiomarinol G**

[194154-33-1]

C₃₀H₄₄N₂O₈S₂ 624.818

Prod. by *Alteromonas rava*. λ_{max} 210;

300; 385 (MeOH).

[156098-43-0]

Shiozawa, H. *et al.*, *J. Antibiot.*, 1993, **46**, 1834; 1995, **48**, 907; 1997, **50**, 449-452 (*isol, uv, ir, pmr, cmr, props*)

Gao, X. *et al.*, *J.A.C.S.*, 2005, **127**, 1628-1629 (*synth*)

Thiomarinol D

T-386

[182154-99-0]

As Thiomarinol A, T-385 with

R = CH₃, n = 1

C₃₁H₄₆N₂O₉S₂ 654.844

Stereochemical identity with Thiomarinol A, T-385 not confirmed. Prod. by *Alteromonas rava*. Inhibitor of isoleucyl transferase and RNA synthetase. [α]_D²⁵ +1.5 (c, 0.8 in MeOH). λ_{max} 210 (ε 20900); 300 (ε 2600); 385 (ε 8900) (MeOH).

Shiozawa, H. *et al.*, *J. Antibiot.*, 1997, **50**, 449-452 (*isol, uv, ir, pmr, cmr*)

Thiomarinol E

T-387

[182155-00-6]

As Thiomarinol A, T-385 with

R = H, n = 3

C₃₂H₄₈N₂O₉S₂ 668.871


Stereochemical identity with Thiomarinol A, T-385 not confirmed. Prod. by *Alteromonas rava*. λ_{max} 210; 300; 385 (MeOH).

Shiozawa, H. *et al.*, *J. Antibiot.*, 1997, **50**, 449-452 (*isol, uv, pmr, cmr*)

Thiomycolide A

T-388

[207227-80-3]



C₅₇H₈₁N₇O₂₀S 1216.368

Isol. from a sponge *Mycale* sp. Cytotoxic agent. λ_{max} 238 (MeOH) (Berdy).

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 663-666 (*isol, pmr*)

Thiomycolide B

T-389

[207227-81-4]

As Thiomycolide A, T-388 with

R = H, –OOCCH(OMe)CH₂OMe

C₆₂H₉₁N₇O₂₃S 1334.5

Isol. from a sponge *Mycale* sp.

Cytotoxic agent. Brownish powder. [α]_D –3.3 (c, 0.1 in MeOH). λ_{\max} 238 (ε 33500) (MeOH).

Matsunaga, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 663-666 (isol, uv, pmr, cmr, ms)

Thiomycin**T-390**

B 870. Antibiotic B 870

[1393-47-1]

Pyrrhothine-related antibiotic. Prod. by *Streptomyces phaeochromogenes-chloromyceticus*. Antibacterial agent, platelet aggregation inhibitor. Golden-yellow needles (EtOAc). Sol. MeOH, C₆H₆; fairly sol. H₂O; poorly sol. Et₂O, hexane. Mp 176-178° dec. λ_{\max} 223 (E1%/1cm 140); 305; 370 (E1%/1cm 375) (MeOH) (Berdy). λ_{\max} 370 (E1%/1cm 690) (HCl) (Berdy). λ_{\max} 300 (E1%/1cm 555) (NaOH) (Berdy).

▶ LD₅₀ (mus, scu) 10 mg/kg.

Hinuma, Y. *et al.*, *J. Antibiot., Ser. A*, 1955, **8**, 118-119; *CA*, **53**, 15201a

Thiosporamicin**T-391**

Antibiotic 46192. Antibiotic CP 46192.

CP 46192

[67383-02-2]

C₇₅H₈₈N₁₈O₂₁S₆ 1770.024

Thiazole-peptide antibiotic. Struct. unknown. Prod. by *Streptosporangium roseum-incarnatum*. Active against gram-positive bacteria. Cryst. (MeOH). Sol. CHCl₃, EtOAc; fairly sol. MeOH; poorly sol. H₂O, hexane. Mp 210-216°. [α]_D –129 (c, 1 in CHCl₃). λ_{\max} 238 (E1%/1cm 350); 303 (E1%/1cm 80) (MeOH) (Berdy).

U.S. Pat., 1978, 4 083 963; *CA*, **89**, 105892h


Thiostrepton, 9CI, 8CI**T-392**

Thiostrepton A. Bryamycin. Thiactin.

Antibiotic 6739-15. Antibiotic 6761-31.

Antibiotic A 43791. A 43791

[1393-48-2]



C₇₂H₈₅N₁₉O₁₈S₅ 1664.91

Prod. by *Streptomyces azureus* ATCC

14921, *Streptomyces hawaiiensis*

ATCC 12236 and *Streptomyces*

laurentii ATCC 31255. Exhibits

antimalarial and cytotoxic props. and

active against gram-positive bacteria.

Protein synthesis inhibitor. Cryst.

(CHCl₃/EtOH). Mp 246-255° dec.

[α]_D²³ –98.5 (AcOH). λ_{\max} 240 (sh);

245 (ε 50000); 280 (sh); 294 (ε 18000);

305 (sh); 306 (ε 14900) (MeOH)

(Derep).

▶ XN6300100

26-Thione, 1,2R-dihydro: Antibiotic Sch 18640. Sch 18640

[79053-93-3]

C₇₂H₈₇N₁₉O₁₇S₆ 1682.992

Isol. from *Micromonospora arborensis* NRRL8041; major component of 68-1147 antibiotic complex. Active against gram-positive and anaerobic bacteria. Fairly sol. butanol; poorly sol. hexane, H₂O. Mp 225-229°. [α]_D²⁶ +81.8 (c, 0.3 in CHCl₃). λ_{\max} 240 (sh); 245 (ε 50000); 280 (sh); 294 (ε 18000); 305 (sh); 306 (ε 14900) (MeOH) (Derep).

7'-Parent amide: Thiostrepton B. Thiostrepton A₂

[75524-61-7]

C₆₆H₇₉N₁₇O₁₆S₅ 1526.784

An artifact formed from

Thiostrepton A on standing. Sol.

CHCl₃, Py; fairly sol. MeOH;

poorly sol. H₂O. λ_{\max} 245 (sh) (ε

60000); 290 (ε 20000) (MeOH) (De-

rep).

Tori, K. *et al.*, *Tet. Lett.*, 1976, **17**, 185-188

(pmr, cmr)

Florey, K. *et al.*, *Anal. Profiles Drug Subst.*,

1978, **7**, 423 (rev)

U.S. Pat., 1978, 4 078 056; *CA*, **89**, 105873c

(Sch 18640)

Puar, M.S. *et al.*, *J.A.C.S.*, 1981, **103**, 5231-

5233 (Sch 18640, struct)

Hensens, O.D. *et al.*, *J. Antibiot.*, 1983, **36**,

799-813; 814-831; 831-845 (struct, bibl)

Mocek, U. *et al.*, *J. Antibiot.*, 1989, **42**, 1649-

1652 (pmr, cmr)

Mocek, U. *et al.*, *J.A.C.S.*, 1993, **115**, 7992-

8001 (biosynth)

Bond, C.S. *et al.*, *Acta Cryst. D*, 2001, **57**, 755-

758 (cryst struct)

Nicolaou, K.C. *et al.*, *J.A.C.S.*, 2005, **127**,

11159-11175; 11176-11183 (synth, pmr)


Lewis, R.J. *et al.*, *Sax's Dangerous Properties*

of Industrial Materials, 8th edn., Van

Nostrand Reinhold, 1992, TFQ275

Thiotipin**T-393**

[158792-26-8]



C₅₅H₅₀N₁₆O₁₇S₂ 1271.229

Cyclic peptide antibiotic related to

Sulfomycin I in S-638. Prod. by

Streptomyces sp. DT31. TipA promotor

inducer. Powder. Sol. EtOAc, MeOH,

CHCl₃; poorly sol. Et₂O, hexane, H₂O.

Mp 265-270° dec. [α]_D –9.5 (c, 0.1 in

CHCl₃/MeOH). λ_{\max} 215 (sh) (ε 53000);

247 (ε 67000); 315 (sh) (ε 9000)

(EtOH).


Yun, B.-S. *et al.*, *Tetrahedron*, 1994, **50**, 11659

(isol, uv, ir, pmr, cmr)

2-Thioxo-4-thiazolidinecarboxylic acid**T-394**

4,5-Dihydro-2-thioxo-4(3H)-thiazolecarboxylic acid. 4-Carboxy-2-thiazolidinethione

[20933-67-9]



(S)-form

C₄H₅NO₂S₂ 163.221

(R)-form

Raphanusamic acid

[98169-56-3]

Isol. from etiolated seedlings of *Raphanus*

sativus var. *hortensis* (Japanese radish

Daikon). Plant growth inhibitor. [α]_D²⁹ –86

(c, 2.5 in 0.5M HCl).

Me ester: [80963-80-0]

C₅H₇NO₂S₂ 177.248

Light yellow oil. [α]_D²⁰ –68.2 (CHCl₃).

(S)-form

Et ester: [157832-46-7]

C₆H₉NO₂S₂ 191.275

Orange solid. Mp 44-46°.

(±)-form

Me ester: [114180-04-0]

Prisms (CH₂Cl₂). Mp 95-96°.

[101799-35-3; 64014-03-5]

Hase, T. *et al.*, *Rev. Latinoam. Quim.*, 1985, **16**,

1 (synth, isol, props)

Hsiao, C.-N. *et al.*, *J.O.C.*, 1987, **52**, 2201

(synth, ester, ir, pmr)

Nagao, Y. *et al.*, *Chem. Pharm. Bull.*, 1988, **36**,

495 (synth, ester, uv, pmr, cmr)

Inamori, Y. *et al.*, *Chem. Pharm. Bull.*, 1992,

40, 2854 (synth, activity)

U.S. Pat., 1993, 5 210 094; *CA*, **119**, 22809y

(use)

Calo, V. *et al.*, *Tetrahedron*, 1994, **50**, 7283


(synth, ester, pmr)

Dobson, A.J. *et al.*, *Acta Cryst. C*, 1998, **54**,

1634-1637 (R-form, cryst struct)

Thomasic acid**T-395**

3,4-Dihydro-6-hydroxy-4-(4-hydroxy-3,5-dimethoxyphenyl)-3-hydroxymethyl-5,7-dimethoxy-2-naphthalenecarboxylic acid, 9CI



C₂₂H₂₄O₉ 432.426

Possible artifact.

(ξ)-form

9'-Carboxylic acid, bis(4-hydroxyphenylethylamide):

$C_{38}H_{40}N_2O_{10}$ 684.741


Alkaloid from *Porcelia macrocarpa*. Amorph. powder. $[\alpha]_D^{25}$ -20 (c, 0.06 in MeOH). λ_{max} 210 (log ϵ 4.87); 240 (log ϵ 4.5); 282 (log ϵ 4.11); 325 (log ϵ 4.23) (MeOH).

Chaves, M.H. *et al.*, *Phytochemistry*, 1997, **46**, 879-881 (amide deriv)

Thorectandramine

[459165-95-8]

T-396



$C_{27}H_{24}N_3O_3^+$ 438.505


Quaternary alkaloid isol. from the marine sponge *Thorectandra* sp. Yellow solid (as NH_4 salt). $[\alpha]_D^{25}$ +4.9 (c, 0.08 in MeOH) (NH_4 salt). λ_{max} 224 (log ϵ 4.16); 270 (log ϵ 3.97); 338 (log ϵ 3.87); 407 (log ϵ 3.22) (MeOH) (NH_4 salt).

Charan, R.D. *et al.*, *Tet. Lett.*, 2002, **43**, 5201-5204 (isol, pmr, cmr)

Thuggacin A

[190895-98-8]

T-397



$C_{35}H_{53}NO_7S$ 631.872

Prod. by *Sorangium cellulosum* strain So ce895. Antimycobacterial agent. Cryst. (Et_2O /petrol). Mp 92-94°. $[\alpha]_D^{22}$ -148.4 (c, 0.4 in MeOH). λ_{max} 224 (log ϵ 4.74); 231 (sh); 239 (sh); 289 (log ϵ 4.11) (MeOH).

13-Methyl-13-Methylthuggacin A

[949574-78-1]

$C_{36}H_{55}NO_7S$ 645.899

Prod. by *Sorangium cellulosum* So ce895. $[\alpha]_D^{22}$ -127.7 (c, 0.94 in MeOH). Stereochem. not determined. λ_{max} 227 (log ϵ 4.62); 286 (log ϵ 4.04) (MeOH).

Pat. Coop. Treaty (WIPO), 1997, 18 217; *CA*, **127**, 32937 (isol, pmr, cmr, ms)


Steinmetz, H. *et al.*, *Chem. Eur. J.*, 2007, **13**, 5822-5832 (isol, pmr, cmr, ms, biosynth)

Bock, M. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 2308-2311 (abs config)

Thuggacin B

[190895-99-9]

T-398



Absolute Configuration

$C_{35}H_{53}NO_7S$ 631.872

Prod. by *Sorangium cellulosum* So ce895. $[\alpha]_D^{22}$ +88.3 (c, 1.1 in MeOH). λ_{max} 226 (log ϵ 4.7); 231 (sh); 240 (sh); 287 (log ϵ 4.13) (MeOH).

Pat. Coop. Treaty (WIPO), 1997, 18 217; *CA*, **127**, 32937 (isol, pmr, cmr, ms)


Steinmetz, H. *et al.*, *Chem. Eur. J.*, 2007, **13**, 5822-5832 (isol, pmr, cmr, ms)

Bock, M. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 2308-2311 (abs config)

Thuggacin C

[949574-77-0]

T-399



Absolute Configuration

$C_{35}H_{53}NO_7S$ 631.872

Prod. by *Sorangium cellulosum* strain So ce895. $[\alpha]_D^{22}$ -35.7 (c, 0.7 in MeOH). Rearr. prod. of Thuggacin A, T-397. λ_{max} 228 (log ϵ 4.55); 286 (log ϵ 4.05) (MeOH).


Steinmetz, H. *et al.*, *Chem. Eur. J.*, 2007, **13**, 5822-5832 (isol, pmr, cmr)

Bock, M. *et al.*, *Angew. Chem., Int. Ed.*, 2008, **47**, 2308-2311 (abs config)

Thuggacin cmc-A

[949574-79-2]

T-400



$C_{30}H_{43}NO_7S$ 561.738

Prod. by *Chondromyces crocatus* strain Cm c5. Antimycobacterial agent. $[\alpha]_D^{22}$ -160.6 (c, 1.24 in MeOH). λ_{max} 224 (log ϵ 4.67); 285 (log ϵ 4.08) (MeOH).

(1→18)-Lactone isomer: **Thuggacin cmc-C**

[949574-80-5]

$C_{30}H_{43}NO_7S$ 561.738

Prod. by *Chondromyces crocatus* strain Cm c5. $[\alpha]_D^{22}$ -47.4 (c, 0.5 in MeOH). λ_{max} 227 (log ϵ 4.64); 285 (log ϵ 4.13) (MeOH).

(1→17)-Lactone isomer: **Thuggacin cmc-B**

$C_{30}H_{43}NO_7S$ 561.738

Prod. by *Chondromyces crocatus* strain Cm c5. Struct. not confirmed.

Steinmetz, H. *et al.*, *Chem. Eur. J.*, 2007, **13**, 5822-5832 (isol, pmr, cmr, ms)


Thymine, 8CI

T-401

5-Methyl-2,4(1H,3H)-pyrimidinedione, 9CI. 4-Hydroxy-5-methyl-2(1H)-pyrimidinone, 9CI. 5-Methyluracil. 2,4-Dihydroxy-5-methylpyrimidine. 5-Methyl-2,4-pyrimidinediol

[65-71-4]

[27942-00-3]



$C_5H_6N_2O_2$ 126.115

Dioxo tautomer predominates in solid, soln. and gas phases. Intermediate tautomers also possible. Component of nucleic acids. Also detected in the free state in bacteria. Used as a 0.25% aq. soln for pptn. sepn. of Hg. Plates (H_2O) or parallelepipeds + H_2O (H_2O). Spar. sol. H_2O (0.6-0.8 g per 100 cm^3). Mp 326° Mp 340°. pK_{a1} 9.9 (25°).

▶ LD₅₀ (mus, orl) 3500 mg/kg. XP2100000

N^1 -(2-Hydroxyethyl): N^1 -(2-Hydroxyethyl)thymine. **Thyminol**

[22441-51-6]

$C_7H_{10}N_2O_3$ 170.168

Isol. from a colonial Zoanthid. Mp 179-181°. λ_{max} 263 (log ϵ 2.99) (MeOH).

[41935-71-1]

Rice, J.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4569 (ms)

Ozeki, K. *et al.*, *Acta Cryst. B*, 1969, **25**, 1038 (cryst struct)

Wong, J.L. *et al.*, *J.O.C.*, 1970, **35**, 3786 (synth, pmr, uv)

Ellis, P.D. *et al.*, *J.A.C.S.*, 1973, **95**, 4398 (cmr)

Les, A. *et al.*, *Spectrochim. Acta A*, 1992, **48**, 1385 (ir)

Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1999, **13**, 255-261 (Thyminol)

Tienmulilmine

T-402

$C_{27}H_{43}NO$ 397.643

Steroidal alkaline alkaloid. Struct. unknown. Isol. from the Chinese drug Li-Lu (*Veratrum* spp.) (Liliaceae). Mp 172-174°. $[\alpha]_D^{16}$ -99.3 (MeOH).

Hydrochloride: Mp 302°.

Hydrobromide: Mp 306°.

Methiodide: Mp 244°.

Chu, J.-H. *et al.*, *CA*, 1958, **52**, 6716

Chu, J.-H. *et al.*, *CA*, personal communication cited by Tomko, J. *et al.*, *Alkaloids*, N.Y., 1973, **14**, 32

Tienmuliminine T-403

C₃₄H₅₁NO₈ 601.779

Struct. unknown. Alkaloid from the Chinese drug Li-Lu (*Veratrum*) (Liliaceae). Mp 231°. [α]_D¹³ -24 (c, 1 in MeOH).

Hydrobromide: Mp 226°.

Hydrochloride: Mp 215°.


Picrate: Mp 201° dec.

Methiodide: Mp 224° dec.

Chu, J.-H. *et al.*, *Huaxue Xuebao*, 1956, **52**, 229-304; *CA*, **52**, 6716f

3-Tigloyloxytropane T-404

2-Methyl-2-butenic acid 8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester, 8CI. Tropine tiglate



C₁₃H₂₁NO₂ 223.314

Antiparkinsonian, anticholinergic agent. CNS depressant. Log P 2.03 (calc). Both α - and β -forms reported to possess pharmacol. activity.

3 α -form

Tigloyltropine. Tropigline, BAN, INN [495-83-0]

Alkaloid from *Datura ferox* and other *Datura* spp., *Physalis alkekengi* (winter cherry) roots, *Mandragora vernalis*, *Mandragora autumnalis*, *Hyoscyamus* spp., etc. (Solanaceae). Mp 181.5-183° Mp 204-207°.

Hydrochloride: Mp 214.5-217.5°.

N-Oxide: 3 α -Tigloyloxytropane *N-oxide*

C₁₃H₂₁NO₃ 239.314

Alkaloid from *Physalis alkekengi* (winter cherry) (Solanaceae). Mp 185-186° dec.

3 β -form

Tigloidine, BAN, INN

[533-08-4]

Alkaloid from *Datura innoxia*, *Datura metel*, *Datura pruinosa*, *Datura meteloides*, *Physalis alkekengi* (winter cherry), *Mandragora* spp. and *Duboisia myoporoides* (Solanaceae). Syrup.

Hydrochloride:

Tabular cryst. (H₂O). Sol. CHCl₃, mod. sol. H₂O. Mp 234-235°.

Picrate: Mp 239°.

Barger, G. *et al.*, *J.C.S.*, 1937, 1820 (*isol*, *synth*) Sanghvi, I.S. *et al.*, *Eur. J. Pharmacol.*, 1968, **4**, 246 (*pharmacol*)

Evans, W.C. *et al.*, *J.C.S. Perkin 1*, 1972, 2017 (*isol*)

Evans, W.C. *et al.*, *Phytochemistry*, 1973, **12**, 2077; 2505 (*isol*)

Beresford, P.J. *et al.*, *Phytochemistry*, 1974, **13**, 2143; 1975, **14**, 2205 (*isol*, *biosynth*)

Yamaguchi, H. *et al.*, *Yakugaku Zasshi*, 1974, **94**, 1115 (*oxide*)

Martindale, *The Extra Pharmacopoeia*, 28th/29th edn., Pharmaceutical Press, 1982, 419

Witte, L. *et al.*, *Planta Med.*, 1987, **53**, 192 (*glc-ms*)

Tiliacoridine T-405

[11076-69-0]


C₃₉H₄₀N₂O₈ 664.754

Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the leaves of *Tiliacora racemosa* (Menispermaceae). Mp 153-156° dec.

Barua, A.K. *et al.*, *J. Indian Chem. Soc.*, 1970, **47**, 920 (*isol*, *uv*, *pmr*, *ms*)

Tiliacorine T-406

[27073-72-9]



C₃₆H₃₆N₂O₅ 576.691

Diastereomeric with Tiliacorinine, T-407. Alkaloid from the roots of *Tiliacora racemosa* and *Tiliacora funifera* (Menispermaceae). Cryst. + 1Me₂CO (CHCl₃/Me₂CO). Mp 262-264° dec. [α]_D +71.2 (c, 1.5 in Py).

Ac:

Needles (Et₂O). Mp 232-235°. [α]_D²¹ +86.73 (c, 1.13 in CHCl₃).

N,N'-*Di-Me*:

C₃₈H₄₂N₂O₅[⊕] 606.76

Needles + 2H₂O (Me₂CO/MeOH) (as diiodide). Mp 294-295° dec. (diiodide). [α]_D +136.7 (c, 0.3 in H₂O).

2'-*N-De-Me*: *Nortiliacorine A*. *Isotiliarine* [27577-49-7]

C₃₅H₃₄N₂O₅ 562.664

Alkaloid from the roots of *Tiliacora funifera* (Menispermaceae). Mp 258-260°. [α]_D +194.5 (c, 0.77 in CHCl₃).

Me ether: Obt. by methylation of both Tiliacorine and Dinklacorine. Cryst. (Et₂O). Mp 210-212° dec. [α]_D²¹ +38.5 (c, 0.65 in CHCl₃) (+32.6).

Me ether, *N,N'*-*di-Me*: Obt. by methylation of both Tiliacorine and Dinklacorine. Pale yellow needles (MeOH) (as diiodide). Mp 310° (diiodide). [α]_D³⁰ +113.6 (c, 0.22 in MeOH) (+108.3).

O¹²-*De-Me*, O^{12'}-*Me*: *Dinklacorine*.

Alkaloid TD-2

[60579-86-4]

C₃₆H₃₆N₂O₅ 576.691

Alkaloid from the roots of *Tiliacora dinklagei* (Menispermaceae). Cryst. (Et₂O/EtOAc). Mp 238-240°. [α]_D²¹ +42.55 (c, 0.47 in CHCl₃).

O¹²-*De-Me*, O^{12'}-*Me*, *Ac*: Mp 168-170°. [α]_D¹ +20.45 (c, 0.88 in CHCl₃).

5-Hydroxy, N²-*de-Me*: *Norisoyanangine* [119067-89-9]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from aerial parts of *Tiliacora triandra* (Menispermaceae). Cryst. + 1CH₂Cl₂. [α]_D²⁰ +139 (c, 1.2 in CHCl₃).

Tackie, A.N. *et al.*, *Planta Med.*, 1968, **16**, 158-165 (*Tiliacorine*, *Nortiliacorine A*, *isol*)

Anjaneyulu, B. *et al.*, *Tetrahedron*, 1969, **25**, 3091-3105; 1971, **27**, 439-443 (*Tiliacorine*, *isol*, *uv*, *ms*, *pmr*, *struct*, *synth*)

Baldas, J. *et al.*, *J.C.S. Perkin 1*, 1972, 592-596 (*ms*)

Tackie, A.N. *et al.*, *Phytochemistry*, 1973, **12**, 203-205 (*Nortiliacorine A*)

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1975, **38**, 210-212 (*Dinklacorine*, *isol*)

Dwuma-Badu, D. *et al.*, *J. Nat. Prod.*, 1976, **39**, 213-217; 1979, **42**, 116-119 (*Tiliacorine*, *Dinklacorine*, *uv*, *ir*, *pmr*, *ms*, *cd*, *struct*)


Shamma, M. *et al.*, *J.O.C.*, 1976, **41**, 1293-1294 (*struct*)

Bhakuni, D.S. *et al.*, *J.C.S. Perkin 1*, 1981, 2598-2603 (*abs config*, *biosynth*)

Pachaly, P. *et al.*, *Planta Med.*, 1988, **54**, 433-437 (*Norisoyanangine*)

Tiliacorinine T-407

[27073-73-0]



C₃₆H₃₆N₂O₅ 576.691

Diastereomeric with Tiliacorine, T-406. Alkaloid from the roots of *Tiliacora racemosa*, *Tiliacora dinklagei* and *Tiliacora triandra* (Menispermaceae). Cryst. + 1Me₂CO (Me₂CO). Mp 195° dec. (shrinks above 172°). [α]_D +310 (c, 2.6 in Py).

Ac:

Cryst. (CH₂Cl₂/hexane). Mp 170-172° dec. [α]_D +363.8 (c, 2.5 in Py).

2'-*N-Oxide*: *Tiliacorinine 2'-N-oxide*

[80161-67-7]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from the roots of *Tiliacora triandra* (Menispermaceae). Pale yellow glass. Mp 215-217° dec. [α]_D¹⁹ +238.3 (c, 1.1 in CHCl₃).

N,N'-*Di-Me*:

Needles (MeOH) (as diiodide). Mp 300° (diiodide). [α]_D +194.7 (c, 0.3 in H₂O).

N²-*De-Me*: *Nortiliacorinine A*. *Pseudotiliarine*

[26426-60-8]

C₃₅H₃₄N₂O₅ 562.664

Alkaloid from the roots of *Tiliacora funifera*, *Tiliacora racemosa*, *Tiliacora dinklagei* and *Tiliacora triandra* (Menispermaceae). Mp 220-222° Mp 252-254° dec Mp 264-268° dec.

N²-*De-Me*: *Nortiliacorinine B*

[26777-94-6]

C₃₅H₃₄N₂O₅ 562.664

A minor alkaloid from the roots of *Tiliacora racemosa* (Menispermaceae).

Cryst. + 1H₂O (Me₂CO/MeOH). Mp 218-220° dec. [α]_D²⁵ +356.2 (c, 1.25 in Py).

Di-N-de-Me: Pachyovatamine

[96910-84-8]

C₃₄H₃₂N₂O₅ 548.637

Alkaloid from the leaves and stems of *Pachygone ovata* (Menispermaceae). Amorph. Mp 182-185°. [α]_D²⁵ +259 (c, 0.29 in CHCl₃).

Me ether, N,N'-di-Me:

Cryst. + 3MeOH (MeOH) (as diiodide). Mp 270-275° dec. (diiodide).

O¹²-De-Me, O^{12'}-Me: Yanangcorinine

[102571-99-3]

C₃₆H₃₆N₂O₅ 576.691

Alkaloid from *Tiliacora triandra* (Menispermaceae). Amorph. + 1CH₂Cl₂. Mp 174-180°. [α]_D¹⁹ +368 (c, 0.6 in CHCl₃).

O¹²-De-Me, O^{12'}-Me, N^{2'}-de-Me: Tiliarine

[7221-73-0]

C₃₅H₃₄N₂O₅ 562.664

Alkaloid from the roots of *Tiliacora racemosa* (Menispermaceae). Mp 203-207°. [α]_D²⁵ +283.6 (MeOH).

Rao, K.V.J. et al., *J. Sci. Ind. Res., Sect. B*, 1959, **18**, 247 (Tiliarine, isol)

Tackie, A.N. et al., *Planta Med.*, 1968, **16**, 158 (Tiliacarinine, isol)

Anjaneyulu, B. et al., *Tetrahedron*, 1969, **25**, 3091 (isol, uv, pmr, Nortiliacarinines)

Tackie, A.N. et al., *J. Nat. Prod.*, 1975, **38**, 210 (Nortiliacarinine A)

Shamma, M. et al., *J.O.C.*, 1976, **41**, 1293 (struct)

Wiryachitra, P. et al., *Aust. J. Chem.*, 1981, **34**, 2001 (Tiliacarinine N-oxide)


Bhakuni, D.S. et al., *J.C.S. Perkin I*, 1981, 2598 (abs config, biosynth)

Bhakuni, D.S. et al., *Tetrahedron*, 1981, **37**, 2651 (Nortiliacarinines)

Guinaudeau, H. et al., *J. Nat. Prod.*, 1985, **48**, 651 (Tiliarine, uv, pmr, cd, ms, struct)

Sultanbawa, M.U.S. et al., *Phytochemistry*, 1985, **24**, 589 (Pachyovatamine)

Pachaly, P. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1986, **319**, 126 (Yanangcorinine)

Tiliapunimine**T-408**C₃₆H₃₆N₂O₆ 592.69**(S)-form**

Alkaloid from the leaves of *Tiliacora junifera* (Menispermaceae). Needles. Mp 198-200°. [α]_D²⁵ +294 (c, 0.52 in CHCl₃).

Ayim, J.S.K. et al., *J. Nat. Prod.*, 1977, **40**, 561 (isol, uv, ir, pmr, ms, struct)

Tiliandrine**T-409**C₃₄H₃₄N₂O₅ 550.653


Bisbenzylisoquinoline alkaloid. Struct. unknown. Alkaloid from the roots of *Tiliacora triandra* (Menispermaceae). Mp

175°. [α]_D²⁵ +408 (EtOH).

Paris, R.R. et al., *Ann. Pharm. Fr.*, 1967, **25**, 627; *CA*, **68**, 33140c (isol)

Tiliarepine**T-410**

[128562-89-0]


C₃₆H₃₆N₂O₅ 576.691

Alkaloid from the leaves of *Tiliacora racemosa* (Menispermaceae). Amorph. [α]_D²² +75.17 (CHCl₃).

Ray, A.K. et al., *Phytochemistry*, 1990, **29**, 1020; 1023 (isol, uv, ir, pmr, cmr, struct)

Tilivalline**T-411**

1,2,3,10,11,11a-Hexahydro-9-hydroxy-11-(1H-indol-3-yl)-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI [80279-24-9]

C₂₀H₁₉N₃O₂ 333.389

Alkaloid from *Klebsiella pneumoniae* var. *oxytoca*. Mp 168° Mp 180-185° (synthetic). [α]_D²⁵ +126.8 (MeOH). [α]_D²⁵ +210 (c, 1 in MeOH) (synthetic). λ_{max} 220 (ε 28500); 239 (sh); 255 (sh); 278 (sh); 287 (ε 3548); 323 (ε 2950); 332 (ε 2950) (MeOH) (Derep).


Budzikiewicz, H. et al., *Org. Mass Spectrom.*, 1981, **16**, 329 (ms)

Mohr, N. et al., *Tetrahedron*, 1982, **38**, 147 (isol, uv, pmr, cmr, struct, synth)

Nagasaka, T. et al., *J.O.C.*, 1998, **63**, 6797-6801 (synth)

Tinctarine**T-412**

1,2,3,4,5,6-Hexahydro-3-methyl-4-(2-propenyl)-1,5-methano-5H-pyrido[1,2-a][1,5]diazocin-8-one, 9CI. *N-Methyl-11-(2-propenyl)cytisine*. *Alteramine* [33023-11-9]

C₁₅H₂₀N₂O 244.336

Identity of Tinctarine and Alteramine has not been proven but is v. probable.

Alkaloid from *Genista tinctoria*, *Baptisia australis* and *Thermopsis alterniflora* (Fabaceae). Mp 112-113°. Bp_{0.0045} 125-128°. [α]_D²⁵ -43 (EtOH). [α]_D²⁵ -49 (c, 1.385 in EtOH).

Hydrochloride:

Cryst. (MeOH/Me₂CO). Mp 185-186°.

Picrate:

Cryst. (EtOH). Mp 215-216° (211-212° dec.).

N-De-Me: 11-Allylcytisine

[73615-75-5]

[73710-87-9]

C₁₄H₁₈N₂O 230.309

Alkaloid from seeds of *Calotropis brachypetala* and fruits of *Sophora secundiflora* (Fabaceae). Brown oil; needles (as perchlorate). Mp 183-184° (perchlorate). [α]_D²⁶ -215 (c, 0.006 in MeOH) (perchlorate). Abs. config. not detd.

Stereoisomer: Isotinctarine

[77254-90-1]

C₁₅H₂₀N₂O 244.336

Alkaloid from *Baptisia australis* (Fabaceae). Undetd. config. No phys. props. reported.

[33530-05-1]

Knöfel, D. et al., *J. Prakt. Chem.*, 1970, **312**, 887 (isol, ir, uv, ms, struct)

Shaimardanov, R.A. et al., *Khim. Prir. Soedin.*, 1970, **6**, 276; 1971, **7**, 160; *Chem. Nat. Compd. (Engl. Transl.)*, 160; 277 (isol, struct)


Keller, W.J. et al., *Phytochemistry*, 1979, **18**, 2068 (N-Allylcytisine)

Keller, W.J. et al., *J. Nat. Prod.*, 1980, **43**, 164 (N-Allylcytisine)

Wink, M. et al., *J. Nat. Prod.*, 1981, **44**, 14 (isol, ms, Tinctarine, Isotinctarine)

Tinctormine**T-413**

[149475-43-4]

C₂₇H₃₁NO₁₄ 593.54

Pigment of the flowers of *Carthamus tinctorius* (safflower). Potent calcium antagonist. Yellow amorph. powder. [α]_D²⁵ -260 (c, 0.1 in MeOH) (-206). Two different opt. rotn. values given in the ref. λ_{max} 275 (log ε 4.5); 405 (log ε 4.5) (MeOH).

Meselhy, M.R. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1796-1802 (isol, pmr, cmr, ms)

Tinosporine†**T-414**

[80210-02-2]


Struct. unknown. Alkaloid from *Tinospora cordifolia* (Menispermaceae).

Picrate: Mp 174-178°.

Quadrat-i-Khuda, M. et al., *Sci. Res. (Dacca)*, 1964, **1**, 177-183; *CA*, **61**, 12331c

Tintamine

[211311-10-3]


C₂₀H₂₁N₃O₂S 367.471

Numbered by analogy with pyridoacridine alkaloids to which it appears to be related. Tautomeric with the 4-oxo-3NH-form. Alkaloid from the tunicate *Cystodytes violatinctus*. λ_{\max} 265 (log ϵ 3.68); 308 (log ϵ 3.67) (MeOH) (as di-Ac).

Koren-Goldshlager, G. *et al.*, *J.O.C.*, 1998, **63**, 4601-4603 (*isol, uv, ir, pmr, cmr*)

Tirandalydigin

[114118-91-1]

C₂₂H₂₇NO₆ 401.458

Prod. by a *Streptomyces tirandis-umidus*. Sol. MeOH, bases, CHCl₃, C₆H₆; poorly sol. H₂O, hexane. $[\alpha]_D^{26}$ -4 (c, 0.5 in MeOH). λ_{\max} 236 (ϵ 8100); 353 (ϵ 32700); 366 (sh) (ϵ 30000) (EtOH/HCl) (Derep). λ_{\max} 253 (ϵ 12500); 287 (ϵ 16200); 331 (ϵ 16700) (EtOH/KOH) (Derep). λ_{\max} 254; 289; 332 (MeOH).


Brill, G.M. *et al.*, *J. Antibiot.*, 1988, **41**, 36 (*pmr, cmr, ir, struct*)

Karwowski, J.P. *et al.*, *J. Antibiot.*, 1992, **45**, 1125 (*isol, props*)

Iwata, Y. *et al.*, *Angew. Chem., Int. Ed.*, 2005, **44**, 1532-1536 (*synth*)

Tirandamycin A

[34429-70-4]

C₂₂H₂₇NO₇ 417.458

Tetramic acid antibiotic. Isol. from *Streptomyces tirandis* NRRL3689 and

T-415

Streptomyces flaveolus. Active against gram-positive bacteria. Potent inhibitor of RNA-polymerase. Sol. MeOH, Et₂O; poorly sol. H₂O, hexane. Mp 98-102°. Related to Streptolydigin. λ_{\max} 236 (ϵ 8100); 353 (ϵ 32700); 366 (sh) (ϵ 30000) (EtOH/HCl) (Derep). λ_{\max} 253 (ϵ 12500); 287 (ϵ 16200); 331 (ϵ 16700) (EtOH/KOH) (Derep). λ_{\max} 253 (E1%/1cm 295); 286 (E1%/1cm 366); 332 (E1%/1cm 400) (NaOH) (Berdy). λ_{\max} 287 (ϵ 16200); 331 (ϵ 16700); 353 (ϵ 32700) (EtOH/NaOH) (Berdy).

► LD₅₀ (mus, scu) 370 mg/kg. UY7625500

Na salt:

Cryst. $[\alpha]_D^{25}$ +51 (EtOH).

11-Hydroxy: **Tirandamycin B**

[60587-14-6]

C₂₂H₂₇NO₈ 433.457

Isol. from *Streptomyces flaveolus*. Active against gram-positive bacteria. RNA polymerase inhibitor. Yellow powder. Sol. MeOH, CH₂Cl₂, EtOH; poorly sol. H₂O, hexane, C₆H₆. Mp 92-96°. λ_{\max} 236 (ϵ 8100); 353 (ϵ 32700); 366 (sh) (ϵ 30000) (EtOH/HCl) (Derep). λ_{\max} 253 (ϵ 12500); 287 (ϵ 16200); 331 (ϵ 16700) (EtOH/KOH) (Derep). λ_{\max} 250 (E1%/1cm 460); 288 (E1%/1cm 520); 348 (E1%/1cm 650) (EtOH) (Berdy).

MacKellar, F.A. *et al.*, *J.A.C.S.*, 1971, **93**, 4943 (*uv, ir, pmr, ms*)

Meyer, C.E. *et al.*, *J. Antibiot.*, 1971, **24**, 558 (*isol*)

Duchamp, D.J. *et al.*, *J.A.C.S.*, 1973, **95**, 4077 (*cryst struct*)

Hagenmaier, H. *et al.*, *Arch. Microbiol.*, 1976, **109**, 65 (*deriv*)

Reusser, F. *et al.*, *Antibiotics (N.Y.)*, 1979, **5**, 361 (*rev*)

Lee, V.J. *et al.*, *J. Antibiot.*, 1980, **33**, 408 (*cmr*)

Nolte, M.J. *et al.*, *J.C.S. Perkin 1*, 1980, 1057 (*struct*)

Schlessinger, R.H. *et al.*, *J.A.C.S.*, 1985, **107**, 1777 (*synth*)


DeShong, P. *et al.*, *J.A.C.S.*, 1985, **107**, 5219 (*synth*)

Martin, S.F. *et al.*, *Tetrahedron*, 1988, **44**, 3171 (*synth, bibl*)

Shimshock, S.J. *et al.*, *J.A.C.S.*, 1991, **113**, 8791 (*synth, bibl*)

Tiruchanduramine

T-418

C₁₇H₁₈N₆O 322.369


(ξ)-form

Alkaloid from the ascidian *Synoicum macroglossum*. α -Glucosidase inhibitor. Semisolid. $[\alpha]_D$ +31 (c, 0.5 in MeOH). λ_{\max} 215; 234; 270; 334; 347 (MeOH).

Ravinder, K. *et al.*, *Tet. Lett.*, 2005, **46**, 5475-5478 (*isol, synth, pmr, cmr*)

Tjipanazole J

[139112-21-3]


C₂₀H₁₁Cl₂N₃O₂ 396.231

Alkaloid from the blue-green alga *Tolypothrix tjipanasensis*. λ_{\max} 238 (ϵ 11000); 259 (ϵ 8570); 303 (ϵ 20200); 339 (ϵ 4610); 369 (ϵ 2410) (MeOH/CHCl₃) (Derep).

Bonjouklian, R. *et al.*, *Tetrahedron*, 1991, **47**, 7739 (*isol, uv, pmr, cmr, struct*)

Toddacoumalone

[139750-79-1]


C₃₁H₃₁NO₆ 513.589

Alkaloid from *Toddalia asiatica* (*Toddalia aculeata*) (Rutaceae). Prisms (EtOAc). Mp 202-204°. λ_{\max} 225 (ϵ 6760); 262 (ϵ 2040); 285 (ϵ 2950); 316 (ϵ 2090) (MeOH) (Derep).

Ishii, H. *et al.*, *Tet. Lett.*, 1991, **32**, 6907-6910 (*isol, pmr, cmr, struct*)

Toddaliamide

[189295-01-0]

C₂₂H₃₃NO₆ 407.506

Isol. from root wood of *Toddalia asiatica*. Prisms (MeOH/EtOAc). Mp 227-228°. $[\alpha]_D^{23}$ +19 (c, 0.1 in MeOH). λ_{\max} 205 (log ϵ 4.48); 225 (log ϵ 4.27); 245 (sh) (log ϵ 4.22); 300 (log ϵ 4.21); 320 (log ϵ 4.19) (EtOH).

3''-Me ether: **Methyltoddaliamide**

[189295-02-1]

C₂₃H₃₅NO₆ 421.533

From rootwood of *Toddalia asiatica*. Prisms (MeOH/EtOAc). Mp 210-211°.


T-419

T-420

T-421

λ_{\max} 220 (log ϵ 4.22); 240 (log ϵ 4.21); 260 (log ϵ 3.73); 300 (log ϵ 4.25); 320 (log ϵ 4.23) (EtOH).

Tsai, I.-L. *et al.*, *Phytochemistry*, 1997, **44**, 1383 (*isol, uv, ir, pmr, cmr, ms, struct*)

Tokaradine A**T-422**

$C_{28}H_{31}Br_4N_4O_4^{\oplus}$ 807.193

Quaternary alkaloid from *Pseudoceratina purpurea*. Amorph. yellow solid. Counterion not specified. λ_{\max} 216 (ϵ 21800); 258 (sh) (ϵ 4000) (MeOH).

Regioisomer: Tokaradine B


$C_{28}H_{31}Br_4N_4O_4^{\oplus}$ 807.193

Quaternary alkaloid from *Pseudoceratina purpurea*. Amorph. yellow solid. Has the groups attached to C-4 and C-16 interchanged. Counterion not specified. λ_{\max} 214 (ϵ 20600); 258 (sh) (ϵ 3800) (MeOH).

Fusetani, N. *et al.*, *Tetrahedron*, 2001, **57**, 7507-7511

Tomatillidine**T-423**

21-(3,4-Dihydro-3-methyl-2H-pyrrol-5-yl)-3-hydroxy-20-methylpregn-5-en-21-one, 9CI
[986-45-8]



Absolute Configuration

$C_{27}H_{41}NO_2$ 411.626

Alkaloid from *Solanum tomatillo* leaves (Solanaceae). Cryst. (Me₂CO). Mp 219-222°. [α]_D -18.1 (CHCl₃).

5 α ,6-Dihydro: Dihyrotomatillidine

[986-44-7]

$C_{27}H_{43}NO_2$ 413.642

Alkaloid from *Solanum tomatillo* (Solanaceae). Mp 179-181°. [α]_D²⁰ +21.4 (CHCl₃).


Bianchi, E. *et al.*, *J.O.C.*, 1965, **30**, 754-760

(*isol, pmr, ord*)

Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1976, **24**, 661-666 (*struct, cd, pmr, ms, synth*)

Tomaymycin**T-424**

2-Ethylidene-1,2,3,10,11,11a-hexahydro-8-hydroxy-7,11-dimethoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, 9CI
[35050-55-6]



(E)-form

$C_{16}H_{20}N_2O_4$ 304.345

Anthramycin-type antibiotic. Isol. from *Streptomyces achromogenes* and *Nocardia C-15003*. Antitumour agent. Platelets + MeOH. Mp 145-146°.

[α]_D²⁰ +423 (c, 0.5 in Py). Strain also produces minor analogs (hydroxy, ethoxy, propoxy). λ_{\max} 234 (ϵ 19600); 252 (sh) (ϵ 16700); 289 (ϵ 11400); 325 (ϵ 9270) (MeOH/KOH) (Derep). λ_{\max} 224 (ϵ 26500); 237 (ϵ 24500); 260 (sh); 262 (ϵ 7840); 322 (ϵ 4500) (MeOH) (Derep).

▶ LD₅₀ (mus, ipr) 1 mg/kg. UY8533000

11-O-De-Me: 11-De-O-methyltomaymycin

[55511-85-8]

$C_{15}H_{18}N_2O_4$ 290.318

From *Streptomyces achromogenes*.

Shows antitumour activity.

11-O-De-Me, 11-ketone: Oxotomaymycin

[35050-54-5]

$C_{15}H_{16}N_2O_4$ 288.302

From *Streptomyces achromogenes*.

Biol. inactive. Mp 230° dec. Not obtained completely free of tomaymycin.

11-Demethoxy: Antibiotic SEN 215. 11-Demethoxytomaymycin. SEN 215

[67731-61-7]

$C_{15}H_{18}N_2O_3$ 274.319

Prod. by *Streptomyces cylindrosporus*

and *Streptomyces sakaiensis*. Shows

antitumour, analgesic, antispasmodic

and sedative props. Needles. Mp 216-

219°. [α]_D²³ +177 (c, 0.5 in CHCl₃).

Lacks the alicyclic methoxy group.

λ_{\max} 231 ; 238 ; 269 ; 340 (MeOH)

(Berdy).

▶ LD₅₀ (mus, ipr) 60 mg/kg. UY8534580

11-Demethoxy, 11-ethoxy: Tomaymycin I

$C_{17}H_{22}N_2O_4$ 318.372

Prod. by *Nocardia* sp. ATCC31281.

Active against bacteria, phages, and

tumours. Pale-yellow needles. Mp 134-

136°.

11-Demethoxy, 10,11-didehydro: Preto-maymycin. Pretomeimycin

[28797-41-3]

$C_{15}H_{16}N_2O_3$ 272.303

Anthramycin antibiotic. Prod. by

Streptomyces achromogenes tomaymy-

ceticus ATCC21353. Biosynthetic in-

termediate for Tomaymycin. Converts

on standing to Tomaymycin, useful for

prep. of analogs. Pale-yellow powder

(EtOAc). Sol. MeOH, DMSO, CHCl₃;

fairly sol. Et₂O; poorly sol. hexane,

H₂O. [α]_D²⁵ +215 (c, 0.08 in Py). λ_{\max}

220 (ϵ 26000); 316 (ϵ 3500) (MeOH)

(Berdy).

Kariyone, K. *et al.*, *Chem. Pharm. Bull.*, 1971, **19**, 2289 (*isol, struct*)

Arima, K. *et al.*, *J. Antibiot.*, 1972, **25**, 437

(*isol, props*)

Japan. Pat., 1973, 79 73195; 1974, 74 25 956; 74

25 957; 1978, 78 56 693; 1981, 81 121 495;

C.A. **83**, 58899; 77004; **89**, 161536; **91**,

209360 (*isol, props, derivs*)

Hurley, L. *et al.*, *Antimicrob. Agents*

Chemother., 1979, **15**, 42 (*biosynth*)

Hurley, L.H. *et al.*, *Acc. Chem. Res.*, 1980, **13**,

263 (*rev*)

Tozuka, Z. *et al.*, *J. Antibiot.*, 1983, **36**, 142;

276 (*synth, struct, pmr, cmr*)

Kaneko, T. *et al.*, *Tet. Lett.*, 1983, **24**, 5165 (*synth*)


Barkley, M.D. *et al.*, *Biochemistry*, 1986, **25**, 3021 (*pmr, struct*)

Mori, M. *et al.*, *Tetrahedron*, 1986, **42**, 3793 (*synth*)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, THG500

Tombozine**T-425**

Sarpagan-17-ol, 9CI. Normacusine B. Vellosiminol. Desformoakuammidinol. Desoxysarpagine. Tombozine
[604-99-9]



$C_{19}H_{22}N_2O$ 294.396

Alkaloid from *Geissospermum vellosii*,

several *Rauwolfia* and *Strychnos* spp.

(Apocynaceae, Loganiaceae). Shows se-

dativ and ganglion blocking activity.

Dimorphic cryst. Mp 245° Mp 270-272°.

O-Ac:

Prisms (Et₂O). Mp 219-221° (205-

212°). [α]_D +12 (c, 0.33 in MeOH).

O-Benzoyl: O-Benzoyltombozine

[54357-60-7]

$C_{26}H_{26}N_2O_2$ 398.504

Alkaloid from *Vinca erecta* epigeal

parts (Apocynaceae). Cryst. (Et₂O).

[α]_D²⁰ +93 (c, 0.3 in CHCl₃).

N¹-Me: Affinisine. Dehydroxymethylvo-

chaloitolin. N⁴-Methyldeoxysarpagine

[2912-11-0]

$C_{20}H_{24}N_2O$ 308.422

Alkaloid from *Peschiera affinis*, *Ta-*

bernaemontana fuchsiaeifolia and *Al-*

stonia macrophylla (Apocynaceae).

Shows weak antibacterial activity and

mod. analgesic and CNS depressant

props. Active against gram-positive

bacteria and fungi. Mp 194-196°

(185°). [α]_D²⁰ +19 (c, 0.78 in CHCl₃)

(+30).

▶ AW4000000

N¹-Me, N⁴-oxide: Affinisine N^b-oxide

[138989-35-2]

$C_{20}H_{24}N_2O_2$ 324.422

Alkaloid from *Ervatamia hirta* (Apoc-

ynaceae). [α]_D +3 (c, 1 in MeOH).

N¹-Me, O-Ac: Mp 179-180° (172°). [α]_D³⁰

+8 (c, 0.82 in CHCl₃).

N¹-Me, N⁴-chloromethyl: Chloromethyle-

neaffinisinium

[167696-85-7 (chloride)]

$C_{21}H_{26}ClN_2O^{\oplus}$ 357.902

Alkaloid from stem bark of *Peschiera*

buchtieni (Apocynaceae). [α]_D +4 (c,

0.4 in MeOH) (as chloride). Artifact.

N⁴-Me: Macusine B

[6792-07-0]

$C_{20}H_{25}N_2O^{\oplus}$ 309.43

Quaternary alkaloid from

- Aspidosperma peroba*, *Pleiocarpa pycnantha*, *Pleiocarpa tubicina*, *Strychnos toxifera* and several other *Strychnos* spp. (Apocynaceae, Loganiaceae). Cryst. (MeOH) (as chloride). Mp 249-250° dec. (chloride). $[\alpha]_D^{25} +14$ (c, 0.35 in H₂O) (chloride).
- Me ether: 17-Methoxysarpagan. O-Methylnormacusine B** [68160-78-1]
C₂₀H₂₄N₂O 308.422
Alkaloid from the stem bark of *Rauwolfia cumminsii* (Apocynaceae). Off-white amorph. powder.
- Me ether, N⁴-oxide: 17-Methoxysarpagan N-oxide** [70319-20-9]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from the stem bark of *Rauwolfia cumminsii* (Apocynaceae). Cream amorph. powder.
- Me ether, N⁴-Me: O-Methylmacusine B** [55249-53-1]
C₂₁H₂₇N₃O[⊕] 323.457
Alkaloid from *Strychnos usambarensis* roots and the stem bark of *Strychnos decussata* (Loganiaceae).
- 17-Aldehyde: Vellosimine. Alkaloid RB19** [6874-98-2]
C₁₉H₂₀N₂O 292.38
Alkaloid from *Geissospermum vellosii*, *Rauwolfia vomitoria*, *Rauwolfia macrophylla*, *Rauwolfia nitida* and *Astonia yunnanensis* (Apocynaceae). Cryst. (MeOH). Mp 260° Mp 305-306°. λ_{max} 226 (log ϵ 4.52); 283 (log ϵ 3.82) (EtOH).
- 17-Aldehyde, N¹-Me: N⁴-Methylvellosimine** [81525-53-3]
C₂₀H₂₂N₂O 306.407
Alkaloid from the root bark of *Rauwolfia nitida* (Apocynaceae). Plates. Mp 255-260°. $[\alpha]_D^{22} +23$ (c, 0.01 in CHCl₃).
- 17-Carboxylic acid, N¹-Me, Me ester: N-Methyl-16-epipericyclivine** [2912-15-4]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from the leaves of *Astonia undulata* (Apocynaceae). Cryst. (MeOH). Mp 231-232°. $[\alpha]_D +22.8$ (c, 0.5 in CHCl₃) (synthetic). $[\alpha]_D 0^\circ$ (c, 1 in CHCl₃).
- 17-Carboxylic acid, N⁴-Me (zwitterion): Panarine** [119736-90-2]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from a Venezuelan curare prepd. from the bark of *Strychnos toxifera*. Powder.
- 17-Carboxylic acid, N⁴-Me, Me ester: Alkaloid Q₃** [80151-85-5]
C₂₁H₂₅N₃O₂[⊕] 337.441
Quaternary alkaloid from *Aspidosperma peroba* (Apocynaceae). Tentative struct.
- 19,20 ξ -Dihydro, Me ether, N⁴-Me: Dihydro-O-methylmacusine B** [55249-54-2]
C₂₁H₂₉N₂O[⊕] 325.473
- Alkaloid from the roots of *Strychnos usambarensis* (Loganiaceae).
- 18-Hydroxy, N¹-Me: 18-Hydroxyaffinisine** [167696-86-8]
C₂₀H₂₄N₂O₂ 324.422
Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae).
- 16-Epimer: 16-Epitombozine. 16-Epinormacusine B** [126640-98-0]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Ervatamia hirta* (Apocynaceae). $[\alpha]_D +3$ (c, 0.25 in MeOH).
- 16-Epimer, N¹-Me: 16-Epiaffinisine** [139067-48-4]
C₂₀H₂₄N₂O 308.422
Alkaloid from *Ervatamia hirta* (Apocynaceae). $[\alpha]_D -18$ (c, 0.5 in MeOH).
- 16-Epimer, 17-carboxylic acid, Me ester: Pericyclivine. Methyl sarpagan-17-oate** [975-77-9]
C₂₀H₂₂N₂O₂ 322.406
Alkaloid from *Tabernaemontana holstii*, *Tabernaemontana johnstonii*, *Vinca rosea*, *Catharanthus lanceus*, *Catharanthus ovalis*, *Catharanthus roseus*, *Hazunia modesta*, *Gabunia odoratissima* (preferred genus name *Tabernaemontana*) and *Rauwolfia cumminsii* (Apocynaceae). Shows weak cytotoxic activity vs. P-388 rat leukaemia cells. Needles (Me₂CO). Mp 232-233° (226-228°). $[\alpha]_D^{25} +5.2$ (c, 1.0 in CHCl₃). $[\alpha]_D +2.9$ (CHCl₃) (synthetic). Pmr and cmr revised in 1996. λ_{max} 225 (log ϵ 4.51); 281 (log ϵ 3.88); 289 (log ϵ 3.79) (MeOH).
▶ VQ5835000
- 16-Epimer, 17-carboxylic acid, N¹-Me, Me ester: N-Methylpericyclivine** [160497-66-5]
C₂₁H₂₄N₂O₂ 336.433
Alkaloid from stem bark of *Peschiera buchtieni* (Apocynaceae). $[\alpha]_D +7.5$ (c, 0.29 in CHCl₃).
- 16-Epimer, 17-carboxylic acid, N⁴-Me (zwitterion): 16-Epipanarine** [135355-84-9]
C₂₀H₂₂N₂O₂ 322.406
Constit. of *Stemmadenia minima* (Apocynaceae). Needles (MeOH). Mp 226°. $[\alpha]_D^{20} -29$ (c, 0.7 in MeOH).
- 16-Epimer, 6 ξ -hydroxy: Ervincidine** [32075-02-8]
C₁₉H₂₂N₂O₂ 310.395
Alkaloid from epigeal parts of *Vinca erecta* (Apocynaceae). Cryst. (MeOH). Mp 279-280° dec. $[\alpha]_D^{25} +29.5$ (c, 0.6 in MeOH). Probable struct. MF incorr. given as C₁₉H₂₄N₂O₂.
- 16-Epimer, 19Z-isomer: Koumidine** [1358-75-4]
C₁₉H₂₂N₂O 294.396
Alkaloid from *Gelsemium elegans* and *Gelsemium sempervirens* (Loganiaceae). Cryst. (Me₂CO). Mp 201-203°. $[\alpha]_D -11$ (c, 0.07 in MeOH). Originally assigned the common 19E-config. later shown to be (19Z-).
Battersby, A.R. et al., *J.C.S.*, 1960, 1848-1854; 1964, 4419-4427 (*Macusine B*, isol)
- Gosset, J. et al., *Bull. Soc. Chim. Fr.*, 1961, 1033-1035 (*Affinisine, synth, O-Ac*)
Anatonaccio, L.D. et al., *J.A.C.S.*, 1962, **84**, 2161-2169 (*Tombozine, isol, ms*)
Rapoport, H. et al., *J.O.C.*, 1962, **27**, 2981-2985 (*Tombozine, Vellosimine, isol, synth, struct*)
Cava, M.P. et al., *Chem. Ind. (London)*, 1964, 1193-1194 (*Affinisine, isol, Ac, ir, uv, pmr*)
Farnsworth, N.R. et al., *J. Pharm. Sci.*, 1964, **53**, 1558 (*Pericyclivine, isol, struct*)
Achenbach, H. et al., *Tet. Lett.*, 1966, 4405-4407 (*Affinisine, ms, struct*)
Khan, Z.M. et al., *Helv. Chim. Acta*, 1967, **50**, 625-627 (*Macusine B*)
Gorman, A.A. et al., *Helv. Chim. Acta*, 1969, **52**, 33-55 (*Macusine B, uv*)
Mitscher, L.A. et al., *J. Nat. Prod.*, 1972, **35**, 157-176 (*Affinisine, activity*)
Malikov, V.M. et al., *Khim. Prir. Soedin.*, 1972, **8**, 760-761; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **8**, 741-742 (*Ervincidine*)
Banerji, A. et al., *Phytochemistry*, 1972, **11**, 2605-2607 (*Affinisine, isol, ms*)
Marini-Bettòlo, G.B. et al., *Gazz. Chim. Ital.*, 1973, **103**, 591 (*Tombozine, cmr*)
Patel, M.B. et al., *Phytochemistry*, 1973, **12**, 451-456 (*Tombozine, isol, uv, ir, pmr, ms*)
Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1974, **39**, 3168-3176 (*Pericyclivine, uv, cd*)
Sharipov, M.R. et al., *Khim. Prir. Soedin.*, 1974, **10**, 413-414; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **10**, 422-423 (*O-Benzoyltombozine*)
Angenot, L. et al., *Planta Med.*, 1975, **27**, 24-30 (*O-Methylmacusine B, Dihydro-O-methylmacusine*)
Kingston, D.G.I. et al., *J. Pharm. Sci.*, 1977, **66**, 1135-1138; 1978, **67**, 249-251 (*Pericyclivine, isol, activity*)
Iwu, M.M. et al., *Phytochemistry*, 1978, **17**, 1651-1654 (*17-Methoxysarpagan, 17-Methoxysarpagan N-oxide*)
Chatterjee, A. et al., *Indian J. Chem., Sect. B*, 1979, **18**, 87-88 (*Vellosimine, isol, uv, ir, pmr, ms, struct*)
Marini-Bettòlo, G.B. et al., *Phytochemistry*, 1980, **19**, 992-994 (*O-Ac, cmr*)
Amer, M.A. et al., *Phytochemistry*, 1981, **20**, 2569-2573 (*N⁴-Methylvellosimine*)
Schun, Y. et al., *Phytochemistry*, 1987, **26**, 2875-2876 (*Koumidine, struct*)
Dupont, L. et al., *Acta Cryst. C*, 1988, **44**, 2156-2158 (*Panarine, cryst struct*)
Quetin-Leclercq, J. et al., *Phytochemistry*, 1988, **27**, 4002-4004 (*Panarine*)
Ponglux, D. et al., *Tetrahedron*, 1988, **44**, 5075-5094 (*Koumidine, struct*)
Magnus, P. et al., *J.A.C.S.*, 1989, **111**, 786-789; 1990, **112**, 5220-5230 (*Koumidine, synth*)
Takayama, H. et al., *J.C.S. Perkin 1*, 1989, 1075-1076 (*Koumidine, synth, abs config*)
Pinchon, T.M. et al., *Phytochemistry*, 1990, **29**, 3341-3344 (*N-Methyl-16-epipericyclivine*)
Achenbach, H. et al., *J. Nat. Prod.*, 1991, **54**, 473-476 (*16-Epipanarine*)
Kitajima, M. et al., *J.C.S. Perkin 1*, 1991, 1773-1779 (*synth*)
Clivio, P. et al., *Phytochemistry*, 1991, **30**, 3785-3792 (*16-Epitombozine, Affinisine N-oxide*)
Bailey, P.D. et al., *J.C.S. Perkin 1*, 1993, 441-449 (*synth*)
Azoug, M. et al., *Phytochemistry*, 1995, **39**, 1223-1228 (*18-Hydroxyaffinisine, Chlormethleaffinisinium, N-Methylpericyclivine*)
Jokela, R. et al., *Heterocycles*, 1996, **43**, 1015-1020 (*Pericyclivine, pmr, cmr*)
Liu, X. et al., *Tet. Lett.*, 2000, **41**, 6299-6303 (*Affinisine, synth*)
Deiters, A. et al., *J.A.C.S.*, 2003, **125**, 4541-4550 (*N-Methylvellosimine, synth*)


Yu, J. *et al.*, *J.O.C.*, 2003, **68**, 5852-5859; 7565-7581 (*Tombozine, Vellosimine, Panarine, 16-Epiaffinisine, Alkaloid Q3, N-Methyl-16-epipericyclivine, synth, pmr, cmr*)
 Sarma, P.V.S. *et al.*, *Org. Lett.*, 2006, **8**, 1017-1020 (*N-Methylpericyclivine, synth*)

Tomentocurine T-426
 [1361-62-2]

C₃₆H₃₈N₂O₆ 594.706
 Bisbenzylisoquinoline alkaloid. Struct. unknown. Contains 2 OMe groups and at least one OH. Prob. a close relative of Chondocurine, C-420. Alkaloid from the stems and leaves of *Chondodendron tomentosum* (Menispermaceae). Cryst. + 2³/₄ H₂O. Mp 265° (efferv.). [α]_D¹⁷ +210 (0.1M HCl).

King, H. *et al.*, *J.C.S.*, 1948, 1945-1949 (*isol*)
 Bick, I.R.C. *et al.*, *J.C.S.*, 1960, 2402-2407 (*isol*)


Tongolinine T-427
 19,21-Didehydro-6,21-secohetisan-5,15-diol
 [179234-13-0]



C₂₀H₂₇NO₂ 313.439
 Alkaloid from *Delphinium tongolense*. Needles. Mp 233-234°.

He, L. *et al.*, *Chin. Chem. Lett.*, 1996, **7**, 557-560 (*isol, pmr, cmr*)

Tonkinensine A T-428
 [1044253-68-0]




Absolute Configuration

C₂₈H₂₆N₂O₆ 486.523
 Alkaloid from the roots of *Sophora tonkinensis*. Gum (CHCl₃). [α]_D²⁰ -334 (c, 0.11 in CHCl₃). λ_{max} 205 (log ε 3.89); 231 (log ε 3.45); 308 (log ε 3) (MeOH).

Li, X.-N. *et al.*, *Tet. Lett.*, 2008, **49**, 3797-3801 (*isol, cd, pmr, cmr, ms*)

Tonkinensine B T-429
 [1044253-69-1]




Absolute Configuration

C₂₈H₂₆N₂O₆ 486.523
 Alkaloid from the roots of *Sophora tonkinensis*. Cryst. (MeOH). Mp 258-260°. [α]_D²⁰ -327 (c, 0.11 in CHCl₃). λ_{max} 204 (log ε 3.85); 233 (log ε 3.51); 309 (log ε 3.18) (MeOH).

Li, X.-N. *et al.*, *Tet. Lett.*, 2008, **49**, 3797-3801 (*isol, cd, pmr, cmr, ms, cryst struct*)

Topostin B553 T-430
 [129204-42-8]



R¹ = H, R² = OH

C₃₃H₆₃NO₅ 553.864
 Struct. established 1998. Earlier tentative structs. shown to be incorrect. Prod. by *Flexibacter topostinus*. Inhibitor of mammalian DNA topoisomerase I. Cytotoxic. Powder. [α]_D²² +1.5 (c, 0.5 in CHCl₃).

[126603-04-1, 126603-02-9, 126603-03-0]

Suzuki, K. *et al.*, *J. Antibiot.*, 1990, **43**, 154; 158 (*isol, activity*)

Noguchi, H. *et al.*, *Tetrahedron*, 1995, **51**, 10531; 10545 (*synth, analogues*)

Nemoto, T. *et al.*, *Tetrahedron*, 1998, **54**, 2683-2690 (*isol, ir, pmr, cmr*)


Topostin D 640 T-431
 [205250-26-6]

As Topostin B553, T-430 with R¹ = H, R² = -NHCH(CH₂OH)-COOH(S-)

C₃₆H₆₈N₂O₇ 640.942
 Prod. by *Flexibacter topostinus*. Inhibitor of mammalian DNA topoisomerase I. Powder. [α]_D²² +14.2 (c, 2 in CHCl₃).

Nemoto, T. *et al.*, *Tetrahedron*, 1998, **54**, 2683-2690 (*isol, ir, pmr, cmr*)

Toposentin A T-432
 1*H*-Indol-3-yl[4-(1*H*-indol-3-yl)-1*H*-imidazol-2-yl]methanone, 9*CI*. 4-(3-Indolyl)-2-(3-indolylcarbonyl)imidazole. Deoxytoposentin
 [112515-42-1]



C₂₀H₁₄N₄O 326.357
 Shows tautomerism of the imidazole ring in soln., tautomer shown predominates. Naming and numbering systems vary. Alkaloid from the Mediterranean sponge *Toposentia genitrix*. Shows weak piscicidal activity. Mp 290-292°.

13,14-Dihydro: Toposentin D
 [122889-35-4]

C₂₀H₁₆N₄O 328.373
 Alkaloid from *Toposentia* sp.

6-Hydroxy: **Toposentin B1**. *Toposentin* [112515-43-2]
 C₂₀H₁₄N₄O₂ 342.356
 Alkaloid from *Toposentia genitrix* and from the Caribbean deep-sea sponge *Spongosorites* sp. Weak piscicide. Amorph. bright yellow solid. Mp 270°. λ_{max} 208 (ε 1800); 246 (ε 5100); 300 (ε 3400); 375 (ε 3100) (EtOH/KOH) (Derep). λ_{max} 202 (ε 41000); 220 (sh) (ε 31500); 240 (sh) (ε 19200); 280 (ε 13500); 378 (ε 17300) (95% EtOH) (Derep).

▶ PC4957000

6-Bromo: 6-Bromotoposentin A. **Isobromodeoxytoposentin**
 [223596-72-3]
 C₂₀H₁₃BrN₄O 405.253
 Alkaloid from *Spongosorites genitrix*. Moderate cytotoxic agent. Amorph. yellow solid. Mp 225-230°. λ_{max} 213 (log ε 4.41); 252 (log ε 4.04); 280 (log ε 4.06); 375 (log ε 4.02) (MeOH).

6-Bromo, 13,14-dihydro: **Dihydrodeoxybromotoposentin**. *Spongotine B*
 [116747-40-1]
 C₂₀H₁₅BrN₄O 407.269
 Isol. from *Spongosorites* sp., *Toposentia* sp. and *Rhaphisia lacazei*. Antiviral and antitumour agent. Amorph. yellow powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. [α]_D²⁴ +198 (c, 2.0 in MeOH). λ_{max} 196 (ε 102000); 198 (ε 29300); 214 (ε 34000); 215 (ε 76000); 274 (ε 8800); 294 (ε 109000); 328 (ε 5700); 330 (ε 7600) (MeOH) (Berdy).

6-Bromo, 22-hydroxy: **Isobromotoposentin**
 [173220-97-8]
 C₂₀H₁₃BrN₄O₂ 421.252
 Alkaloid from *Spongosorites* sp. Amorph. yellow solid. Mp 225-228° dec. λ_{max} 207 (ε 11150); 284 (ε 2900); 415 (ε 4260) (EtOH).

21-Bromo: 21-Bromotoposentin A. **Bromodeoxytoposentin**
 [180633-55-0]
 C₂₀H₁₃BrN₄O 405.253
 Alkaloid from *Spongosorites genitrix* and *Rhaphisia lacazei*. Moderate cytotoxic agent. Amorph. yellow solid. Mp 240-243°. λ_{max} 209 (log ε 4.61); 235 (log ε 4.4); 250 (sh) (log ε 4.36); 367 (log ε 4.17) (MeOH).

21-Bromo, 13,14-dihydro: **Spongotine A**
 C₂₀H₁₅BrN₄O 407.269
 Alkaloid from a *Spongosorites* sp. Amorph. yellow powder. [α]_D²⁵ -14 (c, 0.2 in MeOH). λ_{max} 224 (log ε 4.27); 268 (log ε 3.92); 320 (log ε 3.75) (MeOH).

21-Bromo, 6-hydroxy: **Toposentin B2**. *Bromotoposentin*
 [112515-44-3]
 C₂₀H₁₃BrN₄O₂ 421.252
 Alkaloid from *Toposentia genitrix*, *Spongosorites* sp. and another sponge *Hexadella* sp. Shows antitumour and antiviral activity. Weak piscicide. Bright yellow cryst. (CHCl₃/MeOH) or yellow-green oil. Sol. MeOH, CH₂Cl₂. Mp 260° Mp 296-297°. λ_{max} 209 (ε 19000); 234 (sh) (ε 9700); 300 (ε 4200); 375 (ε 3500) (EtOH/KOH) (Derep). λ_{max} 208 (ε 40000); 237

(ϵ 28800); 254 (sh) (ϵ 22300); 286 (ϵ 15300); 378 (ϵ 17200) (95% EtOH) (Derep). λ_{\max} 208 (ϵ 40000); 237 (ϵ 28800); 286 (ϵ 15300); 328 (ϵ 17200); 382 (ϵ 15500) (EtOH) (Berdy).

6,21-Dibromo-6,21-Dibromotopsentin A. Dibromodeoxytopsentin

$C_{20}H_{12}Br_2N_4O$ 484.149

Alkaloid from the sponge *Spongosorites* sp. Amorph. yellow powder.

6,21-Dibromo-13,14-dihydro- Spongotine C

$C_{20}H_{14}Br_2N_4O$ 486.165

Alkaloid from a *Spongosorites* sp. Cryst. $[\alpha]_D^{25}$ -9 (c, 0.8 in MeOH). λ_{\max} 221 (log ϵ 4.21); 276 (log ϵ 3.85); 315 (log ϵ 3.57) (MeOH).

6,21-Dibromo-13,14-dihydro, N¹²-Me: Topsisent C

[128364-30-7]
 $C_{21}H_{16}Br_2N_4O$ 500.192

Alkaloid from the Pacific Ocean sponge *Hexadella* sp. Pale yellow powder.

Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*, 1987, **96**, 809-812 (*Topsisent A*, synth)

Bartik, K. *et al.*, *Can. J. Chem.*, 1987, **65**, 2118-2121 (*Topsisentins A, B, B2*)

Tsujii, S. *et al.*, *J.O.C.*, 1988, **53**, 5446-5453 (*Topsisent*, *Bromotopsent*, *Dihydrodeoxybromotopsent*)

Morris, S.A. *et al.*, *Can. J. Chem.*, 1989, **67**, 677-681 (*Topsisent B2*)

Morris, S.A. *et al.*, *Tetrahedron*, 1990, **46**, 715-720 (*Topsisent C*)

Murray, L.M. *et al.*, *Aust. J. Chem.*, 1995, **48**, 2053-2058 (*Isobromotopsent*)

Achab, S. *et al.*, *Tet. Lett.*, 1996, **37**, 5503-5506 (*Topsisent*, *Deoxytopsent*, *Bromotopsent*, synth)

Kawasaki, I. *et al.*, *Heterocycles*, 1998, **48**, 1887-1901 (synth)

Shin, J. *et al.*, *J. Nat. Prod.*, 1999, **62**, 647-649 (*Bromodeoxytopsent*, *Isobromodeoxytopsent*)

Casapullo, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 447-451 (isol)

Miyake, F.Y. *et al.*, *Org. Lett.*, 2000, **2**, 2121-2123 (synth)

Bao, B. *et al.*, *J. Nat. Prod.*, 2005, **68**, 711-715; 2007, **70**, 2-8 (*Dibromodeoxytopsent*, *Spongotines A-C*)

Guinchard, X. *et al.*, *J.O.C.*, 2007, **72**, 3972-3975 (synth)


Murai, K. *et al.*, *J.O.C.*, 2007, **72**, 8947-8949 (*Spongotine A*, synth)

Mal, S.K. *et al.*, *Tetrahedron*, 2008, **64**, 5904-5914 (synth)

Tortuosamine

T-433

6-(3,4-Dimethoxyphenyl)-5,6,7,8-tetrahydro-N-methyl-6-quinolineethanamine, 9CI. 7-(3,4-Dimethoxyphenyl)-7-(2-methylaminoethyl)-5,6,7,8-tetrahydroquinoline



(R)-form

$C_{20}H_{26}N_2O_2$ 326.438

Modified type of mesembrenoid alkaloid.

(R)-form [35722-04-4]

Alkaloid from *Sceletium tortuosum* and *Sceletium namaquense* (Aizoaceae). Oil. $[\alpha]_D^{20}$ -29 (c, 1.04 in MeOH).

N-Formyl: N-Formyltortuosamine

[51934-14-6]

$C_{21}H_{26}N_2O_3$ 354.448

Alkaloid from *Sceletium namaquense* (Aizoaceae). Oil.

N-Ac: N-Acetyltortuosamine

[82545-10-6]

$C_{22}H_{28}N_2O_3$ 368.475

Alkaloid from *Sceletium namaquense* (Aizoaceae). Oil.

(±)-form [79517-20-7]

Synthetic. Oil.

N-Formyl: [79517-21-8]

Synthetic. Oil.

Snyckers, F.O. *et al.*, *Chem. Comm.*, 1971, 1467 (isol, uv, ir, pmr, ms, struct)

Jeffs, P.W. *et al.*, *J.O.C.*, 1974, **39**, 2703; 1982, **47**, 3611 (isol, synth, deriv)

Martin, N.H. *et al.*, *Org. Mass Spectrom.*, 1976, **11**, 1 (ms, deriv)

Capps, T.M. *et al.*, *J.C.S. Perkin 2*, 1977, 1098 (isol, uv, ir, pmr, ms)

Koyama, J. *et al.*, *Heterocycles*, 1981, **16**, 969; 1984, **22**, 1973 (synth)

Okatani, T. *et al.*, *Heterocycles*, 1989, **29**, 1809 (synth)

Goehring, R.R. *et al.*, *Tet. Lett.*, 1994, **35**, 8145 (synth)

Kamikubo, T. *et al.*, *Chem. Comm.*, 1998, 783-784 (synth)


Yamada, O. *et al.*, *Tet. Lett.*, 1998, **39**, 7747-7750 (synth)

Hayashi, M. *et al.*, *Tet. Lett.*, 2002, **43**, 1461-1464 (synth)

Torulosine

T-434

[156953-74-1]



$R^1R^2 = CH_2$

$C_{20}H_{17}NO_7$ 383.357

(±)-form

Alkaloid from whole plants of *Dactylis-capnos torulosa* and *Hypecoum leptocarpum*. Orange powder. Mp 190-192°. λ_{\max} 287 (log ϵ 4.8); 405 (log ϵ 4.29) (MeOH/NaOH). λ_{\max} 223 (log ϵ 5); 247 (log ϵ 5.09); 302 (log ϵ 4.88) (MeOH/HCl).

O-De-Me, N-Me: O-Demethyl-N-methyltorulosine. N-Methylde-O-methyltorulosine

[401892-71-5]

$C_{20}H_{18}NO_7$ 384.365

Alkaloid from *Hypecoum leptocarpum*. Amorph. yellow powder (MeOH) (as chloride). Mp 100-105° (chloride).

λ_{\max} 250 (log ϵ 4.02); 310 (log ϵ 3.83); 371 (log ϵ 3.76) (MeOH).

Rücker, G. *et al.*, *Phytochemistry*, 1994, **36**, 519-523 (isol, uv, ir, pmr, cmr, ms)

Li, B.-G. *et al.*, *Indian J. Chem., Sect. B*, 2001, **40**, 1215-1218 (*N-Methylde-O-methyltorulosine*)

Torulosinine

T-435

[156953-75-2]

As Torulosine, T-434 with

$R^1 = R^2 = Me$

$C_{21}H_{21}NO_7$ 399.399

(±)-form

Alkaloid from whole plants of *Dactylis-capnos torulosa* (Papaveraceae). Yellow amorph. powder. Mp 207-208°.


Rücker, G. *et al.*, *Phytochemistry*, 1994, **36**, 519 (isol, uv, ir, pmr, ms, struct)

Toxiferine I

T-436

C-Toxiferine I. Toxiferine V. Toxiferine XI [6888-23-9]

[6696-58-8]



Absolute configuration

$C_{40}H_{46}N_4O_2^{\oplus}$ 614.829

Dimer showing 2-fold rotational symmetry. Alkaloid from calabash curare, *Strychnos toxifera* and *Strychnos froesii* (Loganiaceae). Neuromuscular blocking agent. The most potent of the physiologically active calabash-curare alkaloids. $[\alpha]_D^{25}$ -511 (MeOH) (as dichloride). Red-viol. col. with $Ce(SO_4)_2$, becoming colourless on standing. Equilibrates in mild acid conds. with Caracurine V dimethosalt.

▶ LD₅₀ (mky, ivn) 8900 mg/kg. LD₅₀ (mky, ims) 0.018 mg/kg. Very toxic by intravenous and intramuscular routes.

XW5520000

Monodeoxy: C-Alkaloid H

$C_{40}H_{46}N_4O_2^{\oplus}$ 598.83

Alkaloid from calabash curare (Loganiaceae). Mp 189-192° (as picrate). Red-viol. col. with $Ce(SO_4)_2$, becoming colourless on standing. No CAS no. found to 2007.

Monodeoxy, di-N-de-Me: Caracurine VI.

Bisnor-C-alkaloid H. 18-Hydroxynor-dihydrotoxiferine. Dinor-C-alkaloid H

[67739-70-2]

$C_{38}H_{40}N_4O$ 568.76

Alkaloid from *Strychnos longicaudata*, *Strychnos dolichothyrsa*, *Strychnos afzelii*, *Strychnos malacoclados*, *Strychnos matopensis* and *Strychnos urceolata* (Loganiaceae). Shows antimicrobial props. Purple col. with $Ce(SO_4)_2$, becoming brown on standing. Picrate darkening >260°.

does not melt <300°. Has been confused with Longicaudatine, L-249. λ_{\max} 286 ; 292 (MeOH) (Berdy).

18-Deoxy, di-N-de-Me, N⁴-oxide: **Bisnor-C-alkaloid H N-oxide**

C₃₈H₄₀N₄O₂ 584.76

Alkaloid from *Strychnos dolichothyrsa*.

Monodeoxy, di-N-de-Me, di-N-oxide:

Bisnor-C-alkaloid H di-N-oxide

[69320-13-4]

C₃₈H₄₀N₄O₃ 600.759

Alkaloid from *Strychnos dolichothyrsa*.

Bis(deoxy): **18,18'-Dideoxytoxiferine I. C-Deoxytoxiferine. C-Alkaloid K. C-Dihydrotoxiferine I**

[664-27-7]

C₄₀H₄₆N₄²⁺ 582.83

Alkaloid from calabash curare, *Strychnos toxifera* and a few other *Strychnos* spp. (Loganiaceae). Curarising agent. $[\alpha]_D$ -611 (H₂O) (as dichloride). Blue-viol. col. with Ce(SO₄)₂, becoming colourless on standing.

Bis(deoxy), di-N-de-Me: **Bisnordihydrotoxiferine. 4,4'-Didemethyl-18,18'-dideoxytoxiferine I. Nordihydrotoxiferine. Nordihydrotoxiferine I**

[24163-58-4]

C₃₈H₄₀N₄ 552.761

Alkaloid from *Strychnos dolichothyrsa*, *Strychnos afzelii*, *Strychnos toxifera*, *Strychnos variabilis*, *Strychnos longicaudata* and others. Major alkaloid of *Strychnos pseudoquina* (Loganiaceae). Shows antimicrobial props. Noncryst. Violet col. with Ce(SO₄)₂, becoming brown on standing. λ_{\max} 293 ; 310 (MeOH) (Berdy).

▶ XW5525000

Bis(deoxy), di-N-de-Me, N⁴-oxide: **Bisnordihydrotoxiferine N-oxide**

[62569-70-4]

[69345-26-2]

C₃₈H₄₀N₄O 568.76

Alkaloid from *Strychnos dolichothyrsa* and *Strychnos afzelii* (Loganiaceae). Noncryst. Poss. artifact.

Bis(deoxy), di-N-de-Me, N⁴, N^{4'}-dioxide: **Bisnordihydrotoxiferine di-N-oxide**

[62520-57-4]

C₃₈H₄₀N₄O₂ 584.76

Alkaloid from *Strychnos dolichothyrsa* (Loganiaceae). Active against gram-positive bacteria and fungi. Noncryst. Poss. artifact.

Wieland, H. et al., *Annalen*, 1941, **547**, 156-179 (C-Dihydrotoxiferine, isol)

King, H. et al., *J.C.S.*, 1949, 3263-3271 (isol)

Schmid, H. et al., *Helv. Chim. Acta*, 1952, **35**, 1864-1879; 1953, **36**, 102-121 (isol, C-Alkaloid H)

Asmis, H. et al., *Helv. Chim. Acta*, 1954, **37**, 1983-1992; 1955, **38**, 1661-1668

(Bisnordihydrotoxiferine, Caracurine VI)

Bernaer, K. et al., *Helv. Chim. Acta*, 1958, **41**, 2293-2308; 1959, **42**, 461-463 (ir, struct, synth, C-Dihydrotoxiferine)

Battersby, A.R. et al., *J.C.S.*, 1960, 736-741 (struct, synth)

Grdinic, M. et al., *J.A.C.S.*, 1964, **86**, ; 3357-3363 (pmr)

Verpoorte, R. et al., *J. Nat. Prod.*, 1976, **39**, 357-362 (*Strychnos dolichothyrsa constits*)

Rosato, R.R. et al., *Toxicol. Appl. Pharmacol.*, 1976, **35**, 107-111 (tox)

Verpoorte, R. et al., *Org. Magn. Reson.*, 1977, **9**, 567 (cmr)

Wenkert, E. et al., *J.O.C.*, 1978, **43**, 1099-1105 (cmr)

Verpoorte, R. et al., *Planta Med.*, 1978, **33**, 237-242 (*Bisnordihydrotoxiferine di-N-oxide, activity*)

Tits, M. et al., *Planta Med.*, 1978, **34**, 57-61 (*Nordihydrotoxiferine*)

Verpoorte, R. et al., *Planta Med.*, 1982, **44**, 21-27 (*Bisnor-C-alkaloid H N-oxide, Bisnor-C-alkaloid H di-N-oxide*)

Massiot, G. et al., *J.O.C.*, 1983, **48**, 1869-1872 (*Caracurine VI, bibl*)

Massiot, G. et al., *Tetrahedron*, 1983, **39**, 3645-3656 (*Bisnordihydrotoxiferine*)

Massiot, G. et al., *Phytochemistry*, 1988, **27**, 3293-3304 (isol, cmr)

Delaude, C. et al., *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (occur, derivs, *Strychnos*)

Zlotos, D.P. et al., *Eur. J. Org. Chem.*, 2004, 2375-2380 (pmr, conformm)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, TH1000

Toxiferine II T-437

Strychnotoxine II

C₄₀H₄₈N₄O₄²⁺ 648.844

Struct. unknown. Not the same as C-Toxiferine II (see Calebassine, C-31). Alkaloid from calabash curare and *Strychnos toxifera* (Loganiaceae). Hexagonal prisms (Me₂CO aq.) (as dipicrate). $[\alpha]_D^{214}$ +216 (dec. (rapid heating) (as dipicrate)).

Wieland, H. et al., *Annalen*, 1941, **547**, 156

Wieland, T. et al., *Chem. Ber.*, 1952, **85**, 731-743

Toxiferine III T-438

C₂₀H₂₇N₂O⁺ 311.446

Struct. unknown. Alkaloid from bark of *Strychnos toxifera* (Loganiaceae). Tablets (H₂O) (as chloride). Mp 285° chloride.

King, H. et al., *J.C.S.*, 1949, 3263-3271

Toxiferine VIII T-439

C₂₂H₂₅N₂O₃⁺ 365.451

Minimum formula. Struct. unknown. Alkaloid from bark of *Strychnos toxifera* (Loganiaceae). Balls of needles (Me₂CO) (as picrate). Mp 300° (picrate).

King, H. et al., *J.C.S.*, 1949, 3263-3271

Battersby, A.R. et al., *J.C.S.*, 1960, 1848

Toxiferine XII T-440

C₃₉H₄₆N₄O²⁺ 586.819


Struct. unknown. Alkaloid from *Strychnos toxifera* bark (Loganiaceae). Mp 333° (darkens at ca. 230°) (as picrate). A methyleneindoline belonging to the Toxiferine group of alkaloids.

King, H. et al., *J.C.S.*, 1949, 3263-3271

Battersby, A.R. et al., *J.C.S.*, 1960, 1848

Toxin MG 30 T-441

MG 30




C₂₅H₄₅N₅O₂⁺ 447.663

Isol. from the venom of the spider, *Macrothele gigas*. $[\alpha]_D$ +3.5 (c, 0.43 in H₂O). Counterion not specified.

Yamaji, N. et al., *Tet. Lett.*, 2004, **45**, 5371-5373 (isol, synth, cd, pmr, cmr, ms)

Trabzonine T-442

[303175-76-0]




C₂₂H₃₃NO₃ 359.508

Alkaloid from the roots of *Aconitum nasutum*. Cryst. Mp 123-126°. $[\alpha]_D^{25}$ -5 (c, 1.2 in MeOH).

Mericli, A.H. et al., *Heterocycles*, 2000, **53**, 1987-1996

Trachelanthamine T-443

[14140-18-2]



C₁₅H₂₇NO₄ 285.383

Trachelanthamine and its stereoisomers are esters of 1-Hydroxymethylpyrrolizidine, H-629, various stereoisomers, with 2,3-Dihydroxy-2-isopropylbutanoic acid, various stereoisomers. Posn. 8 is 7a in CAS numbering. Alkaloid from *Trachelanthus korolkovii*, *Trachelanthus hissaricus* and *Rindera baldschuanica* (Boraginaceae). Cryst. (Me₂CO/petrol). Mp 92-93°. $[\alpha]_D$ -18.14 (c, 1 in EtOH).

▶ EK7787800

Picrate: Mp 155-156°.

N-Oxide: **Trachelanthine. Trachelanthamine N-oxide**

[510-19-0]

C₁₅H₂₇NO₅ 301.382

Alkaloid from *Trachelanthus korolkovii* (Boraginaceae). Needles (Me₂CO). Mp 166-167°. $[\alpha]_D$ -22.5.

▶ EK7788500

3'-Ac: **3'-O-Acetyltrachelanthamine**

[200062-29-9]

C₁₇H₂₉NO₅ 327.42

Alkaloid from *Heliotropium floridum*

(Boraginaceae). Oil. $[\alpha]_D^{25} +7$ (c, 0.04 in CH_2Cl_2).

8-Epimer: Lindelofine

[487-99-0]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from *Lindelofia anchusoides* air-dried plants and *Lindelofia macrostyla* (Boraginaceae). Cryst. (Me_2CO). Mp 106-107°. $[\alpha]_D^{25} +50$ (EtOH).

8-Epimer, picrate:

Cryst. (EtOH), Mp 123-124°.

8-Epimer, N-oxide: Lindelofine N-oxide

$\text{C}_{15}\text{H}_{27}\text{NO}_5$ 301.382

Alkaloid from *Lindelofia macrostyla* (Boraginaceae). Mp 195-196.5°. $[\alpha]_D^{25} +20.6$.

8-Epimer, O-tigloyl: Lindelofamine

$\text{C}_{20}\text{H}_{33}\text{NO}_5$ 367.484

Alkaloid from *Lindelofia anchusoides* air-dried plants (Boraginaceae). Cryst. (petrol). Mp 88°. Posn. of esterification not known.

2'-Epimer: Coromandaline

[68473-86-9]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from *Heliotropium curassavicum* (Boraginaceae). Pale yellow gum. $[\alpha]_D^{25} -6.9$ (c, 0.5 in EtOH). Contains the rare enantiomer (+)-Viridifloric acid.

3'-Epimer: Viridiflorine

[551-57-5]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from *Cynoglossum viridiflorum*, *Lindelofia olgae*, *Lindelofia pterocarpa*, *Lindelofia stylosa*, *Paracaryum himalayense*, *Symphytum officinale* and *Trachelanthus hissaricus* (Boraginaceae). Prisms (Me_2CO). Mp 102-103°. $[\alpha]_D^{25} -11.73$ (c, 1 in EtOH).

3'-Epimer, N-oxide: Viridiflorine N-oxide

[19038-28-9]

$\text{C}_{15}\text{H}_{27}\text{NO}_5$ 301.382

Alkaloid from *Lindelofia macrostyla* and *Lindelofia olgae* (Boraginaceae). Mp 197-198°.

1,2'-Diepimer: Heliocoromandaline

[82354-33-4]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Isol. from *Heliotropium curassavicum* (Boraginaceae). Gum.

2',3'-Diepimer: Heliovicine

[68473-85-8]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from *Heliotropium curassavicum* (Boraginaceae). Pale yellow gum. $[\alpha]_D^{25} -2.7$ (c, 0.6 in EtOH). Contains the rare enantiomer (-)-trachelanthic acid.

3',8-Diepimer: Cynaustaline

[17958-37-1]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from *Cynoglossum australe* (dried plants) (Boraginaceae). Gum. Mp 149-150° (as picrolonate). Apparently the enantiomer of Heliocoromandaline.

1,2',3'-Triepimer: Heliocurassavicine

[82354-34-5]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Minor alkaloid from *Heliotropium curassavicum* (Boraginaceae). Gum.

$[\alpha]_D^{25} +0.3$ (c, 0.0035 in CHCl_3).

Should be the enantiomer of Lindelofine, although opt. rotns. do not agree.

1,2',3'-Triepimer, N-oxide: Heliocurassavicine N-oxide

[156714-77-1]

$\text{C}_{15}\text{H}_{27}\text{NO}_5$ 301.382

Alkaloid from aerial parts of *Heliotropium curassavicum* (Boraginaceae).

1,3',8-Triepimer: Neocoromandaline

[283168-04-7]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Alkaloid from the roots of *Cynoglossum furcatum*. Gum. $[\alpha]_D^{25} -6.8$ (EtOH). Synonym misspelt in paper.

Tetraepimer: Heliocurassavicine

[82374-02-5]

$\text{C}_{15}\text{H}_{27}\text{NO}_4$ 285.383

Minor alkaloid from *Heliotropium curassavicum* (Boraginaceae). Gum. $[\alpha]_D^{25} +0.3$ (c, 0.0035 in CHCl_3). Should be the enantiomer of Trachelanthamine, but apparently is of v. low opt. purity.

Men'shikov, G.P. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1941, **11**, 209; 1945, **15**, 225; 1947, **17**, 343; 1948, **18**, 1736; *CA*, **35**, 7111; **40**, 2141; **42**, 556; **46**, 2625

(*Trachelanthamine, Viridiflorine*)

Labenskii, A.S. *et al.*, *J. Gen. Chem. USSR (Engl. Transl.)*, 1948, **18**, 1836; *CA*, **43**, 3827

(*Lindelofine, Lindelofamine*)

Tsyru'nikova, L.G. *et al.*, *Zh. Obshch. Khim.*, 1962, **32**, 2705-2709; *J. Gen. Chem. USSR (Engl. Transl.)*, 1962, **32**, 2663-2666

(*Lindelofine N-oxide*)

Akramov, S.T. *et al.*, *CA*, 1964, **61**, 4700;

11005; 1968, **68**, 47001 (*Viridiflorine, Viridiflorine N-oxide*)

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1967, **20**, 2499-2503 (*Cynaustaline*)

Akramov, S.T. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 351; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 296-297 (*Trachelanthine, Viridiflorine oxide*)

Kiyamitdinova, F. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 411-412; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 343-344 (*Viridiflorine*)

Kochetkov, N.K. *et al.*, *Tetrahedron*, 1969, **25**, 2313-2323 (*abs config, synth*)

Subramanian, P.S. *et al.*, *Aust. J. Chem.*, 1980, **33**, 1357-1363 (*Heliovicine, Coromandaline*)

Mohanraj, S. *et al.*, *Phytochemistry*, 1982, **21**, 1775-1779 (*Heliocurassavicine, Heliocurassavicine, Heliocoromandaline, Coromandaline, Heliocurassavicine*)

Roeder, E. *et al.*, *Phytochemistry*, 1990, **29**, 11-29 (*rev, cmr*)

Marquina, G. *et al.*, *Rev. Cubana Farm.*, 1992, **26**, 52; *CA*, **121**, 104063f (*Heliocurassavicine N-oxide*)

Reina, M. *et al.*, *Phytochemistry*, 1997, **46**, 845-853 (*3'-Acetyltrachelanthamine*)


Weber, S. *et al.*, *Phytochemistry*, 1999, **50**, 1005-1014 (*pmr, cmr, biosynth*)

Ravi, S. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 80-82 (*Neocoromandaline*)

Trachyone

T-444

1-[15-(3,4-Methylenedioxyphenyl)-14-pentadecenyl]pyrrolidine. 15-(3,4-Methylenedioxyphenyl)-14-pentadecenoic acid pyrrolidide



$\text{C}_{26}\text{H}_{39}\text{NO}_3$ 413.599

(E)-form [102965-07-1]


Alkaloid from *Piper amalago* and *Piper nigrum* (white pepper).

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12-18 (*isol*)

Reddy, S.V. *et al.*, *Phytomedicine*, 2004, **11**, 697-700 (*isol*)

Transvalencin A

T-445



$\text{C}_{23}\text{H}_{29}\text{ClN}_4\text{O}_6\text{S}_3$ 589.156

Isol. as Zn complex. Isol. from *Nocardia transvalensis* (IFM 10065). Antifungal agent.

Zn complex:


$\text{C}_{23}\text{H}_{27}\text{ClN}_4\text{O}_6\text{S}_3\text{Zn}$ 652.53

Prisms. Mp 280° dec. (shattered at 258°). λ_{max} 225 (log ϵ 4.47); 251 (log ϵ 4.05); 358 (log ϵ 3.65) (MeOH).

Hoshino, Y. *et al.*, *J. Antibiot.*, 2004, **57**, 797-802; 803-807 (*isol, cd, pmr, cmr, ms, cryst struct*)

Transvalencin Z

T-446



$\text{C}_{17}\text{H}_{21}\text{N}_3\text{O}_6$ 363.369


Isol. from *Nocardia transvalensis* (IFM 10065). Active against gram-positive bacteria. Powder. $[\alpha]_D^{25} +15.3$ (c, 1 in MeOH). λ_{max} 205 (log ϵ 3.77); 241 (log ϵ 3.28); 301 (log ϵ 2.91) (MeOH).

Mukai, A. *et al.*, *J. Antibiot.*, 2006, **59**, 366-369 (*isol, ir, pmr, cmr*)

Trehalamine

T-447

2-Amino-3a,5,6,6a-tetrahydro-4-(hydroxymethyl)-4H-cyclopentoxazole-4,5,6-triol, 9CI [144811-33-6]



$\text{C}_7\text{H}_{12}\text{N}_2\text{O}_5$ 204.182

Tautomeric with the 2-imino form. Found in culture broths of *Amycolatopsis* sp. and *Micromonospora* sp. Sol. H_2O ; poorly sol. Me_2CO , CHCl_3 . Mp 163-165°. $[\alpha]_D^{25} +13.5$ (c, 0.74 in H_2O).

N²-β-D-Glucopyranosyl: Trehalostatin.
Trehazolin. A 70615. Antibiotic A 70615
[132729-37-4]
C₁₃H₂₂N₂O₁₀ 366.324
Aminoglycoside antibiotic. Prod. by *Amycolatopsis trehalostatica* and *Micromonospora* strain SANK 62390. Potent and specific inhibitor of trehalase, active against blowfly (*Aldrichna grahami*). Amorph. powder. [α]_D²⁵ +115 (c, 1 in H₂O) (+ 99.5).

[147059-34-5]

- Murao, S. *et al.*, *Agric. Biol. Chem.*, 1991, **55**, 895 (*isol*, *pmr*, *props*)
Nakayama, T. *et al.*, *Chem. Comm.*, 1991, 919 (*pmr*, *cmr*, *struct*)
Ando, O. *et al.*, *J. Antibiot.*, 1991, **44**, 1165; 1993, **46**, 1116 (*isol*, *struct*)
Kobayashi, Y. *et al.*, *J.A.C.S.*, 1992, **114**, 10065; 1994, **59**, 813 (*synth*, *abs config*)
Ogawa, S. *et al.*, *J.C.S. Perkin 1*, 1992, **57**, 1939 (*synth*, *struct*)
Kobayashi, Y. *et al.*, *J. Antibiot.*, 1994, **47**, 932
Uchida, C. *et al.*, *J.C.S. Perkin 1*, 1994, 589 (*synth*, *pmr*, *abs config*)
Shiozaki, M. *et al.*, *J.O.C.*, 1994, **59**, 4450
Li, J. *et al.*, *J.O.C.*, 1998, **63**, 3403-3410 (*synth*, *Trehazolin*)
Boiron, A. *et al.*, *J.O.C.*, 1998, **63**, 5877-5883 (*synth*)
Kobayashi, Y. *et al.*, *Carbohydr. Res.*, 1999, **315**, 3-15 (*rev*, *synth*, *props*)
Berecibar, A. *et al.*, *Chem. Rev.*, 1999, **99**, 779-844 (*rev*, *synth*)
Storch de Gracia, I. *et al.*, *Org. Lett.*, 1999, **1**, 1705-1708 (*synth*)
Crimmins, M.T. *et al.*, *J.O.C.*, 2001, **66**, 4012-4018 (*Trehazolin*, *synth*)
Sugiyama, Y. *et al.*, *J. Antibiot.*, 2002, **55**, 263-269 (*Trehazolin*, *biosynth*)

Tremine

T-448

Struct. unknown. Alkaloid from fruits of *Trema micrantha* (Urticaceae). Rods (EtOH). Mp 103°. [α]_D +50 (EtOH).
Hydrochloride: Mp 194°.

Sulfate:

Powder. Mp 252°.

Glycoside: Tremidine


Alkaloid from *Trema micrantha* (Urticaceae). Plates (EtOH). Mp 53°. [α]_D²¹ +43 (EtOH). Neutral.

Ribeiro, O. *et al.*, *CA*, 1954, **48**, 1490

Trenidine

T-449

[79101-56-7]



C₃₆H₄₈ClN₃O₁₃ 766.24

Alkaloid from *Trewia nudiflora* (Euphorbiaceae). Cryst. (CH₂Cl₂/hexane). Sol. CHCl₃, MeOH; poorly sol. hexane, H₂O. Mp 200-205° dec. [α]_D²³ -114 (c, 0.24 in CHCl₃). λ_{max} 233 (ε 26400); 248 (sh) (ε 21500); 253 (ε 22600); 282 (ε 6130); 288 (ε 6130) (EtOH).

6'-Deoxy: Treflorine, 9CI

[79101-55-6]
C₃₆H₄₈ClN₃O₁₂ 750.241
Alkaloid from *Trewia nudiflora* (Euphorbiaceae). Cryst. (CH₂Cl₂/hexane). Sol. CHCl₃, MeOH; poorly sol. hexane, H₂O. Mp 205-208° dec. [α]_D²³ -138 (c, 0.045 in CHCl₃). λ_{max} 233 (ε 24000); 243 (sh) (ε 18500); 253 (ε 19850); 282 (ε 5060); 288 (ε 5060) (EtOH) (Derep). λ_{max} 233 (ε 24000); 253 (ε 19850); 282 (ε 5060); 288 (ε 5060) (EtOH) (Berdy).

6'-Ketone, N^{3'}-Me: N-Methyltrenudone

[82400-19-9]
C₃₇H₄₈ClN₃O₁₃ 778.251
Alkaloid from *Trewia nudiflora* (Euphorbiaceae). Cryst. (CH₂Cl₂/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 192-197° dec. [α]_D²³ -110 (c, 0.183 in CHCl₃). λ_{max} 233 (ε 27000); 247 (sh) (ε 21300); 252 (ε 21900); 282 (ε 5270); 289 (ε 5470) (EtOH).

15-Desmethoxy, 6'-ketone, N^{3'}-Me: Maytanbicyclinol

[221170-49-6]
C₃₆H₄₆ClN₃O₁₂ 748.225
Constit. of *Maytenus buchananii*. Amorph. solid. Mp 185-190°. Stereochem. at C-2' is S-.


Powell, R.G. *et al.*, *J.A.C.S.*, 1982, **104**, 4929-4934 (*Trenudine*, *Treflorine*, *N-Methyltrenudone*)

Larson, G.M. *et al.*, *J. Nat. Prod.*, 1999, **62**, 361-363 (*Maytanbicyclinol*)

Trewiasine

T-450

[78987-26-5]



C₃₇H₅₂ClN₃O₁₁ 750.284

Closely related to Maytansine, M-135. Alkaloid from *Trewia nudiflora* and mosses *Isotheicum subdiversiforme* and *Thamnobryum sandei*. Antineoplastic agent. Cryst. (CH₂Cl₂/hexane). Mp 182-185°. [α]_D²³ -94 (c, 0.15 in CHCl₃). Log P 3.03 (calc). λ_{max} 233 (ε 27000); 254 (ε 26200); 282 (ε 6700); 288 (ε 6000) (EtOH) (Berdy).

N²²-De-Me: Nortrewiasine

[88147-94-8]

C₃₆H₅₀ClN₃O₁₁ 736.257

Alkaloid from *Trewia nudiflora*. Cryst. (CH₂Cl₂/hexane). Mp 155-158°. [α]_D²³ -58 (c, 0.04 in CHCl₃).

N'-De-Me: Demethyltrewiasine

[78987-28-7]

C₃₆H₅₀ClN₃O₁₁ 736.257

Alkaloid from *Trewia nudiflora*. Cryst. (CH₂Cl₂/hexane). Mp 129-142°.

O¹⁵-De-Me: Colubrinol. NSC 196520

[50657-33-5]

C₃₆H₅₀ClN₃O₁₁ 736.257

Alkaloid from *Colubrina texensis*.

Antineoplastic agent. Mp 194-196°.

[α]_D²² -94 (c, 0.035 in CHCl₃). Log P

2.22 (calc). λ_{max} 233 (ε 17100); 253 (ε

14700); 280 (ε 4040); 289 (ε 3860)

(EtOH) (Berdy).

O¹⁵-De-Me, 15-Ac: Colubrinol acetate

[50499-79-1]

C₃₈H₅₂ClN₃O₁₂ 778.294

Alkaloid from *Colubrina texensis*.

Antineoplastic agent. Mp 179-182°.

[α]_D²² -127 (c, 0.073 in CHCl₃). Log P

3.07 (calc).

2'',3''-Didehydro: Dehydrotrewiasine

[78987-27-6]

C₃₇H₅₀ClN₃O₁₁ 748.268

Alkaloid from *Trewia nudiflora*. Cryst.

(CH₂Cl₂/hexane). Sol. MeOH, CHCl₃;

poorly sol. H₂O, hexane. Mp 165-

170°. [α]_D²³ -90 (c, 0.12 in CHCl₃).

λ_{max} 233 (ε 23600); 254 (ε 21000);

282 (ε 5240); 289 (ε 4870) (EtOH)

(Berdy).

10-Epimer: 10-Epitrewiasine

[88198-82-7]

C₃₇H₅₂ClN₃O₁₁ 750.284

Alkaloid from *Trewia nudiflora*. Cryst.

(CH₂Cl₂/hexane). Mp 159-162°. [α]_D²³ -

48 (c, 0.1 in CHCl₃).

Wani, M.C. *et al.*, *Chem. Comm.*, 1973, 390

(*Colubrinol*, *Colubrinol acetate*)

Powell, R.G. *et al.*, *J.O.C.*, 1981, **46**, 4398-4403

(*Trewiasine*, *Demethyltrewiasine*,

Dehydrotrewiasine)

Powell, R.G. *et al.*, *J. Nat. Prod.*, 1983, **46**,

660-666 (*10-Epitrewiasine*, *Nortrewiasine*)

Cassady, J.M. *et al.*, *Chem. Pharm. Bull.*, 2004,

52, 1-26 (*rev*)

Triacanthine


T-451

3-(3-Methyl-2-butenyl)-3H-purin-6-

amine, 9CI. 6-Amino-3-prenylpurine. To-

gholamine. Chidlovine

[10091-84-6]



C₁₀H₁₃N₅ 203.246

Alkaloid from leaves of *Holarrhena floribunda*, *Holarrhena mitis*, *Holarrhena congolensis* and another *Holarrhena* sp. (poss. *Holarrhena wulfsbergii*), also from *Gleditsia triacanthos* (Caesalpiaceae,

Apocynaceae), *Chidlowia sanguinea* (Fabaceae). Cryst. by subl. Mp 228-229°. The presence of Triacanthine in the plant is strongly dependent on the season of harvesting.

Hydrochloride:

Needles (EtOH). Mp 232-234° (218-219°).

Picrate: Mp 246° dec. (249-251°).

Belikov, A.S. *et al.*, *Zh. Obshch. Khim.*, 1954, **24**, 919 (*isol*)

Janot, M.-M. *et al.*, *Bull. Soc. Chim. Fr.*, 1959, 896

Leonard, N.J. *et al.*, *J.A.C.S.*, 1962, **84**, 2148 (*isol, ms, struct, synth*)

Leonard, J. *et al.*, *J.O.C.*, 1962, **27**, 1778 (*struct*)

Nellé, S. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1970, **271**, 153 (*isol*)

Leboeuf, M. *et al.*, *Phytochemistry*, 1972, **11**, 843 (*isol, occur*)

Kistenmacher, T.J. *et al.*, *J. Cryst. Mol. Struct.*, 1978, **7**, 219 (*cryst struct*)

1-Triacontylamine T-452

1-Triacontylamine. 1-Aminotriacontane
H₃C(CH₂)₂₈CH₂NH₂

C₃₀H₆₃N 437.834

Pentadecanoyl: N-Triacontylpentadecanamide. Platyamide A


C₄₅H₉₁NO 662.22

Alkaloid from *Platyaenia multicaule*. Cryst. [α]_D²⁵ +80 (c, 0.02 in Py). Optical rotation unaccounted for.

Ahmad, V.U. *et al.*, *Nat. Prod. Res.*, 2006, **20**, 69-74 (*Platyamide A*)

$\alpha, \alpha', 2$ -Triamino- $\gamma, \gamma', 3$ -trioxo-3H-phenoxazine-1,9-dibutanoic acid, 9CI T-453

[105334-59-6]



C₂₀H₁₈N₄O₈ 442.384

Ommochrome pigment from the eyes and skin of cephalopods *Loligo vulgaris*, *Sepia officinalis* and *Octopus vulgaris*.


Yellow cryst. Insoluble. Closely related to Xanthommatin, X-11. Readily cyclises to Dihydroxanthommatin, D-527. λ_{\max} 430 (no solvent reported).

Bolognese, A. *et al.*, *CA*, 1987, **106**, 15949 (*isol*)

Bolognese, A. *et al.*, *J. Het. Chem.*, 1988, **25**, 1243-1246 (*bibl*)

Triangularine T-454

[87340-27-0]



C₁₈H₂₅NO₅ 335.399

Diester of Retronecine, in T-188. Alkaloid from *Senecio triangularis* and *Alkanna tinctoria* (Asteraceae, Boraginaceae).

Pale-yellow oil. [α]_D²⁵ +2.2 (c, 1 in CHCl₃).

2'E-Isomer: Neotriangularine

[87392-67-4]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio triangularis* (Asteraceae). Yellow oil.

2'E-Isomer: Triangularine

[136173-28-9]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio hydrophyllus* and *Senecio mikanoides* (Asteraceae).

(2'E, 2'E)-Isomer: Neotriangularine

[136173-29-0]

C₁₈H₂₅NO₅ 335.399

Alkaloid from *Senecio hydrophyllus* (Asteraceae).

[136173-29-0]

Roitman, J.N. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1203 (*isol, pmr, cmr, ms*)

Röder, E. *et al.*, *Phytochemistry*, 1984, **23**, 2125 (*isol, cmr*)

Stelljes, M.A. *et al.*, *J. Nat. Prod.*, 1991, **54**, 759 (*Triangularine, Neotriangularine*)

Triantheme T-455

C₃₂H₃₆N₂O₆ 544.646

Struct. unknown. Alkaloid from *Trianthema monogyna*. Mp 127° Mp 112° (as picrate).

Basu, N.K. *et al.*, *Q. J. Pharm. Pharmacol.*, 1947, **20**, 38-42; *C.A.*, **41**, 7671f

4,8,12-Triazapentadecane-1,15-diamine T-456

N-(3-Aminopropyl)-N'-[3-[(3-aminopropyl)amino]propyl]-1,3-propanediamine, 9CI. 1,15-Diamino-4,8,12-triazapentadecane. 1,5,9,13,17-Pentaazaheptadecane.

Caldopentamine

[13274-42-5]

H₂N(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH(CH₂)₃NH₂

C₁₂H₃₁N₅ 245.41

Prod. by *Thermophilum album*, *Thermophilum minutum*, *Hydrogenobacter thermophilus*, *Acidothermus* spp. and *Thermus* spp. Also found in various insects and spiders. Bp_{0.2} 150-154°.

[107886-40-8]

Oshima, T. *et al.*, *J. Biol. Chem.*, 1982, **257**, 9913-9914 (*isol, struct*)

Niitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 1032-1038 (*synth, cmr*)

Osvath, P. *et al.*, *Aust. J. Chem.*, 1987, **40**, 347-360 (*synth, pmr, cmr, ms*)

Hamana, K. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1991, **100**, 399-402 (*isol*)

Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-747 (*Thermophilum constits, Hydrogenobacter constits*)

Tribenzylamine T-457

N,N-Bis(phenylmethyl)benzenemethanamine, 9CI. TBA

[620-40-6]

(PhCH₂)₃N

C₂₁H₂₁N 287.404

Isol. from hops *Humulus lupulus*. Used as 3% soln. in CHCl₃ for extraction separation of ion-associates with anionic complexes (e.g. SbCl₆⁻, Cr₂O₇²⁻); extraction-sepn. of V(III), Cd; extraction-photometric detn. of Mo (λ_{\max} 465 nm,

CHCl₃). Plates or prisms (Et₂O). Sol. Et₂O, CHCl₃. Mp 92°. Bp₁₃ 230°.

Hydrochloride: [7673-07-6]

Prisms (EtOH). Mp 227-228°.

Picrate: [67747-71-1]

Mp 191°.

N-Oxide: [6852-46-6]

C₂₁H₂₁NO 303.403

Cryst. (hexane/Me₂CO). Mp 138°.

N-Me: Tribenzylmethylammonium(1+)
[18265-23-1]

C₂₂H₂₄N⁺ 302.438

Needles (H₂O) (as iodide). Mp 184° (as iodide).

[134636-42-3]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 1279D (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **2**, 590B (*nmr*)

Sadtler Standard C-13 NMR Spectra, 2247 (*cmr*)

Sekiya, M. *et al.*, *Chem. Pharm. Bull.*, 1967, **15**, 802 (*synth*)

Suga, K. *et al.*, *Chem. Ind. (London)*, 1969, 78 (*synth*)

Iwasaki, F. *et al.*, *Acta Cryst. B*, 1972, **28**, 3370 (*cryst struct*)

Yokoi, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 1942 (*pmr*)

Yatirajam, V. *et al.*, *Mikrochim. Acta*, 1974, 671 (*detn, Mo*)

Singh, O.V. *et al.*, *J. Inorg. Nucl. Chem.*, 1975, **37**, 609 (*detn, Cd*)

Yatirajam, V. *et al.*, *Anal. Chim. Acta*, 1976, **86**, 209 (*detn, V*)

Sandell, E.B. *et al.*, *Photometric Determination of Traces of Metals*, General Aspects: Analytical Separations, Wiley, New York, 1978, (*use*)

Donaldson, E.M. *et al.*, *Talanta*, 1980, **27**, 779 (*detn, Cr*)

Cheng, K.L. *et al.*, *Handbook of Organic Analytical Reagents*, CRC Press Inc., Boca Raton, Florida, 1983, (*use*)

Vorbrüggen, H. *et al.*, *Chem. Ber.*, 1984, **117**, 1523 (*synth*)

Qu, Y. *et al.*, *Z. Naturforsch., C*, 2003, **58**, 640-641 (*isol*)

Tribromoacetic acid, 9CI, T-458 8CI

[75-96-7]

Br₃CCOOH

C₂HBr₃O₂ 296.741

Monoclinic prisms. Sol. H₂O, EtOH, Et₂O. Mp 131°. Bp 245° dec. pK_a 0.66 (25°, H₂O).

► Irritant.

Me ester: [3222-05-7]

C₃H₃Br₃O₂ 310.768

d 2.33. Bp₁₅ 103°. n_D²⁰ 1.5580.

Et ester: [599-99-5]

C₄H₅Br₃O₂ 324.794

Constit. of *Asparagopsis armata*. Liq. Bp 178-180° Bp₇₃ 148°.

► AJ7600000

Chloride: [34718-47-3]

C₂Br₃ClO 315.186

Liq. Bp₁₂ 80-82°.

Bromide: [10588-31-5]

C₂Br₄O 359.637

Liq. Bp 210-215° Bp₁₂ 88-90°.

Amide: Tribromoacetamide, 9CI

[594-47-8]

C₂H₂Br₃NO 295.756Isol. from the alga *Wrangelia* sp. Sol. Et₂O, hot EtOH. Mp 122°. Sublimes.

Nitrile: Tribromoacetonitrile, 9CI. Tribromocyanomethane

[75519-19-6]

C₂Br₃N 277.741Oil. Sol. EtOH, Et₂O, C₆H₆. Bp 170°.

Anhydride: [41936-89-4]

C₄Br₆O₃ 575.466Solid (CCl₄). Mp 63-64°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 1, 509B (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 1, 793C (nmr)

Sadtler Standard C-13 NMR Spectra, 12470 (cmr)

Sadtler Standard Proton NMR Spectra, 41588 (pmr)

U.S. Pat., 1936, 2 057 964; CA, 31, 112 (synth)

Köbrich, G. et al., Chem. Ber., 1968, 101, 3208 (synth)

McConnell, O. et al., Phytochemistry, 1977, 16, 367-374 (Et ester, isol)

Parkash, R. et al., Bull. Chem. Soc. Jpn., 1991, 64, 1443 (anhydride)

Márquez, F. et al., Spectrosc. Lett., 1992, 25, 821 (ir, Raman)


Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, 7, 4992-4993 (bromide, use)

Kigoshi, H. et al., Chem. Lett., 2004, 98-99 (amide, isol)

Sax, N.I. et al., Dangerous Properties of Industrial Materials, 5th edn., Van Nostrand Reinhold, 1979, 1040

2,4,7(3,4,7)-Tribromo-3(2)-chloro-1H-indole, 9CI T-459


[68124-91-4]

C₈H₃Br₃ClN 388.283Alkaloid from the marine red alga *Rhodophyllis membranacea*.

Brennan, M.R. et al., Tet. Lett., 1978, 1637-1640 (isol, pmr, struct)

2,4,6-Tribromo-3-hydroxyphenethylamine T-460

3-(2-Aminoethyl)-2,4,6-tribromophenol, 9CI. 2-(2,4,6-Tribromo-3-hydroxyphenyl)ethylamine

C₈H₈Br₃NO 373.869Me ether, N-formyl: 2,4,6-Tribromo-N-formyl-3-methoxyphenethylamine. **Lutamide A**

[294210-56-3]

C₁₀H₁₀Br₃NO₂ 415.907Alkaloid from the marine bryozoan *Amathia convoluta*. Needles (EtOH).

Mp 105°.

N₃O-Di-Me, N-formyl: 2,4,6-Tribromo-N-formyl-3-methoxy-N-methylphenethylamine. **Lutamide C**

[294210-58-5]


C₁₁H₁₂Br₃NO₂ 429.933Alkaloid from the marine bryozoan *Amathia convoluta*. Needles (EtOH). Mp 69-70°.

Hashima, H. et al., Bioorg. Med. Chem., 2000, 8, 1757-1766 (Lutamides)

2,4,5-Tribromo-1H-imidazole, 9CI T-461

Tribromoglyoxaline

[2034-22-2]

C₃HBr₃N₂ 304.766Isol. from the egg masses of the molluscs *Ceratosoma erinaceum*, *Trophon geversianus* and *Trunculariopsis trunculus*. Silky needles (AcOH). Mp 221°.► Exp. neurotoxic props. LD₅₀ (rat, orl) 34 mg/kg. NI8660000

N-Me: 2,4,5-Tribromo-1-methyl-1H-imidazole

[1003-91-4]

C₄H₃Br₃N₂ 318.793

Cryst. (AcOH aq.). Mp 88-89°.

N-4-Methylbenzenesulfonyl: [6595-50-2]

C₁₀H₇Br₃N₂O₃S 474.955

Solid. Mp 157°.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 616D (ir)

Balaban, E. et al., J.C.S., 1922, 947 (synth)

Stensiö, K.-E. et al., Acta Chem. Scand., 1973, 27, 2179 (synth)

Verschoyle, R.D. et al., Arch. Toxicol., 1984, 56, 109 (neurotox)

Iddon, B. et al., Chem. Comm., 1985, 1428

(deriv, synth)

Iddon, B. et al., Tet. Lett., 1986, 27, 1635 (haz, use)

Iddon, B. et al., J.C.S. Perkin I, 1987, 1445

(tox, props)


O'Connell, J.F. et al., Synthesis, 1988, 767

(synth, ir, pmr)

Benkendorff, K. et al., Nat. Prod. Res., 2004, 18, 427-431 (isol, synth, ms)

Schmidt, A. et al., Heterocycles, 2006, 68, 1393-1400 (N-tosyl)

Lewis, R.J. et al., Sax's Dangerous Properties of Industrial Materials, 8th edn., Van Nostrand Reinhold, 1992, THV450

2,3,5-Tribromo-1H-indole, 9CI T-462C₈H₄Br₃N 353.838

N-Me: 2,3,5-Tribromo-1-methyl-1H-indole

[70063-25-1]

C₉H₆Br₃N 367.865Alkaloid from the alga *Laurencia brongniartii* and from *Aplysia dactylomela*. Sol. MeOH, C₆H₆; fairly sol. hexane; poorly sol. H₂O. Mp 120-122°. λ_{max} 228 (ε 34000); 282 (ε 7300); 290 (ε 10000); 297 (ε 8000); 300 (sh) (EtOH) (Derep).

Carter, G.T. et al., Tet. Lett., 1978, 4479-4482 (isol, uv, pmr, ms, struct)

Liu, Y. et al., J. Nat. Prod., 2002, 65, 748-749 (synth)

Suárez-Castillo, O.R. et al., J. Nat. Prod., 2006, 69, 1596-1600 (synth)

2,3,6-Tribromo-1H-indole, 9CI T-463C₈H₄Br₃N 353.838Alkaloid from *Laurencia similis*. Cryst. Mp 105-107°. λ_{max} 216 (log ε 5.15); 240 (log ε 4.44) (CHCl₃).

N-Me: 2,3,6-Tribromo-1-methyl-1H-indole

[70063-24-0]

C₉H₆Br₃N 367.865Alkaloid from the alga *Laurencia brongniartii*. Mp 90.5-91°. λ_{max} 230 (ε 39000); 288 (ε 10000); 294 (ε 10000) (EtOH) (Derep).

Carter, G.T. et al., Tet. Lett., 1978, 19, 4479-4482 (N-Me, isol, pmr, cmr)

Ji, N.-Y. et al., Helv. Chim. Acta, 2007, 90, 385-391 (isol, pmr, cmr, ms)

2,3,7-Tribromo-1H-indole, 9CI T-464

[68234-21-9]

C₈H₄Br₃N 353.838Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 115-116°.

Brennan, M.R. et al., Tet. Lett., 1978, 19, 1637-1640 (isol, pmr, struct)

Erickson, K.L. et al., Synth. Commun., 1981, 11, 253-259 (synth, ir, pmr, ms)

2,4,6-Tribromo-1H-indole, 9CI T-465

[128367-88-4]

C₈H₄Br₃N 353.838Alkaloid from the Okinawan red alga *Laurencia brongniartii*. Cryst. (hexane/CCl₄). Mp 106-113°.

Tanaka, J. et al., Tetrahedron, 1989, 45, 7301-7310 (isol, ir, pmr, ms, struct)

3,4,6-Tribromo-1H-indole, 9CI T-466**Balanoglossol**

[74076-58-7]

C₈H₄Br₃N 353.838Isol. from *Balanoglossus carnosus*. Mp 95° (89-90° dec.).

Ac:

C₁₀H₆Br₃NO 395.876

Mp 240-242°.

Higa, T. et al., Comp. Biochem. Physiol., B: Comp. Biochem., 1980, 65, 525-530 (isol, pmr, ms, struct)

Ohta, T. et al., Heterocycles, 1987, 26, 2817-2822 (synth)

Martin, P. et al., Tet. Lett., 1987, 28, 1645-1647 (synth)

Higa, T. et al., Tetrahedron, 1987, 43, 1063-1070 (isol, pmr)

Martin, P. *et al.*, *Helv. Chim. Acta*, 1988, **71**, 344-347 (*synth, pmr*)

3,5,6-Tribromo-1H-indole T-467
[929624-69-1]

$C_8H_4Br_3N$ 353.838
Alkaloid from *Laurencia similis*. Cryst. Mp 123-124°. λ_{max} 218 (log ϵ 5.15); 241 (log ϵ 4.64) ($CHCl_3$).

N-Me: 3,5,6-Tribromo-1-methyl-1H-indole
[929624-70-4]
 $C_9H_6Br_3N$ 367.865
Alkaloid from *Laurencia similis*. Cryst. Mp 166-168°. λ_{max} 218 (log ϵ 5.17); 240 (log ϵ 4.51) ($CHCl_3$).

Ji, N.-Y. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 385-391 (*isol, pmr, cmr, ms*)


3,5,7-Tribromo-1H-indole, 9CI T-468
[59025-61-5]

$C_8H_4Br_3N$ 353.838
Isol. from the hemichordates *Ptychodera flava laysanica* and *Ptychodera flava*. Mp 120-122°.

Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 231-233 (*synth, pmr, ms*)

Higa, T. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1980, **65**, 525-530 (*occur*)

2,5,6-Tribromo-1H-indole-3-carboxaldehyde T-469
2,5,6-Tribromo-3-formylindole




$C_9H_4Br_3NO$ 381.849

N-Me: 2,5,6-Tribromo-1-methyl-1H-indole-3-carboxaldehyde
[85908-67-4]

$C_{10}H_6Br_3NO$ 395.875
Alkaloid from the marine bryozoan *Zoobotryon verticillatum*. Delays the metamorphosis in fertilised sea urchin eggs at low concentrations. Mp 228.5-229.5°. λ_{max} 223 (ϵ 18100); 256 (ϵ 13700); 309 (ϵ 6900) (MeOH) (Berdy).

Ortega, M.J. *et al.*, *J. Nat. Prod.*, 1993, **56**, 633-636 (*isol, uv, ir, pmr, cmr, ms, struct*)

5,6,7-Tribromo-6-methoxyindigotin T-470
[58933-46-3]




$C_{17}H_9Br_3N_2O_3$ 528.982

Pigment from the marine invertebrate *Ptychodera flava laysanica*. Purple-blue powder. Mp 300°.

Higa, T. *et al.*, *Heterocycles*, 1976, **4**, 227-230 (*isol, uv, struct*)

3,5,7-Tribromo-6-methoxy-1H-indole, 9CI T-471
6-Methoxy-3,5,7-tribromoindole
[74076-57-6]




$C_9H_6Br_3NO$ 383.864

Isol. from the hemichordate *Ptychodera flava*. Mp 110-112°.

Higa, T. *et al.*, *Comp. Biochem. Physiol., B: Comp. Biochem.*, 1980, **65**, 525-530 (*isol, pmr, ms, struct*)

2,5,6-Tribromo-N-methylgramine T-472
2,5,6-Tribromo-N,N,1-trimethyl-1H-indole-3-(methanamine, 9CI). 2,5,6-Tribromo-3-(dimethylaminomethyl)-1-methylindole
[64945-29-5]



$C_{12}H_{13}Br_3N_2$ 424.96

Alkaloid from the subtropical marine bryozoan *Zoobotryon verticillatum* and *Zoobotryon pellucidum*. Toxic to brine shrimp. Larval settlement inhibitor. Antifouling agent. Cryst. (MeOH). Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 112-113°. λ_{max} 232 (ϵ 46800); 298 (ϵ 9900); 308 (ϵ 9100) (MeOH) (Berdy). λ_{max} 232 (ϵ 46000); 298 (ϵ 9700); 307 (ϵ 9000) (EtOH) (Berdy).


l'-N- Oxide: 2,5,6-Tribromo-N-methylgramine N-oxide
[85908-65-2]

$C_{12}H_{13}Br_3N_2O$ 440.959

Alkaloid from *Zoobotryon verticillatum*. Toxic to brine shrimp. Cryst. (MeOH aq.). Sol. MeOH, EtOAc; poorly sol. H_2O . Mp 116-120° dec. λ_{max} 230 (ϵ 38800); 294 (ϵ 11800); 306 (ϵ 10700) (MeOH) (Berdy).

DaSettino, A. *et al.*, *Chim. Ind. (Milan)*, 1977, **59**, 454; *CA*, **87**, 201244c (*synth*)
Sato, A. *et al.*, *Tet. Lett.*, 1983, **24**, 481-484 (*isol, uv, ir, pmr, cmr, ms, struct, synth*)
Kon-Ya, K. *et al.*, *Biosci., Biotechnol., Biochem.*, 1994, **58**, 2178-2181

2,4,6-Tribromo-3-methylthio-1H-indole T-473
[128351-83-7]



$C_9H_6Br_3NS$ 399.931

Isol. from a Taiwanese collection of the red alga *Laurencia brongniartii*. Also found in the Okinawan red alga *Laurencia grevilleana* (possibly synonymous with *Laurencia brongniartii*).

S-Oxide: 2,4,6-Tribromo-3-(methylsulfinyl)-1H-indole

$C_9H_6Br_3NOS$ 415.931

Isol. from *Laurencia brongniartii*. Mp 102-104°. λ_{max} 236 (log ϵ 4.7); 312 (log ϵ 4.1) (MeOH).


5-Bromo: 2,4,5,6-Tetrabromo-3-methylthio-1H-indole
[128351-84-8]

$C_9H_5Br_4NS$ 478.827

Isol. from *Laurencia brongniartii* and *Laurencia grevilleana* (? *Laurencia brongniartii*).

Tanaka, J. *et al.*, *Tetrahedron*, 1989, **45**, 7301-7310 (*Laurencia brongniartii* constits)
El-Gamal, A.A. *et al.*, *J. Nat. Prod.*, 2005, **68**, 815-817 (*S-oxide*)

2,3,4-Tribromo-1H-pyrrole T-474
[69624-12-0]



$C_4H_2Br_3N$ 303.778

Isol. from the marine polychaete *Polypsiphia crassa* and *Saccoglossus kowalevskii*. Possesses antibacterial and antifeedant activity. Pale yellow oil. Dec. slowly above -18° in soln.

N-Sulfonic acid: [162821-17-2]

$C_4H_2Br_3NO_3S$ 383.843

Isol. from *Saccoglossus kowalevskii*. Isol. as Na salt, to which CAS no. refers.


Emrich, R. *et al.*, *J. Nat. Prod.*, 1990, **53**, 703-705 (*isol, pmr, ms, synth*)

Martin, P. *et al.*, *Marine Ecol.: Progr. Ser.*, 1995, **116**, 125-136; *CA*, **122**, 261393 (*l-sulfonic acid*)

John, E.A. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1929-1931 (*synth*)

Kicklighter, C.E. *et al.*, *Limnol. Oceanogr.*, 2004, **49**, 430-441 (*isol, activity*)

3,4,5-Tribromo-1H-pyrrole-2-carboxylic acid T-475
[74039-30-8]



$C_5H_2Br_3NO_2$ 347.788

Isol. from an Australian sponge *Axinella* sp. Plates (H_2O). Mp 192-195°. λ_{max} 258 (ϵ 12800) (MeOH).

Me ester: [1198-67-0]


$C_6H_4Br_3NO_2$ 361.815

Cryst. (hexane). Mp 212-214°.

N-Me: 3,4,5-Tribromo-1-methyl-1*H*-pyrrole-2-carboxylic acid
[150314-26-4]
C₆H₄Br₃NO₂ 361.815
Isol. from an *Axinella* sp. Cryst. Mp 175-176° dec. λ_{max} 254 (ε 13700); 272 (ε 4400) (MeOH).

Hodge, P. et al., *J.C.S.*, 1965, 459-470 (*synth, ester*)
Tashiro, M. et al., *J. Chem. Res., Synop.*, 1988, 136-137 (*synth*)
Barrow, R.A. et al., *Nat. Prod. Lett.*, 1993, **1**, 243-250 (*isol, synth, cmr*)

3,3',4-Tribromo-4',5,5'-trichloro-2,2'-bi-1*H*-pyrrole T-476




C₈H₂Br₃Cl₃N₂ 472.187

N,N'-Di-Me: 3,3',4-Tribromo-4',5,5'-trichloro-1,1'-dimethyl-2,2'-bi-1*H*-pyrrole.
DBP-Br₃Cl₃
[400766-93-0]
C₁₀H₆Br₃Cl₃N₂ 500.241
Marine natural product of unknown biogenic origin. Present in marine or freshwater fish, shrimps, seabirds and seal blubber.

Tittlemier, S.A. et al., *Environ. Pollut.*, 2002, **116**, 85-93 (*N,N'*-di-Me, *occur, detm*)
Tittlemier, S.A. et al., *J. Agric. Food Chem.*, 2004, **52**, 2010-2015 (*N,N'*-di-Me, *occur, detm*)

Trichamide T-477

[905455-92-7]




C₄₆H₆₆N₁₆O₁₂S₂ 1099.259
Isol. from the cyanobacterium *Trichodesmium erythraeum*.

Sudek, S. et al., *Appl. Environ. Microbiol.*, 2006, **72**, 4382-4387 (*isol, struct, ms*)

Trichanthin A T-478

[211184-05-3]



C₂₂H₂₇NO₇S 449.524

Related to Gerambullin, G-68. Constit. of *Glycosmis trichanthera*. Cryst. (Et₂O). Mp 154-156°. λ_{max} 224; 240 (sh); 296 (MeOH).

4'-Deoxy: *Glyparvin A*
[204641-51-0]

C₂₂H₂₇NO₆S 433.524
Constit. of *Glycosmis parva*. Cryst. (Et₂O). Mp 134-137°. λ_{max} 226; 239 (sh); 288 (MeOH).

4'-Deoxy, 2',3'-dihydro: *Dihydroglyparvin A*
[204591-96-8]

C₂₂H₂₉NO₆S 435.54
Constit. of *Glycosmis parva*. Cryst. (Et₂O). Mp 93-94°. [α]_D²⁰ +51 (c, 0.2 in CHCl₃). λ_{max} 223; 260 (MeOH).


(2'E)-Isomer: *Trichanthin B*

[211184-04-2]
C₂₂H₂₇NO₇S 449.524
Constit. of *Glycosmis trichanthera*. Cryst. (Et₂O). Mp 112-114°. λ_{max} 220; 240 (sh); 286 (sh); 292 (MeOH).

Hofer, O. et al., *Monatsh. Chem.*, 1998, **129**, 213-219 (*Glyparvin A, Dihydroglyparvin A*)
Vajrodaya, S. et al., *Phytochemistry*, 1998, **48**, 897-902 (*isol, uv, ir, pmr, cmr, ms*)

3α-(4,4,4-Trichloro-2-butyroxy)tropane T-479

[78416-85-0]




C₁₇H₂₂Cl₃N 346.726

Isol. from roots of *Scopolia tanguticus* (Solanaceae) along with Apoptropine. Almost certainly an artifact derived from Apoptropine, A-1359 by reactn. with CHCl₃.

Yang, J.S. et al., *CA*, 1981, **95**, 138446t

2,3,4-Trichloro-1*H*-indole, 9CI T-480

[68234-24-2]



C₈H₄Cl₃N 220.484

Alkaloid from the marine red alga *Rhodophyllis membranacea*.

Brennan, M.R. et al., *Tet. Lett.*, 1978, **19**, 1637-1640 (*isol, pmr, struct*)

2,3,7-Trichloro-1*H*-indole, 9CI T-481

[68234-18-4]

C₈H₄Cl₃N 220.484

Alkaloid from the marine red alga *Rhodophyllis membranacea*. Mp 86°.

Brennan, M.R. et al., *Tet. Lett.*, 1978, **19**, 1637-1640 (*isol, pmr, struct*)
Erickson, K.L. et al., *Synth. Commun.*, 1981, **11**, 253-259 (*synth, ir, pmr, ms*)

1,1,1-Trichloro-5-undecylamine T-482

1,1,1-Trichloro-5-undecanamine. 5-Amino-1,1,1-trichloroundecane
H₃C(CH₂)₅CH(NH₂)(CH₂)₃CCl₃
C₁₁H₂₂Cl₃N 274.66


(+)-form

N-Ac: 5-Acetamido-1,1,1-trichloroundecane
[345261-12-3]
C₁₃H₂₄Cl₃NO 316.697
Isol. from the cyanobacterium *Microcoleus lyngbyaceus*. Oil. [α]_D +60 (c, 0.02 in CHCl₃).

Orsini, M.A. et al., *J. Nat. Prod.*, 2001, **64**, 572-577 (*isol, ir, pmr, cmr*)

Trichochrome B T-483

Trichosiderin B
[35394-49-1]




C₂₃H₂₀N₄O₉S₂ 560.564

Constit. of human and mammalian red hair and birds' feathers. Found in urine of melanoma patients. Amorph. red-orange solid. Insol. H₂O, org. solvs. λ_{max} 243 (log ε 4.6); 329 (log ε 4.01); 454 (log ε 3.98) (0.2*M* NaOH).

Prota, G. et al., *Experientia*, 1971, **27**, 1145; 1381-1383 (*isol, struct*)
Prota, G. et al., *Experientia*, 1976, **32**, 1122-1124 (*isol*)
Agrup, G. et al., *Acta Derm. Venereol., Suppl.*, 1978, **58**, 269-270 (*isol*)

Trichochrome C T-484

Trichosiderin C
[32214-49-6]



C₂₃H₂₀N₄O₉S₂ 560.564


Constit. of red cock feathers. Found in the urine of melanoma patients. λ_{max} 240 (log ε 4.56); 327 (log ε 4.04); 452 (log ε 4.13) (0.2*M* NaOH).

Prota, G. et al., *Experientia*, 1976, **32**, 1122 (*isol, uv*)

Agrup, G. *et al.*, *Acta Derm. Venereol.*, Suppl., 1978, **58**, 269-270; Suppl., 1979, **59**, 453 (isol, synth, epr)

Trichochrome E**T-485**

[25942-18-1]


 $R^1 = \text{CH}_2\text{CH}(\text{NH}_2)\text{COOH}$, $R^2 = \text{H}$

$\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_6\text{S}_2$ 500.555
Isol. from chicken feathers.

Prota, G. *et al.*, *Gazz. Chim. Ital.*, 1969, **99**, 1193

Trichochrome F**T-486**

[25942-17-0]

As Trichochrome E, T-485 with

 $R^1 = \text{H}$, $R^2 = -\text{CH}_2\text{CH}(\text{NH}_2)\text{COOH}$

$\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_6\text{S}_2$ 500.555
Isol. from hair and feathers.


[131579-52-7]

Hansson, C. *et al.*, *Acta Derm. Venereol.*, Suppl., 1979, **59**, 453 (synth, epr)

Constantini, C. *et al.*, *Tetrahedron*, 1990, **46**, 6831 (synth, bibl)

Trichoderamide A**T-487**

Penicillazine
[508218-11-9]



Absolute Configuration

$\text{C}_{20}\text{H}_{20}\text{N}_2\text{O}_9$ 432.386

Isol. from the marine-derived fungi *Penicillium* sp. (strain No. 386), *Spicaria elegans* and *Trichoderma virens*. Also prod. by *Aspergillus unilateralis*. Cytotoxic. Needles (Me_2CO). Mp 258-260° (224-226°). $[\alpha]_{\text{D}}^{15} +128$ (c, 0.15 in MeOH). $[\alpha]_{\text{D}}^{25} -11.9$ (c, 2.2 in CHCl_3). λ_{max} 250 (sh) (log ϵ 3.98); 334 (log ϵ 4.23) (MeOH). λ_{max} 205 (log ϵ 4.6); 266 (log ϵ 4); 343 (log ϵ 4) (CHCl_3).

5-Deoxy, 5R-chloro: *Trichoderamide B*
[508218-12-0]

$\text{C}_{20}\text{H}_{19}\text{ClN}_2\text{O}_8$ 450.831

Isol. from *Trichoderma virens*. Cytotoxic. Oil. $[\alpha]_{\text{D}}^{15} +110.7$ (c, 0.15 in MeOH). λ_{max} 252 (log ϵ 3.56); 344 (log ϵ 3.86) (MeOH).

Lin, Y. *et al.*, *Tetrahedron*, 2000, **56**, 9607-9609 (*Trichoderma*)

Garo, E. *et al.*, *J. Nat. Prod.*, 2003, **66**, 423-426 (*Trichoderma*, pmr, cmr, cryst struct)


Liu, R. *et al.*, *Arch. Pharmacol Res.*, 2005, **28**, 1042-1046 (*Spicaria*, pmr, cmr, ms)

Capon, R.J. *et al.*, *Org. Biomol. Chem.*, 2005, **3**, 123-129 (*Aspergillus*, isol, struct)

Wan, X. *et al.*, *J.A.C.S.*, 2008, **130**, 17236-17237 (synth)

Trichodesmine**T-488**

14,19-Dihydro-12,13-dihydroxy-19-methylcrotalanan-11,15-dione, 9CI [548-90-3]



Absolute Configuration

$\text{C}_{18}\text{H}_{27}\text{NO}_6$ 353.414

Cyclic diester of 2,3,5,7a-Tetrahydro-1-hydroxy-1H-pyrrolizine-7-methanol, T-188 and trichodesmic acid. Alkaloid from *Trichodesma incanum*, *Heliotropium arguzioides*, *Crotalaria juncea*, *Crotalaria tetragona* and *Crotalaria rubiginosa* (Boraginaceae, Fabaceae). Prisms (Me_2CO). Mp 160-161° dec. $[\alpha]_{\text{D}} +38$ (10% EtOH aq.).

Methiodide:

Needles (EtOH). Mp 202° dec.

N-Oxide: *Trichodesmine N-oxide*

[55727-46-3]

$\text{C}_{18}\text{H}_{27}\text{NO}_7$ 369.414

Alkaloid from *Heliotropium arguzioides* and *Trichodesma incanum*. Cryst. (EtOH/ Me_2CO). Mp 169-170° dec.

13-Deoxy: *Incanine*†

[480-77-3]

$\text{C}_{18}\text{H}_{27}\text{NO}_5$ 337.415

Alkaloid from *Trichodesma incanum* and *Heliotropium olgae* (Boraginaceae). Mp 96-97°. $[\alpha]_{\text{D}} -58.4$ (H_2O). Originally reported as $\text{C}_{18}\text{H}_{29}\text{NO}_5$. Hydrol. gives retronecine and a mixt. of incanic and isoincanic acids.

13-Deoxy, N-Oxide: *Incanine N-oxide*

[55740-49-3]

$\text{C}_{18}\text{H}_{27}\text{NO}_6$ 353.414

Alkaloid from *Heliotropium olgae* and *Trichodesma incanum*. Cryst. (EtOH). Mp 168-169° dec.

17-Hydroxy: *Junceine*†

[480-53-5]

$\text{C}_{18}\text{H}_{27}\text{NO}_7$ 369.414

Alkaloid from *Crotalaria juncea* and *Crotalaria rubiginosa* (Fabaceae). Mp 191-192°. $[\alpha]_{\text{D}} -3$ (Py). Cyclic ester of retronecine with junceic acid.

19-Hydroxy: *Globiferine*

[90706-03-9]

$\text{C}_{18}\text{H}_{27}\text{NO}_7$ 369.414

Alkaloid from *Crotalaria globifera* seeds (Fabaceae). Cryst. (petrol). Mp 126-129°. $[\alpha]_{\text{D}}^{18} -8.6$ (c, 0.232 in CHCl_3).

Men'shikov, G.P. *et al.*, *Ber.*, 1935, **68**, 2039 (isol)

Adams, R. *et al.*, *J.A.C.S.*, 1956, **78**, 1919; 1922; 1926 (isol, struct, *Trichodesmine*, *Junceine*)

Yunusov, S.Yu. *et al.*, *CA*, 1957, **51**, 1539; 1958, **52**, 13017; 1959, **15**, 315 (*Incanine*, isol, struct)

Akramov, S.T. *et al.*, *CA*, 1964, **60**, 16209 (isol) Atal, C.K. *et al.*, *Aust. J. Chem.*, 1966, **19**,

2189 (isol, pmr, *Trichodesmine*, *Junceine*) Robins, D.J. *et al.*, *J.C.S.(C)*, 1969, 1386 (abs config)

Culvenor, C.C.J. *et al.*, *J.C.S.(C)*, 1971, 3653 (cd)


Rashkes, Ya.V. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 40; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 34 (ms)

Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 363; 363; 368; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 315; 319

(*Trichodesmine*, *Incanine*, cryst struct)

Stoekli-Evans, H. *et al.*, *Acta Cryst. B*, 1982, **38**, 1614 (cryst struct, *Junceine*)

Brown, K. *et al.*, *Phytochemistry*, 1984, **23**, 457 (*Globiferine*)

Trichoguttine**T-489**

$\text{C}_{19}\text{H}_{15}\text{NO}_3$ 305.332


Alkaloid from the leaves of *Guatteria sagotiana* (Annonaceae). Amorph. Doubtful structural assignment; props. of authentic synthetic material are not in accord with those reported for the nat. prod.

Rasamizafy, S. *et al.*, *J. Nat. Prod.*, 1986, **49**, 1078 (isol, uv, ir, pmr, ms)

Atanes, N. *et al.*, *Heterocycles*, 1987, **26**, 1183

Tricholomic acid**T-490**

α -Amino-3-oxo-5-isoxazolidineacetic acid, 9CI. 2-(3-Oxo-5-isoxazolidinyl)glycine [2644-49-7]



Absolute Configuration

$\text{C}_5\text{H}_8\text{N}_2\text{O}_4$ 160.129

Constit. of the mushroom *Tricholoma muscarium*. Said to be useful as a flavouring substance. Insecticide. Pesticide. Mycotoxin. Narcotic intoxicant. Cryst. (H_2O). Sol. H_2O . Mp 207° dec. $[\alpha]_{\text{D}}^{20} +80$ (c, 0.20 in H_2O). λ_{max} 215 (ϵ 3470) (H_2O) (Berdy).

Takemoto, T. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1183; 1230; *CA*, **62**, 7859b; 8121c (isol, struct)

Iwasaki, H. *et al.*, *Chem. Pharm. Bull.*, 1965, **13**, 753 (synth)

Kamiya, K. *et al.*, *Chem. Pharm. Bull.*, 1966, **14**, 1307 (synth)


Japan. Pat., 1969, 69 16 354; *CA*, **71**, 111463p (*mamuf*)

Japan. Pat., 1979, 79 148 782; *CA*, **92**, 198384u (synth)

Conti, P. *et al.*, *Tetrahedron*, 2007, **63**, 2249-2256 (synth)

Trichonine**T-491**

1-(1-Oxo-2,4-eicosadienyl)pyrrolidine, 9CI. N-Pyrrolidyleicosa-2,4-dienamide. 2,4-Eicosadienoic acid pyrrolidide. 2,4-Icosadienoic acid pyrrolidide. N-(2,4-Eicosadienyl)pyrrolidine



C₂₄H₄₃NO 361.61**(E,E)-form** [33169-28-7]

Alkaloid from the leaves of *Piper trichostachyon* (Piperaceae). Cryst. (petrol). Mp 65–67°.

Tetrahydro: N-Eicosanoylpyrrolidine. Eicosanoic acid pyrrolidide [33117-75-8]

C₂₄H₄₇NO 365.641

Alkaloid from *Piper amalago*. Cryst. (petrol). Mp 38–40°.

4,5-Dihydro: N-(2-Eicosenoyl)pyrrolidine. 2-Eicosenoic acid pyrrolidide [102934-32-7]

C₂₄H₄₅NO 363.626

Alkaloid from *Piper amalago*.

Singh, J. *et al.*, *Tet. Lett.*, 1971, 2119 (*uv, ir, pmr, ms, struct*)

Vig, B. *et al.*, *Indian J. Chem.*, 1972, **10**, 564 (*synth, uv, ir, pmr*)

Trost, B.M. *et al.*, *Tet. Lett.*, 1983, **24**, 4525 (*synth*)


Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**, 12–18 (*isol, derivs*)

Kaga, H. *et al.*, *Synlett*, 1994, 607 (*synth*)

Trichophylline

T-492

[88524-54-3]

C₂₁H₂₂N₂O₄ 366.416


Indole alkaloid prob. most closely related to the Aspidosperma type, poss. by oxidative cleavage and rearrangement of a didehydrotaberionine. Alkaloid from the roots of *Catharanthus trichophyllus* (Apocynaceae). Shows antitumour activity. Cryst. (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 230°. [α]_D²⁶ -209.5. λ_{max} 238 ; 306 ; 362 (EtOH) (Berdy).

Mukhopadhyay, S. *et al.*, *Tetrahedron*, 1983, **39**, 3639 (*isol, uv, ir, pmr, ms, cryst struct*)

Trichoponamic acid

T-493

4-Methyl-6-(2-oxobutyl)-2-piperidinecarboxylic acid, 9CI [74235-25-9]



Absolute configuration

C₁₁H₁₉NO₃ 213.276

Isol. from *Trichoderma polysporum* as artifact derived from Trichopolyn. Mp 206–214° dec. [α]_D²¹ 0. [α]_D²⁸⁶ +193.

6-Epimer: Epitrichoponamic acid [79026-24-7]

C₁₁H₁₉NO₃ 213.276


Isol. from *Trichoderma polysporum*.

Fujita, T. *et al.*, *Chem. Comm.*, 1981, 585

Trichosanatine

T-494

[169626-16-8]

C₂₇H₂₈N₂O₄ 444.529


Alkaloid from *Trichosanthes rosthornii*. Cryst. Mp 175.5–176°.

Chao, Z.M. *et al.*, *Yaoxue Xuebao*, 1995, **30**, 517–520 (*isol*)

Trichostachine

T-495

1-[5-(1,3-Benzodioxol-5-yl)]-1-oxo-2,4-pentadienylpyrrolidine, 9CI. 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid pyrrolidide. 1-Piperoylpyrrolidine. Pyrroperine. Piperylene



(E,E)-form

C₁₆H₁₇NO₃ 271.315**(2E,4E)-form** [25924-78-1]

Alkaloid from *Piper nigrum* (pepper) and *Piper guineense* (Piperaceae). Cryst. (EtOAc/hexane or C₆H₆/petrol). Mp 143–145°. λ_{max} 245 (ε 10500); 261 (ε 10800); 309 (ε 19200); 343 (ε 33500) (MeOH).

(2E,4Z)-form

Alkaloid from the leaves of *Piper arboreum*. Amorph. solid. λ_{max} 280 (ε 2890) (MeOH).

(2Z,4E)-form

Alkaloid from the roots of *Piper nigrum* (pepper). Oil. λ_{max} 263 (log ε 3.92); 309 (log ε 4.01); 346 (log ε 4.12) (MeOH).

Grewe, R. *et al.*, *Chem. Ber.*, 1970, **103**, 3752 (*isol, uv, ir, pmr, ms, struct, synth*)

Okogun, J.I. *et al.*, *J.C.S. Perkin 1*, 1974, 2195 (*isol*)

Olsen, R.A. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 942 (*synth, uv, ir, pmr*)

Jacobs, H. *et al.*, *J. Indian Chem. Soc.*, 1999, **76**, 713–717 (*isol, pmr, cmr*)


Vasques da Silva, R. *et al.*, *Phytochemistry*, 2002, **59**, 521–527 (**2E,4Z-form**)

Wei, K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1005–1009 (**2Z,4E-form**)

Trichostatin A

T-496

7-[4-(Dimethylamino)phenyl]-N-hydroxy-4,6-dimethyl-7-oxo-2,4-heptadienamide, 9CI. A 300I. Antibiotic A 300I [58880-19-6]

C₁₇H₂₂N₂O₃ 302.372

Isol. from *Streptomyces hygroscopicus* and *Streptomyces sioyaensis*.

Antifungal antibiotic. Potent differentiation inducer of Friend leukaemia.

Inhibits histone deacetylase. Cell cycle inhibitor. Powder (EtOAc). Sol. MeOH, Et₂O; poorly sol. hexane, H₂O. Mp 172–174° (150.1°) dec. [α]_D^{20.5} +62.8 (c, 1.007 in EtOH). λ_{max} 252 (sh) (ε 16000); 265 (ε 17600); 341 (ε 19600) (EtOH) (Derep). λ_{max} 253 (E1%/1cm 477); 263 (ε 23600); 267 (E1%/1cm 495); 341 (E1%/1cm 776); 342 (ε 25100) (MeOH) (Berdy). λ_{max} 249 (E1%/1cm 500); 343 (E1%/1cm 800) (MeOH-NAOH) (Berdy). λ_{max} 264 (ε 24100); 265 (E1%/1cm 582); 341 (E1%/1cm 648); 342 (ε 27000) (EtOH) (Berdy).

▶MI5215000

Fe complex: Trichostatin B. A 300-II. Antibiotic A 300-II

[58895-00-4]

C₅₁H₆₆FeN₆O₉ 962.964

Dark reddish-purple prisms + H₂O (MeOH). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 192° dec. λ_{max} 252 (sh) (ε 16000); 265 (ε 17600); 341 (ε 19600) (EtOH) (Derep). λ_{max} 262 (E1%/1cm 680); 345 (E1%/1cm 920); 450 (E1%/1cm 50) (MeOH) (Berdy). λ_{max} 277 (E1%/1cm 651); 351 (E1%/1cm 918); 450 (E1%/1cm 50) (EtOH) (Berdy).

O-α-D-Glucopyranoside: Trichostatin DC₂₃H₃₂N₂O₈ 464.514

From *Streptomyces violaceusniger*.

Antitumour agent. Powder. Mp 131–135°. [α]_D²¹ +183 (c, 0.56 in MeOH). λ_{max} 267 (ε 27000); 345 (ε 27000) (MeOH).

O-β-D-Glucopyranoside: Trichostatin C.*Antibiotic 145-A*

[68676-88-0]

C₂₃H₃₂N₂O₈ 464.514

From *Streptomyces hygroscopicus* and *Streptomyces platensis*. Active against trichophytons, fungi and tumours.

Prisms (MeOH). Mp 171–173°. [α]_D²⁴ +50.5 (c, 0.987 in MeOH). λ_{max} 268 (ε 14600); 343 (ε 13200) (MeOH) (Derep).

Di-Ac:

Amorph. powder. [α]_D^{20.5} +70.8 (c, 0.891 in CHCl₃).

N-Deoxy: Antibiotic FL 657C. FL 657C

[127349-34-2]

C₁₇H₂₂N₂O₂ 286.373

Incorrectly named in CA, although the struct. shown is correct. Isol. from a *Streptomyces* sp. Possesses antitumour props. λ_{max} 250 ; 350 (MeOH) (Berdy).

N-Deoxy, N-Me: Trichostatin RKC₁₈H₂₄N₂O₂ 300.4

Prod. by *Streptomyces* sp. RK98-A74. λ_{max} 265 ; 350 (no solvent reported).

Tsuji, N. *et al.*, *J. Antibiot.*, 1976, **29**, 1; 1978, **31**, 939 (*isol, ir, uv, pmr, cmr, struct*)

Fleming, I. *et al.*, *Tetrahedron*, 1983, **39**, 841 (*synth*)


Yoshida, M. *et al.*, *Agric. Biol. Chem.*, 1985, **49**, 563 (*Trichostatin C*)

Morioka, H. *et al.*, *Agric. Biol. Chem.*, 1988, **52**, 251 (*isol, props*)

- Mori, K. *et al.*, *Tetrahedron*, 1988, **42**, 6013 (synth)
 Japan. Pat., 1989, 89 299 264; *CA*, **112**, 234003 (FL 657C)
 Yoshida, M. *et al.*, *Ann. N.Y. Acad. Sci.*, 1999, **886**, 23-36 (*Trichostatin A*, activity)
 Hayakawa, Y. *et al.*, *J. Antibiot.*, 2000, **53**, 179-183 (*Trichostatin D*)
 Ueki, M. *et al.*, *J. Antibiot.*, 2001, **54**, 1093-1095 (*Trichostatin RK*)
 Hosokawa, S. *et al.*, *Tet. Lett.*, 2005, **46**, 333-337 (*Trichostatin D*, synth)
 Zhang, S. *et al.*, *Adv. Synth. Catal.*, 2006, **348**, 1228-1234 (*Trichostatin A*, synth)

Trichotomine**T-497**

[53472-14-3]

C₃₀H₂₀N₄O₆ 532.511

Blue pigment from the fruits of *Clerodendrum trichotomum* (Verbenaceae). Shows bronchodilator, hypotensive and sedative props. Amorph. Mp 300°. The sp. *Clerodendron* is used in the paper.

N-β-D-Glucopyranosyl: **Trichotomine G₁** [56079-56-2]

C₃₆H₃₀N₄O₁₂ 710.653

Alkaloid from fruit of *Clerodendrum trichotomum* (Verbenaceae). Amorph. blue powder. Mp 300°.

N,N'-Di-D-glucopyranosyl: **N,N'-Di(D-glucopyranosyl)trichotomine**

C₄₂H₄₀N₄O₁₇ 872.795

Alkaloid from fruit of *Clerodendrum trichotomum* (Verbenaceae). Mixt. of anomers.

Di-Me ester: Mp 285-287°.

N,N'-Di-Ac, di-Me ester: Mp 246-248°.


Iwadare, S. *et al.*, *Tetrahedron*, 1974, **30**, 4105 (isol, ir, uv, pmr, cd, struct)

Kapadia, G.J. *et al.*, *Tet. Lett.*, 1977, 975 (synth)

Iwadare, S. *et al.*, *Tetrahedron*, 1978, **34**, 1457 (synth)

Triclisine**T-498**

5,6-Dimethoxyindeno[1,2,3-ij]isoquinoline, 9CI. 5,6-Dimethoxy-1-azafluoranthene [60888-60-0]

C₁₇H₁₃NO₂ 263.295

Alkaloid from *Triclisia gillettii* (Menispermaceae). Yellow cryst. (C₆H₆/petrol). Mp 155°.

9-Hydroxy: 9-Hydroxy-5,6-dimethoxyindeno[1,2,3-ij]isoquinoline. 5,6-Dimethoxyindeno[1,2,3-ij]isoquinolin-9-ol, 9CI. **Telitoxine**

[78416-88-3]


C₁₇H₁₃NO₃ 279.295

Alkaloid from the stems of *Telitoxicum peruvianum* (Menispermaceae). Yellow flakes (MeOH/CH₂Cl₂/CHCl₃). Mp 273-275°.

Huls, R. *et al.*, *Bull. Soc. R. Sci. Liege*, 1976, **45**, 40-45; *CA*, **85**, 160372f (isol, uv, ir, pmr, ms, struct)

Menachery, M.D. *et al.*, *J. Nat. Prod.*, 1981, **44**, 320-323; 1987, **50**, 726-729 (*Telitoxine*, isol, uv, pmr, struct, synth)

Ramana, M.M.V. *et al.*, *Tet. Lett.*, 2005, **46**, 4385-4386 (synth)

3-(7,16-Tricosadienyl)-1H-pyrrole-2-carboxaldehyde, 9CI**T-499**C₂₈H₄₇NO 413.685

(E,E)-form [57992-58-2]

Isol. from the marine sponge *Oscarella lobularis*.


7',8'-Dihydro: 3-(16-Tricosenyl)-1H-pyrrole-2-carboxaldehyde, 9CI

[57992-56-0]

C₂₈H₄₉NO 415.701

Isol. from *Oscarella lobularis*.

Cimino, G. *et al.*, *Experientia*, 1975, **31**, 1387 (isol, uv, pmr, ms, struct)

Trideacetylpyripyropene A**T-500**C₂₅H₃₁NO₇ 457.522

Terpenoid numbering shown. Parent substance of a large group of closely related Pyripyropenes prod. by *Aspergillus fumigatus* and *Eupenicillium reticulosporum*. Pyripyropenes are potent inhibitors of cholesterol acyltransferase and acylcoenzyme A (mostly investigated as mixtures).

3,7,19-Tri-Ac: **Pyripyropene A**

[147444-03-9]

C₃₁H₃₇NO₁₀ 583.634

Shows insecticide and MDR reversing activity. Powder. [α]_D²⁸ +65.8 (c, 1 in CHCl₃). λ_{max} 231 (ε 24300); 320 (ε 13400) (MeOH).

3-Propanoyl, 7,19-di-Ac: **Pyripyropene D** [151519-46-9]

C₃₂H₃₉NO₁₀ 597.661

Powder. [α]_D¹⁸ +64.5 (c, 1 in CHCl₃).

7-Propanoyl, 3,19-di-Ac: **Pyripyropene C** [151519-45-8]

C₃₂H₃₉NO₁₀ 597.661

Powder. [α]_D¹⁸ +9.4 (c, 1 in CHCl₃).

19-Propanoyl, 3,7-di-Ac: **Pyripyropene B** [151519-44-7]

C₃₂H₃₉NO₁₀ 597.661

Powder. [α]_D¹⁸ +62 (c, 1 in CHCl₃). λ_{max} 231 ; 320 (MeOH).

3,7-Dipropanoyl, 19-Ac: **Pyripyropene J** [165467-60-7]

C₃₃H₄₁NO₁₀ 611.688

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 248-250° dec. [α]_D²⁸ +65 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

3,19-Dipropanoyl, 7-Ac: **Pyripyropene K** [165467-61-8]

C₃₃H₄₁NO₁₀ 611.688

Cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁸ +58 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

7,19-Dipropanoyl, 3-Ac: **Pyripyropene L** [165467-62-9]

C₃₃H₄₁NO₁₀ 611.688

Cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 148-150°. [α]_D²⁸ +58 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

3,17,19-Tripropanoyl: **Pyripyropene I** [165467-59-4]

C₃₄H₄₃NO₁₀ 625.714

Cryst. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁸ +69 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

7,19-Dideoxy, 3-Ac: **Pyripyropene G** [165467-57-2]

C₂₇H₃₃NO₆ 467.561

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. Mp 248-250° dec. [α]_D²⁸ +102 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

7,19-Dideoxy, 3-propanoyl: **Pyripyropene H**

[165467-58-3]

C₂₈H₃₅NO₆ 481.588

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁸ +104 (c, 1 in MeOH). λ_{max} 231 ; 320 (MeOH).

7-Deoxy, 3,19-dipropanoyl: **Pyripyropene N**

[176385-03-8]

C₃₁H₃₉NO₈ 553.651

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. [α]_D²⁸ +97 (c, 0.05 in MeOH). λ_{max} 231 ; 320 (MeOH).

7-Deoxy, 19-propanoyl, 3-Ac: **Pyripyropene O**

[176385-06-1]

C₃₀H₃₇NO₈ 539.624

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{max} 231 ; 320 (MeOH).

11-Deoxy, 7-propanoyl, 3,19-di-Ac: **Pyripyropene M**

[176385-01-6]

C₃₂H₃₉NO₉ 581.661

Powder. Sol. MeOH, EtOAc; poorly sol. H₂O. λ_{max} 231 ; 320 (MeOH).

7,11-Dideoxy, 3,19-di-Ac: **Pyripyropene O**
[176385-04-9]
C₂₉H₃₅NO₇ 509.598
Powder. Sol. MeOH, EtOAc; poorly
sol. H₂O. [α]_D²⁸ +103 (c, 0.2 in MeOH).
λ_{max} 231 ; 320 (MeOH).

7,11-Dideoxy, 3-propanoyl, 19-Ac: **Pyri-
pyropene R**
C₃₀H₃₇NO₇ 523.625
Powder. Sol. MeOH, EtOAc; poorly
sol. H₂O. [α]_D²⁸ +146 (c, 0.1 in MeOH).
λ_{max} 231 ; 320 (MeOH).

7,11-Dideoxy, 19-propanoyl, 3-Ac: **Pyri-
pyropene P**
[176385-05-0]
C₃₀H₃₇NO₇ 523.625
Powder. Sol. MeOH, EtOAc; poorly
sol. H₂O. [α]_D²⁸ +95 (c, 0.4 in MeOH).
λ_{max} 231 ; 320 (MeOH).

7,11,19-Trideoxy, 3-Ac: **Pyripyropene E.**
**GERI-BP001 M. Antibiotic GERI-
BP001 M**
[156980-53-9]
C₂₇H₃₃NO₅ 451.561
Isol. from *Aspergillus fumigatus*. Inhi-
bitor of acyl-CoA : cholesterol acyl
transferase (ACAT). Cryst. (MeOH).
Mp 174-176°. [α]_D²⁸ +113 (c, 1 in
MeOH). [α]_D¹⁸ +146 (c, 0.5 in CHCl₃).
λ_{max} 232 (ε 19200); 322 (ε 11100)
(MeOH).

7,11,19-Trideoxy, 3-propanoyl: **Pyripyro-
pene F. GERI-BP001 B. Antibiotic**
GERI-BP001 B
[165467-56-1]
C₂₈H₃₅NO₅ 465.588
Isol. from *Aspergillus fumigatus*.
Inhibitor of ACAT. Powder. [α]_D²⁸
+122 (c, 1 in MeOH) (Pyripyropene
F). [α]_D¹⁸ +59.5 (c, 0.5 in CHCl₃).
λ_{max} 232 (ε 20200); 322 (ε 11500)
(MeOH).

11-Epimer, 7,19-dideoxy,3-Ac: **GERI-
BP001 A. Antibiotic GERI-BP001 A**
[168959-29-3]
C₂₇H₃₃NO₆ 467.561
Isol. from *Aspergillus fumigatus*. Inhi-
bitor of ACAT. Powder. Sol. MeOH,
EtOAc, CHCl₃; poorly sol. hexane,
H₂O. [α]_D¹⁸ +39.6 (c, 0.5 in CHCl₃).
λ_{max} 232 (ε 8400); 322 (ε 3600)
(MeOH).

Tomoda, H. *et al.*, *J. Antibiot.*, 1994, **47**, 148-
153; 154-162; 1995, **48**, 495-503; 1996, **49**,
292-298 (*isol, uv, ir, pmr, cmr, ms, cryst
struct*)

Tomoda, H. *et al.*, *J.A.C.S.*, 1994, **116**, 12097-
12098 (*abs config*)

Wang, H.-J. *et al.*, *Appl. Environ. Microbiol.*,
1995, **61**, 4429-4435 (*activity, Pyripyropene
A*)

Obata, R. *et al.*, *J. Antibiot.*, 1995, **48**, 749-750;
49, 1149-1156 (*activity, Pyripyropene A*)

Jeong, T.-S. *et al.*, *J. Antibiot.*, 1995, **48**, 751-
755 (*GERI BP001*)

Nagamitsu, T. *et al.*, *J.O.C.*, 1995, **60**, 8126-
8127 (*synth*)

Tomoda, H. *et al.*, *J.O.C.*, 1996, **61**, 882-886
(*biosynth*)

Smith, A.B. *et al.*, *Tet. Lett.*, 1996, **37**, 6461-
6464 (*synth*)

Aggarwal, V.K. *et al.*, *J.C.S. Perkin 1*, 1999,
3315-3321 (*synth, bibl*)

**2,7-Tridecadiene-10,12-diy-
noic acid** T-501
HC≡CC≡CCH₂CH=CH(CH₂)₃CH=
CHCOOH
C₁₃H₁₄O₂ 202.252

(2E,7Z)-form

2-Methylpropylamide: **2,7-Tridecadiene-
10,12-dienoic acid isobutylamide.** N-(2-
Methylpropyl)-2,7-tridecadiene-10,12-
diynamide, 9CI
[87797-74-8]
C₁₇H₂₅NO 257.375
Alkamide from the aerial parts of
Echinacea purpurea, *Echinacea angu-
stifolia* (Asteraceae) and flower heads
of *Spilanthes acmella*. Cryst. (petrol) or
gum. Mp 40°.

2-Methylbutylamide: **2,7-Tridecadiene-
10,12-dienoic acid 2-methylbutylamide**
[87797-76-0]
C₁₈H₂₅NO 271.402
Isol. from aerial parts of *Echinacea
purpurea*. Gum.

Bohlmann, F. *et al.*, *Phytochemistry*, 1983, **22**,
1173 (*isol, ir, pmr, ms, struct*)

Bauer, R. *et al.*, *Phytochemistry*, 1988, **27**,
2339; 1989, **28**, 505 (*isol, ir, pmr, ms,
struct*)

Nakatani, N. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 1992, **56**, 759 (*isol, uv, ir, pmr,
cmr*)


2,4-Tridecadienoic acid T-502
H₃C(CH₂)₇CH=CHCH=CHCOOH
C₁₃H₂₂O₂ 210.316

(2E,4E)-form

2-Methylpropylamide: **2,4-Tridecadienoic
acid isobutylamide.** N-(2-Methylpro-
pyl)-2,4-tridecadienamamide
C₁₇H₃₁NO 265.438
Alkaloid from *Leucocyclus formosus*
(Asteraceae). Obt. only in admixture
with homologues.

Gregor, H. *et al.*, *Phytochemistry*, 1981, **20**,
2579 (*isol, ms, struct*)

Tridecanochelerythrine T-503
8-(2-Oxotridecyl)dihydrochelerythrine
[152753-15-6]



C₃₄H₄₃NO₅ 545.717
Alkaloid from bark of *Zanthoxylum
integrifolium* (Rutaceae). Mp 68-72°.
Numbering systems vary. λ_{max} 229 (ε
4073); 283 (ε 4785); 320 (sh) (ε 1514)
(MeOH).

Jen, C.-M. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2019-
2021 (*isol, uv, ir, pmr, cmr, ms, struct*)

**2,6,8-Tridecatriene-10,12-
dienoic acid** T-504
HC≡CC≡CCH=CHCH=
CHCH₂CH₂CH=CHCOOH
C₁₃H₁₂O₂ 200.237

(2E,6E,8Z)-form

2-Methylpropylamide: **2,6,8-Tridecatriene-
10,12-dienoic acid isobutylamide.** N-(2-
Methylpropyl)-2,6,8-tridecatriene-
10,12-diynamide, 9CI
[87797-78-2]
C₁₇H₂₁NO 255.359
Alkaloid from the aerial parts of
Echinacea purpurea (Asteraceae).
Gum.

Bohlmann, F. *et al.*, *Phytochemistry*, 1983, **22**,
1173 (*isol, ir, pmr, ms, struct*)

**7-Tridecene-10,12-dienoic
acid, 9CI** T-505
HC≡CC≡CCH₂CH=
CH(CH₂)₅COOH
C₁₃H₁₆O₂ 204.268

(Z)-form [142505-97-3]

2-Methylpropylamide: **N-Isobutyl-7-tride-
cene-10,12-diynamide**
C₁₇H₂₅NO 259.391
Isol. from flower heads of *Spilanthes
acmella*. Oil.

Nakatani, N. *et al.*, *Biosci., Biotechnol.,
Biochem.*, 1992, **56**, 759 (*isol, ms, ir, cmr, pmr*)


6-Tridecylamine T-506
**6-Tridecanamine. 1-Pentyl-octylamine. 6-
Aminotridecane**
H₃C(CH₂)₆CH(NH₂)(CH₂)₄CH₃
C₁₃H₂₉N 199.379

(ξ)-form

N-Ac: **6-Acetamidotridecane**
[345261-17-8]
C₁₅H₃₁NO 241.416
Isol. from the cyanobacterium *Micro-
coleus lyngbyaceus*. Cryst.

Orsini, M.A. *et al.*, *J. Nat. Prod.*, 2001, **64**,
572-577 (*isol, ir, pmr, cmr*)

Tridentatol A T-507
[185548-48-5]



C₁₁H₁₃NOS₂ 239.362
Metab. from the marine hydroid *Triden-
tata marginata*. Amorph. solid. λ_{max} 337
(ε 18000) (MeOH).

O-Sulfate: **Tridentatol E**
C₁₁H₁₃NO₄S₃ 319.426
Isol. from *Tridentata marginata*.
Amorph. solid (as Na salt). λ_{max} 326
(log ε 4.41) (MeOH) (Na salt).

Z-Isomer: **Tridentatol B**
[185548-49-6]
C₁₁H₁₃NOS₂ 239.362
Isol. from *Tridentata marginata*.

Amorph. solid. λ_{\max} 332 (ϵ 11000) (MeOH).

Z-Isomer, O-sulfate: Tridentatol F

$C_{11}H_{13}NO_4S_3$ 319.426

Isol. from *Tridentata marginata*.


Amorph. solid (as Na salt). λ_{\max} 325 (log ϵ 3.78) (MeOH) (Na salt).

Lindquist, N. *et al.*, *Tet. Lett.*, 1996, **37**, 9131-9134 (*Tridentatols A,B*)

Lindquist, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 681-684 (*Tridentatols E,F*)

Tridentatol C T-508

4-[2-(Methylthio)-5-thiazolyl]phenol, 9CI. 2-(4-Hydroxyphenyl)-5-(methylthio)thiazole [185548-50-9]



$C_{10}H_9NOS_2$ 223.319

Metab. from the marine hydroid *Tridentata marginata*. Plates. λ_{\max} 313 (ϵ 3000) (MeOH).

O-Sulfate: Tridentatol H

[439255-40-0]

$C_{10}H_9NO_4S_3$ 303.383


Isol. from *Tridentata marginata*.

Amorph. powder (as Na salt).

Lindquist, N. *et al.*, *Tet. Lett.*, 1996, **37**, 9131-9134 (*isol, uv, ir, pmr, cryst struct*)

Jayatilake, G.S. *et al.*, *Org. Lett.*, 1999, **1**, 661-662 (*synth*)

Lindquist, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 681-684 (*Tridentatol H*)

Tridentatol D T-509

$C_{10}H_{11}NOS_2$ 225.335

Isol. from the marine hydroid *Tridentata marginata*. Amorph. solid. λ_{\max} 342 (log ϵ 4.35) (MeOH).

O-Sulfate: Tridentatol G

$C_{10}H_{11}NO_4S_3$ 305.399


Isol. from *Tridentata marginata*.

Amorph. powder (as Na salt). λ_{\max} 289 (sh); 339 (log ϵ 4.32) (MeOH) (Na salt).

Lindquist, N. *et al.*, *J. Nat. Prod.*, 2002, **65**, 681-684 (*isol, pmr, cmr*)

5,6,11-Trideoxytetradotoxin T-510

[172228-91-0]



$C_{11}H_{17}N_3O_5$ 271.272

Slightly misleading name. In Tetrodotoxin, T-298 the 5-OH is present as an internal acetal. Isol. from the fish *Fugu poecilonotus*. $[\alpha]_D^{23}$ -17.4 (c, 0.5 in AcOH aq.).

► Toxic.

N¹,6-Dihydroxy: 1-Hydroxy-5,11-dideoxytetradotoxin

[146689-16-9]

$C_{11}H_{17}N_3O_7$ 303.271

Isol. from the newt *Taricha granulosa* and *Taricha torosa*. Amorph. powder.

$[\alpha]_D$ +14.5 (c, 0.3 in AcOH aq.).

Misleading synonym. λ_{\max} 225 (H₂O) (Berdy).

► Neurotoxin.

6,11-Dihydroxy: 5-Deoxytetradotoxin

$C_{11}H_{17}N_3O_7$ 303.271

Isol. from *Fugu poecilonotus*. Amorph. solid.

4-Epimer: 5,6,11-Trideoxy-4-epitetrodotoxin

[172339-07-0]

$C_{11}H_{17}N_3O_5$ 271.272

Isol. from *Fugu poecilonotus*. $[\alpha]_D^{23}$ -60.2 (c, 0.65 in AcOH aq.).

► Toxic.


Kotaki, Y. *et al.*, *J.A.C.S.*, 1993, **115**, 827-830 (*1-Hydroxy-5,11-dideoxytetradotoxin*)

Yotsu-Yamashita, M. *et al.*, *Tet. Lett.*, 1995, **36**, 9329-9332 (*isol, struct*)

Yotsu-Yamashita, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 961-963 (*5-Deoxytetradotoxin*)

Tridictyophylline T-511

14-Hydroxyisostephodeline. Alkaloid TGL3 [52038-19-4]



$C_{21}H_{27}NO_6$ 389.447

Alkaloid from *Pachygone dasycarpa*, *Triclisia dictyophylla* and *Triclisia gillettii* (Menispermaceae). Long feathery needles or prisms (MeOH). Mp 204° (softens at 190°). $[\alpha]_D^{28}$ +159 (c, 0.16 in CHCl₃). λ_{\max} 204 (log ϵ 4.6); 226 (sh) (log ϵ 4.04); 279 (log ϵ 3.96) (MeOH).


Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 1 (*isol*)

Spiff, A.L. *et al.*, *J. Nat. Prod.*, 1981, **44**, 160 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Guinaudeau, H. *et al.*, *Phytochemistry*, 1998, **49**, 2561-2563 (*isol, uv, pmr, cmr, ms, cd*)

Trididemnic acid A T-512

4,8-Dihydroxy-5-(4-hydroxybenzoyl)-2-quinolinecarboxylic acid, 9CI [142808-50-2]



$C_{17}H_{11}NO_6$ 325.277

Alkaloid from the marine ascidian *Trididemnum* sp. Pale yellow solid.

3'-Hydroxy: **Trididemnic acid B**. 5-(3,4-Dihydroxybenzoyl)-4,8-dihydroxy-2-quinolinecarboxylic acid, 9CI [142808-51-3]

$C_{17}H_{11}NO_7$ 341.276


Alkaloid from *Trididemnum* sp.

Dilip de Silva, E. *et al.*, *Tet. Lett.*, 1992, **33**, 2917-2920 (*isol, pmr, cmr, struct*)

Trigamine T-513

[59272-74-1]

[62278-79-9]



$C_{19}H_{27}NO_3$ 317.427

Alkaloid from *Merendera trigina*. Mp 169-170°. $[\alpha]_D$ -7 (c, 1.0 in CHCl₃).

N-Oxide: Trigamine N-oxide

[97763-03-6]

$C_{19}H_{27}NO_4$ 333.427

Alkaloid from *Merendera jolantae* (Liliaceae). Mp 201-202°. $[\alpha]_D$ +10.

Yusupov, M.K. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 824-825 (*isol, uv, ir, pmr, ms*)


Timbekov, E.K. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 1-9 (*ms*)

Chommadov, B. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 395-397 (*N-oxide*)

Trigilletimine T-514

Alkaloid TGS-1

[52038-21-8]



$C_{33}H_{30}N_2O_5$ 558.632

Alkaloid from the stems and roots of *Triclisia gillettii* and *Triclisia patens*, and from *Triclisia dictyophylla* (whole plant) (Menispermaceae). Needles (EtOH). Mp 284° dec. $[\alpha]_D^{25}$ -285.7 (c, 0.7 in CH₂Cl₂).

Tackie, A.N. *et al.*, *J. Nat. Prod.*, 1974, **37**, 1 (isol)


Dwuma-Badu, D. *et al.*, *Experientia*, 1975, **31**, 1251 (uv, pmr, ms, struct)

Spiff, A.I. *et al.*, *J. Nat. Prod.*, 1981, **44**, 160 (isol, uv, ir, ms)

Triglochinin

T-515

[28876-11-1]



C₁₄H₁₇NO₁₀ 359.289

Constit. of *Triglochin maritimum*, *Alocasia macrorrhiza* and *Arum maculatum*.

Hygroscopic amorph. powder. $[\alpha]_D^{16.5}$ +5.5 (c, 0.2 in MeOH).

2E-Isomer: Isotriglochinin

[56816-21-8]

C₁₄H₁₇NO₁₀ 359.289

Constit. of *Alocasia macrorrhiza* and *Arum maculatum*. Poss. artifact.

2E-Isomer, 8-Me ester: [39026-83-0]

C₁₅H₁₉NO₁₀ 373.316

Constit. of *Thalictrum aquilegifolium*.

Eyjölfsson, R. *et al.*, *Phytochemistry*, 1970, **9**, 845

Ettlinger, M. *et al.*, *Chem. Comm.*, 1972, 572 (struct)

Sharples, D. *et al.*, *Phytochemistry*, 1972, **11**, 2999-3002 (Me ester)

Nahrstedt, A. *et al.*, *Phytochemistry*, 1975, **14**, 1339-1340; 1870-1871 (isol)


Jaroszewski, J.W. *et al.*, *Phytochemistry*, 1981, **20**, 819 (biosynth)

Seigler, D.S. *et al.*, *Phytochemistry*, 2005, **66**, 1567-1580 (biosynth, occur)

1,2,3-Trihydroxyacridone

T-516

1,2,3-Trihydroxy-9(10H)-acridinone, 9CI [263007-73-4]



C₁₃H₉NO₄ 243.218

Isol. from root bark of *Severinia buxifolia*.

2,3-Di-Me ether: Xanthoxoline. 1-Hydroxy-2,3-dimethoxyacridone

[17014-43-6]

C₁₅H₁₃NO₄ 271.272

Alkaloid from leaves of *Evodia xanthoxyloides* (Rutaceae). Mp 256-257°.

2,3-Di-Me ether, O-Ac:

C₁₇H₁₅NO₅ 313.309

Mp 238-241°.

2,3-Di-Me ether, N-Me: Arborinine

[5489-57-6]

C₁₆H₁₅NO₄ 285.299

Alkaloid from *Glycosmia arborea*, *Fagaria leprieurii*, *Evodia xanthoxyloides*,

Ruta graveolens (rue), and other spp. in the Rutaceae. Antihistamine, antiinflammatory, spasmolytic agent. Cryst. (EtOH). Mp 175-176°. Log P 1.11 (calc).

2,3-Di-Me ether, N-Me, O-Ac:

C₁₈H₁₇NO₅ 327.336

Mp 215-216°.

2,3-Di-Me ether, N-(acetyloxymethyl):**Toddaliopsin C**

[865103-33-9]

C₁₈H₁₇NO₆ 343.335

Alkaloid from the leaves of *Toddaliopsis bremekampii*. Yellow glass.

Tri-Me ether, 1,2,3-Trimethoxyacridone.**Toddaliopsin A**

[33130-09-5]

C₁₆H₁₅NO₄ 285.299

Alkaloid from the leaves of *Toddaliopsis bremekampii*. Yellow glass.

Tri-Me ether, N-Me: 1,2,3-Trimethoxy-**10-methylacridone**

[13082-16-1]

C₁₇H₁₇NO₄ 299.326

Alkaloid from *Evodia alata*, *Vepris bilocularis*, *Melicope leratii* and *Almeidea rubra*. Mp 169-171°.

Tri-Me ether, N-(methoxymethyl): Toddaliopsin D

[865103-34-0]

C₁₈H₁₉NO₅ 329.352

Alkaloid from the leaves of *Toddaliopsis bremekampii*.

Tri-Me ether, N-(acetyloxymethyl): Toddaliopsin B

[865103-32-8]

C₁₉H₁₉NO₆ 357.362

Alkaloid from the leaves of *Toddaliopsis bremekampii*. Yellow glass.

3-(3-Methyl-2-butenyl) ether, 2-Me ether, N-Me: 1-Hydroxy-2-methoxy-10-**methyl-3-prenyloxyacridone. Evoprenine**

[16584-45-5]

C₂₀H₂₁NO₄ 339.39

Alkaloid from the bark of *Evodia alata*. Yellow needles (EtOH). Mp 143°.

2,3-Methylene, 1-Me ether: 11-Methoxy-**1,3-dioxolo[4,5-b]acridin-10-one. 1-Methoxy-2,3-methylenedioxyacridone.****Evoxanthidine**

[668-35-9]

C₁₅H₁₁NO₄ 269.256

Alkaloid from *Evodia xanthoxyloides* (Rutaceae). Yellow needles (EtOH). Mp 310-312° dec.

2,3-Methylene, 1-Me ether, N-Me: 1-**Methoxy-10-methyl-2,3-methylenedioxyacridone. Evoxanthine**

[477-82-7]

C₁₆H₁₃NO₄ 283.283

Alkaloid from *Evodia xanthoxyloides*, *Balfouriodendron riedelianum* and *Oriacia suaveolens* (Rutaceae). Yellow needles (EtOH). Mp 217-218°.

Hughes, G.K. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1949, **A2**, 429-436; 1952, **5**, 401-405

(*Evoxanthine*, *Evoxanthidine*, *Xanthoxoline*, *isol*, *struct*)

Cannon, J.R. *et al.*, *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 406-411 (*Evoxanthine*,

Evoxanthidine, *Xanthoxoline*, *struct*)

Chakravati, D. *et al.*, *J.C.S.*, 1953, 3337-3340 (*Arborinine*, *isol*)

Gell, R.J. *et al.*, *Aust. J. Chem.*, 1955, **8**, 114-120 (*Evodia elata constis*, *struct*)

Pakrashi, S.C. *et al.*, *Chem. Ind. (London)*, 1961, 464-465 (*Arborinine*, *pmr*)

Banerjee, S.K. *et al.*, *Tetrahedron*, 1961, **16**, 251-254 (*Arborinine*)

Bowie, J. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1179-1193 (ms)

Diment, J.A. *et al.*, *Aust. J. Chem.*, 1967, **20**, 1719-1727 (*Evoprenine*)

Govindachari, T.R. *et al.*, *Tetrahedron*, 1967, **23**, 1827-1829 (*Evoxanthidine*, *synth*, *ir*, *uv*)

Bergenthal, D. *et al.*, *Phytochemistry*, 1979, **18**, 161-163 (*cmr*)

Santos, C.S. *et al.*, *J. Braz. Chem. Soc.*, 1988, **9**, 39-42 (*Almeidea rubra constit*)

Coppola, G.M. *et al.*, *J. Het. Chem.*, 1989, **26**, 957-964 (*Evoxanthine*, *1,2,3-Trimethoxy-10-methylacridone*, *synth*, *ir*, *pmr*)

Rasoanaivo, P. *et al.*, *Fitoterapia*, 1999, **70**, 625-627 (*Evoxanthine*, *cmr*)

Wu, T.-S. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1040-1043 (*Severinia buxifolia constit*)

Naidoo, D. *et al.*, *Phytochemistry*, 2005, **66**, 1724-1728 (*Toddaliopsins A-D*)

1,3,4-Trihydroxyacridone

T-517

1,3,4-Trihydroxy-9(10H)-acridinone, 9CI C₁₃H₉NO₄ 243.218

3,4-Di-Me ether, N-Me: 1-Hydroxy-3,4-**dimethoxy-10-methylacridone**

[96935-29-4]

C₁₆H₁₅NO₄ 285.299

Alkaloid from the wood and bark of *Sarcomelicope leiocarpa* (Rutaceae). Yellow prisms (CH₂Cl₂). Mp 129-131°.

Tri-Me ether, N-Me: 1,3,4-Trimethoxy-**10-methylacridone**

[66642-48-6]

C₁₇H₁₇NO₄ 299.326

Alkaloid from leaves of *Teclea boivini-ana* and bark of *Acronychia baueri*. Pale yellow needles (CHCl₃/Et₂O). Mp 137°.

3-(3,7-Dimethyl-2,6-octadienyl) ether, 4-**Me ether, N-Me: 1-Hydroxy-3-geranyloxy-4-methoxy-10-methylacridone**

[96935-30-7]

C₂₅H₂₉NO₄ 407.508

Alkaloid from the wood and bark of *Sarcomelicope leiocarpa* (Rutaceae).

Yellow prisms (CHCl₃). Mp 114°.

Vaquette, J. *et al.*, *Planta Med.*, 1978, **33**, 78

(*1,3,4-Trimethoxy-10-methylacridone*)

Funayama, S. *et al.*, *J. Nat. Prod.*, 1984, **47**,

285 (*1,3,4-Trimethoxy-10-methylacridone*)

Baudouin, G. *et al.*, *J. Nat. Prod.*, 1985, **48**,

260 (*isol*, *uv*, *ir*, *pmr*, *ms*, *struct*)

1,3,5-Trihydroxyacridone

T-518

1,3,5-Trihydroxy-9(10H)-acridinone, 9CI [85990-00-7]

C₁₃H₉NO₄ 243.218

Bright yellow prisms (Me₂CO aq.). Mp >320°.

N-Me: 1,3,5-Trihydroxy-10-methylacri-**done**

[679835-33-7]

C₁₄H₁₁NO₄ 257.245

Alkaloid from roots of Yalaha [several hybrid seedlings resulting from a cross of Duncan grapefruit (*Citrus paradisi*) x Dancy tangerine (*Citrus tangerina*)]. Yellow oil.

3-Me ether, N-Me: 1,5-Dihydroxy-3-methoxy-10-methylacridone. Citrusamine

[108598-30-7]

C₁₅H₁₃NO₄ 271.272

Alkaloid from the leaves of *Atalantia ceylanica* and from roots of *Citrus tamurana* and *Citrus natsudaidai* (Rutaceae). Orange-yellow needles (CHCl₃/petrol), yellow prisms (Me₂CO). Mp 140-141° Mp 243-246°. Large discrepancy in Mp between the two isolates may indicate an incorrect identity.

5-Me ether: 1,3-Dihydroxy-5-methoxyacridone

[63110-81-6]

C₁₄H₁₁NO₄ 257.245

Bright yellow needles (EtOH aq.). Mp 278-280°.

5-Me ether, N-Me: 1,3-Dihydroxy-5-methoxy-10-methylacridone. Yukodine

[145940-33-6]

C₁₅H₁₃NO₄ 271.272

Alkaloid from the roots of *Citrus yuko* (Rutaceae). Yellow needles. Mp 246-250°.

3,5-Di-Me ether: 1-Hydroxy-3,5-dimethoxyacridone

[85990-01-8]

C₁₅H₁₃NO₄ 271.272

Yellow prisms (DMSO aq.). Mp 220-222° (208-210°).

3,5-Di-Me ether, N-Me: 1-Hydroxy-3,5-dimethoxy-10-methylacridone. Yukodine

[145940-34-7]

C₁₆H₁₅NO₄ 285.299

Alkaloid from roots of Swingle citrus-melo (*Poncirus trifoliata* x *Citrus paradisi*) (Rutaceae). Pale yellow oil.

Tri-Me ether, N-Me: 1,3,5-Trimethoxy-10-methylacridone

[52911-52-1]

C₁₇H₁₇NO₄ 299.326

Alkaloid from *Teclea boiviniana* (Rutaceae). Beige needles (CHCl₃/Et₂O). Mp 141°.

Vaquette, J. et al., *Plant. Med. Phytother.*, 1974, **8**, 57-62 (1,3,5-Trimethoxy-10-methylacridone, struct)

Ramesh, K. et al., *Indian J. Chem., Sect. B*, 1986, **25**, 684-687 (3,5-di-Me ether, synth, pmr)

Ju-Ichi, M. et al., *Heterocycles*, 1987, **26**, 1873-1876 (*Citrusamine*, struct)

Bowen, I.H. et al., *Planta Med.*, 1987, **53**, 73-75 (*Citrusamine*, struct)

Takemura, Y. et al., *Heterocycles*, 1992, **34**, 2123-2130 (*Yukodine*, *Yukodine*)

Takemura, Y. et al., *Chem. Pharm. Bull.*, 1996, **44**, 804 (1,3,5-Trihydroxy-10-methylacridone)

Lowden, C.T. et al., *J. Med. Chem.*, 2003, **46**, 5015-5020 (5-Me ether, 3,5-di-Me ether, synth, pmr)

Herath, H.M.T.B. et al., *J. Het. Chem.*, 2004, **41**, 23-28 (synth, pmr, cmr, ms, N-Me, 5-Me ether, 3,5-di-Me ether)

1,3,6-Trihydroxyacridone T-519

1,3,6-Trihydroxy-9(10H)-acridinone, 9CI

C₁₃H₉NO₄ 243.218

3-Me ether, N-Me: 1,6-Dihydroxy-3-

methoxy-10-methylacridone. **Pumme-****line**

[145940-36-9]

C₁₅H₁₃NO₄ 271.272

Alkaloid from roots of several hybrid seedlings resulting from a cross of Pummelo (*Citrus grandis* cv. May Pummelo × Marsh grapefruit (*Citrus paradisi*) (Rutaceae)). Yellow cubes. Mp 235-240°.

Takemura, Y. et al., *Heterocycles*, 1992, **34**, 2123 (isol, uv, ir, pmr, cmr, ms, struct)

1,3,8-Trihydroxyacridone T-520

1,3,8-Trihydroxy-9(10H)-acridinone, 9CI

[197007-31-1]

C₁₃H₉NO₄ 243.218

Yellow solid (EtOH aq.). Mp >300° dec.

N-Me: 1,3,8-Trihydroxy-10-methylacridone

[151077-56-4]

C₁₄H₁₁NO₄ 257.245

Alkaloid from aerial parts of *Boronia lanceolata* (Rutaceae). Yellow needles (CHCl₃/MeOH). Mp 285-290°.

3-Me ether: 1,8-Dihydroxy-3-methoxyacridone

[156547-74-9]

C₁₄H₁₁NO₄ 257.245

Alkaloid from aerial parts of *Boronia bowmanii* (rutaceae). Yellow needles (CHCl₃). Mp 275°.

3-Me ether, N-Me: 1,8-Dihydroxy-3-methoxy-10-methylacridone. Oligophyllidine

[94443-43-3]

C₁₅H₁₃NO₄ 271.272

Alkaloid from the roots of *Acronychia oligophylebia* (Rutaceae).

Xu, W. et al., *Huaxue Xuebao*, 1984, **42**, 899; *CA*, **102**, 59257s


Ahsan, M. et al., *Phytochemistry*, 1993, **33**, 1507 (1,3,8-Trihydroxy-10-methylacridone)

Ahsan, M. et al., *J. Nat. Prod.*, 1994, **57**, 670 (1,8-Dihydroxy-3-methoxyacridone)

Chen, J.-J. et al., *Tetrahedron*, 1997, **53**, 12717-12728 (synth, pmr, cmr, N-Me, 1,3-di-Me ether)

1,2,3-Trihydroxyaporphine T-521

5,6,6a,7-Tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-1,2,3-triol, 9CI

C₁₇H₁₇NO₃ 283.326**(R)-form**

1,2-Di-Me ether: 3-Hydroxy-1,2-dimethoxyaporphine. **3-Hydroxynuciferine. Lirinine**

[41645-66-3]

C₁₉H₂₁NO₃ 311.38

Alkaloid from the leaves of *Liriodendron tulipifera* (Magnoliaceae). Active

against gram-positive bacteria. Needles (Et₂O). Mp 150-152°. [α]_D²⁸ -83.7 (c, 1.0 in CHCl₃). Struct. revised in 1975.

1,2-Di-Me ether, N-oxide: Lirinine N-oxideC₁₉H₂₁NO₄ 327.379

Alkaloid from the leaves of *Liriodendron tulipifera* (Magnoliaceae).

Amorph. Mp 162-164° dec. λ_{max} 218 (log ε 4.4); 284 (log ε 4.03) (EtOH).

1,2-Di-Me ether, N-de-Me: 3-Hydroxy-1,2-dimethoxynoraporphine. 3-Hydroxynuciferine

[82644-36-8]

C₁₈H₁₉NO₃ 297.353

Alkaloid from the stem bark of *Guatteria melosma*, the bark and leaves of *Polyalthia acuminata*, and the stem and root bark of *Hexalobus crispiflorus* (Annonaceae). Cryst. (Me₂CO). Mp 194-195°. [α]_D²⁰ -68 (c, 0.2 in EtOH). Struct. revised in 1975. An alkaloid from *Monanthes cauliflora* considered to be 9-Hydroxy-1,2-dimethoxynoraporphine was shown to give *O*-Methylirinine on methylation and was therefore prob. identical with 3-Hydroxynuciferine. λ_{max} 219 (log ε 4.39); 240 (sh) (log ε 3.9); 280 (log ε 4.14); 292 (sh) (log ε 4.04) (no solvent reported).

1,3-Di-Me ether: 2-Hydroxy-1,3-dimethoxyaporphine. Liridinine

[58998-24-6]

C₁₉H₂₁NO₃ 311.38

Alkaloid from *Liriodendron tulipifera* (Magnoliaceae). Mp 142-144°. [α]_D²⁰ -38 (c, 0.095 in CHCl₃). λ_{max} 221 (log ε 4.41); 281 (log ε 4.16) (EtOH).

1,3-Di-Me ether, N-de-Me: 2-Hydroxy-1,3-dimethoxynoraporphine. Norliridinine

[83694-77-3]

C₁₈H₁₉NO₃ 297.353

Alkaloid from the bark and leaves of *Polyalthia acuminata* (Annonaceae). Amorph. λ_{max} 230 (sh) (log ε 4.05); 274 (log ε 3.95); 282 (sh) (log ε 3.89); 304 (sh) (log ε 3.38) (no solvent reported).

2,3-Di-Me ether: 1-Hydroxy-2,3-dimethoxyaporphine. N-Methylisopiline

[65953-84-6]

C₁₉H₂₁NO₃ 311.38

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae). Prisms (MeOH). Mp 220-222° dec. [α]_D -56 (c, 0.2 in MeOH).

2,3-Di-Me ether, N-de-Me: 1-Hydroxy-2,3-dimethoxynoraporphine. Isopiline

[65953-83-5]

C₁₈H₁₉NO₃ 297.353

Alkaloid from the trunk bark of *Isolona pilosa*, the stem bark of *Guatteria ouregou* and the bark and leaves of *Polyalthia acuminata* (Annonaceae). Shows antimicrobial activity. Prisms (Et₂O). Mp 153°. [α]_D -55 (c, 1.06 in MeOH).

Tri-Me ether: 1,2,3-Trimethoxyaporphine. 3-Methoxynuciferine. O,N-Dimethylisopiline. O-Methylirinine

[5531-86-2]

C₂₀H₂₃NO₃ 325.407

Alkaloid from the leaves of *Liriodendron tulipifera* and the bark of *Polyalthia acuminata* (Magnoliaceae, Annonaceae). Pale yellow needles (hexane). Mp 105-106°. [α]_D²⁸ -112.15 (c, 0.214 in CHCl₃). λ _{max} 212 (ε 43000); 228 (sh) (ε 25100); 275 (ε 21400) (EtOH).

Tri-Me ether, N-de-Me: 1,2,3-Tri-methoxynoraporphine. O-Methylisopiline. O-Methylnorlirinine

[65953-85-7]

C₁₉H₂₁NO₃ 311.38

Alkaloid from bark of *Liriodendron tulipifera*, bark and leaves of *Polyalthia acuminata*, stem bark of *Guatteria ouregou* and leaves of *Guatteria scandens* (Magnoliaceae, Annonaceae). Shows antimicrobial activity. Non-cryst. Negative opt. rotn. in MeOH. Struct. revised in 1973. λ _{max} 222 (log ε 4.46); 283 (log ε 4.22) (EtOH).

Tri-Me ether, N-de-Me, N-formyl: Formouregine. N-Formyl-O-methylisopiline

[107633-66-9]

C₂₀H₂₁NO₄ 339.39

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae). [α]_D -146 (c, 0.02 in EtOH). λ _{max} 222 (log ε 4.27); 233 (sh) (log ε 4.12); 274 (log ε 4.05) (EtOH).

Tri-Me ether, N-de-Me, N-Ac: Tuliferoline

[60755-82-0]

C₂₁H₂₃NO₄ 353.417

Alkaloid from the heartwood of *Liriodendron tulipifera* (Magnoliaceae). Cryst. (EtOH/hexane). Mp 145-146°. [α]_D²⁵ -330 (c, 0.83 in CHCl₃). λ _{max} 224 (log ε 4.06); 273 (log ε 4.1) (MeOH).

6a,7-Didehydro, 1,2-di-Me ether: 3-Hydroxy-6a,7-dehydronuciferine

[82628-33-9]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the stem bark and root bark of *Hexalobus crispiflorus* (Annonaceae).

6a,7-Didehydro, 1,2-di-Me ether, 3-Ac: Cryst. (MeOH). Mp 134-135°.

6a,7-Didehydro, 2,3-di-Me ether, N-de-Me, N-formyl: 1-De-O-methyldehydroformouregine

[938433-34-2]

C₁₉H₁₇NO₄ 323.348

Alkaloid from *Sabia yunnanensis*. Needles (Me₂CO). Mp 203-205°. λ _{max} 256 ; 279 ; 318 ; 347 ; 365 (MeOH).

6a,7-Didehydro, tri-Me ether, N-de-Me: O-Methyldehydroisopiline

[107633-67-0]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the leaves of *Guatteria ouregou* (Annonaceae). Not obt. pure. λ _{max} 214 (log ε 4.48); 260 (log ε 4.69); 327 (log ε 3.96) (EtOH).

6a,7-Didehydro, tri-Me ether, N-de-Me, N-formyl: Dehydroformouregine

[107633-69-2]

C₂₀H₁₉NO₄ 337.374

Alkaloid from the leaves of *Guatteria*

ouregou (Annonaceae). Cryst.

(MeOH). Mp 120-122°. λ _{max} 221 (log ε 3.92); 251 (sh) (log ε 4.29); 263 (log ε 4.32); 310 (log ε 3.67) (EtOH).

(S)-form

2,3-Di-Me ether: (+)-N-Methylisopiline

[119060-85-4]

C₁₉H₂₁NO₃ 311.38

Trace alkaloid from roots and stems of *Neostenanthera gabonensis* (Annonaceae). Gum. [α]_D²⁰ +53 (c, 0.05 in MeOH) (semisynthetic).

2,3-Di-Me ether, N-de-Me: (+)-Isopiline

[119060-84-3]

C₁₈H₁₉NO₃ 297.353

Alkaloid from the roots and stems of *Neostenanthera gabonensis* (Annonaceae). Oil. [α]_D²⁰ +55 (c, 0.15 in MeOH). λ _{max} 273 ; 291 (sh) ; 311 (MeOH). λ _{max} 256 (sh) ; 275 ; 341 (MeOH/NaOH).

Ziayev, R. et al., *Khim. Prir. Soedin.*, 1973,

9, 67-70; 505-506; *Chem. Nat. Compd.*

(Engl. Transl.), 59-61; 475-476

(Lirinine, Lirinine N-oxide, O-Methylirinine)

Abdusamatov, A. et al., *Khim. Prir. Soedin.*,

1975, 11, 812-813; *Chem. Nat. Compd.*

(Engl. Transl.), 1975, 11, 829-830

(Liridinine)

Hufford, C.D. et al., *Phytochemistry*, 1976, 15,

1169-1171 (*Tuliferoline*)

Hocquemiller, R. et al., *C. R. Hebd. Seances*

Acad. Sci. Ser. C, 1977, 285, 447-450 (*O-**Methylirinine, Isopiline*)

Wiriyachitra, P. et al., *J.O.C.*, 1977, 42,

2274-2277 (*Lirinine, O-Methylirinine,**synth*)

Chen, C.-L. et al., *Phytochemistry*, 1978,

17, 779-782 (*O-Methylirinine, isol,**struct*)

Achenbach, H. et al., *Annalen*, 1982, 1132-

1141; 1623-1633 (*3-Hydroxynornuciferine, 3-**Hydroxy-6a,7-dehydronuciferine*)

Zarga, M.H.A. et al., *J. Nat. Prod.*, 1982, 45,

471-475 (*O-Methylirinine, 3-**Hydroxynornuciferine, Isopiline,**Norlirinidine*)

Abd-El Atti, S. et al., *J. Nat. Prod.*, 1982, 45,

476-480 (*3-Hydroxynornuciferine*)

Hoquemiller, R. et al., *J. Nat. Prod.*, 1983, 46,

335-341 (*O-Methylisopiline*)

Leboeuf, M. et al., *Planta Med.*, 1983, 48, 234-

245 (*Isopiline, isol, cmr*)

Villar, A. et al., *Farm. Tijdschr. Belg.*, 1984, 61,

300 (*Isopiline, O-Methylisopiline, activity*)

Cortes, D. et al., *J. Nat. Prod.*, 1986, 49, 878-

884 (*Guatteria ouregou constits*)

Villar, A. et al., *Planta Med.*, 1986, 52, 556-557

(*3-Hydroxynuciferine, activity*)


Renner, C. et al., *J. Nat. Prod.*, 1988, 51, 973-

976 (*(+)-Isopiline, (+)-N-methylisopiline*)

Deng, Y. et al., *Nat. Prod. Res.*, 2007, 21, 28-32

(*1-De-O-methyldehydroformouregine*)**1,2,8-Trihydroxyaporphine**

T-522

C₁₇H₁₇NO₃ 283.326**(R)-form**

1,2-Methylene, 8-Me ether: 8-Methoxy-1,2-methylenedioxyaporphine. Stephanine

[517-63-5]

C₁₉H₁₉NO₃ 309.364

Alkaloid from *Stephania capitata* and *Stephania japonica* (Menispermaceae). Mp 155-157°. [α]_D -92.5 (CHCl₃).

1,2-Methylene, 8-Me ether, N-Me: Mp 211-215° (as iodide).

6a,7-Didehydro, 1,2-methylene, 8-Me ether: Dehydrostephanine

[76907-76-1]

C₁₉H₁₇NO₃ 307.348

Alkaloid from the root tubers of *Stephania kwangsiensis* and *Stephania micrantha*, and from the seeds of *Stephania cepharantha* (Menispermaceae). Mp 161-163°.

7R-Hydroxy, 1,2-methylene, 8-Me ether: 7-Hydroxy-8-methoxy-1,2-methylenedioxyaporphine. Ayuthianine

[82413-18-1]

C₁₉H₁₉NO₄ 325.363

Alkaloid from the tuberous roots of *Stephania venosa* (Menispermaceae). Amorph.

(±)-form

1,2-Methylene, 8-Me ether: Synthetic.

Prisms (Me₂CO). Mp 136-137° (131-133°).

1,2-Methylene, 8-Me ether, hydrochloride: Needles (MeOH/Et₂O). Mp 269-270° dec.

1,2-Methylene, 8-Me ether, N-Me:

Needles (as iodide). Mp 212-214°

(iodide).

Shirai, H. et al., *Yakugaku Zasshi*, 1956, 76,

1287; *CA*, 51, 9646f (*Stephanine*)

Tomita, M. et al., *Yakugaku Zasshi*, 1957, 77,

290; 1963, 83, 996; *CA*, 51, 11362h; 60,
4202a (*Stephanine, isol, synth, struct*)

Hey, D.H. et al., *J.C.S.*, 1958, 1876 (*synth*)

Watanabe, Y. et al., *Phytochemistry*, 1975, 14,

2522 (*synth, uv, ir, pmr, ms*)

Min, Z.-D. et al., *Yaoxue Xuebao*, 1980, 15,

532; 1981, 16, 557; *CA*, 94, 117773m; 97,
3595m (*Dehydrostephanine*)

Kunitomo, J. et al., *Yakugaku Zasshi*, 1981,

101, 951; *CA*, 96, 31640v (*Dehydrostephanine*)

Kunitomo, J. et al., *Heterocycles*, 1982, 19,

1883 (*Stephanine, cd, ord, cryst struct, abs**config*)

Guinaudeau, H. et al., *J. Nat. Prod.*, 1982, 45,

355 (*Ayuthianine, isol, uv, pmr, ms, cd, struct*)


Bartley, J.P. et al., *Phytochemistry*, 1994, 36,

1327-1331 (*Stephanine, Ayuthianine, isol,**pmr, cmr*)

Blanchfield, J.T. et al., *Phytochemistry*, 2003,

63, 711-720 (*Stephanine, Ayuthianine, pmr,**cmr*)**1,2,9-Trihydroxyaporphine**

T-523



C₁₇H₁₇NO₃ 283.326**(S)-form**

2,9-Di-Me ether: **1-Hydroxy-2,9-dimethoxyaporphine. Orientinine** [92008-87-2]
C₁₉H₂₁NO₃ 311.38
Alkaloid from *Papaver orientale* (Papaveraceae). Amorph. [α]_D +62 (c, 0.18 in MeOH). λ_{max} 275 ; 315 (sh) (no solvent reported).

2,9-Di-Me ether, N-de-Me: **1-Hydroxy-2,9-dimethoxynoraporphine. 2,9-Dimethoxy-1-hydroxynoraporphine (incorr.)**. N-Nororientinine [126622-47-7]
C₁₈H₁₉NO₃ 297.353
Alkaloid from trunk wood of *Ocotea caesia* (Lauraceae). Amorph. [α]_D²⁵ +61 (c, 0.11 in MeOH). λ_{max} 245 (log ε 4.23); 289 (log ε 4.36); 326 (log ε 4.38) (MeOH).


Tri-Me ether: **1,2,9-Trimethoxyaporphine. Orientine†** [90275-85-7]
C₂₀H₂₃NO₃ 325.407
Alkaloid from *Papaver orientale* epigeal parts (Papaveraceae). [α]_D +70 (c, 0.16 in MeOH). λ_{max} 278 (log ε 4.25); 310 (sh) (log ε 3.21) (no solvent reported).

Israilov, I.A. et al., *Khim. Prir. Soedin.*, 1984, **20**, 81; 258; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 76-78; 243-244 (*Orientine, Orientinine*)

Yariwake, J.H. et al., *Phytochemistry*, 1989, **28**, 3577-3578 (*1-Hydroxy-2,9-dimethoxynoraporphine*)

1,2,10-Trihydroxyaporphine T-524

5,6a,7-Tetrahydro-6-methyl-4H-dibenzo[de,g]quinoline-1,2,10-triol, 9CI

C₁₇H₁₇NO₃ 283.326**(R)-form**

1-Me ether, N-de-Me: **Apocrotonosine** C₁₇H₁₇NO₃ 283.326
Mp 218-220° (hydrochloride hemihydrate). [α]_D -148 (c, 0.42 in H₂O). Rearrangement prod. of Crotonosine, in P-643.

2-Me ether: **1,10-Dihydroxy-2-methoxyaporphine. Apoglaziovine. N-Methylsparsiflorine** [2128-77-0]
C₁₈H₁₉NO₃ 297.353
Alkaloid from the leaves of *Ocotea glaziovii* (Lauraceae). Hypotensive agent. Cryst. + 1H₂O. Mp 149-152° dec. [α]_D²⁶ -35 (c, 0.2 in CHCl₃). This product must be partly racemic.

2-Me ether, N-de-Me, tri-Ac: Mp 196-

197°.

1,2-Di-Me ether: **10-Hydroxy-1,2-dimethoxyaporphine. Nuciferoline** [1862-49-3]
C₁₉H₂₁NO₃ 311.38
Alkaloid from *Papaver causicum* (Papaveraceae). Also obt. by the dienone-phenol rearr. of Pronuciferine, P-643. Cryst. (EtOH). Mp 227-229°. [α]_D²⁰ -157 (c, 0.180 in EtOH).

1,2-Di-Me ether, N-Me: **Amurenine** [169626-39-5]
[169211-34-1]
C₂₀H₂₄NO₃[⊕] 326.414
Quaternary alkaloid from young shoots of *Berberis amurensis*. Cryst. (Me₂CO) (as chloride). Mp 194-195° (chloride). [α]_D -164 (c, 0.03 in MeOH) (chloride).

1,2-Di-Me ether, N-de-Me: **10-Hydroxy-1,2-dimethoxynoraporphine. Tuduranine** [517-97-5]
C₁₈H₁₉NO₃ 297.353
Alkaloid from *Sinomenium acutum*, the root tubers of *Stephania rotunda* and the bark and leaves of *Polalthia acuminata* (Menispermaceae, Annonaceae). Poorly cryst. Mp 105-125° Mp 204°.

1,2-Di-Me ether, N-de-Me, hydrochloride: Prisms (H₂O). Mp 286° dec. [α]_D^{14,5} -148 (MeOH aq.).

1,2-Di-Me ether, N-de-Me, N,O-di-Ac: Prisms (MeOH or EtOH). Mp 170°. [α]_D¹⁴ -321.71 (MeOH).

1,10-Di-Me ether: **2-Hydroxy-1,10-dimethoxyaporphine. Phoebe base II** C₁₉H₂₁NO₃ 311.38
Alkaloid from *Phoebe* sp. (Lauraceae). Mp 198-200°. [α]_D +160 (CHCl₃).

2,10-Di-Me ether: **1-Hydroxy-2,10-dimethoxyaporphine. Pulchine. N-Methylzenkerine** [114579-00-9]
C₁₉H₂₁NO₃ 311.38
Constit. of *Ocotea caesia* and *Ocotea pulchella* (Lauraceae). Cryst. (MeOH). Mp 58-60°. [α]_D²⁵ -128 (c, 0.19 in MeOH).

2,10-Di-Me ether, N-Me: **N,N-Dimethylzenkerine. N-Methylpulchine** [866269-27-4]
C₂₀H₂₄NO₃[⊕] 326.415
Quaternary alkaloid from *Monodora grandidieri* and *Monodora junodii*.

2,10-Di-Me ether, N-de-Me: **1-Hydroxy-2,10-dimethoxynoraporphine. Zenkerine†. 10-Methoxycaaverine** [65953-82-4]
C₁₈H₁₉NO₃ 297.353
Alkaloid from the trunk bark of *Isolona pilosa* and from *Isolona zenkeri* (Annonaceae). Cryst. (MeOH). Mp 73-76°. [α]_D²⁵ -103 (c, 0.17 in MeOH).

Tri-Me ether: **1,2,10-Trimethoxyaporphine. N,O,O-Trimethylsparsiflorine** C₂₀H₂₃NO₃ 325.407
Alkaloid from the rhizomes of *Thalictrum foliolosum* (Ranunculaceae).

Tri-Me ether, N-Me:

Cryst. (MeOH) (as iodide). Mp 210-212° dec. (iodide).

Tri-Me ether, N-de-Me, N-Ac: Prisms (EtOH). Mp 189°. [α]_D¹⁸ -400.17 (CHCl₃).

(S)-form

2-Me ether: **(+)-Apoglaziovine** [18058-59-8]
C₁₈H₁₉NO₃ 297.353

Alkaloid from *Ocotea variabilis* (Lauraceae). Mp 249-252° dec. [α]_D²⁵ +165 (CHCl₃).

2-Me ether, N-de-Me: **1,10-Dihydroxy-2-methoxynoraporphine. Sparsiflorine** [2128-61-2]
C₁₇H₁₇NO₃ 283.326
Alkaloid from the leaves of *Croton sparsiflorus* (Euphorbiaceae). Needles (EtOH). Mp 228° dec. [α]_D³⁰ +43 (H₂O) (hydrochloride).

1,2-Di-Me ether: **(+)-Nuciferoline** [6874-94-8]
C₁₉H₂₁NO₃ 311.38

Alkaloid from *Papaver causicum* (Papaveraceae). Also obt. by the dienone-phenol rearr. of Pronuciferine (S)-form, P-643. Prisms (EtOH). Mp 225-227°. [α]_D²¹ +160 (c, 0.31 in EtOH).

1,2-Di-Me ether, hydrobromide: [20827-79-6]
Needles (MeOH/Et₂O). Mp 222-226° dec.

1,2-Di-Me ether, N-Me: Needles (MeOH/Et₂O). Mp 220-222° dec.

1,2-Di-Me ether, N-de-Me, N-methoxycarbonyl: **Romucosine C** [356048-25-4]
C₂₀H₂₁NO₅ 355.39
Alkaloid from *Rollinia mucosa* (biri-ba). Amorph. powder. [α]_D²⁵ +140 (c, 0.05 in CHCl₃). λ_{max} 265 (log ε 4.25); 280 (log ε 4.02); 305 (log ε 4.35) (MeOH).

Tri-Me ether: [82444-06-2]
Alkaloid from the rhizomes of *Thalictrum foliolosum* (Ranunculaceae). Mp 124-126°.

Tri-Me ether, N-de-Me, N-methoxycarbonyl: **Romucosine D** [356048-26-5]
C₂₁H₂₃NO₅ 369.416
Alkaloid from *Rollinia mucosa* (biri-ba). Amorph. powder. [α]_D²⁵ +145 (c, 0.05 in CHCl₃). λ_{max} 263 (log ε 4.33); 275 (log ε 4.22); 307 (log ε 3.75) (MeOH).

(±)-form

2-Me ether: **(±)-Apoglaziovine** [56261-23-5]
C₁₈H₁₉NO₃ 297.353

Alkaloid from the leaves of *Ocotea glaziovii* (Lauraceae).

2-Me ether, N-de-Me: [33074-89-4 (HCl salt)] Synthetic. Pale brown cryst. (MeOH) (as hydrochloride). Mp 300° (hydrochloride). CAS no. refers to hydrochloride.


1,2-Di-Me ether: [28277-74-9]
Synthetic. Plates (C₆H₆). Mp 211-213°.

1,2-Di-Me ether, N-de-Me: Synthetic.
Cryst. (MeOH) (as hydrochloride). Mp
282° (hydrochloride).

Tri-Me ether: [28277-71-6]
C₂₀H₂₃NO₃ 325.407
Synthetic. Oil.

- Goto, K. *et al.*, *Annalen*, 1936, **521**, 175-184
(*Tuduranine, isol, uv*)
Bernauer, K. *et al.*, *Helv. Chim. Acta*, 1964, **47**,
2122-2129 (*Nuciferoline, synth*)
Gilbert, B. *et al.*, *J.A.C.S.*, 1964, **86**, 694-696
(*Pulchine, isol, uv, ir, pmr, struct, Tuduranine,*
pmr)
Chatterjee, A. *et al.*, *Tet. Lett.*, 1965, 1539-
1544 (*uv, pmr, struct, Sparsiflorine*)
Haynes, L.J. *et al.*, *J.C.S. (C)*, 1966, 1676-1679
(*Apocrotosine, synth*)
Tomita, M. *et al.*, *Yakugaku Zasshi*, 1966, **86**,
871-873 (*Tuduranine*)
Narayanaswami, S. *et al.*, *Indian J. Chem.*,
1969, **7**, 945-947; 1971, **9**, 509-511
(*Tuduranine, Sparsiflorine, Apoglaziovine,*
synth)
Ishiwata, S. *et al.*, *Chem. Pharm. Bull.*, 1970,
18, 1219-1223 ((±)-form, synth)
Raffauf, R. *et al.*, *Handb. Alkaloids Alkaloid-*
Containing Plants, Wiley, N.Y., 1970,
(*Phoebe base II*)
Bhakuni, D.S. *et al.*, *Phytochemistry*, 1970, **9**,
2573-2580; 1974, **13**, 2767-2769 (*isol, uv, ir,*
pmr, biosynth, Sparsiflorine)
Cava, M.P. *et al.*, *Tet. Lett.*, 1972, 4647-4649
(*isol, (+)-Apoglaziovine*)
Bhakuni, D.S. *et al.*, *Tetrahedron*, 1972, **28**,
4579-4582 (*Sparsiflorine, abs config*)
Casagrande, C. *et al.*, *Farmaco, Ed. Sci.*, 1975,
30, 479; *CA*, **83**, 93873p ((±)-
Apoglaziovine)
Cava, M.P. *et al.*, *J.O.C.*, 1975, **40**, 1553-1556
(*Nuciferoline, synth*)
Hoquemiller, R. *et al.*, *C. R. Hebd. Seances*
Acad. Sci. Ser. C, 1977, **285**, 447
(*Zenkerine*)
Guinaudeau, H. *et al.*, *J. Nat. Prod.*, 1979, **42**,
325-360 (*rev, cmr, Apoglaziovine*)
Hara, H. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**,
1083 (*synth*)
Bhakuni, D.S. *et al.*, *J. Nat. Prod.*, 1982, **45**,
252-255 (*S-form, isol, uv, ir, pmr, ms*)
Zarga, M.H.A. *et al.*, *J. Nat. Prod.*, 1982, **45**,
471-475 (*Tuduranine, isol*)
Cassels, B.K. *et al.*, *J. Nat. Prod.*, 1987, **50**,
297-300 (*cmr, pmr, Apoglaziovine, Zenkerine,*
Nuciferoline)
Vilegas, J.H.Y. *et al.*, *Phytochemistry*, 1989, **28**,
3577-3578 (*Zenkerine, Pulchine*)
Yusupov, M.M. *et al.*, *Khim. Prir. Soedin.*,
1993, **29**, 401-404; *Chem. Nat. Compd.*
(*Engl. Transl.*), 1993, **29**, 338-340
(*Amurenine*)
Kuo, R.Y. *et al.*, *Phytochemistry*, 2001, **57**, 421-
425 (*Romucosines*)
Nishiyama, Y. *et al.*, *Nat. Med. (Tokyo)*, 2004,
58, 303-306; *CA*, **143**, 363319h (*N,N-*
Dimethylzenkerine)

1,2,11-Trihydroxyaporphine T-525
5,6,6a,7-Tetrahydro-6-methyl-4H-diben-
zo[de,g]quinoline-1,2,11-triol, 9CI



C₁₇H₁₇NO₃ 283.326

► CE1017000

(S)-form

2-Me ether: 1,11-Dihydroxy-2-methoxya-
porphine. **Isothebaidine**
[68156-56-9]
C₁₈H₁₉NO₃ 297.353
Alkaloid from the aerial parts of
flowering *Papaver orientale* (Papaver-
aceae). Mp 236-237° dec. [α]_D +321 (c,
0.5 in MeOH).

1,2-Di-Me ether: **11-Hydroxy-1,2-di-**
methoxyaporphine
[95652-98-5]
C₁₉H₂₁NO₃ 311.38
Alkaloid from the twigs of *Discaria*
serratifolia var. *montana* (Rhamna-
ceae). Amorph. solid (EtOH). Mp 230-
231°. [α]_D²⁰ +244 (c, 0.037 in CHCl₃).

2,11-Di-Me ether: 1-Hydroxy-2,11-di-
methoxyaporphine. **Isothebaine**
[568-21-8]
C₁₉H₂₁NO₃ 311.38
Alkaloid from *Papaver orientale* and
Papaver bracteatum (Papaveraceae).
Convulsive agent in high doses. Small
doses increase muscle tonus in intesti-
nal tissue, higher doses produce re-
laxation. Increases tonus in uterine
tissue. Respiratory depressant, shows
analgesic and some antiinflammatory
action. Mp 164-166° (203-204°). [α]_D²¹
+281 (c, 1.11 in CHCl₃). λ_{max} 228 (log
ε 4.12); 271 (log ε 4.07); 294 (sh) (log ε
3.81) (MeOH).

► LD₅₀ (mus, scu) 26 mg/kg.

2,11-Di-Me ether, N-oxide: **Isothebaine**
N-oxide
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Papaver pseudo-orien-*
tale. Amorph. [α]_D +260 (CHCl₃).

2,11-Di-Me ether, N-Me: **N-Methyl-**
isothobainium
[25651-02-9]
C₂₀H₂₄NO₃⁺ 326.414
Alkaloid from *Papaver orientale*, *Pa-*
paver pseudo-orientale and *Papaver*
bracteatum. Needles (MeOH) (as io-
dide). Mp 254-256° (249-253°) (io-
dide). [α]_D²¹ +194 (c, 0.51 in MeOH).
CAS no. refers to iodide. λ_{max} 210 (log
ε 4.5); 271 (log ε 4.24); 314 (log ε 3.66)
(MeOH).

Tri-Me ether: 1,2,11-Trimethoxyaporphine. **O-Methylisothobaine**
[90275-86-8]
C₂₀H₂₃NO₃ 325.407
Alkaloid from *Papaver orientale* (Pa-
paveraceae). Mp 98°. [α]_D +26 (c, 0.45
in CHCl₃). λ_{max} 273 (log ε 4.26); 303
(log ε 3.31) (EtOH).

Tri-Me ether, N-de-Me: **1,2,11-Tri-**
methoxynoraporphine
[111261-80-4]
C₁₉H₂₁NO₃ 311.38
Alkaloid from leaves and stems of
Discaria chacaya.

Tri-Me ether, N-Me: **Zanthoxyphylline**.
Xanthoxyphylline
[68711-44-4]
C₂₁H₂₆NO₃⁺ 340.441
Quaternary alkaloid from the roots of
Zanthoxylum oxyphyllum (Rutaceae).

Cryst. (MeOH). Mp 202° dec. (as
hydroxide) Mp 300° (as iodide). [α]_D²²
+51.7 (MeOH). λ_{max} 230 ; 278 (sh) ;
310 (EtOH).

6a,7-Didehydro, 2,11-di-Me ether: **Dehy-**
droisothobaine
[25984-81-0]
C₁₉H₁₉NO₃ 309.364
Alkaloid from *Papaver orientale* (Pa-
paveraceae). λ_{max} 267 ; 340 ; 391 ; 439
(no solvent reported).

6a,7-Didehydro, tri-Me ether: **Orientidine**
[90275-84-6]
C₂₀H₂₁NO₃ 323.391
Alkaloid from *Papaver orientale*
(Papaveraceae). λ_{max} 215 (log ε 4.45);
271 (log ε 4.39); 340 (log ε 3.38)
(EtOH).

(R)-form

1,2-Methylene ether: see Pukateine, P-781

(±)-form

Tri-Me ether:

Cryst. (petrol). Mp 98°. CAS no. not
found 8-14 CI. λ_{max} 230 (log ε 4.1); 270
(log ε 4.1); 300 (log ε 3.9) (no solvent
reported).

Govindachari, T.R. *et al.*, *J.O.C.*, 1953, **18**,
1352-1355 (*tri-Me ether, synth*)

Battersby, A.R. *et al.*, *J.C.S.*, 1965, 4550-4556
(*synth, stereochem*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1969,
464-465 (*biosynth*)

Israelov, I.A. *et al.*, *Khim. Prir. Soedin.*,
1978, **14**, 474-475; *Chem. Nat. Compd.*
(*Engl. Transl.*), 1978, **14**, 402-403
(*Isothebaidine*)

Tiwari, K.P. *et al.*, *Phytochemistry*, 1978, **17**,
1068-1069 (*Zanthoxyphylline*)

Slavik, J. *et al.*, *Coll. Czech. Chem. Comm.*,
1980, **45**, 914-920; 1985, **50**, 1216-1226;
1991, **56**, 1534-1538 (*N-*
Methylisothobainium)

Dolejš, L. *et al.*, *Coll. Czech. Chem. Comm.*,
1984, **49**, 2816-2818 (*N-Methylisothobaine,*
ms)

Rivera, A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 1040
(*11-Hydroxy-1,2-dimethoxyaporphine*)

Israelov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1984,
20, 81-83; 258-259; *Chem. Nat. Compd.*
(*Engl. Transl.*), 76-78; 243-244 (*Orientidine,*
O-Methylisothobaine, Dehydroisothobaine)

Věžník, F. *et al.*, *Coll. Czech. Chem. Comm.*,
1986, **51**, 1752-1763 (*N-*
Methylisothobainium, N-Methylisothobaine)


Correa, C. *et al.*, *Bol. Soc. Chil. Quim.*, 1987,
32, 105; *CA*, **107**, 214795q (*1,2,11-*
Trimethoxynoraporphine)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1996,
32, 761-863; *Chem. Nat. Compd. (Engl.*
Transl.), 1996, **32**, 737-858 (*Isothebaine N-*
oxide)

Marek, R. *et al.*, *Magn. Reson. Chem.*, 1999,
37, 195-202; 2002, **40**, 687-692 (*Isothebaine,*
N-15 nmr, cryst struct)

2,6,7-Trihydroxy-2H-1,4-
benzoxazin-3(4H)-one

T-526



C₈H₇NO₅ 197.147

(R)-form

N-Hydroxy, 6,7-di-Me ether, 2,4-Dihydroxy-6,7-dimethoxy-2H-1,4-benzoxazin-3(4H)-one, 9CI
[67467-45-2]

C₁₀H₁₁NO₆ 241.2

Constit. of sweet corn (*Zea mays*).

Hofman, J. et al., *Phytochemistry*, 1973, **12**, 207-208

2,7,8-Trihydroxy-2H-1,4-benzoxazin-3(4H)-one

T-527

C₈H₇NO₅ 197.147

(R)-form

N-Hydroxy, 7,8-di-Me ether, 2,4-Dihydroxy-7,8-dimethoxy-2H-1,4-benzoxazin-3(4H)-one, 9CI
[60032-92-0]

C₁₀H₁₁NO₆ 241.2

Constit. of wheat and sweet corn.

N-Hydroxy, 7,8-di-Me ether, 2-O-β-D-glucopyranoside: [40246-08-0]

C₁₆H₂₁NO₁₁ 403.342


Isol. from sweet corn (*Zea mays*).

Hofman, J. et al., *Phytochemistry*, 1973, **12**, 207-208

2,3,5-Trihydroxybenzylamine

T-528

6-(Aminomethyl)-1,2,4-benzenetriol, 9CI.
α-Amino-2,3,5-trihydroxytoluene



C₇H₉NO₃ 155.153

N-Me, 3-O-β-D-glucopyranoside: **Zinolol**
[942626-27-9]

C₁₄H₂₁NO₈ 331.322

Constit. of *Anagallis monelli*. Antioxidant and mutagenic agent. Amorph. solid. Mp 164-166°. [α]_D²⁵ -375 (c, 0.04 in MeOH). λ_{max} 244 (ε 10450) (MeOH).


▶ Mutagen.

Ammar, S. et al., *Chem. Pharm. Bull.*, 2007, **55**, 385-388 (*Zinolol*)

1,2,7-Trihydroxy-3,5-bis(hydroxymethyl)-1H-pyrrolizidine

T-529

Hexahydro-1,2,7-trihydroxy-1H-pyrrolizidine-3,5-dimethanol



(1R*,2S*,3S*,5R*,7R*,7aR*)-form

C₉H₁₇NO₅ 219.237

(1R*,2S*,3S*,5R*,7R*,7aR*)-form Hyacinthacine C₂

[944408-25-7]

Alkaloid from the bulbs of *Scilla socialis*. Glycosidase inhibitor. Powder. [α]_D +12.9 (c, 0.22 in H₂O).

(1R*,2S*,3S*,5R*,7S*,7aR*)-form

Hyacinthacine C₃

[944408-26-8]


Alkaloid from the bulbs of *Scilla socialis*. Powder. [α]_D +3.5 (c, 0.23 in H₂O).

Kato, A. et al., *J. Nat. Prod.*, 2007, **70**, 993-997 (isol, pmr, cmr)

1,10,11-Trihydroxycanthin-6-one

T-530

1,10,11-Trihydroxy-6H-indolo[3,2,1-de][1,5]naphthyrin-6-one, 9CI



C₁₄H₈N₂O₄ 268.228

1,11-Di-Me ether: 10-Hydroxy-1,11-dimethoxycanthin-6-one. **Bruceoline C**
[165967-62-4]

C₁₆H₁₂N₂O₄ 296.282

Alkaloid from root bark of *Brucea mollis* var. *tonkinensis* (Simaroubaceae). Amorph. yellow powder. λ_{max} 276 (sh) (log ε 3.37); 286 (log ε 3.45); 350 (log ε 3.35); 370 (log ε 3.35) (MeOH).

Ouyang, Y. et al., *Phytochemistry*, 1995, **39**, 911-913 (isol, uv, ir, pmr, cmr, ms, struct)

4,5,10-Trihydroxycanthin-6-one

T-531

C₁₄H₈N₂O₄ 268.228

5-Me ether: 4,10-Dihydroxy-5-methoxycanthin-6-one. **Picrasidine W**
[155416-28-7]

C₁₅H₁₀N₂O₄ 282.255

Alkaloid from the wood of *Picrasma quassioides* (Simaroubaceae). Pale yellow needles (MeOH). Mp 225-227°. λ_{max} 230 (log ε 4.53); 242 (log ε 4.5); 254 (sh) (log ε 4.44); 282 (log ε 4.4); 304 (sh) (log ε 4.13); 355 (log ε 4.21); 370 (log ε 4.21) (MeOH).

Tri-Me ether:


Pale yellow needles. Mp 258-260°.

Li, H.-Y. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1807-1811 (isol, uv, ir, pmr, cmr, ms, struct)

1,6,7-Trihydroxy-9H-carbazole-3-carboxaldehyde

T-532

3-Formyl-1,6,7-trihydroxycarbazole



C₁₃H₉NO₄ 243.218

6-Me ether: 1,7-Dihydroxy-6-methoxy-9H-carbazole-3-carboxaldehyde. **Clausine J**

[186002-91-5]

C₁₄H₁₁NO₄ 257.245

Alkaloid from stem bark of *Clausena excavata*. Powder (Me₂CO). Mp >290°. λ_{max} 235; 255 (sh); 289; 305; 347 (sh) (MeOH).

Wu, T.-S. et al., *Phytochemistry*, 1996, **43**, 1427 (*Clausine J*)

2,6,8-Trihydroxy-9H-carbazole-3-carboxaldehyde

T-533

6-Formyl-1,3,7-trihydroxycarbazole

C₁₃H₉NO₄ 243.218

6,8-Di-Me ether: 2-Hydroxy-6,8-dimethoxy-9H-carbazole-3-carboxaldehyde. 6-Formyl-7-hydroxy-1,3-dimethoxycarbazole. **Clausine B**

[182261-81-0]


C₁₅H₁₃NO₄ 271.272

Alkaloid from stem bark of *Clausena excavata*. Yellow needles (Me₂CO). Mp 228-229°. λ_{max} 210 (log ε 4.28); 235 (log ε 4.3); 269 (sh) (log ε 4.24); 278 (log ε 4.32); 304 (log ε 4.36); 357 (log ε 3.83) (MeOH).

Wu, T.-S. et al., *Phytochemistry*, 1996, **43**, 133-140 (*Clausine B*)

14,16,20-Trihydroxyceve-4-en-3-one

T-534



C₂₇H₄₁NO₄ 443.625

(16β,20β)-form

Veratrenone


[55839-66-2]

Alkaloid from *Veratrum album* var. *grandiflorum* (*Veratrum grandiflorum*) (Liliaceae). Mp 217-218°. [α]_D +60.5 (EtOH). λ_{max} 240 (14000) (EtOH).

Takasugi, M. et al., *Chem. Lett.*, 1974, 1477-1480 (isol, ir, uv, pmr, cryst struct)

6,7,8-Trihydroxy-3,11,13-clerodatrien-15,16-olide

T-535



C₂₀H₂₈O₅ 348.438**(ent-6β,7α,8α,11E)-form****Barbatin C**

[905929-94-4]

Constit. of *Scutellaria barbata*. Cryst. Mp 156-158°. $[\alpha]_D^{20}$ -103.8 (c, 0.14 in MeOH). λ_{\max} 220 ; 257 (CDCl₃).

6-O-(3-Pyridinecarbonyl), 7-benzoyl:

Scutebarbatine B

[905929-95-5]

C₃₃H₃₅NO₇ 557.642


Constit. of *Scutellaria barbata*. Cryst. Mp 151-153°. $[\alpha]_D^{20}$ -109.6 (c, 0.13 in MeOH). λ_{\max} 217 ; 222 ; 257 (CDCl₃).

6,7-Bis(3-pyridinecarbonyl): **Scutebarbatine A**


[176520-13-1]

C₃₂H₃₄N₂O₇ 558.63

Constit. of *Scutellaria barbata*. Needles (Me₂CO). Mp 148-150°. $[\alpha]_D^{20}$ -128.7 (c, 0.2 in MeOH).

Wang, Z.-Q. et al., *Chin. Chem. Lett.*, 1996, 7, 333-334 (isol, pmr, cmr)Dai, S.-J. et al., *Phytochemistry*, 2006, 67, 1326-1330 (Barbatin C, Scutebarbatine B)**1,4,5-Trihydroxy-2-cyclopentene-1-carboxamide T-536**C₆H₉NO₄ 159.141

Alkaloid from *Lindackeria dentata*. Cubes (MeOH). Mp 208-210°. $[\alpha]_D^{20}$ -10 (c, 0.1 in MeOH). Shown to be a mixt. of diastereoisomers by synth.

Gibbons, S. et al., *Phytochemistry*, 1998, 49, 2395-2396 (isol, ir, pmr, cmr, ms)Jaroszewski, J.W. et al., *Planta Med.*, 2004, 70, 1001-1003 (synth, pmr, cmr)**1,2,3-Trihydroxydibenz[cd,f]indol-4(5H)-one, 9CI T-537**C₁₅H₉NO₄ 267.24

CAS numbering shown. Alternative (phenanthrene) numbering frequently encountered, in which 1,2,3 are 4,3,2-respectively.

1,2-Di-Me ether: **Piperolactam B**

[128718-51-4]

C₁₇H₁₃NO₄ 295.294

Alkaloid from *Piper attenuatum* and *Piper boehmerifolium*. Yellow cryst. (C₆H₆/MeOH). Mp 226-227°. This struct. was originally (1990) assigned to 1,2,3-Trihydroxydibenz[cd,f]indol-

4(5H)-one, T-537.

1,3-Di-Me ether: **Goniopedaline**. *Aristolactam F II*. *Goniopetaline* (incorr.)

[112501-43-6]

C₁₇H₁₃NO₄ 295.294

Alkaloid from the leaves and twigs of *Goniotalamus sesquipedalis* and the root of *Pararistolochia flos-avis*. Shows platelet aggregation inhibitory activity. Fine greenish-yellow cryst. (CHCl₃/petrol). Mp 225-227° (218°).

2,3-Di-Me ether: 1-Hydroxy-2,3-dimethoxydibenz[cd,f]indol-4(5H)-one, 9CI. 10-Amino-2,3-dimethoxy-4-hydroxyphenanthrene-1-carboxylic acid lactam. **Piperolactam D**

[116084-93-6]

C₁₇H₁₃NO₄ 295.294

Alkaloid from the roots of *Piper longum* (long pepper), stems of *Piper acutislegium* and whole plants of *Piper boehmerifolium* (Piperaceae). Greenish-yellow cryst. (MeOH). Mp 222-224°. This struct. was originally assigned to Piperolactam B by Desai et al (1988).

2,3-Di-Me ether, N,O-di-Ac:

C₂₁H₁₇NO₆ 379.368

Cryst. (C₆H₆). Mp 260-262° (194-195°).

2,3-Di-Me ether, N-Me: N-Methylpiperolactam D. **Piperolactam E**

[219687-22-6]

C₁₈H₁₅NO₄ 309.321

Alkaloid from *Piper augustum* and *Piper taiwanense*. Greyish needles (C₆H₆/MeOH). Mp 179-181°. λ_{\max} 228 (log ϵ 3.9); 269 (sh) (log ϵ 4.11); 296 (log ϵ 3.73); 336 (log ϵ 3.54); 373 (log ϵ 3.31) (MeOH).

Tri-Me ether: 1,2,3-Trimethoxydibenz[cd,f]indol-4(5H)-one, 9CI. **Piperolactam C**


[116064-76-7]

C₁₈H₁₅NO₄ 309.321

Alkaloid from whole plants of *Piper longum* (long pepper) and *Piper boehmerifolium* (Piperaceae). Shows platelet aggregation inhibitory activity. Yellow cryst. (C₆H₆/MeOH). Mp 187-188°.

Priestap, H.A. et al., *Phytochemistry*, 1985, 24, 849-852 (*Goniopedaline*)Villar, A. et al., *Planta Med.*, 1986, 52, 556-557 (activity)Sun, N.J. et al., *J. Nat. Prod.*, 1987, 50, 843 (*Goniopedaline*)Talapatra, S.K. et al., *Phytochemistry*, 1988, 27, 903-906 (*Goniopedaline*)Desai, S.J. et al., *Indian J. Chem., Sect. B*, 1989, 28, 775 (*Piperolactam C*)Olsen, C.E. et al., *Phytochemistry*, 1993, 33, 518-520 (*Piperolactam B*, *Piperolactam D*)Delgado, A.W. et al., *Rev. Colomb. Quim.*, 1998, 27, 13-21 (*Piperolactam E*)Chia, Y.-C. et al., *J. Nat. Prod.*, 2000, 63, 1160-1163 (isol, activity)Rys, V. et al., *Eur. J. Org. Chem.*, 2003, 1231-1237 (*Goniopedaline*, *Piperolactam C*, synth)Chen, Y.-C. et al., *Planta Med.*, 2004, 70, 174-177 (*Piperolactam E*)**1,2,6-Trihydroxydibenz[cd,f]indol-4(5H)-one, 9CI**

T-538

C₁₅H₉NO₄ 267.24

CAS numbering shown. Alternative (phenanthrene) numbering frequently used, in which 1,2,6- are 4,3,9- respectively.

1,2-Methylene, 6-Me ether: **9-Methoxyaristolactam II**

[133485-40-2]

C₁₇H₁₁NO₄ 293.278

Alkaloid from roots of *Aristolochia auricularia* (Aristolochiaceae). Pale yellow solid.

Houghton, P.J. et al., *Phytochemistry*, 1991, 30, 253**1,2,7-Trihydroxydibenz[cd,f]indol-4(5H)-one, 9CI T-539**C₁₅H₉NO₄ 267.24

CAS numbering. Alternative (phenanthrene) numbering frequently used in which 1,2,7- become 4,3,8- respectively.

1-Me ether: **Aristolactam A Ia**

[97399-90-1]

C₁₆H₁₁NO₄ 281.267

Alkaloid from the rhizomes of *Aristolochia argentina* (Aristolochiaceae). Cryst. (AcOH). Mp 350°.

1,2-Di-Me ether: **Velutinam**

[146428-62-8]

Alkaloid from leaves of *Goniotalamus velutinus*. Shows platelet aggregation inhibitory activity. Brownish-yellow needles. Mp 267-270°.

1,7-Di-Me ether: **Enterocarpam II**

[102719-96-0]

C₁₇H₁₃NO₄ 295.294

Alkaloid from stem bark of *Orophea enterocarpa*. Mp 268-272°.

1,7-Di-Me ether, N,O-di-Ac:

Cryst. (MeOH). Mp 219-220°.

2,7-Di-Me ether: **Griffithinam**. *Goniofithine*. *Uvarilactam*

[240122-32-1]

C₁₇H₁₃NO₄ 295.294

Alkaloid from the roots of *Goniotalamus griffithii* and the stems of *Uvaria microcarpa*. Pale yellow needles (MeOH). Mp 262-264° Mp 312-314°. λ_{\max} 239 (log ϵ 4.43); 257 (log ϵ 4.51); 289 (log ϵ 4.25); 325 (log ϵ 4.05); 340 (log ϵ 3.94); 385 (log ϵ 3.92) (MeOH).

Tri-Me ether: **Aristolactam B I**. *Taliscanine*

[7688-86-0]


C₁₈H₁₅NO₄ 309.321

Alkaloid from the roots of *Aristolochia taliscana* (Aristolochiaceae). Mp 272-

273° (260°).

Tri-Me ether, N-Ac: Mp 236-237°.

1,2-Methylene, 7-Me ether: see Aristolactam I, A-1427

Maldonado, L.A. et al., *Ciencia (Mexico City)*, 1966, **24**, 237; *CA*, **65**, 15438e (Taliscanine)Priestap, H.A. et al., *Phytochemistry*, 1985, **24**, 849 (Aristolactams)Mahmood, K. et al., *Phytochemistry*, 1986, **25**, 965-967 (Enterocarpan II, Velutinam)Omar, S. et al., *Phytochemistry*, 1992, **31**, 518 (Velutinam)Zhang, Y.-J. et al., *J. Nat. Prod.*, 1999, **62**, 1050-1052 (Griffithinam)Yu, D.-L. et al., *Zhiwu Xuebao (Acta Bot. Sin.)*, 1999, **41**, 1104-1107 (Uvarilactam)Chia, Y.-C. et al., *J. Nat. Prod.*, 2000, **63**, 1160-1163 (isol, activity)Hu, Z. et al., *Yaoxue Xuebao*, 2000, **35**, 277-278 (Goniofithine)**1,2,9-Trihydroxydibenz[cd,-f]indol-4(5H)-one, 9CI** T-540C₁₅H₉NO₄ 267.24

CAS numbering shown. Alternative (phenanthrene) numbering frequently used, in which 1,2,9- become 4,3,6- respectively.

N-β-D-Glucopyranosyl: [208774-76-9]

C₂₁H₁₉NO₉ 429.382Alkaloid from *Aristolochia constricta*. Yellow needles. Mp 240-248°. [α]_D²⁵ -12 (c, 1 in MeOH).1-Me ether: **Aristolactam A IIIa**

[97399-91-2]

C₁₆H₁₁NO₄ 281.267Alkaloid from *Aristolochia argentina* and *Fissistigma* spp. Mp >350°. λ_{max} 212 (log ε 4.39); 237 (log ε 4.53); 252 (log ε 4.38); 260 (sh) (log ε 4.37); 279 (log ε 4.27); 292 (log ε 4.28); 321 (log ε 4.08); 402 (log ε 3.87) (EtOH).

1-Me ether, 9-O-β-D-glucopyranoside:

Aristolactam F

[710319-84-9]

C₂₂H₂₁NO₉ 443.409Alkaloid from *Aristolochia elegans*. Yellow syrup. [α]_D²⁵ -7.8 (c, 0.7 in MeOH). λ_{max} 240 (log ε 4.55); 246 (log ε 4.33); 280 (log ε 4.23); 290 (log ε 4.15); 324 (log ε 4.02) (MeOH).1,2-Di-Me ether: **Goniothalactam**

[204975-46-2]

C₁₇H₁₃NO₄ 295.294Alkaloid from *Goniothalamus borneensis* (Annonaceae). Constit. of *Fissistigma* spp. Shows platelet aggregation inhibitory activity. Light yellow needles. Mp 257-259°. λ_{max} 236 (log ε 4.58); 256 (log ε 4.45); 262 (log ε 4.48); 294 (log ε 4.35); 320 (log ε 4.21)

(EtOH).

1,2-Di-Me ether, N-β-D-glucopyranosyl:

[208774-72-5]

C₂₃H₂₃NO₉ 457.436Alkaloid from *Aristolochia constricta*. Yellow needles. Mp 250-259°. [α]_D²⁵ -14.5 (c, 1 in MeOH).

1,2-Di-Me ether, N-[4-hydroxy-E-cinnamoyl-(→6)-β-D-glucopyranosyl]:

[208774-73-6]

C₃₂H₂₉NO₁₁ 603.581Alkaloid from *Aristolochia constricta*. Yellow needles. Mp 270-275°. [α]_D²⁵ -21.5 (c, 1 in MeOH).1,9-Di-Me ether: **Aristolactam A III**

[53948-08-6]

C₁₇H₁₃NO₄ 295.294Alkaloid from the roots of *Aristolochia argentina* (Aristolochiaceae) and *Fissistigma* spp. Shows platelet aggregation inhibitory activity. Yellow cryst. (AcOH). Mp 275°.Tri-Me ether: **Aristolactam B III**

[53948-10-0]

C₁₈H₁₅NO₄ 309.321Alkaloid from the roots of *Aristolochia argentina* (Aristolochiaceae) and *Fissistigma* spp. Shows platelet aggregation inhibitory activity. Yellow-green cryst. (1-propanol/1-butanol). Mp 264-265°.

Tri-Me ether, N-β-D-glucopyranosyl:

Aristolactam B III N-glucoside

[208774-74-7]

C₂₄H₂₅NO₉ 471.463Alkaloid from *Aristolochia constricta*. Yellow needles. Mp 260-269°. [α]_D²⁵ -13.5 (c, 1 in MeOH).

1,2-Methylene ether: see Aristolactam

IIIa, A-1429

Crohare, R. et al., *Phytochemistry*, 1974, **13**,

1957-1962 (Aristolactams)

Priestap, H.A. et al., *Phytochemistry*, 1985, **24**,

849-852 (Aristolactams, pmr, cmr)

Sun, N.J. et al., *J. Nat. Prod.*, 1987, **50**, 843-846

(Goniothalactam)

Desai, S.J. et al., *Phytochemistry*, 1988, **27**,

1511-1515 (Aristolactam BIII)

Estévez, J.C. et al., *Tet. Lett.*, 1989, **30**, 5785

(Aristolactam BIII, synth)

De Tomassi, N. et al., *Nat. Prod. Lett.*, 1998,**11**, 263-270 (*Aristolochia constricta*

alkaloids)

Cao, S.-G. et al., *Tetrahedron*, 1998, **54**, 2143-

2148 (Goniothalactam)

Couture, A. et al., *Nat. Prod. Lett.*, 1999, **13**,


33-40 (Goniothalactam, synth)

Chia, Y.-C. et al., *J. Nat. Prod.*, 2000, **63**, 1160-

1163 (isol, activity)

Shi, L.-S. et al., *Bioorg. Med. Chem.*, 2004, **12**,439-446 (*Aristolactam AIIIa*)**2,10,13-Trihydroxydibenz[ol,a]quinolinizinium(1+)** T-541

[169706-67-8]


C₁₇H₁₂NO₃[⊕] 278.287Alkaloid from roots of *Aristolochia arcuata*. Red amorph. solid + 2H₂O (MeOH)(as hydroxide). Mp 250° dec. (as hydroxide). CAS no. refers to hydroxide.

10-O-Glucopyranoside: [169700-68-9]

C₂₃H₂₂NO₈[⊕] 440.429From roots of *Aristolochia arcuata*.

Amorph. red solid (MeOH) (as hydroxide). Mp 248° dec. (hydroxide).

CAS no. refers to hydroxide.

Watanabe, L.Y. et al., *Phytochemistry*, 1995,**40**, 991 (isol, uv, ir, pmr, cmr, ms, struct)**1,6,9-Trihydroxydihydro-β-agarofuran** T-542C₁₅H₂₆O₄ 270.368**(1α,6β,9β)-form****Celorbicol**

[59812-41-8]

Parent alcohol of esters from the seed oil of *Celastrus orbiculatus*. Cryst. (CHCl₃/Me₂CO). Mp 222-223°. [α]_D²⁶ -24 (c, 0.5 in CHCl₃).


6-(3-Pyridinecarbonyl), 9-benzoyl, 1-Ac:

Triptogelin F1

[145940-86-9]

C₃₀H₃₅NO₇ 521.609Constit. of *Tripterygium wilfordii*.Needles. Mp 193-195°. [α]_D²⁵ +67.7 (c, 0.4 in MeOH).Takaishi, Y. et al., *Phytochemistry*, 1992, **31**,3943-3947 (*Triptogelins*)**3,4,6-Trihydroxy-1,2-dimethylcarbazole** T-543

1,2-Dimethyl-9H-carbazole-3,4,6-triol, 9CI

C₁₄H₁₃NO₃ 243.2623,6-Di-Me ether: 4-Hydroxy-3,6-dimethoxy-1,2-dimethylcarbazole. **Carbazomycin C**

[108073-62-7]

C₁₆H₁₇NO₃ 271.315Isol. from *Streptovorticillium chimense*. Pale-yellow prisms (EtOAc/hexane).Mp 198-198.5°. λ_{max} 228 (ε 25000); 246

(ε 24000); 260 (sh) (ε 12500); 288 (sh)

(ε 7600); 297 (ε 12200); 341 (ε 3600);

355 (ε 4200) (MeOH) (Derep).

Tri-Me ether: 3,4,6-Trimethoxy-1,2-di-

methylcarbazole. **Carbazomycin D**

[108073-63-8]

C₁₇H₁₉NO₃ 285.342
From *Streptovorticillium ehimensis*.
Needles (CH₂Cl₂/hexane). Sol. MeOH,
CHCl₃; poorly sol. H₂O. Mp 129.5-
130°. λ_{max} 228 (ε 25000); 246 (ε 24000);
260 (sh) (ε 12500); 288 (sh) (ε 7600);
297 (ε 12200); 341 (ε 3600); 355 (ε
4200) (MeOH) (Derep). λ_{max} 229 (ε
30500); 247 (ε 27800); 300 (ε 1700); 340
(ε 4300) (MeOH) (Berdy).


Naid, T. et al., *J. Antibiot.*, 1987, **40**, 157 (*isol. struct. props*)

Kameda, M. et al., *J. Antibiot.*, 1988, **41**, 602 (*cryst. struct.*)

Knölker, H.J. et al., *J.C.S. Perkin 1*, 1997, 349 (*synth.*)

1,6,8-Trihydroxy-3,4-dimethylisoquinoline T-544

6,8-Dihydroxy-3,4-dimethyl-1(2H)-isoquinolinone, 9CI. 3,4-Dimethyl-1,6,8-isoquinolinetriol. *Siaminine A* [92446-24-7]



C₁₁H₁₁NO₃ 205.213
Alkaloid from the leaves of *Cassia siamea* (Fabaceae). Mp 215° (as hydrochloride).

El-Sayyad, S.M. et al., *J. Nat. Prod.*, 1984, **47**, 708 (*isol. uv, ir, pmr, ms, struct.*)

7,18,19-Trihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione T-545



Relative Configuration

C₂₈H₃₇NO₅ 467.604
Isol. from *Microporellus subsessilis* and an unidentified *Daldinia* fungus. Mycotoxin. Amorph. solid. Mp 217-219°. [α]_D +2.6 (c, 2.56 in CHCl₃).

19-Ac:

C₃₀H₃₉NO₆ 509.641
From *Daldinia* sp. Needles (C₆H₆). Mp 138-140°. [α]_D -23.7 (c, 0.38 in CHCl₃).

19-O-(4ξ-Methyl-6E,8E-hexadecadienyl):

C₄₅H₆₅NO₆ 716.012
Isol. from *Microporellus subsessilis*. Amorph. solid (MeOH). [α]_D²⁸ -22 (c, 0.002 in CHCl₃). λ_{max} 232 (ε 55410) (MeOH).

19-Me ether: 7,18-Dihydroxy-19-methoxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione

C₂₉H₃₉NO₅ 481.631
From *Daldinia* sp. and *Microporellus*

subsessilis. Amorph. solid. Mp 124-126°. [α]_D -24.8 (c, 1.29 in CHCl₃).

18-Deoxy: 7,19-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione

C₂₈H₃₇NO₄ 451.605
From *Daldinia* sp. Cryst. (EtOAc). Mp 196-199°. [α]_D +70.3 (c, 0.13 in MeOH). Also descr. as an amorph. solid to which the opt. rotn. refers.

19-Deoxy: 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione

C₂₈H₃₇NO₄ 451.605
From *Daldinia* sp. and *Microporellus subsessilis*. Amorph. solid. Mp 120-122°. [α]_D -12.6 (c, 1.04 in CHCl₃).

19-Deoxy, 19,20-didehydro(E-): 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-triene-1,21-dione

C₂₈H₃₅NO₄ 449.589
From *Daldinia* sp. Needles (EtOAc). Mp 179-181°. [α]_D -14.8 (c, 0.88 in CHCl₃).

18,19-Dideoxy: 7-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione

C₂₈H₃₇NO₃ 435.605
From *Daldinia* sp. Amorph. solid. [α]_D +66.5 (c, 0.18 in MeOH).

18,19-Dideoxy, 19,20-didehydro(E-): 7-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-triene-1,21-dione


C₂₈H₃₅NO₃ 433.589
From *Daldinia* sp. Amorph. solid. [α]_D -16.1 (c, 0.31 in CHCl₃).

Buchanan, M.S. et al., *Phytochemistry*, 1995, **40**, 135; 1996, **41**, 821; **42**, 173

Kurnia, D. et al., *Phytochemistry*, 2007, **68**, 697-702 (*Microporellus constits*)

1,3,5-Trihydroxy-2,4-diprenylacridone T-546

1,3,5-Trihydroxy-2,4-bis(3-methyl-2-butenyl)-9(10H)-acridinone, 9CI. *Atalaphylline* [28233-35-4]



C₂₃H₂₅NO₄ 379.455
Alkaloid from root bark of *Atalantia monophylla* (Rutaceae). Yellow needles (C₆H₆/EtOAc). Mp 246°. λ_{max} 255 (log ε 4.53); 265 (log ε 4.54); 284 (log ε 4.48); 312 (log ε 4.2); 402 (log ε 3.75) (EtOH).

N-Me: 1,3,5-Trihydroxy-9-methyl-2,4-diprenylacridone. N-*Methylatalaphylline*

[28233-34-3]
C₂₄H₂₇NO₄ 393.482

Constit. of *Atalantia monophylla* (Rutaceae). Mp 192-193°. λ_{max} 260 (sh) (log ε 4.36); 273 (log ε 4.48); 336 (log ε 4.06); 415 (log ε 3.65) (EtOH).

3-Me ether: 1,5-Dihydroxy-3-methoxy-2,4-diprenylacridone. *Buxifoliadine B* [263007-66-5]

C₂₄H₂₇NO₄ 393.482
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 238-240°. λ_{max} 259 (log ε 4.71); 309 (log ε 4.17); 322 (log ε 4.13); 408 (log ε 3.82) (MeOH).

3-Me ether, N-Me: 1,5-Dihydroxy-3-methoxy-10-methyl-2,4-diprenylacridone. *Buxifoliadine A* [263007-65-4]

C₂₅H₂₉NO₄ 407.508
Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 155-157°. λ_{max} 278 (log ε 4.54); 326 (log ε 3.9); 425 (log ε 3.65) (MeOH).

3,5-Di-Me ether: 1-Hydroxy-3,5-dimethoxy-2,4-diprenylacridone. 3,5-Di-O-methylatalaphylline. *Atalaphylline dimethyl ether*

[28233-37-6]
C₂₅H₂₉NO₄ 407.508
Alkaloid from *Atalantia monophylla*. Yellow needles (Et₂O/hexane). Mp 145-147°.

A^{3'}-Isomer, 2''-ξ-hydroxy, N-Me: *Bosistine*

C₂₄H₂₇NO₅ 409.481
Alkaloid from leaves and stem bark of *Bosistoa transversa*. Amorph. [α]_D -4.1 (c, 0.0013 in CHCl₃). Incorrectly indexed by CA.

Govindachari, T. et al., *Tetrahedron*, 1970, **26**, 2905-2910 (*N-Methylatalaphylline, isol. pmr, ms, uv, ir*)

Desai, H. et al., *Indian J. Chem., Sect. B*, 1976, **14**, 473-475 (*isol.*)

Banerji, J. et al., *Indian J. Chem., Sect. B*, 1981, **20**, 835-838 (*cmr*)

Kulkarni, G.H. et al., *Phytochemistry*, 1981, **20**, 867 (*di-Me ether*)

Shah, J.S. et al., *Indian J. Chem., Sect. B*, 1982, **21**, 16-19 (*N-Methylatalaphylline*)

Wu, T.-S. et al., *Phytochemistry*, 1982, **21**, 1771-1773 (*N-Methylatalaphylline*)

Bahar, M.H. et al., *Phytochemistry*, 1982, **21**, 2729-2731 (*synth.*)

Auzi, A.A. et al., *Phytochemistry*, 1996, **42**, 235-238 (*Bosistine*)

Wu, T.-S. et al., *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*Buxifoliadines*)

1,3,5-Trihydroxy-2,8-diprenylacridone T-547

1,3,5-Trihydroxy-2,8-bis(3-methyl-2-butenyl)-9H-acridin-10-one

C₂₃H₂₅NO₄ 379.455

N-Me: 1,3,5-Trihydroxy-9-methyl-2,8-bis(3-methyl-2-butenyl)-9H-acridin-10-one. 1,3,5-Trihydroxy-9-methyl-2,8-di-

prenylacridoneC₂₄H₂₇NO₄ 393.482

Alkaloid from the stem bark of *Swin-glea glutinosa*. Amorph. powder. λ_{\max} 262 (log ϵ 3.49); 288 (log ϵ 3.12); 320 (log ϵ 3.03); 409 (log ϵ 2.69) (no solvent reported).

Weniger, B. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1221-1223

1,3,8-Trihydroxy-2,7-diprenyl-lacridone T-548

1,3,8-Trihydroxy-2,7-bis(3-methyl-2-but-2-enyl)-9(10H)-acridinone, 9CI.

Atalaphyllidine[†]

[60776-36-5]


C₂₃H₂₅NO₄ 379.455

Constit. of the roots of *Atalantia monophylla* (Rutaceae). Yellow needles (C₆H₆/EtOAc). Mp 222°.

1,3-Di-Me ether: 8-Hydroxy-1,3-dimethoxy-2,7-diprenylacridone

Yellow needles (Me₂CO). Mp 118°.


Chatterjee, A. *et al.*, *Phytochemistry*, 1976, **15**, 1303 (*isol, uv, ir, pmr, struct*)

7,11,14-Trihydroxyhetisan-2,13-dione T-549C₂₀H₂₃NO₅ 357.405**(7 α ,11 α)-form****Orientinine**[†]

[174293-69-7]


Alkaloid from aerial parts of *Aconitum orientale*. $[\alpha]_D^{25}$ +42 (c, 0.1 in CHCl₃).

Ulubelen, A. *et al.*, *Phytochemistry*, 1996, **41**, 957 (*isol, ir, pmr, cmr, ms, struct*)

1,2,6-Trihydroxyhetisan-13-one T-550C₂₀H₂₅NO₄ 343.422**(1 β ,2 α)-form***1-Ac*: [943912-59-2]C₂₂H₂₇NO₅ 385.459

Alkaloid from *Aconitum coreanum*.

Tang, Q. *et al.*, *Yaoxue Xuebao*, 2005, **40**, 640-643 (*isol*)

6,9,13-Trihydroxyhetisan-2-one T-551C₂₀H₂₅NO₄ 343.422**(13S)-form****13-Ac: Delnuttaline**


[153331-26-1]

C₂₂H₂₇NO₅ 385.459

Minor alkaloid from *Delphinium nuttalianum* (Ranunculaceae). Cryst. Mp 269-271°.

Parvez, M. *et al.*, *Acta Cryst. C*, 1993, **49**, 1871-1873 (*cryst struct*)

Bai, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1717-1724 (*isol, ir, pmr, cmr*)

6,11,13-Trihydroxyhetisan-2-one T-552C₂₀H₂₅NO₄ 343.422**(11 α ,13R)-form****Delbidine**

[123497-82-5]

Alkaloid from a *Delphinium occidentale* - *Delphinium barbeyi* hybrid (Ranunculaceae). Cubes (MeOH). Mp 360°. $[\alpha]_D^{25}$ +22.3 (c, 0.268 in MeOH).

13-Ac: Geyeridine. 6-Hydroxy-13-O-acetylhetisinone

[100045-29-2]

C₂₂H₂₇NO₅ 385.459

Minor alkaloid from the flowers, small stems and leaves of *Delphinium geyeri* (Ranunculaceae). Gum.

11-O-(2 ξ -Methylbutanoyl): Geyerine

[100045-28-1]


C₂₅H₃₃NO₅ 427.539

A major alkaloid from the flowers, small stems and leaves of *Delphinium geyeri* (Ranunculaceae). Glassy amorph. solid which turns translucent on standing. $[\alpha]_D^{25}$ +9.6 (c, 0.36 in EtOH).

Grina, J.A. *et al.*, *J.O.C.*, 1986, **51**, 390

(*Geyerine, Geyeridine*)

Joshi, B.S. *et al.*, *Phytochemistry*, 1989, **28**, 1561 (*Delbidine*)

1,2,7-Trihydroxy-5-(3-hydroxybutyl)-3-(hydroxymethyl)-1H-pyrrolizidine T-553(1R,2R,3R,3' ξ ,5R,7R,7aR)-formC₁₂H₂₃NO₅ 261.317**(1R,2R,3R,3' ξ ,5R,7R,7aR)-form 5-(3-Hydroxybutyl)-7-epiaustraline** [710948-64-4]

Alkaloid from the bulbs of *Scilla peruviana*. Glucosidase inhibitor. $[\alpha]_D$ -20.3 (c, 1.35 in H₂O).

7-Deoxy: 1,2-Dihydroxy-5-(3-hydroxybutyl)-3-(hydroxymethyl)-1H-pyrrolizidine. 5-(3-Hydroxybutyl)hyacinthacine

A₂

[710948-65-5]

C₁₂H₂₃NO₄ 245.318

Alkaloid from the bulbs of *Scilla socialis*. Syrup. $[\alpha]_D$ +1.2 (c, 0.25 in H₂O).

(1R*,2S*,3S*,3' ξ ,5S*,7aS*)-form

7-Deoxy: 5-(3-Hydroxybutyl)hyacinthacine A₁

C₁₂H₂₃NO₄ 245.318

Alkaloid from the bulbs of *Scilla peruviana*. Syrup. $[\alpha]_D$ -20.3 (c, 1.35 in H₂O).

7-Deoxy, 1' ξ -hydroxy: 5-(1,3-Dihydroxybutyl)-1,2-dihydroxy-3-(hydroxymethyl)-1H-pyrrolizidine. 5-(1,3-Dihydroxybutyl)hyacinthacine A₁

[710948-66-6]

C₁₂H₂₃NO₅ 261.317

Alkaloid from the bulbs of *Scilla peruviana* and *Scilla socialis*. Glucosidase inhibitor. Powder. $[\alpha]_D$ +42 (c, 1 in H₂O).

7-Deoxy, 1' ξ ,4'-dihydroxy: 1,2-Dihydroxy-3-(hydroxymethyl)-5-(1,3,4-trihydroxybutyl)-1H-pyrrolizidine. 5-(1,3,4-Trihydroxybutyl)hyacinthacine A₁

[710948-67-7]


C₁₂H₂₃NO₆ 277.317

Alkaloid from the bulbs of *Scilla peruviana*. Powder. $[\alpha]_D$ +33.4 (c, 0.67 in H₂O).

Asano, N. *et al.*, *J. Nat. Prod.*, 2004, **67**, 846-850; 2007, **70**, 993-997 (*isol, pmr, cmr*)

1,2,7-Trihydroxy-3-hydroxy-methyl-5-methyl-1H-pyrrolizidine T-554

Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizidine-1,2,7-triol, 9CI



(1R*,2R*,3R*,5R*,7S*,7aR*)-form

C₉H₁₇NO₄ 203.238**(1R*,2R*,3R*,5R*,7S*,7aR*)-form****Hyacinthacine B₄**

[479348-19-1]

Alkaloid from the bulbs of *Scilla sibirica* and *Scilla socialis*. Powder. $[\alpha]_D^{25}$ -6.7 (c, 1.2 in H₂O).**(1R*,2R*,3S*,5S*,7R*,7aS*)-form****Hyacinthacine B₆**

[479348-21-5]

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. $[\alpha]_D^{25}$ -61.2 (c, 1 in H₂O).**(1R*,2S*,3S*,5R*,7S*,7aS*)-form****Hyacinthacine B₇**


[944408-24-6]

Alkaloid from the bulbs of *Scilla socialis*. Powder. $[\alpha]_D^{25}$ -4.4 (c, 0.2 in H₂O).**(1R*,2S*,3S*,5S*,7R*,7aS*)-form****Hyacinthacine B₅**

[479348-20-4]

Alkaloid from the bulbs of *Scilla sibirica*. Syrup. $[\alpha]_D^{25}$ -25.4 (c, 0.26 in H₂O).**(1R*,2S*,3S*,5S*,7S*,7aS*)-form****Hyacinthacine B₃**

[268209-91-2]

Alkaloid from the bulbs of *Muscari armeniacum*, *Scilla sibirica* and *Scilla socialis*. Glycosidase inhibitor. $[\alpha]_D^{25}$ +3.1 (c, 0.33 in H₂O).Asano, N. *et al.*, *Tetrahedron: Asymmetry*, 2000, **11**, 1-8 (isol, pmr, cmr)Yamashita, T. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1875-1881 (isol, pmr, cmr)Kato, A. *et al.*, *J. Nat. Prod.*, 2007, **70**, 993-997 (*Hyacinthacine B₇*)**1,2,7-Trihydroxy-3-(hydroxymethyl)-1H-pyrrolizidine T-555****Hexahydro-3-(hydroxymethyl)-1H-pyrrolizine-1,2,7-triol, 9CI**

(1R,2R,3R,7R,7aR)-form

C₈H₁₅NO₄ 189.211**(1R,2R,3R,7R,7aR)-form****7-Epiaustraline. 7,7a-Diepiealexine. 7,8-Diepiealexine**

[126655-21-8]

[126655-01-4]

Weak inhibitor of glucosidase activity. Cryst. Mp 193-194°. $[\alpha]_D^{25}$ -13 (c, 0.55 in H₂O). Synthetic. This config. originally assigned to a nat. alkaloid but was shown to be the same as Australine (2003).**(1R,2R,3R,7R,7aS)-form****7-Epiealexine**

[121588-45-2]

Synthetic. Syrup. $[\alpha]_D^{25}$ -11 (c, 1 in H₂O).**(1R,2R,3R,7S,7aR)-form****Australine. 8-Epiealexine. 7a-Epiealexine**

[118396-02-4]

Alkaloid from seeds of *Castanospermum australe*. Potent and specific inhibitor ofamyloglucosidase. Prisms (Me₂CO). Sol. H₂O, MeOH. Mp 148-149°. $[\alpha]_D^{26}$ +19.3 (c, 2.09 in MeOH).**l'-Carboxylic acid: 7a-Epiealexiflorine. 8-Epiealexiflorine. Australiflorine**C₈H₁₃NO₅ 203.194Alkaloid from leaves of *Alexa grandiflora* (Fabaceae).**(1R,2R,3R,7S,7aS)-form****Alexine. 7a-Epiaustraline**

[116174-63-1]

Alkaloid from the pods of *Alexa leiopetala*. Weak inhibitor of glucosidase activity. Cryst. (EtOH aq.). Sol. MeOH, H₂O. Mp 162-163°. $[\alpha]_D^{25}$ +40.4 (c, 0.3 in H₂O).**(1R,2R,3S,7S,7aR)-form****3-Epiaustraline. 3,7a-Diepiealexine. 3,8-Diepiealexine**

[119065-82-6]

Alkaloid from the seeds of *Castanospermum australe*. Weak inhibitor of glycosidase activity. Oil; cryst. (EtOH aq.) (as hydrochloride). Mp 148-152° (hydrochloride). $[\alpha]_D^{20}$ -3.5 (c, 1.35 in H₂O) (hydrochloride).**(1R,2R,3S,7S,7aS)-form [121605-56-9]**Synthetic. Mp 190-192°. $[\alpha]_D^{20}$ +63.8 (c, 0.11 in H₂O).**(1R,2S,3S,7R,7aR)-form****2,3,7-Triepiaustraline**

[529496-76-2]

Alkaloid from the seeds of *Castanospermum australe*. $[\alpha]_D^{25}$ +59.7 (c, 0.58 in H₂O).**(1R,2S,3S,7S,7aR)-form****2,3-Diepieastraline**

[529496-75-1]


Alkaloid from the seeds of *Castanospermum australe*. $[\alpha]_D^{25}$ +38.2 (c, 0.51 in H₂O).**(1S,2R,3R,7S,7aR)-form****1-Epiaustraline. 1,7a-Diepiealexine. 1,8-Diepiealexine**

[126594-77-2]

[126654-99-7]

Alkaloid from the seeds of *Castanospermum australe*. Weak inhibitor of glycosidase activity. Oil or hygroscopic solid. Mp 28-30°. $[\alpha]_D^{20}$ +13.3 (c, 0.1 in H₂O). $[\alpha]_D^{20}$ -3 (c, 0.4 in H₂O) (as hydrochloride).**2-O-β-D-Glucopyranoside: 1-Epiaustraline 2-glucoside**


[528886-31-9]

C₁₄H₂₅NO₉ 351.353Alkaloid from the seeds of *Castanospermum australe*. $[\alpha]_D^{25}$ +35.8 (c, 0.42 in H₂O).Molyneux, R.J. *et al.*, *J. Nat. Prod.*, 1988, **51**, 1198 (*Australine*)Nash, R.J. *et al.*, *Tet. Lett.*, 1988, **29**, 2487 (isol, pmr, cmr, ms, cryst struct, abs config)Fleet, G.W.J. *et al.*, *Tet. Lett.*, 1988, **29**, 5441 (synth, cmr)Nash, R.J. *et al.*, *Tetrahedron*, 1988, **44**, 5959 (3,8-Diepiealexine)Tropea, J.E. *et al.*, *Biochemistry*, 1989, **28**, 2027 (pharmacol)Nash, R.J. *et al.*, *Phytochemistry*, 1990, **29**, 111 (1,8-Diepiealexine, 7,8-Diepiealexine)Pearson, W.H. *et al.*, *Tet. Lett.*, 1991, **32**, 5513 (7,7a-Diepiealexine, synth)Choi, S. *et al.*, *Tet. Lett.*, 1991, **32**, 5517 (1,7a-Diepiealexine, synth)Pereira, A.C. de S. *et al.*, *Tetrahedron*, 1991, **47**, 5637 (8-Epiealexiflorine)Taylor, D.L. *et al.*, *Antiviral Chem. Chemother.*, 1992, **3**, 273 (7,8-Diepiealexine, anti-HIV activity)Furneaux, R.H. *et al.*, *Tetrahedron*, 1994, **50**, 2131 (*Australine*, synth)Denmark, S.E. *et al.*, *J.A.C.S.*, 1998, **120**, 7357-7358 (synth)White, J.D. *et al.*, *J.A.C.S.*, 1998, **120**, 7359-7360 (synth)Ikota, N. *et al.*, *Tetrahedron*, 1998, **54**, 8985-8998 (synth)Pearson, W.H. *et al.*, *J.O.C.*, 2000, **65**, 5785-5793 (synth)Romero, A. *et al.*, *J.O.C.*, 2000, **65**, 8264-8268 (synth)White, J.D. *et al.*, *J.O.C.*, 2000, **65**, 9129-9142 (synth)Yoda, H. *et al.*, *Tet. Lett.*, 2000, **41**, 7661-7665 (synth)Denmark, S.E. *et al.*, *J.O.C.*, 2001, **66**, 4276-4284 (1-Epiaustraline, synth)Tang, M. *et al.*, *J.O.C.*, 2003, **68**, 7818-7824 (7-Epiaustraline, 1,7-Diepieastraline, synth)Kato, A. *et al.*, *Tetrahedron: Asymmetry*, 2003, **14**, 325-331 (synth, cmr, abs config)Donohoe, T.J. *et al.*, *J.O.C.*, 2005, **70**, 7297-7304 (1-Epiaustraline, synth)Lauritsen, A. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 2898-2905 (*Australine*, synth)Donohoe, T.J. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 3896-3898 (7-Epiealexine, synth)Takahashi, M. *et al.*, *Tetrahedron*, 2008, **64**, 5254-5261 (*Alexine*, synth)**3,5,15-Trihydroxy-6(17),12-lathyradien-14-one T-556**

(2α,3β,5α,12E,15β)-form

C₂₀H₃₀O₄ 334.455**(2β,3β,5β,12E,15β)-form****Lathyrol**

[34420-19-4]

Cryst. Mp 168-169°. $[\alpha]_D^{25}$ +101 (c, 1.3 in MeOH).**(3-Pyridinecarbonyl), di-Ac:**C₃₀H₃₇NO₇ 523.625Constit. of *Euphorbia lathyris*. Position of esters not established.Adolf, W. *et al.*, *Experientia*, 1971, **27**, 1393-1394 (*Euphorbia lathyris* constits)**3,4,13-Trihydroxylupanine T-557**

(3β,4α,13α)-form

C₁₅H₂₄N₂O₄ 296.366**(3β,4α,13α)-form**Alkaloid from the leaves of *Calpurnia aurea* ssp. *aurea* (Fabaceae).

13-O-(2-Pyrrolicarbonyl): *Calpaurine*
[103847-10-5]
C₂₀H₂₇N₃O₅ 389.45
Alkaloid from the leaves of *Calpurnia aurea* ssp. *aurea* (Fabaceae).

(3β,4α,13β)-form

13-Me ether: *3,4-Dihydroxy-13-methoxylupanine*
[193559-15-8]
C₁₆H₂₆N₃O₄ 310.392
Minor alkaloid from bark of *Acosmium panamense*.


13-Me ether, 4-angeloyl: *4-Angeloyloxy-3-hydroxy-13-methoxylupanine*
C₂₁H₃₂N₃O₅ 392.494
Alkaloid from the seeds of *Acosmium panamense*. Amorph. yellow solid. [α]_D²⁵ -72 (c, 0.57 in CHCl₃).

Asres, K. et al., *Phytochemistry*, 1986, **25**, 1443
(*Trihydroxylupanine, Calpaurine, isol, ir, cmr, ms, struct*)

Mascagni, P. et al., *Tetrahedron*, 1987, **43**, 149
(*Calpaurine, pmr, cmr, struct*)

Veitch, N.C. et al., *Phytochemistry*, 1997, **45**, 847-850 (*3,4-Dihydroxy-13-methoxylupanine*)

Nuzillard, J.-M. et al., *Tetrahedron*, 1999, **55**, 11511-11518
(*Angeloyloxyhydroxymethoxylupanine*)

3,8,13-Trihydroxylupanine T-558

C₁₅H₂₄N₂O₄ 296.366

(3β,8α,13α)-form [138680-24-7]

Alkaloid from the leaves of *Pearsonia cajanifolia* ssp. *cryptantha* (Fabaceae). [α]_D²² +2 (c, 2.2 in CHCl₃).

13-Angeloyl: *Pearsonine*

[126420-99-3]
C₂₀H₃₀N₂O₅ 378.467


Alkaloid from *Pearsonia cajanifolia* ssp. *cryptantha* (Fabaceae). Mp 93-96°. [α]_D²² +7 (c, 0.6 in CHCl₃).

Verdoorn, G.H. et al., *Phytochemistry*, 1990, **29**, 1297 (*Pearsonine*)

Verdoorn, G.H. et al., *Phytochemistry*, 1991, **30**, 3631 (*isol*)

1,6,7-Trihydroxy-3-methyl-carbazole T-559

3-Methyl-9H-carbazole-1,6,7-triol




C₁₃H₁₁NO₃ 229.235

6,7-Di-Me ether: *6,7-Dimethoxy-3-methyl-9H-carbazol-1-ol. 1-Hydroxy-6,7-dimethoxy-3-methylcarbazole*
[211106-13-7]
C₁₅H₁₅NO₃ 257.288

Alkaloid from the leaves of *Murraya koenigii*. Antibacterial agent. Needles (C₈H₆). Mp 216°. λ_{max} 235 (log ε 4.65); 285 (log ε 4.18); 305 (log ε 4.32); 330 (log ε 3.5) (MeOH).

Saha, C. et al., *Phytochemistry*, 1998, **48**, 363-366 (*synth*)

Chowdhury, B.K. et al., *Indian J. Chem., Sect. B*, 2001, **40**, 490-494 (*isol, pmr, cmr*)

3,6,7-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI T-560

C₁₃H₉NO₄ 243.218

3,6-Di-Me ether: *7-Hydroxy-3,6-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 7-Hydroxy-2,8-dimethoxy-1-methyl-4-aza-9-fluorenone. 7-Hydroxy-2,8-dimethoxyonychine*
[161196-98-1]
C₁₅H₁₃NO₄ 271.272

Alkaloid from stem bark of *Piptostigma fugax* (Annonaceae). Orange-red needles. Mp 206°.

Achenbach, H. et al., *Phytochemistry*, 1995, **38**, 1037 (*7-Hydroxy-2,8-dimethoxyonychine*)

3,7,8-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one T-561

C₁₃H₉NO₄ 243.218

3,7-Di-Me ether: *8-Hydroxy-3,7-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 6-Hydroxy-2,7-dimethoxy-1-methyl-4-azafluorenone. 6-Hydroxy-2,7-dimethoxyonychine. Oxylpidine*
[101899-52-9]
C₁₅H₁₃NO₄ 271.272

Alkaloid from stem bark and twigs of *Oxandra xylopioides* (Annonaceae). Mp 271-273°. Struct. revised in 1994. Orig. descr. as 7-Hydroxy-2,6-dimethoxyonychine.

Zhang, J. et al., *J. Nat. Prod.*, 1987, **50**, 800-806 (*isol*)

Achenbach, H. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1994, **327**, 755 (*synth, struct, Oxylpidine*)

3,8,9-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one T-562

C₁₃H₉NO₄ 243.218

3,8-Di-Me-ether: *9-Hydroxy-3,8-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 5-Hydroxy-2,6-dimethoxy-1-methyl-4-aza-9-fluorenone. 2,6-Dimethoxy-5-hydroxyonychine*
[126660-44-4]
C₁₅H₁₃NO₄ 271.272

Alkaloid from the stem bark of *Alphonsea mollis* (Annonaceae). Mp 211-

212°.

Xie, N. et al., *Zhongguo Yaokexue Xuebao*, 1989, **20**, 321; *CA*, **112**, 195252

Guinaudeau, H. et al., *J. Nat. Prod.*, 1994, **57**, 1033 (*rev*)

6,7,8-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one T-563

6,7,8-Trihydroxy-1-methyl-4-azafluorenone

C₁₃H₉NO₄ 243.218

6-Me ether: *7,8-Dihydroxy-6-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one.*

Penduline†

C₁₄H₁₁NO₄ 257.245

Alkaloid from the roots of *Polyalthia longifolia* var. *pendula*. Orange cryst. Mp 188-189°. λ_{max} 205 ; 261 ; 349 (MeOH).

Faizi, S. et al., *Planta Med.*, 2003, **69**, 350-355 (*isol, pmr, cmr, ms*)

6,8,9-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one T-564

5,6,8-Trihydroxy-1-methyl-4-azafluorenone

C₁₃H₉NO₄ 243.218

6,9-Di-Me ether: *8-Hydroxy-6,9-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 6-Hydroxy-5,8-dimethoxy-1-methyl-4-azafluorenone. 6-Hydroxy-5,8-dimethoxyonychine. Kinabaline*
[109028-37-7]
C₁₅H₁₃NO₄ 271.272

Trace alkaloid from the trunk bark of *Meiogyne virgata* (Annonaceae). Amorph. yellow solid. λ_{max} 208 (log ε 3.96); 222 (log ε 3.91); 231 (log ε 3.93); 246 (sh) (log ε 4.01); 254 (log ε 4.12); 280 (sh) (log ε 3.72); 292 (log ε 3.77); 304 (log ε 3.73); 388 (log ε 3.32) (MeOH).

Tadč, D. et al., *Phytochemistry*, 1987, **26**, 537-541 (*isol, uv, pmr, ms, struct*)

7,8,9-Trihydroxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one T-565

C₁₃H₉NO₄ 243.218

8-Me ether: *7,9-Dihydroxy-8-methoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 5,7-Dihydroxy-6-methoxy-1-methyl-4-azafluorenone. 5,7-Dihydroxy-6-methoxyonychine. Cyathocaline*
[164176-16-3]
C₁₄H₁₁NO₄ 257.245

Alkaloid from stem bark of *Cyathocalyx zeylanica* (Annonaceae). Orange cryst. Mp 222-224°. λ_{max} 232 (ε 2630); 274 (ε 3980); 354 (ε 2190); 448 (ε 2000) (EtOH/NaOH) (Derep). λ_{max} 224 (ε 2510); 261 (ε 6030); 313 (ε 1910); 348 (ε 1820) (EtOH) (Derep). λ_{max} 224 (ε 2754); 261 (ε 6025); 313 (ε 1905); 348 (ε 1820) (EtOH) (Berdy). λ_{max} 232 (ε 2690); 274 (ε 2187); 354 (ε 3980); 448 (ε 1995) (EtOH-NaOH) (Berdy).

8-Me ether, di-Ac: Mp 107-108°.

7,8-Di-Me ether: 9-Hydroxy-7,8-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one, 9CI. 5-Hydroxy-6,7-dimethoxy-1-methyl-4-azafloren-9-one. 5-Hydroxy-6,7-dimethoxyonychine [250597-10-5]

C₁₅H₁₃NO₄ 271.272

Alkaloid from *Alphonsea monogyna*. Yellow needles. Mp 224-226°. λ_{max} 206 (log ε 4.09); 235 (log ε 4.03); 266 (log ε 4.33); 292 (log ε 4.02) (no solvent reported).

7,8-Di-Me ether, N-oxide: 5-Hydroxy-6,7-dimethoxyonychine N-oxide
C₁₅H₁₃NO₅ 287.271

Alkaloid from *Alphonsea monogyna*. Orange-red needles. Mp 287-288°. λ_{max} 215 (log ε 4.09); 240 (log ε 4.03); 272 (log ε 4.33); 299 (log ε 4.02); 310 (sh) (log ε 4) (no solvent reported).

8,9-Di-Me ether: 7-Hydroxy-8,9-dimethoxy-4-methyl-5H-indeno[1,2-b]pyridin-5-one. 7-Hydroxy-5,6-dimethoxy-1-methyl-4-azafloren-9-one. 7-Hydroxy-5,6-dimethoxyonychine.

Darientine
[111316-27-9]

C₁₅H₁₃NO₄ 271.272

Alkaloid from the stem bark of *Oxandra* cf. *major* (Annonaceae). Amorph. yellow solid. λ_{max} 206 (log ε 4.09); 235 (log ε 4.03); 266 (log ε 4.33); 292 (log ε 4.02); 302 (sh) (log ε 4) (EtOH). λ_{max} 218 (log ε 4.05); 226 (sh) (log ε 4.04); 285 (log ε 4.18); 322 (log ε 4.12); 360 (log ε 3.78) (EtOH/MeOH). λ_{max} 206 (log ε 4.09); 226 (log ε 4.03); 265 (log ε 4.27); 292 (log ε 3.96); 302 (sh) (log ε 3.95) (EtOH/HCl).

8,9-Di-Me ether, 7-Ac: Mp 143°.

Arango, G.J. et al., *Phytochemistry*, 1987, **26**, 2093-2098 (*Darientine*)

Wijeratne, E.M.K. et al., *J. Nat. Prod.*, 1995, **58**, 459-462 (*Cyathocaline*)


Yang, N. et al., *CA*, 1999, **131**, 349043j (5-Hydroxy-6,7-dimethoxyonychine)

Xie, N. et al., *Chin. Chem. Lett.*, 1999, **10**, 671-672 (7,8-di-Me ether)

Yang, N.Y. et al., *Chin. Chem. Lett.*, 2000, **11**, 409-410 (7,8-di-Me ether N-oxide)

1,6,8-Trihydroxy-3-methylisoquinoline T-566

6,8-Dihydroxy-3-methyl-1(2H)-isoquinolinone, 9CI. 3-Methyl-1,6,8-isoquinoline-triol. 6,8-Dihydroxy-3-methylisocarboxystyryl. **Siamine** [60352-12-7]




C₁₀H₉NO₃ 191.186

Alkaloid from the seeds and leaves of *Cassia siamea* (Fabaceae). Cryst. (MeOH). Mp 264-268°.

Ahn, B.Z. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1978, **311**, 569 (isol, uv, ir, pmr, cmr, struct)

2,3,5-Trihydroxy-6-methylpyridine T-567

3,5-Dihydroxy-6-methyl-2(1H)-pyridinone. 6-Methyl-2,3,5-pyridinetriol [95508-57-9]




C₆H₇NO₃ 141.126

Tentative struct. assigned. Prod. by *Streptovercillium hiroshimensense*. Dopamine β-hydroxylase inhibitor, antihypertensive agent. Cryst. (H₂O). Sol. H₂O; fairly sol. MeOH; poorly sol. hexane. λ_{max} 272 (HCl) (Berdy). λ_{max} 226; 308 (NaOH) (Berdy).

U.S. Pat., 1984, 4 487 761; *CA*, **102**, 130458r

3,4,6-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid T-568



C₁₅H₉NO₇ 315.239

4-Me ether, Me ester: Methyl 3,6-dihydroxy-4-methoxy-10-nitro-1-phenanthrenecarboxylate. **Ariskanin B** [157207-61-9]

C₁₇H₁₃NO₇ 343.292

From roots and stems of *Aristolochia kankauensis* (Aristolochiaceae). Yellowish needles (CHCl₃). Mp 274-276°. λ_{max} 211 (ε 27542); 243 (ε 38884); 253 (ε 38904); 260 (ε 38904); 288 (ε 15135); 365 (ε 7244); 386 (ε 8128) (MeOH) (Berdy).

3,4-Di-Me ether, Me ester: Methyl 6-hydroxy-3,4-dimethoxy-10-nitro-1-phenanthrenecarboxylate. **Ariskanin C** [157207-62-0]

C₁₈H₁₅NO₇ 357.319

From roots and stems of *Aristolochia kankauensis* (Aristolochiaceae). Orange needles (CHCl₃/MeOH). Mp 186-190°. λ_{max} 253 (ε 33884); 289 (ε 12590); 363 (ε 7586); 383 (ε 9550); 393 (ε 7586) (MeOH) (Berdy).

Methylene ether: see 6-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, H-600

Wu, T.S. et al., *Phytochemistry*, 1994, **36**, 1063-1068 (isol, pmr)

Lou, F.C. et al., *Yaoxue Xuebao*, 1995, **30**, 588; *CA*, **123**, 251411n

3,4,8-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid T-569

C₁₅H₉NO₇ 315.239

4,8-Di-Me ether, Me ester: Methyl 3-hydroxy-4,8-dimethoxy-10-nitro-1-phenanthrenecarboxylate. **Ariskanin D**

[157207-63-1]

C₁₈H₁₅NO₇ 357.319

From roots and stems of *Aristolochia kankauensis* (Aristolochiaceae). Yellowish needles (CHCl₃/MeOH). Mp 145-148°. λ_{max} 239 (ε 39810); 257 (ε 38020); 279 (ε 21380); 308 (ε 13182); 390 (ε 9772) (MeOH) (Berdy).

Tri-Me ether, Me ester: Methyl 3,4,8-trimethoxy-10-nitro-1-phenanthrenecarboxylate. **Ariskanin E**

[157207-64-2]

C₁₉H₁₇NO₇ 371.346

From roots and stems of *Aristolochia kankauensis* (Aristolochiaceae). Yellowish needles (CHCl₃/MeOH). Mp 158-160°.

3,4-Methylene, 8-Me ether: see 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262

Wu, T.S. et al., *Phytochemistry*, 1994, **36**, 1063

Lou, F.C. et al., *Yaoxue Xuebao*, 1995, **30**, 588; *CA*, **123**, 251411n

3,4,9-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid T-570

C₁₅H₉NO₇ 315.239

3,4-Methylene, 9-Me ether: 3,4-Methylenedioxy-9-methoxy-10-nitro-1-phenanthrenecarboxylic acid. 9-Methoxyaristolochic acid II


[134306-18-6]

C₁₇H₁₁NO₇ 341.276

Alkaloid from *Aristolochia ponticum* (Aristolochiaceae).

Houghton, P.J. et al., *Phytochemistry*, 1991, **30**, 717

1,2,3-Trihydroxy-5-oxonoraporphine T-571



C₁₆H₁₃NO₄ 283.283

(R)-form

1,2-Di-Me ether: 3-Hydroxy-1,2-dimethoxy-5-oxonoraporphine

C₁₈H₁₇NO₄ 311.337

Alkaloid from the bark of *Mitrephora* cf. *maingayi*. Light straw-coloured needles (CHCl₃). Mp 160-162°. [α]_D²⁵ -52 (c, 0.07 in CHCl₃). λ_{max} 228 (log ε 4.08); 282 (log ε 4.08); 320 (log ε 2.85) (MeOH).


Tri-Me ether: 1,2,3-Trimethoxy-5-oxonoraporphine

C₁₉H₁₉NO₄ 325.363

Alkaloid from the bark of *Mitrephora* cf. *maingayi*. Yellow needles (CHCl₃). Mp 185-187°. [α]_D²⁵ -31.6 (c, 0.04 in CHCl₃). λ_{max} 228 (log ε 4.1); 274 (log ε

4.04); 316 (log ϵ 3.26) (MeOH).Lee, N.H.S. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1158-1159 (*isol, uv, ir, pmr, cmr, ms*)**4',5,7-Trihydroxy-6-(5-oxo-2-pyrrolidinyl)flavanone** T-572

6-(5-Oxo-2-pyrrolidinyl)naringenin



(2R,2''R)-form

C₁₉H₁₇NO₆ 355.346**(2R,2''R)-form****Dracocephalin A₄**

[1023993-43-2]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: 3',4',5,7-Tetrahydroxy-6-(5-oxo-2-pyrrolidinyl)flavanone. 6-(5-Oxo-2-pyrrolidinyl)eriodictyol. **Dracocephalin C₄**

[1029363-44-7]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.**(2R,2''S)-form****Dracocephalin A₁**

[1029363-26-5]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin C₁**

[1029363-38-9]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.**(2S,2''R)-form****Dracocephalin A₃**

[1029363-32-3]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin C₃**

[1029363-42-5]

C₁₉N₁₇NO₇ 592.325Constit. of *Dracocephalum rupestre*.**(2S,2''S)-form****Dracocephalin A₂**


[1029363-29-8]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin C₂**

[1029363-40-3]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.Ren, D.-M. *et al.*, *Phytochemistry*, 2008, **69**, 1425-1433 (*isol, cd, pmr, cmr, crystal*)**4',5,7-Trihydroxy-8-(5-oxo-2-pyrrolidinyl)flavanone** T-573

8-(5-Oxo-2-pyrrolidinyl)naringenin



(2R,2''R)-form

C₁₉H₁₇NO₆ 355.346**(2R,2''R)-form****Dracocephalin B₃**

[1029363-36-7]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: 3',4',5,7-Tetrahydroxy-8-(5-oxo-2-pyrrolidinyl)flavanone. 8-(5-Oxo-2-pyrrolidinyl)eriodictyol. **Dracocephalin D₃**

[1029363-50-5]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.**(2R,2''S)-form****Dracocephalin B₂**

[1029363-34-5]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin D₂**

[1029363-48-1]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.**(2S,2''R)-form****Dracocephalin B₄**

[1023993-45-4]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin D₄**

[1029363-52-7]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.**(2S,2''S)-form****Dracocephalin B₁**


[1023993-44-3]

Constit. of *Dracocephalum rupestre*.3'-Hydroxy: **Dracocephalin D₁**


[1029363-46-9]

C₁₉H₁₇NO₇ 371.346Constit. of *Dracocephalum rupestre*.Ren, D.-M. *et al.*, *Phytochemistry*, 2008, **69**, 1425-1433 (*isol, cd, pmr, cmr*)**2,3,9-Trihydroxy-1-phenazinicarboxylic acid** T-574


[128723-47-7]

C₁₃H₈N₂O₅ 272.217Isol. from *Pseudomonas fluorescens* grown under beryllium stress. No phys. props. reported.Taraz, K. *et al.*, *Z. Naturforsch., B*, 1990, **45**, 552 (*isol, pmr, cmr, ms, struct*)**2,3,7-Trihydroxy-1,6-phenazinedicarboxylic acid, 9CI** T-575

[71670-86-5]

C₁₄H₈N₂O₇ 316.226Isol. from *Pseudomonas aureofaciens*.Römer, A. *et al.*, *Tet. Lett.*, 1979, 509 (*pmr, struct*)**3,4,5-Trihydroxyphenethylamine** T-576

5-(2-Aminoethyl)-1,2,3-benzenetriol, 9CI. 5-(2-Aminoethyl)pyrogallol, 8CI. 2-(3,4,5-Trihydroxyphenyl)ethylamine. 5-Hydroxydopamine [1927-04-4]

C₈H₁₁NO₃ 169.18Alkaloid from *Acacia rigidula*.

Hydrochloride: [5720-26-3]

Cryst. (MeOH/EtOAc). Mp 215-219°.

Hydrobromide: [20555-57-1]

Mp 192-196°.

Picrate:

Cryst. Mp 193-194°.

N-Me: 5-[2-(Methylamino)ethyl]-1,2,3-benzenetriol, 9CI. N-Methyl-3,4,5-trihydroxyphenethylamine [73917-91-6]

C₉H₁₃NO₃ 183.207Alkaloid from *Acacia rigidula*.3'-Me ether: **3,4-Dihydroxy-5-methoxyphenethylamine**. 5-(2-Aminoethyl)-3-methoxy-1,2-benzenediol, 9CI

[16032-86-3]

C₉H₁₃NO₃ 183.207Alkaloid present in *Lophophora williamsii* (Cactaceae). Major intermed. in biosynth. of peyote phenethylamine and tetrahydroisoquinoline alkaloids.

3',4'-Di-Me ether: see 3-Hydroxy-4,5-dimethoxyphenethylamine, H-465

3',5'-Di-Me ether: **4-Hydroxy-3,5-dimethoxyphenethylamine**. 4-(2-Aminoethyl)-2,6-dimethoxyphenol, 9CI. 3,5-Dimethoxytyramine [2413-00-5]C₁₀H₁₅NO₃ 197.233Alkaloid present in *Acacia rigidula*, *Trichocereus werdermannianus*, *Trichocereus pachanoi* and *Lophophora williamsii* (mescal) (Cactaceae). Cryst. Mp 180-181° (152-154°).

3',5'-Di-Me ether, hydrochloride: [2176-14-9]

Cryst. Mp 258-259°.

3',5'-Di-Me ether, N-(4-hydroxy-3-methoxy-E-cinnamoyl): N-trans-Feruloyl-3,5-dimethoxytyramine. **Hibiscusamide** [925211-84-3]C₂₀H₂₃NO₆ 373.405Alkaloid from the stem wood of *Hibiscus tiliaceus* and from *Alternanthera philoxeroides*.Cytotoxic. Needles (EtOH) or yellow solid. Mp 126-128°. [α]_D²⁵ -18 (c, 0.68 in MeOH). λ_{\max} 220 (log ϵ 3.87); 288 (log ϵ 3.76); 316 (log ϵ 3.79) (MeOH). λ_{\max} 232 (log ϵ 4.13); 295 (log ϵ 3.98); 319 (log ϵ 4.09) (MeOH).


Tri-Me ether: see Mescaline, M-219

[5090-25-5 , 2176-14-9]

Hahn, G. *et al.*, *Ber.*, B, 1938, **71**, 2141 (*synth*)
 Agurell, S. *et al.*, *J. Nat. Prod.*, 1969, **32**, 40
 (*isol*, *synth*)
 Lunstrom, J. *et al.*, *Acta Chem. Scand.*, 1971,
25, 3489 (*occur*, *deriv*)
 Andersen, A.M. *et al.*, *Acta Chem. Scand.*,
 1972, **26**, 2670-2680 (*cryst struct*)
 Short, J.H. *et al.*, *Tetrahedron*, 1973, **29**, 1931-
 1939 (*synth*)
 Pardani, J.H. *et al.*, *J. Nat. Prod.*, 1977, **40**,
 585-590 (*isol*, *deriv*)
 Basmaljian, G.P. *et al.*, *J. Nat. Prod.*, 1978, **41**,
 375
 Severin, T. *et al.*, *Chem. Ber.*, 1980, **113**, 970-
 978 (*synth*, *deriv*, *pmr*, *ms*)
 Battersby, A.R. *et al.*, *J.C.S. Perkin 1*, 1981,
 2016-2029 (*synth*, *pmr*)
 Ma, W.W. *et al.*, *J. Nat. Prod.*, 1986, **49**, 735-
 737 (*isol*, *deriv*)
 Clement, B.A. *et al.*, *Phytochemistry*, 1998, **49**,
 1377-1380 (*isol*)
 Chen, J.-J. *et al.*, *Planta Med.*, 2006, **72**, 932-
 935 (*Hibiscusamide*)
 Fang, J.-B. *et al.*, *J. Asian Nat. Prod. Res.*,
 2007, **9**, 511-515 (*N-*
Feruloyldimethoxytyramine)

3-(2,4,5-Trihydroxyphenyl)- 2-propenoic acid T-577

3-(2,4,5-Trihydroxyphenyl)acrylic acid.
 2,4,5-Trihydroxycinnamic acid. *Aesculetic*
acid. *Esculetic acid*
 [56437-15-1]

C₉H₈O₅ 196.159


(E)-form

Tri-Me ether, pyrrolidide: N-(2,4,5-Tri-
 methoxycinnamoyl)pyrrolidine. **Sar-**
mentamide C
 C₁₆H₂₁NO₄ 291.346
 Constit. of the roots of *Piper sarment-*
osum. Amorph. solid. Mp 159-162°. λ_{\max}
 203 (log ϵ 4.25); 230 (log ϵ 4.29);
 305 (log ϵ 4.26) (MeOH).

Tuntiwachwuttikul, P. *et al.*, *Chem. Pharm.*
Bull., 2006, **54**, 149-151 (*Sarmentamide C*)

3-(3,4,5-Trihydroxyphenyl)- 2-propenoic acid T-578

3-(3,4,5-Trihydroxyphenyl)acrylic acid.
 3,4,5-Trihydroxycinnamic acid
 [6093-59-0]



(E)-form

C₉H₈O₅ 196.159

Does not appear to have been found in

nature in the free state despite extensive
 searches.

(E)-form

Needles + 1H₂O (H₂O) becoming anhyd.
 at 120°. Mp 207-208° dec. CAS no. not
 found 8-14 Cl.

Tri-Me ether, 3-(3,4,5-trimethoxyph-
enyl)-2-propenoic acid, 9CI. 3,4,5-Tri-
methoxycinnamic acid
 [20329-98-0]

C₁₂H₁₄O₅ 238.24
 Constit. of *Piper tuberculatum*, *Tecoma*
radicans, *Polygala tenuifolia* and
 others. Needles (H₂O). Mp 126.5-127°. λ_{\max}
 222 (log ϵ 4.408); 282 (log ϵ 4.255)
 (MeOH).

Tri-Me ether, 2-methylpropylamide: 3,4,5-
 Trimethoxycinnamic acid isobutyla-
 mide. *N-Isobutyl-3,4,5-trimethoxycin-*
namide
 [10511-98-5]
 C₁₆H₂₃NO₄ 293.362
 Alkaloid from *Piper amalago*.

Tri-Me ether, pyrrolidide: N-(3,4,5-Tri-
 methoxycinnamoyl)pyrrolidine. 3,4,5-
 Trimethoxycinnamic acid pyrrolidide.

Piperlotine C

[1703-35-1]
 C₁₆H₂₁NO₄ 291.346
 Constit. of *Piper amalago* and *Piper*
lolot. Cryst. (H₂O). Mp 158° (148-
 150°). λ_{\max} 231 ; 304 (MeOH).

Tri-Me ether, piperidide: 3,4,5-Tri-
 methoxycinnamoyl piperidide. **Puberul-**
lumine
 [188546-66-9]
 C₁₇H₂₃NO₄ 305.373
 Alkaloid from leaves and stems of
Piper puberulum. Needles. Mp
 94-96°. λ_{\max} 200 ; 224 ; 296
 (MeOH).

3,4-Methylene, 5-Me ether, 2-methylpro-
 pylamide: **N-Isobutyl-3-methoxy-4,5-**
methylenedioxcinnamide. N-(3-Meth-
 oxy-4,5-methylenedioxcinnamoyl)iso-
 butylamine
 [102934-27-0]
 C₁₅H₁₉NO₄ 277.319
 Alkaloid from *Esenbeckia almawilla*
 and *Piper amalago*. Amorph. solid.

3,4-Methylene, 5-Me ether, pyrrolidide:
N-(3-Methoxy-4,5-methylenedioxcin-
namoyl)pyrrolidine
 [102934-26-9]
 C₁₅H₁₇NO₄ 275.304
 Alkaloid from *Piper amalago*. Config.
 not determined, may have contained
 some Z-isomer.

(Z)-form

CAS no. not found to 14 Cl.

3,4,5-Tri-Me ether, pyrrolidide: **Piperlo-**
tine D
 [958296-13-4]
 C₁₆H₂₁NO₄ 291.346
 Alkaloid from the leaves of *Piper lolot*.
 Syrup. λ_{\max} 226 ; 289 (MeOH).

3,4-Methylene, 5-Me ether, piperidide: N-
 (3-Methoxy-4,5-methylenedioxcinma-
 moyl)piperidide
 C₁₆H₁₉NO₄ 289.33


Alkaloid from the roots of *Piper*
nigrum (pepper). Oil.

[54976-67-9]

Achenbach, H. *et al.*, *Planta Med.*, 1986, **52**,
 12-18 (*Piper amalago amides*)
 Wu, Q.-L. *et al.*, *Phytochemistry*, 1997, **44**, 727-
 730 (*Puberullumine*)
 Wei, K. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1005-
 1009 (*Piper nigrum piperidide*)
 Li, C.-Y. *et al.*, *J. Agric. Food Chem.*, 2007, **55**,
 9436-9442 (*Piperlotines C-D*)
 Barros-Filho, B.A. *et al.*, *Quim. Nova*, 2007,
30, 1589-1591 (3,4-methylene 5-Me ether 2-
 methylpropylamide)

3,4,5-Trihydroxy-2-piperidi- necarboxylic acid T-579

3,4,5-Trihydroxypipercolic acid

C₆H₁₁NO₅ 177.157

(2S,3R,4R,5S)-form

2,6-Dideoxy-2,6-imino-L-gulonic acid,
 9CI
 [96861-04-0]
 Isol. from seeds of *Baphia racemosa*
 (Fabaceae). Glucuronidase and iduroni-
 dase inhibitor. Needles (EtOH aq.). Mp
 228-232°. [α]_D +18.5 (c, 1 in H₂O).


(2R,3R,4R,5S)-form

2,6-Dideoxy-2,6-imino-L-idonic acid, 9CI
 [108428-44-0]
 Mp 198-202°. [α]_D²⁰ +34 (c, 0.25 in
 H₂O).

DiBello, I.C. *et al.*, *FEBS Lett.*, 1984, **176**, 61
 (*biochem*)
 Manning, K.S. *et al.*, *Chem. Comm.*, 1985, 127
 (*isol*, *ms*, *pmr*)
 Bernotas, R.C. *et al.*, *Tet. Lett.*, 1985, **26**, 4981
 (*synth*, *cmr*)
 Bashyal, B.P. *et al.*, *Tet. Lett.*, 1986, **27**, 3205
 (*synth*, *cmr*)
 Lee, B.W. *et al.*, *Synthesis*, 2000, 1305-1309
 (2R,3R,4R,5S-form, *synth*, *pmr*, *cmr*)
 Tsimilaza, A. *et al.*, *Tetrahedron: Asymmetry*,
 2007, **18**, 1585-1588 (2S,3R,4R,5S-form,
synth)
 Booth, K.V. *et al.*, *Acta Cryst. E*, 2008, **63**,
 o3783-o3784 (*cryst struct*)

1,3,5-Trihydroxy-2-prenyl- acridone T-580

1,3,5-Trihydroxy-2-(3-methyl-2-butenyl)-
 9(10H)-acridinone, 9CI

C₁₈H₁₇NO₄ 311.337

N-Me: 1,3,5-Trihydroxy-10-methyl-2-pre-
 nylacridone. **Junosine**


[103956-34-9]

C₁₉H₁₉NO₄ 325.363Alkaloid from the bark of *Citrus junos* (yuzu) (Rutaceae). Light yellow prisms. Mp 210-213°.**3-Me ether, N-Me: 1,5-Dihydroxy-10-methyl-2-prenylacridone. Buxifoliadine C**
[263007-67-6]C₁₉H₁₉NO₄ 325.363Alkaloid from the root bark of *Severinia buxifolia*. Yellow needles (Me₂CO). Mp 275-278° dec. λ_{max} 231 (log ε 3.23); 237 (sh) (log ε 3.19); 262 (log ε 3.66); 273 (log ε 3.55); 275 (sh) (log ε 3.53); 283 (log ε 3.63); 394 (log ε 2.74) (MeOH).**5-Me ether, N-Me: 1,3-Dihydroxy-5-methoxy-10-methyl-2-prenylacridone.****Yukomine**

[139219-97-9]

C₂₀H₂₁NO₄ 339.39Alkaloid from *Citrus wilsonii* and *Citrus yuko* (Rutaceae). Yellow needles (CH₂Cl₂). Mp 214-217°.**Δ³-Isomer, 2'ξ-hydroxy, N-Me: 1,3,5-Trihydroxy-2-(2-hydroxy-3-methyl-3-butenyl)-10-methylacridone. Bosistidine**
[176391-58-5]C₁₉H₁₉NO₅ 341.363Alkaloid from stem bark of *Bosistoa transversa* (Rutaceae). Amorph. [α]_D -15.1 (c, 0.0018 in CHCl₃).Ju-ichi, M. *et al.*, *Heterocycles*, 1986, **24**, 1595-1597; 1991, **32**, 1781-1784 (*Junosine, Yukomine*)Auzi, A.A. *et al.*, *Phytochemistry*, 1996, **42**, 235-238 (*Bosistidine*)Wu, T.-S. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 85-90 (*Buxifoliadine C*)**1,3,5-Trihydroxy-4-prenylacridone**

T-581

1,3,5-Trihydroxy-4-(3-methyl-2-butenyl)-9(10H)-acridinone, 9CI
[85990-02-9]C₁₈H₁₇NO₄ 311.337Alkaloid from the stem bark of *Oriopsis glaberrima*. α-Glucosidase inhibitor. Yellow powder (MeOH). Mp 250°. [α]_D²⁵ +51.6 (c, 0.62 in MeOH). λ_{max} 233 (log ε 1.49); 253 (log ε 1.99); 276 (log ε 1.73); 278 (log ε 1.75); 289 (log ε 1.5); 303 (log ε 1.56); 341 (log ε 0.86); 397 (log ε 1.2) (MeOH).**N-Me: Oriaciacidone E**

[886227-05-0]

C₁₉H₁₉NO₄ 325.363Alkaloid from the stem bark of *Oriopsis glaberrima*. Amorph. yellow powder (MeOH). [α]_D²⁵ +63.6 (c, 0.7 in MeOH). λ_{max} 235 (log ε 1.5); 253 (log ε 2); 275 (log ε 1.8); 280 (log ε 1.75); 290


(log ε 1.5); 303 (log ε 1.6); 342 (log ε 1); 420 (log ε 1.21) (MeOH).

3-Me ether, N-Me: 1,5-Dihydroxy-3-methoxy-10-methyl-4-prenylacridone.**Glycoctrine I**


[82354-36-7]

Alkaloid from the root and stem bark of *Glycosmis citrifolia* and from the root bark of *Citrus grandis* f. *hakuni-kuyu*. Orange-yellow needles (CHCl₃). Mp 210-212°. λ_{max} 228 (log ε 4.17); 268 (log ε 4.57); 322 (sh) (log ε 4.01); 337 (log ε 4.07); 415 (log ε 3.58) (MeOH).**Δ³-Isomer, 2'ξ-hydroxy, 3-Me ether, N-Me: 1,5-Dihydroxy-4-(2-hydroxy-3-methyl-3-butenyl)-3-methoxy-10-methylacridone. Marshmine**
[160927-88-8]C₂₀H₂₁NO₅ 355.39Alkaloid from roots of Marsh grapefruit (*Citrus paradisi*). Yellow oil. [α]_D -28 (c, 0.15 in CHCl₃).Wu, T.-S. *et al.*, *J.C.S. Perkin 1*, 1983, 1681-1688 (*Glycoctrine I*)Takemura, Y. *et al.*, *Heterocycles*, 1994, **39**, 315-318 (*Marshmine*)Wansi, J.D. *et al.*, *Chem. Pharm. Bull.*, 2006, **54**, 292-296 (*Oriopsis glaberrima* constits, *Oriaciacidone E*)**2,4,7-Trihydroxy-3-prenylquinoline**

T-582

4,7-Dihydroxy-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CIC₁₄H₁₅NO₃ 245.277**NH-form****O⁷,N-Di-Me, 2'ξ,3'-epoxide: 3-(2,3-Epoxy-3-methylbutyl)-4-hydroxy-7-methoxy-1-methyl-2(1H)-quinolinone**
C₁₆H₁₉NO₄ 289.33Alkaloid from *Toddalia aculeata*. Yellow cryst. Mp 110-112°. λ_{max} 232; 238; 243; 248; 250; 299; 324 (MeOH).**2',3'-Dihydro, 2'ξ,3'-dihydroxy, O⁴,O⁷,N-tri-Me: 3-(2,3-Dihydroxy-3-methylbutyl)-4,7-dimethoxy-1-methyl-2(1H)-quinolinone**C₁₇H₂₃NO₅ 321.372Alkaloid from *Toddalia aculeata*. Yellow cryst. Mp 89-92°. λ_{max} 232; 238; 257; 284; 331 (MeOH).Jain, S.C. *et al.*, *Phytochemistry*, 2006, **67**, 1005-1010 (*isol, pmr, cmr, ms*)**2,4,8-Trihydroxy-3-prenylquinoline**

T-583

4,8-Dihydroxy-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CIC₁₄H₁₅NO₃ 245.277**4,8-Di-Me ether: 4,8-Dimethoxy-3-(3-****methyl-2-butenyl)-2(1H)-quinolinone. 2-Hydroxy-4,8-dimethoxy-3-prenylquinoline. Glycolone†**

[41303-26-8]

C₁₆H₁₉NO₃ 273.331Alkaloid from the leaves of *Glycosmis pentaphylla*. Cryst. (Me₂CO/petrol). Mp 118°.**NH-form****O⁶,N-Di-Me: 8-Hydroxy-4-methoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CI. Glycosolone**

[67879-81-6]

C₁₆H₁₉NO₃ 273.331Alkaloid from the root bark of *Glycosmis pentaphylla* (Rutaceae). Light yellow cryst. (C₆H₆/petrol). Mp 159°.**O⁶,N-Di-Me: 4-Hydroxy-8-methoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CI. Glycophylone**

[41303-24-6]

C₁₆H₁₉NO₃ 273.331Alkaloid from the seeds of *Glycosmis pentaphylla* (Rutaceae). Mp 151°.**O⁴,O⁸,N-Tri-Me: 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone. O-Methylglycosolone. O-Methylglycophylone**

[41303-25-7]

C₁₇H₂₁NO₃ 287.358Alkaloid from the roots of *Glycosmis arborea* and *Glycosmis mauritiana*. Oil. Bp_{0.0001} 160°. λ_{max} 240; 258; 285; 295; 335 (MeOH).**O⁴-Et, N-Me: 4-Ethoxy-8-hydroxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, 9CI. Homoglycosolone**

[107030-41-1]


C₁₇H₂₁NO₃ 287.358Alkaloid from root bark of *Glycosmis pentaphylla* (Rutaceae). Mp 168°.**2'-Oxo, 2',3'-dihydro, O⁴,O⁸,N-tri-Me: 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-oxobutyl)-2(1H)-quinolinone. Oriaxalone A**C₁₇H₂₁NO₄ 303.357Alkaloid from the stems of *Orixa japonica*. Oil. λ_{max} 212; 238; 256; 284; 292; 332 (MeOH).**Δ³-Isomer, 2'-oxo, O⁴,O⁸-di-Me: 4,8-Dimethoxy-3-(3-methyl-2-oxo-3-butenyl)-2(1H)-quinolinone. Oriaxalone C**C₁₆H₁₇NO₄ 287.315Alkaloid from the stems of *Orixa japonica*. Oil. λ_{max} 226; 234 (sh); 254; 280; 292; 326 (MeOH).**Δ³-Isomer, 2'-oxo, O⁴,O⁸,N-tri-Me: 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-oxo-3-butenyl)-2(1H)-quinolinone. Oriaxalone B**C₁₇H₁₉NO₄ 301.341Alkaloid from the stems of *Orixa japonica*. Oil. λ_{max} 218; 230; 256; 284; 294; 328 (MeOH).Clarke, E.A. *et al.*, *J.C.S.*, 1964, 438 (*synth, N,O,O-tri-Me*)Rastogi, K. *et al.*, *Phytochemistry*, 1980, **19**, 945-948 (*isol, uv, ir, pmr, ms*)Das, B.P. *et al.*, *Indian J. Chem., Sect. B*, 1982, **21**, 176 (*isol, uv, ir, pmr, cmr, ms, struct, Glycosolone*)Wu, T.-S. *et al.*, *J.C.S. Perkin 1*, 1983, 1681 (*isol, deriv*)

Bhattacharyya, P. *et al.*, *Chem. Ind. (London)*, 1984, 352 (*Glycopholone*)
 Ramesh, M. *et al.*, *Tetrahedron*, 1984, **40**, 4041 (*synth. ir, pmr, deriv*)
 Bhattacharyya, P. *et al.*, *Phytochemistry*, 1985, **24**, 634 (*Glycolone*)
 Kumar, P. *et al.*, *Chem. Ind. (London)*, 1986, 669 (*Homoglycosolone*)
 Chakravarty, A.V. *et al.*, *Phytochemistry*, 1999, **50**, 1263-1266 (*O-Methylglycosolone*)
 Ito, C. *et al.*, *J. Nat. Prod.*, 2004, **67**, 1800-1803 (*Orixalones*)

3,4,7-Trihydroxy-8-prenyl-2-quinolinecarboxylic acid

T-584

1,4-Dihydro-3,7-dihydroxy-8-(3-methyl-2-butenyl)-4-oxo-2-quinolinecarboxylic acid



$C_{15}H_{15}NO_5$ 289.287

NH-form

O^3,N -Di-Me, Me ester: *Megistonine I*


$C_{18}H_{21}NO_5$ 331.368

Alkaloid from the bark of *Sarcomelicope megistophylla*. Amorph. yellow powder. λ_{max} 243 (log ϵ 3.91); 267 (sh); 335 (log ϵ 3.48) (MeOH).

Fokialakis, N. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 2009-2010

3,4,5-Trihydroxy-2-[[2-[(3-pyridinylcarbonyl)amino]benzoyl]amino]benzoic acid, 9CI

T-585



$C_{20}H_{15}N_3O_7$ 409.354

Tri-Me ether, Me ester: *Methyl 3,4,5-trimethoxy-2-[[2-[(3-pyridinylcarbonyl)amino]benzoyl]amino]benzoate* [81469-77-4]

$C_{24}H_{23}N_3O_7$ 465.462


Metab. of *Aspergillus terreus* var. *africanus*. Needles (MeOH). Mp 141-143°. Alkaloid not named in the paper.

Arai, K. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1005 (*uv, ir, pmr, ms, struct, synth*)

4,5,8-Trihydroxy-2-quinolinecarboxylic acid

T-586

1,4-Dihydro-5,8-dihydroxy-4-oxo-2-quinolinecarboxylic acid, 9CI [115525-96-7]




$C_{10}H_7NO_5$ 221.169

Depicted as the quinolone tautomer which prob. predominates in soln. Pigment from the Antarctic sponge *Dendrilla membranosa*. Exhibits antimicrobial activity. Yellow solid. Sol. H_2O . Mp 295-300° dec. λ_{max} 236 (ϵ 16300); 258 (ϵ 13300); 280 (sh) (ϵ 6550); 353 (ϵ 4250) (H_2O at pH 10) (Derep). λ_{max} 232 (ϵ 17100); 250 (ϵ 14600); 270 (sh) (ϵ 6550); 331 (ϵ 3900); 347 (ϵ 3080) (H_2O) (Derep). λ_{max} 232 (ϵ 17100); 250 (ϵ 14600); 331 (ϵ 3900); 347 (ϵ 3080) (H_2O) (Berdy).

Molinski, T.F. *et al.*, *Tet. Lett.*, 1988, **29**, 2137 (*isol, uv, ir, pmr, ms, struct*)

3,5,6-Trihydroxy-6-undecyl-2-piperidinecarboxylic acid

T-587



$C_{17}H_{33}NO_5$ 331.451

6- O - β -*D*-Glucopyranoside: [194099-50-8]

$C_{23}H_{43}NO_{10}$ 493.593


Alkaloid from *Cyclamen coum*.

Yayli, N. *et al.*, *Turk. J. Chem.*, 1997, **21**, 139-143 (*isol, pmr, cmr, ms*)

3,13,23-Trihydroxyveratra-7,12(14)-dien-6-one

T-588

7,8,12,14-Tetradehydro-5,6,12,13-tetrahydro-3,13,23-trihydroxyveratraman-6-one



$C_{27}H_{41}NO_4$ 443.625

Nomenclature of these compounds is confusing and not well established.

(3 β ,5 α ,13 β OH,23R,25S)-form [946412-11-9]

Alkaloid from the bulbs of *Fritillaria hupehensis*. Amorph. powder (MeOH). Mp 132-133.1°. Small negative optical rotation. λ_{max} 242 (log ϵ 4.28); 348 (log ϵ 3.56) (no solvent reported).

13-Deoxy: 3,23-Dihydroxyveratra-7,12(14)-dien-6-one

[1011725-08-8]

$C_{27}H_{41}NO_3$ 427.626


Alkaloid from the bulbs of *Fritillaria hupehensis*. Powder (MeOH). Mp 132-133°. $[\alpha]_D^{20}$ -0.01 (c, 0.001 in $CHCl_3$). λ_{max} 242 (log ϵ 4.28); 348 (log ϵ 3.56) (MeOH).

Zhang, Y.H. *et al.*, *Chin. J. Chem.*, 2007, **25**, 1728-1731 (*isol, pmr, cmr*)

Zhang, Y.H. *et al.*, *Chem. Biodiversity*, 2008, **5**, 259-266 (*isol, pmr, cmr*)

1,1,3-Tri-(1*H*-indol-3-yl)butane

T-589



$C_{28}H_{25}N_3$ 403.526


(ξ)-form [639079-45-1]

Isol. from the marine bacterium *Vibrio parahaemolyticus* Bio249. Yellowish solid. λ_{max} 276 (sh) (log ϵ 4); 282 (log ϵ 4.03); 290 (log ϵ 3.99) (MeOH).

Veluri, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1520-1523 (*isol, pmr, cmr, ms*)

1,1,1-Tri-(1*H*-indol-3-yl)ethane

T-590



$C_{26}H_{21}N_3$ 375.472


Isol. from *Escherichia coli* supplemented with indole.

Garbe, T.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 596-598

Tri-1*H*-indol-3-ylmethane

T-591

3,3',3''-Methylidynetris-1*H*-indole, 9CI [518-06-9]



$C_{25}H_{19}N_3$ 361.445

Isol. from the marine bacterium *Vibrio parahaemolyticus* Bio249. Cryst. (MeOH). Mp 244-246°. λ_{max} 248 (sh) (log ϵ 3.56); 283 (log ϵ 3.32); 388 (log ϵ 3.65) (MeOH).

N,N',N''-Tri-Me: *Tris(1-methyl-3-indolyl)methane*. 3,3',3''-Methylidynetris[1-methyl-1*H*-indole]


[27065-95-8]
 $C_{28}H_{25}N_3$ 403.526
 Cryst. (Py). Mp 268-270°.

Harley-Mason, J. *et al.*, *Biochem. J.*, 1952, **51**, 430 (*synth*)
 Bergman, J. *et al.*, *J. Het. Chem.*, 1971, **8**, 329; 1985, **22**, 341 (*synth, deriv*)
 Akgün, E. *et al.*, *J. Het. Chem.*, 1983, **20**, 1303 (*synth, ir, pmr*)
 Koshima, H. *et al.*, *J. Het. Chem.*, 2002, **39**, 1089-1091 (*synth, ir, pmr*)
 Veluri, R. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1520-1523 (*isol, pmr, cmr*)

Trikendiol

T-592

[158734-27-1]



Tentative Relative Configuration


$C_{38}H_{46}N_2O_4$ 594.792
 Red pigment from the sponge *Trikentrion loeve*. Sows anti-HIV props. Red cryst. (Me_2CO). Mp 160-162°. $[\alpha]_D +102$ (c, 0.02 in $CHCl_3$). λ_{max} 210 (ϵ 37400); 265 (ϵ 8050); 337 (sh); 510 (ϵ 8360) (MeOH assumed, not reported) (Derep).

Loukaci, A. *et al.*, *Tet. Lett.*, 1994, **35**, 6869-6872 (*isol, uv, pmr, cmr, struct*)

Trikentramine

T-593

2-(5-Ethyl-1,3,4,7-tetramethyl-1H-inden-6-yl)-1H-pyrrole, 9CI
 [129536-24-9]




$C_{19}H_{23}N$ 265.397
 Alkaloid from the Senegalese sponge *Trikentrion loeve*. Cryst. (hexane). Mp 145°. $[\alpha]_D^{27} +175$ (c, 0.81 in $CHCl_3$). λ_{max} 209 (ϵ 21300); 222 (ϵ 39700); 271 (ϵ 10700) (no solvent reported).

Aknin, M. *et al.*, *Tet. Lett.*, 1990, **31**, 2979 (*isol, uv, pmr, cmr, cryst struct*)

Trikentrin A

T-594

4-Ethyl-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, 9CI



(1''R,3''S)-form

R = H, R' = CH_2CH_3

$C_{15}H_{19}N$ 213.322

(1''R,3''S)-form

(+)-*cis*-form
 [107368-92-3]

Isol. from *Trikentrion flabelliforme*. Possesses antimicrobial activity. Unstable oil which darkens on storage. Sol. MeOH, $CHCl_3$. $[\alpha]_D +48$ (c, 2.47 in $CHCl_3$). λ_{max} 241 (ϵ 11350); 271 (ϵ 11200) (MeOH) (Berdy).

(1''S,3''S)-form

(+)-*trans*-form
 [107438-53-9]

Isol. from the marine sponge *Trikentrion flabelliforme*. Possesses antimicrobial activity. Stable oil which cryst. on standing. Sol. MeOH, $CHCl_3$. $[\alpha]_D +23.3$ (c, 1.0 in $CHCl_3$). λ_{max} 222 (ϵ 68400); 271 (ϵ 12600) (MeOH) (Berdy).

Capon, R.J. *et al.*, *Tetrahedron*, 1986, **42**, 6545 (*isol, uv, ir, pmr, cmr, ms*)

Macleod, J.K. *et al.*, *Tet. Lett.*, 1988, **29**, 391 (*synth, pmr*)

Muratake, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 854 (*abs config, bibl*)

Huntley, R.J. *et al.*, *Org. Lett.*, 2006, **8**, 3403-3406 (*synth*)


Jackson, S.K. *et al.*, *J.O.C.*, 2007, **72**, 1405-1411 (*synth*)

Silva, L.F. *et al.*, *Org. Lett.*, 2008, **10**, 5417-5420 (*synth*)

Trikentrin B

T-595

5-(1-Butenyl)-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, 9CI



(6*R*,8*S*)-form

$C_{17}H_{21}N$ 239.36

CAS numbering shown. Isol. as an inseparable mixt. of *cis*- and *trans*-Trikentrin B's. Shows antimicrobial activity. Unstable oil which darkens on storage. $[\alpha]_D -13$ (c, 1.97 in $CHCl_3$). λ_{max} 249 (ϵ 24200); 275 (ϵ 8800) (MeOH) (Derep).

(6*R*,8*S*)-form

cis-form
 [107438-54-0]

Isol. from the marine sponge *Trikentrion flabelliforme*.

(6*S*,8*S*)-form

trans-form
 [107368-93-4]

Isol. from *Trikentrion flabelliforme*.

Capon, R.J. *et al.*, *Tetrahedron*, 1986, **42**, 6545 (*isol, uv, ir, pmr, cmr, ms, struct*)

Muratake, H. *et al.*, *Chem. Pharm. Bull.*, 1994, **42**, 854 (*abs config, bibl*)

Macleod, J.K. *et al.*, *Aust. J. Chem.*, 1998, **51**, 177-187 (*synth, pmr, cmr*)


Jackson, S.K. *et al.*, *Org. Lett.*, 2005, **7**, 1215-1218 (*synth*)

Huntley, R.J. *et al.*, *Org. Lett.*, 2006, **8**, 3403-3406 (*synth*)

1,2,8-Trimethoxy-6-methyl-9H-carbazole, 9CI

T-596

Murrayastine
 [104778-02-1]



$C_{16}H_{17}NO_3$ 271.315


Alkaloid from the stem bark of *Murraya euchrestifolia* (Rutaceae). Syrup.

Furukawa, H. *et al.*, *Chem. Pharm. Bull.*, 1986, **34**, 2672 (*uv, ir, pmr, ms, struct, synth*)

***ar*-Trimethoxy-1-methylisoquinoline**

T-597

[93474-22-7]



$C_{13}H_{15}NO_3$ 233.266


Alkaloid detected in *Pachycereus weberi* by tandem mass spectrom. (Cactaceae).

Roush, R.A. *et al.*, *Anal. Chem.*, 1985, **57**, 109-114 (*occur*)

4,8,9-Trimethoxy-N-methyl-2,3-methylenedioxy-pavinane

T-598

[60941-19-7]



$C_{21}H_{23}NO_5$ 369.416

Probable struct., rel. config. only. Numbered by the authors as 2,3,7-trimethoxy-8,9-methylenedioxy (the pavin nucleus shows twofold rotational symmetry).


Alkaloid from *Thalictrum strictum* epigeal parts (Ranunculaceae). Prisms (Et_2O). Mp 144-145°. $[\alpha]_D^{22} -174$ (c, 0.977 in MeOH).

Maekh, S.Kh. *et al.*, *Khim. Prir. Soedin.*, 1976, **116**; *Chem. Nat. Compd. (Engl. Transl.)*, **110** (*isol, uv, pmr, ms*)

1,2,4-Trimethoxy-3H-phenoxazin-3-one

T-599

AV toxin E
 [112899-86-2]



$C_{15}H_{13}NO_5$ 287.271

Prod. by leaf spot fungus *AcrospERMUM viticola*. Phytotoxin. Red needles (MeOH). Mp 160-162°. Related to 2-Amino-3H-phenoxazin-3-one, A-862 and Michigazone, M-590.

4-Demethoxy: 1,2-Dimethoxy-3H-phenoxazin-3-one. **AV toxin D** [112899-85-1]

C₁₄H₁₁NO₄ 257.245

From *AcrospERMUM viticola*. Phytotoxin. Yellow-brown plates (MeOH). Mp 148-150°.

Kinjo, J. *et al.*, *Tet. Lett.*, 1987, **28**, 3697 (*isol. struct*)

Kajimoto, T. *et al.*, *Chem. Lett.*, 1988, 1113 (*synth*)

Trimethylamine, 8CI T-600

N,N-Dimethylmethanamine, 9CI. FEMA 3241

[75-50-3]

Me₃N

C₃H₉N 59.111

Manuf. by catalytic reaction of ammonia with MeOH. Widespread in higher plants, fungi and bacteria. Volatile component of anal sac secretions from dog *Canis familiaris* and coyote *Canis latrans*. Used for manuf. of choline salts. Volatile liq. with fishy odour. Misc. H₂O, EtOH; sol. Et₂O, C₆H₆, CHCl₃. d₄²⁰ 0.67. Mp -117.2°. Bp_{746.6} 3.2-3.8°. pK_a 9.81. Crit. pt. 160.5°/41 atm.

- ▶ Extremely flammable. Eye, skin and respiratory tract irritant. High vapour conc. may affect CNS. LD₅₀ (mus, ivn) 90 mg/kg. LC₅₀ (mus, ihl) 19000 mg m⁻³. OES: long-term 10 ppm; short-term 15 ppm. PA0350000

Hydrochloride: [593-81-7]

Needles (EtOH). Sol. EtOH; insol.

Et₂O. Mp 277-278° dec. Subl. 200.

- ▶ LD₅₀ (mus, ivn) 325 mg/kg. YH2700000

Hydrobromide: [2840-24-6]

Prisms (EtOH). Mp 243-245°.

Hydroiodide: [20230-89-1]

Prisms (EtOH). Sol. H₂O, insol. Et₂O. Mp 263°.

SO₃ complex: [3162-58-1]

C₃H₉NO₃S 139.175

Moisture-sensitive solid. Mp 232° dec Mp 240°.

N-Oxide: see Trimethylamine oxide, T-601

N-Me: see Tetramethylammonium(1+), T-280

[16962-53-1]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 298D; 299A (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 479C (*nmr*)

Aldrich Library of FT-IR Spectra: Vapor Phase, 1989, **3**, 383B (*ir*)

Koepfen, A. *et al.*, *Ber.*, 1905, **38**, 882 (*synth*) *Org. Synth., Coll. Vol.*, **1**, 1932, 531 (*synth, props*)

Moede, J.A. *et al.*, *J.A.C.S.*, 1949, **71**, 852 (*SO₃ complex*)

Ger. Pat., 1969, 2 053 709; *CA*, **75**, 35113q (*synth*)

Hvistendahl, G. *et al.*, *Org. Mass Spectrom.*, 1970, **3**, 821 (*ms*)

Karrer, W. *et al.*, *Konstitution und Vorkommen der Organischen Pflanzenstoffe*, 2nd edn., Birkhäuser Verlag, 1972, no. 2446 (*occur*)

Price, C.C. *et al.*, *J.O.C.*, 1973, **38**, 615 (*pmr*) Sarneski, J.E. *et al.*, *Anal. Chem.*, 1975, **47**, 2116 (*cmr*)

Preti, G. *et al.*, *J. Chem. Ecol.*, 1976, **2**, 177

Blake, A.J. *et al.*, *Acta Cryst. C*, 1984, **40**, 413 (*cryst struct*)

Halpern, A.M. *et al.*, *J.A.C.S.*, 1986, **108**, 3907 (*uv*)

Ullmann's Encycl. Ind. Chem., 5th Ed., VCH, Weinheim, 1990, **A16**, 535 (*rev*)

Murphy, W.F. *et al.*, *J. Phys. Chem.*, 1993, **97**, 581 (*ir, Raman*)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **7**, 5195-5197 (*use*)

Fenaroli's Handbook of Flavor Ingredients, 3rd edn., (ed. Burdock, G.A.), CRC Press, 1995, 767

Encyclopedia of Food and Color Additives, (ed. Burdock, G.A.), CRC Press, 1997, 2835; 2836

Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 1280

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TLD500; TLE000; TLD750; TLF750

Trimethylamine oxide, 8CI T-601

N,N-Dimethylmethanamine N-oxide, 9CI.

Trimethylloxamine. Triox

[1184-78-7]

Me₃N⁺-O⁻

C₃H₉NO 75.11

Widely distrib. in animal tissues, esp. fish. Oxidising agent used in synthesis. Cleaves Si-C bonds. Large cryst. (DMF). Sol. H₂O, MeOH. Mp 224-226° dec. Hygroscopic.

- ▶ Potentially explosive.

Dihydrate: [62637-93-8]

Cryst. (H₂O). Mp 96°.

- ▶ YH2850000

Hydrochloride: [7651-88-9]

Needles (EtOH). Sol. H₂O, hot

MeOH. Mp 218°.

Hydroiodide: [81752-56-9]

Prisms (EtOH). Sol. H₂O, EtOH. Mp 130° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **1**, 299B (*ir*)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **1**, 480B (*nmr*)

Dunstan, W.R. *et al.*, *J.C.S.*, 1899, **75**, 792 (*synth*)

Caron, A. *et al.*, *Acta Cryst.*, 1964, **17**, 102 (*cryst struct*)

Craig, J.C. *et al.*, *J.O.C.*, 1970, **35**, 1721 (*synth, props*)

Bravo, R. *et al.*, *Org. Magn. Reson.*, 1973, **5**, 357 (*pmr*)

Olah, G.A. *et al.*, *J.O.C.*, 1978, **43**, 2268 (*cmr*) *Fieser and Fieser's Reagents for Organic Synthesis*, Wiley, 1980, **8**, 507; 1981, **9**, 31; 489; 1982, **10**, 423; 1986, **12**, 533; 1988, **13**, 325 (*use*)

Harmon, K.M. *et al.*, *J. Mol. Struct.*, 1982, **78**, 43 (*ir*)

Beugelmans, R. *et al.*, *Can. J. Chem.*, 1985, **63**, 725 (*use*)

Chastanet, J. *et al.*, *J.O.C.*, 1985, **50**, 2910 (*use*)

Sakurai, H. *et al.*, *Tet. Lett.*, 1986, **27**, 75 (*use*)

Soderquist, J.A. *et al.*, *Tet. Lett.*, 1986, **27**, 3961 (*synth*)

Mak, T.C.W. *et al.*, *J. Mol. Struct.*, 1988, **178**, 169 (*dihydrate, cryst struct*)

Encyclopaedia of Reagents for Organic Synthesis, (ed. Paquette, L.A.), Wiley, 1995, **7**, 5197-5201 (*use*)

Knoelker, H.-J. *et al.*, *J. Prakt. Chem.*, 1996, **338**, 190 (*use*)

Bretherick, L. *et al.*, *Handbook of Reactive Chemical Hazards*, 4th edn., Butterworths, 1990, 1236

Luxon, S.G. *et al.*, *Hazards in the Chemical Laboratory*, 5th edn., Royal Society of Chemistry, 1992, 1281

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TLE100; TLE250

(2-Trimethylammonioethyl)- phosphonic acid betaine T-602

N,N,N-Trimethyl-2-phosphonoethanaminium hydroxide, inner salt, 9CI. Phosphonyldeoxycholine

[14596-57-7]

Me₃N⁺CH₂CH₂PO₃H⁻

C₅H₁₄NO₃P 167.144

Constit. of the sea anemone *Anthopleura xanthogrammica*. Needles (EtOH). Mp 252° dec. Hygroscopic.


Myers, T.C. *et al.*, *J.O.C.*, 1957, **22**, 180 (*synth*) Rosenthal, A.F. *et al.*, *J.A.C.S.*, 1958, **80**, 5240 (*synth*)

Kittredge, J.S. *et al.*, *Biochemistry*, 1967, **6**, 289 (*isol, synth, ir, pmr*)

Kittredge, J.S. *et al.*, *Comp. Biochem. Physiol.*, 1969, **29**, 859 (*biosynth*)

Richard, H. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1974, **278**, 1275 (*pmr, conformm*)

2,4,6-Trimethyl-2,4-decadienoic acid T-603



C₁₃H₂₂O₂ 210.316

(2E,4E,6R)-form [112406-54-9]

Prod. by *Streptomyces parvulus*.

Amide: 2,4,6-Trimethyl-2,4-decadienamide. **Antibiotic SW-B. SW-B. Antibiotic 64p-A** [149837-44-5]

C₁₃H₂₃NO 209.331

Prod. by *Streptomyces flavescens* and *Streptomyces parvulus*. Also obt. by directed biosynthesis. Antifungal agent. λ_{max} 208 ; 260 (MeOH).


[112406-55-0]

Kaiser, D. *et al.*, *Appl. Microbiol. Biotechnol.*, 1994, **41**, 309 (*isol*)




Hwang, B.K. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3653 (*isol, bibl, amide*)

2-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1H-indole T-604


2-Farnesylindole



C₂₃H₃₁N 321.505

(E,E)-form [113801-52-8]Prod. by *Escherichia coli*.Brady, S.F. *et al.*, *J.A.C.S.*, 2007, **129**, 12102-12103 (*isol*, *pmr*, *ms*)**3-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1H-indole, 9CI**
3-Farnesylindole
[102720-33-2] $C_{23}H_{31}N$ 321.505Alkaloid from the leaves of *Uvaria pandensis* (Annonaceae). Oil.**8',9'-Dihydroxy: 3-(8,9-Dihydroxyfarnesyl)indole** $C_{23}H_{31}NO_2$ 353.503Alkaloid from the roots of *Uvaria pandensis* (Annonaceae). Viscous light brown oil. $[\alpha]_D^{20} +23.5$ (c, 0.17 in MeOH).**8',9'-Dihydroxy; 6',7'-dihydro: 3-(6,7-Dihydro-8,9-dihydroxyfarnesyl)indole** $C_{23}H_{33}NO_2$ 355.519Alkaloid from the roots of *Uvaria pandensis* (Annonaceae). Viscous light brownish oil. $[\alpha]_D^{20} +12.1$ (c, 0.34 in MeOH).Nkunya, M.H.H. *et al.*, *Phytochemistry*, 1987, **26**, 2402-2403; 1989, **28**, 2217-2218 (*isol*, *uv*, *ir*, *pmr*, *cmr*, *ms*)**2-(2,4,6-Trimethyl-1-heptenyl)-4(1H)-quinazolinone**
2-(2,4,6-Trimethyl-1-heptenyl)-4-quinazolinone. 4-Hydroxy-2-(2,4,6-trimethyl-1-heptenyl)quinazolinone
[564444-35-5] $C_{18}H_{24}N_2O$ 284.4Alkaloid from the leaves of *Zanthoxylum budrunga*. Mp 160-162°. λ_{max} 192 ; 207 ; 283 ; 309 (MeOH).Ahmad, M.U. *et al.*, *Fitoterapia*, 2003, **74**, 191-193 (*isol*, *pmr*)**2,4,5-Trimethyl-1H-imidazole, 9CI**
[822-90-2] $C_6H_{10}N_2$ 110.158Spar. sol. H_2O . Mp 132.5-133°. pK_a 8.92

(25°). Strong base.

Hydrochloride: [70807-89-5]
Mp 316°.**1,3-Dibenzyl: 1,3-Dibenzyl-2,4,5-trimethylimidazolium(1+). Lepidiline B**
[596093-97-9] $C_{20}H_{23}N_5^{\oplus}$ 291.415Alkaloid from the roots of *Lepidium meyenii* (maca). Plates ($Me_2CO/MeOH$) (as chloride). Mp 220-222° (chloride). λ_{max} 201 (log ϵ 3.32); 258 (log ϵ 2.29); 278 (log ϵ 2.1) (MeOH (chloride)).Schaubum, M.L. *et al.*, *Tet. Lett.*, 1971, 2205 (*synth*)Gnichtel, H. *et al.*, *Chem. Ber.*, 1972, **105**, 1865 (*synth*)Cui, B. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1101-1103 (*Lepidiline B*)**1,3,7-Trimethylisoguanine** T-608**1,3,6,7-Tetrahydro-6-imino-1,3,7-trimethyl-2H-purin-2-one, 9CI**
[290821-46-4] $C_8H_{11}N_5O$ 193.208Isol. from a New Zealand ascidian *Pseudodistoma* sp. Constit. of the cuticular wax of *Osmunda regalis*. Light tan oil. λ_{max} 214 (log ϵ 3.6); 288 (log ϵ 3.4) (MeOH).Copp, B.R. *et al.*, *J. Nat. Prod.*, 2000, **63**, 1168-1169 (*isol*, *pmr*, *cmr*, *uv*)**N',N'',N'''-Trimethyl-N-(3-methyl-2,4-dodecadienoyl)spermidine** T-609

N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methyl-2,4-dodecadienamamide. 2,7,14-Trimethyl-2,7,11-triaza-13,15-tricosadien-12-one

 $H_3C(CH_2)_6CH=CHC(CH_3)=CHCONH(CH_2)_3NMe(CH_2)_4NMe_2$ $C_{23}H_{45}N_3O$ 379.628**(2'E,4'E)-form** [583850-68-4]Isol. from a *Simularia* sp. Oil.**4',5'-Dihydro:** N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methyl-2-dodecenamamide. 1-(Dimethylamino)-5,12-dimethyl-5,9-diazaheneicos-11-en-10-one. N',N'',N'''-Trimethyl-N-(3-methyl-2-dodecenoil)spermidine. 2,7,14-Trimethyl-2,7,11-triaza-13-tricosen-12-one
[73710-47-1]
 $C_{23}H_{47}N_3O$ 381.644
Isol. from *Simularia brongersmai*.**(2'Z,4'E)-form** [583850-67-3]Isol. from a soft coral *Simularia* sp. Pale yellow oil (rapidly darkens on standing).**4',5'-Dihydro:** [81419-45-6]Isol. from the soft coral *Simularia**brongersmai*. Cytotoxic agent. Pale yellow oil. Rapidly darkens on standing. Obt. as a 9:1 mixt. with its dihydro deriv.**2',3',4',5'-Tetrahydro: N',N'',N'''-Trimethyl-N-(3-methyl-dodecanoyl)spermidine.** N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methyl-dodecanamide. 2,7,14-Trimethyl-2,7,11-triaza-12-tricosanone
[73710-48-2] $C_{23}H_{49}N_3O$ 383.66Minor constit. of *Simularia brongersmai*. Cytotoxic.Schmitz, F.J. *et al.*, *Tet. Lett.*, 1979, 3387 (*isol*, *ir*, *pmr*, *ms*, *struct*, *Simularia brongersmai constits*)Chantrapromma, K. *et al.*, *Tet. Lett.*, 1980, **21**, 2605 (*synth*, *ir*, *pmr*, *ms*)Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1982, **35**, 69-75 (*isol*, *pmr*, *cmr*, *ms*, *struct*, *synth*)Ojika, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2003, **67**, 1410-1412 (2'E,4'E-form, 2'Z,4'E-form, *isol*, *pmr*, *cmr*)**N',N'',N'''-Trimethyl-N-(5-methyl-2,4-tetradecadienoyl)spermidine** T-610

N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-5-methyl-2,4-tetradecadienamamide, 9CI. 2,7,16-Trimethyl-2,7,11-triaza-13,15-pentacosadien-12-one

 $H_3C(CH_2)_8C(CH_3)=CHCONH(CH_2)_3NMe(CH_2)_4NMe_2$
 $C_{25}H_{49}N_3O$ 407.682**(2E,4E)-form** [583850-70-8]Isol. from the soft coral *Simularia* sp. Cytotoxic. Oil.**(2Z,4E)-form** [583850-69-5]Isol. from a *Simularia* sp. Cytotoxic. Oil. Ojika, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 2003, **67**, 1410-1412 (*isol*, *pmr*, *cmr*)**N',N'',N'''-Trimethyl-N-(5-methyl-3-tetradecenoil)spermidine** T-611

N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-5-methyl-3-tetradecenamamide. 2,7,16-Trimethyl-2,7,11-triaza-14-pentacosen-12-one

 $H_3C(CH_2)_8CH(CH_3)CH=CHCH_2CONH(CH_2)_3NMe(CH_2)_4NMe_2$
 $C_{25}H_{51}N_3O$ 409.697**(3'E,5'ε)-form** [129722-93-6]Isol. from the soft coral *Simularia* sp. Cytotoxic agent. Oil. $[\alpha]_D +7$ (c, 0.2 in CH_2Cl_2). λ_{max} 200 (MeOH).Choi, Y.-H. *et al.*, *J. Nat. Prod.*, 1997, **60**, 495-496**Trimethyl[3-(methylthio)propyl]ammonium(1+), 8CI** T-612N,N,N-Trimethyl-3-(methylthio)-1-propanaminium(1+), 9CI. *TA Toxin A* [61672-50-2] $MeSCH_2CH_2CH_2NMe_3^{\oplus}$ $C_7H_{18}NS^{\oplus}$ 148.292Isol. from the reef shell *Marmarostoma argyrostoma* and the turban shell *Turbo argyrostoma*. Mp 223-225° (as iodide).


S-Me: 3-(Dimethylsulfonio)-N,N,N-trimethylpropanaminium(2+), 9CI
[61672-51-3]
C₈H₂₁NS²⁺ 163.327
Isol. from *Marmarostoma argyrostoma*. Mp 246° (as diiodide).

[54289-29-1, 14617-80-2]

Kirchner, F.D. *et al.*, *J.A.C.S.*, 1955, **77**, 4599 (synth)
Tadashi, S. *et al.*, *Yakugaku Zasshi*, 1964, **84**, 1012; *CA*, **62**, 6552g (synth)
Yasumoto, T. *et al.*, *Nippon Suisan Gakkaishi*, 1974, **40**, 217; *CA*, **80**, 129589e (isol)
Yasumoto, T. *et al.*, *Rinsho Eiyō*, 1976, **49**, 115; *CA*, **86**, 84458s (isol)

Trimethylpyrazine, 9CI T-613

FEMA 3244
[14667-55-1]




C₇H₁₀N₂ 122.169

Urinary signalling pheromone in tree shrews *Tupaia belangeri* and pine vole *Microtus pinetorum*. Isol. from an arctic marine bacterium. Flavouring ingredient. Liq. with roast coffee odour. Bp₇₃₅ 171-172°. pK_{a1} 2.65; pK_{a2} -3.6 (25°, H₂O). Steam-volatile. Odour threshold 9000ppb in H₂O.

- ▶ LD₅₀ (rat, orl) 806 mg/kg. UQ3907000
- Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 841B (ir)
- Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, **3**, 401A (nmr)
- Aldrich Library of FT-IR Spectra: Vapor Phase*, 1989, **3**, 1558B (ir)
- Das, B.K. *et al.*, *J. Indian Chem. Soc.*, 1968, **45**, 1075 (uv)
- Bus, J. *et al.*, *Rec. Trav. Chim. (J. R. Neth. Chem. Soc.)*, 1973, **92**, 123 (ir)
- Calabretta, P.J. *et al.*, *Cosmet. Perfum.*, 1975, **90**, 74 (synth)
- Gumbley, S.J. *et al.*, *J. Het. Chem.*, 1985, **22**, 1143 (props)
- von Stralendorff, F. *et al.*, *J. Chem. Ecol.*, 1987, **13**, 655
- Boyer, M.L. *et al.*, *J. Chem. Ecol.*, 1989, **15**, 649
- Dickschat, J.S. *et al.*, *Chem. Biodiversity*, 2005, **2**, 318-353 (marine isol)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, TME270

2,2,6-Trimethyl-4-(4-quinolinyl)-3-azabicyclo[3.3.1]non-6-ene, 9CI T-614

[151677-05-3]



C₂₀H₂₄N₂ 292.423

Alkaloid from aerial parts of *Aristolotelia chilensis* (Elaeocarpaceae).

Céspedes, C. *et al.*, *Phytochemistry*, 1993, **34**, 881-882 (isol)

Trimethylvinylammonium(1+), 8CI T-615

N,N,N-Trimethylethenaminium(1+), 9CI. Ethenyltrimethylammonium(1+).

Neurine

[13448-18-5]

H₂C=CHN⁺Me₃

C₅H₁₂N⁺ 86.156

Chloride: [6018-82-2]

C₅H₁₂ClN 121.609

Cryst. Mp 193-194°.

Bromide: [10603-92-6]

C₅H₁₂BrN 166.061

Has curare-like activity. Plates (EtOH/Et₂O). Sol. H₂O, EtOH, insol. Et₂O. Mp 194° dec.

▶ LD₅₀ (mus, ipr) 52 mg/kg. BT3675000

Picrate:

Golden-yellow needles (H₂O). Sol. H₂O, hot EtOH, spar. sol. CHCl₃, cold EtOH. Mp 264°.

Hydroxide: [463-88-7]

C₅H₁₃NO 103.164

Formed during putrefaction by dehydration of choline. Occurs free and combined in many animal and vegetable products, e.g., bile, brain, egg yolk, cadavers. Found in *Cannabis sativa*. One of the ptomaine poisons. One of the ptomaine poisons. Syrupy liq. with fishy odour. Sol. H₂O, EtOH. Reacts alkaline and readily absorbs CO₂ from air. Forms hygroscopic cryst. hydrate with 3H₂O. Easily dec. → Me₃N.

▶ Highly toxic. BT3850000

Meyer, K.H. *et al.*, *Ber.*, 1921, **54**, 2265 (synth, hydroxide)

Renshaw, R.R. *et al.*, *J.A.C.S.*, 1925, **47**, 2989 (synth, bromide)

Gardner, C. *et al.*, *J.C.S.*, 1949, 789 (synth, hydroxide, chloride, picrate)

Grabowska, M. *et al.*, *Diss. Pharm.*

Pharmacol., 1967, **19**, 43 (synth, pharmacol)

Ohtsuru, M. *et al.*, *J.A.C.S.*, 1969, **91**, 1187


(bromide, pmr)

Kroeller, E. *et al.*, *CA*, 1973, **79**, 16990 (isol)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VQR300

Trinervine T-616

17,19-Epoxy-sarpagan-19(20H)-ol, 9CI
[132242-49-0]



Absolute Configuration

C₁₉H₂₂N₂O₂ 310.395

Alkaloid from the roots of *Strychnos trinervis* (Loganiaceae). Needles (MeOH/EtOAc). Mp 219-220°. [α]_D²² -7 (c, 0.37

in CHCl₃).

N⁴-Me: Venecurine

[122908-09-2]

C₂₀H₂₅N₂O₂⁺ 325.43

Quaternary alkaloid from Venezuelan curare.

Quetin-Leclercq, J. *et al.*, *Phytochemistry*, 1989, **28**, 2221-2223 (Venecurine)


Mukherjee, R. *et al.*, *Heterocycles*, 1990, **31**,

1819-1822 (Trinervine)

Liu, X. *et al.*, *Org. Lett.*, 2001, **3**, 4023-4026 (synth)

Triophamine T-617

N,N'-Bis(2,4-diethyl-4-hexenyl)guanidine
[81256-25-9]



C₂₁H₃₇N₃O₂ 363.542


Isol. from skin extracts of the dorid nudibranch *Triopha catalinae*. Light yellow oil. [α]_D -7 (c, 1.7 in MeOH). λ_{max} 214 (ε 14500) (MeOH/HCl) (Derep). λ_{max} 232 (ε 10000) (KOH) (Derep). λ_{max} 251 (ε 12000) (MeOH) (Derep).

Gustafson, K. *et al.*, *J.O.C.*, 1982, **47**, 2167-2169 (isol, uv, pmr, cmr, ms, struct)

Piers, E. *et al.*, *Can. J. Chem.*, 1984, **62**, 1 (synth)

Triostin T-618

[12795-78-7]



Triostin A R = -CH(CH₃)₂
B R = -CH₂CH(CH₃)₂ R- or S-
B₀ R = -CH₂CH(CH₃)₂ R- or S-
C R = -CH(CH₃)CH(CH₃)₂

Cyclic depsipeptide antibiotic complex. Prod. by *Streptomyces* spp. Shows broad activity against gram-positive bacteria and Hela cells.

▶ YJ8950000

Triostin A [13758-27-5]

C₅₀H₆₂N₁₂O₁₂S₂ 1087.245

Needles (CHCl₃/MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 245-248° dec. [α]_D^{23.5} -157 (c, 0.97 in CHCl₃). λ_{max} 243 (ε 78800); 323 (ε 12700) (MeOH) (Derep). λ_{max} 243 (ε 70800); 320 (ε 12900) (MeOH) (Berdy).

Triostin B [39421-38-0]

C₅₂H₆₆N₁₂O₁₂S₂ 1115.299

Prod. by: *Streptomyces aureus* in small quantities in medium supplemented with isoleucine. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 209-213°. [α]_D -

144.

Triostin B₀

C₅₄H₆₆N₁₂O₁₂S₂ 1115.299
Sol. MeOH, C₆H₆; poorly sol. H₂O,
hexane. λ_{max} 243 (ε 71000); 323
(ε 13100) (MeOH) (Derep). λ_{max} 243
(ε 61600); 320 (ε 10200) (MeOH)
(Berdy).


Triostin C [10382-35-1]

C₅₄H₇₀N₁₂O₁₂S₂ 1143.352
Isol. from *Streptomyces* sp. Needles
(CHCl₃/MeOH). Sol. MeOH, C₆H₆;
poorly sol. H₂O, hexane, Et₂O. Mp
260° dec. [α]_D²⁵ -143.9 (c, 1.121 in
CHCl₃). λ_{max} 243 (log ε 4.87) and
315-26 nm (4.13). λ_{max} 243 (ε 71000);
323 (ε 13100) (MeOH) (Derep). λ_{max}
243 (ε 74170); 320 (ε 13500) (EtOH)
(Berdy).

- LD₅₀ (mus, ipr) 100 mg/kg. YJ8950400
Otsuka, M. *et al.*, *Tetrahedron*, 1965, **21**, 2931;
1967, **23**, 1535 (*struct*)
Otsuka, M. *et al.*, *J. Antibiot., Ser. A*, 1966, **19**,
128 (*isol, struct*)
U.S. Pat., 1972, 3 647 631; CA, **77**, 3780e
(*manuf*)
Shoji, J. *et al.*, *J. Antibiot.*, 1973, **26**, 302
(*struct*)
Otsuka, M. *et al.*, *J. Antibiot.*, 1976, **29**, 107
(*nmr*)
Blake, T.J. *et al.*, *Tet. Lett.*, 1977, 2621
(*conformm*)
Chakravarty, P.K. *et al.*, *Tet. Lett.*, 1978, 1613
(*synth*)
Shin, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1984, **57**,
2203; 2211 (*synth*)
Sheldrick, G.M. *et al.*, *Acta Cryst. B*, 1995, **51**,
987-999 (*cryst struct*)

Triptonine A†
[226979-90-4]

T-619


C₄₅H₅₅N₂₁ 945.924

Not to be confused with Triptonine A
in P-253. Alkaloid from *Tripterygium*
hypoglaucom. Anti-HIV agent. Needles.
Mp 284-285.5°. [α]_D²⁵ -24.1 (c, 1 in
MeOH). λ_{max} 224 (log ε 3.96); 264
(log ε 3.58) (MeOH).

Duan, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 357-
361 (*isol*)

Triptonine B†
[226980-25-2]

T-620

C₄₅H₅₅N₂₂ 961.923


Alkaloid from *Tripterygium hypoglaucum*.
Anti-HIV agent. Amorph. powder.
[α]_D²⁵ +15.5 (c, 0.6 in MeOH). Not to be
confused with Triptonine B in P-253.
λ_{max} 261 (log ε 3.4) (MeOH).

Duan, H. *et al.*, *J. Nat. Prod.*, 2000, **63**, 357-
361

Triptotin J

[368423-83-0]

T-621

C₂₀H₃₁N₂O₂ 317.47

Alkaloid from the root bark of *Tripterygium*
wilfordii. Yellow oil. [α]_D²⁰ +21 (c,
0.27 in CHCl₃).

Yang, G.-Z. *et al.*, *Helv. Chim. Acta*, 2002, **85**,
168-174 (*isol, pmr, cmr, ms*)

Tris(3-aminopropyl)amine

T-622

N,N-Bis(3-aminopropyl)-1,3-propanedi-
amine, 9CI. 3,3',3''-Triaminotripropyla-
mine, 8CI. N⁴-(3-
Aminopropyl)norspermidine. Trpn
[4963-47-7]
(H₂NCH₂CH₂CH₂)₃N

C₉H₂₄N₄ 188.315

Isol. from the thermophilic bacteria *Hydrogenobacter halophilus* and *Hydrogenobacter thermophilus*. Liq. Bp_{1.5} 134-135°.

Hemihydrate:

Deliquescent cryst. Mp 227-229° (with
foaming).

Hydrochloride (1:4): [66322-79-0]

Cryst. (EtOH). Mp 230-231°.

Picrate:

Yellow needles. Mp 222° dec. Softens
at 190°.

N-(3-Aminopropyl): 3-Amino-N,N,N-tris(3-aminopropyl)-1-propanaminium(1+), 9CI. Tetrakis(3-aminopro-

pyl)ammonium(1+), N⁴,N⁴-Bis(3-
aminopropyl)norspermidine

[111216-37-6]

C₁₂H₃₂N₅[⊕] 246.418

Isol. from *Hydrogenobacter halophilus*,
Hydrogenobacter thermophilus and
Thermus thermophilus. Counterion not
specified.

v. Winkle, J.L. *et al.*, *J.O.C.*, 1966, **31**, 3300-
3306 (*synth*)

Oshima, T. *et al.*, *J. Biol. Chem.*, 1987, **262**,
11979-11981 (*Thermus thermophilus constit*)

Chin, J. *et al.*, *J.A.C.S.*, 1989, **111**, 186-190

(synth, pmr)

Hamana, K. *et al.*, *Biochem. J.*, 1992, **284**, 741-
747 (*isol*)

Nitsu, M. *et al.*, *Chem. Pharm. Bull.*, 1992, **40**,
2958-2961 (*synth*)

Hoveyda, H.R. *et al.*, *Can. J. Chem.*, 1998, **76**,
414-425 (*synth*)

N,N',N''-Tris(3-methylbutyl)guanidine

T-623

N,N',N''-Triisopentylguanidine


(H₃C)₂CHCH₂CH₂N=C[NHCH₂CH₂CH(CH₃)₂]₂C₁₆H₃₅N₃ 269.473

Alkaloid from *Alchornea cordifolia*.
Antibacterial agent.

Lamikanra, A. *et al.*, *Phytother. Res.*, 1990, **4**,
198-200

Trisulcusine

T-624

C₁₈H₁₄N₂O₂ 290.321**(ε)-form**


Alkaloid from the aerial parts of *Anisotes*
trisulcus. Needles (CHCl₃). Mp 235-236°
dec. [α]_D²⁰ -20 (c, 0.02 in CHCl₃). λ_{max} 232
(log ε 3.35); 259 (log ε 2.75); 301 (log ε
2.76); 396 (log ε 2.32) (CHCl₃).

Al-Rehaily, A.J. *et al.*, *Indian J. Chem., Sect. B*,
2002, **41**, 2385-2389 (*isol, pmr, cmr, ms*)

Triticone A

T-625

6-Hydroxy-2-methoxy-3-methylene-10-
propenylidene-2-azaspiro[4.5]dec-8-ene-
1,4,7-trione, 9CI. Spirostaphylotrichin C
[114582-74-0]

C₁₄H₁₅NO₅ 277.276

Isol. from *Drechslera tritici-repentis*,
Curvularia pallescens and *Staphylotri-*
chum coccosporum. Phytotoxin. Cryst.
Mp 133-138°. λ_{max} 224 (ε 19300); 289 (ε

15400) (EtOH) (Derep). λ_{\max} 208 (ε 12900); 288 (ε 12000) (MeOH) (Berdy).

4-Alcohol: Triticone C

$C_{14}H_{17}NO_5$ 279.292

Isol. from *Drechslera tritici-repentis*.

Pale yellow oil. $[\alpha]_D^{25} +2$ (c, 1.0 in $CHCl_3$).

6-Epimer: Triticone B. Spirostaphylotrichin D

[114613-32-0]

$C_{14}H_{15}NO_5$ 277.276

From *Drechslera tritici-repentis*, *Curvularia pallescens* and *Staphylotrichum coccosporum*. Phytotoxin. Enhances fibrinolysis. Cryst. The configurational assignments of Spirostaphylotrichins C and D may be reversed. λ_{\max} 224 (ε 19300); 289 (ε 15400) (EtOH) (Derep).

4-Alcohol, 6-epimer: Triticone D

$C_{14}H_{17}NO_5$ 279.292

From *Drechslera tritici-repentis*. Cryst. (EtOAc). $[\alpha]_D^{25} +22$ (c, 0.64 in MeOH).

Sugawara, F. et al., *J.A.C.S.*, 1988, **110**, 4086 (isol, struct)

Sandmeier, P. et al., *Helv. Chim. Acta*, 1989, **72**, 774; 784 (isol, pmr, cmr, uv, cd, struct, biosynth)

Hallock, Y.F. et al., *J. Nat. Prod.*, 1993, **56**, 747 (Triticones)


Abraham, W.-R. et al., *Tetrahedron*, 1995, **51**, 4947 (isol)

Shinohara, C. et al., *J. Antibiot.*, 2000, **53**, 262-268 (activity)

Triticone E

T-626

[149564-25-0]



$C_{14}H_{19}NO_6$ 297.307

Isol. from culture broth of the plant pathogenic fungus *Drechslera tritici-repentis*. Pale yellow oil. $[\alpha]_D^{25} -70$ (c, 1.0 in MeOH). Insep. mixt. with Triticone F (ratio 10:9).

6-Epimer: Triticone F

[149654-98-8]

$C_{14}H_{19}NO_6$ 297.307

From *Drechslera tritici-repentis*.

Hallock, Y.F. et al., *J. Nat. Prod.*, 1993, **56**, 747 (isol, uv, ir, pmr, cmr, ms, struct)

Triuret, 8CI

T-627

Diimidotricarbonic diamide, 9CI. *Dicarbamoylurea*. *Carbonyldiurea*. *Carbonic acid diureide* [556-99-0]

$H_2NCONHCONHCONH_2$

$C_3H_6N_4O_3$ 146.105

Formed by uv irradiation of Urea.

Identified in cytoplasm of *Amoeba* spp. Leaflets (NH_3 aq.). Sol. acid, alkali, hot H_2O . Mp 233° dec.

Haworth, R.C. et al., *J.C.S.*, 1943, 603 (synth)

Werner, R.A. et al., *CA*, 1947, **41**, 710 (synth)


Griffin, J.L. et al., *Biochim. Biophys. Acta*, 1961, **47**, 433 (occur, ir)

Furrar, W.V. et al., *Chem. Ind. (London)*, 1972, 682 (synth)

Trivalvone

T-628

7-(5,6-Dihydro-1,2-dimethoxy-6-methyl-4H-dibenzo[de,g]quinolin-7-yl)-2-methoxy-1H-dibenzo[de,g]quinolin-1-one, 9CI [131989-90-7]



$C_{36}H_{28}N_2O_4$ 552.628

Alkaloid from *Trivalvaria macrophylla* and *Piptostigma fugax* (Annonaceae). Brown cryst. Mp 258°.


Cortes, D. et al., *J. Nat. Prod.*, 1990, **53**, 862 (isol, uv, ir, pmr, cmr, ms, struct)

Achenbach, H. et al., *Phytochemistry*, 1995, **38**, 1037 (isol, pmr, cmr)

Trolline

T-629

1,5,6,10b-Tetrahydro-8,9-dihydroxypyrrrolo[2,1-a]isoquinolin-3(2H)-one, 9CI. *Oleracein E*. *Salsoline A*



(R)-form

$C_{12}H_{13}NO_3$ 219.24

(R)-form

Alkaloid from *Portulaca oleracea* (purslane). Powder (MeOH). Mp 238-240°. $[\alpha]_D^{26} +61.1$ (c, 0.32 in MeOH). λ_{\max} 290 (log ε 4.44); 318 (log ε 4.43) (MeOH).

(S)-form [724706-17-6]

Alkaloid from the flowers of *Trollius chinensis* and *Salsola collina*. Plates (MeOH). Mp 165-167°. $[\alpha]_D^{20} -197$ (c, 0.8 in MeOH). λ_{\max} 255; 289 (MeOH).

(±)-form

8-Me ether: **Erythrinarine**

[245348-39-4]

$C_{13}H_{15}NO_3$ 233.266

Alkaloid from the stems of *Erythrina arboreascens*. Plates (MeOH). Mp 193-194°. λ_{\max} 286 (MeOH).

Yu, D.L. et al., *Chin. Chem. Lett.*, 1999, **10**, 139-142 (*Erythrinarine*)

Wang, R.F. et al., *Heterocycles*, 2004, **63**, 1443-1448 (*Trolline*)


Zhao, Y. et al., *Yaoxue Xuebao*, 2004, **39**, 598-600 (*Salsoline A*)

Xiang, L. et al., *Phytochemistry*, 2005, **66**, 2595-2601 (*Oleracein E*)

He, Q.Q. et al., *Chin. Chem. Lett.*, 2007, **18**, 651-652 (synth)

Tronocarpine

T-630



$C_{20}H_{20}N_2O_3$ 336.39


Prob. related to the Tacaman alkaloids. Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Oil. $[\alpha]_D +231$ (c, 0.08 in CH_2Cl_2). λ_{\max} 226 (log ε 4.3); 290 (log ε 3.71); 296 (log ε 3.72) (no solvent reported).

Kam, T.-S. et al., *Tet. Lett.*, 2000, **41**, 2733-2736

Tronoharine

T-631

[246136-19-6]



$C_{21}H_{24}N_2O_2$ 336.433


Abs. config prob. as shown, on biogenetic grounds. Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Amorph. $[\alpha]_D +35$ (c, 0.1 in $CHCl_3$). λ_{\max} 202 (log ε 4.43); 250 (log ε 3.88); 272 (log ε 3.83); 296 (log ε 3.58) (no solvent reported).

Kam, T.-S. et al., *Tet. Lett.*, 1999, **40**, 5409-5412

Tropine tropate

T-632

8-Methyl-8-azabicyclo[3.2.1]oct-3-yl α-(hydroxymethyl)benzeneacetate, 9CI



(S)-form

$C_{17}H_{23}NO_3$ 289.374

Log P 1.32 (calc).

(S)-form

Hyoscyamine, BAN, USAN. Daturine.

Duboisine

[101-31-5]

Alkaloid from *Atropa*, *Datura*, *Duboisia*, *Hyoscyamus*, *Anthocercis* and *Scopolia* spp., *Mandragora officinarum* and several other genera in the Solanaceae, esp. some strains of *Duboisia myoporoides*. Antimuscarinic agent with approx. twice potency of atropine. Spasmolytic agent. Bronchodilator, sedative, analgesic. Mp 108-111°. $[\alpha]_D^{15} -22$ (50% EtOH aq.). Log P 1.32 (calc). Racemises slowly in EtOH

to Atropine, more rapidly in alkali or on melting; sometimes occurs naturally as partial racemate. Mandragorine from *M. officinarum* was an alkaloid mixture including Scopolamine.

- ▶ Adverse effects on CNS and peripheral nerves. Adverse ocular effects. Can cause death. LD₅₀ (mus, ivn) 95 mg/kg. NH0875000

Hydrobromide: Hyoscyamine hydrobromide, USAN

[306-03-6]

Mp 151°. Deliquescent. Component of Dolonil and Pyridium Plus.

Sulfate (2:1): Hyoscyamine sulfate, USAN. Anaspaz. Peptard

[6835-16-1] pK_a 9.68 (21°). Component of Donnagel, Donnazyme, Kinesed and Spalix.

Picrate: [5934-49-6]

Mp 165°.

N-Oxide (equatorial O): Hyoscyamine N-oxide 1

[56362-73-3]

C₁₇H₂₃NO₄ 305.373

Alkaloid from *Atropa*, *Datura*, *Hyoscyamus*, *Scopolia* and *Mandragora* spp. (Solanaceae). Mp 165-170° dec. (as hydrochloride).

N-Oxide (axial O): Hyoscyamine N-oxide 2

[56362-74-4]

C₁₇H₂₃NO₄ 305.373

Alkaloid from *Atropa*, *Datura*, *Hyoscyamus*, *Scopolia* and *Mandragora* spp. Mp 175-180° dec. (as hydrochloride).

N-De-Me: Norhyoscyamine

[537-29-1]

C₁₆H₂₁NO₃ 275.347

Alkaloid from aerial parts of *Datura meteloides*. The major alkaloid of some strains of *Duboisia myoporoides* (Solanaceae).

(±)-form

Atropine, BAN, INN, USAN

[51-55-8]

Alkaloid from *Atropa belladonna*, *Datura stramonium*, *Anthocercis* and other Solanaceae. Mydriatic, antispasmodic and cycloplegic agent. Smooth muscle relaxant. Used in preanaesthetic medication to prevent reflex bradycardia and bronchospasm and to decrease gland secretions. Reduces rigidity in parkinsonism. Antidote to poisoning with parasympathomimetic agents, e.g. nerve gases, organophosphorus insecticides. Cause of deliberate and accidental poisoning of humans and livestock. Nontoxic to some spp., e.g. rabbits. Reference material used in elemental microanalysis. Sol. MeOH, CHCl₃, acids; fairly sol. Et₂O, C₆H₆; poorly sol. H₂O, hexane. Mp 116-117°. pK_a 10.2 (25°). Log P 1.32 (calc). In many cases, Atropine may be an artifact of the extraction procedure.

- ▶ Adverse human effects by ingestion, and other routes of administration, incl. dryness of the mouth, difficulty swallowing and thirst. Adverse ocular effects. Can cause death. Variable human lethal

dose, but 100 mg can be fatal. LD₅₀ (rat, orl) 500 mg/kg. Exp. reprod. and teratogenic effects. CK0700000

Hydrobromide: [6415-90-3]

Mp 163-164°.

Sulfate (2:1): Atropine sulfate, JAN, USAN. Atropisol

[55-48-1]

Cryst. Mp 190-194°. Component of Antrocol, Colonaïd, Donnagel, Kinesed, Lomotil and Hydrapred.

- ▶ Human systemic effects. LD₅₀ (rat, orl) 600 mg/kg. Exp. reprod. and teratogenic effects. CK2450000

Picrate: Mp 175-176°.

Methobromide: 8-Methylatropinium bromide. Mydrasine

[2870-71-5]

C₁₈H₂₆BrNO₃ 384.312

Cryst. Mp 222-223°.

Methonitrate: Methyلاتropine nitrate, USAN. Atropine methonitrate, BAN, INN, JAN. Metropine. Harvatrate. Metanite. Ekomine. Eumydrin

[52-88-0]

C₁₈H₂₆N₂O₆ 366.413

Mp 166-168°.

- ▶ CK2800000

N-Oxide: Aminoxytropine tropate. Atropine oxide, INN. Genatropine

[4438-22-6]

C₁₇H₂₃NO₄ 305.373

Mp 127-128°. Dec. at 135°.

- ▶ YM6105000

N-Oxide; hydrochloride: Atropine oxide hydrochloride, USAN. Atropigen. Tropinox. Xtro

[4574-60-1]

Prisms (EtOH). Mp 192-193°.

N-De-Me: Noratropine

[16839-98-8]

C₁₆H₂₁NO₃ 275.347

Used in treatment of peptic ulcers.

Cryst. (Me₂CO). Mp 113-114°.

Propanoyl: Prampine, BAN, INN. Atropine propionate. Tropan-3-yl-O-propionyl tropate. PAMN

[7009-65-6]

C₂₀H₂₇NO₄ 345.438

Used in treatment of peptic ulcers.

(ξ)-form

O-Formyl: O-Formyltropine tropate

[645419-61-0]

C₁₈H₂₃NO₄ 317.384

Alkaloid from *Datura ceratocaula*.

Stereochem. not studied.

O-Ac: O-Acetyltropine tropate

[16655-60-0]

C₁₉H₂₅NO₄ 331.411

Alkaloid from *Datura ceratocaula* and *Datura stramonium*.

[620-61-1, 56362-77-7, 56362-78-8, 31610-87-4, 5908-99-6]

Warren, R.E. et al., *Rep. Lab. Am. Med. Assoc.*, 1921, **14**, 65-67; *CA*, **16**, 2580 (methobromide)

Leete, E. et al., *Can. J. Chem.*, 1954, **32**, 1116-1123; 1955, **33**, 1853-1854 (isol, biosynth)

Staub, H. et al., *Helv. Chim. Acta*, 1962, **45**, 2297-2305 (Mandragorine)

Liebisch, H.W. et al., *Annalen*, 1969, **721**, 163-167 (biosynth)

Cannon, J.R. et al., *Aust. J. Chem.*, 1969, **22**, 221-227 (*Anthocercis constits*, pmr)

Baralle, F.E. et al., *Chem. Comm.*, 1969, 721 (biosynth)

O'Donovan, D.G. et al., *J.C.S. Perkin I*, 1969, 223-226 (biosynth)

Fales, H.M. et al., *J.A.C.S.*, 1970, **92**, 1590-1597 (ms)

Takeuchi, Y. et al., *Chem. Pharm. Bull.*, 1971, **19**, 2603-2608 (synth)

Kussäther, E. et al., *Acta Cryst. B*, 1972, **28**, 2896-2899 (cryst struct)

Analyst (London), 1972, **97**, 740-756 (microanal)

Innes, I.R. et al., *Pharmacol. Basis Ther.*, 5th edn., 1975, 514 (rev. pharmacol)

Phillipson, J.D. et al., *Phytochemistry*, 1975, **14**, 999-1003 (oxides)

Sternberg, V.I. et al., *J. Het. Chem.*, 1977, **14**, 225-230 (cmr)

Pfister, J.R. et al., *J.O.C.*, 1978, **43**, 4373-4374 (*Noratropine synth*)

Shutt, L.E. et al., *Anaesthesia*, 1979, **34**, 476-490 (rev. pharmacol)

Kanto, J. et al., *Int. J. Clin. Pharmacol., Ther. Toxicol.*, 1983, **21**, 92-94 (rev)

Al-Badr, A.A. et al., *Anal. Profiles Drug Subst.*, 1985, **14**, 325-389 (rev. uv, ir, pmr, cmr, ms, anal)

Seeger, R. et al., *Dtsch. Apoth. -Ztg.*, 1986, **126**, 1930-1934 (rev. pharmacol)

Muhtadi, F.J. et al., *Anal. Profiles Drug Subst.*, 1994, **23**, 153-228 (*Hyoscyamine*, rev)

Negwer, M. et al., *Organic-Chemical Drugs and their Synonyms*, 7th edn., Akademie-Verlag, 1994, 5975; 5976 (synonyms)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 455; 457; 464; 468

Philipov, S. et al., *Z. Naturforsch., C*, 2002, **57**, 559-561 (O-Ac)


Berkov, S. et al., *Z. Naturforsch., C*, 2003, **58**, 455-458 (O-formyl)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, ARR000; HOU000; MGR500; ARR500

Trunkamide A

[181758-83-8]

T-633



Absolute Configuration

C₄₃H₆₃N₇O₈S 838.079

Cyclic heptapeptide. Isol. from an ascidian *Lissoclinum* sp. Shows promising antitumor activity. Oil. [α]_D²⁰ -231 (c, 0.06 in CHCl₃). λ_{max} 248 (ε 14300) (EtOH).


Carroll, A.R. et al., *Aust. J. Chem.*, 1996, **49**, 659-667 (isol, uv, ir, pmr, cmr, ms)

Pat. Coop. Treaty (WIPO), 1997, 039 025; *CA*, **127**, 290919m

Wipf, P. et al., *J.O.C.*, 2000, **65**, 1037-1049 (synth, abs config)

Caba, J.M. *et al.*, *J.O.C.*, 2001, **66**, 7568-7574 (synth)
 Salvatella, X. *et al.*, *J.O.C.*, 2003, **68**, 211-215 (pmr, struct)
 McKeever, B. *et al.*, *Tetrahedron*, 2003, **59**, 2713-2727 (synth)

α-Truxilline T-634
 γ-Isotropylcocaine. Cocamine
 [490-17-5]




C₃₈H₄₆N₂O₈ 658.79
 Ester of α-truxillic acid with ecgonine methyl ester. Alkaloid from the leaves of *Erythroxylum coca*. Obt. as a mixt. with β-Truxilline. Obt. pure by synth. (Erythroxylaceae). Mp 63° (sinters). The mixt. with β-Truxilline was also known as Cocamine. Hydrochloride: Mp 274°.

- 1-Epimer: γ-Truxilline**
 [113350-56-4]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.
- 2-Epimer: epi-Truxilline. Epitruixilline**
 [113351-64-7]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.
- 1,2-Diepimer: peri-Truxilline**
 [113350-53-1]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.
- 2,3-Diepimer: ε-Truxilline**
 [113350-55-3]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

Liebermann, C. *et al.*, *Ber.*, 1889, **22**, 130; 680 (synth)
 Hesse, O. *et al.*, *J. Prakt. Chem.*, 1902, **66**, 401 (bibl)
 Novak, M. *et al.*, *J. Ethnopharmacol.*, 1984, **10**, 261-274 (rev)
 Moore, J.M. *et al.*, *J. Chromatogr.*, 1987, **410**, 297-318; 1990, **504**, 391-401 (isol, chromatog)

β-Truxilline T-635
 Isococamine. β-Truxinic acid bis(2-carbomethoxypropyl) ester. δ-Isotropylcocaine
 [490-15-3]



C₃₈H₄₆N₂O₈ 658.79
 Alkaloid from the leaves of *Erythroxylum coca* (Erythroxylaceae). Amorph. solid. Mp 45° (sinters). [α]_D -29.3.
1-Epimer: neo-Truxilline. Neotruixilline
 [113350-54-2]

C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

3-Epimer: ξ-Truxilline
 [113350-58-6]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

1,2-Diepimer: ω-Truxilline
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

1,3-Diepimer: μ-Truxilline
 [113350-57-5]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

1,4-Diepimer: δ-Truxilline
 [113350-52-0]
 C₃₈H₄₆N₂O₈ 658.79
 Constit. of *Erythroxylum coca*.

Liebermann, C. *et al.*, *Ber.*, 1889, **22**, 680 (synth)
 Hesse, O. *et al.*, *Annalen*, 1892, **271**, 180; 1893, **272**, 238 (struct)
 Hesse, O. *et al.*, *J. Prakt. Chem.*, 1902, **66**, 401 (bibl)
 Novak, M. *et al.*, *J. Ethnopharmacol.*, 1984, **10**, 261-274 (rev)
 Moore, J.M. *et al.*, *J. Chromatogr.*, 1987, **410**, 297-318; 1990, **504**, 391-401 (isol, chromatog)

Trypanothione T-636

N⁷,N⁸-Bis(glutathionyl)spermidine. Trypanothione disulfide
 [96304-42-6]
 Glu-Cys-Gly-NH(CH₂)_nNH(CH₂)₃NH-Gly-Cys-Glu (n = 3)
 C₂₇H₄₇N₉O₁₀S₂ 721.855
 Reduced form shown. Metab. of *Crithidia fasciculata* and *Trypanosoma brucei*. Glutathione reductase cofactor. Trypanocidal agent.
Homologue: Homotrypanothione
 [159240-52-5]
 C₂₈H₄₉N₉O₁₀S₂ 735.881
 Metab. of *Trypanosoma cruzi*. Has n = 5.

Fairlamb, A.H. *et al.*, *Science (Washington, D.C.)*, 1985, **227**, 1485-1487 (isol, synth)
 Shames, S.L. *et al.*, *Biochemistry*, 1986, **25**, 3519 (biochem)
 Henderson, G.B. *et al.*, *J.C.S. Perkin 1*, 1990, 911-914 (synth, pmr, cmr, N-15 nmr)
 Fairlamb, A.H. *et al.*, *Annu. Rev. Microbiol.*, 1992, **46**, 695-729 (rev)
 Fauchet, V. *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 2559-2562 (synth)
 Hunter, K.J. *et al.*, *Eur. J. Biochem.*, 1994, **226**, 1019-1027 (Homotrypanothione)
 Krauth-Siegel, R.L. *et al.*, *Methods Enzymol.*, 1995, **251**, 287-294 (rev, isol, detm)

Trypargine T-637

3-[(2,3,4,9-Tetrahydro-1H-pyrido[3,4-b]indol-1-yl)propyl]guanidine, 9CI. 1-(3-Guanidinopropyl)-1,2,3,4-tetrahydro-β-carboline
 (S)-form

C₁₅H₂₁N₅ 271.364
(S)-form [82054-21-5]
 [82372-68-7]
 Obt. from the skin of the African frog *Kassina senegalensis*. Needles (as hydrochloride). Mp 211-212° (hydrochloride). [α]_D¹⁵ +37.2 (c, 0.54 in MeOH) (hydrochloride).
 ▶ Highly toxic.

6-Hydroxy: 6-Hydroxytrypargine
 C₁₅H₂₁N₅O 287.364
 Alkaloid from the venom of the spider *Parawixia bistriata*. λ_{max} 224 (log ε 4.52); 275 (log ε 3.71); 289 (log ε 3.42) (MeOH).

(±)-form [82264-57-1]
 [82264-59-3, 82264-58-2]
 Alkaloid from a *Eudistoma* sp. Pale yellow prisms (MeOH/Et₂O) (as hydrochloride). Mp 202-206° (hydrochloride).
1,2-Didehydro: Trypargimine
 [228113-43-7]
 C₁₅H₁₉N₅ 269.349
 Alkaloid from a *Eudistoma* sp. Light yellow solid. λ_{max} 206 (log ε 4.5); 236 (log ε 4.3); 254 (log ε 4); 316 (log ε 4.3); 348 (log ε 3.9) (MeOH).

Akizawa, T. *et al.*, *Biomed. Res.*, 1982, **3**, 232-234 (isol, pmr, cmr, ms)
 Shimizu, M. *et al.*, *Chem. Pharm. Bull.*, 1982, **30**, 909-914; 3453-3456; 4529-4533; 1984, **32**, 1313-1325 (synth, resoln, cryst struct, abs config)
 Van Wagoner, R.M. *et al.*, *J. Nat. Prod.*, 1999, **62**, 794-797 (Trypargimine)
 Cesar, L.M.M. *et al.*, *Helv. Chim. Acta*, 2005, **88**, 796-801 (pmr, cmr, 6-Hydroxytrypargine)
 Czarnocki, S.J. *et al.*, *Tetrahedron*, 2008, **64**, 3176-3182 (synth)

Tryprostatin A T-638

[171864-80-5]

 Absolute Configuration


C₂₂H₂₇N₃O₃ 381.474
 Prod. by the marine-derived *Aspergillus fumigatus*. Mammalian cell cycle inhibitor. Mycotoxin. Microtubule polym. inhibitor. Pale yellow cryst. Sol. MeOH, EtOAc, Me₂CO. Mp 120-123°. [α]_D²⁷ -69.7 (c, 0.7 in CHCl₃). λ_{max} 227 (ε 24540); 270 (ε 5450); 297 (ε 6590) (MeOH).

Demethoxy: Tryprostatin B
 [179936-52-8]
 C₂₁H₂₅N₃O₂ 351.447
 Prod. by the marine-derived *Aspergillus fumigatus*. Cell cycle progression and microtubule polym. inhibitor. Mycotoxin. Pale yellow cryst. Sol. MeOH, Me₂CO, EtOAc. Mp 102-105°. [α]_D²⁷ -71.1 (c, 0.6 in CHCl₃). λ_{max} 226 (23780); 277 (8690); 298 (sh) (7180) (MeOH).

- Cui, C.-B. *et al.*, *J. Antibiot.*, 1995, **48**, 1382-1384; 1996, **49**, 527-533; 534-540 (*isol. uv. ir. props*)
- Gan, T. *et al.*, *J.O.C.*, 1997, **62**, 9298-9304 (*synth*)
- Kondoh, M. *et al.*, *J. Antibiot.*, 1998, **51**, 801-804 (*activity*)
- Zhao, S. *et al.*, *Tet. Lett.*, 1998, **39**, 7009-7012 (*synth*)
- Schkeryantz, J.M. *et al.*, *J.A.C.S.*, 1999, **121**, 11964-11975 (*synth*)
- Wang, B. *et al.*, *Tet. Lett.*, 2001, **42**, 1463-1466 (*Tryprostatin B, synth*)
- Caballero, E. *et al.*, *J.O.C.*, 2003, **68**, 6944-6951 (*Tryprostatin B, synth*)
- Cardoso, A.S.P. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3966-3972 (*Tryprostatin B, synth*)

Tryptamine**T-639**

1H-Indole-3-ethanamine, 9CI. 3-(2-Aminoethyl)indole. 2-(3-Indolyl)ethylamine [61-54-1]



$C_{10}H_{12}N_2$ 160.218

Occurs widely in plants, esp. *Acacia* spp., *Lens esculenta* (lentil) and *Prosopis juliflora* (Fabaceae) and the fungi *Panaeolus foenicicii* and *Coprinus micaceus* (glistening ink cap). Also obt. from the gorgonian *Paramuricea chamaeleon*. Shows psychotropic effects. Needles (petrol). V. spar. sol. Et₂O, CHCl₃, C₆H₆. Mp 118°. Bp_{0.15} 137°. pK_{a1} -6.31; pK_{a2} 10.2 (25°). Log P 1.27 (calc). Metab. precursor of 1H-Indole-3-acetic acid, I-81.

- LD₅₀ (rat, ipr) ca. 223 mg/kg. NL4020000
Hydrochloride: [343-94-2]

Needles (EtOH/EtOAc). Mp 248°.

- LD₅₀ (mus, ipr) 197 mg/kg. NL4375000

Picrate:

Dark red prisms (Et₂O). Mp 247° dec.

N^b-Ac: N^b-*Acetyltryptamine*

[1016-47-3]

$C_{12}H_{14}N_2O$ 202.255

Alkaloid from leaves of *Prosopis nigra*. Prod. by the marine-derived bacteria *Cytophaga marinoflava* sp. AM13.1, *Roseivirga echinicomitans* KMM 6058^T and marine-derived fungus *isol.* from *Gracilaria verrucosa*. Mp 77°.

N^b-Di-Ac: N^b-*Diacetyltryptamine*

[934391-06-7]

$C_{14}H_{16}N_2O_2$ 244.293

Isol. from the marine bacterium *Roseivirga echinicomitans* KMM 6058^T. Cryst. λ_{max} 274 ; 281 ; 290 (MeOH).

N^b-(3-Methylbutanoyl): *Madugin*

[179027-43-1]

$C_{15}H_{20}N_2O$ 244.336

Alkaloid from the leaves of *Clausena indica*. Prod. by *Cytophaga marinoflava* sp. AM13.1. Oil. λ_{max} 224 ; 274 (sh) ; 283 ; 292 (Et₂O).

N^b-(3-Methyl-2-oxopentanoyl): *Nemato-pin*. N-[2-(1H-Indol-3-yl)ethyl]-3-methyl-2-oxopentanamide, 9CI

[183314-23-0]

[183314-24-1]

$C_{16}H_{20}N_2O_2$ 272.346

Prod. by *Xenorhabdus nematophilus*. Exhibits antibacterial and antifungal activities. Shows strong anti-MRSA activity. Powder. [α]_D²⁵ +0.1 (c, 0.6 in CHCl₃).

N^b-(3-Methyl-2Z-dodecenoyl): N^b-(3-Methyl-2-dodecenoyl)tryptamine.

Granulatamide A

[881417-85-2]

$C_{23}H_{34}N_2O$ 354.534

Constit. of *Eunicella granulata*. Cytotoxic. Pale yellow oil.

N^b-(3,5-Dimethyl-2Z,4E-dodecadienoyl):

N^b-(3,5-Dimethyl-2,4-dodecadienoyl)tryptamine. *Granulatamide B*

[881417-94-3]

$C_{24}H_{34}N_2O$ 366.545

Constit. of *Eunicella granulata*. Cytotoxic. Pale yellow oil.

N^b-Hexadecanoyl: N^b-*Hexadecanoyltryptamine*. N^b-*Palmitoyltryptamine*

[21469-15-8]

$C_{26}H_{42}N_2O$ 398.631

Alkaloid from seeds of *Annona reticulata* (custard apple) and *Rollinia mucosa* (biriba).

N^b-Octadecanoyl: N^b-*Octadecanoyltryptamine*. N^b-*Stearoyltryptamine*

[21469-14-7]

$C_{28}H_{46}N_2O$ 426.684

Alkaloid from seeds of *Annona reticulata* (custard apple) and *Rollinia mucosa* (biriba) (Annonaceae).

N^b-Nonadecanoyl: N^b-*Nonadecanoyltryptamine*

$C_{29}H_{48}N_2O$ 440.711

Alkaloid from the seeds of *Annona atemoya* (custard apple).

N^b-Eicosanoyl: N^b-*Eicosanoyltryptamine*. N^b-*Arachidoyltryptamine*

[152766-92-2]

$C_{30}H_{50}N_2O$ 454.738

Alkaloid from seeds of *Annona reticulata* (custard apple) and *Rollinia mucosa* (biriba) (Annonaceae).

N^b-Docosanoyl: N-[2-(1H-Indol-3-yl)ethyl]docosanamide. 3-[2-(Docosanoylaminoethyl)]indole. Behenic acid tryptamide. N^b-*Behenoyltryptamine*. N^b-*Docosanoyltryptamine*

[7367-79-5]

$C_{32}H_{54}N_2O$ 482.791

Alkaloid from seeds of *Annona reticulata* (custard apple) and *Rollinia mucosa* (biriba) (Annonaceae). Cryst. (Me₂CO/Me₂CO). Mp 121-123° (116-117°).

N^b-Tricosanoyl: N^b-*Tricosanoyltryptamine*

[152766-93-3]

$C_{33}H_{56}N_2O$ 496.818

Alkaloid from seeds of *Annona reticulata* and *Rollinia mucosa*.

N^b-Tetracosanoyl: N^b-*Tetracosanoyltryptamine*. N^b-*Lignoceroyltryptamine*

[152766-94-4]

$C_{34}H_{58}N_2O$ 510.845

Alkaloid from seeds of *Annona reticulata* and *Rollinia mucosa* cocoa shell.

Mp 120-123°.

N^b-Pentacosanoyl: N^b-*Pentacosanoyltryptamine*

[152766-95-5]

$C_{35}H_{60}N_2O$ 524.872

Alkaloid from seeds of *Annona reticulata* and *Rollinia mucosa*.

N^b-Hexacosanoyl: N^b-*Hexacosanoyltryptamine*. N^b-*Cerotoyltryptamine*

[152766-96-6]

$C_{36}H_{62}N_2O$ 538.899

Alkaloid from seeds of *Annona reticulata* and *Rollinia mucosa*.

N^b-Octacosanoyl: N^b-*Octacosanoyltryptamine*

$C_{38}H_{66}N_2O$ 566.952

Constit. of the seeds of *Annona atemoya* (custard apple).

N^b-(25Z-Triacontenoyl): N^b-(25-Triacontenoyl)tryptamine. *Cheritamine*

[223696-37-5]

$C_{40}H_{68}N_2O$ 592.99

Alkaloid from *Annona cherimola* (cherimoya). Powder (MeOH). Mp 121-123°. λ_{max} 222 ; 280 (EtOH).

N^b-(2S-Hydroxy-3S-methylpentanoyl):

N^b-(2-Hydroxy-3-methylpentanoyl)tryptamine. *Dihydroneumatophin*

[184031-87-6]

$C_{16}H_{22}N_2O_2$ 274.362

Metab. from the culture broth of *Xenorhabdus nematophilus*. [α]_D²⁵ -0.2 (c, 0.32 in CHCl₃). Config. revised in 2003.

N^b-(7S-Methoxy-4E-tetradecenoyl): N-

(7-Methoxy-4-tetradecenoyl)tryptamine. *Hermitamide B*

$C_{25}H_{38}N_2O_2$ 398.587

Alkaloid from *Lynghya majuscula*. Cytotoxic agent. Pale yellow oil. [α]_D²⁶ -4.5 (c, 0.1 in CHCl₃). λ_{max} 224 (ε 13600); 282 (ε 3600); 292 (ε 3000) (EtOH).

N^b-Benzoyl: N-[2-(1H-Indol-3-yl)ethyl]benzamide, 9CI. *Benztryptamide*

[4753-09-7]

$C_{17}H_{16}N_2O$ 264.326

Prisms. Mp 137-138°.

N^b-(2-Methylaminobenzoyl): N-[2-(1H-Indol-3-yl)ethyl]-2-(methylamino)benzamide, 9CI. N^b-(2-Methylaminobenzoyl)tryptamine

[72502-82-0]

$C_{18}H_{19}N_3O$ 293.368

Alkaloid from the fruits of *Evodia rutaecarpa*. Needles (C₆H₆), cryst. (toluene). Mp 126-127° (114-117°). λ_{max} 223 (log ε 4.4); 259 (log ε 3.93); 284 (log ε 3.47); 292 (log ε 3.45); 346 (log ε 3.55) (EtOH).

N^b-(4-Hydroxycinnamoyl): N^b-*p-Coumaroyltryptamine*

[53905-12-7]

$C_{19}H_{18}N_2O_2$ 306.363

Found in kernels of sweet corn (*Zea mays*).

N^b-(4-Hydroxy-3-methoxycinnamoyl):

N^b-*Feruloyltryptamine*

[53905-13-8]

$C_{20}H_{20}N_2O_3$ 336.39

Found in kernels of sweet corn (*Zea*

mays).

N^b-Me: see *N*-Methyltryptamine, M-582

N^b,N^b-Di-Me: see *N,N*-Dimethyltryptamine, D-785

[54268-28-9]

Aldrich Library of FT-IR Spectra, 1st edn., 1985, **2**, 660C; 660D (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, **3**, 131C (nmr)

Ewins, A.J. *et al.*, *J.C.S.*, 1911, **99**, 270-273 (synth)

White, E.P. *et al.*, *N.Z. J. Sci. Technol., Sect. B*, 1944, **25**, 137-162; 1951, **33**, 54-60; 1957, **38**, 718-725 (isol, occur)

Cohen, L.A. *et al.*, *J.A.C.S.*, 1960, **82**, 2184-2187 (pmr)

Jackson, A.H. *et al.*, *J.C.S.*, 1965, 3498-3500 (synth, struct)

Sacher, H. *et al.*, *Z. Lebensm.-Unters.-Forsch.*, 1965, **128**, 264-267 (docosanoyl)

Kametani, T. *et al.*, *Synthesis*, 1972, 475 (synth)

Wakahara, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1973, **46**, 2481-2486 (cryst struct)

Ehmann, A. *et al.*, *Phytochemistry*, 1974, **13**, 1979-1983 (N^b-p-Coumaroyltryptamine, N^b-Feruloyltryptamine)

Cimino, G. *et al.*, *Comp. Biochem. Physiol., C: Comp. Pharmacol.*, 1978, **61**, 361-362 (isol)

Bergman, J. *et al.*, *J.O.C.*, 1985, **50**, 1246-1255 (N-2-methylaminobenzoyl, synth)

Morales-Rios, M.S. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 377-395 (cmr)

Shoji, N. *et al.*, *J. Nat. Prod.*, 1988, **51**, 161-162 (N-2-methylaminobenzoyl, isol, ir, uv, pmr, cmr, ms)

Szelag, H. *et al.*, *Nahrung*, 1988, **32**, 285-290 (N-docosanoyl)

Maeda, U. *et al.*, *Phytochemistry*, 1993, **34**, 1633-1635 (N-Acyltryptamines)

Li, J. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1157-1158 (Dihydronematophin)

Li, J. *et al.*, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 1349-1355 (Nematophin, synth)

Li, J. *et al.*, *Can. J. Microbiol.*, 1997, **43**, 770-773 (Nematophin)

Riemer, B. *et al.*, *Phytochemistry*, 1997, **45**, 337-341 (Madugin)

Himmler, T. *et al.*, *Bioorg. Med. Chem. Lett.*, 1998, **8**, 2045-2049 (Nematophin, activity)

Chen, C.Y. *et al.*, *J. Chin. Chem. Soc. (Taipei)*, 1999, **46**, 77-86 (Cheritamine)

Chavez, D. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1119-1122 (N-Acyltryptamines)

Li, J. *et al.*, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 1751-1754 (Nematophin, activity)

Tan, L.T. *et al.*, *J. Nat. Prod.*, 2000, **63**, 952-955 (Hermitamide B)

Li, Y. *et al.*, *Arch. Pharmacol. Res.*, 2003, **26**, 21-23 (N^b-Ac, marine, isol)

Paik, S. *et al.*, *Bull. Korean Chem. Soc.*, 2003, **24**, 623-626 (Dihydronematophin, Nematophin)

Czerwinski, K.M. *et al.*, *Synth. Commun.*, 2003, **33**, 1225-1231 (N^b-benzoyl)

Shabaan, M. *et al.*, *Dissertation*, Univ. of Göttingen, 2004, (Cytophaga marinoflava derivs)

Wu, Y.-C. *et al.*, *J. Nat. Prod.*, 2005, **68**, 406-408 (N-Acyltryptamines)

Schmitt, H.M. *et al.*, *J.A.C.S.*, 2005, **127**, 10356-10364 (conformm)

Oleinikova, G.K. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2006, **42**, 713-717 (N-Acyltryptamines)

Hug, B. *et al.*, *J. Agric. Food Chem.*, 2006, **54**, 3199-3203 (N-docosanoyl, N-tetracosanoyl, detn, occur)

Reyes, F. *et al.*, *J. Nat. Prod.*, 2006, **69**, 668-670 (Granulatamides)


Virolleaud, M.-A. *et al.*, *Tet. Lett.*, 2006, **47**, 5127-5130 (Hermitamide B, synth)

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials, 8th edn.*, Van Nostrand Reinhold, 1992, AJX000; AJX250

Tryptophan, 9CI, 8CI, USAN

T-640

2-Amino-3-(3-indolyl)propanoic acid. α -Amino-1H-indole-3-propanoic acid. Indole-3-alanine. 3-Indolylalanine. Trp

C₁₁H₁₂N₂O₂ 204.228

In CAS the N atoms labelled a and b are referred to as N- and I- respectively.

(S)-form

L-form

[73-22-3]

Constit. of many plants. Enzymatic hydrol. prod. of most plant and animal proteins. Dietary supplement, nutrient, antidepressant. Plates (EtOH aq.). Sol. boiling H₂O, hot EtOH; spar. sol. cold H₂O, EtOH; insol. CHCl₃. Soly. 1.14 g/100 g in H₂O at 25°. Mp ca. 282° dec. [α]_D²⁵ -68.8 (H₂O). [α]_D²⁰ -32.5 (c, 1.1 in H₂O). [α]_D²¹ -5.7 (5M HCl). pK_{a1} 2.46; pK_{a2} 9.41; pK_{a3} 16.82 (25°.NH, KOH aq.). pK_a 2.38 (COOH, α -NH₃⁺). Log P -1.57 (uncertain value) (calc). Isoelectric point 5.89. Bitter taste.

► LD₅₀ (rat, ipr) 1634 mg/kg. Exp. reprod. and teratogenic effects (large dose). Adverse effects when used therapeutically. YN6130000

Hydrochloride: [6159-33-7]

Needles (MeOH). Mp 257° dec.

Picrate:

Red needles and plates. Mp 195-196° slight dec.

N^a-Hexanoyl, N^b-inosityl: [124772-47-0]C₂₃H₃₂N₂O₈ 464.514

Constit. of green gram flowers (*Phaseolus aureus*). Antihyperglycaemic agent. Exact struct. not indicated.

N^a-(Carboxyacetyl): N-Malonyltryptophan

[29399-11-9]

C₁₄H₁₄N₂O₅ 290.275

Constit. of various plants incl. *Lycopersicon esculentum* (tomato), *Malus esculentum*, *Medicago sativa* (alfalfa) and *Melilotus albus* (white melilot). Originally assigned to the D-form.

N^a-(4-Aminobenzoyl): N^a-(4-Aminobenzoyl)tryptophan

[39545-05-6]

C₁₈H₁₇N₃O₃ 323.351

Isol. from the myxomycete *Fuligo candida*. Mp 213-214°. [α]_D²³ -6.8 (c, 0.76 in MeOH). λ _{max} 282 (ε 12000) (MeOH).

N^a-(4-Hydroxycinnamoyl): N^z-p-HydroxycoumaroyltryptophanC₂₀H₁₈N₂O₄ 350.373Constit. of green coffee beans (*Coffea*

canephora var. *robusta*) (Rubiaceae). Amorph. powder. Mp 208° dec.

N^a-(3,4-Dihydroxycinnamoyl): N-Caffeoyltryptophan

[109163-69-1]

C₂₀H₁₈N₂O₅ 366.373

Constit. of green coffee beans (*Coffea canephora*) (Rubiaceae). Amorph. powder.

N^a-(4-Amino-E-cinnamoyl):C₂₀H₁₉N₃O₃ 349.388

Prod. by *Fuligo aurea*. Amorph. solid. [α]_D²⁵ +49 (c, 0.7 in MeOH). λ _{max} 222 (log ε 4.2); 291 (log ε 3.8); 325 (log ε 3.8) (MeOH).

N^a-(4-Amino-Z-cinnamoyl):C₂₀H₁₉N₃O₃ 349.388

Prod. by *Fuligo aurea*. Amorph. solid. [α]_D²³ +97 (c, 0.8 in MeOH). λ _{max} 222 (log ε 4.3); 291 (log ε 3.9); 325 (log ε 3.9) (MeOH).

N^a,N^a-Di-Me: N,N-Dimethyltryptophan

[2812-40-0]

C₁₃H₁₆N₂O₂ 232.282

Isol. from *Abrus precatorius*, *Desmodium triflorum* and *Erythrina lithosperma*. Plant growth inhibitor. Small prisms (EtOH aq.). Mp 243-245°.

N^a,N^a-Di-Me, Me ester: [35214-77-8]C₁₄H₁₈N₂O₂ 246.308

Alkaloid present in *Pultenaea altissima*, seeds of *Abrus precatorius*, *Erythrina variegata* and others. Plant growth inhibitor. Needles (hexane). Mp 97-98°. [α]_D²⁰ +65 (c, 2 in EtOH). [α]_D²⁰ +70 (c, 0.5 in dil. HCl). λ _{max} 203; 222; 283; 290 (EtOH).

N^a,N^a,N^a-Tri-Me, Me ester: N,N-Dimethyltryptophan methocation methyl ester

[33440-04-9]

C₁₅H₂₁N₂O₂⁺ 261.343

Isol. from the seeds of *Abrus precatorius*.

N^a-(4-Carboxy-3-hydroxy-3-methylbutanoyl): N-(3-Hydroxy-3-methylglutaryl)tryptophan. Achillamide

[223924-17-2]

C₁₇H₂₀N₂O₆ 348.355


Constit. of the roots of *Astragalus trojanus*. [α]_D²⁵ -29 (c, 0.1 in MeOH). λ _{max} 221; 270; 281; 290 (MeOH).

[7401-26-5, 3184-74-5]

Ghosal, S. *et al.*, *Phytochemistry*, 1971, **10**, 195; 3312 (Dimethyltryptophan methocation methyl ester)Mandava, N. *et al.*, *Phytochemistry*, 1974, **13**, 2853 (Dimethyltryptophan)Gamburg, K.Z. *et al.*, *Plant Sci. (Limerick, Ire.)*, 1991, **77**, 149 (N-Malonyltryptophan)Sakagami, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 1362 (N-Malonyltryptophan, abs config)Murata, M. *et al.*, *Biosci., Biotechnol., Biochem.*, 1995, **59**, 1887 (cinnamoyltryptophans)Bedir, E. *et al.*, *J. Nat. Prod.*, 1999, **62**, 563-568 (Achillamide)Nakatani, S. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 368-370 (N-4-Aminobenzoyltryptophan)Hosoya, T. *et al.*, *Heterocycles*, 2006, **69**, 463-468 (4-aminocinnamates)

Tryptophan tryptophylquinone T-641

α, α' -Diamino-6',7'-dihydro-6',7'-dioxo-[2,4'-bi-1H-indole]-3,3'-dipropionic acid, 9CI. T7Q [134645-25-3]



$C_{22}H_{20}N_4O_6$ 436.423

Amino acid derived redox cofactor isol. from bacterial methylamine dehydrogenases. Very difficult to isolate.

McIntire, W.S. *et al.*, *Science (Washington, D.C.)*, 1991, **252**, 817-824 (isol)


Hartmann, C. *et al.*, *Methods Enzymol.*, 1995, **258**, 149-164; 1997, **280**, 98-150 (rev, biosynth)

Itoh, S. *et al.*, *Nat. Prod. Rep.*, 1995, **12**, 45-53 (rev)

Tryptophanbetaxanthin T-642

[419555-21-8]

[1007894-71-4]




$C_{20}H_{19}N_3O_6$ 397.387

Yellow pigment from *Celosia argentea*. Also found in beet (*Beta vulgaris*).

Schliemann, W. *et al.*, *Phytochemistry*, 2001, **58**, 159-165 (isol, biosynth)

Stintzing, F.C. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 2302-2307 (occur)

Kugler, F. *et al.*, *Z. Naturforsch., C*, 2007, **62**, 311-318 (isol)

Tryptoquivaline A T-643

$C_{27}H_{26}N_4O_7$ 518.525

Lower homologue of Tryptoquivaline A,

Absolute Configuration

R = CH₃

T-645. Metab. of *Penicillium digitatum* on citrus fruits. $[\alpha]_D^{20}$ 0 (EtOH). λ_{max} 222 (log ϵ 4.48); 225 (log ϵ 4.49); 273 (sh) (log ϵ 3.9); 289 (sh) (log ϵ 3.69); 298 (sh) (log ϵ 3.54); 310 (sh) (log ϵ 3.39) (EtOH).

Ariza, M.R. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 6361-6365 (isol, cd, pmr, cmr)

Tryptoquivaline B T-644

As Tryptoquivaline A, T-643 with R = H


$C_{26}H_{24}N_4O_7$ 504.498

Metab. of *Penicillium digitatum* on citrus fruits. $[\alpha]_D^{22}$ +57.1 (c, 0.14 in EtOH).

Ariza, M.R. *et al.*, *J. Agric. Food Chem.*, 2002, **50**, 6361-6365 (isol, cd, pmr, cmr)

Tryptoquivaline A T-645

Tryptoquivaline C. Tryptoquivaline [55387-45-6]



Absolute Configuration

$C_{29}H_{30}N_4O_7$ 546.579

Metab. of *Aspergillus clavatus*. Tremorogenic toxin. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 155-157°. $[\alpha]_D^{25}$ +130 (c, 0.22 in CHCl₃). λ_{max} 228 (ϵ 37000); 275 (ϵ 8550); 305 (ϵ 3800); 317 (ϵ 3040) (EtOH) (Derep). λ_{max} 228 (ϵ 41200); 279 (ϵ 9500); 307 (ϵ 3700); 319 (ϵ 3000) (MeOH) (Berdy).

N-Deoxy: **Deoxytryptoquivaline**

[60676-57-5]

$C_{29}H_{30}N_4O_6$ 530.579

Prod. by *Aspergillus clavatus*. Needles (CH₂Cl₂/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 150-152°. $[\alpha]_D^{25}$ +56.8 (c, 0.78 in CHCl₃). λ_{max} 227 (ϵ 44500); 232 (sh) (ϵ 41900); 252 (sh) (ϵ 18500); 267 (sh) (ϵ 12000); 278 (sh) (ϵ 10300); 304 (ϵ 3300); 318 (sh) (ϵ 2700) (EtOH).

N-Deoxy, N-acetoxy, O-de-Ac: **Isotryptoquivaline**

[61897-83-4]

$C_{29}H_{30}N_4O_7$ 546.579

Metab. of *Aspergillus fumigatus*. Mp 215-217° dec. $[\alpha]_D^{28}$ +168 (c, 0.23 in CHCl₃).

O-De-Ac, 27-ketone: **Tryptoquivaline I**

[66180-23-2]

$C_{27}H_{26}N_4O_6$ 502.526

Isol. from *Aspergillus fumigatus*. Tremorogenic toxin. Leaflets (MeOH/CH₂Cl₂). Mp 232-235.5° dec. $[\alpha]_D^{14}$ +239 (c, 0.16 in CHCl₃). λ_{max} 235 (ϵ 31700); 250 (ϵ 21400); 292 (ϵ 9600); 321 (ϵ 6100) (MeOH) (Berdy).

► Toxic.

27-Epimer: **27-Epitryptoquivaline**

[182967-49-3]

$C_{29}H_{30}N_4O_7$ 546.579

Metab. from *Corynascus setosus*. Tremorogenic agent. Needles (MeOH). Mp

225-227°. $[\alpha]_D^{25}$ +138 (c, 0.02 in CHCl₃). λ_{max} 209 (sh) (log ϵ 4.67); 289 (sh) (log ϵ 3.83); 306 (sh) (log ϵ 3.54); 318 (sh) (log ϵ 3.44) (MeOH).

Clardy, J. *et al.*, *J.A.C.S.*, 1975, **97**, 663-665 (ir, uv, pmr, cryst struct)

Yamazaki, M. *et al.*, *Tet. Lett.*, 1976, **17**, 2861-2864 (Isotryptoquivaline)

Büchi, G. *et al.*, *J.O.C.*, 1977, **42**, 244-246 (Tryptoquivaline, Deoxytryptoquivaline, isol, uv, ir, pmr)

Yamazaki, M. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 111-117 (Tryptoquivaline I)

Springer, J.P. *et al.*, *Tet. Lett.*, 1979, **20**, 339-342 (abs config)

Hermkens, P.H.H. *et al.*, *Tetrahedron*, 1988, **44**, 1991-2000 (synth)


Fujimoto, H. *et al.*, *Chem. Pharm. Bull.*, 1996, **44**, 1843-1848 (27-Epitryptoquivaline)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 414

Tryptoquivaline F T-646

Fumitremorgin F. FTF

[61897-89-0]



$C_{22}H_{18}N_4O_4$ 402.409

Isol. from *Aspergillus fumigatus*. Tremorogenic toxin. Fine needles (MeOH). Mp 277° dec. $[\alpha]_D^{15.5}$ -109 (c, 0.006 in CHCl₃). λ_{max} 226 (ϵ 33400); 303 (ϵ 2800); 315 (ϵ 2200) (MeOH) (Berdy).

► Toxic.

N-Ac: [61897-90-3]

Needles (MeOH). Mp 280-283° dec. $[\alpha]_D^{12}$ -116 (c, 0.12 in CHCl₃). Stereochem. requires confirmation.

N¹⁶-Hydroxy: **Tryptoquivaline H. Fumitremorgin H. FTH**

[61949-67-5]

$C_{22}H_{18}N_4O_5$ 418.408

Isol. from *Aspergillus fumigatus*. Tremorogenic. Fine needles (MeOH). Mp 274° dec. $[\alpha]_D^{11}$ -155 (c, 0.02 in Me₂CO). λ_{max} 226 (ϵ 33100); 303 (ϵ 3100); 315 (ϵ 2500) (MeOH) (Berdy).

► Toxic.

N¹⁶-Acetoxy: [61897-88-9]

Prisms (MeOH/CH₂Cl₂). Mp 231-234°. $[\alpha]_D^{16.5}$ +243 (c, 0.0095 in Me₂CO).

12-Epimer: **Tryptoquivaline J. FTJ**

[66212-51-9]

$C_{22}H_{18}N_4O_4$ 402.409

Isol. from *Aspergillus fumigatus*. Tremorogenic. Fine needles (Me₂CO/MeOH). Mp 254-258° dec. $[\alpha]_D^{14}$ +135 (c, 0.024 in Me₂CO). λ_{max} 225 (ϵ 41000); 302 (ϵ 3900); 310 (ϵ 3100) (MeOH) (Berdy).

► Toxic.


12-Epimer, N¹⁶-hydroxy: **Tryptoquivaline E. Fumitremorgin E** [61897-87-8]

$C_{22}H_{18}N_4O_5$ 418.408
Metab. from *Aspergillus fumigatus*.
Tremorgenic toxin. Mp 257° dec. $[\alpha]_D^{25}$
+257 (c, 0.009 in $CHCl_3$). λ_{max} 225 (ε
32300); 303 (ε 3000); 315 (ε 2600)
(MeOH) (Berdy).

Yamazaki, M. *et al.*, *Chem. Pharm. Bull.*,
1978, **26**, 111-117 (*Tryptoquivalines E,F,H,J*)
Handbook of Secondary Fungal Metabolites,
(ed. Cole, R.J. *et al.*), Academic Press, 2003,
1, 391; 395; 399
Afiyatullo, S.S. *et al.*, *Chem. Nat. Compd.*
(*Engl. Transl.*), 2005, **41**, 236-238
(*Tryptoquivaline J, marine isol*)

Tryptoquivaline G
Fumitremorgin G
[61897-91-4]

T-647

Absolute
Configuration

$C_{23}H_{20}N_4O_5$ 432.435
Metab. of *Aspergillus fumigatus*. Myco-
toxin. Prisms (Me_2CO). Mp 240-241.5°.
 $[\alpha]_D^{25}$ +215 (c, 0.011 in Me_2CO). λ_{max} 226
(ε 34300); 302 (ε 3000); 315 (ε 2500)
(MeOH) (Berdy).

Ac: [61897-92-5]
Prisms ($CH_2Cl_2/MeOH$). Mp 231-
234°. $[\alpha]_D^{16.5}$ +243 (c, 0.0095 in
 Me_2CO).


*12-Epimer: Tryptoquivaline L. Fumitre-
morgin L*
[69483-20-1]
 $C_{23}H_{20}N_4O_5$ 432.435
Prod. by *Aspergillus fumigatus*. Leaf-
lets (Me_2CO). Mp 275-277° dec. $[\alpha]_D^{25}$ -
230 (c, 0.02 in Me_2CO).

Yamazaki, M. *et al.*, *Chem. Pharm. Bull.*,
1978, **26**, 111-117; 1979, **27**, 1611-1617
(*Tryptoquivalines G,L, isol, struct, ir, uv, ms,
pmr, cd*)

Büchi, G. *et al.*, *J.A.C.S.*, 1979, **101**, 5084-5086
(*Tryptoquivaline G, synth*)
Ohnuma, T. *et al.*, *Tet. Lett.*, 1981, **22**, 4969-
4972 (*±-Tryptoquivaline G, synth*)
Nakagawa, M. *et al.*, *J.A.C.S.*, 1983, **105**,
3709-3710 (*(+)-Tryptoquivaline G, (-)-
Tryptoquivaline G, synth, biosynth*)
Handbook of Secondary Fungal Metabolites,
(ed. Cole, R.J. *et al.*), Academic Press, 2003,
1, 393

Tscheschamine
[120186-26-7]

T-648




$C_{29}H_{36}N_4O_5$ 520.627
Alkaloid from the stem bark of *Zizyphus
sativa* (Rhamnaceae). Amorph. powder.
Mp 197-198°.

Shah, A.H. *et al.*, *Heterocycles*, 1988, **27**, 2777
(*isol, uv, ir, pmr, ms, struct*)

Tsilanine, 9CI
[29028-11-3]

T-649

Absolute
Configuration

$C_{22}H_{26}N_2O_3$ 366.459
Alkaloid from *Strychnos henningsii* twigs
(Loganiaceae). Cryst. (Me_2CO /hexane).
Mp 218-225°. $[\alpha]_D^{20}$ +61 ($CHCl_3$).

O-De-Me: O-Demethylsilanine
[29028-13-5]

$C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from *Strychnos henningsii*
twigs and leaves (Loganiaceae). Not
obt. pure.

10-Methoxy: 10-Methoxysilanine

$C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Strychnos henningsii*
twigs (Loganiaceae). Cryst. ($CHCl_3$ /
 Et_2O). Mp 160-170°. $[\alpha]_D^{20}$ +50
($CHCl_3$).


*10-Methoxy, O-de-Me: O-Demethyl-10-
methoxysilanine. 10-Methoxy-O-de-
methylsilanine*

[29028-14-6]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Strychnos henningsii*
twigs and leaves (Loganiaceae). Not
obt. pure.

Sarfati, R. *et al.*, *Phytochemistry*, 1970, **9**,
1107-1113 (*isol, uv, pmr, ms, struct*)
Wenkert, E. *et al.*, *J.O.C.*, 1978, **43**, 1099 (*cmr,
deriv*)

Tsitsikammamine A
[183114-86-5]

T-650



$C_{18}H_{13}N_3O_2$ 303.32
Related to Wakayin, W-3. Alkaloid from
the South African latruncid sponge
Tsitsikamma favus and another sponge.
Cytotoxic. Exhibits antimicrobial activ-
ity. λ_{max} 242 ; 317 ; 377 (MeOH) (Berdy).

N¹³-Me: Tsitsikammamine B
[183114-88-7]
 $C_{19}H_{15}N_3O_2$ 317.346

Alkaloid from *Tsitsikamma favus* and a
marine sponge. Cytotoxic. Exhibits
antimicrobial activity. λ_{max} 242 ; 317 ;
374 (MeOH) (Berdy).

*N¹⁸-Hydroxy: Tsitsikammamine A N¹⁸-
oxime*

$C_{18}H_{14}N_3O_3$ 320.327
Alkaloid from *Tsitsikamma favus*.
Bright orange-red solid.


*N¹⁸-Hydroxy, N¹³-Me: Tsitsikammamine
B N¹⁸-oxime*

$C_{19}H_{16}N_3O_3$ 334.354
Alkaloid from *Tsitsikamma favus*.
Bright orange-red solid.

Hooper, G.J. *et al.*, *Tet. Lett.*, 1996, **37**, 7135-
7138 (*isol, uv, ir, pmr, cmr*)
Antunes, E.M. *et al.*, *J. Nat. Prod.*, 2004, **67**,
1268-1276 (*oximes*)

Tsukushinamine A
[70711-82-9]

T-651

Absolute
configuration

$C_{15}H_{20}N_2O$ 244.336
Alkaloid from *Sophora franchetiana* (Fa-
baceae). Liq. $[\alpha]_D^{25}$ -72.3 (c, 0.56 in
 $EtOH$).

Hydrobromide: Mp 260°.

Δ³-Isomer: Tsukushinamine C
[73575-31-2]

$C_{15}H_{20}N_2O$ 244.336
Minor alkaloid from *Sophora fran-
chetiana* (Fabaceae). Oil. Opt. rotn. not
dtd.

14-Epimer: Tsukushinamine B
[73610-25-0]

$C_{15}H_{20}N_2O$ 244.336
Alkaloid from various parts of *So-
phora franchetiana* (Fabaceae). Oil.
 $[\alpha]_D^{25}$ -144.4 ($EtOH$).


Ohmiya, S. *et al.*, *Chem. Pharm. Bull.*, 1979,
27, 1055 (*isol, uv, ir, pmr, cmr, ms, struct*)

Bordner, J. *et al.*, *Chem. Pharm. Bull.*, 1980,
28, 1965 (*cryst struct*)

Ohmiya, S. *et al.*, *Phytochemistry*, 1981, **20**,
1997 (*derivs*)

Tubacetine
[144442-82-0]

T-652



R = Ph

$C_{29}H_{37}NO_7$ 511.614

Closely related to Acidissiminol, A-77. Alkaloid from aerial parts of *Haplophyl- lum tuberculatum* (Rutaceae). Microcryst. (Et₂O/hexane). Mp 71.7-72.7°. $[\alpha]_D^{22} +12$ (c, 0.1 in CHCl₃).

Al-Yahya, M.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 899 (isol, uv, ir, pmr, cmr, struct)

Tubasencine T-653

[144425-13-8]

As Tubacetine, T-652 with

R = -CH=C(CH₃)₂


C₂₇H₃₉NO₇ 489.608

Alkaloid from aerial parts of *Haplophyl- lum tuberculatum* (Rutaceae). Pale yellow gum. $[\alpha]_D^{25} +12$ (c, 0.1 in CHCl₃).

Al-Yahya, M.A. *et al.*, *J. Nat. Prod.*, 1992, **55**, 899 (isol, uv, ir, pmr, cmr, struct)

Tubastraine T-654

[114216-90-9]




C₃₀H₂₅Br₂NO₇ 671.338

Alkaloid from the non-symbiotic marine coral *Tubastrea micrantha*. Glassy compd. Mp 122°. First chromone isol. from a marine invertebrate. Possibly derived from dietary sources.

Alam, M. *et al.*, *Heterocycles*, 1988, **27**, 719 (isol, ir, pmr, cmr, ms, struct)

Tubastrindole C T-655

[524067-26-3]



C₂₈H₂₇N₇O₃ 509.566

Alkaloid from a stony coral *Tubastrea aurea*. $[\alpha]_D -20$ (c, 0.04 in MeOH).

11'-Imide: **Tubastrindole B**

[524067-25-2]

C₂₈H₂₈N₈O₂ 508.582

Alkaloid from *Smenospongia cerebriformis* and a *Tubastrea aurea*. $[\alpha]_D -26$ (c, 0.07 in MeOH).

6-Bromo, 11'-imide: **Tubastrindole A**

[524067-24-1]


C₂₈H₂₇BrN₈O₂ 587.478

Alkaloid from *Smenospongia cerebriformis* and *Tubastrea aurea*. $[\alpha]_D -38$ (c, 0.13 in MeOH). λ_{max} 284 (ε 2800); 294 (ε 2100) (no solvent reported).

Iwagawa, T. *et al.*, *Tet. Lett.*, 2003, **44**, 2533-2535 (isol, pmr, cmr)

Tubastrine T-656

2-[(3,4-Dihydroxyphenyl) ethenyl]guanidine, 9CI. 3,4-Dihydroxy-β-guanidinostyrene. β-(Aminoiminomethyl) amino-3,4-dihydroxystyrene [107585-47-7]



C₉H₁₁N₃O₂ 193.205

Constit. of *Aplidium orthium*, *Dendrodoa grossularia* and *Tubastrea aurea*. Shows antiviral props. Light yellow solid. Sol. Me₂CO, MeOH; fairly sol. EtOAc; poorly sol. H₂O. Mp 173-175°. λ_{max} 205 (log ε 3.79); 221 (log ε 3.79); 287 (log ε 3.79); 306 (sh) (log ε 3.68) (MeOH).

1,2-Dihydro: **7,8-Dihydrotubastrine**

C₉H₁₃N₃O₂ 195.221

Alkaloid from the sponge *Petrosia cf. contignata*. Brown viscous oil. λ_{max} 222 (ε 3900); 284 (ε 1900) (MeOH).

1,2-Dihydro, di-Me ether: 2-[(3,4-Dimethoxyphenyl) ethyl]guanidine, **7,8-Dihydro-3,4-di-O-methyltubastrine**

[46719-17-9]

C₁₁H₁₇N₃O₂ 223.274

Alkaloid from *Aplidium orthium*. Pale yellow gum. λ_{max} 205 (log ε 4.07); 229 (log ε 3.52); 281 (log ε 3.11) (MeOH).

3'-Deoxy: **3'-Deoxytubastrine**

[160525-08-6 (monohydrochloride)]

C₉H₁₁N₃O 177.205

Alkaloid from the Australian marine sponge *Spongosorites* sp. Shows modest antibiotic activity. Oil (as monohydrochloride trihydrate). λ_{max} 219 (ε 13000); 282 (ε 19000); 289; 308 (EtOH) (Berdy).

4'-Deoxy, 1,2-dihydro: **4-Deoxy-7,8-dihydrotubastrine**

C₉H₁₃N₃O 179.221

Isol. from the sponge *Petrosia cf. contignata*. Brown viscous oil. λ_{max} 214 (ε 4700); 274 (ε 1400) (MeOH).

Sakai, R. *et al.*, *Chem. Lett.*, 1987, 127-128 (isol)


Urban, S. *et al.*, *Aust. J. Chem.*, 1994, **47**, 2279 (3'-Deoxytubastrine)

Sperry, S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 859-861 (*Dihydrotubastrines*)

Pearce, A.N. *et al.*, *Tetrahedron*, 2008, **64**, 5748-5755 (isol, pmr, cmr, ms)

Tuberactidine T-657

α,2-Diamino-1,4,5,6-tetrahydro-6-hydroxy-4-pyrimidineacetic acid, 9CI. α-(Hexahydro-4-hydroxy-2-iminopyrimidin-6-yl)-glycine [28945-12-2]



Absolute configuration


C₆H₁₂N₄O₃ 188.186

Amino acid component of Tuberactinomycins. Mp 182° dec. (as hydrobromide). $[\alpha]_D^{15} -25.8$ (c, 0.5 in H₂O) (hydrobromide).

Shiba, T. *et al.*, *Tet. Lett.*, 1970, 3497 (struct) Kitagawa, T. *et al.*, *J. Biochem. (Tokyo)*, 1977, **81**, 1759

Tubercidin T-658

7-Ribofuranosyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine, 9CI. 4-Amino-7-ribofuranosyl-7H-pyrrolo[2,3-d]pyrimidine, 8CI. 7-Deazaadenosine. Sparsomycin A. B 79D. NSC 56408. U 10071. XK 101-1. Antibiotic B 79D. Antibiotic XK 101-1 [69-33-0]



C₁₁H₁₄N₄O₄ 266.256

Nucleoside antibiotic. Isol. from *Streptomyces tubericidus*. Major cytotoxin of *Tolypothrix hyssoides*. Also prod. by *Micromonospora chalcea tubericidica*. Constit. of the sponge *Caulospongia biflabellata*. Antitumour, antifungal and antiviral agent. Nucleoside transporter substrate. Cryst. Mp 247° dec. $[\alpha]_D^{17} -67$ (50% AcOH). pK_a 5.3. λ_{max} 227 (ε 25000); 272 (ε 12200) (dil HCl) (Derep). λ_{max} 270 (ε 12100) (dil. NaOH) (Derep). λ_{max} 270 (ε 12100) (H₂O) (Derep).

▶ LD₅₀ (rat, orl) 16 mg/kg. UY8870000

5'-O-Sulfamoyl: **Antibiotic SF 2494**. SF 2494

[114746-65-5]

C₁₁H₁₅N₅O₆S 345.335

From *Streptomyces mirabilis*. Herbicide.

5'-Deoxy: **5'-Deoxytubercidin**

[41107-17-9]

C₁₁H₁₄N₄O₃ 250.257

Isol. from *Didemnum voeltzkowi*.

Cryst. (EtOH/C₆H₆). Mp 203-207°.

$[\alpha]_D^{23} -70.3$ (c, 0.48 in DMSO). λ_{max} 270 (ε 12700) (MeOH).

5'-α-D-Glucopyranosyl: **Tubercidin 5'-α-D-glucopyranose**

[117456-78-7]

C₁₇H₂₄N₄O₉ 428.398

Found in *Plectonema radius* and *Tolypothrix distorta*. Cytotoxic. Antifungal. Sol. H₂O. $[\alpha]_D^{25} +10$ (c, 0.01 in H₂O). λ_{max} 268 (ε 9300) (H₂O) (Derep).

5-Bromo, 5'-deoxy: **5-Bromo-5'-deoxytubercidin, 9CI**

[95931-82-1]

C₁₁H₁₃BrN₄O₃ 329.153

Isol. from *Didemnum voeltzkowi*.

5-Iodo, 5'-deoxy: 5'-Deoxy-5-iodotubercidin

[85209-84-3]

C₁₁H₁₃IN₄O₃ 376.153

Constit. of *Hypnea valentiae* and *Didemnum voeltzowi*. Smooth muscle relaxant, hypothermic agent, inhibitor of adenosine kinase. Cryst. (Py). Mp 227-228° dec. [α]_D²⁵ -55 (c, 0.2 in MeOH).

1'-Epimer, 5-iodo, 5'-deoxy: [85209-85-4]C₁₁H₁₃IN₄O₃ 376.153

Constit. of *Hypnea valentiae* and *Didemnum voeltzowi*. Muscle relaxant. Pale yellow gum.

2'-Epimer: Aratubercidin. Ara Tb

[64526-34-7]

C₁₁H₁₄N₄O₄ 266.256


Shows antiviral props. Mp 125-126°. [α]_D²⁴ +6.9 (c, 0.5 in DMF).

► UY8820000

- Anzai, K. *et al.*, *J. Antibiot., Ser. A*, 1957, **10**, 201-204 (*isol*)
- Mizuno, Y. *et al.*, *J.O.C.*, 1963, **28**, 3329-3331 (*struct, uv, config*)
- Smulson, M.E. *et al.*, *J. Biol. Chem.*, 1967, **242**, 2872-2876 (*biosynth*)
- Tolman, R.L. *et al.*, *J.A.C.S.*, 1969, **91**, 2102-2108 (*synth, uv, ir, nmr*)
- Abola, J. *et al.*, *Acta Cryst. B*, 1973, **29**, 697-703 (*cryst struct*)
- Chenon, M.-T. *et al.*, *J.A.C.S.*, 1975, **97**, 4627-4636 (*pmr, cmr*)
- Robbins, M.J. *et al.*, *Can. J. Chem.*, 1977, **55**, 1260 (*Aratubercidin*)
- Ektova, L.V. *et al.*, *Bioorg. Khim.*, 1978, **4**, 1250-1255; *Sov. J. Bioorg. Chem. (Engl. Transl.)*, 1978, **4**, 907-910 (*synth*)
- Fukushima, K. *et al.*, *J. Antibiot.*, 1978, **31**, 377-378 (*ms*)
- Japan. Pat., 1978, 78 124 685; *CA*, **90**, 136241v (*isol*)
- Uzawa, J. *et al.*, *Org. Magn. Reson.*, 1979, **12**, 612-615 (*cmr*)
- Seela, F. *et al.*, *Angew. Chem., Int. Ed.*, 1981, **20**, 97 (*Aratubercidin*)
- Kazlauskas, R. *et al.*, *Aust. J. Chem.*, 1983, **36**, 165-170 (*5-iodo-5'-deoxy, 1'-epimer 5-iodo-5'-deoxy*)
- Barchi, J.J. *et al.*, *Phytochemistry*, 1983, **22**, 2851-2852 (*isol*)
- Seela, F. *et al.*, *Annalen*, 1984, 1972-1980 (*synth, cmr*)
- Yoo, J.C. *et al.*, *J. Liq. Chromatogr.*, 1984, **7**, 151-158 (*hplc*)
- Bergstrom, D.E. *et al.*, *J. Med. Chem.*, 1984, **27**, 285-292 (*props*)
- Cook, A.F. *et al.*, *Nucleosides Nucleotides*, 1984, **3**, 401-411 (*5-Bromo-5'-deoxytubercidin, 5-iodo-5'-deoxy, synth*)
- Ramasamy, K. *et al.*, *Tet. Lett.*, 1987, **28**, 5107-5110 (*synth*)
- Iwata, M. *et al.*, *CA*, 1988, **109**, 3509 (*SF2494*)
- Stewart, J.B. *et al.*, *J. Antibiot.*, 1988, **41**, 1048-1056 (*5'-glucosyl*)
- Isono, K. *et al.*, *J. Antibiot.*, 1988, **42**, 1711-1739 (*rev*)
- Reddy, A.M. *et al.*, *J. Het. Chem.*, 1990, **27**, 1297-1305 (*ms*)
- Plagemann, P.G. *et al.*, *Biochem. Pharmacol.*, 1991, **42**, 247-252 (*pharmacol*)
- Mitchell, S.S. *et al.*, *J. Nat. Prod.*, 1996, **59**, 1000-1001 (*5'-deoxy, 5-bromo-5'-deoxy, 5-iodo-5'-doxy*)
- Biabani, M.F. *et al.*, *Pharm. Biol.*, 2002, **40**, 302-303 (*isol*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., *J. Wiley*, 2000, TNY500

Tuberine

[97400-75-4]

C₂₇H₃₅NO₆ 469.577

Alkaloid from the aerial parts of *Haplophyllum tuberculatum* (Rutaceae). Shows strong antibacterial effects on *Staphylococcus aureus* and *E. coli*. Needles (EtOAc/petrol). Mp 150-152°. [α]_D²⁵ +12.5 (c, 0.12 in CHCl₃).

Sheriha, G.M. *et al.*, *Phytochemistry*, 1985, **24**, 884 (*uv, ir, pmr, cmr, ms, struct, stereochem*)

McPhail, A.T. *et al.*, *Phytochemistry*, 1990, **29**, 3055 (*abs config*)

Tubermycin AC₁₇H₁₆N₂O₂ 280.326

Struct. unknown; prob. an alkyl-substit. phenazine carboxylic acid. Isol. from *Streptomyces misakiensis*. Antibiotic; active against *Staphylococcus aureus* and *Mycobacterium*. Yellow cryst. (Me₂CO aq.). Sol. AcOH, dioxan, DMF, bases, Me₂CO; fairly sol. MeOH, EtOH, EtOAc, Et₂O, C₆H₆; poorly sol. H₂O. Mp 174°. λ_{\max} 254 (ε 85000); 370 (ε 16500) (MeOH/HCl) (Derep). λ_{\max} 256 (ε 98300); 365 (ε 16000) (MeOH) (Derep).

► LD₅₀ (mus, ivn) 160 mg/kg.


Isono, K. *et al.*, *J. Antibiot., Ser. A*, 1958, **11**, 264-267 (*isol, uv, ir, struct*)

Tuberosinone

T-661

11-Hydroxy-5H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinoline-5,6(7H)-dione, 9CI

[81451-87-8]

C₁₇H₉NO₅ 307.262

Alkaloid from the roots of *Aristolochia tuberosa* (Aristolochiaceae). Red needles. Mp 340°. λ_{\max} 238 (ε 37200); 277 (ε 13200); 286 (ε 14100); 318 (ε 11200); 330 (ε 14500); 372 (ε 8130); 390 (ε 8710); 476 (ε 16600) (EtOH) (Derep).

O-Ac:

Yellow needles. Mp 340°.

N-β-D-Glucopyranoside: Tuberosinone N-β-D-glucoside

[81451-88-9]

C₂₃H₁₉NO₁₀ 469.404

Alkaloid from the roots of *Aristolochia tuberosa* (Aristolochiaceae). Red needles. Mp 235-237°.

T-659

Me ether, N-Me: Tuberosinone BC₁₉H₁₃NO₅ 335.315


Alkaloid from *Aristolochia tuberosa*. Red needles. Mp 151-153°.

Zhu, D. *et al.*, *Heterocycles*, 1982, **17**, 345 (*uv, ir, pmr, ms, struct*)

Zhu, D.-Y. *et al.*, *Planta Med.*, 1990, **56**, 514 (*Tuberosinone B*)

Tuberosinone C

T-662

C₂₀H₁₇NO₅ 351.358

Alkaloid from *Aristolochia tuberosa*. Red needles. Mp 254-256°.

O²-De-Me: Aristoliukine C

[310396-77-1]

C₁₈H₁₃NO₅ 323.304

Alkaloid from *Aristolochia kaempferi*. Red needles (MeOH). Mp 220-221° dec. λ_{\max} 219; 237; 246; 274 (sh); 327; 472 (MeOH).

O¹,O¹⁰-Di-de-Me: Triangularine A

[197006-51-2]

C₁₇H₁₁NO₅ 309.278

Alkaloid from the roots of *Aristolochia triangularis*.

O²,O¹⁰-Di-de-Me: Aristoliukine B

[217310-32-2]

C₁₇H₁₁NO₅ 309.278

Alkaloid from *Aristolochia kaempferi*. Red needles. Mp 249-250° (dec.). λ_{\max} 219; 235; 246; 275 (sh); 321 (sh); 333; 373 (sh); 480 (MeOH).

Zhu, D.-Y. *et al.*, *Planta Med.*, 1990, **56**, 514 (*Tuberosinone C*)

Lin, W. *et al.*, *J. Chin. Pharm. Sci.*, 1997, **6**, 8-13 (*Triangularine A*)


Wu, T.S. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**, 1624-1626 (*Aristoliukine B*)

Wu, T.S. *et al.*, *Biol. Pharm. Bull.*, 2000, **23**, 1216-1219 (*Aristoliukine C*)

Tuberosmoenone

T-663

[220784-37-2]


C₂₂H₂₉NO₅ 387.475

Constit. of the roots of *Stemona tuberosa*.

Lin, W. *et al.*, *J. Chin. Pharm. Sci.*, 1999, **8**, 1-7

Tuberostemonine**T-664**

2-(Tetrahydro-4-methyl-5-oxo-2-furanyl)-stenine, 9CI
[6879-01-2]



Absolute
Configuration

$C_{22}H_{33}NO_4$ 375.507

Numbering shown as given by Lin *et al.* Alkaloid from *Stemona tuberosa*, *Stemona japonica* and *Stemona sessilifolia* (Stemonaceae). Anthelmintic agent. Needles (EtOH). Poorly sol. hexane. Mp 86-88°. $[\alpha]_D^{20}$ -25.4 (Me₂CO). pK_a 6.4 (60% EtOH).

Hydrobromide: Mp 120° dec.

Perchlorate: Mp 242° dec.

N-Oxide: **Tuberostemonine N-oxide**

[232921-40-3]

$C_{22}H_{33}NO_5$ 391.506

Alkaloid from the roots of *Stemona tuberosa*.

1,2,3,9a-Tetrahydro: **Bisdehydrotuberostemonine**. Didehydrotuberostemonine [106861-40-9]

$C_{22}H_{29}NO_4$ 371.475

Alkaloid from roots of *Stemona tuberosa* (Stemonaceae). Needles (Et₂O). Mp 176-178° (172-174°). $[\alpha]_D^{20}$ +105.96 (c, 0.1 in C₆H₆).

9-Hydroxy, 1,9a-didehydro: **Tuberostemonol**

[142905-24-6]

$C_{22}H_{31}NO_5$ 389.491

Alkaloid from roots of *Stemona tuberosa* (Stemonaceae). Amorph. $[\alpha]_D^{20}$ +33.54 (c, 0.3 in MeOH).

3-Epimer: **Tuberostemonine A**

[876313-35-8]

$C_{22}H_{33}NO_4$ 375.507

Minor alkaloid from the roots of *Stemona sessilifolia* (Stemonaceae). Needles (MeOH). Mp 118-120°. $[\alpha]_D^{20}$ -65 (c, 1.0 in EtOH). pK_a 5.45 (80% MeOH).

3-Epimer, hydrochloride:

Cryst. + 1H₂O. Mp 140-141°.

10-Epimer, 1,2,3,9a-tetrahydro: **Bisdehydrotuberostemonine B**

[263863-25-8]

$C_{22}H_{29}NO_4$ 371.475

Alkaloid from the roots of *Stemona japonica*.

13-Epimer: **Tuberostemonine N**

[942579-85-3]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from *Stemona tuberosa*. Amorph. powder. $[\alpha]_D^{20}$ -24 (c, 0.5 in CHCl₃).

13-Epimer, 4β,9a-epoxy: **Sessilifoline A**

[929637-35-4]

$C_{22}H_{31}NO_5$ 389.491

Alkaloid from the stems of *Stemona sessilifolia*. Amorph. powder. $[\alpha]_D^{20}$ +31.4 (c, 0.8 in CHCl₃).

1,10-Diepimer: **Tuberostemonine B**

[263870-88-8]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from the roots of *Stemona japonica*.

9,10-Diepimer, 1,2,3,9a-tetrahydro:

Bisdehydrotuberostemonine C

[263863-26-9]

$C_{22}H_{29}NO_4$ 371.475

Alkaloid from the roots of *Stemona japonica*.

11,15-Diepimer: **Tuberostemonine K**

[885056-80-4]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from the roots of *Stemona tuberosa*. Powder. $[\alpha]_D^{20}$ -12.6 (c, 0.1 in MeOH).

11,15-Diepimer, 1,2,3,9a-tetrahydro:

Bisdehydroneotuberostemonine

[160333-27-7]

$C_{22}H_{29}NO_4$ 371.475

From roots of *Stemona tuberosa* (Stemonaceae). Needles (EtOH). Mp 172-174°. $[\alpha]_D^{18.5}$ -32 (c, 1.0 in EtOH).

11,15-Diepimer, 1-hydroxy, 9,9a-didehydro:

Neotuberostemonol

[473925-49-4]

$C_{22}H_{31}NO_5$ 389.491

Alkaloid from *Stemona tuberosa*. Prisms (hexane/EtOAc). Mp 195-197°.

1,9,10-Triepimer: **Tuberostemonine C**

[263870-89-9]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from the roots of *Stemona japonica*.

1,11,15-Triepimer: **Neotuberostemonine**.

Tuberostemonine LG. **Stemonine LG**

[143120-46-1]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from roots of *Stemona tuberosa*. Antitussive agent. Cryst. (EtOH). Mp 160.5-162°. $[\alpha]_D^{20}$ +83 (c, 0.1 in MeOH).

5',9,13-Triepimer, 1,2,3,9a-tetrahydro:

Isobisdehydrotuberostemonine

[232921-41-4]

$C_{22}H_{29}NO_4$ 371.475

Alkaloid from the roots of *Stemona tuberosa*.

1,9a,11,15-Tetraepimer: **Tuberostemonine H**

[670254-76-9]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from the roots of *Stemona tuberosa*. Needles (hexane/EtOAc). Mp 183-185°. $[\alpha]_D^{20}$ +77.6 (c, 0.1 in MeOH).

9,11,15,18-Tetraepimer, 1,2,3,9a-tetrahydro:

Epibisdehydroneotuberostemonine J

[670254-77-0]

$C_{22}H_{29}NO_4$ 371.475

Alkaloid from the roots of *Stemona tuberosa*. Prisms (hexane/EtOAc). Mp 186-188°. $[\alpha]_D^{20}$ -16.1 (c, 0.1 in MeOH).

1,9,9a,11,15-Pentaepimer: **Tuberostemonine J**

[670254-75-8]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from the roots of *Stemona tuberosa*. Prisms (hexane/EtOAc). Mp 180-182°. $[\alpha]_D^{20}$ +36.4 (c, 0.1 in MeOH).

1,11,15,18,20-Pentaepimer: **Tuberostemo-**

nine L

[947730-01-0]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from a *Stemona* sp.

1,3,9a,11,15,18-Hexaepimer: **Tuberostemonine M**

[947730-02-1]

$C_{22}H_{33}NO_4$ 375.507

Alkaloid from a *Stemona* sp.

Edwards, O.E. *et al.*, *Can. J. Chem.*, 1962, **40**, 455; 2416 (*isol, ir, pmr, struct, epimer*)

Harada, H. *et al.*, *Chem. Comm.*, 1967, 460 (*cryst struct, abs config*)

Götz, M. *et al.*, *Tetrahedron*, 1968, **24**, 2631 (*ir, pmr, struct*)

Lin, W.-H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 571 (*Tuberostemonol*,

Didehydrotuberostemonine)

Dao, C.N. *et al.*, *Acta Cryst. C*, 1994, **50**, 1612-1615 (*Neotuberostemonine, cryst struct*)

Ye, Y. *et al.*, *Phytochemistry*, 1994, **37**, 1201 (*Neotuberostemonine*,

Bisdehydroneotuberostemonine)

Lin, W. *et al.*, *J. Chin. Pharm. Sci.*, 1999, **8**, 1-7 (*N-oxide, Isodidehydrotuberostemonine*)

Zou, C. *et al.*, *J. Chin. Pharm. Sci.*, 1999, **8**,

185-190 (*Tuberostemonines B,C*)

Bisdehydrotuberostemonines B,C)

Jiang, R.-W. *et al.*, *Tetrahedron*, 2002, **58**,

6705-6712 (*Neotuberostemonol*)

Chung, H.-S. *et al.*, *Planta Med.*, 2003, **69**,

914-920 (*Tuberostemonines H,J*,

Epibisdehydroneotuberostemonine J)

Wipf, P. *et al.*, *J.A.C.S.*, 2005, **127**, 225-235

(*synth*)

Jiang, R.-W. *et al.*, *J. Nat. Prod.*, 2006, **69**, 749-754 (*Tuberostemonine K, cryst struct*)

Sastrarajii, T. *et al.*, *Nat. Prod. Commun.*, 2006, **1**, 813-818 (*Tuberostemonines L,M*)


Greger, H. *et al.*, *Planta Med.*, 2006, **72**, 99-113 (*rev. occur, activity*)

Qian, J. *et al.*, *Helv. Chim. Acta*, 2007, **90**, 326-331 (*Sessilifoline A*)

Schinnerl, J. *et al.*, *Phytochemistry*, 2007, **68**, 1417-1427 (*Tuberostemonine N*)

Tuberostemoninol**T-665**

[156280-88-5]



$C_{22}H_{31}NO_6$ 405.49

Alkaloid from the roots of *Stemona tuberosa*. Mp 217-219°. $[\alpha]_D^{20}$ +124 (c, 0.83 in MeOH).

1-Deoxy, 8α-hydroxy: **Maireistemoninol**

[954379-67-0]

$C_{22}H_{31}NO_6$ 405.49

Alkaloid from the roots of *Stemona mairei*. Needles. Mp 155-157°. $[\alpha]_D^{24}$ +98.5 (c, 0.24 in CHCl₃).

11,12,13-Triepimer: **Neotuberostemoninol**

[473928-70-0]

$C_{22}H_{31}NO_6$ 405.49

Alkaloid from *Stemona tuberosa*. Prisms (hexane/EtOAc). Mp 190-192°. C-13 stereochem. incorrectly drawn in ref.


1,9,10,13-Tetraepimer: **Sessilifoliamide I**

[953391-39-4]
 $C_{22}H_{31}NO_6$ 405.49
 Alkaloid from the roots of *Stemona sessilifolia*. Prisms (MeOH aq.). Mp 273-275°. $[\alpha]_D^{20}$ -0.2 (c, 0.78 in MeOH).

- Lin, W.H. *et al.*, *Phytochemistry*, 1994, **36**, 1333-1335 (*isol, ir, pmr, cmr, ms*)
 Jiang, R.-W. *et al.*, *Tetrahedron*, 2002, **58**, 6705-6712 (*Neotuberostemoninol*)
 Hitotsuyanagi, Y. *et al.*, *Heterocycles*, 2007, **71**, 2035-2040 (*Sessilifoliamide I*)
 Cai, X.-H. *et al.*, *Planta Med.*, 2007, **73**, 170-173 (*Maireistemoninol*)

Tuberostemonone**T-666**

[134822-46-1]



$C_{22}H_{31}NO_6$ 405.49
 Config. at C-12 incorrectly shown in earlier papers. Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Cubes (Me₂CO). Mp 208-209°. $[\alpha]_D^{20}$ +134.8 (c, 0.1 in CHCl₃).

11-Epimer, 9,10-didehydro, 9ξ,10ξ-epoxide: Epoxytuberostemonone
 [954379-69-2]

$C_{22}H_{29}NO_7$ 419.474
 Alkaloid from the roots of *Stemona mairei*. Cryst. Mp 286-288°. $[\alpha]_D^{24}$ -67.5 (c, 0.16 in CHCl₃).

12-Epimer, 9-hydroxy: Sessilifoliamide H
 [922500-45-6]

$C_{22}H_{31}NO_7$ 421.489
 Alkaloid from the roots of *Stemona sessilifolia*. Prisms. Mp 240-243°. $[\alpha]_D^{20}$ -42 (c, 0.1 in CHCl₃).


13-Epimer: Neotuberostemonone

[954379-68-1]
 $C_{22}H_{31}NO_6$ 405.49
 Alkaloid from the roots of *Stemona mairei*. Cryst. Mp 228-230°. $[\alpha]_D^{24}$ +129.6 (c, 0.38 in CHCl₃).

- Lin, W.-H. *et al.*, *J. Crystallogr. Spectrosc. Res.*, 1991, **21**, 189-194 (*cryst struct*)
 Lin, W.-H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 571-576 (*isol, ir, pmr, cmr, ms*)
 Cai, X.-H. *et al.*, *Planta Med.*, 2007, **73**, 170-173 (*Neotuberostemonone, Epoxytuberostemonone*)
 Hitotsuyanagi, Y. *et al.*, *Tetrahedron*, 2007, **63**, 1008-1013 (*Sessilifoliamide H*)

Tuberostemospironone**T-667**

[142905-26-8]



Relative configuration


 $C_{13}H_{19}NO_4$ 253.297

Alkaloid from the roots of *Stemona tuberosa* (Stemonaceae). Cubes (Me₂CO). Mp 245-246°. $[\alpha]_D^{16}$ -30 (c, 0.02 in MeOH).

- Lin, W.-H. *et al.*, *J. Nat. Prod.*, 1992, **55**, 571-576 (*isol, ir, pmr, cmr, ms, struct*)

Tubifolidine**T-668**

17-Norcuran, 9CI
 [6883-33-6]



Absolute Configuration

$C_{18}H_{24}N_2$ 268.401
 Alkaloid from *Pleiocarpa pycnantha* var. *tubicina* (Apocynaceae) and from *Strychnos angolensis*. Prisms (Et₂O/pentane, Me₂CO aq. or by subl.). Mp 176-177°. $[\alpha]_D^{23}$ -67 (c, 0.61 in CHCl₃).

N-Formyl: Mp 148-149°.
1,2-Didehydro: Tubifoline, 1,2-Didehydro-17-norcuran, 9CI, Decarbomethoxy-19,20-dihydroakuammicine
 [2912-06-3]
 $C_{18}H_{22}N_2$ 266.385
 Alkaloid from *Pleiocarpa pycnantha* var. *tubicina* (Apocynaceae) and from *Strychnos angolensis*. Cryst. (pentane or by subl.). Mp 124-126°. $[\alpha]_D^{23}$ -342 (c, 0.55 in CHCl₃).

1,2-Didehydro; methiodide: Mp 251-253° dec.

12-Methoxy, 19,20-didehydro: 19,20-Didehydro-12-methoxytubifolidine, 19,20-Didehydro-12-methoxy-17-norcuran, 9CI, 16-Decarbomethoxyvinervinine (incorr.)
 [179461-86-0]

$C_{19}H_{24}N_2O$ 296.411
 Alkaloid from *Haplophyton crooksii* (Apocynaceae). Glass. Erroneously named as 16-Decarbomethoxyvinervinine in the paper; it is actually 16-Decarbomethoxy-2,16-dihydrovinervinine. 19,20-Didehydro-12-methoxytubifolidine is a better name.


- [20823-98-7]
 Weissmann, C. *et al.*, *Helv. Chim. Acta*, 1961, **44**, 1877 (*synth, deriv*)
 Kump, W.G. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 1497 (*isol, uv, ir, struct*)
 Dadson, B.A. *et al.*, *Chem. Comm.*, 1968, 1233 (*synth*)

- Wu, A. *et al.*, *Tet. Lett.*, 1975, 2057 (*synth*)
 Takano, S. *et al.*, *Tet. Lett.*, 1982, **23**, 881 (*synth*)
 Ban, Y. *et al.*, *Tetrahedron*, 1983, **39**, 3657 (*synth*)
 Amat, M. *et al.*, *J.O.C.*, 1990, **55**, 6299 (*synth, pmr, cmr*)
 Alvarez, M. *et al.*, *Tet. Lett.*, 1990, **31**, 5089 (*synth*)
 Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1995, **64**, 243-246 (*isol*)
 Mroue, M.A. *et al.*, *J. Nat. Prod.*, 1996, **59**, 890-893 (*19,20-Didehydro-12-methoxytubifolidine*)
 Soli, D. *et al.*, *Tet. Lett.*, 1996, **37**, 5213 (*synth*)

- Bonjoch, J. *et al.*, *J.A.C.S.*, 1997, **119**, 7230-7240 (*synth*)
 Amat, M. *et al.*, *Tetrahedron: Asymmetry*, 1997, **8**, 935-948 (*synth*)
 Shimizu, S. *et al.*, *J.O.C.*, 1998, **63**, 7547-7551 (*synth*)
 Mori, M. *et al.*, *J.A.C.S.*, 2003, **125**, 9801-9807 (*Tubifoline, synth*)
 Ishikura, M. *et al.*, *Heterocycles*, 2008, **75**, 107-118 (*Tubifoline, synth*)

Tubingensin A**T-669**

[122951-44-4]



$C_{28}H_{35}NO$ 401.591
 Biogenetically related to Aflavinine, A-172. Alkaloid from the sclerotia of *Aspergillus tubingensis* NRRL4700. Exhibits activity against the widespread crop pest *Heliothis zea* and displays *in vitro* antiviral activity against herpes simplex virus type 1. Cytotoxic. Light yellow solid. Mp 95-98°. $[\alpha]_D^{20}$ +13.6 (c, 1.0 in CHCl₃). λ_{max} 218 (ε 14900); 239 (ε 18200); 262 (ε 6930); 302 (ε 6780); 326 (ε 480); 340 (ε 480) (MeOH) (Derep).


10,27-Dihydro: 10,27-Dihydrotubingensin A
 [354154-75-9]

$C_{28}H_{37}NO$ 403.606
 Alkaloid from *Aspergillus tubingensis*. Solid.

- TePaske, M.R. *et al.*, *J.O.C.*, 1989, **54**, 4743-4746 (*isol, uv, pmr, cmr, ms, struct*)
 Sings, H.L. *et al.*, *J. Nat. Prod.*, 2001, **64**, 836-838 (*10,27-Dihydrotubingensin A*)

Tubingensin B**T-670**

[126663-32-9]



$C_{28}H_{35}NO$ 401.591
 Isol. from the sclerotia of *Aspergillus tubingensis* NRRL 4700. Cytotoxic. Light yellow cryst. Sol. MeOH, hexane; poorly sol. H₂O. Mp 152-154°. $[\alpha]_D^{20}$ -6.7 (c, 0.8 in CHCl₃). λ_{max} 218 (ε 17200); 237 (ε 25500); 260 (ε 10100); 299 (ε 10100); 325 (ε 2200); 338 (ε 6700) (MeOH) (Berdy).

10,27-Dihydro: 10,27-Dihydrotubingensin B

[354154-76-0]


C₂₈H₃₇NO 403.606

Alkaloid from *Aspergillus tubingensis*. Solid. $[\alpha]_D^{20}$ -50 (c, 0.1 in MeOH). λ_{\max} 207 (sh) (log ϵ 4.02); 227 (log ϵ 4.26); 281 (log ϵ 3.57); 290 (log ϵ 3.52) (MeOH).

TePaske, M.R. et al., *Tet. Lett.*, 1989, **30**, 5965-5968 (isol, pmr, cmr, struct)
Sings, H.L. et al., *J. Nat. Prod.*, 2001, **64**, 836-838 (10,27-Dihydrotubingensin B)

Tuboflavine**T-671**

5-Ethyl-4H-indolo[3,2-de][1,5]naphthyridin-4-one. 5-Ethylcanthin-4-one

C₁₆H₁₂N₂O 248.284

Alkaloid from *Pleiocarpa tubicina* (Apocynaceae). Yellow needles. Mp 207-208°.


Methiodide: Mp 210-212° dec.

Kump, C. et al., *Helv. Chim. Acta*, 1963, **46**, 498 (isol, uv, ir, pmr, struct)

Achenbach, H. et al., *J.A.C.S.*, 1965, **87**, 4177 (ms)

Tubotaiwinol**T-672**

[88721-01-1]



Absolute Configuration


C₁₉H₂₂N₂O 294.396

Alkaloid from the root bark and stem bark of *Strychnos ngouniensis* (Loganiaceae). $[\alpha]_D$ +894 (c, 1.17 in CHCl₃). $[\alpha]_D$ +613 (c, 1.2 in MeOH).

Massiot, G. et al., *Tetrahedron*, 1983, **39**, 3645 (isol, uv, ir, pmr, cmr, ms, struct)

Tubotaiwine**T-673**

14,19-Dihydrocondylocarpine
[67111-69-9]



Absolute Configuration

C₂₀H₂₄N₂O₂ 324.422

Alkaloid from *Tabernaemontana* sp., *Amsonia brevifolia*, *Pleiocarpa tubicina*, *Pleiocarpa pycnantha*, *Aspidosperma lima*, *Ervatamia heyneana* and many other spp. in the Apocynaceae. Also from several *Strychnos* spp. Weak convulsant. Shows mod. cytotoxic activity vs. mouse P388 carcinoma cells. Cryst. (Et₂O) (difficult to cryst.). Mp 101-103°. Bp_{0.001} 120-

125°. $[\alpha]_D^{22}$ +628 (c, 0.8 in CHCl₃). λ_{\max} 204 (log ϵ 4.12); 232 (log ϵ 4.01); 298 (log ϵ 3.96); 328 (log ϵ 4.12) (EtOH) (Derep).

Picrate:

Cryst. (Me₂CO aq.). Mp 171-172° dec.

N⁴-Oxide: Tubotaiwine N⁴-oxide

[40169-69-5]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from *Melodinus aeneus*, *Strychnos mitis*, the roots of *Tabernaemontana holstii*, *Tabernaemontana pachysyphon*, and the root bark of *Conopharyngia johnstonii* (Apocynaceae). Shows stronger cytotoxic activity than Tubotaiwine. Cryst. (Me₂CO). Mp 170° (sinters at 130°) Mp 224° dec. (double Mp). $[\alpha]_D^{22}$ +588 (c, 1.2 in CHCl₃). λ_{\max} 230 (ϵ 11500); 289 (ϵ 6310); 324 (ϵ 7410) (EtOH) (Derep). λ_{\max} 208 ; 224 ; 293 ; 327 (EtOH) (Berdy).

19,20E-Didehydro: Condylocarpine

[4939-81-5]

C₂₀H₂₂N₂O₂ 322.406

Alkaloid from *Diplorhynchus condylocarpon*, *Craspidospermum verticillatum*, *Vallesia antillana*, *Melodinus australis* and some other spp. in the Apocynaceae, also in *Strychnos angolensis* and *Strychnos dolichothyrsa*. Prisms (Et₂O). Mp 167-168° (159-162°). $[\alpha]_D^{20}$ +900 (CHCl₃). $[\alpha]_D$ +870 (EtOH). λ_{\max} 228 (log ϵ 4.04); 295 (log ϵ 4.01); 328 (log ϵ 4.17) (MeOH).

19,20E-Didehydro, N⁴-oxide: Condylocarpine N⁴-oxide

[74918-28-8]

C₂₀H₂₂N₂O₃ 338.405

Minor alkaloid from the stems of *Tabernaemontana olivacea* (Apocynaceae). Needles (MeOH). Mp 203-204° dec. $[\alpha]_D^{20}$ +694 (CHCl₃). λ_{\max} 225 (log ϵ 4.16); 294 (log ϵ 4); 328 (log ϵ 4.03) (MeOH).

11-Hydroxy: 11-Hydroxytubotaiwine

[160598-53-8]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from cultured cells of *Aspidosperma quebracho-blanco* (quebracho) (Apocynaceae). Amorph. powder. $[\alpha]_D$ +207 (c, 0.04 in EtOH).

14R-Hydroxy, 19,20E-didehydro: 14-Hydroxycondylocarpine

[180869-67-4]

C₂₀H₂₂N₂O₃ 338.405

Minor alkaloid from the stem bark of *Kopsia deverrei*. Amorph. $[\alpha]_D^{20}$ +430 (c, 1.2 in CHCl₃). λ_{\max} 228 (log ϵ 4.01); 294 (log ϵ 3.95); 332 (log ϵ 4.21) (EtOH).

19S-Hydroxy: Lagunamine. 19-Hydroxytubotaiwine

[126721-42-4]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Prisms (MeOH). Mp 230-235°. $[\alpha]_D^{25}$ +578.7 (c, 0.9 in MeOH).

11-Methoxy: 11-Methoxytubotaiwine.

14,19-Dihydro-11-methoxycondylocarpine
[2122-32-9]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the bark of *Aspidosperma populifolium* and from *Aspidosperma excelsum* (Apocynaceae). Shows weak antibacterial activity against *Bacillus subtilis* and *Staphylococcus aureus*. Amorph. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. $[\alpha]_D^{29}$ +213 (c, 0.49 in CHCl₃). Blue fluor. in soln. λ_{\max} 255 (log ϵ 4.17); 286 (log ϵ 4.04); 327 (log ϵ 4.05) (EtOH).

12-Methoxy: 12-Methoxytubotaiwine

[123064-75-5]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from leaves of *Alstonia congesta* (Apocynaceae). $[\alpha]_D$ +305 (c, 0.15 in CHCl₃).

19R-Methoxy: 19R-Methoxytubotaiwine

[949932-18-7]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D$ +442 (c, 0.14 in CHCl₃). λ_{\max} 229 (log ϵ 3.88); 297 (log ϵ 3.75); 329 (log ϵ 3.87) (EtOH).

19S-Methoxy: 19S-Methoxytubotaiwine

[949932-17-6]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the stem bark of *Kopsia arborea*. Oil. $[\alpha]_D$ +429 (c, 0.24 in CHCl₃). λ_{\max} 229 (log ϵ 3.95); 297 (log ϵ 3.89); 327 (log ϵ 3.99) (EtOH).

11-Nitro: 11-NitrotubotaiwineC₂₀H₂₃N₃O₄ 369.419

Alkaloid from the stems of *Ervatamia flabelliformis*. Yellow powder (MeOH). $[\alpha]_D^{25}$ -13.9 (c, 0.2 in EtOH).

Stauffer, D. et al., *Helv. Chim. Acta*, 1961,**44**, 2006-2015 (*Condylocarpine, isol, uv, ir*)Sandoval, A. et al., *Tet. Lett.*, 1962, 409-414(*Condylocarpine, uv, ir, pmr, struct, synth*)Biemann, K. et al., *Tet. Lett.*, 1962, 527(*Condylocarpine, uv, ms, struct*)Schumann, D. et al., *Helv. Chim. Acta*, 1963,**46**, 1996 (*Condylocarpine, config*)Budzikiewicz, H. et al., *Tetrahedron*, 1963, **19**,

1265-1276 (ms)

Kump, W.E. et al., *Helv. Chim. Acta*, 1964, **47**,

1497-1503 (isol)


Linde, H.H.A. et al., *Helv. Chim. Acta*, 1965,**48**, 1822-1842 (*Condylocarpine*)Gilbert, B. et al., *Tetrahedron*, 1965, **21**, 1141-1166 (*14,19-Dihydro-14-methoxycondylocarpine*)Pinar, M. et al., *Helv. Chim. Acta*, 1972, **55**,2972-2974 (*Tubotaiwine N-oxide*)Harley-Mason, J. et al., *Pure Appl. Chem.*,1975, **41**, 167-174 (*synth*)Wang, A.H.J. et al., *Acta Cryst. B*, 1977, **33**,2977-2979 (*Condylocarpine, cryst struct*)Kingston, D.G.I. et al., *J. Nat. Prod.*, 1977, **40**,215-216 (*isol, oxide*)Baassou, S. et al., *Phytochemistry*, 1978, **17**,1449-1450 (*isol, oxide*)Laguna, A. et al., *Coll. Czech. Chem. Comm.*,1980, **45**, 1419-1423 (*isol, uv, ir, pmr, ms*)Achenbach, H. et al., *Z. Naturforsch.*, **B**, 1980,**35**, 885 (*Condylocarpine N-oxide*)Verpoorte, R. et al., *Planta Med.*, 1983, **48**,283-289 (*11-Methoxytubotaiwine*)Caron, C. et al., *Phytochemistry*, 1989, **28**,1241-1244 (*12-Methoxytubotaiwine*)Yamauchi, T. et al., *Phytochemistry*, 1990, **29**,3321-3325 (*Lagunamine*)Kuehne, M.E. et al., *J.O.C.*, 1991, **56**, 2696-2700 (*synth, uv, ir, pmr, cmr, ms*)Nkiliza, J. et al., *Tet. Lett.*, 1991, **32**, 1787-1790(*synth, 19-Hydroxytubotaiwine*)

- Aimi, N. *et al.*, *Heterocycles*, 1994, **38**, 2411-2414 (*11-Hydroxytubotaivine*)
 Grácia, J. *et al.*, *J.O.C.*, 1994, **59**, 3939-3951 (*synth*)
 Kuehne, M.E. *et al.*, *Pure Appl. Chem.*, 1994, **66**, 2094-2098 (*Lagunamine, synth*)
 Kan, C. *et al.*, *Nat. Prod. Lett.*, 1995, **7**, 275 (*14-Hydroxycondylocarpine*)
 Lim, K.-H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 1302-1307 (*19-Methoxytubotaivine*)
 Liang, S. *et al.*, *Helv. Chim. Acta*, 2008, **91**, 239-243 (*11-Nitrotubotaivine*)

Tuboxenine

T-674

2,20-Cycloaspidospermidine, 9CI
 [26242-59-1]



C₁₉H₂₄N₂ 280.412

Indole alkaloid most closely related to the Aspidofractine type. Minor alkaloid from leaves of *Pleiocarpa pycnantha* var. *tubinnica* (Apocynaceae). Cryst. (pentane). Mp 139-140°. [α]_D²⁰ +5.4 (c, 0.17 in CHCl₃).

Picrate:

Cryst. (MeOH aq. or MeOH/Et₂O).
 Mp 163-166°.

14,15-Didehydro, N¹-Me: N¹-Methyl-14,15-didehydrotuboxenine

[21196-98-5]
 C₂₀H₂₄N₂ 292.423

Alkaloid from the roots of *Vinca sardoa* (Apocynaceae). Mp 246-250° (hydrobromide).


[26242-58-0, 116839-59-9, 116839-58-8]

- Kump, C. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 358; 1497 (*isol, ms, pmr*)
 Ahoud, A. *et al.*, *J.A.C.S.*, 1974, **96**, 633 (*struct*)
 Hugel, G. *et al.*, *Tet. Lett.*, 1987, **28**, 1773 (*synth*)
 Cartier, D. *et al.*, *Heterocycles*, 1988, **27**, 657 (*synth*)
 Crippa, S. *et al.*, *Heterocycles*, 1990, **31**, 1663 (*N-Methyl-didehydrotuboxenine*)

Tubulosine

T-675

10,11-Dimethoxytubulosan-8'-ol, 9CI



C₂₉H₃₇N₃O₃ 475.63

Several numbering systems have been used. Log P 4.41 (calc).

(-)-form**Marckine**

[2632-29-3]

Alkaloid from the root bark and stem bark of *Alangium lamarckii* and the bark of *Pogonopus tubulosus* (Alangiaceae,

Rubiaceae). Shows amoebicidal and antineoplastic activity. Inhibitor of RNA, DNA and protein biosynth. Cryst. (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 259-261° Mp 261-263° Mp 282-283° (evacuated capillary). Pharmacol. active isomer. λ_{max} 228 (1350); 285 (1380) (MeOH). λ_{max} 281 (ε 14500) (MeOH) (Berdy). λ_{max} 228 (ε 1200); 285 (ε 1220) (EtOH) (Berdy). λ_{max} 283 (log ε 3.94); 292 (log ε 3.84) (EtOH). λ_{max} 283 (log ε 4) (0.1M NaOH).

O,N-Di-Ac: Mp 149-151°.

Me ether: Mp 183°. [α]_D²⁰ +15.4 (c, 1 in MeOH).

O⁹-De-Me: 9-Demethyltubulosine

[96948-27-5]

C₂₈H₃₅N₃O₃ 461.603

Alkaloid from the trunk bark of *Alangium vittense* (Alangiaceae). Cryst. (CHCl₃). Mp 200°. [α]_D²⁰ -40 (c, 1 in Py). λ_{max} 275 (ε 11700) (MeOH/HCl) (Derep). λ_{max} 282 (ε 10500); 305 (ε 8510); 322 (ε 4470) (MeOH/NaOH) (Derep). λ_{max} 279 (ε 12600) (MeOH) (Derep). λ_{max} 282 (ε 12800) (MeOH) (Berdy). λ_{max} 276 (ε 11500) (HCl) (Berdy). λ_{max} 282 (ε 10800) (NaOH) (Berdy).

▶ LD₅₀ (mus, ipr) 150 mg/kg.

O¹⁰-De-Me: 10-Demethyltubulosine

[86709-00-4]

C₂₈H₃₅N₃O₃ 461.603

Alkaloid from the bark and root bark of *Alangium lamarckii* (Alangiaceae). Needles + 2H₂O (CH₂Cl₂/MeOH). Mp 198-200°. [α]_D²³ -51.9 (c, 1 in Py).

1',2'-Didehydro: 1',2'-Didehydrotubulosine

[262369-76-6]

C₂₉H₃₅N₃O₃ 473.614

Alkaloid from *Alangium lamarckii*. Amorph. powder. [α]_D¹⁸ +2.1 (c, 0.42 in MeOH). λ_{max} 230 (sh) (log ε 4.2); 291 (log ε 3.69); 328 (log ε 4.04); 370 (sh) (log ε 3.67) (MeOH).

1',2',3',4'-Tetradehydro: 1',2',3',4'-Tetra-dehydrotubulosine

[2753-27-7]

C₂₉H₃₃N₃O₃ 471.598

Alkaloid from *Pogonopus speciosus*. Amorph. powder. [α]_D -20.4 (c, 0.2 in MeOH). λ_{max} 206 (log ε 3.13); 231 (log ε 2.88); 296 (log ε 2.6); 367 (log ε 2.2) (MeOH).

Deoxy: Deoxytubulosine

[2632-30-6]

C₂₉H₃₇N₃O₂ 459.63

Alkaloid from *Cassinopsis ilicifolia* and from the fruits and seeds of *Alangium lamarckii* (Icacinaeae, Alangiaceae). Weak amoebicide (*ca.* 1/20th the activity of Emetine). Mp 230-232°. [α]_D -24 (CHCl₃). [α]_D -17 (CHCl₃). Log P 5.08 (calc).

Deoxy, 8-hydroxy: Alangimarckine

[13849-53-1]

C₂₉H₃₇N₃O₃ 475.63

Alkaloid from the leaves of *Alangium lamarckii* (Alangiaceae). Cryst. + 0.5 H₂O (EtOAc/diisopropyl ether). Mp 185-187° dec. [α]_D¹⁸ -68.5 (c,

0.34 in Py). λ_{max} 283 (3.94); 292 (3.84) (EtOH). λ_{max} 283 (4) (0.1M NaOH).

1'-Epimer: Isotubulosine

[5610-39-9]

C₂₉H₃₇N₃O₃ 475.63

Alkaloid from the root bark of *Alangium lamarckii* (Alangiaceae). Cryst. (EtOH). Mp 177-178°. [α]_D²⁵ -84 (Py).

1'-Epimer, Me ether: Mp 163°. [α]_D²⁰ -18.8 (MeOH).

(±)-form [16049-31-3]

Synthetic. Cryst. (EtOH aq.). Mp 259-260° (with previous sintering at 255°).

Me ether: [74112-40-6]

Needles (MeOH). Mp 146-148°.

O⁹-De-Me: [74059-21-5]

Synthetic. Cryst. + 2H₂O. Mp 213-214° dec.

O¹⁰-De-Me: [74059-22-6]

Dihydrate. Mp 199-201° dec.

Deoxy: [70428-43-2]

Synthetic. Needles + 1H₂O (CHCl₃/MeOH). Mp 156-158°.

Deoxy, 8-hydroxy: [65847-00-9]

Synthetic. Cryst. + 0.75 H₂O (EtOH). Mp 157-159° dec.

Brauchli, P. *et al.*, *J.A.C.S.*, 1964, **86**, 1895-1896 (*isol, uv, ir, pmr, ms, struct*)

Battersby, A.R. *et al.*, *Chem. Comm.*, 1965, 315-317 (*Deoxytubulosine, struct, synth*)
 Monteiro, H. *et al.*, *Chem. Comm.*, 1965, 317-318 (*Deoxytubulosine*)

Albright, J.D. *et al.*, *J. Nat. Prod.*, 1965, **28**, 212-217 (*isol, uv, ir, pmr*)

Szántay, C. *et al.*, *Acta Chim. Acad. Sci. Hung.*, 1966, **49**, 427-428 (*struct, synth*)

Openshaw, H.T. *et al.*, *Chem. Comm.*, 1966, 131-132 (*synth*)

Salgar, S.S. *et al.*, *Curr. Sci.*, 1966, **35**, 281 (*isol, uv, ir*)

Popelak, A. *et al.*, *Tet. Lett.*, 1966, 1081-1085; 5077-5080 (*Isotubulosine, 10-Demethyltubulosine, struct, uv, pmr, ms*)

Battersby, A.R. *et al.*, *Tet. Lett.*, 1966, 4965-4971 (*Alangimarckine, isol, ir, pmr, ms*)

Szántay, C. *et al.*, *Chem. Ber.*, 1969, **102**, 2270-2272 (*synth, Tubulosine, Isotubulosine*)

Pakrashi, S.C. *et al.*, *Experientia*, 1970, **26**, 933-934 (*isol*)

Pareye, C. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. D*, 1973, **277**, 2689-2691 (*Deoxytubulosine, pharmacol*)

Kametani, T. *et al.*, *Can. J. Chem.*, 1979, **57**, 1679-1681 (*Deoxytubulosine, synth*)

Kametani, T. *et al.*, *Heterocycles*, 1979, **13**, 209-215 (*Me ether, synth, ir, pmr*)

Ohba, M. *et al.*, *Heterocycles*, 1980, **14**, 299-301 (*9-Demethyltubulosine, synth, pmr*)

Fujii, T. *et al.*, *Heterocycles*, 1980, **14**, 971-974 (*10-Demethyltubulosine, struct, synth*)

Achari, B. *et al.*, *Planta Med. (Suppl.)*, 1980, **5**
 Fujii, T. *et al.*, *Alkaloids (Academic Press)*, 1983, **22**, 31-32 (*bibl, pharmacol*)

Brown, R.T. *et al.*, *Tet. Lett.*, 1984, **25**, 3127-3130 (*Deoxytubulosine, synth*)

Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 1946-1954 (*Alangimarckine, struct, abs config, uv, ir, pmr, cd*)

Ohba, M. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 3724-3730 (*synth, uv, pmr, cmr, ms*)

Kan-Fan, C. *et al.*, *Heterocycles*, 1985, **23**, 1089-1092 (*9-Demethyltubulosine*)


Fujii, T. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 2355-2359 (*9-Demethyltubulosine, synth, cd, abs config*)

Ihara, M. *et al.*, *J.C.S. Perkin 1*, 1990, 1469-1476 (*synth*)

- Ito, A. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1346-1348 (Tetradehydrotubulysine)
 Itoh, A. *et al.*, *J. Nat. Prod.*, 2000, **63**, 723-725 (Didehydrotubulysine)
 Tietze, L.F. *et al.*, *Chem. Eur. J.*, 2004, **10**, 2722-2731 (*synth*)

Tubulysins

T-676



Tubulysin A R = -COCH₂CH(CH₃)₂
 B R = -COCH₂CH₂CH₃
 C R = -COCH₂CH₃
 G R = -COCH=C(CH₃)₂
 I R = -COCH₃
 Absolute Configuration

Prod. by two different species of myxobacteria. Inhibitors of tubulin polymerisation. Potent cytotoxins.

Tubulysin A [205304-86-5]

C₄₃H₆₅N₅O₁₀S 844.08
 Prod. by *Archangium gephyra* Ar 315 and *Angiococcus disciformis* An d48. Cryst. (MeOH aq.). Mp 106-108°. [α]_D²⁵ +15.3 (c, 5 in MeOH). λ_{\max} 205 (log ϵ 4.44); 225 (log ϵ 4.2); 250 (sh) (log ϵ 3.86); 276 (log ϵ 3.25); 287 (log ϵ 3.08) (MeOH).

4'-Deoxy: Tubulysin D

[309935-57-7]
 C₄₃H₆₅N₅O₉S 828.081
 Prod. by *Angiococcus disciformis* An d48.

Tubulysin B [205304-87-6]

C₄₂H₆₃N₅O₁₀S 830.053
 Prod. by *Archangium gephyra* Ar 315.

4'-Deoxy: Tubulysin E

[309935-58-8]
 C₄₂H₆₃N₅O₉S 814.054
 Prod. by *Angiococcus disciformis* An d48.

Tubulysin C [205304-88-7]

C₄₁H₆₁N₅O₁₀S 816.026
 Prod. by *Archangium gephyra* Ar 315.

4'-Deoxy: Tubulysin F

C₄₁H₆₁N₅O₉S 800.027
 Prod. by *Angiococcus disciformis* An d48.

Tubulysin G

C₄₃H₆₃N₅O₁₀S 842.064
 Prod. by *Archangium gephyra* Ar 315.

Tubulysin I

C₄₀H₅₉N₅O₁₀S 802
 Prod. by *Archangium gephyra* Ar 315.

4'-Deoxy: Tubulysin H

C₄₀H₅₉N₅O₉S 786
 Prod. by *Angiococcus disciformis* An d48.

Sasse, F. *et al.*, *J. Antibiot.*, 2000, **53**, 879-885 (*isol, activity*)

Steinmetz, H. *et al.*, *Angew. Chem., Int. Ed.*, 2004, **43**, 4888-4892 (*isol, pmr, cmr, cryst struct*)


Sandmann, A. *et al.*, *Chem. Biol.*, 2004, **11**, 1071-1079 (*prodn*)

- Khalil, M.W. *et al.*, *ChemBioChem*, 2006, **7**, 678-683 (*activity*)
 Peltier, H.M. *et al.*, *J.A.C.S.*, 2006, **128**, 16018-16019 (*Tubulysin D, synth*)

Tunichrome An1

T-677

[116050-17-0]



C₂₆H₂₅N₃O₁₁ 555.497
 Isol. from the tunicate *Ascidia nigra*. Blood pigment which selectively accumulates vanadium. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep).

1'Z-Isomer: Tunichrome B1

[97689-87-7]
 C₂₆H₂₅N₃O₁₁ 555.497
 Isol. from *Ascidia nigra*. Sol. MeOH. Dec. in air on warming. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep).

3''-Deoxy: Tunichrome An2

[115982-31-5]
 C₂₆H₂₅N₃O₁₀ 539.498
 Isol. from *Ascidia nigra*. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep).

1'Z-Isomer, 3''-deoxy: [133695-74-6]

C₂₆H₂₅N₃O₁₀ 539.498
 Isol. from *Ascidia nigra*. Unstable.

3'',3'''-Dideoxy: Tunichrome An3

[115982-32-6]
 C₂₆H₂₅N₃O₉ 523.498
 Isol. from *Ascidia nigra*. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep).

1'Z-Isomer, 3'',3'''-dideoxy: [133695-75-7]

C₂₆H₂₅N₃O₉ 523.498
 Isol. from *Ascidia nigra*. Unstable.

Bruening, R.C. *et al.*, *J.A.C.S.*, 1985, **107**, 5298-5300 (*Tunichrome B1*)

Bruening, R.C. *et al.*, *J. Nat. Prod.*, 1986, **49**, 193 (*Tunichrome B1*)

Oltz, E.M. *et al.*, *J.A.C.S.*, 1988, **110**, 6162-6172 (*isol*)

Horenstein, B.A. *et al.*, *J.A.C.S.*, 1989, **111**, 6242 (*synth*)


Kim, D. *et al.*, *Chem. Comm.*, 1991, 9 (*isol*)

He, X. *et al.*, *Experientia*, 1992, **48**, 367 (*biosynth*)

Tunichrome Pm 1

T-678

3,5-Dihydroxy-L-tyrosyl-3,5-dihydroxy-N-[2-(3,4,5-trihydroxyphenyl)ethenyl]-L-tyrosinamide, 9CI
 [137824-58-9]



C₂₆H₂₇N₃O₁₁ 557.513
 Isol. from the tunicate *Phallusia mammillata*. Readily chelates vanadium. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep). λ_{\max} 301 (MeOH) (Berdy).

3'-Deoxy: Tunichrome Pm 2

[137824-59-0]
 C₂₆H₂₇N₃O₁₀ 541.513
 Isol. from *Phallusia mammillata*. Readily chelates vanadium. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep). λ_{\max} 301 (MeOH) (Berdy).

3''-Deoxy: Tunichrome Pm 3


[137844-97-4]
 C₂₆H₂₇N₃O₁₀ 541.513
 Isol. from *Phallusia mammillata*. Readily chelates vanadium. λ_{\max} 210 (ϵ 68000); 245 (sh); 285 (sh); 340 (ϵ 19600) (MeOH) (Derep).

Bayer, E. *et al.*, *Angew. Chem., Int. Ed.*, 1992, **31**, 52-54 (*isol*)

Turbinatine

T-679

[575432-21-2]



C₂₇H₃₄N₂O₉ 530.574
 Alkaloid from the leaves of *Chimarrhis turbinata*. Amorph. yellow powder. [α]_D²⁷ -8.5 (c. 0.04 in MeOH). λ_{\max} 235 (ϵ 5875) (MeOH).


Cardoso, C.L. *et al.*, *J. Nat. Prod.*, 2003, **66**, 1017-1021 (*isol, pmr, cmr, conformn*)

Turbotoxin A

T-680

[245128-77-2]

[245128-78-3]



C₁₇H₃₀I₂N₂O₂⁺ 532.246
 Alkaloid from the Japanese gastropod *Turbo marmorata*. Acetylcholinesterase inhibitor. Isol. as the bis(trifluoroacetate). λ_{\max} 226 (ϵ 19000); 239 (sh) (ϵ 9500); 275 (sh) (ϵ 2000) (MeOH) (bistrifluoroacetate).


N⁹-De-Me: Turbotoxin B

[245128-79-4]
 [245128-80-7]
 C₁₆H₂₇I₂N₂O⁺ 517.211
 Alkaloid from *Turbo marmorata*. Isol. as the bis(trifluoroacetate). λ_{\max} 225 (ϵ

11000); 239 (sh) (ε 5500); 279 (sh) (ε 1500) (MeOH) (bistrifluoroacetate).


Kigoshi, H. *et al.*, *Tetrahedron*, 2000, **56**, 9063-9070 (*isol, synth, uv, pmr, cmr*)

Turcberine T-681
[169626-34-0]



$C_{23}H_{30}N_2O_4$ 398.501
Alkaloid from *Berberis turcomanica*. Oil.
Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **77**; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 63

Turconidine T-682
[66113-64-2]



$C_{23}H_{30}N_2O_4$ 398.501
Alkaloid from young shoots of *Berberis turcomanica*. Amorph. λ_{max} 225 (log ε 3.87); 284 (log ε 3.44) (EtOH).

Karimov, A. *et al.*, *Khim. Prir. Soedin.*, 1993, **29**, 866; *Chem. Nat. Compd. (Engl. Transl.)*, 1993, **29**, 771

Turkestanine T-683
[1361-67-7]


Struct. unknown. Pyrrolizidine alkaloid (?). Alkaloid from *Solenanthes turkestanicus* and *Rindera baldschuanica* (Boraginaceae). Mp 179-180° (?). Co-occurs with Rinderine, in E-12 and Echinatine, E-12. Opt. inactive. The name Turkestanine is not used in the first paper and it is not clear that all of the data given here refers to the same alkaloid.

N-Oxide: Turkestanine N-oxide

Alkaloid from above-ground parts of *Rindera austrochinata* (Boraginaceae). Mp 196-197°.

Akramov, S.T. *et al.*, *Dokl. Akad. Nauk SSSR*, 1962, **19**, 29-31; 1965, **22**, 35-38

Turkiyenine T-684



$C_{20}H_{15}NO_6$ 365.342

(+)-form [92219-92-6]
Alkaloid from *Hypecoum procumbens* (Papaveraceae). Amorph. $[\alpha]_D^{23}$ +72 (c, 0.053 in $CHCl_3$). λ_{max} 217 (ε 23400); 254 (sh) (ε 17800); 259 (ε 19500); 308 (ε 5890) (MeOH).


15-Oxo: Oxoturkiyenine
[115598-75-9]

$C_{20}H_{13}NO_7$ 379.325
Alkaloid from *Hypecoum pendulum* (Papaveraceae). Amorph. powder. $[\alpha]_D$ +38 (c, 0.11 in MeOH). $[\alpha]_D$ +75 (c, 0.08 in $CHCl_3$). λ_{max} 217 (ε 23400); 254 (sh) (ε 17800); 259 (ε 19500); 308 (ε 5890) (MeOH) (Derep).

(-)-form [128778-78-9]
Alkaloid from *Chelidonium majus* (Papaveraceae). Amorph. $[\alpha]_D$ -99 (c, 0.12 in $CHCl_3$). $[\alpha]_D$ -76 (c, 0.11 in MeOH).

Gözler, T. *et al.*, *J.A.C.S.*, 1984, **106**, 6101-6102 ((+)-form, *uv, ir, pmr, ms, cd, struct*)
Mete, I.E. *et al.*, *J. Nat. Prod.*, 1988, **51**, 272-274 ((+)-form, *Oxoturkiyenine, isol*)
Kadan, G. *et al.*, *J. Nat. Prod.*, 1990, **53**, 531-532 ((-)-form, *isol*)


Turraparvin D T-685
[566935-02-2]



$C_{28}H_{37}NO_5$ 467.604
Constit. of *Turraea parvifolia*. Cryst. Mp 131-133°. $[\alpha]_D^{23}$ +29.8 (c, 0.08 in $CHCl_3$).

Cheplogoi, P.K. *et al.*, *Phytochemistry*, 2003, **62**, 1173-1178 (*isol, pmr, cmr*)

Turrapubesin B T-686
[909410-15-7]



$C_{33}H_{41}NO_{10}$ 611.688
Constit. of *Turraea pubescens*. Amorph. solid. $[\alpha]_D^{20}$ +96.1 (c, 0.23 in MeOH). λ_{max} 231 (log ε 4.16) (MeOH).

Wang, X.-N. *et al.*, *Org. Lett.*, 2006, **8**, 3845-3848 (*Turrapubesin B*)

Turumiquirensine T-687
[1361-68-8]

$C_{42}H_{52}N_2O_9^{2+}$ 728.881

Struct. unknown. Quaternary alkaloid from *Croton turumiquirensis* bark (Euphorbiaceae).

Diiodide:


$C_{42}H_{52}I_2N_2O_9$ 982.69
Mp 235° dec.

Dinitrate:

$C_{42}H_{52}N_4O_{15}$ 852.891
Mp 240-241° dec.

Burnell, R.H. *et al.*, *Nature (London)*, 1964, **203**, 296-297 (*isol, uv*)

Tylocrebrine T-688
9,11,12,13,13a,14-Hexahydro-2,3,5,6-tetramethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinoline, 9CI, 2,3,5,6-Tetramethoxyphenanthroindolizidine



$C_{24}H_{27}NO_4$ 393.482
Possesses antineoplastic activity in mouse test systems. Log P 3.96 (calc).
▶ Toxic. Powerful vesicant.

(R)-form [6879-02-3]
Alkaloid from *Ficus septica* (Moraceae). Pale yellow cryst. (MeOH). Mp 220-222°. $[\alpha]_D^{22}$ +20.5 (c, 0.8 in $CHCl_3$).
▶ LD₅₀ (rat, orl) 65 mg/kg. LD₅₀ (rat, ivn) 32 mg/kg. YP0175000

Picrate:

Cryst. (MeOH aq.). Mp 144-146°.

Methodide:

Cryst. (MeOH). Mp 271-274°.

α-N-Oxide: (10R,13aR)-Tylocrebrine N-oxide

$C_{24}H_{27}NO_5$ 409.481
Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 205-212° dec. $[\alpha]_D^{25}$ -66.6 (c, 0.02 in MeOH). λ_{max} 244 (log ε 4.14); 265 (log ε 4.29); 276 (log ε 3.71); 310 (log ε 3.46); 320 (log ε 3.41) (MeOH).

β-N-Oxide: (10S,13aR)-Tylocrebrine N-oxide

$C_{24}H_{27}NO_5$ 409.481
Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 215-218° dec. $[\alpha]_D^{25}$ -61.4 (c, 0.08 in MeOH). λ_{max} 245 (log ε 3.14); 264 (log ε 4.39); 279 (log ε 3.91); 285 (log ε 3.82); 307 (log ε 3.41); 319 (log ε 3.38) (MeOH).

(S)-form [61302-92-9]
Alkaloid from *Tylophora crebriflora*,

Tylophora hirsuta and *Ficus septica*. Possesses antineoplastic activity in mouse test systems. Antiinflammatory agent. Cryst. (MeOH). Mp 218-220° dec. $[\alpha]_D^{24}$ -45 (c, 0.74 in CHCl₃). Log P 3.96 (calc). Pharmacol. active isomer. λ_{\max} 263 (ε 66200); 342 (ε 1780); 360 (ε 1230) (EtOH) (Berdy).

► CNS toxicity observed in humans.

Hydroiodide:

Yellow cryst. + 1MeOH (MeOH aq.). Mp 214-217° dec.

Picrate:

Cryst. (Me₂CO/MeOH). Mp 134-136°.

Methiodide:

Yellow cryst. + 1/2 H₂O (MeOH). Mp 255-258° dec. $[\alpha]_D^{21}$ -30 (c, 0.30 in MeOH).

(±)-**form** [30061-35-9]

Synthetic. Needles (CHCl₃/MeOH). Mp 219-221°.

Methiodide:

Cryst. (MeOH). Mp 264-266° dec.

Gellert, E. *et al.*, *J.C.S.*, 1962, 1008-1014 (*isol, uv, ir, struct, synth*)

Russel, J.H. *et al.*, *Naturwissenschaften*, 1963, **50**, 443-444 (*isol, struct*)

Gellert, E. *et al.*, *J. Med. Chem.*, 1964, **7**, 361-362 (*tox, pharmacol*)

Chauncy, B. *et al.*, *Aust. J. Chem.*, 1970, **23**, 2503-2516 (*synth, uv, ir, pmr*)


Gellert, E. *et al.*, *Aust. J. Chem.*, 1978, **31**, 2095-2097 (*ord, cd, abs config*)

Suffness, M. *et al.*, *Alkaloids (Academic Press)*, 1985, **25**, 156-163 (*pharmacol*)

Damu, A.G. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1071-1075 (*N-oxides*)

Tylohirsuticine

T-689



C₂₄H₂₉NO₄ 395.497

(S)-**form** [111316-24-6]

Minor alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae).

Light-yellow beads (Me₂CO). Mp 215-217°. $[\alpha]_D^{18}$ +20.8 (c, 0.8 in MeOH).

Ac: Mp 203-204°.

Me ether: Mp 199-202°.

6-Demethoxy, O⁷-de-Me, 7-Ac: **Tyloindicine B**

[126624-13-3]

C₂₄H₂₇NO₄ 393.482


Alkaloid from the aerial parts of *Tylophora indica*. Cryst. (CHCl₃). Mp 183-185°. $[\alpha]_D^{35}$ +14.3 (c, 1.05 in MeOH).

Ali, M. *et al.*, *Phytochemistry*, 1987, **26**, 2089-2092; 1989, **28**, 3513-3517 (*isol, uv, ir, pmr, ms, struct, Tyloindicine B*)

Tylohirsutinine

[95066-38-9]

T-690



C₂₄H₂₅NO₄ 391.466

Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Cryst. (Me₂CO). Mp 200-202°.

14-Hydroxy, O⁴-de-Me: **Tylohirsutinidine** [95066-41-4]

C₂₃H₂₃NO₅ 393.438


Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Cryst. (Me₂CO). Mp 235-237°. Opt. inactive (racemic).

Bhutani, K.K. *et al.*, *Phytochemistry*, 1984, **23**, 1765-1769 (*isol, uv, ir, pmr, ms, struct*)

Tyloindicine I

[138935-52-1]

T-691



C₂₄H₂₇NO₅ 409.481

Alkaloid from *Tylophora indica* (Asclepiadaceae). Cryst. (Me₂CO/MeOH). Mp 215-217° dec. $[\alpha]_D^{30}$ -2.2 (c, 0.44 in AcOH).

4-Deoxy, O³,O⁷-di-de-Me, O⁷-Ac: **Tyloindicine J**

[138935-55-4]

C₂₄H₂₅NO₅ 407.465


Alkaloid from *Tylophora indica* (Asclepiadaceae). Light yellow cryst. Mp 180-181° dec. $[\alpha]_D^{30}$ -2.85 (c, 0.57 in AcOH).

Ali, M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1271-1278 (*isol, uv, ir, pmr, ms, struct, Tyloindicine J*)

Tyloindicine F

[138964-86-0]

T-692



C₂₃H₂₇NO₄ 381.471


Alkaloid from *Tylophora indica* (Asclepiadaceae). Cryst. (Me₂CO/MeOH). Mp 245-247° dec. $[\alpha]_D^{30}$ -0.5 (c, 0.05 in AcOH).

Ali, M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1271-1278 (*isol, uv, ir, pmr, ms, struct*)

Tyloindicine G

[138935-58-7]

T-693



Absolute Configuration

C₂₄H₂₇NO₅ 409.481


Alkaloid from *Tylophora indica* (Asclepiadaceae). Cryst. (Me₂CO/MeOH). Mp 237-238° dec. $[\alpha]_D^{30}$ -2.35 (c, 0.47 in AcOH).

Ali, M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1271-1278 (*isol, uv, ir, pmr, ms, struct*)

Tyloindicine H

[138964-88-2]

T-694



C₂₃H₂₅NO₄ 379.455


Alkaloid from *Tylophora indica* (Asclepiadaceae). Cryst. Mp 225-226° dec. $[\alpha]_D^{30}$ -5.5 (c, 0.55 in AcOH).

Ali, M. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1271-1278 (*isol, uv, ir, pmr, ms, struct*)

Tylophorine

T-695

9,11,12,13,13a,14-Hexahydro-2,3,6,7-tetramethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinoline, 9CI. 2,3,6,7-Tetramethoxyphenanthroindolizidine



(R)-form

C₂₄H₂₇NO₄ 393.482

Natural (-)-Tylophorine was formerly erroneously assigned (S)-config. Antiasthmatic, antiallergic agent. Shows weak antineoplastic activity. Log P 3.96 (calc).

(R)-form [111408-21-0]

Alkaloid from *Tylophora asthmatica*, *Tylophora indica*, *Tylophora arebriflora*, *Tylophora hirsuta*, *Tylophora tanaka*, *Tylophora flava* and other *Tylophora* spp.; also from *Cynanchum vincetoxicum*, *Pergularia pallida*, *Vincetoxicum officinale* and *Ficus septica* (Asclepiadaceae, Moraceae). Cryst. (CHCl₃/EtOH). Mp 286-287° dec. $[\alpha]_D^{25}$ -76.5 (c, 0.04 in CHCl₃). $[\alpha]_D^{27}$ -11.6 (c, 1.07 in CHCl₃) (synth.). Pharmacol. active isomer. Highly toxic to frogs but low toxicity to mice and guinea pigs. λ_{max} 222 ; 241 ; 250 ; 257 ; 283 ; 288 ; 302 ; 323 (MeOH) (Berdy). λ_{max} 255 ; 290 ; 340 ; 352 (EtOH) (Berdy).

► Powerful vesicant.

Hydrochloride:

Cryst. (EtOH aq.). Mp 276-278° dec.

Hydrobromide: Mp 252-255° dec.

N-Oxide: Tylophorine N-Oxide

[209684-47-9]

C₂₄H₂₇NO₅ 409.481

Alkaloid from *Tylophora tanakae*. Fine yellowish prisms (MeOH). Mp 225-235° dec. $[\alpha]_D^{29}$ -8.6 (c, 0.2 in CHCl₃/MeOH). λ_{max} 220 (log ϵ 4.39); 238 (log ϵ 4.49); 250 (sh) (log ϵ 4.65); 257 (log ϵ 4.85); 282 (sh) (log ϵ 4.44); 287 (log ϵ 4.48); 302 (log ϵ 4.17) (MeOH).

 α -N-Oxide:

C₂₄H₂₇NO₅ 409.481

Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 210-220° dec. $[\alpha]_D^{25}$ -77.7 (c, 0.02 in MeOH). λ_{max} 241 (log ϵ 4.09); 258 (log ϵ 4.13); 287 (log ϵ 3.84); 303 (log ϵ 3.56); 340 (log ϵ 2.52) (MeOH).

 β -N-Oxide:

C₂₄H₂₇NO₅ 409.481

Alkaloid from the stems of *Ficus septica*. Pale yellow needles. Mp 215-218° dec. $[\alpha]_D^{25}$ -61.4 (c, 0.08 in MeOH). λ_{max} 245 (log ϵ 3.14); 264 (log ϵ 4.39); 279 (log ϵ 3.91); 285 (log ϵ 3.82); 307 (log ϵ 3.41); 319 (log ϵ 3.38) (MeOH).

O⁷-De-Me: 7-Demethyltylophorine

[165606-55-3]

C₂₃H₂₅NO₄ 379.455

Alkaloid from leaves of *Tylophora tanakae*. Mp 250-260° dec. $[\alpha]_D^{27}$ -49.3 (c, 0.25 in CHCl₃). λ_{max} 220 ; 237 ; 257 ; 288 ; 302 (MeOH) (Berdy).

O⁷-De-Me, N-oxide: 7-Demethyltylophorine N-oxide

[209684-48-0]

C₂₃H₂₅NO₅ 395.454

Alkaloid from *Tylophora tanakae*. Fine brownish prisms (MeOH). Mp 194-204° dec. $[\alpha]_D^{28}$ -30.9 (c, 0.1 in CHCl₃/MeOH). λ_{max} 220 (log ϵ 4.51); 237 (log ϵ 4.57); 250 (sh) (log ϵ 4.72); 257 (log ϵ 4.9); 282 (sh) (log ϵ 4.5); 288 (log ϵ 4.46); 302 (log ϵ 4.26) (MeOH).

13 α -Hydroxy: 13 α -Hydroxytylophorine

[95066-46-9]

C₂₄H₂₇NO₅ 409.481

Alkaloid from the aerial parts of *Tylophora hirsuta* (Asclepiadaceae). Mp 274-275°. $[\alpha]_D^{25}$ +45 (c, 0.5 in

CHCl₃).

14 β -Hydroxy: Tylophorinicine.**14-Hydroxytylophorine**

[87302-57-6]

C₂₄H₂₇NO₅ 409.481

Minor alkaloid from the roots of *Tylophora asthmatica* and *Pergularia pallida* (Asclepiadaceae). Mp 210-212° dec. $[\alpha]_D^{25}$ -9.4 (c, 0.04 in CHCl₃). λ_{max} 258 (log ϵ 4.4); 287 (log ϵ 4.2); 302 (log ϵ 3.6); 339 (log ϵ 3.1) (MeOH).

(S)-form [482-20-2]

Obt. by resoln. of racemate or by synth. Mp 292° dec. $[\alpha]_D^{30}$ +12.25 (c, 0.733 in CHCl₃). $[\alpha]_D^{25}$ +73 (c, 0.7 in CHCl₃). Opt. rotn. gradually decreases in soln.

(±)-form [25908-92-3]

Synthetic. Cryst. (CHCl₃/EtOH). Mp 292° (263-265°, 287°).

(ξ)-form**O-De-Me: Demethyltylophorine**

[36250-91-6]

C₂₃H₂₅NO₄ 379.455

Alkaloid from *Tylophora indica* (Asclepiadaceae). Lepidopteran oviposition stimulant. Mp 235-237° dec. Location of the phenolic group and stereochemistry not detd.

Govindachari, T.R. *et al.*, *J.C.S.*, 1954, 2801-2803 (*isol, uv*)

Govindachari, T.R. *et al.*, *Tetrahedron*, 1958, **4**, 311-324; 1960, **9**, 53-57; 1961, **14**, 284-287 (*uv, struct, synth, resoln*)

Pailer, M. *et al.*, *Monatsh. Chem.*, 1965, **96**,

1094-1102 (*isol, uv, ms*)

Chandrashekar, V. *et al.*, *Curr. Sci.*, 1968, **37**,

432-433 (*isol, synth, oxide*)

Wiegrebbe, W. *et al.*, *Annalen*, 1969, **721**, 154-

162 (*isol, uv*)

Chauncy, B. *et al.*, *Aust. J. Chem.*, 1970, **23**,

2503-2516 (*synth, uv, ir*)

Herbert, R.B. *et al.*, *Chem. Comm.*, 1970, 121

(*synth*)

Rao, K.V. *et al.*, *J. Pharm. Sci.*, 1971, **60**, 1725-

1726 (*isol, Tylophorine,*

Demethyltylophorine)

Govindachari, T.R. *et al.*, *J. Indian Chem. Soc.*, 1973, **50**, 1-9 (*rev*)

Govindachari, T.R. *et al.*, *J.C.S. Perkin I*,

1974, 1161-1165 (*ord*)

Mulchandani, N.B. *et al.*, *Phytochemistry*,

1976, **15**, 1561-1563 (*isol, uv, ir, ms*)

Liepa, A.J. *et al.*, *Chem. Comm.*, 1977, 826-827

(*synth*)

Herbert, R.B. *et al.*, *Chem. Comm.*, 1977, 955-

956 (*biosynth*)

Gellert, E. *et al.*, *Aust. J. Chem.*, 1978, **31**,

2095-2097 (*ord*)

Weinreb, S.M. *et al.*, *J.A.C.S.*, 1979, **101**, 5073-

5074 (*synth*)

Mangla, V.K. *et al.*, *Tetrahedron*, 1980, **36**,

2489-2490 (*synth*)

Bick, I.R.C. *et al.*, *Alkaloids (Academic*

Press), 1981, **19**, 217 (*tox*)

Khatri, N.A. *et al.*, *J.A.C.S.*, 1981, **103**, 6387-

6393 (*synth*)

Bhakuni, D.S. *et al.*, *Tetrahedron*, 1981, **37**,

401-407 (*biosynth*)

Cragg, J.E. *et al.*, *J.C.S. Perkin I*, 1982, 2477-

2485 (*synth*)

Buckley, T.F. *et al.*, *J.O.C.*, 1983, **48**, 4222-4232

(*synth, uv, ir, pmr, ms*)

Iida, H. *et al.*, *J.O.C.*, 1984, **49**, 2412-2418

(*synth, ir, pmr, ms*)

Mulchandani, N.B. *et al.*, *Phytochemistry*,

1984, **23**, 1206 (*Tylophorinicine*)

Bhutani, K.K. *et al.*, *Phytochemistry*, 1985, **24**,

2778-2780 (*13 α -Hydroxytylophorine*)

Nordlander, J.E. *et al.*, *J.O.C.*, 1987, **52**, 1627-

1630 (*synth, uv, ir, pmr*)

Iwasa, K. *et al.*, *J. Nat. Prod.*, 1988, **51**, 172-

175 (*synth, uv*)

Ihara, M. *et al.*, *J.C.S. Perkin I*, 1990, 2287-

2292 (*synth, pmr, cd, abs config*)

Pearson, W.H. *et al.*, *Tetrahedron*, 1994, **50**,

12293-12304 (*synth*)

Abe, F. *et al.*, *Phytochemistry*, 1995, **39**, 695-

699 (*7-Demethyltylophorine, Tylophorine*)

Ciufolini, M.A. *et al.*, *J.A.C.S.*, 1996, **118**,

12082-12089 (*synth*)

Comins, D.L. *et al.*, *J.O.C.*, 1997, **62**, 7435-

7438 (*synth*)

Abe, F. *et al.*, *Chem. Pharm. Bull.*, 1998, **46**,

767-769 (*N-oxides*)

Jin, Z. *et al.*, *Chin. Chem. Lett.*, 2004, **15**,

1164-1166 (*S-isomer, synth*)

Damu, A.G. *et al.*, *J. Nat. Prod.*, 2005, **68**,

1071-1075 (*N-oxides*)

Fürstner, A. *et al.*, *Chem. Eur. J.*, 2006, **12**,

7398-7410 (*synth*)

Zeng, W. *et al.*, *J.O.C.*, 2008, **73**, 6045-6047


(*synth*)

Wang, K.-L. *et al.*, *Tetrahedron*, 2008, **64**,

7504-7510 (*synth*)

Tylophorinine**T-696**

9,11,12,13,13a,14-Hexahydro-3,6,7-trimethoxydibenzo[f,h]pyrrolo[1,2-b]isoquinolin-14-ol, 9CI. Pergularinine



C₂₃H₂₅NO₄ 379.455

Shows weak antineoplastic activity. Log P 3.5 (calc).

(+)-form [52437-07-7]

Alkaloid from *Ficus hispida* (Moraceae). Cryst. (CHCl₃/EtOH or CH₂Cl₂/Et₂O). Mp 215-216° dec., 229-231° dec. $[\alpha]_D$ +116 (c, 0.25 in CHCl₃). $[\alpha]_D$ +161 (c, 1.5 in CHCl₃).

Ac:

Cryst. (CH₂Cl₂/hexane). Mp 176-177°.

Deoxy, O⁶-de-Me: Tylophoridicine A

[457057-53-3]

C₂₂H₂₃NO₃ 349.429

Alkaloid from the roots of *Tylophora ovata*. Needles (Me₂CO). Mp 226-228°. $[\alpha]_D^{23}$ +176 (c, 0.25 in CHCl₃). λ_{max} 258 (log ϵ 4.53); 286 (log ϵ 4.39); 311 (sh) (log ϵ 3.81); 343 (log ϵ 2.91) (MeOH).


(-)-form [571-70-0]

Alkaloid from *Tylophora asthmatica*, *Tylophora cordifolia*, *Tylophora flava*, *Tylophora ovata*, *Tylophora atrafoliculata* and *Tylophora mollissima* and *Pergularia pallida* (Asclepiadaceae). Cryst. (CHCl₃ or CHCl₃/MeOH). Mp 248-249° (233-235° dec.). $[\alpha]_D^{27}$ -14.2 (c, 1.76 in CHCl₃). Largely racemic. Pharmacol. active isomer. λ_{max} 258 (c 40500); 287 (c 21900); 340 (c 820) (EtOH) (Berdy).

Fujise, Y. *et al.*, *Chem. Lett.*, 1980, 631-632
(isol, uv, ir, pmr, cmr, struct)
Cooksey, C.J. *et al.*, *Molecules*, 2001, 6, 736-769 (rev)

Tyroscherin

[832150-38-6]

C₂₁H₃₅NO₂ 333.513

Abs. config. revised in 2008. Prod. by *Pseudallescheria* sp. Antitumour agent. Powder. Mp 123-127°. [α]_D²⁵ -21 (c, 0.35 in MeOH). λ_{max} 225 (ε 5500); 278 (ε 1100) (MeOH). λ_{max} 225 (ε 4300); 240 (ε 3600); 286 (ε 1100) (MeOH/NaOH).

Stereoisomer (?): **Antibiotic JM 971B**.

JM 971B

[92802-16-9]

C₂₁H₃₅NO₂ 333.513

From *Leptographium lundbergii* NRRL 12527. Inhibits yeast-like fungi. Sol. MeOH, CHCl₃, DMF, DMSO; fairly sol. Et₂O; poorly sol. H₂O, hexane. Stereochem. not determined. λ_{max} 277 (ε 1700); 283 (ε 1450) (CH₂Cl₂) (Berdy).

Stereoisomer (?), N-Me: **Antibiotic JM 971A**.

JM 971A

[92802-17-0]

C₂₂H₃₇NO₂ 347.54

From *Leptographium lundbergii* NRRL 12527. Inhibits yeast-like fungi. Sol. MeOH, DMSO, DMF, CHCl₃; fairly sol. Et₂O; poorly sol. H₂O, hexane. Stereochem. not determined. λ_{max} 278 (ε 1400); 284 (ε 1200) (CH₂Cl₂) (Berdy).

Stereoisomer (?), N-de-Me: **Antibiotic JM 971C**.

JM 971C

[92802-15-8]

C₂₀H₃₃NO₂ 319.486

Isol. from *Leptographium lundbergii* NRRL 12527. Inhibits yeast-like fungi. Sol. MeOH, CHCl₃, DMF, DMSO; fairly sol. Et₂O; poorly sol. H₂O, hexane. Stereochem. not determined. λ_{max} 277 (ε 1450); 283 (ε 1230) (CH₂Cl₂) (Berdy).


Ger. Pat., 1984, 3 333 553; *CA*, 102, 4458 (Antibiotics JM 971)

Hayakawa, Y. *et al.*, *J. Antibiot.*, 2004, 57, 634-638 (isol, uv, pmr, cmr)

Katsuta, R. *et al.*, *Tet. Lett.*, 2008, 49, 7042-7045 (synth, struct)

Tyrosine, 9CI, 8CI**T-700**

α -Amino-4-hydroxybenzenepropanoic acid. 2-Amino-3-(4-hydroxyphenyl)propanoic acid. 4-Hydroxy- α -aminohydrocinamic acid. p-Tyrosine. 3-p-Hydroxyphenyl- α -alanine. Tyr [55520-40-6]

C₉H₁₁NO₃ 181.191**(S)-form***L-form*. *FEMA 3736*

[60-18-4]

Widely distributed in plant and animal proteins. Needles (H₂O). Almost insol. H₂O (0.05 g/100 g at 25°). Mp 290-295° dec. (slow heat) Mp 314-318° dec. approx. (rapid heat). [α]_D²⁰ -11.8 (c, 4.0 in 5M HCl) (opt. pure). [α]_D²⁰ -9.01 (11.6% KOH). pK_{a1} 2.2; pK_{a2} 9.11; pK_{a3} 10.07 (phenolic OH). Isoelectric point 5.66. Bitter taste.

► Exp. reprod. and teratogenic effects (large doses). YP2275600

Hydrochloride: [16870-43-2]Mp 239° dec. [α]_D²⁰ -8 (c, 4 in 1M HCl).

O- β -D-Glucopyranoside: **Tyrosinyl glucoside**

[38292-17-0]

C₁₅H₂₁NO₈ 343.333

Isol. from the seeds of *Entada pursaetha* and *Entada scandens*. Also found in the insects *Drosophila* sp. and *Manduca* sp. Needles (EtOH aq.). Mp 282° dec. [α]_D²⁰ -62.5 (c, 1.3 in 0.1M HCl). [α]_D²⁰ -77 (c, 1 in 0.1M HCl).

N-Octanoyl: N-Octanoyltyrosine. **Antibiotic CSL 12A**. *CSL 12A*

C₁₇H₂₅NO₄ 307.389

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Nonanoyl: N-Nonanoyltyrosine. **Antibiotic CSL 12B**. *CSL 12B*

C₁₈H₂₇NO₄ 321.416

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Decanoyl: N-Decanoyltyrosine. **Antibiotic CSL 12C**. *CSL 12C*

C₁₉H₂₉NO₄ 335.442

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Undecanoyl: N-Undecanoyltyrosine. **Antibiotic CSL 12D**. *CSL 12D*

C₂₀H₃₁NO₄ 349.469

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Dodecanoyl: N-Dodecanoyltyrosine. **Antibiotic CSL 12E**. *CSL 12E*

C₂₁H₃₃NO₄ 363.496

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-(7Z-Dodecenoyl): N-(7-Dodecenoyl)-tyrosine. **Antibiotic CSL 12J**. *CSL 12J*

C₂₁H₃₁NO₄ 361.48

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Tridecanoyl: N-Tridecanoyltyrosine. **Antibiotic CSL 12F**. *CSL 12F*

C₂₂H₃₅NO₄ 377.523

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Tetradecanoyl: N-Tetradecanoyltyrosine. **Antibiotic CSL 12G**. *CSL 12G*

C₂₃H₃₇NO₄ 391.55

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-(8Z-Tetradecenoyl): N-(8-Tetradecenoyl)tyrosine. **Antibiotic CSL 12K**.

*CSL 12K*C₂₃H₃₅NO₄ 389.534

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Pentadecanoyl: N-Pentadecanoyltyrosine. **Antibiotic CSL 12H**. *CSL 12H*

C₂₄H₃₉NO₄ 405.576

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-Hexadecanoyl: N-Hexadecanoyltyrosine. **Antibiotic CSL 12I**. *CSL 12I*

C₂₅H₄₁NO₄ 419.603

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-(9Z-Hexadecenoyl): N-(9-Hexadecenoyl)tyrosine. **Antibiotic CSL 12L**.

*CSL 12L*C₂₅H₃₉NO₄ 417.587

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-(11Z-Octadecenoyl): N-(11-Octadecenoyl)tyrosine. **Antibiotic CSL 12M**.

*CSL 12M*C₂₇H₄₃NO₄ 445.641

Prod. by a cosmid DNA library obt. from soil microorganisms.

N-(4,6R-Dimethyl-2E,4E-dodecadienoyl): **Gymnastatin N**

C₂₃H₃₃NO₄ 387.518

Prod. by the fungus *Arachniotus punctatus*. Plk1 inhibitor. Oil. [α]_D³⁰ -32.5 (c, 0.2 in EtOH). Isol. as a mixt. of (2R,6'R)- and (2S,6'R)-diastereoisomers to which data refers.

N-(4,6R-Dimethyl-2E,4E-dodecadienoyl), Me ester: **Gymnastatin H**

[229334-21-8]

C₂₄H₃₅NO₄ 401.545

Prod. by *Gymnascella dankaliensis* isol. from the sponge *Halichondria japonica*. Oil. [α]_D²⁶ +42.3 (c, 0.14 in CHCl₃). λ_{max} 238 (log ε 4.18); 266 (log ε 4.51) (EtOH).

N-(12-Hydroxy-4,6R-dimethyl-2E,4E-dodecadienoyl): **12'-Hydroxygymnastatin N**

C₂₃H₃₃NO₅ 403.517

Prod. by the fungus *Arachniotus punctatus*. Oil.

N-Me: 3-(4-Hydroxyphenyl)-2-(methylamino)propanoic acid. **N-Methyltyrosine, 9CI**. *Geoffroyin*. *Angelin*.

Andirin. *Surinamine*. *Ratanhin*

[537-49-5]

C₁₀H₁₃NO₃ 195.218

Present in bark of legumes *Geoffroea surinamensis*, *Andira anthelmintica* and *Andira retusa*. Also from *Ratanhia* root (*Krameria* sp.) and seeds of *Combretum zeyheri*. Needles. Mp 257°. [α]_D²³ +16 (c, 0.42 in 1M HCl). [α]_D²⁵ +31.45 (c, 0.42 in 1M NaOH). Dec. at 280°.

N,N-Di-Me: N,N-Dimethyltyrosine

[17350-74-2]

C₁₁H₁₅NO₃ 209.244


Aminoacid from *Lobaria laetevirens*. Cryst. Mp 274°.

N,N,N-Tri-Me, betaine: α -Carboxy-4-hydroxy-N,N,N-trimethylbenzeneethana-

- minium*(1+), 9CI. Tyrosine betaine.
Maokonine
[69168-08-7]
C₁₂H₁₈NO₃[⊕] 224.279
Constit. of *Lobaria laetevirens* and
Ephedra sp. Hypertensive agent. Mp
257-259°.
[36546-50-6, 16978-66-8, 81158-85-2, 126060-58-0]
- Ebata, M. *et al.*, *Bull. Chem. Soc. Jpn.*, 1966, **39**, 2535-2538 (*N-Me*, synth)
Tamada, M. *et al.*, *Planta Med.*, 1978, **34**, 291-293 (*Maokonine*)
Van Heerden, F.R. *et al.*, *Phytochemistry*, 1980, **19**, 2125-2129 (*N-Coumaroyltyrosine*)
Bernard, T. *et al.*, *Phytochemistry*, 1981, **20**, 2325-2326 (*N,N-Dimethyltyrosine*)
Brueckner, C. *et al.*, *Phytochemistry*, 1986, **25**, 2236-2237 (*N-Jasmonoyltyrosine*)
- Brady, S.F. *et al.*, *J.A.C.S.*, 2000, **122**, 12903-12904 (*Antibiotics CSL 12*)
Phoon, C.W. *et al.*, *Tetrahedron*, 2004, **60**, 11619-11628 (*Gymnastatin N*)
Kramell, R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1345-1349 (*N-Jasmonoyltyrosine*)
Amagata, T. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1384-1388 (*Gymnastatin H*)


Ugibohlin

[391936-28-0]

C₁₁H₁₁Br₂N₅O 389.049

Isol. from the marine sponge *Axinella carteri*. Amorph. solid. λ_{\max} 208 (log ϵ 3.83); 240 (log ϵ 4.25); 242 (log ϵ 4.27); 269 (log ϵ 3.6) (MeOH).

Goetz, G.H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1581-1582 (*isol*, *pnr*, *cmr*)

Ulapualides

Ulapualide A R = O
Ulapualide B R = H, COCOH(OMe)CH₂OMe

Ulapualide A [100045-73-6]C₄₆H₆₄N₄O₁₃ 881.031

Isol. from *Hexabranchnus sanguineus* egg masses. Shows antitumour and antifungal props. Oil. $[\alpha]_D^{25}$ -43.3 (c, 0.3 in MeOH). λ_{\max} 246 (ϵ 34000) (MeOH) (Derep).

Ulapualide B [100045-74-7]C₅₁H₇₄N₄O₁₆ 999.163

Isol. from *Hexabranchnus sanguineus* egg masses. Shows antitumour and antifungal props. Sol. MeOH, CHCl₃. $[\alpha]_D^{25}$ -21.7 (c, 0.138 in MeOH). λ_{\max} 246 (ϵ 33000) (MeOH) (Derep).

Roesener, J.A. *et al.*, *J.A.C.S.*, 1986, **108**, 846-847 (*isol*, *pnr*)


Chattopadhyay, S.K. *et al.*, *J.C.S. Perkin 1*, 2000, 2415-2428; 2429-2454 (*synth*, *config*)

Allingham, J.S. *et al.*, *Org. Lett.*, 2004, **6**, 597-599 (*cryst struct*, *abs config*)

Pattenden, G. *et al.*, *Org. Biomol. Chem.*, 2008, **6**, 1478-1497 (*synth*, *abs config*)

Ulbactin A

[184971-57-1]



U-1

C₁₃H₁₆N₂O₂S₂ 296.414

Similar to Pyochelin I, P-832. Prod. by *Vibrio* sp. B-93. Uv absorbant for cosmetics.

Kikuchi, K. *et al.*, *CA*, 1997, **126**, 44673w


(isol)

Japan. Pat., 1998, 98 101 657; *CA*, **129**, 3913t

(isol)

Ulbactin B

[184763-21-1]



U-4

C₁₇H₂₀N₂O₃S₂ 364.489

Prod. by *Vibrio* sp. B-93. Uv absorbant for cosmetics.

Kikuchi, K. *et al.*, *CA*, 1997, **126**, 44673w


(isol)

Japan. Pat., 1998, 98 101 657; *CA*, **129**, 3913t

(isol)

Ulbactin C

[184971-60-6]



U-5

C₁₉H₂₆N₄O₄S₃ 470.637

Similar to Yersiniabactin, Y-9. Prod. by *Alteromonas* sp. D-12. Uv absorbant for cosmetics.

Kikuchi, K. *et al.*, *CA*, 1997, **126**, 44673w

(isol)


Japan. Pat., 1998, 98 101 657; *CA*, **129**, 3913t

(isol)

Ulbactin D

3-[4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolyl]tetrahydro-6,9-imino-1H,3H,5H-thiazolo[4,3-c][1,4]thiazepin-5-one, 9CI
[213185-50-3]

U-6

C₁₆H₁₇N₃O₂S₃ 379.527


Prod. by *Alteromonas* sp. D-12 (PERM P-15733). Uv absorbant.

Japan. Pat., 1998, 98 245 377; *CA*, **129**, 244217z

Ulbactin E

2-(4,4',5,5'-Tetrahydro[4,4'-bithiazol]-2-yl)phenol, 9CI. 4,4',5,5'-Tetrahydro-2-(2-hydroxyphenyl)-4,4'-bithiazole
[213185-52-5]

U-7

C₁₂H₁₂N₂OS₂ 264.372


Prod. by *Alteromonas* sp. D-12 (FERM P-15733). Uv absorbant.

Japan. Pat., 1998, 98 245 377; *CA*, **129**, 244217z

Uleine

1-Methylenedascarpidan, 9CI
[517-81-7]

U-8



Absolute Configuration

C₁₈H₂₂N₂ 266.385

Alkaloid from *Aspidosperma ulei*, *Aspidosperma australe*, *Aspidosperma formosanum* and *Plumeria lancifolia*. Shows antitumour activity vs. Walker 256 carcinosarcoma. Poorly cryst. substance showing solvation and wide melting range. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 76-78° Mp 115-118° (double Mp; imprecise). $[\alpha]_D^{25}$ +18.5 (CHCl₃). λ_{\max} 307 (ϵ 17000); 316 (ϵ 17000) (MeOH).

Hydrochloride:

Cryst. (2-propanol/Et₂O). Mp 241-242° dec.

Methiodide:

Cryst. (EtOH). Mp 204-206°.

N-Oxide: Uleine N-oxide. Pausperidine A [878007-11-5]

C₁₈H₂₂N₂O 282.385

Alkaloid from the bark of *Geissospermum vellosii*. Amorph. solid. $[\alpha]_D^{25}$ +2.3 (c, 0.5 in MeOH). λ_{\max} 208 (log ϵ 4.03); 242 (sh) (log ϵ 3.65); 307 (log ϵ 3.65) (MeOH).

N-De-Me: De-N-methyluleine

[3620-83-5]

C₁₇H₂₀N₂ 252.358

Alkaloid from the bark of *Aspidosperma dasycarpon* (Apocynaceae). Cryst. (CHCl₃). Mp 143-145°. $[\alpha]_D^{25}$ -20 (c, 1.18 in EtOH). λ_{\max} 305 (log ϵ 4.24); 312 (log ϵ 4.23) (EtOH).

Didehydro, N-de-Me: Dehydrode-N-methyluleine

[3620-82-4 ($\Delta^{3(18)}$ -isomer), 3620-77-7 (Δ^{17} -isomer)]

$C_{17}H_{18}N_2$ 250.343

Alkaloid from the bark of *Aspidosperma dasycarpon* (Apocynaceae). Mp 220°. Full struct. not known. Double bond located between C_{3-18} or C_{17-18} . λ_{max} 303 (log ϵ 4.21); 310 (log ϵ 4.21) (EtOH).

1\alpha,13-Dihydro, 13-hydroxy: 1,13-Dihydro-13-hydroxyuleine. Dasycarpidan-1-methanol. 1-Hydroxymethyl-dasycarpidan

[60865-02-3]

$C_{18}H_{24}N_2O$ 284.4

Alkaloid from *Aspidosperma dasycarpon* and *Aspidosperma formosanum* (Apocynaceae). Amorph. $[\alpha]_D$ -96 (c, 0.25 in EtOH). λ_{max} 211 (log ϵ 4.56); 282 (log ϵ 3.91); 289 (log ϵ 3.87) (EtOH).

3-Epimer: Epiuleine

[17463-46-6]

$C_{18}H_{22}N_2$ 266.385

Alkaloid from *Aspidosperma formosanum* and *Aspidosperma subincanum* (Apocynaceae). Amorph. Racemate is crystalline. λ_{max} 213 (log ϵ 4.38); 307 (log ϵ 4.28); 315 (log ϵ 4.24) (MeOH).

[19775-51-0 (3-epimer \pm -form), 19775-50-9 (\pm -form)]

Schmutz, J. et al., *Helv. Chim. Acta*, 1957, **40**, 1189-1200 (*isol, uv, ir*)

Büchi, G. et al., *J.A.C.S.*, 1959, **81**, 4433-4434; 1971, **93**, 2492-2501 (*Uleine, Epiuleine, ir, pmr, ms, struct, synth*)

Ondetti, M.A. et al., *Tetrahedron*, 1961, **15**, 160-166 (*isol, uv*)

Ohashi, M. et al., *Experientia*, 1964, 363-364 (*De-N-methyluleine, isol*)

Joule, J.A. et al., *J.C.S.*, 1964, 2777-2790 (*ms, struct*)

Joule, J.A. et al., *Tetrahedron*, 1965, **21**, 1717-1734 (*De-N-methyluleine, Dehydrode-N-methyluleine, 1,13-Dihydro-13-hydroxyuleine*)

Gaskell, A.J. et al., *Chem. Ind. (London)*, 1967, 1089-1090 (*Uleine, Epiuleine, uv, ir, pmr, struct*)

Jackson, A. et al., *J.C.S. (C)*, 1969, 2738-2747 (*Epiuleine, synth*)

Dolby, L.J. et al., *J.O.C.*, 1970, **35**, 3843-3845 (*Epiuleine, synth*)

Barth, G. et al., *Helv. Chim. Acta*, 1972, **55**, 2168-2178 (*uv, cd*)

Garcia, M. et al., *Phytochemistry*, 1976, **15**, 1093-1095 (*Epiuleine, 1,13-Dihydro-13-hydroxyuleine*)

Natsume, M. et al., *Tet. Lett.*, 1980, **21**, 839-840 (*Epiuleine, synth*)

Harris, M. et al., *Tet. Lett.*, 1981, **22**, 331-334 (*Epiuleine, synth*)

Borris, R.P. et al., *J. Nat. Prod.*, 1983, **46**, 200-205 (*cmr*)

Grierson, D.S. et al., *Tetrahedron*, 1983, **39**, 3683-3694 (*Epiuleine, synth*)

Bonjoch, J. et al., *Chem. Comm.*, 1991, 1687-1688 (*synth*)

Saito, M. et al., *Chem. Comm.*, 1997, 765-766 (*synth, abs config*)


Amat, M. et al., *J.O.C.*, 2004, **69**, 8681-8693 (*synth*)

Ishiyama, H. et al., *Heterocycles*, 2005, **66**, 651-658 (*Pausperadine A*)

Baggio, C.H. et al., *Planta Med.*, 2005, **71**, 733-738 (*Uleine, isol, pmr, cmr*)

Ulithyclamide, 9CI

[74839-81-9]



$C_{33}H_{39}N_7O_5S_2$ 677.847

Numbering systems differ. The closely related Lissoclinamides are numbered differently from Ulithyclamide in CAS. Stereochem. given here is tentative as it is not completely clear in the lit. Isol. from the marine tunicate *Lissoclinum patella*. Cytotoxic. Oil. Sol. MeOH, C_6H_6 ; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ +35.7 (c, 2.3 in CH_2Cl_2). λ_{max} 248 (ϵ 7900) (MeOH) (Derep).

7,14-Diepimer, 10,11\alpha-dihydro: Lissoclinamide 3

[87393-59-7]

[121209-52-7]

$C_{33}H_{41}N_7O_5S_2$ 679.863

Isol. from *Lissoclinum patella*. Sol. $CHCl_3$, MeOH; poorly sol. H_2O . λ_{max} 248 (ϵ 7900) (MeOH) (Derep).

7,32-Diepimer, 10,11\alpha-dihydro: Lissoclinamide 2

[87393-58-6]

$C_{33}H_{41}N_7O_5S_2$ 679.863

Isol. from *Lissoclinum patella*.

Ireland, C. et al., *J.A.C.S.*, 1980, **102**, 5688-5691 (*isol, uv, ir, pmr, ms*)


Wasyluk, J.M. et al., *J.O.C.*, 1983, **48**, 4445-4449 (*isol, ir, pmr, cmr, ms, struct, derivs*)

Schmidt, U. et al., *Angew. Chem., Int. Ed.*, 1985, **24**, 569-571 (*synth*)

Sugiura, T. et al., *Tet. Lett.*, 1987, **28**, 2251-2254 (*synth*)

Ulithiacyclamide

[74847-09-9]



$C_{32}H_{42}N_8O_6S_4$ 762.998

Isol. from the marine tunicate *Lissoclinum patella* and from an unidentified sp. of ascidian collected from Rodda Reef, Queensland, Australia. Shows selective

U-9

metal binding props. Cytotoxic agent. Oil. Sol. MeOH, C_6H_6 ; poorly sol. H_2O , hexane. $[\alpha]_D^{25}$ +62.4 (c, 2.9 in CH_2Cl_2). λ_{max} 247 (ϵ 7000) (MeOH) (Derep).

Ireland, C. et al., *J.A.C.S.*, 1980, **102**, 5688-5691 (*uv, ir, pmr, cmr, ms, struct*)

Hamamoto, Y. et al., *Chem. Comm.*, 1983, 323-324 (*occur*)

Biskupiak, J.E. et al., *J.O.C.*, 1983, **48**, 2302-2304 (*abs config*)

Kato, S. et al., *Tet. Lett.*, 1986, **27**, 2653-2656 (*synth*)


Schmidt, U. et al., *Tet. Lett.*, 1986, **27**, 3495-3496 (*synth*)

Ishida, T. et al., *J.O.C.*, 1989, **54**, 5337-5343 (*pmr, conformn*)

Morris, L.A. et al., *Tetrahedron*, 2001, **57**, 3185-3197 (*activity*)

Ulithiacyclamide B

[122759-67-5]




$C_{35}H_{40}N_8O_6S_4$ 797.015

Isol. from the ascidian *Lissoclinum patella*. Potent cytotoxin. Amorph. solid. $[\alpha]_D^{24.5}$ +117 (c, 0.17 in MeOH). λ_{max} 202 (ϵ 154897); 235 (ϵ 63291) (MeOH) (Berdy).

Williams, D.E. et al., *J. Nat. Prod.*, 1989, **52**, 732-739 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ulithiacyclamide E

[218916-92-8]



$C_{35}H_{44}N_8O_8S_4$ 833.045

Isol. from the ascidian *Lissoclinum patella*. Amorph. solid. $[\alpha]_D$ +4.9 (c, 0.8 in MeOH).

Fu, X. et al., *J. Nat. Prod.*, 1998, **61**, 1547-1551 (*isol, ir, pmr, cmr, ms*)


U-10

Absolute configuration

Ulithiacyclamide F

U-13

[218916-93-9]




$C_{35}H_{42}N_8O_7S_4$ 815.03
Isol. from the ascidian *Lissoclinum patella*. Amorph. solid. $[\alpha]_D^{25} +29.6$ (c, 0.27 in MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1547-1551 (isol, ir, pmr, cmr)

Ulithiacyclamide G

U-14

[218916-94-0]




$C_{35}H_{42}N_8O_7S_4$ 815.03
Isol. from the ascidian *Lissoclinum patella*. Amorph. solid. $[\alpha]_D^{25} +25.6$ (c, 0.18 in MeOH).

Fu, X. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1547-1551 (isol, ir, pmr, cmr, ms)

Ulongamide B

U-15



R = CH(CH₃)₂

$C_{32}H_{45}N_5O_7S$ 643.803
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +10$ (c, 0.1 in MeOH). λ_{max} 201 (log ϵ 4.24); 230 (log ϵ 3.87) (MeOH).

4''-Deoxy: Ulongamide A
 $C_{32}H_{45}N_5O_6S$ 627.803
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +12$ (c, 0.73 in MeOH). λ_{max} 201 (log ϵ 4.25); 230 (log ϵ 3.88) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 996-1000 (isol, pmr, cmr)
Alvaredo, C. *et al.*, *Tet. Lett.*, 2007, **48**, 603-607 (synth)

Ulongamide C


U-16

As Ulongamide B, U-15 with R = CH₂Ph
 $C_{36}H_{45}N_5O_7S$ 691.847
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +9$ (c, 0.33 in MeOH). λ_{max} 202 (log ϵ 4.54); 230 (log ϵ 4.12) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 996-1000 (isol, pmr, cmr)

Ulongamide D

U-17



R¹ = -CH₂-C₆H₄-OH, R² = CH₃

$C_{34}H_{49}N_5O_7S$ 671.856
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +22$ (c, 0.12 in MeOH). λ_{max} 201 (log ϵ 4.24); 230 (log ϵ 3.89) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 996-1000 (isol, pmr, cmr)

Ulongamide E

U-18

As Ulongamide D, U-17 with R¹ = CH₂Ph-4OH, R² = CH₂CH₃
 $C_{35}H_{51}N_5O_7S$ 685.883
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +20$ (c, 0.4 in MeOH). λ_{max} 202 (log ϵ 4.24); 230 (log ϵ 3.89) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 996-1000 (isol, pmr, cmr)

Ulongamide F

U-19

As Ulongamide D, U-17 with R¹ = CH(CH₃)₂, R² = CH₂CH₃
 $C_{31}H_{51}N_5O_6S$ 621.84
Isol. from a Palauan *Lyngbya* sp. Amorph. solid. $[\alpha]_D^{25} +25$ (c, 0.2 in


MeOH). λ_{max} 202 (log ϵ 3.85); 230 (log ϵ 3.5) (MeOH).

Luesch, H. *et al.*, *J. Nat. Prod.*, 2002, **65**, 996-1000 (isol, pmr, cmr)

Ulosantoin

U-20

[145644-07-1]




$C_5H_9N_2O_5P$ 208.11
Exists as zwitterion. Isol. from the sponge *Ulosa ruetzleri*. Shows marked insecticidal activity. Acetylcholinesterase inhibitor. Cryst. (Me₂CO/isooctane). Mp 127-128°. λ_{max} 224 (ϵ 5000) (MeOH) (Derep).

Van Wagenen, B.C. *et al.*, *J.O.C.*, 1993, **58**, 335-337 (isol, uv, ir, pmr, cmr, ms, crystal struct)

Umbellamine

U-21

Hunterine
[21851-24-1]



Absolute Configuration

$C_{41}H_{48}N_4O_4$ 660.855
Alkaloid from the root bark of *Hunteria umbellata*, *Hunteria eburnea* and *Hunteria congolana* (Apocynaceae). Needles (CHCl₃/MeOH). $[\alpha]_D^{24} -217$ (c, 0.45 in CHCl₃). Dec. >250°.


Neus, N. *et al.*, *Experientia*, 1960, **16**, 302 (*Hunterine*)

Morita, Y. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 89-103 (isol, uv, ir, pmr, ms, struct)

Uncaramine

U-22

[110883-27-7]



$C_{43}H_{52}N_4O_7$ 736.906
Alkaloid from the leaves of *Uncaria gambier* (Rubiaceae). Cream amorph. powder. Mp 300° dec. $[\alpha]_D^{20} +20$ (c, 0.2 in CHCl₃).

17-Epimer: Callophylline A
[141610-96-0]

$C_{43}H_{52}N_4O_7$ 736.906
Alkaloid from the leaves of *Uncaria*

callophylla (Rubiaceae).

16,17-Diepimer: **Calophylline**

[111322-40-8]

C₄₃H₅₂N₄O₇ 736.906

Alkaloid from leaves of *Uncaria callophylla* (Rubiaceae). Yellow powder. Mp 330° dec.


Goh, S.H. *et al.*, *Malays. J. Sci.*, 1986, **8**, 109-113; *CA*, **107**, 214793n (*Calophylline*)

Arnone, A. *et al.*, *J.C.S. Perkin 1*, 1987, 571-578 (*Uncaramine*)

Kam, T.-S. *et al.*, *Phytochemistry*, 1991, **30**, 3441-3444 (*Calophylline A*)

Uncialamycin

[870471-83-3]



Absolute Configuration

C₂₆H₁₇NO₆ 439.423


Prod. by an unidentified *Streptomyces* sp. Antibacterial agent. Bright purple oil. [α]_D +3300 (c, 0.005 in MeOH). λ_{max} 206 (ε 25000); 254 (ε 33000); 280 (sh); 320 (sh); 539 (ε 9400) (MeOH).

Davies, J. *et al.*, *Org. Lett.*, 2005, **7**, 5233-5236 (*isol, pmr, cmr*)

Nicolaou, K.C. *et al.*, *Angew. Chem., Int. Ed.*, 2007, **46**, 4704-4707; 2008, **47**, 185-189 (*synth, config*)

Uncinine

U-24



C₁₂H₁₅NO₃ 221.255

Alkaloid from *Artabotrys uncinatus*. Amorph. powder. λ_{max} 204 (log ε 3.76); 291 (log ε 4.27) (EtOH).

Hsieh, T.-J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1157-1161

2,4-Undecadiene-7,9-diynoic acid

U-25

H₃CC≡CC≡CCH₂CH=CHCH=CHCOOH

C₁₁H₁₀O₂ 174.199

(2E,4Z)-form

2-Methylpropylamide: 2,4-Undecadiene-7,9-diynoic acid isobutylamide. **Neopeltitorine A**

C₁₅H₁₉NO 229.321

Alkaloid from *Artemisia dracunculus* (tarragon). Yellow oil. λ_{max} 257 (MeOH).

Saadali, B. *et al.*, *Phytochemistry*, 2001, **58**, 1083-1086

2,4-Undecadiene-8,10-diynoic acid

U-26

HC≡CC≡CCH₂CH=CHCH=CHCOOH

C₁₁H₁₀O₂ 174.199

(2E,4E)-form

trans,trans-form

4,6-Heptadiynyl ester: [55309-64-3]

C₁₈H₁₆O₂ 264.323

Isol. from *Chrysanthemum macrotum*. Mp 35°.

2-Methylpropylamide: 2,4-Undecadiene-8,10-diynoic acid isobutylamide. N-(2-Methylpropyl)-2,4-undecadiene-8,10-diynamide

[13891-74-2]

C₁₅H₁₉NO 229.321

Isol. from *Chrysanthemum frutescens*, *Achillea macrophylla*, *Achillea ptarmica*, *Achillea millefolium* (yarrow), *Otanthus maritimus*, *Anacyclus clavatus* and other spp. Cryst. (Et₂O/petrol). Mp 99-100°. λ_{max} 252 (Et₂O).

2-Methylbutylamide: N-(2-Methylbutyl)-2,4-undecadiene-8,10-diynamide

[99615-78-8]

C₁₆H₂₁NO 243.348

Isol. from *Acmella ciliata*.

3-Methylbutylamide: 2,4-Undecadiene-8,10-diynoic acid isopentylamide. 2,4-Undecadiene-8,10-diynecarboxy-N-isopentylamide

[112663-71-5]

C₁₆H₂₁NO 243.348

Isol. from the root, leaf and flowers of *Achillea ptarmica* (Asteraceae). Non-cryst.

2-Phenylethylamide: N-(2-Phenylethyl)-2,4-undecadiene-8,10-diynamide

[37064-11-2]

C₁₉H₁₉NO 277.365

Isol. from the roots of *Anacyclus pyrethrum*, *Achillea ptarmica* and *Achillea falcata*. Cryst. (Et₂O/petrol). Mp 117°. λ_{max} 250 (no solvent reported).

Piperidine: 2,4-Undecadiene-8,10-diynoic acid piperidine

C₁₆H₁₉NO 241.332

Alkaloid from *Achillea sudetica* (Asteraceae) and *Achillea ptarmica*. Oil. Not obt. completely pure.

2,3-Didehydropiperidine: 2,4-Undecadiene-8,10-diynoic acid 2,3-dehydropiperidine

[52704-38-8]

C₁₆H₁₇NO 239.316

Alkamide from *Otanthus maritimus*. Isol. from *Achillea millefolium* (yarrow) and *Achillea ptarmica* (Asteraceae). Oil.

(2E,4Z)-form

Shows light-mediated antifungal activity.

2-Methylpropylamide: [113817-70-2]

Identified in *Echinacea angustifolia* and *Echinacea purpurea*. Cryst. (hexane). Mp 61°. λ_{max} 259 (no solvent reported).

2-Methylbutylamide: [99615-79-9]

Isol. from *Acmella ciliata* and roots of *Echinacea purpurea*. Oil. λ_{max} 261 (no solvent reported).

2-Phenylethylamide: [37064-15-6]

Cryst. (Et₂O/petrol). Mp 78°.

(2Z,4E)-form

2-Methylpropylamide: [13894-69-4]

Isol. from roots of *Echinacea purpurea*, *Echinacea angustifolia*, *Spilanthes alba*. Cryst. (petrol). Mp 60-61°.

2-Phenylethylamide: [99615-80-2]

Isol. from *Acmella ciliata*.

[52997-51-0, 92264-88-5]

Bohlmann, F. *et al.*, *Chem. Ber.*, 1966, **99**, 3197; 1967, **100**, 104; 3861; 1972, **105**, 1694; 1973, **103**, 1328; 1974, **107**, 1038; 1975, **108**, 739 (*isol, synth, uv, ir, pmr*)

Jente, R. *et al.*, *Chem. Ber.*, 1972, **105**, 1694 (*synth, isol, ms, struct, pmr, ir, uv*)

Bohlmann, F. *et al.*, *Phytochemistry*, 1980, **19**, 1535 (*isol*)

Greger, H. *et al.*, *Phytochemistry*, 1984, **23**, 1503; 1989, **28**, 2363 (*isol*)

Martin, R. *et al.*, *Phytochemistry*, 1985, **24**, 2295 (*isol, ms, pmr*)

Kuroopka, G. *et al.*, *Planta Med.*, 1986, **52**, 244; 1987, **53**, 440 (*isol, uv, ir, pmr, ms, struct*)

Hofer, O. *et al.*, *Tetrahedron*, 1986, **42**, 2707 (*cmr*)

Jakupovic, J. *et al.*, *Phytochemistry*, 1988, **27**, 1135 (*isol*)

Bauer, R. *et al.*, *Phytochemistry*, 1988, **27**, 2339; 1989, **28**, 505 (*isol, pmr, ms, ir, uv*)

Bauer, R. *et al.*, *Planta Med.*, 1989, **55**, 367 (*hplc*)

Binns, S.E. *et al.*, *Planta Med.*, 2000, **66**, 241-242 (*activity, 2-methylpropylamide*)

Kraus, G.A. *et al.*, *Molecules*, 2006, **11**, 758-767 (*2Z,4E-isobutylamide, synth*)

3,5-Undecadiene-8,10-diynoic acid, 9CI

U-27

HC≡CC≡CCH₂CH=CHCH=CHCOOH

C₁₁H₁₀O₂ 174.199

(3E,5E)-form

2-Phenylethylamide: N-(2-Phenylethyl)-3,5-undecadiene-8,10-diynamide, 9CI.

3,5-Undecadiene-8,10-diynoic acid 2-phenylethylamide

[99615-82-4]

C₁₉H₁₉NO 277.365

Isol. from flower heads of *Acmella ciliata*.

Martin, R. *et al.*, *Phytochemistry*, 1985, **24**, 2295-2300 (*isol, struct, pmr, ms*)

2,4-Undecadienoic acid

U-28

H₃C(CH₂)₅CH=CHCH=CHCOOH

C₁₁H₁₈O₂ 182.262

(2E,4E)-form

2-Methylpropylamide: 2,4-Undecadienoic acid isobutylamide. N-(2-Methylpropyl)-2,4-undecadienamamide

[74267-82-6]

C₁₅H₂₇NO 237.384

Alkaloid from *Leucocyclus formosus* (Asteraceae). Yellow cryst. Obt. only in admixture with homologues.

Greger, H. *et al.*, *Phytochemistry*, 1981, **20**,

2579 (*isol, ms, struct*)
Plobeck, N.A. *et al.*, *J.O.C.*, 1991, **56**, 4508
(*synth, cmr*)

2,7,9-Undecatrienoic acid U-29

$\text{H}_3\text{CCH}=\text{CHCH}=\text{CH}(\text{CH}_2)_3\text{CH}=\text{CHCOOH}$
 $\text{C}_{11}\text{H}_{16}\text{O}_2$ 180.246

(2E,7Z,9E)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2,7,9-undecatrienamamide. *2,7,9-Undecatrienoic acid isobutylamide*
 $\text{C}_{15}\text{H}_{25}\text{NO}$ 235.369
Alkaloid from the dried flower buds of *Spilanthes acmella*.

Ramsewak, R.S. *et al.*, *Phytochemistry*, 1999, **51**, 729-732 (*isol, amide, pmr, cmr, ms*)

2-Undecene-8,10-diynoic acid U-30

$\text{HC}\equiv\text{CC}\equiv\text{C}(\text{CH}_2)_4\text{CH}=\text{CHCOOH}$
 $\text{C}_{11}\text{H}_{12}\text{O}_2$ 176.215

(E)-form

2-Methylpropylamide: N-(2-Methylpropyl)-2-undecene-8,10-diynamide. *2-Undecene-8,10-diynoic acid isobutylamide*
[99615-81-3]

$\text{C}_{15}\text{H}_{21}\text{NO}$ 231.337
Isol. from flowers of *Acemella ciliata* and *Echinacea angustifolia*. Cryst. (hexane). Mp 69°.

2-Methylbutylamide: N-(2-Methylbutyl)-2-undecene-8,10-diynamide

Isol. from flower heads of *Spilanthes acmella*. $[\alpha]_D^{25} +4.9$ (c, 0.41 in CHCl_3).

(Z)-form

2-Methylpropylamide: [120727-28-8]
Isol. from the roots of *Echinacea angustifolia*. Oil.

2-Methylbutylamide: [120727-30-2]
 $\text{C}_{16}\text{H}_{23}\text{NO}$ 245.364
Isol. from the roots of *Echinacea angustifolia*. Oil.

Martin, R. *et al.*, *Phytochemistry*, 1985, **24**, 2295-2300 (*isol, pmr, ms*)

Bauer, R. *et al.*, *Phytochemistry*, 1989, **28**, 505-508 (*isol, pmr, ms*)


Nakatani, N. *et al.*, *Biosci., Biotechnol., Biochem.*, 1992, **56**, 759 (*isol*)

Ramsewak, R.S. *et al.*, *Phytochemistry*, 1999, **51**, 729-732 (*isobutylamide, pmr, cmr*)

Wu, L. *et al.*, *Phytochemistry*, 2004, **65**, 2477-2484 (*isobutylamide, synth, occur*)

Undecylprodiginine U-31

Undecylprodigosine. Prodigiosin 25c
[13129-81-2]



$\text{C}_{25}\text{H}_{35}\text{N}_3\text{O}$ 393.571

Isol. from various *Streptomyces* spp., *Actinomadura pelletieri*, *Streptovorticillum rubriviculi* and *Nocardia pelletieri*.

Also from a marine-derived *Hahella chejuensis*. Antimalarial agent. Weakly active against gram-positive bacteria. Immunosuppressant, bone resorption inhibitor; gastric proton pump inhibitor and antiulcer agent. Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O, acids, bases. Exhibits dimorphism. λ_{max} 273 (ε 3640); 297 (ε 6900); 362 (ε 5000); 500 (sh) (ε 31500); 530 (ε 75900) (MeOH/HCl) (Derep). λ_{max} 285 (ε 6050); 325 (ε 4850); 467 (ε 30600) (MeOH/KOH) (Derep). λ_{max} 273 (ε 3640); 297 (ε 6900); 362 (ε 5000); 500 (sh) (ε 31500); 530 (ε 75900) (hydrochloride) (Derep).

▶ LD₅₀ (mus, ivn) 26.7 mg/kg.

Hydrochloride: [15377-21-6]

Red needles (heptane). Mp 76-78° Mp 106-107° (dimorph.).

Lower homologue (n = 8): **Nonylprodiginine**

[34852-34-1]

$\text{C}_{23}\text{H}_{31}\text{N}_3\text{O}$ 365.517

Prod. by *Nocardia madurae*, *Nocardia pelletieri* and *Streptomyces* sp. Y-42. Mp 150-155° (as perchlorate salt).

[14960-80-6, 56247-02-0, 52340-48-4]

Gerber, N.N. *et al.*, *Appl. Microbiol.*, 1969, **18**, 1-3 (*Nonylprodiginine*)

Gerber, N.N. *et al.*, *J. Antibiot.*, 1971, **24**, 636-640; 1975, **28**, 194-199 (*isol*)

Wasserman, H.H. *et al.*, *Tetrahedron*, 1976, **32**, 1851 (*isol, uv, ir, pmr, struct*)


D'Alessio, R. *et al.*, *Synlett*, 1996, 513 (*synth*)
Fürstner, A. *et al.*, *Angew. Chem., Int. Ed.*, 2003, **42**, 3582-3603 (*rev*)

Kim, D. *et al.*, *J. Appl. Microbiol.*, 2007, **102**, 937-944 (*isol*)

Jolicœur, B. *et al.*, *Can. J. Chem.*, 2008, **86**, 213-218 (*synth*)

Undulatine† U-32

Alkaloid 20
[123871-90-9]



$\text{C}_{44}\text{H}_{52}\text{N}_4\text{O}_6$ 732.918

Alkaloid from the trunk bark of *Alstonia sphaerocapitata* and the roots of *Alstonia undulata* (Apocynaceae). Amorph. $[\alpha]_D^{25} -32$ (c, 1 in CHCl_3).

De(hydroxymethyl): **Deformoundulatine**
[123901-45-1]

$\text{C}_{43}\text{H}_{50}\text{N}_4\text{O}_5$ 702.892


Alkaloid from the leaves and roots of *Alstonia undulata* (Apocynaceae). Amorph.

Caron, C. *et al.*, *Phytochemistry*, 1984, **23**, 2355 (*isol, uv, ir, pmr, ms*)

Nuzillard, J.-M. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. 2*, 1989, **309**, 195 (*isol, uv, cmr, ms, struct*)

Undulifoline U-33

[142750-31-0]



Absolute Configuration

$\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_3$ 340.421


Alkaloid from the stem bark of *Alstonia undulifolia* (Apocynaceae). $[\alpha]_D^{25} -33$ (c, 0.16 in CHCl_3). λ_{max} 218; 282; 290 (MeOH).

Massiott, G. *et al.*, *Phytochemistry*, 1992, **31**, 1078-1079 (*isol, uv, ir, pmr, cmr, ms, struct*)

Ungeremine U-34

Lycobetaine. AT 1840

[2121-12-2]



$\text{C}_{16}\text{H}_{11}\text{NO}_3$ 265.268

Alkaloid from the leaves of *Ungernia minor* and the fruits of *Crinum asiaticum*. Also isol. from the Texas grasshopper *Brachystola magna*. Natural metab. of Lycorine, L-348. DNA intercalator. Shows antineoplastic props. *in vitro*. Mp 270-272° dec.

▶ LD₅₀ (rat, orl) 90 mg/kg. Exp. reprod. effects (testicular). OL2827500

Hydrochloride: [2121-16-6]

Mp 314-315° dec.

Hydrobromide: Mp 327-329° dec.

Picrate: Mp 280-282° dec.

8-Methoxy: Zeflabetaine

[105708-77-8]

$\text{C}_{17}\text{H}_{13}\text{NO}_4$ 295.294

Alkaloid from fresh mature seeds of *Zephyranthes flava* (Amaryllidaceae). Light-brown powder (EtOH). Sol. MeOH, CHCl_3 ; poorly sol. H₂O. λ_{max} 262 (ε 47860); 292 (ε 7940); 305 (ε 9550); 415 (ε 3550) (MeOH) (Berdy). λ_{max} 267; 275; 282; 292; 368 (MeOH- CHCl_3) (Berdy). λ_{max} 267; 280; 435 (MeOH-NaOH) (Berdy).

8-Methoxy; hydrochloride:

Yellow powder + 1H₂O. Mp 300°.

[19488-20-1, 21326-01-2, 61221-41-8]

Normatov, M. *et al.*, *Uzb. Khim. Zh.*, 1965, **9**, 25; *CA*, **63**, 7061f (*struct*)


Allayarov, K. *et al.*, *Khim. Priir. Soedin.*, 1970, **6**, 143; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 143 (*isol*)

Ru-yun, J. *et al.*, *Drugs of the Future*, 1983, **8**, 118 (*rev*)

- Ghosal, S. *et al.*, *J. Chem. Res., Synop.*, 1986, 112 (*isol, uv, pmr, ms, synth*)
 Ghosal, S. *et al.*, *Phytochemistry*, 1986, **25**, 1975 (*Zeflabetaine*)
 Ghosal, S. *et al.*, *Planta Med.*, 1988, **54**, 114 (*tox*)
 Siddiqui, M.A. *et al.*, *Tet. Lett.*, 1990, **31**, 1523 (*synth*)
 Lauk, U. *et al.*, *Tet. Lett.*, 1991, **32**, 65 (*synth*)
 Stark, L.M. *et al.*, *J.O.C.*, 2000, **65**, 3227-3230 (*synth*)
 Pettit, G.R. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1256-1258 (*isol*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LJB800

Ungminorine

U-35



(-)-form

C₁₇H₁₉NO₅ 317.341

(-)-form [27857-09-6]

Alkaloid from the leaves of *Ungernia minor* and from the leaves and bulbs of *Leucojum aestivum* (Amaryllidaceae). Acetylcholinesterase inhibitor. Prisms (Me₂CO). Mp 208-211° (206-208° dec.). [α]_D²⁴ -49 (c, 0.67 in EtOH). [α]_D²⁴ -31.6 (c, 0.73 in CHCl₃).

Methiodide:

Cryst. (Me₂CO/MeOH). Mp 246-248° dec. [α]_D²⁰ -42.7 (c, 2.34 in EtOH).

Picrate: Mp 174-175° (170-172°).

N-Oxide (α-): *Ungminorine N-oxide*

[119308-28-0]

C₁₇H₁₉NO₆ 333.34

Alkaloid from bulbs of *Pancreatum maritimum* (Amaryllidaceae). Cryst. (MeOH). Mp 182-184°.

O³-Ac: *3-O-Acetylungminorine*

[101899-55-2]

C₁₉H₂₁NO₆ 359.378

Alkaloid from the leaves and bulbs of *Leucojum aestivum* (Amaryllidaceae). Prisms (C₆H₆). Mp 157-160°. [α]_D²⁴ -257.1 (c, 0.70 in CHCl₃).

Di-Ac:

Needles (Et₂O). Mp 172-174°. [α]_D²⁴ -126.8 (c, 0.47 in CHCl₃).

O-De-Me: *Pancreassinine*

[69787-51-5]

C₁₆H₁₇NO₅ 303.314

Alkaloid from the bulbs of *Pancreatum biflorum* infected with *Imperata cylindrica*. Straw-coloured microcryst. (MeOH). Mp 206-208°. [α]_D²² -28.2 (c, 0.51 in MeOH).

O-De-Me, tri-Ac: Mp 201-204°.

2-Epimer, O-de-Me: *2-Epipancreassinine*

[126381-95-1]

C₁₆H₁₇NO₅ 303.314

Alkaloid from the flower-stem fluid of *Crinum latifolium*. Straw-coloured microcryst. (2-propanol). Mp 207-210° dec. [α]_D²⁸ -112.5 (c, 0.7 in

MeOH).

16-Epimer: *Siculinine*

[126371-96-8]

C₁₇H₁₉NO₅ 317.341

Alkaloid from *Sternbergia sicula* (Amaryllidaceae). Amorph. [α]_D -34 (c, 1.9 in MeOH).

(±)-form

Alkaloid from *Ungernia vvedenskyi*. Mp 210-212°.

Normatov, M. *et al.*, *Uzb. Khim. Zh.*, 1965, **9**, 25-30; *CA*, **63**, 7061f (*isol*)

Razakov, R. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 19-22; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 15-17 (*ms*)

Clardy, J.C. *et al.*, *J.A.C.S.*, 1970, **92**, 1781-1782 (*struct*)

Kadyrov, K.A. *et al.*, *Khim. Prir. Soedin. (Engl. Transl.)*, 1979, **15**, 370-371 (*isol*)

Kobayashi, S. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 5258-5263 (*isol, ir, pmr, ms, deriv*)

Ghosal, S. *et al.*, *Phytochemistry*, 1986, **25**, 1097-1102; 1989, **28**, 2535-2537 (*Pancreassinine, 2-Epipancreassinine*)

Suau, R. *et al.*, *Phytochemistry*, 1988, **27**, 3285-3287 (*Ungminorine N-oxide*)


Richomme, P. *et al.*, *J. Nat. Prod.*, 1989, **52**, 1150-1152 (*Siculinine*)

Ingkaninan, K. *et al.*, *J. Nat. Prod.*, 2000, **63**, 803-806 (*isol, pmr, activity*)

Unonopsine

U-36

7,7'-Bis-(dehydroanonaine). *Bidebiline A* [112523-82-7]

C₃₄H₂₄N₂O₄ 524.575

Alkaloid from the roots of *Polyalthia debilis* and the bark of *Unonopsis spectabilis*. Yellowish amorph. powder. Mp 240° dec. λ_{max} 261 (log ε 4.65); 335 (log ε 4.1); 384 (log ε 3.89) (CHCl₃).

8-Methoxy: *Bidebiline B*

[543706-57-6]

C₃₅H₂₆N₂O₅ 554.601

Alkaloid from the roots of *Polyalthia debilis*. Yellow-green amorph. powder. Mp 245° dec. λ_{max} 260 (log ε 4.69); 335 (log ε 4.22); 384 (log ε 4.08) (CHCl₃).

3,3'-Dimethoxy: *Artabonatine F*

[368422-67-7]

C₃₆H₂₈N₂O₆ 584.627

Alkaloid from *Artabotrys uncinatus*. Amorph. powder. λ_{max} 244 (log ε 4.11); 264 (log ε 3.94); 324 (log ε 3.76) (EtOH).

8,8'-Dimethoxy: *Bidebiline C*

[543706-58-7]

C₃₆H₂₈N₂O₆ 584.627

Alkaloid from the roots of *Polyalthia debilis*. Exhibits moderate antimalarial

activity. Yellow-green amorph. powder. Mp 210° dec. λ_{max} 266 (log ε 4.72); 334 (log ε 4.29); 384 (log ε 4.16) (CHCl₃).

9,9'-Dimethoxy: *Bidebiline E*

[113425-61-9]

C₃₆H₂₈N₂O₆ 584.627

Alkaloid from the roots of *Polyalthia cerasoides*. Pale yellow solid. [α]_D²⁹ -42.8 (c, 0.2 in CHCl₃). Dec. at 250°. λ_{max} 272 (log ε 4.7); 343 (log ε 4) (MeOH).

10,10'-Dimethoxy: *Bidebiline D*

[543706-59-8]

C₃₆H₂₈N₂O₆ 584.627

Alkaloid from the roots of *Polyalthia debilis*. Yellow-green amorph. powder. Mp 205° dec. λ_{max} 256 (log ε 4.92); 333 (log ε 4.48); 384 (log ε 4.36) (CHCl₃).

Laprévôte, O. *et al.*, *J. Nat. Prod.*, 1987, **50**, 984-988 (*isol, uv, ir, pmr, ms*)


Hsieh, T.-J. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1157-1161 (*Artabonatine F*)

Kanokmedhakul, S. *et al.*, *J. Nat. Prod.*, 2003, **66**, 616-619; 2007, **70**, 1536-1538 (*Bidebilines*)

Unsevine

U-37

[4838-99-7]



Absolute configuration

C₁₈H₂₁NO₅ 331.368

Alkaloid from the roots and bulbs of *Ungernia severtzovii* (Amaryllidaceae). Mp 173-174°. [α]_D²⁰ +163 (c, 2.5 in CHCl₃).

Oxalate:

Cryst. (EtOH). Mp 195-196°.

Methiodide:

Cryst. (EtOH). Mp 249-250°. [α]_D²⁰ +146 (c, 1.56 in H₂O).

O-De-Me, O⁷-Me: *5-Hydroxy-7-O-methyloduline. 2-Hydroxy-6-O-methyloduline*C₁₈H₂₁NO₅ 331.368

Alkaloid from bulbs of *Narcissus* cv. Salome. Mp 245-248°. [α]_D²² +91.1 (c, 0.34 in CHCl₃). Authors' numbering scheme is nonstandard.

O-De-Me, O⁷-Et: *Radiatine*†C₁₉H₂₃NO₅ 345.394

Alkaloid from the rhizomes of *Lycoris radiata*. Cryst. (Me₂CO). Mp 171-172.5°. Assumed artifact from processing of the plant with EtOH. 7-config. not defined.

Smirnova, L.S. *et al.*, *Khim. Prir. Soedin.*, 1965, **1**, 322; *Chem. Nat. Compd. (Engl. Transl.)*, 1965, **1**, 252 (*uv, ir*)


Uyeo, S. *et al.*, *Yakugaku Zasshi*, 1965, **85**, 615; *CA*, **63**, 11632e (*Radiatine*)

Razahov, R. *et al.*, *Khim. Prir. Soedin.*, 1967, **3**, 23; *Chem. Nat. Compd. (Engl. Transl.)*, 1967, **3**, 18 (*ms*)

Yagudaev, M.R. *et al.*, *Khim. Prir. Soedin.*, 1970, **6**, 94; *Chem. Nat. Compd. (Engl. Transl.)*, 1970, **6**, 88 (*pmr, struct*)
 Clardy, J. *et al.*, *J.O.C.*, 1972, **37**, 49 (*config*)
 Almanza, G.R. *et al.*, *Phytochemistry*, 1996, **43**, 1375 (*Hydroxydemethyloduline*)

Uoamine A

[356550-05-5]

C₂₂H₄₁NO₃S 399.637

Alkaloid from the ascidian *Aplidium uouo*. Oil. [α]_D +5 (c, 0.16 in CHCl₃). λ_{max} 275 (ε 10740) (EtOH).

Z-Isomer: Uoamine B

[356550-06-6]


C₂₂H₄₁NO₃S 399.637

Alkaloid from *Aplidium uouo*. Oil. [α]_D +8 (c, 0.15 in CHCl₃). λ_{max} 283 (ε 1372) (EtOH).

McCoy, M.C. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1087-1089 (*Uoamines A,B, isol, pmr, cmr, uv*)

Upenamidine

U-39

C₃₂H₄₆N₂O₄ 522.726


Alkaloid from the Indonesian sponge *Echinocalina* sp. Amorph. solid. [α]_D -9.4 (c, 2.3 in MeOH).

Jimenez, J.I. *et al.*, *J.O.C.*, 2000, **65**, 8465-8469

Uplandicine

U-40

7-Acetyl-9-echimidinylretronecine
 [74202-10-1]


C₁₇H₂₇NO₇ 357.403

Ester of *O*-acetylretronecine with echimidinic acid. Alkaloid from the leaves of *Symphytum uplandicum* (Boraginaceae). Noncryst. [α]_D²⁰ +0.1 (c, 1.11 in EtOH).

Culvenor, C.C.J. *et al.*, *Aust. J. Chem.*, 1980, **33**, 1105 (*isol, pmr, ms, struct*)

Urbaine

5,5',6,6'-Tetrahydro-1,1',2,2'-tetramethoxy-7,7'-bi-4H-dibenzo[de,g]quinoline, 9CI
 [108906-93-0]

C₃₆H₃₂N₂O₄ 556.66

Alkaloid from the trunk bark of *Oxandra* cf. *major* and roots of *Piptostigma fugax* (Annonaceae). Also isol. from stem bark of *Polyalthia bullata*. Green powder. Mp 280°.

N-Me: N-Methylurbaine

[108906-94-1]

C₃₇H₃₄N₂O₄ 570.687

Alkaloid from *Oxandra* cf. *major* and *Polyalthia fugax* (Annonaceae). Cryst. (MeOH). Mp 262°.

N,N'-Di-Me: N,N'-Dimethylurbaine

[108906-95-2]

C₃₈H₃₆N₂O₄ 584.713

Alkaloid from *Oxandra* cf. *major* and *Polyalthia fugax* (Annonaceae). Mp 254°.

3-Methoxy: 7-Dehydronornuciferine-7-dehydro-O-methylisopiline. 3-Methoxyurbaine

[182295-42-7]

C₃₇H₃₄N₂O₅ 586.686

Alkaloid from stem bark of *Polyalthia bullata*. Yellowish-brown microcryst. (MeOH). Mp >310°.

3,3'-Dimethoxy: 7,7'-Bisdehydro-O-methylisopiline. 3,3'-Dimethoxyurbaine

[182295-40-5]

C₃₈H₃₆N₂O₆ 616.712

Alkaloid from stem bark of *Polyalthia bullata*. Yellowish-brown microcryst. (MeOH). Mp 268-270°.

Jossang, A. *et al.*, *Heterocycles*, 1987, **26**, 2191 (*synth, pmr*)


Arango, G. *et al.*, *Phytochemistry*, 1987, **26**, 1227 (*isol, uv, pmr, cmr, ms, struct, derivs*)

Achenbach, H. *et al.*, *Phytochemistry*, 1995, **38**, 1037 (*isol, pmr, cmr*)

Connolly, J.D. *et al.*, *Phytochemistry*, 1996, **43**, 295 (*7,7'-Bisdehydro-O-methylisopiline, 7-Dehydronornuciferine-7-dehydro-O-methylisopiline*)

Uracil, 8CI, USAN

2,4(1H,3H)-Pyrimidinedione, 9CI. 2,4-Dihydroxypyrimidine. 2,4-Pyrimidinediol. Hybar X
 [66-22-8]



U-41

C₄H₄N₂O₂ 112.088

Dioxo tautomer predominates in solid, soln. and gas phases. Two monooxo tautomers also possible. Widely occurring component of nucleic acids. Normally present in RNA only; an abnormal constit. of DNA as a consequence of cytosine deamination and reincorporation. Removed from DNA by action of DNA-glycosylases. Isol. in free state from ferns *Colysis hemionitidea* and *Microsorium fortunei* in large amts. Used in combination with Tegafur as a potentiator. Inhibits degradn. of the metab. 5-Fluoro-2,4-dihydroxypyrimidine, F-104. Needles (H₂O). Mp 335°. pK_{a1} -3.38; pK_{a2} 9.45; pK_{a3} 13 (25°). ▶YQ8650000

3-Oxide: Uracil N³-oxide. 3-Oxyuracil

[766-44-9]

Prod. by *Streptomyces lydicus* (Mk233, P8541). Antibacterial agent. Cryst. Sol. H₂O. Mp 290-293°.

1-β-D-Riburonofuranoside: Antibiotic AJP 117510. AJP 117510C₁₀H₁₂N₂O₇ 272.214

Prod. by an unidentified fungus AJ 117510. Inhibitor of integrin α₂β₁-collagen binding. Needles. Mp 202-205°. [α]_D²³ +30.8 (c, 0.5 in H₂O). λ_{max} 261 (ε 19100) (H₂O).

Kloetzer, W. *et al.*, *Monatsh. Chem.*, 1964, **95**, 1729-1739 (*oxides*)

Rice, J.M. *et al.*, *J.A.C.S.*, 1965, **87**, 4569 (*ms*)
 Stewart, R.F. *et al.*, *Acta Cryst.*, 1967, **23**, 1102 (*cryst struct*)

Ellis, P.D. *et al.*, *J.A.C.S.*, 1973, **95**, 4398 (*cmr*)
 Murakami, T. *et al.*, *Yakugaku Zasshi*, 1985, **105**, 655 (*isol*)


Japan. Pat., 1987, 87 143 692; *CA*, **107**, 216184p (*isol, 3-oxide*)

Sato, S. *et al.*, *J. Antibiot.*, 2006, **59**, 251-253 (*AJP 117510*)

Urdamycin D

U-43

[104443-44-9]

C₅₃H₆₁NO₁₈ 1000.061


Anthracycline antibiotic. Prod. by *Streptomyces fradiae*. Active against gram-positive bacteria and tumours. Blue powder. Sol. MeOH, EtOAc; poorly sol. H₂O, hexane. Mp 220° dec. λ_{max} 222 (ε 32400); 270 (sh); 324 (ε 7500); 412 (ε 4200); 600 (ε 8500); 625 (sh) (MeOH/NaOH) (Derep). λ_{max} 328 (ε 11600); 575 (ε 11200) (MeOH) (Derep).

Rohr, J. *et al.*, *J. Antibiot.*, 1988, **41**, 126; 1989, **42**, 1151 (*isol, struct, biosynth*)

Ureidosorbicillinol

U-44

[923582-49-4]

Absolute
ConfigurationC₁₇H₂₀N₂O₅ 332.355


Related to Rezhishanone A and Trichotetrone. Isol. from a marine-derived *Paecilomyces marquandii*. Yellow oil. $[\alpha]_D^{25} +50$ (c, 0.1 in MeOH). λ_{\max} 362 (log ϵ 4.3) (MeOH).

Cabrera, G.M. *et al.*, *J. Nat. Prod.*, 2006, **69**, 1806-1808 (isol, cd, pmr, cmr, ms)

Uric acid, 8CI

U-45

7,9-Dihydro-1H-purine-2,6,8(3H)-trione, 9CI. 2,6,8-Trihydroxypurine. 2,6,8(1H,3H,9H)-Purinetrione [69-93-2]

C₅H₄N₄O₃ 168.112

Exists in several tautomeric forms. Chief end-product of purine metab. Constit. of urine of carnivorous animals, bird excrement (guano), excrement of reptiles, insects etc. Produced by *Mamestra brassicae*. Present in small amts. in higher plants, esp. seeds. Prod. by marine bacterium strain Cr 11 from *Crassostrea gigas* and by a range of microorganisms from fish and shellfish. Sperm-release pheromone of *Platynereis dumerilii*. Odourless, tasteless, rhombic prisms or plates. Sol. alkalis, glycerol; spar. sol. min. acids; v. spar. sol. H₂O; insol. EtOH, Et₂O. pK_a 5.75; pK_{a2} 10.6. Dec. without melting.

► Exp. reprod. effects (male). May evolve HCN when heated. YU7050080

1,3,7,9-Tetra-Me: 1,3,7,9-Tetramethyluric acid. **Theacrine** [2309-49-1]

C₉H₁₂N₄O₃ 224.219

Isol. from *Camellia sinensis*, *Coffea* spp. and *Theobroma* spp. (tea, coffee and cocoa). Needles (H₂O). Spar. sol. Et₂O; sol. hot CHCl₃. Mp 229°. Distils without dec. λ_{\max} 211 (ϵ 11860); 231 (ϵ 6980); 292 (ϵ 9960) (MeOH).

[1198-77-2]

Johnson, T.B. *et al.*, *J.A.C.S.*, 1937, **59**, 1261-1264 (*Theacrine*)

Sutor, D.J. *et al.*, *Acta Cryst.*, 1963, **16**, 97-104 (*Theacrine, cryst struct*)

Lifschitz, C. *et al.*, *Isr. J. Chem.*, 1968, **6**, 827-831 (ms)


Kirk-Othmer *Encycl. Chem. Technol.*, 3rd edn., Wiley, 1978, **23**, 608 (rev, bibl)

Zheng, X.-Q. *et al.*, *Phytochemistry*, 2003, **60**, 129-134 (*Theacrine*)

Urocanic acid

U-46

3-(1H-Imidazol-4-yl)-2-propenoic acid, 9CI. 4-Imidazoleacrylic acid, 8CI. Urocanic acid [104-98-3]



(E)-form

C₆H₆N₂O₂ 138.126

Zwitterionic. Found in the stratum corneum of human skin. Sunscreen. Shows antineoplastic activity. Mp 243-245°.

► NI3425200

(E)-form [3465-72-3]

Degradn. prod. of histidine. Prod. by *Bacillus subtilis*, *Bacillus cereus*, *Micrococcus lysodeikticus*, *Achromobacter liquidum*, *Micrococcus lysodeikticus* and basidiomycete fungi *Coprinus atramentarius* (common ink cap), *Phallus impudicus* (common stinkhorn). Forms 0.5% of dry weight of human epidermis. Cryst. (dihydrate). Sol. Py, bases; fairly sol. H₂O, butanol, EtOH; poorly sol. EtOAc, hexane. Mp 225° (218-224°). λ_{\max} 260 (H₂O) (Berdy).

► LD₅₀ (mus, ivn) 200 - 400 mg/kg.*Hydrochloride*: Mp 189-191°.*Picrate*: Mp 218-220°.*Me ester*: [70346-51-9]C₇H₈N₂O₂ 152.152

Solid (EtOAc). Mp 100-101° (79-81°).

Also descr. as viscous liq.

N¹-(1,1-Dimethyl-3-oxobutyl): 3-[1-(1,1-Dimethyl-3-oxobutyl)imidazol-4-yl]-2-propenoic acid, 9CI [117082-14-1]

C₁₂H₁₆N₂O₃ 236.27

Isol. from rabbit skin tissue. Cryst. (EtOH). Mp 148-149° dec. Poss. artifact.

(Z)-form [7699-35-6]

Mp 175-176°.

Me ester: [88181-49-1]

Cryst. (toluene). Mp 100-101°.

[6198-55-6]

Aldrich Library of NMR Spectra, 2nd edn., 1983, **2**, 492A (nmr)*Aldrich Library of FT-IR Spectra*, 1st edn., 1985, **2**, 621A (ir)

Edlbacher, S. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1942, **276**, 126; 1943, **279**, 63 (isol, biosynth)

Biochem. Prep., 1955, **4**, 50 (synth)

Gregoire, J. *et al.*, *Bull. Soc. Chim. Belg.*, 1958, **40**, 767 (isol)

List, F.H. *et al.*, *Hoppe-Seyler's Z. Physiol. Chem.*, 1960, **319**, 17 (isol)Japan. Pat., 1961, 61 23 094; CA, **60**, 2302 (isol)

Baden, H. *et al.*, *Nature (London)*, 1966, **210**, 732 (pmr)

Shibatari, T. *et al.*, *Appl. Microbiol.*, 1974, **27**, 688 (biosynth)

Svinning, T. *et al.*, *Acta Cryst. B*, 1979, **35**, 2813 (*cryst struct*)

Quinn, R. *et al.*, *J.A.C.S.*, 1984, **106**, 4136

(synth, esters)

Ienaga, K. *et al.*, *J. Het. Chem.*, 1988, **25**, 1037 (*deriv*)

Shukla, M.K. *et al.*, *Spectrochim. Acta A*, 1995, **51**, 831 (uv, fluorescence, photochem)

Blake, A. J. *et al.*, *Acta Cryst. C*, 1997, **53**, 1093-1097 (*cryst struct, Me esters*)


Hanson, K.M. *et al.*, *J.A.C.S.*, 1997, **119**, 2715 (*spectra*)

Mohammad, T. *et al.*, *Org. Prep. Proced. Int.*, 2000, **32**, 581-584 (*Z-form, synth, ir, uv, pmr, cmr*)

Urochordamine A

U-47

[151756-66-0]

Relative
configurationC₂₂H₂₆BrN₇O 484.398

Alkaloid from the tunicates *Ciona savignyi* and *Botrylloides* sp. Promotes larvae settlement and metamorphosis in the tunicate. Exhibits antibacterial activity. $[\alpha]_D +11.7$ (c, 0.263 in CHCl₃). λ_{\max} 211 (ϵ 33100); 253 (ϵ 20000); 313 (ϵ 4170); 357 (ϵ 5500) (MeOH) (Derep).

9-Epimer: **Urochordamine B**

[151851-37-5]

C₂₂H₂₆BrN₇O 484.398


Alkaloid from *Ciona savignyi* and *Botrylloides* sp. Shows antibacterial activity. Metamorphosis promoter and larval settlement inhibitor. $[\alpha]_D -36.6$ (c, 0.174 in CHCl₃). λ_{\max} 211 (ϵ 33100); 253 (ϵ 20000); 313 (ϵ 4170); 357 (ϵ 5500) (MeOH) (Derep).

Tsukamoto, S. *et al.*, *Tet. Lett.*, 1993, **34**, 4819-4822 (isol, uv, ir, pmr, cmr, struct)

Urothione

U-48

2-Amino-7-(1,2-dihydroxyethyl)-6-(methylthio)thieno[3,2-g]pteridin-4(3H)-one, 9CI [17801-77-3]

C₁₁H₁₁N₅O₃S₂ 325.372

(R)-form

Orange-red pigment from human urine. Orange cryst. $[\alpha]_D -12$ (c, 0.24 in 0.05M NaOH).

Tri-Ac: [27498-07-3]

Yellow cryst. (Me₂CO/petrol). Mp 220°.

(±)-form [19295-31-9]

Yellow cryst. Mp 260-270° dec.


Koschara, W. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1943, **279**, 44 (isol, synth)
 Goto, M. et al., *Tet. Lett.*, 1967, 4507 (struct)
 Sakuri, A. et al., *Tet. Lett.*, 1968, 2941 (synth, struct, uv)
 Horibe, H. et al., *Tet. Lett.*, 1995, **36**, 2631 (abs config)

Ursuline† U-49

Struct. unknown. Alkaloid from the stem bark of *Oxandra cf. major* (Annonaceae).
 Arango, G.J. et al., *Phytochemistry*, 1987, **26**, 2093-2098 (isol, uv, ir, pmr)

Urukthapelstatin A U-50

[890085-24-2]




C₃₄H₃₀N₈O₆S₂ 710.793

Related to Mechercharstatin A, M-142 and Antibiotic YM 216391, A-1322. Isol. from a marine-derived *Mechercharimyces asporophorigenens* YM11-542. Cytotoxic. Powder. Mp 311° dec. [α]_D²² +38 (c, 0.5 in CHCl₃). λ_{max} 267 (log ε 4.35); 291 (log ε 4.52) (MeCN).

Matsuo, Y. et al., *J. Antibiot.*, 2007, **60**, 251-255; 256-260 (isol, pmr, cmr, cryst struct)

Usambanoline U-51

[174204-34-3]




C₂₁H₂₂NO₅[⊕] 368.408

Alkaloid from stems of *Zanthoxylum usambarensense*. Orange cryst. (MeOH/Et₂O)(as perchlorate). Mp 191° (perchlorate).

Kato, A. et al., *J. Nat. Prod.*, 1996, **59**, 316 (isol, ir, pmr, cmr, ms, struct)

Usambarensine U-52

[36150-14-8]



C₂₉H₂₈N₄ 432.567

Numbering systems vary. Alkaloid from the roots of *Strychnos usambarensis*, also from *Strychnos dale* and *Strychnos memecycloides* (Loganiaceae). Muscarinic receptor antagonist in isolated rat intestine. Shows antiparasitodal, antiamebic and antimitotic activities. Active against gram-positive bacteria. Powder. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 233 (ε 37153); 291 (ε 15850); 339 (ε 3390); 349 (ε 3390) (MeOH) (Berdy).

N^b-Oxide: **Usambarensine N^b-oxide**

C₂₉H₂₈N₄O 448.566

Alkaloid from *Strychnos dale*. Unpubl. results from 1996 review.

N^b-Me: **N^b-Methylusambarensine**

[36150-17-1]

C₃₀H₃₁N₄[⊕] 447.602

Quaternary alkaloid from *Strychnos usambarensis* (Loganiaceae).

3',4'-Dihydro: **Tchibangensine**. 3',4'-Dihydrousambarensine. 5',6'-Dihydrousambarensine

[36150-15-9]

C₂₉H₃₀N₄ 434.583

Major alkaloid from the root and stem bark of *Strychnos tchibangensis*. Also from *Strychnos dale* and *Strychnos usambarensis* (Loganiaceae). Mp 143-145°. [α]_D +109.7 (c, 0.7 in CHCl₃).

3',4'-Dihydro, N^b-oxide: 3',4'-Dihydrousambarensine N^b-oxide. 5',6'-Dihydrousambarensine N-oxide

C₂₉H₃₀N₄O 450.582

Alkaloid from *Strychnos dale*. Unpublished results from 1996 review.

3',4'-Dihydro, N^b-Me: **N^b-Methyl-3',4'-dihydrousambarensine**. N^b-Methyltchibangensine

[36150-16-0]

C₃₀H₃₃N₄[⊕] 449.617

Quaternary alkaloid from *Strychnos usambarensis* (Loganiaceae).

3',4'-Dihydro, N²-Me: **N²-Methyl-3',4'-dihydrousambarensine**. N⁴-Methyl-5',6'-dihydrousambarensine

C₃₀H₃₃N₄[⊕] 449.618

Quaternary alkaloid from *Strychnos usambarensis*.

19,20β-Dihydro: **19,20-Dihydrousambarensine**

[144548-37-8]

C₂₉H₃₀N₄ 434.583

Alkaloid from root bark of *Strychnos potatorum* (Loganiaceae). Also from *Strychnos tchibangensis* and *Strychnos usambarensis*. Active against gram-positive bacteria.

1'R,2',3',4'-Tetrahydro: **4',17-Dihydro-17β-tchibangensine**. Tetrahydro-17R-usambarensine

[73034-15-8]

C₂₉H₃₂N₄ 436.599

Isol. from the root bark and stem bark of *Strychnos ngouniensis* (Loganiaceae). Shows antimicrobial props. Active against gram-positive bacteria. [α]_D +40 (c, 0.5 in EtOH).

1'R,2',3',4'-Tetrahydro, N^b-oxide(α-):

4',5',6',17-Tetrahydro-17R-usambarensine N-oxide

[88607-57-2]

C₂₉H₃₂N₄O[⊕] 452.598

Alkaloid from *Aspidosperma marcgravianum*. Amorph. powder. [α]_D +15 (c, 0.51 in MeOH).

1'R,2',3',4'-Tetrahydro, N²-methoxy-carbonyl(β-): **N^b-Carbomethoxy-4',5',6',17-tetrahydro-17R-usambarensine**

[88607-55-0]

C₃₁H₃₄N₄O₂ 494.635

Alkaloid *Aspidosperma marcgravianum* (Apocynaceae). Amorph. [α]_D -7 (c, 0.62 in MeOH).

1'R,2',3',4'-Tetrahydro, N²-ethoxycarbonyl(β-): **N^b-Carboethoxy-4',5',6',17-tetrahydro-17R-usambarensine**

[88607-56-1]

C₃₂H₃₆N₄O₂ 508.662

Alkaloid from *Aspidosperma marcgravianum* (Apocynaceae). Amorph. powder. [α]_D -50 (CHCl₃). [α]_D -6 (c, 0.75 in MeOH).

1'S,2',3',4'-Tetrahydro: **4',17-Dihydro-17α-tchibangensine**. Tetrahydro-17S-usambarensine

[73034-12-5]

C₂₉H₃₂N₄ 436.599

Isol. from the stem bark of *Strychnos ngouniensis* and from *Strychnos dale* (Loganiaceae). Active against gram-positive bacteria. [α]_D -11 (c, 1 in EtOH).

6'-Hydroxy: **10'-Hydroxyusambarensine**

[225939-61-7]

C₂₉H₂₈N₄O 448.566

Alkaloid from *Strychnos usambarensis*. Amorph. powder. λ_{max} 227 (log ε 4.18); 253 (log ε 3.69); 291 (log ε 3.67); 298 (log ε 3.75); 362 (log ε 3.03) (MeOH).

6'-Hydroxy, 1'R,2',3',4'-tetrahydro: **10'-Hydroxy-4',17-dihydro-17β-tchibangensine**. 10'-Hydroxytetrahydro-17R-usambarensine

[88721-08-8]

C₂₉H₃₂N₄O 452.598

Isol. from the root bark and stem bark of *Strychnos ngouniensis* (Loganiaceae). Active against gram-positive bacteria. [α]_D +47 (c, 0.5 in EtOH).

6'-Hydroxy, 1'S,2',3',4'-tetrahydro: **10'-Hydroxy-4',17-dihydro-17α-tchibangensine**. 10'-Hydroxytetrahydro-17S-usambarensine

[88721-07-7]

C₂₉H₃₂N₄O 452.598

Isol. from the root bark and stem bark of *Strychnos ngouniensis* (Loganiaceae). Active against gram-positive bacteria. [α]_D 0 (c, 0.5 in EtOH).

10-Methoxy, 1'R,2',3',4'-tetrahydro: **Ramiflorine B**. 10-Methoxy-4',17-dihydro-17β-tchibangensine. 10-Methoxytetrahydrousambarensine

[173693-53-3]

C₃₀H₃₄N₄O 466.625

Alkaloid from bark of *Aspidosperma ramiflorum*. Mp 194°. [α]_D²⁵ +59.2 (c, 1.0 in EtOH).

10-Methoxy, 1'S,2',3',4'-tetrahydro: **Ramiflorine A**. 10-Methoxy-4',17-dihydro-


- 17 α -*tchibangensine*. 10-Methoxytetrahydro-17S-usambarensine
[173693-52-2]
C₃₀H₃₄N₄O 466.625
Alkaloid from bark of *Aspidosperma ramiflorum* (Apocynaceae). Mp 159°. [α]_D²⁵ +23.52 (c, 1.0 in EtOH).
- 6',10-Dihydroxy, 1'S,2',3',4'-tetrahydro: 10,10'-Dihydroxytetrahydrousambarensine
[118625-67-5]
C₂₉H₃₂N₄O₂ 468.597
Alkaloid from *Strychnos dale*.
- 6',10-Dihydroxy, 1'S,2',3',4'-tetrahydro, N²-Me: 10,10'-Dihydroxy-N⁴-methyltetrahydrousambarensine
[118625-69-7]
C₃₀H₃₄N₄O₂ 482.624
Alkaloid from *Strychnos dale* and *Strychnos ngouniensis*.
- 6'-Methoxy, 10-hydroxy, 1'S,2',3',4'-tetrahydro: 10-Hydroxy-10'-methoxytetrahydrousambarensine
[118625-66-4]
C₃₀H₃₄N₄O₂ 482.624
Alkaloid from *Strychnos dale* and *Strychnos ngouniensis*.
- 6'-Methoxy, 10-hydroxy, 1'S,2',3',4'-tetrahydro, N²-Me: 10-Hydroxy-10'-methoxy-N⁴-methyltetrahydrousambarensine
C₃₁H₃₆N₄O₂ 496.651
Alkaloid from *Strychnos ngouniensis* and *Strychnos dale*.
- 10-Methoxy, 6'-hydroxy, 1'S,2,3,4-tetrahydro, N²-Me: 1',2',3',4'-Tetrahydro-6'-hydroxy-10-methoxy-N²-methylusambarensine. 10'-Hydroxy-10-methoxy-N⁴-methyl-17,4',5',6'-tetrahydrousambarensine
C₃₁H₃₆N₄O₂ 496.651
Alkaloid from *Strychnos dale*. Unpubl. results from 1996 review.
- 6',10-Dimethoxy, 1'S,2',3',4'-tetrahydro: 10,10'-Dimethoxytetrahydrousambarensine
[101303-79-1]
C₃₁H₃₆N₄O₂ 496.651
Alkaloid from the leaves of *Strychnos dale*.
- 6',10-Dimethoxy, 1'S,2',3',4'-tetrahydro, N²-Me: 10,10'-Dimethoxy-N⁴-methyltetrahydrousambarensine
[101303-80-4]
C₃₂H₃₈N₄O₂ 510.678
Isol. from the leaves of *Strychnos dale* (Loganiaceae).
- Angenot, L. *et al.*, *J. Pharm. Belg.*, 1971, **26**, 585-588; *CA*, **76**, 72694w (*Usambarensine*, *Tchibangensine*, N^b-Methylusambarensine)
Dideberg, O. *et al.*, *Acta Cryst. B*, 1975, **31**, 1571-1575 (*cryst struct, Usambarensine*)
Yamada, K. *et al.*, *J.O.C.*, 1975, **40**, 2572-2573 (*synth, config, Tchibangensine*)
Richard, C. *et al.*, *Phytochemistry*, 1978, **17**, 539-541 (*struct, uv, ir, pmr, cmr, ms, Tchibangensine*)
Coune, C.A. *et al.*, *Phytochemistry*, 1980, **19**, 2009-2011 (*cmr*)
Robert, G.M.T. *et al.*, *J. Nat. Prod.*, 1983, **46**, 694-707 (*Tetrahydrousambarensine oxide, Carbalkoxytetrahydrousambarensines*)
Massiot, G. *et al.*, *Tetrahedron*, 1983, **39**, 3645-3656 (*derivs*)

- Verpoorte, R. *et al.*, *Tet. Lett.*, 1986, **27**, 239-242 (*Dimethoxytetrahydrousambarensine, Dimethoxy-N-methyltetrahydrousambarensine*)
Caron, C. *et al.*, *Planta Med.*, 1988, **54**, 409-412 (*derivs, activity*)
Massiot, G. *et al.*, *Phytochemistry*, 1992, **31**, 2873 (*19,20-Dihydrousambarensine*)
Wright, C.W. *et al.*, *Phytother. Res.*, 1994, **8**, 149-151 (*activity, isol*)
Bonjeau, K. *et al.*, *Anticancer Res.*, 1996, **16**, 1129-1138 (*activity*)
Marques, M.F.S. *et al.*, *Phytochemistry*, 1996, **41**, 963-967 (*Ramiflorines*)
Delaude, C. *et al.*, *Bull. Soc. R. Sci. Liege*, 1997, **66**, 183-286 (*occur, derivs*)
Frederich, M. *et al.*, *J. Nat. Prod.*, 1999, **62**, 619-621 (*10'-Hydroxyusambarensine, activity*)

Uscharin

[24211-81-2]

U-53



- C₃₁H₄₁NO₈S 587.733
Constit. of *Asclepias fruticosa* and *Calotropis procera*. Mp 270-271°. [α]_D +29.

► Cardiac poison. YV0995600

3',4'-Dihydro: Voruscharin

- [27892-03-1]
C₃₁H₄₃NO₈S 589.749
Constit. of the latex of *Calotropis procera*. Cryst. Mp 165-166°. [α]_D¹⁹ -60.8 (EtOH).

3',4'-Dihydro, 3'-oxo: 2''-Oxovorucharin

- [676541-57-4]
C₃₁H₄₁NO₉S 603.732
Constit. of *Calotropis procera* root bark. Numbering systems vary.
3',4'-Dihydro, 5,6-didehydro, 3'-oxo:
C₃₁H₃₉NO₉S 601.716
Constit. of *Asclepias tuberosa*. Mp 260-280° dec. [α]_D²⁹ -9.6 (c, 1.9 in MeOH).

19-Deoxo: 19-Deoxyuscharin

- [88972-32-1]
C₃₁H₄₃NO₇S 573.749
Constit. of *Asclepias fruticosa*. Pale yellow cryst. (CHCl₃). Mp 243.5-244.5°.

19-Deoxo, 15 β -hydroxy: 19-Deoxy-15 β -hydroxyuscharin

- [159336-96-6]
C₃₁H₄₃NO₈S 589.749
Constit. of *Asclepias fruticosa*. Amorph. powder. [α]_D²² -31.4 (c, 1.1 in MeOH).

16 α -Hydroxy: 16 α -Hydroxyuscharin

- [159337-02-7]
C₃₁H₄₁NO₉S 603.732
Constit. of *Asclepias fruticosa*. Amorph. powder. [α]_D²² -3.3 (c, 0.4 in MeOH).

5'-Epimer, 3', 4'-dihydro, 5,6-didehydro, 3'-oxo, S-oxide:

- C₃₁H₃₉NO₁₀S 617.716
Constit. of *Asclepias tuberosa*. Mp 240-245° dec. [α]_D²⁶ +21.1 (c, 1.1 in MeOH).

17-Epimer, 17-hydroxy: 17 β -Hydroxyuscharin


- [159337-00-5]
C₃₁H₄₁NO₉S 603.732
Constit. of *Asclepias fruticosa*. Amorph. powder. [α]_D²² +11.4 (c, 0.7 in MeOH).

- Bruschweiler, F. *et al.*, *Helv. Chim. Acta*, 1969, **52**, 2276 (*isol*)
Cheung, H.T.A. *et al.*, *J.C.S. Perkin I*, 1983, 2827-2835 (*struct, deriv*)
Warashina, T. *et al.*, *Phytochemistry*, 1994, **37**, 217-226 (*derivs*)
Abe, F. *et al.*, *Chem. Pharm. Bull.*, 2000, **48**, 991-993
Van Quaquebeke, E. *et al.*, *J. Med. Chem.*, 2005, **48**, 849-856 (*2''-Oxovorucharin*)

Usimine B

[1005771-14-1]

U-54



- C₂₃H₂₃NO₁₀ 473.435
Deriv. of Usnic acid. Constit. of *Ramalina terebrata*. Yellow gum. [α]_D²⁵ +159 (c, 0.46 in MeOH). Organism initially reported incorrectly as *Stereocaulon alpinum*. λ _{max} 216 (log ϵ 4.3); 297 (log ϵ 4.2) (MeOH).

5'-Me ester: Usimine A

- [1005771-13-0]
C₂₄H₂₅NO₁₀ 487.462
Constit. of *Ramalina terebrata*. Yellow gum. [α]_D²⁵ +39 (c, 0.77 in CH₂Cl₂). λ _{max} 216 (log ϵ 4); 297 (log ϵ 4.1) (MeOH).

Z-Isomer: Usimine C


- [1005771-15-2]
C₂₃H₂₃NO₁₀ 473.435
Constit. of *Ramalina terebrata*. Yellow gum. [α]_D²⁵ +162 (c, 0.43 in MeOH). λ _{max} 216 (log ϵ 4.3); 297 (log ϵ 4.2) (MeOH).

- Seo, C. *et al.*, *J. Nat. Prod.*, 2008, **71**, 710-712; 1322 (*isol, pmr, cmr*)

Uskudaramine

[83983-89-5]

U-55




- C₃₉H₄₄N₂O₈ 668.785
Alkaloid from *Thalictrum minus* var. *microphyllum* (Ranunculaceae). Amorph. [α]_D²⁵ +84 (c, 0.15 in MeOH). The first aporphine-benzylisoquinoline dimer bonded through C-C coupling.

Guinaudeau, H. *et al.*, *J.O.C.*, 1982, **47**, 5406
(*uv, cd, pmr, ms, struct*)

Ussuriedine

[125201-07-2]

U-56

C₂₇H₃₇NO₃ 423.594

Alkaloid from the Chinese herbal drug "ping-bei-mu" (dried bulbs of *Fritillaria ussuriensis*) (Liliaceae). Amorph. powder. Mp 189.5-192.5°. [α]_D +19 (c, 0.79 in CHCl₃).

O¹⁸-Me: Ussurienine

[116085-12-2]

C₂₈H₃₉NO₃ 437.621

Alkaloid artifact from dried bulbs of *Fritillaria ussuriensis* (Liliaceae). Needles (MeOH). Mp 300°. [α]_D +20 (c, 0.92 in CHCl₃). λ_{\max} 271 (ε 1280); 280 (ε 1280) (MeOH) (Derep).

6-Ketone: Ussuriedinone

[125201-08-3]

C₂₇H₃₅NO₃ 421.578

Alkaloid from dried bulbs of *Fritillaria ussuriensis* (Liliaceae). Needles (Me₂CO). Mp 268-272°. [α]_D +12 (c, 0.80 in CHCl₃).

6-Ketone, O¹⁸-Me: Ussurienone

[125941-96-0]


C₂₈H₃₇NO₃ 435.605

Alkaloid artifact from dried bulbs of *Fritillaria ussuriensis* (Liliaceae). Amorph. powder. Mp 110-116°. [α]_D +8 (c, 0.71 in CHCl₃).

Kitamura, Y. *et al.*, *Tetrahedron*, 1989, **45**, 5755-5766 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Ustalic acid

α, α' -(1,3-Dioxolane-4,5-diylidene)bis[benzeneacetic acid], 9CI. 3,4-(Methylenedioxy)-2,5-diphenyl-2,4-hexadienedioic acid
[470699-70-8]

C₁₉H₁₄O₆ 338.316

Isol. from *Tricholoma ustale*. ATPase inhibitor. Toxic principle. Cryst. Mp

200-202°.

2-Hydroxyethylamide: **Ustalic ethanolamide**

[470699-73-1]

C₂₁H₁₉NO₆ 381.384

Isol. from *Tricholoma ustale*. Cryst. Mp 128-131°.

Glycine amide: **Glycoustalic acid**. Ustalic glycine amide

[470699-72-0]

C₂₁H₁₇NO₇ 395.368

Isol. from *Tricholoma ustale*. Cryst. Mp 149-152°.

L-Proline amide: **Ustalic prolinamide**

[470699-71-9]

C₂₄H₂₁NO₇ 435.432

Isol. from *Tricholoma ustale*. Cryst. Mp 200-202°. [α]_D²⁰ -28 (c, 0.1 in MeOH).

Imide: see 4,8-Diphenyl-5H-1,3-dioxolo[4,5-d]azepine-5,7(6H)-dione, D-816

Sano, Y. *et al.*, *Chem. Comm.*, 2002, 1384-1385 (*isol, pmr, cmr, cryst struct*)


Sawayama, Y. *et al.*, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 2998-3003 (*synth*)

Hayakawa, I. *et al.*, *Tetrahedron*, 2008, **5873**, 5877 (*synth*)

Uvarindole A

[94977-29-4]

U-59



R = H

C₂₉H₂₅NO₃ 435.521

Alkaloid from the stem bark of *Uvaria angolensis* (Annonaceae). Needles (EtOAc/cyclohexane). Mp 66-68°.

1''-Oxo, N-de(hydroxybenzyl): **Uvarindole C**

[94977-31-8]

C₂₂H₁₇NO₃ 343.381

Alkaloid from the stem bark of *Uvaria angolensis* (Annonaceae). Needles (CHCl₃). Mp 151-153°.

Waterman, P.G. *et al.*, *Chem. Comm.*, 1984, 1280 (*uv, pmr, cmr, ms, struct*)

Uvarindole B

[94977-30-7]

As Uvarindole A, U-59 with

R = 2-Hydroxybenzyl

C₃₆H₃₁NO₄ 541.645

Alkaloid from the stem bark of

U-60


Uvaria angolensis (Annonaceae). Gummy solid.

Waterman, P.G. *et al.*, *Chem. Comm.*, 1984, 1280 (*pmr, cmr, struct*)

Uvarindole D

[94977-32-9]

U-61

C₂₉H₂₅NO₄ 451.521


Alkaloid from the stem bark of *Uvaria angolensis* (Annonaceae). Yellow plates (CHCl₃/MeOH). Mp 170-174°.

Waterman, P.G. *et al.*, *Chem. Comm.*, 1984, 1280 (*uv, ir, pmr, cmr, ms, struct*)

Uvarindole E

[113689-35-3]

U-62

C₃₆H₃₁NO₅ 557.645


Tentative struct. Alkaloid from stem bark of *Uvaria angolensis* (Annonaceae).

Muhammad, I. *et al.*, *J. Bangladesh Acad. Sci.*, 1987, **11**, 195; *CA*, **108**, 143090p

Uvariopsamine

3,4,7,8-Tetramethoxy-N,N-dimethyl-1-phenanthreneethanamine, 9CI. 1-(2-Dimethylaminoethyl)-3,4,7,8-tetramethoxyphenanthrene
[38764-73-7]

U-63

C₂₂H₂₇NO₄ 369.46Alkaloid from the bark of *Uvariopsis*

guineensis (Annonaceae). Yellow-orange needles (Me₂CO)(as picrate). Mp 181-182° (picrate).

N-Oxide: Uvariopsamine N-oxide

[38820-51-8]

C₂₂H₂₇NO₅ 385.459

Alkaloid from the bark of *Uvariopsis guineensis* (Annonaceae). Yellow-orange cryst. (Me₂CO)(as picrate). Mp 184-185° (picrate).

N-De-Me: Noruvariopsamine

[38764-74-8]


C₂₁H₂₅NO₄ 355.433

Alkaloid from the bark of *Uvariopsis guineensis* (Annonaceae). Mp 224-225° (picrate).

Leboeuf, M. *et al.*, *Phytochemistry*, 1972, **11**, 2833 (*isol, pmr, ms, struct*)

Uvariopsine

U-64
9-Methoxy-N,N-dimethylphenanthro[3,4-d]-1,3-dioxole-5-ethanamine, 9CI. 1-(2-Dimethylaminoethyl)-7-methoxy-3,4-methylenedioxyphenanthrene
[30147-99-0]



C₂₀H₂₁NO₃ 323.391

Aporphine-derived numbering shown. Alkaloid from the bark of *Uvariopsis guineensis* and from *Uvariopsis solheidii* (Annonaceae). Mp 95-96°. λ_{max} 216 (log ε 3.09); 236 (log ε 5); 258 (log ε 5); 284 (log ε 2.25); 322 (log ε 1.44); 353 (log ε 0.43) (MeOH).

8-Methoxy: 8-Methoxyuvariopsine

[38820-55-2]

C₂₁H₂₃NO₄ 353.417

Alkaloid from the bark of *Uvariopsis guineensis* (Annonaceae). Cryst. (hexane). Mp 99-100°.

Bouquet, A. *et al.*, *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1970, **271**, 1100-1102 (*uv, pmr, ms, struct*)

Leboeuf, M. *et al.*, *Phytochemistry*, 1972, **11**, 2833-2840 (*8-Methoxyuvariopsine*)

Lopez-Martin, J. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1613-1615 (*isol, pmr, cmr*)

Vakatidine

V-1

[1361-77-9]

C₂₂H₃₃NO₂ 343.508

Diterpenoid alkaloid. Struct. unknown. Alkaloid from the roots of *Aconitum palmatum* (Ranunculaceae). Mp 295-296°. *Hydrochloride*: Mp 304°. *Hydrobromide*: Mp 272-274°.

Singh, N. *et al.*, *J. Indian Chem. Soc.*, 1965, **42**, 49-50 (*uv, ir*)

Vakatisine

V-2

[1361-79-1]

[39389-34-9 (chloride)]

C₂₂H₃₄NO₂[⊕] 344.516

Struct. unknown. Quaternary alkaloid from the roots of *Aconitum palmatum* (Ranunculaceae). Mp 306° (as chloride). [α]_D²² +17.76 (EtOH) (chloride). CAS no. refers to chloride.

Singh, N. *et al.*, *J. Indian Chem. Soc.*, 1965, **42**, 49-50 (*uv, ir*)

Singh, N. *et al.*, *Indian J. Chem.*, 1972, **10**, 953-954 (*ir, pmr*)

Vakatisinine

V-3

[1361-78-0]

C₂₂H₃₃NO₄ 375.507

Diterpenoid alkaloid. Struct. unknown. Alkaloid from the roots of *Aconitum palmatum* (Ranunculaceae). Mp 308°.


Hydrochloride: Mp 303°.

Picolonate: Mp 302° dec.

Singh, N. *et al.*, *J. Indian Chem. Soc.*, 1965, **42**, 49-50 (*uv, ir*)

Valachine

V-4

C₂₅H₂₅NO₅ 419.476(±)-*form* [95307-59-8]


Alkaloid from *Berberis aldiviana* (Berberidaceae). Cryst. (EtOAc/MeOH/Et₂O). Mp 237-238°.

Firdous, S. *et al.*, *Chem. Comm.*, 1984, 1371 (*isol, uv, ir, pmr, ms, struct*)

Valanimycin

V-5

2-[(2-Methylpropyl)-ONN-azoxy]-2-propenoic acid, 9CI
[101961-60-8]

C₇H₁₂N₂O₃ 172.183

Prod. by *Streptomyces viridifaciens* (MG456.4F10; P8146). Active against some bacteria (gram-positive and -negative) and leukaemia and carcinoma. Oil; powder (as NH₃ adduct). Sol. EtOAc, MeOH, H₂O; poorly sol. CHCl₃, hexane. Mp 150.5° dec. (NH₃ adduct). pK_a 4.7. λ_{max} 230 (sh) (H₂O) (Derep). λ_{max} 230 (MeOH) (Berdy).

▶AT2140000


Yamato, M. *et al.*, *J. Antibiot.*, 1986, **39**, 184; 1263 (*isol, biosynth, props*)

Tao, T. *et al.*, *Org. Lett.*, 2003, **5**, 1213-1215 (*biosynth*)

Valindolmycin

V-6

[102141-01-5]


C₁₇H₂₅N₃O₂ 303.403

Indole antibiotic. Prod. by *Streptomyces kitasatoensis wenshanensis*. Active against gram-positive bacteria and anaerobes. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{max} 233 (ε 22900); 274 (ε 4670); 282 (ε 5000); 290 (ε 4350) (MeOH) (Berdy). Jin, W. *et al.*, *CA*, 1989, **111**, 20567d (*isol, struct*)

Valivine

V-7

20-(Hexahydro-1-methyl-1H-azepin-2-yl)pregn-5-ene-3,15,17-triol, 9CI
[85403-65-2]

C₂₈H₄₇NO₃ 445.684


Struct. dubious. Prob. a piperidine related to Sevcoridinine, S-275 or a solanidane. No further reports to 2006. Alkaloid from *Fritillaria valujevii* (Liliaceae). Cryst. (MeOH). Mp 256-258°. [α]_D -48 (c, 1 in CHCl₃/MeOH 1:1).

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 644-646; *Chem. Nat. Compd. (Engl. Transl.)*, 1983, **18**, 610-611 (*isol, ir, ms, pmr*)

Vallesamidine

V-8

[19637-77-5]



Absolute Configuration

C₂₀H₂₈N₂ 296.455

Alkaloid from *Vallesia dichotoma* (Apocynaceae). Noncryst. [α]_D -55 (CHCl₃). λ_{max} 206 (log ε 4); 257 (log ε 3.65); 301 (log ε 3.15) (CHCl₃).

Wälsler, A. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 391-404 (*isol, uv, ir, pmr, ms*)

Brown, S.H. *et al.*, *Ann. Chim. Farm.*, 1968, **90**, 2445 (*cryst struct, ms*)

Dickman, D.A. *et al.*, *J.A.C.S.*, 1989, **111**, 1528-1530 (*synth*)

Heathcock, C.H. *et al.*, *J.O.C.*, 1990, **55**, 798-811 (*synth, ir, pmr, cmr*)

Costa, P.R.R. *et al.*, *Tetrahedron: Asymmetry*, 1993, **4**, 1499-1500 (*synth*)


Tanino, H. *et al.*, *Tetrahedron*, 2004, **60**, 3273-3282 (*synth*)

Ho, T.-L. *et al.*, *Helv. Chim. Acta*, 2006, **89**, 249-257 (*synth*)

Vallesamine

V-9

Methyl 4-ethylidene-1,3,4,5,6,7-hexahydro-6-(hydroxymethyl)-2,5-ethano-2H-azocino[4,3-b]indole-6-carboxylate, 9CI
[3368-87-4]



Absolute Configuration

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from *Vallesia dichotoma*, *Hazunta modesta*, *Tabernaemontana amblyocarpa*, *Tabernaemontana dichotoma* and *Alstonia scholaris* (Apocynaceae). Mp 163-165°. [α]_D²⁷ +165 (CHCl₃). pK_a 8.2 (33% DMF aq.).

Methiodide: Mp 175-180° dec. [α]_D²⁸ +103 (EtOH).

N^b-Oxide: *Vallesamine N^b-oxide*

[126594-73-8]

C₂₀H₂₄N₂O₄ 356.421

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Amorph. solid. [α]_D²³ +76.1 (c, 1.14 in MeOH).

O-Ac: *O-Acetylvallesamine*

[56293-11-9]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the wood stem and stem bark of *Ervatamia heyneana* and from *Vallesia dichotoma* (Apocynaceae). Shows mod. cytotoxic activity vs. mouse P-388 carcinoma. Cryst. (Et₂O/hexane or C₆H₆/hexane). Mp 168-171°. [α]_D²⁵ +151 (CHCl₃). pK_a 7.85 (33% DMF aq.).

Z-Isomer: *Z-Vallesamine*

[107657-44-3]

C₂₀H₂₄N₂O₃ 340.421

Alkaloid from the leaves of *Alstonia scholaris* (Apocynaceae). Amorph. solid. [α]_D +182 (CHCl₃).

Wälsler, A. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 2072 (*isol, uv, ir, pmr, ms, struct*)

Scott, A.I. *et al.*, *Chem. Comm.*, 1978, 947 (*synth*)

Gunasekera, S.P. *et al.*, *Phytochemistry*, 1980, **19**, 1213 (*O-Acetylvallesamine*)


Perera, P. *et al.*, *Planta Med.*, 1984, **50**, 251 (*isol, uv, ir, pmr, cmr, ms*)

Atta-ur-Rahman, *et al.*, *Heterocycles*, 1987, **26**, 413 (*Z-Vallesamine*)

Yamauchi, T. *et al.*, *Phytochemistry*, 1990, **29**, 3547 (*N⁹-oxide*)

Vallesiachotamine

V-10

*(E)*-formC₂₁H₂₂N₂O₃ 350.416**(E)-form** [5523-37-5]

Alkaloid originally isol. from *Vallesia dichotoma* as a mixt. of geom. isomers and subsequently reported (probably also as a mixt. of isomers) from the roots of *Rhazya orientalis* and the leaves of *Rhazya stricta*. Also present in the seeds of *Strychnos tricalysioides* and from *Amsonia brevifolia*. Shows cytotoxic activity against mouse P-388 and human nasopharynx KB carcinoma cells. Clusters (CHCl₃/MeOH). Poorly sol. hexane. Mp 245-247°. [α]_D²⁵ +204 (c, 0.1 in CHCl₃). λ_{\max} 220 (ε 35500); 290 (ε 79400) (EtOH) (Berdy).

(Z)-form**Isovallesiachotamine**

[34384-71-9]

Alkaloid from the seeds of *Strychnos tricalysioides* and *Rhazya stricta* legumes (without seeds) (Loganiaceae, Apocynaceae). Prisms (CHCl₃/MeOH). Mp 235-237°. [α]_D²⁵ -54 (c, 0.06 in CHCl₃).

Djerassi, C. *et al.*, *J.A.C.S.*, 1966, **88**, 1792
Evans, D.A. *et al.*, *Phytochemistry*, 1968, **7**, 1429 (*isol*)

Waterman, P.G. *et al.*, *Planta Med.*, 1982, **45**, 28-30 (*isol, uv, ir, pmr, ms*)

Spitzner, D. *et al.*, *Angew. Chem., Int. Ed.*, 1984, **23**, 984 (*synth*)

Atta-ur-Rahman, *et al.*, *J. Nat. Prod.*, 1984, **47**, 388 (*isol, uv, ir, ms*)


Spitzner, D. *et al.*, *J.O.C.*, 1988, **53**, 2274 (*synth*)

Amann, R. *et al.*, *Annalen*, 1996, 349 (*synth*)
Vencato, I. *et al.*, *Acta Cryst. E*, 2006, **62**, 429-431 (*cryst struct*)

Vallesiachotamine lactone

V-11

[191288-40-1]

C₂₁H₂₀N₂O₄ 364.4

Alkaloid from aerial parts of *Cephaelis dichroa* (Rubiaceae).

Solis, P.N. *et al.*, *Phytochemistry*, 1993, **33**,

1117-1119 (*isol, uv, ir, pmr, ms, struct*)
Engler, A. *et al.*, *Nat. Prod. Lett.*, 1997, **9**, 225-228 (*synth*)

Valloporphine

V-12

[1361-83-7]

C₁₈H₂₃NO₅ 333.383

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Vallota purpurea* (preferred genus name *Cyrtanthus*) (Amaryllidaceae). Mp 244-247° dec. [α]_D -90 (c, 0.2 in EtOH).

Picrate:

Fine needles (Me₂CO). Mp 193° dec.

Döpke, W. *et al.*, *Naturwissenschaften*, 1965, **52**, 347 (*isol, uv, ir*)

Vallotidine

V-13

C₁₈H₂₁NO₅ 331.368

Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Vallota purpurea* (preferred genus name *Cyrtanthus*) (Amaryllidaceae). Prisms (Me₂CO). Mp 172-173°. [α]_D²² +57 (c, 0.25 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129-1134 (*isol*)

Vallotine

V-14

C₁₇H₁₉NO₅ 317.341


Amaryllidaceae alkaloid. Struct. unknown. Isol. from *Vallota purpurea* (preferred genus name *Cyrtanthus*) (Amaryllidaceae). Prisms (Me₂CO). Mp 217-218° dec. [α]_D²² -53 (c, 0.2 in CHCl₃).

Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 1129-1134 (*isol*)

Valoneic acid dilactone

V-15

Valoneic dilactone. Valonaeic dilactone
[60202-70-2]

C₂₁H₁₀O₁₃ 470.302

Ellagitannin. Constit. of *Elaeagnus umbellata*, *Epilobium hirsutum*, *Lagerstroemia* sp. and *Lythrum salicaria*. Xanthine oxidase inhibitor.

Amide: Epilobamide A. Valoneic acid amide dilactone

[194298-95-8]

C₂₁H₁₁NO₁₂ 469.317


Constit. of the whole plant of *Epilobium hirsutum*.

Schmidt, O.T. *et al.*, *Annalen*, 1955, **591**, 156 (*struct*)

Nawwar, M.A.M. *et al.*, *J. Mass Spectrom.*, 1997, **32**, 645-654 (*Epilobamide A*)

Valparicine

V-16

C₁₉H₂₀N₂ 276.38


Alkaloid from the stem bark of *Kopsia arborea*. Oil. [α]_D -40 (c, 0.22 in CHCl₃). λ_{\max} 228 ; 297 (EtOH).

Lim, K.-H. *et al.*, *Tet. Lett.*, 2006, **47**, 5037-5039 (*isol, pmr, cmr*)

Vancoresmycin

V-17

[268728-82-1]

C₇₁H₁₂₆N₂O₂₁ 1343.777

Prod. by *Amycolatopsis* sp. ST 101170.


Powder. λ_{\max} 234 (log ε 4.39); 280 (log ε 4.29) (MeOH).

Pat. Coop. Treaty (WIPO), 2000, 00 28 064; *CA*, **132**, 346708q

Hopmann, C. *et al.*, *Tet. Lett.*, 2002, **43**, 435-438 (*isol, pmr, cmr*)

Vanillobiocin

V-18

C₃₁H₃₁ClN₂O₁₂ 659.045

Coumermycin-type antibiotic. Closely related to Chlorobiocin, C-395 and Novobiocin. Prod. by a mutant of *Streptomyces roseochromogenes*.

Dechloro: DeclovannoillobiocinC₃₁H₃₂N₂O₁₂ 624.6

Prod. by a mutant of *Streptomyces roseochromogenes*.

2''-(5-Methyl-2-pyrrolicarbonyl) isomer: IsovanillobiocinC₃₁H₃₁ClN₂O₁₂ 659.045


Prod. by a mutant of *Streptomyces roseochromogenes*.

Freitag, A. *et al.*, *J. Antibiot.*, 2004, **57**, 205-209 (*isol, pmr*)


Vanuatine

V-19

[87183-74-2]

C₃₉H₄₆N₂O₈ 670.801Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). [α]_D²⁵ +138 (c, 0.12 in MeOH).*Di-Me ether*: [α]_D²⁵ +78 (c, 0.12 in MeOH).Bruneton, J. et al., *J.O.C.*, 1983, **48**, 3957 (*isol, uv, cd, pmr, ms, struct*)**Varacin A**

V-20

6,7-Dimethoxy-4-benzotrithioleethanamine, 9CI
[162857-71-8]C₁₀H₁₃NO₂S₃ 275.416Alkaloid from the Far Eastern ascidian *Polycitor* sp.**1-S-Oxide: Varacin B**

[162901-84-0]

C₁₀H₁₃NO₃S₃ 291.416From *Polycitor* sp.**3-S-Oxide: Varacin C**

[162857-72-9]

C₁₀H₁₃NO₃S₃ 291.416From *Polycitor* sp.**Ac:**

Exhibits potent antifungal and antimicrobial activity. Thin yellow cryst. (MeOH). Mp 140-142°.

Ac, 1-S-oxide:

Exhibits potent antifungal and antimicrobial activity. Thin yellow cryst. (MeOH). Mp 131-133°.


Ac, 3-S-oxide:

Exhibits potent antifungal and antimicrobial activity. Thin yellow cryst. (MeOH). Mp 70-73°.

5-(Methylthio): 6-(2-Aminoethyl)-3,4-dimethoxy-5-(methylthio)benzotrithiane.**5-(Methylthio)varacin A**C₁₁H₁₅NO₂S₄ 321.509Isol. from a *Lissoclinum* sp. Protein kinase C inhibitor. Not in CA. λ_{\max} 211 (€ 34800); 240 (€ 21000); 271 (€ 14700) (MeOH) (Berdy).**5-(Methylthio), N,N-di-Me: 6,7-Dimethoxy-N,N-dimethyl-5-(methylthio)-4-benzotrithioleethanamine. 6-(2-Aminoethyl)-3,4-dimethoxybenzotrithiane**C₁₃H₁₉NO₂S₄ 349.563Isol. from *Lissoclinum japonicum*. Protein kinase C inhibitor. Pale yellow oil (as trifluoroacetate). λ_{\max} 209 (€ 15900); 238 (€ 7410); 277 (€ 5960) (MeOH).Campagnone, R.S. et al., *Tetrahedron*, 1994,**50**, 12785-12792 (*methylthio derivs*)Makarjeva, T.N. et al., *J. Nat. Prod.*, 1995, **58**, 254 (*isol, ir, pmr, ms, struct*)Lee, A.H.F. et al., *J.A.C.S.*, 2002, **124**, 13972-13973 (*Varacin C, synth*)**Varadine**

V-21

[71639-34-4]




Relative Configuration

C₁₈H₁₉NO₅ 329.352Unique subtype of Amarylidiaceae alkaloid. Alkaloid from *Hymenocallis arenicola* (Amarylidiaceae). Prisms (Me₂CO). Mp 95°.Döpke, W. et al., *Z. Chem.*, 1979, **19**, 215-216**Varamine A**

V-22

[122093-14-5]

R = CH₂CH₃C₂₂H₂₃N₃O₂S 393.509Alkaloid from the tunicate *Lissoclinum vareau*. Cytotoxic agent (about 1 order of magnitude more toxic than the cystodytins which lack the thiomethyl group). Topoisomerase inhibitor. Orange solid. Sol. MeOH, CHCl₃. λ_{\max} 224 (€ 24400); 228 (sh); 240 (sh); 263 (sh); 280 (sh); 291 (€ 31100); 339 (€ 20400); 359 (€ 4520); 376 (€ 4640); 527 (€ 5670); 552 (sh) (pH 2) (Derep). λ_{\max} 232 (€ 31500); 275 (€ 25800); 292 (sh); 324 (sh); 382 (€ 3530); 464 (€ 5170); 494 (sh) (MeOH) (Derep).

► Toxic.

Molinski, T.F. et al., *J.O.C.*, 1989, **54**, 4256-4259 (*isol, uv, ir, pmr, cmr, ms, struct*)**Varamine B**

V-23

[122093-16-7]

As Varamine A, V-22 with


R = CH₃C₂₁H₂₁N₃O₂S 379.482Alkaloid from the tunicate *Lissoclinum vareau*. Cytotoxic agent. Topoisomerase inhibitor. Orange solid. Sol. MeOH, CHCl₃. λ_{\max} 224 (€ 24400); 228 (sh); 240 (sh); 263 (sh); 280 (sh); 291 (€ 31100); 339 (€ 20400); 359 (€ 4520); 376 (€ 4640); 527 (€ 5670); 552 (sh) (pH 2) (Derep). λ_{\max} 232 (€ 31500); 275 (€ 25800); 292 (sh); 324 (sh); 382 (€ 3530); 464 (€ 5170); 494 (sh) (MeOH) (Derep). λ_{\max} 234 (€ 25900); 274 (€ 21400); 382 (€ 3040); 462 (€ 4320) (MeOH/HCl) (Berdy). λ_{\max} 223 (€ 20570); 294 (€ 27400); 311 (€ 17300); 379 (€ 4190); 529 (€ 4720) (MeOH/HCl) (Berdy).

► Toxic.

Molinski, T.F. et al., *J.O.C.*, 1989, **54**, 4256-4259 (*isol, uv, ir, pmr, cmr, struct*)**Variabiline†**

V-24

10-[Bis(phenylmethyl)amino]-5,6,6a,7-tetrahydro-2-methoxy-6-methyl-4H-dibenzo[de,g]quinolin-1-ol, 9CI. 10-(Dibenzylamino)-1-hydroxy-2-methoxyaporphine [40374-54-7]

C₃₂H₃₂N₂O₂ 476.617**(±)-form**Alkaloid from *Ocotea variabilis* (Lauraceae). Mp 116-117°.*Hydrochloride*: Mp 230-232° dec.*Hydrochloride (1:2)*: Mp 185-187° dec.Cava, M.P. et al., *Tet. Lett.*, 1972, 4647 (*uv, ir, pmr, ms, synth, struct*)**Variceramides**

V-25

H₃C(CH₂)_nCOCH(OH)CH(-CH₂OH)NHCO(CH₂)₂₀CH₃
Isol. from the marine sponge *Ircinia variabilis*.**Variceramide 1** [128733-00-6]C₄₁H₈₁NO₄ 652.095

Has n=14.

Variceramide 2 [128733-01-7]C₄₀H₇₉NO₄ 638.068

Has n=13.


Variceramide 3 [128733-02-8]C₃₉H₇₇NO₄ 624.042

Has n=12.

Cafieri, F. et al., *Annalen*, 1990, 1141-1142 (*isol, pmr, ms*)**Variicolorin G**

V-26

[957780-76-6]

C₂₄H₂₉N₃O₂ 391.512

Isol. from *Aspergillus variegator* B-17. Amorph. powder. $[\alpha]_D^{25}$ -16 (c, 0.1 in MeOH). λ_{\max} 201 (log ϵ 3.5); 230 (log ϵ 3.6); 281 (log ϵ 3.1); 338 (log ϵ 3.1) (MeOH).


2'',3''-Dihydro, 3''-chloro, 2''R-hydroxy: **Variolorin F** [957780-73-3]

$C_{24}H_{30}ClN_3O_3$ 443.972

Isol. from *Aspergillus variegator* B-17. Amorph. powder. $[\alpha]_D^{25}$ -28 (c, 0.1 in MeOH). λ_{\max} 197 (log ϵ 3.7); 228 (log ϵ 3.8); 285 (log ϵ 3.2); 337 (log ϵ 3.3) (MeOH).

Wang, W.-L. et al., *J. Nat. Prod.*, 2007, **70**, 1558-1564 (isol, cd, pmr, cmr)

Variolorin J V-27
[957780-83-5]




$C_{23}H_{25}N_3O_3$ 391.469

Regioisomer of Neoechinulin. Isol. from *Aspergillus variegator* B-17. Red powder. λ_{\max} 208 (log ϵ 3.7); 232 (log ϵ 3.7); 285 (log ϵ 3.2); 420 (log ϵ 3.1) (MeOH).

Wang, W.-L. et al., *J. Nat. Prod.*, 2007, **70**, 1558-1564 (isol, pmr, cmr)

Variolorin K V-28
[957780-85-7]




$C_{27}H_{35}N_3O_4$ 465.591

Isol. from *Aspergillus variegator* B-17. Amorph. powder. $[\alpha]_D^{25}$ -49 (c, 0.1 in MeOH). λ_{\max} 210 (log ϵ 3.8); 228 (log ϵ 3.8); 275 (log ϵ 3.3); 340 (log ϵ 3.2) (MeOH).

Wang, W.-L. et al., *J. Nat. Prod.*, 2007, **70**, 1558-1564 (isol, cd, pmr, cmr)

Variolorin L V-29
[957780-87-9]




$C_{29}H_{39}N_3O_2$ 461.646

Regioisomer of Echinulin. Isol. from *Aspergillus variegator* B-17. Amorph. powder. $[\alpha]_D^{25}$ -21 (c, 0.05 in $CHCl_3$). λ_{\max} 197 (log ϵ 3.5); 235 (log ϵ 3.5); 293 (log ϵ 3.5)

3.1) (MeOH).

Wang, W.-L. et al., *J. Nat. Prod.*, 2007, **70**, 1558-1564 (isol, cd, pmr, cmr)

Variocolortide A V-30



R = $-CH_2CH=C(CH_3)_2$

$C_{39}H_{35}N_3O_7$ 657.721

Prod. by *Aspergillus variegator* B-17. Weakly antioxidant and cytotoxic. Yellow cryst. (MeOH). Mp 182° dec. $[\alpha]_D^{25}$ +5.5 (c, 0.1 in $CHCl_3$). λ_{\max} 229 (log ϵ 4.1); 265 (log ϵ 3.5); 291 (log ϵ 3.6); 368 (log ϵ 3.4) (MeOH).

Wang, W.-L. et al., *Chem. Biodiversity*, 2007, **4**, 2913-2919 (isol, pmr, cmr, cryst struct)

Variocolortide B V-31

As Variocolortide A, V-30 with R = H

$C_{34}H_{27}N_3O_7$ 589.603

Prod. by *Aspergillus variegator* B-17. Weakly antioxidant and cytotoxic. Amorph. yellow powder. $[\alpha]_D^{25}$ +4.5 (c, 0.1 in $CHCl_3$). λ_{\max} 230 (log ϵ 4); 264 (log ϵ 3.4); 290 (log ϵ 3.5); 359 (log ϵ 3.4) (MeOH).


8-Me ether: **Variocolortide C**

$C_{35}H_{29}N_3O_7$ 603.63

Prod. by *Aspergillus variegator* B-17. Amorph. yellow powder. $[\alpha]_D^{25}$ +4 (c, 0.1 in $CHCl_3$). λ_{\max} 228 (log ϵ 4.1); 266 (log ϵ 3.4); 289 (log ϵ 3.5); 367 (log ϵ 3.3) (MeOH).

Wang, W.-L. et al., *Chem. Biodiversity*, 2007, **4**, 2913-2919 (isol, pmr, cmr)

Variolin B V-32
[156790-85-1]



$C_{14}H_{11}N_7O$ 293.287

Alkaloid from the Antarctic sponge *Kirkpatrickia varialosa*. Cyclin-dependent kinase inhibitor. Antineoplastic agent. Yellow prisms (TFA aq.). Fairly sol. MeOH, DMSO; poorly sol. H_2O . Mp 45° dec. λ_{\max} 221 (log ϵ 14100); 324 (log ϵ 3900); 404 (log ϵ 9100); 422 (log ϵ 8400) (MeOH) (Berdy). λ_{\max} 213 (log ϵ 15500); 311 (log ϵ 4400);

394 (log ϵ 6300) (MeOH-HCl) (Berdy).

3',4',5',6'-Tetrahydro, N^{3'}-Me: N^{3'}-**Methyltetrahydrovariolin B** [155205-63-3]

$C_{15}H_{17}N_7O$ 311.346

From *Kirkpatrickia varialosa*. Shows antifungal activity and moderate cytotoxicity. Light yellow solid. Fairly sol. MeOH, DMSO. Mp 226° dec. $[\alpha]_D$ -22.4 (c, 3.5 in MeOH). λ_{\max} 256 (log ϵ 9680); 311 (log ϵ 8660); 324 (log ϵ 9885); 351 (log ϵ 6995); 367 (log ϵ 6840); 385 (log ϵ 3190) (MeOH) (Berdy).

5'-Hydroxy, 1'-Me (zwitterion): **Variolin A**

[155416-43-6]

$C_{15}H_{13}N_7O_2$ 323.313

Alkaloid from *Kirkpatrickia varialosa*. Weakly cytotoxic. Red solid. Fairly sol. MeOH, DMSO; poorly sol. H_2O . Mp 196° dec. Unstable in base. λ_{\max} 254 (log ϵ 8945); 315 (log ϵ 5375); 469 (log ϵ 14570) (MeOH) (Berdy).

Perry, N.B. et al., *Tetrahedron*, 1994, **50**, 3987-3982 (isol, uv, ir, pmr, cmr, cryst struct)

Trimurtulu, G. et al., *Tetrahedron*, 1994, **50**, 3993-4000 (N^{3'}-Methyltetrahydrovariolin B, Variolin A)

Molina, P. et al., *Tet. Lett.*, 2002, **43**, 1005-1007 (synth)

Ahaidar, A. et al., *J.O.C.*, 2003, **68**, 10020-10029 (synth)


Simone, M. et al., *Eur. J. Cancer*, 2005, **41**, 2366-2377 (pharmacol)

Anderson, R.J. et al., *J.O.C.*, 2005, **70**, 6204-6212 (synth)

Fresneda, P.M. et al., *J. Med. Chem.*, 2006, **49**, 1217-1221 (pharmacol)

Baeza, A. et al., *Tet. Lett.*, 2008, **49**, 4073-4077 (synth)

Variolin D V-33
[156790-86-2]




$C_{12}H_{10}N_4O_3$ 258.236

Possibly an artifact. Alkaloid from the Antarctic sponge *Kirkpatrickia varialosa*. Fairly sol. MeOH, DMSO. Mp 248°. λ_{\max} 226 (log ϵ 11700); 240 (log ϵ 14700); 246 (log ϵ 14800); 368 (log ϵ 12000); 388 (log ϵ 9700) (MeOH) (Berdy).

Perry, N.B. et al., *Tetrahedron*, 1994, **50**, 3987 (isol, uv, ir, pmr, cmr, struct)

Variotin, JAN V-34

1-(8-Hydroxy-6-methyl-1-oxo-2,4,6-dodecatrienyl)-2-pyrrolidinone, 9Cl. 1-(8-Hydroxy-6-methyl-2,4,6-dodecatrienyl)-2-pyrrolidinone, 8Cl. **Pecilocin, BAN, INN**. *Leofungine. Supral* [19504-77-9]



C₁₇H₂₅NO₃ 291.389


Isol. from *Paecilomyces variotii* var. *antibioticus*. Antifungal antibiotic. Hygroscopic needles + 1H₂O (Et₂O/petrol). Sol. MeOH, CS₂, dioxan, CCl₄, Et₂O. Mp 41.5-42.5°. [α]_D²⁸ -5.68 (MeOH). Log P 1.73 (uncertain value) (calc). Unstable, v. air-sensitive. λ_{max} 320 (ε 46000) (MeOH) (Derep). λ_{max} 321 (E1%/1cm 1198) (MeOH) (Berdy).

- LD₅₀ (mus, ipr) 320 mg/kg. UY5777000
Takeuchi, S. *et al.*, *J. Antibiot.*, 1969, **22**, 179 (pmr, struct)
Sakakibara, M. *et al.*, *Agric. Biol. Chem.*, 1973, **37**, 911 (ir, uv, ms, nmr, synth)
Ishida, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1978, **51**, 2077 (synth, ir, uv, ms, nmr)
Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 330

Vasconine

V-35

4,5-Dihydro-9,10-dimethoxyppyrolo[3,2,1-de]phenanthridinium(1+), 9CI [139955-90-1]

C₁₇H₁₆NO₂[⊕] 266.319

Alkaloid from whole plants of *Narcissus vasconicus* (Amaryllidaceae). Mp 233-235°. Counterion not specified.

O⁸-De-Me: **8-O-Demethylvasconine**

[170129-28-9]

C₁₆H₁₄NO₂[⊕] 252.292

Alkaloid from bulbs of *Crinum kirkii* (Amaryllidaceae). Mp 200-202°. Counterion not specified.

2-Methoxy: **Tortuosine**

[109644-38-4]

C₁₈H₁₈NO₃[⊕] 296.345

Alkaloid from whole plants of *Narcissus tortuosus* (Amaryllidaceae). Mp 242-243°. Counterion not specified.

2-Methoxy, O⁹-de-Me: **4,5-Dihydro-10-hydroxy-2,9-dimethoxyppyrolo[3,2,1-de]phenanthridinium(1+)**. **9-O-Demethyltortuosine**

[353246-57-8]

C₁₇H₁₆NO₃[⊕] 282.318

Alkaloid from *Zephyranthes carinata*.

Bastida, J. *et al.*, *J. Nat. Prod.*, 1992, **55**, 122 (isol, ir, pmr, cmr, ms, struct)

Parnes, J.S. *et al.*, *J.O.C.*, 1994, **59**, 3497 (synth)

Bastida, J. *et al.*, *Phytochemistry*, 1995, **38**, 549; **40**, 1291 (*Tortuosine*, 8-O-Demethylvasconine)

Rosa, A.M. *et al.*, *Tetrahedron*, 1997, **53**, 299 (synth)

Stark, L.M. *et al.*, *J.O.C.*, 2000, **65**, 3227-3230 (*Tortuosine*, synth)

Matsuga, M. *et al.*, *Nat. Med. (Tokyo)*, 2001, **55**, 201-204 (9-O-Demethyltortuosine)

Vasicinone

V-36

Struct. unknown. Prob. quinazoline alkaloid. Alkaloid from the young inflorescences of *Adhatoda vasica* (Acanthaceae). Bronchodilator, expectorant. Mp 208°. Ir 1492, 1590 and 1628

cm⁻¹.

Hydrochloride: Mp 258-260°.

Picrate: Mp 207-210°.


Ac: Mp 239-240°.

Ikram, M. *et al.*, *Pak. J. Sci. Ind. Res.*, 1965, **8**, 76; *CA*, **64**, 9721d (isol, ir)

Vasicinol

V-37

1,2,3,9-Tetrahydropyrrolo[2,1-b]quinazoline-3,7-diol, 9CI. 7-Hydroxypeganine. 7-Hydroxyvasicine



(R)-form

C₁₁H₁₂N₂O₂ 204.228

Abs. config. revised in 1996.

(R)-form

Alkaloid from *Linaria vulgaris*. Needles (MeOH). Mp 260° dec. [α]_D¹⁸ -35.5 (c, 0.01 in MeOH). λ_{max} 204 ; 292 (MeOH).

(S)-form [5081-51-6]

Alkaloid from the roots, leaves and seeds of *Adhatoda vasica*, the roots of *Sida cordifolia* and from other *Sida* spp. (Acanthaceae, Malvaceae). Transient hypotensive agent, cardiac depressant. Histamine antagonist, shows mild anticholinesterase activity. Mp 272-273° (260° dec.). [α]_D²⁵ +2.5 (c, 0.32 in AcOH). [α]_D +51 (c, 0.5 in 5% AcOH). The low specific rotn. recorded by Ghosal *et al* suggests that their compd. was largely racemic.

(±)-form

Synthetic. Mp 270-273°.

O⁷-Me:

Cryst. (H₂O, EtOAc or EtOH). Mp 216-218°.

Späth, E. *et al.*, *Ber.*, 1936, **69**, 384 (isol)

Southwick, P.L. *et al.*, *J.A.C.S.*, 1958, **80**, 1168 (deriv, synth)

Späth, E. *et al.*, *Monatsh. Chem.*, 1960, **91**, 1150 (isol)

Kuffner, F. *et al.*, *Monatsh. Chem.*, 1960, **91**, 1152 (synth)

Bhatnagar, A.K. *et al.*, *Indian J. Chem.*, 1966, **4**, 291 (ms)

Ghosal, S. *et al.*, *Phytochemistry*, 1975, **14**, 830 (isol)

Johns, S. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 129 (rev, pharmacol)


Joshi, B.S. *et al.*, *Tetrahedron: Asymmetry*, 1996, **7**, 25 (abs config)

Hua, H. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 1393-1394 (isol, pmr, cmr)

Vasicinone

V-38

2,3-Dihydro-3-hydroxyppyrolo[2,1-b]quinazolin-9(1H)-one, 9CI

C₁₁H₁₀N₂O₂ 202.212

The enantiomer originally isol. was the (-)-form which was originally assigned R config. Abs. stereochem. reversed to S- in 1996. The R-enantiomer has been isolated once.

(R)-form [119364-42-0]

Alkaloid from *Adhatoda vasica*. Cryst. (CHCl₃/EtOAc). Mp 201-202°. [α]_D +148 (EtOH).

(S)-form [486-64-6]

Alkaloid from *Biebersteimia multifida*, the seeds and above-ground parts of *Peganum harmala*, the foliage of *Peganum nigellastrum*, the roots of *Sida cordifolia* and the leaves of *Adhatoda vasica*. Also isol. from *Nitraria sibirica* (Biebersteiniaceae, Zygophyllaceae, Malvaceae, Acanthaceae). Shows bronchodilator and weak hypotensive inotropic action. Cryst. (EtOH). Mp 200-201°. [α]_D -129 (CHCl₃). [α]_D -100 (CHCl₃). [α]_D -78 (CHCl₃). These are probably partial racemates.

Hydrochloride: Mp 232-234° dec.

Hydrobromide: Mp 254-255° dec.

Hydroiodide: Mp 222-226° dec.

N⁴-Oxide: **Vasicinone N-oxide**

[168781-19-9]

C₁₁H₁₀N₂O₃ 218.212

Alkaloid from *Nitraria komarovii*.

Cryst. (Me₂CO). Mp 203-204°.

5-Methoxy: **5-Methoxyvasicinone**

[178740-33-5]

C₁₂H₁₂N₂O₃ 232.238

Alkaloid from leaves of *Adhatoda vasica*. Cryst. (MeOH). Mp 220-222°.

7-Hydroxy: **Vasicinolone**

[84847-50-7]

C₁₁H₁₀N₂O₃ 218.212

Alkaloid from the roots of *Adhatoda vasica* (Acanthaceae). Mp 279°.

7-Methoxy: **7-Methoxyvasicinone**

[178991-89-4]

[178631-31-7, 178991-88-3]

C₁₂H₁₂N₂O₃ 232.238

Alkaloid from leaves of *Adhatoda vasica*. Mp 199-202°.

(±)-form [35387-16-7]

Alkaloid from the inflorescence of *Adhatoda vasica* (Acanthaceae) and from aerial parts of *Galium aparine* collected at flowering (Rubiaceae). Cryst. (MeOH). Mp 209-211° dec. Also obt. by racemisation of (-)-Vasicinone.

5-Methoxy: **Adhavasinsonone**

[107783-56-2]

C₁₂H₁₂N₂O₃ 232.238

Alkaloid from the leaves of *Adhatoda vasica* (Acanthaceae). Mp 230-232°. Opt. inactive.

Morris, R.C. *et al.*, *J.A.C.S.*, 1935, **57**, 951 (synth)

Mehta, D.R. *et al.*, *J.O.C.*, 1963, **28**, 445 (isol, uv, ir)

Onaka, T. *et al.*, *Tet. Lett.*, 1971, 4387 (synth)

Ghosal, S. *et al.*, *Phytochemistry*, 1975, **14**, 830 (isol)


Szulzewsky, K. *et al.*, *J. Prakt. Chem.*, 1976, **318**, 463 (abs config)

Johns, S. *et al.*, *J. Prakt. Chem.*, 1977, **319**, 919 (cmr)

- Rashkes, Y.V. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 378; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 320 (ms)
- Choudhury, M.K. *et al.*, *Naturwissenschaften*, 1979, **66**, 205 (*isol, uv, synth*)
- Jain, M.P. *et al.*, *Planta Med.*, 1982, **46**, 250 (*Vasicolinone*)
- Johne, S. *et al.*, *Alkaloids (Academic Press)*, 1986, **29**, 129 (*rev, pharmacol*)
- Chowdhury, B.K. *et al.*, *Chem. Ind. (London)*, 1987, 35 (*Adhavasinsonone*)
- Poi, R. *et al.*, *J. Indian Chem. Soc.*, 1988, **65**, 814 (*isol, R-form*)
- Sener, B. *et al.*, *CA*, 1989, **110**, 111673f (*isol*)
- Tulyaganov, T.S. *et al.*, *Khim. Prir. Soedin. (Engl. Transl.)*, 1993, **29**, 509-511 (*Vasicinone N-oxide*)
- Magotra, D.K. *et al.*, *Acta Cryst. C*, 1996, **52**, 1491 (*cryst struct, 7-Methoxyvasicinone*)
- Eguchi, S. *et al.*, *J.O.C.*, 1996, **61**, 7316-7319 (*synth*)
- Thappa, R.K. *et al.*, *Phytochemistry*, 1996, **42**, 1485 (*5-Methoxyvasicinone, 7-Methoxyvasicinone*)
- Joshi, B.S. *et al.*, *Tetrahedron: Asymmetry*, 1996, **7**, 25 (*cryst struct, abs config*)
- Atta-ur-Rahman, *et al.*, *Nat. Prod. Lett.*, 1997, **10**, 249-256 (*cryst struct*)
- Mhaske, S.B. *et al.*, *J.O.C.*, 2002, **66**, 9038-9040 (*synth*)

Vasicol**V-39**

1-[(2-Aminophenyl)methyl]-3-hydroxy-2-pyrrolidinone, 9CI. Alkaloid AV2 [78720-03-3]



Absolute Configuration

$C_{11}H_{14}N_2O_2$ 206.244

Struct. finally established in 2003. Alkaloid from the roots of *Adhatoda vasica* and from *Peganum harmala*. Oil. $[\alpha]_D^{20}$ -90.4 (c, 0.95 in $CHCl_3$).

N,O-Di-Ac:

Cryst. (Et_2O /petrol). Mp 133-134° (130°).


Dhar, K.L. *et al.*, *Phytochemistry*, 1981, **20**, 319-321 (*isol, pmr, ms*)

Telezhenetskaya, M.V. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 1989, **25**, 14-18 (*isol, cryst struct*)

Huang, P.-Q. *et al.*, *Heterocycles*, 2003, **60**, 1833-1841 (*synth, pmr, cmr, abs config*)

Vasicoline**V-40**

N,N-Dimethyl-2-(1,2,3,9-tetrahydropyrrolo[2,1-b]quinazolin-3-yl)benzenamine, 9CI. 3-(2-Dimethylaminophenyl)-1,2,3,9-tetrahydropyrrolo[2,1-b]quinazoline [33903-13-8]



$C_{19}H_{21}N_3$ 291.395

Alkaloid from *Adhatoda vasica* (Acanthaceae). Cryst. (Me_2CO /heptane). Mp 135°.

9-Oxo: **Vasicolinone**. 3-[2-(Dimethylamino)phenyl]-2,3-dihydropyrrolo[2,1-b]quinazolin-9(1H)-one, 9CI [33903-15-0]

$C_{19}H_{19}N_3O$ 305.379

Alkaloid from *Adhatoda vasica* (Acanthaceae). Cryst. (hexane). Mp 152°.

Johne, S. *et al.*, *Helv. Chim. Acta*, 1971, **54**, 826 (*isol, uv, ir, pmr, ms, struct*)

Kaneko, C. *et al.*, *Heterocycles*, 1985, **23**, 1385 (*synth, ir, pmr, Vasicolinone*)

Kökösi, J. *et al.*, *Tet. Lett.*, 1992, **33**, 2995 (*synth, Vasicolinone*)

Wiedemann, S.H. *et al.*, *J.O.C.*, 2006, **71**, 1969-1976 (*synth*)

Vatamidine, 9CI**V-41**

[129741-48-6]

As Vatine, V-44 with

$n = 6$

$C_{88}H_{98}N_{16}$ 1379.849

Alkaloid from aerial parts of *Calycodendron milnei* (Rubiaceae). Platelet aggregation inhibitor, analgesic, sedative.

Amorph. powder. Sol. $MeOH, CHCl_3$; poorly sol. H_2O . $[\alpha]_D^{20}$ +77 (c, 1 in $EtOH$). λ_{max} 244 (log ϵ 4.56); 306 (log ϵ 4.33) ($EtOH$).

Adjibadé, Y. *et al.*, *Planta Med.*, 1990, **56**, 212-215 (*isol, uv, ir, pmr, cmr, ms, struct*)

Vatine, 9CI**V-42**

[129741-49-7]

As Vatine, V-44 with

$n = 5$


$C_{77}H_{86}N_{14}$ 1207.62

Alkaloid from aerial parts of *Calycodendron milnei* (Rubiaceae). Platelet aggregation inhibitor, analgesic, sedative. Fine needles ($MeOH$). Sol. $MeOH, CHCl_3$; poorly sol. H_2O . Mp 210°. $[\alpha]_D^{20}$ +64 (c, 1 in $EtOH$). λ_{max} 244 (log ϵ 4.65); 306 (log ϵ 4.42) ($EtOH$).

Adjibadé, Y. *et al.*, *Planta Med.*, 1990, **56**, 212-215 (*isol, uv, ir, pmr, cmr, ms, struct*)

Vateamine**V-43**

[87183-75-3]



$C_{38}H_{44}N_2O_8$ 656.774

Alkaloid from the bark of *Hernandia peltata* (Hernandiaceae). $[\alpha]_D^{25}$ +204 (c, 0.14 in $MeOH$).

N²-Oxide (β -): **Vateamine 2'- β -N-oxide** [178765-70-3]

$C_{38}H_{44}N_2O_9$ 672.774

Alkaloid from trunk bark of *Hernandia nymphaeifolia*. Shows strong anti-platelet aggregation activity. Needles ($MeOH$). Mp 133-135°. $[\alpha]_D^{24}$ +341 (c,

0.11 in $CHCl_3$).

Tri-Me ether: $[\alpha]_D^{25}$ +118 (c, 0.2 in $MeOH$).


Bruneton, J. *et al.*, *J.O.C.*, 1983, **48**, 3957 (*isol, uv, cd, pmr, ms, struct*)

Chen, J.-J. *et al.*, *Phytochemistry*, 1996, **42**, 1479 (*oxide*)

Chen, J.-J. *et al.*, *Planta Med.*, 2000, **66**, 251-256 (*oxide, activity*)

Vatine, 9CI**V-44**

[129741-50-0]



$C_{66}H_{74}N_{12}$ 1035.391

Alkaloid from aerial parts of *Calycodendron milnei* (Rubiaceae). Fine needles ($MeOH$). Mp 206-208°. $[\alpha]_D^{20}$ +57 (c, 1 in $EtOH$). λ_{max} 244 (log ϵ 4.49); 306 (log ϵ 4.25) ($EtOH$).

Stereoisomer: **Vatine A, 9CI**

[129829-05-6]


$C_{66}H_{74}N_{12}$ 1035.391

Alkaloid from aerial parts of *Calycodendron milnei* (Rubiaceae). Amorph. powder. $[\alpha]_D^{20}$ +45 (c, 1 in $EtOH$). λ_{max} 244 (log ϵ 3.43); 306 (log ϵ 4.18) ($EtOH$).

Adjibadé, Y. *et al.*, *Planta Med.*, 1990, **56**, 212-215 (*isol, uv, ir, pmr, cmr, ms, struct*)

Veatchine, 9CI**V-45**

[76-53-9]



Absolute Configuration

$C_{22}H_{33}NO_2$ 343.508

Exists as mixt. of C-20 epimers in soln. and in the solid state. Alkaloid from the bark of *Garrya veatchii* (Garryaceae). Prismatic cryst. (Me_2CO). Mp 125.5-128.5° (122-126°). $[\alpha]_D^{30}$ -67.5 (c, 1.35 in $CHCl_3$). Unlike Garryfoline, stable to acid-catalysed rearr.

Hydrochloride: **Veatchinium chloride**

Cryst. ($EtOH/Et_2O$). Mp 273-276° (267-271°). $[\alpha]_D^{29}$ -56.2 (c, 1.45 in $EtOH$). Exists as a ternary iminium salt rather than a protonated salt.

16 α ,17-Dihydro, 15-ketone: Cuauchichicine

[545-60-8]

C₂₂H₃₃NO₂ 343.508

Alkaloid from the bark of *Garrya laurifolia* and the leaves of *Garrya ovata* var. *lindheimeri* (Garryaceae). Cryst. (Et₂O). Mp 152-154°. [α]_D¹⁸ -69 (c, 1.0 in CHCl₃). Unusually, does not exist in the epimeric form at C-20 either in soln. or in the solid state.

▶ YX0800000

16 α ,17-Dihydro, 15-ketone, hydrochloride:Cryst. (MeOH/Et₂O). Mp 259-262°.**15-Epimer: Garryfoline. Laurifoline†**

[509-30-8]

C₂₂H₃₃NO₂ 343.508

Alkaloid from the bark of *Garrya laurifolia* and *Garrya ovata* var. *lindheimeri* (Garryaceae). Cryst. (Et₂O or hexane). Mp 130-133°. [α]_D -60 (CHCl₃). pK_a 11.81. Exists as a mixt of C-20-epimers in soln. Rearr. to Garryine, G-25 in hot MeOH. Undergoes acid-cat. rearr. to Cuauchichicine.

15-Epimer, Ac: Ovatine†. (15 β)-Veatchine acetate (ester), 9CI

[68719-14-2]

C₂₄H₃₅NO₃ 385.545

Alkaloid from the bark and leaves of *Garrya ovata* var. *lindheimeri* (Garryaceae). Mp 113-114°. [α]_D²² -79.4 (c, 1.0 in CHCl₃). Mixt. of C-20-epimers in soln.

Oneto, J.F. et al., *J. Am. Pharm. Assoc.*, 1946, **35**, 204 (isol)

Wiesner, K. et al., *Can. J. Chem.*, 1952, **30**, 608 (isol, struct, ir)

Wiesner, K. et al., *Chem. Ber.*, 1953, **86**, 800 (isol, struct, ir)

Wiesner, K. et al., *Experientia*, 1955, **11**, 255 (isol, struct, ir)

Djerassi, C. et al., *J.A.C.S.*, 1955, **77**, 4801; 6633 (*Garryfoline, Cuauchichicine, isol, ir, nomencl*)

Verbrueggen, H. et al., *J.A.C.S.*, 1962, **84**, 2990-2997 (abs config)

Pelletier, S.W. et al., *J.A.C.S.*, 1965, **87**, 761; 1977, **99**, 284; 1978, **100**, 7976; 1979, **101**, 6741 (cmr, cryst struct, abs config, isol, struct)

Pelletier, S.W. et al., *Tetrahedron*, 1968, **24**, 2019 (pmr)

Scott, A.I. et al., *Tetrahedron*, 1971, **27**, 4787 (cd, ord)

Pelletier, S.W. et al., *Heterocycles*, 1978, **9**, 1409; 1979, **13**, 277 (*Garryfoline, Ovatine*)


Mody, N.V. et al., *Tetrahedron*, 1978, **34**, 2421 (cmr)

Pelletier, S.W. et al., *J.A.C.S.*, 1979, **101**, 6741-6742 (*Cuauchichicine, pmr, cmr, cryst struct*)

Pelletier, S.W. et al., *J. Nat. Prod.*, 1980, **43**, 41 (rev)

Veitamine

[188010-55-1]

**V-46**C₁₇H₁₆N₃O₂[⊕] 294.332


Alkaloid from the Fijian sponge *Zyzzya fuliginosa*. Potent *in vitro* cytotoxin. Green solid (as trifluoroacetate). λ_{\max} 246 ; 344 ; 550 (MeOH). λ_{\max} 228 ; 330 ; 470 (MeOH/KOH).

Venables, D.A. et al., *Tet. Lett.*, 1997, **38**, 721 (isol, uv, ir, pmr, cmr, struct)

Moro-oka, Y. et al., *Tet. Lett.*, 1999, **40**, 1713-1716 (synth, pmr, cmr)

Venacarpine A

[768388-29-0]

R,R' = -OCH₂O-C₂₂H₂₂N₂O₄ 378.427

Alkaloid from *Kopsia fruticosa*. Oil. [α]_D +18 (c, 0.43 in CHCl₃). λ_{\max} 219 (log ϵ 4.46); 244 (log ϵ 4); 280 (log ϵ 3.53) (EtOH).

Kam, T.-S. et al., *Phytochemistry*, 2004, **65**, 2119-2122 (isol, pmr, cmr, ms)

Venacarpine B

As Venacarpine A, V-47 with

R = H, R' = OMe


C₂₂H₂₄N₂O₃ 364.443

Alkaloid from leaves of *Kopsia fruticosa*. [α]_D +18 (c, 0.43 in CHCl₃). λ_{\max} 216 (log ϵ 4.27); 243 (log ϵ 3.93); 284 (log ϵ 3.5) (EtOH).

Kam, T.S. et al., *Phytochemistry*, 2004, **65**, 2119-2122 (isol, pmr, cmr, ms)

Venalstonine**14,15-Dehydrokopsinine. $\Delta^{14,15}$ -Kopsinine**

[5001-20-7]

C₂₁H₂₄N₂O₂ 336.433

Alkaloid from *Alstonia venenata*, *Catharanthus ovalis*, *Catharanthus roseus*, *Melodinus australis*, *Vinca erecta*, *Vinca libanotica* and several other spp. in the Apocynaceae. Prisms (Et₂O/pentane). Mp 132-134° Mp 140-142°. [α]_D²³ -88.8 (c, 1.406 in CHCl₃). λ_{\max} 245 (log ϵ 3.87); 292 (log ϵ 3.46) (EtOH).

14 α ,15 α -Epoxide: Venalstonidine. 6,7-**Epoxykopsinine. Ld 63**

[26619-93-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from *Alstonia venenata*, *Melodinus australis*, *Melodinus balsansae*,

V-47

Absolute Configuration

V-48**V-49**

Absolute Configuration

Melodinus celastroides, *Melodinus polyadenus* and *Catharanthus ovalis* (Apocynaceae). Mp 230-236° (210-214°, 223-225°). [α]_D -96 (CHCl₃). λ_{\max} 244 (ϵ 7150); 292 (ϵ 2800) (EtOH).

3-Oxo: 14,15-Didehydro-3-oxokopsinine. Ld 65

[4880-84-6]

C₂₁H₂₂N₂O₃ 350.416

Alkaloid from *Vinca erecta* and *Melodinus australis*. Mp 231-232°. [α]_D -93 (c, 0.3 in CHCl₃). λ_{\max} 244 (log ϵ 3.93); 292 (log ϵ 3.42) (EtOH).

3-Oxo, N⁴-oxide: 14,15-Didehydro-3-oxokopsinine N-oxide. Ld 85C₂₁H₂₂N₂O₄ 366.416

Alkaloid from *Vinca erecta* and *Melodinus australis*. Cryst. (Et₂O). Mp 245-248°. [α]_D -55.

3-Oxo, 14 α ,15 α -epoxide: 3-Oxovenalstonidine

[87387-68-6]

C₂₁H₂₂N₂O₄ 366.416

Alkaloid from stems and leaves of *Melodinus reticulatus* (Apocynaceae). [α]_D²² -15 (c, 1 in CHCl₃).

17 α -Hydroxy: 17 α -Hydroxy- $\Delta^{14,15}$ -kopsinine

[149355-60-2]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the stem bark of *Kopsia teoi*.

17 α -Hydroxy, Et ester analogue: Kopsiloscine JC₂₂H₂₆N₂O₃ 366.459Alkaloid from *Kopsia singapurensis*.

Oil. [α]_D +22 (c, 0.23 in CHCl₃). λ_{\max} 207 (log ϵ 4.41); 241 (log ϵ 3.96); 289 (log ϵ 3.58) (EtOH).

17 β -Hydroxy, N-methoxycarbonyl: N-Carbomethoxy-17 β -hydroxy- $\Delta^{14,15}$ -kopsinine

[116988-04-6]

C₂₃H₂₆N₂O₅ 410.469

Alkaloid from the stem bark of *Kopsia deverrei* (Apocynaceae). Amorph. [α]_D²⁰ -110 (c, 0.50 in CHCl₃). λ_{\max} 245 (log ϵ 4.21); 280 (log ϵ 3.63); 288 (log ϵ 3.59) (EtOH).

19 β -Hydroxy: 19 β -Hydroxyvenalstonine

[87387-66-4]

C₂₁H₂₄N₂O₃ 352.432

Alkaloid from stems and leaves of *Melodinus reticulatus* (Apocynaceae). [α]_D²² -32 (c, 1 in CHCl₃).

19 β -Hydroxy, 14 α ,15 α -epoxide: 19-Hydroxyvenalstonidine

[87387-67-5]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from stems and leaves of *Melodinus reticulatus* (Apocynaceae). [α]_D²² -25 (c, 1 in CHCl₃).

16-Epimer, 17 α -hydroxy: 16-Epi-17 α -hydroxy- $\Delta^{14,15}$ -kopsinine

[152243-72-6]

C₂₁H₂₄N₂O₃ 352.432


Alkaloid from the stem bark of *Kopsia teoi*. Amorph. powder. [α]_D²⁰ -23 (c, 1.48 in CHCl₃).

Linde, H.H.A. et al., *Helv. Chim. Acta*, 1965, **48**, 1822-1842 (*Venalstonidine*)

- Das, B. *et al.*, *Tet. Lett.*, 1965, 2239-2244 (*Venalstonine, Venalstonidine*)
 Aynilian, G.H. *et al.*, *J. Nat. Prod.*, 1974, **37**, 299-308 (*Venalstonine*)
 Ahond, A. *et al.*, *J.A.C.S.*, 1974, **96**, 633-634 (*cmr*)
 Sharipov, M. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 263-264; 1976, **12**, 401-402; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 281; 1976, **12**, 355-356 (*14,15-Didehydro-3-oxokopsinine, 14,15-Didehydro-3-oxokopsinine N-oxide*)
 Mehri, M.H. *et al.*, *Phytochemistry*, 1978, **17**, 1451-1452 (*Venalstonine*)
 Mehri, H. *et al.*, *Planta Med.*, 1983, **48**, 72-76 (*19-Hydroxyvenalstonine, 19-Hydroxyvenalstonidine, 3-Oxovenalstonidine*)
 Kan-Fan, C. *et al.*, *J. Nat. Prod.*, 1988, **51**, 703-707 (*N-Carbomethoxy-17-hydroxy- $\Delta^{14,15}$ -kopsinine*)
 Varea, T. *et al.*, *J. Nat. Prod.*, 1993, **56**, 2166-2169 (*16-Epi-17 α -hydroxy- $\Delta^{14,15}$ -kopsinine*)
 Kam, T.S. *et al.*, *Phytochemistry*, 1999, **50**, 171-175 (*17 α -Hydroxy- $\Delta^{14,15}$ -kopsinine*)
 Subramaniam, G. *et al.*, *J. Nat. Prod.*, 2008, **71**, 53-57 (*Kopsilosine J*)

Venenatine V-50

Methyl 17-hydroxy-9-methoxyyohimban-16-carboxylate, 9CI
 [1055-75-0]



Absolute Configuration

$C_{22}H_{28}N_2O_4$ 384.474
 Alkaloid from *Alstonia venenata* (Apocynaceae). Antihypertensive agent showing similar action to, and synergism with, Reserpine, R-52. Shows antifungal activity. Cryst. + MeOH (MeOH). Mp 123-126° dec. $[\alpha]_D^{24}$ -76.1 (CHCl₃). pK_a 7.2 (2-Methoxyethanol).

▶ ZG0820000

Hydroiodide:

Cryst. (Me₂CO). Mp 255-257° dec.

N⁴-Oxide: Venoxidine

[17206-07-4]

$C_{22}H_{28}N_2O_5$ 400.474

Alkaloid from *Alstonia venenata* (Apocynaceae). Mp 218-219° dec. $[\alpha]_D^{25}$ -58.2 (H₂O). pK_a 4.6 (H₂O).

O-Ac:

Cryst. (MeOH). Mp 98-101° dec.

Parent acid: Venenatic acid

[4837-83-6]

Cryst. + 2H₂O (EtOH). Mp 245° dec.

3-Epimer: Alstovenine. Isovenenatine

[4837-79-0]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Alstonia venenata* (Apocynaceae). Monoamine oxidase inhibitor. Needles (MeOH). Mp 169-171°. $[\alpha]_D^{24}$ +9.42 (CHCl₃).

▶ ZG0830000

3-Epimer, O-Ac:

Needles (MeOH). Mp 143° dec.

16-Epimer: 16-Epivenenatine

[80735-17-7]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Alstonia venenata* (Apocynaceae). Cryst. (Me₂CO/MeOH). Mp 188-190°. $[\alpha]_D^{25}$ -52.68 (EtOH).

17-Epimer: 17-Epivenenatine. 9-Methoxy-3-epi- α -yohimbine

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from the stem bark of *Mitragyna africanus*. Amorph. powder. λ_{max} 224 (log ϵ 4.27); 270 (log ϵ 3.6) (MeOH).

3,16-Diepimer: 16-Epialstovenine

[80734-61-8]

$C_{22}H_{28}N_2O_4$ 384.474

Alkaloid from *Alstonia venenata* (Apocynaceae). Cryst. (Me₂CO/MeOH). Mp 227-230°. $[\alpha]_D^{25}$ +72.02 (EtOH).

[1056-57-1]

Chatterjee, A. *et al.*, *Tet. Lett.*, 1965, 159-162 (*Venenatine, Venoxidine, isol, uv, ir, pmr, ms, struct*)

Govindachari, T.R. *et al.*, *Tetrahedron*, 1965, **21**, 2951-2956 (*isol, uv, ir, pmr, struct, Venenatine, Alstovenine*)

Chatterjee, A. *et al.*, *Phytochemistry*, 1981, **20**, 1981-1985 (*16-Epivenenatine, 16-Epialstovenine*)

Verpoorte, R. *et al.*, *Org. Magn. Reson.*, 1984, **22**, 328-334 (*cmr*)

Singh, U.P. *et al.*, *Folia Microbiol. (Prague)*, 2000, **45**, 173-176 (*activity*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 2004, **52**, 359-361 (*17-Epivenenatine*)

C-Venezueline V-51


Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). An oxidn. prod. of C-Alkaloid D, A-476.

Gorman, A.A. *et al.*, *Alkaloids (London)*, 1971, **1**, 200-338

Venoterpine V-52

Gentialutine. Alkaloid RW47

[17948-42-4]



$C_9H_{11}NO$ 149.192

Config. revised in 1985. Alkaloid from *Alstonia venenata*, *Rauwolfia verticillata*, *Lonicera japonica*, *Neisosperma glomerata*, *Melodinus aeneus*, *Gentiana lutea* (yellow gentian), *Strychnos dinklagei*, *Strychnos pungens* and *Camptotheca acuminata*. Mp 130-132°. $[\alpha]_D$ +27 (CHCl₃).

O-Ac: O-Acetylvenoterpine

[189322-84-7]

$C_{11}H_{13}NO_2$ 191.229

Alkaloid from seeds of *Strychnos pungens*. $[\alpha]_D$ +45 (c, 1.5 in CHCl₃).

Arthur, H.R. *et al.*, *Phytochemistry*, 1966, **5**, 977-983 (*isol*)

Arthur, H.R. *et al.*, *Aust. J. Chem.*, 1967, **20**, 2505-2508 (*uv, ir, pmr, ms, struct*)

Ray, A.B. *et al.*, *Tet. Lett.*, 1968, 2763-2765 (*uv, ir, pmr, struct*)

Mitscher, L.A. *et al.*, *Experientia*, 1971, **27**, 16-17 (*config*)


Michel, S. *et al.*, *J. Nat. Prod.*, 1982, **45**, 489-494 (*isol*)

Ravao, T. *et al.*, *Tet. Lett.*, 1985, **26**, 837-838 (*config, cmr*)

Thépenier, P. *et al.*, *Bull. Soc. R. Sci. Liege*, 1996, **65**, 379-382 (*Ac*)

Venturamide A V-53

[312958-58-0]



R = CH₃

$C_{21}H_{24}N_6O_4S_2$ 488.59

Cyclic peptide. Related to Bistratamide A, B-223. Isol. from an *Oscillatoria* sp. Antimalarial agent. Glass. $[\alpha]_D^{25}$ +53.4 (c, 0.001 in MeOH). λ_{max} 224 (log ϵ 4.17) (MeOH).

Linnington, R.G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 397-401 (*isol, pmr, cmr*)

Venturamide B V-54

[936098-82-7]

As Venturamide A, V-53 with

R = -CH(OH)CH₃(R-)


$C_{22}H_{26}N_6O_5S_2$ 518.617

Isol. from an *Oscillatoria* sp. Antimalarial agent. Glass. $[\alpha]_D^{25}$ +53.6 (c, 0.004 in MeOH). λ_{max} 224 (log ϵ 4.08) (MeOH).

Linnington, R.G. *et al.*, *J. Nat. Prod.*, 2007, **70**, 397-401 (*isol, pmr, cmr*)

Vepridimerine C V-55

[82841-72-3]



$C_{34}H_{38}N_2O_8$ 602.683

Alkaloid from the bark of *Vepris louisii* and *Orcicia renieri* (Rutaceae). Mp 272°. Racemic.

7 α -Epimer: Vepridimerine E

[101541-27-9]

Synthetic. Cryst. (MeOH). Mp 287-288°.


16 α -Epimer: Vepridimerine D

[82864-61-7]
 $C_{34}H_{38}N_2O_8$ 602.683
 Alkaloid from the bark of *Vepris louisii* (Rutaceae). Racemic.
 Ngadjui, T.B. *et al.*, *Tet. Lett.*, 1982, **23**, 2041 (pmr, cmr, struct)
 Ayafor, J.F. *et al.*, *Tet. Lett.*, 1985, **26**, 4529 (synth)

Veracevine

V-56

4,9-Epoxy-3,4,12,14,16,17,20-cevaneheptol. Protocevine
 [5876-23-3]



Absolute configuration

$C_{27}H_{43}NO_8$ 509.639
 Alkaloid from *Schoenocaulon officinale* (Liliaceae). Sabadilla (Veratrum D), a mixt. of Cevadine and Veratridine has been used as an insecticide. Mp 181-183°. $[\alpha]_D^{25}$ -24 (EtOH).

▶ FL4740000

Perchlorate: Mp 228-230°.

3-Ac: **Cevacine**

[28111-33-3]
 $C_{29}H_{45}NO_9$ 551.676

Alkaloid from *Schoenocaulon officinale* (Liliaceae). Mp 205-207°. $[\alpha]_D^{21}$ -27 (c, 2.67 in $CHCl_3$).

3-Angeloyl: **Cevadine**. Veratrine

[62-59-9]
 $C_{32}H_{49}NO_9$ 591.74

Alkaloid from *Schoenocaulon officinale*, *Helleborus viridis*, *Veratrum lobelianum* and *Veratrum viride*. Insecticide. Weak hypotensive agent. Shows antifungal activity. Mp 213-214.5°. $[\alpha]_D^{28}$ +13 (c, 0.97 in EtOH).

▶ LD₅₀ (mus, ipr) 3.5 mg/kg. FL56000003-O-(4-Hydroxy-3-methoxybenzoyl): **Vanilloveracevine**

[187237-90-7]
 $C_{35}H_{49}NO_{11}$ 659.772

Alkaloid from *Schoenocaulon officinale*. Antihypertensive agent. Shows insecticidal activity. Silky needles (EtOH). Mp 257.5-258.5° dec. $[\alpha]_D^{25}$ +43.6 (c, 1.87 in Py). Log P -0.33 (uncertain value) (calc).

3-O-(3,4-Dimethoxybenzoyl): **Veratridine**

[71-62-5]
 $C_{36}H_{51}NO_{11}$ 673.799

Alkaloid from *Schoenocaulon officinale* and *Veratrum album*. Hypotensive agent. Depolarising agent for neurones and muscle cells via specific sodium channel interaction, similar activity to Batrachotoxin, but weaker. Shows antifungal activity. Noncryst. $[\alpha]_D^{22}$ +8 (EtOH).

▶ LD₅₀ (mus, ipr) 1.35 mg/kg. YX56000003-Epimer: **Cevine**. Sabadinine
 [124-98-1]

$C_{27}H_{43}NO_8$ 509.639
 Alkaloid from seeds of *Veratrum sabadilla* and product of hydrol. of ester alkaloids (Liliaceae). Mp 172-176° dec. (anhyd.) Mp 110° (hydrate). $[\alpha]_D^{17}$ -17.5 (EtOH).

▶ LD₅₀ (rat, ipr) 67 mg/kg. FL4550000

3-Epimer, hydrochloride: Mp 247° Mp 239-240° (dihydrate).

Pelletier, S.W. *et al.*, *J.A.C.S.*, 1953, **75**, 3248-3252 (Cevadine, Veratridine)Kupchan, S.M. *et al.*, *J.A.C.S.*, 1953, **75**, 5519-5524 (Cevacine)Barton, D.H.R. *et al.*, *Experientia*, 1954, **10**, 81-90 (struct, bibl)Stuart, D.M. *et al.*, *J. Am. Pharm. Assoc.*, 1956, **45**, 252-256 (Vanilloveracevine)Kupchan, S.M. *et al.*, *Tetrahedron*, 1959, **7**, 47-61 (struct)Eeles, W.T. *et al.*, *Tet. Lett.*, 1960, **No. 7**, 24-26 (Cevine, cryst struct)Wolters, B. *et al.*, *Planta Med.*, 1970, **19**, 189-193 (Veracevine, activity)Carey, F.A. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 141-144 (Veracevine, cmr)Holan, G. *et al.*, *J. Chromatogr.*, 1984, **288**, 479-483 (Veracevine, hplc)Ujváry, I. *et al.*, *Phytochemistry*, 1997, **44**, 1257-1260 (Vanilloveracevine, synth)


Pesticide Manual, 13th edn., 2003, 725

Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, CDG000; Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VHU000; EBL000

Veractintine

V-57

22,25-Epimino-27-norcholesta-5,25(N)-dien-3-ol
 [33596-06-4]

 $C_{26}H_{41}NO$ 383.616

Alkaloid from *Veratrum album* (Liliaceae). Shows cytotoxic and antitumour activity. Mp 196-201°. $[\alpha]_D^{25}$ +7.5 (c, 0.825 in $CHCl_3$).

3-O-β-D-Glucopyranoside: **Glucoveractintine**

[67006-43-3]
 $C_{32}H_{51}NO_6$ 545.758

Alkaloid from the aerial parts of *Veratrum album* ssp. *lobelianum* (Liliaceae). Amorph. $[\alpha]_{11g}^{19}$ +26 (c, 0.69 in MeOH).

3-O-α-L-Rhamnocide: **Rhamnoveractintine**

[110934-18-4]
 $C_{32}H_{51}NO_5$ 529.759

Alkaloid from the aerial parts of *Veratrum album* ssp. *lobelianum* (Liliaceae).

O,N-Di-Ac (of 5,24-diene isomer):

Amorph. $[\alpha]_D^{25}$ -98 (EtOH).Tomko, J. *et al.*, *Tet. Lett.*, 1971, 3041 (isol, ms, ir, pmr)Vassova, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1976, **41**, 2964 (cmr)Grančai, D. *et al.*, *Chem. Zvesti*, 1978, **32**, 120 (glucoside)Fuska, J. *et al.*, *CA*, 1982, **96**, 173914m (pharmacol)Pavelcik, F. *et al.*, *Chem. Zvesti*, 1983, **37**, 145 (cryst struct)Grančai, D. *et al.*, *Chem. Zvesti*, 1986, **40**, 835; *CA*, **107**, 194870e (Rhamnoveractintine)**Veragenine†**

V-58

 $C_{31}H_{53}NO_{13}$ 647.759

Struct. unknown. Approx. MF, may be H₅₅. Alkaloid from *Veratrum officinale* (*Veratrum sabadilla*) (Liliaceae). Mp 262-264°. $[\alpha]_D^{20}$ -4.87 (EtOH).

Vejdělek, Z.J. *et al.*, *Chem. Listy*, 1955, **49**, 1538-1545; *CA*, **50**, 4991**Veralbidine**

V-59

 $C_{37}H_{61}NO_{12}$ 711.888

Steroidal alkaloid. Struct. unknown. Alkaloid from *Veratrum album* (Liliaceae). Prisms (MeOH aq.), pentagonal plates (Me₂CO aq.), fine needles (Et₂O). Mp 181-183°. $[\alpha]_D^{20}$ -11.7 (Py). $[\alpha]_D^{20}$ +5.4 ($CHCl_3$).


Hydrochloride: Mp 250-251° dec.

Thiocyanate salt: Mp 235-236° dec.

Stoll, A. *et al.*, *Science (Washington, D.C.)*, 1952, **115**, 678**Veralinine**

V-60

17-Methyl-18-nor-16,28-secosolanida-5,12-dien-3-ol, 9CI. 16-Deoxyveralkamine
 [21233-19-2]



Absolute Configuration

 $C_{27}H_{43}NO$ 397.643

Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Needles (hexane). Mp 124-126°. $[\alpha]_D^{25}$ -80 (c, 0.42 in $CHCl_3$).

N,O-Di-Ac:

Cryst. (Et₂O). Mp 159-161°. $[\alpha]_D^{21}$ -52.8 (c, 0.56 in $CHCl_3$).Stereoisomer: **Veralomidine**

[58165-72-3]

Hydrol. prod. of Veralomine. Mp 174-175°. $[\alpha]_D$ -52.23. Said to be epimeric with Veralinine at C₁₇.

Stereoisomer, 3-β-D-glucopyranoside:

Veralomine

[58078-63-0]

 $C_{33}H_{53}NO_6$ 559.785

Isol. from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 275-277°. $[\alpha]_D$ -54.11.

Stereoisomer, 3,4-didehydro: **Veralomidene**

[58031-31-5]

Hydrol.-dehydration prod. of Veralo-


mine. Mp 175-178°. $[\alpha]_D$ -49.5.

Tomko, J. *et al.*, *Tetrahedron*, 1968, **24**, 6839-6843 (*isol, ms, ir, pmr, struct*)

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 527-528; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 558-559 (*Veralomine*)

Veralkamine**V-61**

17-Methyl-18-nor-16,28-secosolanida-5,12-diene-3,16-diol, 9CI. *Veralkamine* [17155-31-6]



Absolute Configuration

$C_{27}H_{43}NO_2$ 413.642

Alkaloid from *Veratrum album* ssp. *lobelianum* and from *Fritillaria camtschaticensis* (Liliaceae). Cryst. (EtOH). Mp 119-123° Mp 165-169°. $[\alpha]_D^{24}$ -84.1 (c, 0.55 in $CHCl_3$).

Hydroiodide: Mp 264° dec.

O,O-Di-Ac: [195244-85-0]

$C_{31}H_{47}NO_4$ 497.717

Alkaloid from *Veratrum lobelianum*.

N,O,O-Tri-Ac: Mp 152-154°. $[\alpha]_D^{22}$ -8 ($CHCl_3$).

16-Deoxy: see Veralinine, V-60

5 α ,6,12,13-Tetrahydro: **Tetrahydroveralkamine**

[17155-36-1]

$C_{27}H_{47}NO_2$ 417.674

Alkaloid from *Veratrum lobelianum*.

Cryst. (EtOH). Mp 219-221°. $[\alpha]_D^{25}$ +6.1 (c, 1.2 in $CHCl_3$).

Tomko, J. *et al.*, *Tetrahedron*, 1968, **24**, 4865-4873 (*isol, pmr, ms, ir, struct*)

Höhne, E. *et al.*, *Tetrahedron*, 1968, **24**, 4875-4880 (*cryst struct*)


Tomko, J. *et al.*, *CA*, 1981, **94**, 205391r (*isol*)

Kaneko, K. *et al.*, *Phytochemistry*, 1981, **20**, 327-329 (*isol*)

Foldesiova, V. *et al.*, *CA*, 1997, **127**, 231880u (*isol, derivs*)

Veralobine**V-62**

18-Hydroxysolanid-4-en-3-one [6242-49-5]



Absolute Configuration

$C_{27}H_{41}NO_2$ 411.626

Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 238°. $[\alpha]_D^{20}$ +67 (c, 0.1 in MeOH).


Tomko, J. *et al.*, *Pharmazie*, 1965, **20**, 385 (*isol, struct*)

Tomko, J. *et al.*, *Arch. Pharm. (Weinheim, Ger.)*, 1966, **299**, 347-350 (*struct*)

Höhne, E. *et al.*, *Tetrahedron*, 1966, **22**, 673-678 (*cryst struct*)

Veralodisine**V-63**

16-(Acetyloxy)-3-hydroxy-16,28-secosolanida-5,22(28)-dien-24-one, 9CI [52617-23-9]



Absolute Configuration

$C_{29}H_{43}NO_4$ 469.663

Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 172-174°. $[\alpha]_D$ -92.8 (c, 0.49 in $CHCl_3$).

3-O- β -D-Glucopyranoside: **Veralodinine** [56598-27-7]

$C_{35}H_{53}NO_9$ 631.805


Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Cryst. (MeOH). Mp 226-228°. $[\alpha]_D$ -95.4 (c, 0.524 in $CHCl_3$).

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 44-48; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 38-41 (*ir, uv, pmr, ms, struct*)

Samikov, K. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 183-188; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 193-196 (*Veralodinine*)

Veralosidinine**V-64**

16-Acetoxy-16,28-secosolanida-5,22(28)-diene-3,5-diol [52389-14-7]



Absolute configuration

$C_{29}H_{45}NO_4$ 471.679


Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 220-221°. $[\alpha]_D$ -173 (c, 0.59 in $CHCl_3$).

Shakirov, R. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 501; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 472 (*ir, uv, pmr, ms, struct*)

Moiseeva, G.P. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 623; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 557 (*cd*)

Veramanine**V-65**

[182816-87-1]



$C_{27}H_{43}NO_5$ 461.64


Alkaloid from rhizomes of *Veratrum*

album. Yellow solid. $[\alpha]_D^{25}$ -33.5 (c, 0.08 in $CHCl_3$). λ_{max} 247 (MeOH).

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1996, **43**, 907 (*isol, uv, ir, pmr, cmr, ms, struct*)

Veramine**V-66**

17-Methyl-18-norspirosola-5,12-dien-3-ol, 9CI [21059-48-3]



$C_{27}H_{41}NO_2$ 411.626

Alkaloid from *Veratrum album* ssp. *lobelianum*, *Veratrum nigrum* and *Veratrum oxysepalum* (Liliaceae). Amorph. $[\alpha]_D^{24}$ -85.6 (c, 0.74 in EtOH). $[\alpha]_D^{24}$ -93.9 (c, 1.04 in $CHCl_3$).

Tomko, J. *et al.*, *Chem. Zvesti*, 1964, **18**, 266 (*isol*)

Adam, G. *et al.*, *CA*, 1972, **77**, 114667t (*struct, ir, uv, pmr, ms*)

Bondarenko, N.V. *et al.*, *Khim. Prir. Soedin.*, 1981, **17**, 527 (*isol*)


Bondarenko, N.V. *et al.*, *Khim. Prir. Soedin.*, 1984, **20**, 263; *Chem. Nat. Compd. (Engl. Transl.)*, 1984, **20**, 248 (*isol*)

Veraminine**V-67**

Prob. steroidal alkaloid. Struct. unknown. Alkamine from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 201-206°. Tomko, J. *et al.*, *Chem. Zvesti*, 1964, **18**, 266

Veramitaline**V-68**

16,28-Secosolanida-5,22(28)-diene-3,12-diol. 22,26-Epiminocholesta-5,22-diene-3,12-diol [313697-00-6]



$C_{27}H_{43}NO_2$ 413.642

Alkaloid from the roots and rhizomes of *Veratrum taliense*. Needles (MeOH). Mp 217-218°.

Zhou, C.X. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 283-286 (*isol, pmr, cmr*)

Veranovine**V-69**

Struct. unknown. Alkaloid from *Veratrum viride* (Liliaceae). Prob. a mixt. *Hydrochloride*: Mp 190-200°.

U.S. Pat., 1957, 2 787 616; *CA*, **51**, 9102e

Verareine

V-70


C₂₇H₄₁NO 395.627

Steroidal alkaloid. Struct. unknown. Alkaloid from *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 150-155°. [α]_D²⁷ -13 (EtOH). [α]_D²² +6.1 (CHCl₃).

Tomko, J. et al., *Chem. Zvesti*, 1964, **18**, 266**Veratramine**

V-71

14,15,16,17-Tetrahydroveratraman-3,23-diol, 9CI
[60-70-8]

C₂₇H₃₉NO₂ 409.611

Alkaloid from *Veratrum album* var. *grandiflorum* and several other *Veratrum* spp. (Liliaceae). Shows antifungal activity. Active against *Helicobacter pylori*. Powerful cardiac stimulant. Effect not antagonised by atropine. Mp 209.5-210.5°. [α]_D¹⁹ -70 (c, 1.56 in MeOH). pK_a 7.49. λ_{\max} 268 (MeOH) (Berdy).

► LD₅₀ (mus, scu) 4.5 mg/kg. Exp. teratogen. YX5094000

Hydrochloride: Mp 310° (201-202°).

O³- β -D-Glucopyranoside: **Veratrosine**
[475-00-3]

C₃₃H₄₉NO₇ 571.753

Alkaloid from *Veratrum viride* and *Veratrum eschscholtzii* (Liliaceae). Mp 242-243° dec. (236-239°). [α]_D²⁵ -53 (c, 0.26 in CHCl₃/EtOH 1:1).

► YX7050000

23-Deoxy: **Verarine**. 23-Deoxyveratramine

C₂₇H₃₉NO 393.611

From *Veratrum album* ssp. *lobelianum* (Liliaceae). Mp 174-176°. [α]_D²⁵ -68.5 (EtOH).

23-Deoxy, N-Ac: Mp 195-196°. [α]_D²⁸ -14.4 (c, 1.02 in EtOH).

23-Deoxy, O,N-di-Ac: Mp 189-190°. [α]_D²⁶ -21.3 (c, 1.33 in EtOH).

20-Epimer: **20-Isoveratramine**C₂₇H₃₉NO₂ 409.611

Alkaloid from *Veratrum patulum*. Prisms. Mp 267-270°. [α]_D²³ +102.2 (c, 0.09 in MeOH). λ_{\max} 207 (log ϵ 4.32) (EtOH).

20-Epimer, 23-O- β -D-glucopyranoside: **23-O- β -D-Glucopyranosyl-20-isoveratramine**
[148440-62-4]

C₃₃H₄₉NO₇ 571.753

Alkaloid from roots of *Veratrum patulum* (Liliaceae). Amorph. powder. Mp 165°. [α]_D²⁰ -95 (c, 1.0 in Py).

Jacobs, W.A. et al., *J. Biol. Chem.*, 1944, **155**, 565 (isol, Veratrosine)

Tamm, C. et al., *J.A.C.S.*, 1952, **74**, 3842 (uv, ir, struct)

Tomko, J. et al., *Coll. Czech. Chem. Comm.*, 1964, **29**, 2570 (Verarine)

Budzikiewicz, H. et al., *Tetrahedron*, 1964, **20**, 2267 (ms)

Johnson, W.S. et al., *J.A.C.S.*, 1967, **89**, 4523 (synth)

Kupchan, S.M. et al., *J.A.C.S.*, 1968, **90**, 2730 (config)

Masamune, T. et al., *Tetrahedron*, 1971, **27**, 3387 (pmr)

Sprague, P.W. et al., *Tetrahedron*, 1971, **27**, 4852 (cmr)

Han, Y.B. et al., *Yakhak Hoeji*, 1973, **17**, 13-16; 137-140 (activity)

Kutney, J.P. et al., *Can. J. Chem.*, 1975, **53**, 1796 (synth)

Irsch, E.M. et al., *Annalen*, 1993, 281 (23-O- β -D-Glucopyranosyl-20-isoveratramine)


Tezuka, Y. et al., *J. Nat. Prod.*, 1998, **61**, 1078-1081 (20-Isoveratramine)

Tezuka, W. et al., *Wakan Iyagakugaku Zasshi*, 1999, **16**, 196; *CA*, **132**, 276648u (activity)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VHP500

Veratra-5,11,13-triene-3,23-diol

V-72

C₂₇H₄₁NO₂ 411.626**(3 β ,22S,23R,25S)-form**23-O- β -D-Glucopyranoside: [148440-63-5]C₃₃H₅₁NO₇ 573.768


Alkaloid from roots of *Veratrum patulum* (Liliaceae). Amorph. powder.

Irsch, E.-M. et al., *Annalen*, 1993, 281 (isol, pmr, cmr, struct)

Verazine

V-73

16,28-Secosolanida-5,22(28)-dien-3-ol, 9CI. 22,26-Iminocholesta-5,22(N)-dien-3-ol. Verasine
[14320-81-1]

C₂₇H₄₃NO 397.643

Alkaloid from *Veratrum album* ssp. *lobelianum*, *Veratrum grandiflorum*, *Veratrum nigrum* (Liliaceae) and *Solanum surinamense*. Shows antifungal activity. Mp 176-178°. [α]_D²⁴ -89.7 (EtOH). [α]_D -91 (c, 0.35 in CHCl₃). λ_{\max} 206 (ϵ 39800); 243 (ϵ 630) (MeOH) (Berdy).

3-O- β -D-Glucopyranoside: **Verazimine**

[97763-02-5]

C₃₃H₅₃NO₆ 559.785

Alkaloid from aerial parts of *Zygadenus sibiricus* (Liliaceae). Cryst. (Me₂CO). [α]_D -112.6 (c, 0.49 in

CHCl₃).

22S,N-Dihydro: 16,28-Secosolanid-5-en-3-ol, 9CI. 22,26-Epiminocholest-5-en-3-ol. **Veramiline**. Oblonginine
[17463-47-7]

C₂₇H₄₅NO 399.659

Alkaloid from *Veratrum album* ssp. *lobelianum* aerial parts (Liliaceae) and *Veratrum oblongum*. Cryst. (Me₂CO). Mp 198-200°. [α]_D²² -49 (c, 0.79 in EtOH). Stereochem. of Oblonginine revised in 1998.

22S,N-Dihydro, 3-O- β -D-glucopyranoside: **Veramiline 3-O- β -D-glucopyranoside**
[128351-76-8]

C₃₃H₅₅NO₆ 561.801

Alkaloid from hypogaeal parts of *Veratrum taliense* (Liliaceae). Powder (EtOAc/MeOH). Mp 303-305°. [α]_D¹⁸ -41.9 (c, 0.6 in MeOH).

22S,N-Dihydro, N,O-di-Ac:

Cryst. (Et₂O). Mp 156-157°. [α]_D²² -21 (c, 0.72 in EtOH).

4 β -Hydroxy: **4 β -Hydroxyverazine**

[215720-25-5]

C₂₇H₄₃NO₂ 413.642

Alkaloid from *Solanum surinamense*. Cryst. (MeOH). Mp 163-165°. [α]_D²⁶ -12.8 (c, 0.7 in MeOH). λ_{\max} 207 (log ϵ 3.4) (MeOH).

12 β -Hydroxy, 22S,N-dihydro: **Veramivirine**12 β -Hydroxyveramiline

[164178-47-6]

C₂₇H₄₅NO₂ 415.658

Alkaloid from roots and rhizomes of *Veratrum viride* (Liliaceae). Needles (MeOH). Mp 229-231°. [α]_D -81 (c, 0.1 in CHCl₃).

25 β -Hydroxy: **25 β -Hydroxyverazine**

[215720-27-7]

C₂₇H₄₃NO₂ 413.642

Alkaloid from *Solanum surinamense*. Cryst. (MeOH). Mp 179-182°. [α]_D²⁶ -28 (c, 0.5 in MeOH). λ_{\max} 211 (log ϵ 3.6) (MeOH).

20-Epimer: **20-Epiverazine**

[145033-50-7]

C₂₇H₄₃NO 397.643

Alkaloid from *Solanum surinamense* and *Veratrum maackii*. Amorph. [α]_D²⁶ +6.5 (c, 1.9 in MeOH). λ_{\max} 206 (log ϵ 4.8) (MeOH).

20-Epimer, Δ^4 -isomer, 3-ketone: **4,5-Dehydro-5,6-dihydro-3-oxo-20-epiverazine**. 16,28-Secosolanida-4,22(28)-dien-3-one
[215804-90-3]

C₂₇H₄₁NO 395.627Alkaloid from *Solanum surinamense*.

Amorph. [α]_D²⁶ +68 (c, 1 in MeOH). λ_{\max} 209 (log ϵ 4.9); 240 (log ϵ 5.1) (MeOH).

20-Epimer, 4 β -hydroxy: **4 β -Hydroxy-20-epiverazine**

[215720-24-4]

C₂₇H₄₃NO₂ 413.642

Alkaloid from *Solanum surinamense*. Powder (MeOH). Mp 107-109°. [α]_D²⁶ -20.5 (c, 2 in MeOH). λ_{\max} 206 (log ϵ 3.4) (MeOH).


20-Epimer, 25 β -hydroxy: **25 β -Hydroxy-20-epiverazine**

[215720-26-6]

C₂₇H₄₃NO₂ 413.642Alkaloid from *Solanum surinamense*.Adam, G. *et al.*, *Tetrahedron*, 1967, **23**, 167-171 (*isol, struct, ir, ms, uv, pmr*)Kessar, S.V. *et al.*, *Indian J. Chem.*, 1974, **12**, 1245-1248 (*synth*)Vassová, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 3643-3645 (*Veramiline*)Bondarenko, N.V. *et al.*, *Khim. Prir. Soedin.*, 1979, 415; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 366 (*isol*)Tomko, J. *et al.*, *CA*, 1981, **94**, 205391r (*isol*)Taskhanova, G.M. *et al.*, *Khim. Prir. Soedin.*, 1985, **21**, 368-369; *Chem. Nat. Compd. (Engl. Transl.)*, 1985, **21**, 343-344 (*Verazine*)Kusano, G. *et al.*, *Chem. Pharm. Bull.*, 1987, **35**, 4862-4867 (*activity*)Mizuno, M. *et al.*, *Phytochemistry*, 1990, **29**, 359-361 (*Veramiline 3-O-β-D-glucopyranoside*)Kadota, S. *et al.*, *Phytochemistry*, 1995, **38**, 777-781 (*Oblonginine*)El Sayed, K.A. *et al.*, *Phytochemistry*, 1995, **38**, 1547-1550 (*Veramivirine*)Abdel-Kader, M.S. *et al.*, *J. Nat. Prod.*, 1998, **61**, 1202-1208; 2000, **63**, 1184 (*Solanum surinamense alkaloids*)Lowe, P.R. *et al.*, *Phytochemistry*, 1998, **47**, 887-890 (*Oblonginine, stereochem*)**Verdine**

V-74

17,23-Epoxy-5,6-dihydro-1,3,6-trihydroxyveratraman-11-one. 1,3,6-Trihydroxyjervanin-12-en-11-one [73667-53-5]

C₂₇H₄₁NO₅ 459.625Alkaloid from *Veratrum lobelianum* epigeal parts (Liliaceae). Mp 218-220°.Nakhatov, I. *et al.*, *Khim. Prir. Soedin.*, 1984, 395; *Chem. Nat. Compd. (Engl. Transl.)*, 375 (*isol, uv, ir, pmr*)Tashkhodzhaev, B. *et al.*, *Khim. Prir. Soedin.*, 1984, 753; *Chem. Nat. Compd. (Engl. Transl.)*, 711 (*cryst struct*)**Verine**


V-75

[1361-89-3]

C₂₅H₃₉NO₂ 385.589Steroidal alkaloid. Struct. unknown. Alkamine from *Veratrum album* (Liliaceae). Mp 221-222° dec. [α]_D¹⁸ -21.12 (c, 1 in CHCl₃).Cionga, E. *et al.*, *Acta Pol. Pharm.*, 1957, **14**, 73-76; *CA*, **52**, 12882c**Vermelhotin**

V-76

3-[6-(1-Propenyl)-2H-pyran-2-ylidene]-2,4-pyrrolidinedione


C₁₂H₁₁NO₃ 217.224

Tetramic acid. *Isol.* from an unidentified fungus, IFM 52672. Red needles (CH₂Cl₂/MeOH). Mp 212-214°. λ_{max} 228 (log ε 3.82); 234 (log ε 3.8); 276 (log ε 4.11); 319 (log ε 3.5); 332 (log ε 3.4); 439 (log ε 3.96); 456 (log ε 3.95) (MeOH).

Hosoe, T. *et al.*, *Heterocycles*, 2006, **68**, 1949-1953 (*isol, pmr, cmr, cryst struct*)**Vermisporin**

V-77

[122301-98-8]



Relative Configuration


C₂₅H₃₇NO₄ 415.572

Tetramic acid deriv. Similar to Antibiotic PF 1052, A-1225. Prod. by *Ophiobolus vermisporus* and *Phoma* sp. FK1-1840. Antibacterial agent. Cryst. Sol. MeOH, Et₂O; poorly sol. hexane, H₂O. [α]_D²⁰ +73.8 (CHCl₃). [α]_D +55.9 (MeOH). λ_{max} 229 (ε 6180); 291 (ε 12280) (MeOH).

Japan. Pat., 1988, 190 894; *CA*, **111**, 95600t (*isol*)Koyama, N. *et al.*, *J. Antibiot.*, 2005, **58**, 338-345 (*isol, struct*)**Verongamine**

V-78

[150036-88-7]

C₁₅H₁₇BrN₄O₃ 381.228

Alkaloid from the marine sponge *Verongula gigantea*. Specific histamine H₃-receptor antagonist. Yellow semi-solid or oil. λ_{max} 206 (ε 33600); 280 (ε 3040); 388 (ε 630) (MeOH) (Berdy).


5-Bromo- 5-Bromoverongamine

[218931-98-7]

C₁₅H₁₆Br₂N₄O₃ 460.124Metab. of the sponge *Pseudoceratina* sp. Shows antifouling props. Oil. λ_{max} 207 (ε 44140) (MeOH).Mierzwa, R. *et al.*, *J. Nat. Prod.*, 1994, **57**, 175-177 (*isol, uv, ir, pmr, cmr, struct*)Wasserman, H.H. *et al.*, *J.O.C.*, 1998, **63**, 5581-5586 (*synth*)Thirionet, I. *et al.*, *Nat. Prod. Lett.*, 1998, **12**, 209-214 (5-Bromoverongamine, *activity*)Boehlow, T.R. *et al.*, *J.O.C.*, 2001, **66**, 3111-3118 (*synth*)**Verpacamide C**

V-79

3-(Aminoiminoethyl)-1,2,3,7,8,9a-hexahydro-5H,10H-dipyrrolo[1,2-a:1',2'-d]pyrazine-5,10-dione [890152-49-5]



Relative Configuration


C₁₁H₁₅N₅O₂ 249.272

Alkaloid from *Axinella vaceleti*. Amorph. solid (as formate salt). [α]_D²⁴ +17.4 (c, 0.15 in H₂O) (formate salt).

Vergne, C. *et al.*, *Org. Lett.*, 2006, **8**, 2421-2424 (*isol, pmr, cmr*)**Verpacamide D**

V-80

[890152-48-4]



Relative Configuration

C₁₂H₁₉N₅O₄ 297.313

Alkaloid from *Axinella vaceleti*. Amorph. solid (as formate salt).

Vergne, C. *et al.*, *Org. Lett.*, 2006, **8**, 2421-2424 (*isol, pmr, cmr*)**Verrucarin F**

V-81

C₂₁H₁₈N₂O₆ 394.383

Pyrrole antibiotic. Struct. unknown. Alkaloid from *Myrothecium verrucaria*. Cryst. (Me₂CO/Et₂O or MeOH). Sol. Py; fairly sol. MeOH, butanol; poorly sol. CHCl₃, hexane. Mp 237-238°. [α]_D²³ -1 (c, 0.64 in Py). λ_{max} 202 ; 233 ; 308 (EtOH).

Haerri, E. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 839-853 (*isol*)**Verrucarin G**

V-82

C₁₅H₁₂N₂O₆ 316.27


Pyrrole antibiotic. Struct. unknown. Alkaloid from *Myrothecium verrucaria*. Cryst. (Me₂CO/Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 118° Mp 131-135° double Mp). λ_{max} 208 ; 254 ; 300 (EtOH).

Haerri, E. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 839-853**Verrucine A**

V-83

[252032-22-7]

[366787-64-6 (Verrucine D), 366787-58-8 (Verrucine E), 366786-27-8 (Verrucine C)]



Absolute
Configuration

$C_{21}H_{20}N_4O_3$ 376.414

Related to Fumiquinazoline A, F-185 and Anacine, A-966. Prod. by *Penicillium verrucosum*. Amorph. solid. $[\alpha]_D^{25} +37$ (c, 0.1 in EtOH) (natural). $[\alpha]_D^{30} +57$ (c, 0.25 in EtOH) (synthetic). Verrucine C, Verrucine D and Verrucine E have also been isol. but not characterized. λ_{max} 227 (sh) (log ϵ 3.65); 272 (log ϵ 3.17); 278 (log ϵ 3.17); 306 (log ϵ 2.82); 317 (log ϵ 2.68) (EtOH).

3-Epimer: Verrucine B

[252032-23-8]

$C_{21}H_{20}N_4O_3$ 376.414

Prod. by *Penicillium verrucosum*.

Amorph. solid. $[\alpha]_D^{22} +124$ (c, 0.08 in EtOH) (natural). $[\alpha]_D^{29} +183$ (c, 0.19 in EtOH) (synthetic). λ_{max} 227 (sh) (log ϵ 3.56); 272 (log ϵ 2.9); 278 (log ϵ 2.87); 306 (log ϵ 2.69); 317 (log ϵ 2.65) (EtOH).

Larsen, T.O. *et al.*, *J. Nat. Prod.*, 1999, **62**, 1578-1580 (isol, pmr, cmr)


Larsen, T.O. *et al.*, *Appl. Environ. Microbiol.*, 2001, **67**, 3630-3635 (isol)

Wang, H. *et al.*, *J. Nat. Prod.*, 2001, **64**, 1497-1501 (synth, pmr, cmr, abs config)

Verruculotoxin

V-84

Hexahydro-3-(phenylmethyl)-2H-pyrido[1,2-a]pyrazin-1(6H)-one. 3-Benzyl-1,4-diazabicyclo[4.4.0]decan-5-one [56092-63-8]



Absolute
Configuration

$C_{15}H_{20}N_2O$ 244.336

Prod. by *Penicillium verrucosum*. Mycotoxin. Large needles (CH₂Cl₂/heptane), cryst. (C₆H₆). Sol. Me₂CO, Et₂O; fairly sol. MeOH, toluene; poorly sol. H₂O, hexane. Mp 152°. $[\alpha]_D^{25} -56$ (c, 0.25 in MeOH) (synthetic).

▶ LD₅₀ (ckn, orl) 20 mg/kg; LD₅₀ (mus, orl) 20 mg/kg. YX9840000

10-Epimer: 10-Epiverruculotoxin

[69651-21-4]

$C_{15}H_{20}N_2O$ 244.336

Prod. by *Penicillium brasilianum*. Powder (EtOAc/hexane). Mp 143-144° (synthetic). $[\alpha]_D^{25} +59.6$ (c, 0.5 in MeOH) (synthetic).

Cole, R.J. *et al.*, *Toxicol. Appl. Pharmacol.*, 1975, **31**, 465-468 (isol, uv, ir, pmr, ms)

MacMillan, J.G. *et al.*, *J.A.C.S.*, 1976, **98**, 246-247 (synth, cryst struct)

Frisvad, J.C. *et al.*, *J. Chromatogr.*, 1987, **392**, 333-347; **404**, 195-214 (chromatog)

Williams, R.M. *et al.*, *Synthesis*, 1988, 963-966 (synth, ir, pmr)


Martens, J. *et al.*, *Tet. Lett.*, 1991, **32**, 1417-1418 (synth)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 838

Versimide

V-85

Methyl 3-methyl- α -methylene-2,5-dioxo-1-pyrrolidineacetate, 9CI



(R)-form

$C_9H_{11}NO_4$ 197.19

(R)-form [30270-17-8]

Metab. of *Aspergillus versicolor*. Has contact insecticidal and antibiotic props. Viscous oil. Sol. MeOH, Et₂O; poorly sol. H₂O, hexane. Bp_{0.003} 60°. $[\alpha]_D^{20} +22.3$ (c, 2 in CHCl₃). n_D^{21} 1.4896. λ_{max} 204 (ϵ 10260) (EtOH) (Berdy).

(±)-form

Yellow oil.

Brown, A.G. *et al.*, *J.C.S. (C)*, 1970, 2572 (isol, struct, ir, pmr)


Cole, M. *et al.*, *Appl. Microbiol.*, 1972, **24**, 660 (use)

Brown, A.G. *et al.*, *J.C.S. Perkin 1*, 1972, 65 (synth, ir, pmr)

Vertaline

V-86

(10 α)-Decaline, 9CI. 10-Epidecaline



$C_{26}H_{31}NO_5$ 437.535

(-)-form [32886-91-2]

Alkaloid from *Decodon verticillatus* (Lythraceae). Mp 194-196°. $[\alpha]_D^{26} -170$ (c, 1.25 in CHCl₃).

Hydrobromide:

Cryst. (MeOH). Mp 275-278°.

Methiodide:

Cryst. + 1H₂O (THF/CH₂Cl₂). Mp 221°.

O²³-De-Me: Demethylvertaline

[34437-41-7]

$C_{25}H_{29}NO_5$ 423.508

Alkaloid from *Decodon verticillatus* (Lythraceae). Mp 120-160°.

5-Epimer: Decaline. Epivertaline

[14727-56-1]

$C_{26}H_{31}NO_5$ 437.535

Alkaloid from *Decodon verticillatus*

(Lythraceae). Mp 81-82° (MeOH solvate) Mp 102-118° (dry).

5-Epimer, methiodide:

Cryst. (THF/CH₂Cl₂). Mp 215°.

5-Epimer, O²³-de-Me: Demethyldecaline

[34437-42-8]

$C_{25}H_{29}NO_5$ 423.508

Alkaloid from *Decodon verticillatus* (Lythraceae).

(±)-form [53494-86-3]

Synthetic. Prisms (MeOH). Mp 224-225°.

5-Epimer: [50412-65-2]

Synthetic. Needles (MeOH). Mp 196-197°.

5-Epimer, O²³-de-Me: [54422-29-6]

Synthetic. Mp 259-261°.

Ferris, J.P. *et al.*, *J.O.C.*, 1962, **27**, 2985 (isol, uv)

Hamilton, J.A. *et al.*, *J.A.C.S.*, 1971, **93**, 2939 (cryst struct)

Ferris, J.P. *et al.*, *J.A.C.S.*, 1971, **93**, 2953

(Vertaline, Decaline, Demethyldecaline, Demethylvertaline, isol, pmr, struct, synth)

Wróbel, J.T. *et al.*, *Tet. Lett.*, 1973, 4293 (Decaline, synth)

Hanaoka, M. *et al.*, *Chem. Pharm. Bull.*, 1974, **22**, 973; 1945; 1975, **23**, 2140; 1976, **24**, 1045 (synth, ir, pmr, uv)

Corey, E.J. *et al.*, *J.A.C.S.*, 1975, **97**, 654 (synth)


Hart, D.H. *et al.*, *J.O.C.*, 1982, **47**, 1555 (synth, ir, pmr)

Shishido, K. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 532 (synth, ir, pmr)

Vertaline B

V-87

[118985-28-7]



Absolute
configuration

$C_{27}H_{45}NO_3$ 431.657

Structurally unrelated to Vertaline. Alkaloid from the whole plant of *Veratrum taliense* (Liliaceae).

16-Deoxy: Stenophylline B

[91421-75-9]

$C_{27}H_{45}NO_2$ 415.658

Alkaloid from whole plants of *Veratrum stenophyllum*. Cryst. (MeOH). Mp 225-228°.

16-Deoxy, 3-O- β -D-glucopyranoside: Stenophylline B 3-O- β -D-glucopyranoside

[128351-77-9]

$C_{33}H_{55}NO_7$ 577.8

Alkaloid from *Veratrum taliense*. Plates (MeOH). Mp 286-288°. $[\alpha]_D^{18} -42.5$ (c, 0.1 in MeOH).

Liang, G. *et al.*, *Yaoxue Xuebao*, 1984, **19**, 131-136; *CA*, **101**, 97524p (Stenophylline B)


Min, Z.D. *et al.*, *Yaoxue Xuebao*, 1988, **23**, 584-587; *CA*, **110**, 92057q (Vertaline B)

Mizuno, M. *et al.*, *Phytochemistry*, 1990, **29**, 359-361 (Stenophylline B 3-glucoside)

Verticillatine†

V-88

2',6''-Dihydroxy-5''-methoxylythran-12-one, 9CI
[10247-54-8]



$C_{25}H_{27}NO_5$ 421.492

CAS numbering differs from that shown. Alkaloid from *Decodon verticillatus* (Lythraceae). Mp 312°.

Hydrochloride:

Cryst. + 0.5 MeOH (MeOH/Et₂O). Mp 240° dec. $[\alpha]_D^{26}$ +119 (c, 1.93 in MeOH).

Di-O-Ac:

Cryst. (cold MeOH). Mp 176-178°. $[\alpha]_D^{26}$ +36 (c, 2.41 in CHCl₃).

O²¹-Me: Sinicuichine

[31946-81-3]
 $C_{26}H_{29}NO_5$ 435.519

Alkaloid from *Heimia salicifolia* (Lythraceae). Tranquilliser. Mp 187-188° (in vacuo). $[\alpha]_D$ +77.8 (CHCl₃). Log P 4.17 (calc).

O²¹-Me; hydrochloride: Mp 265-266°.**Di-Me ether:** Mp 252° (in vacuo).**12,13-Dihydro: Dihydroverticillatine**

[10215-02-8]
 $C_{25}H_{29}NO_5$ 423.508

Alkaloid from *Lagerstroemia indica* (Lythraceae). Cryst. (MeOH). Mp 260-263° dec.

12,13-Dihydro, O²¹-Me: Lagerstroemine.

Indicamine†. *Lagerstroemine*

[10247-53-7]

$C_{26}H_{31}NO_5$ 437.535

Alkaloid from *Lagerstroemia indica*. Also detected in *Plantago psyllium* seeds (African plantain) (Lythraceae, Plantaginaceae). Cryst. (EtOH). Mp 240°. $[\alpha]_D$ -137 (CHCl₃). pK_a 7.7 (EtOH).

12,13-Dihydro, di-Me ether:

Cryst. (Me₂CO/Et₂O). Mp 187-188°. $[\alpha]_D^{26}$ -99 (c, 1.13 in CHCl₃).

5-Epimer: Dehydrodecodine

[35323-19-4]
 $C_{25}H_{27}NO_5$ 421.492

Alkaloid from *Heimia salicifolia* (Lythraceae). Pale yellow powder. Mp 181-183°. $[\alpha]_D^{20}$ +149.3 (c, 0.15 in MeOH). Dec. above 184° if Mp. is determined in vacuo. Stereoisomer of Verticillatine, V-88. λ_{max} 285 (MeOH).

5-Epimer, O²¹-Me: Nesodine

[34628-60-9]
 $C_{26}H_{29}NO_5$ 435.519

Alkaloid from *Heimia salicifolia* (Lythraceae). Needles (CH₂Cl₂ or 2-propanol). Mp 190°.

5-Epimer, 12,13-dihydro: Decodine

[26996-01-0]

$C_{25}H_{29}NO_5$ 423.508

Alkaloid from *Decodon verticillatus* (Lythraceae). Cryst. (MeOH). Mp 193-197°. $[\alpha]_D^{26}$ -97 (c, 1.78 in CHCl₃).

5-Epimer, 12,13-dihydro, di-Ac:

Cryst. (Et₂O). Mp 202-203°. $[\alpha]_D^{26}$ -66 (c, 1.18 in CHCl₃).

Ferris, J.P. et al., *J.O.C.*, 1962, **27**, 2985; 1963, **28**, 817 (isol, uv, pmr, Verticillatine, Dihydroverticillatine, Lagerstroemine, Decodine)

Blomster, R.N. et al., *J. Nat. Prod.*, 1964, **27**, 15 (isol, uv, ir, struct, Sinicuichine)

Appel, H. et al., *J. Nat. Prod.*, 1965, **28**, 84 (isol)

Hörhammer, R.B. et al., *J. Nat. Prod.*, 1970, **33**, 483 (pmr, struct, Sinicuichine)

Ferris, J.P. et al., *J.A.C.S.*, 1971, **93**, 2942; 2958 (Lagerstroemine, Dihydroverticillatine, Nesodine, Decodine)

Balboa, S.I. et al., *U. A. R. J. Pharm. Sci.*, 1971, **12**, 35; *CA*, **77**, 156311c (Lagerstroemine, occur)

Karawya, M.S. et al., *U. A. R. J. Pharm. Sci.*, 1971, **12**, 53; *CA*, **78**, 13729e (Lagerstroemine, occur)


Hörhammer, R.B. et al., *Z. Naturforsch., B*, 1971, **26**, 970 (Dehydrodecodine)

Rumalla, C.S. et al., *Phytochemistry*, 2008, **69**, 1756-1762 (Dehydrodecodine)

Verticillin A†

V-89

[32164-16-2]



Absolute Configuration

$R^1 = R^2 = CH_3$

$C_{30}H_{28}N_6O_6S_4$ 696.852

Epidithiodioxopiperazine-type antibiotic. Isol. from *Verticillium* sp. TM759, *Gliocladium* sp. SCF1168 and a marine *Penicillium* sp. CNC-350. Mycotoxin. Cytotoxin. Yellow needles (Py). Sol. MeOH, dioxan, CHCl₃, Py; poorly sol. H₂O, hexane. Mp 202-217° dec. $[\alpha]_D$ +703.7. $[\alpha]_D$ +727 (dioxan). λ_{max} 203 (ε 50100); 240 (sh) (ε 14100); 301 (ε 5370) (Ac₂O/MeOH) (Derep). λ_{max} 306 (ε 6300) (dioxan) (Berdy).

▶ LD₅₀ (mus, ipr) 7.6 mg/kg. YY1830000**6-Deoxy: 11-Deoxyverticillin A**

[240480-95-9]
 $C_{30}H_{28}N_6O_5S_4$ 680.852

Isol. from a marine *Penicillium* sp. CNC-350. Cytotoxic. Yellow cryst. (CH₂Cl₂/MeOH).

6,6'-Dideoxy: 11,11'-Dideoxyverticillin A

[240480-94-8]
 $C_{30}H_{28}N_6O_4S_4$ 664.853

Isol. from a marine *Penicillium* sp. CNC-350. Cytotoxic. Solid. $[\alpha]_D$

+624.1.

19-Hydroxy: Verticillin B

[52212-86-9]
 $C_{30}H_{28}N_6O_7S_4$ 712.851

From *Verticillium* sp. TM759. Pale-yellow prisms (CHCl₃). Sol. MeOH, CHCl₃, dioxan, Py; poorly sol. H₂O, Et₂O, hexane. Mp 254-256° dec. $[\alpha]_D$ +704.7 (c, 0.493 in dioxan). λ_{max} 306 (ε 5600) (dioxan) (Berdy).

Mono (trisulfide) homologue: Gliocladine A†

[871335-05-6]
 $C_{30}H_{28}N_6O_6S_5$ 728.918

Not to be confused with Gliocladine A, G-97. Prod. by *Gliocladium roseum* 1A. Amorph. powder. $[\alpha]_D^{18}$ +553.6 (c, 0.45 in Py). λ_{max} 202 (log ε 0.7); 250 (log ε 0.37); 256 (log ε 0.29); 261 (log ε 0.31) (Py).

Mono (tetrasulfide) homologue: Gliocladine B†

[871335-06-7]
 $C_{30}H_{28}N_6O_6S_6$ 760.984

Not to be confused with Gliocladine B in G-97. Prod. by *Gliocladium roseum* 1A. Amorph. powder. $[\alpha]_D^{18}$ +555.8 (c, 0.44 in Py). λ_{max} 202 (log ε 0.81); 250 (log ε 0.57); 256 (log ε 0.59); 261 (log ε 0.42) (Py).

Katagiri, K. et al., *J. Antibiot., Ser. B*, 1970, **23**, 420-422 (Verticillin A)

Minato, H. et al., *J.C.S. Perkin 1*, 1973, 1819-1825 (Verticillin B)

Son, B.W. et al., *Nat. Prod. Lett.*, 1999, **13**, 213-222 (Deoxyverticillin A, Dideoxyverticillin A)

Dong, J.-Y. et al., *J. Nat. Prod.*, 2005, **68**, 1510-1513 (Gliocladine A, B)

Liu, F. et al., *Acta Cryst. E*, 2006, **62**, 974-976 (cryst struct)

Cole, R.J. et al., *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 606

Verticillin C

V-90

As Verticillin A, V-89 with
 $R^1 = SME, R^2 = CH_2OH$

$C_{30}H_{28}N_6O_7S_5$ 744.917

From *Verticillium* sp. strain TM-759. Shows antibiotic props. Pale-yellow amorph. powder (MeOH aq.). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 230-235° dec. $[\alpha]_D$ +765 (c, 0.506 in dioxan). λ_{max} 303 (ε 5500) (dioxan) (Berdy).

Minato, H. et al., *J.C.S. Perkin 1*, 1973, 1819 (isol, ir, nmr)

Verticillin D

V-91

As Verticillin A, V-89 with
 $R^1 = R^2 = -^{13}CH(OH)CH_3$

$C_{32}H_{32}N_6O_8S_4$ 756.904

Epidithiodioxopiperazine-type antibiotic. Prod. by *Gliocladium catenulatum*. Powder. Mp 244-247°. $[\alpha]_D$ +220 (c, 0.1 in MeOH). λ_{max} 214 (ε 19000); 304 (ε 1900) (MeOH).


13-Ketone: Verticillin F

$C_{32}H_{30}N_6O_8S_4$ 754.888


Prod. by *Gliocladium catenulatum*. Yellow powder. Mp 221-224°. $[\alpha]_D$ -80 (c, 0.1 in MeOH). λ_{max} 214 (ε 18000); 302 (ε 1400) (MeOH).

13,13'-Diketone: Verticillin EC₃₂H₂₈N₆O₈S₄ 752.873Prod. by *Gliocladium catenulatum*.Yellow powder. Mp 211-214°. [α]_D²⁵ +440 (c, 0.1 in MeOH). λ_{max} 218 (ε 26000); 304 (ε 4400) (MeOH).Joshi, B.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 730-733 (*isol, uv, pmr, cmr*)**Verticillin G**

[933785-95-6]

V-92C₃₀H₂₈N₆O₇S₄ 712.851Prod. by *Bionectra byssicola* F120. Antibacterial agent. Powder. [α]_D²⁵ +467.6 (c, 0.2 in MeOH). λ_{max} 215 (log ε 4.79); 301 (log ε 3.79) (MeOH).Zheng, C.-J. *et al.*, *J. Antibiot.*, 2007, **60**, 61-64 (*isol, pmr, cmr*)**Verticilline†****V-93**C₁₉H₃₃N₂O₂ 307.475Steroidal alkaloid. Struct. unknown. Alkaloid from *Fritillaria verticillata* var. *thunbergii* (Liliaceae). Antihypertensive agent. Smooth muscle relaxant. Mp 148-150°. Sinters from 130°; resolidifies and remelts at 212-213°. Co-occurs with Verticine in C-298.Fukuda, M. *et al.*, *Nippon Kagaku Zasshi*, 1929, **50**, 74-78; *CA*, **23**, 2930Ito, S. *et al.*, *Chem. Pharm. Bull.*, 1963, **11**, 1337-1340**Vertilecanine A****V-94**


5-(Hydroxyphenylmethyl)-2-pyridinecarboxylic acid. 5-(α-Hydroxybenzyl)-2-pyridinecarboxylic acid

C₁₃H₁₁N₃ 229.235**(R)-form**Isol. from *Verticillium lecanii*. Solid. Mp 155-157°. [α]_D²⁵ -11 (c, 0.04 in MeOH). λ_{max} 208 (ε 5800); 228 (ε 1900); 270 (ε 1100) (MeOH).**Me ester:**C₁₄H₁₃N₃O₃ 243.262Isol. from *Verticillium lecanii*. Solid. [α]_D²⁴ -7 (c, 0.02 in MeOH). λ_{max} 210 (ε 2400); 228 (ε 1700); 270 (ε 1000)

(MeOH).

Glycine amide: Vertilecanine BC₁₅H₁₄N₂O₄ 286.287Isol. from *Verticillium lecanii*. Solid. [α]_D²⁴ +13 (c, 0.03 in MeOH). λ_{max} 210 (ε 3300); 228 (ε 2500); 270 (ε 1500) (MeOH).**Glycine amide, Me ester:**C₁₆H₁₆N₂O₄ 300.313Isol. from *Verticillium lecanii*. Solid. Mp 119-121°. [α]_D²⁵ +9.7 (c, 0.03 in MeOH). λ_{max} 210 (ε 3600); 232 (ε 2900); 270 (ε 1600) (MeOH).**7-Ketone, glycine amide, Me ester: Vertilecanine C**C₁₆H₁₄N₂O₄ 298.298Isol. from *Verticillium lecanii*. Oil. λ_{max} 210 (ε 5800); 254 (ε 4800); 272 (ε 4400) (MeOH).Soman, A.G. *et al.*, *J. Nat. Prod.*, 2001, **64**, 189-192 (*isol, pmr, cmr*)Tumer, F. *et al.*, *Russ. Chem. Bull. (Engl. Transl.)*, 2005, **54**, 2466 (*synth*)**Vertine**

[10308-13-1]

V-95

Absolute Configuration

C₂₆H₂₉NO₅ 435.519CAS numbering is not the same as that shown. Alkaloid from *Decodon verticillatus*, *Heimia salicifolia*, *Heimia myrtifolia* and *Lagerstroemia fauriei* (Lythraceae). Antiinflammatory, hyperglycaemic and hypotensive agent. Sedative. Cryst. (MeOH/CHCl₃). Mp 250-252° (245-247°). [α]_D²⁵ +69.9 (c, 0.1 in MeOH). [α]_D²⁰ +61 (c, 1 in CHCl₃). λ_{max} 285 (MeOH).

▶OL6150000

Hydrochloride: Mp 322-323°.**Ac:**Cryst. (Et₂O). Mp 183-185°. [α]_D²⁶ -92.7 (c, 0.96 in CHCl₃).**Me ether:** Mp 231-233° (225-228°).**O²³-De-Me: 5-Epilyfoline. 10-Epilyfoline.****Demethylvertine**

[128502-80-7]

C₂₅H₂₇NO₅ 421.492Alkaloid from leaves and stems of *Heimia montana*. Yellow powder. [α]_D²⁰ -75.9 (c, 0.15 in MeOH). λ_{max} 285 (MeOH).**12,13-Dihydro: Decamine. Dihydrovertine**

[17349-05-2]

C₂₆H₃₁NO₅ 437.535Alkaloid from *Decodon verticillatus* and *Lagerstroemia indica* (Lythraceae). Shows fungicidal props. Cryst. (MeOH). Mp 223-224°. [α]_D²⁶ -145 (c, 1.0 in CHCl₃). Log P 2.89 (calc).**12,13-Dihydro, Ac:**Cryst. (cold MeOH). Mp 197-198°. [α]_D²⁶ -188 (c, 1.16 in CHCl₃). Deacetylated by hot MeOH.**12,13-Dihydro, 13ξ-hydroxy: Heimidine.****Hydroxydihydrovertine**

[36531-11-0]

C₂₆H₃₁NO₆ 453.534Minor alkaloid from *Heimia salicifolia* (Lythraceae). Mp 221-223°. [α]_D²⁰ -101.3 (c, 0.15 in MeOH). λ_{max} 285 (MeOH).**12,13-Dihydro, 13ξ-hydroxy, hydrochloride:**

Mp 316-318°.

6R-Hydroxy: 6-Hydroxyvertine. 9-HydroxyvertineC₂₆H₂₉NO₆ 451.518Alkaloid from the leaves of *Heimia salicifolia*. Powder. [α]_D²⁰ +12 (c, 0.15 in MeOH). λ_{max} 283 (MeOH).**5-Epimer: Lythrine**

[5286-10-2]

C₂₆H₂₉NO₅ 435.519Alkaloid from *Heimia salicifolia*, *Heimia myrtifolia* and *Lagerstroemia fauriei* (Lythraceae). Cryst. (MeOH). Mp 243-245°. [α]_D²⁰ +32.5 (CHCl₃). [α]_D²⁰ +40.6 (CHCl₃).**5-Epimer, hydrochloride:** Mp 325-331°.**5-Epimer, Ac:** Mp 172-173°.**5-Epimer, O²³-de-Me: Lyfoline**

[30356-01-5]

C₂₅H₂₇NO₅ 421.492Alkaloid from *Heimia salicifolia*. Yellow powder. Mp 223-224°. [α]_D²⁰ +70.6 (c, 0.15 in MeOH). λ_{max} 281 (MeOH).**5-Epimer, 12,13-dihydro: Decinine. Dihydrolythrine. Cryogenine†**

[10183-64-9]

C₂₆H₃₁NO₅ 437.535Alkaloid from *Decodon verticillatus*, *Lagerstroemia indica* and *Lythrum lanceolatum* (Lythraceae). Antiinflammatory agent, shows diuretic activity. Cryst. (MeOH). Mp 222-224°. [α]_D²⁶ -142 (c, 1.37 in CHCl₃). pK_a 7.4 (50% MeOH).**5-Epimer, 12,13-dihydro, Ac:** Mp 197-198°.**5-Epimer, 12,13-dihydro, 13S-hydroxy:****Lythridine. Sinine**

[15299-77-1]

C₂₆H₃₁NO₆ 453.534Alkaloid from *Heimia salicifolia*, *Heimia myrtifolia* and *Lagerstroemia fauriei* (Lythraceae). Cryst. (EtOAc). Mp 218-219°. [α]_D²⁰ -174.2 (CHCl₃). [α]_D²⁰ +49.3 (c, 0.15 in MeOH). λ_{max} 280 (MeOH).**5-Epimer, 12,13-dihydro, 13S-hydroxy, hydrochloride:** Mp 330° dec.**Stereoisomer (1): Alkaloid ALC 1**

[57572-82-4]


C₂₆H₂₉NO₅ 435.519Alkaloid from *Heimia salicifolia* (Lythraceae). Mp 335-345° dec. [α]_D²² +115.6 (CHCl₃). Stereochem. undetermined.**Stereoisomer (1), Ac:** Mp 126-128°.**Stereoisomer (1), Me ether:** Mp 230-233°.

Stereoisomer (2): Alkaloid ALC 2
[57572-83-5]
C₂₆H₂₉NO₅ 435.519
Alkaloid from *Heimia salicifolia*. Mp 309-310°. [α]_D +72.3 (CHCl₃). Stereochem. undetermined.

Stereoisomer (2), Me ether: Mp 235-237°.

Ferris, J.P. *et al.*, *J.O.C.*, 1962, **27**, 2985 (isol, uv, ir, synth, Vertine, Decamine, Decinine)
Blomster, R.N. *et al.*, *J. Nat. Prod.*, 1964, **27**, 15 (Vertine, Lythrine, Lythridine)
Douglas, B. *et al.*, *J. Nat. Prod.*, 1964, **27**, 25 (Vertine, Lythrine, Sinine)
Zacharias, D.E. *et al.*, *Experientia*, 1965, **21**, 247 (cryst struct, Lythrine)
Appel, H. *et al.*, *J. Nat. Prod.*, 1965, **28**, 84 (isol, Lyfoline)
Chu, S.C. *et al.*, *Chem. Ind. (London)*, 1966, 1795 (cryst struct, Sinine)
Appel, H. *et al.*, *Tet. Lett.*, 1966, 5789 (Sinine, uv, ms)
Ferris, J.P. *et al.*, *J.A.C.S.*, 1971, **93**, 2942; 2958; 2963 (isol, abs config, Vertine, Lythrine, Lyfoline, Decinine)
Wright, H. *et al.*, *J.A.C.S.*, 1973, **95**, 6467 (occur, Decinine)
Dominguez, X.A. *et al.*, *Phytochemistry*, 1975, **14**, 1883 (isol, ord, ir, pmr, ms, struct, Lythrine)
Lantos, I. *et al.*, *J.O.C.*, 1977, **42**, 228 (synth, ir, pmr, ms, Decamine, Decinine)
Fuji, K. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 2515 (Vertine, Sinine, isol, uv, ir)
Hedges, S.H. *et al.*, *Chem. Comm.*, 1983, 145 (biosynth)
Rother, A. *et al.*, *Phytochemistry*, 1990, **29**, 1683 (Demethylvertine)
Xie, X.-Q. *et al.*, *J. Nat. Prod.*, 1995, **58**, 1876 (10-Epilyfoline)
Rumalla, C.S. *et al.*, *Phytochemistry*, 2008, **69**, 1756-1762 (isol, pmr, cmr, cryst struct, 6-Hydroxyvertine, Vertine, 5-Epilyfoline, Heimidine, Lyfoline, Lythridine)

Vibriobactin **V-96**
[88217-23-6]



C₃₅H₃₉N₅O₁₁ 705.72
Isol. from *Vibrio cholerae*. Siderophore. Powder. Mp 153-155°. Darkens at ca. 130°. MF given as C₃₅H₄₁N₅O₁₁ in paper. λ_{\max} 252 ; 316 (MeOH) (Berdy).


3',3''-Dideoxy: Vulnibactin
[156368-88-6]

C₃₅H₃₉N₅O₉ 673.721
Isol. from *Vibrio vulnificus*. Siderophore. Mp 94-97°. [α]_D²⁵ +92.5 (c, 0.80 in MeOH).

Griffiths, G.L. *et al.*, *J. Biol. Chem.*, 1984, **259**, 383 (isol, struct)
Bergeron, R.J. *et al.*, *Tetrahedron*, 1985, **41**, 507 (synth)
Okujo, N. *et al.*, *BioMetals*, 1994, **7**, 109 (Vulnibactin)

Bergeron, R.J. *et al.*, *Synthesis*, 2007, 1033-1037 (Vulnibactin, synth)


Vibrioferriin **V-97**
[157568-17-7]



C₁₆H₂₂N₂O₁₂ 434.356
Prod. by *Vibrio parahaemolyticus* and a *Marinobacter* sp. Siderophore. Oil. Sol. H₂O, MeOH, Me₂CO, MeCN, DMSO; fairly sol. Et₂O; poorly sol. EtOAc, CHCl₃, [α]_D²⁰ +13.3 (c, 1.4 in MeOH). λ_{\max} 200 (ϵ 4000) (H₂O).

Yamamoto, S. *et al.*, *J. Biochem. (Tokyo)*, 1994, **115**, 868-874 (isol, struct)
Yamamoto, S. *et al.*, *Microbiol. Immunol.*, 1994, **38**, 687-693 (isol, activity)
Takeuchi, Y. *et al.*, *Chem. Pharm. Bull.*, 1999, **47**, 459-460; 1284-1287 (synth)
Amin, S.A. *et al.*, *J.A.C.S.*, 2007, **129**, 478-479 (isol, props)

Vicenistatin aglycone **V-98**



Absolute configuration


C₂₃H₃₅NO₂ 357.535
O-(2,4,6-Trideoxy-4-methylamino- β -D-ribo-hexopyranoside): **Vicenistatin**
[150999-05-6]
C₃₀H₄₈N₂O₄ 500.72
Prod. by *Streptomyces halstedii* HC34. Antitumour agent. Powder. Sol. MeOH, Me₂CO, CHCl₃; poorly sol. H₂O, hexane. Mp 151-153°. [α]_D²³ -3 (c, 0.1 in MeOH). λ_{\max} 235 (ϵ 35000); 240 (ϵ 35800); 268 (ϵ 13700) (MeOH).

O-(2,6-Dideoxy-3-C-methyl- β -D-ribo-hexopyranoside): **Vicenistatin M**
[312532-50-6]
C₃₀H₄₇NO₅ 501.705
Minor metab. of *Streptomyces halstedii* HC34. Powder. Mp 201-210°. [α]_D²⁷ -4.2 (c, 0.05 in MeOH). λ_{\max} 237 (37200); 241 (38000); 268 (13650) (MeOH).

Shindo, K. *et al.*, *J. Antibiot.*, 1993, **46**, 1076 (isol, pmr, cmr, struct)
Matsushima, Y. *et al.*, *J. Antibiot.*, 1998, **51**, 688-691 (synth)
Arai, H. *et al.*, *Tet. Lett.*, 1998, **39**, 3181-3184 (synth, abs config)
Otsuka, M. *et al.*, *Tetrahedron*, 2000, **56**, 8281-8286 (biosynth)
Matsushima, Y. *et al.*, *J. Antibiot.*, 2001, **54**, 211-219 (Vicenistatin M)

Nishida, H. *et al.*, *Tetrahedron*, 2001, **57**, 8237-8242 (biosynth)
Matsushima, Y. *et al.*, *J.C.S. Perkin 1*, 2002, 949-958 (synth)
Ogasawara, Y. *et al.*, *Chem. Biol.*, 2004, **11**, 79-86 (biosynth)

Vignatic acid A **V-99**
[181485-19-8]



R = CH₂Ph

C₃₀H₃₉N₃O₇ 553.654
Constit. of *Vigna radiata* (mung bean). Insecticidal agent. Powder. [α]_D -99 (c, 0.11 in MeOH).

Sugawara, F. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3360-3364 (isol, pmr, cmr, hplc)


Vignatic acid B **V-100**
[181485-20-1]

As Vignatic acid A, V-99 with
R = -CH₂CH(CH₃)₂

C₂₇H₄₁N₃O₇ 519.637
Constit. of *Vigna radiata* (mung bean). Insecticidal agent. Powder. [α]_D -79 (c, 0.16 in MeOH).

Sugawara, F. *et al.*, *J. Agric. Food Chem.*, 1996, **44**, 3360-3364 (isol, pmr, cmr, hplc)

Viguine **V-101**
1,2,3,4-Tetrahydro-2-(4-methoxybenzyl)-6,7-methylenedioxyisoquinoline
[107602-51-7]




C₁₈H₁₉NO₃ 297.353
Alkaloid from *Corydalis claviculata* (Papaveraceae). Cryst. (C₆H₆/hexane). Mp 63-64°.

Hydrochloride:
Cryst. (EtOH). Mp 232-233°.

N-Me: N-Methylviguine
[136092-29-0]
C₁₉H₂₂NO₃⁺ 312.388
Alkaloid from *Sarcocapnos saetabensis* (Papaveraceae). Powder.
Boente, J.M. *et al.*, *Heterocycles*, 1986, **24**, 3359 (uv, pmr, ms, struct, synth)
Blanco, O. *et al.*, *Phytochemistry*, 1991, **30**, 2071 (N-Methylviguine)

Villagorgin A V-102

4,6,7,12,12b,13-Hexahydro-1H-imidazo[4,5-g]indolo[2,3-a]quinazoline, 9CI [152606-59-2]



C₁₆H₁₆N₄ 264.329

Miscellaneous indole alkaloid, not seco-loganin-derived. CAS numbering shown.

(R)-form

Alkaloid from the gorgonian *Villogorgia rubra*. Shows calmodulin-related antagonist activity. Red amorph. solid. $[\alpha]_D^{20} +7.8$. Genus name incorrectly given as Villagorgia. λ_{max} 224 (ε 25600); 280 (ε 52409) (MeOH) (Berdy).

4,N,12b,13-Dehydro: 1,6,7,12-Tetrahydroimidazo[4,5-g]indolo[2,3-a]quinolin-5-ium(1+), 9CI. Villagorgin B [152606-60-5]

C₁₆H₁₃N₄⁺ 261.305


Alkaloid from the gorgonian *Villogorgia rubra*. Red amorph. solid. Counterion not specified. λ_{max} 228; 244; 280; 338 (MeOH) (Berdy). λ_{max} 222; 242; 332 (MeOH/HCl) (Berdy).

Espada, A. et al., *Tet. Lett.*, 1993, **34**, 7773-7776 (isol, uv, ir, pmr, cmr, ms, struct)

Grazul, R.M. et al., *Nat. Prod. Lett.*, 1994, **5**, 187 (Villagorgin B, synth)

Villalstonine

Alkaloid B⁺ [2723-56-0]



Absolute Configuration

C₄₁H₄₈N₄O₄ 660.855

Alkaloid from *Alstonia muelleriana*, *Alstonia macrophylla*, *Alstonia spectabilis*, *Alstonia villosa*, *Alstonia somersetensis* and *Alstonia verticillosa* (Apocynaceae). Cytotoxic. Noncryst. Mp 235-270° dec. $[\alpha]_D +79$ (CHCl₃).

Hydrochloride (1:2):

Cryst. + 4H₂O (MeOH/Et₂O). Mp 270° dec. $[\alpha]_D^{25} +46.6$ (c, 0.995 in 50% MeOH aq.). $[\alpha]_D +56.3$ (c, 0.5 in H₂O).

Sulfate:

Cryst. + 6H₂O. Mp 310°. $[\alpha]_D +52.94$ (c, 1.02 in H₂O).

4'-N-Oxide: Villalstonine 4'-N-oxide

[120396-42-1]

C₄₁H₄₈N₄O₅ 676.854

Alkaloid from the stem bark of *Alstonia angustifolia* (Apocynaceae).

10-Methoxy: 10-Methoxyvillalstonine

[120396-40-9]

C₄₂H₅₀N₄O₅ 690.881

Alkaloid from the leaves of *Alstonia angustifolia* (Apocynaceae). $[\alpha]_D +39$ (c, 0.66 in CHCl₃).

10-Methoxy, 4'-N-oxide: 10-Methoxyvillalstonine 4'-N-oxide

[120396-41-0]

C₄₂H₅₀N₄O₆ 706.88

Alkaloid from the leaves of *Alstonia angustifolia* (Apocynaceae).

Sharp, T.M. et al., *J.C.S.*, 1934, 1227 (isol)

Hesse, M. et al., *Helv. Chim. Acta*, 1965, **48**, 689 (uv, ms, struct)

Nordman, C.E. et al., *J.A.C.S.*, 1965, **87**, 2059 (cryst struct)

Kishi, T. et al., *Helv. Chim. Acta*, 1966, **49**, 946 (isol)

Hesse, M. et al., *Helv. Chim. Acta*, 1966, **49**, 1173 (struct)

Hart, N.K. et al., *Aust. J. Chem.*, 1972, **25**, 2739 (isol)

Elderfield, R.C. et al., *Phytochemistry*, 1972, **11**, 339 (isol)

Burke, D.E. et al., *J.A.C.S.*, 1973, **95**, 546 (synth)

Das, B.C. et al., *Tet. Lett.*, 1974, 4299 (cmr)

Ghedira, K. et al., *Phytochemistry*, 1988, **27**, 3955 (Villalstonine N-oxide, 10-Methoxyvillalstonine N-oxide)


Bi, Y. et al., *J.A.C.S.*, 1994, **116**, 9027 (synth)
Keawpradub, N. et al., *Planta Med.*, 1999, **65**, 311-315 (isol, activity)

Villatamine A

V-104

2-(5,7-Dodecadien-3-ynyl)-1-ethylpyrrolidine

[168434-14-8]



C₁₈H₂₉N 259.434

Alkaloid from the predatory flatworm *Prostheceraeus villatus* and its tunicate prey *Clavelina lepadiformis*. Oil. $[\alpha]_D +49$ (MeOH). *Prostheceraeus villatus* appears to be an Authors' misspelling of *P. vittatus*. λ_{max} 266 (ε 18700) (MeOH) (Berdy).

3',4'E, 7',8'-Tetrahydro: 2-(3,5-Dodecadienyl)-1-ethylpyrrolidine. Villatamine B [168434-15-9]

C₁₈H₃₃N 263.465


Alkaloid from *Prostheceraeus villatus* and *Clavelina lepadiformis*. Exhibits significant *in vitro* cytotoxicity against human cancer cells. Oil. $[\alpha]_D +15$ (MeOH). λ_{max} 230 (ε 19500) (MeOH) (Berdy).

Kubaneck, J. et al., *Tet. Lett.*, 1995, **36**, 6189-6192 (isol, uv, pmr, cmr, struct)

Vilmoridine

V-105

[159474-79-0]



1993

C₂₂H₂₉NO₆ 403.474


Alkaloid from roots of *Aconitum vilmorinianum* (Ranunculaceae).

Ding, L.-S. et al., *Huaxue Xuebao*, 1994, **52**, 932; *CA*, **122**, 5461y (isol, struct)

Vilmoritrine

V-106

[143086-35-5]



C₂₄H₃₅NO₅ 417.544

Alkaloid from the roots of *Aconitum vilmorinianum*.


Ding, L.S. et al., *Huaxue Xuebao*, 1992, **50**, 405-408; *CA*, **117**, 108100f

Vinamidine

V-107

Catharinine

[58511-83-4]



C₄₆H₅₆N₄O₁₀ 824.969

Alkaloid from *Catharanthus roseus*, *Catharanthus longifolius* and *Catharanthus ovalis* (Apocynaceae). Amorph. $[\alpha]_D -32$ (c, 1.1 in CHCl₃).

Tafur, S. et al., *J. Pharm. Sci.*, 1975, **64**, 1953 (isol, uv, ir, cmr, ms)

Andriamialisoa, R.Z. et al., *Tetrahedron*, 1978, **34**, 677 (isol, uv, ir, pmr, cmr, ms, cd, cryst struct)

Bornmann, W.G. et al., *J.O.C.*, 1992, **57**, 1752 (synth)

Vinaphamine

V-108

[1361-98-4]

Bisindole alkaloid. Struct. unknown.

Alkaloid from *Catharanthus roseus* (Apocynaceae). Thin blades (MeOH). Mp 229-235°. Mol. formula not recorded.

Svoboda, G.H. et al., *J. Pharm. Sci.*, 1964, **53**, 1227-1231 (isol, uv, ir)

Vinaspine

V-109


[1361-99-5]

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Blades (MeOH). Mp 235-238°. pK_a 7.85 (33% DMF aq.).

Svoboda, G.H. et al., *J. Pharm. Sci.*, 1964, **53**,

1227-1231 (*isol, ir, uv*)**Vinblastine, BAN, INN** **V-110**

Vincalcalekoblastine, 9CI. Vincalcalekoblastine. Velbe. VLB
[865-21-4]



$C_{46}H_{58}N_4O_9$ 810.986

Alkaloid from *Catharanthus roseus* (Apocynaceae) and from *Catharanthus trichophyllus*. Antineoplastic agent, used widely in the treatment of Hodgkin's disease and other lymphomas. Microtubule assembly inhibitor. Shows trypano-somical effect. Mp 180-182° (Et₂O solvate) Mp 216° (dry). $[\alpha]_D^{25} +42$ (CHCl₃). Log P 4.29 (uncertain value) (calc). λ_{max} 214 (ϵ 53700); 259 (ϵ 16220) (EtOH) (Berdy).

- ▶ Adverse human effects reported when used therapeutically or from occup. exposure. Skin and respiratory tract irritant. Causes corneal damage in contact with eyes. Central and peripheral neurotoxic effects. Other effects incl. bone-marrow depression and gastrointestinal effects. LD₅₀ (rat, ipr) 1 mg/kg. Exp. reprod. and teratogenic effects. YY8050000

Sulfate salt: Vinblastine sulfate, JAN, USAN. Blastovin. Exal. Periblastine. Rozevin. Velban. Velbe. LE 29060. NSC 49842
[143-67-9]
Mp 284-285°. $[\alpha]_D^{26} -28$ (MeOH).

- ▶ YY8400000

O-De-Ac: Deacetylvinblastine

[3352-69-0]
 $C_{44}H_{56}N_4O_8$ 768.948
Alkaloid from *Catharanthus roseus*. Antitumour agent. λ_{max} 214 (E1%/1cm 576); 266 (E1%/1cm 186); 294 (E1%/1cm 126) (EtOH) (Berdy).

- ▶ YY8150000

O-De-Ac, O¹⁷-(2-dimethylaminoacetyl): Vinglycinate, INN. Lilly 49040
[865-24-7]

$C_{48}H_{63}N_5O_9$ 854.054
Antineoplastic agent. Cryst. (Et₂O). Log P 4.19 (uncertain value) (calc).

- ▶ YY8180300

O-De-Ac, O¹⁷-(2-dimethylaminoacetyl); sulfate salt (2:3): Vinglycinate sulfate, USAN
[7281-31-4]
Cryst. (MeOH/EtOH).

O,O-Di-Ac: Mp 168-170°. $[\alpha]_D -26.4$ (CHCl₃).

N-De-Me: N-Demethylvinblastine. N-De-formylvincristine

$C_{45}H_{56}N_4O_9$ 796.959

Alkaloid from *Catharanthus roseus* (Apocynaceae). Cryst. (EtOH) (as sul-

fate). Mp 250° dec. (sulfate).

N-De-Me, N-formyl: see Vincristine, V-128

20'-Deoxy: Isoleurosine. 20'-Deoxyvinblastine. Deoxyvinblastine A. Deoxy-vincalcalekoblastine A
[20072-25-7]

$C_{46}H_{58}N_4O_8$ 794.986

Isol. from *Catharanthus roseus* (Apocynaceae). Blades (MeOH). Mp 202-206° dec. $[\alpha]_D^{26} +61.2$ (c, 1 in CHCl₃).

20'-Deoxy, hydroxy: Roseamine

[87571-01-5]

$C_{46}H_{58}N_4O_9$ 810.986

Alkaloid from *Catharanthus roseus* (Apocynaceae). Shows antineoplastic props. Struct. not fully elucidated.

20'-Deoxy, 15',20'-didehydro: 15',20'-Anhydrovinblastine. 3',4'-Anhydrovinblastine
[38390-45-3]

$C_{46}H_{56}N_4O_8$ 792.97

Alkaloid from shoot cultures and freshly picked leaves of *Catharanthus roseus* (Apocynaceae). Antineoplastic agent. Unstable in soln. Undergoes ready aerial oxidn. to Leurosine, L-162 and Catharine, C-197 as well as to the other main antitumour alkaloids of *Catharanthus* spp., Vinblastine, Leurosine and their deoxy derivs.

Deacetoxy: Deacetoxyvinblastine

$C_{44}H_{56}N_4O_7$ 752.949

Alkaloid from *Catharanthus roseus* (Apocynaceae). Cryst. (MeOH). Mp 183-190° dec. $[\alpha]_D^{26} +95.3$ (CHCl₃).

14'-Hydroxy: Leurocolumbine

[56974-17-5]

$C_{46}H_{58}N_4O_{10}$ 826.985

From *Catharanthus roseus* (Apocynaceae). Antineoplastic and antimitotic agent. Log P 3.08 (uncertain value) (calc). λ_{max} 217 (ϵ 51000); 265 (15600) (MeOH).

15' α -Hydroxy: Vincadioline

[56897-74-6]

$C_{46}H_{58}N_4O_{10}$ 826.985

Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Antimitotic agent. Mp 218-221°. Log P 3.05 (uncertain value) (calc). Proposed struct. No details concerning the elucidation of its struct. have been publ. λ_{max} 217 (ϵ 51090); 265 (ϵ 15660) (MeOH) (Berdy).

16-De(methoxycarbonyl), 16-carbamoyl, O-de-Ac: Vindesine, BAN, INN, USAN. NSC 24567

[53643-48-4]

$C_{43}H_{55}N_5O_7$ 753.937

Alkaloid from *Catharanthus roseus*. Antineoplastic agent. Cryst. (EtOH/MeOH). Mp 230-232°. $[\alpha]_D^{25} +39.4$ (c, 1 in MeOH). Log P 2.41 (uncertain value) (calc). λ_{max} 214 (ϵ 53400); 266 (ϵ 17450); 288 (ϵ 13950); 296 (ϵ 12500) (MeOH) (Berdy).

- ▶ LD₅₀ (mus, ivn) 6.3 mg/kg. YY8080000

16-De(methoxycarbonyl), 16-carbamoyl, O-de-Ac; sulfate: Vindesine sulfate, JAN, USAN. Eldisine. Elsedine. Fildesine. Gesidine. Lilly 112531. LY 99094.

NSC 245467

[59917-39-4]

Amorph. solid (2-propanol). Mp 250°.

- ▶ YY8090000

20'-Epimer: Leurosidine. Vinrosidine. 20'-Epivinblastine. Lilly 36781
[15228-71-4]

$C_{46}H_{58}N_4O_9$ 810.986

Alkaloid from *Vinca rosea* (*Catharanthus roseus*) (Apocynaceae). Anti-

neoplastic agent, toxic side effects in clinical trials. Mp 210-212° dec. $[\alpha]_D^{25} +55.8$ (c, 1 in CHCl₃). Log P 4.29 (uncertain value) (calc). λ_{max} 214; 265 (EtOH) (Berdy).

- ▶ LD₅₀ (mus, ipr) 50 mg/kg. OH6350000

20'-Epimer, sulfate salt: Vinrosidine sulfate
[18556-44-0]

- ▶ OK6126000

20'-Epimer, N^{4'}-oxide: Leurosidine N^{b'}-oxide

$C_{46}H_{58}N_4O_{10}$ 826.985

Isol. from *Catharanthus roseus* (Apocynaceae). Cryst. (Me₂CO). Poorly sol. hexane. Mp 215-218°. $[\alpha]_D^{25} +26$ (c, 0.5 in CHCl₃). λ_{max} 218; 288; 298; 315 (EtOH) (Berdy).

20'-Epimer, tri-Ac: Mp 154-159° dec.

$[\alpha]_D^{25} +12.6$ (CHCl₃).

20'-Epimer, 20'-deoxy: 20'-Deoxyleurosidine. Deoxyvincalcalekoblastine B
[21631-00-5]

$C_{46}H_{58}N_4O_8$ 794.986

Alkaloid from the aerial parts of *Catharanthus ovalis* (Apocynaceae).

20'-Epimer, 20'-deoxy, N-de-Me, N-formyl: Vinepidine, INN. LY 119863

[68170-69-4]

[83200-11-7 (sulfate salt)]

$C_{46}H_{56}N_4O_9$ 808.97

Antineoplastic agent. Log P 4.11 (uncertain value) (calc).

14',16',20'-Triepimer: Vincovaline

[61949-65-3]

$C_{46}H_{58}N_4O_9$ 810.986

Alkaloid from *Catharanthus ovalis* (Apocynaceae). $[\alpha]_D -118$ (c, 1.0 in CHCl₃).

Aldrich Library of Infrared Spectra, 3rd edn., 1981, 1511H (ir)

Neuss, N. et al., *J.A.C.S.*, 1959, **81**, 4754; 1962, **84**, 1509; 1964, **86**, 1440 (*uv, ir, pmr, struct*)

Svoboda, G.H. et al., *J. Nat. Prod.*, 1961, **24**, 213 (*isol, uv, ir*)

Svoboda, G.H. et al., *J. Pharm. Sci.*, 1961, **50**, 409-413 (*Isoleurosine*)

Svoboda, G.H. et al., *J. Pharm. Sci.*, 1964, **53**, 1227

Bommer, P. et al., *J.A.C.S.*, 1964, **86**, 1439-1440 (*ms*)

Moncrief, J.W. et al., *J.A.C.S.*, 1965, **87**, 4963-4964 (*Leurocristine, cryst struct*)

Johnson, I.S. et al., *Cancer Res.*, 1966, **26**, 2431 (*Vinglycinate, activity*)

Netherlands Pat., 1966, 6 501 121; *CA*, **65**, 18642f (*Vinglycinate*)

Armstrong, J.G. et al., *Cancer Res.*, 1967, **27A**, 221-227 (*Vinglycinate, activity*)

Neuss, N. et al., *Tet. Lett.*, 1967, **8**, 811-816; 1968, 783 (*Leurosidine, struct*)

U.S. Pat., 1968, ((*Roussel-UCLAF*)) 3 387 001; *CA*, **69**, 59476n (*synth, vinglycinate*)

Burns, J.H. et al., *Anal. Profiles Drug Subst.*, 1972, **1**, 443 (*rev, uv, ir, pmr, anal*)


Wenkert, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4990-4995 (*cmr*)
Catharanthus Alkaloids, (Eds, Taylor, W.I. *et al.*), M. Dekker, 1975, (*pharmacol*)
 Neuss, N. *et al.*, *Experientia*, 1975, **31**, 18 (*Deacetoxyvinblastine*)
 Wenkert, E. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1560 (*cmr, struct*)
 Tafur, S. *et al.*, *J. Pharm. Sci.*, 1975, **64**, 1953 (*Leurocolumbine*)
U.S. Pat., 1975, ((*Eli Lilly*)) 3 887 565; *CA*, **84**, 5230u (*Vincadioline*)
 Sieber, S.M. *et al.*, *Cancer Treat. Rep.*, 1976, **60**, 1127 (*pharmacol*)
 Dorman, D.E. *et al.*, *Org. Magn. Reson.*, 1976, **8**, 413 (*cmr*)
 Langlois, N. *et al.*, *Tet. Lett.*, 1976, **17**, 1099-1102 (*Leurosidine, synth, ms, pmr*)
 Andriamialisoa, R.Z. *et al.*, *Tet. Lett.*, 1976, **17**, 2849-2852 (*Vincovaline*)
 Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B.*, 1976, **31**, 1416 (*rev*)
 Kutney, J.P. *et al.*, *J. Nat. Prod.*, 1977, **40**, 107 (*rev*)
 Kutney, J.P. *et al.*, *Can. J. Chem.*, 1978, **56**, 2560 (*Vinepidine*)
 Langlois, N. *et al.*, *Chem. Comm.*, 1978, 102; 1979, 582 (*Anhydrovinblastine*)
Ger. Pat., 1978, 2 801 748; *CA*, **89**, 197778 (*Vinepidine*)
 Burnett, C.J. *et al.*, *J. Med. Chem.*, 1978, **21**, 88 (*Vindesine*)
 Scott, A.I. *et al.*, *J.A.C.S.*, 1978, **100**, 6253 (*Anhydrovinblastine*)
 Hassam, S.B. *et al.*, *Tet. Lett.*, 1978, 1681 (*biosynth*)
 Potier, P. *et al.*, *J.A.C.S.*, 1979, **101**, 2243 (*synth*)
 Dana, W.J. *et al.*, *Drug Intell. Clin. Pharm.*, 1980, **14**, 28 (*Vindesine, rev*)
 Langlois, N. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 793 (*20'-Deoxyleurosidine*)
 Brade, W. *et al.*, *Contrib. Oncol.*, (Eds), Vindesine. S. Karger, Basel, Switz., 1981, **6**, (*book*)
 Mukhopadhyay, S. *et al.*, *J. Nat. Prod.*, 1981, **44**, 611-613 (*Leurosidine N^b oxide*)
 Baxter, R.L. *et al.*, *Chem. Comm.*, 1982, 791 (*biosynth*)
 Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2088 (*biosynth*)
 El-Sayed, A. *et al.*, *J. Nat. Prod.*, 1983, **46**, 517-527 (*Roseamine*)
 McLauchlan, W.R. *et al.*, *Tetrahedron*, 1983, **39**, 3777 (*biosynth*)
 Simonds, R. *et al.*, *Planta Med.*, 1984, **50**, 274 (*N-Demethylvinblastine*)
 Magnus, P. *et al.*, *Stud. Org. Chem. (Amsterdam)*, 1986, **25**, 203; 257 (*synth, revs*)
 Endo, T. *et al.*, *Planta Med.*, 1987, **53**, 479-482 (*isol*)
 Kutney, J.P. *et al.*, *Heterocycles*, 1988, **27**, 1845-1853 (*Vinblastine, Leurosidine, synth*)
 Goodbody, A.E. *et al.*, *Phytochemistry*, 1988, **27**, 1713 (*isol, Anhydrovinblastine*)
 Kuehne, M.E. *et al.*, *J.O.C.*, 1989, **54**, 3407; 1991, **56**, 513 (*Vinblastine, 20'-Deoxyvinblastine, 20'-Deoxyleurosidine, Vincovaline, synth, uv, ir, pmr, ms*)
 Kuehne, M.E. *et al.*, *Alkaloids (Academic Press)*, 1990, **37**, 77 (*synth, rev*)
 Neuss, N. *et al.*, *Alkaloids (Academic Press)*, 1990, **37**, 229 (*pharmacol*)
 Magnus, P. *et al.*, *J.A.C.S.*, 1990, **112**, 8210; 1992, **114**, 10232 (*synth*)
 Muhtadi, F.J. *et al.*, *Anal. Profiles Drug Subst.*, 1992, **21**, 611 (*rev*)
 Borrmann, W.G. *et al.*, *J.O.C.*, 1992, **57**, 1752 (*synth*)
 Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 504

Andrews, C.W. *et al.*, *J. Het. Chem.*, 1995, **32**, 1011 (*pmr, conformn, Anhydrovinblastine*)
 Parish, C.A. *et al.*, *Tetrahedron*, 1998, **54**, 15739-15759 (*cd*)
 Bau, R. *et al.*, *J.C.S. Perkin 1*, 2000, 2079-2082 (*cryst struct*)
 Schneider, C. *et al.*, *Angew. Chem., Int. Ed.*, 2002, **41**, 4217-4219 (*synth*)
 Hardouin, C. *et al.*, *J.O.C.*, 2002, **67**, 6571-6574 (*Anhydrovinblastine, synth, pmr, cmr*)
 Hardouin, C. *et al.*, *Org. Lett.*, 2002, **4**, 1151-1153 (*synth*)
 Ramnath, N. *et al.*, *Cancer Chemother. Pharmacol.*, 2003, **51**, 227-230 (*Anhydrovinblastine, pharmacol*)
 Yokoshima, S. *et al.*, *Pure Appl. Chem.*, 2003, **75**, 29-38 (*synth, bibl*)
 Miyazaki, T. *et al.*, *Org. Lett.*, 2007, **9**, 4737-4740 (*synth*)
 Ishikawa, H. *et al.*, *J.A.C.S.*, 2008, **130**, 420-421 (*synth*)
 Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VGU750; VLA000; VKZ000

Vincadifformine

V-111

Methyl 2,3-didehydroaspidospermidine-3-carboxylate, 9CI



(-)-form

C₂₁H₂₆N₂O₂ 338.449**(+)-form** [15539-10-3]

Alkaloid from *Amsonia tabernaemontana*, *Amsonia angustifolia*, *Rhazya stricta*, *Tabernaemontana riedelii* (Apocynaceae). Mp 96°. [α]_D²⁵ +605 (c, 0.2 in EtOH).

(-)-form [3247-10-7]

Alkaloid from *Vinca minor* (Apocynaceae). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 96°. [α]_D²⁰ -540 (EtOH) (natural). [α]_D -600 (EtOH) (semisynthetic).

N-Me: Minovine

[19074-77-2]
 [19621-72-8, 70469-81-7]
 C₂₂H₂₈N₂O₂ 352.475

Alkaloid from *Vinca minor* (Apocynaceae). Cryst. (MeOH). Mp 80-82°. [α]_D 0 (MeOH).

11-Hydroxy: 11-Hydroxyvincadifformine

[80993-50-6]
 C₂₁H₂₆N₂O₃ 354.448

Alkaloid from aerial parts of *Melodinus hemsleyanus* (Apocynaceae). Amorph. [α]_D²³ -502 (c, 0.1 in CHCl₃).

15β-Hydroxy: 15β-Hydroxyvincadifformine

[119478-89-6]
 C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the leaves of *Rhazya stricta* (Apocynaceae). [α]_D +240.57 (CHCl₃).

11-Methoxy: Ervinceine. 11-Methoxyvincadifformine

[25858-80-4]
 C₂₂H₂₈N₂O₃ 368.475

Alkaloid from the epigeal parts of

Vinca erecta (Apocynaceae). Cryst. (MeOH). Mp 99-100°. [α]_D²⁵ -448 (c, 0.09 in CHCl₃). An earlier isoln. (1968) of this compd. consisted of impure or incorrectly identified material.

11-Methoxy, N-oxide: 11-Methoxyvincadifformine N-oxide

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from epigeal parts of *Vinca erecta* (Apocynaceae). Cryst. (Me₂CO). Sol. H₂O. Mp 150-152°. Struct. unconfirmed; struct. proof was by reference to the impure 1968 isol. of Ervinceine.

5-Oxo: 5-Oxovincadifformine. Ervinidinine

[23107-01-9]
 C₂₁H₂₄N₂O₃ 352.432

Alkaloid from the leaves of *Pterotaberna inconspua* and *Vinca erecta* (Apocynaceae). Cryst. (MeOH). Mp 265-266° dec. [α]_D -152 (c, 0.25 in MeOH). Identity of Ervinidinine and 5-Oxovincadifformine not certain. Mp. refers to Ervinidinine.

Stereoisomer (?): Ervamine

[6878-14-4]
 C₂₁H₂₆N₂O₂ 338.449

Alkaloid from *Vinca erecta* and *Catharanthus roseus* (Apocynaceae). Cryst. (EtOH)(as hydroiodide). Mp 198-200° (hydroiodide). [α]_D -502 (c, 0.29 in MeOH). Stereochem. not clear from the abstract.

(±)-form [18374-17-9]

Alkaloid from *Vinca difformis*, *Rhazya stricta*, *Tabernaemontana riedelii* (Apocynaceae). Shows hypotensive activity, about half that of Reserpine, R-52. Mp 124-125°.

Plat, M. *et al.*, *Bull. Soc. Chim. Fr.*, 1962, 2237 (*isol*)

Mokry, J. *et al.*, *Experientia*, 1962, **18**, 564 (*Minovine, isol, uv, ir*)

Djerassi, C. *et al.*, *Tet. Lett.*, 1962, 235 (*isol, ms, uv, ir*)

Malikov, V.M. *et al.*, *CA*, 1963, **59**, 11584b (*Ervamine*)

Zachystalová, D. *et al.*, *Chem. Ind. (London)*, 1963, 610 (*Minovine, isol*)

Klyne, W. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 443 (*abs config*)

Rakhimov, D.A. *et al.*, *Khim. Prir. Soedin.*, 1969, **5**, 330; 1970, **6**, 226; *Chem. Nat. Compd. (Engl. Transl.)*, 1969, **5**, 221; 1970, **6**, 280 (*Ervinceine, isol, pmr, struct*)

Ziegler, F.E. *et al.*, *J.A.C.S.*, 1970, **92**, 3492; 1973, **95**, 7146 (*synth*)

Szadon, B. *et al.*, *Tet. Lett.*, 1970, 4615 (*isol*)
 Malikov, V.M. *et al.*, *Khim. Prir. Soedin.*, 1971, **7**, 640; *Chem. Nat. Compd. (Engl. Transl.)*, 619 (*Ervinidinine*)

Wenkert, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4990 (*cmr*)

Khalmirzaev, M.M. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 806; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 776 (*Ervinceine N-oxide*)

Laronze, J.Y. *et al.*, *Tet. Lett.*, 1974, 491 (*synth*)
 Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1648 (*Minovine, synth*)

Kutney, J.P. *et al.*, *J.A.C.S.*, 1978, **100**, 4220 (*Ervinceine, synth, ir, uv, pmr*)


Kuehne, M.E. *et al.*, *J.O.C.*, 1979, **44**, 1063; 2477 (*Minovine, Ervinceine, synth, pmr, ir, ms, uv*)

Kuehne, M.E. *et al.*, *J.O.C.*, 1986, **51**, 2913 (*15β-Hydroxyvincadifformine*)

- Massiot, G. et al., *Phytochemistry*, 1988, **27**, 1085 (5-Oxovincadiformine)
 Torrenegra, R. et al., *Phytochemistry*, 1988, **27**, 1843 (3-Oxovincadiformine)
 Atta-ur-Rahman, et al., *Phytochemistry*, 1988, **27**, 3721 (15 β -Hydroxyvincadiformine)
 Takano, S. et al., *Chem. Lett.*, 1989, 87 (synth)
 Guo, L.-W. et al., *Phytochemistry*, 1993, **34**, 563 (11-Hydroxyvincadiformine)
 Kobayashi, S. et al., *Tet. Lett.*, 1999, **40**, 1519-1522 (synth)
 Yuan, Z.Q. et al., *Org. Lett.*, 2005, **7**, 741-744 (synth)
 Kalaus, G. et al., *Heterocycles*, 2006, **68**, 257-270 (15 β -Hydroxyvincadiformine, synth)
 Ishikawa, H. et al., *J.A.C.S.*, 2006, **128**, 10596-10612 (Minovine, synth)

Vincadine V-112

Methyl 7-ethyl-1,4,5,6,7,8,9,10-octahydro-2H-3,7-methanoazacycloundecino[5,4-b]indole-9-carboxylate, 9CI. 16-Carbomethoxyquebrachamine



(16R,20S)-form

C₂₁H₂₈N₂O₂ 340.464**(16R,20S)-form** [2896-91-5]

[32790-08-2]

Alkaloid from *Vinca minor*, also identified in the leaves of *Amsonia tabernaemontana* (Apocynaceae). Mp 76°. [α]_D²⁰ +92 (EtOH).

N-Me: **Vincaminoreine**. N-Methylvincadine

[2411-51-0]

C₂₂H₃₀N₂O₂ 354.491

Minor alkaloid from *Vinca minor* (Apocynaceae). Needles (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 138-139° (126°). [α]_D +26.5 (CHCl₃). pK_a 7.55 (50% EtOH aq.). λ_{\max} 230 (log ϵ 4.59); 288 (log ϵ 3.93); 296 (log ϵ 3.91) (EtOH).

14,15-Didehydro: **14,15-Didehydrovincadine**. 14,15-Dehydrovincadine. 6,7-Dehydrovincadine

[41222-15-5]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from the leaves of *Amsonia tabernaemontana* (Apocynaceae). Amorph. solid. [α]_D²⁰ +65 (c, 0.5 in EtOH). pK_a 6.75 (50% EtOH). Epimerised by silica gel at r.t. to 14,15-Dehydroepivincadine. λ_{\max} 226 (log ϵ 4.58); 284 (log ϵ 3.9); 292 (log ϵ 3.86) (EtOH).

(16R,20R)-form

(-)-Epivincadine

[67968-78-9]

Constit. of the leaves of *Amsonia tabernaemontana* (Apocynaceae). Mp 156°. [α]_D²⁰ -24 (EtOH). Positively identified but not isol. Constants are for synthetic material.

(16S,20R)-form [36132-39-5]

Alkaloid from the leaves of *Amsonia tabernaemontana* (Apocynaceae). Cryst. (heptane). Mp 76°. [α]_D²⁰ -92 (c, 0.5 in EtOH). pK_a 7.2 (50% EtOH).

N-Me: [52553-75-0]

Synthetic. Cryst. (EtOH). Mp 129°.

[α]_D²⁰ -27 (c, 1. in EtOH). pK_a 7.5 (50% EtOH).**(16S,20S)-form**

(+)-Epivincadine

[41221-70-9]

Constit. of the leaves of *Amsonia tabernaemontana* (Apocynaceae). Mp 156°. [α]_D²⁰ +24 (EtOH). Positively identified, but not isol. Constants are for synthetic material.

N-Me: **Vincaminorine**. N-Methylepivincadine

[1935-07-5]

C₂₂H₃₀N₂O₂ 354.491

Alkaloid from *Vinca minor* (Apocynaceae). Needles (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 130-131°. [α]_D²⁰ +46 (EtOH). pK_a 5.45 (50% EtOH). λ_{\max} 226; 282; 291 (EtOH) (Berdy).

14,15-Didehydro: **14,15-Didehydroepivincadine**. Dehydroepivincadine

[41222-16-6]

C₂₁H₂₆N₂O₂ 338.449

Alkaloid from leaves of *Amsonia tabernaemontana* (Apocynaceae). Solid or foam. [α]_D²⁰ +90 (c, 0.5 in EtOH) (+65). λ_{\max} 226 (log ϵ 4.57); 285 (log ϵ 3.94); 292 (log ϵ 3.91) (EtOH).

11-Methoxy, N-Me: **Vincaminoridine**

[21197-03-5]

C₂₃H₃₂N₂O₃ 384.517

Minor alkaloid from *Vinca minor* (Apocynaceae). Cryst. (heptane). Mp 99-100°. [α]_D²⁰ +57.7 (c, 1.05 in CHCl₃). λ_{\max} 232 (log ϵ 4.57); 288 (sh) (log ϵ 3.79); 300 (log ϵ 3.84) (no solvent reported).

(16RS,20RS)-form(\pm)-Epivincadine

[58072-80-3]

Alkaloid from the leaves of *Amsonia tabernaemontana* (Apocynaceae). Cryst. (MeOH). Mp 156-157°. pK_a 5.2 (50% EtOH). λ_{\max} 227 (log ϵ 4.52); 287 (log ϵ 3.91); 293 (log ϵ 3.87) (EtOH).

(16RS,20SR)-form(\pm)-Vincadine

[19621-74-0]

Alkaloid from the leaves of *Amsonia tabernaemontana* and the leaves and stems of *Amsonia angustifolia* (Apocynaceae). Cryst. (MeOH). Mp 126°. λ_{\max} 227 (log ϵ 4.45); 286 (log ϵ 3.89); 292 (log ϵ 3.87) (EtOH).

Trojánek, J. et al., *Coll. Czech. Chem. Comm.*, 1959, **24**, 526-529; 1960, **25**, 2045-2048 (*Vincaminorine*, *Vincaminoreine*, isol, uv, ir)

Mokry, J. et al., *Tet. Lett.*, 1962, 1185-1188 (*Vincadine*, *Vincaminoreine*, isol, uv, ir, struct)

Mokry, J. et al., *Chem. Ind. (London)*, 1964, 1988-1989 (*Vincaminorine*, *Vincaminoreine*, uv, ir, struct, config)

Klyne, W. et al., *Helv. Chim. Acta*, 1967, **50**, 115-127 (ord)

Kompiš, J. et al., *Coll. Czech. Chem. Comm.*, 1968, **33**, 4328-4336 (*Vincaminoridine*)

Kutney, J.P. et al., *J.A.C.S.*, 1968, **90**, 3891-3893 (*Vincaminoreine*, synth)

Zsadon, B. et al., *Acta Chim. Acad. Sci. Hung.*, 1973, **78**, 207-216; 1978, **96**, 167-181 (*Amsonia tabernaemontana constitis*)

Zsadon, B. et al., *Chem. Ind. (London)*, 1973, 229-230 (*Epivincadine*, 14,15-Dehydrovincadine, 14,15-Dehydroepivincadine)

Kutney, J.P. et al., *Helv. Chim. Acta*, 1975, **58**, 1648-1671 (*Vincadine*, *Vincaminorine*, *Vincaminoridine*, uv, ir, pmr, ms)


Wenkert, E. et al., *Helv. Chim. Acta*, 1976, **59**, 2711-2723 (emr)

Wenkert, E. et al., *Tetrahedron*, 1981, **37**, 4017 (synth)

Vincamajine

V-113

[2506-26-5]



Absolute Configuration

C₂₂H₂₆N₂O₃ 366.459

Alkaloid from *Vinca major*, *Vinca herbarcea*, *Vinca libanotica*, *Alstonia spectabilis*, *Alstonia quaternata*, *Alstonia legouixiae*, *Alstonia odontophora*, *Alstonia constricta* and *Cabucala torulosa* (Apocynaceae). Needles (MeOH). Mp 226.5-227°. [α]_D²⁵ -58 (c, 0.5 in MeOH).

Hydrochloride: Mp 259-261° dec. [α]_D -40 (c, 0.38 in H₂O).

O-Ac: **Vincamedine**

[912-27-6]

C₂₄H₂₈N₂O₄ 408.496

Alkaloid from *Vinca difformis* and *Vinca major* (Apocynaceae). Mp 185.5-186.5°. [α]_D²⁰ -75 (c, 1 in CHCl₃). λ_{\max} 250 (log ϵ 3.92); 295 (log ϵ 3.45) (no solvent reported).

O-Benzoyl: **O-Benzoylvincamajine**

[24190-04-3]

C₂₉H₃₀N₂O₄ 470.567

Constit. of the leaves of *Alstonia macrophylla* (Apocynaceae). Mp 267-269°. [α]_D³⁰ -147 (CHCl₃).

O-(3,4-Dimethoxybenzoyl): **Vincamajine****17-O-veratrate**

[154849-47-5]

C₃₁H₃₄N₂O₆ 530.619

Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. [α]_D²⁵ -129.6 (c, 0.68 in MeOH).

O-(4-Hydroxy-3,5-dimethoxybenzoyl):

O-(4-Hydroxy-3,5-dimethoxybenzoyl)-vincamajine

[144379-35-1]

C₃₁H₃₄N₂O₇ 546.619

Minor alkaloid from roots of *Alstonia angustifolia* (Apocynaceae). λ_{\max} 248; 290 (MeOH).

O-(3,4,5-Trimethoxybenzoyl): **O-(3,4,5-Trimethoxybenzoyl)vincamajine**

[71385-80-3]

C₃₂H₃₆N₂O₇ 560.646

Alkaloid from *Alstonia macrophylla*, *Alstonia boullindaensis* and *Alstonia villosa*.

O-(3,4,5-Trimethoxybenzoyl), N^d-oxide:

- [214263-33-9]
C₃₂H₃₆N₂O₈ 576.645
Alkaloid from *Alstonia macrophylla* and *Alstonia villosa*.
- O-(3,4,5-Trimethoxycinnamoyl): 3,4,5-Trimethoxycinnamoylvincamajine**
[57800-03-0]
C₃₄H₃₈N₂O₇ 586.683
Alkaloid from the root bark of *Alstonia constricta* and from *Alstonia lanceolifera* (Apocynaceae). Noncryst. [α]_D²⁵ -68 (c, 2.6 in CHCl₃). λ_{max} 236 (log ε 4.34); 305 (log ε 3.71) (EtOH).
- N-De-Me: Quebrachidine. Vincarine**
[4835-69-2]
[21641-60-1]
C₂₁H₂₄N₂O₃ 352.432
Alkaloid from *Aspidosperma quebracho-blanco*, *Rauwolfia discolor*, several *Cabucala* spp., *Vinca libanotica*, *Vinca herbacea*, *Vinca erecta*, *Alstonia odontophora*, *Alstonia spectabilis*, *Alstonia macrophylla*, *Alstonia muelleriana*, *Tabernaemontana undulata* and *Tabernaemontana macrocalyx*. Hypotensive agent showing CNS effects. Exp. sedative. Mp 276-278° (264-265°). [α]_D²⁶ +54 (CHCl₃). [α]_D²⁵ +14 (c, 0.78 in MeOH). pK_a 6.7. λ_{max} 242 (ε 6390); 290 (ε 2920) (EtOH).
- ▶ LD₅₀ (mus, ipr) 330 mg/kg.
- N-De-Me, N¹-(3,4,5-trimethoxybenzoyl): N¹-Demethylvincamajine N¹-tri-O-methylgallate**
C₃₁H₃₄N₂O₇ 546.619
Alkaloid from leaves of *Alstonia macrophylla* (Apocynaceae). Solid. [α]_D²⁵ -98.4 (c, 0.56 in MeOH). Erroneously named Vincamajine N¹-tri-O-methylgallate in paper. Incorrectly indexed in CA.
- N-De-Me, O-(3,4,5-trimethoxybenzoyl): (3,4,5-Trimethoxybenzoyl)quebrachidine**
[31148-63-7]
C₃₁H₃₄N₂O₇ 546.619
Alkaloid from the root bark of *Alstonia constricta* (Apocynaceae). Needles (MeOH). Mp 231-232°. [α]_D²⁵ -36 (c, 1.3 in CHCl₃).
- 10-Hydroxy, O¹⁷-(3,4,5-trimethoxybenzoyl): 10-Hydroxy-O¹⁷-(3,4,5-trimethoxybenzoyl)vincamajine**
[57800-05-2]
C₃₂H₃₆N₂O₈ 576.645
Alkaloid from the aerial parts of *Alstonia lanceolifera* (Apocynaceae). Amorph. [α]_D²⁵ -43 (c, 1 in CHCl₃).
- 10-Hydroxy, O¹⁷-(3,4,5-trimethoxycinnamoyl): 10-Hydroxy-O¹⁷-(3,4,5-trimethoxycinnamoyl)vincamajine**
[57808-41-0]
C₃₄H₃₈N₂O₈ 602.683
Alkaloid from the aerial parts of *Alstonia lanceolifera* (Apocynaceae). [α]_D²⁵ -114 (c, 1 in CHCl₃).
- 19ξ-Hydroxy, 19,20-dihydro: 19-Hydroxyvincamajine. Dihydro-19-hydroxyvincamajine**
[108195-74-0]
C₂₂H₂₈N₂O₄ 384.474
Alkaloid from the leaves of *Alstonia macrophylla* (Apocynaceae). λ_{max} 175 (log ε 3.5); 247 (log ε 3.8) (no solvent reported).
- 10-Methoxy: 10-Methoxyvincamajine**
[57800-02-9]
C₂₃H₂₈N₂O₄ 396.485
Alkaloid from the aerial parts of *Alstonia lanceolifera* and *Alstonia bouldaensis* (Apocynaceae). Amorph. [α]_D²⁵ 0 (c, 1 in EtOH). λ_{max} 248 (log ε 3.97); 307 (log ε 3.63) (EtOH).
- 10-Methoxy, O-Ac: 10-Methoxyvincamide**
[94444-30-1]
C₂₅H₃₀N₂O₅ 438.522
Alkaloid from leaves, fruit and stem bark of *Alstonia sphaerocapitata* (Apocynaceae). [α]_D²⁵ -9 (c, 1 in CHCl₃). λ_{max} 247; 313 (MeOH).
- 10-Methoxy, O-Ac, N⁴-oxide: 10-Methoxyvincamide N(4)-oxide**
[94444-31-2]
C₂₅H₃₀N₂O₆ 454.522
Alkaloid from the leaves, fruit and stem bark of *Alstonia sphaerocapitata* (Apocynaceae). [α]_D²⁵ -2 (c, 1 in CHCl₃). λ_{max} 212; 248; 305 (MeOH).
- 10-Methoxy, O-(3,4,5-trimethoxycinnamoyl): 10-Methoxy-O-(3,4,5-trimethoxycinnamoyl)vincamajine. (3,4,5-Trimethoxycinnamoyl)-10-methoxyvincamajine**
[57800-04-1]
C₃₅H₄₀N₂O₈ 616.71
Alkaloid from the aerial parts of *Alstonia lanceolifera* (Apocynaceae). [α]_D²⁵ -134 (c, 1 in CHCl₃).
- 11-Methoxy: 11-Methoxyvincamajine**
[132242-26-3]
C₂₃H₂₈N₂O₄ 396.485
Alkaloid from the stem of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae). [α]_D²⁵ -14 (c, 0.25 in CHCl₃). λ_{max} 214; 249; 296 (MeOH).
- 11-Methoxy, O-Ac: 11-Methoxyvincamide**
[142750-28-5]
C₂₅H₃₀N₂O₅ 438.522
Alkaloid from leaves of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae). [α]_D²⁵ -7.5 (c, 1 in CHCl₃).
- 2-Epimer, 17-ketone, N-de-Me: Leepacine**
[135649-95-5]
C₂₁H₂₂N₂O₃ 350.416
Alkaloid from *Rhazya stricta* (Apocynaceae). Air- and light-sensitive. [α]_D²⁵ -91 (MeOH). λ_{max} 207 (log ε 4.23); 250 (log ε 3.59); 298 (log ε 3.12) (MeOH).
- 17-Epimer: Vincaminine**
[1748-05-6]
C₂₂H₂₆N₂O₃ 366.459
Alkaloid from aerial parts of *Vinca major* (Apocynaceae). Mp 257-259° Mp 274-275°.
- 17-Epimer, 11-methoxy: 17-Epi-11-methoxyvincamajine. 11-Methoxy-17-epi-vincamajine**
[132268-03-2]
C₂₃H₂₈N₂O₄ 396.485
Alkaloid from the stem bark of *Tonduzia pittieri* (*Alstonia pittieri*) (Apocynaceae). [α]_D²⁵ -12 (c, 0.5 in CHCl₃).
- Janot, M.-M. et al., *C. R. Hebd. Seances Acad. Sci.*, 1956, **243**, 85-87 (*Vincamedine*)
Gorman, M. et al., *Tet. Lett.*, 1963, 39-46 (*Quebrachidine, isol, uv, ir, pmr, ms, struct*)
Yuldashev, P.K. et al., *Dokl. Chem. (Engl. Transl.)*, 1964, **154**, 212-213; 1965, **163**, 644-645 (*Vincarine*)
Yuldashev, P.K. et al., *Khim. Prir. Soedin.*, 1966, **1**, 110-113; *Chem. Nat. Compd. (Engl. Transl.)*, 1966, **1**, 85-87 (*Vincarine*)
Crow, W.D. et al., *Aust. J. Chem.*, 1970, **23**, 2489-2501 (*Trimethoxycinnamoylvincamajine, O-Trimethoxybenzoylquebrachidine*)
Aynilian, G.H. et al., *J. Nat. Prod.*, 1974, **37**, 299-308 (*Vincamajine, Quebrachidine, isol, uv, ir, ms*)
Lewin, G. et al., *Phytochemistry*, 1975, **14**, 2067-2071 (*10-Methoxyvincamajine, trimethoxycinnamoyl, trimethoxybenzoyl*)
Chatterjee, A. et al., *Tet. Lett.*, 1978, 3879-3882 (*cmr*)
Stöckigt, J. et al., *Tet. Lett.*, 1979, 2615-2618 (*biosynth*)
Khalimirzaev, M.M. et al., *Khim. Prir. Soedin.*, 1980, 426 (*Vincarine*)
Yagudaev, M.R. et al., *Khim. Prir. Soedin.*, 1981, 608-613; *Chem. Nat. Compd. (Engl. Transl.)*, 1982, **17**, 442-446 (*Vincarine*)
Caron, C. et al., *Phytochemistry*, 1984, **23**, 2355-2357 (*10-Methoxyvincamide, 10-Methoxyvincamide N-oxide*)
Zhukovich, E.N. et al., *Khim. Prir. Soedin.*, 1985, **21**, 720; *Chem. Nat. Compd. (Engl. Transl.)*, 1986, **21**, 682 (*Vincaminine*)
Solans, X. et al., *Acta Cryst. C*, 1987, **43**, 1981-1983 (*cryst struct, abs config, Vincamedine*)
Ratnayake, C.K. et al., *Phytochemistry*, 1987, **26**, 868-870 (*19-Hydroxyvincamajine*)
Morfaux, A.-M. et al., *Phytochemistry*, 1990, **29**, 3345-3349 (*11-Methoxyvincamajine, 11-Methoxy-17-epi-vincamajine*)
Atta-ur-Rahman, et al., *Phytochemistry*, 1991, **30**, 1285-1293 (*Leepacine*)
Said, I.M. et al., *J. Nat. Prod.*, 1992, **55**, 1323-1324 (*O-(4-Hydroxy-3,5-dimethoxybenzoyl)vincamajine*)
Abe, F. et al., *Phytochemistry*, 1994, **35**, 249-252 (*Vincamajine 17-O-veratrate, N¹-Demethylvincamajine N¹-tri-O-methylgallate*)
Abe, F. et al., *Chem. Pharm. Bull.*, 1998, **46**, 1235-1238 (*17-trimethoxybenzoyl, 17-trimethoxybenzoyl N-oxide*)
Yu, J. et al., *J.O.C.*, 2005, **70**, 3963-3979 (*synth*)


Vincamicine V-114

[1362-02-3]

Bisindole alkaloid. Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Needles (MeOH). Mp 224-228° dec. [α]_D²⁶ +418 (c, 1 in CHCl₃). Mol. formula not recorded.
Svoboda, G.H. et al., *J. Pharm. Sci.*, 1961, **50**, 409-413 (*isol, uv, ir*)

Vincamine, BAN, INN V-115

Methyl 14,15-dihydro-14-hydroxyburnamenine-14-carboxylate, 9CI. Vincamarine. Minorine. Pervincamine. Vincafor



- $C_{21}H_{26}N_2O_3$ 354.448
Log P 3.82 (uncertain value) (calc).
- (+)-form** [1617-90-9]
Alkaloid from *Vinca minor*, *Vinca major*, *Vinca erecta*, *Vinca difformis* and *Tabernaemontana rigida* (Apocynaceae). Used in a wide variety of disorders to increase cerebral blood circulation. Antihypertensive agent. Mp 232-233° dec. (190-220°, 274° dec.). $[\alpha]_D^{25} +15.3$ (c, 0.11 in AcOH) (synthetic). Mp behaviour variable owing to phase change.
▶ LD₅₀ (mus, orl) 1000 mg/kg. YY8575000
Hydrochloride: Vincametrine
Mp 202-212°. Difficult to prepare owing to acid sensitivity of the alkaloid.
Picrate: Mp 228-229°.
- 14,15-Didehydro: 14,15-Dehydrovincamine. 14,15-Didehydrovincamine. Δ^{14} -Vincamine**
[32790-09-3]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from *Crioceras longiflorus*, *Crioceras dipladeniiflorus*, *Voacanga chaloniana* and *Amsonia elliptica* (Apocynaceae). Mp 221-223° (218°). $[\alpha]_D +116$ (c, 0.75 in $CHCl_3$).
- 16-Deoxy, 16,17-didehydro: Apovincamine, INN.** *Apv*
[4880-92-6]
[65826-02-0]
 $C_{21}H_{24}N_2O_2$ 336.433
Alkaloid from *Tabernaemontana rigida* (Apocynaceae) and prod. of dehydration of Vincamine. Cerebral vasodilator. Mp 160-161°. $[\alpha]_D^{26} +118$ (c, 2.11 in $CHCl_3$). Log P 4.89 (uncertain value) (calc).
- 16-Deoxy, 14,15,16,17-tetradehydro: Apo-14,15-dehydrovincamine. Apo- Δ^{14} -vincamine**
[50298-88-9]
 $C_{21}H_{22}N_2O_2$ 334.417
Alkaloid from the root bark of *Crioceras dipladeniiflorus* (Apocynaceae). Amorph. $[\alpha]_D +150$ (c, 1.13 in $CHCl_3$).
- 19-Hydroxy: 19-Hydroxyvincamine**
 $C_{21}H_{26}N_2O_4$ 370.447
Alkaloid from *Vinca minor* (Apocynaceae). Mp 222-225°.
- 19-Oxo: Vincaminine. Vincareine. 19-Oxovincamine**
[6880-35-9]
 $C_{21}H_{24}N_2O_4$ 368.432
Alkaloid from *Vinca minor* (Apocynaceae). Mp 208-210° dec. (205-206°). $[\alpha]_D +29.5$ (Py).
- 19-Oxo, methiodide:** Mp 222° dec.
- 11-Methoxy: Vincine. 11-Methoxyvincamine**
[4752-37-8]
 $C_{22}H_{28}N_2O_4$ 384.474
Alkaloid from *Vinca minor* (Apocynaceae). Needles. Mp 232° (212-214°) (block). $[\alpha]_D +39$ (c, 1 in Py). $[\alpha]_D -10$ (c, 1 in $CHCl_3$).
- 11-Methoxy, 14,15-didehydro: 14,15-Dehydrovincine. Δ^{14} -Vincine. 11-Methoxy- Δ^{14} -vincamine**
[32790-04-8]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Craspidospermum verticillatum* (Apocynaceae). Cryst. (EtOH). Mp 211°. $[\alpha]_D +70$ (c, 1 in $CHCl_3$).
- 11-Methoxy, 19-oxo: Vincinine. 11-Methoxy-19-oxovincamine**
[6880-41-7]
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Vinca minor* (Apocynaceae). Cryst. (Me₂CO). Mp 202-204° dec. $[\alpha]_D^{20} +24$ (Py).
- 11-Methoxy, 10-hydroxy, 14,15-didehydro: 14,15-Dehydro-10-hydroxy-11-methoxyvincamine. 14,15-Didehydro-10-hydroxy-11-methoxyvincamine**
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Ervatamia officinalis*. Amorph. powder. $[\alpha]_D^{20} +81.9$ (c, 0.11 in $CHCl_3$). λ_{max} 228 (log ϵ 4.29); 276 (log ϵ 3.72); 306 (log ϵ 3.87) (MeOH).
- 12-Methoxy, 14,15-didehydro: 14,15-Didehydro-12-methoxyvincamine. 12-Methoxy-14,15-didehydrovincamine. 12-Methoxy- Δ^{14} -vincamine**
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from the root bark of *Crioceras dipladeniiflorus* (Apocynaceae). Cryst. (Me₂CO). Mp 210-211°. $[\alpha]_D +96$ (c, 1 in $CHCl_3$).
- 10,11-Dimethoxy, 14,15-didehydro: 14,15-Dehydro-10,11-dimethoxyvincamine. 14,15-Didehydro-10,11-dimethoxyvincamine**
 $C_{23}H_{28}N_2O_5$ 412.485
Alkaloid from *Ervatamia officinalis*. Amorph. powder. $[\alpha]_D^{20} +86.3$ (c, 0.9 in $CHCl_3$). λ_{max} 229 (log ϵ 4.41); 276 (log ϵ 3.83); 299 (log ϵ 3.92); 303 (log ϵ 3.93) (MeOH).
- 16-Epimer: 16-Epivincamine. Base TR1**
[6835-99-0]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Vinca minor* and *Tabernaemontana rigida* (Apocynaceae). Mp 181-185°. $[\alpha]_D^{22} -36.4$ (c, 1.036 in $CHCl_3$). $[\alpha]_D^{26} -5$ (c, 2.69 in Py) (Base TR1). Base TR1, isol. from *Tabernaemontana rigida*, was prob. largely racemic.
- 16-Epimer, 14,15-didehydro: 14,15-Dehydro-16-epivincamine. 16-Epi- Δ^{14} -vincamine**
[32790-10-6]
 $C_{21}H_{24}N_2O_3$ 352.432
Alkaloid from the root bark of *Crioceras longiflorus* and *Crioceras dipladeniiflorus* and from seeds of *Amsonia elliptica* (Apocynaceae). Cryst. (Me₂CO). Mp 185° (181-182°). $[\alpha]_D +30$ (c, 1.2 in $CHCl_3$).
- 16-Epimer, 11-methoxy, 14,15-didehydro: 16-Epi-14,15-dehydrovincine. 16-Epi- Δ^{14} -vincine**
[32790-05-9]
 $C_{22}H_{26}N_2O_4$ 382.458
Alkaloid from *Craspidospermum verticillatum* (Apocynaceae). Noncryst.
- 16-Epimer, 11-methoxy, 10-hydroxy, 14,15-didehydro: 14,15-Dehydro-10-hydroxy-11-methoxy-16-epivincamine. 14,15-Didehydro-10-hydroxy-11-methoxy-16-epivincamine**
 $C_{22}H_{26}N_2O_5$ 398.458
Alkaloid from *Ervatamia officinalis*. Amorph. powder. $[\alpha]_D^{20} +8.4$ (c, 0.19 in $CHCl_3$). λ_{max} 228 (log ϵ 4.25); 277 (log ϵ 3.71); 301 (log ϵ 3.8); 306 (log ϵ 3.81); 311 (log ϵ 3.77) (MeOH).
- 16-Epimer, 10,11-dimethoxy, 14,15-didehydro: 14,15-Dehydro-10,11-dimethoxy-16-epivincamine. 14,15-Didehydro-10,11-dimethoxy-16-epivincamine**
 $C_{23}H_{28}N_2O_5$ 412.485
Alkaloid from *Ervatamia officinalis*. Amorph. powder. $[\alpha]_D^{20} +19$ (c, 0.3 in $CHCl_3$). λ_{max} 228 (log ϵ 4.42); 275 (log ϵ 3.86); 299 (log ϵ 3.91); 303 (log ϵ 3.92) (MeOH).
- 21-Epimer: 21-Epivincamine. Base TR2**
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Tabernaemontana rigida* (Apocynaceae). Cryst. (EtOH/ CH_2Cl_2). Mp 209-211°. $[\alpha]_D^{26} +1.8$ (c, 1.45 in Py). Biogenetically most probable struct. (partial racemate) for this substance.
- (-)-form** [38990-16-8]
Synthetic. Mp 227-228.5°. $[\alpha]_D^{20} -14.6$ (c, 0.5 in AcOH).
- (±)-form** [2122-39-6]
Alkaloid from *Tabernaemontana rigida* (Apocynaceae). Mp 235-236°. [54615-44-0]
Aldrich Library of NMR Spectra, 2nd edn., 1983, 2, 942A (nmr)
Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 1066D (ir)
Schlittler, E. et al., *Helv. Chim. Acta*, 1953, 36, 2017 (isol, uv, ir)
Trojaneček, J. et al., *Coll. Czech. Chem. Comm.*, 1961, 26, 867; 1962, 27, 2801 (Vincine, Vincaminine, Vincinine)
Trojaneček, J. et al., *Tet. Lett.*, 1961, 2, 702-706 (Vincamine, struct)
Plat, M. et al., *Bull. Soc. Chim. Fr.*, 1962, 1082 (ms, uv)
Strouf, O. et al., *Chem. Ind. (London)*, 1962, 2037 (Vincine)
Mokry, J. et al., *Tet. Lett.*, 1963, 999-1000; 1917-1919 (Vincamine, pmr, abs config, 16-Epivincamine)
Holubek, J. et al., *Tet. Lett.*, 1963, 4, 897-900 (Vincaminine, Vincinine, struct)
Machova, J. et al., *Arch. Int. Pharmacodyn. Ther.*, 1964, 150, 516 (Apovincamine, pharmacol)
Mokry, J. et al., *J. Nat. Prod.*, 1964, 27, 428 (19-Hydroxyvincamine)
Cava, M.P. et al., *J.O.C.*, 1968, 33, 1055-1059 ((+)-Vincamine, (±)-Vincamine, Apovincamine, Base TR1, Base TR2)
Cavé, A. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, 272, 1367 (14,15-Dehydrovincamine, 14,15-Dehydro-16-epivincamine)
Kan-Fan, C. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1971, 272, 1431 (14,15-Dehydrovincine, 16-Epi-14,15-dehydrovincine)
Hugel, G. et al., *C. R. Hebd. Seances Acad. Sci. Ser. C*, 1972, 274, 1350 (synth)
Neuss, N. et al., *Helv. Chim. Acta*, 1973, 56, 2660 (nmr)
Bruneton, J. et al., *Phytochemistry*, 1973, 12, 1475-1480 (Δ^{14} -Vincamine, 16-Epi- Δ^{14} -vincamine, 12-Methoxy- Δ^{14} -vincamine, Apo- Δ^{14} -vincamine)

- Taylor, W.I. *et al.*, *The Vinca Alkaloids*, M. Dekker, 1973, (pharmacol)
- Pfäffli, P. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1131-1145 (*Vincamine, synth, pmr, uv, ir*)
- Oppolzer, W. *et al.*, *Helv. Chim. Acta*, 1977, **60**, 1801-1810 (*Vincamine, synth*)
- Gibson, K.H. *et al.*, *Tetrahedron*, 1977, **33**, 833-836 ((±)-*Vincamine, synth*)
- Aimi, N. *et al.*, *Chem. Pharm. Bull.*, 1978, **26**, 1182 (*14,15-Dehydrovincamine*)
- Paracchini, S. *et al.*, *Farmaco, Ed. Sci.*, 1978, **33**, 573 (*synth, 14,15-Dehydrovincamine*)
- Rossey, G. *et al.*, *J.O.C.*, 1982, **47**, 4745-4749 (*Vincamine, synth*)
- Govindachari, T.R. *et al.*, *Indian J. Chem., Sect. B*, 1983, **22**, 531 (*synth*)
- Szabó, L. *et al.*, *Tetrahedron*, 1983, **39**, 3737-3747 ((+)-*Vincamine, synth*)
- Vereczkey, L. *et al.*, *Eur. J. Drug Metab. Pharmacokinet.*, 1985, **10**, 89 (*rev*)
- Takano, S. *et al.*, *Chem. Comm.*, 1986, 156-158 ((+)-*Vincamine, synth*)
- Lounasmaa, M. *et al.*, *Heterocycles*, 1986, **24**, 1663-1665 ((±)-*Vincamine, synth*)
- Negwer, M. *et al.*, *Organic-Chemical Drugs and their Synonyms*, 6th edn., Akademie-Verlag, 1987, 6203 (*synonyms*)
- Jucker, M. *et al.*, *Neurobiol. Aging*, 1990, **11**, 39 (*Apovincamine, pharmacol*)
- Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 1427
- Nagy, T. *et al.*, *Heterocycles*, 1997, **45**, 2007-2013 ((+)-*Vincamine, synth*)
- Martin, G.E. *et al.*, *J. Het. Chem.*, 1997, **34**, 695-699 (*pmr, N-15 nmr*)
- Schultz, A.G. *et al.*, *J.O.C.*, 1997, **62**, 1223-1229 ((+)-*Apovincamine, (+)-Vincamine, synth*)
- Desmaele, D. *et al.*, *J.O.C.*, 1997, **62**, 3890-3901 ((+)-*Vincamine, synth*)
- Alves, J.C.F. *et al.*, *Tetrahedron: Asymmetry*, 1999, **10**, 297-306 (*synth*)
- Kalaus, G. *et al.*, *Tetrahedron*, 2003, **59**, 5661-5666 (*Vincaminine, synth*)
- Zhang, H. *et al.*, *J. Nat. Prod.*, 2007, **70**, 54-59 (*Ervatamia officinalis constits*)
- England, D.B. *et al.*, *Org. Lett.*, 2007, **9**, 3249-3252 ((±)-21-*Epivincamine, synth*)
- Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, VLF000


Vincanovine V-116

Struct. unknown. Alkaloid from the aerial parts of *Vinca major* (Apocynaceae). Cryst. (MeOH). Mp 330° dec. $[\alpha]_D^{20}$ -20 (c, 0.49 in EtOH).

Kaul, J.L. *et al.*, *J. Nat. Prod.*, 1966, **29**, 26-34 (*isol, uv, ir*)

Vincarodine V-117

[1362-08-9]



Absolute Configuration

C₂₂H₂₆N₂O₅ 398.458

Trace alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Blades (CHCl₃/MeOH). Mp 253-256° dec. (236-238° dec., 246-247°). $[\alpha]_D^{26}$ -197.4 (c, 1 in CHCl₃). λ_{max} 230 (log ϵ 2.82); 272 (log ϵ 2.3); 298 (log ϵ 2.04) (EtOH).

Ac:

Cryst. (MeOH). Mp 180-185°.

14-Deoxy, 3,14-didehydro: Craspidospermine

[59373-42-1]

C₂₂H₂₄N₂O₄ 380.443

Alkaloid from the leaves of *Craspidospermum verticillatum* (Apocynaceae). Noncryst. $[\alpha]_D^{20}$ -59 (c, 1 in CHCl₃). λ_{max} 230 (log ϵ 4.58); 272 (log ϵ 3.95); 300 (log ϵ 3.57) (EtOH).

Demethoxy: Vincapusine

[80248-97-1]

C₂₁H₂₄N₂O₄ 368.432

Alkaloid from the roots and leaves of *Vinca pusilla*. Cryst. (EtOAc/petrol). Mp 263°. $[\alpha]_D$ -122 (c, 0.04 in CHCl₃). λ_{max} 228 (log ϵ 4.4); 271 (log ϵ 3.89); 280 (sh) (log ϵ 3.82); 292 (sh) (log ϵ 3.56) (EtOH).

Demethoxy, 14-deoxy, 3,14-didehydro: Criocerine

[55872-13-4]


C₂₁H₂₂N₂O₃ 350.416

Alkaloid from *Crioceras dipladeniiflorus* (Apocynaceae). Cryst. (MeOH). Mp 161-162°. $[\alpha]_D$ -27 (c, 0.85 in CHCl₃). λ_{max} 269 (log ϵ 3.96); 277 (log ϵ 4.51); 281 (log ϵ 3.86); 391 (log ϵ 3.59) (EtOH).

- Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*Vincarodine, isol, uv, ir*)
- Neuss, N. *et al.*, *Helv. Chim. Acta*, 1973, **56**, 2660-2666 (*Vincarodine, ir, uv, pmr, cmr, struct*)
- Kutney, J.P. *et al.*, *Heterocycles*, 1974, **2**, 73-78 (*Vincarodine, cryst struct*)
- Cordell, G.A. *et al.*, *J.O.C.*, 1974, **39**, 431-434 (*Vincarodine, isol, uv, ir, pmr, ms, struct*)
- Bruneton, J. *et al.*, *Phytochemistry*, 1975, **14**, 569-571 (*Criocerine, isol*)
- Kan-Fan, C. *et al.*, *Bull. Soc. Chim. Fr.*, 1976, 1227-1228 (*Craspidospermine*)
- Beugelmans, R. *et al.*, *Tet. Lett.*, 1976, **17**, 435-438 (*Craspidospermine, synth*)
- Mitra, A.K. *et al.*, *Phytochemistry*, 1981, **20**, 865-866 (*Vincapusine, isol*)
- Moldvai, I. *et al.*, *Synth. Commun.*, 1991, **21**, 965-967; 1992, **22**, 509-512 (*Criocerine, Craspidospermine, synth*)
- Moldvai, I. *et al.*, *J.O.C.*, 2006, **71**, 3768-3772 (*Vincapusine, synth*)

Vincarubine V-118

[107290-03-9]

C₄₃H₅₀N₄O₆ 718.891


Alkaloid from the leaves of *Vinca minor* (Apocynaceae). Cytotoxic agent. Dark-red solid. Mp 170° dec. $[\alpha]_D^{23}$ -550 (EtOH). λ_{max} 273 (ϵ 16200); 340 (ϵ

19330); 480 (ϵ 2450) (MeOH) (Berdy).

Proksa, B. *et al.*, *Planta Med.*, 1988, **54**, 214 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

Vincathicine V-119

[57665-10-8]

C₄₆H₅₆N₄O₉ 808.97

Alkaloid from *Catharanthus roseus* (Apocynaceae). Noncryst. λ_{max} 213 ; 264 ; 300 (MeOH) (Berdy).

▶ YY8620000

Sulfate:

Blades (MeOH). Mp 320° dec.

Methodide:

Cryst. (2-propanol). Mp 320° dec.


Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1964, **53**, 1227 (*isol, uv, ir, pmr*)

Tafur, S.S. *et al.*, *J.O.C.*, 1976, **41**, 1001 (*pmr, cmr, struct*)

Langlois, N. *et al.*, *Helv. Chim. Acta*, 1980, **63**, 793 (*struct*)

Vincatine V-120

[23185-53-7]



Absolute Configuration

C₂₂H₃₀N₂O₃ 370.491

Alkaloid from the leaves of *Vinca minor* (Apocynaceae). Prisms (EtOH). Mp 111-112°. $[\alpha]_D$ -13.5. Enantiomeric config. originally proposed.

Döpke, W. *et al.*, *Tet. Lett.*, 1969, 1701 (*isol, uv, ir, pmr, ms, struct*)


Ali, E. *et al.*, *Heterocycles*, 1982, **19**, 1667 (*synth, ir, pmr, cmr*)

Ali, E. *et al.*, *Tet. Lett.*, 1983, **24**, 2497 (*synth, cd*)

Danielli, B. *et al.*, *J.O.C.*, 1984, **49**, 547 (*abs config*)

Vincawajine V-121

[164176-14-1]



Absolute Configuration


C₂₄H₂₈N₂O₅ 424.496

Alkaloid from aerial parts of *Vinca major* (Apocynaceae). $[\alpha]_D^{25}$ -10.9 (c, 0.009 in CHCl_3).

Atta-ur-Rahman, et al., *Phytochemistry*, 1995, **38**, 1057 (isol, uv, ir, pmr, cmr, ms, struct)

Vincetene V-122

4-(7,9,10,11,11a,12-Hexahydro-2,3-dimethoxybenzo[f]pyrrolo[1,2-b]isoquinolin-6-yl)-3-buten-2-ol, 9CI
[72169-92-7]



$\text{C}_{22}\text{H}_{27}\text{NO}_3$ 353.46

Trace alkaloid from *Cynanchum vince-toxicum* (Asclepiadaceae). Unstable liq.

1',2'-Dihydro, 3'-ketone: 4-

(7,9,10,11,11a,12-Hexahydro-2,3-dimethoxybenzo[f]pyrrolo[1,2-b]isoquinolin-6-yl)-2-butanone, 9CI
[402943-22-0]

$\text{C}_{22}\text{H}_{27}\text{NO}_3$ 353.46

Alkaloid from *Cynanchum komarovii*. Needles. Mp 164-166°. $[\alpha]_D^{25}$ -71 (c, 0.001 in CHCl_3). λ_{max} 235 (log ϵ 3.93); 315 (log ϵ 3.5); 330 (log ϵ 3.54) (MeOH).

Budzikiewicz, H. et al., *Annalen*, 1979, 1212-1231 (isol, uv, ir, ms, pmr, struct)

Jirovsky, D. et al., *Monatsh. Chem.*, 2001, **132**, 765-768 (config)

Vincolidine V-123

[1362-11-4]


$\text{C}_{23}\text{H}_{26}\text{N}_2\text{O}_3$ 378.47

Struct. unknown. Alkaloid from *Catharanthus roseus* (Apocynaceae). Blades (Me_2CO). Mp 165-170°. $[\alpha]_D^{25}$ -25.4 (c, 1 in CHCl_3). pK_a 5.45 (33% DMF aq.).

Svoboda, G.H. et al., *J. Nat. Prod.*, 1964, **27**, 203-213 (isol, ir, uv)

Vincoline V-124

[11034-66-5]



Absolute configuration

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Catharanthus roseus*, *Vinca libanotica* and *Melodinus balansae* (Apocynaceae). CNS depressant. Mp 230-233°. $[\alpha]_D^{25}$ -32.6 (c, 1 in CHCl_3).

► CJ0140000

Me ether: **Kitraline**

[59086-88-3]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Catharanthus ovalis* (Apocynaceae). $[\alpha]_D$ -9 (c, 0.47 in CHCl_3).

19-Epimer: **Melobaline**

[38225-08-0]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from *Melodinus balansae* (Apocynaceae). Cryst. (MeOH). Mp 200-204°. $[\alpha]_D^{25}$ +205 (c, 1 in CHCl_3). Incorr. struct. originally proposed.

19-Epimer, Me ether: **Kitramine**

[59129-64-5]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Catharanthus ovalis* (Apocynaceae). $[\alpha]_D$ -46 (c, 0.45 in CHCl_3).

Svoboda, G.H. et al., *J. Nat. Prod.*, 1964, **27**, 203 (isol, uv, ir)

Mehri, M.H. et al., *Bull. Soc. Chim. Fr.*, 1972, 3291 (isol, Melobaline)

Taylor, W.I. et al., *Catharanthus Alkaloids*, M. Dekker, 1975, (pharmacol)

Andriamialisoa, R.Z. et al., *Tet. Lett.*, 1976, 163 (pmr, ms, struct)

Damak, M. et al., *Tet. Lett.*, 1976, 167 (pmr, cmr, ms)

Langlois, N. et al., *J.O.C.*, 1979, **44**, 2468

(synth, Kitraline, Kitramine)


Langlois, N. et al., *Phytochemistry*, 1979, **18**, 467 (Kitraline, Kitramine)

Vincorine V-125

Methyl 10-methoxy-1-methyl-2,4(1H)-cyclo-3,4-secoakuammilan-17-oate, 9CI.

Vincovine. Desformocabuamine

[33023-08-4]



Absolute Configuration

$\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_3$ 368.475

Alkaloid from *Vinca minor*, *Alstonia vitiensis* and *Cabucala erythrocarpa* (Apocynaceae). Cryst. (EtOH). Mp 93-94°. $[\alpha]_D^{20}$ -139 (c, 1 in EtOH). λ_{max} 254 (log ϵ 4.01); 326 (log ϵ 3.58) (EtOH).

N-De-Me: **1-Norvincorine**

[109028-35-5]

$\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3$ 354.448

Alkaloid from *Vinca minor* (Apocynaceae). Cryst. (Me_2CO). Mp 219-222°. $[\alpha]_D^{20}$ -165 (CHCl_3).

N-De-Me, N-formyl: **Demethoxyalstonamide**

[135824-72-5]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Alstonia macrophylla* (Apocynaceae). $[\alpha]_D$ +74 (c, 0.005 in CHCl_3).

11-Methoxy, N-de-Me, N-formyl: **Alstonamide**

[135792-64-2]

$\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_5$ 412.485

Alkaloid from *Alstonia macrophylla* (Apocynaceae). $[\alpha]_D$ +82 (c, 0.062 in CHCl_3).

Demethoxy, 3-oxo: **Vincoridine**

[21290-53-9]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$ 352.432

Alkaloid from *Vinca minor* (Apocynaceae). Cryst. (heptane). Mp 159-160°. $[\alpha]_D^{25}$ -158 (c, 0.87 in CHCl_3). 16-Config. uncertain.

Mokry, J. et al., *Experientia*, 1962, **18**, 564-565 (isol, uv, ir)

Kompiš, I. et al., *Coll. Czech. Chem. Comm.*, 1968, **33**, 4328-4336 (Vincoridine)

Meisel, H. et al., *Tet. Lett.*, 1971, 1285-1286 (uv, ir, pmr, ms, struct)

Douzoua, L. et al., *Phytochemistry*, 1974, **13**, 1994-1995 (isol, uv, pmr, ms)

Mansour, M. et al., *Phytochemistry*, 1974, **13**, 2861-2863 (struct)

Das, B.C. et al., *Tet. Lett.*, 1974, 4299-4302 (cmr)

Mamatas-Kalamaras, S. et al., *Phytochemistry*, 1975, **14**, 1637-1639 (uv, ms, pmr, struct)


Proska, B. et al., *Planta Med.*, 1987, **53**, 120 (Norvincorine)

Atta-ur-Rahman, et al., *J. Nat. Prod.*, 1991, **54**, 750-754 (Alstonamide, Demethoxyalstonamide)

Vincosamide V-126

Vincoside lactam

[23141-27-7]



$\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_8$ 498.532

Alkaloid from *Adina rubescens* (Rubiaceae). Plates + 2.5H₂O (MeOH aq.). Mp 201-202°. $[\alpha]_D$ -118 (c, 2.8 in MeOH).

N¹-β-D-Glucopyranosyl: **N-β-D-Glucopyranosylvincosamide**

$\text{C}_{32}\text{H}_{40}\text{N}_2\text{O}_{13}$ 660.674

Alkaloid from the leaves of *Psychotria leiocarpa*. Amorph. yellow powder. λ_{max} 225 ; 275 (sh) (MeOH).

3'-(3,4-Dihydroxycinnamoyl): **Rubescine**

3'-Caffeoylvincoside lactam

[34382-43-9]

$\text{C}_{35}\text{H}_{36}\text{N}_2\text{O}_{11}$ 660.676

Alkaloid from *Adina rubescens* (Rubiaceae). Yellow amorph. powder. $[\alpha]_D^{25}$ -121 (c, 0.90 in MOH). Subst. in the glucosyl residue.

6'-(3-Hydroxy-4-methoxycinnamoyl):

Rhynchophine

[84638-29-9]

$\text{C}_{36}\text{H}_{38}\text{N}_2\text{O}_{11}$ 674.703

Alkaloid from leaves of *Uncaria rhynchophylla* (Rubiaceae). Amorph. powder. $[\alpha]_D^{27}$ -45 (c, 0.7 in MeOH). Subst. in the glucosyl residue.

10-Hydroxy: **10-Hydroxyvincoside lactam**

10-Hydroxyvincosamide

[135531-63-4]

$\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_9$ 514.531

Alkaloid from fruits of *Alangium lamarkii*. Amorph. powder. $[\alpha]_D^{24}$ -120 (c, 0.79 in MeOH).

10-Hydroxy, 10-O-β-D-glucopyranoside:

10-Glucosyloxyvincoside lactam

[38971-68-5]

C₃₂H₄₀N₂O₁₄ 676.673Alkaloid from fresh shoots of *Adina rubescens* (Rubiaceae). Noncryst. [α]_D²⁵ -86 (CHCl₃) (as octa-Ac).**19-Hydroxy, 3,14-didehydro: Naucleoside A**
C₂₆H₂₈N₂O₉ 512.515Alkaloid from the bark of *Nauclea orientalis*. Amorph. orange-yellow solid (MeOH). Mp 171-172°. [α]_D²⁵ -48.6 (c, 0.15 in MeOH). λ_{\max} 216 (log ϵ 4.02); 260 (log ϵ 3.57); 372 (log ϵ 3.92); 390 (log ϵ 3.89) (MeOH).**19-Hydroxy, 14,15-didehydro: Naucleoside**
C₂₆H₂₈N₂O₉ 512.515Alkaloid from the stems of *Nauclea officinalis* (Rubiaceae). Cryst. [α]_D²⁵ -149 (c, 0.1 in EtOH aq.). Mp >310°.**3-Epimer: Strictosamide. Strictosidine lactam**



[23141-25-5]

C₂₆H₃₀N₂O₈ 498.532Alkaloid from *Rhazya stricta* (Apocynaceae) and *Nauclea latifolia* (Rubiaceae). Foam. [α]_D -71 (c, 1.4 in MeOH). [α]_D -77 (CHCl₃).**3-Epimer, 2'-Ac: 2'-O-Acetylstrictosamide**C₂₈H₃₂N₂O₉ 540.569Alkaloid from *Mostuea brunonis*. [α]_D -63 (c, 0.18 in MeOH). λ_{\max} 226 (log ϵ 4.5); 313 (log ϵ 3.2); 320 (log ϵ 3.2) (MeOH).**3-Epimer, 10-hydroxy: 10-Hydroxystrictosamide**

[169871-22-1]

C₂₆H₃₀N₂O₉ 514.531Alkaloid from leaves of *Nauclea orientalis* (Rubiaceae). Extremely unstable in basic soln.**3-Epimer, 19-hydroxy, 14,15-didehydro: Naucleoside B**C₂₆H₂₈N₂O₉ 512.515Alkaloid from the bark of *Nauclea orientalis*. Amorph. orange-yellow solid (MeOH). Mp 189-190°. [α]_D²⁵ -58.2 (c, 0.15 in MeOH). λ_{\max} 212 (log ϵ 3.68); 260 (log ϵ 3.2); 372 (log ϵ 3.55); 390 (log ϵ 3.52) (MeOH).**3-Epimer, aglycone, 21-Me ether: 21-O-Methylstrictosamide aglycone**C₂₁H₂₂N₂O₃ 350.416Alkaloid from *Sarcocephalus latifolius*. Oil. [α]_D -0.1 (c, 0.06 in CHCl₃).**3-Epimer, aglycone, 21-Et ether: 21-O-Ethylstrictosamide aglycone**C₂₂H₂₄N₂O₃ 364.443Alkaloid from *Sarcocephalus latifolius*. Oil.Battersby, A.R. *et al.*, *J.C.S.(C)*, 1969, 1193 (synth, uv, ir, *Vincosamide*, *Strictosamide*)De Silva, K.T.D. *et al.*, *Chem. Comm.*, 1971, 905 (occur, struct, *Strictosamide*)Blackstock, W.P. *et al.*, *Chem. Comm.*, 1971, 910 (occur, struct)Blackstock, W.P. *et al.*, *Tet. Lett.*, 1971, 3727 (*Rubescine*)Brown, R.T. *et al.*, *Tet. Lett.*, 1972, 3063 (uv, ir, pmr, ms, cd, struct, *10-Glucosyloxyvincoside lactam*)Aimi, N. *et al.*, *Chem. Pharm. Bull.*, 1982, 30, 4046 (*Rhynchophine*)Lin, M. *et al.*, *Yaouxue Xuebao*, 1989, 24, 32-36; *CA*, 111, 74776q (*Naucleoside*)Erdelmeier, C.A.J. *et al.*, *Planta Med.*, 1991, 57, 149 (*10-Hydroxystrictosamide*)Itoh, A. *et al.*, *J. Nat. Prod.*, 1995, 58, 1228 (*10-Hydroxyvincoside lactam*)Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1997, 45, 1231-1233 (*Strictosamide*, pmr, cmr)Takayama, H. *et al.*, *Heterocycles*, 1998, 49, 49-52 (*10-Hydroxystrictosamide*)Dai, J.-R. *et al.*, *J. Nat. Prod.*, 1999, 62, 1427-1429 (*2'-Acetylstrictosamide*)Zhang, Z. *et al.*, *J. Nat. Prod.*, 2001, 64, 1001-1005 (*Naucleosides*)Abreu, P. *et al.*, *Nat. Prod. Lett.*, 2001, 15, 43-48 (*Strictosamide aglycone ethers*)Henriques, A.T. *et al.*, *Phytochemistry*, 2004, 65, 449-454 (*N-Glucosylvincosamide*)**Vincovalicine**

[61991-58-0]

C₄₆H₅₄N₄O₁₀ 822.953Alkaloid from the aerial parts of *Catharanthus ovalis* (Apocynaceae). [α]_D -45 (c, 0.45 in CHCl₃).Langlois, N. *et al.*, *Helv. Chim. Acta*, 1980, 63, 793-805 (isol, uv, ir, pmr, ms, cd, struct)**Vincristine, BAN, INN**22-Oxovincalcoloblastine, 9CI. *Leurocristine*. *Cristovin*. *Kyocristine*. *Leucid*. *Oncovin*. *Pericristine*. *Vincosid*. *Vincrisul*. *L 37231*. *Onkovin* [57-22-7]C₄₆H₅₆N₄O₁₀ 824.969Alkaloid from *Catharanthus roseus* (Apocynaceae). Antineoplastic agent. Used in combination therapies for treatment of acute lymphoblastic leukaemia, Hodgkin's disease and other neoplastic diseases. Blades (MeOH). Mp 218-220° dec. [α]_D²⁵ +26.2 (1,2-dichloroethane). Log P 2.9 (uncertain value) (calc). λ_{\max} 220 (log ϵ 44670); 255 (log ϵ 16220); 296 (log ϵ 15135) (EtOH) (Berdy).► Adverse human effects reported when used therapeutically or from occup. exposure. Skin irritant. Exp. reprod. and teratogenic effects. LD₅₀ (mus, ipr) 5.2 mg/kg. LD₅₀ (rat, ivn) 1 mg/kg. OH6300000**V-127**Sulfate salt: *Vincristine sulfate*, JAN, USAN. *Marqibo*. NSC 67574

[2068-78-2]

Cryst. (EtOH).


► OH6340000

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 1068C (ir)

Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1961, 24, 173 (isol, uv, ir)Neuss, N. *et al.*, *J.A.C.S.*, 1962, 84, 1509; 1964, 86, 1440 (uv, ir, pmr, struct)Moncrief, J.W. *et al.*, *J.A.C.S.*, 1965, 87, 4963 (cryst struct)Johnson, I.S. *et al.*, *Cancer Chemother. Rep.*, 1968, 52, 455 (numerous papers)Burns, J.H. *et al.*, *Anal. Profiles Drug Subst.*, 1972, 1, 463 (rev, uv, ir, pmr, anal)Taylor, W.I. *et al.*, *Catharanthus Alkaloids*, M. Dekker, 1975, (pharmacol)Sieber, S.M. *et al.*, *Cancer Treat. Rep.*, 1976, 60, 1127-1139 (pharmacol)Potier, P. *et al.*, *J.A.C.S.*, 1979, 101, 2243 (synth)Brade, W. *et al.*, *Contrib. Oncol.*, Eds. S. Karger, Basel, Switz., 1981, 6, (bibl)Rao, K.S.P.B. *et al.*, *J. Med. Chem.*, 1985, 28, 1079 (tox)Kuehne, M.E. *et al.*, *Alkaloids (Academic Press)*, 1990, 37, 77 (synth, rev)Neuss, N. *et al.*, *Alkaloids (Academic Press)*, 1990, 37, 229 (pharmacol)Martindale, *The Extra Pharmacopoeia*, 30th edn., Pharmaceutical Press, 1993, 505Kuboyama, T. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, 101, 11966-11970 (synth)Lewis, R.J. *et al.*, *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, LEY000; LEZ000**Vindolicine**

10,10'-Methylenebisvindoline


[1362-14-7]

V-129

Absolute Configuration

C₅₁H₆₄N₄O₁₂ 925.086Alkaloid from *Catharanthus longifolius* and *Catharanthus roseus* (Apocynaceae). Plates (Et₂O). Mp 248-251° Mp 265-267° dec. (double Mp).Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1961, 50, 409 (isol, uv, ir)Rabaron, A. *et al.*, *Plant. Med. Phytother.*, 1973, 7, 53; *CA*, 79, 63524a (uv, ir, ms, isol, struct)**Vindolidine†***Vindorosine*, 8CI. *Demethoxyvindoline*

[5231-60-7]

V-130

(+) -form

C₂₄H₃₀N₂O₅ 426.511

▶ NM2410000

(+)-formAlkaloid from *Vinca pusilla* (Apocynaceae). Cryst. (C₆H₆). Mp 164-166°. [α]_D +23.5 (CHCl₃).**11-Methoxy: Vindoline**

[2182-14-1]

C₂₅H₃₂N₂O₆ 456.538Main alkaloid of *Catharanthus roseus* also in *Vinca pusilla* (Apocynaceae). Active against gram-positive bacteria and fungi. Mp 154-155° (172-174°). [α]_D²⁷ +42 (CHCl₃). λ_{max} 212; 250; 304 (EtOH) (Berdy).

▶ CJ0120000

11-Methoxy, O-de-Ac: Deacetylvindoline
[3633-92-9]C₂₃H₃₀N₂O₅ 414.5Alkaloid from *Catharanthus roseus* (Apocynaceae). Cryst. (C₆H₆/petrol or Me₂CO/petrol). Mp 163-165°.**11-Methoxy, deacetoxy: Desacetoxylvindoline**C₂₃H₃₀N₂O₄ 398.501Minor alkaloid from the cotyledons of dark-grown seedlings of *Catharanthus roseus* (Apocynaceae). Biosynth. precursor of Vindoline.**(-)-form**Alkaloid from *Catharanthus roseus*, *Catharanthus longifolius* and *Catharanthus trichophyllus* (Apocynaceae). Needles (C₆H₆/petrol). Mp 167°. [α]_D¹⁶ -31 (CHCl₃).**(±)-form** [33190-27-1]

Mp 182-185°.

O-De-Ac: Catharosine


[2564-23-0]

C₂₂H₂₈N₂O₄ 384.474Alkaloid from *Catharanthus roseus* (Apocynaceae). Needles (C₆H₆/petrol). Mp 141-143°. [α]_D²⁰ 0 (CHCl₃).**11-Methoxy: [57794-53-3]**

Synthetic. Mp 203-205°.

Gorman, M. *et al.*, *J.A.C.S.*, 1962, **84**, 1058-1059 (*ms, pmr, struct*)Neuss, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, 1509 (*pmr*)Moza, B.K. *et al.*, *Coll. Czech. Chem. Comm.*, 1963, **28**, 1419; 1427; 1964, **29**, 1913 (*isol, uv, ir, pmr, ms, ord, struct*)Gröger, D. *et al.*, *Naturwissenschaften*, 1964, **51**, 637; 1965, **52**, 132 (*isol, ir, uv, ms, synth, deriv*)McCapra, F. *et al.*, *Chem. Comm.*, 1965, 537 (*biosynth*)Moza, B.K. *et al.*, *Chem. Ind. (London)*, 1965, 1260 (*Catharosine*)Büchi, G. *et al.*, *J.A.C.S.*, 1971, **93**, 3299 (*synth*)Rasoanaivo, P. *et al.*, *Phytochemistry*, 1972, **11**, 2616 (*isol*)Chatterjee, A. *et al.*, *Indian J. Chem.*, 1973, **11**, 7 (*isol, uv, ir*)Wenkert, E. *et al.*, *J.A.C.S.*, 1973, **95**, 4990 (*cmr*)Taylor, W.I. *et al.*, *Catharanthus Alkaloids*, Marcel Dekker, 1975, (*pharmacol*)Ando, M. *et al.*, *J.A.C.S.*, 1975, **97**, 6880 (*synth*)Cordell, G.A. *et al.*, *J. Pharm. Sci.*, 1976, **65**, 366 (*isol*)Takano, S. *et al.*, *Heterocycles*, 1977, **6**, 1699 (*synth*)Hernandez, N.M.R. *et al.*, *Rev. Cubana Farm.*, 1977, **11**, 249-255 (*activity*)Kutney, J.P. *et al.*, *J.A.C.S.*, 1978, **100**, 4220 (*synth*)Nagakura, N. *et al.*, *J.C.S. Perkin I*, 1979, 2308 (*biosynth*)Hernandez, N.M.R. *et al.*, *Rev. Cubana Med. Trop.*, 1979, **31**, 199-204 (*activity*)Wenkert, E. *et al.*, *Heterocycles*, 1981, **15**, 255 (*cmr*)Veenstra, S.J. *et al.*, *J.A.C.S.*, 1981, **103**, 4645 (*synth*)Kutney, J.P. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 2088 (*biosynth*)Balsevich, J. *et al.*, *Heterocycles*, 1986, **24**, 2415 (*Desacetoxylvindoline*)Winkler, J.D. *et al.*, *J.A.C.S.*, 1990, **112**, 8971 (*synth*)Zhou, S.-Z. *et al.*, *Org. Lett.*, 2002, **4**, 443-445 (*synth*)Ruszkowska, J. *et al.*, *Acta Cryst. C*, 2004, **60**, 377-380 (*Vindoline, Vindolidine, cryst struct*)Choi, Y. *et al.*, *Org. Lett.*, 2005, **7**, 4539-4542 (*synth*)Ishikawa, H. *et al.*, *J.A.C.S.*, 2006, **128**, 10596-10612 (*synth*)**Vindolidine†****V-131**C₄₈H₆₄N₄O₁₀ 857.054Struct. unknown. Bisindole alkaloid. Not the same as Vindolidine, V-130. Alkaloid from *Catharanthus roseus* (Apocynaceae). Microcryst. (MeOH). Mp 244-250° dec. [α]_D²⁶ -113.2 (c, 1 in CHCl₃).Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1962, **51**, 518-523 (*isol, uv, ir*)**Vindolinine****V-132***Methyl 6,7-didehydro-2,20-cycloaspido-permidine-3-carboxylate, 9CI*

[5980-02-9]



Absolute configuration

C₂₁H₂₄N₂O₂ 336.433Alkaloid from *Catharanthus roseus*, *Echium lycopsis* and *Melodinus balansae* (Apocynaceae). Antihyperglycaemic agent. Active against fungi. Mp 214-218°. Log P 1.55 (uncertain value) (calc).*Hydrochloride (1:2)*: Mp 210-212° dec
Mp 250-252°.*Picrate*: Mp 268-272°.**N⁴-Oxide: 19R-Vindolinine N-oxide. Vindolinine N-oxide**

[41682-54-6]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from *Catharanthus roseus* and *Melodinus* spp.**14,15-Dihydro: Pseudokopsinine. 14,15-Dihydrovindolinine**

[17172-16-6]

C₂₁H₂₆N₂O₂ 338.449Alkaloid from *Vinca erecta* epigeal parts (Apocynaceae). Shows antihypertensive activity. [α]_D +30.4 (c, 1.51 in MeOH). Log P 2.04 (uncertain value) (calc).**14,15-Dihydro, N⁴-oxide: Pseudokopsinine N^b-oxide**

[54387-75-6]

C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Vinca erecta*. Cryst. (petrol). Mp 186-188°.**14,15-Dihydro, 15α-hydroxy: 15α-Hydroxy-14,15-dihydrovindolinine**C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Melodinus morsei* (Apocynaceae). Granular cryst. (EtOH). Mp 172-174°. [α]_D¹⁵ -26 (c, 0.88 in EtOH).**16β-Hydroxy: 16β-Hydroxy-19R-vindolinine**C₂₁H₂₄N₂O₃ 352.432Alkaloid from aerial parts of *Melodinus hemsleyanus* (Apocynaceae). Needles (Me₂CO). Mp 185° dec. [α]_D¹⁰ -42.9 (c, 0.2 in CHCl₃).**16β-Hydroxy, N^b-oxide: 16β-Hydroxy-19R-vindolinine N-oxide**

[222557-89-3]

C₂₁H₂₄N₂O₄ 368.432Alkaloid from *Melodinus hemsleyanus*.**19-Epimer: 19S-Vindolinine. 19-Epivindolinine**C₂₁H₂₄N₂O₂ 336.433Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Cryst. (EtOH). Mp 200°. [α]_D +40 (MeOH). Erroneously descr. as 16-epi-19S-vindolinine in the original ref. Isomerises in CHCl₃ to Vindolinine.**19-Epimer, N^b-oxide: 19S-Vindolinine N-oxide. 19-Epivindolinine N-oxide**

[88720-99-4]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from *Catharanthus roseus* (Apocynaceae). Descr. in the lit. as 16-Epi-19S-vindolinine N-oxide but this should presumably be revised.**16-Epimer, 14,15-dihydro, 15α-hydroxy: 15-Hydroxy-14,15-dihydro-16-epivindolinine**C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Melodinus morsei* (Apocynaceae). Amorph. powder + 1H₂O. [α]_D¹⁵ +28 (c, 1.00 in EtOH).**19-Epimer, 16β-hydroxy: 16β-Hydroxy-19S-vindolinine. 16β-Hydroxy-19-epivindolinine**C₂₁H₂₄N₂O₃ 352.432Alkaloid from aerial parts of *Melodinus hemsleyanus* (Apocynaceae). Prisms (Me₂CO). Mp 220° dec. [α]_D¹⁰ -87.5 (c, 0.03 in CHCl₃).**19-Epimer, 16β-hydroxy, N⁴-oxide: 16β-Hydroxy-19S-vindolinine N-oxide. 16β-Hydroxy-19-epivindolinine N-oxide**

[222557-90-6]


C₂₁H₂₄N₂O₄ 368.432Alkaloid from *Melodinus hemsleyanus*.Janot, M.M. *et al.*, *Bull. Soc. Chim. Fr.*, 1959, 891-892 (*isol*)Djerassi, C. *et al.*, *Helv. Chim. Acta*, 1964, **47**, 827-836 (*ms, struct*)Nasyrov, S.-M. *et al.*, *Chem. Comm.*, 1974, 979 (*cryst struct, Pseudokopsinine*)Ahond, A. *et al.*, *J.A.C.S.*, 1974, **96**, 633-634 (*cmr, struct*)Sharipov, M.R. *et al.*, *Khim. Prir. Soedin.*, 1974, **10**, 413-414; *Chem. Nat. Compd. (Engl. Transl.)*, 1974, **10**, 422-423(*Pseudokopsinine N-oxide*)

- Rasonaivo, P. *et al.*, *Tet. Lett.*, 1974, 3669-3672 (abs config)
- Riche, C. *et al.*, *Acta Cryst. B*, 1976, **32**, 1975-1980 (cryst struct)
- Nasyrov, S.-M. *et al.*, *Khim. Prir. Soedin.*, 1976, **12**, 197-206; *Chem. Nat. Compd. (Engl. Transl.)*, 1976, **12**, 176-183 (abs config, Pseudokopsinine)
- Hernandez, N.M.R. *et al.*, *Rev. Cubana Farm.*, 1977, **11**, 249-255 (Vindolinine, activity)
- Hernandez, N.M.R. *et al.*, *Rev. Cubana Med. Trop.*, 1979, **31**, 199-204 (Vindolinine, activity)
- Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1983, **22**, 1021-1023 (epimer)
- Atta-ur-Rahman, *et al.*, *Planta Med.*, 1983, **49**, 124-125 (diepimer oxide)
- Atta-ur-Rahman, *et al.*, *Z. Naturforsch., B*, 1986, **41**, 386-392 (epimer)
- Guo, L.-W. *et al.*, *Phytochemistry*, 1993, **34**, 563-566 (16 β -Hydroxy-19R-vindolinine, 16 β -Hydroxy-19S-vindolinine)
- He, Y.-L. *et al.*, *Phytochemistry*, 1994, **37**, 1055-1057 (15 α -Hydroxy-14,15-dihydrovindolinine, 15 α -Hydroxy-14,15-dihydro-16-epivindolinine)
- Yan, K. *et al.*, *Zhongcaoyao*, 1998, **29**, 793-795 (16-Hydroxyvindolinine N-oxides)

Vingramine

V-133

[215316-92-0]

C₄₆H₅₆N₄O₈ 792.97

Alkaloid from the seeds of *Catharanthus roseus*. Shows cytotoxic activity. Amorph. [α]_D -270.4 (c, 0.5 in CHCl₃). λ_{\max} 217 (sh) (log ϵ 4.69); 231 (log ϵ 4.8); 280 (log ϵ 4.33); 303 (sh) (log ϵ 4.3) (MeOH).

N¹-Me: Methylvingramine

[215316-95-3]

C₄₇H₅₈N₄O₈ 806.997

Alkaloid from the seeds of *Catharanthus roseus*. Amorph. [α]_D -278 (c, 0.5 in CHCl₃). λ_{\max} 216 (log ϵ 4.89); 234 (sh) (log ϵ 4.57); 280 (log ϵ 4.12); 297 (log ϵ 4.12); 310 (sh) (log ϵ 4.11) (MeOH).

Jossang, A. *et al.*, *J.O.C.*, 1998, **63**, 7162-7167 (isol, props, struct, activity)

Vinomine

V-134

[11075-79-9]

C₁₉H₂₂N₂O₃ 326.394

Struct. unknown. Alkaloid from *Vinca minor* (Apocynaceae). Prisms (MeOH). Mp 267-268°.

Meisel, H. *et al.*, *Pharmazie*, 1971, **26**, 182 (isol, uv)

Vinosidine

V-135

[1362-19-2]

C₂₂H₂₆N₂O₅ 398.458

Struct. unknown. Alkaloid from the roots of *Catharanthus roseus* and from *Catharanthus lanceus* (Apocynaceae). Needles (MeOH).


Mp 253-257° dec. Cordell *et al* list Vinosidine as a bisindole alkaloid (C₄₄H₅₂N₄O₁₀).

- Svoboda, G.H. *et al.*, *J. Nat. Prod.*, 1963, **26**, 141-153 (isol, uv, ir)
- Svoboda, G.H. *et al.*, *J. Pharm. Sci.*, 1963, **52**, 407-408 (isol, uv, ir)
- Blomster, R.N. *et al.*, *J. Pharm. Sci.*, 1967, **56**, 284-286 (isol)
- Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 267 (rev)

Vinoxine

V-136

[55856-81-0]



Absolute Configuration

C₂₀H₂₄N₂O₃ 340.421

Indole alkaloid most closely related to an Akuammicine but with C-6/C-7 bond cleavage and N¹/C-16 bond formation. Alkaloid from *Vinca minor* (Apocynaceae). Noncryst. [α]_D²⁴ -18.6 (CHCl₃). λ_{\max} 221 (log ϵ 4.56); 273 (log ϵ 3.95); 282 (log ϵ 3.9); 291 (log ϵ 3.77) (EtOH).

Hydrochloride: Mp 216-218° dec. [α]_D²⁰ -25 (c, 0.89 in CHCl₃).

Ac:

Noncryst. [α]_D²⁴ -30.8 (CHCl₃).

- Mokry, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1967, **32**, 2523-2531 (isol, uv, ir, ms)
- Volticky, Z. *et al.*, *Coll. Czech. Chem. Comm.*, 1977, **42**, 548-552 (uv, ms, pmr, cmr, struct)
- Bosch, J. *et al.*, *Tet. Lett.*, 1984, **25**, 3119-3122 (synth, stereochem)
- Bennasar, M.-L. *et al.*, *J.O.C.*, 1993, **58**, 7756-7767 (synth)

Vinsedicine

V-137

Bisindole alkaloid. Struct. unknown. Alkaloid from the seeds of *Catharanthus roseus* (Apocynaceae). Mol. formula not reported.

- Gorman, M. *et al.*, *J. Nat. Prod.*, 1965, **28**, 259 (occur)
- Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 267 (rev)

Vinsedine

V-138

Bisindole alkaloid. Struct. unknown. Alkaloid from the seeds of *Catharanthus roseus* (Apocynaceae). Mol. formula not reported.


- Gorman, M. *et al.*, *J. Nat. Prod.*, 1965, **28**, 259 (occur)
- Cordell, G.A. *et al.*, *Alkaloids (Academic Press)*, 1981, **20**, 1; 267 (rev)

1-Vinyl- β -carboline

V-139

1-Ethenyl-9H-pyrido[3,4-b]indole, 9CI.

Pavettine
[20127-60-0]

C₁₃H₁₀N₂ 194.235

Alkaloid from *Pavetta lanceolata* (Rubiaceae), the bryozoans *Cribricellina cribraria* and *Costaticella hastata* and from *Soulamea raxinifolia* and *Pavetta lanceolata*. Yellowish cryst. Mp 167-169°.

N-Methoxy: N-Methoxy-1-vinyl- β -carboline

[69355-01-7]

C₁₄H₁₂N₂O 224.262

Alkaloid from *Picrasma excelsa* (Jamaican quassia wood) (Simarouba-ceae). Mp 185-190° dec.

7-Bromo: 7-Bromo-1-ethenyl-9H-pyrido[3,4-b]indole. 7-Bromo-1-vinyl- β -carboline. Plakortamine BC₁₃H₉BrN₂ 273.132

Alkaloid from the sponge *Plakortis nigra*. Cytotoxic. Yellow oil. λ_{\max} 214 (ϵ 32000) (no solvent reported).

Jordaan, A. *et al.*, *J. S. Afr. Chem. Inst.*, 1968, **21**, 22 (isol, uv, ir, pmr, ms, synth)

Wagner, H. *et al.*, *Tet. Lett.*, 1978, 2777 (deriv)

Blackman, A.J. *et al.*, *J. Nat. Prod.*, 1987, **50**, 494-496 (isol, Costaticella)

Bracher, F. *et al.*, *Annalen*, 1992, 1315 (synth, ir, pmr, cmr, ms)


Rocca, P. *et al.*, *Tetrahedron*, 1993, **49**, 3325 (synth, ir, pmr, cmr)

Sandler, J.S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1258-1261 (Plakortamine B)

1-Vinyl- β -carboline-3-carboxylic acid

V-140

1-Ethenyl-9H-pyrido[3,4-b]indole-3-carboxylic acid, 9CI. 3-Carboxy-1-vinyl- β -carboline
[85105-51-7]

C₁₄H₁₀N₂O₂ 238.245


Prod. by actinomycete *Nocardioopsis dassonvillei*. Antiinflammatory and antitumour agent. Cryst. (MeOH/MeCN).

Japan. Pat., 1982, 82 169 481; CA, **98**, 126057r

5-Vinyl-3-pyridinecarboxylic acid

V-141

5-Ethenyl-3-pyridinecarboxylic acid
[209915-44-6]

C₈H₇NO₂ 149.149

Me ester: 3-Methoxycarbonyl-5-vinylpyridine

[38940-67-9]

C₉H₉NO₂ 163.176

Constit. of *Nauclea diderrichii* (Rubiaceae). Minor constit. of *Jasminum grandiflorum*. Needles (hexane). Mp 50-60°.

Et ester: [68686-60-2]


C₁₀H₁₁NO₂ 177.202

Minor constit. of *Jasminum grandiflorum*.

- McLean, S. *et al.*, *Can. J. Chem.*, 1972, **50**, 1478-1485 (*Me ester, isol, synth*)
 Toyoda, T. *et al.*, *Agric. Biol. Chem.*, 1978, **42**, 1901-1905 (*esters, isol, ms*)
 Bracher, F. *et al.*, *Monatsh. Chem.*, 1995, **126**, 805-809 (*Me ester, synth, spectra*)

Violacein V-142

3-[1,2-Dihydro-5-(5-hydroxy-1H-indol-3-yl)-2-oxo-3H-pyrrol-3-ylidene]-1,3-dihydro-2H-indol-2-one, 9CI. 3-[2-(5-Hydroxyindol-3-yl)-5-oxo-2-pyrrolin-4-ylidene]-2-indolinone, 8CI. Anorosin† [548-54-9]



C₂₀H₁₃N₃O₃ 343.341
 Violet phenolic pigment produced by *Chromobacterium violaceum*. Also isol. from *Janthinobacterium lividum* and *Pseudoaerachniotus roseus*. Antibacterial and antiprotozoal agent. Shows potential as a photochemotherapeutic drug. Antioxidant. Purplish-black needles. Mp 350° dec. λ_{max} 255 (ε 14600); 278 (sh) (ε 11800); 295 (sh) (ε 6900); 370 (ε 5200); 568 (ε 18600) (MeOH) (Derep). λ_{max} 260 (ε 20000); 377 (ε 7760); 585 (ε 20000) (EtOH) (Derep).

- ▶ LD₅₀ (mus, ipr) 100 mg/kg. NM2203500
 3,3'-Dihydro: [83555-57-1]

C₂₀H₁₅N₃O₃ 345.357
 Prod. by *Chromobacterium violaceum*. Active against gram-positive and -negative bacteria.

Deoxy: Deoxyviolacein

[5839-61-2]
 Minor prod. from *Chromobacterium violaceum*. Dark blue powder. Mp 310° dec. λ_{max} 255 (ε 14600); 278 (sh) (ε 11800); 295 (sh) (ε 6900); 370 (ε 5200); 568 (ε 18600) (MeOH) (Derep). λ_{max} 260 (ε 20000); 377 (ε 7760); 585 (ε 20000) (EtOH) (Derep).


5-Hydroxy: Oxyviolacein

[132185-16-1]
 C₂₀H₁₃N₃O₄ 359.34
 Blue pigment from *Chromobacterium violaceum*. Sol. EtOH, Me₂CO, dioxan; poorly sol. H₂O. λ_{max} 255 (ε 15000); 278 (ε 14600); 295 (sh) (ε 9000); 370 (ε 7000); 560 (ε 18000) (MeOH) (Derep). λ_{max} 366; 370; 426 (MeOH) (Berdy).

- Beer, R.J.S. *et al.*, *J.C.S.*, 1949, 885-889; 1958, 755-760 (*isol*)
 Ballantine, J.A. *et al.*, *J.C.S.*, 1960, 2292-2299 (*synth*)
 de Moss, R.D. *et al.*, *Antibiotics (N.Y.)*, 1967, **2**, 77-81 (*rev*)
 Kimmel, K.E. *et al.*, *Can. J. Microbiol.*, 1969, **15**, 111-116 (*biosynth*)
 Duran, N. *et al.*, *Ann. Acad. Bras. Cinc.*, 1980, **52**, 297-302; 1983, **55**, 231-234 (*props, isol*)

- Hoshino, T. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 965-968; 2733-2741; 1990, **54**, 2339-2346 (*biosynth*)
Eur. Pat., 1989, ((Hoechst))341 649; *CA*, **112**, 233994y (*Anorosin*)
 Noble, J.A. *et al.*, *Diss. Abstr. Int.*, **B**, 1990, **50**, 2919 (*isol, struct*)
 Hoshino, T. *et al.*, *J.C.S. Perkin 1*, 1995, 1565-1567 (*biosynth*)
 Duran, N. *et al.*, *Crit. Rev. Microbiol.*, 2001, **27**, 201-222 (*pharmacol, rev*)
 Wille, G. *et al.*, *Synthesis*, 2001, 759-761 (*synth, ir, uv, pmr, cmr, ms, deoxy*)
 Konzen, M. *et al.*, *Bioorg. Med. Chem.*, 2006, **14**, 8307-8313 (*activity*)
 Sánchez, C. *et al.*, *ChemBioChem*, 2006, **7**, 1231-1240 (*biosynth*)
 Asamizu, S. *et al.*, *Tet. Lett.*, 2007, **48**, 2923-2926 (*biosynth*)

Violatinctamine V-143




C₂₀H₂₁N₃O₂S 367.471
 Alkaloid from *Cystodytes* cf. *violatinctus*. Orange oil.

Chill, L. *et al.*, *Tet. Lett.*, 2004, **45**, 7925-7928 (*isol*)

Viomycinidene V-144

3-Amino-2,4,6-triazabicyclo[3.2.1]oct-2-ene-7-carboxylic acid, 9CI. 2,4,6-Triaza-3-iminobicyclo[3.2.1]octane-7-carboxylic acid [24250-74-6]



C₆H₁₀N₄O₂ 170.171
 Acid hydrol. prod. of Viomycin. Mp 200-208° dec. (as hydrochloride). [α]_D³⁰ -78 (c, 1.78 in H₂O) (hydrochloride). pK_{a1} 2.8; pK_{a2} 5.87; pK_{a3} 13.4 (66% DMF aq.). pK_{a1} 1.3; pK_{a2} 5.5; pK_{a3} 12.6 (H₂O).


Ac: Mp 256-257°. [α]_D²⁸ +41.5 (c, 2.4 in H₂O).

- Dyer, J.R. *et al.*, *J.A.C.S.*, 1964, **86**, 5363
 Bowie, J.H. *et al.*, *Tet. Lett.*, 1964, 863
 Floyd, J.C. *et al.*, *Chem. Comm.*, 1968, 998 (*cryst struct*)

- Koyama, G. *et al.*, *J. Antibiot.*, 1969, **22**, 34 (*cryst struct*)
 Büchi, G. *et al.*, *J.O.C.*, 1971, **36**, 873

Virantmycin V-145

3-Chloro-2-(3,4-dimethyl-3-pentenyl)-1,2,3,4-tetrahydro-2-(methoxymethyl)-6-quinolinecarboxylic acid, 9CI [76417-04-4]



C₁₉H₂₆ClNO₃ 351.872
 Isol. from *Streptomyces nitrosporeus*. Active against viruses and weakly active against fungi. Needles. Sol. MeOH, DMSO, C₆H₆; fairly sol. hexane; poorly sol. H₂O. Mp 59°. [α]_D¹⁸ -0.05 (c, 1 in CHCl₃). λ_{max} 226 (ε 3500); 306 (ε 8100) (EtOH) (Derep). λ_{max} 223 (ε 9900); 283 (ε 14900); 300 (ε 14000) (MeOH) (Berdy). λ_{max} 214 (E1%/1cm 1950); 270 (E1%/1cm 306) (NaOH aq.) (Berdy). ▶ LD₅₀ (mus, ipr) 5 mg/kg. ZA4826100

Amide: Benzastatin C

[173429-77-1]
 C₁₉H₂₇ClN₂O₂ 350.887
 Prod. by *Streptomyces nitrosporeus*. Free radical scavenger, neuroprotective agent. Powder. Sol. MeOH, Me₂CO, EtOAc, DMSO; fairly sol. CHCl₃; poorly sol. H₂O. Mp 152°. [α]_D¹⁸ -23 (c, 0.1 in MeOH). λ_{max} 206 (ε 35480); 301 (ε 15488) (MeOH) (Berdy).


3-Dechloro, 3β-hydroxy: Benzastatin D

[173429-78-2]
 C₁₉H₂₇NO₄ 333.427
 Prod. by *Streptomyces nitrosporeus*. Free radical scavenger. Powder. Mp 220°. [α]_D -30 (c, 0.1 in MeOH).

- Omura, S. *et al.*, *J. Antibiot.*, 1980, **33**, 1395; 1996, **52**, 10631 (*isol*)
 Nakagawa, A. *et al.*, *J. Antibiot.*, 1981, **34**, 1408 (*isol*)
 Omura, S. *et al.*, *Tet. Lett.*, 1981, **22**, 2199 (*struct*)
 Pearce, C.M. *et al.*, *J.C.S. Perkin 1*, 1990, 409 (*pmr, abs config*)
 Hill, M.L. *et al.*, *Tetrahedron*, 1990, **46**, 4587 (*synth*)
 Kim, W.-G. *et al.*, *J. Antibiot.*, 1996, **49**, 20; 26 (*Benzastatin, isol, uv, ir, pmr, cmr, props*)
 Morimoto, Y. *et al.*, *Tetrahedron*, 1996, **52**, 10609 (*synth*)
 Morimoto, Y. *et al.*, *J. Het. Chem.*, 1998, **35**, 279-284 (*conformn*)
 Steinhagen, H. *et al.*, *Org. Lett.*, 1999, **1**, 823-824 (*synth*)
 Ori, M. *et al.*, *Tetrahedron*, 2005, **61**, 2075-2104 (*synth*)
 Keck, D. *et al.*, *Eur. J. Org. Chem.*, 2006, 4916-4923 (*synth*)

Virenamide A V-146

[176666-59-4]



C₃₁H₄₆N₄O₂S 538.796

Peptide antibiotic. Isol. from the ascidian *Diplosoma virens*. Cytotoxic agent. Oil. [α]_D -341 (c, 0.1 in CHCl₃). λ_{max} 203 (ε 7700); 237 (ε 4300) (EtOH).

N-Deprenyl: **Virenamide D**

[193264-93-6]

C₂₆H₃₈N₄O₂S 470.678

Isol. from the ascidian *Diplosoma virens*. Oil. [α]_D -65 (c, 0.07 in CHCl₃). λ_{max} 202 (ε 13700); 239 (ε 3050) (MeOH).


Carroll, A.R. *et al.*, *J.O.C.*, 1996, **61**, 4059-4061 (isol, uv, ir, pmr, cmr, ms)

Feng, Y. *et al.*, *Aust. J. Chem.*, 1997, **50**, 337-339 (Virenamide D)

Virenamide B

V-147

[176666-60-7]

R = CH(CH₃)₂C₃₀H₃₈N₄O₂S 518.722

Peptide antibiotic. Isol. from the ascidian *Diplosoma virens*. Cytotoxic agent. Oil. [α]_D -775 (c, 0.1 in CHCl₃). λ_{max} 209 (ε 13600); 240 (ε 4500) (EtOH).

Carroll, A.R. *et al.*, *J.O.C.*, 1996, **61**, 4059-4061 (isol, uv, ir, pmr, cmr, ms)

Moody, C.J. *et al.*, *J.O.C.*, 1999, **64**, 8715-8717 (synth)

Virenamide C

V-148

[176666-61-8]

As Virenamide B, V-147 with

R = -CH₂PhC₃₄H₃₈N₄O₂S 566.766


Peptide antibiotic. Isol. from the ascidian *Diplosoma virens*. Cytotoxic agent. Oil. [α]_D -634 (c, 0.05 in CHCl₃). λ_{max} 202 (ε 10900); 240 (ε 4800) (EtOH).

Carroll, A.R. *et al.*, *J.O.C.*, 1996, **61**, 4059-4061 (isol, uv, ir, pmr, cmr, ms)

Virenamide E

V-149

[193264-95-8]

C₂₁H₂₇N₃O₂S 385.529


Isol. from the ascidian *Diplosoma virens*. Oil. [α]_D -40 (c, 0.04 in CHCl₃). λ_{max} 216 (ε 12700); 222 (ε 10800); 274 (ε 14150) (MeOH).

Fang, Y. *et al.*, *Aust. J. Chem.*, 1997, **50**, 337-339 (isol, uv, ir, pmr, cmr, ms)

Virgidivarine

V-150

1'-(3-Butenyl)-[2,3'-bipiperidine]-5'-carboxylic acid, 9CI. 1-(3-Butenyl)-5-(2-piperidyl)-3-piperidinecarboxylic acid [81633-42-3]



Relative configuration

C₁₅H₂₆N₂O₂ 266.383

Biogenetically related to the Sparteine group. Isol. from *Virgilia divaricata* (Fabaceae). Tetragonal cryst. Mp 157°.

de Kok, A.J. *et al.*, *Acta Cryst. B*, 1982, **38**, 466 (cryst struct)

Van Eijk, J.L. *et al.*, *Planta Med.*, 1982, **44**, 221 (isol, ms, ir, pmr, cmr)

Virgildine

V-151

C₁₀H₁₉NO 169.266

Struct. unknown. Alkaloid from *Virgilia capensis* (Fabaceae). Bp_{0.01} 90°. [α]_D +12 (EtOH). Co-occurs with Virgiline, V-153. Not identical with Octahydro-2H-quinolizine-1-methanol, O-58 or Epilupinine, E-95.

Picrate: Mp 203°.


Methiodide: Mp 256-259°.

White, E.P. *et al.*, *N.Z. J. Sci. Technol., Sect. B*, 1946, **27**, 478-485

Virgildone

V-152

Decahydro-1,5-methano-6H-pyrido[1,2-a][1,5]diazocin-6-one, 9CI [135442-76-1]

C₁₁H₁₈N₂O 194.276

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

N-(3-Butenyl): **Virgiboidine**

[82464-31-1]

C₁₅H₂₄N₂O 248.367

Alkaloid from leaves of *Virgilia oroboides* and *Virgilia divaricata* (Fabaceae).

Didehydro: **Dehydrovirgildone**

[135442-77-2]

C₁₁H₁₆N₂O 192.26

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae). Posn. of double bond not determined.

Didehydro, N-(3-Butenyl): **Dehydrovirgiboidine**

[135556-70-6]

C₁₅H₂₂N₂O 246.352

Alkaloid from *Virgilia divaricata* and *Virgilia oroboides* (Fabaceae).

Position of double bond not determined.


Van Eijk, J.L. *et al.*, *Planta Med.*, 1982, **44**, 224 (Virgiboidine)

Veen, G. *et al.*, *Phytochemistry*, 1991, **30**, 1891 (Dehydrovirgiboidine, Virgildone)

Virgiline

V-153

Dodecahydro-9-hydroxy-7,14-methano-2H,6H-dipyrido[1,2-a:1',2'-e]diazocin-6-one, 9CI. 13-Hydroxyaphylline. 13-Hydroxy-10-oxosparteine [2636-61-5]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from seeds of *Virgilia oroboides*, tops of *Virgilia capensis* and from *Lupinus verbasciformis* and *Lupinus hartwegii* (Fabaceae). Prisms (Me₂CO/MeOH). Mp 249-250°. [α]_D²⁰ -29.

Methiodide:

Prisms (Me₂CO/MeOH). Mp 173-174° (170-171°).

Ac:

Prisms (Et₂O/Me₂CO). Mp 175-176°.

Tigloyl: **13α-Tigloyloxyaphylline**. O-Tigloylvirgiline

C₂₀H₃₀N₂O₃ 346.469

Minor alkaloid from the seeds of *Lupinus hartwegii*. Oil. [α]_D²⁵ -17.3 (c, 0.1 in CHCl₃).

2-Pyrrolocarboxylate: **O-(2-Pyrrolylcarbonyl)virgiline**. Virgiline 2-pyrrolocarboxylate, 8CI

[18526-91-5]

C₂₀H₂₇N₃O₃ 357.452

Alkaloid from branches of *Virgilia oroboides*, leaves of *Readea membranacea* and *Calpurnia aurea* (Fabaceae, Rubiaceae). Plates (EtOH). Mp 271° (265°). [α]_D²⁰ -23 (c, 2.4 in CHCl₃).

2-Pyrrolocarboxylate; hydrochloride:

[18526-92-6]

Needles (EtOH/Me₂CO). Mp 237-238° dec Mp 293°.

2,3-Didehydro, 2-pyrrolocarboxylate: **2,3-Dehydro-O-(2-pyrrolylcarbonyl)virgiline**

[18361-82-5]

C₂₀H₂₅N₃O₃ 355.436

Alkaloid from leaves of *Readea membranacea* (Rubiaceae). Shows strong molluscicidal activity. Cryst. (Et₂O). Mp 145° Mp 186-193°. [α]_D²⁰ -161 (c, 0.14 in CHCl₃).

13-Epimer: **Lindenianine**

[52539-65-8]

C₁₅H₂₄N₂O₂ 264.367

Alkaloid from *Lupinus lindenianus*

and *Lupinus verbasciformis* (Fabaceae). Cryst. (CHCl₃/Et₂O). Mp 212–213°.

13-*Epimer, picrate*: [52539-66-9]

Cryst. (Me₂CO/Et₂O). Mp 160–162°.

13-*Epimer, Ac*: [52539-67-0]

Viscous oil.

White, E.P. *et al.*, *N.Z. J. Sci. Technol., Sect. B*, 1946, **27**, 478 (*isol*)

Gerrans, G.C. *et al.*, *J.C.S.*, 1964, 2202 (*isol, struct*)

White, E.P. *et al.*, *J.C.S.*, 1964, 5243 (*isol, deriv*)

Manchanda, A.H. *et al.*, *J.C.S.(C)*, 1968, 615 (*isol, struct, uv, derivs*)

Nakano, T. *et al.*, *J.O.C.*, 1974, **39**, 3584 (*Lindenianine, isol, struct, cd*)

Anderson, J.N. *et al.*, *J.O.C.*, 1976, **41**, 3441 (*isol*)


Nakano, T. *et al.*, *Planta Med.*, 1976, **29**, 241 (*Virgiline, Lindenianine*)

Kubo, I. *et al.*, *Agric. Biol. Chem.*, 1984, **48**, 2839 (*isol, uv, pmr, derivs*)

Mohamed, M.H. *et al.*, *Pharmazie*, 1999, **54**, 778–780 (*tigloyl*)

Viridic acid V-154

N-[2-(Dimethylamino)benzoyl]glycyl-N-(2-carboxyphenyl)-N²-methylvalinamide, 9CI
[106283-19-6]




C₂₄H₃₀N₄O₅ 454.525

Isol. from *Penicillium viridicatum*.

Mycotoxin. Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. λ_{max} 260 (ε 15690); 301 (ε 5440) (MeOH) (Berdy). λ_{max} 260 (ε 15590); 295 (ε 4563) (MeOH/HCl) (Berdy). λ_{max} 262 (ε 16250); 305 (ε 5950) (MeOH/NaOH) (Berdy).

Holzappel, C.W. *et al.*, *S. Afr. J. Chem.*, 1986, **39**, 75 (*isol, struct, synth*)

Viriditin V-155



C₁₈H₂₉NO₃ 307.432

Prod. by *Aspergillus viridi-nutans*. Yellowish oil. [α]_D -65 (c, 1.2 in CHCl₃). λ_{max} 303 (log ε 4.53) (MeOH).

8-*Me ether, O-Methylviriditin*


C₁₉H₃₁NO₃ 321.459

Prod. by *Aspergillus viridi-nutans*. Yellowish oil. [α]_D -67 (c, 0.4 in CHCl₃). λ_{max} 302 (log ε 4.62) (MeOH).

Omolo, J.O. *et al.*, *J. Nat. Prod.*, 2000, **63**, 975–977

Viridomycin A V-156

[52970-22-6]



C₂₁H₁₂FeN₃O₉ 506.188

Fe complex of 4-Hydroxy-3-nitrosobenzaldehyde. Prod. by *Streptomyces viridans*. Moderate antibacterial agent. Mp 300°.

Monoalcohol, Viridomycin F

[221394-79-2]

C₂₁H₁₄FeN₃O₉ 508.203

Prod. by *Streptomyces* sp. K96-0188. Shows moderate antibacterial, insecticidal and nematocidal activity. Green powder. [α]_D²⁵ +6 (c, 0.1 in MeOH). Mp >300°. Has one -CHO group reduced to -CH₂OH. λ_{max} 207 (ε 21800); 226 (sh) (ε 13700); 288 (ε 7620); 435 (ε 2290); 680 (ε 2790) (MeOH).

Blinova, I.N. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 490; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 506 (*isol*)

Yang, C.C. *et al.*, *Antimicrob. Agents Chemother.*, 1981, **20**, 558 (*props*)

Kurobane, I. *et al.*, *J. Antibiot.*, 1987, **40**, 1131 (*isol, struct, bibl*)

Omura, S. *et al.*, *J. Antibiot.*, 1999, **52**, 61–64 (*Viridomycin F*)

Viridomycin B V-157

[57425-73-7]

Struct. unknown. Co-occurs with 4-Hydroxy-3-nitrosobenzaldehyde. Prob. an aromatic nitroso-Fe complex (uv v. similar to Viridomycin A). Prod. by a number of *Actinomyces* spp., isol. from cultures of *Actinomyces viridis* strain 1876. Amorph. yellow-green powder. Sol. Py, AcOH, DMF; fairly sol. MeOH, H₂O, butanol, EtOH, Me₂CO; poorly sol. EtOAc, hexane. Prob. an aromatic nitroso-Fe complex (uv v. similar to Viridomycin A). λ_{max} 222; 294; 435; 680 (EtOH) (Berdy). λ_{max} 294; 430; 680 (EtOH/HCl) (Berdy). λ_{max} 296; 430; 680 (EtOH/NaOH) (Berdy).

Blinova, I.N. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 490; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 506

Viridomycin C V-158


[57425-74-8]

Struct. unknown. Co-occurs with 4-Hydroxy-3-nitrosobenzaldehyde. Prob. an aromatic nitroso Fe complex (uv v. similar to Viridomycin A). Prod. by a number of *Actinomyces* spp., isol. from cultures of *Actinomyces viridis* strain

1876. Amorph. yellow-green powder. Sol. AcOH, DMF, Py; fairly sol. MeOH, H₂O, Me₂CO; poorly sol. EtOAc, hexane. λ_{max} 228; 294; 435; 680 (EtOH). λ_{max} 294; 435; 680 (EtOH-HCl) (Berdy). λ_{max} 294; 438; 680 (EtOH-NaOH) (Berdy).

Blinova, I.N. *et al.*, *Khim. Prir. Soedin.*, 1975, **11**, 490; *Chem. Nat. Compd. (Engl. Transl.)*, 1975, **11**, 506

Viridomycin E V-159




C₁₄H₈FeN₂O₇ 372.073

Fe complex of 4-Hydroxy-3-nitrosobenzaldehyde and 4-Hydroxy-3-nitrosobenzoic acid. Isol. from *Streptomyces griseus*. Dark-green cryst. Mp 300°. Related to Viridomycin A and Actinoviridin A. λ_{max} 268 (ε 22800); 295 (ε 21300); 434 (ε 6500); 685 (ε 6400) (MeOH) (Berdy).

Kurobane, I. *et al.*, *J. Antibiot.*, 1987, **40**, 1131 (*isol, struct*)

Virotoxin V-160



Viroisin, R¹ = CH₂OH, R² = CH(CH₃)₂
Viroidin, R¹ = CH₃, R² = CH(CH₃)₂
[Ala¹]Viroidin, R¹ = R² = CH₃

A series from *Amanita virosa*.

Viroisin [74113-57-8]

C₃₈H₅₆N₈O₁₆S 912.97

Major toxin from *Amanita virosa*. [α]_D²⁰ -22.4. λ_{max} 291 (ε 15000) (0.01N NaOH) (Derep). λ_{max} 278 (ε 15500); 293 (sh) (ε 9900); 305 (sh) (ε 6300) (H₂O/pH 7) (Derep).

▶ LD₅₀ (mus, ipr) 2.5mg/kg. ZA5300000

S-Deoxy, Desoxoviroisin. 2-[2-(Methylsulfinyl)-L-tryptophan]viroisin, 9CI [74125-17-0]

C₃₈H₅₆N₈O₁₅S 896.971

Toxin from *Amanita virosa*.

▶ ZA5360000

Viroidin

3-(4,5-Dihydroxy-L-leucine)viroisin, 9CI, VD

[53568-33-5]

C₃₈H₅₆N₈O₁₅S 896.971

Toxin from *Amanita virosa*. [α]_D²⁰ -13.6. λ_{max} 291 (ε 15000) (0.01N NaOH)

(Derep). λ_{\max} 278 (ϵ 15500); 293 (sh) (ϵ 9900); 305 (sh) (ϵ 6300) ($\text{H}_2\text{O}/\text{pH}$ 7) (Derep).

▶ ZA5320000

S-Deoxy: **Desoxoviroidin**. 2-[2-(Methylsulfinyl)-L-tryptophan]-3-(4,5-dihydroxy-L-leucine) viroisin, 9CI [74125-14-7]
 $\text{C}_{38}\text{H}_{56}\text{N}_8\text{O}_{14}\text{S}$ 880.972
 Toxin from *Amanita virosa*.

▶ ZA5380000

[Ala¹]Viroidin

3-(4,5-Dihydroxy-L-leucine)-4-L-alanine-viroisin [74125-15-8]
 $\text{C}_{36}\text{H}_{52}\text{N}_8\text{O}_{15}\text{S}$ 868.917
 Toxin from *Amanita virosa*.


▶ ZA5340000

S-Deoxy: [Ala¹]Desoxoviroidin. 2-[2-(Methylsulfinyl)-L-tryptophan]-3-(4,5-dihydroxy-L-leucine)-4-L-alanine-viroisin, 9CI [74125-16-9]
 $\text{C}_{36}\text{H}_{52}\text{N}_8\text{O}_{14}\text{S}$ 852.918
 Toxin from *Amanita virosa*.

Faulstich, H. et al., *Biochemistry*, 1980, **19**, 3334-3343 (struct, pmr, uv, cd)

Viscosaline

V-161



$\text{C}_{39}\text{H}_{65}\text{N}_3\text{O}_2$ 607.961

Alkaloid from the sponge *Haliclona viscosa*. CAS no. not found CA 141. λ_{\max} 204 ; 264 (MeOH).


Volk, C.A. et al., *Org. Biomol. Chem.*, 2004, **2**, 1827-1830 (isol, pmr, cmr)

Shorey, B.J. et al., *Tetrahedron*, 2007, **63**, 5587-5592 (synth)

Viscosamine

V-162

[606136-89-4]



$\text{C}_{54}\text{H}_{90}\text{N}_3^{\oplus}$ 781.325

Related to Cyclostelletamines, C-910. Alkaloid from the sponge *Haliclona viscosa*. Solid (as trifluoroacetate). λ_{\max} 203 ; 267 (MeOH) (TFA salt).

Volk, C. et al., *Org. Lett.*, 2003, **5**, 3567-3569


(isol, pmr, cmr, ms)

Timm, C. et al., *Synthesis*, 2006, 2580-2584 (synth, ir, uv, pmr, cmr)

Vitamycin A

V-163

Prodigiosin 25b [22373-48-4]



$\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}$ 391.555

Pyrrole antibiotic. Isol. from *Actinomyces aureovorticillatus*. Active against gram-positive bacteria. Orange-red cryst. powder (C_6H_6). Sol. MeOH, Et₂O; fairly sol. hexane; poorly sol. H₂O. Mp 192-194°. λ_{\max} 295 ; 322 ; 465 ; 495 ; 532 (hexane) (Berdy).

Korenyako, A.I. et al., *CA*, 1962, **57**, 17202 (biosynth)

Khokhlova, Y.M. et al., *Biokhimiya (Moscow)*, 1964, **29**, 841 (pmr, ms, struct)


Khokhlova, Y.M. et al., *Khim. Prir. Soedin.*, 1968, **4**, 307; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 260 (pmr, ms, struct)

Makukhina, A.M. et al., *CA*, 1976, **85**, 44853 (manuf)

Vitidoamine A

V-164

[819861-41-1]



$\text{C}_{20}\text{H}_{17}\text{NO}_5$ 351.358

Alkaloid from the seeds of *Vitex negundo*. Antioxidant. Powder.

Ono, M. et al., *J. Nat. Prod.*, 2004, **67**, 2073-2075 (isol, pmr, cmr)

Vitricine

V-165

[1362-23-8]

$\text{C}_{17}\text{H}_{15}\text{NO}_3$ 281.31

Struct. unknown. Poss. an aporphine. Alkaloid from *Vitex trifoliata* (Verbenaceae). Conts. 1 OH and one -OCH₂O-group but no OMe or NMe.

Perchlorate: Mp 178° dec.

Picrate: Mp 228°.


Döpke, W. et al., *Naturwissenschaften*, 1962, **49**, 375 (isol)

Shamma, M. et al., *Chem. Rev.*, 1964, **64**, 59-79 (rev)

Vittacarboline

V-166

9-[(5-Methylfuranyl)methyl]- β -carboline



$\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}$ 262.31


Alkaloid from the flowers of *Hippeastrum vittatum*. Yellow needles. Mp 157-159°. λ_{\max} 240 (log ϵ 4.2); 256 (log ϵ 4.7); 268 (log ϵ 4.26); 291 (log ϵ 3.76); 305 (log ϵ 4.05); 350 (log ϵ 3.45); 370 (log ϵ 3.88); 385 (log ϵ 3.25) (MeOH).

Youssef, D.T.A. et al., *J. Nat. Prod.*, 2001, **64**, 839-841

Voacamidine

V-167

[6897-04-7]



$\text{C}_{43}\text{H}_{52}\text{N}_4\text{O}_5$ 704.908

Alkaloid from *Voacanga africana*, *Voacanga schweinfurthii*, *Peschiera echinata* and *Tabernaemontana accedens* (Apocynaceae). Shows v. weak activity against P-388 lymphocytic leukaemia. Needles (C_6H_6). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 128-130° dec. [α]_D²⁴ -174.5 (CHCl₃).

Hydrochloride (1:2):

Needles (Me₂CO/MeOH). Mp 265-267° dec. [α]_D^{16.5} -166.5 (MeOH).

Renner, U. et al., *Experientia*, 1957, **13**, 468-469 (uv, ir)

Renner, U. et al., *Tet. Lett.*, 1964, 283-287 (struct)

Achenbach, H. et al., *Chem. Ber.*, 1976, **109**, 3527-3536 (isol, pmr)

Richard, B. et al., *J. Nat. Prod.*, 1983, **46**, 283-284 (isol)


Braga, R.M. et al., *Phytochemistry*, 1984, **23**, 175-178 (cmr)

Voacamine

V-168

Voacangimine

[3371-85-5]



C₄₃H₅₂N₄O₅ 704.908

Alkaloid from *Voacanga africana*, other *Voacanga* spp., *Tabernaemontana accedens*, *Tabernaemontana dichotoma*, *Tabernaemontana johnstonii*, *Tabernaemontana pachysiphon*, *Peschiera fuchsiaeifolia*, *Peschiera laeta*, *Ervatamia orientalis* and several other spp. in the Apocynaceae. Antineoplastic agent, shows strong activity vs. P-388 lymphocytic leukaemia. Also strongly active against gram-positive bacteria and mod. active against gram-negative organisms. Shows antiparasitic activity. Antimalarial agent. Sol. CHCl₃, Me₂CO; fairly sol. MeOH, EtOH. Mp 224°. [α]_D -52 (c, 1 in CHCl₃). Log P 7.36 (uncertain value) (calc). λ_{max} 226; 288; 295 (MeOH) (Berdy). λ_{max} 222; 225 (ε 30900); 285; 286 (ε 8700); 293; 294 (ε 8300) (EtOH) (Berdy).

► LD₅₀ (mus, ivn) 21.5 mg/kg (as sulfate). NH6737230

N^b-Oxide: **Voacamine N^b-oxide**

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from *Voacanga africana*, *Voacanga bracteata* and *Tabernaemontana accedens* (Apocynaceae). Mp 216–219° dec. [α]_D²⁰ -55 (c, 0.24 in CHCl₃).

N-De-Me: **N-Demethylvoacamine. 13'-Perivoacangine**

[61589-94-4]

C₄₂H₅₀N₄O₅ 690.881

Alkaloid from the root bark of *Tabernaemontana accedens* (Apocynaceae). Antineoplastic agent. Shows stronger activity than Voacamine vs. P-388 lymphocytic leukaemia. Cryst. (MeOH). Mp 227° dec. [α]_D²⁰ -67 (c, 0.12 in CHCl₃). Log P 6.51 (uncertain value) (calc).

19,20α-Dihydro: **Ervadivaricatine B**

[204581-20-4]

C₄₃H₅₄N₄O₅ 706.923

Alkaloid from *Ervatamia divaricata*.

19,20β-Dihydro: **Ervadivaricatine A**

[11047-69-1]

C₄₃H₅₄N₄O₅ 706.923

Alkaloid from *Ervatamia divaricata*.

3'-Hydroxy: **3-Hydroxyvoacamine**

C₄₃H₅₂N₄O₆ 720.907

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Active against gram-positive and gram-negative bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. Approx. 1:1 mixt. of 3-epimers. λ_{max} 224; 286; 294 (MeOH) (Berdy).

19'R-Hydroxy: **Voacorine**

[5130-80-3]

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from the trunk bark of *Tabernaemontana brachyantha*, and the root and stem bark of *Voacanga africana*, also from *Voacanga schweinfurthii* (Apocynaceae). Antineoplastic agent. Prisms (MeOH or MeOH/Et₂O). Mp 240° Mp 273° Mp 275–280°. Log P 5.15 (uncertain value) (calc).

► NH6737000

19'S-Hydroxy: **Epivoacorine**

[4835-65-8]

C₄₃H₅₂N₄O₆ 720.907

Alkaloid from *Tabernaemontana arborea*, *Tabernaemontana brachyantha*, *Voacanga bracteata* and *Voacanga africana* (Apocynaceae). Shows cytotoxic activity vs. mouse P-388 carcinoma. Cryst. (MeOH). Mp 265° (260° dec.). [α]_D -80 (c, 0.8 in CHCl₃). pK_a 6.12 (80% MCS aq.). Log P 5.15 (uncertain value) (calc). λ_{max} 226 (ε 42600); 288 (ε 14800); 295 (ε 13200) (EtOH) (Berdy).

Demethoxy, 3'-oxo: **Ervahaimine A**

[126398-88-7]

C₄₂H₄₈N₄O₅ 688.865

Minor alkaloid from the roots of *Ervatamia hainanensis* (Apocynaceae).

Janot, M.-M. et al., C. R. Hebd. Seances Acad. Sci., 1957, **244**, 1955 (*Voacorine*, uv, ir, struct)

Budzikiewicz, H. et al., Bull. Soc. Chim. Fr., 1963, 1899 (*Voacorine*, pmr, ms, struct)

Büchi, G. et al., J.A.C.S., 1964, **86**, 4631 (*Voacorine*, Voacamine, uv, ms, pmr, ir, struct, synth)

Poisson, J. et al., Bull. Soc. Chim. Fr., 1965, 3549 (*Voacorine*, uv, ir, ms, pmr, struct)

Thomas, D.W. et al., J.A.C.S., 1965, **87**, 5447 (*Voacamine*, ms)

Patel, M.B. et al., Phytochemistry, 1973, **12**, 451 (*Epivoacorine*)

Achenbach, H. et al., Chem. Ber., 1976, **109**, 3527 (*Voacamine N-oxide*, N-Demethylvoacamine)

Suffness, M. et al., Alkaloids (Academic Press), 1985, **25**, 216 (*Voacamine*, antileukaemic props)

Van Beek, T.A. et al., J. Ethnopharmacol., 1985, **14**, 315-318 (*3-Hydroxyvoacamine*, activity)

Van Beek, T.A. et al., J. Nat. Prod., 1985, **48**, 400 (*3-Hydroxyvoacamine*)

Feng, X.Z. et al., J. Nat. Prod., 1989, **52**, 928 (*Ervahaimine A*)

Huang, L. et al., Zhongcaoyao, 1997, **28**, 451-454; CA, **128**, 228465c (*Ervadivaricatines*)

Medeiros, W.L.B. et al., Magn. Reson. Chem., 1999, **37**, 676-681 (*Voacamine*, pmr, cmr)

Federici, E. et al., Planta Med., 2000, **66**, 93 (*Voacamine*, activity)

Voacaminine

V-169

[1362-29-4]


Struct. unknown. Alkaloid from *Voacanga africana* (Apocynaceae). Cryst. (MeOH). Mp 242°. [α]_D -45 (c, 0.44 in CHCl₃).

U.S. Pat., 1958, 2 823 204; CA, **52**, 11976a
La Barre, J. et al., Actual. Pharmacol., 1961, **14**, 109-124

Voacangine

V-170

[510-22-5]



Absolute Configuration

C₂₂H₂₈N₂O₃ 368.475

Alkaloid from *Voacanga africana*, several other *Voacanga* spp., *Tabernaemontana johnstonii*, *Tabernaemontana divaricata*, *Tabernanthe pubescens*, *Ervatamia heyneana*, *Pagiantha cerifera* (preferred

genus name *Tabernaemontana*), several *Stemmadenia* spp. and other spp. in the Apocynaceae. Probably the most widely distributed of the Iboga alkaloids. Shows mod. cytotoxic activity, also some CNS, brachycardial and hypotensive action. Mp 137–138°. [α]_D -42 (c, 1 in CHCl₃). λ_{max} 224 (ε 32580); 288 (ε 10220) (EtOH).

Picrate: Mp 148°.

O-De-Me: **10-Hydroxycoronaridine**

[76129-67-4]

C₂₁H₂₆N₂O₃ 354.448

Alkaloid from the leaves of *Tabernanthe pubescens* and from the wood and stem bark of *Ervatamia heyneana* (Apocynaceae). Shows cytotoxic activity vs. mouse P-388 carcinoma cells. Shows estrogen-like activity. Amorph. [α]_D -33 (c, 0.5 in CHCl₃).

3-Oxo: **3-Oxovoacangine**

[3306-59-0]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from whole plants of *Ervatamia coronaria* var. *plena*. Also obt. by oxidn. of Voacangine. Prisms (C₆H₆/EtOAc). Mp 252°. [α]_D -52 (c, 0.05 in CHCl₃).

19-Oxo: **Voacryptine. 19-Oxovoacangine.**

20-Oxovoacangine

[4995-51-1]

C₂₂H₂₆N₂O₄ 382.458

Alkaloid from the bark of *Voacanga africana* and from *Tabernaemontana* sp. (Apocynaceae). Needles (Et₂O). Mp 175–176°. [α]_D²² +24.7 (c, 1 in CHCl₃). λ_{max} 224 (log ε 4.46); 285 (log ε 3.98) (no solvent reported).

3R-Hydroxy: **10-Methoxyeglandine**

[80981-96-0]

[95840-38-3]

C₂₂H₂₈N₂O₄ 384.474

Alkaloid from the stem bark and seeds of *Peschiera echinata* and seeds of *Voacanga schweinfurthii* var. *puberula*. Amorph. Originally descr. as 3,6-Oxidocoronaridine. Struct. revised here in accordance with the new struct. of Eglantine, but this is not confirmed. No further work since 1982.

3R-Hydroxy, N-oxide: **10-Methoxyeglandine N-oxide**

[76129-66-3]

C₂₂H₂₈N₂O₅ 400.474

Alkaloid from the wood and stem bark of *Ervatamia heyneana* (Apocynaceae). Cytotoxic agent. [α]_D²⁶ +86.6 (c, 0.76 in CHCl₃). Struct. and MF revised from that shown in the lit., in accordance with the new struct. of Eglantine. λ_{max} 224 (ε 28900); 282 (ε 9600) (MeOH) (Berdy).

3R-Methoxy: **3R-Methoxyvoacangine**

[162229-95-0]

C₂₃H₃₀N₂O₄ 398.501

Minor alkaloid from bark of *Tabernaemontana markgrafiana* (Apocynaceae). Amorph. solid. [α]_D -39 (c, 0.4 in CHCl₃). Erroneous struct. diag. in paper.

3S-Hydroxy: 3S-Hydroxyvoacangine

[56867-76-6]

C₂₂H₂₈N₂O₄ 384.474Alkaloid from tissue cultures of *Tabernaemontana pandacaqui*. λ_{\max} 218 ; 279 ; 310 (sh) (no solvent reported).**19R-Hydroxy: 19-Epivoacristine. 19-Epivoacangine**

[6883-77-8]

C₂₂H₂₈N₂O₄ 384.474Alkaloid from *Voacanga bracteata*, *Trachelospermum jasminoides*, *Tabernaemontana arborea* and *Tabernaemontana dichotoma*. Mp 115°. $[\alpha]_{\text{D}}^{25}$ -47 (CHCl₃). λ_{\max} 224 ; 282 ; 298 (MeOH).**19R-Hydroxy, 3-oxo: 19-Epi-3-oxovoacristine. 3-Oxo-19-epivoacristine**

[524729-86-0]

C₂₂H₂₆N₂O₅ 398.458Alkaloid from *Tabernaemontana cal-carea*. Pale yellow oil. $[\alpha]_{\text{D}}^{20}$ -10 (c, 0.42 in CHCl₃). λ_{\max} 224 (log ϵ 4.14); 280 (log ϵ 3.82); 294 (log ϵ 3.45) (MeOH).**19S-Hydroxy: Voacristine. 19-Hydroxyvoacangine. Voacangine**

[545-84-6]

C₂₂H₂₈N₂O₄ 384.474Alkaloid from *Voacanga africana*, *Ervatamia orientalis*, several *Tabernaemontana* and *Pandaca* spp. (Apocynaceae). Shows weak cytotoxicity vs. P-388 leukaemia cells. Also shows CNS, brachycardial and hypotensive activity. Cryst. Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 112-114° Mp 163-165° (dimorph.). λ_{\max} 222 (log ϵ 4.45); 282 (log ϵ 3.98); 302 (log ϵ 3.9); 314 (sh) (log ϵ 3.64) (MeOH).

▶ NH6737100

19S-Hydroxy, O-de-Me: 10-Hydroxyheyanine

[77795-14-3]

C₂₁H₂₆N₂O₄ 370.447Alkaloid from the leaves of *Tabernaemontana pubescens*, *Peschiera echinata*, *Anartia* cf. *meyeri* (preferred genus name *Tabernaemontana*), and the roots of *Ervatamia hainanensis* (Apocynaceae). Amorph. $[\alpha]_{\text{D}}^{25}$ -42 (c, 0.4 in CHCl₃).**19S-Hydroxy, 3-oxo: 3-Oxovoacristine.****19-Oxovoacristine**

[84453-51-0]

C₂₂H₂₆N₂O₅ 398.458Alkaloid from the stem bark of *Tabernaemontana citrifolia* (Apocynaceae). Cryst. (MeOH). Mp 251-252°. $[\alpha]_{\text{D}}^{22}$ -35.5 (c, 0.1 in CHCl₃).**19S-Acetoxy: Mp 191-193°. $[\alpha]_{\text{D}}^{23}$ -27 (CHCl₃).****3-Ethoxy: 3-Ethoxyvoacangine**


[746643-03-8 (R-form), 746643-04-9 (S-form)]

C₂₄H₃₂N₂O₄ 412.528Alkaloid from the stem bark of *Tabernaemontana divaricata*. Pale yellow oil. Obt. as a mixt. of 3R- and 3S-epimers. λ_{\max} 224 (log ϵ 4.4); 287 (log ϵ 4.09); 299 (log ϵ 4.06)

(EtOH).

3 ξ -Cyano: Voacangine 3-carbonitrile

[84306-86-5]

C₂₃H₂₇N₃O₃ 393.485Alkaloid from the seeds of *Pterotaberna inconspicua* (preferred genus name *Tabernaemontana*). $[\alpha]_{\text{D}}^{25}$ -68 (c, 1.1 in CHCl₃). λ_{\max} 225 ; 285 ; 300 (MeOH).Janot, M.-M. et al., *C. R. Hebd. Seances Acad. Sci.*, 1955, **240**, 1800-1801 (*isol*)Stauffer, D. et al., *Helv. Chim. Acta*, 1958, **41**, 169-180 (*Voacristine, isol, uv, ir*)Dickel, D.F. et al., *J.A.C.S.*, 1958, **80**, 123-125 (*isol*)Bartlett, M.F. et al., *J.A.C.S.*, 1958, **80**, 126-136 (*struct*)Walls, F. et al., *Tetrahedron*, 1958, **2**, 173-182 (*uv*)Renner, U. et al., *Experientia*, 1959, **15**, 185-186; 456-457; 1961, **17**, 106 (*Voacristine, Voacryptine, struct*)Budzikiewicz, H. et al., *Bull. Soc. Chim. Fr.*, 1963, 1899-1905 (*ms*)Guisse, G.B. et al., *Aust. J. Chem.*, 1965, **18**, 1279-1286 (*synth, 3-Oxovoacangine*)Poisson, J. et al., *Bull. Soc. Chim. Fr.*, 1965, 3549-3352 (*19-Epivoacristine*)Govindachari, T.R. et al., *Tet. Lett.*, 1965, 3873-3878 (*Voacristine, pmr, ms*)Goldblatt, A. et al., *Phytochemistry*, 1970, **9**, 1293-1298 (*abs config*)Bláha, K. et al., *Coll. Czech. Chem. Comm.*, 1974, **39**, 2258-2266 (*Voacristine, abs config*)Agwada, V.G. et al., *Helv. Chim. Acta*, 1975, **58**, 1001 (*synth, 3-Hydroxyvoacangine*)Harmouche, A. et al., *Ann. Pharm. Fr.*, 1976, **34**, 31-35 (*pmr*)Wenkert, E. et al., *Helv. Chim. Acta*, 1976, **59**, 2437-2442 (*cmr*)Kingston, D.G.I. et al., *J. Pharm. Sci.*, 1978, **67**, 271-272; 272-274 (*activity*)Gunasekera, S.P. et al., *Phytochemistry*, 1980, **19**, 1213-1218 (*10-Methoxyeglandine N-oxide, 10-Hydroxycoronaridine*)Mulamba, T. et al., *J. Nat. Prod.*, 1981, **44**, 184-189 (*10-Hydroxycoronaridine, 10-Hydroxyheyanine*)Ghorbel, N. et al., *J. Nat. Prod.*, 1981, **44**, 717-721 (*10-Methoxyeglandine*)Kutney, J.P. et al., *Helv. Chim. Acta*, 1982, **65**, 2242-2250 (*3-Oxovoacristine*)Morfaux, A.M. et al., *Phytochemistry*, 1982, **21**, 1767-1769 (*Voacangine 3-carbonitrile, 3,6-Oxidovoacangine*)Perera, P. et al., *Planta Med.*, 1983, **47**, 148-150 (*19-Epivoacristine*)Danieli, B. et al., *Alkaloids (Academic Press)*, 1986, **27**, 1-130 (*rev*)Atta-ur-Rahman, et al., *Planta Med.*, 1987, **53**, 57-59 (*19-Epivoacristine*)Sharma, P. et al., *J. Nat. Prod.*, 1988, **51**, 528-531 (*isol, 3-Oxovoacangine*)Soriano-Garcia, M. et al., *J. Crystallogr. Spectrosc. Res.*, 1989, **19**, 725-732 (*cryst struct*)Sierra, M.I. et al., *Planta Med.*, 1991, **57**, 543-547 (*isol, 3S-Hydroxyvoacangine*)Nielsen, H.B. et al., *Phytochemistry*, 1994, **37**, 1729-1735 (*3R-Methoxyvoacangine*)Masuda, K. et al., *Planta Med.*, 2000, **66**, 169-173 (*10-Hydroxycoronaridine, activity*)Chaturvedula, V.S.P. et al., *J. Nat. Prod.*, 2003, **66**, 528-531 (*19-Epi-3-oxovoacristine*)Kam, T.-S. et al., *Chem. Biodiversity*, 2004, **1**, 646-656 (*3-Ethoxyvoacangine*)Kam, T.-S. et al., *Heterocycles*, 2004, **63**, 845-850 (*3-Oxovoacangine*)Pereira, P.S. et al., *Quim. Nova*, 2008, **31**, 20-24 (*pmr, cmr*)**Voacarpine***Methyl 3,17-dihydroxysarpagan-16-carboxylate, 9CI*
[3650-53-1]

Absolute Configuration

C₂₁H₂₄N₂O₄ 368.432Alkaloid from *Voacanga chalitiana* and *Hazunta* spp. (Apocynaceae). Prisms (C₆H₆). Mp 227-228° dec. $[\alpha]_{\text{D}}^{20}$ +43.5 (c, 0.4 in MeOH). λ_{\max} 225 (ϵ 31350); 282 (ϵ 6260); 292 (sh) (ϵ 5150) (MeOH).O,N-Di-Ac: Mp 212-213°. $[\alpha]_{\text{D}}^{20}$ -22 (c, 0.7 in MeOH).N^d-Me: **Vincadiffine**. N^b-Methylvoacarpine. 3-Oxo-N^b-methyl-3,4-secoakummidine

[3238-64-0]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from *Vinca difformis* (Apocynaceae). Mp 230° (222°). $[\alpha]_{\text{D}}^{22}$ -121 (CHCl₃). $[\alpha]_{\text{D}}^{22}$ -70 (MeOH). Zwitterionic, tautomeric with the 3-oxo ring-cleaved amine form.N^d-Me, hydrochloride: Mp 202°.N^d-Me, stereoisomer 1 (?): **Voacafrine**

[7080-82-2]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from the tree bark of *Voacanga africana* (Apocynaceae). Plates (CH₂Cl₂/Et₂O). Mp 135-137° dec. Suggested struct. 16-Config, undetermined. Either Voacafrine or Voacafricine should be identical with Vincadiffine above, but there has been no further work on these alkaloids. λ_{\max} 240 (ϵ 17070); 315 (ϵ 20670) (no solvent reported).N^d-Me, stereoisomer 1 (?), hydrochloride:Plates (Me₂CO/MeOH). Mp 165-167° dec. $[\alpha]_{\text{D}}^{25}$ -107 (c, 1 in MeOH).N^d-Me, stereoisomer 2 (?): **Voacafricine**

[6817-40-9]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from the bark of *Voacanga africana* (Apocynaceae). Prisms (CH₂Cl₂/Et₂O). Mp 196-198°. Tentative struct. MF may be C₂₂H₂₄N₂O₄. See note under Voacafrine above. λ_{\max} 238 (ϵ 18570); 315 (ϵ 22500) (no solvent reported).N^d-Me, stereoisomer 2 (?), hydrochloride:

Plates. Mp 200-202° dec.

16-Epimer: 16-Epivoacarpine

[114027-38-2]

C₂₁H₂₄N₂O₄ 368.432Alkaloid from the roots of *Gelsemium elegans* (Loganiaceae). Prisms or plates + 1/8 H₂O (CH₂Cl₂). Mp 162-165° dec. $[\alpha]_{\text{D}}^{22}$ +42.3 (c, 0.20 in CHCl₃).**16-Epimer, N^l-Me: 3-Hydroxyvoachalotine**

[53632-75-0]

C₂₂H₂₆N₂O₄ 382.458Alkaloid from the root bark of *Voa-*

canga chalitiana (Apocynaceae).
Cryst. (EtOAc). Mp 247°. $[\alpha]_D^{22} +9$ (c, 1
in CHCl_3). λ_{max} 229 (log ϵ 4.62); 286
(log ϵ 3.88) (MeOH).

16-Epimer, N¹-Me, di-Ac:
Cryst. (EtOAc). Mp 232°.

16-Epimer, 19Z-isomer: (19Z)-16-Epi-
voacarpine

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ 368.432

Alkaloid from the radix of *Gelsemium
sempervirens*. Amorph. powder. λ_{max}
225 (log ϵ 4.49); 282 (log ϵ 3.79); 290
(log ϵ 3.71) (MeOH).

16-Epimer, N⁴-Me: *Gelsempervine A*

[865187-17-3]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from the radix of *Gelsemium
sempervirens*. Amorph. powder. Sol-
vent-dependent tautomeric equilib.
between cyclic zwitterionic and ring-
cleaved ketoamine forms. λ_{max} 220 (log
 ϵ 4.44); 282 (log ϵ 3.71); 290 (log ϵ 3.65)
(MeOH) (zwitterion). λ_{max} 224 (log ϵ
4.17); 311 (log ϵ 3.99) (MeCN) (ke-
toamine).

16-Epimer, N⁴-Me, 17-Ac: *Gelsempervine
B*

[865187-29-7]

$\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_5$ 424.496

Alkaloid from the radix of *Gelsemium
sempervirens*. Amorph. powder. λ_{max}
219 (log ϵ 4.46); 282 (log ϵ 3.74); 290
(log ϵ 3.72); 316 (log ϵ 3.65) (MeOH).
 λ_{max} 221 (log ϵ 4.07); 312 (log ϵ 3.92)
(MeCN).

16-Epimer, (19Z)-isomer, N⁴-Me: *Gel-
sempervine C*

[865187-18-4]

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from the radix of *Gelsemium
sempervirens*. Amorph. powder. λ_{max}
221 (log ϵ 4.54); 281 (log ϵ 3.8); 290
(log ϵ 3.7) (MeOH). λ_{max} 227 (log ϵ
4.27); 309 (log ϵ 4.02) (MeCN).

16-Epimer, (19Z)-isomer, N⁴-Me, 17-Ac:
Gelsempervine D

[865187-20-8]

$\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_5$ 424.496

Alkaloid from the radix of *Gelsemium
sempervirens*. Amorph. powder. λ_{max}
220 (log ϵ 4.48); 282 (log ϵ 3.77); 290
(log ϵ 3.71); 315 (log ϵ 3.36) (MeOH).
 λ_{max} 221 (log ϵ 4.21); 311 (log ϵ 4.01)
(MeCN).

Rao, K.V. *et al.*, *J.O.C.*, 1958, **23**, 1455-1456
(*Voacarpine, Voacarpine*)

Denayer-Tournay, M. *et al.*, *Bull. Soc. Chim.
Belg.*, 1965, **74**, 170-186 (*Voacarpine, isol, uv,
ir, pmr, ms, struct*)

Das, B.C. *et al.*, *Bull. Soc. Chim. Fr.*, 1965,
1903-1904 (*Vincadiffine*)

Weisbach, J.A. *et al.*, *Chem. Ind. (London)*,
1965, 623-633 (*Voacarpine, Voacarpine*)

Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*,
1966, **75**, 465-472 (*Voacarpine, Vincadiffine,
abs config*)

Gabetta, B. *et al.*, *Fitoterapia*, 1974, **45**, 32-36
(*3-Hydroxyvoachalotine*)

Bombardelli, E. *et al.*, *Phytochemistry*, 1974,
13, 2857-2859 (*3-Hydroxyvoachalotine*)


Achenbach, H. *et al.*, *Annalen*, 1982, 1223-
1227 (*Vincadiffine, ms*)

Ponglux, D. *et al.*, *Tetrahedron*, 1988, **44**, 5075-
5094 (*16-Epivoacarpine*)

Kogure, N. *et al.*, *Tet. Lett.*, 2005, **46**, 5857-
5861 (*Gelsempervines A-D, 19Z-16-
Epivoacarpine*)

Voachalotine oxindole, 9CI V-172

[26126-84-1]



$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458


Minor alkaloid from *Voacanga chalo-
tiana* (Apocynaceae). Mp 280-281°. $[\alpha]_D$ -
33 (CHCl_3).

Braekman, J.-C. *et al.*, *Bull. Soc. Chim. Belg.*,
1969, **78**, 523

Voacinol V-173

14,14'-Methylenebis-18-hydroxytabersone-
nine

[113141-69-8]




$\text{C}_{43}\text{H}_{48}\text{N}_4\text{O}_6$ 716.875

Alkaloid from the leaves of *Voacanga
grandifolia* (Apocynaceae). Yellow
amorph. solid. $[\alpha]_D^{25}$ -105.5 (CHCl_3).

Govindachari, T.R. *et al.*, *Chem. Comm.*, 1987,
1137 (*uv, ir, pmr, cmr, ms, struct*)

Voacoline V-174

[5539-91-3]



Absolute
Configuration

$\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_4$ 382.458

Alkaloid from *Voacanga chalitiana*
(Apocynaceae). Needles (Et₂O). Mp
147.5°. $[\alpha]_D^{20}$ -29.8 (c, 0.5 in CHCl_3). pK_a
5.7. λ_{max} 230 (ϵ 35000); 285 (ϵ 7300)
(MeOH).

3-Hydroxy, N-de-Me: *Voamonine*

[5296-84-4]

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_5$ 384.431

Minor alkaloid from *Voacanga cha-*

lotiana (Apocynaceae). Needles
(MeOH). Mp 258-250°. λ_{max} 225 (ϵ
41500); 282 (ϵ 7150); 292 (sh) (ϵ 5800)
(MeOH).


Braekman, J.C. *et al.*, *Bull. Soc. Chim. Belg.*,
1965, **74**, 170-186; 1969, **78**, 523-538
(*Voamonine*)

Lhoest, G. *et al.*, *Bull. Soc. Chim. Belg.*, 1965,
74, 534-550 (*Voacoline*)

Voafolidine

[32063-91-5]

V-175



$\text{C}_{42}\text{H}_{48}\text{N}_4\text{O}_5$ 688.865

Minor alkaloid from leaves of
Voacanga africana (Apocynaceae). Cryst.
(MeOH). Mp 200° dec. $[\alpha]_D$ -300
(CHCl_3).

2-Deoxy: *Voafoline, 9CI*

[31947-67-8]

$\text{C}_{42}\text{H}_{48}\text{N}_4\text{O}_4$ 672.866

Alkaloid from the leaves of *Voacanga
africana* (Apocynaceae). Cryst.
(MeOH). Mp 325° dec. $[\alpha]_D$ -310
(CHCl_3).

14'-Epimer: *Isovoafolidine*

[33055-38-8]

Semisynthetic, by redn. of Folicangine,
F-117. Mp 225-232° dec. $[\alpha]_D$ -642
(CHCl_3).

14'-Epimer, 2-deoxy: *Isovoafoline*

[31947-66-7]

$\text{C}_{42}\text{H}_{48}\text{N}_4\text{O}_4$ 672.866

Alkaloid from the leaves of *Voacanga
africana* (Apocynaceae). Cryst.
(MeOH). Mp 232° dec. $[\alpha]_D$ -567
(CHCl_3).

Kunesch, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1970,
4370 (*isol, uv, ir, pmr, ms, struct*)


Rolland, Y. *et al.*, *J.O.C.*, 1976, **41**, 3270-3275
(*cmr*)

Kunesch, N. *et al.*, *Helv. Chim. Acta*, 1977, **60**,
2854 (*cd, config*)

Voafrine A

[88761-30-2]

V-176





C₄₂H₄₆N₄O₄ 670.85Trace alkaloid from cell suspension cultures of *Voacanga africana* (Apocynaceae). Amorph. [α]_D²⁰ -8 (c, 0.62 in MeOH).**3ξ-Hydroxy: 3-Hydroxyvoafrine A**Obt. by dimerisation of Tabersonine using a crude enzyme prep. from *Catharanthus roseus*. Amorph. [α]_D²⁰ -21 (c, 0.4 in MeOH).**3'-Epimer: Voafrine B**

[88824-28-6]

C₄₂H₄₆N₄O₄ 670.85Trace alkaloid from cell suspension cultures of *Voacanga africana* (Apocynaceae). [α]_D²⁰ -265 (c, 0.83 in MeOH).**3'-Epimer, 3ξ-hydroxy: 3-Hydroxyvoafrine B**Obt. by dimerisation of Tabersonine using a crude enzyme prep. from *Catharanthus roseus*. Amorph. [α]_D²⁰ -68 (c, 0.45 in MeOH).Stöckigt, J. et al., *Helv. Chim. Acta*, 1983, **66**, 2525 (isol, uv, ir, pmr, cmr, ms, cd, struct)
Fahn, W. et al., *Phytochemistry*, 1990, **29**, 127**Voaharine****V-177**

[142741-23-9]

C₁₉H₂₂N₂O₃ 326.394Alkaloid from the leaves of *Tabernaemontana divaricata* (Apocynaceae). Prisms (EtOAc). Mp 176-178°.Kam, T.-S. et al., *Tet. Lett.*, 1992, **33**, 969-972 (isol, pmr, cmr, struct)**Voaluteine****V-178***Voacangine pseudoindoxyl. Rupicoline*† [3306-58-9]

Absolute Configuration

C₂₂H₂₈N₂O₄ 384.474Alkaloid from *Rejoua aurantiaca* (preferred genus name *Tabernaemontana*), *Tabernaemontana heyneana* and *Tabernaemontana rupicola* (Apocynaceae). Mp 208° (202-203°). [α]_D¹⁸ -115 (c, 1 in CHCl₃).*Hydrochloride*: Mp 265° dec. [α]_D²⁵ -228 (c, 1.2 in H₂O).**19S-Hydroxy: Voacristine pseudoindoxyl.***19-Hydroxyvoaluteine. Montanine*†

[18646-15-6]

C₂₂H₂₈N₂O₅ 400.474Alkaloid from *Tabernaemontana rupicola*, *Tabernaemontana apoda* and *Peschiera lundii* (Apocynaceae). Amorph. Mp 137-138°.**19S-Hydroxy; hydrochloride:**Cryst. (2-propanol). Mp 261°. [α]_D²⁵ -223.5 (c, 0.47 in EtOH).**11-Methoxy: Conopharyngine pseudoindoxyl**

[52579-71-2]

C₂₃H₃₀N₂O₅ 414.5Alkaloid from the leaves of *Tabernaemontana pachysiphon* var. *cumminsii* (Apocynaceae). Yellow oil. 7-Config. undetermined.**Demethoxy: Coronaridine pseudoindoxyl**


[16620-29-4]

C₂₁H₂₆N₂O₃ 354.448Alkaloid from *Peschiera affinis*, *Tabernaemontana quadrangularis* and *Tabernaemontana olivacea* (Apocynaceae). Yellow oil. Mp 278-279° dec. (as hydrochloride). [α]_D²⁰ -163 (c, 0.1 in CHCl₃).**Demethoxy, 15,20-didehydro: Rosamine**

[94061-94-6]

C₂₁H₂₄N₂O₃ 352.432Alkaloid from the leaves of *Catharanthus roseus* (Apocynaceae). Amorph. [α]_D²⁰ +100 (CHCl₃).**10-Demethoxy, 11-methoxy, 19R-hydroxy: Ervatine**C₂₂H₂₈N₂O₅ 400.474Alkaloid from the fruit of *Tabernaemontana heyneana*. Amorph. yellow powder. λ_{\max} 228 (log ϵ 3.06); 252 (log ϵ 1.7); 281 (log ϵ 1.3) (MeOH).Guise, G.B. et al., *Aust. J. Chem.*, 1965, **18**, 927; 1279 (isol, uv, ir, pmr, struct synth)Niemann, C. et al., *J.O.C.*, 1966, **31**, 2265 (isol, ir, uv, pmr, struct, deriv)Thomas, D.W. et al., *Tetrahedron*, 1968, **24**, 4223 (ms)Hwang, B. et al., *J.O.C.*, 1969, **34**, 412

(Voacristine pseudoindoxyl, synth)

Crooks, P.A. et al., *J. Pharm. Pharmacol.*, 1973, **25**, 820 (Coronaridine pseudoindoxyl, Conopharyngine pseudoindoxyl, isol, uv, ir, pmr, struct synth)Abreu Matos, F.J. et al., *Phytochemistry*, 1976, **15**, 551 (Coronaridine pseudoindoxyl, isol)Achenbach, H. et al., *Z. Naturforsch., B*, 1980, **35**, 885 (Coronaridine pseudoindoxyl, isol, uv, ir, pmr, ms)Atta-ur-Rahman, et al., *Z. Naturforsch., B*, 1984, **39**, 1292 (Rosamine)Srivastava, S. et al., *Planta Med.*, 2001, **67**, 577-579 (Ervatine)**Voaphylline****V-179***13-Ethyl-1a,4,5,10,11,12,13,13a-octahydro-2H-3,13-methanooxireno[9,10]azacycloundecino[5,4-b]indole, 9CI.**Conoflorine. Ervayunine*

(+) -form

C₁₉H₂₄N₂O 296.411

Ervayunine was the (-)-form.

(+) -form [15266-46-3]Alkaloid from *Voacanga africana*, *Conopharyngia longiflora*, several *Tabernaemontana* spp. and other spp. in theApocynaceae. Cryst. (MeOH). Mp 166-168°. [α]_D²⁴ +24.4 (c, 1.04 in CHCl₃). pK_a 5.27 (MCS).**N-Me: N-Methylvoaphylline. Hecubine**

[62874-52-6]

C₂₀H₂₆N₂O 310.438Alkaloid from the leaves and flowers of *Ervatamia coronaria* (Apocynaceae).**12-Methoxy: 12-Methoxyvoaphylline**


[53904-98-6]

C₂₀H₂₆N₂O₂ 326.438Alkaloid from the leaves of *Crioceras dipladeniiflorus* (Apocynaceae). Cryst. (MeOH). Mp 149-151°. [α]_D²⁰ +61 (c, 1 in CHCl₃).**16β-Hydroxy: Voafinine**

[181284-52-6]

C₁₉H₂₄N₂O₂ 312.411Trace alkaloid from leaves of *Tabernaemontana divaricata* (double flower variety). [α]_D²⁰ -13.5 (c, 0.021 in CHCl₃). λ_{\max} 237 (log ϵ 3.75); 282 (log ϵ 3.72); 298 (log ϵ 2.84) (EtOH).**16β-Hydroxy, N¹-Me: N¹-Methylvoafinine**

[181284-82-2]

C₂₀H₂₆N₂O₂ 326.438Trace alkaloid from leaves of *Tabernaemontana divaricata* (double flower variety). [α]_D²⁰ -39.1 (c, 0.069 in CHCl₃). λ_{\max} 217 (log ϵ 3.83); 275 (log ϵ 3.6); 303 (log ϵ 3.04) (EtOH).**(-) -form [120202-64-4]**Alkaloid from the roots of *Ervatamia yunnanensis* (Apocynaceae). Mp 153-154°. [α]_D¹⁵ -26 (c, 0.05 in CHCl₃).Kunesch, N. et al., *Bull. Soc. Chim. Fr.*, 1967, 2155 (uv, ir, pmr, ms, struct)Dugan, J.J. et al., *Helv. Chim. Acta*, 1967, **50**, 60 (isol, uv, ir, pmr, ms, struct)Bruneton, J. et al., *Phytochemistry*, 1974, **13**, 1963 (isol, uv, pmr, ms, struct, 12-Methoxyvoaphylline)Gomez-Gonzalez, C. et al., *CA*, 1977, **87**, 2343j; 1979, **90**, 138069u; 1982, **96**, 214301e (isol, spectra, Hecubine)Wenkert, E. et al., *Heterocycles*, 1979, **12**, 1439 (cmr)Stoekigt, J. et al., *Z. Naturforsch., C*, 1982, **37**, 857 (biosynth)Atta-ur-Rahman, et al., *Z. Naturforsch., B*, 1983, **38**, 117 (ms, cmr)Gui, L. et al., *Planta Med.*, 1988, **54**, 519 (Ervayunine)Éles, J. et al., *J.O.C.*, 2002, **67**, 7255-7260 (synth)Kam, T.-S. et al., *Org. Biomol. Chem.*, 2003, **1**, 1292-1297 (Voafinine, N-Methylvoafinine)**Voaphylline hydroxyindolenine****V-180***13-Ethyl-4,5,11,12,13,13a-hexahydro-2H-3,13-methanooxireno[9,10]azacycloundecino[5,4-b]indol-5a(1aH)-ol, 9CI* [18269-16-4]

Absolute Configuration

C₁₉H₂₄N₂O₂ 312.411

The hydrochloride has a quaternary hexacyclic struct. (cf. Rhazidigenine, R-72). Alkaloid from the leaves of *Voacanga africana* and *Tabernaemontana pubescens* (Apocynaceae). Amorph. [α]_D²⁵ -123 (CHCl₃).

Hydrochloride:

[34787-55-8 quaternary struct]

[α]_D²⁵ +4 (EtOH).**16-Oxo: Voalenine**


[180059-78-3]

C₁₉H₂₂N₂O₃ 326.394

Alkaloid from the leaves of *Tabernaemontana divaricata*. Light yellowish oil. [α]_D -445 (c, 0.04 in CHCl₃). λ_{\max} 204 (log ϵ 4.21); 226 (log ϵ 4.12); 296 (log ϵ 3.79) (EtOH).

Kunesch, N. *et al.*, *Bull. Soc. Chim. Fr.*, 1967, 3551 (*uv, ir, ms, struct*)Guilhem, J. *et al.*, *Acta Cryst. B*, 1970, **26**, 2029 (*cryst struct*)Wenkert, E. *et al.*, *Heterocycles*, 1979, **12**, 1439 (*cmr*)Mulamba, T. *et al.*, *J. Nat. Prod.*, 1981, **44**, 184 (*isol*)Kam, T.S. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 1292-1297 (*Voalenine*)**14,15-Voaphyllinediol**

V-181

(14*R*,15*S*)-formC₁₉H₂₆N₂O₂ 314.427

Stereochem. of isolates not certain. Voafinidine was definitely the 14*S*,15*S*-isomer, but the assignments of TC-A and TC-C may be reversed.

(14*R*,15*S*)-form**Alkaloid TC-A**

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae).

(14*S*,15*S*)-form**Voafinidine. Alkaloid TC-C**

[180059-77-2]


Minor alkaloid from leaves of *Tabernaemontana divaricata* and root bark of *Tabernaemontana chippii*. Light yellowish oil. [α]_D +57 (c, 0.18 in CHCl₃). λ_{\max} 233 (log ϵ 4.49); 287 (log ϵ 4.08); 295 (log ϵ 4.07) (EtOH).

19 ξ -Hydroxy: Alkaloid TC-BC₁₉H₂₆N₂O₃ 330.426

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Tentative struct. and stereochem.

van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400 (*TC alkaloids*)Kam, T.S. *et al.*, *Org. Biomol. Chem.*, 2003, **1**, 1292-1297 (*Voafinidine*)**Voastriectine**

V-182



Relative Configuration


C₁₉H₂₀N₂O 292.38

Alkaloid from *Tabernaemontana corymbosa*. Oil. [α]_D +321 (c, 0.23 in CHCl₃). λ_{\max} 209 (log ϵ 4.37); 231 (log ϵ 4.32); 283 (log ϵ 3.4); 294 (log ϵ 3.36); 307 (log ϵ 3.27); 321 (log ϵ 3.19) (no solvent reported).

Kam, T.S. *et al.*, *Tet. Lett.*, 2001, **42**, 4721-4723**Vobasenal**

V-183

[131653-77-5]



Absolute Configuration

C₂₀H₂₀N₂O₄ 352.389

Alkaloid from the leaves and stem bark of *Ervatamia polyneura* (Apocynaceae). [α]_D -62 (c, 0.5 in CHCl₃).

16-Epimer: 16-Epivobasenal

[131724-58-8]


C₂₀H₂₀N₂O₄ 352.389

Alkaloid from the leaves and stem bark of *Ervatamia polyneura* (Apocynaceae). [α]_D +26 (c, 0.4 in CHCl₃).

Clivio, P. *et al.*, *Phytochemistry*, 1990, **29**, 3007 (*isol, uv, ir, pmr, cmr, ms, struct*)**Vobasonidine**

V-184

[442156-08-3]

C₄₄H₅₄N₄O₇ 750.933


Alkaloid from the leaves of *Tabernaemontana corymbosa*. Light yellow oil.

[α]_D -80 (c, 0.31 in CHCl₃). λ_{\max} 229 (log ϵ 4.2); 286 (log ϵ 3.66); 295 (log ϵ 3.64) (EtOH).

Kam, T.S. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1027-1032 (*isol, uv, pmr, cmr*)**Vobatricine**

V-185

[442156-09-4]


C₄₀H₄₄N₄O₃ 628.813

Alkaloid from the stem bark of *Tabernaemontana corymbosa*. Light yellow oil. [α]_D -575 (c, 0.13 in CHCl₃). λ_{\max} 225 (log ϵ 4.62); 287 (log ϵ 4.16); 295 (log ϵ 4.15); 369 (log ϵ 4.36) (EtOH).

Kam, T.S. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1027-1032 (*isol, uv, pmr, cmr*)**Vobparicine**

V-186

[90930-64-6]



Absolute Configuration

C₃₉H₄₄N₄O₂ 600.802

Alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Active against gram-positive and gram-negative bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O. λ_{\max} 222; 290 (sh); 296 (sh); 304 (MeOH) (Berdy).

N⁴-Oxide: Vobparicine N-oxide

[98776-10-4]


C₃₉H₄₄N₄O₃ 616.802

Trace alkaloid from the root bark of *Tabernaemontana chippii* (Apocynaceae). Active against gram-positive and gram-negative bacteria. Sol. MeOH, CHCl₃; poorly sol. H₂O.

Van Beek, T.A. *et al.*, *Tet. Lett.*, 1984, **25**, 2057-2065 (*uv, cd, pmr, cmr, ms, struct*)Van Beek, T.A. *et al.*, *J. Ethnopharmacol.*, 1985, **14**, 315-318 (*activity*)Van Beek, T.A. *et al.*, *J. Nat. Prod.*, 1985, **48**, 400-403 (*N-oxide*)

Vobtusamine, 9CI

[84009-34-7]


C₄₃H₅₀N₄O₇ 734.891

Alkaloid from *Voacanga chalongiana* (Apocynaceae). Cryst. (MeOH). Mp 262° dec. [α]_D²⁵ -153 (c, 1.0 in CHCl₃). Eburnea-Aspidosperma dimer.

Danieli, B. et al., *J.O.C.*, 1983, **48**, 381 (*uv, ir, pmr, cmr, ms, cd, struct*)

Vobtusine

Papuanine
[19772-79-3]

C₄₃H₅₀N₄O₆ 718.891

Alkaloid from *Voacanga africana* and many other *Voacanga* spp., *Hedranthera (Callichilia) barteri* and some other *Callichilia* spp., and a number of other spp. in Apocynaceae. Mp 302° dec. [α]_D²² -352 (c, 0.96 in CHCl₃).

Hydrochloride (1:2): Mp 242-244°.

O-De-Me: 12-Demethylvobtusine. Alkaloid F†

[38790-81-7]

C₄₂H₄₈N₄O₆ 704.864

Alkaloid from the roots of *Hedranthera barteri* and the leaves of *Voacanga thouarsii* (Apocynaceae). Cryst. (Me₂CO/hexane/MeOH). Mp 295° (dec.). [α]_D²⁸ -273 (c, 0.035 in MeOH).

2-Deoxy: Goziline. 2-Deoxyvobtusine

[31148-61-5]

C₄₃H₅₀N₄O₅ 702.892

Alkaloid from leaves and root bark of *Hedranthera barteri*, leaves of *Voacanga africana* and *Voacanga grandifolia* (Apocynaceae). Cryst. (MeOH). Mp 305° (260°, 270°, 290°) dec. [α]_D -337 (-301, -355) (CHCl₃).

V-187

3'-Hydroxy: 3'-Hydroxyvobtusine

[65967-13-7]

C₄₃H₅₀N₄O₇ 734.891

Alkaloid from the root bark of *Voacanga chalongiana* (Apocynaceae). Amorph. solid.

3'-Oxo: 3'-Oxovobtusine. *Vobtusine lactam*

[50924-04-4]

C₄₃H₄₈N₄O₇ 732.875

Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). [α]_D -130 (CHCl₃).

3'-Oxo, N⁴-oxide: 3'-Oxovobtusine N-oxide

[50924-05-5]

C₄₃H₄₈N₄O₈ 748.874

Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). [α]_D -142 (c, 1 in CHCl₃).

3'-Oxo, 2-deoxy: 2-Deoxy-3'-oxovobtusine. 2-Deoxyvobtusine lactam. Alkaloid D†

[50924-02-2]

C₄₃H₄₈N₄O₆ 716.875

Alkaloid from the leaves of *Voacanga thouarsii* (Apocynaceae). Noncryst. Stereochem. of C-16 and C-14' unknown.

18-Oxo: 18-Oxovobtusine. *Vobtusine lactone*

[19772-81-7]

C₄₃H₄₈N₄O₇ 732.875

Alkaloid from *Voacanga africana*, *Voacanga grandifolia* and *Voacanga thouarsii* (Apocynaceae). Mp 310° dec. [α]_D²² -320 (CHCl₃).

18-Oxo, 2-deoxy: 2-Deoxy-18-oxovobtusine. 2-Deoxyvobtusine lactone

[19772-80-6]

C₄₃H₄₈N₄O₆ 716.875

Alkaloid from the leaves of *Voacanga africana* (Apocynaceae). Mp 305° dec. [α]_D²² -348 (CHCl₃). Stereochem. at C-16 and C-14' not rigidly assigned.

14'-Epimer, 2-deoxy, 2,16-didehydro: OVERRINE. *Anhydrodihydroamataine*

[31222-09-0]

C₄₃H₄₈N₄O₅ 700.876

Trace alkaloid from the root bark of *Hedranthera barteri* (Apocynaceae). Cryst. (MeOH). Mp 265° dec. [α]_D²⁷ -496 (c, 0.613 in CHCl₃).

Gorman, A.A. et al., *Helv. Chim. Acta*, 1966, **49**, 2072 (*isol, ir, uv, pmr, ms, struct*)

Poisson, J. et al., *Tetrahedron*, 1966, **22**, 1075 (*uv, ir, pmr, ms, struct*)

Kunesch, N. et al., *Bull. Soc. Chim. Fr.*, 1970, 4370 (18-Oxovobtusine, 2-Deoxy-18-oxovobtusine)

Agwada, V. et al., *Helv. Chim. Acta*, 1970, **53**, 1567; 1977, **60**, 2830 (*Goziline, OVERRINE*)

Naranjo, J. et al., *Helv. Chim. Acta*, 1972, **55**, 1849 (12-Demethylvobtusine, *Goziline*)

Lefebvre-Soubeyran, O. et al., *Acta Cryst. B*, 1973, **29**, 2855 (*cryst struct*)

Rolland, Y. et al., *Phytochemistry*, 1973, **12**, 2039 (12-Demethylvobtusine)

Majumder, P.L. et al., *Phytochemistry*, 1974, **13**, 1261 (*Goziline*)

Rolland, Y. et al., *Bull. Soc. Chim. Fr.*, 1975, 2503 (3'-Oxovobtusine, 3'-Oxovobtusine N-oxide, 12-Demethylvobtusine)


Rolland, Y. et al., *J.O.C.*, 1976, **41**, 3270 (*cmr*)

Danieli, B. et al., *Heterocycles*, 1980, **14**, 201 (3-Hydroxyvobtusine)

Vochysine

V-189

3,4-Dihydro-2-(4-hydroxyphenyl)-7-methoxy-8-(2-pyrrolidinyl)-2H-1-benzopyran-5-ol, 9CI
[88607-66-3]

C₂₀H₂₃NO₄ 341.406


Alkaloid from the fruits of *Vochysia guianensis* (Vochysiaceae). Cryst. (MeOH). Mp 138°. [α]_D²⁰ 0 (CHCl₃).

Baudouin, G. et al., *J. Nat. Prod.*, 1983, **46**, 681 (*isol, uv, ir, pmr, cmr, ms, struct*)

Volutamide A

V-190

[176181-95-6]


C₁₅H₂₂Br₂N₂O₂ 422.159

Alkaloid from the Atlantic bryozoan *Amathia convoluta*. Feeding deterrent to hydroid larvae. Oil. Sol. MeOH, butanol. λ_{\max} 285 (ε 4200); 292 (ε 4100) (MeOH) (Berdy).

Montanari, A.M. et al., *Tetrahedron*, 1996, **52**, 5371-5380 (*isol, uv, ir, pmr, cmr, ms, struct*)

Volutamide B

V-191

C₂₀H₃₃Br₂N₃O₂ 507.308**(S)-form** [176181-96-7]

Alkaloid from the Atlantic bryozoan *Amathia convoluta*. Exhibits antifeedant activity to hydroid larvae. Oil. Sol. MeOH, butanol. [α]_D 0 (c, 0.053 in MeOH). λ_{\max} 287 (ε 1600); 292 (ε 1500) (MeOH) (Berdy).

N¹⁵-Me: Volutamide C

[176181-97-8]


C₂₁H₃₅Br₂N₃O₂ 521.334

From *Amathia convoluta*. Exhibits antifeedant activity to hydroid larvae. Oil. Sol. MeOH, butanol. [α]_D 0 (c, 0.032 in MeOH). λ_{\max} 285 (ε 2100); 293 (ε 1900) (MeOH) (Berdy).

Montanari, A.M. et al., *Tetrahedron*, 1996, **52**, 5371-5380 (*isol, uv, ir, pmr, cmr, ms, cd, struct*)

Volutamide D

V-192



$C_{22}H_{27}Br_3N_2O_3$ 607.179
Sol. MeOH, butanol. λ_{max} 287 (ϵ 3500)
(MeOH) (Berdy).


(R)-form [176182-10-8]

Alkaloid from the Atlantic bryozoan *Amathia convoluta*. Feeding deterrent to hydroid larvae, shows antiinflammatory props. Cryst. (MeOH). $[\alpha]_D^{20}$ -12.9 (c, 0.010 in MeOH). Abs. config. not certain.

Montanari, A.M. *et al.*, *Tetrahedron*, 1996, **52**, 5371 (*isol, uv, ir, pmr, cmr, ms, cryst struct*)

Volutamide E

V-193



$C_{25}H_{34}Br_3N_3O_3$ 664.274
Sol. MeOH, butanol.

(R)-form [176182-11-9]


Alkaloid from the Atlantic bryozoan *Amathia convoluta*. Feeding deterrent to hydroid larvae, shows antiinflammatory props. Oil. $[\alpha]_D^{20}$ -15.8 (c, 0.014 in MeOH). Abs. config. not certain. λ_{max} 287 (ϵ 3800) (MeOH).

Montanari, A.M. *et al.*, *Tetrahedron*, 1996, **52**, 5371-5380 (*isol, uv, ir, pmr, cmr, struct*)

Vomilene

V-194

1,2,19,20-Tetrahydro-1-demethylajmalan-17,21-diol 17-acetate, 9CI
[6880-50-8]



Absolute
Configuration

$C_{21}H_{22}N_2O_3$ 350.416
Alkaloid from roots of *Rauwolfia vomitoria* (Apocynaceae). Cryst. (MeOH). Mp 207°. $[\alpha]_D^{20}$ -72 (c, 0.5 in Py).

O-β-D-Glucopyranoside: Raucaffricine
[31282-07-2]

$C_{27}H_{32}N_2O_8$ 512.558
Alkaloid from the roots of *Rauwolfia caffra*. A major constit. of cell cultures of *Rauwolfia serpentina* (Apocynaceae). Plates (MeOH/EtOAc). Mp 186° (frothing) Mp 220° (double Mp). Originally considered to be the galactoside of Vomilene.

19,20-Dihydro, Ac: 21-Acetyl-19,20-dihydrovomilene. Alkaloid RCLM 1

$C_{23}H_{26}N_2O_4$ 394.469
Alkaloid tentatively identified in leaves of *Rauwolfia caffra* (Apocynaceae).

Deoxy: Vinorine. Alkaloid RCL 2. Alkaloid RP7. *21-Deoxyvomilene*
[34020-07-0]

$C_{21}H_{22}N_2O_2$ 334.417
Alkaloid from *Rauwolfia caffra*, *Rauwolfia perakensis* and *Vinca minor* (Apocynaceae). Amorph.; cryst. (MeOH)(as nitrate). Mp 235-237° (nitrate). Identity of the various alkaloids descr. under this struct. not well establ. One sample was opt. inactive and prob. racemic. The Mp of the nitrate refers to the salt of Alkaloid RP 7.

10-Methoxy: Majorinine. *10-Methoxy-21-hydroxyvinorine*

[64986-27-2]
 $C_{22}H_{24}N_2O_4$ 380.443
Alkaloid from *Vinca minor* (Apocynaceae). Mp 195-196°.

10-Methoxy, deoxy: 10-Methoxyvinorine

$C_{22}H_{24}N_2O_3$ 364.443
Alkaloid from aerial parts of *Vinca major* (Apocynaceae) and from *Vinca erecta* (Apocynaceae). Amorph. solid. $[\alpha]_D^{20}$ +23.5 (c, 0.2 in $CHCl_3$).

12-Methoxy, deoxy, O-de-Ac: O-Deacetyl-12-methoxyvinorine

[70509-81-8]
 $C_{20}H_{22}N_2O_2$ 322.406
Alkaloid from stem bark of *Rauwolfia vomitoria* (Apocynaceae). Off-white amorph. powder.

Stereoisomer: Raucaffricine

[107585-43-3]
 $C_{21}H_{22}N_2O_3$ 350.416
Alkaloid from *Rauwolfia caffra* (Apocynaceae). Cryst. (EtOAc). Mp 200-201°. Config. not given in CAS.

[1354-08-1]

Hofmann, A. *et al.*, *Helv. Chim. Acta*, 1957, **40**, 1866

Taylor, W.I. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 611 (*ir, uv, pmr, struct*)

Kiang, A.K. *et al.*, *Tetrahedron*, 1966, **22**, 3293 (*Alkaloid RP7*)

Khan, M.A. *et al.*, *Tet. Lett.*, 1970, 5137 (*Raucaffricine, isol, uv, pmr*)

Meisel, H. *et al.*, *Tet. Lett.*, 1971, 1291 (*Vinorine*)

Khalmirzhaev, M.M. *et al.*, *Khim. Prir. Soedin.*, 1973, **9**, 681; *Chem. Nat. Compd. (Engl. Transl.)*, 1973, **9**, 657 (*10-Methoxyvinorine*)

Il'yashenko, L.I. *et al.*, *Khim. Prir. Soedin.*, 1977, **13**, 382; *Chem. Nat. Compd. (Engl. Transl.)*, 1977, **13**, 324 (*Majorinine*)

Sabri, N.N. *et al.*, *Phytochemistry*, 1978, **17**, 2023 (*O-Deacetyl-12-methoxyvomilene*)

Libot, F. *et al.*, *Phytochemistry*, 1980, **19**, 989 (*cmr*)

Khan, M.A. *et al.*, *Z. Naturforsch., B*, 1982, **37**, 494 (*Raucaffricine, isol, cmr*)

Nasser, A.M.A.G. *et al.*, *Phytochemistry*, 1983, **22**, 2297 (*derivis*)

Schübel, H. *et al.*, *Helv. Chim. Acta*, 1984, **62**, 2078 (*Raucaffricine, pmr, cmr, ms, struct*)

Khan, M.A. *et al.*, *CA*, 1987, **106**, 153050y (*Raucaffricine*)

Takayama, H. *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 266 (*synth, config*)


Ruyter, C.M. *et al.*, *Helv. Chim. Acta*, 1991, **74**, 1707 (*Raucaffricine, synth*)

Atta-ur-Rahman, *et al.*, *Phytochemistry*, 1995, **38**, 1057 (*10-Methoxyvinorine*)

Vulgaxanthin II

V-195

4-[(1,3-Dicarboxypropyl)imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, 9CI. Glutamic acid betaxanthin
[1047-87-6]



$C_{14}H_{16}N_2O_8$ 340.289

Yellow pigment from beetroot (*Beta vulgaris*), *Portulaca grandiflora* and *Chenopodium rubrum* cell cultures (Chenopodiaceae, Portulacaceae).

5-Amide: Vulgaxanthin I. Glutamidebetaxanthin

[904-62-1]
 $C_{14}H_{17}N_3O_7$ 339.304

Yellow pigment from *Beta*, *Portulaca* and *Conophytum* spp. and other plants (Chenopodiaceae, Aizoaceae, Portulacaceae). Food colourant.

5-Alcohol: Humilixanthin

[111534-70-4]
 $C_{14}H_{18}N_2O_7$ 326.305

Isol. from the fruits of *Rivina humilis*. Also detected in *Phytolacca acinosa*, *Phytolacca bogotensis*, *Delosperma luteum*, *Lampranthus aurantiacus*, *Lampranthus peersii* and *Portulaca grandiflora*, and in the yellow-coloured root of *Beta vulgaris* (Phytolaccaceae, Aizoaceae, Portulacaceae, Chenopodiaceae).

Piattelli, M. *et al.*, *Phytochemistry*, 1965, **4**, 121 (*isol, ir, uv, struct*)

Singer, J.W. *et al.*, *J. Food Sci.*, 1980, **45**, 489 (*isol, use*)

Strack, D. *et al.*, *Phytochemistry*, 1987, **26**, 2285 (*isol, uv, pmr, ms, struct, Humilixanthin*)

Vulracine

V-196


$C_{18}H_{19}NO_3$ 297.353

Struct. unknown. Alkaloid from *Berberis vulgaris* (barberry) (Berberidaceae). Fine needles (Me_2CO). Mp 164°. $[\alpha]_D^{24}$ -114 (c, 0.2 in $CHCl_3$).

Döpke, W. *et al.*, *Naturwissenschaften*, 1963, **50**, 595

Waianacamine A

[256472-04-5]




C₁₇H₁₉Br₂ClN₂O₃ 494.609
Alkaloid from an undescribed Verongid sponge. [α]_D²⁷ +2.6 (c, 0.23 in MeOH). λ_{max} 209 (log ε 4.36); 310 (log ε 3.89) (MeOH).

Lacy, C. et al., *J. Nat. Prod.*, 2000, **63**, 119-121 (isol, pmr, cmr, uv)

Waianacamine B

[256472-05-6]




C₁₇H₁₉Br₂ClN₂O₃ 494.609
Alkaloid from an undescribed Verongid sponge. [α]_D²⁷ -20.9 (c, 0.07 in MeOH). λ_{max} 209 (log ε 4.31); 309 (log ε 3.84) (MeOH).

Lacy, C. et al., *J. Nat. Prod.*, 2000, **63**, 119-121 (isol, pmr, cmr, uv)

Wakayin

2,3,5,7-Tetrahydro-9-(1H-indol-3-yl)-6H-dipyrrolo[4,3,2-de:2',3'-h]quinolin-6-one, 9CI
[134781-25-2]




C₂₀H₁₄N₄O 326.357
Related to Tsitsikammamine A, T-650. Alkaloid from the ascidian *Clavelina* sp. Cytotoxic agent. Exhibits antimicrobial activity. Topoisomerase inhibitor. λ_{max} 226 (ε 24600); 299 (ε 14000); 413 (ε 5100); 474 (ε 4600) (MeOH/KOH) (Derep). λ_{max} 222 (ε 33500); 243 (ε 28200); 311 (ε 15000) (MeOH) (Derep). λ_{max} 226 (ε 24600); 270 (ε 14000); 412 (ε 5100); 473 (ε 4600) (MeOH/NaOH) (Berdy).

W-1

Copp, B.R. et al., *J.O.C.*, 1991, **56**, 4596-4597 (isol, uv, ir, pmr, cmr, struct)

Wallemia A

5-Hydroxy-2,6-dimethyl-11-(1H-pyrrol-2-yl)-4,6,8,10-undecatetraen-3-one, 9CI
[50657-01-7]



C₁₇H₂₁NO₂ 271.358
Pigment from *Wallemia sebi*. Orange needles (cyclohexane). Mp 107°. Struct. revised in 1984.

3'-Chloro: Wallemia E

[50656-96-7]
C₁₇H₂₀ClNO₂ 305.803
Isol. from *Wallemia sebi*. Orange cryst. (Et₂O/petrol). Mp 122-124°.


6Z-Isomer: Wallemia B

[50657-00-6]
C₁₇H₂₁NO₂ 271.358
Isol. from *Wallemia sebi*. Tentative struct. The struct. given in CAS is erroneous following the structural revision of Wallemia A in 1984.

Badar, Y. et al., *J.C.S. Perkin I*, 1973, 1416; 1977, 1372 (isol, struct, uv, pmr, ir, ms, cryst struct)
Ahmed, F.R. et al., *J. Chem. Res., Synop.*, 1984, 178 (struct)
Ahmed, F.R. et al., *J.C.S. Perkin I*, 1984, 1577 (synth)

Wallemia C

2,2-Dimethyl-5-[5-methyl-6-(1H-pyrrol-2-yl)-1,3,5-hexatrienyl]-3(2H)-furanone, 9CI
[50656-99-0]



C₁₇H₁₉NO₂ 269.343
Pigment from *Wallemia sebi*. Orange prisms (CHCl₃/cyclohexane). Mp 206-208°.

4'-Chloro: Wallemia F

[50656-98-9]
C₁₇H₁₈ClNO₂ 303.787
Isol. from *Wallemia sebi*. Red cryst. (cyclohexane). Mp 206-208°.


Geom. isomer: Wallemia D

[50656-97-8]
C₁₇H₁₉NO₂ 269.343
Pigment from *Wallemia sebi*. Config. undetermined.
Badar, Y. et al., *J.C.S. Perkin I*, 1973, 1416 (isol, struct, uv, ms, pmr, ir)
Ito, M. et al., *J.C.S. Perkin I*, 1981, 3255 (synth)
Ahmed, F.R. et al., *J. Chem. Res., Synop.*, 1984, 178 (pmr, struct)

W-4

Waltherine A

[243464-59-7]




Absolute Configuration

C₃₁H₄₂N₄O₄ 534.697
Alkaloid from the root bark of *Waltheria douradinha*. Needles (CHCl₃/Et₂O). Mp 234-235°. [α]_D²⁰ -229.8 (c, 0.24 in MeOH).

Morel, A.F. et al., *Phytochemistry*, 1999, **51**, 473-477 (isol, pmr, cmr, ms)

Waltherine C

[256445-94-0]




W-7

C₃₀H₃₇N₅O₄ 531.653
Alkaloid from the bark of *Waltheria douradinha*. Amorph. powder. [α]_D -182 (c, 0.2 in CHCl₃).

Morel, A.F. et al., *Tet. Lett.*, 1999, **40**, 9205-9209 (isol, pmr, cmr, ms)

Waltherione A



Relative Configuration

C₂₃H₂₃NO₅ 393.438
Stereochem. revised in 2008. Alkaloid from the root bark of *Waltheria douradinha*. Solid (CHCl₃/MeOH). Mp 206-207.5°. [α]_D²⁵ -25.5 (c, 0.04 in CHCl₃).

10,13-Diepimer: Waltherione B


C₂₃H₂₃NO₅ 393.438
Alkaloid from the stems of *Waltheria douradinha*. Cryst. (CHCl₃/MeOH). Mp 170.5-171°. [α]_D²⁵ +2.1 (c, 0.01 in CHCl₃).

Hoelzel, S.C.S.M. et al., *Phytochemistry*, 2005, **66**, 1163-1167 (isol, pmr, cmr)
Gressier, V. et al., *Phytochemistry*, 2008, **69**, 994-999 (cryst struct, Waltherione B)

W-8

Warifteine

[30996-86-2]

C₃₆H₃₆N₂O₆ 592.69Alkaloid from the rhizomes of *Cissampelos ovalifolia* (Menispermaceae).O⁷-Me: **Cissampareine**. *Methylwarifteine* [32728-54-4]C₃₇H₃₈N₂O₆ 606.717Alkaloid from *Cissampelos pareira* whole plant and *Cissampelos ovalifolia* rhizomes (Menispermaceae). Shows antineoplastic props. Rods (Me₂CO). Mp 239-240° dec. [α]_D²⁶ -111 (c, 1.05 in CHCl₃). Log P 6.78 (uncertain value) (calc). λ_{max} 282 (ε 10000); 320 (ε 4000) (EtOH) (Berdy).Di-Me ether: **Dimethylwarifteine**. *O*-*Methylcissampareine*

[7678-91-3]

C₃₈H₄₀N₂O₆ 620.744Alkaloid from the rhizomes of *Cissampelos ovalifolia* (Menispermaceae). Rods (Me₂CO). Mp 192-194°. [α]_D²⁶ -121 (c, 1.36 in CHCl₃).1ξ,2-Dihydro: **Dihydrowarifteine**

[30996-85-1]

C₃₆H₃₈N₂O₆ 594.706Alkaloid from the rhizomes of *Cissampelos ovalifolia* (Menispermaceae).1ξ,2-Dihydro, O⁷-Me: **Methyldihydrowarifteine**

[3114-26-8]


C₃₇H₄₀N₂O₆ 608.733Alkaloid from the rhizomes of *Cissampelos ovalifolia* (Menispermaceae). Prisms (MeOH). Mp 208-212°. [α]_D²⁷ -157 (c, 0.7 in CHCl₃). Phys. props. refer to synthetic material.1ξ,2-Dihydro, di-Me ether: **Dimethyldihydrowarifteine**. *O*-*Methylcissampareine*

[30994-04-8]


C₃₈H₄₂N₂O₆ 622.76Alkaloid from the rhizomes of *Cissampelos ovalifolia* (Menispermaceae).α-Oxo, O⁷-Me: **Roraimine**C₃₇H₃₆N₂O₇ 620.701Alkaloid from the roots of *Cissampelos sympodialis*. Amorph. dark yellow powder.Kupchan, S.M. *et al.*, *J.A.C.S.*, 1966, **88**, 4212 (*Cissampareine* isol, uv, ir, pmr, ms, struct, abs config)Snedden, W. *et al.*, *Org. Mass Spectrom.*, 1970, **4**, 607 (*struct, ms, derivs*)Borkakoti, N. *et al.*, *Acta Cryst. B*, 1978, **34**, 482; 490 (*cryst struct, Dimethylwarifteine, Cissampareine*)Ralph, I. *et al.*, *Heterocycles*, 1981, **16**, 2105 (*biosynth*)de Lira, G.A. *et al.*, *Fitoterapia*, 2002, **73**, 356-358 (*isol, pmr, cmr*)Mukherjee, R. *et al.*, *Magn. Reson. Chem.*, 2003, **41**, 213-218 (*pmr, cmr*)

W-9

Wasalexin A

C₁₃H₁₄N₂O₂S₂ 294.398Constit. of *Thlaspi arvense* (stinkweed) and *Wasabia japonica* (Japanese horseradish). Phytoalexin. Yellow solid.*(Z)*-Isomer: **Wasalexin B**C₁₃H₁₄N₂O₂S₂ 294.398Constit. of *Wasabia japonica* (Japanese horseradish). Phytoalexin. Yellow solid.Pedras, M.S.C. *et al.*, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 3015-3020 (*isol, synth, pmr, cmr, ms*)Pedras, M.S.C. *et al.*, *Phytochemistry*, 2003, **64**, 949-956 (*isol, synth, activity*)Pedras, M.S.C. *et al.*, *J.O.C.*, 2005, **70**, 1828-1834 (*synth*)Pedras, M.S.C. *et al.*, *Org. Biomol. Chem.*, 2006, **4**, 3526-3535 (*metab*)


Watasemycin A

C₁₆H₂₀N₂O₃S₂ 352.478Related to Thiazostatin A, T-373. Prod. by the marine-derived *Streptomyces* sp. TP-A0597. Active against gram-positive and -negative bacteria. Pale yellow powder. Mp 62-65°. [α]_D²⁸ +20.5 (c, 0.2 in CHCl₃). λ_{max} 212 (log ε 4.17); 252 (log ε 3.78); 319 (log ε 3.45) (MeOH).2-Epimer: **Watasemycin B**C₁₆H₂₀N₂O₃S₂ 352.478Prod. by *Streptomyces* sp. TP-A0597.Active against gram-positive and -negative bacteria. Pale yellow powder. Mp 58-60°. [α]_D²⁸ -2.5 (c, 0.2 in CHCl₃). λ_{max} 216 (log ε 4.21); 286 (log ε 3.87); 322 (log ε 3.85) (MeOH).Sasaki, T. *et al.*, *J. Antibiot.*, 2002, **55**, 249-255 (*isol, uv, pmr, cmr*)

Welwitindolinone A isonitrile

W-10

[159934-03-9]



Absolute Configuration


W-12

C₂₁H₂₁ClN₂O 352.862Alkaloid from the freshwater blue-green alga *Hapalosiphon welwitschii*, also from *Westiella intricata* and *Fischerella muscicola*. [α]_D +377 (c, 0.078 in CH₂Cl₂). λ_{max} 212 (ε 1720); 250 (ε 4000); 290 (ε 900) (MeOH).Stratmann, K. *et al.*, *J.A.C.S.*, 1994, **116**, 9935-9942 (*isol, uv, ir, pmr, cmr*)Reisman, S.E. *et al.*, *J.A.C.S.*, 2008, **130**, 2087-2100 (*synth*)Richter, J.M. *et al.*, *J.A.C.S.*, 2008, **130**, 17938-17954 (*synth*)

Welwitindolinone B isothiocyanate

W-13

[159993-29-0]



Absolute Configuration

C₂₁H₂₁ClN₂O₂S 400.928The abs. config. shown is based on that determined crystallographically for Welwistatin. Alkaloid from the blue-green algae *Hapalosiphon welwitschii* (freshwater) and *Westiella intricata* (terrestrial).N-Me: **N-Methylwelwitindolinone B isothiocyanate**

[159189-03-4]

C₂₂H₂₃ClN₂O₂S 414.955From *Hapalosiphon welwitschii*. [α]_D -149 (c, 0.071 in CH₂Cl₂).13,14-Didehydro: **Welwitindolinone C isothiocyanate**

[159934-04-0]

C₂₁H₁₉ClN₂O₂S 398.912From *Hapalosiphon welwitschii* and *Westiella intricata*. [α]_D -283 (c, 0.148 in CH₂Cl₂).13,14-Didehydro, N-Me: **Welwistatin**. *N*-*Methylwelwitindolinone C isothiocyanate*

[159189-05-6]

C₂₂H₂₁ClN₂O₂S 412.939Major alkaloid in *Hapalosiphon welwitschii* and *Westiella intricata*. MDR inhibitor. Cytostatic agent. [α]_D -278 (c, 0.77 in CH₂Cl₂). λ_{max} 210 (ε 24500); 258 (ε 8930); 285 (ε 2300) (MeOH) (Berdy).13,14-Didehydro, N-Me, isocyanide: **N-Methylwelwitindolinone C isonitrile**

[159189-04-5]

C₂₂H₂₁ClN₂O₂ 380.873Alkaloid from terrestrial *Hapalosiphon welwitschii*. [α]_D -117 (c, 0.25 in CH₂Cl₂). Has -NC replacing -NCS. λ_{max} 218 (ε 26200); 260 (ε 16900); 292 (ε 16900) (MeOH) (Berdy).3-Hydroxy, 13,14-didehydro, N-Me: **3-Hydroxy-N-methylwelwitindolinone C isothiocyanate**C₂₂H₂₁ClN₂O₃S 428.938

Alkaloid from terrestrial *Fischerella major*. $[\alpha]_D^{25}$ -290 (c, 2.6 in CH_2Cl_2). λ_{max} 208 (ϵ 31100); 212 (ϵ 26500); 259 (ϵ 5680) (MeOH).

3-Hydroxy, 13,14-didehydro, N-Me, isocyanide: 3-Hydroxy-N-methylwelwitindolinone C isonitrile

$\text{C}_{22}\text{H}_{21}\text{ClN}_2\text{O}_3$ 396.872

Alkaloid from terrestrial *Fischerella muscicola*. $[\alpha]_D^{25}$ -103 (c, 0.4 in CH_2Cl_2). λ_{max} 220 (ϵ 43300); 266 (ϵ 13200); 301 (ϵ 3600) (MeOH).

3-Epimer: 3-Epiwelwitindolinone B isothiocyanate

[159249-49-7]

$\text{C}_{21}\text{H}_{21}\text{ClN}_2\text{O}_2\text{S}$ 400.928


From *Hapalosiphon welwitschii*. Struct. diag. in paper erroneously shows N-Me group.

Stratmann, K. *et al.*, *J.A.C.S.*, 1994, **116**, 9935-9942 (*isol, uv, ir, pmr, cmr*)

Jimenez, J.I. *et al.*, *J. Nat. Prod.*, 1999, **62**, 569-572 (3-Hydroxy-N-methylwelwitindolinone C derivs)

Avedaño, C. *et al.*, *Curr. Org. Synth.*, 2004, **1**, 65-82 (*Welwistatin, rev*)

Welwitindolinone D isonitrile W-14



$\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_4$ 362.384

N-Me: N-Methylwelwitindolinone D isonitrile

[224559-35-7]


$\text{C}_{22}\text{H}_{20}\text{N}_2\text{O}_4$ 376.411

Alkaloid from terrestrial *Fischerella major*. $[\alpha]_D^{25}$ -30 (c, 0.37 in CH_2Cl_2). λ_{max} 207 (ϵ 3910); 213 (ϵ 3390); 267 (ϵ 824) (MeOH).

Jimenez, J.I. *et al.*, *J. Nat. Prod.*, 1999, **62**, 569-572 (*isol, uv, ir, pmr, cmr, ms*)

Wewakazole W-15

[494798-69-5]



$\text{C}_{59}\text{H}_{72}\text{N}_{12}\text{O}_{12}$ 1141.291


Isol. from a *Lyngbya majuscula*. Glassy oil. $[\alpha]_D^{25}$ -46.8 (c, 0.41 in MeOH). λ_{max} 213 (log ϵ 4.55); 221 (log ϵ 4.39)

(MeOH).

Nogle, L.M. *et al.*, *Org. Lett.*, 2003, **5**, 3-6 (*isol, pmr, cmr*)

Wisandine W-16

1-[5-(6-Methoxy-1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]pyrrolidine, 9CI. 1-[5-(2-Methoxy-4,5-methylenedioxyphenyl)pentadienyl]pyrrolidine. Okolasine. 6-Methoxytrichostachine



$\text{C}_{17}\text{H}_{19}\text{NO}_4$ 301.341

λ_{max} 250 (ϵ 10000); 304 (ϵ 14500); 309 (ϵ 13200); 375 (ϵ 21000) (prob. EtOH) (Derep).

(E,E)-form [63424-62-4]

Alkaloid from the seeds and roots of *Piper guineense* (Piperaceae). Yellow cryst. (EtOAc/Et₂O). Mp 171-173°.

4,5-Dihydro: α,β -Dihydrowisandine

[64494-92-4]

$\text{C}_{17}\text{H}_{21}\text{NO}_4$ 303.357

Alkaloid from seeds of *Piper guineense* (Piperaceae). Cryst. (hexane). Poorly sol. hexane. Mp 82-84°. λ_{max} 300 (ϵ 6020) (MeOH) (Berdy).

Addae-Mensah, I. *et al.*, *Phytochemistry*, 1977, **16**, 757 (*isol, uv, ir, pmr, ms, struct, synth*)

Sondengam, B.L. *et al.*, *Phytochemistry*, 1977, **16**, 1121 (*deriv*)

Sondengam, B.L. *et al.*, *Tet. Lett.*, 1977, 367 (*isol, pmr, struct*)

Scharf, H.D. *et al.*, *Annalen*, 1978, 573 (*synth, ir, pmr, ms*)

Linke, S. *et al.*, *Tetrahedron*, 1978, **34**, 1979 (*synth*)

Vig, O.P. *et al.*, *Indian J. Chem., Sect. B*, 1979, **17**, 427-429 (*synth, deriv*)

Döpke, W. *et al.*, *Z. Chem.*, 1981, **21**, 358 (*deriv*)

Wisanine W-17

1-[5-(6-Methoxy-1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine, 9CI. 1-[5-(2-Methoxy-4,5-methylenedioxyphenyl)pentadienyl]piperidine. 2'-Methoxy-piperine

As Wisandine, W-16 with n = 2

$\text{C}_{18}\text{H}_{21}\text{NO}_4$ 315.368

(E,E)-form [61756-56-7]

Alkaloid from the roots of *Piper guineense* (Piperaceae). Active against gram-positive bacteria. Yellow prismatic cryst. (EtOAc/Et₂O). Mp 179-181°. λ_{max} 250 (ϵ 10000); 304 (ϵ 14150); 309 (ϵ 13180); 318 (ϵ 20890); 371 (ϵ 26300) (MeOH) (Berdy).

4,5-Dihydro: 4,5-Dihydro-2'-methoxy-piperine. 4,5-Dihydrowisanine

[62926-59-4]

$\text{C}_{18}\text{H}_{23}\text{NO}_4$ 317.384

Alkaloid from *Piper guineense* (Piperaceae). Active against gram-positive

bacteria. Cryst. (C_6H_6 /hexane). Poorly sol. hexane. Mp 99-100°. λ_{max} 300 (ϵ 6000) (MeOH) (Berdy).

(2E,4Z)-form

Piperx

[69699-10-1]

Minor alkaloid from the root bark of *Piper guineense* (Piperaceae). Yellow needles (EtOAc). Mp 193-195° (189-190°).

Vig, O.P. *et al.*, *Indian J. Chem., Sect. B*, 1977, **15**, 950 (*synth*)

Addae-Mensah, I. *et al.*, *Phytochemistry*, 1977, **16**, 757 (*isol, uv, ir, pmr, ms, struct*)

Sondengam, B.L. *et al.*, *Tet. Lett.*, 1977, 69 (*deriv*)

Scharf, H.D. *et al.*, *Annalen*, 1978, 573 (*synth, ir, uv, pmr*)

Linke, S. *et al.*, *Tetrahedron*, 1978, **34**, 1979 (*synth, ir, pmr, activity*)

Addae-Mensah, I. *et al.*, *Planta Med.*, 1981, **41**, 200 (*isol, ir, pmr, ms, struct*)

Woode, K.A. *et al.*, *J. Nat. Prod.*, 1984, **47**, 1024 (*cryst struct*)

Okwute, S.K. *et al.*, *Tetrahedron*, 1984, **40**, 2541 (*synth*)

Withananine W-18

Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Mp 75-80° (35-40°).

Picrate: Mp 176-178°.

Aurichloride: Mp 115-119°.

Majumdar, D.N. *et al.*, *Curr. Sci.*, 1952, **21**, 46; *CA*, **47**, 2184b

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f

Withanine W-19

$\text{C}_{44}\text{H}_{80}\text{N}_2\text{O}_{12}$ 829.122

Struct. unknown. Alkaloid from *Withania somnifera* (Solanaceae). Cryst. + 1CHCl₃. Mp 87-88° dec.


Aurichloride: Mp 170-175°.

Majumdar, D.N. *et al.*, *Curr. Sci.*, 1952, **21**, 46; *CA*, **47**, 2184b

Majumdar, D.N. *et al.*, *Indian J. Pharm.*, 1955, **17**, 158-161; *CA*, **50**, 3713f

Wondonine A W-20

[336825-31-1]



$\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_8\text{S}$ 503.532

Isol. from the association of sponges, *Jaspis* sp. and *Poecillastra wondoensis*. Yellow gum (as Na salt). $[\alpha]_D^{25}$ -4.8 (c, 0.12 in MeOH) (Na salt). λ_{max} 210 (log ϵ 4.16); 264 (log ϵ 3.94) (MeOH) (Na salt).

Diastereomer: Wondonine B

[336825-33-3]

$\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_8\text{S}$ 503.532


Isol. from the association of sponges, *Jaspis* sp. and *Poecillastra wondoensis*. Yellow gum (as Na salt). $[\alpha]_D^{25} +3.6$ (c, 0.2 in MeOH) (Na salt). λ_{\max} 211 (log ϵ 4.23); 264 (log ϵ 3.25) (MeOH) (Na salt).

Shin, J. et al., *Tet. Lett.*, 2001, **42**, 1965-1968 (*Wondonines A,B*)

Woodinine

W-21

[116339-96-9]



Absolute configuration

C₁₇H₂₂BrN₃ 348.285

Alkaloid from the ascidian *Eudistoma fragum*. Mp 112-113°. $[\alpha]_D -38$ (c, 0.6 in MeOH). $[\alpha]_D^{23} -80.8$ (c, 0.6 in MeOH) (synthetic). λ_{\max} 230 (ϵ 38000); 291 (ϵ 8510); 300 (sh) (ϵ 7080) (EtOH) (De-rep).

Debitus, C. et al., *J. Nat. Prod.*, 1988, **51**, 799 (isol, uv, pmr, cmr, ms, cd, struct)

McNulty, J. et al., *Tet. Lett.*, 1991, **32**, 4875 (synth, abs config)

Mahboobi, S. et al., *Arch. Pharm. (Weinheim, Ger.)*, 1992, **325**, 249; 1993, **326**, 33 (synth, pmr, abs config)


McNulty, J. et al., *J.C.S. Perkin 1*, 1994, 1329 (synth)

Mahboobi, S. et al., *J. Nat. Prod.*, 1997, **60**, 587 (pmr, cmr, cryst struct)

Worenine

W-22

5,6-Dihydro-14-methylbis[1,3]benzodioxolo[5,6-a:5',6'-g]quinolinizinium(1+), 9CI. *13-Methylpseudocoptisine* [38763-29-0]

C₂₀H₁₆NO₄⁺ 334.351

Probable struct. Originally proposed to have the struct. now given to Corysamine, C-699. Synthetic 13-methylpseudocoptisine has props. closely similar to those of Worenine, although direct comparison has not been possible. Quaternary alkaloid from *Coptis japonica* and *Coptis chinensis* (Ranunculaceae).

Chloride: [38763-54-1]

C₂₀H₁₆ClNO₄ 369.803

Yellow needles (H₂O). Mp 295° (nat.) Mp 288-289° (synthetic).

Iodide: [38763-55-2]

C₂₀H₁₆I₂NO₄ 461.255

Cryst. (EtOH). Mp 300° Mp 297-299°

(synthetic).

Picrate: [38763-56-3]

Cryst. (MeOH). Mp 233°.

Kitasato, Z. et al., *Yakugaku Zasshi*, 1927, **542**, 315 (isol)

Schramm, G. et al., *Pharmazie*, 1959, **14**, 405 (isol)

Govindachari, T.R. et al., *Indian J. Chem.*, 1971, **9**, 1313 (synth, struct)

Wuchuyine

W-23

[11034-73-4]

C₁₀H₁₃NO₂ 179.218

Struct. unknown. Alkaloid from the fruits of *Evodia rutaecarpa* (Rutaceae). Shows antiviral activity. Cryst. (Me₂CO). Mp 237.5°. $[\alpha]_D -68$.


Chen, A.L. et al., *J. Am. Pharm. Assoc.*, 1933, **22**, 716-719; *CA*, **27**, 5151 (isol)

Chou, S.C. et al., *Med. Pharmacol. Exp.*, 1967, **16**, 407-413; *CA*, **67**, 61622j

Wuchuyamide I

W-24

[263842-26-8]

C₁₉H₁₇N₃O₄ 351.361

Alkaloid from the fruit of *Evodia rutaecarpa*. Component of Wu Zhu Yu. Needles (CHCl₃/MeOH). Mp 261-262°. Racemic.

3-Deoxy: Wuchuyamide II

[263842-27-9]

C₁₉H₁₇N₃O₃ 335.362


Alkaloid from the fruit of *Evodia rutaecarpa*. Component of Wu Zhu Yu. Needles (CHCl₃/MeOH). Mp 199-200°. Racemic.

Zuo, G.Y. et al., *Chin. Chem. Lett.*, 2000, **11**, 127-128

Wuchuyamide III

W-25

[1016984-76-1]

C₁₈H₁₇N₃O₃ 323.351


Alkaloid from the fruit of *Evodia officinalis*. Cytotoxic. Prisms. Mp 175.6-176.6°. $[\alpha]_D^{25} -17$ (c, 0.6 in CHCl₃). Unaccounted for optical rotation. λ_{\max} 226 (log ϵ 3.97); 243 (log ϵ 4.33); 315 (log ϵ 3.38); 325 (log ϵ 3.39); 361 (log ϵ 3.4) (CHCl₃).

Jim, H.-Z. et al., *J. Asian Nat. Prod. Res.*, 2007, **9**, 685-688 (isol, pmr, cmr)

Wuzhuyurutine A

W-26

5,5a-Dihydro-6H-indolo[2',3':3,4]pyrrolo[2,1-b]-quinazoline-11,13-dione, 9CI [912537-94-1]

C₁₇H₁₁N₃O₂ 289.293


Alkaloid from the unripe fruit of *Evodia rutaecarpa*. Amorph. solid. λ_{\max} 223; 239; 264; 323 (MeOH).

Teng, J. et al., *Heterocycles*, 2006, **68**, 1691-1698 (isol, pmr, cmr, ms)

Wuzhuyurutine B

W-27

2-(1,4-Dihydro-4-oxo-2-quinazoliny)-1H-indole-3-carboxylic acid, 9CI [912537-95-2]

C₁₇H₁₁N₃O₃ 305.292


Alkaloid from the unripe fruit of *Evodia rutaecarpa*. Amorph. solid. λ_{\max} 220; 240; 347; 364 (MeOH).

Teng, J. et al., *Heterocycles*, 2006, **68**, 1691-1698 (isol, pmr, cmr, ms)

Wybutine

W-28

Y-base. Y-wye



(S)-form

C₁₃H₁₆N₆O₃ 304.308

1H-Form shown. Prob. in equilib. with 3H-form.

(S)-form [35693-91-5]

Base obt. from eukaryotic tRNAs. Mp 200-204° dec. $[\alpha]_D^{26} -40$ (c, 0.14 in MeOH).

3-β-D-Ribofuranosyl:

C₁₈H₂₄N₆O₇ 436.424

Nucleoside present in tRNAs. Foam. $[\alpha]_D^{23} -53.6$ (c, 0.344 in MeOH).

2''-Hydroxy:

C₁₃H₁₆N₆O₄ 320.307

Base from eukaryotic tRNAs.

(±)-form


Mp 214-215° dec. (204-206°).

Itaya, T. et al., *Tet. Lett.*, 1985, **26**, 347 (synth,

uv, pmr, abs config, bibl
 Itaya, T. *et al.*, *J.C.S. Perkin 1*, 1994, 2759
 (ribofuranosyl)

Wybutoxine**W-29**

Methyl 4,9-dihydro-β-hydroperoxy-α-[(methoxycarbonyl)amino]-4,6-dimethyl-9-oxo-1H-imidazo[1,2-a]purine-7-butanolate, 9CI. Peroxywybutine [36238-44-5]



$C_{16}H_{20}N_6O_7$ 408.37


Constit. of phenylalanine tRNA of beef, chicken, calf and rat livers. Also from *Geotrichum candidum* and *Lupinus luteus*.
 Fluorescent.

Feinberg, A.M. *et al.*, *J.A.C.S.*, 1974, **96**, 7797
 (*isol, struct*)

Mochizaki, A. *et al.*, *Bull. Chem. Soc. Jpn.*, 1980, **53**, 813 (*isol, pmr*)

Wye base**W-30**

1,4-Dihydro-4,6-dimethyl-9H-imidazo[1,2-a]purin-9-one, 9CI. 4,9-Dihydro-4,6-dimethyl-9-oxo-1H-imidazo[1,2-a]purine. Yt base [33359-03-4]



$C_9H_9N_5O$ 203.203

1*H*-Form shown in equilib. with 3*H*-form. Nucleotide present in t-RNA from *Torulopsis utilis* and other sources. The most highly modified of the minor nucleotides of t-RNA which are important for codon recognition. Cryst. (H_2O).

λ_{max} 227 (ϵ 37000); 231 (ϵ 37100); 255 (ϵ 5000); 284 (ϵ 9700) (H_2O) (pH 1).

3H-form

3-O-β-D-Ribofuranosyl: **Wyosine**. *Wyo.*

Yt

[52662-10-9]

 $C_{14}H_{17}N_5O_5$ 335.319Hypermodified nucleoside from *Torulopsis utilis* phenylalanine-tRNA.

Cryst. (H_2O). Mp 233° dec. $[\alpha]_D^{20}$ -57 (c, 0.08 in H_2O). Glycosidic bond extremely labile. λ_{max} 236 ; 295 (H_2O) (pH 4.2-12.1).

Kasai, H. *et al.*, *Biochemistry*, 1976, **15**, 898
 (*synth, struct, bibl*)

Nakatsuga, S. *et al.*, *Tet. Lett.*, 1978, 2579
 (*synth*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1985, **33**, 2339 (*synth, bibl*)

Bazin, H. *et al.*, *Tet. Lett.*, 1987, **28**, 275
 (*synth, uv, pmr, cmr*)

Glemarec, C. *et al.*, *Tetrahedron*, 1988, **44**, 1273 (*props, bibl*)

Nagamatsu, T. *et al.*, *Chem. Comm.*, 1995, 2041 (*synth*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 2002, **50**, 547-548 (*Wyosine, isol, struct*)

α-Xantherin


X-1

C₂₄H₂₃NO₆ 421.449
Struct. unknown. Alkaloid from *Zanthoxylum ochroxylum*. Mp 186-187°. Leprince, M. et al., *Bull. Sci. Pharmacol.*, 1912, **18**, 337-345; *CA*, **6**, 2287 (isol)

Xanthevodine

X-2

4,11-Dimethoxy-1,3-dioxolo[4,5-b]acridin-10(5H)-one, 9CI. 1,4-Dimethoxy-2,3-methylenedioxy-9(10H)-acridone. *Nor-melicopidine* [477-78-1]



C₁₆H₁₃NO₅ 299.282
Alkaloid from *Evodia xanthoxyloides*, *Acronychia baueri* and *Melicope leratii* (Rutaceae). Mp 217-218°. λ_{max} 250 (ε 22800); 284 (ε 28100); 312 (ε 9290) (EtOH/NaOH) (Derep). λ_{max} 218 (ε 15000); 251 (sh) (ε 23400); 282 (ε 33900); 304 (sh) (ε 10200) (EtOH) (Derep).

Picrate: Mp 194-196°.

N-Me: Melicopidine

[475-91-2]
C₁₇H₁₅NO₅ 313.309
Alkaloid from *Melicope fareana*, *Teclea natalensis*, *Acronychia baueri* and *Melicope leptococca* (Rutaceae). Mp 121-122°.


▶ J14630000

N-Me; hydrochloride: Mp 88-90°. Price, J.R. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 249 (isol)
Crow, W.D. et al., *Aust. J. Sci. Res., Ser. A*, 1949, **2**, 255; 282 (struct)
Hughes, G.K. et al., *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 401 (isol)
Cannon, J.R. et al., *Aust. J. Sci. Res., Ser. A*, 1952, **5**, 406 (struct)
Dallacker, F. et al., *Annalen*, 1966, **691**, 138 (synth)
Bowie, J. et al., *Aust. J. Chem.*, 1967, **20**, 1179 (ms)
Bert, M. et al., *Phytochemistry*, 1974, **13**, 301 (isol, pmr, ms)
Skaltsounis, A.L. et al., *J. Nat. Prod.*, 1983, **46**, 732 (isol)

Xanthiazone

X-3

4,8-Dihydro-7-(hydroxymethyl)-8,8-dimethyl-2H-1,4-benzothiazine-3,5-dione [212701-97-8]



C₁₁H₁₃NO₃S 239.295
Alkaloid from the fruit of *Xanthium strumarium*. Cubic cryst. (MeOH). Mp 159-161°.

11-O-β-D-Glucopyranoside: Xanthiside

C₁₇H₂₃NO₈S 401.437
Alkaloid from the fruits of *Xanthium pungens* and *Xanthium strumarium*. Amorph. powder; oil (as tetra-Ac). Mp 180-182°. [α]_D²⁵ -3.2 (c, 0.67 in CHCl₃) (tetra-Ac). λ_{max} 280 (MeOH).

11-O-[3,4-Dihydroxy-E-cinnamoyl-(→2)-β-D-glucopyranoside]:

C₂₆H₂₉NO₁₁S 563.581
Alkaloid from the fruit of *Xanthium strumarium*. Yellow cryst. Mp 186-188°. λ_{max} 248 ; 332 (MeOH).

2ξ-Hydroxy: 4,8-Dihydro-2-hydroxy-7-(hydroxymethyl)-8,8-dimethyl-2H-1,4-benzothiazine-3,5-dione. 2-Hydroxyxanthiazone

C₁₁H₁₃NO₄S 255.294

2ξ-Hydroxy, 11-O-β-D-glucopyranoside:

C₁₇H₂₃NO₉S 417.436
Constit. of the fruit of *Xanthium strumarium*. Amorph. powder. Mp 190-192°. λ_{max} 280 (MeOH).

Ma, Y.-T. et al., *Phytochemistry*, 1998, **48**, 1083-1085 (isol, pmr, cmr, cryst struct)
Mahmoud, A.A. et al., *Nat. Prod. Res.*, 2005, **19**, 585-589 (*Xanthiside*)
Han, T. et al., *Chem. Nat. Compd. (Engl. Transl.)*, 2006, **42**, 567-570 (glucosides)
Qin, L. et al., *Fitoterapia*, 2006, **77**, 245-246 (*caffeoylglucoside*)
Dai, Y.-H. et al., *J. Asian Nat. Prod. Res.*, 2008, **10**, 303-305 (isol, pmr, cmr)

Xanthicin


X-4

C₁₃H₁₅NO₅ 265.265
Phenoxazine deriv. Struct. unknown. Prod. by *Streptomyces xanthochromogenes* ATCC 19818. Antifungal agent. Yellow needles (EtOAc). Sol. Me₂CO, EtOAc, MeOH. Mp 211-213° dec. [α]_D²⁵ +319 (Me₂CO). λ_{max} 270 (ε 35230) (MeOH) (Berdy). λ_{max} 260 (ε 33930); 325 (ε 23300) (NaOH) (Berdy).
▶ LD₅₀ (mus, ipr) 105 mg/kg.
Arishima, M. et al., *CA*, 1958, **52**, 17388ab

Xanthine, 8CI

X-5

3,7-Dihydro-1H-purine-2,6-dione, 9CI. 2,6-Dihydroxypurine. 2,6-Purinediol [69-89-6]



C₅H₄N₄O₂ 152.112
Exists mainly as 7H-dioxo-form in aq. solns. with small amount of 9H-dioxo-form. Found in potatoes, coffee beans, etc. Cryst. Mp 350°. pK_{a1} 7.7; pK_{a2} 11.94 (20°). pK_{a1} 0.8; pK_{a2} 7.45; pK_{a3} 11.13 (20°).
▶ LD₅₀ (mus, ipr) 500 mg/kg. ZD7700000
1-Me: 3,7-Dihydro-1-methyl-1H-purine-2,6-dione. 1-Methylxanthine [6136-37-4] [28109-92-4]
C₆H₆N₄O₂ 166.139

Cryst. (H₂O). Mp 350°.
▶ LD₅₀ (mus, ipr) 510 mg/kg. ZD8575000
3-Me: 3,7-Dihydro-3-methyl-1H-purine-2,6-dione. 3-Methylxanthine [1076-22-8] [28109-92-4]
C₆H₆N₄O₂ 166.139
Found in human urine. Isol. from the ascidian *Sympyegma rubra*. Needles (H₂O). Mp 360° dec. λ_{max} 204 (log ε 3.7); 272 (log ε 3.3) (MeOH).
▶ LD₅₀ (mus, ipr) 894 mg/kg. ZD8750000
1,3-Di-Me: see Theophylline, T-360
1-Et: 1-Ethyl-3,7-dihydro-1H-purine-2,6-dione. 1-Ethylxanthine [104285-81-6]
C₇H₈N₄O₂ 180.166
Cryst. (MeOH). Mp 300°.
3-Propyl: 3,7-Dihydro-3-propyl-1H-purine-2,6-dione, 9CI. 3-Propylxanthine. *Enprofylline*, INN, USAN. Nilyph. *Oxeze*. D 4028 [41078-02-8]
C₈H₁₀N₄O₂ 194.193
Diuretic, antiasthmatic agent, bronchodilator. Lacks adenosine receptor antagonism of Xanthine. Mp 287-289°. Log P 0.36 (calc).
▶ LD₅₀ (rat, orl) 481 mg/kg. UO8439700
3-(2-Methylpropyl), 1-Me: 3,7-Dihydro-1-methyl-3-(2-methylpropyl)-1H-purine-2,6-dione. 3-Isobutyl-1-methylxanthine [28822-58-4]
C₁₀H₁₄N₄O₂ 222.246
Inhibitor of cyclic AMP phosphodiesterase. Mp 199-201°.

7H-form

7-(2-Deoxy-β-D-ribofuranosyl): N⁷-2'-Deoxy pseudoxanthosine [35912-09-5]
C₁₀H₁₂N₄O₅ 268.229
Constit. of the starfish *Asterias roll-estoni*. Mp 176-178°.
7-Me: 3,7-Dihydro-7-methyl-1H-purine-2,6-dione, 9CI. 7-Methylxanthine. *Heteroxanthine* [552-62-5] [28109-92-4]
C₆H₆N₄O₂ 166.139
Found in urine, sugar cane and other biol. sources, a urinary metab. of caffeine in man. Needles (H₂O). Mp 380° dec.
▶ ZD8925000
1,7-Di-Me: 3,7-Dihydro-1,7-dimethyl-1H-purine-2,6-dione, 9CI. 1,7-Dimethylxanthine. *Paraxanthine* [611-59-6]
C₇H₈N₄O₂ 180.166
Present in human urine. Plates (H₂O). Mp 298-299°. pK_{a1} 8.6.
▶ RV9380000
3,7-Di-Me: see Theobromine, T-355
1,3,7-Tri-Me: see Caffeine, C-15
1,3-Dipropyl, 7-Me: 3,7-Dihydro-7-methyl-1,3-dipropyl-1H-purine-2,6-dione, 9CI [31542-63-9]
C₁₂H₁₈N₄O₂ 250.3
Adenoside A₂-antagonist. Depressant.

Cryst. (Me₂CO/DMF). Mp 116°.

1,3-Bis(3-methyl-2-butenyl), 7-Me: 7-Methyl-1,3-diprenylxanthine

C₁₆H₂₂N₄O₂ 302.375

Alkaloid from leaves of *Bosistoa floydii* (Rutaceae). Powder.

1,7-Dibenzyl: [140396-45-8]

C₁₉H₁₆N₄O₂ 332.361

Cryst. (MeOH). Mp 223.5-225°.

9H-form [51953-26-5]

9-Me: 3,9-Dihydro-9-methyl-1H-purine-2,6-dione. 9-Methylxanthine

[1198-33-0]

C₆H₆N₄O₂ 166.139

Needles (H₂O). Mp 384° dec.

1,9-Di-Me: 1,9-Dimethyl-3,9-dihydro-1H-purine-2,6-dione, 9CI. 1,9-Dimethylxanthine, 8CI. Isoparaxanthine

[33073-01-7]

C₇H₈N₄O₂ 180.166

Cryst. (HCl aq.). Mp 35°. pK_{a1} 5.99

(20°).

3,9-Di-Me: 3,9-Dimethyl-3,9-dihydro-1H-purine-2,6-dione, 9CI. 3,9-Dimethylxanthine, 8CI

[15837-08-8]

C₇H₈N₄O₂ 180.166

Cryst. (H₂O). Mp 321-322°. pK_{a1} 10.14 (20°).

(7H,9H)-form

Imidazolium betaine-form

Minor tautomer.

7,9-Di-Me: 2,3,6,7-Tetrahydro-7,9-dimethyl-2,6-dioxo-1H-purinium hydroxide inner salt, 9CI

[5752-21-6]

C₇H₈N₄O₂ 180.166

Cryst. (MeOH aq.). Mp 378-379° dec.

Aldrich Library of FT-IR Spectra, 1st edn., 1985, 2, 709C (ir)

Aldrich Library of 13C and 1H FT NMR Spectra, 1992, 3, 213C; 214A (nmr)

Traube, W. et al., *Ber.*, 1913, **46**, 3839-3852 (1,7-di-Me)

Blitz, H. et al., *Annalen*, 1921, **423**, 200-226 (1,9-di-Me)

Sarasin, E. et al., *Helv. Chim. Acta*, 1924, **7**, 713-719 (7-Me)

Cook, A.H. et al., *J.C.S.*, 1949, 1069-1071; 1950, 1884-1888 (1,7-di-Me, 1,9-di-Me)

U.S. Pat., 1952, 2602795; CA, **47**, 4920g (3-(2-methylpropyl)-1-Me)

Blicke, F.F. et al., *J.A.C.S.*, 1954, **76**, 3653-3655 (1,7-di-Me)

Koppel, H.C. et al., *J.A.C.S.*, 1958, **80**, 2751-2755 (9-Me)

Pfleiderer, W. et al., *Annalen*, 1960, **631**, 168-174; 1961, **647**, 155-160 (9-Me, 1,9-di-Me, 3,9-di-Me)

Elion, G.B. et al., *J.O.C.*, 1962, **27**, 2478-2491 (3-Me)

Birkofer, L. et al., *Chem. Ber.*, 1964, **97**, 934-945 (7-Me)

Mizuno, H. et al., *Bull. Chem. Soc. Jpn.*, 1969, **42**, 3099-3105 (cryst struct)

Kwiatkowski, J.S. et al., *Theor. Chim. Acta*, 1969, **13**, 149-154 (uv)

Montague, W. et al., *Biochem. J.*, 1971, **122**, 115-120 (3-(2-methylpropyl)-1-Me, pharmacol)

Lister, J.H. et al., *Chem. Heterocycl. Compd.*, (Weissberger, A. et al., Ed.), 1971, (rev)

Bergmann, F. et al., *J.C.S.(C)*, 1971, 1676-1682; 1939-1941 (1,9-di-Me, 3,9-di-Me, pmr)

Lichtenberg, D. et al., *J.C.S.(C)*, 1971, 1676-1682 (tautom)

Ohtsuka, Y. et al., *Bull. Chem. Soc. Jpn.*, 1973, **46**, 506-509 (Enprofylline, synth)

Twanmoh, L.-M. et al., *J. Het. Chem.*, 1973, **10**, 187-190 (pmr)

Takayama, S. et al., *Chem. Pharm. Bull.*, 1974, **22**, 1200-1202 (3-Me, synth)

Kistenmacher, T.J. et al., *Acta Cryst. B*, 1975, **31**, 489-492 (7-Me)

Marzilli, L.G. et al., *J.A.C.S.*, 1975, **97**, 3351-3358 (7-Me)

Tanabe, T. et al., *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3224-3226 (3-Me)

Sekiya, M. et al., *Chem. Pharm. Bull.*, 1976, **24**, 1331-1335 (synth)

Coxon, B. et al., *J.O.C.*, 1977, **42**, 3132-3140 (cmr)

Saha, S.K. et al., *Indian J. Chem., Sect. B*, 1980, **19**, 325-327 (ms)

Stalhandske, C. et al., *Acta Cryst. C*, 1985, **41**, 1517-1520 (Enprofylline, cryst struct)

Andersson, K.E. et al., *Curr. Clin. Pract. Ser.*, 1985, **19**, 41 (Enprofylline, rev)

Daly, J.W. et al., *J. Med. Chem.*, 1986, **29**, 1305-1308; 1989, **32**, 1231-1237 (1,3-dipropyl-7-Me, synth)

Evoniuk, G. et al., *J. Pharmacol. Exp. Ther.*, 1987, **242**, 882-887 (1,3-dipropyl-7-Me, pharmacol)

Itaya, T. et al., *Chem. Pharm. Bull.*, 1989, **37**, 1235-1238 (3,9-di-Me)

Black, T.H. et al., *Synth. Commun.*, 1989, **19**, 843-850 (3,7-di-Me)

Fujii, T. et al., *Chem. Pharm. Bull.*, 1991, **39**, 2855-2862 (1,7-di-Me)

Chapman, K.R. et al., *Chest*, 1994, **106**, 1407 (Enprofylline, clin trial, rev)

Auzi, A.A. et al., *Phytochemistry*, 1994, **36**, 535-536 (7-Methyl-1,3-diprenylxanthine)

Sponer, J. et al., *Struct. Chem.*, 1995, **6**, 281-286 (tautom)

Mueller, C.E. et al., *Synthesis*, 1998, 1428-1436 (Paraxanthine, Isoparaxanthine, synth)

Lindsay, B.S. et al., *J. Nat. Prod.*, 1999, **62**, 1573-1575 (3-Me, isol, pmr, cmr, ms)

Martindale, *The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 1645

Civcir, P.U. et al., *THEOCHEM*, 2001, **545**, 7-15 (tautom)

Burnett, F.N. et al., *Tetrahedron*, 2002, **58**, 9567-9578 (1,7-dibenzyl)


Li, G.-Q. et al., *J. Chin. Pharm. Sci.*, 2004, **13**, 81-86 (2'-Deoxyseudoxanthosine)

Goswami, S. et al., *Eur. J. Org. Chem.*, 2007, 4056-4064 (synth, ir, pmr)

Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, PAK300; XCA000

Xanthobaccin A

[227596-81-8]



C₂₉H₃₈N₂O₆ 510.629

Macrolactam-tetramic acid antibiotic similar to Ikarugamycin. Enolised β-trioxo compd. Prod. from *Stenotrophomonas* sp. SB-K88. Antifungal agent. Amorph.

Relative Configuration

powder (as O-22 Na salt). Xanthobaccin B and C have also been isol. but not characterised. λ_{max} 239 (ε 18900); 321 (ε 9800) (MeOH) (Na salt).

Stereoisomer (?): **Maltophilin**. A 90931B. Antibiotic A 90931B

[184877-66-5]

C₂₉H₃₈N₂O₆ 510.629

Prod. by *Stenotrophomonas maltophilia* (formerly *Pseudomonas maltophilia*) and a *Streptomyces* sp. Antifungal agent. λ_{max} 320 (MeOH).

Stereoisomer, 14-alcohol: **Dihydromaltophilin**. Antibiotic A 90931A. Antibiotic TAN 883B. A 90931A. TAN 883B

[138634-40-9]

C₂₉H₄₀N₂O₆ 512.645

Prod. by *Streptomyces* sp. Antifungal agent, angiogenesis inhibitor, antiulcer agent, blood vessel growth promotor. Solid. Sol. MeOH. [α]_D +161 (DMF). Identities of Dihydromaltophilin/A 90931A and TAN 883B not certain. λ_{max} 233 (E1%/1cm 410); 318 (E1%/1cm 240) (MeOH) (Berdy). λ_{max} 218 (E1%/1cm 485); 320 (E1%/1cm 340) (MeOH/HCl) (Berdy). λ_{max} 233 (E1%/1cm 445); 315 (E1%/1cm 240) (MeOH/NaOH) (Berdy).

[227596-82-9]

Japan. Pat., 1991, 03 178 988; CA, **116**, 57524 (TAN 883B)

Jakobi, M. et al., *J. Antibiot.*, 1996, **49**, 1101-1104 (Maltophilin)

Graupner, P.R. et al., *J. Antibiot.*, 1997, **50**, 1014-1019 (Dihydromaltophilin)

Nakayama, T. et al., *Appl. Environ. Microbiol.*, 1999, **65**, 4334-4339 (isol, activity)

Hashidoko, Y. et al., *CA*, 1999, **131**, 184771e (struct)

Hashidoko, Y. et al., *Tet. Lett.*, 1999, **40**, 2957-2960 (isol, uv, pmr, cmr)


Xanthocillin X

X-7

4,4'-(2,3-Diisocyano-1,3-butadiene-1,4-diyl)bisphenol, 9CI. Xanthocillin, BAN.

Xantocillin, INN. Brevicide. Ophthocillin. Trianthil. Xantyril

[580-74-5]



C₁₈H₁₂N₂O₂ 288.305

Metab. of *Penicillium notatum* and *Eu-penicillium egyptiacum*. Cytotoxic. Antibacterial formerly used in feed supplements. Yellow cryst. Sol. MeOH, Et₂O, bases; poorly sol. H₂O, CHCl₃, hexane, C₆H₆. Mp 200° dec. λ_{max} 240 (ε 14000); 295 (sh) (ε 14000); 368 (ε 40000); 380 (sh) (ε 35000) (2-propanol) (Derep). λ_{max} 243 (ε 12020); 374 (ε 39800) (EtOH) (Berdy). λ_{max} 252 (ε 12300); 398 (ε 9550); 430 (ε 38900) (EtOH/NaOH) (Berdy).

▶ LD₅₀ (mus, ivn) 20 mg/kg; LD₅₀ (mus, ipr) 35 mg/kg; LD₅₀ (mus, orl) 40 mg/kg. KX1625000

Mono-Me ether: 4-[2,3-Diisocyano-4-(4-methoxyphenyl)-1,3-butadienyl]phenol, 9CI

[19559-24-1]

C₁₉H₁₄N₂O₂ 302.332

Prod. by *Basipetospora* sp., *Dichotomomyces albus* and *Dichotomomyces ceipii*. Potent inhibitor of prostaglandin biosynth. Active against some gram-positive and -negative bacteria. Yellow cryst. (C₆H₆). Sol. Me₂CO, CHCl₃, Et₂O, EtOAc; fairly sol. C₆H₆, EtOH; poorly sol. MeOH, H₂O, hexane. λ_{max} 236 (ε 14200); 295 (ε 17200); 360 (ε 30000) (MeOH) (Derep). λ_{max} 243 (ε 10960); 303 (ε 10000); 368 (ε 16600); 385 (ε 14120) (MeOH) (Berdy).

▶ LD₅₀ (mus, ipr) 40 mg/kg. KX1700000**4-Me ether, 4'-sulfate: Antibiotic BU****4704. BU 4704**

[149997-61-5]

C₁₉H₁₄N₂O₅S 382.396

Prod. by an *Aspergillus* sp. Off-white needles (as K salt). Sol. DMSO; fairly sol. MeOH, EtOAc. Mp 165-170° dec. (as K salt). λ_{max} 232 (ε 17200); 283 (sh) (ε 11000); 356 (ε 61900) (MeOH) (Derep). λ_{max} 356 (ε 61900) (MeOH/HCl) (Berdy). λ_{max} 283 (ε 11000) (MeOH/NaOH) (Berdy).

Di-Me ether: Xanthocillin X dimethyl ether

[4464-33-9]

C₂₀H₁₆N₂O₂ 316.359

Metab. of *Aspergillus clavatus* and a *Basipetospora* sp. Yellow needles (CH₂Cl₂/hexane). Sol. Me₂CO, EtOAc, acids, Et₂O; poorly sol. H₂O, hexane. Mp 186° dec. λ_{max} 230 (ε 2000); 296 (ε 1600); 363 (ε 2800); 380 (sh) (ε 2000) (MeOH) (Derep). λ_{max} 230 (ε 11200); 296 (ε 10700); 363 (ε 16600); 380 (ε 13800) (MeOH) (Berdy). λ_{max} 238 (ε 13600); 297 (ε 12200); 364 (ε 46400) (EtOH) (Berdy).

▶ KX1650000

1'',2''-Dihydro, di-Me ether: Antibiotic NK 372135A. NK 372135AC₂₀H₁₈N₂O₂ 318.374

Prod. by *Neosartorya fischeri*. Antifungal agent. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. λ_{max} 225; 297 (MeOH) (Berdy).

3-Hydroxy: Xanthocillin Y₁

[38965-69-4]

C₁₈H₁₂N₂O₃ 304.304

From *Penicillium notatum*. Yellow needles or rhombs. (EtOAc). Sol. MeOH, Et₂O, EtOH; poorly sol. H₂O, C₆H₆, hexane, CHCl₃. λ_{max} 250 (E1%/1cm 520); 375 (E1%/1cm 1650) (MeOH) (Berdy). λ_{max} 243; 379 (EtOH) (Berdy).

▶ LD₅₀ (mus, ivn) 14 mg/kg; LD₅₀ (mus, ipr) 15 mg/kg; LD₅₀ (mus, orl) 100 mg/kg. CZ8985000**3'-Hydroxy, 1'',2''-dihydro, di-Me ether: Antibiotic NK 372135C. NK 372135C**C₂₀H₁₈N₂O₃ 334.374

Prod. by *Neosartorya fischeri*. Antifungal agent. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. λ_{max} 238; 291; 318 (MeOH) (Berdy). λ_{max} 224; 285; 290; 318 (MeOH) (Berdy).

3,3'-Dihydroxy: Xanthocillin Y₂

[38965-70-7]

C₁₈H₁₂N₂O₄ 320.304

From *Penicillium notatum*. Sol. EtOH, Et₂O, MeOH; poorly sol. H₂O, hexane, CHCl₃, C₆H₆. λ_{max} 260 (E1%/1cm 320); 382 (E1%/1cm 1500) (MeOH) (Berdy).

▶ LD₅₀ (mus, ivn) 18 mg/kg; LD₅₀ (mus, ipr) 20.5 mg/kg; LD₅₀ (mus, orl) 100 mg/kg. CZ8937750**3-Methoxy, di-Me ether: Methoxy-xanthocillin X dimethyl ether**

[19764-51-3]

C₂₁H₁₈N₂O₃ 346.385

Isol. from a *Aspergillus* sp. and a *Basipetospora* sp. Sol. Me₂CO, EtOAc, Et₂O, CHCl₃; poorly sol. H₂O, hexane. λ_{max} 236 (ε 14200); 295 (ε 17200); 360 (ε 30000) (MeOH) (Derep). λ_{max} 238 (ε 13600); 297 (ε 12200); 364 (ε 46400) (EtOH) (Berdy).

3'-Methoxy, 1'',2''-dihydro, di-Me ether: Antibiotic NK 372135B. NK 372135BC₂₁H₂₀N₂O₃ 348.401

Prod. by *Neosartorya fischeri*. Antifungal agent. Sol. MeOH, EtOAc, CHCl₃; poorly sol. H₂O. [α]_D²⁰ +188.7 (c, 1 in MeOH). λ_{max} 226; 288; 315 (MeOH) (Berdy).

3,5-Dimethoxy, di-Me ether: [663918-81-8]C₂₂H₂₀N₂O₄ 376.411

Isol. from the marine fungus *Basipetospora* sp. Yellow solid.

[149026-37-9]

Hagedorn, I. et al., *Angew. Chem., Int. Ed.*, 1962, **1**, 212 (*synth*)Ando, K. et al., *J. Antibiot.*, 1968, **21**, 582; 587 Takatsuki, A. et al., *J. Antibiot.*, 1968, **21**, 671; 676 (*Methoxyxanthocillin*)Zarnack, J. et al., *Pharmazie*, 1971, **26**, 503 (*isol*)Achenbach, H. et al., *Chem. Ber.*, 1972, **105**,784; 3061 (*biosynth, derivs*) Takahashi, C. et al., *Chem. Pharm. Bull.*, 1976, **24**, 2317 (*pmr, derivs*)Büchi, G. et al., *J.O.C.*, 1977, **42**, 244 (*isol, uv, ir, pmr, deriv*)Vesonder, R.F. et al., *J. Nat. Prod.*, 1979, **42**, 232 (*isol*)Britton, D. et al., *Cryst. Struct. Commun.*, 1981, **10**, 1497 (*cryst struct*)Kitahara, N. et al., *J. Antibiot.*, 1981, **34**, 1556Herbert, R.B. et al., *Tet. Lett.*, 1984, **25**, 4263Cable, K.M. et al., *J.C.S. Perkin 1*, 1991, 595 (*biosynth*)Tsunakawa, M. et al., *J. Antibiot.*, 1993, **46**, 687 (*BU 4704*)Morino, T. et al., *J. Antibiot.*, 1994, **47**, 1546 (*NK 372135*)Sakai, R. et al., *Bioorg. Med. Chem.*, 2005, **13**, 6388-6393 (*3,5-dimethoxy di-Me ether*)Tatsuta, K. et al., *Tet. Lett.*, 2005, **46**, 5017-5020 (*di-Me ether, synth*)Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 8th edn., Van Nostrand Reinhold, 1992, DVU000; XCS680; XCS700,**C-Xanthocurine****X-8**C₂₀H₂₁N₂O⁺ 305.399


Struct. unknown. Alkaloid from calabash curare (*Strychnos* spp.) (Loganiaceae). Mp 275° (as picrate). [α]_D +813 (MeOH) (as chloride). Blue-green col. with Ce(SO₄)₂.

Giesbrecht, E. et al., *Helv. Chim. Acta*, 1954, **37**, 1974-1982 (*isol, uv*)**Xanthofagarine****X-9**C₁₈H₂₂NO₈ 380.374

Struct. unknown. Alkaloid from *Fagara microphylla* (Rutaceae). Pale-yellow needles (MeOH). Mp 270-272° dec. (block). Prob. opt. inactive.

Hydrochloride: Mp 245°.*Picrate*: Mp 270°.Paris, M.R. et al., *Ann. Pharm. Fr.*, 1951, **9**, 479-493 (*isol, uv*)**Xantholipin****X-10**


[405214-19-9]

C₂₇H₁₈ClNO₁₀

Prod. by a *Streptomyces* sp. Inhibitor of HSP47 gene expression. Yellow powder. [α]_D -303.8 (c, 0.53 in dioxan). λ_{max} 244 (log ε 4.54); 271 (log ε 4.47); 312 (log ε 4.12); 394 (log ε 4.03) (MeOH).

Terui, Y. et al., *Tet. Lett.*, 2003, **44**, 5427-5430 (*isol, uv, pmr, cmr, abs config*)**Xanthommatin****X-11**

α-Amino-3-carboxy-1-hydroxy-γ,5-dioxo-5H-pyrido[3,2-a]phenoxazine-11-butanoic acid, 9CI. Urechochrome [521-58-4]


C₂₀H₁₃N₃O₈ 423.338**(S)-form**

Insect metab. found in secretions of *Vanessa urticae* and in eyes of grasshoppers and blowflies (*Calliphora erythrocephala*). Also obt. from eggs of the marine worm *Urechis caupo*. Yellow-brown powder.

Dihydro: see Dihydroxanthommatin, D-527Butenandt, A. et al., *Annalen*, 1954, **586**, 217-228; **588**, 106-116; **590**, 75-90 (*isol, struct, synth*)Linzen, B. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1959, **314**, 12-14 (*isol*)Butenandt, A. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1960, **321**, 258-275Linzen, B. et al., *Adv. Insect Physiol.*, 1974, **10**, 117-246 (*biosynth*)Kayser, H. et al., *Compr. Insect Physiol.**Biochem. Pharmacol.*, 1985, **10**, 367-415 (*rev*)Martel, R.R. et al., *J. Biol. Chem.*, 1991, **266**, 21392-21398 (*bibl*)

Xanthopterincarboxylic acid X-12

2-Amino-1,4,5,6-tetrahydro-4,6-dioxo-7-pteridincarboxylic acid, 9CI. 2-Amino-4,6-dihydroxy-7-pteridincarboxylic acid, 8CI
[2757-91-7]



$C_7H_5N_5O_4$ 223.148

Coenzyme isol. from L-glutamate oxidase of wheat seeds. Orange cryst. + $1H_2O$. Mp $>340^\circ$.


Tschesche, R. *et al.*, *Chem. Ber.*, 1955, **88**, 81-90 (*synth*)

Pfleiderer, W. *et al.*, *Chem. Ber.*, 1957, **90**, 2624-2631 (*synth*)

Gil'manov, M.K. *et al.*, *Dokl. Akad. Nauk SSSR, Ser. Khim.*, 1989, **309**, 484-487 (*isol*)

Xanthosine, 8CI X-13

3,9-Dihydro-9- β -D-ribofuranosyl-1H-purine-2,6-dione, 9CI. 9- β -D-Ribofuranosyl-xanthine
[146-80-5]



$C_{10}H_{12}N_4O_6$ 284.228

Prod. by guanine-free mutants of bacteria e.g. *Bacillus subtilis*, *Aerobacter aerogenes*. Also reported from seeds of *Trifolium alexandrinum*. Prismatic cryst. (H_2O). Spar. sol. cold H_2O ; v. sol. hot H_2O . $[\alpha]_D^{30}$ -51.2 (c, 8.0 in 0.3M NaOH). pK_{a1} 5.67; pK_{a3} 12.85 (25°). Dec. on heating with no distinct melting range. λ_{max} 253 (ϵ 8790) (H_2O).

3'-Phosphate: 3'-Xanthylic acid, 9CI, 8CI
[21089-32-7]

$C_{10}H_{13}N_4O_9P$ 364.208

Mp 200° (as brucine salt). $[\alpha]_D^{20}$ -61.66 (c, 5.0 in NaOH).

5'-Phosphate: Xanthosine 5'-(dihydrogen phosphate), 9CI. 5'-Xanthylic acid, 8CI. Xanthosine monophosphate. Xanthosine 5'-phosphate. XMP

[523-98-8]

[25899-70-1]

$C_{10}H_{13}N_4O_9P$ 364.208

Inhibitor of isoforms I and II of human inosine 5'-monophosphate dehydrogenase. No phys. props.

reported.

5'-Triphosphate: [6253-56-1]

[50801-64-4, 90011-94-2]

$C_{10}H_{15}N_4O_{15}P_3$ 524.168

No phys. props. reported.

5'-O-Sulfate: Xanthosine 5'-sulfate

$C_{10}H_{12}N_4O_9S$ 364.292

Isol. from the venom of the spider *Tegenaria agrestis*.

2',3'-O-Isopropylidene: [4137-57-9]

$C_{13}H_{16}N_4O_6$ 324.293

Mp 240° .

3-Me: 3-Methylxanthosine

[3080-28-2]

$C_{11}H_{14}N_4O_6$ 298.255

Needles (H_2O). Mp 200° dec.

[5968-90-1]

Magasanik, B. *et al.*, *J. Biol. Chem.*, 1954, **206**, 83 (*isol*)

Bolis, M.E. *et al.*, *J. Biol. Chem.*, 1956, **219**, 917 (*biosynth*)

Jardetzky, C.D. *et al.*, *J.A.C.S.*, 1960, **82**, 222 (*pmr*)

Krishnaswamy, N.P. *et al.*, *Curr. Sci.*, 1966, **35**, 11 (*isol*)

Yoshikawa, M. *et al.*, *Bull. Chem. Soc. Jpn.*,

1967, **40**, 2849-2853 (5'-phosphate, *synth*)

Yamazaki, A. *et al.*, *J.O.C.*, 1967, **32**, 3258 (*synth*)

Holy, A. *et al.*, *Coll. Czech. Chem. Comm.*, 1968, **33**, 2259 (3'-Xanthylic acid)

Jones, A.J. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1970, **65**, 27 (*cmr*)

De Pamphilis, M.L. *et al.*, *Biochemistry*, 1973, **12**, 3714-3724 (*synth*, *uv*, 5'-triphosphate)

Sowa, T. *et al.*, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 2084-2090 (*phosphate*)

Westhof, E. *et al.*, *Z. Naturforsch., C*, 1975, **30**, 131 (*pmr*)

Koyama, G. *et al.*, *Acta Cryst. B*, 1976, **32**, 969 (*cryst struct*)

Itaya, T. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1235 (3-Methylxanthosine)


Ortiz, B. *et al.*, *Eur. J. Biochem.*, 1993, **212**, 263-270 (*pmr*, 5'-triphosphate)

Carr, S.F. *et al.*, *J. Biol. Chem.*, 1993, **268**, 27286-28290 (5'-phosphate, *biol*)

Taggi, A.E. *et al.*, *J.A.C.S.*, 2004, **126**, 10364-10369 (5'-sulfate)

Xenocoumacin 1 X-14

[105688-01-5]



$C_{22}H_{35}N_5O_6$ 465.548

Metab. of *Xenorhabdus* bacteria.

Shows antiulcer, antiinflammatory and antifungal activity. Antiparasitic agent.

Sol. H_2O , MeOH. $[\alpha]_D^{23}$ -57 (c, 0.7 in H_2O). λ_{max} 248 (sh) (ϵ); 347 (ϵ)

(0.1N NaOH) (Derep). λ_{max} 246 (ϵ 5700); 314 (ϵ 4080) (MeOH) (Derep).

λ_{max} 249 (ϵ 5460); 314 (ϵ 3300) (H_2O) (Berdy).


[105762-00-3]

McInerney, B.V. *et al.*, *J. Nat. Prod.*, 1991, **54**,

785 (*isol*)

Xestamine A X-15

N-Methoxy-N-methyl-18-(3-pyridinyl)-11-octadecen-13-yn-1-amine, 9CI
[131985-15-4]



$C_{25}H_{40}N_2O$ 384.604

Isol. from the marine sponges *Xestospongia wiedenmayeri* and *Calyx podotyta*. Oil. λ_{max} 260 (ϵ 2500); 265 (ϵ 3100); 270 (ϵ 2000) (MeOH) (Derep). λ_{max} 227 (ϵ 12200); 257 (ϵ 2100); 263 (ϵ 2400); 269 (ϵ 1600) (MeOH) (Berdy).

N¹-Me: Xestamine F

[136945-78-3]

$C_{26}H_{43}N_2O^{\oplus}$ 399.638


Alkaloid from the sponge *Calyx podotyta*. Oil. No counterion specified, presumably chloride. λ_{max} 226; 257; 269 (MeOH) (Berdy).

Sakemi, S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 995 (*isol*, *struct*)

Stierle, D.B. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1134 (Xestamine F)

Xestamine B X-16

N-Methoxy-N-methyl-14-(3-pyridinyl)-9-tetradecen-1-amine, 9CI
[131985-16-5]




$C_{21}H_{36}N_2O$ 332.528

Alkaloid from the marine sponge *Xestospongia wiedenmayeri*. Oil. λ_{max} 260 (ϵ 2500); 265 (ϵ 3100); 270 (ϵ 2000) (MeOH) (Derep). λ_{max} 205 (ϵ 9700); 257 (ϵ 4300); 262 (ϵ 4900); 269 (ϵ 3700) (MeOH) (Berdy).

Sakemi, S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 995 (*isol*, *struct*)

Xestamine C X-17

N-Methoxy-N, ζ -dimethyl-3-pyridinetetradecanamine, 9CI
[131985-17-6]



$C_{22}H_{40}N_2O$ 348.571


Alkaloid from the marine sponge *Xestospongia wiedenmayeri*. Oil. $[\alpha]_D^{20}$ -12.9 (c, 0.34 in MeOH). λ_{max} 260 (ϵ 2500); 265 (ϵ 3100); 270 (ϵ 2000) (MeOH) (Derep). λ_{max} 205 (ϵ 5200); 256 (ϵ 2700); 262 (ϵ 3100); 269 (ϵ 2300) (MeOH) (Berdy).

Sakemi, S. *et al.*, *J. Nat. Prod.*, 1990, **53**, 995

(isol, struct)

Xestamine D X-18

N-Methoxy-*N*-methyl-3-pyridinetetradecanamine, 9CI
[136945-76-1]



C₂₁H₃₈N₂O 334.544

Alkaloid from the sponge *Calyx podatypa*. Obt. as a mixt. with Xestamine E. λ_{max} 226 (ε 8000); 257 (ε 2200); 262 (ε 2500); 269 (ε 2800) (MeOH) (Berdy).

*N*¹-Me: Xestamine G

[136945-79-4]

C₂₂H₄₁N₂O[⊕] 349.579


Alkaloid from the sponge *Calyx podatypa*. Obt. as a mixt. with Xestamine H. λ_{max} 226 ; 257 ; 269 (MeOH) (Berdy).

Stierle, D.B. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1134 (isol, pmr, cmr, struct)

Wang, Y. *et al.*, *J.O.C.*, 2003, **68**, 3090-3098 (synth)

Xestamine E X-19

N-Methoxy-*N*-methyl-3-pyridinepentadecanamine
[136945-77-2]



C₂₂H₄₀N₂O 348.571

Alkaloid from the sponge *Calyx podatypa*. Obt. as a mixt. with Xestamine D. λ_{max} 226 ; 257 ; 269 (MeOH) (Berdy).

*N*¹-Me: Xestamine H

[136945-80-7]


C₂₃H₄₃N₂O[⊕] 363.605

Alkaloid from the sponge *Calyx podatypa*. Obt. as a mixt. with Xestamine G. λ_{max} 226 ; 257 ; 269 (MeOH) (Berdy).

Stierle, D.B. *et al.*, *J. Nat. Prod.*, 1991, **54**, 1134 (isol, pmr, cmr, struct)

Xestoamine X-20

[145237-04-3]




C₁₄H₁₉N₃ 229.324

Alkaloid from the New Caledonian sponge *Xestospongia* sp. Amorph. Opt. inactive.

Quirion, J.-C. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1505 (isol, ir, pmr, cmr, ms, struct)

Xestocyclamine A

[151232-83-6]



C₂₆H₄₀N₂O 396.615


Alkaloid from the marine sponge *Xestospongia* sp. Protein kinase C inhibitor. [α]_D -13.5 (c, 0.019 in MeOH).

Rodríguez, J. *et al.*, *J.A.C.S.*, 1993, **115**, 10436 (isol, ir, pmr, cmr, ms)

Rodríguez, J. *et al.*, *Tet. Lett.*, 1994, **35**, 4719 (struct)

Xestocyclamine B

[157207-89-1]




C₂₈H₄₄N₂O 424.668

Alkaloid from the sponge *Xestospongia* sp. Protein kinase inhibitor.

Rodríguez, J. *et al.*, *Tet. Lett.*, 1994, **35**, 4719 (isol, pmr, cmr, struct)

Xestomanzamine A X-23

(1-Methyl-1*H*-imidazol-5-yl)-9*H*-pyrido[3,4-*b*]indol-1-ylmethanone, 9CI
[164301-25-1]



C₁₆H₁₂N₄O 276.297

Alkaloid from the Okinawan marine sponge *Xestospongia* sp. Yellow needles (CHCl₃/MeOH). Mp 185-186°. λ_{max} 221 (ε 7400); 257 (ε 1700); 300 (ε 3900); 395 (ε 1600) (MeOH) (Berdy).

N-De-Me: *N*-Demethylxestomanzamine A

[581782-69-6]

C₁₅H₁₀N₄O 262.27

Alkaloid from an Indonesian sponge. Yellow powder. Mp 192° dec. λ_{max} 218 ; 258 ; 298 ; 356 ; 395 (MeOH).

3,4-Dihydro: Xestomanzamine B

X-21

[164301-24-0]

C₁₆H₁₄N₄O 278.313

Alkaloid from *Xestospongia* sp. Exhibits weak cytotoxicity against KB cells. Yellow oil. λ_{max} 222 (ε 27200); 270 (ε 10300); 298 (ε 16500); 388 (ε 4500) (MeOH) (Berdy).

Kobayashi, M. *et al.*, *Tetrahedron*, 1995, **51**, 3727-3736 (isol, uv, ir, pmr, cmr, cryst struct, Xestomanzamine B)

Molina, P. *et al.*, *Tet. Lett.*, 1996, **37**, 9353-9356 (synth, Xestomanzamine B)

Panosyan, F.B. *et al.*, *Can. J. Chem.*, 2001, **79**, 1110-1114 (synth)

Burm, B.E.A. *et al.*, *Heterocycles*, 2001, **55**, 495-503 (synth, Xestomanzamine B)


Rao, K.V. *et al.*, *J. Nat. Prod.*, 2003, **66**, 823-828 (*N*-Demethylxestomanzamine A)

Xestospongine A

X-24

Araguspongine D

[88840-02-2]



Absolute Configuration

C₂₈H₅₀N₂O₂ 446.715

The abs. configs. of Xestospongins A and C and Araguspongine B were revised in 1998. The work suggested that the abs. configs. of the other nat. prods. in the series required revision. Araguspongine D was found to be partially racemic Xestospongine A. Isol. from the marine sponge *Xestospongia exigua* and from an Okinawan *Xestospongia* sp. Shows vasodilator props. Cryst. (Et₂O). Mp 135-136°. [α]_D +6.9 (c, 0.84 in CHCl₃) (+ 10). Related to Petrosine, P-290.

7*S*-Hydroxy: 7*S*-Hydroxyxestospingine A

C₂₈H₅₀N₂O₃ 462.715

Isol. from *Xestospongia* sp. Cryst. (Me₂CO). Mp 166-167°. [α]_D +4.7 (c, 0.25 in CHCl₃).

5,9-Diepimer: Xestospingine C. Araguspongine E

[88903-69-9]

C₂₈H₅₀N₂O₂ 446.715

Isol. from *Xestospongia exigua*. Calcium channel blocker. Inositol triphosphate Ins(1,4,5)P₃ receptor antagonist. Shows vasodilator props. Cryst. (Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 149-150°. [α]_D -2.4 (c, 0.54 in CHCl₃). Struct. of Araguspongine E revised in 1998.

5,9-Diepimer, 9-hydroxy: Xestospingine D. Araguspongine A

[88840-00-0]

$C_{28}H_{50}N_2O_3$ 462.715

Isol. from *Xestospongia exigua* and *Haliclona* sp. Also isol. as racemate from sponge *Niphates* sp. Shows vasodilator props, human cancer cell inhibitory and antimicrobial activity. Cryst. (Et₂O). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 156-157°. [α]_D +18.43 (c, 1.08 in CHCl₃).

5,9-Diepimer, 9-hydroxy, N³-oxide: Araguspangine K

$C_{28}H_{50}N_2O_4$ 478.714

Alkaloid from *Xestospongia exigua*. Needles. [α]_D²⁵ +2.6 (c, 0.35 in CHCl₃).

2,2',5,5'-Tetraepimer: Araguspangine M

$C_{28}H_{50}N_2O_2$ 446.715

Alkaloid from the sponge *Neopetrosia exigua*. [α]_D -4.6 (c, 0.5 in CHCl₃).

5,5',9,9'-Tetraepimer: Araguspangine B [123000-02-2]

$C_{28}H_{50}N_2O_2$ 446.715

Alkaloid from an Okinawan sponge *Xestospongia* sp. Shows vasodilator activity. Somatostatin inhibitor. Sol. MeOH, CHCl₃; poorly sol. H₂O. Isol. as racemate, both enantiomers characterized. Abs. config. indicated refers to (-)-form ([α]_D -15°). Abs. config. revised in 1998.

[123048-14-6, 123000-03-3, 123000-06-6, 123000-07-7]

Nakagawa, M. *et al.*, *Tet. Lett.*, 1984, **25**, 3227 (*ir, pmr, cmr, struct, cryst struct*)

Kobayashi, M. *et al.*, *Chem. Pharm. Bull.*, 1989, **37**, 1676 (*isol, ir, pmr, cmr, struct, abs config*)

Hoye, T.R. *et al.*, *J.A.C.S.*, 1994, **116**, 2617 (*synth*)

Pettit, G.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 1313 (*isol, cryst struct, rac-Xestospongins D*)

Gafni, J. *et al.*, *Neuron*, 1997, **19**, 723-733 (*Xestospongins C, pharmacol*)

Kobayashi, M. *et al.*, *Heterocycles*, 1998, **47**, 195-203 (*pmr, cmr, ms, struct*)

Schultz, A.G. *et al.*, *J.A.C.S.*, 1998, **120**, 8259-8260 (*synth, abs config*)

Matzanke, N. *et al.*, *Org. Prep. Proced. Int.*, 1998, **30**, 3-51 (*rev, synth*)


Moon, S.-S. *et al.*, *J. Nat. Prod.*, 2002, **65**, 249-254 (*Xestospongins D, abs config, 7-Hydroxyxestospongins A*)

Orabi, K.Y. *et al.*, *J. Nat. Prod.*, 2002, **65**, 1782-1785 (*Araguspangines A,K*)

Liu, H. *et al.*, *Mar. Drugs*, 2004, **2**, 154-163 (*Araguspangine M*)

Xestospongins B

[88840-01-1]



X-25

$C_{29}H_{52}N_2O_3$ 476.741

Isol. from the marine sponge *Xestospongia exigua*. Shows vasodilator props. Cryst. (Et₂O). Mp 179-181°. [α]_D +7.1 (c, 0.91 in CHCl₃).

Demethylxestospongins B

$C_{28}H_{50}N_2O_3$ 462.715


Alkaloid from a New Caledonian sponge *Xestospongia* sp. Amorph. [α]_D²⁰ +6 (c, 0.8 in CHCl₃).

Nakagawa, M. *et al.*, *Tet. Lett.*, 1984, **25**, 3227 (*ir, pmr, cmr, struct*)

Quirion, J.-C. *et al.*, *J. Nat. Prod.*, 1992, **55**, 1505 (*Demethylxestospongins B*)

Xylactam

X-26



$C_{23}H_{31}NO_6$ 417.501

Similar to Xylaral. Prod. by *Xylaria euglossa*. Powder. λ_{max} 242; 277; 325 (CHCl₃).

Wang, X.-N. *et al.*, *J. Antibiot.*, 2005, **58**, 268-270 (*isol, pmr, cmr, ms*)

Xylaramide†

X-27

2-Hydroxy-N-(2-phenylethenyl)propanamide. N-(2-Phenethenyl)-2-hydroxypropanamide. N-Styryllactamide

PhCH=CHNHCOCH(OH)CH₃

$C_{11}H_{13}NO_2$ 191.229

(±)-(E)-form


Isol. from culture fluid of the ascomycete *Xylaria longipes* A19-91. Oil. Not found in CA indexes. λ_{max} 279 (ε 10600) (MeOH).

Schneider, G. *et al.*, *Z. Naturforsch., C*, 1996, **51**, 802-806 (*isol, uv, ir, pmr, cmr, ms*)

Xyloallenolide A

X-28

[329901-31-7]



$C_{29}H_{40}N_4O_5$ 524.659


Isol. from *Xylaria* sp. No. 2508. Cryst. Mp 82-85°. [α]_D²⁵ -34.6 (c, 0.06 in CHCl₃). λ_{max} 217 (ε 38000); 291 (ε 29000) (CHCl₃).

Lin, Y. *et al.*, *Tet. Lett.*, 2001, **42**, 449-451 (*isol, ir, pmr, cmr, uv*)

Xylogranatinine

X-29

9-Hydroxy-3-methoxy-6H-pyrido[1,2-a]pyrazin-6-one
[1000974-03-7]



$C_9H_8N_2O_3$ 192.174


Alkaloid from the fruit of the mangrove *Xylocarpus granatum*. Powder. λ_{max} 204; 227; 296; 343 (MeCN).

Zhou, Y. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 426-428 (*isol, pmr, cmr*)

Xylopinine

X-30

5,8,13,13a-Tetrahydro-2,3,10,11-tetra-methoxy-6H-dibenzo[a,g]quinolizine, 9CI. Norcoralydine. O,O-Dimethylcoreximine. O,O-Dimethylgovanine
[4216-86-8]



(S)-form

$C_{21}H_{25}NO_4$ 355.433

(S)-form [523-02-4]

Alkaloid from *Xylophia discreta*, *Xylophia buxifolia*, *Stephania* spp., *Polyalthia* spp., *Guatteria* spp., *Duguetia* spp., first known as a semisynthetic product. Shows anti-tussive, central adrenergic and soporific activities. Inhibitor of atrial activity in guinea pig heart. Light yellow prismatic cryst. (CHCl₃/Me₂CO). Mp 182-183° (177°). [α]_D -297 (c, 1 in CHCl₃).

▶HQ1793000

Hydrochloride: Mp 216-219° dec.

Hydrobromide: [51873-48-4]

Needles (MeOH/Et₂O). Mp 236-237° (222-225° dec.). [α]_D -227 (c, 1.00 in EtOH).

N-Oxide (α-): cis-Xylopinine N-oxide
[109063-91-4]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from the roots of *Stephania suberosa*. [α]_D²⁵ -96 (c, 0.41 in MeOH).

N-Oxide (β-): trans-Xylopinine N-oxide
[109063-90-3]

$C_{21}H_{25}NO_5$ 371.432

Alkaloid from the roots of *Stephania suberosa*. [α]_D²⁵ -200 (c, 0.44 in MeOH).

O³,O¹⁰,O¹¹-Tri-de-Me: Artavenustine
[88668-27-3]

$C_{18}H_{19}NO_4$ 313.352

Minor alkaloid from the stem bark of *Artabotrys venustus*. Amorph. [α]_D -127 (c, 0.26 in EtOH).


O³,O¹⁰,O¹¹-Tri-de-Me, 11-O-β-D-gluco-

- pyranoside: **Dauricoside**. 11-O- β -D-Glucopyranosylartavenustine. Artavenustine 11-O- β -D-glucopyranoside [155416-25-4]
C₂₄H₂₉NO₉ 475.494
Alkaloid from rhizomes of *Menispermum dauricum*. Inhibits blood-platelet aggregation. Prisms + 1H₂O (MeOH) (as hydrochloride). Mp 216-217° (hydrochloride). [α]_D²² -185.5 (c, 0.5 in MeOH).
- O²,O¹⁰,O¹¹-Tri-de-Me: **Pessoine** [88668-29-5]
C₁₈H₁₉NO₄ 313.352
Alkaloid from trunk bark of *Ammona spinescens*. Amorph. [α]_D²⁰ -160 (c, 0.2 in MeOH).
- O¹⁰,O¹¹-Di-de-Me: **Spinosine** [175274-51-8]
C₁₉H₂₁NO₄ 327.379
Alkaloid from trunk bark of *Ammona spinescens*. Amorph. [α]_D²⁰ -35 (c, 0.3 in EtOH).
- O³,O¹¹-Di-de-Me: **Isocoreximine**
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Toddalia asiatica*. Light yellowish prisms (CHCl₃/MeOH). Mp 238-242°. [α]_D²⁵ -361 (c, 0.1 in MeOH). λ_{\max} 225 (log ϵ 3.18); 286 (log ϵ 2.93) (EtOH).
- O³,O¹⁰-Di-de-Me: **10-Demethylidiscetretine**. 11-O-Methylartavenustine [91200-36-1]
C₁₉H₂₁NO₄ 327.379
Alkaloid from the stem bark of *Guatteria discolor* and *Artabotrys venustus*. Noncryst. [α]_D -178 (c, 0.27 in MeOH). λ_{\max} 208 (log ϵ 4.49); 226 (sh) (log ϵ 4.06); 289 (log ϵ 3.71) (EtOH).
- O²,O¹⁰-Di-de-Me: **Govadine** [60342-43-0]
C₁₉H₂₁NO₄ 327.379
Alkaloid from the leaves and stems of *Corydalis govaniana*. Mp 238-240°. [α]_D -128 (Py).
- O²-De-Me: **Govanine**. *Corydalis L* [59444-66-5]
C₂₀H₂₃NO₄ 341.406
Alkaloid from leaves and stems of *Corydalis govaniana*. Cryst. (MeOH). Mp 168-170°. [α]_D²⁶ -319 (CHCl₃). Incorr. descr. as (R-).
- O³-De-Me: **Discretine** [6872-75-9]
C₂₀H₂₃NO₄ 341.406
Alkaloid from *Xylopiya discreta*, *Pachypodanthium staudtii*, *Guatteria discolor* and *Guatteria scandens*. Claimed to have analgesic, antiinflammatory, antipyretic and vasodepressor activity. Fine needles (Me₂CO/Et₂O). Mp 180-181° dec. [α]_D -300 (c, 1.44 in CHCl₃).
- O²,O¹¹-Di-de-Me, N-Me: **Phellodendrine** [6873-13-8]
C₂₀H₂₄NO₄⁺ 342.414
Quaternary alkaloid from *Phellodendron amurense*. Prisms (as iodide). Mp 258-258.5° dec. (iodide). [α]_D³⁰ -130.
- O²,O¹¹-Di-de-Me: **Coreximine**. *Coramine*. Alkaloid F29 [483-45-4]
C₁₉H₂₁NO₄ 327.379
Alkaloid from *Dicentra eximia*, *Papaver somniferum* (opium poppy) and *Corydalis* spp. Prisms (CHCl₃). Mp 254-256°. [α]_D²⁵ -420 (c, 0.69 in Py).
- O¹¹-De-Me, hydrobromide: Mp 245-246°. [α]_D²⁰ -240 (c, 1.0 in MeOH).
- O¹¹-De-Me: **Corytenchine** [18090-68-1]
C₂₀H₂₃NO₄ 341.406
Alkaloid from *Corydalis ochotensis*. Cryst. (MeOH). Mp 257-258°. [α]_D³⁰ -268 (c, 0.89 in CHCl₃).
- (±)-form [13407-95-9]
Cryst. (MeOH/Et₂O). Mp 148-149°.
- HQ1791000
N-Oxide (α -): [75821-48-6]
Synthetic. Cryst. (MeOH/CHCl₃). Mp 130-135°.
- N-Oxide (β -): [75821-47-5]
Synthetic. Cryst. (MeOH/CHCl₃). Mp 167-168°.
- O³,O¹⁰-Di-de-Me: [62744-19-8]
Synthetic. Mp 218-220°.
- O²,O¹⁰-Di-de-Me: Synthetic. Cryst. (CHCl₃/MeOH). Mp 240-241°.
- O²-De-Me: Synthetic. Mp 170-172°.
- O³-De-Me, hydrochloride:
Powder (EtOH). Mp 212-213°.
- O³-De-Me: [40225-82-9]
Powder (MeOH or Me₂CO/Et₂O). Mp 182-184° dec. Could not be resolved.
- O²,O¹¹-Di-de-Me: [6719-48-8]
Cryst. (EtOH). Mp 233-234°.
- O¹¹-De-Me: [26863-94-5]
Synthetic. Cryst. (CHCl₃/MeOH). Mp 236-237°.
- Manske, R.H.F. et al., *Can. J. Res., Sect. B*, 1938, **16**, 81 (*Coreximine*)
Corradi, H. et al., *Helv. Chim. Acta*, 1956, **39**, 889 (*synth, abs config*)
Schmutz, J. et al., *Helv. Chim. Acta*, 1959, **42**, 335 (*Xylopinine, Discretine, isol, uv, struct*)
Tomita, M. et al., *Yakugaku Zasshi*, 1960, **80**, 880; 1138 (*Phellodendrine*)
Battersby, A.R. et al., *Tetrahedron*, 1961, **14**, 46 (*Coreximine, synth, uv*)
Bernouilli, F. et al., *Helv. Chim. Acta*, 1963, **46**, 323 (*Discretine, struct*)
Ninomiya, I. et al., *Chem. Comm.*, 1973, 137 (*synth*)
Irie, H. et al., *Chem. Pharm. Bull.*, 1973, **21**, 855 (*synth*)
Kametani, T. et al., *Chem. Pharm. Bull.*, 1973, **21**, 907 (*Discretine, synth*)
Tani, C. et al., *Chem. Pharm. Bull.*, 1975, **23**, 313 (*synth*)
Mehra, K. et al., *Indian J. Chem., Sect. B*, 1976, **14**, 58; 216; 844 (*Govadine, Govanine*)
Lu, S.T. et al., *J.C.S. Perkin I*, 1976, 63 (*Corytenchine, uv, ms, pmr*)
Khamdamov, I. et al., *CA*, 1977, **87**, 15836x (*pharmacol*)
Govindachari, T.R. et al., *Indian J. Chem., Sect. B*, 1977, **15**, 873 (*Corytenchine, synth, uv, pmr, ms*)
Kametani, T. et al., *J.C.S. Perkin I*, 1977, 1151; 1981, 2830 (*synth*)
Chiang, H.-C. et al., *J.O.C.*, 1977, **42**, 3190 (*synth, 10-Demethylidiscetretine*)
Brockmann-Hanssen, E. et al., *J.O.C.*, 1977, **42**, 3588 (*Coreximine, pmr*)
- Tourwe, O. et al., *Org. Magn. Reson.*, 1977, **9**, 341 (*pmr*)
Chinnasamy, P. et al., *Tetrahedron*, 1980, **36**, 1515 (*oxides*)
Lin, Z. et al., *Fudan Xuebao, Ziran Kexuebau*, 1981, **20**, 446-449; *CA*, **97**, 39192s (*Corydalis L*)
Ohiri, F.C. et al., *Planta Med.*, 1983, **49**, 162 (*Discretine, pmr*)
Meyers, A.I. et al., *Angew. Chem., Int. Ed.*, 1984, 458 (*synth*)
Mali, R.S. et al., *Indian J. Chem., Sect. B*, 1984, **23**, 268 (*Discretine, synth, uv, pmr*)
Hocquemiller, R. et al., *J. Nat. Prod.*, 1984, **47**, 353 (*10-Demethylidiscetretine*)
Cavé, A. et al., *J. Nat. Prod.*, 1986, **49**, 602 (*Artavenustine, 10-Demethylidiscetretine*)
Patra, A. et al., *Phytochemistry*, 1987, **26**, 547 (*oxides*)
Dai-Ho, G. et al., *J.O.C.*, 1988, **53**, 5113 (*synth, ir, pmr, cmr*)
Orito, K. et al., *Org. Prep. Proced. Int.*, 1989, **21**, 309 (*synth*)
Czarnocki, Z. et al., *J. Chem. Res., Synop.*, 1992, 334 (*synth*)
Cobas, A. et al., *J.O.C.*, 1992, **57**, 6765 (*synth*)
Hu, S.-M. et al., *Chem. Pharm. Bull.*, 1993, **41**, 1866 (*Dauricoside*)
Takano, S. et al., *Heterocycles*, 1993, **35**, 47 (*synth*)
Queiroz, E.F. et al., *J. Nat. Prod.*, 1996, **59**, 438 (*Spinosine, Pessoine*)
Comins, D.L. et al., *Tetrahedron*, 1997, **48**, 16327-16340 (*synth*)
Tsai, I.-L. et al., *Phytochemistry*, 1998, **48**, 1377-1382 (*Isocoreximine*)
Davis, F.A. et al., *J.O.C.*, 2002, **67**, 1290-1296 (*synth*)
Mujahidin, D. et al., *Eur. J. Org. Chem.*, 2005, 2689-2693 (*synth*)

Xylostosidine

X-31

[74518-57-3]



C₁₈H₂₅NO₈S 415.463
Alkaloid from *Lonicera xylosteum* (Caprifoliaceae). Mp 190-191°. [α]_D²⁰ -289 (c, 0.56 in MeOH).

S-Oxide (R-): **Loxylostosidine B**

[78184-23-3]
C₁₈H₂₅NO₉S 431.463
Trace alkaloid from *Lonicera xylosteum* (Caprifoliaceae). Noncryst. [α]_D²⁰ -287 (c, 0.42 in MeOH).

S-Oxide (S-): **Loxylostosidine A**


[78119-20-7]
C₁₈H₂₅NO₉S 431.463
Alkaloid from *Lonicera xylosteum* (Caprifoliaceae). Noncryst. [α]_D²⁰ -248 (c, 0.65 in MeOH).

Chaudhuri, R.K. et al., *Helv. Chim. Acta*, 1980, **63**, 1045 (*uv, ir, pmr, cmr, ms, struct*)
Chaudhuri, R.K. et al., *Tet. Lett.*, 1981, 559 (*oxides*)

Tietze, L.F. et al., *Annalen*, 1989, 1241 (*synth, Xylostosidine, Loxylostosidine A*)

Yaequinolone B

[889659-55-6]


Relative
ConfigurationC₂₁H₂₁NO₆ 383.4Prod. by *Penicillium*

um sp. FKI-2140. Insecticidal agent. Pale yellow powder. $[\alpha]_D^{23} +41.2$ (c, 0.1 in EtOH). λ_{\max} 228 (ε 15600); 334 (ε 7900); 357 (ε 8900) (EtOH).

Uchida, R. *et al.*, *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*isol*, *pmr*, *cmr*)

Yaequinolone C

[889659-56-7]



Relative Configuration

C₂₇H₃₃NO₇ 483.56


Prod. by *Penicillium* sp. FKI-2140. Insecticidal agent. Pale yellow powder. $[\alpha]_D^{23} +32.4$ (c, 0.1 in EtOH). λ_{\max} 220 (ε 25900); 279 (ε 13800); 290 (ε 10400); 324 (ε 12900) (EtOH).

23-Deoxy, 23,24-didehydro: **Yaequinolone F** [889659-59-0]

C₂₇H₃₁NO₆ 465.545

Prod. by *Penicillium* sp. FKI-2140. Insecticidal agent. Pale yellow powder. λ_{\max} 219 (ε 20600); 279 (ε 11800); 287 (ε 9100); 324 (ε 10800) (EtOH).

Uchida, R. *et al.*, *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*isol*, *pmr*, *cmr*)

Yaequinolone J₁Relative
ConfigurationC₂₇H₃₁NO₅ 449.546

Prod. by *Penicillium* sp. FKI-2140. Insecticidal agent. Pale yellow powder. $[\alpha]_D^{23} -65.6$ (c, 0.1 in EtOH). λ_{\max} 218 (ε 21700); 284 (ε 7900); 295 (ε 8600); 318 (ε 8000); 324 (ε 7200) (EtOH).

2'-Epimer: **Yaequinolone J₂**C₂₇H₃₁NO₅ 449.546

Y-1


Prod. by *Penicillium* sp. FKI-2140. Pale yellow powder. $[\alpha]_D^{23} +181.7$ (c, 0.1 in EtOH). λ_{\max} 218 (ε 17900); 284 (ε 9300); 295 (ε 9800); 324 (ε 8700) (EtOH).

Uchida, R. *et al.*, *Org. Lett.*, 2005, **7**, 5701-5704 (*pmr*, *cmr*, *struct*)

Uchida, R. *et al.*, *J. Antibiot.*, 2006, **59**, 646-651; 652-658 (*isol*, *activity*)

Yagonine

[95377-96-1]

C₂₀H₁₇NO₆ 367.357

Alkaloid from *Sarcocapnos enneaphylla* (Papaveraceae). Red needles (EtOH). Mp 226-227°.

O^{3'}-De-Me: **3,4-Dioxosarcocapnidine**

[120186-25-6]

C₁₉H₁₅NO₆ 353.331

Trace alkaloid from aerial parts of *Sarcocapnos baetica* ssp. *integrifolia* (Papaveraceae). Orange amorph. solid.


Campello, M.J. *et al.*, *Tet. Lett.*, 1984, **25**, 5933 (*uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)

Castedo, L. *et al.*, *Heterocycles*, 1988, **27**, 2783 (3,4-Dioxosarcocapnidine)

Yanangine

5-Hydroxytiliacorinine

[105705-86-0]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from *Tiliacora triandra* (Menispermaceae). Cryst. + 1/2 CH₂Cl₂. Mp 237-240°. $[\alpha]_D^{19.2} +356$ (c, 0.6 in CHCl₃).

N^{2'}-De-Me: **Noryanangine**

[119002-88-9]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the aerial parts of *Tiliacora triandra* (Menispermaceae). Cryst. + 1/2 CH₂Cl₂. Mp 196-198°. $[\alpha]_D^{20} +295$ (c, 0.5 in CHCl₃).

O⁵-Me: **N-Methyltiliamosine**

[87686-96-2]

C₃₇H₃₈N₂O₆ 606.717

Alkaloid from leaves of *Tiliacora racemosa* (Menispermaceae). Mp 142-145° (synth). $[\alpha]_D^{26} +281$ (c, 0.89 in CHCl₃) (synth). $[\alpha]_D^{25} +510$ (c, 1.5 in

CHCl₃).O⁵-Me, N^{2'}-de-Me: **Tiliamosine**

[62592-71-6]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from leaves and stems of *Pachygone ovata* and leaves of *Tiliacora racemosa* (Menispermaceae). Amorph. solid (CHCl₃/MeOH). Mp 167-170°. $[\alpha]_D^{25} +267$ (c, 0.48 in CHCl₃).

O⁵-Me, N^{2'}-de-Me, O^{12'}, N^{2'}-di-Ac:

Cryst. (CHCl₃). Mp 182-184°. $[\alpha]_D^{25} +423$ (c, 0.35 in CHCl₃).

O⁵, O^{12'}-Di-Me:

Semisynthetic; O,O,N,N-tetra-Me deriv. of Pachygonamine. Amorph. Mp 157-160°. $[\alpha]_D^{24} +230$ (c, 0.59 in CHCl₃).

O⁶-De-Me, O⁵-Me:

Semisynthetic; di-N-Me deriv. of Pachygonamine. Cryst. (CHCl₃/MeOH). Mp 199-202° dec. $[\alpha]_D^{25} +327$ (c, 0.43 in CHCl₃).

O⁶-De-Me, O⁵-Me, N^{2'}-de-Me: **N-****Methylpachygonamine**

[87686-94-0]

C₃₅H₃₄N₂O₆ 578.663

Alkaloid from the leaves and stems of *Pachygone ovata* (Menispermaceae). Amorph. (CHCl₃/MeOH). Mp 182-185° dec. $[\alpha]_D^{27} +287$ (c, 0.23 in MeOH).

O⁶-De-Me, O⁵-Me, di-N-de-Me: **Pachygonamine**

[87686-93-9]

C₃₄H₃₂N₂O₆ 564.637

Alkaloid from the leaves and stems of *Pachygone ovata* (Menispermaceae). Amorph. (CHCl₃). Mp 225-227° dec. $[\alpha]_D^{25} +257$ (c, 0.28 in MeOH).

O^{12'}-De-Me, O^{12'}-Me: **Tilianangine. 5-****Hydroxyyanangorinine**

[105801-15-8]

C₃₆H₃₆N₂O₆ 592.69

Alkaloid from *Tiliacora triandra* (Menispermaceae). Amorph. Mp 143-149°. $[\alpha]_D^{19.7} +258.6$ (c, 0.6 in CHCl₃).

13'-Hydroxy, O⁵-Me: **Tiliacosine**

[412913-17-8]

C₃₇H₃₈N₂O₇ 622.716

Alkaloid from the leaves of *Tiliacora racemosa*.

13'-Hydroxy, O⁵-Me, N^{2'}-de-Me: **Tiliasine**

[412913-18-9]

C₃₆H₃₆N₂O₇ 608.69

Alkaloid from the leaves of *Tiliacora racemosa*.

Guha, K.P. *et al.*, *Tet. Lett.*, 1976, 4241(Tiliamosine, *uv*, *ir*, *pmr*, *ms*, *struct*)Cassels, B.K. *et al.*, *Heterocycles*, 1980, **14**, 211(Tiliamosine, *abs config*)Sultanbawa, M.U.S. *et al.*, *Heterocycles*, 1983,**20**, 1927 (Pachygonamine, N-

Methylpachygonamine)


Pachaly, P. *et al.*, *Arch. Pharm. (Weinheim,**Ger.*), 1986, **319**, 841; 872 (Yanangine, *isol*,*uv*, *ir*, *pmr*, *cmr*, *ms*, *struct*)Pachaly, P. *et al.*, *Planta Med.*, 1988, **54**, 433

(Noryanangine, Norisoyanangine)

Ray, A.K. *et al.*, *Phytochemistry*, 1989, **28**, 675;1990, **29**, 1023 (Tiliamosine, N-Methyltiliamosine, *cmr*)Schiff, P.L. *et al.*, *J. Nat. Prod.*, 1997, **60**, 934-953 (Tiliamosine, *pmr*)

Seal, T. *et al.*, *Nat. Prod. Sci.*, 2001, **7**, 83-86;
CA, **136**, 322041 (*Tiliacosine*, *Tiliasine*)

Yaoundamine A Y-6
[188417-17-6]



$C_{24}H_{25}NO_4$ 391.466
Alkaloid from *Ancistrocladus korupensis* (Ancistrocladaceae). Antimalarial agent. Light yellow solid. $[\alpha]_D^{25} +240$ (c, 0.3 in MeOH). λ_{max} 230 (log ϵ 4.64); 310 (log ϵ 4.08); 322 (log ϵ 4.04); 337 (log ϵ 4.03); 381 (log ϵ 4.3) (MeOH).

6-O- α -L-Rhamnopyranoside: **Yaoundamine B**

[188131-26-2]


$C_{30}H_{35}NO_8$ 537.608

Alkaloid from *Ancistrocladus korupensis*. Antimalarial agent. Light yellow solid. $[\alpha]_D^{25} -82$ (c, 0.3 in MeOH). λ_{max} 230 (log ϵ 4.72); 269 (log ϵ 4.06); 309 (log ϵ 4.11); 317 (log ϵ 4.1) (MeOH). λ_{max} 230 (ϵ 63650); 269 (ϵ 15020); 309 (ϵ 12960); 317 (ϵ 12400) (MeOH) (Berdy).

Bringmann, G. *et al.*, *Tetrahedron*, 1997, **53**, 2817 (*cd, abs config*)

Hallock, Y.F. *et al.*, *Tetrahedron*, 1997, **53**, 8121 (*isol, uv, ir, pmr, cd, cd*)

Yatakemycin Y-7
[606136-98-5]



Absolute Configuration

$C_{35}H_{29}N_5O_8S$ 679.709

Related to Duocarmycin A, D-952 and Rachelmycin, R-4. Struct. revised in 2004. Prod. by *Streptomyces* sp. TP-A0356. Antifungal and cytotoxic agent. Yellow powder. Mp > 220° dec. $[\alpha]_D^{24} +99.9$ (c, 0.39 in DMF). λ_{max} 210 (ϵ 31200); 278 (sh); 310 (sh); 382 (ϵ 32600) (MeOH).

Igarashi, Y. *et al.*, *J. Antibiot.*, 2003, **56**, 107-113 (*isol, pmr, cmr, ms, activity*)

Tichenor, M.S. *et al.*, *J.A.C.S.*, 2006, **128**, 15683-15696 (*synth, abs config*)

Okano, K. *et al.*, *Chem. Asian J.*, 2008, **3**, 296-309 (*synth*)

Di Micco, S. *et al.*, *Eur. J. Org. Chem.*, 2008, 2454-2462 (*complex struct*)

Tichenor, M. *et al.*, *Nat. Prod. Rep.*, 2008, **25**, 220-226 (*rev*)

Yemensine Y-8
[1362-35-2]

$C_{18}H_{19}NO_4$ 313.352

Amaryllidaceae alkaloid. Struct. unknown. Isol. from the bulbs of *Crinum yemensense* (Amaryllidaceae). Prisms + 1H₂O (Me₂CO). Mp 193° dec. $[\alpha]_D^{25} +100$ (c, 0.2 in CHCl₃).

Hydroiodide:

Microcryst. powder (H₂O). Mp 205-206° dec.

Picrate:


Needles (Me₂CO/MeOH). Mp 262-263° dec.

Methiodide:

Microcryst. powder (H₂O). Mp 235° dec.

Boit, H.-G. *et al.*, *Chem. Ber.*, 1957, **90**, 2203-2206 (*isol*)

Yersiniabactin Y-9
Yersiniophore
[176707-32-7]



Absolute Configuration

$C_{21}H_{27}N_3O_4S_3$ 481.66

Natural prod. is a mixt. of C-10 epimers. Prod. by *Yersinia enterocolitica*. Siderophore. λ_{max} 255 (log ϵ 3.72); 310 (log ϵ 3.4) (H₂O). λ_{max} 210; 250; 280 (MeOH) (Berdy). λ_{max} 210; 260; 320; 390 (H₂O) (Berdy).


Drechsel, H. *et al.*, *Annalen*, 1995, 1727-1733 (*isol, uv, cd, pmr, cmr, ir*)

Chambers, C.E. *et al.*, *BioMetals*, 1996, **9**, 157-167 (*isol, struct, props*)

Ino, A. *et al.*, *Tetrahedron*, 2001, **57**, 1897-1902 (*synth, abs config*)

Pfeifer, B.A. *et al.*, *Appl. Environ. Microbiol.*, 2003, **69**, 6698-6702 (*biosynth*)

Yesonine Y-10
Deveratrolyyesoline
[116139-64-1]



$C_{21}H_{29}NO_3$ 343.465

Alkaloid from rhizomes of *Aconitum yesoense* var. *macroyoense* (Ranunculaceae). Amorph. $[\alpha]_D^{25} +2.4$ (c, 0.17 in EtOH).

15-O-(3,4-Dimethoxybenzoyl): **Yesoline**.


15-Veratrolyyesoline
[116159-74-1]

$C_{30}H_{37}NO_6$ 507.625

Alkaloid from rhizomes of *Aconitum yesoense* var. *macroyoense* (Ranunculaceae). Amorph. $[\alpha]_D^{25} -10.6$ (c, 0.34 in EtOH).

Wada, K. *et al.*, *Heterocycles*, 1988, **27**, 1249; 1989, **29**, 2141 (*isol, uv, ir, pmr, cmr, ms, struct*)

Yibeissine Y-11
17,23-Epoxy-3,11-dihydroxyveratraman-6(5H)-one, 9CI
[143502-51-6]




$C_{27}H_{41}NO_4$ 443.625

Alkaloid from bulbs of *Fritillaria pallioflora* (Liliaceae). Cryst. Mp 164.5-166°. $[\alpha]_D^{25} -47.6$ (c, 0.13 in EtOH).

Xu, Y.J. *et al.*, *Yaouxue Xuebao*, 1992, **27**, 121; *CA*, **117**, 147162x (*isol*)

Yohambinine Y-12
5-Methyl-yohimban, 9CI. 5-Methylpseudo-yohimbane
[109305-83-1]



Absolute Configuration


$C_{20}H_{26}N_2$ 294.439

The struct. shown is almost certainly incorrect. All stereoisom have been synth. and the nat. alkaloid was not identical with any of them. Alkaloid from the roots of *Rauwolfia serpentina* (Apocynaceae). Elongated rods (MeOH). Mp 188-190°. $[\alpha]_D^{20} +98$ (MeOH). λ_{max} 218; 278 (no solvent reported).

Siddiqui, S. *et al.*, *Tet. Lett.*, 1987, **28**, 1311-1312 (*isol, uv, ir, pmr, cmr, ms, struct*)

Lohse, C. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 945-961

Yohimbine Y-13
Methyl 17-hydroxy-yohimban-16-carboxylate, 9CI. Aphrodine. Corymbin†. Corynine. Hydroergotocin. Quebrachine. Yohimex. Yohimvetol
[146-48-5]
[24252-70-8 (\pm -form)]



Absolute Configuration

- $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Alchornea*, *Alstonia*, *Amsonia*, *Aspidosperma*, *Catharanthus*, *Corynanthe*, *Diplorhynchus*, *Rauwolfia*, *Vinca*, *Pausinystalia*, *Pouteria*, *Hunteria* and *Pseudocinchona* (preferred genus name *Corynanthe*) spp. (Apocynaceae, Rubiaceae). Local anaesthetic. Selective α_2 -adrenoceptor antagonist. Yohimbine and its stereoisomers are used as tools for study of different adrenoceptor sites. Antidepressant agent used clinically. Shows antihypertensive activity, causes vasopressin release and shows antidiuretic activity. Alleged aphrodisiac. Anxiogenic activity in rodents. Needles (EtOH aq.). Mp 241° (235-236°). $[\alpha]_D^{20} +108$ (Py). Log P 2.24 (calc).
- Adverse human effects by ingestion and skin absorption e.g. cardiotoxic. Transient anxiety also seen following large overdose. LD₅₀ (mus, orl) 51 mg/kg. ZG1000000
- Hydrochloride*: Yohimbine hydrochloride. Gynimbine. Menolysin. Parkimbine. Tosanpin. Yobinol. Yocon. Yohydrol [65-19-0]
Plates. Mp 302°. $[\alpha]_D^{22} +103$ (H₂O). Component of Potensan, Vikonon.
- LD₅₀ (rat, ipr) 55 mg/kg. ZG1015000
- Et ester analogue*: Ethyl yohimbate [73840-32-1]
Mp 190°.
- LD₅₀ (mus, orl) 36 mg/kg. LD₅₀ (mus, ipr) 17 mg/kg. ZG0800000
- Parent acid*: Yohimbic acid, INN. Yohimbic acid [522-87-2]
 $C_{20}H_{24}N_2O_3$ 340.421
Alkaloid from the roots of *Rauwolfia serpentina*. α_2 -Adrenoceptor antagonist. Cryst. (MeOH or EtOH); cryst. (H₂O) (as monohydrate). Mp 280-300° Mp 265° (monohydrate). $[\alpha]_D +148.7$. λ_{max} 221 (log ϵ 4.28); 273 (sh) (log ϵ 3.63); 279 (log ϵ 3.64); 288 (log ϵ 3.57) (MeOH).
- O-Ac*: O-Acetylyohimbine [15471-41-7]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Aspidosperma excelsum* (Apocynaceae). Mp 150°.
- O-Ac*; hydrochloride: [14430-18-3]
Mp 274-275° dec.
- Me ether*: 17-O-Methylyohimbine [132339-47-0]
 $C_{22}H_{28}N_2O_3$ 368.475
Alkaloid from the roots of *Amsonia elliptica* (Apocynaceae). Prisms (CHCl₃). Mp 147°. $[\alpha]_D +12$ (c, 0.188 in MeOH). λ_{max} 224 ; 288 (sh) ; 292 (MeOH).
- 3-Epimer*: Pseudoyohimbine. ψ -Yohimbine [84-37-7]
[25920-66-5 (\pm -form)]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Alstonia quaternata*, *Catharanthus trichophyllus*, *Corynanthe yohimbe*, *Voacanga africana*, *Rauwolfia* spp., *Uncaria attenuata* ssp. *attenuata* and *Uncaria callophylla* (Apocynaceae, Rubiaceae). Mp 293° (268°). $[\alpha]_D^{19} +27$ (c, 1.0 in Py).
- 3-Epimer, hydrochloride*: Mp 233°. $[\alpha]_D^{20} -10$ (c, 1.0 in H₂O).
- 16-Epimer*: Corynanthine. Rauhimbine [483-10-3]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Pseudocinchona africana*, *Corynanthe yohimbe*, *Corynanthe pachyceras*, *Rauwolfia canescens* and *Pausinystalia macroceras* (Rubiaceae, Apocynaceae). Strong α_1 -adrenoceptor antagonist, sympatholytic agent. Antileishmanial. Mp 225-226°. $[\alpha]_D -123$ (EtOH). $[\alpha]_D -85$ (Py). Log P 2.24 (calc). λ_{max} 226 ; 283 ; 290 (MeOH) (Berdy).
- ZG1025000
- 16-Epimer, hydrochloride*: Mp 285-290°. $[\alpha]_D -63$ (H₂O).
- 16-Epimer, O-Ac*: Mp 147°.
- 16-Epimer, parent acid*: Corynanthic acid Mp 337° (284°). $[\alpha]_D -85.9$ (c, 1 in Py) (-70.7).
- 17-Epimer*: β -Yohimbine. Amsonine [549-84-8]
[2516-78-1 (\pm -form)]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Amsonia*, *Aspidosperma*, *Corynanthe*, *Diplorhynchus*, *Rauwolfia*, *Voacanga* and *Pausinystalia* spp. (Apocynaceae, Rubiaceae). Mp 236° (260-262°). $[\alpha]_D -47$ (-54) (Py). γ -Yohimbine was a mixt. of β -Yohimbine with Yohimbine.
- ZG1000030
- 17-Epimer, hydrochloride*: Mp 297°. $[\alpha]_D +29$ (H₂O).
- 17-Epimer, N⁴-oxide*: β -Yohimbine N-oxide
 $C_{21}H_{26}N_2O_4$ 370.447
Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. solid (MeOH). Mp 230°. Config. apparently wrongly assigned in CA.
- 17-Epimer, 3(N)-dehydro*: 3,4-Dehydro- β -yohimbine [88607-65-2]
 $C_{21}H_{25}N_2O_3^{\oplus}$ 353.44
Quaternary alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). $[\alpha]_D +114$ (c, 0.95 in MeOH).
- 20-Epimer*: Alloyohimbine [522-94-1]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Corynanthe yohimbe*, *Corynanthe paniculata*, *Pausinystalia yohimbe*, *Pausinystalia angolensis* and *Rauwolfia* spp. (Apocynaceae). Mp 165-170° (135-140°). $[\alpha]_D -84$ (Py).
- 20-Epimer, hydrochloride*: Mp 280°. $[\alpha]_D^{19} +33$ (H₂O).
- 20-Epimer, O-Ac*: Acetylalloyohimbine [81571-22-4]
 $C_{23}H_{28}N_2O_4$ 396.485
Alkaloid from *Rauwolfia nitida* root bark (Apocynaceae). Mp 136°. $[\alpha]_D^{20} -25$ (Py) (synthetic). $[\alpha]_D^{22} -89$ (c, 0.01 in CHCl₃) (nat.). λ_{max} 225 (log ϵ 4.58); 284 (log ϵ 3.77); 290 (log ϵ 3.65) (MeOH).
- 3,17-Diepimer*: 3-Epi- β -yohimbine [41904-78-3]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from the root of *Rauwolfia linearifolia* (Apocynaceae). Cryst. Mp 201-203°.
- 16,20-Diepimer*: α -Yohimbine. Chalcupine A. Isoyohimbine. Mesoyohimbine. Rauwolsine [131-03-3]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Alstonia*, *Aspidosperma*, *Corynanthe*, *Pseudocinchona*, *Rauwolfia* and *Vinca* spp. Also from *Pausinystalia croceras* (Apocynaceae, Rubiaceae). Strong selective α_2 -adrenoceptor antagonist, also reported to possess aphrodisiac props. Mp 243° (235-236°). $[\alpha]_D^{19} -27$ (c, 1.0 in EtOH). Log P 2.24 (calc). Serpine was a mixt. of α -Yohimbine with Yohimbine, Y-13. λ_{max} 227 (log ϵ 4.5); 281 (log ϵ 3.93) (no solvent reported).
- LD₅₀ (rat, ipr) 50 mg/kg. ZG1005000
- 16,20-Diepimer, hydrochloride*: [6211-32-1]
Mp 288°. $[\alpha]_D^{19} +58$ (c, 1.0 in H₂O).
- ZG1035000
- 17,20-Diepimer*: 17-Epi-alloyohimbine [59952-56-6]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from seeds of *Aspidosperma oblongum*. Amorph. $[\alpha]_D -8$ (c, 0.74 in CHCl₃). λ_{max} 227 (log ϵ 4.78); 274 (log ϵ 3.7); 283 (log ϵ 3.92); 291 (log ϵ 3.85) (EtOH).
- 3,16,20-Triepimer*: Isorauyohimbine. 3-Epi-*isoraufolsine*. Isoraufolsine. Alkaloid Su 3078. 3-Epi- α -yohimbine [483-09-0]
 $C_{21}H_{26}N_2O_3$ 354.448
Alkaloid from *Rauwolfia serpentina* and *Voacanga africana* (Apocynaceae). Prisms (MeOH aq.). Mp 225°. $[\alpha]_D^{20} -104$ (c, 0.5 in Py). Polymorphs having lower Mps are formed from other solvs.
- 3,16,20-Triepimer, hydrochloride*: Prisms (MeOH/Et₂O). Mp 235-240° dec. $[\alpha]_D^{26} -75$ (c, 1.0 in H₂O).
- 3,16,20-Triepimer, parent acid*: Isorauyohimbic acid
 $C_{20}H_{24}N_2O_3$ 340.421
Alkaloid from the roots of *Rauwolfia serpentina*. Cryst. (MeOH). Mp 234-236°. $[\alpha]_D^{28} -131$ (c, 0.25 in Py). λ_{max} 220 (log ϵ 4.47); 272 (log ϵ 3.82); 279 (log ϵ 3.82); 289 (log ϵ 3.74) (MeOH).
- Stereoisomer*: Vinca pusilla Alkaloid A [41787-60-4]
 $C_{21}H_{26}N_2O_3$ 354.448
Isol. from *Vinca pusilla*. Mp 269° dec. Of undetd. stereochem., except that it appears to be (3 β -).
- Aldrich Library of FT-IR Spectra, 1st edn.*, 1985, 2, 1068B; 1069B (ir)
- Aldrich Library of 13C and 1H FT NMR Spectra*, 1992, 3, 614C (nmr)
- Spiegel, L. et al., *Ber.*, 1903, 36, 169 (Yohimbic acid, isol)
- Janot, M.M. et al., *Bull. Soc. Chim. Fr.*, 1949, 509 (synth)
- Swan, G.A. et al., *J.C.S.*, 1950, 1534-1539 (struct)


- Marion, L. et al., *Alkaloids (Academic Press)*, 1952, **2**, 369-498 (rev, bibl, occur)
- Le Hir, A. et al., *Bull. Soc. Chim. Fr.*, 1953, 1027-1032 (*α*-Yohimbine, β-Yohimbine, Alloyohimbine, struct)
- Hofmann, A. et al., *Helv. Chim. Acta*, 1954, **37**, 314-320 (*Isorauhimbine*, isol)
- Bader, F.E. et al., *J.A.C.S.*, 1955, **77**, 3547-3550 (*Isorauhimbine*, struct)
- Goutarel, R. et al., *Helv. Chim. Acta*, 1957, **40**, 156-160 (*Isorauhimbine*, struct, config)
- Antonaccio, L.D. et al., *J.A.C.S.*, 1962, **84**, 2161-2169 (ms)
- Raymond-Hamet, et al., *C. R. Hebd. Seances Acad. Sci.*, 1963, **257**, 2351 (*Yohimbic acid*, pharmacol)
- Albright, J.D. et al., *J.O.C.*, 1963, **28**, 38-45 (pmr)
- Ban, Y. et al., *Tetrahedron*, 1964, **20**, 2877-2884 (sterochem)
- Benoin, P.R. et al., *Can. J. Chem.*, 1967, **45**, 725-730 (*O-Acetylyohimbine*)
- Klyne, W. et al., *Helv. Chim. Acta*, 1967, **50**, 115-125 (ord)
- Van Tamelen, E.E. et al., *J.A.C.S.*, 1969, **91**, 7315-7333 (*Pseudoyohimbine*, synth, bibl)
- Fales, H.M. et al., *J.A.C.S.*, 1970, **92**, 1590-1597 (ms)
- Stork, G. et al., *J.A.C.S.*, 1972, **94**, 5109-5110 (*β*-Yohimbine, *Pseudoyohimbine*, synth)
- Chatterjee, A. et al., *Indian J. Chem.*, 1973, **11**, 7-8 (*Vinca Pusilla Alkaloid A*)
- Ambady, G. et al., *J. Cryst. Mol. Struct.*, 1973, **3**, 37-45 (cryst struct, abs config)
- Levin, R.H. et al., *J.O.C.*, 1973, **38**, 1983-1986 (cmr)
- Töke, L. et al., *J.O.C.*, 1973, **38**, 2496-2500 (*Alloyohimbine*, *Isorauhimbine*, synth, abs config)
- Catharanthus Alkaloids*, (Eds., Taylor, W.I. et al), Marcel Dekker, 1975, (rev)
- Szantay, C. et al., *Chem. Ber.*, 1976, **109**, 1737-1748 (*α*-Yohimbine, β-Yohimbine, synth)
- Miet, C. et al., *Phytochemistry*, 1977, **16**, 803-805 (pmr)
- Fanso-Free, S.N.Y. et al., *J.A.C.S.*, 1979, **101**, 1549-1553 (*N nmr*)
- Wenkert, E. et al., *J.A.C.S.*, 1979, **101**, 5370-5376 (*Yohimbine*, *Pseudoyohimbine*, *α*-Yohimbine, β-Yohimbine, synth, cmr)
- Amer, M.A. et al., *Phytochemistry*, 1981, **20**, 2569-2573 (*Acetylalloyohimbine*)
- Honty, K. et al., *J.O.C.*, 1982, **47**, 5111-5114 (*Isorauhimbine*, synth)
- Miyata, O. et al., *Chem. Comm.*, 1983, 1231-1232 (*Alloyohimbine*, synth)
- Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 708-722 (*Aspidosperma oblongum constits*)
- Goldberg, M.R. et al., *Pharmacol. Rev.*, 1983, **35**, 143-180 (pharmacol, rev)
- Goh, S.H. et al., *Phytochemistry*, 1985, **24**, 880-881 (*Pseudoyohimbine*)
- Martin, S.F. et al., *Tet. Lett.*, 1985, **26**, 5227-5230 (*α*-Yohimbine, synth)
- Szantay, C. et al., *Alkaloids (Academic Press)*, 1986, **27**, 131-268 (rev)
- Blasko, G. et al., *Annalen*, 1986, 655-663 (*β*-Yohimbine, synth)
- Mekkawi, A.G. et al., *Anal. Profiles Drug Subst.*, 1987, **16**, 731-768 (rev, synth, anal, pharmacol)
- Sauerwein, M. et al., *Phytochemistry*, 1990, **29**, 3377-3379 (*17-O-Methyl-yohimbine*)
- Hirai, Y. et al., *Tet. Lett.*, 1990, **31**, 4755-4756 (synth)
- Martinez Perez, J.A. et al., *Phytochemistry*, 1991, **30**, 1352-1353 (*3-Epi-β*-yohimbine)
- Friesen, K. et al., *J. Emergency Med.*, 1993, **11**, 287 (tox)
- Biaggioni, I. et al., *J. Clin. Pharmacol.*, 1994, **34**, 418-423 (pharmacol, human, rev)
- Heidelbaugh, T.M. et al., *Tet. Lett.*, 1998, **39**, 4757-4760 (*Alloyohimbine*, synth)

- Martindale, The Extra Pharmacopoeia*, 32nd edn., Pharmaceutical Press, 1999, 941; 1645
- Staerk, D. et al., *Planta Med.*, 2000, **66**, 531-536 (*α*-Yohimbine, *Corynanthine*, isol, pmr, cmr)
- Tanaka, M. et al., *Tetrahedron*, 2004, **60**, 2271-2281 (*Alloyohimbine*, synth)
- Itoh, A. et al., *J. Nat. Prod.*, 2005, **68**, 848-852 (*Isorauhimbic acid*, isol, pmr, cmr)
- Mergott, D.J. et al., *Org. Lett.*, 2008, **10**, 745-748 (synth)
- Lewis, R.J. et al., *Sax's Dangerous Properties of Industrial Materials*, 10th edn., J. Wiley, 2000, YBJ000; YBS000; YCA000

β-Yohimbine oxindole

Y-14

[88668-03-5]




Absolute Configuration

C₂₁H₂₆N₂O₄ 370.447Alkaloid from the seeds of *Aspidosperma oblongum* (Apocynaceae). Amorph. [α]_D -38 (c, 0.32 in MeOH).Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 708-722 (isol, uv, ir, pmr, ms, cd, struct, synth)

β-Yohimbine pseudoindoxyl

Y-15

[88668-02-4]




Absolute Configuration

C₂₁H₂₆N₂O₄ 370.447Alkaloid from the seeds of *Aspidosperma oblongum*, from leaves of *Ervatamia corymbosa* and from leaves and root bark of *Ervatamia hirta*. Amorph. [α]_D -431 (c, 1.75 in MeOH). λ_{max} 234 (log ε 4.56); 255 (log ε 3.91); 407 (log ε 3.67) (EtOH).Robert, G.M.T. et al., *J. Nat. Prod.*, 1983, **46**, 708-722 (isol, uv, ir, pmr, ms, cd, struct, synth)
Takayama, H. et al., *Heterocycles*, 1996, **42**, 87-92 (abs config, cd)

Yohimbol

Y-16

Yohimbol, 9CI. *Corynanthol* [523-13-7]

Absolute Configuration

C₁₉H₂₄N₂O 296.411Alkaloid from *Hunteria zeylanica* (Apocynaceae). Fine needles. Mp 251-253° (243°). [α]_D -63.4 (-55.4, -38) (MeOH).N^b-Me: Yohimbol methosalt

[100816-28-2]

C₂₀H₂₇N₂O[⊕] 311.446Alkaloid from *Hunteria eburnea*(Apocynaceae). Cryst. + ½ H₂O(Me₂CO aq.) (as chloride). Mp 264-265° (chloride). [α]_D +53 (MeOH)

(synthetic). Opt. rotn. of -16° was recorded for natural alkaloid, but this was thought due to contamination with a small amt. of the highly laevorotatory akuammicine methiodide.

10-Methoxy: 10-Methoxyyohimbol-17α-ol

10-Methoxyyohimbol

C₂₀H₂₆N₂O₂ 326.438Alkaloid from the roots of *Ochrosia**acuminata*. Amorph. powder. [α]_D²⁵

+3.3 (c, 0.14 in MeOH).

17-Epimer: Epiyohimbol

[439-70-3]

C₁₉H₂₄N₂O 296.411Alkaloid from *Hunteria zeylanica*(Apocynaceae). Mp 258°. [α]_D²⁰ -80.1 (c,

2.075 in MeOH).

▶ ZG1255000

Witkop, B. et al., *Annalen*, 1943, **554**, 83-126

(synth)

Bartlett, M.F. et al., *J.O.C.*, 1963, **28**, 1445-1449 (*methochloride*, isol)Mori, K. et al., *Agric. Biol. Chem.*, 1972, **36**,

2605-2606 (synth)

Arambewala, L.S.R. et al., *Phytochemistry*,1981, **20**, 349-350 (*Yohimbol*, *Epiyohimbol*,

isol)

Kametani, T. et al., *J. Het. Chem.*, 1982, **19**,

1217-1219 (synth)

Lavaud, C. et al., *Phytochemistry*, 1982, **21**,


445-447 (isol)

Salim, A.A. et al., *J. Nat. Prod.*, 2004, **67**,1719-1721 (*10-Methoxyyohimbol*)

Ypaoamide

Y-17

[180965-25-7]

C₂₆H₃₆N₂O₅ 456.581

Isol. from a mixed cyanobacteria assem-

blage, actual source believed to be

Lyngbya majuscula. Feeding deterrent.Ichthyotoxic. [α]_D¹⁹ +197 (c, 1.0 inCHCl₃). λ_{max} 222 (log ε 4.2); 270 (log ε

4.1) (MeOH).


Nagle, D.G. et al., *Tet. Lett.*, 1996, **37**, 6263

(isol, uv, ir, pmr, cmr, struct)

Yuanamide

Y-18


[102421-42-1]



C₂₂H₂₃NO₅ 381.427Alkaloid from aerial parts of *Corydalis bulbosa* (Papaveraceae). Prisms (MeOH). Mp 228-230°. Also descr. as an oil.Saá, C. et al., *J.O.C.*, 1986, **51**, 2781 (synth)Ito, C. et al., *Phytochemistry*, 1990, **29**, 2044 (isol, uv, ir, pmr, ms)**Yuehchukene**

[96624-37-2]

Y-19


C₂₆H₂₆N₂ 366.505Alkaloid from the roots of *Murraya paniculata*, *Micromelum falcatum* and *Micromelum integerrimum* (Rutaceae). Exhibits potent antiimplantation activity in rats. Racemic. λ_{max} 225 (ε 14800); 283 (ε 3470); 292 (ε 3090) (EtOH) (Derep).

▶ NK8927150


Kong, Y.-C. et al., *Chem. Comm.*, 1985, 47 (uv, ir, pmr, cmr, ms, struct)Kong, Y.-C. et al., *Jiegou Huaxue*, 1985, **4**, 30; *CA*, **104**, 31728p (cryst struct)Kong, Y.-C. et al., *Biochem. Syst. Ecol.*, 1988, **16**, 485 (isol)Kutney, J.P. et al., *Can. J. Chem.*, 1991, **69**, 949 (synth)Cheng, K.-F. et al., *J.C.S. Perkin I*, 1991, 2955-2959 (synth, abs config)Bergman, J. et al., *Tetrahedron*, 1992, **48**, 759 (synth)Henry, K.J. et al., *Chem. Comm.*, 1993, 510 (synth)Sheu, J.-H. et al., *J.C.S. Perkin I*, 1998, 1959-1965 (synth)Ishikura, M. et al., *Heterocycles*, 2000, **53**, 553-556; 2201-2220 (synth)**Yunnandaphnine E**

[917955-56-7]

Y-20

C₂₃H₃₁NO₄ 385.502Alkaloid from *Daphniphyllum yunnanense*. Powder. [α]_D²⁵ -88 (c, 0.16 in CHCl₃).Di, Y.-T. et al., *J. Nat. Prod.*, 2006, **69**, 1745-1748 (isol, pmr, cmr)**Yunnanensine**

Y-21


C₁₉H₂₄N₂O₃ 328.41Alkaloid from the stems of *Ervatamia yunnanensis*. Pale yellow powder (CHCl₃). Sol. CHCl₃. Mp 183-184°.N⁴-(2-Cyanoethyl)-6-Cyano-6,7-secostemmadenine

[112448-52-9]

C₂₂H₂₇N₃O₃ 381.474Alkaloid from *Alstonia angustiloba* and *Alstonia pneumatophora* (Apocynaceae). [α]_D +19 (c, 0.5 in CHCl₃). Stereochem. not determined.Zeches, M. et al., *J. Nat. Prod.*, 1987, **50**, 714-720 (6-Cyano-6,7-secostemmadenine)Luo, X.G. et al., *Chin. Chem. Lett.*, 2007, **18**, 697-699 (Yunnanensine)**Yuremamine**

[883973-98-6]

Y-22




Relative Configuration

C₂₇H₂₈N₂O₆ 476.528Alkaloid from the stem bark of *Mimosa tenuiflora*. Purple-red solid.Vepsäläinen, J.J. et al., *Planta Med.*, 2005, **71**, 1053-1057 (isol, pmr, cmr, ms)**Yuzurimine C**

[57520-21-5]

Y-23



Absolute Configuration


C₂₃H₂₉NO₅ 399.486Minor alkaloid from the leaves and bark of *Daphniphyllum macropodum* (Daphniphyllaceae). Cryst. (hexane/Me₂CO). Mp 186-187°. [α]_D²⁵ +28 (c, 1.33 in CHCl₃).*Methodide*:Needles (Me₂CO/Et₂O). Mp 209-210° dec.21-Alcohol: **Macropodimine F**. *Pordamacrine A* [949172-76-3]C₂₃H₃₁NO₅ 401.502Alkaloid from the leaves of *Daphniphyllum macropodum*. Vasorelaxant. Oil or amorph. solid. [α]_D²³ -13.8 (c, 0.53 in CHCl₃) (Macropodimine F). [α]_D²⁷ -27 (c, 0.2 in MeOH) (Pordamacrine A).14,15-Didehydro, 21-alcohol: **Macropodimine G** [949172-77-4]C₂₃H₂₉NO₅ 399.486Alkaloid from the leaves of *Daphniphyllum macropodum*. Oil. [α]_D²³ +73.8 (c, 0.42 in CHCl₃). λ_{max} 303 (log ε 3.76) (MeOH).2-Deoxy, 21-alcohol: **Pordamacrine B** [951677-64-8]C₂₃H₃₁NO₄ 385.502Alkaloid from the leaves of *Daphniphyllum macropodum*. Amorph. solid. [α]_D²⁰ +37 (c, 0.3 in MeOH).Toda, M. et al., *Tetrahedron*, 1972, **28**, 1477 (isol)Yamamura, S. et al., *Bull. Chem. Soc. Jpn.*, 1975, **48**, 2120 (uv, ir, pmr, cmr, ms, struct)Li, Z.Y. et al., *Helv. Chim. Acta*, 2007, **90**, 1353-1359 (Macropodimines F,G)Matsuno, Y. et al., *J. Nat. Prod.*, 2007, **70**, 1516-1518 (Pordamacrines A-B, cryst struct)**Yuzurimine D**

Y-24

C₂₄H₃₂NO₅ 414.52Struct. unknown. Alkaloid from *Daphniphyllum macropodum* (Daphniphyllaceae). Mp 194-195°.Toda, M. et al., *Nippon Kagaku Zasshi*, 1970, **91**, 103-108; *CA*, **73**, 22137j (isol)

Zangezurine

[114216-91-0]

C₂₂H₂₅NO₇ 415.442

Probable struct. Biogenetic numbering shown; other schemes also used. Alkaloid from flowering *Papaver zangezuricum* (Papaveraceae). Amorph.

12-Demethoxy: Triniifoline

[201136-31-4]

C₂₁H₂₃NO₆ 385.416

Alkaloid from the aerial parts of *Papaver triniifolium*. Amorph. powder. $[\alpha]_D^{20} +100$ (c, 0.5 in CHCl₃). λ_{max} 231 (log ϵ 3.97); 288 (log ϵ 3.67) (MeOH).

12-Demethoxy, Et ether: O-Ethyltriniifoline

[201136-32-5]

C₂₃H₂₇NO₆ 413.469


Alkaloid from *Papaver triniifolium*. Amorph. solid. $[\alpha]_D^{20} +120$ (c, 0.5 in CHCl₃). Possible artifact. λ_{max} 234 (log ϵ 4.01); 285 (log ϵ 3.56) (MeOH).

Israilov, I.A. *et al.*, *Khim. Prir. Soedin.*, 1987, **23**, 858; *Chem. Nat. Compd. (Engl. Transl.)*, 715 (*Zangezurine*)

Sari, A. *et al.*, *Planta Med.*, 1997, **63**, 575-576 (*Triniifoline, O-Ethyltriniifoline, isol, uv, ir, pmr, ms*)

Zanthobisquinolone

[57147-67-8]


C₂₁H₁₈N₂O₄ 362.384

Alkaloid from root wood of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Needles (CHCl₃/MeOH). Mp 295-298°.

Chen, I.-S. *et al.*, *Phytochemistry*, 1994, **36**, 237 (*isol, uv, ir, pmr, cmr, ms, struct*)

Zanthocadinanine A

[1006378-72-8]



Relative Configuration

Z-1

C₃₇H₄₅NO₅ 583.766

Structurally derived from Dihydrochelerythrine, D-407 and the methyl ether of 4-Cadinene-10-ol. Alkaloid from the stem bark of *Zanthoxylum nitidum*. Needles. Mp 190-191°. $[\alpha]_D^{20} -71.5$ (c, 0.13 in CHCl₃). λ_{max} 229 (log ϵ 3.9); 284 (log ϵ 4.2); 320 (sh) (log ϵ 3.1) (MeOH).

1'-Epimer: Zanthomurolanine

[1006378-70-6]

C₃₇H₄₅NO₅ 583.766

Alkaloid from the stem bark of *Zanthoxylum nitidum*. Needles. Mp 198-199°. $[\alpha]_D^{20} -51.3$ (c, 0.1 in CHCl₃). λ_{max} 229 (log ϵ 3.9); 284 (log ϵ 4.53); 322 (sh) (log ϵ 4.05) (MeOH).

10'-Epimer: Zanthocadinanine B

[1006378-73-9]

C₃₇H₄₅NO₅ 583.766

Alkaloid from the stem bark of *Zanthoxylum nitidum*. Needles (MeOH). Mp 160-161°. $[\alpha]_D^{20} -52.5$ (c, 0.16 in CHCl₃). λ_{max} 229 (log ϵ 4.67); 284 (log ϵ 4.81); 322 (sh) (log ϵ 4.33) (MeOH).

1',8-Diepimer: 8-Epizanthomurolanine

[1006378-71-7]

C₃₇H₄₅NO₅ 583.766

Alkaloid from the stem bark of *Zanthoxylum nitidum*. Oil. $[\alpha]_D^{20} +86.9$ (c, 0.22 in CHCl₃). λ_{max} 227 (log ϵ 4.37); 283 (log ϵ 4.39); 324 (sh) (log ϵ 3.96) (MeOH).

8,10'-Diepimer: 8-Epizanthocadinanine B

[1006378-74-0]


C₃₇H₄₅NO₅ 583.766

Alkaloid from the stem bark of *Zanthoxylum nitidum*. Needles. Mp 162-163°. $[\alpha]_D^{20} +108.1$ (c, 0.87 in CHCl₃). λ_{max} 228 (log ϵ 4.73); 283 (log ϵ 4.75); 322 (sh) (log ϵ 4.32) (MeOH).

Yang, C.-H. *et al.*, *J. Nat. Prod.*, 2008, **71**, 669-673 (*isol, cd, pmr, cmr, cryst struct*)

Zanthomamide

[99933-31-0]

C₁₉H₁₉NO₃ 309.364

Alkaloid from the stem bark of *Zanthoxylum thomense* (Rutaceae). Amorph.

N-De-Me: Zanthomamine

[121817-67-2]

C₁₈H₁₇NO₃ 295.337


Synthetic. Mp 112-114°.

Simeray, J. *et al.*, *Phytochemistry*, 1985, **24**, 2720 (*isol, uv, ir, pmr, ms, struct*)

Adesina, S.K. *et al.*, *Phytochemistry*, 1989, **28**, 839 (*Zanthomamine*)

Zanthosimuline

[155416-20-9]

C₂₀H₂₃NO₂ 309.407

Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Solid. Mp 127-128°. $[\alpha]_D^{20} +7$ (c, 0.1 in CHCl₃).

3',4'-Dihydro, 3'-oxo: Huajiaosimuline

[155416-21-0]

C₂₀H₂₃NO₃ 325.407

Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil.

3',4'-Dihydro, 3'-hydroxy: Simulansine

[176520-66-4]

C₂₀H₂₅NO₃ 327.422

Alkaloid from root bark of *Zanthoxylum simulans* (Szechuan pepper). Oil.

Δ^{2'}-Isomer, 4'-hydroxy: Simulenoline

[198336-56-0]

C₂₀H₂₃NO₃ 325.407

Alkaloid from the bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil. $[\alpha]_D^{23} +6.9$ (c, 0.2 in CHCl₃). λ_{max} 228 (log ϵ 4.33); 256 (sh) (log ϵ 3.86); 336 (log ϵ 3.68); 350 (log ϵ 3.73); 367 (log ϵ 3.56) (EtOH).

Δ^{2'}-Isomer, 4'-hydroperoxy: Peroxysimulenoline

[198336-57-1]

C₂₀H₂₃NO₄ 341.406

Alkaloid from the bark of *Zanthoxylum simulans* (Szechuan pepper) (Rutaceae). Oil. λ_{max} 228 (log ϵ 4.33); 256 (sh) (log ϵ 3.86); 336 (log ϵ 3.68); 350 (log ϵ 3.73); 367 (log ϵ 3.56) (EtOH).

Wu, S.-J. *et al.*, *Phytochemistry*, 1993, **34**, 1659-1661 (*isol, uv, ir, pmr, cmr, ms, struct*)

Chen, I.-S. *et al.*, *Phytochemistry*, 1996, **42**, 217-219 (*Simulansine*)

Chen, I.-S. *et al.*, *Phytochemistry*, 1997, **46**, 525-529 (*Simulenoline, Peroxysimulenoline*)

McLaughlin, M.J. *et al.*, *J.O.C.*, 2001, **66**, 1049-1053 (*Huajiaosimuline, Simulenoline, synth*)


Lee, Y.R. *et al.*, *Synthesis*, 2001, 1851-1855 (*synth, ir, pmr*)

Wang, X. *et al.*, *Synthesis*, 2007, 3044-3050 (*synth, ir, pmr*)

Zanthosin

Z-6

3-(1,3-Benzodioxol-5-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-propenamide. 3,4-Methylenedioxybenzoic acid 3,4-dimethoxyphenethylamide

C₂₀H₂₁NO₅ 355.39

Alkaloid from the stem bark of *Zanthoxylum rubescens* (Rutaceae). Needles. Mp 159-161°.

N-Me: **Zanthosinamide**

[119060-90-1]

C₂₁H₂₃NO₅ 369.416Alkaloid from the pericarps of *Zanthoxylum rubescens* (Rutaceae).3'-Demethoxy: **Armatamide**

[220458-86-6]

C₁₉H₁₉NO₄ 325.363Alkaloid from the bark of *Zanthoxylum armatum*. Cryst. Mp 159.4°. λ_{max} 202 ; 226 ; 288 ; 320 (MeOH).3'-Demethoxy, N-Me: **Podocarpamide**

[121880-09-9]


C₂₀H₂₁NO₄ 339.39Alkaloid from the bark of *Zanthoxylum podocarpum*, other *Zanthoxylum* spp. and *Esenbeckia leiocarpa*.Bis(demethoxy), N-Me: **Demethoxy-podocarpamide**

[132113-92-9]

C₁₉H₁₉NO₃ 309.364Isol. from *Esenbeckia leiocarpa*. Mp 139-141°.Adesina, S.K. *et al.*, *Pharmazie*, 1988, **43**, 517 (*Zanthosinamide*)Adesina, S.K. *et al.*, *Planta Med.*, 1989, **55**, 324 (*Zanthosin*)Ren, L.J. *et al.*, *Yaoxue Xuebao*, 1989, **24**, 67; *CA*, **111**, 74777r (*Podocarpamide*)Xie, J.X. *et al.*, *CA*, 1990, **111**, 142968 (*Podocarpamide*, *isol*, *synth*)Delle Monache, F. *et al.*, *Gazz. Chim. Ital.*, 1990, **120**, 387 (*isol*, *pmr*, *cmr*)Kalia, N.K. *et al.*, *J. Nat. Prod.*, 1999, **62**, 311-312 (*Armatamide*)**Zanthoxylines**

Z-7

[202002-14-0]

C₁₉H₁₃NO₄ 319.316Alkaloid from the bark of *Zanthoxylum rhoifolium*. Cryst. (CHCl₃/Et₂O). Mp 220-222°.Me ether, N-Me: **Broussonpapyrine**


[942137-65-7]

C₂₁H₁₈NO₄⁺ 348.378Quaternary alkaloid from the fruit of *Broussonetia papyrifera*. Yellow needles (CHCl₃/MeOH). Mp 201-203°. Counterion not specified.De Moura, N.F. *et al.*, *Phytochemistry*, 1997, **46**, 1443-1446 (*Zanthoxylines*)Pang, S.-Q. *et al.*, *Chem. Nat. Compd. (Engl. Transl.)*, 2007, **43**, 100-102 (*Broussonpapyrine*)**Zapotidine**

Z-8


6-Methylimidazo[1,5-c]tetrahydropyrimidine-5-thione

[525-84-8]

C₇H₉N₃S 167.234Minor alkaloid from the seeds of *Casimiroa edulis* (Mexican apple) (Rutaceae). Shows little pharmaceutical activity. Plates (Et₂O). Mp 96-98°.▶ LD₅₀ (mus, ipr) 50 mg/kg.*Picrate*: Mp 195-196°.Kincl, F.A. *et al.*, *J.C.S.*, 1956, 4163 (*isol*, *uv*, *ir*)Mechoulam, R. *et al.*, *J.A.C.S.*, 1961, **83**, 2022 (*pmr*, *struct*)Mechoulam, R. *et al.*, *Tetrahedron*, 1967, **23**, 239 (*synth*)Wróbel, J.T. *et al.*, *Alkaloids (Academic Press)*, 1985, **26**, 56 (*props*, *synth*)**Zascanol epoxide**

Z-9

[149998-44-7]



Relative configuration

C₁₇H₂₁NO₅ 319.357Alkaloid from roots of *Zanthoxylum scandens* (Rutaceae). Oil.Brader, G. *et al.*, *Annalen*, 1993, 355 (*isol*, *uv*, *ir*, *pmr*, *ms*, *cd*, *struct*)**Zeagenin**


Z-10

C₁₁H₂₀N₂O₅ 260.289Struct. unknown. Isol. from *Gibberella zeae*. Lowers blood lipoprotein levels in mice. Cryst. (MeOH). Mp 169-171°. [α]_D²⁵ +4.3 (c, 2.9 in H₂O).*Japan. Pat.*, 1970, 70 21 634; *CA*, **73**, 108242b**Zeatin**

Z-11

2-Methyl-4-(1H-purin-6-ylamino)-2-buten-1-ol, 9CI. 6-(4-Hydroxy-3-methyl-2-butenylamino)purine. N⁶-(ω-Hydroxyisopentenyl)adenine

[13114-27-7]



(E)-form

C₁₀H₁₃N₅O 219.246**(E)-form****trans-Zeatin**

[1637-39-4]

Isol. from sweet corn *Zea mays* (Poaceae) and numerous other plants. Also from algae, bacteria and basidiomycetes. Induces cell division; most effective of all known natural cytokinins. Sol. H₂O. Mp 212-213° (207-208°). λ_{max} 212 (ε 17050); 270 (ε 16150) (H₂O) (Berdy). λ_{max} 207 (ε 14500); 275 (ε 14650) (HCl) (Berdy). λ_{max} 220 (ε 15900); 276 (ε 14650) (NaOH) (Berdy).

▶ EM9506000

Picrate: Mp 180-190°.

O-β-D-Glucopyranoside: [56329-06-7]

C₁₆H₂₃N₅O₆ 381.388Isol. from *Lupinus angustifolius* and other plants. Cryst. (1-butanol). Mp 124-128°.

O-Ac: Mp 168-169°.

7-β-D-Glucopyranosyl: **Raphanatin**

[38165-56-9]

C₁₆H₂₃N₅O₆ 381.388Prod. by *Raphanus sativus* (radish) and crown galls of *Catharanthus roseus*. Cytokinin. Cryst. (MeOH). Mp 266-271° (dec.). [α]_D²⁶ -14.8 (c, 3.5 in DMSO).

9-β-D-Glucopyranosyl: [51255-96-0]

C₁₆H₂₃N₅O₆ 381.388Cytokinin, widely distributed in plants. λ_{max} 269 (no solvent reported).9-β-D-Ribofuranosyl: 9-(β-D-Ribofuranosyl)zeatin. **Ribosylzeatin**

[6025-53-2]

C₁₅H₂₁N₅O₅ 351.361Isol. from many plants incl. bacteria. Cytokinin, phytotoxin, plant hormone. Sol. H₂O. Mp 180-182°. λ_{max} 265 (ε 19200) (pH 1 H₂O) (Derep). λ_{max} 269 (ε 19400) (pH 13 H₂O) (Derep). λ_{max} 270 (ε 18400) (95% EtOH) (Derep).9-β-D-Ribofuranosyl, 5'-phosphate: **Ribosylzeatin phosphate**

[25615-16-1]

[15075-52-2]

C₁₅H₂₂N₅O₈P 431.341Isol. from *Zea mays* (sweet corn). Cytokinin. Solid.

9-β-D-Ribofuranosyl, O-β-D-glucopyranoside: [62512-97-4]

C₂₁H₃₁N₅O₁₀ 513.503

Cytokinin, widely distributed in plants. No phys. props. accessible.

9-β-D-Ribofuranosyl, O-[α-L-arabinopyranosyl-(1→4)]-[β-D-galactopyranosyl-(1→2)]-α-D-galactopyranosyl-(1→3)-α-L-arabinofuranosyl-(1→3)]-β-D-galactopyranoside]: **Gazer**

[171628-14-1]

C₄₃H₆₇N₅O₂₈ 1102.019Constit. of coconut milk (*Cocos nucifera*). Cytokinin. [α]_D²⁵ -5.2 (c, 1 in H₂O). λ_{max} 210 (ε 16600); 267 (ε 18200) (H₂O).9-(2-Deoxy-β-D-ribofuranosyl): **Zeatin 2-deoxyribofuranoside**. 2'-Deoxyzeatin riboside

[124325-91-3]

C₁₅H₂₁N₅O₄ 335.362Prod. by *Pseudomonas amygdali*. Cytokinin. Inhibitor of DNA polymerase I. Oil. [α]_D²⁵ -13.5 (c, 0.35 in EtOH).2',3'-S-Dihydro: 2-Methyl-4-(1H-purin-6-ylamino)-1-butanol, 9CI. 6-[(4-Hydroxy-3-methylbutyl)amino]purine. **Dihydrozeatin**

[19272-91-4]

[23599-75-9 (general no.)]

C₁₀H₁₅N₅O 221.261Cytokinin from immature *Lupinus luteus* seeds (Fabaceae). Constit. of numerous plant spp. Present in the brown alga *Sargassum heterophyllum*. Prisms (EtOH/MeCN). Mp 165-166° (154-156°). [α]_D²¹ -12.1 (MeOH). [α]_D²² -

15.4 (50% EtOH aq.).

2',3'-Dihydro, picrate:

Yellow needles (EtOH). Mp 187-189°.

2',3'-Dihydro, O-β-D-glucopyranoside:

Dihydrozeatin O-glucoside

[62512-96-3]

C₁₆H₂₅N₅O₆ 383.403

Constit. of various plant spp. incl.

Alnus spp., *Lupinus* sp. and *Phaseolus* sp.

2',3'-Dihydro, 9-β-D-ribofuranosyl: Dihydrozeatin riboside

[22663-55-4]

C₁₅H₂₃N₅O₅ 353.377

Constit. of numerous plants esp. in the Fabaceae. Prod. by *Pseudomonas* sp. Cytokinin. Sol. H₂O.

Deoxy: N-(3-Methyl-2-butenyl)-1H-purin-6-amine. 6-(3-Methyl-2-butenylamino)purine. N⁶-Prenyladenine. 6-(γ,γ-Dimethylallylamino)purine. N⁶-(3-Methyl-2-butenyl)adenine. N⁶-Isopentenyladenine. 2iP. Bryokinin

[2365-40-4]

C₁₀H₁₃N₅ 203.246

Widespread in plant spp. Prod. by *Corynebacterium fascians* and *Pseudomonas amygdali* and prob. present in brown alga *Sargassum heterophyllum*. Potent cytokinin. Mp 208-209° Mp 237-239° dec. Appears to be difficult to purify. λ_{max} 269 (EtOH). λ_{max} 218; 275 (0.1M NaOH).

2-Hydroxy: 2-Hydroxyzeatin

[29736-32-1]

C₁₀H₁₃N₅O₂ 235.245

Prod. by *Pseudomonas syringae*. Cytokinin. Oil. Sol. H₂O. λ_{max} 270 (MeOH).

(Z)-form

cis-Zeatin

[32771-64-5]

A modified nucleoside present in tRNAs. Isol. from *Corynebacterium fascians*. At least 50 times less active than the (E)-form. Mp 206-208°.

9-β-D-Glucopyranosyl: cis-Zeatin 9-glucoside

[169565-72-4]

C₁₆H₂₃N₅O₆ 381.388

Isol. from *Solanum tuberosum* (potatoes).

9-β-D-Ribofuranosyl: [15896-46-5]

Cryst. (EtOH). Mp 202-205°. λ_{max} 267 (ε 20280) (no solvent reported).

Shaw, G. et al., *J.C.S. (C)*, 1966, 921 (synth, uv, ir)

Letham, D.S. et al., *Life Sci.*, 1966, 5, 1999-2004 (isol, ribosylzeatin phosphate)

Klambt, D. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1966, 56, 52 (isol, deriv)

Helgeson, J.P. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1966, 56, 60 (deriv)

Koshimizu, K. et al., *Tet. Lett.*, 1967, 8, 1317-1320 (-)-Dihydrozeatin, isol, (synth)

Letham, D.S. et al., *Tetrahedron*, 1967, 23, 479 (ms, uv, pmr, struct)

Hecht, S.M. et al., *Phytochemistry*, 1970, 9, 1173-1180 (2-Hydroxyzeatin)

Playtis, A.J. et al., *Biochem. Biophys. Res. Commun.*, 1971, 45, 1-5 (synth, 9-ribosyl)

Leonard, N.J. et al., *J.A.C.S.*, 1971, 93, 3056 (synth, uv, pmr, ms)

Bugg, C.E. et al., *Biochem. Biophys. Res. Commun.*, 1972, 46, 779-784 (deoxy, cryst struct)

Corse, J. et al., *Synthesis*, 1972, 11, 618 (synth, pmr)

Fujii, T. et al., *Tet. Lett.*, 1972, 13, 3075-3078 (-)-Dihydrozeatin, (synth)

Parker, C.W. et al., *Biochem. Biophys. Res. Commun.*, 1973, 55, 1370-1376 (synth, 9-glucoside)

Parker, C.W. et al., *Planta*, 1973, 114, 199-218 (*Raphanatin*, isol, struct, metab)

Scarborough, E. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1973, 70, 3825 (cis-Zeatin)

Parker, C.W. et al., *Planta*, 1974, 115, 337-344 (isol, 9-glucosyl)

Parker, C.W. et al., *Ann. Bot. (London)*, 1975, 39, 375-376 (O-glucoside)

Letham, D.S. et al., *Ann. Bot. (London)*, 1976, 41, 261-263 (9-ribosyl O-glucoside)

Mackenzie, G. et al., *J.C.S. Perkin I*, 1976, 1446 (synth)

Matsubara, S. et al., *Phytochemistry*, 1977, 16, 933-937 (-)-Dihydrozeatin, Dihydrozeatin riboside, (synth)

Wang, T.L. et al., *Planta*, 1977, 135, 285; 1978, 140, 151 (glycosides)

Cowley, D.E. et al., *Aust. J. Chem.*, 1978, 31, 1095-1111 (7-glucopyranosyl, 9-glucopyranosyl)

Duke, C.C. et al., *Aust. J. Chem.*, 1978, 31, 1291-1301; 2219 (O-glucopyranoside, isol, synth, uv, pmr, cmr)

Yokota, T. et al., *Phytochemistry*, 1980, 19, 2367-2369 (deoxy, isol)

Chen, S.C. et al., *Agric. Biol. Chem.*, 1982, 46, 2361 (cmr)

Corse, J. et al., *J. Plant Growth Regul.*, 1983, 2, 47; *CA*, 100, 34316 (synth, resoln)

Evidente, A. et al., *Phytochemistry*, 1989, 28, 2603-2607 (2-deoxyribosyl, Dihydrozeatin, trans-Zeatin, isol)

De Napoli, L. et al., *Phytochemistry*, 1990, 29, 701-709 (2-deoxyribosyl, synth)

Shadid, B. et al., *Tetrahedron*, 1990, 46, 901-912 (synth, ribosylzeatin phosphate)

Evidente, A. et al., *Chem. Pharm. Bull.*, 1992, 40, 1937 (synth, 2-deoxyribosyl)

Nicander, B. et al., *Plant Physiol.*, 1995, 109, 513 (9-glucoside)


Kobayashi, H. et al., *Chem. Pharm. Bull.*, 1997, 45, 260 (Gazer)

Jordis, U. et al., *Org. Prep. Proced. Int.*, 1997, 29, 549-560 (synth, pmr)

Tsai, Y.-H. et al., *Helv. Chim. Acta*, 2003, 86, 2452-2457 (deoxy, isol)

Zefbetaine

[105708-78-9]



C₁₆H₁₃NO₃ 267.284

Alkaloid from the fresh, mature seeds of *Zephyranthes flava* (Amaryllidaceae). Brown powder (EtOH). Sol. MeOH, CHCl₃; poorly sol. H₂O. Mp 300°. λ_{max} 262 (ε 50120); 408 (ε 6165) (MeOH) (Berdy). λ_{max} 263; 278; 283; 378 (MeOH/HCl) (Berdy). λ_{max} 265; 458 (MeOH/NaOH) (Berdy).

Ac:

Straw-coloured solid (Me₂CO/petrol).

Z-12

Mp 222-225°.

Me ether: Criasbetaine

[103246-12-4]

C₁₇H₁₅NO₃ 281.31

Alkaloid from the fruits of *Crinum asiaticum*. Shows antitumour props. Yellow amorph. powder.

Fales, H.M. et al., *J.A.C.S.*, 1956, 78, 4145 (*Criasbetaine*, synth)

Ghosal, S. et al., *J. Chem. Res., Synop.*, 1986, 112 (*Criasbetaine*)


Ghosal, S. et al., *Phytochemistry*, 1986, 25, 1975 (isol, uv, ir, pmr, ms, struct)

Stark, L.M. et al., *J.O.C.*, 2000, 65, 3227-3230 (*Criasbetaine*, synth)

Zelkovamycin

[221197-33-7]

Z-13



C₃₆H₄₅N₉O₉S 779.872

Cyclic peptide antibiotic. Possible identity with Antibiotic CP 21635.

Prod. by *Streptomyces* sp. K96-0670.

Antibacterial agent. Powder. [α]_D²⁸ +84.3 (c, 0.1 in MeOH). λ_{max} 205 (ε 57800); 222 (ε 59300); 244 (sh) (ε 24000); 284 (ε 6000); 292 (ε 5300) (MeOH).

Zhang, H. et al., *J. Antibiot.*, 1999, 52, 29-33; 34-39 (isol, uv, ir, pmr, cmr, ms)

Martin, G.E. et al., *J. Nat. Prod.*, 2000, 63, 543-585 (N-15 mm)

Zenkerene


[56053-17-9]

Z-14

16,17,19,20-Tetrahydrocuran, 9CI.

Zenkerine†

[56053-17-9]



Absolute Configuration

C₁₉H₂₂N₂ 278.396

Alkaloid from the root bark of *Strychnos zenkeri*.

N⁴-Oxide: **Zenkerene N^b-oxide.**

Zenkerine N^b-oxide

[197390-41-3]


C₁₉H₂₂N₂O 294.396

Alkaloid from *Strychnos zenkeri*.

Thepenier, P. et al., *Bull. Soc. R. Sci. Liege*, 1997, 66, 341-344; *CA*, 127, 305384j

Zephyranthine

Trianthine. Ungminoridine

C₁₆H₁₉NO₄ 289.33

Historically, Zephyranthine was the (-)-form, Trianthine the (+)-form and Ungminoridine the (±)-form.

(+)-form [84236-20-4]Alkaloid from *Pancratium trianthum*. Cryst. (MeOH). Mp 205-206° (179-180°). [α]_D²⁵ +51.2.**(-)-form** [2030-55-9]Alkaloid from *Zephyranthes candida* (Amaryllidaceae). Cryst. + 1/2 H₂O. Mp 201-202° dec. [α]_D²⁴ -43.17 (c, 0.47 in CHCl₃).*Picrate*: Mp 194-195° dec.*Methiodide*: Mp 294-295° dec.*Di-Ac*: Mp 68-70°.**2-Epimer: Dihydrolycorine**C₁₆H₁₉NO₄ 289.33Alkaloid from *Pancratium maritimum* (Amaryllidaceae).

Protozoicide which has been used against amoebic dysentery. Shows antifungal activity. Mp 255° (247°) dec.

2-Epimer, di-Ac: Nartazine. Diacetyldihydrolycorine


[475-90-1]

C₂₀H₂₃NO₆ 373.405Alkaloid from the bulbs of *Narcissus tazetta*. Also occurs in *Galanthus nivalis* (Amaryllidaceae). Long needles (Me₂CO). Mp 185-186°. [α]_D²⁵ -120 (c, 0.25 in CHCl₃).**(±)-form** [21851-15-0]Alkaloid from *Ungernia minor* and *Ungernia vvedenskyi* (Amaryllidaceae). Mp 193-194° Mp 224-225°.*Di-Ac*:Cryst. (Me₂CO). Mp 143-145°.Boit, H.-G. *et al.*, *Chem. Ber.*, 1956, **89**, 2462 (*Nartazine*)Nakagawa, Y. *et al.*, *J.C.S.*, 1959, 3736 (*Dihydrolycorine, stereochem*)Boit, H.-G. *et al.*, *Naturwissenschaften*, 1960, **47**, 109 (*Nartazine*)Ozeki, S. *et al.*, *Chem. Pharm. Bull.*, 1964, **12**, 253 (*uv, ir, struct*)Shiro, M. *et al.*, *Chem. Ind. (London)*, 1966, 1229 (*Dihydrolycorine, cryst struct*)Kotera, K. *et al.*, *Tet. Lett.*, 1966, 2009 (*stereochem*)Sandberg, F. *et al.*, *Acta Pharm. Suec.*, 1968, **5**, 61; *CA*, **69**, 25043d (*Dihydrolycorine, isol*)Abduazimov, K.A. *et al.*, *Khim. Prir. Soedin.*, 1968, **4**, 263; *Chem. Nat. Compd. (Engl. Transl.)*, 1968, **4**, 227 (*isol, uv, pmr, ms, struct*)Miyakado, M. *et al.*, *Phytochemistry*, 1975, **14**, 2717 (*Dihydrolycorine, activity*)Yamaki, M. *et al.*, *Heterocycles*, 1976, **5**, 163 (*synth*)

Z-15

Tsuda, Y. *et al.*, *J.C.S. Perkin 1*, 1979, 1358 (*synth*)Kadyrov, K.A. *et al.*, *Khim. Prir. Soedin.*, 1979, **15**, 418; *Chem. Nat. Compd. (Engl. Transl.)*, 1979, **15**, 370 (*isol*)Frederik, D.M. *et al.*, *Khim. Prir. Soedin.*, 1982, **18**, 534; *Chem. Nat. Compd. (Engl. Transl.)*, 1992, **18**, 510-511 (*Trianthine*)Oppolzer, W. *et al.*, *J.A.C.S.*, 1994, **116**, 3139-3140 (*Trianthine, synth*)Herrera, M.R. *et al.*, *Fitoterapia*, 2001, **72**, 444-448 (*isol, pmr, cmr*)Wang, Q. *et al.*, *Org. Lett.*, 2004, **6**, 2189-2192 (*synth*)**Zeraconine**

[111313-35-0]

C₃₀H₄₀N₂O 444.659Alkaloid from *Aconitum zeraconicum* (Ranunculaceae). Mp 130-131°.**N-Oxide: Zeraconine N-oxide**


[111261-81-5]

C₃₀H₄₀N₂O₂ 460.658Alkaloid from *Aconitum zeraconicum* (Ranunculaceae). Mp 94-95°.Vaisov, Z.M. *et al.*, *Khim. Prir. Soedin.*, 1987, 407; *Chem. Nat. Compd. (Engl. Transl.)*, 337**Zeravschanidine**

Z-17

Struct. unknown. Alkaloid from the above-ground parts of *Aconitum zeravschanicum* collected during budding and blooming (Ranunculaceae). Amorph. Mp 85°.*Hydrochloride*: Mp 271°. [α]_D +30.61.Progressov, N.N. *et al.*, *Dokl. Akad. Nauk UzSSR*, 1953, **No. 11**, 19-21; *CA*, **51**, 1539h**Zetekitoxin AB**


Z-18

Atelopidtoxin
[62996-38-7]

Relative Configuration


C₁₆H₂₄N₈O₁₂S 552.478Related to Saxitoxin, S-109. Toxin from the golden frog *Atelopus zeteki* of Panama. Extremely potent sodium channel blocker.Yotsu-Yamashita, M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2004, **101**, 4346-4351 (*isol, pmr, cmr, N-15 nmr, ms*)**Zijinlongine**

Z-19

4,6,7,13-Tetrahydro-12-methoxy-12bH-bis[1,3]benzodioxolo[5,6-a:4',5'-g]quinolizine-12b-methanol, 9CI
[133086-83-6]C₂₁H₂₁NO₆ 383.4Alkaloid from the whole plant of *Dactylicapnos torulosa* (Papaveraceae). Cryst. Mp 166-167°.Zhang, G.L. *et al.*, *Yaoxue Xuebao*, 1990, **25**, 604-607 (*isol*)**Zippeline**

Z-20

[123750-35-6]



Relative Configuration

C₂₀H₂₇NO₄ 345.438Alkaloid from aerial parts of *Stephania zippeliana* (Menispermaceae). [α]_D +20 (c, 0.60 in CHCl₃).Charles, B. *et al.*, *Can. J. Chem.*, 1989, **67**, 1257**Zizyphine I**


Z-21

[159509-34-9]

As Zizyphine A, Z-22 with

R = -CH(CH₃)CH₂CH₃, R' = -CH₂PhC₃₆H₄₇N₅O₆ 645.797Stereochem. not determined. Alkaloid from bark of *Zizyphus oenoplia* (Rhamnaceae). Mp 135°.Khokhar, I. *et al.*, *Pak. J. Sci.*, 1993, **45**, 54; *CA*, **122**, 5434s (*isol, pmr, ms*)**Zizyphine A**

Z-22

Zizyphine[†]
[51059-42-8]

Absolute Configuration

R = R' = CH(CH₃)CH₂CH₃ (S-)C₃₃H₄₉N₅O₆ 611.78

Alkaloid from the root bark and stem bark of *Zizyphus oenoplia* and from *Zizyphus jujuba* (Rhamnaceae). Cryst. (CHCl₃/petrol). Mp 124-126°. [α]_D²⁰ -411 (c, 0.086 in CHCl₃).

Hydrochloride: Mp 184-184.5°.

N-De-Me: **Zizyphine B**. Zizyphinine [24604-79-3]

C₃₂H₄₇N₅O₆ 597.753

Alkaloid from root and stem bark of *Zizyphus oenoplia* (Rhamnaceae). Amorph. [α]_D²⁰ -457 (c, 1 in CHCl₃). Revised struct. is consequent on the revised struct. of Zizyphine A.

N-De-Me; hydrochloride:

Cryst. + 1H₂O. Mp 242-243° dec.

O-De-Me: **Zizyphine F**

[55839-64-0]

C₃₂H₄₇N₅O₆ 597.753

Alkaloid from stem bark of *Zizyphus oenoplia* and *Zizyphus spina-christi* (Rhamnaceae). Needles (Et₂O). Mp 235°. [α]_D²⁰ -277 (c, 0.15 in MeOH).

Ménard, E.L. *et al.*, *Helv. Chim. Acta*, 1963, **46**, 1801 (*isol, uv, ir, pmr*)

Zbiral, E. *et al.*, *Helv. Chim. Acta*, 1965, **48**, 404 (*uv, ir, pmr, ms*)

Tschesche, R. *et al.*, *Tet. Lett.*, 1973, 2577; 1974, 2941 (*ms, struct, Zizyphine F*)

Cassels, B.K. *et al.*, *Tetrahedron*, 1974, **30**, 2461 (*isol*)

Hindenlang, D.M. *et al.*, *Annalen*, 1980, 447 (*cmr*)

Schmidt, U. *et al.*, *Angew. Chem., Int. Ed.*, 1981, **20**, 1026 (*synth, ms*)

Haslinger, E. *et al.*, *Monatsh. Chem.*, 1982, **113**, 95 (*pmr, cmr*)

Schmidt, U. *et al.*, *J.O.C.*, 1983, **48**, 2680 (*synth, cd, pmr, ms*)

Zizyphine C

Z-23

[55323-67-6]

As Zizyphine A, Z-22 with

R = -CH₂Ph, R' = -CH(CH₃)CH₂CH₃

C₃₆H₄₇N₅O₆ 645.797

Alkaloid from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Amorph. [α]_D -331 (c, 0.10 in CHCl₃). [α]_D -343 (c, 1.0 in MeOH).


Cassels, B.K. *et al.*, *Tetrahedron*, 1974, **30**, 2461 (*isol, uv, ir, pmr, ms, struct*)

Zizyphine G

Z-24

1-(2-Amino-3-methyl-1-oxopentyl)-2,3,3a,12a,13,14,15,17a-octahydro-5,8-ethenodipyrrolo[3,2-b:1',2'-e][1,5,8]-oxadiazacyclotetradecine-12,17(1H,11H)-dione, 9CI

[55839-65-1]



Absolute Configuration

C₂₄H₃₂N₄O₄ 440.541

Alkaloid from the stem bark of *Zizyphus oenoplia* (Rhamnaceae). Cryst. (Me₂CO).

Mp 130°. [α]_D²⁰ -185 (c, 0.19 in MeOH).

Tschesche, R. *et al.*, *Tet. Lett.*, 1974, **15**, 2941-2944 (*isol, pmr, ms, struct*)

Zizyphine K

Z-25

[164176-41-4]

As Zizyphine A, Z-22 with

R = CH(CH₃)₂, R' = CH(CH₃)CH₂CH₃, O-de-Me

C₃₁H₄₅N₅O₆ 583.726

Alkaloid from *Zizyphus oenoplia*. Mp 230°.

Khokhar, I. *et al.*, *J. Nat. Sci. Math.*, 1994, **34**, 171-175; *CA*, **123**, 29541w

Zizyphine N

Z-26

As Zizyphine A, Z-22 with

R = -CH(CH₃)CH₂CH₃(S-), R' = -CH₂CH(CH₃)₂

C₃₃H₄₉N₅O₆ 611.78

Alkaloid from the roots of *Zizyphus oenoplia* var. *brunoniana*. Shows antimicrobial and antiplasmodial activity. Mp 117-119°. [α]_D³⁰ -326.6 (c, 0.18 in CHCl₃). λ _{max} 266 (log ϵ 4.4); 319 (log ϵ 4.28) (EtOH).

N-De-Me: **Zizyphine O**

C₃₂H₄₇N₅O₆ 597.753

Alkaloid from the roots of *Zizyphus oenoplia* var. *brunoniana*. Mp 106-108°. [α]_D³¹ -380.2 (c, 0.15 in CHCl₃). λ _{max} 268 (log ϵ 4.01); 319 (log ϵ 3.89) (EtOH).

O-De-Me: **Zizyphine P**

C₃₂H₄₇N₅O₆ 597.753

Alkaloid from the roots of *Zizyphus oenoplia* var. *brunoniana*. Mp 127-129°. [α]_D³¹ -385.4 (c, 0.15 in CHCl₃). λ _{max} 265 (log ϵ 4.73); 321 (log ϵ 4.59) (MeOH).

Suksamrarn, S. *et al.*, *Tetrahedron*, 2005, **61**, 1175-1180 (*isol, pmr, cmr, ms*)

He, G. *et al.*, *Org. Lett.*, 2007, **9**, 1367-1369 (*synth*)

Zizyphine Q

Z-27

As Zizyphine A, Z-22 with

R = -CH(CH₃)CH₂CH₃(S-), R' = -CH(CH₃)₂

C₃₂H₄₇N₅O₆ 597.753

Alkaloid from the roots of *Zizyphus oenoplia* var. *brunoniana*. Shows antimicrobial and antiplasmodial props. Mp 140-142°. [α]_D²⁹ -345 (c, 0.16 in CHCl₃). λ _{max} 265 (log ϵ 3.64); 319 (log ϵ 3.5) (MeOH).

Suksamrarn, S. *et al.*, *Tetrahedron*, 2005, **61**, 1175-1180 (*isol, pmr, cmr, ms*)

Zoamides

Z-28



R¹

R²

Zoamide A:	-COCH=C(CH ₃) ₂	-COCH=C(CH ₃) ₂
B:	-COPh	-COCH=C(CH ₃) ₂
C:	-COPh	-COCH(CH ₃) ₂
D:	-COPh	-COCH(CH ₃)CH ₂ CH ₃ (S ⁻)

Related to Palyzoanthoxanthins in P-94.

Zoamide A [188558-01-2]

C₂₀H₂₆N₆O₂ 382.464

Alkaloid from an undescribed marine zoanthid, *Parazoanthus* sp., collected in the Philippines. Gummy solid. [α]_D 0. λ _{max} 200 (ϵ 20500); 290 (ϵ 18500) (MeOH).

Zoamide C [188558-97-6]

C₂₁H₂₄N₆O₂ 392.46

From *Parazoanthus* sp. Gummy solid. [α]_D 0. λ _{max} 225 (ϵ 19600); 270 (ϵ 13600) (MeOH).

Zoamide B [188558-50-1]

C₂₂H₂₄N₆O₂ 404.471

From *Parazoanthus* sp. Solid. [α]_D 0. λ _{max} 220 (ϵ 29000); 290 (ϵ 23500) (MeOH).

Zoamide D [188559-04-8]

C₂₂H₂₆N₆O₂ 406.486


From *Parazoanthus* sp. Gummy solid. [α]_D +15.1. λ _{max} 230 (ϵ 17000); 270 (ϵ 14700) (MeOH).

D'Ambrosio, M. *et al.*, *Tet. Lett.*, 1997, **38**, 717 (*isol, uv, ir, pmr, cmr, struct*)

Zoanthamide

Z-29

[97877-70-8]



C₃₀H₃₇NO₇ 523.625


Alkaloid from an unidentified marine zoanthid, *Zoanthus* sp. Shows antiinflammatory props. Mp 278-280°. [α]_D +133 (c, 0.83 in CHCl₃). λ _{max} 235 (ϵ 23900) (MeCN) (Derep).

Rao, C.B. *et al.*, *J.O.C.*, 1985, **50**, 3757 (*isol, uv, ir, pmr, cmr, struct*)

Zoanthamine

Z-30

[93426-90-5]



$C_{30}H_{41}NO_5$ 495.658
Alkaloid from a marine zoanthid, *Zoanthus* sp. Cytotoxic agent, antiinflammatory. Interleukin 6 production inhibitor. Mp 306-308°. $[\alpha]_D^{25} +18$ (c, 0.48 in $CHCl_3$). λ_{max} 233 (ϵ 11000) (MeCN) (Derep).

13,14,17,18-Tetradehydro, 17-alcohol:

Zoanthenol

[231945-73-6]

$C_{30}H_{39}NO_5$ 493.642

Alkaloid from a zoanthid. Amorph. solid. $[\alpha]_D^{25} +7.1$ (c, 0.24 in $CHCl_3$).

11-Oxo: **Zoanthenone**

[123853-69-0]

$C_{30}H_{39}NO_6$ 509.641

Alkaloid from a marine zoanthid, *Zoanthus* sp. Cryst. $[\alpha]_D +30$ (c, 0.1 in $CHCl_3$). Mp not recorded. λ_{max} 235 (ϵ 23900) (MeCN) (Derep).

11 β -Hydroxy: **11-Hydroxyzoanthenamine**

[231945-77-0]

$C_{30}H_{41}NO_6$ 511.657

Alkaloid from a zoanthid. Amorph. solid. $[\alpha]_D^{25} +1.8$ (c, 0.23 in $CHCl_3$).

16 β -Hydroxy, 17-deoxo, 15 β ,16-dihydro:

Zoosanthellamine

[208256-91-1]

$C_{30}H_{45}NO_5$ 499.689

Alkaloid from cultures of the dinoflagellate *Symbiodinium* sp. Solid. Mp > 300°. $[\alpha]_D^{20} +40$ (c, 0.05 in MeOH).

26-Hydroxy: **26-Hydroxyzoanthenamine**.

Oxyzoanthenamine

[159509-36-1]

$C_{30}H_{41}NO_6$ 511.657

Alkaloid from a *Zoanthus* sp. Cytotoxic agent. Interleukin 6 production inhibitor. Oil.

19-Epimer, 26-hydroxy: **26-Hydroxy-19-epizoanthenamine**.

Epioxyzoanthenamine

[211049-29-5]

$C_{30}H_{41}NO_6$ 511.657

Alkaloid from a *Zoanthus* sp. Amorph. solid. $[\alpha]_D^{25} -17.5$ (c, 0.24 in $CHCl_3$).

$A^{15,27}$ -Isomer, 16 β -hydroxy, 17-deoxo:

Loboanthenamine

$C_{30}H_{43}NO_5$ 497.673

Alkaloid from a *Lobophytum* sp. Amorph. solid. $[\alpha]_D^{25} +26$ (c, 0.15 in $CHCl_3$).

Rao, C.B. et al., *J.A.C.S.*, 1984, **106**, 7983-7984

(*uv*, *ir*, *pmr*, *cmr*, *cryst struct*)

Atta-ur-Rahman, et al., *Tet. Lett.*, 1989, **30**, 6825-6828 (*Zoanthenone*)

Fukuzawa, S. et al., *Heterocycl. Commun.*, 1995, **1**, 207-214 (*Oxyzoanthenamine*)

Nakamura, H. et al., *Bull. Chem. Soc. Jpn.*, 1998, **71**, 781-787 (*Zoosanthellamine*)

Daranas, A.H. et al., *Tetrahedron*, 1998, **54**, 7891-7896; 1999, **55**, 5539-5546


(*Epioxyzoanthenamine*, 11-Hydroxyzoanthenamine, *Zoanthenol*)

Behenna, D.C. et al., *Angew. Chem., Int. Ed.*, 2008, **47**, 2365-2386 (*rev. biol. chem*)

Fattorusso, E. et al., *Tet. Lett.*, 2008, **49**, 2189-2192 (*Loboanthenamine*)

Zoanthenamine

[97877-69-5]



$C_{30}H_{39}NO_6$ 509.641

The abs. config. is prob. as shown by analogy with Norzoanthenamine, N-317. Alkaloid from an unidentified marine zoanthid, *Zoanthus* sp. Shows anti-inflammatory props. Powder. Mp 238-240°. Slowly dec. in soln. λ_{max} 233 (ϵ 11000) (MeCN) (Derep).

20,28-Deepoxy, 20-ketone: **28-Deoxyzoanthenamine**

[120314-15-0]

$C_{30}H_{39}NO_5$ 493.642

Alkaloid from the unidentified colonial zoanthid of the genus *Zoanthus*. Potent antiinflammatory and analgesic agent. Mp 305-307°. $[\alpha]_D +216$ (c, 3.7 in $CHCl_3$). λ_{max} 238 (ϵ 16250) (MeOH) (Berdy).

22-Epimer, 20,28-deepoxy, 20-ketone: **28-Deoxy-22-epizoanthenamine**. 22-Epi-28-deoxyzoanthenamine

[120409-36-1]

$C_{30}H_{39}NO_5$ 493.642

Alkaloid from an unidentified *Zoanthus* spp. Antiinflammatory and analgesic agent. Oil. $[\alpha]_D +85$ (c, 2.36 in $CHCl_3$). λ_{max} 237 (ϵ 14200) (MeOH) (Berdy).


Rao, C.B. et al., *J.O.C.*, 1985, **50**, 3757 (*isol. uv, ir, pmr, cmr, struct*)

Rao, C.B. et al., *Heterocycles*, 1989, **28**, 103 (*Deoxyzoanthenamine*, 28-Deoxy-22-epizoanthenamine)

Zooanemonin

Z-32

4-(Carboxymethyl)-1,3-dimethyl-1H-imidazolium hydroxide inner salt, 9CI. 1,3-Dimethyl-4-imidazoleacetic acid betaine. *Anemonin*† [584-91-8]



$C_7H_{10}N_2O_2$ 154.168

Isol. from the mussel *Arca noae*, the sea anemone *Anemonia sulcata* and the sponge *Axinella verrucosa*. Antibacterial, antineoplastic agent. Extremely hygro-

scopic.

Hydrochloride: Mp 184° (178°).

Picrate: Mp 149-153°.


Kutscher, F. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1933, **221**, 38 (*isol*)

Ackermann, D. et al., *Hoppe-Seyler's Z. Physiol. Chem.*, 1953, **294**, 93; **295**, 1; 1954, **296**, 286; 1960, **318**, 281 (*synth. isol, nomencl. struct*)

Zopfiellamide A

Z-33

[478945-64-1]



$C_{25}H_{35}NO_6$ 445.555

Tetramic acid deriv. Incorr. MF given in the ref. Prod. by the marine fungus *Zopfiella latipes*. Yellowish cryst. (EtOH). Mp 225-230°. $[\alpha]_D^{22} +5.2$ (c, 1.1 in $CHCl_3$). λ_{max} 247 (ϵ 11000); 320 (ϵ 10000) (MeOH).

Daferner, M. et al., *Tetrahedron*, 2002, **58**, 7781-7784 (*isol, pmr, cmr*)

Zopfiellamide B

Z-34

[478945-65-2]

As Zopfiellamide A, Z-33 with

R = CH_2CH_3

$C_{26}H_{37}NO_6$ 459.581

Tetramic acid deriv. Prod. by the marine fungus *Zopfiella latipes*. Yellowish oil. $[\alpha]_D^{25} -24$ (c, 0.3 in $CHCl_3$). λ_{max} 247 (ϵ 3800); 321 (ϵ 3400) (MeOH).


Daferner, M. et al., *Tetrahedron*, 2002, **58**, 7781-7784 (*isol, pmr, cmr*)

Zygadenilic acid δ -lactone

Z-35

Zygadenine δ -lactone

[10060-03-4]



$C_{27}H_{41}NO_7$ 491.623


Alkamine from *Veratrum* spp., *isol. from Veratrum oxysepalum* after hydrol. Mp 252° dec. $[\alpha]_D^{15} -59.4$ (c, 1.70 in $CHCl_3$).

16-Angeloyl: Angeloylzygadenilic acid δ -lactoneC₃₂H₄₇NO₈ 573.725Alkaloid from the rhizomes of *Veratrum grandiflorum*. Monohydrate. Mp 235°. [α]_D¹⁵ -10.9 (c, 1.65 in CHCl₃). CAS no. not found to 2006.Tsukamoto, T. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 1102-1106 (*isol. struct*)Yagi, A. *et al.*, *Chem. Pharm. Bull.*, 1962, **10**, 519-521 (*struct, bibl*)**Zygadenine****Z-36**

4,9-Epoxy-3,4,14,15,16,20-cevanehexol.

7-Deoxygermine

[545-45-9]



Absolute Configuration

C₂₇H₄₃NO₇ 493.639Alkaloid from *Zygadenus intermedius* and several other *Zygadenus* spp. (Liliaceae). Shows antitumour activity. Mp 218-220° (201-204°, 214°). [α]_D²⁰ -48.4 (c, 1.26 in CHCl₃).

Hydrochloride: Mp 231-234°.

3-O- β -D-Glucopyranoside: Zygadenine 3-glucoside

[169907-14-6]

C₃₃H₅₃NO₁₂ 655.781Alkaloid from aerial parts of *Astragalus polycanthus*. Cryst. Mp 310-311°. [α]_D³⁰ -67 (c, 0.3 in MeOH).

3-Ac: Zygacine

[2777-79-9]

C₂₉H₄₅NO₈ 535.676Alkaloid from *Veratrum album*, *Zygadenus paniculatus*, *Zygadenus venenosus* and *Zygadenus gramineus*. Noncryst. [α]_D -22 (CHCl₃).

3-O-(2R-Methylbutanoyl): 2-Methylbutyrylzygadenine

[105765-19-3]

C₃₂H₅₁NO₈ 577.757Alkaloid from *Veratrum stamineum*. Mp 175°. [α]_D -7.8 (CHCl₃).

3-Angeloyl: Angeloylzygadenine

[67370-02-9]

C₃₂H₄₉NO₈ 575.741Alkaloid from *Veratrum stenophyllum*, *Veratrum maackii* and *Veratrum stamineum* (Liliaceae). Mp 222-224°. [α]_D -35 (CHCl₃).3-Angeloyl, β -N-oxide: Angeloylzygadenine β -N-oxide

[313677-61-1]

C₃₂H₄₉NO₉ 591.74Alkaloid from *Veratrum taliense*. Needles (MeOH). Mp 196-198°.

3-O-(4-Hydroxy-3-methoxybenzoyl): Vanilloylzygadenine

[67370-01-8]

C₃₅H₄₉NO₁₀ 643.773Alkaloid from *Zygadenus venenosus* and *Zygadenus paniculatus*. Mp 258-259°. [α]_D²³ -27.5 (CHCl₃).

3-O-(3,4-Dimethoxybenzoyl): Veratroylzygadenine


[31329-58-5]

C₃₆H₅₁NO₁₀ 657.8Alkaloid from *Zygadenus venenosus*, *Zygadenus paniculatus*, *Veratrum album* and several other *Veratrum* spp. Moderate hypotensive agent. Mp 278-280° dec. (270-271°). [α]_D -27 (CHCl₃).

▶ FL5660000

Heyl, F.W. *et al.*, *J.A.C.S.*, 1949, **71**, 1751-1752 (*isol*)Stoll, A. *et al.*, *Helv. Chim. Acta*, 1953, **36**, 1570-1575 (*Veratroylzygadenine*)Kupchan, S.M. *et al.*, *J.A.C.S.*, 1953, **75**, 1025-1029; 1959, **81**, 1925-1928(*Vallinoylzygadenine*, *Veratroylzygadenine*, *struct*)Kupchan, S.M. *et al.*, *J.A.C.S.*, 1955, **77**, 689-691 (*Zygacine*)Suzuki, M. *et al.*, *Yakugaku Zasshi*, 1959, **79**, 619-623 (*Angeloylzygadenine*)Tomko, J. *et al.*, *Planta Med.*, 1962, **10**, 138-142 (*Veratroylzygadenine*)Yagi, A. *et al.*, *Yakugaku Zasshi*, 1962, **82**, 210-213; *C.A.*, **58**, 5748b (2-Methylbutyrylzygadenine)Budzikiewicz, H. *et al.*, *Tetrahedron*, 1964, **20**, 2267-2268 (*ms*)Carey, F.A. *et al.*, *Org. Magn. Reson.*, 1980, **14**, 141-144 (*cmr*)Liang, G. *et al.*, *Yaoxue Xuebao*, 1984, **19**, 131-136 (*isol. Angeloylzygadenine*, *Veratroylzygadenine*)Gupta, R.K. *et al.*, *Indian J. Chem., Sect. B*, 1995, **34**, 1021-1022 (*glucoside*)Zhou, C.X. *et al.*, *Indian J. Chem., Sect. B*, 2000, **39**, 283-286 (*Angeloylzygadenine N-oxide*)**Zygosporin D****Z-37**

[25374-67-8]



Absolute Configuration

C₂₈H₃₅NO₅ 465.588Prod. by *Zygosporium masonii* and *Metarhizium anisopliae*. Antibiotic possessing antitumour and antiinflammatory props. Plates (MeOH). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 180-190°. [α]_D²⁵ -15.5 (c, 0.58 in dioxan). [α]_D +20.6 (c, 0.25 in EtOH).▶ LD₅₀ (mus, ipr) 10 mg/kg. HA5304000

21-Ac: Cytochalasin D. Zygosporin A

[22144-77-0]

C₃₀H₃₇NO₆ 507.625Metab. of *Engleromyces goetzii*, *Me-**tarhizium anisopliae* and *Hypoxyylon terricola*. Shows antibiotic and cytotoxic props. Inhibits lipid droplet formn. Specific cholesteryl ester synthesis inhibitor. Needles (Me₂CO/petrol). Mp 268-271°. [α]_D²⁵ -7.5 (c, 0.55 in dioxan). Log P 1.54 (uncertain value) (calc). λ_{\max} 286 (ε 330) (MeOH) (Derep).▶ LD₅₀ (rat, ipr) 0.9 mg/kg. Exp. teratogen. GZ4850000

7,21-Di-Ac: Zygospurin F

[25374-68-9]

C₃₂H₃₉NO₇ 549.663Minor metab. of *Zygosporium masonii*. Shows antibiotic props. similar to Zygosporin D. Prisms (diisopropyl ether). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 126-129°. [α]_D²⁴ -12 (c, 0.78 in dioxan).

21-Ketone: 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalas-6(12),13,19-triene-1,17,21-trione

C₂₈H₃₃NO₅ 463.572From *Daldinia* sp. Amorph. solid. [α]_D -38.1 (c, 0.05 in CHCl₃).19 α ,20 β -Epoxide, 21-Ac: 19,20-Epoxy-cytochalasin DC₃₀H₃₇NO₇ 523.625C-19 stereochem. revised in 2007. Metab. of *Engleromyces goetzii* and *Xylaria hypoxylon*. Cytotoxic. [α]_D -228 (c, 0.03 in CHCl₃).

18-Deoxy, 21-Ac: Zygosporin E

[26399-27-9]

[114651-74-0]

C₃₀H₃₇NO₅ 491.626Minor metab. of the fungus *Zygosporium masonii*. Shows antibiotic props. similar to Zygosporin D. Needles (Me₂CO/hexane). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 218-223.5°. [α]_D²⁴ +6.2 (c, 0.97 in dioxan).

18-Deoxy, di-Ac:

Amorph. powder. Mp 86-89°.

17-Deoxo: Cytochalasin J. Deacetylcytochalasin H. Paspalin PII. Kodocytoc

chalin 2

[53760-20-6]

C₂₈H₃₇NO₄ 451.605Metab. of a *Phomopsis* sp. Cryst. (Et₂O). Sol. MeOH; poorly sol. H₂O. Mp 274-276° part dec. [α]_D²⁷ +47.8 (c, 0.43 in MeOH). Identity with Kodocytocchalin 2 not firmly establ. λ_{\max} 260 (MeOH) (Berdy).▶ LD₅₀ (mus, ivn) 5 mg/kg. HA5320000

17-Deoxo, 21-Ac: Cytochalasin H. Kodocytocchalin 1. Paspalin P1

[53760-19-3]

C₃₀H₃₉NO₅ 493.642Metab. of *Phomopsis* spp. Shows antibiotic and cytotoxic props. Long needles (CHCl₃/Et₂O). Mp 258-263°. [α]_D -9 (CHCl₃). [α]_D +91.2 (MeOH). Log P 2.74 (uncertain value) (calc). λ_{\max} 286 (ε 330) (MeOH) (Derep).▶ LD₅₀ (mus, ivn) 2 mg/kg. HA5306000

17-Deoxo, 18-deoxy, 21-Ac: Antibiotic L

696474. L 696474C₃₀H₃₉NO₄ 477.642

Metab. of *Hypoxylon fragiforme* ATCC 20995. HIV-1 protease inhibitor. Cryst. (MeOH). Sol. DMSO, CH₂Cl₂, MeOH; poorly sol. H₂O, hexane. Mp 201°. Log P 4.83 (uncertain value) (calc).

4-Methoxy, 17-deoxy, 21-Ac: Pyrichalasin H

[111631-97-1]

C₃₁H₄₁NO₆ 523.668

Isol. from *Pyricularia grisea*. Phytotoxin. Plates (EtOAc). Sol. MeOH, Et₂O; poorly sol. H₂O. Mp 207-209°. [α]_D²³ -18.4 (c, 1.05 in CHCl₃). [α]_D²³ +59 (c, 1.05 in MeOH). Has a 4-methoxy substit. on the phenyl ring. λ_{max} 203 (ε 22000); 224 (ε 17000); 276 (sh) (ε 2000); 283 (sh) (ε 1640) (EtOH) (Derep).

Aldridge, D.C. *et al.*, *J.C.S. (C)*, 1969, 923 (isol, struct, nmr)

Minato, H. *et al.*, *J.C.S. (C)*, 1970, 45 (Zygosporins)

Binder, M. *et al.*, *J.C.S. Perkin I*, 1973, 1146 (Zygosporins E,F)

Graf, W. *et al.*, *Helv. Chim. Acta*, 1974, **57**, 1801 (cmr, biosynth)

Patwardhan, S.A. *et al.*, *Phytochemistry*, 1974, **13**, 1985 (isol, ir, pmr)

Veders, J.C. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 1886; 1976, **59**, 558 (biosynth, cmr)

McMillan, J.A. *et al.*, *Chem. Comm.*, 1977, 105 (cryst struct)

Beno, M.A. *et al.*, *J.A.C.S.*, 1977, **99**, 4123 (cryst struct, pmr, stereochem)

Cole, R.J. *et al.*, *J. Agric. Food Chem.*, 1981, **29**, 205 (deriv)

Chappuis, G. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 521 (synth, props)

Vedejs, E. *et al.*, *J.O.C.*, 1982, **47**, 4384 (synth)

Magin, D.F. *et al.*, *J. Agric. Food Chem.*, 1984, **32**, 544 (ms)

Nukina, M. *et al.*, *Agric. Biol. Chem.*, 1987, **51**, 2625 (Pyrichalasin F)

Vedejs, E. *et al.*, *J.A.C.S.*, 1988, **110**, 4822 (synth)

Edwards, R.L. *et al.*, *J.C.S. Perkin I*, 1989, 57 (isol)

Thomas, E.J. *et al.*, *J.C.S. Perkin I*, 1989, 507; 1999, 3269-3283; 3285-3290 (synth)

Ondeyka, J. *et al.*, *J. Antibiot.*, 1992, **45**, 671-678; 679-685; 686-691 (L 696474)

Maccotta, A. *et al.*, *J.C.S. Perkin 2*, 1993, 729 (pmr, cmr, conformn)

Buchanan, M.S. *et al.*, *Phytochemistry*, 1995, **40**, 135; 1996, **41**, 821; **42**, 173 (21-ketone)

Espada, A. *et al.*, *Tetrahedron*, 1997, **53**, 6485 (19,20-Epoxychochalsin D)

Namatame, I. *et al.*, *J. Antibiot.*, 2000, **53**, 19-25 (activity)

Fujii, Y. *et al.*, *J. Nat. Prod.*, 2000, **63**, 132-135 (isol, pmr, cmr)

Jikai, L. *et al.*, *Helv. Chim. Acta*, 2002, **85**, 1439-1442 (Cytochalsin D, 19,20-Epoxychochalsin D)

Haidle, A.M. *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 2006, **101**, 12048-12053 (L 696474, synth)


Shi, L.-M. *et al.*, *J. Chem. Res.*, 2007, 144-145 (19,20-Epoxychochalsin D, struct)

Tao, Y. *et al.*, *Magn. Reson. Chem.*, 2008, **46**, 501-505 (Cytochalsin H)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 338; 340

Zygosporin G

[25374-69-0]

C₃₀H₃₇NO₅ 491.626

Minor metab. of the fungus *Zygosporium masonii*. Shows antibiotic props. similar to Zygosporin D, Z-37. Prisms (diisopropyl ether). Sol. MeOH, CHCl₃; poorly sol. H₂O, hexane. Mp 115-125°. [α]_D²⁴ -82 (c, 0.87 in dioxan).

► LD₅₀ (mus, ipr) 100 mg/kg. HA5302000

Minato, H. *et al.*, *J.C.S. (C)*, 1970, 45 (isol, ir, ur, nmr, struct)


Binder, M. *et al.*, *J.C.S. Perkin I*, 1973, 1146 (struct)

Chappuis, G. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 521 (synth)

Cole, R.J. *et al.*, *Handbook of Toxic Fungal Metabolites*, Academic Press, 1981, 341

Zyzzin

[159509-39-4]

C₁₁H₇N₃OS 229.262

Metab. from the sponge *Zyzzya massalis*. Deep orange amorph. solid. Undergoes exchange of the thioxo for the oxo group during aqueous workup. λ_{max} 212 (ε 15800); 270 (ε 5200); 345 (ε 2200); 420 (ε 1800); 475 (ε 1200) (MeOH) (Berdy).

3,4-Dihydro, 4-hydroxy: 4-Hydroxy-4-(1H-indol-3-yl)-5-thioxo-2-imidazolidinone

[159308-50-6]

C₁₁H₉N₃O₂S 247.277

Artifact from *Zyzzya massalis*. Racemic.

3,4-Dihydro, 4-methoxy: 4-(1H-Indol-3-yl)-4-methoxy-5-thioxo-2-imidazolidinone

[159308-48-2]

C₁₂H₁₁N₃O₂S 261.304

Artifact from *Zyzzya massalis*. Opt. inactive (racemic). The 5-thioxo analogue of Polyandrocarpamide D, P-542.

5-Oxo analogue: 4-(1H-Indol-3-yl)-1H-imidazole-2,5-dione

[159308-52-8]

C₁₁H₇N₃O₂ 213.195**Z-38**

Metab. from *Zyzzya massalis*. Yellow amorph. solid. Almost certainly an artifact derived from Zyzzin during the isol. procedure.

5-Oxo analogue, C(4),N-dihydro: 5-(1H-Indol-3-yl)-2,4-imidazolidinedione

[159308-51-7]

C₁₁H₉N₃O₂ 215.211

Metab. of *Zyzzya massalis*. Powder. [α]_D²⁵ +10 (c, 0.13 in MeOH). Almost certainly a workup artifact originating from the corresp. thioamide.

5-Oxo analogue, 3,4-dihydro, 4-hydroxy: 5-Hydroxy-5-(1H-indol-3-yl)-2,4-imidazolidinedione

[159308-49-3]


C₁₁H₉N₃O₃ 231.21

Artifact from *Zyzzya massalis*. Solid (EtOH/hexane). Mp 140° dec. Racemic.

Mancini, I. *et al.*, *Helv. Chim. Acta*, 1994, **77**, 1886-1894 (isol, pmr, cmr, ms, struct, *Zyzzya massalis* consists)

Zyzzyanone A

[784152-58-5]

Z-40C₂₀H₁₉N₃O₃ 349.388

Alkaloid from the sponge *Zyzzya fuliginosa*. Cytotoxic. Purple solid (as trifluoroacetate salt). Mp 300° dec. (TFA salt). λ_{max} 242 (log ε 4.26); 286 (log ε 3.9); 342 (log ε 3.59); 484 (log ε 3.44) (MeOH) (TFA salt). λ_{max} 243 (log ε 4.28); 286 (log ε 3.96); 346 (log ε 3.6); 505 (log ε 3.48) (MeOH/KOH) (TFA salt).

N²-Formyl: Zyzzyanone C

[866363-60-2]

C₂₁H₁₉N₃O₄ 377.399

Alkaloid from *Zyzzya fuliginosa*. Brownish-red solid (as TFA salt). λ_{max} 242 (log ε 3.67); 294 (log ε 3.24); 349 (log ε 2.49); 486 (log ε 2.85) (MeOH) (TFA salt).

N⁴-De-Me: Zyzzyanone B

[866363-59-9]

C₁₉H₁₇N₃O₃ 335.362

Alkaloid from *Zyzzya fuliginosa*. Purple solid (as TFA salt). λ_{max} 240 (log ε 3.99); 281 (log ε 3.64); 328 (log ε 3.33); 483 (log ε 3.08) (MeOH) (TFA salt).

N⁴-De-Me, N²-formyl: Zyzzyanone D

[866363-61-3]

C₂₀H₁₇N₃O₄ 363.372

Alkaloid from *Zyzzya fuliginosa*. Brownish-red solid (as TFA salt). λ_{max} 241 (log ε 3.67); 288 (log ε 3.23); 337 (log ε 2.95); 487 (log ε 2.84) (MeOH) (TFA salt).

Utkina, N.K. *et al.*, *Tet. Lett.*, 2004, **45**, 7491-7494 (Zyzzyanone A)

Utkina, N.K. *et al.*, *J. Nat. Prod.*, 2005, **68**, 1424-1427 (Zyzzyanones B,C,D)

- 2-(Acetoxyacetyl)-1-(2-acetoxyethyl)-1*H*-pyrrole, H-401
 2-(Acetoxyacetyl)-1-(2-acetoxypropyl)-1*H*-pyrrole, H-401
 7-Acetoxy-9-acetylspicatamine, T-46
 10-Acetoxyactinidine, A-125
 13-Acetoxyanagyrine, B-20
 N⁶-Acetoxyaphelandrine, A-1334
 2 α -Acetoxyaustrospicatin, T-46
 28-Acetoxy-N³-benzoylbuxidienine, B-466
 9-Acetoxy-1-benzoyloxy-4,8-dihydroxy-6-nicotinoyloxydihydro- β -agarofuran, P-219
 6-Acetoxy-7-benzoyloxy-3-(3,4,5-trimethoxycinnamoyloxy)tropane, M-392
 2-*O*-(3-Acetoxybutanoyl)glycorine, L-348
 16-Acetoxybuxabenzamidiene, B-456
 16-Acetoxycardiopetaline, M-355
 7 β -Acetoxycauabine E, M-392
 3-Acetoxy-3-(2-chloro-2-isocyanovinyl)-2-oxindole, I-74
 4-Acetoxycycloheximide, D-765
 5-Acetoxycycloheximide, D-765
 N-Acetoxyctisine, C-938
 2-Acetoxy-2'-deacetylaustrospicatin, T-46
 2-Acetoxy-2-deacetyl-1-hydroxyaustrospicatin, T-43
 18-Acetoxy-N³-desacetylisoretuline, R-63
 19-Acetoxydihydrogelsevirine, G-49
 5-Acetoxy-5,6-dihydro-1-(3-phenylpropanoyl)-2(1*H*)-pyridinone, P-463
 16-Acetoxy-3,2,3-dihydroxy-16,28-secosolanidan-4-one, S-349
 4-Acetoxy-20-(dimethylamino)-3-phthalimidoprogan-2-ol, D-262
 Acetoxyeduline (incorr.), E-40
 23-Acetoxy-25-episuladulcine, S-442
 1-Acetoxy-24,25-epoxy-7,22-dihydroxyergost-4-en-3-one, P-294
 11-Acetoxyerysotrine, E-206
 2-Acetoxyethylamine, A-767
 14-Acetoxygelsedilam, G-43
 14-Acetoxygelsegine, G-47
 14-Acetoxygelsenine, G-44
 11-Acetoxygraciline, G-151
 3-Acetoxyhexadecanoic acid choline ester, P-12
 14-Acetoxy-15-hydroxygelsenine, G-44
 2-Acetoxy-1-(hydroxynitrosoamino)heptadecane, P-530
 23-Acetoxy-12-hydroxysolasodine, S-453
 3-Acetoxy-1*H*-indole, I-104
 6-Acetoxy-3-isobutyryloxytropane, M-391
 3-Acetoxy-6-isobutyryloxytropane, M-391
 2-Acetoxyisotaxodine, H-331
 13-Acetoxyilupanine, H-554
 4 β -Acetoxyilupanine, H-551
 14 α -Acetoxymatrine, M-121
 14 β -Acetoxymatrine, M-121
 1-Acetoxy-3-methoxy-10-methylacridone, D-528
 7-Acetoxy-1-methoxymethyl-1,2-dehydropyrrolizidine, T-188
 4'-*N*-(Acetoxy-methoxy)staurosporine, S-498
 1-Acetoxy-methyl-2-(10-acetoxyundecyl)-4(1*H*)-quinolinone, H-757
 1-Acetoxy-methyl-2,3-dimethyl-4(1*H*)-quinolinone, H-477
 N-Acetoxy-methylflindersine, F-90
 1-(Acetoxy-methyl)-2-formyl-1,2-dihydro-5-hydroxy-7-methoxyisoquinoline, A-39
 N-(1-Acetoxy-methyl-2-methoxyethyl)-7-methoxy-4-eicosenamide, H-605
 1-Acetoxy-methyl-2-methyl-4(1*H*)-quinolinone, H-633
 1-Acetoxy-methyl-2-nonyl-4(1*H*)-quinolinone, H-642
 3-(2-Acetoxy-4-methyl-3-oxohexyl)-1*H*-indole, H-535
 3-(2-Acetoxy-4-methyl-3-oxopentyl)-1*H*-indole, H-536
 1-Acetoxy-methyl-2-propyl-4(1*H*)-quinolinone, H-707
 3-Acetoxy-2-methyl-4(1*H*)-pyridinone, D-632
 2-(Acetoxy-methyl)quinoline, Q-37
 3-Acetoxy-norerythrosumamine, E-218
 18-Acetoxy-norfluorourarine, N-301
 19-Acetoxy-3-oxotabersonine, O-210
 Acetoxytelefoliarine (incorr.), E-40
 4-Acetoxy-2-pyrrolidone, H-718
 α -Acetoxyquinaldine, Q-37
 4-Acetoxy-2(1*H*)-quinolinone, Q-33
 25-Acetoxyrobustine, S-451
 Acetoxysecodaphniphylline, S-181
 16-Acetoxy-16,28-secosolanida-5,22(28)-diene-3,5-diol, V-64
 18-Acetoxy-senkirine, S-245
 23-Acetoxy-soladulcine, S-442
 2-Acetoxytaxodine, H-331
 6-Acetoxy-3-tiglyloxytropane, M-391
 23*R*-Acetoxytomatidine, S-442
 23*R*-Acetoxytomatine, S-442
 3 α -Acetoxytropane, M-393
 3-Acetoxytropane-6-ol, M-391
 10-Acetyl-17-*O*-acetylajmaline, A-217
 1-Acetyl-17-(acetyloxy)-20,21-didehydro-19-hydroxy-10,11-dimethoxy-4-methyl-3,4-securan-3-one, A-667
 3-Acetylaconitine, A-104
 13-Acetylacridine, H-189
 6-*O*-Acetylacoseptine, A-92
O-Acetylacosmine, A-107
 Acetylacrifoline, L-328
 5-Acetylalaine B, A-199
 10-Acetylalaine B, A-199
 1-*O*-Acetylalaxoxin A, P-72
 N¹-Acetylalaxoxin, A-719
 17-*O*-Acetylajmaline, A-216
 N²-Acetylalanylthranilamide, A-33
 Acetylalloyohimbine, Y-13
 11-*O*-Acetylbambinine, A-683
 2-(Acetylamino)benzamide, A-31
 2-(Acetylamino)benzoic acid, A-31
 4-(Acetylamino)-2-butenic acid, A-717
 6-Acetyl-2-amino-7,8-dihydro-4(3*H*)-pteridinone, A-40
 3-(Acetylamino)-6,7-dimethoxycoumarin, A-738
 2-(Acetylamino)-5,6-epoxy-2-cyclohexene-1,4-dione, A-804
 7-[5-[2-(Acetylamino)ethyl]-2-hydroxyphenyl]-9,10-dihydro-3,5,6,8-tetrahydro-9,10-dioxo-1,2-anthracenedicarboxylic acid, L-3
 5-[(1-Acetylamino)ethyl]-2-methylpyridine, B-248
 2-Acetyl-2-amino-4-hydroxyindeno[1,2-*b*]oxet-3(2*H*)-one, C-358
 2-Acetyl-3-amino-5-hydroxy-1,4-naphthalene-dione, A-41
 2-Acetyl-3-amino-5-hydroxy-1,4-naphthoquinone, A-41
 N-(*N*-Acetylamino)methylcytisine, C-938
 3-(Acetylamino)-7-oxabicyclo[4.1.0]hept-3-ene-2,5-dione, A-804
 5-(Acetylamino)-4-oxo-5-hexenamide, P-630
 2-[4-(Acetylamino)-3-oxo-2-isoxazolidinyl]tetrahydro-5-oxo-2-furancarboxylic acid, L-8
 N²-[2-(Acetylamino)propanoyl]thranilamide, A-33
 8-(Acetylamino)-5,6,7,8-tetrahydro-6,7-dihydroxy-5-(hydroxymethyl)imidazo[1,2-*a*]pyridine-2-acetic acid, N-9
 6-Acetyl-2-amino-1,7,8,9-tetrahydro-4*H*-pyrimido[4,5-*b*][1,4]diazepin-4-one, A-52
 Acetyl-*trans*-anacrotine, A-967
 Acetylanacrotine, A-967
 3-Acetyl-15-angeloylgermine, G-70
O-Acetylanguistoline, A-1020
 N-Acetylanhalamine, T-236
 N-Acetylanhaloline, A-1021
 14-Acetyl-8-anisoylchamanine, M-323
 2-Acetyl-3-anisoylthetidine, H-161
 8-Acetyl-14-anisoylpseudoanone, M-336
 8-Acetyl-14-anisoyltalatzamine, M-352
 14-Acetyl-8-anisoyltalatzamine, M-352
 Acetylannofoline, F-29
 N-Acetylanolobine, A-1066
 N-Acetylanonaine, R-120
 Acetylanonamine, S-245
 N-Acetylanthranilic acid, A-31
 5-*O*-Acetylaponinchonamine, A-1360
 5-*N*-Acetylardeemin, A-1399
 N-Acetylasiimilobine, D-531
 N-Acetylaspidoalbidine, A-1490
 N-Acetylaspidospermatidine, A-1497
 N-Acetylaspidospermidine, A-1498
 1-Acetylaspidospermidine-6,17-diol, A-1067
 2-Acetyl-9-azabicyclo[4.2.1]non-2-ene, A-974
O-Acetylbaptifoline, B-20
 14-Acetylbearline, G-91
 3-*O*-Acetylbeiwutine, M-302
 4-Acetylbenzoxazolin-2-one, A-42
 4-Acetyl-2(3*H*)-benzoxazolone, A-42
 3-Acetyl-14-benzoyloxazine, M-331
O-Acetyl-*N*-benzoylbuxalongifoline, B-466
 16-*O*-Acetyl-*N*-benzoylbuxidienine, B-466
 8-Acetyl-14-benzoylchamanine, M-323
 14-Acetyl-8-benzoylchamanine, M-323
 6-Acetyl-14-benzoyldictyocarpine, M-307
 8-Acetyl-14-benzoylzeochamanine, M-303
 14-Acetyl-8-benzoyltalatzamine, M-352
 3-Acetyl-5-benzyl-2,4-pyrrolidinedione, A-43
 N-Acetylbongardine, B-248
 14-Acetylbrowniine, M-309
 11-*O*-Acetylbulgarsenine *N*-oxide, B-401
 11-*O*-Acetylbulgarsenine, B-401
 3-Acetyl-5-*sec*-butyltetramic acid, T-74
 N²⁰-Acetylbumamine E, B-455
 N²⁰-Acetylbumamine G, B-455
 28-Acetylbumanoldine, B-460
 18-Acetylcammaconine, M-352
O-Acetylcampthoecine, C-70
 17-*O*-Acetylcampthoecinic acid, C-71
 Acetylcandescence, E-12
 Acetylcandescine, E-12
 15-*N*-Acetylcapparisine, C-97
 Acetylcaranine, F-7
 1-Acetyl- β -carboline, A-44
 1-Acetyl- β -carboline-3-carboxylic acid, A-45
 1-Acetyl-3-carboxy- β -carboline, A-45
 11-Acetylcardionine, H-192
 15-Acetylcardiopetaline, H-190
 (+)-Acetylcephalotaxine, C-271
 (-)-Acetylcephalotaxine, C-271
 Acetylcedvedine, C-295
 19-*O*-Acetylchaetoglobosin A, C-325
 19-*O*-Acetylchaetoglobosin B, C-326
 19-*O*-Acetylchaetoglobosin D, C-327
 14-Acetylchamanine, M-323
 5-Acetyl-2-chlorotetrahydro-4-methyl-2*H*-1,3-oxazine, A-46
 Acetylcholine chloride, A-47
 Acetylcholine(1+), A-47
 N-Acetylchlonemorphone, P-601
 N-Acetylchonicinoline, Q-17
O-Acetylcinnabarine, C-452
 N¹-Acetyl-N²-cinnamoyl-N¹-dihydrocinnamoylpur-trescine, D-362
 14-Acetyl-8-*trans*-cinnamoyltalatzamine, M-352
 14-Acetylcolumbidine, M-352
 8-Acetylcondelphine, M-352
O-Acetylcorymine, D-410
 Acetylcorynoline, C-692
 8-*O*-Acetylcorysoline, R-9
O-Acetylcrotaverine, S-245
 6'-*O*-Acetylcupreine, Q-20
 31-Acetylcyclopropylamine A, C-883
 N-Acetylcyclopropobuxine C, C-902
 N-Acetylcyclopropobuxine D, C-902
 N-Acetylcyclopropocarpinol, L-177
O-Acetylcyclopholine, C-935
 N-Acetylcytisine, C-938
 7-*O*-Acetylcytochalasin B, C-943
 9-Acetyl-10-deacetylspicatamine, T-47
 7-Acetyl-10-deacetyltaxol, T-52
 Acetyldebenzoylallopecurine, L-335
 N-Acetyl-*N*-debenzoyltaxol, T-52
 1'-Acetyl-1,1',2,3,4,4',5,5',6,6'-decahydro-2,3'-bipyridine, A-930
 13-Acetyl-5,6,7*a*,8,8*a*,9,12,12*a*,12*b*,13-decahydro-7*a*-hydroxy-10-methyl-10*H*-furo[2,3-*d*]pyra-no[3,4-*a*]carbazole-9-carboxaldehyde, M-122
 13-Acetyldecarbamoylegonautoxin II, H-731
 13-Acetyldecarbamoylegonautoxin III, H-731

- O*-Acetyldecarine, C-356
N-Acetyldehydroanonaine, D-166
 15-Acetyl-13-dehydrocardiopetamine, H-190
N-Acetyldehydrodopamine, D-157
 16-Acetyl-2,3-dehydroelastine, M-362
N-Acetyl-5,6-dehydrotololine, L-234
 12-Acetyldehydrogluculine, D-162
 12-Acetyldehydroglucidusculine, D-162
 14-Acetyldelcosine, M-309
N-Acetyldelectine, M-309
 14-Acetyldelectine, M-309
 14-*O*-Acetyldelectinine, M-309
 14-Acetyldelosine, M-309
 6-Acetyldepheline (incorr.), M-321
 1-Acetyldephisine, M-323
N-Acetyldemethoxycylindrocaryne, C-929
 3-*O*-Acetyldemethylchlidanthine, G-3
 2-Acetyl-2-demethylcolchifoline, C-571
 17-*O*-Acetyl-*N*-demethylechitamine, D-410
 6-*O*-Acetyldemethylenedecorine, M-309
N-Acetyl-3-demethylmescaline, H-465
 1-*O*-Acetyl-9-*O*-demethylpluviine, P-528
 15-Acetyldenudatine, D-221
N-Acetyldeoxycinchonincol, Q-17
 13-*O*-Acetyl-9-deoxyglanduline, H-178
 14-*O*-Acetyl-9-deoxyglanduline, H-178
N^m-Acetyl-3-deoxystrychnosplendine, S-400
N-Acetyl-*N*-deoxymayfoline, P-356
N-Acetyl-*N*-depropionylaspidoalbine, A-1490
 17-*O*-Acetyl-*N^m*-desacetyl-18-hydroxyisoretuline,
 R-63
O-Acetyldiabolone B, C-115
 Acetyldiabolone, C-115
 3-Acetyl-2,5-diamino-1,4-benzoquinone, A-48
 7-Acetyl-3,15-diangeloylgermine, G-70
 Acetyldicrotaline, D-363
 14-Acetyl dictyocarpine, M-307
 6-Acetyl dictyocarpine, M-307
 1-Acetyl-19,20-didehydrocuran-17-ol, R-63
 1-Acetyl-19,20-didehydro-16-(23-deoxy-11-oxa-
 12,24-secostrychinidin-10-yl)-17-norcuran,
 S-591
 1-Acetyl-14,19-didehydro-10,11-dimethoxy-4-
 methyl-3,4-secocondyfolan-3-one, G-36
N-Acetyl-1,2-didehydrodopamine, D-157
 1-Acetyl-19,20-didehydro-17-norcuran-16-one,
 S-603
 1-Acetyl-20,21-didehydro-19-oxocuran-17-ol,
 S-607
O-Acetyl-19,20-didehydroyohimbine, D-378
 1-Acetyl-17,18,17,19-diepoxy-11-methoxycuran-
 20-ol, M-278
N-Acetyl-2,16-dihydroakummicinal, R-63
 9-Acetyl-3,4-dihydro- β -carboline, C-130
 7-Acetyl-6,7-dihydro-1,7-dihydroxy-3-methoxy-
 5*H*-cyclohepta[*b*]quinoline-8,11-dione,
 C-486
 14-Acetyldihydrogadesine, M-321
 3-Acetyl-1,5-dihydro-4-hydroxy-5-(1-methylpro-
 pyl)-2*H*-pyrrol-2-one, T-74
 Acetyldihydrolycopodine, L-341
 1-Acetyl-1,5-dihydro-4-methoxy-2*H*-pyrrol-2-one,
 P-954
 9-Acetyl-4,9-dihydro-3*H*-pyridof[3,4-*b*]indole,
 C-130
 5-Acetyl-3,4-dihydro-2*H*-pyrrole, A-49
O-Acetyldihydroalutarinine, I-318
 17-*O*-Acetyl-19,20-dihydrovoachalotine, P-562
 21-Acetyl-19,20-dihydrovomilenine, V-194
N-Acetyl-3,4-dihydroxyphenylalanine, A-745
 1-Acetyl-16,17-dimethoxyaspidoaspermidine, P-924
 1-Acetyl-16,17-dimethoxyaspidoaspermidin-6-ol,
 A-1067
 2-Acetyl-4,6-dimethoxyfuro[2,3-*b*]quinoline, A-50
 2-Acetyl-4,7-dimethoxyfuro[2,3-*b*]quinoline, A-50
N-Acetyl-3,4-dimethoxyphenethylamine, D-718
N-[2-(2-Acetyl-4,5-dimethoxyphenyl)ethyl]aceta-
 mide, D-219
 6-Acetyl-1,3-dimethylumazine, A-51
 6-Acetyl-1,3-dimethyl-2,4(1*H*,3*H*)-pteridinedione,
 A-51
N-Acetyl-3,5-dinitrotyramine, H-481
 8-Acetyldolaconine, A-99
 3'-*O*-Acetylechihumiline *N*-oxide, E-10
 3'-Acetylechimidine, E-11
 7-Acetyl-9-echimidinylretroreine, U-40
 7-Acetylchinateine, E-12
 17-*O*-Acetylechitamine, D-410
 2'-*O*-Acetyledulinine, E-40
 10-*O*-Acetyledulirin A, T-349
 Acetylenedicarboxylic acid, B-447
 2'-Acetyl-7-epicephalomannine, T-52
*O*¹⁷-Acetyl-3-epidihydrocorymine, D-410
 12-Acetyl-12-epinapelline, N-30
 14-Acetylepineoline, M-323
 2'-Acetyl-7-epitaxol, T-52
 11-*O*-Acetyl-1,2-epoxyambelline, A-683
 1-Acetyl-17,19-epoxycuran, S-400
 11-Acetyl-1,19-epoxydenudatine, K-48
 1-Acetyl-17,19-epoxy-4-methyl-3,4-secocuran-3-
 one, S-594
O-Acetylerucifoline, E-184
O-Acetylethanolamine, A-767
 2-Acetyl-1-ethylpyrrole, A-74
 16-*O*-Acetyletioline, S-199
N-Acetyleudistomine L, E-287
 5'-Acetyleuropine, E-299
 7-Acetyleuropine, E-299
 2-Acetylevolitrine, A-50
 2'-Acetylevoxine, F-223
 8-Acetylexcelsine, D-187
 1-Acetylexcelsine, D-187
 Acetylfaucettine, F-29
N-Acetylflavaconitine, M-332
 14-Acetylgadesine, G-152
O-Acetylgalanthamine, G-3
O-Acetylgeibalansine, G-29
O-Acetylgeissoschizol, G-33
 14-*O*-Acetylgeniculine B, M-349
 9-*O*-Acetylglanduline, H-170
 13-*O*-Acetylglanduline, H-170
N-Acetylglucobrassicin, I-122
 14-*O*-(*N*-Acetylglucosaminyl)teleocidin A,
 L-367
 13-Acetylgomandonine, G-139
 Acetylgyneuramine, G-233
 7-Acetylhackelidine, H-4
 12-*O*-Acetylhadiesine, P-516
 11-*O*-Acetylhaemanthamine, H-5
 3-*O*-Acetylhamayne, H-5
 Acetylhaplatine, F-223
 Acetylhaplophyllidine, H-57
 3'-Acetylheliosupine *N*-oxide, H-94
 Acetylheliosupine, H-94
 2'-*O*-Acetylheliotrine, H-95
O-Acetylhenningsamide, H-110
O-Acetylhenningsine, H-111
 4-*O*-Acetylhennoxazole A, H-112
 9-Acetyl-2-(1-heptenyl)-1,5,9-triazacyclotridecan-
 4-one, L-230
 6-Acetylheteratisine, H-157
 8-Acetylheterophyllisine, H-157
 11-Acetylhetisine, H-202
 13-Acetylhetisine, H-202
 13-Acetylhetisinone, H-202
 4-Acetyl-2a,3,5,6,7,7*b*-hexahydro-3,5-dihydroxy-
 1*H*-2-oxa-7*a*-azacycloprop[*cd*]inden-1-one,
 H-741
 3-Acetyl-1,4,5,6,7,7*a*-hexahydro-7-methyl-2*H*-pyr-
 rolo[3,2-*b*]pyridin-2-one, L-4
N-Acetylhistamine, H-307
N-Acetylhistidinol, H-308
 6'-*O*-Acetylholocalin, H-519
 Acetylhomogmatine, A-859
 6-Acetylhomoperin, A-52
N-Acetylhomoveratrylamine, D-718
N-Acetylhordenine, H-357
 14-*O*-Acetylhumantenidine, G-44
 [*N'*]-5-[[4-[[4-(Acetylhydroxyamino)butyl]amino]-
 1,4-dioxobutyl]hydroxyamino]pentyl]-*N*-(5-ami-
 nopentyl)-*N*-hydroxybutanediamidato(3-)]iron
 (*III*), F-38
 [*N'*]-5-[[4-[[5-(Acetylhydroxyamino)pentyl]amino]-
 1,4-dioxobutyl]hydroxyamino]pentyl]-*N*-(5-ami-
 nopentyl)-*N*-hydroxybutanediamidato(3-)]iron,
 F-39

- O*-Acetyllyngbyatoxin A, L-367
O-Acetylmacralstonine, M-8
O-Acetylmacranthine, M-9
 Acetyl-*cis*-madurensine, M-36
 Acetylmadurensine, M-36
 Acetylmaymyrsine, P-217
 3-*O*-Acetylmaytansinol, M-136
N-Acetylmescaline, M-219
O-Acetylmesebrenol, M-224
 7-*O*-Acetyl-3-methoxyaegeline, A-153
 1-*O*-Acetyl-3'-methoxyaglaroxin A, P-72
 1-Acetyl-17-methoxyaspidospermidine, A-1499
 1-Acetyl-16-methoxyaspidospermidine-6,17-diol, A-1067
 20-*O*-Acetyl-10-methoxycamptothecin, C-70
 20-*O*-Acetyl-9-methoxycamptothecin, C-70
 17-*O*-Acetyl-9-methoxycamptothecin acid, C-71
 1-Acetyl-4-methoxy- β -carboline, A-53
 1-Acetyl-7-methoxy- β -carboline, A-55
 1-Acetyl-3-methoxycarbonyl- β -carboline, A-45
O-Acetyl-11-methoxyisoretuline, R-63
 20-Acetyl-5-methoxy-7-oxa-15,20,24,27-tetraazatracyclo[13.9.6.2^{6,11}.1^{2,6}]tritaconta-2,4,6(33),8,10,12,31-heptaene-14,26-dione, C-317
 2-Acetyl-3-methoxy-1*H*-pyrrole, A-58
N⁶-Acetyl-11-methoxystrychnosplendine, S-605
N-Acetyl-5-methoxytryptamine, M-288
 4-[2-(Acetylmethylamino)ethyl]-4-(3,4-dimethoxyphenyl)cyclohexanone, A-60
 4-[2-(Acetylmethylamino)ethyl]-4-(4-hydroxy-3-methoxyphenyl)-2,4-cyclohexadien-1-one, A-60
 1-(*N*-Acetyl-*N*-methylamino)ethyl-7-hydroxy-3,4,6-trimethoxyphenanthrene, M-559
 2-Acetyl-7-methyl-7-azabicyclo[3.2.1]oct-2-ene, F-47
 6-*O*-Acetyl-14-*O*-methyldephinifoline, M-309
N-(2-Acetyl-5,6-methylenedioxybenzyl)-6,7-methylenedioxy-1(2*H*)-isoquinolinone, I-305
 6-Acetyl-14-*O*-methylforensicine, M-323
 2-Acetyl-3-methylindole, A-61
 3-Acetyl-1-methylindole, A-59
 14-*O*-Acetyl-8-*O*-methyl-18-*O*-[2-(2-methyl-4-oxo-4*H*-quinazolin-3-yl)benzoyl]cammaconine, M-501
 3-Acetyl- ω -methyl- ω -(4-oxodecanoyl)histidine, H-307
N-Acetyl- β -methyl-L-phenylalanine 7-carboxy-1-(2-methyloxiranyl)-2,4,6-heptatrienyl ester, A-220
N⁶-Acetyl-*N⁶*-methylphlegmarine, P-361
 2-Acetyl-3-methylpyrazine, A-62
 4-Acetyl-2-methylpyridine, A-67
 4-Acetyl-3-methylpyridine, A-68
 5-Acetyl-2-methylpyridine, A-69
 2-Acetyl-4-methylpyridine, A-63
 2-Acetyl-6-methylpyridine, A-64
 3-Acetyl-2-methylpyridine, A-65
 3-Acetyl-4-methylpyridine, A-66
 4-Acetyl-6-methyl-2(1*H*)-pyridinone, A-70
 2-Acetyl-1-methylpyrrole, A-74
 3-Acetyl-4-methylpyrrole, A-71
N-Acetyl-*N*-methyl-*N,7a*-secomesembrine, A-60
N⁶-Acetyl-*O*-methylstrychnosplendine, S-605
N⁶-Acetyl-*N⁶*-methyltryptamine, M-582
O-Acetylmillaurine, M-613
N-Acetylmitiphylline, M-651
 Acetylmonocrotaline, M-688
O-Acetylmontanine, P-57
 12-Acetylmapelline *N*-oxide, N-30
 1-Acetylmapelline, N-30
 12-Acetylmapelline, N-30
 3-Acetyl-2,7-naphthyridine, A-72
 3-*O*-Acetylnarcissidine *N*-oxide, N-43
 3-*O*-Acetylnarcissidine, N-43
 18-*O*-Acetylnargenicin A₁, N-276
 12-*O*-Acetylneoehadiensine, P-516
 8-Acetylnesoline, M-323
 14-Acetylnesoline, M-323
 3-*O*-Acetylnerbodine, N-169
O-Acetylnitrarine, N-243
N⁶-Acetyl-2-nitrotryptamine, A-771
N-Acetyl-4-nitrotryptophan, N-257
 18-*O*-Acetylnodumscin, N-276
N-Acetylnoradrenaline, N-298
 17-*O*-Acetylnorajmaline, A-216
N-Acetylnorloline, L-234
N-Acetylnornantene, N-29
N-Acetylnornicotine, P-957
N-Acetylnornuciferine, D-531
 1-*O*-Acetyl-10-norpluviine, P-528
 1-*O*-Acetyl-9-norpluviine, P-528
 Acetylnorrauglucine, A-216
 17-*O*-Acetylnortetraphyllicine, T-288
 14-Acetylnudicauldine, M-321
 4-Acetyloctahydro-2,5-methano-5*H*-pyrido[3,2-*b*]-1,4-oxazine, L-111
 2-*O*-Acetylrochlorine, H-198
O-Acetyl-8-oxocorynoline, C-692
 22-*O*-Acetyl-19-oxodihydroatisine, A-1533
N-Acetyl-2-oxodopamine, A-736
N-Acetyl- β -oxotryptamine, I-132
 2-Acetyloxy-*N*-benzoyl-2-phenylethylamine, A-864
 3-(Acetyloxy)-3-(2-chloro-2-isocyanoethenyl)-1,3-dihydro-2*H*-indol-2-one, I-74
 3-(Acetyloxy)-3-(2-chloro-2-isocyanoethenyl)-2-oxoindole, I-74
 17-(Acetyloxy)-1,2-didehydro-1-demethyl-21-methyl-18-norajmalan-19-al, P-240
 12-(Acetyloxy)-3,8-didehydro-14-methyl-21-norse-necianan-5,11,16-trione, P-770
 12-(Acetyloxy)-3,8-didehydro-2-norse-necianan-5,11,16-trione, S-232
 7-Acetyloxy-2,9*a*-diethyloctahydro-6,9-dimethyl-propyl-2-*H*-pyrido[1,2-*d*]pyrazine-1-carboxaldehyde, C-385
 3-[5-(Acetyloxy)-4,5-dihydro-3-furanyl]-*N*-(2-hydroxy-5-oxo-1-cyclopenten-1-yl)-2-propenamide, R-34
 2-(Acetyloxy)-7,9-dihydroxy-21-methyl-6,21-secorhettisan-6,13-dione, M-665
 16-(Acetyloxy)-3-hydroxy-16,28-secosolanida-5,22(28)-dien-24-one, V-63
 2-[[3-(Acetyloxy)-1-oxohexadecyl]oxy]-*N,N,N*-trimethylethanaminium(1+), P-12
 2-Acetyloxy-*N,N,N*-trimethylethanaminium, A-47
N-Acetylpachypodanthine, O-87
N³-Acetylpachysamine A, P-601
 1-*O*-Acetylpannellin, P-72
N-Acetylparthenolidine, P-99
O-Acetylparadaphnine B, P-139
 1'-*O*-Acetylparaxilline, P-140
 14-*O*-Acetylperegrine, M-347
 6-Acetyl-1-phenazinecarboxylic acid, H-496
 4-Acetyl-2-picoline, A-67
 4-Acetyl-3-picoline, A-68
 5-Acetyl-2-picoline, A-69
 6-Acetyl-2-picoline, A-64
 2-Acetyl-4-picoline, A-63
 3-Acetyl-2-picoline, A-65
 3-Acetyl-4-picoline, A-66
 2-Acetyl- Δ^1 -piperidine, A-76
 2-Acetyl- Δ^2 -piperidine, A-76
O-Acetylplatydesmine, P-515
O-Acetylporanthicine, P-569
O-Acetylpreperaksine, D-491
 1-*O*-Acetylpsudolycorine, P-716
 2-*O*-Acetylpsudolycorine, P-716
O-Acetylptelefoliarine, E-40
 2-Acetylpteleine, A-50
 2-Acetylpyridine, A-73
 1-Acetyl-2-(3-pyridinyl)pyrrolidine, P-957
 1-Acetyl-9*H*-pyrido[3,4-*b*]indole-3-carboxamide, A-45
 1-Acetyl-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, A-45
 2-Acetylpyrrole, A-74
 2-Acetyl-1-pyrrolidine, A-49
N-Acetylpyrrothine, A-759
N-Acetylquestionin A, A-862
 2-Acetyl-4(3*H*)-quinazolinone, A-75
 Acetylrauglucine, A-216
O-Acetylretuline, R-63
 7-*O*-Acetylrinderine, E-12
 12-*O*-Acetylrosmarinine, R-133
 14-Acetylsachaonitine, M-355
O-Acetylsamandarine, S-44
 3-*O*-Acetylsanguinine, G-3
 17-Acetylsarpagine, S-89
 7-Acetylscorpioidine, S-163
N-Acetylseco-*N*-methylauroretanine, M-559
 14-*O*-Acetylsenbusine A, M-323
 12-Acetylseneciphylline, S-240
O-Acetylsenkirkine, S-245
N-Acetylsepaconitine, A-87
 2-Acetylseptentriose, H-180
 Acetylsevedine, C-295
O-Acetylsolasodine, S-456
 15-Acetylsongoramine, D-162
 Acetylsongorine, N-30
 3-*O*-Acetyl(-)-spectaline, S-388
N¹-Acetylspermidine, S-398
N⁵-Acetylspermidine, S-398
N⁸-Acetylspermidine, S-398
N-Acetylspermine, S-399
 3-Acetylstachyflin, S-492
N-Acetylstepharine, P-643
 2'-*O*-Acetylstrictosamide, V-126
N⁶-Acetylstrychnosplendine, S-605
O-Acetylsukhodiane, C-733
 Acetylsyneilesine, S-658
 Acetyltabascanine, S-604
 14-Acetyltakaosamine, M-309
 14-Acetylaltatizamine, M-352
 7-Acetylaxine A, A-6
 2-*O*-Acetylaxine II, T-273
 2-Acetylaxuspine Z, T-47
 14-*O*-Acetylteleocidin A₂, L-367
 14-*O*-Acetylteleocidin B₃, T-63
N-Acetyl- Δ^2 -tetrahydroanabasine, A-930
 1-Acetyl-1,2,3,4-tetrahydro-5-[1-(1*H*-indol-3-ylmethyl)-2-piperidinyl]pyridine, G-155
 1-Acetyl-1,2,3,4-tetrahydro-5-(2-piperidinyl)pyridine, A-930
 2-Acetyl-1,2,3,4-tetrahydro-1-propyl- β -carboline, T-219
 6-Acetyl-1,2,3,4-tetrahydropyridine, A-76
 2-Acetyl-3,4,5,6-tetrahydropyridine, A-76
 1-Acetyl-1,2,3,4-tetrahydropyridine, T-220
 10-*O*-Acetylthapsakin A, T-349
 2-(Acetylthiomethyl)quinoline, Q-36
N-Acetyltortuosamine, T-433
 3'-*O*-Acetyltrachelanthamine, T-443
 6-Acetyl-1,2,3-trimethoxy-9,10-methylenedioxy-noraporphine, P-369
O-Acetyltropine tropate, T-632
N⁶-Acetyltryptamine, T-639
N-Acetyltryptophanol, A-830
N-Acetyltyramine, H-667
 6-Acetylmumbrophine, A-92
 3-*O*-Acetylungminorine, U-35
 13-*O*-Acetylvakhamatine, H-191
O-Acetylvalllesamine, V-9
O-Acetylvenoterpine, V-52
 8-Acetyl-14-veratrolychasmamine, M-323
 17-Acetyl-15-veratrolyditycine, D-370
 3-Acetyl-15-veratrolygermine, G-70
 17-Acetyl-15-veratroly-19-oxodictysine, D-370
 14-Acetyl-8-veratrolytalatizamine, M-352
 3-*O*-Acetylverticine, C-298
 3-*O*-Acetylverticinone, C-298
 3-Acetylvilmorrianine A, M-303
N-Acetylvincoside, S-583
N-Acetylvinerine, C-1
 14-Acetylviresenine, M-325
 9-(3'-Acetylviridifloryl)retronecine, I-68
O-Acetylamataimine *N*-oxide, B-194
O-Acetylamataimine, B-194
O-Acetylyohimbine, Y-13
 3-Acetylyunaconitine, M-336
 Achillamide, T-640
 Achillein, H-476
 Achyranthine, P-952
 β -Acid, D-672
 Acidissiminin epoxide, A-77
 Acidissiminol, A-77
 Acidissiminol epoxide, A-77
 Acidissiminol, A-77

- Acidomycin, A-78
 Aciergocristine, E-155
 Aciergocristinine, E-155
 Aciergotamine, E-172
 Aciergotaminine, E-172
 Acinetobactin, A-79
 Acinetoferrin, A-80
 Acivicin, A-81
 Aclidinomycin A, A-82
 Aclidinomycin B, A-82
 Aclidinomycin C, A-83
 Aclidinomycin D, A-82
 Acobretine A, M-352
 Acobretine B, M-352
 Acobretine C, M-352
 Acobretine D, M-352
 Acobretine E, M-352
 Acofamine A, M-337
 Acofamine B, M-335
 Acofine, A-105
 Acoforesticine, M-323
 Acoforestine, M-312
 Acoforestinine, M-336
 Acoforine, M-352
 Acomonine, M-309
 Aconasutine, M-380
 Aconcaguine, A-84
 Aconifine, M-302
 Aconine 8-linoleate, M-331
 Aconine, M-331
 Aconitane-1,4,6,7,8,14,16-heptol, A-85
 Aconitane-1,4,7,8,9,14,16-heptol, A-86
 Aconitane-1,4,8,9,10,14,16-heptol, A-87
 Aconitane-1,6,7,8,10,14,16-heptol, A-88
 Aconitane-1,4,6,8,14,16-hexol, A-89
 Aconitane-1,4,7,8,14,16-hexol, A-90
 Aconitane-1,4,8,9,14,16-hexol, A-91
 Aconitane-1,6,7,8,14,16-hexol, A-92
 Aconitane-1,3,4,7,8,9,14,16-octol, A-93
 Aconitane-1,4,7,8,9,10,14,16-octol, A-94
 Aconitane-1,4,8,14,16-pentol, A-95
 Aconitane-1,6,7,14,16-pentol, A-96
 Aconitane-1,8,13,14,16-pentol, A-98
 Aconitane-1,8,9,14,16-pentol, A-97
 Aconitane-1,8,14,16-tetrol, A-99
 Aconitane-1,14,16-triol, A-100
 Aconit-19(*N*)-ene-1,6,7,8,14,16-hexol, A-101
 Aconit-19(*N*)-ene-1,8,13,14,16-pentol, A-102
 Aconit-19(*N*)-ene-1,8,14,16-tetrol, A-103
 Aconitilearine, M-314
 Aconitine, A-104
 Aconitinine, M-320
 Aconoridine, M-352
 Aconorine, M-352
 Aconosine, A-99
 Acophine, A-105
 Acordine, H-189
 Acorientine, H-206
 Acorine, H-189
 Acosanine, M-309
 Acosepticine, A-92
 Acoseptine, A-106
 Acoseptidine, M-359
 Acoseptridinine, M-352
 Acoseptrigine, M-323
 Acoseptriginine, M-323
 Acoseptrine, A-88
 Acoseptrinine, M-309
 Acosmine acetate, A-107
 Acosmine, A-107
 Acotoxinine, M-323
 Acovulparine, M-359
 Acozerine, A-108
 Acrawax C, E-225
 Acrawax CT, E-225
 Acremoauxin A, A-109
 Acremoauxin B, A-110
 Acremolactone B, A-111
 9-Acridinol, A-112
 9(10*H*)-Acridinone, A-112
 Acridol, A-112
 Acridone, A-112
 Acrifoline, C-472
 Acrifoline, L-328
 Acrifolinol, L-328
 Acriginine A, A-113
 Acrimarine A, A-114
 Acrimarine B, A-114
 Acrimarine C, A-115
 Acrimarine D, A-115
 Acrimarine E, A-114
 Acrimarine F, A-114
 Acrimarine G, A-114
 Acrimarine H, A-114
 Acrimarine I, A-116
 Acrimarine J, A-116
 Acrimarine K, A-114
 Acrimarine L, A-115
 Acrimarine M, A-114
 Acrimarine N, A-114
 Acrodontiolamide, A-117
 Acromelic acid A, A-118
 Acromelic acid B, A-119
 Acromelic acid C, A-119
 Acromelic acid D, A-118
 Acromelic acid E, A-118
 Acromelinic acid A, A-118
 Acromelinic acid B, A-119
 Acronine, A-120
 Acronycidine, F-216
 Acronycine epoxide, A-120
 Acronycine, A-120
 Acronyidine, A-121
 Acrophyllidine, F-212
 Acrophylline, F-212
 Acrylamide, P-646
 Acrylic acid, P-646
 6-*O*-Acryloyl-2,3-bis-*O*-(3-nitropropanoyl)- α -D-glucopyranose, G-104
 2-*O*-Acryloyl-3,6-bis-*O*-(3-nitropropanoyl)- α -D-glucopyranose, G-104
 2-*O*-Acryloyl-3,4,6-tris-*O*-(3-nitropropanoyl)- α -D-glucopyranose, G-104
 Acsinate, H-201
 Acsine, A-122
 Aconine, A-123
 Actaline, A-210
 Acti-Aid, D-765
 Actidione, D-765
 Actiketal, A-124
 Actinac, A-625
 Actinidine, A-125
 Actinine, A-126
 Actinodaphnine, D-346
 Actinoleukin, Q-58
 Actinophyllic acid, A-127
 Actinoplanone A, A-128
 Actinoplanone B, A-128
 Actinoplanone C, A-128
 Actinoplanone D, A-128
 Actinoplanone E, A-128
 Actinoplanone F, A-128
 Actinoplanone G, A-128
 Actinoviridin A, H-641
 Actiphenol, A-129
 Actithiazic acid, A-78
 γ -Activity X, I-300
 Aculeacin B, A-130
 Aculeamine, D-99
 Acuminaminoside, A-131
 Acuminatopyrone, M-250
 Acutiaporberine, A-132
 Acutifolidine, L-288
 Acutifolidine, P-210
 Acutifoline palmitate, L-288
 Acutifoline, L-288
 Acutine, H-133
 Acutudaurine, A-133
 Acutumidine, A-134
 Acutumine, A-134
 Acutuminine, A-134
 Acutupyrocoline, A-135
 Adaline, P-235
 Adalimine, A-136
 Adenine, A-137
 3-(9-Adeniny)propionic acid, A-900
 Adenocarpine, A-138
 Adenophorine, E-245
 Adermin, P-923
 Aderoxin, P-923
 Adhatodine, A-1045
 Adhavasinsonone, V-38
 Adiantifoline, A-139
 Adifoline, A-140
 Adipamide, H-258
 Adipic acid, H-258
 Adirubine, A-141
 A₃-118D, L-374
 Adlumiceine enol lactone, A-142
 Adlumiceine, B-120
 Adlumidiceine enol lactone, A-144
 Adlumidiceine imide, F-172
 Adlumidiceine, A-143
 Adlumidine, A-145
l-Adlumine, A-146
 Adlumine, A-146
 Adociaquinone B, A-147
 Adonifoline, A-148
 Adouetine X, A-149
 Adouetine Y', A-150
 Adouetine Y, C-600
 Adouetine Z, A-151
 Adouetine, A-151
 Adrenaline, A-152
 Adrenoglomerulotropin, S-285
 Adrenolone, A-736
 Adrenone, A-736
 Adrenor, N-298
 Aducril, F-104
 Aducaine, H-751
 Aduncamide, H-566
 Aegeline, H-12
 Aegeline, A-153
 A2-E, O-65
 AEP, A-773
 Aequale, D-856
 Aerophobin 1, A-154
 Aerophobin 2, A-155
 Aeroplysinillin, A-156
 Aeroplysinin 1, A-157
 Aerothionin, A-158
 Aeruginascine, H-480
 Aerugine, H-695
 Aeruginic acid, H-695
 Aeruginol, H-695
 Aeruginosamide, A-159
 Aeruginosin 101, A-160
 Aeruginosin 205A, A-161
 Aeruginosin 98A, A-160
 Aeruginosin A, A-860
 Aeruginosin 205B, A-161
 Aeruginosin 98B, A-160
 Aeruginosin B, A-162
 Aeruginosin 98C, A-160
 Aervine, H-445
 Aervolanine, C-134
 Aervoside, H-445
 Aesculetic acid, T-577
 Aestivine, A-163
 Aestivophoenin A, A-164
 Aestivophoenin B, A-165
 Aestivophoenin C, A-165
 Aethalic acid, H-221
 AF 140, A-1115
 AF 110, F-197
 Affinine, A-166
 Affinine, D-124
 Affinisine N⁰-oxide, T-425
 Affinisine oxindole, A-167
 Affinisine, T-425
 Aflastatin A, A-168
 Aflastatin B, A-168
 Aflatrem B, A-170
 Aflatrem, A-169
 β -Aflatrem, A-170
 Aflavazole, A-171
 Aflavinine, A-172
 Afragilimycin A, S-566
 Africanine, A-173

- Afrinol, M-386
 Afrocurarine, A-174
 Afzelechin-(4β→2'')-*N*-(*p*-coumaroyl)-6''-hydroxy-tryptamine, L-265
 AG 452, A-176
 AG 468, A-176
 AG 452a, A-176
 AG 379a, H-413
 AG 395a, H-413
 AG 416a, I-82
 AG 432a, I-82
 AG 448a, I-82
 AG 464a, I-82
 AG 379b, H-413
 AG 395b, H-413
 AG 416b, I-82
 AG 432b, I-82
 AG 448b, I-82
 AG 489b, I-82
 AG 505b, I-82
 AG 379c, H-413
 AG 395c, H-413
 AG 416c, I-82
 AG 432c, I-82
 AG 448c, I-82
 AG 432d, I-82
 AG 432e, I-82
 AG 432f, I-82
 AG 432g, I-82
 AG 379, H-413
 AG 395, H-413
 AG 432h, I-82
 AG 416, I-82
 AG 432, I-82
 AG 448, I-82
 AG 464, I-82
 AG 489, I-82
 AG 505, I-82
 AG 521, I-82
 AG 432i, I-82
 AG 55, T-373
 Agaricone, A-175
 Agardiol, A-807
 Agarin, A-837
 Agel 452, A-176
 Agel 468, A-176
 Agel 452a, A-176
 Agel 395a, H-413
 Agel 432a, I-82
 Agel 448a, I-82
 Agel 464a, I-82
 Agel 395b, H-413
 Agel 432b, I-82
 Agel 448b, I-82
 Agel 489b, I-82
 Agel 505b, I-82
 Agel 395c, H-413
 Agel 432c, I-82
 Agel 448c, I-82
 Agel 432d, I-82
 Agel 432e, I-82
 Agel 432f, I-82
 Agel 432g, I-82
 Agel 395, H-413
 Agel 432h, I-82
 Agel 432, I-82
 Agel 448, I-82
 Agel 464, I-82
 Agel 489, I-82
 Agel 505, I-82
 Agel 521, I-82
 Agel 432i, I-82
 Ageladine A, A-177
 Agelasidine A, A-178
 Agelasidine B, A-179
 Agelasidine C, A-180
 Agelasidine D, A-180
 Agelasimine A, A-181
 Agelasimine B, A-181
 Agelazine A, A-184
 Agelazine B, A-184
 Agelazine C, A-185
 Agelazine D, A-186
 Agelazine E, A-187
 Agelazine F, A-191
 Agelazine G, A-184
 Agelazine H, A-184
 Agelazine I, A-183
 Agelazine, A-182
 Agelaspongin, A-188
 Agelastatin A, A-189
 Agelastatin B, A-189
 Agelastatin C, A-189
 Agelastatin D, A-189
 Ageliferin, A-190
 Ageline A, A-191
 Ageline B, A-184
 Agelocaissarine A₁, A-192
 Agelocaissarine A₂, A-192
 Agelocaissarine B₁, A-192
 Agelocaissarine B₂, A-192
 Agelongine, A-193
 Agelolin A, A-194
 Agelolin B, A-194
 Agesamide A, A-195
 Agesamide B, A-195
 Aglaforbesine A, A-196
 Aglaforbesine B, A-196
 Aglaiastatin, A-197
 Aglaiathioduline, L-114
 Aglaidithioduline, A-198
 Aglaiduline, B-422
 Aglaiformosanin, A-197
 Aglaine A, A-199
 Aglaine B, A-199
 Aglaine C, A-199
 Aglairubine, C-456
 Aglamide A, L-115
 Aglamide B, L-115
 Aglamide C, O-67
 Aglamide D, P-955
 Aglanthine, B-422
 Aglaroxin A, P-72
 Aglaroxin B, P-72
 Aglaroxin C, A-200
 Aglaroxin D, A-197
 Aglaroxin F, P-72
 Aglaroxin G, A-200
 Aglaroxin H, A-200
 Aglaroxin I, A-200
 Aglaroxin J, A-199
 Aglaroxin K, A-199
 Aglaroxin L, A-199
 Aglatenin, A-716
 Aglatenol, A-716
 Aglaxiflorin A, A-202
 Aglaxiflorin B, A-202
 Aglaxiflorin C, A-201
 Aglaxiflorin D, A-202
 Agleptine, L-115
 Agmatine, A-719
 Agrobactin, A-203
 Agrochelin, A-204
 Agroclavine I, A-205
 Agroclavine, A-205
 Agrocybenine, A-206
 Agrocybin, H-647
 Agropine, A-207
 AH 23491X, C-100
 AH 135Y, A-129
 2088A, H-514
 Ailanindole, A-208
 Ailantcanthinoside A, H-442
 Ailantcanthinoside B, H-438
 Ailanthoidine, A-209
 AIR, A-827
 AIS 2M3, R-33
 Ajabicine, A-210
 Ajacine, M-309
 Ajaconine, A-211
 Ajacusine, G-91
 Ajadelphine, M-325
 Ajadelphinine, M-325
 Ajadine, M-309
 Ajadinine, M-359
 Ajanine, M-309
 Aji 9561, A-1116
 Ajimaline, A-216
 Ajmalan-17,21-diol, A-216
 Ajmalicine, A-214
 Ajmalicine hydroxyindolenine, A-213
 Ajmalicine oxindole A, M-661
 Ajmalicine oxindole B, M-661
 Ajmalicine, A-212
 Ajmalicinial, A-214
 Ajmalicinine, A-215
 Ajmalidine, A-216
 Ajmalimine, A-216
 Ajmaline, A-216
 Ajmalinimine, A-217
 Ajmalinine, A-218
 Ajmalinol, A-216
 AJP 117510, U-42
 Ajudazol A, A-219
 Ajudazol B, A-219
 AK toxin I, A-220
 AK toxin II, A-221
 Akagerine lactone, A-223
 Akagerine, A-222
 Akalone, A-820
 Akanthomycin, A-224
 Akarpine, P-424
 Akashin A, D-353
 Akashin B, D-353
 Akashin C, A-225
 AK-40-II, O-69
 AK-40-I, O-68a
 Akiradine, D-187
 Akiramidine, A-89
 Akiramine, A-89
 Akirane, A-89
 Akiranine, A-89
 Akirine, A-226
 Aknadicine, H-70
 Aknadilactam, H-70
 Aknadine, E-102
 Aknadinine, H-70
 AK-Ps, O-124
 Akuammenine, A-227
 Akuammicine *N*^b-methosalt, A-228
 Akuammicine *N*^b-oxide, A-228
 Akuammicine, A-228
 Akuammidine, A-229
 (*Z*)-Akuammidine, A-229
 Akuammigine (4*R*)-*N*-oxide, A-212
 Akuammigine (4*S*)-*N*-oxide, A-212
 Akuammigine pseudoindoxyl, T-289
 Akuammigine, A-212
 ψ-Akuammigine, A-232
 Akuammiginone, A-230
 Akuammiline *N*¹-oxide, A-231
 Akuammiline, A-231
 Akuammine *N*-oxide, A-232
 Akuammine, A-232
 [Ala¹]Desoxoviroidin, V-160
 Alafine, H-629
 Alalakine, A-233
 Alamarckine, E-72
 Alamaridine, A-234
 Alamarine, A-235
 Alancine, A-1047
 Alangamide, E-72
 Alangicine, E-72
 Alangimarckine, T-675
 Alangimaridine, A-235
 Alangimarine, A-235
 Alangimarinone, A-235
 Alangine A, A-236
 Alangine B, E-72
 Alangine, A-1328
 Alanginine, A-237
 Alangiobussine, A-238
 Alangiobussinine, A-238
 Alangiside, A-239
 Alanine anhydride, D-769
 Alanine cyclic anhydride, D-769
 Alaninol, A-894
 Alantrypinone, A-240
N-β-Alanyldehydrodopamine, D-157

- N*- β -Alanyldopamine, A-241
O-Alanylgriseoliteic acid, H-514
N- β -Alanylhistamine, C-157
 4-Alanyl-3-isovalin-5-one, A-733
 2-Alanyl-3-isoxazolin-5-one, A-850
N- β -Alanylnoradrenaline, N-298
*N*⁶- β -Alanyl-8-oxoadenine, E-177
 Alatamide, A-242
 Alatamine, P-253
 Alaternifoline, M-392
 Alatol, H-248
 Alatusamine, P-243
 Alatusinine, P-253
 Albertamine, D-79
 Albertidine, M-121
 Albertine, L-94
 Albiflomanthine, H-5
 Albifloranine, C-657
 Albine, A-243
 Albizzine A, P-40
 Albofungin B, A-244
 Albofungin, A-244
 Alboinone, A-245
 Albomaculine, N-171
 Albomitomycin A, A-246
 Albomitomycin C, A-246
 Alborine, A-247
 Alboviolaconitine A, M-309
 Alboviolaconitine B, L-317
 Alboviolaconitine C, L-317
 Alboviolaconitine D, A-248
 Albovionitine, A-249
 Alcaligin, A-250
 Alchorneine, A-251
 Alchorneinone, A-252
 Alchornidine, A-253
 Alchornine, A-254
 Aldehydeollidine, E-256
 Aldisine, A-255
 Aldohypaconitine, M-343
 α -Aldotripiperidine, A-256
 Aldrithiol 2, D-831
 Aleicide B, A-257
 Aleicide C, A-258
 Aleurodiscoester, A-259
 Aleurodisconitrile, A-259
 Alexine, T-555
 Alfileramine, A-260
 Alginidine, A-261
 Alginine, A-262
 Alihirsutine A, A-263
 Alioline, A-264
 Aljesaconitine A, M-331
 Aljesaconitine B, M-331
Geotrichum Alkaloid A 25822A, D-742
 Alkaloid A 58365B, A-1104
Buxus Alkaloid 1, A-275
Buxus Alkaloid 2, A-276
Buxus Alkaloid 3, A-277
Buxus Alkaloid 4, A-278
Buxus Alkaloid 5, A-279
Dendrobates Alkaloid 185, A-287
Dendrobates Alkaloid 197, A-288
Dendrobates Alkaloid 203, A-289
Dendrobates Alkaloid 205, A-289
Dendrobates Alkaloid 217, A-290
Dendrobates Alkaloid 225, A-291
Dendrobates Alkaloid 233, A-292
Dendrobates Alkaloid 241, A-361
Dendrobates Alkaloid 245, A-509
Dendrobates Alkaloid 247, A-509
Dendrobates Alkaloid 295, A-293
Dendrobates Alkaloid 301, A-294
Dendrobates Alkaloid 325, A-295
Dendrobates Alkaloid 351, A-296
Fritillaria Alkaloid 5, A-297
Fritillaria Alkaloid 6, A-298
Leuconotis griffithii Alkaloid 308, A-309
Leuconotis Alkaloid 376, A-308
Lycopodium Alkaloid 168, A-312
Lycopodium Alkaloid 258, A-313
Mantella Alkaloid 161, A-318
Mantella Alkaloid 191, A-319
Mantella Alkaloid 317, A-320
Pancreatium Alkaloid 31, A-324
Pancreatium Alkaloid 6, A-323
Rosmarinus officinalis Alkaloid 2, A-329
Senecio renardi Alkaloid 4, A-332
Sophora Alkaloid 9, A-333
 Alkaloid 196, A-907
 Alkaloid 302, A-1548
Abrus precatorius Alkaloid A, A-347
Buxus Alkaloid A1, A-348
Buxus Alkaloid A2, A-349
Buxus Alkaloid A3, A-350
Buxus Alkaloid A4, A-351
Buxus Alkaloid A5, A-352
Chondrodendron limaciiifolium Alkaloid A, A-353
Daphniphyllum Alkaloid A1, A-345
Dendrobates Alkaloid 167A, A-354
Dendrobates Alkaloid 181A, A-355
Dendrobates Alkaloid 209A, A-356
Dendrobates Alkaloid 219A', A-357
Dendrobates Alkaloid 219A, A-357
Dendrobates Alkaloid 223A, A-357
Dendrobates Alkaloid 231A, A-358
Dendrobates Alkaloid 237A, A-359
Dendrobates Alkaloid 239A, A-360
Dendrobates Alkaloid 241A, A-361
Dendrobates Alkaloid 243A', A-362
Dendrobates Alkaloid 243A, A-362
Dendrobates Alkaloid 251A, A-363
Dendrobates Alkaloid 253A, A-359
Dendrobates Alkaloid 257A, A-364
Dendrobates Alkaloid 265A, A-365
Dendrobates Alkaloid 269A, A-366
Dendrobates Alkaloid 275A', A-367
Dendrobates Alkaloid 275A, A-367
Dendrobates Alkaloid 277A, A-367
Dendrobates Alkaloid 281A, A-368
Dendrobates Alkaloid 289A, A-367
Dendrobates Alkaloid 297A, A-369
Dendrobates Alkaloid 309A, A-370
Erythrophleum Alkaloid A, A-371
Euonymus Alkaloid A, A-372
Fritillaria Alkaloid A, A-373
Mantella Alkaloid 384A, A-374
Scelietum Alkaloid A4, A-346
Securidaca longipedunculata Alkaloid A, A-375
Sophora alopecuroides Alkaloid A1, A-637
Sophora alopecuroides Alkaloid A2, A-637
Stephania glabra Alkaloid A, A-376
Stephania glabra Alkaloid A, A-377
Stephania japonica Alkaloid A, A-378
Taxus baccata Alkaloid A, A-379
Thermopsis lanceolata Alkaloid A, A-380
 Alkaloid A1, A-1337
 Alkaloid AA2, F-2
Knightia Alkaloid A, B-96
Dendrobates Alkaloid 223AB, O-41
Dendrobates Alkaloid 239AB, O-41
 C-Alkaloid A, C-31
 Alkaloid A, C-355
 Alkaloid A, C-587
 Alkaloid A8, C-115
Dendrobates Alkaloid 223A, D-394
 Alkaloid A2, D-407
 Alkaloid AD-IV, D-412
 Alkaloid AD-VI, G-33
Dendrobates Alkaloid 233A, E-264
 Alkaloid A, G-49
 Alkaloid A4, G-87
Dendrobates Alkaloid 235A, H-309
Dendrobates Alkaloid 259A, H-309
Dendrobates Alkaloid 283A, H-309
Dendrobates Alkaloid 285A, H-309
Dendrobates Alkaloid 287A, H-309
Dendrobates Alkaloid 291A, H-309
 Alkaloid A, I-342
Alangium Alkaloid AL64, A-381
 Alkaloid ALC 1, V-95
 Alkaloid ALC 2, V-95
 Alkaloid AM 1, A-150
 Alkaloid AM 3, A-382
 Alkaloid AM 2, C-600
Dendrobates Alkaloid 217A, M-507
Hamelia patens Alkaloid A, M-661
 Alkaloid 20A, M-693
Dendrobates Alkaloid 203A, O-52
Dendrobates Alkaloid 205A, O-52
Dendrobates Alkaloid 207A, O-52
Dendrobates Alkaloid 195A, P-794
Dendrobates Alkaloid 211A, P-794
Dendrobates Alkaloid 307A, P-791
Dendrobates Alkaloid 323A, P-791
Dendrobates Alkaloid 341A, P-792
Pseudophryne Alkaloid 258A, P-720
Pseudophryne Alkaloid 272A, P-720
Pseudophryne Alkaloid 286A, P-720
Pseudophryne Alkaloid 346A, P-720
 Alkaloid APAL-VII, P-550
 Alkaloid AQC2, A-1491
 Alkaloid A6, R-167
 Alkaloid AV2, V-39
Vinca pusilla Alkaloid A, Y-13
Alangium Alkaloid B1, A-383
Alangium Alkaloid B2, A-384
Alangium Alkaloid B3, A-385
Alangium Alkaloid B4, A-386
Alangium Alkaloid B5, A-387
Asteracantha Alkaloid B1, A-388
Asteracantha Alkaloid B2, A-389
Buxus Alkaloid B1, A-391
Buxus Alkaloid B387, A-397
Buxus Alkaloid B398, A-398
Buxus Alkaloid B426, A-399
Buxus Alkaloid B4, A-392
Buxus Alkaloid B5, A-393
Buxus Alkaloid B6, A-394
Buxus Alkaloid B7, A-395
Buxus Alkaloid B8, A-396
Buxus Alkaloid B, A-390
 C-Alkaloid B, A-400
Dendrobates Alkaloid 181B, A-401
Dendrobates Alkaloid 205B, A-402
Dendrobates Alkaloid 219B, A-403
Dendrobates Alkaloid 223B, A-404
Dendrobates Alkaloid 225B, A-405
Dendrobates Alkaloid 231B, A-406
Dendrobates Alkaloid 237B, A-407
Dendrobates Alkaloid 239B, A-408
Dendrobates Alkaloid 249B, A-509
Dendrobates Alkaloid 253B, A-409
Dendrobates Alkaloid 257B, A-410
Dendrobates Alkaloid 265B, A-509
Dendrobates Alkaloid 267B, A-411
Dendrobates Alkaloid 269B, A-412
Dendrobates Alkaloid 279B, A-509
Dendrobates Alkaloid 281B, A-413
Dendrobates Alkaloid 283B, A-414
Dendrobates Alkaloid 297B, A-415
Dendrobates Alkaloid 307B, A-416
Dendrobates Alkaloid 309B, A-417
Euonymus Alkaloid B, A-372
Gentiana macrophylla Alkaloid B, A-418
Mantella Alkaloid 243B, A-511
Mantella Alkaloid 245B, A-511
Mantella Alkaloid 253B, A-419
Mantella Alkaloid 271B, A-420
Mantella Alkaloid 293B, A-421
Mantella Alkaloid 295B, A-422
Pseudophryne Alkaloid 252B, A-423
Skytanthus acutus Alkaloid B, A-424
Sophora griffithii Alkaloid B, A-425
Strychnos camptoneura Alkaloid B, A-426
Thermopsis lanceolata Alkaloid B, A-427
 Alkaloid 205B, A-402
 Alkaloid 241B, A-428
Dendrobates Alkaloid 195B, B-434
Knightia Alkaloid B, B-96
Dendrobates Alkaloid 197B, D-246
Dendrobates Alkaloid 275B', D-109
Dendrobates Alkaloid 275B, D-109
 Alkaloid B3, D-810
u-Alkaloid B, E-67
Dendrobates Alkaloid 289B, G-65
 Alkaloid B, G-181
Dendrobates Alkaloid 235B', H-127
Dendrobates Alkaloid 235B, H-127

- Dendrobates* Alkaloid 251B, H-127
Dendrobates Alkaloid 285B, H-309
Dendrobates Alkaloid 287B, H-309
 Alkaloid B, I-342
 Alkaloid B', I-12
 C-Alkaloid BL, A-429
 Alkaloid B, M-309
 Alkaloid BNC 1, D-161
Dendrobates Alkaloid 167B, O-56
Dendrobates Alkaloid 209B, O-52
Dendrobates Alkaloid 259B, O-51
 Alkaloid B1, O-21
Dendrobates Alkaloid 341B, P-792
Pseudophryne Alkaloid 258B, P-720
Pseudophryne Alkaloid 272B, P-722
Pseudophryne Alkaloid 286B, P-720
Psychotria Alkaloid B, Q-3
 Alkaloid B, T-51
 Alkaloid B, V-103
Buxus Alkaloid BX10, A-431
Buxus Alkaloid BX6, A-430
 Alkaloid 5, C-588
Buxus Alkaloid C, A-435
 C-Alkaloid C, A-436
Chamaecytisus Alkaloid C₁, A-432
Chamaecytisus Alkaloid C₂, A-433
Crotalaria gorensis Alkaloid C, A-437
Dendrobates Alkaloid 223C, A-438
Dendrobates Alkaloid 225C, A-439
Dendrobates Alkaloid 237C, A-440
Dendrobates Alkaloid 239C, A-441
Dendrobates Alkaloid 251C, A-442
Dendrobates Alkaloid 279C, A-509
Dendrobates Alkaloid 283C, A-443
Dendrobates Alkaloid 307C, A-444
Dendrobates Alkaloid 309C, A-445
Erythrophleum Alkaloid C, A-446
Euonymus Alkaloid C, A-372
Gentiana macrophylla Alkaloid C, A-447
Leucojum Alkaloid C, A-448
Mantella Alkaloid 205C, A-449
Mantella Alkaloid 211C, A-450
Mantella Alkaloid 235C, A-451
Mantella Alkaloid 243C, A-511
Mantella Alkaloid 245C, A-486
Mantella Alkaloid 261C, A-452
Mantella Alkaloid 321C, A-453
Rauwolfia obscura Alkaloid C₂, A-434
Salicornia herbacea Alkaloid C, A-454
Stephania glabra Alkaloid C, A-455
Voacanga thouarsii Alkaloid C, A-456
 Alkaloid C, A-1330
 Alkaloid C, A-654
 Alkaloid C, A-669
 Alkaloid C₂, A-457
 Alkaloid 1 (Calabash curare), A-342
 Alkaloid 2 (Calabash curare), A-343
 Alkaloid C, B-94
 Alkaloid CB2, A-458
 Alkaloid CC 11, A-461
 Alkaloid CC 13, A-462
 Alkaloid CC 14, A-463
 Alkaloid CC 20, A-459
 Alkaloid CC 23, A-464
 Alkaloid CC 2, A-459
 Alkaloid CC 3, A-460
 Alkaloid CC 10, A-1005
 Alkaloid CC 3b, A-465
 Alkaloid CC 12, C-568
 Alkaloid CC 19, D-92
 Alkaloid CC 5, D-92
 Alkaloid CC 6, D-92
 Alkaloid CC 7, D-92
 Alkaloid CC 1, N-29
 Alkaloid CC 24, P-221
 Alkaloid C, C-356
 Alkaloid CC22, A-1558
 Alkaloid CC21, K-99
Lycopodium Alkaloid C, D-114
 Alkaloid C, D-510
 Alkaloid C1, D-391
 Alkaloid C3, D-391
Dendrobates Alkaloid 239CD, O-41
 Alkaloid CF 15, C-260
 Alkaloid CF 5, C-271
 Alkaloid CG 1, T-188
Dendrobates Alkaloid 287C, G-65
u-Alkaloid C, G-211
 Alkaloid C, G-44
Dendrobates Alkaloid 285C, H-309
Mantella Alkaloid 275C, H-260
Hortia Alkaloid C₂₂H₁₉N₃O₃, A-467
Papaver Alkaloid C₂₁H₂₁NO₆, A-466
 Alkaloid C, I-12
 Alkaloid C, I-342
Dendrobates Alkaloid 195C, M-532
 Alkaloid C, M-309
 Alkaloid C, M-571
 Alkaloid Cm-4, C-684
 Alkaloid CM1, S-54
 Alkaloid CN1, A-468
 Alkaloid CN2, A-469
Aegle marmelos Alkaloid C, O-63
 Alkaloid C, P-606
 Alkaloid CPC 1, A-626
 Alkaloid CPC 2, T-139
 Alkaloid CP-P, E-270
Cinchona ledgeriana Alkaloid C, Q-12
 Alkaloid C, Q-2
 Alkaloid CR1, A-470
 Alkaloid CRB 11, A-471
 Alkaloid C, S-125
 Alkaloid C, S-89
 Alkaloid CSA 3, A-472
Buxus Alkaloid D, A-475
 C-Alkaloid D, A-476
Cassia Alkaloid D, A-477
Coelidium Alkaloid D, A-478
Dendrobates Alkaloid 223D, A-479
Dendrobates Alkaloid 251D, A-480
Dendrobates Alkaloid 253D, A-481
Dendrobates Alkaloid 285D, A-482
Dendrobates Alkaloid 307D, A-483
Leucojum Alkaloid D, A-484
Mantella Alkaloid 211D, A-485
Mantella Alkaloid 243D, A-486
Mantella Alkaloid 259D, A-452
Mantella Alkaloid 279D, A-487
Melodinus aeneus Alkaloid D, A-488
Narcissus Alkaloid D, A-489
Pandaca caducifolia Alkaloid D, A-490
Rauwolfia obscura Alkaloid D₁, A-473
Rauwolfia obscura Alkaloid D₂, A-474
Salicornia herbacea Alkaloid D, A-491
Stephania glabra Alkaloid D, A-492
 Alkaloid 267D, A-495
 Alkaloid D, A-649
 Alkaloid D1, A-493
 Alkaloid D2, A-494
Knightsia Alkaloid D, B-95
 Alkaloid D2b, D-391
 Alkaloid D, C-138
 Alkaloid D1, D-391
Darlingia Alkaloid DF, H-429
 Alkaloid D₂, G-31
Dendrobates Alkaloid 209D, H-264
Dendrobates Alkaloid 233D, H-127
Dendrobates Alkaloid 237D, H-127
Dendrobates Alkaloid 241D, H-611
Dendrobates Alkaloid 287D, H-309
 Alkaloid D3, H-434
 Alkaloid D, I-288
 Alkaloid D, P-228
 Alkaloid D, S-126
Mantella Alkaloid 193D, T-204
 Alkaloid D, V-188
Buxus Alkaloid E, A-496
Coelidium Alkaloid E, A-497
Croton linearis Alkaloid E, A-498
Dendrobates Alkaloid 223E, A-499
Dendrobates Alkaloid 237E, A-500
Dendrobates Alkaloid 239E, A-438
Dendrobates Alkaloid 251E, A-359
Lycopodium Alkaloid E, A-501
Muntafara sessilifolia Alkaloid E, A-502
Pandaca caducifolia Alkaloid E, A-503
Skytanthus acutus Alkaloid E, A-504
Sophora microphylla Alkaloid E, A-505
Voacanga thouarsii Alkaloid E, A-506
 Alkaloid E, A-638
 C-Alkaloid E, C-809
 Alkaloid ED, F-2
Dendrobates Alkaloid 285E, H-309
Knightsia Alkaloid E, H-428
 Alkaloid E, P-228
 Alkaloid E, P-742
 Alkaloid E, S-126
 Alkaloid E7, T-137
 Alkaloid 7, F-143
Buxus Alkaloid F, A-507
Dendrobates Alkaloid 223F, A-508
Dendrobates Alkaloid 239F, A-438
Dendrobates Alkaloid 251F', A-509
Dendrobates Alkaloid 251F, A-509
Mantella Alkaloid 221F, A-510
Mantella Alkaloid 233F, A-451
Mantella Alkaloid 241F, A-511
Mantella Alkaloid 249F, A-512
Mantella Alkaloid 265F, A-513
Mantella Alkaloid 281F, A-514
Pandaca caducifolia Alkaloid F, A-515
 Alkaloid F45, A-143
 C-Alkaloid F, C-31
 Alkaloid F10, C-805
 Alkaloid F22, C-76
 Alkaloid F30, C-805
 Alkaloid F33, C-172
 Alkaloid F35, C-172
 Alkaloid F3, C-298
 Alkaloid F4, C-298
 Alkaloid F, D-92
 Alkaloid F37, F-174
 Alkaloid F38, F-180
 Alkaloid F46, F-172
 Alkaloid F_k-5, F-174
Knightsia Alkaloid F, H-428
 Alkaloid FI-A, L-88
 Alkaloid FK 1000, S-311
 Alkaloid FK 3000, S-311
 Alkaloid 6F, M-693
 Alkaloid F17, O-16
 Alkaloid F48, O-16
 Alkaloid FR 5, P-168
 Alkaloid F, S-126
 Alkaloid F28, S-234
 Alkaloid F, V-188
 Alkaloid F29, X-30
Dendrobates Alkaloid 239G, A-479
Dendrobates Alkaloid 275G, A-367
Lycopodium Alkaloid G, A-516
Mantella Alkaloid 207G, A-517
Mantella Alkaloid 251G, A-451
Mantella Alkaloid 263G, A-452
Mantella Alkaloid 277G, A-452
Pandaca caducifolia Alkaloid G, A-518
Voacanga thouarsii Alkaloid G, A-519
 Alkaloid 251G, A-520
 Alkaloid GB14, A-521
 Alkaloid GB15, A-522
 Alkaloid GB16, A-523
 Alkaloid GB17, A-524
 Alkaloid GB18, A-525
 Alkaloid GB10, H-277
 Alkaloid GB11, H-277
 Alkaloid GB12, H-277
 Alkaloid GB13, H-281
 Alkaloid GB1, H-277
 Alkaloid GB2, H-277
 Alkaloid GB3, H-277
 Alkaloid GB4, H-277
 Alkaloid GB5, H-277
 Alkaloid GB6, H-277
 Alkaloid GB7, H-277
 Alkaloid GB8, H-277
 Alkaloid GB9, H-277
 C-Alkaloid G, C-809
 Alkaloid G, E-130
 Alkaloid GF, H-444
Dendrobates Alkaloid 223G, H-345

- Alkaloid G, S-125
Dendrobates Alkaloid 255, H-611
Dendrobates Alkaloid 235H, A-509
Mantella Alkaloid 277H, A-452
Alkaloid H, A-676
Dendrobates Alkaloid 249H, E-240
Alkaloid HF1, S-164
Alkaloid HL1, P-667
Alkaloid H, S-126
C-Alkaloid H, T-436
C-Alkaloid I, A-280
Gentiana Alkaloid I, A-301
Voacanga thouarsii Alkaloid I, A-340
Dendrobates histrionicus Alkaloid I, B-428
Dendrobates Alkaloid 207I, E-265
Alkaloid I, E-53
Gentiana Alkaloid II, A-526
Dendrobates histrionicus Alkaloid III, A-366
Sternbergia Alkaloid III, A-527
Alkaloid III, A-970
Dendrobates Alkaloid 209I, O-44
Lycopodium Alkaloid IV, A-344
Solanum Alkaloid IV, A-528
Ungernia Alkaloid IV, A-529
Gentiana cruciata Alkaloid IVb, A-530
Lycopodium annotinum Alkaloid IX, A-531
C-Alkaloid J, A-532
Dendrobates Alkaloid 251J, A-509
H-Alkaloid J, A-1328
Mantella Alkaloid 207J, A-533
Mantella Alkaloid 235J, A-534
Dendrobates Alkaloid 223J, B-443
Alkaloid 195J, D-112
Dendrobates Alkaloid 267J, E-263
Alkaloid J, G-21
Alkaloid J, M-293
Alkaloid J, S-125
Mantella Alkaloid 235K, A-535
Alkaloid K10, A-542
Alkaloid K11, A-543
Alkaloid K2, A-536
Alkaloid K3, A-537
Alkaloid K4, A-537
Alkaloid K5, A-538
Alkaloid K6, A-539
Alkaloid K7, A-540
Alkaloid K9, A-541
Alkaloid KD1, B-96
Alkaloid K8, J-43
Alkaloid K13, L-287
Alkaloid K, S-125
C-Alkaloid K, T-436
Alkaloid 2, L-278
Alkaloid LA 4, A-548
Alkaloid LA 5, A-549
Alkaloid LA 6, A-550
Alkaloid LA 9, A-551
Alkaloid LA 1, E-269
Alkaloid LA 2, E-257
C-Alkaloid L, A-544
Colubrina Alkaloid L, A-545
Mantella Alkaloid 251L, A-546
Alkaloid L10, A-547
Alkaloid L11, A-1062
Alkaloid L7, A-1063
Dendrobates Alkaloid 237L, B-430
Alkaloid LB1, A-552
Alkaloid LBX, A-553
Alkaloid LBY, L-293
Alkaloid LBZ, A-553
Alkaloid LC2, A-554
Alkaloid LC7, A-555
Alkaloid LC1, E-95
Alkaloid LC4, E-95
Alkaloid LC5, E-95
Alkaloid LC8, E-95
Alkaloid L5, D-92
Alkaloid LE1, A-556
Alkaloid L34, F-29
Alkaloid L5, F-71
Alkaloid L9, F-29
Alkaloid L, G-21
Alkaloid L12, L-328
Alkaloid L14, L-341
Alkaloid L1, L-341
Alkaloid L20, L-322
Alkaloid L21, L-280
Alkaloid L23, L-325
Alkaloid L24, L-341
Alkaloid L25, L-325
Alkaloid L26, L-341
Alkaloid L27, L-328
Alkaloid L29, L-328
Alkaloid L2, L-341
Alkaloid L30, L-325
Alkaloid L32, L-321
Alkaloid L33, L-321
Alkaloid L3, L-341
Alkaloid L6, L-287
Alkaloid L8, L-325
Alkaloid LO1, A-557
Alkaloid LO2, A-558
Alkaloid LO3, A-559
Alkaloid LS14, A-560
Alkaloid LV1, A-561
Alkaloid LV1, A-562
Alkaloid LV2, A-563
Alkaloid LV3, A-555
Alkaloid LV3, A-564
Alkaloid LV4, A-565
Alstonia quaternata Alkaloid 3, M-292
Buxus Alkaloid M, A-566
Buxus Alkaloid M, A-567
C-Alkaloid M, A-568
Alkaloid MB 10, A-569
Dendrobates Alkaloid 251M, E-263
Alkaloid M, G-21
Alkaloid MJ1, J-43
Colchicum autumnale Alkaloid M, L-287
Alkaloid MQ-A₃, P-949
Alkaloid MQ-B₂, P-949
Alkaloid MQ-B₄, P-949
Buxus Alkaloid N, A-570
Alkaloid NB, C-754
Alkaloid ND 370, A-571
Alkaloid ND 305B, N-76
Alkaloid ND 363C, N-76
Alkaloid N, G-21
Alkaloid N3, H-290
Alkaloid NP, O-63
C-Alkaloid O, A-572
Alkaloid Or₂, A-573
Alkaloid Or₁, S-39
Dendrobates Alkaloid 357, P-792
Pseudophryne Alkaloid 242, P-720
Pseudophryne Alkaloid 256, P-720
Pseudophryne Alkaloid 270, P-720
Pseudophryne Alkaloid 300, P-722
Pseudophryne Alkaloid 302, P-720
Pseudophryne Alkaloid 316, P-720
Pseudophryne Alkaloid 330, P-722
Pseudophryne Alkaloid 332, P-720
Pseudophryne Alkaloid 512, P-721
Pseudophryne Alkaloid 524, P-721
Pseudophryne Alkaloid 528, P-721
Baptisia Alkaloid P₂, A-574
Baptisia Alkaloid P₄, A-575
C-Alkaloid P, A-576
Lupinus Alkaloid P1, A-577
Lupinus Alkaloid P2, A-578
Mucuna pruriens Alkaloid P, A-579
Alkaloid PAR 1, A-580
Alkaloid PAR 2, A-581
Alkaloid PAR 3, A-582
Alkaloid P3, B-20
Alkaloid P61, D-407
Alkaloid P, E-250
Alkaloid P7, I-242
Alkaloid P4, L-322
Alkaloid PN1, A-955
Alkaloid PO3, A-583
Alkaloid PO4, A-584
Alkaloid PO5, A-247
Alkaloid P2, P-71
Alkaloid P3, P-40
Alkaloid P5, P-71
Alkaloid PP1, A-585
Alkaloid PP1, A-586
C-Alkaloid Q, A-587
Mucuna pruriens Alkaloid Q, A-588
Veratrum Alkaloid Q, A-589
Alkaloid Q, S-200
Alkaloid Q₃, T-425
C-Alkaloid R, A-590
Fritillaria imperialis Alkaloid R, A-591
Lycopodium fawcettii Alkaloid R, A-592
Mucuna pruriens Alkaloid R, A-593
Alkaloid RA, R-86
Alkaloid RB20, P-241
Alkaloid RB19, T-425
Alkaloid RCL 2, V-194
Alkaloid RCLM 12, M-10
Alkaloid RCLM 1, V-194
Alkaloid RC, R-86
Alkaloid R6, E-99
Alkaloid R1, G-32
Alkaloid R-K, A-247
Alkaloid RL-A, S-327
Alkaloid R-L, G-87
Alkaloid RMB 10, A-594
Alkaloid RMB 1, C-689
Alkaloid RMB 5, C-689
Alkaloid RMB 7, C-689
Alkaloid RP5, P-241
Alkaloid RP7, V-194
Alkaloid RS, R-86
Alkaloid RW47, V-52
Dendrobates Alkaloid 222, S-437
Dendrobates Alkaloid 236, S-437
Alkaloid 3, S-126
C-Alkaloid S, A-597
Mucuna pruriens Alkaloid S, A-598
Voacanga schweinfurthii Alkaloid S₂, A-595
Voacanga schweinfurthii Alkaloid S₃, A-596
Alkaloid SC, E-184
Alkaloid SC2, S-54
Alkaloid SD, S-238
Alkaloid S-F, A-599
Alkaloid SN-A, N-89
Alkaloid SN-b, S-447
Alkaloid SN-c, S-454
Alkaloid SN-d, S-454
Alkaloid SN-e, S-447
Alkaloid SN-f, S-454
Alkaloid Su 3078, Y-13
Mantella Alkaloid 189, T-204
Alkaloid T, C-616
Alkaloid TC-A, V-181
Alkaloid TC-B, V-181
Alkaloid TC-C, V-181
Alkaloid TC-D, A-600
Alkaloid TD-2, T-406
Alkaloid TGL4, C-552
Alkaloid TGL3, T-511
Alkaloid TGS-1, T-514
C-Alkaloid T, S-89
Alkaloid 20, U-32
Veratrum Alkaloid U, A-601
C-Alkaloid UB, A-602
Alkaloid Us 7, A-603
Alkaloid Us 8, A-603
Gentiana Alkaloid V, A-302
Helleborus Alkaloid V, A-305
Lycopodium Alkaloid V, A-310
Pachysandra Alkaloid V, A-322
Dendrobates Alkaloid 223V, B-437
Alkaloid V, H-521
Gentiana Alkaloid VI, A-604
Lycopodium annotinum Alkaloid VI, A-605
Pachysandra terminalis Alkaloid VI, A-607
Rhodophiala bifida Alkaloid VI, A-608
Lycopodium annotinum Alkaloid VII, A-606
Lycopodium annotinum Alkaloid VIII, A-609
Alkaloid V, J-36
Alkaloid V3, M-631
Alkaloid V, T-51
Alkaloid W102, A-1013
Alkaloid W95, H-554
Lycopodium Alkaloid X, A-311

- Mucuna pruriens* Alkaloid X, A-321
Veratrum Alkaloid X, A-339
 Alkaloid X, C-616
Lycopodium annotinum Alkaloid XI, A-610
Pachysandra terminalis Alkaloid XI, A-612
Lycopodium annotinum Alkaloid XII, A-611
Rhodophiala bifida Alkaloid XIII, A-613
 Alkaloid XI, T-149
 Alkaloid X₁, M-731
 Alkaloid X₂, M-731
Pachysandra terminalis Alkaloid XVI, A-614
Rhodophiala bifida Alkaloid XXV, A-615
Pachysandra terminalis Alkaloid XXVII, A-616
 C-Alkaloid Y, A-617
 Alkaloid YC₂, A-618
 Alkaloid Y, D-567
 Alkaloid YS, P-270
 Alkaloid Z, R-125
 Alkaloid ZT₁, A-34
 Alkaloid ZT₃, A-34
 Alkaloid ZT₂, D-408
Aconitum kirinense Alkaloid, A-268
Aconitum koreanum Alkaloid, A-269
Aconitum nemorosum C₂₀ Alkaloid, A-265
Aconitum nemorosum C₂₃ Alkaloid, A-266
Alstonia constricta Alkaloid, A-270
Ammოდendron karelinii Alkaloid, A-271
Aplysina archeri Alkaloid, A-272
Artemisia rutifolia Alkaloid, A-273
Astragalus berterianus Alkaloid, A-274
Callyspongia fibrosa Alkaloid, A-281
Chamaecytisus Alkaloid, A-282
Colchicum luteum Alkaloid, A-283
Crotalaria retusa Alkaloid, A-284
Cytisus laburnum Alkaloid, A-285
Cytisus ruthenicus Alkaloid, A-286
Erythrophleum C₂₄ Alkaloid, A-267
Fromia monilis Alkaloid, A-299
Galanthus caucasicus Alkaloid, A-300
Gynura segetum Alkaloid, A-303
Heliotropium europaeum Alkaloid, A-304
Hovea longipes Alkaloid, A-306
Jacmaia incana Alkaloid, A-307
Lycopodium annotinum Alkaloid, A-314
Lycopodium annotinum Alkaloid, A-315
Lycopodium clavatum Alkaloid, A-316
Lycopodium thyoides Alkaloid, A-317
Pancreatum sickenbergi Alkaloid, A-325
Papaver rhoesas Alkaloid, A-326
Pseudaxinyssa Alkaloid, A-327
Rhynchosia pyramidalis Alkaloid, A-328
Sceletium namaquense Alkaloid, A-330
Senecio borysthenicus Alkaloid, A-331
Strepeliopsis strepelioides Alkaloid, A-334
Thermopsis lanceolata Alkaloid, A-335
Ulex europaeus Alkaloid, A-336
Uncaria callophylla Alkaloid, A-337
Ungernia severtzovii Alkaloid, A-338
Zephyranthes Alkaloid, A-341
Pseudoceratina crassa Alkaloid, B-323
 2-(3-Alkyl-5-chloro-2,6-dihydroxybenzoyl)-4,5-dichloro-1H-pyrroles, A-619
 2-Alkyl-3,4-dihydroxy-5-hydroxymethylpyridines, A-620
 2-Alkyl-3,4-dihydroxyquinolines, A-621
 2-Alkyl-5-formylpyrroles, A-624
 3-Alkyl-2-formylpyrroles, A-623
 2-Alkyl-5-hydroxymethyl-3,4-pyridinediols, A-620
 2-Alkyl-3-hydroxy-4(1H)-quinolinones, A-621
 2-Alkyl-5-(1H-indol-3-yl)oxazole, A-622
 1-Alkyl-2-methyl-3H-carbazole-3,4(9H)-dione, C-129
 3-(2-Alkyl-5-oxazolyl)-1H-indoles, A-622
 5-Alkyl-1H-pyrrole-2-carboxaldehydes, A-624
 3-Alkyl-1H-pyrrole-2-carboxaldehydes, A-623
 2-Alkyl-3,4-quinolinediols, A-621
 Allantoin, A-625
 Alliarinoside, H-431
 Alline, A-626
 Allithiamine, A-627
 Alloanodendrine, A-1065
 Alloaristoteline, A-1432
d-Allobisnorbiotin, B-214
 Allocolchicine, A-537
 Allocryptopine, F-2
 Alloculantaraminol, C-803
 Allocuspareine, G-9
 Allodihydrohistrionicotoxin, H-309
 Alloelaeocarpiline, E-55
 Alloenduracididine, E-80
 α-Allokainic acid, K-4
 α-Allokainimic acid, K-4
 Allomatrine, A-628
 5-Allomuscaine, M-773
 Allopterin, A-637
 Allopumiliotoxin 267A, A-480
 Allopumiliotoxin 339A, P-791
 Allopumiliotoxin 341A, P-792
 Allopumiliotoxin 323B', P-791
 Allopumiliotoxin 323B'', P-791
 Allopumiliotoxin 339B, P-791
 Allopumiliotoxin 321C, A-453
 Allopumiliotoxin 309D, A-629
 Allopumiliotoxin 225E, P-795
 Allosecurinine, A-630
 Allosedamine, S-210
 (±)-Allosedamine, S-210
 (+)-Allosedamine, S-210
 (±)-Allosedridine, S-216
 (+)-Allosedridine, S-216
 (-)-Allosedridine, S-216
 3β-Allosolanidan-3-ol, D-205
 Allosophoramine, S-376
 Allotetrahydrohistrionicotoxin, H-309
 Allothiobinupharidine, A-631
 Alloxazine, A-632
 Allohimbine, Y-13
N-Allylloperine, A-637
 11-Allylcytisine, T-412
 1-Allyl-4,5-dibromo-1H-pyrrole-2-carboxylic acid, D-339
 4-Allyl-6,7-dihydro-5H-2-pyridin-5-one, L-272
 2-Allyl-5-(2,3-dihydroxypropyl)decahydroquinoline, A-481
 4-Allyl-1-ethylquinolizidine, E-265
N-Allylnitarine, N-246
N-Allylnorgalanthamine, G-3
 2-Allyl-5-(2-penten-4-ynyl)decahydroquinoline, A-362
 Almazole A, A-633
 Almazole B, A-633
 Almazole C, A-634
 Almazole D, A-635
 Almazolone, A-636
 Almeine, A-1377
 Almocarpin, P-424
 Aloycopine, F-28
 Alopecuridine, F-30
 Alopecurine, L-335
 Aloperine, A-637
 Alphosyl, A-625
 Alpinigene, A-638
 Alpinine, A-638
 Alpinine, G-91
 Alpinone, M-734
 Alpkinidine, A-639
 Al-R 2081, P-974
 Al-R2397, M-136
 Alschomine, A-640
 Alstocraline, A-641
 Alstofoline, A-654
 Alstogustine, E-30
 Alstohentine, T-20
 Alstolactone, A-642
 Alstolagumine, A-228
 Alstolenine, A-231
 Alstomacrine, A-643
 Alstomacrine, A-644
 Alstomacrophyllyne, A-645
 Alstomaline, A-646
 Alstomicine, A-647
 Alstonal, A-648
 Alstonamide, V-125
 Alstonamine, A-1015
 Alstonerinal, T-20
 Alstonerine, A-649
 Alstonidine, A-650
 Alstonilidine, A-651
 Alstoniline, O-140
 Alstonine, A-652
 Alstonisidine, A-653
 Alstonisine, A-654
 Alstonoxine A, A-655
 Alstonoxine B, A-655
 Alstophyllal, A-656
 Alstophyllan-19-one, A-649
 Alstophylline, A-649
 Alstopicralamine, P-417
 Alstoscholarine, A-657
 Alstoumerine, A-658
 Alstovenine, V-50
 Alstovine, C-589
 Alstozine *N*-oxide, S-577
 Altaconitine, M-301
 Altamycin A, A-659
 Altemicidin, A-660
 Alteramide A, A-661
 Alteramide B, A-661
 Alteramine, T-412
 Alternatamide A, A-662
 Alternatamide B, A-662
 Alternatamide C, A-662
 Alternatamide D, A-662
 Alternidine, C-938
 Alternine, T-362
 Altersetin, A-663
 Althiomycin, A-664
 Alvanidine, A-665
 Alvanine, A-666
 Alvimine, A-667
 Alviminine, A-668
 Alzegir, L-162
 AM 5344A₁, C-287
 AM 5344A₂, C-287
 AM 2498, B-216
 AM 2, C-600
 AM 404, E-48
 AM 6201, R-34
 AM 2282, S-498
 Amabiline *N*-oxide, A-669
 Amabiline, A-669
 Amabile, A-670
 Amaminol A, A-671
 Amaminol B, A-671
 Amamistatin A, A-672
 Amamistatin B, A-672
 Amanin, A-673
 Amaninamide, A-673
 Amanitin, A-673
 Amanitin, C-419
 Amanitin, A-673
 Amaranthin, B-110
 Amaranthin, B-110
 Amarbellisine, A-674
 Amaroridine, H-446
 Amarorine, H-446
 Amaryllydine, F-7
 Amaryllylsine, A-675
 Amataine, A-676
 Amathamide A, A-677
 Amathamide B, A-677
 Amathamide C, A-678
 Amathamide D, A-678
 Amathamide E, A-678
 Amathamide F, A-678
 Amathamide G, A-678
 Amathaspiramide A, A-679
 Amathaspiramide B, A-679
 Amathaspiramide C, A-679
 Amathaspiramide D, A-679
 Amathaspiramide E, A-679
 Amathaspiramide F, A-679
 Amatoxin, A-673
 Amauromine, A-680
 Amazonine, E-99
 Ambaline, A-681
 Ambalinine, A-682
 Ambelline, A-683
 Ambiguine A isonitrile, A-684

- Ambiguine B isonitrile, A-684
 Ambiguine C isonitrile, A-684
 Ambiguine D isonitrile, A-685
 Ambiguine E isonitrile, A-686
 Ambiguine F isonitrile, A-687
 Ambiguine G nitrile, A-688
 Ambiguine H isonitrile, A-684
 Ambiguine I isonitrile, A-686
 Ambiguine J isonitrile, A-685
 Ambiguine, M-309
 Ambinine, A-689
 Ambrimine, E-42
 Amelizol, C-420
 Americine, A-690
 Amerovolficine, P-10
 Amerovolfine, A-27
 Ametycine, M-654
 AMG 1, H-713
 Amianthane, A-691
 Amicin, N-250
 Amide C-18, O-32
 Amidenin, D-104
N-Amidino- β -alanine, G-208
 1-Amidino-3-(3-carboxypropyl)urea, G-140
N-(2-Amidinoethyl)-3-aminocyclopentanecarboxamide, A-692
N-(2-Amidinoethyl)-5-imino-2-pyrrolidinecarboxamide, N-278
*N*⁶-Amidinolysine, H-326
 Amidinomycin, A-692
 3-Amidinopyrrole, P-943
N-Amidinotaurine, G-204
 5-(3-Amidinoureido)-2-aminovaleric acid, G-75
 γ -(Amidinoureido)butyric acid, G-140
 1-(3-Amidinoureido)-4-(*N*-methylacetamido)butane, N-187
 15-Amido-3-demethoxy-2,3-methylenedioxyerythroline, D-200
 Amigal, H-616
 Amination, A-715
 Aminazin, C-417
 C₂₄ Amine β -acetate, A-267
 Amin-Glaukosan, H-307
 2-Amino-12-methyl-1,3,12-tetradecanetriol, A-693
 Aminoacetamide trimethylbetaine, C-119
 Aminoacetophenone, A-694
 (Aminoacetyl)benzene, A-694
 2-Aminoadenine, P-817
 4-Amino-*N*-(4-aminobutyl)-*N,N*-bis(3-aminopropyl)-1-butanaminium, B-149
 α -Amino-5-[[[2-aminocarbonyl-1-carboxyethyl]amino]carbonyl]-2-pyrrolopropanoic acid, A-695
 α -Amino- β -(aminocarbonyl)-1,4-dihydro-4-methoxy-1-methyl-4-quinolinepropanoic acid, E-24
 8-Amino-9-(aminocarbonyl)-7-oxo-7*H*-phenoxazine-2-carboxylic acid, A-854
 α -Amino-2-[(4-amino-4-carboxybutyl)amino]-4*H*-imidazo[4,5-*b*]pyridine-4-hexanoic acid, P-233
 2-Amino- α -(2-amino-1-chloroethyl)-1*H*-imidazole-4-methanol, G-79
 5-Amino-3-[5-amino-1,6-dihydro-2,6-dioxo-3(2*H*)-pyridinylidene]-2,6(1*H*,3*H*)-pyridinedione, I-70
 2-Amino-6-(2-aminoethyl)-3-(3,4-dihydroxyphenyl)-1,4-benzodioxan, A-696
 2-Amino-7-(2-aminoethyl)-3-(3,4-dihydroxyphenyl)-1,4-benzodioxan, A-744
 2-Amino-5-(2-amino-1-hydroxyethyl)-4,5-dihydro-1*H*-imidazole-4-carboxylic acid, S-565
 3-Amino-1-(2-amino-3-hydroxyphenyl)-1-propanone, A-697
 4-Amino-1-(2-amino-4-imidazolyl)-2-chloro-1-butanol, G-79
 3-Amino-1-(2-aminoimidazolyl)-1-propene, A-701
 1-Amino-2-[[[aminoiminomethyl]amino]methyl]cyclopropanecarboxylic acid, C-163
 2-Amino-5-[[[aminoiminomethyl]amino]pentanoic acid, A-1413
 3-Amino-5-[[[aminoiminomethyl]amino]pentanoic acid, A-779
 5-Amino-*N*-(3-amino-3-iminopropyl)-3,4-dihydro-2*H*-pyrrole-2-carboxamide, N-278
 2-Amino-3-(3-amino-1*H*-inden-2-yl)propanoic acid, A-698
 2-Amino-5-(aminomethyl)pyrrolo[2,3-*d*]pyrimidin-4(3*H*)-one, A-699
 3-Amino-1-(2-aminophenyl)-1-propanone, A-700
 2-Amino-3-(3-aminopropionyl)phenol, A-697
 2-Amino-4-(3-amino-1-propenyl)-1*H*-imidazole, A-701
 2-Amino-*N*-(3-aminopropyl)-4,7-dihydro-4-oxo-1*H*-pyrrolo[2,3-*d*]pyrimidine-5-carboxamide, E-9
 2-Amino-7-[[2-amino-1,4,5,6-tetrahydro-4,6-dioxo-7-pteridiny]methylene]-1,5,7,8-tetrahydro-4,6-pteridinedione, P-771
 3-Aminoandrost-4-en-17-ol, A-702
 2-Aminoanisic acid, A-788
 4-Amino-1-arabinofuranosyl-2-oxo-1,2-dihydropyrimidine, C-962
 4-Amino-1-arabinofuranosyl-2(1*H*)-pyrimidinone, C-962
 1-Amino-1(10)-aristolene-2,8,9-trione, A-703
 3'-Aminoavarone, A-1560
 1-Amino-8-azabicyclo[3.2.1]octane-2,3,4-triol, A-704
 2-Aminobenzamide, A-706
 4-Amino-1,3-benzenediol, A-705
 2-Aminobenzoic acid, A-706
 9-Amino-8*H*-benzo[*b*]pyrido[4,3,2-*de*][1,10]phenanthroline-8-one, H-542
 2-Aminobenzoyl-6-deoxy- α -L-talopyranoside, A-706
 2-Amino-8-benzoyl-6-hydroxy-3*H*-phenoxazin-3-one, A-707
*N*⁶-(4-Aminobenzoyl)tryptophan, T-640
 2-(2-Aminobenzylamino)benzyl alcohol, A-708
 2-Amino-3-benzyl-5-(4-hydroxyphenyl)pyrazine, C-559
 4-Amino-1-benzyl-1*H*-imidazole, A-826
 7-Amino-2,10-bisaboladiene, A-710
 3-Amino-8,10-bisaboladiene, A-709
 7-Amino-2,11-bisaboladien-10*R*-ol, A-710
 7-Amino-2,11-bisaboladien-10*S*-ol, A-710
 7-Amino-2,9-bisaboladien-11-ol, A-710
 7-Aminobisabolene, A-710
 Aminobisabololene, A-710
 7-Amino-2,8-bis(2,4-dihydroxyphenyl)-3*H*-phenoxazin-3-one, A-711
 2-Amino-1,4-bis(*p*-hydroxybenzyl)imidazole, I-252
 2-Amino-3-(3-bromo-5-chloro-4-hydroxyphenyl)propanoic acid, A-712
 1-Amino-11-bromo-1,2,7,8,13,13*b*-hexahydro[1,6,2]oxathiazepino[2',3':1,2]pyrido[3,4-*b*]indol-10-ol, E-287
 3-Amino-2-bromo-5-hydroxy-5-vinyl-2-cyclopenten-1-one, A-824
 4-Amino-5-bromopyrrolo[2,3-*d*]pyrimidine, A-713
 4-Amino-1,3-butanediol, A-714
 4-Aminobutanoic acid betaine, A-126
 4-Aminobutanoic acid lactam, P-956
 4-Aminobutanoic acid trimethylbetaine, A-126
 4-Aminobutanoic acid, A-715
 4-Amino-1-butanol, A-716
N-(4-Aminobutanoyl)glutamic acid, G-107
 4-Amino-2-butenic acid, A-717
 Amino-*tert*-butyl alcohol, A-840
N-(4-Aminobutyl)- β -alanine, P-826
N-(4-Aminobutyl)-*N'*-[4-[(4-aminobutyl)amino]butyl]-1,4-butanediamine, D-290
N-(4-Aminobutylamino)propanoic acid, P-826
N-[3-[(4-Aminobutyl)amino]propyl]acetamide, S-398
N-(4-Aminobutyl)-*N*-(3-aminopropyl)acetamide, S-398
 3-[[[3-(4-Aminobutyl)amino]propyl]amino]-7-hydroxycholesterol-25-en-24-one, S-485
N-(4-Aminobutyl)-*N*-(3-aminopropyl)benzamide, S-398
N-(4-Aminobutyl)-*N*-(3-aminopropyl)-1,4-butanediamine, A-718
*N*¹-(4-Aminobutyl)-*N*¹,*N*¹⁶-bis(3-aminopropyl)-*N*¹⁶-[4-[(3-aminopropyl)amino]butyl]hexadecanediamine, T-79
N-(4-Aminobutyl)-1,4-butanediamine, D-252
N'-(4-Aminobutyl)canavalmine, D-291
N-(4-Aminobutyl)-2,3-dihydroxybenzamide, B-422
N-(4-Aminobutyl)-3-(3,4-dihydroxyphenyl)-2-propanamide, C-456
N-(4-Aminobutyl)glutamic acid, G-107
 (4-Aminobutyl)guanidine, A-719
*N*¹⁰-(4-Aminobutyl)homohexamine, D-286
*N*⁵-(4-Aminobutyl)homohexamine, D-286
*N*¹⁰-(4-Aminobutyl)homopentamine, D-290
*N*⁵-(4-Aminobutyl)homopentamine, D-290
*N*⁵-(4-Aminobutyl)homospermine, B-150
N-(4-Aminobutyl)-*N*-(3-methyl-2-butenyl)guanidine, S-406
N-(4-Aminobutyl)-*N'*-methylguanidine, A-719
 4-(4-Aminobutyl)octadecahydroprido[1,2-*e*][1,5]diazacycloheptadecim-5(6*H*)-one, O-91
N-(4-Aminobutyl)-*N'*-prenylguanidine, A-719
N-(4-Aminobutyl)-*N*-prenylguanidine, S-406
 1-(4-Aminobutyl)tetrahydro- β -carboline, N-80
 (4-Aminobutyl)urea, A-720
 γ -Aminobutyric acid, A-715
 β -Amino- γ -butyrolactone, A-792
 3-Amino- ϵ -caprolactam, A-785
 2-[(Aminocarbonyl)amino]pentanediamide, G-108
 3-(Aminocarbonyl)-1,4-dihydro-1-methyl-4-oxo-2-quinolinecarboxylic acid, H-728
 2-(Aminocarbonyl)hydrazinecarboxylic acid, H-395
 2-(Aminocarbonyl)-3-hydroxy-6-methoxy-5-methyl-4-(1-propenyl)benzoic acid, N-194
N-(Aminocarbonyl)-1*H*-indole-3-carboxamide, I-84
 11-[5-(Aminocarbonyl)-4-oxo-4*H*-pyran-2-yl]-10-undecenoic acid, H-282
 1-(Aminocarbonyl)pyrrolidine, P-951
 3-(Aminocarbonyl)-1- β -D-ribofuranosylpyridinium(1+), N-189
 α -Amino-3-carboxy-1,5-dihydroxy- γ -oxo-12*H*-pyrido[3,2-*a*]phenoxazine-11-butanolic acid, D-527
 4-(2-Amino-2-carboxyethyl)-1-(5-amino-5-carboxy-2-hydroxypropyl)-3-(3-amino-3-carboxypropyl)-5-hydroxypyridinium(1+), P-907
 4-(2-Amino-2-carboxyethyl)-1*H*-pyrrole-2-carboxylic acid, A-721
 (4-Amino-4-carboxy-2-hydroxybutyl)dimethylsulfonium hydroxide inner salt, D-784
 α -Amino-3-carboxy-1-hydroxy- γ ,5-dioxo-5*H*-pyrido[3,2-*a*]phenoxazine-11-butanolic acid, X-11
 7-Amino-1-carboxy-5-methylphenazinium hydroxide inner salt, A-860
 7-Amino-1-carboxy-5-methyl-3-sulfophenazinium betaine, A-162
 7-Amino-1-carboxy-5-methyl-3-sulfophenazinium hydroxide inner salt, A-162
 11-(Aminocarbonylmethyl)tetrodotoxin, C-390
 α -Amino-6-carboxy-2-oxo-3-pyridinepropanoic acid, A-722
 1-(1-Amino-1-carboxypentyl)-3-[1-(1-amino-1-carboxypentyl)-2-piperidinyl]pyridinium(1+), A-964
 1-(5'-Amino-5'-carboxypentyl)pyrrole, A-904
 2-Amino-1-carboxy-3*H*-phenoxazin-3-one, A-852
N-[[[(3-Amino-3-carboxypropoxy)amino]imino]methyl]aspartic acid, C-75
 2-Amino-3-(2-carboxy-4-pyrrolyl)propanoic acid, A-721
 5-Amino-2-carboxy-1,4,5,6-tetrahydro-4-oxo-3-pyridineacetic acid, G-146
 17-Amino-3,7,11-cembratrien-16,2-olid-19-oic acid, A-723
 Aminocheilin, B-422
 α -Amino-3-chloro-4,5-dihydro-5-isoxazoleacetic acid, A-81
 7-Amino-6-chloro-3,4-dihydro-1-methyl-2-(methylthio)pyrrolo[4,3,2-*de*]quinolin-8(1*H*)-one, I-193
 α -Amino-4-(2-chloroethenyl)-1*H*-pyrrole-3-propanoic acid, A-1277
 2-Amino-3-(3-chloro-4-hydroxyphenyl)propanoic acid, A-724
 3-(2-Amino-3-chlorophenyl)-4-chloro-1*H*-pyrrole, A-725

- 4-(2-Amino-3-chlorophenyl)-2,3-dichloro-1*H*-pyrrole, A-725
 Aminochloropyrrolinitrin, A-725
 3-Aminoconan-12-ol, H-318
 α -Amino-*m*-cresol, H-416
 α -Amino-*o*-cresol, H-415
 α -Amino-*p*-cresol, H-417
 4-Aminocrotonic acid, A-717
 6-Amino-9-(2-cyanoethyl)purine, A-900
 2-Amino-4-cyanopyrimidine, A-903
 Aminocyclohexane, C-868
 7-Amino-7-demethoxymimosamycin, M-624
 6-Amino-6-demethoxymitomycin B, M-654
 5-Amino-5-deoxy-1,2,3,6-di-*O*-isopropylidene- α -D-glucofuranose, A-727
 5-Amino-5-deoxygalactopyranose, A-726
 1-*C*-(5-Amino-5-deoxygalactopyranosyl)butane, B-432
 5-Amino-5-deoxygalactose, A-726
 5-Amino-5-deoxyglucose, A-727
 5-Amino-5-deoxy-1,2-*O*-isopropylidene- α -D-galactofuranose, A-726
 5-Amino-5-deoxy-1,2-*O*-isopropylidene- α -D-glucofuranose, A-727
 5-Amino-5-deoxy-1,2-*O*-isopropylidene-6-*O*-trityl- α -D-galactofuranose, A-726
 2-Amino-2-deoxykealiquinone, K-25
 5-Amino-5-deoxymannose, A-728
 8-Amino-1,15-dichloro-1,14-pentadecadiene-3,12-diyne, D-357
 2-Amino-4,6-dichloropyridine, A-729
 4-Amino-1,13-diguanylidene-5-tridecanone, A-730
 7-Amino-7,8-dihydro- α -bisabolene, A-710
 3-Aminodihydro-4,5-dimethyl-2(3*H*)-furanone, A-800
 α -Amino-3,6-dihydro-2,6-dioxo-1(2*H*)-pyrimidine-propanoic acid, I-345
 1-(7-Amino-5,8-dihydro-5,8-dioxo-2-quinolinyl)-4-methyl-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, L-78
 4-Aminodihydro-2(3*H*)-furanone, A-792
 2-Amino-4,5-dihydro-1-[(hydroxymethoxyphosphinyl)oxy]-*N,N*-dimethyl-1*H*-imidazole-5-methanamine, A-975
 2-Amino-7,8-dihydro-6-(2-hydroxy-1-oxopropyl)-4(1*H*)-pteridinedione, S-246
 2-Amino-4,5-dihydro-1*H*-imidazole-1-acetic acid, A-731
 2-Amino-4,5-dihydro-1*H*-imidazole-4-acetic acid, A-732
 6-Amino-6,7-dihydroimidazo[4,5-*d*]pyrimidin-7-one, P-820
 2-Amino-7,9-dihydro-6-methoxy-7-methyl-8*H*-purin-8-one, A-753
 2-Amino-3,5-dihydro-5-[(4-methoxyphenyl)methylidene]-3-methyl-4*H*-imidazol-4-one, L-152
 2-(5-Amino-4,7-dihydro-1-methyl-4,7-dioxo-1*H*-indol-3-yl)-4-thiazolecarboxamide, A-1126
 2-Amino-1,5-dihydro-1-methyl-4*H*-imidazol-4-one, C-732
 1-Amino-2,3-dihydro-2-methyl-1*H*-indole, D-472
 2-Amino-1,5-dihydro-7-methyl-4,6-pteridinedione, C-432
 6-Amino-3,7-dihydro-3-methyl-8*H*-purin-8-one, A-819
 6-Amino-1,3-dihydro-1-methyl-2*H*-purin-2-one, A-818
 6-Amino-7,9-dihydro-9-methyl-8*H*-purin-8-one, A-819
 2-Amino-1,7-dihydro-1-methyl-6*H*-purin-6-one, M-467
 2-Amino-1,7-dihydro-7-methyl-6*H*-purin-6-one, M-469
 2-Amino-3,7-dihydro-3-methyl-6*H*-purin-6-one, M-468
 2-Amino-8,9-dihydrooxazolo[4,5-*g*]pyrido[2,1-*c*][1,4]benzodiazepine-7,13(6*H*,7*aH*)-dione, I-20
 α -Amino-2,3-dihydro-3-oxo-5-isoxazoleacetic acid, I-8
 α -Amino-2,5-dihydro-5-oxo-4-isoxazolepropanoic acid, A-733
 2-Amino-4-(2,5-dihydro-5-oxo-2-isoxazolyl)butanoic acid, A-734
 2-Amino-1,4-dihydro-4-oxo-6-pteridinecarboxylic acid, A-817
N-[4-[(2-Amino-1,4-dihydro-4-oxo-6-pteridinyl)methyl]amino]benzoyl]glutamic acid, P-772
 4-[(2-Amino-1,4-dihydro-4-oxo-6-pteridinyl)methyl]formylaminobenzoic acid, R-82
 2-Amino-4,7-dihydro-4-oxo-7- β -D-ribofuranosyl-1*H*-pyrrolo[2,3-*d*]pyrimidine-5-carboximidamide, A-1387
 2-Amino-4,7-dihydro-4-oxo-7- β -D-ribofuranosyl-1*H*-pyrrolo[2,3-*d*]pyrimidine-5-carboxylic acid, C-7
 2-Amino-1,7-dihydro-7-(phenylmethyl)-6*H*-purin-6-one, G-210
 2-Amino-1,5-dihydro-4,6-pteridinedione, A-750
 2-Amino-5,8-dihydro-4,6,7(1*H*)-pteridinetriene, A-925
 2-Amino-7,9-dihydro-1*H*-purine-6,8-dione, A-753
 6-Amino-1,3-dihydro-2*H*-purin-2-one, A-818
 6-Amino-1,7-dihydro-8*H*-purin-8-one, A-819
 2-Amino-1,7-dihydro-6*H*-purin-6-one, G-210
 4-Amino-1,2-dihydro-3*H*-pyrazolo[3,4-*d*]pyrimidin-3-one, A-820
 5-Amino-1,6-dihydro-7*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7-one, A-735
 2-Amino-3',4'-dihydroxyacetophenone, A-736
 2-Amino-4,5-dihydroxybenzoic acid, A-737
 3-Amino-6,7-dihydroxy-2*H*-1-benzopyran-2-one, A-738
 4-Amino-2,5-dihydroxybenzyl alcohol, A-739
 26-Amino-3,16-dihydroxycholestan-22-one, A-740
 26-Amino-3,16-dihydroxycholest-5-en-22-one, A-741
 3-Amino-6,7-dihydroxycoumarin, A-738
 2-Amino-5-[[4,5-dihydroxy-2-cyclopenten-1-yl]amino]methyl]-1,7-dihydro-4*H*-pyrrolo[2,3-*d*]pyrimidin-4-one, Q-9
 2-Amino-5-(4,5-dihydroxycyclopent-1-en-3-ylaminomethyl)pyrrolo[2,3-*d*]pyrimidine, Q-9
 2-Amino-4,7-dihydroxy-6-(1,2-dihydroxypropyl)pteridine, I-13
 2-Amino-7-(1,2-dihydroxyethyl)-6-(methylthio)thieno[3,2-*g*]pteridin-4(3*H*)-one, U-48
 2-Amino-3,4-dihydroxy-5-methoxybenzoic acid, A-922
 2-Amino-1,4-dihydroxy-18-methyl-4-aza-3,7,14-nonadecanetriene, L-189
 4-Amino-2-(2,3-dihydroxy-3-methylbutyl)benzoic acid, A-742
 2-Amino-*N*,3-dihydroxy-*N*-(14-methyl-13,10-dioxopentadecyl)propanamide, L-189
 2-Amino-6,8-dihydroxy-7-methylpurine, A-753
 2-Amino-6,8-dihydroxy-9-methylpurine, A-753
 α -Amino- γ ,5-dihydroxy- β -methyl-2-pyridinebutanoic acid, A-796
 2-Amino-1,7-dihydroxyphenanthrene, A-743
 2-Amino-3-(3,4-dihydroxyphenyl)-2,3-dihydro-1,4-benzodioxin-6-ethanamine, A-696
 3-Amino-2-(3,4-dihydroxyphenyl)-2,3-dihydro-1,4-benzodioxin-6-ethanamine, A-744
 2-Amino-1-(3,4-dihydroxyphenyl)ethanol, N-298
 2-Amino-1-(3,4-dihydroxyphenyl)ethanone, A-736
 3-Amino-*N*-[2-(3,4-dihydroxyphenyl)ethyl]propanamide, D-157
 3-Amino-*N*-[2-(3,4-dihydroxyphenyl)ethyl]propanamide, A-241
 6-Amino-9-[1-(3,4-dihydroxyphenyl)ethyl]purine, A-137
 2-Amino-3-(3,4-dihydroxyphenyl)propanoic acid, A-745
 6-Amino-4,5-dihydroxy-3-piperidinecarboxylic acid, A-746
 3-Amino-11,12-dihydroxypregnan-20-one, A-747
 2-Amino-7-(1,2-dihydroxypropyl)-1,5-dihydro-4,6-pteridinedione, A-748
 2-Amino-6-(1,2-dihydroxypropyl)-3-methylpteridin-4-one, A-749
 2-Amino-6-(1,2-dihydroxypropyl)-4,7(1*H*,8*H*)-pteridinedione, I-13
 2-Amino-6-(1,2-dihydroxypropyl)-4(1*H*)-pteridinone, A-749
 2-Amino-6-(1,2-dihydroxypropyl)-5,6,7,8-tetrahydro-4(1*H*)-pteridinone, T-142
 2-Amino-4,6-dihydroxypteridine, A-750
 2-Amino-4,7-dihydroxypteridine, A-751
 2-Amino-4,7-dihydroxy-6-pteridinecarboxylic acid, A-752
 2-Amino-4,6-dihydroxy-7-pteridinecarboxylic acid, X-12
 6-Amino-2,8-dihydroxypurine, A-754
 2-Amino-6,8-dihydroxypurine, A-753
 4-Amino-3,6-dihydroxy-1*H*-pyrazolo[3,4-*d*]pyrimidine, A-755
 25-Amino-3,23-dihydroxystigmast-5-ene-7,22-dione, A-756
 3-Amino-6,7-dimethoxy-2*H*-1-benzopyran-2-one, A-738
 3-Amino-6,7-dimethoxycoumarin, A-738
 4-Amino-1,2-dimethoxy-3*H*-dibenzo[*de,g*]quinoline-3,5(6*H*)-dione, T-64
 1-Amino-2,3-dimethoxy-7*H*-dibenzo[*de,g*]quinolin-7-one, T-62
 6-Amino-8,9-dimethoxy-*N,N*-dimethyl-7-(methylthio)benzopentathiepin, L-205
 10-Amino-2,3-dimethoxy-4-hydroxyphenanthrene-1-carboxylic acid lactam, T-537
 10-Amino-2,4-dimethoxy-1-phenanthrenecarboxylic acid lactam, D-568
 2-Amino-7,8-dimethoxy-3*H*-phenoxazin-3-one, A-862
 2-Amino-1-(3,4-dimethoxyphenyl)ethanol, M-18
 3-Amino-20-dimethylaminopregnone, P-601
 7-Amino-1,6-dimethyl-5,8-isoquinolinedione, A-757
 2-Amino-14,16-dimethyl-3-octadecanol, A-758
 6-Amino-1,3-dimethyl-1*H*-purine-2,8(3*H*,8*H*)-dione, A-754
 2-Amino-*N*-(3,10-dioxohexadecyl)-*N*,3-dihydroxypropanamide, N-113
 1-Aminodiscorhabdin D, D-852
 6-Amino-1,2-dithiolo[4,3-*b*]pyrrolo-5(4*H*)-one, A-759
 2-Amino-3,5-dodecadiene, D-881
 2-Amino-5,7,9-dodecatrien-3-ol, D-345
 2-(12-Amino-3,7,9-dodecatrienyl)pyridine, A-760
 2-Amino-11-dodecen-3-ol, A-761
 4-Amino-2-dodecyl-3-hydroxytetrahydrofuran, A-762
 4-Amino-2-dodecyltetrahydro-3-furanol, A-762
 3-Amino-22,26-epimino-5 α -cholest-22(*N*)-en-16-ol, S-335
 3-Amino-22,26-epimino-16,23-epoxycholestan-23-ol, S-346
 2-Amino-5,6-epoxy-4-hydroxy-2-cyclohexen-1-one, A-804
 3-Amino-5,6-epoxy-4-hydroxy-6-hydroxymethyl-2-cyclohexen-1-one, C-289
 3-Amino-16,23-epoxy-16,28-secosolanidan-23-ol, S-346
 3-Aminoestra-1,3,5(10)-trien-17-ol, A-763
 2-Aminoethanesulfonic acid, A-764
 2-Aminoethanesulfonic acid, A-765
 2-Aminoethanesulfonothioic acid, A-766
 2-Aminoethanol, A-767
 3-Amino-5-ethenyl-5-hydroxy-2-cyclopenten-1-one, A-824
 4-(2-Aminoethyl)phenol, H-677
 2-Amino-5-ethoxybenzoic acid, A-789
 2-Amino-6-ethoxypurine, G-210
 2-Aminoethyl dihydrogen phosphate, A-768
S-(2-Aminoethyl) thiosulfate, A-766
N-(2-Aminoethyl)acetamide, E-225
N-(2-Aminoethyl)benzamide, E-225
 4-(2-Aminoethyl)-1,2-benzenediol, D-920
 α -(1-Aminoethyl)benzenemethanol, A-869
 β -Aminoethylbenzene, P-342
 5-(2-Aminoethyl)-1,2,3-benzenetriol, T-576
 9-(2-Aminoethyl)-6,7-benzopentathiepiindol, L-205
 3-(2-Aminoethyl)-6-bromo-2-(1,1-dimethyl-2-propenyl)-1*H*-indole, B-316
 3-(2-Aminoethyl)-6-bromoindole, B-356
 3-(2-Aminoethyl)-5-bromoindole, B-355
 4-(2-Aminoethyl)-2-bromophenol, A-769

- 3-(1-Aminoethyl)-5-carbomethoxy pyridine, A-774
 (2-Aminoethyl)cyclohexane, C-869
 Aminoethylcysteine ketimine decarboxylated dimer, H-241
 3-(2-Aminoethyl)-5,6-dibromoindole, D-343
 4-(2-Aminoethyl)-2,6-dibromophenol, D-322
N-[3-[4-(2-Aminoethyl)-2,6-dibromophenoxy]propyl]-7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxamide, A-1378
N-[3-[4-(2-Aminoethyl)-2,6-dibromophenoxy]propyl]- α -(hydroxyimino)benzenepropanamide, P-821
 2-Amino-9-ethyl-6,8-dihydroxypurine, A-753
 6-(2-Aminoethyl)-3,4-dimethoxybenzotriathiane, V-20
 9-(2-Aminoethyl)-6,7-dimethoxy-8-(methylthio)benzopentathiepin, L-205
 6-(2-Aminoethyl)-3,4-dimethoxy-5-(methylthio)benzotriathiane, V-20
 5-(2-Aminoethyl)-2,3-dimethoxyphenol, H-465
 4-(2-Aminoethyl)-2,6-dimethoxyphenol, T-576
 6-(2-Aminoethyl)-2,2-dimethyl-2*H*-1-benzopyran, A-770
 4-(2-Aminoethyl)-2,6-dinitrophenol, H-481
N-(2-Aminoethyl)dodecanamide, E-225
 4-(2-Aminoethyl)guaiaicol, H-566
 3-(2-Aminoethyl)-5-hydroxyindole, H-755
 4-(2-Aminoethyl)-1*H*-imidazole, H-307
 3-(2-Aminoethyl)-1*H*-indole-4,5-diol, D-683
 3-(2-Aminoethyl)-1*H*-indole-5,6-diol, D-684
 3-(2-Aminoethyl)indole, T-639
 3-(2-Aminoethyl)-1*H*-indol-5-ol, H-755
 2-(1-Aminoethyl)-*N*-[2-(1*H*-indol-3-yl)ethyl]-4-thiazolecarboxamide, A-1286
 2-(2-Aminoethyl)-3-isoxazolin-5-one, I-349
 5-(2-Aminoethyl)-3-methoxy-1,2-benzenediol, T-576
 9-(2-Aminoethyl)-6-methoxy-7-benzopentathiepinol, L-205
 5-(2-Aminoethyl)-2-methoxy-3,4-bis(methylthio)phenol, L-207
 3-(2-Aminoethyl)-5-methoxyindole, M-288
 4-(2-Aminoethyl)-2-methoxyphenol, H-566
 5-(2-Aminoethyl)-2-methoxyphenol, H-565
 5-(1-Aminoethyl)-2-methylpyridine, B-248
 3-(2-Aminoethyl)-2-nitroindole, A-771
 4-(2-Aminoethyl)-2-nitrophenol, H-640
 4-(2-Aminoethyl)phenol, H-667
 3-(2-Aminoethyl)phenol, H-678
 3-[4-(2-Aminoethyl)phenoxy]bulbispermine, H-5
 (2-Aminoethyl)phosphonic acid, A-772
 (2-Aminoethyl)phosphonic acid, A-773
 (2-Aminoethyl)phosphonic acid, A-772
 2-Amino-9-ethyl-6,8(1*H*,9*H*)-purinedione, A-753
 5-(1-Aminoethyl)-3-pyridinecarboxylic acid, A-774
 5-(2-Aminoethyl)pyrogallol, T-576
 3-Amino-2-ethyl-4(3*H*)-quinazolinone, E-247
 Aminoethylsulfonic acid, A-765
 3-(2-Aminoethyl)-2,4,6-tribromophenol, T-460
 1-(2-Aminoethyl)-3,6,7-trimethoxy-4-phenanthrenol, M-559
 4-(β -Aminoethyl)veratrole, D-718
 4-Amino-11-eudesmene, A-775
 6-Amino-4(15)-eudesmene, A-776
 1-Amino-2,6,10-farnesatriene, F-12
 Aminofornamide, G-199
N-[4-[(2-Amino-5-formyl-1,4,5,6,7,8-hexahydro-4-oxo-6-pteridiny)methyl]amino]benzoylglutamic acid, F-121
 2-Amino-3-formyl-6-hydroxypyridine, A-821
 2-Amino-3-formyl-6-methoxypyridine, A-821
 2-Amino-1-formyl-3-oxo-3*H*-phenoxazine-9-carboxylic acid, C-452
 2-Amino-9-formyl-3-oxo-3*H*-phenoxazine-1-carboxylic acid, C-452
 11-Amino-1(10),4-germacradiene, G-69
 α -Aminoglutaric acid lactam, O-206
 2-Aminoglutaric acid, G-107
 4-Aminoglyoxaline, A-826
 2-Aminoglyoxaline, A-825
 4-Amino-11-gorgonene, A-777
 14-Amino-1(5),7,10(14)-guaiaatriene-6,9-dione, A-778
 1-Amino-4-guanidinobutane, A-719
 1-Amino-19-guanidino-11,15-dihydroxy-4,9,12-triazanonadecane-10,13-dione, S-394
 2-Amino-6-guanidinohexanoic acid, H-326
 1-Amino-19-guanidino-11-hydroxy-4,9,12-triazanonadecane-10,13-dione, S-394
 1-Amino-2-(guanidinomethyl)-1-cyclopropanecarboxylic acid, C-163
 2-Amino-4-(guanidinooxy)butanoic acid, C-74
 1-Amino-5-guanidinopentane, A-859
 3-Amino-5-guanidinopentanoic acid, A-779
 1-Amino-20-guanidino-4,8,12,16-tetrazaicosane, A-780
 1-Amino-16-guanidino-4,8,12-triazahexadecane, A-781
 2-Amino-5-guanidinovaleric acid, A-1413
 1-Aminoheptane, H-130
 8-Amino-1,1,1,15,15,15-hexachloro-3,12-pentadecadiene, H-213
 2-Amino-11,15-hexadecadien-3-ol, A-783
 2-Amino-7,15-hexadecadien-3-ol, A-782
 2-Amino-3,5,10,13-hexadecatetraene, H-222
 2-Amino-11-hexadecen-3-ol, A-783
 2-Amino-15-hexadecen-3-ol, A-783
 2-Amino-9-hexadecen-3-ol, A-784
 3-(16-Amino-9-hexadecynyl)pyridine, N-224
 3-Amino-6-hydroxy-2*H*-azepin-2-one, A-785
 2-Amino-3,4,5,6,7,8-hexahydro- α -hydroxy-4,6-dioxo-7-pteridinepropanoic acid, E-216
N-[4-[(2-Amino-1,4,5,6,7,8-hexahydro-4-oxo-6-pteridiny)methyl]amino]benzoylglutamic acid, T-172
 1-Aminohexane, H-261
 2-Amino-4'-hydroxyacetophenone, A-786
 2-Amino-5-hydroxy-*p*-anisic acid, A-737
 1-Amino-2-hydroxybenzene, A-861
 β -Amino- α -hydroxybenzenepropanoic acid, A-808
 α -Amino-4-hydroxybenzenepropanoic acid, T-700
 2-Amino-3-hydroxybenzoic acid, A-787
 2-Amino-4-hydroxybenzoic acid, A-788
 2-Amino-5-hydroxybenzoic acid, A-789
 3-Amino-5-hydroxybenzoic acid, A-790
 2-Amino-5-(*p*-hydroxybenzyl)-4-(*p*-methoxybenzyl)-1-methylimidazole, N-3
 2-Amino-4-(4-hydroxybenzyl)-1-methyl-1*H*-imidazole, A-791
 3-Amino-4-hydroxybutanoic acid, A-792
 1-Amino-3-hydroxy-8-decen-5-one, A-793
 2-Amino-4-hydroxy-6-(1,2-dihydroxypropyl)pterin, A-749
 2-Amino-6-hydroxy-7,9-dimethylpurinium, H-143
 β -Amino- α -hydroxyethylbenzene, A-864
 4-(2-Amino-1-hydroxyethyl)-1,2-benzenediol, N-298
 4-(β -Amino- α -hydroxyethyl)catechol, N-298
p-(β -Amino- α -hydroxyethyl)phenol, O-63
 (2-Amino-1-hydroxyethyl)phosphonic acid, A-794
 2-Amino-6-hydroxy-8-(4-hydroxybenzoyl)-3*H*-phenoxazin-3-one, A-707
 4-Amino-5-hydroxy-1-hydroxymethyl-7-oxabicyclo[4.1.0]hept-3-en-2-one, C-289
 2-Amino-4-hydroxy-6-(hydroxymethyl)pteridine, A-802
 4-Amino-2-hydroxy-5-(hydroxymethyl)pyrimidine, A-795
 2-Amino-4-hydroxy-6-(3-hydroxy-4-phosphonoxy-1-butenyl)pteridine, A-809
 2-Amino-4-hydroxy-4-(5-hydroxy-2-pyridinyl)-3-methylbutanoic acid, A-796
 3-Amino-2-hydroxy-2-(1*H*-imidazol-4-ylmethyl)-5-methylhexanoic acid, L-160
 2-Amino-3-hydroxy-3-(4-imidazolyl)propanoic acid, A-797
 2-Amino-4-hydroxy- α -imino- $\Delta^{7(3*H*)\beta}$ -pteridinepropionic acid, L-109
 2-Amino-5-hydroxy-4-methoxybenzoic acid, A-737
 2-[4-Amino-3-hydroxy-5-methoxy-2,6-bis(3-methyl-2-butenyl)phenyl]-5,7-dihydroxy-4*H*-1-benzopyran-4-one, P-483
 2-Amino-8-hydroxy-6-methoxy-7-methylpurine, A-753
 2-Amino-1-(4-hydroxy-3-methoxyphenyl)ethanol, N-303
 2-Amino-5-(hydroxymethyl)-1,4-benzenediol, A-739
 4-Amino-5-(hydroxymethyl)-2-methoxypyrimidine, A-795
 3-Amino-4-hydroxymethyl-14-methyl-9,19-cyclopregn-4-en-20-one, A-798
 7-Amino-1-hydroxymethyl-6-methyl-5,8-isoquinolinedione, A-799
 2-Amino-9-hydroxymethyl-3-oxo-3*H*-phenoxazine-1-carboxylic acid, C-452
 2-Amino-1-hydroxy-4-methylpentane, A-838
 2-Amino-4-hydroxy-3-methylpentanoic acid, A-800
 2-Amino-8-(hydroxymethyl)-3*H*-phenoxazin-3-one, A-801
 2-Amino-6-(hydroxymethyl)-4(1*H*)-pteridinone, A-802
 4-Amino-2-hydroxy-5-methylpyrimidine, A-803
 4-Amino-5-hydroxymethyl-2(1*H*)-pyrimidinone, A-795
 3-Amino-5-hydroxy-7-oxabicyclo[4.1.0]hept-3-en-2-one, A-804
 8-Amino-1-hydroxyphenazine, A-805
 2-Amino-6-hydroxy-3*H*-phenoxazin-3-one, A-862
N-[3-[2-Amino-3-(4-hydroxyphenoxy)phenyl]-3-oxopropyl]-4-hydroxybenzamide, B-289
 2-Amino-1-(4-hydroxyphenyl)ethanol, O-63
 2-Amino-1-(4-hydroxyphenyl)ethanone, A-786
 2-Amino-5-[[[3-(4-hydroxyphenyl)-2-methoxy-2-propenoyl]amino](imino)methyl]amino]pentanoic acid, A-806
 1-(3-Amino-4-hydroxyphenyl)-1,2-propanediol, A-807
 3-Amino-2-hydroxy-3-phenylpropanoic acid, A-808
 2-Amino-3-(4-hydroxyphenyl)propanoic acid, T-700
 2-Amino-6-[3-hydroxy-4-(phosphonoxy)-1-butenyl]-4(1*H*)-pteridinone, A-809
 20-Amino-18-hydroxypregna-1,4-dien-3-one, A-810
 3-Amino-20-hydroxypregnan-20-one, A-880
 3-Amino-12-hydroxypregnan-20-one, A-811
 3-Amino-14-hydroxypregnan-20-one, A-812
 20-Amino-3-hydroxypregna-5-ene, A-886
 3-Amino-12-hydroxypregna-15-en-20-one, A-811
 3-Amino-15-hydroxypregna-5-en-20-one, A-888
 3-Amino-2-hydroxypregna-5-en-16-one, A-813
 3-Amino-2-hydroxypregna-5-en-20-one, A-814
 2-Amino-3-hydroxy-1-propanesulfonic acid, A-815
 2-Amino-6-(1-hydroxypropyl)-8-methyl-4,7(1*H*,8*H*)-pteridinedione, I-13
 2-Amino-4-hydroxypteridine, A-816
 2-Amino-4-hydroxy-6-pteridinecarboxylic acid, A-817
 3-[2-Amino-4-hydroxy-7(3*H*)-pteridinyldene]-2-iminopropanoic acid, L-109
 6-Amino-2-hydroxypurine, A-818
 6-Amino-8-hydroxypurine, A-819
 2-Amino-6-hydroxypurine, G-210
 4-Amino-3-hydroxy-1*H*-pyrazolo[3,4-*d*]pyrimidine, A-820
 2-Amino-6-hydroxy-3-pyridinecarboxaldehyde, A-821
 α -Amino-5-hydroxy-2-pyridinepropanoic acid, A-822
 α -Amino-3-hydroxy-1(4*H*)-pyridinepropanoic acid, L-148
 2-Amino-4-hydroxypyrimidine, A-823
 4-Amino-2-hydroxypyrimidine, C-961
 4-Amino-2-hydroxy-5-pyrimidinemethanol, A-795
 3-Amino-2-hydroxy-4(3*H*)-quinazolinone, Q-13
 3-Amino-23-hydroxysolanidane, A-908
N-(20-Amino-4-hydroxy-4,8,12,17-tetrazaeicos-1-yl)-4-hydroxybenzamide, A-176
 2-Amino-4-hydroxy-6-(1,2,3,4-tetrahydroxybutyl)pteridine, A-915
 3-Amino-5-hydroxy-5-vinyl-2-cyclopenten-1-one, A-824
 2-Aminohypoxanthine, G-210
 1-Aminohypoxanthine, P-820
 5-Aminoimidazole ribonucleotide, A-827

- 5-Aminoimidazole riboside, A-827
 4(5)-Aminoimidazole, A-826
 2-Aminoimidazole, A-825
 4,4'-(2-Amino-1*H*-imidazole-1,4-diyl)bis(methylene)bisphenol, I-252
 β -Amino-1*H*-imidazole-4-propanol, H-308
 2-[2-Amino-2-imidazol-4-yl]acetic acid, A-732
N-[2-(2-Amino-1*H*-imidazol-4-yl)ethyl]-3,5-dibromo-2-hydroxy- α -(hydroxyimino)-4-methoxybenzenepropanamide, P-814
N-[2-(2-Amino-1*H*-imidazol-4-yl)ethyl]-3-(3,5-dibromo-4-methoxyphenyl)-2-propenamamide, A-828
 3-Amino-*N*-[2-(1*H*-imidazol-4-yl)ethyl]propanamide, C-157
 2-Amino-3-(4-imidazolyl)propanol, H-308
 3-(2-Amino-1*H*-imidazol-4-yl)-2-propen-1-amine, A-701
 2-Amino-6-[[3-(1*H*-imidazol-4-yl)-2-propenoyl]amino]hexanoic acid, I-42
N-[3-(2-Amino-1*H*-imidazol-4-yl)-2-propenyl]-4,5-dibromo-1*H*-pyrrole-2-carboxamide, O-123
 2-Amino-1*H*-imidazo[4,5-*b*]pyrazine, A-829
 3-Amino-22,26-imincholestane-16,23-diol, S-336
N-(Aminoiminomethyl)- β -alanine, G-208
 [(Aminoiminomethyl)amino]acetic acid, G-200
 7-(Aminoiminomethyl)amino]-*N*-[2-[[4-[(3-amino-propyl)amino]butyl]amino]-1-hydroxy-2-oxoethyl]heptanamamide, S-394
 2-[4-[(Aminoiminomethyl)amino]-1-azabicyclo[3.1.0]hex-2-yl]-*N*-valylglycine, F-52
 4-[(Aminoiminomethyl)amino]butanoic acid, G-201
 3-[[[1-[[4-[(Aminoiminomethyl)amino]butyl]amino]carbonyl]-3-methylbutyl]amino]carbonyl]oxiranecarboxylic acid, R-69
 3-[[[4-[(Aminoiminomethyl)amino]butyl]amino]propyl]guanidine, H-305
N-[4-[(Aminoiminomethyl)amino]butyl]-3,5-dibromo-2-hydroxy- α -(hydroxyimino)-4-methoxybenzenepropanamide, P-815
 4-[[[(Aminoiminomethyl)amino]carbonyl]amino]butanoic acid, G-140
N-[4-[[[(Aminoiminomethyl)amino]carbonyl]amino]butyl]-*N*-methylacetamide, N-187
*N*²-[[[(Aminoiminomethyl)amino]carbonyl]ornithine, G-75
 β -(Aminoiminomethyl)amino-3,4-dihydroxystyrene, T-656
 2-[(Aminoiminomethyl)amino]ethanesulfinic acid, G-203
 2-[(Aminoiminomethyl)amino]ethanesulfonic acid, G-204
O-[(Aminoiminomethyl)amino]homoserine, C-74
 4-[(Aminoiminomethyl)amino]-3-hydroxybutanoic acid, G-206
 3-[(Aminoiminomethyl)amino]-*N*-(2-hydroxyheptyl)propanamide, P-313
 3-[(Aminoiminomethyl)amino]-*N*-(2-methoxyheptyl)-2-methylpropanamide, P-314
 5-[(Aminoiminomethyl)amino]-3-(methylamino)pentanoic acid, A-779
 5-[(Aminoiminomethyl)amino]-2-oxopentanoic acid, G-207
N-(Aminoiminomethyl)-4-bromo-1*H*-pyrrole-2-carboxamide, B-319
*N*⁶-(Aminoiminomethyl)-*N*²-[4-bromo-1*H*-pyrrol-2-yl]carbonyl]lysine, B-346
N-[*N*-(Aminoiminomethyl)-2-[3,5-dihydroxy-4-(1-methylethyl)phenyl]glycyl]glycine, R-54
N-[*N*-(Aminoiminomethyl)-2-(4-ethyl-3,5-dihydroxyphenyl)glycyl]glycine, R-55
N-(Aminoiminomethyl)glycine, G-200
 3-(Aminoiminomethyl)-1,2,3,7,8,9a-hexahydro-5*H*,10*H*-dipyrrolo[1,2-*a*:1',2'-*a'*]pyrazine-5,10-dione, V-79
*N*⁶-(Aminoiminomethyl)lysine, H-326
 4-[(Aminoiminomethyl)(3-methyl-2-butenyl)amino]butanoic acid, C-592
N-(Aminoiminomethyl)-*N*-methylglycine, C-731
 3-[3-(Aminoiminomethyl)propyl]hexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione, C-841
 [3-(Aminoiminomethyl)propylidene]butanedioic acid, L-373
N-(Aminoiminomethyl)-2-pyrrolidineaetic acid, P-950
 3-(3-Amino-1*H*-inden-2-yl)alanine, A-698
 α -Amino-1*H*-indole-3-propanoic acid, T-640
 β -Amino-1*H*-indole-3-propanol, A-830
 2-Amino-*N*-[[2-[(1*H*-indol-3-yl)ethenyl]amino]-2-oxoethyl]benzamide, P-189
 2-Amino-5-(1*H*-indol-3-ylmethyl)-4(5*H*)-oxazolone, A-1092
 2-Amino-3-(3-indolyl)propanoic acid, T-640
 2-Amino-3-(3-indolyl)-1-propanol, A-830
 9-Aminoisoascididemnin, H-542
 2-Aminoisocaproic acid, L-153
 7-Amino-5,8-isoquinolinedione, A-831
 7-Amino-5,8-isoquinolinequinone, A-831
 4-Amino-3-isoxazolidinone, A-832
 2-Amino-4-(isoxazolin-5-on-2-yl)butanoic acid, A-734
 α -Amino- γ -(isoxazolin-5-on-2-yl)butyric acid, A-849
o-Aminomandelic lactam, I-91
 Aminomethanamide, G-199
 Aminomethane, M-383
 2-Amino-4'-methoxyacetophenone, A-786
 2-Amino-3-methoxybenzoic acid, A-787
 2-Amino-4-methoxybenzoic acid, A-788
 2-Amino-5-methoxybenzoic acid, A-789
 2-Amino-3-methoxybenzonitrile, A-787
 2-Amino-5-methoxybenzonitrile, A-789
 3-Amino-5-methoxybenzonitrile, A-790
 1-Amino-4-methoxybutane, A-716
 6-Amino-5-methoxy-7*H*-dibenzof[*de,h*]quinolin-7-one, L-19
 1-Amino-2-methoxy-7*H*-dibenzof[*de,g*]quinolin-7-one, T-62
 2-Amino-6-methoxy-7,9-dimethylpurinium(1+), H-143
 4-Amino-3-[3-(methoxymethyl)-6,7-dimethyl-2,6-octadienyl]benzamide, B-68
 10-Amino-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid lactam, A-1427
 2-Amino-*N*-[2-(methoxymethyl)phenyl]benzethanamine, A-708
 8-Amino-1-methoxyphenazine, A-805
 4-Amino-3-methoxyphenol, A-705
 4-[[2-Amino-4-[(4-methoxyphenyl)methyl]-1-methyl-1*H*-imidazol-5-yl]methyl]phenol, N-3
 2-Amino-6-methoxy-3-pyridinecarboxaldehyde, A-821
 4-Amino-2-methoxypyrimidine, C-961
 4-Amino-2-methoxy-5-pyrimidinemethanol, A-795
 8-Amino-4-methylaconitane-1,3,6,13,14,16,18-heptol, A-833
 8-Amino-4-methylaconitane-1,3,6,13,14,15,16,18-octol, A-834
 2-Amino-4-(methyl-*NNO*-azoxy)butanoic acid, A-1616
 α -Aminomethylbenzenemethanol, A-864
 6-(Aminomethyl)-1,2,4-benzenetriol, T-528
 α -(Aminomethyl)benzyl alcohol, A-864
 1-Amino-3-methyl-2,3-butanediol, A-835
 1-Amino-3-methylbutane, M-420
 α -(1-Amino-3-methylbutyl)- α -hydroxy-1*H*-imidazole-4-propanoic acid, L-160
 5-(Aminomethyl)-4-(carboxymethyl)-1*H*-pyrrole-3-propanoic acid, P-574
 7-Aminomethyl-7-deazaguanine, A-699
 α -(Aminomethyl)-3,4-dihydroxybenzyl alcohol, N-298
 α -(Aminomethyl)-3,4-dimethoxybenzenemethanol, M-18
 6-Amino-4-methyl-1,2-dithiolo[4,3-*b*]pyrrol-5(4*H*)-one, A-759
 3-(12-Amino-10-methyldodecyl)pyridine, N-226
 3-(12-Amino-11-methyldodecyl)pyridine, N-225
 3-(Aminomethylene)-1,3-dihydro-2*H*-indole-2-thione, A-836
 3-(Aminomethylene)dihydro-2*H*-pyran-2,4(3*H*)-dione, G-62
 10-Amino-3,4-methylenedioxy-1-phenanthrenecarboxylic acid lactam, D-567
 3-Aminomethyleneindole-2-thione, A-836
 3-Amino-2-methylenepropanoic acid, A-841
 2-(1-Amino-1-methylethyl)-7*H*-furo[3,2-*g*][1]benzopyran-7-one, P-585
 α -(Aminomethyl)-4-hydroxybenzenemethanol, O-63
 α -(Aminomethyl)-*p*-hydroxybenzyl alcohol, O-63
 α -(Aminomethyl)-4-hydroxy-3-methoxybenzenemethanol, N-303
 5-Amino-1-methyl-1*H*-imidazole, A-826
 2-Amino-1-methyl-2-imidazol-4-one, C-732
 4-[(2-Amino-1-methyl-1*H*-imidazol-4-yl)methyl]phenol, A-791
 3-(Aminomethyl)indole, I-97
 1-Amino-2-methylindoline, D-472
 5-(Aminomethyl)-3(2*H*)-isoxazolone, A-837
 α -(Aminomethyl)-4-methoxybenzenemethanol, O-63
 2-Amino-1-methyl-4-(3,4-methylenedioxybenzyl)-1*H*-imidazole, P-594
 3-(Aminomethyl)-1-methylindole, I-97
 7-Amino-2-methyl-6-(methylthio)-3,5,8(2*H*)-isoquinolinetriene, P-255
 (S)-2-Amino-4-methyl-*N*-2-naphthalenylpentanamide, L-153
 1-(2-Amino-3-methyl-1-oxopentyl)-2,3,3*a*,12*a*,13,14,15,17*a*-octahydro-5,8-ethenodipyrrolo[3,2-*b*:3',2'-*e*][1,5,8]-oxadiazacyclotetradecene-12,17(1*H*,11*H*)-dione, Z-24
 2-Amino-4-methylpentanamide, L-153
 2-Amino-4-methylpentanoic acid, L-153
 2-Amino-4-methyl-1-pentanol, A-838
 4-Amino-4-methyl-2-pentanone, A-839
 4-(Aminomethyl)phenol, H-417
 2-(Aminomethyl)phenol, H-415
 3-(Aminomethyl)phenol, H-416
 (Aminomethyl)phenylcarbinol, A-864
 1-Amino-2-methyl-2-propanol, A-840
 2-(Aminomethyl)-2-propanoic acid, A-841
 2-Amino-1-methyl-4(1*H*)-pteridinone, A-816
 6-Amino-1-methylpurine, A-842
 6-Amino-2-methylpurine, A-843
 6-Amino-3-methylpurine, A-844
 2-Amino-7-methyl-6,8-purinediol, A-753
 2-Amino-9-methyl-6,8-purinediol, A-753
 2-Amino-4-methylpyridine, A-845
 4-Amino-5-methyl-2-pyrimidinol, A-803
 4-Amino-5-methyl-2(1*H*)-pyrimidinone, A-803
 4-Amino-1-methyl-2(1*H*)-pyrimidinone, C-961
 6-Amino-1-methyl-2(1*H*)-pyrimidinone, C-961
 3-[(4-Amino-2-methyl-5-pyrimidinyl)methyl]-5-(2-hydroxyethyl)-4-methylthiazolium(1+), T-366
N-[(4-Amino-2-methyl-5-pyrimidinyl)methyl]-*N*-[4-hydroxy-1-methyl-2-(2-propenylthio)-1-butenyl]formamide, A-627
 3-Amino-*N*^b-methylsecovoacarpine, A-846
 1-Amino-12-methyltetradecane, M-566
 2-Amino- α -(methylthio)- γ -oxobenzenebutanoic acid, T-381
 1-Amino-12-methyltridecane, M-579
 α -(Aminomethyl)-3,4,5-trimethoxybenzenemethanol, H-557
 α -(Aminomethyl)vanillyl alcohol, N-303
 4-Aminomimosamycin, M-624
 4-Aminomorpholine, M-705
 14-Aminomyristic acid, A-911
 2-Amino-3-(4-nitro-1*H*-indol-3-yl)propanoic acid, N-257
 9-Aminononyl methyl sulfone, M-564
 3-Amino-23-norcon-5-ene, C-595
 2-Amino-5,9,12,15,17-octadecapentaen-3-ol, A-847
 2-Amino-6,9,12,15-octadecapentaen-3-ol, A-848
 2-Amino-5,9,12,15-octadecapentaen-3-ol, A-847
 1-Amino-2,4-octadiene, O-38
 6-Amino-octahydro-1-indolizol acetate, S-327
 4-(2-Amino-4-oxo-2-imidazolidin-5-ylidene)-2-bromo-4,5,6,7-tetrahydropyrrolo[2,3-*c*]azepin-8-one, H-762
 α -Amino-5-oxo-2(5*H*)-isoxazolebutanoic acid, A-849
 α -Amino-5-oxo-2(5*H*)-isoxazolepropanoic acid, A-850

- α -Amino-3-oxo-5-isoxazolidineacetic acid, T-490
 α -Amino-5-oxo-3-isoxazoline-4-propionic acid, A-733
 8-Amino-7-oxononanoic acid, A-851
 α -Amino-5-oxo-3-oxazoline-2-propionic acid, A-850
 2-Amino-3-oxo-3*H*-phenoxazine-1-carboxylic acid, A-852
 2-Amino-3-oxo-3*H*-phenoxazine-8-carboxylic acid, A-853
 2-Amino-3-oxo-3*H*-phenoxazine-1,8-dicarboxylic acid, A-854
 2-Amino-3-oxo-3*H*-phenoxazine-1,9-dicarboxylic acid, C-452
 1-(2-Amino-1-oxopropyl)-2,3,3a,13a,14,15,16,18a-octahydro-8-methoxy-5,9-metheno-9*H*-dipyrrolo[3,2-*b*:1',2'-*e*] [1,5,8]oxadiazacyclopentadecine-13,18(1*H*,12*H*)-dione, A-855
 α -Amino-*N*-[2-oxo-2-[[1-(1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-2-methyl-1-isoquinolinyl)ethylamino]ethyl]benzenepropanamide, P-339
 22-Amino-2,7,11,15,19-pentaazadocosanimidamide, A-780
 1-Aminopentadecane, P-203
 5-Amino-2-(1,3,8,10-pentadecatetraenyl)-4-piperidinol, P-713
 2-Amino-6-(1,2,3,4,5-pentahydroxypentyl)-4(1*H*)-pteridinone, A-856
 2-Aminopentanedioic acid, G-107
 2-Amino-3-pentanol, A-857
 2-Amino-3-pentanone, A-858
 5-Amino-2-pentenoic acid γ -lactam, D-495
N-[3-[[3-[[3-(5-Aminopentyl)amino]propyl]amino]propyl]amino]propyl]-4-hydroxy-1*H*-indole-3-acetamide, C-536
 (5-Aminopentyl)guanidine, A-859
 2-Amino-1,7-phenanthrenediol, A-743
 7-Amino-1-phenazinecarboxylic acid, A-860
 8-Amino-1-phenazolinol, A-805
 2-Aminophenol, A-861
 2-Amino-3*H*-phenoxazin-3-one, A-862
 2-[(4-Aminophenyl)amino]pyrimidine, A-863
 2-Amino-1-phenylethanol, A-864
 2-Amino-1-phenylethanone, A-694
 3-Amino-3-phenyllactic acid, A-808
 2-[[2-(Aminophenyl)methyl]amino]benzenemethanol, A-708
 1-[(2-Aminophenyl)methyl]-3-hydroxy-2-pyrrolidinone, V-39
 4-[(5-Amino-6-phenylmethyl)pyrazinyl]phenol, C-559
 4-(2-Aminophenyl)-2-(methylthio)-4-oxobutanoic acid, T-381
 1-Amino-4-phenyl-2,7-naphthyridine, A-865
 5-(4-Aminophenyl)-2,4-pentadienamide, A-866
 5-(4-Aminophenyl)-2,4-pentadienoic acid, A-866
 4-Amino-1-phenyl-1-penten-3-ol, A-867
 4-Amino-1-phenyl-1-penten-3-one, A-868
 2-Amino-1-phenyl-1-propanol, A-869
 2-Amino-1-phenyl-1-propanone, A-870
 5-(3-Amino-3-phenylpropanoyl)nicotaxine, N-188
 2-Amino- γ -picoline, A-845
 4-Aminopipicolinic acid, A-871
 4-Amino-2-piperidinecarboxylic acid, A-871
 3-Amino-2-piperidinone, A-872
 20-Aminopregna-5,16-diene-2,3-diol, A-874
 3-Aminopregna-5,17(20)-diene-2,16-diol, A-873
 20-Aminopregna-5,16-dien-3-ol, A-876
 3-Aminopregna-5,16-dien-20-ol, A-875
 20-Aminopregna-1,4-dien-3-one, A-878
 3-Aminopregna-5,17(20)-dien-16-one, A-877
 20-Aminopregnane-3,4-diol, A-879
 20-Aminopregnan-3-ol, A-881
 3-Aminopregnan-20-ol, A-880
 20-Aminopregnan-3-one, A-883
 3-Aminopregnan-20-one, A-882
 20-Aminopregn-5-ene-3,16-diol, A-884
 20-Aminopregn-5-en-3-ol, A-886
 3-Aminopregn-5-en-20-ol, A-885
 3-Aminopregn-5-en-18,20-olide, P-93
 20-Aminopregn-4-en-3-one, A-891
 3-Aminopregn-16-en-20-one, A-889
 3-Aminopregn-17(20)-en-16-one, A-890
 3-Aminopregn-5-en-16-one, A-887
 3-Aminopregn-5-en-20-one, A-888
 6-Amino-3-prenylpurine, T-451
 4-Aminopropine, A-905
 3-Aminopropanal, A-892
 2-Amino-1,3-propanediol, A-893
 1-Aminopropane, P-648
 2-Amino-1-propanol, A-894
 4-(3-Amino-1-propenyl)-1*H*-imidazol-2-amine, A-701
 3-Aminopropionaldehyde, A-892
 α -Aminopropiophenone, A-870
 4-(3-Aminopropoxy)-3,5-dibromobenzeneethanamine, M-679
 (3-Aminopropoxy)guanidine, A-895
 3-Aminopropyl methyl sulfide, M-578
*N*¹-(3-Aminopropyl)-*N*⁴-acetyl-1,4-butanediamine, S-398
N-[4-[(3-Aminopropyl)amino]butyl]acetamide, S-398
*N*¹-[3-[[4-[(3-Aminopropyl)amino]butyl]amino]propyl]bleomycinamide, B-236
N-[[[4-[(3-Aminopropyl)amino]butyl]carbonyl]hydroxymethyl]-7-guanidinoheptanamide, S-394
N-[5-[[3-[[3-[(3-Aminopropyl)amino]propyl]amino]propyl]amino]pentyl]-4-hydroxy-1*H*-indole-3-acetamide, C-535
N-(3-Aminopropyl)-*N*'-[3-[(3-aminopropyl)amino]propyl]-1,4-butanediamine, D-289
N-[3-[[3-[(3-Aminopropyl)amino]propyl]amino]propyl]-1,4-butanediamine, D-288
N-(3-Aminopropyl)-*N*'-[3-[(3-aminopropyl)amino]propyl]-1,3-propanediamine, T-456
N'-[3-[(3-Aminopropyl)amino]propyl]-1,4-butanediamine, D-249
 4-(3-Aminopropyl)-1,2-benzenediol, A-896
N-(3-Aminopropyl)-1,4-butanediamine, S-398
N-(3-Aminopropyl)cadaverine, A-897
N'-(3-Aminopropyl)canavalmine, D-287
 5-(3-Aminopropyl)-1,9-diamino-5-azanonane, A-718
*N*²-(3-Aminopropyl)homospermidine, A-718
*N*¹-(3-Aminopropyl)homospermidine, D-250
N-[3-[(3-Aminopropyl)methylamino]propyl]-1,4-butanediamine, D-249
N-(3-Aminopropyl)-*N*-methyl-1,4-butanediamine, S-398
N-(3-Aminopropyl)-*N*-methyl-1,3-propanediamine, D-255
*N*⁴-(3-Aminopropyl)norspermidine, T-622
 10-(3-Aminopropyl)octadecahydro-2*H*-pyrido[1,2-*f*][1,6]diazacyclooctadecine-11(6*H*)-one, N-130
N-(3-Aminopropyl)-1,5-pentanediamine, A-897
p- β -Aminopropylphenol, A-898
 4-(2-Aminopropyl)phenol, A-898
 4-(3-Aminopropyl)phenol, H-350
N-(3-Aminopropyl)-1,3-propanediamine, D-255
*N*⁴-(3-Aminopropyl)spermidine, B-152
 7'-*N*-(3-Aminopropyl)squalamine, S-485
 6-Aminoprotocatechuic acid, A-737
 2-Amino-4,7(1*H*,8*H*)-pteridinedione, A-751
 2-Amino-4-pteridinol, A-816
 2-Amino-4(1*H*)-pteridinone, A-816
 2-Amino-4(3*H*)-pteridinone-6-carboxylic acid, A-817
 Aminopteroic acid, A-899
 6-Aminopurine, A-137
 6-Aminopurine-2,8-diol, A-754
 2-Aminopurine-6,8-diol, A-753
 6-Amino-1*H*-purine-2,8(3*H*,7*H*)-dione, A-754
 6-Amino-9*H*-purine-9-propanoic acid, A-900
 6-Amino-8-purinol, A-819
 2-Amino-6-purinol, G-210
 3-(6-Amino-9*H*-purin-9-yl)-1-(hydroxymethyl)-6-oxabicyclo[3.1.0]hexane-2,4-diol, N-168
 4-(6-Amino-9*H*-purin-9-yl)-4-methyl-2-pentanone, G-15
 6-[[6-Amino-9*H*-purin-9-yl]methyl]-3-pyridinol, P-151
 4-Amino-1*H*-pyrazolo[3,4-*d*]pyrimidine-3,6(2*H*,5*H*)-dione, A-755
 1-*C*-(7-Amino-1*H*-pyrazolo[4,3-*d*]pyrimidin-3-yl)-1,4-anhydro-*D*-ribitol, F-131
 4-Amino-2,3-pyridinedicarboxylic acid, A-901
 α -Amino-2-pyridinepropanoic acid, A-902
 2-Amino-3-(2-pyridinyl)propanoic acid, A-902
 9-Aminopyrido[1,2-*d*]indole-6,10-dione, A-208
 2-Amino-4-pyrimidinecarboxylic acid, A-903
 2-Amino-4-pyrimidinol, A-823
 2-Amino-4(1*H*)-pyrimidinone, A-823
 4-Amino-2(1*H*)-pyrimidinone, C-961
 1-(2-Amino-4-pyrimidinyl)- β -carboline, A-1051
 1-(2-Amino-4-pyrimidinyl)-6-methoxy- β -carboline, A-1051
 α -Amino-1*H*-pyrrole-1-hexanoic acid, A-904
 4-Amino-2-pyrrolidinecarboxylic acid, A-905
 1-Aminopyrrolidine, P-949
 1-Aminopyrrolizidine, A-906
 Aminopyrrolitrin, A-725
 3-Amino-2,4(1*H*,3*H*)-quinazolinedione, Q-13
 2-Aminoquinoline, Q-28
 α -Aminoquinoline, Q-28
 1-Aminoquinolizidine, A-907
 4-Aminoresorcinol, A-705
 4-Amino-7- β -*D*-ribofuranosyl-7*H*-pyrrolo[2,3-*d*]pyrimidine-5-carbonitrile, S-53
 4-Amino-7- β -*D*-ribofuranosyl-7*H*-pyrrolo[2,3-*d*]pyrimidine-5-carboxylic acid, S-53
 4-Amino-7-ribofuranosyl-7*H*-pyrrolo[2,3-*d*]pyrimidine, T-658
 3-Amino-16,28-secosolanida-22,24,26(28)-triene-16,23-diol, S-351
 3-Amino-16,28-secosolanid-22(28)-en-16-ol, S-335
 Aminosin, I-148
 3-Aminosolanidan-23-ol, A-908
 6-Amino-1-spiroxene, S-425
 3-Aminospirosolane, S-439
 3-Aminospirosol-5-ene, S-446
 3-Aminospirostane, S-458
 3-Aminospirostan-6-ol, A-909
 3-Aminospirostan-9-ol, A-910
N-[2-(Aminosulfonyl)ethyl]-3-bromo-4-hydroxy- α -(hydroxymino)benzenepropanamide, P-688
 18-Amino-2,7,11,15-tetraazaoctadecanimidamide, A-781
 2-Amino-3,5-tetradecadiene, T-131
 14-Aminotetradecanoic acid, A-911
 2-Amino-3,5,13-tetradecatriene, T-131
 2-Amino-13-tetradecen-3-ol, A-913
 2-Amino-5-tetradecen-3-ol, A-912
 3-(14-Amino-3-tetradecenyl)pyridine, T-357
 3-(11-Amino-3-tetradecenyl)pyridine, T-357
 7-Amino-3,4,4a,10*b*-tetrahydro-2,4-dimethylbenzo[*f*]quinolin-6(5*H*)-one, A-914
 2-Amino-1,4,7,8-tetrahydro-4,7-dioxo-6-pteridinecarboxylic acid, A-752
 2-Amino-1,4,5,6-tetrahydro-4,6-dioxo-7-pteridinecarboxylic acid, X-12
 3-(2-Amino-4,5,6,8-tetrahydro-4,6-dioxo-7(3*H*)-pteridinylidene)-2-oxopropanoic acid, E-216
 2-Amino-3a,5,6,6a-tetrahydro-4-(hydroxymethyl)-4*H*-cyclopentoxazole-4,5,6-triol, T-447
 7-Amino-3,4,4a,10*b*-tetrahydro-2-(hydroxymethyl)-4-methylbenzo[*f*]quinolin-6(5*H*)-one, A-914
 3-Amino-1,2,5,6-tetrahydro-5-imino-2,6-pyrazine-dibutanamine, B-35
 α -Amino-3,4,5,6-tetrahydro-3-methyl-2-(methylamino)-4-pyrimidineacetic acid, S-527
 6-Amino-1,2,3,4-tetrahydro-4-oxo-1,3,5-triazine-2-carboxylic acid, T-104
 2-Amino-6-(1,2,3,4-tetrahydroxybutyl)-4(1*H*)-pteridinone, A-915
 16-Amino-2,6,10,14-tetramethyl-5,10,14-hexadecatrien-4-one, A-916
 α -Aminotoluene, B-84
 1-Aminotriacotane, T-452
 3-Amino-2,4,6-triazabicyclo[3.2.1]oct-2-ene-7-carboxylic acid, V-144
 5-Amino-1,1,1-trichloroundecane, T-482
 1-Amino-4,12-tridecadien-2-ol, A-917
 5-Amino-2-(6,8-tridecaadienyl)-4-piperidinol, P-712
 6-Aminotridecane, T-506
 1-Amino-3-tridecen-2-ol, A-918
 1-Amino-5-tridecen-2-ol, A-919
 5-(12-Aminotridecyl)hexahydropyrrolo[2,1-*b*]oxazole, A-920

- 3-(13-Aminotridecyl)pyridine, T-358
 2-(12-Aminotridecyl)pyrrolidine, A-921
 2-(12-Aminotridecyl)-1-pyrrolidineethanol, A-921
 2-Amino-2,3,5-trideoxy-3-methyl-D-arabinonic acid, A-800
 2-Amino-2,3,5-trideoxy-3-methyl-L-arabinonic acid, A-800
 2-Amino-2,3,5-trideoxy-3-methyl-D-lyxonic acid, A-800
 2-Amino-2,3,5-trideoxy-3-methylpentonic acid, A-800
 2-Amino-2,3,5-trideoxy-3-methyl-L-ribonic acid, A-800
 2-Amino-2,3,5-trideoxy-3-methyl-D-xyloxic acid, A-800
 2-Amino-5-(trifluoromethoxy)benzoic acid, A-789
 2-Amino-3,4,5-trihydroxybenzoic acid, A-922
 11-Amino-4,5,9-trihydroxy-2-(hydroxymethyl)-10*H*-benzo[*b*]fluoren-10-one, S-500
 4'-Amino-3',5',7-trihydroxy-5'-methoxy-2',8-diprenylisoflavone, I-267
 4'-Amino-3',5',7-trihydroxy-5'-methoxy-2',6'-diprenylisoflavone, P-483
 11-Amino-4,5,9-trihydroxy-2-methyl-10*H*-benzo[*b*]fluoren-10-one, S-501
 1-Amino-2,3,4-trihydroxynortropane, A-704
 2-Amino-7-(1,2,3-trihydroxypropyl)-4(1*H*)-pteridine, A-924
 2-Amino-6-(1,2,3-trihydroxypropyl)-4(1*H*)-pteridine, A-923
 2-Amino-4,6,7-trihydroxypteridine, A-925
 α-Amino-2,3,5-trihydroxytoluene, T-528
 2-Amino-3,4,5-trimethoxybenzoic acid, A-922
 6-Amino-4,5,9-trimethoxy-7*H*-dibenzo[*de,h*]quinolin-7-one, D-89
 7-Amino-2,9,10-trimethoxy-1*H*-dibenzo[*de,g*]quinolin-1-one, P-56
 2-Amino-1-(3,4,5-trimethoxyphenyl)ethanol, H-557
 Aminotrimethylcarbinol, A-840
 3-Amino-4,4,14-trimethyl-9,19-cyclopregn-17(20)-en-16-one, C-848
 4-Amino-3-(3,6,7-trimethyl-2,6-octadienyl)benzamide, B-68
 2-Amino-*N,N,N*-trimethyl-2-oxoethanaminium(1+), C-119
 5-Amino-*N,N,N*-trimethylpentanaminium(1+), P-229
 5-Amino-*N,N,N*-trimethyl-1-pentenaminium(1+), P-232
 4-Amino-*N,N,N*-tris(3-aminopropyl)-1-butanaminium, A-926
 3-Amino-*N,N,N*-tris(3-aminopropyl)-1-propanaminium(1+), T-622
 5-(10-Aminoundecyl)hexahydropyrrolo[2,1-*b*]oxazole, A-927
 2-(10-Aminoundecyl)-1-pyrrolidineethanol, P-958
 2-(10-Aminoundecyl)pyrrolidine, P-958
 3-(2-Aminovinyl)-1*H*-indole, D-172
 Aminoxypitropate, T-632
 Aminozaanemonin, A-928
 Amiroside, D-535
 Amisine, J-49
 Ammocalline, A-929
 Ammodendrine, A-930
 Ammorosine, A-931
 Ammothamine, M-121
 Amphasterin C₁, A-932
 Amphasterin C₂, A-933
 Amphasterin C₃, A-932
 Amphasterin C₄, A-933
 Amphasterin D₁, A-932
 Amphasterin D₂, A-933
 Amphasterin D₃, A-933
 Amphasterin E₁, A-932
 Amphibine A, A-935
 Amphibine B, A-936
 Amphibine C, A-937
 Amphibine D, A-938
 Amphibine E, A-939
 Amphibine F, A-940
 Amphibine G, A-941
 Amphibine H, A-942
 Amphibine I, A-934
 Amphikuemin, A-943
 Amphimedine, A-944
 Amphimedoside A, A-945
 Amphimedoside B, A-946
 Amphimedoside C, A-947
 Amphimedoside D, H-3
 Amphimedoside E, A-948
 Amphistin, A-949
 Amphitoxin, A-950
 Amphorogynine A, H-720
 Amphorogynine B, H-720
 Amphorogynine C, H-720
 Amphorogynine D, H-720
 Ampullicin, A-951
 Amsoniaefoline, A-952
 Amsonine, Y-13
 Amsosinine, A-953
 Amurenine, T-524
 Amurensine, A-954
 Amurensinine *N*-oxide A, A-954
 Amurensinine *N*-oxide B, A-954
 Amurensinine, A-954
 Amurine, A-955
 Amurinol, A-956
 Amurinol I, A-955
 Amuroline, A-957
 Amuroline, A-957
 Amycomycin, A-958
 Amygdalin, H-673
 Amygdaloidin A, H-121
 Amygdaloidin B, H-121
 Amygdaloidin C, H-121
 Amygdaloidine, H-673
 Amygdonitrile, H-673
 Amythiamicin A, A-959
 Amythiamicin B, A-960
 Amythiamicin C, A-961
 Amythiamicin D, A-962
 Anabasamine, A-963
 Anabesine, T-144
 Anabasine, JMAF, P-460
 Anabilsine, A-964
 Anachelin H, A-965
 Anacine, A-966
trans-Anacrotine, A-967
 Anacrotine, A-967
 Anacycline, A-968
 Anadoline *N*-oxide, S-163
 Anadoline, S-163
 Anaferine, A-969
 Anagyryne, A-970
 Anahyryne, A-971
 Analeptin, S-659
 Analobine, A-1066
 Anandamide, E-48
 Anantine, A-972
 Anaprel, R-50
 Anapterin, A-924
 Anarkotin, N-44
 Anaspaz, T-632
 Anatabine, T-143
 Anatalline, A-973
 Anatoxin a, A-974
 Anatoxin a(s), A-975
 Anaxagoreine, P-4
 Anchinopeptolide A, A-976
 Anchinopeptolide B, A-976
 Anchinopeptolide C, A-976
 Anchinopeptolide D, A-976
 Anchustringosine, A-977
 Ancisheynine, A-978
 Ancistine, A-981
 Ancistrine, A-990
 Ancistrobenomine A, A-982
 Ancistrobertsonine A, A-980
 Ancistrobertsonine B, A-982
 Ancistrobertsonine C, A-980
 Ancistrobrevine A, A-979
 Ancistrobrevine B, A-980
 Ancistrobrevine C, A-981
 Ancistrobrevine D, A-990
 Ancistrocladeine, A-982
 Ancistrocladidine, A-992
 Ancistrocladine, A-982
 Ancistrocladinine, A-982
 Ancistrocladinium A, A-983
 Ancistrocladinium B, A-984
 Ancistrocladisine, A-985
 Ancistrocladonine, A-982
 Ancistrocline, A-982
 Ancistrocongine, A-986
 Ancistrocongolensine, A-987
 Ancistrocongoline A, K-91
 Ancistrocongoline B, K-91
 Ancistrocongoline C, K-91
 Ancistrocongoline D, A-990
 Ancistroealaensine, A-982
 Ancistroealaine A, A-980
 Ancistroealaine B, A-980
 Ancistrogriffine A, A-979
 Ancistrogriffine B, A-990
 Ancistrogriffine C, A-979
 Ancistrogriffithine A, A-988
 Ancistroguineine A, A-980
 Ancistroguineine B, A-980
 Ancistroheynine A, A-989
 Ancistroheynine B, A-992
 Ancistrolikokine A, K-91
 Ancistrolikokine B, K-91
 Ancistrolikokine C, K-91
 Ancistrolikokine D, A-980
 Ancistrorobertsonine D, A-990
 Ancistrotrozanine A, A-991
 Ancistrotrozanine B, A-980
 Ancistrotrozanine C, A-992
 Ancistrotectoriline A, A-980
 Ancistrotectoriline B, A-990
 Ancistrotectorine, A-992
 Ancorinazole, A-993
 Ancorinolate A, C-402
 Ancorinolate B, I-93
 Ancorinolate C, C-402
 Ancorinoside A, A-994
 Ancorinoside B, A-995
 Ancorinoside C, A-996
 Ancorinoside D, A-997
 Andersobine, H-204
 Andersoline, G-91
 Andersonidine, M-309
 Andersonine, A-998
 Andesine, A-999
 Andirin, T-700
 Andrachamine, B-187
 Andrachcine, H-433
 Andrachcinidine, A-1000
 Andrachcine, H-433
 Andrachmine, A-1001
 Andrangine, A-1002
 Andrangine, A-1003
 Andrimide, A-1004
 Androbine, P-221
 Androbiphenylene, C-567
 Androcimine, P-221
 Androcine, P-221
 Androcymbine, A-1005
 Anemonin, Z-32
 Aneurin diphosphate, T-366
 Aneurine, T-366
 Angchibangkine, A-1006
 Angelicastigmine, A-1007
 Angelin, T-700
 3-(Angeloylamino)-20-(dimethylamino)-5-pregnen-4-one, D-282
 Angeloyl-*trans*-anacrotine *N*-oxide, A-967
 7-Angeloyldihydroheliotridine, H-233
 9-Angeloyldihydroheliotridine, H-233
 15-Angeloylgermine, G-70
 9-Angeloylhastaneceine, H-233
 7-Angeloylheliotridine, T-188
 6-Angeloyl-3-(*trans*-4-hydroxyseneciyl)tropane, M-391
 7-Angeloyl-1-methylenepyrrolizidine *N*-oxide, H-601

- 7-Angeloyl-1-methylenepyrrolizidine, H-601
6-Angeloyloxyditropan-3-yl itaconate, A-1008
13-(Angeloyloxy)-4-hydroxylupanine, D-606
4-Angeloyloxy-3-hydroxy-13-methoxylupanine, T-557
4-(Angeloyloxy)lupanine, H-551
13-Angeloyloxyylupanine, H-554
13-Angeloyloxy-17-oxolupanine, O-188
6-Angeloyloxyplatynecine, D-595
6-Angeloyloxytropan-3-ol, M-391
9-Angeloylplatynecine *N*-oxide, H-233
7-Angeloylplatynecine, H-233
9-Angeloylplatynecine, H-233
9-Angeloylretronecine *N*-oxide, T-188
7-Angeloylretronecine, T-188
9-Angeloylretronecine, T-188
7-Angeloylturneforcidine, H-233
9-Angeloylturneforcidine, H-233
Angeloylturneforcidine, H-233
15-Angeloyl-3-veratroylgermine, G-70
Angeloylzygadenilic acid δ -lactone, Z-35
Angeloylzygadenine β -*N*-oxide, Z-36
Angeloylzygadenine, Z-36
7-Angelylheliotridine trachelanthate, E-31
7-Angelylheliotridine viridiflorate, E-31
7-Angelylheliotrine, L-47
7-Angelyl-9-irididiflorylheliotridine, E-31
Angoline, D-407
Angolinine, N-239
Anguibactin, A-1009
Anquivine, S-456
Angularine, R-133
Angulatamine, P-247
Angulatueoid A, P-218
Angulatueoid B, P-218
Angulatueoid C, T-259
Angulatueoid D, T-259
Angulatueoid E, T-259
Angulatueoid F, T-259
Angulatusine A, P-218
Angusteine, A-1010
Angusticaline, A-1011
Angustidine, A-1012
Angustifoline *N*-carboxyethyl ester, A-1013
Angustifoline *N*-carboxymethyl ester, A-1013
Angustifoline, A-1013
Angustilobine A, A-1014
Angustilobine B acid, A-1015
Angustilobine B *N*^b-oxide, A-1015
Angustilobine B, A-1015
Angustilocine, L-264
Angustilodine, A-1016
Angustimalal, A-1017
Angustimaline, A-1018
Angustimicine, A-228
Angustine, A-1019
Angustmycin B, A-137
Angustoline, A-1020
Angustureine, P-238
Anhalamine, T-236
Anhalidine, T-236
Anhaline, H-357
Anhalinine, T-236
Anhalonidine, T-237
Anhalonine, A-1021
Anhalotine, T-236
Anhweiaconitine, A-1022
Anhweidelphinine, A-1023
Anhydroadrubine, A-141
Anhydroalstonatine, N-132
Anhydroaposerratinine, F-28
Anhydroberberilic acid, A-1024
Anhydrobuxpiene, C-882
Anhydrocannabisativine, C-83
14,15-Anhydrocapuronidine, D-377
Anhydrodeacetylpaniculine, P-71
Anhydrodehydrotylophorinidine, A-1025
Anhydrodehydrotylophorimine, A-1025
4,9-Anhydro-11-deoxytetrodotoxin, A-1035
Anhydrosdihydroamataine, V-188
14,15-Anhydrosdihydrocapuronidine, D-377
Anhydrosdihydrocypodine, L-341
Anhydroecgonine methyl ester *N*-oxide, M-395
Anhydroecgonine methyl ester, M-395
Anhydroecgonine, M-395
4,9-Anhydro-6-epitetrodotoxin, A-1035
Anhydroevoxine, F-223
Anhydrohapaloxindole A, A-1026
Anhydrohapaloxindole B, A-1026
Anhydrohapaloxindole M, A-1026
Anhydroharringtonine, A-1027
Anhydrohazuntiphyllidine, H-80
Anhydroholantogenin, A-1028
Anhydroignavinol, H-184
Anhydrolycaconitine, A-106
Anhydrolycoeceruleine, L-321
Anhydrolycodoline, A-1029
Anhydrolycorinium, A-1030
Anhydrolycorin-7-one, H-290
Anhydromazethramycin, M-137
Anhydromethylpseudolycorine, A-1508
Anhydronupharamine, N-340
4,6'-Anhydrooxysporidinone, A-1031
Anhydroperforine, A-1032
Anhydroperipentamine, A-1033
Anhydropilosine, P-428
Anhydroseudolycorine, A-1508
Anhydroshumannifine, S-145
Anhydroscopin A, A-1044
16,17-Anhydrotacamine, T-12
Anhydrotetracycline, A-1034
4,9-Anhydrotetrodotoxin, A-1035
15',20'-Anhydrovinblastine, V-110
3',4'-Anhydrovinblastine, V-110
Anhydrovobasinediol, T-10
Anhydroscopin B, S-53
Anibacanine, M-81
Anibamine, A-1036
Anibine, A-1037
Anicanine, A-1038
Aniflorine, A-1039
2-Anilino-1,4-naphthoquinone, A-1040
2-Anilinoquinoline, Q-28
Anisamide, M-240
Anisamitocin PHO 3, M-136
Anisessine, A-1041
p-Anisic acid, M-240
Anisocycline, A-1042
Anisodamine, H-521
Anisodine, A-1043
Anisoezochasmaconitine, M-323
Anisomycin C, A-1044
Anisomycin D, A-1044
Anisomycin, A-1044
Anisotine, A-1045
14-Anisoylaconine, M-331
14-Anisoylneoline, M-323
14-Anisoyltalatzamine, M-352
N-(2-*p*-Anisylethyl)-*N*-methylcinnamide, H-144
Ankoline, A-1046
Ankorine, A-1047
Annapawine, A-1048
Annobrine, L-145
Annocherine A, A-1049
Annocherine B, A-1049
Annocherine C, A-1049
Annofoline, F-29
Annolatine, A-1050
Annomontine, A-1051
Annomurine, A-1051
Annonelliptine, A-1052
Annonidine A, A-1053
Annonidine B, A-1054
Annonidine C, A-1055
Annonidine D, A-1056
Annonidine E, A-1055
Annonidine F, A-1057
Annopoline, A-1058
Annopodine, A-1059
Annoretine, A-1060
Annosqualine, A-1061
Annotine, A-1062
Annotimine, A-1063
Annotoxine, A-1062
Annuloline, A-1064
Anodendrine, A-1065
Anolobine, A-1066
Anomaline, A-1067
Anomoian A, A-1068
Anomoline, A-1052
Anomuricine, A-1069
Anomurine, A-1069
Anonaine, R-120
Anonamine, S-245
Anopterimine *N*-oxide, A-1070
Anopterimine, A-1070
Anopteryne, A-1071
Anopteryl alcohol, A-1071
Anopteryl 11,12-ditiglate, A-1071
Anopteryl 11-4'-hydroxybenzoate 12-tiglate, A-1071
Anopteryl 12-tiglate, A-1071
Anorldianine *N*²⁷-oxide, A-1072
Anorldianine, A-1072
Anorosin, V-142
Anosmine, O-45
Anrakorinine, A-1073
Ansamitocin PHO 0, M-136
Ansamitocin PHO 1, M-136
Ansamitocin PHO 2, M-136
Ansamitocin PHO 4, M-136
Ansamitocin P1, M-136
Ansamitocin P2, M-136
Ansamitocin P3', M-136
Ansamitocin P3, M-136
Ansamitocin P4, M-136
Ansamitocin P₆, M-136
Ansamitocin PND 1, M-136
Ansamitocin PND 2, M-136
Ansamitocin PND 3, M-136
Ansamitocin PND 4, M-136
Ansamitocinoside P1, M-136
Ansamitocinoside P2, M-136
Antazirine, A-1074
Antemovis, H-755
Anthcolorin A, A-1075
Anthcolorin B, A-1075
Anthcolorin C, A-1075
Anthcolorin D, A-1075
Anthcolorin E, A-1076
Anthcolorin F, A-1076
Anthcolorin G, A-1075
Anthcolorin H, A-1075
Anthelvincin A, A-1077
Anthemovister, H-755
Anthirine lactone, A-1329
Anthirine, A-1328
Anthocero Diazonin, A-1078
Anthorine, A-1534
Anthoroidine, H-181
Anthosamine A, A-1079
Anthosamine B, A-1079
Anthracylcinone blue A, A-1080
Anthracylcinone blue B, A-1080
Anthramycin, A-1081
Anthranilamide, A-706
Anthranilic acid, A-706
4-Anthranoyllappaconidine, A-91
Anthranoyllycoctonine, M-309
 β -Anthrapyridinequinone, B-71
Anthraserpine, R-50
Anthriscifolcine A, A-92
Anthriscifolcine B, A-92
Anthriscifolcine C, A-88
Anthriscifolcine D, A-88
Anthriscifolcine E, A-88
Antibiotic A 10947, A-1097
Antibiotic A 33853, A-1098
Antibiotic A 6413, A-1202
Antibiotic A 65636, A-1099
Antibiotic A 6984, A-1095
Antibiotic A 7413, A-1096
Antibiotic A 75943, A-1100
Antibiotic A 500359A, A-1102
Antibiotic A 503083A, A-1102
Antibiotic A 58365A, A-1101
Antibiotic A 72363A1, A-746
Antibiotic A 72363A2, A-746

- Antibiotic A 89271A, A-685
 Antibiotic A 25822A, D-742
 Antibiotic A 83094A, I-64
 Antibiotic A 2315A, M-35
 Antibiotic A 90931A, X-6
 Antibiotic A 10255B, A-1103
 Antibiotic A 25822B, A-1602
 Antibiotic A 58365B, A-1104
 Antibiotic A 72363B, A-746
 Antibiotic A 89271B, A-684
 Antibiotic A 500359B, C-103
 Antibiotic A 503083B, C-103
 Antibiotic A 90931B, X-6
 Antibiotic A 23187, C-26
 Antibiotic A 10255C, A-1105
 Antibiotic A 500359C, A-1102
 Antibiotic A 58365C, A-1106
 Antibiotic A 72363C, A-746
 Antibiotic A 89271C, A-684
 Antibiotic A 19C, C-900
 Antibiotic A 17002C, M-35
 Antibiotic A 25822D, A-1602
 Antibiotic A 500359D, A-1102
 Antibiotic A 7413D, A-1107
 Antibiotic A 89271D, A-686
 Antibiotic A 10255E, A-1108
 Antibiotic A 89271E, A-684
 Antibiotic A 10255F, A-1109
 Antibiotic A 89271F, A-687
 Antibiotic A 17002F, M-35
 Antibiotic A 10073, G-179
 Antibiotic A 500359G, C-103
 Antibiotic A 32256, H-496
 Antibiotic A 10255H, A-1110
 Antibiotic A 25822H, G-64
 Antibiotic A 15104II, M-35
 Antibiotic A 300-II, T-496
 Antibiotic A 300I, T-496
 Antibiotic A 25822L, A-1602
 Antibiotic A 25822M, A-1602
 Antibiotic A 77543, N-265
 Antibiotic A 25822N, A-1602
 Antibiotic A 102338, P-961
 Antibiotic A 23812, P-891
 Antibiotic A 23813, P-891
 Antibiotic A 43791, T-392
 Antibiotic A 70615, T-447
 Antibiotic A 15104V, M-35
 Antibiotic A 15104Y, D-324
 Antibiotic 106-7, A-832
 Antibiotic 1270, A-1084
 Antibiotic 17452, A-832
 Antibiotic 176, A-1195
 Antibiotic 2158, A-1084
 Antibiotic 35763, A-1151
 Antibiotic 41043, A-1152
 Antibiotic 41494, A-1153
 Antibiotic 5102-2, A-1085
 Antibiotic 5915, A-832
 Antibiotic 6604-4, A-78
 Antibiotic 68-1147, A-1082
 Antibiotic 69, A-1083
 Antibiotic 8217, A-832
 Antibiotic 9408, A-1086
 Antibiotic 0231A, A-1088
 Antibiotic 116A, A-664
 Antibiotic 2096A, A-1093
 Antibiotic 2197A, A-1094
 Antibiotic 593A, A-1089
 Antibiotic 859A, A-1090
 Antibiotic 875A, A-1091
 Antibiotic 927A, A-1092
 Antibiotic AB 97, A-1111
 Antibiotic AB 3217A, A-1112
 Antibiotic AB 4063A, A-1114
 Antibiotic AB 3217B, A-1112
 Antibiotic AB 4015B, A-1113
 Antibiotic AB 4063B, A-1114
 Antibiotic AB 3217C, A-1112
 Antibiotic AB 85, F-115
 Antibiotic AB 116, K-8
 Antibiotic AC 7230, C-26
 Antibiotic 167A, C-287
 Antibiotic 2562A, C-395
 Antibiotic 460A, C-936
 Antibiotic 49A, C-830
 Antibiotic A₃-118D, L-374
 Antibiotic AF 140, A-1115
 Antibiotic AF 110, F-197
 Antibiotic AG 55, T-373
 Antibiotic AH 135Y, A-129
 Antibiotic 2088A, H-514
 Antibiotic AIL-B, N-131
 Antibiotic AIS 2M3, R-33
 Antibiotic AJI 9561, A-1116
 Antibiotic AJP 117510, U-42
 Antibiotic AK-Ps, O-124
 Antibiotic AI-R 2081, P-974
 Antibiotic AI-R2397, M-136
 Antibiotic AM 5344A₁, C-287
 Antibiotic AM 5344A₂, C-287
 Antibiotic AM 6201, R-34
 Antibiotic AM 2282, S-498
 Antibiotic A, M-653
 Antibiotic 2814A, N-180
 Antibiotic 1494A, O-134
 Antibiotic A528, Q-60
 Antibiotic ASK 753, A-1117
 Antibiotic AS-N-7A, A-1118
 Antibiotic AT 125, A-81
 Antibiotic AT 2433, A-1119
 Antibiotic 145-A, T-496
 Antibiotic ATF 606, M-159
 Antibiotic AX 1, A-246
 Antibiotic AX 2, I-246
 Antibiotic Azx, M-396
 Antibiotic B 4317, A-1122
 Antibiotic B 90063, A-1123
 Antibiotic B 4607A, H-514
 Antibiotic B 41A₃, M-607
 Antibiotic B 41A₄, M-608
 Antibiotic B 4607B, H-514
 Antibiotic B 41B₂, M-607
 Antibiotic B 41B₃, M-608
 Antibiotic B 79B, S-384
 Antibiotic B 4607C, H-514
 Antibiotic B 41C₁, M-607
 Antibiotic B 41D, M-609
 Antibiotic B 79D, F-658
 Antibiotic B 1625FA_{2p-1}, A-1124
 Antibiotic B 41F, M-609
 Antibiotic B 41G, M-609
 Antibiotic B 371, I-216
 Antibiotic B 14437, S-53
 Antibiotic B 870, T-390
 Antibiotic BA 181314A, A-1125
 Antibiotic BA 90912, S-53
 Antibiotic 0231B, A-1088
 Antibiotic 1401B, A-1120
 Antibiotic 2096B, A-1093
 Antibiotic 4181B, A-1121
 Antibiotic 460B, A-1309
 Antibiotic B057-250A, M-588
 Antibiotic BBM 2040A, C-373
 Antibiotic BBM 928A, L-312
 Antibiotic BBM 2040B, C-374
 Antibiotic BBM 928B, L-312
 Antibiotic BBM 928C, L-312
 Antibiotic B057-250B, M-588
 Antibiotic BC 41C₂, M-608
 Antibiotic 167B, C-287
 Antibiotic 2562B, C-395
 Antibiotic B057-250C, M-588
 Antibiotic B057-250D, M-588
 Antibiotic BE 10988, A-1126
 Antibiotic BE 14106, A-1127
 Antibiotic BE 22179, A-1128
 Antibiotic BE 48021, A-1129
 Antibiotic BE 52211, A-1130
 Antibiotic BE 54017, A-1131
 Antibiotic BE 54476, A-1132
 Antibiotic BE 55051, A-1133
 Antibiotic BE 67251, A-1134
 Antibiotic BE 19412A, A-1135
 Antibiotic BE 54238A, A-1136
 Antibiotic BE 45722A, D-126
 Antibiotic BE 19412B, A-1135
 Antibiotic BE 39907B, A-853
 Antibiotic BE 52211B, A-1137
 Antibiotic BE 54238B, A-1136
 Antibiotic BE 45722B, D-126
 Antibiotic BE 13793C, A-1393
 Antibiotic BE 52211C, A-1137
 Antibiotic BE 45722C, D-126
 Antibiotic BE 12233, H-747
 Antibiotic BE 18591, T-22
 Antibiotic B057-250E, M-588
 Antibiotic 2088B, H-514
 Antibiotic 10204-BII, F-48
 Antibiotic BK 97B1, B-64
 Antibiotic BK 97B2, B-65
 Antibiotic BM 1, A-1606
 Antibiotic BM 123 γ ₂, C-460
 Antibiotic BM 123 γ ₁, C-459
 Antibiotic BM 123 β ₁, G-33
 Antibiotic 434B, M-413
 Antibiotic BMG 162 aF₂, S-394
 Antibiotic BMS 199687, N-265
 Antibiotic BMY 41219, A-1393
 Antibiotic BMY 42428, A-1138
 Antibiotic BMY 28565, F-33
 Antibiotic BMY 40660, H-417
 Antibiotic BMY 28886, L-7
 Antibiotic BMY 28566, M-158
 Antibiotic BMY 28121, P-573
 Antibiotic BMY 42448, P-969
 Antibiotic BMY 28662, Q-11
 Antibiotic BMY 41950, S-498
 Antibiotic BN 229A, H-563
 Antibiotic BN 227F, H-563
 Antibiotic BN 227, H-563
 Antibiotic 4738 B, P-833
 Antibiotic BR 040, A-1139
 Antibiotic BR 042, A-1139
 Antibiotic BS 75A, C-574
 Antibiotic BT 38, A-801
 Antibiotic BU 3557A₃, A-1315
 Antibiotic BU 4601A, A-789
 Antibiotic BU 2349A, G-131
 Antibiotic BU 3557A₁, K-17
 Antibiotic BU 3557A₂, K-18
 Antibiotic BU 2313A, N-264
 Antibiotic BU 4601B, A-789
 Antibiotic BU 2349B, G-132
 Antibiotic BU 3557B₁, K-17
 Antibiotic BU 3557B₂, K-17
 Antibiotic BU 3557B₃, K-18
 Antibiotic BU 2313B, N-264
 Antibiotic BU 620, C-711
 Antibiotic BU 4601C, A-789
 Antibiotic BU 2349C, G-133
 Antibiotic BU 3557C₁, K-17
 Antibiotic BU 3557C₂, K-19
 Antibiotic BU 3557C₃, K-20
 Antibiotic BU 3557C₄, K-20
 Antibiotic BU 3557C₅, K-17
 Antibiotic BU 3557D₁, A-1315
 Antibiotic BU 3557D₂, K-21
 Antibiotic BU 3557D₃, K-20
 Antibiotic BU 3557D₄, K-21
 Antibiotic BU 4664L, D-297
 Antibiotic BU 4514N, A-1140
 Antibiotic BU 4146T, L-7
 Antibiotic BU 2916T, P-573
 Antibiotic BU 3845T, Q-11
 Antibiotic BU 4704, X-7
 Antibiotic C 73, A-129
 Antibiotic C 14482A₁, D-876
 Antibiotic C 304A, H-408
 Antibiotic C 14482B₁, D-877
 Antibiotic C 5507, C-954
 Antibiotic C 15003 deClQND0, M-136
 Antibiotic C 15003 deClQO, M-136
 Antibiotic C 15003 P 4 β HY, M-136
 Antibiotic C 15003 P 4 γ HY, M-136
 Antibiotic C 9154, P-337
 Antibiotic C 15003 PHM1, M-136
 Antibiotic C 15003 PHM2, M-136
 Antibiotic C 15003 PHM3, M-136

- Antibiotic C 15003 PHM4, M-136
 Antibiotic C 15003 PHO 0, M-136
 Antibiotic C 15003 PHO 1, M-136
 Antibiotic C 15003 PHO 4, M-136
 Antibiotic C 15003P0, M-136
 Antibiotic C 15003P1, M-136
 Antibiotic C 15003P2, M-136
 Antibiotic C 15003P3', M-136
 Antibiotic C 15003P3, M-136
 Antibiotic C 15003P4, M-136
 Antibiotic C 15003 PND 4 βHY, M-136
 Antibiotic C 15003 PND 1, M-136
 Antibiotic C 15003 PND 2, M-136
 Antibiotic C 15003 PND 3, M-136
 Antibiotic C 15003 PND 4, M-136
 Antibiotic C 15003 QNDO, M-136
 Antibiotic C 73X, A-129
 Antibiotic 354, C-521
 Antibiotic 900447, C-427
 Antibiotic CA 39A, S-500
 Antibiotic CA 39B, S-500
 Antibiotic CC 1065, R-4
 Antibiotic 2096C, D-498
 Antibiotic CGP 39360, S-498
 Antibiotic CGP 58546, S-498
 Antibiotic CH 777A, N-180
 Antibiotic 2088C, H-514
 Antibiotic CI 4, C-841
 Antibiotic CJ 12662, A-1141
 Antibiotic CJ 12663, A-1141
 Antibiotic CJ 15696, A-1145
 Antibiotic CJ 16169, A-1145
 Antibiotic CJ 16170, A-1142
 Antibiotic CJ 16171, A-1143
 Antibiotic CJ 16173, A-1144
 Antibiotic CJ 16174, A-1145
 Antibiotic CJ 16196, A-1146
 Antibiotic CJ 16197, A-1146
 Antibiotic CJ 16264, A-1147
 Antibiotic CJ 16367, A-1148
 Antibiotic CJ 17572, A-1149
 Antibiotic CJ 17665, A-1470
 Antibiotic CJ 21058, A-1150
 Antibiotic CJ 13136, D-760
 Antibiotic CJ 13217, D-760
 Antibiotic CJ 13536, D-760
 Antibiotic CJ 13564, D-760
 Antibiotic CJ 13565, D-762
 Antibiotic CJ 13566, D-762
 Antibiotic CJ 13567, D-762
 Antibiotic CJ 13568, D-762
 Antibiotic CJ 14877, D-666
 Antibiotic CJ 14897, D-666
 Antibiotic CJ 15335, D-666
 Antibiotic 2908C, K-47
 Antibiotic 547C, O-131
 Antibiotic CP 35763, A-1151
 Antibiotic CP 41043, A-1152
 Antibiotic CP 41494, A-1153
 Antibiotic CP 47444, N-276
 Antibiotic CP 51467, N-48
 Antibiotic CP 52726, N-49
 Antibiotic CP 52748, N-50
 Antibiotic CP 57820, N-276
 Antibiotic CP 61405, R-136
 Antibiotic CP 46192, T-391
 Antibiotic CS 79A, C-169
 Antibiotic CS 79B, C-169
 Antibiotic CSL 12A, T-700
 Antibiotic CSL 12B, T-700
 Antibiotic CSL 12C, T-700
 Antibiotic CSL 12D, T-700
 Antibiotic CSL 12E, T-700
 Antibiotic CSL 12F, T-700
 Antibiotic CSL 12G, T-700
 Antibiotic CSL 12H, T-700
 Antibiotic CSL 12I, T-700
 Antibiotic CSL 12J, T-700
 Antibiotic CSL 12K, T-700
 Antibiotic CSL 12L, T-700
 Antibiotic CSL 12M, T-700
 Antibiotic CV 1, A-1154
 Antibiotic CW 69A, C-129
 Antibiotic CW 69B, C-129
 Antibiotic CW 69, C-129
 Antibiotic CW 69C, C-129
 Antibiotic CW 69D, C-129
 Antibiotic CW 69E, C-129
 Antibiotic CW 69F, C-129
 Antibiotic 0089D, A-1155
 Antibiotic DC 102, A-1157
 Antibiotic DC 105, A-1158
 Antibiotic DC 81, A-1156
 Antibiotic DC 8118A, A-1159
 Antibiotic DC 88A, D-952
 Antibiotic DC 89A₁, D-953
 Antibiotic DC 89A₂, D-954
 Antibiotic DC 59A, S-231
 Antibiotic DC 8118B, A-1159
 Antibiotic DC 89B₁, D-953
 Antibiotic DC 86C, H-496
 Antibiotic DC 113, D-956
 Antibiotic DC 52d, Q-22
 Antibiotic DC 59H, S-231
 Antibiotic DC 107, L-82
 Antibiotic DC 86M, H-496
 Antibiotic DC 62, P-660
 Antibiotic DC 52, Q-21
 Antibiotic DC 86R, H-496
 Antibiotic DC 86Y, H-496
 Antibiotic 2096D, D-522
 Antibiotic DO 248A, R-54
 Antibiotic DO 248B, R-55
 Antibiotic DOB 41, A-1160
 Antibiotic E 733A, A-832
 Antibiotic E 212, S-53
 Antibiotic 1252, E-61
 Antibiotic E129A, O-131
 Antibiotic EC 40, D-761
 Antibiotic ECO 02301, A-1161
 Antibiotic EGS 1300-1, A-1162
 Antibiotic EM 4940A, S-295
 Antibiotic EM 5596, N-101
 Antibiotic EM 5519, S-2
 Antibiotic F 1778, A-1163
 Antibiotic F 1839A, A-1164
 Antibiotic F 1839B, A-1164
 Antibiotic F 10C, A-1165
 Antibiotic F 1839C, A-1164
 Antibiotic F 1839D, A-1164
 Antibiotic F 1839E, A-1164
 Antibiotic F 1839F, A-1164
 Antibiotic F 2787, H-729
 Antibiotic F 1839J, A-1164
 Antibiotic F 4708, S-490
 Antibiotic 4-215, F-131
 Antibiotic FA 4283, A-1166
 Antibiotic 31F508α1, B-140
 Antibiotic 31F508α2, B-140
 Antibiotic 31F508β1, B-140
 Antibiotic 31F508β2, B-140
 Antibiotic 4849F, B-73
 Antibiotic F6, B-403
 Antibiotic FCRC A48, F-152
 Antibiotic FD 838, A-1167
 Antibiotic FD 839, A-1168
 Antibiotic FD 991, Q-58
 Antibiotic 610FF, E-235
 Antibiotic FKI 0550, A-758
 Antibiotic FL 120B', A-1169
 Antibiotic FL 120B, A-1169
 Antibiotic FL 657C, T-496
 Antibiotic FO 7314, A-1407
 Antibiotic FO 2047, P-333
 Antibiotic FR 225656, A-1172
 Antibiotic FR 225659, A-1173
 Antibiotic FR 252921, A-1174
 Antibiotic FR 252922, A-1175
 Antibiotic FR 256523, A-1175
 Antibiotic FR 3383, A-1170
 Antibiotic FR 66979, A-1171
 Antibiotic FR 900148, A-1176
 Antibiotic FR 900220, A-680
 Antibiotic FR 900482, A-1177
 Antibiotic FR 901483, A-1178
 Antibiotic FR 900216B, R-83
 Antibiotic FR 900216C, R-83
 Antibiotic FR 900483, D-420
 Antibiotic FR 900411, E-242
 Antibiotic FR 900216F, R-83
 Antibiotic FR 177391, H-76
 Antibiotic FR 900494, K-41
 Antibiotic G 1499-2, A-1179
 Antibiotic G 1549, H-563
 Antibiotic 3780, G-210
 Antibiotic GE 2270, A-1180
 Antibiotic GE 37468A, A-1181
 Antibiotic GE 2270B₁, A-1180
 Antibiotic GE 37468B, A-1181
 Antibiotic GE 37468C, A-1181
 Antibiotic GE 2270D₁, A-1180
 Antibiotic GE 2270E, A-1180
 Antibiotic GE 2270T, A-1180
 Antibiotic GERI-BP001 A, T-500
 Antibiotic GERI-BP001 B, T-500
 Antibiotic GERI-BP001 M, T-500
 Antibiotic GKK 1032A₁, A-1182
 Antibiotic GKK 1032A₂, A-1182
 Antibiotic GKK 1032B, A-1183
 Antibiotic GM 95, T-66
 Antibiotic GT 32A, A-1127
 Antibiotic GT 32B, A-1127
 Antibiotic HA 23, A-1184
 Antibiotic HA 9, O-124
 Antibiotic HQ 24A, T-370
 Antibiotic HS 3, P-974
 Antibiotic I 1431, A-832
 Antibiotic I 639A, D-769
 Antibiotic I 693B, D-769
 Antibiotic IA 887, N-180
 Antibiotic IC 202A, A-1185
 Antibiotic IC 202B, A-1186
 Antibiotic IC 202C, A-1187
 Antibiotic ICM 0201, A-1188
 Antibiotic IM 8443T, C-815
 Antibiotic IT 143C, P-421
 Antibiotic IT 143D, P-421
 Antibiotic IV-3, M-70
 Antibiotic J 104303, A-1393
 Antibiotic JBIR-02, P-421
 Antibiotic JM 971A, T-699
 Antibiotic JM 971B, T-699
 Antibiotic JM 971C, T-699
 Antibiotic K 300, A-832
 Antibiotic K 252a, A-1189
 Antibiotic K 82A, L-78
 Antibiotic K 16A, M-76
 Antibiotic K 252b, A-1189
 Antibiotic K 73, C-901
 Antibiotic K 252c, A-1190
 Antibiotic K 252d, A-1190
 Antibiotic K 93-0711-I₁, M-34
 Antibiotic K 93-0711-I₂, M-34
 Antibiotic K 117, N-180
 Antibiotic K01-0509B, G-185
 Antibiotic KC 7367A, M-85
 Antibiotic KC 7367B, M-85
 Antibiotic KF 8940, H-133
 Antibiotic KG 431A, A-1191
 Antibiotic KG 431B, I-100
 Antibiotic KR 025, M-807
 Antibiotic KSB 1939H3, M-607
 Antibiotic KSB 193944, M-607
 Antibiotic KSM 2690B, O-155
 Antibiotic KSM 2690C, O-155
 Antibiotic K₃-Ye, P-327
 Antibiotic L 11-1, A-1192
 Antibiotic L 13365, A-1193
 Antibiotic L 654040, A-1194
 Antibiotic L 681176, A-1195
 Antibiotic L 696475, A-1196
 Antibiotic L 697318, A-1196
 Antibiotic L 755807, A-1197
 Antibiotic L 970843, A-1198
 Antibiotic L 970844, A-1198
 Antibiotic L 52-71A, A-1304
 Antibiotic L 657398, B-89
 Antibiotic L 52-71B, A-1304
 Antibiotic L 892-4, C-837

- Antibiotic L 954967, N-269
 Antibiotic L 156586, O-131
 Antibiotic L 156587, O-131
 Antibiotic L 156588, O-131
 Antibiotic L 746628, P-584
 Antibiotic L 696474, Z-37
 Antibiotic LK6A, L-354
 Antibiotic LL C10037 α , A-804
 Antibiotic LL-AP-191 γ , L-90
 Antibiotic LL-BH872 α , A-1199
 Antibiotic LL-BM 123 γ ₁, C-459
 Antibiotic LL-BM 123 γ ₂, C-460
 Antibiotic LL-BM 123 β ₁, G-117
 Antibiotic LL-BO 2964, A-1200
 Antibiotic LL-E19085 α , C-473
 Antibiotic LL-14E605 α , G-126
 Antibiotic LL-E605 β , G-126
 Antibiotic LL-49F233 α , A-1201
 Antibiotic LL-F42248 α , P-974
 Antibiotic LU 1, H-616
 Antibiotic M 259, A-1202
 Antibiotic M 146791, C-837
 Antibiotic M 4582, H-408
 Antibiotic M 193, S-498
 Antibiotic MBH 001, A-1203
 Antibiotic MBP 039-06, A-1204
 Antibiotic MBP 049-13, A-1205
 Antibiotic MDL 62879, A-1180
 Antibiotic Mer-A 2026A, P-421
 Antibiotic Mer-A 2026B, P-421
 Antibiotic Mer-WF 1726, C-945
 Antibiotic MF 5810, O-134
 Antibiotic 610MF, A-1206
 Antibiotic MG 883-12F γ ₂, S-569
 Antibiotic MH 072, A-1199
 Antibiotic MH 435A, E-144
 Antibiotic MH 435B, E-144
 Antibiotic MI 481-42F4A, A-959
 Antibiotic MI 481-42F4B, A-960
 Antibiotic MI 481-42F4C, A-961
 Antibiotic MI 481-42F4D, A-962
 Antibiotic MI 19827, M-607
 Antibiotic MI 198Z1, M-607
 Antibiotic MJ 347-81F4-A, N-266
 Antibiotic MJ 347-81F4-B, N-266
 Antibiotic MK 61, N-278
 Antibiotic ML 1532II, C-900
 Antibiotic MLR 52, S-498
 Antibiotic MM 14201, A-804
 Antibiotic MM 21801, A-759
 Antibiotic MMC 4, M-654
 Antibiotic MP 105, A-772
 Antibiotic MPC 1001F, A-1207
 Antibiotic MR 93A, M-159
 Antibiotic MR 93B, O-156
 Antibiotic MRL 3120, F-97
 Antibiotic MS 681, A-1208
 Antibiotic MSD 819, C-416
 Antibiotic MSD 92, D-498
 Antibiotic MT 35214, A-804
 Antibiotic MT 36531, A-804
 Antibiotic MT 1882-II, P-421
 Antibiotic MT 1882-I, P-421
 Antibiotic MY 336a, A-1209
 Antibiotic MY 12-62a, H-133
 Antibiotic MY 12-62c, H-134
 Antibiotic MYC 8005, A-1210
 Antibiotic N 461, C-102
 Antibiotic N 13, R-160
 Antibiotic NA 22598A₁, A-1211
 Antibiotic NA 30851A, A-1212
 Antibiotic NA 337, T-213
 Antibiotic NFAT 68, A-1213
 Antibiotic NG 311, A-1214
 Antibiotic NG 391, A-1215
 Antibiotic NG 392, A-1215
 Antibiotic NG 393, A-1215
 Antibiotic NG 129, S-490
 Antibiotic NG 242, S-490
 Antibiotic NG 243, S-490
 Antibiotic NG 245, S-490
 Antibiotic NI 15501A, A-33
 Antibiotic NJ 21, A-832
 Antibiotic NK 30424A, A-1216
 Antibiotic NK 170204A, P-421
 Antibiotic NK 372135A, X-7
 Antibiotic NK 170204B, A-1217
 Antibiotic NK 30424B, A-1216
 Antibiotic NK 372135B, X-7
 Antibiotic NK 372135C, X-7
 Antibiotic NK 26588, I-179
 Antibiotic NP 522, A-1218
 Antibiotic NP 101A, A-31
 Antibiotic NP 25302, B-242
 Antibiotic NP 25301, D-224
 Antibiotic NPI 0047, S-28
 Antibiotic NPI 0052, S-28
 Antibiotic NRCS 15, A-1219
 Antibiotic 14752-2, O-131
 Antibiotic OM 704A, D-295
 Antibiotic OM 704B, D-295
 Antibiotic OM 5714, I-161
 Antibiotic OPC 15161, A-1220
 Antibiotic P 42-1, A-244
 Antibiotic P 725C, A-16791
 Antibiotic PA 106, A-1044
 Antibiotic PA 94, A-832
 Antibiotic PA 95, A-78
 Antibiotic PA 155A, I-102
 Antibiotic PA 114A, O-131
 Antibiotic PA 371e, I-300
 Antibiotic PA 371 γ , I-300
 Antibiotic 64p-A, T-603
 Antibiotic PBA, A-1081
 Antibiotic PD 116152, A-1221
 Antibiotic PDE I, A-1222
 Antibiotic PDE II, A-1223
 Antibiotic PF 1018, A-1224
 Antibiotic PF 1052, A-1225
 Antibiotic PF 1140, A-1226
 Antibiotic PF 1163A, A-1227
 Antibiotic PF 1198A, A-240
 Antibiotic PF 1126A, C-208
 Antibiotic PF 1138A, C-209
 Antibiotic PF 1163B, A-1227
 Antibiotic PF 1198B, A-240
 Antibiotic PF 1126B, C-208
 Antibiotic PF 1138B, C-209
 Antibiotic PF 1101B, S-282
 Antibiotic PF 1191, K-5
 Antibiotic 15003 PHO 2, M-136
 Antibiotic 15003 PHO 3, M-136
 Antibiotic PI 091, A-1228
 Antibiotic PI 290, B-29
 Antibiotic PM 94128, A-1229
 Antibiotic PM 93135, T-384
 Antibiotic PO 1227A, Q-37
 Antibiotic PO 1227B, Q-37
 Antibiotic PSA, A-759
 Antibiotic PSB, A-759
 Antibiotic PSC₁, A-759
 Antibiotic PSC₂, A-759
 Antibiotic Q 11270A, A-1164
 Antibiotic Q 20547A, A-1311
 Antibiotic Q 11270B, A-1164
 Antibiotic Q 20547C, A-1310
 Antibiotic 1491, Q-58
 Antibiotic 59266, Q-58
 Antibiotic 6270, Q-61
 Antibiotic R 39FF, M-430
 Antibiotic R 39MF, A-1230
 Antibiotic R 39SF, A-1231
 Antibiotic RK 4415, A-124
 Antibiotic RK 1441A, A-1232
 Antibiotic RK 1441B, A-1232
 Antibiotic RK 1409B, S-498
 Antibiotic RK 286C, S-498
 Antibiotic RK 286D, A-1190
 Antibiotic RK 1061G, L-188
 Antibiotic RK 1061H, L-188
 Antibiotic RK 1409, S-498
 Antibiotic RK-F 1010, T-96
 Antibiotic RKS 1778, A-1196
 Antibiotic Ro 1-9213, A-832
 Antibiotic Ro 5-9000, A-1081
 Antibiotic Ro 09-0049, P-337
 Antibiotic Ro 21-4958, R-151
 Antibiotic RP 13252, A-1234
 Antibiotic RP 6798, A-1233
 Antibiotic RP 18631, C-395
 Antibiotic RP 55185, C-545
 Antibiotic RP 59451, C-545
 Antibiotic RP 5278, S-569
 Antibiotic RUG 13A, R-157
 Antibiotic RUG 13B, R-157
 Antibiotic S 433, A-1235
 Antibiotic S 53210A, A-1236
 Antibiotic S 54832A, A-1237
 Antibiotic S 54832A-IV, A-1237
 Antibiotic S 632A₂, M-561
 Antibiotic S 632A₃, M-561
 Antibiotic S 632A₁, S-563
 Antibiotic S 632B₁, M-561
 Antibiotic S 632B₂, M-561
 Antibiotic S 5511I, R-34
 Antibiotic 1037, S-53
 Antibiotic 230, S-498
 Antibiotic 6741-21, S-321
 Antibiotic SA 3097B₁, A-1044
 Antibiotic SA 3097B₂, A-1044
 Antibiotic SA 3097C₁, A-1044
 Antibiotic SA 3097C₂, A-1044
 Antibiotic SA 3097D₁, A-1044
 Antibiotic SA 3097D₂, A-1044
 Antibiotic SA4-3, A-5
 Antibiotic SAB 711, A-1238
 Antibiotic SB 202327, A-1486
 Antibiotic SB 203207, A-660
 Antibiotic SB 203208, A-660
 Antibiotic SB 212305, A-1239
 Antibiotic SB 217452, A-1240
 Antibiotic SB 219383, A-1241
 Antibiotic SB 253514, A-1242
 Antibiotic SB 253517, A-1242
 Antibiotic SB 253518, A-1242
 Antibiotic SB 291071, A-1242
 Antibiotic SB 291072, A-1242
 Antibiotic SB 291073, A-1242
 Antibiotic SB 315021, A-1243
 Antibiotic SB 212021, H-665
 Antibiotic SB 200437, P-90
 Antibiotic SB 203105, P-90
 Antibiotic 446-S3-1, C-103
 Antibiotic Sch 210971, A-1250
 Antibiotic Sch 210972, A-1250
 Antibiotic Sch 213766, A-1250
 Antibiotic Sch 40832, A-1244
 Antibiotic Sch 40873, A-1245
 Antibiotic Sch 42137, A-1246
 Antibiotic Sch 52900, A-1247
 Antibiotic Sch 52901, A-1247
 Antibiotic Sch 538415, A-1251
 Antibiotic Sch 54445, A-1248
 Antibiotic Sch 56036, A-1249
 Antibiotic Sch 725424, A-1252
 Antibiotic Sch 18640A, A-1082
 Antibiotic Sch 18640B, A-1082
 Antibiotic Sch 575948, C-721
 Antibiotic Sch 38513, F-115
 Antibiotic Sch 38516, F-115
 Antibiotic Sch 38518, F-115
 Antibiotic Sch 39185, F-115
 Antibiotic Sch 42282, F-115
 Antibiotic Sch 42729, F-115
 Antibiotic Sch 18640, T-392
 Antibiotic SEN 34, H-562
 Antibiotic SEN 215, T-424
 Antibiotic SF 1346, A-822
 Antibiotic SF 1508, A-1253
 Antibiotic SF 1739, A-1254
 Antibiotic SF 2140, A-1255
 Antibiotic SF 2312, A-1256
 Antibiotic SF 2364, A-1157
 Antibiotic SF 2369, A-1257
 Antibiotic SF 2370, A-1189
 Antibiotic SF 2425, A-1258
 Antibiotic SF 2538, A-902
 Antibiotic SF 2695, A-1259
 Antibiotic SF 2768, A-1260

- Antibiotic SF 2771, A-1261
 Antibiotic SF 2809, A-1262
 Antibiotic SF 337, A-735
 Antibiotic SF 973A, A-14
 Antibiotic SF 2738A, C-574
 Antibiotic SF 2080A, D-356
 Antibiotic SF 2582A, D-954
 Antibiotic SF 2583A, O-157
 Antibiotic SF 2741A, P-641
 Antibiotic SF 2197B, A-1263
 Antibiotic SF 973B, A-14
 Antibiotic SF 2738B, C-574
 Antibiotic SF 2582B, D-953
 Antibiotic SF 2420B, M-491
 Antibiotic SF 2583B, O-157
 Antibiotic SF 2080B, P-963
 Antibiotic SF 2741B, P-642
 Antibiotic SF 2738, C-574
 Antibiotic SF 2738C, C-575
 Antibiotic SF 2582C, D-954
 Antibiotic SF 2080C, P-964
 Antibiotic SF 1917, D-73
 Antibiotic SF 2441, D-765
 Antibiotic SF 2738D, C-576
 Antibiotic SF 2080D, P-964
 Antibiotic SF 2738E, C-577
 Antibiotic SF 2080E, P-965
 Antibiotic SF 2738F, C-578
 Antibiotic SF 2080F_{2a}, P-966
 Antibiotic SF 2080F_{2b}, P-966
 Antibiotic SF 2080F₁, P-966
 Antibiotic SF 2080F₃, P-966
 Antibiotic SF 1739HP, N-32
 Antibiotic SF 2636, I-216
 Antibiotic SF 2494, T-658
 Antibiotic S-GI, H-616
 Antibiotic SL 3238, F-197
 Antibiotic SM 35, A-853
 Antibiotic SM 76, A-853
 Antibiotic SM 43, G-180
 Antibiotic SMTP 7, A-1264
 Antibiotic SMTP 8, A-1265
 Antibiotic SMTP 7D, A-1264
 Antibiotic SMTP 8D, A-1265
 Antibiotic SMTP 4D, S-332
 Antibiotic SMTP 5D, S-333
 Antibiotic SMTP 6D, S-334
 Antibiotic SMTP O, S-490
 Antibiotic SMTP 1, S-490
 Antibiotic SMTP 2, S-490
 Antibiotic SN 198B, D-203
 Antibiotic SN 198C, P-421
 Antibiotic SN 198D, D-203
 Antibiotic SN 198E, P-421
 Antibiotic SPF 32629A, A-1266
 Antibiotic SPF 32629B, A-1267
 Antibiotic SQ 02-S-L1, A-1268
 Antibiotic SQ 02-S-L2, S-492
 Antibiotic SQ 02-S-V2, A-1269
 Antibiotic SQ 02-S-V1, S-492
 Antibiotic SS 20846A, H-658
 Antibiotic SS 46506A, P-974
 Antibiotic SS 228B, A-1270
 Antibiotic SW 163C, A-1271
 Antibiotic SW 163D, A-1304
 Antibiotic SW 163E, A-1304
 Antibiotic SW 163F, A-1304
 Antibiotic SW 163G, A-1304
 Antibiotic SW-B, T-603
 Antibiotic T 988A, G-99
 Antibiotic T 988B, A-1272
 Antibiotic T 988C, G-99
 Antibiotic T 41348, H-514
 Antibiotic T 1384, N-180
 Antibiotic 46192, T-391
 Antibiotic 6739-15, T-392
 Antibiotic 6761-31, T-392
 Antibiotic TA 3037A, A-1273
 Antibiotic TAN 1263, A-1275
 Antibiotic TAN 1280, A-1276
 Antibiotic TAN 1307, A-1277
 Antibiotic TAN 1813, A-1278
 Antibiotic TAN 2020, A-1279
 Antibiotic TAN 883, A-1274
 Antibiotic TAN 1030A, A-1281
 Antibiotic TAN 1057A, A-1282
 Antibiotic TAN 1251A, A-1284
 Antibiotic TAN 876A, A-1280
 Antibiotic TAN 950A, A-733
 Antibiotic TAN 1771A, D-574
 Antibiotic TAN 868A, K-43
 Antibiotic TAN 1169A, P-400
 Antibiotic TAN 749A, S-389
 Antibiotic TAN 1057B, A-1282
 Antibiotic TAN 1251B, A-1284
 Antibiotic TAN 876B, A-1283
 Antibiotic TAN 1771B, D-572
 Antibiotic TAN 868B, K-43
 Antibiotic TAN 1169B, P-400
 Antibiotic TAN 883B, X-6
 Antibiotic TAN 1142, C-325
 Antibiotic TAN 1057C, A-1285
 Antibiotic TAN 1251C, A-1284
 Antibiotic TAN 749C, S-389
 Antibiotic TAN 1057D, A-1285
 Antibiotic TAN 1251D, A-1284
 Antibiotic TAN 588, L-8
 Antibiotic TAN 999, S-498
 Antibiotic TK 57-164A, D-98
 Antibiotic TK 57-164B, D-98
 Antibiotic TM 64, A-1286
 Antibiotic TMC 169, A-1502
 Antibiotic TMC 260, A-1290
 Antibiotic TMC 52, A-1287
 Antibiotic TMC 66, A-1288
 Antibiotic TMC 69, A-1289
 Antibiotic TMC 120A, A-1294
 Antibiotic TMC 135A, A-1295
 Antibiotic TMC 1A, A-1291
 Antibiotic TMC 2A, A-1292
 Antibiotic TMC 95A, A-1293
 Antibiotic TMC 49A, P-343
 Antibiotic TMC 120B, A-1294
 Antibiotic TMC 135B, A-1296
 Antibiotic TMC 2B, A-1292
 Antibiotic TMC 95B, A-1293
 Antibiotic TMC 34, C-634
 Antibiotic TMC 120C, A-1294
 Antibiotic TMC 1C, A-1297
 Antibiotic TMC 2C, A-1292
 Antibiotic TMC 52C, A-1287
 Antibiotic TMC 95C, A-1293
 Antibiotic TMC 1D, A-1297
 Antibiotic TMC 52D, A-1287
 Antibiotic TMC 95D, A-1293
 Antibiotic TMC 1E, M-89
 Antibiotic TMC 1F, M-87
 Antibiotic TMC 1G, M-88
 Antibiotic TMC 205, M-408
 Antibiotic TPU 0037A, L-353
 Antibiotic TPU 0037B, L-353
 Antibiotic TPU 0037C, L-353
 Antibiotic TPU 0037D, L-353
 Antibiotic TS 885, M-561
 Antibiotic Tu 2480F₁, P-893
 Antibiotic Tu 2480F₂, P-893
 Antibiotic Tu 2480F₃, P-893
 Antibiotic Tu 2480F₄, P-893
 Antibiotic U 106305, A-1299
 Antibiotic U 42126, A-81
 Antibiotic U 43795, A-81
 Antibiotic U 62162, A-1298
 Antibiotic U 27810, B-104
 Antibiotic U 77779, B-229
 Antibiotic U 54703, C-521
 Antibiotic U 11994, D-232
 Antibiotic U 9361, D-765
 Antibiotic U 54702D, C-521
 Antibiotic U 20207, E-229
 Antibiotic U 47929, F-52
 Antibiotic U 48266, F-33
 Antibiotic U 15965, L-166
 Antibiotic U 24792, L-241
 Antibiotic U 77863, M-513
 Antibiotic U 59760, N-276
 Antibiotic U 59761, N-276
 Antibiotic U 61732, N-276
 Antibiotic U 22662, P-876
 Antibiotic U 24544, P-919
 Antibiotic U 48160, Q-61
 Antibiotic U 19183, S-384
 Antibiotic UCA 1064A, A-1602
 Antibiotic UCA 1064B, A-1602
 Antibiotic UCF1B, M-88
 Antibiotic UCF1C, M-87
 Antibiotic UCH 15A, A-1300
 Antibiotic UCH 15B, A-1300
 Antibiotic UCN 01, S-498
 Antibiotic UCN 02, S-498
 Antibiotic UCS 1025A, A-1301
 Antibiotic UCS 1025B, A-1301
 Antibiotic UK 1, A-1302
 Antibiotic UK 2, A-1303
 Antibiotic UK 63052, A-1304
 Antibiotic UK 88051, A-1141
 Antibiotic UK 3A, A-1303
 Antibiotic UK 111866, P-90
 Antibiotic USF 19A, A-1305
 Antibiotic USF 142A, M-512
 Antibiotic USF 406C, D-681
 Antibiotic USF 406D, D-681
 Antibiotic vD 844, O-124
 Antibiotic vD 846, O-124
 Antibiotic VM 55595, A-1306
 Antibiotic VM 55598, A-1486
 Antibiotic VM 55599, A-1307
 Antibiotic VM 29919, P-90
 Antibiotic VM 54158, P-91
 Antibiotic VM 54159, P-90
 Antibiotic VM 55594, P-91
 Antibiotic VM 55596, P-90
 Antibiotic VM 55597, P-90
 Antibiotic 10381V, S-638
 Antibiotic WB 2838, A-725
 Antibiotic WF 14865, A-1308
 Antibiotic WF 6237, A-680
 Antibiotic WF 11231A, A-1178
 Antibiotic WF 14861A, A-1309
 Antibiotic WF 16775A₁, A-1535
 Antibiotic WF 16775A₂, A-1535
 Antibiotic WF 1360B, R-83
 Antibiotic WF 1360C, R-83
 Antibiotic WF 1360F, R-83
 Antibiotic WF 1360, R-83
 Antibiotic WIN 64745, A-1310
 Antibiotic WIN 64821, A-1311
 Antibiotic WJ 35, A-1312
 Antibiotic WJ 85, A-1313
 Antibiotic WJ 85B, A-1313
 Antibiotic WK 5344A, F-45
 Antibiotic WK 5344B, F-45
 Antibiotic 10381W, P-641
 Antibiotic WS 1921, A-1314
 Antibiotic WS 30581A, A-622
 Antibiotic WS 75624A, A-1315
 Antibiotic WS 9659A, L-76
 Antibiotic WS 30581B, A-622
 Antibiotic WS 75624B, A-1316
 Antibiotic WS 63967B, C-129
 Antibiotic WS 9659B, L-76
 Antibiotic WS 63967, C-129
 Antibiotic WS 30581C, A-622
 Antibiotic WS 63967D, C-129
 Antibiotic WS 63967E, C-129
 Antibiotic WS 63967F, C-129
 Antibiotic X 14885A, A-1087
 Antibiotic X 14547A, I-64
 Antibiotic X 53111, Q-58
 Antibiotic X 948, Q-58
 Antibiotic XK 211, A-1317
 Antibiotic XK 101-1, T-658
 Antibiotic 10381X, P-642
 Antibiotic XR 587, S-571
 Antibiotic Y 03559J-A, A-1318
 Antibiotic Y 05460MA, A-1319
 Antibiotic Y 16482 α , S-2
 Antibiotic Y 16482 β , S-2
 Antibiotic Y 16601, S-2
 Antibiotic Y 16760, S-2

- Antibiotic YC 73, F-97
 Antibiotic YCM 1008A, A-1320
 Antibiotic Y1, K-37
 Antibiotic YL 03709B-A, D-574
 Antibiotic YL 0358M-A, H-77
 Antibiotic YL 0358M-B, H-77
 Antibiotic YL 02729S, M-491
 Antibiotic YM 193221, A-1321
 Antibiotic YM 215343, A-1340
 Antibiotic YM 216391, A-1322
 Antibiotic YM 47522, D-574
 Antibiotic YM 30059, M-491
 Antibiotic YM 266183, T-383
 Antibiotic YM 266184, T-383
 Antibiotic YN 0165JA, A-1323
 Antibiotic YP 0298L-C, L-76
 Antibiotic 10381Y, S-638
 Antibiotic YSI 40-2, M-177
 Antibiotic Y-T 0678H, B-72
 Antibiotic YUA 001, H-667
 Antibiotic 1745Z₃A, O-131
 Antibiotic ZG 1494 α , O-134
 Antibiotic ZHD 0501, A-1324
 Antibiotic 10381Z₁, S-638
 Antibiotic 10381Z₂, S-638
 3C-Antibiotic, P-321
 Antidesmone, A-1325
 Antidysentericine, C-595
 Antiepilepsirine, M-444
 Antifungin, A-1326
 Antilirium, P-400
 Antillaridine, S-458
 Antillidine, S-458
 Antineoplaston A10, D-808
 (-)-Antioquine, C-643
 (+)-Antioquine, R-114
 Antiostatin B, A-1327
 Antipellagra factor, P-895
 Antirhine lactone, A-1329
 Antirhine α -*N*-methosalt, A-1328
 Antirhine β -*N*-methosalt, A-1328
 Antirhine, A-1328
 Antofine α -*N*-oxide, A-1330
 Antofine β -*N*-oxide, A-1330
 Antofine, A-1330
 Antramycin, A-1081
 Antrodrin B, H-684
 Antrodrin C, H-684
 Antrodrin D, H-684
 Antrodrin E, H-684
 Antropyrrroledione, H-684
 AnTx, A-974
 Anzatax, T-52
 1494A, O-134
 Aobamidine, A-144
 Aobamine, C-676
 Apateline, A-1331
 Apc, A-1332
 APEA, A-892
 Aphanorphine, A-1333
 APHE 1, A-622
 APHE 2, A-622
 APHE 3, A-622
 APHE 4, A-622
 Aphelandrine, A-1334
 Aphrodine, Y-13
 Aphyllic acid, A-1335
 Aphyllidine, A-1336
 Aphyllidinic acid, A-1335
 Aphylline *N*¹⁶-oxide, A-1337
 Aphylline, A-1337
 Aphyllinic acid, A-1335
 Apicularen A, A-1338
 Apicularen B, A-1338
 Apidionene, A-1339
 Apiosporamide, A-1340
 6'-Apiosyllotaustralin, H-572
 Aplaminal, A-1341
 Aplaminone, A-1342
 Aplicyanin A, A-1343
 Aplicyanin B, A-1343
 Aplicyanin C, A-1343
 Aplicyanin D, A-1343
 Aplicyanin E, A-1343
 Aplicyanin F, A-1343
 Aplidiamine, A-819
 Aplidinone A, A-1344
 Aplidinone B, A-1345
 Aplidinone C, A-1345
 Aplidite A, A-1346
 Aplidite B, A-1346
 Aplidite C, A-1346
 Aplidite D, A-1346
 Aplidite E, A-1346
 Aplidite F, A-1346
 Aplidite G, A-1346
 Aplysamine 1, M-679
 Aplysamine 2, P-823
 Aplysamine 3, P-823
 Aplysamine 4, P-823
 Aplysamine 5, P-823
 Aplysepine, A-1347
 Aplysidine, T-360
 Aplysellamide A, A-719
 Aplysellamide B, A-719
 Aplysinafulvin, A-1348
 Aplysinamisine I, A-1349
 Aplysinamisine II, A-1350
 Aplysinamisine III, A-1378
 Aplysinellin A, A-1351
 Aplysinellin B, A-1352
 Aplysinketal A, A-1353
 Aplysinketal B, A-1354
 Aplysinone A, A-1355
 Aplysinone B, A-192
 Aplysinone C, A-1355
 Aplysinone D, A-1356
 Aplysinopsin, A-1357
 Aplyzanzine A, A-1358
 Apoptropine, A-1359
 Apocavidine, C-222
 Apochlorine, A-1361
 Apocinchonamine, A-1360
 Apocrotonosine, T-524
 Apo-14,15-dehydrovincamine, V-115
 Apodihydrocinchonamine, A-1360
 Apodine, M-666
 Apodinine, M-666
 Apoferrerosamine, D-504
 Apogalanthamine, A-1361
 Apogeissoschizine, A-1362
 Apoglaziovine, T-524
 (\pm)-Apoglaziovine, T-524
 (+)-Apoglaziovine, T-524
 Apohaemanthamine, A-1363
 Apohemanthamine, A-1363
 Apohyosine, A-1364
 Apokhlorin, A-1361
 Aponoratropine, A-1359
 Aponorhyosine, A-1364
 Apophedrin, A-864
 Apoplone, R-52
 Aporheine *N*-methosalt, R-120
 Aporheine, R-120
 Aporphinium DVT, H-751
 Aposcopolamine, A-1364
 Apotacamine, T-12
 Apoterin, R-50
 Apovincamine, V-115
 Apo- Δ ¹⁴-vincamine, V-115
 Apparicine, A-1365
 Apratoxin A, A-1366
 Apratoxin B, A-1366
 Apratoxin C, A-1367
 Aprv, V-115
 A528, Q-60
 AQC-3, R-76
 Aquamycin, B-447
 Aquaticine, A-1368
 Aquifoline, I-328
 Aquifoliunine EIII, P-247
 Aquifoliunine EII, P-247
 Aquifoliunine EI, P-247
 Aquifoliunine EIV, P-247
 Aquiledine, A-1369
 Aquilegine, A-1370
 Ara Tb, T-658
 Arabinofuranosylcytosine, C-962
 Arabinosylcytosine, C-962
 Arachidonic acid, E-48
O-Arachidonoyl ethanolamine, E-48
 1-*O*-Arachidonoyl glycerol, L-348
 Arachidonyl ethanolamide, E-48
N^b-Arachidonyl tryptamine, T-639
N-Arachidylanthranilic acid, A-706
 Arachine, C-419
 Ara-doridosine, A-818
 Aragupetrosine A, A-1371
 Araguspongine A, X-24
 Araguspongine B, X-24
 Araguspongine C, A-1372
 Araguspongine D, X-24
 Araguspongine E, X-24
 Araguspongine F, A-1373
 Araguspongine G, A-1373
 Araguspongine H, A-1374
 Araguspongine J, A-1374
 Araguspongine K, X-24
 Araguspongine L, A-1372
 Araguspongine M, X-24
 Aralionine A, A-1375
 Aralionine B, C-600
 Aralionine C, A-1375
 Aralionine, A-1375
 Araliopdimerine A, A-1376
 Araliopdimerine B, A-1376
 Araliopdimerine C, P-87
 Araliopsine, A-1377
 Araliopsinine, F-90
 Araplysellin I *N*²⁰-sulfamate, A-1378
 Araplysellin I, A-1378
 Araplysellin II, A-1378
 Araplysellin III, A-1379
 Araquine, C-419
 Araregai toxin, T-298
 Aratubercidin, T-658
 Arboflorine, A-1380
 Arbolosine, A-1381
 Arbophylline, A-1382
 Arborescidine A, O-48
 Arborescidine B, A-1383
 Arborescidine C, A-1383
 Arborescidine D, A-1383
 Arboreumine, A-1384
 Arboricine, A-1385
 Arboricine, D-657
 Arboricine, A-1386
 Arborine, B-93
 Arborinine, T-516
 Arcaine, D-400
 Archaeosine, A-1387
 Archazolide A, A-1388
 Archazolide B, A-1389
 Archazolide C, A-1388
 Archazolide D, A-1388
 Archerine, A-1390
 Archil, O-102
 Arcutine, A-1391
 Arcutinidine, A-1391
 Arcutinine, A-1391
 Arcyriacyanin A, A-1392
 Arcyriaflavin A, A-1393
 Arcyriaflavin B, A-1393
 Arcyriaflavin C, A-1393
 Arcyriaflavin D, A-1393
 Arcyriarubin A, A-1394
 Arcyriarubin B 6'-sulfate, A-1394
 Arcyriarubin B, A-1394
 Arcyriarubin C, A-1394
 Arcyriaverdin C, A-1395
 Arcyrin A, A-1396
 Arcyrin B, A-1396
 Arcyroxepin A, A-1398
 Arcyroxepin B, A-1398
 Arcyroxindole A, A-1397
 Arcyroxocin A, A-1398
 Arcyroxocin B, A-1398
 Ardeemin, A-1399
 Ardine, M-436

- Arecaidine, T-205
 Arecaine, T-205
 Arecolidine, A-1400
 Arecoline, T-205
 Arenaine, A-1401
 Arenarine A, A-1402
 Arenarine B, A-1402
 Arenarine C, A-55
 Arenarine D, A-55
 Arenine, A-1403
 Arenosclerin A, A-1404
 Arenosclerin B, A-1404
 Arenosclerin C, A-1404
 Arenosclerin D, A-1404
 Arenosclerin E, A-1405
 Arestrictin A, A-1406
 Arestrictin B, A-1406
 Arg, A-1413
 Argadin, A-1407
 Argemexicaine A, A-1408
 Argemexicaine B, A-1409
 (-)-Argemone *N*-methosalt, A-1410
 (-)-Argemone *N*-oxide, A-1410
 Argemone, A-1410
 Argenaxine, C-72
 Argentamine, B-20
 Argentinamine, A-1411
 Argentine, A-1412
 Argentinine *N*-oxide, A-1532
 Argentinine, A-1532
 L-Arginine, A-1413
 β -Arginine, A-779
 Arginine, A-1413
N-(*N*²-L-Argininyl)cantharimide, C-89
 Argiopine, A-1414
 Argiopinin I, A-1415
 Argiopinin II, A-1415
 Argiopinin III, A-1417
 Argiopinin IV, A-1416
 Argiopinin V, A-1416
 Argiotoxin 636, A-1414
 Argiotoxin 659, A-1417
 Argiotoxin 673, A-1418
 Arglecin, A-1419
 Argutane A, A-1420
 Argutane B, A-1420
 Argvalin, A-1421
 Argyrolobine, A-1336
 Ariakemicin A, A-1422
 Ariakemicin B, A-1422
 Aribine, M-426
 Aricine pseudoindoxyl, T-289
 Aricine, A-1423
 Ariskanin A, D-636
 Ariskanin B, T-568
 Ariskanin C, T-568
 Ariskanin D, T-569
 Ariskanin E, T-569
 Aristinic acid, M-262
 Aristo red, A-1424
 Aristoaristone, A-1425
 Aristocarinol, A-1442
 Aristofolin A, M-263
 Aristofolin B, M-263
 Aristofolin C, M-263
 Aristofolin D, M-263
 Aristofrucosine, A-1426
 Aristolactam A Ia, T-539
 Aristolactam A II, D-567
 Aristolactam A IIIa, T-540
 Aristolactam A III, T-540
 Aristolactam B II, D-567
 Aristolactam B III *N*-glucoside, T-540
 Aristolactam B III, T-540
 Aristolactam B I, T-539
 Aristolactam C II, A-1428
 Aristolactam C III, A-1428
 Aristolactam C IV, A-1428
 Aristolactam D II, A-1428
 Aristolactam D III, A-1428
 Aristolactam E, T-252
 Aristolactam F I, D-567
 Aristolactam F II, T-537
 Aristolactam F, T-540
 Aristolactam I *N*-(6'-*trans-p*-coumaroyl)- β -D-glucopyranoside, A-1427
 Aristolactam I *N*-glucoside, A-1427
 Aristolactam Ia *N*- β -D-glucoside, A-1427
 Aristolactam Ia *O*-glucoside, A-1427
 Aristolactam I, A-1427
 Aristolactam Ia, A-1427
 Aristolactam II, D-567
 Aristolactam IIIa *N*-glucoside, A-1429
 Aristolactam IIIa *O*-glucoside, A-1429
 Aristolactam III, A-1429
 Aristolactam IIIa, A-1429
 Aristolactam IV, T-253
 Aristolactam, A-1427
 Aristolamide, M-263
 Aristolarine, A-1430
 Aristolasene, A-1431
 Aristolasicol, A-1432
 Aristolasicolone, A-1432
 Aristolasicone, A-1432
 Aristolasol, A-1433
 Aristolic acid, M-263
 Aristolin, A-1434
 Aristolinic acid, M-263
 Aristoliukine A, A-1428
 Aristoliukine B, T-662
 Aristoliukine C, T-662
 Aristolochia yellow, M-262
 Aristolochic acid A, M-262
 Aristolochic acid BII methyl ester, D-636
 Aristolochic acid B, M-447
 Aristolochic acid C, H-600
 Aristolochic acid D, T-266
 Aristolochic acid E, T-267
 Aristolochic acid Ia methyl ester, M-262
 Aristolochic acid Ia, M-262
 Aristolochic acid II methyl ester, M-447
 Aristolochic acid III methyl ester, H-600
 Aristolochic acid IIIa, H-600
 Aristolochic acid III, H-600
 Aristolochic acid II, M-447
 Aristolochic acid I, M-262
 Aristolochic acid IV methyl ester, T-266
 Aristolochic acid IVa, T-266
 Aristolochic acid IV, T-266
 Aristolochic acid Va, T-265
 Aristolochic acid VIa, T-264
 Aristolochic acid VII methyl ester, T-267
 Aristolochic acid VIIa, T-267
 Aristolochic acid VII, T-267
 Aristolochic acid V, T-265
 Aristolochic acid, M-262
 Aristolochine, C-812
 Aristolochin, M-262
 Aristolochin, T-266
 Aristolodione, C-273
 Aristoloin I, A-1435
 Aristoloin II, A-1435
 Aristololactam, A-1427
 Aristololide, T-266
 Aristoloterpenate I, A-1436
 Aristoloterpenate II, A-1436
 Aristoloterpenate III, A-1436
 Aristoloterpenate IV, A-1436
 Aristomakine, A-1437
 Aristomakinine, A-1437
 Aristomanoside, H-566
 Aristone, A-1438
 Aristophyllide A, A-1439
 Aristophyllide B, A-1439
 Aristophyllide C, A-1439
 Aristophyllide D, A-1439
 Aristoquinoline A, A-1440
 Aristoquinoline B, A-1440
 Aristoquinoline C, A-1440
 Aristoserratenine, A-1441
 Aristoserratine, A-1442
 Aristoteline, A-1442
 Aristotelinine, S-259
 Aristotelinone, A-1442
 Aristotelone, A-1430
 Aristoyagonine, A-1443
 Arizonine, T-163
 Armatamide, Z-6
 Armepravine, A-1444
 Armillarisin B, H-688
 Arnoamine A, A-1445
 Arnoamine B, A-1445
 Arnottianamide, A-1446
 Arolycoricidine, C-744
 Arolycoricidinol, C-744
 Aromadendrane 1-isonitrile, A-1447
 Aromadendrane 1-isothiocyanate, A-1447
 Aromoline, O-1
 Arosine, G-93
 Arosinine, G-93
 Artabonatine A, A-1448
 Artabonatine B, G-213
 Artabonatine C, A-1449
 Artabonatine D, A-1449
 Artabonatine E, A-1448
 Artabonatine F, U-36
 Artabotrine, C-272
 Artabotrinine, A-1066
 Artacinatine, A-1450
 Artarine, D-407
 Artavenustine 11-*O*- β -D-glucopyranoside, X-30
 Artavenustine, X-30
 Artemicidine, A-660
 Arterenol, N-298
 Arterenone, A-736
 Arthrobactin, A-1451
 Artrichine, C-569
 Arudonine, S-456
 Arundacine, A-1452
 Arundamine, A-1452
 Arundanine, A-1452
 Arundarine, A-1453
 Arundine, M-436
 Arundinine, A-1454
 Arvelexin, H-524
 Aryapavine, M-140
 Asaramide, A-1455
 Ascidiacyclamide, A-1456
 Ascidiathiazone A, A-1457
 Ascidiathiazone B, A-1458
 Ascididemine, A-1459
 Ascididemnin, A-1459
 Ascochalin, A-1460
 Ascophylline, P-229
 Ascorbigen A, A-1461
 Ascorbigen B, A-1461
 Ascorbigen, A-1461
 Ascosalipyrrolidinone A, A-1462
 Ascosalipyrrolidinone B, A-1462
 Ascosonchine, O-205
 Asimicilone, A-1463
 Asimilobine 2-glucoside, D-531
 Asimilobine, D-531
 ASK 753, A-1117
 ASL 279, D-920
 Asmarine A, A-1465
 Asmarine B, A-1465
 Asmarine C, A-1466
 Asmarine D, A-1466
 Asmarine E, A-1466
 Asmarine F, A-1466
 Asmarine G, A-1465
 Asmarine H, A-1465
 Asmarine I, A-1464
 Asmarine J, A-1464
 Asmarine K, A-1465
 AS-N-7A, A-1118
 Asparagamine A, S-515
 Aspartic acid betaxanthin, M-646
 1-L-Aspartic acid- α -amanitin, A-673
 1-L-Aspartic acid-3-(*S*)-4-hydroxy-L-isoleucine- α -amanitin, A-673
 1-L-Aspartic acid-3-isoleucine- α -amanitin, A-673
 1-L-Aspartic acid-4-(2-mercapto-L-tryptophan)- α -amanitin, A-673
 L-Aspartyl-L-*N*²-hydroxyaspartyl-D-cycloserine, A-1467
*N*²-Aspartyllobmbricine, L-240
 Asperazine, A-1468

- Asperfumoid, A-1469
 Aspergamide A, A-1470
 Aspergamide B, A-1470
 Aspergillamide A, A-1471
 Aspergillamide B, A-1471
 Aspergillazine A, A-1472
 Aspergillazine B, A-1473
 Aspergillazine C, A-1473
 Aspergillazine D, A-1474
 Aspergillazine E, A-1474
 Aspergillilic acid, A-1475
 Aspergillimide, A-1486
 Aspergillin PZ, A-1476
 Aspergillin, A-1475
 Aspergillitine, A-1477
 Asperlicin B, A-1478
 Asperlicin C, A-1479
 Asperlicin D, A-1480
 Asperlicin E, A-1481
 Asperlicin, A-1478
 Aspernigerin, A-1482
 Aspernigrin A, A-1483
 Aspernigrin B, A-1484
 Aspernomine, A-1485
 Asperparaline A, A-1486
 Asperparaline B, A-1486
 Asperparaline C, A-1486
 Aspertine A, H-433
 Aspertine B, A-1487
 Aspertine C, B-187
 Aspertine D, H-433
 Asperumine *N*-oxide, T-188
 Asperumine, T-188
 Aspidexcelcine, A-1488
 Aspidexcine, A-1489
 Aspidoalbidine, A-1490
 Aspidoalbine, A-1490
 Aspidocarpine, P-924
 Aspidochibine, A-1491
 Aspidodascarpine, A-1492
 Aspidodispermine, A-1493
 Aspidofendlerine, A-1490
 Aspidofiline, A-1494
 Aspidofractine, K-78
 Aspidofractinine, A-1494
 Aspidofractinin-16-ol, H-406
 Aspidolimidine, A-1490
 Aspidolimidinol, L-177
 Aspidolimine, P-924
 Aspidophylline A, A-1495
 Aspidophytine, A-1496
 Aspidosine, A-1499
 Aspidospermatidine, A-1497
 Aspidospermatine, A-1497
 Aspidospermidine, A-1498
epi- Δ^1 - ψ -Aspidospermidine, D-377
 Δ^1 - ψ -Aspidospermidine, D-377
 Aspidospermidose, A-1498
 Aspidospermine, A-1499
 Aspidospermiose, A-1498
 Aspochalamin A, A-1500
 Aspochalamin B, A-1500
 Aspochalamin C, A-1500
 Aspochalamin D, A-1500
 Aspochalasin A, A-1502
 Aspochalasin B, A-1502
 Aspochalasin C, A-1502
 Aspochalasin D, A-1502
 Aspochalasin E, A-1502
 Aspochalasin F, A-1503
 Aspochalasin G, A-1502
 Aspochalasin H, A-1502
 Aspochalasin I, A-1501
 Aspochalasin J, A-1501
 Aspochalasin K, A-1502
 Aspochalasin Z, A-1502
 Aspoquinolone A, A-1504
 Aspoquinolone B, A-1504
 Aspoquinolone C, A-1505
 Aspoquinolone D, A-1505
 Asposterol, A-1502
 Aspyridone A, A-1506
 Aspyridone B, A-1506
 Assamicadine, A-1507
 Assoanine, A-1508
 Astechrome, A-1509
 Asteranthine, A-1510
 Asterin A, A-1511
 Asterin B, A-1511
 Asterinin A, A-1512
 Asterinin B, A-1512
 Asterinin C, A-1513
 Asterinin D, A-1523
 Asterinin E, A-1523
 Asterinin F, A-1523
 Asterionellin A, A-1514
 Asterionellin B, A-1515
 Asterionellin C, A-1515
 Asteropterine, A-1516
 Asterredione, A-1517
 Asterridinone, A-1518
 Asterriquinol D, A-1519
 Asterriquinone A1 hydroquinone, A-1519
 Asterriquinone A1, A-1520
 Asterriquinone A2, A-1521
 Asterriquinone A3, A-1521
 Asterriquinone A4, A-1521
 Asterriquinone B1, A-1521
 Asterriquinone B2, A-1521
 Asterriquinone B3, A-1521
 Asterriquinone B4, A-1521
 Asterriquinone C1 hydroquinone, A-1519
 Asterriquinone C1, A-1521
 Asterriquinone C2, A-1521
 Asterriquinone CT1, A-1521
 Asterriquinone CT2, A-1521
 Asterriquinone CT3, A-1521
 Asterriquinone CT4, A-1521
 Asterriquinone CT5, A-1521
 Asterriquinone D hydroquinone, A-1519
 Asterriquinone D, A-1521
 Asterriquinone E, A-1521
 Asterriquinone monoacetate, A-1520
 Asterriquinone PM 53, A-1521
 Asterriquinone SU 5501, A-1521
 Asterriquinone SU 5503, A-1521
 Asterriquinone SU 5220, I-191
 Asterriquinone SU5228, A-1521
 Asterriquinone SU5500, A-1521
 Asterriquinone SU5504, A-1520
 Asterriquinone, A-1520
 Asterubin, A-1522
 Astin J, A-1523
 Astramorph P/F, M-704
 Astrimalvine A *N*-oxide, M-393
 Astrimalvine B, M-393
 Astrocasine, A-1524
 Astrophylline, A-1525
 AT 125, A-81
 AT 2433, A-1119
 AT 2433A₁, A-1119
 AT 2433A₂, A-1119
 AT 2433B₁, A-1119
 AT 2433B₂, A-1119
 AT 1001, H-616
 AT 1840, U-34
 Atalafoline B, P-207
 Atalafoline, P-206
 Atalanine, A-1526
trans-Atalantine, A-1526
 Atalaphyllidine, A-1527
 Atalaphyllidine, T-548
 Atalaphylline dimethyl ether, T-546
 Atalaphylline, T-546
 Atalaphyllinine, S-278
 Ataline, A-1528
 Atanine, A-1529
 Atanisatine, A-1530
 Atelopidtoxin, Z-18
 Atemoine, K-6
 ATF 606, M-159
 Atheroline, O-178
 Atherophylline, P-923
 Atherospermidine, A-1531
 Atherosperminine *N*-oxide, A-1532
 Atherosperminine, A-1532
 Atherospermoline, T-286
 Athrocupressine, C-588
 Atidine, A-1533
 Atisine azomethine, A-1609
 Atisine, A-1534
 Atisinium chloride, A-1534
 Atpenin, A-1535
 Atractylenolactam, A-1536
 Atropamine, A-1359
 Atropigen, T-632
 Atropine methonitrate, T-632
 Atropine oxide hydrochloride, T-632
 Atropine oxide, T-632
 Atropine propionate, T-632
 Atropine sulfate, T-632
 Atropine, T-632
 Atropisol, T-632
 Atropoylnortropan-3-ol, A-1359
 Atropurpursine, M-329
 Atropyltrophein, A-1359
 Atroscine, S-159
 Aucherine, A-1538
 Aucubinine A, C-562
 Aucubinine B, C-562
 Audouine, D-401
 Augmentin, G-201
 Augustamine, A-1539
 Augustine, C-748
 Aulamine, L-348
 Aulosirazole, A-1540
 Aurachin A, A-1541
 Aurachin B, A-1542
 Aurachin C, A-1543
 Aurachin D, A-1543
 Aurachin E, A-1544
 Aurachin F, A-1545
 Aurachin G, A-1546
 Aurachin H, A-1546
 Aurachin I, A-1546
 Auranthine, A-1547
 Aurantiamine, A-1548
 Aurantioclavine, A-1549
 Aurantiomide A, A-966
 Aurantiomide B, A-966
 Aurantiomide C, A-966
 Aurantoside A, A-1550
 Aurantoside B, A-1550
 Aurantoside C, A-1551
 Aurantoside D, A-1552
 Aurantoside E, A-1552
 Aurantoside F, A-1553
 Aurantoside G, A-1554
 Aurantoside H, A-1554
 Aurantoside I, A-1554
 Aureine, S-238
 Aureothricin, A-759
 Aureoverticillactam, A-1555
 Auricularine, B-253
 Auriculine, P-39
 Auroramine, S-192
 Aurotensine, S-164
 Austamide, A-1556
 Australiflorine, T-555
 Australine, T-555
 Austroconitine A, M-355
 Austroconitine B, M-303
 Austrodimerine, A-1557
 Austronine, P-252
 Austrospicatin, T-48
 Austrotaxine, P-224
 Austyn, T-360
 Autumnaline, A-1558
 Auxarconjugatin A, A-1559
 Auxarconjugatin B, A-1559
 Auxarconjugatin C, A-1559
 Auxoperan, C-420
 AV toxin C, A-862
 AV toxin D, T-599
 AV toxin E, T-599
 Avadharidine, P-776
 Avarol, A-1560
 Avenalumin I, A-789
 Avenalumin II, A-1561

- Avenalumin III, A-1562
 Avenanthramide 1, A-789
 Avenanthramide 2, A-737
 Avenanthramide A, A-789
 Avenanthramide B, A-789
 Avenanthramide C, A-789
 Avenanthramide D, A-1563
 Z-Avenanthramide E, A-1563
 Avenanthramide E, A-1563
 Avenanthramide G, A-788
 Avenanthramide L, A-1564
 Avicennamine, A-1565
 Avicine, A-1566
 Avinosol, A-1567
 Avrainvillamide, A-1470
 Awaine, A-1568
 Awajanomycin, A-1569
 AX 2, I-246
 Axamide 1, A-1570
 Axamide 2, A-1571
 Axamide 4, A-1570
 Axamide 3, S-425
 Axialit, E-157
 Axillaridine A, D-274
 Axillaridine, A-1572
 Axillarine A, D-264
 Axillarine B, D-263
 Axillarine C, D-262
 Axillarine D, D-262
 Axillarine E, D-262
 Axillarine F, D-262
 Axillarine, A-1572
 Axinellamide, A-1573
 Axinellamine A, A-1574
 Axinellamine A, M-496
 Axinellamine B, A-1574
 Axinellamine B, A-1575
 Axinellamine C, A-1574
 Axinellamine D, A-1574
 Axinohydantoin, A-1576
 (Z)-Axinohydantoin, A-1576
 Axinyssimide A, A-1577
 Axinyssimide B, A-1578
 Axinyssimide C, A-1578
 Axisonitrile 1, A-1570
 Axisonitrile 2, A-1571
 Axisonitrile 4, A-1570
 Axisonitrile 3, S-425
 Axisothiocyante 1, A-1570
 Axisothiocyante 2, A-1571
 Axisothiocyante 4, A-1570
 Axisothiocyante 3, S-425
 Axsain, C-100
 Ayuthianine, T-522
 Azaacridone A, A-1579
 2-Azaanthraquinone, B-71
 Azabenzene, P-894
 6-Azabenzob[*b*]carbazole, Q-19
 1-Azabicyclo[4.3.0]nonane-7,8-diol, O-46
 9-Azabicyclo[3.3.1]nonan-3-one, A-1580
 1-(9-Azabicyclo[4.2.1]non-2-en-2-yl)ethanone, A-974
 1-(9-Azabicyclo[4.2.1]non-2-en-2-yl)-1-propanone, H-325
 8-Azabicyclo[3.2.1]octane-2,3-diol, A-1581
 8-Azabicyclo[3.2.1]octane-2,6-diol, A-1582
 8-Azabicyclo[3.2.1]octane-2,7-diol, A-1583
 8-Azabicyclo[3.2.1]octane-3,6-diol, A-1584
 8-Azabicyclo[3.2.1]octane-3,7-diol, A-1585
 8-Azabicyclo[3.2.1]octane-1,2,3,4,6-pentol, A-1586
 8-Azabicyclo[3.2.1]octane-1,2,3,4-tetrol, A-1587
 8-Azabicyclo[3.2.1]octane-1,2,3,6-tetrol, A-1588
 8-Azabicyclo[3.2.1]octane-1,2,4,7-tetrol, A-1589
 8-Azabicyclo[3.2.1]octane-1,2,3-triol, A-1590
 8-Azabicyclo[3.2.1]octane-1,2,4-triol, A-1591
 8-Azabicyclo[3.2.1]octane-1,2,7-triol, A-1592
 8-Azabicyclo[3.2.1]octane-1,3,4-triol, A-1593
 8-Azabicyclo[3.2.1]octane-2,3,4-triol, A-1594
 8-Azabicyclo[3.2.1]octane-2,3,6-triol, A-1595
 8-Azabicyclo[3.2.1]octane-3,6,7-triol, A-1596
 8-Azabicyclo[3.2.1]octan-3-ol, A-1597
 1-Azabicyclo[3.3.0]oct-3-en-2-one, T-229
 8-Azabicyclo[3.2.1]oct-3-yl 3,4-dimethoxybenzoate, C-633
 Azacridone A, A-1579
 Azacyclohexane, P-454
 Azacyclopentane, P-949
 N-(3-Azacyclotetradec-1-ylpropyl)-1,3-propanediamine, M-715
 N-(3-Azacyclotridec-1-ylpropyl)-1,3-propanediamine, M-714
 Azacyclo-6-undecene, A-1598
 4-Aza-1,8-diguanidinooctane, H-305
 1-Aza-1,2-dihydro-7,8,9,10-tetramethoxy-4-methyl-2-oxoanthracene, T-277
 9-Azafluorene, C-120
 8-Azaguanine, A-735
 8-Azaguanosine 5'-phosphate, A-735
 8-Azaguanosine, A-735
 3-Aza-A-homoandrostan-17-ol 1,4-epoxide, E-127
 3-Aza-A-homoandrostan-16-ol, S-47
 15-Aza-D-homoergosta-8,14,24(28)-trien-3-ol, A-1602
 3-Aza-6-hydroxybicyclo[3.3.0]oct-1(5)-ene-2,4-dione, M-70
 1-Azaindene, I-79
 8-O-Azaloyl-14-benzoylaconine, M-331
 Azamerone, A-1599
 Azamonosporascone, D-534
 Azan, A-735
 2-Azanaphthalene, I-301
 1-Azanaphthalene, Q-29
 4-Azaskatole, M-551
 Azaspiracid 10, A-1600
 Azaspiracid 11, A-1600
 Azaspiracid 12, A-1600
 Azaspiracid 1, A-1600
 Azaspiracid 2, A-1600
 Azaspiracid 3, A-1600
 Azaspiracid 4, A-1600
 Azaspiracid 5, A-1600
 Azaspiracid 6, A-1600
 Azaspiracid 7, A-1600
 Azaspiracid 8, A-1600
 Azaspiracid 9, A-1600
 Azaspirene, A-1601
 2-Azapiro[5.5]undecan-7-ol, N-241
 15-Azasterol, A-1602
 1-Aza-2,5,8-trioxacyclononane, A-1603
 Azatyrosine, A-822
 Azcarpine, A-1604
 6*H*-Azepino[3',2':4,5]cyclopent[1,2-*g*]indole-7,11-dione, M-694
 Azepinomycin, A-1605
 Azepinostatin, B-9
 3,3-Azetidinediol, A-1606
 3-Azetidinone, A-1606
 6-Azidotetrazolo[5,1-*a*]phthalazine, A-1607
 Azimine, A-1608
 Azine, P-894
 Azitine, A-1609
 Azomycin, N-250
 4-Azoniaspiro[3.3]heptane-2,6-diol, A-1610
 Azotobactin D, A-1611
 Azotobactin G173, A-1612
 Azotobactin P19, A-1613
 Azoverdin A, A-1614
 Azoverdin G, A-1615
 Azoverdin, A-1614
 Azoxybacilin, A-1616
 Aztequine, A-1617
 Azx, M-396
 B 4317, A-1122
 B 90063, A-1123
 B 4607A, H-514
 B 41A₃, M-607
 B 41A₄, M-608
 B 4607B, H-514
 B 41B₂, M-607
 B 41B₃, M-608
 B 79B, S-384
 B 4607C, H-514
 B 41C₁, M-607
 B 41D, M-609
 B 79D, T-658
 B 1625FA_{2p-1}, A-1124
 B 41F, M-609
 B 41G, M-609
 B 371, I-216
 B 14437, S-53
 B 870, T-390
 BA 180265A, A-244
 BA 181314A, A-1125
 Ba 29837, D-142
 Ba 33112, D-142
 BA 90912, S-53
 4181B, A-1121
 460B, A-1309
 1401B, A-1120
 0231B, A-1088
 Bacillamide A, B-1
 Bacillamide B, B-1
 Bacillamide C, B-1
 Bacillibactin, B-2
 Bacimethrin, A-795
 Backebergine, I-302
 Bactropone, B-72
 Badianic acid, M-240
 Baecocystine, H-480
 Bagougeramine A, B-3
 Bagougeramine B, B-4
 Bagren, E-157
 Bahamamide, B-5
 Bahienoside A, B-6
 Bahienoside B, B-6
 Baicalidine, B-7
 Baicaline, B-7
 Baikaidine, H-11
 Baikaine, H-11
 Baikian betaine, T-221
 Baikian, T-221
 Baimonidine, C-298
 Baiyumine A, A-120
 Baiyumine B, P-616
 Bakankoside, B-8
 Bakankosine, B-8
 Balanoglossol, T-466
 Balanol, B-9
 Balasubramide, B-10
 Baldrianic acid, M-409
 Baleabuxaline I, B-11
 Baleabuxidienine I, B-466
 Baleabuxidine F, B-12
 Baleabuxidine I, B-12
 Baleabuxidine, B-12
 Baleabuxine, C-920
 Baleabuxoxazine C, C-851
 Balfourine, M-366
 Balfouridine, B-13
 Balfouronium, B-14
 Balfourone, L-288
 Baloxine, M-631
 Balsamide, M-18
 Balsoxine, D-650
 Baluchistanamine, S-192
 Baluchistine, O-1
 B057-250A, M-588
 Banisterine, H-592
 Banlangenjasu, R-11
 Bannucine, B-15
 Banyascyclamide A, D-381
 Banyascyclamide B, B-16
 Banyascyclamide C, B-17
 Banyaside A, B-18
 Banyaside B, B-19
 Baogongteng A, A-1582
 Baogongteng C, A-1582
 BAP, A-137
 Baptifoline, B-20
 Baptitoxine, C-938
 Barbaleucamide A, B-21
 Barbaleucamide B, B-21
 Barbamide, B-22
 Barbarin, P-346
 Barbatin C, T-535
 Barbeline, M-358
 Barbinidine, M-321

- Barbinine, G-91
 Barbitol, B-23
 Barbitone, B-23
 Baretin bromobenzisoxazolone, B-25
 Baretin, B-24
 Bargustanine, B-26
 Barrenazine A, B-27
 Barrenazine B, B-27
 Barterin, D-565
 Barterine, S-595
Fritillaria Base 5, A-297
Fritillaria Base 6, A-298
Lycopodium Base 168, A-312
Lycopodium Base 258, A-313
Pancreatium Base 31, A-324
Pancreatium Base 6, A-323
Senecio renardi Base 4, A-332
Sophora Base 9, A-333
Sophora alopecuroides Base A₁, A-637
Sophora alopecuroides Base A₂, A-637
Stephania glabra Base A, A-376
Stephania Base A, A-378
 Base A, D-888
 Base B, D-888
Crotalaria goroensis Base C, A-437
Leucojum Base C, A-448
Leucojum Base D, A-484
Narcissus Base D, A-489
Lycopodium Base E, A-501
Sophora Base E, A-505
Lycopodium Base G, A-516
Sternbergia Base III, A-527
Lycopodium Base IV, A-344
Ungernia Base IV, A-529
Lycopodium Base IX, A-531
 Base LV1, M-731
Mucuna pruriens Base P, A-579
Mucuna pruriens Base Q, A-588
Fritillaria Base R, A-591
Lycopodium Base R, A-592
Mucuna pruriens Base R, A-593
Mucuna pruriens Base S, A-598
 Base T, D-114
 Base TR1, V-115
 Base TR2, V-115
Lycopodium Base V, A-310
Pachysandra Base V, A-322
Lycopodium Base VI, A-605
Pachysandra Base VI, A-607
Rhodophiala Base VI, A-608
Lycopodium Base VII, A-606
Lycopodium Base VIII, A-609
Lycopodium Base X, A-311
Mucuna pruriens Base X, A-321
Lycopodium Base XI, A-610
Lycopodium Base XII, A-611
Rhodophiala Base XIII, A-613
Pachysandra Base XVI, A-614
Rhodophiala Base XXV, A-615
Pachysandra Base XXVII, A-616
 Basiliskamide A, D-572
 Basiliskamide B, D-572
 Bassianin, B-28
 Bassiatin, B-29
 Bassinic acid, O-32
 Bastadin 10, B-32
 Bastadin 11, B-32
 Bastadin 12, B-32
 Bastadin 13, B-33
 Bastadin 14, B-32
 Bastadin 15, B-32
 Bastadin 16, B-34
 Bastadin 17, B-34
 Bastadin 18, B-34
 Bastadin 19, B-33
 Bastadin 1, B-30
 Bastadin 20, B-33
 Bastadin 21, B-33
 Bastadin 22, B-32
 Bastadin 23, B-32
 Bastadin 24, B-32
 Bastadin 2, B-30
 Bastadin 3, B-31
 Bastadin 4, B-32
 Bastadin 5, B-32
 Bastadin 6, B-32
 Bastadin 7, B-32
 Bastadin 8, B-32
 Bastadin 9, B-32
 Bastadin 12 (obsol.), B-33
 Bastadin 9 (obsol.), B-32
 Bataconine, M-323
 Batrachamine, B-35
 Batrachotoxin, B-36
 Batrachotoxinin A *cis*-crotonate, B-36
 Batrachotoxinin A *trans*-crotonate, B-36
 Batrachotoxinin A homocrotonate, B-36
 Batrachotoxinin A, B-36
 Batzelladine A, B-37
 Batzelladine B, B-38
 Batzelladine C, B-39
 Batzelladine D, B-40
 Batzelladine E, B-41
 Batzelladine F, B-42
 Batzelladine G, B-43
 Batzelladine H, B-44
 Batzelladine I, B-44
 Batzelladine J, B-45
 Batzelladine K, B-46
 Batzelladine L, B-47
 Batzelladine M, B-48
 Batzelladine N, B-42
 Batzellaside A, B-49
 Batzellaside B, B-50
 Batzellaside C, B-51
 Batzelline A, B-52
 Batzelline B, B-52
 Batzelline C, B-52
 Batzelline D, B-52
 Bauerine A, C-399
 Bauerine B, D-355
 Bauerine C, D-351
 Bauhinin, L-213
 Baytopine, P-221
 BB 3, A-55
 BB 4, A-54
 BBM 2040A, C-373
 BBM 928A, L-312
 BBM 2040B, C-374
 BBM 928B, L-312
 BBM 928C, L-312
 B057-250B, M-588
 BC 41C₂, M-608
 B057-250C, M-588
 5-*epi-cis*-275B', D-109
 5-*epi-trans*-275B, D-109
 B057-250D, M-588
 BE 10988, A-1126
 BE 14106, A-1127
 BE 22179, A-1128
 BE 48021, A-1129
 BE 52211, A-1130
 BE 54017, A-1131
 BE 54476, A-1132
 BE 55051, A-1133
 BE 67251, A-1134
 BE 19412A, A-1135
 BE 54238A, A-1136
 BE 45722A, D-126
 BE 19412B, A-1135
 BE 39907B, A-853
 BE 52211B, A-1137
 BE 54238B, A-1136
 BE 45722B, D-126
 BE 13793C, A-1393
 BE 52211C, A-1137
 BE 45722C, D-126
 BE 12233, H-747
 BE 18591, T-22
 Bearline, G-91
 Bebeerine, C-812
 Bebuxine, C-917
 Becatecarin, R-32
 Beccapoline, B-53
 Beccapolinium, B-53
 Beccapolydione, B-53
 Bechuanine, P-221
 Becilan, P-923
 Beecheyamide, M-280
 Begonanline, H-458
 Behenic acid tryptamide, T-639
 N^b-Behenoyltryptamine, T-639
 Beilupeimine, B-54
 Beiwucine, M-302
 Beiwudine, F-147
 Beiwusine A, A-1533
 Beiwusine B, A-1533
 Beiwutine, M-302
 Belamarine, F-7
 Belarine, I-329
 Belemine, B-55
 Belladine, N-288
 α-Belladonnine, B-56
 Belladonnine, B-56
 β-Belladonnine, B-56
 Bellamarine, F-7
 Bellaradine, C-818
 Bellendine, B-57
 B057-250E, M-588
 Bendopa, A-745
 Benethenamine, P-342
 Bengacarboline, B-58
 Bengamide A, B-60
 Bengamide B, B-60
 Bengamide C, B-60
 Bengamide D, B-60
 Bengamide E, B-60
 Bengamide F, B-60
 Bengamide G, B-60
 Bengamide H, B-60
 Bengamide I, B-60
 Bengamide J, B-60
 Bengamide K, B-59
 Bengamide L, B-60
 Bengamide M, B-60
 Bengamide N, B-60
 Bengamide O, B-60
 Bengamide P, B-60
 Bengamide Q, B-60
 Bengamide R, B-60
 Bengamide Y, B-60
 Bengamide Z, B-60
 Bengazole A, D-95
 Bengazole B, D-95
 Bengazole B₁, D-95
 Bengazole C₂, B-61
 Bengazole C₃, B-61
 Bengazole C₄, B-61
 Bengazole C₆, B-61
 Bengazole C, D-95
 Bengazole D₂, B-61
 Bengazole D₃, B-61
 Bengazole D₄, B-61
 Bengazole D, D-95
 Bengazole E, D-95
 Bengazole F, D-95
 Bengazole G, D-95
 Bengazole Z, B-61
 Benhamycin, B-62
 Beninine, B-63
 Benthocyanin A, B-64
 Benthocyanin B, B-64
 Benthocyanin C, B-65
 Benthophoenin, B-66
 Benzaldehyde cyanohydrin, H-673
 2-Benzamidobenzoic acid, B-67
 6-(2-Benzamidoethyl)-2,2-dimethyl-2H-1-benzopyran, A-770
 2-Benzamido-4-hydroxybenzoic acid, B-67
 2-Benzamido-4-methoxybenzoic acid, B-67
 Benzastatin A, B-68
 Benzastatin B, B-68
 Benzastatin C, V-145
 Benzastatin D, V-145
 Benzastatin E, B-69
 Benzastatin F, B-69
 Benzastatin G, B-70
 Benzastatin H, B-68
 Benzastatin I, B-68

- Benzathine tenuazonate, T-74
 2-Benzazine, I-301
 1-Benzazine, Q-29
 1-Benzazole, I-79
 Benzeneacetamide, P-336
 Benzeneacetic acid, P-336
 Benzeneacetonitrile, P-336
 Benzeneethanamine, P-342
 Benzenemethanamine, B-84
 Benzenepropanenitrile, P-350
 Benzenepropanoic acid, P-350
 Benz[*g*]isoquinoline-5,10-dione, B-71
 1,2-Benzisoxazole-3,6-diol, B-72
 Benzof[*l*]-1,3-benzodioxol[6,5,4-*cd*]indol-5(6*H*)-one, D-567
 8*H*-Benzof[*g*]-1,3-benzodioxol[6,5,4-*de*]quinolin-8-one, L-194
 3,4-Benzo- γ -carboline, I-110
 Benzoc[*cinnoline*-1,10-diol, B-73
 1,3-Benzodioxole-5-carboxylic acid, M-442
 1,3-Benzodioxole-5-nonanoic acid, M-454
 [1,3]Benzodioxol[5,6-*c*]-1,3-dioxol[4,5-*l*]phenanthridine, S-56
 [1,3]Benzodioxol[5,6-*h*]quinolin-3-ol, H-599
 3-(1,3-Benzodioxol-5-yl)-*N*-[2-(1,3-benzodioxol-5-yl)ethyl]-2-propenamide, D-799
 3-(1,3-Benzodioxol-5-yl)-*N*-[2-(3,4-dimethoxyphenyl)ethyl]-2-propenamide, Z-6
 2-[2-(1,3-Benzodioxol-5-yl)ethenyl]-4-methoxyquinoline, M-265
N-[2-(1,3-Benzodioxol-5-yl)ethyl]cyclopropanecarboxamide, R-142
 2-[2-(1,3-Benzodioxol-5-yl)ethyl]-4-methoxyquinoline, C-822
 2-[2-(1,3-Benzodioxol-5-yl)ethyl]quinoline, D-645
 1-[2-(1,3-Benzodioxol-5-yl)ethyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, D-978
 2-[2-(1,3-Benzodioxol-5-yl)ethyl]-1,2,3,4-tetrahydro-1-methylquinoline, G-9
 2-(1,3-Benzodioxol-5-yl)-4-methoxyquinoline, M-264
 13-(1,3-Benzodioxol-5-yl)-*N*-(3-methylbutyl)-2,4,12-tridecatrienamide, G-224
 5-(1,3-Benzodioxol-5-ylmethyl)-1,3-dioxol[4,5-*g*]isoquinoline, M-672
 1-(1,3-Benzodioxol-5-ylmethyl)-6-methoxy-7-isoquinolinol, S-274
 4-(1,3-Benzodioxol-5-ylmethyl)-1-methyl-1*H*-imidazol-2-amine, P-594
 7-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4-heptadienamide, C-387
 7-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,6-heptadienamide, F-232
 16-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4,15-hexadecatrienamide, B-266
 15-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4-pentadecadienamide, B-265
 15-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4,13-pentadecatrienamide, B-265
 15-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4,12-pentadecatrienamide, B-265
 15-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4,14-pentadecatrienamide, B-265
 5-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4-pentadienamide, P-462
 5-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2-pentenamide, P-462
 5-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-4-pentenamide, P-462
 3-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2-propenamide, M-457
 13-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4,12-tridecatrienamide, G-224
 11-(1,3-Benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,8,10-undecatrienamide, P-469
 2-(1,3-Benzodioxol-5-yl)-1-methyl-4(1*H*)-quinolinone, G-171
 5-(1,3-Benzodioxol-5-yl)-*N*-(10-methyl-5-undecenyl)-2,4-pentadienamide, S-653
 2-(1,3-Benzodioxol-5-yl)-1,2,4*a*,5,6,7,8,8*a*-octahydro-*N*-(2-methylpropyl)-1-naphthalenecarboxamide, C-896
 1-[[2-(1,3-Benzodioxol-5-yl)-1,2,4*a*,5,6,7,8,8*a*-octahydro-1-naphthalenyl]carbonyl]pyrrolidine, C-909
 3-[5-(1,3-Benzodioxol-5-yl)-2-oxazolyl]pyridine, T-299
 1-[10-(1,3-Benzodioxol-5-yl)-1-oxo-9-decenyl]pyrrolidine, I-266
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4-heptadienyl]piperidine, P-452
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,6-heptadienyl]pyrrolidine, M-452
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,6-heptatrienyl]piperidine, P-452
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,6-heptatrienyl]pyrrolidine, P-453
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-6-heptenyl]piperidine, P-465
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-2-heptenyl]piperidine, P-465
 1-[7-(1,3-Benzodioxol-5-yl)-1-oxo-6-heptenyl]pyrrolidine, M-452
 1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,8-nonadienyl]piperidine, P-464
 1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,8-nonadienyl]pyrrolidine, M-455
 1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,8-nonatrienyl]pyrrolidine, M-455
 1-[9-(1,3-Benzodioxol-5-yl)-1-oxo-8-nonenyl]pyrrolidine, M-455
 1-[8-(1,3-Benzodioxol-5-yl)-1-oxo-7-octadecenyl]piperidine, P-470
 1-[5-(1,3-Benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine, P-467
 1-[5-(1,3-Benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]pyrrolidine, T-495
 1-[5-(1,3-Benzodioxol-5-yl)-1-oxo-2-pentenyl]piperidine, P-467
 1-[3-(1,3-Benzodioxol-5-yl)-1-oxo-2-propenyl]piperidine, M-444
 1-[13-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,12-tridecatrienyl]piperidine, P-448
 1-[13-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,12-tridecatrienyl]pyrrolidine, B-263
 1-[11-(1,3-Benzodioxol-5-yl)-1-oxo-2,4,10-undecatrienyl]piperidine, P-471
 5-(1,3-Benzodioxol-5-yl)-2-phenyloxazole, D-650
 3-(1,3-Benzodioxol-5-yl)-2-propenoic acid, M-457
 2-(1,3-Benzodioxol-5-yl)quinoline, M-458
 1-(1,3-Benzodioxol-5-yl)-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, C-796
 6,7-Benzolumazine, A-632
 Benzomalvin A, B-74
 Benzomalvin B, B-74
 Benzomalvin C, B-74
 Benzomalvin D, B-74
 Benzoc[*c*][2,7]naphthyridin-4-ol, B-75
 Benzoc[*c*][2,7]naphthyridin-4(3*H*)-one, B-75
 Benzof[*g*]pteridine-2,4(3*H*,10*H*)-dione, A-632
 Benzof[*g*]pteridine-2,4(1*H*,3*H*)-dione, A-632
 Benzopyridine, I-301
 2,3-Benzopyrrole, I-79
 4-(3*H*-Benzof[*a*]pyrrolo[2,3-*c*]phenazin-1-yl)phenol, H-674
 Benzosimuline, B-76
 Benzothiazole, B-77
 2,6-Benzothiazole-diol, B-78
 2-Benzothiazolethiol, B-79
 2(3*H*)-Benzothiazolethione, B-79
 2-Benzothiazolinethione, B-79
 2-Benzothiazolinone, B-80
 2-Benzothiazolol, B-80
 2(3*H*)-Benzothiazolone, B-80
 2,4-Benzoxazole-diol, B-81
 2,6-Benzoxazole-diol, B-82
 2-Benzoxazolinone, B-83
 2-Benzoxazolol, B-83
 2(3*H*)-Benzoxazolone, B-83
N-Benzoyl-4-(4-acetoxyneroxy)phenethylamine, A-77
N-Benzoyl-*O*-acetylbutoxidienine E, B-456
N-Benzoyl-16-acetylcylohexobuxidine, B-12
N-Benzoyl-*O*-acetylcylohexobuxoline F, C-921
 14-Benzoylaconine, M-331
 10-Benzoyl-9(10*H*)-acridinone, A-112
 2-(Benzoylamino)acetophenone, A-694
 2-(Benzoylamino)benzoic acid, B-67
 1-[4-(Benzoylamino)butyl]hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1*H*)-one, D-927
 2-(Benzoylamino)ethylphosphonic acid, A-773
 2-(Benzoylamino)-1-phenylethanol, A-864
N-Benzoylanthranilic acid, B-67
N-Benzoylbaleabuxidienine F, B-466
 Benzoylbaleabuxidine F, B-12
 14-Benzoylbrownine, M-309
 8-Benzoyl-2-[3-(1-butanyl)oxiranyl]-9-hydroxy-8-methoxy-3-methyl-1-oxa-7-azaspiro[4.4]non-2-ene-4,6-dione, S-660
N-Benzoylbuxahyrcanine, B-451
N-Benzoylbuxidienine F, B-466
N-Benzoylbuxidienine, B-466
N-Benzoylbuxocyclamine F, B-474
N-Benzoylbuxodienine E, B-456
N-Benzoyl-*N'*-cinnamoylputrescine, C-456
N-Benzoylcyclobuxophylline M, C-848
N-Benzoylcyclocimicrophylline C, C-883
 3-Benzoylcyclocimicrophylline F, C-883
N-Benzoylcycloprotobuxine C, C-902
N-Benzoylcycloprotobuxoline C, C-903
N-Benzoylcycloprotobuxoline D, C-903
*N*³-Benzoylcyclohexobuxidine F, B-12
*N*³-Benzoylcyclohexobuxine F, C-920
N-Benzoylcyclohexobuxoline F, C-921
*N*¹-Benzoylcylindrocarpine, C-929
 18-Benzoyldavisinol, H-168
 5-Benzoyl-5-deacetylwilforidine, P-253
 14-Benzoyldelcosine, M-309
 14-*O*-Benzoyldelphinone, M-312
 Benzoyldeoxyaconine, M-343
O^{2b}-Benzoyl-16-deoxybuxidienine C, B-466
 14-Benzoyldictyocarpine, M-307
 2-Benzoyl-1,2-dihydro-6-acetylpyridine, S-212
*N*³-Benzoyldihydrocyclocimicrophylline F, C-883
 14-Benzoyldihydrogadesine, M-321
 1-(6-Benzoyl-1,6-dihydro-2-pyridinyl)-2-propa-none, S-212
N-(7-Benzoyl-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl)-2-(dimethylamino)-3-methylpentanamide, A-1375
 Benzoyllegonine methyl ester, C-537
 Benzoyllegonine, E-8
 Benzoylneurea, Q-13
 14-Benzoylepiscopalisinine, A-97
 14-Benzoyl-8-*O*-ethylaconine, M-331
 α -Benzoylethylamine, A-870
 14-*O*-Benzoyl-8-*O*-ethylbikhaconine, M-312
 14-Benzoylgadesine, G-152
 6-Benzoylheteratisine, H-157
N-Benzoyl-4-hydroxyanthranilic acid, B-67
N-Benzoyl-20-hydroxycylindrocarpine, C-929
 11-*O*-Benzoyl-4-hydroxyepilupinine, H-482
N-Benzoyl-2-hydroxy-2-(4-methoxyphenyl)ethylamine, T-67
N-Benzoyl-2-hydroxy-2-phenylethylamine, A-864
 Benzoylhpaconine, M-343
O-Benzoyl-6-isocassine, C-183
 1-Benzoylkarasamine, M-355
 Benzoyllaburnine, H-629
 14-Benzoylmesaconine, M-331
N-Benzoylmesaline, M-219
N-Benzoyl-4-methoxyanthranilic acid, B-67
 14-Benzoyl-8-*O*-methylaconine, M-331
 Benzoylmethylamine, A-694
 14-*O*-Benzoyl-8-*O*-methylbikhaconine, M-312
 14-Benzoylneoline, M-323
 14-Benzoylnudicaulidine, M-321
 5'-*O*-Benzoyloxyanthin, H-673
 3-Benzoyloxy-6-hydroxy-7-(2-hydroxy-3-phenylpropanoyloxy)nortropine, A-1596
 2-Benzoyloxy-3-hydroxynortropine, A-1581
 6-Benzoyloxy-3-isobutanoyloxynortropine, A-1584
 13-Benzoyloxylupanine, H-554
 3 α -Benzoyloxynortropine, A-1597
 3-Benzoyloxynortropine-6-ol, A-1584
 13-Benzoyloxy-17-oxolupanine, O-188

- 7-Benzoyloxy-3-(3,4,5-trimethoxycinnamoyloxy)-tropan-6-ol, M-392
 7-Benzoyloxytropane-3,6-diol, M-392
 3-Benzoyloxytropan-6-ol, M-391
 6-Benzoyloxy-3-tropanone, M-391
 14-*O*-Benzoylperegrine, M-347
N-Benzoyl-*N*-phenylaminomethylcarbinol, H-485
 15-Benzoylpseudokobusine, H-205
 14-Benzoylsachaconitine, M-355
O-Benzoylsaphenic acid, H-496
*N*⁵-Benzoylspermidine, S-398
N-Benzoylstaurosporine, S-498
 14-*O*-Benzoyltaecosamine, M-309
 14-Benzoyltalatizamine, M-352
 3-*O*-Benzoylteloidine, M-392
O-Benzoyltombozine, T-425
 2 α -Benzoyltropane, F-46
 Benzoyltropein, M-393
N-Benzoyltyramine methyl ether, M-280
N-Benzoyltyramine, H-667
O-Benzoylvincamajine, V-113
 Benztryptamide, T-639
 Benzyl cyanide, P-336
 Benzyl nitrile, P-336
 Benzylacetic acid, P-350
 2-Benzyl-*O*-acetyl tropanol, B-96
N-Benzyladenine, A-137
 Benzylamine, B-84
 6-(Benzylamino)-2-(2-hydroxyethylamino)-9-methylpurine, O-90
 6-Benzylaminopurine, A-137
N-Benzylbenzamide, B-84
N-Benzylbenzeneacetamide, P-336
 2-Benzyl-3-*O*-benzoyltropanol, B-96
 5-Benzyl-3-benzylidene-1-hydroxy-2,6(1*H*,3*H*)-pyrazinedione, S-152
N-Benzylbiotin, B-139
 Benzylcarbamide, B-97
 2-Benzyl-5-carbamoyl-4(1*H*)-pyridinone, A-1483
 3-Benzyl-1,4-diazabicyclo[4.4.0]decan-5-one, V-84
 5-Benzyl-2,5-dihydro-4-hydroxy-3-(2,4,6-octatrienoyl)-1*H*-pyrrol-2-one, B-85
 6-Benzyl-1,2-dihydro-4-hydroxy-2-oxo-3-pyridine-carboxaldehyde, C-142
 3-Benzyl-1,2-dihydro-1-hydroxy-4-pyridinecarboxaldehyde, M-2
 3-Benzyl-3,4-dihydro-4-methyl-1*H*-1,4-benzodiazepine-2,5-dione, C-890
 2-Benzyl-2,3-dihydro-1-methyl-4(1*H*)-quinazolinone, G-127
 4-Benzyl-1,4-dihydro-3-oxo-3*H*-pyrrolo[2,1-*c*][1,4]oxazine-6-carboxaldehyde, B-86
 6-Benzyl-2,4-dihydroxy-3-pyridinecarboxaldehyde, C-142
 2-Benzyl-3,6-dihydroxytropane, B-94
 2-Benzyl-3,7-dihydroxytropane, B-95
 5-Benzyl-3-(2,4,6,8,10-dodecapentaenoyl)-2,5-dihydro-4-hydroxy-1*H*-pyrrol-2-one, B-87
 3-Benzyl-4-formyl-1,2-dihydro-1-hydroxypyridine, M-2
 α -Benzyl-2-formyl-5-hydroxymethyl-1*H*-pyrrole-1-acetic acid, F-138
 6-Benzyl-3-formyl-4-hydroxy-2-pyridone, C-142
N-Benzylhexadecanamide, H-221
 3-Benzylhexahydro-7-hydroxypyrrolo[1,2-*a*]pyrazine-1,4-dione, C-873
 3-Benzylhexahydro-7-hydroxypyrrolo[1,2-*a*]pyrazine-1,4-dione, C-891
 5-Benzyl-1-hydroxy-3-(hydroxybenzylidene)-2,6(1*H*,3*H*)-pyrazinedione, S-152
 5-Benzyl-3-hydroxy-2-isopropylpyrazine, B-88
 2-Benzyl-3-hydroxy-1-methyl-5-nonylpyrrolidine, B-89
 2-Benzyl-3-hydroxypyrazine, B-90
 2-Benzyl-4-hydroxyquinazoline, B-93
 2-Benzyl-3-hydroxytropane, B-96
 3-Benzylidene-3,4-dihydro-2-oxo-2*H*-1,4-benzoxazine-5-carboxylic acid, A-1273
 3-Benzylidene-6-(hydroxymethyl)-1,4-dimethyl-6-(methylthio)-2,5-piperazinedione, B-91
 2-Benzyl-5-(3-indolyl)oxazole, A-622
N-Benzylisopropylamine, B-84
 3-Benzyl-6-isopropyl-4-methyl-2,5-morpholine-dione, B-29
 6-Benzyl-3-isopropyl-2-pyrazinone, B-88
 6-Benzyl-3-isopropyl-2(1*H*)-pyrazinone, B-88
N-Benzyl-*N'*-(4-methoxybenzyl)urea, D-302
 2-Benzyl-3-methoxypyrazine, B-90
 2-Benzyl-4-methoxyquinazoline, B-93
 2-Benzyl-8-methyl-8-azabicyclo[3.2.1]octan-3-ol, B-96
 Benzylmethylnitrosamine, B-84
N-Benzyl-*N*-methyl-1-pentadecylamine, P-203
 2-Benzyl-1-methyl-4(1*H*)-quinazolinone, B-93
N-Benzyl-13-oxo-9,11-octadecadienamide, O-191
N-Benzyl-5-oxo-6,8-octadecadienamide, O-189
N-Benzyl-9-oxo-12,15-octadecadienamide, O-190
N-Benzyl-9-oxo-12-octadecadienamide, O-193
 6-Benzyl-4-oxo-4*H*-pyran-3-carboxamide, B-92
 6-Benzyl-4-oxo-4*H*-pyran-3-carboxylic acid, B-92
 4-Benzyl-1,5-dihydro-2*H*-pyrrol-2-one, P-954
 6-Benzylxypurine, P-820
N-Benzyl-2-phenylacetamide, P-336
 Benzyl-2-phenylethylamine, P-342
N-Benzylphthalimide, P-383
 3-Benzylpyrazinone, B-90
 3-Benzyl-2(1*H*)-pyrazinone, B-90
 3-Benzyl-2,4(1*H*,3*H*)-quinazolinone, Q-13
 2-Benzyl-4(3*H*)-quinazolinone, B-93
N-Benzyl-15-tetracosenamide, T-120
 4-Benzyl-3,6-tropanediol, B-95
 2-Benzyl-3,6-tropanediol, B-94
 2-Benzyl-3-tropanol, B-96
 Benzylurea, B-97
 18-*O*-Benzylwilfortrine, P-253
 Berbacolorflamme, C-171
 Berbamine 2'- β -*N*-oxide, I-328
 Berbamine, I-328
 Berbamunine, B-98
 Berbenine, I-328
 Berberal, B-101
 Berberal, B-99
 Berberastine, T-321
 Berbericine, B-101
 Berbericine, P-34
 Berberilycine, B-100
 Berberine chloroform, P-35
 Berberine, B-101
 Berberinol, O-216
 Berberrubine, B-101
 Berbervirine, C-72
 Berbibuxine, C-23
 Berbidine, B-102
 Berbilaurine, L-67
 Berbithine, B-103
 Berbivaldine, P-15
 Berisol, B-101
 Berlamine, O-216
 Berninamycin A, B-104
 Berninamycin B, B-104
 Bernumicine, B-105
 Bernumidine, B-106
 Bernumine, B-106
 Berpodine, B-107
 Bervulcine, B-108
 Betalamic acid, B-109
 Betalin S, T-366
 Betanidin 6-(2^Q-glucosylrutinoside), B-110
 Betanidin rhamnosophoroside, B-110
 Betanidin 5-sophoroside, B-110
 Betanidin 6-sophoroside, B-110
 Betanidin, B-110
 Betanin, B-110
 Betaxin, T-366
 Betonicine, H-476
 Bewon, T-366
 Bgugaine, T-135
 2088B, H-514
 Bharatamine, B-111
 Bharhingine, B-112
 Bhesine, T-139
 Bhimamycin C, B-113
 Bhimamycin D, B-114
 Bhimberine, I-319
 10,10'-Bi[*N*-acetyl-11,12-dihydroxyaspidospermidine], B-115
 Bianfugecine, B-116
 Bianfugedine, B-117
 Bianfugenine, D-90
 Bicarbamie acid, H-395
N,N'-Bi- β -carboline, B-146
 Bicoloridine alcohol, M-347
 Bicoloridine, M-347
 Bicolorine 14-*O*-acetate, M-347
 Bicolorine 6-*O*-acetate, M-347
 Bicolorine, M-347
 Bicolorine, M-448
 Bicucine, B-118
 Bicuculline, B-119
 Bicucullinidine, B-120
 Bicucullinine, A-143
 Bicyclomahanimbicine, B-121
 Bicyclomahanimbine, B-122
 Bidebiline A, U-36
 Bidebiline B, U-36
 Bidebiline C, U-36
 Bidebiline D, U-36
 Bidebiline E, U-36
 3,3'-Bi(2,5-dibromoindole), T-108
 3,3'-Bi(2,6-dibromoindole), T-109
 3,3'-Bi[4,6-dibromo-2-methylthio-1*H*-indole], T-110
 Biemnadin, B-123
 Biemnidin, D-929
 Bietaserpine, R-52
 Biflorine, B-124
 Biflorine, P-667
 Biflorone, B-125
 4,4''-Bi[*N*-4-hydroxycinnamoylserotonin], B-126
 12,12''-Bi[11-hydroxycoronaridine], B-127
 2,2''-Bi(1-hydroxy-6,8-dimethoxy-3-methylcarbazole), C-503
 4,4''-Bi[*N*-(4-hydroxy-3-methoxycinnamoyl)serotonin], B-126
 1,1'-Bi[2-hydroxy-3-methylcarbazole], B-128
 Bi[5-hydroxyxonoracronyline], B-129
 2,2''-Bi[3-hydroxy-4(1*H*)-pyridinone], T-249
 3,3'-Bi-1*H*-indole-5,5',6,6'-tetrol, B-130
 [3,3'-Biindoline]-2,2'-dione, B-131
 $\Delta^{2,2'}$ -Biindoline-3,3'-dione, I-72
 $\Delta^{2,3'}$ -Biindoline-2',3'-dione, I-73
 8',11'*O*-Biisocorydine, B-132
 8,8''-Biisocorydine, B-133
 Bikhacconine, M-312
 Bikhacconitine, M-312
 8,8''-Bikoeningine, B-134
 Bikoeninginone A, B-135
 Bilain A, B-136
 Bilain C, B-136
 Bilineurine, C-419
Cypridina Biluciferol, B-137
Cypridina Biluciferyl, B-137
 Bimanol, D-730
 2,3''-Bi(1-methylpyrrolidine), B-147
 2,2''-Bimorphine, M-703
 4,4''-Binecatorone, N-85
 Biochanin C, B-138
 Biocolina, C-419
 Bionectin A, G-99
 Bionectin B, G-99
 Bionectin C, G-97
 Biopten, T-142
 Biopterin glucose, A-749
 7-Biopterin, A-924
 Biopterin, A-749
 Biotin sulfoxide, B-139
 Biotinamide, B-139
 γ -Biotin, B-139
 Biotin, B-139
 Bioxalomycin α 1, B-140
 Bioxalomycin α 2, B-140
 Bioxalomycin β 1, B-140
 Bioxalomycin β 2, B-140
 1,1'-Bipiperidine, B-141
 Bipowine, B-142
 Bipowinone, B-143
 $\Delta^{2,2'}$ -Bipseudoindoxyl, I-72

- Bipy, B-144
 2,2'-Bipyridine, B-144
 2,3'-Bipyridine, B-145
 [2,2'-Bipyridine]-3,3',4,4'-tetrol, T-249
 9,9'-Bi-9*H*-pyrido[3,4-*b*]indole, B-146
 α,α'-Bipyridyl, B-144
 2,2'-Bipyridyl, B-144
 2,3'-Bipyrrolidine, B-147
 Biquidone, B-71
 Birnbaum A, B-148
 Birnbaum B, B-148
N,1-Bis[3-(acetylhydroxyamino)propyl]-3-hydroxy-2,5-dioxo-3-pyrrolidoneacetamide, S-134
N,N'-Bis(agmatine)oxamide, B-178
 Bisaknadimine, H-70
 Bis(4-aminobutyl)bis(3-aminopropyl)ammonium(1+), B-149
N,N'-Bis(4-aminobutyl)-1,4-butanediamine, B-150
 2,6-Bis(4-aminobutyl)-3,5-diiminopiperazine, B-35
*N*³,*N*¹⁰-Bis(4-aminobutyl)homopentamine, D-290
*N*³,*N*¹⁵-Bis(4-aminobutyl)homopentamine, D-290
*N*⁵,*N*¹⁰-Bis(4-aminobutyl)homospermine, B-150
 3,3'-Bis(aminocarbonyl)-5,5'-dicarboxydiphenyl ether, D-815
 Bis[3-(2-aminoethyl)-6-bromo-1*H*-indol-2-yl] disulfide, B-163
 3,3'-Bis(2-aminoethyl)-5,5'-dibromo-4,4'-bi-1*H*-indole-7,7'-diol, D-206
 4,10-Bis(2-aminoethyl)-1,7-dimethoxydibenzo[*c,g*][1,2,5,6]-tetrathioicene-2,8-diol, L-208
 1,28-Bis[(3-amino-2-hydroxypropyl)amino]-3,6,9,12,15,18,25-octacosaeptaene, R-71
N,N'-Bis[4-[(aminoiminomethyl)amino]butyl]ethanediamine, B-178
 1,3-Bis[(4-aminomethyl)phenyl]-2-octadecanoylglycerol, B-151
 Bis(3-aminopropyl)amine, D-255
N,N'-Bis(3-aminopropyl)-1,4-butanediamine, B-152
N,N'-Bis(3-aminopropyl)-1,4-butanediamine, S-399
N,N'-Bis(3-aminopropyl)-1,2-ethanediamine, B-153
*N*³,*N*⁵-Bis(3-aminopropyl)homospermidine, B-149
N,N'-Bis(3-aminopropyl)methylamine, D-255
*N*³,*N*⁴-Bis(3-aminopropyl)nospermidine, T-622
N,N'-Bis(3-aminopropyl)-1,3-propanediamine, B-154
N,N'-Bis(3-aminopropyl)-1,3-propanediamine, T-622
*N*⁴,*N*⁴-Bis(3-aminopropyl)spermidine, A-926
 Bis(3-aminopropyl)tetramethylenediamine, S-399
 Bisancorinolate B, B-155
 Bisaprasin 11'-sulfate, P-686
 Bisaprasin, P-686
 Bisaristone A, B-156
 Bisavenanthramide B₁, B-157
 Bisavenanthramide B₂, B-158
 Bisavenanthramide B₃, B-160
 Bisavenanthramide B₄, B-160
 Bisavenanthramide B₅, B-159
 Bisavenanthramide B₆, B-160
 8*H*-Bis[1,3-benzodioxolo[6,5,4-*de*:4',5'-*g*]quinolin-8-one, H-150
 4,5-Bis(1,3-benzodioxol-5-ylmethyl)-1-methyl-1*H*-imidazol-2-amine, L-150
N,N'-Bis(2,10-bisaboladien-7-yl)urea, B-161
 3,3'-Bis(5-bromo-4-chloro-7-methoxy-1*H*-indole), D-309
N,N'-Bis[3-(3-bromo-4-hydroxyphenyl)-2-oximido-propanoyl]cystamine, P-686
 2,2-Bis(6-bromo-3-indolyl)ethylamine, B-162
 2,4-Bis(6-bromo-3-indolyl)imidazole, N-312
 2,5-Bis(6-bromo-3-indolyl)-1-methylpiperazine, D-929
 2,5-Bis(6-bromo-3-indolyl)piperazine, D-929
 Bis(6-bromo-2-tryptaminy) disulfide, B-163
N,N'-Bis[3-(carboxyamino)-4-methylphenyl]urea, B-164
*N*¹,*N*⁴-Bis(2-carboxyethyl)-1,4-butanediamine, S-397
 Biscephalezomine A, B-165
 Biscephalezomine B, B-166
 Biscephalezomine C, B-167
 Biscephalezomine D, B-165
 Biscephalezomine E, B-166
*N*²,*N*²-Bis(chloromethyl)tetrandrinium, T-286
 3,6-Bis(5-chloro-2-piperidinyl)-2,5-piperazinedione, A-1089
 4,4'-Bis(*N*-*p*-coumaroylserotonin), B-126
 Biscryptolepine, B-168
 2',7'-Bisdeacetoxyaustrospicatin, T-49
 7,7'-Bis(dehydroanonaime), U-36
 7,7'-Bisdehydro-*O*-methylisopiline, U-41
 Bisdehydroneostemonine, N-140
 Bisdehydroneostemoninine, B-169
 Bisdehydroneotuberostemonine, T-664
 Bisdehydroprotostemonine, P-670
 Bisdehydrostemochochinine, P-670
 Bisdehydrostemonine, S-519
 Bisdehydrostemoninine A, B-170
 Bisdehydrostemoninine B, U-170
 Bisdehydrostemoninine, S-521
 Bisdehydrotuberostemonine B, T-664
 Bisdehydrotuberostemonine C, T-664
 Bisdehydrotuberostemonine, T-664
N,O-Bis(demethyl)buchenavianine, D-662
N,N'-Bis(2,3-dibromo-4,5-dihydroxybenzyl)-*N*-(3-methoxycarbonylpropyl)urea, D-312
*N*¹,*N*⁴-Bis(4,4-dicarboxy-3-hydroxy-1-oxobutyl)-1,4-butanediamine, R-80
 7,7'-Bis(6*a*,7-didehydrodrolisnorine), B-142
N,N'-Bis(2,4-diethyl-4-hexenyl)guanidine, T-617
*N*¹,*N*¹⁰-Bis(dihydrocaffeoyl)spermidine, S-398
 1,10-Bis(dihydrocaffeoyl)spermine, S-399
 1,14-Bis(dihydrocaffeoyl)spermine, S-399
 1,3-Bis(13,14-dihydro-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridin-14-yl)-2-propanone, S-54
*N*²,*N*⁹-Bis(2,3-dihydroxybenzoyl)lysine, M-806
N,N'-Bis(2,3-dihydroxybenzoyl)spermidine, S-398
*N*¹,*N*¹⁰-Bis(3,4-dihydroxybenzoyl)spermidine, S-398
N,N'-Bis(3,4-dihydroxycinnamoyl)-1,4-butanediamine, D-362
N,N'-Bis[2-(3,5-diiodo-4-methoxyphenyl)ethyl]urea, B-171
 6,7-Bis(3,4-dimethoxyphenyl)-1,2,3,5,8,8*a*-hexahydroindolizine, S-248
 3,5-Bis(γ,γ-dimethylallyl)indole, B-200
 3,6-Bis(γ,γ-dimethylallyl)indole, B-201
 3,7-Bis(γ,γ-dimethylallyl)indole, B-202
 1,4-Bis(dimethylamino)butane, T-281
 1,4-Bis(dimethylamino)-2-butene, B-423
N,N'-Bis[4-(dimethylamino)butyl]hexanamide, S-340
 4,4'-Bis(dimethylamino)diethylamine, S-340
 3,3'-Bis(dimethylamino)dipropylamine, D-255
 3,3'-Bis(dimethylamino)-*N*-methylpropylamine, D-255
 3,20-Bis(dimethylamino)pregnane, P-601
 3,20-Bis(dimethylamino)pregnan-4-ol, D-266
N,N'-Bis(20-dimethylaminopregnan-3-yl)-*N*-methylthiourea, J-23
 1,5-Bis(1,3-dimethyl-6-lumaziny)-2-methyl-1,5-pentanedione, B-172
 Bis(2,4-dimethyl-6-nitrophenyl)diazene, T-282
N,N'-Bis(3,7-dimethyl-2,6-octadienyl)guanidine, B-173
N,N'-Bis(3,7-dimethyl-2,6-octadienyl)guanidine, B-174
 3,5-Bis[(3,3-dimethylxiranyl)methyl]-1*H*-indole, B-200
 3,6-Bis[(3,3-dimethylxiranyl)methyl]-1*H*-indole, B-201
N,N'-Bis(2,6-dimethylphenyl)-2,2,4,4-tetramethyl-1,3-cyclobutanediimine, B-175
 2,5-Bis[1-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dihydroxy-2,5-cyclohexadiene-1,4-dione, A-1520
 2,5-Bis[2-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dihydroxy-2,5-cyclohexadiene-1,4-dione, H-288
 2,5-Bis[1-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dimethoxy-1,4-benzenediol, A-1519
 Biselide A, H-76
 Biselide B, H-76
 Biselide C, H-76
 Biselide D, H-76
 1,1-Bis(8,9-epoxy-6,8-dimethylergoline), B-176
N,N'-Bis[4-(ethylamino)butyl]-1,4-butanediamine, B-150
 1,2-Bis(ethylamino)ethane, E-225
N,N'-Bis(2-ethylhexanoyl)guanidine, L-174
 4,4''-Bis(*N*-feruloyl)serotonin, B-126
*N*¹,*N*⁸-Bis(glutathionyl)spermidine, T-636
 Bis(gorgiacerol)amine, B-177
N,N'-Bis(4-guanidinobutyl)oxamide, B-178
 Bishaconitine, B-179
 Bishatisine, B-180
 Bishomodeoxyharringtonine, H-66
 Bishomopolytoxin, P-42
 1,14-Bis(homovanillyl)spermine, S-399
 1,3-Bis(11-hydrochelerythriny)acetone, C-355
 1,3-Bis(6-hydrochelerythriny)acetone, C-355
 1,3-Bis(8-hydrochelerythriny)acetone, C-355
 1,3-Bis(11-hydrosanguinariny)acetone, S-54
 1,3-Bis(8-hydrosanguinariny)acetone, S-54
 2,3-Bis(4-hydroxybenzoyl)-5-(4-hydroxyphenyl)pyrazine, B-258
N,N'-Bis(4-hydroxybenzyl)hydroxylamine, B-181
 3,11-Bis-*O*-(3-hydroxybutanoyl)hamayne, H-5
 2,6-Bis(2-hydroxybutyl)-1-methyl-Δ³-piperidine, D-391
 2,6-Bis(2-hydroxybutyl)-1-methylpiperidine, D-391
N,N'-Bis(4-hydroxycinnamoyl)-1,4-butanediamine, D-362
*N*¹,*N*¹⁰-Bis(4-hydroxycinnamoyl)spermidine, S-398
N,N'-Bis(4-hydroxy-3,5-dimethoxycinnamoyl)-1,4-butanediamine, D-362
 Bis(2-hydroxyethyl)amine, D-388
 Bis(7-hydroxygirinimbine A), B-212
 5,5'-Bis(6-hydroxygirinimbine A), K-117
 Bis(7-hydroxygirinimbine B), B-211
 Bis(5-hydroxy-1*H*-indol-3-yl)ethanedione, H-782
 3,4-Bis(5-hydroxy-1*H*-indol-3-yl)-1*H*-pyrrole-2-carboxylic acid, B-182
 3,4-Bis(6-hydroxy-1*H*-indol-3-yl)-1*H*-pyrrole-2,5-dione, A-1394
N,N'-Bis(4-hydroxy-3-methoxycinnamoyl)-1,4-butanediamine, D-362
 3,3'-Bis-3-(4-hydroxy-2-methoxyphenyl)-2-propanamide, B-183
 2,5-Bis(1-hydroxy-3-methylbutyl)-3,6-dimethylpyrazine, B-184
 Bis(*N*-hydroxy-*N*-methylmethanethioamidato-*O,S*)copper, F-97
 Bis(*N*-hydroxy-*N*-methylmethanethioamidato-*O,S*)nickel, F-98
 2,6-Bis(hydroxymethyl)-3,4-piperidinediol, B-185
 2,6-Bis(hydroxymethyl)-3,4,5-piperidinetriol, B-186
 6,8-Bis[10-(5-hydroxy-6-methyl-2-piperidinyl)decyl]-1*H*-indolizinium, J-57
 2,6-Bis(1-hydroxy-2-methylpropyl)-1*H*,5*H*-pyrrolo[2,3-*b*]indole-4,8-dione, T-105
 3-[2,5-Bis(hydroxymethyl)-1*H*-pyrrol-1-yl]-4,5-dimethyl-2(5*H*)-furanone, F-193
N,N'-Bis(2-hydroxy-5-oxo-1-cyclopenten-1-yl)-2,4,6,8,10,12,14-hexadecaheptaenediamide, L-178
 2,6-Bis(2-hydroxypentyl)-1-methylpiperidine, B-187
 2,6-Bis(2-hydroxypentyl)piperidine, B-187
 3,6-Bis(4-hydroxyphenyl)-1,2,4,5-benzenetetrol, T-97
 2,6-Bis(2-hydroxy-2-phenylethyl)-1-methylpiperidine, L-218
 3,3'-Bis[(4-hydroxyphenyl)methylene]-[1,1'-bicyclopent[*b*]indole]-2,2'(3*H*,3'*H*)-dione, S-168
 3,4-Bis(4-hydroxyphenyl)-1*H*-pyrrole-2,5-dicarboxylic acid, B-188
 3,4-Bis(8-hydroxy-4-quinolinyl)-γ-butyrolactone, B-358
 Bisindigotin, B-189
Strepeliopsis Bisindole base, A-334
 7,7-Bis(3-indolyl)-*p*-cresol, D-692
 2,5-Bis(1*H*-indol-3-yl)-3,6-dimethoxy-1,4-benzenediol, A-1519
 2,2-Bis(3-indolyl)indoxyl, T-87

- 3,4-Bis(1*H*-indol-3-yl)maleic anhydride, P-491
 2,3-Bis(1*H*-indol-3-yl)maleimide, A-1394
 3,3'-Bis(indolylmethyl)dimethylammonium, B-190
 2,5-Bis(1*H*-indol-3-ylmethyl)pyrazine, B-191
 2,2'-Bisnomenine, S-313
 1,1'-Bisnomenine, S-313
 8',11*O*-Bisiscocorydine, B-132
 8',11*O*-Bisiscocorydine, B-133
 2,5-Bis[(3-isocyanobutanoyl)amino]-1-pentanol, A-1257
 20,25-Bisisoetioline, S-199
 Bisisomahanine, B-192
 Bisjatrorrhizine, B-193
 8,8'-Biskoenigine, B-134
 Bisline, B-194
 Bismahanine, B-195
 2,4-Bis(5-methoxy-1,3-benzodioxol-6-yl)-1,3-cyclobutanedicarboxylic acid bispyrrolidide, B-196
 1,1'-[[2,4-Bis(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispiperidine, B-198
 1,1'-[[2,4-Bis(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidide, B-196
N,N'-Bis(4-methoxybenzyl)urea, D-302
 Bis(9-methoxy-9*H*-carbazol-3-ylmethyl) ether, O-219
N-[1,3-Bis(methoxycarbonyl)propyl]-2,3-dihydro-2-(4-hydroxy-3-methoxyphenyl)-3-hydroxymethyl-7-methoxy-5-benzofuranpropenamidine, B-197
 Bis(7-methoxygirinimbine A), B-212
 2,4-Bis(2-methoxy-4,5-methylenedioxyphenyl)-1,3-cyclobutanedicarboxylic acid dipiperidide, B-198
 4,5-Bis[(4-methoxyphenyl)methyl]-1*H*-imidazol-2-amine, N-6
 1,4-Bis(methylamino)butane, B-422
N,N-Bis(3-methyl-2-butenyl)guanidine, B-199
N,N'-Bis(3-methyl-2-butenyl)guanidine, P-769
 3,5-Bis(3-methyl-2-butenyl)-1*H*-indole, B-200
 3,6-Bis(3-methyl-2-butenyl)-1*H*-indole, B-201
 3,7-Bis(3-methyl-2-butenyl)-1*H*-indole, B-202
 3,6-Bis[5-(3-methyl-2-butenyl)-1*H*-indol-3-yl]furo[3,2-*b*]furan-2,5-dione, C-548
 3,15-Bis(2-methylbutyl)germine, G-70
 1,2,9,10-Bis(methylenedioxy)aporphine, N-125
 1,2:10,11-Bis(methylenedioxy)aporphine, O-143
 2,3,7,8-Bis(methylenedioxy)benzo[*c*]phenanthridine, S-56
 1,2,9,10-Bismethylenedioxyaporphine, N-125
 1,2:10,11-Bismethylenedioxyaporphine, O-143
 1,2:10,11-Bis(methylenedioxy)oxoaporphine, H-150
 2,3,9,10-Bis(methylenedioxy)-8-oxoprotuberberine, D-409
 2,3,8,9-Bis(methylenedioxy)pavinane, E-221
 2,3,9,10-Bismethylenedioxyprotuberberine, C-638
N-[3,7-Bis(1-methylethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-2-(dimethylamino)-4-methylpentanamide, M-794
 2,5-Bis(1-methylethyl)pyrazine, D-700
 2,6-Bis(1-methylethyl)pyrazine, D-701
 6,6'-[1,2-Bis(2-methyl-1-propenyl)-1,2-ethanediyl]-bis-1*H*-indole, I-248
 2,5-Bis(2-methylpropyl)pyrazine, B-203
 3,6-Bis(2-methylpropyl)-2,5-pyrazinediol 1,4-dioxide, P-788
 3,6-Bis(1-methylpropyl)-2-pyrazinol, B-204
 3,6-Bis(1-methylpropyl)-2(1*H*)-pyrazinone, B-204
 3,6-Bis(2-methylpropyl)-2(1*H*)-pyrazinone, F-76
 1,3-Bis(1-methyl-2-pyrrolidinyl)-2-propanol, D-413
 1,3-Bis(1-methyl-2-pyrrolidinyl)-2-propanone, C-818
 3,5-Bis(1-methyl-2-pyrrolidinyl)pyridine, D-832
N,N'-Bis[6-(methylsulfinyl)hexyl]urea, D-829
 2,6-Bis(methylthio)-1,4-benzoquinone 1-semicarbazone, D-229
 2,2'-Bis(methylthio)-6,6'-dibromoindogotin, T-698
 Bis(*N*-methylthioformohydroxamate)copper, F-97
N,N'-Bis[6-(methylthio)hexyl]urea, D-829
N,N'-Bis[3-(methylthio)-2-propenoyl]putrescine, A-198
 Bismorphine A, B-205
 Bismorphine B, B-206
 Bismurrayafoline A, B-207
 Bismurrayafoline B, B-208
 Bismurrayafoline C, B-209
 Bismurrayafoline D, B-209
 Bismurrayafoline E, B-210
 Bismurrayafolinol, B-207
 6,8'-Bismurrayamine A, B-211
 8,8'-Bismurrayamine A, B-212
 Bismurrayaquinone A, B-213
 Bis(naamidinato A)zinc(II), N-1
 Bis(naamidinato G)zinc(II), N-1
 Bisnor-C-alkaloid D, A-476
 Bisnor-C-alkaloid H di-*N*-oxide, T-436
 Bisnor-C-alkaloid H *N*-oxide, T-436
 Bisnor-C-alkaloid H, T-436
 17,18-Bisnorantidesmone, A-1325
 Bisnorargemone, A-1410
N,N'-Bisnoraromoline, O-1
d-Bisnorbiotin, B-214
 Bisnorbiotin, B-214
 2,2'-Bisnorcoesulinine, C-552
 Bisnor-C-curarine, C-809
 Bisnordihydrotoxiferine di-*N*-oxide, T-436
 Bisnordihydrotoxiferine *N*-oxide, T-436
 Bisnordihydrotoxiferine, T-436
 Bisnorephedrine, A-864
N-Bisnorgramine, I-97
 2,2'-Bisnorquattaguanine, G-212
 Bisnorobamegine, I-328
 2,2'-Bisnorphaeanthine, P-301
 Bisnortalphenine, T-345
 Bisnortalphosine, I-328
 1,3-Bis(octahydro-6-hydroxy-7-methyl-6-phenyl-5-indolizinyloxy)-2-propanone, D-212
 2,6-Bis(2-oxobutyl)piperidine, D-391
 1,2-Bis(3-oxo-2-piperidyl)ethane, E-226
 7,7'-Bispancoridine, B-143
 Bisparthenolidine, B-215
N,N'-Bis(phenylacetyl)putrescine, B-422
N,N'-Bis(2-phenylethyl)urea, B-216
 10-[Bis(phenylmethyl)amino]-5,6,6a,7-tetrahydro-2-methoxy-6-methyl-4*H*-dibenzo[*de,g*]quinolin-1-ol, V-24
N,N-Bis(phenylmethyl)benzenemethanamine, T-457
 2,5-Bis(phenylmethyl)hexahydropyrazine, D-301
 3,6-Bis(phenylmethyl)-2,5-pyrazinediol, D-300
N,N'-Bis(phenylmethyl)urea, D-302
 Bispasmaplin A, B-217
 Bispyrafoline D, B-192
 Bispyrithione magsulfex, D-831
 Bissecodehydrocyclopiazonic acid, B-218
 Bistabersonine, T-11
 Bistelletadine A, B-219
 Bistelletadine B, B-220
 2,5-Bis(1,2,3,4-tetrahydroxybutyl)pyrazine, B-221
 Bistratamide A, B-223
 Bistratamide B, B-223
 Bistratamide C, B-224
 Bistratamide D, B-225
 Bistratamide E, B-226
 Bistratamide F, B-225
 Bistratamide G, B-225
 Bistratamide H, B-226
 Bistratamide I, B-222
 Bistratamide J, B-227
 Bistratamide K, B-227
 Bistratamide L, B-227
 Bistratamide M, B-227
 Bistratamide N, B-227
 Bistratamide O, B-227
 Bistratamide P, B-227
 Bistratamide Q, B-227
 Bistratamide R, B-227
 Bistratamide S, B-227
 Bistratamide T, B-227
 Bistratamide U, B-227
 Bistratamide V, B-227
 Bistratamide W, B-227
 Bistratamide X, B-227
 Bistratamide Y, B-227
 Bistratamide Z, B-227
 Bistratamide AA, B-227
 Bistratamide AB, B-227
 Bistratamide AC, B-227
 Bistratamide AD, B-227
 Bistratamide AE, B-227
 Bistratamide AF, B-227
 Bistratamide AG, B-227
 Bistratamide AH, B-227
 Bistratamide AI, B-227
 Bistratamide AJ, B-227
 Bistratamide AK, B-227
 Bistratamide AL, B-227
 Bistratamide AM, B-227
 Bistratamide AN, B-227
 Bistratamide AO, B-227
 Bistratamide AP, B-227
 Bistratamide AQ, B-227
 Bistratamide AR, B-227
 Bistratamide AS, B-227
 Bistratamide AT, B-227
 Bistratamide AU, B-227
 Bistratamide AV, B-227
 Bistratamide AW, B-227
 Bistratamide AX, B-227
 Bistratamide AY, B-227
 Bistratamide AZ, B-227
 Bistratamide BA, B-227
 Bistratamide BB, B-227
 Bistratamide BC, B-227
 Bistratamide BD, B-227
 Bistratamide BE, B-227
 Bistratamide BF, B-227
 Bistratamide BG, B-227
 Bistratamide BH, B-227
 Bistratamide BI, B-227
 Bistratamide BJ, B-227
 Bistratamide BK, B-227
 Bistratamide BL, B-227
 Bistratamide BM, B-227
 Bistratamide BN, B-227
 Bistratamide BO, B-227
 Bistratamide BP, B-227
 Bistratamide BQ, B-227
 Bistratamide BR, B-227
 Bistratamide BS, B-227
 Bistratamide BT, B-227
 Bistratamide BU, B-227
 Bistratamide BV, B-227
 Bistratamide BW, B-227
 Bistratamide BX, B-227
 Bistratamide BY, B-227
 Bistratamide BZ, B-227
 Bistratamide CA, B-227
 Bistratamide CB, B-227
 Bistratamide CC, B-227
 Bistratamide CD, B-227
 Bistratamide CE, B-227
 Bistratamide CF, B-227
 Bistratamide CG, B-227
 Bistratamide CH, B-227
 Bistratamide CI, B-227
 Bistratamide CJ, B-227
 Bistratamide CK, B-227
 Bistratamide CL, B-227
 Bistratamide CM, B-227
 Bistratamide CN, B-227
 Bistratamide CO, B-227
 Bistratamide CP, B-227
 Bistratamide CQ, B-227
 Bistratamide CR, B-227
 Bistratamide CS, B-227
 Bistratamide CT, B-227
 Bistratamide CU, B-227
 Bistratamide CV, B-227
 Bistratamide CW, B-227
 Bistratamide CX, B-227
 Bistratamide CY, B-227
 Bistratamide CZ, B-227
 Bistratamide DA, B-227
 Bistratamide DB, B-227
 Bistratamide DC, B-227
 Bistratamide DD, B-227
 Bistratamide DE, B-227
 Bistratamide DF, B-227
 Bistratamide DG, B-227
 Bistratamide DH, B-227
 Bistratamide DI, B-227
 Bistratamide DJ, B-227
 Bistratamide DK, B-227
 Bistratamide DL, B-227
 Bistratamide DM, B-227
 Bistratamide DN, B-227
 Bistratamide DO, B-227
 Bistratamide DP, B-227
 Bistratamide DQ, B-227
 Bistratamide DR, B-227
 Bistratamide DS, B-227
 Bistratamide DT, B-227
 Bistratamide DU, B-227
 Bistratamide DV, B-227
 Bistratamide DW, B-227
 Bistratamide DX, B-227
 Bistratamide DY, B-227
 Bistratamide DZ, B-227
 Bistratamide EA, B-227
 Bistratamide EB, B-227
 Bistratamide EC, B-227
 Bistratamide ED, B-227
 Bistratamide EE, B-227
 Bistratamide EF, B-227
 Bistratamide EG, B-227
 Bistratamide EH, B-227
 Bistratamide EI, B-227
 Bistratamide EJ, B-227
 Bistratamide EK, B-227
 Bistratamide EL, B-227
 Bistratamide EM, B-227
 Bistratamide EN, B-227
 Bistratamide EO, B-227
 Bistratamide EP, B-227
 Bistratamide EQ, B-227
 Bistratamide ER, B-227
 Bistratamide ES, B-227
 Bistratamide ET, B-227
 Bistratamide EU, B-227
 Bistratamide EV, B-227
 Bistratamide EW, B-227
 Bistratamide EX, B-227
 Bistratamide EY, B-227
 Bistratamide EZ, B-227
 Bistratamide FA, B-227
 Bistratamide FB, B-227
 Bistratamide FC, B-227
 Bistratamide FD, B-227
 Bistratamide FE, B-227
 Bistratamide FF, B-227
 Bistratamide FG, B-227
 Bistratamide FH, B-227
 Bistratamide FI, B-227
 Bistratamide FJ, B-227
 Bistratamide FK, B-227
 Bistratamide FL, B-227
 Bistratamide FM, B-227
 Bistratamide FN, B-227
 Bistratamide FO, B-227
 Bistratamide FP, B-227
 Bistratamide FQ, B-227
 Bistratamide FR, B-227
 Bistratamide FS, B-227
 Bistratamide FT, B-227
 Bistratamide FU, B-227
 Bistratamide FV, B-227
 Bistratamide FW, B-227
 Bistratamide FX, B-227
 Bistratamide FY, B-227
 Bistratamide FZ, B-227
 Bistratamide GA, B-227
 Bistratamide GB, B-227
 Bistratamide GC, B-227
 Bistratamide GD, B-227
 Bistratamide GE, B-227
 Bistratamide GF, B-227
 Bistratamide GG, B-227
 Bistratamide GH, B-227
 Bistratamide GI, B-227
 Bistratamide GJ, B-227
 Bistratamide GK, B-227
 Bistratamide GL, B-227
 Bistratamide GM, B-227
 Bistratamide GN, B-227
 Bistratamide GO, B-227
 Bistratamide GP, B-227
 Bistratamide GQ, B-227
 Bistratamide GR, B-227
 Bistratamide GS, B-227
 Bistratamide GT, B-227
 Bistratamide GU, B-227
 Bistratamide GV, B-227
 Bistratamide GW, B-227
 Bistratamide GX, B-227
 Bistratamide GY, B-227
 Bistratamide GZ, B-227
 Bistratamide HA, B-227
 Bistratamide HB, B-227
 Bistratamide HC, B-227
 Bistratamide HD, B-227
 Bistratamide HE, B-227
 Bistratamide HF, B-227
 Bistratamide HG, B-227
 Bistratamide HH, B-227
 Bistratamide HI, B-227
 Bistratamide HJ, B-227
 Bistratamide HK, B-227
 Bistratamide HL, B-227
 Bistratamide HM, B-227
 Bistratamide HN, B-227
 Bistratamide HO, B-227
 Bistratamide HP, B-227
 Bistratamide HQ, B-227
 Bistratamide HR, B-227
 Bistratamide HS, B-227
 Bistratamide HT, B-227
 Bistratamide HU, B-227
 Bistratamide HV, B-227
 Bistratamide HW, B-227
 Bistratamide HX, B-227
 Bistratamide HY, B-227
 Bistratamide HZ, B-227
 Bistratamide IA, B-227
 Bistratamide IB, B-227
 Bistratamide IC, B-227
 Bistratamide ID, B-227
 Bistratamide IE, B-227
 Bistratamide IF, B-227
 Bistratamide IG, B-227
 Bistratamide IH, B-227
 Bistratamide II, B-227
 Bistratamide IJ, B-227
 Bistratamide IK, B-227
 Bistratamide IL, B-227
 Bistratamide IM, B-227
 Bistratamide IN, B-227
 Bistratamide IO, B-227
 Bistratamide IP, B-227
 Bistratamide IQ, B-227
 Bistratamide IR, B-227
 Bistratamide IS, B-227
 Bistratamide IT, B-227
 Bistratamide IU, B-227
 Bistratamide IV, B-227
 Bistratamide IW, B-227
 Bistratamide IX, B-227
 Bistratamide IY, B-227
 Bistratamide IZ, B-227
 Bistratamide JA, B-227
 Bistratamide JB, B-227
 Bistratamide JC, B-227
 Bistratamide JD, B-227
 Bistratamide JE, B-227
 Bistratamide JF, B-227
 Bistratamide JG, B-227
 Bistratamide JH, B-227
 Bistratamide JI, B-227
 Bistratamide JJ, B-227
 Bistratamide JK, B-227
 Bistratamide JL, B-227
 Bistratamide JM, B-227
 Bistratamide JN, B-227
 Bistratamide JO, B-227
 Bistratamide JP, B-227
 Bistratamide JQ, B-227
 Bistratamide JR, B-227
 Bistratamide JS, B-227
 Bistratamide JT, B-227
 Bistratamide JU, B-227
 Bistratamide JV, B-227
 Bistratamide JW, B-227
 Bistratamide JX, B-227
 Bistratamide JY, B-227
 Bistratamide JZ, B-227
 Bistratamide KA, B-227
 Bistratamide KB, B-227
 Bistratamide KC, B-227
 Bistratamide KD, B-227
 Bistratamide KE, B-227
 Bistratamide KF, B-227
 Bistratamide KG, B-227
 Bistratamide KH, B-227
 Bistratamide KI, B-227
 Bistratamide KJ, B-227
 Bistratamide KK, B-227
 Bistratamide KL, B-227
 Bistratamide KM, B-227
 Bistratamide KN, B-227
 Bistratamide KO, B-227
 Bistratamide KP, B-227
 Bistratamide KQ, B-227
 Bistratamide KR, B-227
 Bistratamide KS, B-227
 Bistratamide KT, B-227
 Bistratamide KU, B-227
 Bistratamide KV, B-227
 Bistratamide KW, B-227
 Bistratamide KX, B-227
 Bistratamide KY, B-227
 Bistratamide KZ, B-227
 Bistratamide LA, B-227
 Bistratamide LB, B-227
 Bistratamide LC, B-227
 Bistratamide LD, B-227
 Bistratamide LE, B-227
 Bistratamide LF, B-227
 Bistratamide LG, B-227
 Bistratamide LH, B-227
 Bistratamide LI, B-227
 Bistratamide LJ, B-227
 Bistratamide LK, B-227
 Bistratamide LL, B-227
 Bistratamide LM, B-227
 Bistratamide LN, B-227
 Bistratamide LO, B-227
 Bistratamide LP, B-227
 Bistratamide LQ, B-227
 Bistratamide LR, B-227
 Bistratamide LS, B-227
 Bistratamide LT, B-227
 Bistratamide LU, B-227
 Bistratamide LV, B-227
 Bistratamide LW, B-227
 Bistratamide LX, B-227
 Bistratamide LY, B-227
 Bistratamide LZ, B-227
 Bistratamide MA, B-227
 Bistratamide MB, B-227
 Bistratamide MC, B-227
 Bistratamide MD, B-227
 Bistratamide ME, B-227
 Bistratamide MF, B-227
 Bistratamide MG, B-227
 Bistratamide MH, B-227
 Bistratamide MI, B-227
 Bistratamide MJ, B-227
 Bistratamide MK, B-227
 Bistratamide ML, B-227
 Bistratamide MM, B-227
 Bistratamide MN, B-227
 Bistratamide MO, B-227
 Bistratamide MP, B-227
 Bistratamide MQ, B-227
 Bistratamide MR, B-227
 Bistratamide MS, B-227
 Bistratamide MT, B-227
 Bistratamide MU, B-227
 Bistratamide MV, B-227
 Bistratamide MW, B-227
 Bistratamide MX, B-227
 Bistratamide MY, B-227
 Bistratamide MZ, B-227
 Bistratamide NA, B-227
 Bistratamide NB, B-227
 Bistratamide NC, B-227
 Bistratamide ND, B-227
 Bistratamide NE, B-227
 Bistratamide NF, B-227
 Bistratamide NG, B-227
 Bistratamide NH, B-227
 Bistratamide NI, B-227
 Bistratamide NJ, B-227
 Bistratamide NK, B-227
 Bistratamide NL, B-227
 Bistratamide NM, B-227
 Bistratamide NN, B-227
 Bistratamide NO, B-227
 Bistratamide NP, B-227
 Bistratamide NQ, B-227
 Bistratamide NR, B-227
 Bistratamide NS, B-227
 Bistratamide NT, B-227
 Bistratamide NU, B-227
 Bistratamide NV, B-227
 Bistratamide NW, B-227
 Bistratamide NX, B-227
 Bistratamide NY, B-227
 Bistratamide NZ, B-227
 Bistratamide OA, B-227
 Bistratamide OB, B-227
 Bistratamide OC, B-227
 Bistratamide OD, B-227
 Bistratamide OE, B-227
 Bistratamide OF, B-227
 Bistratamide OG, B-227
 Bistratamide OH, B-227
 Bistratamide OI, B-227
 Bistratamide OJ, B-227
 Bistratamide OK, B-227
 Bistratamide OL, B-227
 Bistratamide OM, B-227
 Bistratamide ON, B-227
 Bistratamide OO, B-227
 Bistratamide OP, B-227
 Bistratamide OQ, B-227
 Bistratamide OR, B-227
 Bistratamide OS, B-227
 Bistratamide OT, B-227
 Bistratamide OU, B-227
 Bistratamide OV, B-227
 Bistratamide OW, B-227
 Bistratamide OX, B-227
 Bistratamide OY, B-227
 Bistratamide OZ, B-227
 Bistratamide PA, B-227
 Bistratamide PB, B-227
 Bistratamide PC, B-227
 Bistratamide PD, B-227
 Bistratamide PE, B-227
 Bistratamide PF, B-227
 Bistratamide PG, B-227
 Bistratamide PH, B-227
 Bistratamide PI, B-227
 Bistratamide PJ, B-227
 Bistratamide PK, B-227
 Bistratamide PL, B-227
 Bistratamide PM, B-227
 Bistratamide PN, B-227
 Bistratamide PO, B-227
 Bistratamide PP, B-227
 Bistratamide PQ, B-227
 Bistratamide PR, B-227
 Bistratamide PS, B-227
 Bistratamide PT, B-227
 Bistratamide PU, B-227
 Bistratamide PV, B-227
 Bistratamide PW, B-227
 Bistratamide PX, B-227
 Bistratamide PY, B-227
 Bistratamide PZ, B-227
 Bistratamide QA, B-227
 Bistratamide QB, B-227
 Bistratamide QC, B-227
 Bistratamide QD, B-227
 Bistratamide QE, B-227
 Bistratamide QF, B-227
 Bistratamide QG, B-227
 Bistratamide QH, B-227
 Bistratamide QI, B-227
 Bistratamide QJ, B-227
 Bistratamide QK, B-227
 Bistratamide QL, B-227
 Bistratamide QM, B-227
 Bistratamide QN, B-227
 Bistratamide QO, B-227
 Bistratamide QP, B-227
 Bistratamide QQ, B-227
 Bistratamide QR, B-227
 Bistratamide QS, B-227
 Bistratamide QT, B-227
 Bistratamide QU, B-227
 Bistratamide QV, B-227
 Bistratamide QW, B-227
 Bistratamide QX, B-227
 Bistratamide QY, B-227
 Bistratamide QZ, B-227
 Bistratamide RA, B-227
 Bistratamide RB, B-227
 Bistratamide RC, B-227
 Bistratamide RD, B-227
 Bistratamide RE, B-227
 Bistratamide RF, B-227
 Bistratamide RG, B-227
 Bistratamide RH, B-227
 Bistratamide RI, B-227
 Bistratamide RJ, B-227
 Bistratamide RK, B-227
 Bistratamide RL, B-227
 Bistratamide RM, B-227
 Bistratamide RN, B-227
 Bistratamide RO, B-227
 Bistratamide RP, B-227
 Bistratamide RQ, B-227
 Bistratamide RR, B-227
 Bistratamide RS, B-227
 Bistratamide RT, B-227
 Bistratamide RU, B-227
 Bistratamide RV, B-227
 Bistratamide RW, B-227
 Bistratamide RX, B-227
 Bistratamide RY, B-227
 Bistratamide RZ, B-227
 Bistratamide SA, B-227
 Bistratamide SB, B-227
 Bistratamide SC, B-227
 Bistratamide SD, B-227
 Bistratamide SE, B-227
 Bistratamide SF, B-227
 Bistratamide SG, B-227
 Bistratamide SH, B-227
 Bistratamide SI, B-227
 Bistratamide SJ, B-227
 Bistratamide SK, B-227
 Bistratamide SL, B-227
 Bistratamide SM, B-227
 Bistratamide SN, B-227
 Bistratamide SO, B-227
 Bistratamide SP, B-227
 Bistratamide SQ, B-227
 Bistratamide SR, B-227
 Bistratamide SS, B-227
 Bistratamide ST, B-227
 Bistratamide SU, B-227
 Bistratamide SV, B-227
 Bistratamide SW, B-227
 Bistratamide SX, B-227
 Bistratamide SY, B-227
 Bistratamide SZ, B-227
 Bistratamide TA, B-227
 Bistratamide TB, B-227
 Bistratamide TC, B-227
 Bistratamide TD, B-227
 Bistratamide TE, B-227
 Bistratamide TF, B-227
 Bistratamide TG, B-227
 Bistratamide TH, B-227
 Bistratamide TI, B-227
 Bistratamide TJ, B-227
 Bistratamide TK, B-227
 Bistratamide TL, B-227
 Bistratamide TM, B-227
 Bistratamide TN, B-227
 Bistratamide TO, B-227
 Bistratamide TP, B-227
 Bistratamide TQ, B-227
 Bistratamide TR, B-227
 Bistratamide TS, B-227
 Bistratamide TT, B-227
 Bistratamide TU, B-227
 Bistratamide TV, B-227
 Bistratamide TW, B-227
 Bistratamide TX, B-227
 Bistratamide TY, B-227
 Bistratamide TZ, B-227
 Bistratamide UA, B-227
 Bistratamide UB, B-227
 Bistratamide UC, B-227
 Bistratamide UD, B-227
 Bistratamide UE, B-227
 Bistratamide UF, B-227
 Bistratamide UG, B-227
 Bistratamide UH, B-227
 Bistratamide UI, B-227
 Bistratamide UJ, B-227
 Bistratamide UK, B-227
 Bistratamide UL, B-227
 Bistratamide UM, B-227
 Bistratamide UN, B-227
 Bistratamide UO, B-227
 Bistratamide UP, B-227
 Bistratamide UQ, B-227
 Bistratamide UR, B-227
 Bistratamide US, B-227
 Bistratamide UT, B-227
 Bistratamide UU, B-227
 Bistratamide UV, B-227
 Bistratamide UW, B-227
 Bistratamide UX, B-227
 Bistratamide UY, B-227
 Bistratamide UZ, B-227
 Bistratamide VA, B-227
 Bistratamide VB, B-227
 Bistratamide VC, B-227
 Bistratamide VD, B-227
 Bistratamide VE, B-227
 Bistratamide VF, B-227
 Bistratamide VG, B-227
 Bistratamide VH, B-227
 Bistratamide VI, B-227
 Bistratamide VJ, B-227
 Bistratamide VK, B-227
 Bistratamide VL, B-227
 Bistratamide VM, B-227
 Bistratamide VN, B-227
 Bistratamide VO, B-227
 Bistratamide VP, B-227
 Bistratamide VQ, B-227
 Bistratamide VR, B-227
 Bistratamide VS, B-227
 Bistratamide VT, B-227
 Bistratamide VU, B-227
 Bistratamide VV, B-227
 Bistratamide VW, B-227
 Bistratamide VX, B-227
 Bistratamide VY, B-227
 Bistratamide VZ, B-227
 Bistratamide WA, B-227
 Bistratamide WB, B-227
 Bistratamide WC, B-227
 Bistrat

- Bosistidine, T-580
 Bosistine, T-546
 Bosmin, A-152
 Bostrycoidin, B-255
 Botryllamide A, B-256
 Botryllamide B, B-256
 Botryllamide C, B-256
 Botryllamide D, B-256
 Botryllamide E, B-256
 Botryllamide F, B-256
 Botryllamide G, B-256
 Botryllamide H, B-257
 Botryllazine A, B-258
 Botryllazine B, B-259
 Bouchardatine, B-260
 Bougainvillein r-I, B-110
 Bougainvillein r-II, B-110
 Bougainvillein r-III, B-110
 Bougainvillein r-IV, B-110
 Bougainvillein r-V, B-110
 Bougainvillein-v, B-110
 Bowdensine, B-261
 Bowdichine, A-107
 Boxazomycin A, B-262
 Boxazomycin B, B-262
 Boxazomycin C, B-262
 BP 2, O-197
 BP 179, T-52
 BR 040, A-1139
 BR 042, A-1139
 Brachyaconitine, M-331
 Brachyamide A, B-263
 Brachyamide B, M-455
 Brachycerine, B-264
 Brachystamide A, B-265
 Brachystamide B, B-265
 Brachystamide C, B-265
 Brachystamide D, B-266
 Brachystamide E, B-266
 Brachystemidine A, B-267
 Brachystemidine B, B-268
 Brachystemidine C, B-268
 Brachystemidine D, B-268
 Brachystemidine E, B-268
 Brachystemidine F, B-269
 Brachystemidine G, B-270
 Brachystine, B-271
 Bractamine, B-272
 Bractavine, O-110
 Bractazonine, B-273
 Bracteatine, B-274
 Bracteine, B-275
 Bracteoline, B-276
 Bradyoxetin, B-277
 Brafoedine, A-1365
 Brasiliamide A, B-278
 Brasiliamide B, B-279
 Brasiliamide C, B-279
 Brasiliamide D, B-279
 Brasiliamide E, B-279
 Brasilibactin A, N-265
 Brasilicardin A, B-280
 Brasilicardin B, B-280
 Brasilicardin C, B-280
 Brasilicardin D, B-280
 Brasilidine A, B-281
 Brassicanal A, M-201
 Brassicanal B, B-282
 Brassicanal C, B-283
 Brassicanate A, M-202
 Brassidic acid, D-880
 Brasilexin, I-336
 Brassinin, B-284
 Brassitin, B-285
 Bravomicin A, B-286
 Bravomicin B, B-286
 Bravomicin C, B-286
 Bravomicin D, B-287
 Bravomicin E, B-287
 Bravomicin F, B-287
 Braznitidumine, B-288
 Bredilonine, P-962
 Brefelamide, B-289
 Breoganine, C-805
 Brevianamide E, B-290
 Brevianamide F, C-899
 Brevicarine, B-291
 Brevicide, X-7
 Brevicolline, B-292
 Brevicompanine A, B-293
 Brevicompanine B, F-160
 Brevicompanine C, B-294
 Brevigellin, B-295
 Brevinic acid, B-296
 Brevioxime, B-297
 Bripiodionene, B-298
 Brocadopa, A-745
 Brochodine, C-555
 Brolitene, H-488
 Bromcholitin, T-246
 Bromergocryptine, E-157
 Bromergon, E-157
 2-Bromoageliferin, A-190
 2-Bromoaldisine, A-255
 6-Bromoaplysinopsin, A-1357
 Bromobenzisoxazolone Baretin, B-25
 6-Bromo-2-(6-bromo-1,3-dihydro-3-oxo-2*H*-indol-2-ylidene)-1,2-dihydro-3*H*-indol-3-one, D-325
 6-Bromo-1-(3-bromo-4-hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411
 3-Bromo-2-(5-bromo-2-hydroxyphenyl)-4,5-dichloropyrrole, P-966
 6-Bromo- β -(6-bromo-1*H*-indol-3-yl)-1*H*-indole-3-ethanamine, B-162
 3-Bromo-4-(3-bromo-2-nitrophenyl)-1*H*-pyrrole, D-305
 6-Bromo- β -carboline, B-299
 7-Bromo- β -carboline, B-300
 7-Bromocavernicolone, B-301
 5-Bromocavernicolin, C-221
 Bromochelonin B, C-362
 Bromochloroacetamide, B-302
 Bromochloroacetic acid, B-302
 Bromochloroacetonitrile, B-302
 7-Bromo-5-chlorocavernicolin, C-221
 Bromochlorocyanomethane, B-302
 6-Bromo-3-chloro-1*H*-indole, B-303
 7-Bromo-5-(2-chlorophenyl)-1,3-dihydro-3-hydroxy-2*H*-1,4-benzodiazepin-2-one, B-304
 3-Bromo-5-chlorotyrosine, A-712
 3-Bromo-5-chlororongiaquinol, H-650
 Bromocriptine mesilate, E-157
 Bromocriptine, E-157
 Bromocryptine, E-157
 6-Bromo-3-(cyanomethyl)indole, B-332
 6-Bromo-6-dechloropentrem A, P-194
 6-Bromo-3'-deimino-2'-4'-bis(dimethyl)-3'-oxoaplysinopsin, A-1357
 6-Bromo-3'-deimino-3'-oxoaplysinopsin, A-1357
 6-Bromo-2'-*N*-demethylaplysinopsin, A-1357
 6-Bromo-4'-*N*-demethylaplysinopsin, A-1357
 6-Bromo-2'-*N*-demethyl-*N*-3'-methylaplysinopsin, A-1357
 Bromodeoxytopsentin, T-432
 5-Bromo-5'-deoxytubercidin, T-658
 9-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]adenine, B-305
 7-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]-3,7-dihydro-1*H*-purine-2,6-dione, B-306
N-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]-*N'*-(3-methoxycarbonylpropyl)urea, B-307
 6-Bromo-1-(3,5-dibromo-4-hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411
 7-Bromo-1-(3,5-dibromo-4-hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411
 7-Bromo-2,3-dichloro-1*H*-indole, B-308
 14-Bromo-7,8-didehydro-3-dihydrodiscorhabdin C, D-851
 6-Bromo-1',8-dihydroaplysinopsin, A-1357
 6-Bromo-1,2-dihydro-2,2-bis(methylthio)-3*H*-indol-3-one, B-309
 14-Bromo-3-dihydro-7,8-didehydrodiscorhabdin C, D-851
 14-Bromo-3-dihydrodiscorhabdin C, D-851
 19-Bromo-3,4-dihydroeudistomin U, I-119
 10-Bromo-6,7-dihydrohomofascaplysate A, H-332
 6-Bromo-1',8-dihydro-1'-hydroxyaplysinopsin, A-1357
 6-Bromo-1,2-dihydro-2-hydroxymethyl-2-methoxy-3*H*-indol-3-one, M-115
 5-Bromo-1,2-dihydro-3*H*-indol-3-one, B-324
 6-Bromo-1,2-dihydro-3*H*-indol-3-one, B-325
 6-Bromo-1',8-dihydro-1'-methoxyaplysinopsin, A-1357
 7-Bromo-2,3-dihydro-2-methyl-3-[(methylthio)methyl]-1*H*-2,4,10*b*-triazafuoranthene-8-ol, E-283
 6-Bromo-1-(3,4-dihydro-2*H*-pyrrol-5-yl)-9*H*-pyrido[3,4-*b*]indole, E-288
 7-Bromo-1-(3,4-dihydro-2*H*-pyrrol-5-yl)-9*H*-pyrido[3,4-*b*]indole, E-285
 6-Bromo-1-(3,4-dihydro-2*H*-pyrrol-5-yl)-9*H*-pyrido[3,4-*b*]indol-6-ol, E-285
 6-Bromo-4,5-dihydroxyindole, B-310
 6-Bromo-4,7-dihydroxyindole, B-311
 2-Bromo-6,8-dihydroxy-7-propyl-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, S-571
 7-Bromo-2,4-dihydroxyquinazoline, B-312
 6-Bromo-2,4-dihydroxyquinoline, B-313
 11-Bromo-3,12-dihydroxy-2,2,4,10-tetramethyl-4,6,8,10,14,16-eicosahexaen-18-ynamide, C-497
 6-Bromo-2-(1,1-dimethylallyl)tryptamine, B-316
 7-Bromo-1-[2-(dimethylamino)ethyl]- β -carboline, P-507
 12-[3-Bromo-2-[2-(dimethylamino)ethyl]-5-hydroxy-4-methoxyphenyl]-2,6,10-trimethyl-2,10-dodecadien-5-one, A-1342
 5-Bromo-3-(2-dimethylaminoethyl)indole, B-355
 5-Bromo-*N,N*-dimethyl-1*H*-indole-3-ethanamine, B-355
 6-Bromo-3a-(3,7-dimethyl-2,6-octadienyl)-1,2,3,3a,8,8a-hexahydro-1-methylpyrrolo[2,3-*b*]indol-7-ol, B-314
 6-Bromo-2-(1,1-dimethyl-2-propenyl)-1*H*-indole-3-carboxaldehyde, B-315
 6-Bromo-2-(1,1-dimethyl-2-propenyl)-1*H*-indole-3-ethanamine, B-316
N-[2-[6-Bromo-2-(1,1-dimethyl-2-propenyl)-1*H*-indol-3-yl]ethyl]-*N*-methylmethanesulfonamide, B-316
 6-Bromo-2-(1,1-dimethyl-2-propenyl)tryptamine, B-316
 7-Bromo-*N,N*-dimethyl-9*H*-pyrido[3,4-*b*]indole-1-ethanamine, P-507
 5-Bromo-*N^b,N^b*-dimethyltryptamine, B-355
p-Bromo- α,α -diphenylacetanilide, B-344
 14-Bromodiscorhabdin C, D-851
 7-Bromo-1-ethenyl-9*H*-pyrido[3,4-*b*]indole, V-139
 6-Bromo-1'-ethoxy-1',8-dihydroaplysinopsin, A-1357
 7-Bromo-4-(2-ethoxyethyl)quinoline, B-317
 6-Bromo-1-ethyl- β -carboline, B-318
 6-Bromo-1-ethyl-9*H*-pyrido[3,4-*b*]indole, B-318
 10-Bromofascaplysin, F-13
 3-Bromofascaplysin, F-13
 2-Bromofestucalvin, F-49
 6-Bromo-3-formyl-2-(1,1-dimethyl-2-propenyl)indole, B-315
 6-Bromo-3-formyl-5-hydroxyindole, B-327
 6-Bromo-3-formylindole, B-333
 5'-Bromo-5-formyl-4-methoxy-2,2'-bipyrrrole, B-339
 3-Bromo-5'-formyl-4'-methoxy-2,2'-bipyrrrole, B-338
 6-Bromo-*N^b*-formyl-*N^b*-methyltryptamine, B-356
 6-Bromogranulatimide, G-169
 4-Bromo-2-(guanidinomethyl)pyrrole, B-319
 3-Bromohomofascaplysin B1, H-335
 3-Bromohomofascaplysin B, H-334
 3-Bromohomofascaplysin C, H-336
 6-Bromo-3-(hydroxyacetyl)-1*H*-indole, B-320
 6-Bromo-1-(4-hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411
 1-(3-Bromo-4-hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411

- 5-Bromo-6-hydroxy- β -carboline, B-321
 7-Bromo-6-hydroxy- β -carboline, B-322
 14-Bromo-1-hydroxydiscorhabdin V, D-849
 3-Bromo-4-hydroxy- α -(hydroxyimino)-*N*-[2-(1*H*-imidazol-4-yl)ethyl]benzenepropanamide, B-323
 5-Bromo-3-hydroxy-1*H*-indole, B-324
 6-Bromo-3-hydroxy-1*H*-indole, B-325
 6-Bromo-5-hydroxy-1*H*-indole, B-326
 6-Bromo-5-hydroxy-1*H*-indole-3-carboxaldehyde, B-327
 5-Bromo-*N*-hydroxy-1*H*-indole-3-ethanamine, B-355
N-(3-Bromo- β -hydroxy-4-methoxyphenethyl)tryptamine, C-362
 6-Bromo-3-hydroxy-2-(methylthio)indole, B-328
 2-(3-Bromo-4-hydroxyphenyl)ethylamine, A-769
 [2-(3-Bromo-4-hydroxyphenyl)ethyl]ethanedi-
 amide, B-329
N-[2-(3-Bromo-4-hydroxyphenyl)ethyl]- α -oxo-1*H*-
 indole-3-acetamide, P-541
 2-[[3-(3-Bromo-4-hydroxyphenyl)-2-(hydroxyimi-
 no)-1-oxopropyl]amino]ethyl thiocyanate, P-687
 5-Bromo-6-hydroxy-9*H*-pyrido[3,4-*b*]indole, B-321
 7-Bromo-6-hydroxy-1-(2-pyrrolyl)- β -carboline,
 E-285
 6-Bromo-4-hydroxy-2(1*H*)-quinolinone, B-313
 3-Bromo-4-hydroxy-*N,N,N*-trimethylbenzeneetha-
 naminium, A-769
 5-Bromo-*N'*-hydroxytryptamine, B-355
 3-Bromohymenialdisine, H-762
 5-Bromohypaphorine, H-764
 6-Bromohypaphorine, H-764
 6-Bromoindigo, D-325
 6-Bromoindigotin, D-325
 6-Bromoindirubin, I-73
 6-Bromo-1*H*-indole-3-acetamide, B-332
 5-Bromo-1*H*-indole-3-acetic acid, B-331
 6-Bromo-1*H*-indole-3-acetic acid, B-332
 6-Bromo-1*H*-indole-3-acetonitrile, B-332
 3-Bromo-1*H*-indole, B-330
 6-Bromoindole-3-carbaldehyde, B-333
 6-Bromo-1*H*-indole-3-carboxaldehyde, B-333
 6-Bromo-1*H*-indole-3-carboxylic acid, B-334
 6-Bromo-1*H*-indole-4,5-diol, B-310
 6-Bromo-1*H*-indole-4,7-diol, B-311
 6-Bromo-1*H*-indole-2,3-dione, B-335
 5-Bromo-1*H*-indole-3-ethanamine, B-355
 6-Bromo-1*H*-indole-3-ethanamine, B-356
 5-Bromo-1*H*-indol-3-ol, B-324
 6-Bromo-1*H*-indol-5-ol, B-326
 6-Bromo-1*H*-indol-3-ol, B-325
 5-Bromo-3-indolone, B-324
 6-Bromo-3-indolone, B-325
 4-(6-Bromo-1*H*-indol-3-yl)-3-buten-2-one, I-117
*N*²-(6-Bromo-1*H*-indol-3-ylcarbonyl)arginine,
 I-120
*N*²-(6-Bromo-1*H*-indol-3-ylcarbonyl)enduracidi-
 dine, E-80
*N*²-(6-Bromo-1*H*-indol-3-ylcarbonyl)histidine,
 I-121
 3-Bromo- α -[[2-(1*H*-indol-3-yl)ethyl]amino]-
 methyl]-4-methoxybenzenemethanol, C-362
N-[2-(6-Bromo-1*H*-indol-3-yl)ethyl]-*N*-methylfor-
 mamide, B-356
 (6-Bromo-1*H*-indol-3-yl)glyoxylic acid, B-342
 1-(6-Bromo-1*H*-indol-3-yl)-2-hydroxyethanonone,
 B-320
 3-(6-Bromo-1*H*-indol-3-yl)-2-propenoic acid,
 B-336
 6-Bromoindoxyl, B-325
 Bromoiodoacetamide, B-337
 Bromoiodoacetic acid, B-337
N-[2-[3-Bromo-5-iodo-4-[3-[(3-methyl-1-oxo-2-
 butenyl)amino]propoxy]phenyl]ethyl]-4-hydro-
 xybenzeneacetamide, D-333
 6-Bromoisatin, B-335
 19-Bromoisoeudistomin U (incorr.), I-119
 2-Bromoleptoclimidinone, A-1459
 2-Bromomaleimide, B-350
 5'-Bromo-4-methoxy-2,2'-bipyrrrole-5-carboxalde-
 hyde, B-339
 3'-Bromo-4-methoxy-2,2'-bipyrrrole-5-carboxalde-
 hyde, B-338
 4-Bromo-*N*²-methoxymethyl-1*H*-pyrrole-2-carbox-
 amide, B-348
 5-Bromo-*N*²-(methoxymethyl)-1*H*-pyrrole-2-car-
 boxamide, B-349
 3-[2-[[2-(3-Bromo-4-methoxyphenyl)-2-hydro-
 xyethyl]amino]ethyl]indole, C-362
 6-Bromo-7-(methylamino)-5,8-isoquinolinedione,
 A-831
 5-Bromo-1-[1-(methylamino)-2-(methylthio)ethyl]-
 9*H*-pyrido[3,4-*b*]indol-6-ol, E-282
 6-Bromo-1-methyl- β -carboline, B-340
N-Bromo-*N*-methylformamide, M-463
 4-Bromo-*N*⁷-methylphakellin, B-343
 6-Bromo-*N*^b-methyl-*N*^a-prenyltryptamine, B-356
 7-Bromo- α -(2-methylpropyl)-9*H*-pyrido[3,4-*b*]in-
 dole-1-methanamine, E-278
 6-Bromo-1-methyl-9*H*-pyrido[3,4-*b*]indole, B-340
 7-Bromo-9-[[4-(methylthio)-1*H*-imidazol-3-yl]-
 methyl]-9*H*-pyrido[3,4-*b*]indole, D-380
 6-Bromo-2-(methylthio)-1*H*-indol-3-ol, B-328
 6-Bromo-2-(methylthio)-3*H*-indol-3-one, B-341
 7-Bromo-5-methyl-1,3,7-triazecine-
 2,6,8(1*H*,3*H*,7*H*)-trione, R-70
 2-Bromomyrothenone B, A-824
 Bromonitrin A, D-305
 Bromonitrin B, D-305
 Bromonitrin C, D-305
 3-(3-Bromo-2-nitrophenyl)-1*H*-pyrrole, P-961
 Bromonorharman, B-299
 6-Bromo- α -oxo-1*H*-indole-3-acetamide, B-342
 6-Bromo- α -oxo-1*H*-indole-3-acetic acid, B-342
 4-Bromopalauamine, P-18
 6-Bromopenitrem E, P-194
 4-Bromophakellin, B-343
N-(4-Bromophenyl)-2,2-diphenylacetamide,
 B-344
N-(4-Bromophenyl)- α -phenylbenzeneacetamide,
 B-344
 6-Bromo-*N*¹-propionylaplysinsinopsin, A-1357
 5-Bromopsammaplinsin A, P-686
 7-Bromo-9*H*-pyrido[3,4-*b*]indole, B-300
 6-Bromo-9*H*-pyrido[3,4-*b*]indole, B-299
 5-Bromo-9*H*-pyrido[3,4-*b*]indol-6-ol, B-321
 7-Bromo-9*H*-pyrido[3,4-*b*]indol-6-ol, B-322
 1-(7-Bromo-9*H*-pyrido[3,4-*b*]indol-1-yl)-3-methyl-
 1-butanone, E-279
*N*²-(4-Bromo-1*H*-pyrrole-2-carbonyl)arginine, B-345
*N*²-(4-Bromo-1*H*-pyrrole-2-carbonyl)homoargi-
 nine, B-346
*N*⁶-(4-Bromo-1*H*-pyrrole-2-carbonyl)lysine, B-347
 5-Bromo-1*H*-pyrrole-2-carboxamide, B-349
 4-Bromo-1*H*-pyrrole-2-carboxylic acid, B-348
 5-Bromo-1*H*-pyrrole-2-carboxylic acid, B-349
 3-Bromo-1*H*-pyrrole-2,5-dione, B-350
 6-Bromo-1-(1-pyrrolin-2-yl)- β -carboline, E-288
 Bromopyrrolnitrin A, D-305
 Bromopyrrolnitrin B, D-305
 Bromopyrrolnitrin C, D-305
 5-Bromo-1*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine,
 A-713
 5-Bromo-7*H*-pyrrolo[2,3-*d*]pyrimidine, B-351
 7-Bromo-1-(1*H*-pyrrol-2-yl)-9*H*-pyrido[3,4-*b*]in-
 dol-6-ol, E-285
 7-Bromo-2,4-quinazolinediol, B-312
 7-Bromo-2,4(1*H*,3*H*)-quinazolinedione, B-312
 6-Bromo-2,4-quinolinediol, B-313
 Bromorebeccamycin, R-32
 14-Bromoreticulate, R-57
 14-Bromoreticulate, R-57
 14-Bromoreticulatol, R-58
 Bromosalinosporamide, S-28
 Bromosceptrin, S-124
 3-Bromosecofascaplysin A, S-186
 3-Bromosecofascaplysin B, S-186
 3-Bromostyloquinidine, S-611
 6-Bromo-1,2,3,4-tetrahydro-7,8-dihydroxy-3-iso-
 quinolinecarboxylic acid, B-352
 8-Bromo-1,2,3,4-tetrahydro-6,7-dihydroxy-3-iso-
 quinolinecarboxylic acid, B-353
 6-Bromo-1,2,3,4-tetrahydro-8-hydroxy-7-methoxy-
 3-isoquinolinecarboxylic acid, B-352
 8-Bromo-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-
 3-isoquinolinecarboxylic acid, B-353
 7-Bromo-2,3,4,9-tetrahydro-1-(1*H*-imidazol-4-yl-
 methyl)-1*H*-pyrido[3,4-*b*]indole, L-199
 6-Bromo-2,3,4,9-tetrahydro-2-methyl- α -[(4-methyl-
 phenyl)methyl]-1*H*-pyrido[3,4-*b*]indole-1-metha-
 namine, E-281
 5-(2-Bromo-5,6,7,8-tetrahydro-8-oxopyrrolo[2,3-
 c]azepin-4(1*H*)-ylidene)-2,4-imidazolidinedione,
 A-1576
 6-Bromo-1,3,4,5-tetrahydropyrrolo[4,3,2-*de*]quino-
 line-7,8-dione, D-13
 6-Bromotopsentin A, T-432
 21-Bromotopsentin A, T-432
 Bromotopsentin, T-432
 4(7)-Bromo-2,3,7(2,3,4)-trichloro-1*H*-indole,
 B-354
 3-Bromo-*N,N,N*-trimethyltyramine, A-769
 5-Bromotryptamine, B-355
 6-Bromotryptamine, B-356
 3-Bromotyramine, A-769
Oceanapia Bromotyrosine alkaloid, B-357
 5-Bromoverongamine, V-78
 3-Bromoverongiaquinol, H-650
 7-Bromo-1-vinyl- β -carboline, V-139
 Bronkaid, A-152
 Broussonetine A, B-367
 Broussonetine B, B-367
 Broussonetine C, B-367
 Broussonetine D, B-367
 Broussonetine E, B-367
 Broussonetine F, B-367
 Broussonetine G, B-360
 Broussonetine H, B-361
 Broussonetine I, B-359
 Broussonetine J, B-359
 Broussonetine J₁, B-359
 Broussonetine J₂, B-359
 Broussonetine K, B-367
 Broussonetine L, B-367
 Broussonetine M, B-367
 Broussonetine M₁, B-367
 Broussonetine N, B-362
 Broussonetine O, B-367
 Broussonetine P, B-367
 Broussonetine Q, B-367
 Broussonetine R, B-363
 Broussonetine S, B-367
 Broussonetine T, B-367
 Broussonetine U, B-364
 Broussonetine U₁, B-367
 Broussonetine V, B-367
 Broussonetine W, B-365
 Broussonetine X, B-365
 Broussonetine Y, B-367
 Broussonetine Z, B-366
 Broussonetine, B-358
 Broussonetine A, B-367
 Broussonetine B, B-367
 Broussonetine C, Z-7
 Brovon, P-79
 Brownine, M-309
 Bruceanthinoside, H-442
 Bruceolline A, H-446
 Bruceolline B, H-442
 Bruceolline C, T-530
 Bruceolline D, B-368
 Bruceolline E, B-368
 Bruceolline F, T-25
 Bruceolline G, H-446
 Brucine *N*-oxide, S-589
 Brucine, S-589
 Brugine, B-369
 Bruneomycin, S-569
 Brunfelsamidine, P-943
 Brunfelsine, B-370
 Brunnein A, T-206
 Brunnein B, T-206
 Brunnein C, T-224
 Brunonine, B-371
 Brunsbelline, A-683
 Brunsdonnine, B-372
 Brunsvigine, P-57
 Brunsvinine, B-373
 Brussalexin A, B-374

- Bryamycin, T-392
 Bryokinin, Z-11
 BS 75A, C-574
 BS Factor, T-284
 BT 38, A-801
 BU 3557A₃, A-1315
 BU 4601A, A-789
 BU 2349A, G-131
 BU 3557A₁, K-17
 BU 3557A₂, K-18
 BU 2313A, N-264
 BU 4601B, A-789
 BU 2349B, G-132
 BU 3557B₁, K-17
 BU 3557B₂, K-17
 BU 3557B₃, K-18
 BU 2313B, N-264
 BU 620, C-711
 BU 4601C, A-789
 BU 2349C, G-133
 BU 3557C₁, K-17
 BU 3557C₂, K-19
 BU 3557C₃, K-20
 BU 3557C₄, K-20
 BU 3557C₅, K-17
 BU 3557D₁, A-1315
 BU 3557D₂, K-21
 BU 3557D₃, K-20
 BU 3557D₄, K-21
 BU 4664L, D-297
 BU 4514N, A-1140
 BU 3420T, D-959
 BU 4146T, L-7
 BU 2916T, P-573
 BU 3845T, Q-11
 BU 4704, X-7
 Bubbialidine, B-375
 Bubbialine, B-375
 Buchananine, B-376
 Buchapine, B-377
 Bucharaine, B-378
 Bucharamine, B-379
 Bucharidine, B-380
 Buchenavianine, D-662
 Buchnerine, P-673
 Buchtienine, B-381
 Buddamine, B-382
 Budelphine, H-123
 Budmunchiamine A, B-383
 Budmunchiamine B, B-384
 Budmunchiamine C, B-387
 Budmunchiamine D, B-387
 Budmunchiamine E, B-387
 Budmunchiamine F, B-383
 Budmunchiamine G, B-387
 Budmunchiamine H, B-387
 Budmunchiamine I, B-387
 Budmunchiamine K, B-389
 Budmunchiamine L1, B-385
 Budmunchiamine L2, B-386
 Budmunchiamine L3, B-383
 Budmunchiamine L4, B-385
 Budmunchiamine L4, B-387
 Budmunchiamine L5, B-388
 Budmunchiamine L5, B-389
 Budmunchiamine L6, B-389
 Buesgeniine, B-390
 Bufagin, E-132
 Buflavine, A-1361
 Bufobutanoic acid, H-755
 Bufochrome, A-923
 Bufopyramide, B-391
 Bufoserotonin A, H-755
 Bufoserotonin B, H-755
 Bufoserotonin C, B-392
 Bufotenidine *O*-sulfate, B-393
 Bufotenidine, B-393
 Bufotenine *O*-glucoside *N*-oxide, B-393
 Bufotenine *O*-glucoside, B-393
 Bufotenine *N*-oxide, B-393
 Bufotenine, B-393
 Bufothionine, D-153
 Bufotoxin, B-394
 Bufoviridine, B-393
 Bujeine, B-395
 Bukittinggine, B-396
 Bulbisine, B-261
 Bulbispermine, H-5
 Bulbocapnine β-*N*-oxide, B-397
 Bulbocapnine, B-397
 Bulbocodine, B-398
 Bulbodieine, B-399
 Bulgaramine, B-400
 Bulgarsenine, B-401
 Bulgecin A, B-402
 Bulgecin B, B-403
 Bulgecin C, B-404
 Bulgecinine, H-516
 Bullatine A, B-405
 Bullatine B, M-323
 Bullatine C, M-323
 Bullatine D, B-406
 Bullatine E, M-313
 Bullatine F, M-313
 Bullatine G, N-30
 Bulleyanaline, B-407
 Bulleyanine, B-407
 Bulleyanitine A, M-359
 Bulleyanitine B, B-408
 Bulleyanitine C, B-409
 Bungeanine, B-410
 Bungeanool, T-129
 Buntanbismine, B-411
 Buntanine, T-271
 Buntanmine A, P-616
 Buphacetine, B-412
 Buphanamine, B-413
 Buphanidine, P-581
 Buphanine, B-414
 (+)-Buphanisine, C-748
 (-)-Buphanisine, C-748
 Buphanitine, N-169
 Bupicomide, B-445
 Buprestin A, B-415
 Buprestin B, B-416
 Burmannaline, B-417
 Burmannine, B-418
 Burnamicine, B-419
 Burnamine, B-420
 Bursanine, A-139
 Buscopan, S-159
N,N'-(1,3-Butadiene-1,4-diyl)bis[2-hydroxy-3-methylbutanamide], P-375
 1,3-Butadiene-1,1,4,4-tetracarboxylic acid, B-421
 1,4-Butanediamine, B-422
 Butane-1,4-dicarboxylic acid, H-258
N,N'-1,4-Butanediylobisacetamide, B-422
N,N'-1,4-Butanediylobisalanine, S-397
N,N'-1,4-Butanediylobisbenzamide, B-422
N,N'-1,4-Butanediylobis[benzeneacetamide], B-422
N,N'''-1,4-Butanediylobisguanidine, D-400
 1,1'-(1,4-Butanediylobis[hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1*H*)-one], H-324
N,N'-1,4-Butanediylobis[3-(4-hydroxyphenyl)-2-propenamide], D-362
 2,2'-[1,4-Butanediylobis[imino(2-oxo-2,1-ethanediylobis[2-hydroxybutanedioic acid], R-80
N,N'-1,4-Butanediylobis[*N*-methylbenzamide], B-422
 Butanoyllupinine, O-58
N-Butanoylpyrrothine, A-759
 2-Butene-1,4-diamine, B-423
 (*E*)-2-Butenedioic acid, F-173
 3-*O*-(2*E*-Butenyl)-11-*O*-(3-hydroxybutanoyl)hamayne, H-5
 1'-(3-Butenyl)-[2,3'-bipiperidine]-5'-carboxylic acid, V-150
 3-(2-Butenyl)-4-ethoxy-8-hydroxy-1-methyl-2(1*H*)-quinolinone, G-124
 5-(2-Butenylidene)-3-ethyl-1,2,3,4,5,7*a*-hexahydro-4*aH*-1-pyridine-4,4*a*-diol, B-424
 3-(3-Butenyl)-1,2,3,4,5,6,11,11*a*-octahydro-1,5-methano-10*H*-pyrido[1,2-*a*][1,5]diazocin-10-one, A-554
 8-(3-Butenyl)octahydro-5-(2-penten-4-ynyl)indolizine, A-511
 8-(3-Butenyl)octahydro-5-(4-pentenyl)indolizine, A-511
 5-(3-Butenyl)picolinic acid, B-425
 1-(3-Butenyl)-5-(2-piperidyl)-3-piperidinecarboxylic acid, V-150
 5-(3-Butenyl)-2-pyridinecarboxylic acid, B-425
 4-(1-Butenyl)-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, I-338
 5-(1-Butenyl)-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, T-595
 5-(1-Butenyl)-3,4,6,7-tetrahydro-3-methyl-2*H*-cyclopenta[*b*]pyridine, P-932
 4-(1-Butenyl)-1,6,7,8-tetrahydro-5,6,8-trimethylcyclopent[*g*]indole, H-142
 1-Butoxycarbonyl-β-carboline, C-132
 3-(Butoxymethyl)-6,7-dihydro-8(5*H*)-indolizinone, D-447
 Butoprine, M-393
tert-Butyl *N*-(2-aminoethyl)carbamate, E-225
 Butyl *N*-(2-phenylethyl)carbamate, P-343
 2-Butyl-1-azacyclohexene, B-446
 7-Butyl-2-chloro-6,8-dihydroxy-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, B-429
 4(5)-Butyl-2-chloro-5(4)-formyl-1*H*-imidazole, B-426
 5(4)-Butyl-2-chloro-1*H*-imidazole-4(5)-carboxaldehyde, B-426
 Butylcycloheptylprodigiosine, B-427
 2-Butyldecahydroquinoline, B-428
 1β-*C*-Butyl-1-deoxygalactonojirimycin, B-432
N-Butyldeoxynojirimycin, B-616
 7-Butyl-1,2-dichloro-6,8-dihydroxy-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, B-429
 5-Butyl-6,8-diethylindolizidine, B-430
 5-Butyl-6,8-diethyloctahydroindolizine, B-430
 7-Butyl-5,10-dihydro-4,6,9-trihydroxy-8-methoxy-5,10-dioxobenzo[*g*]quinoline-2-carboxylic acid, I-264
 2-*sec*-Butyl-3,4-dihydroxy-5-hydroxymethylpyridine, A-620
 3-Butyl-2,5-dimethylpyrazine, B-431
 Butyl-DNJ, H-616
N,N'-Butylenebis(cantharimide), P-650
 4-Butyl-*N*-feruloylphenethylamine, B-441
 2-Butyl-5-heptyl-3,4-dihydro-2*H*-pyrrole, D-246
 2-Butyl-5-heptyl-1-methylpyrrolidine, D-246
 5-Butyl-2-heptyl-Δ¹-pyrroline, D-246
 2-Butyl-6-hydroxymethyl-3,4,5-piperidinetriol, B-432
 6-*tert*-Butyl-1-hydroxy-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, B-433
 3-Butyl-8-hydroxy-5-propylindolizidine, B-438
 2-Butyl-5-(3-indolyl)oxazole, A-622
 3-Butyllehmizidine, B-439
 2-*sec*-Butyl-3-methoxy-pyrazine, M-272
 3-Butyl-5-methylindolizidine, B-434
N-Butyl-*O*-methylachanthopyridone, L-5
 4-(4-Butyl-3-methyl-5-oxo-1-imidazolidinyl)-7-(1-methylethyl)-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraene-5,8-dione, S-93
 2-Butyl-1-methylpiperidine, B-442
 3-Butyl-2-methylprodigiosine, P-636
 2-Butyl-5-(8-nonenyl)pyrrolidine, D-246
 3-Butyloctahydro-8-hydroxy-5-propylindolizine, B-438
 1-(3-Butyloctahydro-5-indolizinyloxy)-1-propanone, B-436
 3-Butyloctahydro-5-methylindolizine, B-434
 3-Butyloctahydro-5-(4-pentenyl)indolizine, B-435
 3-Butyloctahydro-5-propanoylindolizine, B-436
 8-Butyloctahydro-5-propyl-1*H*-indolizine, B-437
 3-Butyloctahydro-5-propyl-8-indolizino[1,2-*a*]zepine, B-439
 4-*tert*-Butyl-2-oxazolindione, B-440
 3-Butyl-5-(4-pentenyl)indolizidine, B-435
N-[2-(4-Butylphenyl)ethyl]-3-(4-hydroxy-3-methoxyphenyl)-2-propenamide, B-441
 5-Butylpicolinic acid, B-445
 2-Butylpiperidine, B-442
 3-Butyl-5-propanoylindolizidine, B-436
 5-Butyl-8-propylindolizidine, B-443
 8-Butyl-5-propylindolizidine, B-437

- 3-Butyl-5-propylindolizidine, O-41
 3-Butylpyridine, B-444
 5-Butyl-2-pyridinecarboxylic acid, B-445
 Butylscopolammonium bromide, S-159
 2-Butyl-3,4,5,6-tetrahydropyridine, B-446
 2-*sec*-Butyl-2-thiazoline, D-475
 2-Butyl-3,4,5-trihydroxy-6-hydroxymethylpiperidine, B-432
 Butynediamide, B-447
 2-Butynedioic acid, B-447
 8-(3-Butynyl)octahydro-5-(2-penten-4-ynyl)indolizine, A-511
 8-(3-Butynyl)octahydro-5-(4-pentynyl)indolizine, A-511
 Butyrobetaine, A-126
 γ -Butyrobetaine, A-126
 Butyrolactam, P-956
 6-*O*-Butyrylcastanospermine, C-188
 Butyrylcholine, C-419
N-Butyryl-*N*-depropionyldecorticasine, L-234
 8-Butyryl-1,2,3,5,6,8 α -hexahydroindolizine, E-58
 8-Butyryl-7-hydroxyindolizidine, E-59
 13-Butyryloxylupanine, H-554
 3-Butyryloxytropane, M-393
 Buxabenzacine, B-460
 Buxabenzacinine, B-460
 Buxabenzamidienine, B-455
 Buxabenzamidine, B-455
 Buxadienine, B-460
 Buxafuranamide, B-448
*O*¹⁰-Buxafuranamine, B-449
*O*⁶-Buxafuranamine, B-450
 Buxahejramine, B-466
 Buxahyrcanine, B-451
 Buxakarachiamine, B-466
 Buxakashmiramine, C-888
 Buxalfine, B-452
 Buxalfol, B-452
 Buxaline C, C-902
 Buxalongifolamidine, B-453
 Buxalongifoline, B-466
 Buxalphine, B-452
 Buxaltine H, B-454
 Buxaltine, B-454
 Buxamideine K, B-457
 Buxamine A, B-455
 Buxamine B, B-455
 Buxamine C, B-455
 Buxamine E, B-455
 Buxamine F, B-455
 Buxamine G, B-455
 Buxamine I, B-455
 Buxamine, B-455
 Buxaminol A, B-456
 Buxaminol B, B-456
 Buxaminol C, B-456
 Buxaminol E, B-456
 Buxaminol G, B-466
 Buxaminol I, B-456
 Buxaminol, B-456
 Buxaminone, B-457
 Buxanaldimine, B-458
 Buxandonine L, B-459
 Buxandonine, B-459
 Buxandrine F, C-883
 Buxandrine, C-883
 Buxanine M, C-848
 Buxanine, C-848
 Buxanoldine, B-460
 Buxapapilinine, B-477
 Buxapapinolamine, B-461
 Buxaprogestine, A-891
 Buxaquamarine, B-462
 Buxarine F, C-917
 Buxarine, C-917
 Buxasamarine, C-917
 Buxatine, C-920
 Buxazidine B, C-883
 Buxazidine I, C-883
 Buxazine, B-463
 Buxbarbarine K, B-457
 Buxdeltine, B-464
 Buxene O, C-848
 Buxene, C-848
 Buxenine G, B-455
 Buxenone M, C-848
 Buxenone, C-848
 Buxepidine, C-883
 Buxeridine C, C-902
 Buxeridine, C-902
 Buxetine, B-465
 Buxidienine F, B-466
 Buxidienine I, B-466
 Buxidine F, B-12
 Buxidine F, C-883
 Buxidine, C-883
 Buxifoliadine A, T-546
 Buxifoliadine B, T-546
 Buxifoliadine C, T-580
 Buxifoliadine D, B-467
 Buxifoliadine E, B-468
 Buxifoliadine F, B-469
 Buxifoliadine G, B-470
 Buxifoliadine H, P-207
 Buxifoline, B-471
 Buxippine K, B-482
 Buxiramine D, B-472
 Buxiramine, B-472
 Buxirane, B-460
 Buxisine, B-474
 Buxithienine M, C-848
 Buxitrienine C, B-473
 Buxizine, C-883
 Buximicrophylline A, C-917
 Buximicrophylline B, C-848
 Buximicrophylline C, C-883
 Buximicrophylline D, C-881
 Buxocyclamine A, B-474
 Buxocyclamine I, B-474
 Buxodienine E, B-456
 Buxodienine I, B-456
 Buxomegine, A-886
 Buxomeline, B-475
 Buxotrienine, B-476
 Buxoxybenzamine, B-477
 Buxozine C, B-478
 Buxpapamine, B-479
 Buxpapine, B-480
 Buxpapinine, B-481
 Buxpiine K, B-482
 Buxpiine, B-482
 Buxpsiine K, B-457
 Buxpsiine, B-457
 Buxtaune M, B-482
 Buxtaune, B-482
 Buxupapine, B-483
 Buzonamine, B-484
 C 73, A-129
 C 14482A₁, D-876
 C 304A, H-408
 C 14482B₁, D-877
 C 5507, C-954
 C 15003 deClQNDQ, M-136
 C 15003 deClQO, M-136
 C 15003 P 4 β HY, M-136
 C 15003 P 4 γ HY, M-136
 C 9154, P-337
 C 15003 PHM1, M-136
 C 15003 PHM2, M-136
 C 15003 PHM3, M-136
 C 15003 PHM4, M-136
 C 15003 PHO 0, M-136
 C 15003 PHO 1, M-136
 C 15003 PHO 2, M-136
 C 15003 PHO 3, M-136
 C 15003 PHO 4, M-136
 C 15003P0, M-136
 C 15003P1, M-136
 C 15003P2, M-136
 C 15003P3', M-136
 C 15003P3, M-136
 C 15003P4, M-136
 C 15003 PND 4 β HY, M-136
 C 15003 PND 1, M-136
 C 15003 PND 2, M-136
 C 15003 PND 3, M-136
 C 15003 PND 4, M-136
 C 73X, A-129
 CA 39A, S-500
 CA 39B, S-500
 (+)-Caaverine, D-531
 (-)-Caaverine, D-531
 Cabbage identification factor 1, D-517
 Caberine, C-191
 Caberoline, R-21
 Caboxine A, C-1
 Cabuamine, D-410
 Cabucine, A-1423
 Cabucinine, R-28
 Cabucraline 10-carboxaldehyde, C-191
 Cabucraline *N*⁴-oxide, C-191
 Cabucraline, C-191
 Cabudine, C-2
 Cabufile, C-3
 Cactine, H-357
 Cadabicine diacetate, C-4
 Cadabicine methyl ether, C-4
 Cadabicine, C-4
 Cadabine, S-491
 Cadambine acid, C-5
 Cadambine, C-5
 Cadamine, C-6
 Cadaverine, P-229
 Cadeguomycin, C-7
 Cadiaine, H-554
 Cadiamine hydroxyphenylacetate, C-8
 Cadiamine 2-pyrrolicarboxylate, C-8
 Cadiamine, C-8
 Caeruleine, M-379
 Caerulomycin A, C-9
 Caerulomycin B, C-9
 Caerulomycin C, C-9
 Caerulomycin D, C-10
 Caerulomycin E, C-11
 Caerulomycin, C-9
 Caerunine, M-361
 Caesalpinine A, C-12
 Cafamycin, C-13
 Cafcit, C-15
 Caffaeschizine, C-14
 Caffedyimine, D-920
 Caffaic acid, D-653
 Caffeine citrate, C-15
 Caffeine, C-15
N-Caffeoyl DOPA, C-534
*N*⁵-Caffeoyl-*N*¹-*p*-coumaroyl-*N*¹⁰-feruloylspermidine, S-398
*N*¹⁰-Caffeoyl-*N*¹,*N*⁵-di-*p*-coumaroylspermidine, S-398
*N*⁵-Caffeoyl-*N*¹,*N*¹⁰-di-*p*-coumaroylspermidine, S-398
*N*¹-Caffeoyl-*N*¹⁰-feruloylspermidine, S-398
 Caffeoylferuloylspermidine, S-398
N-*trans*-Caffeoyl-*O*-methyltyramine, M-280
 1-[Caffeoyloxy(carboxy)methoxy]-1*H*-pyrrole-2,3,5-tricarboxylic acid, C-16
 4'-*O*-Caffeoylprunasin, H-673
 Caffeoylputrescine, C-456
 Caffeoylspermidine, S-398
N-Caffeoyltryptophan, T-640
N-*cis*-Caffeoyltyramine, H-667
N-*trans*-Caffeoyltyramine, H-667
 3'-Caffeoylvincoside lactam, V-126
 Caffipel, C-15
 Cairomycin C, C-17
 Caissarine A, P-813
 Caissarine B, C-18
 Caissarine C, A-158
 Caissarone, C-19
 Cajanifoline, D-605
 Calabacine, C-20
 Calabatine, C-21
 Calafatimine, C-22
 Calafatine 2' α -*N*-oxide, C-23
 Calafatine 2' β -*N*-oxide, C-23
 Calafatine, C-23
 Calafianine, C-24

- Calanthoside, H-534
 Calcaridine A, C-25
 Calcimycin, C-26
 Calcium folinate, F-121
 Calcium levofolinate, F-121
 Caldaphnidine A, C-27
 Caldaphnidine B, P-137
 Caldaphnidine C, C-28
 Caldaphnidine D, C-29
 Caldaphnidine E, C-30
 Caldaphnidine F, D-37
 Caldensine, D-567
 Caldine, D-255
 Caldoheptamine, P-199
 Caldohexamine, T-107
 Caldopentamine, T-456
 C-Calebassine A, C-31
 C-Calebassine, C-31
 Calebassine, C-31
 Calebassinine I, C-32
 Caledonin, C-33
 Caledonine, Q-1
 Californidine, E-221
 Californine, E-221
 Calimycin, C-26
 Calipamine, M-18
 Calliactine, C-34
 Callichiline, C-35
 Calligonidine, T-202
 Calligonine, T-202
 Callimorphine, C-36
 Callipeltoside A, C-37
 Callophylline A, U-22
 Callophylline B, C-38
 Callosine, C-39
 Callyspongamide A, H-117
 Calophylline, U-22
 Calothrixin A, I-107
 Calothrixin B, I-107
 Calpaurine, T-557
 Calpurmenine, C-40
 Calpurnine, H-554
 Calvine, C-41
 Calycanthidine, C-382
 Calycanthine, C-42
 Calycilactone A, C-43
 Calycine, C-44
 Calycinine A, D-33
 Calycinine, D-846
 Calyciphylline A, D-55
 Calyciphylline B, C-45
 Calyciphylline C, C-46
 Calyciphylline D, C-47
 Calyciphylline E, C-48
 Calyciphylline F, C-49
 Calyciphylline G, C-50
 Calyciphylline H, C-51
 Calyciphylline I, C-51
 Calyciphylline J, C-52
 Calyciphylline K, D-34
 Calyciphylline L, H-330
 Calyciphylline M, D-24
 Calycosidine, C-53
 Calycotomine, C-54
 Calyculin A, C-55
 Calyculin B, C-55
 Calyculin C, C-56
 Calyculin D, C-56
 Calyculin E, C-55
 Calyculin F, C-55
 Calyculin G, C-56
 Calyculin H, C-56
 Calyculin J, C-57
 Calyculinamide A, C-55
 Calyculinamide B, C-55
 Calyculinamide F, C-55
 Calydaphninone, C-58
 Calystegine A₃, A-1590
 Calystegine A₅, A-1593
 Calystegine A₆, A-1592
 Calystegine A₇, A-1591
 Calystegine B₁, A-1588
 Calystegine B₂, A-1587
 Calystegine B₃, A-1587
 Calystegine B₄, A-1587
 Calystegine B₅, A-1589
 Calystegine C₁, A-1586
 Calystegine C₂, A-1586
 Calystegine N₁, A-704
 Calystegine, C-59
 Calyxamine A, C-60
 Calyxamine B, C-60
 Calyxinin, D-292
 Calyxoside, D-292
 Camalexin, T-372
 Cammaconine, M-352
 Camoensidine *N*-oxide, C-61
 Camoensidine, C-61
 Camoensine, C-62
 Campedine, C-63
 Campestrine, C-64
 Camphorataimide B, H-684
 Camphorataimide C, H-684
 Camphorataimide D, H-684
 Camphorataimide E, H-684
 Camptacumanine, C-65
 Camptacumotine, C-66
 Camptine, C-67
 Camptinine, C-68
 Camptoneurine, C-69
 Camptothecin, C-70
 Camptothecin acid, C-71
 Campylocine, G-152
 Campylotheine, M-309
 Camtschatcanidine, S-345
 Canadalin, C-72
 α -Canadine, C-73
 Canadine, C-73
 β -Canadine, C-73
 Canadinic acid, C-72
 Canangine, I-65
 Cananodine, G-190
 Canavalmine, D-251
 Canavanine, C-74
 Canavaninosuccinic acid, C-75
 Canavanosuccinic acid, C-75
 Cancentrine, C-76
 Candicine, H-357
 Candicusine, L-175
 Candidine, C-77
 Candimine, C-78
 Caneine, C-79
 Canelilline, A-1038
 Canelillinoxine, C-80
 Canembine, C-81
 Canescenine, E-12
 Canescensterone, H-244
 Canescic acid, D-687
 Canescine, D-234
 Canescine, E-12
 Canforemetina, E-72
 Cangorin A, H-119
 Cangorin B, H-119
 Cangorin C, H-119
 Cangorin D, H-119
 Cangorin E, H-119
 Cangorin F, H-119
 Cangorin G, H-119
 Cangorin H, H-119
 Cangorin I, H-119
 Cangorin J, H-248
 Cangorinine EI, P-247
 Cangorinine WII, P-252
 Cangorinine WI, P-252
 Cannabiloid B, C-82
 Cannabisativine, C-83
 Cannabisin A, C-84
 Cannabisin B, C-85
 Cannabisin C, C-85
 Cannabisin D, C-85
 Cannabisin E, C-86
 Cannabisin F, C-86
 Cannabisin G, C-87
 Cannabisin H, C-88
 Cantabricine, M-108
 Cantharimide, C-89
 Canthin-2,6-dione, I-105
 Canthin-6-one *N*-oxide, C-90
 Canthin-6-one, C-90
 Canthiumine, C-91
 Cantleyine, C-92
 Cantuzumab mertansine, M-135
 Capauridine, C-93
 Capaurimine, C-93
 Capaurine, C-93
 Capitavine, C-94
 Capitelline, C-95
 Capnoidine, A-145
 Capnosine, S-234
 Capnosinine, S-234
 Cappamensine, C-96
 Capparidisine, C-97
 Cappariloside A, H-524
 Cappariloside B, H-524
 Capparine A, S-426
 Capparine B, M-201
 Capparisine, C-97
 Capparisinine, C-97
 Caprazamycin, C-98
 Capreoline, F-176
 Capreomycidine, C-99
 Capric acid, D-115
 Caprinic acid, D-115
 Caprolactin A, A-785
 Caprolactin B, A-785
 13-Caproyloxylupanine, H-554
 Capsacutin, C-100
 Capsaicin glucoside, C-100
 Capsaicin, C-100
 Capsaicinol, C-101
 Capsaiene, C-100
 Capsiamide, A-30
 Capsicastrine, S-200
 Capsimine 3-glucoside, S-200
 Capsimine, S-200
 Capsimycin, C-102
 Captax, B-79
 Capuramycin, C-103
 Capuronidine, D-377
 Capurone, C-104
 Capuvosidine, C-105
 Capuvosine, C-106
 Caracasanamide, C-107
 Caracasan diamide, C-108
 Caracurine I, C-109
 Caracurine II dimethosalt, C-111
 Caracurine II, C-111
 Caracurine III, C-112
 Caracurine IV, C-113
 Caracurine IX, C-114
 Caracurine V dimethosalt, C-110
 Caracurine V di-*N*-oxide, C-110
 Caracurine V *N*-oxide, C-110
 Caracurine V, C-110
 Caracurine VII, C-115
 Caracurine VIII, C-116
 Caracurine VI, T-436
 Caranine, F-7
 Carapanaubine *N*-oxide, C-117
 Carapanaubine, C-117
 Carategine, C-118
 Carazostatin, H-132
 Carbalexin A, D-616
 Carbalexin B, D-618
 Carbalexin C, D-617
 Carbamamidine, G-199
 Carbamic acid nitrile, C-823
 Carbamidine, G-199
 Carbamite, D-818
 Carbamonitrile, C-823
N-Carbamoylanonaine, R-120
N-Carbamoylasimilobine, D-531
 1-Carbamoyl- β -carboline, C-132
 1-*O*-Carbamoyl-3-deoxy-3-guanidino-*scytlo*-inositol, B-237
N'-(2-Carbamoylethyl)-4-[(2-guanidinoacetimidoyl)amino]-1,1'-dimethyl-*N*,4'-bipyrrrole-2-carboxamide], N-180
*N*²-Carbamoylglutamine amide, G-108

- 2-Carbamoyl-3-methoxy-1,4-naphthoquinone, H-638
N-Carbamoyl-2-methoxypyrrolidine, P-955
 4-(Carbamoylmethyl)-4-hydroxy-2,5-cyclohexadien-1-one, H-650
 2-Carbamoyl-4-methylquinazoline, M-555
 (Carbamoylmethyl)trimethylammonium(1+), C-119
N-Carbamoylputrescine, A-720
N-Carbamoylpyrrolidine, P-951
 1-Carbamoyl-2-pyrrolidone, S-487
N-Carbamoylserotonin, H-755
N-Carbamoylstaurosporine, S-498
N-Carbamoylstepharine, P-643
 6-Carbamoyluracil, D-669
 Carbanilide, D-818
 Carbazole, C-120
 9*H*-Carbazole-3-carboxaldehyde, C-121
 2-Carbazolecarboxylic acid (obsol.), C-122
 9*H*-Carbazole-3-carboxylic acid, C-122
 9*H*-Carbazole-2,7-diol, C-123
 9*H*-Carbazole-3-methanol, C-124
 Carbazoline, C-130
 Carbazomadurin A, C-125
 Carbazomadurin B, C-126
 Carbazomarin A, C-127
 Carbazomycin A, D-571
 Carbazomycin B, D-571
 Carbazomycin C, T-543
 Carbazomycin D, T-543
 Carbazomycin E, H-561
 Carbazomycin F, H-561
 Carbazomycin G, C-128
 Carbazomycin H, C-128
 Carbazomycinal, H-561
 Carbazochinocin, C-129
 2-Carbothoxycarbazole, C-122
 1-Carbothoxy- β -carboline, C-132
 3-Carbothoxyharman, M-427
 9-Carbothoxy-3-methyl-9*H*-carbazole, M-424
 Carbimide, C-823
 Carbinamine, M-383
 1-Carbobutoxy- β -carboline, C-132
 Carbodiimide, C-823
 N^b -Carboethoxy-4',5',6',17-tetrahydro-17*R*-usambarensine, U-52
 Carbohydroquinonic acid, D-536
 β -Carboline, C-130
 2-Carboline, C-130
 β -Carboline-1-carboxaldehyde, C-131
 β -Carboline-1-carboxamide, C-132
 β -Carboline-1-carboxylic acid, C-132
 β -Carboline-3-carboxylic acid, C-133
 β -Carboline-1-propanoic acid, C-134
 β -Carboline-1,3,4-trione, C-135
 β -Carbolin-1-yl 4,8-dimethoxy- β -carbolin-1-ylethyl ketone, P-406
 β -Carbolin-1-yl 3-(4,8-dimethoxy- β -carbolin-1-yl)-1-methoxypropyl ketone, P-408
 2-(β -Carbolin-1-yl)-5-hydroxymethylfuran, P-270
 1-(β -Carbolin-1-yl)-3-hydroxy-1-propanone, C-136
 4-(β -Carbolin-1-yl)-4-oxobutanoic acid, C-137
 1-(β -Carbolin-1-yl)-3,4,5-trihydroxy-1-pentanone, C-138
 2-Carbomethoxycarbazole, C-122
 1-Carbomethoxy- β -carboline, C-132
 16-Carbomethoxy-18,19-dihydronaufoline, N-63
 3-Carbomethoxy-1,7-dihydroxy-9*H*-carbazole, D-559
 1-Carbomethoxy-4,8-dimethoxy- β -carboline, D-561
 3-Carbomethoxyharman, M-427
 16-Carbomethoxy-16-hydroxy-14,15-epoxy-3-oxo-1,2-dehydroaspido-permidine, M-297
N-Carbomethoxy-17-hydroxycopsinine, K-78
N-Carbomethoxy-17 β -hydroxy- $\Delta^{14,15}$ -copsinine, V-49
 3-Carbomethoxy-1-hydroxy-6-methoxycarbazole, D-558
 Carbomethoxyibogamine, C-657
 3-Carbomethoxy-7-methoxycarbazole, H-455
 16-Carbomethoxyaufoline, N-63
 16-Carbomethoxyquebrachamine, V-112
 N^b -Carbomethoxy-4',5',6',17-tetrahydro-17*R*-usambarensine, U-52
 Carbonarin C, C-139
 Carbonarin D, C-140
 Carbonarin E, C-141
 Carbonarone A, B-92
 Carbonarone B, C-142
 Carbonic acid diureide, T-627
N,N'-Carbonimidoylbis[2-ethylhexanamide], L-174
 2,2'-(Carbonimidoyldi-4,1-phenylene)bis-3-oxetanamine, B-277
 3,3'-Carbonylbiscytisine, A-1412
 1,1'-Carbonylbis[3-ethylpiperidine], S-581
 3,3'-Carbonylbis[1,2,3,4,5,6-hexahydro-1,5-methano-2*H*-pyrido[1,2-*a*][1,5]diazocin-5-one], A-1412
 6,6'-[Carbonylbis(imino-1*H*-indole-5,2-dylcarbonyl)]bis[8-(chloromethyl)-3,6,7,8-tetrahydro-1-methylbenzo[1,2-*b*:4,3-*b'*]dipyrrol-4-ol], B-229
 [Carbonylbis(imino(6-methyl-3,1-phenylene))]bis-carbamic acid, B-164
N,N'-Carbonylbis-3-veratrolyxynortropane, S-622
 Carbonyldiurea, T-627
 Carbostyryl-4-carboxylic acid, H-723
 Carbostyryl, Q-50
N-Carboxamidostaurosporine, S-498
N-Carboxamidostepharine, P-643
 2-[(Carboxyacetyl)amino]benzoic acid, A-706
 2-Carboxy-4-acetylamino-piperidine, A-871
 5-Carboxyamino-2,4-imidazolidinedione, D-804
 11-Carboxyanatoxin a, C-143
 1-(*o*-Carboxyanilino)-1-deoxyfructose, F-161
 11-(4-Carboxybutyramido)-11-deoxy-1-epilupinine methyl ester, O-59
 2-Carboxy-4-(5-carboxy-1-methyl-1,3-hexadienyl)-3-pyrrolidineacetic acid, D-912
 2-Carboxy-4-(5-carboxy-1-methyl-1,4-hexadienyl)-3-pyrrolidineacetic acid, I-228
 2-Carboxy-4-(5-carboxy-1-methyl-2-hexenylidene)-3-pyrrolidineacetic acid, I-229
 2-Carboxy-4-(5-carboxy-1-methyl-1,3-pentadienyl)pyrrolidineacetic acid, N-297
 3-Carboxy-1-(carboxymethyl)pyridinium betaine, C-144
 5-[5-Carboxy-4-(carboxymethyl)-3-pyrrolidinyl]-1,6-dihydro-6-oxo-2-pyridinecarboxylic acid, A-118
 3-[5-Carboxy-4-(carboxymethyl)-3-pyrrolidinyl]-1,6-dihydro-6-oxo-2-pyridinecarboxylic acid, A-119
 4-[[[1-Carboxy-2-(6-carboxy-2-oxo-2*H*-pyran-4-yl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, M-769
 5-Carboxycorynanthine, C-145
 7-Carboxy-7-deazaquanosine, C-7
 16-Carboxy-3,4,14,15,16,17,18,19,20,21-decadehydrohimbantium methyl ester, O-140
 2-Carboxy-1-[(2,6-dicarboxy-2,3-dihydro-4(1*H*)-pyridinylidene)ethylidene]-2,3-dihydro-5,6-dihydroxy-1*H*-indolium hydroxide inner salt, B-110
 5-Carboxy-3,4-didehydrostrictosidine, C-153
 2-Carboxy-3,4-dihydro- α -hydroxy-5-(2-pyridinyl)-2*H*-pyrrole-4-acetic acid, P-637
 α -Carboxy-2,3-dihydro- β -hydroxy-*N,N,N*-trimethyl-2-thioxo-1*H*-imidazole-4-ethanaminium hydroxide inner salt, H-484
 4-[[[Carboxy(2,3-dihydro-3-oxo-5-isoxazolyl)-methyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, M-768
 α -Carboxy-2,3-dihydro-*N,N,N*-trimethyl-2-thioxo-1*H*-imidazole-4-ethanaminium hydroxide inner salt, E-173
 4-[[[1-Carboxy-2-(3,4-dihydroxyphenyl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, D-921
 α -Carboxy-3,4-dihydro-*N,N,N*-trimethylbenzene-ethanaminium(1+), A-745
 4-Carboxy-1,3-dimethyl-1*H*-imidazolium hydroxide inner salt, N-318
 2-Carboxy-1,1-dimethylpiperidinium hydroxide inner salt, H-348
 3-Carboxy-1,1-dimethylpyrrolidinium hydroxide inner salt, P-952
 2-Carboxy-1,1-dimethylpyrrolidinium hydroxide inner salt, S-491
 5-(2-Carboxy-1-ethenyl)-4-methoxyfuro[2,3-*b*]pyridine-6-carboxylic acid, R-88
 58-(1-Carboxyethenyl)sulfomycin I, P-640
 N^2 -(2-Carboxyethyl)arginine, A-1413
 N^{16} -Carboxyethylidihydroroquefortine C, R-125
 3-(2-Carboxyethyl)-1,4-dimethylpyridinium, C-146
 1-(1-Carboxyethyl)-3,4-dimethyl-1*H*-pyrrole, D-780
N-(Carboxyethyl)glycine, C-147
 2-Carboxy-3-ethyl-12*H*-indolo[2,3-*a*]quinolizin-5-ium inner salt, F-82
 2-(2-Carboxyethyl)-3-isoxazolin-5-one, O-185
N-(1-Carboxyethyl)mescaline, M-219
 1-(1-Carboxyethyl)-2-methylpropyl 3-[[5-[(aminomethyl)amino]-1-oxopentyl]oxy]-2,4-dimethylpentanoate, A-1279
 7-Carboxyevodiamine, E-310
 16-Carboxyherbavine, M-50
 1-Carboxyhexahydro-4-(3-methyl-2-butenyl)-1*H*-pyrrolizinium, A-1065
 1-Carboxy-3-[(4-hydroxybenzoyl)oxy]-*N,N,N*-trimethyl-2-propanaminium inner salt, A-792
 2-Carboxy-3-hydroxy-1,1-dimethylpyrrolidinium hydroxide inner salt, H-740
 2-Carboxy-4-hydroxy-1,1-dimethylpyrrolidinium hydroxide inner salt, H-476
 3-Carboxy-1-(2-hydroxyethyl)pyridinium betaine, P-897
 4-[(2-Carboxy-2-hydroxyethyl)thio]-2-piperidinecarboxylic acid, P-787
 2-Carboxy-4-(1-hydroxy-1-methylethyl)-3-pyrrolidineacetic acid, K-4
 2-Carboxy-5-hydroxy-6-methylpiperidine, H-613
 N^2 -(3-Carboxy-2-hydroxy-1-oxopropyl)arginine, A-1413
 4-[[[Carboxy-(4-hydroxyphenyl)ethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, P-576
 2-Carboxy-3-hydroxypiperidine, H-696
 α -Carboxy-4-hydroxy-*N,N,N*-trimethylbenzene-ethanaminium(1+), T-700
 3-Carboxyhyrtiomanzamine, D-933
 16-Carboxyisoherbavine, M-50
 4-(Carboxymethoxymethyl)-1,3-dimethyl-1*H*-imidazolium hydroxide inner salt, E-15
 17-Carboxy-12-methoxy- N^b -methylechitamide, E-30
 Carboxymethyl 5,10-dihydro-1-phenazinecarboxylate, P-319
N-(Carboxymethyl)- β -alanine, C-147
 3-(Carboxymethylamino)propanoic acid, C-147
 3-(Carboxymethylamino)propionitrile, C-147
 N^2 -(Carboxymethyl)arginine, A-38
 3-Carboxy-2-methyl- β -carbolinium, C-133
 8-Carboxymethylidihydrochelerythrine, D-408
 4-(Carboxymethyl)-1,3-dimethyl-1*H*-imidazolium hydroxide inner salt, Z-32
 17-Carboxy- N^b -methylechitamide, E-30
 2-Carboxy-4-(1-methylethenyl)-3-pyrrolidineacetic acid, K-4
 (Carboxymethyl)guanidine, G-200
 N^2 -(2-Carboxymethyl-2-hydroxysuccinoyl)arginine, A-1413
 4-[[[Carboxymethyl]imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, P-577
 1-(Carboxymethyl)-2-iminoimidazolidine, A-731
 1-(Carboxymethyl)-2-imino-3-phosphonoimidazolidine, A-731
 3-Carboxymethyl-4-isopropenylproline, K-4
 2-Carboxy-1-methylisoquinolinium hydroxide inner salt, Q-31
 2-Carboxymethyl-3-isoxazolin-5-one, O-184
N-(Carboxymethyl)mescaline, M-219
 1-Carboxymethyl-4-methoxy-2-(8-nonyl)pyridinium(1+), M-699
 1-Carboxymethylnicotinic acid, C-144
 1-(1-Carboxy-2-methylpropyl)-3,4-dimethyl-1*H*-pyrrole, D-779
 2-Carboxy-1-methylpyridinium betaine, C-148
 3-Carboxy-1-methylpyridinium betaine, C-149

- 3-Carboxy-1-methylpyridinium hydroxide inner salt, C-149
- 2-Carboxy-1-methylpyridinium hydroxide inner salt, C-148
- 3-Carboxy-2-methyl-9*H*-pyrido[3,4-*b*]indolinium inner salt, C-133
- 2-Carboxy-1-methylquinolinium betaine, Q-31
- 4-[[*N*-[1-Carboxy-3-(methylsulfinyl)propyl]formimido]methylene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, M-645
- 4-[[[1-Carboxy-3-(methylsulfinyl)propyl]imino]ethylene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, M-645
- N*-(Carboxymethyl)tryptophan, K-62
- 12-Carboxymonascorubramine, C-150
- Carboxymycobactins, C-151
- 3-(2-Carboxy-5-oxo-1-piperidinyl)thioalanylglycine, C-831
- 3-(6-Carboxy-2-oxo-4-pyridinyl)alanine, A-722
- 2-(5-Carboxypentyl)-4-thiazolidone, A-78
- 3-(2-Carboxyphenyl)-4(3*H*)-quinazolinone, Q-14
- 4-Carboxyproline, P-953
- 5-[(3-Carboxypropyl)amino]-2-hydroxy-3-tridecyl-1,4-benzoquinone, C-152
- N*-(3-Carboxypropyl)-*N*-(3-methyl-2-butenyl)guanidine, C-592
- 5-Carboxy-2-(2-pyridyl)-1-pyrroline-3-glycolic acid, P-637
- 4-[2-(2-Carboxy-1-pyrrolidinyl)ethenyl]-2,3-dihydro-2,6-pyridinedicarboxylic acid, I-67
- 3-(2-Carboxy-4-pyrrolyl)alanine, A-721
- 5 α -Carboxystrictosamide, T-150
- 5-Carboxystrictosidine, C-153
- 5-Carboxytetrahydroalstonine, C-154
- 3-Carboxy-1,2,3,4-tetrahydro- β -carboline-1-propanoic acid, C-155
- 3-Carboxy-1,2,3,5-tetrahydro-8-hydroxy-5-oxo-6-indolizinepropanoic acid, A-1101
- 4-Carboxy-1,3,4,6-tetrahydro-9-hydroxy-6-oxo-2*H*-quinolizine-7-propanoic acid, A-1104
- 6-Carboxy-1,2,3,4-tetrahydro-7-phenylquinolizinium inner salt, C-496
- 1-*C*-(3-Carboxy-2,3,4,9-tetrahydro-1*H*-pyrido[3,4-*b*]indol-1-yl)pentitol, T-215
- 4-Carboxy-2-thiazolidinethione, T-394
- α -Carboxy-*N,N,N*-trimethyl-1*H*-indole-3-ethanaminium hydroxide inner salt, H-764
- 3-Carboxy-*N,N,N*-trimethyl-1-propanaminium hydroxide inner salt, A-126
- 1-Carboxytryptargine, C-156
- 5 α -Carboxyvincosamide, T-150
- 3-Carboxy-1-vinyl- β -carboline, V-140
- 3-(1-Carboxyvinyl)oxybenzoic acid, D-224
- Carcinine, C-157
- Carcisine A, C-756
- Carcisine B, C-757
- Cardinalic acid, D-677
- Cardiochrysin, C-158
- Cardiodine, H-169
- Cardionidine, C-159
- Cardionine, H-192
- Cardiopetalidine, M-353
- Cardiopetaline, M-356
- Cardiopetamine, H-190
- Cardiopidine, H-174
- Cardiopimine, H-174
- Cardiopine, H-174
- Cardiopinine, H-174
- Cardiospermin 5-*p*-coumarate, D-622
- Cardiospermin 5-(4-hydroxybenzoate), D-622
- Cardiospermin 5-sulfate, D-622
- Cardiospermin, D-622
- Cardiosteril, D-920
- Carduchorone, T-317
- Caribaeolin, C-161
- Caribaeorane, C-160
- Caribaeoside, C-161
- Caribine, C-162
- Carinatine, P-716
- Carinatamine A, H-375
- Carinatamine B, H-376
- Carinatamine C, H-380
- Carmichaeline, M-355
- Carnavaline, C-183
- Carnegine, T-167
- Carnine, I-148
- Carnosadine, C-163
- Carococculine, C-164
- Carolinianine, L-321
- Carolisterol A, C-165
- Carolisterol B, C-165
- Carolisterol C, C-165
- Carosidine, C-166
- Carosine, C-167
- Carpacil, R-52
- ψ -Carpaine, C-168
- Carpaine, C-168
- Carpidine, P-428
- Carpiline, P-428
- Carpoxidine, O-94
- Carquinostatin A, C-169
- Carquinostatin B, C-169
- Cartermine A, S-611
- Cartormin, C-170
- Caryachine *N*-methosalt, E-220
- Caryachine *N*-oxide, E-220
- Caryachine, E-220
- (\pm)-Caryachine, E-220
- Caryoline, C-171
- cis*-Caseadine *N*-oxide, C-172
- Caseadine, C-172
- Caseadinium(1+), C-173
- Casealutine, I-210
- cis*-Caseamine *N*-oxide, C-172
- Caseamine, C-172
- Caseanadine, C-174
- Caseanine, T-212
- Cashmiradelphine, P-776
- Casimidine, H-307
- Casimirine, C-175
- Casimiroedine, C-176
- Casimiroin, C-177
- Cassaide, N-289
- Cassaidine, C-178
- Cassaine, C-178
- Cassamedine, C-179
- Cassameridine, C-179
- Cassamide (obsol.), N-299
- Cassamidine, N-300
- Cassamine, N-300
- Cassiadine, C-180
- Cassiarine A, C-181
- Cassiarine B, C-182
- Cassine, C-183
- Cassinicine, S-388
- Cassinine, P-251
- Cassipoureamide A, C-184
- Cassipoureamide B, C-185
- Cassipourine, C-186
- Cassipurine, C-186
- Cassyfiline, O-27
- Cassyformine, P-209
- Cassythic acid, C-187
- Cassythicine, D-346
- (-)-Cassythicine, D-346
- Cassythidine, N-125
- Cassythine, O-27
- Castanospermine, C-188
- Castoramine, D-227
- Casuarine 6- α -*D*-glucoside, T-262
- Casuarine, T-262
- Catacandin, C-189
- Catalauesmine, D-606
- Catalauperine, D-606
- Cataline, C-190
- Catalpifoline, T-248
- Catechol-4-carboxylic acid, D-536
- Catechol-4,5-dicarboxylic acid, D-533
- Cathafiline, D-346
- Cathafoline *N*⁺-oxide, C-191
- Cathafoline, C-191
- Cathafornine, O-27
- Cathalanceine, C-192
- Cathanneine, C-213
- Cathaphylline, C-193
- Catharanthamine, C-194
- Catharanthine, C-195
- ψ -Catharanthine, P-733
- Catharicine, C-196
- Catharine, C-197
- Catharinensine, R-90
- Catharinine, V-107
- Catharosine, V-130
- Catharsitoxin A, C-198
- Catharsitoxin B, C-199
- Catharsitoxin C, C-200
- Catharsitoxin D, M-528
- Catharsitoxin E, M-527
- Catharsitoxin F, I-280
- Cathedulin E3, C-201
- Cathedulin E4, C-201
- Cathedulin E2, P-218
- Cathedulin E5, P-247
- Cathedulin E6, P-247
- Cathedulin E8, P-218
- Cathedulin K11, C-201
- Cathedulin K17, C-202
- Cathedulin K19, C-203
- Cathedulin K20, C-201
- Cathedulin K12, P-247
- Cathedulin K15, P-247
- Cathedulin K1, P-247
- Cathedulin K2, P-247
- Cathedulin K6, P-247
- Cathedulin 2, P-218
- Cathedulin Y10, C-206
- Cathedulin Y7, C-204
- Cathedulin Y9, C-205
- Cathedulin Y1, P-247
- Cathedulin Y8, P-218
- Cathenamine, C-207
- Cathestatin A, C-208
- Cathestatin B, C-208
- Cathestatin C, C-209
- Cathidine A, C-210
- Cathidine B, C-211
- Cathindine, C-212
- Cathine, A-869
- Cathinone, A-870
- Cathovaline, C-213
- Cathovalinine, H-743
- Catuabine A, M-391
- Catuabine B, M-391
- Catuabine C, M-391
- Catuabine D, M-391
- Catuabine E *N*-oxide, M-391
- Catuabine E, M-391
- Catuabine F, M-391
- Catuabine G, M-392
- Catuabine H, M-391
- 3 α *H*-Catuabine H, M-391
- Catuabine I, M-391
- Caulerpicin, C-214
- Caulerpin diacid, D-415
- Caulerpin, D-415
- Caulerpinic acid, D-415
- Caulersin, C-215
- Caulibugulone A, A-831
- Caulibugulone B, A-831
- Caulibugulone C, A-831
- Caulibugulone D, A-831
- Caulibugulone E, A-831
- Caulibugulone F, A-831
- Caulidine A, C-216
- Caulidine B, C-217
- Caulilexin A, M-434
- Caulilexin B, I-97
- Caulilexin C, I-81
- Caulindole A, C-218
- Caulindole B, C-218
- Caulindole C, C-219
- Caulindole D, C-219
- Caulitenin, A-716
- Caulophylline, C-938
- Caulophyllumine A, P-522
- Caulophyllumine B, C-220
- Cavanine, S-149
- Cavernicolin 1, C-221

- Cavernicolin 2, C-221
 Cavidine, C-222
 Cavincidine, C-223
 Cavincine, C-224
 Cavinine, A-683
 Caymandimerine, C-225
 CB 154, E-157
 CC 3, D-229
 CC 1065, R-4
 β -CCA methylamide, C-133
 CCU 18, D-229
 2096C, D-498
 Ceanothamine A, F-151
 Ceanothamine B, A-149
 Ceanothine A, C-226
 Ceanothine B, S-165
 Ceanothine C, C-227
 Ceanothine D, C-228
 Ceanothine E, C-229
 Cebin V, O-131
 Ceciline, C-230
 Ceflatonin[®], H-66
 Celabenzine, P-356
 Celacinnine, P-356
 Celafurine, P-356
 Celahinine A, P-247
 Celalocinnine, P-356
 Celapagine, T-258
 Celapanigine, T-258
 Celapanine, T-258
 Celastramycin A, C-231
 Celastromelidone, C-232
 Celastromeline, C-233
 Celenamide A, C-234
 Celenamide B, C-235
 Celenamide C, C-236
 Celenamide D, C-237
 Celenamide E, C-238
 Celeromycalin, P-774
 Celesticetin C, C-239
 Celesticetin, C-239
 Celgосivir hydrochloride, C-188
 Celgосivir, C-188
 Celliamine, C-240
 Cellocidin, B-447
 Celogentin A, C-241
 Celogentin B, C-241
 Celogentin C, C-242
 Celogentin D, C-241
 Celogentin E, M-702
 Celogentin F, M-702
 Celogentin G, C-243
 Celogentin H, C-243
 Celogentin J, C-243
 Celogentin K, C-244
 Celorbicol, T-542
 Celosianin I, B-110
 Celosianin II, B-110
 Celtine, C-805
 Celtisine, C-805
 Cenocladamide, C-245
cis-Centcyamine, M-288
 Centcyamine, M-288
 Centralite I, D-818
 Centralite II, D-818
 Centralite III, D-818
 Cepabactin, H-563
 Cepaciachelin, C-246
 Cepaciamide A, C-247
 Cepaciamide B, C-248
 Cepafungin I, G-96
 Cepafungin II, G-96
 Cepafungin III, G-96
 Cephaeline, E-72
cis-Cephaeloside, I-167
trans-Cephaeloside, I-167
 Cephacicine, S-311
 Cephalandole A, C-249
 Cephalandole B, C-250
 Cephalandole C, C-251
 Cephalezomine A, C-252
 Cephalezomine B, C-252
 Cephalezomine C, I-238
 Cephalezomine D, I-238
 Cephalezomine E, H-66
 Cephalezomine F, H-66
 Cephalezomine G, C-253
 Cephalezomine H, C-253
 Cephalezomine J, C-271
 Cephalezomine K, H-66
 Cephalezomine L, H-66
 Cephalezomine M, T-51
 Cephalimysin A, C-254
 Cephalinone A, C-255
 Cephalinone B, C-256
 Cephalinone C, C-257
 Cephalinone D, C-258
 Cephalocyclidine A, C-259
 Cephalofortuneine, C-260
 Cephalomannine, T-52
 Cephalostatin 10, C-261
 Cephalostatin 11, C-261
 Cephalostatin 12, C-268
 Cephalostatin 13, C-268
 Cephalostatin 14, C-261
 Cephalostatin 15, C-262
 Cephalostatin 16, C-269
 Cephalostatin 17, C-261
 Cephalostatin 18, C-261
 Cephalostatin 19, C-261
 Cephalostatin 1, C-261
 Cephalostatin 2, C-261
 Cephalostatin 3, C-262
 Cephalostatin 4, C-261
 Cephalostatin 5, C-263
 Cephalostatin 6, C-264
 Cephalostatin 7, C-265
 Cephalostatin 8, C-266
 Cephalostatin 9, C-267
 Cephalotaxidine, C-270
 Cephalotaxinamide, C-271
 Cephalotaxine 4-methyl 2-hydroxy-2-(3-hydroxy-3-methylbutyl)butanedioate (ester), H-66
 Cephalotaxine α -*N*-oxide, C-271
 Cephalotaxine β -*N*-oxide, C-271
 Cephalotaxine, C-271
 Cephalotaxinone, C-271
 Cephamonine, S-313
 Cephamorphinanine, S-311
 Cephamine, S-313
 Cepharradione A, C-272
 Cepharradione B, C-273
 Cepharamine, C-274
 Cepharranoline, C-275
 Cepharranone A *N*- β -*D*-glucoside, D-567
 Cepharranone A, D-567
 Cepharranone B, D-567
 Cepharranone C, T-252
 Cepharranthine 2'- β -*N*-oxide, C-275
 Cepharranthine, C-275
 Cephassamine, C-276
 Cephasugine, S-311
 Cephatonine, R-161
 Cerasodine, O-202
 Cerasonine, O-202
 Ceratamine A, C-277
 Ceratamine B, C-277
 Ceratinamide A, P-695
 Ceratinamide B, P-695
 Ceratinamine, M-679
 Ceratinic acid, H-214
 Ceratocapnidine, C-278
 Ceratocapnine, C-278
 Ceratonicine, C-279
 Ceratospongamide, C-280
 Cerespan, P-79
 Ceric acid, H-214
 Ceridimine, C-281
 Ceric acid, H-214
 Cermizine A, C-282
 Cermizine B, C-283
 Cermizine C, D-783
 Cermizine D, C-284
 Ceruine *N*-oxide, L-321
 Ceruine, L-321
 Cerotic acid, H-214
 N^b-Cerotoyltryptamine, T-639
 Cerpegin, C-285
 Cerulenin, C-286
 Cerulomycin, C-9
 Cervinomycin A₁, C-287
 Cervinomycin A₂, C-287
 Cervoxan, E-6
 Cespitulactam A, C-288
 Cespitulactam B, C-288
 Cespitulactam C, C-288
 Cespitulactam D, C-288
 Cespitulactam E, C-288
 Cespitulactam F, C-288
 Cespitulactam G, C-288
 Cespitulactam H, C-288
 Cespitulactam I, C-288
 Cespitulactam J, C-288
 Cespitulactam K, C-288
 Cetoniacytone A, C-289
 Cetoniacytone B, C-289
 Cevacine, V-56
 Cevadine, V-56
 3,6-Cevanediol, C-290
 3,4,12,14,16,17,20-Cevaneheptol, C-291
 3,6,12,14,16,20-Cevanehexol, C-292
 3,6,14,16,20-Cevanepentol, C-293
 3,6,14-Cevanetriol, C-295
 3,6,15-Cevanetriol, C-296
 3,6,16-Cevanetriol, C-297
 3,6,20-Cevanetriol, C-298
 3,6,7-Cevanetriol, C-294
 5-Cevaniden-3-ol, P-633
 Cevanine, C-299
 12,14,16-Cevatriene-3,6-diol, C-300
 Cevedine *N*-oxide, C-295
 Cevedine, C-295
 Cev-13(17)-ene-3,6-diol, C-308
 Cev-5-ene-3,20-diol, C-301
 Cev-8(14)-ene-1,3-diol, C-304
 Cev-8(14)-ene-3,11-diol, C-306
 Cev-8(14)-ene-3,15-diol, C-307
 Cev-8(14)-ene-3,6-diol, C-305
 Cev-8-ene-3,11-diol, C-303
 Cev-8-ene-3,6-diol, C-302
 Cev-12-ene-3,6,15-triol, C-312
 Cev-23-ene-3,6,16-triol, C-297
 Cev-5-ene-3,16,20-triol, C-309
 Cev-8-ene-3,6,20-triol, C-310
 Cev-8-ene-3,6,23-triol, C-311
 Cev-5-en-3-ol, C-313
 Ceverine *N*-oxide, C-295
 Ceverine, C-295
 Cevine, V-56
 Ceylanicine, C-314
 Ceylanine, C-315
 Cezomycin, C-26
 CGP 39360, S-498
 CGP 41251, S-498
 CGP 58546, S-498
 CG-SI-1, E-248
 CG-SI-2, E-248
 CGX 635, H-66
 CH 777A, N-180
 CH 846, E-6
 2088C, H-514
 Chabamide, C-316
 Chaboside, C-70
 α -Chaconine, S-345
 β ₁-Chaconine, S-345
 β ₂-Chaconine, S-345
 γ -Chaconine, S-345
 Chaenorhine, C-317
 Chaenorpine, C-318
 Chaetochalasin A, C-319
 Chaetocin B, C-321
 Chaetocin C, C-322
 Chaetocin, C-320
 Chaetocochin A, C-323
 Chaetocochin B, C-323
 Chaetocochin C, C-324
 Chaetoglobosin A, C-325
 Chaetoglobosin B, C-326
 Chaetoglobosin 510, C-329

- Chaetoglobosin 540, C-329
 Chaetoglobosin 542, C-329
 Chaetoglobosin C, C-325
 Chaetoglobosin D, C-327
 Chaetoglobosin E, C-326
 Chaetoglobosin F, C-325
 Chaetoglobosin F_{ex}, C-327
 Chaetoglobosin G, C-326
 Chaetoglobosin J, C-328
 Chaetoglobosin K, C-329
 Chaetoglobosin L, C-329
 Chaetoglobosin M, C-329
 Chaetoglobosin N, C-330
 Chaetoglobosin O, C-326
 Chaetoglobosin P, C-330
 Chaetoglobosin Q, C-331
 Chaetoglobosin R, C-331
 Chaetoglobosin T, C-328
 Chaetoglobosin U, C-332
 Chaetoindicin A, C-333
 Chaetoindicin B, C-334
 Chaetoindicin C, C-334
 Chaetomin B, C-335
 Chaetomin, C-335
 Chaetominine, C-336
 Chaetracin A, C-372
 Chairamidine, C-337
 Chairamine, C-338
 Chakranine, I-208
 Chaksine, C-339
 Chalcioporone, C-340
 Chalcostrobamine, C-341
 Chalcupine A, Y-13
 Chaloridone, C-342
 Chamaedrone, C-343
 Chamaetin, H-551
 Chamobtusine A, C-344
 Chandrananimycin A, A-862
 Chandrananimycin B, A-862
 Chandrananimycin C, C-345
 Chandrine, C-346
 Channaine, C-347
 Chanoclavine I acid, C-348
 Chanoclavine I aldehyde, C-348
 Chanoclavine I fructoside, C-348
 Chanoclavine I fructosylfructoside, C-348
 Chanoclavine I, C-348
 Chanoclavine II, C-348
 (±)-Chanoclavine II, C-348
 Chanoclavine, C-348
 Chanofrucitocine methyl ester, M-296
 Charamin, A-1610
 Charine, D-253
 Chartellamide A, C-349
 Chartellamide B, C-349
 Chartelline A, C-350
 Chartelline B, C-350
 Chartelline C, C-350
 Chasmaconitine, M-312
 Chasmanine, M-323
 Chasmanthine, M-312
 Chatinine, C-351
 Chavicine, P-467
 Cheilanthifoline, S-307
 Chelamidine, C-352
 Chelamine, C-353
 Chelelactam, C-354
 Chelerythridimerine, C-355
 Chelerythrine pseudoethanolate, D-407
 Chelerythrine pseudomethanolate, D-407
 Chelerythrine, C-356
 ψ-Chelerythrine, S-56
 Chelidamine, S-612
 Chelidimerine, S-54
 Chelidonine, C-357
 Cheliensisamine, C-358
 Cheliensisine, C-359
 Chelilutine, C-356
 Chelirubine, C-360
 Chellespontine, A-1533
 Chelonin A, C-361
 Chelonin B, C-362
 Chelonin C, C-363
 Chemetron 100, E-225
 Chemosan, A-736
 Chenabine, J-37
 Chenabinol methyl ether, J-37
 Chenabinol, J-37
 Chenoalbicin, C-364
 Chenopodine, L-153
 Chenorrhine, C-317
 Cheratamine, C-365
 Cherianoine, C-366
 Cherimoline, C-367
 Cherinonaine, C-368
 Cheritamine, T-639
 Cherylline, C-369
 Chestnutamide, C-370
 Chetocin B, C-321
 Chetocin C, C-322
 Chetocin, C-320
 Chetomin B, C-335
 Chetomin, C-335
 Chetoseminudin A, C-335
 Chetoseminudin B, C-371
 Chetoseminudin C, C-371
 Chetracin A, C-372
 Chevalieramide, B-421
 Chiapenine ES III, P-253
 Chiapenine ES II, P-253
 Chiapenine ES I, P-253
 Chiapenine ES IV, P-253
 Chicamycin A, C-373
 Chicamycin B, C-374
 Chidlovine, T-451
 Chilenamine, L-91
 Chilenine, C-375
 Chileninone, B-101
 Chillanamine, C-376
 Chilocorine A, C-377
 Chilocorine B, C-378
 Chilocorine C, C-379
 Chilocorine D, C-380
 Chilocorine, C-377
 Chiloenamine, C-381
 Chiloenine, C-381
 Chimanine A, H-707
 Chimanine B, P-647
 Chimanine C, P-647
 Chimanine D, P-647
 Chimonamidine, D-916
 Chimonanthidine, C-382
 Chimonanthine, C-382
 Chinentaxunine, A-7
 Chinese bittersweet alkaloid I, C-383
 Chinese bittersweet alkaloid II, C-384
 Chinese bittersweet alkaloid III, C-385
 Chingazumianine, C-386
 Chingchengenamide A, C-387
 Chingchengenamide B, C-387
 Chinpeimine, C-388
 Chippiine, C-389
 Chiriquitoxin, C-390
 Chisitine 1, C-391
 Chisitine 2, C-392
 Chitosenine, G-21
 Chitraline, P-16
 Chivosazole A, C-393
 Chivosazole A₁, C-393
 Chivosazole B, C-393
 Chivosazole C, C-393
 Chivosazole D, C-393
 Chivosazole E, C-393
 Chivosazole F, C-393
 Chlidanthine, G-3
 Chloptosin, C-394
 Chlordelazin, C-417
 Chloralbofungin, A-244
 Chlorobiocin, C-395
 6-Chlorobohemamine C, B-242
 3-Chloro-5-bromotyrosine (incorr.), A-712
 4-Chlorobutyl veratrate, D-706
 N-(4-Chlorobutyl)butanamide, C-396
 3-Chloro-9H-carbazole, C-397
 6-Chloro-β-carboline, C-398
 7-Chloro-β-carboline, C-399
 7-Chlorocavernicolone, B-301
 5-Chlorocavernicolin, C-221
 2-Chloro-5-(3-chloro-2,2-dimethyl-6-methylenecyclohexyl)-3-methylene-pentyl carbonimidic dichloride, D-349
 3-Chloro-4-(3-chloro-4-fluoro-2-nitrophenyl)-1H-pyrrole, P-961
 [2-Chloro-2-[4-chloro-5-hydroxy-3-methyl-3-(4-methyl-3-pentenyl)cyclohexylidene]ethyl]carbo-nimidic dichloride, C-400
 3-Chloro-4-(3-chloro-6-hydroxy-2-nitrophenyl)-1H-pyrrole, P-961
 4-Chloro-1-(chloromethyl)octahydro-5-methylenazocine, O-224
 3-Chloro-4-(3-chloro-2-nitrophenyl)-1H-pyrrole, P-961
 2-Chloro-6-(4-chloro-1H-pyrrol-3-yl)benzenamine, A-725
 4-Chloro-2-(4-chloro-1H-pyrrol-3-yl)-3-nitrophenol, P-961
 7-Chloro-6-demethylcepharadione B, C-273
 Chlorodeoxyoxovine, F-223
 6'-Chlorodeoxyfloridanine, O-92
 Chlorodeoxyscleratin, S-122
 Chlorodesnkolbisine, F-221
 Chlorodesoxyoxovine, F-223
 3-Chloro-2,7-dibromindole, D-308
 5-Chloro-2-(3,5-dichloro-2-hydroxyphenyl)-3-nitropyrrole, P-965
 20-Chloro-15,20-dihydro-12,15-dihydroxysenecio-nan-11,16-dione, J-4
 2-Chloro-2,3-dihydro-5-imino-1H-pyrrolizin-7a(5H)-ol, C-521
 3-Chloro-11,12-dihydroindolo[2,3-a]carbazole, C-401
 4-Chloro-5,6-dihydroxy-1H-indole, C-402
 4-Chloro-5,6-dihydroxy-1H-indole-1-sulfonic acid, C-402
 2-Chloro-1,6-dihydroxyphenazine, C-415
 2-Chloro-6,8-dihydroxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, S-571
 4-Chloro-6,7-dimethoxy-2-benzoxazolinone, D-708
 2-Chloro-10-(3-dimethylaminopropyl)phenothiazine, C-417
 3-Chloro-2-(3,4-dimethyl-3-pentenyl)-1,2,3,4-tetrahydro-2-(methoxymethyl)-6-quinolinecarboxylic acid, V-145
 2-Chloro-N,N-dimethyl-10H-phenothiazine-10-propanamine, C-417
 Chlorodysinosin A, D-975
 3-Chloro-2-ethenyl-1-isocyanato-2-methyl-5-(1-methylethenyl)spiro[bicyclo[4.1.0]heptane-7,3'-[3H]indol]-2'(1'H)-one, H-51
 8-Chloro-9-ethenyl-2,6,6a,7,8,9,10,10a-octahydro-10-isocyanato-6,6,9-trimethylnaphth[1,2,3-cd]indole, H-48
 (3-Chloro-5-ethyl-2,6-dihydroxyphenyl)(4,5-dichloro-1H-pyrrol-3-yl)methanone, A-1283
 2-Chloro-7-ethyl-6,8-dihydroxy-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, C-403
 2-Chloro-7-ethyl-8-hydroxy-6-methoxy-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, C-403
 2-Chloro-7-ethyl-8-hydroxy-6-methyl-9H-pyrrolo[1,2-b][1,3]benzoxazin-9-one, C-404
 3-Chloro-N-formylnormantennine, N-29
 Chlorofusin, C-405
 2-Chlorohuperzine E, H-378
 3-Chloro-4-hydroxybenzenaeetic acid, C-407
 12-Chloro-11-hydroxydibromoisophakellin, D-328
 erythro-3''-Chloro-2''-hydroxyechiumine, E-31
 2-[1-Chloro-2-hydroxy-1-(hydroxymethyl)ethyl]-1,11-dihydro-5-hydroxy-11-methylfuro[2,3-c]jaccridin-6(2H)-one, R-167
 5-Chloro-2-hydroxy-6-methoxybenzoxazole, C-406
 2-Chloro-8-hydroxy-6-methoxy-7-propyl-9H-pyrrolo[2,1-b][1,3]benzoxazin-9-one, S-571
 7-(3-Chloro-2-hydroxy-3-methylbutoxy)-4-methoxyfuro[2,3-b]quinoline, F-212
 7-O-(3-Chloro-2-hydroxy-3-methylbutyl)haplopin, F-223
 (3-Chloro-4-hydroxyphenyl)acetamide, C-407
 (3-Chloro-4-hydroxyphenyl)acetic acid, C-407

- 3-Chloro-4-hydroxy-2-piperidinone, C-408
 Chlorohyellazole, M-270
 4-Chloro-1*H*-indole-3-acetic acid, C-410
 3-Chloro-1*H*-indole, C-409
 4-Chloro-1*H*-indole-5,6-diol, C-402
 4-Chloro-5-(3-indolyl)oxazole, O-157
 Chloroiodoacetamide, C-411
 Chloroiodoacetic acid, C-411
 Chlorolisoclimide, D-354
 5-Chloro-6-methoxy-2-benzoxazolinone, C-406
 5-Chloro-6-methoxy-2(3*H*)-benzoxazolone, C-406
 5-Chloro-6-methoxy-1-methyl-1*H*-indole-2,3-dione, C-412
 5-Chloro-6-methoxy-1-methylisatin, C-412
 6-Chloro-3-methoxy-2-methyl-1-phenyl-9*H*-carbazole, M-270
 6-Chloro-7-(methylamino)-5,8-isoquinolinedione, A-831
N-Chloromethylbulgarsenine, B-401
 7-Chloro-9-methyl- β -carboline, C-399
 6-Chloro-2-methyl- β -carbolinium, C-398
 4-(Chloromethyl)-2,16-didehydro-16-(methoxycarbonyl)condyfolanium, C-413
N-Chloromethyl-*N,N*-dimethyltryptamine, D-785
 Chloromethyleneaffinisinium, T-425
 4-Chloromethylene-1,2,3,4-tetrahydro-6-hydroxy-8-methylquinoline, C-414
 4-Chloromethylene-1,2,3,4-tetrahydro-8-methyl-6-quinolinol, C-414
N-Chloromethylnarcissidinium, N-43
*N*²-Chloromethyltetrandrinium, T-286
 3-Chloro-4-(2-nitrophenyl)pyrrole, P-961
 6-Chloronorharman, C-398
 7-Chloronorharman, C-399
 3-(4-Chloro-5-oxazolyl)-1*H*-indole, O-157
 2-Chloro-1,6-phenazinediol, C-415
 2-(6-Chloro-3-pyridinyl)-7-azabicyclo[2.2.1]heptane, E-87
 7-Chloro-9*H*-pyrido[3,4-*b*]indole, C-399
 6-Chloro-9*H*-pyrido[3,4-*b*]indole, C-398
 Chloropyrrolnitrin, P-961
 6-Chloro-2-quinoxalinecarboxylic acid, C-416
 Chlororaphin, P-319
 8-Chlororugulovasine A, R-157
 8-Chlororugulovasine B, R-157
 6-Chloro-1,3,4,5-tetrahydro-1-methyl-2-(methylthio)pyrrolo[4,3,2-*de*]quinoline-7,8-dione, B-52
 1-(2-Chlorotetrahydro-4-methyl-2*H*-1,3-oxazin-5-yl)ethanone, A-46
 6-Chloro-8*H*-thieno[2,3-*b*]indole-2-carboxamide, T-374
 3-Chlorotyrosine, A-724
 Chloroxylophine, F-223
 Chlorpromazine, C-417
 Choisyine, C-418
 Cholergol, D-669
 Choline acetate, A-47
 Choline chloride, C-419
 Choline fenofibrate, C-419
 Choline imidazole-4-acrylate, M-739
 Choline orotate, D-669
 Choline 4-hydroxy-3,5-dimethoxycinnamate, S-308
 Choline imidazole-4-acrylate, M-739
 Choline perchlorate nitrate, C-419
 Choline theophyllinate, C-419
 Choline urocanate, M-739
 Choline, C-419
 Cholesulfuric acid, C-419
 Cholinophylline, C-419
 Chondocurarine, C-420
 Chondocurine, C-420
 Chondodendrine, C-812
 Chondriamide A, C-421
 Chondriamide B, C-421
 Chondriamide C, C-421
 Chondrochloren A, C-422
 Chondrochloren B, C-422
 Chondrocurarine, C-420
 Chondrocurine, C-420
 Chondrofoline, C-812
 Chonemorphine, P-601
 Chrestifoline A, C-423
 Chrestifoline B, C-424
 Chrestifoline C, C-425
 Chrestifoline D, B-207
 Chromopyrrolic acid, L-329
 Chromoviridan, C-426
 Chromoxymycin, C-427
 Chrycentrine, C-428
 Chrysanthemine, S-491
 Chrysanthone, P-892
 Chrysogine, H-499
 Chrysopentamine, C-429
 Chrysophysarin A, C-430
 Chrysopine, C-431
 Chrysopterin, C-432
 Chrysotricine, C-433
 Chuanbeinone, C-290
 Chuanfumine, C-434
 Chuanghsinmycin, D-480
 Chuangxinmycin, D-480
 Chuan-wu base A, C-435
 Chuan-wu base B, C-436
 Chuangxiongine, T-284
 Chuchuhuanine EIII, P-244
 Chuchuhuanine EII, P-244
 Chuchuhuanine EI, P-244
 Chuchuhuanine EIV, P-244
 Chuchuhuanine EVII, P-244
 Chuchuhuanine EVI, P-244
 Chuchuhuanine EV, P-244
 Chuchuhuanine WI, P-243
 Chysin A, P-960
 Chysin B, P-960
 Chysine, P-960
 C.I. 75160, B-101
 CI 4, C-841
 CI 75800, D-325
 C.I. 75780, I-72
 C.I. 75790, I-73
 C.I. Natural Blue 1, I-72
 C.I. Natural Red 33, B-110
 C.I. Natural Red 28, O-102
 C.I. Natural Violet 1, D-325
 C.I. Natural White 1, G-210
 C.I. Vat Blue 1, I-72
 Ciba 12669A, D-92
 Ciba 33112, D-142
 Cibarian, G-104
 Cichorine, C-437
 Cicloheximide, D-765
 Cicutine, P-652
 CIF 1, D-517
 CIF 2, D-517
 CIF 3, D-517
 Ciliaphylline *N*-oxide, R-134
 Ciliaphylline, R-134
 Ciliapterin, A-749
 Ciliatamide A, C-438
 Ciliatamide B, C-438
 Ciliatamide C, C-439
 Ciliatine, A-773
 Cimicidine, D-144
 Cimiciduphytine, C-440
 Cimicifine A, N-247
 Cimicifugadine, N-247
 Cimicifugamide, H-566
 Cimicine, D-144
 Cimiciphytine, C-441
 Cimilophytine, C-442
 Cimipronidine, P-950
 Cinachyramine, C-443
 Cinchamidine, C-446
 Cinchonamine, C-444
 Cinchonamine, C-444
 Cinchonaminone, C-445
 Cinchonon-9-ol, C-446
 Cinchonon-9-ol, C-447
 Cinchonon-9-one, C-446
 Cinchonine, Q-17
 Cinchoninicin, Q-17
 Cinchonidicinol, Q-17
 Cinchonidine, C-446
 Cinchonidinone, C-446
 Cinchonifine, C-447
 Cinchoninaldehyde, Q-30
 Cinchonine, C-447
 ψ -Cinchonine, C-447
 Cinchoninic acid, Q-32
 Cinchoninone, C-446
 Cinchononitrile, Q-32
 Cinchophyllamine, C-448
 3 α ,17 α -Cinchophylline, C-448
 3 α ,17 β -Cinchophylline, C-448
 3 β ,17 α -Cinchophylline, C-448
 3 β ,17 β -Cinchophylline, C-448
 Cinchotine, C-447
 Cinchotoxine, Q-17
 Cinchovatine, A-1423
 Cinegalleine, H-554
 Cinegalline, H-554
 Cinereain, C-449
 Cinereapyrrole A, B-182
 Cinereapyrrole B, B-182
 Cineractone, H-554
 Cinevanine, H-554
 Cineverine, H-554
 Cinnabaramide A, C-450
 Cinnabaramide B, C-450
 Cinnabaramide C, C-450
 Cinnabaramide D, C-450
 Cinnabaramide E, C-450
 Cinnabaramide F, C-451
 Cinnabaramide G, C-451
 Cinnabaric acid, C-452
 Cinnabarine, C-452
 Cinnabarinic acid, C-452
 Cinnaloid, R-50
 Cinnamic acid 2-hydroxy-2-(*p*-methoxyphenyl)ethylamide, A-153
 Cinnamic acid *p*-methoxyphenethylamide, H-144
 Cinnamic acid pyrrolide, C-457
 Cinnamic acid pyrrolidide, C-458
 Cinnamolaurine, C-453
 Cinnamonin, A-78
 3-[2-(Cinnamoylamino)ethyl]-3-hydroxy-2-indolone, C-454
 1'-Cinnamoyl-2,3'-bipiperidine, A-1525
 Cinnamoylcoaine, E-8
 (*Z*)-Cinnamoylcoaine, E-8
 1'-Cinnamoyl-1,1',2,3,4,4',5,5',6,6'-decahydro-2,3'-bipyridine, A-138
N-Cinnamoyl-2,4'-dihydroxyphenethylamine, O-63
 Cinnamoyllegonine methyl ester, E-8
 8-*O*-Cinnamoylgraciline, G-152
*N*²-Cinnamoylhistamine, C-455
N-Cinnamoyl-20-hydroxycylindrocaine, C-929
N-Cinnamoyl-2-hydroxy-2-(4-methoxyphenyl)ethylamine, A-153
*N*¹-Cinnamoyl-*N*⁴-(4-hydroxytylglyl)putrescine, C-456
N-Cinnamoyl-2-methoxypyrrolidine, P-955
N-Cinnamoyl-2-(2-methylbutanoylamino)pyrrolidine, O-67
*N*¹-Cinnamoyl-*N*⁴-(2-methylbutanoyl)putrescine, C-456
*N*²-Cinnamoyl-*N*²-methylhistamine *N*-glucoside, C-176
cis-*N*²-Cinnamoyl-*N*¹-methylhistamine, C-455
trans-*N*²-Cinnamoyl-*N*¹-methylhistamine, C-455
N-Cinnamoyl-2-(2-methylpropanoylamino)pyrrolidine, P-479
*N*¹-Cinnamoyl-*N*⁴-(2-methylpropanoyl)putrescine, C-456
 8-*O*-*trans*-Cinnamoylneoline, M-323
 14-Cinnamoylneoline, M-323
N-Cinnamoylnorcuskygrine, D-211
 14-*cis*-Cinnamoylnudicaulidine, M-321
 14-*trans*-Cinnamoylnudicaulidine, M-321
N-Cinnamoyloctopamine, O-63
cis-13-Cinnamoyloxylyupanine, H-554
trans-13-Cinnamoyloxylyupanine, H-554
cis-13-Cinnamoyloxy-17-oxolupanine, O-188
trans-13-Cinnamoyloxy-17-oxolupanine, O-188
 6-*trans*-Cinnamoyloxy-3-propanoyloxynortropane, A-1584
 3-Cinnamoyloxy-6,7-tropanediol, M-392
 3-Cinnamoyloxytropan-6-ol, M-391
N-Cinnamoylphenethylamine, P-344

- 9-Cinnamoyl-4-phenyl-1,5,9-triazacyclotridec-12-en-2-one, P-265
N-Cinnamoyl-*O*-prenyloctopamine, O-63
N-Cinnamoyl-*p*-prenyloxyphenethylamine, H-667
N-Cinnamoylputrescine, C-456
 1-Cinnamoyl-1*H*-pyrrole, C-457
 1-Cinnamoylpyrrolidine, C-458
N-Cinnamoyl-2-pyrrolidinone, P-956
 14-*O*-*trans*-Cinnamoyltalatzamine, M-352
N-Cinnamoyl- Δ^2 -tetrahydroanabasine, A-138
*N*¹-Cinnamoyl-*N*⁴-tigloylputrescine, C-456
 1-Cinnamoyl-2,4,5-trimethylpiperazine, D-769
N-Cinnamoyltyramine, H-667
 Cinobufotenine, B-393
 Cinodine hydrochloride, C-459
 Cinodine I, C-459
 Cinodine II, C-460
 CinQuin, Q-18
 Circinadine A, M-304
 Circinadine B, M-304
 Circinamide, C-461
 Circinasine A, M-306
 Circinasine B, M-334
 Circinasine C, M-304
 Circinasine D, M-319
 Circinasine E, M-319
 Circinasine F, M-334
 Circinatin, C-462
 Circumdatin A, C-463
 Circumdatin B, C-463
 Circumdatin C, C-464
 Circumdatin D, C-465
 Circumdatin E, C-465
 Circumdatin F, C-464
 Circumdatin G, C-464
 Circumdatin H, C-465
 Circumdatin J, C-465
 Cirsiumamide, P-456
 Cissaglaberrimine, C-466
 Cissamine, C-833
 Cissampareine, W-9
 Cissampentin, C-467
 Cissampine, C-833
 Cissetin, C-468
 Citbismine A, C-469
 Citbismine B, C-469
 Citbismine C, C-469
 Citbismine D, C-470
 Citbismine E, C-469
 Citbismine F, C-469
 Citbrasine, P-205
 Citisine, C-938
 Citorellamine, C-471
 Citpressine II, T-243
 Citpressine I, T-243
 Citraconimide, M-542
 Citracridone I, C-472
 Citracridone II, C-472
 Citracridone III, C-472
 Citramine, P-206
 Citreamicin α , C-473
 Citreamicin β , C-474
 Citreamicin η , C-476
 Citreamicin γ , C-475
 Citreamicin ξ , C-473
 Citroazopyrone, C-477
 Citridone A, C-478
 Citridone B', C-479
 Citridone B, C-479
 Citridone C, C-480
 Citridone D, C-481
 Citrinadin A, C-482
 Citrinadin B, C-482
 Citriodorol, C-483
 Citropone A, C-484
 Citropone B, C-485
 Citropone C, C-486
 Citrovorin, P-772
 Citrovorum factor, F-121
 Citrusamine, T-518
 Citrusinine II, T-242
 Citrusinine I, T-242
 Cividiclavine, C-487
 CJ 12662, A-1141
 CJ 12663, A-1141
 CJ 15696, A-1145
 CJ 16169, A-1145
 CJ 16170, A-1142
 CJ 16171, A-1143
 CJ 16173, A-1144
 CJ 16174, A-1145
 CJ 16196, A-1146
 CJ 16197, A-1146
 CJ 16264, A-1147
 CJ 16367, A-1148
 CJ 17572, A-1149
 CJ 17665, A-1470
 CJ 21058, A-1150
 CJ 13136, D-760
 CJ 13217, D-760
 CJ 13536, D-760
 CJ 13564, D-760
 CJ 13565, D-762
 CJ 13566, D-762
 CJ 13567, D-762
 CJ 13568, D-762
 CJ 14877, D-666
 CJ 14897, D-666
 CJ 15335, D-666
 CL 98984, C-459
 CL 307782, F-121
 Cladioxazole, C-488
 Cladobotryal, A-1145
 Clarkeanidine, C-174
 Clathculine A, C-489
 Clathculine B, C-489
 Clathramide A, C-490
 Clathramide B, C-490
 Clathramide C, C-490
 Clathramide D, C-490
 Clathridine A, C-491
 Clathridine B, C-492
 Clathridine C, C-493
 Clathridine Zn, C-491
 Clathridine-9-*N*-(2-sulfoethyl)imine, C-494
 Clathrodine, O-123
 Clathrotropine, C-495
 Clathryimine A, C-496
 Clathryimine B, C-496
 Clathrynamide A, C-497
 Clathrynamide B, C-497
 Clathrynamide C, C-497
 (6*E*)-Clathrynamide A, C-497
 Clausamide II, P-348
 Clausamide I, P-348
 Clausamine A, C-498
 Clausamine B, C-498
 Clausamine C, C-498
 Clausamine D, C-507
 Clausamine E, C-499
 Clausamine F, C-499
 Clausamine G, C-499
 Clausaniline, C-500
 Clausenal, D-554
 Clausenalene, D-615
 Clausenamidine, C-502
 ζ -Clausenamidine, C-501
 Clausenamidine A, C-503
 Clausenapin, C-504
 Clausenaquinone A, C-505
 Clausenatone A, M-729
 Clausenine, D-613
 Clausenlactam, D-294
 Clausenol, D-613
 Clausevatine D, C-506
 Clausevatine E, C-506
 Clausevatine F, C-506
 Clausevatine G, C-506
 Clausine A, D-557
 Clausine B, T-533
 Clausine C, H-455
 Clausine D, C-507
 Clausine E, H-452
 Clausine F, C-507
 Clausine G, D-558
 Clausine H, D-560
 Clausine I, D-552
 Clausine J, T-532
 Clausine K, D-560
 Clausine L, H-453
 Clausine M, H-455
 Clausine N, H-455
 Clausine O, D-556
 Clausine P, D-614
 Clausine Q, D-553
 Clausine R, D-559
 Clausine S, H-125
 Clausine T, C-508
 Clausine TY, D-560
 Clausine U, H-125
 Clausine V, C-123
 Clausine W, C-508
 Clauszoline A, C-509
 Clauszoline B, C-510
 Clauszoline C, D-560
 Clauszoline D, H-125
 Clauszoline E, M-757
 Clauszoline F, M-729
 Clauszoline G, M-757
 Clauszoline H, C-511
 Clauszoline I, H-452
 Clauszoline J, D-560
 Clauszoline K, H-450
 Clauszoline L, H-455
 Clauszoline M, D-557
 Clavamine, C-512
 Clavatine, F-73
 Clavatoxine, F-73
 Clavepictine A, C-513
 Clavepictine B, C-513
 Clavicipitic acid I, C-514
 Clavicipitic acid II, C-514
cis-Clavicipitic acid, C-514
trans-Clavicipitic acid, C-514
 Clavicularine, S-80
 Clavizepine, C-515
 Clavolonine, F-29
 Clavosine A, C-516
 Clavosine B, C-516
 Clavosine C, C-516
 Clavulazine, C-517
 Clavulazol A, C-518
 Clavulazol B, C-519
 Clavulinin, C-520
 Clazamycin A, C-521
 Clazamycin B, C-521
 Clazamycin, C-521
 7-Cl-DIBOA-Glc, H-409
 Cleavamine, C-522
 Cleistopholine, M-399
 Clemaine, C-523
 Cleomycin B₂, C-524
 Cleregil, D-730
 3,14-Clerodadiene-13,18-diol, C-525
 Clinacoside C, C-526
 Clionamide, C-527
 Clivacetine, C-532
 Clivatine, C-532
 Clivialine, C-528
 Cliviahaksine, C-528
 Cliviamartine, C-532
 Cliviasindhine, L-348
 Cliviasine, C-532
 Cliviasyline (incorr.), C-529
 Clividine, C-532
 Clivimine, M-629
 Clivisyline, C-529
 Clivojuline, C-530
 Clivonidine, C-531
 Clivonine, C-532
 Clivorine, C-533
 Clolimalongine, L-176
 Clorobiocin, C-395
 Closina, A-832
 Clovamide, C-534
 CNS 2103, C-535
 CNS 2104, C-536
 CO 61, H-715
 547C, O-131

- Cocaine hydrochloride, C-537
 Cocaine, C-537
 ψ -Cocaine, C-537
 Cocamine, T-634
 Cocarboxylase, T-366
 Coccinelline, D-886
 Coccinine, P-57
 Coccoline, E-202
 Coccolimine, E-202
 Coccludienone, E-199
 Cocculidine *N*-oxide, C-538
 Cocculidine, C-538
 Cocculidinone, C-539
 Cocculimine, I-207
 Cocculine, C-538
 Cocculitine, C-539
 Cocculitine, C-539
 Cocculolidine, C-540
 Coccuorbiculatine A, C-541
 Coccutrine, C-542
 Coccuvinine, C-543
 Coccuvinine, C-543
 Cochinchistemonine, C-544
 Cochinchistemoninone, C-544
 Cochinnicin I, C-545
 Cochinnicin II, C-545
 Cochinnicin III, C-545
 Cochinnicin IV, C-545
 Cochinnicin V, C-545
 Cochlearine, C-546
 Cochlearine, M-393
 Cochliodinol, C-547
 Cochliodinone, C-548
 Coclafine, C-538
 Coclamine, C-549
 Coclanoline, R-59
 Coclaurine, H-275
 Coclifoline, C-550
 Coclobine, C-551
 Cocsarmine, H-752
 Cocsiline, C-552
 Cocsilinine, C-552
 Cocsoline 2'- β -*N*-oxide, I-339
 Cocsoline, I-339
 Cocsoline 2-*N*-oxide, I-339
 Cocsuline, I-339
 Cocsulinine, C-552
 Cocsupendine, C-553
 Codamine *cis*-*N*-oxide, C-554
 Codamine, C-554
 Codaphniphylline, D-68
 Codeigene, C-555
 Codeine aminoxyde, C-555
 Codeine *N*-oxide, C-555
 Codeine phosphate, C-555
 Codeine sulfate, C-555
 Codeine, C-555
 β -Codeine, N-134
 Codeinfos, C-555
 Codeinone methyl enol ether, T-350
 Codeinone, C-555
 Codeisan, C-555
 Codeophen, C-555
 Codethyline, M-704
 Codicept, C-555
 Codlin, C-555
 Codonocarpine, C-556
 Codonopsine, C-557
 Codonopsinine, C-557
 Codonopsinol, R-10
 Codopiloic acid, O-152
 Coelenteramide, C-558
 Coelenteramine, C-559
 Coelenterazine, C-560
 Coelibactin, C-561
 Coelobillardierine, C-562
 Coelosperminone, C-562
 Coenzyme F, P-772
 Coenzyme PQQ, M-236
 Coenzyme R, B-139
 Coerulescine, H-359
 Coffearin, C-149
 Coffeine, C-15
 CoF, P-772
 Cogmine, P-400
 Coharmine, H-591
 Cohirsine, C-563
 Cohirsinine, C-564
 Cohirsitine, C-565
 Cohirsitinine, C-566
 Coixlactam B, D-451
 Coixol, B-82
 Colamine, A-767
 Colemid, D-92
 Colchameine, D-92
 Colchamine, D-92
 Colchibiphenylene, C-567
 Colchiceine methyl ether, C-569
 Colchiceine, D-92
 Colchiciline, C-568
 Colchicine, C-569
 Colchicoside, D-92
 Colchiethanamine, C-570
 Colchiethine, C-570
 Colchifoleine, C-571
 Colchifoline, C-571
 Colchilutine, C-572
 Colchiritchine, C-573
 Colchisol, C-569
 Colletine, H-275
 Collismycin A, C-574
 Collismycin B, C-574
 Collismycin C, C-575
 Collismycin D, C-576
 Collismycin E, C-577
 Collismycin F, C-578
 Collismycin, C-574
 Collutine *N*-oxide, A-1005
 Collutine, A-1005
 Colorflamme, S-544
 Colsaloid, C-569
 Colubricidin A, C-579
 β -Colubrine *N*-oxide, S-589
 α -Colubrine, S-589
 β -Colubrine, S-589
 Colubrinol acetate, T-450
 Colubrinol, T-450
 Columbamine, P-34
 Columbianine, M-352
 Columbidine, M-352
 Columellarine, C-580
 Combretine A, H-476
 Combretine B, H-476
 Commersonine, D-205
 Communesin A, C-581
 Communesin B, C-582
 Communesin C, C-582
 Communesin D, C-581
 Communesin D, C-582
 Communesin E, C-581
 Communesin F, C-581
 Communesin G, C-583
 Communesin H, C-584
 Comoramide A, C-585
 Comoramide B, C-586
 Comosidine, L-278
 Comosiline, F-143
 Comosimine, S-125
 Comosine, C-587
 Comosivine, C-588
 Compactinervine, C-589
 Complanadine A, C-590
 Complanadine B, C-590
 Complanadine C, C-591
 Complanadine D, C-590
 Complanatin, C-592
 Complanatine, L-341
 Compound 42339, A-120
 Compound CC 3, D-229
 Compound CCU 18, D-229
 Comptonine, T-48
 ComX_{RO-E-2} pheromone, C-593
 ComX_{RO-H-1} pheromone, C-594
 Conaconitine, M-351
 Cona-1,4-dienin-3-one, H-314
 3,5-Conadien-7-one, F-198
 Conambine, M-309
 Conamine, C-595
 Conan-5-en-3-ol, L-56
 Conarrhimine, C-595
 Cona-1,4,14-trienin-3-one, H-314
 Concavine, C-596
 Conchairamidine, C-597
 Conchairamine, C-598
 Conchicine, Q-18
 Concinnamide, H-471
 Concneorine, M-393
 Concurchine, C-610
 Concuressine, C-601
 Concusconine, C-599
 Condaline A, C-600
 Condelphine, M-352
 Condensamine, C-115
 Condurangamine A, P-603
 Condurangamine B, P-603
 Condyllocarpine *N*⁴-oxide, T-673
 Condyllocarpine, T-673
 Con-5-enin-3-amine, C-595
 Con-5-enin-3-ol, L-56
 Conessidine, C-610
 Conessimine, C-595
 Conessine, C-601
 Confolidine, C-633
 Confoline, C-633
 Confusadine, F-212
 Confusameline, F-212
 Conglobatin, C-602
 Congocidine, N-180
 α -Conhydrine, C-603
 Conhydrine, C-603
 ψ -Conhydrine, P-653
 Conhydrinone, C-604
 Conicamine, D-172
 Conicaquinone A, C-605
 Conicaquinone B, C-606
 Conicine, P-652
 Coniine, P-652
 Conimine, C-595
 Conioidine A, C-607
 Conioidine B, C-607
 Coniosetin, C-608
 Coniothyriomycin, C-609
 Konkurchine, C-610
 Konkuressine, C-601
 Conodiparine A, C-611
 Conodiparine B, C-612
 Conodiparine C, C-611
 Conodiparine D, C-612
 Conodiparine E, C-611
 Conodiparine F, C-612
 Conodirinine A, C-613
 Conodirinine B, C-614
 Conoduramine, C-615
 Conodurine, C-616
 Conodurinine, C-616
 Conodusarine, E-189
 Conodutarine A, C-617
 Conodutarine B, C-617
 Conoflorine, V-179
 Conofoline, C-618
 Conolidine, E-191
 Conolline, C-619
 Conolobine A, E-191
 Conolobine B, E-191
 Cononitarine A, C-612
 Cononitarine B, C-612
 Conopharyngine hydroxyindolenine, C-658
 Conopharyngine oxindole, C-726
 Conopharyngine pseudoindoxyl, V-178
 Conopharyngine, A-886
 Conopharyngine, C-620
 Conophyllidine, C-621
 Conophylline, C-621
 Conophyllinine, C-621
 Conquinamine, Q-12
 Conquinine, Q-18
 Consabatine, C-622
 Consiculine, C-622
 Consolarine, M-321

- Consolidine, C-623
 Consolidine, M-309
 Consolinine, M-323
 Consorientaline, A-1533
 Consperine, B-407
 Constrictosine, C-624
 Contomin, C-417
 Contorine, H-161
 Contorsine, H-161
 Contortine, H-161
 Contortumine, A-98
 Converginine, D-886
 Convolacine, M-391
 Convolamine *N*-oxide, C-633
 Convolamine, C-633
 Convolicine, C-633
 Convolidine, C-633
 Convoline, C-633
 Convolutamide A, D-96
 Convolutamide B, D-96
 Convolutamide C, D-96
 Convolutamide D, D-96
 Convolutamide E, D-96
 Convolutamide F, D-96
 Convolutamine A, C-625
 Convolutamine B, C-625
 Convolutamine C, C-625
 Convolutamine D, C-626
 Convolutamine E, C-627
 Convolutamine F, C-625
 Convolutamine G, C-625
 Convolutamine H, C-628
 Convolutamylidines, C-629
 Convolutindole A, C-630
 Convolvicine, C-631
 Convolvidine, C-632
 Convolvine, C-633
 Convosine, C-633
 Copiamycin, C-634
 Coprogen B, C-635
 Coprogen, C-635
 Coproverdine, C-636
 Copsinine, K-78
 Copsomylin, C-637
 Coptisine, C-638
 Corallistine, C-639
 Coramine, X-30
 Coranicine, C-640
 Corcovadine, P-462
 Cordatanine, H-440
 Cordatine A, S-201
 Cordatine B, S-201
 Cordianin, A-625
 Cordifoline, C-641
 Cordiline, C-303
 Cordizine, C-642
 Cordobimine, O-28
 Cordobine, C-643
 (±)-Cordrastine I, C-644
 (-)-Cordrastine I, C-644
 (±)-Cordrastine II, C-644
 (-)-Cordrastine II, C-644
 Cordrastine, C-644
 Cordyformamide, C-645
 Cordypyridone A, C-646
 Cordypyridone B, C-646
 Cordypyridone C, C-647
 Cordypyridone D, C-647
 Corepanine, M-309
 Coreximine, X-30
 Corftaline, H-394
 Corgoine, S-234
 Coriacenine A, C-648
 Coriacenine B, C-648
 Coriacenine C, C-649
 Coriacenine D, C-649
 Coriacenine E, C-650
 Corialstonidine, C-651
 Corialstonine, C-652
 Corifine, C-697
 Coriphidine, C-696
 Corledine, A-146
 Corlumidine, C-653
 Corlumine, C-653
 Cornigereine, C-654
 Cornigerine, C-654
 Cornocentin, E-164
 Cornucervine, C-655
 Corollin, G-104
 Coromandaline, T-443
 Coromandalinine, A-669
 Coronafacic acid, C-656
N-Coronafacylalloisoleucine, C-656
N-Coronafacyloisoleucine, C-656
N-Coronafacylserine, C-656
N-Coronafacylthreonine, C-656
N-Coronafacylvaline, C-656
 Coronarian, G-104
 Coronaridine hydroxyindolenine, C-658
 Coronaridine pseudoindoxyl, V-178
 Coronaridine, C-657
 Coronillin, G-104
 Corpaine, R-9
 Corphthaline, H-394
 Correantine A, C-659
 Correantine B, C-660
 Correantine C, C-661
 Correantoxide, C-662
 Corsevine, K-88
 Corsevinine, K-89
 Corsutine, C-663
 Cortamidine oxide, C-664
 Corticrocin, T-126
 Cortistatin A, C-665
 Cortistatin B, C-665
 Cortistatin C, C-665
 Cortistatin D, C-665
 Cortistatin J, C-665
 Cortistatine E, C-667
 Cortistatine F, C-666
 Cortistatine G, C-667
 Cortistatine H, C-667
 Cortistatine K, C-668
 Cortistatine L, C-668
 Corumdefine, M-325
 Corumdizine, D-370
 Corumdizimine, D-370
 Corunnine, O-178
 Coruscine, C-669
 Corybrachylobine, C-686
 Corybulbine, C-675
 Corycavamine, C-670
 Corycavidine, C-671
 Corycavine, C-670
 Corycidine, C-672
 Corycularicine, C-804
 Corydaine, C-673
 Corydalactam, E-250
 Corydaldine, I-305
 Corydalic acid methyl ester, C-674
 Corydalic acid, C-674
 Corydalidzine, C-675
 Corydaline, C-675
 Corydalis A, C-675
 Corydalis B, T-212
 Corydalis C, P-667
 Corydalis E, S-612
 Corydalis G, C-675
 Corydalis L, X-30
 Corydalisol, C-676
 Corydalispirone, H-766
 Corydalmine α -*N*-oxide, C-677
 Corydalmine β -*N*-oxide, C-677
 Corydalmine, C-677
 Corydamine, C-678
 Corydecumbine, C-679
 Corydendramine A, C-680
 Corydendramine B, C-680
 Corydicine, C-681
 Corydine *N*-oxide, C-682
 Corydine, C-682
 Corydinemethine, C-683
 Corydinine, P-667
 Corydione, C-684
 Corygovanine, C-685
 Coryincine, C-692
 Corymbin, Y-13
 Corymine hemiacetal, D-410
 Corymine, D-410
 Corymotine, C-686
 Corynantheal, D-412
 Corynantheidinaline, C-687
 Corynantheidinalinic acid, C-688
 Corynantheidine, C-689
 Corynantheidol, D-412
 Corynantheine, C-689
 Corynantheol, D-412
 Corynanthic acid, Y-13
 Corynanthine, Y-13
 Corynanthol, Y-16
 Corynebactin, B-2
 Coryneine, C-690
 Corynine, Y-13
 Corynocarpin, G-104
 Corynolamine, C-691
 Corynoline *O*-sulfate, C-692
 Corynoline, C-692
 Corynoloxine, C-693
 Corynoxine, R-90
 Corynoxidine, T-212
 Corynoxine A, R-90
 Corynoxine B, R-90
 Corynoxine, R-90
 Corypalline, T-158
 Corypalmine, C-694
 Coryphanthine, C-695
 Coryphenanthrine, M-559
 Coryphidine, C-696
 Coryphine, C-697
 Coryrutine, C-698
 Corysamine, C-699
 Corysolidine, R-9
 Corystewartine, C-700
 Corytenchine, X-30
 Corytenchirine, C-701
 Corytensine, E-43
 Corytuberine methosalt, C-702
 Corytuberine, C-702
 Coryximine, C-676
 Coryzeylamine, C-703
 Cosaldon, T-355
 Coscinamide A, C-704
 Coscinamide B, C-704
 Coscinamide C, C-704
 Cossonidine, H-162
 Cossonine, H-185
 Costaclavine, F-49
 Costaricine, D-88
 Costinone A, C-705
 Costinone B, C-706
 Cotarnine, N-292
 Cotarnoline, C-707
 Cotinine fumarate, C-708
 Cotinine, C-708
 Coulteroberbinone, C-709
 Coulteropine, P-667
 Coumamylin, C-711
 Coumaperine, C-710
m-Coumaric acid, H-690
p-Coumaric acid, H-691
*N*¹-*cis-p*-Coumaroylagmatine, A-719
*N*¹-*trans-p*-Coumaroylagmatine, A-719
 4-*O-p*-Coumaroylbetonicine, H-476
*N*⁵-*p*-Coumaroyl-*N*¹,*N*¹⁰-diferuloylspermidine, S-398
*N*⁵-*trans-p*-Coumaroyl-*N*¹,*N*¹⁴-di-*trans*-feruloylspermidine, S-399
N-cis-p-Coumaroyl-DOPA, A-745
N-trans-p-Coumaroyl-DOPA, A-745
*N*¹-*p*-Coumaroyl-*N*⁴-feruloylputrescine, D-362
N-cis-(p-Coumaroyl)glutamic acid, G-107
 1-Coumaroyl-2-hydroxyputrescine, D-248
N-E-p-Coumaroyl-5-methoxytryptamine, M-288
N-Z-p-Coumaroyl-5-methoxytryptamine, M-288
 5-*O-p*-Coumaroylnigrumin, H-509
N-cis-p-Coumaroyloctopamine, O-63
N-trans-p-Coumaroyloctopamine, O-63
N-trans-p-Coumaroyl-2-phenylethylamine, P-342
N-p-Coumaroylpipecolic acid, P-456

- 4'-*O*-*p*-Coumaroylprunasin, H-673
 4-Coumaroylputrescine, C-456
N^b-*cis*-*p*-Coumaroylserotonin, H-755
N^b-*p*-Coumaroylserotonin, H-755
 4-[*N*-(*p*-Coumaroyl)serotonin-4'-yl]-*N*-feruloylserotonin, B-126
p-Coumaroylspermidine, S-398
N^b-*p*-Coumaroyltryptamine, T-639
N-*p*-*cis*-Coumaroyltyramine, H-667
N-*p*-*trans*-Coumaroyltyramine, H-667
 Coumermycin A₁, C-711
 Coumermycin A₂, C-712
 Coumermycin, C-711
 Coumidine, C-178
 Coumingaine, C-713
 Coumingidine, C-178
 Coumingine, C-178
 Courouppitine A, I-108
 Courouppitine B, I-73
 Coxylase, T-366
 Coyhaiquine, C-714
 Coyhaiquinine, C-714
 Cozymase II, T-366
 CP 35763, A-1151
 CP 41043, A-1152
 CP 41494, A-1153
 CP 20, D-632
 CP 47444, N-276
 CP 51467, N-48
 CP 52726, N-49
 CP 52748, N-50
 CP 57820, N-276
 CP 61405, R-136
 CP 46192, T-391
 CPC 1, A-626
 CPT, C-70
 Crabbine, I-208
 Crambene, H-659
 Crambescidic acid, P-774
 Crambescidin acid, C-716
 Crambescidin 359, C-715
 Crambescidin 431, C-716
 Crambescidin 800, C-717
 Crambescidin 816, C-717
 Crambescidin 826, C-718
 Crambescidin 830, C-719
 Crambescidin 844, C-720
 Crambescidin 657 methyl ester, N-115
 Crambescidin 657, N-115
 Crambescin A, C-721
 Crambescin B, C-722
 Crambescin C1, C-723
 Crambescin C2, C-724
 Crambidine, C-725
 Crambine A, C-721
 Crambine B, C-722
 Crambine C1, C-723
 Crambine C2, C-724
 Craniformin, D-229
 Craspidospermine, V-117
 Crassanine, C-726
 Crassicaudine, M-323
 Crassicaulidine, M-337
 Crassicauline A, M-312
 Crassicauline B, H-196
 Crassicauline II, H-196
 Crassicauline I, M-312
 Crassicaulisine, M-313
 Crassicausine, M-312
 Crassicautine, M-336
 Crassifolazonine, C-727
 Crassifoline methine, C-729
 Crassifoline, C-728
 Craugsodine, C-730
 Creatine, C-731
 Creatinine, C-732
 Creatonotine *N*-oxide, T-188
 Crebanine *N*-oxide, C-733
 Crebanine, C-733
 Cremastrine, H-629
 Crenatidine, D-686
 Crenatine A, D-838
 Crenatine, M-290
 Crenulatine, C-734
 Crepidamine, C-735
 Crepidine, C-736
 Criamide A, C-737
 Criamide B, C-737
 Criasbetaine, Z-12
 Criasiaticidine A, P-586
 Cribetamine, N-169
 Cribochaline A, H-3
 Cribrochalinamine oxide A, C-738
 Cribrochalinamine oxide B, C-739
 Cribronic acid, D-661
 Cribrostatin 1, A-757
 Cribrostatin 3, A-799
 Cribrostatin 5, A-799
 Cribrostatin 7, A-799
 Cribrostatin 6, C-740
 Cribrostatin 2, M-624
 Cribrostatin 4, R-42
 Crinafolidine, C-741
 Crinafoline, A-683
 Crinalbine, C-743
 Crinamabine, C-742
 Crinamidine, C-743
 Crinamine, H-5
 Crinan-1,2-diol, A-670
 Crinan-3-ol, C-748
 Crinasiadine, C-744
 Crinasiatine, C-745
 Crinatine, C-746
 Crinatine, C-747
 Crinidine, C-748
 Crinine, C-369
 Crinine, C-748
 Crinisine, C-749
 Crinosine, C-750
 Criocerine, V-117
 Criophylline, C-751
 Cripaline, P-579
 Cripowellin A, C-752
 Cripowellin B, C-753
 Crispanine, C-754
 Crispatine, F-168
 Crispine A, C-756
 Crispine B, C-756
 Crispine C, C-757
 Crispine D, C-758
 Crispine E, C-759
 Crispine, C-755
 Crispulidine, M-346
 Cristadine, P-80
 Cristovin, V-128
 Criwelline, T-53
 Croaegyptine, C-760
 Croalbidine, C-761
 Croalbinecine, D-597
 Crobarbatine, C-762
 Crobruhine, C-771
 Crocacin A, C-763
 Crocacin B, C-763
 Crocacin C, C-763
 Crocacin D, C-763
 Crocandine, C-764
 Croceaine A, B-264
 Croceaine B, B-264
 Croceaurine, C-765
 Crociflorine, C-766
 Crociflorinine, C-766
 Crociflorinone, J-43
 Cronaburmine, C-767
 Crooksidine, D-120
 Crooksiine, C-441
 Croomine, S-523
 Croomionidine, P-606
 Cropodine, C-768
 Crosemperine, C-769
 Crotafloquine, M-36
 Crotafoline, C-770
 Crotalaburmine, A-967
 Crotalarine lactone, C-772
 Crotalarine, C-771
 Crotaleschenine, C-773
 Crotanamine, C-774
 Crotanecine, C-775
 Crotastriatine, N-219
 Crotaverrine, S-245
 Crotoflorine, P-643
 Crotonine, C-776
 Crotonosine, P-643
 Crotsparine, P-643
 Crotsparinine, C-781
 Crucigasterin 275, A-847
 Crucigasterin 277, A-847
 Cruentaren A, C-777
 Cruentaren B, C-778
 Cruentine A, C-779
 Cruentine B, C-780
 Crychine, E-221
 Crykonisine, H-424
 Cryogenine, V-95
 Cryprochine, C-781
 Cryptadine A, C-782
 Cryptadine B, C-782
 Cryptanthine, D-607
 Cryptaustoline, C-783
 Cryptocavine, C-790
 Cryptocin, C-784
 Cryptocurine, K-103
 Cryptodorine, N-125
 Cryptoheptine, C-785
 Cryptolepicarboline, C-786
 Cryptolepine, I-109
 Cryptolepinoic acid, I-111
 Cryptolepinone, D-505
 Cryptomisirine, C-787
 Cryptopalmatine, M-734
 Cryptophorine, C-788
 Cryptophorinine, C-789
 Cryptopine, C-790
 Cryptopleuridine, C-791
 Cryptopleurine, C-792
 Cryptopleurospermine, C-793
 Cryptoquindoline, C-794
 Cryptosanguinolentine, I-110
 Cryptospirolepine, C-795
 Cryptostyline I, C-796
 Cryptostyline II, C-797
 Cryptostyline III, C-797
 Cryptotackeieine, Q-19
 Cryptowolidine, C-798
 Cryptowoline, C-798
 Cryptowolinol, C-798
 Crispatine, F-168
 Crystalloiodinine A, P-321
 Crystalloiodinine B, P-321
 Crystamidine, C-799
 CS 79A, C-169
 CS 79B, C-169
 CSA 3, A-472
 CSL 12A, T-700
 CSL 12B, T-700
 CSL 12C, T-700
 CSL 12D, T-700
 CSL 12E, T-700
 CSL 12F, T-700
 CSL 12G, T-700
 CSL 12H, T-700
 CSL 12I, T-700
 CSL 12J, T-700
 CSL 12K, T-700
 CSL 12L, T-700
 CSL 12M, T-700
 CT 2103, T-52
 Cuanzine, C-800
 Cuauchichicine, V-45
 Cucoline, S-313
 Cucubalactam, C-801
 Cucumopine, C-802
 Cucurbitoside L, H-414
 Culacorine, C-805
 Culantraramine *N*-oxide, C-803
 Culantraramine, C-803
 Culantraraminol *N*-oxide, C-803
 Culantraraminol, C-803
 Cularicine, C-804
cis-Cularidine *N*-oxide, C-805

- trans*-Cularidine N-oxide, C-805
 Cularidine, C-805
 Cularimine, C-805
cis-Cularine N-oxide, C-805
 Cularine, C-805
 Cultithalminine, T-340
m-Cumarinic acid, H-690
p-Cumarinic acid, H-691
 Cupreine, Q-20
 Curacautine, C-806
 Curacin A, C-807
 Curacin B, C-807
 Curacin C, C-807
 Curacin D, C-808
 Curane B, C-420
 16 α -Curan-17-ol, G-34
 C-Curarine I, C-809
 C-Curarine II, C-31
 C-Curarine III, N-301
 C-Curarine, C-809
 Curarine, C-420
 Curassaneceine, H-517
 Curassavine N-oxide, C-810
 Curassavine, C-810
 Curassavinine, C-810
 Curicycleatjenine, C-811
 Curicycleatjine, C-811
 Curine, C-812
 Currayangine, C-813
 Currayanine, C-814
 Curromycin A, C-815
 Curromycin B, C-815
 Curryangine, C-813
 Curryanine, C-814
 Curtisin, D-634
 Curvupallide A, C-816
 Curvupallide B, C-816
 Curvupallide C, C-816
 Cuscamidine, C-817
 Cuscohygrine, C-818
 Cusconidine, C-819
 Cusconine, C-820
 Cusculine, P-206
 Cuscutamine, H-237
 Cuskhygrine, C-818
 Cuspanine, P-206
 Cuspareine, C-821
 Cusparine, C-822
 Cuspidaline, D-88
 CV 1, A-1154
 CW 69A, C-129
 CW 69B, C-129
 CW 69, C-129
 CW 69C, C-129
 CW 69D, C-129
 CW 69E, C-129
 CW 69F, C-129
 CX 59, H-480
 CY 39, H-480
 Cyanamide, C-823
 5-(ω -Cyanoalkyl)-2-pyrrolicarboxaldehydes, F-141
 4-Cyano-1,3-benzenediol, D-535
 2-Cyano-2-butanol, H-572
 1-Cyano-3-buten-2-ol, H-659
 3-Cyanocarbazole, C-122
 1-Cyano- β -carbolone, C-132
 3-(Cyanocarbonyl)indole, O-182
 3S-Cyanocoronaridine, C-657
 3-Cyano-N-[6-[[[(3-cyano-1-oxobutyl)amino]-methyl]tetrahydro-2-hydroxy-2H-pyran-3-yl]butanamide, A-1260
 Cyanocycline A, C-830
 Cyanocycline B, C-830
 Cyanocycline C, C-830
 Cyanocycline D, C-824
 Cyanocycline F, C-830
 5-Cyano-7-deazaguanosine, C-7
 5-Cyano-2,3-dibromopyrrole, D-339
 4-Cyano-1,2-dimethoxybenzene, D-706
 6-Cyano-1,3-dimethyluracil, D-669
 5-(12-Cyano-6-dodecenyloxy)-2-pyrrolicarboxaldehyde, C-825
 5-(20-Cyanoicosyloxy)-1H-pyrrole-2-carboxaldehyde, F-141
 N-(2-Cyanoethyl)glycine, C-147
 3-(2-Cyanoethyl)indole, I-100
 2-(2-Cyanoethyl)-3-isoxazolin-5-one, O-185
 5-(21-Cyanoheptacosyloxy)-1H-pyrrole-2-carboxaldehyde, F-141
 5-(17-Cyanoheptadecyloxy)-1H-pyrrole-2-carboxaldehyde, F-141
 3-Cyano-2-hydroxy-4-methoxypyridine, D-668
 3-Cyano- μ -hydroxy-3-methyloxiranetridecanoic acid, A-1318
 3-Cyano-2-hydroxymethyl-2-propen-1-ol, H-510
 3-Cyano-4-hydroxy-1-methyl-2-pyridone, D-668
 N-[1-Cyano-2-(4-hydroxyphenyl)ethenyl]formamide, C-826
 3-(2-Cyano-3-hydroxyphenyl)-2-hydroxy-N-[2-(4-hydroxyphenyl)ethenyl]-2-propenamide, B-257
 2-Cyano-3-hydroxyquinoline, H-724
 5-(23-Cyano-23-hydroxy-6-tricosenyloxy)-1H-pyrrole-2-carboxaldehyde, C-827
 3-Cyanoindole, I-90
 4-Cyanoindoloquinolizidine, O-49
 2-[Cyano(3-indolyl)methylene]-3-indolone, D-487
 Cyanoiodomethane, I-163
 4-Cyanoisoxazole, I-346
 2-Cyano-4-methoxyaniline, A-789
 2-Cyano-6-methoxyaniline, A-787
 3-Cyano-5-methoxyaniline, A-790
 5-Cyano-6-methoxyindolo[2,3-*a*]carbazole, C-828
 5-Cyano-6-methoxy-11-methylindolo[2,3-*a*]carbazole, C-828
 3-Cyano-4-methoxy-1-methyl-2-pyridone, R-96
 6-Cyano-4-methoxy-5-(methylthio)-2,2'-bipyridine, C-576
 3-Cyano-4-methoxypyridine 1-oxide, H-709
 3-Cyano-4-methoxypyridine, H-709
 3-Cyano-6-methoxypyridine, H-711
 3-Cyano-4-methoxy-2-pyridone, D-668
 4-Cyano-6-methoxyquinoline, H-726
 4-(Cyanomethyl)-1,3-benzenediol, D-646
 (Cyanomethyl)benzene, P-336
 5-Cyano-7-(3-*O*-methyl-5-deoxy- β -D-erythro-pent-4-enofuranosyl)pyrrolo[2,3-*d*]pyrimidin-4-one, M-778
 4-Cyano-1,2-methylenedioxybenzene, M-442
 2-Cyanomethylene-1-glucosyloxy-3-hydroxy-4,5-dimethoxycyclohexane, S-302
 2-(Cyanomethylene)-1,3-propanediol, H-510
 2-(Cyanomethyl)-6-ethoxypyridine, H-708
 3-(Cyanomethyl)-4-hydroxyindole, H-524
 3-Cyanomethyl-3-hydroxy-1H-indol-2(3H)-one, C-829
 3-Cyanomethyl-3-hydroxyoxindole, C-829
 3-Cyano-1-methyl-1H-indole, I-90
 3-Cyanomethyl-1H-indole, I-81
 3-Cyano-5-methylisoxazole, M-479
 3-(Cyanomethyl)-4-methoxyindole, H-524
 3-(Cyanomethyl)-6-methoxyindole, H-526
 2-Cyanomethyl-1-methylpyrrolidine, P-950
 N'-Cyano-N'-methylmoloaiamine, M-679
 4-(Cyanomethyl)phenol, H-672
 3-Cyanomethyl-6-prenylindole, H-497
 1-Cyano-2-methyl-2-propanol, H-573
 3-Cyano-2-methyl-2-propen-1-ol, H-576
 3-Cyano-1-methyl-4(1H)-pyridinone, H-709
 5-Cyano-1-methyl-2(1H)-pyridinone, H-711
 3-Cyano-1-methyl-2-pyridone, R-95
 5-Cyano-2-methylpyrimidine, M-539
 2-Cyano-1-methyl-1H-pyrrole, P-944
 1-Cyano-4-(methylthio)butane, M-203
 Cyanomycin, P-833
 Cyanonaphthyridinomycin, C-830
 5-(19-Cyano-nonadecyloxy)-1H-pyrrole-2-carboxaldehyde, F-141
 9-Cyano-nonyl methyl sulfone, M-563
 1-Cyanophenazine, P-319
 1-Cyano-2-phenylethane, P-350
 Cyanophenylmethane, P-336
 3-(Cyanophenylmethylene)-5-(3,7-dimethyl-2,6-oc-tadienyl)-2,3,5,10-tetrahydro-2-oxo-1-phenazinecarboxylic acid, B-65
 5-Cyano-1-phenyl-1H-pyrazole, P-890
 3-Cyano-1-phenyl-1H-pyrazole, P-890
 2-Cyano-1-phenyl-1H-pyrrole, P-944
 2-Cyano-2-propanol, H-617
 N-Cyano-sec-pseudobrucine, I-12
 N-Cyano-sec-pseudostrychnine, I-12
 Cyanopterin, A-802
 3-Cyanopyrazole, P-890
 2-Cyanopyrrole, P-944
 4-Cyanoquinoline, Q-32
 Cyanoquinoline, S-2
 4-Cyanoresorcinol, D-535
 6-Cyano-6,7-secostemmadenine, Y-21
 5-(23-Cyano-6-tricosenyloxy)-1H-pyrrole-2-carboxaldehyde, C-827
 Cyanotubercidin, S-53
 6-Cyanouracil, D-669
 Cyathocaline, T-565
 CYB 3, H-717
 Cycasindene, A-698
 Cycasin, M-397
 Cycasthioamide, C-831
 Cycemanine, C-677
 Cyclamenol, C-832
 Cyclamidomycin, P-876
 α -Cyclanoline, C-833
 Cyclanoline, C-833
 β -Cyclanoline, C-833
 Cyclebarbatine, I-328
 Cycleacurine, C-812
 Cycleadrine, T-286
 Cycleahomine, T-286
 Cycleaneonine, C-834
 Cycleanine N²-oxide, C-835
 Cycleanine, C-835
 Cycleanorine, T-286
 Cycleapeltine, R-46
 Cycleatjeheneine, C-836
 Cycleatjehine, C-836
 γ -Cyclic diglutamic acid, D-803
 Cyclizidine, C-837
 Cyclo(alanylalanyl), D-769
 Cyclo(alanyl-4-hydroxypropyl), C-838
 Cyclo(alanylpropyl), C-839
 Cycloanchinopeptolide C, C-840
 Cycloanthranilylproline, D-502
 Cyclo(arginylprolyl), C-841
 2,20-Cycloaspidospermidine, T-674
 Cyclobaleabuxine, C-883
 Cyclobastadin 1, B-32
 Cyclobastadin 2, B-32
 Cyclobastadin 3, B-32
 Cyclobastadin 4, B-32
 Cyclobassinin sulfoxide, C-842
 Cyclobassinin, C-842
 Cyclobassinone, C-843
 Cyclobullatine A, C-846
 Cyclobuxamidine, B-474
 Cyclobuxamine A, C-846
 Cyclobuxamine H, C-846
 Cyclobuxamine I, C-846
 Cyclobuxamine, C-846
 Cyclobuxaphylamine, C-844
 Cyclobuxargentine G, C-845
 Cyclobuxidine F, C-883
 Cyclobuxine A, C-846
 Cyclobuxine B, C-846
 Cyclobuxine D, C-846
 Cyclobuxine I, C-846
 Cyclobuxine, C-846
 Cyclobuxomicreine K, C-882
 Cyclobuxomicreine, C-882
 Cyclobuxomicreinein, C-847
 Cyclobuxophylline K, C-848
 Cyclobuxophylline M, C-848
 Cyclobuxophylline O, C-848
 Cyclobuxophylline, C-848
 Cyclobuxophyllinine M, C-848
 Cyclobuxophyllinone, C-848
 Cyclobuxosuffrine K, C-912
 Cyclobuxosuffrine, C-912
 Cyclobuxoviramine, C-849
 Cyclobuxoviricine, C-850
 Cyclobuxoviridine B, C-850

- Cyclobuxoviridine L, C-850
 Cyclobuxoviridine, C-850
 Cyclobuxoxazine A, C-851
 Cyclobuxoxazine C, C-851
 Cyclobuxoxazine, C-851
 Cyclobuxoxine, B-482
 Cyclobuxupaline C, C-902
 Cyclocapitelline, C-433
 Cyclocarbamide A, C-852
 Cyclocarbamide B, C-852
 Cyclocelabenzine, C-853
 Cyclocide, C-962
 Cyclocinamide A, C-854
 Cyclocinamide B, C-854
 Cycloclausenamide, C-855
 Cycloclavine, C-856
 Cyclocreatine, A-731
 Cyclo(dehydroprolyltryptophyl), C-899
 Cycloclercitine, C-857
 Cyclo(2,3-didehydroprolyl)[2-(1,1-dimethyl-2-propenyl)tryptophyl], C-861
 Cyclo(didemnamide B, C-859
 Cyclo(didemnamide, C-858
 Cyclo(didemniserinol, C-860
N,α',N',α-Cyclodi(4-methylaminobenzeneethanol), D-745
 Cyclo[2-(1,1-dimethyl-2-propenyl)tryptophyl]prolyl, C-861
 Cyclo(dopa glucoside, C-862
 Cyclo(dopa, C-862
 14,18-Cyclo-21,23-epoxyaprotirucallane-3,7,21,24,25-pentol, C-863
 Cyclofoveoglin, C-864
 Cyclo(glycylprolyl), C-865
 Cycloguineense A, C-866
 Cycloguineense B, C-866
 Cyclohaliclonamines, C-867
 Cycloheptaphylline, M-757
 Cycloheptazoline, M-757
 Cyclohexanamine, C-868
 Cyclohexaneethanamine, C-869
 Cycloheximide, D-765
 Cyclohexylamine, C-868
 2-Cyclohexylethylamine, C-869
*O*⁶-Cyclohexylmethylguanidine, G-210
 Cyclo(homoleucylprolyl), C-870
 Cyclo(4-hydroxyprolylleucyl), C-871
 Cyclo(2-hydroxyprolylphenylalanyl), C-872
 Cyclo(4-hydroxyprolylphenylalanyl), C-873
 Cyclo(4-hydroxyprolyltyrosyl), C-874
 Cyclo(isoleucylprolyl), C-875
 Cyclokoreanine A, C-917
 Cyclokoreanine B, C-917
 Cyclo(leucylprolyl), C-876
 Cyclomahanimbine, C-814
 Cyclomalayanine B, C-915
 Cyclomalayanine, C-915
 Cyclomarazine A, C-877
 Cyclomarazine B, C-877
 Cyclomegistine B, C-879
 Cyclomegistine, C-878
 Cyclo(methionylprolyl), C-880
 Cyclomethoxazine B, C-881
 Cyclomethyldecylprodiginine, M-429
 Cyclomicrobuxamine, B-482
 Cyclomicrobuxeine K, C-882
 Cyclomicrobuxeine, C-882
 Cyclomicrobuxine, B-482
 Cyclomicrobuxinine, B-482
 Cyclomicrophyllidine A, C-883
 Cyclomicrophylline A, C-883
 Cyclomicrophylline B, C-883
 Cyclomicrophylline C, C-883
 Cyclomicrophylline I, C-883
 Cyclomicrosine C, C-883
 Cyclomicrosine, C-883
 Cyclomicuranine L, C-884
 Cyclomikuranine, C-884
 Cyclomycin, A-832
 Cyclomyrsinol, C-885
 Cycloleosamandione, C-886
 Cyclononylprodiginine, N-285
 Cyclononylprodigosin, N-285
 Cycloornithine, A-872
 Cycloooridin, C-887
 Cyclopamine, J-36
 Cyclopapilosine D, C-888
 Cyclopapilosine I, C-888
 Cyclophenin, C-889
 Cyclophenol, C-889
 Cyclopeptide, C-890
 Cyclo(phenylalanylprolyl), C-891
 Cyclopiamide, C-892
 Cyclopiamine A, C-893
 Cyclopiamine B, C-893
 Cyclopiazonic acid imine, C-895
 β-Cyclopiazonic acid, B-218
 α-Cyclopiazonic acid, C-894
 Cyclopiiperstachine, C-896
 Cycloposine, J-36
 Cycloprodigosin, C-897
 Cyclo(prolylglycyl), C-865
 Cyclo(prolylmethionyl), C-880
 Cyclo(prolyl-3-nitrotyrosyl), C-900
 Cyclo(prolylprolyl), C-898
 Cyclo(prolyltryptophyl), C-899
 Cyclo(prolyltyrosyl), C-900
 Cyclo(prolylvalyl), C-901
 Cycloprotobuxinamine, C-883
 Cycloprotobuxine A, C-902
l-Cycloprotobuxine C, C-902
 Cycloprotobuxine C, C-902
 Cycloprotobuxine D, C-902
 Cycloprotobuxine F, C-902
 Cycloprotobuxine I, C-902
 Cycloprotobuxoline I, C-903
 Cyclorocaglamide, C-904
 Cyclorolfeibuxine A, C-883
 Cyclorolfeibuxine C, C-883
 Cyclorolfeine, C-905
 Cyclorolfoxazine, C-906
 2,11-Cyclo-11,12-secoaspidospermidine, M-188
 12,16-Cyclo-1,12-secoaphnan-23-oic acid, H-346
 18,28-Cyclo-16,28-secosolanid-5-en-3-ol, P-633
 Cycloserine, A-832
 Cycloshermilamine D, C-907
 Cycloshermilamine, C-908
 Cyclostachine A, C-909
 Cyclostachine B, C-909
 Cyclostelletamines, C-910
 Cyclostrychnine, C-911
 Cyclovirobuxine K, C-912
 Cyclosuffrobuxine, C-912
trans-Cyclosuffrobuxinine M, C-912
 Cyclosuffrobuxinine M, C-912
 Cyclosuffrobuxinine, C-912
 Cyclothiazomycin, C-913
 Cyclotryprostatin A, C-914
 Cyclotryprostatin B, C-914
 Cyclotryprostatin C, C-914
 Cyclotryprostatin D, C-914
 Cyclo(tryptophanylprolyl), C-899
 Cyclo(tryptophylprolyl), C-899
 Cyclovirobuxeine A, C-915
 Cyclovirobuxeine B, C-915
 Cyclovirobuxeine F, C-916
 Cyclovirobuxeine I, C-915
 Cyclovirobuxine A, C-917
 Cyclovirobuxine C, C-917
 Cyclovirobuxine D, C-917
 Cyclovirobuxine F, C-917
 Cyclovirobuxine I, C-917
 3,5'-Cycloxanthosine, C-918
 Cycloxazoline, C-919
 Cycloxobuxidine F, B-12
 Cycloxobuxidine I, B-12
 Cycloxobuxine I, C-920
 Cycloxobuxoline I, C-921
 Cycloxobuxoviricine, C-884
 Cycloxobuxoxazine C, C-851
 Cycloxyisoschelhammericine, S-125
 Cyclozoanthosine, C-922
 Cygnine, C-923
 Cyldramide, C-924
 Cyldricine A, C-926
 Cyldricine B, C-925
 Cyldricine C, C-926
 Cyldricine D, C-926
 Cyldricine E, C-926
 Cyldricine F, C-926
 Cyldricine G, C-927
 Cyldricine H, L-102
 Cyldricine I, L-102
 Cyldricine J, C-928
 Cyldricine K, C-926
 Cyldrocarpine, C-929
 Cyldrocarpine, C-929
 Cyldrocarpine, C-929
 Cyldrocarpinol, L-177
 Cyldropermopsin, C-930
 Cynaustine, A-669
 Cynaustaline, T-443
 Cynodine, C-931
 Cynoglossamine, E-12
 Cynoglossophine, H-94
 Cynolujine, C-932
 Cynometrine, C-933
 Cynthichlorine, C-934
 Cypholophine, C-935
 Cyphoplectine, M-352
 Cypridina luciferin, L-281
 Cypridina oxyluciferin, O-222
 Cyprino-Pourpre B, A-752
 Cystamidin A, P-948
 Cystaminesulfinic acid, A-764
Aspergillus oryzae Cysteine protease inhibitors, C-936
 Cysteinolic acid, A-815
 4-*S*-Cysteinyltetradotoxin, T-298
 Cystodamine, A-1459
 Cystodytin A, D-97
 Cystodytin B, D-97
 Cystodytin C, D-97
 Cystodytin D, D-97
 Cystodytin E, D-97
 Cystodytin F, D-97
 Cystodytin G, D-97
 Cystodytin H, D-97
 Cystodytin I, D-97
 Cystodytin J, D-97
 Cystodytin K, D-97
 Cystothiazole A, M-177
 Cystothiazole B, M-177
 Cystothiazole C, M-177
 Cystothiazole D, M-177
 Cystothiazole E, C-937
 Cystothiazole F, M-177
 Cystothiazole G, M-181
 L-Cytidine, C-961
 Cytisine *N*-oxide, C-938
 12-Cytisineacetamide, C-939
 12-Cytisineacetic acid, C-939
 Cytisine, C-938
 5-(12-Cytisinylmethyl)-6-hydroxylupanine, C-940
 Cytiton, C-938
 Cytoblastin, C-941
 Cytochalasin A, C-943
 Cytochalasin B, C-943
 Cytochalasin C, C-944
 Cytochalasin D, Z-37
 Cytochalasin E, C-945
 Cytochalasin F, C-946
 Cytochalasin G, C-947
 Cytochalasin H, Z-37
 Cytochalasin J, Z-37
 Cytochalasin K, C-948
 Cytochalasin K, C-949
 Cytochalasin L, C-950
 Cytochalasin M, C-951
 Cytochalasin N, C-944
 Cytochalasin N, C-953
 Cytochalasin N_{phos}, C-953
 Cytochalasin O, C-952
 Cytochalasin O, C-953
 Cytochalasin O_{phos}, C-953
 Cytochalasin P, C-952
 Cytochalasin P_{phos}, C-952
 Cytochalasin Q, C-954

- Cytochalasin R, C-954
 Cytochalasin S, C-955
 Cytochalasin T, C-943
 Cytochalasin U, A-1196
 Cytochalasin U, C-956
 Cytochalasin V, C-942
 Cytochalasin W, C-957
 Cytochalasin X, C-958
 Cytochalasin Y, C-958
 Cytochalasin Z, C-947
 Cytochalasin Z₁, C-943
 Cytochalasin Z₂, C-943
 Cytochalasin Z₃, C-943
 Cytochalasin Z₄, C-943
 Cytochalasin Z₅, C-943
 Cytochalasin Z₆, C-946
 Cytochalasin Z₇, C-959
 Cytochalasin Z₈, C-959
 Cytochalasin Z₉, C-959
 Cytokinin B, A-137
 Cytolipin, M-485
 Cytomycin, C-960
 Cytosine arabinoside, C-962
 Cytosine, C-961
 Cytosporone A, C-963
 Cytosporone B, C-963
 Cytosporone F, C-963
 Cytosporone G, C-963
 Cytosporone H, C-963
 Cytosporone I, C-963
 Cytostal, D-92
 Cytosylglucuronic acid, C-961
 Cytozoxone, C-964
 D 4028, X-5
 DAB-1, D-593
 Dactylamide A, D-1
 Dactylamide B, D-2
 Dactylicapnosine, D-3
 Dactylicapnosinine, D-4
 Dactylidine, D-5
 Dactyline, D-6
 Daechualkaloid A, M-77
 Daechualkaloid C, L-193
 Daechualkaloid E, D-531
 Daechucyclopeptide I, P-25
 Daechuine S27, A-942
 Daechuine S3, D-7
 Daechuine S7, D-8
 Daechuine S8-1, D-9
 Daechuine S1, F-150
 Daechuine S2, F-151
 Daechuine S4, F-149
 Daechuine S5, M-189
 Daechuine S9, M-720
 Daechuine S10, N-338
 Daechuine S26, P-25
 Daechuine S6, P-25
 Daijisong, C-812
 Dakaramine, D-10
 Dalomycin T, D-11
 Damascenic acid, A-787
 Damascenic acid, A-787
 Damascenine, A-787
 Damascinine, D-256
 Damazor, A-736
 Dambullin, D-12
 Daminine, A-193
 Damipipecoline, H-697
 Damirone A, D-13
 Damirone B, D-13
 Damirone C, D-13
 Damituricine, H-476
 Danaidal, D-14
 Danaidone, L-263
 Danguyelline, P-209
 Dankastatin A, D-15
 Dankastatin B, D-15
 Danomycin, D-16
 Danoxamine, D-16
 Danshensuan A, D-648
 Danshensu, D-648
 Danuphylline A, D-17
 Danuphylline, D-17
 Daphcalycic acid, D-18
 Daphcalycine, D-19
 Daphcalycinosidine A, D-18
 Daphcalycinosidine B, D-20
 Daphcalycinosidine C, D-55
 Daphgraciline, D-50
 Daphgracine, D-51
 Daphlongeramine A, D-21
 Daphlongeramine A, D-22
 Daphlongeramine B, D-23
 Daphmacrine, D-24
 Daphmacropodine, D-24
 Daphmacropodosidine A, D-20
 Daphmacropodosidine B, D-20
 Daphmanidine A, D-25
 Daphmanidine B, D-26
 Daphmanidine C, D-27
 Daphmanidine D, D-27
 Daphmanidine E, D-25
 Daphmanidine F, D-25
 Daphnandrine, O-1
 Daphnan-23-oic acid, H-330
 Daphnarcine, D-28
 Daphnezomic acid, D-51
 Daphnezomine A, D-29
 Daphnezomine B, D-29
 Daphnezomine C, D-30
 Daphnezomine D, D-31
 Daphnezomine E, D-68
 Daphnezomine F, D-32
 Daphnezomine G, D-32
 Daphnezomine H, D-53
 Daphnezomine I, D-53
 Daphnezomine J, D-33
 Daphnezomine K, D-33
 Daphnezomine L, D-34
 Daphnezomine M, D-35
 Daphnezomine N, D-35
 Daphnezomine O, D-36
 Daphnezomine P, D-37
 Daphnezomine Q, D-38
 Daphnezomine R, D-51
 Daphnezomine S, D-53
 Daphnicadine, D-39
 Daphnicaline, D-40
 Daphnicamine, D-41
 Daphnicyclidine A, D-42
 Daphnicyclidine B, D-42
 Daphnicyclidine C, D-42
 Daphnicyclidine D, D-43
 Daphnicyclidine E, D-43
 Daphnicyclidine F, D-43
 Daphnicyclidine G, D-43
 Daphnicyclidine H, D-44
 Daphnicyclidine J, D-45
 Daphnicyclidine K, D-46
 Daphnicyclidine L, D-47
 Daphniglaucine A, D-48
 Daphniglaucine B, D-48
 Daphniglaucine C, D-49
 Daphniglaucine D, D-55
 Daphniglaucine E, D-55
 Daphniglaucine F, D-55
 Daphniglaucine G, D-55
 Daphniglaucine H, D-55
 Daphniglaucine J, D-33
 Daphniglaucine K, D-33
 Daphnigraciline, D-50
 Daphnigracine, D-51
 Daphnijsmine, D-33
 Daphnilactone A, D-52
 Daphnilactone B *N*-oxide, D-53
 Daphnilactone B, D-53
 Daphnilongeramine A, D-54
 Daphnilongeramine B, D-69
 Daphnilongeramine C, D-55
 Daphnilongeramine D, D-68
 Daphnilongeridine, D-63
 Daphnilongerine, D-56
 Daphnilongertone, D-57
 Daphnimacrine, D-58
 Daphnimacropine, D-68
 Daphnimacropodine A, D-59
 Daphnimacropodine B, D-49
 Daphnimacropodine C, D-49
 Daphnimacropodine D, D-53
 Daphnine, D-60
 Daphnioldhanine A, D-61
 Daphnioldhanine B, D-61
 Daphnioldhanine C, D-62
 Daphnioldhanine D, D-63
 Daphnioldhanine E, D-63
 Daphnioldhanine F, D-63
 Daphnioldhanine G, D-63
 Daphnioldhanine H, D-63
 Daphnioldhanine I, D-68
 Daphnioldhanine J, D-64
 Daphnioldhanine K, D-56
 Daphnipaxianine A, D-65
 Daphnipaxianine B, D-65
 Daphnipaxianine C, D-69
 Daphnipaxianine D, D-66
 Daphnipaxinine, D-67
 Daphniphyllamine, D-68
 Daphniphyllidine, D-68
 Daphniphylline *N*-oxide, D-68
 Daphniphylline, D-68
 Daphnitejsmine, S-181
 Daphniyunnine A, D-55
 Daphniyunnine B, L-246
 Daphniyunnine C, L-245
 Daphniyunnine D, D-69
 Daphniyunnine E, D-69
 Daphnoline, O-1
 Daphtenidine A, D-70
 Daphtenidine B, D-71
 Daphtenidine C, D-72
 Daphtenidine D, D-33
 Dapiramicin A, D-73
 Dapiramicin B, D-73
 Darcyribeirine, R-51
 Dardanin, D-730
 Darienine, T-565
 Darlingianine, D-74
 Darlingine *N*-oxide, D-75
 Darlingine, D-75
 Darlinine, D-76
 Darlucin A, D-77
 Darlucin B, D-78
 Darvasamine, D-79
 Darvasine, D-80
 Darvasoline, D-81
 Daryamide A, D-82
 Daryamide B, D-82
 Daryamide C, D-82
 Dasycarine, D-83
 Dasycarpamine, P-625
 Dasycarpidan-1-methanol, U-8
 Dasycarpidan-1-one, D-84
 Dasycarpidol, D-84
 Dasycarpidone, D-84
 Dasycarpine, D-85
 Dasycarponilide, L-213
 Dasycarponin, L-213
 Dasyclamide, C-456
 Dasymachaline α -*N*-oxide, D-948
 Dasymachaline, D-948
 Dasyrachine, D-86
 Datumetine, M-393
 Daturamine, A-1043
 Daturine, T-632
 Daucine, D-87
 Dauricicoline, D-88
 Dauriciline, D-88
 Dauricine, D-88
 Dauricinoline, D-88
 Dauricoline, D-88
 Dauricoside, X-30
 Dauricumidine, A-134
 Dauricumine, A-134
 Daurine, Q-47
 Daurinoline, D-88
 Daurioxoisoporphine A, T-697
 Daurioxoisoporphine B, D-89
 Daurioxoisoporphine C, D-89
 Daurioxoisoporphine D, M-199

- Dauriporphine, D-90
 Dauriporphinoline, D-90
 Daurisoline, D-88
 Davallioside A, D-91
 Davallioside B, D-91
 Davidisine A, M-309
 Davidisine B, M-309
 Davisine, H-162
 Davisinol, H-168
 DBP-Br₂Cl, P-201
 DBP-Br₃Cl₃, T-476
 DC 102, A-1157
 DC 105, A-1158
 DC 81, A-1156
 DC 8118A, A-1159
 DC 88A, D-952
 DC 89A₁, D-953
 DC 89A₂, D-954
 DC 59A, S-231
 DC 8118B, A-1159
 DC 89B₁, D-953
 DC 86C, H-496
 DC 113, D-956
 DC 52d, Q-22
 DC 59H, S-231
 DC 107, L-82
 DC 86M, H-496
 DC 62, P-660
 DC 52, Q-21
 DC 86R, H-496
 DC 86Y, H-496
 2096D, D-522
 2'-Deacetoxyaustrospicatin, T-48
 2'-Deacetoxyaustrotaxine, P-224
 Deacetoxy-*trans*-1'-epierioaustralasine hydrate, E-180
 Deacetoxy-*trans*-erioaustralasine hydrate, E-180
 Deacetoxy-*trans*-erioaustralasine, E-179
 Deacetoxyleurosine, L-162
 5-Deacetoxy-5-oxosenampeline G, S-233
 7-Deacetoxytaxuspine J, A-7
 Deacetoxyvinblastine, V-110
 Deacetylglaine A, A-199
 14-Deacetylajadine, M-309
 Deacetylakuumilline, A-231
 Deacetylambiguine, M-309
 14-Deacetyl andersoline, G-91
 Deacetylanisomycin, A-1044
 Deacetylaspidospermine, A-1499
 7-Deacetylaustrospicatin, T-48
 2'-Deacetylaustrospicatin, T-48
 2'-Deacetylaustrotaxine, P-224
 14-Deacetyl-18-benzoyl-18-demethylpubescenine, M-309
 Deacetyl-*O*-benzoyl-5,10-dimethoxyakuammiline, A-231
 6-Deacetyl bicoloridine, M-347
 Deacetylbowdensine, B-261
 Deacetyl cathovaline, C-213
 10-Deacetylcephalomannine, T-52
N-Deacetylcolchicine, D-92
N-Deacetylcolchicine, D-92
 Deacetylcytochalasin C, C-944
 Deacetylcytochalasin H, Z-37
 Deacetyldaphnijsmine, D-33
 Deacetyldeformylpicraline, P-417
 10-Deacetyl-10-dehydro-7-acetyl taxol A, T-52
 Deacetyl delphinine, M-312
 Deacetylde-*O*-methylanisomycin, A-1044
N-Deacetyl-2-demethyl-*N*-formylcolchicine, D-92
N-Deacetyl-3-demethyl-*N*-formylcolchicine, D-92
 14-Deacetyl-18-demethylpubescenine, M-309
 Deacetyl diaboline, C-115
N-Deacetyl-2,3-didemethylcolchicine, D-92
 Deacetyl-1,2-β-dihydroakuammiline, A-231
 Deacetyl-1,2-dihydro-5-oxoakuammiline, A-231
 Deacetyl-5β,11-dimethoxyakuammiline, A-231
N-Deacetyl-6,7-di-*O*-methylisoepicosidic acid, I-167
N-Deacetyl-*N,N*-dimethyl-β-lumicolchicine, L-287
N-Deacetyl-*N,N*-dimethyl-γ-lumicornigerine, A-382
 Deacetyl eutherobin, E-64
 De-*O*-acetyl-1-epiaconitine, M-331
 Deacetylepipachysamine A, P-601
 Deacetyl-6-epipubescenine, M-309
 10-Deacetyl-7-epitaxol, T-52
 Deacetyl-19,20-epoxycytochalasin C, C-944
 Deacetylepoxycytochalasin H, E-129
 Deacetyl-19,20-epoxycytochalasin Q, C-954
 6-*O*-Deacetylleuonine, P-252
 8-*O*-Deacetylleuonine, P-252
 2-*O*-Deacetylleuonine, P-252
*O*²-Deacetylleuonine, P-248
 Deacetyl fawcettiine, F-29
N-Deacetylflnaconitine, A-94
N-Deacetyl-*N*-formylcolchicine, D-92
N-Deacetyl-*N*-formyl-2-*O*-demethyl-β-lumicolchicine, L-287
N-Deacetyl-*N*-formyl-β-lumicolchicine, L-287
N-Deacetyl-*N*-formyl-γ-lumicolchicine, L-287
 20-Deacetyl-20-formyl-17-methylalstonisine, A-648
 Deacetylgeyerline, G-91
 Deacetylghanamisine, H-195
 Deacetylhetochlorin, H-84
 Deacetylhemslayanine, M-303
 Deacetylheterophylloidine, H-161
 10-Deacetyl-10-(β-hydroxybutyryl)cephalomannine, T-52
 10-Deacetyl-10-(β-hydroxybutyryl)taxol B, T-52
 10-Deacetyl-10-(3-hydroxybutyryl)taxol, T-52
 6-Deacetyl-10-hydroxyperegrine, M-322
 Deacetylhypanitine, M-343
 2-*De*-*O*-acetylhyponine C, P-247
 Deacetylisoepicoside, I-167
 14-Deacetyl-14-isobutyrylajadine, M-309
 14-Deacetyl-14-isobutyrylinducalaine, G-91
 Deacetylisoepicoside, I-167
N-Deacetylkuanoniamine D, D-93
N-Deacetyl lappaconitine, A-91
 Deacetyl lutessine, L-310
 Deacetyllycloclavine, L-322
 Deacetyllythramine, L-378
*N*²-Deacetyl-*N*²-(3-mercapto-1-oxopropyl)maytansine, M-135
 Deacetylmetaplexigenin, P-223
O-Deacetyl-12-methoxyvinorine, V-194
 2'-*De*(acetylmethylamino)maytansine, M-136
 3-*De*[2-(acetylmethylamino)-1-oxopropoxy]-3-hydroxymaytansine, M-136
N-Deacetyl-*N*-methylcolchicine, D-92
N-Deacetyl-6-*O*-methylisoepicosidic acid, I-167
N-Deacetyl-6-*O*-methylisoepicosidic acid, I-167
N-Deacetyl-7-*O*-methylisoepicosidic acid, I-167
 14-Deacetylinducalaine, G-91
N-Deacetyl-*N*-3-oxobutyrylcolchicine, D-92
 10-Deacetyl-10-oxo-7-epicephalomannine, T-52
 10-Deacetyl-10-oxo-7-*epi*-taxol, T-52
 8-Deacetyl-8-palmitoyl vilmorrianine C, M-323
 Deacetylpaniculine, P-71
 6-*O*-Deacetylperegrine, M-347
 Deacetylpicraline 3,4,5-trimethoxybenzoate, B-420
 Deacetylpicraline, B-420
 14-Deacetylpubescenine 14-methyl ether, M-309
 14-*O*-Deacetylpubescenine, M-309
 Deacetylpyrifolidine, P-924
N-Deacetylranaconitine, A-86
N-Deacetylretuline, R-63
 Deacetylrigidusol, C-525
N-Deacetylscaconitine, M-352
 Deacetylsiwanine A, M-362
 Deacetylsolaphyllidine, S-349
N-Deacetylspermostrychnine, S-400
 Deacetylspiramine F, A-211
 Deacetylspiramine S, A-1534
 Deacetylstrychnobrasiline, S-594
 Deacetylsubcumine, M-323
 Deacetylsungpaconitine, M-303
 Deacetyltatsiensine, M-371
 2-Deacetyl taxine A, A-6
 10-Deacetyl taxol C, T-52
 10-Deacetyl taxol, T-52
 10-Deacetyl taxuyunnanine A, T-52
O-Deacetyl-1,2,4,5-tetrahydro-2-hydroxy-1-methylakuammilinium(1+), D-94
 Deacetylveralosine, S-199
 Deacetylvinblastine, V-110
 Deacetylvindoline, V-130
 10-Deacetyl-7-xylosylcephalomannine, T-52
 10-Deacetyl-7-xylosyltaxol B, T-52
 10-Deacetyl-7-xylosyltaxol C, T-52
 10-Deacetyl-7-xylosyltaxol, T-52
 8-Deacetyllyuonitine, M-336
 Deacetyluzurimine A, D-33
 Deacylbengazole C, D-95
 Deacylconvolutamide, D-96
 Deacylcynanchogenin, T-268
 Deacylcystodytin, D-97
 Deacyldrevogenin, P-602
 Deacylisariotin, D-98
O-Deacylpachysandrine B, D-266
 1-Deacylwillfordine, P-253
 Deaminocavananine, H-235
 3-Deamino-22-*O*-ethyl-3-hydroxysolanocapsine, D-99
 3-Deamino-3-hydroxysolanocapsine, D-99
 Deaner, D-730
 Deanol aceglumate, D-730
 Deanol acetamidobenzoate, D-730
 Deanol, D-730
 7-Deazaadenosine, T-658
 Debenzoylalopecurine, L-335
 Debenzoylaraloniine A, D-100
N-Debenzoyl-*N*-butanoyl-10-deacetylpaclitaxel, T-52
N-Debenzoyl-*N*-cinnamoylpaclitaxel, T-52
 Debenzoyllhmandrine, H-277
N-Debenzoyllucentamycin A, D-101
N-Debenzoyl-*N*-(α-methylbutyryl)paclitaxel, T-52
 2-Debenzoyl-2-nicotinoylwillforine, P-252
N-Debenzoyl-*N*-propanoyl-10-deacetylpaclitaxel, T-52
 2-Debenzoyl-2-tigloylpaclitaxel, T-52
 Debenzoylzucchini factor B, M-730
 Debric acid, D-102
 14-Debromoaplysamine 4, P-823
 14-Debromoaraplysin I, A-1378
 Debromoaxinohydantoin, A-1576
 (Z)-Debromoaxinohydantoin, A-1576
 Debromocathrynamide A, C-497
 (4*E*,6*E*)-Debromocathrynamide A, C-497
trans-6''-Debromo-3,4-dihydrohamacanthin A, H-40
trans-6''-Debromo-3,4-dihydrohamacanthin A, H-40
cis-6''-Debromo-3,4-dihydrohamacanthin B, H-41
cis-6''-Debromo-3,4-dihydrohamacanthin B, H-41
 2-Debromodispacamide A, D-863
 Debromodispacamide B, D-863
 Debromodispacamide D, D-863
 Debromodistomin K, E-287
 Debromoflustramine B, F-108
 Debromogrenadiene, H-131
 (R)-6''-Debromohamacanthin A, H-40
 (R)-6''-Debromohamacanthin A, H-40
 6''-Debromohamacanthin B, H-41
 (R)-6''-Debromohamacanthin B, H-41
 (R)-6''-Debromohamacanthin B, H-41
 Debromohymenialdisine, H-762
 (10*E*)-Debromohymenialdisine, H-762
 2-Debromohymenine, O-66
 6-Debromologamide A, L-242
 5-Debromomidpacamide, M-604
 2-Debromoodiline, O-66
 Debromooxysceptrine, O-225
 14-Debromopreaplysin I, P-821
 Debromosceptrine, S-124
 Debromoshermillamine A, S-286
 2-Debromostevensine, O-66
 2-Debromotauridispacamide A, T-39
 2,4-Decadienamide, D-104
 2,8-Decadiene-4,6-dienoic acid isobutylamide, D-103
 2,8-Decadiene-4,6-dienoic acid isopentylamide, D-103
 2,8-Decadiene-4,6-dienoic acid 2-phenylethylamide, D-103
 2,8-Decadiene-4,6-dienoic acid, D-103
 2,4-Decadienoic acid dehydropiperidide, D-125

- 2,4-Decadienoic acid *p*-hydroxyphenethylamide, H-668
- 2,4-Decadienoic acid isobutylamide, D-106
- 2,4-Decadienoic acid isopentylamide, D-104
- 2,4-Decadienoic acid 2-methylbutylamide, D-104
- 2,4-Decadienoic acid 2-phenylethylamide, D-104
- 2,4-Decadienoic acid piperidide, D-104
- 2,4-Decadienoic acid pyrrolidide, D-104
- 6,8-Decadienoic acid, D-105
- 2,4-Decadienoic acid, D-104
- N^o-2,4-Decadienylhistamine, H-307
- 1-(2,4-Decadienyl)pyrrolidine, D-104
- 5-(1,3-Decadienyl)indolizidine, D-107
- 5-(1,3-Decadienyl)octahydroindolizine, D-107
- 6-(1,3-Decadienyl)octahydro-4-methyl-2*H*-quinolin-3-ol, C-513
- 2,8-Decadien-6-ynoic acid, D-108
- 2,8-Decadien-6-ynoic 2-phenylethylamide, D-108
- Decahydro-2,5-bis(4-pentenyl)quinoline, D-109
- Decahydro-2,5-bis(2-propenyl)quinoline, A-357
- 1-(Decahydro-4,6-dihydroxy-5-(5-hydroxyoctyl)-2-methylquinoline, L-104
- Decahydro-3,3-dimethyl-2*H*-oxireno[*d*]pyrrolo[2,1-*a*]isoquinoline, B-484
- Decahydro-2,5-dipropylquinoline, A-357
- Decahydro-3-hydroxy-5-(5-hydroxyoctyl)-2-methylquinoline, L-104
- Decahydro-10-[(3-hydroxy-2,4,6-octadecatrienoyl)amino]-1,11*a*,11*b*-trihydroxy-2*H*-pyrazino[1,2-*a*]pyrrolo[2,1-*c*]pyrazine-3,6,9-trione, D-110
- 7-[5-(Decahydro-4*a*-hydroxy-1,2,5,5-tetramethyl-1-naphthalenyl)-3-methyl-2-pentenyl]-3,7-dihydro-2,3-dimethyl-6*H*-purin-6-one, D-111
- Decahydro-5-(1*H*-indol-3-ylmethyl)-1,4*a*-dimethyl-6-methylene-1-(4-methyl-3-pentenyl)-2-naphthalenol, E-74
- 3,4,7,7*a*,8,9,10,11,13,14-Decahydro-7,14-methano-2*H*,6*H*-dipyrido[1,2-*a*:1',2'-*e*][1,5]diazocin-6-one, A-1336
- 1,7,7*a*,8,9,10,11,13,14,14*a*-Decahydro-7,14-methano-2*H*,6*H*-dipyrido[1,2-*a*:1',2'-*e*][1,5]diazocin-2-one, M-731
- Decahydro-1,5-methano-6*H*-pyrido[1,2-*a*][1,5]diazocin-6-one, V-152
- Decahydro-8'-methoxy-4-methyl-3'-(tetrahydro-4-methyl-5-oxo-2-furanyl)spiro[furan-2(5*H*),9'-[9*H*]pyrrolo[1,2-*a*]azepin]-5-one, S-523
- Decahydro-1-methyl-5,7-diphenyl-6-(3,4,5,6-tetrahydro-2-pyridinyl)quinoline, L-220
- Decahydro-5-methyl-2-(2,4-pentadienyl)quinoline, A-403
- Decahydro-9*a*-methyl-2-propyl-2*H*-[1,3]oxazino[2,3,4-*de*]quinolinizine, P-570
- Decahydro-2-methyl-5-propylquinoline, D-112
- Decahydro-5-methyl-2-propylquinoline, P-794
- Decahydro-3-methyl-1*H*-pyrano[2,3-*g*]indolizin-1-one, E-60
- 1,2,3,3*a*,4,5,6,6*a*,7,9*a*-Decahydro-8-methylpyrido[2,1,6-*de*]quinolinizine, H-295
- 1,2,4,5,6,6*a*,7,8,9,9*a*-Decahydro-2-methylpyrido[2,1,6-*de*]quinolinizine, P-649
- Decahydro-3-methylpyrrolo[2,1-*f*][1,6]naphthyridin-1(2*H*)-one, E-57
- Decahydro-1-methyl-8-(tetrahydro-4-methyl-5-oxo-2-furanyl)-2*H*-furo[3,2-*c*]pyrrolo[1,2-*a*]azepin-2-one, S-519
- Decahydro-2-(3,4-pentadienyl)-5-(2-penten-4-ynyl)quinoline, A-366
- Decahydro-5-(2-penten-4-ynyl)-2-(2-propenyl)quinoline, A-362
- Decahydro-2-(2-penten-4-ynyl)quinoline, A-289
- Decahydro-5-pentyl-5*H*-pyrido[1,2-*c*]pyrrolo[1,2-*a*]pyrimidine, T-293
- 3,4,5,6,8,9,10,11,12,13-Decahydro-9-phenyl-2,13-methano-2*H*-2,6,10-benzotriazacyclopentadecine-1,7-dione, I-222
- Decahydro-8-(1-piperidinylmethyl)isoquinoline, D-113
- Decahydro-4-(2-propenyl)-1,5-methano-8*H*-pyrido[1,2-*a*][1,5]diazocin-8-one, A-1013
- Decahydro-2-propylquinoline, A-401
- 1,5,6,7,7*a*,8,9,10,11,12*a*-Decahydro-2-(2-pyrrolidinylmethyl)-2*H*-pyrano[2,3,4-*ij*]pyrido[1,2-*b*][2,7]naphthyridine, H-352
- 1,2,3,3*a*,5,6,11,11*b*,12,13-Decahydro-2-pyrrolo[2'',1'':2',3']pyrimido[1',6':1,2]pyrido[3,4-*bj*]indole, E-54
- Decahydro-2,5,8-trimethyl-2*H*-1-oxa-4,7,9*b*-triazaphenylene, D-114
- Decaisnine, P-221
- (10*α*)-Decaline, V-86
- Decaline, V-86
- Decamine, V-95
- Decanoic acid isobutylamide, D-115
- Decanoic acid, D-115
- 3-Decanoyl-5-methylpyridine, M-538
- 5-Decanoyl-2-nonylpyridine, D-116
- N*-Decanoyltyrosine, T-700
- Decarbamoyl-12-deoxysaxitoxin, S-109
- Decarbamoylgonyauroxin I, H-731
- Decarbamoylgonyauroxin II, H-731
- Decarbamoylgonyauroxin III, H-731
- Decarbamoylgonyauroxin IV, H-731
- Decarbamoyl-11*α*-hydroxyneosaxitoxin, H-731
- Decarbamoyl-11*β*-hydroxyneosaxitoxin, H-731
- Decarbamoyl-11*β*-hydroxysaxitoxin, H-731
- Decarbamoylneosaxitoxin, S-109
- 10-Decarbamoyloxy-9-dehydromitomycin B, M-656
- 10-Decarbamoyloxy-9-dehydroporfirofomycin, M-655
- Decarbamoyloxygonyauroxin II, H-731
- Decarbamoyloxygonyauroxin III, H-731
- Decarbamoyloxyneosaxitoxin, S-109
- Decarbamoylsaxitoxin, S-109
- Decarbomethoxyapocuanzine, D-117
- Decarbomethoxy-19,20-dihydroakummicine, T-668
- 16'-Decarbomethoxy-19,20-dihydroconoduramine, D-118
- Decarbomethoxydihydrogambirtannine, D-434
- 16'-Decarbomethoxy-19,20-dihydrovoacamine, T-8
- 16-Decarbomethoxy-19,20-dihydrovobasine, D-119
- 16'-Decarbomethoxy-20-epi-19,20-dihydroconoduramine, D-118
- 16'-Decarbomethoxy-20-epidihydrovoacamine, T-8
- 16-Decarbomethoxy-20-epi-19,20-dihydrovobasine, D-119
- Decarbomethoxyerymelanthine, E-198
- 16'-Decarbomethoxy-3'-hydroxyconodurine, C-616
- N*-Decarbomethoxykopsaporine, K-66
- Decarbomethoxykopsifine, K-69
- Decarbomethoxykopsine, K-76
- Decarbomethoxynauclechione, N-63
- Decarbomethoxytetrahydrosecamine, S-174
- Decarbomethoxytetrahydrosecodine, D-120
- 16-Decarbomethoxyvinerivine (incorr.), T-668
- 16-Decarbomethoxyvoacamine pseudoindoxyl, D-121
- Decarbomethoxyvoacangarine, I-4
- 2-Decarboxybetanidin, D-122
- 2-Decarboxybetanin, D-122
- Decarboxydehydrodanguinone A, S-58
- Decarboxydehydrodanguinone A, S-58
- 2-Decarboxyphylllocactin, D-122
- Decarine acetate, C-356
- Decarine, C-356
- 2,4,6,8-Decatetraenoic acid dehydropiperidide, D-125
- 6-(1,3,5,7-Decatetraenyl)-1,2-dimethyl-3-piperidinol, C-788
- 2,4,6-Decatrienoic acid dehydropiperidide, D-125
- 2,4,8-Decatrienoic acid isobutylamide, D-123
- 2,4,8-Decatrienoic acid, D-123
- 2,6,8-Decatrienoic acid, D-124
- 6-(1,3,5-Decatrienyl)-3-hydroxy-2-hydroxymethyl-1-methylpiperidine, M-600
- 6-(1,3,5-Decatrienyl)-3-hydroxy-1-methyl-2-piperidinemethanol, M-600
- 5-(1,3,5-Decatrienyl)indolizidine, D-107
- 6-(1,3,5-Decatrienyl)-3-methoxy-1,2-dimethylpiperidine, C-788
- 5-(1,3,5-Decatrienyl)octahydroindolizine, D-107
- Decatromicin A, D-126
- Decatromicin B, D-126
- Decaturin A, D-127
- Decaturin B, D-127
- Decaturin C, D-127
- Decaturin D, D-128
- 2-Decene-6,8-diyonoic acid isobutylamide, D-130
- 2-Decene-6,8-diyonoic acid, D-130
- 2-Decene-4,6-diyonoic acid, D-129
- 2-Decene-4,6,8-triynoic acid isopentylamide, D-131
- 2-Decene-4,6,8-triynoic acid 2-phenylethylamide, D-131
- 2-Decene-4,6,8-triynoic acid, D-131
- 2-Decenoic acid isobutylamide, D-132
- 2-Decenoic acid, D-132
- 2-(3-Decenyl)-1,6-diazacyclododec-8-ene-7,12-dione, B-5
- 2-(1-Decenyl)-4-hydroxyquinoline, D-139
- 5-(2-Decenyl)indolizidine, D-133
- 5-(2-Decenyl)octahydroindolizine, D-133
- 2-(1-Decenyl)-4(1*H*)-quinolinone, D-139
- N*-(3-Decenyl)sulfamic acid, D-134
- Dechloroacutumine, A-134
- Dechlorobarbamide, B-22
- 6-Dechlorobazelline C, D-13
- Dechlorodauricumine, A-134
- Dechloro-4,5-deepoxy-*N*-demethylmaytansinol, M-136
- Dechloro-*N*-deepoxymaytansinol, M-136
- 10-Dechlorodysideathiazole, D-970
- Dechlorofontomamine, F-126
- 10-Dechloro-*N*-methyllysidiathiazole, D-970
- 6-Dechloropenitrem A, P-194
- Decinine, V-95
- Decipinol ester A, H-255
- Decipinone A, H-255
- Declovanillobicin, V-18
- Decodine, V-88
- Decoic acid (obsol.), D-115
- Decorticasine, L-234
- Decumbenine B, D-136
- Decumbenine C, H-767
- Decumbenine, D-135
- Decumbensine, E-43
- Decursivine, D-137
- Decussine, D-138
- 21-Decyano-25-dihydrosafamycin A, S-4
- 2-Decyl-4-hydroxyquinoline, D-139
- Decylic acid, D-115
- 2-Decyl-4-quinolinol, D-139
- 2-Decyl-4(1*H*)-quinolinone *N*-oxide, D-139
- 2-Decyl-4(1*H*)-quinolinone, D-139
- 2,5-De(epidithio)-2,5-(epitritio)chaetocin, C-321
- Deepoxybohemamine, B-242
- 17,18-Deepoxyacarurine II, D-910
- Deet, ESA, M-398
- N*-Deethylaconitine, M-331
- N*-Deethylanthranoylcoctonine, M-309
- N*-Deethyldehydrolicidusculine, D-162
- N*-Deethyldephatine, M-309
- N*-Deethyldeleine, M-309
- N*-Deethyl-*N*,19-didehydrosachaconitine, M-378
- N*-Deethyl-*N*-formylbrownine, M-309
- N*-Deethyl-*N*-formylsachaconitine, L-317
- Deethylbophyllidine, D-140
- 16-Deethylindanomycin, I-64
- N*-Deethylsachaconitine-*N*-aldehyde, L-317
- N*-Deethyllycoctonine, M-309
- N*-Deethyl-*N*-methyl-12-epinapelline, N-30
- 8-*O*-Deethyl-8-*O*-methylkongboenine, M-312
- N*-Deethylmethylsachaconitine, G-91
- N*-Deethyl-14-*O*-methylperegrine, M-347
- N*-Deethylsongorine, D-162
- N*-Deethylsongorine, N-30
- 19-*O*-Deethylspiramine N, B-371
- O*-Deethylalroconvolutin C, D-141
- Deferiprone, D-632
- Deferoxamine hydrochloride, D-142
- Deferoxamine, D-142
- Deformoundatine, U-32

- Deformylcorymine, D-236
 Deformylcoryzylamine, D-143
N-Deformyldichotamine, D-144
 Deformylflustrabromine B, B-356
 Deformylflustrabromine, B-316
 Deformylstemmedenine, S-510
 Deformyltalbotinic acid methyl ester, D-145
 Deformylvallesine, A-1499
N-Deformylvincristine, V-110
 Deglucopterocereine *N*-oxide, T-178
 Deglucopterocereine, T-178
Sambucus nigra Degraded cyanogenic glycosides, D-146
 Dehassiline, D-147
 Dehatridine, C-171
 Dehatrine, D-148
 Dehatriphine, D-149
 DEHSPM, B-150
 19,20-Dehydro-*O*-acetylöhimbine, D-378
 Dehydroacosanine, M-309
 Dehydroactidione, D-765
 19,20-Dehydrodirubrine acetate, A-141
 Dehydroaglaiaastatin, A-197
 Dehydroajmalicine, C-207
 Dehydroalbaine, A-243
 Δ^5 -Dehydroalbaine, A-243
 Dehydroanacycline, A-968
 Dehydroangustifoline, A-1013
 4,5-Dehydroanhydrolycorine, H-290
 4,5-Dehydroanhydroseuodolcorine, A-1508
 Dehydroanonaine, D-166
 Dehydroantofine, A-1025
 1,2-Dehydroapateline, K-115
 Dehydroapocavidine, D-150
 Dehydroaporheine, D-166
 9,10-Dehydroaristoteline, A-1442
 1,2-Dehydroaspidospermidine *N*-oxide, E-3
 1,2-Dehydro- ψ -aspidospermidine, D-377
 1,2-Dehydroaspidospermidine, E-3
 Dehydroazasiosterol, D-151
 Dehydrobatzelladine C, D-152
 1,2-Dehydrobeninine, B-63
 Dehydrobhesine, T-139
 Dehydrobicoloridine, G-152
 α -Dehydrobiotin, L-166
 Dehydroboldine, D-159
 Dehydroborrecapine, B-251
 Dehydrobrevicolline, B-292
 14-Dehydrobrowniine, M-309
 Dehydrobufotenine, D-153
 18,19-Dehydrocampothecin, C-70
 Dehydrocancrine A, C-76
 Dehydrocancrine B, C-76
 3,4-Dehydro-5-carboxystrictosidine, C-153
 Dehydrocardiopetaline, G-152
 Dehydrocarpaine I, C-168
 Dehydrocarpaine II, C-168
 Dehydrocavidine, D-150
 Dehydrochelanthifoline, G-181
 17,4'-Dehydro-3 α -cinchophylline, C-448
 2-Dehydrococcinelline, H-295
 Dehydrocoelenterazine, C-560
 7,8-Dehydrocoelobillardierine, C-562
 Dehydrocommersonine, D-205
 Dehydrocoreximine, P-719
 5,6-Dehydrocoronaridine, C-657
 Dehydrocorybulbine, D-154
 Dehydrocorydaline, D-154
 Dehydrocorydalmine, P-34
 Dehydrocorydine, C-682
 1,2-Dehydrocorypalline, D-421
 Dehydrocorypalline, P-34
 Dehydrocorytenchine, P-719
 Dehydrocrambescine A, D-155
 Dehydrocrambescine A, D-155
 Dehydrocrebanine, C-733
 Dehydrocrenatidine, D-686
 Dehydrocrenatine, M-290
 3,4-Dehydrocycleanine, C-835
 Dehydrocycloheximide, D-765
 Dehydrocyclopeptide, C-890
 Dehydrocyclostelletamine D, C-910
 Dehydrocyclostelletamine E, C-910
 11,12-Dehydrocytisine, C-938
 12,13-Dehydrocytisine, C-938
 Dehydrodaphnigraciline, D-50
 Dehydrodaphnigraciline, D-51
 Dehydrodarlingianine, D-74
 Dehydrodarline, D-76
 2-Dehydrodeacetylheterophyllidine, H-161
 Dehydrodecodine, V-88
 6-Dehydrodelcorine, M-309
 14-Dehydrodelcosine, M-309
 2,3-Dehydrodelcosine, M-363
 6-Dehydrodelpeline, M-321
 1-Dehydrodelpisine, M-323
 Dehydrodelsoline, G-152
 Dehydrodeltamine, M-307
 Dehydrodeltatsine, G-152
 2,3-Dehydro-6-*O*-demethyldecosine, M-363
 Dehydrode-*N*-methyluleine, U-8
 5,6-Dehydrodemissine, S-345
 Dehydrodeoxynupharidine, D-227
 1,2-Dehydrodesacetylretuline, R-63
 2,16-Dehydrodiaboline, C-115
 Dehydrodicentrine, D-156
 2,3-Dehydrodictyocarpine, M-362
 15-Dehydrodictysine, D-243
 3,4-Dehydro-8,10-diethyllobelidiol, D-391
 5,8'-Dehydrodiferulic acid, D-395
 24,25-Dehydro-10,11-dihydroaflavinine, A-172
 24,25-Dehydro-10,11-dihydro-20-epi-20-hydroxyaflavinine, A-172
 24,25-Dehydro-10,11-dihydro-20-hydroxyaflavinine, A-172
 24,25-Dehydro-10,11-dihydro-13-oxoaflavinine, A-172
 4,5-Dehydro-5,6-dihydro-3-oxo-20-epiverazine, V-73
 14,15-Dehydro-10,11-dimethoxy-16-epivincamine, V-115
 14,15-Dehydro-10,11-dimethoxyvincamine, V-115
 Dehydrodioxolide A, D-807
 Dehydrodioxolide B, D-807
 Dehydrodiscretamine, P-34
 Dehydrodiscretine, P-719
 Dehydrodisecocyclopiazonic acid, B-218
 Dehydrodopamine, D-157
 19,20-Dehydroedulirin A, T-349
 19,20-Dehydroedulirone A, E-41
 Dehydroeldelidine, M-307
 Dehydroelymoclavine, A-205
 Dehydro-16-epiaffinisine, G-23
 2,3-Dehydroepiaphylline, A-1337
 6,7-Dehydro-20-epiminovincinine, H-743
 14,15-Dehydro-19-epiminovincinine, H-743
 Dehydroepivincadine, V-112
 14,15-Dehydro-16-epivincamine, V-115
 19,20-Dehydroervatamine, E-190
 10,11-Dehydroerysodine, E-203
 10,11-Dehydroerysovine, E-203
 Dehydro- α -erythroidine, E-215
 Dehydroevodiamine, D-158
 13,13*b*-Dehydroevodiamine, D-158
 6*a*,7-Dehydrofloripavidine, D-164
 1,2-Dehydrofluorodaturatin (incorr.), H-444
 4,5-Dehydrofluorodaturatin, H-444
 1,2-Dehydrofluorodaturine (incorr.), H-444
 Dehydroformouregine, T-521
 6*a*,7-Dehydro-*N*-formylnormantene, N-29
 1,2-Dehydrofuchsisenecionine, T-188
 4-Dehydrofuntumafribe B, A-891
 Dehydrofusaric acid, B-425
 4,21-Dehydrogeissoschizine, G-32
 1,2-Dehydrogeissoschizoline, G-34
 Dehydrogentianine lactam, G-60
 Dehydroglaucone, D-159
 Dehydroguatessine, G-214
 5',6'-Dehydroguaiachrysin, S-593
 5',6'-Dehydroguiaflavine, G-219
 Dehydroheliamine, D-421
 Dehydrohernandaline, H-147
 Dehydrohetisine, H-202
 14,15-Dehydroholaleniene, H-314
 Dehydrohuangshanine, H-366
 19,20-Dehydro-6-hydroxyervatamine, E-190
 14,15-Dehydro-10-hydroxy-11-methoxy-16-epivincamine, V-115
 14,15-Dehydro-10-hydroxy-11-methoxyvincamine, V-115
 Δ^5 -Dehydro-13-hydroxymultiflorine *N*-oxide, M-731
 Δ^5 -Dehydro-13-hydroxymultiflorine, M-731
 18-Dehydro-10-hydroxynigritanine, O-21
 3,4-Dehydro-5-hydroxy-2-piperidone, D-458
 8*a*,8*b*-Dehydro-8-hydroxyptilocalin, P-773
 7,8-Dehydro-1-hydroxyrutaecarpine, R-168
 6,7-Dehydro-19-hydroxyschizozogine, S-137
 14-Dehydro-10-hydroxytermine, T-90
 1',2-Dehydrohygrine, H-759
 14-Dehydroliensine, M-309
 Dehydroindoxyl, I-106
 2,3-Dehydro- α -isoaphylline, A-1337
 Dehydroisoapocavidine, D-150
 Dehydroisoboldine, D-159
 Dehydroisochalciporone, C-340
 Dehydroisocinchophyllamine, C-448
 Dehydroisocorydione, S-367
 Dehydroisocorypalline, P-34
 Dehydroisolaurine, D-175
 Dehydroisolongistrobine, D-160
 5,6-Dehydro- α -isolupanine, D-161
 5,17-Dehydroisomatrine, D-80
 18,19-Dehydro-3-isoreserpiline, R-51
 Dehydroisosenaeatine, S-232
 (+)-Dehydroisosenaeatine, S-232
 5,6-Dehydro- α -isosparteine, D-168
 16,17-Dehydroisostrychnobiline, S-591
 Dehydroisothebaine, T-525
 Dehydrojoubertamine, J-49
 11-Dehydrokobusine, H-167
 1',2'-Dehydrokohatine 2*B*-*N*-oxide, K-56
 Dehydrokohatine, K-56
 14,15-Dehydrokopsinine, V-49
 Dehydrokuanoniamine B, D-93
 Dehydrolirinidine, D-164
 Dehydrolirioferine, D-159
 5,6-Dehydrololinine, L-234
 Dehydroluciculine, D-162
 Dehydrolucidusculine, D-162
 5,6-Dehydrolopamine, D-161
 2,3-Dehydrolycaonic acid piperidide, O-194
 2,3-Dehydrolycaonic acid pyrrolidide, O-194
 2,3-Dehydrolycaonic acid, O-194
 Dehydrolyccaroline, L-320
 Dehydrolycopecurine, L-335
 5,6-Dehydromalouphyllinol, D-279
 Dehydromatricaric acid, D-131
 Dehydromatricaric acid isobutylamide, D-131
 Dehydromatricaric acid, D-131
 5,17-Dehydromatrine *N*-oxide, L-93
 5,17-Dehydromatrine, L-93
 7,11-Dehydromatrine, L-94
 12,13-Dehydromatrine, L-81
 Dehydro-4-methoxycyclobassinin, M-276
 19,20-Dehydro-10-methoxydihydrocorynantheol, G-33
 7,8-Dehydro-1-methoxyrutaecarpine, R-168
 19,20-Dehydro-10-methoxytalcarpine, T-20
 14,15-Dehydro-*N*-methylaspidospermidine, A-1498
 Dehydro-8-methyl-10-ethyllobelidiol, H-434
 1,2-Dehydromicranthine, K-115
 3,4,5,6-Dehydromitragynine, M-657
 3-Dehydromitragynine, M-657
 Dehydromonocrotaline, M-688
 Dehydromorphine, M-703
 5-Dehydromultiflorine, M-731
 Dehydromultiflorine, M-731
 Dehydronantene, N-29
 Dehydronapelline, D-162
 3,4-Dehydroneoipecoside, I-167
 Dehydronolitsine, N-125
 18,19-Dehydrogrigritanine, O-21
 Dehydroniranin A, N-235
 Dehydroniranin B, N-235
 Dehydronoryclodercitine, C-857
 Dehydronorythrosuamide, N-299
 Dehydronorerythrosuamine, N-299

- 6,6 α -Dehydronorglaucine, D-159
6 α ,7-Dehydronorglaucine, D-159
6,6 α -Dehydronorlaureline, L-72
1,2-Dehydro-2-norlimacusine, C-551
Dehydronormacrostromine, D-163
Dehydronornuciferine, D-164
7-Dehydronornuciferine-7'-dehydro-*O*-methylisopiline, U-41
3,4-Dehydronoroxyhydrastinine, I-305
7(11)-Dehydro-13-norpilocarpine, N-295
Dehydronortehaunine, I-307
1,2-Dehydro-2'-nortelobine, K-115
Dehydronorweberine, T-276
Dehydronuciferine, D-164
18-Dehydroochrolifuanine A, O-21
18-Dehydroochrolifuanine E, O-21
18-Dehydroochrolifuanine F, O-21
Dehydroocopodine, L-158
Dehydroocoteine, O-27
Dehydroodorine, O-67
Dehydroorientalidine, A-584
19,20-Dehydro-5-oxoervatamine, E-190
10,11-Dehydro-8-oxoerythraline, C-799
14-Dehydro-18-oxoholadienine, H-314
2,3-Dehydro-10-oxo- α -isopartaine, A-1337
Dehydropachycereine, T-233
Dehydropfanostenine, D-156
Dehydropphenylahistin, P-338
Dehydrophomin, C-943
3,4-Dehydro-2-piperidone, D-495
Dehydropipernonaline, P-464
Dehydropredicentrine, D-159
Watasenia Dehydropreluciferin, C-560
12,13-Dehydropropyl-2-(1,1-dimethylallyltryptophyl)diketopiperazine, C-861
12,13-Dehydropropyltryptophyldiketopiperazine, C-899
Dehydroprotostemonine, P-670
Dehydropseudocheilanthifoline, T-325
8 α ,8 β -Dehydroptilocaulin, P-773
Dehydropuntarenine, D-165
2,3-Dehydro-*O*-(2-pyrrolylcarbonyl)virgiline, V-153
19,20-Dehydroreserpiline, R-51
19-Dehydroreserpine, D-233
Dehydroretrofractamide C (incorr.), R-60
Dehydoroemerine, D-166
7,8-Dehydrotutaecarpine, R-168
Dehydroyanodine, R-175
Dehydrosalsolidine, D-627
Dehydro- γ -sanshool, T-127
16-Dehydrosarcovagine D, D-260
Dehydrosaulatine, D-167
Dehydroschoberine, S-140
Dehydroscoulerine, P-34
Dehydrosenaetnine, S-232
Dehydrosinomenine, S-313
 Δ^4 -Dehydroskytanthine, S-326
 Δ^{4a} -Dehydroskytanthine, S-326
Dehydrosolenopsin B, M-580
Dehydrosolenopsin C, M-504
1-Dehydrosongorine, N-30
 Δ^7 -Dehydrosophoramine, S-376
13,14-Dehydrosophoridine *N*-oxide, S-371
13,14-Dehydrosophoridine, S-371
11,12-Dehydrosparteine, D-168
2,3-Dehydro- β -stachydrine, D-169
Dehydrostephalagine, S-536
Dehydrostephanine, T-522
Dehydrostepholidine, P-34
Dehydrostesakine, C-733
3,4-Dehydrostrictosidine, S-583
3,4-Dehydrostrictosidinic acid, S-583
14-Dehydrotalatzamine, M-352
1,2-Dehydrotelobine, K-115
4,6-Dehydro-1,2,4,5-tetrahydro-2,5-dioxopilocarpine, E-272
14-Dehydrotetrastachyne, T-294
14,15-Dehydrotetrastachynine, T-295
Dehydrothalebanin A, T-303
Dehydrothalebanin B, T-303
Dehydrothalicarpine, D-170
Dehydrothalicmine, O-27
Dehydrothalicimidine, P-208
Dehydrothalictricavine, D-171
Dehydrothalifaberine, T-322
Dehydrothaliporphine, D-159
Dehydrothallmelatine, D-170
Dehydrotomatine, S-456
Dehydrotrewiasine, T-450
Dehydrotryptamine, D-172
1',2'-Dehydrotubulosine, T-675
Dehydrotylophorine, A-1025
Dehydroverrucosine, F-160
6,7-Dehydrovincadine, V-112
14,15-Dehydrovincadine, V-112
14,15-Dehydrovincamine, V-115
14,15-Dehydrovincine, V-115
Dehydrovirgiboidine, V-152
Dehydrovirgilidone, V-152
Dehydrovoachalotine, D-173
1-Dehydroxybaccatin III, E-138
6-Dehydroxy-*O*¹⁰-buxafuranamine, B-449
2-Dehydroxy-*O*²-buxafuranamine, D-174
Dehydroxycapuvosine, C-106
4 α -Dehydroxycrinamabine, A-670
Dehydroxyervataminol, E-190
10-Dehydroxyflavaconitine, M-343
Dehydroxyhaplocidine, A-1490
17-Dehydroxykopsingine, K-66
Dehydroxylophine, D-175
16-De(hydroxymethyl)-4-demethyl-16-(methoxycarbonyl)fluorocurine, F-99
6-De(hydroxymethyl)-6-epitrodoxin, T-298
Dehydroxymethylflazine, F-85
Dehydroxymethylvoachalotinol, T-425
Dehydroxypaxilline, P-140
19,20-Dehydro- α -yohimbine, D-378
19,20-Dehydro- β -yohimbine, D-378
19-Dehydroyohimbine, D-378
3,4-Dehydro- β -yohimbine, Y-13
Delafrine, C-297
Delafirinone, C-297
Delagoenine, D-176
Delagoensine, D-177
Delajacine, M-309
Delajacrine, M-309
Delajadine, M-309
Delamide, D-178
Delaminomycin A, D-179
Delaminomycin B, D-179
Delaminomycin C, D-179
Delartine, G-91
Delatine, H-202
Delatisine, D-180
Delavaconitine C, A-99
Delavaconitine D, A-99
Delavaconitine E, A-98
Delavaconitine F, A-98
Delavaconitine G, A-102
Delavaconitine, A-98
Delavaine A, D-182
Delavaine B, D-183
Delavaine, D-181
Delavayine A, D-184
Delavayine B, D-185
Delavayine C, D-186
Delavayine, D-181
Delavine, C-290
Delavinone, C-290
Delbidine, T-552
Delbine, A-85
Delbine, F-221
Delbiterine, M-309
Delbonine, M-309
Delbotine, M-309
Delboxine, D-187
Delbruline, M-309
Delbrunine, M-309
Delbruninol, M-309
Delbrusine, M-309
Delcaine, C-537
Delcarduchol, S-170
Delcaroline, M-339
Delcoline, M-309
Delcoridine, M-309
Delcorine, M-309
Delcorinine, M-309
Delcosine, M-309
Delectine, M-309
Delectinine, M-309
Delelatine, M-321
Delfissinol, H-207
Delflexine, D-188
Delfrenine, D-189
Delgramine, H-175
Delimmun, I-148
Delmenzine, M-321
Delnudine, D-190
Delnuttaline, T-551
Delnuttidine, H-198
Delnuttine, H-208
Delorine, D-191
Delosine, M-309
Delpetine, M-307
Delphamine, M-309
Delphatamine, M-665
Delphatine, M-309
Delphatisine A, A-1534
Delphatisine B, D-192
Delphelaine, M-321
Delpheline, M-321
Delphicripuline, A-95
Delphidine, M-323
Delphigraciline, H-169
Delphimifoline, M-309
Delphinine, M-312
Delphiperegrine, M-321
Delphirine, M-323
Delphisine, M-323
Delphoccine, D-193
Delphonine, M-312
Delphoorine, D-194
Delpoline, M-355
Delporphine, P-208
Delpyrine, D-195
Delretine, M-307
Delsemidine, G-91
Delsemine, D-196
Delsine, M-309
Delsoline, M-309
Delsonine, M-309
Delstaphidine, G-152
Delstaphigine, M-312
Delstaphinine, G-152
Delstaphisagnine, M-323
Delstaphisagrine, M-323
Delstaphisine, M-323
Delstaphisinine, M-323
Deltaline, M-307
Deltamine, M-307
Deltatsine, M-309
Delterine, M-307
Delvestidine, M-309
Delvestine, M-309
Demalonylcopiamycin, C-634
Demalonylmethylcopiamycin, C-634
Demanol aceglumate, D-730
Demanyl phosphate, D-197
Demecolceine, D-92
Demecolcine, D-92
Demecolcinone, D-198
Demerarine, R-46
5-Demethoxy IT 143B, P-421
Demethoxyabresoline, L-53
Demethoxyalstonamide, V-125
Demethoxyalstoniline, O-140
Demethoxyamiflorine, A-1039
Demethoxyanomaline, A-1067
Demethoxyaspidospermine, A-1498
16-De(methoxycarbonyl)-17-deoxy-19,20-dihydro-19-hydroxy-21-methyl-18-nortalbotine, S-600
Demethoxycarbonyldihydrogambirtannine, D-434
16-De(methoxycarbonyl)-19,20-dihydrovobasine, D-119
16-Demethoxycarbonyl-20-epiervatamine, M-293
Demethoxycarbonyl-3,4,15,16,17,18-hexahydrogambirtannine, D-434
Demethoxycarbonylkopsifine, K-69

- 1-Demethoxycarbonyl-12-methoxyfruticosine, F-163
 16-Demethoxycarbonylpagicerine, P-10
 10,11-Demethoxychippine, C-389
 Demethoxychoisyne, C-418
 17-Demethoxycorynoxine B, R-90
 30-Demethoxycurromycin A, C-815
 (±)-Demethoxycyclindrocarpidine, C-929
 Demethoxycylindrocarpidine, C-929
 16-Demethoxydelavaine A, D-182
 16-Demethoxydelavaine B, D-183
 10-Demethoxy-21-deoxykopsidasine, K-71
 Demethoxy-4,5-dioxodehydroasimilobine, C-273
 11-Demethoxy-3-epimyrtoidine, M-803
 3-Demethoxyerythradinone, D-199
 3-Demethoxyfaberonine, T-333
 Demethoxyfunitremorgin C, F-190
 (19*E*)-18-Demethoxygardfloramine, G-18
 (19*E*)-18-Demethoxygardneramine, G-20
 Demethoxyguadisine, G-188
 α-Demethoxyhaemanthamine, H-9
 β-Demethoxyhaemanthamine, H-9
 Demethoxyhoerhammerine, H-743
 8-Demethoxyhostasine, H-362
 1-Demethoxyhumanenine, H-367
 4'-Demethoxy-6'-hydroxydecaline, L-11
 18-Demethoxy-16-hydroxygardneramine oxindole, G-21
 Demethoxyisomigrastatin, I-245
 17-Demethoxysorhynchophylline, R-90
 10-Demethoxykopsidasine, K-68
 Demethoxykopsingine, K-66
 10-Demethoxy-12-methoxykopsidasine, K-68
 3-Demethoxy-2,3-methylenedioxyerythroculine, D-200
 8-Demethoxy-10-*O*-methylhostasine, H-362
 16-Demethoxymethyllycaconitine, G-91
 4-Demethoxymichigazone, M-590
 11-Demethoxymyrtoidine, M-803
 Demethoxyपालosine, A-1498
 3'-Demethoxyplaptine, P-473
 Demethoxypodocarpamide, Z-6
 Demethoxyprodigosin, P-636
 18-Demethoxypubescence, M-321
 Demethoxypurpleline, T-288
 6-Demethoxyppyrifoline, K-78
 11-Demethoxyreserpine, D-234
ar-Demethoxyrindline, H-323
 8-Demethoxyrunanine, R-161
 10'-Demethoxystreptonigrin, S-569
 21-Demethoxysulfomycin I, S-638
 12-Demethoxytabernulosine, P-417
 11-Demethoxytomaymycin, T-424
 Demethoxyvandrikinine, M-666
 Demethoxyvindoline, V-130
 Demethyl Antibiotic UK 1, A-1302
 Demethyl UK 1, A-1302
 9-*De-O*-methylaaptamine, A-2
N-Demethylacedine, A-27
 Demethylacedinsine, A-28
 14-*O*-Demethyl-14-acetylanhweidelphinine, A-1023
 9-*O*-Demethyl-9-*O*-acetylhomolycorine, L-347
De-N-methylacronycine, A-120
*O*⁶-Demethyladiantifoline, A-139
N-Demethyladouetine X, A-149
N-Demethyladouetine Y', A-150
N-Demethyladouetine Z, A-151
 Demethylalangside, A-239
*N*⁶-Demethylalfileramine, A-260
*N*⁶-Demethylalfileramine, A-260
 2-Demethylalcolchicine, A-537
N-Demethylalpinine, A-638
*N*⁶-Demethylalstogustine *N*-oxide, E-30
*N*⁶-Demethylalstogustine, E-30
N-Demethylalstonal, A-648
*N*¹-Demethylalstonerinal, T-20
*N*⁴-Demethylalstonerinal, T-20
*N*⁴-Demethylalstonerine, A-649
N-Demethylalstonisine, A-654
*N*⁶-Demethylalstophyllal oxindole, A-648
*N*⁶-Demethylalstophylline oxindole, A-654
*N*¹-Demethylalstophylline, A-649
 4'-Demethylamino-4',5'-dihydroxystaurosporine, S-498
 4'-Demethylamino-4'-hydroxy-3'-epistaurosporine, S-498
 4'-Demethylamino-4-hydroxystaurosporine, S-498
 4'-Demethylamino-4'-nitrostaurosporine, S-498
N-Demethylamphibine H, A-942
 6-*O*-Demethylancistrobenomine A, A-982
 4'-*O*-Demethylancistrocladine, A-982
 6-*O*-Demethylancistrocladisine, A-985
 5'-*O*-Demethylancistrocline, A-982
 6-*O*-Demethylancistroalaine A, A-980
N-Demethylanisamitocin P3, M-136
De-O-methylanisomycin, A-1044
N-Demethylansamitocin P1, M-136
N-Demethylansamitocin P2, M-136
N-Demethylansamitocin P4, M-136
 6-Demethylanthranoyllycotonine, M-309
 6-Demethylantofine, A-1330
 2'-*N*-Demethylaplysinopsin, A-1357
N-Demethylarmepavine, A-1444
O-Demethylaspidocarpine, P-924
 Demethylaspidospermine, A-1499
 Demethylasterriquinone B1, A-1521
N-Demethylbelladine, N-288
 6-*O*-Demethylberbamine, I-328
 10-*O*-Demethylbrucine, S-589
N-Demethylbuchenavianine, D-662
O-Demethylbuchenavianine, D-662
 8-*O*-Demethylbulflavine, A-1361
 7-Demethylcalafatine, C-23
N-Demethylcalyculin A, C-55
N-Demethylcapitavine, C-94
N-Demethylcapuvosine, C-106
N-Demethylcarboethoxymacronine, P-626
O-Demethylcarnegine, T-162
N-Demethylcelesticetin, C-239
O-Demethylcelesticetin, C-239
 7-*O*-Demethylcephaeline, E-72
 9-*O*-Demethylcephaeline, E-72
 Demethylcephaeline, E-72
 10-*O*-Demethylcephaeline, E-72
 Demethylcephalotaxine, C-271
 Demethylcephalotaxinone, I-201
 Demethylceridimine, H-275
De-O-methylcervinomycin A₂, C-287
 11-Demethylcezymocin, C-26
N-Demethylchelidonine, C-357
N-Demethyl-*meso*-chimonanthine, C-382
N-Demethylcitracridone I, C-472
 Demethylcoculaurine, H-275
 12-*O*-Demethylcoclobine, C-551
 2-Demethylcolchicine, D-92
 3-Demethylcolchicine, D-92
 2-Demethylcolchiciline, C-568
 10-Demethylcolchicine, D-92
 2-Demethylcolchicine, D-92
 3-Demethylcolchicine, D-92
 2-Demethylcolchifoline, C-571
N-Demethylcolletine, H-275
N-Demethylconoduramine, C-615
 Demethylconoduramine, C-615
N-Demethylconodurine, C-616
N-Demethylcorymine, D-410
 Demethylcorynantheidine, C-689
 Demethylcorynantheine, C-689
 Demethylcrinamine, H-5
O-Demethylcristine, T-53
N-Demethylcrychine, E-221
 1-*O*-Demethylcryprochine, C-781
 10-*De-O*-methylcryptaustoline, C-783
 6'-*O*-Demethyl(-)-curine, C-812
N-Demethylcyanocycline A, C-830
N-Demethylcyclomicurarine L, C-884
N-Demethylcyclomikuranine, C-884
*N*¹-Demethylcynodine, C-931
*N*¹-Demethylcynometrine, C-933
De-N-methyldasycarpidone, D-84
N-Demethyldasymachaline, D-948
*N*⁷-Demethyldauricine, D-88
N-Demethyldauricine, D-88
 3-Demethyl-*N*-deacetylcolchicine, D-92
 Demethyldecaline, V-86
N-Demethyl-4,5-deepoxymaytansinol, M-136
N-Demethyl-20-deethylsuaveoline, S-220
 1-*De-O*-methyldehydroformouregine, T-521
 14-*O*-Demethyldeboxine, D-187
 18-*O*-Demethyldecosine, M-309
*O*¹-Demethyldelelatine, M-321
 6-Demethyldephatine, M-309
 16-Demethyldephatine, M-309
 16-*O*-Demethyldeolsone, M-309
 16-Demethyldeotaline, M-307
 2-Demethyldemecolcine, D-92
 3-Demethyldemecolcine, D-92
N-Demethyldendrobine, D-210
 7-*O*-Demethyl-6-deoxybostrycoidin, B-255
 30-Demethyl-8-deoxylydicamycin, L-353
 7-Demethyldeoxynupharidine, D-201
*O*⁶-Demethyl-9,12-dideoxy-12-oxolythrancine, L-379
 4-Demethyl-20,21-dihydroalstophyllan-17-ol, R-22
 Demethylidihydrocorynantheine, C-689
N-Demethylidihydrogalanthamine, G-3
O-Demethylidihydrogalanthamine, G-3
 Demethylidihydrogalanthamine, G-3
O-Demethyl-2,7-dihydro-*C*-homo-α-erythroidine, 10CI, P-315
N-Demethyl-1,2-dihydro-*N*-hydroxy-1-oxoacronycine, D-417
N-Demethyl-5,6-dihydrooxysanguinarine, D-510
*N*²-Demethylidihydropurpeline, T-288
 5'-*O*-Demethyl-*ent*-dioncophylleine A, D-793
 5'-*O*-Demethylidioncophylline A, D-793
 7'-*O*-Demethyl-α,α'-dioxostebisimine, S-502
N-Demethyldiscarine B, D-836
 10-Demethylidiscrine, X-30
N-Demethylidoryphornine, I-305
 13-Demethylidysidenin, D-972
N-Demethyllechitamine *N*-oxide, D-410
N-Demethyllechitamine, D-410
 Demethyleleutherobin, E-64
 Demethyleberberine, P-34
 9,10-Demethyle-9,10-dehydrosanguinarine, C-356
 Demethylenedelcorine, M-309
 Demethylenedelpheline, M-321
 Demethylenedelidone, M-307
 9,10-Demethylenesanguinarine, C-356
N-Demethyl-16-epiaccedine, A-27
 6-*O*-Demethyl-7-epi-*ancistrobreve* D, A-990
N-Demethyl-3-epimacronine, P-626
O-Demethyl-3-epimacronine, P-626
 5'-*O*-Demethyl-7-epi-8-*O*-methylidioncophylline A, D-793
N-Demethylequisetin, E-142
*O*³-Demethylerysodine, E-203
N-Demethyleschscholtzine, E-221
N-Demethyl-*N*-ethoxycarbonylmacronine, P-626
 9-*O*-Demethyl-7-*O*-ethyllycorenine, L-347
 7-*O*-Demethylevolitrine, F-212
N-Demethyl-*N*-formyldehydronuciferine, D-164
N-Demethyl-*N*-formylleurosine, L-162
N-Demethyl-*N*-formylmesembrenone, M-224
*N*²⁰-Demethyl-*N*²⁰-formylsalonine B, A-876
N-Demethylfranguloline, F-150
N-Demethylfrangulamine, F-151
N-Demethylgalanthamine, G-3
O-Demethylgalanthamine, G-3
 9-*O*-Demethylgalwesine, H-504
*O*¹⁸-Demethylgardneramine oxindole, G-21
 16*R*-*O*¹⁸-Demethylgardneramine oxindole, G-21
*O*¹⁸-Demethylgardneramine, G-20
N-Demethylglaudine, G-87
 Demethylgrossamide, G-182
O-Demethylhaemanthamine, H-5
 5'-*O*-Demethylhamatine, A-982
 5'-*O*-Demethylhamatine, A-982
*N*⁶-Demethylharappamine, M-667
N-Demethylharzianic acid, H-67
 4-Demethylhasubanone, H-70
*O*³-Demethylhasubanone, H-70
N-Demethylhemisine B, H-105
 5-*O*-Demethylhernandezine, H-148

- 2'-Demethylhernandezine, H-148
*O*³-Demethylhernandifoline, L-244
*O*³-Demethylhippeastidine, H-293
 18-De-*O*-methylholinidine, C-588
 9-*O*-Demethylhomolycorine α -*N*-oxide, L-347
 9-*O*-Demethylhomolycorine, L-347
 4-Demethylhumantenine, H-367
 11-*O*-Demethylhumantenirine, H-367
N-Demethyluperzine, H-388
 3-De-*O*-methyl-14 α -hydroxyisotylocrebrine *N*-oxide, I-342
 3-De-*O*-methyl-14 α -hydroxyisotylocrebrine, I-342
N-Demethyl-3-(3-hydroxyisovaleryl)maytansinol, M-136
N-Demethylhygrine, H-759
 1-Demethylhyaconitine, M-343
 De-*N*-methylcaecine, I-11
N-Demethylintegerrine, I-154
N-Demethylintegerrine, I-156
N-Demethylisaomurone, C-781
 Demethylisoalansidine, A-239
O-Demethylisobrucine, P-672
 14-*O*-Demethyl-14-isobutyrylanhweidelphinine, A-1023
 7'-*O*-Demethylisocephaline, E-72
 10-*O*-Demethylisocephaline, E-72
N-Demethylisocorydione, S-367
 13-Demethylisodysidenin, D-972
 8-*O*-Demethylisomigrastatin, I-245
 De-*N*-methyl-1-isoremerine, S-546
N-Demethylisorhoeadine, R-86
 6-*O*-Demethylisotetrandrine, I-328
 7-*O*-Demethylisothalichubine, I-329
 3-De-*O*-methylisotylocrebrine, I-342
N-Demethyljubanine A, J-51
*O*¹²-Demethylkesselringine, K-36
 4-Demethyl-9-ketoeopothilone C, E-115
 1-De-*O*-methylappaconitine, A-91
 Demethylasubine II, L-53
 Demethylasubine I, L-53
 12-*O*-Demethylalauerine, L-67
 Demethylalauterine, O-187
 12-*O*-Demethylimacusine, L-175
 De-*O*-methylisocloclitoxin A, L-205
 2-*O*-Demethyl- β -lumiocolchicine, L-287
 2-*O*-Demethyl- γ -lumiocolchicine, L-287
 3-*O*-Demethyl- β -lumiocolchicine, L-287
 3-*O*-Demethyl- γ -lumiocolchicine, L-287
N-Demethylunidonine, L-295
 14-*O*-Demethyllycaconitine, L-317
N-Demethyllycoramine, G-3
O-Demethyllycoramine, G-3
 30-Demethyllydicamycin, L-353
N-Demethylmacronine, P-626
O-Demethylmacronine, P-626
 8-*O*-Demethylmaritidine, M-108
 9-*O*-Demethylmaritidine, M-108
N-Demethylmasonine, O-68
N-Demethylmaytanacine, M-136
*N*²-Demethylmaytanbutine, M-135
 11-Demethylmecambrine, M-140
O-Demethylmenisarine, M-197
 3-Demethylmescaline, H-465
N-Demethylmesembranol, M-226
 4'-*O*-Demethylmesembranol, M-224
N-Demethylmesembrenol, M-224
 4'-*O*-Demethylmesembrenol, M-224
 4'-*O*-Demethylmesembrenone, M-224
*O*¹-Demethylmetaphanine, M-233
N^b-Demethyl-12-methoxyalstogustine, E-30
 9-Demethyl-8-methoxydihydrochelerythrine, D-407
 4-Demethyl-11-methoxyhumantenine, H-367
O-Demethyl-10-methoxysilanine, T-649
 9-De-*O*-methyl-*N*¹-methylaaptamine, A-2
 3-*O*-Demethyl-2-*O*-methylalangiside, A-239
 4'-*O*-Demethyl-6-*O*-methylancistrocladine, A-982
 2'-*N*-Demethyl-*N*³-methylaplysinopsin, A-1357
 4-Demethyl-*N*-methylatanine, A-1529
 3-Demethyl-*N*-methyldecemcoline, D-92
 5'-*O*-Demethyl-8-*O*-methylidioncophylline A, D-793
 6-*O*-Demethyl-8-*O*-methyl-7-epiancistrobrevine D, A-990
 5'-*O*-Demethyl-8-*O*-methyl-7-epidioncophylline A, D-793
 4'-*O*-Demethyl-6-*O*-methylhamatine, A-982
 3-*O*-Demethyl-2-*O*-methylisoalangiside, A-239
 9-*O*-Demethyl-7-*O*-methyllycorine, L-347
*O*¹²-Demethyl-*O*¹²-methylrodiasine, R-114
O-Demethyl-*N*-methyltuluzosine, T-434
*O*³-Demethyl-17-methyltuluzosine, D-51
N-Demethylmitiphylline, M-651
 8-*O*-Demethylmucosine, M-678
O-Demethylmucronine D, M-720
 Demethylmurrayanine, H-447
N-Demethylmyrianthine C, M-794
 Demethylmyxothiazole A, M-807
 Demethylneoalangiside, A-239
 Demethylneodrupacine, D-202
N-Demethylnicaustrine, T-47
N-Demethylnitidine, N-239
 De-*N*-methylnoracronyline, A-120
 4-Demethylnorhasubanone, H-70
 8-*O*-Demethylnormitidine, N-239
N-Demethylnuciferine, D-531
N-Demethylnummularine D, M-720
 De-*N*-methyl- α -obscurene, O-5
 De-*N*-methyl- β -obscurene, O-5
 Demethylolcypine, A-38
N-Demethyloligophylline, O-82
 De-*O*-methylolivetretin B, T-63
N-Demethyloreodine, G-87
 4-Demethyl-9-oxoeopothilone D(1), E-117
 4-Demethyl-9-oxoeopothilone D(2), E-117
 4-Demethyl-16-oxohasubanone, H-70
 Demethylolaxureline, O-187
 Demethyloxyaaptamine, M-241
 9-Demethyloxychelerythrine, D-407
N-Demethyloxysanguinarine, D-510
*O*¹⁴-Demethylpacidine, M-321
O-Demethylpalposine, A-1499
N-Demethylpamamycin 593A, P-43
N-Demethylpamamycin 593B, P-43
 2'-Demethylpanurensine, P-76
N^b-Demethylpapillotrienine, B-473
N-Demethylparaensidimerine C, P-87
 Demethylpeceyline, P-148
 7-*O*-Demethylpeinamine, P-171
 7-Demethylpericidin A₁, D-203
 4'-*O*-Demethylpipilartine, P-473
N-Demethylpleiocorine, P-519
 De-*O*-methylprodigiosin, P-636
 9-Demethylprotoemetinol, P-664
 10-Demethylprotoemetinol, P-664
 De-*O*-methylpseurotin A, P-738
 9-Demethylpsychotrine, E-72
 18-Demethylpubescence, M-309
N-Demethylpugetinone, P-807
O-Demethylpureine, P-208
N^a-Demethylquaternine, P-417
O-Demethylrenierol acetate, R-45
O-Demethylrenierone, R-45
 7-Demethyl-4'-rhamnopericidin A₁, D-203
 17-De-*O*-methylrhizoxin, R-83
N-Demethylrhoeadine, R-86
N-Demethylricinine, D-668
O-Demethylricinine, D-668
 2'-Demethylrodiasine, R-114
N-Demethylrohitukine 3'-acetate, R-121
N-Demethylrugulovasine A, R-157
N-Demethylrugulovasine B, R-157
 16-Demethylsachaconitine, M-355
 26-Demethylsafaframycin Y₃, S-13
N-Demethylsambutoxin, S-48
N-Demethylsanjoinine A, F-150
*N*³-Demethylsaracodine, P-601
N-Demethylscutianine A, S-165
 6-*O*-Demethylsecoantofine, S-248
 Demethylsecologanol, D-204
N-Demethylsemperviraminone, S-227
 4-*O*-Demethylsimmondsin, S-302
N-Demethylsinomenine, S-313
N-Demethylsonodione, S-367
 1-Demethylsonodione, S-367
 13-Demethylspiroside C, S-432
 13-Demethylspiroside D, S-432
 De-*O*-methylstaurosporine, S-498
 4'-*N*-Demethylstaurosporine, S-498
 7'-*O*-Demethylstebisimine, S-502
 2-*O*-Demethylstephodeline, T-28
 6-Demethylstreptonigrin, S-569
 10'-Demethylstreptonigrin, S-569
*N*⁴-Demethylsternamine, T-8
 6-De-*O*-methyltakatonine, T-16
 17-*O*-Demethyltaxodine, T-51
O-Demethyltazettine, T-53
*N*¹³-Demethylteleocidin A₁, L-367
N-Demethylteleocidin B₁, T-63
N-Demethylteleocidin B₄, T-63
 De-*O*-methyltenuicausine, T-77
 2'-*N*-Demethyltetrandrine, T-286
 2'-Demethylthalibrunine, T-311
 7-*O*-Demethylthalicberine, T-313
 2-*O*-Demethylthalichubine, T-318
 3-*O*-Demethylthalichubine, T-318
*N*²-Demethylthalidasine, T-320
 6'-*O*-Demethylthalifaboramine, T-322
 9-Demethylthallimonine, T-330
 2-Demethylthallimonine, T-330
 5-*O*-Demethylthallistylone, T-337
N-Demethylthallistylone, T-337
N-Demethylthalphenine, T-345
 5-*O*-Demethylthalrugosamine, T-346
N-Demethylthalrugosine, T-320
 7-*O*-Demethylthalrugosinone, T-320
 (Demethylthio)cystodytin B, D-823
N-Demethylthiolutin, A-759
 11-De-*O*-methyltomaymycin, T-424
 9-*O*-Demethyltortuosine, V-35
 1-Demethyltoxoflavin, R-66
 Demethyltrevasine, T-450
 3-Demethyltrichocereine, H-465
 1-*O*-Demethyltricornine, M-309
 12-*O*-Demethyltrilobine, I-339
O-Demethyltsilanine, T-649
 9-Demethyltubulosine, T-675
 10-Demethyltubulosine, T-675
 14-*O*-Demethyltugaconitine, D-187
 7-Demethyltylophorine *N*-oxide, T-695
 7-Demethyltylophorine, T-695
 Demethyltylophorine, T-695
 Demethyltylophorinone, T-696
 De-*N*-methyluleine, U-8
 8-*O*-Demethylvasconine, V-35
 Demethylvertaline, V-86
 Demethylvertine, V-95
N-Demethylvinblastine, V-110
*N*¹-Demethylvincamajine *N*¹-tri-*O*-methylgallate, V-113
 18-Demethylvirescence, M-325
N-Demethylvoacamine, V-168
 12-Demethylvobtusine, V-188
 1-Demethylwinkleridine, M-325
N-Demethylxestomanzamine A, X-23
 Demethylxestospongine B, X-25
 Demissidine, D-205
 Demissine, D-205
 β ₁-Demissine, D-205
 Δ^5 -Demissine, S-345
 Dendramine, D-210
 Dendridine A, D-206
 Dendrine, D-207
 Dendroamide A, D-208
 Dendroamide B, D-209
 Dendroamide C, D-209
 Dendroban-12-one, D-210
 Dendrobine *N*-oxide, D-210
 Dendrobine, D-210
cis-Dendrochrysanine, H-759
trans-Dendrochrysanine, H-759
 Dendrochrysanine, D-211
 Dendrocrepine, D-212
 Dendrodoine, D-213
 Dendronobilin A, D-214
 Dendroprimine, O-42
 Dendrowardine, D-215
 Dendroxine, D-216

- Densiberine, P-79
 Densiflorine, D-217
 Densiflorine, D-218
 Densinine, D-219
 Denudatidine, D-220
 Denudatine, D-221
 Deoxaphomin, A-1460
 13-Deoxo-13-acetyloxy-1-deoxytaxine B, T-47
 13-Deoxo-13-acetyloxytaxine B, T-45
 Deoxoamphimedine, D-222
 Deoxoangustifoline, A-1013
 8-Deoxoantidesmone, A-1325
 Deoxoapodine, M-666
 Deoxoaralione A, A-1375
 Deoxoaristolasicone, A-1432
 15-Deoxo-13-deoxy-15-hydroxymadumycin I, M-35
 3-Deoxo-3-hydroxylunarine, L-293
 9-Deoxo-9 α -hydroxytaxol, T-52
 11-Deoxojervine, J-36
 2-Deoxoparaherquamide A, P-90
 13-Deoxy-13-acetyloxy-1-deoxynortaxine B, T-47
 3-Deoxyaconine 8-linoleate, M-343
 Deoxyaconitine, M-343
 5-Deoxyadenophorine, E-244
 1-Deoxyadenophorine, E-254
 Deoxyadifoline, A-140
 Deoxyadrenaline, E-98
 1-Deoxy-D-allonjirimycin, H-616
 1-Deoxy-D-altronojirimycin, H-616
 1-Deoxy-L-altronojirimycin, H-616
 2-Deoxy-2-aminokealiquinone, K-25
 25-Deoxy-25-aminosafrafrimycin A, S-11
 Deoxyamphimedine, D-222
 Deoxyangustifoline, A-1013
 Deoxyaniflorine, A-1039
 Deoxyaralione A, A-1375
 Deoxyaspergillid acid, A-1475
 Deoxyaspidodispermine, A-1493
 17-Deoxyaspochalasin C, A-1502
 12-Deoxyaxillarine, A-1572
 10-Deoxy-4,4'-binectorone, N-85
 3-Deoxyblastmycetin B, B-232
 3-Deoxyblastmycetin C, B-232
 6-Deoxybostrycoidin, B-255
 Deoxybrevianamide E, C-861
 3-Deoxybuxandone, B-459
 28-Deoxybuxanoldine, B-460
 16-Deoxybuxidienine C, B-466
 Deoxycalyciphyllyne B, C-45
 20-Deoxycamptothecin, C-70
 7-Deoxycephalofortuneine, H-331
 13-Deoxychilenine, C-375
 Deoxychromoviridan, C-426
 13-Deoxycopiamycin, C-634
 Deoxycordifoline lactam, D-223
 Deoxycordifoline, C-641
 3-Deoxycrassicauldine, M-313
 4 α -Deoxycrinamabine, A-670
 9-Deoxycurtisin, H-635
 7-Deoxycylindrospermopsin, C-930
 2-Deoxydaphnezomine S, D-53
 8-Deoxy-14-dehydroaconosine, A-100
 Deoxydehydrochorismic acid, D-224
 Deoxydehydromutaaspergillid acid, M-462
 8-Deoxy-13-dehydroseratinine, S-257
 Deoxydelcorine, M-325
 6-Deoxydelpheline, M-348
 Deoxydelsoleine, M-328
 18-Deoxydiaboline, R-63
 1-Deoxydiacetylaxine B, T-273
 7-Deoxy-*trans*-dihydronarciclasine, N-37
 4-Deoxy-7,8-dihydrotubastrine, T-656
 Deoxydiphthocarpaine, D-828
 Deoxydiphthocarpamine, D-828
 Deoxydiptocarpaine, D-828
 Deoxydiptocarpamine, D-828
 Deoxydynamicin A, D-959
 7-Deoxy-6-epicastanospermine, C-188
 17-Deoxyepilamprolobine, L-26
 11-Deoxy-4-epitetradotoxin, T-298
 28-Deoxy-22-epizoanthamine, Z-31
 Deoxyepothilone A, E-115
 Deoxyepothilone B, E-117
 18-Deoxy-19,20-epoxycytochalasin R, C-954
 11-Deoxy-6,11-epoxy-13-methylchelidonine, C-693
 4'-Deoxy-3',4'-epoxy-22-oxovincalcolchicine, L-162
 4'-Deoxy-3',4'-epoxyvincalcolchicine, L-162
 4-Deoxyester A, R-175
 4-Deoxyeuonymine, P-244
 6-Deoxyfagomine, M-516
 11-Deoxyfistularin 3, F-70
 19-Deoxyfistularin 3, F-70
 10-Deoxyflavaconitine, M-343
 2'-N-(1-Deoxy- β -D-fructopyranos-1-yl)cephaline, E-72
 2'-N-(1-Deoxy- β -D-fructopyranos-1-yl)neocephaline, E-72
 Deoxyfructosazine, T-250
 1-Deoxy-D-galactonojirimycin, H-616
 1-Deoxy-D-galactostatin, H-616
 7-Deoxygermine, Z-36
 9-Deoxyglanduline, H-178
 Deoxyguadiscidine, G-188
 1-Deoxy-D-gulonojirimycin, H-616
 1-Deoxy-L-gulonojirimycin, H-616
 Deoxyharringtonic acid, H-66
 Deoxyharringtonine, H-66
 1-Deoxy-1-(1,2,3,6,7,8-hexahydro-2,6,8-trioxo-4H-imidazo[4,5-g]pteridin-4-yl)-D-ribose, R-166
 Deoxyhomoharringtonine, H-66
 7-Deoxy- α -homomannonojirimycin, H-607
 7-Deoxy- β -homomannonojirimycin, H-607
 7-Deoxy- α -homonojirimycin, H-607
 10-Deoxy-17-hydroxyakuammine, A-232
 Deoxyhydroxyaspergillid acid, A-1475
 4-Deoxy-20-hydroxyryanodine, R-175
 19-Deoxy-15 β -hydroxyuscharin, U-53
 9-Deoxyhypogonavine, H-195
 1-Deoxyhypogonavine, H-200
 1-Deoxyhypogonavinol, H-200
 1-Deoxy-D-idojirimycin, H-616
 1-Deoxy-L-idojirimycin, H-616
 1-Deoxy-1,5-imino-D-allitol, H-616
 1-Deoxy-1,5-imino-D-gulitol, H-616
 Deoxyindolizomycin, I-101
 5'-Deoxy-5-iodotubercidin, T-658
 Deoxyisocalyciphyllyne B, C-45
 Deoxyjesaconitine, M-343
 11-Deoxykobusine, H-172
 Deoxykopsijasminilam, K-72
 9-Deoxylappaconitine, A-95
 20'-Deoxyleurosine, V-110
 13-Deoxyludaconitine, M-303
 18-Deoxylycoctonine, M-321
 Deoxylycorenine, L-347
 8-Deoxylydicamycin, L-353
 9-Deoxylythrancine, L-377
 8'-Deoxymalngamide C, D-225
 Deoxymalngamide C, D-225
 1-Deoxymannonojirimycin, H-616
 6-Deoxymanzamine X, M-91
 4-Deoxymayteine, P-244
 Deoxymesaconitine, M-343
 2'-Deoxy-3-methyladenosine, A-844
 6-Deoxy-8-methylbostrycoidin, D-226
 N-Deoxymilitarinone A, M-610
 5-Deoxymiriamide, D-639
 Deoxymutaaspergillid acid, M-462
 7-Deoxynarciclasine, N-37
 18-Deoxynargenicin A₁, N-276
 Deoxyneo- β -hydroxyaspergillid acid, F-76
 1-Deoxynojirimycin 6-phosphate, H-616
 1-Deoxynojirimycin, H-616
 1-Deoxy-5-nornojirimycin, P-458
 N-Deoxynortryptoquivaline, N-315
 Deoxynortryptoquivalone, N-315
 Deoxynupharidine, D-227
 Deoxynupharidine-6,7-diol, D-227
 Deoxynybomycin, N-346
 15-Deoxyoxalicine A, O-148
 15-Deoxyoxalicine B, O-148
 19-Deoxy-11-oxofistularin 3, F-70
 2-Deoxy-2-(5-oxo-2-pyrrolidinedicarboxamido)glucopyranose, P-937
 1-Deoxy-1-oxosongorine, N-30
 2-Deoxy-18-oxovobtusine, V-188
 2-Deoxy-3'-oxovobtusine, V-188
 6'-Deoxyoxysporidinone, O-226
 1-Deoxyacitaxel, E-138
 Deoxyalustrine, P-40
 Deoxyalpytoxin, P-42
 7-Deoxypancratistatine, P-58
 14-Deoxyparsonsianidine, P-96
 14-Deoxyparsonsianine, P-97
 16-Deoxyparsonsianine, P-97
 Deoxypeganidine, P-164
 Deoxypeganone N-oxide, T-231
 Deoxypeganone, T-231
 14-Deoxypergularinine, T-696
 4-Deoxyphorbol, T-275
 1'-Deoxypsammalylin D, P-695
 N⁷-2'-Deoxypseudoanthosine, X-5
 7-Deoxypubescence, M-323
 Deoxypumiliotoxin 251H, D-228
 (3R)-Deoxypumilolide, P-796
 (3S)-Deoxypumilolide, P-796
 Deoxypridinoline, P-907
 1-Deoxy-1-[2-(3-pyridinyl)-1-pyrrolidinyl]- β -D-fructopyranose, P-957
 Deoxyrhexifoline, B-254
 Deoxyrubroflavin, D-229
 1-Deoxysecofascaplysin A, R-57
 Deoxysepiapterin, S-246
 8-Deoxyserratinine, S-256
 8-Deoxyserratinine, S-257
 15-Deoxyspergualin, S-394
 1-Deoxy-D-talonojirimycin, H-616
 3'-Deoxytalopiericidin A₁, P-421
 9-Deoxytauroacidin B, T-39
 1-Deoxytaxol, E-138
 1-Deoxy-1-[1,3,4,7-tetrahydro-6-(hydroxymethyl)-2,4,7-trioxo-8(2H)-pteridinyl]-D-ribose, P-382
 5-Deoxytetradotoxin, T-510
 11-Deoxytetradotoxin, T-298
 Deoxythalidastine, T-321
 Deoxytoposentin, T-432
 C-Deoxytoxiferine, T-436
 5'-Deoxytoyocamycin, S-53
 Deoxytryptoquivaline, T-645
 3'-Deoxytubastrine, T-656
 5'-Deoxytubercidin, T-658
 Deoxytubulosine, T-675
 Deoxytylophorinine N-oxide, T-696
 19-Deoxyuscharin, U-53
 Deoxyvasicine, T-231
 Deoxyvasicinone N-oxide, D-503
 Deoxyvasicinone, D-503
 16-Deoxyveralkamine, V-60
 23-Deoxyveratramine, V-71
 Deoxyverrucarin E, A-71
 11-Deoxyverticillin A, V-89
 Deoxyvinblastine A, V-110
 20'-Deoxyvinblastine, V-110
 Deoxyvincalcolchicine A, V-110
 Deoxyvincalcolchicine B, V-110
 Deoxyviolacein, V-142
 2-Deoxyvobtusine lactam, V-188
 2-Deoxyvobtusine lactone, V-188
 2-Deoxyvobtusine, V-188
 21-Deoxyvomilenine, V-194
 16-Deoxy-Wieland-Gumlich aldehyde, R-63
 Deoxyyuzurimine, D-33
 2'-Deoxyzeatin riboside, Z-11
 28-Deoxyzoanthamine, Z-31
 Dephosphonocalyculin A, C-55
 Deplancheine, D-230
 Deppeanol, T-141
 N-Depropionyldecorticasine, L-234
 Deracil, D-518
 Dercitamide, D-93
 Dercitamine, D-93
 Dercitine, D-231
 Dermalex, A-625
 Derquantel, P-90
 Desacetoxylaphiphylline, D-68
 Desacetoxylvindoline, V-130
 Desacetylcolchicine, D-92

- Desacetylcolchicine, D-92
 Desacetyldaphnijsmine, D-33
 Desacetyldesformoakuumilline, S-576
 Desacetyldoronine, O-92
 N^a-Desacetyl-18-hydroxyisoretuline, R-63
 Desacetylisoretulinol, R-63
 N^a-Desacetylisoretuline, R-63
 Desacetylisoretuline, R-63
 Desacetylneoprotoveratrine, P-674
 O-Desacetyl pachysandrine B, D-266
 Desacetylprotoveratrine A, P-674
 Desacetylprotoveratrine B, P-674
 Desacetylretuline, R-63
 Desacetylpermostrychnine, S-400
 Desacetylacraline, T-15
 1-Desacetylwilforine, P-253
 Desacetylcynanchogenin, T-268
 Desacetylcondurangenin C, P-603
 O,N-Desacetyl-N-methylpachysandrine A, D-266
 Desaminocanavanine, H-235
 3-Desamino-22-O-ethyl-3-hydroxysolanocapsine, D-99
 3-Desamino-3-hydroxysolanocapsine, D-99
 Desanthraserpidine, D-687
 Desatrine, P-674
 Desbenzoylaralione A, D-100
 Descarbomethoxyconopharyngine, I-4
 16-Descarbomethoxy-16,17-dihydro-17-hydroxy-19-epiajmalicine, A-214
 16R-Descarbomethoxytacamine, T-13
 16S-Descarbomethoxytacamine, T-13
 2-Descarboxybetanidin, D-122
 2-Descarboxybetanin, D-122
 2-Descarboxyphyllolactin, D-122
 Deschloroanhydroaureomycin, A-1034
 11-Deschlororebeccamycin, R-32
 Desclidium, Q-17
 Descrespt, M-262
 Desdamethine, D-232
 Desdanine, P-876
 Deserpidine, D-233
 Deserpidic acid, D-687
 Deserpidine, D-234
 Deserpidinic acid, D-687
 N-Desethyl-N-methylaconitine, M-331
 N-Desethyl-19-oxoanthranoyllycoctonine, H-123
 N-Desethylpacifiline, H-123
 Desferal, D-142
 Desferrin, D-142
 Desferrioxamine mesilate, D-142
 Desferrioxamine, D-142
 Desferrihiocin, D-235
 Desferroferrithiocin, D-235
 Desformoakuumidinol, T-425
 Desformocubamine, V-125
 Desformocorymine, D-236
 Desformylstemmadenine, S-510
 Deshydroxyacetylhenningssine, H-110
 Deshydroxyacetylhenningssine, H-111
 Desmethoxychoisyne, C-418
 Desmethoxyyboluteine, I-5
 N-Desmethoxyrankinidine, H-367
 Des-N-methylacronycine, A-120
 Des-N^{1a}-methylanhydromacralstonine, M-8
 N-Desmethyllumepavine, A-1444
 Des-N-methylavicine, A-1566
 Desmethylcephalotaxinone, I-201
 Des-N-methylcherythrine, C-356
 Des-N-methylcitpressine II, T-243
 Des-N-methylcitpressine I, T-243
 2-Desmethylcolchicine, D-92
 N²-Desmethylcycleanine, C-835
 O-Desmethylidihydrochelerithrinyll-11-acetone, A-34
 O-Desmethylidihydrochelerithrinyll-8-acetone, A-34
 N-Desmethyl-1,2-dihydro-1-hydroxyacronycine, D-417
 Desmethylleutherobin, E-64
 Desmethylemetine, E-72
 3-Desmethylerysoline, E-203
 Des-N-methylfastigiataine, F-17
 5'-Des-O-methylharringtonine, H-66
 3''-Des-O-methylharringtonine, H-66
 5'-Des-O-methylhomoharringtonine, H-66
 3''-Des-O-methylhomoharringtonine, H-66
 7-Desmethylhuangshanine, H-366
 N-Desmethylindolactam V, I-78
 5'-Des-O-methylisoharringtonine, I-238
 N-Desmethyljubanine B, J-52
 Desmethylmyxothiazole A, M-807
 N-Desmethylnitidine, N-239
 Des-N-methylnoracronycine, A-120
 N-Desmethylnuciferine, D-531
 Des-O-methylolivetin C, O-88
 Desmethylphidolopin, P-357
 13-Desmethylspirolide C, S-432
 N-Desmethylthalidezine, H-148
 Desmethylzanthophylline, F-90
 Desmodilactone, A-800
 Desmodimine, D-237
 Desmotrichinine, D-238
 Desoxoviroidin, V-160
 Desoxoviroisin, V-160
 12-Desoxyaxillarine, A-1572
 16-Desoxybutadienine, B-466
 Desoxycabufiline, C-3
 Desoxycordifoline, C-641
 Desoxycordifolinic acid, C-641
 Desoxycyclobuxoxazine A, C-851
 Desoxyepinephrine, E-98
 4-Desoxyevonine, P-248
 17-Desoxy-cis-lamprolobine, L-26
 14-Desoxy-13a-methyltylohisutinidine, M-583
 2-Desoxyryosenaminol, H-166
 Desoxysarpagine, T-425
 Desoxy-Wieland-Gumlich aldehyde, C-115
 Desthiobiotin, M-499
 Detamide, M-398
 Dethiobiotin, M-499
 Dethiotetrakis(methylthio)chetomin, D-239
 Dethiotetra(methylthio)chetomin, D-239
 Detoxinine, D-240
 Deveratrolylesoline, Y-10
 DGCC, D-243
 DGTA, D-244
 DGTS, D-245
 DHA-paclitaxel, T-52
 DHBOA-Glc, D-538
 DHQD-CLB, Q-18
 Dhurrin 6'-glucoside, H-520
 Dhurrin, H-520
 Diaboline N-oxide, C-115
 Diaboline, C-115
 Diabroticin A, D-241
 Diabroticin B, D-241
 Diacetoneamine, A-839
 2,5-Diacetonyl-1-methylpyrrolidine, H-704
 1,14-Diacetoxy-9-benzoyloxy-4,6-dihydroxy-2-isobutanoyloxy-8-nicotinoyloxydihydro-β-agarofuran, H-119
 8,9-Diacetoxy-1-benzoyloxy-4-hydroxy-6-nicotinoyloxydihydro-β-agarofuran, P-219
 3,4-Diacetoxy-N-cinnamoylpyrrolidine, S-87
 2,4-Diacetoxyepipachysamine D, D-262
 2,16-Diacetoxy-9,11-epoxybuxamidine, B-477
 2,14-Diacetoxy-1-furoyloxydihydro-6-(2-methylbutanoyloxy)-9-nicotinoyloxy-β-agarofuran, H-247
 2,6-Diacetoxy-9-furoyloxydihydro-14-(2-methylbutanoyloxy)-1-nicotinoyloxy-β-agarofuran, H-247
 2,14-Diacetoxy-1-furoyloxydihydro-6-(2-methylpropanoyloxy)-9-nicotinoyloxy-β-agarofuran, H-247
 2,6-Diacetoxy-9-furoyloxydihydro-14-(2-methylpropanoyloxy)-1-nicotinoyloxy-β-agarofuran, H-247
 8,14-Diacetylaconosine, A-99
 3,7-Diacetyl-15-angeloylgermine, G-70
 Diacetylanthranyllic acid, A-31
 16,28-Diacetylbusanoldine, B-460
 Diacetylcevedine, C-295
 8,14-Diacetylchasanine, M-323
 O,N-Diacetyl-N-demethylgalanthamine, G-3
 11,13-Di-O-acetyl-9-deoxyglanaduline, H-178
 Diacetyldihydrolycorine, Z-15
 N,2-Diacetyl-4,5-dimethoxyphenethylamine, D-219
 Di-O-acetyl-3-epidihydrocorymine, D-410
 N,N-Diacetylepipachysamine C, P-601
 11,13-Di-O-acetylglanaduline, H-170
 Diacetylglycoperine, F-223
 6,8-Diacetylheteratisine, H-157
 11,13-Diacetylhetisine, H-203
 1,15-Di-O-acetylhyppoglavine, H-179
 3',7-Diacetylintermedine, I-160
 Diacetylohyppoglavine, H-203
 Di-O-acetylleptimidine, L-116
 Diacetyllycofoline, L-328
 3',7-Diacetyllycopsamine, I-160
 1,2-Di-O-acetyllycorine, L-348
 O,O-Diacetylmacranthine, M-9
 8,14-Diacetylneoline, M-323
 1,14-Diacetylneoline, M-323
 N,O-DiacetylNorornuciferidine, P-4
 N,O-DiacetylNoroliveroline, O-87
 1,10-Diacetyl-10-norpluviin, P-528
 Diacetylpsuedaconitine, M-336
 6,7-Di-O-acetylsinococuline, S-311
 7,10-Diacetylaxine A, A-6
 Diacetylaxine B, T-45
 2',7-Diacetylaxol, T-52
 N^b-Diacetyltryptamine, T-639
 3,5-Diacetyl-1,4-dihydro-4-nonylpyridines, D-242
 1,2-Diacetylglyceryl-3-(O-carboxyhydroxymethyl)-choline, D-243
 1,2-Diacetylglyceryl-O-2'-(hydroxymethyl)-N,N,N-trimethyl-β-alanine, D-244
 4-O-(1,2-Diacetylglyceryl)-N,N,N-trimethylhomoserine, D-245
 Dialanine anhydride, D-769
 Dialanine dianhydride, D-769
 2,5-Dialkylpyrrolidines, D-246
 2,4-Dialkylthiazoles, D-247
 2,5-Diallyldecahydroquinoline, A-357
 Diamethine, C-420
 Diamid Y, D-889
 Diamidinocadaverine, D-401
 Diamidinospemidine, H-305
 1,7-Diamino-4-azapeptane, D-255
 1,9-Diamino-4-azanonane, A-897
 1,9-Diamino-5-azanonane, D-252
 1,8-Diamino-4-azaocetane, S-398
 1,4-Diaminobutane, B-422
 1,4-Diaminobutane-N,N'-dipropanoic acid, S-397
 1,4-Diamino-2-butanol, D-248
 1,4-Diamino-2-butene, B-423
 3,20-Diamino-9,19-cyclopregnan-2-ol, C-903
 1,10-Diamino-4,7-diazadecane, B-153
 1,12-Diamino-4,8-diazadodecane, D-249
 1,12-Diamino-4,9-diazadodecane, S-399
 1,14-Diamino-5,10-diazatetradecane, B-150
 1,13-Diamino-4,9-diazatridecane, D-250
 1,13-Diamino-5,9-diazatridecane, D-251
 1,11-Diamino-4,8-diazadodecane, B-154
 4,4'-Diaminodibutylamine, D-252
 α,α'-Diamino-6',7'-dihydro-6',7'-dioxo-[2,4'-bi-1H-indole]-3,3'-dipropanoic acid, T-641
 α,2-Diamino-4,5-dihydro-1H-imidazole-4-propanoic acid, E-80
 5,5'-Diamino-4,4'-dihydroxy-3,3'-diaz-2,2'-diphenone, I-70
 4,5-Diamino-2,6-dihydroxypyrimidine, D-254
 2,4-Diamino-5,6-dihydroxypyrimidine, D-253
 3,20-Diamino-4,14-dimethyl-9,19-cyclopregnane, B-474
 3,20-Diamino-4,14-dimethyl-9,19-cyclopregnane-4-methanol, C-888
 3,20-Diamino-4,14-dimethyl-9,19-cyclopregnan-16-ol, C-846
 3,3'-Diaminodipropylamine, D-255
 4,5-Diaminoecicosane, E-47
 1,2-Diaminoethane, E-225
 2,4-Diamino-3-hydroxybenzoic acid, D-256
 3,20-Diamino-16-hydroxy-4,14-dimethyl-9,19-cyclopregnane-4-methanol, C-883
 3,20-Diamino-16-hydroxy-4,14-dimethyl-9,19-cyclopregn-6-ene-4-methanol, C-883
 3,20-Diamino-16-hydroxy-4-hydroxymethyl-4,14-dimethyl-9,19-cyclopregnan-11-one, B-12

- 3,20-Diamino-4-hydroxymethyl-4,14-dimethyl-9,19-cyclopregnan-16-one, C-883
- 3,20-Diamino-4-hydroxypregnane, D-266
- 3,20-Diamino-15-hydroxypregn-5-ene, D-278
- 3,20-Diamino-18-hydroxypregn-5-ene, D-279
- 3,20-Diamino-18-hydroxypregn-5-ene-11,16-dione, D-257
- 2',3'-Diamino-3'-hydroxypropiofenone, A-697
- 2,6-Diamino-5-hydroxy-4(1H)-pyrimidinone, D-253
- 3,20-Diamino-28-hydroxy-4,4,14-trimethyl-9,19-cyclopregnan-11-one, C-921
- α ,3-Diamino-1H-indene-2-propanoic acid, A-698
- 2,6-Diamino-4-methyl-1H-cyclohepta[1,2-d:4,5-d']diimidazole, P-94
- 2,8-Diamino-4-methyl-1H-cyclohepta[1,2-d:3,4-d']diimidazole, P-737
- 3,20-Diamino-4-methylene-14-methyl-9,19-cyclopregnan-16-ol, C-846
- 3,20-Diamino-11-methylenepregnane, M-460
- 3,3'-Diamino-N-methylpropylamine, D-255
- 1,17-Diamino-9-octadecene, O-35
- 2,5-Diamino-5-oxopentanoic acid, G-108
- 1,5-Diaminopentane, P-229
- 1,5-Diamino-1-pentene, P-232
- 3,20-Diaminopregna-14,16-diene, P-600
- 3,20-Diaminopregna-4,14-diene, P-598
- 3,20-Diaminopregna-5,16-diene, P-599
- 3,20-Diaminopregna-4,14-dien-2-ol, D-258
- 3,20-Diaminopregna-4,16-dien-2-ol, D-259
- 3,20-Diaminopregna-2,16-dien-4-one, D-260
- 3,20-Diaminopregna-5,14-dien-4-one, D-261
- 3,20-Diaminopregnan-18-ol, D-269
- 3,20-Diaminopregnan-2,4-diol, D-262
- 3,20-Diaminopregnan-4,12-diol, D-263
- 3,20-Diaminopregnan-4,16-diol, D-264
- 3,20-Diaminopregnan-16-diol
- 3,20-Diaminopregnan-11-ol, D-267
- 3,20-Diaminopregnan-12-ol, D-268
- 3,20-Diaminopregnan-18-ol, D-269
- 3,20-Diaminopregnan-2-ol, D-265
- 3,20-Diaminopregnan-4-ol, D-266
- 3,20-Diaminopregn-14-ene-2,4-diol, D-272
- 3,20-Diaminopregn-16-ene-2,4-diol, D-273
- 3,20-Diaminopregn-5-ene-15,18-diol, D-270
- 3,20-Diaminopregn-5-ene-16,18-diol, D-271
- 3,20-Diaminopregn-14-ene, P-607
- 3,20-Diaminopregn-16-ene, P-608
- 3,20-Diaminopregn-4-ene, P-605
- 3,20-Diaminopregn-5-ene, P-606
- 3,20-Diaminopregn-16-ene-4-ol, D-280
- 3,20-Diaminopregn-2-ene-4-ol, D-274
- 3,20-Diaminopregn-4-ene-2-ol, D-275
- 3,20-Diaminopregn-5-ene-15-ol, D-278
- 3,20-Diaminopregn-5-ene-18-ol, D-279
- 3,20-Diaminopregn-5-ene-2-ol, D-276
- 3,20-Diaminopregn-5-ene-4-ol, D-277
- 3,20-Diaminopregn-2-ene-4-one, D-281
- 3,20-Diaminopregn-5-ene-4-one, D-282
- 2',3'-Diaminopropiophenone, A-700
- 4-[[[(2,4-Diamino-6-pteridiny)methyl]amino]benzoic acid, A-899
- 4-[[[(2,4-Diamino-6-pteridiny)methyl]methylamino]benzoic acid, A-899
- p*-[N-2,4-Diamino-6-pteridyl]methylaminobenzoic acid, A-899
- 2,6-Diaminopurine, P-817
- 5,6-Diamino-2,4-pyrimidinediol, D-254
- 2,4-Diamino-5,6-pyrimidinediol, D-253
- 5,6-Diamino-2,4(1H,3H)-pyrimidinedione, D-254
- 1,20-Diamino-4,8,12,16-tetrazaeicosane, D-283
- 1,20-Diamino-4,8,12,17-tetrazaeicosane, D-284
- 1,20-Diamino-4,8,13,17-tetrazaeicosane, D-285
- 1,24-Diamino-5,10,15,20-tetraazatetracosane, D-286
- α ,2-Diamino-1,4,5,6-tetrahydro-6-hydroxy-4-pyrimidineacetic acid, T-657
- α ,2-Diamino-1,4,5,6-tetrahydro-4-pyrimidineacetic acid, C-99
- α ,2-Diamino-1,4,5,6-tetrahydro-4-pyrimidinepropanoic acid, T-199
- α , α' -Diamino-5,5',6,6'-tetrahydroxy-3,3'-biphenyldipropionic acid, A-745
- 1,17-Diamino-4,9,13-triazaheptadecane, D-287
- 1,16-Diamino-4,8,12-triazaheptadecane, D-288
- 1,16-Diamino-4,8,13-triazaheptadecane, D-289
- 1,19-Diamino-6,11,16-triazaonadecane, D-290
- 1,18-Diamino-5,9,14-triazaoctadecane, D-291
- 1,15-Diamino-4,8,12-triazapentadecane, T-456
- 2,27-Diamino-3,26,28-trihydroxy-11-octacosanone, D-292
- 3,20-Diamino-4,4,14-trimethyl-9,19-cyclopregnan-16-ol, C-917
- 3,20-Diamino-4,4,14-trimethyl-9,19-cyclopregnan-11-one, C-920
- 3,20-Diamino-4,4,14-trimethyl-9,19-cyclopregn-11-en-16-ol, C-917
- 3,20-Diamino-4,4,14-trimethyl-9,19-cyclopregn-6-en-16-ol, C-915
- 5,6-Diaminouracil, D-254
- 7,9-Diangeloylheliotridine, T-188
- 7,9-Diangeloyl-1-hydroxy-N-methylplatynecium(1+), D-596
- Diangeloylplatynecine, H-233
- Diangoutengian I, A-953
- Dianthalexine, H-675
- Dianthramide A (obsol.), B-67
- Dianthramide B (obsol.), B-67
- Dianthramide B, B-67
- Dianthramide P, A-1563
- Dianthramide R, B-67
- Dianthramide S methyl ester, B-67
- Dianthramide S, B-67
- Dianthramine, I-45
- Diaporthichalasin, P-376
- Diatom polyamines, D-293
- 1,4-Diazabicyclo[4.3.0]nonane-2,5-dione, C-865
- 1,13-Diazacyclotetracosane-2,14-dione, D-294
- 4,9-Diazadodecanedioic acid, S-397
- 1,6-Diazafluoranthene, I-65
- 2,9-Diazafluorene, C-130
- 2,9-Diazaphenanthren-1(2H)-one, B-75
- Diazaquinomycin A, D-295
- Diazaquinomycin B, D-295
- Diazaquinomycin C, D-296
- Diazepinomicin, D-297
- 1,3-Diazole, I-35
- 6-Diazo-3-methyl-4-(1,3,5-trimethyl-1-hexenyl)-2,5,7,8(1H,6H)-quinolinetetrone, L-12
- Diazonamide A, D-298
- Diazonamide B, D-298
- Diazonamide C, D-298
- Diazonamide D, D-298
- Diazonamide E, D-298
- Dibenzo[*b,h*]dipyrido[4,3,2-*de*:2',3',4'-*gh*][1,10]phenanthroline, D-299
- Dibenzopyrrole, C-120
- 3,6-Dibenzoyloxytropane, M-391
- 3,7-Dibenzoyloxytrop-6-ol, M-392
- N^1,N^{10} -Dibenzoylspermidine, S-398
- 10-(Dibenzylamino)-1-hydroxy-2-methoxyaporphine, V-24
- 3,6-Dibenzyl-2,5-dihydroxypyrazine, D-300
- 1,3-Dibenzyl-4,5-dimethylimidazolium(1+), D-750
- 2,5-Dibenzyl-1,4-dimethylpiperazine, D-301
- N,N'*-Dibenzyl-2-hydroxybutanediamide, M-72
- 2,5-Dibenzyl-3-hydroxy-6-methoxypyrazine 1-oxide, D-300
- 2,5-Dibenzyl-3-hydroxy-6-methoxypyrazine, D-300
- 3,6-Dibenzyl-5-hydroxy-2(1H)-pyrazinone, D-300
- N,N'*-Dibenzylmalamide, M-72
- 2,5-Dibenzylpiperazine, D-301
- 1,3-Dibenzyl-2,4,5-trimethylimidazolium(1+), T-607
- N,N'*-Dibenzylurea, D-302
- DIBOA-Glc, H-409
- DIBOA, H-409
- Dibromoacetamide, D-303
- Dibromoacetic acid, D-303
- Dibromoacetone, D-303
- Dibromoacetonitrile, D-303
- Dibromoagelaspongins methyl ether, A-188
- Dibromoagelaspongins, A-188
- 2,2'-Dibromoageliferin, A-190
- 4,6-Dibromo-2,3-bis(methylsulfanyl)-1H-indole, D-304
- 4,6-Dibromo-2,3-bis(methylthio)-1H-indole, D-304
- 2,4-Dibromo-5-(5-bromo-2-hydroxybenzoyl)-3-chloropyrrole, P-966
- 3,4-Dibromo-2-(5-bromo-2-hydroxybenzoyl)-5-chloropyrrole, P-966
- 2,3-Dibromo-4-(3-bromo-2-nitrophenyl)-1H-pyrrole, D-305
- 2,3-Dibromo-4-(5-bromo-2-nitrophenyl)-1H-pyrrole, D-305
- Dibromocantharelline, D-328
- 2,7-Dibromo-9H-carbazole, D-306
- 3,6-Dibromo-9H-carbazole, D-307
- 4,6-Dibromo-3-(2-chloroethyl)-1,3-dihydro-3-hydroxy-2H-indol-2-one, C-629
- 2,7-Dibromo-3-chloro-1H-indole, D-308
- Dibromocyanomethane, D-303
- 2,3-Dibromo-4-cyano-5-methoxycarbonylpyrrole, D-340
- 2,6-Dibromo-4-(cyanomethyl)anisole, D-321
- 3,5-Dibromo-1-cyanomethyl-4-methoxy-3,5-cyclohexadiene-1,2-diol, A-157
- 2,6-Dibromo-4-(cyanomethyl)phenol, D-321
- 5,6-Dibromo-2'-N-demethylaplysinopsin, A-1357
- Dibromodeoxytopsentin, T-432
- 5,5'-Dibromo-4,4'-dichloro-7,7'-dimethoxy-3,3'-bi-1H-indole, D-309
- 4,7-Dibromo-2,3-dichloro-1H-indole, D-310
- 4,6-Dibromo-1,3-dihydro-3-hydroxy-3-(2-hydroxyethyl)-2H-indol-2-one, C-629
- 4,6-Dibromo-1,3-dihydro-3-hydroxy-3-methyl-2H-indol-2-one, C-629
- 4,6-Dibromo-1,3-dihydro-3-hydroxy-3-(2-oxopropyl)-2H-indol-2-one, C-629
- 6,7-Dibromo-3,4-dihydro-4-hydroxypyrrolo[1,2-*a*]pyrazin-1(2H)-one, L-242
- 2,4-Dibromo-3,6-dihydroxybenzeneacetamide, D-313
- 2,4-Dibromo-3,6-dihydroxybenzeneacetic acid, D-313
- 3,5-Dibromo-2,4-dihydroxybenzeneacetic acid, D-314
- 7-(2,3-Dibromo-4,5-dihydroxybenzyl)-3,7-dihydro-1H-purine-2,6-dione, D-311
- N*-(2,3-Dibromo-4,5-dihydroxybenzyl)-*N'*-(3-methoxycarbonylpropyl)urea, D-312
- 3,5-Dibromo-1,6-dihydroxy-4-methoxy-2,4-cyclohexadiene-1-acetonitrile, A-157
- 2,4-Dibromo-3,6-dihydroxyphenylacetamide, D-313
- 2,4-Dibromo-3,6-dihydroxyphenylacetic acid, D-313
- 3,5-Dibromo-2,4-dihydroxyphenylacetic acid, D-314
- 6,7-Dibromo-2,4-dihydroxyquinoline, D-315
- 3,5-Dibromo- β ,4-dimethoxy-N-methylbenzenepropanamine, D-331
- 3,5-Dibromo-4-[3-(dimethylamino)propoxy]cinamic acid, D-323
- 5,6-Dibromo-N,N-dimethyl-1H-indole-3-ethanamine, D-343
- 5,6-Dibromo- N^b,N^b -dimethyltryptamine, D-343
- 4,6-Dibromo-3-ethenyl-1,3-dihydro-3-hydroxy-2H-indol-2-one, C-629
- 3,5-Dibromo-4-ethoxy-1-hydroxy-4-methoxy-2,5-cyclohexadien-1-acetamide, D-316
- 6,8-Dibromo-1-ethyl- β -carboline, B-318
- 4,5-Dibromo-1-ethyl-1H-pyrrole-2-carboxylic acid, D-339
- 3,10-Dibromofascaplysin, F-13
- 4,5-Dibromo-2-furancarboxamide, D-317
- 4,5-Dibromo-2-furancarboxylic acid, D-317
- 4,5-Dibromo-2-furoic acid, D-317
- 3,5-Dibromo-4-hydroxybenzeneacetic acid, D-321
- 3,5-Dibromo-4-hydroxybenzeneacetoneitrile, D-321
- 3,5-Dibromo-4-hydroxybenzeneethanamine, D-322
- 1-(3,5-Dibromo-4-hydroxybenzoyl)-9H-pyrido[3,4-*b*]indole, H-411
- 3,5-Dibromo-4-hydroxybenzyl cyanide, D-321
- 3,5-Dibromo-4-hydroxycinnamic acid, D-323
- 3,5-Dibromo-1-hydroxy-4,4-dimethoxy-2,5-cyclohexadiene-1-acetamide, D-318
- 7,9-Dibromo-10-hydroxy-8-methoxy-6-methyl-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxylic acid, D-319

- 7,9-Dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxylic acid, D-320
- 3,5-Dibromo-2-hydroxy-4-methoxyphenylacetamide, D-314
- 2,3-Dibromo-5-hydroxymethylpyrrole, D-342
- 3,5-Dibromo-4-hydroxyphenethyltrimethylammonium, D-322
- 3,5-Dibromo-4-hydroxyphenylacetamide, D-321
- 1,3-Dibromo-5-[2-[(4-hydroxyphenyl)acetamido]ethyl]-2-[3-(3-methyl-2-butenamido)propoxyl]benzene, D-333
- (3,5-Dibromo-4-hydroxyphenyl)acetic acid, D-321
- 2-(3,5-Dibromo-4-hydroxyphenyl)ethylamine, D-322
- 3-(3,4-Dibromo-5-hydroxyphenyl)-3-hydroxy-2-pyrrolidinone, D-96
- 3-(3,5-Dibromo-4-hydroxyphenyl)-2-propenoic acid, D-323
- 2-(3,5-Dibromo-2-hydroxyphenyl)-3,4,5-tribromopyrrole, D-324
- 6,7-Dibromo-4-hydroxy-2(1*H*)-quinolinone, D-315
- 3,5-Dibromo-4-hydroxy-*N,N,N*-trimethylbenzeneethaniminium, D-322
- 6,6'-Dibromoindigo, D-325
- 6,6'-Dibromoindigotin, D-325
- 6,6'-Dibromoindirubin, I-73
- 4,6-Dibromo-1*H*-indole, D-327
- 3,6-Dibromo-1*H*-indole, D-326
- 5,6-Dibromo-1*H*-indole-3-ethanamine, D-343
- Dibromoisophakellin, D-328
- Dibromomaleimide, D-341
- 3,5-Dibromo-4-methoxybenzeneacetoneitrile, D-321
- 2,4-Dibromo-5-methoxybenzeneethanamine, D-330
- 3,5-Dibromo-4-methoxybenzeneethanamine, D-322
- 3,5-Dibromo-4-methoxybenzyl cyanide, D-321
- N*¹⁰-(3,5-Dibromo-4-methoxycinnamoyl)spermidine, S-398
- 5,7-Dibromo-6-methoxy-1*H*-indole, D-329
- 4,5-Dibromo-*N*²-methoxymethyl-1*H*-pyrrole-2-carboxamide, D-339
- 2,3-Dibromo-5-(methoxymethyl)-1*H*-pyrrole, D-342
- N*-[2-(3,5-Dibromo-4-methoxyphenyl)ethenyl]-3-(4-hydroxyphenyl)-2-methoxy-2-propenamide, B-256
- 2-(2,4-Dibromo-5-methoxyphenyl)ethylamine, D-330
- (3,5-Dibromo-4-methoxyphenyl)ethylamine, D-322
- 3-(3,5-Dibromo-4-methoxyphenyl)-2-methoxy-*N*-methylpropylamine, D-331
- 3,5-Dibromo-4-methoxy-*N,N,N*-trimethylbenzeneethaniminium, D-322
- 4,6-Dibromo-2-methyl-1*H*-indole, D-332
- 5,6-Dibromo-*N*-methyl-1*H*-indole-3-ethanamine, D-343
- Dibromo-*N*¹-methylisophakellin, D-328
- 2,3-Dibromo-*N*-methylmaleimide, D-341
- 4,5-Dibromo-1-methyl-*N*-[3-(1-methyl-2,5-dioxo-4-imidazolidinyl)propyl]-1*H*-pyrrole-2-carboxamide, M-604
- N*-[2-[3,5-Dibromo-4-[3-(3-methyl-1-oxo-2-butenyl)amino]propoxy]phenyl]ethyl]-4-hydroxybenzeneacetamide, D-333
- 4,5-Dibromo-1-methyl-1*H*-pyrrole-2-carboxylic acid, D-339
- 3,4-Dibromo-1-methyl-1*H*-pyrrole-2,5-dione, D-341
- 4,6-Dibromo-2-(methylsulfinyl)-3-(methylthio)-1*H*-indole, D-304
- 4,6-Dibromo-3-(methylsulfinyl)-2-(methylthio)-1*H*-indole, D-304
- 4,6-Dibromo-2-methylthio-1*H*-indole, D-334
- 5,6-Dibromo-*N*⁶-methyltryptamine, D-343
- 5-[3,5-Dibromo-4-[(2-oxo-5-oxazolidinyl)methoxyl]phenyl]-2-oxazolidinone, D-335
- 4,5-Dibromopalauamine, P-18
- (+)-Dibromophakellin, B-343
- (-)-Dibromophakellin, B-343
- Dibromophakellin, D-336
- 2,3-Dibromo-*N*-phenylmaleimide, D-341
- 3,4-Dibromo-1-phenyl-1*H*-pyrrole-2,5-dione, D-341
- 4,5-Dibromopyromucic acid, D-317
- 4,5-Dibromo-1*H*-pyrrole-2-carbonitrile, D-339
- 4,5-Dibromo-1*H*-pyrrole-2-carboxamide, D-339
- 4,5-Dibromo-1*H*-pyrrole-2-carboxylic acid, D-339
- 2,5-Dibromo-1*H*-pyrrole-3-carboxylic acid, D-338
- 2,3-Dibromo-1*H*-pyrrole, D-337
- 4,5-Dibromo-1*H*-pyrrole-2,3-dicarboxylic acid, D-340
- 3,4-Dibromo-1*H*-pyrrole-2,5-dione, D-341
- 4,5-Dibromo-1*H*-pyrrole-2-methanol, D-342
- 6,7-Dibromo-2,4-quinolinediol, D-315
- Dibromosceptiline, S-124
- 2,3-Dibromostyloguanidine, S-611
- 6,7-Dibromo-1,2,3,4-tetrahydro-1-oxopyrrolo[1,2-*a*]pyrazine-4-acetic acid, L-243
- 3,5-Dibromo-*N,N,N*,*O*-tetramethyltyramine, D-322
- 6,21-Dibromotopsentin A, T-432
- 2,4-Dibromo-6-(3,4,5-tribromo-1*H*-pyrrol-2-yl)-phenol, D-324
- 3,5-Dibromo-*N,N,N*-trimethyltyramine, D-322
- 5,6-Dibromotryptamine, D-343
- 3,5-Dibromotyramine, D-322
- 3,5-Dibromoverongiaquinol, H-650
- 3,6-Di-*sec*-butyl-2-hydroxypyrazine, B-204
- N*⁵,*N*¹⁰-Dicafeoyl-*N*¹-*p*-coumaroylspermidine, S-398
- N*⁵,*N*¹⁰-Dicafeoyl-*N*¹-feruloylspermidine, S-398
- N*¹,*N*⁴-Dicafeoylputrescine, D-362
- Dicafeoylspermidine, S-398
- 5,5''-Dicapsaicin, D-344
- Dicarbamoylurea, T-627
- 2,3-Dicarbomethoxy-1-methyl-4(1*H*)-quinolinone, H-728
- 4-[[[(1,2-Dicarboxyethyl)imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, M-646
- 4-[[[(1,3-Dicarboxypropyl)imino]ethylidene]-1,2,3,4-tetrahydro-2,6-pyridinedicarboxylic acid, V-195
- 4,6-Dicarbonyl-4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine-4-propanoic acid, C-802
- Dicarprine A, D-345
- Dicarprine B, D-345
- Dicarprine C, D-345
- Dicentrine methine, M-559
- Dicentrine, D-346
- Dicentrinone, D-347
- Dichloroacetamide, D-348
- Dichloroacetic acid, D-348
- 2,3-Dichloro-4-(3-chloro-5-ethyl-2,6-dihydroxybenzoyl)pyrrole, A-1283
- 2,3-Dichloro-4-(3-chloro-5-hexyl-2,6-dihydroxybenzoyl)pyrrole, C-231
- 2,3-Dichloro-4-(3-chloro-2-nitrophenyl)-1*H*-pyrrole, P-961
- 2,4-Dichloro-6-(5-chloro-3-nitro-1*H*-pyrrol-2-yl)-phenol, P-965
- 2,10-Dichloro-6,11-cyclo-3(15),7(14)-farnesadien-1-yl carbonimidic dichloride, D-349
- 2,3-Dichloro-5-(3,5-dichloro-2-hydroxybenzoyl)pyrrole, P-964
- 2,3-Dichloro-5-(3,5-dichloro-2-hydroxybenzyl)-4-nitropyrrole, P-963
- 2,4-Dichloro-6-[(4,5-dichloro-3-nitro-1*H*-pyrrol-2-yl)methyl]phenol, P-963
- 3,8-Dichloro-11,12-dihydroindolo[2,3-*a*]carbazole, D-350
- 1,11-Dichloro-12,13-dihydro-12-(4-*O*-methyl-β-D-glucopyranosyl)-5*H*-indolo[2,3-*a*]pyrrolo[3,4-*c*]carbazole-5,7(6*H*)-dione, R-32
- 7,8-Dichloro-2,9-dihydro-9-methylpyridid[3,4-*b*]indol-1-one, D-351
- 2,3-Dichloro-5-(2,6-dihydroxybenzoyl)pyrrole, P-834
- 1,2-Dichloro-6,8-dihydroxy-7-propyl-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, S-571
- 2,4-Dichloro-6-(dimethylamino)pyridine, A-729
- 4,6-Dichloro-*N,N*-dimethyl-2-pyridinamine, A-729
- Dichloroethanoic acid, D-348
- 1,2-Dichloro-7-ethyl-6,8-dihydroxy-9*H*-pyrrolo[1,2-*b*][1,3]benzoxazin-9-one, C-403
- 6,14-Dichloro-5-hydroxy-3(14),9-axinysadien-15-yl carbonimidic dichloride, C-400
- 1,2-Dichloro-8-hydroxy-6-methoxy-7-propyl-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, S-571
- 7,8-Dichloro-1-hydroxy-9-methyl-β-carboline, D-351
- 1,2-Dichloro-8-hydroxy-6-methyl-7-propyl-9*H*-pyrrolo[1,2-*b*][1,3]benzoxazin-9-one, D-352
- (3,5-Dichloro-2-hydroxyphenyl)(4,5-dichloro-1*H*-pyrrol-2-yl)methanone, P-964
- 2-(3,5-Dichloro-2-hydroxyphenyl)-3,4,5-trichloropyrrole, D-324
- 6,6'-Dichloroindogotin, D-353
- Dichlorolissoclimide, D-354
- α-(Dichloromethoxymethyl)-β-hydroxy-4-nitrobenzeneopropanamide, A-117
- 7,8-Dichloro-9-methyl-β-carboline, D-355
- 5-(3,3-Dichloro-2-hydroxypropyl)-4-hydroxy-3,3-dimethyl-2-pyrrolidinone, D-966
- 7,8-Dichloro-9-methylpyridid[3,4-*b*]indole, D-355
- 7,8-Dichloro-9-methyl-9*H*-pyridid[3,4-*b*]indol-1-ol, D-351
- 2,3-Dichloro-4-(2-nitrophenyl)-1*H*-pyrrole, P-961
- 2,3-Dichloro-4-nitro-1*H*-pyrrole, D-356
- 1,15-Dichloro-1,14-pentadecadiene-3,12-diyn-8-amine, D-357
- 4,6-Dichloro-2-pyridinamine, A-729
- (4,5-Dichloro-1*H*-pyrrol-2-yl)(2,6-dihydroxyphenyl)methanone, P-834
- 2,3-Dichloro-1-(2,3,4-trichloro-6-hydroxyphenyl)pyrrole, N-136
- 2,4-Dichloro-6-(2,3,4-trichloro-1*H*-pyrrol-5-yl)-phenol, D-324
- Dichloroverongiaquinol, H-650
- Dichomine, D-358
- Dichotamine, D-144
- Dichotine, D-359
- Dichotomide I, A-45
- Dichotomide II, D-360
- Dichotomine A, A-45
- Dichotomine B, D-579
- Dichotomine C, D-579
- Dichotomine D, D-579
- Dichroidine, D-361
- Dichroine A, I-232
- Dichroine B, F-31
- β-Dichroine, F-31
- α-Dichroine, I-232
- N*¹,*N*⁴-Di-*E*-cinnamoyl-2,3-dehydroputrescine, B-423
- N*¹,*N*⁴-Di-*trans*-cinnamoyl-2-hydroxyputrescine, D-248
- N,N'*-Dicinnamoylputrescine, D-362
- N*¹,*N*⁶-Dicinnamoylspermidine, S-398
- N*⁵,*N*¹⁰-Di-*p*-coumaroyl-*N*¹-feruloylspermidine, S-398
- N*¹,*N*⁴-Di-*p*-coumaroylputrescine, D-362
- N*¹,*N*¹⁰-Dicoumaroylspermidine, S-398
- N*¹,*N*⁵-Di-*p*-coumaroylspermidine, S-398
- Dicrotaline, D-363
- Dictamine, F-224
- Dictangustine A, F-211
- Dictizine, D-370
- Dictyocarpine, M-307
- Dictyocarpine, M-307
- Dictyodendrine A, D-364
- Dictyodendrine B, D-365
- Dictyodendrine C, D-366
- Dictyodendrine D, D-366
- Dictyodendrine E, M-247
- Dictyodiamine, P-601
- Dictyolomide A, D-367
- Dictyolomide B, D-367
- Dictyolucidamine, D-368
- Dictyolucidine, D-368
- Dictyophlebine, P-601
- Dictyopterin, A-749
- Dictyoquinazole A, D-369
- Dictyoquinazole B, D-369
- Dictyoquinazole C, D-369
- Dictysine acetone, D-370
- Dictysine, D-370
- Dictyzine, D-370

- 4,5-Dicyano-1,2-benzenediol, D-533
 3,5-Dicyano-2,6-dimethylpyridine, D-776
 2,5-Dicyanopyrrole, P-946
 2',7-Dideacetoxyaustrospicatine, T-49
 7,9-Dideacetylaustrospicatine, T-48
*O*⁶,*O*⁶-Dideacetyllevonine, P-248
 1,6-Dideacetyllevonine, P-248
 6',6''-Didebromo-3,4-dihydrohamacanthin B, H-41
 (*S*)-6',6''-Didebromohamacanthin A, H-40
 (*R*)-6',6''-Didebromohamacanthin B, H-41
 Didebromosceptrine, S-124
 3,5-Didecanoylpyridine, D-371
 6,8-Di-1-decenyl-2,3-dihydro-5,7-dimethyl-1*H*-indolizinium, D-372
 9,11-Didechloro-13-demethylisodysidenin, D-972
 Didechloro-4'-*O*-demethylrebeccamycin, A-1393
 9,10-Didechloro-*N*-methylsidsideathiazole, D-970
 2,5:2',5'-Dide(epidithio)-2,5:2',5'-bis(epitetrathio)-6,6'-dihydroxyacetocin, C-372
 2,5:2',5'-Dide(epidithio)-2,2',5,5'-tetrakis(methylthio)chetomin, D-239
 8,9-Didehydroyanodine, R-175
 19,20-Didehydroaconitane-1,8,14,16-tetrol, A-103
 Didehydroaglaiaastatin, A-197
 19,20-Didehydroaojimalan-17-ol, T-288
 1,2-Didehydro-*N*-β-alanyldopamine, D-157
 11,12-Didehydroaloperine, A-637
 12,13-Didehydroanacycline, A-968
 1,2-Didehydroancistrobertsonine D, A-985
 Didehydroaporphine, D-166
 1,2-Didehydroaspidospermidine, E-3
 1,2-Didehydrobeninine, B-63
 13,14-Didehydro-[14,14'-bimatridine]-15,15'-dione, G-137
 Didehydrobrevicolline, B-292
 12,13-Didehydrocarpine, C-168
 5,6-Didehydro-3,20-cevanediol, C-301
 8,14-Didehydro-3,11-cevanediol, C-306
 8,14-Didehydro-3,15-cevanediol, C-307
 8,9-Didehydro-3,6-cevanediol, C-302
 8,14-Didehydro-1,3-cevanediol, C-304
 8,14-Didehydro-3,6-cevanediol, C-305
 8,9-Didehydro-3,11-cevanediol, C-303
 5,6-Didehydro-3,16,20-cevanetriol, C-309
 8,9-Didehydro-3,6,23-cevanetriol, C-311
 12,13-Didehydro-3,6,15-cevanetriol, C-312
 23,24-Didehydro-3,6,16-cevanetriol, C-297
 8,9-Didehydro-3,6,20-cevanetriol, C-310
 5,6-Didehydrocevan-3-ol, C-313
 14,19-Didehydrocondyfolan, A-1497
 16,17-Didehydrodemethoxyisomigrastatin, I-245
 10,11-Didehydro-4-demethylepothilone D, E-117
 10,11-Didehydro-6-demethylepothilone D, E-117
 16,17-Didehydro-8-*O*-demethylisomigrastatin, I-245
 1',2'-Didehydro-2'-demethylpheanthine, D-148
 13',19'-Didehydro-19'-deoxy-28,44-dihydro-41,44-dihydroxymicrococin P, T-383
 16,17-Didehydro-17-deoxygardneramine oxindole, G-21
 14,15-Didehydro-8-deoxylydicamycin, L-353
 16,17-Didehydro-9-de-2-piperidinylormosanine, A-637
 15,16-Didehydro-14,16-diformylcorynan-17-oic acid, C-688
 7,8-Didehydro-3-dihydrodiscorhabdin C, D-851
 12,13-Didehydro-13,14-dihydro-α-erythroidine, 10CI, E-215
 11,12-Didehydro-10,11-dihydro-20-hydroxyflavinine, A-172
 24,25-Didehydro-10,11-dihydro-20-hydroxyflavinone, H-541
 1,6-Didehydro-12,17-dihydro-3-hydroxy-*C*-homo-16(15*H*)-oxaerythrinan-15-one, 11CI, P-315
 11,12-Didehydro-12,13-dihydro-9-oxoepothilone D, E-117
 1,2-Didehydro-3,11-dihydroxyrincane, H-5
 7,8-Didehydro-4,7-dihydroxy-3,8-dimethoxy-17-methylmorphinan-6-one, C-164
 14,15-Didehydro-10,11-dimethoxy-16-epivincamine, V-115
 1,2-Didehydro-3,15-dimethoxyerythrinan, I-207
 1,6-Didehydro-3,15-dimethoxyerythrinan-2-ol, C-539
 1,6-Didehydro-3,17-dimethoxyerythrinan-15-ol, C-542
 1,6-Didehydro-3,15-dimethoxyerythrinan-16-ol, D-431
 1,6-Didehydro-3,16-dimethoxyerythrinan-15-ol, D-431
 1,6-Didehydro-3,16-dimethoxy-*C*-homoerythrinan-15-ol, T-51
 8,14-Didehydro-3,8-dimethoxymorphinan-4,6,7-triol, S-311
 14,15-Didehydro-10,11-dimethoxyvincamine, V-115
 8,9-Didehydro-6,8-dimethylergoline, A-205
 14,15-Didehydro-10,22-dioxokopsane, D-805
 Didehydrodopamine, D-157
 19,20-Didehydroedulirine A, T-349
 19,20-Didehydroedulirone A, E-41
 6,16-Didehydro-20-episilicine, M-293
 14,15-Didehydroepivincadine, V-112
 10,11-Didehydroepothilone D, E-117
 19,20-Didehydro-17,18-epoxycuran, C-115
 19,20-Didehydro-17,18-epoxycuran-17-ol, C-115
 19,20-Didehydro-17,18-epoxy-11,17-dimethoxycuran, C-115
 6,7-Didehydro-8,10-epoxy-3,7-dimethoxy-17-methylhasubanan-4,8-diol, P-658
 7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6,14-diol, H-462
 7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-ol, C-555
 8,14-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-ol, N-134
 7,8-Didehydro-4,5-epoxy-3-methoxy-17-methylmorphinan-6-one, C-555
 7,8-Didehydro-4,5-epoxy-17-methylmorphinan-3,6-diol, M-704
 6,7-Didehydro-8,10-epoxy-3,4,7-trimethoxy-17-methylhasubanan-8-ol, P-639
 19,20-Didehydroervatamine, E-190
 1,2-Didehydro-7-ethyl-20,21-dinoraspidospermidine, D-377
 16,17-Didehydro-21-ethyl-4-methyl-7,20-cycloatidane-11,15-diol, D-221
 1,2-Didehydrogeissoschizoline, G-34
 4,20-Didehydrogelsedine, G-44
 4,20-Didehydrogelsemicine, G-44
 4,5-Didehydroguadalcine, P-188
 Didehydro-3'-hydroxyaglaiaastatin, A-197
 1,2-Didehydro-12-hydroxyaspidospermidine, E-3
 8,9-Didehydro-20-hydroxy-3-cevanone, H-461
 6*a*,7-Didehydro-7-hydroxy-1,2-dimethoxyaporphine, D-373
 3,4-Didehydro-19-hydroxy-16,17-dimethoxy-1-methylaspidospermidin-21-oic acid, γ-lactone, A-1496
 7,8-Didehydro-4-hydroxy-3,7-dimethoxy-17-methylhasubanan-6-one, C-274
 5,6-Didehydro-4-hydroxy-3,6-dimethoxy-17-methylmorphinan-7-one, I-318
 5,6-Didehydro-2-hydroxy-3,6-dimethoxy-17-methylmorphinan-7-one, P-31
 7,8-Didehydro-4-hydroxy-3,7-dimethoxy-17-methylmorphinan-6-one, S-313
 6*a*,7-Didehydro-7-hydroxy-1,2-dimethoxynoraporphine, D-373
 20,21-Didehydro-17-hydroxy-1,4-dimethyl-4,21-secosarpagan-19(20*H*)-one, M-17
 19,20-Didehydro-6-hydroxyervatamine, E-190
 4,20-Didehydro-14-hydroxygelsedine, G-44
 4,20-Didehydro-14-hydroxygelsemicine, G-44
 16,17-Didehydro-17-hydroxy-16-(hydroxymethyl)-corynan-5-carboxylic acid, D-374
 7,11-Didehydro-13-hydroxymatridin-15-one, L-94
 14,15-Didehydro-10-hydroxy-11-methoxy-16-epivincamine, V-115
 14,15-Didehydro-10-hydroxy-11-methoxyvincamine, V-115
 14,15-Didehydro-12-hydroxy-*N*¹-methylaspidofractinine, A-1494
 19,20-Didehydro-17-hydroxy-4-methyl-1,16-cyclocorynanium, M-130
 9,10-Didehydro-*N*-(2-hydroxy-1-methylethyl)-6-methylergoline-8-carboxamide, E-164
 1,6-Didehydro-12-(hydroxymethyl)-3-methoxy-*c*-homo-*D*-nor-16-oxaerythrinan-15(12*H*)-one, I-263
 1,2-Didehydro-20-hydroxypseudoaspidospermidine, D-377
 1,3*a*-Didehydro-8-hydroxyptilocaulin, D-375
 2,16-Didehydro-23-hydroxyretuline, N-301
 6,7-Didehydro-19-hydroxyschizozogine, S-137
 13,19-Didehydro-12-hydroxysenecionan-11,16-dione, S-240
 7,8-Didehydro-4-hydroxy-3,7,8-trimethoxy-17-methylhasubanan-6-one, H-70
 6,6*a*-Didehydro-11-hydroxy-1,2,10-trimethoxynoraporphine, H-97
 14,15-Didehydroisoburnamine, E-6
 16,17-Didehydroisomigrastatin, I-245
 5,6-Didehydro-α-isoparastine, D-168
 1',2'-Didehydroisostemofoline, S-515
 Didehydrojoubertamine, J-49
 1',2'-Didehydrokohatamine, K-56
 16,17-Didehydrokopsinine, K-71
 Didehydrokrigenamine, K-102
 17,18-Didehydroloesenerine, L-230
 16,17-Didehydroloesenerin-18-ol, L-230
 5,6-Didehydrolopamine, D-161
 8,15-Didehydro-1(18*H*)-lycodinone, H-376
 11,12-Didehydrolycopodane-5,8-diol, L-328
 11,12-Didehydrokamakamin-10-one, H-310
 5,17-Didehydromatridin-15-one, D-80
 5,17-Didehydromatridin-15-one, I-241
 5,17-Didehydromatridin-15-one, L-93
 7,11-Didehydromatridin-15-one, L-94
 12,13-Didehydromatridin-15-one, L-81
 13,14-Didehydromatridin-15-one, S-371
 5,17-Didehydromatrine, L-93
 Didehydro-3'-methoxyaglaiaastatin, A-197
 19,20-Didehydro-9-methoxy-4,17-cyclocorynanium, A-572
 2,16-Didehydro-11-methoxydiaboline, C-115
 19,20-Didehydro-*N*¹-methoxyervatamine, E-190
 1,6-Didehydro-3-methoxyerythrinan-15-ol, C-538
 4,20-Didehydro-11-methoxygelsedine, G-44
 14,15-Didehydro-12-methoxy-*N*¹-methylaspido-fractinine, A-1494
 1,6-Didehydro-3-methoxy-15,16-[methylenebis(oxy)]erythrinan-2-ol, E-208
 19,20-Didehydro-12-methoxy-17-norcuran, T-668
 19,20-Didehydro-12-methoxytubifolidine, T-668
 14,15-Didehydro-12-methoxyvincamine, V-115
ent-14,15-Didehydro-*N*¹-methylaspidospermidine, A-1498
 6,7-Didehydro-1-methylaspidospermidine, A-1498
 14,15-Didehydro-11,12-methylenedioxykopsinaxline, K-74
 9,10-Didehydro-6-methylergoline-8-carboxylic acid, L-370
 19,21-Didehydro-4-methyl-16-methylene-14,20-cyclooctachane-11,12-diol 12-(2-methyl-2-butenolate), A-1070
 3,14-Didehydro-19-methylnormalindine, D-376
 Didehydromirabazole A, M-634
 1,3-Didehydrotriazole, N-246
 4*a*,*N*-Didehydronoraugustamine, A-1539
 1,2-Didehydro-17-norcuran, T-668
 Didehydroocoteine, O-27
 16,17-Didehydroormosanine, P-529
 10,11-Didehydro-9-oxoepothilone D, E-117
 19,20-Didehydro-5-oxoervatamine, E-190
 4,20-Didehydro-19-oxogelsedine, G-44
 14,15-Didehydro-3-oxokopsinine *N*-oxide, V-49
 14,15-Didehydro-3-oxokopsinine, V-49
 14,15-Didehydro-3-oxorhazinilam, R-76
 16,17-Didehydropanamine, O-114
 1,5-Didehydroproline, D-501
 1,2-Didehydropseudoaspidospermidine, D-377
 Didehydroremerine, D-166
 9,21-Didehydroyanodine, R-175
 19,21-Didehydro-6,21-secohetisan-5,15-diol, T-427
 1,13-Didehydro-10,11-secomultiflorine, A-554
 4,4*a*-Didehydroskytanthine, S-326

- 4a,5-Didehydroskytanthine, S-326
 7,8-Didehydrophoramine, S-376
 13,14-Didehydrosporidone, S-371
 11,12-Didehydrosparteine, D-168
 1',2'-Didehydrostemofoline *N*-oxide, S-515
 1',2'-Didehydrostemofoline, S-515
 Didehydrostemonine, S-519
 1-*O*,19-Didehydrotakaosamine, G-152
 3,4-Didehydrotalbotine, T-19
 19,20-Didehydrotalcarpine, T-20
 12,13-Didehydro-2,7,13,14-tetrahydro- α -erythro-
 idine, E-215
 7,8-Didehydro-2,3,7,8-tetramethoxyhasubanan-6-
 one, R-161
 7,8-Didehydro-3,4,7,8-tetramethoxy-17-methylha-
 subanan-6,16-dione, H-70
 7,8-Didehydro-3,4,7,8-tetramethoxy-17-methylha-
 subanan-6-one, H-70
 6'-[(6 α ,7-Didehydro-1,2,10-trimethoxyaporphin-9-
 yl)oxy]codamine, D-170
 1,6-Didehydro-3,15,16-trimethoxyerythran-2-ol,
 E-207
 1,6-Didehydro-3,15,16-trimethoxy-*C*-homoerythri-
 nan-2,7-diol, C-260
 7,8-Didehydro-3,7,8-trimethoxy-17-methylhasuba-
 nan-4,6-diol, H-70
 Didehydrotryptamine, D-172
 Didehydrotriberostemonine, T-664
 1',2'-Didehydrotribulosine, T-675
 14,15-Didehydrovincadine, V-112
 14,15-Didehydrovincamine, V-115
 14,15-Didehydrovincamone, E-6
 14,15-Didehydrovincanol, E-6
 16,17-Didehydrovohimban-17-methanol, N-243
 19,20-Didehydro- α -yohimbine, D-378
 19,20-Didehydro- β -yohimbine, D-378
 19,20-Didehydroyohimbine, D-378
 Didemethoxycarbonyltetrahydrosecamine, S-174
 Dide-*O*-methylaaptamine 9-sulfate, A-2
 Dide-*O*-methylaaptamine, A-2
 N^a,N^b-Didemethylalfileramine, A-260
 5',6-Di-*O*-demethylancistroalaine A, A-980
 Didemethylasterriquinone D, A-1521
 N,*O*-Didemethylcelesticetin, C-239
 1,2-Didemethylcolchicine, D-92
 2,3-Didemethylcolchicine, D-92
 N,N'-Didemethylculantramine, C-803
 O',O'-Didemethylcycleanine, C-835
 2,3-Didemethyl-*N*-deacetylcolchicine, D-92
 1,14-Didemethyldepheline, M-321
 2,3-Didemethyldemecolcine, D-92
 4,4'-Didemethyl-18,18'-dideoxytoxiferine I,
 T-436
 Dide-*O*-methyl-4-*O*- α -D-glucopyranosylsimmondsin,
 S-302
 N,N'-Didemethylgrossularine 1, G-183
 3,4-Dide-*O*-methyl-14 α -hydroxyisotylocrebrine *N*-
 oxide, I-342
 3,6-Dide-*O*-methyl-14 α -hydroxyisotylocrebrine,
 I-342
 N^a,18-Didemethyl-19-hydroxy-N^b-methylsuaveo-
 line, M-10
 13,19-Didemethyl-27-hydroxyspirolide C, S-432
 C,N-Didemethylindolmycin, A-1092
 N^a,N^b-Didemethylisoalfileramine, I-188
 3,6-Dide-*O*-methylisotylocrebrine, I-342
 4',4''-Dide-*O*-methyljulandine, J-54
 O',O''-Didemethylnorthalibrine, T-309
 Dide-*O*-methylsimmondsin, S-302
 13,19-Didemethylspirolide C, S-432
 4'-*N*,*O*-Didemethylstaurosporine, S-498
 Dide-*O*-methylvaracin, L-205
 Didemnimide A, D-379
 Didemnimide B, D-379
 Didemnimide C, D-379
 Didemnimide D, D-379
 Didemnimide E, D-379
 Didemnoline A, D-380
 Didemnoline B, D-380
 Didemnoline C, D-380
 Didemnoline D, D-380
 13,15-Dideoxyaconitine, M-303
 11,17-Dideoxyagelarin A, A-194
 11,17-Dideoxyagelarin B, A-194
 1,4-Dideoxyallonojirimycin, H-615
 1,4-Dideoxyaltronojirimycin, H-615
 10,10'-Dideoxy-4,4'-bincatorone, N-85
 18,18'-Dideoxy-2,2'-epoxytoxiferine I, C-809
 11,19-Dideoxyfistularin 3, F-70
 2,6-Dideoxy-7-*O*- β -D-glucopyranosyl-2,6-imino-D-
glycero-L-*gulo*-heptitol, B-186
 2,5-Dideoxy-2,5-iminoallitol, D-542
 2,5-Dideoxy-2,5-imino-L-altritol, D-542
 1,5-Dideoxy-1,5-imino-D-altritol, H-616
 1,5-Dideoxy-1,5-imino-L-altritol, H-616
 1,4-Dideoxy-1,4-imino-D-arabinitol, D-593
 1,4-Dideoxy-1,4-imino-L-arabinitol, D-593
 1,5-Dideoxy-1,5-imino-D-arabinitol, P-458
 1,5-Dideoxy-1,5-imino-L-arabinitol, P-458
 2,5-Dideoxy-2,5-imino-D-fucitol, D-591
 2,5-Dideoxy-2,5-imino-galactitol, D-542
 1,5-Dideoxy-1,5-imino-D-galactitol, H-616
 1,5-Dideoxy-1,5-imino-L-galactitol, H-616
 2,5-Dideoxy-2,5-imino-D-glucitol, D-542
 1,5-Dideoxy-1,5-imino-D-glucitol, H-616
 1,5-Dideoxy-1,5-imino-L-glucitol, H-616
 1,4-Dideoxy-1,4-imino-5-*O*- β -D-glucopyranosyl-D-
 arabinitol, D-593
 1,5-Dideoxy-1,5-imino-L-gulitol, H-616
 2,6-Dideoxy-2,6-imino-L-gulonic acid, T-579
 2,6-Dideoxy-2,6-iminoheptitol, B-186
 2,5-Dideoxy-2,5-imino-D-*glycero*-D-*talo*-heptitol,
 D-580
 2,5-Dideoxy-2,5-imino-*glycero*-D-*galacto*-heptitol,
 D-580
 2,5-Dideoxy-2,5-imino-*glycero*-D-*manno*-heptitol,
 D-580
 2,5-Dideoxy-2,5-iminoheptitol, D-580
 1,5-Dideoxy-1,5-iminohexitol, H-616
 2,5-Dideoxy-2,5-imino-L-iditol, D-542
 1,5-Dideoxy-1,5-imino-D-iditol, H-616
 1,5-Dideoxy-1,5-imino-L-iditol, H-616
 2,6-Dideoxy-2,6-imino-L-idonic acid, T-579
 1,4-Dideoxy-1,4-imino-D-lyxitol, D-593
 1,4-Dideoxy-1,4-imino-L-lyxitol, D-593
 1,5-Dideoxy-1,5-imino-D-lyxitol, P-458
 1,5-Dideoxy-1,5-imino-L-lyxitol, P-458
 2,5-Dideoxy-2,5-imino-D-mannitol, D-542
 1,5-Dideoxy-1,5-imino-D-mannitol, H-616
 1,5-Dideoxy-1,5-imino-L-mannitol, H-616
 1,4-Dideoxy-1,4-iminopentitol, 11CI, D-593
 1,5-Dideoxy-1,5-iminopentitol, P-458
 1,4-Dideoxy-1,4-imino-D-ribitol, D-593
 1,4-Dideoxy-1,4-imino-L-ribitol, D-593
 1,5-Dideoxy-1,5-iminoribitol, P-458
 1,5-Dideoxy-1,5-imino-D-talitol, H-616
 1,4-Dideoxy-1,4-imino-D-xylitol, D-593
 1,4-Dideoxy-1,4-imino-L-xylitol, D-593
 2,5-Dideoxy-2,5-imino-D-xylitol, D-593
 1,5-Dideoxy-1,5-iminoxylitol, P-458
 Dideoxylyzopeptin C, L-312
 Dideoxymalngamide C, D-225
 1,4-Dideoxymannonojirimycin, H-615
 2',3'-Dideoxy-5-methylcytidine, A-803
 1,5-Dideoxy-1,5-(methylimino)-D-glucitol,
 H-616
 1,6-Dideoxyynojirimycin, M-517
 18,18'-Dideoxytoxiferine I, T-436
 11,11'-Dideoxyverticillin A, V-89
 Didesacetylprotoveratrine A, P-674
 Didmolamide A, D-381
 Didmolamide B, D-382
 Didymeline, D-383
 Dielsine, D-384
 Dielsine, D-385
 Dielsinol, D-385
 Dielsiquinone, M-97
 Dienomycin A, D-386
 Dienomycin B, D-386
 Dienomycin C, D-386
 Dienone II, K-100
 Dienone I, K-100
 7,7a-Diepialexine, T-555
 7,8-Diepialexine, T-555
 1,7a-Diepialexine, T-555
 1,8-Diepialexine, T-555
 3,7a-Diepialexine, T-555
 3,8-Diepialexine, T-555
 2,3-Diepiasturaline, T-555
 6,7-Diepicastanospermine, C-188
 3,4-Diepifagomine, H-614
 3,4-Diepi- α -homonojirimycin, B-186
 2,8-Diepilentiginosine, O-46
 16,17-Diepi pseudoreserpine, R-52
 16,20-Diepisilicine, M-293
 22,25-Diepiscophantine, S-456
 11,13:11,16-Diepoxy-16,17-dihydro-11,12-secoheti-
 san-2-ol, D-387
 14,20:18,20-Diepoxypregnan-3-ol, A-1028
 8,9:8',9'-Diepoxy-6,6',8,8'-tetramethyl-1,1'-bi ergo-
 line, B-176
 Diethanolamine, D-388
 4,4-Diethoxy-1-hydroxy-2,5-cyclohexadiene-1-acet-
 amide, H-650
 3,3-Diethoxy-1-propylamine, A-892
 2,4-Diethoxyquinoline, Q-33
 6,7-Diethoxy-1,2,3,4-tetrahydroisoquinoline,
 T-158
 Diethoxythiobinupharidine, T-380
 Diethyl 1*H*-indole-1,3-dicarboxylate, I-90
 Diethyl 4,4'-(methylenediimino)bisbenzoate,
 D-919
 Diethyl (2-phthalimidoethyl)phosphonate, A-773
 2-Diethylaminoethylamine, E-225
 1-(2-Diethylaminoethyl)-8-hydroxy-4-methoxy- β -
 carboline, P-412
 5,5-Diethylbarbituric acid, B-23
 N,N-Diethylbenzeneacetamide, P-336
 N,N-Diethyl-1,4-butanediamine, B-422
 2,5-Diethyl-3,6-dimethylpyrazine, D-389
 2,6-Diethyl-3,5-dimethylpyrazine, D-390
 N,N'-Diethyl-N,N'-diphenylurea, D-818
 N,N'-Diethyl-1,2-ethanediamine, E-225
 N,N-Diethyl-1,2-ethanediamine, E-225
 N¹,N⁴-Diethylhomospermine, B-150
 8,10-Diethyllobelidol, D-391
 8,10-Diethyllobelionol, D-391
 Diethylmalonylurea, B-23
 α,α' -Diethyl-1-methyl-2,6-piperidinediethanol,
 D-391
 2,3-Diethyl-5-methylpyrazine, D-392
 3,5-Diethyl-2-methylpyrazine, D-393
 8,10-Diethylnorlobelidione, D-391
 8,10-Diethylnorlobelionol, D-391
 6,8-Diethyloctahydro-5-pentylindolizine, E-263
 6,8-Diethyloctahydro-5-propyl-1*H*-indolizine,
 D-394
 6,8-Diethyl-5-pentylindolizidine, E-263
 6,8-Diethyl-5-propylindolizidine, D-394
 5,5-Diethyl-2,4,6(1*H*,3*H*,5*H*)-pyrimidinetrione,
 B-23
 α,α' -Diethyl-1,2,5,6-tetrahydro-1-methyl-2,6-pyri-
 dinediethanol, D-391
 Diethyltoluamide, M-398
 5,8'-Diferulic acid, D-395
 N¹,N⁴-Diferuloylputrescine, D-362
 N¹,N¹⁰-Diferuloylspermidine, S-398
 N⁵,N¹⁰-Diferuloylspermidine, S-398
 Diferuloylspermine, S-399
 Difforine, D-396
 Difforlemenine, D-397
 Difforlemenitine, D-398
 Diffusomycin, O-155
 Diforene, D-730
 2',6'-Di-*O*-galloylprunasin, H-673
 3',6'-Di-*O*-galloylprunasin, H-673
 4',6'-Di-*O*-galloylprunasin, H-673
 Digenic acid, K-4
 Digenin, K-4
 N,N'-Digeranylguanidine, B-174
 Digitine, D-606
 N,N'-Di(D-glucopyranosyl)trichotomine, T-497
 Digonazole, B-61
 Digracine, D-399
 1,4-Diguanidinobutane, D-400
 N-(3,4-Diguanidinobutanoyl)dopa, H-46
 1,5-Diguanidinopentane, D-401
 3,5-Di-5-hexenylpyrrolizidine, D-402
 Dihomogammalinolenic acid, E-52

- Dihomo- γ -linolenic acid, E-52
 Dihomosedinone, D-403
 Dihydroabikoviromycin, A-14
 Dihydroacacipetalin, A-20
 Dihydroaeruginic acid, H-695
 Dihydroagroclavine, F-49
 Dihydroajacanine, A-1533
 19,20-Dihydroakuummicine, A-228
 2 β ,16 β -Dihydroakuummicinol, R-63
 1,2 β -Dihydroakuummiline, A-231
 Dihydroalamarine, A-235
 Dihydroalalamatide, M-280
 14,15-Dihydroallosecurinine-15-ol, A-630
 Dihydroambelline, A-683
 Dihydroampullicine, A-951
 8,14-Dihydroamurine, A-955
 13b,14-Dihydroangustine, A-1019
 18,19-Dihydroangustine, A-1019
 3,14-Dihydroangustine, A-1019
 3,14-Dihydroangustoline, A-1020
 Dihydroantirrhine α -methochloride, A-1328
 Dihydroantirrhine β -*N*-methylsulfate, A-1328
 Dihydroantirrhine, A-1328
 (20*S*)-Dihydroantirrhine, A-1328
 1',8-Dihydroaplysinopsin, A-1357
 Dihydroarcyriacyanin A, A-1392
 Dihydroarcyriarubin B, A-1394
 Dihydroarcyriarubin C, A-1394
 Dihydroarcyroxocin A, A-1398
 Dihydroaspidospermatine, A-1497
 Dihydroatisine, A-1533
 12,13-Dihydroaustamide, A-1556
 Dihydroavicine, D-404
 24,28-Dihydro-15-azasterol, A-1602
 5,6-Dihydroazoverdin G, A-1615
 8,9-Dihydrobaretin, B-24
 2,3-Dihydrobellendine, B-57
 9,10-Dihydro-5*H*,7*H*-benzo[*f*][1,3]dioxolo[6,7]iso-
 quino[8,1,2-*hi*][3,1]benzoxazine, D-946
 3,4-Dihydro-2*H*-1,3-benzoxazin-2-one, D-405
 Dihydroberberine, L-23
 3',4'-Dihydro-3,5'-bi(*N*-methylcytisine), D-704
 7,8-Dihydrobiopterin, A-749
 6,7-Dihydrobis[1,3]benzodioxolo[5,6-*a*:4',5'-*g*]qui-
 nolizinium, C-638
 2,3-Dihydro-6,8-bis[10-(5-hydroxy-6-methyl-2-pi-
 peridinyl)decyl]-1*H*-indolizinium, J-57
 Dihydro-4,5-bis(8-hydroxy-4-quinolinyl)-2(3*H*)-
 furanone, B-358
 7,8-Dihydro-1,1'-bisinomenine, S-313
 2,3-Dihydro-6,8-bis(4-methoxyphenyl)-1*H*-indoli-
 zinium(1+), F-54
 3',4'-Dihydro-3,5'-bithermopsine, D-871
 Dihydrobocconine, D-510
 Dihydrobufofthionine, B-393
 Dihydrobungeanool, T-133
 Dihydrocadambine, D-406
 3 β -Dihydrocadambine, D-406
N-Dihydrocaffeoyltyramine, H-667
 4,5-Dihydrocanthin-6-one, C-90
 2,3-Dihydrocapitavine, C-94
 Dihydrocapsaicin glucoside, C-100
 Dihydrocapsaicin, C-100
 Dihydrocapuvosidine, C-105
 20-Dihydrochaetoglobosin A, C-325
 Dihydrochanoclavine I, C-348
 Dihydrochelerythrine, D-407
 Dihydrochelerythrinyll-8-acetaldehyde, D-408
 Dihydrochelilutine, D-407
 Dihydrochelirubine, D-510
 Dihydrocheminortoxiferine, R-63
 10,11-Dihydrocinchonon-9-ol, C-446
 (9*S*)-10,11-Dihydrocinchonon-9-ol, C-447
 Dihydrocinchonincol, Q-17
 10,11-Dihydrocinchonidine, C-446
 10,11-Dihydrocinchonine, C-447
 18,19-Dihydrocinchophyllamine, C-448
 Dihydrocinnamic acid, P-350
N-Dihydrocinnamoyl-20-hydroxycylindrocaine,
 C-929
 (+)-Dihydrocleavamine, C-522
 (-)-Dihydrocleavamine, C-522
 1,2-Dihydrocomosidine, L-278
 Dihydroconcuressine, C-601
 14,19-Dihydrocondylocarpine, T-673
 Dihydroconessine, C-601
 Dihydroconkuressine, C-601
 4'-Dihydroconsabatine, C-622
 5,6-Dihydroconstrictosine, C-624
 Dihydrocoptisine, D-409
 Dihydrocorymine, D-410
 Dihydrocorynantheine *N*-oxide, C-689
 Dihydrocorynantheine pseudoindoxyl, D-411
 Dihydrocorynantheine, C-689
 Dihydrocorynantheol *N*-metho salt, D-412
 Dihydrocorynantheol *N*-oxide, D-412
 Dihydrocorynantheol, D-412
 Dihydro-*p*-coumaric acid, H-689
 Dihydrocrinine, C-748
 Dihydrocupreine, Q-20
 Dihydrocuscohygrine, D-413
 Dihydrocycloakagerine, D-414
 5,6-Dihydro-7*H*-cyclohepta[*c*]pyridin-7-one, P-157
 Dihydrocycloclimicrophyllidine A, C-883
 Dihydrocycloclimicrophylline A, C-883
 Dihydrocycloclimicrophylline C, C-883
 Dihydrocycloclimicrophylline F, C-883
 Dihydrocycloclimicrophylline I, C-883
 5,12-Dihydrocycloocta[1,2-*b*:5,6-*b'*]diindole-6,13-
 dicarboxylic acid, D-415
 1,3-Dihydrocyclopentapyrimidin-2-one, D-416
 Dihydrodarlingianine, D-74
 5,11-Dihydrodarlingine, D-75
 2,3-Dihydrodarlingine, D-75
 2,3-Dihydrodauriporphine, D-90
 3,14-Dihydrodecussine, D-138
 3-Dihydro-7,8-dehydrodiscorhabdin C, D-851
 16,17-Dihydro-17-demethoxyisorhynchophylline *N*-
 oxide, R-90
 16,17-Dihydro-17-demethoxyisorhynchophylline,
 R-90
 8,9-Dihydro-3'-demethoxyplartine, P-473
 Dihydrodeoxybromotopsentin, T-432
 Dihydrodesoxyceruinine, L-321
 Dihydrodesoxylycoeruinine, L-321
 6'',7''-Dihydro-5,5''-dicapsaicin, D-344
 1,2-Dihydro-1,2-dihydroxyacronycine, D-417
cis-1,2-Dihydro-1,2-dihydroxyacronycine, A-121
trans-1,2-Dihydro-1,2-dihydroxyacronycine, A-121
 3,4-Dihydro-3,4-dihydroxy-1,4-benzoxazin-2-one,
 D-436
 1,2-Dihydro-1,2-dihydroxy-*N*-demethylacronycine,
 D-417
 4,5-Dihydro-1,9-dihydroxy-2,11-dimethoxy-7*H*-di-
 benzo[*de,g*]quinolin-7-one, D-418
 4,5-Dihydro-1,9-dihydroxy-2,11-dimethoxy-7-oxo-
 oaporphine, D-418
 14,19-Dihydro-8,12-dihydroxy-4,19-dimethyl-
 11,15-dioxocrotalananium, C-769
 2,3-Dihydro-5,6-dihydroxy-1,1-dimethylindol-
 ium(1+), D-419
 2,11-Dihydro-5,10-dihydroxy-2,11-dimethyl-2-(4-
 methyl-3-pentenyl)-6*H*-pyrano[3,2-*b*]acridin-6-
 one, G-123
 3,12-Dihydro-6,11-dihydroxy-3,3-dimethyl-7*H*-pyr-
 ano[2,3-*c*]acridin-7-one, A-1527
 3-(6,7-Dihydro-8,9-dihydroxyfarnesyl)indole,
 T-605
 3,4-Dihydro-3,4-dihydroxy-2-(hydroxymethyl)-2*H*-
 pyrrole, D-420
 2,3-Dihydro-5,6-dihydroxy-1*H*-indole-2-carboxylic
 acid, C-862
 3,4-Dihydro-6,7-dihydroxyisoquinoline, D-421
 3,4-Dihydro-6,7-dihydroxy-1(2*H*)-isoquinolinone,
 I-305
 32,33-Dihydro-6,31-dihydroxymanzamine A,
 M-91
cis-1,2-Dihydro-1,2-dihydroxymanzamine, A-121
 3,4-Dihydro-4,5-dihydroxy-3-methoxy-4-(4-meth-
 oxyphenyl)-2(1*H*)-quinolinone, D-422
 7,8-Dihydro-6,10-dihydroxy-7-methoxy-3-methyl-
 benz[*g*]isoquinolin-9(6*H*)-one, P-892
 3,4-Dihydro-3,4-dihydroxy-4-(4-methoxyphenyl)-
 2(1*H*)-quinolinone, D-422
 3,12-Dihydro-6,11-dihydroxy-10-methoxy-3,3,12-
 trimethyl-7*H*-pyrano[2,3-*c*]acridin-7-one, C-472
 2,11-Dihydro-5,10-dihydroxy-12-methoxy-2,2,11-
 trimethyl-6*H*-pyrano[3,2-*b*]acridin-6-one,
 P-879
 1,4-Dihydro-3,7-dihydroxy-8-(3-methyl-2-butenyl)-
 4-oxo-2-quinolinecarboxylic acid, T-584
 14,19-Dihydro-12,13-dihydroxy-14-methylcrotala-
 nan-11,15-dione, C-771
 14,19-Dihydro-12,13-dihydroxy-19-methylcrotala-
 nan-11,15-dione, T-488
 3,4-Dihydro-3,4-dihydroxy-1-methyl-3-(3-methyl-
 2-butenyl)-2(1*H*)-quinolinone, P-440
 15,20-Dihydro-12,14-dihydroxy-4-methyl-4,8-seco-
 senecionan-8,11,16-trione, S-658
 14,19-Dihydro-12,13-dihydroxy-20-norcrotalan-
 an-11,15-dione, M-688
 2,3-Dihydro-3,5-dihydroxy-2-oxo-1*H*-indole-3-
 acetic acid, D-423
 2,3-Dihydro-3,7-dihydroxy-2-oxo-1*H*-indole-3-
 acetic acid, D-424
 1,4-Dihydro-5,8-dihydroxy-4-oxo-2-quinolinecar-
 boxylic acid, T-586
 2,7-Dihydro-2,7-dihydroxypleiocarpamine, A-617
 15,20-Dihydro-12,15-dihydroxysenecionan-11,16-
 dione, B-194
 1,2-Dihydro-2,12-dihydroxysenecionan-11,16-
 dione, R-133
N-(6,7-Dihydro-4,9-dihydroxy-3,10,11-trimethoxy-
 5*H*-dibenzo[*a,c*]cyclohepten-5-yl)acetamide,
 C-567
 9,9*a*-Dihydro-3,9-di-1*H*-indol-3-ylfur-
 o[3',2':3,4]cyclopent[1,2-*b*]indol-2(8*H*)-one,
 M-60
 5,6-Dihydro-9,10-dimethoxybenzo[*g*]-1,3-benzo-
 dioxolo[5,6-*a*]quinolizinium(1+), B-101
 11,12-Dihydro-8,9-dimethoxybenzo[*a*]-1,3-benzo-
 dioxolo[4,5-*g*]quinolizinium, G-181
 12,13-Dihydro-1,2-dimethoxy- α ,12-dimethyl[1,3]-
 benzodioxolo[5,6-*c*]phenanthridine-13-metha-
 nol, H-490
 5,6-Dihydro-1,2-dimethoxy-6,7-dimethyl-4*H*-di-
 benzo[*de,g*]quinoline-3,9-diol, G-147
 1,2-Dihydro-4,10-dimethoxy- α , α -dimethyl-di-
 furo[2,3-*b*:3',2']quinoline-2-methanol, C-418
 3,4-Dihydro-6,8-dimethoxy-1,3-dimethylisoquino-
 line, D-425
 8,13-Dihydro-2,3-dimethoxyindo-
 lo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one,
 E-305
 3,4-Dihydro-6,7-dimethoxyisocarbostyryl, I-305
 3,9-Dihydro-6,8-dimethoxy-2-isopropenyl-9-
 methylfuro[2,3-*b*]quinolin-4(4*H*)-one, P-756
 3,4-Dihydro-6,7-dimethoxyisoquinoline, D-421
 3,4-Dihydro-6,7-dimethoxy-1-isoquinolinol,
 I-305
 3,4-Dihydro-6,7-dimethoxy-1(2*H*)-isoquinolinone,
 I-305
 5,6-Dihydro-9,10-dimethoxy-13-methylbenzo[*g*]-
 1,3-benzodioxolo[5,6-*a*]quinolizinium, D-171
 12,13-Dihydro-1,2-dimethoxy-12-methyl[1,3]ben-
 zodioxolo[5,6-*c*]phenanthridine-13-acetalde-
 hyde, D-408
 12,13-Dihydro-1,2-dimethoxy-12-methyl[1,3]ben-
 zodioxolo[5,6-*c*]phenanthridine, D-407
 12,13-Dihydro-2,3-dimethoxy-12-methyl[1,3]ben-
 zodioxolo[5,6-*c*]phenanthridine, D-482
 12,13-Dihydro-1,2-dimethoxy-12-methyl[1,3]ben-
 zodioxolo[5,6-*c*]phenanthridine-13-methanol,
 B-239
 7-(5,6-Dihydro-1,2-dimethoxy-6-methyl-4*H*-diben-
 zo[*de,g*]quinolin-7-yl)-2-methoxy-1*H*-dibenzo[*de,g*]
 quinolin-1-one, T-628
 10,11-Dihydro-7,8-dimethoxy-13-methyl-10,5-(imi-
 nomethano)-5*H*-benzo[4,5]cyclohepta[1,2-*f*]-1,3-
 benzodioxole, R-36
 8,14-Dihydro-2,3-dimethoxy-14-methylindo-
 lo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one,
 E-309
 3,4-Dihydro-6,7-dimethoxy-1-methylisoquinoline,
 D-627
 3,4-Dihydro-6,7-dimethoxy-2-methylisoquinoli-
 nium(1+), D-421
 3,4-Dihydro-6,7-dimethoxy-1-methyl-8-isoquinoli-
 nol, D-439

- 3,4-Dihydro-6,7-dimethoxy-2-methyl-1(2*H*)-isoquinolinone, I-305
- 3,9-Dihydro-6,8-dimethoxy-9-methyl-2-(1-methylethyl)furo[2,3-*b*]quinolin-4(2*H*)-one, P-756
- 2,3-Dihydro-4,8-dimethoxy-9-methyl-2-(1-methylethyl)furo[2,3-*b*]quinolinium, B-14
- 3,4-Dihydro-5-(4,5-dimethoxy-2-methyl-1-naphthalenyl)-8-hydroxy-6-methoxy-1,3-dimethylisoquinoline, D-444
- 3,4-Dihydro-5-(4,5-dimethoxy-2-methyl-1-naphthalenyl)-6-methoxy-1,3-dimethyl-8(2*H*)-isoquinolinone, D-444
- 12,13-Dihydro-1,2-dimethoxy-12-methyl-13-(2-oxopropyl)-1,3-benzodioxolo[5,6-*c*]phenanthridine, A-34
- 1,2-Dihydro-3,4-dimethoxy-1-methylpyridine, A-1400
- 6-(1,3-Dihydro-6,7-dimethoxy-3-oxo-1-isobenzofuranyl)-7,8-dihydro-1,3-dioxolo[4,5-*g*]isoquinolin-5(6*H*)-one, B-99
- 2,3-Dihydro-8,9-dimethoxy-1*H*-pyrrolo[2,1-*a*]isoquinolinium, C-756
- 4,5-Dihydro-9,10-dimethoxy-7*H*-pyrrolo[3,2,1-*de*]phenanthridine, A-1508
- 4,5-Dihydro-9,10-dimethoxypyrrrolo[3,2,1-*de*]phenanthridinium(1+), V-35
- 3,5-Dihydro-6,7-dimethoxy-2,3,3-trimethylfuro[3,2-*c*]quinolin-4(2*H*)-one, O-82
- 6,8-Dihydro-8,8-dimethyl-5*H*-benzo[*g*]1,3-benzodioxolo[6,5,4-*de*]quinolin-10-ol, G-188
- 5,6-Dihydro-3,10-di-*O*-methylconstrictosine, C-624
- 6,7-Dihydro-2,3-dimethyl-5*H*-cyclopentapyrazine, D-426
- Dihydro-4,6-dimethyl-4*H*-1,3,5-dithiazine, D-427
- 4,5-Dihydro-2,4(5)-dimethyl-1*H*-imidazole, D-428
- 1,4-Dihydro-4,6-dimethyl-9*H*-imidazo[1,2-*a*]purin-9-one, W-30
- 2,9-Dihydro-2,2-dimethyl-7-(3-methylbutyl)pyrano[3,2-*g*]indole, A-1510
- 3,7-Dihydro-*N*,3-dimethyl-2-(methylimino)-2*H*-purin-6-amine, I-48
- 3,11-Dihydro-3,8-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-*a*]carbazole-7,10-dione, P-888
- 3,7-Dihydro-3,10-dimethyl-3-(4-methyl-3-pentenyl)pyrano[2,3-*c*]carbazole, E-301
- 3,11-Dihydro-3,8-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-*a*]carbazole, M-46
- 3,11-Dihydro-3,5-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-*a*]carbazol-9-ol, M-47
- 4,9-Dihydro-4,6-dimethyl-9-oxo-1*H*-imidazo[1,2-*a*]purine, W-30
- 3,7-Dihydro-1,3-dimethyl-1*H*-purine-2,6-dione, T-360
- 3,7-Dihydro-3,7-dimethyl-1*H*-purine-2,6-dione, T-355
- 3,7-Dihydro-1,7-dimethyl-1*H*-purine-2,6-dione, X-5
- 1,7-Dihydro-1,7-dimethyl-6*H*-purin-6-one, P-820
- 1,9-Dihydro-1,9-dimethyl-6*H*-purin-6-one, P-820
- 3,7-Dihydro-3,7-dimethyl-6*H*-purin-6-one, P-820
- 3,9-Dihydro-3,9-dimethyl-6*H*-purin-6-one, P-820
- 3,11-Dihydro-3,3-dimethylpyrano[3,2-*a*]carbazole-5-carboxaldehyde, M-757
- 2,6-Dihydro-2,2-dimethyl-5*H*-pyrano[3,2-*c*]quinolin-5-one, F-90
- 2,8-Dihydro-6,8-dimethylpyrimido[5,4-*e*]-1,2,4-triazine-3,5,7(6*H*)-trione, D-498
- 4,5-Dihydro-2,3-dimethylthiazolium(1+), D-479
- 7,8-Dihydro-3,4-di-*O*-methyltubastrine, T-656
- Dihydrodioscorine, D-798
- 4-(3,4-Dihydro-1,1-dioxido-2*H*-1,4-thiazin-6-yl)-phenol, D-457
- 6,12-Dihydro-6,12-dioxindolo[2,1-*b*]quinazoline, I-108
- 5,15-Dihydro[1,3]dioxolo[4,5-*g*]indolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-8(6*H*)-one, E-306
- 7,8-Dihydro-1,3-dioxolo[4,5-*g*]isoquinolin-5(6*H*)-one, I-305
- 4,5-Dihydro-4,5-dioxo-1*H*-pyrrolo[2,3-*f*]quinoline-2,7,9-tricarboxylic acid, M-236
- 5,6-Dihydro-3,6-dioxo-1,2,4-triazine-4(3*H*)-carboxylic acid, D-429
- 4,5-Dihydro-2,5-diphenyloxazole, D-430
- 3-Dihydrodiscorhabdin C, D-851
- 7,8-Dihydro-1,1'-disinomenine, S-313
- Dihydroburnamenine, E-5
- Dihydroelaeocarpidine, T-34
- 15,16-Dihydroelaeocarpine, E-55
- 13,16-Dihydroelaeocarpine, P-714
- 3,4-Dihydroellipticine, E-67
- Dihydroelymoclavine, F-49
- 1,2-Dihydro-3-epimacronine, P-626
- Dihydroepistephamiacrine 6-acetate, L-244
- Dihydroergocornine, E-153
- Dihydroergocristine mesilate, E-155
- Dihydroergocristine, E-155
- Dihydroergosine, E-169
- Dihydroergotamine mesylate, E-172
- Dihydroergotamine, E-172
- 19,20-Dihydroervahanine A, E-189
- Dihydroerysodine, D-431
- Dihydroerysotrine, D-431
- Dihydroerysovine, D-431
- Dihydroerythraline, E-208
- 3,4-Dihydroeudistomin U, I-119
- Dihydroevocarpine, H-750
- N*-Dihydroferuloyltyramine, H-667
- Dihydroferupurine, C-710
- 6,7-Dihydroflavopereirine, F-84
- Dihydroflustramine C *N*-oxide, F-106
- Dihydroflustramine C, F-106
- 7,8-Dihydrofolic acid, P-772
- 2,3-Dihydro-1-formyl-2-hydroxy-5-methoxy-6-methyl-1*H*-indole, D-432
- 2,3-Dihydro-1-formyl-5-methoxy-6-methyl-1*H*-indole, D-432
- Dihydrofumariline 1, F-176
- Dihydrofumariline 2, F-176
- Dihydrofumariline, F-176
- 4-Dihydrofuntumafurine C, A-891
- 6,7-Dihydro-4*H*-furo[2,3-*c*]azepine-4,8(5-*H*)-dione, D-433
- Dihydrogadesine, M-321
- Dihydrogagaminine, P-602
- Dihydrogambirtannine, D-434
- Dihydrogephyrotoxin, G-65
- Dihydroglyparvin A, T-478
- Dihydrohaemanthamine, H-5
- Dihydrohalichondramide, H-15
- cis*-3,4-Dihydrohamacanthin A, H-40
- trans*-3,4-Dihydrohamacanthin A, H-40
- cis*-3,4-Dihydrohamacanthin B, H-41
- Dihydrohaplamine, F-90
- Dihydrohaplocine (incorr.), L-177
- Dihydroharman, M-426
- Dihydroharmine, H-592
- Dihydroheliotridine, H-233
- 3,4-Dihydro-5-(5-hexenyl)-2-nonyl-2*H*-pyrrole, D-435
- Dihydrohistrionicotoin, H-309
- Dihydroholaphyllamine, A-882
- 6,7-Dihydrohomoerysotrine, L-278
- 2,7-Dihydrohomoerysotrine, T-51
- 2,7-Dihydrohomoerysotrine, T-51
- 6,7-Dihydrohomoerythraline, C-587
- 2,7-Dihydrohomo- β -erythroidine, P-315
- Dihydro-*N*-hydroxyabikoviromycin, A-14
- 10,11-Dihydro-24-hydroxyflavinine, A-172
- 19,20-Dihydro-19-hydroxyakuammicine, E-30
- 16,22-Dihydro-16-hydroxyapparine, A-1365
- 8,9-Dihydro-11-hydroxyascididemin, A-1459
- 12,13-Dihydro-12-hydroxyaustamide, A-1556
- 10,11-Dihydro-8-hydroxy-9*H*-benzo[*b*]pyrido[4,3,2-*de*][1,10]phenanthroline-9-one, L-2
- 4,5-Dihydro-2-(6-hydroxy-2-benzothiazolyl)-4-thiazolecarboxylic acid, L-282
- 3,4-Dihydro-3-hydroxy-1,4-benzoxazin-2-one, D-436
- 7,8-Dihydro-8-hydroxyberberine, O-216
- 1,2-Dihydro-9-hydroxy-6-canthinone, H-444
- 2,3-Dihydro-4'-hydroxycapitavine, C-94
- 3,4-Dihydro-7-hydroxy- β -carboline-1-carboxylic acid, H-459
- 16,17-Dihydro-17-hydroxycathenamine, C-207
- 1,3-Dihydro-3-[(4-hydroxy-3-cinnolinyl)methylene]-2*H*-indol-2-one, S-132
- 18,19-Dihydro-9-hydroxycorynantheine, G-13
- 5,6-Dihydro-4-hydroxycyclopenta[*c*]pyrrolo-1,3(2*H*,4*H*)-dione, M-70
- 1,2-Dihydro-2-hydroxy-*N*-demethyl-1-oxonoracrocynine, A-120
- 2,3-Dihydro-1-(4-hydroxy-3,5-dimethoxycinnamoyl)-4*H*-pyridin-4-one, C-245
- 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinolinium inner salt, D-439
- 3,4-Dihydro-8-hydroxy-6,7-dimethoxyisoquinoline, D-437
- 1,3-Dihydro-3-hydroxy-4,9-dimethoxy-3-methyl-2*H*-benz[*f*]indol-2-one, D-438
- 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-1-methylisoquinoline, D-439
- 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-2-methylisoquinolinium inner salt, D-437
- 3,4-Dihydro-1-[(2-hydroxy-5,7-dimethoxyphenyl)methylene]-6,7-dimethoxy-2(1*H*)-isoquinoline-carboxaldehyde, P-554
- 4,5-Dihydro-10-hydroxy-2,9-dimethoxypyrrrolo[3,2,1-*de*]phenanthridinium(1+), V-35
- 2,3-Dihydro-6-hydroxy- α,α -dimethylfuro[2,3-*b*]quinoline-2-methanol, R-97
- 3,12-Dihydro-6-hydroxy-3,3-dimethyl-5-(3-methyl-2-butenyl)-7*H*-pyrano[2,3-*c*]jacridin-7-one, S-278
- 4-(2,3-Dihydro-2-hydroxy-5,7-dimethyl-3-oxo-2-benzofuranyl)-2,6-piperidinedione, A-124
- 1,4-Dihydro-3-hydroxy-1,4-dioxo-2-naphthalene-carboxylic acid, H-638
- 16,17-Dihydro-17-hydroxy-19-epicathenamine, A-215
- 14,19-Dihydro-12-(1-hydroxyethyl)-17,20-dinorcro-talanan-11,15-dione, C-767
- 2,3-Dihydro-2-(1-hydroxyethyl)-2-methylquinazolin-4(1*H*)-one, D-440
- 2,3-Dihydro-3-(1-hydroxyethyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, M-19
- 3',4'-Dihydro-4'-hydroxyflindersine, F-90
- 18,19-Dihydro-19-hydroxygelsemine, G-49
- 3,4-Dihydro-3-hydroxygraciline, G-151
- 3,4-Dihydro-7-hydroxy-1-(4-hydroxybenzoyl)-6-methoxyisoquinoline, L-252
- 3,4-Dihydro-6-hydroxy-4-(4-hydroxy-3,5-dimethoxyphenyl)-3-hydroxymethyl-5,7-dimethoxy-2-naphthalenecarboxylic acid, T-395
- 2,3-Dihydro-4-hydroxy-2-(2-hydroxyethyl)-1*H*-isoindol-1-one, D-442
- 3,4-Dihydro-7-hydroxy-6-(5-hydroxy-1*H*-indol-3-yl)-1*H*-azepino[5,4,3-*cd*]indole-4-carboxylic acid, H-778
- 4,8-Dihydro-2-hydroxy-7-(hydroxymethyl)-8,8-dimethyl-2*H*-1,4-benzothiazine-3,5-dione, X-3
- 2,3-Dihydro-8-hydroxy-2-(1-hydroxy-1-methylethyl)-4-methoxy-9-methylfuro[2,3-*b*]quinolinium, B-14
- 2,3-Dihydro-6-hydroxy-2-(1-hydroxy-1-methylethyl)-4-methoxy-9-methylfuro[2,3-*b*]quinolinium(1+), R-93
- 3,9-Dihydro-6-hydroxy-2-(1-hydroxy-1-methylethyl)-9-methylfuro[2,3-*b*]quinolin-4(2*H*)-one, R-91
- 2,3-Dihydro-7-hydroxy-2-(4-hydroxyphenyl)-4*H*-1-benzopyran-4-one, D-581
- 3,6-Dihydro-6-hydroxy-6-(hydroxyphenylmethyl)-3,5-dimethoxy-3-(1-methylethyl)-2(1*H*)-pyrazinone, T-85
- 3,4-Dihydro-4-hydroxy-5-(3-hydroxy-2-pyridinyl)-4-methyl-2*H*-pyrrole-2-carboxamide, S-295
- 2,3-Dihydro-5-hydroxy-1*H*-indole, D-465
- 1,3-Dihydro-3-hydroxy-2*H*-indole-2-thione, H-534
- 1,3-Dihydro-3-hydroxy-2*H*-indol-2-one, I-91
- 1,3-Dihydro-5-hydroxy-2*H*-indol-2-one, I-92
- 3-(2,3-Dihydro-3-hydroxy-1*H*-indol-3-yl)-2-oxopropanoic acid, D-441
- 3-[1,2-Dihydro-5-(5-hydroxy-1*H*-indol-3-yl)-2-oxo-3*H*-pyrrol-3-ylidene]-1,3-dihydro-2*H*-indol-2-one, V-142
- 2,3-Dihydro-4-hydroxy-1*H*-isoindol-1-one, D-442
- 16,17-Dihydro-17-hydroxyisomitrphylline, M-661

- 3,4-Dihydro-1-hydroxyisoquinoline, D-469
 3,4-Dihydro-2-hydroxy-1(2*H*)-isoquinolinone, D-469
 19,20-Dihydro-10-hydroxyisostiriricine, I-319
 12,13-Dihydro-12-hydroxyisostyrichine, P-672
 5 α ,6-Dihydro-1 α -hydroxyjervine, J-36
 5 β ,6-Dihydro-1 β -hydroxyjervine, J-36
 8,9-Dihydro-9-hydroxylactimidomycin, I-245
 8,9-Dihydro-8-hydroxylactimidomycin, L-7
 3,4-Dihydro-6-hydroxymanzamine A, M-91
 32,33-Dihydro-31-hydroxymanzamine A, M-91
 11,12-Dihydro-9-hydroxy-8-methoxybenzo[*a*]-1,3-benzodioxolol[4,5-*g*]quinolinizium, G-181
 1,2-Dihydro-1-(α -hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline, M-242
 1,9-Dihydro-1-hydroxy-3-methoxy-1,2-dimethyl-4*H*-carbazol-4-one, C-128
 4-(3,4-Dihydro-8-hydroxy-6-methoxy-1,3-dimethylisoquinolin-5-yl)-8-methoxy-3-methyl-1-naphthalenol, D-444
 8,13-Dihydro-2-hydroxy-3-methoxyindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, E-308
 3,4-Dihydro-7-hydroxy-6-methoxy-1(2*H*)-isoquinolinone, I-305
 3,4-Dihydro-4-hydroxy-3-methoxy-4-(4-methoxyphenyl)-2(1*H*)-quinolinone, D-422
 2,3-Dihydro-2-hydroxy-5-methoxy-6-methyl-1*H*-indole-1-carboxaldehyde, D-432
 1,3-Dihydro-3-hydroxy-4-methoxy-3-methyl-2*H*-indol-2-one, D-443
 2,3-Dihydro-6-hydroxy-4-methoxy-5-methyl-1*H*-isoindol-1-one, C-437
 3,4-Dihydro-6-hydroxy-7-methoxy-2-methylisoquinolinium(1+), D-421
 3,4-Dihydro-7-hydroxy-6-methoxy-2-methylisoquinolinium(1+), D-421
 3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-6,8-dimethoxy-1,3-dimethylisoquinoline, D-444
 3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-8-hydroxy-6-methoxy-1,3-dimethylisoquinoline, D-444
 3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-6-methoxy-1,3-dimethyl-8(2*H*)-isoquinolinone, D-444
 1,2-Dihydro-4-hydroxy-3-methoxy-1-methylquinoline, E-13
 1,5-Dihydro-4-[(4-hydroxy-3-methoxyphenyl)methyl]-3-(4-hydroxyphenyl)-2*H*-pyrrol-2-one, C-141
 1-[2,3-Dihydro-2-(4-hydroxy-3-methoxyphenyl)-3-(2-pyrrolidinyl)-1,4-benzodioxin-6-yl]-2-(1-pyrrolidinyl)ethanone, H-770
 2,11-Dihydro-5-hydroxy-10-methoxy-2,2,11-trimethyl-6*H*-pyrano[3,2-*b*]acridin-6-one, J-59
 3*a*,4-Dihydro-2-hydroxymethyl-1,3-benzodioxole-5-carboxamide, D-807
 2,3-Dihydro-3-(3-hydroxy-2-methyl-1-butanyl)-3-methyl-7-phenylfuro[3,2-*c*]pyridine-2,4-diol, A-1146
 7,8-Dihydro-2-hydroxy-*N*⁶-(3-methylbutyl)adenine, A-818
 7,8-Dihydro-2-hydroxy-6-(3-methylbutylamino)purine, A-818
 3,4-Dihydro-7-hydroxy-1-methyl- β -carboline-3-carboxylic acid, T-206
 3,4-Dihydro-7-hydroxy-1-methyl- β -carboline, H-592
 4,8-Dihydro-7-(hydroxymethyl)-8,8-dimethyl-2*H*-1,4-benzothiazine-3,5-dione, X-3
 7,8-Dihydro-4-hydroxy-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolinium(1+), C-707
 3,4-Dihydro-3-hydroxy-1-methyl-4,5-diphenyl-2(1*H*)-pyridinone, D-445
 2,3-Dihydro-2-(1-hydroxy-1-methylethyl)-4,8-dimethoxy-9-methylfuro[2,3-*b*]quinolinium, B-14
 3,9-Dihydro-2-(1-hydroxy-1-methylethyl)-8-methoxy-9-methylfuro[2,3-*b*]quinolin-4(2*H*)-one, B-13
 7,10-Dihydro-8-(1-hydroxy-1-methylethyl)-10-methyl-1,3-dioxolo[4,5-*h*]furo[2,3-*b*]quinolin-6(8*H*)-one, H-549
 3,5-Dihydro-2-(1-hydroxy-1-methylethyl)-5-methylfuro[3,2-*c*]quinolin-4(2*H*)-one, A-1377
 3,9-Dihydro-2-(1-hydroxy-1-methylethyl)-9-methylfuro[2,3-*b*]quinolin-4(2*H*)-one, I-268
 3,9-Dihydro-2-(1-hydroxy-1-methylethyl)-9-methyl-7,8-methylenedioxyfuro[2,3-*b*]quinolin-4(2*H*)-one, H-549
 7,8-Dihydro-8-hydroxymethylfagaridine, B-239
 6,7-Dihydro-4-(hydroxymethyl)-2-(*p*-hydroxyphenethyl)-7-methyl-5*H*-2-pyrindinium(1+), D-446
 6,7-Dihydro-3-hydroxymethyl-8(5*H*)-indolizone, D-447
 3,4-Dihydro-5-hydroxy-2-methyl-1(2*H*)-isoquinolinone, I-304
 5,6-Dihydro-3-hydroxymethyl-9-methoxy-6-methyl-1,2-methylenedioxy-4*H*-dibenzo[*de,g*]quinoline, C-2
 3,4-Dihydro-8-hydroxy-2-methyl-6,7-methylenedioxyisoquinolinium(1+), C-707
 1,11-Dihydro-5-hydroxy-11-methyl-2-(1-methylethyl)furo[2,3-*c*]acridin-6(2*H*)-one, R-167
 [3,4-Dihydro-5-(hydroxymethyl)-4-methyl-3-oxopyrazinyl]guanidine, S-553
 1,5-Dihydro-5-hydroxy-3-methyl-1-(4-methyl-5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, H-467
 1-[[1,5-Dihydro-4-(hydroxymethyl)-1-methyl-5-oxo-2*H*-pyrrol-2-ylidene]acetyl]-1,5-dihydro-4-methoxy-2*H*-pyrrol-2-one, P-783
 2,3-Dihydro-4-hydroxy-3-methyl-2-(1-methyl-1-propenyl)-7-phenylfuro[3,2-*c*]pyridine-3-methanol, A-1142
 10-(20,21-Dihydro-21-hydroxy-21-methyl-18-norallstophyllan-19-yl)-11-methoxyalstophyllan-19-one, M-8
 1,5-Dihydro-5-hydroxy-5-(6-methyl-2,4-octadecyl)-2*H*-pyrrol-2-one, A-1573
 2,5-Dihydro-4-hydroxy-1-methyl-3-(1-oxo-2,4,6-octatrienyl)-5-(phenylmethyl)-1*H*-pyrrol-2-one, B-85
 1,2-Dihydro-4-hydroxy-1-methyl-2-oxo-3-pyridinecarbonitrile, D-668
 1,5-Dihydro-5-hydroxy-3-methyl-1-(5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, D-448
 5,6-Dihydro-5-hydroxy-3-methyl-6-phenyl-3-benzazocin-4(3*H*)-one, C-501
 2,3-Dihydro-4-hydroxy-2-(1-methyl-1-propenyl)-5-phenylfuro[2,3-*b*]pyridine-3,3-dimethanol, A-1145
 3,6-Dihydro-10-hydroxy-9-methyl-4*H*-pyrido[4,3-*f*][1,4]oxazocin-4-one, P-159
 6,7-Dihydro-7-hydroxy-7-methyl-5*H*-2-pyridine-4-carboxaldehyde, E-296
 6,7-Dihydro-7-(hydroxymethyl)-5*H*-2-pyridin-5-one, C-562
 3,4-Dihydro-2-(hydroxymethyl)-2*H*-pyrrole-3,4-diol, D-420
 2,3-Dihydro-7-(hydroxymethyl)-1*H*-pyrrolizin-1-one, L-263
 1,5-Dihydro-5-hydroxy-3-methyl-2*H*-pyrrol-2-one, D-449
 7,8-Dihydro-8'-hydroxy-6-methylspiro[1,3-dioxolo[4,5-*g*]isoquinoline-5(6*H*),7'-[7*H*]indeno[4,5-*d*][1,3]dioxol]-6'(8'*H*)-one, C-673
 2,3-Dihydro-3-hydroxy-3-methylthiazolo[3,2-*a*]indole-9-carboxaldehyde, B-282
 5*a*,9*a*-Dihydro-5*a*-hydroxymillaurine, M-613
 16,17-Dihydro-17-hydroxymitraphylline, M-661
 3,7-Dihydro-7-[(4-hydroxy-3-nitrophenyl)methyl]-1,3-dimethyl-1*H*-purine-2,6-dione, P-357
 14,19-Dihydro-13-hydroxy-20-norcrotalanan-11,15-dione, F-168
N-[5-[(1,6-Dihydro-2-hydroxy-5-octanamido-6-oxo-3-pyridyl)imino]-1,2,5,6-tetrahydro-2,6-dioxo-3-pyridyl]octanamide, D-450
 1*a*,2-Dihydro-1*a*-hydroxy-6*H*-oxireno[*b*]indole-6*b*-acetic acid, M-758
 2,5-Dihydro-4-hydroxy-3-(1-oxo-2,4,6,8,10-dodecapentaenyl)-5-(phenylmethyl)-1*H*-pyrrol-2-one, B-87
 2,3-Dihydro-3-hydroxy-2-oxo-1*H*-indole-3-acetic acid, D-451
 2,3-Dihydro-5-hydroxy-2-oxo-1*H*-indole-3-acetic acid, D-453
 2,3-Dihydro-4-hydroxy-2-oxo-1*H*-indole-3-acetic acid, D-452
 2,3-Dihydro-3-hydroxy-2-oxo-1*H*-indole-3-acetonitrile, C-829
 2,3-Dihydro-4-hydroxy-2-oxo-1*H*-indole-3-acetonitrile, D-452
 2,3-Dihydro-3-hydroxy- α -oxo-1*H*-indole-3-propionic acid, D-441
N-[2-(2,3-Dihydro-3-hydroxy-2-oxo-1*H*-indol-3-yl)ethyl]-3-phenyl-2-propenamide, C-454
 32,33-Dihydro-6-hydroxy-35-oxomanzamine A, M-91
 2,5-Dihydro-4-hydroxy-3-(1-oxo-2,4,6-octatrienyl)-5-(phenylmethyl)-1*H*-pyrrol-2-one, B-85
 2,5-Dihydro-3-hydroxy-5-oxo-4-(1-oxo-2,4,6-octatrienyl)-1*H*-pyrrole-2-acetic acid, A-1204
 1,2-Dihydro-4-hydroxy-2-oxo-3-pyridinecarboxylic acid, D-668
 1,4-Dihydro-6-hydroxy-4-oxo-2-quinolinecarboxylic acid, D-675
 1,2-Dihydro-6-hydroxy-2-oxo-4-quinolinecarboxylic acid, D-672
 1,2-Dihydro-8-hydroxy-2-oxo-4-quinolinecarboxylic acid, D-673
 1,4-Dihydro-3-hydroxy-4-oxo-2-quinolinecarboxylic acid, D-674
 1,4-Dihydro-7-hydroxy-4-oxo-2-quinolinecarboxylic acid, D-676
 1,4-Dihydro-8-hydroxy-4-oxo-2-quinolinecarboxylic acid, D-677
 7,8-Dihydro-8-hydroxypalmitate, O-213
 20*R*,21*S*-Dihydro-10-hydroxyperaksine, D-491
 3,4-Dihydro-6-(4-hydroxyphenyl)-1,1-dioxo-2*H*-1,4-thiazine, D-457
 3,4-Dihydro-2-(4-hydroxyphenyl)-7-methoxy-8-(2-pyrrolidinyl)-2*H*-1-benzopyran-5-ol, V-189
 3,4-Dihydro-3-[(4-hydroxyphenyl)methylene]cyclopent[*b*]indole-1,2-dione, N-323
 2,3-Dihydro-6-(4-hydroxyphenyl)-7-methyl-1*H*-indolizinium, I-170
 4,5-Dihydro-2-(2-hydroxyphenyl)-4-methyl-4-thiazolecarboxylic acid, D-454
 4,5-Dihydro-2-(2-hydroxyphenyl)-4-oxazolecarboxylic acid, D-455
 5,6-Dihydro-3-[2-(4-hydroxyphenyl)-2-oxoethyl]-2(1*H*)-pyridinone, D-456
 3,4-Dihydro-6-(4-hydroxyphenyl)-2*H*-1,4-thiazine, D-457
 4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolemethanol, H-695
 2-[4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolyl]-3-methyl-4-thiazolidinecarboxylic acid, P-832
 3-[4,5-Dihydro-2-(2-hydroxyphenyl)-4-thiazolyl]tetrahydro-6,9-imino-1*H*,3*H*,5*H*-thiazolo[4,3-*c*][1,4]thiazepin-5-one, U-6
 6,7-Dihydro-4-(1-hydroxypropyl)-5*H*-2-pyridin-5-one, L-272
 5,6-Dihydro-5-hydroxy-2(1*H*)-pyridinone, D-458
 4,5-Dihydro-2-(3-hydroxy-2-pyridinyl)-4-methyl-4-thiazolecarboxylic acid, D-235
 2,3-Dihydro-1-hydroxy-1*H*-pyrrolizine-7-carboxaldehyde, D-14
 1,3-Dihydro-3-hydroxy-2*H*-pyrrol-2-one, D-459
 1,5-Dihydro-5-hydroxy-2*H*-pyrrol-2-one, D-460
 1,5-Dihydro-4-hydroxy-2*H*-pyrrol-2-one, P-954
 2,3-Dihydro-5-hydroxypyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, D-503
 2,3-Dihydro-3-hydroxypyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, V-38
 3,4-Dihydro-6-hydroxy-2(1*H*)-quinolinone, D-461
 14,15-Dihydro-14-hydroxysecuranin-11-one, S-208
 15,20-Dihydro-12-hydroxysecuranin-11,16-dione, B-194
 1,2-Dihydro-12-hydroxysecuranin-11,16-dione, P-516
 5,6-Dihydro-1-hydroxy-2,3,10,11-tetramethoxydibenzo[*a,g*]quinolinizium(1+), S-531
 1,13-Dihydro-13-hydroxyuleine, U-8
 Dihydro-19-hydroxyvincamajimine, V-113
 7,8-Dihydroimidazo[1,5-*c*]pyrimidin-5(6*H*)-one, D-462

- 2,3-Dihydro-2-imino-1,3-dimethyl-1*H*-imidazole-4-acetic acid, A-928
 Dihydroimpranine, I-54
 8,9-Dihydroindanomycin, I-64
 6,7-Dihydro-8(5*H*)-indolizone, D-463
 11,12-Dihydroindolo[2,3-*a*]carbazole, D-464
 2,3-Dihydro-1*H*-indol-5-ol, D-465
 1,2-Dihydro-3*H*-indol-3-one, I-104
 1,3-Dihydro-2*H*-indol-2-one, I-103
 8,13-Dihydroindolo[2',3':3,4]pyrido[1,2-*b*][2,7]naphthyridin-5(7*H*)-one, N-70
 8,13-Dihydroindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, R-168
 12,13-Dihydro-5*H*-indolo[2,3-*a*]pyrrolo[3,4-*c*]carbazole-5,7(6*H*)-dione, A-1393
 2,9-Dihydro-1*H*-indolo[3,2-*b*]pyrrolo[2,3,4-*gh*]phenanthridine-1,8-diol, A-1396
 5,5*a*-Dihydro-6*H*-indolo[2',3':3,4]pyrrolo[2,1-*b*]quinazoline-11,13-dione, W-26
 5*a*,6-Dihydro-3-(1*H*-indol-3-yl)-1,4-dimethoxy-10*b**H*-benzofuro[2,3-*b*]indole-2,10*b*-diol, D-466
 3,4-Dihydro-3-(1*H*-indol-3-yl)methyl-1*H*-1,4-benzodiazepine-2,5-dione, D-467
 6,7-Dihydro-7-(1*H*-indol-3-ylmethyl)quinazolinol[3,2-*a*][1,4]benzodiazepine-5,13-dione, A-1479
 5,6-Dihydro-3-[2-(1*H*-indol-3-yl)-2-oxoethyl]-2(1*H*)-pyridinone, D-468
 3-[1,5-Dihydro-4-(1*H*-indol-3-yl)-5-oxo-2*H*-pyrrol-2-ylidene]-1,3-dihydro-2*H*-indol-2-one, P-711
 5*a*,6-Dihydro-3-(1*H*-indol-3-yl)-1,2,4-trimethoxy-10*b**H*-benzofuro[2,3-*b*]indol-10*b*-ol, D-466
 Dihydroisoalamarine, A-235
 5,11-Dihydroisobellendine, I-194
 3,4-Dihydroisocarbostyryl, D-469
 Dihydroisoconessimine, C-595
 Dihydroisokomarovine, Q-53
 8,9-Dihydroisoorientalinone, O-112
 Dihydroisopelletierine, S-216
 3,5-Dihydro-2-isopropenyl-5-methylfuro[3,2-*c*]quinolin-4(2*H*)-one, A-1377
 2,3-Dihydro-2-isopropyl-4,8-dimethoxy-9-methylfuro[2,3-*b*]quinolinium, B-14
 3,9-Dihydro-2-isopropyl-7,8-methylenedioxy-9-methylfuro[2,3-*b*]quinolin-4(2*H*)-one, L-296
 3,4-Dihydro-6,7-isokinolinediol, D-421
 3,4-Dihydro-1,6,7-isokinolinetriol, I-305
 3,4-Dihydro-1-isokinolinol, D-469
 3,4-Dihydro-1(2*H*)-isoquinolinone, D-469
 8,9-Dihydroisoroemerialinone, O-112
 Dihydroisosakerol, G-68
 Dihydroisocloavine, S-273
 19,20-Dihydroisostyrychne, P-672
 4,5-Dihydro-3(2*H*)-isoxazolone, I-347
 Dihydrojasminine, J-25
 Dihydrojaspamide, H-15
 Dihydrojoubertamine, J-49
 9,10-Dihydrokramadine, K-29
 2,3-Dihydrokoniamborine, K-64
 Dihydrokoumine, K-96
 Dihydrokreysiginone, K-100
 Dihydrokurchessine, P-601
 8,9-Dihydrolactimidomycin, L-7
 Dihydrolemellarin B, L-24
 Dihydrolatumcidin, A-14
 5,6-Dihydrolepidine, L-116
 Dihydroleptopine, L-122
 Dihydrolinaresine, R-156
 Dihydrolongicaudatine Y, L-250
 Dihydrolongicaudatine L, L-249
 DihydroLuciduline, L-280
 Dihydrolycolucine, L-277
 Dihydrolycopodine, L-341
 Dihydrolycorine, Z-15
 α -Dihydrolysergol, F-49
 Dihydrolysergol, F-49
 Dihydrolythrine, V-95
 Dihydromacarpine, D-510
 Dihydromaleimide β -*D*-glucoside, D-460
 Dihydromaleimide, D-460
 Dihydromaltophilin, X-6
 3,4-Dihydromanzamine A *N*-oxide, M-91
 3,4-Dihydromanzamine A, M-91
 3,4-Dihydromanzamine J, M-94
 Dihydromelosmine, M-194
 2,3-Dihydromenisporphine, M-199
 6,7-Dihydro-10-methoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, D-175
 5,6-Dihydro-1-[3-(7-methoxy-1,3-benzodioxol-5-yl)-1-oxopropyl]-2(1*H*)-pyridinone, D-470
 5,6-Dihydro-*N*-[3-(7-methoxy-1,3-benzodioxol-5-yl)-2-propenoyl]-2(1*H*)-pyridinone, D-470
 1,2-Dihydro-1-(4-methoxybenzoyl)-6,7-methylenedioxyisoquinoline, M-242
 3,4-Dihydro-3-(4-methoxybenzyl)-4-methyl-1*H*-1,4-benzodiazepine-2,5-dione, C-890
 2,3-Dihydro-2-(4-methoxybenzyl)-1-methyl-4(1*H*)-quinazolinone, G-127
 7,8-Dihydro-8-methoxyberberine, O-216
 3,11-Dihydro-11-[(1-methoxy-9*H*-carbazol-3-yl)-methyl]-3,5-dimethyl-3-(4-methyl-3-pentenyl)-pyrano[3,2-*a*]carbazole, C-425
 3,11-Dihydro-11-[(1-methoxy-9*H*-carbazol-3-yl)-methyl]-3,3,5-trimethylpyrano[3,2-*a*]carbazole, C-424
 8,13-Dihydro-1-(methoxycarbonyl)-7*H*-benzo[*g*]indolo[2,3-*a*]quinolin-6-ium(1+), O-140
 14,19-Dihydro-11-methoxycondycarpine, T-673
 19,20-Dihydro-11-methoxydiaboline, C-115
 2,3-Dihydro-4-methoxy- α,α -dimethylfuro[2,3-*b*]quinoline-2-methanol, P-515
 3,11-Dihydro-10-methoxy-3,5-dimethyl-3-(4-methyl-3-pentenyl)pyrano[3,2-*a*]carbazole, M-47
 7,9-Dihydro-2-methoxy-1,9-dimethyl-1*H*-purine-6,8-dione, L-165
 3,7-Dihydro-2-methoxy-3,7-dimethyl-6*H*-purin-6-one, T-355
 3,4-Dihydro-6(8)-methoxy-5,8(5,6)-dimethyl-1*H*-pyrano[3,4-*c*]pyridin-1-one, G-57
 3,4-Dihydro-5-methoxy-2,2-dimethyl-2*H*-pyrano[2,3-*b*]quinoline, M-251
 7,8-Dihydro-4-methoxy-1,3-dioxolo[4,5-*g*]isoquinoline, N-292
 1,2-Dihydro-9-methoxyellipticine, M-252
 2-(2,3-Dihydro-4-methoxyfuro[2,3-*b*]quinolin-2-yl)-1,2-propanediol, D-945
 2,3-Dihydro-5-methoxy-1*H*-indole, D-465
 11,12-Dihydro-6-methoxyindolo[2,3-*a*]carbazole-5-carbonitrile, C-828
 1,3-Dihydro-5-methoxy-2*H*-indol-2-one, I-92
 8,13-Dihydro-10-methoxyindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, H-361
 7,12-Dihydro-10-methoxyindolo[2,3-*a*]quinolizine-4(6*H*)-one, H-63
 3,4-Dihydro-6-methoxy-1,7-isokinolinediol, I-305
 14,15-Dihydro-10-methoxykopsinone, K-80
 2,3-Dihydro-5-methoxy-3-[4-(methylamino)phenyl]pyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, S-264
 6,7-Dihydro-10-methoxy-7-methoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-4-methanol, C-2
 3,11-Dihydro-5-[(1-methoxy-3-methyl-9*H*-carbazol-9-yl)methyl]-3,3-dimethylpyrano[3,2-*a*]carbazole, M-745
 3,11-Dihydro-5-[(1-methoxy-3-methyl-9*H*-carbazol-9-yl)methyl]-3-methyl-3-(4-methyl-3-pentenyl)pyrano[3,2-*a*]carbazole, M-753
 3,4-Dihydro-6-methoxy-1-methyl- β -carboline, H-591
 3,4-Dihydro-7-methoxy-1-methyl- β -carboline, H-592
 8,9-Dihydro-4-methoxy-7-methyl-1,3-dioxolo[4,5-*f*]isoquinolin-6(7*H*)-one, T-306
 5,6-Dihydro-1-(3-methoxy-4,5-methylenedioxyaminamoyl)-2(1*H*)-pyridinone, D-470
 3,4-Dihydro-8-methoxy-6,7-methylenedioxyisoquinoline, N-292
 3,4-Dihydro-3-methoxy-8,9-methylenedioxyphenanthridine, M-448
 1-[(4,5-Dihydro-3-methoxy-4-methylene-1*H*-pyrrol-2-yl)thio]-1-tridecanone, I-171
 1,5-Dihydro-4-methoxy-5-(1-methylethyl)-1-(6,6,6-trichloro-3-methoxy-5-methyl-1-oxo-2-hexenyl)-2*H*-pyrrol-2-one, D-973
 10,11-Dihydro-7-methoxy-13-methyl-11,5-(imino-methano)-5*H*-benzo[4,5]cyclohepta[1,2-*f*]-1,3-benzodioxol-8-ol, A-954
 2,3-Dihydro-5-methoxy-6-methyl-1*H*-indole-1-carboxaldehyde, D-432
 1,3-Dihydro-5-methoxy-1-methyl-2*H*-indol-2-one, I-92
 8,14-Dihydro-10-methoxy-14-methylindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, D-158
 7,12-Dihydro-10-methoxy-2-methylindolo[2,3-*a*]quinolizine-4(6*H*)-one, H-62
 2,3-Dihydro-5-methoxy-6-methyl-7-[(3-methyl-2-butenyl)oxy]-1-oxo-4(1*H*)-isoindolecarboxylic acid, D-957
 3,4-Dihydro-7-methoxy-2-methyl-5,6-methylenedioxyisoquinolinium(1+), T-344
 3,4-Dihydro-7-methoxy-2-methyl-5,6-methylenedioxy-1(2*H*)-isoquinolinone, T-306
 1,2-Dihydro-5-(4-methoxy-2-methyl-1-naphthalenyl)-3-methyl-6,8-isokinolinediol, A-986
 6,7-Dihydro-3-methoxy-2-methyl-5-octyl-4,8(1*H*,5*H*)-quinolinedione, A-1325
 1,2-Dihydro-4-methoxy-1-methyl-2-oxo-3-pyridinecarbonitrile, R-96
 4,9-Dihydro-7-methoxy-1-methyl-3*H*-pyrido[3,4-*b*]indole, H-592
 1,5-Dihydro-4-methoxy-1-methyl-2*H*-pyrrol-2-one, P-954
 1,2-Dihydro-3-methoxy-1-methyl-4-quinolinol, E-13
 3,4-Dihydro-6-methoxy-2-methyl-7-[(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-8-isoquinolinyl)oxy]-1(2*H*)-isoquinolinone, B-102
 7,8-Dihydro-8-methoxynitidine, D-482
 1,2-Dihydro-4-methoxy-2-oxo-3-pyridinecarbonitrile, D-668
 1,4-Dihydro-7-methoxy-4-oxo-2-quinolinecarboxylic acid, D-676
 1,2-Dihydro-8-methoxy-2-oxo-4-quinolinecarboxylic acid, D-673
 1,4-Dihydro-6-methoxy-4-oxo-2-quinolinecarboxylic acid, D-675
 13,14-Dihydro-4-methoxyoxypalmitate, T-301
 14,15-Dihydro-15-methoxyphyllorchrysin, A-630
 4,5-Dihydro-2'-methoxypiperine, W-17
 1,5-Dihydro-5-methoxy-2*H*-pyrrol-2-one, D-460
 1,3-Dihydro-3-methoxy-2*H*-pyrrol-2-one, D-459
 1,5-Dihydro-4-methoxy-2*H*-pyrrol-2-one, P-954
 1,2-Dihydro-2-methoxyrhazimine, R-73
 3,6-Dihydro-4-methoxy- α,α,α -6-trimethyl-2*H*-furo[2,3-*f*][2,3]benzoxazine-2-methanol, E-294
 3,5-Dihydro-6-methoxy-2,3,3-trimethylfuro[3,2-*c*]quinolin-4(2*H*)-one, O-82
 7,9-Dihydro-2-methoxy-1,7,9-trimethyl-1*H*-purine-6,8-dione, L-165
 3,12-Dihydro-6-methoxy-3,3,12-trimethyl-7*H*-pyrano[2,3-*c*]acridin-7-one, A-120
 2,11-Dihydro-9-methoxy-2,2,8-trimethylpyrano[2,3-*a*]carbazole, C-511
 3,11-Dihydro-5-methoxy-3,3,8-trimethylpyrano[3,2-*a*]carbazole, P-883
 3,11-Dihydro-9-methoxy-3,3,8-trimethylpyrano[3,2-*a*]carbazole, P-883
 3,4-Dihydro-3-methoxytrisphaeridine, M-448
 10,11-Dihydro-7-(methylamino)benzo[*b*]pyrrolo[4,3,2-*de*][1,10]phenanthroline-8(9*H*)-one, P-503
 6,7-Dihydro-7-methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-11,12-dione, M-445
 5,6-Dihydro-5-methyl-1,3-benzodioxolo[5,6-*c*][1,3]dioxolo[4,5-*f*]phenanthridine, D-404
 13,14-Dihydro-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridine, D-510
 13,14-Dihydro-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridine-14-methanol, H-596
 9,14-Dihydro-14-methyl-5*H*-benzo[2,3][1,7]-naphthyridino[8,7-*b*]quinazolin-5-one, E-311
 7,8-Dihydro-*N*-methylberberine, L-23
 6,7-Dihydro-13-methylbis[1,3]benzodioxolo[5,6-*a*:4',5'-*g*]quinolinium, C-699

- 5,6-Dihydro-14-methylbis[1,3]benzodioxolo[5,6-*a*:5',6'-g]quinolizinium(1+), W-22
- 5,6-Dihydro-5-methyl-2,3,7,8-bis(methylenedioxy)-benzo[*c*]phenanthridine, D-510
- 1,3-Dihydro-3-(3-methyl-2-butenylidene)-2*H*-indol-2-one, D-471
- 1,3-Dihydro-6-(3-methyl-2-butenyl)-2*H*-indol-2-one, H-701
- 3,4-Dihydro-1-methyl-β-carboline, M-426
- 3-(3,4-Dihydro-12-methyl-5-cytisyl)-12-methylcytisine, D-704
- 7,8-Dihydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5(6*H*)-one, I-305
- 3,7-Dihydro-7-methyl-1,3-dipropyl-1*H*-purine-2,6-dione, X-5
- 3,4-Dihydro-2-methyllellipticine, E-67
- 3,4-Dihydro-6,7-methylenedioxy-1(2*H*)-isoquinolinone, I-305
- 6,7-Dihydro-7-methylene-5*H*-2-pyridin-5-ol, C-562
- 10,11-Dihydro-8-(1-methylethyl)-11-(2-methylpropylidene)-6*H*-oxepino[2,3-*d*]pyrazino[1,2-*a*]pyrimidine-6,9(8*H*)-dione, C-449
- [3-[1,6-Dihydro-5-(1-methylethyl)-6-oxopyrazinyl]-propyl]guanidine, A-1421
- 4,5-Dihydro-6-(*N*-methylformamido)-5-oxo-1,2-dithiolo[4,3-*b*]pyrrole, O-124
- N*-(2,5-Dihydro-3-methyl-2-furanyl)-4-hydroxy-L-glutamine, L-256
- 6,7-Dihydro-3-methylimidazo[1,2-*a*]pyridin-8(5*H*)-one, S-293
- 11,12-Dihydro-14-methyl-11,5-(iminomethano)-5*H*-cyclohepta[1,2-*f*:4,5-*f'*]bis[1,3]benzodioxole, R-35
- 2,3-Dihydro-2-methyl-1*H*-indol-1-amine, D-472
- 2,3-Dihydro-2-methyl-1*H*-indole-1-carboxaldehyde, D-472
- 2,3-Dihydro-2-methylindole, D-472
- 8,13-Dihydro-2-methylindolo[2',3':3,4]pyrido[1,2-*b*]2,7]naphthyridin-5(7*H*)-one, A-1012
- Dihydro-*O*-methylmacusine B, T-425
- 7,9-Dihydro-9-methyl-6-(methylamino)-8*H*-purin-8-one, A-819
- 5,10-Dihydro-5-methyl-9-(3-methyl-2-butenyl)-1-phenazinecarboxylic acid, M-418
- 3,4-Dihydro-2-methyl-6,7-methylenedioxy-2(1*H*)-isoquinolinone, I-305
- 3,5-Dihydro-5-methyl-2-(1-methylethyl)furo[3,2-*c*]quinolin-4(2*H*)-one, A-1377
- 7,10-Dihydro-10-methyl-8-(1-methylethyl)-1,3-dioxolo[4,5-*h*]furo[2,3-*b*]quinolin-6(8*H*)-one, L-296
- 2,3-Dihydro-7-methyl-2-(1-methylethyl)furo[3,2-*h*]isoquinolin-3-one, A-1294
- 1,5-Dihydro-3-methyl-1-(4-methyl-5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, H-467
- 3,11-Dihydro-3-methyl-3-(4-methyl-3-pentenyl)-pyranof[3,2-*a*]carbazole-5-carboxaldehyde, M-47
- 5,6-Dihydro-4-methyl-6-[(1-methyl-2-piperidinyl)-methyl]-2*H*-pyran-2-one, D-951
- 1,5-Dihydro-4-methyl-5-(2-methylpropylidene)-2*H*-pyrrol-2-one, P-784
- 3,7-Dihydro-1-methyl-3-(2-methylpropyl)-1*H*-purine-2,6-dione, X-5
- 1,4-Dihydro-1-methyl-2,7-naphthyridin-3(2*H*)-one, J-24
- 20,21-Dihydro-21-methyl-18-noralstophyllan-19-ol, T-20
- 19,20-Dihydro-21-methyl-18-norsarpagan-17,19-diol, D-491
- N*-[3-[[3-(3,4-Dihydro-5-methyl-4-oxo-2*H*-1-benzopyran-2-yl)propyl]amino]propyl]hexanamide, A-1033
- 2,3-Dihydro-1-(2-methyl-3-oxodecanoyl)pyrrole, D-473
- 2,3-Dihydro-1-(2-methyl-3-oxo-8-decenoyl)-1*H*-pyrrole, M-553
- N*-(4,5-Dihydro-4-methyl-5-oxo-1,2-dithiolo[4,3-*b*]pyrrol-6-yl)butanamide, A-759
- N*-(4,5-Dihydro-4-methyl-5-oxo-1,2-dithiolo[4,3-*b*]pyrrol-6-yl)-3-methylbutanamide, A-759
- 1,4-Dihydro-1-methyl-4-oxo-3-pyridinecarbonitrile, H-709
- 1,2-Dihydro-1-methyl-2-oxo-3-pyridinecarbonitrile, R-95
- 1,6-Dihydro-1-methyl-6-oxo-3-pyridinecarboxylic acid, H-711
- 1,5-Dihydro-3-methyl-1-(5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, D-448
- 3-(1,2-Dihydro-2-methyl-1-oxopyrrolo[1,2-*a*]pyrazin-3-yl)propylguanidine, P-242
- (1,5-Dihydro-3-methyl-2-oxo-2*H*-pyrrol-5-yl)-1,5-dihydro-5-hydroxy-3-methyl-2*H*-pyrrol-2-one, H-467
- 1-(2,5-Dihydro-4-methyl-5-oxo-1-*H*-pyrrol-2-yl)-4-methyl-5-oxoproline, D-474
- 5,6-Dihydro-5-methyl-2,3,7,8,10-pentamethoxybenzo[*c*]phenanthridine, D-509
- 1,5-Dihydro-5-methyl-1-phenazinone, P-833
- 3,4-Dihydro-4-methyl-3-(phenylmethyl)-1*H*-1,4-benzodiazepine-2,5-dione, C-890
- 6,7-Dihydro-6-methyl-7-(phenylmethylene)quinazolino[3,2-*a*][1,4]benzodiazepine-5,13-dione, B-74
- 3-[5,6-Dihydro-5-methyl-6-(1-propenyl)-2*H*-pyran-2-ylidene]-2,4-pyrrolidinedione, A-1339
- [3-[1,6-Dihydro-5-(2-methylpropyl)-6-oxopyrazinyl]propyl]guanidine, A-1419
- 4,5-Dihydro-2-(1-methylpropyl)thiazole, D-475
- 3,7-Dihydro-7-methyl-1*H*-purine-2,6-dione, X-5
- 3,7-Dihydro-1-methyl-1*H*-purine-2,6-dione, X-5
- 3,7-Dihydro-3-methyl-1*H*-purine-2,6-dione, X-5
- 3,9-Dihydro-9-methyl-1*H*-purine-2,6-dione, X-5
- 1,7-Dihydro-1-methyl-6*H*-purin-6-one, P-820
- 1,7-Dihydro-7-methyl-6*H*-purin-6-one, P-820
- 3,7-Dihydro-3-methyl-6*H*-purin-6-one, P-820
- 1,9-Dihydro-9-methyl-6*H*-purin-6-one, P-820
- 2,7-Dihydro-7-methyl-3*H*-pyrazolo[4,3-*c*][1,2,4]triazin-3-one, N-320
- 4,9-Dihydro-1-methyl-3*H*-pyrido[3,4-*b*]indole, M-426
- 4,9-Dihydro-1-methyl-3*H*-pyrido[3,4-*b*]indol-7-ol, H-592
- 6,7-Dihydro-7-methyl-5*H*-2-pyridine-5,7-diol, O-160
- 6,7-Dihydro-7-methyl-5*H*-2-pyridine, N-287
- 6,7-Dihydro-7-methyl-5*H*-2-pyridin-5-ol, C-562
- 6,7-Dihydro-7-methyl-5*H*-pyrindin-5-one, C-562
- 3,4-Dihydro-3-methyl-2*H*-pyrrole-2-carboxylic acid, D-477
- 2,3-Dihydro-3-methyl-1*H*-pyrrole, D-476
- 2,3-Dihydro-3-methyl-1*H*-pyrrolizin-1-one, L-263
- 7,8-Dihydro-3-methylpyrrolo[1,2-*a*]pyrimidin-2(6*H*)-one, D-478
- 21-(3,4-Dihydro-3-methyl-2*H*-pyrrol-5-yl)-3-hydroxy-20-methylpregnane-6,21-dione, P-279
- 21-(3,4-Dihydro-3-methyl-2*H*-pyrrol-5-yl)-3-hydroxy-20-methylpregnane-4,21-dione, S-339
- 21-(3,4-Dihydro-3-methyl-2*H*-pyrrol-5-yl)-3-hydroxy-20-methylpregn-5-en-21-one, T-423
- 20-(3,4-Dihydro-5-methyl-2*H*-pyrrol-2-yl)pregn-4-en-3-one, M-550
- 5,10-Dihydro-5-methyl-11*H*-quindolin-11-one, D-505
- 1,4-Dihydro-1-methylquinoline, D-507
- 1,3-Dihydro-1-methyl-4-(2-quinolinyl)-2*H*-imidazol-2-one, M-28
- 7,8-Dihydro-6-methylspiro[1,3-dioxolo[4,5-*g*]isoquinoline-5(6*H*),7'-[7*H*]indeno[4,5-*d*][1,3]dioxol]-8'(6'*H*)-one, F-176
- 8,9-Dihydro-*N*-methylstepharine, C-781
- 4,9-Dihydro-2-(methylsulfinyl)-1,3-thiazino[6,5-*b*]indole, C-842
- Dihydro-6-(methylsulfonyl)-1,2,3-dithiazin-4(3*H*)-one, S-162
- 1,2-Dihydro-*O*-methylazettine, T-53
- 4,5-Dihydro-2-methylthiazole, D-479
- 3,5-Dihydro-3-methyl-2*H*-thiopyranof[4,3,2-*cd*]indole-2-carboxylic acid, D-480
- 4,9-Dihydro-2-(methylthio)-1,3-thiazino[6,5-*b*]indole, C-842
- 1-[4,9-Dihydro-2-(methylthio)-1,3-thiazino[6,5-*b*]indol-4-yl]-2-propanone, D-481
- Dihydromontamicaine, A-95
- Dihydromurrayacaine, M-757
- Dihydromyricoidine, L-230
- 9,9*a*-Dihydronaphtho[2,3-*c*]isoxazol-3(1*H*)-one, N-31
- cis*-Dihydronarciclasine, N-37
- trans*-Dihydronarciclasine, N-37
- Dihydronematophin, T-639
- Dihydronitidine, D-482
- Dihydronitraraine, N-243
- Dihydronitraridine, Q-52
- 1-[1,4-Dihydro-4-nonyl-5-(1-oxodecyl)-3-pyridinyl]-1-dodecanone, D-242
- 1,1'-(1,4-Dihydro-4-nonyl-3,5-pyridinediyl)bis[1-dodecanone], D-242
- 1,1'-(1,4-Dihydro-4-nonyl-3,5-pyridinediyl)bis[1-dodecanone], D-242
- Dihydronorpupeline, T-288
- 8,14-Dihydronorsalutaridine, I-318
- 14,15-Dihydro-*A*-norsecurinan-11-one, N-307
- Dihydronorsecurinine, N-307
- Dihydronudaurine, A-955
- Dihydroobscurinervidine, O-7
- Dihydroobscurinervine, O-8
- 1,2-Dihydro-2-octadecyl-3*H*-indol-3-one, H-646
- 11,12-Dihydroorientalinone, O-112
- 2,3-Dihydro-4-oxa-1,3*a*-diazacyclohepta[*def*]fluoren-10(1*H*)-one, H-772
- 7,8-Dihydro-8-oxoadenine, A-819
- 12,13-Dihydro-8-oxoberberine, C-73
- 7,9-Dihydro-1-(3-oxobutyl)-1*H*-purine-6,8-dione, P-818
- 7,9-Dihydro-9-(3-oxobutyl)-1*H*-purine-6,8-dione, P-818
- N*-(4,5-Dihydro-5-oxo-1,2-dithiolo[4,3-*b*]pyrrol-6-yl)hexanamide, A-759
- N*-(4,5-Dihydro-5-oxo-1,2-dithiolo[4,3-*b*]pyrrol-6-yl)propanamide, A-759
- 1-(2,3-Dihydro-2-oxo-3-furanyl)-5-(hydroxymethyl)-1*H*-pyrrole-2-carboxaldehyde, D-483
- 7,8-Dihydro-8-oxoguanine, A-753
- 2,3-Dihydro-1-(1-oxo-4,6,8,12-hexadecatraen-10-ynyl)-1*H*-pyrrole, H-223
- 2,3-Dihydro-2-oxo-1*H*-imidazole-4-carboxylic acid, D-484
- 2,3-Dihydro-2-oxo-1*H*-indole-3-acetic acid, D-485
- Dihydro-3-oxoindole, I-104
- 5,11-Dihydro-*α*-oxoindolo[3,2-*b*]carbazole-6-acetic acid, M-62
- 3-(1,2-Dihydro-2-oxo-3*H*-indol-3-ylidene)-1,3-dihydro-2*H*-indol-2-one, B-131
- 3-(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)-1,3-dihydro-2*H*-indol-2-one, I-73
- 2-(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)-7-hydroxycyclopent[*g*]indol-3(2*H*)-one, D-486
- α*-(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)-1*H*-indole-3-acetonitrile, D-487
- 6-(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)indolo[2,1-*b*]quinazolin-12(6*H*)-one, C-77
- 11,33-Dihydro-10-oxopenitrem B, P-194
- 1,4-Dihydro-4-oxo-6-(phenylmethyl)-3-pyridinecarboxamide, A-1483
- 1,5-Dihydro-1-(1-oxo-3-phenylpropyl)-2*H*-pyrrol-2-one, S-86
- 1,8-Dihydro-1-(1-oxopropyl)lycodin-8-one, G-228
- N*-(7,8-Dihydro-8-oxo-*H*-purin-6-yl)-β-alanine, E-177
- 1,6-Dihydro-6-oxo-2-pyridineacetic acid, H-708
- 1,6-Dihydro-6-oxo-3-pyridinecarboxylic acid, H-711
- 1,4-Dihydro-4-oxo-3-pyridinecarboxylic acid, H-709
- 12,13-Dihydro-13-oxopyrido[1,2-*a*:3,4-*b'*]diindol-5-ium(1+), F-13
- 2,3-Dihydro-1-oxo-1*H*-pyrrolizine-7-carboxaldehyde, D-14
- 2,3-Dihydro-1-oxo-1*H*-pyrrolizine-3-carboxylic acid, D-488
- 1,2-Dihydro-1-oxo-4-(1*H*-pyrrol-2-yl)-3-isoquinolinecarboxylic acid, M-104
- 2-(1,4-Dihydro-4-oxo-2-quinazolinyl)-1*H*-indole-3-carboxylic acid, W-27
- 1,2-Dihydro-2-oxo-4-quinolinecarboxaldehyde, H-721
- 1,4-Dihydro-4-oxo-2-quinolinecarboxylic acid, H-725

- 1,2-Dihydro-2-oxo-4-quinolinecarboxylic acid, H-723
 1,4-Dihydro-4-oxo-2,3-quinolinedicarboxylic acid, H-728
 1,4-Dihydro-4-oxo-2-quinolinehexanoic acid, M-68
 1,4-Dihydro-4-oxo-1- β -D-ribofuranosyl-3-pyridinecarboxamide, H-709
 5,6-Dihydro-1-[1-oxo-3-(3,4,5-trimethoxyphenyl)-2-propenyl]-2(1*H*)-pyridinone, P-473
 1,5-Dihydro-1-[1-oxo-3-(3,4,5-trimethoxyphenyl)propyl]-2*H*-pyrrol-2-one, D-489
 16,16'-Dihydro-17,17'-oxybisnordihydrotoxiciferine, M-120
 5,6-Dihydro-2-(1,3-pentadienyl)-4(1*H*)-pyridinone, D-490
 Dihydroptentagynine, M-347
 5,6-Dihydro-2,3,4,9,10-pentamethoxydibenzo[*a,g*]-quinolizinium(1+), A-1042
 20*R*,21*S*-Dihydroperaksin-17-*al*, D-491
 Dihydroperaksine, D-491
 20*R*,21*S*-Dihydroperaksine, D-491
 Dihydroperamine, P-254
 (+)-Dihydroperamine, P-254
 Dihydroperaminole, H-57
 Dihydroperimagine, P-259
 Dihydroperiphylline, P-265
 5,10-Dihydrophencomycin methyl ester, P-320
 5,6-Dihydro-1-(3-phenylpropanoyl)-2(1*H*)-pyridinone, P-463
 2,3-Dihydro-1-phenyl-1*H*-pyrrole, D-499
 5,6-Dihydro-3-phenyl-4*H*-pyrrolo[1,2-*b*]pyrazole, D-492
 Dihydropimidine, M-529
 10,11-Dihydroperiperide, R-60
 4,5-Dihydropiperettine, P-452
 6,7-Dihydropiperettine, P-452
 $\Delta^{\alpha,\beta}$ -Dihydropiperine, P-467
 4,5-Dihydropiperlonguminine, P-462
 $\Delta^{\alpha,\beta}$ -Dihydropiperlonguminine, P-462
 8,9-Dihydropiplartine, P-473
 24,25-Dihydroplakinamine A, P-496
 Dihydroplakinamine K, P-496
 19,20-Dihydropolyneuridine, P-562
 Dihydropontevodrine, P-566
 Dihydropresecamine, P-624
 20-Dihydroprochaetoglobosin II, C-328
 3,4-Dihydro-5-propanoyl-2*H*-pyrrole, D-493
 6,7-Dihydro-4-(2-propenyl)-5*H*-cyclopenta[*c*]pyridin-5-one, L-272
 3,7-Dihydro-3-propyl-1*H*-purine-2,6-dione, X-5
 20-Dihydroprotochaetoglobosin III, C-325
 Dihydroprotoemine, P-664
 Dihydroprotopine, P-667
 Dihydropsychothrine, E-72
 1,5-Dihydro-2,4,6(3*H*)-pteridinetriene, D-494
 7,8-Dihydropterin-6-carboxylic acid, A-817
 7,8-Dihydropteroylglutamic acid, P-772
 Dihydropumiliotoxin A, A-370
 7,9-Dihydro-1*H*-purine-6,8-dione, P-818
 3,7-Dihydro-1*H*-purine-2,6-dione, X-5
 7,9-Dihydro-1*H*-purine-2,6,8(3*H*)-trione, U-45
 1,7-Dihydro-6*H*-purin-6-one, P-820
 Dihydropyoverdin Pp 2, P-869
 Dihydropyraacrimycin A, P-876
 7,8-Dihydro-5*H*-pyrano[3,4-*b*]pyrazine-7-methanol, C-517
 3,4-Dihydro-1*H*-pyrano[3,4-*c*]pyridin-1-one, G-54
 5,6-Dihydropyridin-2-ol, D-495
 5,6-Dihydro-2(1*H*)-pyridinone, D-495
 3,4-Dihydro-5-(2-pyridinyl)-2*H*-pyrrole-2-carboxylic acid, D-496
 2,9-Dihydro-1*H*-pyrido[3,4-*b*]indol-1-one, P-914
 5,6-Dihydro-2(1*H*)-pyridone, D-495
 Dihydro-2,4(1*H*,3*H*)-pyrimidinedione, D-497
 4,8-Dihydropyrimido[5,4-*c*]-1,2,4-triazine-3,5,7(6*H*)-trione, D-498
 3,4-Dihydro-2*H*-pyrrole-2-carboxylic acid, D-501
 2,3-Dihydro-1*H*-pyrrole, D-499
 3,4-Dihydro-2*H*-pyrrole, D-500
 2,3-Dihydro-1*H*-pyrrolizine-7-carboxaldehyde, D-14
 11,11*a*-Dihydro-1*H*-pyrrolizino[3,2-*b*]quinoline-3,10(2*H*,5*H*)-dione, C-370
 6,7-Dihydropyrrolo[2,3-*c*]azepine-4,8(1*H*,5*H*)-dione, A-255
 2,3-Dihydro-1*H*-pyrrolo[2,1-*c*][1,4]benzodiazepine-5,11(10*H*,11*aH*)-dione, D-502
 5,9-Dihydro-1*H*-pyrrolo[3',4':6',7']cyclohepta[2,1-*b*:4,3-*c'*:*d'*]diindole-1,3(2*H*)-dione, A-1392
 5,10-Dihydro-1*H*-pyrrolo[3',4':4,5]oxocino[2,3-*b*:6',7,8-*c'*:*d'*]diindole-1,3(2*H*)-dione, A-1398
 2,3-Dihydropyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, D-503
 1-(3,4-Dihydro-2*H*-pyrrol-5-yl)ethanone, A-49
 3-(3,4-Dihydro-2*H*-pyrrol-5-yl)-2-hydroxy-2-propanamide, P-877
 1-(2,3-Dihydro-1*H*-pyrrol-1-yl)-2-methyl-1,3-decanedione, D-473
 1-(2,3-Dihydro-1*H*-pyrrol-1-yl)-2-methyl-8-decene-1,3-dione, M-553
 2-(3,4-Dihydro-2*H*-pyrrol-5-yl)-1-piperidinecarboxaldehyde, S-330
 1-(3,4-Dihydro-2*H*-pyrrol-5-yl)-1-propanone, D-493
 3-(3,4-Dihydro-2*H*-pyrrol-5-yl)-2-propenamide, P-876
 2-(3,4-Dihydro-2*H*-pyrrol-5-yl)pyridine, D-504
 3-(3,4-Dihydro-2*H*-pyrrol-5-yl)pyridine, M-792
 1-(3,4-Dihydro-2*H*-pyrrol-5-yl)-9*H*-pyrido[3,4-*b*]indole, E-288
 Dihydroquinamine, Q-12
 5,10-Dihydro-11*H*-quindolin-11-one, D-505
 5,6-Dihydro-11*H*-quindolinol-11-one, D-506
 1,4-Dihydroquinoline, D-507
 6,7-Dihydroquinolino[3,2-*a*][1,4]benzodiazepine-5,13-dione, S-154
 Dihydroretrofractamide B, R-60
 Dihydroretrofractamide C, R-60
 Dihydroretrosine, P-516
 5,21-Dihydrorhazinilam, R-76
 3,9-Dihydro-9- β -D-ribofuranosyl-1*H*-purine-2,6-dione, X-13
 1,9-Dihydro-9- β -D-ribofuranosyl-6*H*-purin-6-one, I-148
 1,4-Dihydro-3- β -D-ribofuranosyl-7*H*-pyrazolo[4,3-*d*]pyrimidin-7-one, F-132
 1,9-Dihydro-9- β -D-ribofuranosyl-6*H*-purin-6-one, P-820
 3,12-Dihydroroquefortine, R-125
 Dihydrorostratine, P-602
 Dihydrorugosinone, R-156
 Dihydroruine, D-621
 Dihydrorutaecarpine, D-508
 13*b*,14-Dihydrorutaecarpine, D-508
 25-Dihydrosaframycin A, S-4
 25-Dihydrosaframycin B, S-4
 8,14-Dihydrosalutaridine, I-318
 Dihydrosanguilline, D-509
 Dihydrosanguinarine, D-510
 5,6-Dihydrosarconidine, P-608
 Dihydrosarcostin, P-602
 Dihydrosauvagnine, S-108
 6,7-Dihydroschelhammeridine, C-587
 Dihydroshocherone, D-113
 Dihydrosecamine, S-174
 Dihydrosecocepharanthine, S-179
 15,20-Dihydrosecodine, S-184
 16,17-Dihydrosecodine, S-184
 16,17-Dihydrosecodin-17-ol, S-184
 Dihydrosecoquettamine, S-197
 Dihydrosecurinine, S-206
 Dihydroseminine, S-214
 Dihydrosenkirkine, S-245
 Dihydroshihunine, D-511
N-(Dihydrosinapoyl)tyramine, D-512
 (16*R*)-Dihydrositsirikine, S-325
 (16*S*)-Dihydrositsirikine, S-325
 Dihydrosolanidine T, D-205
 Dihydrosolasone A, S-200
 Dihydrosolasodine, S-445
 Dihydrosolasuaveoline, S-445
 Dihydrosongarine, N-30
 6',7'-Dihydrospiro[cyclohexane-1,8'(5*H*)-imidazo[1,2-*a*]pyridin]-2-ol, N-240
 11,12-Dihydrostemofofline, S-515
 Dihydrostemonine, S-521
 8,9-Dihydrostepharine, C-781
 3',4'-Dihydrostephasubine, E-102
 Dihydrostrobiline, S-587
 19,20-Dihydrotabernamine, T-8
 Dihydrotaxilamine, T-50
 Dihydrotaxol, T-52
 4',17-Dihydro-17*\alpha*-tchibangensine, U-52
 4',17-Dihydro-17*\beta*-tchibangensine, U-52
 5,8-Dihydro-6-(5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)furo[3,4-*e*]-1,3-benzodioxol-8-ol, E-43
 5,6-Dihydro-2,3,10,11-tetramethoxydibenzo[*a,g*]-quinolizinium(1+), P-719
 5,6-Dihydro-2,3,9,10-tetramethoxydibenzo[*a,g*]quinolizinium(1+), P-34
 5,6-Dihydro-2,3,10,11-tetramethoxy-8*H*-dibenzo[*a,g*]quinolizinium-8-one, O-202
 3,4-Dihydro-5,6,7,8-tetramethoxyisoquinoline, T-276
 5,6-Dihydro-2,3,7,8-tetramethoxy-5-methylbenzo[*c*]phenanthridine-6-acetic acid, D-513
 5,6-Dihydro-2,3,7,8-tetramethoxy-5-methylbenzo[*c*]phenanthridin-6-one, D-514
 5,6-Dihydro-1,2,9,10-tetramethoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline, D-159
 3,4-Dihydro-5,6,7,8-tetramethoxy-1-methylisoquinoline, T-233
 5,6-Dihydro-2,3,7,8-tetramethoxy-5-methyl-6-oxobenzo[*c*]phenanthridine, D-514
 3,9-Dihydro-2,3,3,9-tetramethylfuro[2,3-*b*]quinolin-4(2*H*)-one, I-19
 3,9-Dihydro-2,2,3,9-tetramethylfuro[2,3-*b*]quinolin-4(2*H*)-one, L-88
 3,5-Dihydro-2,3,3,5-tetramethylfuro[3,2-*c*]quinolin-4(2*H*)-one, O-82
 6,7-Dihydro-1,3,6,6-tetramethyl-8(5*H*)-isoquinolinone, D-515
 Dihydrothalietrinine, T-310
 Dihydrothalthemine, T-313
 4-(3,4-Dihydro-2*H*-1,4-thiazin-6-yl)phenol, D-457
 3-(4,5-Dihydro-2-thiazolecarbonyl)-1*H*-indole, D-516
 1,2-Dihydrothiazolo[2',3':2,3]pyrimido[4,5-*b*]indole-1-carboxylic acid, D-517
 (4,5-Dihydro-2-thiazolyl)-1*H*-indole-3-ylmethanone, D-516
 2,3-Dihydro-2-thioxo-4(1*H*)-pyrimidinone, D-518
 3,4-Dihydro-4-thioxo-2(1*H*)-pyrimidinone, H-556
 4,5-Dihydro-2-thioxo-4(3*H*)-thiazolecarboxylic acid, T-394
 Dihydrotoddaline, D-407
 Dihydrotomatillidine, T-423
 C-Dihydrotoxiciferine I, T-436
 13-Dihydro-2,9,13-triacetylaxine B, T-45
 3,4-Dihydro-1,6,7-trihydroxyisoquinoline, I-305
 15,20-Dihydro-12,15,18-trihydroxy-14-methyl-21-norsenecionan-11,16-dione, S-122
 15,20-Dihydro-12,15,20-trihydroxy-4-methyl-4,8-secosenecionan-8,11,16-trione, O-92
 1,4-Dihydro-3,7,8-trihydroxy-4-oxo-2-quinolinecarboxylic acid, T-272
 1,5-Dihydro-3,4,5-trihydroxy-2*H*-pyrrol-2-one, D-519
 15,20-Dihydro-12,15,20-trihydroxysenecionan-11,16-dione, J-3
 5,10-Dihydro-4,6,8-trihydroxy-10-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-11*H*-dibenzo[*b,e*][1,4]diazepin-11-one, D-297
 3,12-Dihydro-6,10,11-trihydroxy-3,3,12-trimethyl-7*H*-pyrano[2,3-*c*]acridin-7-one, C-472
 5,6-Dihydro-1-(3,4,5-trimethoxycinnamoyl)-2(1*H*)-pyridinone, P-473
 3,4-Dihydro-5,8,9-trimethoxy-2,2-dimethyl-2*H*-pyrano[2,3-*b*]quinoline, M-251
 8,13-Dihydro-2,3,10-trimethoxyindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, E-307
 3,4-Dihydro-5,6,7-trimethoxyisoquinoline, I-307
 10,11-Dihydro-3,7,8-trimethoxy-12-methyl-10,5-(iminomethano)-5*H*-dibenzo[*a,d*]cyclohepten-4-ol, I-333

- 10,11-Dihydro-3,7,8-trimethoxy-12-methyl-10,5-(iminomethano)-5*H*-dibenzo[*a,d*]cyclohepten-2-ol, T-336
- 3,4-Dihydro-*ar*-trimethoxy-1-methylisoquinoline, D-520
- 3,4-Dihydro-5,6,7-trimethoxy-2-methyl-1(2*H*)-isoquinolinone, I-304
- 2,3-Dihydro-4,7,8-trimethoxy-9-methyl-2-(1-methylethenyl)furo[2,3-*b*]quinolinium, D-720
- 2,3-Dihydro-4,6,8-trimethoxy-9-methyl-2-(1-methylethenyl)furo[2,3-*b*]quinolinium(1+), P-757
- 1,2-Dihydro-4,7,8-trimethoxy-2-oxo-3-quinoline-carboxaldehyde, G-119
- 1,2-Dihydro-4,6,7-trimethoxy-2-oxo-3-quinoline-carboxaldehyde, M-175
- 1,9-Dihydro-5-[(2,4,4-trimethyl-1-cyclohexen-1-yl)methyl]phenazin-1-one, L-76
- 5,6-Dihydro-3,6,6-trimethyl-7*H*-cyclopenta[*c*]pyridin-7-one, D-525
- 1,4-Dihydro-4,6,7-trimethyl-9*H*-imidazo[1,2-*a*]purin-9-one, D-521
- 2,7-Dihydro-2,5,7-trimethyl-5*H*-imidazo[4,5-*e*]1,2,4-triazine-3,6-dione, D-522
- 7,8-Dihydro-2,7,7-trimethyl-1*H*-isoindolo[4,5,6-*cd*]indole-1,9(2*H*)dione, C-892
- 3,9-Dihydro-1,3,9-trimethyl-8-nitroso-1*H*-purine-2,6-dione, D-523
- 3,7-Dihydro-1,3,7-trimethyl-1*H*-purine-2,6-dione, C-15
- 3,7-Dihydro-3,3,10-trimethylpyrano[2,3-*c*]carbazole, E-300
- 3,11-Dihydro-3,3,5-trimethylpyrano[3,2-*a*]carbazole, G-81
- 5,8-Dihydro-1,8,8-trimethyl-1*H*-pyrano[3,4-*b*]pyridine-4,6-dione, D-524
- 4,8-Dihydro-4,6,8-trimethylpyrimido[5,4-*e*]1,2,4-triazine-3,5,7(6*H*)-trione, D-498
- 2,8-Dihydro-2,6,8-trimethylpyrimido[5,4-*e*]1,2,4-triazine-3,5,7(6*H*)-trione, D-498
- 5,6-Dihydro-3,6,6-trimethyl-7*H*-2-pyridin-7-one, D-525
- 2,5-Dihydro-2,4,5-trimethylthiazole, D-526
- 7,8-Dihydrotrubastrine, T-656
- 10,27-Dihydrotribingensin A, T-669
- 10,27-Dihydrotribingensin B, T-670
- Dihydrouracil, D-497
- 5',6'-Dihydrousambarensine *N*-oxide, U-52
- 3',4'-Dihydrousambarensine *N*^b-oxide, U-52
- 5',6'-Dihydrousambarensine, U-52
- 19,20-Dihydrousambarensine, U-52
- 3',4'-Dihydrousambarensine, U-52
- 18,19-Dihydrousambaridine Br, O-21
- 18,19-Dihydrousambaridine Vi, O-21
- 18,19-Dihydrousambarine, O-21
- Dihydroverticillatine, V-88
- Dihydrovertine, V-95
- Dihydrovincarpine, F-82
- 14,15-Dihydrovindolinine, V-132
- Dihydrovomifoline, D-491
- Dihydrowarifteine, W-9
- α,β -Dihydrowisanidine, W-16
- 4,5-Dihydrowisanine, W-17
- Dihydroxanthommatin, D-527
- Dihydroxyacidissiminol, A-77
- 1,3-Dihydroxy-9(10*H*)-acridinone, D-528
- 1,7-Dihydroxy-9(10*H*)-acridinone, D-529
- 1,8-Dihydroxy-9(10*H*)-acridinone, D-530
- 1,3-Dihydroxyacridone, D-528
- 1,7-Dihydroxyacridone, D-529
- 1,8-Dihydroxyacridone, D-530
- 2,8-Dihydroxyadenine, A-754
- Dihydroxyaerotheronin, A-158
- 20,25-Dihydroxyaflavinine, A-172
- 3',19-Dihydroxyaglaïne C, A-199
- 2,4-Dihydroxyaniline, A-705
- Dihydroxyanopterine, A-1071
- 4,5-Dihydroxyanthranilic acid, A-737
- 2*R*,9*R*-Dihydroxyaphyllidine, A-1336
- 2*S*,9*R*-Dihydroxyaphyllidine, A-1336
- 2,7-Dihydroxyapogeissoschizine, A-1362
- 1,2-Dihydroxyaporphine, D-531
- 2,3-Dihydroxyaricyriaflavin A, A-1393
- 2',3'-Dihydroxyasteranthine, A-1510
- 3,3-Dihydroxyazetidide, A-1606
- 2-(2,5-Dihydroxybenzamido)benzoic acid, D-532
- 2,4-Dihydroxybenzeneacetic acid, D-646
- $\alpha,3$ -Dihydroxybenzeneacetic acid, H-519
- $\alpha,4$ -Dihydroxybenzeneacetic acid, H-520
- 4,5-Dihydroxy-1,2-benzenedicarboxylic acid, D-533
- α,β -Dihydroxybenzenepropanoic acid, D-651
- 3,4-Dihydroxybenzenepropanoic acid, D-652
- 5,7-Dihydroxy-2*H*-benz[*b*]isoindole-4,9-dione, D-534
- 3,6-Dihydroxy-1,2-benzisoxazole, B-72
- 2,4-Dihydroxybenzoic acid, D-535
- 3,4-Dihydroxybenzoic acid, D-536
- 2,4-Dihydroxybenzotrile, D-535
- 2,6-Dihydroxybenzothiazole, B-78
- 2,7-Dihydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, D-538
- 2,5-Dihydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, D-537
- 2,4-Dihydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, H-409
- 2,4-Dihydroxybenzoxazole, B-81
- 2,6-Dihydroxybenzoxazole, B-82
- 2-[2-(2,5-Dihydroxybenzoyl)amino]-5-hydroxybenzoic acid, D-539
- N*-(2,4-Dihydroxybenzoyl)anthranilic acid, B-67
- 5-(3,4-Dihydroxybenzoyl)-4,8-dihydroxy-2-quinolinocarboxylic acid, T-512
- 3,4-Dihydroxybenzoylformic acid, D-649
- N*-(3,4-Dihydroxybenzoyl)glutamic acid, G-107
- N*-(2,4-Dihydroxybenzoyl)-4-hydroxyanthranilic acid, B-67
- N*-(2,5-Dihydroxybenzoyl)-5-hydroxyanthranilic acid, D-539
- N*-(2,4-Dihydroxybenzoyl)-4-methoxyanthranilic acid, B-67
- N*-(2,6-Dihydroxybenzoyl)-*O*-methyltyramine, M-280
- N*-(2,3-Dihydroxybenzoyl)putrescine, B-422
- 8-(3,4-Dihydroxybenzyl)-2,3,10,11-berbintetrol, D-541
- 8-(3,4-Dihydroxybenzyl)-2,3,9,10-berbintetrol, D-540
- 1-(3,4-Dihydroxybenzyl)-6,7-dihydroxyisoquinoline, P-80
- 1-(3,4-Dihydroxybenzyl)-6,7-dimethoxyisoquinoline, P-80
- 1-(3,4-Dihydroxybenzyl)-7-hydroxy-6-methoxyisoquinoline, P-80
- N*-(3,4-Dihydroxybenzyl)tyramine, N-288
- 3,4-Dihydroxy-2,5-bis(hydroxymethyl)pyrrolidine, D-542
- 1,2-Dihydroxy-3,5-bis(hydroxymethyl)-1*H*-pyrrolizidine, D-543
- 2,5-Dihydroxy-3,6-bis[5-(3-methyl-2-butenyl)-1*H*-indol-3-yl]-2,5-cyclohexadiene-1,4-dione, C-547
- 1,8-Dihydroxy-2,3,9,10-bis(methylenedioxy)protoberberine, D-409
- 5-(1,3-Dihydroxybutyl)-1,2-dihydroxy-3-(hydroxymethyl)-1*H*-pyrrolizidine, T-553
- 5-(1,3-Dihydroxybutyl)hyacinthacine A₁, T-553
- 4,5-Dihydroxycanthin-6-one, D-546
- 5,11-Dihydroxycanthin-6-one, D-548
- 5,9-Dihydroxycanthin-6-one, D-547
- 8,9-Dihydroxycanthin-6-one, D-549
- 9,10-Dihydroxycanthin-6-one, D-550
- 1,9-Dihydroxycanthin-6-one, D-544
- 10,11-Dihydroxycanthin-6-one, D-551
- 1,11-Dihydroxycanthin-6-one, D-545
- 2,7-Dihydroxycarbazole, C-123
- 2,7-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-556
- 2,8-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-557
- 1,6-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-552
- 1,7-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-553
- 1,8-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-554
- 2,6-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-555
- 1,6-Dihydroxy-9*H*-carbazole-3-carboxylic acid, D-558
- 1,7-Dihydroxy-9*H*-carbazole-3-carboxylic acid, D-559
- 2,7-Dihydroxy-9*H*-carbazole-3-carboxylic acid, D-560
- 4,8-Dihydroxy- β -carboline-1-carboxylic acid, D-561
- 2,4-Dihydroxycelorbicol, P-215
- 1,3-Dihydroxy- $\Delta^{8(14)}$ -cevanone, C-304
- 3,6-Dihydroxy- $\Delta^{12,14,16}$ -cevanone, C-300
- 3,6-Dihydroxycevanine, C-290
- 3,14-Dihydroxy-6-cevanone, C-295
- 3,16-Dihydroxy-6-cevanone, C-297
- 3,20-Dihydroxy-6-cevanone, C-298
- 6,12-Dihydroxycevan-3-one, D-563
- 3,25-Dihydroxycevan-6-one, D-562
- 2,6-Dihydroxycinchoninic acid, D-672
- 2,8-Dihydroxycinchoninic acid, D-673
- 3,4-Dihydroxycinnamic acid, D-654
- 3,5-Dihydroxycinnamic acid, D-654
- N*-(3,4-Dihydroxycinnamoyl)-1,4-butanediamine, C-456
- N*-(3,4-Dihydroxycinnamoyl)DOPA, C-534
- N*-(3,4-Dihydroxycinnamoyl)-5-hydroxyanthranilic acid, A-789
- Dihydroxycitraconone I, C-472
- 18,19*R*-Dihydroxycoconaridine, C-657
- 2,3-Dihydroxycycloarta-22,24-dien-26-oic acid, D-564
- 2,3-Dihydroxy-9,19-cyclolanost-22,24-dien-26-oic acid, D-564
- 1,4-Dihydroxy-2-cyclopentene-1-carboxamide, D-565
- 1,4-Dihydroxy-2-cyclopentene-1-carboxylic acid, D-565
- 7-(4,5-Dihydroxy-2-cyclopent-1-en-3-ylamino-methyl)-7-deazaguanine, Q-9
- 4,5-Dihydroxy-2-decenoic acid piperidide, D-566
- 4,5-Dihydroxy-2-decenoic acid, D-566
- 3,11-Dihydroxy-1,2-dehydrocrinane, H-5
- 3,10-Dihydroxydibenz[*c,g*]azecine-13(14*H*)-one, C-624
- 1,2-Dihydroxydibenz[*cd,f*]indol-4(5*H*)-one, D-567
- 1,3-Dihydroxydibenz[*cd,f*]indol-4(5*H*)-one, D-568
- 1,2-Dihydroxy-7*H*-dibenzo[*de,g*]quinolin-7-one, L-193
- 2,2'-Dihydroxydiethylamine, D-388
- 1,2-Dihydroxy-1,2-dihydroacronyline, D-417
- cis*-1,2-Dihydroxy-1,2-dihydroacronyline, A-121
- trans*-1,2-Dihydroxy-1,2-dihydroacronyline, A-121
- 3',4'-Dihydroxy-3',4'-dihydroflindersine, F-90
- 2,3-Dihydroxydihyromaleimide, D-519
- cis*-1,2-Dihydroxy-1,2-dihydromedicosmine, A-121
- 2,5-Dihydroxy-3,6-diisobutylpyrazine *N,N*-dioxide, P-788
- 1,6-Dihydroxy-3,5-dimethoxyacridone, T-243
- 2,9-Dihydroxy-1,10-dimethoxyaporphine, B-243
- 1,10-Dihydroxy-2,9-dimethoxyaporphine, B-276
- 1,11-Dihydroxy-2,10-dimethoxyaporphine, C-702
- 1,9-Dihydroxy-2,10-dimethoxyaporphine, I-196
- 1,2-Dihydroxy-9,10-dimethoxyaporphine, L-52
- 2,10-Dihydroxy-1,9-dimethoxyaporphine, L-195
- 1,9-Dihydroxy-2,3-dimethoxyaporphine, T-244
- 10,11-Dihydroxy-1,2-dimethoxyaporphine, T-248
- 2,10-Dihydroxy-1,11-dimethoxyaporphine, T-248
- 2,11-Dihydroxy-1,10-dimethoxyaporphine, T-248
- 1,10-Dihydroxy-2,11-dimethoxyaporphine, T-248
- 2,4-Dihydroxy-6,7-dimethoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, T-526
- 2,4-Dihydroxy-7,8-dimethoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, T-527
- 4,4'-Dihydroxy-3,5'-dimethoxy- α,β -3'-bicinnamic acid, D-395
- 3,3'-(6,6'-Dihydroxy-2,2'-dimethoxy-[1,1'-biphenyl]-3,3'-diyl)bis[2-propenamide], B-183
- 4,4'-Dihydroxy-2,2'-dimethoxy-3,3'-dicinnamide, B-183
- 3,11-Dihydroxy-2,10-dimethoxy-5,6-dihydroprotoberberine, P-719

- 5,8-Dihydroxy-4,7-dimethoxy-2,6-dimethylisoquinolinium(1+), D-569
- 1,6-Dihydroxy-3,5-dimethoxy-4-(2-hydroxy-3-methyl-3-butenyl)-10-methylacridone, P-616
- 1,3-Dihydroxy-5,6-dimethoxy-10-methylacridone, T-243
- 1,5-Dihydroxy-2,3-dimethoxy-10-methylacridone, T-241
- 1,5-Dihydroxy-3,4-dimethoxy-10-methylacridone, T-242
- 1,5-Dihydroxy-3,6-dimethoxy-10-methylacridone, T-243
- 1,6-Dihydroxy-3,5-dimethoxy-10-methylacridone, T-243
- 3,9-Dihydroxy-1,2-dimethoxy-7-methyldehydroaporphine, G-147
- 1,6-Dihydroxy-3,5-dimethoxy-10-methyl-4-(3-methyl-2-butenyl)-9(10*H*)-acridinone, P-616
- 1,11-Dihydroxy-2,10-dimethoxy-6-methyl-7-oxo-7*H*-dibenzo[*de,g*]quinolinium, G-93
- 2,9-Dihydroxy-1,10-dimethoxynoraporphine, B-243
- 1,10-Dihydroxy-2,9-dimethoxynoraporphine, B-276
- 1,11-Dihydroxy-2,10-dimethoxynoraporphine, C-702
- 1,9-Dihydroxy-2,10-dimethoxynoraporphine, I-196
- 2,10-Dihydroxy-1,9-dimethoxynoraporphine, L-195
- 1,9-Dihydroxy-2,3-dimethoxynoraporphine, T-244
- 10,11-Dihydroxy-1,2-dimethoxynoraporphine, T-248
- 2,10-Dihydroxy-1,11-dimethoxynoraporphine, T-248
- 2,11-Dihydroxy-1,10-dimethoxynoraporphine, T-248
- 3,9-Dihydroxy-1,2-dimethoxynoraporphine, T-244
- 1,10-Dihydroxy-2,11-dimethoxynoraporphine, T-248
- 2,9-Dihydroxy-3,8-dimethoxypavinane, A-1410
- 1,8-Dihydroxy-2,9-dimethoxypavinane, P-514
- 3,4-Dihydroxy-2-(3,4-dimethoxyphenyl)-1,5-dimethylpyrrolidine, C-557
- 2,4-Dihydroxy-7,8-dimethoxyquinoline, Q-43
- 9,10-Dihydroxy-2,3-dimethoxytetrahydroprotoberberine, D-570
- 3,9-Dihydroxy-2,10-dimethoxytetrahydroprotoberberine, I-315
- 3,4-Dihydroxy-1,2-dimethylcarbazole, D-571
- 7,9-Dihydroxy-8,10-dimethyl-2,4-dodecadienoic acid, D-572
- 6,8-Dihydroxy-3,4-dimethyl-1(2*H*)-isoquinolinone, T-544
- 3,5-Dihydroxy-2,6-dimethyl-7-(2-methyl-4-thiazolyl)-6-heptenoic acid, D-628
- 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione, T-545
- 7,19-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione, T-545
- 12,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13,19-triene-1,21-dione, H-473
- 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-triene-1,21-dione, T-545
- 7,18-Dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-triene-1,17,21-trione, Z-37
- 7,17-Dihydroxy-16,18-dimethyl-10-phenyl-22-oxa[12]cytochalasa-6(12),13,19-triene-1,21-dione, C-959
- 7,18-Dihydroxy-16,18-dimethyl-10-phenyl-22-oxa[12]cytochalasa-6(12),13,19-triene-1,21-dione, D-573
- 7,9-Dihydroxy-8,10-dimethyl-2,4-tridecadienoic acid, D-574
- 5,5'-Dihydroxydiphenylamine-2,2'-dicarboxylic acid, I-45
- 3,6-Dihydroxy-4,5-diphenyl-2-piperidinone, D-575
- 6,11-Dihydroxy-2,7,9-dodecatrienoic acid, D-576
- 8,11-Dihydroxy-2,4,9-dodecatrienoic acid, D-577
- 10,11-Dihydroxy-2,6,8-dodecatrienoic acid, D-578
- threo*-2'',3''-Dihydroxyechiumine, E-31
- 16,17-Dihydroxy-11-epilepine, L-51
- 1-(1,2-Dihydroxyethyl)- β -carboline-3-carboxylic acid, D-579
- 2-(1,2-Dihydroxyethyl)-3,4-dihydroxy-5-(hydroxymethyl)pyrrolidine, D-580
- 1-(1,2-Dihydroxyethyl)-3-hydroxymethyl- β -carbazole, P-893
- 1-(1,2-Dihydroxyethyl)-4-methoxy- β -carboline, H-493
- 1-(1,2-Dihydroxyethyl)-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, D-579
- 3-(8,9-Dihydroxyfarnesyl)indole, T-605
- 8*S*,11*S*-Dihydroxyfawcettidine, F-28
- 2*S*,11*S*-Dihydroxyfawcettidine, F-28
- 4',7-Dihydroxyflavanone, D-581
- 3 α ,13-Dihydroxyforesaconitine, M-336
- 12,13-Dihydroxyfunitremorgin C, F-190
- 4,5-Dihydroxyfuro[2,3-*b*]quinoline, F-210
- 4,6-Dihydroxyfuro[2,3-*b*]quinoline, F-211
- 4,7-Dihydroxyfuro[2,3-*b*]quinoline, F-212
- 4,8-Dihydroxyfuro[2,3-*b*]quinoline, F-213
- 5,7-Dihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-219
- 6,7-Dihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-221
- 6,8-Dihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-222
- 7,8-Dihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-223
- 14,15-Dihydroxygelsenicine, G-44
- Dihydroxygirinimbine, D-582
- 3,4-Dihydroxy- β -guanidinostyrene, T-656
- 2-(3,6-Dihydroxyheptyl)-3,4-dihydroxy-5-hydroxymethylpyrrolidine, D-583
- 13-(1,2-Dihydroxyheptyl)-1,4,5,6,7,8,9,10,11,13,16,16*a*-dodecahydroxyrido[2,1-*d*][1,5,9]triazacyclotridecin-2(3*H*)-one, C-83
- 2-(3,6-Dihydroxyheptyl)-5-hydroxymethyl-3,4-pyrrolidinediol, D-583
- 11,15-Dihydroxyhetisan-2,13-dione, H-190
- 6,13-Dihydroxyhetisan-11-one, D-584
- 9,13-Dihydroxyhetisan-2-one, D-585
- 13,15-Dihydroxyhetisan-2-one, D-586
- 6,13-Dihydroxyhetisan-2-one, H-198
- 11,13-Dihydroxyhetisan-2-one, H-202
- 2,6-Dihydroxyhetisan-13-one, H-198
- 6-(4,15-Dihydroxyhexadecyl)-3-hydroxy-2-methylpiperidine, A-477
- 3,4-Dihydroxyhydrocinnamic acid, D-652
- α,β -Dihydroxyhydrocinnamic acid, D-651
- 4,8-Dihydroxy-5-(4-hydroxybenzoyl)-2-quinolinecarboxylic acid, T-512
- 6,7-Dihydroxy-1-(4-hydroxybenzyl)isoquinoline, H-424
- 1,2-Dihydroxy-5-(3-hydroxybutyl)-3-(hydroxymethyl)-1*H*-pyrrolizidine, T-553
- 3,4-Dihydroxy-1-(2-hydroxyethyl)-2-(hydroxymethyl)pyrrolidine, D-593
- 3,4-Dihydroxy-2-(1-hydroxyethyl)-5-(hydroxymethyl)pyrrolidine, D-587
- 3,4-Dihydroxy-2-(2-hydroxyethyl)-5-(hydroxymethyl)pyrrolidine, H-492
- 3,4-Dihydroxy-2-(1-hydroxyethyl)pyrrolidine, D-588
- 3,4-Dihydroxy-2-(3-hydroxy-4-methoxyphenyl)-5-hydroxymethylpyrrolidine, R-10
- 2,5-Dihydroxy-4-(hydroxymethyl)acetanilide, A-739
- 3,4-Dihydroxy-3-hydroxymethylbutanoic acid, D-589
- 1,5-Dihydroxy-4-(2-hydroxy-3-methyl-3-butenyl)-3-methoxy-10-methylacridone, T-581
- 6,7-Dihydroxy-1-hydroxymethyl-1,2-dehydro-8 α -pyrrolizidine, C-775
- 3,4-Dihydroxy-2-hydroxymethyl-5-(1-hydroxypentyl)pyrrolidine, H-602
- 3,4-Dihydroxy-2-(hydroxymethyl)-5-(2-hydroxypropyl)pyrrolidine, D-665
- 3,4-Dihydroxy-5-hydroxymethyl-2-isobutylpyridine, A-620
- 6,7-Dihydroxy-1-hydroxymethylisoquinoline, D-590
- 3,4-Dihydroxy-2-(hydroxymethyl)-5-methylpyrrolidine, D-591
- 1,2-Dihydroxy-3-hydroxymethyl-5-methyl-1*H*-pyrrolizidine, D-592
- 3,4-Dihydroxy-5-(hydroxymethyl)-3-[3-(3-nonyloxiranyl)-1-oxo-2-propenyl]-2-pyrrolidinone, P-584
- 3-[4,5-Dihydroxy-4-(hydroxymethyl)-*L*-norvaline]phallacidin, P-309
- 7-[4,5-Dihydroxy-4-(hydroxymethyl)-*L*-norvaline]phalloidin, P-310
- 3,4-Dihydroxy-2-(hydroxymethyl)-5-(1,5,7,12,13-pentahydroxytridecyl)pyrrolidine, B-367
- 3,4-Dihydroxy-2-(hydroxymethyl)piperidine, H-614
- 3,4-Dihydroxy-2-(hydroxymethyl)pyrrolidine, D-593
- 3,4-Dihydroxy- α -(hydroxymethyl)-2,5-pyrrolidinedimethanol, D-580
- 3,4-Dihydroxy-5-(hydroxymethyl)-1-pyrrolidineethanol, H-492
- 3,4-Dihydroxy-2-hydroxymethyl-1-pyrrolidinepropanamide, D-593
- 1,2-Dihydroxy-3-hydroxymethyl-1*H*-pyrrolizidine, D-594
- 1,2-Dihydroxy-7-hydroxymethylpyrrolizidine, D-595
- 1,7-Dihydroxy-1-hydroxymethylpyrrolizidine, D-596
- 2,7-Dihydroxy-1-hydroxymethylpyrrolizidine, D-597
- 1,2-Dihydroxy-3-(hydroxymethyl)-5-(1,3,4-trihydroxybutyl)-1*H*-pyrrolizidine, T-553
- 2,4(5)-Dihydroxyimidazole, I-40
- 5,6-Dihydroxy-1*H*-indole-2-carboxylic acid, D-598
- 5,6-Dihydroxyindole, I-93
- 6,7-Dihydroxyindole, I-94
- 2,3-Dihydroxyindole, I-91
- 2,5-Dihydroxy-1*H*-indole, I-92
- α,δ -Dihydroxy-1*H*-indole-3-propanoic acid, H-506
- 3,5-Dihydroxy-2-indolinone-3-acetic acid, D-423
- 3,7-Dihydroxy-2-indolinone-3-acetic acid, D-424
- 1,2-Dihydroxyindolizidine, O-46
- 4,5-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-546
- 5,11-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-548
- 5,9-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-547
- 9,10-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-550
- 1,9-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-544
- 8,9-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-549
- 10,11-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-551
- 1,11-Dihydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-545
- 6,7-Dihydroxy-5-(isobutyliminomethyl)-8-isopropyl-2-methyl-1,4-naphthoquinone, G-145
- 4,5-Dihydroxy-*N*-isobutyl-2-octenamide, D-637
- 6,7-Dihydroxy-3-isocyano-2*H*-1-benzopyran-2-one, D-599
- 6,7-Dihydroxy-3-isocyanocoumarin, D-599
- 19,20-Dihydroxyisoechinulin A, I-230
- 6-*O*-(1,3-Dihydroxyisopropyl)inosine, I-148
- 6,7-Dihydroxyisoquinoline, I-302
- 7,8-Dihydroxyisoquinoline, I-303
- 6,7-Dihydroxy-1,3,4(2*H*)-isoquinolinetrione, D-600
- 6,7-Dihydroxy-1(2*H*)-isoquinolinone, I-305
- 6,8-Dihydroxy-1(2*H*)-isoquinolinone, I-306
- 6,9-Dihydroxy-13-labdene-16,15-lactam, D-601
- 3,18-Dihydroxylanosta-8,22,24-trien-26-oic acid, D-602
- 3,15-Dihydroxy-5,12-lathyradien-14-one, D-603
- 3-(4,5-Dihydroxy-*L*-leucine)-4-*L*-alanineviroisin, V-160
- 3-(4,5-Dihydroxy-*L*-leucine)viroisin, V-160
- 4,5-Dihydroxy-*N*^b-lignoceroyltryptamine, D-683
- N*²-(15,16-Dihydroxylinoleoyl)glutamine, G-108
- 6,7-Dihydroxyliitorine, M-392
- 12,13-Dihydroxylupanine, C-40

- 8,13-Dihydroxylupanine, D-607
 10,13-Dihydroxylupanine, D-608
 3,13-Dihydroxylupanine, D-605
 3,4-Dihydroxylupanine, D-604
 4,13-Dihydroxylupanine, D-606
 4,6 α -Dihydroxylycopodine, F-73
 7,8 β -Dihydroxylycopodine, F-29
 4,12-Dihydroxylycopodine, L-344
 18,18'-Dihydroxymatopensine, M-120
 5,9-Dihydroxymatridin-15-one, S-377
 5 α ,9 α -Dihydroxymatrine, S-377
 2,7-Dihydroxymavacurine, A-617
 3,4-Dihydroxymaytol, P-217
 1,3-Dihydroxy-5-methoxyacridone, T-518
 1,8-Dihydroxy-3-methoxyacridone, T-520
 2,7-Dihydroxy-1-methoxyaporphine, P-4
 1,10-Dihydroxy-2-methoxyaporphine, T-524
 1,11-Dihydroxy-2-methoxyaporphine, T-525
 2,4-Dihydroxy-7-methoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, D-538
 3-(2,6-Dihydroxy-4-methoxybenzoyl)pyridine, D-609
 4,10-Dihydroxy-5-methoxycanthin-6-one, T-531
 1,7-Dihydroxy-6-methoxy-9*H*-carbazole-3-carboxaldehyde, T-532
 3,4-Dihydroxy- α -(methoxycarbonyl)-*N,N,N*-trimethylbenzenethanaminium(1+), A-745
N-(3,4-Dihydroxy-5-methoxycinnamoyl)-1,4-butanediamine, C-456
 12,18-Dihydroxy-19-methoxy-16,18-dimethyl-10-phenyl[1,1]cytocalasa-6,13-diene-1,21-dione, H-473
 7,18-Dihydroxy-19-methoxy-16,18-dimethyl-10-phenyl[1,1]cytocalasa-6(12),13-diene-1,21-dione, T-545
 1,5-Dihydroxy-3-methoxy-2,4-diprenylacridone, T-546
 1,7-Dihydroxy-6-methoxyisoquinoline, I-305
 3,4-Dihydroxy-13-methoxylupanine, T-557
 2',6'-Dihydroxy-5''-methoxyluphan-12-one, V-88
 2,5-Dihydroxy-4-(methoxymethyl)acetanilide, A-739
 1,3-Dihydroxy-5-methoxy-10-methylacridone, T-518
 1,5-Dihydroxy-3-methoxy-10-methylacridone, T-518
 1,6-Dihydroxy-3-methoxy-10-methylacridone, T-519
 1,8-Dihydroxy-3-methoxy-10-methylacridone, T-520
 5,7-Dihydroxy-6-methoxy-1-methyl-4-azafluoren-9-one, T-565
 6,9-Dihydroxy-7-methoxy-3-methylbenz[*g*]isoquinoline-5,10-dione, B-255
 1,5-Dihydroxy-3-methoxy-10-methyl-2,4-diprenylacridone, T-546
 7,8-Dihydroxy-6-methoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-563
 7,9-Dihydroxy-8-methoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-565
 1,3-Dihydroxy-5-methoxy-10-methyl-2-prenylacridone, T-580
 1,5-Dihydroxy-3-methoxy-10-methyl-4-prenylacridone, T-581
 3,6-Dihydroxy-4-methoxy-1-methyl-2(1*H*)-pyridinone, H-146
 2,7-Dihydroxy-1-methoxynoraporphine, P-4
 1,10-Dihydroxy-2-methoxynoraporphine, T-524
 5,7-Dihydroxy-6-methoxyonychine, T-565
 5,7-Dihydroxy-2-[[1-(4-methoxy-2-oxo-2*H*-pyran-6-yl)-2-phenylethyl]amino]-1,4-naphthoquinone, D-610
 2,8-Dihydroxy-6-methoxy-2-pentylisoquinoline, R-163
 3,4-Dihydroxy-5-methoxyphenethylamine, T-576
 3,4-Dihydroxy-2-(4-methoxyphenyl)-1,5-dimethylpyrrolidine, C-557
 (2,6-Dihydroxy-4-methoxyphenyl)-3-pyridinylmethanone, D-609
 4,7-Dihydroxy-8-methoxyquinoline, Q-48
 2,4-Dihydroxy-6-methoxyquinoline, Q-46
 2,4-Dihydroxy-8-methoxyquinoline, Q-47
 3,12-Dihydroxy-11-methoxystrychnine, P-726
 7,18-Dihydroxy-4-methylaconit-8(15)-en-14-one, D-611
 1,3-Dihydroxy-10-methylacridone, D-528
 1,7-Dihydroxy-10-methylacridone, D-529
 1,8-Dihydroxy-10-methylacridone, D-530
 3',4'-Dihydroxy-2-methylaminoacetophenone, A-736
 3,4-Dihydroxy- α -[(methylamino)methyl]benzyl alcohol, A-152
 14,15-Dihydroxy-*N*-methylaspidospermidine, A-1498
 2,15-Dihydroxy-7-methyl-6-azabicyclo[11.3.0]hexadec-3-en-5-one, I-18
 5,6-Dihydroxy-1-methyl-4-azafluorenone, D-626
 6,7-Dihydroxy-1-methyl-4-aza-9-fluorenone, D-625
 7,8-Dihydroxy-1-methyl-4-azafluoren-9-one, D-624
 2,7-Dihydroxy-1-methyl-4-azafluorenone, D-623
 1,2-Dihydroxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridinium(1+), C-356
 1,8-Dihydroxy-3-methylbenzo[*b*]phenanthridine-7,12-dione, D-612
 8,9-Dihydroxy-4-methyl-10*H*-[1]benzopyrano[3,2-*b*]pyridin-10-one, P-638
 1,3-Dihydroxy-4-(3-methyl-2-butenyl)-9(10*H*)-acridone, D-664
 2,7-Dihydroxy-1-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-125
 2,8-Dihydroxy-1-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-125
 4,7-Dihydroxy-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, T-582
 4,8-Dihydroxy-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, T-583
 7-(2,3-Dihydroxy-3-methylbutoxy)-8-methoxydic-tamine, F-223
 8-(2,3-Dihydroxy-3-methylbutoxy)-4-methoxy-1-methyl-2(1*H*)-quinolinone, F-122
 4-(2,3-Dihydroxy-3-methylbutoxy)-1-methyl-2(1*H*)-quinolinone, H-53
N-(2,3-Dihydroxy-3-methylbutyl)acetamide, A-835
 1-(2,3-Dihydroxy-3-methylbutyl)-2,7-dihydroxy-9*H*-carbazole-3-carboxaldehyde, H-125
 3-(2,3-Dihydroxy-3-methylbutyl)-4,8-dimethoxy-1-methyl-2(1*H*)-quinolinone, L-288
 3-(2,3-Dihydroxy-3-methylbutyl)-4,7-dimethoxy-1-methyl-2(1*H*)-quinolinone, T-582
 3-(2,3-Dihydroxy-3-methylbutyl)-5-(2,3-epoxy-3-methylbutyl)-1*H*-indole, B-200
 3-(2,3-Dihydroxy-3-methylbutyl)-4-methoxy-1-methyl-2(1*H*)-quinolinone, E-40
 3-(2,3-Dihydroxy-3-methylbutyl)-6-(3-methyl-2-butenyl)-1*H*-indole, B-201
 3-(2,3-Dihydroxy-3-methylbutyl)-5-(3-methyl-1-oxo-2-butenyl)-1*H*-indole, B-200
 3-(2,3-Dihydroxy-3-methylbutyl)-4,6,8-trimethoxy-1-methyl-2(1*H*)-quinolinone, P-755
 2,7-Dihydroxy-6-methyl-9*H*-carbazole-1-carboxaldehyde, D-619
 1,6-Dihydroxy-3-methyl-9*H*-carbazole, D-613
 1,7-Dihydroxy-6-methyl-9*H*-carbazole, D-614
 2,3-Dihydroxy-6-methyl-9*H*-carbazole, D-615
 2,5-Dihydroxy-3-methyl-9*H*-carbazole, D-616
 2,6-Dihydroxy-3-methyl-9*H*-carbazole, D-617
 2,8-Dihydroxy-3-methyl-9*H*-carbazole, D-618
 6,7-Dihydroxy-1-methyl- β -carboline, D-620
 7,8-Dihydroxy-1-methyl- β -carboline, D-621
 4-(5,7-Dihydroxy-2-methyl-6-chromanyl)-2-piperidone, D-629
 3,6-Dihydroxy-1-methyl-4,5-diphenyl-2-piperidinone, D-575
 2,4-Dihydroxy-3-methylenebutanoic acid, D-622
 4,7-Dihydroxy-1,2-methylenedioxyaporphine, S-534
 7,8-Dihydroxy-2,3-methylenedioxy-5-methylbenzo[*c*]phenanthridinium(1+), C-356
 8,9-Dihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, D-626
 3,7-Dihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, D-623
 6,7-Dihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, D-624
 7,8-Dihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, D-625
 6,8-Dihydroxy-3-methylisocarbostyryl, T-566
 6,7-Dihydroxy-1-methylisoquinoline, D-627
 6,8-Dihydroxy-3-methyl-1(2*H*)-isoquinolinone, T-566
 1,3-Dihydroxy-10-methyl-4-(3-methyl-2-butenyl)-9(10*H*)-acridone, D-664
 6,7-Dihydroxy-2-methyl-5-[[3-methylbutyl]imino]methyl]-8-(1-methylethyl)-1,4-naphthalenedione, G-145
 3,5-Dihydroxy-6-methyl-7-(2-methyl-4-thiazolyl)-6-heptenoic acid, D-628
 4-(5,7-Dihydroxy-2-methyl-4-oxo-4*H*-1-benzopyran-6-yl)-1-methyl-2-piperidinone, D-629
 4-(5,7-Dihydroxy-2-methyl-4-oxo-4*H*-1-benzopyran-6-yl)-2-piperidinone, D-629
 3-(3,5-Dihydroxy-4-methyl-5-oxo-6-octenyl)-2,4-pyrrolidinedione, P-589
 4,5-Dihydroxy-1-methyl-3-oxo-2-(trichloromethyl)-3*H*-indolium(1+), D-630
 4,5-Dihydroxy-1-methyl-2-piperidinecarboxylic acid, D-661
 3,4-Dihydroxy-2-methylpiperidine, M-516
 5,7-Dihydroxy-6-(1-methylpiperidin-2-yl)flavone, C-94
 1,3-Dihydroxy-10-methyl-4-prenylacridone, D-664
 1,5-Dihydroxy-10-methyl-2-prenylacridone, T-580
 2,7-Dihydroxy-6-methyl-1-prenylcarbazole, E-275
 2,6-Dihydroxy-3-methyl-5-prenyl-9*H*-carbazole, G-110
 4,5-Dihydroxy-*N*-(2-methylpropyl)-2-decenamide, D-566
 8,11-Dihydroxy-*N*-(2-methylpropyl)-2,4,9-dodecatrienamide, D-577
 4,5-Dihydroxy-*N*-(2-methylpropyl)-2-octenamide, D-637
 2,5-Dihydroxy-3-methylpyrazine, D-631
 3,4-Dihydroxy-2-methylpyridine, D-632
 3,5-Dihydroxy-6-methyl-2(1*H*)-pyridinone, T-567
 2,4-Dihydroxy-5-methylpyrimidine, T-401
 3,4-Dihydroxy- α -methyl-2,5-pyrrolidinedimethanol, D-587
 5,7-Dihydroxy-6-(1-methylpyrrolidinyl)flavone, I-233
 5,7-Dihydroxy-6-(1-methyl-2-pyrrolidinyl)-2-phenyl-4*H*-1-benzopyran-4-one, I-233
 2,4-Dihydroxy-5-methylquinazoline, D-633
 2,3-Dihydroxy-21-methyl-6,21-secohetisan-6,13-dione, H-161
 10,10'-Dihydroxy-*N*⁺-methyltetrahydrousambar-ensine, U-52
 1,9-Dihydroxy-4-(methylthio)canthin-6-one, D-634
 1,9-Dihydroxy-4-methylthio-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-634
 1,4-Dihydroxymimosamin, M-624
 30,32-Dihydroxymycalolide A, M-779
 5,8-Dihydroxy-2-naphthalenecarboxylic acid, D-635
 5,8-Dihydroxy-2-naphthoic acid, D-635
 6,6'-Dihydroxyneothiobinapharidine, T-380
 2,4-Dihydroxynicotinic acid, D-668
 3,4-Dihydroxy-10-nitro-1-phenanthrenecarboxylic acid, D-636
 3,7-Dihydroxy-1,8-nonadiene-1,9-dicarboxylic acid, D-685
 5,13-Dihydroxynoracronycine, A-1527
 11,12-Dihydroxynorcona-*N*(18),1,4-trienin-3-one, H-322
 2,3-Dihydroxynortropane, A-1581
 2,6-Dihydroxynortropane, A-1582
 2,7-Dihydroxynortropane, A-1583
 3 β ,6 β -Dihydroxynortropane, A-1584
 3,6-Dihydroxynortropane, A-1584
 3,7-Dihydroxynortropane, A-1585
*N*²-(15,16-Dihydroxy-9,12-octadecadienyl)glutamine, G-108
 3,5-Dihydroxy-2-octanoylphenylacetic acid, C-963
 4,5-Dihydroxy-2-octenoic acid, D-637
 6,7-Dihydroxyonychine, D-625
 2,7-Dihydroxyonychine, D-623
 3,4-Dihydroxy- α -oxobenzeneacetic acid, D-649
 3,4-Dihydroxy-6-oxo-1-cyclohexene-1-acetic acid, D-638

- 3,4-Dihydroxy-6-oxo-1-cyclohexene-1-acetonitrile, D-638
 1-(4,5-Dihydroxy-1-oxo-2-decenyl)piperidine, D-566
 3,5-Dihydroxy-2-(1-oxooctyl)benzeneacetic acid, C-963
 (1,5-Dihydroxy-2-oxo-3-pyrrolidinyl)phosphonic acid, A-1256
 3,5-Dihydroxy-2-[[1-oxo-3-(3,4,5-trihydroxyphenyl)-2-propenyl]amino]benzoic acid, D-639
 4,5-Dihydroxy-*N*^b-pentacosanoyltryptamine, D-683
 4-(3,4-Dihydroxy-1-pentenyl)oxazole, M-159
 2,3-Dihydroxy-1-phenazinecarboxylic acid, D-640
 2,6-Dihydroxy-1-phenazinecarboxylic acid, D-641
 2,9-Dihydroxy-1-phenazinecarboxylic acid, D-642
 3,6-Dihydroxy-1-phenazinecarboxylic acid, D-643
 4,9-Dihydroxy-1,6-phenazinedicarboxylic acid, D-644
 1,6-Dihydroxyphenazine, P-321
 1,8-Dihydroxyphenazine, P-322
 1,9-Dihydroxyphenazine, P-323
 2,3-Dihydroxyphenazine, P-324
 Di(2-hydroxy-1-phenaziny)methane, M-440
N-(3,4-Dihydroxyphenethyl)acetamide, D-920
 3,4-Dihydroxyphenethylamine, D-920
 2-(3,4-Dihydroxyphenethyl)quinoline, D-645
 (3,4-Dihydroxyphenethyl)trimethylammonium, C-690
 5-[(3,5-Dihydroxyphenoxy)methyl]-3-pyridinol, D-238
 (2,4-Dihydroxyphenyl)acetic acid, D-646
 α,3-Dihydroxyphenylacetic acid, H-519
 α,4-Dihydroxyphenylacetic acid, H-520
 (2,4-Dihydroxyphenyl)acetonitrile, D-646
 3,4-Dihydroxyphenylalanine, A-745
 2-(3,4-Dihydroxyphenyl)-3,4-dihydro-6,8-bis(octahydro-2,4,7-trimethyl-1*H*-2-pyridin-1-yl)-2*H*-1-benzopyran-3,5,7-triol, K-81
 2-(2,3-Dihydroxyphenyl)-4,5-dihydro-*N*-hydroxy-*N*-[2-(1*H*-imidazol-4-yl)ethyl]-4-thiazolecarboxamide, A-1009
 2-(3,4-Dihydroxyphenyl)ethenamine, D-157
N-[2-(3,4-Dihydroxyphenyl)ethenyl]acetamide, D-157
N-[2-(2,5-Dihydroxyphenyl)ethenyl]formamide, E-144
 2-(3,4-Dihydroxyphenyl)ethenyl]guanidine, T-656
 9-[1-(3,4-Dihydroxyphenyl)ethyl]adenine, A-137
 2-(3,4-Dihydroxyphenyl)ethylamine, D-920
N-[2-(2,5-Dihydroxyphenyl)ethyl]formamide, E-144
 (3,4-Dihydroxyphenyl)glyoxylic acid, D-649
 3-(3,4-Dihydroxyphenyl)-2-hydroxy-2-methylpropanoic acid, D-647
 3-(3,4-Dihydroxyphenyl)-2-hydroxypropanoic acid, D-648
 3-(3,4-Dihydroxyphenyl)lactamide, D-648
 3-(3,4-Dihydroxyphenyl)lactic acid, D-648
 1-(3,4-Dihydroxyphenyl)-2-methylaminoethanol, A-152
 1-(3,4-Dihydroxyphenyl)-2-(methylamino)ethanone, A-736
 1-[(3,4-Dihydroxyphenyl)methyl]-6,7-isoquinolinediol, P-80
 2-(3,4-Dihydroxyphenyl)-2-oxoacetic acid, D-649
N-[3-(3,4-Dihydroxyphenyl)-1-oxo-2-propenyl]-3-hydroxytyrosine, C-534
 5-(3,4-Dihydroxyphenyl)-2-phenylloxazole, D-650
 2,3-Dihydroxy-3-phenylpropanoic acid, D-651
 3-(3,4-Dihydroxyphenyl)propanoic acid, D-652
 3-(3,4-Dihydroxyphenyl)-2-propenoic acid, D-653
 3-(3,5-Dihydroxyphenyl)-2-propenoic acid, D-654
 3α-(2,3-Dihydroxy-2-phenylpropionyl)-6β,7β-epoxytropolane, A-1043
 3-(3,4-Dihydroxyphenyl)-1-propylamine, A-896
N-(2,5-Dihydroxyphenyl)pyridinium(1+), D-655
 4,5-Dihydroxy-2-phenylquinoline, D-658
 4,6-Dihydroxy-2-phenylquinoline, D-659
 4,7-Dihydroxy-2-phenylquinoline, D-660
 2,3-Dihydroxy-4-phenylquinoline, D-656
 2,4-Dihydroxy-3-phenylquinoline, D-657
 7,11α-Dihydroxyphlegmariurine B, P-363
 4,5-Dihydroxyphthalic acid, D-533
 4,5-Dihydroxypipicolinic acid 4-sulfate, D-661
 4,5-Dihydroxypipicolinic acid, D-661
 4,5-Dihydroxy-2-piperidinecarboxylic acid, D-661
 3,4-Dihydroxy-2-piperidinemethanol, H-614
 5,7-Dihydroxy-8-(2-piperidinyl)flavone, D-662
 2,7-Dihydroxypleiocarpamine, A-617
 1,3-Dihydroxy-2-prenylacridone, D-663
 1,3-Dihydroxy-4-prenylacridone, D-664
 2'-*O*-*trans*-[4-(1,3-Dihydroxypropoxy)-3-methoxycinnamoyl]alaginside, A-239
 2,3-Dihydroxypropyl 1-hydroxy-4-methyl-1*H*-pyrrole-2-carboxylate, G-111
 2-(1,2-Dihydroxypropyl)-3,4-dihydroxy-5-(hydroxymethyl)pyrrolidine, D-665
 3,3-(2,3-Dihydroxypropyl)diindole, D-693
 1-(1,2-Dihydroxypropyl)-4-hydroxy-3-methoxy-2-methyl-6-prenyl-9*H*-carbazole, N-103
 6-(1,2-Dihydroxypropyl)isoxanthopterin, I-13
 5-(1,2-Dihydroxypropyl)-2-pyridinecarboxylic acid, D-666
 2,4-Dihydroxypyridine, P-762
 6,8-Dihydroxypurine, P-818
 2,6-Dihydroxypurine, X-5
 6-(6,8-Dihydroxy-9-puriny)suberosanone, D-667
 2,4-Dihydroxy-3-pyridinecarboxylic acid, D-668
 5,10-Dihydroxy-6*H*-pyrido[4,3,2-*k*]acridin-6-one, N-85
 4,8-Dihydroxypyrido[3,4-*b*]indole-1-carboxylic acid, D-561
 2,6-Dihydroxy-4-pyrimidinecarboxylic acid, D-669
 2,4-Dihydroxypyrimidine, U-42
 β-(2,4-Dihydroxy-3-pyrimidinyl)alanine, I-345
 4,5-Dihydroxy-1*H*-pyrrole-3-carboxylic acid, D-670
 3,4-Dihydroxy-2,5-pyrrolidinedimethanol, D-542
 4,6-Dihydroxyquinaldine acid, D-675
 4,7-Dihydroxyquinaldine acid, D-676
 4,8-Dihydroxyquinaldine acid, D-677
 3,4-Dihydroxyquinaldine acid, D-674
 2,4-Dihydroxyquinazoline, Q-13
 4,8-Dihydroxy-2-quinolinecarbothioic acid, D-671
 4,6-Dihydroxy-2-quinolinecarboxylic acid, D-675
 4,7-Dihydroxy-2-quinolinecarboxylic acid, D-676
 4,8-Dihydroxy-2-quinolinecarboxylic acid, D-677
 2,6-Dihydroxy-4-quinolinecarboxylic acid, D-672
 2,8-Dihydroxy-4-quinolinecarboxylic acid, D-673
 3,4-Dihydroxy-2-quinolinecarboxylic acid, D-674
 4,8-Dihydroxyquinoline, Q-35
 2,4-Dihydroxyquinoline, Q-33
 3,8-Dihydroxyquinoline, Q-34
 4,6-Dihydroxy-2(1*H*)-quinolinone, Q-46
 4,8-Dihydroxy-2(1*H*)-quinolinone, Q-47
 7,8-Dihydroxy-4(1*H*)-quinolinone, Q-48
 3,4-Dihydroxy-2(1*H*)-quinolinone, Q-45
 7,8-Dihydroxyrutaeacarpine, R-168
 1,2-Dihydroxyrutaeacarpine, R-168
 3,6-Dihydroxy-11,12-secocevan-18-one, H-370
 2,15-Dihydroxy-6,21-secohetisan-6,13-dione, S-170
 3,23-Dihydroxy-16,28-secosolanida-22,24,26(28)-trien-6-one, P-280
 6,12-Dihydroxysenecionan-11,16-dione, A-967
 12,18-Dihydroxysenecionan-11,15-dione, R-62
 4,6α-Dihydroxyserratidine, A-1029
 12,17-Dihydroxysolasodine, S-454
 17,23-Dihydroxyspermostrychnine, S-400
 3,12-Dihydroxyspirosol-5-en-27-*oic* acid, S-454
 3,11-Dihydroxystaurosporine, S-498
 12,16-Dihydroxystrychnine, P-726
 3,12-Dihydroxystrychnine, P-726
 11,19*R*-Dihydroxytaberanone, H-743
 1,11-Dihydroxy-1,6,11,16-tetraazacycloicosane-2,5,12,15-tetrone, A-250
 4,5-Dihydroxy-*N*^b-tetracosanoyltryptamine, D-683
 10,10'-Dihydroxytetrahydroousambarensine, U-52
 1,6-Dihydroxy-2,3,4,5-tetramethoxy-10-methylacridone, H-242
 4-(3,11-Dihydroxy-2,6,10,12-tetramethyl-9,13-dioxo-1,5-pentadecadienyl)-2-methylthiazole, D-678
 5,13-Dihydroxy-4,6,10,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadiene-3,7-dione, D-678
 5,13-Dihydroxy-2,4,6,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadien-3-one, D-679
 3,5'-Dihydroxythaliaboramine, T-322
syn-6,6'-Dihydroxythiobinupharidine sulfoxide, T-380
 6,6'-Dihydroxythiobinupharidine, T-380
 6,6'-Dihydroxythiobinupharidine B, T-380
 α,4-Dihydroxytoluene, H-414
 α,3-Dihydroxy-α-toluic acid, H-519
 α,4-Dihydroxy-α-toluic acid, H-520
 Dihydroxytriangularine, D-680
 4,5-Dihydroxy-*N*^b-tricosanoyltryptamine, D-683
 2,4-Dihydroxy-1,9,10-trimethoxyaporphine, C-190
 3,11-Dihydroxy-1,2,10-trimethoxyaporphine, P-209
 3,9-Dihydroxy-1,2,10-trimethoxyaporphine, P-208
 1,3-Dihydroxy-2,5,6-trimethoxy-10-methylacridone, P-206
 1,5-Dihydroxy-2,3,4-trimethoxy-10-methylacridone, P-205
 1,6-Dihydroxy-2,3,5-trimethoxy-10-methylacridone, P-206
 1,6-Dihydroxy-3,4,5-trimethoxy-10-methylacridone, P-207
 1,4-Dihydroxy-2,9,10-trimethoxynoraporphine, C-190
 1,9-Dihydroxy-2,3,10-trimethoxynoraporphine, P-208
 3,11-Dihydroxy-1,2,10-trimethoxynoraporphine, P-209
 4,14-Dihydroxy-3,6,7-trimethoxyphenanthroindolizidine, I-342
 3,4-Dihydroxy-*N,N,N*-trimethylbenzenethanaminium(1+), C-690
N-(2,4-Dihydroxy-3,5,6-trimethylbenzoyl)glycine, D-681
 4,5-Dihydroxy-*N,N,N*-trimethyl-1-hexanaminium(1+), M-772
 2,4-Dihydroxy-3,3,4-trimethyl-5-oxoproline, D-682
 2,4-Dihydroxy-3,3,4-trimethyl-5-oxo-2-pyrrolidinecarboxylic acid, D-682
 2,4-Dihydroxy-3,3,4-trimethylpyroglutamic acid, D-682
 2,6-Dihydroxytropolane, A-1582
 3,6-Dihydroxytropolane, M-391
 6,7-Dihydroxytropolane-3-yl benzoate, M-392
 4,5-Dihydroxytryptamine, D-683
 5,6-Dihydroxytryptamine, D-684
 3,5-Dihydroxy-*L*-tyrosyl-β-3,5-dihydroxy-*N*-[2-(3,4,5-trihydroxyphenyl)ethenyl]-*L*-tyrosinamide, T-678
 4,8-Dihydroxy-2,9-undecadienediamide, D-685
 4,8-Dihydroxy-2,9-undecadienedioic acid, D-685
 3,23-Dihydroxyveratra-7,12(14)-dien-6-one, T-588
 3,23-Dihydroxyveratraman-6(5*H*)-one, S-526
 18,19-Dihydroxy-ψ-vincadifformine, P-733
 19,20-Dihydroxy-ψ-vincadifformine, P-733
 4,8-Dihydroxy-1-vinyl-β-carbolone, D-686
 17,18-Dihydroxyyohimban-16-carboxylic acid, D-687
 3,17-Dihydroxyzenkerine, D-688
 Dihydrozeatin *O*-glucoside, Z-11
 Dihydrozeatin riboside, Z-11
 Dihydrozeatin, Z-11
 Diimidotricarbonic diamide, T-627
 Diindogen, I-72
 3,3-Di-1*H*-indol-3-yl-2-butanone, D-689
 4,5-Di-6-indolyl-2,7-dimethyl-2,6-octadiene, I-248
 1,5-Di-6-indolyl-3,7-dimethyl-2,6-octadiene, M-692
 Di-1*H*-indol-3-ylethanedione, H-782
 1,1-Di(3-indolyl)ethane, S-564
 2,2-Di(3-indolyl)ethyl acetate, S-564
 3,4-Di-1*H*-indol-3-yl-2,5-furandione, P-491
 4,5-Di-1*H*-indol-3-yl-2,3-furandione, P-490
 3,4-Di-1*H*-indol-3-yl-2(5*H*)-furanone, D-690
 Di-3-indolylglyoxal, H-782
 2,2-Di-3-indolyl-3-indolone, T-87
 3,3-Di-3-indolyl-2-indolone, T-88
 Di-3-indolylmethane, M-436
 2,2-Di-1*H*-indol-3-yl-4-methylpentanoic acid, D-691
 4-[(Di-1*H*-indol-3-yl)methyl]phenol, D-692

- 3,3-Di-1*H*-indol-3-yl-1,2-propanediol, D-693
 2,2-Di-1*H*-indol-3-ylpropanoic acid, D-694
 1,3-Di-1*H*-indol-3-yl-2-propanone, M-67
 3,4-Di-1*H*-indol-3-yl-1*H*-pyrrole-2,5-dicarboxylic acid, L-329
 3,4-Di-1*H*-indol-3-yl-1*H*-pyrrole-2,5-dione, A-1394
 3,5-Di-1*H*-indol-3-yl-2*H*-pyrrol-2-one, P-675
 Diiodoacetamide, D-695
 Diiodoacetic acid, D-695
 3,6-Diiodo-9*H*-carbazole, D-696
 3,5-Diiodo-4-methoxybenzenethanamine, D-697
 3-(3,5-Diiodo-4-methoxyphenyl)-2-dimethylamino-3'-(3-iodo-4-methoxyphenyl)-2'-methylamino-*N,N'*-(1,5-pentanediyloxy)bispropanamide, D-729
 (3,5-Diiodo-4-methoxyphenyl)ethylamine, D-697
 3-(3,5-Diiodo-4-methoxyphenyl)-3'-(3-iodo-4-methoxyphenyl)-*N,N'*-(1,5-pentanediyloxy)bis[2-dimethylaminopropanamide], D-729
N-[2-[3,5-Diiodo-4-(3-methyl-1-oxo-2-butenyl)amino]propoxy]phenyl]ethyl-4-hydroxybenzeneacetamide, D-333
 3,5-Diisobutyl-6-isopropyl-2-piperazinone, P-445
 2,5-Diisobutylpyrazine, B-203
 3,6-Diisobutyl-2(1*H*)-pyrazinone, F-76
 7,20-Diisocyanoadociane, D-698
 4,4'-(2,3-Diisocyanato-1,3-butadiene-1,4-diyloxy)bisphenol, X-7
 7,20-Diisocyanoisocycloamphilectane, D-698
 4-[2,3-Diisocyanato-4-(4-methoxyphenyl)-1,3-butadienyloxy]phenol, X-7
N,N-Diisopentylguanidine, B-199
N,N'-Diisopentylguanidine, P-769
 3,5-Diisopentylindole, B-200
 3,6-Diisopentylindole, B-201
 3,7-Diisopentylindole, B-202
 2,5-Diisopentyl-3,6-dimethylpyrazine, B-184
 Diisopropyl 2-aminoethyl phosphate, A-768
N,N-Diisopropylethylenediamine, E-225
 2,5-Diisopropyl-3-methoxy-pyrazine, D-699
N,N-Diisopropyl-5-methoxytryptamine, M-288
 2,5-Diisopropylpyrazine, D-700
 2,6-Diisopropylpyrazine, D-701
 Diisoretreonocanyl *p,p'*-dihydroxy- α -truxillate, T-364
Aspergillus niger Diketoterpiperazine dimer, D-702
 Diketotetrahydroglyoxaline, I-40
 Dilemmaone A, D-703
 Dilemmaone B, D-703
 Dilemmaone C, D-703
 Dilupanine, L-302
 DIMBOA, D-538
 DIMBOA-Glc, D-538
 Dimeberine chloride, C-420
 Dimethamine, D-704
 1,3-Dimethoxyacridone, D-528
 10,11-Dimethoxyajmalicine, R-51
 10,11-Dimethoxyajmalicinone, A-215
 5,10-Dimethoxyalstonine, A-231
 10,11-Dimethoxyalstonine, A-652
 2,4-Dimethoxyaniline, A-705
 Dimethoxyanthraserpine, R-50
 1,2-Dimethoxyaporphine, D-531
 11,12-Dimethoxyaspidofractinine, A-1494
 16,17-Dimethoxyaspidofractinine, A-1494
 5,6-Dimethoxy-1-azafluoroanthene, T-498
 3,4-Dimethoxybenzaldehyde *N*-(3,4-dimethoxybenzoyl)hydrazine, D-705
 4,5-Dimethoxy-1,2-benzenedicarboxylic acid, D-533
 3,4-Dimethoxybenzeneethanamine, D-718
 3,4-Dimethoxybenzenepropanamine, A-896
 9,10-Dimethoxy-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-8-one, O-170
 4,10-Dimethoxy-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-8-one, O-168
 1,10-Dimethoxybenzo[*c*]cinnoline, B-73
 1,2-Dimethoxy[1,3]benzodioxolo[5,6-*c*]phenanthridine, C-356
 2,3-Dimethoxy[1,3]benzodioxolo[5,6-*c*]phenanthridine, N-239
 2,13-Dimethoxy[1,3]benzodioxolo[5,6-*c*]phenanthridin-1-ol, C-356
 4-(4,5-Dimethoxy-1,3-benzodioxol-6-yl)-2-(5-methoxy-1,3-benzodioxol-6-yl)-1,3-cyclobutanedicarboxylic acid bispyrrolidide, B-196
 1,1'-[[2-(6,7-Dimethoxy-1,3-benzodioxol-5-yl)-4-(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidine, B-196
 1-[3-(6,7-Dimethoxy-1,3-benzodioxol-5-yl)-1-oxo-2-propenyl]pyrrolidine, P-163
 3,4-Dimethoxybenzoic acid carbonylbis(8-azabicyclo[3.2.1]octane-8,3-diyloxy) ester, S-622
 3,4-Dimethoxybenzoic acid 3,4-dimethoxybenzylidenehydrazide, D-705
 3,4-Dimethoxybenzoic acid [(3,4-dimethoxyphenyl)methylene]hydrazide, D-705
 3,4-Dimethoxybenzoic acid 1,2-ethanediylbis(8-azabicyclo[3.2.1]octane-8,3-diyloxy) ester, C-632
 3,4-Dimethoxybenzoic acid, D-706
 8,9-Dimethoxy-1*H*-benzo[*de*][1,6]naphthyridine, A-2
 8,9-Dimethoxy-6-benzopentathiepinethanamine, L-205
 6,7-Dimethoxy-4-benzotrithioethanamine, V-20
 2,7-Dimethoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, D-538
 5,6-Dimethoxy-2-benzoxazolinone, D-707
 5,6-Dimethoxy-2(3*H*)-benzoxazolone, D-707
 6,7-Dimethoxy-1(3*H*)-benzoxazolone, D-708
N-(3,4-Dimethoxybenzoyloxy)isobutylamine, D-706
N-(2,5-Dimethoxybenzoyloxy)mescaline, M-219
 8-(3,4-Dimethoxybenzoyloxy)neoline, M-323
 14-(3,4-Dimethoxybenzoyloxy)neoline, M-323
 3 α -(3,4-Dimethoxybenzoyloxy)norpropane, C-633
 15-(3,4-Dimethoxybenzoyloxy)pseudokobusine, H-205
 3,4-Dimethoxybenzoylreserpine methyl ester, R-50
 (3,4-Dimethoxybenzyl)(4-methoxyphenethyl)methylamine, N-288
 1-(3,4-Dimethoxybenzyl)-1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline, P-715
 1-(3,4-Dimethoxybenzyl)-1,2,3,4-tetrahydro-6,7-methylenedioxy-2-methylisoquinoline, R-123
N,N'-Di(4-methoxybenzyl)urea, D-302
 3,10-Dimethoxy-1,2:8,9-bis(methylenedioxy)aporphine, H-243
 4,5-Dimethoxycanthin-6-one, D-546
 5,11-Dimethoxycanthin-6-one, D-548
 5,9-Dimethoxycanthin-6-one, D-547
 9,10-Dimethoxycanthin-6-one, D-550
 1,11-Dimethoxycanthin-6-one, D-545
 2,7-Dimethoxycarbazole, C-123
 1,6-Dimethoxy-9*H*-carbazole-3-carboxaldehyde, D-552
 1,8-Dimethoxy-9*H*-carbazole-3-carboxaldehyde, D-554
 2,6-Dimethoxy-9*H*-carbazole-3-carboxaldehyde, D-555
 2,7-Dimethoxy-9*H*-carbazole-3-carboxaldehyde, D-556
 2,7-Dimethoxy-9*H*-carbazole-3-carboxylic acid, D-560
 4,8-Dimethoxy- β -carboline, D-709
 3,4-Dimethoxycinnamic acid isobutylamide, D-653
 3,5-Dimethoxycinnamic acid pyrrolidide, D-654
 3,5-Dimethoxycinnamic acid, D-654
*N*¹-(3,4-Dimethoxycinnamoyl)agmatine, A-719
N-(3,4-Dimethoxycinnamoyl)isobutylamide, D-653
N-(3,4-Dimethoxycinnamoyl)- Δ^3 -piperidin-2-one, P-473
N-(3,5-Dimethoxycinnamoyl)pyrrolidine, D-654
 5,6-Dimethoxycleistopholine, M-399
 6,7-Dimethoxycleistopholine, M-399
 Dimethoxydesantraserpidine, D-687
 1,9-Dimethoxy-8*H*-5,8-diazabenz[*cd*]azulene, A-4
 1,3-Dimethoxydibenz[*cd*]indol-4(5*H*)-one, D-568
 5,9-Dimethoxy-7*H*-dibenzo[*de,h*]quinolin-7-one, B-116
 1,2-Dimethoxydibenz[*de,g*]quinolin-7-one, L-193
 5,6-Dimethoxydictamine, F-218
 5,7-Dimethoxydictamine, F-219
 6,7-Dimethoxydictamine, F-221
 6,8-Dimethoxydictamine, F-222
 7,8-Dimethoxydictamine, F-223
 10,11-Dimethoxy-18,19-dihydro-16*S*-sitsirikine, S-325
 6,8-Dimethoxy-2,4-dihydroxyquinoline, Q-42
 6,7-Dimethoxy-1-(3,4-dimethoxybenzoyl)isoquinoline, P-77
 6,7-Dimethoxy-1-(3,4-dimethoxybenzyl)isoquinoline, P-79
 4,5-Dimethoxy-12,12-dimethyl-12*H*-[1]benzoxepino[2,3,4-*ij*]isoquinoline-6,10-diol, G-148
 1',8-Dimethoxy-3',6-dimethyl-3,9'-bi-9*H*-carbazole, M-751
 1,1'-Dimethoxy-3,3'-dimethyl-4,9'-bi-9*H*-carbazole, M-755
 7,7'-Dimethoxy-3,3'-dimethyl-8,8'-bis(3-methyl-2-butenyl)[2,2'-bi-9*H*-carbazole]-1,1'-diol, B-208
 3,4-Dimethoxy-1,2-dimethyl-9*H*-carbazole, D-571
 2,3-Dimethoxy-7,7-dimethyl-7*H*-dibenzo[*de,g*]quinoline-1,9-diol, M-194
 5,11-Dimethoxy-3,3-dimethyl-3*H*-furan[2,3-*b*]pyrano[3,2-*f*]quinoline, A-121
 5,7-Dimethoxy-*N,N*-dimethyl-1*H*-indole-3-methanamine, D-733
 1,5-Dimethoxy-*N,N*-dimethyl-1*H*-indole-3-methanamine, D-734
 6,8-Dimethoxy-1,3-dimethylisoquinoline, D-710
 6',12'-Dimethoxy-2,2'-dimethyl-6,7-[methylenebis(oxy)]oxyacanthan, C-275
 6,7-Dimethoxy-*N,N*-dimethyl-5-(methylthio)-4-benzotrithioethanamine, V-20
 3,4-Dimethoxy-*N,N*-dimethyl-1-phenanthreneethanamine, A-1532
 1,2-Dimethoxy-*N,N*-dimethylphenanthro[2,3-*dj*][1,3]dioxole-4-ethanamine, T-318
 3,16-Dimethoxy-*N,N*-dimethylpregn-5-en-20-amine, A-884
 6',7'-Dimethoxy-2,2'-dimethylthalman-6,12-diol, D-944
 4,11-Dimethoxy-1,3-dioxolo[4,5-*b*]acridin-10(5*H*)-one, X-2
 4,9-Dimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline, F-217
 9,10-Dimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline, F-215
 Dimethoxyeburnamone, E-6
 3,3-Dimethoxy-22,26-epiminocholestane, S-350
 10,11-Dimethoxy-17-epi- α -yohimbine, S-251
 4,6-Dimethoxyfuro[2,3-*b*]quinoline, F-211
 4,7-Dimethoxyfuro[2,3-*b*]quinoline, F-212
 4,5-Dimethoxyfuro[2,3-*b*]quinoline, F-210
 4,8-Dimethoxyfuro[2,3-*b*]quinoline, F-213
 4,8-Dimethoxyfuro[2,3-*b*]quinolin-5-ol, F-220
 4,6-Dimethoxyfuro[2,3-*b*]quinolin-7-ol, F-221
 4,7-Dimethoxyfuro[2,3-*b*]quinolin-6-ol, F-221
 4,7-Dimethoxyfuro[2,3-*b*]quinolin-8-ol, F-223
 4,8-Dimethoxyfuro[2,3-*b*]quinolin-7-ol, F-223
 1-[(6,7-Dimethoxyfuro[2,3-*b*]quinolin-4-yl)oxy]-3-methyl-2,3-butanediol, F-221
 8,9-Dimethoxygeibalsine, G-29
 1,4-Dimethoxyglucobrassicin, I-122
 5,7-Dimethoxygramine, D-733
 1,5-Dimethoxygramine, D-734
 3',8-Dimethoxygraveoline, G-171
 3,4-Dimethoxyhydrocinnamic isobutylamide, D-652
 2,9-Dimethoxy-1-hydroxynoraporphine (incorr.), T-523
 2,6-Dimethoxy-5-hydroxyonychine, T-562
 2,9-Dimethoxy-3-hydroxypavinane (incorr.), H-464
 3,8-Dimethoxy-9-hydroxypavinane (incorr.), H-464
 7,8-Dimethoxy-4-hydroxy-2(1*H*)-quinolinone, Q-43
 5,6-Dimethoxyindeno[1,2,3-*ij*]isoquinoline, T-498
 5,6-Dimethoxyindeno[1,2,3-*ij*]isoquinolin-9-ol, T-498
 5,6-Dimethoxy-1*H*-indole-2-carboxylic acid, D-598
 1,11-Dimethoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, D-545
 2,3-Dimethoxyindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(13*H*)-one, E-305
 Dimethoxyipomine, I-165
 5,6-Dimethoxy-1,3-isobenzofurandione, D-533

- 5,6-Dimethoxy-1*H*-isoindole-1,3(2*H*)-dione, D-533
 10,11-Dimethoxyisomitraphylline, C-117
 6,7-Dimethoxyisoquinoline, I-302
 7,8-Dimethoxyisoquinoline, I-303
 6,7-Dimethoxy-1(2*H*)-isoquinolinone, I-305
 (6,7-Dimethoxy-1-isoquinolinyl)(2-hydroxy-3,4-dimethoxyphenyl)methanone, T-50
 4-[(6,7-Dimethoxy-1-isoquinolinyl)methyl]phenol, H-424
 11,12-Dimethoxykopsine, K-76
 8,9-Dimethoxyliriodenine, O-170
 5,8-Dimethoxymaculine, F-214
 6,7-Dimethoxy-1-(4-methoxybenzyl)isoquinoline, H-424
 11,12-Dimethoxy-*N*-methoxycarbonylkopsinaline, K-74
 8,9-Dimethoxy-*N*-methoxycarbonyl-1,2-methylenedioxyornoraporphine, C-733
 4,8-Dimethoxy-1-(2-methoxyethyl)- β -carboline, P-407
 4,8-Dimethoxy-1-(2-methoxyethyl)-9*H*-pyrido[3,4-*b*]indole, P-407
 4,6-Dimethoxy-5-(3-methoxy-3-methyl-1-butenyl)furo[2,3-*b*]quinoline, H-578
 4,7-Dimethoxy-3-(methoxymethyl)-2(1*H*)-quinolinone, S-221
 6,7-Dimethoxy-4-(4-methoxyphenyl)-1-methyl-1*H*-naphth[2,3-*d*]imidazole-2,5,8(3*H*)-trione, K-25
 5,8-Dimethoxy-2-(3-methoxyphenyl)-3-propyl-4(1*H*)-quinolinone, D-711
 5,6-Dimethoxy-2-(3-methoxyphenyl)-4(1*H*)-quinolinone, D-712
 5,7-Dimethoxy-4-(4-methoxyphenyl)-2(1*H*)-quinolinone, D-713
 2,3-Dimethoxy-6-(10-methoxy-3,7,11-trimethyl-2,4,7,11-tridecatetraenyl)-5-methyl-4-pyridinol, A-257
 1,3-Dimethoxy-10-methylacridone, D-528
 3,6-Dimethoxy-9-[2-(methylamino)ethyl]dibenz[*b*,*f*]floxepin-4-ol, S-198
 7,9-Dimethoxy-3-methylbenz[*g*]isoquinoline-5,10-dione, B-255
 1,2-Dimethoxy-6-methyl-4*H*-benzo[*de*][1,3]benzodioxolo[5,6-*g*]quinoline-4,5(6*H*)-dione, C-684
 9,10-Dimethoxy-7-methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-5,6(7*H*)-dione, D-802
 5,7-Dimethoxy-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridinium(1+), C-360
 1,2-Dimethoxy-13-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridine, M-494
 1,2-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridinium(1+), C-356
 2,3-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridinium(1+), N-239
 1,2-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridin-13(12*H*)-one, D-407
 2,3-Dimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridin-13(12*H*)-one, D-482
 5,10-Dimethoxy-4-methylbenzo[*g*]quinoline, A-1058
 6,7-Dimethoxy-4-methylbenzo[*g*]quinoline-5,10-dione, M-399
 5,10-Dimethoxy-4-methylbenzo[*g*]quinolin-2(1*H*)-one, K-6
 1,1'-Dimethoxy-3'-methyl-3,9'-bi-9*H*-carbazole, M-752
 2,4-Dimethoxy-3-(3-methyl-2-butenyl)cinnamic acid piperidide, D-721
 6,8-Dimethoxy-7-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-*h*]quinoline, D-715
 4,7-Dimethoxy-8-(3-methyl-2-butenyl)furo[2,3-*b*]quinoline, H-570
 4,8-Dimethoxy-8-(3-methyl-2-butenyl)furo[2,3-*b*]quinolin-7(8*H*)-one, P-254
 4,6-Dimethoxy-8-(3-methyl-2-butenyloxy)furo[2,3-*b*]quinoline, F-222
 4,8-Dimethoxy-7-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline, F-223
 1-[3-[2,4-Dimethoxy-3-(3-methyl-2-butenyl)phenyl]-1-oxo-2-propenyl]piperidine, D-721
 4,8-Dimethoxy-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, T-583
 2,7-Dimethoxy-6-methyl-9*H*-carbazole-1-carboxaldehyde, D-619
 2,6-Dimethoxy-3-methyl-9*H*-carbazole, D-617
 1,6-Dimethoxy-3-methyl-9*H*-carbazole, D-613
 1,7-Dimethoxy-6-methyl-9*H*-carbazole, D-614
 2,5-Dimethoxy-3-methyl-9*H*-carbazole, D-616
 6,7-Dimethoxy-3-methyl-1*H*-carbazole-1,4(9*H*)-dione, K-55
 6,7-Dimethoxy-3-methyl-9*H*-carbazol-1-ol, T-559
 6,7-Dimethoxy-1-methyl- β -carboline, D-620
 10,11-Dimethoxy-1-methyldeacetylpicraline benzoate, B-420
 10,11-Dimethoxy-1-methyldeacetylpicraline 3,4,5-trimethoxybenzoate, B-420
 10,11-Dimethoxy-1-methyldeacetylpicraline, B-420
 1,2-Dimethoxy-6-methyl-4*H*-dibenzo[*de*,*g*]quinoline-4,5(6*H*)-dione, C-273
 1,2-Dimethoxy-7-methyl-4*H*-dibenzo[*de*,*g*]quinoline-4,5(6*H*)-dione, G-175
 6,11-Dimethoxy-5-methyl-1,3-dioxo[4,5-*b*]acridin-10(5*H*)-one, T-56
 4,5-Dimethoxy-11-methyl-1,3-dioxolo[4,5-*c*]acridin-6(11*H*)-one, M-169
 8,9-Dimethoxy-13-methyl-1,3-dioxolo[6,7]indeno[2,1-*a*][3]benzazepinium(1+), L-15
 1,4-Dimethoxy-2,3-methylenedioxy-9(10*H*)-acridone, X-2
 10,11-Dimethoxy-1,2-methylenedioxyaporphine, B-397
 3,9-Dimethoxy-1,2-methylenedioxyaporphine, B-471
 8,9-Dimethoxy-1,2-methylenedioxyaporphine, C-733
 9,10-Dimethoxy-1,2-methylenedioxyaporphine, D-346
 1,2-Dimethoxy-9,10-methylenedioxyaporphine, N-29
 7,9-Dimethoxy-2,3-methylenedioxybenzo[*c*]phenanthridine, D-714
 8,9-Dimethoxy-2,3-methylenedioxybenzo[*c*]phenanthridine, N-239
 8,9-Dimethoxy-2,3-methylenedioxybenzo[*c*]phenanthridin-6(5*H*)-one, D-482
 9,10-Dimethoxy-2,3-(methylenedioxy)berberine, C-73
 9,10-Dimethoxy-2,3-(methylenedioxy)berbin-13-ol, O-94
 2,3-Dimethoxy-4,5-methylenedioxycinnamic acid pyrrolidide, P-163
 1-[2,3-Dimethoxy-4,5-(methylenedioxy)cinnamoyl]pyrrolidine, P-163
 3,4-Dimethoxy-11,12-methylenedioxy-8,14-dioxohexahydroisohomoprotoberberine, P-801
 4,5-Dimethoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline, F-215
 4,8-Dimethoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline, F-217
 6,7-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-265
 6,8-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-266
 7,8-Dimethoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-267
 10,11-Dimethoxy-1,2-methylenedioxyornoraporphine, B-397
 3,9-Dimethoxy-1,2-methylenedioxyornoraporphine, B-471
 9,10-Dimethoxy-1,2-methylenedioxyornoraporphine, D-346
 9,11-Dimethoxy-1,2-methylenedioxyornoraporphine, D-846
 1,2-Dimethoxy-10,11-methylenedioxyornoraporphine, L-10
 1,2-Dimethoxy-9,10-methylenedioxyornoraporphine, N-29
 3,11-Dimethoxy-1,2-methylenedioxyornoraporphine, T-245
 1,2-Dimethoxy-9,10-methylenedioxyoxoaporphine, O-195
 2,3-Dimethoxy-8,9-methylenedioxyapavane, E-220
 6,8-Dimethoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid, M-263
 2,4-Dimethoxy-7,8-methylenedioxy-3-prenylquinoline, D-715
 4,7-Dimethoxy-2,3-methylenedioxyquinoline, Q-40
 6,7-Dimethoxy-*N*-methylflindersine, F-90
 7,8-Dimethoxy-*N*-methylflindersine, F-90
 4,5-Dimethoxy-11-methylfuro[2,3-*c*]acridin-6(11*H*)-one, T-351
 6,8-Dimethoxy-9-methylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-222
 5,7-Dimethoxy-9-methylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-219
 7,8-Dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, D-625
 2,3-Dimethoxy-14-methylindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(14*H*)-one, E-309
 6,7-Dimethoxy-2-methylisocarboxystyryl, I-305
 5,6-Dimethoxy-2-methyl-1*H*-isoindole-1,3(2*H*)-dione, D-533
 6,7-Dimethoxy-1-methylisoquinoline, D-627
 6,7-Dimethoxy-2-methyl-1,3,4(2*H*)-isoquinoline-trione, D-600
 6,7-Dimethoxy-2-methylisoquinolinium(1+), I-302
 6,7-Dimethoxy-2-methyl-1(2*H*)-isoquinolinone, I-305
 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, T-583
 2,3-Dimethoxy-16-methyl-10,11-[methylenebis(oxy)]rheadan-8-ol, G-87
 1,2-Dimethoxy-10-methyl-3,4-methylenedioxyacridone, M-169
 1,5-Dimethoxy-10-methyl-2,3-methylenedioxyacridone, T-56
 7,8-Dimethoxy-10-methyl-2',3'-methylenedioxy-1,2-benzophenanthridinium(1+), C-356
 2,3-Dimethoxy-13-methyl-9,10-(methylenedioxy)berberine, C-222
 6,7-Dimethoxy-2-methyl-1-(3,4-methylenedioxyphenyl)isoquinolinium(1+), C-796
 4,8-Dimethoxy-1-methyl-6,7-methylenedioxy-3-prenyl-2(1*H*)-quinolinone, P-750
 2,6-Dimethoxy-3-methyl-5-(1-methylethyl)pyrazine, I-272
 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-oxo-3-butenyl)-2(1*H*)-quinolinone, T-583
 4,8-Dimethoxy-1-methyl-3-(3-methyl-2-oxobutyl)-2(1*H*)-quinolinone, T-583
 7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-3,4-dihydro-6,8-dimethoxy-1,3-dimethylisoquinoline, A-985
 7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-3,4-dihydro-6-methoxy-1,3-dimethyl-8-isoquinolinol, A-981
 5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2-dihydro-8-methoxy-3-methyl-6-isoquinolinol, A-987
 5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,2,3-trimethylisoquinoline, A-982
 7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-1,3-dimethyl-8-isoquinolinol, D-793
 7-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-8-methoxy-1,3-dimethyl-6-isoquinolinol, A-990
 5-(4,5-Dimethoxy-2-methyl-1-naphthalenyl)-1,2,3,4-tetrahydro-8-methoxy-1,3-dimethyl-6-isoquinolinol, A-982
 4,8-Dimethoxy-3-(3-methyl-2-oxo-3-butenyl)-2(1*H*)-quinolinone, T-583
 3,4-Dimethoxy-*N*-methylphenethylamine, D-718
 3,6-Dimethoxy-1-methyl-2-phenyl-4(1*H*)-quinolinone, D-716
 5,6-Dimethoxy-*N*-methylphthalimide, D-533
 10,11-Dimethoxy-1-methylpicraline, B-420
 1-(Dimethoxymethyl)piperidine, P-455
 6,7-Dimethoxy-1-methyl-9*H*-pyrido[3,4-*b*]indole, D-620
 4,6-Dimethoxy-1-methyl-2(1*H*)-quinolinone, Q-46
 3,4-Dimethoxy-1-methyl-2(1*H*)-quinolinone, Q-45
 4,8-Dimethoxy-1-methyl-2(1*H*)-quinolinone, Q-47
 10,10'-Dimethoxy-*N*^{4'}-methyltetrahydrousambarensine, U-52

- 10,11-Dimethoxymitraphylline, C-117
 7,8-Dimethoxymyrtopside, M-804
 5,8-Dimethoxy-2-naphthalenecarboxamide, D-635
 10,11-Dimethoxynareline, N-47
 10,12-Dimethoxynareline, N-47
 8,9-Dimethoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, T-267
 8,10-Dimethoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, T-266
 1,2-Dimethoxynoraporphine, D-531
 6,7-Dimethoxyonchicine, D-625
 3,4-Dimethoxy- α -oxobenzeneacetic acid, D-649
 3-(1,1-Dimethoxy-2-oxoethoxy)-9,10-dihydro-1-methoxy-9-oxo-4-acridinecarboxaldehyde, D-717
 7,7'-Dimethoxyperonatin B, P-271
 2-(3,4-Dimethoxyphenacyl)pyrrolidine, R-165
N-(1,7-Dimethoxy-2-phenanthrenyl)acetamide, A-743
 1,6-Dimethoxyphenazine, P-321
 1,8-Dimethoxyphenazine, P-322
 1,9-Dimethoxyphenazine, P-323
 2,3-Dimethoxyphenazine, P-324
 3,4-Dimethoxyphenethylamine, D-718
 2-(3,4-Dimethoxyphenethyl)quinoline, D-645
 1,2-Dimethoxy-3*H*-phenoxazin-3-one, T-599
 6-(3,4-Dimethoxyphenyl)-3,4-dihydro-4-oxobenzoc[2,7]naphthridinium(1+), P-269
 2-(2,3-Dimethoxyphenyl)-5,8-dimethoxy-3-propyl-4(1*H*)-quinolinone, D-711
 2-(3,4-Dimethoxyphenyl)-5,6-dimethoxy-4(1*H*)-quinolinone, D-712
 1-(3,4-Dimethoxyphenyl)-2-dimethylaminoethanol, M-18
 4-(3,4-Dimethoxyphenyl)-4-[2-(dimethylamino)ethyl]-2-cyclohexen-1-ol, J-50
 4-(3,4-Dimethoxyphenyl)-4-[2-(dimethylamino)ethyl]-2-cyclohexen-1-one, J-49
 2-(3,4-Dimethoxyphenyl)-1,5-dimethyl-3,4-pyrroli-dinediol, C-557
N-[2-(2,4-Dimethoxyphenyl)ethyl]-3-pyridine-carboxamide, D-719
N-(3,4-Dimethoxyphenylethyl)cinnamide, L-87
N-[2-(3,4-Dimethoxyphenyl)ethyl]-3,4-dimethoxybenzeneacetamide, H-351
 2-[(3,4-Dimethoxyphenyl)ethyl]guanidine, T-656
 2-[2-(3,4-Dimethoxyphenyl)ethyl]-4-hydroxyquinoline, G-11
 4-[2-(3,4-Dimethoxyphenyl)ethyl]-5-methyl-1*H*-imidazole-2-butanol, C-935
 2-[2-(3,4-Dimethoxyphenyl)ethyl]quinoline, D-645
 1-[2-(3,4-Dimethoxyphenyl)ethyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, H-338
 8-(3,4-Dimethoxyphenyl)-1,3,4,6,9,9a-hexahydro-7-(4-methoxyphenyl)-2*H*-quinolizine, J-54
 3*a*-(3,4-Dimethoxyphenyl)-2,3,3*a*,6,7,7*a*-hexahydro-1-methyl-1*H*-indol-6-ol, M-224
 3*a*-(3,4-Dimethoxyphenyl)-1,2,3,3*a*,4,5-hexahydro-1-methyl-6*H*-indol-6-one, M-225
 3*a*-(3,4-Dimethoxyphenyl)-2,3,3*a*,4,5,9*b*-hexahydro-1-methyl-1*H*-pyrrolo[2,3-*f*]quinoline, A-346
 2-(2,3-Dimethoxyphenyl)-4-hydroxy-5,8-dimethoxy-3-propylquinoline, D-711
 2-(3,4-Dimethoxyphenyl)-4-hydroxy-5,6-dimethoxyquinoline, D-712
N-[2-(3,4-Dimethoxyphenyl)-2-hydroxyethyl]benzamide, M-18
N-[2-(3,4-Dimethoxyphenyl)-2-hydroxyethyl]-*N*-methylformamide, M-18
 3-(3,4-Dimethoxyphenyl)-*N*-[4-[[imino(3-methyl-2-butenyl)amino]methyl]amino]butyl]-2-propanamide, C-107
N-[2-(3,4-Dimethoxyphenyl)-2-methoxyethyl]benzamide, M-18
 2-(3,4-Dimethoxyphenyl)-3-(4-methoxyphenyl)quinolizidine, K-23
 4-(3,4-Dimethoxyphenyl)-4-[2-(methylamino)ethyl]-2-cyclohexen-1-ol, J-50
 4-[2-[[3,4-Dimethoxyphenyl)methyl]amino]ethyl]phenol, N-288
 7-(3,4-Dimethoxyphenyl)-7-(2-methylaminoethyl)-5,6,7,8-tetrahydroquinoline, T-433
N-[(3,4-Dimethoxyphenyl)methyl]-4-methoxy-*N*-methylbenzeneethanamine, N-288
 1-[[3,4-Dimethoxyphenyl)methyl]-1,2,3,4-tetrahydro-7-methoxy-2-methyl-6-isoquinolinol, P-715
 5-[[3,4-Dimethoxyphenyl)methyl]-5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinoline, R-123
 4-[6-(3,4-Dimethoxyphenyl)-2-morpholinyl]phenol, C-363
 3*a*-(3,4-Dimethoxyphenyl)octahydro-1-methyl-6*H*-indol-6-one, M-226
 4-(3,4-Dimethoxyphenyl)octahydro-2*H*-quinolizine-2-ol, L-53
 1-[3-(3,4-Dimethoxyphenyl)-1-oxopropyl]-1*H*-pyrrole, P-351
 5-(3,4-Dimethoxyphenyl)-2-phenyloxazole, D-650
 3-(3,4-Dimethoxyphenyl)propanamide, D-652
 3-(3,4-Dimethoxyphenyl)propanoic acid 2-methylpropylamide, D-652
 3-(3,4-Dimethoxyphenyl)propanoic acid pyrrolidide, P-351
 1-[3-(3,4-Dimethoxyphenyl)propanoyl]-5,6-dihydro-2(1*H*)-pyridinone, P-473
N-[3-(3,4-Dimethoxyphenyl)propanoyl]pyrrole, P-351
 3-(3,4-Dimethoxyphenyl)propylamine, A-896
 1-(3,4-Dimethoxyphenyl)-2-(2-pyrrolidinyl)ethanone, R-165
 4,6-Dimethoxy-2-phenylquinoline, D-659
 1-(3,4-Dimethoxyphenyl)-1,2,3,4-tetrahydro-6,7-dimethoxy-2-methylisoquinoline, C-797
 6-(3,4-Dimethoxyphenyl)-5,6,7,8-tetrahydro-*N*-methyl-6-quinolineethanamine, T-433
 4,5-Dimethoxyphthalic acid, D-533
 10,11-Dimethoxypicraphylline, P-404
 3,4-Dimethoxy- ω -(2-piperidyl)acetophenone, P-522
 7,8-Dimethoxyplatydesmine, D-720
 (2,4-Dimethoxy-3-prenylcinnamoyl)piperidine, D-721
 4,6-Dimethoxy-7-prenyloxyfuro[2,3-*b*]quinoline, F-221
 4,7-Dimethoxy-6-prenyloxyfuro[2,3-*b*]quinoline, F-221
 4,7-Dimethoxy-8-prenyloxyfuro[2,3-*b*]quinoline, F-223
 4,8-Dimethoxy-9*H*-pyrido[3,4-*b*]indole, D-709
 β ,4-Dimethoxy-9*H*-pyrido[3,4-*b*]indole-1-ethanol, H-560
 3-(4,8-Dimethoxy-9*H*-pyrido[3,4-*b*]indol-1-yl)-1-(9*H*-pyrido[3,4-*b*]indol-1-yl)-1-propanone, P-406
 2,3-Dimethoxypyrido[2,3-*b*]pyrazine, D-722
 3',4'-Dimethoxy-2-pyrrolidin-2-yl-acetophenone, R-165
 7,8-Dimethoxyquestinocin A, A-862
 2,4-Dimethoxyquinazoline, Q-13
 6,8-Dimethoxy-2,4-quinolinediol, Q-42
 7,8-Dimethoxy-2,4-quinolinediol, Q-43
 2,4-Dimethoxyquinoline, Q-33
 7,8-Dimethoxy-4-quinolinol, Q-48
 1,4-Dimethoxy-2(1*H*)-quinolinone, Q-33
 4,8-Dimethoxy-2(1*H*)-quinolinone, Q-47
 7,8-Dimethoxy-4(1*H*)-quinolinone, Q-48
 3,4-Dimethoxy-2(1*H*)-quinolinone, Q-45
 4,9-Dimethoxysampangine, S-49
 Dimethoxyscytonemin, T-279
 3,3-Dimethoxy-16,28-secosolanidane, S-350
 5,10-Dimethoxystrictamine, S-576
 10,11-Dimethoxystrychnine, S-589
 10,11-Dimethoxystrychnobrasiline, S-594
N,N'-[1-(3,4-Dimethoxystyryl)-2-butenylene]diforamidine, A-25
N-(2,4-Dimethoxystyryl)-3-pyridinecarboxamide, D-719
 6,7-Dimethoxy-3-(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)-1(3*H*)-isobenzofuranone, C-644
 6,7-Dimethoxy-3-(5,6,7,8-tetrahydro-4-methoxy-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)-1(3*H*)-isobenzofuranone, N-44
 10,10'-Dimethoxytetrahydroisambarensine, U-52
 2,3-Dimethoxy-1,5,6-trihydroxy-10-methylacridone, P-206
 5,6-Dimethoxy-2-(2,3,6-trimethoxyphenyl)-4(1*H*)-quinolinone, D-712
 10,11-Dimethoxytubulosan-8'-ol, T-675
 3,5-Dimethoxytyramine, T-576
 10,11-Dimethoxyuncarine D, C-117
 10,11-Dimethoxyuncarine F, C-117
 3,3'-Dimethoxyurabaine, U-41
 4,7-Dimethoxy-1-vinyl- β -carboline, D-723
 4,8-Dimethoxy-1-vinyl- β -carboline, D-686
 4,9-Dimethoxy-1-vinyl- β -carboline, D-724
 10,11-Dimethoxy- α -yohimbine, S-251
 Dimethyl *N*²-creatininylphosphate, D-725
 Dimethyl 4,9-dihydroxy-1,6-phenazinedicarboxylate, D-644
 Dimethyl 6,6'-(1,2-dimethyl-1,2-ethanediy)bis[1-phenazinecarboxylate], P-331
 Dimethyl (dithiodiethylene)dycarbamate, P-622
 Dimethyl lycogalate A, L-329
 Dimethyl lycogalate B, L-329
 Dimethyl lycogalate C, L-329
 Dimethyl 21-oxo-11,21-cycloaspidopermidine-1,2-dicarboxylate, M-296
 Dimethyl (E)-rhoifolate, R-88
 Dimethyl (Z)-rhoifolate, R-88
 3-(β , β -Dimethylacryloylamino)-20-dimethylaminopregnane, P-601
 β , β -Dimethylacrylylcholine, C-419
O- β , β -Dimethylacrylylfuntuphyllamine C, A-881
N,O-Dimethylactinodaphnine, D-346
*N*¹,*N*¹-Dimethylageliferin, A-190
 7,8-Dimethylalloxazine, L-285
 6-(γ , γ -Dimethylallyl)amino)purine, Z-11
 8-(1,1-Dimethylallyl)confusameline, D-771
 3-(3,3-Dimethylallyl)-4-(3,3-dimethylallyloxy)-2-quinolone, A-1529
 3-(1,1-Dimethylallyl)-3-(3,3-dimethylallyl)-1,2,3,4-tetrahydro-2,4-quinolinedione, B-377
O-(3,3-Dimethylallyl)halfordinol, H-12
 3-(γ , γ -Dimethylallyl)indole, M-411
 8-(Dimethylallyl)-7-methoxydictamine, H-570
 8-(3,3-Dimethylallyloxy)-7-methoxy-*N*-methylflindersine, F-90
 8-(3,3-Dimethylallyloxy)-*N*-methylflindersine, F-90
N-2-[4-(3,3-Dimethylallyloxy)phenyl]ethylcinnamide, H-667
 α,α -Dimethylallylpaspalinine, A-169
 3-(3,3-Dimethylallyl)-4,7,8-trimethoxy-2(1*H*)-quinolinone, P-625
 4-(3,3-Dimethylallyl)tryptophan, M-419
 Dimethylamine, D-726
 2-(*N,N*-Dimethylamino)acetophenone, A-694
 1-[(Dimethylamino)acetyl]-2,3,3*a*,13,14,15 α -hexahydro-13-(1-methylpropyl)-5,8-ethenopyrrolo[3,2-*b*][1,5,8]oxadiazacyclotetradecene-12,15(1*H*,11*H*)-dione, N-332
 3-Dimethylaminoacetyl-5-methoxyindole, D-727
N-[2-(Dimethylamino)benzoyl]glycyl-*N*-(2-carboxyphenyl)-*N*²-methylvalinamide, V-154
 4-(Dimethylamino)butanoic acid, A-715
 4-(Dimethylamino)-1-butanol, A-716
N'-[4-(Dimethylamino)butyl]-*N,N*-dimethyl-1,4-butanediamine, S-340
N-[4-(Dimethylamino)butyl]-*p*-hydroxy-*N*-methylcinnamide, C-456
N-[4-(Dimethylamino)butyl]-3-(4-hydroxyphenyl)-*N*-methyl-2-propenamide, C-456
N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methyl-2,4-dodecadienamide, T-609
N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methylododecanamide, T-609
N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-3-methyl-2-dodecenenamide, T-609
N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-5-methyl-2,4-tetradecadienamide, T-610
N-[3-[[4-(Dimethylamino)butyl]methylamino]propyl]-5-methyl-3-tetradecenenamide, T-611
 3-Dimethylamino-5-conanene, C-601
 3-Dimethylaminoconan-5-ol, H-315
 3-(Dimethylamino)con-5-enine, C-601
 3-Dimethylaminocon-5-enin-12-ol, H-318

- 2-(Dimethylamino)-1,5-dihydro-5-(1*H*-indol-3-yl)-5-(2-oxopropyl)-4*H*-imidazol-4-one, D-728
- 17-(Dimethylamino)-3,6-dihydroxy-4-methyl-15-oxaandrost-7-ene-4-carboxylic acid, I-11
- α -(Dimethylamino)-3,5-diiodo-*N*-[5-[[3-(3-iodo-4-methoxyphenyl)-2-(methylamino)-1-oxopropyl]amino]pentyl]-4-methoxybenzenepropanamide, D-729
- 20-(Dimethylamino)-3,16-dimethoxypregn-5-ene, A-884
- 3-(Dimethylamino)-4,14-dimethyl-9,19-cyclopregn-16-en-20-one, C-882
- 1-(Dimethylamino)-5,12-dimethyl-5,9-diazaheneicos-11-en-10-one, T-609
- 2-(Dimethylamino)-*N*-[5,8-dioxo-3-phenyl-7-(phenylmethyl)-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]-3-methylbutanamide, I-155
- 2-(Dimethylamino)ethanol, D-730
- 3-(Dimethylamino)-5-ethenyl-5-hydroxy-2-cyclopenten-1-one, A-824
- 4-[4-[2-(Dimethylamino)ethoxy]phenyl]-4-[2-(dimethylamino)ethyl]cyclohexanone, J-49
- 2-(Dimethylamino)ethyl benzoate, D-730
- 2-Dimethylaminoethyl cinnamate, D-730
- 4-[2-(Dimethylamino)ethyl]-1,2-benzenediol, D-920
- α -[1-(Dimethylamino)ethyl]benzenemethanol, D-738
- 6-[2-(Dimethylamino)ethyl]-1,3-benzodioxole-5-carboxylic acid, F-175
- 5-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]acetyl]-1,3-benzodioxole-4-carboxylic acid, A-143
- 6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]carbonyl]furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, N-51
- 7-[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]-6,7-dihydro-9*H*-1,3-dioxolo[4,5-*h*][2]benzopyran-9-one, P-276
- [6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl](3-hydroxy-4-methoxyphenyl)ethanedione, C-793
- 6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methyl]-6,7-dihydro-6-hydroxy-8*H*-1,3-dioxolo[4,5-*e*]isoindol-8-one, F-191
- 6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]-6,7-dihydro-8*H*-1,3-dioxolo[4,5-*e*]isoindol-8-one, F-172
- 3-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]-6,7-dimethoxy-1(3*H*)-isobenzofuranone, M-472
- 6-[[6-[2-(Dimethylamino)ethyl]-1,3-benzodioxol-5-yl]methylene]furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, A-144
- 2-(Dimethylaminoethyl)-3',6-dihydroxy-4',5-dimethoxystilbene, C-729
- 1-[2-(Dimethylamino)ethyl]-3,8-dimethoxy[1]benzopyrano[5,4,3-*cde*][1]benzopyran-5,10-dione, T-36
- 9-[2-(Dimethylamino)ethyl]-3,6-dimethoxydibenz[*b,h*]oxepin-4-ol, S-198
- 1-(2-Dimethylaminoethyl)-3,4-dimethoxy-6,7-methylenedioxyphenanthrene, T-318
- 1-Dimethylaminoethyl-3,4-dimethoxyphenanthrene, A-1532
- 5-[2-(Dimethylamino)ethyl]-2,3-dimethoxyphenol, H-465
- 5-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]acetyl]-1,3-benzodioxole-4-carboxylic acid, B-120
- 7-[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]-6*H*-indenol[4,5-*d*]-1,3-dioxole-6,8(7*H*)-dione, D-217
- 6-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]methylene]-6,7-dihydro-8*H*-1,3-dioxolo[4,5-*e*]isoindol-8-one, F-171
- 6-[[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]methylene]furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, A-142
- 5-[2-[2-[2-(Dimethylamino)ethyl]-4,5-dimethoxyphenyl]-2-oxoethyl]-1,3-dioxole-4-carboxylic acid, B-120
- 3-(2-Dimethylaminoethyl)-5-hydroxyindole, B-393
- 3-(2-Dimethylaminoethyl)-4-hydroxyindole, H-480
- 1-(2-Dimethylaminoethyl)-3-hydroxy-4-methoxyphenanthrene, A-1532
- 3-[2-(Dimethylamino)ethyl]-2-[2-(3-hydroxy-4-methoxyphenyl)ethenyl]-6-methoxyphenol, C-729
- 4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)-2,5-cyclohexadien-1-one, J-49
- 4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)-cyclohexanone, J-49
- 4-[2-(Dimethylamino)ethyl]-4-(4-hydroxyphenyl)-2-cyclohexen-1-one, J-49
- 4-(2-Dimethylaminoethyl)-2-(4-hydroxyphenyl)-7-methoxybenzofuran, S-197
- 4-[2-(Dimethylamino)ethyl]imidazole, D-731
- 3-(2-Dimethylaminoethyl)indole, D-785
- 3-(2-Dimethylaminoethyl)-1*H*-indol-5-ol, B-393
- 3-(2-Dimethylaminoethyl)-1*H*-indol-4-ol, H-480
- 3-(2-Dimethylaminoethyl)-2*H*-indol-2-one, H-754
- 6-[[6-[2-(Dimethylamino)ethyl]-4-methoxy-1,3-benzodioxol-5-yl]acetyl]-2,3-dimethoxybenzoic acid, N-35
- 3-[[6-[2-(Dimethylamino)ethyl]-4-methoxy-1,3-benzodioxol-5-yl]methylene]-2,3-dihydro-6,7-dimethoxy-1*H*-isoindol-1-one, N-36
- 4-[4-[2-(Dimethylamino)ethyl]-7-methoxy-2-benzofuranyl]phenol, S-197
- 3-[2-(Dimethylamino)ethyl]-1-methoxy-10*H*-[1,3]dioxolo[6,7]phenanthro[4,5-*bcd*]pyran-10-one, T-327
- 3-[2-(Dimethylamino)ethyl]-1-methoxyindole, D-785
- 3-[2-(Dimethylamino)ethyl]-6-methoxy-2-[2-(4-methoxyphenyl)ethenyl]phenol, L-96
- 1-(2-Dimethylaminoethyl)-6-methoxy-3,4-methylenedioxyphenanthrene, I-343
- 1-(2-Dimethylaminoethyl)-7-methoxy-3,4-methylenedioxyphenanthrene, U-64
- 1-[2-(Dimethylamino)ethyl]-4-methoxy-3-phenanthrenol, A-1532
- 4-[2-(Dimethylamino)ethyl]-2-methoxyphenol, H-566
- 4-[2-(Dimethylamino)ethyl]-4-(4-methoxyphenyl)-cyclohexanone, J-49
- 1-[2-(Dimethylamino)ethyl]-4-methoxy-9*H*-pyrido[3,4-*b*]indol-8-ol, P-412
- 2-[2-(Dimethylamino)ethyl]-4,5-methylenedioxybenzoic acid, F-175
- 1-(2-Dimethylaminoethyl)-3,4-methylenedioxyphenanthrene, S-546
- 3-(2-Dimethylaminoethyl)oxindole, H-754
- 4-(2-Dimethylaminoethyl)phenol, H-357
- 2-Dimethylaminoethylphosphonic acid, A-773
- 1-(2-Dimethylaminoethyl)-3,4,6,7-tetramethoxyphenanthrene, M-559
- 1-(2-Dimethylaminoethyl)-3,4,7,8-tetramethoxyphenanthrene, U-63
- 1-(2-Dimethylaminoethyl)-2,3,4-trimethoxy-6,7-methylenedioxyphenanthrene, T-328
- 1-(2-Dimethylaminoethyl)-2,3,4-trimethoxyphenanthrene, M-238
- 4-(Dimethylamino)-1,4,4a,5,12,12a-hexahydro-3,10,11,12a-tetrahydroxy-6-methyl-1,12-dioxo-2-naphthacene-carboxamide, A-1034
- 4-[2-(Dimethylamino)-1-hydroxyethyl]-1,2-benzenediol, A-152
- 20-(Dimethylamino)-16-hydroxy-3-(3-isopropylacetam)pregnan-4-one, P-8
- 20-(Dimethylamino)-16-hydroxy-3-(3-isopropyl-2-oxo-1-azetidiny)pregnan-4-one, P-8
- 20-(Dimethylamino)-16-hydroxy-3-(3-isopropyl-2-oxo-1-azetidiny)pregn-2-en-4-one, P-8
- 2-(Dimethylamino)-3-hydroxy-*N*-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]butanamide, N-331
- 2-(Dimethylamino)-1-(4-hydroxyphenyl)-8,10-dimethyl-6-dodecen-3-one, A-1321
- 20-(Dimethylamino)-2-hydroxy-3-phthalimidopregnan-4-yl acetate, D-262
- 20-(Dimethylamino)-18-hydroxypregna-1,4-dien-3-one, A-810
- 3-(Dimethylamino)-2-hydroxypregn-5-en-18,20-olide, L-30
- 3-(Dimethylamino)-2-hydroxypregn-5-en-16-one, A-813
- 3-(Dimethylamino)-2-hydroxypregn-5-en-20-one, A-814
- 2-Dimethylamino-6-hydroxypurine, G-210
- 20-(Dimethylamino)-16-hydroxy-4,4,14-trimethyl-9,19-cyclopregnan-3-one, C-884
- 3-(Dimethylamino)-5-hydroxy-5-vinyl-2-cyclopenten-1-one, A-824
- 20-(Dimethylamino)-18-ikemaoyl-3-(methylamino)pregn-5-ene-15,18-diol, D-270
- 2-[[[(Dimethylamino)iminomethyl]amino]ethanesulfonic acid, A-1522
- 2-(Dimethylamino)-5-(1*H*-indol-3-yl)-4*H*-imidazol-4-one, D-732
- 2-(Dimethylamino)-*N*-[7-(1*H*-indol-3-ylmethyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]-3-methylbutanamide, I-156
- 2-(Dimethylamino)-*N*-[7-(1*H*-indol-3-ylmethyl)-3-(1-methylethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-4-methylpentanamide, T-300
- 4-(Dimethylamino)-6-(1*H*-indol-3-yl)-2*H*-1,3,5-oxadiazin-2-one, A-245
- 20-(Dimethylamino)-3-(3-isopropyl-2-oxo-1-azetidiny)pregn-2-en-4-one, P-8
- 17-Dimethylaminolobohedleolide (incorr.), A-723
- 2-(Dimethylamino)-1-(5-methoxy-1*H*-indol-3-yl)ethanone, D-727
- 12-(Dimethylamino)-15-methoxy-9-(1-methylpropyl)-6-(2-methylpropyl)-4,7,10-triazabicyclo[12.3.1]octadeca-1(18),2,14,16-tetraene-5,8,11-trione, M-719
- 12-(Dimethylamino)-15-methoxy-9-(1-methylpropyl)-6-(phenylmethyl)-4,7,10-triazabicyclo[12.3.1]octadeca-1(18),2,14,16-tetraene-5,8,11-trione, M-718
- 20-(Dimethylamino)-3-methoxypregna-5,16-diene, A-876
- 20-(Dimethylamino)-3-methoxypregn-5-en-16-ol, A-884
- 20-Dimethylamino-3-methylaminopregnane, P-601
- 20-(Dimethylamino)-3-(methylamino)pregnan-4-ol, D-266
- 20-(Dimethylamino)-3-(methylamino)pregn-5-ene-16,18-diol, D-271
- 20-(Dimethylamino)-3-(methylamino)pregn-14-ene, P-607
- 20-(Dimethylamino)-3-(methylamino)pregn-5-en-15-ol, D-278
- α -[(Dimethylamino)methyl]benzenemethanol, A-864
- α -[(Dimethylamino)methyl]benzyl alcohol, A-864
- 20-Dimethylamino-14-methyl-9,19-cyclopregnan-3-one, B-459
- α -[(Dimethylamino)methyl]-3,4-dimethoxybenzenemethanol, M-18
- 3-[(Dimethylamino)methyl]-1,5-dimethoxyindole, D-734
- 3-[(Dimethylamino)methyl]-5,7-dimethoxyindole, D-733
- 3-(Dimethylaminomethyl)-1-(1,1-dimethyl-2-propenyl)-1*H*-indole, D-736
- 3-(Dimethylaminomethyl)-5-ethoxyindole, D-734
- α -(Dimethylamino)-*N*-[3-(1-methylethyl)-7-(2-methylpropyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-1*H*-indole-3-propanamide, N-334
- 3-[(Dimethylamino)methyl]-5-hydroxyindole, D-734
- 3-[(Dimethylamino)methyl]-7-hydroxyindole, D-735
- 3-[(Dimethylamino)methyl]indole, D-736
- 3-[(Dimethylamino)methyl]-1*H*-indol-5-ol, D-734
- 3-[(Dimethylamino)methyl]-1*H*-indol-7-ol, D-735
- 3-(Dimethylaminomethyl)-5-methoxyindole, D-734
- 3-[(Dimethylamino)methyl]-7-methoxyindole, D-735

- 3-(Dimethylamino)-14-methyl-4-methylene-9,19-cyclopregn-17(20)-en-16-one, C-912
- 3-(Dimethylamino)-14-methyl-4-methylene-9,19-cyclopregn-16-en-20-one, C-882
- 2-(Dimethylamino)-3-methyl-*N*-[7-(1-methylethyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-pentanamide, S-92
- 2-(Dimethylamino)-3-methyl-*N*-[3-(1-methylethyl)-7-(2-methylpropyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-pentanamide, F-151
- 2-(Dimethylamino)-3-methyl-*N*-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,16-tetraen-4-yl]-pentanamide, I-154
- 2-(Dimethylamino)-4-methyl-*N*-[7-(1-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]-pentanamide, M-793
- 3-(Dimethylaminomethyl)phenol, H-416
- 1-(Dimethylamino)-2-methyl-2-propanol, A-840
- α -(Dimethylamino)-*N*-[7-(2-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]benzenepropanamide, C-229
- 3-(Dimethylaminomethyl)trimethoxyindole, D-734
- α -(Dimethylamino)-*N*-(1,2,11,12,14,15,16,16 α -octahydro-1,12-dioxo-10-phenyl-5,8-etheno-10*H*-pyrrolo[1,2-*e*][1,5,8]oxadiazacyclotetradecin-11-yl)benzenepropanamide, C-91
- 1-[2-(Dimethylamino)-1-oxo-3-phenylpropyl]-2,3,3 α ,13 α ,14,15,16,18 α -octahydro-8-methoxy-5,9-metheno-9*H*-dipyrrolo[3,2-*b*:1',2'-*e*][1,5,8]oxadiazacyclopentadecine-13,18(1*H*,12*H*)-dione, D-737
- 1-[2-(Dimethylamino)-1-oxo-3-phenylpropyl]-1,2,3,3 α ,12,14,15,16 α -octahydro-8-methoxy-14-(2-methylpropyl)-5,9-metheno-9*H*-pyrrolo[3,2-*b*][1,5,8]oxadiazacyclopentadecine-13,16-dione, N-330
- 20-(Dimethylamino)-3-oxopregna-1,4-dien-18-al, A-810
- 3-[2-(Dimethylamino)phenyl]-2,3-dihydro-3-hydroxy-5-methoxypyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, A-1039
- 3-[2-(Dimethylamino)phenyl]-2,3-dihydropyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, V-40
- 2-Dimethylamino-1-phenylethanol, A-864
- 3-(Dimethylamino)-1-(2-phenylethyl)-2,6-piperidinedione, P-385
- 1-(1-Dimethylamino-2-phenylethyl)-9*H*-pyrido[3,4-*b*]indol-6-ol, E-284
- 7-[4-(Dimethylamino)phenyl]-*N*-hydroxy-4,6-dimethyl-7-oxo-2,4-heptadienamide, T-496
- 2-Dimethylamino-1-phenyl-1-propanol, D-738
- 3-(2-Dimethylaminophenyl)-1,2,3,9-tetrahydropyrrolo[2,1-*b*]quinazoline, V-40
- 3-(Dimethylamino)pregna-5,17(20)-diene-2,16-diol, A-873
- 20-(Dimethylamino)pregna-1,4-dien-3-one, A-878
- N*-[3-(Dimethylamino)pregna-14,16-dien-20-yl]-*N*-methylformamide, P-600
- 20-(Dimethylamino)pregnan-3-ol, A-881
- 20-(Dimethylamino)pregnan-3-one, A-883
- 3-(Dimethylamino)pregnan-20-one, A-882
- 20-Dimethylaminopregn-5-en-3-ol, A-886
- 20-Dimethylaminopregn-4-en-3-one, A-891
- 3-(Dimethylamino)pregn-17(20)-en-16-one, A-890
- 3-Dimethylaminopregn-16-en-20-one, A-889
- 3-Dimethylaminopregn-5-en-20-one, A-888
- N*'-[3-(Dimethylamino)propyl]-*N,N*-dimethyl-1,3-propanediamine, D-255
- 3-(3-Dimethylaminopropyl)-1(3*H*)-isobenzofuranone, D-739
- 3-(3-Dimethylaminopropyl)phthalide, D-739
- N*-[3-(Dimethylamino)propyl]-*N,N,N'*-trimethyl-1,3-propanediamine, D-255
- 20-(Dimethylamino)-2',3,3',4-tetrahydro-3'-methylpregn-3-eno[3,4-*d*]oxazole, A-612
- 2-Dimethylamino-6-(1,2,3-trihydroxypropyl)-4(3*H*)-pteridinone, A-923
- 20-(Dimethylamino)-4,4,14-trimethyl-9,19-cyclopregn-1-en-3-one, C-850
- Dimethylammonium chloride, D-726
- O,N*-Dimethylancistrocladine, A-982
- N,N*-Dimethylanomurine, A-1069
- N*-[4-(Dimethylarsinoyl)butanoyl]aminoethylsulfonic acid, D-740
- N*-[4-(Dimethylarsinoyl)butanoyl]taurine, D-740
- Di-*O*-methylasterriquinol D, A-1519
- 3,5-Di-*O*-methylatalaphylline, T-546
- 1,9-Dimethyl-9-azabicyclo[3.3.1]nonan-3-one, M-389
- 3,8-Dimethyl-2-azabicyclo[2.2.2]oct-2-en-5-one, M-139
- 4,4-Dimethyl-15-aza-*D*-homoergosta-8,14,24(28)-trien-3-ol, D-742
- 6,6-Dimethyl-2-azaspiro[4.4]non-1-ene, D-741
- 4,4-Dimethyl-15-azasterol, D-742
- 7,8-Dimethylbenzo[*g*]pteridine-2,4(1*H*,3*H*)-dione, L-285
- (2,2-Dimethyl-2*H*-1-benzopyran-6-yl)ethanamine, A-770
- N*-[2-(2,2-Dimethyl-2*H*-1-benzopyran-6-yl)ethyl]benzamide, A-770
- N*-[2-(2,2-Dimethyl-2*H*-1-benzopyran-6-yl)ethyl]-*N*-methyl-3-pyridinecarboxamide, A-770
- N*-[2-(2,2-Dimethyl-2*H*-1-benzopyran-6-yl)ethyl]-3-pyridinecarboxamide, A-770
- 3,3'-Dimethyl-[1,1'-bi-9*H*-carbazole]-2,2'-diol, B-128
- 3,3'-Dimethyl-[2,2'-bi-1*H*-carbazole]-1,1',4,4'(9*H*,9'*H*)-tetrone, B-213
- 1,1'-Dimethyl-2,2'-bipyridinium(2+), B-144
- 1,1'-Dimethyl-3,3'-bipyridinium(2+), D-743
- N,N'*-Dimethyl-3,3'-bipyridyl(2+), D-743
- 1,1'-Dimethyl-2,3'-bipyrridine, B-147
- 1-(1,1'-Dimethyl[2,3'-bipyrrolidin]-2'-yl)-2-propanone, M-548
- 1-(1,1'-Dimethyl[2,3'-bipyrrolidin]-5'-yl)-2-propanone, M-549
- 5,5'-Dimethyl-11,11'-bi-5*H*-quindoline, B-168
- 2,5-Dimethyl-3,6-bis(3-methylbutanoyl)pyrazine, B-184
- 2,5-Dimethyl-3,6-bis(3-methylbutyl)pyrazine, B-184
- 3,6-Dimethyl- α,α' -bis(2-methylpropyl)-2,5-pyrazinemethanol, B-184
- N,N'*-Dimethyl-1,4-butanediamine, B-422
- N,N*-Dimethyl-1,4-butanediamine, B-422
- 2,7-Dimethyl-5,10-[2]butenophenazine-11,14-dione, D-744
- 1,2-Dimethyl-9*H*-carbazole-3,4-diol, D-571
- 1,2-Dimethyl-9*H*-carbazole-3,4,6-triol, T-543
- N,N*-Dimethylclaitine, A-773
- N,O*-Dimethylcosoline, I-339
- N,N*-Dimethylcon-5-enin-3-amine, C-601
- N,N*-Dimethyl-(+)-coniine, P-652
- 3,10-Di-*O*-methylconstrictosine, C-624
- O,O*-Dimethylcoreximine, X-30
- O,O*-Dimethylcortyuberine, T-248
- N,O*-Dimethylcrotsparine, P-643
- O,O*-Dimethyl(-)-curine, C-812
- 2,2'-Dimethylcycleaninium, C-835
- 4,14-Dimethyl-9,19-cyclopregnane-3,20-diamine, B-474
- N,N'*-Dimethylcyclovirobuxine D, C-917
- 6,6'-Di-*O*-methylauricoline, D-88
- O,O*-Dimethyl-1',2'-dehydronorkohatane, K-56
- 2,9-Dimethyl-2,9-diazatricyclo[10.2.2.2^{3,8}]octadeca-5,7,12,14,15,17-hexaene-3,10-diol, D-745
- 1,9-Dimethyl-3,9-dihydro-1*H*-purine-2,6-dione, X-5
- 3,9-Dimethyl-3,9-dihydro-1*H*-purine-2,6-dione, X-5
- Dimethyldihydroarifeite, W-9
- N,N*-Dimethyl-3,4-dihydroxyphenethylamine, D-920
- N,N*-Dimethyl-3,4-dimethoxyphenethylamine, D-718
- 2,6-Dimethyldinicotinic acid, D-776
- 1,3-Dimethyl-2,6-dioxopyrimidine, D-777
- N,N'*-Dimethyl-*N,N'*-diphenylurea, D-818
- 4,6-Dimethyl-2,4-dodecadienamide, D-746
- 4,6-Dimethyl-2,4-dodecadienoic acid, D-746
- N*^b-(3,5-Dimethyl-2,4-dodecadienoyl)tryptamine, T-639
- N,N*-Dimethyl-dopamine, D-920
- O,O*-Dimethyl-dopamine, D-718
- Di(α -methylene- γ -butyrolactonyl)amine, I-46
- Dimethylenedicytisine, T-362
- 6,14-Di-*O*-methyl-6-epiforeosticine, M-323
- 6,8-Dimethylergolin-9-ol, F-181
- N*-Dimethyl ethanolamine, D-730
- 4-(1,1-Dimethylethyl)-2-oxazolidinone, B-440
- Dimethylformamide, D-747
- N,N*-Dimethylfuntumine, A-882
- O,O*-Dimethylgovanine, X-30
- O,O*-Dimethylgrisabine, G-177
- 1,3-Dimethylguanidine, D-746
- N*²,*N*²-Dimethylguanidine, G-210
- 1,7-Dimethylguanidine, M-467
- 1,9-Dimethylguanidine, M-467
- 3,7-Dimethylguanidine, M-468
- 3,9-Dimethylguanidine, M-468
- 7,9-Dimethylguaninium, H-143
- N*-Dimethylheliamine, T-158
- 7-[2'-(1,6-Dimethyl-2,4-heptadienyl)[2,4'-bithiazol]-4-yl]-3,5-dimethoxy-4-methyl-2,6-heptadienamide, M-807
- N,O*-Dimethylhernovine, T-248
- 4-(1,5-Dimethyl-1,3-hexadienyl)-1-isocyano-1-methylcyclohexane, I-213
- N*²,*N*²-Dimethylhistamine, D-731
- Dimethylholamine, A-888
- Dimethylholaphyllamine, A-888
- Dimethylhuperzine A, H-375
- N,N*-Dimethyl-3-hydroxy-4,5-dimethoxyphenethylamine, H-465
- Dimethyl-2-hydroxyethylamine, D-730
- 1,7-Dimethylhypoxanthine, P-820
- 1,9-Dimethylhypoxanthine, P-820
- 3,7-Dimethylhypoxanthine, P-820
- 3,9-Dimethylhypoxanthine, P-820
- 1,3-Dimethyl-4-imidazoleacetic acid betaine, Z-32
- 1,3-Dimethyl-4-imidazolecarboxylic acid betaine, N-318
- 4,5-Dimethylimidazole, D-750
- 2,4(5)-Dimethylimidazole, D-749
- N,N*-Dimethyl-1*H*-imidazole-4-ethanamine, D-731
- 1,4-Dimethyl-1*H*-imidazole, M-473
- 1,5-Dimethyl-1*H*-imidazole, M-473
- 2,4-Dimethyl-2-imidazole, D-428
- 1,9-Dimethyl-6-imino-8-oxopurine, T-194
- 1,7-Dimethyl-1*H*-indole-3-carboxaldehyde, M-475
- 2,4-Dimethyl-1*H*-indole, D-751
- N,N*-Dimethyl-1*H*-indole-3-ethanamine, D-785
- 1,3-Dimethyl-1*H*-indole, M-474
- N,N*-Dimethyl-1*H*-indole-3-methanamine, D-736
- 5,7-Dimethylindolizidine, O-42
- N*¹,*N*¹-Dimethylisogaelferlin, A-190
- 4,4'-Dimethylisoboraverine, I-198
- 7,7'-*O,O*-Dimethylisochondodendrine, C-835
- N,O*-Dimethylisocorydine, T-248
- 1,3-Dimethylisoguanine, A-818
- 3,7-Dimethylisoguanine, A-818
- 1,3-Dimethylisoguaninium, A-818
- N,N*-Dimethyl-L-isoleucyl-*N*-[7-(1*H*-indol-3-ylmethyl)-3,3-dimethyl-5,8,11-trioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-12,14,15-trien-4-yl]-L-valinamide, H-763
- O,N*-Dimethylisopiline, T-521
- 3,4-Dimethyl-1,6,8-isoquinolinetrial, T-544
- N,N*-Dimethylisocarpine, T-248
- N,N'*-Dimethylindoldhamine, D-88
- O,N*-Dimethyliriadendronine, L-193
- O,O*-Dimethylmagloferine, T-248
- N,N*-Dimethylmethanamine *N*-oxide, T-601
- N,N*-Dimethylmethanamine, T-600
- 2,5-Dimethyl-6-methoxy-4,7-isoindoquinone, M-248
- N,N*-Dimethyl-4-methoxyphenethylamine, M-280
- N,N*-Dimethyl-3-methoxytyramine, H-566
- 3,4-Dimethyl-7-(methylamino)-5,8-isoquinoline-dione, D-752
- 3,4-Dimethyl-7-(methylamino)-5,8-isoquinoline-quinone, D-752

- N,N'*-Dimethyl-*N*-[3-(methylamino)propyl]-1,3-propanediamine, D-255
- 2,5-Dimethyl-3-(3-methylbutanoyl)-6-(3-methyl-2-butenyl)pyrazine, B-184
- 3,6-Dimethyl-1-(3-methyl-2-butenyl)-9*H*-carbazole, D-753
- 2,5-Dimethyl-3-(3-methyl-1-butenyl)-6-(3-methylbutyl)pyrazine, B-184
- 2,5-Dimethyl-3-(3-methyl-2-butenyl)-6-(3-methylbutyl)pyrazine, B-184
- 2,5-Dimethyl-3-(3-methyl-2-butenyl)pyrazine, D-754
- 2,5-Dimethyl-3-(3-methylbutyl)pyrazine, D-754
- 2,5-Dimethyl-3-(2-methylbutyl)pyrazine, D-755
- α,α -Dimethyl-6-(1-methylethyl)pyrazinemethanol, D-701
- α,α -Dimethyl-5-(3-methyl-1-oxo-2-butenyl)-1*H*-indole-3-propanal, D-756
- 2,5-Dimethyl-3-(2-methylpropyl)pyrazine, D-757
- 2,2-Dimethyl-5-[5-methyl-6-(1*H*-pyrrol-2-yl)-1,3,5-hexatrienyl]-3(2*H*)-furanone, W-5
- 2,5-Dimethyl-3-(methylthio)pyrazine, D-758
- N,N*-Dimethyl-5-(methylthio)varacin, L-205
- N,O*-Dimethylmicranthine, M-591
- O,O*-Dimethylmunitagine, P-514
- N*¹,*N*³-Dimethylnaamine D, N-6
- N,O*-Dimethylnandigerine, B-397
- α -(4,8-Dimethyl-3,7-nonadienyl)-1,2-dihydro- α ,4-dimethylfuro[2,3-*c*]quinoline-2-methanol 5-oxide, A-1541
- 3-(4,8-Dimethyl-3,7-nonadienyl)-3,5-dimethyl-3*H*-pyrano[2,3-*c*]quinoline 6-oxide, A-1545
- α -(4,8-Dimethyl-1,3,7-nonatrienyl)-2,3-dihydro- α ,4-dimethylfuro[2,3-*c*]quinoline-2-methanol 5-oxide, A-1546
- N*,2-Dimethyl-6-nonylpiperidine, M-492
- O,N*-Dimethylnorbelladine, N-288
- 6-(3,7-Dimethyl-2,6-octadienyl)-2,3-dihydro-7-hydroxy-5-methoxy-2-(2-phenylethyl)-1*H*-isoindol-1-one, H-145
- 5-(3,7-Dimethyl-2,6-octadienyl)-2,5-dihydro-2-oxo-3-phenylfuro[2,3-*b*]phenazine-11-carboxylic acid, B-64
- 5-(3,7-Dimethyl-2,6-octadienyl)-2,5-dihydro-2-oxo-3-phenylfuro[2,3-*b*]phenazine-9-carboxylic acid, B-64
- 2-(3,7-Dimethyl-2,5-octadienyl)-4,6-dihydroxy-5-methyl-3-propylpyridine, I-179
- 2-(3,7-Dimethyl-2,6-octadienyl)-1,3-dimethyl-4(1*H*)-quinolinone, D-760
- 1-(3,7-Dimethyl-2,6-octadienyl)-2-hydroxy-9*H*-carbazole-3-carboxaldehyde, M-729
- 8-(3,7-Dimethyl-2,6-octadienyl)-7-hydroxy-3-methyl-1*H*-carbazole-1,4(9*H*)-dione, M-767
- 6-(3,7-Dimethyl-2,5-octadienyl)-4-hydroxy-3-methyl-5-propyl-2(1*H*)-pyridinone, I-179
- 2-(3,7-Dimethyl-2,6-octadienyl)-4-hydroxy-3-methylquinoline, D-760
- 2-(3,7-Dimethyl-2,6-octadienyl)-4-hydroxyquinoline, D-762
- 3-(3,7-Dimethyl-2,6-octadienyl)-1*H*-indole, D-759
- 8-(3,7-Dimethyl-2,6-octadienyl)-7-methoxy-3-methyl-1*H*-carbazole-1,4(9*H*)-dione, M-767
- 1-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-9*H*-carbazole-2,7-diol, E-277
- 1-(3,7-Dimethyl-2,6-octadienyl)-6-methyl-9*H*-carbazole-2,7-diol, E-276
- 4-(3,7-Dimethyl-2,6-octadienyl)-6-methyl-9*H*-carbazol-3-ol, E-302
- 2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-1-(methylthiomethyl)-4(1*H*)-quinolinone, D-760
- 2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-4-quinolinol, D-760
- 2-(3,7-Dimethyl-2,6-octadienyl)-1-methyl-4(1*H*)-quinolinone, D-762
- 2-(3,7-Dimethyl-2,6-octadienyl)-3-methyl-4(1*H*)-quinolinone, D-760
- 4-(3,7-Dimethyl-2,6-octadienyl)-1*H*-pyrrole-2-carboxylic acid, D-761
- 2-(3,7-Dimethyl-2,6-octadienyl)-4(1*H*)-quinolinone, D-762
- 2-(3,7-Dimethyl-2,6-octadienyl)-1,3,5-trihydroxy-10-methyl-9(10*H*)-acridinone, G-121
- 6-(2,5-Dimethyloctylidene)octahydro-8-methyl-7,8-indolizinediol, A-629
- α -(3,3-Dimethylloxiranyl)-1*H*-indole-7-methanol, M-414
- 4-(3,3-Dimethylloxiranyl)-2-methylindolo[2,3-*a*]quinolizine, H-88
- 2-[5-[(3,3-Dimethylloxiranyl)methyl]-1*H*-indol-3-yl]-3-methyl-1,3-butanediol, E-134
- 3-[[3,3-Dimethylloxiranyl)methyl]-6-(3-methyl-2-butenyl)-1*H*-indole, B-201
- N*⁶,9-Dimethyl-8-oxoadenine, A-819
- 4,8-Dimethyl-15-oxo-16-aza-18-norandrost-13-ene-4-carboxylic acid, D-763
- 4,8-Dimethyl-17-oxo-16-aza-18-norandrost-13-ene-4-carboxylic acid, D-764
- N*⁹-(1,1-Dimethyl-3-oxobutyl)adenine, G-15
- 3-[1-(1,1-Dimethyl-3-oxobutyl)imidazol-4-yl]-2-propenoic acid, U-46
- 4-[2-(3,5-Dimethyl-2-oxocyclohexyl)-2-hydroxyethyl]-2,6-piperidinedione, D-765
- 3-(6,8-Dimethyl-1-oxo-2,4,6-decatrienyl)-1,4-dihydroxy-5-(4-hydroxyphenyl)-2(1*H*)-pyridinone, B-28
- 3-(8,10-Dimethyl-1-oxo-2,4,6,8-dodecetraenyl)-4-hydroxy-5-(4-hydroxyphenyl)-2(1*H*)-pyridinone, F-10
- 1,7-Dimethyl-8-oxo-9*H*-guanine, A-753
- α ,5-Dimethyl-2-oxo-4-imidazolidinedihexanoic acid, M-432
- 1,3-Dimethyl-8-oxoisoguanine, A-754
- 3-(4,6-Dimethyl-1-oxo-2,4-octadienyl)-1,4-dihydroxy-5-(4-hydroxyphenyl)-2(1*H*)-pyridinone, T-72
- 4-[3-(3,5-Dimethyl-6-oxo-2-piperidinylidene)-2-oxopropyl]-2,6-piperidinedione, E-92
- 4-[[2,2-Dimethyl-1-oxopropyl]amino]butanoic acid, A-715
- 1,3-Dimethyl-6-(1-oxopropyl)-2,4(1*H*,3*H*)-pteridinedione, H-702
- O,N*-Dimethylpapaverrubine D, G-87
- 7,12-Di-*O*-methylpedroamine, D-88
- 3,5-Dimethyl-2-pentylpyrazine, D-766
- N,N*-Dimethylphenanthro[3,4-*d*]-1,3-dioxole-5-ethanamine, S-546
- N,N*-Dimethylphenethylamine, P-342
- N,N*-Dimethylphenylalanyl-*N*-[5,8-dioxo-3-phenyl-7-(phenylmethyl)-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]prolinamide, A-151
- N,N*-Dimethyl-1-phenylethanolamine, A-864
- N*,3-Dimethyl-*N*-(2-phenylethenyl)butanamide, T-303
- 2,5-Dimethyl-3-(2-phenylethenyl)pyrazine, D-767
- 3,4-Dimethyl-5-phenyl-2-oxazolidinone, M-511
- 2,6-Dimethyl-4-phenyl-3,5-pyridinedicarboxylic acid, D-768
- N,N'*-Dimethylphlegmarine, P-361
- N,O*-Dimethylphylline, T-177
- Di-*O*-methylphyllostemine β -*N*-oxide, S-248
- 3,6-Dimethyl-2,5-piperazinedione, D-769
- 2,6-Dimethylpiperidine, D-770
- α ,6-Dimethyl-2-piperidinediol, H-705
- 4-[2-(1,6-Dimethyl-2-piperidinyl)ethenyl]decahydro-3-methylnaphtho[2,3-*c*]furan-1(3*H*)-one, H-280
- 21-(1,5-Dimethyl-2-piperidinyl)pregn-20-ene-3,6-diol, S-275
- 20-(1,5-Dimethyl-2-piperidinyl)pregn-5-ene-3,16,18-triol, A-1073
- O,N*-Dimethylporphyrone, G-87
- 3,6-Dimethyl-1-prenyl-9*H*-carbazole, D-753
- 2,5-Dimethyl-3-prenylpyrazine, D-754
- 2-[2-(1,1-Dimethyl-2-propenyl)-1*H*-indol-3-yl]-5-(1*H*-indol-3-yl)-3,6-dimethoxy-1,4-benzenediol, A-1519
- 1-(1,1-Dimethyl-2-propenyl)-3-[4-(1*H*-indol-3-yl)-2,3,5,6-tetramethoxyphenyl]-1*H*-indole, A-1519
- 8-(1,1-Dimethyl-2-propenyl)-4-methoxyfuro[2,3-*b*]quinolin-7-ol, D-771
- 3-(1,1-Dimethyl-2-propenyl)-5-(3-methyl-2-butenyl)-1*H*-indole, C-216
- 3-[[2-(1,1-Dimethyl-2-propenyl)-5-(3-methyl-2-butenyl)-1*H*-indol-3-yl]methylene]-6-methyl-2,5-piperazinedione, I-230
- 3-(1,1-Dimethyl-2-propenyl)-3-(3-methyl-2-butenyl)-2,4(1*H*,3*H*)-quinolinedione, B-377
- 3-(1,1-Dimethyl-2-propenyl)-5-(3-oxo-1-butenyl)-1*H*-indole, C-217
- 3-(1,1-Dimethyl-2-propenyl)-5-prenylindole, C-216
- 1,3-Dimethyl-6-propionylumazine, H-702
- 2,5-Dimethyl-3-propylpyrazine, D-772
- 3,5-Dimethyl-2-propylpyrazine, D-773
- N*²,*N*⁸-Dimethylpseudozoanthoxanthin A, P-737
- N*⁸,*N*⁸-Dimethylpseudozoanthoxanthin A, P-737
- 1,*N*⁸-Dimethylpseudozoanthoxanthin A, P-737
- Dimethylpseudozoanthoxanthin, P-737
- 6,7-Dimethyl-7*H*-purine, M-535
- 6,9-Dimethyl-9*H*-purine, M-535
- 1,6-Dimethyl-1*H*-purine, M-535
- 2',2'-Dimethyl(pyran-5',6':3,4)-1,5-dihydroxy-6-methoxy-10-methylacridone, C-472
- 2,5-Dimethylpyrazine, D-774
- 3,5-Dimethyl-2-pyrazinol, H-474
- 3,6-Dimethyl-2-pyrazinol, H-475
- 3,5-Dimethyl-2(1*H*)-pyrazinone, H-474
- 3,6-Dimethyl-2(1*H*)-pyrazinone, H-475
- 2,6-Dimethylpyridine, D-775
- 2,6-Dimethyl-3,5-pyridinedicarboxylic acid, D-776
- α ,6-Dimethyl-3-pyridinemethanamine, B-248
- 1,2-Dimethylpyridinium iodide, M-537
- 5,11-Dimethyl-6*H*-pyrido[4,3-*b*]carbazole, E-67
- 1,5-Dimethyl-6*H*-pyrido[4,3-*b*]carbazole, O-83
- N*,1-Dimethyl-9*H*-pyrido[3,4-*b*]indole-4-butanamine, B-291
- N*,2-Dimethyl-5-pyrimidinedicarboxamide, M-539
- 1,3-Dimethyl-2,4(1*H*,3*H*)-pyrimidinedione, D-777
- 6,8-Dimethylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(6*H*,8*H*)-dione, F-48
- 1,6-Dimethylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-dione, R-66
- 2,4-Dimethyl-1*H*-pyrrole-3-carboxylic acid, D-778
- 1,3-Dimethyl-1*H*-pyrrole-2,5-dione, M-542
- β ,3-Dimethyl-1-pyrrolidinedebutanol, M-484
- α ,1-Dimethyl-2-pyrrolidinedebutanol, H-760
- 1,3-Dimethylpyrrolidine, M-544
- 2-(3,4-Dimethyl-1*H*-pyrrol-1-yl)-3-methylbutanoic acid, D-779
- 2-(3,4-Dimethyl-1*H*-pyrrol-1-yl)propanoic acid, D-780
- 2,4-Dimethylquinazoline, D-781
- 1,3-Dimethyl-2,4(1*H*,3*H*)-quinazolinone, Q-13
- 2,2-Dimethyl-4(1*H*,3*H*)-quinazolinone, D-782
- 1,2-Dimethyl-4(1*H*)-quinazolinone, H-632
- 4,8-Dimethyl-6-quinolinol, H-478
- 5,7-Dimethyl-6-quinolinol, H-479
- 2,3-Dimethyl-4-quinolinol, H-477
- 2,3-Dimethyl-4(1*H*)-quinolinone, H-477
- 1,2-Dimethyl-4(1*H*)-quinolinone, H-633
- 2,4-Dimethylquinolizidine, D-783
- N*^b,*N*^b-Dimethylserotonin, B-393
- 2,4'-Dimethylspiro[2-azabicyclo[2.2.2]octane-5,2'-[2*H*]pyran]-6'(3'*H*)-one, D-798
- 2,5-Dimethyl-3-styrylpyrazine, D-767
- 5-Dimethylsulfonio 2-amino-4-hydroxypentanoate, D-784
- 3-(Dimethylsulfonio)-*N,N,N*-trimethylpropanaminium(2+), T-612
- N,N*-Dimethyl-2-(1,2,3,9-tetrahydropyrrolo[2,1-*b*]quinazolin-3-yl)benzenamine, V-40
- 2,5-Dimethyl-5,6,7,8-tetrahydroquinoline, T-171
- N,N'*-Dimethyltetrandrinum, T-286
- N,O*-Dimethylthaicanine, T-301
- O,O*-Dimethylthalisopidine, T-346
- N,N'*-Di(6-methylthiohexyl)urea, D-829
- N,N*-Dimethyltryptamine *N*-oxide, D-785
- N,N*-Dimethyltryptamine, D-785
- N,N*-Dimethyltryptophan methocation methyl ester, T-640
- N,N*-Dimethyltryptophan, T-640
- Dimethyltubocurarine, C-420
- Dimethyltubocurarinium chloride, C-420
- N,N*-Dimethyltyramine, H-357

- N,N*-Dimethyltyrosine, T-700
N,N'-Dimethylurabaine, U-41
 1,3-Dimethyluracil, D-777
 14-(*N,N*-Dimethyl-L-valyloxy)paspalinine, P-109
 Dimethylwarifteine, W-9
 1,3-Dimethylxanthine, T-360
 3,7-Dimethylxanthine, T-355
 1,9-Dimethylxanthine, X-5
 3,9-Dimethylxanthine, X-5
 1,7-Dimethylxanthine, X-5
 3*R*,3'*R*-Dimethylxestospongin C, A-1374
N,N-Dimethylxopine, A-1066
N,N-Dimethylxenkerine, T-524
 Dimsulf, D-726
N,N-Dinerolyguanidine, B-173
 Dinicotinic acid, P-902
 Dinicotinic amide, P-902
 3,5-Dinitrotyramine, H-481
 Dinklacorine, T-406
 Dinklageine, D-786
 Dinklageine, D-787
 Dinor-C-alkaloid H, T-436
 Dinorcapsaicin, C-100
 19,23-Dinorconan-1,3,5(10),14-tetraen-3-ol, D-788
N-Dinogramine, I-97
 Dinorperipentadenine, P-262
N,N'-Dioctadecanoylthanediamine, E-225
 Diolamine, D-388
N,N'-Dioleylethylenediamine, E-225
 Diolmycin A1, D-789
 Diolmycin A2, D-789
 Dioncolactone A, D-790
 Dioncopeltine A, D-793
 Dioncophyllacine A, D-791
ent-Dioncophylline A, D-793
 Dioncophylline A, D-793
 Dioncophylline D, D-792
 Dioncophylline A, D-793
 Dioncophylline B, D-794
 Dioncophylline C, D-795
 Dioncophyllinol B, D-796
 Dionina, M-704
 Diopterin, P-772
 Dioscoretine, D-797
 Dioscorine *N*-oxide, D-798
 Dioscorine, D-798
 Dioxamide, D-799
 Dioxamine, D-799
 Dioxapyrrolomycin, P-974
 Dioxibrassinin, D-811
 Dioxindole-3-acetic acid, D-451
 Dioxindole, I-91
 Dioxinoacrimarine A, D-800
 4,5-Dioxoartacinate, A-1450
 3,18-Dioxo-1,4-conadienine, H-314
 5,6-Dioxocoronaridine, C-657
 Dioxocularine, D-801
 1-(1,6-Dioxo-2,4-decadienyl)piperidine, O-174
 4,5-Dioxodehydroasimilobine, C-273
 4,5-Dioxodehydrocrebanine, D-802
 4,5-Dioxodehydronantenine, C-684
 5,10-Dioxo-1,6-diazacyclodecane-2,7-dicarboxylic acid, D-803
 10,11-Dioxo-2-epierythratidine, E-207
 10,11-Dioxoerysotrine, E-203
 10,11-Dioxoerythraline, E-205
 10,11-Dioxoerythratidinone, E-207
 10,11-Dioxoerythratine, E-208
 3,6-Dioxohexahydropyrrolo[1,2-*a*]pyrazine (in-corr.), C-865
 11,16-Dioxoholarrhimine, D-257
 1,14-Dioxo-2-hydroxy-7,8-(methylenedioxy)-12,13-dimethoxyaporphoadane, C-375
 2,4-Dioxoimidazolidine, I-40
 4-(2,5-Dioxo-4-imidazolidinyl)carbamic acid, D-804
 (2,5-Dioxo-4-imidazolidinyl)urea, A-625
 10,17-Dioxo- β -isosparteine, I-322
 10,22-Dioxokopsane, D-805
 Dioxolamycin, D-806
 α,α' -(1,3-Dioxolane-4,5-diylidene)bis[benzeneacet-ic acid], U-57
 5-(1,3-Dioxolan-2-yl)-1*H*-indole, I-87
 Dioxolide A, D-807
 Dioxolide B, D-807
 1,3-Dioxolo[4,5-*g*]isoquinoline, M-446
 1,3-Dioxolo[4,5-*g*]isoquinolin-5(6*H*)-one, I-305
 5-(1,3-Dioxolo[4,5-*f*]isoquinolin-8-yl)-1,3-benzadioxole-4-methanol, D-136
 6-(1,3-Dioxolo[4,5-*g*]isoquinolin-5-ylmethyl)-2,3-dimethoxyphenol, B-103
 [1,3]Dioxolo[4,5-*f*]phenanthridine, M-448
 [1,3]Dioxolo[4,5-*f*]phenanthridin-6(5*H*)-one, C-744
 7*H*-[1,3]Dioxolo[4,5-*f*]pyrrolo[3,2,1-*de*]phenanthridine, H-290
 7*H*-[1,3]Dioxolo[4,5-*f*]pyrrolo[3,2,1-*de*]phenanthridin-7-one, H-290
 1-(1,12-Dioxo-2-octadecenyl)piperidine, O-194
 1-(1,12-Dioxo-2-octadecenyl)pyrrolidine, O-194
 1-(1,12-Dioxooctadecyl)piperidine, O-192
 1-(1,12-Dioxooctadecyl)pyrrolidine, O-192
N-[2,6-Dioxo-1-(2-phenylethyl)-3-piperidinyl]-2-methylbutanamide, J-58
N-(2,6-Dioxo-3-piperidinyl)benzeneacetamide, D-808
 5,6-Dioxopleiocarpine, K-78
 10,11-Dioxopleiocarpine, K-78
 6-(6,8-Dioxo-1-purinyloxy)suberosanone, D-809
N-[5-(2,5-Dioxo-1-pyrrolidinyl)pentyl]-*N*-hydroxybenzeneacetamide, T-100
 3,4-Dioxosarcocapnidine, Y-4
 10,17-Dioxosparteine, D-810
 2,17-Dioxosparteine, O-188
 Dioxybrassinin, D-811
 Dipegine, D-812
 Dipeginol, D-812
 2,6-Di-4-pentenylpiperidine, D-813
 2,6-Diphenacylpiperidine, L-218
 Diphenazithionin, D-814
N,N'-Diphenethylurea, B-216
 4,5-Diphenoxy-1,2-dicyanobenzene, D-533
 Diphenyl ether-3,3',5,5'tetracarboxylic acid, D-815
 4,8-Diphenyl-5*H*-1,3-dioxolo[4,5-*a*]azepine-5,7(6*H*)-dione, D-816
 Diphenylenimine, C-120
 8,10-Diphenyllobelidiol, L-218
 8,10-Diphenyllobelidione, L-218
 8,10-Diphenyllobelionol, L-219
 8,10-Diphenylnorlobelidiol, L-218
 8,10-Diphenylnorlobelidione, L-218
 3,5-Diphenyl-2-oxazolidinone, P-347
 2,5-Diphenyl-2-oxazoline, D-430
 α,α' -Diphenyl-2,6-piperidinediethanol, L-218
 2,3-Diphenyl-1*H*-pyrrole, D-817
N,N'-Diphenylurea, D-818
 Diphosphothiamine, T-366
 Diphyllin, C-357
 Diphylline, C-357
 Dipicolinic acid, P-901
 Dipiperamide A, D-819
 Dipiperamide B, D-819
 Dipiperamide C, D-820
 Dipiperamide D, D-821
 Dipiperamide E, D-822
 1,1'-Dipiperidine, B-141
 3,3'-(2,4-Dipiperidinediyl)bispyridine, A-973
 1,3-Di-2-piperidinyl-2-propanone, A-969
 Diplamine, D-823
 Diploceline, I-319
 Diplothyrcine, M-713
 Dippinine A, C-389
 Dippinine B, C-389
 Dippinine C, D-824
 Dippinine D, D-824
N,N-Diprenylguanidine, B-199
N,N'-Diprenylguanidine, P-769
 3,5-Diprenylindole, B-200
 3,6-Diprenylindole, B-201
 3,7-Diprenylindole, B-202
 5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriacon-ta-10,25-diene-15,30-dione, D-825
 5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriacon-ta-7,10,22,25-tetraene-15,30-dione, D-825
 5,20-Dipropyl-1,16-dioxa-4,19-diazacyclotriacon-ta-7,10,25-triene-15,30-dione, D-825
 Dipropylentriamine, D-255
 5,8-Dipropylindolizidine, O-44
 8,10-Dipropyllobelidiol, B-187
 α,α' -Dipropyl-2,6-piperidinediethanol, B-187
 Diptaline, D-826
 Diptamine, D-827
 Dipterine, M-582
 Diphthaline, D-826
 Dipthamine, D-827
 Dipthocarpamine, D-828
 Dipthocarpidine, D-829
 Dipthocarpilidine, D-830
 Diptocarpaine, D-828
 Diptocarpamine, D-828
 Diptocarpidine, D-829
 Diptocarpilidine, D-830
 Diptocarpiline, D-829
 Di-2-pyridinyl disulfide, D-831
 2,2'-Dipyridyl disulfide, D-831
 2,3'-Dipyridyl, B-145
 2,4-Di-3-pyridylpiperidine, A-973
 Dipyrrithione, D-831
 3,5-Di-2-pyrrolidinylpyridine, D-832
 5*H*,10*H*-Dipyrrrolo[1,2-*a*:1',2'-*d*]pyrazine-5,10-dione, D-833
 Di-10*H*-quindolin-11-ylmethanone, C-787
 Dirin, B-101
 Dirosine, G-212
 Discarene C, D-834
 Discarene D, D-835
 Discarine A, A-935
 Discarine B, D-836
 Discarine C, D-837
 Discarine D, D-838
 Discarine E, D-839
 Discarine F, A-149
 Discarine G, A-150
 Discarine H, F-149
 Discarine I, D-836
 Discarine K, D-836
 Discarine L, F-151
 Discarine M, D-840
 Discarine N, D-841
 Discarine X, N-334
 Discobahamin A, D-842
 Discobahamin B, D-843
 Discodermide, D-844
 Discodermindole, D-845
 Discoguttine, D-846
 Discokiolide A, D-847
 Discokiolide B, D-848
 Discokiolide C, D-848
 Discokiolide D, D-848
 Discolorine, A-957
 Discorhabdin A, D-850
 Discorhabdin B, D-850
 Discorhabdin C, D-851
 Discorhabdin D, D-852
 Discorhabdin E, D-851
 Discorhabdin G, D-851
 Discorhabdin H, D-853
 Discorhabdin I, D-850
 Discorhabdin L, D-852
 Discorhabdin N, D-852
 Discorhabdin P, D-851
 Discorhabdin Q, D-850
 Discorhabdin R, D-850
 Discorhabdin S, D-854
 Discorhabdin T, D-854
 Discorhabdin U, D-854
 Discorhabdin V, D-849
 Discorhabdin W, D-855
 Discretamine, D-856
 Discretine, X-30
 Discretinine, C-694
*N*¹,*N*⁴-Disinapoylputrescine, D-362
 Disinapoylspermidine, S-398
 Disinapoylspermine, S-399
 2,2'-Disinomenine, S-313
 1,1'-Disinomenine, S-313

- Disorazole A₁, D-859
 Disorazole A₂, D-859
 Disorazole A₃, D-859
 Disorazole A₄, D-859
 Disorazole A₅, D-859
 Disorazole A₆, D-859
 Disorazole A₇, D-859
 Disorazole B₁, D-859
 Disorazole B₂, D-859
 Disorazole B₃, D-859
 Disorazole B₄, D-859
 Disorazole C₁, D-857
 Disorazole C₂, D-858
 Disorazole D₁, D-859
 Disorazole D₂, D-859
 Disorazole D₃, D-859
 Disorazole D₄, D-859
 Disorazole D₅, D-859
 Disorazole E₁, D-859
 Disorazole E₂, D-859
 Disorazole E₃, D-859
 Disorazole F₁, D-859
 Disorazole F₂, D-859
 Disorazole F₃, D-859
 Disorazole G₁, D-860
 Disorazole G₂, D-861
 Disorazole G₃, D-862
 Disorazole H, D-859
 Disorazole I, D-859
 Dispacamide A, D-863
 Dispacamide B, D-863
 Dispacamide C, D-863
 Dispacamide D, D-863
 Dispegatrine, D-864
 Dispyrin, D-865
 Distamycin A, D-866
 Distamycin 3, D-866
N,N'-Distearoylethylenediamine, E-225
 Distichamine, D-867
 Distichine, P-581
 Distomadine A, D-868
 Distomadine B, D-869
 15,34-Di-*O*-sulfatobastadin 7, B-32
Cortinarius Disulfide, D-870
 Ditabersonine, T-11
 Ditaine, D-410
 Dithermamine, D-871
 5,5'-Dithiobis[1,4-dihydro-4-oxo-6-(2-pentyl-4-oxazolyl)-2-pyridinecarboxaldehyde], A-1123
 5,5'-Dithiobis[4-hydroxy-6-(2-pentyl-4-oxazolyl)-2-pyridinecarboxaldehyde], A-1123
 [2,2'-Dithiobis[pyridine] 1,1'-dioxide-*O,O',S*]sulfato(2-)-*O*]magnesium, D-831
 2,2'-Dithiobispyridine, D-831
 Dithiocolchicine, D-872
 2-(1,3-Dithiolan-2-yl)-1*H*-pyrrole, P-942
 Dithyreanitrile, D-873
 3,6-Ditigloyloxynortropine, A-1584
 3,6-Ditigloyloxystropane, M-391
 3,6-Ditigloyloxystropan-7-ol, M-392
 3,6-Ditigloylstrompanediol, M-391
 1,4-Ditropine α -isatropate, B-56
 1,4-Ditropine β -isatropate, B-56
 1,4-Ditropine isatropate, B-56
 Ditryptophenaline, D-874
 Diurobromine, T-355
 Divaricine, G-31
 Divarine, D-875
 Divicine, D-253
 DL 152, R-52
 DM SHT, B-393
 DMF, D-747
 DM1, M-135
 DMT, D-785
 Dnacin A₁, D-876
 Dnacin B₁, D-877
 DO 248A, R-54
 DO 248B, R-55
 DOB 41, A-1160
 Docosaheptaenoic acid-paclitaxel, T-52
 Docosahydro-1*H*-pyrido[1,2-*f*][1,5,10]triazacyclocosin-10(1*H*)-one, I-259
 3-[2-(Docosanoylaminoethyl)]indole, T-639
N-Docosanoylanthranilic acid, A-706
N^b-Docosanoylserotonin, H-755
N^b-Docosanoyltryptamine, T-639
 2,4,10-Docosatrienoic isobutylamide, D-878
 6-Docosenamamide, D-879
 13-Docosenamamide, D-880
 6-Docosenoic acid, D-879
 13-Docosenoic acid, D-880
N-(13-Docosenoyl)anthranilic acid, A-706
N^b-(13*Z*-Docosenoyl)serotonin, H-755
 3,5-Dodecadiene-2-amine, D-881
 2,4-Dodecadiene-8,10-diyenoic acid isobutylamide, D-882
 2,4-Dodecadiene-8,10-diyenoic acid 2-methylbutylamide, D-882
 2,4-Dodecadiene-8,10-diyenoic acid, D-882
 2,4-Dodecadienoic acid *p*-hydroxyphenethylamide, H-669
 2,4-Dodecadienoic acid isobutylamide, D-883
 2,4-Dodecadienoic acid pyrrolidide, D-883
 2,4-Dodecadienoic acid, D-883
 2,8-Dodecadienoic acid, D-884
 1-(2,4-Dodecadienyl)pyrrolidine, D-883
 2-(3,5-Dodecadienyl)-1-ethylpyrrolidine, V-104
N-(3,6-Dodecadienyl)sulfamic acid, D-885
 2-(5,7-Dodecadien-3-ynyl)-1-ethylpyrrolidine, V-104
 1-[[Dodecahydro-6,9-bis(methylene)-2-oxoazuleno[4,5-*b*]furan-3-yl]methyl]proline, I-162
 Dodecahydro-1*H*,5*H*,10*H*-dipyrido[2,1-*f*:3',2',1'-*i*][1,6]naphthyridin-10-one, M-121
 Dodecahydro-1*H*,4*H*,9*H*-dipyrido[2,1-*b*:3',2',1'-*i*]-quinazoline, S-140
 Dodecahydro-2-hydroxy-7,14-methano-4*H*,6*H*-dipyrido[1,2-*a*:1',2'-*e*]diazocin-4-one, H-551
 Dodecahydro-9-hydroxy-7,14-methano-2*H*,6*H*-dipyrido[1,2-*a*:1',2'-*e*]diazocin-6-one, V-153
 Dodecahydro-1-hydroxy-11-methyl-1,9-ethanobenzo[*f*]quinolizin-14-one, 13C1, F-73
 1,4,5,6,7,8,9,10,11,13,14,16a-Dodecahydro-13-(1-hydroxypropyl)pyrido[2,1-*d*][1,5,9]triazacyclotridecin-2(3*H*)-one, P-40
 Dodecahydro-7-(1*H*-indol-3-ylmethyl)- $\alpha,\alpha,6a,10b$ -tetramethyl-8-methylene-1*H*-naphtho[2,1-*b*]pyran-3-methanol, E-75
 Dodecahydro-7,14-methano-2*H*,6*H*-dipyrido[1,2-*a*:1',2'-*e*]diazocine, S-385
 Dodecahydro-2-methylpyrido[2,1,6-*de*]quinolizine, D-886
 Dodecahydro-5-(1-oxobutyl)-6*H*-pyrido[2,1-*f*][1,6]naphthyridin-6-one, H-38
 Dodecahydro-2,3,7,10-tetramethylcyclopenta[*b*]quinolizine, A-509
 Dodecahydro-3,4,7,10-tetramethylcyclopenta[*b*]quinolizine-2-methanol, A-509
 Dodecahydro-2,3,7,10-tetramethylcyclopenta[*b*]quinolizine-2-ol, A-509
 Dodecahydro-1*H*,6*H*-[1,2,5,6]tetrathiocino[3,4-*a*:8,7-*a'*]dipyrrolizine, C-186
 1,1a,2,4,4a,5,5a,6,6a,7,8,8a-Dodecahydro-2a*H*-3-thia-4-azacyclopenta[*mno*]pentaleno[2,1,6,5-*fghi*]aceanthrylene-2a-carboxylic acid, D-887
 Dodecahydro-3,7,10-trimethylcyclopenta[*b*]quinolizine-2-carboxaldehyde, A-509
 Dodecahydro-3,7,10-trimethylcyclopenta[*b*]quinolizine-2-methanol, A-509
 Dodecahydro-2,5,8-trimethyl-1,4,7,9*b*-tetraazaphenylene, D-888
 Dodecanamide, D-889
 Dodecanoic acid, D-889
N-Dodecanoyltyrosine, T-700
 2,4,6,8,10-Dodecapentaenedioic acid, D-890
 2,4,8,11-Dodecatetraenoic acid isobutylamide, D-892
 2,4,8,10-Dodecatetraenoic acid, D-891
 2,4,8,11-Dodecatetraenoic acid, D-892
 2,6,8,10-Dodecatetraenoic acid, D-893
 6-(1,3,5,9-Dodecatetraenyl)-2-methyl-3-piperidinol, C-680
 6-(1,3,7,9-Dodecatetraenyl)-2-methyl-3-piperidinol, C-680
 2,4,8-Dodecatrienoic acid isobutylamide, D-895
 2,4,6-Dodecatrienoic acid, D-894
 2,4,8-Dodecatrienoic acid, D-895
 2,4,10-Dodecatrien-8-ynoic acid isobutylamide, D-896
 2,4,10-Dodecatrien-8-ynoic acid, D-896
 2-Dodecene-8,10-diyenoic acid isobutylamide, D-897
 2-Dodecene-8,10-diyenoic acid, D-897
N-(7-Dodecenyl)tyrosine, T-700
 2-(1*E*-Dodecenyl)-4-hydroxyquinoline, D-898
 2-(1*Z*-Dodecenyl)-4-hydroxyquinoline, D-898
 2-(1*E*-Dodecenyl)-4(1*H*)-quinolinone, D-898
 2-(1*Z*-Dodecenyl)-4(1*H*)-quinolinone, D-898
 2-Dodecyl-4-hydroxyquinoline, D-898
 2-Dodecyl-1-methyl-4(1*H*)-quinolinone, D-898
 3-Dodecylpyridine, D-899
 2-Dodecyl-4-quinolinol, D-898
 2-Dodecyl-4(1*H*)-quinolinone, D-898
 Dogo base, H-163
 Doisuthine, D-900
 Dolabellin, D-901
 Dolaconine, A-99
 Dolastatin 10, D-907
 Dolastatin 18, D-908
 Dolastatin 1, D-902
 Dolastatin 2, D-904
 Dolastatin 3, D-905
 Dolastatin 4, D-906
 Dolastatin E, D-909
 Dolastatin I, D-903
 Dolichantoside, S-583
 Dolichocurine, D-910
 Dolichotheline, H-307
 Dolichothine, D-911
 Dolichotine A, M-352
 Dolichotine B, M-352
 Dolichotine C, M-352
 Dolichotine D, M-323
 Dolichotine E, M-312
 Domesticine, N-29
 Domestine, N-29
 Domoic acid, D-912
 Domoilactone A, D-913
 Domoilactone B, D-913
 Donasine, D-914
 Donaxamine, I-97
 Donaxanine, D-915
 Donaxaridine, D-916
 Donaxarine, D-917
 Donaxine, D-736
 Dongbeinine, C-290
 Dongbeirine, C-290
 Donglingine, D-918
 Donine, D-919
 DOPA betaine, A-745
 DOPA, A-745
 Dopamine 3-glucoside, D-920
 Dopamine 4-glucoside, D-920
 Dopamine hydrochloride, D-920
 Dopamine lutine, I-93
 Dopamine methosalt, C-690
 Dopaminebetaxanthin, M-647
 Dopamine, D-920
 Dopar, A-745
 Dopastat, D-920
 Dopaxanthin, D-921
 Doranine, D-736
 Doriasenine, D-922
 Dorimidazole A, A-791
 Doronenine, B-401
 Doronine, O-92
 Dorrigocin A, D-923
 Dorrigocin B, D-924
 Doryfranfine, C-453
 Doryanine, I-305
 Doryflavine, D-925
 Doryphorine, D-926
 Doryphormine, I-305
 Dothiorelone A, C-963
 Dothiorelone B, C-963
 Dothiorelone C, C-963
 Dovyalicin A, D-927
 Dovyalicin B, D-927
 Dovyalicin C, D-928

- Dovyalicin D, D-928
 Dovyalicin E, D-927
 Dovyalicin F, S-398
 Dracocephalin A₁, T-572
 Dracocephalin A₂, T-572
 Dracocephalin A₃, T-572
 Dracocephalin A₄, T-572
 Dracocephalin B₁, T-573
 Dracocephalin B₂, T-573
 Dracocephalin B₃, T-573
 Dracocephalin B₄, T-573
 Dracocephalin C₁, T-572
 Dracocephalin C₂, T-572
 Dracocephalin C₃, T-572
 Dracocephalin C₄, T-572
 Dracocephalin D₁, T-573
 Dracocephalin D₂, T-573
 Dracocephalin D₃, T-573
 Dracocephalin D₄, T-573
 Dragabine, S-411
 Dragmacidin A, D-929
 Dragmacidin B, D-929
 Dragmacidin C, D-929
 Dragmacidin D, D-930
 Dragmacidin E, D-931
 Dragmacidin F, D-932
 Dragmacidin, D-929
 Dragmacidonamine A, D-933
 Dragmacidonamine B, D-933
 Dragonic acid, M-240
 Drazepinone, D-934
 Dregamine, D-935
 Dregaminic acid, D-935
 Dregaminol methyl ether, D-935
 Dregaminol, D-935
 Drewamine, M-705
 Drimentine A, D-936
 Drimentine B, D-937
 Drimentine C, D-937
 Drimentine D, D-938
 Drimentine E, D-939
 Drimiopsine A, D-940
 Drimiopsine B, D-940
 Dripterin, P-772
 Drodrenin, H-566
 Drosoplerin, D-941
 Drupacine, D-942
 Drupangtonine, D-943
 Dryadine, D-944
 Dryadodaphnine, D-944
 Drymaritine, H-441
 Dst 3, D-866
 DTPA, P-421
 Dubamine, M-458
 Dubinidine, D-945
 Dubinine, D-945
 Dubirheine, R-86
 Dubiusine, H-504
 Duboisine, T-632
 Duciformine, B-194
 Duckeine, D-609
 Duclouxine, M-342
 Duguecalyne, D-946
 Duguenaine, D-946
 Duguespixine, D-947
 Duguetine, D-948
 Duguevalline, D-949
 Duguevanine, D-950
 Duguexine *N*-oxide, P-781
 Duguexine, P-781
 Dumetorine, D-951
 Duocarmycin A, D-952
 Duocarmycin B₁, D-953
 Duocarmycin B₂, D-954
 Duocarmycin C₁, D-953
 Duocarmycin C₂, D-954
 Duocarmycin D, D-955
 Duocarmycin SA, D-956
 Duramorph, M-704
 Duraquin, Q-18
 Duraspan, C-555
 Duricaulic acid, D-957
 Dutadrupine, D-958
 Dynemicin A, D-959
 Dynemicin C, D-959
 Dynemicin L, D-960
 Dynemicin M, D-961
 Dynemicin N, D-960
 Dynemicin O, D-961
 Dynemicin P, D-961
 Dynemicin Q, D-961
 Dysazecine, D-962
 Dyshomerythrine, D-963
 Dysibetaine PP, D-965
 Dysibetaine, D-964
 Dysidamide B, D-966
 Dysidamide C, D-966
 Dysidamide D, D-966
 Dysidamide E, D-966
 Dysidamide F, D-966
 Dysidamide G, D-966
 Dysidamide H, D-966
 Dysidamide, D-966
 Dysidazirine, D-967
 Dysideaprolin A, D-968
 Dysideaprolin B, D-968
 Dysideaprolin C, D-968
 Dysideaprolin D, D-968
 Dysideaprolin E, D-968
 Dysideaprolin F, D-968
 Dysideapyrrolidone, D-969
 Dysideathiazole, D-970
 Dysidenamide, D-971
 Dysidenin, D-972
 Dysidine, D-973
 Dysiherbaine, D-974
 Dysinosin A, D-975
 Dysinosin B, D-976
 Dysinosin C, D-976
 Dysinosin D, D-976
 Dysithiazolamide, D-977
 Dysoxyline, D-978
 Dyspropterin, T-142
 E 162, B-110
 E 64, R-69
 E 212, S-53
 E129A, O-131
 EB 82B, H-214
 Ebeidine, C-290
 Ebeienine, C-308
 Ebeiensine, P-168
 Ebeietinone, E-1
 Ebeinone, C-308
 Ebenifoline EIII, P-247
 Ebenifoline EII, P-247
 Ebenifoline EI, P-247
 Ebenifoline EIV, P-244
 Ebenifoline EV, P-247
 Ebenifoline VII, P-252
 Ebenifoline WI, P-252
 Ebornoxin, E-6
 Ebracteolatinoside A, E-2
 Eburcine, E-4
 Eburenine, E-3
 Eburine, E-4
 Eburnal, E-6
 Eburnamenine, E-5
 Eburnamine, E-6
 Eburnaminol, E-6
 Eburnamonine *N*⁴-oxide, E-6
 Eburnamonine, E-6
 (±)-Eburnamonine, E-6
 (-)-Eburnamonine, E-6
 Eburnan, E-5
 Eburnaphylline, E-7
 Eburnoxine, E-6
 EC 40, D-761
 Ecgonidine, M-395
 Ecgonine methyl ester benzoate, C-537
 Ecgonine methyl ester, E-8
 Ecgonine, E-8
 Echcin I, P-198
 Echiguanine B, E-9
 Echihumiline *N*-oxide, E-10
 Echihumiline, E-10
 Echimidine *N*-oxide, E-11
 Echimidine, E-11
 Echinacein, D-893
 Echinatine *N*-oxide, E-12
 Echinatine, E-12
 Echinine, E-13
 Echinatine, E-12
 Echinobetaine A, E-14
 Echinobetaine B, E-15
 Echinocathrine A, E-16
 Echinocathrine B, E-17
 Echinocathrine C, E-17
 Echinomycin, Q-58
 Echinophyllin A, E-18
 Echinophyllin B, E-19
 Echinophyllin C, E-20
 Echinophyllin D, E-21
 Echinophyllin E, E-22
 Echinophyllin F, E-23
 Echinopsidine, M-558
 Echinopsine, M-558
 Echinoramine I, E-24
 Echinorine, M-275
 Echinoserine, E-25
 Echinosporamycin, E-26
 Echinossulfone A, E-27
 Echinossulfonic acid A, E-28
 Echinossulfonic acid B, E-28
 Echinossulfonic acid C, E-28
 Echinossulfonic acid D, E-28
 Echinozolinone, E-29
 Echitamidine *N*^b-oxide 19-glucoside, E-30
 Echitamidine *N*^b-oxide, E-30
 Echitamidine, E-30
 Echitamine, D-410
 Echitaminic acid, D-410
 Echitoserpidine, M-631
 Echitoserpine, M-631
 Echitovenaldine, M-631
 Echitovenidine, M-631
 Echitoveniline, M-631
 Echitovenine, M-631
 Echiumine, E-31
 Echiupinine *N*-oxide, E-32
 Echiupinine, E-32
 Ecliptalbine, E-33
 ECO 02301, A-1161
 ECO 4601, D-297
 Ecteinascidin 759A, E-37
 Ecteinascidin 759B, E-37
 Ecteinascidin 759C, E-37
 Ecteinascidin 583, E-35
 Ecteinascidin 594, E-34
 Ecteinascidin 596, E-35
 Ecteinascidin 597, E-35
 Ecteinascidin 722, E-36
 Ecteinascidin 729, E-37
 Ecteinascidin 736, E-36
 Ecteinascidin 743, E-37
 Ecteinascidin 745, E-37
 Ecteinascidin 770, E-37
 Ecteinascidin 786, E-37
 Ectoine, T-207
 Edeine, E-38
 Edpetilidine, C-290
 Edpetilidine, S-275
 Edpetiline, C-298
 Edpetilinine, S-275
 Edpetine, E-39
 Edpetinosine, C-290
 Edpetisidine, C-310
 Edpetisidine, C-312
 Edpetsine, H-461
 Edpetsinine, C-296
 Eduardine, C-290
 Eduardine, C-290
 Eduleine, D-660
 Edulimide, D-362
 Eduline, D-659
 Edulinine, E-40
 Edulirin A, T-349
 Edulisone A, E-41
 Edulisone B, E-41
 Edulitine, Q-47

- Edwardine, C-290
Edwardine, C-290
E297, F-173
Efatine, E-42
Efirine, I-339
Efudex, F-104
E620, G-107
Egeline, A-153
Egenine, E-43
Eglandine, C-657
Eglandulosine, C-657
Egonine, E-44
EGS 1300-1, A-1162
Egyt 402, N-303
E355, H-258
Ehretinine, H-630
Ehretioside A₁, S-302
Ehretioside A₂, S-302
Ehretioside A₃, S-302
Ehretioside B, D-646
2,4-Eicosadienoic acid isobutylamide, E-45
2,14-Eicosadienoic acid piperidide, E-46
2,4-Eicosadienoic acid piperidide, E-45
2,11-Eicosadienoic acid pyrrolidide, P-949
3,6-Eicosadienoic acid pyrrolidide, P-949
2,4-Eicosadienoic acid pyrrolidide, T-491
2,4-Eicosadienoic acid, E-45
2,14-Eicosadienoic acid, E-46
N-(2,14-Eicosadienoyl)piperidine, E-46
N-(2,4-Eicosadienoyl)piperidine, E-45
N-(2,11-Eicosadienoyl)pyrrolidine, P-949
N-(3,6-Eicosadienoyl)pyrrolidine, P-949
N-(2,4-Eicosadienoyl)pyrrolidine, T-491
4,5-Eicosanediamine, E-47
Eicosanoic acid pyrrolidide, T-491
N-Eicosanoylanthranilic acid, A-706
N-Eicosanoylpyrrolidine, T-491
N^b-Eicosanoylserotonin, H-755
N^b-Eicosanoyltryptamine, T-639
5,8,11,14-Eicosatetraenoic acid, E-48
N-(5,8,11,14-Eicosatetraenoyl)taurine, A-765
2,4,14-Eicosatrienoic acid isobutylamide, E-50
2,4,8-Eicosatrienoic acid isobutylamide, E-49
8,11,14-Eicosatrienoic acid, E-52
2,4,14-Eicosatrienoic acid, E-50
2,4,16-Eicosatrienoic acid, E-51
2,4,8-Eicosatrienoic acid, E-49
2-Eicosenoic acid pyrrolidide, T-491
4-(14*Z*-Eicosenoxyloxy)-3-methyl-2-butenenitrile, H-576
N-(2-Eicosenoyl)pyrrolidine, T-491
N^b-(11*Z*-Eicosenoyl)serotonin, H-755
3-(12*Z*-Eicosenylidene)-2-methyl-1-pyrroline, L-37
2-(Eicosylthio)-4,5-dihydro-3-methoxy-4-methylene-1*H*-pyrrole, I-172
Eilatin, D-299
Ejap 13, P-220
Ejap 14, P-220
Ekapterin, E-216
Ekeberginine, C-507
Ekamine, T-632
EL Base 2, E-297
EL Base 3, E-297
EL Base 1, I-65
Elacomine, E-53
Eladine, M-321
Elaeocarpenine, I-231
Elaeocarpidine, E-54
Elaeocarpiline, E-55
Elaeocarpine, E-56
Elaeokanidine A, E-57
Elaeokanidine B, E-57
Elaeokanidine C, E-57
Elaeokanine A, E-58
Elaeokanine B, E-58
Elaeokanine C, E-59
Elaeokanine D, E-60
Elaeokanine E, E-60
Elaidamide, O-37
Elaidic acid, O-37
Elaiomycin, E-61
Elanine, G-91
Elan, T-360
Elasine, M-307
Elatine, G-91
Elatrine, E-62
Eldelidine, M-307
Eldeline, M-307
Eldisine, V-110
Eleagnine, T-202
Elegansamine, E-63
Elegantine, M-50
Elegantissine, M-50
Eleocarpidine, E-54
Eleocarpiline, E-55
Eleocarpine, E-56
Eleokanidine A, E-57
Eleokanidine B, E-57
Eleokanidine C, E-57
Eleokanine A, E-58
Eleokanine B, E-58
Eleokanine C, E-59
Eleokanine D, E-60
Eleokanine E, E-60
Eletefine, S-541
Z-Eleutherobin, E-64
Eleutherobin, E-64
Eleuthoside A, E-64
Eliamide, E-65
Elisabethamine, E-66
Elixophyllin, T-360
Elliptamine, R-51
Ellipticine *N*²-oxide, E-67
Ellipticine, E-67
Elliptifoline, A-199
Elliptinine, E-68
Elliptinol, E-69
Elloxazinone A, A-854
Elloxazinone B, A-854
Elmerrillicine, E-70
Elsedine, V-110
Elwesine, C-748
Elymoclavine *O*-β-*D*-fructofuranoside, A-205
Elymoclavine *O*-β-*D*-fructofuranosyl-(2→1)-*O*-β-*D*-fructofuranoside, A-205
Elymoclavine, A-205
EM 4940A, S-295
EM 5596, N-101
EM 5519, S-2
E296, M-72
Emarginatine A, P-246
Emarginatine B, P-246
Emarginatine C, P-246
Emarginatine D, P-246
Emarginatine E, P-246
Emarginatine F, P-246
Emarginatine G, P-246
Emarginatine H, P-246
Emarginatine, P-253
Emeheterone, D-300
Emeniveol, E-71
Emetamine, E-72
Emetinal, E-72
Emetine hydrochloride, E-72
Emetine, E-72
Emetoplisce, E-72
Emiline, E-73
Emimycin, P-889
Emindole DA, E-74
Emindole DB, E-75
Emindole PA, E-76
Emindole PB, E-77
Emindole PC, E-76
Emindole SA, E-74
Emindole SB, E-78
Enactin Ia, N-113
Enactin Ib, N-113
Enactin IVa, L-189
Enactin IVb, N-113
Enactin Va, L-189
Enactin Vb, N-113
Enactin VIb, N-113
Enantiorhizoferrin, R-80
Encol, E-72
Endabuse, I-4
Endecaphyllin A, G-104
Endecaphyllin A₁, G-104
Endecaphyllin A₂, G-104
Endecaphyllin B, G-104
Endecaphyllin B₁, G-104
Endecaphyllin C, G-104
Endecaphyllin C₁, G-104
Endecaphyllin D, G-104
Endecaphyllin E, G-104
Endecaphyllin X, G-104
Endolobine, T-288
Endophenazine A, M-418
Endophenazine B, E-79
Endophenazine C, M-418
Endophenazine D, P-319
Enduracidine, E-80
Energlut, G-108
Engleromycin, E-81
Enicoflavine, E-82
Enkleine, E-83
Enneaphylline, C-805
Enprofylline, X-5
Entadamide A β-*D*-glucoside, E-84
Entadamide A, E-84
Entadamide B, E-85
Entadamide C, E-84
Enteramine, H-755
Enterobactin, E-86
Enterocarpam II, T-539
Enterocarpam I, T-251
Enterochelin, E-86
Ephrosine, A-1365
Ephedradine A, A-1334
Ephedradine B, A-1334
Ephedradine C, A-1334
Ephedradine D, A-1334
Ephedralone, D-676
Ephedrine hydrochloride, M-386
Ephedrine sulfate, M-386
Ephedrine, M-386
ψ-Ephedrine, M-386
(±)-ψ-Ephedrine, M-386
(±)-Ephedrine, M-386
(+)-Ephedrine, M-386
(-)-ψ-Ephedrine, M-386
Ephedrone, A-870
Ephedroxane, M-511
12-Epiacetyldehydronapelline, D-162
16-Epiaffinine, A-166
16-Epiaffinisine, T-425
Epiagelastine C, A-185
4-Epiaglaine A, A-199
15-Epiajaconine, A-211
19-Epiajmalicine, A-212
Epiajmaline, A-216
16-Epiakuammiline, A-231
7*a*-Epialexafflorine, T-555
8-Epialexafflorine, T-555
7-Epialexine, T-555
7*a*-Epialexine, T-555
8-Epialexine, T-555
Epiallocorynantheine, C-689
Epialloelaecarpiline, E-55
3-Epi-5-allomuscarine, M-773
17-Epialloyohimbine, Y-13
Epialphinine, A-638
19-Epiastogustine, E-30
19-Epiastonine, A-652
16-Epiastovenine, V-50
Epiamauroamine, A-680
Epiamurine, A-956
7-Epiancistrobrevine D, A-990
1-Epiancistrocladine, A-982
15-Epiansamitocin PHO 0, M-136
15-Epiansamitocin PHO 1, M-136
15-Epiansamitocin PHO 3, M-136
15-Epiansamitocin PHO 4, M-136
20-Epiantirrhine, A-1328
3-Epiantirrhine, A-1328
Epiaphylline, A-1337
Epiapocavidine, T-319
3-Epiaristoserratenine, A-1441
2-Epiaristolone, A-1430

- 16-Epiaspidodasycarpine, A-1492
 20,21-Epiaspidospermidine *N*-oxide, A-1498
 5,19-Epiaspidospermidine, A-1498
 20,21-Epiaspidospermidine, A-1498
 1-Epiaustraline 2-glucoside, T-555
 7-Epiaustraline, T-555
 7*a*-Epiaustraline, T-555
 1-Epiaustraline, T-555
 3-Epiaustraline, T-555
 10-Epiaxisonitrile 3, S-425
 13-Epibaptifoline, B-20
 3-Epibassiatin, B-29
 Epibatidine, E-87
 30-Epibatzelladine D, B-40
 Epiberberine, G-181
 Epiberbivaldine, P-15
 Epibisdehydroneotuberostemonine J, T-664
 1-Epibowdensine, B-261
 Epibubbialine, B-375
 Epibuphanisine, C-748
 2-Epicavine, C-41
 22-Epicapsimine, S-200
 6-Epicastanospermine, C-188
 6-Epicasuarine, T-262
 3-Epicasuarine, T-262
 Epicataline, C-190
 3-Epicatuabine H, M-391
 2-Epiccephalofortuneine, C-260
 7-Epiccephalomannine, T-52
 Epiccephalotaxine, C-271
 6-Epichasmanine, M-323
 14-Epichelidonine, C-357
 Epicinchonicinol, Q-17
 17-Epicinchophyllamine, C-448
 3-Epicinchophyllamine, C-448
 Epicoccamide B, E-88
 Epicoccamide C, E-88
 Epicoccamide, E-88
 Epicoccarine A, E-89
 Epicoccarine B, E-89
 Epicoccin A, E-91
 Epicoccin B, E-90
 Epicoccin C, E-91
 Epicoccin D, E-91
 1'-Epicolchicoside, D-92
 3-Epicomosine, C-587
 3-Epiconamine, C-595
 20-Epicorreatine B, C-660
 6-Epicorynoline, C-692
 14-Epicorynoline, C-692
 Epicorynoxidine, T-212
 Epicostaclavine, F-49
 1-Epicrassicaudine, M-323
 3-Epicrinamine, H-5
 Epicrinine, C-748
 5-Epiculantraminol, C-803
 7-Epicylindrospermopsin, C-930
 Epidapiramicin A, D-73
 Epidarlinine, D-76
 3-Epidasycarpidone, D-84
 16-Epideacetyllakuammiline *N*⁴-oxide, A-231
 16-Epideacetyllakuammiline, A-231
 1-Epideacetylbowdensine, B-261
 7-Epi-10-deacetyl-10-oxotaxuyunnanine A, T-52
 7-Epi-10-deacetyltaxuyunnanine A, T-52
 10-Epidecaline, V-86
 20-Epi-16-decarbomethoxy-19,20-dihydrovobasine, D-119
 Epi- α -Decumbensine, E-43
 16-Epidehydroaffinisine, G-23
 12-Epidehydrolicidusculine, D-162
 12-Epidehydronapelline, D-162
 16-Epi-14,15-dehydrovincine, V-115
 5-Epidemethoxyabresoline, L-53
 1-Epidemethoxybowdensine, A-670
 5-Epi-4'-*O*-demethylancistrobertsonine C, A-980
 5-Epidemissidine, D-205
 8*a*-Epidendroprimine, O-42
 7-Epideoxynupharidine, D-227
 1-Epideoxynupharidine, D-227
 7-Epideoxynupharidine-6,7-diol, D-227
 22-Epi-28-deoxyzoanthenamamine, Z-31
 Epidermin, H-573
 Epiderstatin, E-92
 Epidicentrine, N-29
 19-Epidifforlemenitine, D-398
 6-Epidihomosedinone, D-403
 2,3-Epidihydrobellendine, B-57
 3-Epidihydrocorymine 17-acetate, D-410
 3-Epidihydrocorymine 3-acetate, D-410
 3-Epidihydrocorymine, D-410
 3-Epidihydrocorymantheol *N*-methosalt, D-412
 Epidihydrofawcettidine, F-28
 3-Epi-2,7-dihydrohomoerysotrine, T-51
 3-Epi-3,4-dihydro-3-hydroxygraciline, G-151
 Epidihydropropidine, M-529
 3-Epidihydroquinamine, Q-12
 3-Epi-6,7-dihydroschelhammeridine, C-587
 Epidihydroshihunine, D-511
 11-Epi-16,17-dihydroxylepenine, L-51
 3-Epi-2,18-dimethoxyschelhammeridine, D-963
 7-Epidioncophylline A, D-793
 1-Epidioncophylline B, D-794
 2-Epi-10,11-dioxoerythratidine, E-207
 16-Epidiplotelone, I-319
 5'-Epidomoic acid, D-912
 13-Epidorriginocin A, D-923
 20-Epidregamine, D-935
 5-Epidysidamide G, D-966
 25-Epiebeidonine, C-290
 16-Epieburnamine, E-6
 20-Epi-19 ξ -echitamine, E-30
 19-Epi-(+)-echitoveniline, M-631
 Epielaecarpiline, E-55
 16-Epielamine, M-307
 Epielwesine, C-748
 11-Epiephedradine A, A-1334
 3-Epi-17-epicinchophyllamine, C-448
 1-Epi-7-epideoxynupharidine, D-227
 6-Epi-16-epiormosanine, O-115
 1-Epi-1'-epithiobinupharidine, T-380
 5'-Epiergocryptine, E-158
 Epiervafolidene, E-186
 3-Epiervafolidine, E-186
 20-Epiervatamine, E-190
 11-Epierythratine *N*-oxide, E-206
 11-Epierythratine, E-206
 2-Epierythratidine, E-207
 2-Epierythratine, E-208
 11-Epierythrinine, E-205
 2-Epi-2-*O*-ethylcephalofortuneine, C-260
 25-Epietioline, S-199
 16-Epiexelsinine, M-291
 4-Epifagamine, H-614
 3-Epifagamine, H-614
 12-Epifinetianine, N-30
 12-Epifischerindole G isonitrile, F-63
 12-Epifischerindole I isonitrile, F-63
 12-Epifischerindole U isonitrile, F-63
 12-Epifischerindole U isothiocyanate, F-63
 11-Epifistularin 3, F-70
 6-Epiforesticine, M-323
 Epifortuneine, F-143
 Epifrin, A-152
 Epifuntumidine, A-880
 15-Epifuziline, M-313
 Epigalanthamine, G-3
 Epiglaucamine, G-87
 Epiglaudine, G-87
 Epiglaufidine, C-682
 Epiglucosatisin, G-102
 Epiglucosylgouniensiene, N-186
 Epiguaiipyridine, G-190
 Epiguanine, M-469
 11-Epihaemanthamine, H-5
 3-Epihaemanthidine, H-6
 12-Epihapalindole C, H-49
 12-Epihapalindole D isothiocyanate, H-49
 12-Epihapalindole E, H-49
 12-Epihapalindole F isothiocyanate, H-49
 12-Epihapalindole G, H-48
 12-Epihapalindole H, H-48
 12-Epihapalindole J, H-48
 12-Epihapalindole Q, H-49
 12-Epihapalindole R, H-49
 Epithermandolinol, H-70
 Epiheteroconessine, P-606
 19-Epiheyneanine, C-657
 2-Epihomoerythratine, S-126
 Epihomolycorine, L-347
 6-Epi- α -homomannojirimycin, B-186
 4-Epi- α -homonojirimycin, B-186
 18-Epihomomorsanamine, J-16
 6-Epihomosedinone, H-347
 16-Epi-10-hydroxyaffinine, A-166
 13-Epihydroxyanagyrrine, B-20
 2-Epihydroxyisotaxodine, H-331
 1-Epi-12 β -hydroxykarasamine, M-350
 16-Epi-17 α -hydroxy- $\Delta^{14,15}$ -kopsinine, V-49
 Epihydroxylupanine, H-554
 17-Epi-12-hydroxy-11-methoxydiaboline, C-115
 2-Epi-3-hydroxyquebrachidine, H-139
 3-Epi-12-hydroxyschelhammericine, S-126
 13-Epihydroxysparteine, T-361
 19-Epi-23-hydroxyspermostrychnine, S-400
 2-Epi-3-hydroxyvincamajine, H-139
 20-Epiibophyllidine, I-6
 19-Epiiboxygaine, I-4
 3-Epiignavinol, H-184
 Epiindolactam V, I-78
 19-Epi-3-isoajmalicine, A-212
 19-Epiisoatinsine, I-192
 17-Epiisocinchophyllamine, C-448
 6-Epiisocorynoline, C-692
 α -Epiisocycloheximide, D-765
 Epiisoleaeocarpiline, E-55
 Epiisopolisone, P-428
 Epiisopiloturine, P-430
 16-Epi-Z-isositsirikine, I-319
 16-Epiisositirikine, I-319
 19-Epiisovoacristine, I-344
 Epijoubertinamine, J-50
 3-Epikarounidol, M-730
 22-Epikopsanol, K-65
 Epikopsanol-22-ol-10-lactam, D-805
 Epikopsanol-10-lactam, D-805
 17-Epikopsaporine, K-66
 16-Epikopsinine, K-78
 17-Epikopsinol, K-66
 23-Epikuroyuridinine, K-114
 Epilachnadiene, E-93
 Epilachnene, E-93
 Epilamprolobine *N*-oxide, L-26
 Epilamprolobine, L-26
 Epilapidilectinol, L-43
 16-Epilatrunculin B, L-62
 2-Epilentiginosine, O-46
 11-Epileotidane, E-94
 11-Epilepenine, L-105
 Epileuconolam, L-155
 (\pm)-Epi-LL-C10037 α , A-804
 Epilobamide A, V-15
 Epilobscurinol, L-223
 31-Epilolitre B, L-236
 31-Epilolitre F, L-236
 31-Epilolitre N, L-237
 35-Epilolitre N, L-237
 Epilotaustralin, H-572
 12-Epilucidusculine, N-30
 Epilucumin, H-673
 Epilupinine *cis-p*-coumarate, E-95
 Epilupinine *N*-oxide, E-95
 Epilupinine, E-95
 Epilupinyl acetate *N*-oxide, E-95
 Epilupinyl acetate, E-95
 Epilupinyl *trans-p*-acetoxyinnamate, E-95
 Epilupinyl *trans-p*-coumarate, E-95
 Epilupinyl *trans-ferulate*, E-95
 Epilupinyl *cis-p*-rhamnosylcoumarate, E-95
 Epilupinyl *trans-p*-rhamnosylcoumarate, E-95
 Epilupinyl rhamnosylferulate, E-95
 12-Epilycodoline *N*-oxide, L-325
 12-Epilycodoline, L-325
 2-Epilycorine, L-348
 5-Epilyfoline, V-95
 10-Epilyfoline, V-95
 3-Epimacronine, P-626
 Epimalbrancheamide, M-69
 1-Epimanzamine D, M-91

- 3-Epimaritidine, M-108
 Epimaritinamine, M-108
 10-Epimartensine A, M-110
 19-Epimeloscandoinone, M-192
 3-Epimeloscine *N*-oxide, M-193
 17-Epimeloscine, M-193
 3-Epimeloscine, M-193
 13-Epimethoxylupanine, H-554
 17-Epi-11-methoxy-17-*O*-methyladiabolone, C-115
 Epimethoxy-nauclearine, N-62
 3-Epi-18-methoxyschelhammericine, D-963
 17-Epi-11-methoxyvincamajine, V-113
 17-Epi-11-methoxy- α -yohimbine, M-292
 16-Epimethuenine *N*-oxide, M-293
 16-Epimethuenine, M-293
 16-Epi-17-*O*-methylaffinine, A-166
 16-Epi-*N*-methylaffinine, A-166
 5-Epi-6-*O*-methylancistrobertsonine A, A-980
 9-Epi-*N*-methylhydroquinicol, Q-17
 7-Epi-*N*-methylidioncophylline A, D-793
 Epi-17-*O*-methylkribine, K-101
 Epi-21-*O*-methylkribine, K-101
 1-Epi-2-*N*-methylmanzamine D, M-91
 20-Epi-*N*-methylparavallarine, P-93
 16-Epi-*O*-methyl- Δ^{14} -vincanol, E-6
 22,26-Epiminocholesta-5,22(*N*)-diene-3,16-diol, S-199
 22,26-Epiminocholesta-5,22-diene-3,12-diol, V-68
 22,26-Epiminocholesta-22(*N*)-ene-3,16-diol, S-338
 22,26-Epiminocholesta-5-ene-3,16-diol, S-200
 22,26-Epiminocholesta-22(28)-en-3-ol, S-337
 22,26-Epiminocholesta-5-en-3-ol, V-73
 23,26-Epimino-3-hydroxycholesta-23(*N*)-ene-4,22-dione, S-339
 22,25-Epimino-27-norcholesta-5,25(*N*)-dien-3-ol, V-57
 22,25-Epimino-27-norcholesta-4,24-dien-3-one, M-550
 18,20-Epimino-27-norcholesta-4,6,18(11)-trien-3-one, M-59
 19,20-Epimino-27-norcholesta-5-en-3-amine, C-595
 19-Epiminovincine, M-631
 (19*R*)-Epimisiline, H-743
 (19*S*)-Epimisiline, H-743
 3-Epimuscarine, M-773
 Epimyrine, O-53
 3-Epimyrtoidine, M-803
 15-Epinagarine, M-313
 Epinal, A-152
 12-Epinapelline *N*-oxide, N-30
 1-Epinapelline, N-30
 12-Epinapelline, N-30
 Epinardine A, E-96
 Epinardine B, E-97
 Epinardine C, E-97
 Epinardine D, E-97
 5-Epinareline ethyl ether, N-47
 19-Epinauclearine, N-62
 17-Epinaucleidinal, N-74
 6-Epineoline, M-323
 1-Epineoline, M-323
 6-Epineolinine, M-323
 7-Epineoptilocaulin, P-773
 8-Epineosarracine, S-90
 9-Epineoxaline, N-149
 Epinephrine bitartrate, A-152
 Epinephrine borate, A-152
 Epinephrine, A-152
 Epinephryl borate, A-152
 Epingouniensine, N-186
 Epinine, E-98
 3-Epinitrarine, N-246
 Epinorgalanthamine, G-3
 Epinorlycoramine, G-3
 16-Epinormacusine B, T-425
 17-Epinorseredamine, T-288
 6-Epi-11-nortetrodotoxin, T-298
 27-Epinortryptoquivaline, N-315
 Epinorzoanthamine, N-317
 3-Epinuphamine, N-340
 3-Epinupharamine, N-340
 7-Epinupharidine, D-227
 Epiodorine, O-67
 16-Epiolvoretin A, T-63
 Epiophiocarpine, O-94
 11-Epiorantine, A-1334
 6-Epioreobeiline, O-103
 16-Epiormosanine, O-115
 18-Epiormosanine, O-115
 Epioxodaphnigraciline, D-50
 8-Epi-9-oxoeptophilone D, E-117
 16-Epi-6-oxosilicine, M-293
 19-Epi-3-oxovoacristine, V-170
 2'-Epioxysporidinone, O-226
 Epioxyzoanthamine, Z-30
 Epipachysamine A, P-601
 Epipachysamine B, P-601
 Epipachysamine C, P-601
 Epipachysamine D, P-601
 Epipachysamine E 5-en-4-one, D-282
 Epipachysamine E, P-601
 Epipachysamine F, P-601
 3-Epipachysamine H, P-601
 Epipachysandrine A, D-266
 16-Epipanarine, T-425
 2-Epipanrassidine, U-35
 20-Epipandoline, P-733
 14-Epipapaverrubine B, G-87
 14-Epipapaverrubine D, G-87
 3-Epipapyramine, P-86
 Epipassicoriacin, D-565
 4-Epiphyllanthine, P-386
 24-Epipinnasterone, H-244
 Epipleiocarpamine *N*¹-oxide, P-518
 19'-Epipleiomutinine, P-827
 Epipodopetaline, E-99
 Epipolasin A, E-100
 Epiporphyrroxine, G-87
 Epipseudoconhydrine, P-653
 6-Epipubescence, M-309
 Epipyridone, E-101
 16-Epipyroaconitine, H-122
 16-Epipyrohaconitine, H-254
 16-Epipyrojesaconitine, H-122
 16-Epipyromesaconitine, H-122
 Epiquinamide, A-907
 Epiquinamine, Q-12
 Epiquinidine, Q-18
 Epiquimine, Q-20
 3-Epirauvanine, R-51
 Epirauviridine, R-27
 3-Epirauwolsine, Y-13
 Epiremijinine, R-40
 3-Epirescinnamine, R-50
 16-Epirhazinaline, R-75
 Epirubijervine, R-144
 3-Epiryanodine, R-175
 Episarcondonin α , S-76
 Episarcondonin β , S-76
 Episarcondonin, S-76
 Episarcotrine A, S-82
 Episarcotrine B, S-82
 Episarcotrine C, S-82
 Episarcoviolin α , S-76
 16-Episarpagine, S-89
 8-Episarracine *N*-oxide, S-90
 8-Episarracine, S-90
 Episcandomeline, S-120
 Episcandomelonine, S-121
 3-Epischelhammericine, S-126
 3-Epischelhammeridine, S-125
 3-Epischelhammerine, S-126
 3-Epischelhammericine B, T-51
 19-Epischolaricine, E-30
 Episcopalidine, H-161
 Episcopalisine, A-97
 Episcopalisinine, A-97
 Episcopalitine, A-99
 2-Episecurinine, A-630
 15-Episenbusine C, M-313
 19-Episerpentine, A-652
 6-Episevedine, C-295
 16-Episisilicine, M-293
 20-Episisilicine, M-293
 14-Episiinomenine, S-313
 16-Episisirikine, S-325
 5-Epismenospongiarine, S-329
 5-Epismenospongidine, S-329
 5-Epismenospongine, S-329
 5-Epismenospongiorine, S-329
 Episolacarpine, S-336
 5-Episophocarpine, S-371
 5-Episparsoymycin, S-384
 15-Epistemadenine, S-510
 12-Epistemocochinine, P-670
 15-Epistenine, S-528
 Epistephamiensine, S-537
 Epistephamine, E-102
 Episteporphine, S-549
 3-Epistrictosidinic acid, S-583
 22-Episcyphantine, S-456
 Epitabernaemontanine, D-935
 16-Epitacamine, T-12
 20-Epitalcarpine, T-20
 7-Epitaxol B, T-52
 7-Epitaxol, T-52
 7-Epitaxuyunnanine A, T-52
 3-Epitazattadiol, E-44
 3-Epitazettine, T-53
 17-Epitetraphyllicine, T-288
 Epitetraphyllin B, D-565
 4-Epitetrodotoxin, T-298
 6-Epitetrodotoxin, T-298
 1-Epithiobinupharidine, T-380
 1'-Epithiobinupharidine, T-380
 16-Epitombozine, T-425
 Epitrate, A-152
 10-Epitrewiasine, T-450
 Epitrichoponamic acid, T-493
 Epitrixilline, T-634
 27-Epitryptoquivaline, T-645
 Epiuleine, U-8
 Epivaldiberine, P-15
 Epivaldivianine, P-15
 16-Epivenenatine, V-50
 17-Epivenenatine, V-50
 20-Epiverazine, V-73
 10-Epiverruculotoxin, V-84
 Epivertaline, V-86
 20'-Epivinblastine, V-110
 (\pm)-Epivincadine, V-112
 (+)-Epivincadine, V-112
 (-)-Epivincadine, V-112
 16-Epi- Δ^{14} -vincamine, V-115
 16-Epivincamine, V-115
 21-Epivincamine, V-115
 16-Epi- Δ^{14} -vincine, V-115
 19-Epivindolinine *N*-oxide, V-132
 19-Epivindolinine, V-132
 Epivittatine, C-748
 19-Epivoacangarine, V-170
 16-Epivoacarpine, V-171
 (19*Z*)-16-Epivoacarpine, V-171
 Epivoacarine, V-168
 19-Epivoacristine hydroxyindolenine, C-658
 19-Epivoacristine, V-170
 16-Epivobasinal, V-183
 16-Epivobasine, P-268
 16-Epivobasinic acid, P-268
 Epivolkenin, D-565
 13-Epiwanpeine A, C-298
 3-Epiwelwitindolinone B isothiocyanate, W-13
 3-Epiwilsonine, F-143
 3-Epiwilsonone, F-143
 13-Epiyenusomine, R-9
 3-Epi- α -yohimbine, Y-13
 3-Epi- β -yohimbine, Y-13
 Epiyohimbol, Y-16
 8-Epizanthocadinanin B, Z-3
 8-Epizanthomurolanin, Z-3
 Epizoanthoxanthin A, P-94
 Epizoanthoxanthin B, P-94
 EPO 906, E-111
 Epocarbazolin A, E-103
 Epocarbazolin B, E-104
 Epohelmin A, E-105
 Epohelmin B, E-105
 Epolactaene, E-106
 Epostatin, E-107
 Eptophilone A, E-110

- Epothilone A₁, E-108
 Epothilone A₂, E-108
 Epothilone A₈, E-109
 Epothilone A₉, E-110
 Epothilone B, E-111
 Epothilone B₁₀, E-111
trans-Epothilone C₁, E-112
trans-Epothilone C₂, E-112
 Epothilone C, E-115
 Epothilone C₁, E-112
 Epothilone C₂, E-112
 Epothilone C₃, E-113
 Epothilone C₄, E-114
 Epothilone C₅, E-115
 Epothilone C₆, E-115
 Epothilone C₇, E-115
 Epothilone C₈, E-115
 Epothilone C₉, E-115
 Epothilone D, E-117
 Epothilone D₁, E-116
 Epothilone D₂, E-116
 Epothilone D₅, E-117
 Epothilone E, E-110
 Epothilone F, E-111
 Epothilone G₁, E-118
 Epothilone G₂, E-119
 Epothilone H₁, E-118
 Epothilone H₂, E-119
 Epothilone I₁, E-120
 Epothilone I₂, E-121
 Epothilone I₃, E-122
 Epothilone I₄, E-123
 Epothilone I₅, E-124
 Epothilone I₆, E-124
 Epothilone K, E-125
 Epothilone tetrahydrofuran, E-126
 Epoxyagroclavine I, A-205
 19,20-Epoxyakuammicine, A-228
 1,2-Epoxyambelline, A-683
 19,20-Epoxyapparine, A-1365
 19,21-Epoxyaspidospermidine, A-1490
 2',3'-Epoxyasteranthine, A-1510
 6,7-Epoxy-3-atropolyoxytropene, A-1364
 1,4-Epoxy-3-aza-*A*-homoandrostan-17-ol, E-127
 1,4-Epoxy-3-aza-*A*-homoandrostan-16-ol, S-44
 4,7'-Epoxy-3,8'-bilign-7-ene-3',4',5,9,9'-pentol, E-128
 4,9-Epoxy-3,4,7,14,15,16,20-cevaneheptol, G-70
 4,9-Epoxy-3,4,12,14,16,17,20-cevaneheptol, V-56
 4,9-Epoxy-3,4,14,15,16,20-cevanehexol, Z-36
 4,9-Epoxy-3,4,6,7,14,15,16,20-cevaneoctol, P-674
 4,7-Epoxy-3,14,16,20-cevanetetrol, N-116
 10,11-Epoxycolchicine (obsol.), O-220
 1,6-Epoxycomosivine, C-588
 19,20-Epoxyconoduramine, C-615
 17,19-Epoxycuran-3-ol, S-605
 19β,20β-Epoxycytochalasin C, C-944
 19,20-Epoxycytochalasin C, C-944
 19,20-Epoxycytochalasin D, Z-37
 Epoxycytochalasin H, E-129
 19,20-Epoxycytochalasin N, C-944
 19,20-Epoxycytochalasin Q, C-954
 19,20-Epoxycytochalasin R, C-954
 8,9-Epoxy-2-decene-4,6-diynoic acid, D-103
 2',3'-Epoxy-4-*O*-demethyl-*N*-methylatanine, A-1529
 1,19-Epoxydenudatine, K-48
 19,20-Epoxy-18-deoxycytochalasin Q, C-954
 19,20-Epoxy-18-deoxycytochalasin R, C-954
 17,19-Epoxy-19,20-dihydroakuammilan, L-16
 2,3-Epoxy-2,3-dihydro-2-hydroxy-1*H*-indole-3-acetic acid, M-758
 15,20-Epoxy-15,20-dihydro-12-hydroxysenecionan-11,16-dione, J-2
 17,19-Epoxy-19,20-dihydro-19-hydroxy-1,4,21-trimethyl-18-norsarpaganium, M-29
 17,23-Epoxy-5,6-dihydro-3-hydroxyveratraman-11-one, H-371
 17,19-Epoxy-19,20-dihydro-21-methyl-18-norsarpagan-17-ol, P-241
 6,17-Epoxy-19,20-dihydro-1-methylsarpagan-17,21-diol, E-130
 17,19-Epoxy-19,20-dihydro-1-methylsarpagan-21-ol, T-21
 12,16-Epoxy-16,17-dihydronapelline, E-131
 3,4-Epoxy-8,9-dihydroplartine, P-473
 17,23-Epoxy-5,6-dihydro-1,3,6-trihydroxyveratraman-11-one, V-74
 17,23-Epoxy-5,6-dihydroveratraman-2,3,6-triol, K-114
 14,15-Epoxy-3,5-dihydroxybufa-20,22-dienolide, E-132
 6,7-Epoxy-18,19-dihydroxy-16,18-dimethyl-10-phenyl[1]cytochalasin-13-ene-1,21-dione, H-473
 24,25-Epoxy-7,22-dihydroxyergosta-1,4-dien-3-one, P-295
 19,20-Epoxy-12,15-dihydroxy-11-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 19,20-Epoxy-12,15-dihydroxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 17,23-Epoxy-3,11-dihydroxyveratraman-6(5*H*)-one, Y-11
 19,20-Epoxy-11,12-dimethoxycajine, I-12
 8,10-Epoxy-3,7-dimethoxy-17-methylhasubanan-4,6,8-triol, L-244
 19,20-Epoxy-11,12-dimethoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 8-(2,3-Epoxy-3,3-dimethylallyloxy)-7-methoxy-*N*-methylflindersine, F-90
 16,17-Epoxy-4,21-dimethyl-7,20-cyclootidane-1,13,15-triol, G-139
 3,6-Epoxydodecanoic acid pyrrolide, H-269
 2',3'-Epoxyechiumine, E-31
 7,10-Epoxyepothilone, E-126
 1,19-Epoxy-20-ethyl-4-methylaconitane-7,8,14-triol, G-152
 5,11-Epoxy-1,4,6,8,9,14-eudesmanehexol, H-249
 Epoxyfunitremorgin, C, F-190
 4,7-Epoxyhexahydro-3*a*-methyl-2-phenyl-1*H*-isoin-dole-1,3(2*H*)-dione, P-17
 8,10-Epoxy-8-hydroxy-3,4-dimethoxy-17-methylhasubanan-7-one, M-233
 19,20-Epoxy-14-hydroxy-10,11-dimethoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 6,7-Epoxy-18-hydroxy-16,18-dimethyl-10-phenyl[1]cytochalasin-13,19-diene-1,21-dione, H-473
 19,20-Epoxy-15-hydroxycajine, I-12
 6,7-Epoxy-18-hydroxy-19-methoxy-16,18-dimethyl-10-phenyl[1]cytochalasin-13-ene-1,21-dione, H-473
 19,20-Epoxy-15-hydroxy-12-methoxycajine, I-12
 19,20-Epoxy-12-hydroxy-11-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 19,20-Epoxy-15-hydroxy-12-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 19,20-Epoxy-15-hydroxy-11-methoxyvomicine, I-12
 19,20-Epoxy-12-hydroxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 19,20-Epoxy-15-hydroxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 1,2-Epoxy-1-hydroxymethylpyrrolizidine, E-133
 19,20-Epoxy-15-hydroxynovacine, I-12
 23,26-Epoxy-3-hydroxy-16,28-secosolanid-22(28)-en-6-one, K-89
 12,13-Epoxy-19-hydroxysenecionan-11,16-dione, E-184
 14,15-Epoxy-11-hydroxytabersonine, T-11
 17,23-Epoxy-3-hydroxyveratraman-11-one, J-36
 19,20α-Epoxy-15-hydroxyvomicine, I-12
 1β,2β-Epoxyisoretrotronecanol, E-133
 14,15-Epoxykopsingine, K-66
 6,7-Epoxykopsinine, V-49
 6,7-Epoxylatrunclin A, L-61
 N²-(15,16-Epoxylinoleoyl)glutamine, G-108
 6,7-Epoxyllitorine, S-158
 6,21-Epoxy-17-methoxyaspidospermidine, B-63
 1,19-Epoxy-11-methoxydenudatine, K-48
 19,20-Epoxy-10-methoxycajine, I-12
 19,20-Epoxy-12-methoxycajine, I-12
 19,20-Epoxy-10-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 19,20-Epoxy-12-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
 1,2-Epoxy-1-methoxymethylpyrrolizidine, E-133
 6,17-Epoxy-11-methoxysarpagan, G-23
 19,20-Epoxy-11-methoxyvomicine, I-12
 2,7-Epoxy-1-methylaminopyrrolizidine, L-234
 1,4-Epoxy-20-methyl-3-aza-*A*-homopregn-14-en-16-one, S-45
 2-[3-(3,4-Epoxy-3-methyl-1-butenyl)-4-hydroxyphenyl]-*N*-methylethylamine, E-135
 5-(2,3-Epoxy-3-methylbutyl)-3-[2-hydroxy-1-(hydroxymethyl)-2-methylpropyl]-1*H*-indole, E-134
 3-(2,3-Epoxy-3-methylbutyl)-4-hydroxy-7-methoxy-1-methyl-2(1*H*)-quinolinone, T-582
 3-(2,3-Epoxy-3-methylbutyl)-5-(3-hydroxy-3-methyl-1-butenyl)-1*H*-indole, B-200
 3-(2,3-Epoxy-3-methylbutyl)-2-(3-hydroxy-3-methyl-1-butenyl)-1*H*-indole, H-257
 3-(2,3-Epoxy-3-methylbutyl)-4-hydroxy-*N*-methyl-2-quinolone, A-1529
 3-(2,3-Epoxy-3-methylbutyl)-5-(3-methyl-1,3-butadienyl)-1*H*-indole, B-200
 3-(2,3-Epoxy-3-methylbutyl)-5-(3-methyl-2-butenyl)-1*H*-indole, B-200
 1,17-Epoxy-1-methylasycarpidan, G-76
 1,2-Epoxy-2-methyl-4-(*N*-methyltyraminyl)-3-butene, E-135
 19,20-Epoxy-*N*-methyl-*sec*-pseudobrucine, I-12
 8,9-Epoxy-9-methylstreptimidone, M-561
 13,17-Epoxy-8,10(18)-myrsinadiene-3,5,7,14,15,17-hexol, E-136
 17,18-Epoxyxymythiazole, M-807
 16,17-Epoxyneopapakitamine A, D-273
 2,3-Epoxy-6,8-nonadiynoic acid 2-phenylethylamide, E-137
 1,2-Epoxy-6,8-nonadiynoic acid styrylamide, E-137
 2,3-Epoxy-6,8-nonadiynoic acid, E-137
 14,15-Epoxyneroscurinine, N-307
 19,20-Epoxyenovacine, I-12
 N²-(15,16-Epoxy-9,12-octadecadienoyl)glutamine, G-108
 2,3-Epoxy-4-oxo-7,10-dodecadienamide, C-286
 14,15-Epoxy-3-oxovincadifformine, O-210
 4,7'-Epoxy-3',4',5,9,9'-penta-hydroxy-3,8'-bilign-7-ene, E-128
 5,20-Epoxy-2,4,7,10,13-pentahydroxy-11-taxen-9-one, E-138
 2,3-Epoxy-3-phenylpropanoic acid, P-348
 3,4-Epoxy-2-piperidine, O-146
 3,4-Epoxy-piperimethystine, P-463
 1,2-Epoxy-powelline, C-743
 18,20-Epoxypregnane-3,14,20-triol, H-316
 2-(1,2-Epoxypropyl)quinoline, P-647
 Epoxyquinomicin A, C-289
 Epoxyquinomicin B, C-289
 Epoxyquinomicin C, C-289
 Epoxyquinomicin D, C-289
 6,7-Epoxyracemigerine, R-1
 1,6-Epoxyrobustivine, C-588
 16,17-Epoxyarscovagenine D, D-260
 Epoxysarmentosin, H-509
 17,19-Epoxyasarpagan-19(20*H*)-ol, T-616
 14,15-Epoxy-scandine, S-118
 16,23-Epoxy-16,28-secosolanidane-3,23-diol, D-99
 16,23-Epoxy-16,28-secosolanidane-3,23,27-triol, E-139
 16,23-Epoxy-16,28-secosolanid-22(28)-ene-3,20,23,27-tetrol, E-140
 14α,15α-Epoxytabersonine, T-11
 14β,15β-Epoxytabersonine, T-11
 5,20-Epoxy-11-taxene-1,2,4,7,9,10,13-heptol, E-141
 4,5-Epoxy-2-tetradecenoic acid, T-124
 8,10-Epoxy-3,4,7,8-tetramethoxy-17-methylhasubanan-6,16-dione, S-537
 8,10-Epoxy-3,4,7,8-tetramethoxy-17-methylhasubanan-6-one, S-537
 6,7-Epoxytropan-3-ol, S-158
 6,7-Epoxy-3-tropanyl 2,3-dihydroxy-2-phenylpropionate, A-1043
 Epoxytuberostemonone, T-666
 3,4-Epoxyvalerolactam, O-146
 3,17-Epoxyvobasan, T-10

- N,N'*-1,2-Ethanediyldisododecanamide, E-225
2,2'-(1,2-Ethanediyldi)bis[1-methyl-3-piperidinone], E-226
N,N'-1,2-Ethanediyldis(octadecanamide), E-225
N,N'-1,2-Ethanediyldis(1,3-propanediamine), B-153
Ethanolamine *O*-phosphate, A-768
Ethanolamine, A-767
Ethaphene, S-659
6-Ethenyl-3,4,4*b*,5,6,7,8,9,10*b*,12-decahydro-12-hydroxy-5-methyl-1*H*-pyrido[4',3':4,5]pyrano[3,2-*h*]isoquinoline-1,10(2*H*)-dione, E-227
1-Ethenyl-8,13-dihydroindolo[2',3':3,4]pyrido[1,2-*b*][2,7]naphthyridin-5(7*H*)-one, A-1019
5-Ethenyl-3,4-dihydro-2,7-naphthyridin-1(2*H*)-one, G-60
5-Ethenyl-3,4-dihydro-1*H*-pyrano[3,4-*c*]pyridin-1-one, G-59
1-Ethenyl-4,9-dimethoxy-9*H*-pyrido[3,4-*b*]indole, D-724
1-Ethenyl-4,7-dimethoxy-9*H*-pyrido[3,4-*b*]indole, D-723
1-Ethenyl-4,8-dimethoxy-9*H*-pyrido[3,4-*b*]indole, D-686
3-(2-Ethenyl-4,6-dimethylcyclohexyl)-1,4-dihydroxy-2(1*H*)-pyridinone, P-922
4-Ethenyl-3-ethoxy-1,3,4,5,6,7-hexahydro-8*H*-pyrano[3,4-*c*]pyridin-8-one, S-408
1-Ethenyl-4-methoxy-9*H*-pyrido[3,4-*b*]indole, M-290
[3-Ethenyl-3-methyl-5-(2-methyl-1-propenyl)-2-pyrrolidinyl]-1*H*-indol-3-ylmethanone, B-251
5-Ethenyl-4-methylthiazole, M-585
16-Ethenyloxayohimban-17-ol, E-228
2-(3-Ethenyl-4-piperidinyl)-1-[3-(2-hydroxyethyl)-1*H*-indol-2-yl]ethanone, C-445
3-(3-Ethenyl-4-piperidinyl)-1-(6-methoxy-4-quinolinyl)-1-propanone, Q-17
7-Ethenyl-2-(2-propenyl)-1-azaspiro[5.5]undecan-8-ol, H-309
5-Ethenyl-3-pyridinecarboxylic acid, V-141
1-Ethenyl-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, V-140
1-Ethenyl-9*H*-pyrido[3,4-*b*]indole-4,8-diol, D-686
1-Ethenyl-9*H*-pyrido[3,4-*b*]indole, V-139
1-Ethenyl-9*H*-pyrido[3,4-*b*]indol-8-ol, H-758
12-Ethenyl-5,6,13,13*a*-tetrahydro-2-hydroxy-3-methoxy-8*H*-isoquino[2,1-*b*][2,7]naphthyridin-8-one, A-235
Ethenyltrimethylammonium(1+), T-615
3-(2-Ethenyl-2,4,6-trimethylcyclohexyl)-1,4-dihydroxy-2(1*H*)-pyridinone, C-646
2-[[[(1-Ethenyl-1,5,9-trimethyl-4,8-decadienyl)sulfonyl]ethyl]guanidine], A-178
Ethesdanine, E-229
9-Ethoxyacridine, A-112
9-Ethoxyaristolactam I, T-251
4-Ethoxybenzo[*c*][2,7]naphthyridine, B-75
1-Ethoxycarbonyl- β -carboline, C-132
3-Ethoxycarbonyl- β -carboline, C-133
12-Ethoxycarbonylcytosine, C-938
N-Ethoxycarbonyldemecolcine, D-92
N-Ethoxycarbonyllaurotetanine, L-75
N-Ethoxycarbonylprolinamide, E-230
4-Ethoxycarbonyl-2(1*H*)-quinolinone, H-723
*N*¹⁶-Ethoxycarbonylroquefortine D, R-125
2-Ethoxycarbonyltetrahydroharman, T-202
2-Ethoxycarbonyl-1,2,3,4-tetrahydro-1-methyl- β -carboline, T-202
5-Ethoxycheleerythrine, C-356
3-Ethoxycoronaridine, C-657
8 α -Ethoxy-*N*-demethylpretazettine, P-626
8- β -Ethoxy-*N*-demethylpretazettine, P-626
8-Ethoxydihydrochelerythrine, D-407
Ethoxydihydrosanguinarine, D-510
15 β -Ethoxy-14,15-dihydroviraoallosecurinine, A-630
4-Ethoxy-7,8-dimethoxyfuro[2,3-*b*]quinoline, F-223
6-Ethoxy-1,2-dimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridinium(1+), C-356
12-Ethoxy-7,8-dimethoxy-5-methyl-2,3-methylene-dioxybenzo[*c*]phenanthridinium(1+), C-356
2-Ethoxy-*N,N*-dimethylethanamine, D-730
7-Ethoxy-2,6-dimethyl-3,5,8(2*H*)-isoquinoline-trione, M-624
3-Ethoxy-19-epiheyneanine, C-657
11-Ethoxy-3-formamidothionellin, I-213
4-Ethoxyfuro[2,3-*b*]quinoline, F-224
4-Ethoxyhexahydro-3*a*-hydroxy-1*H*-indole-2,6-dione, E-231
3-Ethoxyheptaneine, C-657
4-Ethoxy-1-hydroxy-4-methoxy-2,5-cyclohexadiene-1-acetamide, H-650
4-Ethoxy-8-hydroxy-1-methyl-3-(3-methyl-2-but-1-yl)-2(1*H*)-quinolinone, T-583
2-Ethoxy-4-hydroxyquinoline, Q-33
3-Ethoxy-1*H*-indole, I-104
16-Ethoxyisomatopenseine, M-120
13-Ethoxylupanine, H-554
4-Ethoxy-8-methoxyfuro[2,3-*b*]quinoline, F-213
N-[2-Ethoxy-2-(4-methoxyphenyl)ethyl]cinamide, A-153
3-(Ethoxymethyl)-6,7-dihydro-8(5*H*)-indolizone, D-447
2-Ethoxy-3-methylpyrazine, H-624
4-Ethoxy-2-methylpyridine, H-625
4-Ethoxy-2-methylquinoline, H-633
8-Ethoxy-3-oxo-1,2-dehydroretrosine, E-232
1-Ethoxyphenazine, P-327
8 α -Ethoxypreceiweline, P-626
6-Ethoxy-1*H*-purin-2-amine, G-210
Ethoxypyrazine, P-889
4-Ethoxy-3-pyridinecarboxylic acid, H-709
6-Ethoxy-3-pyridinecarboxylic acid, H-711
4-Ethoxyquinoline, H-633
4-Ethoxyquinazoline, Q-14
6-Ethoxy-4-quinolinecarboxylic acid, H-726
2-Ethoxy-4-quinolinecarboxylic acid, H-723
2-Ethoxyquinoline, Q-50
3-Ethoxyquinoline, Q-49
2-Ethoxy-4(1*H*)-quinolinone, Q-33
16-Ethoxystrychnine, P-726
Ethoxythiobinupharidine, T-380
3-Ethoxyvoacangine, V-170
Ethyl 2-aminobenzoate, A-706
Ethyl 2-(aminocarbonyl)-1-pyrrolidinecarboxylate, E-230
Ethyl β -carboline-2*N*-oxide-1-propionate, C-134
Ethyl β -carboline-1-propionate, C-134
Ethyl centralite, D-818
Ethyl *N*-docosanoylanthranyl, A-706
Ethyl harman-3-carboxylate, M-427
Ethyl hydrogen 2-aminoethyl phosphate, A-768
Ethyl iodoacetate, Y-163
Ethyl 3-methyl-9*H*-carbazole-9-carboxylate, M-424
Ethyl *N*-(2-phenylethyl)carbamate, P-343
Ethyl 2-pyridyl ketone, P-645
Ethyl (raframidin-15-yl)acetate, R-35
Ethyl 6-[[1,2,3,4-tetrahydro-7-methoxy-2-methyl-1-oxo-8-isoquinolinyl)oxy]-1,3-benzodioxole-5-carboxylate, C-279
Ethyl 3,4,5,6-tetrahydro-2-pyridyl ketone, T-217
Ethyl yohimbate, Y-13
8-*O*-Ethylaconorine, M-352
10-Ethylacridone, A-112
O-Ethylakagerine lactone, A-223
17-*O*-Ethylakagerine, A-222
O-Ethylalpinigenine, A-638
Ethylamine, E-233
Ethylaminesulfonic acid, A-765
2-(Ethylamino)-4,5-dihydroxybenzamide, A-737
2-Ethylaminoethylamine, E-225
19-*O*-Ethylangustoline, A-1020
N-Ethylanthalonine, A-1021
N-3'-Ethylaplysinopsin, A-1357
*N*⁵-Ethylaurantioclavine, A-1549
11-Ethyl-12-azacyclotetradec-5-en-14-olide, E-93
N-Ethyl-1,4-butanediamine, B-422
6-Ethylcanthin-4-one, N-302
5-Ethylcanthin-4-one, T-671
1-Ethyl- β -carboline, E-234
1-Ethyl- β -carboline-2*N*-oxide, E-234
2-*O*-Ethylcephalofortuosine, C-260
Ethylcycloonylprodigiosin, E-235
N-Ethyletylsine, C-938
2-Ethyldecahydro-5-methylquinoline, A-355
5-Ethyldecahydro-2-(2-penten-4-ynyl)quinoline, A-358
2-Ethyldecahydroquinoline, A-354
N-Ethyl-*N*-demethyldietylsine, D-370
N-Ethyl-*N*-desmethyldelphinine, M-312
Ethyladioline, C-115
Ethylidihydrocuprein, Q-20
 α -Ethyl-9,11-dihydro- α -hydroxy-8-hydroxymethyl-9-oxindolizinol[1,2-*b*]quinoline-7-acetic acid, C-71
3-Ethyl-1,3-dihydro-3-methoxy-2*H*-indol-2-one, E-236
3-Ethylidihydro-4-(1-methyl-1*H*-imidazol-5-yl)-methyl-2(3*H*)-furanone, P-424
3-Ethyl-4,5-dihydro-2-methylthiazolium(1+), D-479
5-Ethyl-3,4-dihydro-2-pentyl-2*H*-pyrrole, D-246
2-Ethyl-3,4-dihydro-5-pentyl-2*H*-pyrrole, D-246
1-Ethyl-3,7-dihydro-1*H*-purine-2,6-dione, X-5
3-Ethyl-2,7-dihydroxy-4,8-dimethoxyquinoline, H-60
6-Ethyl-3,4-dihydroxy-2-hydroxymethylpiperidine, E-244
20-Ethyl-6,8-dihydroxy-1-methoxy-4-methylheteratitan-14-one, H-157
1-Ethyl-4,8-dimethoxy- β -carboline, D-686
9-Ethyl-7,8-dimethoxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-223
1-Ethyl-4,8-dimethoxy-9*H*-pyrido[3,4-*b*]indole, D-686
5-Ethyl-2,3-dimethylpyrazine, E-239
2-Ethyl-3,5-dimethylpyrazine, E-237
3-Ethyl-2,5-dimethylpyrazine, E-238
N-Ethyl-*N,N'*-diphenylurea, D-818
1-Ethyldecahydro-9*a*-methylpyrido[2,1,6-*de*]quinolizin-2-ol, P-569
4-Ethyldecahydro-2,3,7,10-tetramethylcyclopenta[*b*]quinolizin-2-ol, A-509
4-Ethyldecahydro-3,7,10-trimethylcyclopenta[*b*]quinolizin-2-methanol, A-509
O-Ethyleburnamine, E-6
N,N'-Ethylenebis(diacetamide), E-225
N,N'-Ethylenebis(octadecanamide), E-225
Ethylene-diamine, E-225
trans-Ethylene-1,2-dicarboxylic acid, F-173
N-Ethyl-1,2-ethanediamine, E-225
8-Ethyl-5-(1-ethyl-1-butanyl)-6-methylindolizidine, E-240
8-Ethyl-5-(1-ethyl-1-butanyl)octahydro-6-methylindolizidine, E-240
2-Ethyl-3-[2-(3-ethylpiperidino)ethyl]indole, D-120
Ethylevoxine, F-223
1 α -*C*-Ethylflagamine, E-244
O-Ethylglauconine, G-87
6-*O*-Ethylguanane, G-210
N-Ethylhastaneine trachelanthate, E-241
8-Ethyl-5-(1,4-heptadien-6-ynyl)octahydroindolizidine, A-486
8-Ethyl-5-(1-hepten-6-ynyl)octahydroindolizidine, A-486
2-Ethyl-5-heptyl-3,4-dihydro-2*H*-pyrrole, D-246
5-Ethyl-2-heptyl- Δ^1 -pyrroline, D-246
2-Ethyl-5-heptyl- Δ^1 -pyrroline, D-246
4-Ethyl-2,4-hexadienoic acid, E-242
5-Ethyl-1,4,7,8,9,10-hexahydro-2*H*-3,7-methanoazacycloundecino[5,4-*b*]indole, C-522
13-Ethyl-4,5,11,12,13,13*a*-hexahydro-2*H*-3,13-methanoxireno[9,10]azacycloundecino[5,4-*b*]indol-5*a*(1*aH*)-ol, V-180
12-Ethyl-1,2,7*b*,8,12*b*,12*c*-hexahydro-2-methoxy-2-methylpyrido[4,3-*b*]pyrrolo[1,2,3-*lm*]carbazole, L-31
1-Ethyl-5-(7-hydroxy-1,3-benzodioxol-5-yl)-2-piperidinone, C-63
3-Ethyl-7-hydroxy-4,8-dimethoxy-2(1*H*)-quinolinone, H-60
22-Ethyl-19-hydroxy-15,16-dimethoxy-4,5-secoobscurinervan-21-*oic* acid γ -lactone, A-233
6-Ethyl-8-(2-hydroxyethyl)-5-pentylindolizidine, E-263
4-Ethyl-2-(hydroxyimino)-5-nitro-3-hexenamide, E-243

- 1-Ethyl-8-hydroxy-4-methoxy- β -carboline, D-686
 1-Ethyl-5-(3-hydroxy-4,5-methylenedioxyphenyl)-2-piperidone, C-63
 α -Ethyl-5-hydroxy-6-methyl-2-piperidinedecanol, P-655
 6-Ethyl-2-hydroxymethyl-3,4-piperidinediol, E-244
 2-Ethyl-6-hydroxymethyl-3,4,5-piperidinetriol, E-245
 4-Ethyl-4-hydroxy-1*H*-pyrano[3',4':6,7]indolizino[1,2-*b*]quinoline-3,14(4*H*,12*H*)-dione, C-70
 2-Ethyl-3-hydroxypyrazine, E-246
 2-Ethyl-4-hydroxyquinazoline, E-247
 3-Ethylidenebutyric acid, M-506
 3-Ethylidene-1,3-dihydro-2*H*-indol-2-one, E-248
 α -Ethylidene-2,5-dihydro-3-methyl-2,5-dioxo-1*H*-pyrrole-1-acetic acid, P-179
 5-Ethylidene-4,4a-epoxy-2,3,4,4a-tetrahydro-1-pyrindene, A-14
N'-Ethylidene-*N*-formyl-*N*-methylhydrazine, G-236
 2-Ethylidene-1,2,3,10,11,11a-hexahydro-8-hydroxy-7,11-dimethoxy-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one, T-424
 4-Ethylidene-2a,3,4,6,7,7*b*-hexahydro-3-hydroxy-1*H*-2-oxa-7a-azacyclopent[*cd*]inden-1-one, S-560
 4-Ethylidene-1,3,4,5,6,7-hexahydro-6-methylene-2,5-ethano-2*H*-azocino[4,3-*b*]indole, A-1365
 4-Ethylidene-1,3,4,5,6,8-hexahydro-2-methyl-13-methylene-1,5-methanoazocino[4,3-*b*]indol-7(2*H*)-one, E-193
 3-Ethylidenehexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione, E-249
 Ethylidenemethylhydrazinecarboxaldehyde, G-236
 5-Ethylidene-1,2,3,4,5,6,7,9-octahydro-6-(2-hydroxyethyl)-3-methyl-8*H*-azecino[5,4-*b*]indol-8-one, B-419
 α -Ethylidene-1,2,3,3a,4,5,6,7-octahydro-7-hydroxy-3-methyl-3,7a-diazacyclohept[*jk*]fluorene-5-acetaldehyde, A-222
 3-Ethylidene-1,2,3,4,6,7,12,12*b*-octahydroindolo[2,3-*a*]quinolizine, D-230
 4-Ethylidene-1,3,4,4a,5,7,12,12a-octahydro-2-methylpyrido[3',4':4,5]cyclohept[1,2-*b*]indol-6(2*H*)-one, M-293
 3-Ethylideneoxindole, E-248
 3-Ethylidene-2-pyrrolidinone, E-250
 7-Ethylidene-1a,2,3,7-tetrahydrocyclopent[*b*]oxireno[*c*]pyridine, A-14
 2-Ethylidene-1,2,3,11a-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one, P-660
 5-Ethyl-4*H*-indolo[3,2-*de*][1,5]naphthyridin-4-one, T-671
 3-Ethyl-12*H*-indolo[2,3-*a*]quinolizin-5-ium, F-84
 2-Ethyl-5-(3-indolyl)oxazole, A-622
 1-Ethyl-3-isobutylpiperidine, M-530
 2-Ethyllactic acid, H-572
 5-Ethyl-9-methoxycanthin-6-one, L-300
 1-Ethyl-4-methoxy- β -carboline, M-290
 9-Ethyl-7-methoxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-212
 3-Ethyl-6-methoxy-5-methyl-2-[2-(3-methyl-4-phenyl-3-butenyl)-4-oxazolyl]-4*H*-pyran-4-one, P-334
 2-Ethyl-11-methoxy-3-methyl-3*H*-[1,6]naphthyridino[6,5,4-*def*]quinoxaline, E-251
 3-Ethyl-3-methoxyoxindole, E-236
 3-Ethyl-2-methoxypyrazine, E-246
 1-Ethyl-4-methoxy-9*H*-pyrido[3,4-*b*]indol-8-ol, D-686
 2-Ethyl-3-methyl-2-butenedioic acid, E-252
 1-Ethyl-3-(2-methylbutyl)piperidine, E-253
N-Ethyl-*N*'-methyl-*N,N'*-diphenylurea, D-818
N-Ethyl-*N*'-methyl-1,2-ethanediamine, E-225
N-Ethyl-*N*'-methylthylenediamine, E-225
 2-Ethyl-3-methylfumaric acid, E-252
 Ethylmethylglycollic acid, H-572
 10-Ethyl-8-methyllobelidol, H-434
 2-Ethyl-3-methylmaleic acid, E-252
 2-Ethyl-3-methylmaleimide, E-252
 21-Ethyl-4-methyl-16-methylene-7,20-cycloveatchane-1,12,15-triol, N-30
 α -Ethyl-1-methyl- α' -phenyl-2,6-piperidinediethanol, L-86
 2-Ethyl-6-methyl-3,4,5-piperidinetriol, E-254
 1-Ethyl-3-(2-methylpropyl)piperidine, M-530
 2-Ethyl-5-methylpyrazine, E-255
 5-Ethyl-4-methyl-3-pyridinecarboxylic acid, E-257
 5-Ethyl-2-methylpyridine, E-256
 1-Ethyl-2-methylpyridinium bromide, M-537
 1-Ethyl-2-methylpyridinium iodide, M-537
 5-Ethyl-2-methyl-1*H*-pyrido[3,4-*a*]carbazolium(1+), E-258
 3-Ethyl-4-methyl-1*H*-pyrrole-2,5-dione, E-252
 2-Ethyl-4-methylquinazoline, E-259
 2-Ethyl-1-methyl-4(1*H*)-quinazolinone, E-247
 1-Ethyl-4-methylsulfonyl- β -carboline, E-260
 1-Ethyl-4-(methylsulfonyl)-9*H*-pyrido[3,4-*b*]indole, E-260
 2-Ethyl-4-methylthiazole, E-261
 1-Ethyl-2-methylthiazolium(1+), M-400
 Ethylmorphine, M-704
 (4-Ethylmorpholino-*N*⁺)trihydroboron, M-705
O-Ethylareline, N-47
 4-Ethyl-5-nitro-2-oxo-3-hexenamide oxime, E-243
O-Ethyl Nordictamine, F-224
O-Ethyl nor- γ -fagarine, F-213
 8-Ethyl norlobelol I, H-435
N-Ethyl normicotine, P-957
O-Ethyl norskimmianine, F-223
O-Ethyl nupharamine, N-340
 7-Ethyl-1,4,5,6,7,8,10,14*b*-octahydro-1,2-dioxo-2*H*-3,7-methanoazacycloundecino[5,4-*b*]indole-9-carboxylic acid methyl ester, E-192
 7-Ethyl-1,4,5,6,7,8,9,10-octahydro-2*H*-3,7-methanoazacycloundecino[5,4-*b*]indole, Q-6
 5-Ethyl-1,4,5,6,7,8,9,10-octahydro-2*H*-methanoazacycloundecino[5,4-*b*]indol-6-ol, C-104
 7-Ethyl-1,2,4,5,6,7,8,9-octahydro-14*bH*-3,7-methanoazacycloundecino[5,4-*b*]indol-14*b*-ol, R-72
 13-Ethyl-1*a*,4,5,10,11,12,13,13*a*-octahydro-2*H*-3,13-methanoaxireno[9,10]azacycloundecino[5,4-*b*]indole, V-179
 1*b*-Ethyl-1*a*,1*b*,3,4,8*c*,9,10,10*a*-octahydro-2*H*-oxireno[4,5-*b*]pyrido[3,2-*c*]carbazole, E-262
 6-Ethyl octahydro-5-pentyl-8-indolizineethanol, E-263
 1-Ethyl octahydro-4-(4-pentynyl)-2*H*-quinolizine, E-264
 1-Ethyl octahydro-4-(2-propenyl)-2*H*-quinolizine, E-265
O-Ethyl oreoginine, G-87
 5-Ethyl-1-oxa-4-azacyclopentadec-10-en-15-one, E-93
 5-Ethyl-2-pentyl- Δ^1 -pyrroline, D-246
 2-Ethyl-5-pentyl- Δ^1 -pyrroline, D-246
 1-Ethyl-4-(4-pentynyl)quinolizidine, E-264
 8-Ethyl-10-phenyldehydrolobelidol, L-221
 8-Ethyl-10-phenyldehydrolobelionol, L-221
 8-Ethyl-10-phenyllobelidol, L-86
 5-Ethyl- α -picoline, E-256
 α -Ethyl-2-piperidineethanol, H-435
 α -Ethyl-2-piperidinemethanol, C-603
 1-Ethyl-4-(2-propenyl)quinolizidine, E-265
 8-Ethyl-10-propyllobelidol, H-433
 2-(1-Ethylpropyl)-4-methoxyquinoline, E-266
 2-(1-Ethylpropyl)-1-methyl-4(1*H*)-quinolinone, E-267
 3-Ethylpyrazinol, E-246
 3-Ethyl-2(1*H*)-pyrazinone, E-246
 5-Ethyl-3-pyridinecarboxylic acid, E-269
 3-Ethylpyridine, E-268
 1-Ethyl-9*H*-pyrido[3,4-*b*]indole, E-234
 1-Ethyl-1*H*-pyrrole-2-carboxaldehyde, P-942
 3-Ethyl-2-pyrrolidinone, E-270
 3-(1-Ethyl-2-pyrrolidinyl)pyridine, P-957
 2-Ethyl-4-quinazolinol, E-247
 2-Ethyl-4(3*H*)-quinazolinone, E-247
 1-Ethyl-4(1*H*)-quinazolinone, Q-14
 3-Ethyl-4(3*H*)-quinazolinone, Q-14
*N*¹-Ethylquinine, Q-20
O-Ethylrhoegenine, R-86
*N*¹⁶-Ethylrhoquefortine C, R-125
 8-*O*-Ethylsachaconitine, M-355
 21-*O*-Ethylstrictosamide aglycone, V-126
 β -*O*-Ethylsinephrine, S-659
N-Ethyltetrahydrocystine, T-149
 2-Ethyl-1,2,3,4-tetrahydro-6,7-dihydroxy-1,1-dimethylisoquinoline, T-152
 4-Ethyl-1,6,7,8-tetrahydro-6,8-dimethylcyclopent[*g*]indole, T-594
 2-Ethyl-1,2,3,4-tetrahydro-1,1-dimethyl-6,7-isoquinolinediol, T-152
 3-Ethyl-1,2,3,4-tetrahydro-1,3-dimethylisoquinoline, E-271
 α^2 -Ethyl-1,2,3,6-tetrahydro- α^6 ,1-dimethyl-2,6-pyridinediethanol, H-434
 2-Ethyl-5,6,7,8-tetrahydro-3-methylimidazo[1,2-*a*]pyridine, C-199
 3-Ethyl-5,6,7,8-tetrahydro-2-methylimidazo[1,2-*a*]pyridine, C-198
 α^6 -Ethyl-1,2,3,6-tetrahydro-1-methyl- α^2 -propyl-2,6-pyridinediethanol, H-433
 5-[(4-Ethyltetrahydro-5-oxo-3-furanyl)methylidene]-1-methyl-2,4-imidazolidinedione, E-272
 4-Ethyl-1,6,7,8-tetrahydro-5,6,8-trimethylcyclopent[*g*]indole, H-141
 2-(5-Ethyl-1,3,4,7-tetramethyl-1*H*-inden-6-yl)-1*H*-pyrrole, T-593
 8-*O*-Ethyl-3,14,15-tribenzoylaconine, M-331
 2-Ethyl-3,4,5-trihydroxy-6-hydroxymethylpiperidine, E-245
 2-Ethyl-3,4,5-trihydroxy-6-methylpiperidine, E-254
 Ethyltrimethylpyrazine, E-273
O-Ethyltriniifoline, Z-1
 1-Ethylxanthine, X-5
 Ethylenedicarboxylic acid, B-447
 Etilevodopa, A-745
 Etioline 3-glucoside, S-199
 Etioline, S-199
 Etioline, S-199
 Eubron, E-6
 Euchrestifoline, E-274
 Euchrestine A, E-275
 Euchrestine B, E-276
 Euchrestine C, E-276
 Euchrestine D, E-277
 Euchrestine E, E-276
 Eudermol, N-190
 Eudisin A, I-119
 Eudisin B, I-119
 Eudistalbin A, E-278
 Eudistalbin B, E-279
 Eudistomidin A, E-280
 Eudistomidin B, E-281
 Eudistomidin C, E-282
 Eudistomidin D, B-321
 Eudistomidin E, E-283
 Eudistomidin F, E-283
 Eudistomin A, E-285
 Eudistomin B, E-286
 Eudistomin C, E-287
 Eudistomin D, B-321
 Eudistomin E, E-287
 Eudistomin F, E-287
 Eudistomin G, E-285
 Eudistomin H, E-288
 Eudistomin I, E-288
 Eudistomin J, B-322
 Eudistomin K sulfoxide, E-287
 Eudistomin K, E-287
 Eudistomin L, E-287
 Eudistomin M, E-285
 Eudistomin N, B-299
 Eudistomin O, B-300
 Eudistomin P, E-285
 Eudistomin Q, E-285
 Eudistomin R, E-289
 Eudistomin S, E-289
 Eudistomin T, E-289
 Eudistomin U, I-119
 Eudistomin V, E-287
 Eudistomin V, E-288
 Eudistomin W, E-290
 Eudistomin X, E-284
 Eudistomin Y₁, H-411
 Eudistomin Y₂, H-411

- Eudistomin Y₃, H-411
 Eudistomin Y₄, H-411
 Eudistomin Y₅, H-411
 Eudistomin Y₆, H-411
 Eudistomin Y₇, H-411
 Eudistone A, E-291
 Eudistone B, E-292
 Eugenine A, E-293
 Eugenine B, E-294
 Eugenine, L-347
 Euglenapterin, A-923
 Eumydrin, T-632
 Euojaonin A, P-247
 Euojaonin C, P-247
 Euojaonin D, P-252
 Euojaonin F, P-252
 Euojaonin G, P-252
 Euojaonin I, P-247
 Euojaonin J, P-243
 Euojaonin K, P-252
 Euojaonin L, P-247
 Euojaonin M, P-247
 Euojaonin N, P-243
 Euonine, P-252
 Euonymine, P-247
 Euonyminol, N-281
 Euophelline, P-247
 Euoverrine A, P-247
 Euoverrine B, P-252
 Euphocharacin A, H-252
 Euphocharacin B, H-252
 Euphocharacin C, H-252
 Euphocharacin D, H-252
 Euphocharacin E, P-222
 Euphocharacin F, P-222
 Euphocharacin G, P-222
 Euphocharacin H, P-222
 Euphocharacin I, P-222
 Euphocharacin J, P-222
 Euphocharacin K, P-222
 Euphocharacin L, H-252
 Euphococcinine, M-389
 Euphodendroidin C, H-120
 Euphodendroidin G, H-120
 Euphodendroidin I, H-120
 Euphorbia factor L₁₁, T-263
 Euphosalicin, E-295
 Euphosine, E-296
 Euphyllin, T-360
 Eupolauramine, E-297
 Eupolauridine *N*-oxide, I-65
 Eupolauridine-di-*N*-oxide, I-65
 Eupolauridine, I-65
 Eupomatidine 1, S-49
 Eupomatidine 2, S-49
 Eupomatidine 3, S-49
 Euquinine, Q-20
 Euroline, E-298
 Europine *N*-oxide, E-299
 Europine, E-299
 Eustifoline A, E-300
 Eustifoline B, E-301
 Eustifoline C, E-302
 Eustifoline D, M-465
 Eusynstyelamide, E-303
 Euthyroideone A, E-304
 Euthyroideone B, E-304
 Euthyroideone C, E-304
 Euxylophoricine A, E-305
 Euxylophoricine B, E-305
 Euxylophoricine C, E-306
 Euxylophoricine D, E-307
 Euxylophoricine E, E-307
 Euxylophoricine F, E-308
 Euxylophorine B, E-309
 Euxylophorine C, E-309
 Euxylophorine D, E-309
 Euxylophorine, E-309
 Evellerine, F-212
 Ever 1, P-217
 Ever 6, H-248
 Ever 7, H-248
 Evizon, S-485
 Evocarpine, H-750
 Evodiamide, M-582
 Evodiamine, E-310
 Evodiaxinine, E-311
 Evodine, F-223
 Evoeuropine, A-1444
 Evoevoline, E-312
 Evolatine, F-221
 Evoline, E-313
 Evolitrine, F-212
 Evomeliaefolin, E-314
 Evomerrine, E-315
 Evomine, E-316
 Evonimine, P-252
 Evonine, P-248
 Evonoline, P-248
 Evopine, E-317
 Evoprenine, T-516
 Evorine, P-248
 Evoxanthidine, T-516
 Evoxanthine, T-516
 Evoxine acetate, F-223
 Evoxine, F-223
 Evoxoidine, F-223
 Evozine, P-248
 Exal, V-110
 Exceconidine, A-92
 Exceconitine, M-309
 Excelsine, D-187
 Excelsinine, M-291
 Excentricine, E-318
 Exfoliazone, A-801
 Exicarbamate, S-425
 Exiguamide, S-425
 Exiguamine A, E-319
 Exigurin, S-425
 Eximidine, E-320
 Eximine, D-346
 Exochomine, E-321
 Exozoline, C-814
 Ezochasmaconitine, M-323
 Ezochasmanine, M-303
 F 1778, A-1163
 F 1839A, A-1164
 F 1839B, A-1164
 F 10C, A-1165
 F 1839C, A-1164
 F 1839D, A-1164
 F 1839E, A-1164
 F 1839F, A-1164
 F 2787, H-729
 F 1839J, A-1164
 F 4708, S-490
 FA 4283, A-1166
 Faberidine, T-333
 Faberonine, H-366
 Fabianine, F-1
 Fabioline, B-141
 Factor M, O-131
 Factor S, B-139
Lactobacillus casei Factor, P-772
 Fademin, F-79
 FAD, F-79
 Fagara base, H-751
cis-Fagaramide, M-457
trans-Fagaramide, M-457
 Fagaridine, C-356
 Fagarine I, F-2
 Fagarine II, F-3
 Fagarine III, F-4
 α -Fagarine, F-2
 β -Fagarine, F-223
 γ -Fagarine, F-213
 Fagaronine, F-5
 Fagomine, H-614
 Fagopyrine, F-6
 Falaconitine, M-365
 Falcarindiol, H-116
 Falcatine, F-7
 Falconericine, M-323
 Falconeridine, M-323
 Falconerine 8-acetate, M-303
 Falconerine, M-303
 Faleoconitine, M-336
 Fangchinoline 2' α -*N*-oxide, T-286
 Fangchinoline 2' β -*N*-oxide, T-286
 Fangchinoline, T-286
 Fangchirine, D-148
 Fannine, A-1490
 Faralaoitrine, R-432
 Farcinicin, A-759
 Farcinicine, A-759
 Fareanine, F-8
 Farfugine, H-233
 Fargesine, F-9
 Farinosone A, F-10
 Farinosone B, F-10
 Farinosone C, F-11
 Farmiserina, A-832
 2,6,10-Farnesatrien-1-amine, F-12
 Farnesylamine, F-12
 2-Farnesylindole, T-604
 3-Farnesylindole, T-605
 Fascaplysin A, F-13
 Fascaplysin B, F-13
 Fascaplysin, F-13
 Fasciospongine A, F-14
 Fasciospongine B, F-14
 Fascicularine, F-15
 Fasmerianamine A, F-16
 Fasmerianamine B, F-16
 Fastigiatine, F-17
 Fastrine, F-18
 Fastudine, F-19
 Fastunine, F-20
 Fastusidine, F-21
 Fastusine, F-22
 Fastusinine, F-23
 Fauridine, F-24
 Faurine, F-25
 Fauripavine, F-26
 Faurithaline, F-27
 Fawcettidine, F-28
 Fawcettiine, F-29
 Fawcettimine, F-30
 4849F, B-73
 F6, B-403
 31F508 α 1, B-140
 31F508 α 2, B-140
 31F508 β 1, B-140
 31F508 β 2, B-140
 FCRC A48, F-152
 FD 838, A-1167
 FD 839, A-1168
 FD 991, Q-58
 Febrifugine, F-31
 Fedamazine, F-32
 Feldamycin, F-33
 Felipealbazine A, B-386
 Felipealbazine B, B-389
 Fellutanine D, G-234
 FEMA 2421, A-706
 FEMA 2682, A-706
 FEMA 3147, A-74
 FEMA 3184, A-74
 FEMA 3202, A-74
 FEMA 3251, A-73
 FEMA 3813, A-765
 FEMA 3819, A-1413
 FEMA 2224, C-15
 FEMA 2787, C-100
 FEMA 3404, C-100
 FEMA 2364, D-115
 FEMA 2614, D-889
 FEMA 3272, D-774
 FEMA 3336, D-392
 FEMA 3540, D-775
 FEMA 3555, D-247
 FEMA 3798, D-535
 FEMA 3898, D-500
 FEMA 3913, D-132
 FEMA 3916, D-393
 FEMA 3917, D-426
 FEMA 3150, E-237
 FEMA 3154, E-255
 FEMA 3394, E-268

- FEMA 3546, E-256
 FEMA 3680, E-261
 FEMA 2488, F-173
 FEMA 3189, F-201
 FEMA 3284, F-200
 FEMA 3285, G-107
 FEMA 3684, G-108
 FEMA 2011, H-258
 FEMA 2832, H-221
 FEMA 3204, H-495
 FEMA 3205, H-495
 FEMA 2593, I-79
 FEMA 2978, I-301
 FEMA 3554, I-283
 FEMA 3940, I-285
 FEMA 3297, L-153
 FEMA 2655, M-72
 FEMA 2718, M-384
 FEMA 3019, M-474
 FEMA 3102, M-409
 FEMA 3132, M-273
 FEMA 3133, M-483
 FEMA 3219, M-420
 FEMA 3309, M-536
 FEMA 3313, M-585
 FEMA 3433, M-272
 FEMA 3463, M-505
 FEMA 3716, M-569
 FEMA 3945, M-240
 FEMA 2815, O-37
 FEMA 3035, O-32
 FEMA 2878, P-336
 FEMA 2908, P-454
 FEMA 2909, P-467
 FEMA 2966, P-894
 FEMA 3220, P-342
 FEMA 3302, P-889
 FEMA 3383, P-237
 FEMA 3523, P-949
 FEMA 2975, Q-20
 FEMA 2976, Q-20
 FEMA 2977, Q-20
 FEMA 3470, Q-29
 FEMA 3237, T-284
 FEMA 3241, T-600
 FEMA 3244, T-613
 FEMA 3322, T-366
 FEMA 3591, T-355
 FEMA 3736, T-700
 Fenactil, C-417
 Fendleridine, A-1490
 Fendlerine, A-1490
 Fendispermine, A-1498
 Fenfangjine A, T-286
 Fenfangjine B, T-286
 Fenfangjine C, T-286
 Fenfangjine D, C-171
 Fenfangjine F, S-546
 Fenfangjine G, S-311
 Fenfangjine H, F-34
 Fenfangjine I, F-34
 Feretine, A-151
 Ferriaspergillin, A-1475
 Ferribactin G173, F-35
 Ferricrocin, F-36
 Ferrimycin A₁, F-37
 Ferrineoaspergillin, F-76
 Ferrioxamine A₂, F-38
 Ferrioxamine B, F-39
 Ferrioxamine C, F-40
 Ferrioxamine D, F-39
 Ferrioxamine D₁, F-39
 Ferrioxamine D₂, F-41
 Ferrioxamine E, F-42
 Ferrioxamine F, F-43
 Ferrioxamine H, F-44
 Ferrithiocin, D-235
 Ferropyrimine, D-496
 Ferrorosamine A, D-496
 Ferroverdin A, F-45
 Ferroverdin B, F-45
 Ferroverdin C, F-45
 Ferroverdin, F-45
 Ferruginine, F-46
 Ferruginine, F-47
 N¹-*trans*-Feruloylagmatine, A-719
 2'-*O-trans*-Feruloylalangiside, A-239
 N-Feruloylanthranilic acid, A-1563
 O-Feruloylcassine, C-183
 Feruloylcholine, C-419
 6'-*O*-Feruloyldemethylalangiside, A-239
 2'-*O-trans*-Feruloyldemethylalangiside, A-239
 2'-*O-trans*-Feruloyl-3-*O*-demethyl-2-*O*-methylalangiside, A-239
 N-*trans*-Feruloyl-3,5-dimethoxytyramine, T-576
 N-*trans*-Feruloyldopamine, D-920
 N-(4'-*O*-Feruloylferuloyl)tyramine, H-271
 N²-Feruloylhistamine, C-455
 N-Feruloyl-5-hydroxyanthranilic acid, A-789
 N-*trans*-Feruloyl-N¹-(4-hydroxybenzoyl)putrescine, C-456
 1-Feruloyl-2-hydroxyputrescine, D-248
 6'-*trans*-Feruloyllyaloside, L-316
 N-*cis*-Feruloyl-3-methoxytyramine, H-566
 N-*trans*-Feruloyl-3-methoxytyramine, H-566
 N-*trans*-Feruloyl-3-*O*-methyl-dopamine, H-566
 N-*trans*-Feruloyl-4-*O*-methyl-dopamine, H-565
 N-*trans*-Feruloyl-β-*O*-methyl-octopamine, O-63
 5-*O*-Feruloylnigrumins, H-509
 N-*trans*-Feruloylnormetanephrine, N-303
 N-*cis*-Feruloyloctopamine, O-63
 N-*trans*-Feruloyloctopamine, O-63
 N-*trans*-Feruloyl-*R*-octopamine, O-63
 N-*trans*-Feruloyl-*S*-octopamine, O-63
 ω-Feruloyloxylupinane, O-58
 N-*trans*-Feruloyl-β-*O*-propyloctopamine, O-63
 Feruloylputrescine, C-456
 N^b-(*E*)-Feruloylserotonin, H-755
 N^b-(*Z*)-Feruloylserotonin, H-755
 2'-*O*-Feruloylsimmondsin, S-302
 Feruloylspermidine, S-398
 N^b-Feruloyltryptamine, T-639
 N-*cis*-Feruloyltyramine, H-667
 N-*trans*-Feruloyltyramine, H-667
 Feruperine, C-710
 Fervenulin, F-48
 Fervenulone, D-498
 Festucine, L-234
 Festuclavine, F-49
 Fetidine, T-333
 Fettingine, P-280
 FG 7142, C-133
 FI 6426, D-866
 Fiancine, H-9
 Fibraminine, F-50
 Fibranine, F-51
 Ficellomycin, F-52
 Ficine, F-53
 Ficuseptine A, P-228
 Ficuseptine B, F-55
 Ficuseptine C, F-55
 Ficuseptine D, I-342
 Ficuseptine, F-54
 Fildesin, V-110
 Filifiline, D-878
 Filifoline, A-683
 Filiformine, T-315
 Finaconitine, A-94
 Finetiadine, P-776
 Finetianine, N-30
 Firefly luciferin, L-282
 Firefly oxyluciferin, F-56
 Fiscalin A, F-57
 Fiscalin B, F-58
 Fiscalin C, F-59
 Fischerellin A, F-60
 Fischerellin B, F-61
 Fischerindole L, F-63
 Fischerin, F-62
 Fisostin, P-400
 Fissiceine, O-170
 Fissicesine *N*-oxide, F-64
 Fissicesine, F-64
 Fissilandione, F-65
 Fissisaine, A-1042
 Fissistigine A, D-846
 Fissistigine B, F-66
 Fissistigine C, F-80
 Fissoldhimine, F-67
 Fissoldine, D-846
 Fissumine, D-585
 Fistularin 1, F-68
 Fistularin 2, F-69
 Fistularin 3, F-70
 Fistulosin, H-646
 FK 1000, S-311
 FK 3000, S-311
 FK317, A-1177
 FK973, A-1177
 FK409, E-243
 FK1 0550, A-758
 FL 120B', A-1169
 FL 120B, A-1169
 FL 657C, T-496
 Flabellazole A, S-610
 Flabellazole B, S-610
 Flabellidine, F-71
 Flabelliformide A, F-72
 Flabelliformide B, F-72
 Flabelliformine, F-73
 Flabelliformine, F-74
 Flabelline, F-75
 Flaconitine, A-104
 Flagecidin, A-1044
 Flamitajin, F-79
 Flamin F, F-79
 Flavacol, F-76
 Flavaconidine, M-332
 Flavaconijine, M-343
 Flavaconitine, M-332
 Flavadine, N-30
Aglaia dasyclada Flavagline compound 1, F-77
Aglaia dasyclada Flavagline compound 2, F-78
 Flavamine, N-30
 Flavin, A-632
 (+)-Flavinantifone, F-80
 (-)-Flavinantifone, F-80
 Flavinat, F-79
 Flavindin, I-111
 Flavine adenine dinucleotide, F-79
 Flavinine, F-80
 Flavipucine, F-81
 Flavisiamine A, M-296
 Flavisiamine C, M-296
 Flavisiamine D, M-296
 Flavitan, F-79
 Flavocarpine, F-82
 Flavocristamide A, S-634
 Flavocristamide B, S-634
 Flavolipids, F-83
 Flavopereirine, F-84
 Flazine methyl ester, F-85
 Flazine methyl ether, F-85
 Flazine, F-85
 Fleephilone, F-86
 Flexamine, F-87
 Flexicorine, F-88
 Flexine, F-89
 Flexinine, C-748
 Flindersiamine, F-217
 Flindersine, F-90
 Floramultine, P-221
 Floramultinone, F-91
 Floribundine, D-531
 Floricaline, O-92
 Floridanine, O-92
 Floridimine, F-92
 Floridine, F-92
 Floridinone, F-92
 Floripavidine, D-531
 Floripavine, S-39
 Florosenine, O-137
 Flueggeine, F-95
 Flueggeine A, F-93
 Flueggeine B, F-94
 Flueggeine ether, N-307
 Flueggeinol, N-307
 Flueggeine, F-95
 Fluopsin B, F-96

- Fluopsin C, F-97
 Fluopsin F, F-96
 Fluopsin N, F-98
 Fluorocarpamine *N*-oxide, F-99
 Fluorocarpamine, F-99
 C-Fluorocurarine, N-301
 C-Fluorocurine, F-100
 Fluorocurine, F-100
 ψ -Fluorocurine, F-101
 C-Fluorocurinine, F-102
 Fluorodaturatin, H-444
 5-Fluoro-3,4-dihydro-2,4-dioxo-1(2*H*)-pyrimidin-2-ylideneacetic acid, F-103
 5-Fluoro-2,4-dihydroxypyrimidine, F-104
 Fluoroindolocarbazole A, R-32
 Fluoroindolocarbazole B, R-32
 Fluoroindolocarbazole C, R-32
 5-Fluoro-1-(methoxycarbonylmethyl)uracil, F-103
 Fluoroplex, F-104
 5-Fluoro-2,4-pyrimidinediol, F-104
 5-Fluoro-2,4(1*H*,3*H*)-pyrimidinedione, F-104
 4-Fluoropyrrolnitrin, P-961
 Fluorosolimoines, F-105
 Fluorouracil, F-104
 5-Fluorouracil, F-104
 Fluracil, F-104
 Flustrabromine, B-316
 Flustramide A, F-106
 Flustramide B, F-108
 Flustramine A, F-106
 Flustramine B, F-108
 Flustramine C, F-106
 Flustramine D *N*-oxide, F-107
 Flustramine D, F-107
 Flustramine E, F-108
 Flustraminol A, F-109
 Flustraminol B, F-110
 Flustrarine B, F-111
 Flutimide, F-112
 Fluvibactin, F-113
 Fluviol A, M-285
 Fluviol B, M-285
 Fluviol C, M-285
 Fluviol D, F-114
 Fluviol E, M-285
 Fluvirucin B, F-115
 Fluvirucin B₂, F-115
 Fluvirucin B₄, F-115
 Fluvirucin B₅, F-115
 Fmoc-L-DOPA, A-745
 FO 7314, A-1407
 FO 2047, P-333
 Foetidine I, F-116
 Foetidine II, F-116
 Foetidine, T-333
 Folicin, P-772
 Foliacraline, A-641
 Foliamin, P-772
 Folic acid conjugate, P-772
 Folic acid, P-772
 Folicangine, F-117
 Folicanthine, C-382
 Folidine, F-122
 Folidine, Q-47
 Folidine, R-92
 Folidine, F-118
 Folidine, F-119
 Folidine, Q-47
 Folidine, F-120
 Folinic acid, F-121
 Folioidine acetone, F-122
 Folioidine, F-122
 Folioidine, G-171
 Foliophorine, F-122
 Folidine, F-123
 Folisine, I-268
 Folvite, P-772
 Fontaineine, F-124
 Fontaphilline, F-125
 Fontonamide, F-126
 Forbagline A, F-127
 Forbagline B, F-127
Aglaia dasyclada Forbagline-like compound, F-128
 Fordimine, H-376
 Fordine, H-375
 Foresaconitine, M-323
 Foresticine, M-323
 Forestine, M-312
 7-Formamidino-7-deazaguanosine, A-1387
 10 α -Formamidoallooromadendrane, A-1571
 1-Formamidoaromadendrane, A-1447
 3-Formamido-8,10-bisaboladiene, A-709
 4-Formamido-11-eudesmene, A-775
 11-Formamido-5-eudesmene, F-129
 6-Formamido-4(15)-eudesmene, I-218
 4-Formamido-11-gorgonene, A-777
 2-(Formamido)-3-(4-hydroxyphenyl)-2-propenitrile, C-826
 3-Formamidotheonellin, I-213
 Formobactin, F-130
 Formosanine, M-661
 Formouregine, F-104
 Formycin A, F-131
 Formycin B, F-132
 Formycin, F-131
N-Formylalbaine, A-243
*N*⁴-Formylalstonisine, A-654
 3-(Formylamino)-6,7-dihydroxycoumarin, A-738
 4-Formylaminomaalane, E-100
 5-(Formylamino)-6-(methylamino)-2,4-pyrimidin-2-one, H-761
 5-(Formylamino)-6-(methylamino)uracil, H-761
 3-(Formylaminomethyl)-1-methoxyindole, I-97
N-Formylammodendrine, A-930
N-Formylangustifoline, A-1013
N-Formylanhalamine, T-236
N-Formylanhalinine, T-236
N-Formylanhalonidine, T-237
N-Formylanhalonine, A-1021
N-Formylanolobine, A-1066
N-Formylanonaine, R-120
N-Formylbuxamine B, B-455
*N*²⁰-Formylbuxaminol E, B-456
N-Formylbuxifoline, B-471
 2-Formylcarbazole (obsol.), C-121
 3-Formylcarbazole, C-121
 1-Formyl- β -carboline, C-131
 4-Formylcarbostyryl, H-721
N-Formylchoneomorphine, P-601
 Formylcinegalleine, H-554
 8-Formylconfusamine, E-315
N-Formylconkurchine, C-610
N-Formylconvolvine, C-633
N-Formylcorydamine, C-678
N-Formylcycloimicrobuxeine (incorr.), C-882
N-Formylcyclovirobuxeine B, C-915
N-Formylcyclovirobuxine B (incorr.), C-915
N-Formylcylindrocarine, C-929
N-Formylcylindrocarinol, L-177
N-Formylcystisine, C-938
N-Formyldehydroanonaine, D-166
 7-Formyldehydrohernangerine, F-133
 7-Formyldehydroornantentine, F-134
 7-Formyldehydroovigerine, F-135
N-Formyldehydroovigerine, O-143
 7-Formyldehydrothalicimidine, F-136
N-Formyldehydrotryptamine, D-172
N-Formyldemecolcine, D-92
N-Formyl-*N*-demethylharappamine, M-667
N-Formyl-3-demethylmescaline, H-465
N-Formyl-14,15-didehydroaspidofractinine, A-1494
*N*¹-Formyl-14,15-didehydro-12-hydroxyaspidofractinine, A-1494
 4-Formyl-2,3-dihydro-1*H*-azepine-2,7-dicarboxylic acid, M-770
 1-Formyl-2,3-dihydro-3-(2-hydroxymethyl-4-methoxyphenyl)-6-methoxy-4(1*H*)-quinazolinone, D-369
 1-Formyl-6,7-dihydro-7-oxo-5*H*-pyrrolizine, D-14
 1-Formyl-6,7-dihydro-5*H*-pyrrolizine, D-14
N-Formyl-1,2-dihydrorenierol acetate, R-45
N-Formyl-1,2-dihydrorenierol propionate, R-45
 (+)-*N*-Formyl-1,2-dihydrorenierone, R-45
 (-)-*N*-Formyl-1,2-dihydrorenierone, R-45
 14-Formyldihydrorutaeacarpine, D-508
 6-Formyl-1,7-dihydroxycarbazole, D-557
 3-Formyl-1,6-dihydroxycarbazole, D-552
 3-Formyl-1,7-dihydroxy-9*H*-carbazole, D-553
 3-Formyl-1,8-dihydroxycarbazole, D-554
 3-Formyl-2,6-dihydroxycarbazole, D-555
 3-Formyl-2,7-dihydroxycarbazole, D-556
 1-Formyl-2,7-dihydroxy-6-methylcarbazole, D-619
 3-Formyl-2,7-dihydroxy-1-prenylcarbazole, H-125
 3-Formyl-2,8-dihydroxy-1-prenylcarbazole, H-125
N-Formyl-11,12-dimethoxyaspidofractinine, A-1494
 3-Formyl-1,6-dimethoxy-9*H*-carbazole, D-552
 3-Formyl-1,8-dimethoxycarbazole, D-554
 3-Formyl-2,6-dimethoxycarbazole, D-555
 3-Formyl-2,7-dimethoxycarbazole, D-556
 1-Formyl-2,7-dihydroxy-6-methylcarbazole, D-619
 9-(2-Formyl-4,5-dimethoxyphenoxy)-1,2,3,10-tetramethoxyxoaoporphine, O-179
 Formyldimethylamine, D-747
 3-Formyl-1,7-dimethylindole, M-475
N-Formyliduguevanine, D-950
*N*⁸-Formylechitamine, E-30
*N*²⁰-Formylepipachysamine F, P-601
N-Formylflavaconitine, M-332
 3-Formyl-1-geranyl-2,8-dihydroxycarbazole, M-729
 3-Formyl-1-geranyl-2-hydroxycarbazole, M-729
 6-Formyl-1-geranyl-7-hydroxy-2-methoxycarbazole, E-276
N-Formylharappamine, M-667
 2-Formyl-5-(6-heneicosenyl)-1*H*-pyrrole, H-109
N-Formylhernangerine, N-25
N-Formyl-4-hydroxybenzeneacetamide, H-672
 6-Formyl-1-hydroxycarbazole, H-451
 6-Formyl-2-hydroxycarbazole, H-450
 3-Formyl-1-hydroxycarbazole, H-447
 3-Formyl-2-hydroxycarbazole, H-448
 3-Formyl-6-hydroxycarbazole, H-449
 1-Formyl-4-hydroxy- β -carboline, H-456
N-Formyl-20-hydroxycylindrocarine, C-929
 8-Formyl-7-hydroxydictamine, E-315
 1-Formyl-7-hydroxy-6,7-dihydro-5*H*-pyrrolizine, D-14
 6-Formyl-7-hydroxy-1,3-dimethoxycarbazole, T-533
 3-Formyl-4-hydroxyindole, H-527
 3-Formyl-5-hydroxyindole, H-528
 3-Formyl-6-hydroxyindole, H-529
 6-Formyl-7-hydroxy-1-methoxycarbazole, D-557
 3-Formyl-1-hydroxy-6-methoxycarbazole, D-552
 3-Formyl-2-hydroxy-6-methoxycarbazole, D-555
 3-Formyl-2-hydroxy-7-methoxycarbazole, D-556
 3-Formyl-7-hydroxy-1-methoxy-9*H*-carbazole, D-553
 8-Formyl-7-hydroxy-4-methoxyfuro[2,3-*b*]quinoxaline, E-315
 1-Formyl-7-hydroxy-2-methoxy-6-methylcarbazole, D-619
 1-Formyl-2-hydroxy-5-methoxy-6-methylindoline, D-432
 3-Formyl-2-hydroxy-6-methoxy-1-prenylcarbazole, H-125
 3-Formyl-2-hydroxy-7-methoxy-1-prenylcarbazole, H-125
 1-Formyl-3-hydroxy-6-methylcarbazole, H-589
 2-Formyl-5-hydroxymethyl-1*H*-pyrrole-1-butanoic acid, F-137
 2-Formyl-5-(hydroxymethyl)pyrrole, H-626
 4-(2-Formyl-5-hydroxymethyl-1*H*-pyrrol-1-yl)butanoic acid, F-137
 2-(2-Formyl-5-hydroxymethyl-1*H*-pyrrol-1-yl)-3-phenylpropanoic acid, F-138
 3-(2-Formyl-5-hydroxymethyl-1*H*-pyrrol-1-yl)-2,5-pyrrolidinedione, F-139
 3-Formyl-1-hydroxy-4-prenylcarbazole, C-507
 3-Formyl-2-hydroxy-7-prenylcarbazole, C-500
 6-Formyl-3-hydroxy-4-prenylcarbazole, H-577
 3-Formyl-2-hydroxy-1-prenylcarbazole, H-125
 4-Formyl-2-hydroxyquinoline, H-721
 4-Formyl-8-hydroxyquinoline, H-722
 3-Formyl-2-hydroxy-4,6,7-trimethoxyquinoline, M-175
N-Formyl-3-indoleacrylamide, I-136
 5-Formylindole, I-87

- 7-Formylindole, I-88
 3-Formylindole, I-86
 3-Formyl-1*H*-indole-2-sulfonic acid, F-140
 3-Formylindolyl methyl disulfide, M-434
N-Formyl-3-(1*H*-indol-3-yl)-2-propenamide, I-136
*N*²⁰-Formylrethdiamine C, P-606
N-Formylrethline, C-610
N-Formylisimine, I-186
*N*²-Formylkopsanol, K-65
N-Formyllooline, L-234
 4-Formyl-*N*-mercaptocarbostyryl, H-721
 3-Formyl-2-mercaptindole, M-201
 4-Formyl-1-mercapto-2(1*H*)-quinolinone, H-721
N-Formylmescaline, M-219
N-Formyl-12-methoxyaspidobalbidine, A-1490
N-Formyl-17-methoxyaspidobalbidine, A-1490
N-Formyl-12-methoxyaspidofractinine, A-1494
 6-Formyl-4-methoxy-2,2'-bipyridine, C-11
 5-Formyl-4-methoxy-2,2'-bipyrrole, M-244
 6-Formyl-1-methoxycarbazole, H-451
 6-Formyl-2-methoxycarbazole, H-450
 3-Formyl-1-methoxycarbazole, H-447
 3-Formyl-2-methoxycarbazole, H-448
 3-Formyl-6-methoxycarbazole, H-449
 3-Formyl-8-methoxycarbazole, H-451
 1-Formyl-4-methoxy-β-carboline, H-456
*N*⁸-Formyl-12-methoxyechitamidine, E-30
 1-Formyl-3-methoxy-6-methylcarbazole, H-589
 1-Formyl-5-methoxy-6-methylindoline, D-432
 2-Formyl-5-methoxymethyl-1*H*-pyrrole-1-butanonic acid, F-137
 2-Formyl-5-(methoxymethyl)-1*H*-pyrrole, H-626
 3-Formyl-2-methoxy-8-prenylcarbazole, A-1530
 3-Formyl-1-methoxy-4-prenylcarbazole, C-507
 3-Formyl-2-methoxy-1-prenylcarbazole, H-125
 4-Formyl-8-methoxyquinoline, H-722
 (*Z*)-16-Formyl-5α-methoxystrictamine, R-75
 Formylmethylamine, M-463
N-Formyl-*O*-methylanhalonidine, T-237
 9-Formyl-3-methyl-9*H*-carbazole, M-424
 3-Formyl-2-(methylthio)indole, M-434
 3-Formyl-2,3'-methylenebisindole, M-65
 3-(Formylmethyl)indole, I-80
 3-Formyl-7-methylindole, M-475
N-Formyl-2-methylindoline, D-472
N-Formyl-*O*-methylisopiline, T-521
 3-(2-Formyl-2-methylpropyl)-5-(3-methyl-1-oxo-2-butenyl)-1*H*-indole, D-756
 3-Formyl-2-(methylthio)indole, M-201
*N*⁶-Formyl-*N*⁶-methyltryptamine, M-582
N-Formyl-*N*-methyltyramine, H-667
N-Formylnandigerine, N-25
N-Formylnorcycloclobuxine, C-882
N-Formylnorephedrine, A-869
N-Formylnorgalanthamine, G-3
N-Formylnorlooline, L-234
N-Formylnormacromerine, M-18
N-Formylnormantenine, N-29
N-Formylnormicotine, P-957
N-Formylnormuciferine, D-531
N-Formylnorpurpureine, P-208
 2-Formyl-2-northalidasine, T-320
N-Formyllovigerine, O-143
 5-Formyloxyecorynoline, C-692
*N*⁶-Formyloxydeoxy-Wieland-Gumlich aldehyde, C-115
 3-Formyloxy-6-tigloyloxytropene, M-391
N-Formylpapilicine, B-455
N-Formylpiperidine, P-455
 2-(1-Formyl-2-piperidyl)-1-pyrroline, S-330
 3-(*N*-Formyl-2-piperidyl)quinolizidine, L-97
N-Formylpsammaphysin A, P-695
 12-Formylpterioic acid, R-82
 Formylpteroylglutamic acid, F-121
N-Formylpurpureine (incorr.), P-208
N-Formylputerine, P-781
 5-Formyl-1*H*-pyrrole-2-alkanenitriles, F-141
N-Formyl-1*H*-pyrrole-2-carboxamide, P-944
 5-Formyl-1*H*-pyrrole-2-carboxylic acid, F-142
 2-Formylpyrrole, P-942
 24-(5-Formyl-1*H*-pyrrol-2-yl)-2-hydroxy-18-tetra-cosenenitrile, C-827
 (5-Formyl-1*H*-pyrrol-2-yl)methyl 4-hydroxybutanoate, H-626
 13-(5-Formyl-1*H*-pyrrol-2-yl)-7-tridecenenitrile, C-825
N-Formylquestiomycin B, A-861
 4-Formylquinoline, Q-30
*N*⁶-Formylroquefortine C, R-125
N-Formylstaurosporine, S-498
N-Formylstapharine, P-643
 16-Formylstrictamine, R-75
N-Formyl-4,5,6*a*,7-tetrahydro-7-hydroxyglau-cine, D-159
 3-Formyl-4,5,6,7-tetrahydro-4,5-dihydroxy-1*H*-indole, T-156
 5-Formyl-5,6,7,8-tetrahydrofolic acid, F-121
 1-Formyl-1,2,3,4-tetrahydro-3-(2-hydroxymethyl-4-methoxyphenyl)-6-methoxyquinazoline, D-369
 9-Formyl-1,2,3,4-tetrahydro-7-methoxy-β-carboline, P-916
N-Formyltortuosamine, T-433
 6-Formyl-1,3,7-trihydroxycarbazole, T-533
 3-Formyl-1,6,7-trihydroxycarbazole, T-532
 5-Formyl-4,6,8-trihydroxy-1-methoxycarbonylphena-zine, L-241
O-Formyltropine tropate, T-632
 11-Formyl-2,4,8,10-undecatetraenoic acid, O-176
N-Formylxylofine, A-1066
 Forrestine, P-247
 Forticine, C-290
 Fortucine, P-528
 Fortuneine, F-143
 Fortunine, S-125
 Foveoglin A, F-144
 Foveoglin B, F-144
 Foxy, M-288
 FR 225656, A-1172
 FR 225659, A-1173
 FR 252921, A-1174
 FR 252922, A-1175
 FR 256523, A-1175
 FR 3383, A-1170
 FR 66979, A-1171
 FR 900148, A-1176
 FR 900220, A-680
 FR 900482, A-1177
 FR 901483, A-1178
 FR 900216B, R-83
 FR 900447, C-427
 FR 900216C, R-83
 FR 900483, D-420
 FR 900409, E-243
 FR 900411, E-242
 FR 900216F, R-83
 FR 177391, H-76
 FR 900494, K-41
 FR 005759, P-961
 Fragilamide, F-145
 Fragin, F-146
 Francheline, F-147
 Franchetine, F-147
 Franchetine, F-148
 Franganine, F-149
 Frangulofoline, F-150
 Frangulanine, F-151
 Frankiamide, C-26
 Fredericamycin A, F-152
 Fredericamycin B, F-153
 Fredericamycin E, F-154
 Fritillarine, C-298
 Fritillarizine, C-301
 Fritilline A, F-155
 Fritilline B, F-156
 Fritimine, F-157
 Fromiamycalin, F-158
 FRT-A, F-162
 FRT-B, F-81
 Fructigenine A, F-159
 Fructigenine B, F-160
 d-Fructosazine, B-221
 2-(Fructosylamino)benzoic acid, F-161
 Fruit rot toxin A, F-162
 Fruit rot toxin B, F-81
 Fruticosamine, F-163
 Fruticosine, F-163
 Fruticosoline, F-164
 Fruticosonine, F-164
 FTB, N-315
 FTF, T-646
 FTH, T-646
 FTJ, T-646
 Fuchsiaefoline, M-237
 Fuchsisenecionine, H-233
 5-FU, F-104
 Fugapavine, M-141
 Fugu poison, T-298
 Fukinotoxin, P-277
 Fulgorubin A, F-165
 Fulvanine A, F-166
 Fulvanine B, F-166
 Fulvanine C, F-166
 Fulvanine D, F-166
 Fulvanine E, F-167
 Fulvine *N*-oxide, F-168
 Fulvine, F-168
 Fumadensine, F-169
 Fumafloquine, F-170
 Fumaramidine, F-171
 Fumaramidmycin, P-337
 Fumaramine, F-172
 Fumaric acid isobutylamide, F-173
 Fumaric acid, F-173
 Fumaricine, F-174
 Fumaridine, N-36
 Fumariflorine ethyl ester, F-175
 Fumariflorine, F-175
 Fumarilicine, F-174
 Fumariline, F-176
 Fumarine, P-667
 Fumarine, F-177
 Fumaritine *N*-methosalt, F-174
 Fumaritine *N*-oxide, F-174
 Fumaritine, F-174
 Fumaritridine, F-178
 Fumaritrine, F-178
 Fumarizine, F-179
 Fumarofine, F-180
 Fumarophycine, F-174
 Fumarostelline, F-180
 Fumigaclavine A, F-181
 Fumigaclavine B, F-181
 Fumigaclavine C, F-182
 Fumimycin, F-183
 Fumiquinazoline A, F-185
 Fumiquinazoline B, F-185
 Fumiquinazoline C, F-186
 Fumiquinazoline D, F-187
 Fumiquinazoline E, F-185
 Fumiquinazoline F, K-188
 Fumiquinazoline G, F-188
 Fumiquinazoline H, F-189
 Fumiquinazoline I, F-184
 Fumitremorgin C, F-190
 Fumitremorgin D, N-315
 Fumitremorgin E, T-646
 Fumitremorgin F, T-646
 Fumitremorgin G, T-647
 Fumitremorgin H, T-646
 Fumitremorgin L, T-647
 Fumitremorgin M, N-315
 Fumitremorgin N, N-315
 Fumjudaine, S-612
 Fumschleicherine, F-191
 Fumvaviline, F-192
 Funebradiol, F-193
 Funebral, F-194
 Funebrine, F-195
 Fungerin, F-196
 Fungicide PF 1140, A-1226
 Funiculosin, F-197
 Funiferine *N,N'*-dimethosalt, R-114
 Funiferine *N*-oxide, R-114
 Funiferine, R-114
 Funtessine, H-318
 Funtudiamine A, P-601
 Funtudiamine B, P-601
 Funtudienine, F-198

- Funtuline, L-56
 Funtumafrine B, A-883
 Funtumafrine C, A-883
 Funtumidine, A-880
 Funtumine, A-882
 Funtuphyllamine A, A-881
 Funtuphyllamine B, A-881
 Funtuphyllamine C, A-881
 Furacridone, F-206
 3-Furancarbinol, F-199
 3-Furanmethanol, F-199
 6-(3-Furanyl)- β ,3-dimethyl-2-piperidinebutanol, N-340
 6-(3-Furanyl)-1,6,7,8,9,9a-hexahydro-9-(hydroxymethyl)-2H-quinolizine-3-carboxaldehyde, N-339
 5-(3-Furanyl)-8-methylindolizidine, F-202
 4-[6-(3-Furanyl)-3-methyl-2-piperidinyl]-2-methyl-2-buten-1-ol, N-340
 1-(2-Furanylmethyl)-1H-pyrrole, F-200
 2-[(2-Furanylmethyl)thio]-6-methylpyrazine, F-201
 4-(3-Furanyl)octahydro-1,7-dimethyl-2H-quinolizine, D-227
 5-(3-Furanyl)octahydro-8-methylindolizine, F-202
 1-[5-(2-Furanyl)pyrazinyl]-1,2,4-butanetriol, C-776
 1-(2-Furanyl)-9H-pyrido[3,4-*b*]indole-3-carboxylic acid, F-85
 2-Furanyl-5-(1,2,4-trihydroxybutyl)pyrazine, C-776
 6-(3-Furanyl)- $\alpha,\alpha,3$ -trimethyl-3-piperidinepropa-
 nol, N-340
 3-Furfuryl alcohol, F-199
 3-Furfuryl 2-pyrrolicarboxylate, F-199
N-(2-Furfuryl)pyrrole, F-200
 Furoclausine A, F-203
 Furoclausine B, F-204
 Furoclausine C, F-205
 Furofoline I, F-206
 Furofoline II, F-207
 Furofoline, F-206
 Furomegistine II, F-208
 Furomegistine I, R-88
 Furoparadine, F-209
 Furo[2,3-*b*]quinoline-4,5-diol, F-210
 Furo[2,3-*b*]quinoline-4,6-diol, F-211
 Furo[2,3-*b*]quinoline-4,7-diol, F-212
 Furo[2,3-*b*]quinoline-4,8-diol, F-213
 Furo[2,3-*b*]quinoline-4,5,6,7,8-pentol, F-214
 Furo[2,3-*b*]quinoline-4,5,7,8-tetrol, F-216
 Furo[2,3-*b*]quinoline-4,5,6,7-tetrol, F-215
 Furo[2,3-*b*]quinoline-4,6,7,8-tetrol, F-217
 Furo[2,3-*b*]quinoline-4,5,6-triol, F-218
 Furo[2,3-*b*]quinoline-4,5,7-triol, F-219
 Furo[2,3-*b*]quinoline-4,5,8-triol, F-220
 Furo[2,3-*b*]quinoline-4,6,7-triol, F-221
 Furo[2,3-*b*]quinoline-4,6,8-triol, F-222
 Furo[2,3-*b*]quinoline-4,7,8-triol, F-223
 Furo[2,3-*b*]quinolin-4-ol, F-224
 Furo[2,3-*b*]quinolin-4(9*H*)-one, F-224
 Furostifoline, M-464
 18-*O*-(3-Furoyl)willfortrine, P-253
 Fusaric acid, B-445
 Fusaricide, F-225
 Fusarin A, F-226
 Fusarin C, F-227
 Fusarin D, F-226
 Fusarin F, F-228
 Fusarin X, F-227
 Fusarin, B-444
 Fusarinin, B-444
 Fusarinine C, F-231
 Fusarinolic acid, H-436
 Fuscaine, D-433
 Fuscin, A-1576
 Fuscoatramide, A-715
 Fuscine, F-229
 Fuseine, F-230
 Fusigen, F-231
 Futoamide, F-232
 Fuziline, M-313
 Fuzitine, T-248
 Fysostigmine, P-400
 G 1499-2, A-1179
 G 1549, H-563
 GABA, A-715
 Gabonine, G-1
 Gabunamine, C-615
 Gabunine, C-616
 Gadeline, G-152
 Gadenine, M-307
 Gadesine, G-152
 Gagamine, P-610
 Gagaminine, P-610
 4-*O*- α -D-Galactopyranosylcalystegine B₂, A-1587
 2-*O*- α -D-Galactopyranosyl-1-deoxymannojirimycin, H-616
 6-*O*- α -D-Galactopyranosyl-1-deoxynojirimycin, H-616
 2-*O*- α -D-Galactopyranosyl-1-deoxynojirimycin, H-616
 5-*O*- α -D-Galactopyranosyl- α -homonojirimycin, B-186
 Galactostatin, A-726
 Galactosylqueuosine, Q-9
 Galantamine hydrobromide, G-3
 Galantamine, G-3
 Galanthamide, G-2
 Galanthamine *N*-oxide, G-3
 Galanthamine, G-3
 Galanthamine, G-3
 Galanthidine, L-348
 Galanthindole, G-4
 Galanthine, P-716
 Galanthusine, G-5
 Galantin I, G-6
 Galantin II, G-7
 Galasine, G-8
 Galbazine, M-273
 Galegine, M-410
 Galipeine, C-821
 Galipine, G-11
 Galipinine, G-9
 Galipoidine, G-10
 Galipoline, G-11
 6'-*O*-Galloylprunasin, H-673
O-Galloyltyramine, H-667
 6'-*O*-Galoylsambunigrin, H-673
 Galwesine, H-504
 Gamalon, A-715
 Gambirdine, M-661
 Gambireine, G-13
 Gambirine, G-13
 Gambirtannine, D-434
 Gancidin W, C-876
 Gandharamine, G-14
 Ganoderpurine, G-15
 Ganodine, H-626
 Ganoine, H-626
 Garceine, G-16
 Gardenamide A, G-17
 Gardfloramine, G-18
 Gardmultine, G-19
 Gardneramine *N*⁴-oxide, G-20
 Gardneramine oxindole, G-21
 16*R*-Gardneramine oxindole, G-21
 Gardneramine, G-20
 Gardnerine, G-22
 Gardnutine, G-23
 Gardquinolone, G-24
 Garryfoline, V-45
 Garryine, G-25
 Gasabiimine, G-26
 Gastrodamine, B-181
 Gastrodigenin, H-414
 Gaultherialine A, G-27
 Gazer, Z-11
 GB 13, H-281
 GB 1, H-277
 GB4 Toxin, G-28
 GC 1, H-731
 GC 2, H-731
 GC 3, S-109
 GD 0039, S-650
 GDIMBOA, H-409
 GE 2270, A-1180
 GE 2270A, A-1180
 GE 37468A, A-1181
 GE 2270B₁, A-1180
 GE 2270B₂, A-1180
 GE 37468B, A-1181
 GE 2270C₁, A-1180
 GE 37468C, A-1181
 GE 2270C_{2a}, A-1180
 GE 2270C_{2b}, A-1180
 GE 2270D₁, A-1180
 GE 2270D₂, A-1180
 GE 2270E, A-1180
 GE 2270T, A-1180
 Geamine, S-565
 Gebeline, G-137
 Geibalansine, G-29
 Gejedimerine, G-30
 Geissolosimine, G-31
 Geissoschizal, G-33
 Geissoschizine methyl ether, G-32
 Geissoschizine, G-32
 Geissoschizol, G-33
 Geissoschizoline *N*⁴-oxide, G-34
 Geissoschizoline, G-34
 Geissospermine, G-35
 Geissovelline, G-36
 Gelliusine A, G-37
 Gelliusine B, G-37
 Gelliusine C, G-38
 Gelliusine D, G-39
 Gelliusine E, G-40
 Gelliusine F, G-40
 Gelsamydine, G-41
 Gelsebanine, G-42
 Gelsedilam, G-43
 Gelsedine, G-44
 Gelsefuranidine, G-45
 Gelseiridone, G-46
 Gelselegine, G-47
 Gelsemamide, G-48
 Gelsemicine, G-44
 Gelsemine *N*-oxide, G-49
 Gelsemine, G-49
 Gelsemoxonine, G-50
 Gelsempervine A, V-171
 Gelsempervine B, V-171
 Gelsempervine C, V-171
 Gelsempervine D, V-171
 Gelsenicine, G-44
 Gelsevirine, G-49
 Genatropine, D-632
 Geneserine, G-51
 Geniconitine, M-323
 Geniculatine A, M-323
 Geniculatine B, M-323
 Geniculatine C, M-352
 Geniculatine D, M-352
 Genicunine A, M-355
 Genicunine B, M-349
 Genicunine C, M-326
 Geninthiocin, B-104
 Genipamide, G-52
 Genistein, I-321
 Genocodine, C-555
 Genomorphine, M-704
 Genostrychnine, S-589
 Gentiacraline, G-53
 Gentialutine, V-52
 Gentianadine, G-54
 Gentianaine, G-62
 Gentianal, G-55
 Gentianamine, G-56
 Gentiananine, G-57
 Gentianidine, G-58
 Gentianine lactam, G-60
 Gentianine, G-59
 Gentiatibetine, G-61
 Gentiocrucine, G-62
 Gentioflavine, G-63
 Gentiobetine, G-61
 Gentrymine A, T-182
 Gentrymine B, T-182
 Geodin A, C-924
 Geoffroyin, T-700
 Geometricin A, S-652

- Geotrichum alkaloid A 25822B, A-1602
 Geotrichum alkaloid A 25822D, A-1602
 Geotrichum alkaloid A 25822H, G-64
 Geotrichum alkaloid A 25822L, A-1602
 Geotrichum alkaloid A 25822M, A-1602
 Geotrichum alkaloid A 25822N, A-1602
 Geovanine, K-6
 Gephyrototoxin 223AB, O-41
 Gephyrototoxin 239AB, O-41
 Gephyrototoxin 167B, O-56
 Gephyrototoxin 239CD, O-41
 Gephyrototoxin, G-65
 Geralbine, G-66
 Geraldoamine, D-88
 Gerambullal, G-67
 Gerambullindiol, G-68
 Gerambullin, G-68
 Gerambullol, G-68
 1-Geranyl-2,7-dihydroxy-3-methylcarbazole, E-277
 1-Geranyl-2,7-dihydroxy-6-methylcarbazole, E-276
 O-Geranylhalfordinol, H-12
 4-Geranyl-3-hydroxy-6-methylcarbazole, E-302
 8-Geranyl-7-hydroxy-3-methylcarbazole-1,4-quinone, M-767
 7-Geranyllindolactam V, I-78
 3-Geranyllindole, D-759
 Geranyllinaloyl formamide, I-221
 Geranyllinaloyl isocyanide, I-221
 Geranyllinaloyl isothiocyanate, I-221
 4-Geranyl-1H-pyrrole-2-carboxylic acid, D-761
 2-Geranyl-1,3,5-trihydroxy-1-methylacridone, G-121
 GERI-BP001 A, T-500
 GERI-BP001 B, T-500
 GERI-BP001 M, T-500
 1(10),4-Germacradien-11-amine, G-69
 Germanitrine, G-70
 Germbudine, G-70
 Germerine, G-70
 Germidine, G-70
 Germinaline, G-70
 Germinalinine, G-70
 Germinine, G-70
 Germinine, G-70
 Germinitrine, G-70
 Germitetrine B, G-70
 Germitetrine, G-70
 Germitrine, G-70
 Gerontine, S-399
 Gerrardamine, G-71
 Gerrardine, G-72
 Gerrardoline, G-73
 Gesashidine A, H-781
 Gesidine, V-110
 Geyeridine, T-552
 Geyerine, T-552
 Geyerinine, H-176
 Geyerline, G-91
 GGBA, G-201
 Gigactonine, M-309
 Gigantamide A, G-74
 Gigantine, T-174
 Gigartinine, G-75
 Gilbertine, G-76
 Gilgite, P-800
 Gillette, C-552
 Gilurytmal, A-216
 Gilvusmycin, R-4
 Gindarine, G-77
 Gindarine, T-212
 Gindarinine, P-34
 Ginkgotoxin, P-923
 Ginsenine, G-78
 Giracodazole, G-79
 Giraldine A, M-371
 Giraldine B, M-371
 Giraldine C, M-371
 Giraldine D, M-371
 Giraldine E, M-363
 Giraldine F, M-371
 Giraldine G, D-196
 Giraldine H, D-196
 Giraldine I, M-356
 Girgensohnine, G-80
 Girgensonine, G-80
 Girinimbilol, M-728
 Girinimbine, G-81
 Girolline, G-79
 Gitingensine, P-93
 GKK 1032A₁, A-1182
 GKK 1032A₂, A-1182
 GKK 1032B, A-1183
 Glabratine, G-82
 Glabredelphinine, M-371
 Glabrin, D-661
 Glaciapyrrole A, G-83
 Glaciapyrrole B, G-84
 Glaciapyrrole C, G-85
 Glanduline, H-170
 Glandulosine, H-178
 Glandulosine, M-309
 Glansreginin A, G-86
 Glansresinoside A, G-86
 Glaucamine, G-87
 Glaucedine, M-309
 Glaucenamide, G-88
 Glaucenine, M-307
 Glaucentrine, C-682
 Glaucephine, M-307
 Glaucerine, M-307
 Glaucescine, G-89
 Glaucescinine, G-90
 Glaucic acid, F-173
 Glaucine methyl methine, M-559
 Glaucine, T-246
 Glaucon, A-152
 Glaudelsine, G-91
 Glau dine, G-87
 Glau fidine, C-682
 Glau fine, T-248
 Glau finine, T-248
 Glau firidine, C-682
 Glau flavine, G-92
 Glau nidine, G-93
 Glau nine, G-93
 Glau pavine, G-87
 Glau vent, T-246
 Glau vine, O-178
 Glazen, P-643
 Glaziovine, P-643
 Glenodinine, G-95
 Glicomarine, S-274
 Glidobactamine, G-96
 Glidobactin A, G-96
 Glidobactin B, G-96
 Glidobactin C, G-96
 Glidobactin D, G-96
 Glidobactin E, G-96
 Glidobactin F, G-96
 Glidobactin G, G-96
 Glidobactin H, G-96
 Glidobactin PF1, G-96
 Gliocladine A, G-97
 Gliocladine A, V-89
 Gliocladine B, G-97
 Gliocladine B, V-89
 Gliocladine C, G-98
 Gliocladine C, G-99
 Gliocladine D, G-99
 Gliocladine E, G-99
 Glioclatine, G-99
 Gln, G-108
 Globiferine, T-488
 Glochidacuminoside D, H-520
 Glochidiatusamide, C-456
 Glochidiatusin, C-456
 Glochidicine, G-100
 Glochidine, G-101
 Glomerine, H-632
 Glomerulatine A, C-42
 Glomerulatine B, C-42
 Glomerulatine C, C-42
 Gloriosamine A, C-654
 Gloriosamine B, C-654
 Gloriosamine C, C-568
 Gloriosamine D, C-568
 Gloriosine, D-92
 Glossobalol, D-327
 Glucobrassicin, I-122
 Glucodichotomine B, D-579
 Glucoerysodine, E-203
 15-Glucoerysopine, E-203
 16-Glucoerysopine, E-203
 Glucohaplopine, F-223
 Glucoindican, I-104
 Glucoisatisin, G-102
 Glucopiericidin A, P-421
 Glucopiericidin B, P-421
 Glucopiericidinol A₁, P-422
 Glucopiericidinol A₂, P-422
 Glucoprunasin, H-673
 α-D-Glucopyranose 1,3,6-tri-3-pyridinecarboxylate, A-458
 β-D-Glucopyranose 1,2,6-tri-1H-pyrrole-2-carboxylate, B-415
 β-D-Glucopyranose-6-(4-hydroxybenzoate) 1,2-di-1H-pyrrole-2-carboxylate, B-416
 β-D-Glucopyranosyl anthranilate, A-706
 6'-O-α-D-Glucopyranosylalanside, A-239
 6'-O-β-D-Glucopyranosylalanside, A-239
 3'-O-β-D-Glucopyranosylalanside, A-239
 11-O-β-D-Glucopyranosylartavenustine, X-30
 20-O-β-D-Glucopyranosylcamptothecin, C-70
 6'-O-α-D-Glucopyranosyl-3-O-demethyl-2-O-methylalanside, A-239
 4-O-β-D-Glucopyranosyl-1-deoxymannonojirimycin, H-616
 2-O-β-D-Glucopyranosyl-1-deoxymannonojirimycin, H-616
 3-O-β-D-Glucopyranosyl-1-deoxymannonojirimycin, H-616
 4-O-α-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 4-O-β-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 6-O-β-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 2-O-α-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 2-O-β-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 3-O-α-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 3-O-β-D-Glucopyranosyl-1-deoxynojirimycin, H-616
 N-(5-β-D-Glucopyranosyl-2,3-dihydroxybenzoyl)-serine, S-33
 4-O-β-D-Glucopyranosylfagomine, H-614
 3-O-β-D-Glucopyranosylfagomine, H-614
 7-O-β-D-Glucopyranosyl-α-homonojirimycin, B-186
 20-O-β-D-Glucopyranosyl-18-hydroxycamptothecin, C-70
 N-[2-(1-β-D-Glucopyranosyl-1H-imidazol-4-yl)ethyl]-N-methyl-3-phenyl-2-propenamide, C-176
 3'-O-α-D-Glucopyranosylinosine, I-148
 6''-O-α-D-Glucopyranosylipicoside, I-167
 6''-O-β-D-Glucopyranosylipicoside, I-167
 23-O-β-D-Glucopyranosyl-20-isoveratramine, V-71
 2-β-D-Glucopyranosyl-3-isoxazolin-5-one, I-349
 4-O-α-D-Glucopyranosylmoranoline, H-616
 1-(3-Glucopyranosyloxybutanoyl)pancratistatine, P-58
 9-β-D-Glucopyranosylcamptothecin, C-70
 11-O-β-D-Glucopyranosylxanthin-6-one, H-446
 (4'-β-D-Glucopyranosyloxycinnamoyl)lupinine, O-58
 N-[[3-(β-D-Glucopyranosyloxy)-2,3-dihydro-2-oxo-1H-indol-3-yl]acetyl]aspartic acid, G-103
 [6-(β-D-Glucopyranosyloxy)-2-hydroxy-3,4-dimethoxycyclohexylidene]acetoneitrile, S-302
 4-(β-D-Glucopyranosyloxy)-2-(hydroxymethyl)-2-butenenitrile, H-509
 4-(β-D-Glucopyranosyloxy)-3-(hydroxymethyl)-2-butenenitrile, H-510
 (4'-O-β-D-Glucopyranosyloxy-3'-methoxycinnamoyl)lupinine, O-58

- 2-(β -D-Glucopyranosyloxy)-3-methyl-2-butenetri-
trile, A-20
2-(β -D-Glucopyranosyloxy)-2-methylpropaneni-
trile, H-617
8-Glucopyranosyloxystylopin, S-612
9- β -D-Glucopyranosyloxystylopin, M-658
3-(β -D-Glucopyranosyloxy)-1,2,3,6-tetrahydro-2-
hydroxy-4-methoxy-1-methyl-6-oxo-3-pyridine-
carbonitrile, A-22
N- β -D-Glucopyranosylquestiomycin A, A-862
N- β -D-Glucopyranosylvinicosamide, V-126
Glucose 3-nitropropanoates, G-104
Glucosylngouniense, N-186
 β -D-Glucosyloxazoxymethane, M-397
8-Glucosyloxymethane, D-621
10-Glucosyloxymethane lactam, V-126
Glucoveracintine, V-57
Glu, G-107
Glumin, G-108
Glusun I, G-105
Glusun II, G-106
Glutamic acid 5-amide, G-108
Glutamic acid betaxanthin, V-195
Glutamic acid, G-107
Glutamicine, F-81
Glutamidedetaxanthin, V-195
Glutamine, G-108
 γ -Glutamine, G-108
Glutaminic acid, G-107
2-(β -Glutaminylaminoethyl)-3-isoxazolin-5-one,
I-349
 γ -Glutaminyl-4-hydroxybenzene, G-108
Glutamycin, F-81
N-(γ -Glutamyl)ethanolamine, G-108
 γ -Glutamyl- β -(isoxazolin-5-on-2-yl)alanine, A-850
 γ -L-Glutamyl-L-pipecolic acid, P-456
5-(Glutarimidomethyl)quinolizidine, L-26
Glutaven, G-108
Glutimic acid lactam, O-206
Glutimic acid, O-206
Glutimic acid, O-206
Glyantrypine, G-109
Glybomine A, D-616
Glybomine B, G-110
Glybomine C, G-110
Glycarpine, F-219
Glycerinopyrin, G-111
2-*O*-Glycerophosphoryllycorine, L-348
N-Glyceryltaurine, A-765
Glycinebetaxanthin, P-577
Glycinyllimaquinone, S-329
Glycoamide A, D-900
Glycoamide B, D-900
Glycobismine A, G-112
Glycobismine B, G-113
Glycobismine C, G-113
Glycobismine D, G-114
Glycobismine E, G-115
Glycobismine F, G-115
Glycobismine G, G-116
Glycoborine, H-588
Glycoborinine, E-300
Glycocinnaspermicidin A, G-117
Glycocinnaspermicidin B, C-459
Glycocinnaspermicidin C, C-460
Glycocinnaspermicidin D, G-117
Glycocitlone A, G-118
Glycocitlone B, G-118
Glycocitlone C, G-118
Glycocitridine, G-119
Glycocitrine III, G-121
Glycocitrine I, T-581
Glycocitrine IV, T-270
Glycocitrine V, G-120
Glycocitrine VI, G-122
Glycocytamine, G-200
Glycofoline, G-123
Glycofolinine, P-207
Glycohaplopin, F-223
Glycolylurea, I-40
Glycolone, G-124
Glycolone, T-583
Glycomarine, S-274
Glycomaurin, E-300
Glycomaurrol, G-125
Glycoperine, F-223
Glycophylone, T-583
Glycophymine, B-93
Glycophymoline, B-93
Glycorine, M-556
Glycosine, B-93
Glycosinine, H-448
Glycosmicine, Q-13
Glycosminine, B-93
Glycosolone, T-583
Glycothiohexide α , G-126
Glycothiohexide β , G-126
Glycothiomine A, M-577
Glycothiomine B, M-577
Glycothalic acid, U-57
Glycozolicine, H-584
Glycozolidal, D-555
Glycozolidine, D-617
Glycozolidol, D-617
Glycozoline A, G-127
Glycozoline B, G-127
Glycozoline, H-587
Glycozolinine, H-587
Glycozolinol, H-587
Glycozalone A, G-127
Glycozalone B, G-127
O-Glycylgriseoliteic acid, H-514
Glyfoline, H-242
Glymontanine A, G-128
Glymontanine B, G-129
Glyoxaline, I-35
Glyoxalone-4-carboxylic acid, D-484
Glyoxylic diureide, A-625
Glyparvin A, T-478
Glypetelotone, G-130
Glysperrin A, G-131
Glysperrin B, G-132
Glysperrin C, G-133
Glyyunnanenine, H-764
GM 95, T-66
Gnidine, G-134
Gnidinine, G-135
Gnidiodine, L-328
 α -Gnoscopine, N-44
Gnoscopine, N-44
 β -Gnoscopine, N-44
Goadsporin, G-136
GOBAB, A-792
Goebelene, G-137
Golceptine, P-716
Goleptine, P-716
Golmaeone, G-138
Gomaline, S-576
Gomandonine, G-139
Gomezine, A-1365
Gomphrenin I, B-110
Gomphrenin II, B-110
Gomphrenin III, B-110
Gomphrenin V, B-110
Gongrine, G-140
Goniofithine, T-539
Goniomine, G-141
Goniomitine, G-142
Goniopedaline, T-537
Goniopetaline (incorr.), T-537
Goniothalactam, T-540
Gonyautoxin I, H-731
Gonyautoxin II, H-731
Gonyautoxin III, H-731
Gonyautoxin IV, H-731
Gonyautoxin VIII, H-731
Gonyautoxin VI, S-109
Gonyautoxin V, S-109
Gorchacoine, G-143
Gortschakoine, G-143
Goshuyamide II, G-144
Goshuyamide I, T-146
Gossyribilone, G-145
Gostatin, G-146
Gosuiamide II, G-144
Gosuiamide I, T-146
Goudotianine, G-147
Gouregine, G-148
Govadine, X-30
Govanine, X-30
Goziline, V-188
Gracicleistanthoside, G-149
Gracilamine, G-150
Graciline, G-151
Graciline, G-152
Gracinine, M-308
Grahamine, G-153
Grahamine, M-688
Gramine methosalt, D-736
Gramine *N*-oxide, D-736
Gramine, D-736
Graminifoline, G-154
Gramminifoline, G-154
Gramodendrine, G-155
Granatan-3-one, A-1580
Granatonine, A-1580
Grandiamide A, G-156
Grandiamide B, C-456
Grandiamide C, C-456
Grandiamide D, C-456
Grandifloricine, G-91
Grandiflorine, G-91
Grandiflorine, R-94
Grandifloritine, G-157
Grandifoline, A-676
Grandifoline, G-158
Grandirubrine, G-159
Grandisine A, G-160
Grandisine B, G-161
Grandisine C, G-162
Grandisine D, G-163
Grandisine E, G-164
Grandisine F, G-165
Grandisine G, G-166
Grandisine III, T-243
Grandisine II, T-243
Grandisine I, T-243
Grandisinine, P-616
Granegillin, A-1475
Granjine, C-643
Grantaline, G-167
Grantianine, G-168
Granulatamide A, T-639
Granulatamide B, T-639
Granulatimide, G-169
Grassypeptolide, R-170
Gravacridonechlorine, R-167
Gravacridonediol acetate, R-167
Gravacridonediol glucoside, R-167
Gravacridonediol methyl ether, R-167
Gravacridonediol, R-167
Gravacridonetriol glucoside, R-167
Gravacridonetriol, R-167
Gravacridonol monoglucoside, R-167
Gravacridonolchlorine, R-167
Gravacridonol, R-167
Graveoline, G-171
Graveolinine, M-264
Gravidox, P-923
Grayanin, H-673
3-Greenwayodendrinol, G-172
3-Greenwayodendrinone, G-172
Grenadadiene, H-131
Grenadamide, H-131
Griffiazanone A, G-174
Griffiazanone B, M-97
Griffinine, G-173
Griffithazanone A, G-174
Griffithdione, G-175
Griffithinam, T-539
Griffithine, G-176
Griffonin, L-213
Grisabine, G-177
Grisabutine, G-177
Griseoluteic acid, H-514
Griseolutein A, H-514
Griseolutein B, H-514
Griseoviridin, G-178

- Grisonomycin, G-179
 Grixazone A, G-180
 Grixazone B, G-180
 Groenlandicine, G-181
 Grossamide K, E-128
 Grossamide, G-182
 Grossularine 1, G-183
 Grossularine 2, G-184
 GS 1, G-44
 GS 2, G-44
 GS 3, G-50
 GT 1012, A-216
 GT 32A, A-1127
 GT 32B, A-1127
 GTX1, H-731
 GTX2, H-731
 GTX3, H-731
 GTX4, H-731
 GTX8, H-731
 GTX5, S-109
 GTX6, S-109
 Guacolidine, G-214
 Guacoline, G-214
 Guadinomic acid, G-185
 Guadinomine A, G-186
 Guadinomine B, G-186
 Guadinomine C₁, G-187
 Guadinomine C₂, G-187
 Guadiscidine, G-188
 Guadiscine, G-188
 Guadiscoline, G-188
 C-Guaianine, G-189
 Guaipyrindine, G-190
 Guamamide, G-191
 Guanacastepene D, G-192
 Guanacastepene H, G-193
 Guanazole, A-735
 Guan-fu base A, H-189
 Guan-fu base B, H-189
 Guan-fu base C, G-194
 Guan-fu base D, G-195
 Guan-fu base E, G-196
 Guan-fu base F *N*-oxide, H-189
 Guan-fu base F, H-189
 Guan-fu base G, H-189
 Guan-fu base H, A-1534
 Guan-fu base I, H-189
 Guan-fu base K, G-197
 Guan-fu base O, H-189
 Guan-fu base P, H-189
 Guan-fu base R, H-189
 Guan-fu base S, G-198
 Guan-fu base Y, H-189
 Guan-fu base Z *N*-oxide, H-189
 Guan-fu base Z, H-189
 Guanidineacetic acid, G-200
 Guanidine, G-199
 Guanidinepropionic acid, G-208
 Guanidinoacetic acid, G-200
 4-Guanidinobutanoic acid, G-201
 4-Guanidino-1-butanol, H-432
 4-Guanidinobutyl syringate, H-432
 4-Guanidinobutylamine, A-719
N-(4-Guanidinobutyl)-2-(4-hydroxyphenyl)-2-oxoacetamide, G-202
 1-(4-Guanidinobutyl)-6-hydroxy- β -carboline, T-225
 4-Guanidinobutyramide, G-201
 2-Guanidinoethanesulfonic acid, G-203
 2-Guanidinoethanesulfonic acid, G-204
*O*³-(2-Guanidinoethoxyphosphinocoserine), L-240
 1-(2-Guanidinoethyl)-1,2,3,4-tetrahydro-6-hydroxymethyl- β -carboline, G-205
 1-(2-Guanidinoethyl)-2,3,4,9-tetrahydro-1*H*-pyrido[3,4-*b*]indole-6-methanol, G-205
 4-Guanidino-3-hydroxybutanoic acid, G-206
 3-Guanidino-6-hydroxymethyl-1-methylpyrazin-2-one, S-553
N-(5-Guanidino-2-hydroxypentanoyl)phenylalanine, M-682
N-(3-Guanidinoisobutyl)-2-methoxyheptylamine, P-314
 5-Guanidino-3-methylaminopentanoic acid, A-779
 5-Guanidino-2-oxopentanoic acid, G-207
 γ -Guanidinoxypropylamine, A-895
 3-Guanidinopropanoic acid, G-208
N-(3-Guanidinopropionyl)-2-hydroxyheptylamine, P-313
 4-[[[3-Guanidinopropyl]amino]butyl]guanidine, H-305
 (3-Guanidinopropylidene)succinic acid, L-373
 1-(3-Guanidinopropyl)-1,2,3,4-tetrahydro- β -carboline, T-637
 Guanidinotaurine, G-204
 3-(3-Guanidinylpropyl)hexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione, C-841
 6-(3-Guanidinylpropyl)-3-isopropyl-2(1*H*)-pyrazinone, A-1421
 3-Guanidinylpyridine, G-209
 Guanine *N*⁷-oxide, G-210
 Guanine, G-210
 Guanopterin, A-818
 Guanosinepropionic acid, H-730
N-Guanylglycine, G-200
 γ -(Guanylureido)butanoic acid, G-140
 Guaranine, C-15
 Guatambuine, G-211
 Guatambuinine, O-83
 Guattaguanine, G-212
 Guattamine, O-28
 Guattaminone, O-28
 Guatteboline, E-102
 Guattegaumerine, D-88
 Guatterine *N*-oxide, G-213
 Guatterine, G-213
 Guattescidine, G-214
 Guattescine, G-214
 Guattouregidine, G-215
 Guattouregine, G-215
 Guayewuanine A, M-316
 Guayewuanine B, M-336
 Guayulamine A, G-216
 Guayulamine B, G-217
 Guenerine, M-367
 Guettardine, G-218
 Guiachrysin, S-593
 Guiaflavine, G-219
 Guianensine, G-220
 Guillauminine, G-221
 Guineamide A, G-222
 Guineamide B, G-223
 Guineensine, G-224
 Guinesine A, M-547
 Guinesine B, M-547
 Guinesine C, M-547
 Gusanlung A, C-73
 Gusanlung B, C-73
 Gusanlung C, G-225
 Gusanlung D, G-226
 Gusperimus trihydrochloride, S-394
 Gusperimus, S-394
 Gutingimycin, G-227
 Guvacine, T-223
 Guvacoline, T-223
 Guyonianin A, H-120
 Gyalanine A, D-182
 Gyalanine B, D-183
 Gymnaconitine, M-352
 Gymnamide, D-746
 Gymnamine, G-228
 Gymnandine, D-221
 Gymnastatin F, G-229
 Gymnastatin G, G-229
 Gymnastatin H, T-700
 Gymnastatin L, G-230
 Gymnastatin N, T-700
 Gymnastatin Q, G-229
 Gymnastatin R, G-229
 Gymnasterone A, G-231
 Gymnodimine B, G-232
 Gymnodimine C, G-232
 Gymnodimine, G-232
 Gynergen, E-172
 Gynesis, C-149
 Gynimbine, Y-13
 Gynuramine, G-233
 Gypsetin, G-234
 Gyroamericine, P-301
 Gyrocarpine, G-235
 Gyrocarpusine, L-175
 Gyrolidine, G-235
 Gyromitrin, G-236
 HA 23, A-1184
 HA 9, O-124
 Habaenine A, H-124
 Habaenine B, M-312
 Habbemine A, H-1
 Habbemine B, H-1
 Habranthine, H-2
 Habropetaline A, D-793
 Hachijodines, H-3
 Hackelidine, H-4
 Hadiensine, P-516
 Hadranthine A, I-31
 Hadranthine B, I-31
 Haemanthamine, H-5
 Haemanthidine, H-6
 Haemanthine, N-169
dibasic-Haematinic acid, E-252
 Haematopodin B, H-8
 Haematopodin, H-7
 Haemodan, A-736
 Haemultine, H-9
 Haiderine, J-17
 Hainanensine, H-10
 Hainaneoside A, P-610
 Hainaneoside B, P-610
 α -Hainanine, C-694
 Haitinosporine, D-570
 Hakurirodine, H-11
 Halaminol A, A-913
 Halaminol B, A-761
 Halaminol C, A-912
 Halaminol D, A-848
 Halfordamine, Q-42
 Halfordine, H-12
 Halfordinine, F-217
 Halfordinol, H-12
 Halfordinone, H-12
 Halichlorine, H-13
 Halichonadin A, H-14
 Halichonadin B, A-776
 Halichonadin C, I-218
 Halichonadin D, A-776
 Halichondramide, H-15
 Halichondramine, A-1404
 Halichondriasulfonic acid, G-204
 Haliclamine A, H-16
 Haliclamine B, H-16
 Haliclamine C, H-17
 Haliclamine D, H-18
 Haliclonacyclamine A, H-19
 Haliclonacyclamine B, H-19
 Haliclonacyclamine C, H-19
 Haliclonacyclamine D, H-19
 Haliclonacyclamine E, A-1404
 Haliclonacyclamine F, A-1404
 Haliclonadamine, P-84
 Haliclonamide A, H-20
 Haliclonamide B, H-20
 Haliclonamide C, H-20
 Haliclonamide D, H-21
 Haliclonamide E, H-21
 Halicloremsin, M-433
 Halicyclamine A, H-22
 Halicyclamine B, H-23
 Haligramide A, H-24
 Haligramide B, H-24
 Halimedine, H-25
 Halipeptin A, H-26
 Halipeptin B, H-26
 Halipeptin C, H-27
 Halipeptin D, H-26
 Halishigamide A, H-15
 Halishigamide B, H-28
 Halishigamide C, H-29
 Halishigamide D, H-30
 Halisulfate 1, H-31
 Halitoxin, H-32

- Halitulin, H-33
 Hallacridone, H-34
 Hallucinex, M-219
 Halocyanine A, H-35
 Halocyanine B, H-36
 Halosaline, H-37
 Halostachine, A-864
 Haloxine, H-38
 Haloxylone A, H-39
 Haloxylone B, H-39
 Haloxymine, H-37
 Halxazone, M-253
 Hamacanthin A, H-40
 Hamacanthin B, H-41
 Hamatine, A-982
 Hamatinine, A-982
 Hamayne, H-5
 Hamigeramide, H-42
 Hamigeramine, H-44
 Hamigeran D, H-43
 Hamigeroxalamic acid, H-44
 Haminol A, P-912
 Haminol B, P-912
 Haminol C, P-910
 Haminol 1, P-911
 Haminol 2, P-911
 Haminol 3, P-911
 Haminol 4, P-911
 Haminol 5, P-913
 Haminol 6, P-913
 Hammarbine, K-28
 Hanadamine, H-45
 Hanamisine, H-195
 Hanamiyama base, H-195
 Hanasanagin, H-46
 Hanishin, L-243
 Hanjison, T-286
 Haouamine A, H-47
 Haouamine B, H-47
 Hapalindole A, H-48
 Hapalindole B, H-48
 Hapalindole C, H-49
 Hapalindole D, H-49
 Hapalindole E, H-49
 Hapalindole F, H-49
 Hapalindole G, H-48
 Hapalindole H, H-48
 Hapalindole I, H-48
 Hapalindole J, H-48
 Hapalindole K, H-48
 Hapalindole L, H-48
 Hapalindole M, H-48
 Hapalindole N, H-48
 Hapalindole O, H-48
 Hapalindole P, H-48
 Hapalindole Q, H-49
 Hapalindole T, H-50
 Hapalindole U, H-48
 Hapalindole V, H-48
 Hapalindolinone A, H-51
 Hapalindolinone B, H-51
 Hapalonamide G, H-52
 Hapalonamide H, H-52
 Hapalonamide V, H-52
 Hapepunine 3- β -cellobioside, S-200
 Hapepunine 3-neohesperidoside, S-200
 Hapepunine, S-200
 Haplafine, H-53
 Haplamide, B-422
 Haplamidine, C-456
 Haplamine, F-90
 Haplahpine, H-53
 Haplatine, F-223
 Haplobine, F-223
 Haplobucharine, H-56
 Haplobungine, Q-43
 Haplocidine, A-1490
 Haplocidiphytine, H-54
 Haplocine, A-1490
 Haplodimerine, H-55
 Haplofidine, F-213
 Haplofoline, H-56
 Haploperine, F-223
 Haplophidine, F-213
 Haplophydine, F-213
 Haplophyllidine, H-57
 Haplophylline, F-90
 Haplophytine A, F-90
 Haplophytine B, F-223
 Haplophytine, H-58
 Haplopine prenyl ether, F-223
 Haplopine, F-223
 Haplosamine, H-59
 Haploscleridamine, L-199
 Haplosidine, F-223
 Haplosine, H-60
 Haplosinine, F-223
 Haplotubine, F-223
 Haplotubinone, H-61
 Haplotusine, Q-33
 Hapovine, H-657
 Harappamine, M-667
 Harepermine, C-290
 Hareperminoside, C-290
 Harmalacidine, T-226
 Harmalacine, H-62
 Harmalan N^B -oxide, M-426
 Harmalanine, H-63
 Harmalan, M-426
 Harmalicine, P-916
 Harmalidine, H-64
 Harmaline, H-592
 Harmalinic acid, H-459
 Harmalol, H-592
 Harman carboxamide, C-132
 Harman-3-carboxylic acid, M-427
 Harmanine, M-426
 Harman, M-426
 Harmic acid methyl ester, H-459
 Harmic amide, H-459
 Harmicine, H-65
 Harmidine, H-592
 Harmine N -oxide, H-592
 Harmine, H-592
 γ -Harmine, M-260
 Harmol, H-592
 Harmonine, O-35
 Harmony, D-234
 Harringtonine, H-66
 Harvatrate, T-632
 Harzianic acid, H-67
 Harzianopyridone, H-68
 Hastacine, H-69
 Hastanecine, H-233
 Hasubanonine, H-70
 Hasunohanin, H-71
 Haterumaimide A, D-354
 Haterumaimide B, H-74
 Haterumaimide C, H-73
 Haterumaimide D, H-74
 Haterumaimide E, H-74
 Haterumaimide F, H-75
 Haterumaimide G, H-75
 Haterumaimide H, H-75
 Haterumaimide I, H-72
 Haterumaimide J, H-75
 Haterumaimide K, H-75
 Haterumaimide L, H-74
 Haterumaimide M, H-74
 Haterumaimide N, D-354
 Haterumaimide O, D-354
 Haterumaimide P, D-354
 Haterumaimide Q, D-354
 Haterumalide A, H-76
 Haterumalide B, H-76
 Haterumalide E, H-76
 Haterumalide NA, H-76
 Haterumalide NB, H-76
 Haterumalide NC, H-76
 Haterumalide ND, H-76
 Haterumalide NE, H-76
 Haterumalide X, H-76
 Hatomamicin, H-77
 Haumanamide, D-763
 Havanine, H-78
 Havanine, S-199
 Hayatidine, C-420
 Hayatine, C-812
 Hayatinine, C-812
 Hazaleamide, T-129
 Hazuntamine, H-79
 Hazuntine, T-11
 Hazuntinine, T-11
 Hazuntiphyllidine, H-80
 Hazuntiphylline, H-81
 HBOA-Glc, H-409
 HBOA, H-409
 HDMBOA, D-538
 HDMBOA-Glc, D-538
 Hebeirubescensin A, H-82
 Hebridamine, H-83
 Hectochlorin, H-84
 Hectorine, H-518
 Hecubine, V-179
 Hederacine A, H-85
 Hederacine B, H-85
 Hediamine, H-86
 Hedonegamin, A-763
 Hedrantherine, M-666
 Hedycarine, C-54
 Hedyocapitelline, H-87
 Hedyocapitine, H-88
 Heilonine, C-300
 Heimidine, V-95
 Heimine, H-89
 Heinsiagenin A, H-90
 Heleurine N -oxide, A-669
 Heleurine, A-669
 Heliamine, T-158
 Helibracteatinine, D-596
 Helibracteatine, D-596
 Helibracteatinine, D-596
 Helibractinecine, D-596
 Helicocerin, C-286
 Helietidine, H-91
 Helifoline, D-597
 Helindicine, H-92
 Heliocoromandaline, T-443
 Heliocurassavicine N -oxide, T-443
 Heliocurassavicine, T-443
 Heliocurassavine, C-810
 Heliocurassavinine, T-443
 Heliohoustine, T-188
 Heliopsin, O-31
 Heliospathine, T-188
 Heliospathuline, H-93
 Heliosupine N -oxide, H-94
 Heliosupine, H-94
 Heliotridine, T-188
 Heliotrine N -oxide, H-95
 Heliotrine, H-95
 Heliotropamide, H-96
 Heliotropic acid, M-442
 Heliovicine, T-443
 Heliovinine, A-669
 Heliparvifoline, F-221
 Heliscabine, D-596
 Hellaradine, C-818
 Helquinoline, T-201
 Hemanthamine, H-5
 Hemanthidine, H-6
 Hemanthine, N-169
 Hemiargine A, T-248
 Hemiargine B, H-97
 Hemiargine C, M-559
 Hemiargine D, E-271
 Hemiargyrine, C-701
 Hemiasterlin A, H-98
 Hemiasterlin C, H-99
 Hemiasterlin, H-98
 Hemibastadin 1, H-100
 Hemibastadin 2, H-100
 Hemibastadin 3, H-100
 Hemibastadinol 1, H-101
 Hemibastadinol 2, H-101
 Hemibastadinol 3, H-101
 Hemicalyculin A, H-102
 Hemifistularin 3, H-103
 Hemileptagine, A-198

- m*-Hemipimide, D-533
m-Hemipinic acid, D-533
 Hemipyrocyanine, P-327
 Hemitoxiferine I, C-115
 Hemometina, E-72
 Hemsine A, H-104
 Hemsine B, H-105
 Hemsine C, H-106
 Hemsine D, H-107
 Hemsleyadine, M-306
 Hemsleyanidine, M-306
 Hemsleyanine A, M-306
 Hemsleyanine B, M-306
 Hemsleyanine C, M-319
 Hemsleyanine D, M-319
 Hemsleyanine, M-303
 Hemsleyanisine (incorr.), M-306
 Hemsleyasine, M-338
 Hemsleyatine, A-833
 Henderine, H-108
 3-(12*Z*-Heneicosenylidene)-2-methyl-1-pyrroline, L-37
 13-(12-Heneicosenyl)-1*H*-pyrrole-2-carboxaldehyde, A-623
 5-(6-Heneicosenyl)-1*H*-pyrrole-2-carboxaldehyde, H-109
 Henningsamide, H-110
 Henningsamine, C-115
 Henningsiine *N*-oxide, H-111
 Henningsiine, H-111
 Henningsoline, C-115
 Hennoxazole A, H-112
 Hennoxazole B, H-112
 Hennoxazole C, H-112
 Hennoxazole D, H-112
 Hennoxazole E, H-112
 Hennoxazole F, H-113
 Hennoxazole G, H-113
 Henrycinol A, H-114
 Henrycinol B, H-114
 Hepacholine, C-419
m-Hepimic anhydride, D-533
 1,5,9,13,17,21,25-Heptaazapentacosane, P-199
 2,4,4',5,5',6,6'-Heptabromo-2',3-bis(methylthio)-1,3'-bi-1*H*-indole, P-200
 2,3,3',4,4',5,5'-Heptachloro-1'-methyl-1,2'-bi-1*H*-pyrrole, H-115
^NB-Heptacosanoyl-4,5-dihydroxytryptamine, D-683
 1,9-Heptadecadiene-4,6-diene-3,8-diol, H-116
 14-Heptadecene-4,16-dienoic acid, H-117
 3-(12*Z*-Heptadecenylidene)-2-methyl-1-pyrroline, L-37
 5-(9*Z*-Heptadecenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 8-(13-Heptadecenyl)-1,5,9,13-tetraazacycloheptadecan-6-one, B-388
 3-Heptadecylidene-2-methyl-1-pyrroline, L-37
 5-(4,6-Heptadienyl)octahydro-8-methylindolizine, H-127
 1,2,3,4,6,9,14-Heptahydroxydihydro-β-agarofuran, H-118
 1,2,4,6,8,9,14-Heptahydroxydihydro-β-agarofuran, H-119
 2,3,5,7,8,9,15-Heptahydroxy-6(17),11-jatropha-dien-14-one, H-120
 3,5,7,8,9,13,15-Heptahydroxy-6(17),11-jatropha-dien-14-one, H-121
 1,3,6,13,14,16,18-Heptahydroxy-4-methylaconitan-15-one, H-122
 1,6,7,8,14,16,18-Heptahydroxy-4-methylaconitan-19-one, H-123
 1,6,8,13,14,16,18-Heptahydroxy-4-methylaconitan-19-one, H-124
 1-Heptanamine, H-130
N-Heptanoylsolamine, S-340
 Heptaphylline, H-125
 2-(1,3,5-Heptatrienyl)-3,4,5,6-tetrahydropyridine, N-213
 Heptazolicine, M-757
 Heptazolidine, P-883
 Heptazoline, H-125
 2-(3-Heptenyl)-2,6-dihydro-3-methyl-5*H*-pyrano[3,2-*c*]pyridin-5-one, A-1166
 2-(2-Heptenyl)-4-hydroxy-3-methylquinoline, H-126
 2-(1-Heptenyl)-4-hydroxyquinoline, H-133
 5-(4-Heptenyl)-8-methylindolizidine, H-127
 5-(6-Heptenyl)-8-methylindolizidine, H-127
 2-(2-Heptenyl)-3-methyl-4-quinolinol, H-126
 2-(2-Heptenyl)-3-methyl-4(1*H*)-quinolinone, H-126
 5-(4-Heptenyl)octahydro-8-methylindolizine, H-127
 2-(6-Heptenyl)-6-(4-pentenyl)piperidine, H-128
 2-(1-Heptenyl)-4(1*H*)-quinolinone, H-133
 2-(4-Heptenyl)-4(1*H*)-quinolinone, H-133
 2-(5-Heptenyl)-6,7,8,8*a*-tetrahydro-3-methyl-4*H*-pyrrolo[2,1-*b*][1,3]oxazin-4-one, H-129
 1-Heptylamine, H-130
 2-Heptyl-1-cyclopropanepropanoic acid, H-131
 2-Heptyl-1,2-dihydro-2-hydroxy-6-methoxy-4-methyl-3*H*-indol-3-one, M-183
 5-Heptyl-6,7-dihydro-3-methoxy-2-methyl-4,8(1*H*,5*H*)-quinolinedione, A-1325
 1-[4-[4-Heptylhexahydro-5-methyl-2-oxo-1,5-diazocin-1(2*H*)-yl]butyl]hexahydro-5-methyl-4-pentyl-1,5-diazocin-2(1*H*)-one, H-354
 2-Heptyl-2-hydroxy-6-methoxy-4-methylpseudooxindole, M-183
 1-Heptyl-3-hydroxy-2-methyl-9*H*-carbazole, H-132
 2-Heptyl-4-hydroxy-3-methylquinoline, H-126
 2-Heptyl-4-hydroxyquinoline *N*-oxide, H-133
 3-Heptyl-3-hydroxy-2,4(1*H*,3*H*)-quinolinedione, H-134
 2-Heptyl-4-hydroxyquinoline, H-133
 2-Heptyl-3-hydroxy-4(1*H*)-quinolinone, A-621
 2-Heptyl-8-methoxy-1-methyl-4(1*H*)-quinolinone, H-133
 2-Heptyl-4-methoxyquinoline, H-133
 1-Heptyl-2-methyl-3*H*-carbazole-3,4(9*H*)-dione, C-129
 1-Heptyl-2-methyl-9*H*-carbazol-3-ol, H-132
 5-Heptyl-8-methylindolizidine, H-127
 3-Heptyl-2-methylprodiginine, P-636
 3-Heptyl-5-methylpyrrolizidine, H-135
 2-Heptyl-3-methyl-4-quinolinol, H-126
 2-Heptyl-1-methyl-4(1*H*)-quinolinone, H-133
 2-Heptyl-3-methyl-4(1*H*)-quinolinone, H-126
 2-Heptylpyrrole sulfamate, H-136
 2-Heptylpyrrole, H-136
 2-Heptyl-3,4-quinolinediol, A-621
 2-Heptyl-4-quinolinol, H-133
 2-Heptyl-4(1*H*)-quinolinone, H-133
 Herbaceine, H-137
 Herbadine, H-139
 Herbaine, H-137
 Herbaline, C-117
 Herbamide A, H-138
 Herbamine, H-139
 Herbavine, M-50
 Herbavinine, M-50
 Herbicidal substance 1328-2, C-852
 Herbicidal substance 1328-3, C-852
 Herbindole A, H-140
 Herbindole B, H-141
 Herbindole C, H-142
 Herbiopline, H-143
 Herboksine, M-50
 Herboxine, M-50
 Herclavine, H-144
 Herculin, D-884
 Hericenone B, H-145
 Hericerin, H-145
 Hermidin, H-146
 Hermitamide A, P-342
 Hermitamide B, T-639
 Hernagine, T-248
 Hernandaline, H-147
 Hernandalinol, H-147
 Hernandezine 2'-*N*-oxide, H-148
 Hernandezine, H-148
 Hernandifoline, L-244
 Hernandine, H-149
 Hernandine, L-244
 Hernandine, P-209
 Hernandoline, H-70
 Hernandolinol, H-70
 Hernandonine, H-150
 Hernangerine, N-25
 Hernanymphine, O-195
 Hernovine, O-143
 Hernovine, T-248
 Herperal, D-866
 Herpestine, H-151
 Herquiline A, H-152
 Herquiline B, H-153
 Herveine A, H-154
 Herveine B, H-154
 Herveine C, H-154
 Herveine D, H-154
 Hervine, I-319
 Hesperaline, C-419
 Heteratisine, H-157
 Heteroauxin, I-81
 Heterocarpine, H-158
 Heterocodeine, M-704
 Heteromine A, H-143
 Heteromine B, H-143
 Heteromine C, H-143
 Heteromine D, H-143
 Heteromine E, H-143
 Heteromine F, H-159
 Heteromine G, H-159
 Heteromine H, H-159
 Heterophyllidine, H-157
 Heterophylline, A-1423
 Heterophylline, H-157
 Heterophylline, M-502
 Heterophylline, T-256
 Heterophyllisine, H-157
 Heterophylloidine, H-161
 Heteropsine, H-160
 Heteroxanthine, X-5
 Het₃₈₉, H-155
 Het₄₀₃, H-156
 Hetidine, H-161
 Hetisan-2,15-diol, H-163
 Hetisan-6,11-diol, H-164
 Hetisan-1,15-diol, H-162
 Hetisan-11,15-diol, H-167
 Hetisan-11,18-diol, H-168
 Hetisan-6,13-diol, H-165
 Hetisan-9,15-diol, H-166
 Hetisan-1,2,3,11,13,14-hexol, H-169
 Hetisan-2,3,9,11,13,14-hexol, H-170
 Hetisan-13-ol, H-171
 Hetisan-15-ol, H-172
 Hetisan-11-one, H-173
 Hetisan-13-one, H-171
 Hetisan-1,2,11,13,19-pentol, H-175
 Hetisan-1,2,3,11,13-pentol, H-174
 Hetisan-2,3,11,13,14-pentol, H-178
 Hetisan-2,3,6,11,13-pentol, H-176
 Hetisan-2,3,7,9,15-pentol, H-177
 Hetisan-1,2,9,15-tetrol, H-179
 Hetisan-1,2,11,13-tetrol, H-181
 Hetisan-1,2,9,19-tetrol, H-180
 Hetisan-2,11,13,14-tetrol, H-189
 Hetisan-2,11,13,15-tetrol, H-190
 Hetisan-2,11,13,19-tetrol, H-191
 Hetisan-2,3,11,13-tetrol, H-185
 Hetisan-2,3,6,13-tetrol, H-182
 Hetisan-2,3,7,13-tetrol, H-183
 Hetisan-2,3,9,15-tetrol, H-184
 Hetisan-2,6,7,13-tetrol, H-186
 Hetisan-2,7,13,15-tetrol, H-188
 Hetisan-2,7,9,15-tetrol, H-187
 Hetisan-6,11,12,15-tetrol, H-192
 Hetisan-7,11,15,19-tetrol, H-193
 Hetisan-2,11,13-triol, H-202
 Hetisan-2,6,13-triol, H-198
 Hetisan-1,2,13-triol, H-194
 Hetisan-1,2,15-triol, H-195
 Hetisan-1,7,13-triol, H-196
 Hetisan-1,7,15-triol, H-197
 Hetisan-11,15,19-triol, H-209
 Hetisan-11,15,20-triol, H-210

- Hetisan-2,11,15-triol, H-203
 Hetisan-2,7,11-triol, H-199
 Hetisan-2,9,15-triol, H-200
 Hetisan-2,9,19-triol, H-201
 Hetisan-3,15,19-triol, H-204
 Hetisan-6,11,15-triol, H-205
 Hetisan-6,13,15-triol, H-206
 Hetisan-7,11,13-triol, H-207
 Hetisan-7,11,15-triol, H-208
 Hetsine, H-202
 Hetsinone, H-202
 1,5,7,13,17,22-Hexaazadocosane, D-283
 1,5,9,13,18,22-Hexaazadocosane, D-284
 1,5,9,14,18,22-Hexaazadocosane, D-285
 1,5,9,13,17,21-Hexaazaheneicosane, T-107
 1,6,11,16,21,26-Hexaazahexacosane, D-286
 2,2',3,4',5,5'-Hexabromo-1,3'-bi-1H-indole, H-211
 3,3',4,4',5,5'-Hexabromo-2,2'-bi-1H-pyrrole, H-212
 2,4,4',5',6,6'-Hexabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole, P-200
 2,4,4',5,6,6'-Hexabromo-2',3-bis(methylthio)-1,3'-bi-1H-indole, P-200
 3,3',4,4',5,5'-Hexabromo-1,1'-dimethyl-2,2'-bi-1H-pyrrole, H-212
 1,1,1,15,15,15-Hexachloro-3,12-pentadecadiyn-8-amine, H-213
 Hexacosanoic acid isobutylamide, H-214
 Hexacosanoic acid, H-214
 N^b-Hexacosanolytryptamine, T-639
 2,9-Hexadecadiene-12,14-dienoic acid isobutylamide, H-215
 2,9-Hexadecadiene-12,14-dienoic acid, H-215
 2,4-Hexadecadienoic acid piperidine, H-216
 2,7-Hexadecadienoic acid pyrrolide, H-217
 3,6-Hexadecadienoic acid pyrrolidide, P-949
 2,4-Hexadecadienoic acid, H-216
 2,7-Hexadecadienoic acid, H-217
 N-(3,6-Hexadecadienyl)pyrrolidine, P-949
 6,8-Hexadecadien-10-ynoic acid pyrrolidide, H-219
 6,8-Hexadecadien-10-ynoic acid, H-219
 2,7-Hexadecadien-10-ynoic acid, H-218
 2,4,6,8,10,12,14-Hexadecaheptaenedioic acid, H-220
 Hexadecanamide, H-221
 Hexadecanoic acid 2-phenylethylamide, P-342
 Hexadecanoic acid pyrrolidide, P-949
 Hexadecanoic acid, H-221
 N-Hexadecanoylarginine, A-1413
 N²-Hexadecanoylglutamine, G-108
 N-Hexadecanoylputrescine, B-422
 N-Hexadecanoylpyrrolidine, P-949
 Hexadecanoylsolamine, S-340
 N^b-Hexadecanolytryptamine, T-639
 N-Hexadecanoyltyrosine, T-700
 3,5,10,13-Hexadecatetraen-2-amine, H-222
 2,6,8,12-Hexadecatetraen-10-ynoic acid dehydro-pyrrolidide, H-223
 2,6,8,12-Hexadecatetraen-10-ynoic acid isobutylamide, H-223
 2,6,8,12-Hexadecatetraen-10-ynoic acid, H-223
 2,6,8-Hexadecatrien-10-ynoic acid pyrrolide, H-224
 2,6,8-Hexadecatrien-10-ynoic acid, H-224
 2,7-Hexadeca-10-ynoic acid pyrrolide, H-218
 2,7-Hexadeca-10-ynoic acid pyrrolidide, H-218
 2-Hexadecenoic acid pyrrolidide, P-949
 N-(9Z-Hexadecenyl)arginine, A-1413
 N²-(9-Hexadecenyl)glutamine, G-108
 N-(2-Hexadecenyl)pyrrolidine, P-949
 N-(2-Hexadecenyl)solamine, S-340
 N-(9-Hexadecenyl)tyrosine, T-700
 7-Hexadecen-10-ynoic acid pyrrolide, H-225
 7-Hexadecen-10-ynoic acid, H-225
 3-Hexadecylidene-2-methyl-1-pyrroline, L-37
 N-Hexadecyl-N-nitrosohydroxylamine, N-254
 8-Hexadecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-385
 6,7,8,9,10,14-Hexadecydro-4,5-epoxy-3,6-dimethoxy-17-methylthebinan, H-226
 7,8,11,12,13,14-Hexadehydromatridin-15-one, S-376
 1,2,6,7,10,11-Hexadecydro-3-methoxy-15,16-[methylenebis(oxy)]erythrinan-8-one, C-799
 3,4,5,6,18,19-Hexadecydrochroprosinine, T-136
 Hexadellin A, A-1378
 Hexadellin B, H-227
 Hexadellin C, H-228
 3-(3,5-Hexadiynyl)oxiranecarboxylic acid, E-137
 3-(3,5-Hexadiynyl)-N-(2-phenylethyl)oxiranecarboxamide, E-137
 3-(3,5-Hexadiynyl)-N-(2-phenylethyl)oxiranecarboxamide, E-137
 Hexahydroaniline, C-868
 3,4,5,6,7,8-Hexahydro-1,7-benzodiazecin-2(1H)-one, H-229
 Hexahydroberberonic acid, P-457
 1',2,2',3,3',9'-Hexahydro[1,9'-bipyrrolo[2,1-b]quinazolin]-9(1H)-one, D-812
 1,2,3,3a,6,7-Hexahydrocyclopenta[d]pyrrolo[2,1-b][1,3]oxazin-8(5H)-one, M-77
 1,4,5,6,7,8-Hexahydro-1,3-diazocine-2,4-dicarboxylic acid, H-230
 1,2,3,10,11,11a-Hexahydro-8,11-dihydroxy-3,7-dimethoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, A-1232
 1a,2,4,9,10,10a-Hexahydro-5,10-dihydroxy-1a-methyl-2,4,9-trioxo-3H-benz[b]oxireno[h]carbazole-3-carbonitrile, K-37
 Hexahydro-1,2-dihydroxy-1H-pyrrolizine-3,5-dimethanol, D-543
 Hexahydro-1,7-dihydroxy-1H-pyrrolizine-1-methanol, D-596
 Hexahydro-6,7-dihydroxy-1H-pyrrolizine-1-methanol, D-595
 4-(7,9,10,11,11a,12-Hexahydro-2,3-dimethoxybenzof]pyrrolo[1,2-b]isoquinolin-6-yl)-2-butanone, V-122
 4-(7,9,10,11,11a,12-Hexahydro-2,3-dimethoxybenzof]pyrrolo[1,2-b]isoquinolin-6-yl)-3-buten-2-ol, V-122
 1,2,3,5,6,10b-Hexahydro-8,9-dimethoxypyrrolo[2,1-a]isoquinoline, C-756
 5b,6,7,12b,13,14-Hexahydro-5b,13-dimethyl[1,3]-benzodioxolo[5,6-c]-1,3-dioxolo[4,5-f]phenanthridin-6-ol, C-692
 Hexahydro-3a,7a-dimethyl-4,7-epoxy-2H-isoin-dole-1,3-dione, C-89
 1,2,3,4,5,13-Hexahydro-5,7-dimethyl-2-(1-methylethynyl)-1,5-methanooxocino[3,2-a]carbazole, C-814
 1,2,3,4,5,13-Hexahydro-5,7-dimethyl-2-(1-methylethyl)-1,5-methanooxocino[3,2-a]carbazole, C-814
 Hexahydro-2,2-dimethyl-7'-nitrospiro[cyclopentane-1,1'-[1H]pyrrolizine], N-252
 Hexahydro-2,4-dioxopyrimidine, D-497
 1,3,4,6,8,9-Hexahydrodipyrano[3,4-b:3',4'-e]pyrazine-3,8-dimethanol, P-41
 2,3,4,5,11,12-Hexahydro-6H,13aH-3a,5a-ethano-1H-indolizino[8,1-cd]carbazole, A-1494
 Hexahydro-7a-(1-ethoxyethoxy)-3H-pyrrolizin-3-one, H-234
 Hexahydrofugapavine, L-214
 1,2,3,3',11,11'-Hexahydro-3,3,3',3',5,5'-hexamethyl-1,8'-bipyranol[3,2-a]carbazole, M-748
 Hexahydro-3-(4-hydroxybenzyl)pyrrolo[1,2-a]pyrazine-1,4-dione, C-900
 1,2,3,5,6,7-Hexahydro-11-hydroxy-2,12-dimethoxydibenzo[b,f]quinolizin-9(14H)-one, C-564
 1-[4-[Hexahydro-4-(2-hydroxyheptyl)-5-methyl-2-oxo-1,5-diazocin-1(2H)-yl]butyl]hexahydro-5-methyl-4-pentyl-1,5-diazocin-2(1H)-one, H-355
 1-[4-[Hexahydro-4-(2-hydroxyheptyl)-5-methyl-2-oxo-1,5-diazocin-1(2H)-yl]butyl]hexahydro-5-methyl-4-phenyl-1,5-diazocin-2(1H)-one, H-353
 Hexahydro-7-hydroxy-3-[(4-hydroxyphenyl)methyl]pyrrolo[1,2-a]pyrazine-1,4-dione, C-874
 α-(Hexahydro-4-hydroxy-2-iminopyrimidin-6-yl)glycine, T-657
 1,2,3,10,11,11a-Hexahydro-9-hydroxy-11-(1H-indol-3-yl)-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, T-411
 1,2,3,10,11,11a-Hexahydro-2-hydroxy-11-methoxy-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, A-5
 Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizine-1,2-diol, D-592
 Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizine-1,2,6,7-tetrol, T-261
 Hexahydro-3-hydroxymethyl-5-methyl-1H-pyrrolizine-1,2,7-triol, T-554
 Hexahydro-7-hydroxy-3-(2-methylpropyl)pyrrolo[1,2-a]pyrazine-1,4-dione, C-871
 Hexahydro-2-hydroxy-2-methyl-1H-pyrrolizine-1-carboxylic acid, H-231
 Hexahydro-3-hydroxymethyl-1H-pyrrolizine-1,2-diol, D-594
 Hexahydro-7-(hydroxymethyl)-1H-pyrrolizine-1,6-diol, D-597
 Hexahydro-3-(hydroxymethyl)-1H-pyrrolizine-1,2,6,7-tetrol, T-262
 Hexahydro-3-(hydroxymethyl)-1H-pyrrolizine-1,2,7-triol, T-555
 Hexahydro-7-hydroxy-3-methylpyrrolo[1,2-a]pyrazine-1,4-dione, C-838
 Hexahydro-3-[(4-hydroxy-3-nitrophenyl)methyl]pyrrolo[1,2-a]pyrazine-1,4-dione, C-900
 1,2,3,4,5,6-Hexahydro-3-(3-hydroxy-2-oxobutyl)-1,5-methano-8H-pyrido[1,2-a][1,5]diazocin-8-one, S-378
 3,4,6,7,8,9-Hexahydro-10-hydroxy-2,2,6,6,9-pentamethyl-2H,14H-dipyrano[2,3-α:2',3'-c]acridin-14-one, M-401
 1,2,3,3a,7,7a-Hexahydro-3a-(4-hydroxyphenyl)-1-methyl-6H-indol-6-one, S-123
 Hexahydro-7-hydroxy-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, C-873
 Hexahydro-8a-hydroxy-3-(phenylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, C-872
 1,2,3,4,7,8-Hexahydro-6-(4-hydroxyphenyl)-8-(2,3,4,5-tetrahydroxypentyl)-2,4,7-pteridine-trione, H-232
 Hexahydro-6-hydroxy-1H-pyrrolizine-1-carboxylic acid, H-720
 Hexahydro-1-hydroxy-1H-pyrrolizine-1-methanol, H-517
 Hexahydro-2-hydroxy-1H-pyrrolizine-1-methanol, H-518
 Hexahydro-7-hydroxy-1H-pyrrolizine-1-methanol, H-233
 Hexahydro-7a-hydroxy-3H-pyrrolizin-3-one, H-234
 4,6,7,12,12b,13-Hexahydro-1H-imidazo[4,5-g]indolo[2,3-a]quinazoline, V-102
 Hexahydro-3-imino-1,2,4-oxadiazepine-3-carboxylic acid, H-235
 α-(Hexahydro-2-imino-4-pyrimidyl)glycine, C-99
 2,3,5,6,11,11b-Hexahydro-1H-indolizino[8,7-b]indole, H-65
 1-(1,2,3,5,6,8a-Hexahydro-8-indoliziny)-1-butanone, E-58
 N-[(Hexahydro-8(5H)-indolizinyldene)methyl]acetamide, T-73
 1,2,3,4,6,7-Hexahydroindolo[2,3-a]quinolizine, I-113
 Hexahydro-3-(1H-indol-3-ylmethyl)pyrrolo[1,2-a]pyrazine-1,4-dione, C-899
 1,2,3,4,5,6-Hexahydro-5-(1H-indol-3-yl)-4-oxoazepino[4,5-b]indole-2-carboxylic acid, M-66
 1,2,3,10,11,11a-Hexahydro-11-[4-(1H-indol-3-yl)-2-oxo-3-butenylidene]-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one, H-236
 Hexahydro-3-isopropylpyrrolo[1,2-a]pyrazine-1,4-dione, C-901
 Hexahydrocambrine, L-214
 1,2,3,4,5,6-Hexahydro-1,5-methano-8H-pyrido[1,2-a][1,5]diazocin-8-one, C-938
 2,3,5,6,7,8-Hexahydro-3'-methoxy-4',9-dimethylspiro[1H-cyclopenta[b]pyrrolo[1,2-a]azepine-11(10H),2'(5'H)-furan]-5',10-dione, S-518
 11,12,13,14,14a,15-Hexahydro-6-methoxy-9H-1,3-dioxolo[6,7]phenanthro[9,10-b]quinolizin-12-ol, C-791
 4a,5,9,10,11,12-Hexahydro-3-methoxy-11-methyl-6H-benzofuro[3a,3,2-e][2]benzazepin-6-ol, G-3
 Hexahydro-7a-methoxy-3H-pyrrolizin-3-one, H-234
 20-(Hexahydro-1-methyl-1H-azepin-2-yl)pregn-5-ene-3,15-diol, R-79

- 20-(Hexahydro-1-methyl-1H-azepin-2-yl)pregn-5-ene-3,15,17-triol, V-7
Hexahydro-3-(3-methylbutyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-870
Hexahydro-1-methylene-1H-pyrrolizine, M-461
2,3,5,6,7,7a-Hexahydro-7-methylene-1H-pyrrolizine-1-ol, H-601
Hexahydro-3-(1-methylethyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-901
6-(Hexahydro-8-methyl-6(5H)-indolizinyldene)-5-methyl-2-hexanol, D-228
4-(1,2,3,5,8,8a-Hexahydro-7-methyl-6-indolizinylophenol, I-165
Hexahydro-2-methyl-4H,7H-methano-2H-1,3-benzoxazocine 3-oxide, S-292
Hexahydro-*N*-methyl-2,4-methano-4H-furo[3,2-*b*]pyrrol-3-amine, L-234
Hexahydro-4-methyl-2,5-methano-2H-furo[3,2-*b*]pyrrol-6-ol, S-160
Hexahydro-6-methyl-3-methylenepyrrolo[1,2-*a*]pyrazine-1,4-dione, A-1261
5,7,8,13,13b,14-Hexahydro-14-methyl-5-oxindolo[2',3':3,4]pyrido[2,1-*b*]quinazoline-7-carboxylic acid, E-310
2,3,6,7,8,9-Hexahydro-10-methyl-2-phenylcyclohepta[*b*]pyran-5,8-imin-4(5H)-one, S-586
1,2,3,4,5,6-Hexahydro-3-methyl-4-(2-propenyl)-1,5-methano-5H-pyrido[1,2-*a*][1,5]diazocin-8-one, T-412
Hexahydro- α -methyl-5-propyl-1H-pyrrolizine-3-hexanol, H-500
Hexahydro-3-(2-methylpropyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-876
Hexahydro-3-(1-methylpropyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-875
2,4a,5,6,7,7a-Hexahydro-7-methyl-1H-2-pyridine-4-carboxaldehyde, L-182
Hexahydro-7-methyl-1H-pyrrolizine-1-ol, H-630
Hexahydro- α -methylpyrrolo[2,1-*b*]oxazole-5-decanamine, A-927
Hexahydro- α -methylpyrrolo[2,1-*b*]oxazole-5-dodecanamine, A-920
Hexahydro-3-methylpyrrolo[1,2-*a*]pyrazine-1,4-dione, C-839
2,5,6,8,9,9b-Hexahydro-2-methyl-3H,4H,7H-1,2,3a,6a-tetraazaphenylene, C-443
Hexahydro-3-[2-(methylthio)ethyl]pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-880
1,2,3,4,7,8-Hexahydro-8-methyl-2,4,7-trioxopteridine, P-763
Hexahydro-6bH-oxireno[*a*]pyrrolizine-6b-methanol, E-133
2,3,5,6,11,11b-Hexahydro-3-oxo-1H-indolizino[8,7-*b*]indole-5-carboxylic acid, H-237
Hexahydro-2-oxo- α -methyl-1H-thieno[3,4-*d*]imidazole-4-pentanoic acid, M-402
1,2,3,10,11,11a-Hexahydro-11-(2-oxopropylidene)-5H-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one, H-238
4-(1,2,3,10,11,11a-Hexahydro-5-oxo-5H-pyrrolo[2,1-*c*][1,4]benzodiazepin-11-ylidene)-3-oxobutanoic acid, H-239
Hexahydro-2-oxo-1H-selenolo[3,4-*d*]imidazole-4-pentanoic acid, S-219
Hexahydro-2-oxo-1H-thieno[3,4-*d*]imidazole-4-pentanoic acid, B-139
Hexahydro-2-oxo-1H-thieno[3,4-*d*]imidazole-4-propanoic acid, B-214
5-(Hexahydro-2-oxo-1H-thieno[3,4-*d*]imidazol-4-yl)-2-pentenoic acid, L-166
9,11,12,13,13a,14-Hexahydro-2,3,4,6,7-pentamethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinoline, P-228
3,4,6,7,8,9-Hexahydrophenazine, H-240
Hexahydro-3-(phenylmethyl)-2H-pyrido[1,2-*a*]pyrazin-1(6H)-one, V-84
Hexahydro-3-(phenylmethyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-891
Hexahydrocolonic acid, P-456
1,2,3,5,6,8a-Hexahydro- α -propyl-8-indolizidine-methanol, E-58
7-(Hexahydro-5-propyl-1H-pyrrolizine-3-yl)-2-heptanone, H-500
Hexahydropyridine, P-454
Hexahydro-1H-pyrrolizine-1-amine, A-906
Hexahydro-1H-pyrrolizine-1-carboxylic acid, P-960
Hexahydro-1H-pyrrolizine-1-methanol, H-629
3,4,7,8,10,10a-Hexahydro-2H,5H-pyrrolo[2,1-*c*:3,4-*b'*]bis[1,4]thiazin-5-one, H-241
Hexahydropyrrolo[1,2-*a*]pyrazine-1,4-dione, C-865
N-[(Hexahydro-2H-quinolizin-1(6H)-ylidene)methyl]acetamide, L-305
1'',2'',3'',4'',5'',6''-Hexahydro-2,3':5,2''-terpyridine, A-963
7a,8,11,12,13,14-Hexahydro-11,14,16,17-tetrahydroxy-9-methoxy-3-methyl-2H-xantheno[2',3':6,7][1,3]benzodioxino[4,5-*f*]isoquinoline-1,15-dione, S-301
9,11,12,13,13a,14-Hexahydro-3,4,6,7-tetramethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinoline, I-342
9,11,12,13,13a,14-Hexahydro-2,3,5,6-tetramethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinoline, T-688
9,11,12,13,13a,14-Hexahydro-2,3,6,7-tetramethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinoline, T-695
9a,10,11,12,13,13a-Hexahydro-2,9,9,12-tetramethyl-1,12-epoxy-9H-indolo[3,2,1-*de*]phenanthridine, C-813
Hexahydro-6,7,8-trihydroxy-5-(hydroxymethyl)imidazo[1,2-*a*]pyridine-2,3-dione, K-41
Hexahydro-1,2,7-trihydroxy-1H-pyrrolizidine-3,5-dimethanol, T-529
9,11,12,13,13a,14-Hexahydro-2,3,6-trimethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinoline, A-1330
9,11,12,13,13a,14-Hexahydro-3,6,7-trimethoxydibenzof[*h*]pyrrolo[1,2-*b*]isoquinolin-14-ol, T-696
11,12,13,14,14a,15-Hexahydro-3,6,7-trimethoxy-9H-phenanthro[9,10-*b*]quinolizine, B-241
11,12,13,14,14a,15-Hexahydro-2,3,6-trimethoxy-9H-phenanthro[9,10-*b*]quinolizine, C-792
1,2,3,3a,4,5-Hexahydro-9,10,11-trimethoxypyrrolo[3,2,1-*de*]phenanthridinium(1+), R-130
2,3,4,4a,9,9a-Hexahydro-2,4a,9-trimethyl-1,2-oxazin[6,5-*b*]indol-6-ol methylcarbamate (ester), G-51
1,2,3,3a,8,8a-Hexahydro-1,3a,8-trimethylpyrrolo[2,3-*b*]indol-5-ol methylcarbamate (ester), P-400
Hexahydro-1,4,7,2-trioxazone, A-1603
1,2,3,4,5,6-Hexahydroxyacridone, H-242
1,2,3,8,9,10-Hexahydroxyaporphine, H-243
2,3,14,20,24,25-Hexahydroxycholesterol-7-en-6-one, H-244
3,5,7,13,15,18-Hexahydroxy-6,12-cyclo-14-lathyrone, H-245
1,2,4,6,8,9-Hexahydroxydihydro- β -agarofuran, H-246
1,2,4,6,9,14-Hexahydroxydihydro- β -agarofuran, H-247
1,2,6,8,9,14-Hexahydroxydihydro- β -agarofuran, H-248
1,4,6,8,9,14-Hexahydroxydihydro- β -agarofuran, H-249
1,3,4,6,8,13-Hexahydroxy-10,11-dimethyl-2,5-di-2-piperidinylphenanthro[1,10,9,8-*opqr*]perylene-7,14-dione, F-6
2,3,5,7,8,15-Hexahydroxy-6(17),11-jatrophiadiene-9,14-dione, H-250
2,3,5,7,9,15-Hexahydroxy-6(17),11-jatrophiadiene-14-one, H-251
2,3,5,8,9,15-Hexahydroxy-6(17),11-jatrophiadiene-14-one, H-252
1,6,13,14,16,18-Hexahydroxy-4-methylnonan-15-one, H-254
1,6,7,8,14,16-Hexahydroxy-4-methylnonan-19-one, H-253
3,5,7,13,15,17-Hexahydroxy-8,10(18)-myrsinadiene-14-one, H-255
3,8,14,15,16,18-Hexahydroxypregnan-20-one, H-256
2',3',4,4'',5',6'-Hexahydroxy-*p*-terphenyl, T-97
3,5-Hexalobine A, B-200
3,6-Hexalobine A, B-201
3,5-Hexalobine B, B-201
ent-3,6-Hexalobine C, B-201
3,5-Hexalobine C, B-200
3,6-Hexalobine C, B-201
3,5-Hexalobine D, B-200
(*E*)-3,6-Hexalobine E, B-201
(*Z*)-3,6-Hexalobine E, B-201
3,5-Hexalobine E, B-200
2,3-Hexalobine E, H-257
Hexalupine, T-363
*N*²,*N*²,*N*⁸,*N*⁸,1,4-Hexamethyl-1H-cyclohepta[1,2-*d*:3,4-*d'*]dimidazole-2,8-diamine, P-737
1,1,2,2,6,6-Hexamethyl-4-oxopiperidinium(1+), T-283
1,1,2,2,6,6-Hexamethyl-4-piperidinone, T-283
1-Hexanamine, H-261
Hexanediamide, H-258
Hexanedioic acid, H-258
20-Hexanoylcampthothecin, C-70
5-Hexanoyl-5-hydroxy-*N*-isobutyl-2-pyrrolidinone, P-449
20-Hexanoyl-10-methoxycampthothecin, C-70
N-Hexanoylnornicotine, P-957
N-Hexanoylsolamine, S-340
*N*⁵-Hexanoylspermidine, S-398
[3-(1-Hexenyl)-*ONN*-azoxy]-1,2-butanediol, A-1199
[3-(1-Hexenyl)-*ONN*-azoxy]-2-butanol, M-85
[3-(1-Hexenyl)-*ONN*-azoxy]-1-hydroxy-2-butanone, A-1199
2-(5-Hexenyl)-3,4-dihydro-5-nonyl-2H-pyrrole, H-259
2-(1-Hexenyl)-4-hydroxyquinoline, H-263
2-(5-Hexenyl)-1-methyl-5-(8-nonyl)pyrrolidine, D-246
2-(5-Hexenyl)-1-methyl-5-nonylpyrrolidine, D-246
2-(5-Hexenyl)-5-(8-nonyl)pyrrolidine, D-246
5-(5-Hexenyl)-2-(8-nonyl)- Δ^1 -pyrroline, D-246
2-(5-Hexenyl)-5-(8-nonyl)- Δ^1 -pyrroline, D-246
2-(5-Hexenyl)-5-nonyl- Δ^1 -pyrroline, D-435
3-(5-Hexenyl)octahydro-5-(4-pentenyl)indolizine, H-260
3-(5-Hexenyl)-5-(4-pentenyl)indolizidine, H-260
2-(5-Hexenyl)-5-pentylpyrrolidine, D-246
2-(1-Hexenyl)-4-quinolinol, H-263
2-(1-Hexenyl)-4(1H)-quinolinone, H-263
2-(3-Hexenyl)-1,2,3,4-tetrahydro-6H-pyrido[1,2-*a*]quinolin-6-one, D-367
22-(2-Hexenyl)-9,10,12-trihydroxy-17-methylazacyclodocosane-3,5,7,13,15,17,19-heptaen-2-one, A-1555
1-Hexylamine, H-261
5-Hexyl-6,7-dihydro-3-methoxy-2-methyl-4,8(1H,5H)-quinolinedione, A-1325
2-Hexyl-3,5-dimethylpyrazine, H-262
N-Hexylhexahydro-1,3-dimethyl-*N*-pentyl-2-pyrimidinedecanamine, A-13
2-Hexyl-4-hydroxyquinoline *N*-oxide, H-263
2-Hexyl-4-hydroxyquinoline, H-263
2-Hexyl-3-hydroxy-4(1H)-quinolinone, A-621
5-Hexylindolizidine, H-264
2-Hexyl-8-methoxy-1-methyl-4(1H)-quinolinone, H-265
1-Hexyl-2-methyl-3H-carbazole-3,4(9H)-dione, C-129
3-Hexyl-5-methylindolizidine, H-266
3-Hexyl-2-methylprodigine, P-636
5-Hexyloctahydroindolizine, H-264
3-Hexyloctahydro-5-methylindolizine, H-266
3-Hexylpyridine, H-267
2-Hexylpyrrole sulfamate, H-268
2-Hexylpyrrole, H-268
2-Hexyl-3,4-quinolindiol, A-621
2-Hexyl-4-quinolinol, H-263
2-Hexyl-4(1H)-quinolinone *N*-oxide, H-263
2-Hexyl-4(1H)-quinolinone, H-263
1-[(5-Hexyltetrahydro-2-furanyl)acetyl]-1H-pyrrole, H-269
Hexylthiobromine, T-355
Heyden Antibiotic 9, O-124
Heyneanine hydroxyindolenine, C-658
Heyneanine, C-657
Heyneatine, H-270

- HH 50, C-100
 Hibiscusamide, T-576
 Hibiscuwanin A, H-271
 Hibiscuwanin B, H-271
 Hibispeptin A, H-272
 Hibispeptin B, H-273
 Hicksoane A, H-274
 Hicksoane B, H-274
 Hicksoane C, H-274
 Hidorin M7, O-32
 Higenamine, H-275
 Himalayamine, H-276
 Himandravine, H-280
 Himandreline, H-277
 Himandridine, H-277
 Himandrine, H-277
 Himanimide A, H-278
 Himanimide B, H-278
 Himanimide C, H-278
 Himanimide D, H-278
 Himanthine, H-279
 Himbacine, H-280
 Himbadine, H-281
 Himbeline, H-280
 Himbosine, H-277
 Himeic acid A, H-282
 Himeic acid B, H-282
 Himeic acid C, H-283
 Himeradine A, H-284
 Himgaline, H-285
 Himgravine, H-280
 Himgrine, H-286
 Hinckdentine A, H-287
 Hindaricine, G-77
 Hindarine, T-212
 Hindarinine, P-34
 Hinnuliquinone, H-288
 Hippacine, H-289
 Hippacine, P-586
 Hippadine, H-290
 Hippamine, L-348
 Hippandrine, H-291
 Hippauline, H-292
 Hippawine, H-6
 Hippeastidine, H-293
 Hippeastrine *N*-oxide, H-294
 Hippeastrine, H-294
 Hippocasine *N*-oxide, H-295
 Hippocasine, H-295
 Hippocrateine III, H-296
 Hippocrateine II, P-246
 Hippocrateine I, P-246
 Hippodamine, D-886
 Hippophaine, H-755
 Hippospongin C, H-297
 Hippospongin E, H-298
 Hippospongin F, H-298
 Hippospongine, D-731
 Hiptagin, G-104
 Hirsutine, C-689
 Hirsutellone A, H-299
 Hirsutellone B, H-300
 Hirsutellone C, H-300
 Hirsutellone D, H-301
 Hirsutellone E, H-302
 Hirsutellone F, H-303
 Hirsutine *N*-oxide, C-689
 Hirsutine, C-689
 Δ^{18} -Hirsutine, C-689
 Hirsutine, J-17
 Hirudinoidine A, H-304
 Hirudinoidine B, H-304
 Hirudinoidine C, H-304
 Hirudonine, H-305
 Hispaconitine, A-85
 Hispidine, S-248
 Hispidospermidine, H-306
 Histamine dihydrochloride, H-307
 Histamine phosphate, H-307
 Histamine, H-307
 Histaminol, I-38
 Histamyl, H-307
 Histapon, H-307
 Histidinol, H-308
 Histidol, H-308
 Δ^{17} -*trans*-Histrionicotoin 283A, H-309
 Histrionicotoin D, G-65
 Histrionicotoin, H-309
 HL 523, G-201
 HMBOA, D-538
 HMBOA-Glc, D-538
 HMQ, H-126
 HO 452, A-176
 HO 416b, I-82
 HO 395, H-413
 Hobartidiol, H-310
 Hobartine, H-310
 Hodgkinsine B, H-311
 Hodgkinsine, H-311
 Hodgsonine, C-533
 Hodorine, H-312
 Hoe 933, H-554
 Hoerhammericine, H-743
 Hoerhammerinine, H-743
 Hoffmannseggiamide A, P-462
 Hoffmannseggiamide B, P-462
 Hoheconsoline, M-323
 Hohenackeridine, D-187
 Hohenackerine, M-325
 Hokbusine A, M-331
 Hokbusine B, M-355
 Holacetine, A-881
 Holacimine, H-313
 Holacine, K-113
 Holadienine, H-314
 Holadysamine, A-875
 Holadysenterine, D-279
 Holadysine, A-887
 Holafebrine, A-886
 Holafrine, H-318
 Holaline, H-315
 Holamide, H-314
 Holamine, A-888
 Holaminol, A-885
 Holantogenin, H-316
 Holantosine A, H-316
 Holantosine B, A-1028
 Holantosine C, H-316
 Holantosine D, A-1028
 Holantosine E, H-316
 Holantosine F, A-1028
 Holaphyllamine, A-888
 Holaphyllidine, A-885
 Holaphylline, A-888
 Holaphyllinol, A-885
 Holaromine, H-317
 Holarrheline, H-318
 Holarrhenine, H-318
 Holarrhesine, H-318
 Holarrhessimine, H-319
 Holarrhetine, H-318
 Holarrhidine, D-279
 Holarrhimine, D-279
 Holarrhine, H-320
 Holarricine, D-257
 Holarrifine, P-606
 Holeinine, R-51
 Holidine, H-321
 Holidimine, C-588
 Holomycin, A-759
 Holonamine, H-322
 Holothin, A-759
 Holstiine, H-323
 Holstiline, H-323
 Holyrine A, A-1190
 Holyrine B, A-1190
 Homaline, H-324
 Homarine, C-148
 Homoerthionin, A-158
 Homoagmatine, A-859
 α -Homoallonjirimycin, B-186
 Homoamericine, T-300
 Homoanatoxin a, H-325
 Homoanisylamine, M-280
 HomoAnTx, H-325
 Homoarecoline, T-205
 Homoarginine, H-326
 Homoaromoline, O-1
 Homobatrachotoxin, B-36
 Homobuxaquamarine, H-327
 Homocaldohexamine, D-283
 Homocaldopentamine, D-288
 Homocapsaicin, C-100
 Homochasmanine, M-323
 α -Homochelidonine, C-352
 Homochelidonine, C-352
 β -Homochelidonine, F-2
 γ -Homochelidonine, F-2
 Homocinchonidine, H-328
 Homoclausenamide, D-445
 Homocryptolepinone, H-329
 Homocylindrocarpidine, C-929
 Homodaphniphylic acid, H-330
 Homodasycarpine, J-16
 Homodeoxylharringtonine, H-66
 Homodihydrocapsaicin, C-100
 Homodolastatin 3, D-905
 Homodopamine, A-896
 Homoepilachnene, E-93
 Homo-6-epipodopetaline, H-341
 Homoerysotine, H-331
 Homoerythratine, S-126
 Homofascaplysate A, H-332
 Homofascaplysin A, H-333
 Homofascaplysin B1, H-335
 Homofascaplysin B, H-334
 Homofascaplysin C, H-336
 Homofluorodaturatin, H-337
 β -Homofuconojirimycin, H-607
 α -Homogalactostatin, B-186
 Homoglomerine, E-247
 Homoglycosolone, T-583
 Homoharringtonine, H-66
 Homoharizianic acid, H-67
 Homohexamine, D-286
 Homohydroxydolabellin, H-503
 Homohygrinic acid, P-950
 Homoindanomycin, I-64
 Homolaudanosine, H-338
 Homoleucylprolyl anhydride, C-870
 Homolinearisine, P-643
 γ -Homolinolenic acid, E-52
 Homo- γ -linolenylethanolamide, E-52
 Homolycorine *N*-oxide, L-347
 Homolycorine, L-347
 α -Homomannoijirimycin, B-186
 β -Homomannoijirimycin, B-186
 Homomoenjoaramine, H-339
 Homomoschatoline, M-712
 Homoneoharringtonine, N-117
 α -Homonoijirimycin, B-186
 β -Homonoijirimycin, B-186
 Homoormosamine, J-16
 Homopahutoxin, P-12
 Homopalytoxin, P-42
 Homopentamine, D-290
 Homophleine, H-340
 Homopiptanthine, H-341
 Homoplatensimide, H-342
 Homopodopetaline, H-341
 β -Homoproline, P-950
 Homoproline, P-456
 Homopseudopteraxazole, H-343
 Homopseudozoanthoxanthin, P-737
 Homopumiliotoxin 317, A-320
 Homopumiliotoxin 319A, H-344
 Homopumiliotoxin 319B, H-344
 Homopumiliotoxin 321B, H-344
 Homopumiliotoxin 249F, A-512
 Homopumiliotoxin 207G, A-517
 Homopumiliotoxin 223G, H-345
 Homopumiliotoxin 235J, A-534
 Homopumiliotoxin 251L, A-546
 26-Homosafraamycin Y₃, S-12
 Homosecodaphniphylic acid, H-346
 Homosedinone, H-347
as-Homospermidine, A-897
sym-Homospermidine, D-252
 Homospermine, B-150

- Homostachydrine, H-348
 Homostephanoline, H-70
 Homosweetinine, H-341
 Homothalricine, O-1
 Homothapoxepine A, T-348
 Homothapsakin A, T-349
 Homothermohexamine, D-285
 Homothermopsine, H-349
 Homotoluic acid, P-350
 Homotrilobine, I-339
 Homotrypanothione, T-636
 Homotyramine, H-350
 Homovanillylamine, H-566
N-Homoveratrolylhomoveratrylamine, H-351
 Homoveratrylamine, D-718
 Honyumine, J-59
 Hookerianamide A, D-273
 Hookerianamide B, D-262
 Hookerianamide C, D-265
 Hookerianamide H, D-281
 Hookerianamide I, P-601
 Hookerianamide A, P-607
 Hopamidine, P-943
 Hopeanine, H-352
 Hopromalinol, H-353
 Hopromine, H-354
 Hoprominol, H-355
 Hordatine A glucoside, H-356
 Hordatine A, H-356
 Hordatine B glucoside, H-356
 Hordatine B, H-356
 Hordenine *O*-(6'-*O*-*trans*-cinnamoyl-3'-*O*-β-D-glucopyranosyl-α-L-rhamnopyranoside), H-357
 Hordenine *O*-(6'-*O*-*p*-coumaroyl-3'-*O*-β-D-glucopyranosyl-α-L-rhamnopyranoside), H-357
 Hordenine *O*-glucoside, H-357
 Hordenine *O*-rhamnoside, H-357
 Hordenine, H-357
 Horhammericine, H-743
 Horhammerinine, H-743
 Hormaomycin, H-358
 Hormex, I-83
 Hormodin, I-83
 Horridine, P-247
 Horsfiline, H-359
 Hortesin, H-360
 Hortiacine, H-361
 Hortiamide, H-667
 Hortiamine, D-158
 Hostasine, H-362
 Hostasinine A, H-363
 Hosukinidine, H-364
 Hoveine, H-365
 Hovenine A, F-151
 HPPA, H-689
 HQ 24A, T-370
 HS 3, P-974
 5 HT, H-755
 HTX-235A, H-309
 HTX-259A, H-309
 HTX-283A, H-309
 HTX-285A, H-309
 HTX-287A, H-309
 HTX-291A, H-309
 Δ¹⁷-*trans*-HTX-283A, H-309
 HTX-285B, H-309
 HTX-287B, H-309
 HTX-285C, H-309
 HTX-287D, H-309
 HTX-285E, H-309
 Huajiaosimuline, Z-5
 Huangshanine, H-366
 Huazhongilexol, L-213
 huC242-DM1, M-135
 Humantenedine, G-44
 Humantenine, H-367
 Humanteniridine, R-15
 Humantenirine, H-367
 Humantenmine, G-44
 Humilixanthin, V-195
 Humosine A, E-43
 huN901-DM1, M-135
 Hunnemannine, F-2
 Huntabrine, G-33
 Hunteracine, H-368
 Hunterburnine α-*N*-metho salt, A-1328
 Hunterburnine β-*N*-metho salt, A-1328
 Hunterburnine, A-1328
 Hunteriamine, H-369
 Hunteriatryptamine, C-281
 Huntericine, E-6
 Hunteridine, E-6
 Hunteriline, E-6
 Hunterine, U-21
 Hunterioside B, S-583
 Hunterioside, S-583
 Huntrabrine *N*-metho salt, G-33
 Huntrabrine, G-33
 Hupehemonoside, C-298
 Hupehenidine, H-370
 Hupehenine, C-290
 Hupeheninoside, C-290
 Hupehenirine, C-290
 Hupehenisine, H-371
 Hupehenizine, C-290
 Hupehenizioside, C-290
 Hupeol, H-372
 Huperserratinine, S-257
 Huperzine A, H-375
 Huperzine B, H-376
 Huperzine C, H-388
 Huperzine D, H-388
 Huperzine E', H-377
 Huperzine E, H-378
 Huperzine F', H-377
 Huperzine F, H-378
 Huperzine G, F-75
 Huperzine H, H-379
 Huperzine I, H-373
 Huperzine J, H-380
 Huperzine K, H-380
 Huperzine L, H-380
 Huperzine O, H-378
 Huperzine P, H-381
 Huperzine Q *N*-oxide, H-382
 Huperzine Q, H-382
 Huperzine R, H-383
 Huperzine S, H-384
 Huperzine T, H-385
 Huperzine U, H-386
 Huperzine V, H-374
 Huperzine W, H-387
 Huperzinine C, H-389
 Huperzinine N¹³-oxide, H-388
 Huperzinine, H-388
 Hurghamide A, A-841
 Hurghamide B, A-841
 Hurghamide C, A-841
 Hurghamide D, A-841
 Hurghamide E, A-841
 Hurghamide F, A-841
 Hurghamide G, A-841
 Hyacinthacine A₁, D-594
 Hyacinthacine A₂, D-594
 Hyacinthacine A₃, D-592
 Hyacinthacine A₄, D-592
 Hyacinthacine A₅, D-592
 Hyacinthacine A₆, D-592
 Hyacinthacine A₇, D-592
 Hyacinthacine B₁, D-543
 Hyacinthacine B₂, D-543
 Hyacinthacine B₃, T-554
 Hyacinthacine B₄, T-554
 Hyacinthacine B₅, T-554
 Hyacinthacine B₆, T-554
 Hyacinthacine B₇, T-554
 Hyacinthacine C₁, T-261
 Hyacinthacine C₂, T-529
 Hyacinthacine C₃, T-529
 Hyacinthacine C₄, T-261
 Hyacinthacine C₅, T-261
 Hyacinthoside, S-345
 Hyalbidone, E-226
 Hyatelactam, H-390
 Hybar X, U-42
 Hydantoin, I-40
 Hyderabadine, H-391
 Hydnellin A, S-76
 Hydnellin B, H-392
 Hydrachine A, N-110
 Hydrakouminol, H-393
 Hydrastidine, H-394
 Hydrastine a, H-394
 Hydrastine b, H-394
 Hydrastine imide, N-36
 Hydrastine, H-394
 (±)-α-Hydrastine, H-394
 (±)-β-Hydrastine, H-394
 (+)-α-Hydrastine, H-394
 (+)-β-Hydrastine, H-394
 (-)-α-Hydrastine, H-394
 (-)-β-Hydrastine, H-394
 1,2-Hydrazinedicarboxylic acid, H-395
 Hydrazobisformic acid, H-395
 Hydrazodicarboxylic acid, H-395
 Hydrazodiformic acid, H-395
 Hydrazoformic acid, H-395
 Hydrocafeic acid, D-652
 1,3-(8-Hydrochelerythrinyl-8'-hydro-*N*-norchelerythrinyl)acetone, C-355
 Hydrocinchonidine, C-446
 Hydrocinchonine, C-447
 Hydrocinnamic acid, P-350
 Hydrocinnamonitrile, P-350
 Hydroconchinene, Q-18
 Hydroconquinine, Q-18
 Hydrocotarnine, H-396
 Hydrocotyline, H-397
 Hydrocupreine, Q-20
 Hydroergotocin, Y-13
 Hydrogen cyanamide, C-823
 Hydrohydrastinine, H-398
 7-Hydro-8-methylpteroylglutamylglutamic acid, P-772
 7-Hydroperoxy-17-hydroxy-16,18-dimethyl-10-phenyl[1]cytochalasa-5,13,19-triene-1,21-dione, H-399
 5-(2-Hydroperoxy-3-methyl-3-butenyl)-4,6,7-trimethoxyfuro[2,3-*b*]quinoline, H-580
 5-(3-Hydroperoxy-3-methyl-1-butenyl)-4,6,7-trimethoxyfuro[2,3-*b*]quinoline, H-578
 7-Hydroperoxyphlegmariurine B, P-363
 11α-Hydroperoxyphlegmariurine B, P-363
 β-Hydropiperilpiperidine, P-467
 Hydroquine, Q-18
 Hydroquinidine, Q-18
 Hydroquinine, Q-20
 Hydrorhombinine, L-302
 Hydrosphilanthol, D-105
 Hydrouracil, D-497
 4-Hydroxyacanthamine, D-436
 2-Hydroxyacetanilide, A-861
 10-Hydroxyacetyl-dihydrolycopodine, P-71
 6-Hydroxy-13-*O*-acetylthetisonone, T-552
 3-(Hydroxyacetyl)-1-*H*-indole, H-400
 15α-Hydroxy-*N*-acetylirehdiamine F, D-278
 15β-Hydroxy-*N*-acetylirehdiamine F, D-278
 2-(Hydroxyacetyl)-1-*H*-pyrrole, H-401
N-Hydroxyacetylquestiomycin A, A-862
 10-Hydroxyaconitine, M-302
 2-Hydroxyaconitine, M-301
 9-Hydroxyacridine, A-112
 1-Hydroxy-9(10*H*)-acridinone, H-402
 1-Hydroxyacridone, H-402
 22-Hydroxyacuminatine, H-403
 8-Hydroxyadenine, A-819
 11-Hydroxyaerotherionin, A-158
 18-Hydroxyaffinisine, T-425
 20-Hydroxyaflavinine, A-172
 3'-Hydroxyaglaïne B, A-199
 3'-Hydroxyaglaïne C, A-199
 2'-Hydroxyaglaroxin B, P-72
 12-Hydroxyajmaline, A-216
 10-Hydroxyakagerine, A-222
 Hydroxyakalone, A-755
 10-Hydroxyakuammicine, A-228
 12-Hydroxyakuammicine, A-228

- 14-Hydroxyakuammicine, M-713
 17-Hydroxy- ψ -akuammicine, A-232
 16-Hydroxyalloibogamine, H-404
 17-Hydroxyalloibogamine, H-404
 4-Hydroxyallosedamine, S-210
 3-Hydroxyallosedamine, S-210
 16-Hydroxyalstonal, A-648
 16-Hydroxyalstonisine, A-654
 Hydroxyamfetamine, A-898
 γ -Hydroxy- β -aminobutyric acid, A-792
 4-Hydroxy- α -aminohydrocinnamic acid, T-700
 Hydroxyamphetamine hydrobromide, A-898
 Hydroxyamphetamine, A-898
 Hydroxyanagryne, A-970
 13-Hydroxyanagryne, B-20
 Hydroxyanantine, A-972
 2 α -Hydroxy-9-angeloyloxy(-)-trachelanthamidine N-oxide, H-518
 23-Hydroxyanguivine, S-450
 15-Hydroxyangustilobine A, A-1014
 10-Hydroxyangustine, A-1019
 4-Hydroxyanhydrolycorine, H-290
o-Hydroxyaniline, A-861
 3-Hydroxyanisotone, A-1045
 N⁹-Hydroxyaninomontine, A-1051
 4-Hydroxyanonaine, S-549
 Hydroxyanopteryne, A-1071
 7-Hydroxyanopteryl 11-benzoate 12-tiglate, A-1071
 7-Hydroxyanopteryl 11-4'-hydroxybenzoate 12-tiglate, A-1071
 15-Hydroxyansamitocin P1, M-136
 15-Hydroxyansamitocin P2, M-136
 15-Hydroxyansamitocin P3, M-136
 15-Hydroxyansamitocin P4, M-136
 15-Hydroxyansamitocin P₆, M-136
 30-Hydroxyansamitocin P1, M-136
 30-Hydroxyansamitocin P2, M-136
 30-Hydroxyansamitocin P3, M-136
 30-Hydroxyansamitocin P4, M-136
 4-Hydroxyanthranilic acid, A-788
 5-Hydroxyanthranilic acid, A-789
 3-Hydroxyanthranilic acid, A-787
 10-Hydroxyantirrhine, A-1328
 14-Hydroxyantifone β -N-oxide, A-1330
 14-Hydroxyantofine, A-1330
 5'-Hydroxyapateline, A-1331
 N⁶-Hydroxyaphelandrine, A-1334
 13 α -Hydroxyaphyllidine, A-1336
 2*R*-Hydroxyaphyllidine, A-1336
 2*S*-Hydroxyaphyllidine, A-1336
 2 α -Hydroxyaphylline, A-1337
 Hydroxyaphylline, H-405
 13-Hydroxyaphylline, V-153
N-Hydroxyapiosporamide, A-1340
 10-Hydroxyapparicine, A-1365
 5-Hydroxyarborinine, T-241
N-Hydroxyarcyroxocin B, A-1398
N-Hydroxyaristolactam BII, D-567
N-Hydroxyaristolactam II, D-567
 7-Hydroxyaristolactam I, T-251
 9-Hydroxyaristolactam I, T-251
 7-Hydroxyaristolochic acid A, T-267
 17-Hydroxyaristoloteline, A-1442
 11-Hydroxyascididemin, A-1459
 6-Hydroxyasimilobine, P-4
 Hydroxyaspergillie acid, A-1475
 16-Hydroxyaspidofractinine, H-406
 12-Hydroxy- Δ^1 -aspidospermidine, E-3
 6-Hydroxyatropine, H-521
 1-(4-Hydroxy-9-azabicyclo[4.2.1]non-2-en-2-yl)-1-propanone, H-325
 5-Hydroxy-1-azabicyclo[3.3.0]octan-2-one, H-234
 5-Hydroxy-1-azabicyclo[3.3.0]oct-3-en-2-one, T-189
 3-Hydroxybaikiain, T-187
 10-Hydroxybarterine, S-595
 4-Hydroxybatrachotoxin, B-36
 (2-Hydroxybenzamido)benzoic acid, B-67
 4-Hydroxybenzeneacetamide, H-672
 α -Hydroxybenzeneacetamide, H-671
 4-Hydroxybenzeneacetic acid, H-672
 α -Hydroxybenzeneacetonitrile, H-673
 4-Hydroxybenzeneethanol, H-676
 4-Hydroxybenzenemethanol, H-414
 α -Hydroxybenzenepropanoic acid 8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester, L-216
 4-Hydroxybenzenepropanoic acid, H-689
 6-Hydroxy-1,2-benzisoxazol-3(2*H*)-one, B-72
 10-Hydroxybenzo[*f*]-1,3-benzodioxolo[6,5,4-*cd*]indol-5(6*H*)-one, A-1429
 11-Hydroxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-5,6(7*H*)-dione, T-661
N-[6-(6-Hydroxy-1,3-benzodioxol-5-yl)-naphtho[2,3-*d*]-1,3-dioxol-5-yl]-*N*-methylformamide, I-159
 4-Hydroxybenzo[*c*][2,7]naphthyridine, B-75
 11-Hydroxy-2*H*-benzo[*c*]pyrrolo[2,3-*h*]cinnolin-2-one, N-84
 9-Hydroxy-8*H*-benzo[*b*]pyrido[4,3,2-*de*][1,10]phenanthroline-8-one, H-542
 7-Hydroxy-2*H*-1,4-benzothiazin-3(4*H*)-one, H-407
 6-Hydroxy-5-benzothiazoleacetic acid, H-408
 2-Hydroxybenzothiazole, B-80
 6-Hydroxy-2(3*H*)-benzothiazolone, B-78
 2-(6-Hydroxy-2-benzothiazolyl)-4(5*H*)-thiazolone, F-56
 2-Hydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, H-409
 2-Hydroxybenzoxazole, B-83
 4-Hydroxy-2-benzoxazolone, B-81
 6-Hydroxy-2-benzoxazolone, B-82
 4-Hydroxy-2(3*H*)-benzoxazolone, B-81
 6-Hydroxy-2(3*H*)-benzoxazolone, B-82
 6-[[4-(4-Hydroxybenzoyl)amino]butyl]amino]carbonyl]-1,3-dimethyl-2,4(1*H*,3*H*)-pteridinedione, H-410
N-[4-(4-Hydroxybenzoylamino)butyl]-1,3-dimethylumazine-6-carboxamide, H-410
N-[5-(4-Hydroxybenzoylamino)pentyl]-1,3-dimethylumazine-6-carboxamide, H-410
N-(2-Hydroxybenzoyl)anthranilic acid, B-67
 N²-(4-Hydroxybenzoyl)arginine, A-1413
 1-(4-Hydroxybenzoyl)- β -carboline, H-411
 (4-Hydroxybenzoyl)choline, C-419
 1-(4-Hydroxybenzoyl)-6,7-dimethoxy-2-methylisoquinolinium, G-14
 4-Hydroxybenzoylformic acid, H-687
N-(4-Hydroxybenzoyl)glutamic acid, G-107
 2-(4-Hydroxybenzoyl)-4(5)-(4-hydroxyphenyl)-1*H*-imidazole, H-412
 2-(4-Hydroxybenzoyl)-6-(4-hydroxyphenyl)pyrazine, B-259
 6-(4-Hydroxybenzoyl)-5-(4-hydroxyphenyl)-1*H*-pyrrolo[2,3-*d*]pyrimidine-2,4(3*H*,7*H*)-dione, R-98
N-(2-Hydroxybenzoyl)-4-methoxyanthranilic acid, B-67
N-(2-Hydroxybenzoyl)-*O*-methyltyramine, M-280
 3 α -(4-Hydroxybenzoyloxy)tropane, M-393
 β -(4-Hydroxybenzoyloxy)tropane, M-393
 3-(*m*-Hydroxybenzoyloxy)tropane, M-393
 3-Hydroxy-6-benzoyloxytropane, M-391
Agelenopsis aperta Hydroxybenzoylpentamine toxins, H-413
 1-(4-Hydroxybenzoyl)-9*H*-pyrido[3,4-*b*]indole, H-411
m-Hydroxybenzoyltropine, M-393
N-(2-Hydroxybenzoyl)tyramine, H-667
 4-Hydroxybenzyl alcohol, H-414
 4-Hydroxybenzyl cyanide, H-672
 6*N*-(3-Hydroxybenzyl)adenine, A-137
 4-Hydroxybenzylamine, H-417
 2-Hydroxybenzylamine, H-415
 3-Hydroxybenzylamine, H-416
 (4-Hydroxybenzyl)aminofornic acid, H-421
 (4-Hydroxybenzyl)aminothioformic acid, H-427
 8-(4-Hydroxybenzyl)-2,3,10,11-berbintetrol, H-419
 8-(4-Hydroxybenzyl)-2,3,9,10-berbintetrol, H-418
 8-(4-Hydroxybenzyl)-2,3,10-berbintetrol, H-420
 (4-Hydroxybenzyl)carbamic acid, H-421
 8-(4-Hydroxybenzyl)-2,3-dimethoxy-10-berbinol, H-420
 1-(4-Hydroxybenzyl)-6,7-dimethoxyisoquinoline, H-424
 5-(*p*-Hydroxybenzyl)hydantoin, H-423
 2-(4-Hydroxybenzyl)imidazole, H-422
 5-(4-Hydroxybenzyl)-2,4-imidazolidinedione, H-423
 1-(4-Hydroxybenzyl)-6,7-isoquinolinediol, H-424
 8-(4-Hydroxybenzyl)-2-methoxy-3,10,11-berbintetrol, H-419
 4-(*p*-Hydroxybenzyl)-6-methoxy- β -carboline, C-230
 1-(4-Hydroxybenzyl)-6,7-methylenedioxyisoquinoline, H-424
 2-(4-Hydroxybenzyl)-4-oxo-2-azetidinedicarboxylic acid, H-425
 2-(4-Hydroxybenzyl)-*N*-phenylmaleimide, O-100
 5-(4-Hydroxybenzyl)-2-pyridinedicarboxylic acid, H-426
 5-(α -Hydroxybenzyl)-2-pyridinedicarboxylic acid, V-94
 (4-Hydroxybenzyl)thiocarbamic acid, H-427
 2-(α -Hydroxybenzyl)-3,6-tropanediol, H-428
 2-(α -Hydroxybenzyl)-3-tropanol, H-429
 7-Hydroxybiopterin, I-13
 6-Hydroxy-3,3-bis(3,7-dimethyl-2,6-octadienyl)-2,4(1*H*,3*H*)-quinolinedione, S-276
 3-Hydroxy-4,5-bis(hydroxymethyl)-2-methylpyridine, P-923
 11-Hydroxy-1,2,9,10-bis(methylenedioxy)noraporphine, P-211
 12'-Hydroxy-2',5'-bis(1-methylethyl)ergotaman-3',6',18-trione, E-153
 3-Hydroxy-2,5-bis(1-methylpropyl)pyrazine, B-204
 1-Hydroxy-3,6-bis(1-methylpropyl)-2(1*H*)-pyrazinone, B-204
 1-Hydroxy-3,6-bis[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-430
 4-Hydroxy-2,6-bis(methylthio)phenyl-1-azoformamide, D-229
 2-[4-Hydroxy-2,6-bis(methylthio)phenyl]diazene-carboxamide, D-229
 6-Hydroxyblepharin, H-409
 6'-Hydroxybudmunchiamine C, B-387
 6'-Hydroxybudmunchiamine K, B-389
 4-Hydroxybulbocapnine, B-397
 6 α -Hydroxybuphanidrine, P-581
 6-Hydroxy-(+)-buphanisine, C-748
 6-Hydroxy-(-)-buphanisine, C-748
 Hydroxybutanedioic acid, M-72
 11-*O*-(3-Hydroxybutanoyl)hamayne, H-5
 1-(3-Hydroxybutanoyl)pancratistatine, P-58
 3-*O*-(3-Hydroxybutanoyl)tazettinol, T-53
 4-Hydroxy-2-butenic acid, H-431
 5-(3-Hydroxybutyl)-7-epiaustraline, T-553
 (4-Hydroxybutyl)guanidine, H-432
 5-(3-Hydroxybutyl)hyacinthacine A₁, T-553
 5-(3-Hydroxybutyl)hyacinthacine A₂, T-553
 2-(2-Hydroxybutyl)-6-(2-hydroxypentyl)-1-methylpiperidine, H-433
 2-(2-Hydroxybutyl)-6-(2-hydroxypentyl)piperidine, H-433
 6-(2-Hydroxybutyl)-2-(2-hydroxypropyl)-1-methyl- Δ^3 -piperidine, H-434
 2-(2-Hydroxybutyl)-6-(2-hydroxypropyl)-1-methylpiperidine, H-434
 6-(2-Hydroxybutyl)-1-methyl-2-(2-oxobutyl)- Δ^3 -piperidine, D-391
 2-(2-Hydroxybutyl)-1-methyl-6-(2-oxobutyl)piperidine, D-391
 2-(2-Hydroxybutyl)-*N*-methylpiperidine, H-435
N-(4-Hydroxybutyl)-3-(methylthio)-2-propenamide, A-716
 2-(2-Hydroxybutyl)-6-(2-oxobutyl)piperidine, D-391
 5-(3-Hydroxybutyl)picolinic acid, H-436
 2-(2-Hydroxybutyl)piperidine, H-435
 5-(3-Hydroxybutyl)-2-pyridinedicarboxylic acid, H-436
 2-(4-Hydroxybutyl)-4-quinazolinol, H-437
 2-(4-Hydroxybutyl)-4(3*H*)-quinazolinone, H-437
 30-Hydroxybuxamine A, B-466
 16-Hydroxybuxaminone, B-457
 18-Hydroxycabucraline, C-191
 12-Hydroxycamoensidine, C-61
 12 α -Hydroxycamoensine, C-62
 10-Hydroxycamptothecin, C-70
 11-Hydroxycamptothecin, C-70

- 18-Hydroxycamptothecin, C-70
 13-Hydroxycanadine, O-94
 8-Hydroxycantharimide, C-89
 9-Hydroxycanthin-6-one *N*³-oxide, H-444
 11-Hydroxycanthin-6-one *N*-oxide, H-446
 4-Hydroxycanthin-6-one, H-440
 5-Hydroxycanthin-4-one, H-441
 5-Hydroxycanthin-6-one, H-442
 8-Hydroxycanthin-6-one, H-443
 9-Hydroxycanthin-6-one, H-444
 1-Hydroxycanthin-6-one, H-438
 10-Hydroxycanthin-6-one, H-445
 11-Hydroxycanthin-6-one, H-446
 2-Hydroxycanthin-6-one, H-439
 8-Hydroxycantleyine *cis-p*-coumarate, C-92
 8-Hydroxycantleyine *trans-p*-coumarate, C-92
 4'-Hydroxycapitavine, C-94
 ω-Hydroxycapsaicin, C-100
 7-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-450
 6-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-449
 8-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-451
 1-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-447
 2-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-448
 8-Hydroxy-2-carbazolecarboxylic acid (obsol.), H-454
 7-Hydroxy-9*H*-carbazole-3-carboxylic acid, H-455
 1-Hydroxy-9*H*-carbazole-3-carboxylic acid, H-452
 2-Hydroxy-9*H*-carbazole-3-carboxylic acid, H-453
 6-Hydroxy-9*H*-carbazole-3-carboxylic acid, H-454
 4-Hydroxy-β-carboline-1-carboxaldehyde, H-456
 4-Hydroxy-β-carboline-1-carboxylic acid, H-457
 6-Hydroxy-β-carboline-1-carboxylic acid, H-458
 7-Hydroxy-β-carboline-1-carboxylic acid, H-459
 6-Hydroxy-β-carboline, P-915
 7-Hydroxy-β-carboline, P-916
 8-Hydroxy-β-carboline, P-917
 1-Hydroxy-β-carboline, P-914
 7-Hydroxy-β-carboline-1-propanoic acid, C-134
 2-Hydroxy-3-carbomethoxycarbazole, H-453
 4-Hydroxycarbostyryl, Q-33
 16-Hydroxycardiopetaline, M-355
 15-Hydroxycaribaeorane, C-160
 19-Hydroxycassaine, C-178
 6-Hydroxycassamine, N-300
 14-Hydroxycathovaline, C-213
 7β-Hydroxycatuabine D, M-392
 7β-Hydroxycatuabine E, M-392
 7β-Hydroxycatuabine F, M-392
 7α-Hydroxycatuabine H, M-392
 7β-Hydroxycatuabine H, M-392
 7β-Hydroxycatuabine I, M-392
 3-Hydroxycelacinnine, P-356
 11β-Hydroxycephalotaxine β-*N*-oxide, C-271
 4-Hydroxycephalotaxine, C-271
 11-Hydroxycephalotaxine, C-271
 3-Hydroxycephalotaxine, C-271
 7-Hydroxyceratinamine, M-679
 20-Hydroxy-3,6-cevanedione, C-298
 6-Hydroxy-3-cevanone, C-290
 3-Hydroxy-6-cevanone, C-290
 7-Hydroxyceve-4-ene-3,18-dione, H-460
 3-Hydroxyceve-13(17)-en-6-one, C-308
 3-Hydroxyceve-8-en-6-one, C-302
 20-Hydroxyceve-8-en-3-one, H-461
 3-Hydroxycevomycin, C-26
 12-Hydroxychasmaconitine, M-342
 12-Hydroxychelerythrine, C-356
 10-Hydroxychelidone, C-353
 5-Hydroxychelirubine, C-360
 12-Hydroxychelirubine, C-360
 3-Hydroxychlorolisoclimide, H-74
 8-Hydroxycinchonaldehyde, H-722
 6-Hydroxycinchoninic acid, H-726
 2-Hydroxycinchoninic acid, H-723
p-Hydroxycinnamic acid, H-691
m-Hydroxycinnamic acid, H-690
N-p-Hydroxycinnamoylanthranilic acid, A-1563
 (4-Hydroxycinnamoyl)lupinine, O-58
 4-Hydroxycinnamoylputrescine, C-456
N'-(4-Hydroxycinnamoyl)-*N,N'*-trimethylputrescine, C-456
 14-Hydroxycodeine, H-462
 14-Hydroxycodeinone, H-462
 6-Hydroxycolchicine, C-568
 3-Hydroxy-α-colubrine, P-726
 3-Hydroxy-β-colubrine, P-726
 10-Hydroxy-α-colubrine, S-589
 2-Hydroxycomosivine, C-588
 11-Hydroxyconacona-1,4-dienine-3,18-dione, H-314
 11α-Hydroxyconacona-1,4-dienin-3-one, H-314
 14-Hydroxycondylocarpine, T-673
 12-Hydroxyconessimine, H-318
 7α-Hydroxyconessine, C-601
 12-Hydroxyconessine, H-318
 19'*R*-Hydroxyconoduramine, C-615
 19'*S*-Hydroxyconoduramine, C-615
 3'-Hydroxyconoduramine, C-615
 19'*R*-Hydroxyconodurine, C-616
 19'*S*-Hydroxyconodurine, C-616
 3'-Hydroxyconodurine, C-616
 3-Hydroxyconopharyngine hydroxyindolenine, C-658
 19-Hydroxyconopharyngine, C-620
 3-Hydroxyconopharyngine, C-620
 (±)-19-Hydroxycoronaridine, C-657
 3-Hydroxycoronaridine, C-657
 11-Hydroxycoronaridine, I-344
 10-Hydroxycoronaridine, V-170
 10-Hydroxycorreantoxide, C-662
 9-Hydroxycorynantheine, G-13
 5-Hydroxycorynoline, C-692
 (±)-5-Hydroxycorynoline, C-692
 12*R*-Hydroxycorynoloxine, C-693
 7-Hydroxycossinidine, H-197
*N*²-*p*-Hydroxycoumaroyltryptophan, T-640
 16-Hydroxycrambesidin 359, C-715
 4-Hydroxycrebanine, C-733
 8-Hydroxycrenatine, D-686
 6α-Hydroxycrinamide, C-743
 6-Hydroxycrinamide, C-743
 6-Hydroxycrinamine, H-6
 6-Hydroxycrinine, C-748
 6α-Hydroxycroamine, S-523
 4-Hydroxycrotonic acid, H-431
 12-Hydroxycrychine, E-221
 1-Hydroxycryptochine, C-781
 Hydroxycryptolepine, D-505
 15*R*-Hydroxycryptoleurine, C-792
 16-Hydroxycyclobuxiridine L, C-884
 11-Hydroxy-4,17-cyclocorynanium(1+), O-96
 5-Hydroxycycloheximide, D-765
 21-Hydroxycycloochermerine, T-21
 30-Hydroxycyclocroboxene, A-798
 20-Hydroxycyclindrocaine, C-929
 12-Hydroxycytisine, C-938
 Hydroxydanaidal, D-14
 Hydroxydaphgraciline, D-50
 11-Hydroxydaphnigracine, D-51
 10-Hydroxydarlingine, D-75
 7-Hydroxydavisine, H-197
 2-Hydroxy-2'-deacetoxyaustrospicatine, T-46
 10-Hydroxydeacetylakuummiline, A-231
 3'-Hydroxy-16'-decarbomethoxyconodurine, C-616
 16-Hydroxy-16-decarbomethoxytetrahydrosecamine, S-174
 6-(5-Hydroxy-1,3,7-decatrienyl)-1,2-dimethyl-3-piperidinol, C-789
 20-Hydroxy-1,2-dehydro-Ψ-aspidospermidine, D-377
 8-Hydroxydehydrocrenatine, D-686
 6-Hydroxydehydrocrenatine, M-290
 9-Hydroxy-2,3-dehydrodeoxypeganine, D-503
 7-Hydroxydehydroglucine, D-159
 13-Hydroxy-7,11-dehydromatrine, L-94
 10-Hydroxy-18,19-dehydronigraninone, O-21
 3-Hydroxy-6α,7-dehydronuciferine, T-521
 23-Hydroxy-2,16-dehydrotretuline, N-301
 6-Hydroxydehydrodroskyanthine, I-62
 7-Hydroxydehydrothalicsimidine, P-208
 17-Hydroxydehydrovoachalotine, D-173
 12-Hydroxydelcosine, M-340
 10-Hydroxydelsoline, M-339
 12-Hydroxy-11-demethoxy-3-epimyrtoidine, M-803
 17-Hydroxydemethoxyisomigrastatin, I-245
 16-Hydroxy-*N*⁶-demethylalstophylline oxindole, A-654
 1'-Hydroxy-9-*O*-demethylcephaline, E-72
 10-Hydroxy-*N*-demethyl-19,20-dehydroraumacline, R-22
 3'-Hydroxy-*N*⁴-demethylervahanine A, E-189
 3'-Hydroxy-*N*⁴-demethylervahanine B, C-615
 16-Hydroxy-9-*O*-demethylgalwesine, H-504
 5-Hydroxy-10-*O*-demethylhomolycorine, H-504
 17-Hydroxy-8-*O*-demethylisomigrastatin, I-245
 3-Hydroxy-6'-demethyl-9-*O*-methylthalifaboramine, T-322
 Hydroxy-de-*N*-methyl-α-obscurene, O-5
 4α-Hydroxy-*N*³-demethylplakinamine B, P-497
 11-Hydroxy-4'-*N*-demethylstaurosporine, S-498
 3-Hydroxy-4'-*N*-demethylstaurosporine, S-498
 3-Hydroxy-*O*-demethylstaurosporine, S-498
 3'-Hydroxy-*N*⁴-demethyltabernamine, T-8
 3-Hydroxy-6'-*O*-demethylthalifaboramine, T-322
 2-Hydroxydendrobine, D-210
 4-Hydroxydendroxine, D-216
 6-Hydroxydendroxine, D-216
 10-Hydroxy-3-deoxyaconitine, M-332
 2-Hydroxy-3-deoxyaconitine, M-330
 10-Hydroxydeoxycamptothecin, C-70
 11β-Hydroxydeoxyharringtonine, H-66
 10-Hydroxy-13-deoxypaxilline, P-140
 14α-Hydroxy-4b-deoxypaxilline, P-140
 3-Hydroxy-4-deoxypaxilline, P-140
 9-Hydroxydeoxypeganine, D-503
 1-Hydroxydeoxypeganine, T-231
 5-Hydroxydeoxyvasicinone, D-503
 Hydroxydesaminohistidine, H-523
 3'*S*-Hydroxy-5'-des-*O*-methylharringtonine, H-66
 1''*S*-Hydroxy-3''-des-*O*-methylharringtonine, H-66
 3-Hydroxydiaboline, C-115
 4-Hydroxydianthramide B methyl ester, B-67
 4-Hydroxydianthramide B, B-67
 4-Hydroxydianthramide R, B-67
 Hydroxydianthramide S methyl ester, B-67
 4-Hydroxydianthramide S, B-67
 4-Hydroxydicentrine, D-346
 7-Hydroxydicentrine, D-948
 7-Hydroxydictamine, F-212
 8-Hydroxydictamine, F-213
 27-Hydroxy-13,19-didemethylspirolide C, S-432
 1-Hydroxy-5,11-dideoxytetradotoxin, T-510
N-Hydroxydihydroabakoviromycin, A-14
 Hydroxy-19,20-dihydroakuummicine, A-228
 18(19)-Hydroxy-19,20-dihydroakuummicine, A-228
 19ε-Hydroxy-19,20*R*-dihydroakuummicine, E-30
 15'-Hydroxy-14',15'-dihydroborreverine, B-253
 8-Hydroxydihydrochelerythrine, D-407
 5-Hydroxydihydrochelirubine, D-510
 10-Hydroxydihydrocorynantheine, C-689
 10-Hydroxydihydrocorynantheol, D-412
 19*S*-Hydroxydihydrocorynantheol, D-412
 10-Hydroxy-3,14-dihydrodeucussine, D-138
 1-Hydroxy-1,2-dihydro-*N*-desmethylacronycine, D-417
 15-Hydroxy-14,15-dihydro-16-epivindolinine, V-132
 4'-Hydroxy-3',4'-dihydroflindersine, F-90
 19-Hydroxydihydrogelsemine, G-49
 19*R*-Hydroxydihydrogelsevirine, G-49
 19*S*-Hydroxydihydrogelsevirine, G-49
 15'-Hydroxy-14',15'-dihydroisoborreverine, I-198
 19-Hydroxy-16,17-dihydroayumbine, A-953
 17-Hydroxydihydroymyrinal, E-136
 7'-Hydroxy-7',8'-dihydropleurostyline, P-523
 20-Hydroxydihydrorankinidine, H-367
 8-Hydroxydihydroarangianine, D-510
 12-Hydroxydihydrosanguinarine, D-510
 10'-Hydroxy-4',17-dihydro-17α-tchibangensine, U-52
 10'-Hydroxy-4',17-dihydro-17β-tchibangensine, U-52
 10-Hydroxy-18,19-dihydrousambarine, O-21
 11-Hydroxy-18,19-dihydrousambarine, O-21

- Hydroxydihydrovertine, V-95
 15 α -Hydroxy-14,15-dihydrovindolinine, V-132
 2-Hydroxy-3-(3,4-dihydroxyphenyl)propanamide, D-648
 1-Hydroxy-2,3-dimethoxyacridone, T-516
 1-Hydroxy-3,5-dimethoxyacridone, T-518
 7-Hydroxy-1,2-dimethoxyaporphine, P-4
 1-Hydroxy-2,10-dimethoxyaporphine, T-524
 1-Hydroxy-2,11-dimethoxyaporphine, T-525
 1-Hydroxy-2,3-dimethoxyaporphine, T-521
 1-Hydroxy-2,9-dimethoxyaporphine, T-523
 10-Hydroxy-1,2-dimethoxyaporphine, T-524
 11-Hydroxy-1,2-dimethoxyaporphine, T-525
 2-Hydroxy-1,10-dimethoxyaporphine, T-524
 2-Hydroxy-1,3-dimethoxyaporphine, T-521
 3-Hydroxy-1,2-dimethoxyaporphine, T-521
 5-Hydroxy-4,7-dimethoxybenz[*g*]isoquinolin-1(2*H*)-one, E-83
 4-Hydroxy-2,3-dimethoxybenzo[*g*]quinoline-5,10-dione, S-161
 2-Hydroxy-4,7-dimethoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, D-538
 2-Hydroxy-6,7-dimethoxybenzoxazole, D-708
 8-Hydroxy-6,9-dimethoxy-12*H*-[1]-benzoxepino[2,3,4-*i*]isoquinolin-12-one, S-73
 1-(2-Hydroxy-3,4-dimethoxybenzoyl)-6,7-dimethoxyisoquinoline, T-50
O-(4-Hydroxy-3,5-dimethoxybenzoyl)vincamajine, V-113
 3-(4-Hydroxy-3,5-dimethoxybenzylidene)-2-indolinone, H-463
 1-(2-Hydroxy-3,4-dimethoxybenzyl)-6,7-methylenedioxyisoquinoline, B-103
 9-(2-Hydroxy-4,5-dimethoxybenzyloxy)-1,2,3,10-tetramethoxyoxoaporphine, O-203
 10-Hydroxy-1,11-dimethoxycanthin-6-one, T-530
 2-Hydroxy-6,8-dimethoxy-9*H*-carbazole-3-carboxaldehyde, T-533
N-(4-Hydroxy-3,5-dimethoxycinnamoyl)- Δ^2 -piperidin-4-one, C-245
 5-Hydroxy-6,7-dimethoxycleistopholine, M-399
 6-Hydroxy-7,8-dimethoxycleistopholine, M-399
 1-Hydroxy-4,4-dimethoxy-2,5-cyclohexadiene-1-acetamide, H-650
 1-Hydroxy-2,3-dimethoxydibenz[*cd,f*]indol-4(5*H*)-one, T-537
 2-Hydroxy-1,3-dimethoxy-7*H*-dibenzo[*de,g*]quinolin-7-one, M-712
 9-Hydroxy-1,2-dimethoxy-7*H*-dibenzo[*de,g*]quinolin-7-one, P-275
 8-Hydroxy-6,7-dimethoxy-3,4-dihydroisoquinolinium inner salt, D-437
 4-Hydroxy-3,6-dimethoxy-1,2-dimethylcarbazole, T-543
 3-Hydroxy-4,5-dimethoxydimethylphenethylamine, H-465
 12*b*-Hydroxy-9,10-dimethoxy-6*H*-1,3-dioxolo[4,5-*h*]isoindolo[1,2-*b*][3]benzazepine-8,13(5*H*,12*bH*)-dione, C-375
 8-Hydroxy-1,3-dimethoxy-2,7-diprenylacridone, T-548
 1-Hydroxy-3,5-dimethoxy-2,4-diprenylacridone, T-546
 5-Hydroxy-4,8-dimethoxyfuro[2,3-*b*]quinoline, F-220
 6-Hydroxy-4,7-dimethoxyfuro[2,3-*b*]quinoline, F-221
 7-Hydroxy-4,6-dimethoxyfuro[2,3-*b*]quinoline, F-221
 7-Hydroxy-4,8-dimethoxyfuro[2,3-*b*]quinoline, F-223
 8-Hydroxy-4,7-dimethoxyfuro[2,3-*b*]quinoline, F-223
 9-Hydroxy-5,6-dimethoxyindeno[1,2,3-*ij*]isoquinoline, T-498
 1-Hydroxy-6,7-dimethoxyisoquinoline, I-305
 4-Hydroxy-5,8-dimethoxy-2-(3-methoxyphenyl)-3-propylquinoline, D-711
 4-Hydroxy-5,6-dimethoxy-2-(3-methoxyphenyl)-quinoline, D-712
 8-Hydroxy-5,7-dimethoxy-4-(4-methoxyphenyl)-2(1*H*)-quinolinone, D-713
 1-Hydroxy-3,4-dimethoxy-10-methylacridone, T-517
 1-Hydroxy-3,5-dimethoxy-10-methylacridone, T-518
 5-Hydroxy-2,6-dimethoxy-1-methyl-4-aza-9-fluorenone, T-562
 5-Hydroxy-6,7-dimethoxy-1-methyl-4-azafluoren-9-one, T-565
 6-Hydroxy-2,7-dimethoxy-1-methyl-4-azafluoren-9-one, T-561
 6-Hydroxy-5,8-dimethoxy-1-methyl-4-azafluoren-9-one, T-564
 7-Hydroxy-2,8-dimethoxy-1-methyl-4-aza-9-fluorenone, T-560
 7-Hydroxy-5,6-dimethoxy-1-methyl-4-azafluoren-9-one, T-565
 6-Hydroxy-7,8-dimethoxy-4-methylbenzo[*g*]quinoline-5,10-dione, M-399
 7-Hydroxy-3,4-dimethoxy-5-methylbenzo[*h*]quinolin-2(1*H*)-one, A-1463
 4-Hydroxy-3,6-dimethoxy-2-methyl-9*H*-carbazole-1-carboxaldehyde, H-561
 1-Hydroxy-6,7-dimethoxy-3-methylcarbazole, T-559
 1-Hydroxy-2,11-dimethoxy-6-methyl-7*H*-dibenzo[*de,g*]quinolinium hydroxide inner salt, A-583
 5-Hydroxy-1,2-dimethoxy-7-methyl-4*H*-dibenzo[*de,g*]quinolin-4-one, G-173
 8-Hydroxy-9,10-dimethoxy-1,2-methylenedioxyaporphine, L-158
 9-Hydroxy-3,10-dimethoxy-1,2-methylenedioxyaporphine, O-27
 11-Hydroxy-1,10-dimethoxy-2,3-methylenedioxyaporphine, O-25
 11-Hydroxy-3,9-dimethoxy-1,2-methylenedioxy-noraporphine, D-950
 9-Hydroxy-3,10-dimethoxy-1,2-methylenedioxy-noraporphine, O-27
 1-Hydroxy-2,3-dimethoxy-9,10-methylenedioxy-noraporphine, P-369
 10-Hydroxy-3,11-dimethoxy-1,2-methylenedioxy-noraporphine, P-209
 11-Hydroxy-3,10-dimethoxy-1,2-methylenedioxy-noraporphine, P-209
 2-Hydroxy-1,3-dimethoxy-9,10-methylenedioxy-noraporphine, P-369
 3-Hydroxy-1,2-dimethoxy-9,10-methylenedioxy-noraporphine, P-369
 9-Hydroxy-2,8-dimethoxy-3,4-methylenedioxy-pavine, T-330
 2-Hydroxy-8,9-dimethoxy-3,4-methylenedioxy-pavine, T-330
 7-Hydroxy-3,6-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-560
 8-Hydroxy-6,9-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-564
 9-Hydroxy-3,8-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-562
 9-Hydroxy-7,8-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-565
 7-Hydroxy-8,9-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-565
 8-Hydroxy-3,7-dimethoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-561
 7-Hydroxy-6,8-dimethoxy-2-methyl-1(2*H*)-isoquinolinone, C-366
 1-Hydroxy-2,3-dimethoxy-6-methyl-7-oxo-7*H*-dibenzo[*de,g*]quinolinium inner salt, M-712
 3-Hydroxy-4,5-dimethoxymethylphenethylamine, H-465
 4-Hydroxy-3,8-dimethoxy-1-methyl-2(1*H*)-quinolinone, Q-41
 1-Hydroxy-2,10-dimethoxynoraporphine, T-524
 1-Hydroxy-2,3-dimethoxynoraporphine, T-521
 1-Hydroxy-2,9-dimethoxynoraporphine, T-523
 10-Hydroxy-1,2-dimethoxynoraporphine, T-524
 2-Hydroxy-1,3-dimethoxynoraporphine, T-521
 3-Hydroxy-1,2-dimethoxynoraporphine, T-521
 18-Hydroxy-11,17-dimethoxyohimban-16-carboxylic acid, R-50
 5-Hydroxy-6,7-dimethoxyonchicine *N*-oxide, T-565
 5-Hydroxy-6,7-dimethoxyonchicine, T-565
 6-Hydroxy-2,7-dimethoxyonchicine, T-561
 6-Hydroxy-5,8-dimethoxyonchicine, T-564
 7-Hydroxy-2,8-dimethoxyonchicine, T-560
 7-Hydroxy-5,6-dimethoxyonchicine, T-565
 3-Hydroxy-1,2-dimethoxy-5-oxonoraporphine, T-571
 2-Hydroxy-3,8-dimethoxypavine, H-464
 3-Hydroxy-4,5-dimethoxyphenethylamine, H-465
 4-Hydroxy-3,5-dimethoxyphenethylamine, T-576
 4-[(2-Hydroxy-4,5-dimethoxyphenyl)amino]-3-hydroxy-4-oxobutanoic acid, H-466
N-[2-(3-Hydroxy-4,5-dimethoxyphenyl)ethyl]acetamide, H-465
N-(2-Hydroxy-4,5-dimethoxyphenyl)- α -malamic acid, H-466
N-[6-(2-Hydroxy-3,4-dimethoxyphenyl)naphtho[2,3-*d*]-1,3-dioxol-5-yl]-*N*-methylformamide, A-1446
 2-[[3-(4-Hydroxy-3,5-dimethoxyphenyl)-1-oxo-2-propenyl]oxy]-*N,N,N*-trimethylethanaminium(1+), S-308
 2-Hydroxy-4-(3,4-dimethoxyphenyl)quinolidine, L-53
 4'-Hydroxy-2',3'-dimethoxy-2-(2-piperidyl)acetophenone, P-522
 2-Hydroxy-4,8-dimethoxy-3-prenylquinoline, T-583
 4-Hydroxy-7,8-dimethoxyquinoline, Q-48
 2-Hydroxy-3,4-dimethoxyquinoline, Q-45
 4-Hydroxy-6,8-dimethoxy-2(1*H*)-quinolinone, Q-42
 12-Hydroxy-10,11-dimethoxystrychnobrasiline, S-594
 4-Hydroxy-5,6-dimethoxy-2-(2,3,6-trimethoxyphenyl)quinoline, D-712
 18-Hydroxy-11,17-dimethoxyohimban-16-methanol, R-49
N-2-Hydroxy-2-[4-(3,3-dimethylallyloxy)phenyl]ethylcinnamide, O-63
 5-Hydroxy-3,4'-dimethyl-[1,2'-bipyrrole]-2,5'(2'*H*,5*H*)-dione, H-467
 6-(4-Hydroxy-1,3-dimethyl-2-butenylamino)purine, M-586
 11-Hydroxy-4,6-dimethyl-2,4-dodecadienoic acid, D-746
 1-Hydroxy-6-(1,1-dimethylethyl)-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, B-433
 2-(14-Hydroxy-14,15-dimethylhexadecyl)-3-methoxy-4(1*H*)-quinolinone, H-468
 2-(14-Hydroxy-14,15-dimethylhexadecyl)-4(1*H*)-quinolinone, H-468
 11-Hydroxy-4,12-dimethyl-13-(2-methyl-4-thiazolyl)-8,12-tridecadien-3-one, H-469
 9-Hydroxy-6,10-dimethyl-11-(2-methyl-4-thiazolyl)-6,10-undecadien-3-one, H-470
 6-Hydroxy-2,6-dimethyl-2,7-octadienoic acid, H-471
 2-(1-Hydroxy-3,7-dimethyl-2,6-octadienyl)-1-methyl-4(1*H*)-quinolinone, D-762
 2-(3-Hydroxy-3,7-dimethyl-1,6-octadienyl)-1-methyl-4(1*H*)-quinolinone, D-762
 4-(2-Hydroxy-5,7-dimethyl-4-oxo-6,8-decadienyl)-2,6-piperidinedione, M-561
 4-(2-Hydroxy-5,7-dimethyl-4-oxo-6,8-nonadienyl)-2,6-piperidinedione, S-563
 3-Hydroxy-2,3-dimethyl-5-oxoproline, H-472
 3-Hydroxy-2,3-dimethyl-5-oxo-2-pyrrolidinedicarboxylic acid, H-472
 4-(3-Hydroxy-2,6-dimethyl-9-oxo-1,5-undecadienyl)-2-methylthiazole, H-470
 3-(2-Hydroxy-3,5-dimethylphenacyl)glutarimide, A-129
 4-Hydroxy-*N,N*-dimethylphenethylamine, H-357
 7-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalsin-6(12),13-diene-1,21-dione, T-545
 17-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalsin-6,13,19-triene-1,21-dione, H-473
 18-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalsin-6,13,19-triene-1,21-dione, H-473

- 7-Hydroxy-16,18-dimethyl-10-phenyl[11]cytocalasa-6(12),13,19-triene-1,21-dione, T-545
4-[2-(2-Hydroxy-3,5-dimethylphenyl)-2-oxoethyl]-2,6-piperidinedione, A-129
5-Hydroxy-1,6-dimethyl-2-piperidinedodecanol, J-55
5-Hydroxy- α ,6-dimethyl-2-piperidinetridecanol, S-388
5-Hydroxy- α ,6-dimethyl-2-piperidineundecanol, C-183
6-Hydroxy-7,9-dimethyl-7*H*-purinium hydroxide inner salt, P-820
2-Hydroxy-3,5-dimethylpyrazine, H-474
3-Hydroxy-2,5-dimethylpyrazine, H-475
3-Hydroxy-1,2-dimethyl-4(1*H*)-pyridinone, D-632
8-Hydroxy-5,11-dimethylpyrido[4,3-*b*]carbazole, E-67
4-Hydroxy-*N,N*-dimethyl-3-pyrrolidinecarboxylate, H-716
4-Hydroxy-1,1-dimethylpyrrolidinium-2-carboxylate, H-476
5-Hydroxy-2,6-dimethyl-11-(1*H*-pyrrol-2-yl)-4,6,8,10-undecatetraen-3-one, W-4
4-Hydroxy-2,3-dimethylquinoline, H-477
6-Hydroxy-4,8-dimethylquinoline, H-478
6-Hydroxy-5,7-dimethylquinoline, H-479
4-Hydroxy-3,3-dimethyl-1-(4,4,4-trichloro-3-methyl-1-oxobutyl)-5-(3,3,3-trichloro-2-methylpropyl)-2-pyrrolidinone, D-966
4-Hydroxy-*N,N*-dimethyltryptamine, H-480
4-Hydroxy-2,6-dinitrophenethylamine, H-481
N-2-[(4-Hydroxy-3,5-dinitrophenyl)ethyl]acetamide, H-481
2-(4-Hydroxy-3,5-dinitrophenyl)ethylamine, H-481
12-Hydroxy-19,20-dinorcroctalan-11,15-dione, C-762
5-Hydroxydioxindole-3-acetic acid, D-423
16-Hydroxy-3,18-dioxo-1,4,14-conatrienine, H-314
3-(2-Hydroxy-1,4-dioxopentyl)indole, H-537
4-[(4-Hydroxy-3,6-dioxo-5-tridecyl-1,4-cyclohexadien-1-yl)amino]butanoic acid, C-152
6-Hydroxydiscodermindole, D-845
1-(22-Hydroxy-2,4-docosadienyl)piperidine, H-39
N-(3-Hydroxydodecanoyl)normicotine, P-957
6-(2-Hydroxydodecyl)-2-(hydroxymethyl)-3,4-piperidinediol, B-49
3-(1-Hydroxydodecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
6-(11-Hydroxydodecyl)-3-methoxy-2-methyl-4(1*H*)-pyridinone, M-184
5-Hydroxydopamine, T-576
2-Hydroxydyshomerythrine, D-963
19*R*-Hydroxyeburnamine, E-6
7-Hydroxyechinazolinone, E-29
N-(3-Hydroxyecosanoyl)anthranilic acid, A-706
14-Hydroxyelegansamine, E-63
8-Hydroxyellipticine, E-67
12-Hydroxyellipticine, E-67
9-Hydroxyellipticine, M-252
9-Hydroxyelymoelavine, M-671
10-Hydroxy-16-epiaffinine, A-166
18-Hydroxyepialloyohimbine, D-687
4-Hydroxy-7-epichuchuhuanine EV, P-247
19'*S*-Hydroxy-3-epiervafolidine, E-186
11-Hydroxyepierythratidine, E-207
3'-Hydroxyepiglucosatisin, G-102
4-Hydroxyepilupinine, H-482
12-Hydroxy-19-epimalagashanine, M-58
9-Hydroxyepimeloscine, M-193
10-Hydroxyepi-17-*O*-methylkribine, K-101
10-Hydroxyepi-21-*O*-methylkribine, K-101
3-Hydroxy-23,26-epiminocholest-23(*N*)-ene-6,22-dione, P-279
3-Hydroxy-22,26-epiminocholest-22(*N*)-en-6-one, H-483
3 α -Hydroxy-12-epinapelline, N-30
8 α -Hydroxy-7-epineoptilocaulin, P-773
2-Hydroxyepipachysamine D, D-265
11-Hydroxyepipachysamine E, D-267
19-Hydroxy-20-epipandoline, P-733
25 β -Hydroxy-20-epiverazine, V-73
4 β -Hydroxy-20-epiverazine, V-73
16 β -Hydroxy-19-epivindoline *N*-oxide, V-132
16 β -Hydroxy-19-epivindoline, V-132
26-Hydroxy-19-epizoanthamine, Z-30
8-Hydroxyergine, L-370
8-Hydroxyerginine, L-370
8-Hydroxy- α -ergocryptine, E-157
8-Hydroxyergotamine, E-172
 β -Hydroxyergothioneine, H-484
19'-Hydroxyervafolene, E-185
19'*R*-Hydroxyervafolidine, E-186
19'-Hydroxyervafoline, E-185
19*S*-Hydroxyervahanine A, E-189
11-Hydroxyerysodine, E-206
11-Hydroxyerysotine, E-207
11-Hydroxyerysotinine, E-207
11 β -Hydroxyerysotramidine, E-202
11-Hydroxyerysotrine, E-206
11-Hydroxyerysovine, E-206
11-Hydroxyerythratidine, E-207
11-Hydroxy-*epi*-erythratine, E-208
11-Hydroxyerythratine, E-208
11 α -Hydroxyerythravine, E-206
12-Hydroxyeryscholtzidine, E-220
12-Hydroxyeryscholtzine, E-221
2-Hydroxyethylamine, A-767
7-[(2-Hydroxyethylamino)-5,8-isoquinolinedione, A-831
1-(2-Hydroxyethyl)-2-(12-aminotridecyl)pyrrolidine, A-921
N-(1-Hydroxyethyl)benzanilide, H-485
N-(2-Hydroxyethyl)-3,3-bis(methylthio)propanamide, E-85
1-(1-Hydroxyethyl)- β -carboline-3-carboxylic acid, A-45
1-(1-Hydroxyethyl)- β -carboline, H-486
1-(2-Hydroxyethyl)- β -carboline, H-487
N-(2-Hydroxyethyl)cinchonamide, H-488
3-(1-Hydroxyethyl)conaridine, H-489
N-(2-Hydroxyethyl)cytisine, C-938
8-(1-Hydroxyethyl)dihydrochelerythrine, H-490
8-(2-Hydroxyethyl)dihydrosanguinarine, H-491
N-(2-Hydroxyethyl)gardenamide A, G-17
N-(2-Hydroxyethyl)hexadecanamide, P-38
2-(2-Hydroxyethyl)-5-(hydroxymethyl)-3,4-pyrrolidinediol, H-492
N-(2-Hydroxyethyl)-3-(4-hydroxyphenyl)-*N*-methyl-2-propenamide, H-488
3-(2-Hydroxyethyl)indole, I-96
N-(2-Hydroxyethyl)lactanthopyridone, L-5
N-(2-Hydroxyethyl)linoleamide, A-767
1-(2-Hydroxyethyl)-4-methoxy- β -carboline, H-493
3-(1-Hydroxyethyl)-5-methoxycarbonylpyridine, H-498
3-(2-Hydroxyethyl)-5-methoxyindole, H-505
3-(2-Hydroxyethyl)-6-(3-methyl-2-butenyl)indole, H-497
N-(2-Hydroxyethyl)-*N*-methylcinnamide, H-488
N-(2-Hydroxyethyl)-*N*-methyl-*p*-hydroxycinnamide, H-488
3-(1-Hydroxyethyl)-4-methyl-5-methoxycarbonylpyridine, H-494
2-(1-Hydroxyethyl)-2-methyl-4-(2-methylpropyl)-5(2*H*)-oxazolone, A-1203
N-(2-Hydroxyethyl)-*N*-methyl-3-phenyl-2-propenamide, H-488
5-(1-Hydroxyethyl)-4-methyl-3-pyridinecarboxylic acid, H-494
5-(2-Hydroxyethyl)-4-methylthiazole, H-495
N-(2-Hydroxyethyl)-3-(methylthio)-2-propenamide, E-84
N-(2-Hydroxyethyl)monascorubramine, M-686
N-(2-Hydroxyethyl)-9,12-octadecadienamide, A-767
2-(1-Hydroxyethyl)-3-(4-oxazolyl)-2-butenal, O-156
N-(2-Hydroxyethyl)palmitamide, P-38
6-(1-Hydroxyethyl)-1-phenazinecarboxylic acid, H-496
N-(1-Hydroxyethyl)-*N*-phenylbenzamide, H-485
N-(2-Hydroxyethyl)-3-phenyl-2-propenamide, H-488
3-(2-Hydroxyethyl)-6-prenylindole, H-497
5-(1-Hydroxyethyl)-3-pyridinecarboxylic acid, H-498
2-(1-Hydroxyethyl)-3,4-pyrrolidinediol, D-588
2-(1-Hydroxyethyl)-4(3*H*)-quinazolinone, H-499
3-(2-Hydroxyethyl)-4(3*H*)-quinazolinone, Q-14
5-(2-Hydroxyethyl)thiazole, T-371
*N*¹-(2-Hydroxyethyl)thymine, T-401
(2-Hydroxyethyl)trimethylammonium, C-419
4-(2-Hydroxyethyl)-5-vinylnicotinic lactone, G-59
Hydroxyepolauramine, E-297
2-Hydroxyeustifoline A, E-300
Hydroxyevodiamine, T-226
11*S*-Hydroxyfawcettidine, F-28
6 α -Hydroxyflabelliformine, F-73
3 α -Hydroxyforesaconitine, M-303
2-Hydroxyformanilide, A-861
13-Hydroxyfranchetine, F-147
3-Hydroxyfranchetine, F-147
Hydroxyfungerin A, F-196
Hydroxyfungerin B, F-196
8-Hydroxy-10*H*-furo[3,2-*a*]carbazole-4-carboxaldehyde, F-203
4-Hydroxyfuro[2,3-*b*]quinoline, F-224
6-Hydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-211
7-Hydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-212
8-Hydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-213
4-Hydroxygalegine, H-579
16-Hydroxygalwesine, H-504
7-Hydroxygancidin W, C-871
16-Hydroxygardneramine oxindole, G-21
Hydroxygardnutine, G-23
10-Hydroxygeissoschizol, G-33
14 α -Hydroxygelsamydine, G-41
19 α -Hydroxygelsamydine, G-41
14-Hydroxygelsedine, G-44
14-Hydroxygelsenicine, G-44
14-Hydroxygelsenicine, G-44
 β -Hydroxygerambullin, G-67
 β -Hydroxygerambullin, G-68
 β -Hydroxygerambullol, G-68
1-Hydroxy-3-geranyloxy-4-methoxy-10-methylacridone, T-517
6-Hydroxygermine, P-674
3-Hydroxyglauicine, P-208
4-Hydroxyglucobrassicin, I-122
3-Hydroxyglucosatisin, G-102
13-Hydroxyglucopericidin A, P-421
5-Hydroxygramine, D-734
7-Hydroxygramine, D-735
3'-Hydroxyguanine, G-171
8-Hydroxyguanine, A-753
7-Hydroxyguanine, G-210
12'-Hydroxygymnastatin N, T-700
22-Hydroxyhalicyclamine A, H-22
10-Hydroxyhapalindole G, H-48
6-Hydroxyharman, H-591
7-Hydroxyharman, H-592
8-Hydroxyharman, H-593
11-Hydroxyhedranthine, M-666
17-Hydroxyhedranthine, M-666
18-Hydroxyheinsiagenin A, H-90
16-Hydroxyhenningsiine, H-111
3-Hydroxyhenningsiine, H-111
7-Hydroxyheptaphylline, H-125
3-(6-Hydroxyheptyl)-5-methylpyrrolizidine, H-135
3-(6-Hydroxyheptyl)-5-propylpyrrolizidine, H-500
N-Hydroxyhermaninger, N-25
6-Hydroxyhetisan-2,13-dione, H-198
6-Hydroxyhetisan-11-one, H-164
15-Hydroxyhetisan-11-one, H-167
13-Hydroxyhetisan-11-one, S-420
14-Hydroxyhetisine, H-189
14-Hydroxyhetisine *N*-oxide, H-189
N-(3-Hydroxyhexadecanoyl)normicotine, P-957
16-Hydroxy-4-hexadecenoic acid, H-501
3-(1-Hydroxyhexadecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
3-(1-Hydroxyhexadecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
4-Hydroxy-2-hexenoic acid, H-502
3-[(4-Hydroxy-1-hexenyl)-*ONN*-azoxy]-1,2-butanediol, A-1199
4-[2-(3-Hydroxy-1-hexenyl)-4-methylphenyl]-3-butenamide, L-262
11-Hydroxyheyneanine, I-344

- 10-Hydroxyheptanamine, V-170
 β -Hydroxyhistidine, A-797
 17-Hydroxyhobartine, H-310
 15-Hydroxyholamine, A-888
 11-Hydroxyhomoaerotherionin, A-158
 12-Hydroxyhomoaerotherionin, A-158
 4-Hydroxyhomoaerotoxin A, H-325
 4-Hydroxyhomobatrachotoxin, B-36
 10-Hydroxyhomochelidonine, C-352
 17-Hydroxyhomodaphniphylic acid, H-330
 11 α -Hydroxyhomodeoxyharringtonine, H-66
 11 β -Hydroxyhomodeoxyharringtonine, H-66
 Hydroxyhomolabellin, H-503
 5-Hydroxyhomolycorine, H-504
 11-Hydroxyhumantenine, H-367
 15-Hydroxyhumantenine, H-367
 6 β -Hydroxyhuperzine A, H-375
 10-Hydroxyhuperzine A, H-375
 11-Hydroxyhuperzine B, H-376
 5-Hydroxyhydantoin, H-522
p-Hydroxyhydrocinamic acid, H-689
 5-Hydroxy-3-(hydroxyacetyl)-1*H*-indole, H-400
 4-Hydroxy-2-(6-hydroxy-2-benzothiazolyl)thiazole, F-56
 5-Hydroxy-*N*-(4-hydroxybenzoyl)anthranilic acid, A-789
 4-Hydroxy-*N*-(2-hydroxybenzoyl)anthranilic acid, B-67
 7-Hydroxy-1-(4-hydroxybenzoyl)-6-methoxyisoquinoline, G-14
 7-Hydroxy-1-(4-hydroxybenzyl)-6-methoxyisoquinoline, H-424
 4-Hydroxy-*N*-(4-hydroxycinnamoyl)anthranilic acid, A-788
 5-Hydroxy-*N*-(4-hydroxycinnamoyl)anthranilic acid, A-789
 3-Hydroxy-6-(5-hydroxy-1,3,7-decatrienyl)-1,2-dimethylpiperidine, C-789
 4-Hydroxy-*N*⁴-(2-hydroxy-4,5-dimethoxyphenyl)glutamine, J-62
 4-Hydroxy-2-(14-hydroxy-14,15-dimethylhexadecyl)quinoline, H-468
 3-Hydroxy-*N*-[10-hydroxy-5,11-dimethyl-2-(1-methylpropylidene)-3,7,12-trioxo-9-(3-pyridylmethyl)-1,4-dioxo-8-azacyclododec-6-yl]-2-pyridinecarboxamide, P-919
 8-Hydroxy-6-(6-hydroxy-2,5-dimethyl-4-octenylidene)-8-methylindolizidine, P-791
 4-Hydroxy-3-(2-hydroxy-1,2-dimethylpropyl)-1-methyl-2(1*H*)quinolinone, R-31
 3-Hydroxy-1-(22-hydroxy-2,4-docosadienyl)piperidine, H-39
 3-Hydroxy-6-(11-hydroxydodecyl)-2-hydroxymethylpiperidine, P-656
 4-Hydroxy-6-(11-hydroxydodecyl)-3-methoxy-2-methylpyridine, M-184
 3-Hydroxy-6-(11-hydroxydodecyl)-2-methylpiperidine, C-183
 3-Hydroxy-6-(10-hydroxydodecyl)-2-methylpiperidine, P-655
 4-Hydroxy-*N*-(2-hydroxyethyl)-1,2-dithiolane-3-carboxamide, C-184
 5-Hydroxy-1-(1-hydroxyethyl)-2-(2-hydroxyethyl)-3-methyl-2*H*-benz[*b*]isoindole-4,9-dione, B-113
 3-Hydroxy-*N*-(2-hydroxyethyl)-2-(hydroxymethyl)pyrrolidine, H-717
 5-Hydroxy-3-(2-hydroxyethyl)-1*H*-indole, H-505
 4-Hydroxy-2-(2-hydroxyethyl)-1-isoindolinone, D-442
 5-Hydroxy-*N*-(2-hydroxyethyl)-1,2,3-trithiane-4-carboxamide, C-185
 5-Hydroxy-4-[[5-hydroxy-5-(hydroxymethyl)-2-methoxy-3-oxo-1-cyclohexen-1-yl]amino]pentanoic acid, M-789
 5-Hydroxy-2-[5-hydroxy-2-(hydroxymethyl)-4-pyrimidinyl]-7-methyl-4-benzoxazolecarboxamide, B-262
 2-Hydroxy-3-(5-hydroxy-1*H*-indol-3-yl)propanoic acid, H-506
 7-Hydroxy-1-(4-hydroxy-3-methoxybenzyl)-6-methoxyisoquinoline, P-80
 4-Hydroxy-2-(2-hydroxy-4-methoxyphenyl)-5,8-dimethoxy-3-propylquinoline, H-567
 5-Hydroxy-4-(4-hydroxy-3-methoxyphenyl)-4-(2-imidazolyl)-1,2,3-trithiane, H-507
 4-Hydroxy- α -[(4-hydroxy-3-methoxyphenyl)methylene]-*N*-[2-(4-hydroxyphenyl)ethyl]-3,5-dimethoxybenzeneacetamide, S-488
 2-Hydroxy-3-(4-hydroxy-3-methoxyphenyl)-2-methylpropanamide, D-647
 7-Hydroxy-4-(hydroxymethyl)aconit-8(15)-en-14-one, D-611
 4-Hydroxy-7-[1-hydroxy-2-(methylamino)ethyl]-2(3*H*)-benzothiazolone, H-508
 7-Hydroxy-8-(2-hydroxy-6-methylbenzoyl)indolizidine, I-231
 2-Hydroxy-3-hydroxymethyl-3-butenenitrile, D-622
 4-Hydroxy-3-hydroxymethyl-2-butenenitrile, H-510
 2-Hydroxy-3-hydroxymethyl-3-butenic acid, D-622
 4-Hydroxy-2-(hydroxymethyl)-2-butenic acid, H-509
 4-Hydroxy-3-hydroxymethyl-2-butenic acid, H-510
 2-Hydroxy-1-(2-hydroxy-3-methyl-3-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-125
 8-Hydroxy-7-(3-hydroxy-3-methylbutyl)-4-methoxyfuro[2,3-*b*]quinoline, F-118
 6-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1*H*)-quinolinone, H-511
 7-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1*H*)-quinolinone, H-512
 7-Hydroxy-1-hydroxymethyl-1,2-didehydropyrrolizidine, T-188
 5-Hydroxy-2-(1-hydroxy-1-methylethyl)-11-methylfuro[2,3-*c*]acridin-6(11*H*)-one, F-207
 1-Hydroxy-6-(1-hydroxy-1-methylethyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, M-462
 7-Hydroxy-4-(5-hydroxymethyl-2-furanyl)-2(1*H*)-quinolinone, H-513
 6-Hydroxy-1-hydroxymethyl-7-methoxyisoquinoline, D-590
 3-Hydroxy-5-hydroxymethyl-4-methoxymethyl-2-methylpyridine, P-923
 5-Hydroxy-6-(hydroxymethyl)- α -methyl-2-piperidineundecanone, P-656
 3-Hydroxy-4-(hydroxymethyl)- α -(2-methylpropyl)-2-azetidineundecanol, P-175
 3-(2-Hydroxy-1-hydroxymethyl-2-methylpropyl)-1*H*-indole, T-24
 3-(2-Hydroxy-1-hydroxymethyl-2-methylpropyl)-6-prenylindole, M-480
 3-Hydroxy-2-hydroxymethyl-6-(7-oxododecyl)piperidine, I-286
 3-Hydroxy-2-hydroxymethyl-6-(8-oxododecyl)piperidine, I-287
 3-Hydroxy-2-hydroxymethyl-6-(10-oxododecyl)piperidine, P-657
 3-Hydroxy-2-hydroxymethyl-6-(11-oxododecyl)piperidine, P-656
 9-Hydroxy-6-hydroxymethyl-1-phenazinecarboxylic acid, H-514
 5-Hydroxy-2-(hydroxymethyl)piperidine, H-699
 12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-5-dodecanone, I-287
 12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-6-dodecanone, I-286
 12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-2-dodecanone, P-656
 12-[5-Hydroxy-6-(hydroxymethyl)-2-piperidinyl]-3-dodecanone, P-657
 4-Hydroxy-5-hydroxymethylproline, H-516
 1-Hydroxy-6-(1-hydroxy-1-methylpropyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, A-1475
 1-Hydroxy-6-(1-hydroxy-1-methylpropyl)-3-(1-methylpropyl)-2(1*H*)-pyrazinone, B-204
 5-Hydroxy-2-(hydroxymethyl)pyridine, H-515
 3-Hydroxy-20-(5-hydroxy-3-methyl-6-pyridinyl)pregnan-6-one, P-280
 4-Hydroxy-5-hydroxymethyl-2-pyrrolidinecarboxylic acid, H-516
 3-Hydroxy-2-(hydroxymethyl)-1-pyrrolidineethanol, H-717
 3-Hydroxy-2-(hydroxymethyl)pyrrolidine, H-717
 1-Hydroxy-1-hydroxymethylpyrrolizidine, H-517
 1-Hydroxy-7-hydroxymethylpyrrolizidine, H-233
 2-Hydroxy-1-hydroxymethylpyrrolizidine, H-518
 6-Hydroxy-7-(hydroxymethyl)-3-(1*H*-pyrrol-3-yl)-1*H*-2-benzopyran-1-one, I-71
 4-Hydroxy-2-(12-hydroxy-12-methyltridecyl)quinoline, H-636
 α -Hydroxy-4-(1-hydroxy-2,4-octadienylidene)-1-methyl- α -(1-methylethyl)-3,5-dioxo-2-pyrrolidinepropanoic acid, H-67
 4-Hydroxy-6-(10-hydroxy-3,7,9,12,13-pentamethyl-2,5,7,11,13-pentadecapentaenyl)-3-methoxy-5-methyl-2(1*H*)-pyridinone, A-1217
 2-Hydroxy-2-(3-hydroxyphenyl)acetic acid, H-519
 2-Hydroxy-2-(4-hydroxyphenyl)acetic acid, H-520
N-[2-Hydroxy-2-(4-hydroxyphenyl)ethyl]cinnamide, O-63
 4-Hydroxy-*N*-[2-(4-hydroxyphenyl)ethyl]-3,5-dimethoxybenzenepropanamide, D-512
N-[2-Hydroxy-2-(4-hydroxyphenyl)ethyl]-3-phenyl-2-propenamide, O-63
 4-Hydroxy-2-(2-hydroxy-2-phenylethyl)piperidine, S-210
 3-Hydroxy-1-(4-hydroxyphenyl)-4-(1*H*-indol-3-yl)-2-butanone, H-538
 4-Hydroxy-1-(3-hydroxyphenyl)-3(2*H*)-isoquinolinone, P-377
 3-Hydroxy-5-(hydroxyphenylmethyl)-1-methyl-4-phenyl-2-pyrrolidinone, C-502
 5-Hydroxy-1-(4-hydroxyphenyl)-2-pyrrolidinone, L-110
 3-Hydroxy-4-(3-hydroxyphenyl)-2(1*H*)-quinolinone, D-656
 2-[3-Hydroxy-2-[[[3-hydroxy-2-pyridinyl]carbonyl]amino]]-4-benzoxazolecarboxylic acid, A-1098
 6-Hydroxy-2-(4-hydroxy-2-thiazolyl)benzothiazole, F-56
 4-Hydroxyhygrinic acid, H-715
 6-Hydroxyhyoscyamine *N*-oxide, H-521
 6-Hydroxyhyoscyamine, H-521
 19*R*-Hydroxybogamine pseudoindoxyl, I-5
 20*R*-Hydroxybogamine pseudoindoxyl, I-5
 19*R*-Hydroxybogamine, I-4
 19*S*-Hydroxybogamine, I-4
 15-Hydroxycajine, I-12
 α -Hydroxy-1*H*-imidazole-4-propanoic acid, H-523
 5-Hydroxy-2,4-imidazolimedione, H-522
 2-Hydroxy-*N*-[2-(1*H*-imidazol-4-yl)ethyl]-4-oxodecanamide, H-307
 2-Hydroxy-3-(4-imidazolyl)propanoic acid, H-523
 2-Hydroxy-5-iminoazacyclopent-3-ene, I-51
 1,1'-[[Hydroxyimino]bis(1-amino-2-hydroxy-2,1-ethanediy)]bis[4(1*H*)-pyridinone], M-623
 4,4'-[[Hydroxyimino]bis(methylene)]bisphenol, B-181
 3-Hydroxyiminoindole, I-106
 4*a*-Hydroxyincarvilleine, I-62
 6-Hydroxy-1*H*-indole-3-acetamide, H-526
 2-Hydroxy-1*H*-indole-3-acetic acid, D-485
 5-Hydroxy-1*H*-indole-3-acetic acid, H-525
 4-Hydroxy-1*H*-indole-3-acetic acid, H-524
 6-Hydroxy-1*H*-indole-3-acetic acid, H-526
 4-Hydroxy-1*H*-indole-3-acetonitrile, H-524
 6-Hydroxyindole-3-acetylphenylalanine, H-526
 6-Hydroxyindole-3-acetylvaline, H-526
 7-Hydroxy-3-indoleacrylic acid, I-136
 1-(5-Hydroxy-1*H*-indole-3-carbonyl)-9*H*-pyrido[3,4-*b*]indol-6-ol, P-488
 4-Hydroxy-1*H*-indole-3-carboxaldehyde, H-527
 5-Hydroxy-1*H*-indole-3-carboxaldehyde, H-528
 6-Hydroxy-1*H*-indole-3-carboxaldehyde, H-529
 5-Hydroxy-1*H*-indole-3-carboxylic acid, H-530
 6-Hydroxy-1*H*-indole-3-carboxylic acid, H-531
 1-Hydroxy-1*H*-indole-3-carboxylic acid, I-90
 6-Hydroxy-1*H*-indole-2,3-dione, H-533
 4-Hydroxy-1*H*-indole-2,3-dione, H-532
 5-Hydroxy-1*H*-indole-3-ethanol, H-505
 2-Hydroxyindole, I-103
 3-Hydroxyindole, I-104
 α -Hydroxy-1*H*-indole-3-propanoic acid, H-539
 3-Hydroxy-1*H*-indole-2-thiol, H-534
 5-Hydroxyindoline, D-465

- 4-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-440
 5-Hydroxy-4*H*-indolo[3,2-*de*][1,5]naphthyridin-4-one, H-441
 5-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-442
 8-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-443
 9-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-444
 1-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-438
 10-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-445
 11-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-446
 2-Hydroxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-439
N-(6-Hydroxy-1*H*-indol-3-ylacetyl)phenylalanine, H-526
N-(6-Hydroxy-1*H*-indol-3-ylacetyl)valine, H-526
 2-Hydroxy-1-(1*H*-indol-3-yl)-9*H*-carbazole-3-carboxylic acid, P-495
 2-Hydroxy-1-(1*H*-indol-3-yl)ethanone, H-400
 5-Hydroxy-5-(1*H*-indol-3-yl)-2,4-imidazolidinedione, Z-39
 5-(5-Hydroxy-1*H*-indol-3-yl)-3-(1*H*-indol-3-yl)-2*H*-pyrrol-2-one, P-675
 3-(5-Hydroxy-3-indolyl)lactic acid, H-506
 4-Hydroxyindol-3-ylmethyl glucosinolate, I-122
 2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-hexanone, H-535
 2-Hydroxy-1-(1*H*-indol-3-yl)-5-methyl-3-hexanone, S-101
 2-(Hydroxy-1*H*-indol-3-ylmethyl)-1-methyl-5-(2-methylbutyl)-4-imidazolidinone, M-110
 2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-pentanone, H-536
 2-(6-Hydroxy-3-indolyl)-2-oxoacetic acid, H-654
N-[1-Hydroxy-2-(1*H*-indol-3-yl)-2-oxoethyl]acetamide, I-132
 4-Hydroxy-5-(3-indolyl)-5-oxo-2-pentanone, H-537
 3-[2-(5-Hydroxyindol-3-yl)-5-oxo-2-pyrrolin-4-ylidene]-2-indolinone, V-142
 2-Hydroxy-1-(1*H*-indol-3-yl)-1,4-pentanedione, H-537
 3-Hydroxy-4-(1*H*-indol-3-yl)-1-phenyl-2-butanone, H-538
 2-Hydroxy-3-(3-indolyl)propanoic acid, H-539
 3-(7-Hydroxy-1*H*-indol-3-yl)-2-propenoic acid, I-136
 2-Hydroxy-3-(1*H*-indol-3-yl)-2-propenoic acid, I-133
 4-Hydroxy-4-(1*H*-indol-3-yl)-5-thio-2-imidazolidinone, Z-39
 6-Hydroxyinfracrine, C-134
 19-Hydroxyingol, H-540
 21-Hydroxyintegerrimine, S-238
 4*R*-Hydroxyipecoside, I-167
 15β-Hydroxyirehdiamine H, D-278
 4-Hydroxyisatin, H-532
 6-Hydroxyisatin, H-533
 20-*epi*-Hydroxyisoaflavinine, H-541
 20-Hydroxyisoaflavinine, H-541
 23-Hydroxyisoanguivine, S-450
 9-Hydroxyisoascididemnin, H-542
 7-Hydroxyisoatisine, I-192
 2-Hydroxyisobutylamine, A-840
 6-Hydroxy-*N*-isobutyl-2,4-decadienamide, D-106
 4-Hydroxy-*N*-isobutyl-2-hexenamide, H-502
 1-Hydroxy-3-isobutyl-6-isopropyl-2(1*H*)-pyrazinone, M-462
 1-Hydroxy-3-isobutyl-6-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-620
 1-Hydroxy-6-isobutyl-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-621
N-(2-Hydroxyisobutyl)-2,4-tetradecadienamide, T-124
N-(2-Hydroxyisobutyl)-2,4,8,10,12-tetradecapentaenamide, T-127
 12-Hydroxy-*N*-isobutyl-2,4,8,10-tetradecatetraenamide, H-744
 11-Hydroxy-*N*-isobutyl-2,4,8-tetradecatrienamide, H-745
N-(2-Hydroxyisobutyl)-2,4,8-tetradecatrienamide, T-133
 2-Hydroxyisobutyric acid, H-617
 2-Hydroxyisobutyronitrile, H-617
 13-Hydroxyisocyclocecalbenzine, I-222
 19*R*-Hydroxyisoeburnamine, E-6
 4-Hydroxy-1-isoindolinone, D-442
 4-Hydroxyisoleucine, A-800
 3-(4-Hydroxy-*L*-isoleucine)- α -amanitin, A-673
 17-Hydroxyisomigrastatin, I-245
 15-Hydroxyisoneolaurerine, N-122
*N*⁶-(ω -Hydroxyisopentenyl)adenine, Z-11
 6-Hydroxy-*N*-isopentylendroxine, D-216
 2-Hydroxyisopropylamine, A-894
 8-Hydroxy-7-isopropyl-3-methylbenz[*cd*]indol-2(1*H*)-one, P-631
 1-Hydroxy-6-isopropyl-3-(1-methylpropyl)-2(1*H*)-pyrazinone, H-543
 1-Hydroxy-6-isopropyl-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-544
 2-Hydroxy-3-isopropylpyrazine, H-545
 2-Hydroxy-5-isopropylpyrazine, H-546
 2-Hydroxy-6-isopropylpyrazine, H-547
 12-Hydroxyisoretinal, R-63
 16-Hydroxyisoretinal, R-63
 18-Hydroxyisoretuline, R-63
 12-Hydroxyisorubijervine, I-310
 Hydroxy- γ -isosanshool, T-127
 7-Hydroxy- β -isoparsteine, H-738
 14-Hydroxyisostephodoline, T-511
 12'-Hydroxyisostyrchnobiline, S-591
 18-Hydroxyisostyrgucine, S-642
 10-Hydroxyisotalatizidine, M-326
 15-Hydroxyisotalatizidine, M-327
 2-Hydroxyisotaxodine, H-331
 5-Hydroxy-3-isothiocyanato-4-methoxyindole, H-548
 10-Hydroxyisotrilobine, I-339
 14 α -Hydroxyisotyrocrebrine *N*-oxide, I-342
 3-Hydroxyisovaleric acid, H-573
 3-(3-Hydroxyisovaleroyl)maytansinol, M-136
 3-(4-Hydroxyisovaleroyl)maytansinol, M-136
 β -Hydroxyisovalerylcassaine, C-178
 3-Hydroxyisovalocangine, I-344
 4-Hydroxyisoxazole, I-348
 5-Hydroxyisoxazole, I-349
 18-Hydroxyjaconine, J-4
 1'-Hydroxykainic acid, K-4
 10β-Hydroxykarakoline, M-349
 12β-Hydroxykarasamine 8-*O*-acetate, M-350
 12β-Hydroxykarasamine, M-350
 22-Hydroxy-10-kopsanone, D-805
 17-Hydroxykopsaporine, K-66
 11-Hydroxykopsingine, K-66
 Hydroxykopsinine II, K-78
 Hydroxykopsinine I, K-78
 17 α -Hydroxy- Δ ^{14,15}-kopsinine, V-49
 3-Hydroxykynuramine, A-697
 6-Hydroxykynurenic acid, D-675
 1-Hydroxylaminopropane, P-648
 9β-Hydroxylamprololine, L-26
 10-Hydroxylanciferine, L-29
 Hydroxylanyamide III, T-129
 Hydroxylanyamide II, O-212
 Hydroxylanyamide I, O-211
 Hydroxylappaconine, A-86
 7-Hydroxylathyril, T-263
 Hydroxylelobanidine, L-86
 2-Hydroxylenticellarine, L-92
 11-Hydroxyleontine, L-99
 11-Hydroxylettowianthine, L-145
 3-(4-Hydroxy-*D*-leucine)phallacidin, P-307
 7-(4-Hydroxy-*L*-leucine)phalloidin, P-312
 21'-Hydroxyleporine, L-162
*N*²-(17-Hydroxylinolenoyl)glutamine, G-108
*N*²-(17-Hydroxylinoleoyl)glutamine, G-108
 Hydroxylipoxamycin, L-189
 10-Hydroxylirioidenine, O-187
 11-Hydroxylirioidenine, O-204
 18-Hydroxylochnerine, S-89
 Hydroxylucilactaene, L-283
 Hydroxylunacridine, L-288
 Hydroxylunacrine, B-13
 Hydroxylunidine, P-759
 Hydroxylunidonine, L-295
 Hydroxylunine, H-549
 Hydroxyluninium(1+), I-290
 13-Hydroxylupanine *p*-hydroxyphenylacetate, H-554
 13-Hydroxylupanine phenylacetate, H-554
 4-Hydroxylupanine, H-551
 6-Hydroxylupanine, H-552
 Hydroxylupanine, H-554
 12-Hydroxylupanine, H-553
 13-Hydroxylupanine, H-554
 3-Hydroxylupanine, H-550
 12-(6-Hydroxy-5-lupanyl(methyl)cytosine, C-940
 Hydroxylyalidine, L-314
 10-Hydroxylyallosidic acid, L-316
 11-Hydroxylycodine, L-324
 6-Hydroxylycodoline, L-325
 6 α -Hydroxylycopodine, L-322
 6β-Hydroxylycopodine, L-322
 7-Hydroxylycopodine, L-341
 11 α -Hydroxylycopodine, L-341
 12-Hydroxylycopodine, L-325
 8*S*-Hydroxylycothunine, F-30
 2*R*-Hydroxylycothunine, F-30
 8-Hydroxylysergic acid amide, L-370
 Hydroxylyspyridinoline, P-907
 α -Hydroxymagnocurarine, H-555
 1*a*-Hydroxymagnocurarine, H-555
 12-Hydroxymalagashanine, M-58
 13β-Hydroxymamanine, M-79
m-Hydroxymandelic acid, H-519
p-Hydroxymandelic acid, H-520
p-Hydroxymandelonitrile glucoside, H-520
 6-Hydroxymanzamine A, M-91
 8-Hydroxymanzamine A, M-91
 8-Hydroxymanzamine B, M-92
 8-Hydroxymanzamine D, M-91
 6-Hydroxymanzamine E, M-91
 8-Hydroxymanzamine J, M-94
 17-Hydroxymappicine glucoside, M-96
 19-Hydroxymappicine, M-96
 3'-Hydroxymarikanin, A-197
 18-Hydroxymatopensine, M-120
 11-Hydroxymatridin-15-one, D-81
 11-Hydroxymatridin-15-one, L-99
 5-Hydroxymatridin-15-one, S-377
 14β-Hydroxymatrine *N*¹-oxide, M-121
 11-Hydroxymatrine, D-81
 9 α -Hydroxymatrine, M-121
 13 α -Hydroxymatrine, M-121
 14 α -Hydroxymatrine, M-121
 14β-Hydroxymatrine, M-121
 3 α -Hydroxymatrine, M-121
 5-Hydroxymatrine, S-377
 12-Hydroxymauiensine, T-288
 6-Hydroxymelatonin, D-684
 β -Hydroxy-2-mercaptohistidine trimethylbetaine, H-484
 3-Hydroxy-2-mercaptoindole, H-534
 4-Hydroxy-2-mercaptopyrimidine, D-518
 2-Hydroxy-4-mercaptopyrimidine, H-556
 β -Hydroxymescaline, H-557
 4-Hydroxy-2-methoxyacetanilide, A-705
 12-Hydroxy-11-methoxy-*N*-acetyl*N*-*C*-fluorocurarine, N-301
 12-Hydroxy-11-methoxy-*N*^{acetyl}strychnosphenidine, S-605
 1-Hydroxy-3-methoxyacridone, D-528
 19-Hydroxy-3'-methoxyaglaïne C, A-199
 16-Hydroxy-11-methoxyalstonine, A-648
 14-Hydroxy-11-methoxyalstonisine, A-654
 1-Hydroxy-2-methoxyaporphine, D-531
 2-Hydroxy-1-methoxyaporphine, D-531
 4-Hydroxy-3-methoxybenzamide, D-536
 5-Hydroxy-7-methoxy-2*H*-benz[*b*]isoindole-4,9-dione, D-534
 10-Hydroxy-12-methoxy-8*H*-benzo[*g*]benzodioxolo[6,5,4-*de*]quinolin-8-one, O-183
 10-Hydroxy-9-methoxybenzo[*g*]-1,3-benzodioxolo[5,6-*a*]quinolinium(1+), T-321

- 2-Hydroxy-7-methoxy-2H-1,4-benzoxazin-3(4H)-one, D-538
 4-Hydroxy-2-methoxy-2H-1,4-benzoxazin-3(4H)-one, H-409
 2-Hydroxy-6-methoxybenzoxazole, B-82
 2-(4-Hydroxy-3-methoxybenzoyl)imidazole, H-558
N-(2-Hydroxy-4-methoxybenzoyl)-4-methoxyanthranilic acid, B-67
N-(2-Hydroxy-6-methoxybenzoyl)-*O*-methyltyramine, M-280
 2-(4-Hydroxy-3-methoxybenzoyl)thiazole, H-569
 1-(3-Hydroxy-4-methoxybenzyl)-6,7-dimethoxyisoquinoline, P-80
N-(3-Hydroxy-4-methoxybenzylidene)-4'-hydroxyphenethylamine, C-730
 2-(4-Hydroxy-3-methoxybenzyl)imidazole, H-422
 1-(3-Hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline, I-317
 1-(α -Hydroxy-4-methoxybenzyl)-6,7-methylenedioxyisoquinoline, M-242
 5-Hydroxy-4-methoxy-2,3-bis(methylthio)phenethylamine, L-207
 10-Hydroxy-3-methoxycanthin-2,6-dione, I-105
 5-Hydroxy-4-methoxy-6-canthinone 3-*N*-oxide, D-546
 4-Hydroxy-5-methoxycanthin-6-one, D-546
 5-Hydroxy-4-methoxycanthin-6-one, D-546
 8-Hydroxy-9-methoxycanthin-6-one, D-549
 1-Hydroxy-11-methoxycanthin-6-one, D-545
 1-Hydroxy-9-methoxycanthin-6-one, D-544
 10-Hydroxy-9-methoxycanthin-6-one, D-550
 11-Hydroxy-1-methoxycanthin-6-one, D-545
 11-Hydroxy-10-methoxycanthin-6-one, D-551
 2-Hydroxy-7-methoxy-9H-carbazole-3-carboxaldehyde, D-556
 2-Hydroxy-8-methoxy-9H-carbazole-3-carboxaldehyde, D-557
 7-Hydroxy-1-methoxy-9H-carbazole-3-carboxaldehyde, D-553
 1-Hydroxy-6-methoxy-9H-carbazole-3-carboxaldehyde, D-552
 2-Hydroxy-6-methoxy-9H-carbazole-3-carboxaldehyde, D-555
 7-Hydroxy-2-methoxy-9H-carbazole-1,6-dicarboxaldehyde, M-760
 12-Hydroxy-*N*-methoxycarbonyl- $\Delta^{16,17}$ -kopsinine, K-71
 2-Hydroxy-8-methoxycepharanone A, T-254
 (4-Hydroxy-3-methoxycinnamoyl)histamine, C-455
 5-Hydroxy-6-methoxycleistopholine, M-399
 6-Hydroxy-7-methoxycleistopholine, M-399
 11-Hydroxy-12-methoxycoptisine, H-559
 8-Hydroxy-9-methoxycrinine, M-108
 12-Hydroxy-11-methoxydiaboline, C-115
 1-Hydroxy-2-methoxydibenz[*cd*,*f*]indol-4(5H)-one, D-567
 2-Hydroxy-1-methoxy-4H-dibenzo[*de*,*g*]quinoline-4,5-dione, C-273
 2-Hydroxy-1-methoxy-7H-dibenzo[*de*,*g*]quinoline-7-one, L-193
 5-Hydroxy-8-methoxydictamnine, F-220
 7-Hydroxy-6-methoxydictamnine, F-221
 7-Hydroxy-8-methoxydictamnine, F-223
 8-Hydroxy-7-methoxydictamnine, F-223
 9-Hydroxy-7-methoxy-3,8-dimethylbenz[*g*]isoquinoline-5,10-dione, D-226
 4-Hydroxy-3-methoxy-1,2-dimethylcarbazole, D-571
 8-Hydroxy-4-methoxy- α , α -dimethylfuro[2,3-*b*]quinoline-7-propanol, F-118
 4-Hydroxy-3-methoxydimethylphenethylamine, H-566
 1-(1-Hydroxy-2-methoxyethyl)- β -carboline, A-1402
 1-(1-Hydroxy-2-methoxyethyl)-4-methoxy- β -carboline, H-560
 7-Hydroxy-9-methoxyflindersine, F-90
 8-Hydroxy-6-methoxyflindersine, F-90
 7-Hydroxy-4-methoxyfuro[2,3-*b*]quinoline-8-carboxaldehyde, E-315
 8-Hydroxy-4-methoxyfuro[2,3-*b*]quinoline, F-213
 19R-Hydroxy-11-methoxygelsegine, G-47
 14-Hydroxy-11-methoxygelsegine, G-44
 12-Hydroxy-11-methoxyhenningsamine, C-115
 1-Hydroxy-11-methoxy-6H-indolo[3,2,1-*de*][1,5]-naphthiridin-6-one, D-545
 6-Hydroxy-7-methoxyisoquinoline, I-302
 6-Hydroxy-7-methoxy-1-isoquinolinemethanol, D-590
 7-Hydroxy-6-methoxy-1(2H)-isoquinolinone, I-305
 4-Hydroxy-13-methoxylupanine, D-606
 1-Hydroxy-3-methoxy-10-methylacridone, D-528
 3-Hydroxy-1-methoxy-10-methylacridone, D-528
 5-Hydroxy-6-methoxy-1-methyl-4-azafluoren-9-ol, D-626
 5-Hydroxy-6-methoxy-1-methyl-4-aza-9-fluorenone, D-626
 7-Hydroxy-2-methoxy-1-methyl-4-aza-9-fluorenone, D-623
 7-Hydroxy-8-methoxy-1-methyl-4-azafluoren-9-one, D-624
 2-Hydroxy-1-methoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridin-13(12H)-one, D-407
 6-Hydroxy-7-methoxy-4-methylbenzo[*g*]quinoline-5,10-dione, M-399
 3-Hydroxy-7-methoxy-2-methyl-4H-1,4-benzoxazine-4-carboxaldehyde, C-96
 2-Hydroxy-6-methoxy-1-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, H-125
 2-Hydroxy-7-methoxy-1-(3-methyl-2-butenyl)-9H-carbazole-3-carboxaldehyde, H-125
 1-Hydroxy-3-methoxy-2-(3-methyl-2-butenyl)-10-methylacridone, D-663
 3-(2-Hydroxy-3-methoxy-3-methylbutyl)-5-(3-methyl-1-oxo-2-butenyl)-1H-indole, B-200
 5-Hydroxy-4-methoxy-3-methylcanthin-2,6-dione, I-105
 7-Hydroxy-2-methoxy-6-methyl-9H-carbazole-1-carboxaldehyde, D-619
 4-Hydroxy-3-methoxy-2-methyl-9H-carbazole-1-carboxaldehyde, H-561
 6-Hydroxy-2-methoxy-3-methyl-9H-carbazole, D-617
 1-Hydroxy-6-methoxy-3-methyl-9H-carbazole, D-613
 2-Hydroxy-5-methoxy-3-methyl-9H-carbazole, D-616
 2-Hydroxy-6-methoxy-3-methyl-9H-carbazole, D-617
 2-Hydroxy-8-methoxy-3-methyl-9H-carbazole, D-618
 7-Hydroxy-3-methoxy-6-methyl-1H-carbazole-1,4(9H)-dione, C-505
 7-Hydroxy-1-methoxymethyl-1,2-dehydropyrrolizidine, T-188
 3-Hydroxy-6-methoxy-1-methyl-4,5-diphenyl-2-piperidinone, D-575
 11-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine, B-397
 9-Hydroxy-8-methoxy-1,2-methylenedioxyaporphine, C-733
 9-Hydroxy-10-methoxy-1,2-methylenedioxyaporphine, D-346
 10-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine, D-346
 11-Hydroxy-9-methoxy-1,2-methylenedioxyaporphine, D-846
 11-Hydroxy-3-methoxy-1,2-methylenedioxyaporphine, E-70
 1-Hydroxy-2-methoxy-9,10-methylenedioxyaporphine, N-29
 10-Hydroxy-11-methoxy-1,2-methylenedioxyaporphine, N-25
 2-Hydroxy-1-methoxy-9,10-methylenedioxyaporphine, N-29
 7-Hydroxy-8-methoxy-1,2-methylenedioxyaporphine, T-522
 7-Hydroxy-6-methoxy-1-(3,4-methylenedioxybenzyl)isoquinoline, S-274
 8-Hydroxy-4-methoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline, F-217
 2-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-264
 6-Hydroxy-7-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-265
 6-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-266
 7-Hydroxy-8-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-267
 8-Hydroxy-7-methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, T-267
 11-Hydroxy-10-methoxy-1,2-methylenedioxy-norporphine, B-397
 8-Hydroxy-9-methoxy-1,2-methylenedioxy-norporphine, C-733
 9-Hydroxy-10-methoxy-1,2-methylenedioxy-norporphine, D-346
 9-Hydroxy-11-methoxy-1,2-methylenedioxy-norporphine, D-846
 10-Hydroxy-9-methoxy-1,2-methylenedioxy-norporphine, D-346
 11-Hydroxy-9-methoxy-1,2-methylenedioxy-norporphine, D-846
 11-Hydroxy-3-methoxy-1,2-methylenedioxy-norporphine, E-70
 2-Hydroxy-1-methoxy-10,11-methylenedioxy-norporphine, L-10
 1-Hydroxy-2-methoxy-9,10-methylenedioxy-norporphine, N-29
 10-Hydroxy-11-methoxy-1,2-methylenedioxy-norporphine, N-25
 2-Hydroxy-1-methoxy-9,10-methylenedioxy-norporphine, N-29
 9-Hydroxy-11-methoxy-1,2-methylenedioxyoxaporphine, O-183
 2-Hydroxy-3-methoxy-8,9-methylenedioxy-pavine, E-220
 1-Hydroxy-2-methoxy-8,9-methylenedioxy-pavine, N-104
 6-Hydroxy-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid, M-263
 7-Hydroxy-8-methoxy-3,4-methylenedioxy-1-phenanthrenecarboxylic acid, M-263
 6-Hydroxy-2-methoxy-3-methylene-10-propenyldiene-2-azaspiro[4.5]dec-8-ene-1,4,7-trione, T-625
 7-Hydroxy-3-methoxy-4-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-623
 7-Hydroxy-6-methoxy-4-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-624
 8-Hydroxy-9-methoxy-4-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-626
 9-Hydroxy-8-methoxy-4-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-626
 7-Hydroxy-8-methoxy-1-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-625
 8-Hydroxy-7-methoxy-1-methyl-5H-indeno[1,2-*b*]pyridin-5-one, D-625
 6-Hydroxy-5-methoxy-2-methyl-1H-isoindole-1,3(2H)-dione, D-533
 7-Hydroxy-6-methoxy-1-methylisoquinoline, D-627
 7-Hydroxy-6-methoxy-2-methyl-1(2H)-isoquinolinone, I-305
 4-Hydroxy-8-methoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, T-583
 8-Hydroxy-4-methoxy-1-methyl-3-(3-methyl-2-butenyl)-2(1H)-quinolinone, T-583
 1-Hydroxy-2-methoxy-10-methyl-3,4-methylenedioxyacridone, M-169
 1-Hydroxy-3-methoxy-10-methyl-5,6-methylenedioxyacridone, T-243
 1-Hydroxy-5-methoxy-10-methyl-2,3-methylenedioxyacridone, T-56
 5-Hydroxy-4-(methoxymethyl)-6-methyl-3-pyridinemethanol, P-923
 3-Hydroxy-4-methoxy-3-methyloxindole, D-443
 4-Hydroxy-3-methoxymethylphenethylamine, H-566
 3-Hydroxy-4-methoxy-*N*-methylphenethylamine, H-565
 6-Hydroxy-5-methoxy-*N*-methylphthalimide, D-533
 6-Hydroxy-4-methoxy-5-methylphthalimidine, C-437
 1-Hydroxy-3-methoxy-10-methyl-2-prenylacridone, D-663

- 7-Hydroxy-2-methoxy-6-methyl-1-prenylcarbazole, E-275
- 2-Hydroxy-6-methoxy-3-methyl-5-prenyl-9*H*-carbazole, G-110
- 1-Hydroxy-7-methoxy-3-methyl-8-prenylcarbazole, M-759
- 1-Hydroxy-2-methoxy-10-methyl-3-prenyloxyacridone, T-516
- 12-Hydroxy-11-methoxy-*N*-methyl-*sec*-pseudostyrychnine, I-12
- 3-Hydroxy-4-methoxy-1-methyl-2,6(1*H*,3*H*)-pyridinedione, H-146
- 6-Hydroxy-5-methoxy-1-methyl-2,3,4(1*H*)-pyridinetrione, H-562
- 1-Hydroxy-5-methoxy-6-methyl-2(1*H*)-pyridinone, H-563
- 5-Hydroxy-4-methoxymethyl-1*H*-pyrrole-3-carboxylic acid, H-564
- 6-Hydroxy-4-methoxy-1-methyl-2(1*H*)-quinolinone, Q-46
- 8-Hydroxy-4-methoxy-1-methyl-2(1*H*)-quinolinone, Q-47
- 17-Hydroxy-10-methoxy-4-methylsarpaginium, S-89
- 10'-Hydroxy-10-methoxy-*N*⁴-methyl-17,4',5',6'-tetrahydrousambarensine, U-52
- 10-Hydroxy-10'-methoxy-*N*⁴-methyltetrahydrousambarensine, U-52
- 5-Hydroxy-3-methoxynaphth[2,3-*d*]isothiazole-4,9-dione, A-1540
- 10-Hydroxy-9-methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, T-265
- 4-Hydroxy-8-methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, T-264
- 10-Hydroxy-8-methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, T-266
- 1-Hydroxy-2-methoxynoraporphine, D-531
- 2-Hydroxy-1-methoxynoraporphine, D-531
- 12-Hydroxy-11-methoxynor-*C*-fluorouracrine, N-301
- 5-Hydroxy-6-methoxyonchicine, D-626
- 6-Hydroxy-7-methoxyonchicine, D-625
- 7-Hydroxy-2-methoxyonchicine, D-623
- 7-Hydroxy-6-methoxyonchicine, D-625
- 4-Hydroxy-26-methoxy-2-oxa-11,15,20-triazatricyclo[2.2.2.2.1^{3,7}]nonacos-3,5,7(29),8,22,24,26,27-octaene-10,21-dione, C-556
- 14-Hydroxy-11-methoxy-19-oxogelsenicine, G-44
- 6-[1-(3-Hydroxy-2-methoxy-1-oxopropoxy)ethyl]-1-phenazinecarboxylic acid, A-1160
- 8-Hydroxy-6-methoxy-3-pentylisocarbastyril, R-163
- 8-Hydroxy-6-methoxy-3-pentyl-1(2*H*)-isoquinolinone, R-163
- 7-Hydroxy-7'-methoxyperonatin B, P-271
- 2-(4-Hydroxy-3-methoxyphenacyl)piperidine, P-522
- 1-Hydroxy-6-methoxyphenazine, P-321
- 4-Hydroxy-3-methoxyphenethylamine, H-566
- 3-Hydroxy-4-methoxyphenethylamine, H-565
- N*-β-Hydroxy-β-*p*-methoxyphenethylcinnamide, A-153
- 4-Hydroxy-3-methoxyphenylalanine, A-745
- 2-(2-Hydroxy-4-methoxyphenyl)-5,8-dimethoxy-3-propyl-4(1*H*)-quinolinone, H-567
- N*-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]benzamide, T-67
- N*-[2-(3-Hydroxy-4-methoxyphenyl)ethyl]-3-methyl-2-dodecanamide, H-679
- N*-[2-(4-Hydroxy-3-methoxyphenyl)ethyl]-3-methyl-2-dodecanamide, H-679
- N*-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]-3-phenyl-2-propanamide, A-153
- 4-(4-Hydroxy-3-methoxyphenyl)-4-(1*H*-imidazol-2-yl)-1,2,3-trithian-5-ol, H-507
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]decanamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-hydroxy-8-methyl-6-nonenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-methyldecenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methyl-6-decenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-9-methyl-7-decenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-6-methyl-4-heptenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methylnonanamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-8-methyl-6-nonenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-7-methyloctanamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-7-methyl-5-octenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-10-methyl-8-undecenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]nonanamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]-6-nonenamide, C-100
- N*-[(4-Hydroxy-3-methoxyphenyl)methyl]octanamide, C-100
- 2-(4-Hydroxy-3-methoxyphenyl)-1-methyl-4(1*H*)-quinolinone, F-119
- 4-Hydroxy-4-(4-methoxyphenyl)-1-methyl-5-thioxo-2-imidazolidinone, H-568
- 2-(4-Hydroxy-3-methoxyphenyl)-2-oxoacetamide, D-649
- 1-(4-Hydroxy-3-methoxyphenyl)-2-(2-piperidinylo)ethanone, P-522
- 1-(4-Hydroxy-3-methoxyphenyl)-2-(2-pyrrolidinyl)ethanone, R-165
- 4-Hydroxy-5-methoxy-2-phenylquinoline, D-658
- α-(4-Hydroxy-3-methoxyphenyl)-2-thiazolemethanol, H-569
- (4-Hydroxy-3-methoxyphenyl)-2-thiazolylmethanone, H-569
- 4'-Hydroxy-3'-methoxy-2-(2-piperidyl)acetophenone, P-522
- 7-Hydroxy-4-methoxy-8-prenylfuro[2,3-*b*]quinoline, H-570
- 2-Hydroxy-4-methoxy-3-prenylquinoline, A-1529
- 9-Hydroxy-3-methoxy-6*H*-pyrido[1,2-*a*]pyrazin-6-one, X-29
- 9-Hydroxy-10-methoxy-7*H*-pyrrolo[3,2,1-*de*]phenanthridin-7-one, P-586
- 8-Hydroxy-4-methoxy-2-quinolinecarbothioic acid, D-671
- 8-Hydroxy-4-methoxy-2-quinolinecarboxylic acid, D-677
- 4-Hydroxy-6-methoxy-2-quinolinecarboxylic acid, D-675
- 4-Hydroxy-8-methoxy-2-quinolinecarboxylic acid, D-677
- 2-Hydroxy-8-methoxy-4-quinolinecarboxylic acid, D-673
- 2-Hydroxy-4-methoxyquinoline, Q-33
- 4-Hydroxy-6-methoxy-2(1*H*)-quinolinone, Q-46
- 4-Hydroxy-8-methoxy-2(1*H*)-quinolinone, Q-47
- 7-Hydroxy-8-methoxy-4(1*H*)-quinolinone, Q-48
- 12-Hydroxy-11-methoxyspermostrychnine, S-400
- 12-Hydroxy-11-methoxystrychnine *N*⁹-oxide, S-589
- 10-Hydroxy-11-methoxystrychnine, S-589
- 12-Hydroxy-11-methoxystrychnine, S-589
- 12-Hydroxy-11-methoxystrychnobrasiline, S-594
- 12-Hydroxy-11-methoxystrychnofendlerine, S-594
- 2-(4-Hydroxy-3-methoxystyryl)pyrrolidine, N-305
- 20-Hydroxy-16-methoxytabersonine, H-743
- 10-Hydroxy-11-methoxytabersonine, T-11
- 15-Hydroxy-16-methoxytabersonine, T-11
- 12-Hydroxy-16-methoxy-11,12,13,14-tetrahydrocamoensine, H-571
- 10-Hydroxy-10'-methoxytetrahydrousambarensine, U-52
- 4-Hydroxy-3-methoxy-*N,N,N*-trimethylbenzeneethanaminium, H-566
- 3-Hydroxy-4-methoxy-*N,N,N*-trimethylbenzeneethanaminium, H-565
- 4-Hydroxy-3-methoxytrimethylphenethylammonium, H-566
- 5-Hydroxy-6-methoxytryptamine, D-684
- 6-Hydroxy-5-methoxytryptamine, D-684
- 8-Hydroxy-4-methoxy-1-vinyl-β-carboline, D-686
- 6-Hydroxy-4-methoxy-1-vinyl-β-carboline, M-290
- 1-Hydroxy-10-methylacridone, H-402
- 8-Hydroxy-3-methyladenine, A-819
- 10-Hydroxy-17-*O*-methyllakagerine, A-222
- 5-Hydroxymethylakumammiline, A-231
- 2-Hydroxy-*ω*-methylallophanic acid, M-385
- Hydroxyl(methylamino)carbonylcarbamic acid, M-385
- 17-Hydroxy-20-(methylamino)-19,21-dinorchola-4,11-diene-3,22-dione, S-415
- 4-[1-Hydroxy-2-(methylamino)ethyl]-1,2-benzenediol, A-152
- p*-(α-Hydroxy-β-methylaminoethyl)phenol, S-659
- 4-Hydroxy-α-[(methylamino)methyl]benzenemethanol, S-659
- 3-Hydroxy-6-(methylamino)-6-(2-methylpropyl)-3-(phenylmethyl)-2,5-morpholinedione, M-231
- 18-Hydroxy-20-(methylamino)pregna-1,4-dien-3-one, A-810
- 20-Hydroxy-3-(methylamino)pregn-5-en-18,20-olide, P-93
- 2-Hydroxy-3-(methylamino)pregn-5-en-16-one, A-813
- 2-Hydroxy-6-methylamino-9*H*-purine, A-818
- 6-Hydroxy-2-methylaminopurine, G-210
- 3-Hydroxy-8-methyl-8-azabicyclo[3.2.1]octane-2-carboxylic acid, E-8
- 6-Hydroxy-8-methyl-8-azabicyclo[3.2.1]octan-3-one, M-391
- 1-(3-Hydroxy-8-methyl-8-azabicyclo[3.2.1]oct-2-en-2-yl)-3-phenyl-2-propen-1-one, C-341
- 6-Hydroxy-8-methyl-8-azabicyclo[3.2.1]oct-3-yl α-(hydroxymethyl)benzeneacetate, H-521
- 10-Hydroxy-19-methylazacyclocoisa-3,5,7,11,13,15,17-heptaen-2-one, C-832
- 6-Hydroxy-1-methyl-4-azafluorenone, O-93
- 2-Hydroxy-1-methyl-4-azafluorenone, O-93
- 7-(Hydroxymethyl)benz[6,7]indolizino[1,2-*b*]quinolin-11(13*H*)-one, H-403
- 6-Hydroxy-3-methyl-2(3*H*)-benzothiazolone, B-78
- 4-[1-[(2-Hydroxy-6-methylbenzoyl)oxy]ethyl]-6-methyl-6*H*,10*H*-pyrido[3,2-*a*:5,6,1-*d'*,*e'*]diphenazine-11,17-dicarboxylic acid, E-223
- 2-Hydroxy-2-methylbutanoic acid, H-572
- 3-Hydroxy-3-methylbutanoic acid, H-573
- 4-Hydroxy-3-methyl-2-butenenitrile, H-576
- 4-Hydroxy-3-methyl-2-butenic acid, H-576
- 4-Hydroxy-2-methyl-2-butenic acid, H-575
- 2-Hydroxymethyl-2-butenic acid, H-574
- 6-(4-Hydroxy-3-methyl-2-butenylamino)purine, Z-11
- 1-Hydroxy-4-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, C-507
- 2-Hydroxy-7-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, C-500
- 2-Hydroxy-1-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-125
- 6-Hydroxy-5-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-577
- 5-(3-Hydroxy-3-methyl-1-butenyl)-4,6-dimethoxyfuro[2,3-*b*]quinoline, H-578
- 3-(2-Hydroxy-3-methyl-3-butenyl)-2,4-dimethoxy-7,8-methylenedioxyquinoline, D-715
- (4-Hydroxy-3-methyl-2-butenyl)guanidine, H-579
- 2-Hydroxy-5-(3-methyl-2-butenyl)-3*H*-indol-3-one, M-415
- 2-Hydroxy-6-(3-methyl-2-butenyl)-3*H*-indol-3-one, M-416
- 3-(3-Hydroxymethyl-2-butenyl)-7-(3-methyl-2-butenyl)-1*H*-indole, B-202
- 3-(2-Hydroxy-3-methyl-3-butenyl)-5-(3-methyl-1-oxo-2-butenyl)-1*H*-indole, B-200
- N*-[2-Hydroxy-2-[4-[(3-methyl-2-butenyl)oxy]phenyl]ethyl]-3-phenyl-2-propanamide, O-63
- 6-(3-Hydroxy-3-methyl-1-butenyl)-1-phenazinecarboxylic acid, M-417
- 5-(2-Hydroxy-3-methyl-3-butenyl)-4,6,7-trimethoxyfuro[2,3-*b*]quinoline, H-580
- 3-(2-Hydroxy-3-methyl-3-butenyl)-4,6,8-trimethoxy-1-methyl-2(1*H*)-quinolinone, P-755
- 3-(2-Hydroxy-3-methyl-3-butenyl)-4,7,8-trimethoxy-2(1*H*)-quinolinone, P-625
- 4-(4-Hydroxy-3-methyl-2-butenyl)tryptophan, M-419

- 4-(2-Hydroxy-3-methyl-3-butenyl)jucocitrine, H-581
- 6-[(4-Hydroxy-3-methylbutyl)amino]purine, Z-11
- 3-(2-Hydroxy-3-methylbutyl)-4,8-dimethoxy-1-methyl-2(1*H*)-quinolinone, L-288
- 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methylbutanoyl)pyrazine, B-184
- 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methyl-2-butenoyl)pyrazine, B-184
- 2-(1-Hydroxy-3-methylbutyl)-3,6-dimethyl-5-(3-methylbutyl)pyrazine, B-184
- 9-(3-Hydroxy-3-methylbutyl)-7-methoxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-212
- 3-(2-Hydroxy-3-methylbutyl)-8-methoxy-1-methyl-4(1*H*)-quinolinone, P-426
- 3-Hydroxy-3-methylbutyric acid, H-573
- 5-Hydroxymethylcanthin-6-one, H-582
- 3-(Hydroxymethyl)-9*H*-carbazole, C-124
- 3-Hydroxy-6-methyl-9*H*-carbazole-1-carboxaldehyde, H-589
- 4-Hydroxy-6-methyl-9*H*-carbazole, H-588
- 1-Hydroxy-3-methyl-9*H*-carbazole, H-583
- 1-Hydroxy-6-methyl-9*H*-carbazole, H-584
- 2-Hydroxy-3-methyl-9*H*-carbazole, H-585
- 2-Hydroxy-7-methyl-9*H*-carbazole, H-586
- 3-Hydroxy-6-methyl-9*H*-carbazole, H-587
- 6-Hydroxy-1-methyl-β-carboline, H-591
- 7-Hydroxy-1-methyl-β-carboline, H-592
- 8-Hydroxy-1-methyl-β-carboline, H-593
- 1-Hydroxymethyl-β-carboline, H-590
- 8-Hydroxy-2-methyl-β-carbolineum, P-917
- 16-Hydroxymethylceridimine, C-281
- 5-Hydroxy-*N*-methylcorydaline, I-304
- 10-Hydroxy-*N*^b-methylcorynantheol, D-412
- 4-Hydroxy-3-methylcrotonic acid, H-576
- 5-Hydroxymethylcytosine, A-795
- 1-Hydroxymethylasycarpidan, U-8
- 3-(1-Hydroxy-4-methyl-2,4,6,8-decatetraenyli-dene)-2,4-pyrrolidinedione, R-30
- 9-Hydroxy-3-methyl-2-decenoic acid, H-594
- 12-Hydroxy-*N*¹-methyl-14,15-didehydroaspidiofractinine, A-1494
- 1-Hydroxymethyl-1,2-didehydropyrrolizidine, S-644
- 8-Hydroxymethylidihydrochelerythrine, B-239
- 6-Hydroxymethylidihydronitidine, H-595
- 8-Hydroxymethylidihydronitidine, H-595
- 8-Hydroxymethylidihydrosanguinarine, H-596
- 1-Hydroxymethyl-6,7-dimethoxyisoquinoline *N*-oxide, D-590
- 3-Hydroxymethyl-6,8-dimethoxy-1-methylisoquinoline, D-710
- 8-(Hydroxymethyl)-2,4-dimethoxy-3*H*-phenoxazin-3-one, M-590
- 8-(Hydroxymethyl)-6,11-dimethyl-2*H*,4*H*-oxazol[5,4,3-*ij*]pyrido[3,2-*g*]quinoline-4,10(11*H*)-dione, N-346
- N*-Hydroxy-*N*-(15-methyl-3,10-dioxopentadecyl)-seramide, L-189
- N*-(3-Hydroxy-10-methylododecanoyl)nicotinic acid, P-957
- N*-(3-Hydroxy-11-methylododecanoyl)nicotinic acid, P-957
- 1-(8-Hydroxy-6-methyl-2,4,6-dodecatrienoyl)-2-pyrrolidone, V-34
- 3-(1-Hydroxy-10-methylododecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-11-methylododecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-4-methylododecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-5-methylododecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 2-(10-Hydroxy-10-methylododecyl)-3-methoxy-4(1*H*)-quinolinone, H-597
- 2-(11-Hydroxy-11-methylododecyl)-3-methoxy-4(1*H*)-quinolinone, H-598
- 9-Hydroxy-1,2-methylenedioxyaporphine, A-1066
- 10-Hydroxy-1,2-methylenedioxyaporphine, L-72
- 7-Hydroxy-1,2-methylenedioxyaporphine, O-87
- 11-Hydroxy-1,2-methylenedioxyaporphine, P-781
- 4-Hydroxy-1,2-methylenedioxyaporphine, S-549
- 3-Hydroxy-8,9-methylenedioxybenzo[*h*]quinoline, H-599
- 8-Hydroxy-6,7-methylenedioxydictamine, F-217
- 7-Hydroxy-1,2-methylenedioxy-9,10-dimethoxyaporphine, D-948
- 1-Hydroxy-6,7-methylenedioxyisoquinoline, I-305
- 7-Hydroxy-1,2-methylenedioxy-9-methoxyaporphine, O-85
- 6-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, H-600
- 8-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262
- 9-Hydroxy-1,2-methylenedioxyaporphine, A-1066
- 3-Hydroxy-1,2-methylenedioxyaporphine, C-466
- 7-Hydroxy-1,2-methylenedioxyaporphine, O-87
- 11-Hydroxy-1,2-methylenedioxyaporphine, P-781
- 3-Hydroxy-8,9-methylenedioxyphenanthridine, M-448
- 2-(3-Hydroxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone, G-171
- 8-Hydroxy-6,7-methylenedioxy-5-prenyldictamine, T-57
- 7-Hydroxy-1-methylenepyrrolizidine *N*-oxide, H-601
- 7-Hydroxy-1-methylenepyrrolizidine, H-601
- 6-Hydroxy-3-*O*-methylpimaridine, P-86
- 1-Hydroxymethyl-1,2-epoxypyrrolizidine, E-133
- 3-(Hydroxymethyl)-9-ethyl-9*H*-carbazole, C-124
- 2-(1-Hydroxy-1-methylethyl)-2,3-dihydrofuro[2,3-*b*]quinolin-6-ol, R-97
- 2-(1-Hydroxy-1-methylethyl)-6-isopropylpyrazine, D-701
- 2-(1-Hydroxy-1-methylethyl)-3-methoxypyrazine, H-545
- 12'-Hydroxy-2'-(1-methylethyl)-5'-(2-methylpropyl)ergotaman-3',6',18-trione, E-157
- 1-Hydroxy-6-(1-methylethyl)-3-(1-methylpropyl)-2(1*H*)-pyrazinone, H-543
- 1-Hydroxy-6-(1-methylethyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, M-462
- 1-Hydroxy-6-(1-methylethyl)-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-544
- 12'-Hydroxy-2'-(1-methylethyl)-5'-(phenylmethyl)ergotaman-3',6',18-trione, E-155
- 2-Hydroxy-3-(1-methylethyl)pyrazine, H-545
- 4-Hydroxy-4-(1-methylethyl)-1-(8,10,12-trimethyl-1-oxo-2,4,6,8-tetradeecatetraenyl)-6-oxa-3-azabicyclo[3,1,0]hexan-2-one, A-1197
- α-1-*C*-(Hydroxymethyl)fagomine, B-185
- N*-Hydroxymethylflindersine, F-90
- 3-(Hydroxymethyl)furan, F-199
- 1-(5-Hydroxymethyl-2-furanyl)-γ-carboline, I-262
- 1-[5-(Hydroxymethyl)-2-furanyl]-9*H*-pyrido[3,4-*bj*]indole-3-carboxylic acid, F-85
- 5-Hydroxy-11-methylfuro[2,3-*c*]acridin-6(11*H*)-one, F-206
- 6-Hydroxy-9-methylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-211
- 18-Hydroxy-14-*O*-methylgadesine, G-152
- 20-(Hydroxymethyl)gelsedine, G-47
- N*-(3-Hydroxy-3-methylglutaryl)tryptophan, T-640
- 8-Hydroxy-7-methylguanaine, A-753
- 8-Hydroxy-9-methylguanaine, A-753
- 16-Hydroxy-*N*-methyl-4-hexadecanamide, H-501
- 2-Hydroxymethyl-5-(1-hydroxypentyl)-3,4-pyrrolidinediol, H-602
- 4-Hydroxymethyl-2-(2-hydroxyphenyl)-2-thiazoline, H-695
- 2-(Hydroxymethyl)-6-(2-hydroxytridecyl)-3,4-piperidinediol, B-51
- 2-(Hydroxymethyl)-6-(2-hydroxyundecyl)-3,4-piperidinediol, B-50
- 1-Hydroxy-4-methyl-1*H*-imidazole, M-473
- 1-Hydroxy-5-methyl-1*H*-imidazole, M-473
- 4-(Hydroxymethyl)-1*H*-indeno[1,2-*b*]pyridine-2,5-dione, D-385
- 8-Hydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, O-93
- 3-Hydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, O-93
- 7-Hydroxymethylindole, I-99
- 3-Hydroxymethylindole, I-98
- α-Hydroxy-1-methyl-1*H*-indole-3-propanoic acid, H-539
- 8-Hydroxymethylindolizidine, O-47
- 5-(Hydroxymethyl)-6*H*-indolo[3,2,1-*de*][1,5]-naphthyridin-6-one, H-582
- 2-Hydroxy-3-(1-methylindolyl)propanoic acid, H-539
- 3-Hydroxymethyl-1-isopentylpyrrolidine, M-421
- 1-Hydroxymethyl-6,7-isoquinolinediol, D-590
- N*¹-Hydroxymethyl-(16*S*)-isosisiririkine, I-319
- 10-Hydroxy-17-*O*-methylkribine, K-101
- 10-Hydroxy-21-*O*-methylkribine, K-101
- 6-Hydroxymethyllumazine, H-623
- 10-Hydroxymethyllycaconitine, G-91
- 14-Hydroxy-15-methyllycopodan-5-one, F-73
- 8-Hydroxy-2-*N*-methylmanzamine D, M-91
- 5-Hydroxymethyl-9-methoxycanthin-6-one, H-582
- 3-(Hydroxymethyl)-9-methoxy-9*H*-carbazole, C-124
- 6-Hydroxymethyl-1-methoxycarbazole, H-604
- 3-Hydroxymethyl-1-methoxycarbazole, H-603
- N*-(1-Hydroxymethyl-2-methoxyethyl)-7-methoxy-4-eicosenamide, H-605
- 5-(Hydroxymethyl)-9-methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-582
- 1-(Hydroxymethyl)-7-methoxy-6-methyl-5,8-isoquinolinedione, R-45
- 6-Hydroxymethyl-4-methoxy-5-(methylthio)-2,2'-bipyridine, C-575
- 6-Hydroxymethyl-9-methoxy-1-phenazinecarboxylic acid, H-514
- 8-(Hydroxymethyl)-2-methoxy-3*H*-phenoxazin-3-one, M-590
- 3-(2-Hydroxymethyl-4-methoxyphenyl)-6-methoxy-4(3*H*)-quinazolinone, D-369
- 5-(Hydroxymethyl)-4-(4-methoxyphenyl)-2-oxazolindione, C-964
- 16-Hydroxy-14-methyl-3-(methylamino)-4-methylene-9,19-cyclopregnan-20-one, H-482
- 2-Hydroxymethyl-2'-methylamino-4,5-methylenedioxybiphenyl, I-186
- 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1*H*-indol-3-yl]butyl hexanoate, M-480
- 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1*H*-indol-3-yl]butyl 9*Z*,12*Z*-octadecadienoate, M-480
- 3-Hydroxy-3-methyl-2-[6-(3-methyl-2-butenyl)-1*H*-indol-3-yl]butyl 9*Z*-octadecenoate, M-480
- N*-[1-(Hydroxymethyl)-3-methylbutyl]acetamide, A-838
- 3-(2-Hydroxymethyl-3,4-methylenedioxyphenyl)-5,6-methylenedioxyisoquinoline, D-136
- 8-Hydroxy-3-methyl-7-(1-methylethyl)benz[*cd*]indol-2(1*H*)-one, P-631
- 5-Hydroxymethyl-2-(1-methylethyl)-3,4-pyridinediol, A-620
- 3-(Hydroxymethyl)-4-(1-methyl-1*H*-imidazol-5-yl)-1-phenyl-1-butanone, H-606
- 4-(Hydroxymethyl)-2-methyl-1*H*-indole, M-478
- 5-Hydroxy-3-methyl-1-(3-methyl-2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one, H-467
- 5-Hydroxy-3-methyl-1-(3-methyl-2-oxo-2*H*-pyrrolin-5-yl)-3-pyrrolin-2-one, H-467
- 8-Hydroxy-8-methyl-6-(2-methylpentylidene)indolizidine, A-359
- 2-(Hydroxymethyl)-6-methyl-3,4,5-piperidinetriol, H-607
- 6-[5-(Hydroxymethyl)-1-methyl-3-piperidinyl]-2(1*H*)-pyridinone, K-108
- 8-[1-(Hydroxymethyl)-2-methyl-2-propenyl]-7-methoxy-2*H*-1-benzopyran-2-one, I-249
- 5-Hydroxymethyl-2-(1-methylpropyl)-3,4-pyridinediol, A-620
- 5-Hydroxymethyl-2-(2-methylpropyl)-3,4-pyridinediol, A-620
- 4-(Hydroxymethyl)-6-methyl-2,3,5-pyridinetriol, H-608
- 5-Hydroxymethyl-1-methylpyrrolicarboxaldehyde, H-626

- 2-(Hydroxymethyl)-5-methyl-3,4-pyrrolidinediol, D-591
- 2-Hydroxymethyl-4-methylquinazoline, H-609
- 3-Hydroxy-5-methyl-1-naphthalenecarboxylic acid, H-610
- 3-Hydroxy-5-methyl-1-naphthoic acid, H-610
- N*-Hydroxy-14-methyl-*N*-nitrosopentadecanamine, N-253
- N*-Hydroxy-12-methyl-*N*-nitrosotridecanamine, N-255
- 4-Hydroxy-3-methyl-2-(2-nonenyl)quinoline, M-491
- 1-Hydroxy-3-methyl-2-(2-nonenyl)-4(1*H*)-quinolinone, M-491
- 4-Hydroxy-2-methyl-6-nonylpiperidine, H-611
- 12-Hydroxy-14-methyl-21-norsenecionan-11,16-dione, S-241
- 12-Hydroxy-14-methyl-18-norsenecionan-8,11,16-trione, C-770
- 5-Hydroxy-5-(6-methyl-2,4-octadienyl)-3-pyrrolin-2-one, A-1573
- 5-Hydroxy-7-*O*-methyldoline, U-37
- 2-Hydroxy-6-*O*-methyldoline, U-37
- 1-(8-Hydroxy-6-methyl-1-oxo-2,4,6-dodecatrienyl)-2-pyrrolidinone, V-34
- 3-Hydroxy-2-methyl-6-(11-oxododecyl)piperidine, C-183
- 3-Hydroxy-2-methyl-6-(10-oxododecyl)piperidine, P-655
- 3-Hydroxy-6-methyl-2-(11-oxododecyl)pyridine, H-612
- 1-(3-Hydroxy-10-methyl-1-oxododecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 1-(3-Hydroxy-11-methyl-1-oxododecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 3-(2-Hydroxy-4-methyl-3-oxohexyl)-1*H*-indole, H-535
- 1-(3-Hydroxy-14-methyl-1-oxopentadecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 3-(2-Hydroxy-4-methyl-3-oxopentyl)-1*H*-indole, H-536
- 6-Hydroxy-2-methyl-5-(1-oxopropyl)-2*H*-1,4-oxazine-3-carboxylic acid, O-151
- 4-Hydroxy-2-methyl-6-(2-oxopropyl)piperidine, H-622
- 5-Hydroxy-3-methyl-1-(2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one, D-448
- 3-Hydroxy-2-methyl-6-(13-oxotetradecyl)piperidine, S-388
- 1-(3-Hydroxy-12-methyl-1-oxotetradecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 1-(3-Hydroxy-13-methyl-1-oxotetradecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 1-(3-Hydroxy-12-methyl-1-oxotridecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 1-(3-Hydroxy-10-methyl-1-oxoundecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- N*-(3-Hydroxy-14-methylpentadecanoyl)nornicotine, P-957
- 3-(1-Hydroxy-13-methylpentadecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-14-methylpentadecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- N,N'*-[1-(Hydroxymethyl)-1,5-pentanediy]bis[2,3-dihydroxybenzamide], M-806
- N*^b-(2-Hydroxy-3-methylpentanoyl)tryptamine, T-639
- N*-[2-(1-Hydroxy-4-methylpentyl)]acetamide, A-838
- 4-Hydroxy-3-methyl-2-pentylquinoline, M-508
- 10-Hydroxy-*N*-methylpericyclivine, S-89
- 1-Hydroxy-5-methylphenazinium hydroxide inner salt, P-833
- β-Hydroxy-*N*-methylphenethylamine, A-864
- p*-Hydroxymethylphenol, H-414
- 3-Hydroxy-8-methyl-α-phenyl-8-azabicyclo[3.2.1]octane-2-methanol, H-429
- 12-Hydroxy-2-methyl-5-(phenylmethyl)ergotaman-3',6',18-trione, E-172
- 3-Hydroxy-2-methyl-6-(9-phenylonyl)piperidine, I-177
- (2-Hydroxy-6-methylphenyl)(octahydro-7-hydroxy-8-indoliziny)methanone, I-231
- N*-[3-[2-[2-(2-Hydroxy-6-methylphenyl)-2-oxoethyl]-1-pyrrolidinyl]propyl]hexanamide, P-262
- 4-Hydroxy-1-methyl-α-phenylpiperidineethanol, S-210
- 6-(3-Hydroxy-2-methyl-3-phenylpropanoiloxy)-3,7-tropanediol, M-392
- 3-Hydroxy-3-(2-methylphenyl)-2-propenamide, M-512
- 3-Hydroxy-2-methyl-1-phenyl-4(1*H*)-pyridinone, D-632
- 4-Hydroxy-1-methyl-3-phenyl-2(1*H*)-quinolinone, D-657
- 5-Hydroxy-1-methyl-2-phenyl-4(1*H*)-quinolinone, D-658
- 5-Hydroxy-6-methylpipercolic acid, H-613
- 5-Hydroxy-6-methyl-2-piperidinecarboxylic acid, H-613
- 6-(Hydroxymethyl)-3,4-piperidinediol, H-615
- 2-(Hydroxymethyl)-3,4-piperidinal, H-614
- 5-Hydroxy-6-methyl-2-piperidinedodecanoic acid, S-409
- 5-Hydroxy-6-methyl-2-piperidinedodecanol, J-55
- 6-(Hydroxymethyl)-2,3,4,5-piperidinetetrol, A-727
- 6-Hydroxymethyl-2,3,4,5-piperidinetetrol, A-726
- 2-(Hydroxymethyl)-3,4,5-piperidinetriol, 12CI, H-616
- 5-Hydroxy-6-methyl-2-piperidineundecanoic acid, L-120
- 6-(Hydroxymethyl)-3-piperidinol, H-699
- 4'-Hydroxy-2-(1-methylpiperidin-2-yl)acetophenone, P-522
- 12-(5-Hydroxy-6-methyl-2-piperidinyl)-1,2-dodecanediol, L-119
- 12-(5-Hydroxy-6-methyl-2-piperidinyl)-2-dodecanone, C-183
- 12-(5-Hydroxy-6-methyl-2-piperidinyl)-3-dodecanone, P-655
- 16-(5-Hydroxy-6-methyl-2-piperidinyl)-2,13-hexadecanediol, A-477
- 1-(4-Hydroxy-6-methyl-2-piperidinyl)-2-propanone, H-622
- 14-(5-Hydroxy-6-methyl-2-piperidinyl)-2-tetradecanone, S-388
- 16-Hydroxymethylpleiocarpamine, P-518
- 3-Hydroxy-6-methyl-4-prenylcarbazole, G-125
- 2-Hydroxy-3-methyl-1-prenylcarbazole, M-728
- 2-Hydroxy-6-methyl-3-prenylcarbazole, S-291
- 1-Hydroxy-10-methyl-3-prenyloxyacridone, D-528
- 3-Hydroxy-10-methyl-1-prenyloxyacridone, D-528
- 3-Hydroxy-4-methylproline, H-627
- 3-Hydroxy-5-methylproline, H-628
- 2-Hydroxy-2-methylpropanenitrile, H-617
- 2-Hydroxy-2-methylpropanoic acid, H-617
- 6-Hydroxy-*N*-(2-methylpropyl)-2,4-decadienamide, D-106
- 4-Hydroxy-*N*-(2-methylpropyl)-2-hexenamide, H-502
- 2-(1-Hydroxy-1-methylpropyl)-3-methoxypyrazine, H-618
- 2-(1-Hydroxy-2-methylpropyl)-3-methoxypyrazine, H-619
- 1-Hydroxy-5-(2-methylpropyl)-3-(2-methylpropylidene)-2,6(1*H*,3*H*)-pyrazinedione, F-112
- 1-Hydroxy-6-(1-methylpropyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, A-1475
- 6-(1-Hydroxy-1-methylpropyl)-3-(1-methylpropyl)-2(1*H*)-pyrazinone, B-204
- 6-(2-Hydroxy-2-methylpropyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, F-76
- 1-Hydroxy-3-(2-methylpropyl)-6-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-620
- 1-Hydroxy-6-(2-methylpropyl)-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-621
- 5-Hydroxy-1-(2-methylpropyl)-5-(1-oxohexyl)-2-pyrrolidinone, P-449
- N*-(2-Hydroxy-2-methylpropyl)-8-oxo-2,4-tetradecadienamide, O-211
- N*-(2-Hydroxy-2-methylpropyl)-12-oxo-2,4,8-tetradecatrienamide, O-212
- N*-(2-Hydroxy-2-methylpropyl)-2,9-pentadecadiene-12,14-diyamide, P-202
- 4-Hydroxy-2-methyl-6-propylpiperidine, H-622
- 5-Hydroxymethyl-2-propyl-3,4-pyridinediol, A-620
- 4-Hydroxy-2-(1-methylpropyl)quinoline, M-531
- 7-Hydroxy-*N*-(2-methylpropyl)-2,8-tridecadiene-10,12-diyamide, H-748
- 15-Hydroxy-*N*-methyl-*sec*-pseudostyrychne, I-12
- 6-(Hydroxymethyl)-2,4(1*H*,3*H*)-pteridinedione, H-623
- 6-Hydroxymethylpterin, A-802
- 2-Hydroxy-3-methylpyrazine, H-624
- 5-Hydroxy-3-methyl-2(1*H*)-pyrazinone, D-631
- 5-Hydroxy-6-methyl-3,4-pyridinedimethanol, P-923
- 4-Hydroxy-2-methylpyridine, H-625
- 3-Hydroxymethylpyridine, P-904
- 3-Hydroxy-2-methyl-4(1*H*)-pyridinone, D-632
- 12-(3-Hydroxy-6-methyl-2-pyridinyl)-2-dodecanone, H-612
- α'-(Hydroxymethyl)-9*H*-pyrido[3,4-*b*]indole-1,3-dimethanol, P-893
- 4-Hydroxy-5-methyl-2(1*H*)-pyrimidinone, T-401
- 5-(Hydroxymethyl)-1*H*-pyrrole-2-carboxaldehyde, H-626
- 1-Hydroxymethyl-2-pyrrolicarboxaldehyde, P-942
- 3-Hydroxy-4-methyl-2-pyrrolidinecarboxylic acid, H-627
- 3-Hydroxy-5-methyl-2-pyrrolidinecarboxylic acid, H-628
- 2-(Hydroxymethyl)-3,4-pyrrolidinediol, 12CI, D-593
- 5-Hydroxy-3-methyl-3-pyrrolin-2-one, D-449
- 1-Hydroxymethylpyrrolizidine methyl 2-*O*-acetyl-2-isopropylmalate, H-631
- 7-Hydroxymethyl-1,2-pyrrolizidinediol, D-595
- 7-Hydroxymethyl-1,6-pyrrolizidinediol, D-597
- 1-Hydroxymethyl-1,7-pyrrolizidinediol, D-596
- 1-Hydroxy-7-methylpyrrolizidine, H-630
- 1-Hydroxymethylpyrrolizidine, H-629
- 1-[4-(Hydroxymethyl)-1*H*-pyrrol-3-yl]ethanone, A-57
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-12,15-heneicosadien-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-12,15,18-heneicosatrien-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-14-heneicosen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-16-methyl-1-heptadecanone, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-1-nonadecanone, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-14-nonadecen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-7,10,13,16,19,22-pentacosahexaen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-10,13,16,19,22-pentacosapentaen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-13,16,19,22-pentacosatetraen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-14,17-tricosadien-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-11,14,17,20-tricosatetraen-1-one, M-776
- 1-[5-(Hydroxymethyl)-1*H*-pyrrol-2-yl]-14,17,20-tricosatrien-1-one, M-776
- 4-Hydroxy-3-methylquinaldine, H-477
- 4-Hydroxy-2-methylquinazoline, H-632
- 4-Hydroxy-2-methylquinoline, H-633
- 4-(Hydroxymethyl)quinoline, Q-38
- 2-(Hydroxymethyl)quinoline, Q-37
- 1-Hydroxymethylquinolizidine, O-58
- 19(*S*)-Hydroxy-*N*^b-methylraumacline, R-22
- 2-Hydroxy-*N*-methylsaligamine, A-874
- 12-Hydroxy-4-methyl-4,8-secosenecionan-8,11,16-trione, S-245
- 3-Hydroxy-28-methyl-16,28-secosolanidan-6-one, P-807
- 7-Hydroxy-*O*-methylsolanocapsine, S-346
- 10-Hydroxymethylsparteine, H-634
- 5'-Hydroxy-4'-*N*-methylstaurosporine, S-498
- 3-Hydroxy-4'-*N*-methylstaurosporine, S-498
- 5-Hydroxy-9-methylstreptimidone, M-561
- 13β-Hydroxy-*N*-methylstylopinium, S-612
- 3-Hydroxy-2-(methylsulfonyl)-1*H*-indole, H-534
- N*-(3-Hydroxy-12-methyltetradecanoyl)nornicotine, P-957

- N*-(3-Hydroxy-13-methyltetradecanoyl)nornicotine, P-957
- 3-(1-Hydroxy-12-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-13-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-5-methyltetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-12-methyltetradecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- 3-(1-Hydroxy-13-methyltetradecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- 1-Hydroxy-4-(methylthio)canthin-6-one, H-635
- 3-Hydroxy-2-(methylthio)-1*H*-indole, H-534
- 1-Hydroxy-4-methylthio-6*H*-indolo[3,2,1-*de*][1,5]-naphthyridin-6-one, H-635
- N*-(3-Hydroxy-12-methyltridecanoyl)nornicotine, P-957
- 3-(1-Hydroxy-12-methyltridecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- 2-(12-Hydroxy-12-methyltridecyl)-3-methoxy-4(1*H*)-quinolinone, H-636
- 2-(12-Hydroxy-12-methyltridecyl)-4(1*H*)-quinolinone, H-636
- 5-Hydroxymethyl-1-[2-(3,4,5-trimethoxyphenyl)ethyl]-2-pyrrolicarboxaldehyde, P-296
- 5-Hydroxy-*N*^α-methyltryptamine, H-755
- 2-Hydroxy-*N*-methyltryptamine, H-754
- N*^β-Hydroxy-*N*^β-methyltryptamine, M-582
- 5-Hydroxy-*O*-methyltylophorimidine, T-696
- N*-(3-Hydroxy-10-methylundecanoyl)nornicotine, P-957
- 3-Hydroxy-*N*-methylwelwitindolinone C isonitrile, W-13
- 3-Hydroxy-*N*-methylwelwitindolinone C isothiocyanate, W-13
- 16-Hydroxymethylhohimb-16-ene, N-243
- 2-Hydroxy-1'-methylzeatin, M-586
- 23-Hydroxymilbemycin α2, M-607
- 23-Hydroxymilbemycin α3, M-608
- 26-Hydroxymilbemycin α1, M-607
- 26-Hydroxymilbemycin α2, M-607
- 26-Hydroxymilbemycin α3, M-608
- 4-Hydroxymimosamycin, M-624
- Hydroxyminaline, H-714
- 7-Hydroxymitragynine, M-657
- 9-Hydroxymukanadin B, M-727
- 13α-Hydroxymultiflorine, M-731
- 13β-Hydroxymultiflorine, M-731
- Hydroxymuscovopyridine A, M-774
- Hydroxymuscovopyridine B, M-774
- 30-Hydroxymycaloidine A, M-779
- 32-Hydroxymycaloidine A, M-779
- 38-Hydroxymycaloidine B, M-779
- Hydroxymyoscorpine, E-31
- 17-Hydroxymyrsinol (incorr.), E-136
- 14-Hydroxynaamidine A, N-1
- 14-Hydroxynaamidine G, N-1
- N*-Hydroxynandigerine, N-25
- Hydroxynantenine, N-29
- 10-Hydroxy-18-(*N*-naphthalenyl-*N*-phenylamino)-betaenone C, H-637
- 3-Hydroxy-1,4-naphthoquinone-2-carboxylic acid, H-638
- β-Hydroxyneosaspergillie acid, F-76
- 3'-Hydroxyneoharringtonine, N-117
- 1''-Hydroxyneoharringtonine, N-117
- 7-Hydroxyneolamellarin A, N-121
- 5-Hydroxyneolamellarin B, N-121
- 10-Hydroxyneoline, M-311
- 15-Hydroxyneoline, M-313
- 11α-Hydroxyneosaxitoxin, H-731
- 11β-Hydroxyneosaxitoxin, H-731
- N*'-Hydroxyneosaxitoxin, S-109
- Hydroxyneosenkirkine, S-245
- 6'-Hydroxyneothiobinupharidine, T-380
- 6-Hydroxyneothiobinupharidine, T-380
- 4'-Hydroxyneobouldine, N-185
- 4-Hydroxynicotinamide, H-709
- 6-Hydroxynicotinic acid, H-711
- 4-Hydroxynicotinic acid, H-709
- 10-Hydroxynigraninone, O-21
- N*⁷-(4-Hydroxy-3-nitrobenzyl)theophylline, P-357
- 17-Hydroxy-10-nitroestr-4-en-3-one, H-639
- 10-Hydroxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, H-600
- 4-Hydroxy-2-nitrophenethylamine, H-640
- N*-(4-Hydroxy-3-nitrophenylethyl)acetamide, H-640
- 2-(4-Hydroxy-3-nitrophenyl)ethylamine, H-640
- N*-[2-(Hydroxynitrosamino)-3-methylbutyl]octanamide, F-146
- 2-(Hydroxynitrosoamino)-12-methyl-1,3,12-tetradecanetriol, A-693
- 4-Hydroxy-3-nitrosobenzoic acid, H-641
- N*-Hydroxy-*N*-nitrosohexadecanamine, N-254
- 6-Hydroxynobiline, N-262
- 9-Hydroxynominine, H-166
- 2-Hydroxynominine, H-163
- 4-Hydroxy-2-(3,6-nonadienyl)quinoline, H-642
- 4-Hydroxy-2-(1-nonenyl)quinoline, H-642
- 3-Hydroxy-3-nonyl-2,4(1*H*,3*H*)-quinolinedione, H-643
- 4-Hydroxy-2-nonylquinoline, H-642
- 3-Hydroxy-2-nonyl-4(1*H*)-quinolinone, A-621
- 2-(9-Hydroxynonyl)-4(1*H*)-quinolinone, H-642
- 5-Hydroxynoracrocynine, A-1527
- 3-Hydroxynoralllosedamine, S-210
- 3-Hydroxy-19-norconca-1,3,5(10),14-tetraene, D-788
- 11-Hydroxy-23-norconca-1,4,18(22)-trienin-3-one, H-322
- 12-Hydroxynorconca-*N*(18),1,4-trienin-3-one, H-322
- 3-Hydroxy-11-norcytisine, H-644
- 7-Hydroxynordicentrine, D-948
- 18-Hydroxynordihydrotoxiferine, T-436
- 3β-Hydroxynorerythrosumamide, N-299
- 18-Hydroxynorfluorourarine, N-301
- 6'-Hydroxy-9-normethylbudmunchiamine K, B-389
- 4-Hydroxynornantenine, N-29
- 3-Hydroxynornuciferine, T-521
- 16-Hydroxy-22-norparsonsine, P-97
- 12-Hydroxy-18-norseneconan-11,16-dione, N-219
- 10-Hydroxynortetraphyllicine, T-288
- N*-Hydroxynorthalictuberine, T-318
- 3-Hydroxynortropane, A-1597
- 11-Hydroxynorzoanthamine, N-317
- 3-Hydroxynorzoanthamine, N-317
- 30-Hydroxynorzoanthamine, N-317
- 15-Hydroxynovocaine, I-12
- 3-Hydroxynuciferine, T-521
- 10-Hydroxynudicaulidine, M-307
- Hydroxynubodycin, N-346
- N*²-(17-Hydroxy-9,12-octadecadienyl)glutamine, G-108
- N*²-(17-Hydroxy-9,12,15-octadecatrienyl)glutamine, H-645
- 3-Hydroxy-2-octadecylindole, H-646
- 8-Hydroxy-2,4,6-octatriynamide, H-647
- 8-Hydroxy-2,4,6-octatriynoic acid, H-647
- 4-Hydroxy-2-(1-octenyl)quinoline, H-648
- 4-Hydroxy-2-octylquinoline *N*-oxide, H-648
- 4-Hydroxy-2-octylquinoline, H-648
- 3-Hydroxy-2-octyl-4(1*H*)-quinolinone, A-621
- 6-Hydroxyonychine, O-93
- 2-Hydroxyonychine, O-93
- N*-Hydroxyovigerine, O-143
- 8-Hydroxy-12,28-oxamanzamine A, M-91
- 6-Hydroxy-12,34-oxamanzamine E, M-91
- 3-Hydroxy-2-(4-oxazolylmethylene)butanal, O-156
- 4-Hydroxy-1-(4-oxazolyl)-1-penten-3-one, M-159
- 6-(2-Hydroxy-2-oxido-4*H*-1,3,2-dioxaphosphorin-6-yl)-2,4(1*H*,3*H*)-pteridinedione, H-649
- 4-Hydroxyoxindole-3-acetic acid, D-452
- 5-Hydroxyoxindole-3-acetic acid, D-453
- 4-Hydroxyoxindole-3-acetonitrile, D-452
- 5-Hydroxyoxindole, I-92
- 3-Hydroxyoxindole, I-91
- 12*R*-Hydroxy-11-oxoaerthionin, A-158
- 12*S*-Hydroxy-11-oxoaerthionin, A-158
- 8-Hydroxy-1-oxo-2-azaspiro[4.5]deca-6,9-diene-3-carboxylic acid, S-424
- 4-Hydroxy-α-oxobenzeneacetic acid, H-687
- N*-(3-Hydroxy-2-oxobutyl)cytisine, S-378
- 3-(1-Hydroxy-3-oxobutyl)-4,7,8-trimethoxy-2(1*H*)-quinolinone, E-314
- 18-Hydroxy-3-oxo-1,4,14-conatrienine, H-314
- 5-Hydroxy-6-oxocoronaridine, C-657
- 19-Hydroxy-3-oxocoronaridine, C-657
- 1-Hydroxy-4-oxo-2,5-cyclohexadiene-1-acetamide, H-650
- 1-Hydroxy-4-oxo-2,5-cyclohexadiene-1-acetic acid, H-650
- N*-(3-Hydroxy-1-oxocyclopent-2-en-2-yl)-3-(4-hydroxy-3-methoxyphenyl)propenamide, H-651
- o-(2-Hydroxy-4-oxodecanoyl)histidine, H-307
- 3-Hydroxy-2-oxodendrobine, D-210
- 1-(3-Hydroxy-1-oxodecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 8-Hydroxy-3-oxo-1,7(11)-eremophiladien-12,8-lactam, H-652
- 8-Hydroxy-1-oxo-7(11)-eremophilene-12,8-lactam, H-653
- 10-Hydroxy-11-oxoerysotrine, E-203
- 14-Hydroxy-19-oxogelsenicine, G-44
- 1-(3-Hydroxy-1-oxohexadecyl)-2-(3-pyridinyl)pyrrolidine, P-957
- 14-Hydroxy-19-oxohumantenmine, G-44
- 6-Hydroxy-α-oxo-1*H*-indole-3-acetic acid, H-654
- 2-Hydroxy-17-oxo-β-isopartine, I-322
- 25-Hydroxy-24-oxokibdelone C, H-655
- 13-Hydroxy-17-oxolupanine, O-188
- 15-Hydroxy-17-oxolupanine, O-188
- N*-(5-Hydroxy-2-oxo-7-oxabicyclo[4.1.0]hept-3-en-3-yl)acetamide, A-804
- 4-(4-Hydroxy-3-oxo-1-pentenyl)oxazole, M-159
- α-(3-Hydroxy-5-oxo-4-phenyl-2(5*H*)-furanlydene)-benzeneacetic acid, P-790
- 4-Hydroxy-2-oxo-6-(phenylmethyl)-1*H*-pyridine-3-carboxaldehyde, C-142
- 7-Hydroxy-Δ¹-3-(2-oxopropyl)coronaridine, O-199
- N*-[2-(2-Hydroxy-1-oxopropyl)phenyl]acetamide, A-32
- 3-Hydroxy-9-oxo-9,10-seco-1,3,5-conatriene, D-383
- 7-Hydroxy-3-oxo-3,7-secorhynchophylline, H-656
- 4-Hydroxy-2-oxosparteine, H-551
- 13-Hydroxy-2-oxosparteine, H-554
- 13α-Hydroxy-4-oxosparteine, T-361
- 13-Hydroxy-10-oxosparteine, V-153
- 2-Hydroxy-5-oxo-4-[[4,6,8-trimethyl-1-oxo-2,7-nonadienyl]amino]-7-oxabicyclo[4.1.0]hept-3-ene-2-pentanoic acid, A-1298
- 20*S*-Hydroxy-7-oxovincadifformine, M-631
- 14β-Hydroxyoxymatrine, M-121
- 12β-Hydroxypachysandrine A, D-263
- 16β-Hydroxypachysandrine A, D-264
- 19-Hydroxypachysiphine, H-743
- 8-Hydroxypalmatine, O-213
- 7α-Hydroxyparavallarine, P-93
- 7β-Hydroxyparavallarine, P-93
- 11α-Hydroxyparavallarine, P-93
- 10-Hydroxy-*cis*-paspalic acid amide, P-107
- 10-Hydroxy-*trans*-paspalic acid amide, P-107
- 14-Hydroxypaspalinine, P-109
- N*-Hydroxypaxdaphnine B, P-139
- 14α-Hydroxypaxilline, P-140
- 7-Hydroxypeganine, V-37
- 6-HydroxyPELLITORINE, D-106
- 3-(1-Hydroxypentadecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
- 4-Hydroxy-2-pentadecylquinoline, H-657
- 4-Hydroxy-2-(1,3-pentadienyl)piperidine, H-658
- 14-Hydroxy-2,3,4,6,7-pentamethoxyphenanthroindolizidine, P-228
- 3-Hydroxy-4-pentenenitrile, H-659
- 3-Hydroxy-4-pentenoic acid, H-659
- 4-Hydroxy-2-(1*E*-pentenyl)quinoline, H-660
- 4-Hydroxy-2-(1*Z*-pentenyl)quinoline, H-660
- 2-(2-Hydroxypentyl)-*N*-methylpiperidine, H-37
- 2-(2-Hydroxypentyl)-6-(2-oxopropyl)piperidine, A-1000
- 2-(2-Hydroxypentyl)piperidine, H-37
- 5-Hydroxy-2-pentylpiperidine, P-236
- 4-Hydroxy-2-pentylquinoline *N*-oxide, H-660
- 4-Hydroxy-2-pentylquinoline, H-660

- 3-Hydroxy-2-pentyl-4(1*H*)-quinolinone, A-621
 10-Hydroxyperegrine, M-322
 10-Hydroxypericyclivine, S-89
p-Hydroxyphenacylamine, A-786
 1-Hydroxyphenazine 10-oxide, P-327
 2-Hydroxy-1-phenazincarboxylic acid, H-661
 4-Hydroxy-1-phenazincarboxylic acid, H-662
 6-Hydroxy-1-phenazincarboxylic acid, H-663
 9-Hydroxy-1-phenazincarboxylic acid, H-664
 4-Hydroxy-1,6-phenazinedicarboxylic acid, H-666
 2-Hydroxy-1,6-phenazinedicarboxylic acid, H-665
 1-Hydroxyphenazine, P-327
 2-Hydroxyphenazine, P-328
p-Hydroxyphenethyl alcohol, H-676
N-(*p*-Hydroxyphenethyl)acetamide, H-667
N-(*p*-Hydroxyphenethyl)acetidine, A-125
 4-Hydroxyphenethylamine, H-667
 3-Hydroxyphenethylamine, H-678
 4-[[*p*-Hydroxyphenethyl]amino]methyl]pyrocatechol, N-288
N-(4-Hydroxyphenethyl)-2,4-decadienamide, H-668
N-(4-Hydroxyphenethyl)-2,4-dodecadienamide, H-669
N-(4-Hydroxyphenethyl)-2,4-tetradecadienamide, H-670
p-Hydroxyphenopyrrozin, P-333
N-(2-Hydroxyphenyl)acetamide, A-861
p-Hydroxyphenylacetamide, H-672
 2-Hydroxy-2-phenylacetamide, H-671
 4-Hydroxyphenylacetic acid, H-672
 4-Hydroxyphenylacetone, H-672
 2-Hydroxy-2-phenylacetone, H-673
 3-(4-Hydroxyphenylacetoxy)tropane, M-393
N-[2-(4-Hydroxyphenyl)acetyl]formamide, H-672
p-Hydroxyphenylacrylic acid, H-691
 3-*p*-Hydroxyphenyl- α -alanine, T-700
N-(4-Hydroxyphenyl)arachidonamide, E-48
 8-Hydroxy-2-phenyl-1-benzazocine, P-340
 5-Hydroxy-7-phenyl-1*H*-benz[*de*]isoquinoline-1,6(2*H*)-dione, L-5
 1-(4-Hydroxyphenyl)-3*H*-benzo[*a*]pyrrolo[2,3-*c*]phenazine, H-674
 7-Hydroxy-2-phenyl-4*H*-3,1-benzoxazin-4-one, H-675
 6-(4-Hydroxyphenyl)-2,8-bis(phenylmethyl)imidazo[1,2-*a*]pyrazin-3(7*H*)-one, C-560
 2-[4-(4-Hydroxyphenyl)-1,3-butadienyl]-4*H*-3,1-benzoxazin-4-one, A-1562
 4-Hydroxyphenyl-di-3-indolylmethane, D-692
 2-(4-Hydroxyphenyl)-6-(3,4-dimethoxyphenyl)morpholine, C-363
N-(4-Hydroxyphenyl)-5,8,11,14-eicosatetraenamide, E-48
 2-(4-Hydroxyphenyl)ethanol, H-676
 2-(4-Hydroxyphenyl)ethanamine, H-677
 19-(4-Hydroxyphenyl)-10*H*-1,10-ethenochromeno[2,3-*b*]indole-6,8,18-triol, N-327
 2-[2-(4-Hydroxyphenyl)ethenyl]-1-methylpiperidine, C-220
 2-Hydroxy-*N*-(2-phenylethyl)propanamide, X-27
 2-(4-Hydroxyphenyl)ethyl carbamate, H-676
N-(*p*-Hydroxyphenyl)ethyl *p*-hydroxycinnamide, H-667
N-[2-(4-Hydroxyphenyl)ethyl]acetamide, H-667
 2-Hydroxy-2-phenylethylamine, A-864
 2-(3-Hydroxyphenyl)ethylamine, H-678
 2-(*p*-Hydroxyphenyl)ethylamine, H-667
 4-[[2-(4-Hydroxyphenyl)ethyl]amino]methyl]-1,2-benzenediol, N-288
N-[2-(4-Hydroxyphenyl)ethyl]benzamide, H-667
N-[2-(4-Hydroxyphenyl)ethyl]cinnamide, H-667
N-[2-(4-Hydroxyphenyl)ethyl]dotriacontanamide, A-21
 9-Hydroxy-*N*-(2-phenylethyl)hexadecanamide, P-342
 5-[[2-(4-Hydroxyphenyl)ethyl]imino]methyl]-2-methoxyphenol, C-730
N-[2-(4-Hydroxyphenyl)ethyl]-2-methylbutanamide, H-667
N-[2-(4-Hydroxyphenyl)ethyl]-3-methyldodecanamide, H-679
N-[2-(4-Hydroxyphenyl)ethyl]-3-methyl-2-dodecanamide, H-679
 2-(2-Hydroxy-2-phenylethyl)-1-methyl-6-(2-oxobutyl)piperidine, H-347
 2-(2-Hydroxy-2-phenylethyl)-1-methyl-6-(2-oxopentyl)piperidine, D-403
 2-(2-Hydroxy-2-phenylethyl)-1-methylpiperidine, S-210
 1-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-2-butanone, H-347
 1-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-2-pentanone, D-403
 2-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-1-phenylethanone, L-219
 1-[6-(2-Hydroxy-2-phenylethyl)-1-methyl-2-piperidinyl]-2-propanone, S-215
 2-(2-Hydroxy-2-phenylethyl)-1-methylpyrrolidine, P-970
 1-(1-Hydroxy-2-phenylethyl)-9*H*-pyrido[3,4-*b*]indol-6-ol, E-290
N-[2-(4-Hydroxyphenyl)ethyl]tetracosanamide, H-667
N-[2-(4-Hydroxyphenyl)ethyl]tiglamide, H-667
N-(2-Hydroxyphenyl)formamide, A-861
 4-Hydroxyphenylglycolic acid, H-520
 3-Hydroxyphenylglycolic acid, H-519
p-Hydroxyphenylglyoxylic acid, H-687
 2-(2-Hydroxyphenyl)-4-(hydroxymethyl)thiazole, H-695
 6-(4-Hydroxyphenyl)-2-[[4-(hydroxyphenyl)methyl]-8-(phenylmethyl)imidazo[1,2-*a*]pyrazin-3(7*H*)-one, C-560
 4-Hydroxy-2-(phenylimino)-1(2*H*)-naphthalenone, A-1040
 1-(4-Hydroxyphenyl)-4-(1*H*-indol-3-yl)-2,3-butanediol, D-789
 2-(4-Hydroxyphenyl)-3-isobutylmaleimide, H-684
 1-(4-Hydroxyphenyl)-2-isocyanoethane, H-680
 Hydroxyphenylmalonic acid, H-688
 2'-*O*-[3-(4-Hydroxyphenyl)-2-methoxypropenyl]-3'-*O*-[3-(1*H*-imidazol-4-yl)propenyl]-5'-methyl-5'-thioadenosine, H-681
 1-(*p*-Hydroxyphenyl)-2-methylaminoethanol, S-659
 3-(4-Hydroxyphenyl)-2-(methylamino)propanoic acid, T-700
 [[4-(Hydroxyphenyl)methyl]carbamic acid, H-421
 [[4-(Hydroxyphenyl)methyl]carbamothioic acid, H-427
 3-(4-Hydroxyphenyl)-*N*-[4-(3-methyl-2,5-dioxo-1-pyrrolidinyl)butyl]-2-propenamide, H-682
 5-[[4-(Hydroxyphenyl)methylene]-2-thioxo-4-imidazolidinone, H-683
 6-(4-Hydroxyphenyl)-7-methyl-1,2,3,5,8,8a-hexahydroindolizine, I-165
 5-[[4-(Hydroxyphenyl)methyl]-2-(1*H*-indol-3-yl)-3-morpholinone, O-154
 2-[[[4-(Hydroxyphenyl)methyl]methylamino]-*N*-[2-(1*H*-indol-3-yl)ethenyl]-4-methylhexanamide, F-145
 2-(Hydroxyphenylmethyl)-8-methyl-8-azabicyclo[3.2.1]octane-3,6-diol, H-428
 2-[[4-(Hydroxyphenyl)methyl]-4-oxo-2-azetidinecarboxylic acid, H-425
 3-[[4-(Hydroxyphenyl)methyl]-1-phenyl-1*H*-pyrrole-2,5-dione, O-100
 3-(4-Hydroxyphenyl)-*N*-methylpropanamide, H-689
 3-(4-Hydroxyphenyl)-4-(2-methylpropyl)-1*H*-pyrrole-2,5-dione, H-684
 5-[[4-(Hydroxyphenyl)methyl]-2-pyridinecarboxylic acid, H-426
 5-(Hydroxyphenylmethyl)-2-pyridinecarboxylic acid, V-94
 6-Hydroxy- α -phenylmethyl-9*H*-pyrido[3,4-*b*]indole-1-methanol, E-290
 2-(4-Hydroxyphenyl)-1-methyl-4(1*H*)-quinolinone, H-685
 2-(4-Hydroxyphenyl)-5-(methylthio)thiazole, T-508
 1-Hydroxy-4-phenyl-2,7-naphthyridine, H-686
 2-(2-Hydroxyphenyl)-2-oxazoline-4-carboxylic acid, D-455
 2-(4-Hydroxyphenyl)-2-oxoacetic acid, H-687
 1-[5-(4-Hydroxyphenyl)-1-oxo-2,4-pentadienyl]piperidine, C-710
 1-[3-(4-Hydroxyphenyl)-1-oxopropyl]-1*H*-pyrrole, P-351
 5-(4-Hydroxyphenyl)-2,4-pentadienoic acid piperidide, C-710
N-[5-(4-Hydroxyphenyl)-2,4-pentadienyl]piperidine, C-710
 3-(4-Hydroxyphenyl)-4-phenyl-1*H*-pyrrole-2,5-dicarboxylic acid, B-188
 2-Hydroxy-2-phenylpropanediamide, H-688
 Hydroxyphenylpropanedioic acid, H-688
 3-(4-Hydroxyphenyl)propanoic acid pyrrolidide, P-351
 3-(4-Hydroxyphenyl)propanoic acid, H-689
 3-(4-Hydroxyphenyl)propanoylethanol, C-419
N-[3-(4-Hydroxyphenyl)propanoyl]pyrrole, P-351
 3-(3-Hydroxyphenyl)-2-propenoic acid, H-690
 3-(4-Hydroxyphenyl)-2-propenoic acid, H-691
 3-(*p*-Hydroxyphenyl)propionic acid, H-689
 3 α -(2-Hydroxy-3-phenylpropionyloxy)tropane, L-216
 1-(4-Hydroxyphenyl)-2-propylamine, A-898
 3-(*p*-Hydroxyphenyl)propylamine, H-350
 5-(4-Hydroxyphenyl)-2-(3-pyridyl)oxazole, H-12
 3-(2-Hydroxyphenyl)-4(3*H*)-quinazolinone, Q-14
 4-Hydroxy-2-phenylquinoline, H-693
 3-(4-Hydroxyphenyl)quinoline, H-692
 4-Hydroxy-3-phenyl-2(1*H*)-quinolinone, D-657
 5-Hydroxy-2-phenyl-4(1*H*)-quinolinone, D-658
 6-Hydroxy-2-phenyl-4(1*H*)-quinolinone, D-659
 7-Hydroxy-2-phenyl-4(1*H*)-quinolinone, D-660
 3-Hydroxy-4-phenyl-2(1*H*)-quinolinone, D-656
 6-(*p*-Hydroxyphenyl)-8-(1-*D*-ribose)-2,4,7-trioxohexahydropteridine, H-232
 6-(4-Hydroxyphenyl)-8-*D*-ribose-2,4,7(1*H*,3*H*,8*H*)-pteridinetriene, H-232
 2-(2-Hydroxyphenyl)-4-thiazolecarboxaldehyde, H-695
 2-(2-Hydroxyphenyl)-4-thiazolecarboxylic acid, H-695
 2-(2-Hydroxyphenyl)thiazole, H-694
 2-(4-Hydroxyphenyl)-*N,N,N*-trimethylethenaminium, H-742
 [2-(4-Hydroxyphenyl)vinylamino]iminoacetic acid, H-44
 [2-(4-Hydroxyphenyl)vinyl]oxamic acid, H-44
 7-Hydroxyphlegmariurine B, P-363
 8 β -Hydroxyphlegmariurine B, P-363
 11 α -Hydroxyphlegmariurine B, P-363
 2 α -Hydroxyphlegmariurine B, P-363
 2-(Hydroxyphenyl)ethylamine, A-772
 4-Hydroxyphthalimidine, D-442
 5-Hydroxypicolinic acid, H-710
 3-Hydroxypicolinic acid, H-696
 4-Hydroxypicolinic acid, H-697
 5-Hydroxypicolinic acid, H-698
 4-Hydroxy-2-piperidinecarboxylic acid, H-697
 5-Hydroxy-2-piperidinecarboxylic acid, H-698
 3-Hydroxy-2-piperidinecarboxylic acid, H-696
 5-Hydroxy-2-piperidinemethanol, H-699
 4-Hydroxy- α -(1-piperidinyl)benzeneacetone, G-80
 3-[3-(3-Hydroxy-2-piperidinyl)-2-oxopropyl]-4(3*H*)-quinazolinone, F-31
N-Hydroxypiperolactam, A, D-567
 11-Hydroxypleiocarpamine, P-518
 7-Hydroxypleurocorine, H-700
 7-Hydroxypleurostyline, P-523
 6 α -Hydroxypowelline, P-581
 2-Hydroxy-6-prenylindole, H-701
 2-Hydroxy-3-prenyl-4-prenyloxyquinoline, A-1529
 4-Hydroxyproline betaine, H-476
 3-Hydroxyproline dimethylbetaine, H-740
 Hydroxyproline leucine anhydride, C-871
 4-Hydroxyproline, H-715
 2-(2-Hydroxypropanoyl)acetanilide, A-32
 2-[(2-Hydroxypropanoyl)amino]benzamide, A-706
 1-(3-Hydroxypropanoyl)- β -carboline, C-136
 6-(β -Hydroxypropionyl)-1,3-dimethylumazine, H-702
 6-(β -Hydroxypropionyl)-3-methylumazine, H-702
N-Hydroxypropylamine, P-648

- 1-(2-Hydroxypropyl)-3,4-dimethoxy-2-methyl-9H-carbazole, S-573
6-(1-Hydroxypropyl)-1,3-dimethylumazine, H-702
6-(1-Hydroxypropyl)lumazine, H-702
Hydroxypropyllycodine, H-703
7-(1-Hydroxypropyl)-8-methylindolizino[1,2-*b*]quinolin-9(11*H*)-one, M-96
6-(1-Hydroxypropyl)-8-methylisoxanthopterin, I-13
6-(1-Hydroxypropyl)-1-methylumazine, H-702
6-(1-Hydroxypropyl)-3-methylumazine, H-702
1-(2-Hydroxypropyl)-2-methyl-6-(3-methyl-2-but-2-enyl)-3*H*-carbazole-3,4(9*H*)-dione, C-169
2-(2-Hydroxypropyl)-1-methyl-5-(2-oxopropyl)pyrrolidine, H-704
2-(2-Hydroxypropyl)-6-methylpiperidine, H-705
2-(2-Hydroxypropyl)-1-methylpyrrolidine, H-760
1-[5-(2-Hydroxypropyl)-1-methyl-2-pyrrolidinyl]-2-propanone, H-704
2-(1-Hydroxypropyl)piperidine, C-603
5-Hydroxy-2-propylpiperidine, P-653
2-(2-Hydroxypropyl)piperidine, S-216
6-(1-Hydroxypropyl)-2,4(1*H*,3*H*)-pteridinedione, H-702
2-(3-Hydroxypropyl)-4(3*H*)-quinazolinone, H-706
4-Hydroxy-2-propylquinoline, H-707
17-Hydroxypseudoakumigmine, A-232
8-Hydroxypseudocoptisine, P-710
8*b*-Hydroxyptilocalaunin, P-773
6-Hydroxypurine, P-820
2-Hydroxyputrescine, D-248
2-Hydroxypyrazine, P-889
6-Hydroxy-2-pyridineacetic acid, H-708
5-Hydroxy-2-pyridinecarboxylic acid, H-710
4-Hydroxy-3-pyridinecarboxylic acid, H-709
6-Hydroxy-3-pyridinecarboxylic acid, H-711
5-Hydroxy-2-pyridinemethanol, H-515
3-Hydroxypyridine, P-906
4-Hydroxy-2-[4-(2-pyridinyl)-1,3-butadienyl]benzoic acid, H-712
N-[5-Hydroxy-2-pyridinyl]methyladenosine, H-713
6-(5-Hydroxy-2-pyridinylmethylamino)-9*β*-ribofuranosylpurine, H-713
2-Hydroxy-3-(4-pyridinyl)-2-propenoic acid, O-205
4-Hydroxy-9*H*-pyrido[3,4-*b*]indole-1-carboxaldehyde, H-456
4-Hydroxy-9*H*-pyrido[3,4-*b*]indole-1-carboxylic acid, H-457
6-Hydroxy-9*H*-pyrido[3,4-*b*]indole-1-carboxylic acid, H-458
7-Hydroxy-9*H*-pyrido[3,4-*b*]indole-1-carboxylic acid, H-459
1-(4-Hydroxy-9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, A-53
1-(6-Hydroxy-9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, A-54
1-(7-Hydroxy-9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, A-55
1-(8-Hydroxy-9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, A-56
3-Hydroxy-1-(9*H*-pyrido[3,4-*b*]indol-1-yl)-1-propanone, C-136
4-Hydroxypyrido[4,3,2-*mm*]pyrrolo[3,2,1-*de*]acridine, A-1445
β-(5-Hydroxy-2-pyridyl)alanine, A-822
4-Hydroxy-11-(1-(2-pyrrolocarbonyl)epilupine, H-482
10-Hydroxy-13-(2-pyrrolocarbonyloxy)lupanine, D-608
4-Hydroxy-1*H*-pyrrole-2-carboxylic acid, H-714
2-Hydroxy-1-pyrrolidinedecarboxaldehyde, P-955
4-Hydroxy-2-pyrrolidinedecarboxylic acid, H-715
4-Hydroxy-3-pyrrolidinedecarboxylic acid, H-716
3-Hydroxy-2-pyrrolidinemethanol, H-717
2-Hydroxypyrrrolidine, P-955
4-Hydroxy-2-pyrrolidinone, H-718
5-Hydroxy-2-pyrrolidinone, H-719
4-Hydroxy-2-pyrrolidone, H-718
5-Hydroxy-2-pyrrolidone, H-719
6-Hydroxy-1-(1-(pyrrolin-2-yl)-*β*-carbolone, E-285
6-Hydroxy-1-pyrrolidinedecarboxylic acid, H-720
6-Hydroxy-1-(2-pyrrolyl)-*β*-carbolone, E-285
1-(3-Hydroxy-1*H*-pyrrol-2-yl)ethanone, A-58
14*β*-Hydroxyquebrachamine, Q-6
6-Hydroxyquestionimycin A, A-862
3-Hydroxyquinaldic acid, H-724
4-Hydroxyquinaldine, H-633
α-Hydroxyquinaldine, Q-37
4-Hydroxyquinaldinic acid, H-725
4-Hydroxyquinazoline, Q-14
9-Hydroxy-10*H*-quindoline, I-112
3-Hydroxyquinine, Q-20
8-Hydroxy-4-quinolinecarboxaldehyde, H-722
2-Hydroxy-4-quinolinecarboxaldehyde, H-721
6-Hydroxy-4-quinolinecarboxamide, H-726
3-Hydroxy-2-quinolinecarboxylic acid, H-724
4-Hydroxy-2-quinolinecarboxylic acid, H-725
6-Hydroxy-4-quinolinecarboxylic acid, H-726
8-Hydroxy-2-quinolinecarboxylic acid, H-727
2-Hydroxy-4-quinolinecarboxylic acid, H-723
4-Hydroxy-2,3-quinolinedicarboxylic acid, H-728
2-Hydroxyquinoline, Q-50
3-Hydroxyquinoline, Q-49
8-Hydroxy-2(1*H*)-quinolinone-4-carboxylic acid, D-673
3-Hydroxy-4(1*H*)-quinolinone-2-carboxylic acid, D-674
4-Hydroxy-2(1*H*)-quinolinone, Q-33
8-Hydroxy-4(1*H*)-quinolinone, Q-35
2-Hydroxy-4(1*H*)-quinolinone, Q-33
10-Hydroxyranacoinine, A-94
11-Hydroxyrankinidine, H-367
6*α*-Hydroxyraumacline, R-22
14*α*-Hydroxyraunitic acid, A-212
4-Hydroxy-*N*-resorcylanthranilic acid, B-67
12*S*-Hydroxyretroisosenine, N-89
13*R*-Hydroxyretroisosenine, N-89
12-Hydroxyretulinal, R-63
13*β*-Hydroxyribasine, R-94
5-Hydroxy-1-*β*-*D*-ribofuranosyl-1*H*-imidazole-4-carboxaldehyde, H-729
N-(6-Hydroxy-9-*β*-*D*-ribofuranosyl-9*H*-purin-2-yl)alanine, H-730
4-Hydroxy-3-ribofuranosyl-1*H*-pyrazole-5-carboxamide, P-891
N-Hydroxyrobustine, S-456
4-Hydroxyroemine, S-549
Hydroxyroxburghiline, O-67
1-Hydroxyrutacidone epoxide, R-167
20-Hydroxyrutacidone epoxide, R-167
7-Hydroxyrutaecarpine, R-168
1-Hydroxyrutaecarpine, R-168
11-Hydroxyrutaecarpine, R-168
3-Hydroxyrutaecarpine, R-168
7-Hydroxyrosenamine, H-187
21-Hydroxysaframycin B, S-8
4-Hydroxy-*N*-salicylanthranilic acid, B-67
2-Hydroxysalignarine E, D-275
8-Hydroxysampangine, S-49
12-Hydroxysanguinarine, C-360
Hydroxy-*α*-sanshool, D-893
Hydroxy-*β*-sanshool, D-893
Hydroxy-*γ*-sanshool, T-127
4-Hydroxysarcocapnine, S-73
4-Hydroxysarcocapnine, S-73
3-Hydroxysarpagine, A-27
11-Hydroxysaxitoxin, H-731
N'-Hydroxysaxitoxin, S-109
10-Hydroxyscandine, S-118
α-Hydroxyscopalamine, A-1043
3-Hydroxy-9,10-seco-1,3,5,8(14)-conatetraen-9-one, D-383
3-Hydroxy-9,10-seco-1,3,5-conatrien-9-one, D-383
3-Hydroxy-3,4-secoconaridine, H-732
3-Hydroxy-16,28-secosolanid-22(28)-ene-6,23-dione, P-281
3-Hydroxy-16,28-secosolanid-22(28)-en-6-one, H-483
12-Hydroxy-12,24-secostrychnidin-10-one, P-672
4-Hydroxysecurinine, P-386
4-Hydroxysedamine, S-210
5-Hydroxysedamine, S-210
Hydroxysedolin, S-215
Hydroxysedione, S-215
9-Hydroxysenbusine A, M-310
12-Hydroxysenecionan-11,16-dione, S-238
3-(*trans*-4-Hydroxyseneciyl)-6-seneciolytropene, M-391
Hydroxysenkirkine, S-245
18-Hydroxysenkirkine, S-245
13*α*-Hydroxyseneciopyne, S-248
4-Hydroxyserratinine, A-1029
6*α*-Hydroxyserratinine, A-1029
Hydroxyskytanthine I, H-735
Hydroxyskytanthine II, H-736
4-Hydroxyskytanthine, H-735
4*α*-Hydroxyskytanthine, H-733
7-Hydroxyskytanthine, H-736
7*α*-Hydroxyskytanthine, H-734
6-Hydroxyskytanthine, I-62
4-Hydroxysmirnovine, S-406
15*α*-Hydroxysoladulcine, S-441
15*β*-Hydroxysoladulcine, S-441
2-Hydroxysoladulcine, S-440
23*R*-Hydroxysoladulcine, S-442
12*α*-Hydroxysolamargine, S-447
12*β*-Hydroxysolamargine, S-447
18-Hydroxysolanid-4-en-3-one, V-62
23-Hydroxysolanthrene, L-116
N-Hydroxysolasodine, S-456
15*α*-Hydroxysolasodine, S-448
15*β*-Hydroxysolasodine, S-448
23*S*-Hydroxysolasodine, S-450
12*α*-Hydroxysolasonine, S-447
12*β*-Hydroxysolasonine, S-447
9-Hydroxysophocarpine *N*-oxide, S-371
5-Hydroxysophocarpine, S-371
9*α*-Hydroxysophocarpine, S-371
12*α*-Hydroxysophocarpine, S-371
12*β*-Hydroxysophocarpine, S-371
9*α*-Hydroxysophoramine *N*-oxide, S-376
7*α*-Hydroxysophoramine, S-376
9*α*-Hydroxysophoramine, S-376
14*β*-Hydroxysophoridine, S-380
3*α*-Hydroxysophoridine, S-380
4-Hydroxysparteine, H-737
7-Hydroxysparteine, H-738
14-Hydroxysparteine, H-739
12*α*-Hydroxysparteine, R-56
13-Hydroxysparteine, T-361
7-Hydroxyspeciociliatine, M-657
7-Hydroxy-(-)-spectaline, S-388
23-Hydroxyspermostrychnine *N*⁴-oxide, S-400
23-Hydroxyspermostrychnine, S-400
12-Hydroxysqualamine, S-485
3-Hydroxystachydrine a, H-740
3-Hydroxystachydrine b, H-740
3-Hydroxystachydrine, H-740
5'-Hydroxystaurosporine, S-498
7-Hydroxystaurosporine, S-498
11-Hydroxystaurosporine, S-498
3-Hydroxystaurosporine, S-498
2-Hydroxystaurosporinone, A-1190
6*S*-Hydroxystemofoline, S-515
16-Hydroxystemofoline, S-515
2'*R*-Hydroxystemofoline, S-515
2'*S*-Hydroxystemofoline, S-515
3'*S*-Hydroxystemofoline, S-515
4'-Hydroxystemofoline, S-515
8-Hydroxystephananthrine *N*-oxide, S-546
8-Hydroxystephananthrine, S-546
9-Hydroxystreptazolin, S-560
13-Hydroxystreptazolin, S-560
8-Hydroxystreptazolone, H-741
10-Hydroxystreptamine, S-576
11-Hydroxystreptamine, S-576
12-Hydroxystreptamine, S-576
18-Hydroxystreptamine, S-576
10-Hydroxystreptosamide, V-126
16-Hydroxystrychnidin-10-one, P-726
12-Hydroxystrychnine *N*^b-oxide, S-589
3-Hydroxystrychnine, P-726
10-Hydroxystrychnine, S-589
12-Hydroxystrychnine, S-589
15-Hydroxystrychnine, S-589
12'-Hydroxystrychnobiline, S-591
14-Hydroxystrychnobrasiline, S-594

- 13-Hydroxystylopine, S-612
 4-Hydroxystyrylamine, H-677
 (4-Hydroxystyryl)trimethylammonium(1+), H-742
 Hydroxysuccinic acid, M-72
 N^2 -(2-Hydroxysuccinoyl)arginine, A-1413
 N^2 -(3-Hydroxysuccinoyl)arginine, A-1413
 18-Hydroxysungucine, S-642
 21-Hydroxyscophantine, S-449
 19'-*R*-Hydroxytabernaeanegantane A, C-616
 3'-Hydroxytabernaeanegantane B, C-615
 19'-*R*-Hydroxytabernamine, T-8
 19'-*S*-Hydroxytabernamine, T-8
 3'-Hydroxytabernamine, T-8
 3-Hydroxytabernanthine, I-4
 19-Hydroxytabersonine, H-743
 11-Hydroxytabersonine, T-11
 16-Hydroxytabersonine, T-11
 19-Hydroxytacamine, T-12
 17 β -Hydroxytacamonine, T-13
 6 β -Hydroxytalatizamine, M-323
 10-Hydroxytalatizamine, M-326
 3 α -Hydroxytalatizamine, M-317
 2'-Hydroxytaxine II, T-273
 Hydroxytaxodine, H-331
 4-Hydroxytecomanine, T-58
 5'-Hydroxytelobine, A-1331
 12-Hydroxy-2,4,8,10-tetradecatetraenoic acid, H-744
 11-Hydroxy-2,4,8-tetradecatrenoic acid, H-745
 3-(1-Hydroxytetradecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
 3-(1-Hydroxytetradecylidene)-1-methyl-2,4-pyrrolidinedione, M-191
 6-(13-Hydroxytetradecyl)-2-methyl-3-piperidinol, S-388
 4-Hydroxy-2-tetradecylquinoline, H-746
 4-Hydroxy-3-[1-(tetrahydro-5-hydroxy-2,6,6-trimethyl-2*H*-pyran-2-yl)ethyl]-2(1*H*)-quinolinone, B-380
 6-Hydroxy-1,2,3,4-tetrahydroisoquinoline, T-196
 7-Hydroxy-1,2,3,4-tetrahydroisoquinoline, T-197
 8-Hydroxy-1,2,3,4-tetrahydroisoquinoline, T-198
 5-[Hydroxy-(5,6,7,8-tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)methyl]-1,3-benzodioxole-4-carboxylic acid, B-118
 3-Hydroxytetrahydrodroilvacine, G-211
 Hydroxytetrahydrohombifoline, T-149
 16-Hydroxytetrahydrosecamine, S-174
 10'-Hydroxytetrahydro-17*R*-usambarensine, U-52
 10'-Hydroxytetrahydro-17*S*-usambarensine, U-52
 4-Hydroxy-6-(1,7,9,10-tetrahydroxybenz[*g*]isoquinolin-3-yl)-2*H*-pyran-2-one, A-1313
 1-Hydroxy-2,3,5,6-tetramethoxyacridone, P-206
 4-Hydroxy-1,2,9,10-tetramethoxyaporphine, C-190
 9-Hydroxy-1,2,3,10-tetramethoxyaporphine, P-208
 1-Hydroxy-2,3,10,11-tetramethoxyaporphine, P-209
 1-Hydroxy-2,3,9,10-tetramethoxyaporphine, P-208
 11-Hydroxy-1,2,3,10-tetramethoxyaporphine, P-209
 2-Hydroxy-1,8,9,10-tetramethoxyaporphine, P-210
 3-Hydroxy-1,2,9,10-tetramethoxyaporphine, P-208
 1-Hydroxy-2,3,5,6-tetramethoxy-10-methylacridone, P-206
 1-Hydroxy-2,3,9,10-tetramethoxynoraporphine, P-208
 11-Hydroxy-1,2,3,10-tetramethoxynoraporphine, P-209
 3-Hydroxy-1,2,9,10-tetramethoxynoraporphine, P-208
 14-Hydroxy-3,4,6,7-tetramethoxyphenanthroindolizidine, I-342
 10-Hydroxy-1,4,6,9-tetramethylpyrido[3,2-*g*]quinoline-2,8(1*H*,9*H*)-dione, H-747
 11-Hydroxytetradotoxin, T-298
 5-Hydroxythalidasine 2 α -*N*-oxide, T-320
 5-Hydroxythalidasine, T-320
 5'-Hydroxythalifaboramine, T-322
 3-Hydroxythalifaboramine, T-322
 5-Hydroxythalamine, T-341
 16-Hydroxythebaine, T-350
 13-Hydroxythermopsine, B-20
 2-(4-Hydroxy-2-thiazolyl)-6-benzothiazolol, F-56
 syn-6'-Hydroxythiobinupharidine sulfoxide, T-380
 syn-6-Hydroxythiobinupharidine sulfoxide, T-380
 6'-Hydroxythiobinupharidine, T-380
 6-Hydroxythiobinupharidine, T-380
 6-Hydroxythionupharidine B, T-380
 3-Hydroxy-13-tigloyloxylylupanine, D-605
 13-(4-Hydroxytigloyloxy)lylupanine, H-554
 6-Hydroxy-7-tigloyloxynortropan-3-yl 2-hydroxy-3-phenylpropionate, M-392
 5-Hydroxytiliacorinine, Y-5
p-Hydroxy- α -toluic acid, H-672
 15 α -Hydroxytomatidenol, S-448
 15 α -Hydroxytomatidine, S-441
 7-Hydroxy-2,8-tridecadiene-10,12-dienoic acid, H-748
 4-Hydroxy-2-(1*E*-tridecenyl)quinoline, H-750
 4-Hydroxy-2-(1*Z*-tridecenyl)quinoline, H-750
 3-(1-Hydroxytridecylidene)-1,5-dimethyl-2,4-pyrrolidinedione, M-191
 2-(12-Hydroxytridecyl)-3-methoxy-4(1*H*)-quinolinone, H-749
 4-Hydroxy-2-tridecylquinoline, H-750
 1-Hydroxy-2,3,4-trimethoxyacridone, T-240
 1-Hydroxy-3,5,6-trimethoxyacridone, T-243
 1-Hydroxy-2,10,11-trimethoxyaporphine, C-682
 1-Hydroxy-2,9,10-trimethoxyaporphine, H-751
 10-Hydroxy-1,2,9-trimethoxyaporphine, H-752
 11-Hydroxy-1,2,10-trimethoxyaporphine, I-208
 9-Hydroxy-1,2,10-trimethoxyaporphine, L-75
 2-Hydroxy-1,9,10-trimethoxyaporphine, P-597
 10-Hydroxy-1,2,11-trimethoxyaporphine, T-248
 11-Hydroxy-1,2,3-trimethoxyaporphine, T-245
 2-Hydroxy-1,10,11-trimethoxyaporphine, T-248
 10-Hydroxy-4,5,6-trimethoxy-9*H*-azuleno[1,2,3-*ij*]isoquinolin-9-one, G-159
 10-Hydroxy-*O*¹⁷-(3,4,5-trimethoxybenzoyl)vincamajine, V-113
 10-Hydroxy-*O*¹⁷-(3,4,5-trimethoxycinnamoyl)vincamajine, V-113
 9-Hydroxy-1,2,3-trimethoxydibenz[*cd,f*]indol-4(5*H*)-one, T-255
 11-Hydroxy-1,2,10-trimethoxy-7*H*-dibenzo[*de,g*]quinolin-7-one, G-93
 9-Hydroxy-4,5,6-trimethoxyindeno[1,2,3-*ij*]isoquinoline, N-304
 1-Hydroxy-2,3,4-trimethoxy-10-methylacridone, T-240
 1-Hydroxy-2,3,5-trimethoxy-10-methylacridone, T-241
 1-Hydroxy-3,5,6-trimethoxy-10-methylacridone, T-243
 2-Hydroxy-3,8,9-trimethoxy-5-methylbenzo[*c*]phenanthridinium(1+), F-5
 1-Hydroxy-3,5,6-trimethoxy-10-methyl-4-(3-methyl-2-butenyl)-9(10*H*)-acridinone, P-616
 1-Hydroxy-3,5,6-trimethoxy-10-methyl-4-prenylacridone, P-616
 3-Hydroxy-2,5',6-trimethoxy-7'-methylspiro[2,5-cyclohexadiene-1,3(2'*H*)-quinoline]-2',4,4'(1'*H*)-trione, A-1469
 1-Hydroxy-2,10,11-trimethoxynoraporphine, C-682
 1-Hydroxy-2,9,10-trimethoxynoraporphine, H-751
 10-Hydroxy-1,2,9-trimethoxynoraporphine, H-752
 9-Hydroxy-1,2,10-trimethoxynoraporphine, L-75
 2-Hydroxy-1,9,10-trimethoxynoraporphine, P-597
 9-Hydroxy-1,2,3-trimethoxynoraporphine, T-244
 10-Hydroxy-1,2,11-trimethoxynoraporphine, T-248
 11-Hydroxy-1,2,3-trimethoxynoraporphine, T-245
 2-Hydroxy-1,10,11-trimethoxynoraporphine, T-248
 2-Hydroxy-3,8,9-trimethoxypavinane, A-1410
 3-Hydroxy-2,8,9-trimethoxypavinane, A-1410
 1-Hydroxy-2,8,9-trimethoxypavinane, P-514
 4-Hydroxy-3,6,7-trimethoxyphenanthroindolizidine, I-342
 β -Hydroxy-3,4,5-trimethoxyphenethylamine, H-557
 2-Hydroxy-4,6,8-trimethoxyquinoline, Q-42
 2-Hydroxy-4,7,8-trimethoxyquinoline, Q-43
 2-Hydroxy-3,9,10-trimethoxytetrahydroprotoberberine, I-210
 2-Hydroxy-*N,N,N*-trimethylethanaminium(1+), C-419
 4-Hydroxy-2-(2,4,6-trimethyl-1-heptenyl)quinazoline, T-606
 16-Hydroxy-4,4,14-trimethyl-3-methylamino-9,19-cyclopropan-20-one, C-905
 11-Hydroxy-4,8,12-trimethyl-13-(2-methyl-4-thiazolyl)-8,12-tridecadien-3-one, H-469
 3-Hydroxy-2,2,4-trimethyl-10-(5-oxazolyl)-4,6,8-decatrienamide, I-161
 13-Hydroxy-17,19,20-trinorcrotalanan-11,15-dione, D-363
 4-Hydroxy-1',3,4'-tris(4-hydroxyphenyl)-3'-[2-(4-hydroxyphenyl)ethyl]-7'-(sulfoxy)spiro[furan-2(5*H*),2'(3'*H*)-pyrrolo[2,3-*c*]carbazole]-5,5'(6'*H*)-dione, H-753
 3-Hydroxytrisphaeridine, M-448
 1-Hydroxytropacocaine, M-390
 3-Hydroxytropane-2-carboxylic acid, E-8
 3-Hydroxytropane, M-393
 1-Hydroxytropane, P-399
 6-Hydroxy-3-tropanone, M-391
 3-(2-Hydroxytropoyloxy)tropane, M-393
 6-Hydroxytryptamine, T-637
 5-Hydroxytryptamine, H-755
 2-Hydroxytryptamine, H-754
 5-Hydroxytryptophol, H-505
 11-Hydroxytubotaiwine, T-673
 19-Hydroxytubotaiwine, T-673
 13 α -Hydroxytylophorine, T-695
 14-Hydroxytylophorine, T-695
 Hydroxytyramine, D-920
 3-Hydroxytyrosine, A-745
 10-Hydroxy-2,4,6,8-undecatetraenoic acid, H-756
 4-Hydroxy-2-(1-undecenyl)quinoline, H-757
 4-Hydroxy-2-undecylquinoline, H-757
 3-Hydroxy-2-undecyl-4(1*H*)-quinolinone, A-621
 6-Hydroxyundulatin, C-743
 10'-Hydroxyusambarensine, U-52
 10-Hydroxyusambarine, O-21
 11-Hydroxyusambarine, O-21
 16 α -Hydroxyuscharin, U-53
 17 β -Hydroxyuscharin, U-53
O-(3-Hydroxyvalyl)griseoliteic acid, H-514
 19-Hydroxyvandrikine, M-666
 20-Hydroxyvandrikine, M-666
 7-Hydroxyvasicine, V-37
 19-Hydroxyvenalstonidine, V-49
 19 β -Hydroxyvenalstonine, V-49
 12 β -Hydroxyveramiline, V-73
 16-Hydroxyverazine, S-199
 25 β -Hydroxyverazine, V-73
 4 β -Hydroxyverazine, V-73
 6-Hydroxyvertine, V-95
 9-Hydroxyvertine, V-95
 11-Hydroxyvincadifformine, V-111
 15 β -Hydroxyvincadifformine, V-111
 19-Hydroxyvincamajine, V-113
 19-Hydroxyvincamine, V-115
 12-Hydroxyvincanine, N-301
 10-Hydroxyvincosamide, V-126
 10-Hydroxyvincoside lactam, V-126
 16 β -Hydroxy-19*R*-vindolinine *N*-oxide, V-132
 16 β -Hydroxy-19*S*-vindolinine *N*-oxide, V-132
 16 β -Hydroxy-19*R*-vindolinine, V-132
 16 β -Hydroxy-19*S*-vindolinine, V-132
 8-Hydroxy-1-vinyl- β -carboline, H-758
 11-Hydroxyvittatine, H-5
 3-Hydroxyvoacamine, V-168
 19-Hydroxyvoacangine, V-170
 3*S*-Hydroxyvoacangine, V-170
 3-Hydroxyvoachalotine, V-171
 3-Hydroxyvoafrine A, V-176
 3-Hydroxyvoafrine B, V-176
 19-Hydroxyvoalutaine, V-178
 17-Hydroxyvobasan-3-one, A-166
 3'-Hydroxyvobtusine, V-188
 4-Hydroxywilsonirine, C-190
 4'-Hydroxywithasomnine, D-492
 2-Hydroxyxanthiazone, X-3
 7*S*-Hydroxyxestospingine A, X-24
 5-Hydroxyyanangcorinine, Y-5
 18-Hydroxyyohimbine (incorr.), D-687

- 18-Hydroxy- α -yohimbine, D-687
 18-Hydroxyyohimbine, D-687
 2-Hydroxyzeatin, Z-11
 11-Hydroxyzoanthamine, Z-30
 26-Hydroxyzoanthamine, Z-30
 Hyellazole, M-270
 Hyeronimone, A-1325
 Hyeronine A, A-1325
 Hyeronine B, A-1325
 Hygric acid methylbetaine, S-491
 Hygrine, H-759
 Hygrolin, H-760
 (\pm)-Hygrolin, H-760
 Hygrophyllin, P-516
 Hylocerin, B-110
 Hymeniacidine, H-761
 Hymenialdisine, H-762
 (10*E*)-Hymenialdisine, H-762
 Hymenidine, O-123
 Hymenine, O-66
 Hymenocardine, H-763
 Hyoscine butyl bromide, S-159
 Hyoscine methobromide, S-159
 Hyoscine *N*-oxide 1, S-159
 Hyoscine, S-159
 Hyoscyamide, C-87
 Hyoscyamine hydrobromide, T-632
 Hyoscyamine *N*-oxide 1, T-632
 Hyoscyamine *N*-oxide 2, T-632
 Hyoscyamine sulfate, T-632
 Hyoscyamine, T-632
 Hyosol, S-159
 Hypaconine, M-343
 Hypaconitine, M-343
 Hypaphorine, H-764
 Hypecoleptopine, H-765
 Hypecorine, H-766
 Hypecorinine, H-766
 Hypecoumine, H-767
 Hypecumine, H-768
 Hyperaspine, H-769
 Hypercratine, H-770
 Hyperectine, H-771
 Hypodemapyrazine, H-772
 Hypodematine, P-340
 Hypoepistephanine, E-102
 Hypoestestatin 1, H-773
 Hypoestestatin 2, H-773
 Hypogallic acid, D-536
 Hypoglaunine A, P-249
 Hypoglaunine B, P-249
 Hypoglaunine C, P-249
 Hypoglaunine D, P-249
 Hypoglaunine, P-249
 Hypognavine, H-179
 Hypognavinol, H-179
 Hyponine A, P-247
 Hyponine B, P-247
 Hyponine C, P-247
 Hyponine D, P-247
 Hyponine E, P-247
 Hyponine F, P-247
 Hypotaurine, A-764
 Hypotaurocyamine, G-203
 Hypotaurocyanaminephosphoric acid, G-203
 Hypotuberoestemonine, H-774
 Hypoxanthine riboside, I-148
 Hypoxanthine, P-820
 Hypoxanthosine, I-148
 Hypserpine, H-775
 Hyrcamine, C-883
 Hyrcamine, H-776
 Hyrcanol, S-226
 Hyrcanone, H-777
 Hycatrienine, B-455
 Hyrtiazepine, H-778
 Hyrtinadine A, H-779
 Hyrtioerectine A, H-780
 Hyrtioerectine B, T-206
 Hyrtioerectine C, H-506
 Hyrtiomanzamine, H-781
 Hyrtiosin A, H-400
 Hyrtiosin B, H-782
 Hyrtiosulawesine, P-488
 Hyrudonine, H-305
 Hysodricanine A, H-783
 Hysodricanine B, H-784
 Hysodricanine C, H-785
 Hystrine, O-40
 Hystrixine, A-166
 I 639A, D-769
 I 639B, D-769
 IA 887, N-180
 IAA, I-81
 Ianthelline, I-1
 Ianthesine A, I-2
 Ianthesine B, I-2
 Ianthesine C, I-3
 Ianthesine D, I-2
 IB 01211, M-142
 IBA, I-83
 Ibogaine, I-4
 Ibogaline, I-4
 Ibogamine pseudoindoxyl, I-5
 Ibogamine, I-4
 Iboluteine, I-5
 Ibophyllidine *N*^b-oxide, I-6
 Ibophyllidine, I-6
 Iboquine, I-7
 Ibotenic acid, I-8
 Iboxygaine hydroxyindolenine, I-9
 Iboxygaine, I-4
 Iboxyphylline, I-10
 Ibukinamine, M-363
 IC 202A, A-1185
 IC 202B, A-1186
 IC 202C, A-1187
 Icaceine, I-11
 Icajine *N*-oxide, I-12
 Icajine, I-12
 Icar, P-961
 Ichthyopterin, I-13
 ICM 0201, A-1188
 2,4-Icosadienoic acid isobutylamide, E-45
 2,4-Icosadienoic acid pyrrolidide, T-491
 2,4-Icosadienoic acid, E-45
 4,5-Icosanediamine, E-47
 5,8,11,14-Icosatetraenoic acid, E-48
 8,11,14-Icosatrienoic acid, E-52
 2,4,14-Icosatrienoic acid, E-50
 2,4,16-Icosatrienoic acid, E-51
 2,4,8-Icosatrienoic acid, E-49
 3-Icosyl-1*H*-pyrrole-2-carboxaldehyde, A-623
 Icumazole A, I-14
 Icumazole B, I-14
 Icumazole F, I-15
 Ideamine A *N*-oxide, I-16
 Ideamine A, I-16
 Ideamine B *N*-oxide, P-97
 Ideamine B, P-97
 Idiospermuline, I-17
 Idrocilamide, H-488
 IFB Lactam 1, I-18
 Ifflaiaimine, I-19
 Iforrestine, I-20
 Ignavine, H-184
 Igzamide, I-21
 Iheyamine A, I-22
 Iheyamine B, I-23
 68-1147-II, A-1082
 2880-II, H-651
 Ikimine A, I-24
 Ikimine B, I-25
 Ikimine C, T-358
 Ikimine D, I-26
 Ilamine *N*-oxide, E-299
 Ilamine, E-299
 Ileabothoxazole, I-27
 Ilepicimide, M-444
 ILEVS 1080, J-9
 Illicolin H, I-28
 Illicifoline, O-202
 Ilidine, M-309
 Iliensine, M-309
 Illudinine, I-29
 Illukumbin A, I-30
 Illukumbin B, I-30
 Illukumbin, I-30
 IM 8443T, C-815
 Imbilin 1, I-31
 Imbilin 2, I-31
 Imbilin 3, I-31
 Imbricatine, I-32
 Imeluteine, I-33
 Imenine, I-34
 Imerubriner, G-159
 1*H*-Imidazol-4-amine, A-826
 1*H*-Imidazol-2-amine, A-825
 1*H*-Imidazole-1-acetic acid, I-36
 1*H*-Imidazole-4(5)-acetic acid, I-37
 4-Imidazoleacrylic acid, U-46
 3,3'-(1*H*-Imidazole-2,4-diyl)bis[6-bromo-1*H*-indole], N-312
 3,3'-(1*H*-Imidazole-2,4-diyl)bis-1*H*-indole, N-312
 1*H*-Imidazole-4-ethanamine, H-307
 1*H*-Imidazole-4(5)-ethanol, I-38
 Imidazole, I-35
 1*H*-Imidazole-4(5)-propanoic acid, I-39
 2,4-Imidazolidinedione, I-40
 2-(4-Imidazolyl)ethanol, I-38
N-(2-Imidazol-4-ylethyl)cinnamide, C-455
N-[2-(1*H*-Imidazol-4-yl)ethyl]-2,4-decadienamide, H-307
N-[2-(1*H*-Imidazol-4-yl)ethyl]-3-methylbutanamide, H-307
N-[2-(1*H*-Imidazol-4-yl)ethyl]-3-phenyl-2-propenamide, C-455
 3-(1*H*-Imidazol-4-yl)-*N*-[2-(1*H*-imidazol-4-yl)ethyl]propanamide, I-43
 3-(1*H*-Imidazol-4-yl)-4-(1*H*-indol-3-yl)-1*H*-pyrrole-2,5-dione, D-379
 3-(4-Imidazolyl)lactic acid, H-523
 4-(1*H*-Imidazol-2-ylmethyl)phenol, H-422
 2-[[3-(1*H*-Imidazol-4-yl)-1-oxo-2-propenyl]oxy]-*N,N,N*-trimethylethanaminium(1+), M-739
 3-(1*H*-Imidazol-4-yl)-2-propenoic acid, U-46
 5-[[3-(1*H*-Imidazol-4-yl)propenyl]amino]pentanoic acid, I-41
*N*⁶-[3-(1*H*-Imidazol-4-yl)propenyl]lysine, I-42
*N*²-[3-(1*H*-Imidazol-4-yl)propenyl]histamine, I-43
 3-(4-Imidazolyl)propionic acid, I-39
 2-(1*H*-Imidazol-4-yl)quinoline, I-44
 1*H*-Imidazo[4,5-*b*]pyrazin-2-amine, A-829
 1*H*-Imidazo[4,5-*a*]pyrrolo[3,4-*c*]carbazole-4,6(5*H*,11*H*)-dione, G-169
 Imido, H-307
 Iminazole, I-35
 2,2'-Iminobisethanol, D-388
 2,2'-Iminobis[4-hydroxybenzoic acid], I-45
 3,3'-[Iminobis(methylene)]bis-2(3*H*)furanone, I-46
 3,3'-Iminobis[5-methyl-2(5*H*)-furanone], P-190
 Iminobispropylamine, D-255
 22,26-Iminocholesta-5,22(*N*)-dien-3-ol, V-73
 22,26-Iminocholest-5-en-3-amine, S-66
 22,26-Iminocholest-22(*N*)-ene-3,6-diol, S-201
 5,5'-(Iminodi-4-butanyl-1-ylidene)bis[3-methyl-2(5*H*)-furanone], P-65
 2,2'-Iminodiethanol, D-388
 23,29-Imino-*B*(9*a*)-homo-19-norstigmastan-1(10),7,9(11),23(*N*)-tetraen-3-amine, I-47
 23,29-Imino-*B*(9*a*)-homo-19-norstigmastan-1(10),7,23(*N*)-trien-3-amine, I-47
 2-Imino-1-imidazolidineacetic acid, A-731
 2-Imino-5-(1*H*-indol-3-ylmethylene)-1,3-dimethyl-4-imidazolidinone, A-1357
 2-Iminomethyl-3-methyl-6-methylamino-9*H*-purine, I-48
 5-Imino-1-methylproline, I-50
 5-Imino-1-methyl-2-pyrrolidinecarboxylic acid, I-50
 18,20-Imino-19-norpregna-1,3,5(10),14-tetraen-3-ol, D-788
 2,5-Imino-1,3,4,6,7-octanepentol, D-665
 2-[[Imino(phosphonoamino)methyl]amino]ethanesulfonic acid, G-203
 2-Iminopiperidine, I-49
 5-Iminoprolin, I-50
 Iminopropionicacetic acid, C-147

- 5-Imino-2-pyrrolidinecarboxylic acid, I-50
 5-Imino-2*H*-pyrrol-2-ol, I-51
 8-Iminosanguinarine, D-510
 1,5-Imino-1,5,6-trideoxy-D-glucitol, M-517
 2,5-Imino-2,5,6-trideoxyheptitol, H-492
 1,5-Imino-1,5,6-trideoxyhexitol, M-517
 Iminourea, G-199
 Immunosuppressant GT 32B, A-1127
 Imperialine *N*-oxide, C-298
 Imperialine, C-298
 Imperialone, C-298
 Impericine, C-297
 Imperoline, I-52
 Imperonine, I-53
 Impranine, I-54
 Impulsin, P-38
 Inactone, D-765
 Inaequidenine, I-55
 Inandenin A, I-58
 Inandenin B, I-59
 Inandenin-10,11-diol, I-56
 Inandenin-10-ol, I-56
 Inandenin-11-ol, I-57
 Inandenin-12-ol, I-58
 Inandenin-13-ol, I-59
 Inandenin-10-one, I-56
 Inandenin-11-one, I-57
 Inandenin-12-one, I-58
 Inandenin-13-one, I-59
 Incanine *N*-oxide, T-488
 Incanine, T-488
 Incanumine, S-456
 Incartine, P-716
 Incarvillateine C, I-60
 Incarvillateine D, I-60
 Incarvillateine E, I-61
 Incarvillateine *N*-oxide, I-60
 Incarvillateine, I-60
 Incarvilline 7-*O*-ferulate, I-62
 Incarvilline, I-62
 Incarvine A *N*-oxide, I-63
 Incarvine A, I-63
 Incarvine B, I-62
 Incarvine C, I-62
 Incarvine D, I-62
 Incasine A', P-673
 Incasine A, P-673
 Incasine B', P-665
 Incasine B, P-665
 Incasine C', P-665
 Incasine C, P-665
 Indaconitine, M-336
 Indanomycin, I-64
 Indeno[1,2,3-*ij*][2,7]naphthyridine, I-65
 Indicaïne, B-254
 Indicaïne, B-254
 Indicaïne, B-254
 Indicaïne, I-66
 Indicaïne, V-88
 Indican, I-104
 Indicanthinone, I-105
 Indicaxanthin, I-67
 Indicine *N*-oxide, I-68
 Indicine, I-68
 Indicine, I-69
 Indigo blue, I-72
 Indigo brown, B-131
 Indigo red, I-73
 Indigo X, C-77
 Indigo, I-72
 Indigoidin, I-70
 Indigotiisocoumarin A, I-71
 Indigotin, I-72
 Indirubin, I-73
 Indisocin, I-74
 Indizoline, I-75
 Indobine, I-100
 Indobinine, I-100
 Indocarbazostatin B, I-76
 Indocarbazostatin C, I-76
 Indocarbazostatin D, I-76
 Indocarbazostatin, I-76
 Indocybin, H-480
 Indolactam I, I-77
 Indolactam V, I-78
 1*H*-Indole-3-acetaldehyde, I-80
 1*H*-Indole-3-acetamide, I-81
 1*H*-Indole-3-acetic acid, I-81
 1*H*-Indole-3-acetonitrile, I-81
Agelenopsis aperta Indoleacetyl-pentamine toxins, I-82
 Indoleacryloylonitrile, I-216
 Indole-3-alanine, T-640
 Indole-3-aldehyde, I-86
 1*H*-Indole-3-butanolic acid, I-83
 Indole-3-carbinol, I-98
 1*H*-Indole-3-carbonitrile *N*-oxide, I-90
 1*H*-Indole-3-carbonitrile, I-90
 1-(1*H*-Indole-3-carbonyl)-β-carboline-3-carboxylic acid, P-489
 1-(1*H*-Indole-3-carbonyl)-β-carboline, P-488
 1-(1*H*-Indole-3-carbonyl)-9*H*-pyrido[3,4-*b*]indole, P-488
N-(1*H*-Indole-3-carbonyl)urea, I-84
 1*H*-Indole-3-carbothioic acid, I-85
 1*H*-Indole-3-carboxaldehyde, I-86
 1*H*-Indole-5-carboxaldehyde, I-87
 1*H*-Indole-7-carboxaldehyde, I-88
 1*H*-Indole-3-carboxamide, I-90
 Indole-β-carboxylic acid, I-90
 1*H*-Indole-2-carboxylic acid, I-89
 1*H*-Indole-3-carboxylic acid, I-90
 1*H*-Indole-5,6-diol, I-93
 1*H*-Indole-6,7-diol, I-94
 1*H*-Indole-2,3-diol, I-91
 1*H*-Indole-2,5-diol, I-92
 1*H*-Indole-2,3-dione, I-95
 1*H*-Indole-3-ethanamine, T-639
 1*H*-Indole-3-ethanol, I-96
 Indole, I-79
 1*H*-Indole-3-methanamine, I-97
 1*H*-Indole-7-methanol, I-99
 1*H*-Indole-3-methanol, I-98
 4-(3-Indolemethyl)-7-isobutyl-1,2,3,5,6-pentathiepane, I-129
 3-(3-Indolemethyl)-5-isobutyl-1,2,4-trithiolane, I-130
 1*H*-Indole-3-propanamide, I-100
 1*H*-Indole-3-propanoic acid, I-100
 1*H*-Indole-3-propenoic acid, I-136
 2,3-Indolinedione, I-95
 2-Indolinone, I-103
 Indolizidine 223AB, O-41
 Indolizidine 239AB, O-41
 Indolizidine 223A, D-394
 Indolizidine 203A, O-52
 Indolizidine 205A, O-52
 Indolizidine 207A, O-52
 Indolizidine 243B, A-511
 Indolizidine 245B, A-511
 Indolizidine 249B, A-509
 Indolizidine 253B, A-419
 Indolizidine 265B, A-509
 Indolizidine 279B, A-509
 Indolizidine 195B, B-434
 Indolizidine 235B', H-127
 Indolizidine 235B, H-127
 Indolizidine 251B, H-127
 Indolizidine 167B, O-56
 Indolizidine 209B, O-52
 Indolizidine 259B, O-51
 Indolizidine 243C, A-511
 Indolizidine 245C, A-486
 Indolizidine 279C, A-509
 Indolizidine 239CD, O-41
 Indolizidine 275C, H-260
 Indolizidine 243D, A-486
 Indolizidine 279D, A-487
 Indolizidine 209D, H-264
 Indolizidine 233D, H-127
 Indolizidine 237D, H-127
 Indolizidine 241F, A-511
 Indolizidine 251F', A-509
 Indolizidine 251F, A-509
 Indolizidine 235H, A-509
 Indolizidine 249H, E-240
 Indolizidine 209I, O-44
 Indolizidine 251J, A-509
 Indolizidine 223J, B-443
 Indolizidine 267J, E-263
 Indolizidine 237L, B-430
 Indolizidine 251M, E-263
 Indolizidine 223V, B-437
 Indolizomycin, I-101
 Indolmycin, I-102
 Indolo[2,3-*a*]carbazole, D-464
 1*H*-Indol-3-ol, I-104
 1*H*-Indol-2-ol, I-103
 3*H*-Indolo[3,2,1-*de*][1,5]naphthyridine-2,6-dione, I-105
 6*H*-Indolo[3,2,1-*de*][1,5]naphthyridin-6-one, C-90
 3*H*-Indol-3-one, I-106
 7*H*-Indolo[3,2,7]phenanthridine-7,13(12*H*)-dione, I-107
 Indolopyridocoline, I-113
 Indolo[2,1-*b*]quinazoline-6,12-dione, I-108
 10*H*-Indolo[3,2-*b*]quinoline-11-carboxylic acid, I-111
 Indolo[3,2-*b*]quinoline, I-109
 Indolo[3,2-*c*]quinoline, I-110
 Indolo[2,3-*b*]quinoline, Q-19
 10*H*-Indolo[3,2-*b*]quinolin-9-ol, I-112
 5*H*,10*H*-Indolo[3,2-*b*]quinolin-11-one, D-505
 5*H*,6*H*-Indolo[2,3-*b*]quinolin-11-one, D-506
 Indoloquinolizidine-4-carboxylic acid, O-49
 Indoloquinolizidine, O-48
 Indolo[2,3-*a*]quinolizine, I-113
 Indolo[3,2,1-*ij*]quino[4,5-*bc*][1,5]naphthyridine, I-114
 3-Indolyl gentiobioside, I-104
 3-Indolyl methyl ketone, A-59
 β-Indolylacetaldehyde, I-80
 3-Indolylacetic acid, I-81
 3-Indolylacetyl-D-arabinitol, A-110
 8-(1*H*-Indol-3-ylacetyl)falcariindiol, H-116
 1*H*-Indol-3-ylacetyl-*myo*-inositol, I-115
 3-(3-Indolyl)acrylamide, I-136
 3-(3-Indolyl)acrylic acid, I-136
 3-Indolylalanine, T-640
 4-(1*H*-Indol-3-yl)-1,2,3-butanetriol, I-116
 4-(1*H*-Indol-3-yl)-3-buten-2-one, I-117
 4-(1*H*-Indol-6-yl)-3-buten-2-one, I-118
 4-Indol-3-ylbutyric acid, I-83
 1-(3-Indolyl)-β-carboline, I-119
*N*²-(1*H*-Indol-3-ylcarbonyl)arginine, I-120
*N*²-(1*H*-Indol-3-ylcarbonyl)histidine, I-121
 1-(1*H*-Indol-3-ylcarbonyl)-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, P-489
 2-(3-Indolyl)ethanol, I-96
 1-(1*H*-Indol-3-yl)ethanone, A-59
 2-(1*H*-Indol-3-yl)ethanamine, D-172
N-[2-(1*H*-Indol-3-yl)ethyl]formamide, D-172
 2-(3-Indolyl)ethyl phenylacetate, I-96
 2-(3-Indolyl)ethylamine, T-639
N-[2-(1*H*-Indol-3-yl)ethyl]benzamide, T-639
N-[2-(1*H*-Indol-3-yl)ethyl]docosanamide, T-639
 3-Indolylethylene oxide, O-165
N-[2-(1*H*-Indol-3-yl)ethyl]-2-(methylamino)benzamide, T-639
 5-[1-(1*H*-Indol-3-yl)ethyl]-2-(methylamino)-4(5*H*)-oxazolone, I-102
N-[2-(1*H*-Indol-3-yl)ethyl]-3-methyl-2-oxopentamide, T-639
 3-[2-(1*H*-Indol-3-yl)ethyl]-1-methyl-2,4(1*H*,3*H*)-quinazolinedione, G-144
 3-(3-Indolyl)glycerol, I-135
 2-(3-Indolyl)glyoxylic acid, O-182
 4-(1*H*-Indol-3-yl)-1*H*-imidazole-2,5-dione, Z-39
 5-(1*H*-Indol-3-yl)-2,4-imidazolidinedione, Z-39
 β-1*H*-Indol-3-yl-1*H*-indole-3-ethyl acetate, S-564
 4-(3-Indolyl)-2-(3-indolylcarbonyl)imidazole, T-432
 3-(1*H*-Indol-3-yl)-*N*-[2-(1*H*-indol-3-yl)ethyl]-2-propenamide, C-421
 1*H*-Indol-3-yl[4-(1*H*-indol-3-yl)-1*H*-imidazol-2-yl]-methanone, T-432
 5-(3-Indolyl)-2-isobutyloxazole, A-622
 3-(3-Indolyl)lactic acid, H-539
 5-(1*H*-Indol-3-yl)-5-methoxy-2,4-imidazolidine-dione, P-542

- 1-(1*H*-Indol-3-yl)-2-methoxy-5-methyl-3-hexanone, S-101
 8-(1*H*-Indol-3-yl)-2-methoxy-6-methyl-1,4-naphthalenedione, M-750
 4-(1*H*-Indol-3-yl)-4-methoxy-5-thioxo-2-imidazolidinone, Z-39
 5-(1*H*-Indol-3-ylmethyl) allylthiocarbamate, B-374
 3-Indolylmethyl glucosinolate, I-122
 5-(1*H*-Indol-3-ylmethyl) 2-propenylthiocarbamate, B-374
 3-Indolylmethylamine, I-97
 1-(1*H*-Indol-3-yl)-3-methyl-2,3-butanediol, T-25
 2-(1*H*-Indol-3-yl)-3-methyl-1,3-butanediol, T-24
 1-(1*H*-Indol-7-yl)-3-methyl-1,2,3-butanetriol, I-123
 4-(3-Indolyl)-2-methylbutanoic acid, I-124
 4-(3-Indolyl)-2-methyl-1-butanol, I-124
 1-(1*H*-Indol-3-yl)-3-methyl-2-buten-1-one, I-125
 1-(1*H*-Indol-7-yl)-3-methyl-2-buten-1-one, M-414
N-(1*H*-Indol-3-ylmethyl)-*N,N*-dimethyl-1*H*-indole-3-methanaminium, B-190
 3-(1*H*-Indol-3-yl)-2-methyl-*N,N*-dimethylpropanamide, H-539
 5-β-Indolylmethyl-8*N*-(*N,N*-dimethylvalyl)-9-isopropylphenylcyclopeptide, A-690
 6-(1*H*-Indol-3-yl)-5-methyl-3,5-heptadien-2-one, I-126
 2-*C*-(1*H*-Indol-3-ylmethyl)-3-hexulofuranosonic acid γ -lactone, A-1461
 2-(1*H*-Indol-3-ylmethyl)-1*H*-indole-3-acetic acid, I-127
 2-[(1*H*-Indol-3-yl)methyl]-1*H*-indole-3-carboxaldehyde, M-65
N-(1*H*-Indol-3-ylmethyl)-5-methoxy-*N*-methyl-1*H*-indole-3-ethanamine, I-128
N^b-(3-Indolylmethyl)-5-methoxy-*N*^b-methyltryptamine, I-128
N-[7-(1*H*-Indol-3-ylmethyl)-3-(1-methylethyl)-5,8-dioxo-2-oxa-6,9-diazabicyclo[10.2.2]hexadecane-10,12,14,15-tetraen-4-yl]-3-methyl-2-(methylamino)butanamide, A-690
 4-(3-Indolylmethyl)-7-(2-methylpropyl)-1,2,3,5,6-pentathiepane, I-129
 3-(3-Indolylmethyl)-5-(2-methylpropyl)-1,2,4-trithiolane, I-130
 5-(3-Indolyl)-2-methylloxazole, A-622
 3-(3-Indolyl)-*N*-methylpropanamide, I-100
 3-(1*H*-Indol-3-yl)-2-methylpropanoic acid, I-131
 7-(1*H*-Indol-3-ylmethyl)quinazolinof[3,2-*d*][1,4]benzodiazepine-6,9(5*H*,7*H*)-dione, A-1480
 4-(1*H*-Indol-3-ylmethyl)-2,2,6-trimethyl-3-azabicyclo[3.3.1]non-6-ene, H-310
 5-(3-Indolyl)oxazole, O-157
N-[2-(1*H*-Indol-3-yl)-2-oxoethyl]acetamide, I-132
 3-(1*H*-Indol-3-yl)-2-oxopropanoic acid, I-133
 5-(1*H*-Indol-3-yl)-3-penten-2-one, I-134
 5-(3-Indolyl)-2-pentylloxazole, A-622
 1-(1*H*-Indol-3-yl)-1,2,3-propanetriol, I-135
 2-(3-Indolyl)propanoyl-*D*-arabinitol, A-109
 3-(1*H*-Indol-3-yl)-2-propenamide, I-136
 3-(1*H*-Indol-3-yl)-2-propenoic acid, I-136
N^a-[3-(1*H*-Indol-3-yl)propenoyl]tryptophan, I-137
 3-(3-Indolyl)propionic acid, I-100
 5-(3-Indolyl)-2-propylloxazole, A-622
 1-(1*H*-Indol-3-yl)-9*H*-pyrido[3,4-*b*]indole, I-119
 3-(3-Indolyl)pyruvic acid, I-133
 3-(1*H*-Indol-3-yl)quinoline, I-138
 3-[4-(1*H*-Indol-3-yl)-2,3,5,6-tetramethoxyphenyl]-7-(3-methyl-2-butenyl)-1*H*-indole, A-1519
 2-(3-Indolyl)thiazole, T-372
 5-(1*H*-Indol-3-yl)-2(3*H*)-thiazolethione, I-139
 2-(3-Indolyl)-6-(3,4,5-trimethoxyphenyl)morpholine, C-361
 2-(1*H*-Indol-3-yl)-*N,N,N*-trimethylethenaminium, D-172
 2-(3-Indolyl)vinyl isocyanide, I-216
 Indoxyl, I-104
 Inflatine, L-219
 Infractine, C-134
 Infractopicrine, I-140
 Infumorph, M-704
 Ingamine A, I-141
 Ingamine B, I-141
 Ingenamine B, I-143
 Ingenamine C acetate, I-143
 Ingenamine D, I-144
 Ingenamine E, I-145
 Ingenamine F, I-145
 Ingenamine G, I-146
 Ingenamine, I-142
 Ingol, I-147
 Innovanamine, D-632
 Inosie, I-148
 Inosine, I-148
 Insulamine, I-149
 Insulanoline, I-150
 Insularine 2β-*N*-oxide, I-150
 Insularine 2β-*N*-oxide, I-150
 Insularine, I-150
 Insularoline, I-150
 Insulopinine, I-151
 Intebriamine, S-234
 Intebrine, I-152
 Intebrinone, I-153
 Integerrenine, I-154
 Integerressine, I-155
 Integerrimine *N*-oxide, S-238
 Integerrimine, S-238
 Integerrine, I-156
 Integramine, I-157
 Integramycin, I-158
 Integramide, I-159
 Integriquinolone, Q-46
 Interacton, T-366
 Intermedine, I-160
 Inthomycin A, I-161
 Inthomycin B, I-161
 Inthomycin C, I-161
 Inthomycin, I-161
 Intropin, D-920
 Inuline, M-309
 Inundatine, L-335
 Involucratine, I-162
 Iodanthine, B-401
 Iodinine, P-321
 Iodoacetamide, I-163
 Iodoacetic acid, I-163
 Iodoacetoneitrile, I-163
 Iodoacetyl chloride, I-163
 (Iodomethyl)trimethylammonium(1+), I-164
 1-Iodo-*N,N,N*-trimethylmethanaminium, I-164
 Iolantamine, J-43
 Iolantidine, K-36
 Iolantimine, J-44
 Iolantine, J-43
 Iolantimine, J-45
 Ipaconitine, M-343
 IPA, I-100
 Ipalbidine, I-165
 Ipalbidinium, I-170
 Ipalbine, I-165
 Ipanguline A₁, I-166
 Ipanguline A₂, I-166
 Ipanguline A₃, I-166
 Ipanguline A₄, I-166
 Ipanguline B₁, I-166
 Ipanguline B₂, I-166
 Ipanguline B₃, I-166
 Ipanguline C₁, I-166
 Ipanguline C₂, I-166
 Ipanguline C₃, I-166
 Ipanguline C₄, I-166
 Ipanguline D₆, H-233
 Ipanguline D₇, H-233
 Ipanguline D₁₀, I-166
 Ipanguline D₁₄, I-166
 Ipanguline D₁₅, I-166
 Ipanguline D₃, I-166
 Ipanguline D₅, I-166
 Ipanguline D₉, I-166
 Ipecine, E-72
 Ipecoside, I-167
 Ipecosidic acid, I-167
 Iperphos, G-108
 Ipobscurine A, H-755
 Ipobscurine B, I-168
 Ipobscurine C, I-169
 Ipobscurine D, I-169
 Ipohardine, I-170
 Ipomine A, P-341
 Ipomine, I-165
 2iP, Z-11
 Ircinal A, I-174
 Ircinal B, I-175
 Ircinamine B, I-172
 Ircinamine, I-171
 Irciniastatin A, I-173
 Irciniastatin B, I-173
 Irciniasulfonic acid A₁, H-594
 Irciniasulfonic acid A₂, H-594
 Irciniasulfonic acid A₃, H-594
 Irciniasulfonic acid B₁, H-594
 Irciniasulfonic acid B₂, H-594
 Ircinol A, I-174
 Ircinol B, I-175
 Irehamine, A-886
 Irehdiamine A, P-606
 Irehdiamine B, P-606
 Irehdiamine C, P-606
 Irehdiamine F, P-606
 Irehdiamine I, P-606
 Irehine, A-886
 Irehline, C-610
 Irenine, I-176
 Iresinin I, B-110
 Iresinin II, B-110
 Iresinin III, B-110
 Iresinin IV, B-110
 Irididine, I-178
 Irnigaine, I-177
 Irniine, I-178
 Iromycin A, I-179
 Iromycin B, I-179
 Iromycin C, I-180
 Iromycin D, I-180
 Isaindigotidione, I-181
 Isaindigotone, I-182
 Isalexin, H-532
 Isariotin A, D-98
 Isariotin B, D-98
 Isariotin C, D-98
 Isariotin D, D-98
 Isatan A, I-104
 Isatan B, I-104
 Isatan C, I-91
 Isatidine, R-62
 Isatin, I-95
 Isatinone A, I-183
 Isatinone B, I-184
 Isatisine A, I-185
 Isatogen, I-106
 Isatropyliditropein, B-56
 Iseluxine, I-305
 Isendryl, C-835
 Ishaematinic acid, I-239
 Ismine, I-186
 Isoaaptamine, A-2
 Isoabsoulone *N*-oxide, A-906
 Isoabsoulone, A-906
 Isoaconitine, M-336
 Isoadlumiceine, B-120
 Isoagleptine, L-115
 3-Isoajmalicine, A-212
 Isoajmaline, A-216
 Isoalamarine, A-235
 Isoalangamarine, A-235
 Isoalangaside, A-239
 Isoalatlol, H-249
 Isoalchorneine, I-187
 Isoalfileramine, I-188
 Isoalloxazine adenine dinucleotide, F-79
 Isoalloxazine, A-632
 Isoalschomine, A-640
 Isoalstonisine, A-654
 Isoamaranthin, B-110
 Isoamarantin, B-110
 Isoaminobisabolol a, A-710
 Isoaminobisabolol b, A-710

- Isoammodendrine, A-930
 Isoampullicin, A-951
 Isoamurone, C-781
N-Isoamylacetamide, M-420
 Isoamylamine, M-420
 Isoanagryne, T-363
 Isoanantine, I-189
 Isoancistrocladine, A-982
 Isoandrocybine, A-1005
 Isoanguivine, S-456
 Isoangustifoline, A-1013
 Isoanhalamine, T-236
 Isoanhalidine, T-236
 Isoanhalonidine, T-237
 Isoanthrine, A-1328
 Isoantirrhine, A-1328
 3-Isoantirrhine, A-1328
 α -Isoaphylline, A-1337
 Isoapocavidine, C-222
 Isoaquiledine, I-190
 Isoaristolochic acid, M-262
 Isoarrottianamide, A-1446
 Isoasterriquinone SU 5220, I-191
 Isoasterriquinone, I-191
 Isoatisine, I-192
 Isoautumnaline, A-1558
 Isoazitine, A-1609
 Isobackebergine, I-303
 Isobaimonidine, C-298
 Isobalfourodine, R-92
 Isobastadin 4, B-32
 Isobatzelline A, I-193
 Isobatzelline B, I-193
 Isobatzelline C, I-193
 Isobatzelline D, I-193
 Isobatzelline E, I-193
 Isobebeerine, C-835
 Isobellendine, I-194
 Isobengamide E, I-195
 Isobetanidin 6-(2^O-glucosylrutinoside), B-110
 Isobetanidin rhamnosophoroside, B-110
 Isobetanidin, B-110
 Isobetanin, B-110
 Isobisdehydrostemocochinine, P-670
 Isobisdehydrostemoninine, S-521
 Isobisdehydrotuberostemonine, T-664
 Isoboldine *N*-oxide, I-196
 Isoboldine, I-196
 Isobonafousine, I-197
 Isoborrerine, M-521
 Isoborreverine, I-198
 Isobougainvillein r-I, B-110
 Isobougainvillein r-II, B-110
 Isobrafouedine, I-199
 Isobromodeoxytopsentin, T-432
 Isobromotopsentin, T-432
 Isobrucine *N*-oxide, P-672
 Isobrucine, P-672
 Isobubbialine, B-375
 Isobulbocapnine, N-25
 Isobungeanol, T-129
 3-Isobutanoyloxy-6-(3,4,5-trimethoxy-*cis*-cinnamoyloxy)norpropane, A-1584
 1-Isobutenyl- β -carboline, M-521
 Isobutyl pidolate, O-206
 Isobutylacetic acid, M-505
 Isobutylamine, M-525
N-Isobutyl-2,4-decadienamide, D-106
N-Isobutyl-6,8-decadienamide, D-105
N-Isobutyl-2,4,8-decatrienamide, D-123
N-Isobutyl-2,6,8-decatrienamide, D-124
N-Isobutyl-4,5-dihydroxy-2-decenamide, D-566
N-Isobutyl-3,4-dimethoxybenzamide, D-706
 3-Isobutyl-2,5-dimethylpyrazine, D-757
N-Isobutyl-2,8-dodecadienamide, D-884
N-Isobutyl-2,4-dodecadiene-8,10-diynamide, D-882
N-Isobutyl-2,6,8,10-dodecatetraenamide, D-893
N-Isobutyl-2,4-eicosadienamide, E-45
N-Isobutyl-2,4,14-eicosatrienamide, E-50
N-Isobutyl-2,4,8-eicosatrienamide, E-49
N-Isobutyl-2,4-hexadecadienamide, H-216
N-Isobutylidenemethylamine, M-383
 2-Isobutyl-5-isopropyl-1*H*-imidazole, I-280
 2-Isobutyl-5-isopropylpyrazine, I-282
 3-Isobutyl-6-isopropyl-2(1*H*)-pyrazinone, M-462
N-Isobutyl-3-methoxy-4,5-methylenedioxy-cinnamide, T-578
 5-Isobutyl-3-methoxy-2-methylpyrazine, M-268
N-Isobutyl-6-*p*-methoxyphenylsorbamide, P-466
 2-Isobutyl-3-methoxypyrazine, M-273
 2-Isobutyl-4-methyl-2,4-azetidinedicarboxylic acid, M-482
N-Isobutyl-3,4-methylenedioxy-cinnamide, M-457
N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,12-pentadecatrienamide, B-265
N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,13-pentadecatrienamide, B-265
N-Isobutyl-15-(3,4-methylenedioxyphenyl)-2,4,14-pentadecatrienamide, B-265
N-Isobutyl-5-(3,4-methylenedioxyphenyl)-2-pentenamide, P-462
N-Isobutyl-5-(3,4-methylenedioxyphenyl)-4-pentenamide, P-462
 2-Isobutyl-3-methylpyrazine, M-483
 3-Isobutyl-1-methylxanthine, X-5
N-Isobutyl-2,4-octadecadienamide, O-29
N-Isobutyl-2,4,12-octadecatrienamide, O-33
N-Isobutyl-2,4-octadienamide, O-39
N-Isobutylotanthusic acid amide, O-133
N-Isobutyl-12-oxo-2,4,8,10-dodecatetraenamide, O-176
N-Isobutyl-8-oxo-2,4-tetradecadienamide, O-211
N-Isobutyl-12-oxo-2,4,8-tetradecatrienamide, O-212
 3-[(7-Isobutyl-1,2,3,5,6-pentathiepan-4-yl)methyl]indole, I-129
N-Isobutyl-16-phenyl-2,4-hexadecadienamide, P-345
 2-Isobutyl-5-phenyl-1*H*-imidazole, M-528
 3-Isobutylpiperidine, M-530
 2-Isobutyl-3-(4-prenyloxyphenyl)maleimide, H-684
 Isobutylpropylamine, M-525
N-Isobutyl-2,4,8,10,12-tetradecapentaenamide, T-127
N-Isobutyl-2,4,10,12-tetradecatetraen-8-ynamide, T-130
N-Isobutyl-2,4,12-tetradecatriene-8,10-diynamide, A-968
 2-Isobutylthiazole, M-533
N-Isobutyl-7-tridecene-10,12-diynamide, T-505
N-Isobutyl-3,4,5-trimethoxycinnamide, T-578
 3-[(5-Isobutyl-1,2,4-trithiolan-3-yl)methyl]indole, I-130
 Isobutyropyrrrothine, A-759
 Isobutyropyrrrothine, I-200
 6-Isobutyrylamino-4,5-dihydro-4-methyl-1,2-dithiolo[4,3-*b*]pyrrole, I-200
N-Isobutyrylbaleabuxaline F, B-11
N-Isobutyrylbaleabuxidienine F, B-466
 Isobutyrylbaleabuxidine F, B-12
N-Isobutyrylbuxahyrcanine, B-451
N-Isobutyrylbuxidienine F, B-466
*N*³-Isobutyrylcyclohexobuxidine F, B-12
 3-*N*-Isobutyrylcyclohexobuxidine H, B-12
*N*³-Isobutyrylcyclohexobuxine F, C-920
N-Isobutyryl-*N*-depropionyldecorticasine, L-234
 14-Isobutyryldictyocarpine, M-307
 2-Isobutyryl-14-hydroxyhetisine, H-189
 7-Isobutyrylintermediate, I-160
 7-Isobutyryllycopsamine, I-160
 14-Isobutyrylnudicaulidine, M-321
 13-Isobutyryloxylopamine, H-554
 3-Isobutyryloxytropin-7-ol, A-1585
 3-Isobutyryloxytropin-6-ol, M-391
N-Isobutyrylpyrrrothine, A-759
 Isocaboxine A, C-1
 Isocaboxine B, C-1
 Isocadamine, C-6
 Isocalcanthine, T-139
 Isocalcimine, D-846
 Isocalciphylline B, C-45
 Isocantleyine, C-92
 Isocaproic acid, M-505
 Isocapsicastrine, S-200
 Isocarapanabine, C-117
 6-Isocarnavaline, C-183
 Isocaryachine *N*-oxide B, E-220
 Isocaryachine *N*-oxide, E-220
 (+)-Isocaryachine *N*-oxide, E-220
 Isocaryachine, E-220
 6-Isocassine, C-183
 Isocastoramine, D-227
 Isocelosianin I, B-110
 Isocelosianin II, B-110
 Isocephaline, E-72
 Isocephalotaxine, I-201
 Isocephalotaxinone, I-201
 Isochaetoglobosin D, C-327
 Isochaetoglobosin J, C-328
 Isochaksine, I-202
 Isochalciporone, C-340
 Isochanoclavine I, C-348
 Isochavicine, P-467
 Isochelidonine, I-203
 Isochondourarine, I-204
 Isochondodendrine, C-835
 Isochromophilone IX, I-205
 Isochromophilone VI, I-205
 Isochrysohermidin, I-206
 Isochrysoetricine, C-433
 Isocimicifugamide, H-566
 Isocinchophyllamine, C-448
 Isocinevanine, H-554
 Isococamine, T-635
 Isococcolimine, E-202
 Isococculidine, I-207
 Isococculine, I-207
 Isococclaurine, H-275
 Isocodonocarpine, C-4
 Isoconessimine, C-595
 Isocorcovadine, P-462
 Isocoreximine, X-30
 3-Isocorreatine A, C-659
 Isocorybulbine, C-675
 Isocorydine *N*-oxide, I-208
 Isocorydine, I-208
 9-*O*-(8-Isocorydiny)-*N*-methyl-lauroretanin, D-149
 Isocorydione, S-367
 Isocorymine, I-209
 3-Isocorynantheidine, C-689
 Isocorynoline, C-692
 Isocorynoxine, R-90
 Isocorynoxine, R-90
 Isocorypalline, T-158
 Isocorypalmine α -*N*-oxide, I-210
 Isocorypalmine β -*N*-oxide, I-210
 Isocorypalmine, I-210
 Isocorytuberine, T-248
 Isocrambescidin 800, C-717
 Isocrambescidin 657, N-115
 Isocraugsodine, C-730
 Isocreatinotone *N*-oxide, T-188
 Isocrocandine, C-764
 Isocromadurine, F-168
 Isocrotsparinine, A-957
 Isocryptochine, I-211
 Isocryptolepine, I-110
 Isocuauchichicine, G-25
 Isoculantramine, C-803
 Isocuricyclatejine, C-811
 Isocuricyclatejine, C-811
 7-Isocyanato-2,10-bisaboladiene, I-212
 7-Isocyanato-7,8-dihydro- α -bisabolene, I-212
 11-Isocyanato-5-eudesmene, F-129
 20-Isocyanato-7-isocyanoadocaine, D-698
 7-Isocyanato-20-isocyanoisocycloamphilectane, D-698
 20-Isocyanato-7-isocyanoisocycloamphilectane, D-698
 10 α -Isocyanooalloaromadendrane, A-1571
 1-Isocyanooaromadendrane, A-1447
 3-Isocyanato-8,10-bisaboladiene, A-709
 7-Isocyanato-2,10-bisaboladiene, I-214
 3-Isocyanato-7,9-bisaboladiene, I-213
 8-Isocyanato-1(12)-cycloamphilectene, I-215

- 7-Isocyano-7,8-dihydro- α -bisabolene, I-214
 3-(2-Isocyanoethenyl)-1*H*-indole, I-216
 4-(2-Isocyanoethenyl)phenol, H-680
 4-Isocyano-11-eudesmene, A-775
 5-Isocyano-6-eudesmene, I-217
 6-Isocyano-4(15)-eudesmene, I-218
 4-Isocyano-11-gorgonene, A-777
 20-Isocyano-7-isothiocyantoisocycloamphilectane, D-698
 4-Isocyanomaalialane, I-219
 3-(2-Isocyano-5-methyl-1,3,5-hexatrienyl)-1-methyl-1*H*-indole, B-281
 4-Isocyano-3'-methylspiro[6-oxabicyclo[3.1.0]hex-3-ene-2,2'-oxirane], I-255
 9-Isocyanoneopupekeanane, I-220
 3-(4-Isocyano-6-oxabicyclo[3.1.0]hex-3-en-1-yl)-2-propenoic acid, I-256
 3-Isocyano-3,7,11,15-tetramethyl-1,6,10,14-hexadecatetraene, I-221
 3-Isocyanothionellin, I-213
 Isocycleanonine, C-834
 Isocycleanine, S-148
 Isocyclocapitelline, C-433
 Isocyclocelabenzine, I-222
 Isocycloheximide, D-765
 Isocycloneosamandaridine, I-223
 Isocynodine, I-224
 Isocynometrine, I-224
 Isocytidine, A-823
 Isocytosine, A-823
 Isodaphnilactone B, D-53
 Isodarlingianine, D-74
 6-Isodaucene 14-isonitrile, I-225
 6-Isodaucene 14-isothiocyante, I-225
 Isodaurisoline, D-88
 Isodeacetylretuline, R-63
 Isodecarine, C-356
 Isodecyl 5-hydroxyanthranilate, A-789
 Isodelectine, M-309
 Isodelpheline, M-321
 Isodelphinine, M-313
 Isodendrill, C-835
 Isodictamine, F-224
 Isodihydrocadambine, I-226
 3 β -Isodihydrocadambine, I-226
 Isodihydrochanoclavine I, C-348
 Isodihydrocyclophyllyne A, C-883
 Isodihydrohistrionicotoxin, H-309
 Isodihydropiperlonguminine, P-462
 C-Isodihydrotoxiferine I, I-227
 Isodiplamine, D-823
 Isodolichantoside, S-583
 Isodomesticine, N-29
 Isodomoic acid A, I-228
 Isodomoic acid B, I-228
 Isodomoic acid C, I-228
 Isodomoic acid D, D-912
 Isodomoic acid E, D-912
 Isodomoic acid F, D-912
 Isodomoic acid G, I-229
 Isodomoic acid H, I-229
 Isodrosopeterin, D-941
 Isodysidenin, D-972
 Isoebedine, C-290
 Isoeburnamine, E-6
 Δ^{14} -Isoeburnamine, E-6
 (-)-Isoeburnamine, E-6
 Isoechinatine, E-12
 Isoechinulin A, I-230
 Isoechinulin B, I-230
 Isoechinulin C, I-230
 Isoelacomine, E-53
 Isoelaecarpicine, I-231
 Isoelaecarpiline, E-55
 Isoelaiocarpine, E-56
 Isoelegantissine, M-50
 Isoeleocarpine, E-56
 Isoeutherobin A, E-64
 D-Isoephedrine sulfate, M-386
 Isoephedrine, M-386
 3-Isopropyl-epicathenamine, C-207
 Isoergine, L-370
 Isoerysopinophorine, E-201
 Isoerythreine, E-210
 Isoesculeogenin A, S-444
 25-Isoetioline, S-199
 Isoeudistomin U, I-119
 Isoevonine, P-252
 Isofagaridine, C-356
 Iso- γ -fagarine, F-213
 Isofangchinoline, I-328
 Isofebrifugine, I-232
 Isofedrol, M-386
 Isoferuloylcholine, C-419
N-*cis*-Isoferuloylglutamic acid, G-107
N-*trans*-Isoferuloylglutamic acid, G-107
N-*trans*-Isoferuloyl-3-*O*-methyltyrosine, H-566
 Isogenicine, I-233
 Isofiliformine, F-315
 Isofistularin 3, F-70
 Isoflavipucine, I-234
 Isoflindersiamine, F-217
 Isoflustramine D, I-235
 Isoformosamine, M-661
 Iso-*N*-formylchoneomorphine, P-601
 Isofoveoglin, P-878
 Isogapavine, L-72
 Isogumiglavine A, F-181
 Isogumiglavine B, F-181
 Isofuntumidine, A-880
 Isogagaminine, P-610
 Isogambirdine, M-661
 Isogambirine, C-689
 Isogarryfoline, G-25
 Isogeissoschizol, G-33
 Isogentialtine, I-236
 Isogermidine, G-70
 Isogillette *N*-oxide, C-552
 Isogomphrenin I, B-110
 Isogomphrenin II, B-110
 Isogomphrenin III, B-110
 Isogranulatimide, I-237
 Isogravacridonechlorine, R-167
 Isoguanine, A-818
 Isoguattouregidine, G-215
 Isoguvacine, T-222
 Isohalichondramide, H-15
 Isohaplopin prenyl ether, F-223
 Isohaplopin, F-223
 Isoharmine, H-591
 Isoharringtonic acid, I-238
 Isoharringtonine, I-238
 Isohematinic acid, I-239
 Isohemsleyanidine, M-306
 Isohemsleyanisine (incorr.), M-306
 Isoherbavine, M-50
 Isohexanoic acid, M-505
 Isohobartine, H-310
 Isohydrastidine, H-394
 Isohylocerenin, B-110
 Isohypercetine, H-771
 Isohypoglavine, H-203
 Isohypoglavinol, H-203
 Isoimerubrine, G-159
 Isoindigo, B-131
 Isoindigotin, B-131
 1*H*-Isoindole-1,3(2*H*)-dione, P-383
 Isoinundatine, L-335
 Isoipanguline A₁, I-166
 Isoipanguline A₂, I-166
 Isoipanguline A₃, I-166
 Isoipanguline B₁, I-166
 Isoipanguline B₂, I-166
 Isoipanguline B₃, I-166
 Isoipanguline C₁, I-166
 Isoipanguline C₂, I-166
 Isoipanguline D₁₀, I-166
 Isoipanguline D₃, I-166
 Isoipanguline D₅, I-166
 Isoipomine, I-165
 Isoiresinin I, B-110
 Isojacpaniculine, J-5
 Isojuliprosine, J-57
 Isojulocrotol, J-58
 Isojuripidine, A-909
 Isojuripine, A-909
 Isojurubidine, S-458
 Δ^5 -Isojurubidine, S-458
 3 α - Δ^5 -Isojurubidine, S-458
 Isokibdelone A, I-240
 Isokibdelone B, I-240
 Isokibdelone C, I-240
 Isokomarovine, Q-53
 Isokuraramine, K-108
 23-Isokuroyuridine, K-114
 Isolapidilectine A, L-43
 Isolappaconitine, A-90
 Isolaureline *N*-oxide, A-1066
 (+)-Isolaureline, A-1066
 (-)-Isolaureline, A-1066
 Isolaurifoline, G-25
 Isolauryl 5-hydroxyanthranilate, A-789
 Isoleontalbine, I-241
 Isoleontine, A-628
 Isoleurosine, V-110
 Isoliensinine, N-87
 Isoline, B-194
 Isolobelanine, L-218
 Isolobinanidine, L-221
 Isolobinine, L-221
 Isolonganone, L-244
 Isolongistrobine, D-160
 Isolucidinine, L-278
 α -Isolupanine, I-242
 11-Isolupanine, I-242
 Isolupinine, E-95
 Isolycodoline, L-325
 Isolycopodine, L-328
 Isolycoposamine, H-93
 Isolysergamide, L-370
 Isolysergic acid amide, L-370
 Isolysergic acid α -hydroxyethylamide, L-370
 Isolysergic acid methylcarbinolamide, L-370
 Isolysergic acid, L-370
 Isolysergol, L-371
 Isomacrorine, I-44
 Isomaculine, F-221
 Isomaculosidine, F-222
 Isomahanimbine, M-46
 Isomahanine, M-46
 Isomaistemonine, M-49
 Isomajdine, M-50
 Isomalindine-16-carboxylate, I-243
 Isomalindine, M-73
 Isomallotusins, M-135
 Isomammillarinnin, B-110
 Isomaruquine, M-661
 Isomatrine, I-244
 Isomeridine, M-206
 Isometeloidine, M-392
 Isomethuenine, M-293
 Iso-18-*O*-methylchaenorpine, C-318
 Isomigrastatin, I-245
 Isomitomycin A, I-246
 Isomitrafoline, R-134
 Isomitraphyllic acid β -D-glucopyranosyl ester, M-661
 Isomitraphyllic acid, M-661
 Isomitraphylline *N*-oxide, M-661
 Isomitraphylline, M-661
 Isomitraphyllinol, I-247
 Isomonodoroinole, I-248
 Isomoschatoline, M-712
 Isomurralonginol acetate, I-249
 Isomurralonginol isovalerate, I-249
 Isomurralonginol nicotinate, I-249
 Isomurralonginol, I-249
 Isomurrayafoline B, E-275
 Isomurrayazoline, I-250
 Isonaamidine A, I-251
 Isonaamidine B, I-251
 Isonaamidine C, I-251
 Isonaamidine D, I-251
 Isonaamidine E, I-251
 Isonaamine A, I-252
 Isonaamine B, I-253
 Isonaamine C, I-252
 Isonargenicin A1, I-254
 Isonaolaurerine, N-122
 Isonestemocochinine, N-140

- Isonicotine, B-145
 Isonicotinic acid, P-898
 Isonitramine, N-241
 Isonitrarine, N-246
 Isonitrin A, I-255
 Isonitric acid E, I-256
 Isonorargemonine, A-1410
 Isonortehaunine, I-307
 Isonorweberine, T-276
 Isonoyaine, I-257
 Isoobscurinine, I-258
 Isoochotensine, O-16
 Isooconovine, P-209
 Isooconotine, I-259
 Isooncotine, D-625
 Isoorensine, A-138
 Isooridine, O-109
 Isoorientalinone, O-112
 Isooxodesmosine, I-260
 Isooxyskytanthine, H-733
 Isopachycereine, T-233
 Isopalauamine, S-611
 Isopalythazine, I-261
 Isopaniculidine, A-910
 Isoparaxanthine, X-5
 Isoparfumine, F-174
 Isopauridianthoside, P-135
 3-Isopayantheine, G-13
 Isopeduncularine, P-162
 Isopeganidine, P-164
 Isopelletierine, P-173
 Isopellotine, T-237
 Isopenangine, M-577
 Isopenniclavine, S-273
 Isopentanoic acid, M-409
 N⁹-Isopentanoylhistamine, H-307
 N⁶-Isopentenyladenine, Z-11
 N-Isopentenylidendrobine, D-210
 N-Isopentenylidendroxine, D-216
 O-Isopentenylhalfordinol, H-12
 5-Isopentenylisatin, M-415
 6-Isopentenylisatin, M-416
 N-Isopentenyllaburninic acid, A-1065
 7-Isopentenyl- γ -fagarine, F-223
 9-Isopentenylpaxilline, P-618
 N-Isopentylacetamide, M-420
 Isopentylamine, M-420
 2-Isopentyl-7-azaindole, M-422
 3-Isopentyl-2,5-dimethylpyrazine, D-754
 N-Isopentyl-3,4-methylenedioxybenzamide, M-457
 Isopentylpropylamine, M-420
 Isoperiphylline, P-265
 Isoperlolyrine, I-262
 Isophakellistatin 3, P-303
 Isophellibilidine, I-263
 Isophellibiline, P-315
 Isophomazarin, I-264
 Isophorbol, P-225
 Isophylloactin II, B-110
 Isophylloactin, B-110
 Isophysostigmene, I-265
 Isopiline, T-521
 (+)-Isopiline, T-521
 Isopilosine, P-428
 22-Isopimpifolidine, D-99
 Isopiperine, P-467
 Isopiperlonguminine, P-462
 Isopiperolein B, I-266
 Isopiscerythramine, I-267
 Isoplatydesmine, I-268
 Isoplectrodorine, P-517
 Isoplysins A, A-1357
 Isopolyalthenol, I-269
 Isoporoidine, A-1597
 2-Isopropenyl-5-isopropylpyrazine, I-270
 2-Isopropenyl-6-isopropylpyrazine, I-271
 Isopropylacetic acid, M-409
 11-Isopropylcryptolepine, I-279
 2-Isopropyl-3,5-dimethoxy-6-methylpyrazine, I-272
 N²-Isopropylglutamine, G-108
 2-Isopropyl-10-methoxybenzimidazo[6,7,1-def][1,6]naphthyridine, I-273
 6-Isopropyl-3-(4-methoxybenzyl)-4-methyl-2,5-morpholinedione, I-274
 2-Isopropyl-3-methoxy-5-(1-methylpropyl)pyrazine, I-275
 2-Isopropyl-3-methoxy-5-(2-methylpropyl)pyrazine, I-276
 3-Isopropyl-2-methoxy-5-methylpyrazine, I-277
 2-Isopropyl-3-methoxypyrazine, H-545
 2-Isopropyl-5-methoxypyrazine, H-546
 2-Isopropyl-6-methoxypyrazine, H-547
 3-Isopropyl-2-methoxyprido[2,3-*b*]pyrazine, I-278
 11-Isopropyl-5-methyl-5*H*-indolo[3,2-*b*]quinoline, I-279
 5-Isopropyl-2-(2-methylpropyl)-1*H*-imidazole, I-280
 2-Isopropyl-5-(1-methylpropyl)pyrazine, I-281
 2-Isopropyl-5-(2-methylpropyl)pyrazine, I-282
 2-Isopropyl-5-methylpyrazine, I-283
 2-Isopropyl-6-methylpyrazine, I-284
 1-Isopropyl-3-methylpyrrolidine, M-544
 11-Isopropyl-5-methyl-5*H*-quinoline, I-279
 N-Isopropyl-N'-[6-(methylthio)hexyl]urea, D-828
 N-Isopropylnormicotine, P-957
 Isopropylpyrazine, I-285
 5-Isopropyl-2-pyrazinol, H-546
 6-Isopropyl-2-pyrazinol, H-547
 3-Isopropyl-2-pyrazinol, H-545
 Isoprosopinine A, I-286
 Isoprosopinine B, I-287
 Isoprostephabyssine, P-658
 Isoprotostemonine, P-670
 Isopsychotridine A, I-288
 Isopsychotridine B, I-288
 Isopsychotridine C, I-288
 Isopsychotridine E, I-289
 Isopsyllaborine A, P-749
 Isopteleflorine, I-290
 Isoptelefolidine, D-715
 Isoptelefoline, P-625
 Isoptelefolonium, D-720
 Isopteleine, F-211
 Isoptelepreline, D-715
 Isopterophorine, P-770
 Isopteropodic acid, M-661
 Isopteropodine N-oxide, M-661
 Isopteropodine, M-661
 Isoptilocaulin, I-291
 Isopuloupone, P-789
 Isopumiliotoxin A, A-416
 Isopycnarrhine, D-421
 Isopyoverdin Pp BTP 1, I-292
 Isopyoverdin Pp CFML 90-33, I-293
 Isopyoverdin Pp CFML 90-44, I-294
 Isopyroine, I-295
 Isopyrrolnitrin, P-961
 Isopyruthaldine, I-296
 Isopyruthaline, I-297
 Isopyruthaldine, I-298
 Isopyruthaline, I-299
 Isoquinocycline A, I-300
 Isoquinocycline B, I-300
 7,8-Isoquinolinediol, I-303
 6,7-Isoquinolinediol, I-302
 Isoquinoline, I-301
 1,5,6,7-Isoquinolinetriol, I-304
 5,6,7-Isoquinolinetriol, I-307
 1,6,7-Isoquinolinetriol, I-305
 1,6,8-Isoquinolinetriol, I-306
 1-(8-Isoquinolinylmethyl)pyridinium(1+), I-308
 Isorauhimbine, Y-13
 Isorauhimbic acid, Y-13
 Isoraumacline, R-22
 Isoraunesine, D-687
 3-Isorauniticine pseudoindoxyl, T-289
 3-Isorauniticine, A-212
 Isoraunitidine, R-53
 Isoraouwolsine, Y-13
 Isoregecoline, R-37
 Isoregelinone, R-38
 Isoremerine, R-120
 Isoreserpine oxindole A, C-117
 Isoreserpine oxindole B, C-117
 Isoreserpine pseudoindoxyl stereoisomer 1, T-289
 Isoreserpine pseudoindoxyl stereoisomer 2, T-289
 Isoreserpine, R-51
 3-Isoreserpine, R-52
 3-Isoreserpinine, R-53
 Isoretrohoustine, T-188
 Isoretrocanol, H-629
 (±)-Isoretrocanol, H-629
 (+)-Isoretrocanol, H-629
 Isoretrocanolic acid, P-960
 Isoretrocanyl *trans*- β -methylthioacrylate, P-510
 Isoretrocanyl tiglate, H-629
 Isoretulinal, R-63
 Isoretuline, R-63
 Isorhazicine, I-309
 Isorheagenine, R-86
 Isorhoeadine, R-86
 Isorhoeagenine, R-86
 Isorhynchophyllic acid, R-90
 Isorhynchophylline N-oxide, R-90
 Isorhynchophylline, R-90
 Isorhynchophyllic acid, R-90
 Isorhynchophylline N-oxide, R-90
 Isorhynchophylline, R-90
 Isorobustiline, F-143
 Isoroemerianine, O-112
 Isoroemerine, R-120
 Isorosibiline, R-131
 Isorosmarinine, R-133
 Isorotundifoleine, R-134
 Isorotundifoline, R-134
 Isorubijervine, I-310
 Isorubijervosine, I-310
 12*E*-Isorumbin, R-160
 Isosalsolidine, D-627
 Isosalsoline, T-181
 Isosalsolinol, T-163
 Isosalviamine A, I-311
 Isosalviamine B, I-312
 Isosandwicine, A-216
 Isosaracodine, P-601
 Isosaraine 1, S-69
 Isosaraine 2, S-70
 Isosaraine 3, S-71
 Isosarcodine, P-601
 Isosarcotrine E, S-83
 Isosarcotrine F, S-84
 Isosaxorumamide, S-112
 Isoschizogaline, I-313
 Isoschizogamine, I-313
 Isoschumanniphytine, I-314
 Isoscoulerine, I-315
 Isosogoline A, I-316
 Isoselagine, H-375
 Isosenaetine, S-232
 Isosendaverine, S-234
 Isosenkirkine, S-245
 Isosepiapterin, S-246
 Isosetoclavine, S-273
 Isosevanine, I-317
 Isosinoacutine, S-39
 Isosinococuline, S-311
 Isosinomenine, I-318
 (16*R*)-Isositsirikine (4*S*)-N-oxide, I-319
 (16*S*)-Isositsirikine (4*S*)-N-oxide, I-319
 Isositsirikine, I-319
 19*Z*-Isositsirikine, I-319
 (16*S*)-Isositsirikine, I-319
 Isosolacapine, S-336
 20-Isosolafloridine, S-338
 25-Isosolafloridine, S-338
 Isosolanogantamine, A-908
 Isosolaseaforthine, S-351
 Isosolasonine, S-456
 Isosolasuaveoline, S-456
 Isosolenopsin A, M-584
 Isosoporphamine, S-376
 Isosoporphidine, I-320
 Isosorelline, S-383
 Isosparsomycin, S-384
 α -Isosparteine, I-321
 β -Isosparteine, I-322
 Isospeciofoline, R-134
 Isospecionoxeine, R-134

- Isospeciophylline, M-661
 6-Isospectraline, S-388
 Isospiropachysine, S-436
 Isosplendine, S-594
 Isosplendoline, S-605
 Isostemofoline, S-515
 Isostemonamide, S-517
 Isostemonamine, S-518
 Isostemonidine, I-323
 Isostemonine, I-324
 Isostemotinine, S-524
 Isostenine, S-528
 Isostephaboline, M-233
 Isostephodeline, I-325
 Isostrychnine I, P-672
 Isostrychnine *N*⁸-oxide, P-672
 Isostrychnine, P-672
 Isostrychnobiline, S-591
 Isostrychnofoline, S-595
 Isostrychnopentamine A, S-601
 Isostrychnopentamine B, I-326
 Isostrychnophylline, S-602
 Isostrychnosplendine, S-605
 Isosuccinimide β-D-glucoside, D-460
 Isosuccinimide, D-460
 Isosungucine, S-642
 Isotachin A, M-577
 Isotachin B, M-577
 Isotachin C, M-577
 Isotaifine, I-327
 Isotalatizidine, M-352
 Isotazettine, P-626
 Isotazettinol, T-53
 Isotecleoxine, F-221
 Isoteinemine galactoside, S-200
 Isoteinemine, S-200
 Isotembetarine, C-728
 Isotenucaulin A, A-716
 Isotenuipine, T-81
 Isoterihanine, N-239
 Isotetrahydrohistrionicotoxin, H-309
 Isotetrandrine *N*-2'-oxide, I-328
 Isotetrandrine, I-328
 1-Isothalicberine, I-329
 Isothalicberine, I-329
 Isothalicrine, I-330
 Isothalictrine, I-331
 Isothalidezine, H-148
 Isothaliphine, T-331
 Isothalirine, I-332
 Isothalisopavine, I-333
 Isothalamidine, I-334
 Isothapsakin B, T-349
 Isothapsakone A, I-335
 Isothebaidine, T-525
 Isothebaine *N*-oxide, T-525
 Isothebaine, T-525
 8*H*-Isothiazolo[5,4-*b*]indole, I-336
 10α-Isothiocyanoalloaromadendrane, A-1571
 1-Isothiocyanoaromadendrane, A-1447
 7-Isothiocyanoato-2,10-bisaboladiene, I-214
 11-Isothiocyanoato-5-eudesmene, F-129
 6-Isothiocyanoato-4(15)-eudesmene, I-218
 4-Isothiocyanoato-11-gorgonene, A-777
 4-Isothiocyanoatomaaliane, E-100
 3-Isothiocyanoato-4-methoxy-1*H*-indole, I-337
 3-Isothiocyanoato-4-methoxy-1*H*-indol-5-ol, H-548
 Isothomandersine, T-195
 Isotiliarine, T-406
 Isotinctrine, T-412
 Isotriglochimin, T-515
 Isotrikentrin B, I-338
 Isotrilobine 2-*N*-oxide, I-339
 Isotrilobine, I-339
 Isotriphyophylline, D-793
 Isotripiperidine, I-340
 δ-Isotropylcocaine, T-635
 γ-Isotropylcocaine, T-634
 Isotryptoquivaline, T-645
 Isotuberostemonine, I-341
 Isotuboflavine, N-302
 Isotubulosine, T-675
 Isotussilage, H-231
 Isotussilaginic acid, H-231
 Isotussilagine, H-231
 Isotylocrebrine *N*-oxide, I-342
 (10*S*)-Isotylocrebrine *N*-oxide, I-342
 (10*S*,13*aR*)-Isotylocrebrine *N*-oxide, I-342
 Isotylocrebrine, I-342
 Isoundecyl 5-hydroxyanthranilate, A-789
 Isoursuline, D-626
 Isouvariopsine, I-343
 Isovaleramide, M-409
 Isovaleric acid, M-409
 7-Isovaleroylintermediate, I-160
 7-Isovaleroylylcoptamine, I-160
 13-Isovaleroyloxylylupanine, H-554
 6-Isovaleroyloxytropan-3-ol, M-391
 9-(3'-Isovaleroyl)viridiflorylretrocine, I-160
N-Isovaleryl-*N*-depropionyldecorticasine, L-234
 7-Isovaleryloxy-3-tigloyloxytropan-6-ol, M-392
 Isovallesiachotamine, V-10
 Isovanillobiocin, V-18
 Isovenenatine, V-50
 20-Isoveratramine, V-71
 Isoverbamedine, P-673
 Isoverbamekrine, P-665
 Isoverbamethine, P-665
 Isoverbametrine, P-665
 Isoverbaskirine, P-673
 Isoverbasitrine, P-673
 Isoverticine β-*N*-oxide, C-298
 Isoverticine, C-298
 Isovincoside, S-583
 Isovoacangine, I-344
 Isovoacristine, I-344
 Isovoacryptine, I-344
 Isovoafolidine, V-175
 Isovoafoline, V-175
 Isovorin, F-121
 Isowilfordine, P-250
 Isowilfortrine, P-249
 Isowillardiine, I-345
 Isoxantholumazine, P-763
 Isoxanthopterin, A-751
 Isoxanthopterincarboxylic acid, A-752
 4-Isoxazolecarbonitrile, I-346
 4-Isoxazolecarboxylic acid, I-346
 3-Isoxazolidinone, I-347
 3-Isoxazolidone, I-347
 2-Isoxazolin-5-one, I-349
 β-(Isoxazolin-5-on-2-yl)alanine, A-850
 3-(Isoxazolin-5-on-4-yl)alanine, A-733
 4-Isoxazolol, I-348
 5-Isoxazolol, I-349
 5(4*H*)-Isoxazolone, I-349
 5(2*H*)-Isoxazolone, I-349
 4(5*H*)-Isoxazolone, I-348
 Isoyohimbine, Y-13
 Istanbulamine, P-684
 IT 143A, P-421
 IT 143B, P-421
 IT 143C, P-421
 IT 143D, P-421
 Itampolin A, I-350
 Itampolin B, I-350
 Itomanindole A, D-304
 Itomanindole B, D-304
 Ivin, D-775
 Ivonine, I-351
 Ivorine, C-178
 Iwamide, A-1446
 Izmirine, C-790
 Iznikine, H-366
 J 104303, A-1393
 Jaborandine, J-1
 Jacobine, J-2
 Jacodine, S-240
 Jacoline, J-3
 Jaconine, J-4
 Jacozine, J-2
 Jacpaniculine, J-5
 Jaculadine, J-6
 Jacularine, A-498
 Jadiffine, J-7
 Jadamycin A, J-9
 Jadamycin Ala, J-10
 Jadamycin B, J-9
 Jadamycin F, J-11
 Jadamycin S, J-12
 Jadamycin T, J-13
 Jadamycin V, J-8
 Jafrine, J-14
 Jaluenine, H-183
 Jamaicamide A, J-15
 Jamaicamide B, J-15
 Jamaicamide C, J-15
 Jamaicensine, A-1013
 Jamaidine, H-554
 Jamine, J-16
 Jamtine *N*-oxide, J-17
 Jamtine, J-17
 Jamtinine, J-17
 Janetine, G-211
 Janolusimide, J-18
 Janoxepin, J-19
 Janthitrem A, J-20
 Janthitrem B, J-21
 Janthitrem C, J-21
 Janthitrem E, J-21
 Janthitrem F, J-21
 Janthitrem G, J-21
 Janussine A, J-22
 Janussine B, J-22
 Japaconitine A, M-331
 Japaconitine B, M-331
 Japaconitine B₁, M-343
 Japaconitine C, M-343
 Japaconitine C₁, M-343
 Japindine, J-23
 Japonine, D-716
 Jasminidine, J-24
 Jasminiflorine, F-163
 Jasminine, J-25
N-Jasmonoyl DOPA, A-745
N-Jasmonoyldopamine, D-920
N-Jasmonoylleucine, L-153
N-Jasmonoyltyramine, H-667
 Jaspamide B, J-26
 Jaspamide C, J-26
 Jaspamide D, J-26
 Jaspamide E, J-26
 Jaspamide F, J-26
 Jaspamide G, J-26
 Jaspamide, J-26
 Jaspisamide A, H-15
 Jaspisamide B, H-15
 Jaspisamide C, H-15
 Jaspikinolide, J-26
 Jateorhizine, P-34
 6(17),11-Jatrophadiene-2,3,5,7,8,9,14,15-octol, J-27
 Jatropham 5-(6-*p*-coumaroylglucoside), D-449
 Jatropham, D-449
 Jatrophine, D-449
 Jatrorrhizine, P-34
 Jatrorrhizrubine, P-34
 JatroRubine, P-34
 Javaberine A, D-541
 Javaberine B, D-540
 Javacarboline, J-28
 Javanicunine A, J-29
 Javanicunine B, J-29
 Javanine, J-30
 Javaniside, J-31
 Javaphylline, J-32
 Jaxartinine, J-33
 JBIR-02, P-421
 Jenamidine A, J-34
 Jenamidine B, J-34
 Jenamidine C, J-34
 Jerusalemine, J-35
 Jervine, J-36
 Jervin-11β-ol, J-36
 Jervin-11-one, J-36
 Jervinone, J-36
 Jesaconine, M-331
 Jesaconitine, M-331
 Jhelumine, J-37

- Jietacin A, J-38
 Jietacin B, J-39
 Jilkon, G-3
 Jineol, Q-34
 Jingsimycin, J-40
 Jiufengdine, M-309
 Jiufengsine, J-41
 Jiufengsine, J-41
 Jiufengtine, M-309
 JM 971A, T-699
 JM 971B, T-699
 JM 971C, T-699
 Joalin, J-42
 Jobertine, C-115
 Johnsonine, R-46
 Joint, H-715
 Jolantamine, J-43
 Jolantidine, K-36
 Jolantimine, J-44
 Jolantine, J-43
 Jolantinine, J-45
 Jollyanine, C-658
 Jollyanine, F-18
 Jonctum, H-715
 Jonquilline, L-348
 Joramine, J-46
 Jordanine, J-47
 Jorumycin, J-48
 Jorunnamycin A, R-41
 Jorunnamycin B, R-41
 Jorunnamycin C, R-41
 Joscina, S-159
 Josephinine, A-670
 Joubertiamine, J-49
 Joubertinamine, J-50
 JSTX-1, N-182
 JSTX-2, N-183
 JSTX-3, N-165
 JSTX-4, N-184
 Jubanine A, J-51
 Jubanine B, J-52
 Jubanine C, J-53
 Jubanine D, P-27
 Julandine, J-54
 Julibrine II, P-923
 Julibrine I, P-923
 Julifloricine, J-56
 Julifloridine, J-55
 Juliflorine, J-56
 Juliflorinine, J-56
 Juliprosine, J-57
 Juliprosinene, J-57
 Juliprosopine, J-56
 Julocrotine, J-58
 Julocrotol, J-58
 Julocrotone, J-58
 Junceine, T-488
 Junosidine, J-59
 Junosine, T-580
 Juripidine, A-909
 Jurubidine, S-458
 Jurubine, J-60
 Jusbetonin, I-112
 Jussiaeiine A, J-61
 Jussiaeiine B, M-79
 Jussiaeiine C, M-79
 Jussiaeiine D, M-79
 Justiciamide, J-62
 Juziphine *N*-oxide, J-63
 Juziphine, J-63
 Juzirine, H-424
 Jynosine, D-221
 K 252a, A-1189
 K 82A, L-78
 K 16A, M-76
 K 252b, A-1189
 K 73, C-901
 K 252c, A-1190
 K 252d, A-1190
 K 93-0711-I₁, M-34
 K 93-0711-I₂, M-34
 K 117, N-180
 Kabiramide A, K-2
 Kabiramide B, K-2
 Kabiramide C, K-2
 Kabiramide D, K-2
 Kabiramide E, K-2
 Kabiramide F, K-2
 Kabiramide G, K-2
 Kabiramide H, K-3
 Kabiramide I, K-1
 Kafamycin, C-13
 Kahakamide A, N-138
 Kahakamide B, N-138
 Kainic acid, K-4
 α -Kainic acid, K-4
 α -Kaininic acid, K-4
 Kaitocephalin, K-5
 Kalashine, K-39
 Kalasinamide, K-6
 Kalbreclasin, N-37
 Kalbretorine, H-290
 Kalidine, C-119
 Kalkitoxin, K-7
 Kamaline, D-531
 Kamassine, Q-6
 Kanagawamicin, K-8
 Kanchanomycin, A-244
 Kapakahine A, K-9
 Kapakahine B, K-10
 Kapakahine C, K-11
 Kapakahine D, K-11
 Kapakahine E, K-12
 Kapakahine F, K-10
 Kapakahine G, K-13
 Kaptax, B-79
 Karachicine, K-14
 Karachine, K-15
 Karacolidine, M-349
 Karacoline, M-355
 Karaconitine, M-330
 Karakanine, G-152
 Karakin, G-104
 Karakolidine, M-349
 Karakoline, M-355
 Karakomine, N-30
 Karakoramine, K-16
 Karantonin, G-3
 Karasamine, M-355
 Karnamicin A₃, A-1315
 Karnamicin A₁, K-17
 Karnamicin A₂, K-18
 Karnamicin B₁, K-17
 Karnamicin B₂, K-17
 Karnamicin B₃, K-18
 Karnamicin C₁, K-17
 Karnamicin C₂, K-19
 Karnamicin C₃, K-20
 Karnamicin C₄, K-20
 Karnamicin C₅, K-17
 Karnamicin D₁, A-1315
 Karnamicin D₂, K-21
 Karnamicin D₃, K-20
 Karnamicin D₄, K-21
 Karounidiol, M-730
 Kashmirine, C-298
 Katine, A-869
 Katsuyama base I, H-163
 Kauluamine, K-22
 Kayawongine, K-23
 K01-0509B, G-185
 KC 7367A, M-85
 KC 7367B, M-85
 Kealiinine A, K-24
 Kealiinine B, K-24
 Kealiinine C, K-24
 Kealiquinone, K-25
 Keayanidine A, S-398
 Keayanidine B, S-398
 Keayanidine C, S-398
 Keayanine A, S-399
 Kedarcidin, K-26
 Keenamidine A, K-27
 Keitaone, K-28
 Keitine, K-28
 Kelfer, D-632
 Kemamide O, O-37
 Kemamide S, O-32
 Kenidine, G-81
 Kenigicine, G-81
 Kenigine, G-81
 Kenimbidine, G-81
 Kenimbine, G-81
 Kenine, G-81
 Kephrine, A-736
 Keramadine, K-29
 Keramide B, K-30
 Keramide D, K-31
 Keramide E, K-32
 Keramide F, K-33
 Keramide G, K-33
 Keramide H, K-34
 Keramide J, K-34
 Keramide K, K-34
 Keramide M, K-31
 Keramide N, K-32
 Keramide A, M-91
 Keramide B, M-91
 Keramide C, M-93
 Keramaphidin B, I-142
 Keramaphidin C, A-1598
 Keronopsin A₁, K-35
 Keronopsin A₂, K-35
 Keronopsin B₁, K-35
 Keronopsin B₂, K-35
 Kesselridine, K-36
 Kesselringine, K-36
 3-Ketoadoaciaquinone B, A-147
 14-Ketoalstonidine, A-650
 Ketoanhydrokinamycin, K-37
 16-Ketoaspergillimide, A-1486
 1-Ketodelphisine, M-323
 11-Ketofistularin 3, F-70
 Ketogaze, A-736
 α -Keto- δ -guanidinovaleic acid, G-207
 Ketole (obsol.), I-79
 α -Keto- γ -methyleneglutamic acid, M-459
 31-Keto-12,34-oxa-32,33-dihydroirincinal A, K-38
 4-Ketopipecolic acid, O-196
 3-Ketopropyl-19*R*-heyneanine, O-199
 Ketotetrahydronorharmine, T-226
 KF 8940, H-133
 KG 431A, A-1191
 KG 431B, I-100
 Khaochamide, P-780
 Khaplofoline, H-56
 Khasianine, S-456
 Khyberine, K-39
 Kibataline, P-93
 20-*epi*-Kibataline, P-93
 Kibdelone A rhamnoside, K-40
 Kibdelone A, K-40
 Kibdelone B rhamnoside, K-40
 Kibdelone B, K-40
 Kibdelone C rhamnoside, K-40
 Kibdelone C, K-40
 Kifunensine, K-41
 Kigamicin A, K-42
 Kigamicin B, K-42
 Kigamicin C, K-42
 Kigamicin D, K-42
 Kigamicin E, K-42
 Kigamicinone, K-42
 Kikemanine, C-677
 Kikumycin A, K-43
 Kikumycin B, K-43
 Killarytoxin 3, A-1600
 Kimbasine A, K-44
 Kimbasine B, K-44
 Kimorexin A, K-45
 Kimvulline, I-4
 Kinabaline, T-564
 Kinabalurine A, I-62
 Kinabalurine B, I-62
 Kinabalurine C, I-62
 Kinabalurine D, I-62
 Kinabalurine E, I-62
 Kinabalurine F, I-62
 Kinabalurine G, H-734

- Kinganone, D-447
 Kiridine, K-46
 Kirigamycin C, K-47
 Kirilowin A, G-104
 Kirilowin B, G-104
 Kirilowin G, G-104
 Kirimine, A-89
 Kirinine A, L-105
 Kirinine B, K-48
 Kirinine C, L-105
 Kiritine, D-187
 Kirrine, P-528
 Kirondrine, K-49
 Kisantamine, A-811
 Kisantine, K-50
 Kisantine, K-51
 Kitagine, K-52
 Kitraline, V-124
 Kitramine, V-124
 Klugine, E-72
 Knightalbinol, H-428
 Knightinol, H-429
 Knightolamine, H-428
 Knightoline, B-95
 Kobusine, H-167
 Kobutimycin A, K-53
 Kobutimycin B, K-54
 Kodein, C-555
 Kodocytocalasin 1, Z-37
 Kodocytocalasin 2, Z-37
 Koenidine, G-81
 Koenigicine, G-81
 Koenigine, G-81
 Koeniginequinone A, K-55
 Koeniginequinone B, K-55
 Koenimbidine, G-81
 Koenimbine, G-81
 Koenine, G-81
 Koenoline, H-603
 Kohatamine, K-56
 Kohatine, K-56
 Kojistatin A, C-936
 Kokusagine, F-223
 Kokusagine, F-221
 Kokusaginoline, K-57
 Kolkamin, D-92
 Komarine, K-58
 Komaroidine, T-219
 Komaroine, K-59
 Komarovicine, Q-55
 Komarovidine, Q-55
 Komarovidinine N^3 -oxide, I-114
 Komarovidinine, I-114
 Komarovine, Q-55
 Komarovinine, Q-54
 Komavicine, K-60
 Komavine, K-61
 Konbamidin, K-62
 Konbuacidin A, P-18
 Konbuacidin B, P-18
 Kondurangamine A, P-603
 Kondurangamine B, P-603
 Kongboenine, M-312
 Kongboensine, M-355
 Kongboentine A, K-63
 Kongboentine B, M-323
 Koniamborine, K-64
 Kopsamidine A, K-74
 Kopsamidine B, K-78
 Kopsamine N^1 -oxide, K-74
 Kopsamine, K-74
 Kopsan-10,22-dione, D-805
 Kopsan-22-ol, K-65
 Kopsanol, K-65
 Kopsan-22-one, K-65
 Kopsanone, K-65
 Kopsaporine, K-66
 Kopsidasine N^7 -oxide, K-67
 Kopsidasine, K-67
 Kopsidasinine, K-68
 Kopsidine A, S-309
 Kopsidine B, S-309
 Kopsidine C, S-309
 Kopsidine D, S-309
 Kopsifine, K-69
 Kopsiflorine N^4 -oxide, K-74
 Kopsiflorine, K-74
 Kopsifoline A, K-70
 Kopsifoline B, K-70
 Kopsifoline C, K-70
 Kopsifoline D, K-70
 Kopsifoline E, K-70
 Kopsifoline F, K-70
 Kopsijasmine, K-71
 Kopsijasminilam, K-72
 Δ^{14} -Kopsijasminilam, K-72
 Kopsijasminine, K-71
 Kopsilactone, K-73
 Kopsilongine N^4 -oxide, K-74
 Kopsilongine, K-74
 Kopsilosine A, K-66
 Kopsilosine B, K-66
 Kopsilosine C, K-66
 Kopsilosine D, K-66
 Kopsilosine E, K-66
 Kopsilosine F, K-66
 Kopsilosine H, K-66
 Kopsilosine I, K-66
 Kopsilosine J, V-49
 Kopsimaline A, K-66
 Kopsimaline B, K-66
 Kopsimaline C, K-66
 Kopsimaline D, K-66
 Kopsimaline E, K-66
 Kopsinaline, K-74
 Kopsinarine, K-75
 Kopsine, K-76
 Kopsinganol, K-66
 Kopsingarine, K-66
 Kopsingine, K-66
 Kopsinginine, K-66
 Kopsingiol, K-80
 Kopsinic acid methosalt, K-78
 Kopsinicine, K-66
 Kopsinidine A, K-77
 Kopsinidine B, K-76
 Kopsinilamine, K-78
 Kopsinilam, K-78
 Kopsinine, K-78
 $\Delta^{16,17}$ -Kopsinine, K-71
 $\Delta^{14,15}$ -Kopsinine, V-49
 Kopsinitarine A, K-79
 Kopsinitarine B, K-79
 Kopsinitarine C, K-79
 Kopsinitarine D, K-79
 Kopsinitarine E, K-79
 Kopsinoline, K-78
 Kopsinol, K-66
 Kopsinone, K-80
 Kopsirachin, K-81
 Kopsoffine, P-521
 Kopsoffinol, P-521
 Kopsophonone, K-66
 Kopsone, K-82
 Kopsonoline, K-80
 Kopsorinine, D-805
 Kordiline, C-303
 Kororamide, K-83
 Korormicin A, K-84
 Korormicin B, K-85
 Korormicin C, K-86
 Korormicin D, K-87
 Korseldine, C-298
 Korselimine, C-304
 Korsemine, K-88
 Korseveramine, C-295
 Korseveridine, C-307
 Korseveridinine, C-307
 Korseveriline, C-295
 Korseverilinone, C-295
 Korseverine, C-302
 Korseverinine, C-302
 Korsevine, K-88
 Korsevinine, K-89
 Korsidine, C-302
 Korsiline, H-483
 Korsinamine, C-311
 Korsine N -oxide, C-311
 Korsine, C-311
 Korsinine, C-302
 Korundamine A, K-90
 Korupensamine A, K-91
 Korupensamine B, K-91
 Korupensamine C, K-91
 Korupensamine D, K-91
 Korupensamine E, K-91
 KOS 862, E-117
 Kosinostatin, I-300
 Kottamide A, K-92
 Kottamide B, K-92
 Kottamide C, K-92
 Kottamide D, K-93
 Kottamide E, K-94
 Koumicine, K-95
 Koumidine, T-425
 Koumine N -oxide, K-96
 Koumine, K-96
 Koumidine, K-97
 19*R*-Kouminol, K-96
 19*S*-Kouminol, K-96
 KR 025, M-807
 Krelagine, K-98
 Krepowine, C-748
 Kreysigine N -oxide, P-221
 (\pm)-Kreysigine, P-221
 (+)-Kreysigine, P-221
 (-)-Kreysigine, P-221
 Kreysiginine, K-99
 Kreysiginone, K-100
 Kribine, K-101
 Krigeine, K-102
 Krigenamine, K-102
 Krokiflorine, C-766
 Krokiflorinine, C-766
 Kromantine, A-1490
 Krukovine, P-301
 Kryptocurine, K-103
 KSB 1939H3, M-607
 KSB 193944, M-607
 KSK, I-163
 KSM 2690B, O-155
 KSM 2690C, O-155
 KT 5720, A-1189
 Kuafumine, K-104
 Kuanoniamine A, P-920
 Kuanoniamine B, D-93
 Kuanoniamine C, D-93
 Kuanoniamine D, D-93
 Kuanoniamine E, D-93
 Kuanoniamine F, D-93
 Kuchinoenamine, K-105
 Kukoamine A, S-399
 Kukoamine B, S-399
 Kukoline, S-313
 Kumokirine, P-39
 Kumujancine, H-456
 Kumujanrine, C-134
 Kumujansine B, K-106
 Kumujansine, K-106
 Kumujantine, K-106
 Kumujian A, C-132
 Kumujian C, C-131
 Kumujian G, D-686
 Kupitengester 1, P-220
 Kupitengester 3, P-218
 Kupitengester 2, T-259
 Kupitengester 4, T-259
 Kuramerine, K-107
 Kuraramine, K-108
 Kurasoin B, H-538
 Kurchaline, A-873
 Kurchamide, K-109
 Kurchamine, P-606
 Kurchenine, K-110
 α -Kurchessine, P-606
 Kurchessine, P-606
 Kurchicine, K-111
 Kurchildine, K-112
 Kurchiline, A-814

- Kurchimine, P-606
 Kurchine, C-595
 Kurchinidine, H-314
 Kurchinine, H-322
 Kurchiphyllamine, A-813
 Kurchiphylline, A-813
 Kurcholessine, K-113
 Kuroyuridine, K-114
 Kurramine 2'- α -*N*-oxide, K-115
 Kurramine 2'- β -*N*-oxide, K-115
 Kurramine, K-115
 Kutzneride 1, K-116
 Kutzneride 2, K-116
 Kutzneride 3, K-116
 Kutzneride 4, K-116
 Kutzneride 5, K-116
 Kutzneride 6, K-116
 Kutzneride 7, K-116
 Kutzneride 8, K-116
 Kutzneride 9, K-116
 Kuvan, T-142
 Kwangsine, K-117
 K₃-Ye, P-327
 Kynuramine, A-700
 Kynurenic acid, H-725
 Kyoberin, B-101
 Kyocristine, V-128
 L 11-1, A-1192
 L 13365, A-1193
 L 654040, A-1194
 L 681176, A-1195
 L 696475, A-1196
 L 697318, A-1196
 L 755807, A-1197
 L 970843, A-1198
 L 970844, A-1198
 L 52-71A, A-1304
 L 657398, B-89
 L 52-71B, A-1304
 L 892-4, C-837
 L 954967, N-269
 L 156586, O-131
 L 156587, O-131
 L 156588, O-131
 L 746628, P-584
 L 37231, V-128
 L 696474, Z-37
 Labiatamide A, L-1
 Labiatamide B, L-1
 Labradorin 1, A-622
 Labradorin 2, A-622
 Labrandine, M-650
 Labuanine A, L-2
 Laburnamine, A-906
 Laburnine acetate, H-629
 Laburnine benzoate, H-629
 Laburnine tiglate, H-629
 Laburnine, H-629
 Laburnoic acid, P-960
 Laccaic acid A, L-3
 Laccaic acid A₁, L-3
 Laccaic acid E, L-3
 Laccarin A, L-4
 Laccarin, L-4
 Lachnanthopyridone, L-5
 Lachnophyllic acid, D-129
 Lactacystin, L-6
 β -Lactam activator F2, B-402
 Lactic acid, O-32
 Lactimide, D-769
 Lactimidomycin, L-7
 Lactivicin, L-8
 Lactonamycin Z, L-9
 Lactonamycin, L-9
 6-Lactoyl-7,8-dihydropterin, S-246
 Laetanine, L-195
 Laetine, L-10
 Laetrile, H-673
 Laevisine A, P-247
 Laevisine B, P-252
 Lagerine, L-11
 Lagerstremine, V-88
 Lagerstroemine, V-88
 Lagumicine, A-228
 Lagumidine, C-589
 Lagunamine, T-673
 Lagunamycin, L-12
 Lahadinine A, L-13
 Lahadinine B, L-14
 Lahoramine, L-15
 Lahoricine, L-16
 Lahorine, L-17
 Lajollamycin, L-18
 Lakshminine, L-19
 Lallemanine, L-20
 Lamarckine, L-21
 Lamarckinine, A-101
 Lamarckinine, L-22
 Lamarkine, L-21
 Lambertine, L-23
 Lamellarin A, L-24
 Lamellarin B 20-sulfate, L-24
 Lamellarin B, L-24
 Lamellarin C 20-sulfate, L-24
 Lamellarin C, L-24
 Lamellarin D, L-24
 Lamellarin E, L-24
 Lamellarin F, L-24
 Lamellarin G 8-sulfate, L-24
 Lamellarin G, L-24
 Lamellarin H, L-24
 Lamellarin I, L-24
 Lamellarin J, L-24
 Lamellarin K, L-24
 Lamellarin L 20-sulfate, L-24
 Lamellarin α , L-24
 Lamellarin β , L-24
 Lamellarin χ , L-24
 Lamellarin ϵ , L-24
 Lamellarin η , L-24
 Lamellarin γ , L-24
 Lamellarin ϕ , L-24
 Lamellarin ζ , L-24
 Lamellarin L, L-24
 Lamellarin M, L-24
 Lamellarin N, L-24
 Lamellarin O, L-25
 Lamellarin P, L-25
 Lamellarin Q, L-25
 Lamellarin R, L-25
 Lamellarin S, L-24
 Lamellarin α 20-sulfate, L-24
 Lamellarin T 20-sulfate, L-24
 Lamellarin T, L-24
 Lamellarin U 20-sulfate, L-24
 Lamellarin U, L-24
 Lamellarin V 20-sulfate, L-24
 Lamellarin V, L-24
 Lamellarin W, L-24
 Lamellarin X, L-24
 Lamellarin Y 20-sulfate, L-24
 Lamellarin Z, L-24
 Lampranthin II, B-110
 Lamprolobine, L-26
 Lamuran, A-212
 Lanceine, L-27
 Lanceolin A, S-302
 Lanceolin B, S-302
 Lanceolin C, S-302
 Lanceomigine *N*⁴-oxide, L-28
 Lanceomigine, L-28
 Lanciferine, L-29
 Lanitine, L-30
 Lanitine, L-30
 Lanitine, L-30
 Lankanesine, L-31
 Lankongensisine A, L-32
 Lankongensisine B, L-33
 Lannotinidine A, L-34
 Lannotinidine B, F-30
 Lannotinidine C, F-29
 Lannotinidine D, F-29
 Lannotinidine E, L-35
 Lannotinidine F, L-35
 Lannotinidine G, L-36
 Lanopylin A₂, L-37
 Lanopylin B₂, L-37
 Lanopylin C₂, L-37
 Lanopylin D₂, L-37
 Lanopylin E₂, L-37
 Lanopylin F₂, L-37
 Lanopylins, L-37
 Lansamide 3, D-575
 Lansamide 4, D-575
 Lansamide 1, L-38
 Lansamide 2, L-39
 Lansimide 3, C-502
 Lansimide 4, C-502
 Lansimide 3, D-575
 Lansimide 4, D-575
 Lansimide 1, L-38
 Lansimide 2, L-39
 Lansine, D-555
 Lansiumamide A, L-38
 Lansiumamide B, L-38
 Lansiumamide C, P-344
 Lanuginosine, L-40
 Lanyuamide III, T-129
 Lanyuamide II, O-212
 Lanyuamide I, O-211
 Lanyuamide IV, H-745
 Lanyuamide VI, H-744
 Lanyuamide V, O-176
 Lapaconidine, A-91
 Lapatin A, L-41
 Lapatin B, L-42
 Lapidilectam, L-43
 Lapidilectine A, L-43
 Lapidilectine B, L-44
 Lapidilectinol, L-43
 Lappaconidine, A-91
 Lappaconine, A-91
 Lappaconitic acid, A-31
 Lappaconitine, A-91
 Largazole, L-45
 Larodopa, A-745
 Larutenine, L-46
 Larutensine, L-46
 Lasianine, A-834
 Lasiansine, M-312
 Lasiocarpine *N*-oxide, L-47
 Lasiocarpine, L-47
 Lasiiodine A, L-48
 Lasiiodine B, L-49
 Lasiocarpine, L-51
 Lastourvilline, L-52
 Lasubine II, L-53
 Lasubine I, L-53
 Latericine, H-275
 Lathyrol, T-556
 Latifine, L-54
 Latifolian A, L-55
 Latifolian B, L-55
 Latifoline *N*-oxide, H-4
 Latifoline, H-4
 Latifoline, L-56
 Latifolinine, L-57
 Latindine, L-58
 Latisodine, N-288
 Latisoline, N-288
 Latonduine A, L-59
 Latonduine B, L-60
 Latrunculin A, L-61
 Latrunculin B, L-62
 Latrunculin C, L-63
 Latrunculin D, L-63
 Latrunculin G, L-61
 Latrunculin H, L-62
 Latrunculin M, L-64
 Latrunculin S, L-65
 Latrunculin T, L-66
 Latumcidin, A-14
 Lauberine, L-67
 Laudanidine, L-68
 Laudanine methyl ether, L-69
 Laudanine, L-68
 Laudanosine, L-69
 Laudanosoline 4',6-dimethyl ether, R-59
 Laudanosoline, L-70
 Lauformine, L-214

- Laughine, L-71
 Launobine, B-397
 Laureline, L-72
 Laurelliptine, I-196
 Laurelliptin-1-hexadecanone, I-196
 Laurelliptin-1-octadecanone, I-196
 Laurepukine, P-781
 Lauric acid, D-889
 Laurifine, L-73
 Laurifinine, L-73
 Laurifoline, I-196
 Laurifoline, V-45
 Laurifonine, L-73
 Laurodionine, L-74
 Lauroitsine, B-243
 (-)-Lauroitsine, B-243
 Lauroscholtzine, L-75
 Laurostearic acid, D-889
 Laurotetanine, L-75
 Laurusin, F-132
 Lauterine, O-187
 Lavanducyanin, L-76
 Lavanduquinocin, L-77
 Lavendamycin, L-78
 Lavendustin A, L-79
 Lavendustin B, L-79
 Laxicymine, G-152
 Laxicyminine, G-152
 Laxicymisine, M-305
 LCB 29, H-488
 Ld 90, K-78
 Ld 91, K-78
 Ld 63, V-49
 Ld 65, V-49
 Ld 85, V-49
 LE 29060, V-110
 Lebeckianine, D-604
 Ledeboridine, R-9
 Ledeborine, R-9
 Ledebouridine, R-9
 Ledebourine, R-9
 Ledecorine, L-80
 Lederbourine, R-9
 Lederfolin, F-121
 Lederine, O-20
 Leepacine, V-113
 Lehmannine *N*-oxide, L-81
 Lehmannine, L-81
 Leinamycin, L-82
 Leiocarpadiol, L-83
 Leiocarpatriol A, I-123
 Leiocarpatriol B, I-123
 Leiocarpol, M-414
 Leiocarpone, M-414
 Leiodelide A, L-84
 Leiodelide B, L-85
 Leiokinine A, M-274
 Leiokinine B, E-267
 (-)-Lelobanidine III, L-86
 (-)-Lelobanidine II, L-86
 (\pm)-Lelobanidine I, L-86
 (-)-Lelobanidine I, L-86
 α -Lelobanidine, L-86
 Lelobanidine, L-86
 Lelobanololine, H-347
 Lemairamide, L-87
 Lemairamine, L-87
 Lemaireocereine, T-159
 Lemobiline, L-88
 Lemonnierin, L-89
 LEMONOMYCIN, L-90
 Lenamycin, B-447
 Lennoxamine, L-91
 Lenticellarine, L-92
 Lenticine, H-764
 Lentiginosine, O-46
 Lentoquine, Q-18
 Leofungine, V-34
 Leontalbamine, A-628
 Leontalbine, L-93
 Leontalbinine *N*-oxide, L-94
 Leontalbinine, L-94
 Leontamine, L-95
 Leonticine, L-96
 Leontidine, C-62
 Leontiformidine, L-97
 Leontiformine, L-97
 Leontine, A-628
 Leontismidine, L-98
 Leontismine, L-99
 Leonuridine, L-100
 Leonurine, H-432
 Leonurinine, L-101
 Lepadiformine A, C-926
 Lepadiformine B, L-102
 Lepadiformine C, L-102
 Lepadin A, L-103
 Lepadin B, L-103
 Lepadin C, L-103
 Lepadin D, L-104
 Lepadin E, L-104
 Lepadin F, L-104
 Lepadin G, L-104
 Lepadin H, L-104
 Lepedine, L-105
 Lepenine *N*-oxide, L-105
 Lepenine, L-105
 Lepetine, L-105
 Lepidamine, A-703
 Lepidiline A, D-750
 Lepidiline B, T-607
 Lepidine B, L-106
 Lepidine C, L-106
 Lepidine D, L-106
 Lepidine E, L-107
 Lepidine F, L-108
 Lepidine, L-106
 Lepidoporphyrin, P-771
 Lepidopterin, L-109
 Lepiotin A, L-110
 Lepiotin B, L-110
 Lepistine, L-111
 Leporine A, L-112
 Leporine B, L-112
 Leptactinine, L-113
 Leptaflorine, T-202
 Leptagline, L-114
 Leptanthine *N*-oxide, M-146
 Leptanthine, L-115
 Leptanthine, M-146
 Leptine III, L-116
 Leptine I, L-116
 Leptine IV, L-116
 Leptinidine 3-glucoside, L-116
 Leptinidine, L-116
 Leptinine III, L-116
 Leptinine II, L-116
 Leptinine I, L-116
 Leptinine IV, L-116
 Leptocarpine, L-117
 Leptocarpine, P-667
 Leptocarpinine, L-118
 Leptocladine, T-202
 Leptoclinidinone, A-1459
 Leptomerine, H-707
 Leptophyllin A, L-119
 Leptophyllin B, L-120
 Leptopidine, L-121
 Leptopidinine, H-766
 Leptopine, L-122
 Leptopinine, L-123
 Leptorhabine, C-562
 Leptosin A, L-124
 Leptosin B, L-125
 Leptosin C, L-126
 Leptosin D, L-127
 Leptosin E, L-128
 Leptosin F, L-129
 Leptosin G, L-132
 Leptosin G₁, L-130
 Leptosin G₂, L-131
 Leptosin H, L-133
 Leptosin I, L-134
 Leptosin J, L-134
 Leptosin K, L-137
 Leptosin K₁, L-135
 Leptosin K₂, L-136
 Leptosin M, L-139
 Leptosin M₁, L-138
 Leptosin N, L-139
 Leptosin N₁, L-140
 Leptosin O, L-141
 Leptosin P, L-141
 Leptosin Q, L-142
 Leptosin R, L-142
 Leptosin S, L-143
 Leptothoracine, M-544
 Lercheine, D-412
 Leroyine 14-acetate, M-348
 Leroyine, M-348
 Lespedamine, D-785
 Lettocine, L-144
 Lettowianthine, L-145
 Leucaenine, L-148
 Leucaenol, L-148
 Leucal, P-772
 Leucamide A, L-146
 Leucanthumsine A, M-323
 Leucanthumsine B, M-323
 Leucanthumsine C, M-323
 Leucanthumsine D, M-369
 Leucanthumsine E, M-336
 Leucascandrolide A, L-147
 Leucenine, L-148
 Leucenol, L-148
 Leucentine B, M-323
 Leucettamidine, L-149
 Leucettamine A, L-150
 Leucettamine B, L-151
 Leucettamine C, L-152
 Leucettidine, H-702
 Leucid, V-128
 Leuciglumer, L-153
 Leucinamide, L-153
 Leucine amide, L-153
n-Leucine homopolymer, L-153
l-Leucine homopolymer, L-153
 Leucine, L-153
 Leucinol, A-838
 Leucoagaricone, L-154
 Leucoatromentin, T-97
 Leucodopachrome, C-862
 Leuconine, A-96
 Leuconolam, L-155
 Leuconoxine, L-156
 Leucopterin, A-925
 Leucosar, F-121
 Leucosolenamide A, L-157
 Leucosolenamine B, L-157
 Leucostine A, M-309
 Leucostine B, M-340
 Leucostine, A-92
 Leucotamine, G-3
 Leucovernine, G-3
 Leucovorin calcium, F-121
 Leucovorin, F-121
 Leucoxine, L-158
 Leucoxylinone *N*-oxide, L-159
 Leucoxylinone, L-159
 Leueandine, F-147
 Leueantine A, M-303
 Leueantine C, M-352
 Leueantine D, M-323
 Leuhistin, L-160
 Leu, L-153
 Leupyrrin A₁, L-161
 Leupyrrin A₂, L-161
 Leupyrrin B₁, L-161
 Leupyrrin B₂, L-161
 Leupyrrin C, L-161
 Leupyrrin D, L-161
 Leurocolumbine, V-110
 Leurocristine, V-128
 Leuroformine, L-162
 Leurosidine *N*^B-oxide, V-110
 Leurosidine, V-110
 Leurosine *N*^B-oxide, L-162
F-Leurosine, L-162
 Leurosine, L-162

- Leurosinone, L-163
 Leurosivine, L-164
 Levarterenol, N-298
 Levycloserine, A-832
 Levodopa, A-745
 Levoglutamide, G-108
 Levoleucovorin calcium, F-121
 Levomefolate calcium, T-172
 Levomefolic acid, T-172
 Levomet, A-745
 Levopa, A-745
 Levophed, N-298
 Liaconitine A, M-364
 Liaconitine B, M-364
 Liaconitine C, M-364
 Liangshanine, N-30
 Liangshanone, N-30
 Liberine, L-165
 Libramycin A, M-432
 Licarpin, P-424
 Lichuanine, C-290
 Lichuanisinine, C-290
 Liconosine A, A-103
 Licorice glycoside E, D-581
 Lidimycin, L-166
 Lienkonine, L-167
 Liensinine, N-87
 N^b-Lignoceroyltryptamine, T-639
 Ligudentine, L-168
 Ligularidine, S-245
 Ligularine, L-169
 Ligularinine, P-516
 Ligularizine, L-170
 Ligustrazine, T-284
 Lihoudine, L-171
 Lilalinone, L-172
 Lilaline, L-173
 Lilidine, D-449
 Lilly 22641, D-234
 Lilly 32645, L-162
 Lilly 52230, P-961
 Lilly 112531, V-110
 Lilly 36781, V-110
 Lilly 49040, V-110
 Limaciamine, L-174
 Limacine 2 β -*N*-oxide, P-301
 Limacine 2' α -*N*-oxide, P-301
 Limacine 2' β -*N*-oxide, P-301
 Limacine, P-301
 Limacusine 2' β -*N*-oxide, L-175
 Limacusine, L-175
 Limalongine, L-176
 Limapodine, L-177
 Limaspermidine, L-177
 Limaspermine, L-177
 Limatine, A-1497
 Limatinine, A-1497
 Limipterin, A-749
 Limnanthenine, L-179
 Limocrocin, L-178
 Limogine, R-94
 Limousamine, C-805
 Linalolic acid, H-471
 Linalool-1-*oic* acid, H-471
 Linamarin gentiobioside, H-617
 Linamarin, H-617
 Linantenine, L-179
 Linaresine, R-156
 Lmarine, P-165
 Lmarinic acid, L-180
 Linavuline, L-181
 Lincangenine, A-1042
 Lindcarpine, T-248
 Lindechunine A, O-180
 Lindechunine B, P-211
 Lindelofamine, T-443
 Lindelofidine acetate, H-629
 Lindelofidine, H-629
 Lindelofine *N*-oxide, T-443
 Lindelofine, T-443
 Lindenialine, L-182
 Lindeniamine, L-183
 Lindenianine, V-153
 Lindheimerine, L-184
 Lindoldhamine, D-88
 Lineariline, A-85
 Linearilobine, M-352
 Linearisine, A-957
 Lineolone, T-268
N-Linolenoylglutamic acid, G-107
*N*²-Linolenoylglutamine, G-108
*N*²-(17-Linolenoyloxylinolenoyl)glutamine, H-645
*N*²-(17-Linolenoyloxylinoleoyl)glutamine, H-645
N-Linoleoyl-*L*-glutamic acid, G-107
*N*²-Linoleoylglutamine, G-108
*N*²-(17-Linoleoyloxylinolenoyl)glutamine, H-645
*N*²-(17-Linoleoyloxylinoleoyl)glutamine, H-645
 Linusitamarin, D-654
 Linustatin, H-617
 Lipoaconitine, M-331
 Lipo-14-*O*-anisoylbikhaconine, M-312
 Lipobikhaconitine, M-312
 Lipodeoxyaconitine, M-343
 Lipoforestine, M-312
 Lipogrammistin A, L-185
 Lipohypaconitine, M-343
 Lipoindaconitine, M-336
 Lipomesaconitine, M-331
 α -Lipomycin, L-186
 β -Lipomycin, L-186
 Lipopsudaconitine, M-336
 Lipopurealins, L-187
 Liposidomycin A-III, L-188
 Liposidomycin A-II, L-188
 Liposidomycin A-IV, L-188
 Liposidomycin B-III, L-188
 Liposidomycin C-III, L-188
 Liposidomycin C-II, L-188
 Liposidomycin C-I, L-188
 Liposidomycin C-IV, L-188
 Liposidomycin G-III, L-188
 Liposidomycin G-I, L-188
 Liposidomycin H-III, L-188
 Liposidomycin K-III, L-188
 Liposidomycin L-III, L-188
 Liposidomycin M-III, L-188
 Liposidomycin M-I, L-188
 Liposidomycin N-III, L-188
 Liposidomycin N-I, L-188
 Liposidomycin V-III, L-188
 Liposidomycin V-I, L-188
 Liposidomycin X-II, L-188
 Liposidomycin Y-III, L-188
 Liposidomycin Z-III, L-188
 Liposidomycin Z-I, L-188
 Liposidomycin, L-188
 Lipotril, C-419
 Lipoxamycin, L-189
 Lipoxazolidinone A, L-190
 Lipoxazolidinone B, L-191
 Lipoxazolidinone C, L-192
 Lipoyunaconitine, M-336
 Liquiritigenin, D-581
 Liridine, M-712
 Liridinine, T-521
 Lirinidine, D-531
 Lirinine *N*-oxide, T-521
 Lirinine, T-521
 Liriiodendron base, O-178
 Liriiodendronine, L-193
 Liriodenine, L-194
 Lirioferine, H-752
 Liritulipiferine, L-195
 Lissoclibadin 1, L-196
 Lissoclibadin 2, L-197
 Lissoclibadin 3, L-198
 Lissoclibadin 4, L-197
 Lissoclibadin 5, L-197
 Lissoclibadin 6, L-198
 Lissoclibadin 7, L-208
 Lissoclin A, D-823
 Lissoclin B, D-823
 Lissoclin C, L-199
 Lissoclin disulfoxide, L-209
 Lissoclinamide 10, L-203
 Lissoclinamide 1, L-200
 Lissoclinamide 4, L-201
 Lissoclinamide 5, L-201
 Lissoclinamide 6, L-201
 Lissoclinamide 7, L-201
 Lissoclinamide 8, L-201
 Lissoclinamide 9, L-202
 Lissoclinamide 2, U-9
 Lissoclinamide 3, U-9
 Lissoclinidide, L-204
 Lissoclin, L-209
 Lissoclinotoxin A, L-205
 Lissoclinotoxin B, L-206
 Lissoclinotoxin C, L-207
 Lissoclinotoxin D, L-208
 Lissoclinotoxin E, L-209
 Lissoclinotoxin F, L-210
 Litcubine, C-783
 Litcubinine, C-783
 Litebamine, L-211
 Lithosenine, L-212
 Lithospermoside, L-213
 Litsedine, B-397
 Litseferine, D-346
 Litseglutine A, D-346
 Litseglutine B, T-248
 Litsericine, L-214
 Litsoene, L-75
 Littoraline, L-215
 Littorine, L-216
 Lituarine A, L-217
 Lituarine B, L-217
 Lituarine C, L-217
 Liwaconitine, M-312
 LK6A, L-354
 LL C10037 α , A-804
 LL-AP-191 γ , L-90
 LL-BH872 α , A-1199
 LL-BM 123 γ ₁, C-459
 LL-BM 123 γ ₂, C-460
 LL-BM 123 β ₁, G-117
 LL-BO 2964 α , A-1200
 LL-BO 2964 β , A-1200
 LL-BO 2964 γ , A-1200
 LL-BO 2964, A-1200
 LL-D 42067 α , S-301
 LL-D 42067 β , S-301
 LL-E19085 α , C-473
 LL-14E605 α , G-126
 LL-E605 β , G-126
 LL-49F233 α , A-1201
 LL-F42248 α , P-974
 LMCA, T-172
 LMSR, T-172
 Lobelandine, L-218
 Lobelanine, L-218
 Lobelidine, L-219
 Lobeline, L-219
 α -Lobeline, L-219
 Lobinaline, L-220
 Lobinamidine, L-221
 Lobine, L-222
 Lobinine, L-221
 Loboofrutoside, S-447
 Lobofoanthamine, Z-30
 Lobran, L-219
 Lobscurinol, L-223
 Lochneram, S-89
 Lochnericine, T-11
 Lochneridine, L-224
 Lochnerine, S-89
 Lochnerinine, T-11
 Lochnerivine, L-225
 Lochrovincine, L-226
 Lochroidine, L-227
 Lochrovine, L-228
 Lochvinerine, S-89
 Locoine, L-229
 Loesenerine, L-230
 α -Lofoline, F-29
 β -Lofoline, F-29
 Lofoline, F-29
 Lokysterolamine A *N*³⁰-oxide, P-496
 Lokysterolamine A, P-496

- Lokysterolamine B, P-496
 Lolicine A, L-231
 Lolicine B, L-231
 Lolidine, L-232
 Lolilline, L-233
 Loline, L-234
 Lolinidine, L-235
 Lolinine *N*-oxide, L-234
 Lolinine, L-234
 Lolitrem A, L-236
 Lolitrem B, L-236
 Lolitrem C, L-236
 Lolitrem E, L-237
 Lolitrem F, L-236
 Lolitrem N, L-237
 Lolitriol, L-237
 Lomaiviticin B, L-238
 Lombrine, L-239
 Lombricine phosphate, L-240
 Lombricine, L-240
 Lomofungin, L-241
 Lomondomyacin, L-241
 Longamide A, L-242
 Longamide B, L-243
 Longamide, H-214
 Longanine, L-244
 Longanine, L-244
 Longanone, L-244
 Longatin, N-44
 Longeraciphylline A, L-245
 Longeraciphylline B, L-246
 Longetherine, L-247
 Longiberine, L-248
 Longicaudatine F, L-250
 Longicaudatine *N*-oxide, L-249
 Longicaudatine Y, L-250
 Longicaudatine Z, L-250
 Longicaudatine, L-249
 Longifolidine, L-251
 Longifolonine, L-252
 β -Longilobine, R-62
 α -Longilobine, S-240
 Longimammamine, T-161
 Longimammatine, T-196
 Longimammidine, T-198
 Longimammine, S-659
 Longimammosine, T-196
 Longistrobine, L-253
 Longistylumphylline A, D-55
 Longistylumphylline B, L-254
 Longistylumphylline C, L-255
 Longitubanine A, L-256
 Longitubanine B, L-256
 Longitubine, H-4
 Longtouconitine B, M-323
 Lonicerine, A-1492
 Lonicerose, D-204
 Lophanterine, L-257
 Lophilacrine, L-258
 Lophiline, L-259
 Lophiroside A₁, S-302
 Lophiroside A₂, S-302
 Lophiroside B₁, S-302
 Lophiroside B₂, S-302
 Lophocerine, L-260
 Lophocine, L-261
 Lophocladine A, H-686
 Lophocladine B, A-865
 Lophophorine, A-1021
 Lophotine, A-1021
 Lorajmine hydrochloride, A-216
 Lorajmine, A-216
 Lorneamide A, L-262
 Lorneamide B, L-262
 Loroquine glucoside, L-263
 Loroquine, L-263
 Losbanine, L-264
 Lotaustralin, H-572
 Lotthanongine, L-265
 Loturine, M-426
 Lotusanine A, A-150
 Lotusanine B, L-266
 Lotusine A, L-267
 Lotusine B, L-268
 Lotusine C, L-269
 Lotusine D, L-267
 Lotusine E, L-270
 Lotusine F, P-25
 Lotusine G, L-271
 Lotusine, H-275
 Louisianin A, L-272
 Louisianin B, L-272
 Louisianin C, L-272
 Louisianin D, L-272
 Louludinium(1+), L-273
 Loveraine, L-274
 Loxylostosidine A, X-31
 Loxylostosidine B, X-31
 L9, T-263
 LU 1, H-616
 Lucaconine, M-309
 Lucentamycin A, D-101
 Lucentamycin B, L-275
 Lucentamycin C, D-101
 Lucentamycin D, D-101
 Luciculine *N*-oxide, N-30
 Luciculine, N-30
 Lucidamine A, L-276
 Lucidine A, L-277
 Lucidine B, L-277
 Lucidinine, L-278
 Lucidioline, L-279
 Luciduline, L-280
 Lucidulinone, L-280
 Lucidusculine, N-30
Diplocardia Luciferin, A-892
Ophophorus Luciferin, C-560
Renilla Luciferin, C-560
Watasenia Luciferin, C-560
Cypridina Luciferin, L-281
Photinus Luciferin, L-282
 Lucilactaene, L-283
 Luciola oxyluciferin, F-56
 Luciopterin, P-763
 Lucuminamide, H-671
 Lucumin, H-673
 Lucuminoside, H-673
 Ludaconitine, M-336
 Lukianol A, L-284
 Lukianol B, L-284
 Lumazine, P-762
 Lumbricine, L-240
 Lumichrome, L-285
 Lumicolchicine II, L-287
 Lumicolchicine I, L-287
 α -Lumicolchicine, L-286
 β -Lumicolchicine, L-287
 Lumicolchicine, L-287
 γ -Lumicolchicine, L-287
 β -Lumispeciosine, L-287
 Lunacridine, L-288
 Lunacrine, B-13
 Lunacrinol, R-92
 Lunamaridine, L-289
 Lunamarine, L-290
 Lunariamine, L-291
 Lunaridine, L-292
 Lunarine, L-293
 Lunarinol II, L-293
 Lunasia II, R-92
 Lunasine, B-14
 Lundurine A, L-294
 Lundurine B, L-294
 Lundurine C, L-294
 Lundurine D, L-294
 Lunidine, L-295
 Lunidonine, L-295
 Lunine, L-296
 Lunolone, L-297
 Luotonine A, L-298
 Luotonine B, L-298
 Luotonine C, L-299
 Luotonine D, L-300
 Luotonine E, L-298
 Luotonine F, L-301
 Lupanidine, M-121
 Lupanine *N*-oxide, L-302
 Lupanine, L-302
 Lupanoline, I-322
 Lupetidine, D-770
 Lupilaxine, L-303
 Lupimidine (obsol.), S-385
 Lupinine, O-58
 Lurenine, L-304
 Lusitanine, L-305
 Lutamide A, T-460
 Lutamide C, T-460
 Luteanine, D-405
 Luteanine, I-208
 Luteicine, L-306
 Luteicinone, L-306
 Luteidine α -*N*-oxide, L-307
 Luteidine β -*N*-oxide, L-307
 Luteidine, L-307
 Luteine, K-36
 Luteinine, L-308
 Lutenurine, L-309
 Lutessine, L-310
 2,6-Lutidine, D-775
 2,6-Lutidine-3,5-dicarboxylic acid, D-776
 Lutine, A-1040
 Lutinine, L-311
 Luvenil, E-6
 Luxandrine, H-275
 Luzopeptin A, L-312
 Luzopeptin B, L-312
 Luzopeptin C, L-312
 Luzopeptin E₂, L-312
 LWTX1, H-731
 LWTX2, H-731
 LWTX3, H-731
 LWTX4, S-109
 LWTX5, S-109
 LWTX6, S-109
 LY 783281, A-1521
 LY 119863, V-110
 LY 99094, V-110
 Lyadine, L-313
 Lyalidine, L-314
 Lyaline, L-315
 Lyaloside, L-316
 Lyalosidic acid, L-316
 Lyaconitine, L-317
 Lycaonic acid piperidide, O-192
 Lycaonic acid pyrrolidide, O-192
 Lycaonic acid, O-192
 Lyceamine, L-318
 Lyciumide A, D-920
 Lyciumins, L-319
 Lycobergine, L-327
 Lycobetaine, U-34
 Lycocaroline, L-320
 Lycocernuine *N*-oxide, L-321
 Lycocernuine, L-321
 Lycoclavine, L-322
 Lycoctinine, M-309
 Lycoctonine, M-309
 Lycodiflexine, L-323
 Lycodine, L-324
 Lycodoline *N*-oxide, L-325
 Lycodoline, L-325
 Lycofawcine, L-326
 Lycoflexine, L-327
 Lycofoline, L-328
 Lycogalic acid, L-329
 Lycogarubin A, L-329
 Lycogarubin B, L-329
 Lycogarubin C, L-329
 Lycognidine, L-322
 Lycolucine, L-277
 Lyconadine A, L-330
 Lyconadine B, L-330
 Lyconesidine A, L-331
 Lyconesidine B, L-331
 Lyconesidine C, L-322
 Lyconnotine, L-332
 Lyconnotinol, L-333
 Lycopaniculatine, L-334
 Lycopecurine, L-335

- Lycoperine A, L-336
 Lycoperidine I, T-224
 Lycoperoside A, S-442
 Lycoperoside B, S-442
 Lycoperoside C, S-442
 Lycoperoside D, S-445
 Lycoperoside F, S-444
 Lycoperoside G, S-443
 Lycoperoside H, D-99
 Lycopersicin, S-445
 Lycophlegmarine, L-337
 Lycophlegmine, A-1029
 Lycoplamine A, L-338
 Lycoplamine B, L-339
 Lycoplamine C, L-339
 Lycoplamine D, L-340
 Lycopodamine A, L-335
 Lycopodamine B, A-1029
 Lycopodamine C, A-1029
 Lycopodine, L-341
 Lycoposerramine A, L-342
 Lycoposerramine B, L-343
 Lycoposerramine C, F-30
 Lycoposerramine D, L-327
 Lycoposerramine E, P-362
 Lycoposerramine F, L-344
 Lycoposerramine G, L-344
 Lycoposerramine H, L-322
 Lycoposerramine I, L-341
 Lycoposerramine J, L-322
 Lycoposerramine K, L-279
 Lycoposerramine L, L-322
 Lycoposerramine M, L-341
 Lycoposerramine N, L-325
 Lycoposerramine O, L-322
 Lycoposerramine P, F-30
 Lycoposerramine Q, F-30
 Lycoposerramine S, L-345
 Lycoposerramine U, L-327
 Lycoposerramine V, L-346
 Lycoposerramine W, L-346
 Lycoposerramine X, H-380
 Lycoposerramine Y, H-380
 Lycoposerramine Z, H-380
 Lycopsamine *N*-oxide, I-160
 Lycopsamine, I-160
 Lycoramine *N*-oxide, G-3
 Lycoramine, G-3
 Lycoremine, G-3
 Lycorenine, L-347
 Lycoricidine, N-37
 Lycoricidinol, N-37
 Lycorimine, G-3
 Lycorine 1,2-di-*O*- β -D-glucoside, L-348
 Lycorine 1-*O*- β -D-glucoside, L-348
 Lycorine, L-348
 Lycoriside, L-348
 Lycoserramine, L-349
 Lycoserrine, L-350
 Lycosinine A, L-351
 Lycosinine B, L-351
 Lycothunine, F-30
 Lycovatine A, L-352
 Lycoverticine, F-75
 Lydicamycin, L-353
 Lydimycin, L-166
 Lyfoline, V-95
 Lymphostin, L-354
 Lyngbyabellin A, L-355
 Lyngbyabellin B, L-356
 Lyngbyabellin C, L-357
 Lyngbyabellin D, L-358
 Lyngbyabellin E, L-359
 Lyngbyabellin F, L-360
 Lyngbyabellin G, L-361
 Lyngbyabellin H, L-359
 Lyngbyabellin I, L-360
 Lyngbyapeptin A, L-362
 Lyngbyapeptin B, L-363
 Lyngbyapeptin C, L-364
 Lyngbyastatin 4, L-365
 Lyngbyastatin 5, L-365
 Lyngbyastatin 6, L-365
 Lyngbyastatin 7, L-366
 Lyngbyatoxin A, L-367
 Lyngbyatoxin B, L-367
 Lyngbyatoxin C, L-367
 Lyophyllin, L-368
 Lysergamide, L-370
 Lysergene, L-369
 Lysergic acid amide, L-370
 Lysergic acid α -hydroxyethylamide, L-370
 Lysergic acid methylcarbinolamide, L-370
 Lysergic acid, L-370
 Lysergine, L-371
 Lysergol, L-371
 Lysergylvaline amide, L-372
 Lysergylvaline methyl ester, L-372
 Lysergylvaline, L-372
 Lysergylvalylleucylproline lactam, E-156
 Lysergylvalylphenylalanylproline lactam, E-154
 Lysergylvalylvalylproline lactam, E-152
 Lysianadioic acid, L-373
 Lysicamine, L-193
D-Lysinegalantin I, G-7
 Lysolipin I, L-374
 Lysolipin X, L-375
N-(*N*²-L-Lysyl)cantharimide, C-89
 Lysylpyridinoline, P-907
 Lythramine, L-376
 Lythrancepine III, L-377
 Lythrancepine II, L-377
 Lythrancepine I, L-377
 Lythrancine III, L-377
 Lythrancine II, L-377
 Lythrancine I, L-377
 Lythrancine IV, L-377
 Lythrancine VII, L-377
 Lythrancine VI, L-377
 Lythrancine V, L-377
 Lythrancine, L-377
 Lythranidine, L-378
 Lythranine, L-378
 Lythridine, V-95
 Lythrine, V-95
 Lythrumine, L-379
 M 259, A-1202
 M 146791, C-837
 M 4582, H-408
 M 193, S-498
 Maackiamine, T-228
 Maackinine, G-70
 Maandrosine, M-1
 Macamide A, O-189
 Macamide B, H-221
 Macaridine, M-2
 Macarpine, C-360
 Machigline, D-347
 Machilaminoside A, M-3
 Machilaminoside B, M-4
 Mackinazolinone, M-5
 Macleanine, M-6
 Macleyne, P-667
 Macolidine, G-235
 Macoline, G-235
 Macondine, D-624
 Macowine, M-108
 Macralstonidine, M-7
 Macralstonine, M-8
 Macrantaldehyde, N-45
 Macrantaline, N-45
 Macranthine, M-9
 Macrantoridine, N-45
 Macrocaffrine, M-10
 Macrocarpamine, M-11
 Macrocarpine A, T-20
 Macrocarpine B, T-20
 Macrocarpine C, T-20
 Macrocentridine, M-309
 Macrocentrine, M-12
 Macrocidin A, M-13
 Macrocidin B, M-13
 Macrodaphnidine, D-33
 Macrodaphnine, D-33
 Macrodaphniophyllamine, D-33
 Macrodaphniophyllidine, D-33
 Macrodasine A, M-14
 Macrodasine B, M-14
 Macrogentine, M-15
 Macroolidine, M-16
 Macroline, M-17
 Macromerine, M-18
 Macronecine, H-518
 Macronine, P-626
 Macrophominol, M-19
 Macrophylline A, M-22
 Macrophylline, H-518
 Macrophylline, M-20
 Macrophylline, M-21
 Macropodumine A, M-23
 Macropodumine B, M-24
 Macropodumine C, M-25
 Macropodumine D, M-26
 Macropodumine E, M-27
 Macropodumine F, Y-23
 Macropodumine G, Y-23
 Macropodumine H, D-33
 Macropodumine I, D-53
 Macrorhynine A, M-369
 Macrorhynine B, M-360
 Macrorine, I-44
 Macrorungine, M-28
 Macrosalbine, M-29
 Macrospegatine, M-30
 Macrostromine α -*N*-oxide, A-1403
 Macrostromine β -*N*-oxide, A-1403
 Macrostromine, A-1403
 Macrotomine, F-92
 Macroxine, M-31
 Macrozamin, M-397
 Maculine, F-221
 Maculosidine, F-222
 Maculosin 1, C-900
 Maculosin 6, C-876
 Maculosine 5, C-901
 Maculosine, F-221
 Maculotoxin, T-298
 Macusine A, P-562
 Macusine B, T-425
 Macusine C, A-229
 Madangamine F, M-32
 Madangamines A-E, M-33
 Madindoline A, M-34
 Madindoline B, M-34
 Madugin, T-639
 Madumicin II, M-35
 Madumicin I, M-35
 Madumycin II, M-35
 Madumycin I, M-35
 Madurensine, M-36
 Maedamine A, M-37
 Maedamine B, M-37
 Maeganedine A, M-38
 Magallanesine, M-39
 Magellanine, M-40
 Magellaninone, M-40
 Magnarcine, M-41
 Magnesidin, M-42
 Magniflorine, M-43
 Magnococline, G-143
 (+)-Magnocurarine, H-275
 (-)-Magnocurarine, H-275
 Magnoflorine, C-702
 Magnolamide, M-44
 Magnolamine, M-45
 Magnoline, G-177
 Magnoporphine, T-248
 Magnosprengerine, H-566
 Mahanimbicine, M-46
 Mahanimbidine, C-813
 Mahanimbilol, E-277
 (\pm)-Mahanimbine, M-47
 (+)-Mahanimbine, M-47
 Mahanimbinine, M-47
 Mahanimbinol, E-277
 Mahanimboline, M-47
 Mahanine, M-47
 Maingayine, H-322
 Maingayinine, M-48

- Maireistemoninol, T-665
 Maistemonine, M-49
 Maitansine, M-135
 Majarine, B-101
 Majdine, M-50
 Majdinine, T-288
 Majoridine, T-288
 Majorine, M-51
 Majorinine, V-194
 Majovine, M-52
 Majvinine, S-89
 Makalika ester, M-53
 Makalikone ester, M-53
 Makaluvamine A, M-54
 Makaluvamine B, M-54
 Makaluvamine C, M-54
 Makaluvamine D, M-54
 Makaluvamine E, M-54
 Makaluvamine F, M-55
 Makaluvamine G, M-54
 Makaluvamine H, M-54
 Makaluvamine I, M-54
 Makaluvamine J, M-54
 Makaluvamine K, M-54
 Makaluvamine L, M-54
 Makaluvamine M, M-54
 Makaluvamine N, M-54
 Makaluvamine O, D-13
 Makaluvamine P, M-54
 Makaluvic acid A, M-56
 Makaluvic acid B, M-56
 Makaluvic acid C, M-56
 Makaluvone, D-13
 Makomakine, H-310
 Makonine, A-1442
 Makrotomine, F-92
 Malacitanine, M-57
 Malagashanine, M-58
 Malagashanol, M-58
 Malagashine, S-604
 Malarboreine, M-59
 Malarborine, M-59
 Malasseziacitrin, M-60
 Malasseziaketone, M-67
 Malassezialactic acid, M-61
 Malasseziazole A, M-62
 Malasseziazole B, M-63
 Malasseziazole C, M-64
 Malassezindole A, M-66
 Malassezindole B, M-66
 Malassezin, M-65
 Malassezzone, M-67
 Malatyamine, M-68
 Malaxine, K-28
 Malbrancheamide A, M-69
 Maleimide monooxime, P-947
 Maleimide, P-947
 Maleimycin, M-70
 Malekulatine, M-71
 Malic acid, M-72
 Malindine, M-73
 Malkangunin, T-259
 Malkanguniol, T-259
 Malloapeltine, H-709
 Mallorepine, H-709
 Mallotusin, M-135
 Malonganenone A, M-74
 Malonganenone B, M-75
 Malonganenone C, A-916
 Malonganenone D, M-74
 Malonganenone E, M-74
 Malonganenone F, M-75
 Malonganenone G, M-75
 Malonganenone H, A-916
 Malonomycin, M-76
N-(*N*-Malonyl- β -alanyl)dopamine 3-glucoside, D-920
 2-(Malonylamino)benzoic acid, A-706
N-Malonyldopamine 3-glucoside, D-920
O-Malonylindolactam V, I-78
 6'-*O*-Malonylprunasin, H-673
N-Malonyltryptophan, T-640
 Malouetafrine, H-322
 Malouetamide, L-57
 Malouetine, P-601
 Malouphyllamine, C-595
 Malouphylline, D-269
 Malouphylline, D-269
 Malouphyllinol, D-269
 Maltophilin, X-6
 Maltoxazine, M-77
 Maltoxin, H-357
 Malyngamide C acetate, D-225
 Malyngamide C, D-225
 Malyngamide K, D-225
 Malyngamide W, M-78
 Mamamine *N*-oxide, M-79
 Mamanine, M-79
 MAM, M-397
 Mammillarinin, B-110
 Manadomanzamine A, M-80
 Manadomanzamine B, M-80
 Mandelamide, H-671
 Mandelonitrile gentiobioside, H-673
 Mandelonitrile rutinoid, H-673
 Mandelonitrile sophoroside, H-673
 Mandelonitrile, H-673
 Mangochinine, C-783
 Manibacanine, M-81
 Manicoline A, M-82
 Manicoline B, M-83
 Manihotoxin, H-617
 Manilamine, M-84
 Maniwamycin A, M-85
 Maniwamycin B, M-85
 Mannojirimycin, A-728
 Mannonojirimycin, A-728
 Mannopeptimycin α , M-86
 Mannopeptimycin β , M-86
 Mannopeptimycin δ , M-86
 Mannopeptimycin ϵ , M-86
 Mannopeptimycin γ , M-86
 Mannosylqueuosine, Q-9
 Manshurienine A, A-1429
 Manshurienine B, A-1427
 Manshuritine, M-331
 Manthidine, P-57
 Manthine, P-57
 Manumycin A, M-87
 Manumycin C, M-88
 Manumycin D, M-89
 Manzacidin A, M-90
 Manzacidin B, M-90
 Manzacidin C, M-90
 Manzacidin D, M-90
 Manzamine A *N*-oxide, M-91
 Manzamine A, M-91
 Manzamine B, M-92
 Manzamine C, M-93
 Manzamine D, M-91
 Manzamine E, M-91
ent-Manzamine F, M-91
 Manzamine F, M-91
ent-Manzamine G, M-91
 Manzamine G, M-91
 Manzamine H, M-94
 Manzamine J *N*-oxide, M-94
 Manzamine J, M-94
 Manzamine L, M-94
 Manzamine M, M-95
 Manzamine X, M-91
 Manzamine Y, M-91
 Maokonine, T-700
 Mappicine gentiobioside, M-96
 Mappicine glucoside, M-96
 Mappicine ketone, M-96
 Mappicine, M-96
 Mappine, B-393
 Marcanine A, M-97
 Marcanine B, M-97
 Marcanine C, M-97
 Marcanine D, M-97
 Marcanine E, M-97
 Marcfortine A, M-98
 Marcfortine B, M-98
 Marckidine, M-99
 Marckine, T-675
 Maremycin A, M-100
 Maremycin B, M-100
 Maremycin C₁, M-100
 Maremycin C₂, M-100
 Maremycin D₁, M-101
 Maremycin D₂, M-101
 Margaritarine, M-102
 Margetine, N-37
 Margrapine A, M-103
 Margrapine B, M-103
 Marikarin, A-197
 Marinamide, M-104
 Marine *Streptomyces* C₃₀H₃₈N₂O₅ lactam, M-105
 Marinobactins, M-106
 Marinobufagin 3-pimeloylarginine ester, E-132
 Marinobufagin 3-suberoylglutamine ester, E-132
 Marinobufagin, E-132
 Marinobufogenin O³-sulfate, E-132
 Marinobufogenin, E-132
 Marinobufotoxin, E-132
 Marinopyrrole A, M-107
 Marinopyrrole B, M-107
 Marinoquinoline A, M-552
 Maritidine, M-108
 Maritimamine, M-108
 Marmeline, O-63
 Maroumine, S-192
 Marqibo, V-128
 Marshaline, L-80
 Marshdine, T-243
 Marshmine, T-581
 Marstomentoside A, P-610
 Marstomentoside D, P-604
 Marstomentoside O, P-610
 Marstomentoside P, P-610
 Marstomentoside T, P-604
 Martefragine A, M-109
 Martensine A, M-110
 Martensine B, M-110
 Martinelliacid, M-111
 Martinelline, M-111
 Martsomentoside C, P-604
 Maruquine, M-661
 Masonine, O-68
 Massadine aziridine, M-113
 Massadine chloride, M-112
 Massadine, M-112
 Matadine, M-114
 Matamycin, A-664
 Matemone, M-115
Saccharomyces kluyveri α -Mating pheromone 1, M-116
 α ^{sk1} Mating pheromone, M-116
 Matlystatin D, M-117
 Matlystatin E, M-118
 Matlystatin F, M-119
 Matopensine mono-*N*-oxide, M-120
 Matopensine *N*-oxide, M-120
 Matopensine, M-120
 Matricaria acid, D-103
trans,trans-Matricaria ester, D-103
trans-Matricaria ester, D-103
 Matricaria ester, D-103
 Matricaric acid, D-103
 (5 β ,11 α)-Matridin-11-one, D-79
 (5 β ,6 β)-Matridin-15-one, I-320
 Matridin-15-one, M-121
 (5 β)-Matridin-15-one, S-380
 Matrine *N*-oxide, M-121
 Matrine, M-121
 Mattogrossine, M-122
 Mauiensine, T-288
 Mauritamide A, M-123
 Mauritiamine, M-124
 Mauritian A, M-125
 Mauritian B, M-126
 Mauritian C, M-127
 Mauritian D, M-128
 Mauritian E, M-125
 Mauritian F, M-125
 Mauritian H, M-129

- Mauritine J, A-939
 C-Mavacurine, M-130
 Mavacurine, M-130
 Maxonine, M-131
 Mayfoline, P-356
 Maymyrsine, P-217
 Mayotamide A, M-132
 Mayotamide B, M-133
 Maysenine, M-134
 Maysine, M-134
 Maytanacine, M-136
 Maytanbicyclinol, T-449
 Maytanbutacine, M-136
 Maytanbutine, M-135
 Maytanprine, M-135
 Maytansine, M-135
 Maytansinol, M-136
 Maytanvaline, M-135
 Mayteine, P-247
 Maytenine, S-398
 Maytine, H-247
 MAYT, M-135
 Maytol, H-118
 Maytolidine, H-118
 Maytoline, H-118
 Mayumbine, A-212
 Mazethramycin A, M-137
 Mazethramycin B, M-137
 Mazethramycin C, M-137
 MBH 001, A-1203
 MBI 3253, C-188
 MBOA, B-82
 MBP 039-06, A-1204
 MBP 049-13, A-1205
 McB17, M-593
 MDL 62879, A-1180
 MDL 28574A, C-188
 MDL 25637, B-186
 MDL 28574, C-188
 Mearsamine, M-138
 Mearsine, M-139
 Mecambridine methosalt, M-140
 Mecambridine, M-140
 Mecambrine, M-141
 Mecambroline, L-72
 Mechercharmynin A, M-142
 Mechercharmynin B, M-143
 Mechercharstatin A, M-142
 Mechercharstatin B, M-143
 Meconium, M-704
 Meconoquintupline, M-144
 Mecoquitupline, M-144
 Mecostrin chloride, C-420
 Mecostrin, C-420
 Medelamine A, M-579
 Medelamine B, M-566
 Medelline, M-145
 Medicosmine, A-121
 Megacarpidine, S-445
 Megacarpine, S-445
 Megalanthonine, M-146
 Megastachine, M-147
 Megistonine II, T-272
 Megistonine I, T-584
 Megistophylline II, M-149
 Megistophylline I, M-148
 Megistoquinone II, M-151
 Megistoquinone I, M-150
 Megistosarcimine, M-152
 Megistosarconine, M-152
 Mehranine, M-153
 Melanacanthine, E-198
 Melandrin, A-789
 Melanocin A, M-154
 Melanocin B, M-155
 Melanocin C, M-156
 Melanocrocine, M-157
 Melanostatin, M-158
 Melanoxazol, O-156
 Melanoxidin, M-159
 Melanthioidine, M-160
 Melatonin, M-288
 Melaxillaridine, M-161
 Meleagrins, M-162
 Melemeleone A, M-163
 Melemeleone B, M-164
 Melevodopa, A-745
 Melicarpine, M-165
 Melicarpinone, M-166
 Melicobisquinolinone A, M-167
 Melicobisquinolinone B, M-168
 Melicopicine, T-240
 Melicopidine, X-2
 Melicopine, M-169
 Melinacidin II, C-320
 Melinacidin III, C-320
 Melinacidin IV, C-320
 Melineurine, F-212
 Melinonine A, A-212
 Melinonine B, D-412
 Melinonine E, S-606
 Melinonine F, M-426
 Melinonine G, F-84
 Melinonine H, M-171
 Melinonine I, M-170
 Melinonine K, M-172
 Melinonine L, M-173
 Melinonine M, M-174
 Melisemine, M-175
 Melithiazole A, M-177
 Melithiazole B, M-177
 Melithiazole C, M-178
 Melithiazole D, M-177
 Melithiazole E, M-177
 Melithiazole F, M-179
 Melithiazole G, M-180
 Melithiazole H, M-181
 Melithiazole I, M-176
 Melithiazole K, M-177
 Melithiazole L, M-177
 Melithiazole M, M-181
 Melithiazole N, M-177
 Melleumin A, M-182
 Melleumin B, M-182
 Mellicetine, G-91
 Melobaline, V-124
 Meloceline, A-1498
 Meloceline, A-1498
 Melochicorine, M-183
 Melochinine, M-184
 Melochininone, M-184
 Melochinone, M-184
 Melochinone, M-185
 Melofoline, M-186
 Melomorsine, M-187
 Melonine, M-188
 Melonovine A, M-189
 Melonovine B, M-190
 Melopflins, M-191
 Melosatin A, P-349
 Melosatin B, P-349
 Melosatin C, P-349
 Melosatin D, P-349
 Meloscandonine, M-192
 Meloscine, M-193
 Melosmidine, M-194
 Melosmine, M-194
 Melovinone, M-195
 Memnobotrin A, M-196
 Memnobotrin B, M-196
 Memoril, G-108
 Menisarine, M-197
 Menisaurin, L-213
 Menisidine, T-286
 Menisine, T-286
 Menismine, M-198
 Menisperine, I-208
 Menisporphine, M-199
 Menolsin, Y-13
 Menthiafoliac acid, H-471
 Meprochol, M-284
 Mequinine, M-200
 Mequiverine, Q-17
 Mer-A 2026A, P-421
 Mer-A 2026B, P-421
 Merabitol, A-216
 2-Mercaptobenzothiazole, B-79
 2-Mercaptohistidine trimethylbetaine, E-173
 2-Mercapto-1*H*-indole-3-carboxaldehyde, M-201
 2-Mercapto-1*H*-indole-3-carboxylic acid, M-202
 2-(Mercaptomethyl)quinoline, Q-36
 5-Mercaptopentanoic acid, M-203
 5-Mercapto-4-pentenoic acid, M-204
 4-Mercapto-2-pyrimidinol, H-556
 3-(2-Mercaptothiazol-5-yl)-1*H*-indole, I-139
 5-Mercaptovaleric acid, M-203
 Merckonine, M-373
 Merenderine *N*-oxide, P-221
 Merenderine, P-221
 Merenskine *N*-oxide, S-122
 Merenskine, S-122
 Meridianin A, M-205
 Meridianin B, M-205
 Meridianin C, M-205
 Meridianin D, M-205
 Meridianin E, M-205
 Meridianin F, M-205
 Meridianin G, M-205
 Meridine, M-206
 12(13*H*)-Meridinone, M-206
 Merobustine, P-221
 Merobustinine, P-221
 Merredissine, M-391
 Merresectine A, A-1597
 Merresectine B, M-207
 Merresectine C, M-207
 3β-Merresectine C, M-207
 Merresectine D glucoside, M-208
 Merresectine D, M-208
 3β-Merresectine D, M-208
 Merresectine E glucoside, M-208
 Merresectine E, M-208
 Merresectine F, M-209
 3β-Merresectine F, M-209
 Mersicarpine, M-210
 Mersidasine A, M-217
 Mersidasine B, M-217
 Mersidasine C, M-211
 Mersidasine D, M-211
 Mersidasine E, M-217
 Mersidasine F, M-217
 Mersidasine G, M-217
 Mersifoline A, M-212
 Mersifoline B, M-212
 Mersifoline C, M-212
 Mersilongine, M-213
 Mersilosine A, M-214
 Mersilosine B, M-214
 Mersilosine, M-214
 Mersinaline, M-215
 Mersingine A, M-216
 Mersingine B, M-216
 Mersinine A, M-217
 Mersinine B, M-217
 Mersinine C, M-217
 Mersirachine, M-218
 Merucathine, A-867
 Merucathinone, A-868
 Mer-WF 1726, C-945
 Mesaconitine, M-331
 Mescaline citrimide, M-220
 Mescaline isocitrimide lactone, M-221
 Mescaline maleimide, M-222
 Mescaline malimide, M-222
 Mescaline succinimide, M-222
 Mescaline, M-219
 Mescalarovic acid, M-219
 Mescalotam, P-297
 Mescaloxylic acid, M-219
 Mescengricin, M-223
 Mesembrane, M-226
 Mesembranol, M-226
 Mesembranone, M-226
 Mesembrinine, M-224
 Mesembrenol, M-224
 Mesembrenone, M-224
 Δ⁷-Mesembrenone, M-225
 Mesembrine, M-226
 Mesembrinine, M-224

- Mesembrinol, M-226
 Mesembryanthemins, M-227
 Mesoyohimbine, Y-13
 Metabolite AM 2498, B-216
 Metachromin G, M-228
 Metachromin H, M-229
 Metacycloprodigiosin, M-230
 Metacytofilin, M-231
 Metadelphene, M-398
 Metadrenaline, N-303
 Metafolin[®], T-172
 Metahempic acid, D-533
 Metanephrine, N-303
 Metanicotine, M-232
 Metanite, T-632
 Metaphanine, M-233
 Metaquat, D-743
 Meteloidine, M-392
 Metenaticin B, M-234
 Metenaticin C, M-234
 Metergotamine, E-172
 Methacryloylcholine, C-419
 Methanamine, M-383
 Methanecarboxamide, A-30
 Methanophenazine, P-328
 Methanopterin, M-235
 Metheph, D-738
 Methothalstyline, T-337
 Methoxatin, M-236
 6-Methoxy-2,3,7,8-bis(methylenedioxy)benzo[*c*]-phenanthridine, P-55
 1-(Methoxyacetyl)- β -carboline, A-1402
 11-Methoxy-*O*-acetylserotiline, R-63
N-(Methoxyacetyl)-3-methylbutylamine, M-259
 3-Methoxy-*N*-acetylnornantene, P-369
 3'-Methoxyacoforestinine, M-336
 9-Methoxyacridine, A-112
 6-Methoxyacronycidine, F-214
 5-Methoxyacronycine, A-1527
 3-Methoxyaegeline, A-153
 12-Methoxyaffinisine, M-237
 10-Methoxyaffinisine, S-89
 1-Methoxyagroclavine I, A-205
 10-Methoxyajmalicine, A-1423
 11-Methoxyakuammicine *N*⁹-oxide, A-228
 11-Methoxyakuammicine, A-228
 12-Methoxyakuammicine, A-228
 17-Methoxy- ψ -akuammigine, A-232
 1-Methoxyalloycryptopine, F-2
 12-Methoxyalloycryptopine, F-2
 (2-Methoxyallyl)trimethylammonium, M-284
 11-Methoxyalstonal, A-648
 10-Methoxyamerovalficine, P-10
 4-Methoxyamphetamine, A-898
N-Methoxyanhydrobasinediol, T-10
 Methoxyannomontine, A-1051
 15-Methoxyansamitocin P3, M-136
 4-Methoxyanthranilic acid, A-788
 5-Methoxyanthranilic acid, A-789
 3-Methoxyanthranilic acid, A-787
 10-Methoxyantirrhine, A-1328
 Methoxyantirrhine, A-1328
 10-Methoxyapparicine, A-1365
 5-Methoxyarborinine, T-241
N-Methoxyaristolactam II, D-567
 9-Methoxyaristolactam II, T-538
 9-Methoxyaristolactam I, T-251
 9-Methoxyaristolactam IV, P-214
 7-Methoxyaristolochic acid A, T-267
 9-Methoxyaristolochic acid II, T-570
 12-Methoxyaspidofractinine, A-1494
 17-Methoxyaspidofractinine, A-1494
 10-Methoxyaspidospermidine, A-1498
 17-Methoxyaspidospermidine, A-1499
 2-Methoxyatherosperminine *N*-oxide, M-238
 2-Methoxyatherosperminine, M-238
 4-Methoxybenzamide, M-240
 β -Methoxybenzeneethanamine, A-864
 4-Methoxybenzeneethanamine, M-280
 10-Methoxybenzimidazo[6,7,1-*de*]1,6)naphthyridine, M-239
 8-Methoxybenzo[*f*]-1,3-benzodioxolo[6,5,4-*cd*]indol-5(6*H*)-one, A-1427
 4-Methoxy-8*H*-benzo[*g*]benzodioxolo[6,5,4-*de*]quinolin-8-one, A-1531
 2-Methoxy-12*H*-benzo[*h*]1,3]benzodioxolo[6,5,4-*de*]quinolin-12-one, B-117
 10-Methoxy-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-8-one, L-40
 9-Methoxy-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-8-one, O-209
 7-Methoxy[1,3]benzodioxolo[4,5-*c*]-1,3-dioxolo[4,5-*c*]phenanthridine, P-81
 14-Methoxy[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridine, P-55
 1-Methoxy[1,3]benzodioxolo[5,6-*c*]phenanthridin-2-ol, C-356
 2-Methoxy[1,3]benzodioxolo[5,6-*c*]phenanthridin-1-ol, C-356
 7-(7-Methoxy-1,3-benzodioxol-5-yl)-*N*-(2-methylpropyl)-2,4-heptadienamide, C-387
 3-[5-(7-Methoxy-1,3-benzodioxol-5-yl)-2-oxazolyl]pyridine, T-299
 1-[5-(6-Methoxy-1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]piperidine, W-17
 1-[5-(6-Methoxy-1,3-benzodioxol-5-yl)-1-oxo-2,4-pentadienyl]pyrrolidine, W-16
 4-Methoxybenzoic acid, M-240
 4-Methoxybenzo[*c*]2,7]naphthyridine, B-75
 8-Methoxy-9*H*-benzo[*de*]1,6]naphthyridin-9-one, M-241
 6-Methoxy-2(3*H*)-benzothiazolone, B-78
 2-Methoxy-2*H*-1,4-benzoxazin-3(4*H*)-one, H-409
 6-Methoxy-2-benzoxazolinone, B-82
 6-Methoxy-2-benzoxazolol, B-82
 6-Methoxy-2(3*H*)-benzoxazolone, B-82
 1-(4-Methoxybenzoyl)-6,7-methylenedioxyisoquinoline, M-242
N-(4-Methoxybenzoyl)-*N'*-methylguanidine, M-466
 14-(4-Methoxybenzoyl)neoline, M-323
 3-Methoxybenzylamine, H-416
*N*⁵-(4-Methoxybenzyl)glutamine, G-108
N-(3-Methoxybenzyl)hexadecanamide, H-221
 1-(4-Methoxybenzyl)-6,7-methylenedioxyisoquinoline, H-424
 1-Methoxyberberine, M-243
 4-Methoxy-[2,2'-bipyridine]-6-carboxaldehyde oxime, C-9
 4-Methoxy-[2,2'-bipyridine]-6-carboxaldehyde, C-11
 4-Methoxy-2,2'-bipyrole-5-carboxaldehyde, M-244
 4-Methoxy-8*H*-bis[1,3]benzodioxolo[6,5,4-*de*:5',6'-*g*]quinolin-8-one, C-179
 10-Methoxy-1,2,8,9-bis(methylenedioxy)aporphine, P-210
 3-Methoxy-1,2,9,10-bis(methylenedioxy)noraporphine, N-125
 3-Methoxy-2,5-bis(1-methylethyl)pyrazine, D-699
 3-Methoxy-2,5-bis(2-methylpropyl)pyrazine, F-76
 7-Methoxy- α,α -bis(methylthio)-1*H*-indole-3-acetonitrile, D-873
 5-Methoxy-3,6-bis(phenylmethyl)-2(1*H*)-pyrazinone 4-oxide, D-300
 5-Methoxy-3,6-bis(phenylmethyl)-2(1*H*)-pyrazinone, D-300
 Methoxybrassenin A, M-245
 Methoxybrassenin B, M-246
 4-Methoxybrassinin, B-284
 Methoxybrassinin, B-284
 Methoxybrassinin, B-285
 6 α -Methoxybuphanidine, P-581
 4-Methoxy-1-butylamine, A-716
 2-Methoxy-3-*sec*-butylpyrazine, M-272
 10-Methoxycaaverine, T-524
 6-Methoxycamalexin, T-372
 7-Methoxycamptothecin, C-70
 9-Methoxycamptothecin, C-70
 10-Methoxycamptothecin, C-70
 5-Methoxycanthin-2,6-dione, I-105
 3-Methoxycanthin-2,6-dione, I-105
 8-Methoxycanthin-6-one (incorr.), H-444
 9-Methoxycanthin-6-one *N*³-oxide, H-444
 1-Methoxycanthin-6-one *N*(3)-oxide, H-438
 4-Methoxycanthin-6-one, H-440
 5-Methoxycanthin-4-one, H-441
 5-Methoxycanthin-6-one, H-442
 9-Methoxycanthin-6-one, H-444
 1-Methoxycanthin-6-one, H-438
 10-Methoxycanthin-6-one, H-445
 11-Methoxycanthin-6-one, H-446
 2-Methoxycanthin-6-one, H-439
 8-Methoxy-2-carbazolecarboxaldehyde (obsol.), H-449
 7-Methoxy-9*H*-carbazole-3-carboxaldehyde, H-450
 8-Methoxy-9*H*-carbazole-3-carboxaldehyde, H-451
 1-Methoxy-9*H*-carbazole-3-carboxaldehyde, H-447
 6-Methoxy-9*H*-carbazole-3-carboxaldehyde, H-449
 2-Methoxy-9*H*-carbazole-3-carboxaldehyde, H-448
 7-Methoxy-9*H*-carbazole-3-carboxylic acid, H-455
 1-Methoxy-9*H*-carbazole-3-carboxylic acid, H-452
 8-Methoxy-9*H*-carbazole-3-methanol, H-604
 1-Methoxy-9*H*-carbazole-3-methanol, H-603
 6-Methoxycarbazomycinal, H-561
 7-Methoxy- β -carboline-1-carboxamide, H-459
 1-Methoxy- β -carboline-3-carboxylic acid, C-133
 7-Methoxy- β -carboline, P-916
 8-Methoxy- β -carboline, P-917
 1-Methoxy- β -carboline, P-914
 6-Methoxy- β -carboline-1-propanoic acid, C-134
 7-Methoxy- β -carboline-1-propanoic acid, C-134
 8-Methoxy- β -carboline-1,3,4-trione, C-135
 2-Methoxy-3-carbomethoxycarbazole, H-453
N-Methoxycarbonylactinodaphnine, D-346
N-Methoxycarbonylanonane, R-120
 5-(3-Methoxycarbonylbutyroyl)aminomethyl-*trans*-quinolizidine *N*-oxide, O-59
 5-[(3-Methoxycarbonylbutyroyl)aminomethyl]quinolizidine, O-59
 1-Methoxycarbonyl- β -carboline, C-132
 3-Methoxycarbonyl- β -carboline, C-133
N-Methoxycarbonylcassifiline, O-27
 12-Methoxycarbonylcytisine, C-938
 37-(Methoxycarbonyl)dictyodendrine E, M-247
*N*¹-Methoxycarbonyl-16,17-didehydrokopsinine, K-71
N-Methoxycarbonyl-11,12-dimethoxykopsinaline, K-74
N-Methoxycarbonyl-10,22-dioxokopsane, D-805
N-Methoxycarbonyl-11-hydroxy-12-methoxykopsinaline, K-74
N-Methoxycarbonyl-12-hydroxy-11-methoxykopsinaline, K-74
N-Methoxycarbonylkopsinaline, K-74
 3-(Methoxycarbonyl)meloscine, S-118
N-Methoxycarbonyl-12-methoxykopsinaline, K-74
 (Methoxycarbonylmethoxy)phenazine, P-327
N-Methoxycarbonyl-11,12-methylenedioxykopsinaline, K-74
N-Methoxycarbonyl-11,12-methylenedioxy- Δ ^{16,17}-kopsinine *N*-oxide, K-71
N-Methoxycarbonyl-11,12-methylenedioxy- Δ ^{16,17}-kopsinine, K-71
 4-Methoxycarbonyl-1-methyl-2,7-naphthyridine, M-488
 5-Methoxycarbonyl-4-methyl-2,7-naphthyridine, M-489
N-Methoxycarbonylnorisocorydine, I-208
N-Methoxycarbonyl-*N*-normuciferine, D-531
 1-Methoxycarbonylpyrrolizidine, P-960
 5-(Methoxycarbonyl)tubercidin, S-53
 3-Methoxycarbonyl-5-vinylpyridine, V-141
 10-Methoxycathafoline *N*⁴-oxide, C-191
 10-Methoxycathafoline, C-191
 3-Methoxycepharadione B, O-139
 10-Methoxychelidonine, C-353
 10-Methoxycinchonamine, C-444
 6'-Methoxycinchonan-9-ol, Q-18
 6'-Methoxycinchonan-9-ol, Q-20
 6'-Methoxycinchonan-9-one, Q-18
m-Methoxycinnamic acid pyrrolidide, H-690
p-Methoxycinnamic acid pyrrolidide, H-691
 2-(3-Methoxycinnamoyl)-4*H*-3,1-benzoxazin-4-one, A-1561
*N*¹-(*Z*)-*p*-Methoxycinnamoylbuchnerine, P-673
N-(4-Methoxycinnamoyl)dopamine, D-920
N-(4-Methoxycinnamoyl)ethylenediamine, E-225
 1-(*m*-Methoxycinnamoyl)pyrrolidine, H-690

- N*-(4-Methoxycinnamoyl)pyrrolidine, H-691
N-(4-Methoxycinnamoyl)-2-pyrrolidinone, P-956
 6-Methoxycleistopholine, M-399
 7-Methoxycleistopholine, M-399
 2-Methoxycomosivine, C-588
 11-Methoxycompactinervine, C-589
 12-Methoxycompactinervine, C-589
 3*R*-Methoxycoronarinine, C-657
 9-Methoxycorynanthine, G-13
 10-Methoxycorynantheol α -*N*-methosalt, D-412
 10-Methoxycorynantheol β -*N*-methosalt, D-412
 10-Methoxycorynantheol, D-412
 10-Methoxycorynanthine, M-291
 6-Methoxycrinamine, H-6
 7-Methoxycrinan-11-ol, H-9
 23-Methoxycyclopamine, J-36
 4'-Methoxycyclopeptine, C-890
 4'-Methoxycytochalasin E, C-945
 Methoxydambullin, D-12
 10-Methoxydeacetylakuammiline, A-231
 Methoxydechlorochartelline A, C-350
 9-Methoxydehydroanonaine, D-175
 11-Methoxy-2,16-dehydrodiaboline, C-115
*N*¹-Methoxy-19,20-dehydroervatamine, E-190
 12-Methoxy-14,15-dehydrovincamine, V-115
 18-Methoxydelterine, M-339
 10-Methoxy-16-demethoxycarbonylpagicerine, P-10
 5-Methoxy-9-*O*-demethylhomolycorine, H-504
 10-Methoxy-*O*-demethylsilanine, T-649
 23-Methoxy-11-deoxojervine, J-36
 10-Methoxydeplancheine, D-230
 11-Methoxydiaboline *N*-oxide, C-115
 11-Methoxydiaboline, C-115
 4-Methoxydianthramide B methyl ester, B-67
 4-Methoxydianthramide B, B-67
 4-Methoxydianthramide M, B-67
 4-Methoxydianthramide R, B-67
 4-Methoxydianthramide S, B-67
 11-Methoxydichotine, D-359
 5-Methoxydictamine, F-210
 6-Methoxydictamine, F-211
 7-Methoxydictamine, F-212
 8-Methoxydictamine, F-213
 6-Methoxydictyolomide A, D-367
 5-Methoxydihydrochelerythrine, D-407
 8-Methoxydihydrochelerythrine, D-407
 12-Methoxydihydrochelerythrine, D-407
 5-Methoxydihydrochelirubine, D-510
 10-Methoxydihydrocorynantheol 4*z*-*N*-oxide, D-412
 10-Methoxydihydrocorynantheol 4*β*-*N*-oxide, D-412
 10-Methoxydihydrocorynantheol, D-412
 11-Methoxydihydrocorynantheol, D-412
 4-Methoxydihydroorsecurinine, N-307
 15-Methoxy-14,15-dihydrophylllochrycine, A-630
 2-Methoxy-1,2-dihydrorhazimine, R-73
 8-Methoxydihydrosanguinarine, D-510
 10-Methoxy-4',17-dihydro-17*α*-tchibangensine, U-52
 10-Methoxy-4',17-dihydro-17*β*-tchibangensine, U-52
 3-Methoxy-4,6-dihydroxymorphinandien-7-one, S-39
 3-Methoxy-20-(dimethylamino)pregna-5,16-dien-2-ol, A-874
 3-Methoxy-20-dimethylaminopregn-5-ene, A-886
 4-Methoxy-*N,N*-dimethylbenzenethanamine, M-280
 3-Methoxy-*N,N*-dimethylbenzenemethanamine, H-416
 1'-Methoxy-3,3'-dimethyl-[2,4'-bi-9*H*-carbazole]-1,4-dione, B-135
 3-Methoxy-1,2-dimethyl-9*H*-carbazol-4-ol, D-571
 1-Methoxy-*N,N*-dimethyl-10*H*-[1,3]dioxolo[6,7]-phenanthro[4,5-*bcd*]pyran-3-ethanamine, T-327
 2-Methoxy-*N,N*-dimethylethanamine, D-730
 2-Methoxy-*N,N*-dimethylethylamine, D-730
 11-Methoxy-3,3-dimethyl-3*H*-furanol[2,3-*b*]pyranol[3,2-*f*]quinoline, A-121
 1-Methoxy-*N,N*-dimethyl-1*H*-indole-3-ethanamine, D-785
 7-Methoxy-*N,N*-dimethyl-1*H*-indole-3-methanamine, D-735
 α -Methoxy-*N,N*-dimethyl-1*H*-indole-3-propanamide, H-539
 5-Methoxy-2,6-dimethyl-2*H*-isoindole-4,7-dione, M-248
 7-Methoxy-1,6-dimethyl-5,8-isoquinolinedione, M-249
 7-Methoxy-2,6-dimethyl-3,5,8(2*H*)-isoquinolinetriene, M-624
 5-Methoxy-*N,N*-dimethyl- β -oxotryptamine, D-727
 10-Methoxy-*N,N*-dimethylphenanthro[3,4-*d*]-1,3-dioxole-5-ethanamine, I-343
 9-Methoxy-*N,N*-dimethylphenanthro[3,4-*d*]-1,3-dioxole-5-ethanamine, U-64
 4-Methoxy-7,8-dimethyl-2*H*-pyranol[4,3-*b*]pyridin-2-one, M-250
 5-Methoxy-2,2-dimethyl-2*H*-pyranol[2,3-*b*]quinoline, M-251
 2-Methoxy-3,5-dimethylpyrazine, H-474
 3-Methoxy-2,5-dimethylpyrazine, H-475
 3-Methoxy- α,α -dimethylpyrazinemethanol, H-545
N-Methoxy-*N,\xi*-dimethyl-3-pyridinetetradecanamine, X-17
 8-Methoxy-5,11-dimethylpyridol[4,3-*b*]carbazole, E-67
 9-Methoxy-5,11-dimethylpyridol[4,3-*b*]carbazole, M-252
 1-Methoxy-*N,N*-dimethyltryptamine, D-785
 5-Methoxy-*N,N*-dimethyltryptamine, M-288
 11-Methoxy-1,3-dioxolo[4,5-*b*]acridin-10-one, T-516
 6-Methoxy-13*H*-[1,3]dioxolo[7,8]benzoxepino[2,3,4-*ij*]isoquinolin-13-one, O-169
 6-Methoxy-1,3-dioxolo[4,5-*b*]furo[2,3-*b*]quinoline, F-223
 9-Methoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline, F-221
 9-Methoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-4-ol, F-217
 4-Methoxy-2,2'-dipyridyl-6-aldoxime, C-9
 1-Methoxydiscorhabdin D, D-852
 9-Methoxydispacamide B, D-863
 Methoxydoisuthine, D-900
 2-Methoxydyshomerythrine, D-963
 11-Methoxydeburnamonine, E-6
 11-Methoxyechitovenidine, M-631
 11-Methoxyechitoveniline, M-631
 16-Methoxyechitoveniline, M-631
 10-Methoxyeglandine *N*-oxide, V-170
 10-Methoxyeglandine, V-170
 8-Methoxyellipticine, E-67
 9-Methoxyellipticine, M-252
 10-Methoxy-17-epialloyohimbine, M-291
 6'-Methoxyepidihydroshihunine, D-511
 11-[10-(11-Methoxy-17-epivincamajiny)]vincorine, M-289
 11-Methoxy-17-epivincamine, V-113
 10-Methoxy-3-epi- α -yohimbine, M-291
 11-Methoxy-17-epi- α -yohimbine, M-292
 11-Methoxy-3-epi- α -yohimbine, M-292
 9-Methoxy-3-epi- α -yohimbine, V-50
 11-Methoxy-19,20*α*-epoxyakuammicine, A-228
 12-Methoxy-19*α*,20*α*-epoxyakuammicine, A-228
 11-Methoxyerysodine, E-206
 11-Methoxyerysopine, E-206
 11*β*-Methoxyerysotramidine, E-202
 11-Methoxyerysotrine, E-206
 11-Methoxyerysotine, E-206
 11*β*-Methoxyerythraline *N*-oxide, E-205
 11-Methoxyerythraline, E-205
 11-Methoxyerythratidine, E-207
 11-Methoxyerythratine, E-208
 4-(2-Methoxyethyl)imidazole, I-38
 3-(1-Methoxyethyl)-5-methoxycarbonylpyridine, H-498
 6-(1-Methoxyethyl)-1-phenazinecarboxylic acid, H-496
 3-Methoxyeupolauridine, I-65
 2-Methoxyeustifoline A, E-300
 5'-Methoxyfagaramide, M-261
 3-Methoxyfaurithaline, F-27
 5-Methoxyflindersiamine, F-214
 5-Methoxyflindersine, F-90
 6-Methoxyflindersine, F-90
 7-Methoxyflindersine, F-90
 8-Methoxyflindersine, F-90
N-Methoxy-3-formylcarbazole, C-121
 1-Methoxy-3-formylindole, I-86
 4-Methoxyfuro[2,3-*b*]quinoline, F-224
 4-Methoxyfuro[2,3-*b*]quinolin-7-ol, F-212
 4-Methoxyfuro[2,3-*b*]quinolin-8-ol, F-213
 18-Methoxygadesine, G-152
 9-Methoxygeissoschizol, G-33
 10-Methoxygeissoschizol, G-33
 11-Methoxygelsemamine, G-48
N-Methoxygelsemine, G-49
 11-Methoxygelsemoxonine, G-50
 Methoxygerambullol, G-68
 4-Methoxyglucobrassicin, I-122
N-Methoxyglucobrassicin, I-122
 11*β*-Methoxyglucoerysodine, E-206
 11*β*-Methoxyglucoerysotine, E-206
 2-Methoxyglycomaurin, E-300
 5-Methoxygramine, D-734
 7-Methoxygramine, D-735
 3'-Methoxygraveoline, G-171
 3-Methoxyguattescidine, G-214
 5-Methoxyhalfordine, F-214
 6-Methoxyharmalin, H-591
 6-Methoxyharmal, H-591
 1'-Methoxyhemibastadin 1, H-100
 1'-Methoxyhemibastadin 2, H-100
 11-Methoxyhenningsamine, C-115
 6-Methoxyheptaphylline, H-125
 7-Methoxyheptaphylline, H-125
 11-Methoxyhumantenine, H-367
 9-Methoxy-10-hydroxy-2,3-methylenedioxybenzophenanthridine, C-356
 1-Methoxy-8-hydroxymethyl-3*H*-phenoxazin-3-one, M-253
 4-Methoxy- β -hydroxyphenethylamine, O-63
 7-Methoxy-4-hydroxy-2-quinolinecarboxylic acid, D-676
 10-Methoxy-21-hydroxyvinorine, V-194
 11-Methoxyyibogamine, I-4
 13-Methoxyyibogamine, I-4
 11-Methoxyyciajine, I-12
 Methoxyincarrillateine, I-60
 4-Methoxyindole 3-isothiocyanate, I-337
 5-Methoxy-1*H*-indole-3-acetic acid, H-525
 6-Methoxy-1*H*-indole-3-acetic acid, H-526
 4-Methoxy-1*H*-indole-3-acetonitrile, H-524
 6-Methoxy-1*H*-indole-3-acetonitrile, H-526
 1-Methoxy-1*H*-indole-3-acetonitrile, I-81
 4-Methoxy-1*H*-indole-3-carboxaldehyde, H-527
 5-Methoxy-1*H*-indole-3-carboxaldehyde, H-528
 1-Methoxy-1*H*-indole-3-carboxaldehyde, I-86
 4-Methoxy-1*H*-indole-2,3-dione, H-532
 6-Methoxy-1*H*-indole-2,3-dione, H-533
 5-Methoxy-1*H*-indole-4,7-dione, M-254
 6-Methoxy-1*H*-indole-4,7-dione, M-255
 5-Methoxy-1*H*-indole-3-ethanamine, M-288
 α -Methoxy-1*H*-indole-3-propanoic acid, H-539
 5-Methoxyindoline, D-465
 5-Methoxy-3*H*-indolo[3,2,1-*de*][1,5]naphthyridine-2,6-dione, I-105
 4-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-440
 9-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-444
 1-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-438
 10-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-445
 11-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-446
 2-Methoxy-6*H*-indolo[3,2,1-*de*][1,5]naphthyridin-6-one, H-439
N-[2-(5-Methoxy-1*H*-indol-3-yl)ethyl]acetamide, M-288
 4-Methoxyindol-3-ylmethyl glucosinolate, I-122
 2-Methoxy-3-(3-indolyl)propanoic acid, H-539

- Methoxyipomine, I-165
 4-Methoxyisatin, H-532
 6-Methoxyisatin, H-533
 8-Methoxyisodectamine, C-356
 6-Methoxyisodictamine, F-211
 10-Methoxyisogeissoschizol, G-33
 16-Methoxyisomatopensine, M-120
 3-Methoxyisoprenaline, N-303
 7-Methoxy-6-isoquinolinol, I-302
 11-Methoxyisoretuline, R-63
 9-Methoxy-(16*R*)-*E*-isositsirikine, I-319
 9-Methoxy-(16*S*)-*E*-isositsirikine, I-319
 8-Methoxy-8*H*-isothiazolo[5,4-*b*]indole, I-336
 5-Methoxyisoxazole, I-349
 8-Methoxykarakoline (incorr.), M-355
 11-Methoxykopsilongine *N*⁴-oxide, K-74
 11-Methoxykopsilongine, K-74
 12-Methoxykopsinaline, K-74
 11-Methoxykopsingine, K-66
 15-Methoxykopsinine, K-78
 10-Methoxykopsinone, K-80
 12-Methoxykopsinone, K-80
 6-Methoxykynurenic acid, D-675
 10-Methoxylanciferine, L-29
 2-Methoxylenticellarine, L-92
 11-Methoxylimapodine, L-177
 11-Methoxylimaspermine, L-177
 11-Methoxylimatine, A-1497
 9-Methoxyliriodenine, L-40
 10-Methoxyliriodenine, O-187
 6-Methoxylunidonine, L-295
 6 α -Methoxylupanine, H-552
 13 β -Methoxylupanine, H-554
 13-Methoxylupanine, H-554
 10-Methoxymacrocarpamine 4'-*N*-oxide, M-11
 10-Methoxymacrocarpamine, M-11
 5-Methoxymaculine, F-215
 11-Methoxymacusine A, P-562
 9-Methoxymappicine gentiobioside, M-96
 9-Methoxymappicine, M-96
 1-Methoxy-9-[(1-methoxy-9*H*-carbazol-3-yl)-methyl]-3-methyl-9*H*-carbazole, B-207
 1-Methoxy-4-[(1-methoxy-9*H*-carbazol-3-yl)-methyl]-3-methyl-9*H*-carbazole, C-423
 1-Methoxy-2-[(9-methoxy-9*H*-carbazol-3-yl)-methyl]-3-methyl-9*H*-carbazole, M-746
 4-Methoxy-1-methoxycarbonyl- β -carboline, H-457
 12-Methoxy-*N*-methoxycarbonylkopsinaline, K-74
 12-Methoxy-*N*-methoxycarbonyl- Δ ^{16,17}-kopsinine *N*⁴-oxide, K-71
 12-Methoxy-*N*-methoxycarbonyl- Δ ^{16,17}-kopsinine, K-71
 9-Methoxy-*N*-methoxycarbonyl-1,2-methylenedioxyaporphine, A-1066
 8-Methoxy-2-(3-methoxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone, G-171
 2-Methoxy-2-(4-methoxyphenyl)ethylamine, O-63
N-[2-Methoxy-2-(4-methoxyphenyl)ethyl]cinamide, A-153
 5-Methoxy-5-(4-methoxyphenyl)-3-methyl-2,4-imidazolidinedione, M-256
 4-Methoxy-4-(4-methoxyphenyl)-1-methyl-5-thioxo-2-imidazolidinone, H-568
 4-Methoxy-5-[(3-methoxy-5-pyrrol-2-yl-2*H*-pyrrol-2-ylidene)methyl]-2,2'-bipyrrrole, M-257
 10-Methoxy-11-[10-(11-methoxyvincamajinyl)]-cathafoline, M-258
*N*¹-Methoxymethuenine, M-293
 1-Methoxy-10-methylacridone, H-402
 12-Methoxy-*N*⁶-methylakuummicine, A-228
 3-Methoxy-2-methylaminobenzoic acid, A-787
 3-Methoxy-20-(methylamino)pregna-5,16-diene, A-876
 6-Methoxy-1-methyl-4-azafuorenone, O-93
 7-Methoxy-1,2-(*N*-methylaziridin)mitosene, M-653
 4-Methoxy- α -methylbenzeneethanamine, A-898
 4-Methoxy-*N*-methylbenzeneethanamine, M-280
 5-Methoxy-13-methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*f*]phenanthridinium(1+), C-360
 6-Methoxy-5-methylbenzo[*h*]pyrrolo[4,3,2-*de*]quinolin-4(5*H*)-one, E-297
 7-Methoxy-4-methylbenzo[*g*]quinoline-5,10-dione, M-399
 3-Methoxy-4-methylbenzo[*g*]quinoline-2,5,10(1*H*)-trione, M-97
 2-Methoxy-4-methyl-2*H*-1,4-benzoxazin-3(4*H*)-one, H-409
 8-Methoxy-16-methyl-2,3:10,11-bis[methylenebis(oxy)]rheadan, R-86
 10-Methoxy-*N*⁶-methylburnamine 17-*O*-benzoate, B-420
 10-Methoxy-*N*⁶-methylburnamine 17-*O*-veratrate, B-420
 4-Methoxy-3-(3-methyl-1,3-butadienyl)-2(1*H*)-quinolinone, G-118
 2-Methoxy-8-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, A-1530
 1-Methoxy-4-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, C-507
 1-Methoxy-2-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, I-75
 9-Methoxy-10-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-4-ol, T-57
 4-Methoxy-8-(3-methyl-2-butenyl)furo[2,3-*b*]quinolin-7-ol, H-570
 7-Methoxy-9-(3-methyl-2-butenyl)furo[2,3-*b*]quinolin-4(9*H*)-one, F-212
 6-Methoxy-3-(3-methyl-2-butenylidene)-2-indolinone, D-471
 7-Methoxy-8-(3-methyl-2-butenyloxy)dictamine, F-223
 4-Methoxy-7-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline, F-212
 4-Methoxy-8-[(3-methyl-2-butenyl)oxy]furo[2,3-*b*]quinoline, F-213
 4-Methoxy-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, A-1529
 2-Methoxy-*N*-(3-methylbutyl)acetamide, M-259
 9-Methoxy-3-methyl-5,6-canthindione, P-413
 4-Methoxy-3-methyl-5,6-canthinedione, P-413
 9-Methoxy-5-methylcanthin-6-one, L-299
 3-Methoxy-6-methyl-9*H*-carbazole-1-carboxaldehyde, H-589
 7-Methoxy-3-methyl-1*H*-carbazole-1,4(9*H*)-dione, K-55
 5-Methoxy-3-methyl-9*H*-carbazole, H-588
 1-Methoxy-3-methyl-9*H*-carbazole, H-583
 3-Methoxy-6-methyl-9*H*-carbazole, H-587
 1-Methoxy-6-methyl-9*H*-carbazole, H-584
 2-Methoxy-3-methyl-9*H*-carbazole, H-585
 7-Methoxy-3-methyl-1,4-carbazolequinone, K-55
 6-Methoxy-1-methyl- β -carboline, H-591
 7-Methoxy-1-methyl- β -carboline, H-592
 7-Methoxy-4-methyl- γ -carboline, M-260
 6-Methoxy-2-methyl- β -carbolinium(1+), P-915
 1-Methoxymethyl-1,2-dehydropyrrolizidine, S-644
 12-Methoxy-*N*¹-methyl-14,15-didehydroaspido-fractinine, A-1494
 11-Methoxy-*N*-methyl-dihydropericyclivine, G-22
 4-Methoxy-5-methyl-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-9(5*H*)-one, F-217
 6-Methoxy-9-methyl-1,3-dioxolo[4,5-*h*]quinolin-8(9*H*)-one, C-177
 1-Methoxy-2,3-methylenedioxyacridone, T-516
 9-Methoxy-1,2-methylenedioxyaporphine, A-1066
 10-Methoxy-1,2-methylenedioxyaporphine, L-72
 7-Methoxy-1,2-methylenedioxyaporphine, O-87
 11-Methoxy-1,2-methylenedioxyaporphine, P-781
 3-Methoxy-1,2-methylenedioxyaporphine, S-536
 8-Methoxy-1,2-methylenedioxyaporphine, T-522
 2-Methoxy-4,5-methylenedioxyacinnamic acid, M-261
 2-Methoxy-4,5-methylenedioxyacinnamoyl isobutylamide, M-261
 2-Methoxy-4,5-methylenedioxyacinnamoyl piperide, M-261
 2-Methoxy-4,5-methylenedioxyacinnamoyl pyrrolidide, M-261
N-(3-Methoxy-4,5-methylenedioxyacinnamoyl)isobutylamine, T-578
N-(3-Methoxy-4,5-methylenedioxyacinnamoyl)piperide, T-578
N-(3-Methoxy-4,5-methylenedioxyacinnamoyl)- Δ ³-pyridin-2-one, D-470
N-(3-Methoxy-4,5-methylenedioxyacinnamoyl)pyrrolidide, T-578
 8-Methoxy-6,7-methylenedioxydictamine, F-217
 4-Methoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline, F-221
 4-Methoxy-7,8-methylenedioxyfuro[2,3-*b*]quinoline, F-223
 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthreneacetic acid, D-102
 6-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthreneacetic acid, H-600
 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthreneacetic acid, M-262
 9-Methoxy-1,2-methylenedioxyaporphine, A-1066
 10-Methoxy-1,2-methylenedioxyaporphine, L-72
 7-Methoxy-1,2-methylenedioxyaporphine, O-87
 11-Methoxy-1,2-methylenedioxyaporphine, P-781
 3-Methoxy-1,2-methylenedioxyaporphine, S-536
 3-Methoxy-1,2-methylenedioxyoxoaporphine, A-1531
 9-Methoxy-1,2-methylenedioxy-7-oxodibenzo[*de,g*]quinoline, L-40
 8-Methoxy-3,4-methylenedioxy-1-phenanthrene-carboxylic acid, M-263
 7-(3-Methoxy-4,5-methylenedioxyphenyl)-2,4-heptadienoic acid isobutylamide, C-387
 4-Methoxy-2-[6-(3,4-methylenedioxyphenyl)hexyl]quinoline, M-453
 2-(3-Methoxy-4,5-methylenedioxyphenyl)-1-methyl-4(1*H*)-quinolinone, G-171
 5-(6-Methoxy-3,4-methylenedioxyphenyl)-2,4-pentadienoic acid isobutylamide, P-462
 1-[5-(2-Methoxy-4,5-methylenedioxyphenyl)pentadienyl]piperidine, W-17
 1-[5-(2-Methoxy-4,5-methylenedioxyphenyl)pentadienyl]pyrrolidide, W-16
 2-(3-Methoxy-4,5-methylenedioxyphenyl)-5-(3-pyridinyl)oxazole, T-299
 4-Methoxy-2-(3,4-methylenedioxyphenyl)quinoline, M-264
 4-Methoxy-6,7-methylenedioxy-8-prenyloxyfuro[2,3-*b*]quinoline, F-217
 4-Methoxy-2-(3,4-methylenedioxyphenyl)quinoline, M-265
 3-(Methoxymethylene)-2-pyrrolidinethione, M-266
 1-Methoxymethyl-1,2-epoxypyrrolizidine, E-133
 3-Methoxy-2-(1-methylethyl)-5-(1-methylpropyl)pyrazine, I-275
 3-Methoxy-2-(1-methylethyl)-5-(2-methylpropyl)pyrazine, I-276
 2-Methoxy-3-(1-methylethyl)pyrido[2,3-*b*]pyrazine, I-278
 6-Methoxy-*N*-methylflindersine, F-90
 7-Methoxy-*N*-methylflindersine, F-90
 6-Methoxy-9-methylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-211
 8-Methoxy-9-methylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-213
 4-Methoxy-9-methyl-7*H*-furo[2,3-*b*]quinolin-7(9*H*)-one, M-166
 9-Methoxy-*N*⁶-methylgeissoschizol, G-33
 10-Methoxy-4-methylgeissoschizol, G-33
 8-Methoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridine-5,9-diol, D-626
 8-Methoxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, O-93
 1-Methoxy- β -methyl-1*H*-indole-3-butanol, I-124
 5-Methoxy-1-methyl-1*H*-indole-3-carboxaldehyde, H-528
 6-Methoxy-1-methyl-1*H*-indole-2,3-dione, H-533
 4-Methoxy-3-methyl-3*H*-indolo[3,2,1-*de*][1,5]-naphthyridine-5,6-dione, P-413
 6-Methoxy-*N*-methylisatin, H-533
 6-Methoxy-1-methyl-7-isoquinolinol, D-627
 3'-Methoxy-4'-*O*-methyljoubertiamine, J-49
 3'-Methoxy-4'-*O*-methyljoubertiaminol, J-50

- 1-Methoxy-3-methyl-2-(3-methyl-2-butenyl)-9*H*-carbazole, C-504
- 7-Methoxy-3-methyl-8-(3-methyl-2-butenyl)-1*H*-carbazole-1,4(9*H*)-dione, M-766
- 7-Methoxy-3-methyl-8-(3-methyl-2-butenyl)-9*H*-carbazol-2-ol, E-275
- 7-Methoxy-3-methyl-8-(3-methyl-2-butenyl)-9*H*-carbazol-1-ol, M-759
- 6-Methoxy-9-methyl-7-(3-methyl-2-butenyl)-1,3-dioxolo[4,5-*h*]quinolin-8(9*H*)-one, P-759
- 4-Methoxy-1-methyl-8-[(3-methyl-2-butenyl)oxy]-2(1*H*)-quinolinone, Q-47
- 1-Methoxy-10-methyl-2,3-methylenedioxyacridone, T-516
- 4-Methoxy-2-methyl-6,7-methylenedioxyisoquinolinium, H-424
- 7-Methoxy-1-methyl-2-(3,4-methylenedioxyphenyl)-4(1*H*)-quinolinone, L-290
- 4-Methoxy-1-methyl-7,8-methylenedioxy-3-prenyl-2-quinolone, P-759
- 4-Methoxy-1-methyl-7,8-methylenedioxy-2(1*H*)-quinolinone, C-177
- 2-Methoxy-5-methyl-3-(1-methylethyl)pyrazine, I-277
- 4-(Methoxymethyl)-2-methylindole, M-478
- 4-Methoxy-1-methyl-8-(3-methyl-2-oxobutoxy)-2(1*H*)-quinolinone, F-122
- 4-Methoxy-1-methyl-3-(3-methyl-2-oxobutyl)-2(1*H*)-quinolinone, E-40
- 3-Methoxy-2-methyl-5-(1-methylpropyl)pyrazine, M-267
- 3-Methoxy-2-methyl-5-(2-methylpropyl)pyrazine, M-268
- 3-Methoxy-4-methyl-5-(2-methyl-4-thiazolyl)-4-pentenoic acid, M-269
- 4-Methoxy-*N*-methyl- α -oxobenzeneethanethioamide, M-282
- 9-Methoxy-7-methyl-8-oxo-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolinium(1+), O-209
- 6-Methoxy-7-methyl-8-oxoguanine, A-753
- 4-Methoxy-5-[(5-methyl-4-pentyl-2*H*-pyrrol-2-ylidene)methyl]-2,2'-bi-1*H*-pyrrole, P-636
- 8-Methoxy-1-methyl-2-pentyl-4(1*H*)-quinolinone, H-660
- 10-Methoxy-*N*-methylpericyclivine, S-89
- 1-Methoxy-5-methylphenazinium(1+), P-327
- 3-Methoxy-2-methyl-1-phenyl-9*H*-carbazole, M-270
- 6-Methoxy-1-methyl-2-phenyl-4(1*H*)-quinolinone, D-659
- 7-Methoxy-1-methyl-2-phenyl-4(1*H*)-quinolinone, D-660
- 7-Methoxy-3-methyl-8-prenyl-1,4-carbazolequinone, M-766
- 7-Methoxy-*N*-methyl-8-prenyloxyflindersine, F-90
- 4-Methoxy-1-methyl-8-prenyloxyquinolone, Q-47
- trans*-4-Methoxy-*N*-methyl-L-proline, H-715
- 1-Methoxy-3-(2-methylpropanal-2-oxo)acridin-9-one-4-carbaldehyde, D-717
- 3-Methoxy-2-(1-methylpropyl)-5-(2-methylpropyl)pyrazine, M-271
- 2-Methoxy-3-(1-methylpropyl)pyrazine, M-272
- 2-Methoxy-3-(2-methylpropyl)pyrazine, M-273
- 3-Methoxy-1-methyl-2-propyl-4(1*H*)-quinolinone, M-274
- 6-Methoxy-7-methyl-7*H*-purine, P-820
- 6-Methoxy-7-methyl-9*H*-purine, P-820
- 7-Methoxy-5-methyl-4*H*-pyrano[3,2-*b*]indol-4(5*H*)-one, K-64
- 2-Methoxy-3-methylpyrazine, H-624
- 3-Methoxy-6-methyl-6*H*-pyrazolo[4,3-*e*][1,2,4]triazine, M-285
- 3-Methoxy-1-methylpyrazolo[4,3-*e*][1,2,4]triazine, M-285
- 3-Methoxy-7-methyl-7*H*-pyrazolo[4,3-*e*][1,2,4]triazine, M-285
- N*-Methoxy-*N*-methyl-3-pyridinopentadecanamine, X-19
- N*-Methoxy-*N*-methyl-3-pyridinetradecanamine, X-18
- 3-Methoxy-2-methyl-4(1*H*)-pyridinone, D-632
- N*-Methoxy-*N*-methyl-18-(3-pyridinyl)-11-octadecen-13-yn-1-amine, X-15
- N*-Methoxy-*N*-methyl-14-(3-pyridinyl)-9-tetradecen-1-amine, X-16
- 7-Methoxy-1-methyl-9*H*-pyrido[3,4-*b*]indole, H-592
- 7-Methoxy-1-methyl-5*H*-pyrido[4,3-*b*]indole, M-260
- α -(Methoxymethyl)-9*H*-pyrido[3,4-*b*]indole-1-methanol, A-1402
- 7-Methoxy-6-methylpyrimido[5,4-*e*]-1,2,4-triazin-5(6*H*)-one, R-66
- 5-(Methoxymethyl)-1*H*-pyrrole-2-carboxaldehyde, H-626
- 4-Methoxy-2-methylquinoline, H-633
- 4-Methoxy-1-methylquinolinium(1+), M-275
- 4-Methoxy-1-methyl-2(1*H*)-quinolinone, Q-33
- 2-Methoxy-1-methyl-4(1*H*)-quinolinone, Q-33
- 2-Methoxy-*N*¹³-methylrutaeacarpine, R-168
- 8-Methoxy-3-methyl-2-(1,2,3,4-tetrahydro-6,8-dimethoxy-1,2,3-trimethyl-7-isoquinolinyl)-1-naphthalenol, A-992
- 4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-carbonitrile, C-576
- 4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-carboxaldehyde oxime, C-574
- 4-Methoxy-5-(methylthio)-[2,2'-bipyridine]-6-methanol, C-575
- 6-Methoxy-2-(methylthio)-1*H*-indole-3-carboxaldehyde, M-201
- 5-Methoxy-2-(methylthio)-1,3-thiazino[6,5-*b*]indole, M-276
- 8-Methoxy-1-methyl-2-tridecyl-4(1*H*)-quinolinone, H-750
- 5-Methoxy-*N*-methyltryptamine, M-288
- 12-Methoxy-*N*⁶-methylvellosimine, M-237
- 12-Methoxy-4-methylvoachalotine ethyl ester, P-562
- 12-Methoxy-4-methylvoachalotine, P-562
- 11-Methoxyminoivincine, M-630
- 11-Methoxyminoivincine, M-631
- 9-Methoxymitralactonine, M-660
- 13-Methoxymultiflorine, M-731
- 7-Methoxymurrayacine, M-757
- 6-Methoxymurrayanine, D-552
- 14-Methoxynaamidine A, N-1
- 14-Methoxynaamidine G, N-1
- 11-Methoxy-3*H*-naphthridino[6,5,4-*def*]quinoxalin-2-ol, M-277
- 11-Methoxy-1*H*-naphthridino[6,5,4-*def*]quinoxalin-2(3*H*)-one, M-277
- 5-Methoxyneomphimedine, N-99
- 11-Methoxyneooxydiaboline, M-278
- 4-Methoxyneosidinocin, N-138
- 4'-Methoxyneobouldine, N-185
- 8-Methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-acetic acid, D-102
- 10-Methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, H-600
- 8-Methoxy-6-nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, M-262
- 2-(4-Methoxy-3-nitrophenyl)ethylamine, H-640
- 5-Methoxynoracronycine, A-120
- N*-Methoxynorcephradione A, C-272
- 8-Methoxynorchelerythrine, C-356
- 3-Methoxynordomesticine, P-369
- 10-Methoxynor-C-fluorocararine, N-301
- 8-Methoxynorsanguinarine, P-55
- 4-Methoxynorsecurinine, N-307
- 3-Methoxynuciferine, T-521
- 4-Methoxy-3-(1-octenyl-*O**N**N*-azoxy)-2-butanol, E-61
- 9-Methoxyolivacine, O-83
- 6-Methoxyonychine, O-93
- 8-Methoxyoxygigidione, O-139
- 5-Methoxyoxindole, I-92
- 8-Methoxyoxoasoanine, A-1508
- 13-Methoxyoxoberberine, M-279
- 8-Methoxy-1-oxo-7(11)-eremophilene-12,8-lactam, H-653
- 4-Methoxyoxohernandaline, O-179
- 3-Methoxy-7-oxohernangerine, O-180
- 25-Methoxy-24-oxokibdelone C, H-655
- 4-Methoxy-2-oxo-1-methyl-1,2-dihydronicotinonitrile, R-96
- 4-Methoxy-2-(8-oxononyl)quinoline, H-642
- 3-Methoxyoxoputerine, O-204
- 11-Methoxy-3-oxotabersonine, O-210
- 4-Methoxy-8-oxotetrahydropalmatine, T-301
- 3-Methoxy-2-(12-oxotridecyl)-4(1*H*)-quinolinone, H-749
- 11-Methoxy-19-oxovincamine, V-115
- 13-Methoxy-8-oxerberberine, M-279
- 3'-Methoxypannellin, P-72
- α -Methoxypapaverine, P-79
- 5-Methoxypeganine, P-165
- 4-Methoxy-2-(1-pentenyl)quinoline, H-660
- 6-Methoxy-3-pentyl-1,8-isoquinolinediol, R-163
- 4-Methoxy-2-pentylquinoline, H-660
- 10-Methoxyperakine, P-240
- 10-Methoxypericycline, S-89
- 8-Methoxyphenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, M-263
- 6-Methoxy-1-phenazinecarboxylic acid methyl ester, H-663
- 9-Methoxy-1-phenazinecarboxylic acid, H-664
- 2-Methoxy-1-phenazinecarboxylic acid, H-661
- 1-Methoxyphenazine, P-327
- 2-Methoxyphenazine, P-328
- 6-Methoxy-1-phenazolinol 5,10-dioxide, P-321
- 6-Methoxy-1-phenazolinol, P-321
- 4-Methoxyphenethylamine, M-280
- N*-(*p*-Methoxyphenethyl)-*N*-methylcinnamide, H-144
- 2-Methoxy-2-phenylacetamide, H-671
- 2-(4-Methoxyphenyl)-1,5-dimethyl-3,4-pyrrolidinediol, C-557
- N*-[2-(4-Methoxyphenyl)ethenyl]benzamide, A-242
- 2-[2-(3-Methoxyphenyl)ethenyl]-4*H*-3,1-benzoxazin-4-one, A-1561
- N*-[2-(4-Methoxyphenyl)ethenyl]-3-phenyl-2-propenamide, M-287
- 2-Methoxy-2-phenylethylamine, A-864
- 2-(4-Methoxyphenyl)ethylamine, M-280
- N*-[2-(4-Methoxyphenyl)ethyl]benzamide, M-280
- N*-[2-(4-Methoxyphenyl)ethyl]-*N*-methyl-3-phenyl-2-propenamide, H-144
- 3-[2-(4-Methoxyphenyl)ethyl]-1-methyl-2,4(1*H*,3*H*)-quinazolinone, M-510
- 6-(4-Methoxyphenyl)-2,4-hexadienoic acid isobutylamide, P-466
- 5-(4-Methoxyphenyl)-2,4-hexadienoic acid isopentylamide, M-281
- 6-(4-Methoxyphenyl)-*N*-(3-methylbutyl)-2-oxo-3,5-hexadienamide, M-281
- 3-[Methoxy(phenyl)methylene]-2-indolinone, I-183
- 3-(4-Methoxyphenylmethyl)-4-methyl-6-(1-methylthyl)-2,5-morpholinedione, I-274
- 2-(4-Methoxyphenyl)-*N*-methyl-2-oxoacetamide, H-687
- (4-Methoxyphenyl)-*N*-methyl-2-oxothioacetamide, M-282
- 6-(4-Methoxyphenyl)-*N*-(2-methylpropyl)-2,4-hexadienamide, P-466
- 11-(4-Methoxyphenyl)-*N*-(2-methylpropyl)-2,4,10-undecatrienamide, P-359
- 2-(4-Methoxyphenylmethyl)-3,4-pyrrolidinediol 3-acetate, A-1044
- 4-Methoxy-2-(phenylmethyl)quinazoline, B-93
- 2-[9-(2-Methoxyphenyl)nonyl]-1-methylpyrrolidine, I-178
- 3-[5-(4-Methoxyphenyl)-2-oxazolyl]pyridine, H-12
- 1-(4-Methoxyphenyl)-2-propylamine, A-898
- 4-Methoxy-2-phenylquinoline, H-693
- 3-(4-Methoxyphenyl)quinoline, H-692
- 3-Methoxy-4-phenyl-2(1*H*)-quinolinone, D-656
- 11-(4-Methoxyphenyl)-2,4,10-undecatrienoic acid isobutylamide, P-359
- 11-Methoxypicraphylline, P-404
- 4-Methoxypericolic acid, H-697
- 4-Methoxy-2-piperidinecarboxylic acid, H-697
- 2'-Methoxyperipine, W-17
- 6-Methoxyperoylisobutylamine, P-462
- 8-Methoxyplatydesmine, B-14
- 12-Methoxypleiocarpine, K-78
- Methoxypolysignine, P-563

- 7-Methoxy-8-prenyldictamine, H-570
 7-Methoxy-9-prenylfuro[2,3-*b*]quinolin-4(9*H*)-one, F-212
 8-Methoxy-7-prenyloxydictamine, F-223
 4-Methoxy-7-prenyloxyfuro[2,3-*b*]quinoline, F-212
 4-Methoxy-8-prenyloxyfuro[2,3-*b*]quinoline, F-213
 6-Methoxy-3-(1-propenyl)-2-pyridinecarboxylic acid, M-283
 4-Methoxy-2-(1-propenyl)quinoline, P-647
N-(2-Methoxy-2-propenyl)trimethylammonium(1+), M-284
 6-(β-Methoxypropionyl)-1,3-dimethylumazine, H-702
 6-(β-Methoxypropionyl)-3-methylumazine, H-702
 4-Methoxy-2-propylquinoline, H-707
 11-Methoxypseudoephedrine, M-292
 6-Methoxypurine, P-820
 3-Methoxyputerine, T-245
 Methoxypyrazine, P-889
 3-Methoxypyrazolo[4,3-*e*][1,2,4]triazine, M-285
 4-Methoxy-3-pyridinecarboxylic acid, H-709
 6-Methoxy-3-pyridinecarboxylic acid, H-711
 3-Methoxypyridine, P-906
N-Methoxy-3-pyridinetridecanamine, T-358
N-Methoxy-16-(3-pyridinyl)-5-hexadecyn-1-amine, N-234
N-Methoxy-16-(3-pyridinyl)-7-hexadecyn-1-amine, N-233
 7-Methoxy-5-(2-pyridinyl)isothiazolo[4,5-*b*]pyridine, C-578
 4-Methoxy-6-(3-pyridinyl)-2*H*-pyran-2-one, A-1037
N-Methoxy-14-(3-pyridinyl)-3-tetradecyn-1-amine, I-26
N-Methoxy-14-(3-pyridinyl)-11-tetradecyn-1-amine, N-227
 8-Methoxy-9*H*-pyrido[3,4-*b*]carbazole-1,3,4(2*H*,9*H*)-trione, C-135
 4-Methoxy-9*H*-pyrido[3,4-*b*]indole-1-carboxaldehyde, H-456
 4-Methoxy-9*H*-pyrido[3,4-*b*]indole-1-ethanol, H-493
 6-Methoxy-9*H*-pyrido[3,4-*b*]indole, P-915
 8-Methoxy-9*H*-pyrido[3,4-*b*]indole, P-917
 7-Methoxy-9*H*-pyrido[3,4-*b*]indole, P-916
 1-Methoxy-9*H*-pyrido[3,4-*b*]indole, P-914
 1-(4-Methoxy-9*H*-pyrido[3,4-*b*]indol-1-yl)-1,2-ethanediol, H-493
 2-Methoxy-1-(9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, A-1402
 4-[(6-Methoxy-9*H*-pyrido[3,4-*b*]indol-1-yl)methyl]phenol, C-230
 4-(6-Methoxy-9*H*-pyrido[3,4-*b*]indol-1-yl)-2-pyrimidamine, A-1051
 4-Methoxypyrido[4,3,2-*mm*]pyrrolo[3,2,1-*de*]acridine, A-1445
 4'-Methoxypyridoxine, P-923
 4-Methoxy-2-pyrrolocarboxylic acid, H-714
 2-Methoxy-1-pyrrolidinecarboxamide, P-955
 4-Methoxy-2-pyrrolidinecarboxylic acid, H-715
 5-Methoxy-2-pyrrolidinone, H-719
 2-Methoxy-4-[2-(2-pyrrolidinyl)ethenyl]phenol, N-305
 2-[[3-Methoxy-4-(1*H*-pyrrol-1-yl)-2*H*-pyrrol-2-ylidene]methyl]-5-methyl-1-pentyl-1*H*-pyrrole, M-286
 11-Methoxyquaternoxine, C-191
 11-Methoxyquebrachamine, Q-6
 4-Methoxyquinoline, H-633
 4-Methoxyquinazoline, Q-14
 8-Methoxy-4-quinolinecarboxaldehyde, H-722
 4-Methoxy-2-quinolinecarboxylic acid, H-725
 6-Methoxy-4-quinolinecarboxylic acid, H-726
 8-Methoxy-2-quinolinecarboxylic acid, H-727
 2-Methoxy-4-quinolinecarboxylic acid, H-723
 6-Methoxy-2,4-quinolinediol, Q-46
 8-Methoxy-2,4-quinolinediol, Q-47
 8-Methoxy-4,7-quinolinediol, Q-48
 2-Methoxyquinoline, Q-50
 3-Methoxyquinoline, Q-49
 4-Methoxy-2(1*H*)-quinolinone, Q-33
 9-(4-Methoxy-2-quinolinyl)-2-nonanone, H-642
 4-Methoxy-*N*-resorcylnthranilic acid, B-67
 11-Methoxyretuline, R-63
 7-Methoxyreumycin, R-66
 (Z)-5α-Methoxyrhazimine, R-73
 10-Methoxyrutaeacarpine, H-361
 1-Methoxyrutaeacarpine, R-168
 2-Methoxyrutaeacarpine, R-168
 4-Methoxy-*N*-salicylanthranilic acid, B-67
 4-Methoxysampangine, S-49
 9-Methoxysampangine, S-49
 3-Methoxysampangine, S-49
 12-Methoxysanguinarine, C-360
 8-Methoxysanguinarine, P-55
 17-Methoxysarpagan *N*-oxide, T-425
 11-Methoxysarpagan-17-ol, G-22
 10-Methoxysarpagan-17-ol, S-89
 17-Methoxysarpagan, T-425
 4-Methoxysecurinan-11-one, P-386
 4-Methoxysecurinine, P-386
N-Methoxyseptorine, S-249
N-Methoxyseptorinol, S-249
 10-Methoxysitsirikine, S-325
 6-Methoxyspirobrassinin, S-426
 1-Methoxyspirobrassinin, S-426
N-Methoxyspirobrassinol methyl ether, S-427
N-Methoxyspirobrassinol, S-427
 13-Methoxystaphimine, S-494
 13-Methoxystaphisagrine, S-495
 10-Methoxystaurosporine, S-498
 4'-Methoxystemofoline, S-515
 Methoxystemokerrin *N*-oxide, S-516
 5-Methoxystrictamine, S-576
 11-Methoxystrictamine, S-576
 16-Methoxystrychnine, P-726
 3-Methoxystrychnine, P-726
 10-Methoxystrychnine, S-589
 11-Methoxystrychnine, S-589
 10-Methoxystrychnobrasiline, S-594
 11-Methoxystrychnofendlerine, S-594
N-(*p*-Methoxystyryl)benzamide, A-242
N-4-Methoxystyrylcinnamide, M-287
 3'-Methoxytabernaemontanine C, C-616
 11-Methoxytabersonine, T-11
 8-Methoxytaifine, F-223
N-(7-Methoxy-4-tetradecenyl)tryptamine, T-639
 10-Methoxy-3,4,5,6-tetrahydrocorynantheol, T-136
 10-Methoxy-3,4,5,6-tetrahydro-18,19-dihydrocorynantheol, T-136
 6-Methoxytetrahydro-β-carboline, T-173
 6-Methoxytetrahydroharman, S-285
 5-Methoxytetrahydroharman, T-202
 1-Methoxytetrahydropropalmine, C-93
 4-Methoxytetrahydrosecurinine, P-386
 10-Methoxytetrahydro-17*S*-usambarensine, U-52
 10-Methoxytetrahydrousambarensine, U-52
 2-Methoxy-1,3-thiazino[6,5-*b*]indol-4(9*H*)-one, C-843
 6-Methoxy-3-(2-thiazolyl)-1*H*-indole, T-372
 4-Methoxy-3,24,25-triazatetracyclo[18.2.1.1^{2,3}.1^{7,10}]pentaacosan-2(25),3,5,7,9,20,22-heptaene, N-285
 6-Methoxy-3,5,7-tribromoindole, T-471
 6-Methoxytrichostachine, W-16
 10-Methoxy-*O*-(3,4,5-trimethoxycinnamoyl)vincamajine, V-113
 β-Methoxy-*N,N,N*-trimethylbenzeneethanaminium, C-695
 5-Methoxytryptamine, M-288
 6-Methoxytryptoline, T-173
 5-Methoxytryptophol, H-505
 10-Methoxysilanine, T-649
 11-Methoxytubotaiwine, T-673
 12-Methoxytubotaiwine, T-673
 19*R*-Methoxytubotaiwine, T-673
 19*S*-Methoxytubotaiwine, T-673
 3-Methoxytyraminebetaxanthin, M-647
 3-Methoxytyramine, H-566
 3-Methoxytyrosine, A-745
 3-Methoxyurabaine, U-41
 8-Methoxyuvariopsine, U-64
 5-Methoxyvasicinone, V-38
 7-Methoxyvasicinone, V-38
 12-Methoxyvellosimine, M-237
 10-Methoxyvellosimine, S-89
 10-Methoxyvillalstonine 4'-*N*-oxide, V-103
 10-Methoxyvillalstonine, V-103
 11-Methoxyvincadifformine *N*-oxide, V-111
 11-Methoxyvincadifformine, V-111
 10-Methoxyvincamajine, V-113
 11-Methoxyvincamajine, V-113
 11-[10-(11-Methoxyvincamajinyl)]vincorine, M-289
 10-Methoxyvincamedine *N*(4)-oxide, V-113
 10-Methoxyvincamedine, V-113
 11-Methoxyvincamedine, V-113
 11-[10-(11-Methoxyvincamedinyl)]vincorine, M-289
 11-Methoxy-Δ¹⁴-vincamenine, E-5
 11-Methoxy-Δ¹⁴-vincamine, V-115
 11-Methoxyvincamine, V-115
 12-Methoxy-Δ¹⁴-vincamine, V-115
 11-Methoxy-Δ¹⁴-vincanol, E-6
 10-Methoxyvinorine, V-194
 4-Methoxy-1-vinyl-β-carboline, M-290
N-Methoxy-1-vinyl-β-carboline, V-139
 3*R*-Methoxyvoacangine, V-170
 12-Methoxyvoaphylline, M-290
 11-Methoxy-Wieland-Gumlich aldehyde, C-115
 4'-Methoxywithasomnine, D-492
 Methoxyxanthocillin X dimethyl ether, X-7
 10-Methoxyyohimbano-17α-ol, Y-16
 10-Methoxy-α-yohimbine, M-291
 10-Methoxy-β-yohimbine, M-291
 10-Methoxyyohimbine, M-291
 11-Methoxy-α-yohimbine, M-292
 11-Methoxyyohimbine, M-292
 10-Methoxyyohimbol, Y-16
 Methscopolamine bromide, S-159
 Methuenine *N*¹⁸-oxide, M-293
 Methuenine, M-293
 Methy-F, D-738
 Methyl acethydroxamylacetate, M-294
 Methyl 7-(3-acetylamino-2,5-dihydroxyphenyl)-7-hydroxy-4,6-dimethyl-2,4-heptadienoate, A-1213
 Methyl [(acetylamino)oxy]acetate, M-294
 Methyl 17-(acetyloxy)akuammilan-16-carboxylate, 10CI, A-231
 Methyl 4-(acetyloxy)-3,6-epoxy-1-methylaspidospermidine-3-carboxylate, C-213
 Methyl 16-[(acetyloxy)methyl]akuammilan-17-oate, A-231
 Methyl akuammilan-17-oate, S-576
 Methyl 3-alkylpyrrole-2-carboxylates, M-295
 Methyl 2-aminobenzoate, A-706
 Methyl 5-(1-aminoethyl)-3-pyridinecarboxylate, A-774
 Methyl 2-[(4-amino-2-methyl-1,4-dioxobutyl)amino]benzoate, D-178
 Methyl allyllate, A-1335
 Methyl aristolochate, M-262
 Methyl aspidofractinine-3-carboxylate, K-78
 Methyl aspidospermidine-3-carboxylate, E-4
 Methyl 3-(benzoyloxy)-8-methyl-8-azabicyclo[3.2.1]octane-2-carboxylate, C-537
 Methyl 3,4-bis(4-hydroxyphenyl)-1*H*-pyrrole-2-carboxylate, L-25
 Methyl *N*'-[3-bromo-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl] γ-ureidobutyrate, B-307
 Methyl 6-bromo-1*H*-indole-3-acetate, B-332
 Methyl 5-bromo-2-pyrrolocarboxylate, B-349
 Methyl 7-bromoreticulatate, R-57
 Methyl 3-carbazolecarboxylate, C-122
 Methyl chanofructosinate, M-296
 Methyl cryptolepinoate, I-111
 Methyl 21,23-deepoxy-23-hydroxyerinin-21-oate, E-182
 Methyl *N*¹-demethoxycarbonylchanofructosinate, M-296
 Methyl *N*¹-demethoxycarbonyl-3-oxo-Δ^{14,15}-chanofructosinate, M-296
 Methyl 4-demethyl-3-oxovobasan-17-oate, P-268
 Methyl deoxydehydrochorismamide, D-224
 Methyl deserpidate, D-687

- Methyl *N'*-(2,3-dibromo-4,5-dihydroxybenzyl) γ -ureidobutyrate, D-312
- Methyl 3,5-dibromo-4-[[3-(dimethylamino)propoxylphenylethyl]carbamate, M-679
- Methyl 4,5-dibromo-1*H*-pyrrole-2-carboxylate, D-339
- Methyl 7,14-dibromoreticulate, R-57
- Methyl 2,3-didehydroaspidospermidine-3-carboxylate, V-111
- Methyl 6,7-didehydro-2,20-cycloaspidospermidine-3-carboxylate, V-132
- Methyl 19,20-didehydro-1,16-cyclocorynan-17-oate, P-518
- Methyl 19,20-didehydro-2-(18,19-didehydroalstophyllan-18-yl)-2,7-dihydro-1,16-cyclocorynan-17-oate, M-11
- Methyl 19,20-didehydro-3,17-dihydroxyajmalan-16-carboxylate, H-139
- Methyl 2,16-didehydro-19,20-dihydroxycuran-17-oate, C-589
- Methyl 2,3-didehydro-6,21-epoxyaspidospermidine-3-carboxylate, M-666
- Methyl 1,2-didehydro-6,7-epoxy-3-hydroxy-8-oxoaspidospermidine-3-carboxylate, M-297
- Methyl 2,3-didehydro-7-ethyl-20,21-dinoraspidospermidine-3-carboxylate, P-733
- Methyl 2,3-didehydro-7-ethyl-*E*,20,21-trinoraspidospermidine-3-carboxylate, I-6
- Methyl 2,3-didehydrohomodaphniphyllate, H-330
- Methyl 2,3-didehydro-20-hydroxyaspidospermidine-3-carboxylate, M-631
- Methyl 2,16-didehydro-19-hydroxycuran-17-oate, E-30
- Methyl 2,16-didehydro-20-hydroxycuran-17-oate, L-224
- Methyl 1,6-didehydro-17-hydroxy-3,16-dimethoxyerythrinan-15-carboxylate, E-204
- Methyl 2,3-didehydro-7-hydroxy-8-methyl-*E*-homo-20,21-dinoraspidospermidine-3-carboxylate, I-10
- Methyl 2,3-didehydro-21-hydroxy-20-oxoaspidospermidine-3-carboxylate, C-193
- Methyl 19,20-didehydro-17-hydroxyohimban-16-carboxylate, D-378
- Methyl 1,6-didehydro-16-methoxy-2,3-[methylenebis(oxy)]erythrinan-15-carboxylate, D-200
- Methyl 16,17-didehydrodromethoxy-19-methyloxayohimban-16-carboxylate, M-298
- Methyl 16,17-didehydro-11-methoxy-19-methyloxayohimban-17-carboxylate, R-53
- Methyl 16,17-didehydro-19-methyloxayohimban-16-carboxylate, A-212
- Methyl 2,3-didehydro-20-oxoaspidospermidine-3-carboxylate, M-630
- Methyl 2,3-didehydro-*E*,20,21-trinoraspidospermidine-3-carboxylate, D-140
- Methyl 15-(14,15-dihydroeburnamenin-14-yl)-1-methylaspidofractinine-3-carboxylate, P-521
- Methyl 4,9-dihydro- β -hydroperoxy- α -[(methoxycarbonyl)amino]-4,6-dimethyl-9-oxo-1*H*-imidazo[1,2-*a*]purine-7-butanoate, W-29
- Methyl 14,15-dihydro-14-hydroxyeburnamenine-14-carboxylate, V-115
- Methyl [(2,3-dihydro-3-hydroxy-2-oxo-1*H*-indol-3-yl)methyl]carbamoithioate, D-811
- Methyl 1,2-dihydro-1-methylakuammilan-17-oate, C-191
- 4-Methyl *N'*[(2,3-dihydro-2-oxo-1*H*-indol-3-yl)acetyl]aspartate, M-299
- Methyl 19,20-dihydro-3-oxovobasan-17-oate, D-935
- Methyl 3,6-dihydroxy-4-methoxy-10-nitro-1-phenanthrenecarboxylate, T-568
- Methyl 3,17-dihydroxy-1-methyl-2,4(1*H*)-cyclo-3,4-secoakuammilan-16-carboxylate, D-410
- Methyl 3,17-dihydroxysarpagan-16-carboxylate, V-171
- Methyl 11,12-dimethoxy- $\Delta^{14,15}$ -chanofruticosinate, M-296
- Methyl 11,12-dimethoxychanofruticosinate, M-296
- Methyl 3,4-dimethoxy-10-nitro-1-phenanthrenecarboxylate, D-636
- Methyl *N'*-[6-[(7,9-dimethoxy-3,6,8-trimethyl-1-oxo-11-phenyl-2,4,10-undecatrienyl)amino]-1-oxo-2,5-hexadienyl]glycinate, C-763
- Methyl [6-[2-(dimethylamino)ethyl]-9-hydroxy-8-methoxy-1-oxonaphtho[1,2-*c*]furan-3(1*H*)-ylidene]acetate, C-381
- Methyl 3,22-dioxovobasan-17-oate, P-268
- Methyl 2,5-epoxy-1,2-dihydroakuammilan-17-oate, P-417
- Methyl 2,5-epoxy-1,2-dihydro-16-(hydroxymethyl)-4,5-secoakuammilan-17-oate, A-1492
- Methyl 17,20-epoxy-19,20-dihydro-18-hydroxysarpagan-16-carboxylate, E-7
- Methyl 4,5-epoxy-5-hydroxy-6,21-cyclo-4,5-secoakuammilan-17-oate, N-47
- Methyl 2,17-epoxy-2-hydroxy-1-methyl-1,17-cyclo-1,2(1*H*,2*H*)-secoakuammilan-16-carboxylate, L-28
- Methyl 6,17-epoxy-1-methylsarpagan-16-carboxylate, D-173
- Methyl 5-ethenyl-4-[2-[(4-hydroxybenzoyl)oxy]ethyl]-3-pyridinecarboxylate, F-125
- Methyl 3-[2-(3-ethyl-5,6-dihydro-1(2*H*)-pyridinyl)ethyl]- α -methylene-1*H*-indole-2-acetate, S-184
- Methyl 4-ethylidene-1,3,4,5,6,7-hexahydro-6-(hydroxymethyl)-2,5-ethano-2*H*-azocino[4,3-*b*]indole-6-carboxylate, V-9
- Methyl 7-ethyl-1,4,5,6,7,8,9,10-octahydro-2*H*-3,7-methanoazacycloundecino[5,4-*b*]indole-9-carboxylate, V-112
- Methyl 4-ethyl-2,3,4,4a,5,6,7,12-octahydro-2-methyl-6-oxopyrido[3',4':4,5]cyclohept[1,2-*b*]indole-12a(1*H*)-carboxylate, E-190
- Methyl 2-ethyl-5,7,12-trihydroxy-4-[(2-hydroxyethyl)amino]-6,11-dioxo-1-naphthacene-carboxylate, A-1080
- Methyl 8-formyl-8,9-dihydro-8,9-dihydroxy-5-oxo-5*H*-carbazole-1-carboxylate, C-636
- Methyl 1-formyl-4,9-dihydro-1,9-dihydroxy-4-oxo-1*H*-carbazole-8-carboxylate, C-636
- Methyl 3-formyl-1*H*-indole-2-sulfinate, B-283
- Methyl 2-formyl-5-methoxymethyl-1*H*-pyrrole-1-butanoate, F-137
- Methyl 2-(2-formyl-5-methoxymethyl-1*H*-pyrrol-1-yl)-3-phenylpropanoate, F-138
- Methyl 6-formyl-4,7,9-trihydroxy-8-methyl-1-phenazinecarboxylate, A-1221
- Methyl 6-formyl-4,7,9-trihydroxy-1-phenazinecarboxylate, L-241
- Methyl harman-3-carboxylate, M-427
- Methyl harmate, H-459
- Methyl 3-heneicosenyl-1*H*-pyrrole-2-carboxylate, M-295
- Methyl 15,16,17,18,19,20-hexahydroohimban-16-carboxylate, D-434
- Methyl 5,7,8,13,13*b*,14-hexahydro-5-(hydroxymethyl)indolo[2',3':3,4]pyrido[1,2-*b*][2,7]naphthyridine-1-carboxylate, C-6
- Methyl 2,3,3a,4,5,6-hexahydro-5-[1-(hydroxymethyl)-1-propenyl]-1*H*-indolo[3,2,1-*de*][1,5]-naphthyridine-6-carboxylate, D-145
- Methyl homodaphniphyllate, H-330
- Methyl homosecodaphniphyllate, H-346
- Methyl 3-hydroxyaspidofractinine-3-carboxylate, K-74
- Methyl 2-[(2-hydroxybenzoyl)amino]-4-hydroxybenzoate, B-67
- Methyl 3-hydroxy-2,4-bis(methylamino)benzoate, D-256
- Methyl 15-hydroxy-*N*¹-demethoxycarbonylchanofruticosinate, M-296
- Methyl 3-hydroxy-4,8-dimethoxy-10-nitro-1-phenanthrenecarboxylate, T-569
- Methyl 6-hydroxy-3,4-dimethoxy-10-nitro-1-phenanthrenecarboxylate, T-568
- Methyl 17-hydroxy-10,11-dimethoxyohimban-16-carboxylate, S-251
- Methyl 17-hydroxyhomodaphniphyllate, H-330
- Methyl [2-[4-hydroxy-2-[2-(4-hydroxyphenyl)ethenyl]phenyl]ethyl]carbamate, G-225
- Methyl 15-hydroxy-12-methoxy-*N*¹-demethoxycarbonylchanofruticosinate, M-296
- Methyl 5-hydroxy-4-methoxymethyl-1*H*-pyrrole-3-carboxylate, H-564
- Methyl 3-hydroxy-4-methoxy-10-nitro-1-phenanthrenecarboxylate, D-636
- Methyl 5-hydroxy-4-methoxy-1*H*-pyrrole-3-carboxylate, D-670
- Methyl 17-hydroxy-10-methoxyohimban-16-carboxylate, M-291
- Methyl 17-hydroxy-9-methoxyohimban-16-carboxylate, V-50
- Methyl 3-hydroxy-1-methyl-2,4(1*H*)-cyclo-3,4-secoakuammilan-17-oate, D-236
- Methyl 15-hydroxy-11,12-methylenedioxy-*N*¹-demethoxycarbonylchanofruticosinate, M-296
- Methyl 19-(hydroxymethyl)-12-methoxyibogamine-18-carboxylate, S-79
- Methyl 19-hydroxy-19-methyloxayohimban-16-carboxylate, A-953
- Methyl 4-[(1-hydroxymethyl-2-phenylethyl)amino]-4-oxobutanoate, M-300
- Methyl 3-(17-hydroxy-1-methylsarpagan-10-yl)vobasan-17-oate, A-28
- Methyl 3-hydroxy-22-oxokopsan-1-carboxylate, K-76
- Methyl 6-(2-hydroxy-2-phenylethyl)-1-methyl-2-piperidinecarboxylate, S-211
- Methyl 17-hydroxysarpagan-16-carboxylate, A-229
- Methyl 17-hydroxysarpagan-16-carboxylate, P-562
- Methyl 17-hydroxyohimban-16-carboxylate, Y-13
- Methyl 17-hydroxyohimban-5,16-dicarboxylate, C-145
- S*-Methyl 1*H*-indole-3-carbothioate, I-85
- Methyl 1*H*-indolethiolcarboxylate, I-85
- S*-Methyl [2-(1*H*-indol-3-yl)ethyl]methylcarbamothioate, G-130
- Methyl (1*H*-indol-3-ylmethyl)carbamoithioate, B-284
- S*-Methyl 1*H*-indol-3-ylmethylcarbamoithioate, B-285
- Methyl iodoacetate, I-163
- Methyl jervine-*N*-3'-propanoate, J-36
- Methyl 8-methoxy-2-carbazolecarboxylate (obso.), H-454
- Methyl 6-methoxy-9*H*-carbazole-3-carboxylate, H-454
- Methyl 4-methoxy- β -carboline-1-carboxylate, H-457
- Methyl 12-methoxychanofruticosinate, M-296
- Methyl 12-methoxy-*N*¹-demethoxycarbonyl- $\Delta^{14,15}$ -chanofruticosinate, M-296
- Methyl 12-methoxy-*N*¹-demethoxycarbonylchanofruticosinate, M-296
- Methyl 12-methoxy-*N*¹-demethoxycarbonyl-3-oxo- $\Delta^{14,15}$ -chanofruticosinate, M-296
- Methyl 1-methoxy-1*H*-indole-3-carboxylate, I-90
- Methyl 10-methoxy-1-methyl-2,4(1*H*)-cyclo-3,4-secoakuammilan-17-oate, V-125
- Methyl 12-methoxy-3-oxochanofruticosinate, M-296
- Methyl 13-(17-methoxy-17-oxovobasan-3 α -yl)ibogamine-18-carboxylate, E-189
- Methyl [(methylamino)carbonyl]carbamate, M-385
- Methyl 2-(methylamino)-5-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-*b*]quinazolin-3-yl)benzoate, A-1045
- Methyl *N*-methylantranilate, M-384
- Methyl 11,12-methylenedioxychanofruticosinate, M-296
- Methyl 11,12-methylenedioxy-*N*¹-demethoxycarbonyl- $\Delta^{14,15}$ -chanofruticosinate, M-296
- Methyl 11,12-methylenedioxy-*N*¹-demethoxycarbonylchanofruticosinate, M-296
- Methyl 11,12-methylenedioxy-*N*¹-demethoxycarbonyl-3-oxo- $\Delta^{14,15}$ -chanofruticosinate, M-296
- Methyl 3-methyl- α -methylene-2,5-dioxo-1-pyrrolidinediacetate, V-85
- S*-Methyl methyl(phenylacetyl)carbamoithioate, N-235
- S*-Methyl methyl(2-phenylethyl)carbamoithioate, N-235
- Methyl (3-methylpyrazinyl) ketone, A-62
- Methyl 2-methyl-3-pyridyl ketone, A-65
- Methyl 2-methyl-4-pyridyl ketone, A-67

- Methyl 3-methyl-4-pyridyl ketone, A-68
Methyl 4-methyl-2-pyridyl ketone, A-63
Methyl 4-methyl-3-pyridyl ketone, A-66
Methyl 6-methyl-2-pyridyl ketone, A-64
Methyl 6-methyl-3-pyridyl ketone, A-69
Methyl 7-methyl-5*H*-2-pyridine-4-carboxylate, R-1
Methyl 4-methylpyrrole-2-carboxylate, M-541
Methyl 2-methyl-2,3,4,9-tetrahydro- β -carboline-3-carboxylate, T-224
Methyl *N*-methyl-5-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-*b*]quinazolin-3-yl)anthranilate, A-1045
Methyl 12-oxodendrobane-2-acetate, D-207
Methyl 3-oxo-2-oxa-7,8-dithia-4,11-diazadodecan-12-oate, P-622
Methyl 1-phenazinecarboxylate, P-319
Methyl *N*-(2-phenylethyl)carbamate, P-343
Methyl platensinoate, P-512
Methyl 2-pyridyl ketone, A-73
Methyl 2-pyrrolyl ketone, A-74
Methyl reserpate, R-50
Methyl reticulate, R-57
Methyl *N*-salicyl-4-hydroxyanthranilate, B-67
Methyl sarpagan-17-oate, T-425
Methyl *N*-succinoylanthranilate, A-706
Methyl 2,3,6,7-tetrahydroaspidospermidine-3-carboxylate, T-11
Methyl 2,16,19,20-tetrahydrocuran-17-oate, A-228
Methyl 2,3,6,7-tetrahydro-10,20-epoxy-19-hydroxyaspidospermidine-3-carboxylate, B-475
Methyl 1,2,19,20-tetrahydro-17-hydroxycuran-16-carboxylate, P-587
Methyl 1,2,14,19-tetrahydro-16-(hydroxymethyl)-condyfolan-16-carboxylate, P-595
Methyl 2,16,19,20-tetrahydro-14-hydroxycuran-17-oate, M-713
Methyl 2,7,19,20-tetrahydro-17-hydroxy-3,7-securan-16-carboxylate, S-510
Methyl tetrahydro-5-(2-methylheptyl)-2-(1*H*-pyrrol-2-yl)-2*H*-1,3-oxazine-4-carboxylate, C-384
Methyl 5,6,7,8-tetrahydro-8-methyl-6-oxo-2,7-naphthyridine-4-carboxylate, J-25
Methyl 2-[(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-*b*]quinazolin-3-yl)amino]benzoate, A-1041
Methyl *N*-(1,2,3,9-tetrahydro-9-oxopyrrolo[2,1-*b*]quinazolin-3-yl)anthranilate, A-1041
Methyl 3,4,5,6-tetrahydro-2-pyridyl ketone, A-76
Methyl tetrahydro-2-(1*H*-pyrrol-2-yl)-2*H*-1,3-oxazine-4-carboxylate, C-383
Methyl 3-(tricosadienyl)-1*H*-pyrrole-2-carboxylate, M-295
Methyl 3-tricosenyl-1*H*-pyrrole-2-carboxylate, M-295
Methyl 3,4,8-trimethoxy-10-nitro-1-phenanthrene-carboxylate, T-569
Methyl 3,4,5-trimethoxy-2-[[2-[(3-pyridinylcarbonyl)amino]benzoyl]amino]benzoate, T-585
*N*³-Methylaaptamine, A-2
4-Methyloconitane-1,2,3,6,8,13,14,15,16,18-decol, M-301
4-Methyloconitane-1,3,6,8,10,13,14,15,16,18-decol, M-302
4-Methyloconitane-1,3,6,8,14,16,18-heptol, M-303
4-Methyloconitane-1,3,8,13,14,16,18-heptol, M-304
4-Methyloconitane-1,5,6,7,8,14,16-heptol, M-305
4-Methyloconitane-1,5,8,13,14,16,18-heptol, M-306
4-Methyloconitane-1,6,7,8,10,14,16-heptol, M-307
4-Methyloconitane-1,6,7,8,12,14,16-heptol, M-308
4-Methyloconitane-1,6,7,8,14,16,18-heptol, M-309
4-Methyloconitane-1,6,8,10,14,16,18-heptol, M-311
4-Methyloconitane-1,6,8,13,14,16,18-heptol, M-312
4-Methyloconitane-1,6,8,14,15,16,18-heptol, M-313
4-Methyloconitane-1,6,8,14,16,18,19-heptol, M-314
4-Methyloconitane-1,6,8,9,14,16,18-heptol, M-310
4-Methyloconitane-1,7,8,9,14,16,18-heptol, M-315
4-Methyloconitane-1,8,10,13,14,15,18-heptol, M-316
4-Methyloconitane-1,3,8,14,16,18-hexol, M-317
4-Methyloconitane-1,5,7,8,14,16-hexol, M-318
4-Methyloconitane-1,5,8,14,16,18-hexol, M-319
4-Methyloconitane-1,6,7,8,12,14-hexol, M-320
4-Methyloconitane-1,6,7,8,14,16-hexol, M-321
4-Methyloconitane-1,6,8,10,14,16-hexol, M-322
4-Methyloconitane-1,6,8,10,14,16-hexol, M-323
4-Methyloconitane-1,7,8,14,16,18-hexol, M-325
4-Methyloconitane-1,7,8,9,14,16-hexol, M-324
4-Methyloconitane-1,8,10,14,16,18-hexol, M-326
4-Methyloconitane-1,8,14,15,16,18-hexol, M-327
4-Methyloconitane-6,7,8,14,16,18-hexol, M-328
4-Methyloconitane-1,2,3,6,8,13,14,16,18-nonol, M-329
4-Methyloconitane-1,2,6,8,13,14,15,16,18-nonol, M-330
4-Methyloconitane-1,3,6,8,13,14,15,16,18-nonol, M-331
4-Methyloconitane-1,6,8,10,13,14,15,16,18-nonol, M-332
4-Methyloconitane-1,2,6,8,13,14,16,18-octol, M-333
4-Methyloconitane-1,3,5,8,13,14,16,18-octol, M-334
4-Methyloconitane-1,3,6,8,13,14,15,16-octol, M-335
4-Methyloconitane-1,3,6,8,13,14,16,18-octol, M-336
4-Methyloconitane-1,3,6,8,14,15,16,18-octol, M-337
4-Methyloconitane-1,3,7,8,13,14,16,18-octol, M-338
4-Methyloconitane-1,6,7,8,10,14,16,18-octol, M-339
4-Methyloconitane-1,6,7,8,12,14,16,18-octol, M-340
4-Methyloconitane-1,6,7,8,14,15,16,18-octol, M-341
4-Methyloconitane-1,6,8,12,13,14,16,18-octol, M-342
4-Methyloconitane-1,6,8,13,14,15,16,18-octol, M-343
4-Methyloconitane-1,7,8,9,10,14,16,18-octol, M-344
4-Methyloconitane-6,8,10,13,14,15,16,18-octol, M-345
4-Methyloconitane-1,3,8,14,16-pentol, M-346
4-Methyloconitane-1,6,8,14,16-pentol, M-347
4-Methyloconitane-1,7,8,14,16-pentol, M-348
4-Methyloconitane-1,8,10,14,16-pentol, M-349
4-Methyloconitane-1,8,12,14,16-pentol, M-350
4-Methyloconitane-1,8,10,14,16-pentol, M-351
4-Methyloconitane-1,8,14,16,18-pentol, M-352
4-Methyloconitane-1,7,8,14-tetrol, M-353
4-Methyloconitane-1,8,10,14-tetrol, M-354
4-Methyloconitane-1,8,14,16-tetrol, M-355
4-Methyloconitane-1,14,16-triol, M-357
4-Methyloconitane-1,8,14-triol, M-356
4-Methyloconit-15-ene-1,3,6,8,13,14,18-heptol, M-366
4-Methyloconit-19(*N*)-ene-1,6,7,8,10,14,16-heptol, M-358
4-Methyloconit-19(*N*)-ene-1,6,7,8,14,16,18-heptol, M-359
4-Methyloconit-19(*N*)-ene-1,6,8,13,14,16,18-heptol, M-360
4-Methyloconit-19(*N*)-ene-1,7,8,9,14,16,18-heptol, M-361
4-Methyloconit-2-ene-1,6,7,8,10,14,16-heptol, M-362
4-Methyloconit-2-ene-1,6,7,8,14,16,18-heptol, M-363
4-Methyloconit-2-ene-1,6,8,13,14,16,18-heptol, M-364
4-Methyloconit-8(15)-ene-1,3,6,13,14,16,18-heptol, M-365
4-Methyloconit-19(*N*)-ene-1,3,6,8,14,16-hexol, M-367
4-Methyloconit-19(*N*)-ene-1,6,7,8,14,16-hexol, M-368
4-Methyloconit-19(*N*)-ene-1,6,8,14,16,18-hexol, M-369
4-Methyloconit-19(*N*)-ene-1,7,8,14,16,18-hexol, M-370
4-Methyloconit-2-ene-1,6,7,8,14,16-hexol, M-371
4-Methyloconit-8(15)-ene-1,6,13,14,16,18-hexol, M-372
4-Methyloconit-19(*N*)-ene-1,3,6,8,13,14,15,16,18-nonol, M-373
4-Methyloconit-19(*N*)-ene-1,6,8,14,16-pentol, M-374
4-Methyloconit-19(*N*)-ene-1,8,14,16,18-pentol, M-375
4-Methyloconit-19(*N*)-ene-7,8,10,14,16-pentol, M-376
4-Methyloconit-8(15)-ene-1,6,14,16,18-pentol, M-377
4-Methyloconit-19(*N*)-ene-1,8,14,16-tetrol, M-378
4-Methyloconit-2-ene-1,6,14,16-tetrol, M-379
4-Methyloconit-8(15)-ene-1,14,16,18-tetrol, M-380
4-Methyloconit-15-ene-1,14,18-triol, M-381
10-Methyl-9-acridanone, M-382
10-Methyl-9(10*H*)-acridinone, M-382
N-Methylacridone, M-382
10-Methylacridone, M-382
N-Methylactinodaphnine, D-346
O-Methylacutifoline, L-288
1-Methyladenine, A-842
2-Methyladenine, A-843
3-Methyladenine, A-844
3-Methyladenosine, A-844
N-Methyladlumine, A-146
N-Methyladrenaline, A-152
*O*³-Methyladrenaline, N-303
7-*O*-Methyalaegeline, A-153
N-Methyalaerophobin 2, A-155
4-Methyalaeruginic acid, D-454
Methyalaerin, H-445
1'-*N*-Methylageliferin, A-190
*N*⁶-Methylagmatine, A-719
N^b-Methylajmaline, A-216
17-*O*-Methylakagerine, A-222
O-Methylaknadimine, H-70
O-Methylakuammine, A-232
N-Methylalbine (incorr.), A-554
O-Methylalfileramine, A-260
 ω -Methylallophanic acid, M-385
N-Methylallosedridine, S-216
N-Methylaloperidine, A-637
O-Methylalpinigine, A-638
N-Methylamericine, A-690
Methylamine, M-383
 α -Methylaminoacetophenone, A-694
3'-Methylaminoavarone, A-1560
4'-Methylaminoavarone, A-1560
2-(Methylamino)benzamide, M-384
2-(Methylamino)benzoic acid, M-384
N^b-(2-Methylaminobenzoyl)tryptamine, T-639
3-[4-(Methylamino)-1-butenyl]pyridine, M-232
4-(4-Methylaminobutyl)- β -carboline, B-291
[(Methylamino)carbonyl]carbamic acid, M-385
4-(Methylamino)cycloheptanone, P-399
6-(Methylamino)-1,3-dihydro-2*H*-purin-2-one, A-818
3-(Methylamino)-15,18-dihydroxy-18-ikemaoyl-20-(dimethylamino)pregn-5-ene, D-270
2-(Methylamino)ethylamine, E-225
4-[2-(Methylamino)ethyl]-1,2-benzenediol, E-98
 α -[1-(Methylamino)ethyl]benzenemethanol, M-386
5-[2-(Methylamino)ethyl]-1,2,3-benzenetriol, T-576
4-(β -Methylaminoethyl)catechol, E-98
3-(2-Methylaminoethyl)indole, M-582
3-[2-(Methylamino)ethyl]-1*H*-indol-2-ol, H-754
3-(2-Methylaminoethyl)oxindole, H-754
4-[2-(Methylamino)ethyl]phenol, H-667
2-(Methylamino)ethylphosphonic acid, A-773
4-[2-(Methylamino)ethyl]pyrocatechol, E-98
1-(2-Methylaminoethyl)-3,4,6,7-tetramethoxyphenanthrene, M-559
1-(2-Methylaminoethyl)-2,3,4-trimethoxy-6,7-methylenedioxyphenanthrene, T-328
2-(Methylamino)hypoxanthine, G-210
2-Methylaminoisocaproic acid, L-153

- 7-(Methylamino)-5,8-isoquinolinedione, A-831
 α -[(Methylamino)methyl]benzenemethanol, A-864
 α -[(Methylamino)methyl]benzyl alcohol, A-864
3-(Methylaminomethyl)indole, I-97
 α -(Methylamino)-*N*-[7-(1-methylpropyl)-5,8-dioxo-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraen-4-yl]benzenepropanamide, C-600
3-Methylamino-*N*-norcona-3,5-diene, R-39
2-(Methylamino)-3-oxo-3*H*-phenoxazine-8-carboxylic acid, A-853
2-(Methylamino)-1-phenethanol, A-864
6-[2-(Methylamino)phenyl]-1,3-benzodioxole-5-methanol, I-186
2-(Methylamino)-1-phenylethanol, A-864
2-(Methylamino)-1-phenyl-1-propanol, M-386
2-Methylamino-1-phenyl-1-propanone, A-870
5-(3-Methylamino-3-phenylpropanoyl)nicotaxine, N-188
2-[(Methylamino)phenyl](2,4,6-trimethoxyphenyl)methanone, T-55
3-Methylaminopregna-5,17(20)-dien-16-one, A-877
20-Methylaminopregnan-3-ol, A-881
20-(Methylamino)pregnan-3-one, A-883
3-(Methylamino)pregnan-20-one, A-882
20-(Methylamino)pregn-5-ene-3,16-diol, A-884
20-Methylaminopregn-5-en-3-ol, A-886
20-Methylaminopregn-4-en-3-one, A-891
3 β -(Methylamino)pregn-17(20)-en-16-one, A-890
3-Methylaminopregn-5-en-16-one, A-887
3-Methylaminopregn-5-en-20-one, A-888
4-[(3-Methylamino)propyl]phenol, H-350
2-(Methylamino)-4(1*H*)-pteridinone, A-816
6-(Methylamino)purine, A-137
(4-Methylamino)-2(1*H*)-pyrimidinone, C-961
7-(Methylamino)-3*H*-pyrrolo[2,3-*c*]isoquinoline-6,9-dione, M-387
2'-Methylamino-2,4,6-trimethoxybenzophenone, T-55
N'-Methylammodendrine, A-930
Methylammonium nitrate, M-383
N-Methylanabasine, P-460
N-Methylanacycline, A-968
N-Methylanatabine, T-143
O-Methylancistrocladine, A-982
6-*O*-Methylancistrocladinine, A-982
N-Methylancistrorobertsonine D, A-990
N-Methylancistroretoriline A, A-980
N-Methylandrachamine, B-187
O-Methylandrocybine, A-1005
N-Methylangustifoline, A-1013
N^B-Methylangustilobine B, A-1015
19-*O*-Methylangustoline, A-1020
N-Methylanhalamine, T-236
O-Methylanhalamine, T-236
O-Methylanhalonidine, T-237
N-Methylanhydroscumannificine, S-145
8-Methylanibacanine, M-388
N-Methylanolobine, A-1066
O-Methylanolobine, A-1066
12-*O*-Methylanomaline, A-1067
N-Methylanonaine, R-120
N-Methylantranilic acid 4-methoxyphenethylamide, D-900
N-Methylantranilic acid *N*-methyl-4-methoxyphenethylamide, D-900
N-Methylantranilic acid *N*-methylphenethylamide, D-900
N-Methylantranilic acid phenethylamide, D-900
N-Methylantranilic acid, M-384
*N*⁴-Methylantirrhine, A-1328
N-Methylapateline, A-1331
N-3'-Methylaplysinopsin, A-1357
Methylaposafrane, P-328
3 α -Methylaraguspangine C, A-1372
N-Methylarmepavine *N*-oxide, A-1444
(+)-*O*-Methylarmepavine, A-1444
(-)-*O*-Methylarmepavine, A-1444
11-*O*-Methylartavenustine, X-30
N-Methylasimilobine glucoside, D-531
N-Methylasimilobine, D-531
*N*¹-Methylaspidofractinine, A-1494
N-Methylaspidospermatidine, A-1497
 α -Methylaspidospermidine, A-1498
N-Methylasooaninium, A-1508
O-Methylasterriquinone, A-1520
N-Methylatalaphyllidine, A-1527
N-Methylatalaphylline, T-546
N-Methylatalaphyllinine, S-278
N-Methylatanine, A-1529
O-Methylatheroline, O-178
N-Methylatherosperminium, A-1532
12-*O*-Methylatherospermine, T-286
O-Methylathrocressine, C-588
Methylatropine nitrate, T-632
8-Methylatropinium bromide, T-632
4-Methyl-1*H*-1-aza-2,9,10-anthracenetrione, M-97
3-Methyl-16-azabicyclo[10.3.1]hexadecan-1(16),12,14-triene, M-774
9-Methyl-9-azabicyclo[3.3.1]nonan-3-one, A-1580
1-Methyl-9-azabicyclo[3.3.1]nonan-3-one, M-389
8-Methyl-8-azabicyclo[3.2.1]octane-2,6-diol, A-1582
8-Methyl-8-azabicyclo[3.2.1]octane-3,6-diol, M-391
8-Methyl-8-azabicyclo[3.2.1]octane-1,3-diol, M-390
8-Methyl-8-azabicyclo[3.2.1]octane-3,6,7-triol, M-392
8-Methyl-8-azabicyclo[3.2.1]octan-3-ol, M-393
8-Methyl-8-azabicyclo[3.2.1]octan-1-ol, P-399
8-Methyl-8-azabicyclo[3.2.1]octan-3-one, M-394
8-Methyl-8-azabicyclo[3.2.1]oct-2-ene-2-carboxylic acid, M-395
1-(8-Methyl-8-azabicyclo[3.2.1]oct-2-en-2-yl)ethanone, F-47
8-Methyl-8-azabicyclo[3.2.1]oct-3-yl 1,2-dithiolane-3-carboxylate, B-369
8-Methyl-8-azabicyclo[3.2.1]oct-3-yl α -(hydroxymethyl)benzeneacetate, T-632
8-Methyl-8-azabicyclo[3.2.1]oct-3-yl α -methylenebenzeneacetate, A-1359
(8-Methyl-8-azabicyclo[3.2.1]oct-2-yl)phenylmethanone, F-46
1-Methyl-4-azafluorenone, O-93
4-Methyl-15-aza-*n*-homoeergosta-8,14,24(28)-trien-3-ol, M-396
4-Methyl-15-azasterol, M-396
(Methyl-*ON*-azoxy)methanol, M-397
Methylazoxymethanol, M-397
*O*⁴-Methylbalfouronium, B-14
O-Methylbalsamide, M-18
5-Methylbarbituric acid, M-540
3-*O*-Methylbatrachotoxin, B-36
7-*O*-Methylbelarine, I-329
7-Methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline-5,6(7*H*)-dione, C-272
5-Methyl-1,3-benzodioxolo[5,6-*c*][1,3]dioxolo[4,5-*j*]phenanthridinium(1+), A-1566
13-Methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*j*]phenanthridinium(1+), S-56
5-Methyl-1,3-benzodioxolo[5,6-*c*][1,3]dioxolo[4,5-*j*]phenanthridin-6(5*H*)-one, D-404
13-Methyl[1,3]benzodioxolo[5,6-*c*]-1,3-dioxolo[4,5-*j*]phenanthridin-14(13*H*)-one, D-510
3-Methylbenzoic acid, M-398
10-Methyl-8*H*-benzo[*b*]pyrido[4,3,2-*de*][1,8]phenanthroline-8,11(10*H*)-dione, A-944
10-Methyl-8*H*-benzo[*b*]pyrido[4,3,2-*de*][1,8]phenanthroline-8,9(10*H*)-dione, N-99
4-Methylbenzo[*g*]quinoline-5,10-dione, M-399
4-Methylbenzo[*g*]quinoline-2,5,10(1*H*)-trione, M-97
2-Methylbenzothiazole, M-400
2-[(4-Methylbenzyl)amino]benzoic acid, A-706
O-Methylberbamine, I-328
2'-*N*-Methylberbamine, I-328
N-Methylbicycloataphylline, M-401
 α -Methylbiotin, M-402
4-Methyl-2,3'-bipyridine, M-403
4-Methyl-3,3'-bipyridine, M-404
5-Methyl-2,3'-bipyridine, M-405
N-Methyl-3,3'-bis(methylamino)propylamine, D-255
1-Methyl-3,3-bis(3-methyl-2-butenyl)-2,4(1*H*,3*H*)-quinolinedione, M-406
5-Methyl-2,3,7,8-bis(methylenedioxy)benzo[*c*]phenanthridinium(1+), S-56
5-Methyl-2,3,7,8-bis(methylenedioxy)benzo[*c*]phenanthridin-6(5*H*)-one, D-510
2-Methyl-6,7-bis(methylthio)-3,5,8(2*H*)-isoquinolinetriene, P-255
N-Methylboldine, B-243
4-Methylborreverine, B-253
9-*O*-Methylbostrycoidin, B-255
1'-*N*-Methyl-2'-bromoageliferin, A-190
1'-*N*-Methyl-2-bromoageliferin, A-190
O-Methylbufotenine *N*-oxide, M-288
O-Methylbufotenine, M-288
O-Methylbulbocarpine α -*N*-oxide, B-397
O-Methylbulbocarpine β -*N*-oxide, B-397
N-Methylbulbocarpine, B-397
O-Methylbulbocarpine, B-397
*N*⁸-Methylburnamine, B-420
6-(3-Methyl-1,3-butadienyl)-1*H*-indole-3-carboxylic acid, M-408
6-(3-Methyl-1,3-butadienyl)-1*H*-indole, M-407
5-(3-Methyl-1,3-butadienyl)-1*H*-indole, M-412
3-Methylbutanamide, M-409
3-Methyl-1-butanamine, M-420
N-Methyl-1,4-butanediamine, B-422
3-Methylbutanoic acid *N*-methyl-*N*-(2-phenylethyl)amide, T-303
3-Methylbutanoic acid, M-409
4-[2-[(2-Methylbutanoyl)amino]ethyl]imidazole, H-307
N-(3-Methylbutanoyl)-3-nitrotyramine, H-640
14-(2-Methylbutanoyl)nudicaulidine, M-321
13-(2-Methylbutanoyloxy)azatine, A-1609
3-(2-Methylbutanoyloxy)tropone-6,7-diol, M-392
N-(3-Methylbutanoyl)pyrrothine, A-759
*O*⁷-(3-Methylbutanoyl)retroecine *N*-oxide, T-188
*O*⁷-(3-Methylbutanoyl)retroecine, T-188
N-(2-Methylbutanoyl)tyramine, H-667
15-*O*-(2-Methylbutanoyl)-3-*O*-veratroylprotoverine, P-674
2-Methyl-2-buten-1-carboxylic acid, M-506
2-Methyl-2-butenic acid 8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester, T-404
3-Methyl-2-butenic acid *N*-methyl-*N*-(2-phenylethyl)amide, T-303
7-(3-Methyl-2-butenyl)-1*H*-indole, M-414
9-(3-Methyl-2-butenyl)adenine, A-137
1-(3-Methyl-2-butenyl)adenine, A-137
*N*⁶-(3-Methyl-2-butenyl)adenine, Z-11
*N*⁶-(3-Methyl-2-butenyl)agmatine, A-719
6-(3-Methyl-2-butenylamino)purine, Z-11
N-(3-Methyl-2-butenyl)guanidine, M-410
O-(3-Methyl-3-butenyl)halfordinol, H-12
3-(3-Methyl-2-butenylidene)-2-indolinone, D-471
6-(3-Methyl-2-butenyl)-1*H*-indole-3-acetonitrile, H-497
6-(3-Methyl-2-butenyl)-1*H*-indole-3-carboxylic acid, P-617
5-(3-Methyl-2-butenyl)-1*H*-indole-2,3-dione, M-415
6-(3-Methyl-2-butenyl)-1*H*-indole-2,3-dione, M-416
6-(3-Methyl-2-butenyl)-1*H*-indole-3-ethanol, H-497
5-(3-Methyl-2-butenyl)-1*H*-indole, M-412
3-(3-Methyl-2-butenyl)-1*H*-indole, M-411
6-(3-Methyl-2-butenyl)-1*H*-indole, M-413
7-(3-Methyl-2-butenyl)-1*H*-indole, M-414
6-(3-Methyl-2-butenyl)-1*H*-indol-2-ol, H-701
3-(3-Methyl-2-butenyl)-4-[(3-methyl-2-butenyl)oxy]-2(1*H*)-quinolinone, A-1529
4-(3-Methyl-2-butenyloxy)phenylacetonitrile, H-672
N-[2-[4-[(3-Methyl-2-butenyl)oxy]phenyl]ethyl]-3-phenyl-2-propenamide, H-667
3-[4-(3-Methyl-2-butenyloxy)phenyl]-4-(2-methylpropyl)-1*H*-pyrrole-2,5-dione, H-684
3-[5-[4-[(3-Methyl-2-butenyl)oxy]phenyl]-2-oxazolyl]pyridine, H-12
4-[(3-Methyl-2-butenyl)oxy]-2(1*H*)-quinolinone, H-53

- 4-(3-Methyl-2-butenyl)paspalinine, P-111
 (3-Methyl-2-butenyl)paspalinine, P-110
 6-(3-Methyl-2-butenyl)-1-phenazinecarboxylic acid, M-417
 9-(3-Methyl-2-butenyl)-1-phenazinecarboxylic acid, M-418
 3-(3-Methyl-2-butenyl)-3*H*-purin-6-amine, T-451
N-(3-Methyl-2-butenyl)-1*H*-purin-6-amine, Z-11
N-(3-Methyl-2-butenyl)putrescine, B-422
 4-(3-Methyl-2-butenyl)tryptophan, M-419
N-(3-Methylbutyl)acetamide, M-420
 3-Methyl-1-butylamine, M-420
 4-[(3-Methylbutyl)amino]benzeneacetamide, J-47
N-(2-Methylbutyl)-2,4-decadienamide, D-104
N-(3-Methylbutyl)-2,8-decadiene-4,6-diynamide, D-103
N-(2-Methylbutyl)-2,4-dodecadiene-8,10-diynamide, D-882
N-(2-Methylbutyl)-2-dodecene-8,10-diynamide, D-897
 1-(3-Methylbutyl)-3-pyrrolidinemethanol, M-421
 2-(3-Methylbutyl)-1*H*-pyrrolo[2,3-*b*]pyridine, M-422
N-(3-Methylbutyl)-2,4-tetradecadiene-8,10-diynamide, T-121
N-(3-Methylbutyl)-2,4,12-tetradecatriene-8,10-diynamide, T-132
N-(2-Methylbutyl)-2,4-undecadiene-8,10-diynamide, U-26
N-(2-Methylbutyl)-2-undecene-8,10-diynamide, U-30
 14-(2-Methylbutyl)browniine, M-309
 14-(2-Methylbutyl)dictyocarpine, M-307
 2-Methylbutylgermine, G-70
 7-(2-Methylbutyl)intermediate, I-160
 7-(3-Methylbutyl)intermediate, I-160
 7-(2-Methylbutyl)lycopsamine, I-160
 7-(3-Methylbutyl)lycopsamine, I-160
 13-(2-Methylbutyl)lupanine, H-554
 6-(α -Methylbutyryloxy)-3-tigloyloxytropene, M-391
 2-Methylbutyrylyzgyadenine, Z-36
N-Methylbuxene M, C-848
N-Methylbuxene, C-848
N-Methylbuxifoline, B-471
*N*³-Methylbuxupapine, B-483
N-Methylcaaverine, D-531
O-Methylcadabicine, C-4
N-(4-*O*-Methyl-*trans*-caffeoyl)-3-*O*-methyl dopamine, H-566
N-Methylcalycinine, D-846
O-Methylcalycinine, D-846
N-Methylcalystegine B₂, A-1587
N-Methylcalystegine C₁, A-1586
 1-Methylcamalexin, T-372
N-Methyl- β -canadine, C-73
O-Methylcandicine, M-280
 3-Methylcanthin-2,6-dione, I-105
 3-Methylcanthin-5,6-dione, P-413
 6-Methylcanthin-4-one, N-302
cis-*N*-Methylcapaurine, C-93
O-Methylcapaurine, C-93
 3-(Methylcarbamoyl)pyridine, P-895
 3-Methyl-9*H*-carbazole-9-carboxaldehyde, M-424
 3-Methyl-9*H*-carbazole-2,5-diol, D-616
 6-Methyl-9*H*-carbazole-1,7-diol, D-614
 6-Methyl-9*H*-carbazole-2,3-diol, D-615
 3-Methyl-9*H*-carbazole-1,6-diol, D-613
 3-Methyl-9*H*-carbazole-2,6-diol, D-617
 3-Methyl-9*H*-carbazole-2,8-diol, D-618
 3-Methyl-1*H*-carbazole-1,4(9*H*)-dione, M-425
 1-Methyl-9*H*-carbazole, M-423
 3-Methyl-9*H*-carbazole, M-424
 3-Methyl-9*H*-carbazole-1,6,7-triol, T-559
 6-Methyl-9*H*-carbazol-1-ol, H-584
 6-Methyl-9*H*-carbazol-3-ol, H-587
 7-Methyl-9*H*-carbazol-2-ol, H-586
 3-Methyl-9*H*-carbazol-2-ol, H-585
 6-Methyl-9*H*-carbazol-4-ol, H-588
 3-Methyl-9*H*-carbazol-1-ol, H-583
N-Methyl- β -carboline-3-carboxamide, C-133
 1-Methyl- β -carboline-3-carboxylic acid, M-427
 1-Methyl- β -carboline, M-426
 1-Methyl-5-carbomethoxy- α -pyridone, H-711
O-Methylcaryachine, E-220
N-Methylcaryachinium, E-220
N-Methylcassine, C-183
O-Methylcassifoline, O-27
O-Methylcassiformine, P-209
 6-*O*-Methyl-*cis*-cephaloside, I-167
 6-*O*-Methyl-*trans*-cephaloside, I-167
 Methylcephaleine, E-72
 12-*O*-Methylcepharanoline, C-275
N-Methylceratinamine, M-679
 18-*O*-Methylchaeropsine, C-318
 1-*O*-Methylchitraline, P-16
N-Methylciliane, A-773
O-Methylcinnamic acid, M-513
O-Methylcinnamylaurine, C-453
 4-Methylcinnoline, M-428
O-Methylcissampereine, W-9
O-Methylcissampereine, W-9
O-Methylcoccoline, E-202
O-Methylcocovine, C-543
N-Methylcocolaurine, H-275
O-Methylcocosoline 2'- β -*N*-oxide, I-339
O-Methylcocosoline, I-339
O-Methylcocosuline, I-339
 6'-*O*-Methylcocosuline, C-552
 6-*O*-Methylcodeine, M-704
N-Methylcodonocarpine, C-556
N-Methylcolchicine, D-92
 13-Methylcolumbamine, D-154
 8-*O*-Methylcolumbianine, M-352
N-Methyl-(+)-coniine, P-652
N-Methyl(-)-coniine, P-652
 8-*O*-Methylconsolarine, M-321
 3-*O*-Methylconstriocytosine, C-624
 13-Methylcoptisine, C-699
O-Methylcorpaine, R-9
N-Methylcorydaldine, I-305
N-Methylcorydalmine, C-677
O-Methylcorydine *N*-oxide, T-248
N-Methylcorydine, C-682
O-Methylcorydine, T-248
N-Methylcorypalline, T-158
O-Methylcorypalline, T-158
 1-Methylcorypalline, T-181
N-Methylcorypallinium, T-158
N-Methylcorypalmine, C-694
N-Methylcrinasidine, C-744
 3-(3-Methylcrotonyl)cassaine, C-178
 6-Methylcrotsparine, P-643
N-Methylcrotsparine, P-643
N-Methylcrotsparine, C-781
O-Methylcularicine, C-804
N-Methylcularine, C-805
 7-*O*-Methyl(-)-curine, C-812
 12-*O*-Methyl(+)-curine, C-812
 12-*O*-Methyl(-)-curine, C-812
 7-*O*-Methylcuspidaline, D-88
N-Methylclobuxoxazine, C-851
 Methylcyclohexylprodigiosin, M-429
 4-Methyl-1*H*-cyclohepta[1,2-*d*:3,4-*d'*]diimidazole-2,8-diamine, P-737
 4-Methyl-1*H*-cyclohepta[1,2-*d*:4,5-*d'*]diimidazole-2,6-diamine, P-94
N-Methyl-2-cyclohexylethylamine, C-869
N-Methylcyclokoreanine B, C-917
 Methylcyclooctylprodigiosin, M-430
N-Methylcycloorfeibuxine C, C-883
O-Methylcyclovirobuxine D, C-917
N^b-Methylcycloobuxoviricine, C-884
*N*¹-Methylcylindrocarine, C-929
N-Methylcytisine, C-938
 5-Methylcytosine, A-803
 1-Methylcytosine, C-961
 3-Methylcytosine, C-961
N-Methylcytosine, C-961
 Methyldambullin, D-12
N-Methyldanguyelline, P-209
N-Methyl daphnandrine, O-1
 7-*O*-Methyl daphnoline, O-1
 12-*O*-Methyl daphnoline, O-1
 2-*N*-Methyl daphnoline, O-1
O-Methyl dauricine 2'-*N*-oxide, D-88
O-Methyl dauricine 2-*N*-oxide, D-88
O-Methyl dauricine, D-88
N-Methyl deacetylcolchicine, D-92
N^b-Methyl deacetylpicraline, B-420
O-Methyl decarine, C-356
N-(9-Methyldecyl)sulfamic acid, M-431
O-Methyl dehydroisopiline, T-521
O-Methyl dehydrojoubertamine, J-49
 12'-*O*-Methyl dehydrodrokhatine, K-56
 14-*O*-Methyl delcaroline, M-339
 6-*O*-Methyl delcorine, M-309
 14-*O*-Methyl delcosine, M-309
 6-*O*-Methyl delpheline, M-321
 1-*O*-Methyl delphisine, M-323
N-Methyl demecolone, D-92
N-Methyl-14-*O*-demethylepiporphyroxine, G-87
N-Methyl-7-*O*-demethylpeinamine, P-171
N-Methyl-4'-*O*-demethyl-*N*,7*a*-seco mesembradione, A-60
N-Methyl de-*O*-methyltorulosine, T-434
N-Methyl dendrobine, D-210
O-Methyl deoxopunjabine, P-800
N-Methyl-1-deoxynojirimycin 6-phosphate, H-616
N-Methyl-1-deoxynojirimycin, H-616
N^b-Methyl deoxysarpagine, T-425
O-Methyl-*N*-depropionylaspadolimine, P-924
 α -Methyl dethiobiotin, M-432
 Methyl diacetamide, M-383
 7-Methyl-1,5-diazacyclotetradecane, M-433
 15-Methyl-15,17-diazatetracyclo[12.2.2.1^{3,7}.1^{3,7}]jicosane-3(20),12(19)-diene-6,9-dione, H-153
 1'-*N*-Methyl-2,2'-dibromoagelliferin, A-190
*N*¹-Methyl dibromoisophakellin, D-328
*N*⁷-Methyl dibromophakellin, B-343
N-Methyl dicitrinium, D-346
*N*¹-Methyl-14,15-didehydroaspido fractinine, A-1494
O-Methyl-1,2-didehydro triphyphylline, D-793
*N*¹-Methyl-14,15-didehydro tuboxenine, D-674
N^a-Methyl-2,16-dihydro kuammicine *N*^b-methosalt, R-63
N-Methyl dihydroberberine, L-23
 8-Methyl dihydrochelerythrine, B-239
 8-*O*-Methyl dihydrochelerythrine, D-407
N-Methyl-3,4-dihydro ellipticine, E-67
 33-Methyl dihydrohalichondramide, H-15
O-Methyl dihydrojoubertamine, J-49
N-Methyl dihydroquinicine, Q-17
N-Methyl dihydroquinicol, Q-17
*N*²-Methyl-3',4'-dihydrodrosambarensine, U-52
*N*⁴-Methyl-5',6'-dihydrodrosambarensine, U-52
N^b-Methyl-3',4'-dihydrodrosambarensine, U-52
 Methyl dihydrowarifteine, W-9
N-Methyl-2-(3,4-dihydroxyphenyl)ethylamine, E-98
N-Methyl-3,4-dimethoxy- β -hydroxyphenethylamine, M-18
N-Methyl-6,7-dimethoxyisoquinolone, I-305
N-Methyl-4-dimethylallyltryptophan, M-419
 5'-*O*-Methyl dioncopeltine A, D-793
N-Methyl dioncophylline A, D-793
O-Methyl dioncophylline A, D-793
 8-*O*-Methyl dioncophyllinol B, D-796
N-Methyl-10,22-dioxokopsane, D-805
 4-Methyl[1,3]dioxolo[4,5-*h*]-1,3-dioxolo[6,7]indeno[2,1-*a*][3]benzazepinium, L-17
 7-Methyl-1,3-dioxolo[4,5-*a*]furo[2,3-*c*]acridin-12(7*H*)one, C-342
 5-Methyl-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinolin-9(5*H*)-one, F-221
 6-Methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5(6*H*)-one, I-305
 5-Methyl[1,3]dioxolo[4,5-*j*]phenanthridinium(1+), M-448
 1-Methyl-2,6-diphenacylpiperidine, L-218
 5-Methyl-3,7-diphenyl-2-oxo-5-azabicyclo[2.2.1]-heptan-6-one, C-855
 1-Methyl- α,α' -diphenyl-2,6-piperidinediethanol, L-218
N-Methyl-*N,N'*-diphenylurea, D-818
 1-Methyl-3,3-diprenyl-2,4-quinolinedione, M-406
 7-Methyl-1,3-diprenylxanthine, X-5

- 2-(Methyldithio)-1*H*-indole-3-carboxaldehyde, M-434
- 3-Methyl-2-dodecenoic acid *p*-hydroxyphenethylamide, H-679
- N*^b-(3-Methyl-2-dodecenoyl)tryptamine, T-639
- N*-Methyldomesticine, N-29
- O*-Methyldomesticine, N-29
- N*-Methyldopamine 4-*O*-dihydrogen phosphate, E-98
- N*-Methyldopamine, E-98
- 4-*O*-Methyldopamine, H-565
- 3-*O*-Methyldopamine, H-566
- 6-*O*-Methyldryadodaphnine, D-944
- 12-*O*-Methyldryadodaphnine, D-944
- N*-Methylduguevanine, D-950
- N*-Methyldysideathiazole, D-970
- O*-Methyleburnamine, E-6
- Methylecgonine, E-8
- 2-Methyl-2,6-eicosadienoic acid, M-435
- N*-Methylellipticine, E-67
- N*-Methylemerrillicine, E-70
- α -Methylene- β -alanine, A-841
- 14,14'-Methylenebis-18-hydroxytabersonine, V-173
- 2,3'-Methylenebisindole-3-carboxaldehyde, M-65
- 3,3'-Methylenebisindole, M-436
- Methylenebismehranine, M-437
- N*^b, *N*^b-Methylenebismeloninium(2+), M-438
- 10,10'-Methylenebis[norvallesamide], M-439
- 1,1'-Methylenebis[2-phenazinol], M-440
- 10,10'-Methylenebisvindoline, V-129
- 1-Methylenedasyrcarpidine, U-8
- 12,12'-Methylenedicytisine, M-441
- 3,3'-Methylenediindole, M-436
- 4-Methylene-2,5-dioxo-3-pyrrolidinepropanoic acid, I-239
- 1,2-Methylenedioxyaporphine, R-120
- 3,4-Methylenedioxybenzamide, M-442
- 3,4-Methylenedioxybenzoic acid, M-442
- 9,10-Methylenedioxycamptothecin, C-70
- 3,4-Methylenedioxy-cinnamic acid 3,4-dimethoxyphenethylamide, Z-6
- 3,4-Methylenedioxy-cinnamic acid 3,4-methylenedioxyphenylethylamide, D-799
- 3,4-Methylenedioxy-cinnamic acid 2-phenylethylamide, M-443
- 3,4-Methylenedioxy-cinnamic acid, M-457
- 3,4-Methylenedioxy-cinnamoyl piperidine, M-444
- N*-(3,4-Methylenedioxy-cinnamoyl)isopentylamine, M-457
- N*-(3,4-Methylenedioxy-cinnamoyl)pentylamine, M-457
- 1-[3,4-(Methylenedioxy)cinnamoyl]piperidine, M-444
- 1,2-Methylenedioxy-6 α ,7-dehydroaporphine-10,11-quinone, M-445
- 6,7-Methylenedioxydictamine, F-221
- 7,8-Methylenedioxydictamine, F-223
- 1,2-Methylenedioxy-8,9-dimethoxyoxoaporphine, O-170
- 3,4-(Methylenedioxy)-2,5-diphenyl-2,4-hexadienedioic acid, U-57
- 6,7-Methylenedioxyisocarbostyryl, I-305
- 6,7-Methylenedioxyisoquinoline, M-446
- 6,7-Methylenedioxy-1(2*H*)-isoquinolinone, I-305
- 11,12-Methylenedioxykopsaporine, K-66
- 11,12-Methylenedioxykopsalsaline *N*^a-oxide, K-74
- 11,12-Methylenedioxykopsalsaline, K-74
- 11,12-Methylenedioxykopsine, K-69
- 11,12-Methylenedioxykopsinol, K-66
- 5,6-Methylenedioxy-9-methoxy-7*H*-dibenzof[*de,h*]quinolin-7-one, B-117
- 3,4-Methylenedioxy-9-methoxy-10-nitro-1-phenanthrenecarboxylic acid, T-570
- 6,7-Methylenedioxy-1-(3,4-methylenedioxybenzyl)isoquinoline, M-672
- 6,7-Methylenedioxy-9-methylfuro[2,3-*b*]quinolin-9(5*H*)-one, F-221
- 6,7-Methylenedioxy-*N*-methylisoquinolinium(1+), M-446
- 3,4-Methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-447
- 1,2-Methylenedioxy-noraporphine, R-120
- 1,2-Methylenedioxyoxoaporphine, L-194
- 3,4-Methylenedioxy-*N*-pentylcinnamide, M-457
- 8,9-Methylenedioxyphenanthridine, M-448
- 8,9-Methylenedioxy-6-phenanthridone, C-744
- 3,4-Methylenedioxy-*N*-phenethylbenzamide, M-442
- N*-(3,4-Methylenedioxyphenethyl)cyclopropanecarboxamide, R-142
- 2-(3,4-Methylenedioxyphenethyl)quinoline, D-645
- 3-(3,4-Methylenedioxyphenyl)acrylic acid, M-457
- 2-[4-(3,4-Methylenedioxyphenyl)butyl]-4(1*H*)-quinolinone, M-449
- 10-(3,4-Methylenedioxyphenyl)-9-decenoic acid pyrrolidide, I-266
- 1-(3,4-Methylenedioxyphenyl)-6,7-dimethoxy-2-methyl-3,4-dihydroisoquinolinium(1+), C-796
- 12-(3,4-Methylenedioxyphenyl)-2,4,9,11-dodecatrienoic acid pyrrolidide, M-450
- 1-[12-(3,4-Methylenedioxyphenyl)-2,4,9,11-dodecatrienoyl]pyrrolidine, M-450
- 2-[2-(3,4-Methylenedioxyphenyl)ethyl]-4-methoxyquinoline, C-822
- 17-(3,4-Methylenedioxyphenyl)-16-heptadecenoic acid pyrrolidide, M-451
- 1-[17-(3,4-Methylenedioxyphenyl)-16-heptadecenoyl]pyrrolidine, M-451
- 7-(3,4-Methylenedioxyphenyl)-2,4-heptadienoic acid isobutylamide, C-387
- 7-(3,4-Methylenedioxyphenyl)-2,6-heptadienoic acid pyrrolidide, M-452
- 7-(3,4-Methylenedioxyphenyl)-2,4-heptadienoic acid pyrrolidide, N-214
- N*-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]isobutylamine, F-232
- 1-[7-(3,4-Methylenedioxyphenyl)-2,4-heptadienyl]piperidine, P-452
- 1-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]piperidine, P-452
- 1-[7-(3,4-Methylenedioxyphenyl)-2,6-heptadienyl]pyrrolidine, M-452
- 1-[7-(3,4-Methylenedioxyphenyl)-2,4-heptadienyl]pyrrolidine, N-214
- 7-(3,4-Methylenedioxyphenyl)-2,4,6-heptatrienoic acid pyrrolidide, P-453
- 1-[7-(3,4-Methylenedioxyphenyl)heptatrienyl]piperidine, P-452
- 7-(3,4-Methylenedioxyphenyl)-6-heptenoic acid pyrrolidide, M-452
- 1-[7-(3,4-Methylenedioxyphenyl)-2-heptenoyl]piperidine, P-465
- 1-[7-(3,4-Methylenedioxyphenyl)-6-heptenoyl]piperidine, P-465
- 1-[7-(3,4-Methylenedioxyphenyl)-6-heptenoyl]pyrrolidine, M-452
- 16-(3,4-Methylenedioxyphenyl)-2,4-hexadecadienoic acid isobutylamide, B-266
- 16-(3,4-Methylenedioxyphenyl)-2,4,15-hexadecatrienoic acid isobutylamide, B-266
- 2-[6-(3,4-Methylenedioxyphenyl)hexyl]-4(1*H*)-quinolinone, M-453
- 9-(3,4-Methylenedioxyphenyl)-2,8-nonadienoic acid piperidide, P-464
- 9-(3,4-Methylenedioxyphenyl)-2,8-nonadienoic acid pyrrolidide, M-455
- 9-(3,4-Methylenedioxyphenyl)-2,4-nonadienoic acid, M-454
- 1-[9-(3,4-Methylenedioxyphenyl)-2,8-nonadienyl]pyrrolidine, M-455
- 9-(3,4-Methylenedioxyphenyl)nonanoic acid, M-454
- 9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienoic acid piperidide, P-464
- 9-(3,4-Methylenedioxyphenyl)-4,6,8-nonatrienoic acid piperidide, P-464
- 9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienoic acid pyrrolidide, M-455
- 1-[9-(3,4-Methylenedioxyphenyl)-2,4,8-nonatrienyl]pyrrolidine, M-455
- 9-(3,4-Methylenedioxyphenyl)-8-nonoic acid piperidide, P-464
- 9-(3,4-Methylenedioxyphenyl)-8-nonoic acid pyrrolidide, M-455
- 1-[9-(3,4-Methylenedioxyphenyl)-8-nonenyl]pyrrolidine, M-455
- 1-[8-(3,4-Methylenedioxyphenyl)-7-octadecenyl]piperidine, P-470
- 15-(3,4-Methylenedioxyphenyl)-2,4-pentadecadienoic acid isobutylamide, B-265
- 15-(3,4-Methylenedioxyphenyl)-2,4,14-pentadecatrienoic acid isobutylamide, B-265
- 15-(3,4-Methylenedioxyphenyl)-14-pentadecenoic acid pyrrolidide, T-444
- 1-[15-(3,4-Methylenedioxyphenyl)-14-pentadecenoyl]pyrrolidine, T-444
- 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid isobutylamide, P-462
- 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid piperidide, P-467
- 5-(3,4-Methylenedioxyphenyl)-2,4-pentadienoic acid pyrrolidide, T-495
- 1-[5-(3,4-Methylenedioxyphenyl)-2,4-pentadienyl]piperidine, P-467
- 1-[5-(3,4-Methylenedioxyphenyl)pentanoyl]piperidine, P-467
- 5-(3,4-Methylenedioxyphenyl)-2-pentenoic acid piperidide, P-467
- 5-(3,4-Methylenedioxyphenyl)-2-pentenoic acid pyrrolidide, M-456
- 1-[5-(3,4-Methylenedioxyphenyl)-2-pentenyl]pyrrolidine, M-456
- 5-(3,4-Methylenedioxyphenyl)-2-phenylloxazole, D-650
- 3-(3,4-Methylenedioxyphenyl)-2-propenoic acid, M-457
- 5-(3,4-Methylenedioxyphenyl)-2-(3-pyridinyl)oxazole, T-299
- 2-(3,4-Methylenedioxyphenyl)quinoline, M-458
- 2-(3,4-Methylenedioxyphenyl)-4(1*H*)-quinolinone, G-171
- 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienoic acid isobutylamide, G-224
- 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienoic acid piperidide, P-448
- 13-(3,4-Methylenedioxyphenyl)-2,4,12-tridecatrienoic acid pyrrolidide, B-263
- 11-(3,4-Methylenedioxyphenyl)-2,10-undecadienoic acid isobutylamide, P-469
- 11-(3,4-Methylenedioxyphenyl)-2,10-undecadienoic acid piperidide, P-471
- 11-(3,4-Methylenedioxyphenyl)-2,8,10-undecatrienoic acid isobutylamide, P-469
- 11-(3,4-Methylenedioxyphenyl)-2,4,10-undecatrienoic acid piperidide, P-471
- 8,9-Methylenedioxy-11*H*-pyrrolo[3,2,1-*de*]phenanthridine, H-290
- 4,5-Methylenedodecanoic acid, H-131
- 3-Methylene-4-(1-methyl-1*H*-imidazol-5-yl)-1-phenyl-1-butanone, H-606
- 2-Methylene-4-oxopentanoic acid, M-459
- 14,10-(Methyleneoxy)dendroban-12-one, D-216
- 11-Methylenepachysamine H, M-460
- 11-Methylenepregnane-3,20-diamine, M-460
- 1-Methylenepyrrolizidine *N*-oxide, M-461
- 1-Methylenepyrrolizidine, M-461
- Methylephedrine, D-738
- N*-Methylephedrine, D-738
- Methyl-*v*-ephedrine, D-738
- N*-Methyl-16-epiaffinine, A-166
- N*-Methylepiamauromine, A-680
- N*-Methyl-7-epidioncophylline A, D-793
- 8-*O*-Methyl-1-epidioncophylline B, D-794
- 14-*O*-Methyl-6-epi-forensicine, M-323
- N*³-Methylepipachysamine D, P-601
- N*-Methyl-16-epipericlyclivine, T-425
- O*-Methyl-3-epischelhammericine B, T-51
- N*-Methylepivincadine, V-112
- O*-Methyl-16-epi- Δ^{14} -vincanol, E-6
- 12'-*O*-Methylergocornine, E-153
- 12'-*O*-Methyl- α -ergocryptine, E-157
- 6-Methyl-8-ergolene-8-carboxylic acid, P-107
- 1-Methylergotamine, E-172
- N*-Methyl-1,2-ethanediamine, E-225
- 2-(1-Methylethenyl)-5-(1-methylethyl)pyrazine, I-270
- 2-(1-Methylethenyl)-6-(1-methylethyl)pyrazine, I-271

- 6-(1-Methylethyl)-3,5-bis(2-methylpropyl)-2-piperazinone, P-445
 8-Methyl-10-ethylbelidiol, H-434
 5-(1-Methylethyl)-2-(2-methylpropyl)-1*H*-imidazole, I-280
 2-(1-Methylethyl)-5-(1-methylpropyl)pyrazine, I-281
 2-(1-Methylethyl)-5-(2-methylpropyl)pyrazine, I-282
 6-(1-Methylethyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, M-462
N-(1-Methylethyl)-*N'*-[7-(methylsulfinyl)heptyl]urea, D-827
N-(1-Methylethyl)-*N'*-[6-(methylsulfinyl)hexyl]urea, D-828
 3-(1-Methylethyl)-6-(phenylmethyl)-2(1*H*)-pyrazinone, B-88
 2-(1-Methylethyl)-4-propylthiazole, D-247
 (1-Methylethyl)pyrazine, I-285
 5-(1-Methylethyl)-2(1*H*)-pyrazinone, H-546
 6-(1-Methylethyl)-2(1*H*)-pyrazinone, H-547
 3-(1-Methylethyl)-2(1*H*)-pyrazinone, H-545
 3-[1-(1-Methylethyl)-2-pyrrolidinyl]pyridine, P-957
*N*¹⁴-Methyleudistomidin C, E-282
*N*²-Methyleudistomin D, B-321
*N*¹⁰-Methyleudistomin E, E-287
*N*²-Methyleudistomin J, B-322
N-Methyleuphoccocinine, M-389
 Methylevoxine, F-223
N-Methylexentricine, E-318
*O*⁷-Methylfaberone, H-366
*N*²-Methylfangchinoline, T-286
O-Methylfaurine, F-25
N-Methylferetine, A-151
N-(4-*O*-Methyl-*trans*-feruloyl)-4-*O*-methyl dopamine, H-565
 4-Methylfervenulone, D-498
 2-Methylfervenulone, D-498
O-Methylfetidine, T-333
O-Methylflavinantine *N*-oxide, F-80
O-Methylflavinantine, F-80
O-Methylflazine, F-85
N-Methylflindersine, F-90
N-Methylformamide, M-463
O-Methylfumarophycine, F-174
N-Methylfuntimine, A-882
 9-[(5-Methylfuryl)methyl]-β-carboline, V-166
 4-Methyl-10*H*-furo[3,2-*cj*]carbazole, M-464
 9-Methyl-6*H*-furo[2,3-*cj*]carbazole, M-465
 9-Methylfuro[2,3-*bj*]quinolin-4(9*H*)-one, F-224
O-Methylgalanthamine, G-3
N-Methylgeibalansine, G-29
 Methylgerambullal, G-67
 Methylgerambullin, G-68
 Methylgerambullone, G-68
 18-*O*-Methylgigactonine, M-309
N-Methylgilletine, C-552
O-Methylglaucamine, G-87
N-Methyl-D-glutamic acid, G-107
N-Methyl-L-glutamic acid, G-107
 3-*O*-Methylglycocitrine II, D-664
O-Methylglycophylone, T-583
O-Methylglycosolone, T-583
 1-Methylgranatan-3-one, M-389
N-Methylgranatone, A-1580
 9-*O*-Methylguacolidine, G-214
 9-*O*-Methylguadiscidine, G-188
N-Methylguanidine, M-466
 Methylguanidinoacetic acid, C-731
*N*²-Methylguanine, G-210
 7-Methylguanine, M-469
 1-Methylguanine, M-467
 3-Methylguanine, M-468
 9-Methylguanin-8(7*H*)-one, A-753
 9-*O*-Methylguattescidine, G-214
 9-*O*-Methylguattouregidine, G-215
 Methylgymnaconitine, M-352
 6-*O*-Methylhaemantidine, H-6
N-Methylhalfordinium, H-12
 Methylhalfordinol, H-12
 6-*O*-Methylhamateine, A-982
 6-*O*-Methylhamatinine, A-982
N-Methylhaplamine, F-90
N-Methylheliamine, T-158
*O*⁹-Methylheliotridine, T-188
 4'-*O*-Methylhemibastadin 1, H-100
 4'-*O*-Methylhemibastadin 2, H-100
 2-Methyl-6-(8-heptadecenyl)piperidine, M-470
O-Methylheptaphylline, H-125
 2-Methylheptyl isonicotinate, P-898
 1-Methylherbipoline, H-143
N-Methylhernagine, T-248
O-Methylhernandine, L-244
N-Methylhernangerine, N-25
N-Methylhernovine, T-248
 10-*O*-Methylhernovine, T-248
N-Methylheteropsine, H-160
 14-Methylhexadecanoic acid pyrrolidide, P-949
N-(14-Methylhexadecanoyl)pyrrolidine, P-949
N-(15-Methyl-9-hexadecenoyl)taurine, A-765
N-(5-Methylhexanoyl)pyrrothine, A-759
 8-(5-Methylhexyl)-1*H*,3*H*-[1,3,4]oxadiazolo[3,4-*a*][1,2,4]triazin-1-one, M-471
N-Methylhigenamine 7-*O*-β-D-glucopyranoside, H-275
N-Methylhigenamine *N*-oxide, H-275
N-Methylhigenamine, H-275
N-Methylhistamine, H-307
tele-Methylhistamine, H-307
N-Methylhistidyl-β-[[1-carboxy-2-(1*H*-imidazol-4-yl)ethyl]amino]-α-aminobutanoic acid, F-33
 Methylholaphylline, A-888
*N*²⁰-Methylholarrhimine, D-279
*N*³-Methylholarrhimine, D-279
 3-*O*-Methylhomobatrachotoxin, B-36
 4-Methylhomoidanomycin, I-64
 Methylhomotryamine, H-350
O-Methylhordenine, M-280
 10-*O*-Methylhostasine, H-362
N-Methylhuperzine B, H-376
 3-Methylhydantoin, I-40
N-Methylhydrasteine methyl ester, N-35
N-Methylhydrasteine, N-35
N-Methylhydrastine, M-472
 (3-Methyl-3-hydroxy-1-butenyl)paspalinine, P-110
 1-Methyl-2-(4-hydroxy-1,2-dithiolan-3-yl)pyrrolidine, M-547
*O*⁴-Methylhydroxyluninium(1+), I-290
*N*¹-Methyl-11-hydroxymacusine A, P-562
N-Methyl-*trans*-4-hydroxy-L-proline, H-715
*N*⁴-Methyl-10-hydroxysambarine, O-21
*N*⁴-Methyl-11-hydroxysambarine, O-21
 8-*O*-Methylhyapaconine, M-343
 1-Methylhypoxanthine, P-820
 7-Methylhypoxanthine, P-820
 9-Methylhypoxanthine, P-820
 3-Methylhypoxanthine, P-820
 3,3',3''-Methyldynetris-1*H*-indole, T-591
 3,3',3''-Methyldynetris[1-methyl-1*H*-indole], T-591
 Methylillukumbin A, I-30
 Methylillukumbin B, I-30
 4(5)-Methylimidazole, M-473
 3-Methyl-2,4-imidazolidinedione, I-40
 4-(1-Methyl-1*H*-imidazol-5-yl)-3-(phenylmethylene)-2-pyrrolidinone, I-189
 (1-Methyl-1*H*-imidazol-5-yl)-9*H*-pyrido[3,4-*bj*]indol-1-ylmethanone, X-23
 2-(1-Methyl-1*H*-imidazol-4-yl)quinoline, I-44
 2-(1-Methyl-1*H*-imidazol-5-yl)quinoline, I-44
 6-Methylimidazo[1,5-*cj*]tetrahydropyrimidine-5-thione, Z-8
 5-Methylimidazo[4,5-*ej*]-1,2-thiazin-4(5*H*)-one, N-81
 1-Methyl-6-iminopurine, A-842
 1-Methylindeno[1,2,3-*ij*][2,7]naphthyridinium, I-65
 4-Methyl-1*H*-indeno[1,2-*bj*]pyridine-2,5-dione, D-385
 4-Methyl-5*H*-indeno[1,2-*bj*]pyridin-5-one, O-93
N-Methylindsocin, I-74
 β-Methyl-1*H*-indole-3-butanolic acid, I-124
 β-Methyl-1*H*-indole-3-butanol, I-124
 1-Methyl-1*H*-indole-3-carbothioamide, I-85
 7-Methyl-1*H*-indole-3-carboxaldehyde, M-475
 1-Methyl-1*H*-indole-3-carboxylic acid, I-90
 3-Methyl-1*H*-indole-1-carboxylic acid, M-474
 3-Methyl-1*H*-indole-2-carboxylic acid, M-476
 5-Methyl-1*H*-indole-4,7-dione, M-477
N-Methyl-1*H*-indole-3-ethanamine, M-582
 3-Methyl-1*H*-indole, M-474
 β-Methylindole, M-474
 1-Methyl-1*H*-indole-3-methanamine, I-97
 2-Methyl-1*H*-indole-4-methanol, D-472
N-Methyl-1*H*-indole-3-propanamide, I-100
 α-Methyl-1*H*-indole-3-propanoic acid, I-131
 2-Methylindoline, D-472
 3-Methyl-3*H*-indolo[3,2,1-*de*][1,5]naphthyridine-5,6-dione, P-413
 6-Methyl-4*H*-indolo[3,2,1-*de*]-1,5-naphthyridin-4-one, N-302
 5-Methylindolo[2',3':3,4]pyrido[1,2-*bj*][2,7]naphthyridin-14(5*H*)one, M-131
 5-Methyl-5*H*-indolo[3,2-*bj*]quinoline-11-carboxylic acid, I-111
 5-Methyl-5*H*-indolo[3,2-*cj*]quinoline, I-110
 5-Methyl-5*H*-indolo[2,3-*bj*]quinoline, Q-19
 1-(3-Methyl-1*H*-indol-2-yl)ethanone, A-61
 2'-*O*-Methylinosine, I-148
O-Methylinsularoline, I-150
 6-*O*-Methylisepicoside, I-167
 7-*O*-Methylisepicoside, I-167
N-Methylirrigaine, I-177
O-Methylismine, I-186
 1'-*N*-Methylisoageliferin, A-190
N^b-Methylisoajmaline, A-216
 Methylisoalngiside, A-239
*O*⁹-Methylisoboldine, H-751
 4-Methylisoborverine, I-198
 Methylisochondendrine, C-835
O-Methylisococculine, I-207
N-Methylisocorydine, I-208
 β-*N*-Methylisocorypalminium, I-210
N-Methylisocrotosparinine, A-957
O-Methylisoburnamine, E-6
 Methylisophehdrine, D-738
 Methylisogerambullone, G-68
 Methylisopelletierine, P-173
O-Methylisophellibiline, P-315
N-Methylisopiline, T-521
O-Methylisopiline, T-521
 (+)-*N*-Methylisopiline, T-521
O-Methylisoplatydesmine, I-268
 1-Methyl-6,7-isoquinolinediol, D-627
 3-Methyl-1,6,8-isoquinolinetriol, T-566
N^b-Methylisoserpinilium, R-51
N-Methylisosaloline, T-181
N-Methylisochumannophytine, I-314
 14-Methylisotalitazidine, M-352
 2'-*N*-Methylisotetrandrine, I-328
O-Methylisothalicberine, I-329
O-Methylisothebaine, T-525
N-Methylisothebainium, T-525
 5-Methyl-3-isoxazolecarboxylic acid, M-479
 (±)-5-*O*-Methyljatropham, D-449
O-Methyljoubertamine, J-49
N-Methyljulifloridine, J-55
 8-*O*-Methylkarakoline, M-355
 8-*O*-Methylkarasamine, M-355
N-Methylkhalpofoline, H-56
 12'-*O*-Methylkohatine, K-56
N-Methylkorupensamine E, K-91
O-Methylkreysigine *N*-oxide, P-221
O-Methylkreysigine, P-221
 17-*O*-Methylkribine, K-101
 21-*O*-Methylkribine, K-101
 2-Methylactic acid, H-617
 Methylagerine, L-11
N-Methylambertine, L-23
 15-*O*-Methylatrunculin B, L-62
N-Methylaudanidinium, L-68
N-Methylauroformine, L-214
N-Methylaunobine, B-397
N-Methylaurelliptine, I-196
N-Methylaurotetanine *N*^α-oxide, L-75
N-Methylaurotetanine *N*^β-oxide, L-75
N-Methylaurotetanine, L-75
N-Methylleucine, L-153
N-Methylleuconolam, L-155
 21-*O*-Methyleuconolam, L-155

- O*-Methylleucotamine, G-3
 Methyliberine, L-165
 12'-*O*-Methylinsinine, N-87
O-Methylimacusine, L-175
N-Methylindcarpine, T-248
 7'-*O*-Methylindoldhamine, D-88
 7-*O*-Methylindoldhamine, D-88
 12-*O*-Methylindoldhamine, D-88
 2'-*N*-Methylindoldhamine, D-88
 2-*N*-Methylindoldhamine, D-88
O-Methylirinine, T-521
N-Methyliriendronine, L-193
N-Methylisericine, L-214
N-Methylolone, L-234
O-Methylongiberine, L-248
*N*¹-Methylumichrome, L-285
N-Methyl-β-lumicolchicine, L-287
O-Methyluninium, L-290
 Methyllycaonitine, G-91
 8-*O*-Methyllycaonitine, L-317
 18-*O*-Methyllycoctonine, M-309
 α-Methyllycodine-1-ethanol, H-703
N-Methyllycodine, L-324
O-Methyllycorenine *N*-oxide, L-347
O-Methyllycorenine, L-347
O-Methylmacralstonine, M-8
O-Methylmacromerine, M-18
O-Methylmacusine B, T-425
 Methylmadugin, M-582
 3-Methylmaleimide 5-oxime, M-542
 2-Methylmaleimide, M-542
*N*¹-Methylmanzacidin C, M-90
O-Methylmaritidine, M-108
 3-Methylmercaptopropylamine, M-578
N-Methylmerenderine, P-221
N-Methylmescaline, M-219
N-Methylmethanamine, D-726
*N*⁸-Methyl-10-methoxykuammidine, A-229
N-Methyl-4-methoxyphenethylamine, M-280
 17-*O*-Methyl-11-methoxy-Wieland-Gumlich aldehyde, C-115
N-Methyl-*N*-(2-methylaminobenzoyl)tryptamine, M-582
 17-Methyl-3-methylamino-D-homoandrostane-17,17a-diol, D-368
 4-Methyl-2-(methylamino)pentanoic acid, L-153
 6-Methyl-1-(3-methyl-2-butenyl)-9*H*-carbazole-2,7-diol, E-275
 3-Methyl-5-(3-methyl-2-butenyl)-9*H*-carbazole-2,6-diol, G-110
 6-Methyl-4-(3-methyl-2-butenyl)-9*H*-carbazol-3-ol, G-125
 3-Methyl-1-(3-methyl-2-butenyl)-9*H*-carbazol-2-ol, M-728
 6-Methyl-3-(3-methyl-2-butenyl)-9*H*-carbazol-2-ol, S-291
 3-Methyl-2-[6-(3-methyl-2-butenyl)-1*H*-indol-3-yl]-1,3-butanediol, M-480
 2-Methyl-4-[7-(3-methyl-2-butenyl)-1*H*-indol-3-yl]-2-buten-1-ol, B-202
 3-Methyl-6-[[7-(3-methyl-2-butenyl)-1*H*-indol-3-yl]methyl]-2,5-piperazinedione, T-86
 5-Methyl-9-(3-methyl-2-butenyl)-7-oxo-1(5*H*)-phenazinecarboxylic acid, E-79
 2-Methyl-*N*-(2-methylbutyl)butanamide, M-481
 3-Methyl-1-(2-methylbutyl)pyrrolidine, M-544
 3-Methyl-1-(3-methylbutyl)pyrrolidine, M-544
O-Methyl-*N*-(*N*-methylcarbamoyl)launobine, B-397
O-Methyl-*N*-(*N*-methylcarbamoyl)nandigerine, B-397
 4'-Methyl-*N*-methylcoclaurine, H-275
 2-Methyl-3-(19-methyl-12*Z*-eicosenylidene)-1-pyrrolone, L-37
 6-Methyl-2,3-methylenedioxy-carbazole, D-615
N-Methyl-2,3-methylenedioxyfuro[2,3-*c*]lacridone, C-342
 2-Methyl-6,7-methylenedioxy-1(2*H*)-isoquinoline, I-305
 5-Methyl-8,9-methylenedioxyphenanthridinium(1+), M-448
 1-Methyl-2-[6-(3,4-methylenedioxyphenyl)hexyl]-4(1*H*)-quinolinone, M-453
 1-Methyl-2-(3,4-methylenedioxyphenyl)-4(1*H*)-quinolinone, G-171
 4-Methyl-6-(1-methylethyl)-3-(phenylmethyl)-2,5-morpholinedione, B-29
 2-Methyl-5-(1-methylethyl)pyrazine, I-283
 2-Methyl-6-(1-methylethyl)pyrazine, I-284
 3-Methyl-1-(1-methylethyl)pyrrolidine, M-544
 4-Methyl-2-(1-methylethyl)thiazole, D-247
 2-Methyl-4-(1-methylethyl)thiazole, D-247
 2-Methyl-3-(16-methyl-12*Z*-heptadecenylidene)-1-pyrrolone, L-37
 2-Methyl-3-(16-methylheptadecylidene)-1-pyrrolone, L-37
 2-Methyl-1-(5-methylheptyl)-3*H*-carbazole-3,4(9*H*)-dione, C-129
 2-Methyl-1-(6-methylheptyl)-3*H*-carbazole-3,4(9*H*)-dione, C-129
 2-Methyl-3-(15-methyl-12*Z*-hexadecenylidene)-1-pyrrolone, L-37
 2-Methyl-3-(15-methylhexadecylidene)-1-pyrrolone, L-37
 2-Methyl-1-(5-methylhexyl)-3*H*-carbazole-3,4(9*H*)-dione, C-129
 3-Methyl-4-[(1-methyl-1*H*-imidazol-5-yl)methyl]-2(5*H*)-furanone, N-295
 5-Methyl-2-[(1-methyl-1*H*-indol-3-yl)methylene]-3,5-hexadienenitrile, A-1155
 2-Methyl-3-(18-methyl-12*Z*-nonadecenylidene)-1-pyrrolone, L-37
 2-Methyl-3-(17-methyl-12*Z*-octadecenylidene)-1-pyrrolone, L-37
 2-Methyl-3-(17-methyloctadecylidene)-1-pyrrolone, L-37
 2-Methyl-1-(7-methyloctyl)-3*H*-carbazole-3,4(9*H*)-dione, C-129
 6-Methyl-2-(2-methyl-1-oxobutyl)-1-oxa-5-azaspiro[2.5]oct-6-ene-4,8-dione, F-162
 6-Methyl-2-(3-methyl-1-oxobutyl)-1-oxa-5-azaspiro[2.5]oct-6-ene-4,8-dione, F-81
 3-Methyl-1-(3-methyl-2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one, H-467
 2-Methyl-3-(14-methylpentadecylidene)-1-pyrrolone, L-37
 2-Methyl-1-(3-methylpentyl)-3*H*-carbazole-3,4(9*H*)-dione, C-129
 6,6'-[3-Methyl-5-(2-methyl-1-propenyl)-2-pentene-1,5-diyl]bis-1*H*-indole, M-692
 2-Methyl-4-(2-methylpropyl)-2,4-azetinedicarboxylic acid, M-482
N-Methyl-*N*-(2-methylpropyl)-2,8-decadiene-4,6-diyamide, D-103
N-Methyl-*N*-(2-methylpropyl)-2-decene-4,6-diyamide, D-129
 4-[3-Methyl-4-(2-methylpropyl)-5-oxo-1-imidazolidinyl]-7-(2-methylpropyl)-3-phenyl-2-oxa-6,9-diazabicyclo[10.2.2]hexadeca-10,12,14,15-tetraene-5,8-dione, N-333
 2-Methyl-3-(2-methylpropyl)pyrazine, M-483
 1-Methyl-2-(1-methylpropyl)-4(1*H*)-quinolinone, M-531
N-Methyl-*N*-(2-methylpropyl)-2,4-tetradecadiene-8,10-diyamide, A-968
 2-Methyl-4-(3-methyl-1-pyrrolidinyl)-1-butanol, M-484
 1-Methyl-4-(1-methyl-2-pyrrolidinyl)-β-carbolone, B-292
 1-Methyl-α-[(1-methyl-2-pyrrolidinyl)methyl]-2-pyrrolidineethanol, D-413
 1-Methyl-4-(1-methyl-2-pyrrolidinyl)-9*H*-pyrido[3,4-*b*]indole, B-292
N-[15-Methyl-3-(13-methyltetradecanoyloxy)hexadecanoyl]glycine, M-485
N-[15-Methyl-3-(13-methyl-4-tetradecenoyloxy)hexadecanoyl]glycine, M-485
 2-Methyl-3-(13-methyltetradecylidene)-1-pyrrolone, L-37
 2-Methyl-6-(methylthio)-3,5,8(2*H*)-isoquinoline-trione, P-255
N-Methyl-*N*-[3-(methylthio)-1-oxo-2-propenyl]benzamide, P-192
N-Methyl-3-(methylthio)-2-propenamide, M-577
 3-Methyl-1-[3-(methylthio)propyl]pyrrolidine, M-486
O-Methylmicranthine, M-591
 1-Methylmitomycin A, M-653
 Methylmitomycin C, M-654
N'-Methylmolkaiamine, M-679
*N*⁷-Methylmonobromophakellin, B-343
 3'-*O*-Methylmontrifoline, F-221
N-Methylmoranoline, H-616
 Methylmorphine, C-555
 α-Methyl-4-morpholineethanol, M-705
N-Methylmorpholine, M-705
O-Methylmoschatoline, M-712
 9-*O*-Methylmukanadin A, D-863
O-Methylmukonal, H-448
O-Methylmukonidine, H-453
N-Methylmurexine, M-739
 2'-*O*-Methylmyxalamide D, M-805
 (6*E*)-2'-*O*-Methylmyxalamide D, M-805
 (6*E*,10*Z*)-2'-*O*-Methylmyxalamide D, M-805
N-Methylnandigerine β-*N*-oxide, N-25
N-Methylnandigerine, N-25
N-Methylnantene, N-29
 5-Methyl-2,7-naphthyridine-4-carboxylic acid, M-489
 1-Methyl-2,7-naphthyridine-4-carboxylic acid, M-488
 4-Methyl-2,6-naphthyridine, M-487
N-Methylnarcemicine (incorr.), A-143
N-Methylnarcemine, A-143
 (+)-*N*-Methyl-α-narcotine, N-44
O-Methylnareline, N-47
 3-Methylnarvenone A, P-908
 7-*O*-Methylnecaryachine *N*-methosalt, N-104
N-Methylnecaryachinium, N-104
 7-Methylnecoposide, I-167
N-Methylnicotinamide, P-895
N-Methylnicotinic betaine, C-149
 Methylnigikonol, D-546
N-Methylnitrarine, N-246
 4-[[[Methyl-*aci*-nitro]acetyl]amino]butanamide, A-1323
N-Methyl-*N*-nitrosoguanidine, M-466
N-Methyl-4-nitrotryptophan, N-257
 3'-*O*-Methylnorkolbicine, F-221
 2-Methyl-3-(12*Z*-nonadecenylidene)-1-pyrrolone, L-37
 2-Methyl-3-nonadecylidene-1-pyrrolone, L-37
 5-Methyl-10-(6,8-nonadienyl)lehmizidine, A-367
 3-Methyl-2-[(2-nonen-4,6,8-triynoyl)amino]butanoic acid, M-490
 5-Methyl-10-(8-nonenyl)lehmizidine, A-367
 3-Methyl-2-(2-nonenyl)-4-quinolinol, M-491
 3-Methyl-2-(2-nonenyl)-4(1*H*)-quinolinone, M-491
 8-Methyl-5-(6-nonen-8-ynyl)indolizidine, O-51
 1-Methyl-5-nonyl-2-(phenylmethyl)-3-pyrrolidinol, B-89
 2-Methyl-6-nonylpiperidine, M-492
 2-Methyl-6-nonyl-4-piperidinol, H-611
 1-Methyl-2-nonyl-4(1*H*)-quinolinone, H-642
 3-Methyl-2-nonyl-4(1*H*)-quinolinone, M-491
N-(8-Methylnonyl)sulfamic acid, M-493
 5-Methyl-10-(8-nonyl)lehmizidine, A-367
N-Methylnorapateline, A-1313
O-Methylnorbelladine, N-288
 8-Methylnorchelerythrine, M-494
 6'-*O*-Methyl-*N*-norcoculinine, C-552
N-Methyl-23-norconacona-3,5-dienin-3-amine, R-39
 4-Methyl-19-norconacona-1,3,5(10)-trienine, H-317
O-Methylnorlirine, T-521
 8-Methylnorlobelol, S-216
 8-Methylnorlobelone, P-173
O-Methylnormacromerine, M-18
O-Methylnormacusine B, T-425
 17-Methyl-18-nor-16,28-secosolanida-5,12-diene-3,16-diol, V-61
 17-Methyl-18-nor-16,28-secosolanida-5,12-dien-3-ol, V-60
 17-Methyl-18-norspirosola-5,12-dien-3-ol, V-66
 2'-*N*-Methylnorthalbrine, T-309
 4-Methylnorvaline, L-153
O-Methylnupharamine, N-340
 8-*O*-Methylolongine, L-251
N-Methylbovanine, P-781

- 2-Methyl-3-(12Z-octadecenylidene)-1-pyrroline, L-37
 2-Methyl-3-octadecylidene-1-pyrroline, L-37
 7-Methyl-2,4-octadienamide, M-495
 7-Methyl-2,4-octadienoic acid, M-495
 2-(6-Methyl-2,4-octadienyl)-1H-pyrrole, M-496
 N-(7-Methyloctyl)sulfamic acid, M-497
 O-Methyloduline, O-68
 O-Methylorantine, A-1334
 N-Methyloroline, O-109
 N-Methyloridine, O-109
 3'-O-Methylorixine, O-113
 1-Methylorotic acid, D-669
 Methylosin B, P-636
 N-Methylovigerine, O-143
 9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]nonan-7-ol, S-158
 9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl α -hydroxy- α -(hydroxymethyl)benzeneacetate, A-1043
 9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl α -(hydroxymethyl)benzeneacetate, S-159
 9-Methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl α -methylbenzeneacetate, A-1364
 16R-Methyloxazolomycin C, O-155
 16R-Methyloxazolomycin, O-155
 16S-Methyloxazolomycin, O-155
 2-(3-Methyloxiranyl)quinoline, P-647
 9-Methyl-8-oxoadenine, A-819
 2-Methyl- β -oxobenzenepropanamide, M-512
 10-Methyl-8-oxo-8H-benzo[*b*]pyrido[4,3,2-*de*][1,8]-phenanthroline, D-222
 3-(3-Methyl-1-oxo-2-butenyl)-1H-indole, I-125
 7-(3-Methyl-2-oxobutoxy)-8-methoxydictamine, F-223
 4-Methyl-6-(2-oxobutyl)-2-piperidinecarboxylic acid, T-493
 N-Methyl-2-(2-oxobutyl)piperidine, H-435
 N-(2-Methyl-3-oxodecanoyl)-1H-pyrrole, M-553
 N-(2-Methyl-3-oxodecanoyl)-2-pyrroline, D-473
 1-(2-Methyl-3-oxo-8-decenoyl)-1H-pyrrole, M-553
 1-Methyl-2-(9-oxodecyl)-4(1H)-quinolinone, M-498
 N³-Methyl-4-oxo-3-epilakinamine B, P-497
 1-[4-(4-Methyl-5-oxo-2(5H)-furanilydene)butyl]-2-pyrrolidinone, P-59
 O-Methyl-7-oxohernangerine, O-180
 4-[4-Methyl-2-(3-oxohexyl)phenyl]-3-butenamide, L-262
 9a-Methyl-3-(1-oxohexyl)-6-(1-propenyl)furo[3,2-g]isoquinoline-2,9(7H,9aH)-dione, R-152
 N-Methyloxohydrasteine, N-35
 5-Methyl-2-oxo-4-imidazolidinehexanoic acid, M-499
 7-Methyl-2-(7-oxo-1,3-nonadienyl)-3H-azepine, C-340
 2-Methyl-7-(7-oxo-1,3-nonadienyl)-2H-azepine, C-340
 7-Methyl-2-(7-oxo-1,3,5-nonatrienyl)-3H-azepine, C-340
 1-Methyl-2-(8-oxononyl)-4(1H)-quinolinone, H-642
 9a-Methyl-3-(1-oxooctyl)-6-(1-propenyl)furo[3,2-g]isoquinoline-2,9(7H,9aH)-dione, M-686
 8-(4-Methyl-2-oxopentyl)dihydrochelerythrine, M-500
 N-Methyl-2-(2-oxopentyl)piperidine, H-37
 2-Methyl-N-[1-(1-oxo-3-phenyl-2-propenyl)-2-pyrrolidinyl]butanamide, O-67
 2-Methyl-N-[1-(1-oxo-3-phenyl-2-propenyl)-2-pyrrolidinyl]propanamide, P-479
 3-Methyl-N-(3-oxopropyl)butanamide, A-892
 2-(1-Methyl-2-oxopropylidene)phosphorohydrazidothioate oxime, G-28
 3-Methyl-6-(1-oxopropyl)-2,4(1H,3H)-pteridine-dione, H-702
 N⁷-(1-Methyl-2-oxopropyl)streptonigrin, S-569
 3-Methyl-1-(2-oxo-5-pyrrolidinyl)-3-pyrrolin-2-one, D-448
 1-(3-Methyl-2-oxo-5-pyrrolinyl)-4-methyl-5-oxo-2-pyrrolidinecarboxylic acid, D-474
 18-O-[2-(2-Methyl-4-oxo-4H-quinazolin-3-yl)benzoyl]lycoctonine, M-501
 1-Methyl-2-(10-oxoundecyl)-4(1H)-quinolinone, H-757
 O-Methoxyacanthine, O-1
 N-Methylpachycerine, T-233
 N-Methylpachygonamine, Y-5
 N-Methylpachypodanthine N-oxide, O-87
 N-Methylpachypodanthine, O-87
 N-Methylpachysamine A, P-601
 1-O-Methylpakistanine, P-16
 N-Methylpalaudinone, P-80
 O-Methylpallidine N-oxide, F-80
 O-Methylpallidine, F-80
 O-Methylpallidine, P-31
 20-Methyl-17-O-palmitoylspirolole G, S-434
 12-O-Methylpampulhamine, D-88
 N-Methylpanocrine, P-55
 O²-Methylpancracine, P-57
 N-Methylpapaveraldinium, P-77
 N-Methylpapaverberbine, P-78
 N-Methylpapaverine, P-79
 N-Methylpapaverubine A, R-86
 N-Methylpapaverubine B, G-87
 O-Methylpapaverubine C, G-87
 N-Methylpapaverubine D, G-87
 O-Methylpapaverubine D, G-87
 N-Methylpapaverubine E, R-86
 N-Methylpapaverubine F, G-87
 N-Methylpapaverubine G, A-638
 O-Methylpapyrmine, P-86
 9-Methylparagraine, P-737
 N-Methylparavallarine, P-93
 N-Methylparavallarine, P-93
 17-Methylparsonsianidine, M-502
 N-Methylpavine, A-1410
 7-O-Methylpedroamine, D-88
 Methylpelletierine, P-173
 α -N-Methylpelletierin- β -one, P-173
 O-Methylpellotine, T-237
 O-Methylpendine, C-552
 16-Methylpendolmycin, M-503
 1-Methyl-2-(6,9-pentadecadienyl)-4(1H)-quinolinone, H-657
 14-Methyl-2-pentadecenethioamide S-oxide, S-633
 2-Methyl-6-(6-pentadecenyl)piperidine, M-504
 1-Methyl-2-(10-pentadecenyl)-4(1H)-quinolinone, H-657
 1-Methyl-2-(6-pentadecenyl)-4(1H)-quinolinone, H-657
 1-Methyl-2-(9-pentadecenyl)-4(1H)-quinolinone, H-657
 2-Methyl-3-pentadecylidene-1-pyrroline, L-37
 N-(14-Methylpentadecyl)-N-nitrosohydroxylamine, N-253
 2-Methyl-6-pentadecylpiperidine, M-504
 1-Methyl-2-pentadecyl-4(1H)-quinolinone, H-657
 4-Methylpentanamide, M-505
 4-Methylpentanoic acid, M-505
 3-Methyl-3-pentenoic acid, M-506
 8-Methyl-5-(4-pentenyl)indolizidine, O-52
 8-Methyl-5-(2-penten-4-ynyl)indolizidine, O-52
 1-Methyl-4-(2-penten-4-ynyl)indolizidine, M-507
 2-Methyl-1-pentyl-3H-carbazole-3,4(9H)-dione, C-129
 3-Methyl-2-(pentylidene)cyclopropyl-4(1H)-quinolinone, A-1179
 8-Methyl-5-pentylindolizidine, O-52
 N-(4-Methylpentyl)-2,12-octadecadienamide, O-30
 N-(4-Methylpentyl)-2,4,12-octadecatrienamide, O-33
 2-Methyl-3-pentylprodigine, P-636
 3-Methyl-2-pentyl-4-quinolinol, M-508
 1-Methyl-2-pentyl-4(1H)-quinolinone, H-660
 3-Methyl-2-pentyl-4(1H)-quinolinone, M-508
 8-Methyl-5-(4-pentenyl)indolizidine, O-52
 14-O-Methylperegrine, M-347
 N-Methylpericyclivine, T-425
 O-Methylpeyrovic acid, P-299
 O-Methylpeyoxylic acid, T-175
 O-Methylphellimine, S-126
 10-Methyl-2(10H)-phenazinone, P-328
 N-Methylphenethylamine, P-342
 2-Methyl-N-phenethylbutyramide, P-342
 3-Methyl-N-phenethylbutyramide, P-342
 3-Methyl-2-(4-phenyl-1,3-butadienyl)-4-piperidinol, D-386
 8-Methyl-10-phenyl-3,4-dehydrolobelidol, S-214
 N-Methyl-N-(2-phenylethenyl)-3-phenyl-2-oxirane-carboxamide, M-509
 2-Methyl-N-(2-phenylethyl)butanamide, P-342
 3-Methyl-N-(2-phenylethyl)butanamide, P-342
 N-Methyl-N-phenylethylcinnamamide, P-344
 2-Methyl-N-(2-phenylethyl)propanamide, P-342
 3-Methyl-1-(2-phenylethyl)pyrrolidine, M-544
 1-Methyl-3-(2-phenylethyl)-2,4(1H,3H)-quinazolinone, M-510
 13-Methyl-N-(2-phenylethyl)tetradecanamide, P-342
 3-Methyl-1-phenyl-1H-indole, M-474
 8-Methyl-10-phenyllobelidol, S-214
 8-Methyl-10-phenyllobelinol, S-215
 2-[[9-Methyl-6-[(phenylmethyl)amino]-9H-purin-2-yl]amino]ethanol, O-90
 8-Methyl-2-(phenylmethyl)-8-azabicyclo[3.2.1]octane-3,6-diol, B-94
 8-Methyl-4-(phenylmethyl)-8-azabicyclo[3.2.1]octane-3,6-diol, B-95
 8-Methyl-2-(phenylmethyl)-8-azabicyclo[3.2.1]octan-3-ol, B-96
 16-Methyl-10-phenylmethyl[13]cytochalasane-6,13,21-triene-1,23-dione, P-677
 1-Methyl-2-(phenylmethyl)-4(1H)-quinazolinone, B-93
 2-Methyl-6-(9-phenylnonyl)-3-piperidinol, I-177
 1-Methyl-2-(9-phenylnonyl)pyrrolidine, I-178
 4-Methyl-5-phenyl-2-oxazolindione, M-511
 N-Methyl-3-phenyloxirane-carboxamide, P-348
 3-(2-Methylphenyl)-3-oxopropanamide, M-512
 N-Methyl-3-phenyl-N-(2-phenylethenyl)-2-propanamide, L-38
 1-Methyl- α -phenyl-2-piperidineethanol, S-210
 2-[Methyl(3-phenylpropanoyl)amino]benzoic acid, M-384
 3-(2-Methylphenyl)-2-propanamide, M-513
 3-(2-Methylphenyl)-2-propenoic acid, M-513
 6-Methyl-9-phenyl-9H-purine, M-535
 3-Methyl-1-phenyl-1H-pyrrole-2,5-dione, M-542
 3-Methyl-4-phenyl-1H-pyrrole, M-514
 1-Methyl- α -phenyl-2-pyrrolidineethanol, P-970
 1-Methyl-2-phenyl-4(1H)-quinolinone, H-693
 4-Methyl-3-phenylspiro[3H-1,4-benzodiazepine-3,2'-oxirane]-2,5(1H,4H)-dione, C-889
 6'-O-Methylphlebicine, R-114
 N^a-Methylphlegmarine, P-361
 N^b-Methylphlegmarine, P-361
 N-Methylphoebine, P-369
 N-Methylphylline, T-177
 N-Methylpicolinic betaine, C-148
 N-Methylpicraline, B-420
 3,3'-(1-Methyl-2,5-piperazinediyl)bis[6-bromo-1H-indole], D-929
 N-Methylpiperidine oxide, M-515
 2-Methyl-3,4-piperidinediol, M-516
 2,2'-(1-Methyl-2,6-piperidinediyl)bis[1-phenylethanol], L-218
 α -Methyl-2-piperidineethanol, S-216
 1-Methylpiperidine, M-515
 2-Methyl-3,4,5-piperidinetriol, M-517
 1-Methyl-3,4,5-piperidinetriol, P-458
 5-(1-Methyl-2-piperidinyl)-2,3'-bipyridine, A-963
 1-(N-Methyl-2-piperidinyl)-2-butanol, H-435
 1-(N-Methyl-2-piperidinyl)-2-butanone, H-435
 4-(1-Methyl-2-piperidinyl)- β -carboline, M-518
 4-[2-(1-Methyl-2-piperidinyl)ethenyl]phenol, C-220
 1-(N-Methyl-2-piperidinyl)-2-pentanol, H-37
 1-(N-Methyl-2-piperidinyl)-2-pentanone, H-37
 2-(1-Methyl-2-piperidinyl)-1-phenylethanol, S-210
 2-(1-Methyl-2-piperidinyl)-1-phenylethane, S-210
 20-(5-Methyl-2-piperidinyl)-pregn-5-ene-3,16-diol, 13CI, S-200
 1-(6-Methyl-2-piperidinyl)-2-propanone, M-519
 4-(1-Methyl-2-piperidinyl)-9H-pyrido[3,4-*b*]indole, M-518
 Methylpiperidylamine, M-515
 N-Methylpiperolactam A, D-567

- N*-Methylpiperolactam D, T-537
O-Methylplatycerine, P-514
N-Methylpleiocarpamine, P-518
N-Methylporphyroxine, G-87
O-Methylporphyroxine, G-87
1-*O*-Methylporveniramine, P-16
O-Methylpratorimine, P-586
O-Methylpratorinine, P-586
O-Methylprechilenine, M-520
N-Methyl-7-prenyloxyflindersine, F-90
N-Methyl-8-prenyloxyflindersine, F-90
O-Methylprecoetine, P-208
N-Methylpreskimmianine, P-625
8-*O*-Methylpretazettine, P-626
N-Methylproline methylbetaine, S-491
5-Methylproline, M-545
2-Methyl-1-propanamine, M-525
N-(2-Methylpropanoyl)pyrrothine, A-759
1-(2-Methyl-1-propenyl)- β -carboline, M-521
N-Methyl-11-(2-propenyl)cytisine, T-412
2-Methyl-6-(1-propenyl)piperidine, M-522
2-Methyl-5-(1-propenyl)pyrazine, M-523
2-Methyl-6-(1-propenyl)pyrazine, M-524
1-(2-Methyl-1-propenyl)-9*H*-pyrido[3,4-*b*]indole, M-521
N-(2-Methylpropenyl)-7-(2-thienyl)-2,4,6-heptatrienamamide, O-133
3-Methyl-6-propionylumazine, H-702
N-(2-Methylpropionyl)-3-nitrotyramine, H-640
2-Methyl-1-propylamine, M-525
4-[(2-Methylpropyl)amino]-4-oxo-2-butenic acid, F-173
2-[(2-Methylpropyl)-*ONN*-azoxy]-2-propenoic acid, V-5
3-Methyl-*N*-propylbutylamine, M-420
N-(2-Methylpropyl)-2,4-decadienamamide, D-106
N-(2-Methylpropyl)-6,8-decadienamamide, D-105
N-(2-Methylpropyl)-2,8-decadiene-4,6-diynamide, D-103
2-Methyl-5-propyldecahydroquinoline, D-112
N-(2-Methylpropyl)decanamide, D-115
N-(2-Methylpropyl)-2,4,8-decatrienamamide, D-123
N-(2-Methylpropyl)-2,6,8-decatrienamamide, D-124
N-(2-Methylpropyl)-2-decenamide, D-132
N-(2-Methylpropyl)-2-decene-6,8-diynamide, D-130
N-(2-Methylpropyl)-2,4-dodecadiene-8,10-diynamide, D-882
N-(2-Methylpropyl)-2-dodecene-8,10-diynamide, D-897
N-(2-Methylpropyl)-2,4-eicosadienamamide, E-45
N-(2-Methylpropyl)hexacosanamamide, H-214
N-(2-Methylpropyl)-2,9-hexadecadiene-12,14-diynamide, H-215
N-(2-Methylpropyl)-2,6,8,12-hexadecatetraenamamide, H-223
4-(2-Methylpropyl)morpholine, M-705
N-(2-Methylpropyl)-2,4-nonadien-8-ynamide, N-280
N-(2-Methylpropyl)-2-nonene-6,8-diynamide, N-282
N-(2-Methylpropyl)-2,4-octadienamamide, O-39
N-(2-Methylpropyl)-3-(3-oxo-1-cyclopenten-1-yl)propanamide, M-526
N-(2-Methylpropyl)-2,9-pentadecadiene-12,14-diynamide, P-202
2-(1-Methylpropyl)-5-phenyl-1*H*-imidazole, M-527
2-(2-Methylpropyl)-5-phenyl-1*H*-imidazole, M-528
2-Methyl-6-propylpiperidine, M-529
3-(2-Methylpropyl)piperidine, M-530
2-Methyl-6-propyl-4-piperidinol, H-622
2-Methyl-3-propylprodimine, P-636
2-Methyl-*N*-propyl-1-propylamine, M-525
2-(1-Methylpropyl)-4-quinolinol, M-531
1-Methyl-2-propyl-4(1*H*)-quinolinone, H-707
2-(1-Methylpropyl)-4(1*H*)-quinolinone, M-531
4-Methyl-6-propylquinolizidine, M-532
N-(2-Methylpropyl)-2,4-tetradecadiene-8,10-diynamide, A-968
N-(2-Methylpropyl)-10,12-tetradecadiene-4,6-diynamide, T-122
N-(2-Methylpropyl)-2,4,10,12-tetradecatetraen-8-ynamide, T-130
N-(2-Methylpropyl)-2,4,8-tetradecatrienamamide, T-133
N-(2-Methylpropyl)-2,4,10-tetradecatrien-8-ynamide, T-134
2-(2-Methylpropyl)thiazole, M-533
2-Methyl-1-propylthiazolium(1+), M-400
N-(2-Methylpropyl)-2,4-tridecadienamamide, T-502
N-(2-Methylpropyl)-2,7-tridecadiene-10,12-diynamide, T-501
N-(2-Methylpropyl)-2,6,8-tridecatriene-10,12-diynamide, T-504
N-(2-Methylpropyl)-2,4-undecadienamamide, U-28
N-(2-Methylpropyl)-2,4-undecadiene-8,10-diynamide, U-26
N-(2-Methylpropyl)-2,7,9-undecatrienamamide, U-29
N-(2-Methylpropyl)-2-undecene-8,10-diynamide, U-30
N-Methyl-*sec*-pseudobrucine, I-12
N-Methyl-*sec*-pseudo- β -colubrine, I-12
N-Methylpseudoconhydrine, P-653
13-Methylpseudooxoptisine, W-22
Methylpseudooecgonine, E-8
Methylpseudoephedrine, D-738
N-Methylpseudolaudanine, P-715
Methylpseudolycorine, P-716
8-Methylpseudomanibacanine, M-534
*N*⁶-Methylpseudophrynaminol, P-720
N-Methyl-*sec*-pseudostyrychine, I-12
5-Methylpseudoyohimbane, Y-12
11-*O*-Methylpsurotin A, P-738
*N*⁶-Methylpsilocybine, H-480
O-Methylpsychochine, E-72
O-Methylptelefolidine, P-759
O-Methylptelefoline, P-755
*O*⁴-Methylptelefolonium, P-757
1-Methyl-2,4-pteridinedione, P-762
8-Methyl-2,4,7(1*H*,3*H*,8*H*)-pteridinetriene, P-763
O-Methylpukateine, P-781
N-Methylpulchine, T-524
O-Methylpunjabin, P-800
1-Methyl-1*H*-purin-6-amine, A-842
2-Methyl-1*H*-purin-6-amine, A-843
3-Methyl-3*H*-purin-6-amine, A-844
6-Methylpurine, M-535
2-Methyl-4(1*H*-purin-6-ylamino)-1-butanol, Z-11
2-Methyl-4(1*H*-purin-6-ylamino)-2-buten-1-ol, Z-11
2-Methyl-4(1*H*-purin-6-ylamino)-2-penten-1-ol, M-586
N-Methylpurpureine, P-208
4-Methylpyochelin, T-373
3-Methyl-2,5-pyrazinediol, D-631
Methylpyrazine, M-536
3-Methylpyrazinol, H-624
3-Methyl-2(1*H*)-pyrazinone, H-624
6-Methyl-2-pyrazinyl-2-furfuryl sulfide, F-201
1-(5-Methylpyrazinyl)-1,2,3,4-butanetetrol, P-154
1-(6-Methylpyrazinyl)-1,2,3,4-butanetetrol, P-155
4-(3-Methylpyrazinyl)-1,2,3-butanetriol, P-153
1-(3-Methylpyrazinyl)ethanone, A-62
1-Methyl-1*H*-pyrazole-3-carboxylic acid, P-890
1-Methyl-1*H*-pyrazole-5-carboxylic acid, P-890
4-Methyl-2-pyridinamine, A-845
N-Methyl-3-pyridinecarboxamide, P-895
2-Methyl-3,4-pyridinediol, D-632
 α -Methyl-3-pyridinedodecanal *O*-methyloxime, I-24
 β -Methyl-3-pyridinedodecanal *O*-methyloxime, I-25
 β -Methyl-3-pyridinedodecanamine, N-225
 γ -Methyl-3-pyridinedodecanamine, N-226
2-Methylpyridine, M-537
 α -Methyl-3-pyridinetriecanal *O*-methyloxime, N-229
6-Methyl-2,3,5-pyridinetriol, T-567
2-Methyl-4-pyridinol, H-625
2-Methyl-4(1*H*)-pyridinone, H-625
N-Methyl-4-(3-pyridinyl)-3-buten-1-amine, M-232
1-(5-Methyl-3-pyridinyl)-1-decanone, M-538
2-Methyl-12-(3-pyridinyl)dodecylamine, N-225
3-Methyl-12-(3-pyridinyl)dodecylamine, N-226
1-(2-Methyl-3-pyridinyl)ethanone, A-65
1-(2-Methyl-4-pyridinyl)ethanone, A-67
1-(3-Methyl-4-pyridinyl)ethanone, A-68
1-(4-Methyl-2-pyridinyl)ethanone, A-63
1-(4-Methyl-3-pyridinyl)ethanone, A-66
1-(6-Methyl-2-pyridinyl)ethanone, A-64
1-(6-Methyl-3-pyridinyl)ethanone, A-69
1-Methyl-5-(3-pyridinyl)-2-pyrrolidinone, C-708
11-Methyl-6*H*-pyrido[4,3-*b*]carbazole-5-carboxaldehyde, E-67
1-Methyl-9*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, M-427
1-Methyl-9*H*-pyrido[3,4-*b*]indole-6,7-diol, D-620
1-Methyl-9*H*-pyrido[3,4-*b*]indole-7,8-diol, D-621
1-Methyl-9*H*-pyrido[3,4-*b*]indole, M-426
 α -Methyl-9*H*-pyrido[3,4-*b*]indole-1-methanol, H-486
2-Methyl-9*H*-pyrido[3,4-*b*]indolium, C-130
1-Methyl-9*H*-pyrido[3,4-*b*]indol-6-ol, H-591
1-Methyl-9*H*-pyrido[3,4-*b*]indol-7-ol, H-592
1-Methyl-9*H*-pyrido[3,4-*b*]indol-8-ol, H-593
5-Methyl-11-(9*H*-pyrido[3,4-*b*]indol-9-yl)-5*H*-quinoline, C-786
8-Methylpyridoxatin, C-646
1-Methyl-2-(3-pyridyl)pyrrolidine, N-190
2-Methyl-5-pyrimidinecarboxylic acid, M-539
5-Methyl-2,4-pyrimidinediol, T-401
5-Methyl-2,4(1*H*,3*H*)-pyrimidinedione, T-401
5-Methyl-2,4,6(1*H*,3*H*,5*H*)-pyrimidinetriene, M-540
6-Methylpyrimido[5,4-*e*]-1,2,4-triazine-5,7(1*H*,6*H*)-dione, R-66
1-Methyl-2-pyrrolicarboxaldehyde, P-942
4-Methyl-1*H*-pyrrole-2-carboxylic acid, M-541
1-Methyl-1*H*-pyrrole-2-carboxylic acid, P-944
3-Methyl-1*H*-pyrrole-2,5-dione, M-542
1-Methyl-2-pyrrolidineaetic acid, P-950
5-Methyl-2-pyrrolidineaetic acid, M-545
 α -Methyl-2-pyrrolidinedecanamine, P-958
3,3'-(1-Methyl-2,5-pyrrolidinediyl)bis-1,2-dithiolan-4-ol, G-72
1,1'-(1-Methyl-2,5-pyrrolidinediyl)bis-2-propanone, H-704
 α -Methyl-2-pyrrolidinedodecanamine, A-921
1-Methylpyrrolidine, M-543
3-Methylpyrrolidine, M-544
1-Methyl-2-pyrrolidinone, M-546
3-(1-Methyl-2-pyrrolidinyl)-1,2-dithiolan-4-ol, M-547
3-Methyl-5-(2-pyrrolidinyl)-2(5*H*)-furanone, P-63
2',3-(*N*-Methylpyrrolidinyl)hygrine, M-548
2',4-(*N*-Methylpyrrolidinyl)hygrine, M-549
1-(1-Methyl-2-pyrrolidinylidene)-2-propanone, H-759
2-[3-(1-Methyl-2-pyrrolidinyl)-2-oxopropyl]-1-(1-oxo-3-phenyl-2-propenyl)pyrrolidine, D-211
2-(1-Methyl-2-pyrrolidinyl)-1-phenylethanol, P-970
1-(1-Methyl-2-pyrrolidinyl)-2-propanol, H-760
1-(1-Methyl-2-pyrrolidinyl)-2-propanone, H-759
3-(1-Methyl-2-pyrrolidinyl)pyridine, N-190
N-Methylpyrrolidone, M-546
4-Methyl-1-pyrroline-5-carboxylic acid, D-477
 β -Methylpyrroline, D-476
20-(2-Methyl-1-pyrroline-5-yl)pregn-4-en-3-one, M-550
7-Methyl-1-pyrrolizidinol, H-630
3-Methyl-1*H*-pyrrolo[3,2-*b*]pyridine, M-551
4-Methyl-3*H*-pyrrolo[2,3-*c*]quinoline, M-552
4-Methyl-1*H*-pyrrolo[3,4-*b*]quinoline-1,3,9(2*H*,4*H*)-trione, H-728
2-Methyl-1-(1*H*-pyrrol-1-yl)-1,3-decanedione, M-553
2-Methyl-1-(1*H*-pyrrol-1-yl)-8-decene-1,3-dione, M-553
1-(4-Methyl-1*H*-pyrrol-3-yl)ethanone, A-71
3-(1-Methyl-1*H*-pyrrol-2-yl)pyridine, P-972
N-Methylquebrachamine, Q-6
N-Methylquinaldic acid, Q-31
4-Methyl-2-quinazolinecarboxamide, M-555
4-Methyl-2-quinazolinecarboxylic acid, M-555
5-Methyl-2,4-quinazolinediol, D-633
5-Methyl-2,4(1*H*,3*H*)-quinazolinedione, D-633
1-Methyl-2,4(1*H*,3*H*)-quinazolinedione, Q-13
3-Methyl-2,4(1*H*,3*H*)-quinazolinedione, Q-13

- 4-Methylquinazoline, M-554
 4-Methyl-2-quinazolinemethanol, H-609
 2-Methyl-4-quinazolinol, H-632
 2-Methyl-4(1*H*)-quinazolinone, H-632
 1-Methyl-4(1*H*)-quinazolinone, M-556
 3-Methyl-4(3*H*)-quinazolinone, Q-14
 5-Methyl-5*H*-quindoline, I-109
 5-Methyl-11-(10*H*-quindolin-10-yl)-5*H*-quindoline, C-794
N-Methylquinicidinol, Q-17
 2-Methylquinoline, M-557
 2-Methyl-4-quinolinol, H-633
 2-Methyl-4(1*H*)-quinolinone, H-633
 1-Methyl-4(1*H*)-quinolinone, M-558
 1-Methyl-2(1*H*)-quinolinone, Q-50
 1-Methyl-2-(2-quinolinyl)oxirane, P-647
 2-Methyl-4(1*H*)-quinolone, H-633
 4-Methylraucubainium(1+), C-191
^{N^b}-Methylraumacline, R-22
O-Methylrepandine, R-46
 4-Methylreserpinine, R-51
^{O^b}-Methylretronecine, T-188
 1-Methylreumycin, R-66
O-Methylrhoegenine, R-86
 24-Methylritterazine D, R-104
O-Methylrobustimine, S-126
O-Methylroefractine *N*-oxide, H-275
O-Methylroehybridine, R-116
O-Methylroemeridine, R-116
N-Methylroeraminium, R-120
 8-*O*-Methylsachaconitine, M-355
O-Methylsakambullin, D-12
O-Methylsakerinol A, G-68
N-Methylsalsolidine, T-167
O-Methylsalsoline, T-167
 1-Methylsarpagan-3,17-diol, A-27
^{N^a}-Methylsarpagine methosalt, S-89
N-Methylsarpagine, S-89
O-Methylsattazolin, S-101
^{N^b}-Methylsaxitoxin, S-109
O-Methylschelhammericine B, T-51
N-Methylschinifoline, G-118
^{N^b}-Methylscholaricine, E-30
N-Methylschumagnine, S-144
N-Methylschumannifoline, S-145
N-Methylschumanniphytine, S-146
O-Methylsciadenine, S-148
 8-*O*-Methylsclerotioramine, S-155
N-Methylsecoglaucine, M-559
 4-Methyl-4,21-secosandwicine, S-52
 28-Methyl-16,28-secosolanid-5-ene-3,16-diol, S-200
 28-Methyl-16,28-secosolanid-5-ene-3,16,25-triol, P-433
 19-Methyl-16,19-secostychnidine-10,16-dione, I-12
^{N^b}-Methyl-^{N^b}-21-secotalpinine, T-20
O-Methyl-*N*-seneciolytyramine, M-280
^{N^b}-Methylserotonin, H-755
N-Methylseverifoline, S-278
N-Methylsinactine, S-307
 Methylsinharine, S-310
N-Methylsinococuline, S-311
O-Methylsolanocapsine, S-346
O-Methylsomniferine, S-362
N-Methylsopholupisine, S-375
N-Methylsarsiflorine, T-524
O-Methylspeciosine, S-387
^{N^b}-Methylspermidine, S-398
 6-Methylspinacemine, T-192
 1'-Methylspiro[isobenzofuran-1(3*H*),2'-pyrrolidin]-3-one, S-288
 1'-Methylspiro[1*H*-isindole-1,2'-pyrrolidin]-3(2*H*)-one, S-287
 20-Methylspirovide G, S-434
N-Methylstaurosoprine, S-498
 9-*O*-Methylstecepharine, C-93
 Methylstemofoline, M-560
 1'-*O*-Methylstemokerrin *N*-oxide, S-516
N-Methylstenantherine, T-245
O-Methylstepharinosine, S-542
 4'-*O*-Methylstephanine, S-545
N-Methylstephuline, L-244
 9-Methylstreptimidone, M-561
^{N^b}-Methylstrictamine, S-576
 21-*O*-Methylstrictosamide aglycone, V-126
^{N^a}-Methylstrychninium, S-589
N-Methylstylopine, S-612
N-Methylstylopinium, S-612
N-Methyl-*N*-*cis*-styrylcinnamide, L-38
N-Methyl-*N*-styrylcinnamide, L-38
N-(4-Methylsuccinimidobutyl)-*p*-coumaramide, H-682
 Methylsuccirubine, M-562
 7-(Methylsulfonyl)heptanenitrile, D-830
N-[6-(Methylsulfonyl)hexyl]-*N*'-[6-(methylthio)hexyl]urea, D-829
 [6-(Methylsulfonyl)hexyl]urea, D-828
 5-(Methylsulfonyl)pentanoic acid, M-203
 3-(Methylsulfonyl)-2-propenoic acid, M-577
 3-(Methylsulfonyl)propylamine, M-578
 2-[2-(Methylsulfonyl)-*L*-tryptophan]-3-(4,5-dihydroxy-*L*-leucine)-4-*L*-alanineviroisin, V-160
 2-[2-(Methylsulfonyl)-*L*-tryptophan]-3-(4,5-dihydroxy-*L*-leucine)viroisin, V-160
 2-[2-(Methylsulfonyl)-*L*-tryptophan]viroisin, V-160
 [11-(Methylsulfonyl)undecyl]urea, D-826
 Methylsulfomycin I, S-638
 10-(Methylsulfonyl)decanamide, M-563
 10-(Methylsulfonyl)decanenitrile, M-563
 10-(Methylsulfonyl)decanoic acid, M-563
 9-(Methylsulfonyl)nonanamine, M-564
 9-(Methylsulfonyl)nonylamine, M-564
 5-(Methylsulfonyl)pentanoic acid, M-203
 3-(Methylsulfonyl)-2-propenoic acid, M-577
 3-(Methylsulfonyl)propylamine, M-578
N-Methylswietenidine B, Q-45
 4-*O*-Methylsynephrine, S-659
 β-*O*-Methylsynephrine, S-659
 8-*O*-Methyltalatzimine, M-352
O-Methyltaxodine, T-51
N-Methyltaxol C, T-52
 6-*O*-Methyltazettinol, T-53
^{N^b}-Methyltchibangensine, U-52
O-Methylteleocidin A₁, L-367
N-Methyltelobine, A-1331
O-Methyltembamide, T-67
 4-Methyl-1,2,6,8-tetraazacycloundeca-4,9-diene-3,7,11-trione, M-565
 12-Methyl-1-tetradecanamine, M-566
 3-Methyl-4-tetradecanoyloxy-2-butenenitrile, H-576
 12-Methyl-1-tetradecylamine, M-566
 1-Methyl-2-tetradecylpyrrolidine, T-135
 1-Methyl-2-tetradecyl-4(1*H*)-quinolinone, H-746
O-Methyltetrahydrodriptriphophylline, D-793
^{N^b}-Methyltetrahydroalstonine, A-212
N-Methyltetrahydrocolumbaminium, I-210
N-Methyltetrahydrocytisine, T-149
N-Methyl-*N*-[1,2,3,4-tetrahydro-1,3-dimethyl-6-(methylamino)-2,4-dioxo-5-pyrimidinyl]formamide, H-761
N-Methyltetrahydroellipticine, E-67
 5-Methyltetrahydrofolic acid, T-172
 (2-Methyltetrahydrofuran)solanidine, M-567
^{N^b}-Methyltetrahydroharmol, H-592
 α-Methyl-1-(2,3,4,9-tetrahydro-2-methyl-1*H*-carbazol-1-yl)-1*H*-indole-3-butanolic acid, M-568
 3-Methyl-*N*-(5*α*,6*α*,7,8-tetrahydro-4-methyl-1,5-dioxo-1*H*,5*H*-pyrrolo[1,2-*c*][1,3]oxazepin-3-yl)butanamide, C-852
N-Methyltetrahydropapaverine, L-69
 2-Methyltetrahydrothiazole, M-570
^{N^b}-Methyltetrahydrovariolin B, V-32
 2-Methyl-5-(1,2,3,4-tetrahydroxybutyl)pyrazine, P-154
 2-Methyl-6-(1,2,3,4-tetrahydroxybutyl)pyridine, P-155
 2'-*N*-Methyltetrandrine, T-286
 2-*N*-Methyltetrandrine, T-286
N-Methylthaicanine, T-301
O-Methylthaicanine, T-301
 12-*O*-Methylthalfotidine, T-320
O-Methylthalibrine, T-309
O-Methylthalibrunimine, T-310
O-Methylthalicberine, T-313
N-Methylthalicsimidine, P-208
N-Methylthalidaldine, I-304
^{O³}-Methylthalifarapine, T-322
 12'-*O*-Methylthaligossinine, T-346
O-Methylthalirabine, T-337
O-Methylthalisopavine, T-336
 5-*O*-Methylthalisopine, T-346
 2'-*N*-Methylthalistyline, T-337
O-Methylthalmethine, T-339
O-Methylthallmine, T-341
O-Methylthalsimidine, T-347
N-Methylthebaine, T-350
 Methyltheobromine, C-15
 2-*O*-Methyltheobromine, T-355
^{N^a}-Methylthermospermine, D-249
 3-Methyl-1,3-thiazino[6,5-*b*]indole-2,4(3*H*,9*H*)-dione, R-170
 4-Methyl-5-thiazoleethanol, H-495
 4-Methylthiazole, M-569
 2-Methylthiazolidine, M-570
 2-Methyl-2-thiazoline, D-479
 2-(4-Methyl-5-thiazolyl)ethanol, H-495
 1-Methyl-3-(2-thiazolyl)-1*H*-indole, T-372
 3-(Methylthio)acrylamide, M-577
 3-(Methylthio)acrylic acid, M-577
 4-(Methylthio)butyl cyanide, M-203
 4-(Methylthio)canthin-6-one, M-571
 5-(Methylthio)canthin-6-one, M-572
 9-(Methylthio)canthin-6-one, M-573
 11-(Methylthio)canthin-6-one, M-574
 7-(Methylthio)heptanenitrile, D-830
 [6-(Methylthio)hexyl]urea, D-828
 2-(Methylthio)-1*H*-indole-3-carboxaldehyde, M-201
 4-Methylthio-6*H*-indolo[3.2.1-*de*][1,5]naphthyridin-6-one, M-571
 5-Methylthio-6*H*-indolo[3.2.1-*de*][1,5]naphthyridin-6-one, M-572
 9-Methylthio-6*H*-indolo[3.2.1-*de*][1,5]naphthyridin-6-one, M-573
 11-Methylthio-6*H*-indolo[3.2.1-*de*][1,5]naphthyridin-6-one, M-574
 2-(Methylthio)-3*H*-indol-3-one, M-575
 3-[(Methylthio)methylene]-2-pyrrolidinethione, R-18
 5-(Methylthio)pentanenitrile, M-203
 5-(Methylthio)pentanoic acid, M-203
 2-(Methylthio)-3*H*-phenoxazin-3-one, M-576
 3-(Methylthio)-*N*-(2-phenylethenyl)-2-propenamide, I-30
 3-(Methylthio)propanamine, M-578
 3-(Methylthio)-2-propenamide, M-577
 3-(Methylthio)-2-propenoic acid, M-577
N-[3-(Methylthio)-2-propenoyl]-*N'*-(phenylacetyl)putrescine, L-114
 3-(Methylthio)propylamine, M-578
 2'-(Methylthio)spiro[2*H*-indole-2,5'(4'*H*)-thiazol]-3(1*H*)-one, E-183
 2'-(Methylthio)spiro[3*H*-indole-3,5'(4'*H*)-thiazol]-2(1*H*)-one, S-426
 4-[2-(Methylthio)-5-thiazolyl]phenol, T-508
 5-(Methylthio)varacin A, V-20
 5-(Methylthio)varacin, L-205
 13-Methylthuggacin A, T-397
N-Methyltiliamosine, Y-5
 Methyltoldaliamide, T-421
N-Methyltrenudone, T-449
 5-Methyl-1,3,7-triazacyclodeca-4,9-diene-2,6,8-trione, R-70
 5-Methyl-1,3,7-triazecine-2,6,8(1*H*,3*H*,7*H*)-trione, R-70
N-Methyl-2-(2,3,4-tribromo-5,6-dimethoxyphenyl)methylamine, C-628
 12-*O*-Methyltricornadine, I-339
 1-Methyl-2-(4,7-tridecadienyl)-4(1*H*)-quinolinone, H-750
 12-Methyl-1-tridecanamine, M-579
 2-Methyl-6-(4-tridecenyloxy)piperidine, M-580
 1-Methyl-2-(7-tridecenyloxy)-4(1*H*)-quinolinone, H-750
 1-Methyl-2-(8-tridecenyloxy)-4(1*H*)-quinolinone, H-750
 12-Methyl-1-tridecylamine, M-579
N-(12-Methyltridecyl)-*N*-nitrosohydroxylamine, N-255

- 2-Methyl-6-tridecylpiperidine, M-580
 1-Methyl-2-tridecyl-4(1*H*)-quinolinone, H-750
 2-Methyl-3-(2,3,4-trihydroxybutyl)pyrazine, P-153
 3-Methyl-5-[3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4-oxo-4*H*-1-benzopyran-8-yl]-2-pyrrolidinone, L-173
N-Methyl-3,4,5-trihydroxyphenethylamine, T-576
N-Methyl-5,6,7-trimethoxyisoquinolinium(1+), I-307
 2-Methyl-4-(3,6,7-trimethoxy-2-naphthalenyl)-2-azabicyclo[2.2.1]heptan-3-one, M-581
 4-Methyl-5-(3,7,11-trimethyl-2,6,10-dodecatrienyl)-6*H*-imidazo[4,5,1-*i*]quinoline-2,6(1*H*)-dione, A-1544
N-Methyltryphophylline, D-793
O-Methyltryphophylline, D-793
N-Methyltryptamine, M-582
 13*a*-Methyltylohirsutine, M-583
 13*a*-Methyltylohirsutinidine, M-583
N-Methyltyramine *O*- α -*L*-rhamnopyranoside, H-667
 Methyltyramine, A-898
N-Methyltyramine, H-667
O-Methyltyramine, M-280
N-Methyltyrosine, T-700
 1-Methyl-2-(5-undecenyl)-4(1*H*)-quinolinone, H-757
 1-Methyl-2-(6-undecenyl)-4(1*H*)-quinolinone, H-757
 2-Methyl-6-undecylpiperidine, M-584
 1-Methyl-6-undecyl- Δ^2 -piperidine, M-584
 1-Methyl-2-undecyl-4(1*H*)-quinolinone, H-757
N-Methylurabaine, U-41
 5-Methyluracil, T-401
N^b-Methylusambarensine, U-52
N-Methylushinsunine, N-316
 4-Methylvaleric acid, M-505
 9-Methyl-*N*-vanillyl-7-decenamide, C-100
 7-Methyl-*N*-vanillyl-5-octenamide, C-100
N^a-Methylvellosimine, T-425
 8-*O*-Methylveratrolypseudaconine, M-336
N-Methylviguine, V-101
N-Methylvincadine, V-112
 Methylvingramine, V-133
 4-Methyl-5-vinylthiazole, M-585
O-Methylviridicatin, D-656
O-Methylviriditin, V-155
N-Methylvisoltricin, F-196
N^b-Methylvoacarpine, V-171
N^a-Methylvoachalotine, P-562
*N*¹-Methylvoafinine, V-179
N-Methylvoaphylline, V-179
 Methylwarifteine, W-9
N-Methylwelwitindolinone B isothiocyanate, W-13
N-Methylwelwitindolinone C isonitrile, W-13
N-Methylwelwitindolinone C isothiocyanate, W-13
N-Methylwelwitindolinone D isonitrile, W-14
 7-Methylwey, D-521
N-Methylxanthalinium, P-77
 7-Methylxanthine, X-5
 9-Methylxanthine, X-5
 1-Methylxanthine, X-5
 3-Methylxanthine, X-5
 7-Methylxanthopterin, C-432
 3-Methylxanthosine, X-13
N-Methylxylopinine, A-1066
 5-Methylxohimban, Y-12
 17-*O*-Methylxohimbine, Y-13
 1'-Methylzeatin, M-586
N-Methylzenkerine, T-524
 Metiprenaline, N-303
 Metocurine chloride, C-420
 Metocurine iodide, C-420
 Metopine, T-632
 Metubine iodide, C-420
 Metvin, N-301
 Mevashuntin, M-587
 Mexamine, M-288
 Mezcaline, M-219
 MF 5810, O-134
 610MF, A-1206
 MG 883-12F₂, S-569
 MG 30, T-441
 M6G, M-704
 MH 072, A-1199
 MH 435A, E-144
 MH 435B, E-144
 MI 481-42F4A, A-959
 MI 481-42F4B, A-960
 MI 481-42F4C, A-961
 MI 481-42F4D, A-962
 MI 19827, M-607
 MI 198Z1, M-607
 34M, I-132
 Micacocidin A, M-588
 Micacocidin B, M-588
 Micacocidin C, M-588
 Micacocidins, M-588
 Michelalbine, N-316
 Michelanugine, O-85
 Micheline A, N-316
 Micheline B, L-194
 Michellamine A, M-589
 Michellamine B, M-589
 Michellamine C, M-589
 Michellamine D, M-589
 Michellamine E, M-589
 Michellamine F, M-589
 Michellamine, M-589
 Michepressine, L-72
 Michigazone, M-590
 Micoserina, A-832
 Micranthine, M-591
 Microbiaeratin, A-1286
 Microbiaeratinin, B-1
 Microcarpine, M-592
 Microcin B17, M-593
 Microcin SF608, M-594
 Micrococcin M, M-595
 Micrococcin P₁, M-596
 Micrococcin P₂, M-596
 Microcylamide, M-597
 Micromeline, H-577
 Micromide, M-598
 Micromonosporin A, M-599
 Micropine, M-600
 Microsphaerone A, M-601
 Microsphaerone B, M-601
 Microsporin A, M-602
 Microsporin B, M-602
 Microxine, M-603
 Micutrin, P-961
 Midostaurin, S-498
 Midpacamide, M-604
 Miersine, M-605
 Migalastat hydrochloride, H-616
 Migalastat, H-616
 Miglustat, H-616
 Migranal, E-172
 Migrastatin, M-606
 Mikamycin A, O-131
 Mikamycin, O-131
 Mikaneicine, H-233
 Mikanoidine, S-90
 Mikimopine, C-802
 Milbemection, M-607
 Milbemycin A₃, M-607
 Milbemycin A₄, M-608
 Milbemycin B₂, M-607
 Milbemycin B₃, M-608
 Milbemycin C₁, M-607
 Milbemycin C₂, M-608
 Milbemycin D, M-609
 Milbemycin F, M-609
 Milbemycin G, M-609
 Milbemycin J, M-607
 Milbemycin K, M-608
 Milbemycin α 10, M-608
 Milbemycin α 11, M-607
 Milbemycin α 12, M-607
 Milbemycin α 13, M-607
 Milbemycin α 14, M-608
 Milbemycin α 15, M-608
 Milbemycin α 18, M-608
 Milbemycin α 19, M-608
 Milbemycin α 1, M-607
 Milbemycin α 20, M-607
 Milbemycin α 21, M-608
 Milbemycin α 22, M-607
 Milbemycin α 23, M-608
 Milbemycin α 24, M-607
 Milbemycin α 25, M-608
 Milbemycin α 26, M-607
 Milbemycin α 27, M-608
 Milbemycin α 2, M-607
 Milbemycin α 3, M-608
 Milbemycin α 4, M-608
 Milbemycin α 9, M-607
 Militarionone A, M-610
 Militarionone B, M-611
 Militarionone C, M-611
 Militarionone D, M-612
 Millaaurine A, M-614
 Millaaurine, M-613
 Millettonine, M-615
 Milliamine A, M-616
 Milliamine B, M-616
 Milliamine C, M-616
 Milliamine D, M-616
 Milliamine E, M-616
 Milliamine F, M-616
 Milliamine G, M-616
 Milliamine H, M-617
 Milliamine I, M-617
 Milliamine J, M-616
 Milliamine K, M-616
 Milliamine L, M-616
 Milliamine M, M-616
 Milliamine N, M-616
 Millingtonine, M-618
 Milnamide A, M-619
 Milnamide B, H-98
 Milnamide C, M-619
 Milnamide D, M-619
 Milonine, I-318
 Miltanthine, M-620
 Miltanthin, P-643
 Miltanthoridinone, M-621
 Miltanthoridinone, M-621
 Miltiromycin C, M-653
 Mimocin, M-622
 Mimopudine, M-623
 Mimosamycin, M-624
 Mimoside, L-148
 Mimosine, L-148
 Minalemine D, M-625
 Minalemine E, M-625
 Minalemine F, M-625
 Minalemine, M-625
 Minaline, P-944
 Minalobine O, M-626
 Minalobine R, M-627
 Minfiensine, M-628
 Miniatine, M-629
 Minoriceine, M-630
 Minoricine, M-630
 Minorine, V-115
 Minovincine, M-630
 Minovincinine, M-631
 Minovine, V-111
 Minpeimine, M-632
 Minpeiminine, M-633
 Miochol, A-47
 Mioton, C-100
 Mirabazole A, M-634
 Mirabazole B, M-635
 Mirabazole C, M-636
 Mirabilin A, M-639
 Mirabilin B, M-638
 Mirabilin C, M-639
 Mirabilin D, M-640
 Mirabilin E, M-641
 Mirabilin F, M-642
 Mirabilin G, M-643
 Mirabilin, M-637
 Mirabimide E, M-644
 Mirabliamide, H-350
 Miraxanthin III, M-647

- Miraxanthin II, M-646
 Miraxanthin I, M-645
 Miraxanthin V, M-647
 Miraziridine A, M-648
 Miramide 5-glucoside, D-639
 Miramide, D-639
 Misenine, M-649
 Misrametine, M-650
 Misramine, M-650
 Misrhybridine, R-116
 Mithaconitine, M-365
 Mitiphylline, M-651
 Mitiromycin A, M-652
 Mitiromycin B, M-653
 Mitiromycin D, M-654
 Mitiromycin E, M-654
 Mitocin C, M-654
 Mitomycin A, M-653
 Mitomycin B, M-653
 Mitomycin C, M-654
 Mitomycin D, M-654
 Mitomycin E, M-654
 Mitomycin F, M-653
 Mitomycin G, M-655
 Mitomycin H, M-656
 Mitomycin J, M-653
 Mitomycin K, M-653
 Mitomycin S, M-654
 Mitomycin, M-654
 Mitoridine, T-288
 Mitraciliatine, M-657
 Mitrafoline, R-134
 Mitragynaline, C-687
 Mitragynalinic acid, C-688
 Mitragynine, M-657
 Mitragynol, R-134
 Mitrajavine, M-658
 Mitralactonal, M-659
 Mitralactonine, M-660
 Mitraphyllic acid glucopyranosyl ester, M-661
 Mitraphyllic acid, M-661
 Mitraphylline *N*-oxide, M-661
 Mitraphylline, M-661
 Mitraspecine, M-662
 Mitrasulglynine, M-663
 Mitraversine, M-664
 Mitrinermine, R-90
 Miyaconitine, M-665
 Miyaconitinone, M-665
 Miyoshianine A, L-344
 Miyoshianine B, L-322
 MJ 286A, T-374
 MJ 347-81F4-A, N-266
 MJ 347-81F4-B, N-266
 MK 61, N-278
 ML 1532II, C-900
 ML 1065, R-4
 MLR 52, S-498
 MM 14201, A-804
 MM 21801, A-759
 MMC 4, M-654
 MOCO, M-681
 Mocrispatine, S-39
 Moderil, R-50
 Modestanine, M-666
 Modus, P-80
 Moenjodaramine, M-667
 Moiramide A, M-668
 Moiramide B, M-669
 Moiramide C, A-1004
 Mollenine A, M-670
 Mollenine B, M-670
 Molliclavine, M-671
 Mollinedine, M-672
 Molline, M-315
 Molliorin A, M-673
 Molliorin B, M-674
 Molliorin C, M-675
 Molliorin D, M-676
 Molliorin E, M-677
 Mollisine, M-678
 Molokaiamine, M-679
 Mololipids, M-680
 Molybdenum cofactor, M-681
 Molybdoenzyme molybdenum cofactor, M-681
 Molybdopterin, M-681
 Monamidocin, M-682
 Monanchorin, M-683
D-Monapterin, A-923
L-Monapterin, A-923
 Monascamine, M-686
 Monascopyridine A, M-684
 Monascopyridine B, M-685
 Monascorubramine, M-686
 Monascumic acid, M-482
 Monaspiloidine, I-96
 Monaspyranoindole, T-169
 Monil, E-6
 Monoacetylglycerine, F-223
 Monoacetylthrumine, L-379
 Monoacetylspermine, S-399
 Monobromoisophakellin, D-328
 Monobromophakellin, B-343
N-Mono(*tert*-butyloxycarbonyl)ethylenediamine, E-225
 Monochlorotyrosine, A-724
 Monocillinol A, M-687
 Monocillinol B, M-687
 Monocrotaline *N*-oxide, M-688
 Monocrotaline pyrrole, M-688
 Monocrotaline, M-688
 Monocrotalinine, M-688
 9-Monodechloro-13-demethylisodysidenin, D-972
 11-Monodechloro-13-demethylisodysidenin, D-972
 Mono-2-dimethylaminoethanol dihydrogen phosphate, D-197
 Monodontamide A, M-689
 Monodontamide B, M-689
 Monodontamide C, M-689
 Monodontamide D, M-690
 Monodontamide E, M-690
 Monodontamide F, M-691
 Monodoroidole, M-692
 Monoethanolamine, A-767
 Monoferuloylamaranthin, B-110
 Monoferuloylbetanin, B-110
 Monogagine, M-693
 Monohydroxyisoaflavine, A-172
 Monolupine, A-970
 Monomargine, M-694
 Monomethyl caulerpinate, D-415
 Monomethylholamine, A-888
 Monomethyltetrandrinium, T-286
 Monomorine I, B-434
 Monophosphothiamine, T-366
 Monospermin, M-385
 Monspessulanine, M-695
 Montabuphine, M-696
 Montamine, M-697
 Montanine, P-57
 Montanine, Q-33
 Montanine, V-178
 Monterine, C-643
 Monticamine, D-187
 Monticoline, D-187
 Montiporyne E, M-698
 Montipyridine, M-699
 Montrifoline, F-221
 Montrutanine, Q-33
 Mooniine A, M-700
 Mooniine B, M-700
 Mooreine, M-701
 Moranoline, H-616
 Morfine, M-704
 Morifoline, I-68
 Moringine, B-84
 Moroidin, M-702
 Morphine glucuronide, M-704
 Morphine *N*-oxide, M-704
 Morphine sulfate, M-704
 Morphine, M-704
 ψ -Morphine, M-703
 4-Morpholinamine, M-705
 4-Morpholinecarboxaldehyde, M-705
 4-Morpholineethanol, M-705
 Morpholine, M-705
 1-Morpholino-1-propanone, M-705
 Mortivinacin B, M-706
 Mortivinacin C, M-707
 Morusimic acid A, M-708
 Morusimic acid B, M-708
 Morusimic acid C, M-709
 Morusimic acid D, M-709
 Morusimic acid E, M-709
 Morusimic acid F, M-709
 Moschamide, M-710
 Moschamindole, S-252
 Moschamindolol, M-711
cis-Moschamine, H-755
 Moschamine, H-755
 Moschatoline, M-712
 Mossabine, M-713
 Mostueine, D-138
 Motuporamine A, M-714
 Motuporamine B, M-715
 Motuporamine C, M-716
 Motuporamine D, M-715
 Motuporamine E, M-716
 Motuporamine F, M-716
 Moupinamide, H-667
 Movellan, S-589
 Moxa, N-303
 MP 105, A-772
 MPC 1001F, A-1207
 MR 93A, M-159
 MR 93B, O-156
 MRL 3120, F-97
 MS 681, A-1208
 MS 681A, A-1208
 MS 681B, A-1208
 MS 681C, A-1208
 MS Contin, M-704
 MS 681D, A-1208
 MSD 819, C-416
 MSD 92, D-498
 MSI 1256F, S-485
 MSI 1436, S-485
 MT 35214, A-804
 MT 36531, A-804
 MT 1882-II, P-421
 MT 1882-I, P-421
 Mubironine A, D-210
 Mubironine B, D-210
 Mubironine C, M-717
 Mucronatine, A-818
 Mucronatine, R-62
 Mucronatinine, R-62
 Mucronine A, M-718
 Mucronine B, M-718
 Mucronine C, M-719
 Mucronine D, M-720
 Mucronine E, M-719
 Mucronine F, M-719
 Mucronine G, A-18
 Mucronine H, M-718
 Mucronine J, M-721
 Mucquadine, M-722
 Mucquadinine, M-723
 Mucquadininine, M-724
 Mucunadine, M-725
 Mucunine, M-726
 Mukanadin A, D-863
 Mukanadin B, M-727
 Mukanadin C, L-242
 Mukanadin D, M-727
 Mukeic acid, H-452
 Mukoic acid, H-452
 Mukoenine A, M-728
 Mukoenine B, M-729
 Mukoenine C, G-81
 Mukolidine, H-451
 Mukoline, H-604
 Mukonal, H-448
 Mukonicine, G-81
 Mukonidine, H-453
 Mukonine, H-452
 Muldamine, S-200
 Mulgediifoline, N-89
 Multhiomycin, N-319

- Multifidin, H-575
 7,9(11)-Multifloradiene-3,29-diol, M-730
 Multifloramine, P-221
 (+)-Multifloramine, P-221
 Multiflorine *N*-oxide, M-731
 Multiflorine, M-731
 Munitagine, P-514
 Munroniamide, M-732
 Munronin D, M-733
 Munzianine, M-347
 Munzianone, M-347
 Mupamine, G-81
 Muramine, M-734
 Murayaanthraquinone, M-735
 Muraymycin A, M-736
 Muraymycin A₃, M-736
 Muraymycin A₄, M-736
 Muraymycin A₅, M-736
 Muraymycin B₁, M-737
 Muraymycin B₂, M-737
 Muraymycin B₃, M-737
 Muraymycin B₄, M-737
 Muraymycin B₅, M-737
 Muraymycin B₆, M-737
 Muraymycin B₇, M-737
 Muraymycin C₁, M-737
 Muraymycin C₂, M-737
 Muraymycin C₃, M-737
 Muraymycin C₄, M-737
 Muraymycin D₁, M-737
 Muraymycin D₂, M-737
 Muraymycin D₃, M-737
 Mureidomycin E, M-738
 Mureidomycin F, M-738
 Murexine, M-739
 Muricatide, A-864
 Muricatisine, A-694
 Muricine, M-740
 Muricinine, C-702
 Murrafoline A, M-741
 Murrafoline B, M-742
 Murrafoline C, M-743
 Murrafoline D, M-744
 Murrafoline E, M-745
 Murrafoline F, M-746
 Murrafoline G, M-747
 Murrafoline H, M-748
 Murranimbine, M-749
 Murrapanine, M-750
 Murrastifoline A, M-751
 Murrastifoline B, M-752
 Murrastifoline C, M-753
 Murrastifoline D, M-754
 Murrastifoline E, M-754
 Murrastifoline F, M-755
 Murrayacarine, M-756
 Murrayacine, M-757
 Murrayacinine, M-47
 Murrayaculatine, M-758
 Murrayafoline A, H-583
 Murrayafoline B, M-759
 Murrayaline A, D-619
 Murrayaline B, D-619
 Murrayaline C, M-760
 Murrayaline D, E-276
 Murrayaline, D-619
 Murrayamine A, G-81
 Murrayamine B, M-47
 Murrayamine C, M-46
 Murrayamine C, M-47
 Murrayamine D, C-814
 Murrayamine E, C-813
 Murrayamine F, M-761
 Murrayamine G, M-761
 Murrayamine H, C-814
 Murrayamine I, G-81
 Murrayamine J, M-762
 Murrayamine K, G-81
 Murrayamine M, M-763
 Murrayamine N, M-762
 Murrayamine O, M-764
 Murrayamine P, M-764
 Murrayanine, H-447
 Murrayanine, M-765
 Murrayanol, E-276
 Murrayaquinone A, M-425
 Murrayaquinone B, M-766
 Murrayaquinone C, M-767
 Murrayaquinone D, M-767
 Murrayaquinone E, M-766
 Murrayastine, T-596
 Murrayazolidine, C-814
 Murrayazoline, C-813
 Murrayazolinine, C-814
 Murrayazolinol, C-813
 Muscaaurin II, M-769
 Muscaaurin I, M-768
 Muscaflavin, M-770
 Muscapurpurin, M-771
 Muscaridine, M-772
 Muscarine, M-773
 Muscarone, M-773
 Muscicapine A, A-778
 Muscicapine B, A-778
 Muscicapine C, A-778
 Muscimol, A-837
 Muscopyridine, M-774
 Muscoride A, M-775
 Musculamine, S-399
 Mussaenoside A, H-90
 Mussaenoside B, H-90
 Mussaenoside C, H-90
 Mussaenoside D, H-90
 Mussaenoside E, H-90
 Mussaenoside F, H-90
 Mussaenoside G, H-90
 Mussaenoside H, H-90
 Mussaenoside M, H-90
 Mussaenoside N, H-90
 Mussaenoside O, H-90
 Mussaenoside P, D-564
 Mussaenoside Q, D-602
 Mussaenoside U, H-90
 Mutaaspergillic acid, M-462
 Mutabillicin, S-321
 Mutabilycin, S-321
 Mutamycin, M-654
 Mx 2BC, S-6
 MY 336a, A-1209
 MY 12-62a, H-133
 MY 12-62c, H-134
 MYC 8005, A-1210
 Mycadrine, A-898
 Mycalazal 10, A-624
 Mycalazal 11, A-624
 Mycalazal 12, A-624
 Mycalazal 13, A-624
 Mycalazal 1, A-624
 Mycalazal 2, A-624
 Mycalazal 3, A-624
 Mycalazal 4, A-624
 Mycalazal 5, A-624
 Mycalazal 6, A-624
 Mycalazal 7, A-624
 Mycalazal 8, A-624
 Mycalazal 9, A-624
 Mycalazole 11, M-776
 Mycalazole 12, M-776
 Mycalazole 5, M-776
 Mycalazole 6, M-776
 Mycalazole 8, M-776
 Mycalazole 9, M-776
 Mycalazoles, M-776
 Mycalenitrile 1, F-141
 Mycalenitrile 2, F-141
 Mycalenitrile 3, F-141
 Mycaleoxime, M-777
 Mycalisine B, M-778
 Mycalolide A, M-779
 Mycalolide B, M-779
 Mycalolide C, M-779
 Mycalolide D, M-780
 Mycalolide E, M-781
 Mycapolyols, M-782
 Myceliothermophin A, M-783
 Myceliothermophin B, M-783
 Myceliothermophin C, M-783
 Myceliothermophin D, M-783
 Myceliothermophin E, M-783
 Mycenarubin A, M-784
 Mycenarubin B, M-785
 Mycenarubin D, M-786
 Mycenarubin E, M-786
 Mycenarubin F, M-784
 Mycobacidin, A-78
 Mycobactin J1, M-787
 Mycobactin T, M-788
 Mycobactin, M-787
 Mycomethoxin A, P-325
 Mycomethoxin B, H-663
 Mycosporin glutamicol, M-789
 Mycosporin glutaminol, M-789
 Mycosporin-Gln, M-789
 Mycosporin-Glu, M-789
 Mycothiazole-4,19-diol, M-790
 Mycothiazole, M-790
 Mycothiol bimane, M-791
 Mydriaside, T-632
 Mydriatine, A-869
 Myoscorpine *N*-oxide, E-31
 Myoscorpine, E-31
 Myosmine, M-792
 Myrianthine A, M-793
 Myrianthine B, A-150
 Myrianthine C, M-794
 Myrianthosine, M-795
 Myricoidine, L-230
 Myrioneurinol, M-796
 Myrionine, M-797
 Myrioxazine A, O-57
 Myrioxazine B, O-57
*N*²-Myristoleoylglutamine, G-108
*N*²-Myristoylglutamine, G-108
 Myrmicaric 237A, B-436
 Myrmicaric 213A, M-801
 Myrmicaric 215A, M-801
 Myrmicaric 233A, M-799
 Myrmicaric 430A, M-800
 Myrmicaric 237B, B-436
 Myrmicaric 213B, M-801
 Myrmicaric 215B, M-801
 Myrmicaric 215C, M-801
 Myrmicaric 217, M-801
 Myrmicaric 663, M-798
 Myrobotinol, M-802
 Myrothenone A, A-824
 Myrothenone B, A-824
 Myrrhine, D-886
 Myrtifoline, F-223
 Myrtine, O-53
 Myrtoidine, M-803
 Myrtopsine, M-804
 MYX 6121, O-37
 Myxalamides, M-805
 Myxin, P-321
 Myxocheilin A, M-806
 Myxocheilin B, M-806
 Myxocheilin C, M-806
 Myxothiazole A, M-807
 17Z-Myxothiazole A, M-807
 Myxothiazole Z, M-807
 Myxovalargin, M-808
 Myxoviromycin, A-692
 N 461, C-102
 N 13, R-160
 NA 22598A₁, A-1211
 NA 30851A, A-1212
 NA 337, T-213
 (Naamidinato A)(naamidinato G)zinc(II), N-1
 Naamidine A, N-1
 Naamidine B, N-1
 Naamidine C, N-1
 Naamidine D, N-1
 Naamidine E, P-938
 Naamidine F, N-2
 Naamidine G, N-1
 Naamidine H, N-1
 Naamidine I, N-1
 Naamine A, N-3

- Naamine B, N-4
 Naamine C, N-5
 Naamine D, N-6
 Naamine E, N-5
 Naamine F, N-5
 Naamine G, N-5
 Nagadine, M-375
 Nagarine, M-302
 Nagarine, M-313
 Nagelamide A, N-7
 Nagelamide B, N-7
 Nagelamide C, N-7
 Nagelamide D, N-7
 Nagelamide E, A-190
 Nagelamide F, A-190
 Nagelamide G, A-190
 Nagelamide H, N-8
 Nagstatin, N-9
 Nairomycin, N-10
 Nakadomarin A, N-11
 Nakamuric acid, N-12
 Nakamurol D, C-525
 Nakijinol, N-13
 Nakijiquinone A, N-14
 Nakijiquinone B, N-15
 Nakijiquinone C, N-16
 Nakijiquinone D, N-16
 Nakinadine A, N-17
 Nakinadine B, N-18
 Nakinadine C, N-19
 Nakinadine D, N-20
 Nakinadine E, N-21
 Nakinadine F, N-22
 Nakirodine A, N-23
 Naloamine, A-760
 Nandazurine, N-24
 Nandigerine, N-25
 Nandinine, C-73
 Nandinin, H-520
 Nangustine, N-26
 Nankakurine A, N-27
 Nankakurine B, N-27
 Nannochelin A, N-28
 Nannochelin B, N-28
 Nannochelin C, N-28
 Nanofin, D-770
 Nanophyn, D-770
 Nantemine, N-29
 Nantoamide, M-442
 Napelline N-oxide, N-30
 Napelline, N-30
 Napellonine, N-30
 Naphthisoxazole A, N-31
 Naphthocyanidine, C-830
 7*H*-Naphtho[1,2,3-*ij*][2,7]naphthyridin-7-one, S-49
 Naphthyridinomycin A, N-32
 Naphthyridinomycin B, N-33
 1-(2,7-Naphthyridin-3-yl)ethanone, A-72
 Napsamycin A, M-738
 Napsamycin B, N-34
 Napsamycin C, M-738
 Napsamycin D, N-34
 Naramycin A, D-765
 Naramycin B, D-765
 Naramycin, D-765
 Narceimicine, A-143
 Narceimine, A-143
 Narceine imide, N-36
 Narceine, N-35
 Narceinone, N-35
 Narciclasine 4-*O*-glucoside, N-37
 Narciclasine, N-37
 Narcicrinine, N-38
 Narcicriptine, N-39
 Narcidine, N-40
 Narcimarkine, H-5
 Narcipoetine, L-347
 Narciprimine, C-744
 Narcisine, N-41
 Narcisline, N-42
 Narcissidine, N-43
 Narcissine, L-348
 Narcotinediol, N-45
 Narcotinehemiacetal, N-46
 Narcotine, N-44
 (\pm)- α -Narcotine, N-44
 (\pm)- β -Narcotine, N-44
 (+)- α -Narcotine, N-44
 (-)- α -Narcotine, N-44
 (-)- β -Narcotine, N-44
 Narcotoline, N-44
 Narcotolinol, N-45
 Nareline, N-47
 Nargenicin A₁, N-276
 Nargenicin B₁, N-48
 Nargenicin B₂, N-49
 Nargenicin B₃, N-50
 Nargenicin C₁, N-276
 Nargin, P-923
 Naringenic acid, H-691
 Naringeninic acid, H-691
 Naritheracin, S-53
 Narlumicine, N-51
 Narlumidine, N-51
 Nartazine, Z-15
 Narvedine, G-3
 Narwedine, G-3
 Narzettine, N-52
 Natalamine, N-53
 Natalensine, H-5
 Natalinine, N-53
 Natrine, N-54
 Natsucitrine II, T-243
 Natsucitrine I, T-243
 Natural capsaicin, C-100
 Natural yellow 18, B-101
 Naucleactonine A, N-55
 Naucleactonine B, N-56
 Nauclealine A, N-57
 Nauclealine B, N-58
 Naucleamide A, N-59
 Naucleamide B, N-59
 Naucleamide C, N-60
 Naucleamide D, N-59
 Naucleamide E, N-61
 Naucleaorine, N-62
 Nauclechine, N-63
 Nauclecoside, V-126
 Naucleosidine, N-64
 Naucleoderine, N-65
 Naucleidine, N-66
 Naucleficine, N-67
 Nauclefidine, N-68
 Nauclefiline, N-69
 Nauclefine, N-70
 Nauclefoline, N-71
 Nauclefoline, N-72
 Nauclefolinine, N-73
 Naucleidinal, N-74
 Naucleocoside, V-126
 Naucleoficine A, N-59
 Naucleoficine B, N-57
 Naucleoficine C, N-75
 Naucleoficine D, N-60
 Naucleoficine E, A-1020
 α -Naucleonidine, N-76
 β -Naucleonidine, N-76
 Naucleonidine, N-76
 α -Naucleonine, N-76
 β -Naucleonine, N-76
 Naucleoside A, V-126
 Naucleoside B, V-126
 Nauclequiniine, N-67
 Nauctetine, A-1020
 Nauclexine, N-77
 Naufoline, N-63
 Naulafine, N-78
 Navenone A, P-908
 Navirine, N-79
 Nazlinine, N-80
 NCI 1136, D-92
 Neamphine, N-81
 Neblinine, O-7
 Nebularine, N-82
 Neburnamine, N-83
 Necatorine, N-84
 Necatorone, N-85
Senecio doria Necine base, N-86
 Necrosamine, E-47
 Nectrisine, D-420
 Neferine, N-87
 Neflexine, N-88
 Neisosposinine, C-117
 Nelumstemine, L-252
 Nematophin, T-639
 Nemertelline, Q-5
 Nemorensine, N-89
 Nemorine, N-90
 Nemuarine, N-91
 Neoacrimarine A, N-92
 Neoacrimarine B, N-92
 Neoacrimarine C, N-93
 Neoacrimarine D, N-94
 Neoacrimarine E, N-95
 Neoacrimarine F, N-96
 Neoacrimarine G, N-96
 Neoacrimarine H, N-93
 Neoacrimarine I, N-96
 Neoacrimarine J, N-96
 Neoacrimarine K, N-97
 Neoacutifolin, B-13
 Neoadifoline, N-98
 Neoalangiside, A-239
 Neolatamine, P-253
 Neoamphimedine Y, N-100
 Neoamphimedine Z, N-100
 Neoamphimedine, N-99
 Neoaplaminone sulfate, A-1342
 Neoaplaminone, A-1342
 Neospergillitic acid, F-76
 Neoasterriquinone, A-1521
 Neoberninamycin, N-101
 Neobetamin, B-110
 Neobudofficide B, N-102
 Neocallactine, A-1459
 Neocarazostatin A, N-103
 Neocarazostatin B, N-103
 Neocarazostatin C, N-103
 Neocaryachine *M*-methosalt, N-104
 Neocaryachine, N-104
 Neocephaline, E-72
 Neochondocurarine, N-105
 Neoclausenamide, C-502
 Neocochliodinol, N-106
 Neocopiamycin A, C-634
 Neocopiamycin B, C-634
 Neocoprogen II, N-108
 Neocoprogen I, N-107
 Neocoromandaline, T-443
 Neocryptolepine, Q-19
 Neocycasin A, M-397
 Neocycasin B, M-397
 Neocycasin B₂, M-397
 Neocycasin C, M-397
 Neocycasin E, M-397
 Neocycasin G, M-397
 Neocycasin H, M-397
 Neocycasin I, M-397
 Neocycasin J, M-397
 Neodaphniphylline, N-109
 Neodichroine, N-110
 Neodihydrohistrionicotoin, H-309
 Neodihydrothebaine, N-111
 Neodrosoplerin, D-941
 Neodysidenin, D-972
 Neodysiherbaine A, N-112
 Neoenactin C, P-444
 Neoenactin M₁, L-189
 Neoenactin M₂, N-113
 Neoenactin, N-113
 Neoengleromycin, N-114
 Neoeuonymine, P-247
 Neoevovine, P-248
 Neofinaconitine, A-95
 Neofolitispatate 1, N-115
 Neofolitispatate 2, N-115
 Neofolitispatate 3, N-115
 Neogermbudine, G-70
 Neogermerine, G-70

- Neogermidine, G-70
 Neogermine, N-116
 Neogermitrine, G-70
 Neoglucobrassicin, I-122
 Neoharringtonine, N-117
 Neoherculin, D-893
 Neohuperzine, N-118
 Neohyacinthoside, S-345
 Neohydroxyaspergillus acid, F-76
 Neohydroxylumine, N-119
 Neopecoside, I-167
 Neoisocycloheximide, D-765
 Neoisotussilag acid, H-231
 Neoisotussilage, H-231
 Neojerminalanine, G-70
 Neojiangyouaconitine, M-343
 Neokauluamine, N-120
 Neolamellarin A, N-121
 Neolamellarin B, N-121
 Neolatifoline, H-4
 Neolaugerine, N-122
 Neoleurocristine, N-123
 Neoleurosidine, N-124
 Neoligularidine, S-245
 Neoline, M-323
 Neoline, M-323
 Neolinustatin, H-572
 Neolitacumomine, H-424
 Neolitsine, N-125
 Neolitsimine, N-126
 Neonaucleoside A, N-127
 Neonaucleoside B, N-128
 Neonaucleoside C, N-129
 Neonicotine, P-460
 Neonorreserpine, R-52
 Neooncinothine, N-130
 Neoxazolomycin, N-131
 Neooxygambirtannine, N-132
 Neopellitorine A, U-25
 Neopellitorine B, D-104
 Neopeltolide, N-133
 Neoperiphylline, P-265
 Neopetasitenine, P-277
 Neopine, N-134
 Neoplaether, O-129
 Neoplatyphylline *N*-oxide, P-516
 Neoplatyphylline, P-516
 Neopolyalcohol, N-135
 Neopolyoxin A, N-216
 Neopolyoxin B, N-216
 Neopolyoxin C, N-218
 Neoprotocuridine, P-663
 Neoprotoveratrine, P-674
 Neopterin 3-*O*- β -*D*-glucoside, A-923
 7-Neopterin, A-924
 Neopterin, A-923
 Neopulchellidine, P-785
 Neopurpuratin, D-496
 Neopyrrolomycin, N-136
 Neoreserpiline, R-51
 Neorosmarinine, R-133
 Neorythmin, A-216
 Neosabadine, C-291
 Neosalvianene, N-137
 Neosarracine, S-90
 Neosarranicine, S-90
 Neosaxitoxin, S-109
 Neosenkirkine, S-245
 Neosidomycin, N-138
 Neosphoramine, S-376
 Neostemocochinine, N-140
 Neostemofoline, N-139
 Neostemonine, N-140
 Neostenine, S-528
 Neostychnine, S-589
 Neosurugatoxin, N-141
 Neosutchuenenine, N-142
 Neosymbioimine, N-143
 Neothalfine, N-144
 Neothalbrine 2' α -*N*-oxide, T-309
 Neothalbrine, T-309
 Neothiobinupharidine sulfoxide, T-380
 Neothiobinupharidine, T-380
 Neothramycin A, N-145
 Neothramycin B, N-145
 Neotomatine, S-445
 Neotriangularicine, T-454
 Neotriangularine, T-454
 Neotrilobine, N-146
 Neotruixilline, T-635
 Neotuberostemonine, T-664
 Neotuberostemoninol, T-665
 Neotuberostemonol, T-664
 Neotuberostemonone, T-666
 Neotussilage, H-231
 Neotussilaginic acid, H-231
 Neovasipyridone A, N-147
 Neovasipyridone B, N-147
 Neovasipyridone C, N-147
 Neovasipyridone D, N-147
 Neovasipyridone E, N-147
 Neovasipyridone F, N-147
 Neovasipyridone, N-147
 Neoverapatuline, J-36
 Neoverataline A, N-148
 Neoverataline B, N-148
 Neowilforine, P-243
 Neoxaline, N-149
 Neoyuzurimine, N-150
 Neozeylanine, M-488
 Nepapakistamine A, D-273
 Nepetalactam, N-151
 Nephilatoxin 10, N-160
 Nephilatoxin 11, N-161
 Nephilatoxin 12, N-162
 Nephilatoxin 1, N-152
 Nephilatoxin 2, N-153
 Nephilatoxin 3, N-154
 Nephilatoxin 473, N-163
 Nephilatoxin 4, N-155
 Nephilatoxin 501, N-164
 Nephilatoxin 565, N-165
 Nephilatoxin 594, N-166
 Nephilatoxin 5, N-156
 Nephilatoxin 643, N-167
 Nephilatoxin 6, N-157
 Nephilatoxin 7, N-158
 Nephilatoxin 8, N-152
 Nephilatoxin 9, N-159
 Neplanocin B, N-168
 Neprotine, P-34
 Nerbowdine, N-169
 Nerifline, N-170
 Neriine, C-601
 Nerinine, N-171
 Nerispine, N-172
 Neronine, K-102
 Nerundine, N-173
 Neruscine, L-347
 Nervobscurine, A-231
 Nervonic acid, T-120
 Nervosine, P-39
 Nesodine, V-88
 Netamine A, N-174
 Netamine B, N-175
 Netamine C, N-176
 Netamine D, N-174
 Netamine E, N-177
 Netamine F, N-178
 Netamine G, N-179
 Netropsin, N-180
 Neumolisina, Q-20
 Neumolysin, Q-20
 Neuridine, S-399
 Neurine, T-615
Nephila clavata Neurotoxin 1 JSTX-1, N-182
Nephila clavata Neurotoxin 2 JSTX-2, N-183
Nephila clavata Neurotoxin 4 JSTX-4, N-184
 Neurotoxin JSTX-3, N-165
Nephila maculata Neurotoxin 3, N-181
 Neurotoxin NPXT 10, N-160
 Neurotoxin NSTX 3, N-181
 Neurotropan, C-419
 Nevadenine, G-152
 Nevadensine, G-152
 Nevalina, G-3
 Nevergor, A-216
 Newbouldine, D-492
 Newbouldine, N-185
 Newtron S, D-880
 NFA Toxin, F-62
 NFAT 68, A-1213
 NG 311, A-1214
 NG 391, A-1215
 NG 392, A-1215
 NG 393, A-1215
 NG 129, S-490
 NG 242, S-490
 NG 243, S-490
 NG 245, S-490
 Ngouniensine, N-186
 NI 15501A, A-33
 34N, I-132
 Niacinamide, P-895
 Niacin, P-897
 Niazicin A, H-427
 Niazicin B, H-427
 Niazicinin A, H-421
 Niazimicin A, H-427
 Niazimicin B, H-427
 Niazimin A, H-421
 Niazimin B, H-421
 Niaziminin A, H-427
 Niaziminin B, H-427
 Niazinin A, H-427
 Niazinin B, H-427
 Niazirin, H-672
 Niazirinin, H-672
 Nicaeensin, N-187
 Nicamin, P-897
 Nicaustrine, T-47
 Nicobid, P-897
 Nicoloxide, P-904
 Nicorette, N-190
 Nicosan 2, P-895
 Nicotaxine, N-188
 Nicotelline, T-98
 Nicotimine, P-460
 Nicotinamide mononucleotide, N-189
 Nicotinamide ribonucleoside, N-189
 Nicotinamide ribonucleotide, N-189
 Nicotinamide riboside, N-189
 Nicotinamide ribotide, N-189
 Nicotinamide, P-895
 Nicotine bitartrate, N-190
 Nicotine 1'-*N*-oxide, N-190
 Nicotine polacrifex, N-190
 Nicotine, N-190
 Nicotinic acid amide, P-895
 Nicotinic acid, P-897
 11-*O*-Nicotinoylambelline, A-683
 2-(Nicotinoylamino)-1-phenylethanol, A-864
*O*²-Nicotinoylcathedulin E8, P-218
 12-*O*-Nicotinoyldeacetylmetaplexigenin, P-223
 Nicotinoyllecgonine methyl ester, E-8
 6-*O*-Nicotinoyl- α -*D*-glucopyranose, B-376
 1-*O*-Nicotinoyl- β -*D*-glucopyranose, P-897
N-Nicotinoyl-2-hydroxy-2-phenylethylamine, A-864
 3 α -Nicotinoyloxytropone, M-393
 6-*O*-Nicotinoylstrychnovoline, T-145
 6-*O*-Nicotinoyltetrahydrocantlyline, T-145
 Nicotiny alcohol, P-904
 Nicotiny salicylate, N-190
 β -Nicotyrine, P-972
 Nicotyrine, P-972
 Nigakinone, D-546
 Nigdenine, F-223
 Nigeglanine, N-191
 Nigelline, N-192
 Nigellidine, N-193
 Nigellimine *N*-oxide, D-627
 Nigellimine, D-627
 Nigerazine A, D-769
 Nigerazine B, D-769
 Nigerine, D-785
 Nigleroxin, N-194
 Nigragilline, D-769
 Nigramide A, N-196

- Nigramide B, N-197
 Nigramide C, N-198
 Nigramide D, N-199
 Nigramide E, N-200
 Nigramide F, N-197
 Nigramide G, N-201
 Nigramide H, N-202
 Nigramide I, N-195
 Nigramide J, N-203
 Nigramide K, N-204
 Nigramide L, N-205
 Nigramide M, N-206
 Nigramide N, N-207
 Nigramide O, N-208
 Nigramide P, N-209
 Nigramide Q, N-210
 Nigramide R, N-211
 Nigramide S, N-212
 Nigrifactin, N-213
 Nigrifortine, A-680
 Nigrinodine, N-214
 Nigrin, S-569
 Nigritanine, O-21
 Nigrumin, H-509
 NIH 7380, G-3
 Nikanine *N*-oxide, N-215
 Nikanine, N-215
 Nikkomycin D, A-796
 Nikkomycin L_x, N-216
 Nikkomycin L_z, N-218
 Nikkomycin Q_x, N-217
 Nikkomycin Q_z, N-216
 Nikkomycin X, N-216
 Nikkomycin Z, N-218
 Nikkomycin, N-218
 Nilgirine, N-219
 Nilyph, X-5
 Ningalin A, N-220
 Ningalin B, N-221
 Ningalin C, N-222
 Ningalin D, P-825
 Ningpeisine, N-223
 Ningpeisinoside, N-223
 Niphatesine A, T-357
 Niphatesine B, N-224
 Niphatesine C, N-225
 Niphatesine D, N-226
 Niphatesine E, N-227
 Niphatesine F, N-228
 Niphatesine G, N-229
 Niphatesine H, N-227
 Niphatoxin A, N-230
 Niphatoxin B, N-231
 Niphatoxin C, N-232
 Niphatyne A, N-233
 Niphatyne B, N-234
 Niphithricin A, C-634
 Niphitricin 1, C-634
 Niranin, N-235
 Niromycin, N-10
 Nirurine, N-236
 Niruroidine, B-375
 Nishindine, N-237
 Nitaphylline, N-238
 Nitensidine A, B-174
 Nitensidine B, B-173
 Nitensidine C, B-174
 Nitidine, N-239
 Nitrabirine *N*-oxide, N-240
 Nitrabirine, N-240
 Nitramarine, Q-51
 Nitramidine, N-246
 Nitramine, N-241
 2-Nitraminoquinoline, Q-28
 Nitraraidine, N-242
 Nitraraine, N-243
 Nitramidine, N-244
 Nitramine *N*-oxide, N-244
 Nitramine, N-244
 Nitraricine, N-245
 Nitraridine, Q-52
 Nitrarine, N-246
 Nitrarisine, N-246
 Nitrarizine, N-246
 Nitroxine, N-244
 Nitricholine perchlorate, C-419
 16,23-Nitrolocycloarta-7,16(*M*),17(20),22-tetraene-3,11,24,25-tetrol, N-247
 3,3',3''-[Nitrilotris(methylene)]tris[dihydro-2(3*H*)furanone], N-248
 1-Nitroaknadinine, H-70
 1-Nitroapocavidine, C-222
 3-(12-Nitrododecyl)pyridine, N-249
 Nitrohedon, H-639
 2-Nitroimidazole, N-250
 2-Nitro-1*H*-indole-3-ethanamine, A-771
 3-Nitro-1*H*-indole, N-251
 4-Nitromorpholine, M-705
 6-Nitrophenanthro[3,4-*d*]-1,3-dioxole-5-carboxylic acid, M-447
*N*²-[(2-Nitrophenyl)thio]-L-glutamine, G-108
N-[(2-Nitrophenyl)thio]-L-leucine, L-153
 Nitropolyzonamine, N-252
 2-[6-(3-Nitropropanoyl)-β-D-glucopyranosyl]-3-isoxazolin-5-one, I-349
 3'-Nitropolyuteorin, P-834
 1-Nitro-12-(3-pyridinyl)dodecane, N-249
 1-Nitro-14-(3-pyridinyl)-11-tetradecene, N-256
 1-Nitro-14-(3-pyridinyl)-11-tetradecyne, N-256
N-Nitrosoanonaine, R-120
 3-Nitrosoindole, I-106
 Nitrostromelin, A-693
N-Nitrosotomatidine, S-445
 Nitrosoxacin A, N-253
 Nitrosoxacin B, N-254
 Nitrosoxacin C, N-255
N-Nitrosoxylopinine, A-1066
 3-(14-Nitro-3-tetradecenyl)pyridine, N-256
 3-(14-Nitro-3-tetradecynyl)pyridine, N-256
 2-Nitrotryptamine, A-771
 4-Nitrotryptophan, N-257
 11-Nitrotubotaiwine, T-673
 3-Nitrotyramine, H-640
 Nitrotyrasacutuminine, A-134
 Nitrotyrasanguinarine, N-258
 Nivalidine, A-1361
 Nivaline, H-294
 Nivalin, G-3
 Njaoamine A, N-259
 Njaoamine B, N-259
 Njaoamine C, N-259
 Njaoamine D, N-259
 Njaoamine E, N-260
 Njaoamine F, N-261
 Njaoamine G, N-259
 Njaoamine H, N-259
 NK 30424A, A-1216
 NK 170204A, P-421
 NK 372135A, X-7
 NK 170204B, A-1217
 NK 30424B, A-1216
 NK 372135B, X-7
 NK 109, C-356
 NK 372135C, X-7
 NK 26588, I-179
 Nkolbisine, F-221
 NKT 01, S-394
 NMQ, M-491
 Nobiline, F-221
 Nobiline, N-262
 Nobilisine, C-532
 Nobilisitine A, C-532
 Nobilisitine B, C-532
 Nobilonine, N-262
 δ-Nobilonine, N-263
 Nocamycin II, N-264
 Nocamycin I, N-264
 Nocardamine ferric salt, F-42
 Nocardimicin C, N-265
 Nocardimicin E, N-265
 Nocardimicin G, N-265
 Nocardimicin H, N-265
 Nocardimicin I, N-265
 Nocardimicin, N-265
 Nocathiacin III, N-266
 Nocathiacin II, N-266
 Nocathiacin I, N-266
 Nocathiacin IV, N-266
 Nocathiacin V, N-266
 Nocobactin NA, N-267
 Nodulisporic acid A, N-269
 Nodulisporic acid A₁, N-268
 Nodulisporic acid A₂, N-268
 Nodulisporic acid A₄, N-269
 Δ²³-Nodulisporic acid A₄, N-269
 Nodulisporic acid B, N-269
 Nodulisporic acid B₁, N-268
 Nodulisporic acid B₂, N-268
 Nodulisporic acid C, N-271
 Nodulisporic acid C₁, N-270
 Nodulisporic acid C₂, N-270
 Δ²³-Nodulisporic acid C₄, N-271
 Nodulisporic acid D, N-273
 Nodulisporic acid D₁, N-272
 Nodulisporic acid D₂, N-272
 Nodulisporic acid D₃, N-273
 Nodulisporic acid E, N-274
 Nodulisporic acid F, N-275
 Nodusmicin, N-276
 Noelaquinone, N-277
 Noformicin, N-278
 Nojirimycin B, A-728
galacto-Nojirimycin, A-726
 Nojirimycin, A-727
 Nokoensine, C-93
 Nomi-base I, H-172
 Nominine, H-172
 Nominine, N-279
 Nomofungin, C-582
 2,9-Nonadecadienoic acid pyrrolidide, B-271
N^b-Nonadecanoyltryptamine, T-639
 5-(12*Z*-Nonadecenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 5-(9*Z*-Nonadecenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 2-(3,6-Nonadienyl)-4-quinolinol, H-642
 2-(3,6-Nonadienyl)-4(1*H*)-quinolinone, H-642
 2,4-Nonadien-8-ynoic acid isobutylamide, N-280
 2,4-Nonadien-8-ynoic acid, N-280
 1,2,3,4,6,8,9,12,14-Nonahydroxydihydro-β-agarofuran, N-281
 Nonalupine, I-322
N-Nonanoyltyrosine, T-700
 2-Nonene-6,8-diyonic acid isobutylamide, N-282
 2-Nonene-6,8-diyonic acid, N-282
 2-Nonene-4,6,8-triynoic acid, N-283
 4-[(2-Nonene-4,6,8-triynoyl)amino]benzoic acid, N-283
N-(2-Nonene-4,6,8-triynoyl)valine, M-490
 3-(8-Nonenyl)-5-(1-propenyl)pyrrolizidine, N-284
 2-(1-Nonenyl)-4-quinolinol, H-642
 2-(1-Nonenyl)-4(1*H*)-quinolinone, H-642
 Nong-kang 101G, A-129
 Nonivamide, C-100
 Nonyl hexacosanoate, H-214
 3-(2-Nonyloxiranyl)-2-propenoic acid, T-124
 Nonylprodiginine, U-31
 Nonylprodigiosin, N-285
 3-Nonyl-1*H*-pyrazole, N-286
 1-(6-Nonyl-3-pyridinyl)-1-decanone, D-116
 2-Nonyl-3,4-quinolinediol, A-621
 2-Nonyl-4-quinolinol, H-642
 2-Nonyl-4(1*H*)-quinolinone, H-642
 Nor-16-acetoxybuxbenzamidienine, B-456
 Noracronine, A-120
 Noracronycine, A-120
 Noractinidine, N-287
 2'-Noradiantifoline, A-139
 Noradrenaline, N-298
 Noradrenalone, A-736
 Norajmaline, A-216
N-Noralkaloid Or₁, S-39
 Norallosedamine, S-210
 Noramurine, A-955
 Noranabasamine, A-963
 Noranantine, A-972
 Norandrobine, P-221
 Noranicanine, A-1038
 Norannuradhapurine, C-733
 18-Norantidesmone, A-1325

- Norarecoline, T-223
 Norargemonine, A-1410
 Noraristolochic acid, M-447
 Noraristolodione, C-273
 Norarmepavine, A-1444
 Norartrinal, N-298
 Norathosperminine, A-1532
 Noratropine, T-632
 Noraugustamine, A-1539
 Noravicine, A-1566
 Norbaeocystine, H-480
 Norbelladine, N-288
 2-*N*-Norberbamine, I-328
 2-Norberbamunine, B-98
 Norboldine, B-243
 Norbracteoline, B-276
 Norbulbocapnine, B-397
 Norbutanguinine, G-3
 30-Norbuxamine A, B-455
 Norbuxamine, B-455
N^b-Norbuxupapine, B-483
 Norcanelliline, A-1038
 Norcapsaicin, C-100
N-Norcarnequine, T-167
O-Norcarnequine, T-162
 Norcassaide, N-289
 Norcassaide 3-acetate, N-289
 Norcassaide, N-289
 Norcassaideine, C-178
 Norcassamide, N-299
 Norcassamide, N-299
 Norcassamidine (obsol.), N-299
 Norcassamidine, N-300
 Norcassamine (obsol.), N-299
 Norcataline, C-190
 Norcepharadione A, C-272
 Norcepharadione B, C-273
 2-Norcepharanoline, C-275
 2'-Norcepharanthine, C-275
 2-Norcepharanthine, C-275
 Norchalciporol, N-290
 Norchalciporol propionate, N-290
 Norchanoclavine I, C-348
 Norchanoclavine II, C-348
 Norchelerythrine, C-356
 (+)-Norchelidonine, C-357
 (-)-Norchelidonine, C-357
 Norcholine, D-730
 Nor-*N*^b-chondocurine, C-420
 Norcimiciphytine, C-441
 Norcinnamolarine, C-453
 Norcocaine, C-537
 2'-Norcocoline, I-339
N-Norcoculinine, C-552
 23-Norcona-5,18(22)-dienin-3-amine, C-610
 23-Norcona-4,6,18(22)-trienin-3-one, M-59
 23-Norcon-18(22)-enin-3-amine, C-610
 Norconessine, C-595
 Norcoralydine, X-30
 Norcoronatine, N-291
 Norcorydine, C-682
 Norcorypalline, T-158
 Norcotarnine, N-292
 Norcoumingide, N-289
 Norcularicine, C-804
 Norcularidine, C-805
 17-Norcuran, T-668
 Norcycleanine, C-835
 Norcyclomicrobuxine, C-882
N-Nordasycarpidone, D-84
 2'-*N*-Nordaurisoline, D-88
 19-Nor-4-dehydrocassaideine, N-293
 Nordehydrocycloclercitine, C-857
N-Nor-1,2-dehydroepistephanine, S-502
 4-Nor-7,8-dehydro-10-hydroxyskytanthine, N-294
 13-Nor-7(11)-dehydropilocarpine, N-295
N-Nordelavaine, D-181
 Nordelporphine, P-208
 Norden, O-63
 Nordeoxyharringtonine, H-66
 Nordercitin, D-93
 Nordesoxycabuffline, C-3
 Nordhagenine A, M-318
 Nordhagenine B, M-307
 Nordhagenine C, M-305
 (+)-Nordicentrine, D-346
 (-)-Nordicentrine, D-346
 8-Nordihydrocapsaicin, C-100
 Nordihydrocapsaicin, C-100
 Nordihydrotoxiferine I, T-436
 Nordihydrotoxiferine, T-436
 Nordine, N-296
 1-Nordistamycin A, D-866
 Nordomesticine, N-29
 Nordomoic acid, N-297
 Nordragabine, S-411
 Nordysidenin, D-972
 Norechitamine *N*-oxide, D-410
 Norechitamine, D-410
 Nor- ψ -ephedrine, A-869
 Norephedrine, A-869
 (\pm)-Nor- ψ -ephedrine, A-869
 (\pm)-Norephedrine, A-869
 (+)-Norephedrine, A-869
 (-)-Nor- ψ -ephedrine, A-869
 Norephedrone, A-870
 Norepilhachene, E-93
 Norepimalindine, M-73
 Norepinephrine bitartrate, N-298
 Norepinephrine, N-298
 Nor-Epirenan, N-298
 27-Norepothilone C, E-115
 Noreripinal, E-181
 Norerythrophlamine, N-299
 Norerythrophlamine 3-acetate, N-300
 Norerythrophlamine, N-300
 Norerythrosthachaldine 3-acetate, N-300
 Norerythrosthachaldine, N-300
 Norerythrosthachamide, N-299
 Norerythrosthachamine 3-acetate, N-300
 Norerythrosthachamine, N-300
 Norerythrosuamide, N-299
 Norerythrosuamide (obsol.), N-299
 Norerythrosuaveolide, N-289
 Norfissilandione, F-65
 Norfluorourarine *N*-oxide, N-301
 Nor-*C*-fluorourarine, N-301
 Norfumarine, F-174
 2'-Norfuniferine, R-114
 Norgeamide A, N-324
 Norgeamide B, N-324
 Norgeamide C, N-325
 Norgeamide D, N-325
 Norglaucine, T-246
 Norglaziovine, P-643
 Norgorchacoine, G-143
N-Nogramine, I-97
 Norgranatan-3-one, A-1580
 Norgraveoline, G-171
 2'-Norguattaguanine, G-212
 Norguattevaline, T-244
 Norharman, C-130
 Norharmancarboxylic acid, C-133
 Norharmane, P-916
N'-Norhernandezine, H-148
 Norhexahydromecambrine, L-214
 Norholadiene, H-314
 Norholadienine, H-314
 Norhygrine, H-759
 Norhyoscine, S-159
 Norhyoscyamine, T-632
 Norimeluteine, I-33
 Norine, C-601
 Norisoboldine, I-196
 2-Norisocepharanthine, C-275
 Norisocorydine, I-208
 Norisocorydione, S-367
 Norisocorymine, I-209
 Norisocorytuberine, T-248
 Norisodomesticine, N-29
 Norisoephedrine, A-869
 Norisohaplophytine, H-58
 Norisomalindine, M-73
 2'-Norisotetrandrine, I-328
 2-Norisotetrandrine, I-328
 Norisotryptoquivaline, N-315
 Norisotuboflavine, N-302
 Norisoyanangine, T-406
 Norjuziphine, J-63
 Norkurchamide, K-109
 Norlatifoline, L-56
 Norlaudanosine, L-69
 Norlaureline, L-72
 Norleobanidine, L-86
 Norleucoxylinone, L-159
 2'-Norlimacine, P-301
 2-Norlimacine, P-301
 2-Norlimacusine, L-175
 Norliridinine, T-521
 Norlirioferine, H-752
 Norlobelanidine, L-218
 Norlobelanine, L-218
 Norloline, L-234
 Norlusitanine, T-73
 15-Norlyngbyapeptin A, L-362
 Normacromerine, M-18
 18-Normacrophylline, M-10
 Normacrorine, I-44
 Normacusine A, P-562
 Normacusine B, T-425
 Normalindine, M-73
 Normaritudine, M-108
 (-)-Normaritudine, M-108
 Normavacurine, M-130
 Normaysine, M-134
 Normaytancyprine, M-135
 Normaytansine, M-135
 Normelicopicine, T-240
 Normelicopidine, X-2
O-Normelicopine, M-169
 Normelinone B, D-412
 Normelinone F, C-130
 Normeniarine, M-197
 Normetadrenaline, N-303
 Normetahemipinic acid, D-533
 Normetanephrine, N-303
 9-Normethylbudmunchiamine K, B-389
 1-Normethylbudmunchiamine K, B-389
 13-Normethylbudmunchiamine K, B-389
 Nor-*O*-methylkreysigine, P-221
 Normethylpseudoinidine, M-285
N-Normethylskytanthine, S-326
 Normicheline A, N-316
 Normitoridine, T-288
 Normycosporin-Gln, M-789
 Normantenine, N-29
 Normarceine, N-35
 Norneolitsine, N-125
 Norneronine, K-102
 Normicotine, P-957
 β -Normicotyrine, P-972
 Normicotyrine, P-972
 Normitidine, N-239
 Normorcapsaicin, C-100
 Normuciferidine, P-4
N-Normuciferine, D-531
O-Normuciferine, D-531
 2-Norobaberine 2'- β -*N*-oxide, O-1
 2'-Norobaberine, O-1
 2-*N*-Norobamegine, I-328
 Noroconovine, P-209
 Noroliveridine, O-85
 Noroliverine, O-85
 Noroliveroline, O-87
N-Nororientaline, O-111
N-Nororientinine, T-523
 Nororipavine, T-350
 Nororixine, O-113
 Noroxedrine, O-63
 2'-Noroxyacanthine, O-1
 Noroxyhydrastinine, I-305
 Norpachyconine, P-4
 Norpachystaudine, S-534
 2'-Norpakistanine, P-16
 Norpallidine, F-80
 Norpandamarilactonine A, P-63
 Norpandamarilactonine B, P-63
 Norpanurensine, P-76
 Norparagracine, P-737

- 22-Norparsonsine, P-97
 Norpenduline, T-286
 Norphedrin, A-864
 Norphen, O-63
 Norpholedrine, A-898
 N⁶-Norphysostigmine, P-400
 2'-Norpisopowiaridine, P-484
 Norpleiomutine, P-521
 9-Norpluviine, P-528
 10-Norpluviine, P-528
 Norpredicentrine, P-597
 Norpreocoteine, P-208
 Norprodigiosin, P-636
 Norprostaphabysine, P-658
 Norprotosinomenine, P-668
 Norpseudoephedrine, A-869
 Norpseudohygroline, H-760
 Norpseudoidinin, M-285
 Norpseudopelletierine, A-1580
 Norpseudotropine, A-1597
 Norpseudooanthoxanthin, P-737
 3-Norpseudooanthoxanthin, P-737
 Norpurpeline, T-288
 Norpurpureine, P-208
 Norquaternine, P-417
 Norrauvomitine, T-288
 Norreframidine, R-35
 Norregeline, K-36
 Norreserpine, R-52
 Norribasine, R-94
 Norrodiasine, R-114
 Norroecarboline, R-115
 Norroehybridine, R-116
 Norrufescine, N-304
 Norruspoline, N-305
 Norruspolinone, R-165
 Norsalutaridine, S-39
 Norsanguinarine, S-56
 Norsanguinine, G-3
 Norsarcocapnidine, S-73
 Norsarcocapnine, S-73
 4-(17-Norsarpagan-16-yl)-3-buten-2-one, D-396
 Norscopolamine, S-159
 Nor-6,7-secoangustilobine A, S-177
 Norsecocularidine, S-180
 Norsecocularine, S-180
 Norsecosarcocapnidine, S-198
 Norsecosarcocapnine, S-198
 a-Norsecurinan-11-one, N-307
 Norsecurinic acid, N-306
 Norsecurinine, N-307
 Norsedamine, S-210
 Norsegoline, N-308
 Norsenecaudatin 9-O-senecioate, N-309
 Norseredamine, T-288
 6-Norsetoclavine, S-273
 Norsinoacutine, S-39
 Norsongoramine, D-162
 Norsongorine, N-30
 Norsonodione, S-367
 Norspermidine, D-255
 Norspermine, B-154
 Norstaminolactone A, N-310
 Norstensusine, M-530
 Norstephalagine, S-536
 Norstephasubine, S-544
 Norsuaveoline, S-618
 Norsympathol, O-63
 Norsynephrine, O-63
 O-Nortecleanthine, T-56
 Nortehaunine, T-238
 Nortenuazonic acid, N-311
 Nortenuipine 2'-β-N-oxide, T-81
 (+)-Nortenuipine, T-81
 (-)-Nortenuipine, T-81
 2-Nortetrandrine, T-286
 Nortetraphyllicine, T-288
 11-Nortetrodotoxin-6R-ol, T-298
 11-Nortetrodotoxin-6S-ol, T-298
 11-Nortetrodotoxin, T-298
 Northalibrine, T-309
 Northalibrolone, T-309
 N⁶-Northalibrunine, T-311
 6-Northallicarpine, T-312
 Northallicarpine, T-312
 Northallicimine, O-27
 Northallicthuberine, T-318
 2-Northalidasine, T-320
 Northalifoline, I-305
 2'-Northaliphyllyne, T-313
 Northaliporphine, H-751
 2-Northalmine, T-341
 2-Northalrugosine, I-328
 Nortiliacorine A, T-406
 Nortiliacorinine A, T-407
 Nortiliacorinine B, T-407
 2'-Nortiliageine, R-114
 Nortopsentin A, N-312
 Nortopsentin B, N-312
 Nortopsentin C, N-312
 Nortopsentin D, N-312
 Nortopsentin D, N-313
 Nortrachelogenin amide, N-314
 Nortrewiasine, T-450
 Nortrilobine, I-339
 N-Nor-2,3,6-trimethoxymorphinandien-7-one, F-80
 Nortropacocaine, A-1597
 2,3-Nortropanediol, A-1581
 2,6-Nortropanediol, A-1582
 2,7-Nortropanediol, A-1583
 3,6-Nortropanediol, A-1584
 3,7-Nortropanediol, A-1585
 1,2,3,4,6-Nortropanepentol, A-1586
 1,2,3,4-Nortropanetetrol, A-1587
 1,2,3,6-Nortropanetetrol, A-1588
 1,2,4,7-Nortropanetetrol, A-1589
 1,2,3-Nortropanetriol, A-1590
 1,2,4-Nortropanetriol, A-1591
 1,2,7-Nortropanetriol, A-1592
 1,3,4-Nortropanetriol, A-1593
 2,3,4-Nortropanetriol, A-1594
 2,3,6-Nortropanetriol, A-1595
 3,6,7-Nortropanetriol, A-1596
 3-Nortropanol, A-1597
 Nortropanoline, A-1587
 Nortropan-3-yl phenylacetate, A-1597
 Nortropeine, A-1597
 Nortropenol, A-1597
 Nor-ψ-tropine, A-1597
 Nortropine, A-1597
 Nortropinyl cinnamate, A-1597
 Nortryptoquivaline, N-315
 Nortryptoquivalone, N-315
 (+)-Norushinsunine N-oxide, N-316
 Norushinsunine, N-316
 Noruvariopsamine, U-63
 Norveritol, A-898
 1-Norvincorine, V-125
 Norweberine, T-276
 Noryanangine, Y-5
 Norzoanthamine, N-317
 Norzoanthaminone, N-317
 Norzooanemonin, N-318
 Noscapine, N-44
 Nosiheptide, N-319
 Nosiheptine, N-319
 Nostocarboline, C-398
 Nostocine A, N-320
 Nostocyclamide M, N-322
 Nostocyclamide, N-321
 Nostodione A, N-323
 Nothapodytine A, M-96
 Nothapodytine B, M-96
 Notoamide A, S-153
 Notoamide B, S-153
 (+)-Notoamide B, S-153
 Notoamide C, N-324
 Notoamide D, N-325
 Notomycin A₁, C-711
 Notomycin A₂, C-712
 Novacine, I-12
 Noyaine, N-326
 NP 522, A-1218
 NP 101A, A-31
 NP 25302, B-242
 NP 25301, D-224
 NPAB, A-216
 NPI 0047, S-28
 NPI 0052, S-28
 NPS 1776, M-409
 NPTX 10, N-160
 NPTX 1, N-152
 NPTX 473, N-163
 NPTX 501, N-164
 NPTX 565, N-165
 NPTX 594, N-166
 NPTX 643, N-167
 NRCS 15, A-1219
 NSC 163501, A-81
 NSC 289487, A-1520
 NSC 403169, A-120
 NSC 51954, A-1202
 NSC 62786, A-152
 NSC 749, A-735
 NSC 76712, A-1044
 NSC 5355, B-101
 NSC 613012, B-60
 NSC 615291, B-229
 NSC 646846, B-60
 NSC 107412, C-711
 NSC 110326, C-634
 NSC 757, C-569
 NSC 94600, C-70
 NSC 107212, D-362
 NSC 169105, D-920
 NSC 185, D-765
 NSC 3096, D-92
 NSC 36354, D-92
 NSC 376128, D-907
 NSC 39147, D-765
 NSC 527604, D-142
 NSC 169774, E-157
 NSC 33669, E-72
 NSC 648766, E-37
 NSC 703147, E-117
 NSC 71795, E-67
 NSC 102811, F-131
 NSC 106486, F-132
 NSC 19893, F-104
 NSC 305263, F-152
 NSC 3590, F-121
 NSC 622398, F-13
 NSC 660649, F-13
 NSC 700409, F-13
 NSC 627434, G-79
 NSC 97925, G-108
 NSC 141633, H-66
 NSC 15780, H-673
 NSC 132319, I-68
 NSC 106995, L-241
 NSC 528004, L-162
 NSC 153858, M-135
 NSC 165014, M-135
 NSC 219970, M-135
 NSC 26980, M-654
 NSC 3051, M-463
 NSC 56410, M-654
 NSC 661755, M-589
 NSC 700647, M-54
 NSC 76779, M-230
 NSC 5366, N-44
 NSC 383228, O-209
 NSC 646282, O-197
 NSC 107654, P-961
 NSC 143095, P-891
 NSC 3073, P-772
 NSC 349156, P-58
 NSC 35443, P-79
 NSC 526046, P-904
 NSC 714361, P-42
 NSC 298223, R-4
 NSC 332598, R-83
 NSC 655649, R-32
 NSC 123124, S-340
 NSC 123125, S-340
 NSC 356894, S-394
 NSC 45383, S-569
 NSC 59729, S-384

- NSC 638850, S-498
 NSC 65346, S-53
 NSC 125973, T-52
 NSC 196520, T-450
 NSC 36080, T-284
 NSC 46451, T-284
 NSC 525816, T-74
 NSC 56408, T-658
 NSC 68075, T-312
 NSC 77037, T-286
 NSC 82260, T-74
 NSC 245467, V-110
 NSC 24567, V-110
 NSC 49842, V-110
 NSC 67574, V-128
 NSCL-96F037, P-338
 NSTX 3, N-181
 NU 2058, G-210
 Nuciferidine, P-4
 Nuciferine, D-531
 Nuciferoline, T-524
 (+)-Nuciferoline, T-524
 Nucleoside Q, Q-9
 Nudaurine, A-955
 (±)-Nudaurine, A-955
 Nudicaulamine, M-325
 Nudicaulidine, M-321
 Nudicaulin aglycone, N-327
 Nudicaulin III, N-327
 Nudicaulin II, N-327
 Nudicaulin I, N-327
 Nudicaulin IV, N-327
 Nudicaulin VIII, N-327
 Nudicaulin VII, N-327
 Nudicauline, G-91
 Nudifloric acid, H-711
 Nudiflorine, H-711
 Nuelin, T-360
 Nuevamine, N-328
 Numismine, N-329
 Nummularine A, M-720
 Nummularine B, A-942
 Nummularine C, N-330
 Nummularine D, I-154
 Nummularine E, N-331
 Nummularine F, N-332
 Nummularine G, N-333
 Nummularine H, J-51
 Nummularine K, N-334
 Nummularine M, N-335
 Nummularine N, N-336
 Nummularine O, J-52
 Nummularine P, N-337
 Nummularine R, N-338
 Nummularine S, P-29
 Nummularine T, A-942
 Nummularine, B-105
 Numoquin, Q-20
 Nuphacristine, N-339
 Nuphamine, N-340
 Nupharamine, N-340
 Nupharidine, D-227
 α-Nupharidine, N-341
 β-Nupharidine, N-342
 Nupharine, N-343
 Nupharolidine, D-227
 Nupharolutine, D-227
 Nupharopumiline, D-227
 Nupharpumilamine A, T-380
 Nupharpumilamine B, T-380
 Nupharpumilamine C, T-380
 Nupharpumilamine D, T-380
 Nuphenine, N-340
 Nupheine, T-380
 Nutrase, T-366
 Nutrestore, G-108
 Nutrifolin®, T-172
 Nuttallianine, M-323
 Nuttalline acetate, H-551
 Nuttalline, H-551
 Nuttingine A, N-344
 Nuttingine B, N-344
 Nuttingine C, N-345
 Nuttingine D, N-345
 Nuttingine E, N-345
 Nuttingine F, N-345
 Nybomycin, N-346
 Nyclin, P-897
 Nymphaedaline, N-347
 Nymphaeine, N-348
 Nympheine, N-348
 Obaberine, O-1
 Obamegine, I-328
 Obesine, O-2
 Oblongamine, O-1
 Oblongine, J-63
 Oblonginine, V-73
 Obovamine, O-3
 Obovanine, P-781
 Obovatine, B-127
 Obscuraminol A, A-848
 Obscuraminol B, A-782
 Obscuraminol C, A-783
 Obscuraminol D, A-784
 Obscuraminol E, A-783
 Obscuraminol F, A-783
 Obscuridine, O-4
 Obscurifoline, O-21
 α-Obscurine, O-5
 β-Obscurine, O-5
 Obscurine, O-6
 Obscurinervidine, O-7
 Obscurinervine, O-8
 Obscurinine, O-9
 Obscuroside C_{2z}, O-10
 Obscuroside C_{2b}, O-10
 Obscurumine A, L-325
 Obscurumine B, F-29
 Obtusine, O-11
 Obtusolactam, O-12
 Obyanamide, O-13
 Occidentaline, M-309
 Occidentaline, M-348
 Oceanalin A, O-14
 Oceanapamine, O-15
 Oceanapiside, D-292
 Oceanin, D-292
 Ochotensidine, C-700
 Ochotensimine, O-16
 Ochotensine, O-16
 Ochrindole A, O-17
 Ochrindole B, O-17
 Ochrindole C, O-17
 Ochrindole D, O-18
 Ochrobactin A, O-19
 Ochrobactin B, O-19
 Ochrobactin C, O-19
 Ochrobirine, O-20
 Ochrolifuanine A, O-21
 Ochrolifuanine B, O-21
 Ochrolifuanine C, O-21
 Ochrolifuanine D, O-21
 Ochrolifuanine E, O-21
 Ochrolifuanine F, O-21
 Ochrolifuanines, O-21
 Ochromianine, D-412
 Ochromianoxine, O-22
 Ochropamine, P-268
 Ochropine, P-268
 Ochroproposine, A-652
 Ochroproposine oxindole, O-23
 Ochroproposine, D-412
 Ochrosandwine, D-412
 Ocobotrine, I-318
 Ocodemerine, O-24
 Ocokryptine, O-25
 Ocominarine, P-210
 Ocominarone, O-26
 Oconovine, P-209
 Ocopodine, L-158
 Ocoteamine, O-1
 Ocoteine, O-27
 Ocotine, R-114
 Ocotominarine, H-243
 Ocotosine, O-28
 Ocoxyloine, L-159
 N^b-Octacosanoyltryptamine, T-639
 2,4-Octadecadienoic acid piperidine, O-29
 2,4-Octadecadienoic acid pyrrolidide, O-29
 3,6-Octadecadienoic acid pyrrolidide, P-949
 2,12-Octadecadienoic acid, O-30
 2,4-Octadecadienoic acid, O-29
 N-(9,12-Octadecadienyl)glutamic acid, G-107
 N²-(9,12-Octadecadienyl)glutamine, G-108
 N-(3,6-Octadecadienyl)pyrrolidine, P-949
 2,4,8,10,12,16-Octadecahexaenoic acid, O-31
 Octadecanamide, O-32
 Octadecanoic acid pyrrolidide, P-949
 Octadecanoic acid, O-32
 N-Octadecanoylpyrrolidine, P-949
 N^b-Octadecanoyltryptamine, T-639
 2,4,8,10,12-Octadecapentaenoic isobutylamide, O-31
 2,4,12-Octadecatrienoic acid isobutylamide, O-33
 2,4,12-Octadecatrienoic acid piperidine, O-33
 2,4,12-Octadecatrienoic acid, O-33
 2,4,14-Octadecatrienoic acid, O-34
 N-(9,12,15-Octadecatrienyl)glutamic acid, G-107
 N²-(9,12,15-Octadecatrienyl)glutamine, G-108
 1-(2,4,12-Octadecatrienyl)piperidine, O-33
 9-Octadecene-1,17-diamine, O-35
 9-Octadecenoic acid pyrrolidide, P-949
 2-Octadecenoic acid pyrrolidide, P-949
 9-Octadecenoic acid, O-37
 2-Octadecenoic acid, O-36
 N²-(9-Octadecenyl)glutamine, G-108
 N-(2-Octadecenyl)piperidine, O-36
 N-(2-Octadecenyl)pyrrolidine, P-949
 N-(9-Octadecenyl)pyrrolidine, P-949
 N-(11-Octadecenyl)tyrosine, T-700
 2-Octadecyl-1*H*-indol-3-ol, H-646
 5,6,11,12,13,14,15,16-Octadehydroaloperine, A-637
 2,4-Octadienamide, O-39
 2,4-Octadien-1-amine, O-38
 2,4-Octadienoic acid isobutylamide, O-39
 2,4-Octadienoic acid 2-phenylethylamide, O-39
 2,4-Octadienoic acid, O-39
 [4-[2-(2,5-Octadienylidene)-5-oxo-1-pyrrolidinyl]-butyl]guanidine, P-545
 [5-[2-(2,5-Octadienylidene)-5-oxo-1-pyrrolidinyl]-pentyl]guanidine, P-544
 1',3,4,4',5,5',6,6'-Octahydro-2,3'-bipyridine, O-40
 Octahydro-5-butyl-8-propyl-1*H*-indolizine, B-443
 Octahydro-3-butyl-5-propyl-1*H*-indolizine, O-41
 1,2,3,6*a*,7,8,9,12*a*-Octahydro-5,11-dimethoxy-1,7-dimethyl-6*a*,12*a*-bis(2-methylpropyl)-6,12-dioxo-1,7-diazadibenzo[*def,mmo*]chrysene, L-261
 Octahydro-2,2-dimethyl-1,3-dioxolo[4,5-*a*]indolizine, O-46
 Octahydro-5,7-dimethylindolizine, O-42
 Octahydro-6,9-dimethyl-4-(1-methylethyl)-2*H*-quinolizin-2-ol, P-526
 2-[Octahydro-4,7-dimethyl-1-oxocyclopenta[*c*]pyran-3-yl]nepetalactam, O-43
 Octahydro-2,4-dimethyl-2*H*-quinolizine, D-783
 1,2,3,6,7,8,9*a*-Octahydro-2,4-dimethylquinolizinium(1+), S-243
 Octahydro-5,8-dipropyl-1*H*-indolizine, O-44
 1,2,3,4,6,7,8,9-Octahydrodipyrido[1,2-*a*:1',2'-*c'*]imidazo[10-*ium*(1+)], O-45
 Octahydro-5*H*,10*H*-dipyrrolo[1,2-*a*:1',2'-*a'*]pyrazine-5,10-dione, C-898
 Octahydrohistrionicotoxin, H-309
 Octahydro-6-(4-hydroxy-2,5-dimethyl-5-octenyldene)-8-methyl-8-indolizino, P-793
 Octahydro-6-(6-hydroxy-2,5-dimethyl-4-octenyldene)-8-methyl-8-indolizino, P-791
 Octahydro-3-(7-hydroxy-2,5-dimethyl-4-octenyldene)-1-methyl-2*H*-quinolizin-1-ol, H-344
 Octahydro-2-hydroxy-2,4-dimethyl-α-phenylfuro[3,2-*b*]pyridine-5-ethanol, S-209
 3-[(Octahydro-2-hydroxyfuro[3,2-*b*]pyridin-2-yl)-methyl]-4(3*H*)-quinazolinone, I-232
 6-[Octahydro-8-hydroxy-1-(hydroxymethyl)-2*H*-quinolizin-3-yl]-2-piperidinone, C-8
 1-(Octahydro-7-hydroxy-8-indoliziny)-1-butanone, E-59

- Octahydro-8-hydroxy-6-isopropyl-1,4-dimethylquinolizine, P-526
- 9,10,13,14,15,16,20,20a-Octahydro-15-hydroxy-19-methoxy-2,6-dimethyl-9-phenylcyclopent[*f*]azacyclononadecine-7,17(8*H*,19*H*)-dione, A-1115
- 1,3,4,6,7,8,9,9a-Octahydro-4-(3-hydroxy-4-methoxyphenyl)-2-quinolizidinone, L-53
- 1,2,5,6,6a,7,8,12*c*-Octahydro-6a-hydroxy-7-methyl-14-methylene-4*H*-6,12*b*-ethanoindolizino[8,1-*cd*][1]benzazepin-13-one, G-141
- 1-(Octahydro-6-hydroxy-7-methyl-6-phenyl-5-indoliziny)-2-propanone, C-735
- Octahydro-6-(3-hydroxy-2-methylpropylidene)-8-methyl-8-indolizino, P-795
- 6-[Octahydro-1-(hydroxymethyl)-2*H*-quinolizin-3-yl]-2(1*H*)-pyridinone, M-79
- 3-(Octahydro-9-hydroxy-2-oxo-2*H*-quinolizin-3-yl)-4(3*H*)-quinazolinone, N-110
- Octahydro-2-hydroxy-2*H*-quinolizine-1-methanol, H-482
- 2,6,6a,7,8,9,10,10a-Octahydro-2-hydroxy-6,6a,8,10-tetramethyl-1*H*-[2]benzopyrano[4,3-*c*]pyridin-1-one, A-1226
- Octahydro-1,2-indolizinediol, O-46
- Octahydro-8-indolizinemethanol, O-47
- Octahydro-1,6,7,8-indolizinetetrol, C-188
- Octahydro-1,2,8-indolizinetriol, S-650
- 1,2,3,4,4a,5,6,12a-Octahydroindolo[1,2-*a*][1,8]naphthyridine-7-ethanol, G-142
- 1,2,3,4,6,7,12,12*b*-Octahydroindolo[2,3-*a*]quinolizine-4-carbonitrile, O-49
- 1,2,3,4,6,7,12,12*b*-Octahydroindolo[2,3-*a*]quinolizine-4-carboxylic acid, O-49
- 1,2,3,4,6,7,12,12*b*-Octahydroindolo[2,3-*a*]quinolizine, O-48
- 7,7a,8,9,10,11,13,14-Octahydro-7,14-methano-4*H*,6*H*-dipyrido[1,2-*a*;'2'-*e*][1,5]diazocin-4-one, A-970
- 1,2,3,5,6,12,13,13a-Octahydro-6,13-methano-10*H*-pyrido[1,2-*a*]pyrrole[1,2-*e*][1,5]diazocin-10-one, C-62
- 1,1',4,4',5,5',6,6'-Octahydro-5-methoxy-2,3'-bipyridine, O-50
- 2,6,6a,7,8,9,10,10a-Octahydro-2-methoxy-10-methyl-4-phenyl-6-(1-propenyl)-1*H*-[2]benzopyrano[4,3-*c*]pyridin-1-one, L-112
- 1,2,3,5,6,6a,12a,12*b*-Octahydro-11-methyl-12*H*-[1]benzopyrano[2,3-*g*]indolizin-12-one, E-56
- 3a,4,5,6,7,8,9,9a-Octahydro-9a-methyl-1*H*-1,6a-ethanopyrido[1,6-*de*]quinolizine, P-572
- Octahydro-5-(5-methyl-1,3,5-heptatrienyl)-6*bH*-cycloprop[*a*]oxireno[*g*]indolizin-6*b*-ol, I-101
- (Octahydro-8-methyl-5-indoliziny)-*sec*-heptanol, A-419
- 7-(Octahydro-8-methyl-5-indoliziny)-3-hepten-2-ol, H-127
- (Octahydro-8-methyl-5-indoliziny)-*sec*-nonenol, A-487
- Octahydro-8-methyl-6-(2-methylhexylidene)-8-indolizino, A-480
- Octahydro-8-methyl-6-(2-methylpentylidene)-8-indolizino, A-359
- Octahydro-8-methyl-6-(2-methylpropylidene)-7,8-indolizinediol, P-795
- Octahydro-8-methyl-6-(2-methylpropylidene)-8-indolizino, P-795
- Octahydro-5-methyl-10-(6,8-nonadienyl)-1*H*-pyrrolo[1,2-*a*]azepine, A-367
- Octahydro-5-methyl-10-(8-nonenyl)-1*H*-pyrrolo[1,2-*a*]azepine, A-367
- Octahydro-8-methyl-5-(6-nonen-8-ynyl)-1*H*-indolizine, O-51
- Octahydro-5-methyl-10-(8-nonyl)-1*H*-pyrrolo[1,2-*a*]azepine, A-367
- Octahydro-4-methyl-6-(1,3-octadienyl)-2*H*-quinolizin-3-ol acetate, P-418
- Octahydro-5-methyl-10-(oxo-8-nonyl)-1*H*-pyrrolo[1,2-*a*]azepine, A-367
- Octahydro-8-methyl-5-(4-pentenyl)-1*H*-indolizine, O-52
- Octahydro-8-methyl-5-(2-penten-4-ynyl)-1*H*-indolizine, O-52
- Octahydro-1-methyl-4-(2-penten-4-ynyl)-2*H*-quinolizine, M-507
- Octahydro-8-methyl-5-pentylindolizine, O-52
- Octahydro-8-methyl-5-(4-pentynyl)-1*H*-indolizine, O-52
- 1,4,4a,5,6,7,8,8a-Octahydro-5-methyl-2-propylquinoline, T-204
- Octahydro-6-methyl- α -propyl-2*H*-quinolizine-4-ethanol benzoate, P-571
- Octahydro-4-methyl-6-propyl-2*H*-quinolizine, M-532
- Octahydro-4-methyl-2*H*-quinolizin-2-one, O-53
- Octahydro-4-methyl-1-[[[4,8,12-trimethyl-1-oxo-2,7,11-tridecatrienyl]amino]methyl]indolizinium(1+), S-504
- 1,2,3,4,6,7,8,9-Octahydrophenazine, O-54
- 4,5,6,7,8,9,12,13-Octahydro-8-phenyl-2,13-methano-2*H*-2,7,11-benzotriazacyclopentadecine-1,10(3*H*,11*H*)-dione, C-853
- 2,3,6,7,8,9,10,11-Octahydro-6-phenyl-1*H*-[1,5,9]triazacyclotridecino[2,1-*a*][2]benzazepine-4,13(5*H*,19*bH*)-dione, P-523
- Octahydro-3-(2-piperidinyl)-2*H*-quinolizine-1-carboxylic acid, A-1335
- Octahydro-3-(2-piperidinyl)-2*H*-quinolizine, L-97
- Octahydro-5-propylindolizine, O-56
- Octahydro-3-propylindolizine, O-55
- Octahydro-1*H*,3*H*-pyrido[3,2,1-*ij*]benzoxazine, O-57
- Octahydropropocoll, C-898
- Octahydro-2*H*-quinolizin-1-amine, A-907
- Octahydro-2*H*-quinolizine-1-methanol, O-58
- 5-[[[Octahydro-2*H*-quinolizin-1-yl)methyl]amino]-5-oxopentanoic acid, O-59
- 1-[[Octahydro-2*H*-quinolizin-1-yl)methyl]-2,6-piperidinedione, L-26
- 2-(Octahydro-2*H*-quinolizin-3-yl)-1-piperidinecarboxaldehyde, L-97
- 7-[(1,2,3,4,4a,7,8,8a-Octahydro-1,2,4a,5-tetramethyl-1-naphthalenyl)methyl]-6-benzoxazolol, O-60
- 1,2,2a,10,10c,11,11a,11*b*-Octahydro-2a,4,11,11-tetramethyl-3-oxa-10-azacyclobut[3,4]indeno[5,6-*a*]fluorene, B-122
- 1,2,2a,10,10c,11,11a,11*b*-Octahydro-2a,7,11,11-tetramethyl-3-oxa-10-azacyclobut[3,4]indeno[5,6-*a*]fluorene, B-121
- 2,6,6a,7,8,9,10,10a-Octahydro-6,8,10-trimethyl-1*H*-[2]benzopyrano[4,3-*c*]pyridin-1-one, A-1214
- Octahydro-2,4,7-trimethyl-1*H*-cyclopenta[*c*]pyridin-7a-ol, H-734
- Octahydro-2,4,7-trimethyl-1*H*-cyclopenta[*c*]pyridine, S-326
- Octahydro-2,4,7-trimethyl-1*H*-cyclopenta[*c*]pyridin-4a-ol, H-733
- 2,6,7,10,11,14,15,27-Octahydroxydotriacontanoic acid, O-61
- Octalupine, H-554
- 5-Octanoylamino-2,3,6(1*H*)-pyridinetrione, O-62
- N*-Octanoylnornicotine, P-957
- N*-Octanoyltyrosine, T-700
- 2-(1-Octenyl)-4(1*H*)-quinolinone, H-648
- Octopamine, O-63
- 2-Octylpyrrole sulfamate, O-64
- 2-Octylpyrrole, O-64
- 2-Octyl-3,4-quinolinediol, A-621
- 2-Octyl-4-quinolinol *N*-oxide, H-648
- 2-Octyl-4-quinolinol, H-648
- 2-Octyl-4(1*H*)-quinolinone *N*-oxide, H-648
- 2-Octyl-4(1*H*)-quinolinone, H-648
- Ocucarpine, P-424
- Ocular age pigment A2-E, O-65
- Ocusert P 20, P-424
- Ocusert Pilo 40, P-424
- Odiline, O-66
- Odoratin, T-283
- Odorine, O-67
- (+)-Odorinol, O-67
- (-)-Odorinol, O-67
- Oduline, O-68
- Oduocine, O-143
- Off, M-398
- OGT 2378, H-616
- OGT 918, H-616
- Ohyamycin, F-132
- Okaramine A, O-68a
- Okaramine B, O-69
- Okaramine C, O-70
- Okaramine D, O-69
- Okaramine E, O-71
- Okaramine F, O-71
- Okaramine G, O-70
- Okaramine H, O-72
- Okaramine I, O-68a
- Okaramine J, O-73
- Okaramine K, O-73
- Okaramine L, O-70
- Okaramine M, O-74
- Okaramine N, O-75
- Okaramine O, O-75
- Okaramine P, O-76
- Okaramine Q, O-69
- Okaramine R, O-76a
- Okolasine, W-16
- Olamine, A-767
- Oldhamactam, T-254
- Oldhamine A, O-77
- Oldhamiphylline A, O-78
- Oldhamlactam, T-254
- Oleamide, O-37
- Oleficin, O-79
- Oleic acid, O-37
- Oleioidine, Q-1
- N*²-Oleoylglutamine, G-108
- Oleracein A, C-862
- Oleracein B, C-862
- Oleracein C, C-862
- Oleracein D, C-862
- Oleracein E, T-629
- Oleracins, O-80
- Oliganine, O-81
- Oligophylcidine, O-82
- Oligophylicine, O-82
- Oligophyllidine, T-520
- Oligophylline, O-82
- Olivacine *N*-oxide, O-83
- Olivacine, O-83
- Oliveramine, O-84
- Oliveridine *N*-oxide, O-85
- Oliveridine, G-61
- Oliveridine, O-85
- Oliverine *N*-oxide, O-85
- Oliverine, O-85
- Oliverine, O-86
- Oliveroline β -*N*-oxide, O-87
- Oliveroline, O-87
- Olividine, M-359
- Olivimine, M-359
- Olivoretin A, T-63
- Olivoretin B, T-63
- Olivoretin C, O-88
- Olivoretin D, T-63
- Olivoretin E, O-89
- Olomoucine, O-90
- OM 704A, D-295
- OM 704B, D-295
- OM 5714, I-161
- Omacetaxine mepesuccinate, H-66
- Omadine disulfide, D-831
- Omadine MDS, D-831
- Omadine-DS, D-831
- Omain, D-92
- OMDS, D-831
- Omeienine, M-309
- Ommatin D, D-527
- Omomycin, I-64
- Oncinotine, O-91
- Oncinotin-11-one, O-91
- Oncinotin-12-one, O-91
- Oncodine, D-625
- Oncovin, V-128
- Onetine, O-92
- Onkovin, V-128
- Onosmin A, A-706

- Onosmin B, A-706
 Onychine, O-93
 Oocydin A, H-76
 OPC 15161, A-1220
 Ophiocarpine *N*-oxide, O-94
 Ophiocarpine, O-94
 Ophiocarpinone, O-94
 Ophiocardin, B-9
 Ophiorine A, O-95
 Ophiorine B, O-95
 Ophiorrhizine-12-carboxylate, O-97
 Ophiorrhizine, O-96
 Ophiuroidine, O-98
 OPHR 17, O-99
 OPHR 23, O-99
 Ophthocillin, X-7
 Opianine, N-44
 Oppositine A, P-251
 Oppositine B, P-251
 Optochin, Q-20
 Optoquine, Q-20
 Optoquinic acid, H-726
 Opuntine B, O-100
 Oramorph SR, M-704
 Orantine, A-1334
 Orbiculamide A, O-101
 Orcein, O-102
 Orchil, O-102
 Oreaconine, L-317
 Orellanine 4,4'-diglucoside, T-249
 Orellanine II, T-249
 Orellanine, T-249
 Orelline, T-249
 Orellinine, T-249
 Orensine, A-138
 Oreobeiline, O-103
 Oreodine, G-87
 Oreogenine, G-87
 Oreoline, O-104
 Oreoline, O-109
 Oreonone, F-2
 Oreophiline, M-140
 Orgetine, H-210
 Oriamide, O-105
 Oriciacridone A, O-106
 Oriciacridone B, O-106
 Oriciacridone C, R-167
 Oriciacridone D, O-107
 Oriciacridone E, T-581
 Oriciacridone F, O-108
 Oricine, F-90
 Oridine 2, O-109
 Oridine, O-109
 Orientalidine, O-110
 Orientaline, O-111
 Orientalinone, O-112
 Orientidine, T-525
 Orientine, T-523
 Orientinine, T-549
 Orientinine, T-523
 Orientmycin, A-832
 Orientomycin, A-832
 Orijanone, E-40
 Orinapterin, A-749
 Oripavidine, T-350
 Oripavine, T-350
 Orixalone A, T-583
 Orixalone B, T-583
 Orixalone C, T-583
 Orixalone D, F-90
 Orixiarine, E-40
 Orixine, O-113
 Orixinone, O-113
 Ormocastrine, P-529
 Ormojanine, O-114
 Ormojine, O-117
 Ormosajine, P-53
 Ormosanine, O-115
 21-Ormosanin-20-ylpanamine, O-117
 Ormosin VI, T-52
 Ormosine, O-116
 Ormosinine, O-117
 Ornamine, O-118
 Ornazamine, O-118
 Ornazidine, O-118
 Ornibactin C4, O-119
 Ornibactin C6, O-120
 Ornibactin C8, O-121
 Ornibactin F, O-121
 Ornithine lactam, A-872
 D-Ornithinegalantin I, G-6
N-(*N*²-L-Ornithyl)cantharimide, C-89
 Orobanchamine, O-122
 Oroboidine, H-554
 Orochrine, H-198
 Oroidin, O-123
 Orosomycin, O-124
 Orotamide, D-669
 Orotic acid dimethylamide, D-669
 Orotic acid, D-669
 Orotosans S, D-669
 Orthidine A, O-125
 Orthidine B, O-125
 Orthidine C, O-126
 Orthidine D, O-126
 Orthidine E, O-127
 Orthidine F, S-399
 Orthocentrine, M-376
 Orthosphenine, P-251
 Oryzanin, T-366
 OS 1998, S-53
 Oscillatorin, O-128
 Oscine atropate, A-1364
 Oscine tropate, S-159
 Oscine, S-160
 Osmaronin epoxide, H-576
 Osmaronin, H-576
 Osoic acid, O-129
 Osornine, O-130
 Ostreogrycin A, O-131
 Ostreogrycin G, O-131
 Ostreogrycin, O-131
 Ostreirine A, O-132
 Otanthus acid piperidide, O-133
 Otanthus acid, O-133
 Otanthusic acid 2,3-dehydropiperidide, O-133
 Otanthusic acid, O-133
 Oteromycin, O-134
 Othonnine, A-1572
 Othosenine, O-137
 Otocamine, O-135
 Otonecine, O-136
 Otosenine, O-137
 Otrun, D-730
 Oubatchensine, O-138
 Ouregidione, O-139
 Oureguattidine, T-244
 Oureguattine, T-244
 Ourouparine, O-140
 Ouvrardiane A, O-141
 Ouvrardiane B, O-141
 Ouvrardiantine, M-333
 Ovalin, H-697
 Ovatine, V-45
 Ovigeridimerine, O-142
 Ovigerine, O-143
 Ovihermangerine, O-144
 Oviisocorydine, O-145
 Ovisot, A-47
 Owerrine, V-188
 7-Oxa-3-azabicyclo[4.1.0]heptan-2-one, O-146
 2-Oxa-5-azabicyclo[2.2.2]octan-3-one, H-698
 1-Oxa-4-azacyclohexane, M-705
 Oxaceprol, H-715
 Oxachelin, O-147
 13-Oxa-4,20-diazatetracyclo[12.2.2.2^{3,6}.1^{8,12}]heneicos-8,10,12(19),14,16,17-hexaen-11-ol, P-444
 OXAIDS, H-616
 12,28-Oxaicinal A, I-174
 Oxalicine A, O-148
 Oxalicine B, O-148
 Oxaline, M-162
*N*²-Oxalylarginine, A-1413
 12,28-Oxamanzamine A, M-91
 12,34-Oxamanzamine A, M-91
ent-12,34-Oxamanzamine E, M-91
 12,28-Oxamanzamine E, M-91
 12,34-Oxamanzamine E, M-91
ent-12,34-Oxamanzamine F, M-91
 Oxamphetamine, A-898
 Oxamycin, A-832
 Oxandrine, O-149
 Oxandriline, O-149
 Oxaphyllidine, A-1336
 Oxaphylline, H-405
 Oxasetin, O-150
 Oxathinecarboxylic acid, O-151
 2*H*-1,4-Oxazine-3-carboxylic acid, O-152
 Oxazine 1, O-153
 Oxazine 2, O-153
 Oxazine 3, O-154
 Oxazine 4, O-153
 Oxazine 5, O-154
 Oxazine 6, O-154
 Oxazolomycin, O-155
 Oxazolomycin B, O-155
 Oxazolomycin C, O-155
 4-(4-Oxazolyl)-3-(hydroxymethyl)-3-buten-2-ol, O-156
 3-(5-Oxazolyl)-1*H*-indole, O-157
 2-[(4-Oxazolyl)methylene]-1,3-butanediol, O-156
 5-(4-Oxazolyl)-4-pentene-2,3-diol, M-159
 Oxedrine, S-659
 Oxepinamide A, O-158
 Oxepinamide B, O-158
 Oxepinamide C, O-159
 Oxerine, O-160
 Oxeze, X-5
 Oxiamine, I-148
 3,19-Oxidocoronaridine, H-270
 6,7-Oxidodeoxynupharidine dimer, O-161
 3,6-Oxidoibogaine, O-162
 3,6-Oxidoiboxygaine, O-162
 5,15-Oxidolycopodane, O-163
 Oximidine III, O-164
 Oximidine II, O-164
 Oximidine I, O-164
 Oxindole-3-acetic acid, D-485
 Oxindole, I-103
 3-Oxiranyl-1*H*-indole, O-165
 8-Oxoacetylcorynoline, C-692
 22-Oxoacuminatine, H-403
 8-Oxoadenine, A-819
 3-Oxoadociaquinone B, A-147
 14-Oxoeroophobin 2, A-155
 11-Oxoeroothionin, A-158
 6-Oxoalstophyllal, A-656
 6-Oxoalstophylline, A-649
 18-Oxoamataine, A-676
 7-Oxoanhydromethylpseudolycorine, A-1508
 Oxanolobine, L-40
 19-Oxoanthranoyllycoctonine, H-123
 Oxaphyllidine, A-1336
 16-Oxoaspergillidine, A-1486
 18-Oxoaspidoalbine, D-144
 Oxoasoanine *N*-oxide, A-1508
 Oxoasoanine, A-1508
 2'-Oxoasterriquinol D di-Me ether, A-1519
 3-Oxoaxisonitrile 3, S-425
 β-Oxo-9-azabicyclo[4.2.1]non-2-ene-2-propanoic acid, C-143
 4-Oxo-1-azabicyclo[3.3.0]octa-5,7-diene-2-carboxylic acid, D-488
 Oxobaicaline, O-166
 8-Oxoberberrubine, C-73
 6-Oxo-7-biopterin, A-748
 α-Oxobrowniine, M-309
N-(3-Oxo-1-butenyl)-1,2-dihydroenerione, R-45
 6-(3-Oxo-1-butenyl)-1*H*-indole, I-118
 3-(3-Oxo-1-butenyl)-1*H*-indole, I-117
N-(3-Oxobutyl)cytosine, C-938
 8-(2-Oxobutyl)dihydrochelerythrine, O-167
N-(3-Oxobutyl)mitomycin C, M-654
 2-(2-Oxobutyl)piperidine, H-435
 9-(3-Oxobutyl)-6,8-purinediol, P-818
 1-(3-Oxobutyl)-6,8-purinediol, P-818
 Oxobuxifoline, O-168
 Oxocalycinine, O-183
 8-Oxocanadine, C-73

- 10-Oxocantrine, C-76
 8-Oxocephalotaxine, C-271
 8-Oxochelethrine, D-407
 8-Oxococcuine, E-202
 8-Oxococcuvinine, E-202
 Oxocompostelline, O-169
 1'-Oxoconiine, C-604
 3'-Oxoconodurine, C-616
 19-Oxoconopharyngine, C-620
 3-Oxoconopharyngine, C-620
 8-Oxocoptisine, D-409
 5-Oxocoronaridine, C-657
 6-Oxocoronaridine, C-657
 3-Oxocoronaridine, C-657
 8-Oxocorynoline, C-692
 Oxocoryrutine, C-698
 Oxocrebanine, O-170
 Oxocrinine, C-748
 13-Oxocryptopine, C-790
 14-Oxocryptopine, C-790
 Oxocularicine, O-169
 Oxocularidine, O-171
 Oxocularine, O-171
 17-Oxocycloprotobuxine, O-172
 Oxocyclostylidol, O-173
 5-Oxoeylindrocarpidine, C-929
 10-Oxocylindrocarpidine, C-929
 11-Oxocytisine, C-938
 7-Oxodanaidal, D-14
 Oxodaphnigraciline, D-50
 Oxodaphnigracine, D-51
 6-Oxo-2,4-decadienoic acid piperidine, O-174
 6-Oxo-2,4-decadienoic acid, O-174
N-(6-Oxo-2,4-decadienyl)piperidine, O-174
 1-(1-Oxo-2,4-decadienyl)piperidine, D-104
 1-(1-Oxo-2,4-decadienyl)pyrrolidine, D-104
*N*²-(4-Oxodecanoyl)histamine, H-307
 19-Oxodecarbomethoxytetrahydrosecodine, D-120
 3-(9-Oxo-1,3,5,7-decatetraenyl)pyridine, P-908
 7-Oxodehydroasimilobine, L-193
 8-Oxo-9-dehydrohobartine, H-310
 8-Oxo-9-dehydromakomakine, H-310
 3-Oxo-14,15-dehydrosazininilam, R-76
 2-Oxo- Δ^2 -dehydrosparteine, D-161
 16-Oxodelavaine, D-181
 19-Oxodelphatine, H-123
 α -Oxodelphinine, M-312
 2-Oxendrobine, D-210
 Oxodesmosine, O-175
 Oxodicentrine, D-347
 4-Oxo-2,3-didehydrosparteine, M-731
 6-Oxo-16,20-diepisilicine, M-293
 19-Oxodihydroatisine, A-1533
 1-Oxo-1,2-dihydro-*N*-demethyl-*N*-hydroxyacronycine, D-417
 1-Oxo-1,2-dihydro-*N*-desmethylacronycine, D-417
 8-Oxodihydroepiberberine, O-177
 Oxodiscoguatine, O-183
 12-Oxo-2,4,8,10-dodecatetraenoic acid, O-176
 2-Oxodopamine, A-736
 Oxoduocine, H-150
 19-Oxoeburnamine, E-6
 1-(1-Oxo-2,4-eicosadienyl)piperidine, E-45
 1-(1-Oxo-2,4-eicosadienyl)pyrrolidine, T-491
 13-Oxoellipticine *N*⁶-oxide, E-67
 13-Oxoellipticine, E-67
 8-Oxoepiberberine, O-177
 3-Oxo-19-epiheyanine, C-657
 6-Oxo-16-episilicine, M-293
 Oxoepesthamiersine, S-537
 Oxoepesthamine, E-102
 3-Oxo-19-epivoacristine, V-170
 9-Oxoepothilone D, E-117
 8-Oxoerymelanthine, E-198
 8-Oxoerysodine, E-202
 11-Oxoerysodine, E-206
 11-Oxoerysopine, E-206
 8-Oxoerysotrine, E-202
 11-Oxoerysovine, E-206
 8-Oxoerythraline 1,2-epoxide, E-205
 8-Oxoerythraline, E-205
 11-Oxoerythraline, E-205
 8-Oxoerythrine, E-205
 8-Oxo- α -erythroidine 1,2-epoxide, E-215
 8-Oxo- α -erythroidine, E-215
 8-Oxo- β -erythroidine, E-215
 Oxofangchirine, D-148
 11-Oxofistularin 3, F-70
 Oxogambirtannine, D-434
 21-Oxogelsemine, G-49
 19-Oxogelsemine, G-44
 21-Oxogelsevirine, G-49
 Oxoglucine, O-178
 8-Oxoguanine, A-753
 18-Oxohaplocidine, D-144
 16-Oxohasubanonine, H-70
 3-(6-Oxoheptyl)-5-propylpyrrolizidine, H-500
 7-Oxohernagine, G-93
 7-Oxohernandaline, O-179
 7-Oxohernangerine, O-180
 1-(1-Oxo-2,7-hexadecadien-10-ynyl)-1*H*-pyrrole, H-218
 1-(1-Oxo-2,7-hexadecadien-10-ynyl)pyrrolidine, H-218
 1-(1-Oxo-6,8-hexadecadien-10-ynyl)pyrrolidine, H-219
 1-(1-Oxo-7-hexadecen-10-ynyl)-1*H*-pyrrole, H-225
 6-Oxo-2,4-hexadienoic acid, O-181
 1-(1-Oxo-2,7-hexadienyl)-1*H*-pyrrole, H-217
 1-(1-Oxoheptyl)-2-(3-pyridinyl)pyrrolidine, P-957
 18-Oxoholadienine, H-314
 11-Oxohomoaerotherionin, A-158
 19-Oxohumantentione, G-44
 3-Oxohydroxykopsinine, K-78
 8-Oxohypecorinine *N*-oxide, H-766
 8-Oxohypecorinine, H-766
 2-Oxo-4-imidazolinecarboxylic acid, D-484
 α -Oxo-1*H*-indole-3-acetamide, O-182
 α -Oxo-1*H*-indole-3-acetic acid, O-182
 α -Oxo-1*H*-indole-3-acetonitrile, O-182
 3-Oxoindole, I-106
 α -Oxo-1*H*-indole-3-propanoic acid, I-133
 2-(2-Oxo-3-indolyl)acetic acid, D-485
 Oxoisocalycine, O-183
 8-Oxoiscorypalmine, I-210
 10-Oxo- β -isosparteine, I-322
 3-Oxoisoavocangine, I-344
 5-Oxo-2(5*H*)-isoxazoleacetic acid, O-184
 5-Oxo-2(5*H*)-isoxazolepropanenitrile, O-185
 5-Oxo-2(5*H*)-isoxazolepropanoic acid, O-185
 2-(3-Oxo-5-isoxazolidiny)glycine, T-490
 Oxojanussine, O-186
 3''-Oxojuliflorine, J-56
 3-Oxojuliprosine, J-57
 3''-Oxojuliprosopine, J-56
 3'-Oxojuliprosine, J-57
 10-Oxokibdelone A, K-40
 10-Oxokopsanone, D-805
 Oxokrigenamine, K-102
 Oxolaureline, O-187
 6-Oxoleuconoxine, L-156
 21'-Oxoleurosine, L-162
 Oxolucidine A, L-277
 Oxolucidine B, L-277
 6-Oxolumazine, D-494
 7-Oxolumazine, P-382
 7-Oxolumazine, P-763
 17-Oxolupanine, O-188
 14-Oxolyalosite, P-135
 Oxomaritidine, M-108
 3-Oxomehranine, M-153
 8-Oxo-11 β -methoxyerythraline, E-205
 6-Oxomethuenine, M-293
 18-Oxo-*O*-methylaspidalbine, D-144
 Oxo-*O*-methylpukateine, O-204
 3-Oxo-*N*^b-methyl-3,4-secoakummidine, V-171
 3-Oxominovincine, O-210
 13-Oxomuramine, M-734
 14-Oxonaamidine G, N-1
 Oxonanthenine, O-195
 1-(1-Oxo-2,9-nonadecadienyl)pyrrolidine, B-271
 3-(1-Oxo-4,7-nonadienyl)oxiranecarboxamide, C-286
 2-(8-Oxononyl)-4(1*H*)-quinolinone, H-642
 19-Oxonorcassidine, N-300
 Oxonuciferine, L-193
 5-Oxo-6,8-octadecadienoic acid, O-189
 9-Oxo-12,15-octadecadienoic acid, O-190
 13-Oxo-9,11-octadecadienoic acid, O-191
 12-Oxo-octadecanoic acid, O-192
 9-Oxo-12-octadecenoic acid, O-193
 12-Oxo-2-octadecenoic acid, O-194
 1-(1-Oxo-2-octadecenyl)piperidine, O-36
 1-(1-Oxo-octyl)-2-(3-pyridinyl)pyrrolidine, P-957
 Oxopachycarpine, O-208
 3-Oxopachysiphine, O-210
 16-Oxoparaheerquamide A, P-90
 2-(2-Oxopentyl)piperidine, H-37
N-(3-Oxo-3*H*-phenoxazin-2-yl)acetamide, A-862
 9-Oxo-*N*-(2-phenylethyl)hexadecanamide, P-342
 4-Oxo-6-(phenylmethyl)-4*H*-pyran-3-carboxylic acid, B-92
N-[4-[(1-Oxo-3-phenyl-2-propenyl)amino]butyl]-benzamide, C-456
 1'-(1-Oxo-3-phenyl-2-propenyl)-2,3'-bipiperidine, A-1525
 9-(1-Oxo-3-phenyl-2-propenyl)-4-phenyl-1,5,9-triazacyclotridec-12-en-2-one, P-265
 1-(1-Oxo-3-phenyl-2-propenyl)-1*H*-pyrrole, C-457
 1-(1-Oxo-3-phenyl-2-propenyl)pyrrolidine, C-458
 1-(1-Oxo-3-phenyl-2-propenyl)-2-pyrrolidinone, P-956
 1-(1-Oxo-3-phenylpropyl)-1*H*-pyrrole, P-351
 11-Oxophlegmariurine B, P-363
 2-Oxophlegmariurine B, P-363
 Oxophoebine, O-195
 4-Oxopiperic acid, O-196
 4-Oxo-2-piperidinecarboxylic acid, O-196
 8-[(2-Oxo-3-piperidinyl)amino]-8-oxooctyl 5,9-anhydro-2,3,8-trideoxy-8-(5-hydroxy-4-methyl-2-hexenyl)-3-methyl-DL-glycero-LD-*allo*-non-2-enoate, O-197
 8-Oxopolythiane, S-543
 6-Oxoprimapterin, A-748
 5-Oxopropine, O-206
 16-Oxoprometaphanine, P-639
 Oxopropaline A, O-198
 Oxopropaline B, O-198
 Oxopropaline D, O-198
 Oxopropaline E, O-198
 Oxopropaline G, O-198
 3-(2-Oxopropyl)conodurine, T-3
 5-(2-Oxopropyl)coronaridine, O-200
 3-(2-Oxopropyl)coronaridine, O-199
 3,3'-(Oxopropyl)diconaridine, O-201
 6-(2-Oxopropyl)-6-pentyl-2-piperidinone, A-136
 6-(1-Oxopropyl)-2,4(1*H*,3*H*)-pteridinedione, H-702
 15-(2-Oxopropyl)reframidine, R-35
 3-(2-Oxopropyl)voacangine, O-199
 8-Oxoprotopine, P-667
 13-Oxoprotopine, P-667
 14-Oxoprotopine, P-667
 8-Oxopseudopalmitine, O-202
 Oxopukateine, O-204
 Oxopurpureine, O-203
 Oxoputerine, O-204
 2-Oxo-3-(4-pyridinyl)propanoic acid, O-205
 4-Oxo-4-(9*H*-pyrido[3,4-*b*]indol-1-yl)butanoic acid, C-137
 2-Oxo-1-pyrrolidinecarboxamide, S-487
 5-Oxo-2-pyrrolidinecarboxylic acid, O-206
 2-Oxopyrrolidine, P-956
 6-(5-Oxo-2-pyrrolidinyl)eriodictyol, T-572
 8-(5-Oxo-2-pyrrolidinyl)eriodictyol, T-573
N-(2-Oxo-1-pyrrolidinylmethyl)cytisine, C-938
 6-(5-Oxo-2-pyrrolidinyl)naringenin, T-572
 8-(5-Oxo-2-pyrrolidinyl)naringenin, T-573
 5-[(5-Oxo-2-pyrrolidinyl)oxy]methyl]-2-furancarboxaldehyde, S-269
 15-Oxorefraxidine, R-35
 3-Oxorhazinilam, R-76
 Oxosanguinarine, D-510
 Oxosarcocapnine, S-73
 Oxosarcocapnine, S-73
 Oxosarcophylline, O-207
 8-Oxoschelhammeridine, S-125
 11 α -Oxoschelhammeridine, S-125

- 12-Oxoschellhammeridine, S-125
6-Oxosilicine, M-293
23-Oxosolacongostidine, S-337
24-Oxosolacongostidine, S-337
15-Oxosoladulcidine, S-441
10-Oxosparteine, A-1337
2-Oxo-11 α -sparteine, L-302
Oxosparteine, O-208
17-Oxosparteine, O-208
13-Oxosparteine, T-361
11-Oxospiradine D, S-421
7-Oxostaurosporine, S-498
12-Oxostearic acid, O-192
2-Oxostenine, S-528
3-Oxostenine, S-528
Oxostephabene, S-529
Oxostephamiersine, S-537
Oxostephanine, O-209
Oxostephanosine, O-209
Oxostephasunoline, L-244
5-Oxostriostidine, S-583
19'-Oxotabernamine, T-8
3-Oxotabersonine, O-210
1-(1-Oxo-2,4-tetradecadiene-8,10-diynyl)piperidine, T-121
1-(1-Oxo-2,4-tetradecadiene-8,10-diynyl)pyrrolidine, T-121
1-(1-Oxo-2,4-tetradecadiene-8,10,12-triynyl)piperidine, T-123
8-Oxo-2,4-tetradecadienoic acid, O-211
1-(1-Oxo-2,4-tetradecadienyl)piperidine, T-124
1-(1-Oxo-2,4-tetradecadien-8-ynyl)pyrrolidine, T-125
1-(1-Oxo-2,4,6,12-tetradecatetraene-8,10-diynyl)piperidine, T-128
1-(1-Oxo-2,4,8,11-tetradecatetraenyl)piperidine, T-129
1-(1-Oxo-2,4,8,11-tetradecatetraenyl)pyrrolidine, T-129
1-(1-Oxo-2,4,12-tetradecatriene-8,10-diynyl)piperidine, T-132
1-(1-Oxo-2,4,12-tetradecatriene-8,10-diynyl)pyrrolidine, T-132
12-Oxo-2,4,8-tetradecatrienoic acid, O-212
1-(1-Oxo-2,4,8-tetradecatrienyl)piperidine, T-133
1-(1-Oxo-2,4,10-tetradecatrien-8-ynyl)piperidine, T-134
8-Oxotetrahydroberberine, C-73
1-Oxo-1,2,3,4-tetrahydroisoquinoline, D-469
8-Oxotetrahydropalmatine, O-213
11-Oxotetrahydrothalifendine, T-149
8-Oxotetrahydrothalifendine, C-73
2-Oxo-6-(1,2,3,4-tetrahydroxybutyl)-3-morpholinepropanamide, A-207
11-Oxotetrodotoxin, T-298
8-Oxothaicanine, T-301
Oxothalibrumimine, T-310
Oxothallicarpine, O-214
4-Oxothermopsamine, T-361
4-Oxo-2-thiazolidinehexanoic acid, A-78
Oxotomaymycin, T-424
8-(2-Oxotridecyl)dihydrochelerythrine, T-503
13,13'-(2-Oxotrimethylene)bis[12,13-dihydrochelerythrine], C-355
Oxotuberostemonine, O-215
Oxoturkiyenine, T-684
2-(10-Oxoundecyl)-4(1H)-quinolinone, H-757
Oxoushinsunine, L-194
3-Oxovenalstonidine, V-49
3-Oxovincadifformine, O-210
5-Oxovincadifformine, V-111
22-Oxovincal leukoblastine, V-128
19-Oxovincamine, V-115
19-Oxovoacangine, V-170
20-Oxovoacangine, V-170
3-Oxovoacangine, V-170
19-Oxovoacristine, V-170
3-Oxovoacristine, V-170
3'-Oxovobtusine N-oxide, V-188
3'-Oxovobtusine, V-188
18-Oxovobtusine, V-188
2''-Oxovorucharin, U-53
Oxylophine, L-40
17-Oxyozurimine, D-33
Oxtriphylline, C-419
Oxyacanthine, O-1
Oxyanthin, H-673
Oxyavicine, D-404
Oxyberberine, O-216
Oxyberberubine, O-216
3,3'-Oxybis[5-(aminocarbonyl)benzoic acid], D-815
6,6'-Oxybis[7-(2-amino-1,4-dihydro-4-oxo-6-pteridiny)-6,7-dideoxy-D-glycero-L-galacto-heptonic acid] 3,3'-bis(hydrogen sulfate), S-637
5,5'-Oxybis-1,3-benzenedicarboxylic acid, D-815
5,5'-Oxybis[1,5-dihydro-3-methyl-2H-pyrrol-2-one], O-217
7,7'-Oxybis[6-methyl-5,8-dioxo-1-isoquinoline-methanol], O-218
3,3'-[Oxybis(methylene)]bis[1-methoxy-9H-carbazole], O-221
3,3'-[Oxybis(methylene)]bis[9-methoxy-9H-carbazole], O-219
Oxybocconine, D-510
Oxychelerythrine, D-407
Oxychelidonine, C-357
Oxychelirubine, D-510
Oxychlororaphine, P-319
10,11-Oxy-10,12a-cyclo-10,11-secocolchicine, O-220
5,5'-Oxydi[3-methyl-3-pyrrolin-2-one], O-217
Oxydimorphine, M-703
Oxydimurrayafoline, O-221
N^b-Oxy-20,21-epiaspidospermidine, A-1498
Oxygambirtannine, D-434
Oxyguberpazine Q, H-382
Oxyhydrastinine, I-305
Oxyisoterihanine, D-482
Oxylophine, T-561
Oxylophine, D-626
Oxylophine, O-93
Renilla Oxyluciferin, C-558
Luciola Oxyluciferin, F-56
Cypridina Oxyluciferin, O-222
Watasenia Oxyluciferin, O-223
Oxymaistemonine, M-49
Oxymatrine, M-121
N^b-Oxymelonine, M-188
Oxymorphine, M-703
Oxymycin, A-832
Oxynarcotine, N-35
Oxynemorensine, N-89
Oxynicotine, N-190
Oxynitidine, D-482
Oxypalmatine, O-213
Oxyphylline A, A-151
Oxyproline, H-715
Oxyprotopine, P-667
Oxyprotostemonine, P-670
8-Oxypseudopalmatine, O-202
Oxypterine, O-224
Oxypyrrolnitrin, P-961
Oxyretroisosenine, N-89
Oxyretuline, R-63
Oxyrubroflavin, D-229
Oxysanguinarine, D-510
Oxysceptrine, O-225
Oxysophocarpine, S-371
Oxysparteine, O-208
Oxysporidinone, O-226
Oxystemofoline, S-515
Oxystemokerrillactone, O-227
Oxystemokerrin N-oxide, S-516
Oxystemokerrin, S-516
Oxyterihanine, D-482
Oxythalifendine, O-216
Oxytrimethylamine, C-419
Oxytriptyline, D-430
Oxytyramine, D-920
3-Oxyuracil, U-42
Oxyviolacein, V-142
Oxyzoanthamine, Z-30
P 42-1, A-244
P 725C, O-151
PA 106, A-1044
PA 95, A-78
PA 155A, I-102
PA 114A, O-131
PA 371e, I-300
PA 371 γ , I-300
PA444, B-150
Pachyaximine A, A-886
Pachyaximine B, A-884
Pachyaxioside A, A-881
Pachyaxioside B, A-886
Pachycarpidine, M-121
Pachycarpine (obsol.), S-385
Pachycarpine N¹⁶-oxide, S-385
Pachycereine, T-233
Pachychaline A, P-1
Pachychaline B, P-2
Pachychaline C, P-3
Pachyconfine N-oxide, P-4
Pachyconfine, P-4
Pachydermin, P-5
Pachygonamine, Y-5
Pachygonine, C-538
Pachyovotamine, T-407
Pachypodanthine, O-87
Pachysamine A, D-266
Pachysamine B, P-601
Pachysamine E, P-601
Pachysamine G, P-601
Pachysamine H, P-601
Pachysanaximine A, D-266
Pachysandrine A, D-266
Pachysandrine B, D-266
Pachysandrine C, D-266
Pachysandrine D, D-266
Pachysanonin, A-747
Pachysantermine A, P-6
Pachysiphine, T-11
Pachystaudine, S-534
Pachysterine, P-7
Pachysterminine A, P-8
Pachysterminine B, P-8
Pacidine, M-321
Pacifarinic acid, S-33
Pacifidine, M-359
Pacifiline, H-123
Pacifinine, H-123
Paciline, M-321
Pacinine, M-321
Paclitaxel ceribate, T-52
Paclitaxel poliglumex, T-52
Paclitaxel, T-52
PA33334, D-283
PA33343, D-284
PA3334, D-288
PA3343, D-289
PA334, D-249
PA3434, D-287
PA4344, D-291
PA434, D-251
Paecilaminol, A-758
Paecilosetin, C-468
Paeciloxazine, P-9
Pagicerine, P-10
Pagsulfine, P-11
Pahutoxin, P-12
Pahybrine, P-13
Paipunine, P-14
Pakistanamine, P-15
Pakistanine, P-16
Palasimide, P-17
Palauamine, P-18
Palaudine, P-80
Palauimide, P-19
Paliclavine, P-20
Palicoside, S-583
Palidimine, P-21
Palinine, P-135
Palinurine A, P-22
Palinurine B, P-22
Paliurine A, P-23
Paliurine B, P-23
Paliurine C, P-24
Paliurine D, P-24

- Paliurine E, P-25
 Paliurine F, P-26
 Paliurine G, P-27
 Paliurine H, P-28
 Paliurine I, P-29
 Pallidiflorine, P-30
 Pallidine *N*-oxide, F-80
 Pallidine, F-80
 Pallidinine, P-31
 Pallimamine, P-32
 Palmadine, H-202
 Palmanine, P-33
 Palmasine, H-202
 Palmatine chloroform, P-35
 Palmatine, P-34
 Palmatisine, P-36
 Palmatrubine, P-34
 Palmerolide A, P-37
 Palmidrol, P-38
 Palmilycorine, L-348
 Palmirine, R-159
 Palmitamide, H-221
 Palmitan, P-38
 Palmitic acid, H-221
*N*²-Palmitoleoylglutamine, G-108
*N*²-Palmitoylglutamine, G-108
 1-*O*-Palmitoyl-2-*O*-(1-*O*-palmitoyl-2-*O*-stearyl)-
 glycerophosphorylpseudolycorine, P-716
 1-*O*-Palmitoyl-2-*O*-(1-*O*-palmitoyl-2-*O*-stearyl/
 oleoyl)glycerophosphoryllycorine, L-348
N-Palmitoylputrescine, B-422
N-Palmitoylpyrrolidine, P-949
*N*⁶-Palmitoyltryptamine, T-639
 Palosine, A-1499
 Paludosine, P-39
 Palustridine, P-40
 Palustrine, P-40
 Palythazine, P-41
 Palytoxin, P-42
 Palyzoanthoxanthin A, P-94
 Palyzoanthoxanthin B, P-94
 Palyzoanthoxanthin C, P-94
 Pamamycin 593A, P-43
 Pamamycin, P-43
 Pamine, S-159
 PAMN, T-632
 Pamolyn 100, O-37
 Pampulhamine, D-88
 Panacosmine, P-44
 Panaefluoroline A, P-45
 Panaefluoroline B, P-46
 Panaefluoroline C, P-47
 Panaefluoroline D, P-48
 Panaefluoroline E, P-49
 Panaefluoroline F, P-50
 Panaefluoroline G, P-51
 Panaefluoroline H, P-52
 Panamine, P-53
 Panarine, T-425
 Panclar, D-197
 Pancordine, P-54
 Pancorine, P-55
 Pancorinine, P-56
 Pancracine, P-57
 Pancrassidine, U-35
 Pancratine, H-6
 Pancratisside, P-58
 Pancratistatin, P-58
 Pancritamine, G-3
 Pandamarilactam 3x, P-59
 Pandamarilactam 3y, P-59
 Pandamarilactone 1, P-60
 Pandamarilactone 31, P-61
 Pandamarilactone 32, P-62
 Pandamarilactonine A, P-63
 Pandamarilactonine B *N*-oxide, P-63
 Pandamarilactonine B, P-63
 Pandamarilactonine C, P-63
 Pandamarilactonine D, P-63
 Pandamarine, P-64
 Pandamine, S-166
 Pandaminine, M-190
 Pandanamine, P-65
 (*E,E*)-Pandanamine, P-65
 Pandicine, P-66
 Pandine, P-67
 Pandoline, P-733
 Panganensine R, P-69
 Panganensine S, P-69
 Panganensine X, P-68
 Panganensine Y, P-68
R-Panganensine, P-69
S-Panganensine, P-69
 Pangkoramine, L-175
 Pangkorimine, E-102
 Pangrenine, A-247
 Panicudine, H-198
 Paniculadine, H-198
 Paniculamine, P-70
 Paniculatin A, P-247
 Paniculatin B, P-247
 Paniculatin, H-181
 Paniculatin, L-334
 Paniculatin, M-325
 Paniculidine A, I-124
 Paniculidine B, I-124
 Paniculidine C, I-124
 Paniculidine, A-910
 Paniculine, M-309
 Paniculine, P-71
 Paniculol, I-124
 Panicutine, H-161
 Pannellin, P-72
 Pantherine, A-837
 Pantherinine, P-73
 Pantocin A, P-75
 Pantocin A₂, P-74
 Pantos, D-197
 Panurensine, P-76
 Papaveraldine, P-77
 Papaverberbine, P-78
 Papaverine hydrochloride, P-79
 Papaverine, P-79
 Papaveroline tetramethyl ether, P-79
 Papaveroline, P-80
 Papaveroxidine, N-45
 Papaveroxine, N-45
 Papaveroxinoline, N-45
 Papaverrubine A, R-86
 Papaverrubine B, G-87
 Papaverrubine C, G-87
 Papaverrubine D, G-87
 Papaverrubine E, R-86
 Papaverrubine F, G-87
 Papaverrubine G, A-638
 Papaverrubine H, G-87
 Papavoriendine, P-81
 Papavorientine, P-82
 Papilamine, B-455
 Papilicine, B-455
 Papilinine, M-667
 Papillamidine, M-667
 Papillotriene, B-473
 Papillozine C, P-83
 Papracine, N-35
 Papracinine, F-174
 Paprafumine, A-143
 Papraine, C-653
 Papraline, M-446
 Paprarine, N-51
 Paprazine, H-667
 Papuamine, P-84
 Papuanine, V-188
 Papyraceabromine A, P-85
 Papyramine, P-86
 Parabactin A, A-203
 Parabactin, A-203
 Paraensidimerine A, P-87
 Paraensidimerine B, P-88
 Paraensidimerine C, P-87
 Paraensidimerine D, P-88
 Paraensidimerine E, P-87
 Paraensidimerine F, P-87
 Paraensidimerine G, P-87
 Paraensine, P-89
 Paraensine, R-31
 Paragraine, P-737
 Paraherquamide A *N*¹⁷-oxide, P-90
 Paraherquamide A, P-90
 Paraherquamide B, P-90
 Paraherquamide C, P-90
 Paraherquamide D, P-90
 Paraherquamide E, P-90
 Paraherquamide F, P-91
 Paraherquamide G, P-91
 Paraherquamide H, P-90
 Paraherquamide I, P-90
 Paramaleic acid, F-173
 Paramorphine, T-350
 Paraquileginine, C-812
 Paratunamide A, P-92
 Paratunamide B, P-92
 Paratunamide C, P-92
 Paratunamide D, P-92
 Paravallaridine, P-93
 Paravallarine, P-93
 Paraxanthine, X-5
 Parazoanthoxanthin A, P-94
 Parazoanthoxanthin B, P-94
 Parazoanthoxanthin C, P-94
 Parazoanthoxanthin D, P-94
 Parazoanthoxanthin E, P-94
 Parazoanthoxanthin F, P-94
 Parazoanthoxanthin G, P-94
 Paredrine, A-898
 Paredrinex, A-898
 Pareirubrine A, G-159
 Pareirubrine B, G-159
 Pareitropone, G-159
 Parfumidine, F-174
 Parfumine, F-174
 Paricine, P-95
 Parilac, E-157
 Parkacine, N-43
 Parkamine, F-7
 Parkimbine, Y-13
 Parlodel, E-157
 Parodel, E-157
 Parsonsianidine, P-96
 Parsonsianine, P-97
 Parsonsine *N*-oxide, P-98
 Parsonsine, P-98
 Parthenolidine, P-99
 Parvifagarine, P-100
 Parviflorine, F-174
 Parvine, N-70
 Parvineostemonine, P-101
 Parvinine, P-102
 Parvistemoamide, P-103
 Parvistemoline, P-104
 Parvistemonine, P-105
 Paspaclavine, P-106
 Paspalic acid, P-107
 Paspalicine, P-109
 Paspalin PII, Z-37
 Paspalin PI, Z-37
 Paspaline B, P-108
 Paspaline, P-108
 Paspalinine, P-109
 Paspalitre A, P-110
 Paspalitre B, P-110
 Paspalitre C, P-111
 Passibiflorin, D-565
 Passicapsin, D-565
 Passicoccin, D-565
 Passicoriacin, D-565
 Passiedulin, H-673
 Passiflorine, M-426
 Passitrifasciatin, D-565
 Patagonine, P-15
 Patchoulipyridine, P-112
 Pateamine A, P-113
 Patellamide A, P-114
 Patellamide B, P-115
 Patellamide C, P-116
 Patellamide D, P-117
 Patellamide E, P-118
 Patellamide F, P-119
 Patellamide G, P-120

- Patellin 1, P-121
 Patellin 3, P-122
 Patellin 4, P-123
 Patellin 5, P-124
 Patellin 6, P-125
 Patentine, M-323
 Pathocidin, A-735
 Patricin, O-131
 Patungensine, P-126
 Patupilone, E-111
 Paucicaline, P-127
 Paucidactine A, P-128
 Paucidactine B, P-128
 Paucidactine C, P-129
 Paucifinine *N*-oxide, K-74
 Paucifinine, K-74
 Pauciflorine A, P-130
 Pauciflorine B, P-131
 Paucine, C-456
 Paucivenine, P-132
 Pauridianthine, P-133
 Pauridianthine, P-134
 Pauridianthiol, P-133
 Pauridianthoside, P-135
 Pausperadine A, U-8
 Pavabid, P-79
 Pavanoline, P-136
 Paveral, C-555
 Pavettine, V-139
 Paxdaphnidine A, P-137
 Paxdaphnidine B, D-34
 Paxdaphnine A, P-138
 Paxdaphnine B, P-139
 Paxidine, C-555
 Paxillarine A, D-264
 Paxillarine B, D-263
 Paxilline, P-140
 Paxinorol, P-141
 Paxiphylline A, P-142
 Paxiphylline B, P-143
 Paynantheine, G-13
 PB 421, A-176
 PB 448, I-82
 PB 490, I-82
 PB4, D-631
 PC 3, M-491
 PC-M4, P-144
 PC-M5, P-140
 PC-M5, P-145
 PC-M6, P-140
 PD 116152, A-1221
 PD 125375, P-146
 PDE I, A-1222
 PDE II, A-1223
 PDMEA, D-197
 PEA, P-38
 Pearsonine, T-558
 Peceylanine, P-147
 Peceyline, P-148
 Pecilocin, V-34
 Pocrassipine A, P-149
 Pocrassipine B, P-150
 Pectenine, T-167
 Pedatisectine A, P-151
 Pedatisectine C, P-152
 Pedatisectine D, P-153
 Pedatisectine E, P-154
 Pedatisectine F, P-155
 Pedatisectine G, T-250
 Pedicularidine, P-156
 Pedicularine, P-156
 Pediculidine, P-157
 Pediculine, P-158
 Pediculine, P-157
 Pedoxin, P-159
 Pedroamine, D-88
 Pedrolon, A-898
 Peduncin, P-160
 Peduncularidine, P-161
 Peduncularine, P-162
 Peduncularistine, A-1442
 Pedunculine, C-618
 Peepuloidine, P-163
 Pegaline, H-697
 Pegamine, H-706
 Peganidine, P-164
 Peganine *N*-oxide, P-165
 Peganine, P-165
 Peganol *N*-oxide, D-503
 Peganol, D-503
 Peg-paclitaxel, T-52
 Peimidine, P-166
 Peimine, C-298
 Peiminine, C-298
 Peiminoside, C-298
 Peimiphine, P-167
 Peimisine, P-168
 Peimitidine, P-169
 Peimunine, P-170
 Peinamine, P-171
 Pelagiomicin A, H-514
 Pelagiomicin B, H-514
 Pelagiomicin C, H-514
 Pelankine, V-117
 Pelargonoyl vanillylamide, C-100
 Pelirine, A-166
 Ψ -Pelletierine, A-1580
 Pelletierine, P-173
 Pellitorine, D-106
 Pelottine, T-237
 Pelonin, P-897
 Pelosine, C-812
 Penangine, M-577
 Penaramides, P-174
 Penarcine, L-347
 Penaresamides, P-174
 Penaresidin B, P-175
 Penaresin, B-336
 Penarolide A₁, P-176
 Penarolide A₂, P-177
 Penarolide sulfate A₁, P-176
 Penarolide sulfate A₂, P-177
 Penasulfate A, P-178
 Pencolide, P-179
 Pendilinine, C-552
 Pendine, C-552
 Pendolmycin, P-180
 Pendulamine A, P-181
 Pendulamine B, P-181
 Penduline, M-313
 Penduline, T-286
 Penduline, T-563
 Pendulinine, C-552
 Pengbeimine A, P-182
 Pengbeimine B, P-182
 Pengbeimine C, P-183
 Pengbeimine C, P-212
 Pengbeimine D, P-182
 Pengbeisine A, K-114
 Pengbeisine B, K-114
 Pengshenine A, P-184
 Pengshenine B, M-375
 Peniamidienone, P-185
 Penicidone A, P-186
 Penicidone B, P-186
 Penicidone C, P-187
 Penicillazine, T-487
 Penicillenol A₁, P-188
 Penicillenol A₂, P-188
 Penicillenol B₁, P-188
 Penicillenol B₂, P-188
 Penicillenol C₁, P-188
 Penicillenol C₂, P-188
 Penidiamide, P-189
 Penidilamine, P-190
 Penigequinolone A, P-191
 Penigequinolone B, P-191
 Penimide A, P-192
 Penimide B, P-192
 Penimide, P-192
 Peniprequinolone, P-193
 Penitrem A, P-194
 Penitrem B, P-194
 Penitrem C, P-194
 Penitrem D, P-194
 Penitrem E, P-194
 Penitrem F, P-194
 Penitrem G, P-194
 Penitremone A, P-194
 Penitremone B, P-194
 Penitremone C, P-194
 Penniclavine, S-273
 Pennigritrem, P-195
 Pennsylvavine, P-196
 Pennsylvavoline, P-196
 Pennsylvanamine, T-312
 Pennsylvanine, T-312
 Penochalasin A, P-197
 Penochalasin B, P-197
 Penochalasin C, P-197
 Penochalasin D, P-197
 Penochalasin E, C-325
 Penochalasin F, C-325
 Penochalasin G, C-328
 Penochalasin H, C-326
 1,5,9,14,18-Pentaazaooctadecane, D-289
 1,11,19,27,35-Pentaamino-3-tetracontanol, P-198
 1,6,11,16,21-Pentaazadocosane, D-290
 1,6,10,15,20-Pentaazaicosane, D-291
 1,5,9,13,17-Pentaazazheptadecane, T-456
 1,5,10,14,19-Pentaazanonadecane, D-287
 1,5,9,13,18-Pentaazaoctadecane, D-288
 4,8,12,16,20-Pentaazatricosane-1,23-diamine, P-199
 2,4,4',6,6'-Pentabromo-2',3-bis(methylthio)-1,3'-bi-1*H*-indole, P-200
 3,3',4,4',5-Pentabromo-5'-chloro-2,2'-bi-1*H*-pyrrole, P-201
 3,3',4,4',5-Pentabromo-5'-chloro-1,1'-dimethyl-2,2'-bi-1*H*-pyrrole, P-201
 2,4,4',6,6'-Pentabromo-2'-(methylthio)-3,3'-bi-1*H*-indole, T-110
 Pentabromopseudilin, D-324
 Pentachloropseudilin, D-324
 N^b-Pentacosanoyltryptamine, T-639
 O¹, O², O⁶, O⁹, O⁵-Pentadecacylevonine, P-248
 2,9-Pentadecadiene-12,14-diyonic acid (2-hydroxyisobutyl)amide, P-202
 2,9-Pentadecadiene-12,14-diyonic acid isobutylamide, P-202
 2,9-Pentadecadiene-12,14-diyonic acid, P-202
 1-Pentadecanamine, P-203
 N-Pentadecanoylarginine, A-1413
 N-Pentadecanoyltyrosine, T-700
 2-(6,9,12-Pentadecatrienyl)-4(1*H*)-quinolinone, H-657
 8-(3,6,9-Pentadecatrienyl)-1,5,9,13-tetraazacycloheptadecan-6-one, B-389
 5-(6-Pentadecenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 8-(11-Pentadecenyl)-1,5,9,13-tetraazacycloheptadecan-6-one, B-389
 1-Pentadecylamine, P-203
 4-Pentadecylpyridine, P-204
 2-Pentadecyl-4-quinolinol, H-657
 2-Pentadecyl-4(1*H*)-quinolinone, H-657
 8-Pentadecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-389
 2-(1,3-Pentadienyl)piperidine, H-658
 2-(1,3-Pentadienyl)-4-piperidinol, H-658
 Pentagyline, G-152
 Pentagyline, M-322
 Pentagynine, G-152
 1,2,3,4,5-Pentahydroxy-9*H*-acridin-9-one, P-205
 1,2,3,5,6-Pentahydroxy-9*H*-acridin-9-one, P-206
 1,2,3,4,5-Pentahydroxyacridone, P-205
 1,2,3,5,6-Pentahydroxyacridone, P-206
 1,3,4,5,6-Pentahydroxyacridone, P-207
 1,2,3,9,10-Pentahydroxyaporphine, P-208
 1,2,8,9,10-Pentahydroxyaporphine, P-210
 1,2,9,10,11-Pentahydroxyaporphine, P-211
 1,2,3,10,11-Pentahydroxyaporphine, P-209
 3,7,14,16,20-Pentahydroxycevan-6-one, P-212
 3,3',4,4',5-Pentahydroxy-2,7'-cyclo lign-7-ene-9,9'-dioic acid, P-213
 1,2,6,7,9-Pentahydroxydibenz[*cd,f*]indol-4(5*H*)-one, P-214
 1,2,4,6,9-Pentahydroxydihydro- β -agarofuran, P-215

- 1,2,6,8,9-Pentahydroxydihydro- β -agarofuran, P-216
 1,2,6,9,14-Pentahydroxydihydro- β -agarofuran, P-217
 1,2,8,9,14-Pentahydroxydihydro- β -agarofuran, P-218
 1,4,6,8,9-Pentahydroxydihydro- β -agarofuran, P-219
 1,6,8,9,14-Pentahydroxydihydro- β -agarofuran, P-220
 4,5,6,7,8-Pentahydroxyfuro[2,3-*b*]quinoline, F-214
 1,2,10,11,12-Pentahydroxyhomoaporphine, P-221
 3,5,8,9,15-Pentahydroxy-6(17),11-jatrophadien-14-one, P-222
 1,2,3,4,6-Pentahydroxynortropane, A-1586
 3,8,12,14,17-Pentahydroxypregn-5-en-20-one, P-223
 2,3,4,7,8-Pentahydroxyquinoline, Q-39
 5,7,9,13,14-Pentahydroxy-4(20),11-taxadien-10-one, P-224
 4,9,12,13,20-Pentahydroxy-1,6-tigliadien-3-one, P-225
 5,9,12,13,20-Pentahydroxy-1,6-tigliadien-3-one, P-226
 Pentalupine, P-227
 1,2,3,5,6-Pentamethoxyacridone, P-206
 1,2,3,9,10-Pentamethoxyaporphine, P-208
 1,2,3,9,10-Pentamethoxy-7*H*-dibenzo[*de,g*]quinolin-7-one, O-203
 6,6',7,10,12-Pentamethoxy-2,2'-dimethylberbaman, C-23
 5,6,6',7,12-Pentamethoxy-2,2'-dimethylberbaman-14-ol, T-311
 5,6,6',7,12'-Pentamethoxy-2,2'-dimethyloxyacanthan, T-346
 6,6',7,7',12-Pentamethoxy-2,2'-dimethylthalidasan, T-320
 4,5,6,7,8-Pentamethoxyfuro[2,3-*b*]quinoline, F-214
 4,5,6,9,10-Pentamethoxyindeno[1,2,3-*ij*]isoquinoline, I-33
 2,3,4,6,7-Pentamethoxy-1-(2-methylaminoethyl)phenanthrene, T-316
 2,3,7,8,10-Pentamethoxy-5-methylbenzo[*c*]phenanthridinium(1+), S-55
 2,3,4,6,7-Pentamethoxy-*N*-methyl-1-phenanthrene-methanamine, T-316
 1,2,3,9,10-Pentamethoxynoraporphine, P-208
 1,2,3,9,10-Pentamethoxyoxoaporphine, O-203
 2,3,4,6,7-Pentamethoxyphenanthroindolizidine, P-228
 3,4,5,6,7-Pentamethoxyphenanthroindolizidine, T-278
 1,2,3,9,10-Pentamethoxytetrahydroprotoberberine, C-93
N,N,N',N'',N'''-Pentamethyldipropylenetriamine, D-255
 Pentamethylenediamine, P-229
 1,1'-Pentamethylenediguandine, D-401
 1,5-Pentanediamine, P-229
N,N'-(1,5-Pentanediy)bis[3-(3,5-diiodo-4-methoxyphenyl)-2-dimethylaminopropanamide], D-729
N,N'-1,5-Pentanediybisguanidine, D-401
N-Pentanoilpyrrothine, A-759
 Pentaphylline, F-223
Racophorus Pentapteridiny skin pigment, P-230
 Pentasine, P-231
 1-Pentene-1,5-diamine, P-232
 2-Pentene-2,3-dioic acid, E-252
cis-2-(4-Pentenyl)-6-pentylpiperidine, D-813
trans-2-(4-Pentenyl)-6-pentylpiperidine, D-813
 2-(4-Penten-1-yl)- Δ^1 -piperidine, T-214
 2-(1-Pentenyl)quinoline, P-238
 2-(1*E*-Pentenyl)-4(1*H*)-quinolinone, H-660
 2-(1*Z*-Pentenyl)-4(1*H*)-quinolinone, H-660
 Pentifylline, T-355
 Pentosidine, P-233
 Pentouregine, P-234
 Pentyl β -carboline-1-propionate, C-134
 1-Pentyl-9-azabicyclo[3.3.1]nonan-3-one, P-235
N,N'-Pentylenebis(cantharimide), P-650
 1-Pentylloctylamine, T-506
 6-Pentyl-3-piperidinol, P-236
 2-Pentylpyridine, P-237
 2-Pentyl-3,4-quinolinediol, A-621
 2-Pentylquinoline, P-238
 2-Pentyl-4-quinolinol, H-660
 2-Pentyl-4(1*H*)-quinolinone *N*-oxide, H-660
 2-Pentyl-4(1*H*)-quinolinone, H-660
 Pentylsmeospongine, S-329
 Peptard, T-632
 Pepuline, P-239
 Perakenine, R-29
 Perakine dimethyl acetal, P-240
 Perakine, P-240
 Peraksine, P-241
 Peramine, P-242
 Per(deacyl)alatusamine, P-243
 Per(deacyl)chuchuhuanine, P-244
 Per(deacyl)-24-demethylemarginatine, P-245
 Per(deacyl)emarginatine, P-246
 Per(deacyl)euojaponin, P-247
 Per(deacyl)evomine, P-248
 Perdeacylhippocrateine, P-246
 Per(deacyl)hyppoglaunine, P-249
 Per(deacyl)isowilfordine, P-250
 Per(deacyl)oppositine A, P-251
 Per(deacyl)wilforine, P-252
 Per(deacyl)wilfortrine, P-253
 Peregrine alcohol, M-347
 Peregrine, M-347
 Peregrinine, M-374
 Pereirine, G-34
 Peremin, H-307
 Perfamine, P-254
 Perfaminole, H-57
 Perforine, H-57
 Perfragilin A, P-255
 Perfragilin B, P-255
 Pergilone, M-321
 Pergularinine, T-696
 Pergumidiene, B-266
 Perhentinine, P-256
 Periblastine, V-110
 13'-Peribogamine, T-8
 Pericalline, A-1365
 Pericampylinone A, I-305
 Pericidine, P-257
 Pericine *N*-oxide, S-624
 Pericine, S-624
 Periconin A, P-266
 Periconin B, P-258
 Pericristine, V-128
 Pericyclivine, T-425
 Periformylne, P-268
 Perimargine, P-259
 Perimivine, P-260
 Perinadine A, P-261
 Peripentadene, P-262
 Peripentone A, P-263
 Peripentone B, P-263
 Peripentone C, P-264
 Periphylline, P-265
 Peristrophamide, D-707
 Peristrophine, A-862
 Peritassine A, P-249
 Peritassine B, P-249
 Peritoxin A, P-266
 Peritoxin B, P-266
 Perividine, P-267
 Perivine, P-268
 13'-Perivoacangine, V-168
 Perlolidine, B-75
 Perloline, P-269
 Perlolylrine, P-270
 Permiran, Q-17
 Peronatin A, P-271
 Peronatin B, P-271
 Perophoramidine, P-272
 Perosine, P-273
 Peroxysimulenoline, Z-5
 Peroxywybutine, W-29
 Persicanidine A, C-290
 Persicanidine B, C-290
 Persicaside, H-671
 Persinine, C-290
 Perspicamide A, P-274
 Perspicamide B, P-274
 Peruvianine, P-275
 Pervilleine A *N*-oxide, M-392
 Pervilleine A, M-392
cis-Pervilleine B, M-391
 Pervilleine B, M-391
 Pervilleine C, M-391
 Pervilleine D, M-392
 Pervilleine E, M-391
cis-Pervilleine F, M-391
 Pervilleine F, M-391
 Pervilleine G, M-391
 Pervilleine H, M-392
 Pervincamine, V-115
 Peshawarine, P-276
 Pessoine, X-30
 Petaline methine, L-96
 Petaline, G-143
 Petasinecine, H-518
 Petasine, H-518
 Petasinoside, H-518
 Petasitenine, P-277
 Petchicine, P-278
 Petilidine, C-290
 Petiline, H-483
 Petilinine, C-290
 Petine *N*-oxide, C-298
 Petine, C-298
 Petisidine, P-279
 Petisidinine, P-280
 Petisidinone, P-279
 Petisine, P-281
 Petisinine, P-281
 Petitianine, P-516
 Petizine, P-281
 Petizinine, P-281
 Petomine, P-282
 Petrobactin sulfonate, P-283
 Petrobactin, P-283
 Petromindole, P-284
 Petromurin A, P-285
 Petromurin B, P-285
 Petromurin C, P-286
 Petromurin D, P-286
 Petromyzonamine disulfate, P-287
 Petromyzonamine, P-287
 Petrosamine B, P-288
 Petrosamine, P-288
 Petrosaspongolide L, P-289
 Petrosine A, P-290
 Petrosine B, P-290
 Petrosine, P-290
 Petunaisterone C 22-nicotinate-7-acetate, P-295
 Petunianine A, P-291
 Petunianine B, P-292
 Petunianine C, P-293
 Petuniasterone B 22-*O*-acetate, P-294
 Petuniasterone B 7,22-dinicotinate, P-294
 Petuniasterone B 22-nicotinate, P-294
 Petuniasterone B, P-294
 Petuniasterone C 7,22-dinicotinate, P-295
 Petuniasterone C 22-*O*-[(methylthio)carbonyl]acetate, P-295
 Petuniasterone C 22-nicotinate, P-295
 Petuniasterone C, P-295
 Petuniasterone-B 22-*O*-[(Methylthio)carbonyl]acetate, P-294
 Peyocactine, H-357
 Peyoglunal, P-296
 Peyoglutam, P-297
 Peyonine, P-298
 Peyophorine, A-1021
 Peyoruvic acid, P-299
 Peyotine, T-237
 Peyotline, T-237
 Peyoxylic acid, T-175
 PF 1018, A-1224
 PF 1052, A-1225
 PF 1140, A-1226
 PF 1163A, A-1227
 PF 1198A, A-240
 PF 1126A, C-208

- PF 1138A, C-209
 PF 1163B, A-1227
 PF 1198B, A-240
 PF 1126B, C-208
 PF 1138B, C-209
 PF 1101B, S-282
 PF 1191, K-5
 PF 00520904, P-90
 P11012, G-3
 P11149, G-3
 PG-TXL, T-52
 Phaeantharine, P-300
 Phaeanthine 2 α -*N*-oxide, P-301
 Phaeanthine, P-301
 Phaeosphaeride A, P-302
 Phaeosphaeride B, P-302
 Phakellistatin 3, P-303
 Phalaenopsine Is, P-304
 Phalaenopsine La, P-304
 Phalaenopsine T, P-304
 Phalaenopsines, P-304
 Phalarine, P-305
 Phallacidin, P-306
 Phallacin, P-307
 Phallasacin, P-309
 Phallin B, P-308
 Phallisacin, P-309
 Phallisin, P-310
 Phalloidin, P-311
 Phalloin, P-312
 Phamine, C-744
 Phanostenine, D-346
 Phascoline, P-313
 Phascosolomine, P-314
 Phaseolunatin, H-617
 Pheantharine, P-300
 Pheanthine, P-301
 Phellibilidine, P-315
 Phellibiline, P-315
 Phellinamide, H-321
 Phellinine, S-126
 Phelloberin, B-101
 Phellodendrine, X-30
 Phenacein, D-643
 Phenacetamide, P-336
 Phenacylamine, A-694
N-Phenacylbenzamide, A-694
 Phenalamide A1, P-316
 Phenalamide A2, P-316
 Phenalamide A3, P-316
 Phenalamide B, P-317
 Phenalamide C, P-316
 2-(Phenanthro[3,4-*d*]-1,3-dioxole-6-nitro-5-carboxamido)propanoic acid, M-447
 Phenanthroviridin, D-612
 Phenatic acid A, P-318
 1-Phenazincarboxamide, P-319
 1-Phenazincarboxylic acid, P-319
 1,6-Phenazinedicarboxylic acid, P-320
 1,6-Phenazinediol 5,10-dioxide, P-321
 1,6-Phenazinediol, P-321
 1,8-Phenazinediol, P-322
 1,9-Phenazinediol, P-323
 2,3-Phenazinediol, P-324
 2,3,7-Phenazinetriol, P-326
 1,2,6-Phenazinetriol, P-325
 1-Phenazinol, P-327
 2-Phenazinol, P-328
 Phenazinomycin, P-329
 2(10*H*)-Phenazinone, P-328
 Phenazostatin A, P-330
 Phenazostatin B, P-331
 Phenazostatin C, P-332
 Phenazostatin D, P-331
 Phenazoviridin, M-417
 Phencomycin, P-320
N-(2-Phenethenyl)-2-hydroxypropanamide, X-27
 Phenethyl 2-pyrrolicarboxylate, P-944
N-Phenethylacetamide, P-342
 Phenethylamine, P-342
 β -Phenethylamine, P-342
N-Phenethylbenzamide, P-342
 2-Phenethylformamide, P-342
N-Phenethylhexanamide, P-342
N-Phenethylisobutyramide, P-342
N-Phenethylisovaleramide, P-342
N-Phenethylpropanamide, P-342
 Phenochalasin A, C-945
 Phenochalasin B, C-945
 Phenomydrol, A-694
 Phenopicolinic acid, H-426
 Phenoptin, T-142
 Phenopyrrozin, P-333
 Phenoxan, P-334
 3*H*-Phenoxazin-3-one, P-335
 3-Phenoxazone, P-335
 9-Phenoxyacridine, A-112
 9-Phenoxy-1-phenazinecarboxylic acid, H-664
 6-Phenoxyurine, P-820
 2-Phenoxyurazine, P-889
N-(2-Phenethyl)-2,4-octadienamide, O-39
 Phenyl hydrogen 2-aminoethyl phosphate, A-768
 Phenylacetic acid, P-336
 Phenylacetone, P-336
 7-Phenylacetoxynortropan-3-ol, A-1585
 3-(*N*-Phenylacetyl)amino)piperidine-2,6-dione, D-808
N-(Phenylacetyl)-2-butenediamide, P-337
 1-Phenylacetyl- β -carboline, E-289
N-Phenylacetylumamide, P-337
 Phenylacetylscopine, S-158
 Phenylahistin, P-338
 Phenylalanylproline anhydride, C-891
 Phenylalanyl-*N*-[1-(1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methyl-1-isoquinolyl)ethyl]glycine, P-339
 2-(Phenylamino)-1,4-naphthalenedione, A-1040
 4-(Phenylamino)-2-(phenylimino)-1(2*H*)-naphthalenone, L-311
 2-Phenyl-1-benzazocin-8-ol, P-340
 Phenylethanoic acid, P-336
 1-Phenylethanolamine, A-864
N-(2-Phenylethyl)-2-decene-6,8-diynamide, D-130
 2-(2-Phenylethyl)indole, P-341
 3-(2-Phenylethyl)-4(3*H*)-quinazolinone, Q-14
N-(2-Phenylethyl)acetamide, P-342
 2-Phenylethylamine, P-342
 3'-(2-Phenylethylamino)avarone, A-1560
N-(2-Phenylethyl)benzamide, P-342
N-(2-Phenylethyl)carbamate, P-343
N-(2-Phenylethyl)cinnamamide, P-344
N-(2-Phenylethyl)-2,8-decadiene-4,6-diynamide, D-103
N-(2-Phenylethyl)-2-decene-6,8-diynamide, D-130
N-(2-Phenylethyl)-2-decene-4,6,8-triynamide, D-131
 1'-(2-Phenylethylene)dityryptophenaline, D-874
 (2-Phenylethyl)formamide, P-342
N-(2-Phenylethyl)-14-heptadecene-4,16-diynamide, H-117
N-(2-Phenylethyl)hexadecanamide, P-342
N-(2-Phenylethyl)hexanamide, P-342
N-(2-Phenylethyl)-2-nonene-6,8-diynamide, N-282
N-(2-Phenylethyl)propanamide, P-342
N-(2-Phenylethyl)-3,5-undecadiene-8,10-diynamide, U-27
N-(2-Phenylethyl)-2,4-undecadiene-8,10-diynamide, U-26
 3-Phenylglyceric acid, D-651
 β -Phenylglycidic acid, P-348
 16-Phenyl-2,4-hexadecadienoic acid, P-345
 3-Phenylisoserine, A-808
 8-Phenyllobelol I, S-210
 8-Phenyllobelol, S-210
N-(Phenylmethyl)acetamide, B-84
 3-[2-(Phenylmethyl)-5-oxazolyl]-1*H*-indole, A-622
 3-(Phenylmethyl)-2(1*H*)-pyrazinone, B-90
 2-(Phenylmethyl)-4-quinazolinol, B-93
 2-(Phenylmethyl)-4(3*H*)-quinazolinone, B-93
 4-Phenyl-2,7-naphthyridin-1-amine, A-865
 4-Phenyl-2,7-naphthyridin-1(2*H*)-one, H-686
 8-Phenylnorlobelol I, S-210
 5-Phenyl-2-oxazolidinethione, P-346
 5-Phenyl-2-oxazolidinone, P-347
 3-Phenyloxiranecarboxylic acid, P-348
 4-(5-Phenylpentyl)-1*H*-indole-2,3-dione, P-349
 4-(5-Phenylpentyl)isatin, P-349
 3-Phenyl-*N*-(2-phenylethyl)propanamide, P-344
 3-Phenyl-*N*-(2-phenylethyl)-2-propenamide, P-344
 3-Phenylpropanoic acid 2,3-didehydro-4-hydroxy-piperidine, A-1568
 3-Phenylpropanoic acid pyrrolidide, P-351
 3-Phenylpropanoic acid, P-350
 Phenylpropanolamine hydrochloride, A-869
 Phenylpropanolamine polistirex, A-869
 Phenylpropanolamine, A-869
N-(3-Phenylpropanoyl)pyrrole, P-351
N-(3-Phenylpropanoyl)- Δ^3 -pyrrolidone, S-86
 10-Phenyl-8-propyllobeliolone, D-403
 1-Phenyl-1*H*-pyrazole-3-carboxylic acid, P-890
 1-Phenyl-1*H*-pyrazole-5-carboxylic acid, P-890
 4-Phenylpyridine, P-352
 4-Phenyl-3-(2-pyridinyl)-2-buten-1-ol, P-353
 2-Phenyl-1-(9*H*-pyrido[3,4-*b*]indol-1-yl)ethanone, E-289
 1-Phenyl-1*H*-pyrrole-2-carboxaldehyde, P-942
 1-Phenyl-1*H*-pyrrole-2-carboxylic acid, P-944
 1-Phenyl-2-pyrroline, D-499
 3-Phenyl-2,4(1*H*,3*H*)-quinazolinone, Q-13
 3-Phenyl-4(3*H*)-quinazolinone, Q-14
 1-Phenyl-4(1*H*)-quinazolinone, Q-14
 2-Phenyl-4,5-quinolinediol, D-658
 2-Phenyl-4,6-quinolinediol, D-659
 2-Phenyl-4,7-quinolinediol, D-660
 2-Phenylquinoline, P-354
 3-Phenylquinoline, P-355
 2-Phenyl-4-quinolinol, H-693
 2-Phenyl-4(1*H*)-quinolinone, H-693
 Phenyltartronic acid, H-688
 8-Phenyl-1,5,9,13-tetrazaheptadecan-6-one, P-673
 2-Phenyl-1,5,9,14-tetraazabicyclo[12.3.1]octadecan-4-one, P-665
 1-Phenyl-1,4-tetralindicarboxylic acid discopine ester, S-157
 2-Phenyl-1,5,9-triazacyclotridecan-4-one, P-356
 Phevalin, B-88
 Phidolopin, P-357
 Philanthotoxin 433, P-358
 δ Philanthotoxin, P-358
 Philippinamide, P-359
 Philogaline, P-360
 Phlebicine, R-114
 Phlegmarine, P-361
 Phlegmariurine A, P-362
 Phlegmariurine B, P-363
 Phlegmariurine C, P-364
 Phlegmariurine M, P-365
 Phlegmariurine N, H-479
 Phloeoictine A (obsol.), P-366
 Phloeoictine A1, P-366
 Phloeoictine A2, P-366
 Phloeoictine A3, P-366
 Phloeoictine A4, P-366
 Phloeoictine A5, P-366
 Phloeoictine A6, P-366
 Phloeoictine A7, P-366
 Phloeoictine A, P-366
 Phloeoictyne A, P-366
 Phloeoictyne B, P-367
 Phloeoictyne C, P-368
 Phloretic acid, H-689
 Phloretinic acid, H-689
 Phocenic acid, M-409
 Phoebe base II, T-524
 Phoebe base 2, L-72
 Phoebe base, T-248
 Phoebebrandine A, R-116
 Phoebebrandine B, R-116
 Phoebebrandine C, R-116
 Phoebebrandine D, R-116
 Phoebebrandine E, O-49
 Phoebescortechiniine, R-116
 Phoebine, P-369
 Phoenicanthusine, P-370
 Pholiosidine, F-122

- Pholiotic acid, D-248
 Phomacin A, P-371
 Phomacin B, P-371
 Phomacin C, P-372
 Phomasetin, P-373
 Phomazarine, P-374
 Phomin, C-943
 Phomoenamamide, P-375
 Phomopsichalasin, P-376
 Phomopsin A, P-377
 Phorbasin E, P-378
 Phorbasin F, P-378
 Phorbazole A, P-379
 Phorbazole B, P-379
 Phorbazole C, P-379
 Phorbazole D, P-379
 4 α -Phorbol, P-225
 Phorboxazole A, P-380
 Phorboxazole B, P-380
 Phormidimine A, P-909
 Phormidimine B, P-909
 Phosphatidylglycorine, L-348
 Phosphatidylmethylglycorinium, L-348
 Phosphatidylpseudoglycorine, P-716
 O-Phosphoethanolamine, A-768
 Phospholombricine, L-240
 2-(3-Phosphoguanidino)ethanesulfonic acid, G-203
 N²-(17-Phosphonoxylinolenoyl)glutamine, H-645
 N²-(17-Phosphonoxylinoleoyl)glutamine, H-645
 Phosphonyldeoxycholine, T-602
 Phosphoric acid mono[2-(dimethylamino)ethyl] ester, D-197
 Phosphoryldimethylaminoethanol, D-197
 Phosphoryldimethylcholamine, D-197
 Phosphoryldimethylethanolamine, D-197
 N-Phosphoryllombricine, L-240
 Phosphothallessemine, L-240
 Photobactin, P-381
 Photolumazine B, P-382
 Photolumazine C, P-382
 Phthalimide, P-383
 3,4-Phthaloylpyridine, B-71
 Phthoxazolin A, I-161
 Phthoxazolin B, I-161
 Phthoxazolin C, I-161
 Phthoxazolin D, I-161
 PhTX-433, P-358
 Phulchowkiamide A, D-281
 Phycodioic acid, D-104
 Phyrgrine, P-384
 Phyllalbine N-oxide, C-633
 Phyllalbine, C-633
 Phyllanthimide, P-385
 Phyllanthine, P-386
 Phyllanthin, H-520
 Phyllanthoside, H-520
 Phyllantidine, P-387
 Phylline, T-177
 Phylloactin II, B-110
 Phylloactin, B-110
 Phyllochrysin, A-630
 Phyllocryptine, R-123
 Phyllocryptonine, P-388
 Phyllospadine, P-389
 Phyllostemine, S-248
 Phyllosteminine, P-390
 Phyllosterone, R-165
 Phyllostictine A, P-391
 Phyllostictine B, P-392
 Phyllostictine C, P-393
 Phyllostictine D, P-394
 Physarigin A, P-395
 Physarigin B, P-396
 Physarigin C, P-395
 Physarorubinic acid A, P-397
 Physarorubinic acid B, P-398
 Physochlaine, M-391
 Physoperuvine, P-399
 Physostigmine oxide, G-51
 Physostigmine salicylate, P-400
 Physostigmine sulfate, P-400
 Physostigmine, P-400
 Physostol, P-400
 Physovenine, P-401
 Phytolaccanin, B-110
 PI 091, A-1228
 PI 290, B-29
 Pibocine A, F-49
 Pibocine B, F-49
 Piccolamine N-oxide, P-402
 Piccolamine, P-402
 Piclavine A, D-133
 Piclavine A₁, D-133
 Piclavine A₂, D-133
 Piclavine A₃, D-133
 Piclavine A₄, D-133
 Piclavine B, D-107
 Piclavine C, D-107
 2-Picoline methiodide, M-537
 α -Picoline, M-537
 Picolinic acid, P-896
 Picraconitine, M-331
 Picralinal, B-420
 (Z)-Picralinal, B-420
 Picraline, B-420
 Picralstonine, P-417
 Picranitine, P-403
 Picraphylline, P-404
 Picrasidine A, P-406
 Picrasidine B, P-407
 Picrasidine C, P-408
 Picrasidine D, D-724
 Picrasidine E, P-409
 Picrasidine F, P-410
 Picrasidine G, P-416
 Picrasidine H, P-411
 Picrasidine I, D-686
 Picrasidine J, D-686
 Picrasidine K, P-412
 Picrasidine L, P-413
 Picrasidine M, P-414
 Picrasidine N, P-414
 Picrasidine O, P-413
 Picrasidine P, D-709
 Picrasidine Q, D-546
 Picrasidine R, P-415
 Picrasidine S, P-416
 Picrasidine T, P-416
 Picrasidine U, P-414
 Picrasidine V, C-135
 Picrasidine W, T-531
 Picrasidine X, P-405
 Picrasidine Y, P-405
 Picratidine, B-420
 Picrinine, P-417
 Pictamine, P-418
 Pictonamine, P-419
 Pictumine, M-323
 Pictumine, P-420
 Pidolic acid, O-206
 Piepunendine A, T-239
 Piepunendine B, A-100
 Piepunensine A, M-352
 Pierardine, D-739
 Piericidin B₅N-oxide, P-421
 Piericidin B₁, P-421
 Piericidin B₂, P-421
 Piericidin B₃, P-421
 Piericidin B₄, P-421
 Piericidin B₅, P-421
 Piericidin C₁, P-421
 Piericidin C₂, P-421
 Piericidin C₃, P-421
 Piericidin C₄, P-421
 Piericidin C₅, P-421
 Piericidin C₆, P-421
 Piericidin C₇, P-421
 Piericidin C₈, P-421
 Piericidin D₁, P-421
 Piericidin D₂, P-421
 Piericidin D₃, P-421
 Piericidin D₄, P-421
 Piericidinol A₁, P-422
 Piericidin, P-421
 Pilagan, P-424
 Pillijanine, P-423
 Pilocarpidine, P-424
 Pilocarpine hydrochloride, P-424
 Pilocarpine, P-424
 Pilocarpol, P-424
 Piloceredine, P-425
 Pilocereine, P-425
 Pilofrin, P-424
 Pilokeanine, P-426
 Pilomiotin, P-424
 Pilopine VS, P-424
 Pilosamycin, P-427
 Pilosine, P-428
 Pilosinine, P-429
 Pilosinone, P-428
 Piloturine, P-430
 Pilovisc, P-424
 Pimpifolidine, D-99
 Pimpinellidine, P-431
 Pimprinaphine, A-622
 Pimprinethine, A-622
 Pimprinine, A-622
 Pingbeidinose, P-432
 Pingbeine, P-433
 Pingbeininose, P-433
 Pingbeinone, P-434
 Pingpeimine A, C-293
 Pingpeimine B, C-292
 Pingpeimine C, P-212
 ent-3-(+)-Pinidine, M-522
 Pimidine, M-522
 Pimidinol, H-705
 Pinnaic acid, P-435
 Pinnamine, P-436
 Pinnasterone, H-244
 Pinnatoxin A, P-437
 Pinnatoxin B, P-438
 Pinnatoxin C, P-438
 Pinnatoxin D, P-439
 Pinoline, T-173
 Pinolinone, P-440
 Pipbinine, P-441
 Pipecolic acid methylbetaine, H-348
 Pipecolic acid, P-456
 Pipecolinic acid, P-456
 Piperadione, C-273
 Piperamide-C5:1 (2E), M-456
 Piperamide-C7:1 (6E), M-452
 Piperamide-C7:2 (2E,6E), M-452
 Piperamide-C9:1 (8E), M-455
 Piperamide-C9:2 (2E,8E), M-455
 Piperamide-C9:3 (2E,4E,8E), M-455
 Piperanine, P-467
 Piperarborenine A, P-474
 Piperarborenine B, P-474
 Piperarborenine C, P-442
 Piperarborenine D, P-474
 Piperarborenine E, P-442
 Piperarboresine, P-443
 Piperarboricoline, P-452
 3,3'-(2,5-Piperazinediyl)bis[6-bromo-1H-indole], D-929
 Piperazinomycin, P-444
 Piperazirum, P-445
 Pipericallosidine, P-446
 Pipericallosine, P-447
 Piperchabamide A, P-463
 Piperchabamide B, P-471
 Piperchabamide C, P-448
 Piperchabamide D, P-469
 Pipericide, R-60
 Pipericitine, O-36
 Pipericyclamide, P-449
 Pipericyclobutanamide A, P-450
 Pipericyclobutanamide B, P-451
 Piperdardine, P-452
 Pipericosalidine, E-51
 Piperettine, P-452
 Piperettyline, P-453
 1-Piperetylpyrrolidine, P-453
 Pipericine, O-29
 Δ^2 -Piperideine, T-220

- Piperidic acid, A-715
 1-Piperidinecarboxaldehyde, P-455
 2-Piperidinecarboxylic acid, P-456
 4,4-Piperidinediol, P-459
 Piperidine, P-454
 2,4,5-Piperidinetricarboxylic acid, P-457
 3,4,5-Piperidinetriol, P-458
 Piperidinic acid, A-715
 Piperidinic lactam, P-956
 2-Piperidinimine, I-49
 4-Piperidinone, P-459
 1-(2-Piperidino)-2-propanol, S-216
 5-(2-Piperidinyl)-2,3'-bipyridine, A-963
 1-(2-Piperidinyl)-2-butanol, H-435
 1-(2-Piperidinyl)-2-butanone, H-435
 1-(2-Piperidinyl)-2-pentanone, H-37
 1-(2-Piperidinyl)-1-propanol, C-603
 1-(2-Piperidinyl)-1-propanone, C-604
 1-(2-Piperidinyl)-2-propanone, P-173
 3-(2-Piperidinyl)pyridine, P-460
 4-Piperidone-2-carboxylic acid, O-196
 γ -Piperidone, P-459
N-Piperidyl-*p*-hydroxyphenylacetoneitrile, G-80
 Piperine S, P-465
 Piperine, P-467
 Piperlactam S, D-567
 Piperlongine, P-461
 Piperlongumine, P-473
 Piperlonguminine, P-462
 Piperlotine A, H-691
 Piperlotine B, H-691
 Piperlotine C, T-578
 Piperlotine D, T-578
 Piperlotine E, P-351
 Piperlotine F, P-956
 Piperlotine G, P-956
 Piperlotine H, P-956
 Piperlotine I, P-955
 Piperlotine J, A-714
 Piperlotine K, A-714
 Piperlotine L, A-714
 Pipermethystine, P-463
 Pipernonaline, P-464
 Piperoctadecalidine, O-34
 Piperolactam A, D-567
 Piperolactam B, T-537
 Piperolactam C, T-537
 Piperolactam D, T-537
 Piperolactam E, T-537
 Piperolein A, P-465
 Piperolein B, P-464
 Piperonylic acid, M-442
 Piperonylideneacetic acid, M-457
 9-Piperonyl-2,4-nonadienoic acid, M-454
 9-Piperonylnonanoic acid, M-454
 Piperovatine, P-466
 1-Piperoylpiperidine, P-467
 1-Piperoylpyrrolidine, T-495
 Piperpense, P-468
 Pipersintenamide, P-452
 Piperstachine, P-469
 Pipertipine, P-470
 Piperumbellactam A, D-567
 Piperumbellactam B, D-567
 Piperumbellactam C, D-567
 Piperumbellactam D, D-567
 Piperundecalidine, P-471
 Piperx, W-17
 Piperylene, T-495
 Pipgulzarine, P-472
 Piplaroxide, P-473
 Piplartine dimer A, P-474
cis-Piplartine, P-473
 Piplartine, P-473
 Pipnoohine, O-33
 Pipsaedine, P-475
 Piptahsine, H-502
 Piptamine, O-115
 Piptamine, P-203
 Piptanthine, P-476
 Piptigrine, P-464
 Pipwaqarine, G-224
 Pipyahyine, P-477
 Pipzorine, O-30
 Pirazofurin, P-891
 Piricardio, A-845
 Piricularin, P-478
 Piridoxilate, P-923
 Piriferine, P-479
 Piriferinol, P-479
 Pisatoside, D-459
 Piscarinine A, P-480
 Piscarinine B, P-481
 Piscerythoxazole, P-482
 Piscerythramine, P-483
 Pisolithin A, H-687
 Pisolithin B, H-520
 Pisopowamine, P-484
 Pisopowetine, P-484
 Pisopowiaridine, P-484
 Pisopowiarine, P-484
 Pisopowidine, P-484
 Pisopowine, P-484
 Pistacidimedor, P-485
 Pistillarine, S-398
 Pitayine, Q-18
 Pithecolobine, P-486
 Pitamide A, P-487
 Pityriacitrin B, P-489
 Pityriacitrin, P-488
 Pityrialactone, P-490
 Pityriahydride, P-491
 Pityriarubin A, P-492
 Pityriarubin B, P-493
 Pityriarubin C, P-494
 Pityriazole, P-495
 Pivagabine, A-715
 PKC 412, S-498
 Plakinamine A, P-496
 Plakinamine B, P-497
 Plakinamine C, P-498
 Plakinamine D, P-498
 Plakinamine E, P-496
 Plakinamine F, P-496
 Plakinamine G, P-499
 Plakinamine H, P-497
 Plakinamine I, P-500
 Plakinamine I, P-501
 Plakinamine J, P-502
 Plakinamine K, P-496
 Plakinidine A, P-503
 Plakinidine B, P-503
 Plakinidine C, P-503
 Plakinidine D, P-503
 Plakinidine E, P-504
 Plakohypaphorine A, H-764
 Plakohypaphorine B, H-764
 Plakohypaphorine C, H-764
 Plakohypaphorine D, H-764
 Plakohypaphorine E, H-764
 Plakohypaphorine F, H-764
 Plakoridine A, P-505
 Plakoridine B, P-506
 Plakortamine A, P-507
 Plakortamine B, V-139
 Plakortamine C, P-508
 Plakortamine D, P-509
 Planchonelline, P-510
 Planomycin, F-48
 Plant indican, I-104
 Plantagonine, B-254
 Platencin, P-511
 Platensic acid, P-512
 Platensimide A, P-512
 Platensimycin, P-513
 Platifillin, P-516
 Platyamide A, T-452
 Platylerine *N*-methosalt, P-514
 Platylerine, P-514
 Platydesmine, P-515
 Platydesminium, P-515
 Platynecine 9-viridiflorate, M-146
 Platynecine, H-233
 Platynerepterin, P-771
 Platyphylline *N*-oxide, P-516
 Platyphylline, P-516
 Plectocomine, T-173
 Plectodorine, P-517
 Pleiaseperin, R-52
 Pleiatensin simplex, R-52
 Pleiocarpamine, P-518
 Pleiocarpine, K-78
 Pleiocarpinidine, E-6
 Pleiocarpinilam, K-78
 Pleiocarpinine, K-78
 Pleiocarpoline, K-78
 Pleiocarpolinine, K-78
 Pleiocorine, P-519
 Pleiocoraline, P-520
 Pleiomutine, P-521
 Pleiomutinine, P-827
 Pleurosine, L-162
 Pleurospermine, P-522
 Pleurostyline, P-523
 Plicamine, P-524
 Plicane, P-525
 Plumerinine, P-526
 Plumocraline, P-527
 Pluviine, P-528
 PM 53, A-1521
 PM 94128, A-1229
 PM 297, P-643
 PM 93135, T-384
 Pneumax MBT, B-79
 PNU 141962, P-90
 PO 1227A, Q-37
 PO 1227B, Q-37
 Podocarpamide, Z-6
 Podopetaline, P-529
 Poecillanosine, P-530
 Poetamine, P-531
 Poetaminine, L-348
 Poetaricine, P-532
 Poeticine, P-533
 Poetinate, C-532
 Pogonopamine, P-534
 Pogonopeine, P-535
 Pogonopidine, P-536
 Pogonopine, P-537
 Pohakuline, P-538
 Polonicumtoxin A, P-539
 Polonicumtoxin B, P-539
 Polonicumtoxin C, P-539
 Polyalthenol, P-540
 Polyalthine, B-471
 Polyandrocarpamide A, P-541
 Polyandrocarpamide B, P-541
 Polyandrocarpamide C, P-541
 Polyandrocarpamide D, P-542
 Polyandrocarpamine A, P-543
 Polyandrocarpamine B, P-543
 Polyandrocarpidine A, P-544
 Polyandrocarpidine B, P-544
 Polyandrocarpidine C, P-545
 Polyandrocarpidine D, P-545
 Polyanine, S-445
 Polyavolensin, G-172
 Polyavolensinol, G-172
 Polyavolensinone, G-172
 Polyavolinamide, P-546
Epilachna borealis Polyazamacrolides, P-547
 Polybeccarine, B-53
 Polyberbine, P-548
 Polycanthidine, P-549
 Polycanthisine, P-550
 Polycarpaurine A, P-553
 Polycarpaurine B, P-551
 Polycarpaurine C, P-552
 Polycarpine, P-553
 Polycarpine, P-554
 Polycartine A, H-240
 Polycartine B, O-54
 Polycephalin B, P-555
 Polycephalin C, P-555
 Polycitone A, P-556
 Polycitone B, P-556
 Polycitorol A, P-557
 Polycitorol B, P-558
 Polycitricin A, P-559

- Polycitrin B, P-559
 Polyervine, C-621
 Polyervinine, P-560
 Polyfothine, D-625
 Polygonapholine, P-561
 Polygonatine A, D-447
 Polygonatine B, D-447
 Polygosperrmine, L-10
 Poly-D-leucine, L-153
 Poly-L-leucine, L-153
 Polylongine, D-626
 Polyneuridine aldehyde, P-562
 Polyneuridine *N*-oxide, P-562
 Polyneuridine, P-562
 Polyschistine A, M-331
 Polyschistine B, M-332
 Polyschistine C, M-345
 Polyschistine D, M-331
 Polysignine, P-563
 Polystictin, C-452
 Polysuavine, O-85
 Polyveoline, P-564
 Polyzonimine, D-741
 Ponapensin, P-565
 Poncevedrine, P-566
 Pool Eyes, B-101
 Popidine, D-88
 Popisidine, D-88
 Popisine, D-88
 Popisonine, D-88
 Popisopine, D-88
 Popolohuanone A, P-567
 Popolohuanone B, P-567
 Popolohuanone C, P-567
 Popolohuanone D, P-568
 Poranthericine, P-569
 Porantheridine, P-570
 Porantherilidine, P-571
 Porantheriline, P-569
 Porantherine, P-572
 Pordamacrine A, Y-23
 Pordamacrine B, Y-23
 Porfiromycin, M-654
 Poroidine, A-1597
 Porothramycin A, P-573
 Porothramycin B, P-573
 Porphobilinogen, P-574
 Porphyroxine, G-87
 Porritoxin, C-437
 Porritoxinsulfonic acid, C-437
 Portamycin, S-566
 Portulacaxanthin III, P-577
 Portulacaxanthin II, P-576
 Portulacaxanthin I, P-575
 Portulaxanthine, P-575
 Porveniramine, P-16
 Potanidine A, D-182
 Potanidine B, L-317
 Potanine, M-309
 Potanisine A, M-309
 Potanisine B, G-91
 Potanisine C, M-341
 Potanisine D, M-341
 Potanisine E, G-91
 Potanisine F, G-91
 Potanisine G, P-578
 Powellamine, P-579
 Powellidine, P-580
 Powelline, P-581
 Poweramine, P-582
 Powerchrine, P-583
 Poweridine, M-292
 Powerine, M-291
 PP-Ala, D-780
 PP-Factor, P-895
 PP-R, M-686
 PP-Val, D-779
 PP-V, C-150
 PQQ, M-236
 PQS, A-621
 Praecoxine, T-248
 Prajmalium bitartrate, A-216
 Pramanicin, P-584
 Prampine, T-632
 Prangosine, P-585
 Pratorimine, P-586
 Pratorine, H-290
 Pratorinine, P-586
 Pratosine, P-586
 Pravidel, E-157
 Preakuammicine, P-587
 Prealmazole C, P-588
 Preapiodionene, P-589
 Preaxinellamine, P-590
 Prebalamide, P-591
 Prebetanin, B-110
 Pre-B, S-638
 Precasine, P-592
 Precatorine, P-593
 Preclathridine A, P-594
 Precocinelline, D-886
 Precondylocarpine, P-595
 Precriwelline, P-626
 Precurarine, P-596
 Predicentrine, P-597
 Pregna-14,16-diene-3,20-diamine, P-600
 Pregna-4,14-diene-3,20-diamine, P-598
 Pregna-5,16-diene-3,20-diamine, P-599
 Pregnane-3,20-diamine, P-601
 Pregnane-3,8,12,14,17,20-hexol, P-602
 Pregnane-3,11,12,14,20-pentol, P-603
 Pregnane-3,12,14,17,20-pentol, P-604
 Pregn-14-ene-3,20-diamine, P-607
 Pregn-16-ene-3,20-diamine, P-608
 Pregn-4-ene-3,20-diamine, P-605
 Pregn-5-ene-3,20-diamine, P-606
 Pregn-6-ene-3,5,8,12,14,17,20-heptol, P-609
 Pregn-5-ene-3,8,12,14,17,20-hexol, P-610
 Preininger alkaloid, D-163
 Prelandrine, P-673
 Prellisoclinamide 2, P-611
Watasenia Preluciferin, C-560
Diaphus Preluciferin, P-612
 Premavacurine, P-613
 Premnazole, M-479
 Premnine, P-614
 Premuscimol, I-8
 Prenirurine, B-375
 Prenisatin, M-415
 Prenostodione, P-615
 9-Prenyladenine, A-137
 1-Prenyladenine, A-137
*N*⁶-Prenyladenine, Z-11
*N*⁶-Prenylagmatine, A-719
 Prenylcitpressine, P-616
N-Prenylguanidine, M-410
 4-(*N*¹-Prenylguanido)butanoic acid, C-592
O-Prenylhalfordinol, H-12
N-Prenylhaplofoline, H-56
 6-Prenyl-1*H*-indole-3-carboxylic acid, P-617
 5-Prenylindole, M-412
 6-Prenylindole, M-413
 7-Prenylindole, M-414
 3-Prenylindole, M-411
 6-Prenyl-2-indolinone, H-701
 5-Prenylisatin, M-415
 6-Prenylisatin, M-416
 6-Prenylloxindole, H-701
 7-Prenylxyflindersine, F-90
N-[2-(4-Prenylxyphenyl)ethyl]tiglamide, H-667
 4-Prenylxy-2-quinolone, H-53
 9-Prenylpaxilline, P-618
 6-Prenyl-1-phenazincarboxylic acid, M-417
 9-Prenyl-1-phenazincarboxylic acid, M-418
 3-Prenyl-4-prenyloxy-2(1*H*)-quinolinone, A-1529
N-Prenylputrescine, B-422
 4-Prenyltryptophan, M-419
 6-Prenyltryptophol, H-497
 Preocoteine *N*-oxide, P-208
 Preocoteine, P-208
 Preorixine, D-715
 Preoxazin 7, P-619
 Prepatellamide A, P-620
 Prepatellamide B formate, P-621
 Preperaksine *O*-acetate, D-491
 Prepsammaplin A, P-622
 Prepseudopalmanine, P-623
 Pre-Q1 base, A-699
 Presecamine, P-624
 Preskimmianine, P-625
 Pretazettine, P-626
 Pretomaymycin, T-424
 Pretomeimycin, T-424
 Preulicyclamide, P-627
 Preulithiacyclamide, P-628
 Preussin, B-89
 pre-W, P-641
 pre-X, P-642
 Prianosin A, D-850
 Prianosin B, D-850
 Prianosin C, P-629
 Prianosin D, D-852
 Primapterin, A-924
 Primocarcin, P-630
 Primofax, N-319
 Prioline, P-631
 Pristinamycin II_A, O-131
 Pristinamycin II_B, O-131
 Pristinamycin, O-131
 Proamanullin, A-673
 Pro-Amid, P-674
 Problaston, H-715
 Probovatine, F-80
 Procerine, P-632
 Procevine, P-633
 Prochaetoglobosin I, C-328
 Prochaetoglobosin II, C-328
 Prochaetoglobosin III, C-325
 Prochaetoglobosin III, C-326
 Prochaetoglobosin IV, C-325
 Procumbine, P-634
 Prodeoxyviolacein, P-675
 Prodigiosene, P-635
 Prodigiosin 25b, V-163
 Prodigiosin 25c, U-31
 Prodigiosin, P-636
 Proferrorosamine A, D-496
 Proferrorosamine B, P-637
 C-Profurocurine, A-617
 Prolifine, P-638
 2-L-Proline-3-isoleucine- α -amanitin, A-673
 β -Proline, P-952
 Propylleucyldiketopiperazine, C-876
 Propylproline anhydride, C-898
 Propylvaline anhydride, C-901
 Prometaphanine, P-639
 Promoinducin, P-640
 Promothiocin A, P-641
 Promothiocin B, P-642
 Promucosine, P-643
 Pronuciferine, P-643
 Prooxocryptochine, P-644
 Propadrine, A-869
 Propanamine, P-648
N,N'-1,3-Propanediylbis[1,4-butanediamine],
 D-251
 1,6-Propano-3-ethylidene-2,5-piperazinedione,
 E-249
 13-Propanoyloxylupanine, H-554
 6-Propanoyloxy-3-tigloyloxytropone, M-391
 7-Propanoyloxy-3-tigloyloxytropan-6-ol, M-392
 2-Propanoylpiperidine, C-604
 2-Propanoylpyridine, P-645
 2-Propenamide, P-646
 2-Propenoic acid, P-646
 2-(1-Propenyl)- Δ^1 -piperideine, T-218
 3-[6-(1-Propenyl)-2*H*-pyran-2-ylidene]-2,4-pyrroli-
 dinedione, V-76
 2-(1-Propenyl)quinoline, P-647
*N*¹-Propionylaplysinopsin, A-1357
 Propionylcholine, C-419
N-Propionyl-*N*-deacetylaspidolimidine, A-1490
N-Propionylholothin, A-759
 7-Propionylintermedine, I-160
 6-Propionylumazine, H-702
 7-Propionyllycopsamine, I-160
 8-(Propionylmethyl)dihydrochelerythrine, O-167
 2-Propionyl-1-pyrroline, D-493
 Propionylpyrrothine, A-759

- N*-Propylacetamide, P-648
 1-Propylamine, P-648
 9-Propyl-10-azacyclododecan-12-olide, P-651
 12-Propyl-13-azacyclopentadec-5-en-15-olide, E-93
 11-Propyl-12-azacyclotetradeca-5,8-dien-14-olide, E-93
 11-Propyl-12-azacyclotetradec-5-en-14-olide, E-93
N-Propylbenzamide, P-648
 Propylene, P-649
N,N'-Propylenebis(cantharimide), P-650
N-Propylhydroxylamine, P-648
 5-Propylindolizidine, O-56
 3-Propylindolizidine, O-55
N-Propylisobutylamine, M-525
 5-Propyl-1-oxa-4-azacyclohexadec-11-en-16-one, E-93
 5-Propyl-1-oxa-4-azacyclopentadeca-7,10-dien-15-one, E-93
 5-Propyl-1-oxa-4-azacyclopentadec-10-en-15-one, E-93
 5-Propyl-1-oxa-4-azacyclotridecan-13-one, P-651
 13-Propyloxylupanine (incorr.), H-554
 1-(3-Propylphenyl)- β -carboline, K-59
 8-Propyl-10-phenyllobeliolone, D-403
 1-(3-Propylphenyl)-9*H*-pyrido[3,4-*b*]indole, K-59
 α -Propyl-2-piperidineethanol, H-37
 2-Propylpiperidine, P-652
 6-Propyl-3-piperidinol, P-653
 2-Propylquinoline, P-654
 2-Propyl-4-quinolinol, H-707
 2-Propyl-4(1*H*)-quinolinone, H-707
 3-Propylxanthine, X-5
 Prosafrine, P-655
 Prosafrinine, P-655
 Prosophylline, P-657
 Prosopine, P-656
 Prosopinine, P-657
 Prosopinone, P-656
 Prosthephanine, P-658
 Prosthephanaberrine, P-659
 Prosurugatoxin, N-141
 Protalba, P-674
 Protaxel, T-52
 Proteinase inhibitor E 64, R-69
 Prothracarcin, P-660
 Protocatechuic acid dimethyl ether, D-706
 Protocatechuic acid, D-536
 Protocevine, V-56
 Protochelin, P-661
 Protochondocurarine, P-662
 Protocuridine, P-663
 Protoemetine, P-664
 Protoemetinol, P-664
 Protogonyautoxin 2, H-731
 Protogonyautoxin 3, H-731
 Protogonyautoxin 4, H-731
 Protogonyautoxin I, H-731
 Protomethine, P-665
 Protomycin, S-563
 Protophomin, P-666
 Protopine *N*-oxide, P-667
 Protopine, P-667
 Protosinomenine, P-668
 Protostemediol, P-669
 Protostemonine, P-670
 Protostemotinine, M-49
 Protostephanine, P-671
 Protostrychnine, P-672
 Protothalpine, M-734
 Protoveratridine, G-70
 Protoveratrine A, P-674
 Protoveratrine B, P-674
 Protoveratrine C, P-674
 Protoveratrine, P-674
 Protoverbine, P-673
 Protoverine, P-674
 Proviolacein, P-675
 Proximicin A, P-676
 Proximicin B, P-676
 Proximicin C, P-676
 Proxiphomin, P-677
 Prulaurasin, H-673
 Prunasin, H-673
 Prunifoline A, M-296
 Prunifoline B, M-296
 Prunifoline C, M-296
 Prunifoline D, M-296
 Prunifoline E, M-296
 Prunifoline F, M-296
 Prurienidine, P-678
 Prurienine, P-679
 Prurieninine, P-680
 Przewalidine, P-681
 Przewaline, P-682
 Przewalskine, P-683
 Przewalskinine, P-683
 Przewalstidine, P-684
 Przewalstidinine, P-684
 Przewalstine, P-684
 Przewalstinine, P-684
 PS 258A, P-720
 PS 272A, P-720
 PS 286A, P-720
 PS 346A, P-720
 PS 258B, P-720
 PS 272B, P-722
 PS 286B, P-720
 PS 242, P-720
 PS 256, P-720
 PS 270, P-720
 PS 300, P-722
 PS 302, P-720
 PS 316, P-720
 PS 330, P-722
 PS 332, P-720
 PS 512, P-721
 PS 524, P-721
 PS 528, P-721
 PSA, A-759
 Psammaplin A 11'-sulfate, P-686
 Psammaplin A trisulfide, P-686
 Psammaplin A, P-686
 Psammaplin A₁, P-686
 Psammaplin A₂, P-686
 Psammaplin B, P-687
 Psammaplin C, P-688
 Psammaplin D, P-689
 Psammaplin E, P-690
 Psammaplin F, P-691
 Psammaplin G, P-692
 Psammaplin H, P-693
 Psammaplin I, P-685
 Psammaplin J, P-686
 Psammaplin K, P-686
 Psammaplin L, P-686
 Psammaplysene A, P-694
 Psammaplysene B, P-694
 Psammaplysene C, P-694
 Psammaplysene D, P-694
 Psammaplysin A, P-695
 Psammaplysin B, P-695
 Psammaplysin C, P-695
 Psammaplysin D, P-695
 Psammaplysin E, P-696
 Psammaplysin F, P-695
 Psammopemmin A, P-697
 Psammopemmin B, P-697
 Psammopemmin C, P-697
 PSB, A-759
 PSC₁, A-759
 PSC₂, A-759
 PSC-A, M-508
 PSC-B, H-126
 PSC-C, H-126
 PSC-D, M-491
 PSC-E, M-491
 Pseudaconine, M-336
 Pseudaconitine, M-336
 Pseudakuammicine, A-228
 Pseudane I, H-633
 Pseudane IX, H-642
 Pseudane VII, H-133
 Δ^1 -Pseudene IX, H-642
 Δ^1 -Pseudene VII, H-133
 Pseudoacetylpyrrole, A-74
 Pseudoaconitine, M-336
 Pseudoakuammicine, A-228
 Pseudoakuammigine *N*-oxide, A-232
 Pseudoakuammigine, A-232
 Pseudoanchnynazine A, P-698
 Pseudoanchnynazine B, P-699
 Pseudoanchnynazine C, P-700
 Pseudoanibacanine, P-717
 Pseudoanthraserpine, R-50
 Pseudoargiopinin I, A-1415
 Pseudoargiopinin II, A-1415
 Pseudoargiopinin III, P-701
 Δ^1 -Pseudoaspidospermidine, D-377
 Pseudobactin A, P-702
 Pseudobactin B, P-702
 Pseudobactin 7SR1, P-703
 Pseudobactin, P-702
 Pseudobaleabuxine F, P-704
 Pseudobaleabuxine, P-704
 Pseudoberberine, P-705
 Pseudobrucine, P-726
 Pseudocapsaicin, C-100
 Pseudocarpaine, C-168
 Pseudocatharanthine, P-733
 Pseudoceramine, M-679
 Pseudoceratin, P-706
 Pseudoceratin A, P-707
 Pseudoceratin B, P-707
 Pseudoceratinine A, P-812
 Pseudoceratinine B, P-708
 Pseudoceratinine C, P-709
 Pseudocheilanthifoline, C-685
 Pseudochelerythrine, S-56
 Pseudocinchonine, C-447
 Pseudococaine, C-537
 Pseudo- α -colubrine, P-726
 Pseudo- β -colubrine, P-726
 Pseudocolumbamine, P-719
 Pseudoconhydrine, P-653
 Pseudocoptisine, P-710
 Pseudocyclobuxine D, C-846
 Pseudodeoxyviolacein, P-711
 Pseudodistamine, O-38
 Pseudodistamin A, P-712
 Pseudodistamin B, P-712
 Pseudodistamin C, P-713
 Pseudodistamin D, P-712
 Pseudodistamin E, P-712
 Pseudodistamin F, P-713
 Pseudodysidenin, D-972
 Pseudoecgonine methyl ester, E-8
 Pseudoephedrine hydrochloride, M-386
 Pseudoephedrine sulfate, M-386
 Pseudoephedrine, M-386
 Pseudoepiisoeleocarpiline, P-714
 Pseudoepistephanine, E-102
 Pseudofluorocurine, F-101
 Pseudohomocorine, G-3
 Pseudohygroline, H-760
 (\pm)-Pseudohygroline, H-760
 3-Pseudoindolone, I-106
 Pseudoindoxyl, I-104
 Pseudoiodinine, M-285
 Pseudoisoplatydesmine, A-1377
 Pseudojatrorrhizine, P-719
 Pseudojervine, J-36
 Pseudokobusine, H-205
 Pseudokopsinine *N*^b-oxide, V-132
 Pseudokopsinine, V-132
 Pseudolaudanine, P-715
 Pseudolycorine 1,2-diglucoside, P-716
 Pseudolycorine-1-glucoside, P-716
 Pseudolycorine, P-716
 Pseudomanibacanine, P-717
 Pseudomerucathine, A-867
 Pseudomonine, P-718
 Pseudomorphone, M-703
 Pseudopalmatine, P-719
 Pseudopelletierine, A-1580
 Pseudophrynamine 258A, P-720
 Pseudophrynamine 272A, P-720
 Pseudophrynamine 286A, P-720
 Pseudophrynamine 346A, P-720
 Pseudophrynamine 512A, P-721

- Pseudophrynamine A, P-721
 Pseudophrynamine 258B, P-720
 Pseudophrynamine 272B, P-722
 Pseudophrynamine 286B, P-720
 Pseudophrynamine 242, P-720
 Pseudophrynamine 256, P-720
 Pseudophrynamine 270, P-720
 Pseudophrynamine 300, P-722
 Pseudophrynamine 302, P-720
 Pseudophrynamine 316, P-720
 Pseudophrynamine 330, P-722
 Pseudophrynamine 332, P-720
 Pseudophrynamine 524, P-721
 Pseudophrynamine 528, P-721
 Pseudophrynaminol, P-720
 Pseudoprotopine, P-723
 Pseudopteroxazole, P-724
 Pseudorepanduline, P-725
 Pseudoreserpine, R-52
 Pseudorine, P-715
 Pseudoronine, S-111
 Pseudoscopine, S-158
 Pseudoselagine, L-325
 Pseudosolanidine, P-633
 Pseudostrychnine, P-726
 Pseudotabersonine, P-733
 Pseudotheonamide A₁, P-727
 Pseudotheonamide A₂, P-727
 Pseudotheonamide B₂, P-728
 Pseudothiobinupharidine, P-729
 Pseudotiliarine, T-407
 Pseudotrienoic acid A, P-730
 Pseudotrienoic acid B, P-731
 Pseudotropine, M-393
 Pseudoverdin, A-738
 Pseudovinblastinediol, P-732
 Pseudovincadifformine, P-733
 Pseudovincal leukoblastine diol, P-732
 Pseudovincamine, T-12
 Pseudovincamone I, T-13
 Pseudoviolacein, P-711
 Pseudovobparicine, P-734
 Pseudowithanine, P-735
 Pseudoxandrine, P-736
 Pseudoxandrinine, P-736
 Pseudoyohimbine, Y-13
 Pseudozoanthoxanthin A, P-737
 Pseudozoanthoxanthin, P-737
 Pseurotin F₁, P-738
 Pseurotin F₂, P-738
 Pseurotins, P-738
 Psilocauline, P-739
 Psilocine, H-480
 Psilocybine, H-480
 Psilopine, A-1531
 PSVA, C-100
 Psycholeine, P-740
 Psychollatine, B-264
 Psychopentamine, P-741
 Psychotridine C, P-743
 Psychotridine, P-742
 Psychotrimine, P-744
 Psychotrine, E-72
 Psychrophilin A, P-745
 Psychrophilin B, P-746
 Psychrophilin C, P-747
 Psychrophilin D, P-748
 Psylloborine A, P-749
 Psymberin, I-173
 Pteleatine, B-14
 Pteleatinium, B-14
 Ptelecortine, P-750
 Ptelecultinium, B-14
 Pteleledimericine, P-751
 Pteleledimeridine, P-752
 Pteleledimerine, P-753
 Pteleflorine, P-754
 Ptelefoliarine, E-40
 Ptelefolidine, P-759
 Ptelefolidone, L-296
 Ptelefolidonium(1+), I-290
 Ptelefoline, P-755
 Ptelefolinol, P-755
 Ptelefolone, P-756
 Ptelefolonium, P-757
 Ptelefructine, P-750
 Pteleine, F-211
 Pteleoline, P-758
 Pteleprenine, P-759
 Pteriatoxin A, P-760
 Pteriatoxin B, P-761
 Pteriatoxin C, P-761
 2,4-Pteridinediol, P-762
 2,4(1*H*,3*H*)-Pteridinedione, P-762
 2,4,6(1*H*,3*H*,5*H*)-Pteridinetrione, D-494
 2,4,7(1*H*,3*H*,8*H*)-Pteridinetrione, P-763
 Pteridoxamine, A-816
 Pterin, A-816
 Pterin-6-carboxylic acid, A-817
 Pterocellin A, P-764
 Pterocellin B, P-765
 Pterocellin C, P-766
 Pterocellin D, P-767
 Pterocellin E, P-764
 Pterocellin F, P-768
 Pterocereine, T-178
 Pterogynidine, P-769
 Pterogynine, B-199
 Pterolactam, H-719
 Pterophorine, P-770
 Pteropodic acid, M-661
 Pteropodine *N*-oxide, M-661
 Pteropodine, M-661
 Pteropterin, P-772
 Pterorhodine, P-771
 Pteroyldiglutamic acid, P-772
 Pteroylglutamic acid, P-772
 Pteroylglutamylglutamic acid, P-772
 Pteroylglutamylglutamylglutamic acid, P-772
 Pteroylheptaglutamic acid, P-772
 Pteroylhexaglutamylglutamic acid, P-772
 Pteroylpentaglutamic acid, P-772
 Pteroyltriglutamic acid, P-772
 Ptilocaulin, P-773
 Ptilomycalin A, P-774
 Ptilomycalin D, P-775
 PTP, T-217
 PTX-A, P-791
 PTX-B, P-791
 PTX-C, P-794
 PTZ, P-626
 Pubamide, H-314
 Pubatriol, H-314
 Puberaconitidine, P-776
 Puberaconitine, P-776
 Puberanidine, A-91
 Puberanine, A-86
 Puberuline A, F-159
 Puberuline, F-159
 Puberullumine, T-578
 Pubesamide A, P-777
 Pubesamide B, P-777
 Pubesamide C, P-777
 Pubescenine, M-309
 Pubescimine, P-778
 Pubescine A, M-189
 Pubescine, P-779
 Pubescine, R-53
 Pubescinine, H-314
 Puertogaline A, S-502
 Puertogaline B, S-502
 Puhinamide, P-780
 Pukateine, P-781
 Pukeensine, P-782
 Pukeleimide A, P-783
 Pukeleimide B, P-783
 Pukeleimide C, P-783
 Pukeleimide D, P-783
 Pukeleimide E, P-783
 Pukeleimide F, P-783
 Pukeleimide G, P-783
 Pukeleimide, P-783
 Pulanomycin, F-48
 Pulchellalactam, P-784
 Pulchellidine, P-785
 Pulchelline, P-786
 Pulcheotine A, M-242
 Pulcherrimine, P-787
 Pulcherriminic acid, P-788
 Pulcherrimin, P-788
 Pulchine, T-524
 Pulouppone, P-789
 Pulvic acid, P-790
 Pulvinamide, P-790
 Pulvinic acid, P-790
 Pumiliotoxin 307A'', P-791
 Pumiliotoxin 307A', P-791
 Pumiliotoxin 341A, P-792
 Pumiliotoxin A, P-791
 Pumiliotoxin 307B, P-793
 Pumiliotoxin B, P-791
 Pumiliotoxin C_{II}, A-357
 Pumiliotoxin C_{III}, B-428
 Pumiliotoxin C, P-794
 Pumiliotoxin 251D, A-480
 Pumiliotoxin 209F, P-795
 Pumiliotoxin 225F, P-795
 Pumiliotoxin 307F, P-791
 Pumiliotoxin 321, P-791
 Pumiloside, P-796
 Pummeline, T-519
 Punarnavine, P-797
 Punctancine, P-798
 Punctatine, F-5
 Pumikathine, P-799
 Punjabine, P-800
 Puntarenine, P-801
 Puqiedine, C-290
 3 α -Puqiedin-7-ol, C-294
 Puqiedinone, C-290
 Puqienine A, P-802
 Puqienine B, P-802
 Puqienine C, P-803
 Puqienine D, P-804
 Puqienine E, P-805
 Puqienine F, P-806
 Puqietinedione, P-807
 Puqietinone, P-807
 Puqietinonoside, P-807
 PUR 1, P-808
 Purapuridine, S-456
 Puralidin A, P-811
 Puralidin B, A-1378
 Puralidin C, P-809
 Puralidin D, P-810
 Puralidin E, P-811
 Puralidin F, M-679
 Puralidin G, M-679
 Puralidin H, P-811
 Puralidin J, P-812
 Puralidin K, P-812
 Puralidin L, P-813
 Puralidin M, P-814
 Puralidin N, P-814
 Puralidin O, P-815
 Puralidin P, A-1378
 Puralidin Q, H-227
 Puralidin R, D-320
 Puralidin S, H-227
 Puraline, P-816
 1*H*-Purin-6-amine, A-137
 2,6-Purinediamine, P-817
 6,8-Purinediol, P-818
 2,6-Purinediol, X-5
Subergorgia suberosa Purine-imidazole alkaloid, P-819
 2,6,8(1*H*,3*H*,9*H*)-Purinetrione, U-44
 6-Purinol, P-820
 6(1*H*)-Purinone, P-820
 Purinosine, N-82
 Puroverin, P-674
 Purpeline, T-288
 Purpuramine A, P-821
 Purpuramine B, P-821
 Purpuramine C, P-822
 Purpuramine D, P-823
 Purpuramine E, P-823
 Purpuramine F, P-823

- Purpuramine G, P-823
 Purpuramine H, P-823
 Purpuramine I, P-823
 Purpuramine J, P-823
 Purpuramine K, P-823
 Purpuramine L, P-821
 Purpurealidin A, H-227
 Purpurealidin B, P-824
 Purpurealidin C, H-227
 Purpurealidin D, H-227
 Purpurealidin E, M-679
 Purpurealidin F, M-679
 Purpurealidin G, M-679
 Purpurealidin H, P-823
 Purpureine, P-208
 Purpuroceratic acid A, D-319
 Purpuroceratic acid B, D-319
 Purpurone, P-825
 Purtherianin, L-213
 Pusilline, I-322
 Puterine, P-781
 Putreanine, P-826
 Putrebactin, A-250
 Putrescine, B-422
 Putterine A, P-247
 Putterine B, P-247
 PwTx II, T-225
 Pycnamilline, C-806
 Pycnamine, P-301
 (+)-Pycnamine, T-286
 Pycnanthine, E-32
 Pycnanthine, P-827
 Pycnanthimine, P-828
 Pycnarrhenamine, P-829
 Pycnarrhenine, P-830
 Pycnarrhine, D-421
 Pycnazanthine, S-544
 Pycnosanguin, P-831
 Pyo Ib, H-133
 Pyo Ic, H-642
 Pyo II, H-133
 Pyo III, H-642
 Pyo V, H-757
 Pyochelin II, P-832
 Pyochelin I, P-832
 Pyocyanine, P-833
 Pyoluteorin, P-834
 Pyoverdin A1, P-838
 Pyoverdin BTB 16, P-870
 Pyoverdin BTP 2, P-839
 Pyoverdin BTP16G, P-842
 Pyoverdin BTP7G, P-841
 Pyoverdin BTP7, P-840
 Pyoverdin C 2A, P-870
 Pyoverdin C 3A, P-870
 Pyoverdin C 2B, P-870
 Pyoverdin C 3B, P-870
 Pyoverdin CFBP 2392, P-844
 Pyoverdin CFBP 2461, P-845
 Pyoverdin CHAO-A, P-846
 Pyoverdin CHAO, P-846
 Pyoverdin C₄₃H₆₀N₁₄O₁₇, P-874
 Pyoverdin C₄₄H₅₉N₁₃O₁₉, P-874
 Pyoverdin C₄₄H₆₁N₁₃O₁₇, P-847
 Pyoverdin C₄₄H₆₂N₁₄O₁₆, P-847
 Pyoverdin C₄₅H₆₃N₁₄O₂₄, P-848
 Pyoverdin C₄₅H₆₄N₁₂O₂₃, P-848
 Pyoverdin C₄₇H₆₉N₁₃O₁₈, P-867
 Pyoverdin C₅₂H₇₅N₁₅O₂₃, P-849
 Pyoverdin C₅₂H₇₆N₁₆O₂₂, P-849
 Pyoverdin C₅₅H₈₃N₁₇O₂₁, P-850
 Pyoverdin C₅₇H₈₅N₁₇O₂₃, P-851
 Pyoverdin C₅₇H₈₅N₁₇O₂₄, P-851
 Pyoverdin C, P-843
 Pyoverdin cyclo-Pf P19, P-703
 Pyoverdin D, P-852
 Pyoverdin E, P-852
 Pyoverdin G 173, P-853
 Pyoverdin GM II, P-854
 Pyoverdin GM I, P-854
 Pyoverdin G4RA, P-855
 Pyoverdin G4R, P-855
 Pyoverdin III, P-856
 Pyoverdin II, P-856
 Pyoverdin I, P-835
 Pyoverdin 11370, P-837
 Pyoverdin 2112, P-836
 Pyoverdin Pa II, P-857
 Pyoverdin Pa I, P-857
 Pyoverdin Pa TII, P-858
 Pyoverdin PaA, P-852
 Pyoverdin PaB, P-843
 Pyoverdin PaC, P-859
 Pyoverdin PaO1, P-852
 Pyoverdin Pa, P-852
 Pyoverdin Pf CCM 2798, P-864
 Pyoverdin Pf G 173, P-853
 Pyoverdin Pf 1.3, P-860
 Pyoverdin Pf 1547, P-863
 Pyoverdin Pf/3 17400, P-861
 Pyoverdin Pf P19, P-865
 Pyoverdin Pf12, P-862
 Pyoverdin PL7, P-866
 Pyoverdin PL8, P-867
 Pyoverdin Pm, P-868
 Pyoverdin Pp 1, P-869
 Pyoverdin Pp 2, P-869
 Pyoverdin PpC 2A, P-870
 Pyoverdin PpC 3A, P-870
 Pyoverdin PpC 2B, P-870
 Pyoverdin PpC 3B, P-870
 Pyoverdin PS 6.10, P-872
 Pyoverdin PS, P-871
 Pyoverdin PSS, P-873
 Pyoverdin R', P-874
 Pyoverdin 51WA, P-875
 Pyoverdin WCS358, P-845
 Pyoverdin 51W, P-875
 Pyoverdin 51W-Suca, P-875
 Pyoverdin 51W-Suc, P-875
 Pyoverdin, P-848
 Pyracrimycin A, P-876
 Pyracrimycin B, P-877
 Pyrafoline D, M-46
 Pyramidaglaïne A, P-878
 Pyramidaglaïne B, P-878
 Pyramidatine, C-456
 Pyranofoline, P-879
 5,6-Pyranoglycozoline, E-300
 Pyranonigrin A, P-880
 Pyranonigrin B, P-881
 Pyranonigrin C, P-881
 Pyranonigrin D, P-882
 Pyranonigrin S, P-880
 Pyraxinine, G-209
 Pyrayafoline A, P-883
 Pyrayafoline B, P-884
 Pyrayafoline C, P-883
 Pyrayafoline D, P-885
 Pyrayaquinone A, P-886
 Pyrayaquinone B, P-887
 Pyrayaquinone C, P-888
Vibrio alginolyticus Pyrazine, B-221
 1,1'-(2,5-Pyrazinediyl)bis-1,2,3,4-butanetetrol, B-221
 Pyrazinol, P-889
 2(1*H*)-Pyrazinone, P-889
 Pyrazofurin B, P-891
 Pyrazofurin, P-891
 3(5)-Pyrazolecarboxylic acid, P-890
 Pyrazomycin B, P-891
 Pyrazomycin, P-891
 Pyrenoline A, P-892
 Pyrenoline B, P-892
 Pyrettrin, D-106
 Pyrichalasin H, Z-37
 Pyriculamide, C-900
 Pyricularin, P-478
 Pyridindolol K1, P-893
 Pyridindolol K2, P-893
 Pyridindolol, P-893
 Pyridinebetaine A, P-897
 Pyridinebetaine B, S-636
 3-Pyridinecarboxamide, P-895
 4-Pyridinecarboxylic acid, P-898
 2-Pyridinecarboxylic acid, P-896
 3-Pyridinecarboxylic acid, P-897
 2,6-Pyridinedicarbothioic acid, P-899
 3,5-Pyridinedicarboxamide, P-902
 2,3-Pyridinedicarboxylic acid, P-900
 2,6-Pyridinedicarboxylic acid, P-901
 3,5-Pyridinedicarboxylic acid, P-902
 Pyridine-2,6-di(monothiocarboxylic acid), P-899
 1,1'-(3,5-Pyridinediyl)bis-1-decanone, D-371
 3-Pyridinedodecanal, P-903
 3-Pyridinemethanol, P-904
 Pyridine, P-894
 3-Pyridinetricecanal, P-905
 3-Pyridinetridecanamine, T-358
 2-Pyridinioethanesulfonic acid, S-636
 Pyridinoline, P-907
 3-Pyridinol, P-906
 3-(2-Pyridinyl)alanine, A-902
 10-(3-Pyridinyl)-3,5,7,9-decatetraen-2-one, P-908
 3-[10-(3-Pyridinyl)-7-decenyl]-1-[11-(3-pyridinyl)-8-decenyl]pyridinium(1+), N-230
 3-[10-(3-Pyridinyl)-7-decenyl]-1-[11-(3-pyridinyl)-9-undecenyl]pyridinium(1+), N-231
 12-(3-Pyridinyl)-4,6-dodecadien-2-ol, P-911
 12-(2-Pyridinyl)-6,8-dodecadien-2-ol, P-909
 12-(3-Pyridinyl)dodecanal, P-903
 12-(2-Pyridinyl)-3,5,9-dodecatrien-1-amine, A-760
 12-(3-Pyridinyl)-2,4,6-dodecatrien-1-ol, P-910
 12-(3-Pyridinyl)-4,6,8-dodecatrien-2-ol, P-911
 12-(3-Pyridinyl)-5,8,10-dodecatrien-2-ol, P-913
 12-(3-Pyridinyl)-4,6,11-dodecatrien-2-ol, P-912
 1-(2-Pyridinyl)ethanone, A-73
 3-Pyridinylguanidine, G-209
 16-(3-Pyridinyl)-7-hexadecynal *O*-methyloxime, N-228
 16-(3-Pyridinyl)-7-hexadecyn-1-amine, N-224
 4-[2-(3-Pyridinyl)-5-oxazolyl]phenol, H-12
 2-(3-Pyridinyl)piperidine, P-460
 1-(2-Pyridinyl)-1-propanone, P-645
 2-(3-Pyridinyl)pyrrole, P-972
 2-(3-Pyridinyl)-1-pyrrolidinedicarboxaldehyde, P-957
 (4-Pyridinyl)pyruvic acid, O-205
 14-(3-Pyridinyl)-11-tetradecen-1-amine, T-357
 14-(3-Pyridinyl)-11-tetradecynal *O*-methyloxime, N-227
 14-(3-Pyridinyl)-11-tetradecyn-1-amine, T-357
 13-(3-Pyridinyl)tridecanal, P-905
 12*H*-Pyrido[1,2-*ac*:3,4-*b'*]diindole-13-carboxaldehyde, H-336
 Pyridofylline, P-923
 9*H*-Pyrido[3,4-*b*]indole, C-130
 9*H*-Pyrido[3,4-*b*]indole-1-carboxaldehyde, C-131
 9*H*-Pyrido[3,4-*b*]indole-1-carboxylic acid, C-132
 9*H*-Pyrido[3,4-*b*]indole-3-carboxylic acid, C-133
 9*H*-Pyrido[3,4-*b*]indole-1-ethanol, H-487
 9*H*-Pyrido[3,4-*b*]indole-1-methanol, H-590
 9*H*-Pyrido[3,4-*b*]indole-1-propanoic acid, C-134
 9*H*-Pyrido[3,4-*b*]indole-1,3,4(2*H*,9*H*)-trione, C-135
 9*H*-Pyrido[3,4-*b*]indol-7-ol, P-916
 9*H*-Pyrido[3,4-*b*]indol-1-ol, P-914
 9*H*-Pyrido[3,4-*b*]indol-6-ol, P-915
 9*H*-Pyrido[3,4-*b*]indol-8-ol, P-917
 1-[4-(9*H*-Pyrido[3,4-*b*]indol-1-ylcarbonyl)-3-pyridinyl]ethanone, P-133
 1-(9*H*-Pyrido[3,4-*b*]indol-1-yl)ethanone, A-44
 5-(5*H*-Pyrido[4,3-*b*]indol-1-yl)-2-furanmethanol, I-262
 5-(9*H*-Pyrido[3,4-*b*]indol-1-yl)-2-furanmethanol, P-270
 4-(9*H*-Pyrido[3,4-*b*]indol-1-yl)-2-pyridinamine, A-1051
 3-Pyridol, P-906
 Pyridomacrolidin, P-918
 Pyridomycinol, P-919
 Pyridomycin, P-919
 2-Pyridone-6-acetic acid, H-708
 6(1*H*)-Pyridone-3-carboxylic acid, H-711
 α-Pyridone-5-carboxylic acid, H-711
 4(1*H*)-Pyridone-3-carboxylic acid, H-709
 Pyrido[4,3,2-*mn*]pyrrolo[3,2,1-*de*]acridin-4-ol, A-1445
 Pyridostemine, S-514
 9*H*-Pyrido[4,3,2-*mn*]thiazolo[4,5-*b*]acridin-9-one, P-920

- Pyridovericin, P-921
 Pyridoxatin, P-922
 Pyridoxine hydrochloride, P-923
 Pyridoxine 5'-phosphate, P-923
 Pyridoxine, P-923
 Pyridoxol dihydrogen phosphate, P-923
 Pyridoxol, P-923
 Pyridoxylate, P-923
 2-(3-Pyridyl)-5-(4-methoxyphenyl)oxazole, H-12
 2-(3-Pyridyl)pyrrolidine, P-957
 2-(2-Pyridyl)- Δ^1 -pyrroline-5-carboxylic acid, D-496
 2-(2-Pyridyl)-1-pyrroline, D-504
 2-(3-Pyridyl)-1-pyrroline, M-792
 3-(3-Pyridyl)tridecylamine, T-358
 Pyrifolidine, P-924
 Pyrifoline, K-78
 2,4-Pyrimidinediol, U-42
 2,4(1*H*,3*H*)-Pyrimidinedione, U-42
N-(2-Pyrimidinyl)-1,4-benzenediamine, A-863
 4*H*,6*H*,8*H*-Pyrimido[5,4-*e*]-*as*-triazine-3,5,7-trione, D-498
 Pyrimine, D-496
 Pyrinadine A, P-925
 Pyrinadine B, P-926
 Pyrinadine C, P-927
 Pyrinadine D, P-928
 Pyrinadine E, P-929
 Pyrinadine F, P-930
 Pyrinadine G, P-931
 Pyrindamycin A, D-954
 Pyrindamycin B, D-953
 Pyrindamycin C, D-954
 Pyrindicine, P-932
 Pyrinodemin A, P-933
 Pyrinodemin B, P-934
 Pyrinodemin C, P-934
 Pyrinodemin D, P-934
 Pyripyropene A, T-500
 Pyripyropene B, T-500
 Pyripyropene C, T-500
 Pyripyropene D, T-500
 Pyripyropene E, T-500
 Pyripyropene F, T-500
 Pyripyropene G, T-500
 Pyripyropene H, T-500
 Pyripyropene I, T-500
 Pyripyropene J, T-500
 Pyripyropene K, T-500
 Pyripyropene L, T-500
 Pyripyropene M, T-500
 Pyripyropene N, T-500
 Pyripyropene O, T-500
 Pyripyropene P, T-500
 Pyripyropene Q, T-500
 Pyripyropene R, T-500
 Pyrisulfoxin A, C-574
 Pyrisulfoxin B, C-576
 Pyrizinostatin, P-935
 Pyro-Ace, P-961
 Pyroaconitine, H-122
 Pyrochasmaconitine, M-372
 Pyrochasmaanine, M-377
 Pyroclavine, F-49
 Pyrocoll, D-833
 Pyrocrassicauline A, M-372
 Pyrodelphinine, M-372
 Pyrodysinoic acid, P-936
 Pyroglutamic acid, O-206
 Pyroglutamylglucosamine, P-937
 Pyrohyapaconitine, H-254
 Pyroiobotic acid, A-837
 Pyrojesaconitine, H-122
 Pyromesaconitine, H-122
 Pyronaamidine 9-*N*-methylimide, P-938
 Pyronaamidine, P-938
 Pyronopaline, P-939
 Pyrostatin A, T-185
 Pyrostatin B, T-207
 Pyrrocidine A, P-940
 Pyrrocidine B, P-940
 Pyrroindomycin A, P-941
 Pyrroindomycin B, P-941
 Pyrrolallosedamine, P-970
 Pyrrolam A, T-229
 Pyrrolam B, H-234
 Pyrrolam C, H-234
 Pyrrolam D, H-234
 3-Pyrroleacrylic acid, P-971
 Pyrrole-2-aldehyde, P-942
 Pyrrole-3-carbamidine, P-943
 1*H*-Pyrrole-2-carbonitrile, P-944
 4-(2-Pyrrolocarbonyloxy)epilupinine, H-482
 1*H*-Pyrrole-2-carboxaldehyde, P-942
 1*H*-Pyrrole-2-carboxamide, P-944
 1*H*-Pyrrole-3-carboximidamide, P-943
 1*H*-Pyrrole-2-carboxylic acid, P-944
 1*H*-Pyrrole-2,3-dicarboxylic acid, P-945
 1*H*-Pyrrole-2,5-dicarboxylic acid, P-946
 1*H*-Pyrrole-2,5-dione, P-947
 Pyrrole-1-norleucine, A-904
 1*H*-Pyrrole-3-propanamide, P-948
 1*H*-Pyrrole-3-propanoic acid, P-948
 Pyrrolezanthine, H-626
 1-Pyrrolidinamine, P-949
 2-Pyrrolidineacetic acid, P-950
 2-Pyrrolidine-2-acrylamide, P-876
 1-Pyrrolidinecarboxamide, P-951
 3-Pyrrolidinecarboxylic acid, P-952
 2,4-Pyrrolidinedicarboxylic acid, P-953
 2,4-Pyrrolidinedione, P-954
 Pyrrolidine, P-949
 2-Pyrrolidinol, P-955
 2-Pyrrolidinone, P-956
 2-(2-Pyrrolidinyl)-3',4'-dimethoxyacetophenone, R-165
 2-(2-Pyrrolidinyl)-4'-hydroxy-3'-methoxyacetophenone, R-165
 3-(2-Pyrrolidinyl)pyridine, P-957
 13-(2-Pyrrolidinyl)-2-tridecylamine, A-921
 11-(2-Pyrrolidinyl)-2-undecylamine, P-958
 2-Pyrrolidone-5-carboxylic acid, O-206
Asteronotus cespitosus Pyrrolidone, P-959
 2-Pyrrolidone, P-956
N-Pyrrolidyleicosa-2,4-dienamide, T-491
 1-Pyrroline-5-carboxylic acid, D-501
 2-Pyrroline, D-499
 1-Pyrroline, D-500
 3-(1-Pyrroline-2-yl)acrylamide, P-876
 1-(1-Pyrroline-2-yl)- β -carboline, E-288
 3-(2-Pyrroline-2-yl)-2-propenamide, P-876
 2-(1-Pyrroline-2-yl)pyridine, D-504
 3-(1-Pyrroline-2-yl)pyridine, M-792
 Pyrrolizidine 223H', H-135
 Pyrrolizidine 265H', H-500
 Pyrrolizidine 267H', H-500
 Pyrrolizidine 239K', H-135
 1-Pyrrolizidinecarboxylic acid, P-960
 10*H*-Pyrrolizino[1,2-*b*]quinoline-9,10(4*H*)-dione, Q-24
 Pyrrolnitrin, P-961
 Pyrrolo[3,4-*d*]imidazole-4,6(1*H*,5*H*)-dione, P-962
 Pyrrolomycin A, D-356
 Pyrrolomycin B, P-963
 Pyrrolomycin C, P-964
 Pyrrolomycin D, P-964
 Pyrrolomycin E, P-965
 Pyrrolomycin F, P-966
 Pyrrolomycin G, P-967
 Pyrrolomycin H, P-967
 Pyrrolomycin I, P-964
 Pyrrolomycin J, P-964
 5*H*-Pyrrolo[3,4-*d*]piperidine-2-carboxylic acid, T-230
 Pyrrolo[1,2-*a*]pyrazine-3,6(4*H*,7*H*)-dione, P-968
 Pyrroloquinolinequinone, M-236
 Pyrrolosporin A, P-969
 Pyrrolostatin, D-761
 Pyrrolotenin, L-115
 Pyrrolsedamine, P-970
 1*H*-Pyrrol-2-ylcarbonyl α -L-rhamnopyranoside, P-944
 13-(2-Pyrrolylcarbonyl)calpurmenine, C-40
O-(2-Pyrrolylcarbonyl)virgiline, V-153
 2-(2-Pyrrolyl)-1,3-dithiolane, P-942
 1-(1*H*-Pyrrol-2-yl)ethanone, A-74
 5-(2*H*-Pyrrol-2-ylidene)methyl)-2,2'-bi-1*H*-pyrrole, P-635
*N*⁶-Pyrrolylnorleucine, A-904
 3-(1*H*-Pyrrol-3-yl)-2-propenoic acid, P-971
 3-(3-Pyrrolyl)propionic acid, P-948
 3-(2-Pyrrolyl)pyridine, P-972
 1-(1*H*-Pyrrol-2-yl)-9*H*-pyrido[3,4-*b*]indol-6-ol, E-285
 Pyrronamycin A, P-973
 Pyrronamycin B, P-973
 Pyrroperine, T-495
 Pyrrothine, A-759
 Pyrroxamycin, P-974
 Pyruvodehydrase, T-366
 6-Pyruvyl-5,6,7,8-tetrahydropterin, T-142
 6-Pyruvyltetrahydropterin, T-142
 Py-tetrahydroserpentine, A-212
 Q 11270A, A-1164
 Q 20547A, A-1311
 Q 11270B, A-1164
 Q base, Q-9
 Q 20547C, A-1310
 Qasmine A, A-874
 Qasmine B, A-876
 Q1, H-115
 Qingdainone, C-77
 Quadrigemine A, Q-2
 Quadrigemine B, Q-3
 Quadrigemine C, Q-2
 Quadrigemine H, Q-4
 Quadrigemine I, Q-1
 Quaternatine, M-292
 Quaternidine, P-417
 Quaternine, P-417
 Quaternoline, R-21
 Quaternoxine, C-191
 3,2':3',4':2'':3''':3''''-Quaterpyridine, Q-5
 Quebrachamine, Q-6
 Quebrachidine, V-113
 Quebrachine, Y-13
 Quench spot, A-52
Aesculus hippocastanum Quercetin indoxylacetyl-glycoside, Q-7
 Questionomycin A, A-862
 Questionomycin B, A-861
 Quettamine, Q-8
 Queuine, Q-9
 Queuosine, Q-9
 Quimbeline, Q-10
 Quinaglate, Q-18
 Quinaldic acid, Q-31
 Quinaldine, M-557
 Quinaldineic acid, Q-31
 α -Quinaldinol, Q-37
 γ -Quinaldine, H-633
 Quinaldopeptin, Q-11
 Quinaldyl acetate, Q-37
*N*²-Quinaldylarginine, Q-56
 Quinamine, Q-12
 Quinanium, Q-20
 2,4-Quinazolinediol, Q-13
 2,4(1*H*,3*H*)-Quinazolinone, Q-13
 4-Quinazolinol, Q-14
 4(1*H*)-Quinazolinone, Q-14
 Quinazolone, Q-14
 Quinazomycin, Q-15
 Quindoline-11-carboxylic acid, I-111
 Quindoline, I-109
 Quindolinocryptotackieine, Q-16
 Quindolinone, D-505
 Quine, Q-20
 Quinethylene, Q-20
 Quinicine, Q-17
 Quinidex, Q-18
 Quinidine gluconate, Q-18
 Quinidine sulfate, Q-18
 α -Quinidine, C-446
 Quinidine, Q-18
 Quinidinone, Q-18
 Quinindoline, Q-19
 Quinindolin-11(5*H*)-one, D-506
 Quinine ascorbate, Q-20
 Quinine biascorbate, Q-20

- Quinine bisulfate, Q-20
 Quinine ethylcarbonate, Q-20
 Quinine hydrochloride, Q-20
 Quinine sulfate, Q-20
 Quinine, Q-20
 β-Quinine, Q-18
 Quinic acid, H-726
 Quinite, Q-20
 Quinocarcinol, Q-22
 Quinocarcin, Q-21
 Quinocitrinine A, Q-23
 Quinocitrinine B, Q-23
 Quinocycline B, I-300
 Quinolactacide, Q-24
 Quinolactacin A (obsol.), Q-25
 Quinolactacin A₁, Q-25
 Quinolactacin A₂, Q-25
 Quinolactacin B, Q-26
 Quinolactacin B₁, Q-26
 Quinolactacin B₂, Q-26
 Quinolactacin C₁, Q-25
 Quinolactacin C₂, Q-25
 Quinolactacin D₁, Q-27
 Quinolactacin D₂, Q-27
 2-Quinolamine, Q-28
 4-Quinolinecarboxaldehyde, Q-30
 4-Quinolinecarboxylic acid, Q-32
 2-Quinolinecarboxylic acid, Q-31
 4,8-Quinolinediol, Q-35
 2,4-Quinolinediol, Q-33
 3,8-Quinolinediol, Q-34
 2-Quinolinemethanethiol, Q-36
 4-Quinolinemethanol, Q-38
 2-Quinolinemethanol, Q-37
 2,3,4,7,8-Quinolinediol, Q-39
 Quinoline, Q-29
 2,4,7,8-Quinolinetetrol, Q-43
 2,3,4,7-Quinolinetetrol, Q-40
 2,3,4,8-Quinolinetetrol, Q-41
 2,4,6,8-Quinolinetetrol, Q-42
 3,4,5,8-Quinolinetetrol, Q-44
 2,3,4-Quinolinetriol, Q-45
 4,7,8-Quinolinetriol, Q-48
 2,4,6-Quinolinetriol, Q-46
 2,4,8-Quinolinetriol, Q-47
 Quinolinic acid, P-900
 3-Quinololinol, Q-49
 2-Quinololinol, Q-50
 2(1*H*)-Quinolinsonone-4-carboxylic acid, H-723
 4(1*H*)-Quinolinsonone-2-carboxylic acid, H-725
 2(1*H*)-Quinolinsonone, Q-50
 1-(2-Quinolinyloxy)-β-carboline, Q-51
 1-(4-Quinolinyloxy)-β-carboline, Q-52
 1-(5-Quinolinyloxy)-β-carboline, Q-53
 1-(6-Quinolinyloxy)-β-carboline, Q-54
 1-(8-Quinolinyloxy)-β-carboline, Q-55
*N*²-[(2-Quinolinyloxy)carbonyl]arginine, Q-56
 3-(3-Quinolinyloxy)-1*H*-indole, I-138
S-(2-Quinolinyloxy)methyl thioacetate, Q-36
 4-(3-Quinolinyloxy)phenol, H-692
 1-(2-Quinolinyloxy)-9*H*-pyrido[3,4-*b*]indole, Q-51
 1-(5-Quinolinyloxy)-9*H*-pyrido[3,4-*b*]indole, Q-53
 1-(8-Quinolinyloxy)-9*H*-pyrido[3,4-*b*]indole, Q-55
 1-(4-Quinolinyloxy)-9*H*-pyrido[3,4-*b*]indole, Q-52
 1-(6-Quinolinyloxy)-9*H*-pyrido[3,4-*b*]indole, Q-54
 Quinolizidine 196, A-907
 Quinolizidine 233A, E-264
 Quinolizidine 217A, M-507
 Quinolizidine 195C, M-532
 Quinolizidine 207I, E-265
 Quinolobactin, D-677
Oceanapia Quinolone alkaloid, Q-57
 Quinolonic acid, H-728
 Quinolonomide, H-728
 α-Quinololcarbinol, Q-37
 2-Quinololnitramine, Q-28
 Quinomycin A, Q-58
 Quinomycin B₀, Q-59
 Quinomycin B, Q-60
 Quinomycin C, Q-61
 Quinomycin D, Q-62
 Quinomycin E, Q-63
 Quinonamine A, S-2
 Quinonamine B, S-2
 9*H*-Quino[4,3,2-*de*][1,10]phenanthroline-9-one, A-1459
 Quino[2',3':3,4]pyrrolo[2,1-*b*]quinazolin-11(13*H*)-one, L-298
 Quinotidine, Q-18
 Quinotoxine, Q-17
 2'-*L*-Quinovosyl saphenate, H-496
 3'-*L*-Quinovosyl saphenate, H-496
 Quinoxapeptin A, Q-64
 Quinoxapeptin B, Q-64
 R 39FF, M-430
 R 39MF, A-1230
 R 39SF, A-1231
 Racemigerine, R-1
 Racemocine, H-233
 Racemodine, S-90
 Racemonecine, H-233
 Racemoline, H-233
 Racemazine, H-233
 Racemulodine, H-161
 Racemuloline A, M-325
 Racemuloline B, M-354
 Racemulosine, R-2
 Racemulotone, R-3
 Racephedrine, M-386
 Racepinefrine, A-152
 Racepinephrine hydrochloride, A-152
 Racepinephrine, A-152
 Rachelmycin, R-4
 Radamycin, R-5
 Radarin A, R-6
 Radarin B, R-6
 Radarin C, R-6
 Radarin D, R-6
 Raddeamine, R-7
 Raddeanamine, R-8
 Raddeanidine, R-9
 Raddeanine, C-298
 Raddeanine, R-9
 Raddeanone, R-9
 Radiatine, U-37
 Radicamine A, R-10
 Radicamine B, R-10
 Radixisatidis A, R-11
 Radpetine, R-12
 Rakicidin C, R-13
 Ramiflorine A, U-52
 Ramiflorine B, U-52
 Ramiopterin, S-246
 Ramosine A, R-14
 Ramosine B, H-105
 Ramosine C, L-267
 Ranachrome 1, A-749
 Ranachrome 3, A-802
 Ranachrome 4, A-751
 Ranachrome 5, A-817
 Ranaconine, A-86
 Ranaconitine, A-86
 Rankinidine, H-367
 Rankiniridine, R-15
 Raocyclamide A, R-16
 Raocyclamide B, R-17
 Rapalexin A, I-337
 Rapalexin B, H-548
 Raphanatin, Z-11
 Raphantin, R-18
 Raphanusamic acid, T-394
 Raphanusamide, M-266
 Raphanusanin, R-18
 Rapinic acid, O-37
 Ratanhin, T-700
 Raubasine, A-212
 Raubasinine, R-53
 Raucaffricine, V-194
 Raucaffridine, R-19
 Raucaffriline, V-194
 Raucaffrine, P-240
 Raucaffrinoline, P-240
 Raucanine, R-20
 Raucubaine, R-21
 Raucubainine, C-191
 Rauflexine, T-288
 Raufloxicine, A-231
 Raufloridine, R-53
 Rauflorine, T-288
 Raugalline, A-216
 Rauglucine, A-216
 Raugustine, R-50
 Rauhimbine, Y-13
 Raujemidine *N*-oxide, D-233
 Raujemidine, D-233
 Raumacline, R-22
 Raumitorine, A-1423
 Raunamine, R-23
 Raunescine, C-81
 Raunescine, D-687
 Rauniticine oxindole A, M-661
 Rauniticine pseudoindoxyl, T-289
 Rauniticine, A-212
 Raunitidine, R-53
 Raunormic acid, D-687
 Raunormine, D-234
 Raupine, S-89
 Raureflexine, R-24
 Rausutranine, R-25
 Rausutrine, R-26
 Ravanine, R-51
 Raviridine, R-27
 Ravolcinine, R-28
 Ravomitine, T-288
 Ravoxine, C-117
 Ravoxinine, C-117
 Rauwolfine, A-216
 Rauwolfinine, R-29
 Rauwolscine, Y-13
 Ravenic acid, R-30
 Ravenine, H-53
 Ravenoline, R-31
 Ravesilone, F-90
 Raveyine, M-352
 Ravifoline, S-456
 Razadyne, G-3
 6*R*-BH₄, T-142
 RCNS1, D-374
 Rebeccamyacin, R-32
 Recanescic acid, D-687
 Recanescine, D-234
 Recinine, R-96
 Reductiline, R-33
 Reductiomycin, R-34
 Reevesianine A, H-685
 Reevesianine B, H-685
 Reflexine, T-288
 Refolinon, F-121
 Refractamine, R-36
 Refractidine, K-78
 Refractine, K-78
 (*R*)-Reframidine *N*-oxide, R-35
 (*S*)-Reframidine *N*-oxide, R-35
 Reframidine, R-35
 Reframine *N*-methosalt, R-36
 Reframine, R-36
 Reframoline, R-36
 Refuin, A-1081
 Regalamine, K-36
 Regaline, M-347
 Regecoline, R-37
 Regelamine, K-36
 Regelidine, T-257
 Regeline, K-36
 Regelimine, K-36
 Regelinone, R-38
 Regholarrhenine A, H-314
 Regholarrhenine B, H-314
 Regholarrhenine C, R-39
 Regholarrhenine D, C-595
 Regholarrhenine E, K-113
 Regholarrhenine F, D-279
 Regilimine, K-36
 Rehybrine, R-117
 Reissantin A, P-219
 Reissantin B, H-246
 Reissantin C, P-219
 Reissantin F, H-249
 Reissantin G, P-219

- Reissantin H, P-219
 Remeridine, R-116
 Remerine, R-120
 Remeroline, A-1066
 Remestyp, A-736
 Remijinine, R-40
 Reminyl, G-3
 Remrefidine, R-120
 Remrefine, R-36
 Renardine, S-245
 Renieramycin A, R-41
 Renieramycin B, R-41
 Renieramycin C, R-41
 Renieramycin D, R-41
 Renieramycin E, R-41
 Renieramycin F, R-41
 Renieramycin G, R-41
 Renieramycin H, R-42
 Renieramycin I, R-41
 Renieramycin J (obsol.), R-44
 Renieramycin J, R-43
 Renieramycin K, R-43
 Renieramycin L, R-43
 Renieramycin M, R-41
 Renieramycin N, R-41
 Renieramycin O, R-41
 Renieramycin P, R-44
 Renieramycin Q, R-41
 Renieramycin R, R-41
 Renieramycin S, R-41
 Renierol acetate, R-45
 Renierol propionate, R-45
 Renierol, R-45
 Renierone, R-45
 Renoxidine, R-52
 Renoxydine, R-52
 Repandine, R-46
 Repandinine, T-81
 Repanduline, R-47
 Resactin, D-765
 Rescaloid, R-50
 Rescidine, R-50
 Rescinnamidine, R-50
 Rescinnamine, R-50
 Rescinnaminol, R-48
 Rescufolin, F-121
 Resedine, P-347
 Resedinine, A-864
 Resedinine, P-346
 Reserpenediol, R-49
 Reserpine acid, R-50
 Reserpine alcohol, R-49
 Reserpidine, D-234
 Reserpiline (4R)-*N*-oxide, R-51
 Reserpiline (4S)-*N*-oxide, R-51
 Reserpiline oxindole A, C-117
 Reserpiline oxindole B, C-117
 Reserpiline, R-51
 Reserpine, R-52
 ψ -Reserpine, R-52
 Reserpinine oxindole, C-1
 Reserpinine, R-50
 Reserpinine, R-53
 Reserpinolic acid, R-50
 Reserpoid, R-52
 Reserpoxidine, R-52
 Resistaphyllin, O-155
 Resorcein, A-711
 Resorcinol-4-carboxylic acid, D-535
 Resorcinomycin A, R-54
 Resorcinomycin B, R-55
N-Resoreylantranilic acid, B-67
 β -Resoreylic acid, D-535
 Retamine, R-56
 Reticulate, R-57
 Reticulatine A, R-57
 Reticulatine B, R-57
 Reticulate, R-57
 Reticulatol, R-58
 Reticuline *N* ^{α} -oxide, R-59
 Reticuline *N* ^{β} -oxide, R-59
 Reticuline, R-59
 Retinylidenetaurine, T-38
 Retrofractamide C, R-60
 Retrofractamides, R-60
 Retrohoustine, T-188
 Retrosiosenine *N*-oxide, N-89
 Retrosiosenine, N-89
 Retronecanol, H-630
 Retronecine 9-(3-acetoxy-2-hydroxy-2-methylbutanoate) 7-senecioate *N*-oxide, R-61
 Retronecine 9-(3-acetoxy-2-hydroxy-2-methylbutanoate) 7-senecioate, R-61
 Retronecine 7-*O*-curassavate, T-188
 Retronecine 9-*O*-curassavate, T-188
 Retronecine 9-(2,3-dihydroxy-2-hydroxymethylbutanoate) 7-senecioate, R-61
 Retronecine 9-(2,3-dihydroxy-2-methylbutanoate) 7-senecioate *N*-oxide, R-61
 Retronecine 9-(2,3-dihydroxy-2-methylbutanoate) 7-senecioate, R-61
 Retronecine *N*-oxide, T-188
 Retronecine 9-sarracinate 7-senecioate *N*-oxide, R-61
 Retronecine 9-sarracinate 7-senecioate, R-61
 Retronecine-7,9-dibenzoate, T-188
 Retronecine, T-188
 Retrorsine *N*-oxide, R-62
 Retrorsine, R-62
 Retulinal, R-63
 Retuline *N* ^{β} -oxide, R-63
 Retuline, R-63
 Retusamine, R-64
 Retusine, R-65
 Reumitsin, R-66
 Reumycin, R-66
 Reutericyclin, R-67
 Revolutinone, R-68
 Revolutopine, T-333
 Rexostatine, R-69
 RGH4406, E-6
 Rhamnoerysodine, E-203
 4-Rhamnopericidin A₁, P-421
 α -L-Rhamnopyranosyl anthranilate, A-706
 6-*O*- α -L-Rhamnopyranosyl-1-deoxymannonojirimycin, H-616
 (4'- α -L-Rhamnosyloxycinnamoyl)lupinine, O-58
 (4'- α -L-Rhamnosyloxy-3'-methoxycinnamoyl)lupinine, O-58
 Rhamnoveracintine, V-57
 Rhapallin A, R-70
 Rhapsamine, R-71
 Rhazicine *N*-oxide, L-28
 Rhazicine, L-28
 Rhazidigenine *N* ^{β} -oxide, R-72
 Rhazidigenine, R-72
 Rhazidine base, R-72
 Rhazidine salt, R-72
 Rhazidine, R-72
 Rhazimal, R-75
 Rhazimanine, I-319
 Rhazimine, R-73
 Rhazimol, A-231
 Rhazinaline *N* ^{β} -oxide, R-75
 Rhazinaline, R-75
 Rhazinal, R-74
 Rhazine, A-229
 Rhazinicine, R-76
 Rhazinilam, R-76
 Rhazinine, A-1328
 Rhazinoiline, S-576
 Rhazinol, S-576
 Rhazizine, R-77
 Rheadine, R-86
 Rheagenine, R-86
 Rhetine, R-168
 Rhetsine, E-310
 Rhetsinine, T-226
 Rhexifoline, R-78
 Rhinolidine, R-79
 Rhinoline, R-79
 Rhinolinine, R-79
 Rhizipon A, I-81
 Rhizochalin C, D-292
 Rhizochalinin C, D-292
 Rhizoferrin, R-80
 Rhizophorine, I-131
 Rhizopin, I-81
 Rhizopodin, R-81
 Rhizopterine, R-82
 Rhizoxin D, R-83
 Rhizoxin D₁, R-83
 Rhizoxin D₃, R-84
 Rhizoxin M₁, R-84
 Rhizoxin M₂, R-84
 Rhizoxin S₁, R-84
 Rhizoxin S₂, R-84
 Rhizoxin Z₁, R-84
 Rhizoxin Z₂, R-84
 Rhizoxin, R-83
 Rhodesiacridone, R-85
 Rhodiocyanogenin, H-575
 Rhodiocyanoside A, H-575
 Rhodiocyanoside B, H-509
 Rhodiocyanoside D, H-574
 Rhodomelanochrome, P-771
 Rhodommatin, D-527
 Rhoeadine, R-86
 Rhoeagenine, R-86
 Rhoifoline A, A-1566
 Rhoifoline B, R-87
 Rhoifolinic acid, R-88
 Rhombifoline, C-938
 Rhombinine, A-970
 Rhopaladin A, R-89
 Rhopaladin B, R-89
 Rhopaladin C, R-89
 Rhopaladin D, R-89
 Rhopalatine, I-208
 Rhosoptein, P-771
 Rhynchociline, R-134
 Rhynchophine, V-126
 Rhynchophyllic acid, R-90
 Rhynchophylline *N*-oxide, R-90
 Rhynchophylline, R-90
 Rhyncociline, R-134
 Rhyncophyllic acid, R-90
 Rhyncophylline *N*-oxide, R-90
 Rhyncophylline, R-90
 Ribaline, R-91
 Ribalinidine, R-92
 Ribalinine, R-92
 Ribalinium, R-93
 Ribaliprenylene, R-92
 Ribasidine, R-94
 Ribasine, R-94
 Riboflavin 5'-(trihydrogen diphosphate) 5' → 5'-diester with adenosine, F-79
 1- β -D-Ribofuranosylbrunfelsamidine, P-943
N ^{β} -D-Ribofuranosyldamirone C, D-13
 9- β -D-Ribofuranosylhypoxanthine, I-148
 1- β -D-Ribofuranosyl-1*H*-imidazol-5-amine, A-827
N ^{β} - α -Ribofuranosyllumichrome, L-285
N ^{β} -D-Ribofuranosylmakaluvamine I, M-54
N ^{β} -D-Ribofuranosylmakaluvic acid C, M-56
 9- β -D-Ribofuranosyl-9*H*-purine, N-82
 1- β -D-Ribofuranosyl-4-pyridone-3-carboxamide, H-709
 7-Ribofuranosyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine, T-658
 9- β -D-Ribofuranosylxanthine, X-13
 9-(β -D-Ribofuranosyl)zeatin, Z-11
 9- β -D-Ribopyranosylhypoxanthine, P-820
 Ribosylzeatin phosphate, Z-11
 Ribosylzeatin, Z-11
 Ricinidine, R-95
 Ricinine, R-96
 Ricinic acid, D-668
 Riddelline, S-240
 Riddelline, S-240
 Rideopine, G-107
 Ridleyamide, B-265
 Riedelianine, R-97
 Rigidin A, R-98
 Rigidin B, R-98
 Rigidin C, R-98
 Rigidin D, R-98
 Rigidin E, R-98
 Rigidin, R-98

- Rigidusol, C-525
 Rinderine, E-12
 Rindline, H-323
 Riparin, M-280
 Ripromycin, R-99
 Risatarun, D-730
 Rischiaril, D-730
 Ritigalin, N-235
 Ritmos, A-216
 Ritmosel, A-216
 Ritterazine A, R-101
 Ritterazine B, R-102
 Ritterazine C, R-103
 Ritterazine D, R-101
 Ritterazine E, R-104
 Ritterazine F, R-102
 Ritterazine G, R-102
 Ritterazine H, R-102
 Ritterazine I, R-102
 Ritterazine J, R-105
 Ritterazine K, R-105
 Ritterazine L, R-105
 Ritterazine M, R-105
 Ritterazine N, R-106
 Ritterazine O, R-106
 Ritterazine P, R-107
 Ritterazine Q, R-107
 Ritterazine R, R-108
 Ritterazine S, R-108
 Ritterazine T, R-101
 Ritterazine U, R-109
 Ritterazine V, R-100
 Ritterazine W, R-110
 Ritterazine X, R-110
 Ritterazine Y, R-102
 Ritterazine Z, R-111
 Rivanicline galactarate, M-232
 Rivanicline, M-232
 Rivinianin, B-110
 Rivularin A, H-211
 Rivularin B, T-117
 Rivularin C, T-108
 Rivularin D₁, T-116
 Rivularin D₂, T-116
 Rivularin D₃, T-116
 Rivularine *N*-oxide, T-188
 Rivularine, T-188
 RJR 2403, M-232
 RK 4415, A-124
 RK 1441A, A-1232
 RK 1441B, A-1232
 RK 1409B, S-498
 RK 286C, S-498
 RK 286D, A-1190
 RK 1061G, L-188
 RK 1061H, L-188
 RK 1409, S-498
 RK-F 1010, T-96
 RKS 1778, A-1196
 Ro 31-6045, A-1394
 Ro 5-9000, A-1081
 Ro 5-4645/10, C-711
 Ro 2-9757, F-104
 Ro 09-0049, P-337
 Ro 21-4958, R-151
 Robecine, R-112
 Robeneoside A, S-447
 Robeneoside B, S-447
 Robustamine *cis-N*-oxide, R-113
 Robustamine, R-113
 Robusticine, C-588
 Robustidine, S-126
 Robustiline, F-143
 Robustimine, S-126
 Robustine, F-213
 Robustine, S-456
 Robustinine, Q-47
 Robustivine, C-588
 Rodiasine, R-114
 Roebamine, R-116
 Roecarboline, R-115
 Roefractine, H-275
 Roeharmine, D-620
 Roehybramine β -*N*-oxide, R-116
 Roehybramine, R-116
 Roehybridine α -*N*-oxide, R-116
 Roehybridine β -*N*-oxide, R-116
 Roehybridine, R-116
 Roehybrine, R-117
 Roehymine, R-116
 Roelactamine, R-35
 Roemebramine, R-116
 Roemecarine *2\alpha-N*-oxide, R-118
 Roemecarine, R-118
 α -Roemehybrine, R-117
 Roemeramine, R-119
 Roemerialinone, O-112
 Roemeridine, R-116
 Roemerine *N*-oxide, R-120
 Roemerine, R-120
 Roemerolidine, O-85
 Roemeroline, A-1066
 Roemerone, R-119
 (+)-Roemrefidine, R-120
 (-)-Roemrefidine, R-120
 Roemrefine, R-36
 Rogersine, L-75
 Rohitukine *N*-oxide, R-121
 Rohitukine, R-121
 Rollipyrrole, R-122
 Romneine, R-123
 Romucosine A, D-531
 Romucosine B, D-531
 Romucosine C, T-524
 Romucosine D, T-524
 Romucosine F, T-246
 Romucosine G, P-208
 Romucosine H, I-208
 Romucosine I, R-124
 Romucosine, R-120
 Roquefortine A, F-181
 Roquefortine B, F-181
 Roquefortine C, R-125
 Roquefortine D, R-125
 Roquefortine E, R-126
 Roquessine, C-601
 Roraimine, W-9
 Rorifamide, M-563
 Rorifone, M-563
 Roripamine, M-564
 Rosamine, V-178
 Roseadine, R-127
 Roseamine, V-110
 Rosellichalasin, R-128
 Roseonine, S-565
 Roseophilin, R-129
 Roserine, R-130
 Rosibiline, R-131
 Rosicine, R-132
 Rosmarinecine, D-597
 Rosmarinine, R-133
 Rostratamine, P-223
 Rostratine, P-610
 Rotundifoleine, R-134
 Rotundifoline *N*-oxide, R-134
 Rotundifoline, R-134
 Rotundine A, R-135
 Rotundine B, R-135
 Rotundine C, R-135
 Rotundine, A-1410
 Rotundine, T-212
 Rouhamine, D-138
 Routiennocin, R-136
 Rovidine, R-137
 Roxburghiline, O-67
 Roxburghine A, R-138
 Roxburghine B, R-138
 Roxburghine C, R-138
 Roxburghine D, R-138
 Roxburghine E, R-138
 Roxburghine X, R-138
 Roxburghine, R-138
 Roxburghines, R-138
 Roxiamine A, R-139
 Roxiamine B, R-139
 Roxiamine C, R-140
 Royleinine, M-347
 Royline, M-309
 Rozevin, V-110
 RP 13252, A-1234
 RP 6798, A-1233
 RP 18631, C-395
 RP 55185, C-545
 RP 59451, C-545
 RP 49532, G-79
 RP 7452, L-153
 RP 9671, N-319
 RP 5278, S-569
 Rs I, A-915
 Rubemamide, D-718
 Rubemamine, D-718
 Rubenine, R-141
 Rubesamide, R-142
 Rubescenamamide, R-143
 Rubescenamier, R-143
 Rubescine, V-126
 Rubijervine, R-144
 Rubiverine, R-145
 Rubivirine, I-310
 Rubradinine, M-661
 Rubradirin B, R-146
 Rubradirin C, R-146
 Rubrifacine, R-147
 Rubrobramide, R-148
 Rubrocurarines, R-149
 Rubroflavin, D-229
 Rubrolone B, R-151
 Rubrolone, R-150
 Rubropunctamine, R-152
 Rubroside A, R-154
 Rubroside B, R-153
 Rubroside C, R-154
 Rubroside D, R-153
 Rubroside E, R-154
 Rubroside F, R-154
 Rubroside G, R-153
 Rubroside H, R-154
 Rudrakine, G-162
 Rufai B, D-669
 Rufescine, N-304
 Rufocromomycin, S-569
 RUG 13A, R-157
 RUG 13B, R-157
 Rugosanine A, N-337
 Rugosanine B, R-155
 Rugosin H, P-961
 Rugosinone, R-156
 Rugulovasine A, R-157
 Rugulovasine B, R-157
 Ruine, D-621
 Rulodine, R-158
 Rumberine, R-159
 12*E*-Rumbrin, R-160
 Rumbrin, R-160
 Runanine, R-161
 Rupanamine, R-162
 Rupicoline, V-178
 Ruprechstyril, R-163
 Rurrebanidine, R-164
 Rurrebanine, R-164
 Ruscopine, A-25
 Ruspolinone, R-165
 Russupteridine yellow I, R-166
 Russupteridine yellow IV, R-166
 Rutacridone epoxide, R-167
 Rutacridone, R-167
 Rutaecarpine, R-168
 Rutagravine, R-169
 Rutalexin, R-170
 Rotalinidine, R-91
 Rotalinium, R-171
 Rutamine, G-171
 Rutecarpine, R-168
 Ruticin, R-172
 Ruwenine, B-194
 Ruzorine, R-173
 Ryanex, R-175
Ryania diterpene ester A, R-175
Ryania diterpene ester B, R-175

- Ryania diterpene ester C₁, R-175
 Ryania diterpene ester C₂, R-174
 Ryania diterpene ester D, R-175
 Ryanicidic, R-175
 Ryanodine, R-175
 Ryanodyl 3-nicotinate, R-175
 Ryllistine, N-288
 Ryno-tox, R-175
 Ryosenamine, H-200
 Ryosenaminol, H-200
 Rytmalin, A-216
 Rzedowskin D, P-215
 S 433, A-1235
 S 53210A, A-1236
 S 54832A, A-1237
 S 53210A-I, A-1236
 S 54832A-I, A-1237
 S 53210A-II, A-1236
 S 54832A-II, A-1237
 S 53210A-III, A-1236
 S 54832A-III, A-1237
 S 54832A-IV, A-1237
 S 632A₂, M-561
 S 632A₃, M-561
 S 632A₁, S-563
 S 632B₁, M-561
 S 632B₂, M-561
 S 1319, H-508
 S 551II, R-34
 S 1210, R-52
 SA 3097B₁, A-1044
 SA 3097B₂, A-1044
 SA 3097C₁, A-1044
 SA 3097C₂, A-1044
 SA 3097D₁, A-1044
 SA 3097D₂, A-1044
 SA4-3, A-5
 SAB 711, A-1238
 Sabadillines, S-1
 Sabadine, C-291
 Sabadinine, V-56
 Sabatine, C-291
 Sabine, C-291
 Sachaconitine, M-355
 Sadosine, H-177
 Safracin A, S-2
 Safracin B, S-2
 Saframycin AH₁, S-4
 Saframycin AH₂, S-4
 Saframycin AR₁, S-4
 Saframycin AR₃, S-4
 Saframycin A_{D-1}, S-3
 Saframycin A, S-4
 Saframycin B, S-4
 Saframycin C, S-4
 Saframycin D, S-4
 Saframycin E, S-4
 Saframycin F, S-4
 Saframycin G, S-4
 Saframycin H, S-5
 Saframycin Mx 1BC, S-6
 Saframycin Mx 2BC, S-6
 Saframycin Mx 1, S-6
 Saframycin Mx 2, S-6
 Saframycin R, S-7
 Saframycin S, S-8
 Saframycin Y_{2b-D}, S-10
 Saframycin Y_{2b}, S-9
 Saframycin Y_{D-1}, S-12
 Saframycin Y_{D-2}, S-13
 Saframycin Y₃, S-11
 Sagitol, S-14
 Sagittamide A, S-15
 Sagittamide B, S-15
 Sagittariol, C-525
 Saifine, S-16
 Saipeimine, S-17
 Saitomyin, C-960
 Sakambullin, D-12
 Sakerine, G-68
 Sakerinol A, G-68
 Sakerinol B, G-68
 Sakerol, G-68
 Sakerone, G-68
 Salacin, S-18
 Salarin A, S-19
 Salarin B, S-20
 Salfredin A₃, S-21
 Salfredin A₄, S-21
 Salfredin A₇, S-21
 Salfredin C₁, S-21
 Salfredin C₂, S-21
 Salfredin C₃, S-21
 Salicifoline, H-565
 Salicilobine, S-22
 Salicornine, S-23
N-Salicylanthranilic acid, B-67
 Salicylamine, H-415
N-Salicyloyl-2-amino-1,3-propanediol, A-893
N-Salicyloyl-2-amino-1-propanol, A-894
 Salidrosol, H-676
 Saligcinnamide, P-601
 Salignamine, A-876
 Salignarine A, D-262
 Salignarine B, D-276
 Salignarine C, D-276
 Salignarine D, P-608
 Salignarine E, P-605
 Salignarine F, D-277
 Salignenamamide A, P-601
 Salignenamamide B, D-258
 Salignenamamide C, D-272
 Salignenamamide D, D-259
 Salignenamamide E, P-608
 Salignenamamide F, P-601
E-Salignone, A-890
Z-Salignone, A-890
 Saliherbine, S-24
 Salimine, S-25
 Saliniketol A, S-26
 Saliniketol B, S-26
 Salinosporamide A, S-28
 Salinosporamide B, S-28
 Salinosporamide C, S-29
 Salinosporamide D, S-30
 Salinosporamide E, S-31
 Salinosporamide F, S-28
 Salinosporamide G, S-30
 Salinosporamide H, S-28
 Salinosporamide I, S-27
 Salinosporamide J, S-28
 Salmochelin 4, S-32
 Salmochelin X, S-33
 Salmycin A, S-34
 Salmycin B, S-34
 Salmycin C, S-35
 Salmycin D, S-35
 Salochimin, Q-20
 Salonine A, D-272
 Salonine B, A-876
 Salonine C, P-598
 Saloquinine, Q-20
 Salsamine, S-36
 Salsolidine, T-167
 Salsoline A, T-629
 Salsoline, T-179
 Salsolinol, T-162
 Saludimerine A, S-37
 Saludimerine B, S-37
 Salutadimerine, S-38
 Salutaridine *N*-oxide, S-39
 Salutaridine, S-39
 Salutarine, S-39
 Salvadoricine, A-61
 Salviadiolone, S-40
 Salviamiltamide, S-41
 Salvianane, S-42
 Salvianene, S-42
 Samandaridine, S-43
 Samandarine, S-44
 Samandarone, S-44
 Samandenone, S-45
 Samandinine, S-46
 Samanine, S-47
 Sambunigrin, H-673
 Sambutoxin, S-48
 Samoquasine A, B-75
 Sampangine, S-49
 Sanasin, P-833
 Sanazin, P-833
 Sandramycin, S-50
 Sandril, R-52
 Sandwicine, A-216
 Sandwicolidine, S-51
 Sandwicoline, S-52
 Sangivamyic acid, S-53
 Sangivamycin, S-53
 Sanguidimerine, S-54
 Sanguilutine, S-55
 Sanguinarine pseudoethanolate, D-510
 Sanguinarine pseudomethanolate, D-510
 Sanguinarine, S-56
 Sanguinarium chloride, S-56
 Sanguinine *N*-oxide, G-3
 Sanguinine, G-3
 Sanguinolentaquinone, S-57
 Sanguinone A, S-58
 Sanguinone B, S-59
 Sanguirubine, S-60
 Sanjoinine, D-840
 Sanjoinine A, F-150
 Sanjoinine Ah₁, F-150
 Sanjoinine B, F-150
 Sanjoinine D, S-61
 Sanjoinine E, D-531
 Sanjoinine F, F-150
 Sanjoinine G₁, S-61
 Sanjoinine G₂, S-62
 Sanjoinine Ia, D-531
 Sanjoinine Ib, I-208
 Sanjoinine K, H-275
 Sankhpuspine, S-63
 α-Sanshool, D-893
 β-Sanshool, D-893
 γ-Sanshool, T-127
 Santavy's Substance F, D-92
 Santheose, T-355
 Santiagonamine, S-64
 Santiaguine, S-65
 Sanyonamine, H-163
 Sapatoxin B, P-226
 Saphenamycin, H-496
 Saphenic acid methyl ether, H-496
 Saphenic acid, H-496
 4α-Sapinine, T-275
 Sapinopyridone, F-162
 Sapintoxin A, T-275
 Sapintoxin B, P-226
 Sapintoxin C, T-274
 α-Sapintoxin, T-275
 Sapropterin dihydrochloride, T-142
 Sapropterin, T-142
 Sarachine, S-66
 Saracocinaene, P-599
 Saracocine, P-606
 Saracodine, P-601
 Saracodinine, P-606
 Saracorine, P-601
 Sarain A, S-67
 Sarain B, S-67
 Sarain C, S-68
 Saraine 1, S-69
 Saraine 2, S-70
 Saraine 3, S-71
 Sarcinapterin, S-72
 Sarcine, P-820
cis-Sarcocapnidine *N*-oxide, S-73
 Sarcocapnidine, S-73
cis-Sarcocapnine *N*-oxide, S-73
 Sarcocapnine, S-73
 Sarcocinaene, P-599
 Sarcodifurine A, S-74
 Sarcodifurine B, S-75
 Sarcodine, P-601
 Sarcodonin α, S-76
 Sarcodonin β, S-76
 Sarcodonin δ, S-76
 Sarcodonin γ, S-76
 Sarcodonin, S-76

- Sarcomegistine, P-254
 Sarcomejine B, S-78
 Sarcomejine, S-77
 Sarconidine, P-599
 Sarcopharyngine, S-79
 Sarcophylline, S-80
 Sarcorine, P-601
 Sarcorucinine A1, P-600
 Sarcorucinine A, P-600
 Sarcorucinine B, A-889
 Sarcorucinine D, A-881
 Sarcosaligmine, A-884
 Sarcostin, P-610
 Sarcotragin A, S-81
 Sarcotragin B, S-81
 Sarcotrine A, S-82
 Sarcotrine B, S-82
 Sarcotrine C, S-82
 Sarcotrine D, S-83
 Sarcotrine E, S-82
 Sarcotrine F, S-84
 Sarcovagenine A, D-273
 Sarcovagenine B, D-273
 Sarcovagenine C, D-260
 Sarcovagenine A, D-262
 Sarcovagine B, D-262
 Sarcovagine C, D-266
 Sarcovagine D, D-274
 Sarcoviolin α , S-76
 Sargassumactam, S-85
 Sarkine, P-820
 Sarmentamide A, S-86
 Sarmentamide B, S-87
 Sarmentamide C, T-577
 Sarmentine, D-104
 Sarmentosin epoxide, H-509
 Sarmentosine, M-452
 Sarmentosin, H-509
 Sarodesmine, H-554
 Sarothamnine, S-88
 Sarpagan-3,7-diol, A-27
 Sarpagan-10,17-diol, S-89
 Sarpagan-17-ol, T-425
 Sarpagine, S-89
 Sarracine *N*-oxide, S-90
 Sarracine, S-90
 Sarracinic acid, H-574
 Sarranicine, S-90
 Sarsalignenone, D-261
 Sarsalignone, D-282
 Sarusubine A, S-91
 Sativanine A, S-92
 Sativanine B, S-93
 Sativanine C, S-94
 Sativanine D, S-95
 Sativanine E, S-96
 Sativanine F, S-97
 Sativanine G, S-98
 Sativanine H, S-99
 Sativanine K, S-98
 Sativanine M, S-94
 Sativanine N, S-98
 Sativanine O, S-100
 Sattazolin, S-101
 Saulatine, S-102
 Sauristolactam, D-567
 Sauroine, F-29
 Sauroxine, O-5
 Saururidine, S-103
 Saururine, S-104
 Saussureamine A, S-105
 Saussureamine B, I-162
 Saussureamine C, S-106
 Saussureamine D, S-107
 Saussureamine E, S-107
 Sauvagnine, S-108
 Saxicolaline A, T-248
 11-Saxitoxinacetic acid, S-110
 11-Saxitoxinethanoic acid, S-110
 Saxitoxin, S-109
 Saxoguattine, S-111
 Saxorumamide, S-112
 SB 202327, A-1486
 SB 203207, A-660
 SB 203208, A-660
 SB 212305, A-1239
 SB 217452, A-1240
 SB 219383, A-1241
 SB 253514, A-1242
 SB 253517, A-1242
 SB 253518, A-1242
 SB 291071, A-1242
 SB 291072, A-1242
 SB 291073, A-1242
 SB 315021, A-1243
 SB 212021, H-665
 SB 204900, M-509
 SB 408075, M-135
 SB 200437, P-90
 SB 203105, P-90
 Scabrine, S-113
 Scabrosin 4-acetate-4'-butyrate, S-114
 Scabrosin 4-acetate-4'-hexanoate, S-114
 Scabrosin 4-butanoate-4'-hexanoate, S-114
 Scabrosin 4,4'-diacetate, S-114
 Scabrosin 4,4'-dibutyrate, S-114
 Scabrosin, S-114
 Scaconine, M-352
 Scaconitine, M-352
 Scaevodimerine A, S-115
 Scaevodimerine B, S-115
 Scaevodimerine C, S-115
 Scaevodimerine D, S-115
 Scaevoline, S-116
 Scalusamide A, S-117
 Scalusamide B, S-117
 Scalusamide C, S-117
 Scandine *N*^b-oxide, S-118
 Scandine, S-118
 Scandomelidine, S-119
 Scandomeline, S-120
 Scandomelonine, S-121
 Scatole, M-474
 Sceleratine *N*-oxide, S-122
 Sceleratine, S-122
 Sceleratinyll chloride, S-122
 Sceletenone, S-123
 Sceptrine, S-124
 Sch 210971, A-1250
 Sch 210972, A-1250
 Sch 213766, A-1250
 Sch 40832, A-1244
 Sch 40873, A-1245
 Sch 42137, A-1246
 Sch 52900, A-1247
 Sch 52901, A-1247
 Sch 538415, A-1251
 Sch 54445, A-1248
 Sch 56036, A-1249
 Sch 725424, A-1252
 Sch 18640A, A-1082
 Sch 10595, B-445
 Sch 18640B, A-1082
 Sch 575948, C-721
 Sch 38513, F-115
 Sch 38516, F-115
 Sch 38518, F-115
 Sch 39185, F-115
 Sch 42282, F-115
 Sch 42729, F-115
 Sch 18640, T-392
 SC48334, H-616
 Schefferine, C-677
 Schelhammeridine, S-125
 Schelhammerine, S-126
 Schellhammericine B, T-51
 Schellhammericine, S-126
 Schinifoline, G-118
 Schinifoline, H-133
 Schischkiniin, S-127
 Schizanthine A, S-129
 Schizanthine B, S-128
 Schizanthine C, S-128
 Schizanthine D, S-130
 Schizanthine E, S-130
 Schizanthine F, S-131
 Schizanthine G, S-131
 Schizanthine H, S-131
 Schizanthine I, S-131
 Schizanthine K, S-131
 Schizanthine L, S-131
 Schizanthine M, S-131
 Schizanthine X, S-128
 Schizanthine Y, S-128
 Schizanthine Z, S-128
 Schizocommunine, S-132
 Schizogaline, S-133
 Schizogamine, S-133
 Schizokinen A, S-134
 Schizoluteine, S-135
 Schizophylline, S-136
 Schizozygine, S-137
 α -Schizozygol, S-137
 β -Schizozygol, S-138
 Schlechtendamine, S-200
 Schobericine, T-219
 Schoberidine, S-139
 Schoberine, S-140
 Scholaricine, E-30
 Scholarine *N*^b-oxide, E-30
 Scholarine, E-30
 Scholarisine A, S-141
 Schopf's base VI, L-91
 Schulzeine A, S-142
 Schulzeine B, S-143
 Schulzeine C, S-143
 Schumagnine, S-144
 Schumannificine, S-145
 Schumannioficine, S-145
 Schumanniophytine, S-146
 Schweinine, S-147
 Sciadenine, S-148
 Sciadoferine, S-149
 Sciadoline, S-149
 Sciaferine, M-712
 Scinnamina, R-50
 Sciodole, S-150
 Scleramin, E-6
 Scleritodermin A, S-151
 Sclerominol, S-152
 Sclerotionine, E-173
 Sclerotiamide, S-153
 Sclerotigenin, S-154
 Sclerotiorinamine, S-155
 Scobutil, S-159
 Scolopendrine, S-156
 Scombrotoxin, H-307
 α -Scopadonnine, S-157
 β -Scopadonnine, S-157
 Scopadonnine, S-157
 Scopaline, A-99
 Scopamin, S-159
 Scoparasin A, C-949
 Scoparasin B, C-949
 Scopine, S-158
 ψ -Scopine, S-158
 Scopinone, S-158
 Scopoderm, S-159
 Scopolamine butyl bromide, S-159
 Scopolamine hydrobromide, S-159
 Scopolamine, S-159
 Scopolan, S-159
 Scopoline tropate, S-159
 Scopoline, S-160
 Scopos, S-159
 Scorazanone, S-161
 Scorodocarpine A, H-755
 Scorodocarpine B, H-755
 Scorodocarpine C, H-755
 Scorodophlone A, S-162
 Scorpionone, B-255
 Scopioidine, S-163
 Scotine, C-708
 Scoulerine, S-164
 Scutebarbatine A, T-535
 Scutebarbatine B, T-535
 Scutianene C, D-841
 Scutianene D, D-841
 Scutianine A, S-165

- Scutianine B, S-167
 Scutianine C, S-166
 Scutianine D, S-167
 Scutianine E, S-167
 Scutianine F, S-165
 Scutianine G, S-167
 Scutianine H, S-166
 Scutianine J, S-167
 Scutianine K, S-167
 Scutianine L, C-600
 Scutianine M, C-600
 Scutianine, S-165
 Scutifoliamide A, P-462
 Scutifoliamide B, P-462
 Scytonemin, S-168
 Scytonine, S-169
 Sczukidine, S-170
 Sczukinine, S-170
 Sczukitine, S-170
 Sebastianine A, S-171
 Sebastianine B, S-172
 Sebiferine *N*-oxide, F-80
 Sebiferine, F-80
 Secaloside A, S-173
 Secaloside B, S-173
 Secaloside C, S-173
 Secaloside D, S-173
 Secamine, S-174
 Secantioquine, S-175
 6,7-Secoagroclavine, S-176
 4,6-Secoangustilobinal, S-177
 6,7-Secoangustilobine A, S-177
 6,7-Secoangustilobine B, L-264
 Secoantofine β -*N*-oxide, S-248
 Secoantofine, S-248
 Secobatzelline A, S-178
 Secobatzelline B, S-178
 Secocepharanthine, S-179
 1,8-Secocinchonine, Q-17
 Secoclausenamide, D-651
 Secocularidine, S-180
 Secocularine, S-180
 6,7-Seco-6-cyanostemmadenine, Y-21
 Secodaphniphylline, S-181
 3,4-Seco-3,14-dehydrocabucraline, S-182
 11,12-Seco-12-dehydromultiflorine, A-554
 Secodemethylclausenamide, D-651
 12-Seco-14-deoxyparsonianine, S-183
 Secodihydrocastoramine, N-340
 Secodine, S-184
 1,2-Secodysidamide H, S-185
 1,2-Secodysidamide, S-185
 6,7-Seco-19,20-epoxyangustilobine, L-264
 Secofascaplysin A, S-186
 Secoglaucine, M-559
 Secohomoaromaline, S-192
 Secoisodorinol, C-456
 Secoisopiriferinol, C-456
 Secoisotetrandrine, C-806
 Secojesaconitine, S-187
 Secojollyanine, P-800
 Secojuliprosopinal, S-188
 Secokaraconitine, S-189
 Secologanin, D-204
 Secologanol, D-204
 Secolucidine, S-190
 Secomartinelline, S-191
 6,7-Seco-6-norangustilobine B, L-264
 Secoobaberine, S-192
 Secoodorine, C-456
 14,15-Seco-3-oxokopsinal, S-193
 Secopenitrem B, S-194
 Secophoebine, T-328
 Secopiriferine, C-456
 Secoplicamine, S-195
 Secopseudopteroxazole, S-196
 Secopyrrolotenin, C-456
 Secoquettamine, S-197
 Secosarcocapnidine, S-198
 Secosarcocapnine, S-198
 16,28-Secosolanida-5,22(28)-diene-3,16-diol, S-199
 16,28-Secosolanida-5,22(28)-diene-3,12-diol, V-68
 16,28-Secosolanida-5,22(28)-diene-3,12,16-triol, H-11
 16,28-Secosolanida-5,22(28)-dien-3-ol, V-73
 16,28-Secosolanida-4,22(28)-dien-3-one, V-73
 16,28-Secosolanid-22(28)-ene-3,16-diol, S-338
 16,28-Secosolanid-22(28)-ene-3,6-diol, S-201
 16,28-Secosolanid-5-ene-3,16-diol, S-200
 16,28-Secosolanid-22(28)-en-3-ol, S-337
 16,28-Secosolanid-5-en-3-ol, V-73
 Secoxanthoplanine, M-559
 Secoyunaconitine, S-187
 Secuamamine A, S-202
 Secuamamine B, P-386
 Secuamamine C, S-206
 Secuamamine D, P-387
 Securamine A, S-203
 Securamine B, S-203
 Securamine C, S-204
 Securamine D, S-204
 Securamine E, S-204
 Securamine F, S-204
 Securamine G, S-204
 Securinegine, S-205
 Securinine, S-206
 Securinol A, S-207
 Securinol B, S-207
 Securinol C, S-208
 Securitinine, P-386
 Sedacrine, S-214
 Sedacryptine, S-209
 Sedamine, S-210
 (\pm)-Sedamine, S-210
 Sedaminone, S-210
 Sedantole, C-555
 Sederine, S-211
 Sediendione, S-212
 Sediene, S-213
 Sedinine, S-214
 Sedinone, S-215
 Sedridine, S-216
 (\pm)-Sedridine, S-216
 Segoline A, S-217
 Segoline B, S-217
 Segoline C, S-217
 Sekisanine, T-53
 Sekisanoline, T-53
 Selacholeic acid, T-120
 Selagine, H-375
 Selaginoidine, S-218
 Selagoline, L-341
 Selenobiotin, S-219
 Sellowiine, S-220
 Semecarpifoline, S-221
 Semecarpine, S-222
 Semiaquilegine A, L-70
 Semicochliodinol A, A-1521
 Semicochliodinol B, A-1521
 Semilepidine A, H-422
 Semilepidine B, H-422
 Semilepidinoside A, H-422
 Semilepidinoside B, H-422
 Semiplenamamide A, M-435
 Semiplenamamide B, M-435
 Semiplenamamide C, S-223
 Semiplenamamide D, S-224
 Semiplenamamide E, S-225
 Semiplenamamide F, S-225
 Semiplenamamide G, S-225
 Semperflorine, T-288
 Sempervine, S-228
 Semperviramidine, C-883
 Semperviraminol, S-226
 Semperviraminone, S-227
 Sempervirine, B-466
 Sempervirine, S-228
 Sempervirone, S-229
 Semperviroxazolidine, S-230
 SEN 34, H-562
 SEN 215, T-424
 Senacarcin A, S-231
 Senacarcin H, S-231
 Senaetnine, S-232
 14,15-*trans*-Senaetnine, S-232
 Senampelines, S-233
 Senbusine A, M-323
 Senbusine B, M-327
 Senbusine C, M-313
 Sencalenine, T-188
 Sendaverine *N*-oxide, S-234
 Sendaverine, S-234
 Senecicaudatin *O*-isopentanoate, S-235
 Senecicaudatin *O*-senecioate, S-235
 Senecicaudatinal hemiacetal, S-236
 Senecicaudatin, S-235
 Seneciocannabinine, S-237
 Senecionine *N*-oxide, S-238
 Senecionine, S-238
 Senecioracene, S-239
 Seneciolycholine, C-419
*O*²-Seneciolyllmacronecine, H-518
*O*⁹-Seneciolyllmacronecine, H-518
 3-Seneciolyloxytropane, M-393
 6-Seneciolyloxytrop-3-ol, M-391
 3-Seneciolyloxytrop-6-ol, M-391
O-Seneciolyllatynecine, H-233
N-Seneciolyllpyrrothine, A-759
*O*⁷-Seneciolyllretronecine *N*-oxide, T-188
*O*⁷-Seneciolyllretronecine, T-188
*O*⁹-Seneciolyllretronecine, T-188
 Seneciphylline epoxide, S-240
 (*E*)-Seneciphylline epoxide, S-240
 Seneciphylline *N*-oxide, S-240
 Seneciphylline, S-240
 (*E*)-Seneciphylline, S-240
 Seneciphyllinine, S-240
 Senecivernine *N*-oxide, S-241
 Senecivernine, S-241
 Senepodine A, S-242
 Senepodine B, S-242
 Senepodine C, S-242
 Senepodine D, S-242
 Senepodine E, S-242
 Senepodine F, S-242
 Senepodine G, S-243
 Senepodine H, S-244
 Senkirkine, S-245
 Sepaconitine, A-87
 Sepeerine, O-1
 Sepiapterin C, A-40
 Sepiapterin, S-246
 Septatisine, S-247
 Septedine, S-247
 Septedine, S-247
 Septefine, A-91
 Septenine, H-180
 Septentriodine, P-776
 Septentriodine, P-776
 Septentriodine, P-776
 Septentriodine, H-180
 Septephine, A-91
 Septerine, M-309
 Septicine, S-248
 Septorine, S-249
 Sequoiamonascin D, S-250
 Seradix, I-83
 (+)-Serantrypinone, A-240
 (-)-Serantrypinone, A-240
 Serecor, Q-18
 Seredamine, T-288
 Seredine, S-251
 Serine 2-[(aminoiminomethyl)amino]ethyl hydro-
 gen phosphate (ester), L-240
 Serinol, A-893
 Sermix, R-52
 Seromycin, A-832
 Serotobenine, S-252
 Serotonin, H-755
 Serpasil, R-52
 Serpenticine, A-652
 Serpentine, A-652
 Serpentinine, S-253
 Serpinine, T-288
 Serratanidine, S-255
 Serratanine A, L-277
 Serratanine B, L-277
 Serratenone, H-310
 Serratezomine A, S-254

- Serratezomine B, S-257
 Serratezomine C, L-325
 Serratidine, A-1029
 Serratine, S-255
 Serratinidine, S-256
 Serratinine *N*-oxide, S-257
 Serratinine, S-257
 Serratiochelin, S-258
 Serratoline, S-259
 Sesbanimide A, S-260
 Sesbanimide B, S-260
 Sesbanimide C, S-261
 Sesbanimide, S-260
 Sesbanine, S-262
 Suspendole, S-263
 Sesquiterpenylamylaminohydroxyquinone, S-329
 Sessiflorine, S-264
 Sessilifoliamide A, S-522
 Sessilifoliamide B, S-265
 Sessilifoliamide C, S-265
 Sessilifoliamide D, S-266
 Sessilifoliamide E, S-267
 Sessilifoliamide F, S-267
 Sessilifoliamide G, S-268
 Sessilifoliamide H, T-666
 Sessilifoliamide I, T-665
 Sessilifoline A, T-664
 Sessilifoline B, S-528
 Sessilifoline, D-604
 Sessiline, S-269
 Sessilistemonamine A, S-270
 Sessilistemonamine B, S-270
 Sessilistemonamine C, S-270
 Sessilistemonamine D, S-271
 Sessilistemonine, S-272
 Setigeridine, S-274
 Setigerine, P-79
 Setoclavine, S-273
 Sevamine, S-274
 Sevcordine, S-275
 Sevcordinine, S-275
 Sevcorine, S-275
 Sevedamine, C-295
 Sevedine *N*-oxide, C-295
 Sevedine, C-295
 Sevedinine, D-563
 Seveline, C-305
 Sevelinone, C-305
 Severcine, A-146
 Severibuxine, S-276
 Severidine, S-277
 Severidinine, S-277
 Severifoline, S-278
 Severine *N*-oxide, C-295
 Severine palmitate, A-77
 Severine, A-77
 Severine, C-295
 Severtzidine, C-295
 Severtzine, A-146
 Severtzine, C-305
 Severtzinine, S-279
 Sewarine, A-228
 Sewedamine, C-295
 Sewercinine, S-279
 Seweridine, S-277
 Seweridinine, S-277
 Sewertzidine, C-295
 Sewerzinine, S-279
 SF 1346, A-822
 SF 1508, A-1253
 SF 1739, A-1254
 SF 2140, A-1255
 SF 2312, A-1256
 SF 2364, A-1157
 SF 2369, A-1257
 SF 2370, A-1189
 SF 2425, A-1258
 SF 2538, A-902
 SF 2695, A-1259
 SF 2768, A-1260
 SF 2771, A-1261
 SF 2809, A-1262
 SF 337, A-735
 SF 2197A, A-1094
 SF 2695A, A-1259
 SF 973A, A-14
 SF 2738A, C-574
 SF 2080A, D-356
 SF 2582A, D-954
 SF 2583A, O-157
 SF 2741A, P-641
 SF 2197B, A-1263
 SF 2695B, A-1259
 SF 973B, A-14
 SF 2738B, C-574
 SF 2582B, D-953
 SF 2420B, M-491
 SF 2583B, O-157
 SF 2080B, P-963
 SF 2741B, P-642
 SF 2738, C-574
 SF 2738C, C-575
 SF 2582C, D-954
 SF 2080C, P-964
 SF 1917, D-73
 SF 2441, D-765
 SF 2738D, C-576
 SF 2080D, P-964
 SF 2738E, C-577
 SF 2080E, P-965
 SF 2738F, C-578
 SF 2080F_{2a}, P-966
 SF 2080F_{2b}, P-966
 SF 2080F₁, P-966
 SF 2080F₃, P-966
 SF 1739HP, N-32
 SF 2636, I-216
 SF 2809-I, A-1262
 SF 2809-II, A-1262
 SF 2809-III, A-1262
 SF 2809-IV, A-1262
 SF 2494, T-658
 SF 2809-V, A-1262
 SF 2809-VI, A-1262
 S-GI, H-616
 Shaheenine, C-563
 Shaoguanmycin A, P-421
 Shaoguanmycin B, P-421
 Shawurensine, D-182
 Shearinine A, S-281
 Shearinine B, J-21
 Shearinine C, S-282
 Shearinine D, S-281
 Shearinine E, S-281
 Shearinine F, S-281
 Shearinine F, S-284
 Shearinine G, S-281
 Shearinine H, S-280
 Shearinine I, S-280
 Shearinine J, S-283
 Shearinine K, S-284
 Shepherdine, S-285
 Shermilamine A, S-286
 Shermilamine B, S-286
 Shermilamine C, S-286
 Shermilamine D, S-286
 Shermilamine E, S-286
 Shihuidine, S-287
 Shihunine, S-288
 Shimoburo base II, M-309
 Shimoburo base I, N-30
 Shinonomenine, C-313
 Shisen 1, H-496
 Shishididemniol A, S-289
 Shishididemniol B, S-289
 Shishididemniol C, S-289
 Shishididemniol D, S-289
 Shishididemniol E, S-289
 Shishijimicin A, S-290
 Shishijimicin B, S-290
 Shishijimicin C, S-290
 Shitakigenin, H-256
 SHP 2, S-637
 Shuazine, C-418
 Siamenol, S-291
 Siamine, T-566
 Siaminine A, T-544
 Siaminine B, I-306
 Siastatin B, A-746
 Sibanomicin, A-1157
 Sibiricine, C-673
 Sibiridine, I-308
 Sibirine, N-241
 Sibirinine, S-292
 Sibyllimycin, S-293
 Sickenbergine, S-294
 Sickingine, T-150
 Siculine, M-108
 Siculinine, U-35
 Siddiquamine, K-56
 Siddiquine, K-56
 Siderochelin A, S-295
 Siderochelin B, S-295
 Siderochelin C, S-296
 Sieboldine A, S-297
 Siechuansine, C-298
 Siechuantine, S-298
 Signatipennine, S-299
 Silicine, M-293
 Silvaticamide, S-300
 Simaomicin α , S-301
 Simaomicin β , S-301
 Simmondsin, S-302
 Simpalon, S-659
 Simplakidine A, S-303
 Simplexine, S-304
 Simulanoquinoline, S-305
 Simulansamide, S-306
 Simulansine, Z-5
 Simulenoline, Z-5
 Sinaconitine A, A-86
 Sinaconitine B, A-91
 Sinactine, S-307
 Sinalbin A, C-842
 Sinalbin B, C-842
 Sinalexin, I-336
 Sinanomycin, N-180
 Sinapine, S-308
 2'-*O*-*trans*-Sinapoylalangiside, A-239
 6'-Sinapoyldemethylalangiside, A-239
 2'-*O*-*trans*-Sinapoyldemethylalangiside, A-239
 2'-*O*-*trans*-Sinapoyl-3-*O*-demethyl-2-*O*-methylalangiside, A-239
 6'-*trans*-Sinapoylalloside, L-316
 Sinapoylputrescine, C-456
 Sinapoylspermidine, S-398
 Sinapoylspermine, S-399
N-*cis*-Sinapoyltyramine, H-667
N-*trans*-Sinapoyltyramine, H-667
 Sincalin, C-419
 Sincamidine, I-68
 Sindamine, C-806
 Singapurensine A, S-309
 Singapurensine B, S-309
 Singapurensine C, S-309
 Singapurensine D, S-309
 Sinharine, S-310
 Sinicuichine, V-88
 Sinine, V-95
 Sinkalin, C-419
 Sinoacutine, S-39
 Sinococuline, S-311
 Sinofranine, S-627
 Sinomendine, S-312
 Sinomenine *N*-oxide, S-313
 Sinomenine, S-313
 Sinomontanine A, A-91
 Sinomontanine B, A-91
 Sinomontanine C, S-315
 Sinomontanine D, A-93
 Sinomontanine E, A-86
 Sinomontanine F, A-86
 Sinomontanine G, D-187
 Sinomontanine H, A-85
 Sinomontanine I, S-314
 Sinomontanine I, S-315
 Sinomontanitine A, S-316
 Sinomontanitine B, S-316
 Sinostemonine, S-317

- Sinpeinine A, C-290
 Sintenpyridone, P-473
 Sinulamide, S-318
 Sinulamine III, S-320
 Sinulamine II, S-319
 Sinulamine I, S-319
 Siomycin A, S-321
 Siomycin B, S-321
 Siomycin C, S-321
 Siomycin D₁, S-322
 Sipeimine, C-298
 Sipeimone, C-298
 Siphonazole, S-323
 Siphonodictidine, S-324
 Sisunine, S-445
 Sitsirikine, S-325
 (16S)-Sitsirikine, S-325
 Siwanine A, M-362
 Siwanine B, M-362
 Siwanine C, M-371
 Siwanine D, M-362
 Siwanine E, M-362
 Siwanine F, M-362
 SK 7, T-355
 Skatole-2-carboxylic acid, M-476
 Skatole, M-474
 SKF 250, D-92
 Skimmianine, F-223
 Skototenin, I-80
 β-Skytanthine *N*-oxide, S-326
 α-Skytanthine, S-326
 β-Skytanthine, S-326
 δ-Skytanthine, S-326
 Skytanthine, S-326
 γ-Skytanthine, S-326
 SL 3238, F-197
 Slaframine, S-327
 Slagenine A, S-328
 Slagenine B, S-328
 Slagenine C, S-328
 SM 35, A-853
 SM 76, A-853
 SM 43, G-180
 Smenospongiarine, S-329
 Smenospongidine, S-329
 Smenospongine, S-329
 Smenospongiorine, S-329
 Smipine, S-330
 Smirnovine, S-406
 Smirnovinine, S-406
 SM-Q, F-190
 SMTP 7, A-1264
 SMTP 8, A-1265
 SMTP 7D, A-1264
 SMTP 8D, A-1265
 SMTP 4D, S-332
 SMTP 5D, S-333
 SMTP 6D, S-334
 SMTP O, S-490
 SMTP 1, S-490
 SMTP 2, S-490
 SMTP 3, S-331
 SMTP 4, S-332
 SMTP 5, S-333
 SMTP 6, S-334
 Smyrnovine, S-406
 Smyrnovinine, S-406
 SN 198B, D-203
 SN 198C, P-421
 SN 198D, D-203
 SN 198E, P-421
 Sodium tenuazonate, T-74
 Solacallinidine, S-335
 Solacapine, S-336
 Solacaproine, S-340
 Solacassine, S-346
 Solacauline, S-345
 Solacongostidine, S-337
 Solacongestine, S-337
 α-Solacongestinine, S-337
 β-Solacongestinine, S-337
 Solaculine A, S-456
 Soladulcamaridine, S-456
 Soladulcidine 3-chacotrioxide, S-445
 Soladulcidine 3-lycotetraoxide, S-445
 Soladulcidine, S-445
 Soladulcine A, S-445
 Soladulcine B, S-445
 α-Soladulcine, S-445
 β-Soladulcine, S-445
 γ-Soladulcine, S-445
 Soladulcidine tetraoxide, S-445
 Soladunalinidine, S-439
 Solafilidine, S-349
 Solafloridine, S-338
 Solakhasianine, S-456
 Solalyratine A, S-445
 Solalyratine B, S-445
 Solamaladine, S-339
 β₁-Solamargine, S-456
 β₂-Solamargine, S-456
 Solamargine, S-456
 α-Solamarine, S-456
 β-Solamarine, S-456
 δ-Solamarine, S-456
 γ-Solamarine, S-456
 γ₁-Solamarine, S-456
 γ₂-Solamarine, S-456
 Solamine, S-340
 Solanapyrone K, S-341
 Solanapyrone L, S-341
 Solanaviol, S-447
 Solancarpidine, S-456
 Solancarpine, S-456
 Solanelagnine, S-456
 Solangustidine, S-342
 Solangustine, S-342
 3,5-Solanadadiene, S-343
 3,5,6-Solanidanetriol, S-344
 Solanidan-3-ol, D-205
 Solanid-5-ene-3,18-diol, I-310
 Solanid-5-ene-3,23-diol, L-116
 22 α H,25 β H-Solanid-5-ene-3 β ,12 α -diol, R-144
 Solanid-5-en-3-ol, S-345
 Solanidine S, S-456
 Solanidine T, S-345
 Solanidine, S-345
 α-Solanigrine, S-456
 β-Solanigrine, S-456
 Solanine D, D-205
 α-Solanine, S-345
 β-Solanine, S-345
 γ-Solanine, S-345
 Solanocapsine, S-346
 Solanocardinol, D-99
 Solanocastrine, S-347
 Solanoforthine, S-346
 Solanogantamine, A-908
 Solanogantane, A-908
 Solanopubamide A, A-908
 Solanopubamide B, A-908
 Solanopubamine, A-908
 Solanthrene, S-343
 Solanudine, S-348
 Solanum Alkaloid SL-c, L-116
 Solanum Alkaloid SL-d, L-116
 Solapalmitene, S-340
 Solapalmitine, S-340
 Solaparnaine, S-452
 Solapersine, S-456
 Solaphyllidine, S-349
 Solaplumbine, S-456
 Solaquidine, S-350
 Solaradine, S-456
 Solaradixine, S-456
 Solaseaforthine, S-351
 Solashabanine, S-456
 Solasodamine, S-456
 5 α -Solasodan-3 β -ol (obsol.), S-445
 Solasodanol, S-445
 Solasodan-3-one, S-457
 Solasodene, S-457
 Solasodine 3-*O*- β -D-glucopyranoside, S-456
 Solasodine, S-456
 Solasonine, S-456
 Solaspiralidine, S-352
 Solasuaveoline, S-456
 Solasurine, S-456
 Solateinimine, S-200
 Solatifoline, S-456
 Solatubine, S-345
 Solaurethine, S-340
 Solaverbascine, S-200
 Solaverine III, S-455
 Solaverine II, S-450
 Solaverine I, S-450
 Solaverol A, S-450
 Solaverol B, S-455
 Solavilline, S-456
 Solcodein, C-555
 Solenopsine A, M-584
 Solenopsine B, M-580
 Solenopsine C, M-504
 Solfapterin, A-923
 Solidaline, S-353
 Solimocurarine, S-354
 Solimoosines, S-355
 Sollasin B, S-356
 Sollasin C, S-357
 Solpalmitene, S-340
 Solsodamine A, S-358
 Solsodamine B, S-359
 Somamide A, S-360
 Somniferine, S-361
 Somniferine, S-362
 Somniferinine, S-363
 Somnine, S-364
 Somocystinamide A, S-365
 Songbeimine, C-290
 Songbeimone, C-290
 Songbeisine, H-371
 Songoramine, D-162
 Songorine *N*-oxide, N-30
 Songorine, N-30
 Songorinine, S-366
 Sonodione, S-367
 Sonpeimine, S-368
 Sophazrine, S-369
 Sophoattarine, S-370
 Sophocarpidine, S-371
 Sophocarpine *N*-oxide, S-371
 Sophocarpine, S-371
 Sophochrysin, S-372
 Sophohejrine, S-373
 Sopholupanizidone, S-374
 Sopholupisine, S-375
 Sophoramine, S-376
 Sophoranol *N*-oxide, S-377
 Sophoranol, S-377
 Sophorasine A, S-378
 Sophorasine B, S-378
 Sophorbenzamine, S-379
 Sophoridine *N*-oxide, S-380
 Sophoridine, S-380
 Sophorine, C-938
 Sophorine, O-59
 Sophosalimine, S-381
 Soraphinol A, H-538
 Sorbicillactone A, S-382
 Sorbicillactone B, S-382
 L-Sorbosazine, B-221
 Sorelline, S-383
 Soulidine, M-309
 Soulieotine, D-471
 Souline A, M-347
 Souline B, H-157
 Souline C, M-321
 Souline D, M-356
 Souline E, M-357
 Souline F, H-199
 Spallidamine, H-491
 Spanidin, S-394
 Sparoxomycin A1, S-384
 Sparoxomycin A2, S-384
 Sparsiflorine, T-524
 Sparsogenin, S-384
 Sparsomycin A, T-658
 Sparsomycin, S-384
 Spartalupine, I-322

- Sparteine sulfate, S-385
 11 β -Sparteine, I-321
 Sparteine, S-385
 Spartioidine *N*-oxide, S-240
 Spartioidine, S-240
 Spartocytisine, S-386
 Spathulatine, I-322
 Speciociatinate, G-13
 Speciocolchicine, S-387
 Speciofoline, R-134
 Speciogynine, G-13
 Specionoxeine, R-134
 Speciophylline *N*-oxide, M-661
 Speciophylline, M-661
 Specioritchine, S-387
 Speciosamine, S-387
 Specioseine, S-387
 Speciosine, S-387
 Spectabiline, L-88
 Spectabiline, M-688
 Spectaline, S-388
 Spectalinine, S-388
 Spectamine A, C-183
 Spectamine B, C-183
 Spegazzinidine, S-578
 Spegazzinine, S-578
 Sperabillin A, S-389
 Sperabillin C, S-389
 Speradine A, C-894
 Speranberculatine A, S-390
 Speranerculatine A, S-391
 Speranerculatine B, S-391
 Speranerculatine C, S-391
 Speranskatine A, S-392
 Speranskatine B, S-392
 Speranskilatine A, S-393
 Spergualin, S-394
 Spermacoceine, S-395
 Spermatheridine, L-194
 Spermatherine, S-396
 Spermic acid, S-397
 Spermidine, S-398
 Spermine, S-399
 Spermostrychnine, S-400
 SPF 32629A, A-1266
 SPF 32629B, A-1267
 Sphaeranthine I, S-402
 Sphaeranthine IV, S-403
 Sphaeranthine VII, S-405
 Sphaeranthine VI, S-404
 Sphaeranthine, S-401
 Sphaerocarpine, A-930
 Sphaerophysine, S-406
 Spheroidine, T-298
 Spherophysine, S-406
 Spicaledonine, S-407
 Spicataxine, T-47
 Spicatine A, M-331
 Spicatine B, M-343
 Spicatine, S-408
 Spicigerine, S-409
 Spidamine, J-46
 Spiduxine, S-410
 Spiganthine, R-175
 Spiguetidine, S-411
 Spiguetine, S-411
 Spilanthic acid, D-124
 Spilanthol, D-124
 Spinaceamine, T-192
 Spinacine, T-193
 Spinanine A, S-412
 Spinosine, X-30
 Spiracine, M-502
 Spiradine A, H-164
 Spiradine B, H-164
 Spiradine C, H-164
 Spiradine D, S-421
 Spiradine F, S-418
 Spiradine G, S-418
 Spiraeaine A, S-413
 Spirafine III, S-414
 Spirafine II, S-414
 Spirajine, S-415
 Spiraline, P-98
 Spiramide, A-1534
 Spiramidine A, S-419
 Spiramidine B, S-419
 Spiramine A, S-416
 Spiramine B, S-416
 Spiramine C, S-416
 Spiramine D, S-416
 Spiramine E, A-211
 Spiramine F, A-211
 Spiramine G, A-1533
 Spiramine H, A-1533
 Spiramine I, A-1533
 Spiramine J, S-417
 Spiramine K, S-417
 Spiramine L, S-417
 Spiramine M, S-417
 Spiramine N, B-371
 Spiramine O, B-371
 Spiramine P, S-418
 Spiramine Q, S-418
 Spiramine R, A-211
 Spiramine S, A-1534
 Spiramine T, S-418
 Spiramine U, S-418
 Spiramine V, A-1534
 Spiramine W, S-418
 Spiramine X, S-413
 Spiramine Y, S-413
 Spiramine Z, S-422
 Spiramine Z₂, S-419
 Spiramine Z₃, S-419
 Spiramine, M-502
 Spirasine III, S-421
 Spirasine II, S-421
 Spirasine I, S-421
 Spirasine IV, H-171
 Spirasine IX, H-173
 Spirasine VIII, S-421
 Spirasine VII, S-421
 Spirasine VI, S-421
 Spirasine V, S-421
 Spirasine XI, H-171
 Spirasine XII, D-584
 Spirasine XIII, D-584
 Spirasine XIV, H-165
 Spirasine X, S-420
 Spirasine XV, H-165
 Spiratine A, A-1533
 Spiratine B, S-422
 Spiredine, S-421
 Spireine, S-423
 Spiroarogenic acid, S-424
 1-Spiroaxen-6-amine, S-425
 Spirobrassinin, S-426
 Spirobrassinol, S-427
 Spirocalcaridine A, S-428
 Spirocalcaridine B, S-428
 Spirodihydrobenzofuranlactam I, A-1164
 Spirodihydrobenzofuranlactam II, A-1164
 Spirodihydrobenzofuranlactam III, A-1164
 Spirodihydrobenzofuranlactam IV, A-1164
 Spirodihydrobenzofuranlactam V, A-1164
 Spirodihydrobenzofuranlactam VI, S-489
 Spiroforabuxine, S-429
 Spiroleucettadine, S-430
 Spirolide A, S-431
 Spirolide B, S-431
 Spirolide C, S-432
 Spirolide D, S-432
 Spirolide E, S-433
 Spirolide F, S-433
 Spirolide G, S-434
 Spirolucidine, S-435
 Spiropachysine, S-436
 Spiropyrrrolizidine 238, N-252
 Spiropyrrrolizidine 222, S-437
 Spiropyrrrolizidine 236, S-437
 Spiropyrrrolizidine 252, S-437
 Spiroquinazoline, S-438
 Spirosolan-3-amine, S-439
 Spirosolane-2,3-diol, S-440
 Spirosolane-3,15-diol, S-441
 Spirosolane-3,23-diol, S-442
 Spirosolane-3,23,24-triol, S-443
 Spirosolane-3,23,27-triol, S-444
 Spirosolan-3-ol, S-445
 Spirosol-5-en-3-amine, S-446
 Spirosol-5-ene-3,23-diol, S-450
 Spirosol-5-ene-3,12-diol, S-447
 Spirosol-5-ene-3,15-diol, S-448
 Spirosol-5-ene-3,21-diol, S-449
 Spirosol-5-ene-3,25-diol, S-451
 Spirosol-5-ene-3,27-diol, S-452
 Spirosol-5-ene-3,12,27-triol, S-454
 Spirosol-5-ene-3,23,27-triol, S-455
 Spirosol-5-ene-3,12,23-triol, S-453
 Spirosol-5-en-3-ol, S-456
 Spirosol-4-en-3-one, S-457
 Spirostan-3-amine, S-458
 Spirostaphylotrichin C, T-625
 Spirostaphylotrichin D, T-625
 Spirostaphylotrichin E, S-459
 Spirostaphylotrichin F, S-459
 Spirostaphylotrichin G, S-460
 Spirostaphylotrichin H, S-460
 Spirostaphylotrichin I, S-466
 Spirostaphylotrichin K, S-461
 Spirostaphylotrichin L, S-459
 Spirostaphylotrichin M, S-462
 Spirostaphylotrichin N, S-463
 Spirostaphylotrichin O, S-463
 Spirostaphylotrichin P, S-463
 Spirostaphylotrichin Q, S-464
 Spirostaphylotrichin R, S-465
 Spirostaphylotrichin S, S-466
 Spirostaphylotrichin T, S-467
 Spirostaphylotrichin U, S-465
 Spirostaphylotrichin V, S-465
 Spirotryprostatin A, S-468
 Spirotryprostatin B, S-468
 Spixianine *N*-oxide, S-469
 Spixianine, S-469
 Splendaboline, S-627
 Splendidine, S-470
 Splendoline, S-605
 Spongiacidin A, H-762
 Spongiacidin B, H-762
 Spongiacidin C, A-1576
 Spongiacidin D, A-1576
 Spongiacysteine, S-471
 Spongidine A, S-472
 Spongidine B, S-472
 Spongidine C, P-289
 Spongidine D, S-473
 Spongolactam A, D-763
 Spongolactam B, D-764
 Spongolactam C, D-763
 Spongopurine, A-842
 Spongotine A, T-432
 Spongotine B, T-432
 Spongotine C, T-432
 Spophyllin retard, T-360
 Sporangiomycin, S-321
 Sporidesmin A, S-474
 Sporidesmin B, S-474
 Sporidesmin C, S-475
 Sporidesmin D, S-476
 Sporidesmin E, S-477
 Sporidesmin F, S-478
 Sporidesmin G, S-479
 Sporidesmin H, S-480
 Sporidesmin J, S-474
 Sporidesmin, S-474
 Sporovexin A, S-481
 Sporovexin C, S-481
 Sprintillamine, S-482
 Sprintilline, S-483
 Spylidone, S-484
 SQ 02-S-L1, A-1268
 SQ 02-S-L2, S-492
 SQ 02-S-V2, A-1269
 SQ 02-S-V1, S-492
 Squalamine lactate, S-485
 Squalamine, S-485
 Squalidine, S-238

- Strictosidinic ketone, S-584
 Strigosine, S-585
 Strobamine, S-586
 Strobiline, S-587
 Strobolamine, S-586
 Struxine, I-12
 Strychnambarine, S-588
 Strychnicine, I-12
 Strychnidine, S-589
 Strychnidin-10-one, S-589
 Strychnine *N*-oxide, S-589
 Strychnine sulfate, S-589
 ψ -Strychnine, P-726
 Strychnine, S-589
 Strychnobaridine, S-590
 Strychnobiline, S-591
 Strychnobrasiline, S-594
 Strychnocarpine, T-226
 Strychnochromine, S-592
 Strychnochryesine, S-593
 Strychnofenderline, S-594
 Strychnofluorine, N-301
 Strychnofoline, S-595
 Strychnofuranine, S-596
 Strychnogucine A, S-597
 Strychnogucine B, S-598
 Strychnogucine C, S-597
 Strychnohexamine, S-599
 Strychnohirsutine, S-600
 Strychnopentamine, S-601
 Strychnophylline, S-602
 Strychnopivotine, S-603
 Strychnorubigine, I-319
 Strychnosilidine, S-604
 Strychnosiline, S-604
 Strychnospermine, S-400
 Strychnosplendine, S-605
C-Strychnotoxine Ia, C-31
C-Strychnotoxine I, C-31
 Strychnotoxine II, T-437
 Strychnovoline, T-145
 Strychnoxanthine, S-606
 Strychnozairine, S-607
 Strynervene, S-589
 Stryphon, A-736
 STX, S-109
 Styelsamine A, S-609
 Styelsamine B, S-609
 Styelsamine C, S-608
 Styelsamine D, S-609
 Stylissadine A, S-610
 Stylissadine B, S-610
 Styloguanidine, S-611
 Stylophorine, C-357
 Stylophylline, H-394
 Stylopine, S-612
 Stylotellane A, S-613
 Stylotellane B, S-613
 Stylotellin, S-614
 Stypnon, A-736
 2-Styrylindole, P-341
N-Styryllactamide, X-27
N-Styryl-4(3*H*)-quinazolinone, Q-14
 Suaedin, S-615
 Suavedol, P-643
 Suaveolenine, S-616
 Suaveolindole, S-617
 Suaveoline, S-618
 Suaveoline, T-248
 Subaphylline, C-456
 Subarine, S-619
 Subcosine II, L-53
 Subcosine I, L-53
 Subcumine, M-323
 Subcusine, M-323
 Subdesculine, D-162
 Subereatensin, S-620
 Suberedamine A, S-621
 Suberedamine B, S-621
 Subhirsine, S-622
 Subincanadine A, S-623
 Subincanadine B, S-623
 Subincanadine C, S-623
 Subincanadine D, S-624
 Subincanadine E, S-624
 Subincanadine F, S-625
 Subincanadine G, S-623
 Subincanine, S-626
 Subsessiline lactone, A-676
 Subsessiline, A-676
 Subsessiline, S-627
 Substance B, D-92
 Substance C, D-92
 Substance E₁, D-92
 Substance I, L-287
 Substance J, L-287
 Substance S, D-92
 Substance T_a, D-92
 Substance U, D-92
 Succinopyoverdin G173, S-628
N-Succinoylanthranilic acid, A-706
*N*²-Succinoylarginine, A-1413
 Succirubine, S-629
 Suffruticodine, S-630
 Suffruticoline, S-631
 Sugordomycin D_{1a}, C-711
 Sugordomycin D_{1d}, C-712
 Suhailamine, A-537
 Suillumide, A-762
 Sunin, S-632
 Sukhodianine β -*N*-oxide, C-733
 Sukhodianine, C-733
 Sulcatine, C-146
 10-*O*-Sulfatobastadin 3, B-31
 Sulfinemycin, S-633
 Sulfobacin A, S-634
 Sulfobacin B, S-635
 34-*O*-Sulfobastadin 9, B-32
 34-Sulfobastadin 13, B-33
N-(2-Sulfoethyl)pyridinium betaine, S-636
 Sulfoglucobrassicin, I-122
 Sulfohalopterin 2, S-637
 4-*O*-Sulfohemibastadin 1, H-100
 4-*O*-Sulfohemibastadin 2, H-100
 1-*O*-Sulfohemibastadin 1, H-100
 1-*O*-Sulfohemibastadin 2, H-100
 (1-Sulfo-1*H*-indol-3-yl)methyl glucosinolate, I-122
 Sulfomycin, S-638
 3-(Sulfoxy)tyrosine, A-745
 7-Sulfopseudobactin A, P-702
 Sulfostin, S-639
 Sulphostin, S-639
 Sulpinine A, S-640
 Sulpinine B, S-640
 Sulpinine C, S-641
 Sun 0588, T-142
 Sungpaconitine, M-303
 Sungucine, S-642
 Suomilide, S-643
 (-)-Supinidine curassavate, C-810
 Supinidine, S-644
 Supinine, A-669
 Supral, V-34
 Suprarenin, A-152
 Surinamine, T-700
 Surugatoxin, S-645
 Sutchueneneone, S-646
 Sutchuenenine, S-647
 Sutchuenensine, S-648
 Sutherlandin *trans-p*-coumarate, H-510
 Sutherlandin epoxide, H-510
 Sutherlandin, H-510
 Suvanine, S-649
 Sventrine, O-123
 SW 163C, A-1271
 SW 163D, A-1304
 SW 163E, A-1304
 SW 163F, A-1304
 SW 163G, A-1304
 Swainsonine *N*-oxide, S-650
 Swainsonine, S-650
 Swatinine, M-339
 Swazine, S-651
 SW-B, T-603
 Sweetinine, E-99
 Swietenidin A, Q-41
 Swietenidin B, Q-45
 Swinhoeamide A, S-652
 Sycophantine, S-456
 Sylvamide, D-566
 Sylvatine, S-653
 Symbioimine, S-654
 Symlandine, E-31
 Sympathol, S-659
 Sympectothione, E-173
 Symphytine, E-31
 Symplostatin 1, D-907
 Symviridine, E-32
 Synaine, S-655
 Synapton, P-400
 Syncarpamide, M-18
 (\pm)-Syncarpamide, M-18
 Syncarpine, P-424
 Syncarpurea, S-656
 Syncothrecin A, O-131
 Syndihydrohistrionicotoxin, H-309
 Synechobactin A, S-657
 Synechobactin B, S-657
 Synechobactin C, S-657
 Syneilesine, S-658
 Synephrine acetone, S-659
 Synephrine, S-659
 Synerazol, S-660
 Synergistin A₁, O-131
 Synthetic capsaicin, C-100
 Syntometriline, E-164
 Syphilobin A, S-661
 Syphilobin F, S-662
 Syriamycin, S-663
 Syringolin A, S-664
 Syringolin B, S-664
 Syringolin C, S-665
 Syringolin D, S-666
 Syringolin E, S-666
 Syringolin F, S-667
 16-*O*-Syringoylbuxaminol E, B-456
 Syringyltrachelanthamidine, H-629
 Szechenyine, M-331
 Szovitsamine *N*-oxide, P-221
 Szovitsamine, P-221
 Szovitsidine, A-1005
 Szovitsinine, P-221
 T 988A, G-99
 T 988B, A-1272
 T 988C, G-99
 T 41348, H-514
 T 1384, N-180
 T 1401, T-142
 TA 3037A, A-1273
 TA Toxin A, T-612
 Tabascanine, S-604
 Taberhanine, T-11
 Tabernaebovine, T-1
 Tabernaeelegantidine, T-2
 Tabernaeelegantine A, C-616
 Tabernaeelegantine B, C-615
 Tabernaeelegantine C, C-616
 Tabernaeelegantine D, C-615
 Tabernaeelegantinine A, T-3
 Tabernaeelegantinine B, T-4
 Tabernaeelegantinine C, T-5
 Tabernaeelegantinine D, C-615
 Tabernaemontabovine, T-6
 Tabernaemontanine, D-935
 Tabernaemontaninol, D-935
 Tabernaemontavine, T-7
 Tabernamine, T-8
 Tabernanthine, I-4
 Tabernoschizine, A-1365
 Tabernoxidine, T-9
 Tabernulosine, P-417
 Taberpsychine, T-10
 (19*Z*)-Taberpsychine, T-10
 Tabersonine α -epoxide, T-11
 Tabersonine β -epoxide, T-11
 Tabersonine *N*^b-oxide, T-11
 ψ -Tabersonine, P-733
 Tabersonine, T-11
 Tabex, C-938

- Tabouensinium, G-29
 Tacamine, T-12
 Tacamonine, T-13
 Tachmalin, A-216
 Tacraline, T-15
 Tadzhaconine, H-181
 Taifine, F-212
 Taipaiemine, D-562
 Taipienine, D-562
 Taiwanamide A, D-718
 Taiwanamide B, D-718
 Taiwanamide C, M-219
 Takaobase I, M-309
 Takaokamycin, H-358
 Takaonine, M-363
 Takaosamine, M-309
 Takatonine, T-16
 Takawobase I, M-309
 Talaroconvolutin A, O-134
 Talaroconvolutin B, O-134
 Talaroconvolutin C, D-141
 Talaroconvolutin D, D-141
 Talarotoxin, T-17
 Talassamine, T-18
 Talassicumine A, M-352
 Talassicumine B, D-611
 Talassicumine C, M-381
 Talassimidine, T-18
 Talassimine, T-18
 Talatisamine, M-352
 Talatisidine, M-352
 Talatizamine, M-352
 Talatizidine, M-352
 Talatizine, H-209
 Talbotine, T-19
 Talcamine, C-806
 Talcarpine, T-20
 Talgic acid, O-32
 Taliscanine, T-539
 Talitine A, M-344
 Talitine B, M-315
 Talitine C, M-344
 Talizopine, T-346
 Talmine, T-341
 Talpinine, T-21
 Talval, H-488
 Tambjamine aldehyde, M-244
 Tambjamine A, T-22
 Tambjamine B, T-22
 Tambjamine C, T-22
 Tambjamine D, T-22
 Tambjamine E, T-22
 Tambjamine F, T-22
 Tambjamine G, T-22
 Tambjamine H, T-22
 Tambjamine I, T-22
 Tambjamine J, T-22
 Tamynine, T-23
 TAN 1263, A-1275
 TAN 1280, A-1276
 TAN 1307, A-1277
 TAN 1813, A-1278
 TAN 2020, A-1279
 TAN 883, A-1274
 TAN 1030A, A-1281
 TAN 1057A, A-1282
 TAN 1251A, A-1284
 TAN 1263A, A-1275
 TAN 876A, A-1280
 TAN 883A, A-1274
 TAN 950A, A-733
 TAN 1771A, D-574
 TAN 868A, K-43
 TAN 1169A, P-400
 TAN 749A, S-389
 TAN 1057B, A-1282
 TAN 1251B, A-1284
 TAN 1263B, A-1275
 TAN 876B, A-1283
 TAN 1771B, D-572
 TAN 868B, K-43
 TAN 1169B, P-400
 TAN 883B, X-6
 TAN 1142, C-325
 TAN 1057C, A-1285
 TAN 1251C, A-1284
 TAN 1263C, A-1275
 TAN 883C, A-1274
 TAN 749C, S-389
 TAN 1057D, A-1285
 TAN 1251D, A-1284
 TAN 1263D, A-1275
 TAN 883D, A-1274
 TAN 588, L-8
 TAN 999, S-498
 Tanakamine, T-24
 Tanakine, T-25
 Tanganil, L-153
 Tangirine, T-26
 Tangutisine, H-189
 Tangutorine, T-27
 Tannagine, T-28
 Tantazole A, T-29
 Tantazole B, T-30
 Tantazole F, T-31
 Tantazole I, T-29
 Tanwusine A, H-188
 Tanwusine, H-203
 Taraktophyllin, D-565
 Taraxacine A, P-914
 Taraxacine B, C-133
 Tardioxopiperazine A, T-32
 Tardioxopiperazine B, T-33
 Tardolyt, M-262
 Tarenine, T-34
 Tarichatoxin, T-298
 Tashiromine, O-47
 Tasmanine, T-35
 Taspine, T-36
 Tasteless Quinine, Q-20
 Tataramide B, E-128
 Tatiopterin, T-37
 Tatsidine, M-324
 Tatsiensine, M-371
 Tatsinine, M-324
 Tatsirine, H-198
 Taumustine, A-765
 Taurenine, M-313
 Tauret, T-38
 Taurine, A-765
 Taurine, P-32
 Tauroacidin A, T-39
 Tauroacidin B, T-39
 Taurocayamine, G-204
 Taurodispacamide A, T-39
 Taurohaterumalide NA, H-76
 Tauropinnaic acid, P-435
 Taveuniamide A, T-40
 Taveuniamide B, T-40
 Taveuniamide C, T-40
 Taveuniamide D, T-40
 Taveuniamide E, T-40
 Taveuniamide F, D-357
 Taveuniamide G, D-357
 Taveuniamide H, D-357
 Taveuniamide I, D-357
 Taveuniamide J, D-357
 Taveuniamide K, D-357
 Tawicyclamide A, T-41
 Tawicyclamide B, T-42
 4(20),11-Taxadiene-1,2,5,7,9,10,13-heptol, T-43
 4(20),12-Taxadiene-2,5,7,9,10,11,13-heptol, T-44
 4(20),11-Taxadiene-1,2,5,9,10,13-hexol, T-45
 4(20),11-Taxadiene-2,5,7,9,10,13-hexol, T-46
 4(20),11-Taxadiene-2,5,9,10,13-pentol, T-47
 4(20),11-Taxadiene-5,7,9,10,13-pentol, T-48
 4(20),11-Taxadiene-5,9,10,13-tetrol, T-49
 Taxcultine, T-52
 Taxicin II, T-273
 Taxilamine, T-50
 Taxine A, A-6
 Taxine B, T-45
 Taxine C, A-6
 Taxine II, T-273
 Taxine I, T-45
 Taxiphyllin, H-520
 Taxodine, T-51
 Taxol A, T-52
 Taxol B, T-52
 Taxol C, T-52
 Taxol D, T-52
 Taxoline, T-52
 Taxol, T-52
 Taxuspinanane A, E-141
 Taxuspinanane E, T-52
 Taxuspinanane I, T-52
 Taxuspinanane J, T-52
 Taxuspine H, S-407
 Taxuspine N, E-141
 Taxuspine P, T-44
 Taxuspine Z, T-47
 Taxuyunnanin A, T-52
 Tazettine, T-53
 Tazettinol, T-53
 TB Toxin, T-17
 TBA, T-457
 TC 2403, M-232
 Tchibangensine, U-52
 Tecleabine, F-222
 Tecleamine, F-217
 Tecleanatalensine A, F-221
 Tecleanatalensine B, F-221
 Tecleanine, T-54
 Tecleanone, T-55
 Tecleanthine, T-56
 Tecleaverdine, T-57
 Tecleaverdoornine, T-57
 Tecleine, F-217
 Tecleoxine, F-221
 Tecomanine, T-58
 Tecomine, T-58
 Tecostanine, T-59
 Tecostidine, T-60
 Tedanalactam, O-146
 Tegerrardin A, D-528
 Tegerrardin B, D-528
 Tehaunine *N*-oxide, T-238
 Tehaunine, T-238
 Teidine, A-138
 Teinemine, S-200
 Tejedine, S-192
 Tejeramycin, T-61
 Teladiazoline, T-62
 Telastaside, P-58
 Telazoline, T-62
 Teleocidin A₁, L-367
 Teleocidin A₂, L-367
 Teleocidin B₁, T-63
 Teleocidin B₂, T-63
 Teleocidin B₃, T-63
 Teleocidin B₄, T-63
 Telepathine, H-592
 Teliglazine, M-712
 Telikovinone, T-64
 Telisatin A, T-65
 Telisatin B, T-65
 Telitoxine, T-498
 Telitoxinone, T-64
 Telobine, A-1331
 Teloidine, M-392
 Telomestatin, T-66
 Tembamide acetate, T-67
 Tembamide, T-67
 Tembetarine, R-59
 Templetine, T-68
 Temuconine, B-98
 (-)-Temuconine, G-177
 Tenacibactin A, T-69
 Tenacibactin B, T-69
 Tenacibactin C, T-70
 Tenacibactin D, T-71
 Tenaglin, A-716
 Tenellin, T-72
 Tensibar, R-52
 Tensiplex, E-6
 Tenuamine, T-73
 Tenuazonic acid, T-74
 Tenucaulin B, A-716
 Tenucyclamide A, T-75

- Tenuencyclamide B, T-75
Tenuencyclamide C, T-76
Tenuencyclamide D, T-76
Tenucaulin A, A-716
Tenucausine, T-77
Tenuidine, T-78
Tenuilobine, T-79
Tenuiphylline, T-80
Tenuipine, T-81
Tenuisine A, L-44
Tenuisine B, L-44
Tenuisine C, L-44
Teonova, T-360
Tepenine, T-166
Teperine, C-419
Tepidopterin, A-749
Terapterin, P-772
Terdefine, M-307
Terengganensine A, T-82
Terengganensine B, T-83
Terezine A, T-84
Terezine B, T-85
Terezine C, T-85
Terezine D, T-86
Terihanine, N-239
[3,2':2'(3'H),3''-Ter-1*H*-indol]-3'-one, T-87
[3,3':3'(2'H),3''-Ter-1*H*-indol]-2'-one, T-88
Terminaline, A-879
Termine, T-89
Termisine, T-90
Ternatidine, H-193
Ternatine, H-193
Ternatoside C, R-168
Ternatoside D, R-168
Teropterin, P-772
Terpendole A, T-93
Terpendole B, T-92
Terpendole C, T-93
Terpendole D, T-92
Terpendole E, P-108
Terpendole F, P-108
Terpendole G, P-108
Terpendole H, T-94
Terpendole I, T-91
Terpendole J, T-91
Terpendole K, T-93
Terpendole L, T-95
Terpendole M, T-93
Terpeptin, T-96
[*p*-Terphenyl]-2',3',4,4'',5',6'-hexol, T-97
3,2':4',3''-Terpyridine, T-98
Terragine A, T-99
Terragine B, T-100
Terragine C, T-101
Terragine D, T-102
Terregens factor, A-1451
Terresoxazine, T-103
Terrestriamide, A-786
Terrestribisamide, D-362
Terrestrial acid, T-104
Terreusinone, T-105
Terricolin, P-922
Terriquinone A, T-106
Tessellatine, H-93
Tetraacetylclionamide, C-527
Tetraacetylthylenediamine, E-225
1,5,8,12-Tetraazadodecane, B-153
4,8,12,16-Tetrazaeicosane-1,20-diamine, D-283
4,8,12,17-Tetrazaeicosane-1,20-diamine, D-284
4,8,13,17-Tetrazaeicosane-1,20-diamine, D-285
1,6,11,16-Tetraazahexadecane, B-150
4,8,12,16-Tetraazanonadecane-1,19-diamine, T-107
1,5,10,15-Tetraazapentadecane, D-250
1,6,10,15-Tetraazapentadecane, D-251
5,10,15,20-Tetraazatetracosane-1,24-diamine, D-286
1,5,9,14-Tetraazatetradecane, D-249
1,5,10,14-Tetraazatetradecane, S-399
1,5,9,13-Tetraazatridecane, B-154
2,2',6,6'-Tetrabromo-3,3'-bi-1*H*-indole, T-109
2,2',5,5'-Tetrabromo-3,3'-bi-1*H*-indole, T-108
4,4',6,6'-Tetrabromo-2,2'-bis(methylsulfinyl)-3,3'-bi-1*H*-indole, T-110
4,4',6,6'-Tetrabromo-2,2'-bis(methylthio)-3,3'-bi-1*H*-indole, T-110
3,3',4,4'-Tetrabromo-5,5'-dichloro-1,1'-dimethyl-2,2'-bi-1*H*-pyrrole, T-111
5,5',7,7'-Tetrabromo-6,6'-dimethoxyindigotin, T-112
2,3,4,6-Tetrabromo-1*H*-indole, T-113
2,3,4,7-Tetrabromo-1*H*-indole, T-114
2,3,5,6-Tetrabromo-1*H*-indole, T-115
2,3',5,5'-Tetrabromo-7'-methoxy-3,4'-bi-1*H*-indole, T-116
3,3',5,5'-Tetrabromo-7'-methoxy-1,4'-bi-1*H*-indole, T-117
2,3,4,6-Tetrabromo-1-methyl-1*H*-indole, T-113
2,3,5,6-Tetrabromo-1-methyl-1*H*-indole, T-115
4,4',6,6'-Tetrabromo-2-(methylsulfinyl)-2-(methylthio)-3,3'-bi-1*H*-indole, T-110
2,4,5,6-Tetrabromo-3-methylthio-1*H*-indole, T-473
2,3,4,5-Tetrabromo-1*H*-pyrrole, T-118
Tetrabromostyloguanidine, S-611
3,5,6,10-Tetrachloro-1-(5-chloro-2-hydroxybenzoyl)-2*H*,8*H*-dipyrrolo[3,4-*b*:1',2'-*d'*][1,4]benzoxazocin-8-one, A-1133
2,3,4,7-Tetrachloro-1*H*-indole, T-119
N-Tetracosanoanthranilic acid, A-706
N^b-Tetracosanoylserotonin, H-755
N^b-Tetracosanoyltryptamine, T-639
N-Tetracosanoilytyramine, H-667
15-Tetracosenoic acid, T-120
N¹,N⁵,N¹⁰,N¹⁴-Tetra-*trans-p*-coumaroylspermine, S-399
3,5-Tetradecadien-2-amine, T-131
2,4-Tetradecadien-8,10-dienoic acid piperidide, T-121
2,4-Tetradecadiene-8,10-dienoic acid isobutylamide, A-968
2,4-Tetradecadiene-8,10-dienoic acid isopentylamide, T-121
2,4-Tetradecadiene-8,10-dienoic acid pyrrolidide, T-121
10,12-Tetradecadiene-4,6-dienoic acid, T-122
2,4-Tetradecadiene-8,10-dienoic acid, T-121
2,4-Tetradecadiene-8,10,12-trienoic acid, T-123
2,4-Tetradecadienoic acid *p*-hydroxyphenethylamide, H-670
2,4-Tetradecadienoic acid isobutylamide, T-124
2,4-Tetradecadienoic acid piperidide, T-124
2,4-Tetradecadienoic acid pyrrolidide, T-124
2,4-Tetradecadienoic acid, T-124
1-(2,4-Tetradecadienyl)piperidine, T-124
N-(2,4-Tetradecadienyl)pyrrolidine, T-124
2,4-Tetradecadien-8-ynoic acid pyrrolidide, T-125
2,4-Tetradecadien-8-ynoic acid, T-125
2,4,6,8,10,12-Tetradecahexaenedioic acid, T-126
Tetradecahydro-2*H*,11*H*-tripyrido[1,2-*a*:1',2'-*c*:3'',2''-*e*]pyrimidine, I-340
N-Tetradecanoylarginine, A-1413
N²-Tetradecanoylglutamine, G-108
N-Tetradecanoylholothin, A-759
N-Tetradecanoyltyrosine, T-700
2,4,8,10,12-Tetradecapentaenoic acid, T-127
2,4,6,12-Tetradecatetraene-8,10-dienoic acid isobutylamide, T-128
2,4,6,12-Tetradecatetraene-8,10-dienoic acid isopentylamide, T-128
2,4,6,12-Tetradecatetraene-8,10-dienoic acid piperidide, T-128
2,4,6,12-Tetradecatetraene-8,10-dienoic acid, T-128
2,4,8,11-Tetradecatetraenoic acid pyrrolidide, T-129
2,4,8,11-Tetradecatetraenoic acid, T-129
2,4,8,11-Tetradecatetraenoic acid piperidide, T-129
(2,4,8,11-Tetradecatetraenyl)piperidine, T-129
2,4,10,12-Tetradecatetraen-8-ynoic acid, T-130
3,5,13-Tetradecatrien-2-amine, T-131
2,4,12-Tetradecatrien-8,10-dienoic acid isopentylamide, T-132
2,4,12-Tetradecatrien-8,10-dienoic acid piperidide, T-132
2,4,12-Tetradecatrien-8,10-dienoic acid pyrrolidide, T-132
2,4,12-Tetradecatrien-8,10-dienoic acid, T-132
2,4,8-Tetradecatrienoic acid isobutylamide, T-133
2,4,8-Tetradecatrienoic acid, T-133
2,4,8-Tetradecatrienoic acid piperidide, T-133
N-(2,4,8-Tetradecatrienyl)piperidine, T-133
N-(5,8,11-Tetradecatrienyl)tyramine, H-667
8-(5,8,11-Tetradecatrienyl)-1,5,9,13-tetraazacycloheptadecan-6-one, B-386
2,4,10-Tetradecatrien-8-ynoic acid isobutylamide, T-134
2,4,10-Tetradecatrien-8-ynoic acid piperidide, T-134
2,4,10-Tetradecatrien-8-ynoic acid pyrrolidide, T-134
2,4,10-Tetradecatrien-8-ynoic acid, T-134
2,4-Tetradeca-8,10,12-trienoic acid piperidide, T-123
N²-(9-Tetradecenyl)glutamine, G-108
N-(8-Tetradecenyl)tyrosine, T-700
2-Tetradecylpyrrolidide, T-135
2-Tetradecyl-4(1*H*)-quinolinone, H-746
8-Tetradecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-386
12,13,25,26-Tetradehydrocarpaine, C-168
Tetradehydrochelanthifoline, G-181
17,4',5',6'-Tetradehydro-3 α -cinchophylline, C-448
2,16,19,20-Tetradehydrocuran-17-al, N-301
16,17,19,20-Tetradehydrocuran-3,17-diol, D-688
16,17,19,20-Tetradehydrocuran, Z-14
1,2,19,20-Tetradehydro-1-demethylajmalan-17,21-diol 17-acetate, V-194
3,4,5,6-Tetradehydrodihydrocorynantheol, T-136
1,2,6,7-Tetradehydro-12,17-dihydro-3-methoxy-16(15*H*)-oxaerythrinan-15-one, 11CI, E-215
1,2,6,7-Tetradehydro-15,16-dimethoxyerythrinan-3-ol, E-203
1,2,6,7-Tetradehydro-3,15-dimethoxyerythrinan-16-ol, E-203
3,4,5,6-Tetradehydrodolicchantoside, L-316
Tetradehydrocinulinin, T-137
1,2,18,19-Tetradehydro-3,17-epoxy-7,20(2*H*,19*H*)-cyclovobasan, K-96
6,7,8,14-Tetradehydro-4,5-epoxy-3,6-dimethoxy-17-methylmorphinan, T-350
6,7,8,14-Tetradehydro-4,5-epoxy-6-methoxy-17-methylmorphinan-3-ol, T-350
3,4,5,6-Tetradehydrogeissoschizine, T-138
3,4,5,6-Tetradehydrogeissoschizol, T-136
Tetradehydrohalicyclamine A, H-22
5,6,8,14-Tetradehydro-3-hydroxy-2,6-dimethoxymorphinan-7-one, F-80
3,4,14,15-Tetradehydro-5-hydroxy-14-methoxy-17-methyl-4,17-secoserratinan-13-one, L-337
Tetradehydroisocalycanthine, T-139
Tetradehydroisocinchophyllamine, C-448
3',4',5',6'-Tetradehydroolongicaudatine Y, L-250
11,12,13,14-Tetradehydrmatridin-15-one, S-376
1,2,6,7-Tetradehydro-3-methoxyerythrinan-15-ol, C-543
1,2,6,7-Tetradehydro-3-methoxy-15,16-[methylenebis(oxy)]erythrinan, E-205
5,6,8,14-Tetradehydro-6-methoxy-17-methyl-2,3-[methylenebis(oxy)]morphinan-7-one, A-955
3,4,5,6-Tetradehydromitragynine, M-657
3,4,5,6-Tetradehydroochroposinine, T-136
Tetradehydroocoteine, O-27
3,4,5,6-Tetradehydroopalicoside, L-316
11,12,13,14-Tetradehydro-14-(phenylmethyl)matridin-15-one, S-379
2,3,4,5-Tetradehydropropylallothreonylseryl-3-phenyl- β -alanyl-2-aminobutanoic acid, A-1512
Tetradehydroprotosetemonine, P-670
Tetradehydroreticuline, P-80
3,4,5,6-Tetradehydrositsirikine, S-325
Tetradehydrostemonine, S-519
Tetradehydrostrychnohirsutine, S-600
3,4,5,6-Tetradehydrototalbotine, T-19
7,8,12,14-Tetradehydro-5,6,12,13-tetrahydro-3,13,23-trihydroxyveratraman-6-one, T-588
7,8,13,13*a*-Tetradehydro-2,3,9,10-tetrahydroxyberbinium(1+), P-34
1,2,6,7-Tetradehydro-3,15,16-trimethoxyerythrinan-11-ol, E-206

- 1,2,6,7-Tetrahydro-3,15,16-trimethoxyerythrinan-8-one, E-202
- 1,2,6,7-Tetrahydro-3,15,16-trimethoxy-C-homocerythrinan, F-143
- 1',2',3',4'-Tetrahydrotribulosine, T-675
- 14,15,16,17-Tetrahydroveratraman-3,23-diol, V-71
- 3,4,5,6-Tetrahydroyohimbine, T-140
- 3,4,5,6-Tetrahydro- β -yohimbinium, T-140
- N,N,N',N'*-Tetraethylthanediamine, E-225
- 2',3',4',6'-Tetra-*O*-galloylprunasin, H-673
- Tetrahydranagyrine, L-302
- Tetrahydrokagerine, T-141
- Tetrahydroalstonial, A-214
- Tetrahydroalstonine (4*R*)-*N*-oxide, A-212
- Tetrahydroalstonine, A-212
- Tetrahydroanabasine, P-460
- 3,14,18,19-Tetrahydroangustine, A-1019
- 2,3,4,13-Tetrahydro-1*H*-benz[*g*]indolol[2,3-*a*]quinolizin-6-ium, S-228
- 5,6,13,13*a*-Tetrahydro-8*H*-benzo[*g*]-1,3-benzodioxolo[5,6-*a*]quinolizin-8-one, G-226
- Tetrahydroberberine, C-73
- Tetrahydroberberubrine, C-73
- 5,6,7,8-Tetrahydrobiopterin, T-142
- 1,2,3,6-Tetrahydro-2,3'-bipyridine, T-143
- 3,4,5,6-Tetrahydro-2,3'-bipyridine, T-144
- 6,7,7*a*,8-Tetrahydro-5*H*-bis[1,3]benzodioxolo[6,5,4-*de*:4',5'-*g*]quinoline, O-143
- 6,7,12*b*,13-Tetrahydro-4*H*-bis[1,3]benzodioxolo[5,6-*a*:4',5'-*g*]quinoline, S-612
- 1,2,5,6-Tetrahydro-2,6-bis(2-hydroxybutyl)-1-methylpyridine, D-391
- 2-(4,4',5,5'-Tetrahydro[4,4'-bithiazol]-2-yl)phenol, U-7
- Tetrahydrobungeanol, T-124
- Tetrahydrocantleyine, T-145
- 1,2,3,4-Tetrahydro- β -carboline-3-carboxylic acid, T-224
- 1,2,3,4-Tetrahydro- β -carboline-1,3-dicarboxylic acid, T-147
- 1,2,3,4-Tetrahydro- β -carboline, T-146
- Tetrahydrocolumbamine, T-1210
- Tetrahydrocoptisine, S-612
- Tetrahydrocorysamine *N*-methosalt, T-148
- Tetrahydrocorysamine, T-148
- Tetrahydrocytisine, T-149
- Tetrahydrodarlingianine, D-74
- 3*z*,5*z*-Tetrahydrodeoxycordifolic acid, C-153
- Tetrahydrodeoxycordifoline lactam, T-150
- Tetrahydrodeoxycordifoline, C-153
- 7,7',8,8'-Tetrahydro-4,5:4',5'-diepoxy-17,17'-dimethyl[2,2'-bimorphinan], M-703
- 5,6,7,8-Tetrahydro-5,7-dihydroxy-1*H*-azocin-2-one, T-151
- 1,2,3,4-Tetrahydro-1,6-dihydroxy- β -carboline, T-225
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-1,1-dimethyl-3-isoquinolinecarboxylic acid, T-153
- 1,2,3,4-Tetrahydro-7,8-dihydroxy-1,1-dimethyl-3-isoquinolinecarboxylic acid, T-154
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-1,1-dimethylisoquinoline, T-152
- 2,3,4,6-Tetrahydro-3,4-dihydroxy-2,2-dimethyl-5*H*-pyrano[3,2-*c*]quinolin-5-one, F-90
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-1-(4-hydroxybenzyl)isoquinoline, H-275
- 1,2,3,4-Tetrahydro-5,8-dihydroxy-1-(4-hydroxybenzyl)-2-methylisoquinoline, T-155
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-2-(2-hydroxy-1-hydroxymethylethyl)-1,1-dimethylisoquinoline, T-152
- 2,3,5,7*a*-Tetrahydro-1,7*a*-dihydroxy-7-(hydroxymethyl)-4-methyl-1*H*-pyrrolizinium, O-136
- 4,5,6,7-Tetrahydro-4,5-dihydroxy-1*H*-indole-3-carboxaldehyde, T-156
- 4,5,6,7-Tetrahydro-6,7-dihydroxy-1*H*-indol-4-one, T-157
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-3-isoquinolinecarboxylic acid, T-160
- 1,2,3,4-Tetrahydro-6,7-dihydroxyisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7,8-dihydroxyisoquinoline, T-159
- [3-(1,2,3,4-Tetrahydro-6,8-dihydroxy-1-isoquinolinyl)propyl]guanidine, F-229
- 3,4,6,7-Tetrahydro-3,4-dihydroxy-6-methoxy-3-methyl-7-methylene-2-pentylpyrano[2,3-*c*]pyrrol-5(2*H*)-one, P-302
- 1,2,3,11*a*-Tetrahydro-3,8-dihydroxy-7-methoxy-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one, N-145
- 1,2,3,12-Tetrahydro-1,2-dihydroxy-6-methoxy-3,3,12-trimethyl-7*H*-pyrano[2,3-*c*]acridin-7-one, D-417
- 1,2,19,20-Tetrahydro-19,20-dihydroxy-1-methylakuumilan-17-oic acid γ -lactone, R-21
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-1-methyl-3-isoquinolinecarboxylic acid, T-164
- 1,2,3,4-Tetrahydro-4,8-dihydroxy-2-methylisoquinoline, T-161
- 1,2,3,4-Tetrahydro-6,7-dihydroxy-1-methylisoquinoline, T-162
- 1,2,3,4-Tetrahydro-7,8-dihydroxy-1-methylisoquinoline, T-163
- 3-(5,10,11,11*a*-Tetrahydro-9,11-dihydroxy-8-methyl-5-oxo-1*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-2-yl)-2-propenamide, A-1081
- 1,5,6,10*b*-Tetrahydro-8,9-dihydroxypyrrolo[2,1-*a*]isoquinolin-3(2*H*)-one, T-629
- 5,5*a*,12,12*a*-Tetrahydro-7,14-dimethoxybenzo[1'',2''-4,5;4',5':4',5']difuro[2,3-*b*:2',3'-*b*]diindole-7*b*,14*b*-diol, T-165
- 6,7,7*a*,8-Tetrahydro-4,10-dimethoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, B-471
- 6,7,7*a*,8-Tetrahydro-10,12-dimethoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, D-846
- 6,7,7*a*,8-Tetrahydro-4,10-dimethoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-12-ol, D-950
- 5,8,13,13*a*-Tetrahydro-9,10-dimethoxy-6*H*-benzo[*g*]-1,3-benzodioxolo[5,6-*a*]quinolizin-13-ol, O-94
- 5,8,13,13*a*-Tetrahydro-9,10-dimethoxy-6*H*-benzo[*g*]-1,3-benzodioxolo[5,6-*a*]quinolizin-13-ol, O-94
- 9,10,15*b*,16-Tetrahydro-5,15-dimethoxy-1*H*,7*H*-[1,3]benzodioxino[5,6-*g*][1,3]benzodioxolo[5,6-*a*]quinolinol, O-110
- 5,8,13,13*a*-Tetrahydro-2,10-dimethoxy-6*H*-dibenzo[*a,g*]quinolizine-3,9-diol, I-315
- 5,8,13,13*a*-Tetrahydro-3,9-dimethoxy-6*H*-dibenzo[*a,g*]quinolizine-2,10-diol, S-547
- 5,8,13,13*a*-Tetrahydro-2,11-dimethoxy-6*H*-dibenzo[*a,g*]quinolizin-9-ol, R-82
- 5,7,8,15-Tetrahydro-3,4-dimethoxy-6,15-dimethylbenzo[*e*][1,3]dioxolo[4,5-*k*][3]benzazecin-14(6*H*)-one, C-671
- 1,2,3,4-Tetrahydro-7,8-dimethoxy-1,2-dimethylisoquinoline, T-166
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethylisoquinoline, T-167
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2,2-dimethylisoquinolinium(1+), T-158
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethyl-5-isoquinolinol, T-174
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethyl-8-isoquinolinol, T-237
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1,2-dimethyl-6-isoquinolinol, T-237
- 3',4',6,8-Tetrahydro-6',7'-dimethoxy-2',6-dimethylspiro[7*H*-indeno[4,5-*d*]-1,3-dioxole-7,1'(2'*H*)-isoquinolin]-6-ol, R-8
- 5,6,12*b*,13-Tetrahydro-9,10-dimethoxy-8*H*-1,3-dioxolo[4,5-*h*]isoindolo[1,2-*b*][3]benzazepin-8-one, L-91
- 5,8,9,15*a*-Tetrahydro-3,4-dimethoxy[1,3]dioxolo[4,5-*h*]isoquinol[1,2-*b*][3]benzazepine-6,15-dione, P-801
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-isoquinolinemethanol, C-54
- 1,2,3,4-Tetrahydro-6,7-dimethoxyisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7,8-dimethoxyisoquinoline, T-159
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-8-isoquinolinol, T-236
- 1,2,3,4-Tetrahydro-7,8-dimethoxy-6-isoquinolinol, T-236
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-[(4-methoxyphenyl)methyl]-5-isoquinolinol, A-1069
- 6,7,7*a*,8-Tetrahydro-9,10-dimethoxy-7-methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, C-733
- 5,6,6*a*,7-Tetrahydro-1,2-dimethoxy-6-methyl-4*H*-benzo[*de*][1,3]benzodioxolo[5,6-*g*]quinoline, N-29
- 6,6*a*,11,14-Tetrahydro-8,9-dimethoxy-6-methyl-12*H*-benzo[*a*]-1,3-benzodioxolo[4,5-*g*]quinolizine, C-222
- 5,6,11,12-Tetrahydro-8,9-dimethoxy-14-methylbenzo[5,6]cycloocta[1,2-*f*]-1,3-benzodioxol-5,11-imine, E-220
- 4,6,7,13-Tetrahydro-9,10-dimethoxy-5-methylbenzo[*e*][1,3-dioxolo[4,5-*l*][2]benzazecin-12(5*H*)-one, C-790
- 5,7,8,15-Tetrahydro-2,3-dimethoxy-6-methylbenzo[*e*][1,3]dioxolo[4,5-*k*][3]benzazecin-14(6*H*)-one, F-3
- 5,7,8,15-Tetrahydro-3,4-dimethoxy-6-methylbenzo[*e*][1,3]dioxolo[4,5-*k*][3]benzazecin-14(6*H*)-one, F-2
- 2,3,12,12*a*-Tetrahydro-5,10-dimethoxy-1-methyl-1*H*-[1]benzoxepino[2,3,4-*ij*]isoquinolin-8-ol, F-66
- 2,3,12,12*a*-Tetrahydro-6,9-dimethoxy-1-methyl-1*H*-[1]benzoxepino[2,3,4-*ij*]isoquinolin-8-ol, S-73
- 4*a*,5,6,7-Tetrahydro-8,13-dimethoxy-5-methyl-4*H*-bis[1,3]benzodioxolo[6,5,4-*de*:5',4'-*g*]quinoline, H-243
- 5,6,7,8-Tetrahydro-4,8-dimethoxy-8-(3-methyl-2-butenyl)furo[2,3-*b*]quinolin-7-ol, H-57
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methyl- β -carboline, D-620
- 5,6,11,12-Tetrahydro-2,9-dimethoxy-13-methyl-dibenzo[*a,e*]cycloocten-5,11-imin-3-ol, H-464
- 5,6,6*a*,7-Tetrahydro-1,10-dimethoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline-2,9-diol, B-243
- 5,6,6*a*,7-Tetrahydro-2,9-dimethoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,10-diol, B-276
- 5,6,6*a*,7-Tetrahydro-1,9-dimethoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline-2,10-diol, L-195
- 5,6,6*a*,7-Tetrahydro-9,10-dimethoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2-diol, L-52
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methylisoquinoline, T-167
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methylisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7,8-dimethoxy-2-methylisoquinoline, T-159
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-1-methyl-8-isoquinolinol, T-237
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-8-isoquinolinol, T-236
- 1,2,3,4-Tetrahydro-7,8-dimethoxy-1-methyl-6-isoquinolinol, T-237
- 1,2,3,4-Tetrahydro-7,8-dimethoxy-2-methyl-6-isoquinolinol, T-236
- 6-(1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, C-653
- 4-[(1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)methyl]phenol, A-1444
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-[2-(3,4-methylenedioxyphenyl)ethyl]isoquinoline, D-978
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-(3,4-methylenedioxyphenyl)isoquinoline, C-796
- 1,2,3,4-Tetrahydro-9,11-dimethoxy-2-methylnaphth[2,1-*f*]isoquinoline-8,12-diol, L-211
- 5,6,7,8-Tetrahydro-3,7-dimethoxy-2-methyl-5-octyl-4(1*H*)-quinolinone, C-343
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-phenylisoquinoline, T-168
- 3',4',6,8-Tetrahydro-6',7'-dimethoxy-2'-methylspiro[7*H*-indeno[4,5-*d*]-1,3-dioxole-7,1'(2'*H*)-isoquinoline]-6,8-diol, R-9

- 3',4',6,8-Tetrahydro-6',7'-dimethoxy-2'-methylspiro[7*H*-indeno[4,5-*d*]-1,3-dioxole-7,1'(2'*H*)-isoquinolin]-8-ol, F-174
- 1,2,3,4-Tetrahydro-6,7-dimethoxy-2-methyl-1-(3,4,5-trimethoxyphenyl)isoquinoline, C-797
- 4,6,7,14-Tetrahydro-5,14-dimethylbis[1,3]benzodioxolo[4,5-*c*:5'-6'-*g*]azecin-13(5*H*)-one, C-670
- 1,2,3,4-Tetrahydro-1,2-dimethyl-β-carboline, T-202
- 6,7,8,9-Tetrahydro-3,10-dimethylcyclohepta[*b*]pyran-5,8-imin-4(5*H*)-one, B-57
- 6,7,8,9-Tetrahydro-2,10-dimethylcyclohepta[*b*]pyran-5,8-imin-4(5*H*)-one, I-194
- 2,3,6,7-Tetrahydro-7,9-dimethyl-2,6-dioxo-1*H*-purinium hydroxide inner salt, X-5
- 1,2,3,6-Tetrahydro-1,3-dimethyl-2,6-dioxo-4-pyrimidinecarboxylic acid, D-669
- 1,5,6,7-Tetrahydro-1,6-dimethyl-5,7-dioxopyrimido[5,4-*e*]-*as*-triazine, R-66
- 3,3*a*,8,8*a*-Tetrahydro-3*a*,8-dimethyl-2*H*-furo[2,3-*b*]indol-5-ol methyl carbamate (ester), P-401
- 1,2,3,4-Tetrahydro-1,1-dimethyl-6,7-isoquinoline-diol, T-152
- 1,2,3,4-Tetrahydro-1,3-dimethyl-8-isoquinolinol, T-177
- 2,3,4,5-Tetrahydro-1,3-dimethyl-1,4-methano-1*H*-benzazepin-8-ol, A-1333
- 6,7,8,9-Tetrahydro-2,5-dimethyl-8-(1-methylethenyl)-5*H*-cyclohepta[*b*]pyridine, G-190
- 3,5,7,9-Tetrahydro-3,9-dimethyl-6-(methylimino)-8*H*-purin-8-one, C-19
- Tetrahydro-3,5-dimethyl-6-[octahydro-3-(tetrahydro-4-methyl-5-oxo-2-furanyl)-1*H*-pyrrolo[1,2-*a*]azepin-9-yl]furo[3,2-*b*]furan-2(3*H*)-one, P-105
- 1,2,3,6-Tetrahydro- α^6 ,1-dimethyl- α^2 -phenyl-2,6-pyridinediethanol, S-214
- 1,3,4,9-Tetrahydro-1,1-dimethylpyrano[3,4-*b*]indole, T-169
- 2,3,4,10-Tetrahydro-2,2-dimethyl-5*H*-pyrano[2,3-*b*]quinolin-5-one, H-56
- 2,3,4,6-Tetrahydro-1,5-dimethyl-1*H*-pyrido[4,3-*b*]carbazole, G-211
- 2,3,4,9-Tetrahydro-1,2-dimethyl-1*H*-pyrido[3,4-*b*]indole, T-202
- 1,3,4,5-Tetrahydro-1,5-dimethylpyrrolo[4,3,2-*de*]quinoline-7,8-dione, D-13
- 5,6,7,8-Tetrahydro-2,4-dimethylquinoline, T-170
- 5,6,7,8-Tetrahydro-2,5-dimethylquinoline, T-171
- 1,2,3,8-Tetrahydro-1,8-dimethyl-1,7*b*,10-triazabenz[5,6]cyclohepta[1,2,3-*k*]fluorene, D-138
- 1,2,3,6-Tetrahydro-2,6-dioxo-4-pyrimidinecarboxylic acid, D-669
- 4,6,7,9-Tetrahydro-1*H*,3*H*-dipyrano[3,4-*b*:4':3'-*e*]pyrazine-3,7-dimethanol, I-261
- 1,2,3,4-Tetrahydroellipticine, E-67
- Tetrahydroepiberberine, S-307
- 1,2,10,10*a*-Tetrahydro-2-ethylidene-10-hydroxydipyrrolo[1,2-*c*:2',1'-*f*]pyrimidin-5-one, P-146
- 5,6,7,8-Tetrahydrofolic acid, T-172
- Tetrahydroglaziovine, P-643
- Tetrahydroalichondramide, H-15
- Tetrahydroharman *N*⁶-oxide, T-202
- Tetrahydroharman, T-202
- Tetrahydroharmine, T-202
- Tetrahydroharmol, H-592
- Tetrahydrohistrionicotoxin, H-309
- 1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-6,7-dimethoxyisoquinoline, A-1444
- 1,2,3,4-Tetrahydro-1-(3-hydroxybenzyl)-6,7-dimethoxy-2-methylisoquinoline, A-1038
- 1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-7,8-dimethoxy-1-methylisoquinoline, L-251
- 1,2,3,4-Tetrahydro-1-(4-hydroxybenzyl)-6,7-methylenedioxyisoquinoline, C-453
- 1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxyhexyl)pyridine, A-1487
- 1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxypentyl)-1-methylpyridine, H-433
- 1,2,5,6-Tetrahydro-2-(2-hydroxybutyl)-6-(2-hydroxypentyl)pyridine, H-433
- 1,2,5,6-Tetrahydro-6-(2-hydroxybutyl)-1-methyl-2-(2-oxobutyl)pyridine, D-391
- 1-[1,2,5,6-Tetrahydro-6-(2-hydroxybutyl)-1-methyl-2-pyridinyl]-2-butanone, D-391
- 1,2,4,5-Tetrahydro-9-hydroxy-6-canthinone, H-444
- 1,2,3,4-Tetrahydro-7-hydroxy-β-carboline-3-carboxylic acid, T-224
- 1,2,3,4-Tetrahydro-6-hydroxy-β-carboline, T-173
- 5,6,7,8-Tetrahydro-7-hydroxy-4,8-dimethoxy- α,α -dimethylfuro[2,3-*b*]quinoline-8-propanol, H-57
- 1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinoline, T-174
- 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-1,2-dimethylisoquinoline, T-237
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1,2-dimethylisoquinoline, T-237
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-2,2-dimethylisoquinolinium(1+), T-236
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-isoquinolinecarboxylic acid, T-175
- 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxyisoquinoline, T-236
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxyisoquinoline, T-236
- 1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-1-(4-methoxybenzyl)isoquinoline, A-1069
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-methyl-1-isoquinolinecarboxylic acid, P-299
- 1,2,3,4-Tetrahydro-5-hydroxy-6,7-dimethoxy-2-methyl-1-isoquinolinemethanol, T-178
- 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-1-methylisoquinoline, T-237
- 1,2,3,4-Tetrahydro-6-hydroxy-7,8-dimethoxy-2-methylisoquinoline, T-236
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1-methylisoquinoline, T-237
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-2-methylisoquinoline, T-236
- 6,7,9,10-Tetrahydro-5-(4-hydroxy-3,5-dimethoxyphenyl)-5*H*-benzo[*g*][1,3]dioxolo[4',5':4,5]benzo[*de*]pyrido[3,2,1-*ij*]quinolin-7-one, T-176
- 1,2,3,4-Tetrahydro-8-hydroxy-6,7-dimethoxy-1,2,2-trimethylisoquinolinium(1+), T-237
- 1,2,3,4-Tetrahydro-6-hydroxy-1,2-dimethyl-β-carboline, S-285
- 1,2,3,4-Tetrahydro-8-hydroxy-1,3-dimethylisoquinoline, T-177
- 1,2,3,4-Tetrahydro-7-hydroxy-2-(4-hydroxybenzyl)-6-methoxyisoquinoline, S-234
- 1,2,3,4-Tetrahydro-7-hydroxy-1-(3-hydroxy-4,5-dimethoxyphenethyl)-6-methoxy-2-methylisoquinoline, A-1558
- 1,2,3,4-Tetrahydro-5-hydroxy-1-hydroxymethyl-6,7-dimethoxy-2-methylisoquinoline, T-178
- 1,2,3,4-Tetrahydro-7-hydroxy-1-hydroxymethyl-6-methoxy-2-methylisoquinoline, C-54
- 1,2,3,4-Tetrahydro-7-[5-hydroxy-2-(hydroxymethyl)-4-methoxy-1-naphthalenyl]-1,3-dimethyl-8-isoquinolinol, D-793
- 1,2,3,4-Tetrahydro-5-hydroxy-4-(4-hydroxyphenyl)-6-methoxy-2-methylisoquinoline, L-54
- 1,2,3,4-Tetrahydro-8-hydroxy-1-(4-hydroxyphenyl)methyl]-7-methoxy-2,2-dimethylisoquinolinium(1+), J-63
- 6,7,8,9-Tetrahydro-1-hydroxy-11-(4-hydroxyphenyl)-3-methylpyridazino[1,2-*a*]indazol-5-ium inner salt, N-193
- 5,6,7,7*a*-Tetrahydro-2-hydroxy-1-(4-hydroxyphenyl)-3*H*-pyrrolizin-3-one, P-333
- 4,5,6,7-Tetrahydro-6-hydroxyimidazo[4,5-*e*][1,4]diazepin-8(1*H*)-one, A-1605
- 13,17*b*,18,18*a*-Tetrahydro-17*b*-hydroxy-11*H*-indolo[3',2':4,5]pyrrolo[2,1-*c*]quinazolinol[3,2-*a*][1,4]benzodiazepine-5,11(12*aH*)-dione, A-1481
- 1,2,3,4-Tetrahydro-7-hydroxy-1-isobutyl-6-methoxy-2-methylisoquinoline, L-260
- 1,2,3,4-Tetrahydro-6-hydroxyisoquinoline, T-196
- 1,2,3,4-Tetrahydro-7-hydroxyisoquinoline, T-197
- 1,2,3,4-Tetrahydro-8-hydroxyisoquinoline, T-198
- 1,2,3,4-Tetrahydro-8-hydroxymanzamine A, M-91
- 2,3,4,5-Tetrahydro-8-hydroxy-1,4-methanopyrido[1,2-*a*][1,4]diazepin-7(1*H*)-one, H-644
- 5,8,13,13*a*-Tetrahydro-2-hydroxy-3-methoxy-6*H*-dibenzo[*a,g*]quinolinize, B-111
- 1,2,3,4-Tetrahydro-6-hydroxy-6-methoxy-1,2-dimethylisoquinoline, T-181
- 1,2,3,4-Tetrahydro-9-hydroxy-8-methoxy-2,4-dimethyl-6*H*-naphtho[1',2':4,5]pyrano[3,2-*h*]isoquinolin-6-one, D-790
- 1,2,3,4-Tetrahydro-6-hydroxy-7-methoxyisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxyisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-(4-methoxybenzyl)isoquinoline, S-234
- 1,2,3,4-Tetrahydro-5-hydroxy-8-methoxy-1-(4-methoxybenzyl)-2-methylisoquinoline, T-155
- 5,6,13,13*a*-Tetrahydro-2-hydroxy-3-methoxy-7-methyl[1,3]dioxolo[5,6]indolo[2,1-*a*]isoquinolinium(1+), C-798
- 1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-6-methyl-1,3-isoquinolinedimethanol, A-1209
- 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-methyl-1-isoquinolinemethanol, C-54
- 1,2,3,4-Tetrahydro-6-hydroxy-7-methoxy-1-methylisoquinoline, T-179
- 1,2,3,4-Tetrahydro-6-hydroxy-7-methoxy-2-methylisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7-hydroxy-5-methoxy-2-methylisoquinoline, T-180
- 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-1-methylisoquinoline, T-181
- 1,2,3,4-Tetrahydro-7-hydroxy-6-methoxy-2-methylisoquinoline, T-158
- 1,2,3,4-Tetrahydro-7-hydroxy-8-methoxy-2-methylisoquinoline, T-159
- 1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-1-methylisoquinoline, T-163
- 1,2,3,4-Tetrahydro-8-hydroxy-7-methoxy-2-methylisoquinoline, T-159
- 1,2,3,4-Tetrahydro-7-(1-hydroxy-8-methoxy-3-methyl-2-naphthalenyl)-6,8-dimethoxy-1,2,3-trimethylisoquinoline, A-992
- 5,6,7,8-Tetrahydro-8-hydroxy-3-methoxy-2-methyl-5-octyl-4(1*H*)-quinolinone, A-1325
- 1',2',3',4'-Tetrahydro-6-hydroxy-10-methoxy-*N*²-methylusambarensine, U-52
- 1,2,3,11*a*-Tetrahydro-8-hydroxy-7-methoxy-5*H*-pyrrolo[2,1-*c*]benzodiazepin-5-one, A-1156
- 1,2,3,4-Tetrahydro-8-hydroxy-6-methoxy-1,2,3-trimethylisoquinoline, T-182
- 1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-β-carboline-3-carboxylic acid, T-206
- 1,2,3,4-Tetrahydro-7-hydroxy-1-methyl-β-carboline-3-carboxylic acid, T-206
- 1,2,3,4-Tetrahydro-7-hydroxy-1-methyl-β-carboline, H-592
- 1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-β-carboline, S-285
- 1,2,3,4-Tetrahydro-6-hydroxy-2-methyl-β-carboline, T-173
- 5,8,13,13*a*-Tetrahydro-8-(hydroxymethyl)-2,10-dimethoxy-6*H*-dibenzo[*a,g*]quinolinize-1,11-diol, M-57
- 1,2,3,4-Tetrahydro-1-hydroxymethyl-6,7-dimethoxyisoquinoline, C-54
- 3,4,5,6-Tetrahydro-6-hydroxymethyl-3,6-dimethyl-4-pyrimidinecarboxylic acid, T-183
- 2',3',4',5'-Tetrahydro-5-hydroxymethyl-2',5'-dioxo-1,3'-bi-1*H*-pyrrole-2-carboxaldehyde, F-139
- 1,2,3,4-Tetrahydro-1-(2-hydroxy-3,4-methylene-dioxybenzyl)-6,7-methylenedioxyisoquinoline, L-80
- 1,2,3,4-Tetrahydro-6-hydroxy-2-methylharman, S-285
- 1,2,3,4-Tetrahydro-6-hydroxy-1-methyl-3-isoquinolinecarboxylic acid, T-184
- 1,2,3,4-Tetrahydro-8-hydroxy-2-methylisoquinoline, T-198
- 1,2,3,4-Tetrahydro-6-hydroxy-2-methylisoquinoline, T-196
- 6,7,8,9-Tetrahydro-2-(hydroxymethyl)-3-methoxy-5-octyl-4(1*H*)-quinolinone, A-1325
- 1,2,3,4-Tetrahydro-1-(2-hydroxymethyl-3,4-methylenedioxybenzyl)-2-methyl-6,7-methylenedioxyisoquinoline, C-676
- Tetrahydro-10*b*-hydroxy-5-(1-methylpropyl)-8*H*-oxazolo[3,2-*a*]pyrrolo[2,1-*c*]pyrazine-3,6(2*H*,5*H*)-dione, E-170

- 6,7,8,9-Tetrahydro-1-hydroxy-3-methylpyridazino[1,2-*a*]indazol-5-ium inner salt, N-191
- 1,4,5,6-Tetrahydro-5-hydroxy-2-methyl-4(6)-pyrimidinecarboxylic acid, T-185
- 2,3,5,7a-Tetrahydro-7-(hydroxymethyl)-1*H*-pyrrolizine-1,2-diol, C-775
- 2,7,19,20-Tetrahydro-3-hydroxy-1-methylvobasanol-17-oic acid δ -lactone, T-186
- 1,2,14,19-Tetrahydro-12-hydroxy-20-norcrotalanan-11,15-dione, R-65
- 1,2,3,4-Tetrahydro-6-hydroxy-1-oxo- β -carboline, T-226
- 4,4',5',5''-Tetrahydro-2-(2-hydroxyphenyl)-4,4'-bithiazole, U-7
- 1,2,3,4-Tetrahydro-4-[[[2-(4-hydroxyphenyl)ethyl]imino]ethylidene]-2,6-pyridinedicarboxylic acid, M-647
- 1,2,3,4-Tetrahydro-1-[2-(4-hydroxyphenyl)ethyl]-7-methoxy-2-methyl-6-isoquinolinol, C-570
- 1-[1,2,5,6-Tetrahydro-6-(2-hydroxy-2-phenylethyl)-1-methyl-2-pyridinyl]-2-propanone, S-214
- 1,2,3,4-Tetrahydro-4-(4-hydroxyphenyl)-6-methoxy-2-methyl-7-isoquinolinol, C-369
- 1,2,3,4-Tetrahydro-4-(4-hydroxyphenyl)-6-methoxy-2-methyl-5-isoquinolinol, L-54
- 3,4,5,6-Tetrahydro-3-[(4-hydroxyphenyl)methylene]-2-pyridinecarboxylic acid, A-1079
- 1,3,4,5-Tetrahydro-5-(4-hydroxyphenyl)-1-methylfuro[3,4-*b*]pyridin-7(2*H*)-one, A-1079
- 1,2,3,4-Tetrahydro-1-[(4-hydroxyphenyl)methyl]-6,7-isoquinolinediol, H-275
- 1,2,3,4-Tetrahydro-2-[(4-hydroxyphenyl)methyl]-6-methoxy-7-isoquinolinol, S-234
- 1,2,3,4-Tetrahydro-1-[(4-hydroxyphenyl)methyl]-2-methyl-5,8-isoquinolinediol, T-155
- 5,6,7,7a-Tetrahydro-2-hydroxy-1-phenyl-3*H*-pyrrolizin-3-one, P-333
- 1,2,3,6-Tetrahydro-3-hydroxypicolinic acid, T-187
- 1,2,3,4-Tetrahydro-6-hydroxy-1-propyl- β -carboline, T-219
- 1,2,3,6-Tetrahydro-3-hydroxy-2-pyridinecarboxylic acid, T-187
- 2,3,4,9-Tetrahydro-7-hydroxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 2,3,4,9-Tetrahydro-6-hydroxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 2,3,5,7a-Tetrahydro-1-hydroxy-1*H*-pyrrolizine-7-methanol, T-188
- 5,6,7,7a-Tetrahydro-7a-hydroxy-3*H*-pyrrolizin-3-one, T-189
- 1-(1,2,3,9-Tetrahydro-3-hydroxypyrrrolo[2,1-*b*]quinazolin-9-yl)-2-propanone, P-164
- 1,2,3,9-Tetrahydro-3-hydroxypyrrrolo[2,1-*b*]quinoline, P-165
- 1,2,3,4-Tetrahydro-4-hydroxy-4-quinolinecarboxylic acid, T-190
- Tetrahydro-4-hydroxy-*N,N,N,5*-tetramethyl-2-furanmethanaminium(1+), M-773
- 5,6,12,12a-Tetrahydro-2-hydroxy-3,9,10-trimethoxy-7-methylindolo[2,1-*a*]isoquinolinium(1+), C-783
- 1,2,3,4-Tetrahydro-*ar*-hydroxy-*ar*-trimethoxy-2-methylisoquinoline, T-191
- 2,3,4,6-Tetrahydro-3-hydroxy-2,2,6-trimethyl-5*H*-pyrano[3,2-*c*]quinolin-5-one, F-90
- 2,3,4,6-Tetrahydro-4-hydroxy-2,2,6-trimethyl-5*H*-pyrano[3,2-*c*]quinolin-5-one, F-90
- 2,3,4,10-Tetrahydro-3-hydroxy-2,2,10-trimethyl-5*H*-pyrano[2,3-*b*]quinolin-5-one, R-92
- 1,6,7,12-Tetrahydroimidazo[4,5-*g*]indolo[2,3-*a*]quinolizin-5-ium(1+), V-102
- 2,3,4,9-Tetrahydro-1-(1*H*-imidazol-4-ylmethyl)-1*H*-pyrido[3,4-*b*]indole, L-199
- 4,5,6,7-Tetrahydro-1*H*-imidazo[4,5-*c*]pyridine-6-carboxylic acid, T-193
- 4,5,6,7-Tetrahydro-1*H*-imidazo[4,5-*c*]pyridine, T-192
- 5,6,7,8-Tetrahydroimidazo[1,5-*c*]pyrimidin-5-one, D-462
- 3,6,7,9-Tetrahydro-6-imino-3,9-dimethyl-8*H*-purin-8-one, A-819
- 1,2,3,7-Tetrahydro-2-imino-1,3-dimethyl-6*H*-purin-6-one, D-748
- 1,6,7,9-Tetrahydro-6-imino-1,9-dimethyl-8*H*-purin-8-one, T-194
- 1,2,3,7-Tetrahydro-2-imino-1,3,7-trimethyl-6*H*-purin-6-one, D-748
- 1,3,6,7-Tetrahydro-6-imino-1,3,7-trimethyl-2*H*-purin-2-one, T-608
- 4,5,6,7-Tetrahydro-1*H*-indol-2(3*H*)-one, T-195
- 5,6,7,7a-Tetrahydro-1*H*-indol-2(3*H*)-one, T-195
- 8,13,13*b*,14-Tetrahydroindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, D-508
- 6,7,12,13-Tetrahydro-5*H*-indolo[2,3-*a*]pyrrolo[3,4-*c*]carbazol-5-one, A-1190
- 2,3,5,7-Tetrahydro-9(1*H*-indol-3-yl)-6*H*-dipyrrolo[4,3,2-*de*:2',3'-*h*]quinolin-6-one, W-3
- 1,2,3,4-Tetrahydro-1-isobutyl-2-methyl- β -carboline, M-521
- 1,2,3,4-Tetrahydro-1-isobutyl- β -carboline, M-521
- Tetrahydroisokomarovine, Q-53
- 5,6,7,8-Tetrahydro-3-isopropylimidazo[1,2-*a*]pyridine, C-200
- 1,2,3,4-Tetrahydro-6,7-isoquinolinediol, T-158
- 1,2,3,4-Tetrahydro-7,8-isoquinolinediol, T-159
- 1,2,3,4-Tetrahydro-5,6,7-isoquinolinetriol, T-236
- 1,2,3,4-Tetrahydro-6-isoquinolinol, T-196
- 1,2,3,4-Tetrahydro-8-isoquinolinol, T-198
- 1,2,3,4-Tetrahydro-7-isoquinolinol, T-197
- Tetrahydrojatrorrhizine, C-694
- Tetrahydrokomarovine, Q-54
- Tetrahydrolythyrine, T-199
- Tetrahydroleontidine, C-61
- 5,6,7,8-Tetrahydrolepidine, T-208
- 1,2,3,4-Tetrahydromanzamine B, M-92
- 10,11,12,12a-Tetrahydro-9*H*-6,12*b*-methanofuro[2,3-*c*]pyrido[1,2-*b*][1,2]oxazocin-2(6*H*)-one, P-387
- 9,10,11,11a-Tetrahydro-8*H*-6,11*b*-methanofuro[2,3-*c*]pyrido[1,2-*a*]pyridoozepin-2(6*H*)-one, S-206
- 6,7,7a,8-Tetrahydro-10-methoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-12-ol, D-846
- 6,7,7a,8-Tetrahydro-4-methoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-12-ol, E-70
- 6a,7,8,9-Tetrahydro-12-methoxy-6*H*-benzo[*de*]-1,3-benzodioxolo[4,5-*g*]quinolin-11-ol, L-10
- 6,7,7a,8-Tetrahydro-12-methoxy-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-11-ol, N-25
- 5,8,14,14a-Tetrahydro-3-methoxy-6*H*-benzo[*a*][1,3]benzodioxolo[5,6-*g*]quinolizin-2-ol, C-685
- 1,2,3,4-Tetrahydro-2-(4-methoxybenzyl)-6,7-methylenedioxyisoquinoline, V-101
- 4,6,7,13-Tetrahydro-12-methoxy-12*H*-bis[1,3]benzodioxolo[5,6-*a*:4',5'-*g*]quinolizin-12*b*-methanol, Z-19
- 2,3,5,6-Tetrahydro-1-methoxy-2,6-bis(1-methylethenyl)-1*H*-imidazo[1,2-*a*]imidazole, I-187
- 1,2,3,4-Tetrahydro-6-methoxy- β -carboline, T-173
- 5,8,13,13a-Tetrahydro-3-methoxy-6*H*-dibenzo[*a,g*]quinolizin-2-ol, B-111
- 4,5,5a,11-Tetrahydro-1-methoxy-5,5-dimethyl-3*H*,6*H*-[1,3]benzodioxolo[4',5':6':4,5][2]benzopyrano[1,8,7-*hij*]isoquinolinium(1+), T-345
- 1,2,4,5-Tetrahydro-8-methoxy-4,4-dimethylbenzo[*b*]pyrido[2,3,4-*gh*]pyrrolizine, H-64
- 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl- β -carboline, S-285
- 1,2,3,4-Tetrahydro-6-methoxy-2,9-dimethyl- β -carboline, T-173
- 6,7,8,9-Tetrahydro-4-methoxy-8,9-dimethyl-1,3-dioxolo[4,5-*h*]isoquinoline, A-1021
- 3,9,10,15-Tetrahydro-5-methoxy-3,3-dimethyl-7*H*-indolo[2',3':3,4]pyrido[2,1-*b*]pyrano[2,3-*h*]quinazolin-7-one, P-89
- 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl-7-isoquinolinol, T-181
- 1,2,3,4-Tetrahydro-6-methoxy-1,2-dimethyl-7,8-methylenedioxyisoquinoline, A-1021
- 1,2,3,7-Tetrahydro-1-methoxy-7,7-dimethyl-2-(1-methylethenyl)imidazo[1,2-*a*]pyrimidine, A-251
- 2,3,8,8a-Tetrahydro-3a-methoxy-1,8-dimethylpyrrolo[2,3-*b*]indole, A-626
- 1,2,3,4-Tetrahydro-9-methoxyellipticine, M-252
- 1,2,3,4-Tetrahydro-6-methoxyisoquinoline, T-196
- 1,2,3,4-Tetrahydro-7-methoxyisoquinoline, T-197
- 1,2,3,4-Tetrahydro-8-methoxyisoquinoline, T-198
- 1,2,3,4-Tetrahydro-6-methoxy-7-isoquinolinol, T-158
- 1,2,3,4-Tetrahydro-7-methoxy-6-isoquinolinol, T-158
- 1,2,3,4-Tetrahydro-6-methoxy-2-[(4-methoxyphenyl)methyl]-7-isoquinolinol, S-234
- 6,7,7a,8-Tetrahydro-11-methoxy-7-methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-12-ol, B-397
- 5,6,11,12-Tetrahydro-9-methoxy-14-methylbenzo[5,6]cycloocta[1,2-*f*]-1,3-benzodioxol-5,11-imin-10-ol, N-104
- 1,2,3,11-Tetrahydro-1-(1-methoxy-3-methyl-9*H*-carbazol-9-yl)-3,3,5-trimethylpyrano[3,2-*a*]carbazol-2-ol, M-754
- 1,2,3,4-Tetrahydro-6-methoxy-1-methyl- β -carboline, S-285
- 1,2,3,4-Tetrahydro-6-methoxy-2-methyl- β -carboline, T-173
- 1,2,3,4-Tetrahydro-7-methoxy-1-methyl- β -carboline, T-202
- 5,6,6a,7-Tetrahydro-1-methoxy-6-methyl-4*H*-dibenzo[*de,g*]quinoline-2,7-diol, P-4
- 6,7,8,9-Tetrahydro-4-methoxy-9-methyl-1,3-dioxolo[4,5-*h*]isoquinoline, A-1021
- 5,6,7,8-Tetrahydro-4-methoxy-6-methyl-1,3-dioxolo[4,5-*g*]isoquinoline, H-396
- 1,2,3,4-Tetrahydro-*ar*-methoxy-1-methylisoquinoline, T-200
- 1,2,3,4-Tetrahydro-5-methoxy-2-methyl-7-isoquinolinol, T-180
- 1,2,3,4-Tetrahydro-6-methoxy-1-methyl-7-isoquinolinol, T-181
- 1,2,3,4-Tetrahydro-7-methoxy-1-methyl-6-isoquinolinol, T-179
- 1,2,3,4-Tetrahydro-6-methoxy-2-methyl-7-isoquinolinol, T-158
- 1,2,3,4-Tetrahydro-7-methoxy-1-methyl-8-isoquinolinol, T-163
- 1,2,3,4-Tetrahydro-7-methoxy-2-methyl-6-isoquinolinol, T-158
- 1,2,3,4-Tetrahydro-6-methoxy-1-methyl-7,8-methylenedioxyisoquinoline, A-1021
- 1,2,3,4-Tetrahydro-8-methoxy-2-methyl-6,7-methylenedioxyisoquinoline, H-396
- 3',4',6,8-Tetrahydro-7'-methoxy-2'-methyl-6-methylenespiro[7*H*-indeno[4,5-*d*]-1,3-dioxole-7,1'(2'*H*)-isoquinolin]-6'-ol, O-16
- 1,2,3,4-Tetrahydro-6-methoxy-2-methyl-1-(2-methylpropyl)-7-isoquinolinol, L-260
- 6,7,8,9-Tetrahydro-3-methoxy-2-methyl-5-octyl-4(1*H*)-quinolinone, A-1325
- 2,3,4,9-Tetrahydro-6-methoxy-2-methyl-1*H*-pyrido[3,4-*b*]indole, T-173
- 2,3,4,9-Tetrahydro-5-methoxy-1-methyl-1*H*-pyrido[3,4-*b*]indole, T-202
- 1,2,3,4-Tetrahydro-4-methoxy-2-methyl-8-quinolinecarboxylic acid, T-201
- 1,2,3,4-Tetrahydro-5-methoxy-1-oxo- β -carboline, T-226
- 1,2,3,4-Tetrahydro-6-methoxy-1-oxo- β -carboline, T-226
- 1,2,3,4-Tetrahydro-7-methoxy-1-oxo- β -carboline, T-226
- 1,2,3,4-Tetrahydro-8-methoxy-1-oxo- β -carboline, T-226
- 8,9,10,11-Tetrahydro-6-methoxy-1,2,3,4,5-pentathiepio[6,7-*f*]isoquinolin-7-ol, L-206
- 1,2,3,4-Tetrahydro-7-methoxy-9*H*-pyrido[3,4-*b*]indole-9-carboxaldehyde, P-916
- 2,3,4,9-Tetrahydro-6-methoxy-1*H*-pyrido[3,4-*b*]indole, T-173
- 2,3,4,9-Tetrahydro-6-methoxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 2,3,4,9-Tetrahydro-8-methoxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 2,3,4,9-Tetrahydro-6-methoxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 2,3,4,9-Tetrahydro-7-methoxy-1*H*-pyrido[3,4-*b*]indol-1-one, T-226

- 1,2,3,9-Tetrahydro-5-methoxypyrrolo[2,1-*b*]quinazolin-3-ol, P-165
- 4,5,6,7-Tetrahydro-3-[[3-methoxy-5-(1*H*-pyrrol-2-yl)-2*H*-pyrrol-2-ylidene]methyl]-1,4-dimethyl-2*H*-isoindole, C-897
- 6,7,8,9-Tetrahydro-4-methoxy-8,8,9-trimethyl-1,3-dioxolo[4,5-*h*]isoquinolinium(1+), A-1021
- 1,2,3,4-Tetrahydro-8-methoxy-1,2,3-trimethylisoquinoline, T-177
- 1,2,3,4-Tetrahydro-6-methoxy-1,2,3-trimethyl-8-isoquinolinol, T-182
- 2,3,4,7-Tetrahydro-5-methoxy-2,2,7-trimethylpyran[2,3-*f*][2,3]benzoxazin-3-ol, E-293
- 3,4,6,6-Tetrahydro-5-methyl-1*H*-azepino[5,4,3-*cd*]indol-7-ol *N*²-oxide, F-9
- 6,7,7*a*,8-Tetrahydro-7-methyl-5*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, R-120
- 6-(5,6,7,8-Tetrahydro-6-methyl-1,3-benzodioxolo[4,5-*g*]isoquinolin-5-yl)furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, B-119
- 5,6,7,7*a*-Tetrahydro-7-methylbenzo[*d*]-1,3-dioxolo[4,5-*g*]pyrido[4,3,2-*jk*][2]benzazepin-11-ol, S-411
- 4,6,7,14-Tetrahydro-5-methylbis[1,3]benzodioxolo[4,5-*c*:5',6'-*g*]azecin-13(5*H*)-one, P-667
- 5,7,8,15-Tetrahydro-6-methylbis[1,3]benzodioxolo[5,6-*c*:5',6'-*g*]azecin-14(6*H*)-one, P-723
- 6,7,12*b*,13-Tetrahydro-13-methyl-4*H*-bis[1,3]benzodioxolo[5,6-*a*:4',5'-*g*]quinolizine, T-148
- 1,2,3,4-Tetrahydro-1-methyl-β-carboline-3-carboxylic acid, T-206
- 1,2,3,4-Tetrahydro-1-methyl-β-carboline, T-202
- 1,2,3,4-Tetrahydro-2-methyl-β-carboline, T-203
- 6,7,8,9-Tetrahydro-10-methylcyclohepta[*b*]pyran-5,8-imin-4(5*H*)-one, S-587
- 5,6,12,13-Tetrahydro-15-methylcycloocta[1,2-*f*:5,6-*f'*]bis[1,3]benzodioxol-5,12-imine, E-221
- 5,6,7,8-Tetrahydro-6-methyldibenz[*c,e*]azocine-1,2-diol, A-1361
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,3,10,11-pentol, P-209
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,3,9,10-pentol, P-208
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,9,10-tetrol, T-246
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,10-triol, T-524
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,11-triol, T-525
- 5,6,6*a*,7-Tetrahydro-6-methyl-4*H*-dibenzo[*de,g*]quinoline-1,2,3-triol, T-521
- 2,3,13,13*a*-Tetrahydro-1-methyl-1*H*-[1,3]dioxolo[7,8][1]benzoxepino[2,3,4-*i*]isoquinolin-6-ol, C-804
- 5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinoline, H-398
- 6-(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)furo[3,4-*e*]-1,3-benzodioxol-8(6*H*)-one, A-145
- 8-(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)furo[3,4-*e*]-1,3-benzodioxol-6(8*H*)-one, D-135
- 5-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)methyl]-1,3-benzodioxole-4-methanol, C-676
- 5-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)methyl]-1,3-benzodioxol-4-ol, L-80
- 4-[(5,6,7,8-Tetrahydro-6-methyl-1,3-dioxolo[4,5-*g*]isoquinolin-5-yl)methyl]phenol, C-453
- 1,2,3,6-Tetrahydro-1-methyl-2,6-dioxo-4-pyrimidin-ecarboxylic acid, D-669
- 1,2,3,6-Tetrahydro-3-methyl-2,6-dioxo-4-pyrimidin-ecarboxylic acid, D-669
- N*-(5*a*,6*a*,7,8-Tetrahydro-4-methyl-1,5-dioxo-1*H*,5*H*-pyrrolo[1,2-*c*][1,3]oxazepin-3-yl)hexanamide, C-852
- Tetrahydro-2-methylleptidine, E-67
- 1,2,3,4-Tetrahydro-2-(3,4-methylenedioxyphenethyl)-1-methylquinoline, G-9
- 1,2,3,4-Tetrahydro-2-*N*-methyl-8-hydroxymanzamine A, M-91
- 1,2,8,13-Tetrahydro-2-methyl-4*H*-indolo[2,3-*a*]pyrano[3,4-*g*]quinolizine-4,5(7*H*)-dione, C-66
- 8,13,13*b*,14-Tetrahydro-14-methylindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(7*H*)-one, E-310
- 4,6,7,8-Tetrahydro-7-methylindolo[4,3-*fg*]quinoline-9-methanol, A-205
- 1,2,3,4-Tetrahydro-1-methyl-6,7-isoquinolinediol, T-162
- 1,2,3,4-Tetrahydro-1-methyl-7,8-isoquinolinediol, T-163
- 1,2,3,4-Tetrahydro-2-methyl-4,8-isoquinolinediol, T-161
- 1,2,3,4-Tetrahydro-2-methyl-6,7-isoquinolinediol, T-158
- 1,2,3,4-Tetrahydro-1-methyl-6,7,8-isoquinoline-triol, T-237
- 1,2,3,4-Tetrahydro-2-methyl-6,7-methylenedioxyisoquinoline, H-398
- 2,3,4,9-Tetrahydro-2-methyl-1-(2-methyl-1-propenyl)-1*H*-pyrido[3,4-*b*]indole, M-521
- 7,8,9,9*a*-Tetrahydro-3-[(3-methyl-1-oxobutyl)amino]-4-methyl-1*H*,5*H*-pyrrolo[1,2-*c*][1,3]oxazepine-1,5-dione, C-852
- 1,2,3,4-Tetrahydro-2-methyl-1-oxo-β-carboline, T-226
- 2-(Tetrahydro-4-methyl-5-oxo-2-furanyl)stenine, T-664
- 5,7,8,13-Tetrahydro-14-methyl-5-oxoindolo[2',3':3,4]pyrido[2,1-*b*]quinazolinium hydroxide inner salt, D-158
- 1-[1,2,5,6-Tetrahydro-1-methyl-6-(2-oxopropyl)-2-pyridinyl]-3-buten-2-one, S-213
- 3,4,6,7-Tetrahydro-1-methyl-6-oxo-1*H*-pyrano[3,4-*c*]pyridine-5-carboxaldehyde, G-55
- 1,2,3,4-Tetrahydro-1-methyl-2-pentylquinoline, P-238
- 1,3,4,12*b*-Tetrahydro-3-(1-methyl-2-piperidinyl)-pyrido[2,1-*a*][2]benzazepin-6(2*H*)-one, A-1524
- 3,4,5,6-Tetrahydro-6-(2-methyl-1-propenyl)-1*H*-azepino[5,4,3-*cd*]indole, A-1549
- 1,2,3,4-Tetrahydro-1-(2-methylpropyl)-9*H*-pyrido[3,4-*b*]indole, M-521
- 5,6,7,8-Tetrahydro-5-methyl-2-propylquinoline, T-204
- 1,2,5,6-Tetrahydro-1-methyl-3-pyridinecarboxylic acid, T-205
- 20-(3,4,5,6-Tetrahydro-5-methyl-2-pyridinyl)-pregn-5-ene-3,16-diol, 13CI, S-199
- 2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, T-206
- 2,3,4,9-Tetrahydro-2-methyl-1*H*-pyrido[3,4-*b*]indole, T-203
- 2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indole, T-202
- 2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indol-7-ol, H-592
- 2,3,4,9-Tetrahydro-1-methyl-1*H*-pyrido[3,4-*b*]indol-6-ol, S-285
- 2,3,4,9-Tetrahydro-2-methyl-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 1,4,5,6-Tetrahydro-2-methyl-4(6)-pyrimidinecarboxylic acid, T-207
- 2,3,8,8*a*-Tetrahydro-1-methylpyrrolo[2,3-*b*]indol-3*a*(1*H*)-ol, A-626
- 5,6,7,8-Tetrahydro-4-methylquinoline, T-208
- Tetrahydro-2-methylthiazole, M-570
- 2,3,4,5-Tetrahydro-6-methyl-2-undecylpyridine, M-584
- 5,6,7,8-Tetrahydro-*D*-monapterin, A-923
- 5,6,7,8-Tetrahydro-*L*-monapterin, A-923
- Tetrahydrooneosporamine, D-79
- Δ³-Tetrahydrocortic acid, T-223
- Tetrahydronitramine, Q-51
- Tetrahydronitradine, Q-52
- 2,3,8,8*a*-Tetrahydro-3*a*-nitro-1*H*-pyrrolo[2,3-*b*]indole, T-209
- 5,6,7,8-Tetrahydro-2-nonylquinoline, T-210
- Tetrahydroilvacine, G-211
- Tetrahydro-1,4-oxazine, M-705
- 5,7,8,13-Tetrahydro-5-oxobenz[*g*]indolo[2,3-*a*]quinolizine-1-carboxaldehyde, N-67
- 1,2,3,4-Tetrahydro-1-oxo-β-carboline, T-226
- 1,2,3,4-Tetrahydro-1-(1-oxo-2,4,6-decatrienyl)pyridine, D-125
- 1,2,3,4-Tetrahydro-4-(oxoethylidene)-2,6-pyridinedicarboxylic acid, B-109
- 7,8,9,9*a*-Tetrahydro-3-[(1-oxohexyl)amino]-4-methyl-1*H*,5*H*-pyrrolo[1,2-*c*][1,3]oxazepine-1,5-dione, C-852
- 5,6,7,8-Tetrahydro-5-oxoimidazo[1,5-*c*]pyrimidine, D-462
- 2,3,6,11-Tetrahydro-3-oxo-1*H*-indolizino[8,7-*b*]indole-5,11*b*(5*H*)-dicarboxylic acid, T-211
- 1,2,3,12-Tetrahydro-12-oxoindolo[3,2,1-*de*]pyrido[3,2,1-*ij*][1,5]naphthyridin-4-ium(1+), I-140
- 7,8,13,13*b*-Tetrahydro-5-oxoindolo[2',3':3,4]pyrido[2,1-*b*]quinazoline-14(5*H*)-carboxaldehyde, D-508
- 1,5,6,8-Tetrahydro-8-oxo-1,5-methano-2*H*-pyrido[1,2-*a*][1,5]diazocine-3(4*H*)-acetic acid, C-939
- 1,2,3,4-Tetrahydro-1-(1-oxo-3-phenyl-2-propenyl)-5-(2-piperidinyl)pyridine, A-138
- Tetrahydroalpmatine, T-212
- 2,3,4,5-Tetrahydro-6-(1,3-pentadienyl)pyridine, T-213
- 1,2,3,4-Tetrahydro-1-(1,2,3,4,5-pentahydroxypentyl)-β-carboline-3-carboxylic acid, T-215
- 1,2,3,5-Tetrahydro-2,2,5,5,7-pentamethyl-6*H*-pyrrolo[2,3-*b*]pyridin-6-one, A-206
- 2,3,4,5-Tetrahydro-6-(4-pentenyl)pyridine, T-214
- Tetrahydropentoxylene, T-215
- 2,3,4,5-Tetrahydro-6-pentylpyridine, T-216
- 2,3,6,7-Tetrahydro-3-(phenylmethyl)pyrrolo[1,2-*a*]pyrazine-1,4-dione, C-872
- 1,2,3,4-Tetrahydro-1-phenyl-1,4-naphthalenedicarboxylic acid bis(8-methyl-8-azabicyclo[3.2.1]oct-3-yl) ester, B-56
- 1,2,3,4-Tetrahydro-1-phenyl-1,4-naphthalenedicarboxylic acid bis(9-methyl-3-oxa-9-azatricyclo[3.3.1.0^{2,4}]non-7-yl) ester, S-157
- 3*a*,4,5,6-Tetrahydro-3-phenyl-3*H*-pyrrolo[1,2-*b*]pyrazole, N-185
- 1,2,3,4-Tetrahydro-1-phenylquinolinizinium(1+), C-496
- 1,2,3,6-Tetrahydropicolinic acid, T-221
- Tetrahydropiperine, P-467
- Tetrahydroplakinamine A, P-496
- Tetrahydropresecamine, P-624
- Tetrahydropronuciferin, P-643
- 1,2,3,4-Tetrahydro-6-propanoylpyridine, T-217
- 3,4,5,6-Tetrahydro-2-propanoylpyridine, T-217
- 2,3,4,5-Tetrahydro-6-(1-propenyl)pyridine, T-218
- 1,2,3,4-Tetrahydro-1-propyl-β-carboline, T-219
- 2,3,4,9-Tetrahydro-1-propyl-1*H*-pyrido[3,4-*b*]indole, T-219
- 1,2,3,6-Tetrahydro-2-pyridinecarboxylic acid, T-221
- 1,2,3,6-Tetrahydro-4-pyridinecarboxylic acid, T-222
- 1,2,5,6-Tetrahydro-3-pyridinecarboxylic acid, T-223
- 1,2,3,4-Tetrahydropyridine, T-220
- 1-(1,4,5,6-Tetrahydro-2-pyridinyl)ethanone, A-76
- 1-(1,4,5,6-Tetrahydro-2-pyridinyl)-1-propanone, T-217
- 1-(3,4,5,6-Tetrahydro-2-pyridinyl)-1-propanone, T-217
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indole-1-butanamide, N-80
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indole-3-carboxylic acid, T-224
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indole-1,3-dicarboxylic acid, T-147
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indole-1,6-diol, T-225
- 1,2,3,4-Tetrahydro-9*H*-pyrido[3,4-*b*]indole, T-146
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indol-6-ol, T-173
- 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
- 3-[(2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indol-1-yl)propyl]guanidine, T-637
- 6,8,9,11-Tetrahydro-7*H*-pyrido[2,1-*b*]quinazoline, T-227

- 6,7,8,9-Tetrahydro-11H-pyrido[2,1-*b*]quinazoline, T-227
- 6,7,8,9-Tetrahydropyrido[2,1-*b*]quinazolin-11-one, M-5
- 1-[2-(3,4,5,6-Tetrahydropyridyl)]-1,3-pentadiene, T-213
- Tetrahydropyrrole, P-949
- 2,3,4,9-Tetrahydro-1-[2-(1-pyrrolidinyl)ethyl]-1H-pyrido[3,4-*b*]indole, T-34
- 1,2,3,4-Tetrahydro-5-(2-pyrrolidinyl)pyridine, T-228
- 1,2,3,4-Tetrahydro-1-(2-pyrrolidylethyl)- β -carboline, T-34
- 2,3,5,7a-Tetrahydro-1H-pyrrolizine-7-methanol, S-644
- 5,6,7,7a-Tetrahydro-3H-pyrrolizine-3-one, T-229
- 4,5,6,7-Tetrahydro-2H-pyrrolo[3,4-*c*]pyridine-6-carboxylic acid, T-230
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazoline-1-carboxylic acid, L-180
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazoline-3,7-diol, V-37
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazoline, T-231
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazolin-9-ol, D-503
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazolin-3-ol, P-165
- 1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazolin-1-ol, T-231
- 1-(1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazolin-9-yl)-2-propanone, P-164
- 1,2,3,4-Tetrahydro-1-(2-quinolinyl)- β -carboline, Q-51
- 2,3,4,9-Tetrahydro-1-(5-quinolinyl)-1H-pyrido[3,4-*b*]indole, Q-53
- Tetrahydrohombifoline, T-149
- 1,2,3,4-Tetrahydroroeharmin, D-620
- Tetrahydrosecamine, S-174
- Tetrahydrosecidine, S-184
- Tetrahydrosecodinol, S-184
- Tetrahydroserpentine, A-212
- Tetrahydrostephaine, S-530
- 5,5',6,6'-Tetrahydro-1,1',2,2'-tetramethoxy-7,7'-bi-4H-dibenzo[*de,g*]quinoline, U-41
- 5,8,13,13a-Tetrahydro-2,3,10,11-tetramethoxy-6H-dibenzo[*a,g*]quinolizine, X-30
- 5,8,13,13a-Tetrahydro-2,3,9,10-tetramethoxy-6H-dibenzo[*a,g*]quinolizim-11-ol, S-543
- 5,8,13,13a-Tetrahydro-2,3,9,10-tetramethoxy-13,13-dimethyl-6H-dibenzo[*a,g*]quinolizine, C-686
- 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-1,2-dimethylisoquinoline, T-233
- 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxyisoquinoline, T-276
- 6,7,7a,8-Tetrahydro-4,9,10,11-tetramethoxy-7-methyl-5H-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinoline, L-159
- 5,7,8,14-Tetrahydro-3,4,10,11-tetramethoxy-6-methyl-dibenzo[*c,g*]azecine-13(6H)-one, M-734
- 6,7,8,9-Tetrahydro-2,3,10,12-tetramethoxy-7-methyl-5H-dibenzo[*d,f*]azozine, P-671
- 5,6,11,12-Tetrahydro-2,3,8,9-tetramethoxy-13-methyl-dibenzo[*a,e*]cycloocten-5,11-imine, A-1410
- 5,6,6a,7-Tetrahydro-1,2,9,10-tetramethoxy-6-methyl-4H-dibenzo[*de,g*]quinolin-4-ol, C-190
- 5,8,13,13a-Tetrahydro-2,3,9,10-tetramethoxy-13-methyl-6H-dibenzo[*a,g*]quinolizine, C-675
- 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-1-methylisoquinoline, T-233
- 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-2-methylisoquinoline, T-276
- 1,1',2,2'-Tetrahydro-2,2',4,4'-tetramethyl[2,2'-bi-3H-indole]-3,3'-dione, P-271
- 1,6,7,8-Tetrahydro-4,5,6,8-tetramethylcyclopent[*g*]indole, H-140
- 2-[2,4a,10,10a-Tetrahydro-1,1,3,13-tetramethyl-1H,5H-4b,9b-(iminoethano)indeno[1,2-*b*]indol-10-yl]-1H-indole-3-ethanol, S-395
- 6,7,8,9-Tetrahydro-2,5,10,10-tetramethyl-5,8-methano-5H-cyclohepta[*b*]pyridine, P-112
- Tetrahydro-*N,N,N,N*-tetramethyl-4-oxo-2-furan-methanaminium(1+), M-773
- 2,3,7b,10-Tetrahydro-7,7,9,9-tetramethylpyrrolo[1,2-*c*]quinazoline-5,8,10(1H,6H,9H)-trione, S-656
- Tetrahydrothalifendine, C-73
- Tetrahydrothalifendine, T-234
- Tetrahydrothermopsine, I-242
- 1,2,3,4-Tetrahydro-1,2-(thio-3-pyrrolidinyl)- β -carboline-3-carboxylic acid, T-235
- 3,4,8a,13-Tetrahydro-3,15,16-trihydroxy-1,4-dimethoxy-12-methyl-1H-xantheno[4',3',2':4,5][1,3]benzodioxino[7,6-*g*]isoquinoline-14,17(2H,9H)-dione, A-128
- 1,2,3,4-Tetrahydro-5,6,7-trihydroxyisoquinoline, T-236
- 1,6,7,8-Tetrahydro-2,10,11-trihydroxy-8-methoxy-2H-benzo[*e*]azecine-3,5-dione, S-41
- 1,2,3,4-Tetrahydro-6,7,8-trihydroxy-1-methylisoquinoline, T-237
- 5,6,6a,7-Tetrahydro-9,10,12-trimethoxy-4H-benzo[*g*]-1,3-benzodioxolo[4,5,6-*de*]quinoline, B-7
- 5,8,13,13a-Tetrahydro-2,3,10-trimethoxy-6H-dibenzo[*a,g*]quinolizine-1,11-diol, S-530
- 5,8,13,13a-Tetrahydro-2,9,10-trimethoxy-6H-dibenzo[*a,g*]quinolizim-1-ol, C-174
- 5,8,13,13a-Tetrahydro-2,3,9-trimethoxy-6H-dibenzo[*a,g*]quinolizim-10-ol, C-677
- 5,8,13,13a-Tetrahydro-3,9,10-trimethoxy-6H-dibenzo[*a,g*]quinolizim-2-ol, I-210
- 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1,2-dimethylisoquinoline, T-174
- 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1,2-dimethylisoquinoline, T-237
- 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-isoquinolinecarboxylic acid, T-175
- 1,2,3,4-Tetrahydro-5,6,7-trimethoxyisoquinoline, T-238
- 1,2,3,4-Tetrahydro-6,7,8-trimethoxyisoquinoline, T-236
- 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-1-[(4-methoxyphenyl)methyl]isoquinoline, A-1069
- 2,3,12,12a-Tetrahydro-6,9,10-trimethoxy-1-methyl-1H-[1]benzoxepino[2,3,4-*ij*]isoquinoline, C-805
- 6,7,8,9-Tetrahydro-2,3,12-trimethoxy-7-methyl-5H-dibenzo[*d,f*]azozine, L-73
- 6,7,8,9-Tetrahydro-2,11,12-trimethoxy-7-methyl-5H-dibenzo[*d,f*]azozin-3-ol, E-194
- 5,6,11,12-Tetrahydro-3,8,9-trimethoxy-13-methyl-dibenzo[*a,e*]cycloocten-5,11-imin-4-ol, P-514
- 5,6,6a,7-Tetrahydro-1,2,9-trimethoxy-6-methyl-4H-dibenzo[*de,g*]quinolin-10-ol, H-752
- 5,6,6a,7-Tetrahydro-2,9,10-trimethoxy-6-methyl-4H-dibenzo[*de,g*]quinolin-1-ol, H-751
- 5,6,6a,7-Tetrahydro-1,2,10-trimethoxy-6-methyl-4H-dibenzo[*d,e*]quinolin-9-ol, L-75
- 5,6,6a,7-Tetrahydro-1,9,10-trimethoxy-6-methyl-4H-dibenzo[*de,g*]quinolin-2-ol, P-597
- 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-methyl-1-isoquinolinecarboxylic acid, P-299
- 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-2-methylisoquinoline, T-238
- 1,2,3,4-Tetrahydro-6,7,8-trimethoxy-1-methylisoquinoline, T-237
- 5,7,8,13-Tetrahydro-2,3,10-trimethoxy-14-methyl-5-oxoindolo[2',3':3,4]pyrido[2,1-*b*]quinazolium hydroxide inner salt, E-309
- 6,7,8,9-Tetrahydro-2,3,10-trimethylcyclohepta[*b*]pyran-5,8-imin-4(5H)-one, D-75
- 2,4,4a,8-Tetrahydro-2,6,8-trimethyl-4a-(2-oxo-propyl)pyrimido[5,4-*e*]-1,2,4-triazine-3,5,7(6H)-trione, P-935
- 2,3,4,6-Tetrahydro-1,2,5-trimethyl-1H-pyrido[4,3-*b*]carbazole, G-211
- N*-(1,2,5,6-Tetrahydro-2,5,6-trioxo-3-pyridyl)octanamide, O-62
- 1,5,6,7-Tetrahydro-4-(1,3-undecadienyl)-2H-azepin-2-one, M-698
- 4',5',6',17-Tetrahydro-17R-usambarensine *N*-oxide, U-52
- Tetrahydro-17R-usambarensine, U-52
- Tetrahydro-17S-usambarensine, U-52
- Tetrahydroveralkamine, V-61
- 1,8,14,16-Tetrahydroxyaconitan-19-one, T-239
- 1,2,3,4-Tetrahydroxyacridone, T-240
- 1,2,3,5-Tetrahydroxyacridone, T-241
- 1,3,4,5-Tetrahydroxyacridone, T-242
- 1,3,5,6-Tetrahydroxyacridone, T-243
- 1,2,3,9-Tetrahydroxyaporphine, T-244
- 1,2,9,10-Tetrahydroxyaporphine, T-246
- 1,2,9,11-Tetrahydroxyaporphine, T-247
- 1,2,10,11-Tetrahydroxyaporphine, T-248
- 1,2,3,11-Tetrahydroxyaporphine, T-245
- 5,5',6,6'-Tetrahydroxy-3,3'-bi-1H-indole, B-130
- 3,3',4,4'-Tetrahydroxy-2,2'-bipyridine, T-249
- 2-(1,2,3,4-Tetrahydroxybutyl)-5-(2,3,4-trihydroxybutyl)pyrazine, T-250
- 1,2,3,7-Tetrahydroxydibenzo[*cd,f*]indol-4(5H)-one, T-254
- 1,2,3,9-Tetrahydroxydibenzo[*cd,f*]indol-4(5H)-one, T-255
- 1,2,6,7-Tetrahydroxydibenzo[*cd,f*]indol-4(5H)-one, T-251
- 1,2,7,9-Tetrahydroxydibenzo[*cd,f*]indol-4(5H)-one, T-253
- 1,2,7,8-Tetrahydroxydibenzo[*cd,f*]indol-4(5H)-one, T-252
- 4,6,8,9-Tetrahydroxydihydro- β -agarofuran-1-one, P-219
- 1,2,6,9-Tetrahydroxydihydro- β -agarofuran, T-256
- 1,4,6,9-Tetrahydroxydihydro- β -agarofuran, T-257
- 1,6,8,9-Tetrahydroxydihydro- β -agarofuran, T-258
- 1,8,9,14-Tetrahydroxydihydro- β -agarofuran, T-259
- 10,10',17,17'-Tetrahydroxy-4,4'-dimethyl-9,9'-bisarpaganium, D-864
- 6,13,14,21-Tetrahydroxy-16,18-dimethyl-10-phenyl[1]cytochalasa-7,19-dien-1-one, T-260
- 4,5,6,7-Tetrahydroxyfuro[2,3-*b*]quinoline, F-215
- 4,5,7,8-Tetrahydroxyfuro[2,3-*b*]quinoline, F-216
- 4,6,7,8-Tetrahydroxyfuro[2,3-*b*]quinoline, F-217
- 5,6,7,8-Tetrahydroxyfuro[2,3-*b*]quinolin-4(9H)-one, F-214
- 1,2,6,7-Tetrahydroxy-3-(hydroxymethyl)-5-methyl-1H-pyrrolizidine, T-261
- 1,2,6,7-Tetrahydroxy-3-(hydroxymethyl)-1H-pyrrolizidine, T-262
- 1,6,7,8-Tetrahydroxyindolizidine, C-188
- 3,5,7,15-Tetrahydroxy-6(17),12-lathyradien-14-one, T-263
- 1,3,4,5-Tetrahydroxy-2-(3-methyl-2-butenyl)-9(10H)-acridinone, T-270
- 1,3,5,6-Tetrahydroxy-2-(3-methyl-2-butenyl)-9H-acridin-9-one, T-271
- 3,4',5,7-Tetrahydroxy-8-(4-methyl-5-oxo-2-pyrrolidinyl)flavone, L-173
- 2,3,4,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-264
- 3,4,6,7-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-265
- 3,4,6,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-266
- 3,4,7,8-Tetrahydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-267
- 1,2,3,4-Tetrahydroxynortropane, A-1587
- 1,2,3,6-Tetrahydroxynortropane, A-1588
- 1,2,4,7-Tetrahydroxynortropane, A-1589
- 3',4',5,7-Tetrahydroxy-6-(5-oxo-2-pyrrolidinyl)flavanone, T-572
- 3',4',5,7-Tetrahydroxy-8-(5-oxo-2-pyrrolidinyl)flavanone, T-573
- 3,11,12,14-Tetrahydroxypregn-5-en-20-one, T-269
- 3,8,12,14-Tetrahydroxypregn-5-en-20-one, T-268
- 1,3,4,5-Tetrahydroxy-2-prenylacridone, T-270
- 1,3,5,6-Tetrahydroxy-2-prenylacridone, T-271
- 3,4,7,8-Tetrahydroxy-2-quinolinecarboxylic acid, T-272
- 1,5,6,7-Tetrahydroxyquinoline, I-304
- 2,3,4,7-Tetrahydroxyquinoline, Q-40
- 2,3,4,8-Tetrahydroxyquinoline, Q-41
- 2,4,6,8-Tetrahydroxyquinoline, Q-42
- 2,4,7,8-Tetrahydroxyquinoline, Q-43
- 3,4,5,8-Tetrahydroxyquinoline, Q-44
- 3,4,7,8-Tetrahydroxy-2(1H)-quinolinone, Q-39
- 5,7,9,10-Tetrahydroxy-4(20),11-taxadien-13-one, T-48

- Thalirevoline, T-333
 Thalirevolutine, T-333
 Thalirugidine, T-334
 Thalirugine, T-335
 Thaliruginine, T-335
 Thalissamine, H-148
 Thalispavine, T-336
 Thalispidine, T-346
 Thalispine, T-346
 Thalispinine, P-208
 Thalispynine, P-208
 Thalisyprine, C-790
 Thalistine, T-337
 Thalistryline, T-337
 Thalivarminine, T-313
 Thaliximine, H-148
 Thalixine, T-327
 Thallusin, T-338
 Thalmelatidine, P-683
 Thalmelatine, T-312
 Thalmethine, T-339
 Thalmiculatimine, T-340
 Thalmiculimine, T-340
 Thalmiculine, T-340
 Thalmidine, T-313
 Thalmine, T-342
 Thalmine, T-341
 Thalmirabine, T-343
 Thalpetaline, T-344
 Thalphenine methine, T-327
 Thalphenine, T-345
 Thalphine, T-304
 Thalpinine, T-305
 Thalpindione, T-320
 Thalprzewalskiinone, P-77
 Thalrugosamine, O-1
 Thalrugosaminine 2 β -*N*-oxide, T-346
 Thalrugosaminine, T-346
 Thalrugosidine, T-320
 Thalrugosine, I-328
 Thalrugosinone, T-320
 Thalsimidine, T-347
 Thalsimine, T-347
 Thalsivasine, T-339
 Thapoxepine A, T-348
 Thapoxepine B, T-348
 Thapsakin B, T-349
 Thapsakone A, T-349
 Thapsakone B, T-349
 Thaspine, T-36
 Theacrine, U-45
 Thebaine *N*-methosalt, T-350
 Thebaine *N*-oxide, T-350
 Thebaine, T-350
 Thehaplosin, T-351
 Theine, C-15
 Thelephantin N, T-97
Agelas nakamurai Thelepogane-purine alkaloid, T-352
 Thelepogidine, T-353
 Thelepogine, T-354
 Theobromine, T-355
 Theocin, T-360
 Theodel, T-360
 Theodoxine, P-923
 Theo-Dur, T-360
 Theograd, T-360
 Theolair, T-360
 Theona P, T-360
 Theoneberine, T-356
 Theonelladine A, T-357
 Theonelladine B, T-357
 Theonelladine C, T-358
 Theonelladine D, T-358
 Theonelline formamide, I-213
 Theonelline isocyanide, I-213
 Theonelline isothiocyanate, I-213
 Theonezolidine, T-359
 Theophylline, T-360
 Theotard, T-360
 Thermine, B-154
 Thermohexamine, D-284
 Thermopentamine, D-289
 Thermopsamine, T-361
 Thermopsidine, T-362
 Thermopsine, T-363
 Thermopterin, T-37
 Thermospermine, D-249
 Thesal, T-355
 Thesine, T-364
 Thesinicine, T-365
 Thesinine 4'-glucoside, H-629
 Thesinine 4'-rhamnoside, H-629
 Thesinine, H-629
 Thiactrat, T-366
 Thiactin, T-392
 Thiamine diphosphate, T-366
 Thiamine hydrochloride, T-366
 Thiamine monochloride, T-366
 Thiamine mononitrate, T-366
 Thiamine pyrophosphate, T-366
 Thiamine triphosphate, T-366
 Thiamine triphosphoric acid, T-366
 Thiamine, T-366
 Thiangazole, T-367
 Thiaplidiaquinone A, T-368
 Thiaplidiaquinone B, T-369
 Thiasine, E-173
 Thiazohalostatin, T-370
 5-Thiazolethanol, T-371
 Thiazolidomycin, A-78
 2-(5-Thiazolyl)ethanol, T-371
 3-(2-Thiazolyl)-1*H*-indole, T-372
 2-(2-Thiazolyl)phenol, H-694
 Thiazomycin, N-266
 Thiazostatin A, T-373
 Thiazostatin B, T-373
 Thienodolin, T-374
 7-(2-Thienyl)-2,4,6-heptatrienoic acid, O-133
 Thiersindole A, T-375
 Thiersindole B, T-375
 Thiersindole C, T-376
 Thiersinine A, T-377
 Thiersinine B, T-377
 Thioactin, T-378
 Thioaurin, O-124
 Thiobideoxynupharidine, T-379
anti-Thiobinupharidine sulfoxide, T-380
 Thiobinupharidine, T-380
 Thiobutacin, T-381
Perophora viridis 1,2,3-Thiocane glycoside, T-382
 Thiocillin III, T-383
 Thiocillin II, T-383
 Thiocillin I, T-383
 Thiocoraline, T-384
 Thiofeed, S-321
 Thiolhistidine trimethylbetaine, E-173
 Thiolutin, A-759
 Thiomarinol A, T-385
 Thiomarinol B, T-385
 Thiomarinol C, T-385
 Thiomarinol D, T-386
 Thiomarinol E, T-387
 Thiomarinol F, T-385
 Thiomarinol G, T-385
 2-(Thiomethyl)quinoline, Q-36
 Thiomycololide A, T-388
 Thiomycololide B, T-389
 Thiomycin, T-390
 Thioneine, E-173
 Thionupharodioline, T-380
 Thionupharoline, T-380
 Thionuphltine A, T-380
 Thionuphltine B α -*S*-oxide, T-380
 Thionuphltine B β -*S*-oxide, T-380
 Thionuphltine B, T-380
 Thiopeptin A_{1a}, S-321
 Thiopeptin A_{3a}, S-321
 Thiopeptin A_{4a}, S-321
 Thiopeptin A_{1b}, S-321
 Thiopeptin A_{3b}, S-321
 Thiopeptin A_{4b}, S-321
 Thiopeptin B_a, S-321
 Thiopeptin B_b, S-321
 Thioplabin B, A-1103
 Thioplabin C, A-1108
 Thioprol, H-715
 Thiosporamicin, T-391
 Thiostrepton A, T-392
 Thiostrepton A₂, T-392
 Thiostrepton B, T-392
 Thiostrepton, T-392
 Thiotaaurine, A-766
 Thiotax, B-79
 Thiotipin, T-393
 Thiouracil, D-518
 2-Thiouracil, D-518
 4-Thiouracil, H-556
 2-Thioxo-4-thiazolidinecarboxylic acid, T-394
 Thomandersine, T-195
 Thomasic acid, T-395
 Thomitrem A, S-194
 Thomitrem E, S-194
 Thomsomycin, A-14
 Thorazine, C-417
 Thoretandramine, T-396
 Thrombocytin, H-755
 Thrombotonin, H-755
 Thuggacin A, T-397
 Thuggacin B, T-398
 Thuggacin cmc-A, T-400
 Thuggacin cmc-B, T-400
 Thuggacin cmc-C, T-400
 Thuggacin C, T-399
 Thymine, T-401
 Thyminol, T-401
 Tiantaishandine, H-208
 Tiantaishanmine, M-368
 Tiantaishannine, M-321
 Tiantaishansine, D-187
 Tienemine 3-glucoside, S-200
 Tienmulimline, T-402
 Tienmuliminine, T-403
 Tiformin, G-201
 Tigloidine, T-404
N-Tigloylbuxahyrcanine, B-451
O-Tigloylcyclovirobuxeine B, C-915
 Tigloyllaburnine, H-629
 13 α -Tigloyloxyaphylline, V-153
 13-Tigloyloxylyupanine, H-554
 13 α -Tigloyloxy multiflorine, M-731
 3-Tigloyloxynortropan-6-ol, A-1584
 13-Tigloyloxy-17-oxolupanine, O-188
 3 α -Tigloyloxytropane *N*-oxide, T-404
 7-Tigloyloxytropane-3,6-diol, M-392
 3-Tigloyloxytropane, T-404
 6-Tigloyloxytrop-3-ol, M-391
 (\pm)-3-*O*-Tigloyloxytrop-6 β -ol, M-391
N-Tigloylpyrrothine, A-759
 Tigloyltropeine, T-404
N-Tigloyltyramine, H-667
O-Tigloylvirgiline, V-153
 Tiliacoridine, T-405
 Tiliacrine, T-406
 Tiliacrine 2'-*N*-oxide, T-407
 Tiliacrine, T-407
 Tiliacosine, Y-5
 Tiliafunimine, T-408
 Tiliageine, R-114
 Tiliamosine, Y-5
 Tilianangine, Y-5
 Tiliandrine, T-409
 Tiliaresine, T-410
 Tiliarine, T-407
 Tiliatine, Y-5
 Tilitriandrine, R-114
 Tilivalline, T-411
 Tinctoramine, T-269
 Tinctorine, T-412
 Tinctorimine, T-413
 Tinosporine, T-414
 Tintamine, T-415
 Tirandalydigin, T-416
 Tirandamycin A, T-417
 Tirandamycin B, T-417
 Tirucalicine, I-147
 Tiruchanduramine, T-418
 Tjipanazole A1, D-350
 Tjipanazole A2, D-350

- Tjipanazole B, D-350
 Tjipanazole C1, C-401
 Tjipanazole C2, C-401
 Tjipanazole C3, C-401
 Tjipanazole C4, C-401
 Tjipanazole D, D-350
 Tjipanazole E, D-350
 Tjipanazole F1, C-401
 Tjipanazole F2, C-401
 Tjipanazole G1, D-464
 Tjipanazole G2, D-464
 Tjipanazole I, C-401
 Tjipanazole J, T-419
 TK 57-164A, D-98
 TK 57-164B, D-98
 TM 64, A-1286
 TMC 169, A-1502
 TMC 260, A-1290
 TMC 52, A-1287
 TMC 66, A-1288
 TMC 69, A-1289
 TMC 120A, A-1294
 TMC 135A, A-1295
 TMC 1A, A-1291
 TMC 2A, A-1292
 TMC 52A, A-1287
 TMC 95A, A-1293
 TMC 49A, P-343
 TMC 120B, A-1294
 TMC 135B, A-1296
 TMC 2B, A-1292
 TMC 52B, A-1287
 TMC 95B, A-1293
 TMC 34, C-634
 TMC 120C, A-1294
 TMC 1C, A-1297
 TMC 2C, A-1292
 TMC 52C, A-1287
 TMC 95C, A-1293
 TMC 1D, A-1297
 TMC 52D, A-1287
 TMC 95D, A-1293
 TMC 1E, M-89
 TMC 1F, M-87
 TMC 1G, M-88
 TMC 205, M-408
 TMPEA, M-219
 Toddacoumalone, T-420
 Toddaliamide, T-421
 Toddalidimerine, C-355
 Toddaline, C-356
 Toddalinine, D-407
 Toddaliopsin A, T-516
 Toddaliopsin B, T-516
 Toddaliopsin C, T-516
 Toddaliopsin D, T-516
 Toddaquinoline, H-599
 Togholumine, T-451
 Tokaradine A, T-422
 Tokaradine B, T-422
 Tokaradine C, S-398
 Tokoro-base II, M-323
m-Toluic acid, M-398
 α -Toluic acid, P-336
 3-*o*-Tolylacrylic acid, M-513
 Tolypocin, P-922
 Tolypocladine, B-255
 5 α -Tomatidan-3 β -ol (obsol.), S-445
 5 α -Tomatidan-3-one, S-445
 Tomatidenol 3-lycotetraoside, S-456
 δ^5 -Tomatidenol, S-456
 Tomatid-5-en-3-ol, S-456
 Tomatidenol, S-456
 Tomatidine, S-445
 Tomatillidine, T-423
 α -Tomatine, S-445
 β_1 -Tomatine, S-445
 δ -Tomatine, S-445
 Tomatine, S-445
 γ -Tomatine, S-445
 Tomaymycin I, T-424
 Tomaymycin, T-424
 Tombosine, T-425
 Tombozine, T-425
 Tomentocurine, T-426
 Tomentogenin, P-604
 Tomentomin, P-604
 Tomentosine, O-137
 Tongolenine C, M-370
 Tongolenine D, H-253
 Tongoline, M-321
 Tongolinine, T-427
 Tonkinensine A, T-428
 Tonkinensine B, T-429
meta-Topolin 9-glucoside, A-137
meta-Topolin, A-137
 Topostin B 567, M-485
 Topostin B553, T-430
 Topostin D 640, T-431
 Toppentin A, T-432
 Toppentin B1, T-432
 Toppentin B2, T-432
 Toppentin C, T-432
 Toppentin D, T-432
 Toppentin, T-432
 Torokonine, H-187
 Torondel, E-6
 Tortifoline, C-290
 Tortugamide, L-356
 Tortumine, M-309
 Tortuosamine, T-433
 Tortuosine, V-35
 Torulin, T-366
 Torulosine, T-434
 Torulosinine, T-435
 Tosanpin, Y-13
N-Tosyl-3-azetidione, A-1606
 Toxicoferine, C-812
C-Toxiferine II, C-31
 Toxiferine III, T-438
 Toxiferine II, T-437
C-Toxiferine I, T-436
 Toxiferine I, T-436
 Toxiferine IV, C-31
 Toxiferine IX, C-111
 Toxiferine VIII, T-439
 Toxiferine V, T-436
 Toxiferine XII, T-440
 Toxiferine XI, T-436
 Toxin A, R-157
 Toxin B, P-478
 Toxin B, R-157
 Toxin B₁, S-109
 Toxin B₂, S-109
 Toxin C₁, H-731
 Toxin C₂, H-731
 Toxin C₃, H-731
 Toxin C₄, H-731
 Toxin MG 30, T-441
 Toxin PX₁, H-731
 Toxin PX₂, H-731
 Toxin PX₃, H-731
 Toxin PX₄, H-731
Aphonopelma chalcodes Toxin, A-1332
 β -Toxin, A-837
 Toxoflavin, R-66
 Toyocamycin 5'- α -D-glucopyranose, S-53
 Toyocamycin, S-53
 TPU 0037A, L-353
 TPU 0037B, L-353
 TPU 0037C, L-353
 TPU 0037D, L-353
 TR 1736, M-262
 Trabectedin, E-37
 Trabzonine, T-442
 Trachelanthamidine curassavate, C-810
 Trachelanthamidine strigosate, S-585
 Trachelanthamidine, H-629
 (\pm)-Trachelanthamidine, H-629
 Trachelanthamidinic acid, P-960
 Trachelanthamine *N*-oxide, T-443
 Trachelanthamine, T-443
 Trachelanthine, T-443
 Trachyone, T-444
 Tramesanguin, C-452
 Tranaxine, S-159
 Transconitine A, M-352
 Transconitine B, M-329
 Transconitine C, M-375
 Transcop, S-159
 Transitorine, H-725
 Transtorine, H-725
 Transvalencin A, T-445
 Transvalencin Z, T-446
 Treflorine, T-449
 Trehalamine, T-447
 Trehalostatin, T-447
 Trehazolin, T-447
 Tremidine, T-448
 Tremine, T-448
 Tremortin A, P-194
 Trenudine, T-449
 Trewiasine, T-450
 Triabunnine, S-259
 Triacanthine, T-451
 Triacetanamine, T-283
 2,6,14-Triacetoxy-1-benzoyloxydihydro-9-nicotinoyloxy- β -agarofuran, H-247
 2,6,14-Triacetoxy-9-(benzoyloxy)dihydro-1-nicotinoyloxy- β -agarofuran, H-247
 2,6,14-Triacetoxy-1-furoyloxydihydro-9-nicotinoyloxy- β -agarofuran, H-247
 2,6,14-Triacetoxy-9-furoyloxydihydro-1-nicotinoyloxy- β -agarofuran, H-247
 2,16,28-Triacetylbutirane, B-460
 2,16,28-Triacetyl-9,10-dihydrobutirane, B-460
 Triacetylglycoperine, F-223
 1-Triacontanamine, T-452
N^b-(25-Triacontenyl)tryptamine, T-639
 1-Triacontylamine, T-452
N-Triacontylpentadecanamide, T-452
 $\alpha,\alpha',2$ -Triamino- $\gamma,\gamma',3$ -trioxo-3*H*-phenoxazine-1,9-dibutanoic acid, T-453
 3,3',3''-Triaminotripropylamine, T-622
 Triangularicine, T-454
 Triangularine A, T-662
 Triangularine B, T-251
 Triangularine, T-454
 Trianthemine, T-455
 Trianthil, X-7
 Trianthine, Z-15
 1,5,10-Triazadecane, S-398
 4,8,12-Triazahexadecane-1,16-diamine, D-288
 2,4,6-Triaza-3-iminobicyclo[3.2.1]octane-7-carboxylic acid, V-144
 1,5,9-Triazanone, D-255
 4,8,12-Triazapentadecane-1,15-diamine, T-456
 1,5,11-Triazaundecane, A-897
 1,6,11-Triazaundecane, D-252
 1,6,14-Tribenzoylsenbusine A, M-323
 Tribenzylamine, T-457
 2,3,5-Tri-*O*-benzyl-1,4-benzylimino-1,4-dideoxy-D-arabinitol, D-593
 2,3,5-Tri-*O*-benzyl-1,4-benzylimino-1,4-dideoxy-L-xylitol, D-593
 Tribenzylmethylammonium(1+), T-457
 Tribromoacetamide, T-458
 Tribromoacetic acid, T-458
 Tribromoacetonitrile, T-458
 4,5,6-Tribromo-2,3-bis(methylthio)-1*H*-indole, D-304
 2,3,4-Tribromo-5-(5-bromo-2-hydroxybenzoyl)pyrrole, P-966
 2,4,7(3,4,7)-Tribromo-3(2)-chloro-1*H*-indole, T-459
 Tribromocyanomethane, T-458
N-(2,3,6-Tribromo-4,5-dihydroxybenzyl)-2-pyrrolidinone, P-956
 2,4,6-Tribromo-1,7-dimethoxy-*N,N*-dimethyl-1*H*-indole-3-ethanamine, C-630
 2,3,4-Tribromo-5,6-dimethoxy-*N*-methylbenzeneethanamine, C-628
 2,4,6-Tribromo-3-(2-dimethylaminoethyl)-1,7-dimethoxyindole, C-630
 2,5,6-Tribromo-3-(dimethylaminomethyl)-1-methylindole, T-472
 2,5,6-Tribromo-3-formylindole, T-469
 2,4,6-Tribromo-*N*-formyl-3-methoxy-*N*-methylphenethylamine, T-460

- 2,4,6-Tribromo-*N*-formyl-3-methoxyphenethylamine, T-460
 Tribromoglyoxaline, T-461
 2,4,6-Tribromo-3-hydroxyphenethylamine, T-460
 2-(2,4,6-Tribromo-3-hydroxyphenyl)ethylamine, T-460
 2,4,5-Tribromo-1*H*-imidazole, T-461
 2,5,6-Tribromo-1*H*-indole-3-carboxaldehyde, T-469
 2,3,5-Tribromo-1*H*-indole, T-462
 2,3,6-Tribromo-1*H*-indole, T-463
 2,3,7-Tribromo-1*H*-indole, T-464
 2,4,6-Tribromo-1*H*-indole, T-465
 3,4,6-Tribromo-1*H*-indole, T-466
 3,5,7-Tribromo-1*H*-indole, T-468
 3,5,6-Tribromo-1*H*-indole, T-467
 2,5,5'-Tribromo-7'-methoxy-3,4'-bi-1*H*-indole, T-116
 3',5,5'-Tribromo-7'-methoxy-3,4'-bi-1*H*-indole, T-116
 5,6',7-Tribromo-6-methoxyindigotin, T-470
 3,5,7-Tribromo-6-methoxy-1*H*-indole, T-471
 2,5,6-Tribromo-*N*-methylgramine *N*-oxide, T-472
 2,5,6-Tribromo-*N*-methylgramine, T-472
 2,4,5-Tribromo-1-methyl-1*H*-imidazole, T-461
 2,5,6-Tribromo-1-methyl-1*H*-indole-3-carboxaldehyde, T-469
 2,3,5-Tribromo-1-methyl-1*H*-indole, T-462
 2,3,6-Tribromo-1-methyl-1*H*-indole, T-463
 3,5,6-Tribromo-1-methyl-1*H*-indole, T-467
 3,4,5-Tribromo-1-methyl-1*H*-pyrrole-2-carboxylic acid, T-475
 2,4,6-Tribromo-3-(methylsulfinyl)-1*H*-indole, T-473
 4,5,6-Tribromo-2-(methylsulfinyl)-3-(methylthio)-1*H*-indole, D-304
 2,4,6-Tribromo-3-methylthio-1*H*-indole, T-473
 3,4,5-Tribromo-1*H*-pyrrole-2-carboxylic acid, T-475
 2,3,4-Tribromo-1*H*-pyrrole, T-474
 3,3',4-Tribromo-4',5,5'-trichloro-2,2'-bi-1*H*-pyrrole, T-476
 3,3',4-Tribromo-4',5,5'-trichloro-1,1'-dimethyl-2,2'-bi-1*H*-pyrrole, T-476
 2,5,6-Tribromo-*N,N,N*,1-trimethyl-1*H*-indole-3-methanamine, T-472
 Tribulusamide A, G-182
 Tribulusamide B, G-182
 Tribulusterine, P-270
 N^1, N^5, N^{10} -Tricafeoylspermidine, S-398
 Trichamide, T-477
 Trichanthin A, T-478
 Trichanthin B, T-478
 3 α -(4,4,4-Trichloro-2-butyryloxy)tropane, T-479
 2,3,4-Trichloro-5-(3,5-dichloro-2-hydroxybenzoyl)-pyrrole, P-964
 3,4,5-Trichloro-2-(2,3-dichloro-1*H*-pyrrol-1-yl)phenol, N-136
 2,3,8-Trichloro-6-ethyl-5-hydroxy[1]benzopyrano[2,3-*b*]pyrrol-4(1*H*)-one, A-1280
 2,3,4-Trichloro-1*H*-indole, T-480
 2,3,7-Trichloro-1*H*-indole, T-481
 8-Trichloromethylidhydroberberine, P-35
 5,5,5-Trichloro-4-methyl-2-[methyl(4,4,4-trichloro-3-methyl-1-oxobutyl)amino]-*N*-[1-(2-thiazolylethyl)]pentanamide, D-972
 4,4,4-Trichloro-3-methyl-*N*-[4,4,4-trichloro-3-methyl-1-(2-thiazolyl)butyl]butanamide, D-970
 1,1,1-Trichloro-5-undecanamine, T-482
 1,1,1-Trichloro-5-undecylamine, T-482
 Trichocereine, M-219
 Trichochrome B, T-483
 Trichochrome C, T-484
 Trichochrome E, T-485
 Trichochrome F, T-486
 Trichoderamide A, T-487
 Trichoderamide B, T-487
 Trichodesmine *N*-oxide, T-488
 Trichodesmine, T-488
 Trichoguatine, T-489
 Tricholein, M-455
 Tricholomic acid, T-490
 Trichonine, T-491
 Trichophidine, A-864
 Trichophydine, A-864
 Trichophylline, T-492
 Trichophysine, A-864
 Trichopanic acid, T-493
 Trichosanatine, T-494
 Trichosetin, E-142
 Trichosiderin B, T-483
 Trichosiderin C, T-484
 Trichostachine, T-495
 Trichostatin A, T-496
 Trichostatin B, T-496
 Trichostatin C, T-496
 Trichostatin D, T-496
 Trichostatin RK, T-496
 Trichotomine G₁, T-497
 Trichotomine, T-497
 Triclisine, T-498
 Tricordatine, I-339
 Tricornine, M-309
 5-(14*Z*,17*Z*-Tricosadienyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 3-(7,16-Tricosadienyl)-1*H*-pyrrole-2-carboxaldehyde, T-499
N-Tricosanoylthranilic acid, A-706
 N^b -Tricosanolytryptamine, T-639
 5-(14*Z*,17*Z*,20*Z*-Tricosatrienyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 5-(14*Z*-Tricosenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 5-(9*Z*-Tricosenyl)-1*H*-pyrrole-2-carboxaldehyde, A-624
 3-(16-Tricosenyl)-1*H*-pyrrole-2-carboxaldehyde, T-499
 N,N',N'' -Tricoumaroylspermidine, S-398
 Tricotonyltetramine, D-888
 2',3,9-Trideacetylaustrospicatine, T-48
 2',13,14-Trideacetylaustrotaxine, P-224
 Trideacetylpyrripropene A, T-500
 2,7-Tridecadiene-10,12-diyonic acid isobutylamide, T-501
 2,7-Tridecadiene-10,12-diyonic acid 2-methylbutylamide, T-501
 2,7-Tridecadiene-10,12-diyonic acid, T-501
 2,4-Tridecadienoic acid isobutylamide, T-502
 2,4-Tridecadienoic acid, T-502
 6-Tridecanamine, T-506
 Tridecanocholelytrine, T-503
N-Tridecanoylserine, T-700
 2,6,8-Tridecatrien-10,12-diyonic acid isobutylamide, T-504
 2,6,8-Tridecatrien-10,12-diyonic acid, T-504
 7-Tridecene-10,12-diyonic acid, T-505
 2-(1*E*-Tridecenyl)-4(1*H*)-quinolinone, H-750
 2-(1*Z*-Tridecenyl)-4(1*H*)-quinolinone, H-750
 6-Tridecylamine, T-506
 2-Tridecyl-4-quinolinol, H-750
 2-Tridecyl-4(1*H*)-quinolinone *N*-oxide, H-750
 2-Tridecyl-4(1*H*)-quinolinone, H-750
 8-Tridecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-387
 1,3,4-Tridehydrofangchinolinium, C-171
 Tridentatol A, T-507
 Tridentatol B, T-507
 Tridentatol C, T-508
 Tridentatol D, T-509
 Tridentatol E, T-509
 Tridentatol F, T-507
 Tridentatol G, T-509
 Tridentatol H, T-508
 5,6,11-Trideoxy-4-epitetradotoxin, T-510
 1,2,5-Trideoxy-2,5-imino-*D*-altritol, D-591
 1,2,5-Trideoxy-2,5-imino-*D*-galactitol, D-591
 1,2,5-Trideoxy-2,5-imino-*D*-glucitol, D-591
 1,2,5-Trideoxy-2,5-imino-*L*-glucitol, D-591
 2,3,6-Trideoxy-2,6-iminoheptitol, B-185
 2,5,7-Trideoxy-2,5-imino-*D*-glycero-*D*-manno-heptitol, D-587
 1,2,6-Trideoxy-2,6-iminoheptitol, H-607
 1,4,5-Trideoxy-1,5-iminoheptitol, H-615
 2,5,6-Trideoxy-2,5-imino-*D*-gulo-heptitol, H-492
 2,5,6-Trideoxy-2,5-imino-*D*-manno-heptitol, H-492
 1,2,5-Trideoxy-1,5-iminoheptitol, H-614
 1,2,4-Trideoxy-1,4-iminopentitol, H-717
 1,3,4-Trideoxyojirimycin, H-699
 5,6,11-Trideoxytetradotoxin, T-510
 Tridictyophylline, T-511
 Trididemnic acid A, T-512
 Trididemnic acid B, T-512
 Tridolgosir hydrochloride, S-650
 Tridolgosir, S-650
 Triedimycin A, O-155
 Triedimycin B, O-155
 2,3,7-Triepiaustraline, T-555
 N^1, N^5, N^{10} -Triferuloylspermidine, S-398
 20-Trifluoroarachidonic acid, E-48
 N -[4-(Trifluoromethyl)benzyl]cinchoninium, C-447
 Trifolioliasine A, M-309
 Trifolioliasine B, D-196
 Trifolioliasine C, D-196
 2',3',6'-Tri-*O*-galloylprunasin, H-673
 3',4',6'-Tri-*O*-galloylprunasin, H-673
 Trigamine *N*-oxide, T-513
 Trigamine, T-513
 Trigilletimine, T-514
 Trigillette, I-339
 Triglochinin, T-515
 Trigonelline, C-149
 Trigoxazonane, A-1603
 1,2,4-Triheptylpyrazolidine, G-16
 1,2,4-Triheptyltetrahydroprazole, G-16
 1,2,3-Trihydroxy-9(10*H*)-acridinone, T-516
 1,3,4-Trihydroxy-9(10*H*)-acridinone, T-517
 1,3,5-Trihydroxy-9(10*H*)-acridinone, T-518
 1,3,6-Trihydroxy-9(10*H*)-acridinone, T-519
 1,3,8-Trihydroxy-9(10*H*)-acridinone, T-520
 1,2,3-Trihydroxyacridone, T-516
 1,3,4-Trihydroxyacridone, T-517
 1,3,5-Trihydroxyacridone, T-518
 1,3,6-Trihydroxyacridone, T-519
 1,3,8-Trihydroxyacridone, T-520
 3,4,5-Trihydroxyanthranilic acid, A-922
 1,2,3-Trihydroxyaporphine, T-521
 1,2,8-Trihydroxyaporphine, T-522
 1,2,9-Trihydroxyaporphine, T-523
 1,2,10-Trihydroxyaporphine, T-524
 1,2,11-Trihydroxyaporphine, T-525
 α ,3,4-Trihydroxybenzenepropanoic acid, D-648
 2,6,7-Trihydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, T-526
 2,7,8-Trihydroxy-2*H*-1,4-benzoxazin-3(4*H*)-one, T-527
 2,3,5-Trihydroxybenzylamine, T-528
 1,2,7-Trihydroxy-3,5-bis(hydroxymethyl)-1*H*-pyrrolizidine, T-529
 1,3,5-Trihydroxy-2,4-bis(3-methyl-2-butenyl)-9(10*H*)-acridinone, T-546
 1,3,8-Trihydroxy-2,7-bis(3-methyl-2-butenyl)-9(10*H*)-acridinone, T-548
 1,3,5-Trihydroxy-2,8-bis(3-methyl-2-butenyl)-9*H*-acridin-10-one, T-547
 5-(1,3,4-Trihydroxybutyl)hyacinthacine A₁, T-553
 3-(2,3,4-Trihydroxybutyl)indole, I-116
 1-[5-(2,3,4-Trihydroxybutyl)pyrazinyl]-1,2,3,4-butanetetrol, T-250
 4,5,10-Trihydroxycanthin-6-one, T-531
 1,10,11-Trihydroxycanthin-6-one, T-530
 1,6,7-Trihydroxy-9*H*-carbazole-3-carboxaldehyde, T-532
 2,6,8-Trihydroxy-9*H*-carbazole-3-carboxaldehyde, T-533
 14,16,20-Trihydroxyceve-4-en-3-one, T-534
 2,4,5-Trihydroxycinnamic acid, T-577
 3,4,5-Trihydroxycinnamic acid, T-578
 6,7,8-Trihydroxy-3,11,13-clerodatrien-15,16-olide, T-535
 1,4,5-Trihydroxy-2-cyclopentene-1-carboxamide, T-536
 1,2,3-Trihydroxydibenz[*cd*,*f*]indol-4(5*H*)-one, T-537
 1,2,6-Trihydroxydibenz[*cd*,*f*]indol-4(5*H*)-one, T-538
 1,2,7-Trihydroxydibenz[*cd*,*f*]indol-4(5*H*)-one, T-539
 1,2,9-Trihydroxydibenz[*cd*,*f*]indol-4(5*H*)-one, T-540

- 2,10,13-Trihydroxydibenzo[*a,g*]quinolizinium(1+), T-541
- 1,6,9-Trihydroxydihydro- β -agarofuran, T-542
- 1,3,6-Trihydroxy-2,5-dimethoxy-10-methylacridone, P-206
- 1,3,6-Trihydroxy-4,5-dimethoxy-10-methylacridone, P-207
- 1,4,5-Trihydroxy-3,6-dimethoxy-10-methylacridone, P-207
- 3,4,6-Trihydroxy-1,2-dimethylcarbazole, T-543
- 1,6,8-Trihydroxy-3,4-dimethylisoquinoline, T-544
- 7,18,19-Trihydroxy-16,18-dimethyl-10-phenyl[11]-cytochalasa-6(12),13-diene-1,21-dione, T-545
- 1,3,5-Trihydroxy-2,4-diprenylacridone, T-546
- 1,3,5-Trihydroxy-2,8-diprenylacridone, T-547
- 1,3,8-Trihydroxy-2,7-diprenylacridone, T-548
- 4,5,6-Trihydroxyfuro[2,3-*b*]quinoline, F-218
- 4,5,7-Trihydroxyfuro[2,3-*b*]quinoline, F-219
- 4,5,8-Trihydroxyfuro[2,3-*b*]quinoline, F-220
- 4,6,7-Trihydroxyfuro[2,3-*b*]quinoline, F-221
- 4,6,8-Trihydroxyfuro[2,3-*b*]quinoline, F-222
- 4,7,8-Trihydroxyfuro[2,3-*b*]quinoline, F-223
- 5,6,7-Trihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-215
- 6,7,8-Trihydroxyfuro[2,3-*b*]quinolin-4(9*H*)-one, F-217
- 7,11,14-Trihydroxyhetisan-2,13-dione, T-549
- 11,13,15-Trihydroxyhetisan-2-one, H-190
- 6,11,13-Trihydroxyhetisan-2-one, T-552
- 6,9,13-Trihydroxyhetisan-2-one, T-551
- 1,2,6-Trihydroxyhetisan-13-one, T-550
- 1,2,7-Trihydroxy-5-(3-hydroxybutyl)-3-(hydroxymethyl)-1*H*-pyrrolizidine, T-553
- 1,3,5-Trihydroxy-2-(2-hydroxy-3-methyl-3-butenyl)-10-methylacridone, T-580
- 2,3,5-Trihydroxy-4-(hydroxymethyl)-6-methylpyridine, H-608
- 1,2,7-Trihydroxy-3-hydroxymethyl-5-methyl-1*H*-pyrrolizidine, T-554
- 3,4,5-Trihydroxy-2-hydroxymethylpiperidine, H-616
- 1,2,7-Trihydroxy-3-(hydroxymethyl)-1*H*-pyrrolizidine, T-555
- 1,10,11-Trihydroxy-6*H*-indolo[3,2,1-*de*][1,5]-naphthyridin-6-one, T-530
- 4,8,9-Trihydroxyindolo[2,1-*b*]quinazoline-6,12-dione, O-98
- 5,6,7-Trihydroxyisoquinoline, I-307
- 1,6,7-Trihydroxyisoquinoline, I-305
- 1,6,8-Trihydroxyisoquinoline, I-306
- 5,6,7-Trihydroxy-1(2*H*)-isoquinolinone, I-304
- 1,3,6-Trihydroxyjervanin-12-en-11-one, V-74
- 3,5,15-Trihydroxy-6(17),12-lathyradien-14-one, T-556
- 3,4,13-Trihydroxylupanine, T-557
- 3,8,13-Trihydroxylupanine, T-558
- 2,10,11-Trihydroxy-1-methoxyaporphine, T-248
- 1,3,6-Trihydroxy-5-methoxy-10-methylacridone, T-243
- 1,3,5-Trihydroxy-4-methoxy-10-methylacridone, T-242
- 4,9,10-Trihydroxy-2-methoxy-3-methylene-6-(1-propenyl)-2-azaspiro[4.5]dec-7-en-1-one, S-463
- 1,3,6-Trihydroxy-5-methoxy-10-methyl-4-(3-methyl-2-butenyl)-9(10*H*)-acridinone, P-616
- 1,3,6-Trihydroxy-5-methoxy-10-methyl-4-*C*-prenylacridone, P-616
- 1,3,5-Trihydroxy-4-methoxy-10-methyl-2-prenylacridone, T-270
- 1,3,6-Trihydroxy-5-methoxy-10-methyl-2-prenylacridone, T-271
- 4',5,7-Trihydroxy-6-methoxy-8-(1-methyl-2-pyrrolidyl)flavone, P-389
- 2,9,10-Trihydroxy-3-methoxytetrahydroprotoberberine, D-570
- 1,3,5-Trihydroxy-10-methylacridone, T-518
- 1,3,8-Trihydroxy-10-methylacridone, T-520
- 5,6,8-Trihydroxy-1-methyl-4-azafloren-9-one, T-564
- 6,7,8-Trihydroxy-1-methyl-4-azaflorenone, T-563
- α ,3,4-Trihydroxy- α -methylbenzenepropanoic acid, D-647
- 1,3,5-Trihydroxy-9-methyl-2,8-bis(3-methyl-2-butenyl)-9*H*-acridin-10-one, T-547
- 1,3,5-Trihydroxy-2-(3-methyl-2-butenyl)-9(10*H*)-acridinone, T-580
- 1,3,5-Trihydroxy-4-(3-methyl-2-butenyl)-9(10*H*)-acridinone, T-581
- 7-(1,2,3-Trihydroxy-3-methylbutyl)-1*H*-indole, I-123
- 1,6,7-Trihydroxy-3-methylcarbazole, T-559
- 1,3,5-Trihydroxy-10-methyl-2,4-diprenylacridone, T-546
- 1,3,5-Trihydroxy-9-methyl-2,8-diprenylacridone, T-547
- 3,6,7-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-560
- 6,7,8-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-563
- 6,8,9-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-564
- 7,8,9-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-565
- 3,7,8-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-561
- 3,8,9-Trihydroxy-4-methyl-5*H*-indeno[1,2-*b*]pyridin-5-one, T-562
- 1,6,8-Trihydroxy-3-methylisoquinoline, T-566
- 3,4,5-Trihydroxy-2-methylpiperidine, M-517
- 3,4,5-Trihydroxy-*N*-methylpiperidine, P-458
- 1,3,5-Trihydroxy-10-methyl-2-prenylacridone, T-580
- 2,3,5-Trihydroxy-6-methylpyridine, T-567
- 3,4,6-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-568
- 3,4,8-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-569
- 3,4,9-Trihydroxy-10-nitro-1-phenanthrenecarboxylic acid, T-570
- 1,2,3-Trihydroxynortropane, A-1590
- 1,2,4-Trihydroxynortropane, A-1591
- 1,2,7-Trihydroxynortropane, A-1592
- 1,3,4-Trihydroxynortropane, A-1593
- 2,3,4-Trihydroxynortropane, A-1594
- 2,3,6-Trihydroxynortropane, A-1595
- 3,6,7-Trihydroxynortropane, A-1596
- 1,2,8-Trihydroxyoctahydroindolizine, S-650
- 9,10,11-Trihydroxy-2-oxo-1,7-dioxo-4-azaspir-[5.5]undecane-3-propanamide, C-431
- 1,2,3-Trihydroxy-5-oxonoraporphine, T-571
- 4',5,7-Trihydroxy-6-(5-oxo-2-pyrrolidinyl)flavone, T-572
- 4',5,7-Trihydroxy-8-(5-oxo-2-pyrrolidinyl)flavone, T-573
- 1-(3,4,5-Trihydroxypentanyloxy)- β -carboline, C-138
- 2,3,9-Trihydroxy-1-phenazinecarboxylic acid, T-574
- 2,3,7-Trihydroxy-1,6-phenazinedicarboxylic acid, T-575
- 1,2,6-Trihydroxyphenazine, P-325
- 2,3,7-Trihydroxyphenazine, P-326
- 3,4,5-Trihydroxyphenethylamine, T-576
- 3-(2,4,5-Trihydroxyphenyl)acrylic acid, T-577
- 3-(3,4,5-Trihydroxyphenyl)acrylic acid, T-578
- 2-(3,4,5-Trihydroxyphenyl)ethylamine, T-576
- 3-(2,4,5-Trihydroxyphenyl)-2-propenoic acid, T-577
- 3-(3,4,5-Trihydroxyphenyl)-2-propenoic acid, T-578
- 3,4,5-Trihydroxypipercolic acid, T-579
- 3,4,5-Trihydroxy-2-piperidinedicarboxylic acid, T-579
- 3,4,5-Trihydroxy-2-piperidinemethanol, H-616
- 3,4,5-Trihydroxypiperidine, P-458
- 1,3,5-Trihydroxy-2-prenylacridone, T-580
- 1,3,5-Trihydroxy-4-prenylacridone, T-581
- 3,4,7-Trihydroxy-8-prenyl-2-quinolinecarboxylic acid, T-584
- 2,4,7-Trihydroxy-3-prenylquinoline, T-582
- 2,4,8-Trihydroxy-3-prenylquinoline, T-583
- 3-(1,2,3-Trihydroxypropyl)indole, I-135
- 7-(1,2,3-Trihydroxypropyl)pterin, A-924
- 2,4,7-Trihydroxypteridine, P-763
- 2,6,8-Trihydroxypurine, U-45
- 3,4,5-Trihydroxy-2-[[2-(3-pyridinylcarbonyl)amino]benzoyl]amino]benzoic acid, T-585
- 3,5,7-Trihydroxypyrimido[5,4-*c*]-*s*-triazine, D-498
- 4,5,8-Trihydroxy-2-quinolinecarboxylic acid, T-586
- 4,7,8-Trihydroxyquinoline, Q-48
- 2,3,4-Trihydroxyquinoline, Q-45
- 2,4,6-Trihydroxyquinoline, Q-46
- 2,4,8-Trihydroxyquinoline, Q-47
- 4,6,8-Trihydroxy-2(1*H*)-quinolinone, Q-42
- 3,5,8-Trihydroxy-4(1*H*)-quinolinone, Q-44
- 4,7,8-Trihydroxy-2(1*H*)-quinolinone, Q-43
- 3,4,7-Trihydroxy-2(1*H*)-quinolinone, Q-40
- 3,4,8-Trihydroxy-2(1*H*)-quinolinone, Q-41
- 1,3,13-Trihydroxy-11,12-secocevane-6,12-dione, S-277
- 3,16,23-Trihydroxy-16,28-secosolanidan-4-one, S-349
- 3,6,7-Trihydroxytropene, M-392
- 3,5,6-Trihydroxy-6-undecyl-2-piperidinedicarboxylic acid, T-587
- 3,13,23-Trihydroxyveratra-7,12(14)-dien-6-one, T-588
- 1,1,3-Tri-(1*H*-indol-3-yl)butane, T-589
- 1,1,1-Tri-(1*H*-indol-3-yl)ethane, T-590
- Tri-1*H*-indol-3-ylmethane, T-591
- N,N',N''*-Triisopentylguanidine, T-623
- Trikendiol, T-592
- Triketramine, T-593
- Triketrin A, T-594
- Triketrin B, T-595
- Trilobamine, O-1
- Trilobine, I-339
- Trilobinine, T-247
- Trilupine, L-302
- 1,2,3-Trimethoxyacridone, T-516
- 1,2,3-Trimethoxyaporphine, T-521
- 1,2,9-Trimethoxyaporphine, T-523
- 1,2,10-Trimethoxyaporphine, T-524
- 1,2,11-Trimethoxyaporphine, T-525
- 3,4,5-Trimethoxybenzenethanamine, M-219
- 4,10,12-Trimethoxy-8*H*-benzo[*g*]benzodioxolo[6,5,4-*de*]quinolin-8-one, D-949
- 4,9,10-Trimethoxy-8*H*-benzo[*g*]-1,3-benzodioxolo[6,5,4-*de*]quinolin-8-one, K-104
- 1,2,3-Trimethoxy-7*H*-benzo[*de*][1,3]benzodioxolo[5,6-*g*]quinolin-7-one, O-195
- 1,2,13-Trimethoxy[1,3]benzodioxolo[5,6-*c*]phenanthridine, C-356
- 2,3,13-Trimethoxy[1,3]benzodioxolo[5,6-*c*]phenanthridine, R-87
- 8,9,9-Trimethoxy-9*H*-benzo[*de*][1,6]naphthyridine, M-241
- 6,9,10-Trimethoxy-12*H*-[1]benzoxepino[2,3,4-*ij*]isoquinolin-12-one, O-171
- 6,8,9-Trimethoxy-12*H*-[1]benzoxepino[2,3,4-*ij*]isoquinolin-12-one, S-73
- O*-(3,4,5-Trimethoxybenzoyl)-(Z)-burnamine, B-420
- 3 α -(3,4,5-Trimethoxybenzoyloxy)nortropane, C-633
- 3-(3,4,5-Trimethoxybenzoyloxy)nortropan-6-ol, A-1584
- 7-(3,4,5-Trimethoxybenzoyloxy)-3-(3,4,5-trimethoxycinnamoyloxy)tropan-6-ol, M-392
- 3-(3,4,5-Trimethoxybenzoyloxy)tropan-6-ol, M-391
- (3,4,5-Trimethoxybenzoyl)quebrachidine, V-113
- 3,4,5-Trimethoxybenzoylsereadamine, T-288
- O*-(3,4,5-Trimethoxybenzoyl)vincamajine, V-113
- 2,9,10-Trimethoxyberbin-1-ol, C-174
- 3,9,10-Trimethoxyberbin-2-ol, I-210
- 3,4,5-Trimethoxycinnamic acid isobutylamide, T-578
- 3,4,5-Trimethoxycinnamic acid pyrrolidide, T-578
- 3,4,5-Trimethoxycinnamic acid, T-578
- 3,4,5-Trimethoxycinnamoyl piperidide, T-578
- (3,4,5-Trimethoxycinnamoyl)-10-methoxyvincamajine, V-113
- N*-(3,4,5-Trimethoxycinnamoyl)- Δ^3 -piperidin-2-one, P-473
- N*-(2,4,5-Trimethoxycinnamoyl)pyrrolidine, T-577
- N*-(3,4,5-Trimethoxycinnamoyl)pyrrolidine, T-578
- 3,4,5-Trimethoxycinnamoylvincamajine, V-113

- 1,2,3-Trimethoxydibenz[cd,f]indol-4(5*H*)-one, T-537
- 1,4,5-Trimethoxy-7*H*-dibenzo[de,g]quinolin-7-one, A-1449
- 5,6,9-Trimethoxy-7*H*-dibenzo[de,h]quinolin-7-one, M-199
- 1,2,4-Trimethoxy-7*H*-dibenzo[de,g]quinolin-7-one, S-470
- 5,6,7-Trimethoxydictamine, F-215
- 5,7,8-Trimethoxydictamine, F-216
- 6,7,8-Trimethoxydictamine, F-217
- β,3,4-Trimethoxy-*N,N*-dimethylbenzeneethanamine, M-18
- 3,4,6-Trimethoxy-1,2-dimethylcarbazole, T-543
- 4,6,7-Trimethoxy-*N,N*-dimethylidibenz[*b*,*f*]oxepin-1-ethanamine, S-198
- 6,6',12'-Trimethoxy-2,2'-dimethyloxyacanthan-7-ol, L-175
- 6,6',7-Trimethoxy-2,2'-dimethyloxyacanthan-12'-ol, R-46
- 3,4,8-Trimethoxy-*N,N*-dimethyl-1-phenanthreneethanamine, F-64
- 2,3,4-Trimethoxy-*N,N*-dimethyl-1-phenanthreneethanamine, M-238
- 1,2,3-Trimethoxy-*N,N*-dimethylphenanthro[2,3-*d*][1,3]dioxole-4-ethanamine, T-328
- 5,8,9-Trimethoxy-2,2-dimethyl-2*H*-pyrano[2,3-*b*]quinoline, M-251
- 1,2,3-Trimethoxy-4,5-dioxo-6*a*,7-dehydroaporphine, O-139
- 4,9,10-Trimethoxy-1,3-dioxolo[4,5-*g*]furo[2,3-*b*]quinoline, F-214
- 4,6,8-Trimethoxyfuro[2,3-*b*]quinoline, F-222
- 4,5,6-Trimethoxyfuro[2,3-*b*]quinoline, F-218
- 4,5,7-Trimethoxyfuro[2,3-*b*]quinoline, F-219
- 4,6,7-Trimethoxyfuro[2,3-*b*]quinoline, F-221
- 4,7,8-Trimethoxyfuro[2,3-*b*]quinoline, F-223
- ar*-Trimethoxygramine, D-734
- 4,5,6-Trimethoxyindeno[1,2,3-*ij*]isoquinolin-9-ol, N-304
- 2,3,10-Trimethoxyindolo[2',3':3,4]pyrido[2,1-*b*]quinazolin-5(13*H*)-one, E-307
- 5,6,7-Trimethoxyisoquinoline, I-307
- 1,2,3-Trimethoxy-10-methylacridone, T-516
- 1,3,4-Trimethoxy-10-methylacridone, T-517
- 1,3,5-Trimethoxy-10-methylacridone, T-518
- β,3,4-Trimethoxy-*N*-methylbenzeneethanamine, M-18
- 8,8,9-Trimethoxy-5-methylbenz[cd]isoindolo[2,1-*a*]indol-1(8*H*)-one, A-1088
- 4,10,11-Trimethoxy-7-methyl-5*H*-benzo[*g*,1,3-benzodioxolo[6,5,4-*de*]quinoline, O-27
- 1,2,4-Trimethoxy-12-methyl[1,3]benzodioxolo[5,6-*c*]phenanthridinium(1+), C-356
- 5,9,10-Trimethoxy-12-methylbenzo[*c*]-1,3-dioxolo[4,5-*f*]phenanthridinium(1+), S-60
- 4,5,8-Trimethoxy-3-(3-methyl-2-butenyl)-2(1*H*)-quinolinone, D-83
- 1,2,8-Trimethoxy-6-methyl-9*H*-carbazole, T-596
- 4,6,7-Trimethoxy-*N*-methylidibenz[*b*,*f*]oxepin-1-ethanamine, S-198
- 6,7,11-Trimethoxy-5-methyl-1,3-dioxolo[4,5-*b*]acridin-10(5*H*)-one, T-56
- 8,9,10-Trimethoxy-1,2-methylenedioxyaporphine, L-158
- 3,9,10-Trimethoxy-1,2-methylenedioxyaporphine, O-27
- 1,10,11-Trimethoxy-2,3-methylenedioxyaporphine, O-25
- 1,2,3-Trimethoxy-9,10-methylenedioxyaporphine, P-369
- 1,2,3-Trimethoxy-9,10-methylenedioxydehydroaporphine, P-369
- 4,5,8-Trimethoxy-6,7-methylenedioxyfuro[2,3-*b*]quinoline, F-214
- 1,9,10-Trimethoxy-2,3-methylenedioxyaporphine, B-7
- 1,2,3-Trimethoxy-10,11-methylenedioxyaporphine, L-10
- 3,9,10-Trimethoxy-1,2-methylenedioxyaporphine, O-27
- 1,2,3-Trimethoxy-9,10-methylenedioxyaporphine, P-369
- 3,10,11-Trimethoxy-1,2-methylenedioxyaporphine, P-209
- 3,9,11-Trimethoxy-1,2-methylenedioxyaporphine, D-949
- 1,2,3-Trimethoxy-9,10-methylenedioxyaporphine, O-195
- 3,9,10-Trimethoxy-1,2-methylenedioxyaporphine, T-315
- 2,8,9-Trimethoxy-3,4-methylenedioxyapavine, T-330
- 4,7,8-Trimethoxy-2,3-methylenedioxyquinoline, Q-39
- ar*-Trimethoxy-1-methylisoquinoline, T-597
- 5,6,7-Trimethoxy-2-methyl-1(2*H*)-isoquinolinone, I-304
- 1,5,6-Trimethoxy-10-methyl-2,3-methylenedioxyacridone, T-56
- 2,3,10-Trimethoxy-5-methyl-7,8-methylenedioxybenzo[*c*]phenanthridinium(1+), S-60
- 4,8,9-Trimethoxy-*N*-methyl-2,3-methylenedioxyapavine, T-598
- 1,2,3-Trimethoxy-*N*-methylphenanthro[2,3-*d*][1,3]dioxole-4-ethanamine, T-328
- 4,6,8-Trimethoxy-1-methyl-2(1*H*)-quinolinone, Q-42
- 4,7,8-Trimethoxy-1-methyl-2(1*H*)-quinolinone, Q-43
- 1,2,3-Trimethoxyaporphine, T-521
- 1,2,11-Trimethoxyaporphine, T-525
- 1,2,3-Trimethoxy-5-oxonoraporphine, T-571
- 2,3,6-Trimethoxyphenanthroindolizidine, A-1330
- 3,4,6-Trimethoxyphenanthroindolizidine, I-342
- 3,6,7-Trimethoxyphenanthroindolizidine, T-696
- 1,2,6-Trimethoxyphenazine, P-325
- 3,4,5-Trimethoxyphenethylamine, M-219
- N*-(3,4,5-Trimethoxyphenethyl)maleimide, M-222
- N*-(3,4,5-Trimethoxyphenethyl)malimide, M-222
- N*-(3,4,5-Trimethoxyphenethyl)succinimide, M-222
- 1,2,4-Trimethoxy-3*H*-phenoazin-3-one, T-599
- N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]acetamide, M-219
- N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]alanine, M-219
- N*-[2-(3,4,5-Trimethoxyphenyl)ethyl]glycine, M-219
- 1-[2-(3,4,5-Trimethoxyphenyl)ethyl]-1*H*-pyrrole-2-carboxylic acid, P-298
- 1-[2-(3,4,5-Trimethoxyphenyl)ethyl]-2,5-pyrrolidinedione, M-222
- 3-(3,4,5-Trimethoxyphenyl)-2-propenoic acid, T-578
- 4,7,8-Trimethoxy-3-prenyl-2(1*H*)-quinolinone, P-625
- 4,5,8-Trimethoxy-3-prenyl-2(1*H*)-quinolinone, D-83
- 4,6,8-Trimethoxy-2(1*H*)-quinolinone, Q-42
- 4,7,8-Trimethoxy-2(1*H*)-quinolinone, Q-43
- 3,4,8-Trimethoxy-2(1*H*)-quinolinone, Q-41
- 5,10,11-Trimethoxystrictamine, S-576
- Trimethylamine oxide, T-601
- Trimethylamine, T-600
- (2-Trimethylammonioethyl)phosphonic acid betaine, T-602
- 1-Trimethylammonio-3-(3-indolyl)propionate, H-764
- N,N,N*-Trimethylbenzeneethanaminium(1+), P-342
- 1,7,8-Trimethylbenzo[*g*]pteridine-2,4(1*H*,3*H*)-dione, L-285
- Tri-*N*-methyl-γ-butyrobetaine, A-126
- 5-[(2,4,4-Trimethyl-1-cyclohexen-1-yl)methyl]-1(5*H*)-phenazinone, L-76
- 4,4,14-Trimethyl-9,19-cyclopregnane-3,20-diamine, C-902
- 2,4,6-Trimethyl-2,4-decadienamide, T-603
- 2,4,6-Trimethyl-2,4-decadienoic acid, T-603
- N,N,N*-Trimethyl-1,2-dehydrocadaverine, P-232
- Trimethyldiazene-carboxamide 2-oxide, L-368
- 3,7,11-Trimethyl-2,6,10-dodecatrien-1-amine, F-12
- 3-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1*H*-indole, T-605
- 2-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1*H*-indole, T-604
- Tri(α-methylene-γ-butyrolactonyl)amine, N-248
- N,N,N*-Trimethylethanaminium(1+), T-615
- 1,1,5-Trimethylfuro[3,4-*c*]pyridine-3,4(1*H*,5*H*)-dione, C-285
- 1,3,7-Trimethylguanidine, D-748
- N*²,*N*²,7-Trimethylguanidine, M-469
- 2-(2,4,6-Trimethyl-1-heptenyl)-4-quinazolinol, T-606
- 2-(2,4,6-Trimethyl-1-heptenyl)-4(1*H*)-quinazolinone, T-606
- 4,4,14-Trimethyl-*B*-(9*a*)-homo-19-norpregna-9(11),9*a*-diene-3,20-diamine, B-455
- 1,2,5-Trimethyl-1*H*-imidazole, D-749
- 1,4,5-Trimethyl-1*H*-imidazole, D-750
- 2,4,5-Trimethyl-1*H*-imidazole, T-607
- 1,3,7-Trimethylisoguanidine, T-608
- N*¹,*N*³,*O*⁷-Trimethylisoxanthopterin, A-751
- N,N,N*-Trimethylmethanaminium(1+), T-280
- N,N,N*'-Trimethyl-*N*'-(4-methoxycinnamoyl)putrescine, C-456
- 4,4,14-Trimethyl-3-(methylamino)-9,19-cyclopregna-7,17(20)-dien-16-one, C-844
- N,N,N,N'*-Trimethyl-*N*'-(3-methyl-2,4-dodecadienoyl)spermidine, T-609
- N,N,N,N'*-Trimethyl-*N*'-(3-methyl-dodecanoyl)spermidine, T-609
- N,N,N,N'*-Trimethyl-*N*'-(3-methyl-2-dodecenoyl)spermidine, T-609
- N,N,N,N'*-Trimethyl-*N*'-(5-methyl-2,4-tetradecadienoyl)spermidine, T-610
- N,N,N,N'*-Trimethyl-*N*'-(5-methyl-2,4-tetradecadienoyl)spermidine, T-611
- N,N,N,N*-Trimethyl-3-(methylthio)-1-propanaminium(1+), T-612
- Trimethyl[3-(methylthio)propyl]ammonium(1+), T-612
- 1,3,9-Trimethyl-8-nitrosoisoxanthine, D-523
- 1,9,13-Trimethyl-8-nonyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-384
- Trimethylloxamine, T-601
- 6,8,11-Trimethyl-2*H*,4*H*-oxazolo[5,4,3-*ij*]pyrido[3,2-*g*]quinoline-4,10(11*H*)-dione, N-346
- 2-(2,6,6-Trimethyl-4-oxo-2-cyclohexen-1-yl)acetamide, G-88
- 1,2,5-Trimethyl-4-(1-oxo-3-phenyl-2-propenyl)piperazine, D-769
- Trimethylphenethylammonium(1+), P-342
- N,N,N*-Trimethyl-2-phosphonoethanaminium hydroxide, inner salt, T-602
- 1,3,6-Trimethyl-2,5-piperazinedione, D-769
- 1,2,6-Trimethylpiperidine, D-770
- 5-(1,1,2-Trimethylpropenylperoxy)-2-pyrrolidione, C-801
- 1,*N*⁸,*N*⁸-Trimethylpseudozoanthoxanthin A, P-737
- Trimethylpyrazine, T-613
- 2,2,6-Trimethyl-4-(4-quinoliny)-3-azabicyclo[3.3.1]non-6-ene, T-614
- N,O,O*-Trimethylspariflorine, T-524
- N,O,O*-Trimethylstaphavanine, S-545
- N,N,N*-Trimethyl-2-(sulfoxy)ethanaminium hydroxide inner salt, C-419
- 2,3,3-Trimethyl-2,3,4,5-tetrahydrofuran[3,2-*c*]quinolin-4-one, O-82
- 2,4,5-Trimethyl-3-thiazoline, D-526
- 2,7,16-Trimethyl-2,7,11-triaza-13,15-pentacosadien-12-one, T-610
- 2,7,16-Trimethyl-2,7,11-triaza-14-pentacosen-12-one, T-611
- 2,7,14-Trimethyl-2,7,11-triaza-13,15-tricosadien-12-one, T-609
- 2,7,14-Trimethyl-2,7,11-triaza-12-tricosanone, T-609
- 2,7,14-Trimethyl-2,7,11-triaza-13-tricosen-12-one, T-609
- 1,9,13-Trimethyl-8-tridecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-387
- N,N,N*-Trimethyltryptamine, D-785
- Trimethyltryptophan betaine, H-764
- 1,9,13-Trimethyl-8-undecyl-1,5,9,13-tetraazacycloheptadecan-6-one, B-383
- O*²,1,9-Trimethyluric acid, L-165
- Trimethylvinylammonium(1+), T-615
- 1,3,7-Trimethylxanthine, C-15
- Trinervine, T-616
- 1,3,6-Tri-*O*-nicotinoyl-α-*D*-glucopyranose, A-458
- Triniifoline, Z-1
- Triophamine, T-617

- Triostin, T-618
 3,16,18-Trioxo-1,4,14-conatrienine, H-314
N,N',N''-(2,6,10-Trioxo-1,5,9-trioxacyclododecane-3,7,11-triyl)tris[2,3-dihydroxybenzamide], E-86
 Triox, T-601
 Tripfordine A, P-253
 Tripfordine B, P-252
 Tripfordine C, P-249
 1,2,3-Triphenyl-1*H*-pyrrole, D-817
 Triphyopeltine, D-793
 Triphyophylline, D-793
 TripterospERMUMCIN A, D-204
 Tripterregeline A, H-249
 Tripterregeline B, H-249
 Tripterregeline C, H-249
 Triptofordinine A1, H-249
 Triptofordinine A2, H-249
 Triptogelin A10, P-216
 Triptogelin A5, P-216
 Triptogelin A6, P-216
 Triptogelin A7, P-216
 Triptogelin A9, P-216
 Triptogelin B2, T-258
 Triptogelin C2, T-256
 Triptogelin C4, T-256
 Triptogelin F1, T-542
 Triptonine A, P-253
 Triptonine A, T-619
 Triptonine B, P-253
 Triptonine B, T-620
 Triptotin J, T-621
 Tris(3-aminopropyl)amine, T-622
N,N',N''-Tris(*E-p*-coumaroyl)spermidine, S-398
N¹,N⁵,N¹⁰-Tris-*trans-p*-coumaroylspermine, S-399
 Tris(4-ethenylphenyl 4-hydroxy-3-nitrosobenzoate-*N,O*)ferrate(1-), F-45
 3,3',11-Tris-*O*-(3-hydroxybutanoyl)hamayne, H-5
N,N',N''-Tris(4-hydroxycinnamoyl)spermidine, S-398
 Tris(*N*-hydroxy-*N*-methylmethanethioamidato-*O,S*)iron, F-96
 3,7,8-Tris(4-hydroxyphenyl)-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-1-one, L-284
 Trisindoline, T-88
N,N',N''-Tris(3-methylbutyl)guanidine, T-623
 Tris(1-methyl-3-indolyl)methane, T-591
 Trisoxazoline, C-919
 Trisphaeridine, M-448
 Trisphaerine, H-294
 Trispheridine, M-448
 Trispherine, H-294
 Trisulcusine, T-624
 Triticone A, T-625
 Triticone B, T-625
 Triticone C, T-625
 Triticone D, T-625
 Triticone E, T-626
 Triticone F, T-626
 Tritopine, L-68
 1-Tritylpyrrolidine, P-949
 Triumferol, I-348
 Triumfettoside A, A-756
 Triuret, T-627
 Trivalvone, T-628
 Trodusquemine, S-485
 Trolline, T-629
 Tronocarpine, T-630
 Tronoharine, T-631
 Tropacocaine, M-393
 Tropane 3,6-diol 3-phenylacetate 6-acetate, M-391
 Tropane 3,6-diol 3-(3,4,5-trimethoxycinnamate) 6-benzoate, M-391
 Tropane 3,6-diol 3-phenylacetate 6-propanoate, M-391
 Tropane 3,6-diol 3-phenylacetate, M-391
 2,3-Tropanediol, A-1581
 2,6-Tropanediol, A-1582
 3,6-Tropanediol, M-391
 Tropane-3,6,7-triol 3-phenylacetate, M-392
 Tropane-3,6,7-triol 3-(3,4,5-trimethoxybenzoate), M-392
 3,6,7-Tropanetriol, M-392
 ψ -3-Tropanol, M-393
 3-Tropanol, M-393
 1-Tropanol, P-399
 3-Tropanone, M-394
 Tropan-6-yl benzoate-3-yl-3,4,5-trimethoxycinnamate, M-391
 Tropanyl *cis*-cinnamate, M-393
 Tropanyl *trans*-cinnamate, M-393
 Tropan-3-yl ferulate, M-393
 Tropanyl furoate, M-393
 Tropanyl 3-hydroxy-4-methoxycinnamate, M-393
 Tropan-3-yl 3-hydroxyphenylacetate, M-393
 Tropan-3-yl isovalerate, M-393
 Tropanyl phenylacetate, M-393
 Tropan-3 α -yl phenyllactate, L-216
 Tropanyl senecioidate, M-393
 Tropan-3 α -yl 3,4,5-trimethoxybenzoate, C-633
 Tropan-3-yl-*O*-propionyl tropate, T-632
 Tropeine, M-393
 Trophicardyl, I-148
 Tropigenin, A-1597
 Tropigline, T-404
 Tropine acetate, M-393
 Tropine atropate, A-1359
 Tropine benzoate, M-393
 Tropine butyrate, M-393
 Tropine isobutyrate, M-393
 Tropine isovalerate, M-393
 Tropine nonanoate, M-393
 Tropine propionate, M-393
 Tropine tiglate, T-404
 Tropine 3,4,5-trimethoxycinnamate, M-393
 Tropine tropate, T-632
 Tropine, M-393
 ψ -Tropine, M-393
 Tropinone, M-394
 Tropinox, T-632
 Trpn, T-622
 Trp, T-640
 Trunkamide A, T-633
epi-Truxilline, T-634
neo-Truxilline, T-635
peri-Truxilline, T-634
 α -Truxilline, T-634
 β -Truxilline, T-635
 δ -Truxilline, T-635
 ϵ -Truxilline, T-634
 γ -Truxilline, T-634
 μ -Truxilline, T-635
 ω -Truxilline, T-635
 ξ -Truxilline, T-635
 β -Truxinic acid bis(2-carbomethoxypropyl) ester, T-635
 Trypanothione disulfide, T-636
 Trypanothione, T-636
 Trypargimine, T-637
 Trypargine, T-637
 Tryprostatin A, T-638
 Tryprostatin B, T-638
 Tryptamine, T-639
 Tryptanthrine, I-108
 Tryptophan betaine, H-764
 Tryptophan tryptophylquinone, T-641
 Tryptophanbetaxanthin, T-642
 Tryptophanol, A-830
 Tryptophan, T-640
 Tryptophol galactoside, I-96
 Tryptophol glucoside, I-96
 Tryptophol, I-96
 Tryptoquialanine A, T-643
 Tryptoquialanine B, T-644
 Tryptoquivaline A, T-645
 Tryptoquivaline B, N-315
 Tryptoquivaline C, T-645
 Tryptoquivaline D, N-315
 Tryptoquivaline E, T-646
 Tryptoquivaline F, T-646
 Tryptoquivaline G, T-647
 Tryptoquivaline H, T-646
 Tryptoquivaline I, T-645
 Tryptoquivaline J, T-646
 Tryptoquivaline L, T-647
 Tryptoquivaline M, N-315
 Tryptoquivaline N, N-315
 Tryptoquivaline, T-645
 Tryptoquivalone, N-315
 TS 885, M-561
 Tscheschamine, T-648
 Tsilanimbine, R-63
 Tsilanine, T-649
 Tsitsikammamine A *N*¹⁸-oxime, T-650
 Tsitsikammamine A, T-650
 Tsitsikammamine B *N*¹⁸-oxime, T-650
 Tsitsikammamine B, T-650
 Tsuduranine, T-524
 Tsukushinamine A, T-651
 Tsukushinamine B, T-651
 Tsukushinamine C, T-651
 TTP, T-366
 TTQ, T-641
 TTX, T-298
 Tu 2480F₁, P-893
 Tu 2480F₂, P-893
 Tu 2480F₃, P-893
 Tu 2480F₄, P-893
 Tubacetine, T-652
 Tubarine, C-420
 Tubasencine, T-653
 Tubastraine, T-654
 Tubastrindole A, T-655
 Tubastrindole B, T-655
 Tubastrindole C, T-655
 Tubastrine, T-656
 Tuberaconitine, M-331
 Tuberactidine, T-657
 Tuberanin, M-331
 Tubercidin 5'- α -D-glucopyranose, T-658
 Tubercidin, T-658
 Tuberin, T-659
 Tubermesaconitine, M-331
 Tubermycin A, T-660
 Tubermycin B, P-319
 Tuberosine B, D-920
 Tuberosine B, T-190
 Tuberosinone B, T-661
 Tuberosinone C, T-662
 Tuberosinone *N*- β -D-glucoside, T-661
 Tuberosinone, T-661
 Tuberosipronine, S-523
 Tuberostemonone, T-663
 Tuberostemonine A, T-664
 Tuberostemonine B, T-664
 Tuberostemonine C, T-664
 Tuberostemonine H, T-664
 Tuberostemonine J, T-664
 Tuberostemonine K, T-664
 Tuberostemonine LG, T-664
 Tuberostemonine L, T-664
 Tuberostemonine M, T-664
 Tuberostemonine N, T-664
 Tuberostemonine *N*-oxide, T-664
 Tuberostemonine, T-664
 Tuberostemoninol, T-665
 Tuberostemonol, T-664
 Tuberostemonone, T-666
 Tuberostemosipronine, T-667
 Tubifolidine, T-668
 Tubifoline, T-668
 Tubingensin A, T-669
 Tubingensin B, T-670
 Tubispacine, C-743
 Tubocurarine chloride, C-420
 Tubocurarine, C-420
 (-)-Tubocurarine, C-420
 Tubocurine, C-420
 Tuboflavine, T-671
 Tubotaiwinal, T-672
 Tubotaiwine *N*⁴-oxide, T-673
 Tubotaiwine, T-673
 Tuboxenine, T-674
 Tubulosine, T-675
 Tubulysin D, T-676
 Tubulysin E, T-676
 Tubulysin F, T-676
 Tubulysin H, T-676
 Tubulysins, T-676

- Tuduranine, T-524
 Tuguaconitine, D-187
 Tuliferoline, T-521
 Tunichrome An1, T-677
 Tunichrome An2, T-677
 Tunichrome An3, T-677
 Tunichrome B1, T-677
 Tunichrome Pm 1, T-678
 Tunichrome Pm 2, T-678
 Tunichrome Pm 3, T-678
 T-Up, B-101
 Turbinamide, O-61
 Turbinatine, T-679
 Turbotoxin A, T-680
 Turbotoxin B, T-680
 Turcamine, T-159
 Turcberine, T-681
 Turcomanidine, P-80
 Turcomanine, P-80
 Turconidine, T-682
 Turcosine, M-339
 Turicine, H-476
 Turkestanine *N*-oxide, T-683
 Turkestanine, T-683
 Turkiyenine, T-684
 Turlosine, M-339
 Turneforcidine, H-233
 (+)-Turneforcidine, H-233
 Turneforcine, H-233
 Turpelline, N-30
 Turraeanthine A, A-1446
 Turraeanthine B, D-482
 Turraparvin D, T-685
 Turrapubesin B, T-686
 Turumiquirensine, T-687
 Tusscapine, N-44
 Tussiglaucin, T-246
 Tussilagic acid, H-231
 Tussilagine, H-231
 Tussilaginine, H-231
 Tussipan, C-555
 Tyformin, G-201
 (10*R*,13*aR*)-Tylocrebrine *N*-oxide, T-688
 (10*S*,13*aR*)-Tylocrebrine *N*-oxide, T-688
 Tylocrebrine, T-688
 Tylohirsuticine, T-689
 Tylohirsutinidine, T-690
 Tylohirsutinine, T-690
 Tyloindicine A, T-278
 Tyloindicine B, T-689
 Tyloindicine C, M-583
 Tyloindicine D, T-278
 Tyloindicine E, T-696
 Tyloindicine F, T-692
 Tyloindicine G, T-693
 Tyloindicine H, T-694
 Tyloindicine I, T-691
 Tyloindicine J, T-691
 Tylophoridicine A, T-696
 Tylophoridicine C, T-696
 Tylophoridicine D, A-1025
 Tylophoridicine E, T-696
 Tylophoridicine F, T-696
 Tylophorine B, A-1330
 Tylophorine *N*-Oxide, T-695
 Tylophorine, T-695
 Tylophorinicine, T-695
 Tylophorinidine, T-696
 Tylophorinine *N*-oxide, T-696
 Tylophorinine, T-696
 Tyraminebetaxanthin, M-647
m-Tyramine, H-678
 Tyramine, H-667
 Tyraminoporphine, T-697
 Tyrian purple, D-325
 Tyrindoleninone, B-341
 Tyrindolinone, B-309
 Tyrindoxol, B-328
 Tyriverdin, T-698
 Tyrosamine, H-667
 Tyroscherin, T-699
 Tyrosine betaine, T-700
 Tyrosinebetaxanthin, P-576
 Tyrosine, T-700
p-Tyrosine, T-700
 Tyrosinyl glucoside, T-700
 Tyrosol carbamate, H-676
 Tyrosol, H-676
 Tyroxindoxyl sulfate, B-328
 Tyr, T-700
 U 106305, A-1299
 U 42126, A-81
 U 43795, A-81
 U 62162, A-1298
 U 27810, B-104
 U 77779, B-229
 U 54703, C-521
 U 11994, D-232
 U 4527, D-765
 U 9361, D-765
 U 54702D, C-521
 U 20207, E-229
 U 47929, F-52
 U 48266, F-33
 U 15965, L-166
 U 24792, L-241
 U 77863, M-513
 U 59760, N-276
 U 59761, N-276
 U 61732, N-276
 U 22662, P-876
 U 24544, P-919
 U 48160, Q-61
 U 19183, S-384
 U 10071, T-658
 Uberine, T-180
 Ubine, A-864
 UCA 1064A, A-1602
 UCA 1064B, A-1602
 UCF1B, M-88
 UCF1C, M-87
 UCH 15A, A-1300
 UCH 15B, A-1300
 UCN 01, S-498
 UCN 02, S-498
 UCS 1025A, A-1301
 UCS 1025B, A-1301
 Udici active substance, P-80
 Ugibohlin, U-1
 UK 1, A-1302
 UK 2, A-1303
 UK 63052, A-1304
 UK 63598, A-1304
 UK 65662, A-1304
 UK 88051, A-1141
 UK 2A, A-1303
 UK 3A, A-1303
 UK 2B, A-1303
 UK 2C, A-1303
 UK 2D, A-1303
 UK 163E, A-1304
 UK 111866, P-90
 Ulapualides, U-2
 Ulbactin A, U-3
 Ulbactin B, U-4
 Ulbactin C, U-5
 Ulbactin D, U-6
 Ulbactin E, U-7
 Uleine *N*-oxide, U-8
 Uleine, U-8
 Ulexine, C-938
 Ulicyclamide, U-9
 Ulithiacyclamide B, U-11
 Ulithiacyclamide E, U-12
 Ulithiacyclamide F, U-13
 Ulithiacyclamide G, U-14
 Ulithiacyclamide, U-10
 Ulongamide A, U-15
 Ulongamide B, U-15
 Ulongamide C, U-16
 Ulongamide D, U-17
 Ulongamide E, U-18
 Ulongamide F, U-19
 Ulosantoin, U-20
 Ulosin A, S-613
 Ultraquinine, Q-20
 Uluganine, L-212
 Ulupyrinone, L-272
 Umanopterin, A-923
 Umbellamine, U-21
 Umbellatine, B-101
 Umbellatine, B-264
 Umbellinic acid, M-240
 Umbrofine, A-92
 Umbrophine, A-92
 Umbrosine, M-325
 Umprel, E-157
 Umycin A, A-853
 Umycin B, G-180
 Umycin C, A-853
 Unamycin B, S-53
 Uncaramine, U-22
 Uncarine A, M-661
 Uncarine B, M-661
 Uncarine C, M-661
 Uncarine D, M-661
 Uncarine E, M-661
 Uncarine F *N*-oxide, M-661
 Uncarine F, M-661
 Uncialamycin, U-23
 Uncinatine, A-1533
 Uncinine, U-24
 2,4-Undecadiene-8,10-diyne-carboxy-*N*-isopentylamide, U-26
 2,4-Undecadiene-8,10-diyneic acid 2,3-dehydropiperidide, U-26
 2,4-Undecadiene-7,9-diyneic acid isobutylamide, U-25
 2,4-Undecadiene-8,10-diyneic acid isobutylamide, U-26
 2,4-Undecadiene-8,10-diyneic acid isopentylamide, U-26
 3,5-Undecadiene-8,10-diyneic acid 2-phenylethylamide, U-27
 2,4-Undecadiene-8,10-diyneic acid piperidide, U-26
 3,5-Undecadiene-8,10-diyneic acid, U-27
 2,4-Undecadiene-7,9-diyneic acid, U-25
 2,4-Undecadiene-8,10-diyneic acid, U-26
 2,4-Undecadieneic acid isobutylamide, U-28
 2,4-Undecadieneic acid, U-28
N-Undecanoyltyrosine, T-700
 2,7,9-Undecatrienoic acid isobutylamide, U-29
 2,7,9-Undecatrienoic acid, U-29
 2-Undecene-8,10-diyneic acid isobutylamide, U-30
 2-Undecene-8,10-diyneic acid, U-30
 2-(1-Undecenyl)-4(1*H*)-quinolinone, H-757
 2-(3-Undecenyl)-4(1*H*)-quinolinone, H-757
 Undecylprodiginine, U-31
 Undecylprodigosine, U-31
 2-Undecyl-3,4-quinolinediol, A-621
 2-Undecyl-4-quinolinol, H-757
 2-Undecyl-4(1*H*)-quinolinone *N*-oxide, H-757
 2-Undecyl-4(1*H*)-quinolinone, H-757
 Undulatine, C-743
 Undulatine, U-32
 Undulifoline, U-33
 Ungeremine, U-34
 Ungerine, H-294
 Ungernine, T-53
 Ungminoridine, Z-15
 Ungminorine *N*-oxide, U-35
 Ungminorine, U-35
 Ungspirolidine, P-626
 Ungspiroline, P-626
 Ungvedine, T-53
 Uniflorine A, T-262
 Uniflorine B, T-262
 Uniphyllin, T-360
 Unonopsine, U-36
 Unsevine, U-37
 Untenine A, N-256
 Untenine B, N-249
 Untenine C, N-256
 Uoamine A, U-38
 Uoamine B, U-38
 Uopenamide, U-39
 Uplandicine, U-40
 Urabaine, U-41

- Uracil N^3 -oxide, U-42
 Uracil-6-carboxylic acid, D-669
 Uracil, U-42
 Uraline, G-91
 Uranidine, Q-44
 Urceoline, N-171
 Urdamycin D, U-43
 Urechochrome, X-11
 3-Ureidoandrost-4-en-17-ol, A-702
 3-Ureidoandrost-4-en-17-one, A-702
 5-Ureidohydantoin, A-625
 5-Ureido-2,4-imidazolidinedione, A-625
 Ureidosorbicillinol, U-44
 7,8-Ureylenenanoic acid, M-499
 Uric acid, U-45
 Urminine, N-171
 Urocanic acid, U-46
 Urocaninic acid, U-46
 Urochordamine A, U-47
 Urochordamine B, U-47
 Uropterin, A-750
 Urothione, U-48
 Uroxanthin, I-104
 Ursuline, D-626
 Ursuline, U-49
 Urukthapelstatin A, U-50
 Us 7, A-603
 Us 8, A-603
 Usambanoline, U-51
 Usambarensine N^9 -oxide, U-52
 Usambarensine, U-52
 Usambaridine Br, O-21
 Usambaridine Vc, O-21
 Usambaridine Vi, O-21
 Usambarine, O-21
 Usaramensine, S-238
 Usaramine, R-62
 Usaramoensine, S-238
 Uscharin, U-53
 USF 19A, A-1305
 USF 142A, M-512
 USF 406C, D-681
 USF 406D, D-681
 Ushinsunine β - N -oxide, N-316
 Ushinsunine, N-316
 Usimine A, U-54
 Usimine B, U-54
 Usimine C, U-54
 Uskudaramine, U-55
 Uspallatine, A-967
 Uspallatinecine, C-775
 Ussuriedine, U-56
 Ussuriedinone, U-56
 Ussurienine, U-56
 Ussurienone, U-56
 Ustalic acid imide, D-816
 Ustalic acid, U-57
 Ustalic ethanolamide, U-57
 Ustalic glycine amide, U-57
 Ustalic prolinamide, U-57
 Usuramine, R-62
 Uthongine, O-170
 Uvarilactam, T-539
 Uvarindole A, U-59
 Uvarindole B, U-60
 Uvarindole C, U-59
 Uvarindole D, U-61
 Uvarindole E, U-62
 Uvariopsamine N -oxide, U-63
 Uvariopsamine, U-63
 Uvariopsine, U-64
 Vaccinine A, A-1581
 Vaccinine B, A-1583
 Vaganine A, D-266
 Vaganine B, D-274
 Vaganine C, D-274
 Vaganine D, D-280
 Vaginadine, M-309
 Vaginaline, M-309
 Vaginatine, M-309
 Vaillantine, M-734
 Vakaidine, V-1
 Vakatisine, V-2
 Vakatisinine, V-3
 Vakhmadine, H-182
 Vakhmatine, H-191
 Valachine, V-4
 Valacidin, S-569
 Valamine, M-525
 Valanimycin, V-5
 Valdiberine, P-15
 Valdivianine, P-15
 Valerianine, T-60
 Valeroidine, M-391
 13-Valeroyloxylupanine, H-554
 Valindolmycin, V-6
 1-Valine-2-(2,3-didehydroalanine)thiostrepton, S-321
 Valivine, V-7
 Vallesamidine, V-8
 Vallesamine N^6 -oxide, V-9
 Z-Vallesamine, V-9
 Vallesamine, V-9
 Vallesiachotamine lactone, V-11
 Vallesiachotamine, V-10
 Vallesine, A-1499
 Valloporfine, V-12
 Vallotidine, V-13
 Vallotine, V-14
 Valonaec dilactone, V-15
 Valoneic acid amide dilactone, V-15
 Valoneic acid dilactone, V-15
 Valoneic dilactone, V-15
 Valparicine, V-16
 Valtropine, M-393
O-Valylgriseoluiteic acid, H-514
 Valyltenuazonic acid, N-311
 L-Valyl-N-[1-(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-1-isoquinolinyl)ethyl]glycinamide, A-934
 Vampirol, S-589
 Vancocresmycin, V-17
 Vandrikidine, H-743
 Vandrikine, M-666
 Vanessine, A-1325
 Vanillobiocin, V-18
O-Vanilloylcyclovirobuxine D, C-917
 2-Vanilloylimidazole, H-558
 13-Vanilloxyloxylupanine, H-554
 3 α -Vanilloxyloxynortropane, C-633
 3 β -Vanilloxyloxytropane, M-393
 Vanilloylveracevine, V-56
 Vanilloylzygadenine, Z-36
 Vanillyl tropate, C-633
N-Vanillylnonanamide, C-100
 Vanuatine, V-19
 Vaponephrin, A-152
 VAPY, B-221
 Varacin A, V-20
 Varacin B, V-20
 Varacin C, V-20
 Varacin, L-205
 Varadine, V-21
 Varamine A, V-22
 Varamine B, V-23
 Variabiline, V-24
 Variceramides, V-25
 Variecolorin A, I-230
 Variecolorin B, I-230
 Variecolorin C, I-230
 Variecolorin D, I-230
 Variecolorin E, I-230
 Variecolorin F, V-26
 Variecolorin G, V-26
 Variecolorin I, I-230
 Variecolorin J, V-27
 Variecolorin K, V-28
 Variecolorin L, V-29
 Variecolortide A, V-30
 Variecolortide B, V-31
 Variecolortide C, V-31
 Variogatine, H-161
 Variolin A, V-32
 Variolin B, V-32
 Variolin D, V-33
 Variotin, V-34
 Varunax, A-845
 Vasconine, V-35
 Vasexeten, Q-17
 Vasicine, P-165
 Vasicinine, V-36
 Vasicinolone, V-38
 Vasicinol, V-37
 Vasicinone N -oxide, V-38
 Vasicinone, V-38
 Vasicoline, V-40
 Vasicolinone, V-40
 Vasicol, V-39
 Vasnetine, A-1041
 Vatamidine, V-41
 Vatamine, V-42
 Vateamine 2'- β - N -oxide, V-43
 Vateamine, V-43
 Vatine A, V-44
 Vatine, V-44
 vD 844, O-124
 vD 846, O-124
 VD, V-160
 (15 β)-Veatchine acetate (ester), V-45
 Veatchine, V-45
 Veatchinium chloride, V-45
 Vebilocine, D-528
 Veiutamine, V-46
 Velbanamine, C-522
 (+)-Velbanamine, C-522
 Velban, V-110
 Velbe, V-110
 Veldopa, A-745
 Vellosimine, T-425
 Vellosiminol, T-425
 Velloisine, K-78
 Velucryptine, L-252
 Velutinam, T-539
 Venacarpine A, V-47
 Venacarpine B, V-48
 Venalstonidine, V-49
 Venalstonine, V-49
 Venartan, P-923
 Venecurine, T-616
 Venenatic acid, V-50
 Venenatine, V-50
 Veneserpine, R-50
 C-Venezueline, V-51
 Vengicide, S-53
 Venoterpine, V-52
 Venoxidine, V-50
 Venturamide A, V-53
 Venturamide B, V-54
 Vendelphine, H-194
 Venulol, H-164
 Venulusone, D-586
 Vepridimerine A, P-87
 Vepridimerine B, P-87
 Vepridimerine C, V-55
 Vepridimerine D, V-55
 Vepridimerine E, V-55
 Veprisilone, P-625
 Veprisine, F-90
 Veprisinium, D-720
 Verabenzamine, G-70
 Veracevine, V-56
 Veracintine, V-57
 Veraflorizine, C-301
 Veragenine, S-1
 Veragenine, V-58
 Veralbidine, V-59
 Veralcamine, V-61
 Veralinine, V-60
 Veralkamine, V-61
 Veralobine, V-62
 Veralodine, H-460
 Veralodimine, V-63
 Veralodisine, V-63
 Veralomidene, V-60
 Veralomidine, V-60
 Veralomine, V-60
 Veralosidine, S-199
 Veralosidimine, V-64
 Veralosine, S-199
 Veralosinine, S-199

- Veralozidine, S-199
 Veralozine, S-199
 Veralozinine, S-199
 Veramanine, V-65
 Veramarine, C-309
 Veramiline 3-*O*- β -*D*-glucopyranoside, V-73
 Veramiline, V-73
 Veramine, V-66
 Veraminine, V-67
 Veramitaline, V-68
 Veramivirine, V-73
 Veranovine, V-69
 Verapatuline, J-36
 Verareine, V-70
 Verarine, V-71
 Verasine, V-73
 Veratensine, G-70
 Veratetrine, P-674
 Veratra-5,12-dien-3-ol, H-364
 Veratraman-3-ol, H-364
 Veratramine, V-71
 Veratra-5,11,13-triene-3,23-diol, V-72
 Veratrenone, T-534
 Veratric acid, D-706
 Veratridine, V-56
 Veratrine, V-56
 Veratrosbasine, J-36
 Veratrole-4,5-dicarboxylic acid, D-533
 Veratrosine, V-71
 Veratroylbikhaconine, M-312
 14-Veratroylchamanine, M-323
 15-Veratroyldictysine, D-370
 Veratroylformic acid, D-649
 15-Veratroylgermine, G-70
 3-Veratroylgermine, G-70
 8-Veratroylneoline, M-323
 14-Veratroylneoline, M-323
 Veratroylnortropine, C-633
 3 α -Veratroyloxy-*N*-acetylnortropine, C-633
 3 α -Veratroyloxy-*N*-hydroxynortropine, C-633
 3 α -Veratroyloxynortropine, C-633
 Veratroylpseudaconine, M-336
 15-Veratroylpseudokobusine, H-205
 Veratroyltropine, C-633
 15-Veratroylyesonine, Y-10
 Veratroylzygadenine, Z-36
 Veratrylidenehydrazide, D-705
 Verazine, V-73
 Verazinine, V-73
 Verbacine, P-673
 Verballocline, P-673
 Verballoscenine, P-673
 Verbamedine, P-673
 Verbamekrine, P-665
 Verbamethine, P-665
 Verbametrine, P-665
 Verbascenine, P-673
 Verbascine, P-665
 Verbasikrine, P-673
 Verbasitrine, P-673
 Verbaskine, P-665
 Verdine, V-74
 Verdinine, P-280
 Verine, V-75
 Vermelhotin, V-76
 Vermisporin, V-77
 Vernamycin A, O-131
 Vernamycin, O-131
 Veronal, B-23
 Veronamine, T-326
 Verongamine, V-78
 Verongiaquinol, H-650
 Verpacamide A, C-841
 Verpacamide B, C-841
 Verpacamide C, V-79
 Verpacamide D, V-80
 Verrucarine E, A-57
 Verrucarine F, V-81
 Verrucarine G, V-82
 Verrucine A, V-83
 Verrucine B, V-83
 Verrucofortine, F-160
 Verrucosine, F-160
 Verrucosinol, F-160
 Verrucozine, F-160
 Verrucozinol, F-160
 Verruculotoxin, V-84
 Versicolamide B, S-153
 Versimide, V-85
 Vertaline B, V-87
 Vertaline, V-86
 Verticillatine, P-241
 Verticillatine, V-88
 Verticillin A, V-89
 Verticillin B, V-89
 Verticillin C, V-90
 Verticillin D, V-91
 Verticillin E, V-91
 Verticillin F, V-91
 Verticillin G, V-92
 Verticilline, V-93
 Verticine *N*-oxide, C-298
 Verticine, C-298
 Verticinone *N*-oxide, C-298
 Verticinone, C-298
 Vertilecanine A, V-94
 Vertilecanine B, V-94
 Vertilecanine C, V-94
 Vertine, V-95
 Verussurine, G-70
 Verussurinine, G-70
 Very fast death factor, A-974
 Vevesca, H-616
 VFDF, A-974
 Viaductor, A-216
 Vibratilicin, N-114
 Vibrindole A, S-564
 Vibriobactin, V-96
 Vibrioferrin, V-97
 Viburnine, P-356
 Viburnum acid, M-409
 Vicenistatin aglycone, V-98
 Vicenistatin M, V-98
 Vicenistatin, V-98
 Vicianin, H-673
 Vicine, D-253
 Vicioside, D-253
 Vietnamine, P-149
 Vignatic acid A, V-99
 Vignatic acid B, V-100
 Viguine, V-101
 Vilaportine, O-214
 Villagorgin A, V-102
 Villagorgin B, V-102
 Villalstonine 4'-*N*-oxide, V-103
 Villalstonine, V-103
 Villatamine A, V-104
 Villatamine B, V-104
 Vilmoridine, V-105
 Vilmoriniamine, K-48
 Vilmorinine, M-352
 Vilmorisine, F-147
 Vilmoritine, V-106
 Vilmorrianine A, M-303
 Vilmorrianine B, M-336
 Vilmorrianine C, M-323
 Vilmorrianine D, M-355
 Vilmorrianone, M-665
 Vinamidine, V-107
 Vinaphamine, V-108
 Vinaspine, V-109
 Vinblastine sulfate, V-110
 Vinblastine, V-110
 Vinburnine, E-6
 Vincadiffine, V-171
 $\Delta^{6,7}$ - ψ -Vincadifformine, P-733
 ψ -Vincadifformine, P-733
 Vincadifformine, V-111
 Vincadine, V-112
 (\pm)-Vincadine, V-112
 Vincadioline, V-110
 Vincacor, V-115
 Vincaherbine, H-137
 Vincaherbinine, H-137
 Vincaine, A-212
 Vincalucoblastine, V-110
 Vincalucoblastine, V-110
 Vincamajine 17-*O*-veratrate, V-113
 Vincamajine, V-113
 Vincamajinine, V-113
 Vincamajoreine, T-288
 Vincamajoridine, A-232
 Vincamarine, V-115
 Vincamedine, V-113
 Δ^{14} -Vincamenine, E-5
 Vincamenine, E-5
 Vincametrine, V-115
 Vincamicine, V-114
 Vincamidine, S-576
 Vincamine, V-115
 Δ^{14} -Vincamine, V-115
 Vincaminine, V-115
 Vincaminoreine, V-112
 Vincaminoridine, V-112
 Vincaminorine, V-112
 Δ^{14} -Vincamone, E-6
 Vincamone, E-6
 Vincanicine, N-301
 Vincanidine, N-301
 Vincanine *N*-oxide, N-301
 Vincanine, N-301
 Δ^{14} -Vincanol, E-6
 Vincanol, E-6
 Vincanorine, E-6
 Vincanovine, V-116
 Vincapsine, V-117
 Vincareine, V-115
 Vincaricine, P-417
 Vincaridine, P-417
 Vincarine, V-113
 Vincarinine, B-420
 Vincarodine, V-117
 Vincarpine, F-82
 Vincarubine, V-118
 Vincathicine, V-119
 Vincatine, V-120
 Vincawajine, V-121
 Vinceine, A-212
 Vincetene, V-122
 Δ^{14} -Vincine, V-115
 Vincine, V-115
 Vincinine, V-115
 Vincolidine, V-123
 Vincoline, V-124
 Vincophylline, D-410
 Vincoridine, V-125
 Vincorine, V-125
 Vincosamide, V-126
 Vincoside lactam, V-126
 Vincoside, S-583
 Vincosid, V-128
 Vincovalicine, V-127
 Vincovaline, V-110
 Vincovalinine, L-162
 Vincovine, V-125
 Vincristine sulfate, V-128
 Vincristine, V-128
 Vincrisul, V-128
 Vincubine, T-283
 Vindesine sulfate, V-110
 Vindesine, V-110
 Vindolicine, V-129
 Vindolidine, V-130
 Vindolidine, V-131
 Vindoline, V-130
 Vindolinine *N*-oxide, V-132
 19*R*-Vindolinine *N*-oxide, V-132
 19*S*-Vindolinine *N*-oxide, V-132
 Vindolinine, V-132
 19*S*-Vindolinine, V-132
 Vindorosine, V-130
 Vinepidine, V-110
 Vineridine *N*-oxide, C-1
 Vineridine, C-1
 Vinerine *N*-oxide, C-1
 Vinerine, C-1
 Vinerinine, M-50
 Vinervidine, N-301
 Vinervine, A-228

- Vinervinine, A-228
 Vinformide sulfate, L-162
 Vinformide, L-162
 Vinylicinate sulfate, V-110
 Vinylicinate, V-110
 Vingramine, V-133
 Vinine, C-117
 Vinleurosine sulfate, L-162
 Vinleurosine, L-162
 Vinomine, V-134
 Vinorine, V-194
 Vinosidine, V-135
 Vinothiam, T-366
 Vinoxine, V-136
 Vinrosidine sulfate, V-110
 Vinrosidine, V-110
 Vinsedicine, V-137
 Vinsedine, V-138
 Vinsosidinic acid, S-583
 1-Vinyl- β -carboline-3-carboxylic acid, V-140
 1-Vinyl- β -carboline, V-139
 6-*O*-(5-Vinylnicotinoyl)tetrahydrocantlyne, T-145
 5-Vinyl-3-pyridinecarboxylic acid, V-141
 Violacein, V-142
 Violapterin, P-763
 Violatinctamine, V-143
 Violydoenamidine, H-667
 Viomycidine, V-144
 Viquidil, Q-17
 Virantmycin, V-145
 Virenamide A, V-146
 Virenamide B, V-147
 Virenamide C, V-148
 Virenamide D, V-146
 Virenamide E, V-149
 Virescenine, M-325
 Virgiboidine, V-152
 Virgivarine, V-150
 Virgilidine, V-151
 Virgilidone, V-152
 Virgiline 2-pyrrolicarboxylate, V-153
 Virgiline, V-153
 Virginiamycin M₁, O-131
 Virginiamycin M₂, O-131
 Virginiamycin, O-131
 Viridic acid, V-154
 Viridicatin, D-656
 Viridicatol, D-656
 Viridiflorine *N*-oxide, T-443
 Viridiflorine, T-443
 Viriditin, V-155
 Viridomycin A, V-156
 Viridomycin B, V-157
 Viridomycin C, V-158
 Viridomycin E, V-159
 Viridomycin F, V-156
 Viroallosecurinine, A-630
 Viroidin, A-14
 Virodhamine, E-48
 Virosecurinine, S-206
 Virosine A, S-207
 Virosine, N-307
 Virotoxin, V-160
 Viscosaline, V-161
 Viscosamine, V-162
 Visoltricin, F-196
 Vitamin B_c conjugate, P-772
 Vitamin B₄, A-137
 Vitamin B₇, B-139
 Vitamin B₁₃, D-669
 Vitamin B₁₇, H-673
 Vitamin B₆, P-923
 Vitamin B_c, P-772
 Vitamin B₃, P-895
 Vitamin B₁, T-366
 Vitamin H, B-139
 Vitamin L₁, A-706
 Vitamin M, P-772
 Vitamycin A, V-163
 Vitaneurin, T-366
 Vitedoamine A, V-164
 Vitelactam A, D-601
 Vitricine, V-165
 Vittacarboline, V-166
 Vittatine, C-748
 Vivotoxin, T-74
 Vixaton, C-555
 VLB, V-110
 VM 55595, A-1306
 VM 55598, A-1486
 VM 55599, A-1307
 VM 29919, P-90
 VM 54158, P-91
 VM 54159, P-90
 VM 55594, P-91
 VM 55596, P-90
 VM 55597, P-90
 Voacafricine, V-171
 Voacafrine, V-171
 Voacamidine, V-167
 Voacamine *N*^b-oxide, V-168
 Voacamine, V-168
 Voacaminine, V-169
 Voacangarine, V-170
 Voacangine 3-carbonitrile, V-170
 Voacangine hydroxyindolenine, C-658
 Voacangine pseudoindoxyl, V-178
 Voacangine, V-170
 Voacanginine, V-168
 Voacarpine, V-171
 Voachalotinal, P-562
 Voachalotine oxindole, V-172
 Voachalotine, P-562
 Voacinal, V-173
 Voacoline, V-174
 Voacorine, V-168
 Voacristine hydroxyindolenine, C-658
 Voacristine pseudoindoxyl, V-178
 Voacristine, V-170
 Voacryptine, V-170
 Voafinidine, V-181
 Voafinine, V-179
 Voafolidine, V-175
 Voafoline, V-175
 Voafrine A, V-176
 Voafrine B, V-176
 Voaharine, V-177
 Voalene, V-180
 Voalutene, V-178
 Voamonine, V-174
 Voaphylline hydroxyindolenine, V-180
 14,15-Voaphyllinediol, V-181
 Voaphylline, V-179
 Voastriptide, V-182
 Vobasenal, V-183
 Vobasine methine, R-27
 Vobasine *N*^a-oxide, P-268
 Vobasine, P-268
 Vobasonidine, V-184
 Vobatricine, V-185
 Vobparicine *N*-oxide, V-186
 Vobparicine, V-186
 Vobtusamine, V-187
 Vobtusine lactam, V-188
 Vobtusine, V-188
 Vobtusine lactone, V-188
 Vochysine, V-189
 Volicitin, G-108
 Volkenin, D-565
 Volkensine, P-417
 Volutamide A, V-190
 Volutamide B, V-191
 Volutamide C, V-191
 Volutamide D, V-192
 Volutamide E, V-193
 Vomalidine, A-216
 Vomicine, I-12
 Vomifoline, P-241
 Vomilenine, V-194
 Voruscharin, U-53
 Vulcanine, M-521
 Vulgarobufotoxin, B-394
 Vulgaxanthin II, V-195
 Vulgaxanthin I, V-195
 Vulkacit M, B-79
 Vulnibactin, V-96
 Vulparine, P-776
 Vulracine, V-196
 W 45, A-845
 Waiakeamide 21-sulfone, H-24
 Waiakeamide, H-24
 Waianaeamine A, W-1
 Waianaeamine B, W-2
 Wakamato, B-101
 Wakayin, W-3
 Wallemia A, W-4
 Wallemia B, W-4
 Wallemia C, W-5
 Wallemia D, W-5
 Wallemia E, W-4
 Wallemia F, W-5
 Waltherine A, W-6
 Waltherine B, A-935
 Waltherine C, W-7
 Waltherione A, W-8
 Waltherione B, W-8
 Wampocap, P-897
 Wanpeinane A, C-298
 Warifteine, W-9
 Warsine A, D-275
 Wasalexin A, W-10
 Wasalexin B, W-10
 Watasemycin A, W-11
 Watasemycin B, W-11
 Watasenia oxyluciferin, O-223
 Wax C, E-225
 Waziristanine, P-16
 WB 2838, A-725
 Weberidine, T-197
 Weberine, T-276
 Welwistatin, W-13
 Welwitindolinone A isonitrile, W-12
 Welwitindolinone B isothiocyanate, W-13
 Welwitindolinone C isothiocyanate, W-13
 Welwitindolinone D isonitrile, W-14
 Westiellamide, C-919
 Wewakazole, W-15
 WF 14865, A-1308
 WF 6237, A-680
 WF 11231A, A-1178
 WF 14861A, A-1309
 WF 14865A, A-1308
 WF 16775A₁, A-1535
 WF 16775A₂, A-1535
 WF 14865B, A-1308
 WF 1360B, R-83
 WF 1360C, R-83
 WF 1360F, R-83
 WF 1360, R-83
 Whey factor, D-669
 Wieland-Gumlich aldehyde *N*-oxide, C-115
 Wieland-Gumlich aldehyde, C-115
 Wieland-Gumlich diol, R-63
 Wilfordicine, T-257
 Wilfordconine, P-249
 Wilfordine, P-253
 Wilfordinine A, P-249
 Wilfordinine B, P-249
 Wilfordinine C, P-249
 Wilfordinine D, P-250
 Wilfordinine E, P-250
 Wilfordinine F, P-250
 Wilfordinine G, P-250
 Wilfordinine H, P-250
 Wilfordinine I, P-249
 Wilfordinine J, P-249
 Wilfordlongine, P-249
 Wilfordsine, P-249
 Wilfordsuine, P-250
 Wilforgine, P-252
 Wilfordidine, P-253
 Wilforine, P-252
 Wilforjine, P-252
 Wilformine, P-252
 Wilformine A, P-253
 Wilformine B, P-253
 Wilformine C, P-253
 Wilformine D, P-253
 Wilformine E, P-253

- Wilforfine F, P-247
 Wilforfine G, P-247
 Wilforfine, P-252
 Wilfortrine, P-253
 Wilforzine, P-252
 Wilfoside G1G, P-610
 Willicourtine, A-216
 Wilsonine, F-143
 Wilsonirine, H-751
 Win 11831, A-216
 WIN 64745, A-1310
 WIN 64821, A-1311
 WIN 5512, O-63
 Winkleridine, M-325
 Winkleriline, M-325
 Wintermin, C-417
 Wisanidine, W-16
 Wisanine, W-17
 Withanamide A, H-755
 Withanamide B, H-755
 Withanamide C, H-755
 Withanamide D, H-755
 Withanamide E, H-755
 Withanamide F, H-755
 Withanamide G, H-755
 Withanamide H, H-755
 Withanamide I, H-755
 Withanamine, W-18
 Withanine, W-19
 Withasomine, D-492
 WJ 35, A-1312
 WJ 85, A-1313
 WJ 85B, A-1313
 WK 5344A, F-45
 WK 5344B, F-45
 Wobezit M, B-79
 Wondonine A, W-20
 Wondonine B, W-20
 Woodinine, W-21
 Worenine, W-22
 10381W, P-641
 Wrightiamine A, C-610
 Wrightiamine B, A-883
 Wrightine, C-601
 WS 1921, A-1314
 WS 30581A, A-622
 WS 75624A, A-1315
 WS 63967A, C-129
 WS 9659A, L-76
 WS 30581B, A-622
 WS 75624B, A-1316
 WS 63967B, C-129
 WS 9659B, L-76
 WS 63967, C-129
 WS 30581C, A-622
 WS 63967C, C-129
 WS 63967D, C-129
 WS 63967E, C-129
 WS 63967F, C-129
 Wuchuyine, W-23
 Wuchuyuamide III, W-25
 Wuchuyuamide II, W-24
 Wuchuyuamide I, W-24
 Wuzhuyurutine A, W-26
 Wuzhuyurutine B, W-27
 Wybutine, W-28
 Wybutoxine, W-29
 Wye base, W-30
 Wyosine, W-30
 Wyo, W-30
 X 14885A, A-1087
 X 14547A, I-64
 X 53III, Q-58
 X 948, Q-58
 Xanthaline, P-77
 α -Xantherin, X-1
 Xanthevodine, X-2
 Xanthiazone, X-3
 Xanthicin, X-4
 Xanthine, X-5
 Xanthiside, X-3
 Xanthobaccin A, X-6
 Xanthocillin X dimethyl ether, X-7
 Xanthocillin X, X-7
 Xanthocillin Y₁, X-7
 Xanthocillin Y₂, X-7
 Xanthocillin, X-7
 C-Xanthocurine, X-8
 Xanthofagarine, X-9
 Xantholipin, X-10
 Xanthommatin, X-11
 Xanthopetaline, A-954
 Xanthoplanine, L-75
 Xanthopterin, A-750
 Xanthopterincarboxylic acid, X-12
 Xanthopterinsulfonic acid, A-750
 Xanthoquinic acid, H-726
 Xanthoquininamide, H-726
 Xanthoquininic acid, H-726
 Xanthoraphine, P-319
 Xanthosine 5'-(dihydrogen phosphate), X-13
 Xanthosine monophosphate, X-13
 Xanthosine 5'-phosphate, X-13
 Xanthosine 5'-sulfate, X-13
 Xanthosine, X-13
 Xanthothricin, R-66
 Xanthoxoline, T-516
 Xanthoxyphylline, T-525
 Xanthurenic acid, D-677
 Xanthuric acid, D-677
 3'-Xanthylic acid, X-13
 5'-Xanthylic acid, X-13
 Xantocillin, X-7
 Xantyrin, X-7
 Xenocoumacin 1, X-14
 Xenorhabdin 1, A-759
 Xenorhabdin 2, A-759
 Xenorhabdin 3, A-759
 Xenorhabdin 4, A-759
 Xenorhabdin 5, A-759
 Xenovenine, H-135
 Xeranthin, H-519
 Xestamine A, X-15
 Xestamine B, X-16
 Xestamine C, X-17
 Xestamine D, X-18
 Xestamine E, X-19
 Xestamine F, X-15
 Xestamine G, X-18
 Xestamine H, X-19
 Xestoamine, X-20
 Xestocyclamine A, X-21
 Xestocyclamine B, X-22
 Xestomanzamine A, X-23
 Xestomanzamine B, X-23
 Xestomycin, R-41
 Xestosine A, P-290
 Xestospongina A, X-24
 Xestospongina B, X-25
 Xestospongina C, X-24
 Xestospongina D, X-24
 XK 211, A-1317
 XK 101-1, T-658
 XL 119, R-32
 XMP, X-13
 10381X, P-642
 X-Pigment, P-319
 XR 587, S-571
 Xtro, T-632
 Xylactam, X-26
 Xylaramide, X-27
 Xylariamide A, A-724
 Xyloallenolide A, X-28
 Xylobovatin, C-954
 Xyloformycin, F-131
 Xylogranatinine, X-29
 Xyloguyelline, P-369
 Xylopinine, A-1066
 Xylopinidine, L-68
cis-Xylopinine *N*-oxide, X-30
trans-Xylopinine *N*-oxide, X-30
 Xylopinine, X-30
 6'-*O*- α -D-Xylopyranosylalanside, A-239
 5-*O*- β -D-Xylopyranosylstreptazolin, S-560
 Xylopyrine A, P-29
 Xylopyrine B, J-52
 Xylostosidine, X-31
 7-Xylosylcephalomannine, T-52
 7-Xylosyl-10-deacetylaxol D, T-52
 7-Xylosyl-*N*-methylaxol C, T-52
 7-Xylosylaxol B, T-52
 7-Xylosylaxol C, T-52
 7-Xylosylaxol D, T-52
 7-Xylosylaxol, T-52
 XYOTAX, T-52
 Y 03559J-A, A-1318
 Y 05460MA, A-1319
 Y 16482 α , S-2
 Y 16482 β , S-2
 Y 16601, S-2
 Y 16760, S-2
 Yaequinolone A₁, D-422
 Yaequinolone A₂, D-422
 Yaequinolone B, Y-1
 Yaequinolone C, Y-2
 Yaequinolone D, P-191
 Yaequinolone E, P-193
 Yaequinolone F, Y-2
 Yaequinolone J₁, Y-3
 Yaequinolone J₂, Y-3
 Yageine, H-592
 Yagonine, Y-4
 Yamataimine, B-194
 Yanangcorinine, T-407
 Yanangine, Y-5
 Yaoundamine A, Y-6
 Yaoundamine B, Y-6
 Yatakemycin, Y-7
 Yatorizine, P-34
 Y-base, W-28
 YC 73, F-97
 YCM 1008A, A-1320
 Yemenine A, H-5
 Yemenine B, H-6
 Yemenine C, H-6
 Yemensine, Y-8
 Yehusomidine, R-9
 (\pm)-Yehusomidine, R-9
 Yehusomine, R-9
 Yersiniabactin, Y-9
 Yersiniophore, Y-9
 Yesodine, H-205
 Yesoensine, G-152
 Yesoline, Y-10
 Yesonine, Y-10
 Yesoxine, G-139
 Yewtaxan, T-52
 Yibeinoside A, C-290
 Yibeinoside B, C-290
 Yibeinoside C, P-807
 Yibeissine, Y-11
 YL 03709B-A, D-574
 YL 0358M-A, H-77
 YL 0358M-B, H-77
 YL 02729S, M-491
 YM 193221, A-1321
 YM 215343, A-1340
 YM 216391, A-1322
 YM 47522, D-574
 YM 30059, M-491
 YM 266183, T-383
 YM 266184, T-383
 YN 0165JA, A-1323
 Yobinol, Y-13
 Yocon, Y-13
 Yohambinine, Y-12
 Yohimban-17-ol, Y-16
 15,17,19-Yohimbatriene, D-434
 Yohimbic acid, Y-13
 Yohimbine hydrochloride, Y-13
 β -Yohimbine *N*-oxide, Y-13
 β -Yohimbine oxindole, Y-14
 β -Yohimbine pseudoindoxyl, Y-15
 δ -Yohimbine, A-212
 α -Yohimbine, Y-13
 β -Yohimbine, Y-13
 Yohimbine, Y-13
 ψ -Yohimbine, Y-13
 Yohimbic acid, Y-13

- Yohimbol methosalt, Y-16
 Yohimbol, Y-16
 Yohimex, Y-13
 Yohimvetol, Y-13
 Yohydrol, Y-13
 Yokonoside, D-539
 Yondelis, E-37
 YP 02978L-C, L-76
 Ypaoamide, Y-17
 YSI 40-2, M-177
 Yt base, W-30
 Y-T 0678H, B-72
 Yt, W-30
 YUA 001, H-667
 Yuanamide, Y-18
 Yuanhunine, C-675
 Yubeinine, C-290
 Yubeiside, C-290
 Yuehchukene, Y-19
 Yukocitrine, J-59
 Yukodine, T-518
 Yukodinine, T-518
 Yukomine, T-580
 Yunaconitine, M-336
 Yunnadelphinine, M-321
 Yunnandaphnine A, D-33
 Yunnandaphnine B, D-33
 Yunnandaphnine C, D-33
 Yunnandaphnine D, D-33
 Yunnandaphnine E, Y-20
 Yunnanensine, Y-21
 Yunnanaxamine, T-52
 Yuremamine, Y-22
 Yuzifine, J-63
 Yuziphine, J-63
 Yuzuric acid, D-50
 Yuzurimic acid B, D-33
 Yuzurimine A, D-33
 Yuzurimine B, D-33
 Yuzurimine C, Y-23
 Yuzurimine D, Y-24
 Yuzurimine E, D-33
 Yuzurimine, D-33
 Yuzurine, D-50
 Y-wye, W-28
 Z 2055, E-98
 Z 203, S-589
 Zaidine, P-716
 Zalliline, G-91
 Zamamistatin, A-157
 Zangezurine, Z-1
 Zanoxyline, A-1444
 Zanthobisquinolone, Z-2
 Zanthobungeanine, F-90
 Zanthocadinanine A, Z-3
 Zanthocadinanine B, Z-3
 Zanthodioline, F-90
 Zanthomamide, Z-4
 Zanthomamine, Z-4
 Zanthomurolanine, Z-3
 Zanthonitrile, H-672
 Zanthophylline, F-90
 Zanthosimuline, Z-5
 Zanthosinamide, Z-6
 Zanthosin, Z-6
 Zanthoxyline, Z-7
 Zanthoxyphylline, T-525
 1745Z₃A, O-131
 Zapotidine, Z-8
 Zarzissine, A-829
 Zascanol epoxide, Z-9
 Zavesca, H-616
 Zeagenin, Z-10
 Zeanic acid, D-673
 Zeanoside A, D-424
 Zeanoside B, D-673
 Zeanoside C, D-424
 Zeatin 2-deoxyribofuranoside, Z-11
cis-Zeatin 9-glucoside, Z-11
cis-Zeatin, Z-11
trans-Zeatin, Z-11
 Zeatin, Z-11
 Zefbetaine, Z-12
 Zeflabetaine, U-34
 Zelkovamycin, Z-13
 Zem-Eserine, P-400
 Zenkerene N^b-oxide, Z-14
 Zenkerene, Z-14
 Zenkerene N^b-oxide, Z-14
 Zenkerine, T-524
 Zenkerine, Z-14
 Zephyranthine, Z-15
 Zeraconine N-oxide, Z-16
 Zeraconine, Z-16
 Zeravschanidine, Z-17
 Zeravschanine, H-157
 Zeravschanisine, H-189
 Zeravschanizine, H-189
 Zetekitoxin AB, Z-18
 Zeylamine, T-53
 ZG 1494 α , O-134
 Z-gal, B-324
 ZHD 0501, A-1324
 Zhebeinine, C-298
 Zhebeininoside, C-298
 Zhebeinone, C-298
 Zhebeirine, C-290
 Zhenguangmycin A₆, B-236
 Z-Hydroxyzinnimidine, C-437
 Ziebeimine, C-308
 Zierin, H-519
 Zierinxylsiole, H-519
 Zijinlongine, Z-19
 Zinnimidine, C-437
 Zinolol, T-528
 Zippelianine, T-28
 Zippeline, Z-20
 Zizyphine A, Z-22
 Zizyphine B, Z-22
 Zizyphine C, Z-23
 Zizyphine D, A-18
 Zizyphine E, A-18
 Zizyphine F, Z-22
 Zizyphine G, Z-24
 Zizyphine I, Z-21
 Zizyphine K, Z-25
 Zizyphine N, Z-26
 Zizyphine O, Z-26
 Zizyphine P, Z-26
 Zizyphine Q, Z-27
 Zizyphine, Z-22
 Zizyphinine, Z-22
 Zizyphusine, T-248
 Zn-Aspergillin, A-1475
 Zoamides, Z-28
 Zoanthamide, Z-29
 Zoanthamine, Z-30
 Zoanthaminone, Z-30
 Zoanthenamamine, Z-31
 Zoanthanol, Z-30
 Zoanthoxanthin, P-94
 Zongorine, N-30
 Zooanemonin, Z-32
 Zoolobelin, L-219
 Zooxanthellabetaine A, A-792
 Zooxanthellamine, Z-30
 Zopfiellamide A, Z-33
 Zopfiellamide B, Z-34
 Zostrich, C-100
 ZP-amide A, D-576
 ZP-amide B, D-576
 ZP-amide C, D-578
 ZP-amide D, D-578
 ZP-amide E, D-576
 ZP-amide F, D-576
 Zucapsaicin, C-100
 Zucchini factor B, M-730
 Zygacine, Z-36
 Zygadenilic acid δ -lactone, Z-35
 Zygadenine 3-glucoside, Z-36
 Zygadenine δ -lactone, Z-35
 Zygadenine, Z-36
 Zygofabagine, M-426
 Zygosporin A, Z-37
 Zygosporin D, Z-37
 Zygosporin E, Z-37
 Zygosporin F, Z-37
 Zygosporin G, Z-38
 Zyzzin, Z-39
 Zyzzyanone A, Z-40
 Zyzzyanone B, Z-40
 Zyzzyanone C, Z-40
 Zyzzyanone D, Z-40

Compound Index Contents

- Simple acyclic amine alkaloids with one N
Guanidine alkaloids
Nitriles and isonitriles
Simple isobutylamide alkaloids
Simple amide alkaloids
Azo and azoxy alkaloids
Simple pyrrolidine alkaloids
Chromone alkaloids
Flavonoid alkaloids
Nicotine-like alkaloids
Miscellaneous pyrrolidine alkaloids
Tetramic acids
Tropane alkaloids
Simple pyrrolizidine alkaloids
Pyrrolizidine alkaloids (macroyclic lactones)
Miscellaneous pyrrolizidine alkaloids
Anabasine-like alkaloids
Lobelia-type alkaloids
Simple piperidine alkaloids incl. piperidides
Xestospongins
Miscellaneous piperidine alkaloids
Lythraceae alkaloids
Quinolizidine alkaloids (two rings)
Quinolizidine alkaloids (three rings)
Quinolizidine alkaloids (four rings)
Cylindricine alkaloids
Miscellaneous quinolizidine alkaloids
Azepine alkaloids
Nicotinic acid derived alkaloids
Other pyridine alkaloids
9B-Azaphenalene alkaloids
Naphthalene-isoquinoline alkaloids
Elaeocarpus alkaloids
Galbulimima alkaloids
Stemona alkaloids
Lycopodium alkaloids
Cytochalasin alkaloids
Indolizidine alkaloids
Simple anthranilic acid alkaloids
Maytansinoid alkaloids
Simple quinoline alkaloids
Furanoquinoline alkaloids
Pyranoquinoline alkaloids
Miscellaneous quinoline alkaloids
Quinazoline alkaloids
Acridone alkaloids
Acridone dimers
Acridone-coumarin dimers
Pyrido[2,3,4-kl]acridines
1,4-Benzoxazin-3-one alkaloids
Benzodiazepine alkaloids
Cryptolepine type alkaloids
Simple phenethylamine alkaloids
 α -Hydroxyphenethylamines
Halogenated tyrosinoids
Miscellaneous phenethylamines
Cinnamic acid amides
Securinega alkaloids
Betalain alkaloids
Simple isoquinoline alkaloids
Manzamine alkaloids
Miscellaneous isoquinoline alkaloids
Benzylisoquinoline alkaloids
Pseudobenzylisoquinoline alkaloids
Bisbenzylisoquinoline alkaloids (aryl links only)
Bisbenzylisoquinoline alkaloids (one ether link)
Bisbenzylisoquinolines (one ether/one aromatic link)
Bisbenzylisoquinoline alkaloids (two ether links)
Bisbenzylisoquinoline alkaloids (two ether/one aromatic link)
Bisbenzylisoquinoline alkaloids (three ether links)
Secobisbenzylisoquinoline alkaloids
Cularine group alkaloids
Secoisoquinoline alkaloids
Quettamine alkaloids
Dibenzopyrrocoline alkaloids
Indenobenzazepine alkaloids
Pavine alkaloids
Isopavine alkaloids
Proaporphine alkaloids
Proaporphine-benzylisoquinoline alkaloid dimers
Aporphine alkaloids
Aporphine-benzylisoquinoline alkaloid dimers
Oxoisoaporphine alkaloids
Aristolochic acid alkaloids
Aristolactam alkaloids
Phenanthrene alkaloids
Azaaporphine alkaloids
Azafluoranthrene alkaloids
Morphine alkaloids
Erythrina alkaloids
Dibenzazecine alkaloids
Hasubanan alkaloids
Protoberberine alkaloids
Isoindolobenzazepine alkaloids
Narceine alkaloids
Protopine alkaloids
Rhoeadine alkaloids
Phthalideisoquinoline alkaloids
Spirobenzylisoquinoline alkaloids
Secoberberine alkaloids
Benzof[*c*]phenanthridine alkaloids
Phenethylisoquinoline alkaloids
Homomorphinandienone alkaloids
Homoaporphine alkaloids
Homoproaporphine alkaloids
Colchicine alkaloids
Cephalotaxus alkaloids
Homoerythrina alkaloids
Apogalanthamine alkaloids
Galanthamine alkaloids
Haemanthidine alkaloids
Graciline alkaloids
Tazettine alkaloids
Pancracine alkaloids
Lycorine alkaloids
Lycorenine alkaloids
Narciclasine alkaloids
Miscellaneous Amaryllidaceae alkaloids
Mesembrenoid alkaloids
Emetine group alkaloids
Phenanthroindolizidine alkaloids
Phenanthroquinolizidine alkaloids
Ericamycins
Simple indole alkaloids
Simple biindoles
Simple tryptamine alkaloids
Cyclotryptamine alkaloids
Chaetocin-like alkaloids
Evodia alkaloids
Indolactam alkaloids
 β -Carboline alkaloids
Carbazole alkaloids
Indolo[2,3-*a*]carbazole and related alkaloids
Indolonaphthyrindine alkaloids
Ergot alkaloids
Indoloterpenoid alkaloids
Strictosidine alkaloids
Camptothecine-like alkaloids
Corynanthean alkaloids
Secologanin-derived sesquimer alkaloids
Akageran and related alkaloids
Ajmalicine-like alkaloids
Macroline alkaloids
Malindan alkaloids
Cadamban alkaloids
Vallesiachotaman alkaloids
Gelsemium alkaloids
Yohimbinoide alkaloids
Sarpagine alkaloids
Ajmaline alkaloids
Vobasine alkaloids
Koumine alkaloids
Ervatamia alkaloids
Akuammiline alkaloids
Pleiocarpaman alkaloids
Cinchona alkaloids
Akuammicine alkaloids
Strychnidine alkaloids
Condylocarpan alkaloids
Secodine alkaloids
Aspidosperma alkaloids
Rhazinilam alkaloids
Melonine alkaloids
Quebrachamine alkaloids
Aspidofractine alkaloids
Vindoline alkaloids
Kopsane alkaloids
Melodinus alkaloids
Iboga alkaloids
Pandoline alkaloids
Pyridocarbazole alkaloids
Ulean alkaloids
Vallesaman alkaloids
Eburna alkaloids
Tacaman alkaloids
Schizozygine alkaloids
Chippiine alkaloids
Secologanin-derived dimer alkaloids
Isoindoles
Tryptotoxivalines
Mitomycins
Pyrrolo[4,3,2-*de*]quinoline alkaloids
Miscellaneous indole alkaloids
Monoterpenoid alkaloids
Sesquiterpene alkaloids
Macroyclic sesquiterpene alkaloids
Dendrobium alkaloids
Nuphar alkaloids
C19-Diterpenoid alkaloids and 4-nor analogues
C20 Diterpenoid alkaloids (Atisine type)
Erythrophleum alkaloids
Penitrens
Miscellaneous diterpene alkaloids
Sesterterpene alkaloids
Daphniphylline alkaloids
Steroidal alkaloids (salamandra type)
Steroidal alkaloids (jerveratrum type)
Steroidal alkaloids (cerveratrum type)
Steroidal alkaloids (conanine type)
Steroidal alkaloids (spirosolane type)
Steroidal alkaloids (solanidine type)
Steroidal alkaloids (buxus type)
Steroidal alkaloids (pregnane type)
Steroidal alkaloids (cephalostatins/ritterazines)
Miscellaneous steroidal alkaloids
Azaanthracene alkaloids
Azafluorene alkaloids
Pyrazole alkaloids
Imidazole alkaloids
Pyrrole-imidazole alkaloids
Cycloheptadiimidazoles
Oxazole and benzoxazole alkaloids
Isoxazole alkaloids
Spirobenzoxazoline alkaloids
Acyclic thiazole and benzothiazole alkaloids
Latrunculins
Macrocyclic thiazole alkaloids
Pyrazine and quinoxaline alkaloids
Benzo[1,2-*b*:3,4-*b'*]dipyrrroles
Pyrrolo[1,2-*a*]pyrazines
Morpholines
Pyrimidines
Ptilocaulins
Triazaacenaphthylene alkaloids

Tetrodotoxins	Polyamine alkaloids PA44 (homospermidines)	Saxitoxins
Phenazine alkaloids	Polyamine alkaloids PA343 (spermines)	Alkaloidal metal complexes
Phenoxazine alkaloids (other than actinomycins)	Other 3-residue polyamine alkaloids	Naphthyridinomycins
Simple pyrrole alkaloids	Polyamine alkaloids with more than 3 residues	Saframycins
Miscellaneous pyrrole alkaloids	Ansa-peptide alkaloids	Miscellaneous acyclic alkaloids
Polyamine alkaloids PA3, PA33 and homologues	Amanita alkaloids	Miscellaneous one-ring system alkaloids
Polyamine alkaloids PA4 (putrescines)	Other cyclopeptide alkaloids	Miscellaneous two-ring system alkaloids
Polyamine alkaloids PA5 (cadaverines)	Pyoverdins	Miscellaneous three-ring system alkaloids
Polyamine alkaloids PA43 (spermidines)	Pyrrolo[2,3-d]pyrimidines	Miscellaneous four- and higher-ring system alkaloids
	Purines	Alkaloids of unknown or partially unknown structure
	Pteridines and analogues	

Type of Compound Index

Simple acyclic amine alkaloids with one N

Acetylcholine(1+), A-47
Actinine, A-126
2-Aminoacetophenone, A-694
2-Amino-14,16-dimethyl-3-octadecanol, A-758
2-Aminoethanesulfonic acid, A-764
2-Aminoethanesulfonic acid, A-765
2-Aminoethanesulfonothioic acid, A-766
2-Aminoethanol, A-767
2-Aminoethyl dihydrogen phosphate, A-768
(2-Aminoethyl)phosphinic acid, A-772
(2-Aminoethyl)phosphonic acid, A-773
4-Amino-4-methyl-2-pentanone, A-839
1-Amino-2-methyl-2-propanol, A-840
8-Amino-7-oxononanoic acid, A-851
14-Aminotetradecanoic acid, A-911
Antibiotic YM 193221, A-1321
Benzylamine, B-84
1,3-Bis[(4-aminomethyl)phenyl]-2-octadecanoyl-glycerol, B-151
3-(Carboxymethylamino)propanoic acid, C-147
5-[(3-Carboxypropyl)amino]-2-hydroxy-3-tridecyl-1,4-benzoquinone, C-152
Choline, C-419
Crassifoline methine, C-729
2-Cyclohexylethylamine, C-869
Demanyl phosphate, D-197
1,2-Diacylglyceryl-3-(*O*-carboxyhydroxymethyl)choline, D-243
1,2-Diacylglyceryl-*O*-2'-(hydroxymethyl)-*N,N,N*-trimethyl- β -alanine, D-244
4-*O*-(1,2-Diacylglyceryl)-*N,N,N*-trimethylhomoserine, D-245
Diethanolamine, D-388
5,7-Dihydroxy-2-[1-(4-methoxy-2-oxo-2*H*-pyran-6-yl)-2-phenylethylamino]-1,4-naphthoquinone, D-610
Dimethylamine, D-726
2-(Dimethylamino)ethanol, D-730
N-[4-(Dimethylarsinoyl)butanoyl]aminoethylsulfonic acid, D-740
5-Dimethylsulfonio 2-amino-4-hydroxypentanoate, D-784
Echinobetaine A, E-14
Ethylamine, E-233
2-Guanidinoethanesulfonic acid, G-203
1-Heptylamine, H-130
1-Hexylamine, H-261
2-Hydroxybenzylamine, H-415
3-Hydroxybenzylamine, H-416
4-Hydroxybenzylamine, H-417
3,3'-[Iminobis(methylene)]bis-2(3*H*)furanone, I-46
(Iodomethyl)trimethylammonium(1+), I-164
Kuramerine, K-107
Lavendustin A, L-79
N-(2-Methoxy-2-propenyl)trimethylammonium(1+), M-284
Methylamine, M-383
3-Methyl-1-butylamine, M-420
N-(9-Methyldecyl)sulfamic acid, M-431
N-(8-Methylnonyl)sulfamic acid, M-493
N-(7-Methylheptyl)sulfamic acid, M-497
2-Methyl-1-propylamine, M-525
9-(Methylsulfonyl)nonylamine, M-564
3-(Methylthio)propylamine, M-578
12-Methyl-1-tridecylamine, M-579
Muscaridine, M-772
3,3',3''-[Nitrilotris(methylene)]tris[dihydro-2(3*H*)furanone], N-248
1-Propylamine, P-648
Sinapine, S-308
Tetramethylammonium(1+), T-280

Tribenzylamine, T-457
Trichosanatine, T-494
Trimethylamine oxide, T-601
Trimethylamine, T-600
(2-Trimethylammonioethyl)phosphonic acid betaine, T-602
Trimethyl[3-(methylthio)propyl]ammonium(1+), T-612
Trimethylvinylammonium(1+), T-615
Usumine B, U-54

Guanidine alkaloids

Acanthoine, A-25
3,5-Acarnidine, A-26
(4-Aminobutyl)guanidine, A-719
1-Amino-20-guanidino-4,8,12,16-tetrazaicosane, A-780
1-Amino-16-guanidino-4,8,12-triazaheptadecane, A-781
2-Amino-5-[[[3-(4-hydroxyphenyl)-2-methoxy-2-propenoyl]amino](imino)methyl]amino]pentanoic acid, A-806
(5-Aminopentyl)guanidine, A-859
(3-Aminopropoxy)guanidine, A-895
Aminoazoanemonin, A-928
Antibiotic L 681176, A-1195
Antibiotic SAB 711, A-1238
Antibiotic Sch 40873, A-1245
Antibiotic SF 2425, A-1258
Antibiotic TAN 2020, A-1279
Aplicyanin A, A-1343
Arglecin, A-1419
Argvalin, A-1421
Asterubin, A-1522
Birnbamin A, B-148
N,N'-Bis(4-guanidinobutyl)oxamide, B-178
N,N-Bis(3-methyl-2-butenyl)guanidine, B-199
Bistelletadine A, B-219
Bluensidine, B-237
4-Bromo-2-(guanidinomethyl)pyrrole, B-319
Canavanosuccinic acid, C-75
Capreomycin, C-99
Caracasandiamide, C-108
Carnosadine, C-163
Clavamine, C-512
Complanatin, C-592
Copiamycin, C-634
Crambescin A, C-721
Crambescin B, C-722
Crambescin C1, C-723
Crambescin C2, C-724
Creatine, C-731
Cycloanchinopeptolide C, C-840
Dehydrocrambine A, D-155
1,4-Diguanidinobutane, D-400
1,5-Diguanidinopentane, D-401
Distomadine A, D-868
Distomadine B, D-869
Ficellomycin, F-52
Fontaineine, F-124
Fuscusine, F-229
Gongrine, G-140
Guadinomic acid, G-185
Guadinomine A, G-186
Guadinomine C₁, G-187
Guanidine, G-199
Guanidinoacetic acid, G-200
4-Guanidinobutanoic acid, G-201
N-(4-Guanidinobutyl)-2-(4-hydroxyphenyl)-2-oxoacetamide, G-202
2-Guanidinoethanesulfonic acid, G-204
4-Guanidino-3-hydroxybutanoic acid, G-206
5-Guanidino-2-oxopentanoic acid, G-207
3-Guanidinopropanoic acid, G-208
3-Guanidinyipyridine, G-209

Hanasanagin, H-46
Hirudonine, H-305
Hordatine B, H-356
(4-Hydroxybutyl)guanidine, H-432
Kealiinine A, K-24
Laughine, L-71
Limaciamine, L-174
Cypridina Luciferin, L-281
Lysianadioic acid, L-373
Martinelline, M-111
N-(3-Methyl-2-butenyl)guanidine, M-410
Miraziridine A, M-648
Monamidocin, M-682
Monanchorin, M-683
Nephilatoxin 9, N-159
Nicaeensin, N-187
Orthidine A, O-125
Orthidine C, O-126
Orthidine E, O-127
Phloeocticyne B, P-367
Polyandrocarpidine A, P-544
Polyandrocarpidine C, P-545
Pterogynidine, P-769
Purealidin O, P-815
Pyronopaline, P-939
Resorcinomycin A, R-54
Resorcinomycin B, R-55
Rexostatine, R-69
Secomartinelline, S-191
Siphonodictidine, S-324
Spergualin, S-394
Spherophysine, S-406
Stelletazole A, S-507
Stelletazole B, S-508
Stendomyidine, S-527
Stizolamine, S-553
Streptolidine, S-565
Tetrahydroalathyrine, T-199
Triophamine, T-617
N,N,N'-Tris(3-methylbutyl)guanidine, T-623
Tubastrine, T-656
Verpacamide C, V-79
Verpacamide D, V-80
Viomycinidine, V-144
Xenocoumacin 1, X-14

Nitriles and isonitriles

Acacipetalin, A-20
Aleurodisconitrile, A-259
Antibiotic 0089D, A-1155
Antibiotic L 970843, A-1198
Antibiotic SF 2369, A-1257
Antibiotic Y 035591-A, A-1318
Cyanamide, C-823
Darlucin A, D-77
Darlucin B, D-78
6,7-Dihydroxy-3-isocyano-2*H*-1-benzopyran-2-one, D-599
Diptocarpilidine, D-830
Hemicalyculin A, H-102
3-Isocyano-7,9-bisaboladiene, I-213
Isonitrin A, I-255
Isonitrinic acid E, I-256
Oxazimine 1, O-153
Preoxazinin 7, P-619
Xanthocillin X, X-7

Simple isobutylamide alkaloids

Antibiotic U 106305, A-1299
Asaramide, A-1455
Brachystamide D, B-266
Chingchenamide A, C-387
2-Methoxy-*N*-(3-methylbutyl)acetamide, M-259

- 6-(4-Methoxyphenyl)-*N*-(3-methylbutyl)-2-oxo-3,5-hexadienamide, M-281
 Nigramide K, N-204
 Nigramide N, N-207
 Nigramide O, N-208
 Pipercollosidine, P-446
 Pipercollosine, P-447
 Pippulzarine, P-472
 Scabrine, S-113
 Spherophrysine, S-406
- Simple amide alkaloids**
- Acalyphamide, A-21
 Acetamide, A-30
 1-(2-Acetamidophenyl)-2-hydroxy-1-propanone, A-32
 Acinetoferrin, A-80
 Acrodontiolamide, A-117
 Ajudazol B, A-219
 Amidinomyacin, A-692
 Anhydroperipentamine, A-1033
 Anthelvencin A, A-1077
 Antibiotic A 75943, A-1100
 Antibiotic BE 52211, A-1130
 Antibiotic BE 52211C, A-1137
 Antibiotic ECO 02301, A-1161
 Antibiotic IC 202A, A-1185
 Antibiotic NA 30851A, A-1212
 Antibiotic NFAT 68, A-1213
 Antibiotic PM 94128, A-1229
 Antibiotic Sch 725424, A-1252
 Antibiotic SF 2369, A-1257
 Antibiotic SF 2768, A-1260
 Antibiotic U 62162, A-1298
 Antibiotic USF 19A, A-1305
 Antibiotic Y 05460MA, A-1319
 Antibiotic YN 0165JA, A-1323
 Apicularen A, A-1338
 Aplysinafulvin, A-1348
 Aplysinketal A, A-1353
 Aplysinketal B, A-1354
 Arboreumine, A-1384
 Ariakemicin A, A-1422
 Aspergillazine B, A-1473
 Aspergillazine D, A-1474
 Aspernigrin A, A-1483
 Avenanthramide L, A-1564
 2-Benzamidobenzoic acid, B-67
 Benzastatin A, B-68
 Benzylurea, B-97
 Bisavenanthramide B₅, B-159
 Bisavenanthramide B₆, B-160
N,N'-Bis(2-phenylethyl)urea, B-216
 Botryllamide A, B-256
 Brefelamide, B-289
N-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-*N'*-(3-methoxycarbonylpropyl)urea, B-307
 [2-(3-Bromo-4-hydroxyphenyl)ethyl]ethanedi-
 amide, B-329
N-(4-Bromophenyl)-2,2-diphenylacetamide,
 B-344
 Capsaicin, C-100
 Caracasandiamide, C-108
 Cassipoureamide A, C-184
 Cassipoureamide B, C-185
 Cerulenin, C-286
 Cheliensisine, C-359
N-(4-Chlorobutyl)butanamide, C-396
 Chondrochloren A, C-422
 Ciliatamide A, C-438
 Ciliatamide C, C-439
 Clathrynamide A, C-497
 Clinacoside C, C-526
 Coniothyriomycin, C-609
 Cordyformamide, C-645
 Cruentaren A, C-777
 Cruentaren B, C-778
 Dambullin, D-12
 Dankastatin A, D-15
 Daryamide A, D-82
N,N'-Dibenzylurea, D-302
N-(2,3-Dibromo-4,5-dihydroxybenzyl)-
N'-(3-methoxycarbonylpropyl)urea,
 D-312
 3,5-Dibromo-4-ethoxy-1-hydroxy-4-methoxy-
 2,5-cyclohexadien-1-acetamide, D-316
 3,5-Dibromo-1-hydroxy-4,4-dimethoxy-2,5-cy-
 clohexadiene-1-acetamide, D-318
N-[2-[3,5-Dibromo-4-[3-(3-methyl-1-oxo-2-bu-
 tenyl)amino]propoxy]phenyl]ethyl]-4-hydro-
 xybenzeneacetamide, D-333
 5,5''-Dicapsaicin, D-344
N-[5-[(1,6-Dihydro-2-hydroxy-5-octanamido-6-
 oxo-3-pyridyl)imino]-1,2,5,6-tetrahydro-2,6-
 dioxo-3-pyridyl]octanamide, D-450
N-(2,4-Dihydroxy-3,5,6-trimethylbenzoyl)gly-
 cine, D-681
 3,4-Dimethoxybenzaldehyde *N*-(3,4-dimethoxy-
 benzoyl)hydrazone, D-705
 Dimethylformamide, D-747
 Dioxolamycin, D-806
 Dioxolide A, D-807
N,N'-Diphenylurea, D-818
 Diptaline, D-826
 Diptamine, D-827
 Diptocarpaine, D-828
 Diptocarpidine, D-829
 Elliptinol, E-69
 Entadamide B, E-85
 Erythrococcamide A, E-212
 Erythrococcamide B, E-213
 Erythrococcamide C, E-214
 4-Ethyl-2-(hydroxyimino)-5-nitro-3-hexenamide,
 E-243
 Farinosone C, F-11
 Fumimycin, F-183
 Gerambullin, G-68
 Glaucenamide, G-88
 Glymontanine A, G-128
 Glymontanine B, G-129
 Golmaenone, G-138
 Guamamide, G-191
N-(4-Guanidinobutyl)-2-(4-hydroxyphenyl)-2-
 oxoacetamide, G-202
 Gymnastatin F, G-229
 Gymnastatin L, G-230
 Hamigeroxalamic acid, H-44
 Himeic acid A, H-282
 Himeic acid C, H-283
 Hippospongion C, H-297
 Hippospongion E, H-298
 Homoplantensimide, H-342
 4-[(2-Hydroxy-4,5-dimethoxyphenyl)amino]-3-
 hydroxy-4-oxobutanoic acid, H-466
N-(1-Hydroxyethyl)benzamide, H-485
N-(2-Hydroxyethyl)cinnamamide, H-488
N-(3-Hydroxy-1-oxocyclopent-2-en-2-yl)-3-(4-
 hydroxy-3-methoxyphenyl)propenamide,
 H-651
 Icumazole F, I-15
 IFB Lactam 1, I-18
 Irciniastatin A, I-173
 Jordanine, J-47
 Justiciaamide, J-62
 Kalkitoxin, K-7
 Korormicin A, K-84
 Korormicin B, K-85
 Korormicin C, K-86
 Korormicin D, K-87
 Laccac acid A₁, L-3
 Limaciamine, L-174
 Limocrocin, L-178
 Lipoxamycin, L-189
 Lituarine C, L-217
 Longitubanine A, L-256
 Lorneamide A, L-262
 Malonganenone B, M-75
 Melanocin A, M-154
 Melanocin B, M-155
 Melanocin C, M-156
 Melanocrocin, M-157
 (4-Methoxyphenyl)-*N*-methyl-2-oxothioaceta-
 mide, M-282
 Methyl [(acetylamino)oxy]acetate, M-294
 4-Methyl *N*-[(2,3-dihydro-2-oxo-1*H*-indol-3-
 yl)acetyl]aspartate, M-299
 [(Methylamino)carbonyl]carbamic acid,
 M-385
N-Methylformamide, M-463
 3-Methyl-2-[(2-nonene-4,6,8-triynoyl)amino]bu-
 tanoic acid, M-490
N-Methyl-*N*-(2-phenylethyl)-3-phenyl-2-oxi-
 ranecarboxamide, M-509
 3-(2-Methylphenyl)-3-oxopropanamide,
 M-512
N-(2-Methylpropyl)-3-(3-oxo-1-cyclopenten-1-
 yl)propanamide, M-526
 Microsphaerone B, M-601
 Mirabilin, M-637
 Moiramide A, M-668
 Mortivinacin B, M-706
 Mortivinacin C, M-707
 Neoenactin, N-113
 Neoengleromyacin, N-114
 Nigleroxin, N-194
 Nirananin, N-235
 Norcoronatine, N-291
 Nortrachelogenin amide, N-314
 Ochrobactin C, O-19
 5-Octanoylamino-2,3,6(1*H*)-pyridinetri-
 one, O-62
 Oximidine II, O-164
 Palmerolide A, P-37
 Palmidrol, P-38
 Palytoxin, P-42
 Penidiamide, P-189
 Perspicamide A, P-274
 Petrobactin, P-283
 Phenalamide A1, P-316
 Phenalamide B, P-317
 Phenatic acid A, P-318
 Phomoenamide, P-375
 Physarigin A, P-395
 Physarigin B, P-396
 Pipyahyne, P-477
 Pitiamide A, P-487
 Platencin, P-511
 Platensimide A, P-512
 Platensimycin, P-513
 Primocarcin, P-630
 Protochelin, P-661
 Proximicin A, P-676
 Pseudotrienoic acid A, P-730
 Pseudotrienoic acid B, P-731
 Purealidin O, P-815
 Reductionmycin, R-34
 Rubesamide, R-142
 Sagittamide A, S-15
 Saliniketal A, S-26
 Salmochelin X, S-33
 1,2-Secodysidamide, S-185
 Semiplenamide C, S-223
 Semiplenamide D, S-224
 Semiplenamide E, S-225
 Somocystinamide A, S-365
 Sorbicillactone A, S-382
 Sperabillin C, S-389
 Spongiacysteine, S-471
 Stelletazole A, S-507
 Stephanosporin, S-539
 Streptomycamide A, S-567
 Streptomycamide B, S-568
 Stresgenin B, S-575
 Suimin, S-632
 Sulfinemycin, S-633
 Sulfobacin A, S-634
 Sulfobacin B, S-635
 Synechobactin A, S-657
 Taveuniamide E, T-40
 Topostin B553, T-430
 Topostin D 640, T-431
 Trichanthin A, T-478
 Trichostatin A, T-496
 1,4,5-Trihydroxy-2-cyclopentene-1-carboxa-
 mide, T-536
 Variceramides, V-25

Azo and azoxy alkaloids

Antibiotic DC 8118A, A-1159
 Antibiotic LL-BH872 α , A-1199
 Deoxyrubroflavin, D-229
 Elaiomycin, E-61
 Jietacin A, J-38
 Jietacin B, J-39
 Lyophyllin, L-368
 Maniwamycin B, M-85
 Valanimycin, V-5

Simple pyrrolidine alkaloids

5-Acetyl-3,4-dihydro-2*H*-pyrrole, A-49
 Agaricone, A-175
 Amphiassterin C₁, A-932
 Amphiassterin C₂, A-933
 Antibiotic PI 091, A-1228
 Broussonetine I, B-359
 Broussonetine R, B-363
 Broussonetine U, B-364
 Broussonetine W, B-365
 Broussonetinine A, B-367
 Codonopsine, C-557
 Conioidine A, C-607
 2,3-Dehydro- β -stachydrine, D-169
 3,4-Dihydro-5-propanoyl-2*H*-pyrrole, D-493
 3,4-Dihydro-2*H*-pyrrole, D-500
 2-(3,4-Dihydro-2*H*-pyrrol-5-yl)pyridine, D-504
 2,4-Dihydroxy-3,3,4-trimethyl-5-oxo-2-pyrrolidinedicarboxylic acid, D-682
 Dysibetaine, D-964
 3-Ethyl-2-pyrrolidinone, E-270
 Fulvanine A, F-166
 Gerrardine, G-72
 3-Hydroxy-2,3-dimethyl-5-oxo-2-pyrrolidinedicarboxylic acid, H-472
 2-(2-Hydroxypropyl)-1-methyl-5-(2-oxopropyl)pyrrolidine, H-704
 5-Hydroxy-2-pyrrolidinone, H-719
 3-Hydroxystachydrine, H-740
 Hygrine, H-759
 Involucratine, I-162
 Kainic acid, K-4
 Lepiotin A, L-110
 Leucoagaricone, L-154
 Mescaline succinimide, M-222
 3-(Methoxymethylene)-2-pyrrolidinedithione, M-266
 1-(3-Methylbutyl)-3-pyrrolidinemethanol, M-421
 2-Methyl-4-(3-methyl-1-pyrrolidinyl)-1-butanol, M-484
 1-Methylpyrrolidine, M-543
 1-Methyl-2-pyrrolidinone, M-546
 2',3-(*N*-Methylpyrrolidinyl)hygrine, M-548
 2',4-(*N*-Methylpyrrolidinyl)hygrine, M-549
 Morusimic acid B, M-708
 Pipericyclamide, P-449
 2-Pyrrolidineacetic acid, P-950
 1-Pyrrolidinedicarboxamide, P-951
 Pyrrolidine, P-949
 2-Pyrrolidinone, P-956
 Sarmentamide B, S-87
 Scalusamide A, S-117
 Sessiline, S-269
 Squamolone, S-487
 Uncinine, U-24
 Villatamine A, V-104

Chromone alkaloids

Anhydroperipentamine, A-1033
 Cassiadinine, C-180
 4-(5,7-Dihydroxy-2-methyl-4-oxo-4*H*-1-benzopyran-6-yl)-2-piperidinone, D-629
 Isoschumanniphytine, I-314
 Rohitukine, R-121
 Schumagnine, S-144
 Schumannificine, S-145
 Schumanniphytine, S-146
 Tubastraine, T-654

Flavonoid alkaloids

Aquiledine, A-1369
 Capitavine, C-94
 Cheliensisine, C-359
 Davallioside A, D-91
 5,7-Dihydroxy-8-(2-piperidinyl)flavone, D-662
 Ficine, F-53
 Isoaquileidine, I-190
 Isoficine, I-233
 Isopiscerythramine, I-267
 Kopsirachin, K-81
 Lilaline, L-173
 Phyllospadine, P-389
 Piscerythramine, P-483
 Vochysine, V-189

Nicotine-like alkaloids

Cotinine, C-708
 Myosmine, M-792

Miscellaneous pyrrolidine alkaloids

Aburatubolactam B, A-17
 Acromelic acid A, A-118
 Acromelic acid B, A-119
 Aglaforbesine A, A-196
 Aglaine B, A-199
 Aglaxiflorin C, A-201
 Aglaxiflorin D, A-202
 Althiomycin, A-664
 Amathamide A, A-677
 Amathamide E, A-678
 Amathaspiramide C, A-679
 Ampullicin, A-951
 Andrimide, A-1004
 Antibiotic AB 3217A, A-1112
 Antibiotic BE 54476, A-1132
 Antibiotic BE 67251, A-1134
 Antibiotic BU 4514N, A-1140
 Antibiotic F 10C, A-1165
 Antibiotic FD 839, A-1168
 Antibiotic FR 900148, A-1176
 Antibiotic MBP 039-06, A-1204
 Antibiotic MBP 049-13, A-1205
 Antibiotic SF 2312, A-1256
 Antibiotic TAN 1813, A-1278
 Antibiotic WS 1921, A-1314
 Ascosalipyrrolidinone B, A-1462
 Axinellamide, A-1573
 Azaspirene, A-1601
 Bisavenanthramide B₁, B-157
 Bisavenanthramide B₂, B-158
 1,1'-[[2,4-Bis(6-methoxy-1,3-benzodioxol-5-yl)-1,3-cyclobutanediyl]dicarbonyl]bispyrrolidine, B-196
 Boneratamide A, B-246
 Boneratamide B, B-247
 Brevigellin, B-295
 Broussonetine G, B-360
 Broussonetine H, B-361
 Broussonetine Z, B-366
 Bulgecin A, B-402
 Bulgecin B, B-403
 Bulgecin C, B-404
 Caesalpinine A, C-12
 Cantharimide, C-89
 Carbonarin C, C-139
 Carbonarin D, C-140
 Carbonarin E, C-141
 Celesticetin, C-239
 Chelelactam, C-354
 Citridone C, C-480
 Cuscohygrine, C-818
 Darlinine, D-76
 Desdamethine, D-232
 2,5-Dialkylpyrrolidines, D-246
 Dihydrocuscocohygrine, D-413
 3,4-Dihydro-5-(5-hexenyl)-2-nonyl-2*H*-pyrrole, D-435
 1,5-Dihydro-5-hydroxy-3-methyl-1-(5-oxo-2-pyrrolidinyl)-2*H*-pyrrol-2-one, D-448
 1-(2,5-Dihydro-4-methyl-5-oxo-1*H*-pyrrol-2-yl)-4-methyl-5-oxoproline, D-474
 1,5-Dihydro-1-[1-oxo-3-(3,4,5-trimethoxyphenyl)propyl]-2*H*-pyrrol-2-one, D-489
 3,4-Dihydro-5-(2-pyridinyl)-2*H*-pyrrole-2-carboxylic acid, D-496
 Dipiperamide C, D-820
 Domoic acid, D-912
 Domoilactone A, D-913
 Donaxamine, D-915
 Dysidamide C, D-966
 Dysideapyrrolidone, D-969
 Dysidine, D-973
 Eliamide, E-65
 Epicoccamide, E-88
 Epolactaene, E-106
 Ethesdanine, E-229
 4-Ethoxyhexahydro-3*a*-hydroxy-1*H*-indole-2,6-dione, E-231
 Fischerellin A, F-60
 Fischerellin B, F-61
 Forbagline A, F-127
 Fusarin A, F-226
 Fusarin C, F-227
 Fusarin F, F-228
 Gigantamide A, G-74
 Heliotropamide, H-96
 2-(5-Hexenyl)-3,4-dihydro-5-nonyl-2*H*-pyrrole, H-259
 Hopeanine, H-352
 Huperzine W, H-387
 Hypercratine, H-770
 5-Imino-2*H*-pyrrol-2-ol, I-51
 Ircinamine, I-171
 Isobutyropyrrrothine, I-200
 Isodomoic acid A, I-228
 Isoficine, I-233
 Isopiperolein B, I-266
 Isothapsakone A, I-335
 Jamaicamide A, J-15
 Kaitocephalin, K-5
 Kikumycin A, K-43
 Leptanthine, L-115
 Lilaline, L-173
 Lucentamycin B, L-275
 Lucilactaene, L-283
 Makalika ester, M-53
 Maltoxazine, M-77
 Marine *Streptomyces* C₃₀H₃₈N₂O₅ lactam, M-105
 Mescaline citrimide, M-220
 Mescaline isocitrimide lactone, M-221
 Mirabimide E, M-644
 Moiramide B, M-669
 Munronin D, M-733
 Nigramide G, N-201
 Nigramide M, N-206
 Nordomoic acid, N-297
 Orosomycin, O-124
 Palasimide, P-17
 Palauimide, P-19
 Pandamarilactam 3*x*, P-59
 Pandamarilactonine A, P-63
 Peripentadenine, P-262
 Phygrine, P-384
 Piriferine, P-479
 Plakoridine A, P-505
 Plakoridine B, P-506
 Polyandrocarpidine A, P-544
 Polyandrocarpidine C, P-545
 Polycitrin A, P-559
 Ponapensin, P-565
 Proferrosamine B, P-637
 Pulchellalactam, P-784
 Pyroglutamylglucosamine, P-937
 Pyronopaline, P-939
 Pyrrocidine B, P-940
Asteronotus cespitosus Pyrrolidone, P-959
 Sargassumolactam, S-85
 Saussureamine A, S-105
 Saussureamine D, S-107

Schizokinen A, S-134
 Spiroarogenic acid, S-424
 Spirostaphylotrichin E, S-459
 Spirostaphylotrichin G, S-460
 Spirostaphylotrichin K, S-461
 Spirostaphylotrichin M, S-462
 Spirostaphylotrichin O, S-463
 Spirostaphylotrichin Q, S-464
 Spirostaphylotrichin R, S-465
 Spirostaphylotrichin S, S-466
 Spirostaphylotrichin T, S-467
 Subereatin, S-620
 Synerazol, S-660
 Terragine A, T-99
 Terragine B, T-100
 Thapoxepine A, T-348
 Thapsakin B, T-349
 Thiomyacin, T-390
 Tinctormine, T-413
 Trikendiol, T-592
 Triticone A, T-625
 Turrapubesin B, T-686
 Variotin, V-34
 Vasicol, V-39
 Vibrioferrin, V-97
 Violacein, V-142
 Viomycin, V-144
 Viriditin, V-155
 Ypaoamide, Y-17
 Zopfiellamide B, Z-34

Tetramic acids

Aburatubolactam A, A-16
 3-Acetyl-5-benzyl-2,4-pyrrolidinedione, A-43
 Aflastatin A, A-168
 Altamycin A, A-659
 Alteramide A, A-661
 Altersetin, A-663
 Amycomycin, A-958
 Anchinopeptide A, A-976
 Ancorinoside A, A-994
 Ancorinoside B, A-995
 Ancorinoside C, A-996
 Ancorinoside D, A-997
 Antibiotic 2158, A-1084
 Antibiotic AB 4015B, A-1113
 Antibiotic AB 4063B, A-1114
 Antibiotic CJ 17572, A-1149
 Antibiotic CJ 21058, A-1150
 Antibiotic F 1778, A-1163
 Antibiotic FD 838, A-1167
 Antibiotic L 755807, A-1197
 Antibiotic LL-49F233 α , A-1201
 Antibiotic NG 391, A-1215
 Antibiotic PF 1052, A-1225
 Antibiotic Sch 210971, A-1250
 Antibiotic TAN 883, A-1274
 Antibiotic TAN 1263, A-1275
 Antibiotic TMC 260, A-1290
 Apidionene, A-1339
 Aurantoside A, A-1550
 Aurantoside C, A-1551
 Aurantoside E, A-1552
 Aurantoside F, A-1553
 Aurantoside G, A-1554
 5-Benzyl-2,5-dihydro-4-hydroxy-3-(2,4,6-octatrienyl)-1H-pyrrol-2-one, B-85
 5-Benzyl-3-(2,4,6,8,10-dodecapentaenyl)-2,5-dihydro-4-hydroxy-1H-pyrrol-2-one, B-87
 Blasticidin A, B-230
 Bripiodionene, B-298
 Cinnabaramide D, C-450
 Cinnabaramide F, C-451
 Cissetin, C-468
 Coniosetin, C-608
 Cryptocin, C-784
 Cylindramide, C-924
 Delaminomycin A, D-179
 Discoderme, D-844
 Epicoccarine B, E-89
 Equisetin, E-142

Erythroskyrin, E-217
 Fuligorubin A, F-165
 Harzianic acid, H-67
 Integramycin, I-158
 Isodomoic acid G, I-229
 Janolusimide, J-18
 Lactacystin, L-6
 α -Lipomycin, L-186
 Lydicamycin, L-353
 Macrocidin A, M-13
 Magnesidin, M-42
 Malonomycin, M-76
 Militarione B, M-611
 Myceliothermophin A, M-783
 Nocamycin I, N-264
 Nortenuazonic acid, N-311
 Oleficin, O-79
 Oteromycin, O-134
 Oxasetin, O-150
 Pachydermin, P-5
 Penicillenol A₁, P-188
 Phomasetin, P-373
 Physarorubinic acid A, P-397
 Physarorubinic acid B, P-398
 Polycephalin C, P-555
 Pramanicin, P-584
 Preapiodionene, P-589
 Ravenic acid, R-30
 Rubroside G, R-153
 Rubroside H, R-154
 Spylidone, S-484
 Streptolydigin, S-566
 Tenuazonic acid, T-74
 Tirandalydigin, T-416
 Tirandamycin A, T-417
 Vancocresmycin, V-17
 Vermelhotin, V-76
 Vermisporin, V-77
 Xanthobaccin A, X-6
 Zopfiellamide A, Z-33

Tropane alkaloids

6-Angeloyloxyditropan-3-yl itaconate, A-1008
 Apoatropine, A-1359
 Apohyoscine, A-1364
 8-Azabicyclo[3.2.1]octane-2,6-diol, A-1582
 Belladonnine, B-56
 Bonabiline A, B-244
 Brugine, B-369
 Chalcostrobamine, C-341
 Consiculine, C-622
 Convolvidine, C-632
 Convolvine, C-633
 Darlingine, D-75
 Fastudine, F-19
 Fastunine, F-20
 Fastusidine, F-21
 Ferrugine, F-46
 Grahamine, G-153
 Merresectine C, M-207
 Merresectine D, M-208
 Merresectine F, M-209
 8-Methyl-8-azabicyclo[3.2.1]octan-3-one, M-394
 Mooniine A, M-700
 Physoperuvine, P-399
 Schizanthine A, S-129
 Schizanthine E, S-130
 Schizanthine G, S-131
 Schizanthine X, S-128
 Scopadonnine, S-157
 Scopine, S-158
 Scopoline, S-160
 Strobamine, S-586
 Strobiline, S-587
 Subhirsine, S-622
 3 α -(4,4,4-Trichloro-2-butyryloxy)tropane, T-479
 α -Truxilline, T-634
 β -Truxilline, T-635

Simple pyrrolizidine alkaloids

Crotalaria gorensis Alkaloid C, A-437
Jacmaia incana Alkaloid, A-307

Amabiline, A-669
 Anchustrigosine, A-977
 Anodendrine, A-1065
 Antibiotic CJ 16264, A-1147
 Antibiotic CJ 16367, A-1148
 Antibiotic UCS 1025B, A-1301
 Assamicadine, A-1507
 Bohemamine, B-242
 Broussonetine N, B-362
 Callimorphine, C-36
 Cornucervine, C-655
 Crotalarine lactone, C-772
 Crotanecine, C-775
 Cruentine A, C-779
 Cruentine B, C-780
 Curassavine, C-810
 Danaidal, D-14
 Dihydroxytriangularine, D-680
 Doriasenine, D-922
 Echihumiline, E-10
 Echimidine, E-11
 Echinatine, E-12
 Echiumine, E-31
 Echiupinine, E-32
 Epohelmin A, E-105
 1,2-Epoxy-1-hydroxymethylpyrrolizidine, E-133
N-Ethylthastanecine trachelanthate, E-241
 Europine, E-299
 Floridimine, F-92
 Grandifoline, G-158
 Hackelidine, H-4
 Heliospathuline, H-93
 Heliosupine, H-94
 Heliotrine, H-95
 1-Hydroxymethylpyrrolizidine methyl 2-*O*-acetyl-2-isopropylmalate, H-631
 Indicine, I-68
 Indicinine, I-69
 Intermedine, I-160
 Ipanguline D₃, I-166
 Jenamidine B, J-34
 Keitine, K-28
 Lankongensisine A, L-32
 Lankongensisine B, L-33
 Lasiocarpine, L-47
 Lithosenine, L-212
 Loline, L-234
 Loroquine, L-263
 Megalanthonine, M-146
 Minalobine O, M-626
 Minalobine R, M-627
 3-(8-Nonenyl)-5-(1-propenyl)pyrrolizidine, N-284
 Otonecine, O-136
 Paludosine, P-39
 Planchonelline, P-510
 Procerine, P-632
 Punctanecine, P-798
 Retronecine 9-(2,3-dihydroxy-2-hydroxymethyl-butanoate) 7-senecioate, R-61
 Sarracine, S-90
 Scorpioidine, S-163
 Strigosine, S-585
 Trachelanthamine, T-443
 Triangularine, T-454
 Uplandicine, U-40

Pyrrolizidine alkaloids (macrocyclic lactones)

Adonifoline, A-148
 Anacrotine, A-967
 Aucherine, A-1538
 Axillarine, A-1572
 Bisline, B-194
 Bulgarsenine, B-401
 Callosine, C-39
 Clivorine, C-533
 Croaegyptine, C-760
 Croalbidine, C-761
 Crobarbatine, C-762

Crocandine, C-764
 Cronaburmine, C-767
 Crodopine, C-768
 Crosemperine, C-769
 Crotafoline, C-770
 Crotalarine, C-771
 Crotaleschenine, C-773
 Crotananine, C-774
 Dicrotaline, D-363
 Emiline, E-73
 Erucifoline, E-184
 8-Ethoxy-3-oxo-1,2-dehydroretrosine, E-232
 Fulvine, F-168
 Grantaline, G-167
 Grantianine, G-168
 Gynuramine, G-233
 Hastacine, H-69
 Helindicine, H-92
 Inaequidenine, I-55
 Jacobine, J-2
 Jacoline, J-3
 Jaconine, J-4
 Ligularine, L-169
 Ligularizine, L-170
 Madurensine, M-36
 17-Methylparsonsianidine, M-502
 Monocrotaline, M-688
 Nemorensine, N-89
 Nilgirine, N-219
 Onetine, O-92
 Orosenine, O-137
 Parsonsianidine, P-96
 Parsonsianine, P-97
 Parsonsine, P-98
 Petasitenine, P-277
 Platyphylline, P-516
 Pterophorine, P-770
 Retrosine, R-62
 Retusamine, R-64
 Retusine, R-65
 Rosmarinine, R-133
 Sceleratine, S-122
 Senaetnine, S-232
 Seneciocannabine, S-237
 Senecionine, S-238
 Senecioracene, S-239
 Seneciphylline, S-240
 Senecivernine, S-241
 Senkirkine, S-245
 Swazine, S-651
 Syneilesine, S-658
 Trichodesmine, T-488

Miscellaneous pyrrolizidine alkaloids

Senecio renardi Alkaloid 4, A-332
 Alkaloid S-F, A-599
Cytisus laburnum Alkaloid, A-285
Gynura segetum Alkaloid, A-303
Heliotropium europaeum Alkaloid, A-304
Senecio borysthenticus Alkaloid, A-331
 Antibiotic PF 1018, A-1224
 Campestrine, C-64
 Carategine, C-118
 Cassipourine, C-186
 2,3-Dihydro-1-oxo-1*H*-pyrrolizine-3-carboxylic acid, D-488
 Franchetine, F-148
 Graminifoline, G-154
 Ligudentine, L-168
 Lolidine, L-232
Senecio doria Necine base, N-86
 Nitropolyzonamine, N-252
 Norsenecicaudatin 9-*O*-senecioate, N-309
 Paucalinaline, P-127
 Pictumine, P-420
 Ruzorine, R-173
 Salinosporamide C, S-29
 Senecicaudatal hemiacetal, S-236
 Spiropyrrrolizidine 252, S-437
 Talarotoxin, T-17
 Thelepogine, T-354

Thesine, T-364
 Turkestanine, T-683
 Yuremamine, Y-22

Anabasin-like alkaloids

Anabasamine, A-963
 Astrocasine, A-1524
 Astrophylline, A-1525
 2,3'-Bipyridine, B-145
 Gramodendrine, G-155
 Jussiaeine A, J-61
 1',3,4,4',5',6,6'-Octahydro-2,3'-bipyridine, O-40
 3,4,5,6-Tetrahydro-2,3'-bipyridine, T-144
 Virgivarine, V-150

Lobelia-type alkaloids

Andrachcinidine, A-1000
 Aspertine B, A-1487
 8,10-Diethyllobelidiol, D-391
 Dihomosedinone, D-403
 Homosedinone, H-347
 2-(2-Hydroxybutyl)-6-(2-hydroxypropyl)-1-methylpiperidine, H-434
 Lobinaline, L-220
 Lobinanidine, L-221
 Pyrrolsedamine, P-970
 Sederine, S-211
 Sediene, S-213
 Sedinine, S-214
 Sedinone, S-215

Simple piperidine alkaloids incl. piperidides

6-Acetyl-1,2,3,4-tetrahydropyridine, A-76
Cassia Alkaloid D, A-477
 5-Amino-5-deoxygalactose, A-726
 5-Amino-5-deoxyglucose, A-727
 5-Amino-5-deoxymannose, A-728
 Anahygrine, A-971
 Anatalline, A-973
 Andrachcinidine, A-1000
 Awaine, A-1568
 Azcarpine, A-1604
 Azimine, A-1608
 Batzallaside A, B-49
 Batzallaside B, B-50
 Batzallaside C, B-51
 Calyxamine A, C-60
 Carpaine, C-168
 3-Chloro-4-hydroxy-2-piperidinone, C-408
 Ciliatamide C, C-439
 Corydendramine A, C-680
 Coumaperine, C-710
 Cryptophorine, C-788
 Cryptophorimine, C-789
 Dienomycin C, D-386
 5,6-Dihydro-2(1*H*)-pyridinone, D-495
 1-Ethyl-3-(2-methylbutyl)piperidine, E-253
 Girgensonine, G-80
 Haloxylone B, H-39
 Hoveine, H-365
 2-Iminopiperidine, I-49
 Irnigaine, I-177
 Juliprosine, J-57
 Jussiaeine A, J-61
 Lansamide 2, L-39
 Leptophyllin A, L-119
 Leptophyllin B, L-120
 2-Methyl-6-(8-heptadecenyl)piperidine, M-470
 2-Methyl-6-nonylpiperidine, M-492
 1-Methylpiperidine, M-515
 Micropine, M-600
 Morusimic acid D, M-709
 7-Oxa-3-azabicyclo[4.1.0]heptan-2-one, O-146
 8-[(2-Oxo-3-piperidinyl)amino]-8-oxooctyl 5,9-anhydro-2,3,8-trideoxy-8-(5-hydroxy-4-methyl-2-hexenyl)-3-methyl-DL-glycero-LD-allo-non-2-enoate, O-197
 Penasulfate A, P-178

6-Pentyl-3-piperidinol, P-236
 Pipbinine, P-441
 Piperettine, P-452
 1-Piperidinecarboxaldehyde, P-455
 Piperidine, P-454
 4-Piperidinone, P-459
 Pipernonaline, P-464
 Piperpense, P-468
 Pipertipine, P-470
 Piperundecaldine, P-471
 Pipsaeedine, P-475
 Pleurospermine, P-522
 Prosopine, P-656
 Pseudodistomin B, P-712
 Pseudodistomin F, P-713
 Secojuliprosopinal, S-188
 Smipine, S-330
 Spicigerine, S-409
 Sulphostin, S-639
 1,2,3,4-Tetrahydro-6-propanoylpiperidine, T-217
 2,2,6,6-Tetramethyl-4-piperidinone, T-283
 Uoamine A, U-38

Xestospongins

Aragupetrosine A, A-1371
 Araguspongine C, A-1372
 Araguspongine F, A-1373
 Araguspongine J, A-1374
 Petrosine, P-290
 Upenamide, U-39
 Xestospongine A, X-24
 Xestospongine B, X-25

Miscellaneous piperidine alkaloids

Actiketol, A-124
 Actiphenol, A-129
 Anthosamine A, A-1079
 Antibiotic 593A, A-1089
 Antibiotic BR 040, A-1139
 Antibiotic HA 23, A-1184
 Antibiotic L 654040, A-1194
 Antibiotic NK 30424A, A-1216
 Antibiotic SB 219383, A-1241
 Arenosclerin A, A-1404
 Arenosclerin E, A-1405
 Awajanomycin, A-1569
 Azaspiracid 1, A-1600
 Banyaside A, B-18
 Banyaside B, B-19
 1,1'-Bipiperidine, B-141
 2,4-Bis(2-methoxy-4,5-methylenedioxyphenyl)-1,3-cyclobutanecarboxylic acid dipiperidide, B-198
 Broussonetine I, B-359
 Calvine, C-41
 Campedine, C-63
 Carpaine, C-168
 Cenocladamide, C-245
 Cepaciamide A, C-247
 Cepaciamide B, C-248
 Chabamide, C-316
 Cortamidine oxide, C-664
 Cycasthioamide, C-831
 Cycloguineense A, C-866
 Decahydro-8-(1-piperidinylmethyl)isoquinoline, D-113
 5,8-Dihydro-1,8,8-trimethyl-1*H*-pyrano[3,4-*b*]pyridine-4,6-dione, D-524
 Dioscorine, D-798
 Dipiperamide A, D-819
 Dipiperamide C, D-820
 Dipiperamide D, D-821
 Dipiperamide E, D-822
 Dorrigocin A, D-923
 Dorrigocin B, D-924
 Dumetorine, D-951
 Epiderstatin, E-92
 Fagopyrine, F-6
 Flabelliformine, F-74
 Grandisine G, G-166
 Griffithine, G-176

Haliclamine B, H-16
 Haliclonyclamine A, H-19
 Halicyclamine A, H-22
 Halicyclamine B, H-23
 Isomigrastatin, I-245
 Juliflorine, J-56
 Julocrotine, J-58
 Kopsone, K-82
 Kuraramine, K-108
 Lactimidomycin, L-7
 4-(1-Methyl-2-piperidinyl)- β -carboline, M-518
 9-Methylstreptimidone, M-561
 Migrastatin, M-606
 Myrionine, M-797
 Nigramide A, N-196
 Nigramide B, N-197
 Nigramide C, N-198
 Nigramide D, N-199
 Nigramide E, N-200
 Nigramide G, N-201
 Nigramide H, N-202
 Nigramide I, N-195
 Nigramide J, N-203
 Nigramide K, N-204
 Nigramide L, N-205
 Nigramide M, N-206
 Nigramide N, N-207
 Nigramide P, N-209
 Nigramide Q, N-210
 Nigramide R, N-211
 Nigramide S, N-212
 Nupharamine, N-340
 Pandamarilactone 1, P-60
 Pandamarilactone 31, P-61
 Pandamarilactone 32, P-62
 Pinnaic acid, P-435
 Piperchabamide C, P-448
 Pipericyclobutanamide A, P-450
 Pipericyclobutanamide B, P-451
 Pipermethystine, P-463
 Piplartine, P-473
 Polonicumtoxin C, P-539
 Pulcherrimine, P-787
 Quinaldopeptin, Q-11
 Sesbanimide C, S-261
 Sollasin B, S-356
 Streptimidone, S-563
 Streptovitacin D, S-574
 Strictimine, S-581
 2,3,4,5-Tetrahydro-6-(1-propenyl)pyridine, T-218
 Trichoponamic acid, T-493
 Tubastraine, T-654

Lythraceae alkaloids

Heimine, H-89
 Lagerine, L-11
 Lasubine I, L-53
 Lythramine, L-376
 Lythrancine I, L-377
 Lythranidine, L-378
 Lythrumine, L-379
 Sarusubine A, S-91
 Verticillatine, V-88
 Vertine, V-95

Quinolizidine alkaloids (two rings)

Aphyllic acid, A-1335
 Cadiamine, C-8
 Cermizine D, C-284
 Clathryimine A, C-496
 Clavepictine B, C-513
 7-Demethyldeoxynupharidine, D-201
 Deoxynupharidine, D-227
 1-Ethyl-4-(4-pentynyl)quinolizidine, E-264
 1-Ethyl-4-(2-propenyl)quinolizidine, E-265
 Himeradine A, H-284
 Homopumiliotoxin 321B, H-344
 Homopumiliotoxin 223G, H-345
 4-Hydroxyepilupinine, H-482
 Lasubine I, L-53

Leontiformine, L-97
 Mamanine, M-79
 1-Methyl-4-(2-penten-4-ynyl)quinolizidine, M-507
 Pictamine, P-418
 Plumerinine, P-526
 Pohakuline, P-538
 Sarusubine A, S-91
 Senepodine A, S-242
 Senepodine G, S-243
 Senepodine H, S-244
 Sophoattarine, S-370
 Sopholupanizidone, S-374

Quinolizidine alkaloids (three rings)

Albine, A-243
 Alkaloid LC2, A-554
 12-Cytisineacetic acid, C-939
 5-(12-Cytisinylmethyl)-6-hydroxylupanine, C-940
 Dimethamine, D-704
 Hupeol, H-372
 3-Hydroxy-11-norcytisine, H-644
 12,12'-Methylenedicytisine, M-441
 Sophoattarine, S-370
 Sophohejrine, S-373
 Sophorasine A, S-378
 Sophosalimine, S-381
 Tetrahydrocytisine, T-149
 Thermopsidine, T-362
 Tinctorine, T-412
 Tonkinensine A, T-428
 Tonkinensine B, T-429
 Tsukushinamine A, T-651
 Virgildone, V-152

Quinolizidine alkaloids (four rings)

Chamaecytisus Alkaloid C₁, A-432
Ammodendron karelinii Alkaloid, A-271
Cytisus ruthenicus Alkaloid, A-286
 Aloperine, A-637
 Anagryne, A-970
 Aphylline, A-1337
 Baptifoline, B-20
 Calpurmenine, C-40
 Camoensidine, C-61
 Camoensine, C-62
 Clathrotropine, C-495
 Conolline, C-619
 5-(12-Cytisinylmethyl)-6-hydroxylupanine, C-940
 Darvasamine, D-79
 Darvasine, D-80
 Darvasoline, D-81
 Dasycarpine, D-85
 5,6-Dehydrolupanine, D-161
 11,12-Dehydrosparteine, D-168
 10,13-Dihydroxylupanine, D-608
 10,17-Dioxosparteine, D-810
 Dithermamine, D-871
 11-Epileontidane, E-94
 Goebeline, G-137
 Hydroxyaphylline, H-405
 13-Hydroxylupanine, H-554
 12-Hydroxy-16-methoxy-11,12,13,14-tetrahydrocamoensine, H-571
 10-Hydroxymethylsparteine, H-634
 7-Hydroxysparteine, H-738
 Isoleontalbine, I-241
 Isomatrine, I-244
 Lehmannine, L-81
 Leontalbine, L-94
 Leontismidine, L-98
 Leontismine, L-99
 Matrine, M-121
 Monspessulanine, M-695
 Multiflorine, M-731
 Ormosine, O-116
 17-Oxolupanine, O-188
 Podopetaline, P-529
 Retamine, R-56

Sophazrine, S-369
 Sophocarpine, S-371
 Sophohejrine, S-373
 Sopholupanizidone, S-374
 Sophoramine, S-376
 Sophoranol, S-377
 Sophorbenzamine, S-379
 Sophosalimine, S-381
 Templetine, T-68
 Termine, T-89
 Thermopsamine, T-361
 Virgiline, V-153

Cylindricine alkaloids

Cohirsine, C-563
 Cohirsinine, C-564
 Cohirsitinine, C-566
 Cylindricine B, C-925
 Cylindricine G, C-927
 Cylindricine J, C-928
 Fascicularine, F-15
 Lepadiformine B, L-102
 Polycitorol A, P-557
 Polycitorol B, P-558

Miscellaneous quinolizidine alkaloids

Sophora Alkaloid 9, A-333
Chamaecytisus Alkaloid C₂, A-433
 Alkaloid LC7, A-555
 Alkaloid LV1, A-561
Lupinus Alkaloid P1, A-577
Lupinus Alkaloid P2, A-578
 Antibiotic A 58365B, A-1104
 Argentinamine, A-1411
 Argentine, A-1412
 Halichlorine, H-13
 Homopiaptanthine, H-341
 Homothermopsine, H-349
 Lupilaxine, L-303
 Ormojanine, O-114
 Ormosinine, O-117
 Panamine, P-53
 Saraine 1, S-69
 Saraine 2, S-70
 Saraine 3, S-71
 Sarothamnine, S-88
 Virgilidine, V-151

Azepine alkaloids

Aaptosamine, A-3
 Amamistatin A, A-672
 Antibiotic A 500359A, A-1102
 Balanol, B-9
 Bengamide K, B-59
 Bengamide Z, B-60
 Capuramycin, C-103
 Chalciporone, C-340
 Circinatin, C-462
 4,8-Diphenyl-5*H*-1,3-dioxolo[4,5-*d*]azepine-5,7(6*H*)-dione, D-816
 Drazepinone, D-934
 Formobactin, F-130
 Isobengamide E, I-195
 Montiporyne E, M-698
 Muscaflavin, M-770
 Mycobactin T, M-788
 Periconin B, P-258
 Peritoxin A, P-266
 Sollasin C, S-357

Nicotinic acid derived alkaloids

Anatalline, A-973
 3-Butylpyridine, B-444
 3-Carboxy-1-(carboxymethyl)pyridinium betaine, C-144
 3-Carboxy-1-methylpyridinium betaine, C-149
 Euphosalicin, E-295
 Himeic acid C, H-283

Metanicotine, M-232
 4-Methyl-2,3'-bipyridine, M-403
 4-Methyl-3,3'-bipyridine, M-404
 5-Methyl-2,3'-bipyridine, M-405
 Miniatine, M-629
 Precatorine, P-593
 3-Pyridinecarboxamide, P-895
 3-Pyridinecarboxylic acid, P-897
 2,3-Pyridinedicarboxylic acid, P-900
 Ricinidine, R-95
 Ricinine, R-96
 Rubradirin B, R-146
 1,2,5,6-Tetrahydro-1-methyl-3-pyridinecarboxylic acid, T-205
 1,2,3,6-Tetrahydro-4-pyridinecarboxylic acid, T-222
 1,2,5,6-Tetrahydro-3-pyridinecarboxylic acid, T-223

Other pyridine alkaloids

Acalyphin, A-22
 2-Acetyl-4-methylpyridine, A-63
 2-Acetyl-6-methylpyridine, A-64
 3-Acetyl-2-methylpyridine, A-65
 3-Acetyl-4-methylpyridine, A-66
 4-Acetyl-2-methylpyridine, A-67
 4-Acetyl-3-methylpyridine, A-68
 5-Acetyl-2-methylpyridine, A-69
 4-Acetyl-6-methyl-2(1*H*)-pyridinone, A-70
 2-Acetylpyridine, A-73
 Acromolactone B, A-111
 Acromelic acid A, A-118
 Acromelic acid B, A-119
 Agelongine, A-193
 Akanthomycin, A-224
 Aleicide B, A-257
 Aleicide C, A-258
 Alkaloid CB2, A-458
 Alkaloid LA 4, A-548
 Alkaloid LA 5, A-549
 Alkaloid LA 6, A-550
 Alkaloid LA 9, A-551
Callispongia fibrosa Alkaloid, A-281
 2-Amino-4,6-dichloropyridine, A-729
 2-Amino-4-methylpyridine, A-845
 4-Amino-2,3-pyridinedicarboxylic acid, A-901
 Amphikuemin, A-943
 Amphimedoside A, A-945
 Amphimedoside B, A-946
 Amphimedoside C, A-947
 Amphimedoside E, A-948
 Amphitoxin, A-950
 4,6'-Anhydrooxysporidinone, A-1031
 Anibine, A-1037
 Antibiotic A 33853, A-1098
 Antibiotic B 90063, A-1123
 Antibiotic CJ 16196, A-1146
 Antibiotic CJ 16170, A-1142
 Antibiotic CJ 16171, A-1143
 Antibiotic CJ 16173, A-1144
 Antibiotic CJ 16174, A-1145
 Antibiotic NG 311, A-1214
 Antibiotic NK 170204B, A-1217
 Antibiotic PF 1140, A-1226
 Antibiotic SPF 32629A, A-1266
 Antibiotic SPF 32629B, A-1267
 Antibiotic TMC 69, A-1289
 Antibiotic WS 75624A, A-1315
 Antibiotic WS 75624B, A-1316
 Apiosporamide, A-1340
 Arecolidine, A-1400
 Aspernigrin A, A-1483
 Aspernigrin B, A-1484
 Aspyridone A, A-1506
 Austrodimerine, A-1557
 Bassianin, B-28
 2,3'-Bipyridine, B-145
 Bongardine, B-248
 Buchananine, B-376
 5-(3-Butenyl)-2-pyridinecarboxylic acid, B-425
 5-Butyl-2-pyridinecarboxylic acid, B-445

2-Butyl-3,4,5,6-tetrahydropyridine, B-446
 Caerulomycin D, C-10
 Caerulomycin E, C-11
 Caerulomycin, C-9
 Carbonarone B, C-142
 3-(2-Carboxyethyl)-1,4-dimethylpyridinium, C-146
 2-Carboxy-1-methylpyridinium betaine, C-148
 Cerpegin, C-285
 Citridone A, C-478
 Citridone B, C-479
 Citridone D, C-481
 Collismycin C, C-575
 Collismycin D, C-576
 Collismycin E, C-577
 Collismycin F, C-578
 Colubricidin A, C-579
 Cordypyridone A, C-646
 Cordypyridone D, C-647
 Cortamidin oxide, C-664
 Cribrochalinamine oxide A, C-738
 Cribrochalinamine oxide B, C-739
 5-Decanoyl-2-nonylpyridine, D-116
 Decaturin A, D-127
 Decaturin D, D-128
 7-Demethylpericidin A₁, D-203
 Desferroferrithiocin, D-235
 Desmotrichinine, D-238
 3,5-Didecanoylpyridine, D-371
N-[5-[(1,6-Dihydro-2-hydroxy-5-octanamido-6-oxo-3-pyridyl)imino]-1,2,5,6-tetrahydro-2,6-dioxo-3-pyridyl]octanamide, D-450
 5,6-Dihydro-3-[2-(4-hydroxyphenyl)-2-oxoethyl]-2(1*H*)-pyridinone, D-456
 5,6-Dihydro-3-[2-(1*H*-indol-3-yl)-2-oxoethyl]-2(1*H*)-pyridinone, D-468
 5,6-Dihydro-*N*-[3-(7-methoxy-1,3-benzodioxol-5-yl)-2-propenoyl]-2(1*H*)-pyridinone, D-470
 3,4-Dihydro-5-(2-pyridinyl)-2*H*-pyrrole-2-carboxylic acid, D-496
 2-(3,4-Dihydro-2*H*-pyrrol-5-yl)pyridine, D-504
 5,6-Dihydro-3,6,6-trimethyl-7*H*-2-pyridin-7-one, D-525
 3-(2,6-Dihydroxy-4-methoxybenzoyl)pyridine, D-609
 3,4-Dihydroxy-2-methylpyridine, D-632
N-(2,5-Dihydroxyphenyl)pyridinium(1+), D-655
 1,1'-Dimethyl-3,3'-bipyridinium(2+), D-743
 2,6-Dimethylpyridine, D-775
Cortinarius Disulfide, D-870
 3-Dodecylpyridine, D-899
 Echinoclastrine A, E-16
 Echinoclastrine C, E-17
 Epipyridone, E-101
 5-Ethyl-2-methylpyridine, E-256
 3-Ethylpyridine, E-268
 Farinosone A, F-10
 Fischerin, F-62
 Fontaphilline, F-125
 Fruit rot toxin A, F-162
 Funiculosin, F-197
 Fusaricide, F-225
 Gostatin, G-146
 Griffithine, G-176
 Guaipyridine, G-190
 3-Guanidinylpyridine, G-209
 Guayulamine A, G-216
 Guayulamine B, G-217
 Halfordinol, H-12
 Haliclamine C, H-17
 Haliclamine D, H-18
 Halitoxin, H-32
 Hermidin, H-146
 3-Hexylpyridine, H-267
 5-(4-Hydroxybenzyl)-2-pyridinecarboxylic acid, H-426
 5-Hydroxy-2-(hydroxymethyl)pyridine, H-515
 6-Hydroxy-5-methoxy-1-methyl-2,3,4(1*H*)-pyridinetrione, H-562
 1-Hydroxy-5-methoxy-6-methyl-2(1*H*)-pyridinone, H-563

4-(Hydroxymethyl)-6-methyl-2,3,5-pyridinetrione, H-608
 3-Hydroxy-6-methyl-2-(11-oxododecyl)pyridine, H-612
 4-Hydroxy-2-methylpyridine, H-625
 5-Hydroxy-2-pyridinecarboxylic acid, H-710
 Ikimine A, I-24
 Ikimine B, I-25
 Ikimine D, I-26
 Ilicicolin H, I-28
 Indigoidin, I-70
 Iromycin A, I-179
 Iromycin C, I-180
 Isoflavipucine, I-234
 1-(8-Isoquinolinylmethyl)pyridinium(1+), I-308
 Jussiaeine A, J-61
 Kedarcidin, K-26
 Kuraramine, K-108
 Latidine, L-58
 Leporine A, L-112
 Lycoplamine A, L-338
 Macaridine, M-2
 Mamanine, M-79
 Melochimine, M-184
 4-Methoxy-7,8-dimethyl-2*H*-pyrano[4,3-*b*]pyridin-2-one, M-250
 6-Methoxy-3-(1-propenyl)-2-pyridinecarboxylic acid, M-283
 2-Methylpyridine, M-537
 1-(5-Methyl-3-pyridinyl)-1-decanone, M-538
 Militarione A, M-610
 Militarione D, M-612
 Mimopudine, M-623
 Montipyridine, M-699
 Nakinadine A, N-17
 Nakinadine D, N-20
 Nakinadine E, N-21
 Nakinadine F, N-22
 Neobudoficidin B, N-102
 Nigrifactin, N-213
 Nikkomycin Q_z, N-217
 Nikkomycin X, N-216
 Nikkomycin Z, N-218
 Niphatesine B, N-224
 Niphatesine D, N-226
 Niphatesine E, N-227
 Niphatesine F, N-228
 Niphatesine G, N-229
 Niphatoxin A, N-230
 Niphatoxin B, N-231
 Niphatoxin C, N-232
 Niphatyne A, N-233
 Niphatyne B, N-234
 3-(12-Nitrododecyl)pyridine, N-249
 5-Octanoylamino-2,3,6(1*H*)-pyridinetrione, O-62
 Ocular age pigment A2-E, O-65
 Oxalicine A, O-148
 2-Oxo-3-(4-pyridinyl)propanoic acid, O-205
 Oxysporidinone, O-226
 Pachychaline B, P-2
 Pachychaline C, P-3
 Patchoulipyridine, P-112
 Patungensine, P-126
 Pedoxin, P-159
 Penicidone A, P-186
 Penicidone C, P-187
 4-Pentadecylpyridine, P-204
 Pentasine, P-231
 2-Pentylpyridine, P-237
 Petrosaspongiolide L, P-289
 Petunianine A, P-291
 Petunianine B, P-292
 Petunianine C, P-293
 4-Phenylpyridine, P-352
 Piperarboresine C, P-442
 Piperarboresine, P-443
 Piplartine dimer A, P-474
 Proferrosamine B, P-637
 Prolifine, P-638
 Promothiocin A, P-641
 Promothiocin B, P-642
 2-Propanoylpyridine, P-645

Puloupone, P-789
 Puralidin D, P-810
 Purpuramine C, P-822
 2-Pyridinecarboxylic acid, P-896
 2,6-Pyridinedicarbothioic acid, P-899
 2,6-Pyridinedicarboxylic acid, P-901
 Pyridine, P-894
 Pyridinoline, P-907
 Pyridomacrolidin, P-918
 Pyridomycin, P-919
 Pyridovericin, P-921
 Pyridoxatin, P-922
 Pyridoxine, P-923
 Pyrinadine A, P-925
 Pyrinadine B, P-926
 Pyrinadine C, P-927
 Pyrinadine D, P-928
 Pyrinadine E, P-929
 Pyrinadine F, P-930
 Pyrinadine G, P-931
 Pyrinodemin A, P-933
 Pyrinodemin B, P-934
 3,2':3',4':2'',3'''-Quaterpyridine, Q-5
 Rotundine B, R-135
 Rubrifacine, R-147
 Sambutoxin, S-48
 Siderochelin A, S-295
 Siderochelin C, S-296
 Simplakidine A, S-303
 Speranculatine A, S-391
 Speranskilatine A, S-393
 Streptonigrin, S-569
 Streptonigrone, S-570
N-(2-Sulfoethyl)pyridinium betaine, S-636
 3,2':4',3''-Terpyridine, T-98
 2,3,4,5-Tetrahydro-6-(4-pentenyl)pyridine, T-214
 2,3,4,5-Tetrahydro-6-pentylpyridine, T-216
 3,3',4,4'-Tetrahydroxy-2,2'-bipyridine, T-249
 2,4,6,6-Tetramethyl-3(6*H*)-pyridinone, T-285
 Texaline, T-299
 Thallusin, T-338
 Theonelladine A, T-357
 Theonelladine C, T-358
 Tokaradine A, T-422
 2,3,5-Trihydroxy-6-methylpyridine, T-567
 Viscosaline, V-161
 Viscosamine, V-162
 Xestamine A, X-15
 Xestamine B, X-16
 Xestamine C, X-17
 Xestamine D, X-18
 Xestamine E, X-19

9B-Azaphenalene alkaloids

Dendrobates Alkaloid 205B, A-402
 Chilocorine A, C-377
 Chilocorine B, C-378
 Chilocorine C, C-379
 Chilocorine D, C-380
 Exochomine, E-321
 Poranthericine, P-569
 Propyleine, P-649
 Psylloborine A, P-749

Naphthalene-isoquinoline alkaloids

Ancisheynine, A-978
 Ancistrobrevine A, A-979
 Ancistrobrevine B, A-980
 Ancistrobrevine C, A-981
 Ancistrocladine, A-982
 Ancistrocladinium A, A-983
 Ancistrocladinium B, A-984
 Ancistrocladisine, A-985
 Ancistrocongine, A-986
 Ancistrocongolensine, A-987
 Ancistrogriffithine A, A-988
 Ancistroheynine A, A-989
 Ancistorobertsonine D, A-990
 Ancistrotanzanine A, A-991
 Ancistrotectorine, A-992

3,4-Dihydro-5-(4-hydroxy-5-methoxy-2-methyl-1-naphthalenyl)-8-hydroxy-6-methoxy-1,3-dimethylisoquinoline, D-444
 Dioncolactone A, D-790
 Dioncophylleine D, D-792
 Dioncophylline A, D-793
 Dioncophylline B, D-794
 Dioncophylline C, D-795
 Dioncophyllinol B, D-796
 Korundamine A, K-90
 Korupensamine A, K-91
 Yaoundamine A, Y-6

Elaeocarpus alkaloids

Elaeocarpidine, E-54
 Elaekonanidine A, E-57
 Grandisine C, G-162
 Grandisine D, G-163
 Grandisine E, G-164
 Grandisine F, G-165
 Grandisine G, G-166
 Pseudoepiiselaecarpiline, P-714

Galbulimima alkaloids

Alkaloid GB14, A-521
 Alkaloid GB15, A-522
 Alkaloid GB16, A-523
 Alkaloid GB17, A-524
 Alkaloid GB18, A-525
 Himandridine, H-277
 Himbacine, H-280
 Himbadine, H-281
 Himgaline, H-285
 Himgrine, H-286

Stemona alkaloids

Bisdehydroneostemoninine, B-169
 Bisdehydrostemoninine A, B-170
 Cochinchistemonine, C-544
 Hodorine, H-312
 Hypotuberostemonine, H-774
 Isostemonidine, I-323
 Isostemonine, I-324
 Isotuberostemonine, I-341
 Maistemonine, M-49
 Methylstemofoline, M-560
 Neostemofoline, N-139
 Neostemonine, N-140
 Oxotuberostemonine, O-215
 Oxystemokerrillactone, O-227
 Paipunine, P-14
 Parvineostemonine, P-101
 Parvistemoamide, P-103
 Parvistemoline, P-104
 Parvistemonine, P-105
 Protostemodiol, P-669
 Protostemonine, P-670
 Saxorumamide, S-112
 Sessilifoliamide B, S-265
 Sessilifoliamide D, S-266
 Sessilifoliamide E, S-267
 Sessilifoliamide G, S-268
 Sessilistemonamine A, S-270
 Sessilistemonamine D, S-271
 Sessilistemonine, S-272
 Sinostemonine, S-317
 Stemoamide, S-511
 Stemoburkilline, S-512
 Stemocochinamine, S-513
 Stemocurtisine, S-514
 Stemofoline, S-515
 Stemokerrin, S-516
 Stemonamide, S-517
 Stemonine, S-519
 Stemonine, S-520
 Stemoninine, S-521
 Stemoninoamide, S-522
 Stemospironine, S-523
 Stemotinine, S-524
 Stemotuberine, S-525

Stenine, S-528
 Stichoneurine A, S-552
 Tuberostemoenone, T-663
 Tuberostemonone, T-664
 Tuberostemoninol, T-665
 Tuberostemonone, T-666
 Tuberostemospironine, T-667

Lycopodium alkaloids

Lycopodium Alkaloid 168, A-312
Lycopodium Alkaloid 258, A-313
Lycopodium Alkaloid E, A-501
Lycopodium Alkaloid G, A-516
Lycopodium Alkaloid IV, A-344
Lycopodium annotinum Alkaloid IX, A-531
 Alkaloid L10, A-547
 Alkaloid LE1, A-556
 Alkaloid LO1, A-557
 Alkaloid LO2, A-558
 Alkaloid LO3, A-559
 Alkaloid LS14, A-560
 Alkaloid LV1, A-562
Lycopodium fawcettii Alkaloid R, A-592
Lycopodium Alkaloid V, A-310
Lycopodium annotinum Alkaloid VI, A-605
Lycopodium annotinum Alkaloid VII, A-606
Lycopodium annotinum Alkaloid VIII, A-609
Lycopodium Alkaloid X, A-311
Lycopodium annotinum Alkaloid XI, A-610
Lycopodium annotinum Alkaloid XII, A-611
Lycopodium annotinum Alkaloid, A-314
Lycopodium annotinum Alkaloid, A-315
Lycopodium clavatum Alkaloid, A-316
Lycopodium thyoides Alkaloid, A-317
 Anhydrolycodoline, A-1029
 Annapodine, A-1059
 Annotine, A-1062
 Annotinine, A-1063
 Borbonicine, B-250
 Cermizine A, C-282
 Cermizine B, C-283
 Cermizine D, C-284
 Compladine A, C-590
 Compladine C, C-591
 Cryptadine A, C-782
 Decahydro-2,5,8-trimethyl-2*H*-1-oxa-4,7,9*b*-triazaphenalene, D-114
 Erythreidine, E-209
 Erythreine, E-210
 Fastigiatine, F-17
 Fawcettidine, F-28
 Fawcettiine, F-29
 Fawcettimine, F-30
 Flabelliformine, F-73
 Flabelline, F-75
 Gnidine, G-134
 Gnidinine, G-135
 Gymnamine, G-228
 Himeradine A, H-284
 Huperzine A, H-375
 Huperzine B, H-376
 Huperzine F', H-377
 Huperzine F, H-378
 Huperzine H, H-379
 Huperzine I, H-373
 Huperzine J, H-380
 Huperzine P, H-381
 Huperzine Q, H-382
 Huperzine R, H-383
 Huperzine S, H-384
 Huperzine T, H-385
 Huperzine U, H-386
 Huperzine W, H-387
 Huperzinine C, H-389
 Huperzinine, H-388
 Hydroxypropyllycodine, H-703
 Isoobscurinine, I-258
 Lannotinidine A, L-34
 Lannotinidine E, L-35
 Lannotinidine G, L-36
 Lobscurinol, L-223
 Lucidine B, L-277

Lucidioline, L-279
 Luciduline, L-280
 Lycocaroline, L-320
 Lycocernuine, L-321
 Lycoclavine, L-322
 Lycodiflexine, L-323
 Lycodine, L-324
 Lycodoline, L-325
 Lycofawcine, L-326
 Lycoflexine, L-327
 Lycofoline, L-328
 Lyconadine A, L-330
 Lyconesidine A, L-331
 Lyconnotine, L-332
 Lyconnotinol, L-333
 Lycopaniculatine, L-334
 Lycopocurine, L-335
 Lycoperine A, L-336
 Lycoplegmarine, L-337
 Lycopladine A, L-338
 Lycopladine B, L-339
 Lycopladine D, L-340
 Lycoposerramine B, L-343
 Lycoposerramine G, L-344
 Lycoposerramine S, L-345
 Lycoposerramine W, L-346
 Lycoserramine, L-349
 Lycoserrine, L-350
 Lycovatine A, L-352
 Macleanine, M-6
 Magellanine, M-40
 Megastachine, M-147
 Nankakurine A, N-27
 Neohuperzidine, N-118
 β-Obscurine, O-5
 Obscurinine, O-9
 5,15-Oxidolycopodane, O-163
 Paniculine, P-71
 Phlegmarine, P-361
 Phlegmariurine A, P-362
 Phlegmariurine B, P-363
 Phlegmariurine M, P-365
 Pillijanine, P-423
 Saururidine, S-103
 Saururine, S-104
 Senepodine A, S-242
 Senepodine G, S-243
 Senepodine H, S-244
 Serratezomine A, S-254
 Serratine, S-255
 Serratidine, S-256
 Serratine, S-257
 Sieboldine A, S-297
 SpiroLucidine, S-435

Cytochalasin alkaloids

Antibiotic L 697318, A-1196
 Ascochalasin, A-1460
 Aspergillin PZ, A-1476
 Aspochalamin A, A-1500
 Aspochalasin C, A-1502
 Aspochalasin F, A-1503
 Aspochalasin I, A-1501
 Chaetochalasin A, C-319
 Chaetoglobosin A, C-325
 Chaetoglobosin B, C-326
 Chaetoglobosin D, C-327
 Chaetoglobosin J, C-328
 Chaetoglobosin K, C-329
 Chaetoglobosin P, C-330
 Chaetoglobosin Q, C-331
 Chaetoglobosin U, C-332
 Cytochalasin B, C-943
 Cytochalasin C, C-944
 Cytochalasin E, C-945
 Cytochalasin F, C-946
 Cytochalasin G, C-947
 Cytochalasin K, C-948
 Cytochalasin L, C-949
 Cytochalasin M, C-950
 Cytochalasin N, C-951
 Cytochalasin O, C-952

Cytochalasin O, C-953
 Cytochalasin Q, C-954
 Cytochalasin S, C-955
 Cytochalasin U, C-956
 Cytochalasin V, C-942
 Cytochalasin W, C-957
 Cytochalasin Y, C-958
 Cytochalasin Z₇, C-959
 7,18-Dihydroxy-16,18-dimethyl-10-phenyl-22-oxa[12]cytochalasa-6(12),13,19-triene-1,21-dione, D-573
 Engleromycin, E-81
 Epoxycytochalasin H, E-129
 7-Hydroperoxy-17-hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-5,13,19-triene-1,21-dione, H-399
 18-Hydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6,13,19-triene-1,21-dione, H-473
 Meleagrins, M-162
 Neoxaline, N-149
 Penochalasin C, P-197
 Phomacin A, P-371
 Phomacin C, P-372
 Phomopsichalasin, P-376
 Protophomin, P-666
 Proxiphomin, P-677
 Rosellichalasin, R-128
 7,18,19-Trihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13-diene-1,21-dione, T-545
 Zygosporin D, Z-37
 Zygosporin G, Z-38

Indolizidine alkaloids

Dendrobates Alkaloid 237A, A-359
Mantella Alkaloid 253B, A-419
Mantella Alkaloid 295B, A-422
Mantella Alkaloid 235C, A-451
Dendrobates Alkaloid 251D, A-480
Mantella Alkaloid 243D, A-486
Mantella Alkaloid 279D, A-487
Mantella Alkaloid 241F, A-511
Mantella Alkaloid 281F, A-514
Mantella Alkaloid 221F, A-510
 Allopumiliotoxin 309D, A-629
 Anibamine, A-1036
 5-Butyl-6,8-diethyloctahydroindolizine, B-430
 8-Butyloctahydro-5-propyl-1*H*-indolizine, B-437
 5-Butyl-8-propylindolizidine, B-443
 Buzonamine, B-484
 Castanospermine, C-188
 Crepidamine, C-735
 Cyclizidine, C-837
 5-(1,3-Decadienyl)octahydroindolizine, D-107
 Dendrorepine, D-212
 Deoxypumiliotoxin 251H, D-228
 6,7-Dihydro-3-hydroxymethyl-8(5*H*)-indolizine, D-447
 6,7-Dihydro-8(5*H*)-indolizine, D-463
 8-Ethyl-5-(1-ethyl-1-butenyl)octahydro-6-methylindolizine, E-240
 6-Ethyl-5-(1-ethyl-1-butenyl)octahydro-6-methylindolizine, E-263
 Ficuseptine, F-54
 5-(3-Furanyl)octahydro-8-methylindolizine, F-202
 Grandisine A, G-160
 Grandisine B, G-161
 Grandisine G, G-166
 3-(5-Hexenyl)octahydro-5-(4-pentenyl)indolizine, H-260
 Indolizomycin, I-101
 Ipohardine, I-170
 Juliflorine, J-56
 Juliprosine, J-57
 Louludinium(1+), L-273
 Myrmicarin 233A, M-799
 Octahydro-8-methyl-5-(6-nonen-8-ynyl)-1*H*-indolizine, O-51
 Octahydro-8-methyl-5-(4-pentenyl)-1*H*-indolizine, O-52
 Phyllosteminine, P-390

Polycanthidine, P-549
 Polycanthisine, P-550
 Pumiliotoxin A, P-791
 Pumiliotoxin 341A, P-792
 Pumiliotoxin 307B, P-793
 Pumiliotoxin 209F, P-795
 Stelletamide A, S-504
 Stelletamide B, S-505
 Swainsonine, S-650
 Tyloindicine F, T-692
 Tyloindicine I, T-691

Simple anthranilic acid alkaloids

3-Amino-5-hydroxybenzoic acid, A-790
 Avenanthramide L, A-1564
 Delamide, D-178
 Doisuthine, D-900
 2,2'-Iminobis[4-hydroxybenzoic acid], I-45
 Tecleanone, T-55
 Thiobutacin, T-381
 Viridic acid, V-154

Maytansinoid alkaloids

Maysenine, M-134
 Maytansine, M-135
 Maytansinol, M-136
 Trenudine, T-449
 Trewasine, T-450

Simple quinoline alkaloids

Dendrobates Alkaloid 203, A-289
Dendrobates Alkaloid 167A, A-354
Dendrobates Alkaloid 181A, A-355
Dendrobates Alkaloid 219A, A-357
Dendrobates Alkaloid 231A, A-358
Dendrobates Alkaloid 243A, A-362
Dendrobates Alkaloid 269A, A-366
Dendrobates Alkaloid 181B, A-401
Dendrobates Alkaloid 219B, A-403
Dendrobates Alkaloid 223B, A-404
Dendrobates Alkaloid 231B, A-406
Dendrobates Alkaloid 269B, A-412
Dendrobates Alkaloid 223C, A-438
Dendrobates Alkaloid 223D, A-479
Dendrobates Alkaloid 253D, A-481
 2-Alkyl-3,4-dihydroxyquinolines, A-621
 Atanine, A-1529
 Aurachin B, A-1542
 Aurachin D, A-1543
 6-Bromo-2,4-dihydroxyquinoline, B-313
 7-Bromo-4-(2-ethoxyethyl)quinoline, B-317
 Buchapine, B-377
 Bucharaine, B-378
 Bucharidine, B-380
 2-Butyldecahydroquinoline, B-428
 Casimiroin, C-177
 Cusparine, C-822
 Dasycarine, D-83
 2-Decyl-4-hydroxyquinoline, D-139
 6,7-Dibromo-2,4-dihydroxyquinoline, D-315
 3,4-Dihydro-4,5-dihydroxy-3-methoxy-4-(4-methoxyphenyl)-2(1*H*)-quinolinone, D-422
 3,4-Dihydro-6-hydroxy-2(1*H*)-quinolinone, D-461
 2,3-Dihydroxy-4-phenylquinoline, D-656
 2,8-Dihydroxy-4-quinolinecarboxylic acid, D-673
 3,4-Dihydroxy-2-quinolinecarboxylic acid, D-674
 4,6-Dihydroxy-2-quinolinecarboxylic acid, D-675
 4,8-Dihydroxy-2-quinolinecarboxylic acid, D-677
 5,8-Dimethoxy-2-(3-methoxyphenyl)-3-propyl-4(1*H*)-quinolinone, D-711
 5,6-Dimethoxy-2-(3-methoxyphenyl)-4(1*H*)-quinolinone, D-712
 5,7-Dimethoxy-4-(4-methoxyphenyl)-2(1*H*)-quinolinone, D-713

2,4-Dimethoxy-7,8-methylenedioxy-3-prenylquinoline, D-715
 3,6-Dimethoxy-1-methyl-2-phenyl-4(1*H*)-quinolinone, D-716
 2-Dodecyl-4-hydroxyquinoline, D-898
 Echinine, E-13
 2-(1-Ethylpropyl)-4-methoxyquinoline, E-266
 2-(1-Ethylpropyl)-1-methyl-4(1*H*)-quinolinone, E-267
 Evomeliaefolin, E-314
 Folimidine, F-119
 Foliosidine, F-122
 Galipinine, G-9
 Galipoline, G-11
 Glansreginin A, G-86
 Glycocitlone A, G-118
 Glycocitridine, G-119
 Glycolone, G-124
 Graveoline, G-171
 Haplafine, H-53
 Haplosamine, H-59
 Haplosine, H-60
 Haplotubinine, H-61
 3-Heptyl-3-hydroxy-2,4(1*H*,3*H*)-quinolinedione, H-134
 2-Heptyl-4-hydroxyquinoline, H-133
 2-Hexyl-4-hydroxyquinoline, H-263
 2-Hexyl-8-methoxy-1-methyl-4(1*H*)-quinolinone, H-265
 4-Hydroxy-2,3-dimethylquinoline, H-477
 6-Hydroxy-4,8-dimethylquinoline, H-478
 6-Hydroxy-5,7-dimethylquinoline, H-479
 6-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1*H*)-quinolinone, H-511
 7-Hydroxy-2-(3-hydroxy-3-methylbutyl)-4(1*H*)-quinolinone, H-512
 7-Hydroxy-4-(5-hydroxymethyl-2-furanyl)-2(1*H*)-quinolinone, H-513
 2-(2-Hydroxy-4-methoxyphenyl)-5,8-dimethoxy-3-propyl-4(1*H*)-quinolinone, H-567
 2-(11-Hydroxy-11-methyldodecyl)-3-methoxy-4(1*H*)-quinolinone, H-598
 2-(12-Hydroxy-12-methyltridecyl)-4(1*H*)-quinolinone, H-636
 3-Hydroxy-3-nonyl-2,4(1*H*,3*H*)-quinolinedione, H-643
 4-Hydroxy-2-nonylquinoline, H-642
 4-Hydroxy-2-octylquinoline, H-648
 4-Hydroxy-2-pentylquinoline, H-660
 2-(4-Hydroxyphenyl)-1-methyl-4(1*H*)-quinolinone, H-685
 3-(4-Hydroxyphenyl)quinoline, H-692
 4-Hydroxy-2-propylquinoline, H-707
 8-Hydroxy-4-quinolinecarboxaldehyde, H-722
 2-Hydroxy-4-quinolinecarboxylic acid, H-723
 3-Hydroxy-2-quinolinecarboxylic acid, H-724
 4-Hydroxy-2-quinolinecarboxylic acid, H-725
 4-Hydroxy-2-tridecylquinoline, H-750
 4-Hydroxy-2-undecylquinoline, H-757
 3-(1*H*-Indol-3-yl)quinoline, I-138
 Lunamarine, L-290
 Lunidonine, L-295
 Malatyamine, M-68
 Megistoquinone II, M-151
 Melisemine, M-175
 Melovinone, M-195
 4-Methoxy-2-(3,4-methylenedioxyphenyl)quinoline, M-264
 3-Methoxy-1-methyl-2-propyl-4(1*H*)-quinolinone, M-274
 4-Methoxy-1-methylquinolinium(1+), M-275
 1-Methyl-3,3-bis(3-methyl-2-butenyl)-2,4(1*H*,3*H*)-quinolinedione, M-406
 2-[4-(3,4-Methylenedioxyphenyl)butyl]-4(1*H*)-quinolinone, M-449
 2-[6-(3,4-Methylenedioxyphenyl)hexyl]-4(1*H*)-quinolinone, M-453
 2-(3,4-Methylenedioxyphenyl)quinoline, M-458
 3-Methyl-2-(2-nonenyl)-4(1*H*)-quinolinone, M-491
 1-Methyl-2-(9-oxodecyl)-4(1*H*)-quinolinone, M-498

3-Methyl-2-pentyl-4(1*H*)-quinolinone, M-508
 2-Methylquinoline, M-557
 1-Methyl-4(1*H*)-quinolinone, M-558
 2-Pentylquinoline, P-238
 Perspicamide A, P-274
 2-Phenylquinoline, P-354
 3-Phenylquinoline, P-355
 Pilokeanine, P-426
 Pinolinone, P-440
 Preskimmianine, P-625
 2-(1-Propenyl)quinoline, P-647
 2-Propylquinoline, P-654
 Ptelecortine, P-750
 Ptelefoline, P-755
 Pteleoline, P-758
 Pteleprenine, P-759
 Pumiliotoxin C, P-794
 2-Quinolamine, Q-28
 4-Quinolinecarboxaldehyde, Q-30
 4-Quinolinecarboxylic acid, Q-32
 3,8-Quinolinediol, Q-34
 4,8-Quinolinediol, Q-35
 2-Quinolinemethanethiol, Q-36
 2-Quinolinemethanol, Q-37
 4-Quinolinemethanol, Q-38
 Quinoline, Q-29
 3,4,5,8-Quinolinetetrol, Q-44
 3-Quinolol, Q-49
 Sarcomejine B, S-78
 Sarcomejine, S-77
 Semecarpifoline, S-221
 Severibuxine, S-276
 Sterculinine I, S-550
 5,6,7,8-Tetrahydro-2,4-dimethylquinoline, T-170
 5,6,7,8-Tetrahydro-2,5-dimethylquinoline, T-171
 5,6,7,8-Tetrahydro-5-methyl-2-propylquinoline, T-204
 5,6,7,8-Tetrahydro-4-methylquinoline, T-208
 5,6,7,8-Tetrahydro-2-nonylquinoline, T-210
 Trididemnic acid A, T-512
 4,5,8-Trihydroxy-2-quinolinecarboxylic acid, T-586
 Zanthobisquinolone, Z-2
 Zascanol epoxide, Z-9

Furanoquinoline alkaloids

2-Acetyl-4,6-dimethoxyfuro[2,3-*b*]quinoline, A-50
 Acronydrine, A-121
 Anhydroperforine, A-1032
 Aurachin A, A-1541
 Aurachin G, A-1546
 Bucharamine, B-379
 8-(1,1-Dimethyl-2-propenyl)-4-methoxyfuro[2,3-*b*]quinolin-7-ol, D-771
 Dubinidine, D-945
 Dutadrupine, D-958
 Evomerrine, E-315
 Folifinine, F-118
 Foliminine, F-120
 Haplophyllidine, H-57
 7-Hydroxy-4-methoxy-8-prenylfuro[2,3-*b*]quinoline, H-570
 Megistoquinone I, M-150
 Megistosarconine, M-152
 Melicarpine, M-165
 Melicarpinone, M-166
 Myrtopsine, M-804
 Oligophylline, O-82
 Ptelefolone, P-756
 Ptelefolonium, P-757
 Radixisatidis A, R-11
 Ribalinium, R-93
 Riedelianine, R-97
 Sarcodifurine A, S-74
 Sarcodifurine B, S-75
 Semecarpine, S-222
 Tecleaverdoornine, T-57

Pyranoquinoline alkaloids

Acronydrine, A-121
 Anhydroperforine, A-1032
 Aurachin F, A-1545
 Azaacridone A, A-1579
 Benzosimuline, B-76
 Dutadrupine, D-958
 Flindersine, F-90
 Foliminine, F-120
 Haplofoline, H-56
 Helietidine, H-91
 5-Methoxy-2,2-dimethyl-2*H*-pyrano[2,3-*b*]quinoline, M-251
 Neohydroxylunine, N-119
 Pteleflorine, P-754
 Rutalinium, R-171
 Simulanoquinoline, S-305
 Yaequinolone J₁, Y-3
 Zanthosimuline, Z-5

Miscellaneous quinoline alkaloids

Acutopyrrocoline, A-135
 Antibiotic BE 22179, A-1128
 Antibiotic FR 225656, A-1172
 Antibiotic FR 225659, A-1173
 Antibiotic G 1499-2, A-1179
 Antibiotic Sch 538415, A-1251
 Antibiotic SW 163C, A-1271
 Antidesmone, A-1325
 Araliopdimerine A, A-1376
 Asperumoid, A-1469
 Aspernigerin, A-1482
 Aspoquinolone A, A-1504
 Aspoquinolone C, A-1505
 Broussonetine, B-358
 Calebassinine 1, C-32
 Cermizine A, C-282
 Cermizine B, C-283
 Chamaedrone, C-343
 Chestnutamide, C-370
 Corialstonidine, C-651
 Corialstonine, C-652
 Cribrostatin 6, C-740
 Cryptadine A, C-782
 Cyclomestine, C-878
 Diazaquinomycin B, D-295
 Diazaquinomycin C, D-296
 Dictyolomide A, D-367
 Donglingine, D-918
 Echinoramamine I, E-24
 Erioaustralasine hydrate, E-180
 Fareanine, F-8
 Folipidine, F-123
 Furoeriaustralasine, F-205
 Geijedimerine, G-30
 Haplodimerine, H-55
 Huperzine J, H-380
 Huperzine V, H-374
 10-Hydroxy-1,4,6,9-tetramethylpyrido[3,2-*g*]quinoline-2,8(1*H*,9*H*)-dione, H-747
 Iboquine, I-7
 2-(1*H*-Imidazol-4-yl)quinoline, I-44
 Isaindigotidione, I-181
 Lagunamycin, L-12
 Lavendamycin, L-78
 Lepadin B, L-103
 Lepadin D, L-104
 Luotonine F, L-301
 Lycoperine A, L-336
 Lycoposerramine W, L-346
 Macrorungine, M-28
 Melicobisquinolinone A, M-167
 Melicobisquinolinone B, M-168
 Melochinone, M-185
 Methoxatin, M-236
 4-Methyl-3*H*-pyrrolo[2,3-*c*]quinoline, M-552
 Myrionine, M-797
 Njaoamine A, N-259
 Njaoamine E, N-260
 Njaoamine F, N-261
 Nybomycin, N-346

Paraensidimerine A, P-87
 Paraensidimerine B, P-88
 Penigequinolone A, P-191
 Peniprequinolone, P-193
 Pteledimericine, P-751
 Pteledimeridine, P-752
 Pteledimerine, P-753
 Quinaldopeptin, Q-11
 Quinocitrinone A, Q-23
 Quinolactacide, Q-24
 Quinolactacin A₁, Q-25
 Quinolactacin D₁, Q-27
 1-(2-Quinoliny)-β-carboline, Q-51
 1-(4-Quinoliny)-β-carboline, Q-52
 1-(5-Quinoliny)-β-carboline, Q-53
 1-(6-Quinoliny)-β-carboline, Q-54
 1-(8-Quinoliny)-β-carboline, Q-55
Oceanapia Quinolone alkaloid, Q-57
 Scolopendrine, S-156
 Secomartinelline, S-191
 Senepodine A, S-242
 Streptonigrin, S-569
 Streptonigrone, S-570
 Syphilobin A, S-661
 Thiocoraline, T-384
 Toddacoumalone, T-420
 2,2,6-Trimethyl-4-(4-quinoliny)-3-azabicyclo[3.3.1]non-6-ene, T-614
 Vepridimerine C, V-55
 Virantmycin, V-145
 Voastricline, V-182
 Waltherione A, W-8
 Yaequinolone C, Y-2
 Yaequinolone J₁, Y-3

Quinazoline alkaloids

2-Acetyl-4(3*H*)-quinazolinone, A-75
 Aniflorine, A-1039
 Anisessine, A-1041
 Anisotine, A-1045
 2-Benzyl-4(3*H*)-quinazolinone, B-93
 Bouchardatine, B-260
 7-Bromo-2,4-dihydroxyquinazoline, B-312
 Candidine, C-77
 Chaetominine, C-336
 Dictyoquinazole A, D-369
 2,3-Dihydropyrrolo[2,1-*b*]quinazolin-9(1*H*)-one, D-503
 2,4-Dihydroxy-5-methylquinazoline, D-633
 2,4-Dimethylquinazoline, D-781
 2,2-Dimethyl-4(1*H*,3*H*)-quinazolinone, D-782
 Dipepine, D-812
 2-Ethyl-4-methylquinazoline, E-259
 Goshuyamide II, G-144
 2-(4-Hydroxybutyl)-4(3*H*)-quinazolinone, H-437
 2-Hydroxymethyl-4-methylquinazoline, H-609
 4-Hydroxy-2-methylquinazoline, H-632
 2-(3-Hydroxypropyl)-4(3*H*)-quinazolinone, H-706
 Indolo[2,1-*b*]quinazoline-6,12-dione, I-108
 Isaindigotone, I-182
 Isofebrifugine, I-232
 Luotonine F, L-301
 Mackinazolinone, M-5
 18-*O*-[2-(2-Methyl-4-oxo-4*H*-quinazolin-3-yl)-benzoyl]lycoctonine, M-501
 1-Methyl-3-(2-phenylethyl)-2,4(1*H*,3*H*)-quinazolinone, M-510
 4-Methyl-2-quinazolinecarboxylic acid, M-555
 4-Methylquinazoline, M-554
 1-Methyl-4(1*H*)-quinazolinone, M-556
 Monodontamide F, M-691
 Nordine, N-296
 Ophiuroidine, O-98
 Peganidine, P-164
 2,4(1*H*,3*H*)-Quinazolinone, Q-13
 4-Quinazolinol, Q-14
 Sessiflorine, S-264
 6,8,9,11-Tetrahydro-7*H*-pyrido[2,1-*b*]quinazolinone, T-227

1,2,3,9-Tetrahydropyrrolo[2,1-*b*]quinazolinone, T-231
 2-(2,4,6-Trimethyl-1-heptenyl)-4(1*H*)-quinazolinone, T-606
 Vasicinine, V-36
 Vasicoline, V-40
 Wuchuyamide III, W-25
 Wuchuyamide I, W-24
 Wuzhuyurutine B, W-27

Acridone alkaloids

Acridone, A-112
 Acriginine A, A-113
 Acronycine, A-120
 Atalaphyllidine, A-1527
 Buxifoliadine D, B-467
 Buxifoliadine E, B-468
 Buxifoliadine F, B-469
 Buxifoliadine G, B-470
 Chaloridone, C-342
 Citracridone III, C-472
 Citropone A, C-484
 Citropone B, C-485
 Citropone C, C-486
 1,7-Dihydroxyacridone, D-529
 1,8-Dihydroxyacridone, D-530
 3-(1,1-Dimethoxy-2-oxoethoxy)-9,10-dihydro-1-methoxy-9-oxo-4-acridinecarboxaldehyde, D-717
 Furofoline II, F-207
 Furofoline, F-206
 Furoparadine, F-209
 Glycocitrine III, G-121
 Glycocitrine V, G-120
 Glycocitrine VI, G-122
 Hallacridone, H-34
 1-Hydroxyacridone, H-402
 4-(2-Hydroxy-3-methyl-3-butenyl)jucocitrine, H-581
 Junosidine, J-59
 Margrapine A, M-103
 Megistophylline II, M-149
 Megistophylline I, M-148
 Melicopine, M-169
 10-Methylacridone, M-382
N-Methylbicycloatalaphylline, M-401
 Oriciacridone A, O-106
 Oriciacridone F, O-108
 Prenylcitpressine, P-616
 Pyranofoline, P-879
 Rutacridone, R-167
 Rutagravine, R-169
 Severifoline, S-278
 Teclanthine, T-56
 Thehaplosin, T-351
 1,2,3-Trihydroxyacridone, T-516
 1,3,5-Trihydroxy-2,4-diprenylacridone, T-546
 1,3,8-Trihydroxy-2,7-diprenylacridone, T-548
 1,3,5-Trihydroxy-4-prenylacridone, T-581
 Xanthevodine, X-2

Acridone dimers

Atalanine, A-1526
 Ataline, A-1528
 Bi[5-hydroxynoracronycine], B-129
 Buntanbismine, B-411
 Citbismine B, C-469
 Citbismine D, C-470
 Glycobismine B, G-113
 Glycobismine D, G-114
 Glycobismine E, G-115
 Glycobismine G, G-116

Acridone-coumarin dimers

Acrimarine A, A-114
 Acrimarine J, A-116
 Dioxinoacrimarine A, D-800
 Neoacrimarine A, N-92

Neoacrimarine C, N-93
 Neoacrimarine D, N-94
 Neoacrimarine E, N-95
 Neoacrimarine F, N-96
 Neoacrimarine K, N-97

Pyrido[2,3,4-*kl*]acridines

Alpkinidine, A-639
 Amphimedine, A-944
 Arnoamine A, A-1445
 Ascididemin, A-1459
 Cycloercitine, C-857
 Cycloshermilamine D, C-907
N-Deacetylkuanoniamine D, D-93
 Deoxyamphimedine, D-222
 Dercitine, D-231
 Dibenzo[*b*,*d*]dipyrido[4,3,2-*de*:2',3',4'-*gh*][1,10]phenanthroline, D-299
 Diplamine, D-823
 Eudistone A, E-291
 Eudistone B, E-292
 9-Hydroxyisoascididemnin, H-542
 Isosegoline A, I-316
 Labuanine A, L-2
 Lissoclinidine, L-204
 Meridine, M-206
 Neoamphimedine Y, N-100
 Neoamphimedine, N-99
 Norsegoline, N-308
 Pantherinine, P-73
 Perophoramidine, P-272
 Petrosamine, P-288
 9*H*-Pyrido[4,3,2-*mn*]thiazolo[4,5-*b*]acridin-9-one, P-920
 Sagitol, S-14
 Sebastianine A, S-171
 Sebastianine B, S-172
 Segoline A, S-217
 Shermilamine A, S-286
 Stelletamine, S-506
 Styelsamine C, S-608
 Styelsamine D, S-609
 Subarine, S-619
 Tintamine, T-415
 Varamine A, V-22
 Varamine B, V-23

1,4-Benzoxazin-3-one alkaloids

Cappamensine, C-96
 Monocillinol A, M-687

Benzodiazepine alkaloids

Abbeymycin, A-5
 Anthramycin, A-1081
 Antibiotic A 65636, A-1099
 Antibiotic DC 81, A-1156
 Antibiotic DC 102, A-1157
 Antibiotic DC 105, A-1158
 Antibiotic M 259, A-1202
 Antibiotic RK 1441A, A-1232
 Antibiotic RP 13252, A-1234
 Aplysepine, A-1347
 Benzomalvin B, B-74
 7-Bromo-5-(2-chlorophenyl)-1,3-dihydro-3-hydroxy-2*H*-1,4-benzodiazepin-2-one, B-304
 Chicamycin A, C-373
 Chicamycin B, C-374
 Cyclophenin, C-889
 Diazepinomicin, D-297
 Iforrestine, I-20
 Mazethramycin A, M-137
 Neothramycin A, N-145
 Porothramycin A, P-573
 Prothracarin, P-660
 Tilivalline, T-411
 Tomaymycin, T-424

Cryptolepine type alkaloids

Biscryptolepine, B-168
 Cryptoheptine, C-785

Cryptolepicarboline, C-786
 Cryptomirsine, C-787
 Cryptoquindoline, C-794
 Cryptospirolepine, C-795
 5,10-Dihydro-11*H*-quindolin-11-one, D-505
 5,6-Dihydro-11*H*-quinindolin-11-one, D-506
 Homocryptolepinone, H-329
 Indolo[3,2-*b*]quinoline, I-109
 11-Isopropyl-5-methyl-5*H*-indolo[3,2-*b*]quinoline, I-279
 Quindolinocryptotackeine, Q-16

Simple phenethylamine alkaloids

N-β-Alanyldopamine, A-241
 4-(2-Aminoethyl)-2-bromophenol, A-769
N,N'-Bis[2-(3,5-diiodo-4-methoxyphenyl)ethyl]urea, B-171
 Cannabisin G, C-87
 Chenoalbicin, C-364
 Convolutamine A, C-625
 Convolutamine H, C-628
 Coryneine, C-690
 Dakaramine, D-10
 Densinine, D-219
 2-(2,4-Dibromo-5-methoxyphenyl)ethylamine, D-330
N-(Dihydrosinapoyl)tyramine, D-512
 (3,5-Diiodo-4-methoxyphenyl)ethylamine, D-697
 3,4-Dimethoxyphenethylamine, D-718
 Dioxamine, D-799
 Doisuthine, D-900
 Dopamine, D-920
 Epinine, E-98
 Fumariflorine, F-175
 Hibiscuwanin A, H-271
N-Homoveratroylhomoveratrylamine, H-351
 Hordenine, H-357
 3-Hydroxy-4,5-dimethoxyphenethylamine, H-465
 3-Hydroxy-4-methoxyphenethylamine, H-565
 4-Hydroxy-3-methoxyphenethylamine, H-566
 2-(4-Hydroxy-3-nitrophenyl)ethylamine, H-640
 4-Hydroxyphenethylamine, H-667
 2-(3-Hydroxyphenyl)ethylamine, H-678
 Lissoclinotoxin C, L-207
 Mescaline, M-219
 2-(4-Methoxyphenyl)ethylamine, M-280
 Mololipids, M-680
 Nakirodine A, N-23
 Norbelladine, N-288
 2-Phenylethylamine, P-342
 Polyandrocarpamide A, P-541
 3,4,5-Trihydroxyphenethylamine, T-576
 Turbotoxin A, T-680
 Volutamide A, V-190
 Zanthosin, Z-6

α-Hydroxyphenethylamines

2-Amino-3',4'-dihydroxyacetophenone, A-736

Halogenated tyrosinoids

Agelolin A, A-194
 Anomoian A, A-1068
 Aplysinellin A, A-1351
 Aplysinellin B, A-1352
 Araplysin III, A-1379
 Bastadin 13, B-33
 Bastadin 16, B-34
 Bastadin 1, B-30
 Bastadin 3, B-31
 Bastadin 4, B-32
 Bispsammaplin A, B-217
 α-(Dimethylamino)-3,5-diiodo-*N*-[5-[[3-(3-iodo-4-methoxyphenyl)-2-(methylamino)-1-oxopropyl]amino]pentyl]-4-methoxybenzenepropanamide, D-729
 Fistularin 3, F-70
 Hemibastadin 1, H-100
 Hemibastadinol 1, H-101

Itampolin A, I-350
 Molokaiamine, M-679
 Psammaplin A, P-686
 Psammaplin B, P-687
 Psammaplin C, P-688
 Psammaplin D, P-689
 Psammaplin E, P-690
 Psammaplin F, P-691
 Psammaplin G, P-692
 Psammaplin H, P-693
 Psammaplin I, P-685
 Psammaplysene A, P-694
 Pseudoceratin A, P-707
 Puralidin C, P-809
 Purpuramine H, P-823
 Purpuralidin B, P-824
 Tokaradine A, T-422

Miscellaneous phenethylamines

Acidissiminol, A-77
 AK toxin II, A-221
 Alfileramine, A-260
 Aplyzanzine A, A-1358
 Avicennamine, A-1565
 Cannabisin A, C-84
 Cannabisin E, C-86
 Cannabisin H, C-88
 Convolutamine D, C-626
 Convolutamine E, C-627
 Dambullin, D-12
 Densiflorine, D-217
N-[2-[3,5-Dibromo-4-[[3-(methyl-1-oxo-2-butenyl)amino]propoxy]phenyl]ethyl]-4-hydroxybenzeneacetamide, D-333
 Dispyrin, D-865
 1,2-Epoxy-2-methyl-4-(*N*-methyltyraminyl)-3-butene, E-135
 Gerambullin, G-68
 Grossamide, G-182
 Heliotropamide, H-96
 Hemibastadin 1, H-100
 Integramine, I-157
 Isoalfileramine, I-188
 Jacpaniculine, J-5
 Kuchinoenamine, K-105
 Lissoclibadin 1, L-196
 Lissoclibadin 2, L-197
 Lissoclibadin 3, L-198
 Lissoclinotoxin A, L-205
 Lissoclinotoxin D, L-208
 Lissoclinotoxin E, L-209
 Lissoclinotoxin F, L-210
 Mescaline citrimide, M-220
 Mescaline isocitrimide lactone, M-221
 Orthidine A, O-125
 Orthidine C, O-126
 Orthidine E, O-127
 Peyoglunol, P-296
 Pubesamide B, P-777
 Puhinamide, P-780
 Purpuramine A, P-821
 Purpuramine H, P-823
 Shishididemniol A, S-289
 Storniamide D, S-556
 Taspine, T-36
 Tridentatol A, T-507
 Tridentatol D, T-509
 Tubacetine, T-652
 Tubasencine, T-653
 Tubastrine, T-656
 Tuberine, T-659
 Varacin A, V-20
 Waianaecamine B, W-2

Cinnamic acid amides

Astrophylline, A-1525
 Botryllamide H, B-257
N-[2-(4-Butylphenyl)ethyl]-3-(4-hydroxy-3-methoxyphenyl)-2-propenamide, B-441
 Cannabisin G, C-87
 Casimiroedine, C-176

Celenamide E, C-238
 Cenocladamide, C-245
 Chenoalbicin, C-364
 Cherinonaine, C-368
 5,6-Dihydro-*N*-[3-(7-methoxy-1,3-benzodioxol-5-yl)-2-propenoyl]-2(1*H*)-pyridinone, D-470
N-(Dihydrosinapoyl)tyramine, D-512
 Dioxamine, D-799
 Echinophyllin B, E-19
 Elliptinol, E-69
 Fontaineine, F-124
 Gigantamide A, G-74
 Hibiscuwanin A, H-271
 Magnolamide, M-44
N-Methyl-*N*-(2-phenylethenyl)-3-phenyl-2-oxiranecarboxamide, M-509
 Montamine, M-697
 Nannochelin C, N-28
 Pipericyclobutanamide A, P-450
 Pipericyclobutanamide B, P-451
 Piperpense, P-468
 Piplartine, P-473
 Piriferine, P-479
 Polygonapholine, P-561
 Psammaplysene A, P-694
 Rubescenamine, R-143
 Sarmenamide B, S-87
 Toddaliamide, T-421
 Tunichrome An1, T-677
 Tunichrome Pm 1, T-678
 Xyloallenolide A, X-28
 Zanthomamide, Z-4
 Zanthosin, Z-6

Securinega alkaloids

Flueggenine A, F-93
 Flueggenine B, F-94
 Margaritarine, M-102
 Norsecurine acid, N-306
 Phyllanthine, P-386
 Secuamamine A, S-202
 Securinegine, S-205
 Securinol A, S-207
 Securinol C, S-208
 Simplexine, S-304
 Suffruticodine, S-630
 Suffruticonine, S-631

Betalain alkaloids

Betalamic acid, B-109
 Betanidin, B-110
 Dopaxanthin, D-921
 Indicaxanthin, I-67
 Mesembryanthemins, M-227
 Miraxanthin III, M-647
 Miraxanthin II, M-646
 Miraxanthin I, M-645
 Muscaaurin II, M-769
 Muscaaurin I, M-768
 Muscapurpurin, M-771
 Oleracins, O-80
 Portulacaxanthin III, P-577
 Portulacaxanthin II, P-576
 Portulacaxanthin I, P-575
 Suaedin, S-615
 Tryptophanbetaxanthin, T-642
 Vulgaxanthin II, V-195

Simple isoquinoline alkaloids

7-Amino-1,6-dimethyl-5,8-isoquinolinedione, A-757
 Anhydroberberillic acid, A-1024
 Antibiotic MY 336a, A-1209
 Antibiotic WJ 35, A-1312
 Berbidine, B-102
 Bernumicine, B-105
 Bernumine, B-106
 Cherianoine, C-366
 Chingazumianine, C-386
 Cotarnoline, C-707
 Crispine C, C-757

Crispine D, C-758
 Crispine E, C-759
 Decahydro-8-(1-piperidinylmethyl)isoquinoline, D-113
 3,4-Dihydro-8-hydroxy-6,7-dimethoxyisoquinoline, D-437
 3,4-Dihydro-8-hydroxy-6,7-dimethoxy-1-methylisoquinoline, D-439
 6,7-Dihydro-1,3,6,6-tetramethyl-8(5*H*)-isoquinolinone, D-515
 3,4-Dihydro-*ar*-trimethoxy-1-methylisoquinoline, D-520
 5,8-Dihydroxy-4,7-dimethoxy-2,6-dimethylisoquinolinium(1+), D-569
 6,8-Dimethoxy-1,3-dimethylisoquinoline, D-710
 3,4-Dimethyl-7-(methylamino)-5,8-isoquinolinedione, D-752
 Hydrocotarnine, H-396
 Hydrohydrastinine, H-398
 Intebrine, I-152
 Intebrinine, I-153
 Isoquinoline, I-301
 1,6,8-Isoquinolinetriol, I-306
 1-(8-Isoquinolinylmethyl)pyridinium(1+), I-308
 7-Methoxy-1,6-dimethyl-5,8-isoquinolinedione, M-249
 6,7-Methylenedioxyisoquinoline, M-444
 Mimocin, M-622
 Mimosamycin, M-624
 Perfragilin A, P-255
 Phomopsin A, P-377
 Renierol, R-45
 Ruprechstylil, R-163
 Salsamine, S-36
 Sendaverine, S-234
 Spartocytisine, S-386
 1,2,3,4-Tetrahydro-6,7-dihydroxy-1,1-dimethylisoquinoline, T-152
 1,2,3,4-Tetrahydro-7,8-dimethoxy-1,2-dimethylisoquinoline, T-166
 1,2,3,4-Tetrahydro-7-hydroxy-5-methoxy-2-methylisoquinoline, T-180
 1,2,3,4-Tetrahydro-*ar*-hydroxy-*ar*-trimethoxy-2-methylisoquinoline, T-191
 1,2,3,4-Tetrahydro-*ar*-methoxy-1-methylisoquinoline, T-200
 1,2,3,4-Tetrahydro-5,6,7,8-tetramethoxy-1-methylisoquinoline, T-233
 1,2,3,4-Tetrahydro-5,6,7-trimethoxy-2-methylisoquinoline, T-238
 5,6,7,8-Tetramethoxyisoquinoline, T-276
 Thalflavine, T-306
 Thalpetaline, T-344
 1,6,8-Trihydroxy-3,4-dimethylisoquinoline, T-544
 1,6,8-Trihydroxy-3-methylisoquinoline, T-566
ar-Trimethoxy-1-methylisoquinoline, T-597
 Turcberine, T-681
 Turconidine, T-682
 Viguine, V-101
 Violatinctamine, V-143

Manzamine alkaloids

Ircinol B, I-175
 Kauluamine, K-22
 31-Keto-12,34-oxa-32,33-dihydroircinal A, K-38
 Maeganedin A, M-38
 Manadomanzamine A, M-80
 Manzamine B, M-92
 Manzamine J, M-94
 Manzamine M, M-95
 Nakadomarin A, N-11
 Neokauluamine, N-120
 Xestocyclamine A, X-21
 Xestocyclamine B, X-22

Miscellaneous isoquinoline alkaloids

Aaptamine, A-2
 Amphibine I, A-934

Annoretine, A-1060
 Annosqualine, A-1061
 Antibiotic EGS 1300-1, A-1162
 Antibiotic TMC 2A, A-1292
 Antibiotic TMC 120A, A-1294
 Aspergillitine, A-1477
 Berberal, B-99
 Chaetoindicin A, C-333
 Chaetoindicin B, C-334
 Chlorofusin, C-405
 Cohirsitine, C-565
 Cortistatin B, C-665
 Cortistatine L, C-668
 Corydamine, C-678
 Decumbenine B, D-136
 Drimiopsine A, D-940
 Excentricine, E-318
 Fuscusine, F-229
 Hernandine, H-149
 Hypecumine, H-768
 Illudinine, I-29
 Isochromophilone VI, I-205
 Lissoclotoxin B, L-206
 Litebamine, L-211
 Lophocine, L-261
 Marinamide, M-104
 7-(Methylamino)-3*H*-pyrrolo[2,3-*c*]isoquinoline-6,9-dione, M-387
 Mureidomycin E, M-738
 Napsamycin B, N-34
 Oceanalin A, O-14
 Panaefluoroline A, P-45
 Panaefluoroline B, P-46
 Panaefluoroline C, P-47
 Panaefluoroline D, P-48
 Panaefluoroline E, P-49
 Panaefluoroline F, P-50
 Panaefluoroline G, P-51
 Panaefluoroline H, P-52
 Peyoglutam, P-297
 Phenylalanyl-*N*-[1-(1,2,3,4-tetrahydro-6-hydroxy-7-methoxy-2-methyl-1-isoquinolinyl)ethyl]-glycinamide, P-339
 Pilocereine, P-425
 Siomycin A, S-321
 Siomycin D₁, S-322
 Stephaoxocamine, S-541
 Thiostrepton, T-392
 Usambanoline, U-51

Benzylisoquinoline alkaloids

Aristoquinoline C, A-1440
 Canelillinixine, C-80
 Dehydronormacrostomine, D-163
 Fumafflorine, F-170
 Gandharamine, G-14
 Hebridamine, H-83
 Herveline C, H-154
 Imbricatine, I-32
 Iosevanine, I-317
 Leptopidine, L-121
 Leptopine, L-122
 Leptopinine, L-123
 Longiberine, L-248
 Longifolonine, L-252
 1-(4-Methoxybenzoyl)-6,7-methylenedioxyisoquinoline, M-242
 Miltanthine, M-620
 Miltanthoridine, M-621
 Mollinedine, M-672
 Nymphaedaline, N-347
 Papaveraldine, P-77
 Papaverine, P-79
 Polysignine, P-563
 Protosinomenine, P-668
 Sauvagnine, S-108
 Sevanine, S-274
 Takatonine, T-16
 Thalimicrinone, T-329
 Theoneberine, T-356

Pseudobenzylisoquinoline alkaloids

Berbitine, B-103
 Fumarizine, F-179
 Polyberbine, P-548
 Polycarpine, P-554
 Rugosinone, R-156
 Taxilamine, T-50

Bisbenzylisoquinoline alkaloids (aryl links only)

Isopyruthaldine, I-296
 Isopythaldine, I-298
 Pisopowine, P-484

Bisbenzylisoquinoline alkaloids (one ether link)

Berbamunine, B-98
 Chillanamine, C-376
 Dauricine, D-88
 Efatine, E-42
 Grisabine, G-177
 Isopyruthaline, I-297
 Isopythaline, I-299
 Isothalicine, I-330
 Isothalictrine, I-331
 Isothalarine, I-332
 Magnolamine, M-45
 Malekulatine, M-71
 Neferine, N-87
 Neosutchuenenine, N-142
 Phaeantharine, P-300
 Sutchueneneonine, S-646
 Sutchuenenine, S-647
 Thaliatrine, T-307
 Thalibrine, T-309
 Thalaricebine, T-332
 Thalirugidine, T-334
 Thalirugine, T-335
 Thalistine, T-337
 Vanuatine, V-19
 Vateamine, V-43

Bisbenzylisoquinolines (one ether/one aromatic link)

Cordobine, C-643
 Isothamidine, I-334
 Oxandrine, O-149
 Philogaline, P-360
 Pseudoxandrine, P-736
 Rodiasine, R-114

Bisbenzylisoquinoline alkaloids (two ether links)

Calafatimine, C-22
 Calafatine, C-23
 Cepharanthine, C-275
 Cheratamine, C-365
 Cissampentin, C-467
 Curicycleatjine, C-811
 Cycleanine, C-835
 Cycleatjehine, C-836
 Daphnine, D-60
 Dryadodaphnine, D-944
 Fastrine, F-18
 Gyrolidine, G-235
 Hernandezine, H-148
 Isotetrandrine, I-328
 Isothalicberine, I-329
 Lauberine, L-67
 Limacusine, L-175
 Nemuarine, N-91
 Neothalfine, N-144
 Osornine, O-130
 Panurensine, P-76
 Peinamine, P-171

Phaeanthine, P-301
 Protocuridine, P-663
 Repandine, R-46
 Sciadenine, S-148
 Sciadoline, S-149
 Stebisimine, S-502
 Stepinonine, S-548
 Sutchenensine, S-648
 Tetrandrine, T-286
 Thalfinine, T-305
 Thalibrunine, T-311
 Thaliberberine, T-313
 Thalidasine, T-320
 Thalmethine, T-339
 Thalmiculine, T-340
 Thalmine, T-341
 Thalmirabine, T-343
 Thalrugosaminine, T-346
 Thalsimidine, T-347
 Warifteine, W-9

Bisbenzylisoquinoline alkaloids (two ether/one aromatic link)

Medelline, M-145
 Tiliacorine, T-406
 Tiliacorinine, T-407
 Tiliarsine, T-410
 Yanangine, Y-5

Bisbenzylisoquinoline alkaloids (three ether links)

Angchibangkine, A-1006
 Apateline, A-1331
 Coculinine, C-552
 Cocsupendine, C-553
 Dielsine, D-384
 Gasabiimine, G-26
 Insularoline, I-150
 Isotrilobine, I-339
 Kohatine, K-56
 Menisarine, M-197
 Micranthine, M-591
 Pseudorepanduline, P-725
 Repanduline, R-47
 Trigilletimine, T-514

Secobisbenzylisoquinoline alkaloids

Bargustanine, B-26
 Fenfangjine H, F-34
 Jhelumine, J-37
 Karakoramine, K-16
 Neotrilobine, N-146
 Punjabine, P-800
 Revolutinone, R-68
 Secantioquine, S-175
 Secocepharanthine, S-179
 Secolucidine, S-190

Cularine group alkaloids

Aristoyagonine, A-1443
 Cancentrine, C-76
 Dioxocularine, D-801
 Fissistigine B, F-66
 Gouregine, G-148
 Henderine, H-108
 Oxocompostelline, O-169
 Oxocularine, O-171
 Oxosarcophylline, O-207
 Sarcocapnidine, S-73
 Yagonine, Y-4

Secoisoquinoline alkaloids

Aconcaguine, A-84
 Ceratocapnidine, C-278
 Ceratonicine, C-279
 Isonoyaine, I-257
 Noyaine, N-326

Secocularidine, S-180
 Secosarcocapnidine, S-198

Quettamine alkaloids

Quettamine, Q-8
 Secoquettamine, S-197

Dibenzopyrrocoline alkaloids

Oubatchensine, O-138

Indenobenzazepine alkaloids

Bulgaramine, B-400
 Fumarofine, F-180
 Himalayamine, H-276
 Lahoramine, L-15
 Lahorine, L-17
 Ribasine, R-94

Pavine alkaloids

Fauripavine, F-26
 Herveline C, H-154
 Neocaryachine, N-104
 Pennsylvavoline, P-196
 Thalimonine, T-330
 4,8,9-Trimethoxy-N-methyl-2,3-methylenedioxy-pavinane, T-598

Isopavine alkaloids

Amurensine, A-954
 Isothalisopavine, I-333
 Przewalidine, P-681
 Reframine, R-36
 Thalisoavine, T-336

Proaporphine alkaloids

Croton linearis Alkaloid E, A-498
 Bracteine, B-275
 Coyhaiquine, C-714
 Cryprochine, C-781
 Isocryprochine, I-211
 Jaculadine, J-6
 Misramine, M-650
 Prooxocryptochine, P-644
 Roehybridine, R-116
 Roehybrine, R-117
 Roemeramine, R-119
 Stepharinosine, S-542

Proaporphine-benzylisoquinoline alkaloid dimers

Coyhaiquine, C-714
 Pakistanamine, P-15
 Rupancamine, R-162

Aporphine alkaloids

Acutiaporberine, A-132
 Alkaloid PO3, A-583
 Annolatine, A-1050
 Artabonatin A, A-1448
 Artabonatin C, A-1449
 Artacinatine, A-1450
 Atherospermidine, A-1531
 Beccapoline, B-53
 Belemine, B-55
 Bipowine, B-142
 Bipowinone, B-143
 Bulbodione, B-399
 Cabudine, C-2
 Cassamedine, C-179
 Cassythic acid, C-187
 Cataline, C-190
 Cepharadione A, C-272
 Cepharadione B, C-273
 Clemaine, C-523

Corydione, C-684
 Dactylidine, D-5
 Dehatriline, D-149
 Dehydrodicentrine, D-156
 Dehydroglauicine, D-159
 Dehydronuciferine, D-164
 Dehydrooemerine, D-166
 Dehydroxylophine, D-175
 Dicentrinone, D-347
 6a,7-Didehydro-7-hydroxy-1,2-dimethoxyaporphine, D-373
 4,5-Dihydro-1,9-dihydroxy-2,11-dimethoxy-7-oxoaporphine, D-418
 4,5-Dioxodehydrocrebanine, D-802
 Duguenaine, D-946
 Dugespixine, D-947
 Duguetine, D-948
 Duguevalline, D-949
 Faurine, F-25
 Fauripavine, F-26
 Fissilandione, F-65
 7-Formyldehydrohernergerine, F-133
 7-Formyldehydronornantentine, F-134
 7-Formyldehydroovigerine, F-135
 7-Formyldehydrothalicimidine, F-136
 Fuseine, F-230
 Glauine, G-93
 Goudotianine, G-147
 Griffinine, G-173
 Griffithdione, G-175
 Guadiscidine, G-188
 Guatterine, G-213
 Guattescidine, G-214
 Guattouregidine, G-215
 Hemiargine B, H-97
 Hernandonine, H-150
 Heteropsine, H-160
 Imenine, I-34
 Kuafumine, K-104
 Lanuginosine, L-40
 Laurodionine, L-74
 Lettowianthine, L-145
 Liriodendronine, L-193
 Liriodenine, L-194
 Melosmine, M-194
 1,2-Methylenedioxy-6a,7-dehydroaporphine-10,11-quinone, M-445
 Mollisine, M-678
 Moschatoline, M-712
 Nandazurine, N-24
 Natalinine, N-53
 Ocokryptine, O-25
 Ocominarone, O-26
 Oliveridine, O-85
 Oliveroline, O-87
 Ouregidione, O-139
 Ovigeridimerine, O-142
 Oviherangerine, O-144
 Oviisocorydine, O-145
 Oxobaicaline, O-166
 Oxobuxifoline, O-168
 Oxocrebanine, O-170
 Oxoglauicine, O-178
 7-Oxohernandaline, O-179
 7-Oxohernangerine, O-180
 Oxoisocalycine, O-183
 Oxolaureline, O-187
 Oxophoebine, O-195
 Oxopurpureine, O-203
 Oxoputerine, O-204
 Oxostephanine, O-209
 Pachyconfine, P-4
 Pancoridine, P-54
 Pancorinine, P-56
 Pennsylvavoline, P-196
 Pentouregine, P-234
 Peruvianine, P-275
 Phoenanthusine, P-370
 Pontevedrine, P-566
 Rurrebanidine, R-164
 Sinomendine, S-312
 Sonodione, S-367
 Spixianine, S-469

Splendidine, S-470
 Stephadione, S-535
 Stephalagine, S-536
 Subsessiline, S-627
 Telazoline, T-62
 Telikovinone, T-64
 Telisatin A, T-65
 6,7,9,10-Tetrahydro-5-(4-hydroxy-3,5-dimethoxyphenyl)-5*H*-benzo[*g*][1,3]dioxolo[4',5':4,5]benzo[*de*]pyrido[3,2,1-*ij*]quinolin-7-one, T-176
 Thalibealine, T-308
 Thaliminine, T-315
 Trichoguattine, T-489
 Trivalvone, T-628
 Tuberosinone C, T-662
 Tuberosinone, T-661
 Unonopsine, U-36
 Urabaine, U-41
 Vitricine, V-165

Aporphine-benzylisoquinoline alkaloid dimers

Adiantifoline, A-139
 Dehydrothalicarpine, D-170
 Fauridine, F-24
 Faurithaline, F-27
 Huangshanine, H-366
 Khyberine, K-39
 Oxothalicarpine, O-214
 Pakistanine, P-16
 Przewaline, P-682
 Przewalskine, P-683
 Przewalstine, P-684
 Thaldimerine, T-302
 Thalicarpine, T-312
 Thalifaberine, T-322
 Thalifabine, T-323
 Thaliphine, T-331
 Thalirevolutine, T-333
 Thalmineline, T-342
 Uskudaramine, U-55

Oxoisoporphine alkaloids

Bianfugecine, B-116
 Bianfugedine, B-117
 Daurioisoporphine B, D-89
 Dauriporphine, D-90
 Lakshminine, L-19
 Menisporphine, M-199
 Tyraminoporphine, T-697

Aristolochic acid alkaloids

Aristolin, A-1434
 Aristoloin I, A-1435
 Aristoloterpenate I, A-1436
 Aristophyllide A, A-1439
 Debilic acid, D-102
 6-Hydroxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, H-600
 8-Methoxy-3,4-methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-262
 3,4-Methylenedioxy-10-nitro-1-phenanthrenecarboxylic acid, M-447

Aristolactam alkaloids

Aristo red, A-1424
 Aristolactam C II, A-1428
 Aristolactam I, A-1427
 Aristolactam IIIa, A-1429
 Doryflavine, D-925

Phenanthrene alkaloids

Andesine, A-999
 Annoretine, A-1060

Atherosperminine, A-1532
 Chiloenine, C-381
 Corydinemethine, C-683
 Fissicesine, F-64
 Hebridamine, H-83
 Isouvariopsine, I-343
 Litebamine, L-211
 2-Methoxyatherosperminine, M-238
N-Methylsecoglaucine, M-559
 Romucosine I, R-124
 Santiagonamine, S-64
 Stephenanthrine, S-546
 Thalicipureine, T-316
 Thalicthuberine, T-318
 Thaliglucine, T-327
 Thalihazine, T-328
 Uvariopsamine, U-63
 Uvariopsine, U-64

Azaaporphine alkaloids

Eupolauramine, E-297
 Imbiline I, I-31
 Sampangine, S-49

Azafluoranthrene alkaloids

Grandirubrine, G-159
 Imeluteine, I-33
 Indeno[1,2,3-*ij*][2,7]naphthyridine, I-65
 Norrufescine, N-304
 Triclisine, T-498

Morphine alkaloids

Acutudaurine, A-133
 Acutumine, A-134
 Alkaloid PP1, A-586
 Amurinine, A-956
 Bismorphine A, B-205
 Bismorphine B, B-206
 Cancentrine, C-76
 Carococculine, C-164
 Cephasamine, C-276
 Codeine, C-555
 Corsutine, C-663
 6,7,8,9,10,14-Hexadehydro-4,5-epoxy-3,6-dimethoxy-17-methylthebinan, H-226
 14-Hydroxycodine, H-462
 Hypserpine, H-775
 Isostephodoline, I-325
 Meconoquintupline, M-144
 ψ -Morphine, M-703
 Neopine, N-134
 Oreobeiline, O-103
 Saludimerine A, S-37
 Salutadimerine, S-38
 Sinococculine, S-311
 Sinomenine, S-313
 Somniferine, S-362
 Tannagine, T-28
 Tridictyophylline, T-511
 Zippeline, Z-20

Erythrina alkaloids

Cocculine, C-538
 Cocculitine, C-539
 Cocculolidine, C-540
 Coccutrine, C-542
 3-Demethoxy-2,3-methylenedioxyerythroline, D-200
 Dihydroerysodine, D-431
 Erymelanthine, E-198
 Erysinophorine, E-200
 Erysofporine, E-201
 Erysotramidine, E-202
 Erysotrine, E-203
 Erythlaurine, E-204
 Erythartine, E-206
 Erythratidine, E-207
 Erythratine, E-208

α -Erythroidine, E-215
 Lenticellarine, L-92

Dibenzazecine alkaloids

Bractazonine, B-273
 Crassifolazonine, C-727
 Dysazecine, D-962
 Erybidine, E-194
 Laurifonine, L-73
 Neodihydrothebaine, N-111
 Protostephanine, P-671

Hasubanan alkaloids

Stephania japonica Alkaloid A, A-378
 Delavaine, D-181
 Hasubanonine, H-70
 Limalongine, L-176
 Longanine, L-244
 Longetherine, L-247
 Miersine, M-605
 Prometaphanine, P-639
 Prosthephyssine, P-658
 Prosthephanaberrine, P-659
 Runanine, R-161
 Stephabeneine, S-529
 Stephadiamine, S-533
 Stephamiersine, S-537
 Stephavanine, S-545

Protoberberine alkaloids

Acutiaporberine, A-132
 Alborine, A-247
 Alkaloid PO4, A-584
 Alkaloid PP1, A-585
 Anisocycline, A-1042
 Berberilycine, B-100
 Berberine, B-101
 Berpodine, B-107
 Bisjatrorrhizine, B-193
 Caseadinium(1+), C-173
 Coptisine, C-638
 Corydaline, C-675
 Corysamine, C-699
 Coulteroberbinone, C-709
 Dehydroapocavidine, D-150
 Dehydrocorydaline, D-154
 Dehydrothalictricavine, D-171
 Dihydrocoptisine, D-409
 Groenlandicine, G-181
 Hediamine, H-86
 11-Hydroxy-12-methoxycoptisine, H-559
 Karachine, K-15
 Lambertine, L-23
 Latifolian A, L-55
 Lienkonine, L-167
 Longiberine, L-248
 Malacitanine, M-57
 Mequinine, M-200
 1-Methoxyberberine, M-243
 13-Methoxyoxoberberine, M-279
 8-Methylanibacanine, M-388
 8-Oxoepiberberine, O-177
 8-Oxopseudopalmatine, O-202
 Oxyberberine, O-216
 Palmatine chloroform, P-35
 Palmatine, P-34
 Pseudoberberine, P-705
 Pseudocoptisine, P-710
 Pseudopalmatine, P-719
 Solidaline, S-353
 Staudine, S-497
 Stephabine, S-531
 Thalibealine, T-308
 Thalifaurine, T-325
 Theoneberine, T-356
 2,10,13-Trihydroxydibenzo[*a,g*]quinolizinium(1+), T-541
 Worenine, W-22
 Yuanamide, Y-18
 Zijinlongine, Z-19

Isoindolobenzazepine alkaloids

Pictonamine, P-419

Narceine alkaloids

Adlumiceine enol lactone, A-142
 Adlumidiceine, A-143
 Bicucullinidine, B-120
 Coryrutine, C-698
 Cryptopleurospermine, C-793
 Fumaramidine, F-171
 Fumschleicherine, F-191
 Microcarpine, M-592
 Narceine, N-35
 Narlumidine, N-51
 Saxoguattine, S-111

Protopine alkaloids

Argemexicaine A, A-1408
 Argemexicaine B, A-1409
 Constrictosine, C-624
 Cryptopine, C-790
 Fagarine I, F-2
 Fagarine II, F-3
 Fagarine III, F-4
 Muramine, M-734
 Protopine, P-667
 Pseudoprotopine, P-723

Rheadine alkaloids

Glaucamine, G-87

Phthalidisoquinoline alkaloids

Bicucine, B-118
 Bicuculline, B-119
 Cordrastine, C-644
 Corydecumbine, C-679
 Decumbenine, D-135
 Egenine, E-43
 Fumadensine, F-169
 Hydrastine, H-394
 Hypecoumine, H-767
 Narcotinehemiacetal, N-46

Spirobenzylisoquinoline alkaloids

Africanine, A-173
 Crystewartine, C-700
 Densiflorine, D-218
 Hyperectine, H-771
 Raddeanamine, R-8
 Severzinine, S-279

Secoberberine alkaloids

Berberal, B-99
 Fumafiorine, F-170
 Leptopidine, L-121
 Leptopine, L-122
 Leptopinine, L-123
 Narcotinediol, N-45
 Procumbine, P-634

Benzo[c]phenanthridine alkaloids

8-Acetyldihydrochelerythrine, A-34
 8-Acetyldihydronitidine, A-35
 Ailanthoidine, A-209
 Ambinine, A-689
 Arnottianamide, A-1446
 Avicine, A-1566
 Bocconoline, B-239
 Buesgeniine, B-390
 Bulleyanaline, B-407
 Bungeanine, B-410
 Caymandimerine, C-225
 Chelelactam, C-354
 Chelerythridimerine, C-355
 Chelerythrine, C-356

Chelirubine, C-360
 Corynoloxine, C-693
 Dihydroavicine, D-404
 Dihydrochelerythrine, D-407
 Dihydrochelerythryl-8-acetaldehyde, D-408
 Dihydronitidine, D-482
 Dihydrosanguilutine, D-509
 Dihydrosanguinarine, D-510
 5,6-Dihydro-2,3,7,8-tetramethoxy-5-methylbenzo[c]phenanthridin-6-one, D-514
 7,9-Dimethoxy-2,3-methylenedioxybenzo[c]phenanthridine, D-714
 Fagaronine, F-5
 8-Hydroxymethyldihydronitidine, H-595
 8-Hydroxymethyldihydrosanguinarine, H-596
 Integriamide, I-159
 Isochelidonine, I-203
 8-Methylnorchelerythrine, M-494
 8-(4-Methyl-2-oxopentyl)dihydrochelerythrine, M-500
 Nitidine, N-239
 Nitrotyrasanguinarine, N-258
 8-(2-Oxobutyl)dihydrochelerythrine, O-167
 Pancorine, P-55
 Papavoriendine, P-81
 Rhoifoline B, R-87
 Sanguilutine, S-55
 Sanguinarine, S-56
 Sanguirubine, S-60
 Simulanoquinoline, S-305
 Simulansamide, S-306
 Tridecanochelerythrine, T-503
 Zanthocadinanin A, Z-3
 Zanthoxyline, Z-7

Phenethylisoquinoline alkaloids

Jolantinine, J-45

Homomorphinandienone alkaloids

Alkaloid CC 2, A-459
 Alkaloid CC 3b, A-465
 Colchiritchine, C-573

Homooporphine alkaloids

Bitlisine, B-228

Homoprooporphine alkaloids

Bulbocodine, B-398
 Colchilutine, C-572
 Jolantamine, J-43
 Jolantimine, J-44
 Kesselridine, K-36
 Luteicine, L-306
 Luteidine, L-307
 Regecoline, R-37
 Regelinone, R-38
 Robustamine, R-113
 Trigamine, T-513

Colchicine alkaloids

Alkaloid AM 3, A-382
 Colchiciline, C-568
 Colchifoline, C-571
 Demecolcinone, D-198
 α -Lumicolchicine, L-286
 β -Lumicolchicine, L-287
 10,11-Oxy-10,12 α -cyclo-10,11-secocolchicine, O-220

Cephalotaxus alkaloids

Anhydroharringtonine, A-1027
 Biscephalezomine A, B-165
 Biscephalezomine B, B-166
 Biscephalezomine C, B-167
 Cephalezomine A, C-252
 Cephalezomine G, C-253

Cephalotaxidine, C-270
 Demethylneodrupacine, D-202
 Drupacine, D-942
 Drupangtonine, D-943
 Hainanensine, H-10
 Harringtonine, H-66
 Isocephalotaxine, I-201
 Isoharringtonine, I-238
 Neoharringtonine, N-117

Homoerythrina alkaloids

Cephalofortuneine, C-260
 Comosine, C-587
 Comosivine, C-588
 Dyshomerythrine, D-963
 Fortuneine, F-143
 Holidine, H-321
 Homoerysotine, H-331
 Isophellibilidine, I-263
 Lucidinine, L-278
 Phellibiline, P-315
 Schelhammeridine, S-125
 Schelhammerine, S-126
 Selaginoidine, S-218
 Taxodine, T-51

Apogalanthamine alkaloids

Craugosodine, C-730

Galanthamine alkaloids

Habranthine, H-2
 Narcisine, N-41
 Pallidiflorine, P-30

Haemanthidine alkaloids

Amabiline, A-670
 Amaryllisine, A-675
 Ambelline, A-683
 Aphaemanthamine, A-1363
 Bowdensine, B-261
 Bujeine, B-395
 Buphanamine, B-413
 Crinamabine, C-742
 Crinatine, C-746
 Crinisine, C-749
 Cripowellin A, C-752
 Cripowellin B, C-753
 Delagoenine, D-176
 Delagoensine, D-177
 Haemanthamine, H-5
 Haemanthidine, H-6
 Haemultine, H-9
 Havanine, H-78
 Hippeastidine, H-293
 Narcidine, N-40
 Nerbowdine, N-169
 Oliganine, O-81
 Papyramine, P-86
 Powelline, P-581

Graciline alkaloids

Digracine, D-399
 Graciline, G-151
 Obesine, O-2

Tazettine alkaloids

Egonine, E-44
 Littoraline, L-215
 Ornazidine, O-118
 Pretazettine, P-626
 Tazettine, T-53

Pancracine alkaloids

Montabuphine, M-696
 Nangustine, N-26
 Pancracine, P-57

Lycorine alkaloids

Amarbellisine, A-674
 Anhydrolycorinium, A-1030
 Assoanine, A-1508
 Caribine, C-162
 Falcatine, F-7
 Galasine, G-8
 Hippadine, H-290
 Latindine, L-58
 Lutessine, L-310
 Mooreine, M-701
 Narcissidine, N-43
 Pluviine, P-528
 Pratorinine, P-586
 Pseudolycorine, P-716
 Roserine, R-130
 Ungeremine, U-34
 Vasconine, V-35
 Zefbetaine, Z-12

Lycorenine alkaloids

Candimine, C-78
 Cliviahaksine, C-528
 Clivisyaline, C-529
 Clivojuline, C-530
 Clivonidine, C-531
 Clivonine, C-532
 Galanthindole, G-4
 Galanthusine, G-5
 Hippeastrine, H-294
 Hostasine, H-362
 5-Hydroxyhomolycorine, H-504
 Krigeine, K-102
 Lycorenine, L-347
 Lycosinine A, L-351
 Miniatine, M-629
 Nerinine, N-171
 Oduline, O-68
 Unsevine, U-37

Narciclasine alkaloids

Crinasidine, C-744
 8,9-Methylenedioxyphenanthridine, M-448
 Narciclasine, N-37
 Narcicirinine, N-38
 Pancratistatin, P-58

Miscellaneous Amaryllidaceae alkaloids

Aestivine, A-163
Pancreatium Alkaloid 6, A-323
Pancreatium Alkaloid 31, A-324
Leucojum Alkaloid C, A-448
Leucojum Alkaloid D, A-484
Narcissus Alkaloid D, A-489
Sternbergia Alkaloid III, A-527
Ungernia Alkaloid IV, A-529
Rhodophiala bifida Alkaloid VI, A-608
Rhodophiala bifida Alkaloid XIII, A-613
Rhodophiala bifida Alkaloid XXV, A-615
Galanthus caucasicus Alkaloid, A-300
Pancreatium sickenbergi Alkaloid, A-325
Ungernia severtzovii Alkaloid, A-338
Zephyranthes Alkaloid, A-341
 Annapawine, A-1048
 Augustamine, A-1539
 Bodamine, B-240
 Brunsdonnine, B-372
 Brunsvinine, B-373
 Buphacetine, B-412
 Buphanine, B-414
 Coranicine, C-640
 Coruscine, C-669
 Crinatine, C-747
 Crinosine, C-750
 Crispanine, C-754
 Crispine, C-755
 Daphnarcine, D-28

Distichamine, D-867
 Flexamine, F-87
 Flexine, F-89
 Galanthamidine, G-2
 Gracilamine, G-150
 Hippacine, H-289
 Hippandrine, H-291
 Hippauline, H-292
 Insulamine, I-149
 Irenine, I-176
 Ismine, I-186
 Krelagine, K-98
 Macranthine, M-9
 Magnarcine, M-41
 Narcicriptine, N-39
 Narcisline, N-42
 Narzettine, N-52
 Neflexine, N-88
 Nerifline, N-170
 Nerispine, N-172
 Nerundine, N-173
 Petomine, P-282
 Plicamine, P-524
 Plicane, P-525
 Poetamine, P-531
 Poetaricine, P-532
 Poeticine, P-533
 Powellidine, P-580
 Punikathine, P-799
 Robecine, R-112
 Rulodine, R-158
 Secoplicamine, S-195
 Sickenbergine, S-294
 Squamigerine, S-486
 Sternine, S-551
 Valloporfine, V-12
 Vallotidine, V-13
 Vallotine, V-14
 Varadine, V-21
 Yemensine, Y-8

Mesembrenoid alkaloids

4-[2-(Acetylmethylamino)ethyl]-4-(4-hydroxy-3-methoxyphenyl)-2,4-cyclohexadien-1-one, A-60
Sceletium Alkaloid A₄, A-346
Sceletium namaquense Alkaloid, A-330
 Channaine, C-347
 Crinafolidine, C-741
 Sceletenone, S-123

Emetine group alkaloids

Alangiside, A-239
Alangium Alkaloid AL64, A-381
 Emetine, E-72
 Ipecosidic acid, I-167

Phenanthroindolizidine alkaloids

Anhydrodehydrotylophorinidine, A-1025
 Hypoestestatin 2, H-773
 Phyllosteminine, P-390
 Polycanthidine, P-549
 Tylohirsutinine, T-690
 Tyloindicine F, T-692
 Tyloindicine G, T-693
 Tyloindicine H, T-694
 Tyloindicine I, T-691
 Vincetene, V-122

Phenanthroquinolizidine alkaloids

Alihirsutine A, A-263
 Cryptopleuridine, C-791
 Kayawongine, K-23

Ericamycins

Actinoplanone D, A-128
 Albofungin, A-244

Antibiotic 4181B, A-1121
 Antibiotic BE 19412A, A-1135
 Antibiotic Sch 42137, A-1246
 Antibiotic Sch 54445, A-1248
 Antibiotic Sch 56036, A-1249
 Antibiotic TMC 66, A-1288
 Cervinomycin A₁, C-287
 Citreamicin α , C-473
 Citreamicin β , C-474
 Citreamicin γ , C-475
 Citreamicin η , C-476
 Echinosporicin, E-26
 Ericamycin, E-175
 Fredericamycin A, F-152
 Fredericamycin B, F-153
 Fredericamycin E, F-154
 25-Hydroxy-24-oxokibdelone C, H-655
 Isokibdelone A, I-240
 Kibdelone A, K-40
 Lysolipin I, L-374
 Lysolipin X, L-375
 Simaomicin β , S-301
 Xantholipin, X-10

Simple indole alkaloids

3-Acetylandole, A-59
 2-Acetyl-3-methylindole, A-61
 Acremoauxin A, A-109
 Acremoauxin B, A-110
 Almazolone, A-636
 3-(Aminomethylene)-1,3-dihydro-2*H*-indole-2-thione, A-836
 Annonidine F, A-1057
 Antibiotic SF 2140, A-1255
 Aplysinopsin, A-1357
 Arcyriarubin A, A-1394
 Asterredione, A-1517
 Bengacarboline, B-58
 Birnbaumin A, B-148
 2,2-Bis(6-bromo-3-indolyl)ethylamine, B-162
 Bis(6-bromo-2-tryptaminyl) disulfide, B-163
 3,3'-Bis(indolylmethyl)dimethylammonium, B-190
 3,5-Bis(3-methyl-2-butenyl)-1*H*-indole, B-200
 3,6-Bis(3-methyl-2-butenyl)-1*H*-indole, B-201
 3,7-Bis(3-methyl-2-butenyl)-1*H*-indole, B-202
 Brasilidine A, B-281
 Brassicanal C, B-283
 Brassinin, B-284
 Brassitin, B-285
 6-Bromo-3-chloro-1*H*-indole, B-303
 7-Bromo-2,3-dichloro-1*H*-indole, B-308
 6-Bromo-1,2-dihydro-2,2-bis(methylthio)-3*H*-indol-3-one, B-309
 6-Bromo-4,5-dihydroxyindole, B-310
 6-Bromo-4,7-dihydroxyindole, B-311
 6-Bromo-2-(1,1-dimethyl-2-propenyl)-1*H*-indole-3-carboxaldehyde, B-315
 6-Bromo-3-(hydroxyacetyl)-1*H*-indole, B-320
 6-Bromo-3-hydroxy-1*H*-indole, B-325
 6-Bromo-5-hydroxy-1*H*-indole, B-326
 6-Bromo-5-hydroxy-1*H*-indole-3-carboxaldehyde, B-327
 5-Bromo-1*H*-indole-3-acetic acid, B-331
 3-Bromo-1*H*-indole, B-330
 6-Bromo-1*H*-indole-3-carboxaldehyde, B-333
 6-Bromo-1*H*-indole-3-carboxylic acid, B-334
 6-Bromo-1*H*-indole-2,3-dione, B-335
 6-Bromo-2-(methylthio)-3*H*-indol-3-one, B-341
 4(7)-Bromo-2,3,7(2,3,4)-trichloro-1*H*-indole, B-354
 Brussalexin A, B-374
 Caulidine A, C-216
 Caulidine B, C-217
 Caulindole A, C-218
 Caulindole C, C-219
 Cephalandole A, C-249
 Cephalandole B, C-250
 Cephalandole C, C-251
 Cephalinone A, C-255
 Cephalinone B, C-256
 Cephalinone C, C-257

- Cephalinone D, C-258
 Chelonin A, C-361
 4-Chloro-1*H*-indole-3-acetic acid, C-410
 3-Chloro-1*H*-indole, C-409
 5-Chloro-6-methoxy-1-methyl-1*H*-indole-2,3-dione, C-412
 Citorellamine, C-471
 Costinone A, C-705
 Costinone B, C-706
 4,6-Dibromo-2,3-bis(methylthio)-1*H*-indole, D-304
 2,7-Dibromo-3-chloro-1*H*-indole, D-308
 4,7-Dibromo-2,3-dichloro-1*H*-indole, D-310
 3,6-Dibromo-1*H*-indole, D-326
 4,6-Dibromo-1*H*-indole, D-327
 5,7-Dibromo-6-methoxy-1*H*-indole, D-329
 4,6-Dibromo-2-methyl-1*H*-indole, D-332
 4,6-Dibromo-2-methylthio-1*H*-indole, D-334
 2,3-Dihydro-5,6-dihydroxy-1,1-dimethylindolium(1+), D-419
 2,3-Dihydro-1-formyl-2-hydroxy-5-methoxy-6-methyl-1*H*-indole, D-432
 1,3-Dihydro-3-hydroxy-4-methoxy-3-methyl-2*H*-indol-2-one, D-443
 2,3-Dihydro-1*H*-indol-5-ol, D-465
 2,3-Dihydro-2-oxo-1*H*-indole-3-acetic acid, D-485
 α -(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)-1*H*-indole-3-acetonitrile, D-487
 5,6-Dihydroxy-1*H*-indole-2-carboxylic acid, D-598
 4,5-Dihydroxy-1-methyl-3-oxo-2-(trichloromethyl)-3*H*-indolium(1+), D-630
 3,3-Di-1*H*-indol-3-yl-2-butanone, D-689
 4-[(Di-1*H*-indol-3-yl)methyl]phenol, D-692
 3,3-Di-1*H*-indol-3-yl-1,2-propanediol, D-693
 3-[(Dimethylamino)methyl]-5,7-dimethoxyindole, D-733
 3-[(Dimethylamino)methyl]indole, D-736
 2,4-Dimethyl-1*H*-indole, D-751
 Dithyranitrile, D-873
 Echinonulfone A, E-27
 Echinonulfone acid C, E-28
 3-Ethyl-1,3-dihydro-3-methoxy-2*H*-indol-2-one, E-236
 3-Formyl-1*H*-indole-2-sulfonic acid, F-140
 Galanthindole, G-4
N-[[3-(β -D-Glucopyranosyloxy)-2,3-dihydro-2-oxo-1*H*-indol-3-yl]acetyl]aspartic acid, G-103
 Glusun I, G-105
 Glusun II, G-106
 Hamigeramide, H-42
 3-(Hydroxyacetyl)-1*H*-indole, H-400
 3-(2-Hydroxyethyl)-6-prenylindole, H-497
 5-Hydroxy-3-(2-hydroxyethyl)-1*H*-indole, H-505
 5-Hydroxy-1*H*-indole-3-acetic acid, H-525
 5-Hydroxy-1*H*-indole-3-carboxaldehyde, H-528
 6-Hydroxy-1*H*-indole-3-carboxaldehyde, H-529
 6-Hydroxy-1*H*-indole-2,3-dione, H-533
 2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-hexanone, H-535
 2-Hydroxy-1-(1*H*-indol-3-yl)-4-methyl-3-pentanone, H-536
 5-Hydroxy-3-isothiocyanato-4-methoxyindole, H-548
 3-Hydroxy-2-octadecylindole, H-646
 2-Hydroxy-6-prenylindole, H-701
 Hyrtinadine A, H-779
 Hyrtiosin B, H-782
 1*H*-Indole-3-acetaldehyde, I-80
 1*H*-Indole-3-acetic acid, I-81
 1*H*-Indole-3-butanolic acid, I-83
N-(1*H*-Indole-3-carbonyl)urea, I-84
 1*H*-Indole-3-carboxaldehyde, I-86
 1*H*-Indole-5-carboxaldehyde, I-87
 1*H*-Indole-7-carboxaldehyde, I-88
 1*H*-Indole-2-carboxylic acid, I-89
 1*H*-Indole-3-carboxylic acid, I-90
 1*H*-Indole-2,3-diol, I-91
 1*H*-Indole-2,5-diol, I-92
 1*H*-Indole-5,6-diol, I-93
 1*H*-Indole-6,7-diol, I-94
 1*H*-Indole-2,3-dione, I-95
 1*H*-Indole-3-ethanol, I-96
 Indole, I-79
 1*H*-Indole-3-methanamine, I-97
 1*H*-Indole-3-methanol, I-98
 1*H*-Indole-7-methanol, I-99
 1*H*-Indole-3-propanoic acid, I-100
 1*H*-Indol-2-ol, I-103
 1*H*-Indol-3-ol, I-104
 3*H*-Indol-3-one, I-106
 1*H*-Indol-3-ylacetyl-*myo*-inositol, I-115
 4-(1*H*-Indol-3-yl)-1,2,3-butanetriol, I-116
 3-Indolylmethyl glucosinolate, I-122
 1-(1*H*-Indol-3-yl)-3-methyl-2-buten-1-one, I-125
 2-(1*H*-Indol-3-ylmethyl)-1*H*-indole-3-acetic acid, I-127
 3-(1*H*-Indol-3-yl)-2-oxopropanoic acid, I-133
 1-(1*H*-Indol-3-yl)-1,2,3-propanetriol, I-135
 3-(1*H*-Indol-3-yl)-2-propenoic acid, I-136
 3-(1*H*-Indol-3-yl)quinoline, I-138
 5-(1*H*-Indol-3-yl)-2(3*H*)-thiazolethione, I-139
 3-Isothiocyanato-4-methoxy-1*H*-indole, I-337
 Lycosinine A, L-351
 Malassezialactic acid, M-61
 Malassezin, M-65
 Malassezine, M-67
 Matemone, M-115
 Melochicorine, M-183
 Methoxybrassenin A, M-245
 Methoxybrassenin B, M-246
 5-Methoxy-1*H*-indole-4,7-dione, M-254
 6-Methoxy-1*H*-indole-4,7-dione, M-255
 5-(3-Methyl-2-butenyl)-1*H*-indole-2,3-dione, M-415
 6-(3-Methyl-2-butenyl)-1*H*-indole-2,3-dione, M-416
 3-(3-Methyl-2-butenyl)-1*H*-indole, M-411
 5-(3-Methyl-2-butenyl)-1*H*-indole, M-412
 6-(3-Methyl-2-butenyl)-1*H*-indole, M-413
 7-(3-Methyl-2-butenyl)-1*H*-indole, M-414
 2-(Methylthio)-1*H*-indole-3-carboxaldehyde, M-434
 3-Methyl-1*H*-indole-2-carboxylic acid, M-476
 5-Methyl-1*H*-indole-4,7-dione, M-477
 3-Methyl-1*H*-indole, M-474
 2-Methyl-1*H*-indole-4-methanol, M-478
 2-(Methylthio)-3*H*-indol-3-one, M-575
 Murrayacarine, M-756
 Murrayaculatine, M-758
 Neosidomycin, N-138
 3-Nitro-1*H*-indole, N-251
 Oxazanine 3, O-154
 3-(5-Oxazolyl)-1*H*-indole, O-157
 Penidiamide, P-189
 4-(5-Phenylpentyl)-1*H*-indole-2,3-dione, P-349
 Pityrialactone, P-490
 Pityriazole, P-495
 Polyandrocarpamide A, P-541
 6-Prenyl-1*H*-indole-3-carboxylic acid, P-617
 Preoxazin 7, P-619
 Streptindole, S-564
 Tanakine, T-25
 2,3,4,6-Tetrabromo-1*H*-indole, T-113
 2,3,4,7-Tetrabromo-1*H*-indole, T-114
 2,3,5,6-Tetrabromo-1*H*-indole, T-115
 2,3,4,7-Tetrachloro-1*H*-indole, T-119
 5,6,7,7*a*-Tetrahydro-1*H*-indol-2(3*H*)-one, T-195
 3-(2-Thiazolyl)-1*H*-indole, T-372
 2,4,7(3,4,7)-Tribromo-3(2)-chloro-1*H*-indole, T-459
 2,3,6-Tribromo-1*H*-indole, T-463
 2,3,7-Tribromo-1*H*-indole, T-464
 2,4,6-Tribromo-1*H*-indole, T-465
 3,4,6-Tribromo-1*H*-indole, T-466
 3,5,6-Tribromo-1*H*-indole, T-467
 3,5,7-Tribromo-1*H*-indole, T-468
 3,5,7-Tribromo-6-methoxy-1*H*-indole, T-471
 2,5,6-Tribromo-*N*-methylgramine, T-472
 2,4,6-Tribromo-3-methylthio-1*H*-indole, T-473
 2,3,4-Trichloro-1*H*-indole, T-480
 2,3,7-Trichloro-1*H*-indole, T-481
 3-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1*H*-indole, T-605
 Uvarindole A, U-59
 Uvarindole B, U-60
 Wuzhuyurutine B, W-27
 Zyzzin, Z-39

Simple biindoles

- Akashin C, A-225
 Annonidine A, A-1053
 Annonidine B, A-1054
 Annonidine C, A-1055
 Annonidine D, A-1056
 3,3'-Bi-1*H*-indole-5,5',6,6'-tetrol, B-130
 [3,3'-Biindoline]-2,2'-dione, B-131
 Bisindigotin, B-189
 Cephalinone C, C-257
 Cephalinone D, C-258
 Dendridine A, D-206
 5,5'-Dibromo-4,4'-dichloro-7,7'-dimethoxy-3,3'-bi-1*H*-indole, D-309
 6,6'-Dibromoindigotin, D-325
 2-(1,3-Dihydro-3-oxo-2*H*-indol-2-ylidene)-7-hydroxycyclopent[*g*]indol-3(2*H*)-one, D-486
 2,2',3,4',5,5'-Hexabromo-1,3'-bi-1*H*-indole, H-211
 Indigotin, I-72
 Indirubin, I-73
 2,4,4',6,6'-Pentabromo-2',3-bis(methylthio)-1,3'-bi-1*H*-indole, P-200
 Peronatin B, P-271
 Pityriarubin A, P-492
 Sciodole, S-150
 [3,2':2'(3'*H*),3'-Ter-1*H*-indol]-3'-one, T-87
 2,2',5,5'-Tetrabromo-3,3'-bi-1*H*-indole, T-108
 2,2',6,6'-Tetrabromo-3,3'-bi-1*H*-indole, T-109
 4,4',6,6'-Tetrabromo-2,2'-bis(methylthio)-3,3'-bi-1*H*-indole, T-110
 3,3',5,5'-Tetrabromo-7'-methoxy-1,4'-bi-1*H*-indole, T-117
 Tryptophan tryptophylquinone, T-641

Simple tryptamine alkaloids

- Alternatamide B, A-662
 Arundamine, A-1452
 Arundarine, A-1453
 Arundinine, A-1454
 Bengacarboline, B-58
 6-Bromotryptamine, B-356
 Buchtienine, B-381
 Bufopyramide, B-391
 Bufoserotonin C, B-392
 Bufotenine, B-393
 Ceridimine, C-281
 Chrysopentamine, C-429
 Cinchophyllamine, C-448
 Convolutindole A, C-630
 5,6-Dibromotryptamine, D-343
 3-Dimethylaminoacetyl-5-methoxyindole, D-727
N,N-Dimethyltryptamine, D-785
 Donasine, D-914
 Fragilamide, F-145
 Gelliusine A, G-37
 Gelliusine C, G-38
 Glypetelotine, G-130
 Hedyocapitelline, H-87
 4-Hydroxy-*N,N*-dimethyltryptamine, H-480
 5-Hydroxytryptamine, H-755
 Igzamide, I-21
N^b-(3-Indolylmethyl)-5-methoxy-*N*^b-methyltryptamine, I-128
N-[2-(1*H*-Indol-3-yl)-2-oxoethyl]acetamide, I-132
 Ipobscurine B, I-168
 Isostrychnopentamine B, I-326
 Janussine A, J-22

Kottamide A, K-92
 Kottamide D, K-93
 Kottamide E, K-94
 Lotthanongine, L-265
 4-Methyl *N*-[(2,3-dihydro-2-oxo-1*H*-indol-3-yl)acetyl]aspartate, M-299
N-Methyltryptamine, M-582
 Montamine, M-697
 Oxojanussine, O-186
 Prealmazole C, P-588
 Prebalamide, P-591
 Strychnofoline, S-595
 Strychnopentamine, S-601
 Strychnophylline, S-602
 Tryptamine, T-639
 Usambarensine, U-52

Cyclotryptamine alkaloids

Alline, A-626
 Angelicastigmine, A-1007
 Arundinine, A-1454
 6-Bromo-3*a*-(3,7-dimethyl-2,6-octadienyl)-1,2,3,3*a*,8,8*a*-hexahydro-1-methylpyrrolo[2,3-*b*]indol-7-ol, B-314
 Calycosidine, C-53
 Drimentine E, D-939
 Flustramine C, F-106
 Flustramine D, F-107
 Flustramine E, F-108
 Flustraminol A, F-109
 Flustraminol B, F-110
 Flustrarine B, F-111
 Hodgkinsine, H-311
 Idiospermuline, I-17
 Isoflustramine D, I-235
 Isopsychotridine A, I-288
 Isopsychotridine E, I-289
 Javanicinine A, J-29
 Madindoline A, M-34
 Phakellistatin 3, P-303
 Physostigmine, P-400
 Pseudophrynamine 512A, P-721
 Pseudophrynamine 272B, P-722
 Psycholeine, P-740
 Psychopentamine, P-741
 Psychotridine C, P-743
 Psychotridine, P-742
 Psychotrimine, P-744
 Quadrigemine A, Q-2
 Quadrigemine B, Q-3
 Quadrigemine H, Q-4
 Tetrahydroisocalycanthine, T-139
 2,3,8,8*a*-Tetrahydro-3*a*-nitro-1*H*-pyrrolo[2,3-*b*]indole, T-209
 Urochordamine A, U-47
 Vatamidine, V-41
 Vatamine, V-42
 Vatine, V-44

Chaetocin-like alkaloids

Amauromine, A-680
 Antibiotic Sch 52900, A-1247
 Antibiotic T 988B, A-1272
 Antibiotic WIN 64745, A-1310
 Antibiotic WIN 64821, A-1311
 Asperazine, A-1468
 Brevianamide E, B-290
 Brevicompanine A, B-293
 Brevicompanine C, B-294
 Chaetocin B, C-321
 Chaetocin C, C-322
 Chaetocin, C-320
 Chaetocochin A, C-323
 Chaetocochin C, C-324
 Chaetomin, C-335
 Chetracin A, C-372
 Dethiotetrakis(methylthio)chetomin, D-239
Aspergillus niger Diketopiperazine dimer, D-702
 Ditryptophenamine, D-874
 Drimentine A, D-936

Drimentine C, D-937
 Drimentine D, D-938
 Epicoccin B, E-90
 Epicoccin D, E-91
 Fructigenine A, F-159
 Fructigenine B, F-160
 Gliocladine A, G-97
 Gliocladine C, G-98
 Gliocladine C, G-99
 Gypsetin, G-234
 Isoechinulin A, I-230
 Leptosin A, L-124
 Leptosin B, L-125
 Leptosin C, L-126
 Leptosin D, L-127
 Leptosin E, L-128
 Leptosin F, L-129
 Leptosin G₁, L-130
 Leptosin G₂, L-131
 Leptosin G, L-132
 Leptosin H, L-133
 Leptosin I, L-134
 Leptosin K₁, L-135
 Leptosin K₂, L-136
 Leptosin K, L-137
 Leptosin M₁, L-138
 Leptosin M, L-139
 Leptosin N₁, L-140
 Leptosin O, L-141
 Leptosin Q, L-142
 Leptosin S, L-143
 Okaramine B, O-69
 Okaramine C, O-70
 Okaramine E, O-71
 Okaramine H, O-72
 Okaramine J, O-73
 Okaramine M, O-74
 Okaramine O, O-75
 Okaramine P, O-76
 Roquefortine C, R-125
 Roquefortine E, R-126
 Sporidesmin C, S-475
 Sporidesmin D, S-476
 Sporidesmin E, S-477
 Sporidesmin F, S-478
 Sporidesmin G, S-479
 Sporidesmin H, S-480
 Sporidesmin, S-474
 Variecolorin G, V-26
 Variecolorin J, V-27
 Variecolorin K, V-28
 Variecolorin L, V-29
 Verticillin A, V-89
 Verticillin C, V-90
 Verticillin D, V-91
 Verticillin G, V-92

Evodia alkaloids

Acetonylevodiamine, A-37
 Dehydroevodiamine, D-158
 Dihydrorutaecarpine, D-508
 Euxylophoricine A, E-305
 Euxylophoricine C, E-306
 Euxylophoricine D, E-307
 Euxylophoricine F, E-308
 Euxylophorine, E-309
 Hortiacine, H-361
 Paraensine, P-89
 Rutaecarpine, R-168
 Wuzhuyurutine A, W-26

Indolactam alkaloids

Antibiotic 69, A-1083
 Blastmycetin A, B-231
 Blastmycetin B, B-232
 Blastmycetin D, B-233
 Blastmycetin E, B-234
 Blastmycetin F, B-235
 Cytoblastin, C-941
 Indolactam I, I-77
 Indolactam V, I-78

Lyngbyatoxin A, L-367
 16-Methylpendolmycin, M-503
 Olivoretin C, O-88
 Olivoretin E, O-89
 Pendolmycin, P-180
 Teleocidin B₁, T-63

β -Carboline alkaloids

1-Acetyl- β -carboline, A-44
 1-Acetyl- β -carboline-3-carboxylic acid, A-45
 1-Acetyl-7-hydroxy- β -carboline, A-55
 1-Acetyl-8-hydroxy- β -carboline, A-56
 Alangiobussine, A-238
 Annomontine, A-1051
 Arborescidine C, A-1383
 Arenarine B, A-1402
 Arundarine, A-1453
 Bengacarboline, B-58
 9,9'-Bi-9*H*-pyrido[3,4-*b*]indole, B-146
 Brachycerine, B-264
 Brevicarine, B-291
 6-Bromo- β -carboline, B-299
 7-Bromo- β -carboline, B-300
 6-Bromo-1-ethyl- β -carboline, B-318
 5-Bromo-6-hydroxy- β -carboline, B-321
 7-Bromo-6-hydroxy- β -carboline, B-322
 6-Bromo-1-methyl- β -carboline, B-340
 Camptine, C-67
 Capitelline, C-95
 β -Carboline, C-130
 β -Carboline-1-carboxaldehyde, C-131
 β -Carboline-1-propanoic acid, C-134
 β -Carboline-1,3,4-trione, C-135
 1-(β -Carboline-1-yl)-3-hydroxy-1-propanone, C-136
 4-(β -Carboline-1-yl)-4-oxobutanoic acid, C-137
 1-(β -Carboline-1-yl)-3,4,5-trihydroxy-1-pentanone, C-138
 3-Carboxy-1,2,3,4-tetrahydro- β -carboline-1-propanoic acid, C-155
 1-Carboxytryptargine, C-156
 Ceciline, C-230
 Chrysotricine, C-433
 Cryptolepicarboline, C-786
 7,8-Dichloro-1-hydroxy-9-methyl- β -carboline, D-351
 7,8-Dichloro-9-methyl- β -carboline, D-355
 Dichotomie II, D-360
 Didemnoline A, D-380
 4,8-Dimethoxy- β -carboline, D-709
 4,7-Dimethoxy-1-vinyl- β -carboline, D-723
 4,9-Dimethoxy-1-vinyl- β -carboline, D-724
 Dragmacidonamine A, D-933
 Elaeocarpidine, E-54
 1-Ethyl- β -carboline, E-234
 1-Ethyl-4-methylsulfonyl- β -carboline, E-260
 Eudistalbin A, E-278
 Eudistalbin B, E-279
 Eudistomidin A, E-280
 Eudistomidin B, E-281
 Eudistomidin C, E-282
 Eudistomidin F, E-283
 Eudistomin A, E-285
 Eudistomin B, E-286
 Eudistomin C, E-287
 Eudistomin H, E-288
 Eudistomin T, E-289
 Flazine, F-85
 1-(2-Guanidinoethyl)-1,2,3,4-tetrahydro-6-hydroxymethyl- β -carboline, G-205
 Harmalidine, H-64
 Homofascaplysin A, H-333
 Homofluorodaturatin, H-337
 1-(4-Hydroxybenzoyl)- β -carboline, H-411
 4-Hydroxy- β -carboline-1-carboxaldehyde, H-456
 1-(2-Hydroxyethyl)- β -carboline, H-487
 1-(2-Hydroxyethyl)-4-methoxy- β -carboline, H-493
 1-(1-Hydroxy-2-methoxyethyl)-4-methoxy- β -carboline, H-560
 1-Hydroxymethyl- β -carboline, H-590

6-Hydroxy-1-methyl- β -carboline, H-591
 7-Hydroxy-1-methyl- β -carboline, H-592
 8-Hydroxy-1-methyl- β -carboline, H-593
 8-Hydroxy-1-vinyl- β -carboline, H-758
 Hyrtioerectine A, H-780
 Hyrtiomanzamine, H-781
 1-(3-Indolyl)- β -carboline, I-119
 Jafrine, J-14
 Kauluamine, K-22
 Kirondrine, K-49
 Komarine, K-58
 Komaroine, K-59
 Komavicine, K-60
 Komavine, K-61
 Kumujansine, K-106
 Lavendamycin, L-78
 Lissoclin C, L-199
 Maeganedin A, M-38
 Manzamine B, M-92
 Manzamine C, M-93
 Manzamine J, M-94
 Manzamine M, M-95
Saccharomyces kluyveri α -Mating pheromone 1, M-116
 Maxonine, M-131
 4-Methoxy-1-vinyl- β -carboline, M-290
 1-Methyl- β -carboline-3-carboxylic acid, M-427
 1-Methyl- β -carboline, M-426
 4-(1-Methyl-2-piperidinyl)- β -carboline, M-518
 1-(2-Methyl-1-propenyl)- β -carboline, M-521
 Milnamide A, M-619
 Myrianthosine, M-795
 Neokauluamine, N-120
 Pauridanthine, P-133
 Pedatisectine C, P-152
 Perlolyrine, P-270
 Picrasidine A, P-406
 Picrasidine B, P-407
 Picrasidine C, P-408
 Picrasidine E, P-409
 Picrasidine H, P-411
 Picrasidine K, P-412
 Picrasidine M, P-414
 Picrasidine R, P-415
 Picrasidine X, P-405
 Pityriacitrin B, P-489
 Pityriacitrin, P-488
 Plakortamine A, P-507
 Plakortamine C, P-508
 1-(2-Quinolonyl)- β -carboline, Q-51
 1-(4-Quinolonyl)- β -carboline, Q-52
 1-(5-Quinolonyl)- β -carboline, Q-53
 1-(6-Quinolonyl)- β -carboline, Q-54
 1-(8-Quinolonyl)- β -carboline, Q-55
 Reticulate, R-57
 Reticulato, R-58
 Roecarboline, R-115
 Secofascaplysin A, S-186
 Shishijimicin A, S-290
 Tetrahydroakagerine, T-141
 1,2,3,4-Tetrahydro-6-hydroxy- β -carboline, T-173
 1,2,3,4-Tetrahydro-2-methyl- β -carboline, T-203
 Tetrahydropentoxylone, T-215
 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indole-1,6-diol, T-225
 2,3,4,9-Tetrahydro-1*H*-pyrido[3,4-*b*]indol-1-one, T-226
 1-Vinyl- β -carboline-3-carboxylic acid, V-140
 1-Vinyl- β -carboline, V-139
 Vittacarboline, V-166
 Woodinine, W-21
 Xestoamine, X-20
 Xestomanzamine A, X-23

Carbazole alkaloids

Antibiotic BE 48021, A-1129
 Anisatine, A-1530
 Bicyclomahanimbicine, B-121
 Bicyclomahanimbine, B-122
 1,1'-Bi[2-hydroxy-3-methylcarbazole], B-128
 8,8'-Bikoengine, B-134

Bikoengine A, B-135
 Bisomahanine, B-192
 Bismahanine, B-195
 Bismurrayafoline A, B-207
 Bismurrayafoline B, B-208
 Bismurrayafoline C, B-209
 Bismurrayafoline E, B-210
 6,8'-Bismurrayamine A, B-211
 8,8'-Bismurrayamine A, B-212
 Bismurrayquinone A, B-213
 Carbazole, C-120
 9*H*-Carbazole-3-carboxaldehyde, C-121
 Carbazomadurin A, C-125
 Carbazomadurin B, C-126
 Carbazomarin A, C-127
 Carbazomycin G, C-128
 Carquinostatin A, C-169
 3-Chloro-9*H*-carbazole, C-397
 Chrestifoline A, C-423
 Chrestifoline B, C-424
 Chrestifoline C, C-425
 Clausamine F, C-499
 Clausaniline, C-500
 Clausenamine A, C-503
 Clausenapin, C-504
 Clausenquinone A, C-505
 Clausevatin E, C-506
 Clausine D, C-507
 Clausoline A, C-509
 Clausoline B, C-510
 Clausoline H, C-511
 Coproverdine, C-636
 Currayanine, C-814
 2,7-Dibromo-9*H*-carbazole, D-306
 3,6-Dibromo-9*H*-carbazole, D-307
 2,7-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-556
 2,8-Dihydroxy-9*H*-carbazole-3-carboxaldehyde, D-557
 Dihydroxygirinimbine, D-582
 3,6-Diiodo-9*H*-carbazole, D-696
 3,6-Dimethyl-1-(3-methyl-2-butenyl)-9*H*-carbazole, D-753
 Epocarbazolin A, E-103
 Epocarbazolin B, E-104
 Euchrestifoline, E-274
 Euchrestine A, E-275
 Euchrestine C, E-276
 Euchrestine D, E-277
 Eustifoline A, E-300
 Eustifoline B, E-301
 Eustifoline C, E-302
 Furoclausine A, F-203
 Furoclausine B, F-204
 Girinimbine, G-81
 Glybomine C, G-110
 Glycomaurrol, G-125
 Heptaphylline, H-125
 1-Heptyl-3-hydroxy-2-methyl-9*H*-carbazole, H-132
 1-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-447
 2-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-448
 7-Hydroxy-9*H*-carbazole-3-carboxaldehyde, H-450
 4-Hydroxy-3-methoxy-2-methyl-9*H*-carbazole-1-carboxaldehyde, H-561
 6-Hydroxy-5-(3-methyl-2-butenyl)-9*H*-carbazole-3-carboxaldehyde, H-577
 1-Hydroxy-3-methyl-9*H*-carbazole, H-583
 2-Hydroxy-3-methyl-9*H*-carbazole, H-585
 2-Hydroxy-7-methyl-9*H*-carbazole, H-586
 3-Hydroxy-6-methyl-9*H*-carbazole, H-587
 3-Hydroxymethyl-1-methoxycarbazole, H-603
 6-Hydroxymethyl-1-methoxycarbazole, H-604
 Indizoline, I-75
 Isomurrayazoline, I-250
 Koeniginequinone A, K-55
 Kwangsine, K-117
 3-Methoxy-2-methyl-1-phenyl-carbazole, M-270
 3-Methyl-1*H*-carbazole-1,4(9*H*)-dione, M-425

1-Methyl-9*H*-carbazole, M-423
 3-Methyl-9*H*-carbazole, M-424
 4-Methyl-10*H*-furo[3,2-*a*]carbazole, M-464
 9-Methyl-6*H*-furo[2,3-*c*]carbazole, M-465
 Mukoenine A, M-728
 Mukoenine B, M-729
 Murrayafoline E, M-745
 Murrayafoline F, M-746
 Murranimbine, M-749
 Murrastifoline A, M-751
 Murrastifoline B, M-752
 Murrastifoline C, M-753
 Murrastifoline D, M-754
 Murrastifoline F, M-755
 Murrayacine, M-757
 Murrayafoline B, M-759
 Murrayaline C, M-760
 Murrayamine F, M-761
 Murrayamine J, M-762
 Murrayamine M, M-763
 Murrayamine O, M-764
 Murrayanine, M-765
 Murrayquinone B, M-766
 Murrayquinone D, M-767
 Neocarazostatin A, N-103
 3,3'-[Oxybis(methylene)]bis[9-methoxy-9*H*-carbazole], O-219
 Oxidimurrayafoline, O-221
 Picraniline, P-403
 Pityriazole, P-495
 Pyrayafoline A, P-883
 Pyrayafoline B, P-884
 Pyrayafoline D, P-885
 Pyrayaquinone A, P-886
 Pyrayaquinone B, P-887
 Pyrayaquinone C, P-888
 Siamenol, S-291
 Streptovercillin, S-573
 1,2,8-Trimethoxy-6-methyl-9*H*-carbazole, T-596

Indolo[2,3-*a*]carbazole and related alkaloids

Antibiotic BE 54017, A-1131
 Antibiotic K 252b, A-1189
 Antibiotic K 252c, A-1190
 Antibiotic TAN 1030A, A-1281
 Antibiotic ZHD 0501, A-1324
 Arcyriacyanin A, A-1392
 Arcyriaflavin A, A-1393
 Arcyriaeverdin C, A-1395
 Arcyrin A, A-1396
 Arcyroxindole A, A-1397
 Arcyroxocin A, A-1398
 3-Chloro-11,12-dihydroindolo[2,3-*a*]carbazole, C-401
 5-Cyano-6-methoxyindolo[2,3-*a*]carbazole, C-828
 3,8-Dichloro-11,12-dihydroindolo[2,3-*a*]carbazole, D-350
 11,12-Dihydroindolo[2,3-*a*]carbazole, D-464
 Indocarbazostatin, I-76
 Rebeccamycin, R-32
 Staurosporine, S-498
 Tjipanazole J, T-419

Indolonaphthyrindine alkaloids

Bonafousine, B-245
 Canthin-6-one, C-90
 Cimicidiphytine, C-440
 Cimiciphytine, C-441
 Cimilophytine, C-442
 Deformyltalbotinic acid methyl ester, D-145
 Haplocidiphytine, H-54
 Haplophytine, H-58
 Henrycinol A, H-114
 4-Hydroxycanthin-6-one, H-440
 5-Hydroxycanthin-6-one, H-442
 8-Hydroxycanthin-6-one, H-443
 9-Hydroxycanthin-6-one, H-444
 10-Hydroxycanthin-6-one, H-445

11-Hydroxycanthin-6-one, H-446
 1-Hydroxycanthin-6-one, H-438
 2-Hydroxycanthin-6-one, H-439
 5-Hydroxymethylcanth-6-one, H-582
 3*H*-Indolo[3,2,1-*de*][1,5]naphthyridine-2,6-dione, I-105
 Isobonafousine, I-197
 Luotonine C, L-299
 Luotonine D, L-300
 4-(Methylthio)canth-6-one, M-571
 5-(Methylthio)canth-6-one, M-572
 9-(Methylthio)canth-6-one, M-573
 11-(Methylthio)canth-6-one, M-574
 Norisotuboflavine, N-302
 Picrasidine L, P-413
 Picrasidine M, P-414
 Strychnohirsutine, S-600
 Talbotine, T-19
 Tuboflavine, T-671

Ergot alkaloids

Securidaca longipedunculata Alkaloid A, A-375
 1,1-Bis(8,9-epoxy-6,8-dimethylergoline), B-176
 Cividiclavine, C-487
 Clavicipitic acid I, C-514
 Cycloclavine, C-856
 β,β-Ergoannam, E-147
 Ergobalansine, E-148
 Ergobine, E-149
 Ergobutine, E-150
 Ergobutyryne, E-151
 Ergocornam, E-152
 Ergocornine, E-153
 Ergocristam, E-154
 Ergocristine, E-155
 Ergocryptam, E-156
 α-Ergocryptine, E-157
 β-Ergocryptine, E-158
 γ-Ergocryptinine, E-159
 Ergogaline, E-160
 Ergoheptine, E-161
 Ergohexine, E-162
 Ergoladinine, E-163
 Ergometriner, E-164
 Ergonine, E-165
 β-Ergoptine, E-166
 Ergoptine, E-167
 Ergosecalinine, E-168
 Ergosine, E-169
 Ergosine, E-170
 Ergostine, E-171
 Ergotamine, E-172
 Ergovaline, E-174
 Festuclavine, F-49
 Fumigaclavine C, F-182
 Lysergol, L-371
 Molliclavine, M-671
 Paspaclavine, P-106
 Paspalic acid, P-107

Indoloterpenoid alkaloids

Ambiguine B isonitrile, A-684
 Ambiguine D isonitrile, A-685
 Ambiguine E isonitrile, A-686
 Ambiguine F isonitrile, A-687
 Ambiguine G nitrile, A-688
 Anhydrohapaloxindole A, A-1026
 Aristoaristone, A-1425
 Aristofrucosine, A-1426
 Aristolarine, A-1430
 Aristolasene, A-1431
 Aristolasicone, A-1432
 Aristolasol, A-1433
 Aristone, A-1438
 Aristoserratenine, A-1441
 Bisaristone A, B-156
 Biscodehydrocyclopiazonic acid, B-218
 Borrecapine, B-251
 Borreline, B-252

Borreverine, B-253
 Cyclopiazonic acid imine, C-895
 α-Cyclopiazonic acid, C-894
 Fontonamide, F-126
 Fruticosonine, F-164
 Hapalindole A, H-48
 Hapalindole E, H-49
 Hapalindole T, H-50
 Hapalindolinone A, H-51
 Hapalonamide G, H-52
 Isoborreverine, I-198
 Isopolyalthenol, I-269
 Monodorindole, M-692
 Neopolyalthenol, N-135
 Peduncularine, P-162
 Polyalthenol, P-540
 Polyavolinamide, P-546
 Polyveoline, P-564
 Serratoline, S-259
 Sorelline, S-383
 Spermacoceine, S-395
 Suaveolindole, S-617
 Tasmanine, T-35
 Thiersindole A, T-375
 Thiersindole C, T-376
 2,2,6-Trimethyl-4-(4-quinolinyl)-3-azabicyclo[3.3.1]non-6-ene, T-614
 Welwitindolinone A isonitrile, W-12
 Welwitindolinone B isothiocyanate, W-13
 Yuehchukene, Y-19

Strictosidine alkaloids

Alstonidine, A-650
 Angustidine, A-1012
 Angustine, A-1019
 Bahienoside A, B-6
 5-Carboxystrictosidine, C-153
 Cordifoline, C-641
 Lyadine, L-313
 Lyalidine, L-314
 Lyaline, L-315
 Lyaloside, L-316
 Macrolidine, M-16
 Myrianthosine, M-795
 Nauclealine A, N-57
 Naucleorine, N-62
 Nauclefine, N-70
 Naulafine, N-78
 Neonaucleoside A, N-127
 Neonaucleoside B, N-128
 Neonaucleoside C, N-129
 Paratunamide D, P-92
 Pauridianthine, P-133
 Pauridianthoside, P-135
 Strictosidinic acid, S-583
 Strictosidinic ketone, S-584
 Vincosamide, V-126

Camptothecine-like alkaloids

22-Hydroxyacuminatine, H-403
 OPHR 17, O-99
 Pumiloid, P-796

Corynanthean alkaloids

Adirubine, A-141
 Afrocurarine, A-174
 C-Alkaloid O, A-572
 Alkaloid RMB 10, A-594
 Alkaloid Us 7, A-603
 Angusteine, A-1010
 Arboricine, A-1385
 Arboricinine, A-1386
 Blumeanine, B-238
 Bracteatine, B-274
 Buchtienine, B-381
 Burnamicine, B-419
 Callophylline B, C-38
 Chrysopentamine, C-429
 Cinchophyllamine, C-448
 Corynantheidinaline, C-687

Corynantheidinalinic acid, C-688
 Corynantheine, C-689
 16,17-Didehydro-17-hydroxy-16-(hydroxymethyl)corynan-5-carboxylic acid, D-374
 Dihydrocorynantheine pseudoindoxyl, D-411
 Dihydrocorynantheol, D-412
 Flavocarpine, F-82
 Flavopereirine, F-84
 Gambirine, G-13
 Geissoschizine, G-32
 Geissoschizol, G-33
 Geissospermine, G-35
 Guettardine, G-218
 Guiaflavine, G-219
 Guianensine, G-220
 Hunteracine, H-368
 7-Hydroxy-3-oxo-3,7-secorhynchophylline, H-656
 Isositsirikine, I-319
 Isostrychnopentamine B, I-326
 Lombine, L-239
 Longicaudatine Z, L-250
 Longicaudatine, L-249
 Mitragnine, M-657
 Mitractalonal, M-659
 Mitralsulgnine, M-663
 Nitraraidine, N-242
 Ochromianoxine, O-22
 Ochroprosinine oxindole, O-23
 Ophiorrhizine-12-carboxylate, O-97
 Ophiorrhizine, O-96
 Rhyncophylline, R-90
 Rotundifoline, R-134
 Rumberine, R-159
 Salacin, S-18
 Serpentinine, S-253
 Sitsirikine, S-325
 Strychnofoline, S-595
 Strychnofuranine, S-596
 Strychnopentamine, S-601
 Strychnophylline, S-602
 Strychnoxanthine, S-606
 Subincanadine A, S-623
 3,4,5,6-Tetrahydrodihydrocorynantheol, T-136
 Turbinatine, T-679
 Uncaramine, U-22
 Usambarensine, U-52

Secologanin-derived sesquimer alkaloids

Alkaloid C2, A-457
 Alkaloid D1, A-493
 Alkaloid D2, A-494
 Bahienoside A, B-6
 Bonafousine, B-245
 Buchtienine, B-381
 Cerdimine, C-281
 Chrysopentamine, C-429
 Cimiciduphytine, C-440
 Cimiciphytine, C-441
 Cinchophyllamine, C-448
 Haplocidiphytine, H-54
 Haplophytine, H-58
 Isobonafousine, I-197
 Isostrychnopentamine B, I-326
 Janussine A, J-22
 Oxojanussine, O-186
 Strychnobaridine, S-590
 Strychnofoline, S-595
 Strychnopentamine, S-601
 Strychnophylline, S-602
 Usambarensine, U-52

Akageran and related alkaloids

Adifoline, A-140
 Akagerine lactone, A-223
 Akagerine, A-222
 Camptoneurine, C-69

Correantine A, C-659
 Correantine B, C-660
 Correantine C, C-661
 Correantoxide, C-662
 Deoxycordifoline lactam, D-223
 Dihydrocycloakagerine, D-414
 Janussine A, J-22
 Kribine, K-101
 Neoadifoline, N-98
 Oxojanussine, O-186
 Tetrahydroakagerine, T-141

Ajmalicine-like alkaloids

Ajmalicine hydroxyindolenine, A-213
 Ajmalicinial, A-214
 Ajmalicine, A-215
 Alstonine, A-652
 Amsosinine, A-953
 Aricine, A-1423
 Carapanaubine, C-117
 5-Carboxytetrahydroalstonine, C-154
 Cathenamine, C-207
 Herbaine, H-137
 Isomitraphyllinol, I-247
 Javaniside, J-31
 Javaphylline, J-32
 Magniflorine, M-43
 Majdine, M-50
 Methyl 16,17-didehydromethoxy-19-methyloxayohimban-16-carboxylate, M-298
 Mitrajavine, M-658
 Mitraphylline, M-661
 Naucleosidine, N-64
 Nauclefiline, N-69
 Nauclefolinine, N-73
 Naucleidinal, N-74
 Picraphylline, P-404
 Rauvolcinine, R-28
 Reserpiline, R-51
 Reserpinine, R-53
 Serpentinine, S-253
 Tetraphylline pseudoindoxyl, T-289

Macroline alkaloids

Alstocraline, A-641
 Alstolactone, A-642
 Alstomacrine, A-643
 Alstomacrine, A-644
 Alstomacrophylline, A-645
 Alstomicine, A-647
 Alstonal, A-648
 Alstonerine, A-649
 Alstonisidine, A-653
 Alstonisine, A-654
 Alstophyllal, A-656
 Angusticraline, A-1011
 Angustimalal, A-1017
 Angustimaline, A-1018
 Jadiffine, J-7
 Macralstonidine, M-7
 Macralstonine, M-8
 Macrocaffrine, M-10
 Macrocarpamine, M-11
 Macrodasine A, M-14
 Macrogentine, M-15
 Macrophylline, M-20
 Macropegatrine, M-30
 Macroxine, M-31
 Pandicine, P-66
 Perhentinine, P-256
 Raumacline, R-22
 Sellowiine, S-220
 Suaveoline, S-618
 Talcarpine, T-20
 Villalstonine, V-103

Malindan alkaloids

3,14-Didehydro-19-methylnormalindine, D-376
 Isodihydrocadambine, I-226

Isomalindine-16-carboxylate, I-243
 Malindine, M-73
 Nauclechine, N-63

Cadamban alkaloids

Alkaloid ND 370, A-571
 Cadambine, C-5
 Dihydrocadambine, D-406
 Nauclefoline, N-71
 Rubenine, R-141

Vallesiachotaman alkaloids

Antirrhine lactone, A-1329
 Antirrhine, A-1328
 Camptacumotine, C-66
 Glabratine, G-82
 Matadine, M-114
 Naucleamide A, N-59
 Naucleofficine C, N-75
 Ophiorine A, O-95
 Vallesiachotamine lactone, V-11

Gelsemium alkaloids

Elegansamine, E-63
 Gelsamydine, G-41
 Gelsebanine, G-42
 Gelsedilam, G-43
 Gelsedine, G-44
 Gelsefuranidine, G-45
 Gelseiridone, G-46
 Gelselegine, G-47
 Gelsemamide, G-48
 Gelsemine, G-49
 Gelsemoxonine, G-50
 Humantenirine, H-367
 Hydrakouminol, H-393
 Ourouparine, O-140
 Rankiniridine, R-15

Yohimbinoïd alkaloids

Alstonilidine, A-651
 Cadamine, C-6
 Callophylline B, C-38
 Canembine, C-81
 Deserpideine, D-233
 Deserpidine, D-234
 19,20-Didehydroyohimbine, D-378
 10-Methoxyyohimbine, M-291
 11-Methoxyyohimbine, M-292
 Naucleactonine A, N-55
 Naucleactonine B, N-56
 Naucleamide C, N-60
 Naucleamide E, N-61
 Naucleficine, N-67
 Neoogyambirtannine, N-132
 Nitrraine, N-243
 Ourouparine, O-140
 Rescinnaminol, R-48
 Reserpenediol, R-49
 Reserpic acid, R-50
 Sempervirine, S-228
 Seredine, S-251
 3,4,5,6-Tetrahydroyohimbine, T-140
 Uncaramine, U-22
 Venenatine, V-50
 Yohambinine, Y-12
 β -Yohimbine oxindole, Y-14
 β -Yohimbine pseudoindoxyl, Y-15
 Yohimbine, Y-13
 Yohimbol, Y-16

Sarpagine alkaloids

Accedine, A-27
 Accedinisine, A-28
 Affinisine oxindole, A-167
 Akuammidine, A-229
 Alstocraline, A-641

Alstonoxine B, A-655
 Alstoumerine, A-658
 3-Amino- N^B -methylsecovocarpine, A-846
 Arbophylline, A-1382
 16-Decarbomethoxy-19,20-dihydrovobasine, D-119
 Dehydrovochalotone, D-173
 Difforine, D-396
 Dihydroperaksine, D-491
 Dispegatrine, D-864
 Eburnaphylline, E-7
 6,17-Epoxy-19,20-dihydro-1-methylsarpagan-17,21-diol, E-130
 Ervitsine, E-193
 Gardfloramine, G-18
 Gardmultine, G-19
 Gardneramine, G-20
 Gardnerine, G-22
 Gardnutine, G-23
 Gardquinolone, G-24
 Geissolosimine, G-31
 Macralstonidine, M-7
 Macrosallhine, M-29
 Macropegatrine, M-30
 12-Methoxyaffinisine, M-237
 Peraksine, P-241
 Polyneuridine, P-562
 Sarpagine, S-89
 Talpinine, T-21
 Tombozine, T-425
 Trinervine, T-616
 Undulatine, U-32
 Voacarpine, V-171
 Vochalotone oxindole, V-172
 Voacoline, V-174

Ajmaline alkaloids

Ajmaline, A-216
 Ajmalinimine, A-217
 Alstomacrine, A-644
 Alstonisidine, A-653
 Flexicorine, F-88
 Herbamine, H-139
 Isorhazicine, I-309
 10-Methoxy-11-[10-(11-methoxyvincamajinyl)]-cathafoline, M-258
 11-[10-(11-Methoxyvincamajinyl)]vincorine, M-289
 Perakine, P-240
 Raureflexine, R-24
 Sandwicolidine, S-51
 Sandwicoline, S-52
 Tetraphyllicine, T-288
 Vincamajine, V-113
 Vincawajine, V-121
 Vomilenine, V-194

Vobasine alkaloids

Accedinisine, A-28
 Affinine, A-166
 Alkaloid TC-D, A-600
 Capuvosidine, C-105
 Capuvosine, C-106
 Ceridimine, C-281
 Conodiparine A, C-611
 Conodiparine B, C-612
 Conodirine A, C-613
 Conodirine B, C-614
 Conoduramine, C-615
 Conodurine, C-616
 Conodutarine A, C-617
 Coryzylamine, C-703
 16'-Decarbomethoxy-19,20-dihydroconoduramine, D-118
 16-Decarbomethoxyvoacamine pseudoindoxyl, D-121
 Deformylcoryzylamine, D-143
 Difforlemenine, D-397
 Difforlemenitine, D-398
 Dregamine, D-935
 Erichsonine, E-176

Ervahainamidine A, E-187
 Ervahainamidine B, E-188
 Ervahanine A, E-189
 Hazuntamine, H-79
 Monogagine, M-693
 Pagicerine, P-10
 Pagisulfine, P-11
 Perimivine, P-260
 Perivine, P-268
 Pseudovobparicine, P-734
 Rauviridine, R-27
 Rauwolfinine, R-29
 Tabernaelegantinine A, T-3
 Tabernaelegantinine B, T-4
 Tabernaelegantinine C, T-5
 Tabernaemontabovine, T-6
 Tabernaemontavine, T-7
 Tabernamine, T-8
 Taberpsychine, T-10
 2,7,19,20-Tetrahydro-3-hydroxy-1-methylvobas-
 san-17-oic acid δ -lactone, T-186
 Voacamidine, V-167
 Voacamine, V-168
 Vobasenal, V-183
 Vobasonidine, V-184
 Vobatricine, V-185
 Vobparicine, V-186

Koumine alkaloids

Koumine, K-96

Ervatamia alkaloids

Ervatamine, E-190
 Flabelliformide A, F-72
 Methuenine, M-293

Akuammiline alkaloids

Abereamine 1, A-8
 Abereamine 2, A-9
 Abereamine 3, A-10
 Abereamine 4, A-11
 Akuammiginone, A-230
 Akuammiline, A-231
 Akuamine, A-232
 Alschomine, A-640
 Alstomaline, A-646
 Angusticaline, A-1011
 Aspidodasycarpine, A-1492
 Aspidophylline A, A-1495
 Burnamine, B-420
 Cabufiline, C-3
 Cathafoline, C-191
 Ceylanicine, C-314
 Ceylanine, C-315
 Coryzeylamine, C-703
 O-Deacetyl-1,2,4,5-tetrahydro-2-hydroxy-1-
 methylakuammilinium(1+), D-94
 Deformylcoryzeylamine, D-143
 Desformocorymine, D-236
 Eripinal, E-181
 Eripine, E-182
 Flexicorine, F-88
 Gentiacriline, G-53
 Isocorymine, I-209
 Lahoricine, L-16
 Lanceomigine, L-28
 Lanciferine, L-29
 10-Methoxy-11-[10-(11-methoxyvincamajinyl)]-
 cathafoline, M-258
 11-[10-(11-Methoxyvincamajinyl)]vincorine,
 M-289
 Nareline, N-47
 Peceylanine, P-147
 Peceyline, P-148
 Pelankine, P-172
 Picrinine, P-417
 Pleiocorine, P-519
 Plumocraline, P-527
 Raucanine, R-20
 Raucubaine, R-21

Rausutranine, R-25
 Rausutrine, R-26
 Rhazimine, R-73
 Rhazinaline, R-75
 3,4-Seco-3,14-dehydrocabucraline, S-182
 Strictamine, S-576
 Strictaminolamine, S-577
 Tacraline, T-15
 Umbellamine, U-21
 Undulatine, U-32
 Vincarubine, V-118
 Vincorine, V-125
 Vingramine, V-133

Pleiocarpaman alkaloids

C-Alkaloid Y, A-617
 Fluorocarpamine, F-99
 C-Fluorocurine, F-100
 Macrocarpamine, M-11
 C-Mavacurine, M-130
 Pleiocarpamine, P-518
 Pleiocorine, P-519
 Pleiocraline, P-520
 Plumocraline, P-527
 Pycnanthine, P-827
 Pycnanthine, P-828
 Strictine, S-582
 Villalstonine, V-103

Cinchona alkaloids

Chairamidine, C-337
 Chairamine, C-338
 Cinchonamine, C-444
 Cinchonaminone, C-445
 Cinchonidine, C-446
 Cinchonine, C-447
 Conchairamidine, C-597
 Conchairamine, C-598
 Cuscamidine, C-817
 Cusconidine, C-819
 Cusconine, C-820
 Homocinchonidine, H-328
 Javanine, J-30
 Paricine, P-95
 Quinamine, Q-12
 Quinicine, Q-17
 Quinidine, Q-18
 Quinine, Q-20
 Remijinine, R-40

Akuammicine alkaloids

Afrocuarine, A-174
 C-Alkaloid D, A-476
 Alvimine, A-667
 Alviminine, A-668
 Bharhingine, B-112
 Calebassine, C-31
 Calebassine 1, C-32
 Caracurine II, C-111
 Caracurine V, C-110
 Caracurine VII, C-115
 Compactinervine, C-589
 C-Curarine, C-809
 3,17-Dihydroxyzenkerine, D-688
 Divarine, D-875
 Dolichocurine, D-910
 Echitamidine, E-30
 Geissolosimine, G-31
 Geissoschizoline, G-34
 Geissospermine, G-35
 Guiaflavine, G-219
 Guianensine, G-220
 Henningsamide, H-110
 Henningsine, H-111
 Longicaudatine Z, L-250
 Longicaudatine, L-249
 Malagashanine, M-58
 Matopense, M-120
 Mattogrossine, M-122
 11-Methoxyneooxydiaboline, M-278

Mossambine, M-713
 Myrtoidine, M-803
 Neolaugerine, N-122
 Panganensine R, P-69
 Panganensine X, P-68
 Pericidine, P-257
 Preakuammicine, P-587
 Retuline, R-63
 Rhazizine, R-77
 Rosibiline, R-131
 Spermostrychnine, S-400
 Stemmadenine, S-510
 Stricticine, S-580
 Strychnobiline, S-591
 Strychnochrysin, S-593
 Strychnofendlerine, S-594
 Strychnohexamine, S-599
 Strychnopivotine, S-603
 Strychnosilidine, S-604
 Strychnosplendine, S-605
 Strychnozairine, S-607
 Subincanadine D, S-624
 Subincanadine F, S-625
 Toxiferine I, T-436
 Tsilanine, T-649
 Tubifolidine, T-668
 Valparicine, V-16
 Vinoxine, V-136
 Voastrictine, V-182
 Vobatricine, V-185
 Yunnanensine, Y-21
 Zenkerene, Z-14

Strychnidine alkaloids

Leuconotis Alkaloid 376, A-308
 C-Alkaloid D, A-476
 Cyclostrychnine, C-911
 Holstiine, H-323
 Icajine, I-12
 Minfiensine, M-628
 Protostrychnine, P-672
 Pseudostrychnine, P-726
 Strychnine, S-589
 Strychnobiline, S-591
 Strychnogucine A, S-597
 Strychnogucine B, S-598
 Strychnohexamine, S-599
 Sungucine, S-642

Condylocarpan alkaloids

Arboflorine, A-1380
 Aspidospermatidine, A-1497
 4-(Chloromethyl)-2,16-didehydro-16-(methoxy-
 carbonyl)condyfolanium, C-413
 Dichotine, D-359
 Geissovelline, G-36
 Goniomine, G-141
 Precondylocarpine, P-595
 Strychnochromine, S-592
 Tronoharine, T-631
 Tubotaivinal, T-672
 Tubotaivine, T-673

Secodine alkaloids

Presecamine, P-624
 Secamine, S-174
 Secodine, S-184

Aspidosperma alkaloids

Alalakine, A-233
 Amataine, A-676
 Anomaline, A-1067
 Arboloscine, A-1381
 Aspidoalbidine, A-1490
 Aspidoispermine, A-1493
 Aspidophytine, A-1496
 Aspidospermine, A-1499
 Bannucine, B-15
 Beninine, B-63

10,10'-Bi[*N*-acetyl-11,12-dihydroxyaspidospermidine], B-115
 Buxomelina, B-475
 Callichiline, C-35
 Cathaphylline, C-193
 Catharanthamine, C-194
 Catharine, C-197
 Cathovaline, C-213
 Cimicidiphytine, C-440
 Cimiciphytine, C-441
 Cimilophytine, C-442
 Conofoline, C-618
 Conophyllidine, C-621
 Criophylline, C-751
 Dichomine, D-358
 Eburine, E-4
 Ervafoleone, E-185
 Ervafoleone, E-186
 Ervafoleone, E-186
 1*b*-Ethyl-1*a*,1*b*,3,4,8*c*,9,10,10*a*-octahydro-2*H*-oxireno[4,5-*b*]pyrido[3,2-*c*]carbazole, E-262
 Folicangine, F-117
 Goniomitine, G-142
 Haplocidiphytine, H-54
 Haplophytine, H-58
 Hazuntiphyllidine, H-80
 Hazuntiphylline, H-81
 Insulopinine, I-151
 Kisantine, K-51
 Leuconoxine, L-156
 Leurosine, L-162
 Leurosine, L-163
 Limaspermidine, L-177
 Melomorsine, M-187
 Mercarpine, M-210
 Methyl 1,2-didehydro-6,7-epoxy-3-hydroxy-8-oxoaspidospermidine-3-carboxylate, M-297
 Methylenebismehranine, M-437
 Modestanine, M-666
 Obscurinervidine, O-7
 Obscurinervine, O-8
 3-Oxotabersonine, O-210
 Pandicine, P-66
 Paucivenine, P-132
 Peduncularidine, P-161
 Petchicine, P-278
 Polyervinine, P-560
 Pseudovincadifformine, P-732
 Quimbeline, Q-10
 Roseadine, R-127
 Rosicine, R-132
 Scandomelidine, S-119
 Scandomelina, S-120
 Scandomelonine, S-121
 14,15-Seco-3-oxokopsinal, S-193
 Strepmpeliopidine, S-558
 Strictanine, S-578
 Suaveolenine, S-616
 Tabernaemontabovine, T-6
 Tabernaemontavine, T-7
 Tabersonine, T-11
 Tenuicausine, T-77
 Tetrastachyne, T-294
 Tetrastachynine, T-295
 Trichophylline, T-492
 Vinamidine, V-107
 Vinblastine, V-110
 Vincarubine, V-118
 Vincathicine, V-119
 Vincoline, V-124
 Vincovalicine, V-127
 Vincristine, V-128
 Vindolicine, V-129
 Voacicol, V-173
 Voafolidine, V-175
 Voafrine A, V-176
 Vobtusamine, V-187
 Vobtusine, V-188

Rhazinilam alkaloids

Leuconolam, L-155
 Rhazinal, R-74

Melonine alkaloids

Celastromelidina, C-232
 Celastromelina, C-233
 Melonine, M-188
N^b,*N*^b-Methylenebismeloninium(2+), M-438

Quebrachamine alkaloids

Aspidochibine, A-1491
 Capuronine, C-104
 Ervinidine, E-192
 Hyderabadine, H-391
 Melaxillaridine, M-161
 Pseudovincadifformine, P-732
 Rhazidigenine, R-72
 Stapfinine, S-493
 Vinamidine, V-107
 Voaharine, V-177
 Voaphylline hydroxyindolenine, V-180

Aspidofractine alkaloids

Aspidofractinina, A-1494
 16-Hydroxyaspidofractinina, H-406
 Kopsaporine, K-66
 Kopsidasine, K-67
 Kopsifoline C, K-70
 Kopsijasmine, K-71
 Kopsijaminilam, K-72
 Kopsinine, K-78
 Kopsinone, K-80
 Lahadinine A, L-13
 Lahadinine B, L-14
 Lapidilectine A, L-43
 Lapidilectine B, L-44
 Lundurine A, L-294
 Nitaphylline, N-238
 Paucidactine A, P-128
 Paucidactine C, P-129
 Pauciflorine A, P-130
 Pauciflorine B, P-131
 Pleiomutina, P-521
 Pycnanthine, P-827
 Pycnanthinine, P-828
 Scandomelidina, S-119
 Singapurenine A, S-309
 Tenuiphylline, T-80
 Venacarpine A, V-47
 Venacarpine B, V-48
 Venalstonine, V-49

Vindolinine alkaloids

Tuboxenine, T-674
 Vindolinine, V-132

Kopsane alkaloids

Danuphylline, D-17
 Dasyrachine, D-86
 10,22-Dioxokopsane, D-805
 Fruticosine, F-163
 Kopsanol, K-65
 Kopsidasine, K-68
 Kopsifine, K-69
 Kopsinarine, K-75
 Kopsine, K-76
 Kopsinidine A, K-77
 Kopsinitarine A, K-79
 Mersingine A, M-216
 Methyl chanofruticosinate, M-296

Melodinus alkaloids

Meloscandonine, M-192
 Meloscine, M-193
 Mersilongine, M-213
 Mersiloscine, M-214
 Mersinaline, M-215
 Mersinine A, M-217
 Mersirachine, M-218
 Scandine, S-118

Scandomelina, S-120
 Scandomelonine, S-121

Iboga alkaloids

Alioline, A-264
 12,12'-Bi[11-hydroxycoronaridine], B-127
 Bonafousine, B-245
 Capuvosine, C-106
 Catharanthine, C-195
 Conodiparine A, C-611
 Conodiparine B, C-612
 Conodiridine A, C-613
 Conodiridine B, C-614
 Conoduramine, C-615
 Conodurine, C-616
 Conodutarine A, C-617
 Conopharyngine, C-620
 Coronaridine hydroxyindolenine, C-658
 Coronaridine, C-657
 Crassanine, C-726
 16'-Decarbomethoxy-19,20-dihydroconoduramine, D-118
 Ervahainamidina A, E-187
 Ervahainamidina B, E-188
 Ervahainine A, E-189
 Heyneatine, H-270
 3-(1-Hydroxyethyl)coronaridine, H-489
 3-Hydroxy-3,4-seccoronaridine, H-732
 Ibogamine, I-4
 Iboluteine, I-5
 Iboquine, I-7
 Iboxygaine hydroxyindolenine, I-9
 Isobonafousine, I-197
 Isovoacangine, I-344
 Kisantine, K-50
 Obovamine, O-3
 3,6-Oxidoiboxygaine, O-162
 3-(2-Oxopropyl)coronaridine, O-199
 3,3'-(Oxopropyl)dicoronaridine, O-201
 Roseadine, R-127
 Sarcopharyngine, S-79
 Tabernaelegantinine A, T-3
 Tabernaelegantinine B, T-4
 Tabernaelegantinine C, T-5
 Tabernamine, T-8
 Tabernoxidine, T-9
 Tetrastachyne, T-294
 Tetrastachynine, T-295
 Voacamidine, V-167
 Voacamine, V-168
 Voacangine, V-170
 Voaluteine, V-178

Pandoline alkaloids

Capuvosidine, C-105
 Catharanthamine, C-194
 Catharine, C-197
 Deethylbophyllidine, D-140
 1,2-Didehydropseudoaspidospermidine, D-377
 Ervafoleone, E-185
 Ervafoleone, E-186
 Leurosine, L-162
 Leurosine, L-163
 Pseudovincadifformine, P-733
 Vinblastine, V-110
 Vincathicine, V-119
 Vincovalicine, V-127
 Vincristine, V-128

Pyridocarbazole alkaloids

Angustilodine, A-1016
 Ellipticine, E-67
 Elliptinine, E-68
 9-Methoxyellipticine, M-252
 Olivacine, O-83
 Strellidimine, S-557

Ulean alkaloids

Dasycarpidone, D-84
 Gilbertine, G-76

Manilamine, M-84
 Uleine, U-8

Vallesaman alkaloids

Angustilobine A, A-1014
 Angustilobine B, A-1015
 Ervaticine, E-191
 Isobrafouedine, I-199
 Losbanine, L-264
 Monogagine, M-693
 Pseudovobparicine, P-734
 6,7-Secoangustilobine A, S-177
 Undulifoline, U-33
 Vallesamine, V-9
 Vobparicine, V-186

Eburna alkaloids

Callichiline, C-35
 Celastromelidine, C-232
 Celastromeline, C-233
 Cuanzine, C-800
 Decarbomethoxyapocuanzine, D-117
 Infrafractopicrine, I-140
 Insulopinine, I-151
 Isoschizogamine, I-313
 Larutensine, L-46
 Melaxillarinine, M-161
 Nauclefidine, N-68
 Paucivenine, P-132
 Pleiomutine, P-521
 Saifine, S-16
 Strempelepidine, S-558
 Tenuicausine, T-77
 Terengganensine A, T-82
 Terengganensine B, T-83
 Umbellamine, U-21
 Vincarodine, V-117
 Vincatine, V-120
 Vobtusamine, V-187

Tacaman alkaloids

Tacamine, T-12
 Tacamonine, T-13
 Tronocarpine, T-630

Schizozygine alkaloids

Andrangine, A-1002
 Criophylline, C-751
 10,10'-Methylenebis[norvallesamidine],
 M-439
 Schizogaline, S-133
 Schizophylline, S-136
 Schizozygine, S-137
 Strempelepine, S-559
 Vallesamidine, V-8

Chippiine alkaloids

Chippiine, C-389
 Dippinine C, D-824

Secologanin-derived dimer alkaloids

Accedinisine, A-28
 Afrocurarine, A-174
Rauwolfia obscura Alkaloid C₂, A-434
Voacanga thouarsii Alkaloid C, A-456
Rauwolfia obscura Alkaloid D₁, A-473
Rauwolfia obscura Alkaloid D₂, A-474
 C-Alkaloid D, A-476
Melodinus aeneus Alkaloid D, A-488
Pandaca caducifolia Alkaloid D, A-490
Mumtafari sessilifolia Alkaloid E, A-502
Pandaca caducifolia Alkaloid E, A-503
Voacanga thouarsii Alkaloid E, A-506
Pandaca caducifolia Alkaloid F, A-515
Pandaca caducifolia Alkaloid G, A-518
Voacanga thouarsii Alkaloid G, A-519
Voacanga thouarsii Alkaloid I, A-340

Alstonia constricta Alkaloid, A-270
Strempeleopsis strempeleoides Alkaloid, A-334
Uncaria callophylla Alkaloid, A-337
 Alstocraline, A-641
 Alstomacroline, A-644
 Alstomacrophylline, A-645
 Alstonisidine, A-653
 Amataine, A-676
 Angusticraline, A-1011
 Aspidexcelcine, A-1488
 Aspidexcine, A-1489
 10,10'-Bi[N-acetyl-11,12-dihydroxyaspidosper-
 midine], B-115
 12,12'-Bi[11-hydroxycoronaridine], B-127
 Cabufile, C-3
 Calebassine, C-31
 Callichiline, C-35
 Callophylline B, C-38
 Capuvosidine, C-105
 Capuvosine, C-106
 Caracurine II, C-111
 Caracurine V, C-110
 Carosidine, C-166
 Carosine, C-167
 Catharanthamine, C-194
 Catharicine, C-196
 Catharine, C-197
 Celastromelidine, C-232
 Celastromeline, C-233
 Ceylanicine, C-314
 Ceylanine, C-315
 Cimilophytine, C-442
 Conodiparine A, C-611
 Conodiparine B, C-612
 Conodirinine A, C-613
 Conodirinine B, C-614
 Conoduramine, C-615
 Conodurine, C-616
 Conodutarine A, C-617
 Conofoline, C-618
 Conophyllidine, C-621
 Coryzeylamine, C-703
 Criophylline, C-751
 C-Curarine, C-809
 16'-Decarbomethoxy-19,20-dihydroconodura-
 mine, D-118
 16-Decarbomethoxyvoacamine pseudoindoxyl,
 D-121
 Deformylcoryzeylamine, D-143
 Dispegatrine, D-864
 Divarine, D-875
 Dolichocurine, D-910
 Ervafole, E-185
 Ervafole, E-186
 Ervahainamidine A, E-187
 Ervahainamidine B, E-188
 Ervahanine A, E-189
 Flexicorine, F-88
 Folicangine, F-117
 Gardmultine, G-19
 Geissolosimine, G-31
 Geissospermine, G-35
 Guiaflavine, G-219
 Guianensine, G-220
 Guillauminiine, G-221
 Hazuntamine, H-79
 Hazuntiphyllidine, H-80
 Hazuntiphylline, H-81
 Hunteramine, H-369
 Insulopinine, I-151
 Isoborreverine, I-198
 Kisantine, K-51
 Leurosine, L-162
 Leurosine, L-163
 Longicaudatine Z, L-250
 Longicaudatine, L-249
 Macralstonidine, M-7
 Macralstonine, M-8
 Macrocarpamine, M-11
 Macrophylline, M-21
 Macropegatrine, M-30
 Matopense, M-120
 Melaxillarinine, M-161

Melomorsine, M-187
 10-Methoxy-11-[10-(11-methoxyvincamajinyl)]-
 cathafoline, M-258
 11-[10-(11-Methoxyvincamajinyl)]vincorine,
 M-289
 Methylenebismehranine, M-437
 N^β,N^β-Methylenebismeloninium(2+), M-438
 10,10'-Methylenebis[norvallesamidine], M-439
 Monogagine, M-693
 Neburnamine, N-83
 Neoleurocristine, N-123
 Neoleurosine, N-124
 Nitaphylline, N-238
 Pandicine, P-66
 Panganensine R, P-69
 Panganensine X, P-68
 Paucivenine, P-132
 Peceylanine, P-147
 Peceyline, P-148
 Peduncularidine, P-161
 Pelankine, P-172
 Perhentinine, P-256
 Pleiocorine, P-519
 Pleiocraline, P-520
 Plumocraline, P-527
 Polyervinine, P-560
 Pseudovinblastinediol, P-732
 Pycnanthine, P-827
 Pycnanthinine, P-828
 Quimbeline, Q-10
 Rausutranine, R-25
 Rausutrine, R-26
 Roseadine, R-127
 Rovidine, R-137
 Scandomelidine, S-119
 Scandomeline, S-120
 Scandomelonine, S-121
 Serpentinine, S-253
 Strellidimine, S-557
 Strempelepidine, S-558
 Strychnobaridine, S-590
 Strychnobiline, S-591
 Strychnochrysin, S-593
 Strychnofuramine, S-596
 Strychnogucine A, S-597
 Strychnogucine B, S-598
 Strychnohexamine, S-599
 Sungucine, S-642
 Tabernaelegantinine A, T-3
 Tabernaelegantinine B, T-4
 Tabernaelegantinine C, T-5
 Tabernaemontabovine, T-6
 Tabernaemontavine, T-7
 Tabernamine, T-8
 Tenuicausine, T-77
 Tenuiphylline, T-80
 Tetrastachyne, T-294
 Tetrastachynine, T-295
 Toxiferine I, T-436
 Umbellamine, U-21
 Uncaramine, U-22
 Undulatine, U-32
 Villalstonine, V-103
 Vinamidine, V-107
 Vinaphamine, V-108
 Vinblastine, V-110
 Vincamicine, V-114
 Vincarubine, V-118
 Vincathicine, V-119
 Vincovalcine, V-127
 Vincristine, V-128
 Vindolicine, V-129
 Vindolidine, V-131
 Vingramine, V-133
 Vinosidine, V-135
 Vinsedicine, V-137
 Vinsedine, V-138
 Voacamidine, V-167
 Voacamine, V-168
 Voacinal, V-173
 Voafolidine, V-175
 Voafrine A, V-176
 Vobasonidine, V-184

Vobatricine, V-185
 Vobparicine, V-186
 Vobtusamine, V-187
 Vobtusine, V-188

Isoindoles

Antibiotic 0231A, A-1088
 Antibiotic F 1839A, A-1164
 Antibiotic SMTP 7, A-1264
 Antibiotic SMTP 8, A-1265
 Antibiotic SQ 02-S-L1, A-1268
 Antibiotic SQ 02-S-V2, A-1269
 Bhimamycin D, B-114
 Cichorine, C-437
 Dactyline, D-6
 2,3-Dihydro-4-hydroxy-1*H*-isoindol-1-one, D-442
 Duricaulic acid, D-957
 Hericerin, H-145
 Lactonamycin, L-9
 Magallanesine, M-39
 5-Methoxy-2,6-dimethyl-2*H*-isoindole-4,7-dione, M-248
 Prioline, P-631
 Salfredin C₁, S-21
 Shihunidine, S-287
 SMTP 3, S-331
 SMTP 4, S-332
 SMTP 5, S-333
 SMTP 6, S-334
 Stachylocin A, S-489
 Stachyflin, S-492
 Vitedoamine A, V-164
 Xylactam, X-26

Tryptoquivalines

Anacine, A-966
 Ardeemin, A-1399
 Asperlicin E, A-1481
 Asperlicin, A-1478
 Fiscalin A, F-57
 Fiscalin B, F-58
 Fiscalin C, F-59
 Fumiquinazoline A, F-185
 Fumiquinazoline C, F-186
 Fumiquinazoline H, F-189
 Fumiquinazoline I, F-184
 Fumitremorgin C, F-190
 Glyantrypine, G-109
 Lapatin A, L-41
 Lapatin B, L-42
 Nortryptoquivaline, N-315
 Spiroquinazoline, S-438
 Tryprostatin A, T-638
 Tryptoquialanine A, T-643
 Tryptoquialanine B, T-644
 Tryptoquivaline A, T-645
 Tryptoquivaline F, T-646
 Tryptoquivaline G, T-647
 Verrucine A, V-83

Mitomycins

Albomitomycin A, A-246
 Antibiotic FR 66979, A-1171
 Antibiotic FR 900482, A-1177
 Mitiromycin A, M-652
 Mitomycin A, M-653
 Mitomycin C, M-654
 Mitomycin G, M-655
 Mitomycin H, M-656

Pyrrolo[4,3,2-de]quinoline alkaloids

Batzelline A, B-52
 Damirone A, D-13
 Dehydrobufotenine, D-153
 Discorhabdin A, D-850
 Discorhabdin C, D-851
 Discorhabdin D, D-852

Discorhabdin H, D-853
 Discorhabdin T, D-854
 Discorhabdin V, D-849
 Discorhabdin W, D-855
 Epinardine A, E-96
 Epinardine B, E-97
 Haematopodin B, H-8
 Haematopodin, H-7
 Isobatzelline A, I-193
 Lymphostin, L-354
 Makaluvamine A, M-504
 Makaluvamine F, M-55
 Mycenarubin A, M-784
 Mycenarubin B, M-785
 Mycenarubin D, M-786
 Plakinidine A, P-503
 Plakinidine E, P-504
 Prianosin C, P-629
 Tsitsikammamine A, T-650
 Veitamine, V-46
 Wakayin, W-3

Miscellaneous indole alkaloids

Abereamine 1, A-8
 Abereamine 2, A-9
 Abereamine 3, A-10
 Abereamine 4, A-11
 Actinophyllic acid, A-127
 Ailanindole, A-208
 Alboinone, A-245
 Alioline, A-264
 Almazole C, A-634
 Almazole D, A-635
 Andranginine, A-1003
 Antibiotic 927A, A-1092
 Antibiotic BE 10988, A-1126
 Antibiotic 0089D, A-1155
 Antibiotic KG 431A, A-1191
 Antibiotic OPC 15161, A-1220
 Antibiotic SS 228B, A-1270
 Antibiotic TMC 2A, A-1292
 Antibiotic TMC 95A, A-1293
 Antibiotic VM 55599, A-1307
 Aplicyanin A, A-1343
 Arboflorine, A-1380
 Arbophylline, A-1382
 Arestrictin A, A-1406
 Argiopin II, A-1415
 Argiopin IV, A-1416
 Argiotoxin 659, A-1417
 Argiotoxin 673, A-1418
 Aristomakine, A-1437
 Aspergillamide A, A-1471
 Asteochrome, A-1509
 Asterridinone, A-1518
 Asterriquinone D, A-1521
 Asterriquinone, A-1520
 Balasubramide, B-10
 Baretin bromobenzisoxazolone, B-25
 Baretin, B-24
 Bengacarboline, B-58
 Benzastatin F, B-69
 Benzastatin G, B-70
Cypridina Biluciferyl, B-137
 Bisancorinolate B, B-155
 2,5-Bis(1*H*-indol-3-ylmethyl)pyrazine, B-191
 Bouchardatine, B-260
 Brassicanal B, B-282
 Braznitidumine, B-288
 Bruceolline D, B-368
 Campactamanine, C-65
 Cartormin, C-170
 Caulersin, C-215
 Cavernicolin I, C-221
 Celenamide A, C-234
 Celenamide B, C-235
 Celenamide C, C-236
 Celenamide E, C-238
 Celogentin A, C-241
 Celogentin C, C-242
 Celogentin G, C-243
 Chondriamide A, C-421
 Chromoviridan, C-426
 CNS 2103, C-535
 CNS 2104, C-536
 Cochliodinol, C-547
 Cochliodinone, C-548
 Coscinamide A, C-704
 Crenulatine, C-734
 Criamide A, C-737
 Cyclobrassinin, C-842
 Cyclobrassinone, C-843
 Cyclomarazine A, C-877
 Dactylamide B, D-2
 Decursivine, D-137
 Dendrodoine, D-213
 Diazonamide B, D-298
 Dichomine, D-358
 Dichotine, D-359
 Didemnimide A, D-379
 5,12-Dihydrocycloocta[1,2-*b*:5,6-*b'*]diindole-6,13-dicarboxylic acid, D-415
 5*α*,6-Dihydro-3-(1*H*-indol-3-yl)-1,4-dimethoxy-10*bH*-benzofuro[2,3-*b*]indole-2,10*b*-diol, D-466
 5,6-Dihydro-3-[2-(1*H*-indol-3-yl)-2-oxoethyl]-2(1*H*)-pyridinone, D-468
 1-[4,9-Dihydro-2-(methylthio)-1,3-thiazino[6,5-*b*]indol-4-yl]-2-propanone, D-481
 3-(4,5-Dihydro-2-thiazolecarbonyl)-1*H*-indole, D-516
 3,4-Di-1*H*-indol-3-yl-2(5*H*)-furanone, D-690
 2,2-Di-1*H*-indol-3-yl-4-methylpentanoic acid, D-691
 2,2-Di-1*H*-indol-3-ylpropanoic acid, D-694
 Dilemmaone A, D-703
 2-(Dimethylamino)-1,5-dihydro-5-(1*H*-indol-3-yl)-5-(2-oxopropyl)-4*H*-imidazol-4-one, D-728
 2-(Dimethylamino)-5-(1*H*-indol-3-yl)-4*H*-imidazol-4-one, D-732
 α,α -Dimethyl-5-(3-methyl-1-oxo-2-butenyl)-1*H*-indole-3-propanal, D-756
 Diolmycin A1, D-789
 Discoderminidole, D-845
 Dragmacidin A, D-929
 Dragmacidin E, D-931
 Dragmacidin F, D-932
 Duocarmycin C₁, D-953
 Emeniveol, E-71
 Emindole DB, E-75
 Eripinal, E-181
 Eriptide, E-182
 5-Ethyl-2-methyl-11*H*-pyrido[3,4-*a*]carbazolum(1+), E-258
 1*b*-Ethyl-1*a*,1*b*,3,4,8*c*,9,10,10*a*-octahydro-2*H*-oxireno[4,5-*b*]pyrido[3,2-*c*]carbazole, E-262
 Eusynstyelamide, E-303
 Fargesine, F-9
 Fascaplysin, F-13
 Fasmerianamine B, F-16
 Fischerindole L, F-63
 Flabelliformine, F-74
 Fumiquinazoline F, F-188
 Gelseiridone, G-46
 Ginsenine, G-78
 Glucoisatisin, G-102
 Goniomitine, G-142
 Gramodendrine, G-155
 Granulatimide, G-169
 Grossularine 1, G-183
 Halocyamine A, H-35
 Halocyamine B, H-36
 Harmalacinine, H-62
 Harmalanine, H-63
 Hedyocapitelline, H-87
 Hemiasterlin C, H-99
 Hemiasterlin, H-98
 Herbindole A, H-140
 Herbindole B, H-141
 Herbindole C, H-142
 Het₃₈₉, H-155
 Het₄₀₃, H-156
 2,3-Hexalobine E, H-257

Hinnuliquinone, H-288
 Hunteracine, H-368
 20-Hydroxyisoaflavinine, H-541
 3-Hydroxy-3,4-seccoronaridine, H-732
 Hyrtiazepine, H-778
 Ibophyllidine, I-6
 Iboxyphylline, I-10
 Iheyamine A, I-22
 Iheyamine B, I-23
 Indisocin, I-74
 Indolmycin, I-102
 Indolo[2,3-*c*]quinolizine, I-113
 1-(3-Indolyl)- β -carboline, I-119
 4-(3-Indolylmethyl)-7-(2-methylpropyl)-1,2,3,5,6-pentathiepane, I-129
 3-(3-Indolylmethyl)-5-(2-methylpropyl)-1,2,4-trithiolane, I-130
 Ipobscurine C, I-169
 Isatinone A, I-183
 Isatinone B, I-184
 Isoasterriquinone, I-191
 Isogranulatimide, I-237
 Isomonodoroindole, I-248
 8*H*-Isothiazolo[5,4-*b*]indole, I-336
 Isotrikintrine B, I-338
 Kapakahine G, K-13
 Koniamborine, K-64
 Kopsijasminilam, K-72
 Kutzneride 1, K-116
 Lapidilectine A, L-43
 Lapidilectine B, L-44
 Leiocarpadiol, L-83
 Leuconoxine, L-156
 Lotthanongine, L-265
 Lucentamycin B, L-275
Cypridina Luciferin, L-281
 Lundurine A, L-294
 Lycogalic acid, L-329
 Malasseziacitrin, M-60
 Malasseziazole A, M-62
 Malasseziazole B, M-63
 Malasseziazole C, M-64
 Malasseziindole A, M-66
 Margaritarine, M-102
 Martefragine A, M-109
 Martensine A, M-110
 Meridianin A, M-205
 Mersicarpine, M-210
 Mersidasine C, M-211
 Mersifoline A, M-212
 3,3'-Methylenebisindole, M-436
 Monodontamide D, M-690
 Moroidin, M-702
 Moschamindolol, M-711
 Murrapanine, M-750
 Nauclealine B, N-58
 Naucedine, N-66
 Nauclefoline, N-72
 Nauclexine, N-77
 Naulafine, N-78
 Neochlodioidol, N-106
 Neolaugerine, N-122
 Nephilatoxin 1, N-152
 Nephilatoxin 2, N-153
 Nephilatoxin 3, N-154
 Nephilatoxin 4, N-155
 Nephilatoxin 5, N-156
 Nephilatoxin 6, N-157
 Nephilatoxin 7, N-158
 Nephilatoxin 9, N-159
 Nephilatoxin 10, N-160
 Nephilatoxin 11, N-161
 Nephilatoxin 12, N-162
 Nephilatoxin 473, N-163
 Nephilatoxin 501, N-164
 Nephilatoxin 643, N-167
 Ngouniense, N-186
 Nortopsentin A, N-312
 Nortopsentin D, N-313
 Nostodione A, N-323
 Notoamide C, N-324
 Ochrindole A, O-17
 Ochrindole D, O-18

Orbiculamide A, O-101
 Oscillatorin, O-128
 Oxazinin 1, O-153
Cypridina Oxyluciferin, O-222
 Pandine, P-67
 Pauciflorine A, P-130
 Pauciflorine B, P-131
 Peduncin, P-160
 Pericidine, P-257
 Petromurin A, P-285
 Petromurin C, P-286
 Phalarine, P-305
 Pityriacitrin B, P-489
 Pityriacitrin, P-488
 Pityriandride, P-491
 Pityriarubin B, P-493
 Pityriarubin C, P-494
 Prenostodione, P-615
 Proviolacein, P-675
 Psammopemmin A, P-697
 Pseudoanchnazine A, P-698
 Pseudoanchnazine C, P-700
 Pseudodeoxyviolacein, P-711
 Psychrophilin A, P-745
 Psychrophilin B, P-746
 Psychrophilin C, P-747
 Psychrophilin D, P-748
Aesculus hippocastanum Quercetin indoxylacetilglycoside, Q-7
 Rhazimine, R-73
 Rhazizine, R-77
 Rhopaladin A, R-89
 Rutalexin, R-170
 Saifine, S-16
 Sanguinolentaquinone, S-57
 Schischkiniin, S-127
 Schizocommunine, S-132
 Schoberidine, S-139
 Scholarisine A, S-141
 Scytonine, S-169
 Secaloside A, S-173
 Secobatzelline A, S-178
 Serotobenine, S-252
 Strictibine, S-579
 Suaveoline, S-618
 Subincanadine A, S-623
 Subincanadine F, S-625
 Subincanine, S-626
 Tangutorine, T-27
 Tardioxopiperazine A, T-32
 Tardioxopiperazine B, T-33
 Terezine D, T-86
 [3,2':2(3'*H*),3''-Ter-1*H*-indol]-3'-one, T-87
 [3,3':3(2'*H*),3''-Ter-1*H*-indol]-2'-one, T-88
 Terpeptin, T-96
 Terriquinone A, T-106
 5,5',7,7'-Tetrabromo-6,6'-dimethoxyindigotin, T-112
 Tetrahydroechinulin, T-137
 5,5a,12,12a-Tetrahydro-7,14-dimethoxybenzo[1'',2'':4,5;4'',5'':4',5']difuro[2,3-*b*:2',3'-*b'*]diindole-7*b*,14*b*-diol, T-165
 1,3,4,9-Tetrahydro-1,1-dimethylpyrano[3,4-*b*]indole, T-169
 Tetramethoxyscytonemin, T-279
 Thienodolin, T-374
 Topsentin A, T-432
 Toxin MG 30, T-441
 5,6',7-Tribromo-6-methoxyindigotin, T-470
 Trichophylline, T-492
 Trichotomine, T-497
 1,1,1-Tri-(1*H*-indol-3-yl)ethane, T-590
 Tri-1*H*-indol-3-ylmethane, T-591
 Tyriverdin, T-698
 Urdamycin D, U-43
 Uvarindole D, U-61
 Uvarindole E, U-62
 Valindolmycin, V-6
 Vinoxine, V-136
 Violacein, V-142
 Wakayin, W-3
 Wasalexin A, W-10
 Wuchuyamide I, W-24

Monoterpenoid alkaloids

Alfileramine, A-260
Gentiana macrophylla Alkaloid B, A-418
Skytanthus acutus Alkaloid B, A-424
Gentiana macrophylla Alkaloid C, A-447
Skytanthus acutus Alkaloid E, A-504
Gentiana Alkaloid I, A-301
Gentiana Alkaloid II, A-526
Gentiana cruciata Alkaloid IVb, A-530
Gentiana Alkaloid V, A-302
Gentiana Alkaloid VI, A-604
 Altemicidin, A-660
 Argutane A, A-1420
 Austrodimerine, A-1557
 Avicennamine, A-1565
 Bakankoside, B-8
 Benthophoenin, B-66
 Brachycerine, B-264
 Cantharimide, C-89
 Cantleyine, C-92
 Caulindole A, C-218
 Caulindole C, C-219
 Coelobillardierine, C-562
 Delavayine A, D-184
 Delavayine B, D-185
 Delavayine C, D-186
 6,7-Dihydro-4-(hydroxymethyl)-2-(*p*-hydroxyphenethyl)-7-methyl-5*H*-2-pyridinium(1+), D-446
 5,6-Dihydro-3,6,6-trimethyl-7*H*-2-pyridin-7-one, D-525
 Dinklageine, D-786
 Enicoflavine, E-82
 Euphrosine, E-296
 Fontaineine, F-124
 Gardenamide A, G-17
 Genipamide, G-52
 Gentianadine, G-54
 Gentianal, G-55
 Gentianamine, G-56
 Gentiananine, G-57
 Gentianidine, G-58
 Gentianine lactam, G-60
 Gentianine, G-59
 Gentiatibetine, G-61
 Gentiocrucine, G-62
 Gentioflavine, G-63
 4*a*-Hydroxyskytanthine, H-733
 4-Hydroxyskytanthine, H-735
 7-Hydroxyskytanthine, H-736
 Incarvilleatine E, I-61
 Incarvilleatine, I-60
 Incarvilline, I-62
 Incarvine A, I-63
 Indicamine, I-66
 Integramine, I-157
 Isoalfileramine, I-188
 Isogentialutine, I-236
 Kopsilactone, K-73
 Kopsone, K-82
 Linavuline, L-181
 Lindennialine, L-182
 Louisianin C, L-272
 Nepetalactam, N-151
 4-Nor-7,8-dehydro-10-hydroxyskytanthine, N-294
 2-[Octahydro-4,7-dimethyl-1-oxocyclopenta[*c*]pyran-3-yl]nepetalactam, O-43
 Oliveramine, O-84
 Oliverine, O-86
 Oxerine, O-160
 Pedicularidine, P-156
 Pediculidine, P-157
 Pediculine, P-158
 Plectrodorine, P-517
N,N'-Propylenebis(cantharimide), P-650
 Puhinamide, P-780
 Rhexifoline, R-78
 Scaevodimerine A, S-115
 Scaevoline, S-116
 Spicatine, S-408
 Tecomanine, T-58

Tecostanine, T-59
Tetrahydrocantlyne, T-145
Venoterpine, V-52
Xylostosidine, X-31

Sesquiterpene alkaloids

Acanthamolide, A-23
Acanthisonitrile 3, A-24
Atractylenolactam, A-1536
Axamide 1, A-1570
Axinyssimide A, A-1577
Axinyssimide B, A-1578
Bisparthenolidine, B-215
Boneratamide A, B-246
Boneratamide B, B-247
[2-Chloro-2-[4-chloro-5-hydroxy-3-methyl-3-(4-methyl-3-pentenyl)cyclohexylidene]ethyl]carbonimidic dichloride, C-400
Cladioxazole, C-488
Clavulinin, C-520
Cyclospinospongine, C-908
2,10-Dichloro-6,11-cyclo-3(15),7(14)-farnesadien-1-yl carbonimidic dichloride, D-349
Drimentine C, D-937
Drimentine D, D-938
Drimentine E, D-939
Eremophilene lactam, E-146
Guaipyridine, G-190
Halichonadin A, H-14
Hederacine A, H-85
Involucratine, I-162
3-Isocyanato-7,9-bisaboladiene, I-213
4-Isocyanomaalane, I-219
9-Isocyanoneopupekeane, I-220
6-Isodaucene 14-isonitrile, I-225
Isopolyalthenol, I-269
Kopsirachin, K-81
Manicoline A, M-82
Metachromin G, M-228
Metachromin H, M-229
Nakijiquinone A, N-14
Nakijiquinone B, N-15
Nakijiquinone D, N-16
Neopolyalthenol, N-135
Oceanapamine, O-15
Paeciloxazine, P-9
Patchoulipyridine, P-112
Phenazinomycin, P-329
Piccolamine, P-402
Polyalthenol, P-540
Polyavolinamide, P-546
Pulchellidine, P-785
Pyrodysinoic acid, P-936
Rotundine B, R-135
Saussureamine A, S-105
Saussureamine C, S-106
Saussureamine D, S-107
Siphonodictidine, S-324
Smenospongine, S-329
Stachybocin A, S-489
Stylotellane A, S-613
Stylotellin, S-614
Suaveolindole, S-617
Triendiol, T-592
Trikenramine, T-593
3-(3,7,11-Trimethyl-2,6,10-dodecatrienyl)-1*H*-indole, T-605

Macrocyclic sesquiterpene alkaloids

Cathedulín E3, C-201
Cathedulín K17, C-202
Cathedulín K19, C-203
Cathidine A, C-210
Hippocrateine III, H-296
Triptonine A, T-619
Triptonine B, T-620

Dendrobium alkaloids

Crepidamine, C-735
Dendrine, D-207

Dendrobine, D-210
Dendrorepine, D-212
Dendronobilin A, D-214
Dendrowardine, D-215
Dendroxine, D-216
Mubironine C, M-717
Nobiline, N-262
 δ -Nobilonine, N-263

Nuphar alkaloids

Allothiobinupharidine, A-631
7-Demethyldioxynupharidine, D-201
Deoxynupharidine, D-227
5-(3-Furanyl)octahydro-8-methylindolizine, F-202
Nuphacristine, N-339
Nupharamine, N-340
 α -Nupharidine, N-341
 β -Nupharidine, N-342
Nupharine, N-343
6,7-Oxidodeoxynupharidine dimer, O-161
Pseudothiobinupharidine, P-729
Thiobideoxynupharidine, T-379
Thiobinupharidine, T-380

C19-Diterpenoid alkaloids and 4-nor analogues

Aconitine, A-104
Acoseptine, A-106
Aconine, A-123
Ajabicine, A-210
Akirine, A-226
Alboviolaconitine D, A-248
Andersonine, A-998
Anhwedelphinine, A-1023
Bishaconitine, B-179
Bulleyanitin B, B-408
Bulleyanitin C, B-409
Delavaine A, D-182
Delavaine B, D-183
Delboxine, D-187
Francheline, F-147
Glaudelsine, G-91
Graciline, G-152
Grandifloritine, G-157
Heteratisine, H-157
Jiufengsine, J-41
Kiridine, K-46
Kongboentine A, K-63
Lycaconitine, L-317
18-*O*-[2-(2-Methyl-4-oxo-4*H*-quinazolin-3-yl)-benzoyl]lycoctonine, M-501
Pengshenine A, P-184
Potanisine G, P-578
Puberaconitine, P-776
Racemulosine, R-2
Secojesaconitine, S-187
Secokaraconitine, S-189
Sinomontanine C, S-315
Sinomontanine I, S-314
Sinomontanitine A, S-316
Tangirine, T-26
Vilmoritine, V-106

C20 Diterpenoid alkaloids (Atisine type)

Acophine, A-105
Acozerine, A-108
Ajaconine, A-211
Albovionitine, A-249
Anopterimine, A-1070
Atidine, A-1533
Atisine, A-1534
Azitine, A-1609
Bishatisine, B-180
Brunonine, B-371
Cardionidine, C-159
Chuanfumine, C-434
Cochlearenine, C-546

Cordizine, C-642
Coryphidine, C-696
Coryphine, C-697
Dehydronapelline, D-162
Delatisine, D-180
Delnudine, D-190
Delphatisine B, D-192
Denudatine, D-221
Dictysine, D-370
12,16-Epoxy-16,17-dihydronapelline, E-131
Garryine, G-25
Gomandonine, G-139
Guan-fu base K, G-197
Guan-fu base S, G-198
Hetidine, H-161
Hetisan-11-one, H-173
Isoatisine, I-192
Kirimine B, K-48
Lassiocarpine, L-51
Lepentine, L-105
Lindheimerine, L-184
Macrocentrine, M-12
Miyaconitine, M-665
Napelline, N-30
Navirine, N-79
Ouvrardiandine A, O-141
Paniculamine, P-70
Pukeensine, P-782
Racemulotine, R-3
Sczukidine, S-170
Septatisine, S-247
Spiraeine A, S-413
Spirafine II, S-414
Spiramine C, S-416
Spiramine J, S-417
Spiramine P, S-418
Spiramine Z₂, S-419
Spirasine III, S-421
Spirasine X, S-420
Spiratine B, S-422
Spireine, S-423
Staphinine, S-494
Staphisagnine, S-495
Staphisine, S-496
Tallassamine, T-18
Tangirine, T-26
Thalicessine, T-317
Tongolinine, T-427
Trabzonine, T-442
Veatchine, V-45
Vilmoridine, V-105
Yesonine, Y-10
Zeraconine, Z-16

Erythrophleum alkaloids

Erythrophleum Alkaloid A, A-371
Erythrophleum Alkaloid C, A-446
Erythrophleum C₂₄ Alkaloid, A-267
Cassaidine, C-178
Coumingaine, C-713
Erythrosumamine, E-218
Erythrosumavine, E-219
Homophleine, H-340
Norcassaidide, N-289
19-Nor-4-dehydrocassaidine, N-293
Norerythrosthachamide, N-299
Norerythrosthachamine, N-300

Penitremes

Aflatrem B, A-170
Aflatrem, A-169
Aflavazole, A-171
Aflavinine, A-172
Anthcolorin A, A-1075
Anthcolorin E, A-1076
Aspernomine, A-1485
Emindole DA, E-74
Emindole PA, E-76
Emindole PB, E-77
Emindole SB, E-78
Janthitrem E, J-21

Lolicine A, L-231
 Lolilline, L-233
 Lolitrem B, L-236
 Lolitrem N, L-237
 Nodulisporic acid A₁, N-268
 Nodulisporic acid A, N-269
 Nodulisporic acid C₁, N-270
 Nodulisporic acid C, N-271
 Nodulisporic acid D₁, N-272
 Nodulisporic acid D, N-273
 Nodulisporic acid E, N-274
 Nodulisporic acid F, N-275
 Nominine, N-279
 Paspaline, P-108
 Paspalinine, P-109
 Paspalitrem A, P-110
 Paspalitrem C, P-111
 Paxilline, P-140
 Paxinorol, P-141
 PC-M4, P-144
 PC-M5, P-145
 Penitrem C, P-194
 Pennigritrem, P-195
 Petromindole, P-284
 9-Prenylpaxilline, P-618
 Radarin B, R-6
 Secopenitrem B, S-194
 Suspendole, S-263
 Shearinine A, S-281
 Shearinine C, S-282
 Shearinine I, S-280
 Shearinine J, S-283
 Shearinine K, S-284
 Sulpinine B, S-640
 Sulpinine C, S-641
 Terpendole B, T-92
 Terpendole C, T-93
 Terpendole H, T-94
 Terpendole I, T-91
 Terpendole L, T-95
 Thiersinine A, T-377
 Tubingensin A, T-669
 Tubingensin B, T-670

Miscellaneous diterpene alkaloids

Aceropterin, A-29
 Agelasidine B, A-179
 Agelasimine A, A-181
 Agelasine A, A-184
 Agelasine C, A-185
 Agelasine D, A-186
 Agelasine E, A-187
 Agelasine I, A-183
 Agelasine, A-182
 Ageline A, A-191
 Aristolin, A-1434
 Aristoloin I, A-1435
 Asmarine B, A-1465
 Asmarine C, A-1466
 Asmarine I, A-1464
 Aminosol, A-1567
 Bis(gorgiacerol)amine, B-177
 Brasilicardin A, B-280
 Cespitulactam D, C-288
 Chamobtusine A, C-344
 Concavine, C-596
 Daphnimacrine, D-58
 7-[5-(Decahydro-4a-hydroxy-1,2,5,5-tetramethyl-1-naphthalenyl)-3-methyl-2-pentenyl]-3,7-dihydro-2,3-dimethyl-6H-purin-6-one, D-111
 Decaturin A, D-127
 Dichlorolissoclimide, D-354
 7,20-Diisocyanoisocycloamphilectane, D-698
 Echinophyllin A, E-18
 Echinophyllin B, E-19
 Echinophyllin C, E-20
 Echinophyllin D, E-21
 Echinophyllin E, E-22
 Echinophyllin F, E-23
 Elisabethamine, E-66

Euphosalicin, E-295
 Guanacastepene D, G-192
 Guanacastepene H, G-193
 Haterumaimide C, H-73
 Haterumaimide E, H-74
 Haterumaimide F, H-75
 Haterumaimide I, H-72
 Hebeirubescensin A, H-82
 Homopseudopteroxazole, H-343
 Icaceine, I-11
 Ileabethoxazole, I-27
 8-Isocyan-1(12)-cycloamphilectene, I-215
 Isosalviamine A, I-311
 Isosalviamine B, I-312
 Joalin, J-42
 Melemeleone A, M-163
 Melemeleone B, M-164
 Memnobotrin A, M-196
 Milliamine C, M-616
 Milliamine H, M-617
 Neosalvianene, N-137
 Nicotaxine, N-188
 Nominine, N-279
 Norstaminolactone A, N-310
 Petrosaspongiolide L, P-289
 Phorbasin E, P-378
 Polyveoline, P-564
 Popolohuanone A, P-567
 Popolohuanone D, P-568
 Pseudopteroxazole, P-724
 Pyrrocidine B, P-940
 Radarin B, R-6
 RYANIA diterpene ester C₂, R-174
 Ryanodine, R-175
 Salviadione, S-40
 Salvianene, S-42
 Secopseudopteroxazole, S-196
 Sinulamide, S-318
 Sinulamine III, S-320
 Sinulamine II, S-319
 Spicaledonine, S-407
 Stelletazole B, S-508
 Stelletazole C, S-509
 Tauret, T-38
 Taxol, T-52
Agelas nakamura Thelepogane-purine alkaloid, T-352
 Thelepogine, T-354
 Thiersindole A, T-375
 Thiersindole C, T-376
 Triptotin J, T-621

Sesterterpene alkaloids

Fasciospongine A, F-14
 Hyatelactam, H-390
 Kimbasine A, K-44
 Molliorin A, M-673
 Molliorin B, M-674
 Molliorin C, M-675
 Molliorin D, M-676
 Molliorin E, M-677
 Palinurine A, P-22
 Sarcotragin A, S-81
 Sarcotrine A, S-82
 Sarcotrine D, S-83
 Sarcotrine F, S-84
 Spongidine A, S-472
 Spongidine D, S-473

Daphniphylline alkaloids

Daphniphyllum Alkaloid A₁, A-345
 Bukittinggine, B-396
 Caldaphnidine A, C-27
 Caldaphnidine C, C-28
 Caldaphnidine D, C-29
 Caldaphnidine E, C-30
 Calycilactone A, C-43
 Calycine, C-44
 Calyciphylline B, C-45
 Calyciphylline C, C-46

Calyciphylline D, C-47
 Calyciphylline E, C-48
 Calyciphylline F, C-49
 Calyciphylline G, C-50
 Calyciphylline H, C-51
 Calyciphylline J, C-52
 Calydaphninone, C-58
 Daphcalycic acid, D-18
 Daphcalycine, D-19
 Daphcalycinosidine B, D-20
 Daphlongeramine A, D-21
 Daphlongeramine A, D-22
 Daphlongeramine B, D-23
 Daphmacrine, D-24
 Daphmanidine A, D-25
 Daphmanidine B, D-26
 Daphmanidine C, D-27
 Daphnezomine A, D-29
 Daphnezomine C, D-30
 Daphnezomine D, D-31
 Daphnezomine F, D-32
 Daphnezomine K, D-33
 Daphnezomine L, D-34
 Daphnezomine M, D-35
 Daphnezomine O, D-36
 Daphnezomine P, D-37
 Daphnezomine Q, D-38
 Daphnicadine, D-39
 Daphnicaline, D-40
 Daphnicamine, D-41
 Daphnicyclidine A, D-42
 Daphnicyclidine D, D-43
 Daphnicyclidine H, D-44
 Daphnicyclidine J, D-45
 Daphnicyclidine K, D-46
 Daphnicyclidine L, D-47
 Daphniglaucine B, D-48
 Daphniglaucine C, D-49
 Daphnigraciline, D-50
 Daphnigracine, D-51
 Daphnilactone A, D-52
 Daphnilactone B, D-53
 Daphnilongeramine A, D-54
 Daphnilongeramine C, D-55
 Daphnilongerine, D-56
 Daphnilongertone, D-57
 Daphnimacropodine A, D-59
 Daphnioldhanine A, D-61
 Daphnioldhanine C, D-62
 Daphnioldhanine J, D-64
 Daphnipaxianine A, D-65
 Daphnipaxianine D, D-66
 Daphnipaxinine, D-67
 Daphniphylline, D-68
 Daphniyunnine D, D-69
 Daphtenidine A, D-70
 Daphtenidine B, D-71
 Daphtenidine C, D-72
 Glaucescine, G-89
 Glaucescine, G-90
 Longercinphylline A, L-245
 Longercinphylline B, L-246
 Longistylumphylline B, L-254
 Longistylumphylline C, L-255
 Macropodumine A, M-23
 Macropodumine B, M-24
 Macropodumine C, M-25
 Macropodumine D, M-26
 Macropodumine E, M-27
 Neodaphniphylline, N-109
 Neoyuzurimine, N-150
 Oldhamine A, O-77
 Oldhamiphylline A, O-78
 PAXDAPHNIDINE A, P-137
 PAXDAPHNINE A, P-138
 PAXDAPHNINE B, P-139
 Paxiphylline A, P-142
 Paxiphylline B, P-143
 Secodaphniphylline, S-181
 Yunnandaphnine E, Y-20
 Yuzurimine C, Y-23
 Yuzurimine D, Y-24

Steroidal alkaloids (salamandra type)

Cycloneosamandione, C-886
 1,4-Epoxy-3-aza-*A*-homoandrostan-17-ol, E-127
 Isocycloneosamandaridine, I-223
 Samandaridine, S-43
 Samandarine, S-44
 Samandenone, S-45
 Samandinine, S-46
 Samanine, S-47

Steroidal alkaloids (jerveratrum type)

Edpetine, E-39
 Hupehenisine, H-371
 Jervine, J-36
 Kuroyurinidine, K-114
 Ningpeisine, N-223
 Peimisine, P-168
 Pengbeimine A, P-182
 Pengbeimine C, P-183
 Veratramine, V-71
 Verdine, V-74
 Yibeissine, Y-11

Steroidal alkaloids (cerveratrum type)

Cevanine, C-299
 3,25-Dihydroxycevan-6-one, D-562
 Germine, G-70
 Hupehendine, H-370
 Neogermine, N-116
 Neoveratraline A, N-148
 Severidine, S-277
 Ussuriedine, U-56
 Veracevine, V-56
 Zygadenilic acid δ -lactone, Z-35
 Zygadenine, Z-36

Steroidal alkaloids (conanine type)

Conarrhimine, C-595
 Conessine, C-601
 Conkurchine, C-610
 Didymeline, D-383
 19,23-Dinorcona-1,3,5(10),14-tetraenine-3-ol, D-788
 Funtudienine, F-198
 Holadienine, H-314
 Holaline, H-315
 Holaromine, H-317
 Holarrhenine, H-318
 Holarrhine, H-320
 Holonamine, H-322
 Kurchamide, K-109
 Kurchilidine, K-112
 Kurcholessine, K-113
 Latifoline, L-56
 Latifolinine, L-57
 Malarboreine, M-59
 Pistacimidelor, P-485
 Pubescimine, P-778
 Pubescine, P-779
 Regholarrhenine C, R-39

Steroidal alkaloids (spirosolane type)

Veramine, V-66

Steroidal alkaloids (solanidine type)

Anrakorinine, A-1073
 Ecliptalbine, E-33
 Hakurirodine, H-11
 Hosukinidine, H-364
 Impranine, I-54

Isorubijervine, I-310
 Korsevinine, K-89
 Leptinidine, L-116
 (2-Methyltetrahydrofuran)solanidine, M-567
 Petisidinine, P-280
 Petisine, P-281
 Pingbeidinioside, P-432
 Pingbeimine, P-433
 Procevine, P-633
 Puqietinone, P-807
 Radpetine, R-12
 Rubijervine, R-144
 Sarachine, S-66
 Solacallinidine, S-335
 Solacapine, S-336
 Solacongostidine, S-337
 Solafloridine, S-338
 3,5-Solanidadiene, S-343
 Solanidine, S-345
 Solanudine, S-348
 Solaphyllidine, S-349
 Solaquidine, S-350
 Solaseaforthine, S-351
 Solaspiralidine, S-352
 Veralinine, V-60
 Veralkamine, V-61
 Veralobine, V-62
 Veralodisine, V-63
 Veralosidinine, V-64
 Veramitaline, V-68
 Verazine, V-73
 Vertaline B, V-87

Steroidal alkaloids (buxus type)

Buxus Alkaloid 1, A-275
Buxus Alkaloid 2, A-276
Buxus Alkaloid 3, A-277
Buxus Alkaloid 4, A-278
Buxus Alkaloid 5, A-279
Buxus Alkaloid A1, A-348
Buxus Alkaloid A2, A-349
Buxus Alkaloid A3, A-350
Buxus Alkaloid A4, A-351
Buxus Alkaloid A5, A-352
Buxus Alkaloid B, A-390
Buxus Alkaloid B1, A-391
Buxus Alkaloid B4, A-392
Buxus Alkaloid B5, A-393
Buxus Alkaloid B6, A-394
Buxus Alkaloid B7, A-395
Buxus Alkaloid B8, A-396
Buxus Alkaloid B387, A-397
Buxus Alkaloid B398, A-398
Buxus Alkaloid B426, A-399
Buxus Alkaloid BX6, A-430
Buxus Alkaloid BX10, A-431
Buxus Alkaloid C, A-435
Buxus Alkaloid D, A-475
Buxus Alkaloid E, A-496
Buxus Alkaloid F, A-507
Buxus Alkaloid M, A-566
Buxus Alkaloid M, A-567
Buxus Alkaloid N, A-570
 Buxafuranamide, B-448
 O¹⁰-Buxafuranamine, B-449
 O⁶-Buxafuranamine, B-450
 Buxalfine, B-452
 Buxalongifolamidine, B-453
 Buxaltine H, B-454
 Buxaminone, B-457
 Buxanaldinine, B-458
 Buxandonine L, B-459
 Buxanoldine, B-460
 Buxapapinolamine, B-461
 Buxaquamarine, B-462
 Buxazine, B-463
 Buxdeltine, B-464
 Buxetine, B-465
 Buxiramine D, B-472
 Buxitrienine C, B-473
 Buxotrienine, B-476
 Buxoxybenzamine, B-477

Buxozine C, B-478
 Buxpapamine, B-479
 Buxpapine, B-480
 Buxpapinine, B-481
 Buxtauine M, B-482
 Buxupapine, B-483
 Cortistatin B, C-665
 Cortistatine F, C-666
 Cortistatine G, C-667
 Cortistatine L, C-668
 Cyclobuxargentine G, C-845
 Cyclobuxomicreanine, C-847
 Cyclobuxophylline O, C-848
 Cyclobuxoviramine, C-849
 Cyclobuxoviridine L, C-850
 Cyclobuxoxazine C, C-851
 Cyclomethoxazine B, C-881
 Cyclomicrobuxeine K, C-882
 Cyclomicurarine L, C-884
 Cyclorolfine, C-905
 Cyclorolfoxazine, C-906
 Cyclosuffrobuxine K, C-912
 Cyclovirobuxeine F, C-916
 2-Dehydroxy-O⁶-buxafuranamine, D-174
 Homobuxaquamarine, H-327
 Homomoenjodaramine, H-339
 Hyrcanine, H-776
 Hyrcanone, H-777
 Karachicine, K-14
 Moenjodaramine, M-667
 17-Oxocycloprotobuxine, O-172
 Papillozine C, P-83
 Pseudobalebuxine F, P-704
 Semperviraminol, S-226
 Semperviraminone, S-227
 Sempervirone, S-229
 Semperviroxazolidine, S-230
 Spiroforabuxine, S-429

Steroidal alkaloids (pregnane type)

Pachysandra terminalis Alkaloid XI, A-612
 Japindine, J-23
 Lanitine, L-30
 Pachysantermine A, P-6
 Pachystermine A, P-8
 Paravallarine, P-93
 Spiropachysine, S-436

Steroidal alkaloids (cephalostatins/ritterazines)

Cephalostatin 2, C-261
 Cephalostatin 3, C-262
 Cephalostatin 5, C-263
 Cephalostatin 6, C-264
 Cephalostatin 7, C-265
 Cephalostatin 8, C-266
 Cephalostatin 9, C-267
 Cephalostatin 13, C-268
 Cephalostatin 16, C-269
 Ritterazine A, R-101
 Ritterazine B, R-102
 Ritterazine C, R-103
 Ritterazine E, R-104
 Ritterazine J, R-105
 Ritterazine N, R-106
 Ritterazine P, R-107
 Ritterazine R, R-108
 Ritterazine U, R-109
 Ritterazine V, R-100
 Ritterazine W, R-110
 Ritterazine Z, R-111

Miscellaneous steroidal alkaloids

15-Azasterol, A-1602
 Batrachotoxinin A, B-36
 Boophiline, B-249
 Bufotoxin, B-394
 Carolisterol A, C-165
 Cephalostatin 7, C-265

3-Deamino-3-hydroxysolanocapsine, D-99
 Dehydroazasterosterol, D-151
 Dictyolucidine, D-368
 4,4-Dimethyl-15-azasterol, D-742
 Gymnasterone A, G-231
 Jurubine, J-60
 Korsevine, K-88
 Kurchenine, K-110
 Machilaminoside A, M-3
 Machilaminoside B, M-4
 4-Methyl-15-azasterol, M-396
 20-(2-Methyl-1-pyrrolin-5-yl)pregn-4-en-3-one,
 M-550
 Mitiphylline, M-651
 Munroniamide, M-732
 Petisidine, P-279
 Petunianine A, P-291
 Petunianine B, P-292
 Petunianine C, P-293
 Pimpinellidine, P-431
 Pingbeinone, P-434
 Plakinamine A, P-496
 Plakinamine B, P-497
 Plakinamine C, P-498
 Plakinamine G, P-499
 Plakinamine I, P-500
 Plakinamine I, P-501
 Plakinamine J, P-502
 Procevine, P-633
 Puqjenine A, P-802
 Puqjenine C, P-803
 Puqjenine D, P-804
 Puqjenine E, P-805
 Puqjenine F, P-806
 Rhinolidine, R-79
 Sevcoridinone, S-275
 Siechuantine, S-298
 Solamaladine, S-339
 Solanocapsine, S-346
 Solanocastrine, S-347
 Spirajine, S-415
 Spiroforabuxine, S-429
 Squalamine, S-485
 Stenanzine, S-526
 Tomatillidine, T-423
 Turraparvin D, T-685
 Uscharin, U-53
 Valivine, V-7
 Veracintine, V-57
 Veramanine, V-65

Azaanthracene alkaloids

Annopholine, A-1058
 Antibiotic WJ 85, A-1313
 Benz[*g*]isoquinoline-5,10-dione, B-71
 Bostrycoidin, B-255
 6-Deoxy-8-methylbostrycoidin, D-226
 Enkleine, E-83
 Griffithazanone A, G-174
 Isophomazarin, I-264
 Kalasinamide, K-6
 Marcanine A, M-97
 4-Methylbenzo[*g*]quinoline-5,10-dione, M-399
 Phomazarine, P-374
 Pyrenoline A, P-892
 Scorazanone, S-161
 5,8,9,10-Tetramethoxy-4-methylbenzo[*g*]quino-
 lin-2(1*H*)-one, T-277

Azafluorene alkaloids

Dielsinol, D-385
 3,7-Dihydro-4-methyl-5*H*-indenol[1,2-*b*]pyri-
 din-5-one, D-623
 Onychine, O-93

Pyrazole alkaloids

Citreozopyrone, C-477
 5,6-Dihydro-3-phenyl-4*H*-pyrrolo[1,2-*b*]pyra-
 zole, D-492

Formycin A, F-131
 Formycin B, F-132
 Garceine, G-16
 3-Methoxy-pyrazolo[4,3-*c*][1,2,4]triazine, M-285
 Nagstatin, N-9
 Newbouldine, N-185
 Nitrabirine, N-240
 3-Nonyl-1*H*-pyrazole, N-286

Imidazole alkaloids

Acinetobactin, A-79
 Aerophobin 2, A-155
 Agesamide A, A-195
 Alchorneinone, A-252
 Allantoin, A-625
 2-Amino-4-(4-hydroxybenzyl)-1-methyl-1*H*-imi-
 dazole, A-791
 5-Aminoimidazole riboside, A-827
 2-Aminoimidazole, A-825
 4(5)-Aminoimidazole, A-826
 Aminozoanemonin, A-928
 Amphistin, A-949
 Anatoxin a(s), A-975
 Anguibactin, A-1009
 Antibiotic CV 1, A-1154
 Antibiotic Sch 40873, A-1245
 Aplysinamisine I, A-1349
 Aplysinopsin, A-1357
 Archerine, A-1390
 Argadin, A-1407
 Asteropterin, A-1516
 Aurantiamine, A-1548
 Azepinomycin, A-1605
 Biotin, B-139
 3-Bromo-4-hydroxy- α -(hydroxyimino)-*N*-[2-
 (1*H*-imidazol-4-yl)ethyl]benzenepropana-
 mide, B-323
 5(4)-Butyl-2-chloro-1*H*-imidazole-4(5)-carbox-
 aldehyde, B-426
 Calcaridine A, C-25
 Carcinine, C-157
 Caribaeorane, C-160
 Caribaeoside, C-161
 Casimiroedine, C-176
 Cassiadinine, C-180
 Catharsitoxin A, C-198
 Catharsitoxin B, C-199
 Catharsitoxin C, C-200
 Celogentin A, C-241
 Celogentin C, C-242
 Celogentin G, C-243
 Chaksine, C-339
 Chartelline A, C-350
 Clathridine A, C-491
 Clathridine B, C-492
 Clathridine C, C-493
 Clathridine-9-*N*-(2-sulfoethyl)imine, C-494
 Corallistine, C-639
 Creatinine, C-732
 Cucumopine, C-802
 Cynodine, C-931
 Cynolujine, C-932
 Cynometrine, C-933
 Cypholophine, C-935
 Dehydroisolongistrobine, D-160
 Didemnoline A, D-380
 7,8-Dihydroimidazo[1,5-*c*]pyrimidin-5(6*H*)-one,
 D-462
 2,3-Dihydro-2-oxo-1*H*-imidazole-4-carboxylic
 acid, D-484
 Dimethyl *N*²-creatininylphosphate, D-725
 2-(Dimethylamino)-1,5-dihydro-5-(1*H*-indol-3-
 yl)-5-(2-oxopropyl)-4*H*-imidazol-4-one, D-728
 4-[2-(Dimethylamino)ethyl]imidazole, D-731
 2-(Dimethylamino)-5-(1*H*-indol-3-yl)-4*H*-imida-
 zol-4-one, D-732
 2,4(5)-Dimethylimidazole, D-749
 Discodermindole, D-845
 Dragmacidin D, D-930
 Dragmacidin F, D-932
 Dragmacidonamine A, D-933
 Echinobetaine B, E-15
 Eleuthoside A, E-64
 Ergothioneine, E-173
 5-[(4-Ethyltetrahydro-5-oxo-3-furanyl)methyl-
 idene]-1-methyl-2,4-imidazolidinedione,
 E-272
 Fasmerianamine B, F-16
 Feldamyacin, F-33
 Fungerin, F-196
 Glusun II, G-106
 Granulatimide, G-169
 Guadinomic acid, G-185
 Guadinomine A, G-186
 Guadinomine C₁, G-187
 Histamine, H-307
 2-(4-Hydroxybenzoyl)-4(5)-(4-hydroxyphenyl)-
 1*H*-imidazole, H-412
 β -Hydroxyergothioneine, H-484
 2-Hydroxy-3-(4-imidazolyl)propanoic acid,
 H-523
 2-(4-Hydroxy-3-methoxybenzoyl)imidazole,
 H-558
 3-(Hydroxymethyl)-4-(1-methyl-1*H*-imidazol-5-
 yl)-1-phenyl-1-butanone, H-606
 2'-*O*-[3-(4-Hydroxyphenyl)-2-methoxyprope-
 noyl]-3'-*O*-[3-(1*H*-imidazol-4-yl)propenoyl]-
 5-methyl-5'-thioadenosine, H-681
 5-Hydroxy-1- β -D-ribofuranosyl-1*H*-imidazole-
 4-carboxaldehyde, H-729
 Hyrtiomanzamine, H-781
 Imbricaine, I-32
 1*H*-Imidazole-1-acetic acid, I-36
 1*H*-Imidazole-4(5)-acetic acid, I-37
 1*H*-Imidazole-4(5)-ethanol, I-38
 Imidazole, I-35
 1*H*-Imidazole-4(5)-propanoic acid, I-39
*N*²-[3-(1*H*-Imidazol-4-yl)propenyl]histamine,
 I-43
 2-(1*H*-Imidazol-4-yl)quinoline, I-44
 Isoalchorneine, I-187
 Isocynometrine, I-224
 Isogranulatimide, I-237
 Isonaamide A, I-251
 Isonaamine A, I-252
 Isonaamine B, I-253
 5-Isopropyl-2-(2-methylpropyl)-1*H*-imidazole,
 I-280
 Kifunensine, K-41
 Kottamide A, K-92
 Kottamide D, K-93
 Lepidine E, L-107
 Lepidine F, L-108
 Lepidine, L-106
 Leucettamide, L-149
 Leucettamine A, L-150
 Leucettamine B, L-151
 Leucettamine C, L-152
 Leucosolenamine A, L-157
 Leuhistin, L-160
 Lidimycin, L-166
 Lissoclin C, L-199
 Longistrobine, L-253
 Macrorungine, M-28
 Malonganone B, M-75
 Mannopeptimycin β , M-86
 Melanostatin, M-158
 Meleagrin, M-162
 α -Methylbiotin, M-402
 α -Methylthiobiotin, M-432
 4(5)-Methylimidazole, M-473
 2-(2-Methylpropyl)-5-phenyl-1*H*-imidazole,
 M-528
 Moroidin, M-702
 Murexine, M-739
 Naamidine F, N-2
 Naamine A, N-3
 Naamine B, N-4
 Naamine C, N-5
 Naamine D, N-6
 Neamphine, N-81
 Neoxaline, N-149
 Nikkomycin X, N-216
 2-Nitroimidazole, N-250
 13-Nor-7(11)-dehydropilocarpine, N-295

Nortopsentin A, N-312
 Nortopsentin D, N-313
 Norzooanemonin, N-318
 Oceanapamine, O-15
 1,2,3,4,6,7,8,9-Octahydrodipyrro[1,2-*a*:1',2'-*c*]imidazol-10-ium(1+), O-45
 Phenylahistin, P-338
 Pilocarpine, P-424
 Pilosine, P-428
 Pilosinine, P-429
 Piloturine, P-430
 Polyandrocarpamine A, P-543
 Polycarpaurine B, P-551
 Polycarpaurine C, P-552
 Polycarpine, P-553
 Preclathridine A, P-594
 Pseudoceratinine B, P-708
 Pseudoceratinine C, P-709
 Pseudomonine, P-718
 Puralidin D, P-810
 Puralidin H, P-811
 Puralidin M, P-814
Subergorgia suberosa Purine-imidazole alkaloid, P-819
 Pyronaamide, P-938
Oceanapia Quinolone alkaloid, Q-57
 Rhopaladin A, R-89
 Roquefortine C, R-125
 Roquefortine E, R-126
 Selenobiotin, S-219
 Sibyllimycin, S-293
 Solsodomine A, S-358
 Spirocalcaridine A, S-428
 Stelletazole A, S-507
 Stelletazole B, S-508
 Stelletazole C, S-509
 Streptolidine, S-565
 4,5,6,7-Tetrahydro-1*H*-imidazo[4,5-*c*]pyridine, T-192
 Topsentin A, T-432
 2,4,5-Tribromo-1*H*-imidazole, T-461
 Ulosantoin, U-20
 Verongamine, V-78
 Wondonine A, W-20
 Xestomanzamine A, X-23
 Zapotidine, Z-8
 Zooanemonin, Z-32
 Zyzzin, Z-39

Pyrrole-imidazole alkaloids

Ageladine A, A-177
 Agelastatin A, A-189
 Ageliferin, A-190
 Axinellamine A, A-1574
 Axinohydantoin, A-1576
 Clathramide A, C-490
 Cyclooroidin, C-887
 Dibromophakellstatin, D-336
 Dispacamide A, D-863
 Hymenialdisine, H-762
 Keramadine, K-29
 Latonduine A, L-59
 Latonduine B, L-60
 Massadine aziridine, M-113
 Massadine, M-112
 Mauritamide A, M-123
 Midpacamide, M-604
 Mukanadin B, M-727
 Nagelamide B, N-7
 Nagelamide H, N-8
 Nakamuric acid, N-12
 Odiline, O-66
 Oroidin, O-123
 Oxocyclostylidol, O-173
 Oxysceptrine, O-225
 Palauamine, P-18
 Preaxinellamine, P-590
 Sceptrine, S-124
 Slagenine A, S-328
 Styloguanidine, S-611
 Tauroacidin A, T-39
 Tubastrindole C, T-655

Cycloheptadiimidazoles

Parazoanthoxanthin A, P-94
 Pseudozoanthoxanthin A, P-737

Oxazole and benzoxazole alkaloids

4-Acetyl-2(3*H*)-benzoxazolone, A-42
 Acinetobactin, A-79
 Ajudazol B, A-219
 Almazole C, A-634
 Almazole D, A-635
 Almazolone, A-636
 Amamistatin A, A-672
 5-(10-Aminoundecyl)hexahydropyrrolo[2,1-*b*]oxazole, A-927
 Annuloline, A-1064
 Antibiotic A 33853, A-1098
 Antibiotic 927A, A-1092
 Antibiotic AJI 9561, A-1116
 Antibiotic B 4317, A-1122
 Antibiotic B 90063, A-1123
 Antibiotic CP 35763, A-1151
 Antibiotic GE 37468A, A-1181
 Antibiotic L 654040, A-1194
 Antibiotic MBH 001, A-1203
 Antibiotic UK 1, A-1302
 Antibiotic X 14885A, A-1087
 Antibiotic YM 216391, A-1322
 Ariakemicin A, A-1422
 Ascidiacyclamide, A-1456
 Bengazole Z, B-61
 2,6-Benzoxazolediol, B-82
 2-Benzoxazolol, B-83
 Berninamycin A, B-104
 Bistratamide C, B-224
 Bistratamide D, B-225
 Bistratamide E, B-226
 Bistratamide I, B-222
 Boxazomycin A, B-262
 Calcimycin, C-26
 Calyculin A, C-55
 Calyculin C, C-56
 Calyculin J, C-57
 Carboxymycobactins, C-151
 Ceratospogonamide, C-280
 Chivosazole E, C-393
 5-Chloro-2-hydroxy-6-methoxybenzoxazole, C-406
 Cladioxazole, C-488
 Clavosine A, C-516
 Coelibactin, C-561
 Comoramide A, C-585
 Conglobatin, C-602
 Curromycin A, C-815
 Cyclodidemnamide, C-858
 Cycloxazoline, C-919
 Cytoxazone, C-964
 Dendroamide A, D-208
 Dendroamide B, D-209
 Diazonamide B, D-298
 Didmolamide A, D-381
 5,6-Dimethoxy-2(3*H*)-benzoxazolone, D-707
 6,7-Dimethoxy-1(3*H*)-benzoxazolone, D-708
 Discobahamin A, D-842
 Discobahamin B, D-843
 Discokiolide A, D-847
 Discokiolide B, D-848
 Disorazole C₁, D-857
 Disorazole C₂, D-858
 Disorazole F₁, D-859
 Disorazole G₁, D-860
 Disorazole G₂, D-861
 Disorazole G₃, D-862
 Dolastatin E, D-909
 Dolastatin I, D-903
 Epothilone H₁, E-118
 Epothilone H₂, E-119
 Fluvibactin, F-113
 Formobactin, F-130
 Goadsporin, G-136
 Griseoviridin, G-178
 Halfordinol, H-12

Halichondramide, H-15
 Haliclonamide B, H-20
 Haliclonamide D, H-21
 Halishigamide B, H-28
 Halishigamide C, H-29
 Halishigamide D, H-30
 Hennoxazole A, H-112
 Hennoxazole F, H-113
 Icumazole A, I-14
 Indolmycin, I-102
 Kabiramide D, K-2
 Kabiramide H, K-3
 Keramamide B, K-30
 Keramamide D, K-31
 Keramamide E, K-32
 Lajollamycin, L-18
 Leiodelide A, L-84
 Leiodelide B, L-85
 Leucamide A, L-146
 Leucascandrolide A, L-147
 Leupyrin C, L-161
 Lipoxazolidinone A, L-190
 Lipoxazolidinone B, L-191
 Lipoxazolidinone C, L-192
 Lissoclinamide 10, L-203
 Lissoclinamide 1, L-200
 Lissoclinamide 4, L-201
 Lissoclinamide 9, L-202
 Madumycin II, M-35
 Martefragine A, M-109
 Mechercharstatin A, M-142
 Mechercharstatin B, M-143
 Melanoxidin, M-159
 Microcin B17, M-593
 Microcyclamide, M-597
 Muscoride A, M-775
 Mycalolide A, M-779
 Mycalolide D, M-780
 Mycalolide E, M-781
 Nakijinol, N-13
 Neooxazolomycin, N-131
 Neopeltolide, N-133
 Nocobactin NA, N-267
 Nostocyclamide M, N-322
 Nostocyclamide, N-321
 7-[(1,2,3,4,4a,7,8,8a-Octahydro-1,2,4a,5-tetramethyl-1-naphthalenyl)methyl]-6-benzoxazolol, O-60
 Orbiculamide A, O-101
 Ostreogrycin A, O-131
 Oxachelin, O-147
 Oxazolomycin, O-155
 4-(4-Oxazolyl)-3-(hydroxymethyl)-3-buten-2-ol, O-156
 3-(5-Oxazolyl)-1*H*-indole, O-157
 Patellamide A, P-114
 Patellamide B, P-115
 Patellamide C, P-116
 Patellamide D, P-117
 Patellamide F, P-119
 Phenoxan, P-334
 Phorbazole D, P-379
 Phorbaxazole A, P-380
 Photobactin, P-381
 Piscerythoxazole, P-482
 Prepatellamide A, P-620
 Promoinducin, P-640
 Promothiocin A, P-641
 Promothiocin B, P-642
 Radamycin, R-5
 Raocyclamide A, R-16
 Raocyclamide B, R-17
 Rhizopodin, R-81
 Rhizoxin S₁, R-84
 Rhizoxin, R-83
 Routiennocin, R-136
 Serratiochelins, S-258
 Siphonazole, S-323
 Swinhoieamide A, S-652
 Telomestatin, T-66
 Texaline, T-299
 Thiangazole, T-367
 Thioactin, T-378

Thiomycalolide A, T-388
 Thiomycalolide B, T-389
 Thiotipin, T-393
 Transvalencin A, T-445
 Transvalencin Z, T-446
 Trehalamine, T-447
 Ullithiacyclamide, U-10
 Urukthapelstatin A, U-50
 Venturamide A, V-53
 Venturamide B, V-54
 Vibriobactin, V-96
 Wewakazole, W-15

Isoxazole alkaloids

2-Amino-4-(2,5-dihydro-5-oxo-2-isoxazolyl)butanoic acid, A-734
 5-(Aminomethyl)-3(2*H*)-isoxazolone, A-837
 α -Amino-5-oxo-2(5*H*)-isoxazolebutanoic acid, A-849
 Antibiotic B 4317, A-1122
 L-Aspartyl-L-N²-hydroxyaspartyl-D-cycloserine, A-1467
 1,2-Benzisoxazole-3,6-diol, B-72
 4-Isoxazolecarboxylic acid, I-346
 3-Isoxazolidinone, I-347
 4-Isoxazolol, I-348
 Lactivicin, L-8
 Muscaaurin I, M-768
 5-Oxo-2(5*H*)-isoxazoleacetic acid, O-184
 5-Oxo-2(5*H*)-isoxazolepropanoic acid, O-185
 Pseudomonine, P-718
 Tricholomic acid, T-490

Spirobenzoxazoline alkaloids

Aerophobin 1, A-154
 Aerophobin 2, A-155
 Aeroplysinillin, A-156
 Aerothionin, A-158
 Agelocaisarin A₁, A-192
 Agelolin A, A-194
Aplysina archeri Alkaloid, A-272
 Aplysinfulvin, A-1348
 Aplysinamisine I, A-1349
 Aplysinamisine II, A-1350
 Aplysinone A, A-1355
 Aplysinone D, A-1356
 Araplysinin I, A-1378
 Archerine, A-1390
Oceanapia Bromotyrosine alkaloid, B-357
 Caissarine B, C-18
 Calafianin, C-24
 Fistularin 1, F-68
 Fistularin 2, F-69
 Fistularin 3, F-70
 Hemifistularin 3, H-103
 Hexadellin B, H-227
 Hexadellin C, H-228
 Ianthesine B, I-2
 Ianthesine C, I-3
 Psammaplysin A, P-695
 Psammaplysin E, P-696
 Pseudoceratinine C, P-709
 Puralidin L, P-813
 Puraline, P-816
 Purpurealidin B, P-824
Oceanapia Quinolone alkaloid, Q-57

Acyclic thiazole and benzothiazole alkaloids

Aeruginosamide, A-159
 Agrochelin, A-204
 Althiomycin, A-664
 Anguibactin, A-1009
 Antibiotic BE 10988, A-1126
 Antibiotic WS 75624A, A-1315
 Antibiotic WS 75624B, A-1316
 Archazolide A, A-1388
 Archazolide B, A-1389
 Barbaleucamide A, B-21
 Barbamide, B-22

Benzothiazole, B-77
 2,6-Benzothiazole-diol, B-78
 2(3*H*)-Benzothiazolethione, B-79
 2-Benzothiazolol, B-80
 Coelibactin, C-561
 Curacin A, C-807
 Curacin D, C-808
 Cystothiazole E, C-937
 4,5-Dihydro-2-methylthiazole, D-479
 3-(4,5-Dihydro-2-thiazolecarbonyl)-1*H*-indole, D-516
 2,5-Dihydro-2,4,5-trimethylthiazole, D-526
 3,5-Dihydroxy-6-methyl-7-(2-methyl-4-thiazolyl)-6-heptenoic acid, D-628
 5,13-Dihydroxy-4,6,10,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadiene-3,7-dione, D-678
 5,13-Dihydroxy-2,4,6,14-tetramethyl-15-(2-methyl-4-thiazolyl)-10,14-pentadecadien-3-one, D-679
 Dolabellin, D-901
 Dolastatin 10, D-907
 Dolastatin 18, D-908
 Dysideaprolin A, D-968
 Dysideathiazole, D-970
 Dysidenin, D-972
 Dysithiazolamide, D-977
 Epothilone A₁, E-108
 Epothilone A₈, E-109
 Epothilone A, E-110
 Epothilone B, E-111
 Epothilone C₁, E-112
 Epothilone C₃, E-113
 Epothilone C₄, E-114
 Epothilone C, E-115
 Epothilone D₁, E-116
 Epothilone D, E-117
 Epothilone I₁, E-120
 Epothilone I₂, E-121
 Epothilone I₃, E-122
 Epothilone I₄, E-123
 Epothilone I₅, E-124
 Epothilone K, E-125
 Epothilone tetrahydrofuran, E-126
 2-Ethyl-4-methylthiazole, E-261
 Firefly oxyluciferin, F-56
 Goadsporin, G-136
 Herbamide A, H-138
 6-Hydroxy-5-benzothiazoleacetic acid, H-408
 11-Hydroxy-4,12-dimethyl-13-(2-methyl-4-thiazolyl)-8,12-tridecadien-3-one, H-469
 9-Hydroxy-6,10-dimethyl-11-(2-methyl-4-thiazolyl)-6,10-undecadien-3-one, H-470
 5-(2-Hydroxyethyl)-4-methylthiazole, H-495
 Hydroxyhomodolabellin, H-503
 4-Hydroxy-7-[1-hydroxy-2-(methylamino)ethyl]-2(3*H*)-benzothiazolone, H-508
 α -(4-Hydroxy-3-methoxyphenyl)-2-thiazole-methanol, H-569
 2-(2-Hydroxyphenyl)-4-thiazolecarboxylic acid, H-695
 2-(2-Hydroxyphenyl)thiazole, H-694
 5-(1*H*-Indol-3-yl)-2(3*H*)-thiazolethione, I-139
 Kalkitoxin, K-7
 Karnamicin A₁, K-17
 Karnamicin A₂, K-18
 Karnamicin C₂, K-19
 Karnamicin C₃, K-20
 Karnamicin D₂, K-21
 Latrunculin T, L-66
 Lyngbyabellin D, L-358
 Lyngbyabellin F, L-360
 Lyngbyapeptin A, L-362
 Lyngbyapeptin B, L-363
 Lyngbyapeptin C, L-364
 Melithiazole B, M-177
 Melithiazole C, M-178
 Melithiazole F, M-179
 Melithiazole G, M-180
 Melithiazole H, M-181
 Melithiazole I, M-176
 3-Methoxy-4-methyl-5-(2-methyl-4-thiazolyl)-4-pentenoic acid, M-269

2-Methylbenzothiazole, M-400
 2-(2-Methylpropyl)thiazole, M-533
 4-Methylthiazole, M-569
 4-Methyl-5-vinylthiazole, M-585
 Microcin B17, M-593
 Micromide, M-598
 Mirabazole B, M-635
 Mirabazole C, M-636
 Mycothiazole, M-790
 Myxothiazole A, M-807
 Pyochelin I, P-832
 Tantazole A, T-29
 Tantazole B, T-30
 Tantazole F, T-31
 Thiamine, T-366
 Thiangazole, T-367
 Thiazohalostatin, T-370
 5-Thiazoleethanol, T-371
 3-(2-Thiazolyl)-1*H*-indole, T-372
 Thiazostatin A, T-373
 Transvalencin A, T-445
 Tridentatol C, T-508
 Ulbactin A, U-3
 Ulbactin B, U-4
 Ulbactin C, U-5
 Ulbactin D, U-6
 Ulbactin E, U-7
 Violatinctamine, V-143
 Virenamide A, V-146
 Virenamide B, V-147
 Virenamide C, V-148
 Virenamide E, V-149
 Watasemycin A, W-11
 Yersiniabactin, Y-9

Latrunculins

Latrunculin A, L-61
 Latrunculin B, L-62
 Latrunculin C, L-63
 Latrunculin M, L-64
 Latrunculin S, L-65
 Latrunculin T, L-66

Macrocylic thiazole alkaloids

Amythiamicin A, A-959
 Amythiamicin B, A-960
 Amythiamicin C, A-961
 Amythiamicin D, A-962
 Antibiotic A 10947, A-1097
 Antibiotic A 10255B, A-1103
 Antibiotic A 10255C, A-1105
 Antibiotic A 7413D, A-1107
 Antibiotic A 10255E, A-1108
 Antibiotic A 10255F, A-1109
 Antibiotic A 10255H, A-1110
 Antibiotic 5102-2, A-1085
 Antibiotic AB 97, A-1111
 Antibiotic 1401B, A-1120
 Antibiotic CP 41043, A-1152
 Antibiotic CP 41494, A-1153
 Antibiotic GE 37468A, A-1181
 Antibiotic L 13365, A-1193
 Antibiotic Sch 40832, A-1244
 Antibiotic YM 216391, A-1322
 Apratoxin A, A-1366
 Apratoxin C, A-1367
 Ascidiacyclamide, A-1456
 Banyascyclamide B, B-16
 Banyascyclamide C, B-17
 Berninamycin A, B-104
 Bistratamide A, B-223
 Bistratamide C, B-224
 Bistratamide D, B-225
 Bistratamide E, B-226
 Bistratamide I, B-222
 Bistratamide J, B-227
 Ceratospongamide, C-280
 Cleomycin B₂, C-524
 Comoramide A, C-585
 Comoramide B, C-586
 Copsomycin, C-637

Cyclodidemnamide B, C-859
 Cyclodidemnamide, C-858
 Cyclothiazomycin, C-913
 Dendroamide A, D-208
 Dendroamide B, D-209
 Didmolamide A, D-381
 Didmolamide B, D-382
 Dolastatin 1, D-902
 Dolastatin 2, D-904
 Dolastatin 3, D-905
 Dolastatin 4, D-906
 Dolastatin E, D-909
 Dolastatin I, D-903
 Glycothiohexide α , G-126
 Grassyepetolide, G-170
 Guineamide A, G-222
 Guineamide B, G-223
 Haligramide A, H-24
 Halipeptin A, H-26
 Halipeptin C, H-27
 Hectochlorin, H-84
 Jingsimycin, J-40
 Keenamide A, K-27
 Keramamide F, K-33
 Keramamide H, K-34
 Kimorexin A, K-45
 Kororamide, K-83
 Largazole, L-45
 Leinamycin, L-82
 Leucamide A, L-146
 Lissoclinamide 1, L-200
 Lissoclinamide 4, L-201
 Lissoclinamide 9, L-202
 Lissoclinamide 10, L-203
 Lyngbyabellin A, L-355
 Lyngbyabellin B, L-356
 Lyngbyabellin C, L-357
 Lyngbyabellin E, L-359
 Lyngbyabellin G, L-361
 Mayotamide A, M-132
 Mayotamide B, M-133
 Mechercharstatin A, M-142
 Micrococcin M, M-595
 Micrococcin P₁, M-596
 Microcyclamide, M-597
 Neoberninamycin, N-101
 Nocathiacin I, N-266
 Nosiheptide, N-319
 Nostocyclamide M, N-322
 Nostocyclamide, N-321
 Obyanamide, O-13
 Oriamide, O-105
 Pateamine A, P-113
 Patellamide A, P-114
 Patellamide B, P-115
 Patellamide C, P-116
 Patellamide D, P-117
 Patellamide E, P-118
 Patellamide F, P-119
 Patellamide G, P-120
 Patellin 1, P-121
 Patellin 3, P-122
 Patellin 4, P-123
 Patellin 5, P-124
 Patellin 6, P-125
 Prelissoclinamide 2, P-611
 Prepatellamide A, P-620
 Prepatellamide B formate, P-621
 Preulicyclamide, P-627
 Preulithiacyclamide, P-628
 Promoinducin, P-640
 Promothiocin A, P-641
 Promothiocin B, P-642
 Radamycin, R-5
 Raocyclamide A, R-16
 Raocyclamide B, R-17
 Scleritodermin A, S-151
 Siomycin A, S-321
 Siomycin D₁, S-322
 Syriamycin, S-663
 Tawicyclamide A, T-41
 Tawicyclamide B, T-42
 Telomestatin, T-66

Tenuecyclamide A, T-75
 Tenuecyclamide C, T-76
 Thioactin, T-378
 Thiocillin I, T-383
 Thiosporamicin, T-391
 Thiostrepton, T-392
 Thiutipin, T-393
 Thuggacin A, T-397
 Thuggacin B, T-398
 Thuggacin cmc-A, T-400
 Thuggacin C, T-399
 Trichamide, T-477
 Trunkamide A, T-633
 Ulicyclamide, U-9
 Ulithiacyclamide B, U-11
 Ulithiacyclamide E, U-12
 Ulithiacyclamide F, U-13
 Ulithiacyclamide G, U-14
 Ulithiacyclamide, U-10
 Ulongamide B, U-15
 Ulongamide C, U-16
 Ulongamide D, U-17
 Ulongamide E, U-18
 Ulongamide F, U-19
 Urukthapelstatin A, U-50
 Venturamide A, V-53
 Venturamide B, V-54
 Zelvomycin, Z-13

Pyrazine and quinoxaline alkaloids

2-Acetyl-3-methylpyrazine, A-62
 Antibiotic OPC 15161, A-1220
 Arglecin, A-1419
 Argvalin, A-1421
 Aspernigerin, A-1482
 Barrenazine A, B-27
 Batrachamine, B-35
 5-Benzyl-3-hydroxy-2-isopropylpyrazine, B-88
 2,5-Bis(1*H*-indol-3-ylmethyl)pyrazine, B-191
 2,5-Bis(2-methylpropyl)pyrazine, B-203
 3,6-Bis(1-methylpropyl)-2(1*H*)-pyrazinone, B-204
 Botryllazine A, B-258
 Botryllazine B, B-259
 Brasilamide D, B-279
 Bravomicin A, B-286
 Bravomicin D, B-287
 3-Butyl-2,5-dimethylpyrazine, B-431
 6-*tert*-Butyl-1-hydroxy-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, B-433
 Cephalostatin 2, C-261
 Cephalostatin 3, C-262
 Cephalostatin 5, C-263
 Cephalostatin 6, C-264
 Cephalostatin 8, C-266
 Cephalostatin 9, C-267
 Cephalostatin 13, C-268
 Cephalostatin 16, C-269
 Clavulazol A, C-518
 Coelenteramide, C-558
 Coelenteramine, C-559
 Crotonine, C-776
 Diabroticin A, D-241
 2,5-Diethyl-3,6-dimethylpyrazine, D-389
 2,6-Diethyl-3,5-dimethylpyrazine, D-390
 2,3-Diethyl-5-methylpyrazine, D-392
 3,5-Diethyl-2-methylpyrazine, D-393
 6,7-Dihydro-2,3-dimethyl-5*H*-cyclopentapyrazine, D-426
 2,5-Dihydroxy-3-methylpyrazine, D-631
 2,5-Diisopropyl-3-methoxypyrazine, D-699
 2,5-Diisopropylpyrazine, D-700
 2,6-Diisopropylpyrazine, D-701
 2,3-Dimethoxypyrido[2,3-*b*]pyrazine, D-722
 2,5-Dimethyl-3-(3-methyl-2-butenyl)pyrazine, D-754
 2,5-Dimethyl-3-(2-methylbutyl)pyrazine, D-755
 2,5-Dimethyl-3-(2-methylpropyl)pyrazine, D-757
 2,5-Dimethyl-3-(methylthio)pyrazine, D-758
 3,5-Dimethyl-2-pentylpyrazine, D-766
 2,5-Dimethyl-3-propylpyrazine, D-772
 3,5-Dimethyl-2-propylpyrazine, D-773
 2,5-Dimethylpyrazine, D-774
 Dragmacidin A, D-929
 Dragmacidin D, D-930
 Dragmacidin F, D-932
 Echinoserine, E-25
 2-Ethyl-3,5-dimethylpyrazine, E-237
 3-Ethyl-2,5-dimethylpyrazine, E-238
 5-Ethyl-2,3-dimethylpyrazine, E-239
 2-Ethyl-5-methylpyrazine, E-255
 Ethyltrimethylpyrazine, E-273
 Flavacol, F-76
 Flutimide, F-112
 2-[(2-Furanylmethyl)thio]-6-methylpyrazine, F-201
 2-Hexyl-3,5-dimethylpyrazine, H-262
 1-Hydroxy-3,6-bis[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-430
 1-Hydroxy-6-isopropyl-3-(1-methylpropyl)-2(1*H*)-pyrazinone, H-543
 1-Hydroxy-6-isopropyl-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-544
 2-(1-Hydroxy-1-methylpropyl)-3-methoxypyrazine, H-618
 2-(1-Hydroxy-2-methylpropyl)-3-methoxypyrazine, H-619
 1-Hydroxy-3-(2-methylpropyl)-6-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-620
 1-Hydroxy-6-(2-methylpropyl)-3-[2-(methylthio)ethyl]-2(1*H*)-pyrazinone, H-621
 Isopalathazine, I-261
 2-Isopropenyl-5-isopropylpyrazine, I-270
 2-Isopropenyl-6-isopropylpyrazine, I-271
 2-Isopropyl-3,5-dimethoxy-6-methylpyrazine, I-272
 2-Isopropyl-3-methoxy-5-(1-methylpropyl)pyrazine, I-275
 2-Isopropyl-3-methoxy-5-(2-methylpropyl)pyrazine, I-276
 3-Isopropyl-2-methoxy-5-methylpyrazine, I-277
 3-Isopropyl-2-methoxypyrido[2,3-*b*]pyrazine, I-278
 2-Isopropyl-5-(1-methylpropyl)pyrazine, I-281
 2-Isopropyl-5-(2-methylpropyl)pyrazine, I-282
 2-Isopropyl-5-methylpyrazine, I-283
 2-Isopropyl-6-methylpyrazine, I-284
 Isopropylpyrazine, I-285
 Luzopeptin C, L-312
 Maedamine A, M-37
 3-Methoxy-2-methyl-5-(2-methylpropyl)pyrazine, M-268
 3-Methoxy-2-(1-methylpropyl)-5-(2-methylpropyl)pyrazine, M-271
 2-Methoxy-3-(1-methylpropyl)pyrazine, M-272
 2-Methoxy-3-(2-methylpropyl)pyrazine, M-273
 6-(1-Methylethyl)-3-(2-methylpropyl)-2(1*H*)-pyrazinone, M-462
 2-Methyl-3-(2-methylpropyl)pyrazine, M-483
 2-Methyl-5-(1-propenyl)pyrazine, M-523
 2-Methyl-6-(1-propenyl)pyrazine, M-524
 Methylpyrazine, M-536
Cypridin Oxyluciferin, O-222
Watasenia Oxyluciferin, O-223
 Palythazine, P-41
 Pedatisectine D, P-153
 Pedatisectine E, P-154
 Pedatisectine F, P-155
 Piperazinomycin, P-444
 Piperazirum, P-445
 Pulcherrimic acid, P-788
 Quinazomycin, Q-15
 Quinomycin A, Q-58
 Quinomycin B₀, Q-59
 Quinomycin B, Q-60
 Quinomycin C, Q-61
 Quinomycin D, Q-62
 Quinomycin E, Q-63
 Quinoxapeptin A, Q-64
 Ritterazine A, R-101
 Ritterazine B, R-102
 Ritterazine C, R-103
 Ritterazine E, R-104
 Ritterazine J, R-105

Ritterazine N, R-106
 Ritterazine P, R-107
 Ritterazine R, R-108
 Ritterazine U, R-109
 Ritterazine V, R-100
 Ritterazine W, R-110
 Ritterazine Z, R-111
 Sandramycin, S-50
 Sclerominol, S-152
 Septorine, S-249
 Stizolamine, S-553
 Terezine A, T-84
 Terezine B, T-85
 Tetramethylpyrazine, T-284
 Trimethylpyrazine, T-613

Benzo[1,2-b:3,4-b']dipyrroles

Antibiotic PDE I, A-1222
 Antibiotic PDE II, A-1223
 Bizelesin, B-229
 Duocarmycin A, D-952
 Duocarmycin C₂, D-954
 Duocarmycin SA, D-956
 Rachelmycin, R-4
 Yatakemycin, Y-7

Pyrrolo[1,2-a]pyrazines

Agesamide A, A-195
 Antibiotic MPC 1001F, A-1207
 Antibiotic SF 2771, A-1261
 Cyclo[2-(1,1-dimethyl-2-propenyl)tryptophyl]-
 prolyl, C-861
 Cyclooroidin, C-887
 Macrophominol, M-19
 Makaluvic acid A, M-56
 Oxocyclostylidol, O-173
 Peramine, P-242
 Pyrrolo[1,2-a]pyrazine-3,6(4*H*,7*H*)-dione, P-968
 Tryprostatin A, T-638

Morpholines

Agropine, A-207
 Bassiatin, B-29
 Chelonin A, C-361
 Chelonin C, C-363
 Convolutamine E, C-627
 Ergosecalinine, E-168
 Metacytofilin, M-231
 Oxathinecarboxylic acid, O-151
 2*H*-1,4-Oxazine-3-carboxylic acid, O-152
 Oxazininine 1, O-153
 Oxazininine 3, O-154
 Polygonapholine, P-561

Pyrimidines

Aberiamine, A-13
 Allithiamine, A-627
 4-Amino-3,6-dihydroxy-1*H*-pyrazolo[3,4-*d*]pyri-
 midine, A-755
 4-Amino-2-hydroxy-5-(hydroxymethyl)pyrimi-
 dine, A-795
 4-Amino-2-hydroxy-5-methylpyrimidine, A-803
 2-[(4-Aminophenyl)amino]pyrimidine, A-863
 2-Amino-4-pyrimidinecarboxylic acid, A-903
 Antibiotic TAN 1057A, A-1282
 Aplicyanin A, A-1343
 Barbitol, B-23
 Boxazomycin A, B-262
 Crambescidin, C-99
 Crambescidin A, C-721
 Crambescidin B, C-722
 Crambescidin C1, C-723
 Crambescidin C2, C-724
 Cytomycin, C-960
 Cytosine, C-961
 Dehydrocrambine A, D-155
 4,5-Diamino-2,6-dihydroxypyrimidine,
 D-254

1,3-Dihydrocyclopentapyrimidin-2-one, D-416
 7,8-Dihydro-3-methylpyrrolo[1,2-*a*]pyrimidin-
 2(6*H*)-one, D-478
 Dihydro-2,4(1*H*,3*H*)-pyrimidinedione, D-497
 2,3-Dihydro-2-thioxo-4(1*H*)-pyrimidinone,
 D-518
 2,6-Dihydroxy-4-pyrimidinecarboxylic acid,
 D-669
 1,3-Dimethyl-2,4(1*H*,3*H*)-pyrimidinedione, D-777
 5-Fluoro-2,4-dihydroxypyrimidine, F-104
 Formycin A, F-131
 Formycin B, F-132
 Glusun I, G-105
 Heteromine H, H-159
 Hortesin, H-360
 2-Hydroxy-4-mercaptopyrimidine, H-556
 Hymeniacidine, H-761
 Hyrtinadine A, H-779
 Leucosolenamine A, L-157
 Manzacidin B, M-90
 Meridianin A, M-205
 5-Methyl-2,4,6(1*H*,3*H*,5*H*)-pyrimidinetrione,
 M-540
 Millettonine, M-615
 Muraymycin C₁, M-737
 Nikkomycin Z, N-218
 Psammopemmin A, P-697
 Sparsomycin, S-384
 Stendomycidine, S-527
 3,4,5,6-Tetrahydro-6-hydroxymethyl-3,6-di-
 methyl-4-pyrimidinecarboxylic acid, T-183
 1,4,5,6-Tetrahydro-5-hydroxy-2-methyl-4(6)-pyr-
 imidinecarboxylic acid, T-185
 Thiamine, T-366
 Thymine, T-401
 Tubercatidine, T-657
 Uracil, U-42

Ptilocaulins

1,3*a*-Didehydro-8-hydroxyptilocaulin, D-375
 Isoptilocaulin, I-291
 Mirabilin B, M-638
 Mirabilin C, M-639
 Mirabilin D, M-640
 Mirabilin E, M-641
 Mirabilin F, M-642
 Mirabilin G, M-643
 Netamine A, N-174
 Netamine B, N-175
 Netamine C, N-176
 Netamine E, N-177
 Netamine F, N-178
 Netamine G, N-179
 Ptilocaulin, P-773

Triazaacenaphthylene alkaloids

Batzelladine A, B-37
 Batzelladine B, B-38
 Batzelladine C, B-39
 Batzelladine D, B-40
 Batzelladine E, B-41
 Batzelladine F, B-42
 Batzelladine G, B-43
 Batzelladine H, B-44
 Batzelladine J, B-45
 Batzelladine K, B-46
 Batzelladine L, B-47
 Batzelladine M, B-48
 Crambescidin 359, C-715
 Crambescidin 431, C-716
 Crambescidin 816, C-717
 Crambescidin 826, C-718
 Crambescidin 830, C-719
 Crambescidin 844, C-720
 Crambidine, C-725
 Dehydrobatzelladine C, D-152
 Fromiamycalin, F-158
 Neofolitispatate 1, N-115
 Ptilomycalin A, P-774
 Ptilomycalin D, P-775

Tetrodotoxins

4,9-Anhydrotetrodotoxin, A-1035
 Chiriquitoxin, C-390
 Tetrodonic acid, T-297
 Tetrodotoxin, T-298
 5,6,11-Trideoxytetrodotoxin, T-510

Phenazine alkaloids

Aeruginosin B, A-162
 Aestivophoenin A, A-164
 Aestivophoenin B, A-165
 8-Amino-1-hydroxyphenazine, A-805
 Antibiotic DOB 41, A-1160
 Antibiotic PD 116152, A-1221
 Antibiotic SB 212305, A-1239
 Antifungin, A-1326
 Benthocyanin A, B-64
 Benthocyanin C, B-65
 Benthophoenin, B-66
 2-Chloro-1,6-phenazinediol, C-415
 2,3-Dihydroxy-1-phenazinecarboxylic acid,
 D-640
 2,6-Dihydroxy-1-phenazinecarboxylic acid,
 D-641
 2,9-Dihydroxy-1-phenazinecarboxylic acid,
 D-642
 3,6-Dihydroxy-1-phenazinecarboxylic acid,
 D-643
 2,7-Dimethyl-5,10-[2]butenophenazine-11,14-
 dione, D-744
 Diphenazithionin, D-814
 Endophenazine B, E-79
 Esmeraldine A, E-222
 Esmeraldine B, E-223
 3,4,6,7,8,9-Hexahydrophenazine, H-240
 6-(1-Hydroxyethyl)-1-phenazinecarboxylic acid,
 H-496
 9-Hydroxy-6-hydroxymethyl-1-phenazinecar-
 boxylic acid, H-514
 2-Hydroxy-1-phenazinecarboxylic acid, H-661
 4-Hydroxy-1-phenazinecarboxylic acid, H-662
 9-Hydroxy-1-phenazinecarboxylic acid, H-664
 1-(4-Hydroxyphenyl)-3*H*-benzo[*a*]pyrrolo[2,3-
c]phenazine, H-674
 Lavanducyanin, L-76
 Lomofungin, L-241
 6-(3-Methyl-2-butenyl)-1-phenazinecarboxylic
 acid, M-417
 9-(3-Methyl-2-butenyl)-1-phenazinecarboxylic
 acid, M-418
 1,1'-Methylenebis[2-phenazinol], M-440
 1,2,3,4,6,7,8,9-Octahydrophenazine, O-54
 1-Phenazinecarboxylic acid, P-319
 1,6-Phenazinedicarboxylic acid, P-320
 1,6-Phenazinediol, P-321
 1,8-Phenazinediol, P-322
 1,9-Phenazinediol, P-323
 2,3-Phenazinediol, P-324
 2,3,7-Phenazinetriol, P-326
 1-Phenazinol, P-327
 2-Phenazinol, P-328
 Phenazinomycin, P-329
 Phenazostatin A, P-330
 Phenazostatin B, P-331
 Phenazostatin C, P-332
 Piricularin, P-478
 Pyocyanine, P-833
 Ruticin, R-172
 Senacarcin A, S-231
 2,3,9-Trihydroxy-1-phenazinecarboxylic acid,
 T-574
 2,3,7-Trihydroxy-1,6-phenazinedicarboxylic
 acid, T-575
 Tubermycin A, T-660

Phenoxazine alkaloids (other than actinomycins)

2-Amino-8-benzoyl-6-hydroxy-3*H*-phenoxazin-
 3-one, A-707

7-Amino-2,8-bis(2,4-dihydroxyphenyl)-3*H*-phenoxazin-3-one, A-711
 2-Amino-3-oxo-3*H*-phenoxazine-1-carboxylic acid, A-852
 2-Amino-3-oxo-3*H*-phenoxazine-8-carboxylic acid, A-853
 2-Amino-3*H*-phenoxazin-3-one, A-862
 Antibiotic B 1625FA_{2p,1}, A-1124
 Chandrananimycin C, C-345
 Cinnabarine, C-452
 Grixazone B, G-180
 1-Methoxy-8-hydroxymethyl-3*H*-phenoxazin-3-one, M-253
 2-(Methylthio)-3*H*-phenoxazin-3-one, M-576
 Michigazone, M-590
 Murayaanthraquinone, M-735
 Orcein, O-102
 3*H*-Phenoxazin-3-one, P-335
 Pycnosanguin, P-831
 $\alpha,\alpha',2$ -Triamino- $\gamma,\gamma',3$ -trioxo-3*H*-phenoxazine-1,9-dibutanoic acid, T-453
 1,2,4-Trimethoxy-3*H*-phenoxazin-3-one, T-599
 Xanthicin, X-4

Simple pyrrole alkaloids

3-Acetyl-4-(hydroxymethyl)pyrrole, A-57
 3-Acetyl-4-methylpyrrole, A-71
 2-Acetylpyrrole, A-74
 Agelongine, A-193
 3-Alkyl-1*H*-pyrrole-2-carboxaldehydes, A-623
 3-(2-Amino-3-chlorophenyl)-4-chloro-1*H*-pyrrole, A-725
 α -Amino-1*H*-pyrrole-1-hexanoic acid, A-904
 Antibiotic TAN 876B, A-1283
 Arcyriarubin A, A-1394
 4-Bromo-2-(guanidinomethyl)pyrrole, B-319
 3'-Bromo-4-methoxy-2,2'-bipyrrole-5-carboxaldehyde, B-338
 5'-Bromo-4-methoxy-2,2'-bipyrrole-5-carboxaldehyde, B-339
 N²-(4-Bromo-1*H*-pyrrole-2-carbonyl)arginine, B-345
 N²-(4-Bromo-1*H*-pyrrole-2-carbonyl)homoarginine, B-346
 N⁶-(4-Bromo-1*H*-pyrrole-2-carbonyl)lysine, B-347
 4-Bromo-1*H*-pyrrole-2-carboxylic acid, B-348
 3-Bromo-1*H*-pyrrole-2,5-dione, B-350
 Buprestin A, B-415
 Buprestin B, B-416
 Butylcycloheptylprodigiosine, B-427
 Cannabiloid B, C-82
 Celastramycin A, C-231
 Crambesin A, C-721
 2,3-Dibromo-4-(3-bromo-2-nitrophenyl)-1*H*-pyrrole, D-305
 2-(3,5-Dibromo-2-hydroxyphenyl)-3,4,5-tribromopyrrole, D-324
 2,5-Dibromo-1*H*-pyrrole-3-carboxylic acid, D-338
 4,5-Dibromo-1*H*-pyrrole-2-carboxylic acid, D-339
 2,3-Dibromo-1*H*-pyrrole, D-337
 3,4-Dibromo-1*H*-pyrrole-2,5-dione, D-341
 2,3-Dichloro-4-nitro-1*H*-pyrrole, D-356
 1,3-Dihydro-3-hydroxy-2*H*-pyrrol-2-one, D-459
 1-(2,3-Dihydro-2-oxo-3-furanyl)-5-(hydroxymethyl)-1*H*-pyrrole-2-carboxaldehyde, D-483
 4-(3,7-Dimethyl-2,6-octadienyl)-1*H*-pyrrole-2-carboxylic acid, D-761
 2,3-Diphenyl-1*H*-pyrrole, D-817
 Ebracteolatinoside A, E-2
 4-(2-Formyl-5-hydroxymethyl-1*H*-pyrrol-1-yl)-butanoic acid, F-137
 5-Formyl-1*H*-pyrrole-2-carboxylic acid, F-142
 1-(2-Furanylmethyl)-1*H*-pyrrole, F-200
 2,3,3',4,4',5,5'-Heptachloro-1'-methyl-1,2'-bi-1*H*-pyrrole, H-115
 3,3',4,4',5,5'-Hexabromo-2,2'-bi-1*H*-pyrrole, H-212
 1-[(5-Hexyltetrahydro-2-furanyl)acetyl]-1*H*-pyrrole, H-269

4-Hydroxy-1*H*-pyrrole-2-carboxylic acid, H-714
 Indigotiscoumarin A, I-71
 Ircinamine B, I-172
 Laughine, L-71
 Magnolamide, M-44
 Marinamide, M-104
 4-Methoxy-2,2'-bipyrrole-5-carboxaldehyde, M-244
 3-Methyl-4-phenyl-1*H*-pyrrole, M-514
 2-Methyl-1-(1*H*-pyrrol-1-yl)-8-decene-1,3-dione, M-553
 Neopyrrolomycin, N-136
 Peyoglunol, P-296
 Peyonine, P-298
 N-(3-Phenylpropanoyl)pyrrole, P-351
 Phorbazole D, P-379
 PUR 1, P-808
 Pyoluteorin, P-834
 1*H*-Pyrrole-2-carboxaldehyde, P-942
 1*H*-Pyrrole-3-carboximidamide, P-943
 1*H*-Pyrrole-2-carboxylic acid, P-944
 1*H*-Pyrrole-2,3-dicarboxylic acid, P-945
 1*H*-Pyrrole-2,5-dicarboxylic acid, P-946
 Pyrrolnitrin, P-961
 Pyrrolomycin B, P-963
 Pyrrolomycin C, P-964
 Pyrrolomycin E, P-965
 Pyrrolomycin G, P-967
 Sarmentamide A, S-86
 Solsodimine A, S-358
 Solsodimine B, S-359
 3,3',4,4'-Tetrabromo-5,5'-dichloro-1,1'-dimethyl-2,2'-bi-1*H*-pyrrole, T-111
 2,3,4,5-Tetrabromo-1*H*-pyrrole, T-118
 3,4,5-Tribromo-1*H*-pyrrole-2-carboxylic acid, T-475
 2,3,4-Tribromo-1*H*-pyrrole, T-474

Miscellaneous pyrrole alkaloids

α -Amino-5-[[[(2-aminocarbonyl-1-carboxyethyl)amino]carbonyl]-2-pyrrolepropanoic acid, A-695
 Anthelvencin A, A-1077
 Antibiotic BE 55051, A-1133
 Antibiotic 610MF, A-1206
 Antibiotic R 39MF, A-1230
 Antibiotic R 39SF, A-1231
 Antibiotic TAN 1307, A-1277
 Antibiotic TAN 876A, A-1280
 Antibiotic UCH 15A, A-1300
 Antibiotic X 14885A, A-1087
 Asterin A, A-1511
 Asterinin A, A-1512
 Asterinin C, A-1513
 Astin J, A-1523
 Axinellamine B, A-1575
 4-Benzyl-1,4-dihydro-3-oxo-3*H*-pyrrolo[2,1-*c*]-[1,4]oxazine-6-carboxaldehyde, B-86
 3,4-Bis(4-hydroxyphenyl)-1*H*-pyrrole-2,5-dicarboxylic acid, B-188
 Brachystemidine A, B-267
 Brachystemidine E, B-268
 Brachystemidine F, B-269
 Brachystemidine G, B-270
 Bufopyramide, B-391
 Cafamycin, C-13
 1-[Caffeoyloxy(carboxy)methoxy]-1*H*-pyrrole-2,3,5-tricarboxylic acid, C-16
 Calcimycin, C-26
 Chinese bittersweet alkaloid I, C-383
 Chinese bittersweet alkaloid II, C-384
 Chlorobiocin, C-395
 Chromoviridan, C-426
 Chromoxymycin, C-427
 Cochinmicin I, C-545
 Colubricidin A, C-579
 Coumermycin A₁, C-711
 Coumermycin A₂, C-712
 Cyclocinamide A, C-854
 Cycloprodigiosin, C-897
 Decatromicidin A, D-126
 Desmodimine, D-237
 Didemnimide A, D-379
 5*H*,10*H*-Dipyrrolo[1,2-*a*:1',2'-*a'*]pyrazine-5,10-dione, D-833
 Dispyrin, D-865
 Distamycin A, D-866
 Ethylcycloonylprodigiosin, E-235
 Eudistomin A, E-285
 Funebrial, F-194
 Funebrine, F-195
 Glaciapyrrole A, G-83
 Glaciapyrrole B, G-84
 Glaciapyrrole C, G-85
 Glycerinopyrin, G-111
 Himanimide A, H-278
 Hormamycin, H-358
 5-Hydroxy-3,4'-dimethyl-[1,2'-bipyrrole]-2,5'(2'*H*,5*H*)-dione, H-467
 3-(4-Hydroxyphenyl)-4-(2-methylpropyl)-1*H*-pyrrole-2,5-dione, H-684
 Hyperaspine, H-769
 Indanomycin, I-64
 Isochrysohermidin, I-206
 Isonargenicin A₁, I-254
 Keronopsin B₁, K-35
 Kikumycin A, K-43
 Lamellarin Q, L-25
 Leupyrrin C, L-161
 Lycogalic acid, L-329
 Manzacidin B, M-90
 Marinopyrrole A, M-107
 Metacycloprodigiosin, M-230
 4-Methoxy-5-[(3-methoxy-5-pyrrol-2-yl)-2*H*-pyrrol-2-ylidene]methyl]-2,2'-bipyrrole, M-257
 2-[[[3-Methoxy-4-(1*H*-pyrrol-1-yl)-2*H*-pyrrol-2-ylidene]methyl]-5-methyl-1-pentyl-1*H*-pyrrole, M-286
 Methylcyclodecylprodigiosin, M-429
 Methylcyclooctylprodigiosin, M-430
 Mycaleoxime, M-777
 Nargenicin B₁, N-48
 Nargenicin B₂, N-49
 Nargenicin B₃, N-50
 Neolamellarin A, N-121
 Netropsin, N-180
 Ningalin B, N-221
 Nonylprodigiosin, N-285
 Opuntine B, O-100
 Orosomycin, O-124
 PD 125375, P-146
 Peripentoinine A, P-263
 Peripentoinine C, P-264
 Polycitone A, P-556
 Porphobilinogen, P-574
 Prodigiosene, P-635
 Prodigiosin, P-636
 Proviolacein, P-675
 Pseudoceratidine, P-706
 Pyranonigrin B, P-881
 Pyranonigrin D, P-882
 Pyrrolosporin A, P-969
 Pyrromycin A, P-973
 Pyrroxamycin, P-974
 Reductiline, R-33
 Roseophilin, R-129
 Routiennocin, R-136
 Rumbrin, R-160
 Rynania diterpene ester C₂, R-174
 Ryanodine, R-175
 Speranberculatine A, S-390
 Speranskilatine A, S-393
 Stawamycin, S-499
 Storniamide D, S-556
 Streptorubin B, S-572
 Stylissadine A, S-610
 Tambjamine A, T-22
 Thiazohalostatatin, T-370
 Trikentramine, T-593
 Undecylprodigimine, U-31
 Vanillobiocin, V-18
 Verrucarin F, V-81
 Verrucarin G, V-82
 Vitamycin A, V-163
 Wallemia A, W-4

Walleimia C, W-5

Polyamine alkaloids PA3, PA33 and homologues

3,5-Acarnidine, A-26
 1-Amino-20-guanidino-4,8,12,16-tetrazaicosane, A-780
 1-Amino-16-guanidino-4,8,12-triazahexadecane, A-781
N-(3-Aminopropyl)-1,5-pentanediamine, A-897
 Argiopinin IV, A-1416
N,N'-Bis(3-aminopropyl)-1,3-propanediamine, B-154
 1,12-Diamino-4,8-diazadodecane, D-249
 3,3'-Diaminodipropylamine, D-255
 1,20-Diamino-4,8,12,16-tetrazaeicosane, D-283
 1,20-Diamino-4,8,12,17-tetrazaeicosane, D-284
 1,20-Diamino-4,8,13,17-tetrazaeicosane, D-285
 1,16-Diamino-4,8,12-triazahexadecane, D-288
 Fluvibactin, F-113
 Motuporamine A, M-714
 Motuporamine B, M-715
 Motuporamine C, M-716
 Muraymycin C₁, M-737
 4,8,12,16,20-Pentaazatricosane-1,23-diamine, P-199
 4,8,12,16-Tetraazanonadecane-1,19-diamine, T-107
 4,8,12-Triazapentadecane-1,15-diamine, T-456
 Tris(3-aminopropyl)amine, T-622
 Vibriobactin, V-96

Polyamine alkaloids PA4 (putrescines)

3,5-Acarnidine, A-26
 Aeroplysinillin, A-156
 Aerothionin, A-158
 Aeruginosin 98A, A-160
 Aeruginosin 205A, A-161
Aplysina archeri Alkaloid, A-272
 (4-Aminobutyl)guanidine, A-719
 (4-Aminobutyl)urea, A-720
 Birnbaum A, B-148
N,N'-Bis(4-guanidinobutyl)oxamide, B-178
 Bistelletadine A, B-219
 Bistelletadine B, B-220
Oceanapia Bromotyrosine alkaloid, B-357
 1,4-Butanediamine, B-422
 Calafianine, C-24
 Caracasandiamide, C-108
 Cepaciachelin, C-246
 Clavamine, C-512
 Cleomycin B₂, C-524
 Cyclofoveoglin, C-864
 1,4-Diguanidinobutane, D-400
 Erylusidine, E-196
 Erylusine, E-197
 Eusynstyelamide, E-303
Aglaia dasyclada Flavagline compound 1, F-77
Aglaia dasyclada Flavagline compound 2, F-78
 Fontaineine, F-124
Aglaia dasyclada Forbagline-like compound, F-128
 Foveoglin A, F-144
 Gigantamide A, G-74
 Grandiamide A, G-156
N-(4-Guanidinobutyl)-2-(4-hydroxyphenyl)-2-oxoacetamide, G-202
 Hordatine B, H-356
N-[4-(4-Hydroxybenzoylamino)butyl]-1,3-dimethylmazine-6-carboxamide, H-410
 Microcin SF608, M-594
 Myxochelin A, M-806
 Ornibactin C4, O-119
 Ornibactin C6, O-120
 Ornibactin F, O-121
 Phomoenamide, P-375
 Photobactin, P-381

Protochelin, P-661
 Puralidin L, P-813
 Puralidin O, P-815
 Putreanine, P-826
 Pyramidaglaine A, P-878
 Rexostatine, R-69
 Spermic acid, S-397
 Spherophysine, S-406
N,N,N',N'-Tetramethyl-1,4-butanediamine, T-281
Perophora viridis 1,2,3-Thiocane glycoside, T-382

Polyamine alkaloids PA5 (cadaverines)

(5-Aminopentyl)guanidine, A-859
N-(3-Aminopropyl)-1,5-pentanediamine, A-897
 Antibiotic IC 202A, A-1185
 Aplysinamisine II, A-1350
 Argiopinin IV, A-1416
 Arthrobactin, A-1451
 Cathestatin C, C-209
 Clavamine, C-512
 1,5-Diguanidinopentane, D-401
 α -(Dimethylamino)-3,5-diiodo-*N*-[[3-(3-iodo-4-methoxyphenyl)-2-(methylamino)-1-oxopropyl]amino]pentyl]-4-methoxybenzenepropanamide, D-729
 Ferrioxamine H, F-44
 Flavolipids, F-83
 Laughine, L-71
 Nephilatoxin 9, N-159
 Nephilatoxin 11, N-161
Nephila clavata Neurotoxin 4 JSTX-4, N-184
 1,5-Pentanediamine, P-229
 Tenacibactin B, T-69
 Terragine C, T-101

Polyamine alkaloids PA43 (spermidines)

Agrobactin, A-203
 Alkaloid LBZ, A-553
N-(4-Aminobutyl)-*N*-(3-aminopropyl)-1,4-butanediamine, A-718
 4-Amino-*N,N,N*-tris(3-aminopropyl)-1-butaniminium, A-926
 Antibiotic WF 14861A, A-1309
 Bagougeramine A, B-3
 Bagougeramine B, B-4
 Bis(4-aminobutyl)bis(3-aminopropyl)ammonium(1+), B-149
N,N-Bis(3-aminopropyl)-1,4-butanediamine, B-152
 Cadabicine, C-4
 Cannabisativine, C-83
 Capparisine, C-97
 Chisitine 1, C-391
 Chisitine 2, C-392
 Cinodine I, C-459
 Cinodine II, C-460
 Codonocarpine, C-556
 Crambescidin 816, C-717
 Crambescidin 830, C-719
 Crambescidin 844, C-720
 Crambidine, C-725
 Cyclocelabenzene, C-853
 Galantin I, G-6
 Galantin II, G-7
 Glycocinnaspermicidin A, G-117
 Glycerin B, G-132
 Hirudonine, H-305
 Hispidospermidine, H-306
 7-Hydroxypleurocorine, H-700
 Inandenin-10-one, I-56
 Inandenin-11-one, I-57
 Inandenin-12-one, I-58
 Inandenin-13-one, I-59
 Isocyclocelabenzene, I-222
 Loesenerine, L-230
 Lunaridine, L-292

Lunarine, L-293
 Monodontamide B, M-689
 Monodontamide D, M-690
 Numismine, N-329
 Palustrine, P-40
 Perimargine, P-259
 Periphylline, P-265
 Petrobactin, P-283
 Pleurostyline, P-523
 Pseudoceratidine, P-706
 Ptilomycalin A, P-774
 Ptilomycalin D, P-775
 Spergualin, S-394
 Spermidine, S-398
 Squalamine, S-485
 Tenuilobine, T-79
 Trypanothione, T-636

Polyamine alkaloids PA44 (homospermidines)

N-(4-Aminobutyl)-*N*-(3-aminopropyl)-1,4-butanediamine, A-718
 Bis(4-aminobutyl)bis(3-aminopropyl)ammonium(1+), B-149
N,N'-Bis(4-aminobutyl)-1,4-butanediamine, B-150
 Circinamide, C-461
 4,4'-Diaminodibutylamine, D-252
 1,24-Diamino-5,10,15,20-tetraazatetracosane, D-286
 1,19-Diamino-6,11,16-triazanonadecane, D-290
 Solamine, S-340

Polyamine alkaloids PA343 (spermines)

Agel 452, A-176
 Bleomycin A₆, B-236
 Budmunchiamine A, B-383
 Budmunchiamine B, B-384
 Budmunchiamine L1, B-385
 Budmunchiamine L2, B-386
 Budmunchiamine L4, B-387
 Budmunchiamine L5, B-388
 Budmunchiamine L5, B-389
 1,20-Diamino-4,8,12,17-tetrazaeicosane, D-284
 1,20-Diamino-4,8,13,17-tetrazaeicosane, D-285
 1,16-Diamino-4,8,13-triazahexadecane, D-289
 Het₃₈₉, H-155
 Homaline, H-324
 Hopromalinol, H-353
 Hopromine, H-354
 Hoprominol, H-355
 Nephilatoxin 1, N-152
 Nephilatoxin 565, N-165
Nephila clavata Neurotoxin 1 JSTX-1, N-182
 Penaramides, P-174
 Pithecolobine, P-486
 Protomethine, P-665
 Schweinine, S-147
 Sinulamide, S-318
 Spermine, S-399
 Tenuilobine, T-79

Other 3-residue polyamine alkaloids

Antibiotic IC 202B, A-1186
 Argiopine, A-1414
 Argiotoxin 659, A-1417
N,N'-Bis(4-aminobutyl)-1,4-butanediamine, B-150
N,N'-Bis(3-aminopropyl)-1,2-ethanediamine, B-153
 CNS 2103, C-535
 1,12-Diamino-4,8-diazadodecane, D-249
 1,13-Diamino-4,9-diazatridecane, D-250
 1,13-Diamino-5,9-diazatridecane, D-251
 1,17-Diamino-4,9,13-triazahexadecane, D-287
 1,18-Diamino-5,9,14-triazaoctadecane, D-291
 Glycerin A, G-131
 Glycerin C, G-133

Het₄₀₃, H-156
 Joramine, J-46
 Lipogrammistin A, L-185
 Nephilatoxin 2, N-153
 Nephilatoxin 10, N-160
 Nephilatoxin 12, N-162
 Nephilatoxin 473, N-163
 Nephilatoxin 501, N-164
 Nephilatoxin 594, N-166
Nephila maculata Neurotoxin 3, N-181
 Philanthotoxin 433, P-358
 Tenacibactin C, T-70
 Tenacibactin D, T-71
 Terragine A, T-99
 Terragine D, T-102
 Toxin MG 30, T-441

Polyamine alkaloids with more than 3 residues

Agel 452, A-176
 Alcaligin, A-250
 1-Amino-20-guanidino-4,8,12,16-tetraazaicosane, A-780
 1-Amino-16-guanidino-4,8,12-triazahexadecane, A-781
 Antibiotic IC 202C, A-1187
 Argiopinine II, A-1415
 Argiotoxin 673, A-1418
N,N'-Bis(3-aminopropyl)-1,3-propanediamine, B-154
 Chaenorhine, C-317
 Chaenorpine, C-318
 CNS 2104, C-536
 Deferoxamine, D-142
 1,20-Diamino-4,8,12,16-tetrazaeicosane, D-283
 1,20-Diamino-4,8,12,17-tetrazaeicosane, D-284
 1,20-Diamino-4,8,13,17-tetrazaeicosane, D-285
 1,24-Diamino-5,10,15,20-tetraazatetracosane, D-286
 1,16-Diamino-4,8,12-triazahexadecane, D-288
 1,16-Diamino-4,8,13-triazahexadecane, D-289
 1,19-Diamino-6,11,16-triazanonadecane, D-290
 Diatom polyamines, D-293
 Nephilatoxin 1, N-152
 Nephilatoxin 3, N-154
 Nephilatoxin 4, N-155
 Nephilatoxin 5, N-156
 Nephilatoxin 6, N-157
 Nephilatoxin 7, N-158
 Nephilatoxin 565, N-165
 Nephilatoxin 643, N-167
Nephila clavata Neurotoxin 2 JSTX-2, N-183
 Penaramides, P-174
 4,8,12,16,20-Pentaazatricosane-1,23-diamine, P-199
 Pseudoargiopinine III, P-701
 4,8,12,16-Tetraazonadecane-1,19-diamine, T-107
 4,8,12-Triazapentadecane-1,15-diamine, T-456

Ansa-peptide alkaloids

Abyssenine B, A-18
 Adouetine X, A-149
 Adouetine Y, A-150
 Adouetine Z, A-151
 Americine, A-690
 1-(2-Amino-1-oxopropyl)-2,3,3a,13a,14,15,16,18a-octahydro-8-methoxy-5,9-metheno-9*H*-dipyrrolo[3,2-*b*:1',2'-*e*][1,5,8]oxadiazacyclopentadecine-13,18(1*H*,12*H*)-dione, A-855
 Amphibine A, A-935
 Amphibine B, A-936
 Amphibine C, A-937
 Amphibine D, A-938
 Amphibine E, A-939
 Amphibine F, A-940
 Amphibine G, A-941
 Amphibine H, A-942
 Anordianine, A-1072
 Aralionine A, A-1375

Canthiumine, C-91
 Ceanothine A, C-226
 Ceanothine C, C-227
 Ceanothine D, C-228
 Ceanothine E, C-229
 Condaline A, C-600
 Daechuine S3, D-7
 Daechuine S7, D-8
 Daechuine S8-1, D-9
 Debenzoylaralioine A, D-100
 1-[2-(Dimethylamino)-1-oxo-3-phenylpropyl]-2,3,3a,13a,14,15,16,18a-octahydro-8-methoxy-5,9-metheno-9*H*-dipyrrolo[3,2-*b*:1',2'-*e*][1,5,8]oxadiazacyclopentadecine-13,18(1*H*,12*H*)-dione, D-737
 Discarene C, D-834
 Discarene D, D-835
 Discarine B, D-836
 Discarine C, D-837
 Discarine D, D-838
 Discarine E, D-839
 Discarine M, D-840
 Discarine N, D-841
 Franganine, F-149
 Franguloline, F-150
 Frangulanine, F-151
 Hemsine A, H-104
 Hemsine B, H-105
 Hemsine C, H-106
 Hemsine D, H-107
 Hymenocardine, H-763
 Hysodricanine A, H-783
 Hysodricanine B, H-784
 Hysodricanine C, H-785
 Integerrenine, I-154
 Integerressine, I-155
 Integerrine, I-156
 Jubanine A, J-51
 Jubanine B, J-52
 Jubanine C, J-53
 Lasiodine A, L-48
 Lasiodine B, L-49
 Lotusanine B, L-266
 Lotusine A, L-267
 Lotusine B, L-268
 Lotusine C, L-269
 Lotusine E, L-270
 Lotusine G, L-271
 Mauritine A, M-125
 Mauritine B, M-126
 Mauritine C, M-127
 Mauritine D, M-128
 Mauritine H, M-129
 Melofoline, M-186
 Melonovine A, M-189
 Melonovine B, M-190
 Mucronine A, M-718
 Mucronine C, M-719
 Mucronine D, M-720
 Mucronine J, M-721
 Myrianthine A, M-793
 Myrianthine C, M-794
 Nummularine C, N-330
 Nummularine E, N-331
 Nummularine F, N-332
 Nummularine G, N-333
 Nummularine K, N-334
 Nummularine M, N-335
 Nummularine N, N-336
 Nummularine P, N-337
 Nummularine R, N-338
 Paliurine B, P-23
 Paliurine C, P-24
 Paliurine E, P-25
 Paliurine F, P-26
 Paliurine G, P-27
 Paliurine H, P-28
 Paliurine I, P-29
 Ramosine A, R-14
 Rugosanine B, R-155
 Sanjoinine G₁, S-61
 Sanjoinine G₂, S-62
 Sativanine A, S-92

Sativanine B, S-93
 Sativanine C, S-94
 Sativanine D, S-95
 Sativanine E, S-96
 Sativanine F, S-97
 Sativanine G, S-98
 Sativanine H, S-99
 Sativanine O, S-100
 Scutianine A, S-165
 Scutianine C, S-166
 Scutianine D, S-167
 Spinanine A, S-412
 Texensine, T-300
 Tscheschamine, T-648
 Venturamide A, V-53
 Vignatic acid A, V-99
 Vignatic acid B, V-100
 Waltherine A, W-6
 Waltherine C, W-7
 Zizyphine A, Z-22
 Zizyphine C, Z-23
 Zizyphine G, Z-24
 Zizyphine I, Z-21
 Zizyphine K, Z-25
 Zizyphine N, Z-26
 Zizyphine Q, Z-27

Amanita alkaloids

Phallacidin, P-306
 Phallacin, P-307
 Phallin B, P-308
 Phallisacin, P-309
 Phallisins, P-310
 Phalloidin, P-311
 Phalloin, P-312

Other cyclopeptide alkaloids

Antibiotic TMC 95A, A-1293
 Celogentin A, C-241
 Celogentin C, C-242
 Celogentin G, C-243
 Celogentin K, C-244
 Discokiolide A, D-847
 Discokiolide B, D-848
 Hibispeptin A, H-272
 Hibispeptin B, H-273
 Melleumin A, M-182
 Moroidin, M-702
 Stephanotic acid, S-540

Pyoverdins

Azotobactin D, A-1611
 Azotobactin G173, A-1612
 Azotobactin P19, A-1613
 Azoverdin A, A-1614
 Azoverdin G, A-1615
 Ferribactin G173, F-35
 Isopyoverdin Pp BTP 1, I-292
 Isopyoverdin Pp CFML 90-33, I-293
 Isopyoverdin Pp CFML 90-44, I-294
 Pseudobactin 7SR1, P-703
 Pyoverdin A1, P-838
 Pyoverdin BTP 2, P-839
 Pyoverdin BTP7G, P-841
 Pyoverdin BTP16G, P-842
 Pyoverdin BTP7, P-840
 Pyoverdin CFBP 2392, P-844
 Pyoverdin CFBP 2461, P-845
 Pyoverdin CHAO, P-846
 Pyoverdin C₄₄H₆₁N₁₃O₁₇, P-847
 Pyoverdin C₄₅H₆₃N₁₁O₂₄, P-848
 Pyoverdin C₅₂H₇₅N₁₅O₂₃, P-849
 Pyoverdin C₅₅H₈₃N₁₇O₂₁, P-850
 Pyoverdin C₅₇H₈₅N₁₇O₂₄, P-851
 Pyoverdin C, P-843
 Pyoverdin D, P-852
 Pyoverdin G 173, P-853
 Pyoverdin GM II, P-854
 Pyoverdin G4RA, P-855
 Pyoverdin II, P-856

Pyoverdin I, P-835
 Pyoverdin 2112, P-836
 Pyoverdin 11370, P-837
 Pyoverdin Pa II, P-857
 Pyoverdin Pa TII, P-858
 Pyoverdin PaC, P-859
 Pyoverdin Pf CCM 2798, P-864
 Pyoverdin Pf 1.3, P-860
 Pyoverdin Pf/3 17400, P-861
 Pyoverdin Pf 1547, P-863
 Pyoverdin Pf P19, P-865
 Pyoverdin Pf12, P-862
 Pyoverdin PL7, P-866
 Pyoverdin PL8, P-867
 Pyoverdin Pm, P-868
 Pyoverdin Pp 1, P-869
 Pyoverdin PpC 3B, P-870
 Pyoverdin PS 6.10, P-872
 Pyoverdin PS, P-871
 Pyoverdin PSS, P-873
 Pyoverdin R', P-874
 Pyoverdin 51WA, P-875
 Succinopyoverdin G173, S-628

Pyrrolo[2,3-*d*]pyrimidines

2-Amino-5-(aminomethyl)pyrrolo[2,3-*d*]pyrimidin-4(3*H*)-one, A-699
 4-Amino-5-bromopyrrolo[2,3-*d*]pyrimidine, A-713
 Archaeosine, A-1387
 5-Bromo-7*H*-pyrrolo[2,3-*d*]pyrimidine, B-351
 Cadeguomycin, C-7
 Cathestatin B, C-208
 Dapiramicin B, D-73
 Echiguanine B, E-9
 Estatin A, E-224
 Kanagawamicin, K-8
 Mycalisine B, M-778
 Rigidin A, R-98
 Tubercidin, T-658

Purines

Adenine, A-137
 Agelasimine A, A-181
 Agelasine A, A-184
 Agelasine C, A-185
 Agelasine D, A-186
 Agelasine E, A-187
 Agelasine I, A-183
 Agelasine, A-182
 Ageline A, A-191
 5-Amino-1,6-dihydro-7*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7-one, A-735
 6-Amino-2,8-dihydroxypurine, A-754
 6-Amino-2-hydroxypurine, A-818
 6-Amino-8-hydroxypurine, A-819
 6-Amino-1-methylpurine, A-842
 6-Amino-2-methylpurine, A-843
 6-Amino-3-methylpurine, A-844
 6-Amino-9*H*-purine-9-propanoic acid, A-900
 Asmarine B, A-1465
 Asmarine C, A-1466
 Asmarine I, A-1464
 9-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]adenine, B-305
 7-[3-Bromo-2-(2,3-dibromo-4,5-dihydroxybenzyl)-4,5-dihydroxybenzyl]-3,7-dihydro-1*H*-purine-2,6-dione, B-306
 Caffeine, C-15
 Caissaron, C-19
 3,5'-Cycloanthosine, C-918
 7-[5-(Decahydro-4*a*-hydroxy-1,2,5,5-tetramethyl-1-naphthalenyl)-3-methyl-2-pentenyl]-3,7-dihydro-2,3-dimethyl-6*H*-purin-6-one, D-111
 7-(2,3-Dibromo-4,5-dihydroxybenzyl)-3,7-dihydro-1*H*-purine-2,6-dione, D-311
 1,4-Dihydro-4,6,7-trimethyl-9*H*-imidazo[1,2-*a*]purin-9-one, D-521

3,9-Dihydro-1,3,9-trimethyl-8-nitroso-1*H*-purine-2,6-dione, D-523
 6-(6,8-Dihydroxy-9-purinyloxy)suberosanone, D-667
 1,3-Dimethylguanaine, D-748
 6-(6,8-Dioxo-1-purinyloxy)suberosanone, D-809
 Erinacean, E-177
 Flavine adenine dinucleotide, F-79
 Ganoderpurine, G-15
 Guanine, G-210
 Gutingimycin, G-227
 Herbipoline, H-143
 2'-*O*-[3-(4-Hydroxyphenyl)-2-methoxypropenoyl]-3'-*O*-[3-(1*H*-imidazol-4-yl)propenoyl]-5-methyl-5'-thioadenosine, H-681
N-[(5-Hydroxy-2-pyridinyl)methyl]adenosine, H-713
N-(6-Hydroxy-9-β-D-ribofuranosyl-9*H*-purin-2-yl)alanine, H-730
 2-Iminomethyl-3-methyl-6-methylamino-9*H*-purine, I-48
 Inosine, I-148
 Liberine, L-165
 Machilaminoside B, M-4
 Malonganenone A, M-74
 1-Methylguanaine, M-467
 7-Methylguanaine, M-469
 Microxine, M-603
 Nebularine, N-82
 Neplanocin B, N-168
 Nuttingine A, N-344
 Nuttingine D, N-345
 Olomoucine, O-90
 Ostreine A, O-132
 Pedatisectine A, P-151
 Phidolopin, P-357
 2,6-Purinediamine, P-817
 6,8-Purinediol, P-818
Subergorgia suberosa Purine-imidazole alkaloid, P-819
 6-Purinol, P-820
 Suvanine, S-649
 1,6,7,9-Tetrahydro-6-imino-1,9-dimethyl-8*H*-purin-8-one, T-194
*Agelas nakamura*i Thelepogane-purine alkaloid, T-352
 Theobromine, T-355
 Theophylline, T-360
 Triacanthine, T-451
 1,3,7-Trimethylsoguanine, T-608
 Uric acid, U-45
 Wybutoxine, W-29
 Wye base, W-30
 Xanthine, X-5
 Xanthosine, X-13

Pteridines and analogues

6-Acetyl-2-amino-7,8-dihydro-4(3*H*)-pteridinone, A-40
 6-Acetyl-1,3-dimethyl-2,4(1*H*,3*H*)-pteridine-dione, A-51
 6-Acetylhomopterin, A-52
 Alloxazine, A-632
 2-Amino-7-(1,2-dihydroxypropyl)-1,5-dihydro-4,6-pteridinedione, A-748
 2-Amino-4,6-dihydroxypteridine, A-750
 2-Amino-4,7-dihydroxypteridine, A-751
 2-Amino-4,7-dihydroxy-6-pteridinecarboxylic acid, A-752
 2-Amino-6-(hydroxymethyl)-4(1*H*)-pteridinone, A-802
 2-Amino-6-[3-hydroxy-4-(phosphonoxy)-1-butenyl]-4(1*H*)-pteridinone, A-809
 2-Amino-4-hydroxypteridine, A-816
 2-Amino-4-hydroxy-6-pteridinecarboxylic acid, A-817
 Aminopteroic acid, A-899
 2-Amino-7-(1,2,3-trihydroxypropyl)-4(1*H*)-pteridinone, A-924
 2-Amino-4,6,7-trihydroxypteridine, A-925
 Antibiotic 2096A, A-1093
 Asteropterin, A-1516

Chrysopterin, C-432
 1,5-Dihydro-2,4,6(3*H*)-pteridinetriene, D-494
 Drosopterin, D-941
 Erythropterin, E-216
 Fervulin, F-48
 Flavine adenine dinucleotide, F-79
 Hirudinoidine A, H-304
N-[4-(4-Hydroxybenzoylamino)butyl]-1,3-dimethylmazine-6-carboxamide, H-410
 6-(Hydroxymethyl)-2,4(1*H*,3*H*)-pteridinedione, H-623
 6-(2-Hydroxy-2-oxido-4*H*-1,3,2-dioxaphosphorin-6-yl)-2,4(1*H*,3*H*)-pteridinedione, H-649
 Ichthyopterin, I-13
 Lepidopterin, L-109
 Lumichrome, L-285
 Methanopterin, M-235
 Molybdenum cofactor, M-681
Racophorus Pentapteridinyl skin pigment, P-230
 Photolumazine B, P-382
 Pseudoanchynazine A, P-698
 Pseudoanchynazine B, P-699
 Pseudoanchynazine C, P-700
 2,4,7(1*H*,3*H*,8*H*)-Pteridinetriene, P-763
 Pterorhodine, P-771
 Pteroylglutamic acid, P-772
 Pyrizinostat, P-935
 Reumycin, R-66
 Rhizopterin, R-82
 Russuapteridine yellow IV, R-166
 Sarcinapterin, S-72
 Sulfohalopterin 2, S-637
 Urochordamine A, U-47
 Xanthopterincarboxylic acid, X-12

Saxitoxins

11-Saxitoxinacetic acid, S-110
 Saxitoxin, S-109
 Zetekitoxin AB, Z-18

Alkaloidal metal complexes

Antibiotic ASK 753, A-1117
 Antibiotic AS-N-7A, A-1118
 Antibiotic NRCS 15, A-1219
 Astechrome, A-1509
 Coprogen, C-635
 Danomycin, D-16
 Ferricrocin, F-36
 Ferrimycin A₁, F-37
 Ferrioxamine A₂, F-38
 Ferrioxamine B, F-39
 Ferrioxamine C, F-40
 Ferrioxamine D₂, F-41
 Ferrioxamine E, F-42
 Ferrioxamine F, F-43
 Ferrioxamine H, F-44
 Ferroverdin, F-45
 Fluopsin B, F-96
 Fluopsin C, F-97
 Fluopsin N, F-98
 Fusigen, F-231
 Grisonomycin, G-179
 Molybdenum cofactor, M-681
 Neocoprogen II, N-108
 Neocoprogen I, N-107
 Pilosamycin, P-427
 Pseudobactin B, P-702
 Salmycin B, S-34
 Salmycin C, S-35
 Viridomycin A, V-156
 Viridomycin E, V-159

Naphthyridinomycins

Aclidinomycin B, A-82
 Aclidinomycin C, A-83
 Antibiotic SF 1739, A-1254
 Bioxalomycin α₁, B-140
 Cyanocycline D, C-824
 Cyanonaphthyridinomycin, C-830
 Dnacin A₁, D-876
 Dnacin B₁, D-877

Naphthyridinomycin A, N-32
Naphthyridinomycin B, N-33

Saframycins

Ecteinasidin 594, E-34
Ecteinasidin 597, E-35
Ecteinasidin 722, E-36
Ecteinasidin 743, E-37
Jorumycin, J-48
Renieramycin A, R-41
Renieramycin H, R-42
Renieramycin K, R-43
Renieramycin P, R-44
Safracin A, S-2
Saframycin A_{D-1}, S-3
Saframycin A, S-4
Saframycin H, S-5
Saframycin Mx 1, S-6
Saframycin R, S-7
Saframycin S, S-8
Saframycin Y_{2b-D}, S-10
Saframycin Y₃, S-11
Saframycin Y_{D-1}, S-12
Saframycin Y_{D-2}, S-13
Saframycin Y_{2b}, S-9

Miscellaneous acyclic alkaloids

Aberiamide, A-12
2-Acetyl-3-amino-5-hydroxy-1,4-naphthoquinone, A-41
3-Acetyl-2,5-diamino-1,4-benzoquinone, A-48
Acuminaminoside, A-131
AK toxin I, A-220
Amaminol A, A-671
2-Amino-3-(3-amino-1*H*-inden-2-yl)propanoic acid, A-698
2-(2-Aminobenzylamino)benzyl alcohol, A-708
Anhydrotetracycline, A-1034
2-Anilino-1,4-naphthoquinone, A-1040
Anthracyclinone blue B, A-1080
Antibiotic SB 217452, A-1240
Arboreumine, A-1384
Bacillibactin, B-2
N,N'-Bis(2,6-dimethylphenyl)-2,2,4,4-tetramethyl-1,3-cyclobutanediimine, B-175
N,N'-Bis(4-hydroxybenzyl)hydroxylamine, B-181
Bradyoxetin, B-277
Brasilamide A, B-278
(Carbamoylmethyl)trimethylammonium(1+), C-119
Celenamide D, C-237
Coriacenine A, C-648
Coriacenine D, C-649
Coriacenine E, C-650
Cyclohexylamine, C-868
Cyclorocaglamide, C-904
Dicarprine C, D-345
Donine, D-919
Dysidenamide, D-971
Dysiherbaine, D-974
4,5-Eicosanediamine, E-47
Enterobactin, E-86
Erbstatin, E-144
Fontaineine, F-124
Foveoglin A, F-144
GB4 Toxin, G-28
Gossyrubilon, G-145
Gusanlung C, G-225
Gyromitrin, G-236
10-Hydroxy-18-(*N*-naphthalenyl-*N*-phenylamino)betaone C, H-637
Lilacinone, L-172
Lomaiviticin B, L-238
Lutinine, L-311
Malynamide W, M-78
Neodysiherbaine A, N-112
Nephila clavata Neurotoxin 4 JSTX-4, N-184
Nitrosoxacina A, N-253
Nitrosoxacina B, N-254
Nitrosoxacina C, N-255
Pandanamine, P-65

Peniamidienone, P-185
Penidilamine, P-190
1,11,19,27,35-Pentaamino-3-tetracontanol, P-198
Poecillanosine, P-530
Prangosine, P-585
Prepsammaplina A, P-622
Rhapsamine, R-71
Salmochelin 4, S-32
Signatipennine, S-299
Solanapyrone L, S-341
Spergualin, S-394
Stealthin A, S-500
Stealthin C, S-501
Stockerine, S-555
Perophora viridis 1,2,3-Thiocane glycoside, T-382
Triuret, T-627

Miscellaneous one-ring system alkaloids

Antibiotic 2197A, A-1094
Antibiotic BE 14106, A-1127
Antibiotic FR 252921, A-1174
Antibiotic FR 252922, A-1175
Antibiotic NA 22598A₁, A-12111
Antibiotic PF 1163A, A-1227
Antibiotic SF 2197B, A-1263
Antibiotic TAN 1057C, A-1285
Antibiotic TMC 135A, A-1295
Antibiotic TMC 135B, A-1296
Antibiotic TMC 1A, A-1291
Antibiotic TMC 1C, A-1297
Aquileline, A-1369
Asterionellin A, A-1514
Asterionellin B, A-1515
Aureoverticillactam, A-1555
Bahamamide, B-5
Caledonin, C-33
Caprazamycin, C-98
Chrysopine, C-431
Cyclamenol, C-832
Cycloclausenamide, C-855
1,13-Diazacyclotetracosane-2,14-dione, D-294
5,20-Dipropyl-1,16-dioxo-4,19-diazacyclotriacontane-7,10,22,25-tetraene-15,30-dione, D-825
Dovyalicin A, D-927
Dovyalicin D, D-928
Epilachnane, E-93
Halimedin, H-25
1,4,5,6,7,8-Hexahydro-1,3-diazocine-2,4-dicarboxylic acid, H-230
Hicksoane A, H-274
Liposidomycin, L-188
Manumycin A, M-87
Manumycin C, M-88
Manumycin D, M-89
4-Methyl-1,2,6,8-tetraazacycloundeca-4,9-diene-3,7,11-trione, M-565
Micromonosporin A, M-599
Obscuroside C_{2a}, O-10
Penaresidin B, P-175
Epilachna borealis Polyazamacrolides, P-547
5-Propyl-1-oxa-4-azacyclotridecan-13-one, P-651
Scorodophlone A, S-162
Syringolin A, S-664
Syringolin C, S-665
Syringolin D, S-666
Syringolin F, S-667
Terrestrial acid, T-104
5,6,7,8-Tetrahydro-5,7-dihydroxy-1*H*-azocin-2-one, T-151
Triticone E, T-626

Miscellaneous two-ring system alkaloids

Abikoviromycin, A-14
3-Acetyl-2,7-naphthyridine, A-72

Acremolactone B, A-111
Ageladine A, A-177
Agrocycbenine, A-206
Alchorinidine, A-253
Alchornine, A-254
Aldisine, A-255
Dendrobates Alkaloid 275A, A-367
Altemicidin, A-660
4-Amino-3,6-dihydroxy-1*H*-pyrazolo[3,4-*d*]pyrimidine, A-755
4-Amino-3-hydroxy-1*H*-pyrazolo[3,4-*d*]pyrimidine, A-820
2-Amino-1*H*-imidazo[4,5-*b*]pyrazine, A-829
1-Amino-4-phenyl-2,7-naphthyridine, A-865
5-(12-Aminotridicycl)hexahydropyrrolo[2,1-*b*]oxazole, A-920
5-(10-Aminoundecyl)hexahydropyrrolo[2,1-*b*]oxazole, A-927
Anthocero Diazonin, A-1078
Antibiotic 2096A, A-1093
Antibiotic AF 140, A-1115
Antibiotic FA 4283, A-1166
Antibiotic SB 291071, A-1242
Antibiotic SB 315021, A-1243
Antibiotic TA 3037A, A-1273
Antibiotic TAN 1251B, A-1284
Antibiotic YCM 1008A, A-1320
Aplaminal, A-1341
Aplidinone A, A-1344
Aplidinone B, A-1345
Arenaine, A-1401
Aspergillazine A, A-1472
Avenalumin II, A-1561
Avenalumin III, A-1562
9-Azabicyclo[3.3.1]nonan-3-one, A-1580
4-Azonia Spiro[3.3]heptane-2,6-diol, A-1610
4-Benzyl-1,4-dihydro-3-oxo-3*H*-pyrrolo[2,1-*c*][1,4]oxazine-6-carboxaldehyde, B-86
Cypridina Biluciferyl, B-137
Brevioxime, B-297
7-Bromocavernicolone, B-301
5-(2-Butenylidene)-3-ethyl-1,2,3,4,5,7*a*-hexahydro-4*aH*-1-pyridine-4,4*a*-diol, B-424
Callipeltoside A, C-37
11-Carboxyanatoxin a, C-143
Catharsitoxin A, C-198
Catharsitoxin B, C-199
Catharsitoxin C, C-200
Cephalandole A, C-249
Cephalimycin A, C-254
Ceratamine A, C-277
Chinese bittersweet alkaloid III, C-385
ζ-Clausenamide, C-501
Coelenterazine, C-560
Curvupallide A, C-816
Cyclocarbamide A, C-852
Dehydrocrambine A, D-155
3,4-Dihydro-2*H*-1,3-benzoxazin-2-one, D-405
6,7-Dihydro-4*H*-furo[2,3-*c*]azepine-4,8(5*H*)-dione, D-433
2,7-Dihydro-2,5,7-trimethyl-5*H*-imidazo[4,5-*e*]1,2,4-triazine-3,6-dione, D-522
Dioscoretine, D-797
Donaxanine, D-915
Donaxarine, D-917
Dysinoin A, D-975
Epostatin, E-107
Fluviol D, F-114
Fusarin A, F-226
Grandisine B, G-161
Hatomamicin, H-77
2-(5-Heptenyl)-6,7,8,8*a*-tetrahydro-3-methyl-4*H*-pyrrolo[2,1-*b*][1,3]oxazin-4-one, H-129
3,4,5,6,7,8-Hexahydro-1,7-benzodiazecine(1*H*)-one, H-229
Homoanatoxin a, H-325
7-Hydroxy-2*H*-1,4-benzothiazin-3(4*H*)-one, H-407
7-Hydroxy-2-phenyl-4*H*-3,1-benzoxazin-4-one, H-675
1-Hydroxy-4-phenyl-2,7-naphthyridine, H-686

Hyperaspine, H-769
 Isooxodesmosine, I-260
 Kobutimycin A, K-53
 Kobutimycin B, K-54
 Laccarin, L-4
 Lansamide 2, L-39
 Leptocarpinine, L-118
Cypridina Luciferin, L-281
 Lucilactaene, L-283
 Lukianol A, L-284
 Manicoline B, M-83
 Matlystatin D, M-117
 Matlystatin E, M-118
 Matlystatin F, M-119
 2-(3-Methylbutyl)-1*H*-pyrrolo[2,3-*b*]pyridine, M-422
 4-Methylcinnoline, M-428
 8-(5-Methylhexyl)-1*H*,3*H*-[1,3,4]oxadiazolo[3,4-*a*][1,2,4]triazin-1-one, M-471
 4-Methyl-2,6-naphthyridine, M-487
 3-Methyl-1*H*-pyrrolo[3,2-*b*]pyridine, M-551
 Monocillinol A, M-687
 Mycothiol bimane, M-791
 Nagstatin, N-9
 Nitrabirine, N-240
 Nostocine A, N-320
 Oxodesmosine, O-175
 Pantocin A₂, P-74
 Pantocin A, P-75
 Pentosidine, P-233
 1-Pentyl-9-azabicyclo[3.3.1]nonan-3-one, P-235
 Phaeosphaeride A, P-302
 2-Phenyl-1-benzazocin-8-ol, P-340
 Phloeodictyne B, P-367
Diaphus Preluciferin, P-612
 Pseudotheonamide A₁, P-727
 Pseudotheonamide B₂, P-728
 Pyranonigrin A, P-880
 Pyridicine, P-932
 Reumycin, R-66
 Salinosporamide A, S-28
 Salinosporamide D, S-30
 Salinosporamide E, S-31
 Salinosporamide I, S-27
 Salviamiltamide, S-41
 Sibyllimycin, S-293
 Spirocalcaridine A, S-428
 Spiroleuccetidine, S-430
 Streptazoline, S-560
 Streptazone B₁, S-561
 Streptazone C, S-562
 Subereatensin, S-620
 Thiomarinol A, T-385
 Thiomarinol D, T-386
 Thiomarinol E, T-387
 Trichochrome B, T-483
 Trichochrome C, T-484
 Trichochrome E, T-485
 Trichochrome F, T-486
 Trichodermamide A, T-487
 Verruculotoxin, V-84
 Xanthiazone, X-3
 Xylogranatinine, X-29

Miscellaneous three-ring system alkaloids

Aaptamine, A-2
 Aaptosamine, A-3
 Aaptosine, A-4
Mantella Alkaloid 261C, A-452
Dendrobates Alkaloid 251F, A-509
 7-Amino-3,4,4*a*,10*b*-tetrahydro-2-(hydroxymethyl)-4-methylbenzo[*f*]quinolin-6(5*H*)-one, A-914
 Antibiotic CJ 12662, A-1141
 Antibiotic FR 901483, A-1178
 Antibiotic ICM 0201, A-1188
 Antibiotic NG 311, A-1214
 Antibiotic PF 1140, A-1226
 Aphanorphine, A-1333

Aplidite A, A-1346
 Ascidiathiazone A, A-1457
 Ascidiathiazone B, A-1458
 Asimicilone, A-1463
 Aulosirazole, A-1540
 Aurachin E, A-1544
 Azamerone, A-1599
 6-Azidotetrazolo[5,1-*a*]phthalazine, A-1607
 Benzo[*c*][2,7]naphthyridin-4-ol, B-75
 Bhimamycin D, B-114
 Brevinic acid, B-296
 Buddamine, B-382
 7-Butyl-1,2-dichloro-6,8-dihydroxy-9*H*-pyrrolo[2,1-*b*][1,3]benzoxazin-9-one, B-429
 Buzonamine, B-484
 12-Carboxymonoscorubramine, C-150
 Cassiarine A, C-181
 Cassiarine B, C-182
 Cheliensisamine, C-358
 2-Chloro-7-ethyl-6,8-dihydroxy-9*H*-pyrrolo[1,2-*b*][1,3]benzoxazin-9-one, C-403
 2-Chloro-7-ethyl-8-hydroxy-6-methyl-9*H*-pyrrolo[1,2-*b*][1,3]benzoxazin-9-one, C-404
 Chlorpromazine, C-417
 Cinachyramine, C-443
 Cinereain, C-449
 ComX_{RO-E-2} pheromone, C-593
 ComX_{RO-H-1} pheromone, C-594
 Conicaquinone A, C-605
 Conicaquinone B, C-606
 Crepidine, C-736
 Cycloanchinopeptolide C, C-840
 Cyclodidemiserinol, C-860
 Cyclopiamine A, C-893
 Cyliodrospermopsin, C-930
 Dactylicapnosinine, D-4
 Decahydro-10-[(3-hydroxy-2,4,6-octadecatrienoyl)amino]-1,1*a*,11*b*-trihydroxy-2*H*-pyrazino[1,2-*a*]pyrrolo[2,1-*c*]pyrazine-3,6,9-trione, D-110
 Diazepinomicin, D-297
 1,2-Dichloro-8-hydroxy-6-methyl-7-propyl-9*H*-pyrrolo[1,2-*b*][1,3]benzoxazin-9-one, D-352
 Dictyolomide A, D-367
 2,9-Dimethyl-2,9-diazatricyclo[10.2.2.2^{s,8}]octadeca-5,7,12,14,15,17-hexaene-3,10-diol, D-745
 Dysibetaine PP, D-965
 Erinacerin A, E-178
 Eugenine A, E-293
 Eugenine B, E-294
 Euthyroideone A, E-304
 Fareanine, F-8
 Fulvanine E, F-167
 Fumiquinazoline F, F-188
 Furomegistine II, F-208
 Gauthierialine A, G-27
 Gephyrotoxin, G-65
 Gracicleistanthoside, G-149
 Gymnodimine, G-232
 Haematopodin B, H-8
 Haloxine, H-38
 Hirudinoidine A, H-304
 Hydnellin B, H-392
 3-Hydroxy-8,9-methylenedioxybenzo[*h*]quinoline, H-599
 8-Hydroxystreptazolone, H-741
 Hyrtiazepine, H-778
 Isoperlylryne, I-262
 Isothapsakone A, I-335
 Janoxepin, J-19
 Kealiimine A, K-24
 Kealiiquinone, K-25
 Lemonnierin, L-89
 Lepistine, L-111
 Lycopladine A, L-338
 Lycoposerramine B, L-343
 Maingayinine, M-48
 Malassezindole A, M-66
 Maltoxazine, M-77
 Martinelline, M-111
 Mearsamine, M-138
 Mescengricin, M-223

8-Methoxy-9*H*-benzo[*de*][1,6]naphthyridin-9-one, M-241
 7-Methoxy-4-methyl-γ-carboline, M-260
 5-Methoxy-2-(methylthio)-1,3-thiazino[6,5-*b*]indole, M-276
 4-Methyl-3*H*-pyrrolo[2,3-*c*]quinoline, M-552
 Millaurine A, M-614
 Millaurine, M-613
 Millettonine, M-615
 Millingtonine, M-618
 Monascopyridine A, M-684
 Monascopyridine B, M-685
 Monascorubramine, M-686
 Moschamide, M-710
 Mycenarubin A, M-784
 Mycenarubin B, M-785
 Myrmicarin 213B, M-801
 Neosymbioimine, N-143
 Nigeglanine, N-191
 Nigelline, N-192
 Ningalin B, N-221
 Ningalin C, N-222
 Ostreogrycin A, O-131
 Oxepinamide A, O-158
 Oxepinamide C, O-159
 Paeciloxazine, P-9
 Perloline, P-269
 Phyllostictine A, P-391
 Phyllostictine B, P-392
 Phyllostictine C, P-393
 Phyllostictine D, P-394
 Pinnamine, P-436
 Prolifine, P-638
 Pterocellin A, P-764
 Pterocellin B, P-765
 Pterocellin C, P-766
 Pterocellin D, P-767
 Pterocellin F, P-768
 Pyrroindomycin A, P-941
 Quinocitrinone A, Q-23
 Quinolactacin A₁, Q-25
 Quinolactacin D₁, Q-27
 Rubrobromide, R-148
 Rubropunctamine, R-152
 Salinosporamide C, S-29
 Sarcodonin, S-76
 Schulzeine A, S-142
 Schulzeine C, S-143
 Scytonemin, S-168
 Scytonine, S-169
 Sequoiamonascin D, S-250
 Sibirinine, S-292
 Silvaticamide, S-300
 Spirotryprostatin A, S-468
 Streptopyrrole, S-571
 Suomilide, S-643
 Symbioimine, S-654
 Syncarpurea, S-656
 Tamynine, T-23
 Terresoxazine, T-103
 Terreusinone, T-105
 1,3,4,9-Tetrahydro-1,1-dimethylpyrrolo[3,4-*b*]indole, T-169
 Tetramethoxyscytonemin, T-279
 Tetraponerine 1, T-290
 Tetraponerine 3, T-291
 Tetraponerine 5, T-292
 Tetraponerine 7, T-293
 Ulbactin D, U-6
 Ureidosorbicillinol, U-44
 Variolin B, V-32
 Variolin D, V-33
 Vitedoamine A, V-164
 Yuremamine, Y-22
 Zyzyanone A, Z-40

Miscellaneous four- and higher-ring system alkaloids

Abyssomicin B, A-19
 Acosmine, A-107
 Acutopyrrocoline, A-135

- Adociaquinone B, A-147
 Aglaiaastatin, A-197
 Aglaroxin I, A-200
 α -Aldotriperideine, A-256
Mantella Alkaloid 384A, A-374
 Ancorinazole, A-993
 Antibiotic BA 181314A, A-1125
 Antibiotic BE 54017, A-1131
 Antibiotic BE 55051, A-1133
 Antibiotic BE 54238A, A-1136
 Antibiotic BMY 42428, A-1138
 Antibiotic FL 120B', A-1169
 Antibiotic GKK 1032A₂, A-1182
 Antibiotic GKK 1032B, A-1183
 Antibiotic S 433, A-1235
 Antibiotic SF 2695, A-1259
 Antibiotic VM 55595, A-1306
 Antibiotic VM 55599, A-1307
 Arboloscine, A-1381
 Asperparaline A, A-1486
 Auranthine, A-1547
 Austamide, A-1556
 Azaacidone A, A-1579
 Benhamycin, B-62
 Benzomalvin B, B-74
 Capsimycin, C-102
 Cephalocyclidine A, C-259
 Chaetominine, C-336
 Chartellamide A, C-349
 Chartelline A, C-350
 Chenoalbicin, C-364
 Chestnutamide, C-370
 Chilocorine D, C-380
 Chloptosin, C-394
 Citridone D, C-481
 Citrinadin A, C-482
 Communesin A, C-581
 Communesin B, C-582
 Communesin G, C-583
 Communesin H, C-584
 Crociflorinine, C-766
 Cyclopiamide, C-892
 Cyclozoanthamine, C-922
 Dactylicapnosine, D-3
 Dalomycin T, D-11
 Daphnipaxinine, D-67
 Decursivine, D-137
 Dehydropuntarenine, D-165
 Dehydrosaulatine, D-167
 Diazonamide B, D-298
 Dibromophakellstatin, D-336
 Dictyodendrine A, D-364
 Dictyodendrine B, D-365
 Dictyodendrine C, D-366
 1,2-Dihydrothiazolo[2',3':2,3]-pyrimido[4,5-*b*]indole-1-carboxylic acid, D-517
 1,8-Dihydroxy-3-methylbenzo[*b*]phenanthridine-7,12-dione, D-612
 Discoderamide, D-844
 Distomadine A, D-868
 Distomadine B, D-869
 Dragmacidin E, D-931
 Dragmacidin F, D-932
 Drimiopsine A, D-940
 Dynamycin A, D-959
 Dynamycin L, D-960
 Dynamycin P, D-961
 6-Ethenyl-3,4,4*b*,5,6,7,8,9,10*b*,12-decahydro-12-hydroxy-5-methyl-1*H*-pyrido[4',3':4,5]pyrano[3,2-*h*]isoquinoline-1,10(2*H*)-dione, E-227
 2-Ethyl-11-methoxy-3-methyl-3*H*-[1,6]naphthyrindino[6,5,4-*def*]quinoxaline, E-251
 Evodiainine, E-311
 Excentricine, E-318
 Fissoldhimine, F-67
 Fleophilone, F-86
 Fumiquinazoline D, F-187
 Gardquinolone, G-24
 Gracilamine, G-150
 Granulatimide, G-169
 Grossularine 1, G-183
 Grossularine 2, G-184
 Hamigeran D, H-43
 Haouamine A, H-47
 Herquiline A, H-152
 Herquiline B, H-153
 Himeradine A, H-284
 Hinckdentine A, H-287
 Hirsutellone A, H-299
 Hirsutellone B, H-300
 Hirsutellone D, H-301
 Hirsutellone E, H-302
 Hirsutellone F, H-303
 Homofascaplysate A, H-332
 Homofascaplysin B, H-334
 Homofascaplysin B1, H-335
 Homofascaplysin C, H-336
 Hopeanine, H-352
 Hostasinine A, H-363
 Huperzine V, H-374
 22-Hydroxyacuminatine, H-403
 1-(4-Hydroxyphenyl)-3*H*-benzo[*a*]pyrrolo[2,3-*c*]-phenazine, H-674
 4-Hydroxy-1',3,4'-tris(4-hydroxyphenyl)-3'-[2-(4-hydroxyphenyl)ethyl]-7'-(sulfoxy)spiro[furan-2(5*H*),2'(3'*H*)-pyrrolo[2,3-*c*]carbazole]-5,5'(6'*H*)-dione, H-753
 Hypodemapyrazine, H-772
 Ileabethoxazole, I-27
 7*H*-Indolo[3,2-*f*]phenanthridine-7,13(12*H*)-dione, I-107
 Indolo[3,2,1-*ij*]quino[4,5-*bc*][1,5]naphthyridine, I-114
 Ingamine A, I-141
 Ingenamine B, I-143
 Ingenamine D, I-144
 Ingenamine E, I-145
 Ingenamine G, I-146
 Ingenamine, I-142
 Isaindigotidione, I-181
 Isatisine A, I-185
 Isogranulatimide, I-237
 Isomitomycin A, I-246
 2-Isopropyl-10-methoxybenzimidazo[6,7,1-*def*][1,6]naphthyridine, I-273
 Isoquinocycline A, I-300
 Isoschizogamine, I-313
 Isotriperideine, I-340
 Jadomycin A, J-9
 Jadomycin Ala, J-10
 Jadomycin F, J-11
 Jadomycin S, J-12
 Jadomycin T, J-13
 Jadomycin V, J-8
 Javanicunine A, J-29
 Kapakahine A, K-9
 Kapakahine B, K-10
 Kapakahine C, K-11
 Kapakahine E, K-12
 Kapakahine G, K-13
 Ketoanhydrokinamycin, K-37
 Lamellarin H, L-24
 Lankanescine, L-31
 Lemonomycin, L-90
 Leptocarpine, L-117
 Lihouidine, L-171
 Lindeniamine, L-183
 Lophocine, L-261
 Luotonine A, L-298
 Luteinine, L-308
 Lycopladiene B, L-339
 Lycopladiene D, L-340
 Lycoposerramine A, L-342
 Madangamine F, M-32
 Malasseziacitrin, M-60
 Malasseziazole A, M-62
 Malasseziazole B, M-63
 Malasseziazole C, M-64
 Malbrancheamide A, M-69
 Marcfortine B, M-98
 Marine *Streptomyces* C₃₀H₃₈N₂O₅ lactam, M-105
 Massadine, M-112
 Megistosarconine, M-152
 Mersilosine, M-214
 Mersinaline, M-215
 Mersinine A, M-217
 Mersirachine, M-218
 Metenaticin C, M-234
 10-Methoxybenzimidazo[6,7,1-*def*][1,6]-naphthyridine, M-239
 37-(Methoxycarbonyl)dictyodendrine E, M-247
 11-Methoxy-3*H*-naphthridino[6,5,4-*def*]quinoxalin-2-ol, M-277
 Mevashuntin, M-587
 Misenine, M-649
 Mollenine A, M-670
 Monomargine, M-694
 Myrioneurinol, M-796
 Myrmicaridin 430A, M-800
 Myrmicaridin 663, M-798
 Myrobotinol, M-802
 Necatorine, N-84
 Necatorone, N-85
 Neosurugatoxin, N-141
 Ningalin A, N-220
 Nirurine, N-236
 Nitraramine, N-244
 Nitrarcine, N-245
 Njaoamine A, N-259
 Njaoamine E, N-260
 Njaoamine F, N-261
 Noelaquinone, N-277
 Norzoanthamine, N-317
 Notoamide D, N-325
 Nybomycin, N-346
 Ophiuroidine, O-98
 Palauamine, P-18
 Panacosmine, P-44
 Papuamine, P-84
 Paraherquamide A, P-90
 Paraherquamide G, P-91
 Perinadine A, P-261
 Phalarine, P-305
 Phlegmariurine C, P-364
 Pinnatoxin A, P-437
 Pinnatoxin B, P-438
 Pinnatoxin D, P-439
 Pteriatoxin A, P-760
 Pteriatoxin B, P-761
 Purpurone, P-825
 Quinocarcinol, Q-22
 Quinocarcin, Q-21
 Quinolactacide, Q-24
 Ripromycin, R-99
 Roehybridine, R-116
 Rubrolone, R-150
 Sanguinone A, S-58
 Sanguinone B, S-59
 Sarain A, S-67
 Sarain C, S-68
 Schischkiniin, S-127
 Schoberine, S-140
 Sclerotigenin, S-154
 Securamine A, S-203
 Securamine C, S-204
 Spirolide B, S-431
 Spirolide D, S-432
 Spirolide E, S-433
 Spirolide G, S-434
 Stephacidin B, S-532
 Stephaoxocanine, S-541
 Stylissadine A, S-610
 Styloguanidine, S-611
 Surugatoxin, S-645
 Syphilobin F, S-662
 Tetrapetalone C, T-287
 Tetrazomine, T-296
 Thiaplidiaquinone A, T-368
 Thiaplidiaquinone B, T-369
 Thorectandramine, T-396
 Uncialamycin, U-23
 Variecolortide A, V-30
 Variecolortide B, V-31

Xanthobaccin A, X-6
 Zoanthamide, Z-29
 Zoanthamine, Z-30
 Zoanthenamine, Z-31

Alkaloids of unknown or partially unknown structure

Abrasine, A-15
 Acscine, A-122
 Aculeacin B, A-130
 Aestivine, A-163
 Agelasine, A-182
 Ajmalinine, A-218
 Akuammenine, A-227
 Alangine A, A-236
 Alanginine, A-237
 Alchornidine, A-253
 Alginidine, A-261
 Alginine, A-262
Mantella Alkaloid 161, A-318
Mantella Alkaloid 191, A-319
Mantella Alkaloid 317, A-320
Pancreatium Alkaloid 6, A-323
Pancreatium Alkaloid 31, A-324
Rosmarinus officinalis Alkaloid 2, A-329
Senecio renardi Alkaloid 4, A-332
Sophora Alkaloid 9, A-333
Buxus Alkaloid 1, A-275
Buxus Alkaloid 2, A-276
Buxus Alkaloid 3, A-277
Buxus Alkaloid 4, A-278
Buxus Alkaloid 5, A-279
Dendrobates Alkaloid 185, A-287
Dendrobates Alkaloid 197, A-288
Dendrobates Alkaloid 217, A-290
Dendrobates Alkaloid 225, A-291
Dendrobates Alkaloid 233, A-292
Dendrobates Alkaloid 295, A-293
Dendrobates Alkaloid 301, A-294
Dendrobates Alkaloid 325, A-295
Dendrobates Alkaloid 351, A-296
Fritillaria Alkaloid 5, A-297
Fritillaria Alkaloid 6, A-298
Leuconotis griffithii Alkaloid 308, A-309
Lycopodium Alkaloid 168, A-312
Lycopodium Alkaloid 258, A-313
Daphniphyllum Alkaloid A₁, A-345
Abrus precatorius Alkaloid A, A-347
Buxus Alkaloid A1, A-348
Buxus Alkaloid A2, A-349
Buxus Alkaloid A3, A-350
Buxus Alkaloid A4, A-351
Buxus Alkaloid A5, A-352
Chondrodendron limacifolium Alkaloid A, A-353
Dendrobates Alkaloid 209A, A-356
Dendrobates Alkaloid 239A, A-360
Dendrobates Alkaloid 241A, A-361
Dendrobates Alkaloid 251A, A-363
Dendrobates Alkaloid 257A, A-364
Dendrobates Alkaloid 265A, A-365
Dendrobates Alkaloid 281A, A-368
Dendrobates Alkaloid 297A, A-369
Dendrobates Alkaloid 309A, A-370
Erythrophleum Alkaloid A, A-371
Euonymus Alkaloid A, A-372
Fritillaria Alkaloid A, A-373
Mantella Alkaloid 384A, A-374
Securidaca longipedunculata Alkaloid A, A-375
Stephania glabra Alkaloid A, A-376
Stephania glabra Alkaloid A, A-377
Stephania japonica Alkaloid A, A-378
Taxus baccata Alkaloid A, A-379
Thermopsis lanceolata Alkaloid A, A-380
Alangium Alkaloid AL64, A-381
Alangium Alkaloid B1, A-383
Alangium Alkaloid B2, A-384
Alangium Alkaloid B3, A-385
Alangium Alkaloid B4, A-386
Alangium Alkaloid B5, A-387
Asteracantha Alkaloid B1, A-388

Asteracantha Alkaloid B2, A-389
Buxus Alkaloid B, A-390
Buxus Alkaloid B1, A-391
Buxus Alkaloid B4, A-392
Buxus Alkaloid B5, A-393
Buxus Alkaloid B6, A-394
Buxus Alkaloid B7, A-395
Buxus Alkaloid B8, A-396
Buxus Alkaloid B387, A-397
Buxus Alkaloid B398, A-398
Buxus Alkaloid B426, A-399
 C-Alkaloid B, A-400
Dendrobates Alkaloid 225B, A-405
Dendrobates Alkaloid 237B, A-407
Dendrobates Alkaloid 239B, A-408
Dendrobates Alkaloid 253B, A-409
Dendrobates Alkaloid 257B, A-410
Dendrobates Alkaloid 267B, A-411
Dendrobates Alkaloid 281B, A-413
Dendrobates Alkaloid 283B, A-414
Dendrobates Alkaloid 297B, A-415
Dendrobates Alkaloid 307B, A-416
Dendrobates Alkaloid 309B, A-417
Gentiana macrophylla Alkaloid B, A-418
Mantella Alkaloid 271B, A-420
Mantella Alkaloid 293B, A-421
Mantella Alkaloid 295B, A-422
Pseudophryne Alkaloid 252B, A-423
Skytanthus acutus Alkaloid B, A-424
Sophora griffithii Alkaloid B, A-425
Strychnos camptoneura Alkaloid B, A-426
Thermopsis lanceolata Alkaloid B, A-427
 Alkaloid 241B, A-428
 C-Alkaloid BL, A-429
Buxus Alkaloid BX6, A-430
Buxus Alkaloid BX10, A-431
Chamaecytisus Alkaloid C₁, A-432
Chamaecytisus Alkaloid C₂, A-433
Rauwolfia obscura Alkaloid C₂, A-434
Buxus Alkaloid C, A-435
 C-Alkaloid C, A-436
Dendrobates Alkaloid 225C, A-439
Dendrobates Alkaloid 237C, A-440
Dendrobates Alkaloid 239C, A-441
Dendrobates Alkaloid 251C, A-442
Dendrobates Alkaloid 283C, A-443
Dendrobates Alkaloid 307C, A-444
Dendrobates Alkaloid 309C, A-445
Erythrophleum Alkaloid C, A-446
Gentiana macrophylla Alkaloid C, A-447
Leucojum Alkaloid C, A-448
Mantella Alkaloid 205C, A-449
Mantella Alkaloid 211C, A-450
Mantella Alkaloid 321C, A-453
Salicornia herbacea Alkaloid C, A-454
Stephania glabra Alkaloid C, A-455
Voacanga thouarsii Alkaloid C, A-456
 Alkaloid C2, A-457
 Alkaloid 1 (Calabash curare), A-342
 Alkaloid 2 (Calabash curare), A-343
 Alkaloid CC 3, A-460
 Alkaloid CC 11, A-461
 Alkaloid CC 13, A-462
 Alkaloid CC 14, A-463
 Alkaloid CC 23, A-464
Papaver Alkaloid C₂₁H₂₁NO₆, A-466
Hortia Alkaloid C₂₂H₁₉N₃O₃, A-467
 Alkaloid CN1, A-468
 Alkaloid CN2, A-469
 Alkaloid CR1, A-470
 Alkaloid CRB 11, A-471
 Alkaloid CSA 3, A-472
Rauwolfia obscura Alkaloid D₁, A-473
Rauwolfia obscura Alkaloid D₂, A-474
Buxus Alkaloid D, A-475
Coelidium Alkaloid D, A-478
Dendrobates Alkaloid 285D, A-482
Dendrobates Alkaloid 307D, A-483
Leucojum Alkaloid D, A-484
Mantella Alkaloid 211D, A-485
Melodinus aeneus Alkaloid D, A-488
Narcissus Alkaloid D, A-489
Pandaca caducifolia Alkaloid D, A-490

Salicornia herbacea Alkaloid D, A-491
Stephania glabra Alkaloid D, A-492
 Alkaloid D1, A-493
 Alkaloid D2, A-494
 Alkaloid 267D, A-495
Buxus Alkaloid E, A-496
Coelidium Alkaloid E, A-497
Dendrobates Alkaloid 223E, A-499
Dendrobates Alkaloid 237E, A-500
Lycopodium Alkaloid E, A-501
Munafara sessilifolia Alkaloid E, A-502
Pandaca caducifolia Alkaloid E, A-503
Skytanthus acutus Alkaloid E, A-504
Sophora microphylla Alkaloid E, A-505
Voacanga thouarsii Alkaloid E, A-506
Buxus Alkaloid F, A-507
Dendrobates Alkaloid 223F, A-508
Mantella Alkaloid 249F, A-512
Mantella Alkaloid 265F, A-513
Mantella Alkaloid 281F, A-514
Pandaca caducifolia Alkaloid F, A-515
Lycopodium Alkaloid G, A-516
Mantella Alkaloid 207G, A-517
Pandaca caducifolia Alkaloid G, A-518
Voacanga thouarsii Alkaloid G, A-519
 Alkaloid 251G, A-520
 Alkaloid GB14, A-521
 Alkaloid GB15, A-522
 Alkaloid GB16, A-523
 Alkaloid GB17, A-524
 Alkaloid GB18, A-525
Voacanga thouarsii Alkaloid I, A-340
 C-Alkaloid I, A-280
Gentiana Alkaloid I, A-301
Gentiana Alkaloid II, A-526
Sternbergia Alkaloid III, A-527
Lycopodium Alkaloid IV, A-344
Solanum Alkaloid IV, A-528
Ungernia Alkaloid IV, A-529
Gentiana cruciata Alkaloid IVb, A-530
Lycopodium annotinum Alkaloid IX, A-531
 C-Alkaloid J, A-532
Mantella Alkaloid 207J, A-533
Mantella Alkaloid 235J, A-534
Mantella Alkaloid 235K, A-535
 Alkaloid K2, A-536
 Alkaloid K5, A-538
 Alkaloid K6, A-539
 Alkaloid K7, A-540
 Alkaloid K9, A-541
 Alkaloid K10, A-542
 Alkaloid K11, A-543
 C-Alkaloid L, A-544
Colubrina Alkaloid L, A-545
Mantella Alkaloid 251L, A-546
 Alkaloid L10, A-547
 Alkaloid LB1, A-552
 Alkaloid LE1, A-556
 Alkaloid LO1, A-557
 Alkaloid LO2, A-558
 Alkaloid LO3, A-559
 Alkaloid LS14, A-560
 Alkaloid LV1, A-561
 Alkaloid LV1, A-562
 Alkaloid LV2, A-563
 Alkaloid LV3, A-564
 Alkaloid LV4, A-565
Buxus Alkaloid M, A-566
Buxus Alkaloid M, A-567
 C-Alkaloid M, A-568
 Alkaloid MB 10, A-569
Buxus Alkaloid N, A-570
 Alkaloid Or₂, A-573
Baptisia Alkaloid P₂, A-574
Baptisia Alkaloid P₄, A-575
 C-Alkaloid P, A-576
Lupinus Alkaloid P1, A-577
Lupinus Alkaloid P2, A-578
Mucuna pruriens Alkaloid P, A-579
 Alkaloid PAR 1, A-580
 Alkaloid PAR 2, A-581
 Alkaloid PAR 3, A-582
 Alkaloid PPI, A-585

- Alkaloid PP1, A-586
 C-Alkaloid Q, A-587
Mucuna pruriens Alkaloid Q, A-588
Veratrum Alkaloid Q, A-589
 C-Alkaloid R, A-590
Fritillaria imperialis Alkaloid R, A-591
Mucuna pruriens Alkaloid R, A-593
Voacanga schweinfurthii Alkaloid S₂, A-595
Voacanga schweinfurthii Alkaloid S₃, A-596
 C-Alkaloid S, A-597
Mucuna pruriens Alkaloid S, A-598
 Alkaloid S-F, A-599
 Alkaloid TC-D, A-600
Veratrum Alkaloid U, A-601
 C-Alkaloid UB, A-602
Pachysandra Alkaloid V, A-322
Gentiana Alkaloid V, A-302
Helleborus Alkaloid V, A-305
Lycopodium Alkaloid V, A-310
Gentiana Alkaloid VI, A-604
Lycopodium annotinum Alkaloid VI, A-605
Pachysandra terminalis Alkaloid VI, A-607
Rhizophiala bifida Alkaloid VI, A-608
Lycopodium annotinum Alkaloid VII, A-606
Lycopodium annotinum Alkaloid VIII, A-609
Mucuna pruriens Alkaloid X, A-321
Veratrum Alkaloid X, A-339
Lycopodium Alkaloid X, A-311
Lycopodium annotinum Alkaloid XI, A-610
Lycopodium annotinum Alkaloid XII, A-611
Rhizophiala bifida Alkaloid XIII, A-613
Pachysandra terminalis Alkaloid XVI, A-614
Rhizophiala bifida Alkaloid XXV, A-615
Pachysandra terminalis Alkaloid XXVII, A-616
 Alkaloid YC2, A-618
Lycopodium clavatum Alkaloid, A-316
Lycopodium thyoides Alkaloid, A-317
Pancreatum sickenbergii Alkaloid, A-325
Papaver rhoas Alkaloid, A-326
Pseudaxinysa Alkaloid, A-327
Rhynchosia pyramidalis Alkaloid, A-328
Senecio borysthenicus Alkaloid, A-331
Strepeliopsis strepelioides Alkaloid, A-334
Thermopsis lanceolata Alkaloid, A-335
Ulex europaeus Alkaloid, A-336
Uncaria callophylla Alkaloid, A-337
Ungernia severtzovii Alkaloid, A-338
Zephyranthes Alkaloid, A-341
Aconitum nemorosum C₂₀ Alkaloid, A-265
Aconitum nemorosum C₂₃ Alkaloid, A-266
Erythrophleum C₂₄ Alkaloid, A-267
Aconitum kirinense Alkaloid, A-268
Aconitum koreanum Alkaloid, A-269
Alstonia constricta Alkaloid, A-270
Ammodendron karelinii Alkaloid, A-271
Artemisia rutifolia Alkaloid, A-273
Astragalus berterianus Alkaloid, A-274
Chamaecytisus Alkaloid, A-282
Colchicum luteum Alkaloid, A-283
Crotalaria retusa Alkaloid, A-284
Cytisus laburnum Alkaloid, A-285
Cytisus ruthenicus Alkaloid, A-286
Galanthus caucasicus Alkaloid, A-300
Gynura segetum Alkaloid, A-303
Hovea longipes Alkaloid, A-306
Lycopodium annotinum Alkaloid, A-314
Lycopodium annotinum Alkaloid, A-315
 Allothiobinupharidine, A-631
 Alvanidine, A-665
 Alvanine, A-666
 Ambaline, A-681
 Ambalinine, A-682
 Amianthine, A-691
 Ammocalline, A-929
 Ammorosine, A-931
 Amsoniaefoline, A-952
 Andrachnine, A-1001
 Angusteine, A-1010
 Anhwiaconitine, A-1022
 Ankoline, A-1046
 Annapawine, A-1048
 Antibiotic A 58365C, A-1106
 Antibiotic 69, A-1083
 Antibiotic 859A, A-1090
 Antibiotic 875A, A-1091
 Antibiotic 2197A, A-1094
 Antibiotic ASK 753, A-1117
 Antibiotic AS-N-7A, A-1118
 Antibiotic BA 181314A, A-1125
 Antibiotic BMY 42428, A-1138
 Antibiotic F 10C, A-1165
 Antibiotic FR 3383, A-1170
 Antibiotic L 11-1, A-1192
 Antibiotic LL-BO 2964, A-1200
 Antibiotic M 259, A-1202
 Antibiotic 610MF, A-1206
 Antibiotic MYC 8005, A-1210
 Antibiotic NP 522, A-1218
 Antibiotic NRCS 15, A-1219
 Antibiotic RP 6798, A-1233
 Antibiotic RP 13252, A-1234
 Antibiotic S 433, A-1235
 Antibiotic SF 1508, A-1253
 Antibiotic SF 2197B, A-1263
 Antibiotic TAN 883, A-1274
 Antibiotic TAN 1263, A-1275
 Antibiotic TAN 1280, A-1276
 Antibiotic WS 1921, A-1314
 Antibiotic XK 211, A-1317
 Antifungin, A-1326
 Apc, A-1332
 Aquaticine, A-1368
 Aquileginine, A-1370
 Arecolidine, A-1400
 Argentinamine, A-1411
 Aristo red, A-1424
 Aspidecelcine, A-1488
 Aspidecine, A-1489
 Aztequine, A-1617
 Beilupeimine, B-54
 Bervulcine, B-108
 Biflorine, B-124
 Biflorone, B-125
 Biochanin C, B-138
 Bishaconitine, B-179
 Bishatisine, B-180
 Bodamine, B-240
 Borbonicine, B-250
 Bractamine, B-272
 Bracteine, B-275
 4(7)-Bromo-2,3,7(2,3,4)-trichloro-1*H*-indole, B-354
 Brunfelsine, B-370
 Brunsdonnine, B-372
 Brunsvinine, B-373
 Bullatine A, B-405
 Bullatine D, B-406
 Buphacetine, B-412
 Buphanine, B-414
 Burmannaline, B-417
 Burmannine, B-418
 Buxalfine, B-452
 Buxazine, B-463
 Buxdeltine, B-464
 Buxetine, B-465
 Buxpapamine, B-479
 Buxpapine, B-480
 Buxpapinine, B-481
 Caffaeschizine, C-14
 Cairomycin C, C-17
 Calabacine, C-20
 Calabatine, C-21
 Calycine, C-44
 Calystegine, C-59
 Campestrine, C-64
 Camptine, C-67
 Campantine, C-68
 Caneine, C-79
 Canembine, C-81
 Caracurine I, C-109
 Caracurine III, C-112
 Caracurine IV, C-113
 Caracurine IX, C-114
 Caracurine VIII, C-116
 Carategine, C-118
 Cardiochryrine, C-158
 Carosidine, C-166
 Carosine, C-167
 Casimirine, C-175
 Cathalanceine, C-192
 Catharine, C-196
 Cathedulin Y7, C-204
 Cathedulin Y9, C-205
 Cathedulin Y10, C-206
 Cathidine A, C-210
 Cathidine B, C-211
 Cathindine, C-212
 Cavincidine, C-223
 Cavincine, C-224
 Ceanothine A, C-226
 Ceanothine C, C-227
 Celliamine, C-240
 Cevanine, C-299
 Chairamideine, C-337
 Chairamine, C-338
 Chandrine, C-346
 Chatinine, C-351
 Cherimoline, C-367
 Chinpeimine, C-388
 Chrycentrine, C-428
 Chuan-wu base A, C-435
 Chuan-wu base B, C-436
 Citriodorol, C-483
 Coclamine, C-549
 Coclilofine, C-550
 Coelibactin, C-561
 Columellarine, C-580
 Conchairamideine, C-597
 Conchairamine, C-598
 Concusconine, C-599
 Conolline, C-619
 Consolidine, C-623
 Convolvicine, C-631
 Copsomycin, C-637
 Coranicine, C-640
 Coriacenine E, C-650
 Coruscine, C-669
 Corycicine, C-672
 Corydicine, C-681
 Coumingaine, C-713
 Crinatine, C-747
 Crinosine, C-750
 Criophylline, C-751
 Crispamine, C-754
 Crispine, C-755
 Croceocurine, C-765
 Cruentine A, C-779
 Cruentine B, C-780
 Cuscamidine, C-817
 Cusconidine, C-819
 Cusconine, C-820
 Cygmine, C-923
 Daphnarcine, D-28
 Daphnicadine, D-39
 Daphnicaline, D-40
 Daphnicamine, D-41
 Daphnimacrine, D-58
 Daucine, D-87
 Delflexine, D-188
 Delfrenine, D-189
 Delorine, D-191
 Delphocine, D-193
 Delphoorine, D-194
 Delpyrrine, D-195
 Denudatidine, D-220
 Dichroidine, D-361
 3,4-Dihydro-*ar*-trimethoxy-1-methylisoquinoline, D-520
 Dinklageine, D-787
 Dithiocolchicine, D-872
 Dolichotheine, D-911
 Doryflavine, D-925
 Doryphorine, D-926
 Echinozolinone, E-29
 Elatrine, E-62
 Elliptimine, E-68
 Equisetonine, E-143
 Ercine, E-145

- Erythredine, E-209
 Erythreine, E-210
 Erythrine, E-211
 Euroline, E-298
 Evovoline, E-312
 Evoline, E-313
 Evomine, E-316
 Evopine, E-317
 Eximidine, E-320
 Fagarine III, F-4
 Fastudine, F-19
 Fastumine, F-20
 Fastusidine, F-21
 Fastusine, F-22
 Fastusine, F-23
 Fedamazine, F-32
 Fibraminine, F-50
 Fibramine, F-51
 Flexamine, F-87
 Flexine, F-89
 Floramultinine, F-91
 Fluggeine, F-95
 ψ -Fluorocurine, F-101
 C-Fluorocurinine, F-102
 Fluorosolimoines, F-105
 Franchetine, F-148
 Fredericamycin E, F-154
 Fritilline A, F-155
 Fritilline B, F-156
 Fritimine, F-157
 Fumarinine, F-177
 Fumarizine, F-179
 Fumvaine, F-192
 Galanthamidine, G-2
 Galipoidine, G-10
 Gentiananine, G-57
 Geotrichum alkaloid A 25822H, G-64
 Geralbine, G-66
 Gerrardamine, G-71
 Gerrardoline, G-73
 Gindaricine, G-77
 Glaucescine, G-89
 Glaucescinine, G-90
 Glauflavine, G-92
 Glenodinine, G-95
 Gnidine, G-134
 Gnidinine, G-135
 Graminifoline, G-154
 Grisonomycin, G-179
 C-Guaianine, G-189
 Guan-fu base C, G-194
 Guan-fu base D, G-195
 Guan-fu base E, G-196
 Guillauminine, G-221
 Hanadamine, H-45
 Hasunohanin, H-71
 Heimine, H-89
 Herpestine, H-151
 Himanthine, H-279
 Himgrine, H-286
 Hippacine, H-289
 Hippandrine, H-291
 Hippauline, H-292
 Hodorine, H-312
 Holacimine, H-313
 Holarrhessimine, H-319
 Holarrhine, H-320
 Homocinchonidine, H-328
 Homophleine, H-340
 Homothermopsine, H-349
 Hortesin, H-360
 Hunteriamine, H-369
 Hydrocotylamine, H-397
 4-(Hydroxymethyl)-6-methyl-2,3,5-pyridinetriol, H-608
 Hypotuberostemonine, H-774
 Imperoline, I-52
 Imperonine, I-53
 Indicamine, I-66
 Indicine, I-69
 1*H*-Indole-6,7-diol, I-94
 Insulamine, I-149
 Irenine, I-176
 Isochaksine, I-202
 Isochondocurarine, I-204
 C-Isodihydrotoxiferine I, I-227
 Isophysostigmine, I-265
 Isopyroine, I-295
 Isostemonidine, I-323
 Isostemonine, I-324
 Isotafine, I-327
 Isotuberostemonine, I-341
 Ivonine, I-351
 Janthitrem A, J-20
 Jaxartinine, J-33
 Kirigamycin C, K-47
 Kitagine, K-52
 Kokusaginine, K-57
 Koumicine, K-95
 Kouminidine, K-97
 Krelagine, K-98
 Kryptocurine, K-103
 Kurchenine, K-110
 Kurchicine, K-111
 Lallemanine, L-20
 Lamarckine, L-21
 Lamarckinine, L-22
 Lanceine, L-27
 Leontamine, L-95
 Leonuridine, L-100
 Leonurinine, L-101
 Leptactinine, L-113
 Lettocine, L-144
 Leurosivine, L-164
 Ligudentine, L-168
 Linantenine, L-179
 Lobine, L-222
 Lochnerivine, L-225
 Lochrovicine, L-226
 Lochrovidine, L-227
 Lochrovine, L-228
 Locoine, L-229
 Lolinidine, L-235
 Lophanterine, L-257
 Lophilacrine, L-258
 Lophiline, L-259
 Loveraine, L-274
 Lunamaridine, L-289
 Lunariamine, L-291
 Lunolone, L-297
 Lupilaxine, L-303
 Lurenine, L-304
 Lutenurine, L-309
 Lyceamine, L-318
 Lycocaroline, L-320
 Lycodiflexine, L-323
 Lycoserramine, L-349
 Lycoserrine, L-350
 Macranthine, M-9
 Macrophylline A, M-22
 Macrophylline, M-21
 Magnarcine, M-41
 Majorine, M-51
 Majorvine, M-52
 Marckidine, M-99
 Melinonine H, M-171
 Melinonine I, M-170
 Melinonine K, M-172
 Melinonine L, M-173
 Melinonine M, M-174
 Menismine, M-198
 Mesembryanthems, M-227
 Methyl 16,17-didehydromethoxy-19-methyl-ox-ayohimban-16-carboxylate, M-298
 Methylsuccirubine, M-562
 Minpeimine, M-632
 Minpeimine, M-633
 Mitraspecine, M-662
 Mitraversine, M-664
 Mooreine, M-701
 Mucudine, M-722
 Mucudanine, M-723
 Mucudanine, M-724
 Mucunadine, M-725
 Mucunine, M-726
 Muricine, M-740
 Myrmicarine 233A, M-799
 Naphthyridinomycin B, N-33
 Narcicriptine, N-39
 Narcisline, N-42
 Narzettine, N-52
 Natrine, N-54
 Neburnamine, N-83
Senecio doria Necine base, N-86
 Neflexine, N-88
 Nemorine, N-90
 Neochondocurarine, N-105
 Neodaphniphylline, N-109
 Neoleurocristine, N-123
 Neoleurosidine, N-124
 Neolitsinine, N-126
 Neoyuzurimine, N-150
 Nerifline, N-170
 Nerispine, N-172
 Nerundine, N-173
 Nikanine, N-215
 Nishindine, N-237
 Nordine, N-296
 α -Nupharidine, N-341
 β -Nupharidine, N-342
 Nupharine, N-343
 Nympeine, N-348
 Obscuridine, O-4
 Obscurine, O-6
 Ocodemerine, O-24
 Oleracins, O-80
 Oliverine, O-86
 Oreoline, O-104
 Ormosine, O-116
 Ornazidine, O-118
 Orobanchamine, O-122
 Otocamine, O-135
 Pachysterine, P-7
 Palybrine, P-13
 Paipunine, P-14
 Palidimine, P-21
 Palmatisine, P-36
 Papyraceabromine A, P-85
 Paricine, P-95
 Parvifagarine, P-100
 Paucicaline, P-127
 Pavanoline, P-136
 Peceylanine, P-147
 Pediculine, P-158
 Peimidine, P-166
 Peimiphine, P-167
 Peimitidine, P-169
 Peimunine, P-170
 Pelankine, P-172
 Pentalupine, P-227
 Pepuline, P-239
 Perimargine, P-259
 Perimivine, P-260
 Perividine, P-267
 Perosine, P-273
 Petomine, P-282
 Pictumine, P-420
 Pillijanine, P-423
 Pilosamycin, P-427
 Pimpinellidine, P-431
 Piperlongine, P-461
 Pircularin, P-478
 Pleiocraline, P-520
 Poetamine, P-531
 Poetaricine, P-532
 Poeticine, P-533
 Pogonopamine, P-534
 Pogonopeine, P-535
 Pogonopidine, P-536
 Pogonopine, P-537
 Powellidine, P-580
 Poweramine, P-582
 Powerchrine, P-583
 Precasine, P-592
 Precurarine, P-596
 Premavacurine, P-613
 Premmine, P-614
 Protochondocurarine, P-662
 Pruriendine, P-678

- Prurienine, P-679
 Prurieninine, P-680
 Pseudobaleabuxine F, P-704
 Pseudothiobinupharidine, P-729
 Pseudovinblastinediol, P-732
 Pseudowithanine, P-735
 Psilocauline, P-739
 Pteledimericine, P-751
 Pulchelline, P-786
 Punarnavine, P-797
 Punikathine, P-799
 Pycnarrhenamine, P-829
 Pycnarrhenine, P-830
 Pyoverdin 2112, P-836
 Pyoverdin A1, P-838
 Pyoverdin PaC, P-859
 Pyoverdin Pm, P-868
 Pyoverdin PS, P-871
 Pyoverdin PSS, P-873
 Pyrinadine C, P-927
 Pyrinadine D, P-928
 Pyrinadine E, P-929
 Pyrinadine F, P-930
 Pyrinadine G, P-931
 Raddeamine, R-7
 Raucaffridine, R-19
 Raunamine, R-23
 Rhinolidine, R-79
 Robecine, R-112
 Rovidine, R-137
 Rubiverine, R-145
 Rubrocurarines, R-149
 Rubrolone B, R-151
 Rulodine, R-158
 Ruticin, R-172
 Ruzorine, R-173
 Saipeimine, S-17
 Salicilobine, S-22
 Salicornine, S-23
 Saliherbine, S-24
 Sankhpuspine, S-63
 Sarothamine, S-88
 Saururidine, S-103
 Saururine, S-104
 Scabrine, S-113
 Schizoluteine, S-135
 β -Schizozogol, S-138
 Securinol C, S-208
 Sessilistemonine, S-272
 Sickenbergine, S-294
 Sinostemonine, S-317
 Solangustidine, S-342
 Solimocurarine, S-354
 Solimoesines, S-355
 Somniferine, S-361
 Somniferinine, S-363
 Somnine, S-364
 Songorinine, S-366
 Sonpeimine, S-368
 Sophochryrine, S-372
 Spartocytisine, S-386
 Spermatherine, S-396
 Sphaeranthine I, S-402
 Sphaeranthine IV, S-403
 Sphaeranthine VII, S-405
 Sphaeranthine VI, S-404
 Sphaeranthine, S-401
 Spireine, S-423
 Sprintillamine, S-482
 Sprintilline, S-483
 Squamigerine, S-486
 Stemonine, S-520
 Stemotuberine, S-525
 Stephanoline, S-538
 Sternine, S-551
 Stizolophine, S-554
 Streptovitacin D, S-574
 Strychnambarine, S-588
 Strychnobarididine, S-590
 Suaedin, S-615
 Subincanine, S-626
 Succirubine, S-629
 Suffruticodine, S-630
 Suffruticonine, S-631
 Synaine, S-655
 Syphilobin A, S-661
 Syphilobin F, S-662
 Tabernaegantididine, T-2
 Tecleanine, T-54
 Tejeramycin, T-61
 Tenuidine, T-78
 1,2,3,4-Tetrahydro-*ar*-hydroxy-*ar*-trimethoxy-2-methylisoquinoline, T-191
 1,2,3,4-Tetrahydro-*ar*-methoxy-1-methylisoquinoline, T-200
 Thaldimerine, T-302
 Thaliclavine, T-314
 Thalifabomine, T-324
 Thelepogidine, T-353
 Thesinicine, T-365
 Thiobideoxynupharidine, T-379
 Thiomycin, T-390
 Tienmulimine, T-402
 Tienmuliminine, T-403
 Tiliacoricidine, T-405
 Tiliandrine, T-409
 Tinosporine, T-414
 Tomentocurine, T-426
 Toxiferine III, T-438
 Toxiferine II, T-437
 Toxiferine VIII, T-439
 Toxiferine XII, T-440
 Tremine, T-448
 Trianthemine, T-455
 2,4,7(3,4,7)-Tribromo-3(2)-chloro-1*H*-indole, T-459
 2,3,5-Trihydroxy-6-methylpyridine, T-567
ar-Trimethoxy-1-methylisoquinoline, T-597
 Turkestanine, T-683
 Turumiquirensine, T-687
 Ursuline, U-49
 Vakatisine, V-2
 Vakatisinine, V-3
 Valivine, V-7
 Vallopurfine, V-12
 Vallotidine, V-13
 Vallotine, V-14
 Vasicinine, V-36
 C-Venezueline, V-51
 Veragenine, V-58
 Veralbidine, V-59
 Veraminine, V-67
 Veranovine, V-69
 Verareine, V-70
 Verine, V-75
 Verrucarin F, V-81
 Verrucarin G, V-82
 Verticilline, V-93
 Vinaphamine, V-108
 Vinaspine, V-109
 Vincamicine, V-114
 Vincanovine, V-116
 Vincolidine, V-123
 Vincovalicine, V-127
 Vindolidine, V-131
 Vinomine, V-134
 Vinosidine, V-135
 Vinsedicine, V-137
 Vinsedine, V-138
 Virgilidine, V-151
 Viridomycin B, V-157
 Viridomycin C, V-158
 Vitricine, V-165
 Voacaminine, V-169
 Vulracine, V-196
 Withananine, W-18
 Withanine, W-19
 Wuchuyine, W-23
 Xanthicin, X-4
 C-Xanthocurine, X-8
 Xanthofagarine, X-9
 Yemensine, Y-8
 Yuzurimine D, Y-24
 Zeagenin, Z-10
 Zeravschanidine, Z-17

DICTIONARY OF ALKALOIDS SECOND EDITION ON CD-ROM

This CD-ROM allows users to search the entire contents of the Dictionary in a fast and efficient manner. Dictionary of Alkaloids can be searched by structure, name, CAS number, molecular formula and numerous other text and numerical fields. The CD-ROM also contains an extensive help system.

Installation

Insert the CD in the CD-ROM drive and click on the Start button and select Run. In the command line type D:\setup.exe (where D is your CD-ROM drive letter) then click OK. If you would like a shortcut added to your start menu, check the appropriate box.

You have two Installation options:

Standard installation: The default installation copies the program files only to your computer's hard drive. Use this option if your computer has limited hard disk space. You will need to insert the CD every time you use the Dictionary of Alkaloids.

Hard disk installation: For faster access you may copy the entire database to your hard drive by checking the option Copy database to HD. This will require approximately 155MB of hard disk space.

If you do not already have Adobe® Acrobat® Reader® installed on your PC you may install it at this point to access this manual and the database introduction in PDF format if you wish.

Check that the path to your CD-ROM drive is correct and then press OK.

You will be prompted to select a folder in which to install the software. If you wish the program to be installed to the default folder C:\Program Files\CRC Press\Alkaloids, just click the Install button. Otherwise, type a new folder location and then click Install.

Updated Data

If you wish to receive regularly updated data on all types of natural products, then why not subscribe to *Dictionary of Natural Products*? Using similar search software to Dictionary of Alkaloids, this comprehensive database of natural products is updated every 6 months.

Please contact the Taylor and Francis Online Subscriptions department for details:

E-mail: e-reference@taylorandfrancis.com

URL: www.chemnetbase.com

Technical Support

Please contact:

Technical Support Desk

Electronic Publishing Division

CRC Press LLC

6000 Broken Sound Parkway NW

Boca Raton FL 33487 USA

Tel: 888 318 2367

Tel: +1 561 361 6020 (Outside US)

Fax: +1 561 361 6075

E-mail: techsupport@crcpress.com

Dictionary of Alkaloids

SECOND EDITION

with CD-ROM

Drawn from the venerable *Dictionary of Natural Products*, the second edition of the **Dictionary of Alkaloids** categorizes all known alkaloids to present their concise chemical, structural, and physical data.

Now More than Double the Size of the Previous Edition

The dictionary embraces a wide definition of alkaloids. Double in number from the prior edition, the number of alkaloids documented is now over 20,000. For this new edition, every existing entry has been inspected for accuracy, and the great majority updated.

The **Dictionary's** extensive introduction has been completely rewritten and expanded to give a concise, authoritative, and fully referenced account of the different types of alkaloids. Geoffrey Cordell, eminent alkaloid chemist and Editor-in-Chief of the prestigious series, *The Alkaloids, Chemistry and Biology*, has provided a foreword that sets the scene for this fascinating group of natural products.

Includes a Fully Searchable CD-ROM

Accompanying the dictionary is a fully searchable CD-ROM that mirrors the book's complete contents and adds valuable functionality. The choice of media allows researchers to conduct high-speed electronic searches, while also giving them the ability to leisurely thumb through the book's printed pages. With either format, researchers and readers will find access to data unparalleled both in scope and depth.

Also available both in print and electronically from CRC Press

Dictionary of Natural Products
Dictionary of Marine Natural Products
Dictionary of Food Compounds
Dictionary of Carbohydrates
Polymers: A Property Database
Properties of Organic Compounds

Visit www.CHEMnetBase.com for a complete listing of CRC's online databases

 **CRC Press**
Taylor & Francis Group
an informa business
www.crcpress.com

6000 Broken Sound Parkway, NW
Suite 300, Boca Raton, FL 33487
270 Madison Avenue
New York, NY 10016
2 Park Square, Milton Park
Abingdon, Oxon OX14 4RN, UK

